#### NOTE



# Lectotypification of Memecylon flavescens Gamble (Memecylaceae DC.)

Moumita Das Das<sup>(1)</sup>, Debabrata Maity<sup>(2\*)</sup> and Arabinda Pramanik<sup>(1)</sup>

1. Central Botanical Laboratory, Botanical Survey of India, A. J.C. Bose Indian Botanic Garden, Howrah 711 103.

2. Taxonomy and Biosystematics Laboratory, Department of Botany, University of Calcutta, 35, Ballygunge Circular Road,

Kolkata 700 019; W.B. \* Corresponding author. Email: debmaity@yahoo.com

(Manuscript received 29 November 2012; accepted 24 May 2013)

ABSTRACT: The Lectotype of *Memecylon flavescens* Gamble is designated here.

KEY WORDS: Lectotypification, Memecylon flavescens Gamble, Memecylaceae.

## INTRODUCTION

The species Memecylon flavescens Gamble was described by J. S. Gamble (1919) under Melastomataceae Juss. - Memecyleae based on his own collections from the evergreen shola forest of Kundahs country of Nilgiri Hills, in the vicinity of Avalanche and Sispara under elevations of 2300-3460 m. The status of 'Memecyleae' has been under much discussion in the past and present (APG II, 2003; Renner, 2004; APG III, 2009; Takhtajan, 2009). In the present paper the 'Memecyleae' is maintained as a distinct family Memecylaceae DC. after the treatment of Renner (2004) and Takhtajan (2009). The family has close affinities to the Melastomataceae but differs in cryptocotylar germination of seedlings, leaf venation, presence of sclerids in leaves, stomatal nature, number and size of the seeds, presence of an elliptic terpenoid-producing gland on the staminal connective and fibrous endothecium.

*Memecylon flavescens* is very close to *Memecylon lushingtonii* Gamble described from Tamil Nadu. Both the species are small tree with knobby, 4-angular branchlets and have similar sized leaves, axillary flower clusters, very short (1–2 mm) pedicels, obscure disc rays and blue petals but the former is distinguished from *M. lushingtonii* by its small, oblanceolate, pale yellowish leaves.

In the protologue of *M. flavescens*, Gamble cited four specimens viz., *Gamble 14268*, *Gamble 16161*, *Gamble 16168* and *Gamble 20581* but did not designated the Holotype. Thus these specimens are to be treated as syntypes. All these specimens are kept in K (acronym after Holmgren *et al.*, 1990). However, in the course of study of different herbaria, one set of duplicate of the former three are found in CAL (!) (acronym after Holmgren *et al.*, 1990) and that of the last one is located in MH(!) (acronym after Holmgren *et*  *al.*, 1990). After careful study of all syntype materials in accordance to the Art 9.2 & 9.10 of the International Code of Botanical Nomenclature (McNeill *et al.*, 2006), *Gamble* 14268 (K) is designated here as the Lectotype of *Memecylon flavescens* Gamble. This specimen is most complete and better preserved specimen along with a hand drawing of a floral dissection on the herbarium sheet made by Gamble himself. The isolectotype at CAL is in a comparatively poor state.

#### **TAXONOMIC TREATMENT**

*Memecylon flavescens* Gamble, Bull. Misc. Inform. Kew 1919: 226.1919 & Fl. Pres. Madras 1:503. 1919 (1: 355. 1957, rep. ed.).

Syntypes: India, Tamil Nadu: Nilgiris District, above Sispara, 7000 ft. (2100 m), May,1884, *Gamble* 14268 (K, CAL); l.c., Avalanche, 7500 ft., May 1885, *Gamble* 16161(K, CAL); l.c., 7500 ft., May 1885, *Gamble* 16168 (K, CAL); l.c., Kundahs, 7000 ft., May 1889, *Gamble* 20581 (K, MH).

Lectotypus: (designated here) (Fig. 1). *Gamble* 14268 (K, CAL, isolecto.)

Flowering and fruiting: February–May.

Distribution: India: W. Ghats: Tamil Nadu, Kanyakumari, Nilgiris; Kerala, Avalanche, Kundah, and Sispara, 1500-3460m (Gamble, 1919; Murugan *et al.*, 2002). Grows in evergreen forests and sholas (Fig. 2).

Note: This species is endemic to Western Ghats, Nilgiris in Tamil Nadu. It is under the Endangered (EN) category of IUCN due to habitat destruction (Ahmedullah and Nayar, 1986; Nayar and Sastry, 1990; Murugan *et al.*, 2002). However, updating of current conservation status of the species in the places of distribution is an urgent need.





Fig.1. Lectotype image of Memecylon flavescens Gamble (Memecylaceae).



#### ACKNOWLEDGEMENT

The authors are grateful to the Director of Botanical Survey of India for providing necessary facilities and also thankful to Prof. G. G. Maiti, University of Kalyani and Dr. Subir Bandyopadhyay, Botanical Survey of India for their help. Sincere thanks are also due to Dr. V. P. Prasad, Scientist, BSI for his suggestions.

## LITERATURE CITED

- APG II. 2003. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. Bot. J. Linn. Soc. 141: 399–436. doi: 10.1046/j.1095-8339.2003.t01-1-00158.x
- **APG III.** 2009. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. Bot. J. Linn. Soc. **161**: 105–121. doi: 10.1111/j.1095-8339.2009.00996.x
- Ahmedullah, M. and M. P. Nayar. 1986. Endemic plants of the Indian region, Vol. 1. Botanical Survey of India, Calcutta, pp. 109.
- Gamble, J. S. 1919. XII. Decades Kewenses: Plantarum Novarum in Herbario Horti Regii Conservatarum. Bull. Misc. Inform. Kew 1919: 226.
- Holmgren, P. K., N. H. Holmgren and L. C. Barnett (eds.). 1990. Index Herbariorum Part I: The Herbaria of the World (8<sup>th</sup> edition), New York, U.S.A.
- McNeill, J., F. R. Barrie, H. M. Burdet, V. Demoulin, D. L. Hawksworth, K. Marhold, D. H. Nicolson, J. Prado, P. C. Silva, J. E. Skog, J. H. Wiersema and
  N. J. Turland (eds). 2006. International Code of Botanical Nomenclature (Vienna Code). Regnum Vegetabile 146. Ruggel: Gantner.
- Murugan, C., V. S. Manickam, M. M. Josephine and V. Sundaresan. 2002. Extended distribution of two rare and endangered taxa from Tirunelveli Hills, Western Ghats, Tamil Nadu. J. Bombay Nat. Hist. Soc. 99: 545–546.
- Nayar, M. P. and A. R. K. Sastry. (eds), 1990. Red Data Book of Indian Plants, Vol. 3. Botanical Survey of India, pp. 179.
- Renner, S.S. 2004. Memecylaceae. *In*: N, Smith, S.A. Mori, A. Henderson, D.W. Stevenson, and SV Heald, (eds). Flowering plants of the Neotropics. The New York Botanical Garden, Princeton University Press, Princeton. pp. 246–247.
- Takhtajan, A. 2009. Flowering Plants. 2<sup>nd</sup> Ed. Springer. pp. 339–340.




Fig. 2. Black dots showing the distribution of *Memecylon flavescens* Gamble.



# Memecylon flavescens (谷木科)的選定模式

#### Moumita Das Das<sup>(1)</sup>, Debabrata Maity<sup>(2\*)</sup> and Arabinda Pramanik<sup>(1)</sup>

- 1. Central Botanical Laboratory, Botanical Survey of India, A. J.C. Bose Indian Botanic Garden, Howrah 711 103.
- 2. Taxonomy and Biosystematics Laboratory, Department of Botany, University of Calcutta, 35, Ballygunge Circular Road, Kolkata 700 019; W.B.
- \* 通信作者。Email: debmaity@yahoo.com
- (收稿日期:2012年11月29日;接受日期:2013年5月24日)
- 摘要:本文指定Memecylon flavescens的選定模式。
- 關鍵詞:選定模式、Memecylon flavescens、谷木科。