



Revista Iberoamericana de Micología

www.elsevier.es/reviberoammicol



Original

Molecular phylogeny of asexual entomopathogenic fungi with special reference to *Beauveria bassiana* and *Nomuraea rileyi*

Nageswara Rao Reddy Neelapu^a, Annette Reineke^b, Uma Maheswara Rao Chanchala^c and Uma Devi Koduru^{d,*}

^aDepartment of Bioinformatics, GITAM Institute of Science, GITAM University, Visakhapatnam, India

^bForschungsanstalt Geisenheim, Fachgebiet Phytomedizin, Geisenheim, Germany

^cBioinformatics Division, Advanced Technology Centre, Tata Consultancy Services, Deccan Park, Hyderabad, India

^dDepartment of Botany, Andhra University, Visakhapatnam 530003, Andhra Pradesh, India

ARTICLE INFO

Article history:

Received June 18, 2007

Accepted September 9, 2008

Keywords:

β-tubulin gene sequence

Beauveria bassiana

Cladistic analysis

Clavicipitaceae

Cryptic speciation

Hyphomycetes

Hypocreales

Mitosporic fungi

Nomuraea rileyi

rRNA gene sequence

Species complex

ABSTRACT

The phylogenetic lineage, taxonomic affiliation and interrelationships of important asexual entomopathogenic fungal genera were studied using the sequences of partial regions of β-tubulin and rRNA genes. The species structures of *Beauveria bassiana* and *Nomuraea rileyi* were also investigated. A total of 147 fungal entries covering 94 species were analysed. Phylogenetic analysis placed all the asexual entomopathogenic fungal species analysed, in the family Clavicipitaceae of the order Hypocreales of Ascomycota. Deep phylogenetic lineages were observed in *B. bassiana* iterating the complex nature of this species. Some of the isolates assigned to this species separated out more distinctly than morphologically distinguishable genera. Cryptic speciation was also evident in *N. rileyi*. It is concluded that the asexual fungi with entomopathogenic habit arose from a single lineage in sexual Clavicipitaceae.

© 2007 Revista Iberoamericana de Micología. Published by Elsevier España, S.L. All rights reserved.

Filogenia molecular de los hongos entomopatógenos asexuales con especial referencia a *Beauveria bassiana* y *Nomuraea rileyi*

RESUMEN

Basándose en el análisis de las secuencias de regiones parciales de los genes β-tubulina y ARN ribosómicos se estudió la filogenia, la taxonomía y las interrelaciones de importantes géneros de hongos entomopatógenos asexuales. Se estudio también la estructura de las especies *Beauveria bassiana* y *Nomuraea rileyi*. Se analizó un total de 174 entradas fúngicas que representaban 94 especies. El análisis filogenético demostró que todas las especies de hongos entomopatógenos asexuales incluidas en el estudio pertenecían a la familia Clavicipitaceae, del orden Hypocreales de los Ascomycota. Se observaron diferentes linajes dentro de *B. bassiana*, lo que demostró la complejidad de dicha especie. Algunos de los aislamientos de dicha especie demostraron estar filogenéticamente más separados que determinados géneros morfológicos. Dentro de la especie *N. rileyi* se pudo observar también la presencia de especiación críptica. Se concluye que los hongos asexuales entomopatógenos han evolucionado de un linaje simple dentro de la familia Clavicipitaceae.

© 2007 Revista Iberoamericana de Micología. Publicado por Elsevier España, S.L. Todos los derechos reservados.

Palabras clave:

Secuencias del gen de la β-tubulina

Beauveria bassiana

Análisis cladístico

Clavicipitaceae

Especiación críptica

Hyphomycetes

Hypocreales

Hongos mitospóricos

Nomuraea rileyi

Secuencias de los genes ARN ribosómicos

Complejos de especies

*Corresponding author.

E-mail address: umadevikoduru@gmail.com (U.D. Koduru).

Mitosporic fungi with entomopathogenic habit have been widely used as pesticides because of their potential in biological control of insect pests. Due to their predominant asexual nature, the taxonomic status of these fungi is not certain. The sexual stages (teleomorphs) of some of these genera have been discovered. The teleomorphs are ascomyceteous species. The asexual entomopathogenic fungal species were found to be closely related to each other in several molecular phylogenetic studies^{3,4,8,9,21,22,28,31}. Further, these studies also confirmed the relation between the anamorph (asexual) and the teleomorph species. Based on these evidences, these fungi are recognized as hypocrealean ascomycetes belonging to the family Clavicipitaceae. However, no phylogenetic study of these mitosporic fungi was done including representative members of sexual Ascomycota to view their position in context to other members of Ascomycota. We took up such a study using β -tubulin and rRNA gene sequences. Many of the economically important hyphomycete entomopathogenic fungi are complex species and the boundaries between different morphologically distinguishable species are not validated. Therefore, inter-relationships between some of the important entomopathogenic genera were also investigated including a sample of isolates of complex species – *Beauveria bassiana* and *Nomuraea rileyi*, two of the five hyphomycetous asexual fungal species registered as bioinsecticides²⁹.

Of the several genes used to serve as 'zeit gebers' or molecular clocks for studying phylogenetic relationships among organisms, the highly conserved and functionally essential β -tubulin gene has been found to be very useful for investigating relationships between fungi at all levels – from studies of complex species groups to deep (geological) time phylogenetic investigations^{5,8,12,16,18-20}. Since it is a protein-coding gene, it has some advantages over the more often used ribosomal RNA gene sequences: the alignment of the sequences is less problematic and the third nucleotide can provide a relatively good estimate of the neutral substitution rate³³. rRNA genes are also frequently used to establish ordinal relationships among fungi^{1,9,17}. Both these gene sequences were employed in the present study.

Material and methods

Fungal sample

A fungal sample constituting 147 fungal entries representing 94 species, 21 of which are asexual and entomopathogenic, was analysed (Table 1). Among them, 28 isolates were sequenced for the partial region of β -tubulin gene and the rest of the sequences were obtained from GenBank database entries (<http://www.ncbi.nlm.nih.gov/Genbank/index.html>) (Table 1).

The partial regions of the three genes, namely β -tubulin, small and large subunits of mitochondrial rRNA genes, were amplified as described by Uma Devi et al³⁵. The fungal sequences related to these sequences were retrieved from Genbank using *B. bassiana* sequences with accession numbers AJ312228 (β -tubulin) and AB027336 (rRNA) as query sequences in blast search with default parameters. An *E*-value cutoff of $2e^{-119}$ (β -tubulin) and $2e^{-95}$ (rRNA) was used to assign homology. In addition, partial nucleotide sequences of β -tubulin gene of seven *N. rileyi* isolates⁸ were included in the sample.

The sequences were initially aligned and later realigned until good alignment was obtained using the multiple alignment program AlnExplorer of MEGA ver. 3.0¹⁴. Flanking regions that were not part of significant multiple alignment were trimmed off. The consistently aligned regions that have shared/derived characters of the homologous regions were used as feed for various tree construction methods. DNA sequence alignments were submitted in TreeBASE (<http://www.treebase.org/treebase/>), a relational database designed

to manage and explore information on phylogenetic relationships with accession number SN2616.

To infer the position of asexual entomopathogenic fungi in the fungal kingdom, a data set including partial amino acid sequences of the β -tubulin gene with 44 entries (Table 1) was analysed. To infer the position of asexual entomopathogenic fungi in the order Hypocreales, a data set including partial nucleotide sequences of the rRNA gene with 66 entries (Table 1) was analysed. To examine the interrelationship among the entomopathogenic asexual fungal genera and members within a species, a data set comprising 43 partial nucleotide sequences with several isolates of the two complex species *B. bassiana* and *N. rileyi* was analysed (Table 1). To examine the cryptic speciation in the complex species *B. bassiana*, a data set comprising seven partial nucleotide sequences of three genes, namely β -tubulin, large and small subunit of mitochondrial rRNA genes, was analysed (Table 1).

The default parameters taken for pairwise alignment were as follows: gap opening penalty (GOP) = 10/15, gap extension penalty (GEP) = 0.1/6.6 and for multiple alignment, GOP=10/15, GEP = 0.2/6.6 with a gap separation distance = 4, transition weight of 0.5/0.5, Gonnet protein/IUB DNA weight matrix and a delay divergent cutoff % = 30/30.

The statistical procedures of minimum evolution (ME), maximum parsimony (MP) and neighbor joining (NJ) of the program MEGA ver. 3.0¹⁴ were used. Distances for neighbor joining tree were calculated under the Kimura 2-parameter model. In MP and ME analyses, a single heuristic search was performed with Close Neighbor Interchange (CNI) branch swapping. For ME, NJ tree was used as the starting tree for heuristic search. For MP, stepwise addition procedure was employed. The maximum number of trees that could be saved during the heuristic search procedures was set to 1000. When multiple trees were found under ME and MP procedures, a single consensus tree was created. Confidence levels for individual branches were determined by bootstrap analysis for 1000 replicates⁶. To address the possible misleading effects of the data set phylogenetic relationships were then inferred using selected evolutionary model and the heuristic search option for maximum likelihood implemented in PAUP ver. 4b. We implemented Kishino-Hasegawa (KH) test¹³ using normal approximation (two-tailed test) to statistically evaluate tree topology against a series of topologies. In addition, trees were inferred with Bayesian inference using MrBayes, ver. 3.04b¹⁰. A Metropolis-Coupled Markov Chain Monte Carlo (MCMCMC) starting tree was initiated at random and run for 10 million generations. Trees were sampled each 1,000 cycles. Four chains were run simultaneously, of which three were heated and one was cold. Stationarity of the log likelihoods was monitored to verify convergence by 1,000,000 cycles. A consensus tree was made with multiple phylogenetic trees to determine the posterior probabilities at different nodes. To assess cryptic speciation in the sequence data of *B. bassiana*, the maximum parsimony analysis (PAUP ver. 4b) was done. A heuristic search analysis was run with 'tree-bisection-reconnection' (TBR) branch swapping with accelerated transformation (ACCTRAN) optimization to infer branch lengths with MULTREES option on, ADDSEQ set at random and 10 randomized replicates. All characters were weighed equally. A single tree was described when multiple trees were found.

Results

Phylogenetic position of entomopathogenic fungi in the fungal kingdom

The phylogenetic trees generated based on β -tubulin sequence-aligned data set had similar topologies in both distance and parsimony methods (ME, MP and NJ) but differed in their branching order. Topological similarity of the trees obtained by different

Table 1
Details of the entomopathogenic mitosporic fungi and other fungi used in the analysis

Species	Genes					
	β-tubulin		rRNA		S rRNA	L rRNA
	Isolate acc. no. ^c	GenBank acc. no.	Isolate acc. no.	GenBank acc. No.	GenBank acc. no.	GenBank acc. no.
<i>Beauveria bassiana</i>	ITCC-913 AJ312228		IFO4848	AB027336	n.a.	n.a.
	ARSEF-501 ^a	DQ092752	n.a.	n.a.	n.a.	n.a.
	ARSEF-1166 ^a	DQ092753	n.a.	n.a.	n.a.	n.a.
	ARSEF-1315 ^a	DQ090731	n.a.	n.a.	n.a.	n.a.
	ARSEF-1316 ^a	DQ090731	n.a.	n.a.	n.a.	n.a.
	ARSEF-3041 ^a	DQ090729	n.a.	n.a.	n.a.	n.a.
	ARSEF-1169	DQ090741	n.a.	n.a.	n.a.	n.a.
	ITCC-4521 ^a	DQ090736	n.a.	n.a.	n.a.	n.a.
	BB 4 ^a	DQ090732	n.a.	n.a.	n.a.	n.a.
	II.a.4. ^a	DQ090735	n.a.	n.a.	n.a.	n.a.
	ARSEF-1533 ^{a,b}	DQ090711	n.a.	n.a.	DQ090724	DQ090748
	ARSEF-1535 ^{a,b}	DQ090712	n.a.	n.a.	DQ090725	DQ090749
	ARSEF-1537 ^{a,b}	DQ090713	n.a.	n.a.	DQ090750	DQ090718
	ARSEF-1538 ^{a,b}	DQ090714	n.a.	n.a.	DQ090720	DQ090719
	ARSEF-1540 ^{a,b}	DQ090715	n.a.	n.a.	DQ090726	DQ090721
	ARSEF-1635 ^{a,b}	DQ090717	n.a.	n.a.	DQ090727	DQ090722
	ARSEF-1640 ^{a,b}	DQ090716	n.a.	n.a.	DQ090728	DQ090723
<i>Beauveria brongniartii</i>	Bbr-9301	AAQ96774	BCMU BB05	AB237659	n.a.	n.a.
<i>Beauveria brongniartii</i>	ARSEF-2660	DQ090733	n.a.	n.a.	n.a.	n.a.
<i>Beauveria amorpha</i>	ARSEF-2641	DQ090742	n.a.	n.a.	n.a.	n.a.
<i>Beauveria</i> sps.	ARSEF-1198	DQ090743	n.a.	n.a.	n.a.	n.a.
<i>Beauveria caledonica</i>	n.a.	n.a.	CCRC 32867	AY245650	n.a.	n.a.
<i>Nomuraea rileyi</i>	UFIF-178	DQ09608	NBRC 8560	AB100361	n.a.	n.a.
	I.a.3. ^{a,b}	DQ090744	n.a.	n.a.	n.a.	n.a.
	II.a.1.1 ^{a,b}	DQ090746	n.a.	n.a.	n.a.	n.a.
	II.b.5. ^{a,b}	DQ090745	n.a.	n.a.	n.a.	n.a.
	II.b.7. ^{a,b}	DQ090747	n.a.	n.a.	n.a.	n.a.
	ARSEF-762 ^a	DQ090738	n.a.	n.a.	n.a.	n.a.
	ARSEF-2345	AF399735	n.a.	n.a.	n.a.	n.a.
	ARSEF-935	AF399730	n.a.	n.a.	n.a.	n.a.
	MSU-27	AF399731	n.a.	n.a.	n.a.	n.a.
	ARSEF-711	AF399729	n.a.	n.a.	n.a.	n.a.
	ARSEF-380	AF399720	n.a.	n.a.	n.a.	n.a.
	ARSEF-482	AF399732	n.a.	n.a.	n.a.	n.a.
	ARSEF-2492	AF399728	n.a.	n.a.	n.a.	n.a.
<i>Metarhizium anisopliae</i>	n.a.	AY995134	NBRC 5940	AB250411	n.a.	n.a.
	ARSEF-538 ^a	DQ090737	n.a.	n.a.	n.a.	n.a.
<i>Metarhizium flavoviride</i>	n.a.	n.a.	ARSEF 1764	AF280632	n.a.	n.a.
<i>Paecilomyces farinosus</i>	ARSEF- 648	DQ079607	IFO 8581	AB080090	n.a.	n.a.
<i>Paecilomyces fumosoroseus</i>	ARSEF-1626 ^a	DQ090739	BCMU PF01	AB250413	n.a.	n.a.
<i>Paecilomyces reniformis</i>	ARSEF-577	DQ079606	n.a.	n.a.	n.a.	n.a.
<i>Paecilomyces nostocoides</i>	n.a.	n.a.	JCM 8437	AB104884	n.a.	n.a.
<i>Paecilomyces viridis</i>	n.a.	n.a.	CBS 348.65	AB023949	n.a.	n.a.
<i>Paecilomyces carneus</i>	n.a.	n.a.	JCM 6870	AB103379	n.a.	n.a.
<i>Paecilomyces marquandii</i>	n.a.	n.a.	NBRC 31966	AB114223	n.a.	n.a.
<i>Paecilomyces isarioides</i>	n.a.	n.a.	IFO 7562	AB023944	n.a.	n.a.
<i>Paecilomyces cateniannulatus</i>	n.a.	n.a.	BCMU IF05	AB263742	n.a.	n.a.
<i>Paecilomyces tenuipes</i>	n.a.	n.a.	BCMU IJ06	AB086208	n.a.	n.a.
<i>Paecilomyces javanicus</i>	n.a.	n.a.	NBRC 8297	AB099944	n.a.	n.a.
<i>Paecilomyces amoeneroseus</i>	n.a.	n.a.	CBS 107.73	AY526464	n.a.	n.a.
<i>Paecilomyces lilacinus</i>	n.a.	n.a.	BCMU PL01	AB124670	n.a.	n.a.
<i>Cordyceps dipterigena</i>	n.a.	n.a.	CCRC 35725	AY245655	n.a.	n.a.
<i>Cordyceps japonica</i>	n.a.	n.a.	IFO 9647	AY245669	n.a.	n.a.
<i>Cordyceps roseostromata</i>	n.a.	n.a.	ARS 4870	AY245662	n.a.	n.a.
<i>Cordyceps scarabaeicola</i>	n.a.	n.a.	ARS 5689	AY245663	n.a.	n.a.
<i>Cordyceps ochraceostromata</i>	n.a.	n.a.	ARS 5691	AY245660	n.a.	n.a.
<i>Cordyceps subsessilis</i>	n.a.	n.a.	CBS 305.95	AY245665	n.a.	n.a.
<i>Cordyceps memorabilis</i>	n.a.	n.a.	CCRC 32218	AY245658	n.a.	n.a.
<i>Cordyceps myrmecophila</i>	n.a.	n.a.	CCRC 35726	AY245672	n.a.	n.a.
<i>Cordyceps valliformis</i>	n.a.	n.a.	DAOM 196368	AY245648	n.a.	n.a.
<i>Cordyceps gracilioides</i>	n.a.	n.a.	ARS 5696	AY245652	n.a.	n.a.
<i>Cordyceps sphingum</i>	n.a.	n.a.	CBS 114.22	AY245673	n.a.	n.a.
<i>Cordyceps caloceroides</i>	n.a.	n.a.	CBS 250.76	AY245654	n.a.	n.a.
<i>Cordyceps ophioglossoides</i>	n.a.	n.a.	NBRC 8992	AB113827	n.a.	n.a.
<i>Cordyceps sinensis</i>	n.a.	n.a.	n.a.	AB067701	n.a.	n.a.
<i>Cordyceps heteropoda</i>	n.a.	n.a.	BCMU CH01	AB084157	n.a.	n.a.
<i>Cordyceps jezoensis</i>	n.a.	n.a.	n.a.	AB027319	n.a.	n.a.
<i>Cordyceps chlamydo sporia</i>	n.a.	n.a.	NBRC 9249	AB100362	n.a.	n.a.
<i>Cordyceps formicarum</i>	n.a.	n.a.	BCMU CF02	AB222679	n.a.	n.a.
<i>Cordyceps tricentri</i>	n.a.	n.a.	n.a.	AB027330	n.a.	n.a.
<i>Cordyceps khaoyaiensis</i>	n.a.	n.a.	885.1	AF327393	n.a.	n.a.

(Continued in next page)

Table 1
Details of the entomopathogenic mitosporic fungi and other fungi used in the analysis (continuation)

Species	Genes					
	β-tubulin		rRNA		S rRNA	L rRNA
	Isolate acc. no. ^c	GenBank acc. no.	Isolate acc. no.	GenBank acc. No.	GenBank acc. no.	GenBank acc. no.
<i>Cordyceps takaomontana</i>	n.a.	n.a.	n.a.	AB044631	n.a.	n.a.
<i>Cordyceps militaris</i>	ARSEF-4082 ^a	DQ090734	CBS 178.59	AY245671	n.a.	n.a.
<i>Verticillium leptobactrum</i>	n.a.	n.a.	IAM 14729	AB214657	n.a.	n.a.
<i>Verticillium fungicola</i>	n.a.	n.a.	NBRC 30728	AB111494	n.a.	n.a.
<i>Verticillium insectorum</i>	n.a.	n.a.	IAM 14728	AB214655	n.a.	n.a.
<i>Epichloe typhina</i>	PRG85	X52616	MAFF 306230	AB105952	n.a.	n.a.
<i>Epichloe festucae</i>	F11	AY722412.1	n.a.	n.a.	n.a.	n.a.
<i>Claviceps purpurea</i>	n.a.	n.a.	NBRC 5782	AB160991	n.a.	n.a.
<i>Claviceps paspali</i>	n.a.	n.a.	MAFF 306124	AB250409	n.a.	n.a.
<i>Claviceps africana</i>	n.a.	n.a.	NE1	AF281176	n.a.	n.a.
<i>Claviceps sorghicola</i>	n.a.	n.a.	MAFF 306574	AB250410	n.a.	n.a.
<i>Claviceps panicoides</i>	n.a.	n.a.	MAFF 511350	AB255604	n.a.	n.a.
<i>Acremonium chrysogenum</i>	H780	X72789	n.a.	n.a.	n.a.	n.a.
<i>Acremonium coenophialum</i> = <i>Neotyphodium coenophialum</i>	e19	X56847	n.a.	n.a.	n.a.	n.a.
<i>Neotyphodium uncatum</i>	n.a.	n.a.	NBRC 32642	AB102783	n.a.	n.a.
<i>Hypomyces odoratus</i>	IMI 267134	Y12256	n.a.	n.a.	n.a.	n.a.
<i>Hypomyces chrysospermus</i>	n.a.	n.a.	n.a.	M89993	n.a.	n.a.
<i>Hypocrea virens</i>	n.a.	AY158203	n.a.	n.a.	n.a.	n.a.
<i>Hypocrea jecorina</i>	n.a.	n.a.	ATCC 13631	AF510497	n.a.	n.a.
<i>Hypocrea lutea</i> n.a.	n.a.	IFO9061	D14407	n.a.	n.a.	n.a.
<i>Hypocrea koningii</i>	n.a.	n.a.	ATCC 64262	HKO301990	n.a.	n.a.
<i>Hypocrea rufa</i> n.a.	n.a.	GJS89-127	AY489694	n.a.	n.a.	n.a.
<i>Trichoderma viride</i>	T9BR47	Z15055	IFFI 13001	AF525230	n.a.	n.a.
<i>Trichoderma asperellum</i>	n.a.	AY390326.1	n.a.	n.a.	n.a.	n.a.
<i>Trichoderma harzianum</i>	n.a.	n.a.	ALI 232	AF548100	n.a.	n.a.
<i>Fusarium cerealis</i>	n.a.	n.a.	n.a.	AF141947	n.a.	n.a.
<i>Fusarium equiseti</i>	n.a.	n.a.	n.a.	AF141949	n.a.	n.a.
<i>Fusarium oxysporum</i>	n.a.	n.a.	26-ene	AB110910	n.a.	n.a.
<i>Gibberella avenacea</i>	n.a.	n.a.	n.a.	AF141946	n.a.	n.a.
<i>Gibberella fujikuroi</i>	A 102	U27303	NBRC 30337	AB237662	n.a.	n.a.
<i>Gibberella pulicaris</i>	n.a.	AF484166.1	n.a.	AF149875	n.a.	n.a.
<i>Gibberella zeae</i> NRRL-31084	AACM01000393	NBRC 9462	AB250414	n.a.	n.a.	n.a.
<i>Aspergillus nidulans</i> = <i>Emericella nidulans</i> .	FGSC A4	XM653694	n.a.	n.a.	n.a.	n.a.
<i>Aspergillus flavus</i>	n.a.	P22012	n.a.	n.a.	n.a.	n.a.
<i>Aspergillus parasiticus</i>	NRRL-5067	L49386	n.a.	n.a.	n.a.	n.a.
<i>Aspergillus fumigatus</i>	AF293	AAHF01000004	n.a.	n.a.	n.a.	n.a.
<i>Penicillium digitatum</i>	PD 5	D78154	n.a.	n.a.	n.a.	n.a.
<i>Penicillium paxilli</i>	n.a.	AY846880	n.a.	n.a.	n.a.	n.a.
<i>Leptospaeria biglobosa</i>	AY7490000	FSU415	n.a.	n.a.	n.a.	n.a.
<i>Leptospaeria maculans</i>	AY7490028	FSU2560	n.a.	n.a.	n.a.	n.a.
<i>Phaeosphaeria avenaria</i>	S-81-W10	AY823527	n.a.	n.a.	n.a.	n.a.
<i>Phaeosphaeria nodorum</i>	98-12981	AY823526	n.a.	n.a.	n.a.	n.a.
<i>Pleospora herbarum</i>	ICMP 562077	Y17077	n.a.	n.a.	n.a.	n.a.
<i>Pleospora welwitschiae</i>	FSU2566	AY749034	n.a.	n.a.	n.a.	n.a.
<i>Botryotinia fuckeliana</i>	SAS56	Z69263	n.a.	n.a.	n.a.	n.a.
<i>Sclerotinia sclerotiorum</i>	TZ25	AY312374	n.a.	n.a.	n.a.	n.a.
<i>Monilinia fructicola</i>	ATF16	AY283677	n.a.	n.a.	n.a.	n.a.
<i>Monilinia laxa</i> 515	AY349149	n.a.	n.a.	n.a.	n.a.	n.a.
<i>Blumeria graminis</i>	n.a.	P16040	n.a.	n.a.	n.a.	n.a.
<i>Erysiphe necator</i>	n.a.	AY074934	n.a.	n.a.	n.a.	n.a.
<i>Erysiphe pisi</i> n.a.	S49328	n.a.	n.a.	n.a.	n.a.	n.a.
<i>Neurospora crassa</i>	OR74A	XM323372	n.a.	n.a.	n.a.	n.a.
<i>Neurospora pannonica</i>	TRTC51327	AY780126.1	n.a.	n.a.	n.a.	n.a.
<i>Sordaria fimicola</i>	SMH4106	AY780138	n.a.	n.a.	n.a.	n.a.
<i>Sordaria macrospora</i>	n.a.	AY780140	n.a.	n.a.	n.a.	n.a.
<i>Saccharomyces cerevisiae</i>	n.a.	n.a.	n.a.	J01353	n.a.	n.a.
<i>Melampsora lini</i>	5261	AF317682	n.a.	n.a.	n.a.	n.a.
<i>Cryptococcus neoformans</i>	JEC21	AE017343	n.a.	n.a.	n.a.	n.a.
<i>Ustilago hordei</i> n.a.	n.a.	n.a.	U00973	n.a.	n.a.	n.a.
<i>Ustilago shiraiana</i>	n.a.	n.a.	IFO8809	AB028189	n.a.	n.a.
<i>Micromucon ramannianus</i>	NRRL-5844	AF162073	n.a.	n.a.	n.a.	n.a.
<i>Rhizopus microsporus</i>	NRRL-2710	AF162066	n.a.	n.a.	n.a.	n.a.

^aIsolates that are sequenced for β-tubulin, large and small subunit of rRNA genes of mitochondria.

^bEpizootic population.

^cARSEF isolates are from USDA-ARS collection of Entomopathogenic Fungal cultures, Ithaca, New York, USA. ITCC isolates are from Indian Type Culture Collection, IARI, New Delhi, India. NRRL isolates are from NRRL culture collection, Peoria, Illinois, USA. BB isolates and II.a.4.; II.a.6.; of *B. bassiana* and I.a.3.; II.a.1.1.; II.b.5.; II.b.7. of *N. rileyi* are isolated in our laboratory and not yet accessioned.




Figure 1. Consensus phylogenetic tree derived by neighbor joining method of a sample of 44 fungal entries to assess the phylogenetic affiliation of the asexual entomopathogenic fungi *Beauveria bassiana* and *Nomuraea rileyi* to each other and other asexual fungi with entomopathogenic habit and sexual (meiosporic) fungi. Taxa underlined are entomopathogenic fungi of interest. Bootstrap values (based on 1,000 replicates) when above 50% are indicated on the branches. The tree was derived from partial sequence (271 amino acids) of the β -tubulin gene of which 150 were conserved, 121 were variable, of which 83 were parsimony informative.

methods (MP, ME and NJ) indicates that these clusters are not incidental. The tree derived through NJ is represented in Figure 1 and is further described. The ME and MP trees are given as supplementary material (Figs. 2 and 3). Branching order reflected the expected pattern with fungi of the three divisions Asco, Basidio and Zygomycota separating into three major clades (Fig. 1). The whole clade of Ascomycetes was well supported with a bootstrap value of 91%. The Ascomycota clade branched into two sub-clades with the separation of members of Hypocreales in one group and the rest in the other (Fig. 1). The sub-clade with non-hypocrealean fungi fissured into five branches each representing members of one taxonomic order – the Sordariales, Heliales, Erysiphales, Pleosporales and Euriotales. There were three sub-clades in the hypocrealean clade each representing one family: Nectriaceae,

Hypocreaceae and Clavicipitaceae (Fig. 1). All the mitosporic entomopathogenic fungi *B. bassiana*, *Beauveria brongniartii*, *N. rileyi*, *Paecilomyces fumosoroseus*, *Paecilomyces farinosus* and *Metarhizium anisopliae* grouped into the clavicipitalean sub-clade of Hypocreales (Fig. 1). Within this sub-clade, each of the entomopathogenic fungal genus separated into a different clade (Fig. 1). *M. anisopliae* was found to be closely related to sexual clavicipitalean fungi (Fig. 1). There was one discrepancy in the distribution of the members of Hypocreaceae. Sexual Ascomycete, *Acremonium chrysogenum* of Hypocreaceae, separated from the other members of that family (Fig. 1). The Bayesian tree of highest likelihood (Fig. 4, supplementary material) has a significant (>95%) posterior probability and strong bootstrap support for all (ME, MP and NJ) trees. *p* value in the KH test ($p = 0.05$) indicates that the maximum likelihood tree (5154.26590)




Figure 2. Consensus phylogenetic tree derived for the same data set as in Figure 1 derived by the minimum evolution method.

is not significantly worse than the original maximum likelihood tree (5155.47481).

The results from the analysis of rRNA gene sequence data (with some differences in the fungal entries included) very closely matched the phylogenetic relationships evident in the analysis of the sequence data of β -tubulin gene. The NJ tree from the rRNA gene is given in Figure 5 (the ME, MP and Bayesian trees are provided as supplementary Figs. 6, 7 and 8, respectively).

With the analysis of both β -tubulin and rRNA gene sequences, enough evidence was not found to reject the monophyly of the entomopathogenic fungal group studied. We therefore conclude that the mitosporic entomopathogenic fungi are members of *Clavicipitaceae* family.

Interrelationships between entomopathogenic fungal species

The five major genera included in the analysis separated into different clades in the trees derived by all the three (NJ, ME and MP) methods; only the position of *M. anisopliae* differed in the three phylogenetic trees. The NJ tree is described and is given in Figure 9; the ME and MP trees are provided as supplementary material in Figures 10 and 11. Two isolates of *B. bassiana* separated from the rest of the *B. bassiana* isolates as another clade (Fig. 9). The two other species of *Beauveria* in the sample – *Beauveria amorpha* and *B. brongniartii* – grouped with isolates of *B. bassiana* (Fig. 9). Deep phylogenetic lineages were observed in the *B. bassiana* clade (Fig. 9). Unlike *B. bassiana*, extensive




Figure 3. Consensus phylogenetic tree derived for the same data set as in Figure 1 derived by maximum parsimony analysis. Component tree statistics: tree length (L) = 269; consistency index (CI) = 0.737; retention index (RI) = 0.818.

phylogenetic speciation was not observed in the *N. rileyi* sample (Fig. 9). Only one isolate diverged into a different lineage among the members (Fig. 9). *M. anisopliae* separated as a different lineage in the main group (Fig. 9), while in the MP tree it grouped with the two *B. bassiana* isolates, which diverged from the main group; in the ME tree it separated out as a different lineage from the main group.

Cryptic speciation in *Beauveria bassiana*

Diagnosing species borders from single-gene phylogenies can be difficult because usually the markers do not indicate a clear cutoff between species and higher taxonomic groupings. Therefore an approach termed genealogical concordance phylogenetic species

recognition (GCPSR) was used²⁴. In fungal taxa, where mating studies are difficult to conduct and when patterns are not detectable from multilocus sequence data, multiple gene genealogies (3-10 genes) were used to uncover cryptic speciation using this powerful approach of GCPSR²⁴. GCPSR detects genetically isolated groups from a number of different loci by comparing the gene trees. Different genes have different genealogies within a species due to recombination. It establishes gene flow delimiting species by identifying the unshared polymorphisms, and thus branches that are incompatible, with all genealogies at all loci.

A consensus maximum parsimony tree was generated (PAUP ver. 4.0)³¹ from the partial sequences of the three genes: β -tubulin gene, large and small subunit of rRNA genes of mitochondria aligned by AlnExplorer (MEGA ver 3.1) derived from the isolates of an epizootic




Figure 4. Bayesian Metropolis-coupled Markov chain Monte Carlo (MCMCMC) fungal tree depicting phylogenetic relationships for the same data set as in Figure 1. This phylogeny resulted from a 50% majority rule consensus of 10001 trees sampled with Bayesian MCMCMC. The resulting posterior probabilities (PP) are shown above internal branches. Taxa underlined are entomopathogenic fungi of interest. Species names are colored according to their respective phyla.

population of *B. bassiana* collected from Burgenland, Austria (Catalogue of the entomopathogenic fungi at ARSEF, Ithaca, New York). The details of the maximum parsimony tree generated from the aligned DNA sequences of each gene are as follows:

β -tubulin gene: The tree was constructed from 296 characters, of which 285 were constant and 11 were variable, and of these 11 variable characters only one was parsimony informative (Fig. 12). The tree length (L) was 11, consistency index (CI) was 1.00, homoplasy index (HI) was 0.00 and retention index (RI) was 1.00.

Gene for large subunit of rRNA of mitochondria: The tree was constructed from 183 characters, of which 105 were constant and 78 were variable and of these 37 were parsimony informative (Fig. 12). The tree length (L) was 109, consistency index (CI) was 0.89, homoplasy index (HI) was 0.10 and retention index (RI) was 0.80.

Gene for small subunit of rRNA of mitochondria: The tree was constructed from 504 characters of which 498 were constant and six were variable and of these six variable characters one was parsimony informative (Fig. 9). The tree length (L) was 6, consistency index (CI) was 1.00, homoplasy index (HI) was 0.00 and retention index (RI) was 1.00.

The tree topology of each species tree indicates the presence of cryptic speciation. Incongruity of gene genealogies within a given

group indicates gene flow and delimits a species. As the approach detects reproductive isolation, the resulting groups also fulfill the criteria of a biological species. Genealogical concordance phylogenetic species recognition (GCPSR) detects genetically isolated groups by comparing the gene trees from a number of different loci. Different genes have different genealogies within a species due to recombination. It establishes gene flow delimiting species by identifying the unshared polymorphisms, and thus branches that are incompatible with all genealogies at all loci, represent different species.

Discussion

The β -tubulin gene- and rRNA gene-based phylogenetic analysis affirmed the traditional taxonomic grouping of fungi, which was earlier substantiated by molecular phylogenetic studies¹⁷. The position of the entomopathogenic fungi *B. bassiana*, *B. brongniartii*, *N. rileyi*, *M. anisopliae* and *Paecilomyces* species in the family Clavicipitaceae of Hypocreales in Ascomycota announced in different phylogenetic analysis was found true.

The entomopathogenic fungi in the present sample have been shown earlier through molecular phylogenetic studies of samples with different fungal isolates to be closely related to another entomopathogenic mitosporic fungus – *Lecanicillium lecanii* and other species of *Paecilomyces*, *Beauveria*, *Cordyceps* and




Figure 5. Consensus phylogenetic tree of a sample of 64 fungal entries to assess the phylogenetic affiliation of the asexual entomopathogenic fungi *Beauveria bassiana* and *Nomuraea rileyi* to each other and other asexual fungi with entomopathogenic habit and sexual (meiosporic) fungi derived by neighbor joining method. The tree was derived from 64 taxa and partial sequence (1359 nucleotides) of the rRNA gene of which 933 were conserved, 408 were variable, of which 293 were parsimony informative.




Figure 6. Consensus phylogenetic tree derived for the same data set as in Figure 5 derived by the minimum evolution method.




Figure 7. Consensus phylogenetic tree derived for the same data set as in Figure 1 derived by maximum parsimony analysis. Component tree statistics: tree length (L) = 574; consistency index (CI) = 0.0721; retention index (RI) = 0.818.




Figure 8. Bayesian Metropolis-coupled Markov chain Monte Carlo (MCMCMC) fungal tree depicting phylogenetic relationships for the same data set as in Figure 5. This phylogeny resulted from a 50% majority rule consensus of 10001 trees sampled with Bayesian MCMCMC. The resulting posterior probabilities (PP) are shown above internal branches. Taxa underlined are entomopathogenic fungi of interest. Species names are colored according to their respective phyla.

Nomuraea^{2,8,21,22,27,29,35}. Thus, all the five important conidiogenous fungal genera with entomopathogenic habit – *Beauveria*, *Nomuraea*, *Metarhizium*, *Paecilomyces* and *Lecanicillium* – are closely related. Among these fungi, *M. anisopliae* diverged further from the rest of the entomopathogenic species and showed closer affiliation to sexual Clavicipitaceae both in the present study and in a previous analysis based on mitochondrial genes by Uribe and Khachatourians³⁶ and rRNA gene of the small subunit²².

In the present analysis, the boundaries between species in the entomopathogenic fungal genera were nebulous. *B. brongniartii* and *B. amorpha* were found dispersed among the *B. bassiana* isolates though in a separate clade. A similar observation was made in the

allozyme analysis of a very large sample of *Beauveria* species by St. Leger et al³⁰ – some of the *B. brongniartii* isolates clustered separately while some grouped along with *B. bassiana* isolates. Phylogenetic analysis with the β -tubulin gene sequence not only iterated the complex nature of *B. bassiana* species²⁷ but also revealed the great genetic distance between the members in the species, the distance being more than between morphologically distinguishable species. Boucias et al³ observed some *N. rileyi* isolates clustering with *M. anisopliae*. Such species overlapping was also observed in molecular phylogenetic studies with different members of this group^{21,22,31}. These observations led to the conclusion that the asexual insect pathogenic fungi must have all diverged from a common clavicipitacean




Figure 9. Consensus phylogenetic tree generated from partial β -tubulin gene sequence through neighbor joining method of a sample of isolates of *Beauveria bassiana*, *Beauveria amorpha*, *Beauveria brongniartii*, *Nomuraea rileyi*, *Paecilomyces fumosoroseus*, *Metarhizium anisopliae*, *Cordyceps militaris* to study the interrelationship by. The aligned data consisted of 257 characters of which 125 were conserved and the rest were variable. Of these, 82 were parsimony informative characters.




Figure 10. Consensus phylogenetic tree derived for the same data as in Figure 9 derived by the minimum evolution method. Bootstrap values when above 50% are indicated on the branches.

ancestor. Loss of sex in these clavicipitalean entomopathogenic fungi is believed to be a derived character² having been attained simultaneously or subsequent to acquisition of entomopathogenic habit. Pathogenicity systems are believed to be flexible enough to continue to provide virulence against a broad range of hosts despite the absence of sexual recombination². In fact, episodic events in host

population dynamics may favour clonal populations because of the rapidity and low energy cost associated with asexuality; the population may indeed benefit from reduced out crossing through the maintenance of well-adapted pathotypes². Muller's Ratchet effect – a population genetic mechanism that describes how asexual populations may undergo an unavoidable and progressive decline in




Figure 11. Consensus phylogenetic tree derived for the same data as in Figure 9 derived by the maximum parsimony method. Bootstrap values when above 50% are indicated on the branches. The description of the tree is as follows: consistency index (CI) = 0.905; retention index (RI) = 0.875 for the 326 parsimonious trees.

fitness (due to absence of recombination mediated through sexual reproduction) eventually leading to extinction – was found to not apply to asexual entomopathogenic fungi². Thus, the imperfect entomopathogenic fungal genera represent different mitotic forms of a perfect fungus of Clavicipitaceae.

Acknowledgements

NNRR and CUM are thankful to CSIR, New Delhi, for a research fellowship. We thank the DAAD-DST PPP-03 for facilitating this research collaboration.




Figure 12. One consensus maximum parsimony tree generated from the sequences of (a) partial sequence of β -tubulin gene, (b) large subunit of mt rRNA gene and (c) small subunit of mt rRNA genes derived from the isolates of an epizootic *B. bassiana* population from Burgenland, Austria. Numbers below branches represent bootstrap values (%) based on 500 replicates. Only values >50% are shown. The tree topology of each species tree indicates the presence of cryptic speciation.

References

- Berbee ML, Taylor JW. Convergence in ascospore discharge mechanism among pyrenomycete fungi based on 18S ribosomal RNA gene sequence. *Mol Phylogenet Evol.* 1992;1:59-71.
- Bidochka MJ, de Koning J. Are teleomorphs really necessary?: modelling the potential effects of Muller's Ratchet on deuteromycetous entomopathogenic fungi. *Mycol Res.* 2001;105:1014-1019.
- Boucias DG, Tigano MS, Sosa-Gomez DR, Glare TR, Inglis PW. Genotypic properties of the entomopathogenic fungus *Nomuraea rileyi*. *Biol Control.* 2000;19:124-138.
- Coates BS, Hellmich RL, Lewis LC. Nuclear small subunit rRNA group 1 intron variation among *Beauveria* species provide tools for strain identification and evidence of horizontal transfer. *Curr Genet.* 2002;41:414-424.
- Einax E, Voigt K. Oligonucleotide primers for the universal amplification of β -tubulin genes facilitate phylogenetic analysis in the regnum Fungi. *Organ Divers Evol.* 2003;3:185-194.
- Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution.* 1985;39:783-791.
- Glass NL, Donaldson GC. Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Appl Environ Microbiol.* 1995;61:323-330.
- Han Q, Inglis GD, Hausner G. Phylogenetic relationships among strains of the entomopathogenic fungus, *Nomuraea rileyi*, as revealed by partial β -tubulin sequences and inter-simple sequence repeat (ISSR) analysis. *Lett Appl Microbiol.* 2002;34:376-383.
- Hegedus DD, Pfeifer TA, Muly K, Khachatourians GG. Characterization and structure of the mitochondrial small rRNA gene of the entomopathogenic fungus *Beauveria bassiana*. *Genome.* 1998;41:471-476.
- Huelsenbeck JP, Ronquist F. MrBayes: bayesian inference of phylogenetic trees. *Bioinformatics.* 2001;17:754-755.
- James TY, Kauff F, Schoch CL, Matheny PB, Hofstetter V, Cox CJ, Celio G, Gueidan C, Fraker E, Miadlikowska J, Lumbsch HT, Rauhut A, Reeb V, Arnold AE, Amtoft A, Stajich JE, Hosaka K, Sung G-H, Johnson D, O'Rourke Crockett BM, Binder M, Curtis JM, Slot JC, Wang Z, Wilson AW, Schüssler A, Longcore JE, O'Donnell K, Mozley-Standridge S, Porter D, Letcher PM, Powell MJ, Taylor JW, White MM, Griffith GW, Davies DR, Humber RA, Morton JB, Sugiyama J, Rossman AY, Rogers JD, Pfister DH, Hewitt D, Hansen K, Hambleton S, Shoemaker RA, Kohlmeyer J, Volkman-Kohlmeyer B, Spotts RA, Serdani M, Crous PW, Hughes KW, Matsuura K, Langer E, Langer G, Untereiner WA, Lücking R, Bu'del B, Geiser DM, Aptroot A, Diederich P, Schmitt I, Schultz M, Yahr R, Hibbett DS, Lutzoni F, McLaughlin DJ, Spatafora JW, Vilgalys R. Reconstructing the early evolution of fungi using a six-gene phylogeny. *Nature.* 2006;443:818-822.
- Keeling PJ. Congruent evidence from alpha-tubulin and beta-tubulin gene phylogenies for a zygomycete origin of microsporidia. *Fungal Genet Biol.* 2003;38:298-309.
- Kishino H, Hasegawa M. Evaluation of the maximum likelihood estimate of the evolutionary tree topologies from DNA sequences, and the branching order of Hominoidea. *J Mol Evol.* 1989;29:170-179.
- Kumar S, Tamura K, Nei M. MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform.* 2004;5:150-163.
- Lake JA. The order of sequence alignment can bias the selection of tree topology. *Mol Biol Evol.* 1991;8:378-385.
- Landvik S, Eriksson OE, Berbee ML. *Neolecta* – a fungal dinosaur? Evidence from beta-tubulin amino acid sequences. *Mycologia.* 2001;93:1151-1163.
- Lutzoni F, Kauff F, Cox CJ, McLaughlin D, Celio G, Dentinger B, Padamsee M, Hibbett D, James TY, Baloch E, Grube M, Reeb V, Hofstetter V, Schoch C, Arnold AE,

- Miadlikowska J, Spatafora J, Johnson D, Hambleton S, Crockett M, Shoemaker R, Sung G-H, Lücking R, Lumbsch T, O'Donnell K, Binder M, Diederich P, Ertz D, Gueidan C, Hall B, Hansen K, Harris RC, Hosaka K, Lim Y-W, Matheny B, Nishida N, Pfister D, Rogers J, Rossman A, Schmitt I, Sipman H, Stone J, Sugiyama J, Yahr R, Vilgalys R. Assembling the fungal tree of life: classification and the evolution of their subcellular traits? *Am J Bot.* 2004;91:1446-1480.
18. Miller AN, Huhndorf SM. Multi-gene phylogenies indicate ascomal wall morphology is a better predictor of phylogenetic relationships than ascospore morphology in the Sordariales (Ascomycota, Fungi). *Mol Phylogenet Evol.* 2005;5:60-75.
19. Moore D, Frazer N, Ann L. Systematics, phylogeny and evolution. In: Moore D, Frazer N, Ann L, editors. *Essential fungal genetics*. Heidelberg: Springer; 2002. p. 244-281.
20. Myllys L, Stenroos S, Thell A. New genes for phylogenetic studies of lichenized fungi: glyceraldehyde-3-phosphate dehydrogenase and beta-tubulin genes. *Lichenologist.* 2002;34:237-246.
21. Nikoh N, Fukatsu T. Interkingdom host jumping underground: phylogenetic analysis of entomoparasitic fungi of the genus *Cordyceps*. *Mol Biol Evol.* 2000;17:629-638.
22. Obornik M, Jirku M, Dolezel D. Phylogeny of mitosporic entomopathogenic fungi: is the genus *Paecilomyces* polyphyletic? *Can J Microbiol.* 2001;47:813-819.
23. Olsen GJ, Woese CR. Ribosomal RNA: a key to phylogeny. *FASEB J.* 1993;7:113-123.
24. Padmavathi J, Uma Devi K, Uma Maheswara Rao C, Nageswara Rao Reddy N. Telomere fingerprinting for assessing chromosome number, isolate typing and recombination in the entomopathogen, *Beauveria bassiana* (Balsamo) Vuillemin. *Mycol Res.* 2003;107:572-580.
25. Pfeifer TA, Hegedus DD, Khachatourians GG. The mitochondrial genome of the entomopathogenic fungus *Beauveria bassiana*: analysis of the ribosomal RNA region. *Can J Microbiol.* 1993;39:25-31.
26. Redecker D. New views on fungal evolution based on DNA markers and the fossil record. *Res Microbiol.* 2002;153:125-130.
27. Rehner SA, Buckley E. A *Beauveria* phylogeny inferred from nuclear ITS and EF1- ζ sequences: evidence for cryptic diversification and links to *Cordyceps* teleomorphs. *Mycologia.* 2005;97:84-98.
28. Rehner SA. Phylogenetics of the insect pathogenic genus *Beauveria*. In: Vega EF, Blackwell M, editors. *Insect-fungal associations, ecology and evolution*. New York: Oxford University Press; 2005. p. 3-27.
29. Shah PA, Goettel MS. *Directory of Microbial Control Products and Services*. Gainesville, Florida: Society for Invertebrate Pathology; 1999.
30. St. Leger RJ, Allee LL, May B, Staples RC, Roberts DW. World-wide distribution of genetic variation among isolates of *Beauveria* spp. *Mycol Res.* 1992;96:1007-1015.
31. Sung GH, Spatafora JW, Zare R, Hodge KT, Gams W. A revision of *Verticillium* sect. *Prostrata*. II. Phylogenetic analysis of SSU and LSU nuclear rDNA sequences from anamorphs and teleomorphs of the Clavicipitaceae. *Nova Hedwigia.* 2001;71:311-328.
32. Swofford DL, Olsen GJ, Waddell PJ, Hillis DM. Phylogenetic inference. In: Hillis DM, Moritz C, Mable BK, editors. *Molecular systematics*. Sunderland, USA: Sinauer Associates; 1996. p. 407-514.
33. Takamatsu S. PCR applications in fungal phylogeny. In: Bridge PD, Arora DH, Reddy CA, Elander RP, editors. *Applications of PCR in mycology*. Wallingford: CAB International; 1998. p. 125-152.
34. Taylor JW, Jacobson DJ, Kroken S, Kasuga T, Geiser DM, Hibbett DS, Fischer MC. Phylogenetic species recognition and species concepts in fungi. *Fungal Genet Biol.* 2000;31:21-32.
35. Uma Devi K, Annette R, Nageswara Rao Reddy N, Uma Maheswara Rao C, Padmavathi J. Genetic diversity, reproductive biology and speciation in the entomopathogenic fungus *Beauveria bassiana* (Balsamo) Vuillemin. *Genome.* 2006;49:1-10.
36. Uribe D, Khachatourians GG. Restriction fragment length polymorphism of mitochondrial genome of the entomopathogenic fungus *Beauveria bassiana* reveals high intraspecific variation. *Mycol Res.* 2004;108:1070-1078.
37. White TJ, Bruns T, Lee S, Taylor J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ, editors. *PCR protocols: a guide to methods and applications*. New York: Academic Press; 1990. p. 315-322.