



Nomenclature of Genera and Subdivisions of Genera within *Scleria* P. J. Bergius (Cyperaceae)

Author(s): Konraad Camelbeke, Gea Zijlstra and Paul Goetghebeur

Source: *Taxon*, Vol. 50, No. 2, Golden Jubilee Part 4 (May, 2001), pp. 479-486

Published by: [International Association for Plant Taxonomy \(IAPT\)](#)

Stable URL: <http://www.jstor.org/stable/1223894>

Accessed: 12/02/2015 11:18

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



International Association for Plant Taxonomy (IAPT) is collaborating with JSTOR to digitize, preserve and extend access to *Taxon*.

<http://www.jstor.org>

NOMENCLATURE

Edited by John McNeill¹

Nomenclature of genera and subdivisions of genera within *Scleria* P.J. Bergius (*Cyperaceae*)

Konraad Camelbeke², Gea Zijlstra³ & Paul Goetghebeur²

Summary

Camelbeke, K., Zijlstra, G. & Goetghebeur, P.: Nomenclature of genera and subdivisions of genera within *Scleria* P.J. Bergius (*Cyperaceae*). – Taxon 50: 479–486. 2001. – ISSN 0040-0262.

A list of effectively published names of genera and subdivisions of genera included or formerly included in the genus *Scleria* is provided, together with their status and types. The following names are here typified: *Hymenolytrum* Schrad. ex Nees, *Macrolomia* Schrad. ex Nees, *Mastigoscleria* Nees, *Omoscleria* Nees, and *Trachylomia* Nees; *Scleria* sect. *Foveolidia* Raf., sect. *Margaleia* Raf., sect. *Paniculatae* Pax, and sect. *Rugunia* Raf.; *Scleria* ser. *Polystachyae* Benth. and ser. *Subspicatae* Benth.; *Scleria* [unranked] *Spicatae* Boeck., and [unranked] *Tuberculatae* Core. Moreover, two excluded names are typified: *Cryptangium* Schrad. ex Nees, and *Scleria* ser. *Axillares* Benth.

Keywords: *Cyperaceae*, generic names, infrageneric names, nomenclature, *Scleria*, typification.

Introduction

While preparing a revision of the neotropical representatives of the genus *Scleria* (Camelbeke & Goetghebeur, in prep.) the need was felt for an exhaustive list of names of genera and subdivisions of genera included or formerly included in the genus *Scleria*. Our study of all relevant and available literature revealed that several of those names were not typified or that their nomenclatural status was unclear. Therefore, if infrageneric taxa are to be recognized in this large genus of *Cyperaceae* (+200 species), the nomenclatural problems of these names have to be resolved and the correct types have to be cited or designated. We have tried to achieve these goals and the results are presented here.

¹Royal Botanic Garden, 20A Inverleith Row, Edinburgh, EH3 5LR, Scotland, U.K. E-mail: jmcneill@rbge.org.uk.

²Ghent University, Department of Biology, Laboratory of Botany, K. L. Ledeganckstraat 35, B-9000 Gent, Belgium. E-mail: konraad.camelbeke@rug.ac.be (author for correspondence); paul.goetghebeur@rug.ac.be.

³Nationaal Herbarium Nederland, Utrecht University branch, Heidelberglaan 2, 3584 CS Utrecht, The Netherlands. E-mail: zijlstra@accu.uu.nl.

Generic names

Acriulus Ridl., J. Linn. Soc., Bot. 20: 336. 1883. – Type: *Acriulus madagascariensis* Ridl. (≡ *Scleria acriulus* C.B. Clarke), lectotype designated by Kern (1964: 44).

[*Anerma* Schrad. ex Nees in Mart., Fl. Bras. 2(1): 170. 1842. *Nom. inval.*: ICBN Art. 34.1.c.]

Catagyna P. Beauv. ex T. Lestib., Essai Cyp. 26. 1819. – Type: *Scleria anomala* (Steud.) J. Raynal, lectotype designated by Raynal (1977: 40).

Chondrolomia Nees in Mart., Fl. Bras. 2(1): 173. 1842. – Type: *Chondrolomia sellowiana* (Kunth) Nees (≡ *Scleria sellowiana* Kunth).

[*Cryptopodium* Schrad. ex Nees in Mart., Fl. Bras. 2(1): 180. 1842. *Nom. inval.*: ICBN Art. 34.1.c.]

Cylindropus Nees, Linnaea 9: 303. 1834. – Type: *Cylindropus junciformis* Nees (≡ *Scleria pilosa* Boeck., new name for *Scleria junciformis* (Nees) Thw. (1864), non Kunth (1837)).

Diaphora Lour., Fl. Cochinch. 578. 1790. – Type: *Diaphora cochinchinensis* Lour. (≡ *Scleria cochinchinensis* (Lour.) Druce).

Diploscyphus Liebm., Mexic. Halvgr. 74. 1850. – Type: *Diploscyphus mexicanus* Liebm. (≡ *Scleria mexicana* (Liebm.) Boeck.).

Hymenolytrum Schrad. ex Nees in Mart., Fl. Bras. 2(1): 174. 1842. – Type: *Hymenolytrum cyperinum* (Willd. ex Kunth) Nees (≡ *Scleria cyperina* Willd. ex Kunth), lectotype here designated.

Hypoporum Nees, Linnaea 9: 303. 1834. – Type: *Hypoporum pergracile* Nees (≡ *Scleria pergracilis* (Nees) Kunth), lectotype designated by Kern (1961: 196).

Macrolomia Schrad. ex Nees in Mart., Fl. Bras. 2(1): 181. 1842. – Type: *Macrolomia bracteata* (Cav.) Schrad. ex Nees (≡ *Scleria bracteata* Cav.), lectotype here designated.

Mastigoscleria Nees in Mart., Fl. Bras. 2(1): 177. 1842. – Type: *Mastigoscleria reflexa* (Kunth) Nees (≡ *Scleria reflexa* Kunth), lectotype here designated.

Omoscleria Nees in Mart., Fl. Bras. 2(1): 180. 1842. – Type: *Omoscleria flagellum* Nees, *nom. illeg.*: ICBN Art. 52.1.; (≡ *Scleria flagellum* Sw., *nom. illeg.*: ICBN Art. 52.1.; ≡ *Scleria flagellum-nigrorum* P.J. Bergius), lectotype here designated. ≡ *Scleria* Bergius 1765 (by lectotypification).

[*Ophrydium* Schrad. ex Nees in Mart., Fl. Bras. 2(1): 183. 1842. *Nom. inval.*: ICBN Art. 34.1.c.]

Ophryoscleria Nees in Mart., Fl. Bras. 2(1): 182. 1842. – Type: *Ophryoscleria racemosa* (Poir.) Nees (≡ *Scleria racemosa* Poir.), lectotype designated by Franklin Hennessy (1985: 514).

Schizolepis Schrad. ex Nees in Mart., Fl. Bras. 2(1): 186. 1842. – Type: *Schizolepis latifolia* (Sw.) Nees (≡ *Scleria latifolia* Sw.), lectotype designated by Franklin Hennessy (1985: 514).

Scleria P. J. Bergius, Kongl. Vetensk. Acad. Handl. 26: 142. 1765. – Type: *Scleria flagellum-nigrorum* P. J. Bergius, nom. cons. prop. (see Camelbeke & Goetghebeur 2000: 295), lectotype designated by Britton (1907: 17).

Trachylomia Nees in Mart., Fl. Bras. 2(1): 174. 1842. – Type: *Trachylomia triglomerata* (Michx.) Nees (≡ *Scleria triglomerata* Michx.), lectotype here designated.

Excluded generic names

[*Adamantogeton* Schrad. ex Nees in Mart., Fl. Bras. 2(1): 165. 1842. *Nom. inval.*: ICBN Art. 34.1.c.]

Acrocarpus Nees in Mart., Fl. Bras. 2(1): 157. 1842, *nom. illeg.*: ICBN Art. 53.1.; (non R. Wight ex Arnott 1838). – Type: *Acrocarpus polyphyllus* Nees (≡ *Lagenocarpus polyphyllus* (Nees) O. Kuntze), lectotype designated by Koyama & Maguire (1965: 51).

Becquerelia Brongn. in Duperrey, Voy. Monde 2: 161. 1833. – Type: *Becquerelia cymosa* Brongn., lectotype designated by Koyama (1967: 25).

[*Cladotheca* Steud., Syn. Pl. Glumac. 2: 178. 1855. *Nom. inval.*: ICBN Art. 34.1.c.].

Cryptangium Schrad. ex Nees in Mart., Fl. Bras. 2(1): 163. 1842. – Type: *Cryptangium uliginosum* Schrad. ex Nees, *nom. illeg.*: ICBN Art. 52.1.; (≡ *Fuirena verticillata* Sprengel. ≡ *Lagenocarpus verticillatus* (Sprengel) T. Koyama & Maguire), lectotype here designated.

[*Cryptanguina* Schrad. ex Lindl., Veg. Kingd. 119. 1846. Orth. var., see *Cryptangium*.]

Diplacrum R. Br., Prodr. 240. 1810. – Type: *Diplacrum caricinum* R. Br.

Lagenocarpus Nees, Linnaea 9: 304. 1834. – Type: *Lagenocarpus guianensis* Nees.

Pteroscleria Nees in Mart., Fl. Bras. 2(1): 196. 1842. – Type: *Pteroscleria guianensis* Nees (≡ *Diplacrum guianensis* (Nees) T. Koyama).

Sphaeropus Boeck., Flora 56: 89. 1873. – Type: *Sphaeropus pygmaeus* Boeck. (= *Diplacrum pygmaeum* (R. Br.) Nees ex Boeck.).

Subgeneric names

Scleria subg. *Acriulus* (Ridl.) C.B. Clarke in Dyer, Fl. Trop. Afr. 8: 495. 1902. – Type: see *Acriulus* Ridl. Note: the names published by Clarke in 1902 and which are preceded by an asterisk should be considered subgenera (see p. 494).

Scleria subg. *Browniae* (C.B. Clarke) C.B. Clarke, Bull. Misc. Inform. Kew, Addit. Ser. 8: 132. 1908 (as “*Browneae*”). – Type: see *Scleria* sect. *Browniae* C.B. Clarke.

[*Scleria* subg. *Euscleria* C.B. Clarke, Fl. Cap. 7: 294. 1898. *Nom. inval.*: ICBN Art. 21.3.]

Scleria subg. *Hypoporum* (Nees) C.B. Clarke, Fl. Brit. India 6: 685. 1894. – Type: see *Hypoporum* Nees.

[*Scleria* subg. *Lithocarpella* Rchb., Consp. Regn. Veg. 56. 1828. *Nom. nud.*]

[*Scleria* subg. *Macrolomia* (Nees) H. Pfeiff., Revista Sudamer. Bot. 10: 219. 1956. *Nom. inval.*: ICBN Art. 33.3.]

Scleria subg. *Ophryoscleria* (Nees) C.B. Clarke in Dyer, Fl. Trop. Afr. 8: 495. 1902. – Type: see *Ophryoscleria* Nees.

Scleria subg. *Primoscleriae* H. Pfeiff., Revista Sudamer. Bot. 10: 219. 1956 (as “*Primoscleria*”). – Type: *Scleria hirtella* Sw. [ICBN Art. 7.7.]. Note: For indication of the rank, see the comment on “Untergattung *Macrolomia*” in *Scleria* [unranked] *Omoscleria* on the same p. 219.

Scleria subg. *Schizolepis* (Schr. ex Nees) C.B. Clarke, Fl. Cap. 7: 294. 1898. – Type: see *Schizolepis* Schr. ex Nees.

[*Scleria* subg. *Sclerella* Rchb., Consp. Regn. Veg. 56. 1828. *Nom. nud.*]

Scleria subg. *Scleria*, autonym.

Scleria subg. *Tessellatae* (C.B. Clarke) C.B. Clarke, Bull. Misc. Inform. Kew, Addit. Ser. 8: 132. 1908. – Type: see *Scleria* sect. *Tessellatae* C.B. Clarke.

[*Scleria* subg. *Veroscleriae* H. Pfeiff., Revista Sudamer. Bot. 10: 219. 1956. *Nom. inval.*: ICBN Art. 36.1.]

Excluded subgeneric names

Scleria subg. *Becquerelia* (Brongn.) C.B. Clarke, Bull. Misc. Inform. Kew, Addit. Ser. 8: 134. 1908. – Type: see *Becquerelia* Brongn.

Scleria subg. *Diplacrum* (R. Br.) T. Koyama, J. Fac. Sci. Univ. Tokyo, Sect. 3, Bot. 8: 135. 1961. – Type: see *Diplacrum* R. Br.

Sectional names

[*Scleria* sect. *Abortivae* Cherm. in Humbert, Fl. Madagascar 29: 262. 1937. *Nom. inval.*: ICBN Art. 36.1.]

Scleria sect. *Acriulus* (Ridl.) R. W. Haines & Lye, Sedges Rushes East Africa 357. 1983. – Type: see *Acriulus* Ridl.

Scleria sect. *Browniae* C.B. Clarke in Urb., Symb. Antill. 2: 137. 1900 (as “*Brownieae*”). – Type: *Scleria brownii* Kunth [ICBN Art. 22.6.]

Scleria sect. *Carphiformes* J. Kern, Blumea 11: 193. 1961. – Type: *Scleria carphiformis* Ridl. [ICBN Art. 22.6.]

Scleria sect. *Corymbosae* Pax in Engler & Prantl, Nat. Pflanzenfam. 2(2): 121. 1888. – Type: *Scleria corymbosa* Roxb. [ICBN Art. 22.6.]

Scleria sect. *Elatae* C.B. Clarke in Hook. f., Fl. Brit. India 6: 689. 1894. – Type: *Scleria elata* Thwaites [ICBN Art. 22.6.]

[*Scleria* sect. *Eu-Scleria* C.B. Clarke in Urb., Symb. Antill. 2: 138. 1900. *Nom. inval.*: ICBN Art. 21.3.]

Scleria sect. *Foveolidia* Raf., Bull. Bot., Geneva 1: 219. 1830. – Type: *Scleria reticularis* Michx., lectotype here designated.

Scleria sect. *Hirtellae* (C.B. Clarke) C.B. Clarke, Bull. Misc. Inform., Kew, Addit. Ser. 8: 132. 1908. – Type: see *Scleria* [unranked] *Hirtellae* C.B. Clarke. Note: See unranked; even though Clarke, in 1908 in his sect. *Hirtellae*, included only species with rhizome (a group of species in which he again erroneously listed “*S. hirtella* Sw.”), *S.* sect. *Hirtellae* can only be used for a section including the type, the annual *S. hirtella* Sw.

Scleria sect. *Hymenolytrum* (Schrad. ex Nees) Core, Brittonia 2: 10. 1936. – Type: see *Hymenolytrum* Schrad. ex Nees.

Scleria sect. *Hypoporum* (Nees) C.B. Clarke in Urb., Symb. Antill. 2: 137. 1900. – Type: see *Hypoporum* Nees.

Scleria sect. *Lithospermae* (C.B. Clarke) C.B. Clarke, Bull. Misc. Inform., Kew, Addit. Ser. 8: 132. 1908 (as “*Lithospermeae*”). – Type: see *Scleria* [unranked] *Lithospermae* C.B. Clarke.

Scleria sect. *Margaleia* Raf., Bull. Bot., Geneva 1: 219. 1830. – Type: *Scleria poaeformis* Retz., lectotype here designated.

Scleria sect. *Ophryoscleria* (Nees) C.B. Clarke in Urb., Symb. Antill. 2: 138. 1900. – Type: see *Ophryoscleria* Nees.

Scleria sect. *Paniculatae* Pax in Engler & Prantl, Nat. Pflanzenfam. 2(2): 121. 1888. – Type: *Scleria latifolia* Sw., lectotype here designated.

Scleria sect. *Primoscleriae* H. Pfeiff., Repert. Spec. Nov. Regni Veg. 52: 162. 1943, *nom. illeg.* – Type: *Scleria hirtella* Sw. [ICBN Art. 52.2.] ≡ *Scleria* sect. *Hirtellae* (C.B. Clarke) C.B. Clarke 1908.

[*Scleria* sect. *Racemosae* Cherm. in Humbert, Fl. Madagascar 29: 265. 1937. *Nom. inval.*: ICBN Art. 36.1.]

Scleria sect. *Rugunia* Raf., Bull. Bot., Geneva 1: 219. 1830. – Type: *Scleria foveolata* Cav., lectotype here designated.

Scleria sect. *Schizolepis* (Schrad. ex Nees) C.B. Clarke in Hook. f., Fl. Brit. India 6: 694. 1894. – Type: see *Schizolepis* Schrad. ex Nees.

Scleria sect. *Scirposcleria* Griseb., Cat. Pl. Cub. 249. 1866. – Type: *Scleria gracilis* Elliott 1824, non L. C. Rich. 1792 (≡ *Scleria georgiana* Core).

Scleria sect. *Scleria*, autonym.

Scleria sect. *Spicatae* (Boeck.) Pax in Engler & Prantl, Nat. Pflanzenfam. 2(2): 121. 1888. – Type: see *Scleria* [unranked] *Spicatae* Boeck.

Scleria sect. *Tessellatae* C.B. Clarke in Hook. f., Fl. Brit. India 6: 686. 1894. – Type: *Scleria tessellata* Willd. [ICBN Art. 22.6.]

Excluded sectional names

Scleria sect. *Becquerelia* (Brongn.) C.B. Clarke in Urb., Symb. Antill. 2: 139. 1900. – Type: see *Becquerelia* Brongn.

Scleria sect. *Diplacrum* (R. Br.) Kern, Blumea 11: 208. 1961. – Type: see *Diplacrum* R. Br.

Scleria sect. *Sphaeropus* (Boeck.) Kern, Blumea 11: 208. 1961. – Type: see *Sphaeropus* Boeck.

Serial names

Scleria ser. *Dioicae* Benth., Fl. Austral. 7: 432. 1878. – Type: *Scleria sphacelata* F. Muell.

Scleria ser. *Elatae* (C.B. Clarke) T. Koyama, J. Fac. Sci. Univ. Tokyo, Sect. 3, Bot. 8: 135. 1961. – Type: see *Scleria* sect. *Elatae* C.B. Clarke.

Scleria ser. *Laxae* Benth., Fl. Austral. 7: 428. 1878. – Type: *Scleria laxa* R. Br. [ICBN Art. 22.6.]

Scleria ser. *Macrostachyae* Benth., Fl. Austral. 7: 432. 1878. – Type: *Scleria oryzoides* Presl.

Scleria ser. *Polystachyae* Benth., Fl. Austral. 7: 430. 1878. – Type: *Scleria hebecarpa* Nees, lectotype here designated.

Scleria ser. *Scleria*, autonym.

Scleria ser. *Subspicatae* Benth., Fl. Austral. 7: 428. 1878. – Type: *Scleria brownii* Kunth, lectotype here designated.

Scleria ser. *Tessellatae* (C.B. Clarke) T. Koyama, J. Fac. Sci. Univ. Tokyo, Sect. 3, Bot. 8: 135. 1961. – Type: see *Scleria* sect. *Tessellatae* C.B. Clarke.

[*Scleria* ser. *Tuberculatae* (Core) Core in Maguire & al., Mem. New York Bot. Gard. 12(3): 55. 1965. *Nom. inval.*: ICBN Art. 33.3.]

Excluded serial names

Scleria ser. *Axillares* Benth., Fl. Austral. 7: 426. 1878. – Type: *Scleria caricina* (R. Br.) Benth. (≡ *Diplacrum caricinum* R. Br.), lectotype here designated.

Unranked names

[*Scleria* [unranked] *Corymbosae* Boeck., Linnaea 38: 536. 1874. *Nom. nud.*]

Scleria [unranked] *Hirtellae* C.B. Clarke in Dyer, Fl. Trop. Afr. 8: 493. 1902. – Type: *Scleria hirtella* Sw. [ICBN Art. 22.6.] Note: *Scleria hirtella* Sw., an annual species, even though cited by Clarke, does not occur in Africa. Clarke (and others)

erroneously applied that name for *S. distans* Poir., a perennial species (Raynal 1976: 216). *Scleria* [unranked] *Hirtellae* C.B. Clarke 1902 included both annuals and perennials.

Scleria [unranked] *Hypoporum* (Nees) Endl., Gen. Pl. 112. 1836. – Type: see *Hypoporum* Nees. Note: Kern (1961: 196) stated that Endlicher's name was intended as sectional, but this view was contradicted by Brizicky (1969: 658) who demonstrated that both subgenera and sections were represented among the infra-generic divisions established by Endlicher. Therefore all Endlicher's generic subdivisions must be regarded and dealt with as taxonomic categories published without indication of rank.

Scleria [unranked] *Lithospermae* C.B. Clarke in Dyer, Fl. Trop. Afr. 8: 494. 1902 (as "*Lithospermeae*"). – Type: *Scleria lithosperma* (L.) Sw. [ICBN Art. 22.6.]

Scleria [unranked] *Omoscleria* (Nees) H. Pfeiff., Revista Sudamer. Bot. 10: 219. 1956. – Type: see *Omoscleria* Nees.

[*Scleria* [unranked] *Paniculatae* Boeck., Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn 1869: 151. 1870. *Nom. nud.*]

Scleria [unranked] *Primoscleriae* H. Pfeiff., Repert. Spec. Nov. Regni Veg. 55: 114. 1953. – Type: *Scleria hirtella* Sw. [ICBN Art. 7.7.] ≡ *Scleria* [unranked] *Hirtellae* C.B. Clarke 1902.

[*Scleria* [unranked] *Sellowianae* H. Pfeiff., Revista Sudamer. Bot. 10: 219. 1956. *Nom. inval.*: ICBN Art. 36.1.]

Scleria [unranked] *Spicatae* Boeck., Linnaea 38: 438. 1874. – Type: *Scleria bulbifera* Hochst. ex Rich., lectotype here designated.

Scleria [unranked] *Tessellatae* (C.B. Clarke) C.B. Clarke in Dyer, Fl. Trop. Afr. 8: 494. 1902. – Type: see *Scleria* sect. *Tessellatae* C.B. Clarke.

Scleria [unranked] *Tuberculatae* Core, Brittonia 1: 241. 1934. – Type: *Scleria alpina* Core, lectotype here designated.

Excluded unranked names

Scleria [unranked] *Becquerelia* (Brongn.) Endl., Gen. Pl. 112. 1836 (as "*Bequerela*"). – Type: see *Becquerelia* Brongn.

[*Scleria* [unranked] *Cymiferae* Boeck., Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn 1869: 155. 1870. *Nom. nud.*]

Acknowledgements

Literature was provided by the librarians of BR, K, MO and NY. We would like to thank Dan H. Nicolson (US) for valuable comments on some questions of typification. John McNeill (E) and two anonymous reviewers provided us with valuable comments on the manuscript.

Literature cited

Britton, N. L. 1907. The sedges of Jamaica. *Bull. Dept. Agric. Jamaica* 5, suppl. 1: 1–19.

- Brizicky, G. K. 1969. Subgeneric and sectional names: their starting points and early sources. *Taxon* 18: 643-660.
- Camelbeke, K. & Goetghebeur, P. 2000. (1455) Proposal to conserve *Scleria flagellum-nigrorum* with a conserved type (*Cyperaceae*). *Taxon* 49: 295-296.
- Clarke, C. B. 1902. *Cyperaceae*. Pp. 266-524 in: Dyer, W. T. (ed.), *Flora of tropical Africa*, vol. 8. Ashford.
- Franklin Hennessy, E. F. 1985. The genus *Scleria* in southern Africa. *Bothalia* 15: 505-530.
- Kern, J. H. 1961. Florae Malesianae precursores XXX. The genus *Scleria* in Malaysia. *Blumea* 11: 140-218.
- 1964. *Acriulus* once more reduced (*Cyperaceae*). *Blumea* 12: 41-44.
- Koyama, T. 1967. *Cyperaceae-Mapanioideae*. *Mem. New York Bot. Gard.* 17(1): 23-79.
- & Maguire, B. 1965. *Cyperaceae* tribe *Lagenocarpeae*. *Mem. New York Bot. Gard.* 12(3): 8-54.
- Raynal, J. 1976. Notes Cypérologiques: 27. Identification de deux *Scleria* de Poiret. *Adansonia* ser. 2. 16: 211-217.
- 1977. Notes Cypérologiques: 30. *Catagyna* Pal. Beauv. ex Lest. enfin identifié. *Adansonia*, ser. 2, 17: 35-41.