

WHAT BUCKWHEAT IS THAT?

Buckwheats in the Mt. Diablo area

1.	Plants perennial			
	a.	Plant lov	w growing, often forming mats. Flowers bright yellow.	
			Sulfur flower (Eriogonum umbellatum var. bahiiforme)	
	b.	Plants t	all with branching stems. Flowers white, pinkish, or pale yellow.	
			Naked-stem buckwheat (<i>Erigonum nudum</i> var. <i>auriculatum</i>)	
2.	Plants annual			
		pink. S	and stems sticky (glandular), not covered with white hairs. Flowers white or tems with many branches	
	b.		and stems usually covered with fine white hairs. Flower color various.	
			vers white to rose pink. Flower clusters spaced along elongate stems	
		_	erally < 3 clusters per branch, occasionally 4) or strictly at ends of branches	
			at the junction of two branches (axes).	
		(α)	Leaves at base and often on the stems. Leaves broadest at tip or middle (oblanceolate to elliptic).	
			(i) Stems with lengthwise ridges. Flowers pink to rose in clusters at	
			ends of branches. Flowers and flower cluster sticky.	
			(ii) Stems without ridges. Flowers light rose pink in clusters at ends of	
			branches and at axes of flower stems. Flower clusters sticky.	
			Flowers smooth, not sticky.	
		(b)	Leaves rounded with heart-shaped base. Leaves only at the base of stems. Flowers rose to white with red veins in clusters along the stems.	
	(2) Flowers white to yellow or red, drying reddish. Flower clusters spaced al			
	elongate stems (generally > 5 clusters per branch).			
			Basal leaves ellipse-shaped, tapering to leaf stem (petiole). Flower cluster	
			with fine white hairs.	
			(i) Flowers white to yellow. Stems with uneven branches.	
			Slender wooly buckwheat (Eriogonum gracile)	
			(ii) Flowers yellowish to red. Stems wand-like with few branches.	
			Wand buckwheat (Eriogonum roseum)	
		(b)	Basal leaves round. Flower clusters and flower stems without fine white	
			hairsEar-shaped buckwheat (Eriogonum luteolum var. luteolum)	



Flower structures



Leaf Shapes





Sulfur Flower (*Eriogonum umbellatum* var. *bahiiforme*)



Naked stem buckwheat (Eriogonum nudum var. auriculatum)









Mt. Diablo buckwheat (Eriogonum truncatum)



Coville's buckwheat (Eriogonum covilleanum)









Slender woolly buckwheat (Eriogonum gracile)




Wand buckwheat (Eriogonum roseum)





Anglestem buckwheat (*Eriogonum angulosum*)



Ear-shaped buckwheat (Eriogonum luteolum var. luteolum)







Unarmed buckwheat (Eriogonum inerme)

Photo Credits

Eriogonum umbellatum var. bahiifolium

Habit - Brother Alfred Brousseau © 1995 Saint Mary's College of California Erigonum umbellatum flower head - © 2003 Christopher L. Christie

Eriogonum nudum var. auriculatum

Habit - © James Reveal

Habit - Brother Alfred Brousseau © 1995 Saint Mary's College of California

Inflorescence - Walter Knight © 1999 California Academy of Sciences

Eriogonum truncatum

All images © Scott Hein

Eriogonum covilleanum

Charles Webber © 1998 California Academy of Sciences

William R. Hewlett © 1999 California Academy of Sciences

Eriogonum gracile

Habit © California State University San Bernardino

Inflorescence © 2001 Thomas M. Elder, M.D.

Eriogonum roseum

Habit and flower head © 2003 Christopher L. Christie

Eriogonum angulosum

©2005 Christopher Christie

Eriogonum luteolum var. luteolum

Inflorescence - Brother Alfred Brousseau @ 1995 Saint Mary's College of California

Flower head - @ 2001 James B. Gratiot

Eriogonum inerme

Photographs of herbarium specimen, courtesy of Jepson Herbarium