

EDWARDS'S
BOTANICAL REGISTER:

OR,

ORNAMENTAL FLOWER-GARDEN
AND SHRUBBERY:

CONSISTING OF

COLOURED FIGURES OF PLANTS AND SHRUBS,
CULTIVATED IN BRITISH GARDENS;

ACCOMPANIED BY THEIR

History, Best Method of Treatment in Cultivation, Propagation, &c.

CONTINUED

By JOHN LINDLEY, Ph. D. F.R.S. L.S. AND G.S.

PROFESSOR OF BOTANY IN THE UNIVERSITY OF LONDON,

AND THE ROYAL INSTITUTION OF GREAT BRITAIN.

&c. &c. &c.

New Series.

VOL. VIII. D

OR VOL. XXI. OF THE ENTIRE WORK.

—viret semper—nec fronde caduca
Carpitur.

MISSOURI
BOTANICAL
GARDEN.

LONDON:

JAMES RIDGWAY AND SONS, PICCADILLY.

M.DCCC.XXXVI.

Rare Book
S13407
E 39
1836
v. 21

New series
v. 8

ALPHABETICAL INDEX

TO

VOL. VIII. OF THE NEW SERIES.

	<i>Folium</i>		<i>Folium</i>
Angræcum distichum	1781	Gesneria allagophylla	1757
Angræcum micranthum	1772	Gesneria faucialis	1785
Arctostaphylos tomentosa	1791	Govenia superba	1795
Arbutus procera	1753	Hibiscus Rosa sinensis	1826
Aristolochia fœtens	1824	Indigofera atropurpurea	1744
Azara dentata	1788	Ipomœa Aitoni	1794
Begonia petalodes	1757	Kennedya Marryattæ	1790
Berberis dealbata	1750	Lælia anceps	1751
Bletia reflexa	1760	Lasthenia californica	1823
Brassia Lanceana	1754	Lasthenia glabrata	1780
Calceolaria angustiflora	1743	Lepanthes tridentata	1762
Calotropis procera	1792	Macradenia triandra	1815
Campanula garganica	1768	Maxillaria cristata	1811
Caprifolium hispidulum	1761	Maxillaria crocea	1799
Cereus triangularis	1807	Maxillaria densa	1804
Chironia peduncularis	1803	Maxillaria picta	1802
Clavija ornata	1764	Mimulus luteus, var. <i>variegatus</i>	1796
Clianthus puniceus	1775	Monachanthus viridis	1752
Coccoloba virens	1816	Myanthus barbatus	1778
Colletia horrida	1776	Ochranthe arguta	1819
Coryanthes maculata	1793	Oncidium citrinum	1758
Cosmelia rubra	1822	Oncidium Lemonianum	1789
Cratægus Douglasii	1810	Oncidium pulchellum	1787
Cycnoches Loddigesii	1742	Orobus atropurpureus	1763
Daubenya aurea	1813	Oxalis Piottæ	1817
Dendrobium Pierardi	1756	Pæonia Moutan lacera	1771
Dendrobium cupreum	1779	Pentstemon staticifolius	1770
Dendrobium densiflorum	1828	Pholidota imbricata	1777
Dracæna terminalis	1749	Physianthus albens	1759
Dyckia rariflora	1782	Pimelia ligustrina	1827
Edwardsia chilensis	1798	Pleurothallis Grobyi	1797
Elichrysum bicolor	1814	Pleurothallis picta	1825
Empetrum rubrum	1783	Prunus japonica	1801
Epidendrum gracile	1765	Psoralea macrostachya	1769
Eriogonum compositum	1774	Rhodochiton volubile	1755
Erythronium grandiflorum	1786	Rhododendron pulcherrimum	1820
Eutoca divaricata	1784	Russellia juncea	1773
Eutoca viscida	1808	Stanhopea oculata	1800
Fernandezia acuta	1806	Tritoma Burchelli	1745
Fuchsia discolor	1805	Vanda teres	1809
Galatella punctata	1818	Verbena multifida contracta	1766
Gardoquia Gilliesii	1812	Verbena sulphurea	1748
Gardoquia Hookeri	1747	Zephyranthes Spofforthiana	1746

GENERAL ALPHABETICAL INDEX

TO

VOLS. I. II. III. IV. V. VI. VII. AND VIII.

OF

THE NEW SERIES.

	<i>Vol. Folium</i>		<i>Vol. Folium</i>
Acacia albida	16 . 1317	Aphelandra cristata	18 . 1477
Acacia leprosa	17 . 1441	Arbutus procera	21 . 1753
Acacia lunata	16 . 1352	Arctostaphylos tomentosa	21 . 1791
Acacia pentadenia	18 . 1521	Argemone grandiflora	15 . 1264
Acacia uncinata	16 . 1332	Argemone ochroleuca	16 . 1343
Acanthophippium bicolor	20 . 1730	Aristolochia caudata	17 . 1453
Acæna pinnatifida	15 . 1271	Aristolochia Chilensis	20 . 1680
Aceras secundiflora	18 . 1525	Aristolochia cymbifera	18 . 1543
Adenotrichia amplexicaulis	14 . 1190	Aristolochia fœtens	21 . 1824
Adesmia Loudonia	20 . 1720	Aristolochia trilobata	17 . 1399
Aerides cornutum	18 . 1485	Asphodelus luteus, var. <i>sibiricus</i>	18 . 1507
Agave geminiflora	14 . 1145	Aster adulterinus	19 . 1571
Alströmeria psitacina	18 . 1540	Aster eminens, var. <i>virgineus</i>	20 . 1656
Alströmeria pulchella, var. <i>pilosa</i>	17 . 1410	Aster amygdalinus	18 . 1517
Amaryllis acuminata, var. <i>longi-</i> <i>pedunculata</i>	14 . 1188	Aster concinnus	19 . 1619
Amaryllis coranica, var. <i>pallida</i>	15 . 1219	Aster cordifolius	19 . 1597
Amaryllis intermedia	14 . 1148	Aster coridifolius	18 . 1487
Amaryllis kermesina	19 . 1638	Aster cyaneus	18 . 1495
Amelanchier florida	19 . 1589	Aster eminens	19 . 1614
Amelanchier sanguinea	14 . 1171	Aster fragilis	18 . 1537
Amygdalus communis, var. <i>macro-</i> <i>carpa</i>	14 . 1160	Aster lævis	18 . 1500
Amygdalus persica, var. <i>alba</i>	19 . 1586	Aster pallens	18 . 1509
Anemone vitifolia	16 . 1385	Aster puniceus, var. <i>demissus</i>	19 . 1636
Angræcum distichum	21 . 1781	Aster spectabilis	18 . 1527
Angræcum eburneum	18 . 1522	Astragalus succulentus	16 . 1324
Angræcum micranthum	21 . 1772	Audibertia incana	17 . 1469
Anomatheca cruenta	16 . 1369	Azalea calendulacea, var. <i>subcu-</i> <i>prea</i>	16 . 1366
Anona laurifolia	16 . 1328	Azalea calendulacea, var. <i>lepida</i>	17 . 1402
Anthocercis viscosa	19 . 1624	Azalea calendulacea, var. <i>Staple-</i> <i>toniana</i>	17 . 1407
Antholyza æthiopica, var. <i>minor</i>	14 . 1159	Azalea Indica, var. <i>lateritia</i>	20 . 1700
Anthurium gracile	19 . 1635	Azalea indica, var. <i>variegata</i>	20 . 1716

GENERAL INDEX TO THE NEW SERIES.

	<i>Vol. Folium</i>		<i>Vol. Folium</i>
Azalea nudiflora, var. <i>scintillans</i>	17 . 1461	Calceolaria Herbertiana, var. <i>par-</i>	
Azalea nudiflora, var. <i>thyrsiflora</i>	16 . 1367	<i>viflora</i>	19 . 1576
Azalea pontica, var. <i>sinensis</i>	15 . 1253	Calceolaria polifolia	20 . 1711
Azalea pontica, var. <i>versicolor</i>	18 . 1559	Calceolaria purpurea	19 . 1621
Azara dentata	21 . 1788	Calceolaria rugosa	19 . 1588
Banksia littoralis	16 . 1363	Calceolaria sesilis	19 . 1628
Banksia prostrata	19 . 1572	Calceolaria viscosissima	19 . 1611
Banksia quercifolia	17 . 1430	Calceolaria, Mr. Young's	17 . 1448
Banksia speciosa	20 . 1728	Calliprora Iutea	19 . 1570
Banksia undulata	16 . 1316	Calochortus macrocarpus	14 . 1152
Barleria lupulina	18 . 1483	Calochortus luteus	19 . 1567
Bartholina pectinata	20 . 1653	Calochortus splendens	20 . 1676
Batemannia Colleyi	20 . 1714	Calochortus venustus	20 . 1669
Bauhinia cumanensis	14 . 1133	Calotropis procera	21 . 1792
Begonia heracleifolia	20 . 1668	Camassia esculenta	18 . 1486
Begonia petalodes	21 . 1757	Camellia japonica, var. <i>imbricata</i>	17 . 1398
Begonia villosa	15 . 1252	Camellia japonica, var. <i>punctata</i>	15 . 1267
Beloperone oblongata	20 . 1657	Camellia japonica, var. <i>Reeve-</i>	
Benthamia fragifera	19 . 1579	<i>siana</i>	18 . 1501
Berberis aquifolium	17 . 1425	Campanula fragilis	20 . 1738
Berberis dealbata	21 . 1750	Campanula garganica	21 . 1768
Berberis glumacea	17 . 1426	Canavalia bonariensis	14 . 1199
Berberis repens	14 . 1176	Canna Achiras	16 . 1358
Bignonia Cherere	15 . 1301	Canna discolor	15 . 1231
Billardiera ovalis	20 . 1719	Canna lagunensis	16 . 1311
Billbergia pyramidalis, var. <i>bicolor</i>	14 . 1181	Canna speciosa	15 . 1276
Blackwellia padiflora	16 . 1308	Capparis acuminata	16 . 1322
Bletia florida	17 . 1401	Caprifolium hispidulum	21 . 1761
Bletia gracilis	20 . 1681	Caprifolium longiflorum	15 . 1232
Bletia reflexa	21 . 1760	Caprifolium occidentale	17 . 1457
Bœbera incana	19 . 1602	Cassia biflora	16 . 1310
Brasavola nodosa	17 . 1465	Cassia Herbertiana	17 . 1422
Brasavola Perrinii	18 . 1561	Castilleja coccinea	14 . 1136
Brassia Lanceana	21 . 1754	Catasetum luridum	20 . 1667
Brodiaea grandiflora	14 . 1183	Catasetum semiapertum	20 . 1708
Browallia grandiflora	16 . 1384	Cattleya crispa	14 . 1172
Brownlowia elata	17 . 1472	Cattleya guttata	17 . 1406
Brugmansia bicolor	20 . 1739	Cereus, crimson creeping	19 . 1565
Brunsvigia ciliaris	14 . 1153	Cereus triangularis	21 . 1807
Brunsvigia grandiflora	16 . 1335	Ceropegia elegans	20 . 1706
Buddleia heterophylla	15 . 1259	Chasmonia incisa	15 . 1244
Burtonia conferta	19 . 1600	Cheiranthus mutabilis	17 . 1431
Cactus Ackermanni	16 . 1331	Chelone centranthifolia	20 . 1737
Cactus speciosissimus, var. <i>lateri-</i>		Chelone nemorosa	14 . 1211
<i>tius</i>	19 . 1596	Chironia peduncularis	21 . 1803
Calandrinia arenaria	19 . 1605	Chorozema ovatum	18 . 1528
Calandrinia grandiflora	14 . 1194	Chorozema triangulare	18 . 1513
Calandrinia speciosa	19 . 1598	Chrysanthemum indicum	15 . 1287
Calanthe densiflora	19 . 1646	Chrysanthemum indicum, var. <i>pla-</i>	
Calathea grandifolia	14 . 1210	<i>num</i>	18 . 1502
Calceolaria angustiflora	21 . 1743	Cirrhæa Loddigesii	18 . 1538
Calceolaria arachnoidea	17 . 1454	Clarkia elegans	19 . 1575
Calceolaria ascendens	14 . 1215	Clavija ornata	21 . 1764
Calceolaria chiloensis	17 . 1476	Clematis chlorantha	18 . 1234
Calceolaria crenatiflora	19 . 1609	Cleome speciosissima	16 . 1312
Calceolaria diffusa	16 . 1374	Clerodendron hastatum	16 . 1307
Calceolaria floribunda	14 . 1214	Clianthus puniceus	21 . 1775
Calceolaria Herbertiana	16 . 1313	Clintonia elegans	15 . 1241
		Clivia nobilis	14 . 1182
		Coburgia fulva	18 . 1497

GENERAL INDEX TO THE NEW SERIES.

	<i>Vol. Folium</i>		<i>Vol. Folium</i>
Coccoloba virens	21 . 1816	Dendrobium speciosum	19 . 1610
Coleus aromaticus	18 . 1520	Deutzia scabra	20 . 1718
Colletia horrida	21 . 1776	Dianthus Libanotis	18 . 1548
Collinsia bicolor	20 . 1734	Digitalis laciniata	14 . 1201
Collomia grandiflora	14 . 1174	Dillwynia glycinifolia	18 . 1514
Collomia coccinea	19 . 1622	Diospyrus Mabola	14 . 1139
Collomia heterophylla	16 . 1347	Diplopappus incanus	20 . 1693
Collomia linearis	14 . 1166	Dracæna surculosa	14 . 1169
Colutea nepalensis	20 . 1727	Dracæna terminalis	12 . 1749
Combretum comosum	14 . 1105	Drimia villosa	16 . 1346
Combretum grandiflorum	19 . 1631	Duvaua dependens	19 . 1573
Conanthera campanulata	14 . 1193	Duvaua latifolia	19 . 1580
Conocephalus naucleiflorus	14 . 1203	Duvaua ovata	19 . 1568
Convolvulus farinosus	16 . 1325	Dyckia rariflora	21 . 1782
Cordia grandiflora	18 . 1491		
Coreopsis Atkinsoniana	16 . 1376	Echeveria gibbiflora	15 . 1247
Coreopsis aurea	15 . 1228	Echinocactus oxygonus	20 . 1717
Correa pulchella	15 . 1224	Echinocactus Eyriesii	20 . 1707
Coryanthes maculata	21 . 1793	Echites stellaris	20 . 1664
Corydalis bracteata	19 . 1644	Edwardsia chilensis	21 . 1798
Cosmelia rubra	21 . 1822	Elæagnus angustifolia	14 . 1156
Costus pictus	19 . 1594	Elichrysum bicolor	21 . 1814
Cotoneaster frigida	15 . 1229	Empetrum rubrum	21 . 1783
Cotoneaster laxiflora	15 . 1305	Epacris nivalis	18 . 1531
Cotoneaster microphylla, var. <i>Uva</i>		Epidendrum gracile	21 . 1765
<i>Ursi</i>	14 . 1187	Epidendrum odoratissimum	17 . 1415
Crassula turrita	16 . 1344	Epidendrum oncioides	19 . 1623
Cratægus cordata	14 . 1151	Eranthemum fœcundum	17 . 1494
Cratægus Douglasii	21 . 1810	Erica codonodes	20 . 1698
Cratægus heterophylla	14 . 1161	Eriogonum compositum	21 . 1774
Crinum latifolium	15 . 1297	Eriophyllum cæspitosum	14 . 1167
Crocus vernus, var. <i>leucorhynchus</i>	17 . 1416	Erythrina carnea	16 . 1327
Crocus vernus, var. <i>pictus</i>	17 . 1440	Erythrina poiianthes	15 . 1246
Crotalaria verrucosa	14 . 1137	Erythrina poiianthes, var. <i>subinermis</i>	19 . 1617
Cuphea Llavea	16 . 1386	Erythronium grandiflorum	21 . 1786
Cyclobothra alba	20 . 1661	Escallonia montevidensis	17 . 1467
Cyclobothra pulchella	20 . 1662	Eschscholtzia californica	14 . 1168
Cyclobothra lutea	20 . 1663	Eschscholtzia crocea	20 . 1677
Cynoches Loddigesii	21 . 1742	Eulophia ensata	14 . 1147
Cymbidium marginatum	18 . 1530	Eulophia Mackaiana	17 . 1433
Cyrtanthus carneus	17 . 1462	Eupatorium glandulosum	20 . 1723
Cyrtochilum flavescens	19 . 1627	Euphoria Longan	20 . 1729
Cyrtopodium Woodfordii	18 . 1508	Eurybia corymbosa	18 . 1532
Cypripedium macranthos	18 . 1534	Eurycles Cunninghamii	18 . 1506
Cypripedium spectabile	20 . 1666	Eutoca divaricata	21 . 1784
Cytisus multiflorus	14 . 1191	Eutoca multiflora	14 . 1180
		Eutoca viscida	21 . 1803
Daphne hybrida	14 . 1177		
Daubinya aurea	21 . 1813	Francoa appendiculata	19 . 1645
Delphinium Menziesii	14 . 1192	Fernandezia acuta	21 . 1806
Delphinium speciosum	18 . 1503	Fuchsia bacillaris	18 . 1480
Dendrobium aggregatum	20 . 1695	Fuchsia discolor	21 . 1805
Dendrobium anceps	15 . 1239	Fuchsia globosa	18 . 1556
Dendrobium chrysanthum	15 . 1299	Fuchsia microphylla	15 . 1269
Dendrobium cupreum	21 . 1779	Fuchsia thymifolia	15 . 1284
Dendrobium densiflorum	21 . 1828		
Dendrobium longicornu	16 . 1315	Gaillardia aristata	14 . 1186
Dendrobium moniliforme	16 . 1314	Galatella punctata	21 . 1818
Dendrobium Pierardi	21 . 1756	Galipea odoratissima	17 . 1420
Dendrobium secundum	15 . 1291		

GENERAL INDEX TO THE NEW SERIES.

	<i>Vol. Folium</i>		<i>Vol. Folium</i>
Gardoquia Gilliesii	21 . 1812	Iris bicolor	17 . 1404
Gardoquia Hookeri	21 . 1747	Iris tenax	15 . 1218
Garrya elliptica	20 . 1686	Ismene Amancaes, var. <i>sulphurea</i>	20 . 1665
Gastrolobium retusum	19 . 1647	Isopogon formosus	15 . 1288
Gaultheria Shallon	17 . 1411	Jasminum acuminatum	15 . 1296
Genista procumbens	14 . 1150	Jasminum Wallichianum	17 . 1409
Geodorum fucatum	20 . 1687	Justicia carnea	17 . 1397
Gesnera allagophylla	21 . 1767	Justicia guttata	16 . 1334
Gesnera faucialis	21 . 1785	Justicia picta	15 . 1227
Gesnera macrostachya	14 . 1202	Justicia quadrangularis	16 . 1340
Gesnera rutila	14 . 1158	Justicia venusta	16 . 1380
Gesnera rutila, var. <i>atrosanguinea</i>	15 . 1279	Kæmpferia Roscoeana	14 . 1212
Gesnera Suttoni	19 . 1637	Kennedyia dilatata	18 . 1526
Geum chilense, var. <i>grandiflorum</i>	16 . 1348	Kennedyia inophylla	17 . 1421
Gilia Achilleæfolia	20 . 1682	Kennedyia Marryattæ	21 . 1790
Gilia coronopifolia	20 . 1691	Kennedyia monophylla, var. <i>longi-</i>	
Gilia tricolor	20 . 1704	<i>racemosa</i>	16 . 1336
Gilia capitata	14 . 1170	Kennedyia nigricans	20 . 1715
Gladiolus psittacinus	17 . 1442	Lachenalia pallida	16 . 1350
Glycine biloba	17 . 1413	Lælia anceps	21 . 1751
Gompholobium capitatum	18 . 1563	Lalage ornata	20 . 1722
Gompholobium Knightianum	17 . 1468	Lasthenia californica	21 . 1823
Gompholobium tenue	19 . 1615	Lasthenia glabrata	21 . 1780
Gompholobium venulosum	19 . 1574	Lathyrus californicus	14 . 1144
Gompholobium marginatum	18 . 1490	Lathyrus tingitanus	16 . 1388
Gompholobium tomentosum	17 . 1474	Ledocarpum pedunculare	17 . 1392
Gongora maculata	19 . 1616	Lepanthes tridentata	21 . 1762
Govenia superba	21 . 1795	Lepechinia spicata	15 . 1292
Grevillea concinna	16 . 1383	Leptosiphon androsaceus	20 . 1710
Grevillea punicea	16 . 1319	Leptosiphon densiflorus	20 . 1725
Grobya Amherstiæ	20 . 1740	Leptotes bicolor	19 . 1625
Guettarda speciosa	17 . 1393	Leucocoryne odorata	15 . 1293
Habranthus Andersoni	16 . 1345	Leucopogon parviflorus	18 . 1560
Habranthus Bagnoldi	17 . 1396	Liatris scariosa	20 . 1654
Habranthus Phycelloides	17 . 1417	Libertia formosa	19 . 1630
Hakea linearis	18 . 1489	Limnanthes Douglasii	20 . 1673
Hamelia ventricosa	14 . 1195	Limnocharis Humboldtii	19 . 1640
Haylockia pusilla	16 . 1371	Linaria Dalmatica	20 . 1683
Hedychium coccineum	14 . 1209	Linum mexicanum	16 . 1326
Helianthus lenticularis	15 . 1265	Linum sibiricum, var. <i>Lewisii</i>	14 . 1163
Helianthus tubæformis	18 . 1519	Liparis elata	14 . 1175
Heliconia pulverulenta	19 . 1648	Liparis guineensis	20 . 1671
Herminium cordatum	18 . 1499	Lissanthe sapida	15 . 1275
Hesperoscordum lacteum	19 . 1639	Lithospermum rosmarinifolium	20 . 1736
Heuchera micrantha	15 . 1302	Loasa ambrosiæfolia	16 . 1390
Hibiscus Lindleii	17 . 1395	Loasa Placei	19 . 1599
Hibiscus palustris	17 . 1463	Lobelia longiflora	14 . 1200
Hibiscus Rosa sinensis	21 . 1826	Lobelia purpurea	16 . 1325
Hibiscus splendens	19 . 1629	Lobelia, Low's purple	17 . 1445
Hosackia bicolor	15 . 1257	Lobelia Tupa	19 . 1612
Hosta cœrulea	14 . 1204	Lonicera involucrata	14 . 1179
Hovea chorozemæfolia	18 . 1524	Lophanthus anisatus	15 . 1282
Hovea lanceolata	17 . 1427	Lophospermum erubescens	16 . 1381
Hovea purpurea	17 . 1423	Lotus arenarius	18 . 1488
Hovea villosa	18 . 1512	Lowea berberifolia	15 . 1261
Indigofera atropurpurea	21 . 1744	Lupinus arbustus	15 . 1230
Ipomœa Aitoni	21 . 1794	Lupinus albifrons	19 . 1642
Ipomopsis elegans	15 . 1281		

GENERAL INDEX TO THE NEW SERIES.

	<i>Vol. Folium</i>		<i>Vol. Folium</i>
Lupinus aridus . . .	15 . 1242	Enothera decumbens . . .	15 . 1221
Lupinus densiflorus . . .	20 . 1689	Enothera densiflora . . .	18 . 1593
Lupinus elegans . . .	18 . 1501	Enothera glauca . . .	18 . 1511
Lupinus laxiflorus . . .	14 . 1140	Enothera pallida . . .	14 . 1142
Lupinus lepidus . . .	14 . 1149	Enothera tenella, var. <i>tenuifolia</i>	19 . 1587
Lupinus leptophyllus . . .	20 . 1670	Enothera viminea . . .	15 . 1220
Lupinus littoralis . . .	14 . 1198	Oncidium citrinum . . .	21 . 1758
Lupinus micranthus . . .	15 . 1251	Oncidium cornigerum . . .	18 . 1542
Lupinus mutabilis . . .	18 . 1539	Oncidium altissimum . . .	19 . 1651
Lupinus nanus . . .	20 . 1705	Oncidium ampliatus . . .	20 . 1699
Lupinus ornatus . . .	14 . 1216	Oncidium ciliatum . . .	20 . 1660
Lupinus plumosus . . .	15 . 1217	Oncidium Harrisonianum . . .	19 . 1569
Lupinus polyphyllus, var. <i>albi-</i>		Oncidium Lemonianum . . .	21 . 1789
<i>florus</i> . . .	16 . 1377	Oncidium pulchellum . . .	21 . 1787
Lupinus rivularis . . .	19 . 1595	Ononis peduncularis . . .	17 . 1447
Lupinus Sabinianus . . .	17 . 1435	Ophrys aranifera, var. <i>limbata</i>	14 . 1197
Macradenia triandra . . .	21 . 1815	Opuntia aurantiaca . . .	19 . 1606
Madia elegans . . .	17 . 1458	Opuntia monacantha . . .	20 . 1726
Magnolia Yulan, var. <i>Soulangiana</i>	14 . 1164	Orchis papilionacea . . .	14 . 1155
Malva Munroana . . .	16 . 1306	Orchis foliosa . . .	20 . 1701
Malva purpurata . . .	16 . 1362	Orobus atropurpureus . . .	21 . 1763
Malva umbellata . . .	14 . 1608	Osbeckia nepalensis, var. <i>albiflora</i>	17 . 1475
Mammillaria pulchra . . .	16 . 1329	Oxalis Bowiei . . .	19 . 1585
Mammillaria tenuis . . .	18 . 1523	Oxalis Cummingi . . .	18 . 1545
Maxillaria ciliaris . . .	14 . 1206	Oxalis divergens . . .	19 . 1620
Maxillaria cristata . . .	21 . 1811	Oxalis Piottæ . . .	21 . 1817
Maxillaria crocea . . .	21 . 1799	Oxalis tortuosa . . .	15 . 1249
Maxillaria decolor . . .	18 . 1549	Oxalis variabilis . . .	18 . 1505
Maxillaria densa . . .	21 . 1804	Pachypodium tuberosum . . .	16 . 1321
Maxillaria picta . . .	21 . 1802	Palavia rhombifolia . . .	16 . 1375
Maxillaria racemosa . . .	19 . 1566	Pæonia albiflora, var. <i>Pottsii</i>	17 . 1436
Maxillaria tetragona . . .	17 . 1428	Pæonia hybrida . . .	14 . 1208
Maxillaria viridis . . .	18 . 1510	Pæonia Moutan . . .	20 . 1678
Maytenus chilensis . . .	20 . 1702	Pæonia Moutan lacera . . .	21 . 1771
Masembryanthemum rubrocinctum	20 . 1732	Pæony, Semidouble tree . . .	17 . 1456
Michauxia lævigata . . .	17 . 1451	Pancratium pedale . . .	19 . 1641
Microstylis ophioglossoides . . .	15 . 1290	Papaver Persicum . . .	17 . 1570
Milla biflora . . .	18 . 1555	Passiflora ligularis . . .	19 . 1339
Mimulus luteus, var. <i>variegatus</i>	21 . 1796	Passiflora gossypifolia . . .	19 . 1634
Mimulus propinquus . . .	16 . 1330	Passiflora kermesina . . .	19 . 1633
Mimulus roseus . . .	19 . 1591	Passiflora phœnicea . . .	19 . 1603
Mimulus Smithii . . .	20 . 1674	Pentstemon acuminatum . . .	15 . 1285
Mirbelia Baxteri . . .	17 . 1434	Pentstemon attenuatum . . .	15 . 1295
Monachanthus discolor . . .	20 . 1755	Pentstemon confertum . . .	15 . 1260
Monachanthus viridis . . .	21 . 1752	Pentstemon deustum . . .	16 . 1318
Moscharia pinnatifida . . .	18 . 1564	Pentstemon diffusum . . .	14 . 1132
Myanthus barbatus . . .	21 . 1778	Pentstemon glandulosum . . .	15 . 1262
Myanthus cernuus . . .	20 . 1721	Pentstemon glaucum . . .	15 . 1286
Nanodes discolor . . .	18 . 1541	Pentstemon pruinatum . . .	15 . 1280
Nemophila aurita . . .	19 . 1601	Pentstemon pulchellum . . .	14 . 1138
Nemophila insignis . . .	20 . 1713	Pentstemon Scouleri . . .	15 . 1277
Nierembergia filicaulis . . .	19 . 1649	Pentstemon speciosum . . .	15 . 1270
Nicotiana persica . . .	19 . 1592	Pentstemon staticifolius . . .	21 . 1770
Ochranthe arguta . . .	21 . 1819	Pentstemon triphyllum . . .	15 . 1245
Enothera anisoloba . . .	18 . 1479	Pentstemon venustum . . .	16 . 1309
Enothera bifrons . . .	17 . 1405	Pereskia Bleo . . .	17 . 1473
Enothera biennis, var. <i>grandiflora</i>	19 . 1504	Perilomia ocymoides . . .	17 . 1394
		Pernetia mucronata . . .	20 . 1675
		Persea gratissima . . .	15 . 1258

GENERAL INDEX TO THE NEW SERIES.

	<i>Vol. Folium</i>		<i>Vol. Folium</i>
Petunia violacea	19 . 1626	Rhododendron Alta-clerense	17 . 1414
Phacelia tanacetifolia	20 . 1696	Rhododendron arboreum, var.	
Pharium fistulosum	18 . 1546	<i>roseum</i>	15 . 1240
Platystemon Californicum	20 . 1679	Rhododendron arboreum, var.	
Phlomis floccosa	15 . 1300	<i>album</i>	20 . 1684
Phlox speciosa	16 . 1351	Rhododendron, Cartons	17 . 1449
Pholidota imbricata	14 . 1213	Rhododendron pulcherrimum	21 . 1820
Pholidota imbricata	21 . 1777	Ribes cereum	15 . 1263
Phycella Herbertiana	16 . 1341	Ribes divaricatum	16 . 1359
Physianthus albens	21 . 1759	Ribes inebrians	17 . 1471
Pimelea humilis	15 . 1268	Ribes niveum	20 . 1692
Pimelea hispida	19 . 1578	Ribes punctatum	26 . 1658
Pimelea intermedia	17 . 1439	Ribes sanguineum	16 . 1349
Pimelea ligustrina	21 . 1827	Ribes setosum	15 . 1237
Pimelea sylvestris	19 . 1582	Ribes speciosum	18 . 1557
Pleurothallis Grobyi	21 . 1797	Ribes tenuiflorum	19 . 1574
Pleurothallis picta	21 . 1825	Rose Clare	17 . 1438
Pleurothallis prolifera	15 . 1298	Rosa multiflora, var. <i>platyphylla</i>	16 . 1372
Plumeria Lambertiana	16 . 1378	Rosa Ruga	16 . 1389
Podolobium trilobatum	16 . 1333	Rubus nutkanus	16 . 1368
Polemonium cœruleum, var. <i>pili-</i>		Rubus roridus	19 . 1607
<i>ferum</i>	15 . 1303	Rubus spectabilis	17 . 1424
Polemonium humile	15 . 1304	Ruellia Sabiniana	15 . 1238
Polygala oppositifolia, var. <i>major</i>	14 . 1146	Russellia juncea	21 . 1773
Polygonum injucundum	15 . 1250		
Portulaca Gilliesii	20 . 1672	Saccolabium papillosum	18 . 1552
Potentilla arguta	16 . 1379	Sagittaria angustifolia	14 . 1141
Potentilla glandulosa	19 . 1583	Salpiglossis atropurpurea	18 . 1518
Potentilla Hopwoodiana	16 . 1387	Salvia angustifolia	18 . 1554
Potentilla laciniata	18 . 1478	Salvia foliosa	17 . 1429
Potentilla missourica	17 . 1412	Salvia fulgens	16 . 1356
Potentilla Russell's	18 . 1496	Salvia Grahmi	16 . 1370
Potentilla viscosa	18 . 1492	Salvia involucrata	14 . 1205
Pothos scandens	16 . 1337	Sarcanthus guttatus	17 . 1443
Pratia begonifolia	16 . 1373	Sauroglossum elatum	19 . 1618
Prunus candicans	14 . 1135	Schizanthus pinnatus, var. <i>humilis</i>	18 . 1562
Prunus dasycarpa	15 . 1243	Schizanthus retusus	18 . 1544
Prunus japonica	21 . 1801	Scilla plumbea	16 . 1355
Psoralea macrostachya	21 . 1769	Scottia angustifolia	15 . 1266
Pultenæa flexilis	20 . 1694	Scottia dentata	15 . 1233
Pultenæa rosmarinifolia	19 . 1584	Scottia lævis	19 . 1652
Pultenæa subumbellata	19 . 1632	Scutellaria alpina	18 . 1460
Purshia tridentata	17 . 1446	Scutellaria alpina, var. <i>lupulina</i>	18 . 1493
Pyrolirion aureum	20 . 1724	Sedum Cepæa	16 . 1391
Pyrus angustifolia	14 . 1207	Selago Gillii	18 . 1504
Pyrus crenata	20 . 1655	Sempervivum villosum	18 . 1553
Pyrus Bollwylleriana	17 . 1437	Sempervivum urbicum	20 . 1741
Pyrus grandifolia	14 . 1154	Senecio lilacinus	16 . 1342
Pyrus nivalis	17 . 1434	Senecio Tussilaginis	18 . 1550
Pyrus salvifolia	18 . 1482	Serapias cordigera, var. <i>longipetala</i>	14 . 1189
Pyrus sinensis	15 . 1248	Silene laciniata	17 . 1444
Pyrus spuria	14 . 1196	Sinningia villosa	14 . 1134
		Sisyrinchium grandiflorum	13 . 1364
Ranunculus creticus, var. <i>macro-</i>		Sisyrinchium odoratissimum	15 . 1283
<i>phyllus</i>	17 . 1432	Solandra guttata	18 . 1551
Raphiolepis rubra	17 . 1400	Solanum crispum	18 . 1516
Reevesia thyrsoidea	15 . 1236	Solanum etuberosum	20 . 1712
Renanthera coccinea	14 . 1131	Sollya heterophylla	17 . 1466
Rhodanthe Manglesii	20 . 1703	Sophora velutina	14 . 1185
Rhodochiton volubile	21 . 1755	Soulangia rubra	18 . 1498

GENERAL INDEX TO THE NEW SERIES.

	<i>Vol. Folium</i>		<i>Vol. Folium</i>
Sparaxis pendula . . .	16 . 1360	Trachymene cœrulea . . .	15 . 1225
Spermadietyon azureum . . .	15 . 1235	Tradescantia undata . . .	17 . 1403
Sphacele campanulata . . .	16 . 1382	Trichopetalum gracile . . .	18 . 1535
Sphærostema propinquum . . .	20 . 1688	Trifolium vesiculosum . . .	17 . 1408
Sphenotoma capitatum . . .	18 . 1515	Triteleia laxa	20 . 1685
Spiræa ariæfolia	16 . 1365	Tritoma Burchelli	21 . 1745
Spiræa chamædrifolia	15 . 1222	Tropæolum pentaphyllum . . .	18 . 1547
Stachys albicaulis	18 . 1558	Tulipa Oculus solis, var. <i>persica</i>	14 . 1143
Stachys germanica, var. <i>pubescens</i>	15 . 1289	Tulipa Oculus solis, var. <i>præcox</i>	17 . 1419
Stachys inflata	20 . 1697	Tupistra nutans	15 . 1223
Stachys Salviæ	15 . 1226	Turraea pinnata	17 . 1413
Stanhopea eburnea	18 . 1529		
Stanhopea oculata	21 . 1800	Ulex genistoides	17 . 1452
Stapelia Gussoneana	20 . 1731		
Statice puberula	17 . 1450	Vaccinium ovatum	16 . 1354
Stemodia chilensis	17 . 1470	Vanda teres	21 . 1809
Stenactis speciosa	19 . 1577	Verbena Melindres	14 . 1184
Sterculia lanceolata	15 . 1256	Verbena multifida contracta . . .	21 . 1766
Sterculia Tragacantha	16 . 1353	Verbena sulphurea	21 . 1748
Stigmaphyllon aristatum	20 . 1659	Vernonia axilliflora	17 . 1464
Streptocarpus Rexii	14 . 1173	Viburnum cotinifolium	19 . 1650
Stylidium fasciculatum	17 . 1459	Villarsia reniformis	18 . 1533
Syringa Josikæa	20 . 1733	Viola præmorsa	15 . 1254
Tabernæmontana densiflora	15 . 1273	Westringia longifolia	18 . 1481
Tacsonia pinnatistipula	18 . 1536		
Talauma Candollii	20 . 1709	Xerophyllum setifolium	19 . 1613
Tellima grandiflora	14 . 1178		
Teucrium orchideum	15 . 1255	Yucca superba	20 . 1690
Thermopsis fabacea	15 . 1272		
Thryallis brachystachys	14 . 1162	Zephyranthes mesochloa	16 . 1361
Tillandsia acaulis	14 . 1157	Zephyranthes Spofforthiana . . .	21 . 1746
Tillandsia rosea	16 . 1357	Zinnia violacea, var. <i>coccinea</i>	15 . 1294
Tillandsia stricta	16 . 1338		

THE END.

LONDON:

NORMAN AND SKEEN, PRINTERS, MAIDEN LANE, COVENT GARDEN.



Miss Drake. del.

Pub. by J. Ridgway 169 Piccadilly March. 1. 1835.

J. Walto. sc.

* CYCNOCHES Loddigesii.

Loddiges' Swanwort.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆE § VANDEÆ, (*Introduction to the Natural System of Botany*, p. 262.)

CYCNOCHES, Lindl. *Perianthium* explanatum. *Sepala* lateralia lanceolata, basi paululùm sub labello connata; supremo angustiore. *Petala* latiora, falcata, decurva. *Labellum* liberum, ecalcaratum, columnâ continuum, lanceolatum, integerrimum, ungue abrupto calloso. *Columna* elongata, arcuata, teres, apice clavata, auriculis 2 falcatis ad latera clinandrii. *Anthera* bilocularis. *Pollinia* 2, postice sulcata, subpedicellata; *caudiculâ* lineari; *glandulâ* grossâ.—*Habitus Cataseti* (sed racemus lateralis). Lindl. *Gen. & Sp. Orch.* p. 154.

Cycnoches Loddigesii, Lindl. *l. c. Bot. Cab. t.* 2000.

Caulis erectus, carnosus, cylindræus, subpedalis, pallidè viridis, corrugatus, basi vaginis membranaceis, apice foliis vestitus. Folia omninò Cataseti cujusdam. Racemus multiflorus, pendulus, e latere caulis erumpens, flexuosus. Flores maximi ordinis, initio inodori, mox Vanillam gratissimè spirantes, præsertim manè. Bractææ ovatæ, cucullatæ, obtusæ, pedicellis breviores. *Sepala* viridia apice fusca, et obscure maculata, mox rufescentes; supremum lineari-oblongum, arcuatum 3-pollicare; lateralia 2-poll. longa imâ basi connata, ovato-oblonga, subundulata, pendula. *Petala* ejusdem coloris cum sepalis, sed minùs conspicuè maculata, oblongo-lanceolata, inæquilatera, decurva, 2¼-poll. longa. *Labellum* patentissimum, cum columnâ continuum, angustè oblongum, convexum, carnosum, medio album, apicem versus tenuius, sordidè lutescens, undique maculis latis inæqualibus sanguineis notatum; ungue brevi alato maculoso, alis lutescentibus, disco convexo albo. *Columna* gracillima, arcuata, utrinque clavata; dimidio inferiore atropurpureo, superiore complanato virescente purpureo maculoso. *Anthera* citò devida, membranacea, bilocularis, ferè pellucida, omninò mutica. *Clinandrium* posticè bicornè, cornubus falcatis compressis supra antheram incurvis.

For the discovery of this most extraordinary Epiphyte the public is indebted to John Henry Lance, Esq. who discovered it, and a great number of other new plants in the woods

* So called from κύκνος a swan, and αὐχὴν the neck; in allusion to the column of this plant, which is gracefully curved, like the neck of the swan.

of Surinam. It was originally sent by that gentleman to Messrs. Loddiges, with whom it flowered imperfectly three or four years ago. Other and more healthy plants were afterwards presented by Mr. Lance to the Horticultural Society, the produce of one of which was the singular raceme now represented.

In size the flowers of this plant are the largest that have yet been met with among Orchideous plants, measuring not less than five inches from the tip of the back sepal, to the point of the lip. They are more singular than beautiful, but they compensate in fragrance for their want of brilliant colours: the purest odour of Vanilla is exhaled by them when they have been open a short time, especially in the morning.

This species flowers in the months of June and July. It resembles a *Catasetum* in habit, but its flowers originate from the side, and not the base, of the fleshy stem. It grows very freely under the same management as *Catasetums*, but seems to require the greatest heat that is ever obtained in a damp stove. It is disposed to increase itself pretty freely, by multiplication of its fleshy stems, and will probably cease, after a few years, to be so rare a plant as it now is.



* CALCEOLÁRIA angustiflora.

Narrow-flowered Calceolaria.

DIANDRIA MONOGYNIA.

*Nat. ord. SCROPHULARINEÆ Juss. (Introduction to the Natural System of Botany, p. 228.)**CALCEOLARIA.—Supra, vol. 9. p. 723.*

C. angustiflora; caule perenni diffuso, foliis cordato-ovatis duplicato-serratis glabris oppositis ternatisque, pedunculis axillaribus paucifloris sæpius paniculam angustissimam foliosam efficientibus, corollæ labio inferiore obtuso basi angustato superiore minimo.

C. angustiflora. Ruiz et Pavon. Fl. Peruv. vol. 1. p. 17. t. 28. f. a. Graham in Bot. Mag. t. 3094.

? *C. verticillata. Hooker Bot. Misc. vol. 2. 233.*

A half hardy perennial, flowering in August and September, and remarkable among its race for its prostrate habit, deep green, somewhat shining leaves, and small flowers. It was found first by the authors of the Flora Peruviana, in the valley of Canta in Peru; subsequently Mr. Cruckshanks discovered it at Culluay in the same district, and gathered the seeds from which the plants in our gardens were obtained.

It is a species of no great attraction, but deserves to be recorded in this work, as one of the genuine wild forms of a genus, which, however beautiful and interesting, has already begun to sink in estimation, in consequence of the ruin that has been brought upon it by the unskilfulness of gardeners. In their haste to improve the works of nature, these gentlemen have converted some of the fairest races in the Vegetable world, into forms in no case more beautiful than the original, and in the majority of instances unhealthy, mongrel, and

* See fol. 1214.

debased. We strongly recommend all those who value this really beautiful and most singular genus, to abandon a pursuit which has as yet led to few results of which good taste can approve, and to apply the same skill which they have used in spoiling Calceolarias to recovering the pure original races, to preserving them uncontaminated, and to increasing their native charms ; not by unnatural combinations, but by those well known methods by which the purity of a species may be maintained while its vigour, health and beauty are augmented.

At least, if the genus must be the subject of hybridizing, let the intermixture be made with some reasonable attention to the only rules by which it is possible to arrive at a really desirable result.



* **INDIGÓFERA atropurpúrea.***Purple-flowered Indigo plant.*

DIADELPHIA DECANDRIA.

*Nat. ord. LEGUMINOSÆ Juss. (Introduction to the Natural System of Botany, p. 86.)**INDIGOFERA.—Supra, vol. 5. fol. 386.*

I. atropurpurea; fruticosa, erecta, foliis 5-6-jugis, foliolis oblongis obtusissimis apiculatis tenuibus glabris, racemis multifloris foliis æqualibus v. longioribus, leguminibus linearibus rectis acustriatis 8-9-spermis.

I. atropurpurea *Hamilt. in Roxb. Fl. Ind. vol. 3. 380. Wall. Cat. No. 5463.*

A native of Nepal, where it was originally found by Hamilton; probably in the hot valleys; for it does not succeed well, unless cultivated in the stove. There it becomes a handsome light green bush, richly ornamented by its numerous bunches of purple and crimson flowers. If planted in the open air, it languishes even on a south wall and in such a season as that of 1834; producing its blossoms in small quantities and imperfectly.

Our specimens were communicated in August, 1833, from the hothouse of James Bateman, Esq. of Knypersley.

Roxburgh says, that when raised in the Botanic Garden at Calcutta, it grew from 3 to 5 feet high in about 8 months.

* Literally *Indigo-bearer*; some species of this genus producing the dye of that name.



Miss Drake. del.

Pub by Ridgway 169 Picadilly March. 1. 1835.

S. Watts. sc.

* TRITOMA Burchéllii.

Burchell's Tritoma.

HEXANDRIA MONOGYNIA.

Nat. ord. ASPHODELEÆ Juss. (Introduction to the Natural System of Botany, p. 273.)

TRITOMA Ker. Link. *Perianthium* tubulosum, eplicatum; limbo brevi 6-dentato regulari. *Stamina* hypogyna, recta, libera, exserta, alterna longiora. *Stigma* simplex. *Capsula* cartilaginea, rigescens, ovata, obtusè trigona. *Semina* biserialia, plurima, triquetra v. angulata, sibi invicem appressa.—*Folia* graminea canaliculata. *Caulis nullus.*

T. Burchellii; foliis lætè viridibus margine* lævibus, racemo oblongo crasso denso, perianthiis clavato-cylindræis unicoloribus.

Tritoma Burchellii. *Herbert. in Sweet's Hort. Brit.*

Racemus 4 poll. longus, densissimus, et vix angustior, nullo modo pyramidatus v. protractus. Flores juniores sanguinei, postèd lutei v. potius luteo-aurantiaci unicolores, 1¼-poll. longi; limbo 4 lin. tubo apice 3 lin. lato.

Communicated by the Honourable and Rev. William Herbert in July last. It was originally introduced from the Cape of Good Hope by Mr. Burchell, after whom it has been named.

It is a beautiful perennial plant, and quite hardy. At Spofforth it always flowers before or soon after Midsummer; all the other species of *Tritoma* are late flowerers.

Increased by the offsets, which it throws up in plenty.

* So named from τρεις three, and τεμνω to cut, because of the leaves ending in three sharp angles. As there was a genus of insects already called *Tritoma*, when this genus of plants was first proposed, Professor Link would change its name to *Tritomanthe*; but there is no end to alterations of this sort.



Herbert. del.

Pub. by J. Ridgway, 169, Piccadilly, March, 1. 1835.

J. Watts. sc.

* ZEPHYRÁNTHES Spofforthiána.

Spofforth Zephyranthes.

HEXANDRIA MONOGYNIA.

Nat. ord. AMARYLLIDÆ, R. Br. (Introduction to the Natural System of Botany, p. 259.)

GARDEN VARIETY.

ZEPHYRANTHES SPOFFORTHIANA. Herbert MSS.

“Scape yellowish green, $5\frac{3}{4}$ inches high. Spathe brownish green. Peduncles green, an inch long. Germ deep green. Tube green. Limb rose-coloured, lined with white in the form of a star. Style very much declined, white and much longer than the filaments. Leaves not $\frac{1}{4}$ of an inch wide, acute, keeled, deep green. Pollen abundant, and anthers quite perfect and apparently fertile.”

“The pretty flower which is the subject of this article, is an hybrid production from the tropical *Z. tubispatha*, which has white flowers, fertilized by the pollen of the Mexican *Z. carinata* with large red flowers. It is in every respect intermediate, having the leaves wider than those of *Z. tubispatha*, and distinguishable by the keel, which is conspicuous on the back of those of the male parent. The flower is also of intermediate dimensions and colour, resembling the male parent most both in form and hue, and in the posture of the

* See folio 1361.

style and filaments. A single bulb of this intermixture was obtained at Spofforth, which has produced a few offsets, piercing the ground, like those of the male parent, at a little distance from the principal bulb, their lateral direction giving the plant a stoloniferous appearance. Its anthers are perfectly formed, and the pollen seems to have all the requisites for fertility. Standing in a cool part of the stove, beside *Z. tubispatha*, it flowers before it, and probably will succeed like *Z. carinata* in a lower temperature. It must be observed that *Z. carinata*, Bot. Mag. the male parent of this plant, is, with respect to the flower only, the plant represented in the Bot. Reg. 902. under the name of *Z. grandiflora*, though the flower in the figure is rather longer and paler than its usual habit; but in consequence of the flower having been produced by a newly imported bulb, before the leaves had shot, and a subsequent mistake of the gardeners, the sketch of the leaves, and the description of the leaves and seeds of *Z. grandiflora* were made erroneously from a pinkish variety of the small flowering *Z. striata*.

“It will have been observed that *Habranthus phycelloides* of this work approaches so near in its general aspect to the genus *Phycella*, as to make it questionable whether the genus *Phycella* should be maintained distinct from *Habranthus*, at least from the many-flowered, non-expanding portion of that genus; and that the one-flowered *Habranthi*, which expand in the sun, are not easily distinguishable from the genus *Zephyranthes*; and it may be found that such species, viz. *H. versicolor*, *robustus*, and *Andersoni*, should be removed to the genus *Zephyranthes*; but an accurate revision of the various species should precede any alteration. It has, however, been as yet found impossible to obtain a mule between any *Habranthus* and *Zephyranthes*, which gives some reason to believe that the division as it stands is correct. Several flowers of *Z. striata* at Spofforth have lately been fertilized by *Z. carinata*, and others in the same pots by *H. Andersoni*, all the former appear to be forming seed, and all the latter have died away.”

For the foregoing interesting memorandum we are indebted to the Honourable and Rev. William Herbert. We have for some time been aware of the error to which he alludes, and are happy to have the present opportunity of making it known. It will be necessary to expunge the name of *Z. grandiflora* from the catalogues, as it is a non-entity.



* GARDOQUÍA Hookéri.

Scarlet Gardoquia.

DIDYNAMIA GYMNOSPERMIA.

Nat. ord. LABIATÆ, Juss. (Introduction to the Natural System of Botany, p. 239.)

GARDOQUIA Fl. Per. Calyx tubulosus, 13-nerviis, subincurvus, ore æquali vel obliquo, dentibus brevibus rectis subæqualibus vel subbilabiatis. *Corollæ tubus* longè exsertus, rectus vel incurvus, intùs nudus; limbus bilabiatus, labium superius erectum subplanum emarginatum, inferius subpatens, lobis planis medio latiore. *Stamina* 4, subdidynama, inferioribus longioribus, laxè adscendentia, apice subdistantia, superiora subinde sterilia. *Filamenta* edentula. *Antheræ* biloculares, loculis distinctis parallelis vel subdivergentibus. *Styli* lobi subæquales. *Achenia* sicca, lævia.—Suffrutices fruticesve *ramosissimi, foliosi, sæpe procumbentes*. Flores *pulchri, sæpiùs coccinei*. Bentham Gen. et Sp. Lab. p. 397.

G. Hookeri; fruticosa, glaberrima, foliis obovatis subintegerrimis basi in petiolum brevem angustatis utrinque viridibus, pedunculis 2-3-floris, calycis glabri subbilabiati labio superiore breve bidentato erecto, fauce intùs villis clausâ. *Bentham l. c. 401.*

Cunila coccinea, Nutt. ex Hooker exot. fl. 3. t. 163.

Melissa coccinea, Spreng. syst. 2. 224.

Gardoquia Hookeri, Don in Sweet. Fl. Gard. ser. 2. t. 271.

A very pretty half hardy and half shrubby plant, native of Florida according to Nuttall, as quoted by Dr. Hooker, and of South Carolina according to Mr. Don.

Our specimens were furnished by Mr. Marshall, gardener to Mrs. Langley of Thames Ditton, in Oct. 1834. It is a neat, and very desirable plant, easily propagated by cuttings.

* Don Diego Gardoqui, Minister of Finance under Charles IV. of Spain, received the compliment of having this beautiful genus named after him, because of his having favoured the South American Expedition of Ruiz and Pavon.



Miss Drake del.

Pub by F. Ridgway

169 Piccadilly

March 1. 1835.

S. Watts sc.

* VERBÉNA sulphúrea.

Sulphur-coloured Vervain.

DIDYNAMIA ANGIOSPERMIA.

Nat. ord. VERBENACEÆ. Juss. (Introduction to the Natural System of Botany, p. 238.)

VERBENA.—Supra, vol. 4. fol. 294.

V. sulphurea; diffusa, pilosa, foliis 3-5-partitis: laciniis simplicibus lobatisque linearibus obtusis inferioribus ab imâ basi provenientiibus, floribus subcapitatis, calycibus hispide pilosis, corollæ laciniis patentibus emarginatis.


V. sulphurea. Don in Sweet's Flower Garden, ser. 2. t. 221.

A pretty hardy perennial, with hairy stems spreading flat upon the ground, and heads of sulphur-coloured flowers.

Although not so conspicuous for its beauty as *Verbena chamædrifolia* (*V. Melindres* of this work, *t.* 1184.) it is nevertheless a neat pretty species, which deserves a place in all select collections.

It is a native of Chile, flowering in August, September, and November. Cuming found it near Valparaiso (No. 519); Bridges on the mountains near the same city. It is very nearly the same as *V. erinoides*, from which it differs in having yellow instead of purple flowers, and more particularly in the lower lobes of its leaves originating from the very base of the leaf, and not from near the point.

* See fol. 1184.



Miss Drake del.

Pub by J. Ridgway 169 Piccadilly Sp. 1. 1825.

2000-2000

* DRACÆNA terminalis.

The Sandwich Island Tee-Plant.

HEXANDRIA MONOGYNIA.

*Nat. ord. ASPHODELEÆ, Juss. (Introduction to the Natural System of Botany, p. 271.)**DRACÆNA.—Supra, vol. 12. fol. 956.*

D. terminalis; caule fruticoso vel arborescente, foliis petiolatis lanceolatis utrinque attenuatis, paniculæ ramis divaricatis simplicibus ramosisve floribus subsessilibus. *Römer & Schultes, Sp. Pl. 7. 343.*

D. terminalis. *Willd. Sp. Pl. 2. 157.*

Asparagus terminalis. *Linn. Sp. Pl. 450.*

One of the most graceful of arborescent stove-plants, where there is sufficient height for it to rear its slender stem to the elevation of ten or twelve feet. In appearance it resembles a Palm, and although, when its structure is carefully examined, it is found not to be one of that princely tribe, yet there can be no doubt that it serves to connect the chain of vegetation, by bringing the *Asparagus* in contact with the Cabbage Palm. Linnæus took it for a species of *Asparagus*.

In place of again describing a plant which has repeatedly been already in the hands of Botanists, we shall confine ourselves on this occasion to an extract from the entertaining work of Ellis on the Sandwich Islands.

“The *Ti* plant is common in all the South Sea Islands. It is a slow-growing plant, with a large, woody, fusiform

* See folio 1169.

root, which, when first dug out of the ground, is hard and fibrous, almost tasteless, and of a white or light yellow colour. The natives bake it in large ovens underground, in the same manner as they dress the *arum* and other roots. After baking it appears like a different substance altogether, being of a yellowish brown colour, soft, though fibrous, and saturated with a highly saccharine juice. It is sweet and pleasant to the taste, and much of it is eaten in this state, but the greater part is employed in making an intoxicating liquor much used by the natives. They bruise the baked roots with a stone, and steep them with water in a barrel, or the bottom of an old canoe, till the mass is in a state of fermentation. The liquor is then drawn off, and sometimes distilled, when it produces a strong spirit; but the greater part of it is drunk in its fermented state without any further preparation. The root is certainly capable of being used for many valuable purposes. A good beer may be made from it; and in the Society Islands, though never able to granulate it, we have frequently boiled its juice to a thick syrup, and used it as a substitute for sugar when destitute of that article.

“ We should think it an excellent antiscorbutic, and as such useful to ships in long voyages. Captains visiting the Society Islands frequently procure large quantities of it to make beer with during their voyage, as it will keep good six weeks or two months after it is baked. On my return in the American ship *Russell*, Captain Coleman, we procured a quantity that had been baked at Rurutu, near the Society Islands, and brought it round Cape Horn. It lasted five or six weeks, and would probably have kept longer, as the only change we perceived during that time was a slight degree of acidity in the taste. Cattle, sheep, and goats are fond of the leaves; and as they contain more nutriment than any other indigenous vegetable, and may be kept on board ships several weeks, they are certainly the best provender that can be procured in the Islands for stock taken to sea. It is not so plentiful in the Sandwich Islands as it was before the natives used it for the above purpose, but in some of the

other Islands of the Pacific it is abundant, and may be easily procured.

“Other parts of the *dracæna* are also useful. The natives frequently plant the roots thickly around their enclosures, interweave the stems of the plant, and form a valuable permanent hedge. The branch was always an emblem of peace, and in times of war, borne, together with a young plantain tree, as a flag of truce, by the messengers who passed between the hostile parties. The leaves, wove together by their stalks, formed a short cloak, which the natives wore in their mountainous journeys; they also make the most durable thatch for the sides and roofs of their best houses, are employed in constructing their tents in war, and their temporary abodes during their inland excursions.”

The specimen from which our drawing was taken was furnished by Mr. Lambert, in March 1834.

It is propagated either by seeds or by truncheons of its stem, which when cut down will throw up suckers from its base, just as the *Asparagus*, when its early shoots are destroyed in the spring, will continue to replace them by fresh bourgeons from its bottom.



* *BÉRBERIS dealbata.**Whitened Barberry.*

HEXANDRIA MONOGYNIA.

Nat. ord. BERBERIDÆ, Juss. (Introduction to the Natural System of Botany, p. 30.)

BERBERIS.—Supra, vol. 12. fol. 1176.

§ *Folia simplicia, sempervirentia (coriacea). Flores racemosi.*

B. dealbata; subinermis, foliis simplicibus coriaceis suborbiculatis spinoso-dentatis subtus dealbatis, racemis oblongis densissimis pedunculatis.

Frutex in horto 4-5-pedalis, verosimiliter orgyalis ad minimum, ramis purpureo-fuscis inermibus parcè foliatis. Folia coriacea, subrotundo-ovata, convexa, glaucescentia, spinoso-dentata, aliquandò subquadrata, imò cuneata 3-dentata, subtus albedine densâ corticata; petiolis brevibus basi articulatis. Flores in racemos oblongos multifloros densissimos nutantes congesti, lutei; sepalis apice sanguinolentis. Petala intus biglandulosa.

A native of Mexico whence it was obtained by the Horticultural Society. Our drawing was made in the Chiswick Garden in April last.

This remarkable species is an evergreen shrub, which is probably hardy; but of this we are not certain, in consequence of its having been always hitherto protected in front of an east wall in winter. It differs in a striking manner from all other Barberries in several particulars. In the first place its leaves are very round and convex, and covered with a sort of bloom; then they are very white on the under surface, not however from the presence of any mealiness or white hairs, but on account of a discoloration of the

* See folio 1176.

cuticle; besides this there are scarcely any spines upon the branches; and finally its flowers are collected in remarkably dense nodding clusters, which are not much longer than the leaves. Its fruit is unknown.

Like the other species of its genus it is most conveniently increased by layers, which will root in the course of a single season.



Miss Drake. del.

Pub. by J. Ridgway 169 Piccadilly Sep. 1. 1835.

L. Watts. sc.

* *LÆLIA* *anceps*.*Two-edged Lælia.*

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆE § EPIDENDRÆE, Lindl. (Introduction to the Natural System of Botany, p. 262.)

LÆLIA, Lindl. Sepala explanata, lanceolata, æqualia. Petala majora paulò difformia. Labellum (posticum), 3-partitum, lamellatum, circa columnam convolutum. Columna aptera carnosâ, anticè canaliculata. Anthera 8-ocularis. Pollinia 8, caudiculis 4 elasticis.—Herbæ epiphytæ, rhizomate pseudobolbophoro. Scapi terminales, pauci v. multiflori. Flores speciosi, odorati. Gen. & Sp. Orch. p. 115.

L. anceps; foliis binis aut solitariis lanceolatis, scapo ancipiti bifloro squamis carinatis vestito, ovario viscoso, labelli disco lineari elevato apice trilobo, pseudobolbis ovatis distantibus tetraquetris.

Rhizoma repens, squamosum. Pseudobulbi ovati, distantes, tetraquetri, 2 poll. longi, juventute squamis acuminatis vestiti. Folia solitaria, rarò bina, lanceolata, acuminata, coriacea, atroviridia. Scapus ex apice pseudobulbi ortum ducens, sesquipedalis, gracilis, anceps, squamis carinatis arcè vaginantibus vestitus, apice biflorus. Ovarium viscosum, bracteam lato-lanceolatam, acuminatissimam, carinatam, membranaceam, fuscam, involvente brevius, apice teres, cuniculo a basi labelli orto brevi instructum. Sepala violacea, membranacea, dorso subherbacea, lanceolata, bipollicaria, patentissima. Petala oblongo-lanceolata, ejusdem coloris texturæ, et longitudinis, sed lineam mediam dorsi tantum herbacea, et duplò latiora. Labellum aut anticum, aut posticum, cucullatum, trilobum, basi columnam involvens et pallidè violaceum, intus luteum sanguineo-venosum; trilobum, lobis lateralibus rotundatis, intermedio atropurpureo basi albo, oblongo, acuto, subundulato, plano, lineam latam elevatam crassam luteam antrorsum trilobam in medio; lobis lateralibus abruptis, intermedio productione in laminam abruptam verticalem attenuato. Columna semiteres, marginata, clavata, cum labello continua. Anthera 8-ocularis. Pollinia 8, cuneata, utraq; extremitate caudicularum quatuor inserta et inflexa.

A very beautiful Orchideous plant, imported from Mexico by Messrs. Loddiges, in whose collection our drawing was made last December. We at first took it for *Lælia grandiflora*, but Mr. George Loddiges has satisfied us that it is

not that species, which it is, however, highly probable that he possesses in a plant from Xalapa, with pear-shaped pseudo-bulbs, each of which has two or three lanceolate purplish leaves. The pseudo-bulbs of this are remarkable for their ovate outline, and for their four-angled figure, produced by a sharp ridge being planted upon each face of a compressed body.

When we say that this plant is equal in beauty to any of the Cattleyas, that it has a far more graceful mode of growth, in consequence of the length of its slender scaly stems, from the point of which the flowers swing, and that it diffuses an agreeable fragrance, we shall have said that it is one of the most interesting of the tribe that has yet made its appearance in our stoves.

It will probably succeed, without difficulty, in any hot-house which is adapted for the cultivation of Maxillarias and plants of that description.



* **MONACHÁNTHUS** *viridis*.*Green-flowered Cowlwort.*

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆE § VANDEÆ, Juss. (Introduction to the Natural System of Botany, p. 262.)

MONACHANTHUS.—Supra, vol. 20. fol. 1735.

M. viridis; racemo multifloro, labello oblongo cuspidato margine lævi, sepalis petalisque rigidis ovatis.

M. viridis, *Lindl. Gen. & Sp. Orch. p. 157.*

Planta facie omnino *Cataseti tridentati* quo foliis et caule tantum convenit ut floribus adeptis eandem diceres. Flores carnosi, herbacei, labelli margine vitellino. Sepala ovata, acuta, rigida, petalis parallela. Petala conformia, sed latiora et maculis quibusdam purpureis obscuris notata. Labellum posticum ventricosum, carnosum, $1\frac{1}{2}$ -poll. longum, ore obliquo lævi cuspidato. Columna brevis, ovata, carnosissima, mutica.

This is the original species on which the genus *Monachanthus* was founded. In habit it is so like *Catasetum tridentatum* that we long doubted whether it ought to be generically separated. The absence of the long tendril-like processes, which are so conspicuous upon the column of *Catasetum*, was, however, so remarkable a circumstance, that we came to the conclusion that it must be an essentially distinct form; and the discovery of the curious *Monachanthus discolor*, figured at tab. 1735, has shewn that we judged correctly.

A native of Brazil, growing upon trees in the Corcovado, whence Dr. Hooker received the drawing and specimen which first made the genus known. Our figure was made from a specimen communicated last November from Went-

* See fol. 1735.

worth, by permission of Lord Fitzwilliam. We are not aware at what time, or by whom it was imported, but it has probably been taken for a green form of the common *Catasetum tridentatum*, and consequently no record has been kept of it.

It requires precisely the same treatment as *Catasetums*, and may be cultivated with facility by any one who has a damp stove.



Miss Drake del.

Pub by J. Ridgway 169 Piccadilly Sp. 1. 1835.

J. Watts sc.

* *ARBUTUS procera*.*Tall Arbutus, or Strawberry Tree.*

DECANDRIA MONOGYNIA.

*Nat. ord. ERICACEÆ, Juss. (Introduction to the Natural System of Botany, p. 182.)**ARBUTUS.—Supra, vol. 2. fol. 113.**A. procera*; foliis oblongis serratis serrulatis integrisque glabris, petiolis calvis, racemis terminalibus paniculatis secundis.*A. procera, Douglas Herb.*

Rami juvenes glaucescentes, adulti rufo-castanei, glabri, salvis vegetissimis qui hispidi sunt; epidermide firmâ non deglubente. Folia in ramis surculosis oblonga, utrinque angustata, duplicato-serrata, petiolis hispidis; ramulorum oblonga, nunc cordata, integra vel irregulariter serrulata; petiolis ferè unciam longis. Racemi tomentosi, paniculati, terminales, secundi; bracteis ovatis, concavis, glabris. Corollæ albæ, leviter herbaceæ, ovatae, medio paululùm constrictæ, basi obtusæ et obscurè decem-gibbosæ; limbo revoluto. Fructus juniores subrotundi, leviter obovati, tuberculati more *Unedonis*.

A small tree found by the late unfortunate Douglas in the mountainous woody parts of the North West Coast of North America, and introduced by the Horticultural Society in 1825.

In the Gardens it forms an evergreen bush, with a fine broad glossy foliage, and a very vigorous appearance; but

* The *Arbutus* of Virgil and Horace was undoubtedly the modern Strawberry Tree. De Théis says the word has been formed from the Celtic *ar* rough, and *boise* a bush, in consequence of the astringency of its fruit. He considers that the term *horrida*, applied by Virgil to his *Arbutus*, has equally, in this case, the signification of austere.

it requires to be carefully protected from extreme cold in winter, and succeeds best against a west wall. Its flowers are of a delicate greenish white, and are arranged after the manner of those of *Arbutus Andrachne*, to which species it is most nearly allied; differing however altogether in the form and serratures of its leaves, and in the form and size of its flowers.

The unripe fruits appear from our dried specimens to be like those of the common *Arbutus*, only obovate in a slight degree; the ripe fruit is unknown. The strong root-shoots are covered with scattered bristles, as also are the leaf-stalks, and the leaves themselves on such shoots are very strongly serrated.

Our drawing was made from a specimen obligingly supplied in May last, by Mr. Osborn, the present proprietor of the celebrated Fulham Nursery, lately occupied by Messrs. Whitley and Co.



Miss Drake del.

Pub by J. Ridgway 769 Piccadilly Apr. 1. 1835

J. Walpole sc.

* BRÁSSIA Lanceána.

Mr. Lance's Brassia.

GYNANDRIA MONANDRIA.

*Nat. ord. ORCHIDÆE § VANDEÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.)**BRASSIA.—Supra, vol. 10. fol. 832.*

B. *Lanceana*; sepalis ovato-lanceolatis acuminatis, petalis minoribus, labello oblongo acuminato undulato sepalis lateralibus duplò brevioribus.

Color totius herbæ amœnè et diffuso viridis. Pseudobulbi densi, oblongi, macri, compressissimi, ancipites, longitudinaliter sulcati, haud rarò arcuati. Folia bina aut solitaria, oblongo-lanceolata, macra, valdè striata. Racemi radicales, foliis longiores, floribus magnis luteis secundis odorem *Primulæ suavissimum* spirantibus onusti. Bracteæ ovatæ, concavæ, breves, membranaceæ. Sepala patentissima, lineari-lanceolata, purpureo paululum maculata, basi virescentia; supremum $1\frac{1}{2}$, lateralia 2 poll. longa. Petala ejusdem formæ et coloris sed sepalo supremo duplo minora. Labellum luteum, immaculatum, oblongum, undulatum, ferè repandum, acuminatum, sepalis lateralibus duplò brevius; basi tuberculis 2 albis oblongis contiguis parallelis inanibus pubescentibus, dentibusque totidem membranaceis in fronte.

A native of Surinam, in woods, where it was found growing upon trees by John Henry Lance, Esq.; and by him presented to the Horticultural Society in 1833. It is also wild in Brazil, where it was found by Dr. von Martius upon the trees on the banks of the river Jui, one of the tributaries

* Thus named by Mr. Brown in due commemoration of the late Mr. Brass, a skilful botanical traveller and draughtsman, who collected seeds, plants, and dried specimens on the Guinea Coast for Sir Joseph Banks, Dr. Fothergill and Dr. Pitcairn, and whose sketches being most liberally lent by Sir Joseph Banks to Dr. Afzelius in his visit to Sierra Leone, were maliciously damaged, and partly destroyed out of characteristic and wanton brutality by some piratical slave-mongers under the French flag, during the late war.—*Smith in Rees' Cyclopædia.*

of the Japurà in the Province of Rio Negro, flowering in February. It blossomed at nearly the same time in the Chiswick Garden, and with Messrs. Loddiges, in whose collection our drawing was made last August.

It is of course, like other Surinam plants, a very tender species, and requires the hottest and dampest part of a stove; but there is no Orchideous plant which is more easy to cultivate, or more ready to multiply. It thrives in decayed leaf-mould, better than in any other compost, and may be readily known when out of flower, by the bright light green colour of its leaves, and by its peculiarly thin pseudobulbs, which often curve down upon themselves, as is represented in our figure.

If this is not equal to *Lælia* in the size of its flowers, and the brilliancy of its colours, it far surpasses that plant in its exquisite fragrance, which is like nothing so much as newly gathered cowslips and primroses.



* RHODOCHÍTON volúbile.

Twining Red-Cloak.

DIDYNAMIA ANGIOSPERMIA.

Nat. ord. SCROPHULARINEÆ, Juss. § ANTIRRHINEÆ (*Introduction to the Natural System of Botany*, p. 273.)

RHODOCHITON Zuccarini. Calyx membranaceus, coloratus, campanulatus, 5-fidus. Corolla: tubus anguloso-clavatus, internè pilis simplicibus reflexis, basi ubique, faucem versus 5-fariam vestitus; limbus 5-partitus, segmenta subæqualia erecta. Stamina didynama, rudimento quinti, erecta, apicibus simplicibus. Stylus sub stigmate rectus *Graham in Bot. Mag. t. 3367.* (Capsula vestita, pergamenea, bilocularis, polysperma, irregulariter dehiscens. Semina alata, radiatim striata; ex Zuccarini.)

Rhodochiton volubile; Zuccarini in litt. 1829, Otto et Dietr. in verhandl. ver. gart. Preuss. 10. 152. t. 1. Graham *l. c.*

Lophospermum atosanguineum. Zuccarini *Plant. nov. et min. cogn. fasc. 1. in Abhandl. Math. Phys. Cl. Monac. vol. 1. p. 306. t. 13.*

Lophospermum Rhodochiton. D. Don in *Sweet's Brit. Fl. Gard. t. 250.*

A beautiful little climbing herbaceous plant, which was introduced late in 1833, by Mr. Low of Clapton, from the Berlin Garden. It is not very usual for us to be behindhand with our contemporaries in the publication of new plants, which are of much Horticultural interest; but on the present occasion we find ourselves forestalled in every direction. We console ourselves with the reflection that we are on that very account enabled to give a more complete history of the plant than would have been possible at an earlier period.

It was originally raised in the Botanic Garden at Munich from seeds collected in Mexico by Count Karwinski, and in the year 1829 was distributed to other gardens under the name of *Rhodochiton volubile*, an appellation which had been given it by Professor Zuccarini.

An account of it was soon after published in the Transactions of the Prussian Horticultural Society, by Messrs. Otto and Dietrich, with a tolerable figure, and the following account of the manner in which it had been treated in the Berlin Garden.

* From *ῥοδός* red, and *χιτῶν* a cloak, in allusion to the colour of its calyx.

“ It is a climber, and may be applied to cover the walls in conservatories, as well as in the open ground during summer. It presents a really surprising aspect, when the outer walls of a greenhouse are covered with a screen formed by its numerous branches, tinged with red, and loaded with the most beautiful purplish-black flowers hanging down from the red calyxes. Out of every axil, to the most remote part of the twigs, a flower takes its origin, hanging down from a long stalk, which like the underside of the leaves and branches is coloured red. There can be no doubt, that it is, of all the known climbing plants with which we cover walls during summer, the most remarkable and beautiful, and ought to be particularly recommended, as it is easily cultivated, and flowers so very freely. Its propagation is by seeds and cuttings. The seeds are sown in March and April, that the young plants may become strong enough in time for planting in the open ground, where they remain during the whole summer, until frost sets in; if these plants are to be preserved, they should be potted off and put in a conservatory, where they prosper in a temperature, from 43° to 48° *Fahr.* They are also easily propagated by cuttings.”

This mode of management appears to have been attended with great success, for when the plant was observed in the Berlin Garden in the autumn of 1833 by Dr. Henderson, it was in a much more thriving state than we have seen it in England.

Its next appearance in print was in the Transactions of the Mathematical and Physical Class (we believe) of the Munich Academy of Sciences, where it was published with an admirable description and figure by Professor Zuccarini; who, however, had altered his opinion of its belonging to a distinct genus, and called it *Lophospermum atrosanguineum*; this must have been in 1833, but our copy of the memoir to which we refer, bears no date.

Then Mr. Don, in August 1834, published it under the name of *Lophospermum Rhodochiton*, and he was followed in the succeeding December by Professor Graham, who adhered to Professor Zuccarini's first opinion, that the plant is distinct from *Lophospermum*, in which we entirely agree. In such groups as the Antirrhineous section of Scrophularineæ, less weighty reasons than the great coloured campanulate calyx, and salver-shaped corolla of *Rhodochiton*, as compared with the five-parted herbaceous calyx, and funnel-shaped corolla of *Lophospermum* are universally admitted as sufficient to justify the separation of such genera as *Antirrhinum*, *Linaria* and *Anarrhinum*, and we cannot conceive upon what principle the former are to be combined while the latter remain disunited.

In this country the *Rhodochiton* appears to require exactly the same treatment as *Lophospermum*; we saw it in great beauty at Mrs. Marryat's in a pot in the greenhouse last September, and it grew very well in the Garden of the Horticultural Society, trained to a pole in the open ground. Its greatest enemy seems to be bright sunlight.



* DENDRÓBIUM Pierárdi.

Mr. Pierard's Dendrobium.

GYNANDRIA MONANDRIA.

*Nat. ord. ORCHIDÆE § MALAXIDÆE, Lindl. (Introduction to the Natural System of Botany, p. 262.)**DENDROBIUM.—Supra, vol. 7. fol. 548.*

- D. Pierardi*; caulibus pendulis glabris, foliis ovato-lanceolatis acutis, floribus geminatis racemum spurium formantibus, sepalis acuminatis membranaceis, petalis sepalo supremo majoribus acuminatis, labello cucullato dilatato subtruncato pubescente ciliato. *Gen. et Sp. Orch. p. 79.*
- D. Pierardi.* *Roxb. Fl. Ind. 3. 482. Hooker Exot. Fl. t. 9.*

Many years since an Orchideous Epiphyte was sent from Chittagong with some other species to the Botanical Garden, Calcutta, when Dr. Roxburgh named one of them *Dendrobium Pierardi*, in compliment to the gentleman who discovered it. In his Flora Indica Dr. Roxburgh adds that it is also a native of various parts of the Delta of the Ganges, where it is generally found on Mangoe Trees.

No one has however been able to discover any plant which answers exactly to Dr. Roxburgh's description of his *D. Pierardi*; and what is now cultivated under that name in the Calcutta Garden appears from Dr. Wallich's specimens to be the plant defined by Dr. Brown in the seventh volume of the first series of this work under the name of *D. cucullatum*.

When the first part of the *Genera and Species of Orchideous Plants* was published, we had satisfied ourselves that in all probability Dr. Roxburgh had confounded two or three different species under the common name of *D. Pierardi*; and accordingly we took the species represented by Dr. Hooker in his Exotic Flora, as the authority for the name, and we considered Dr. Brown's *Dendrobium cucullatum*, to be a mere form of it.

We have since had an opportunity of comparing *D. Pierardi* and *cucullatum* side by side, in the utmost perfection in the stove of the Messrs. Loddiges, where these lovely species flowered in great splendour in January last, forming festoons two or three feet long, quite covered with the most delicate pink and yellow blossoms. At first sight they are so entirely the same that one does not immediately perceive in what their differences consist; but upon a more attentive inspection it is found that *D. cucullatum* has larger flowers, with a broad roundish-ovate lip, the base of which is rolled up into a sort of short stalk, while in *D. Pierardi* the lip is rounded and very blunt, and its base is rolled into a much longer stalk, which quite conceals the column; it is moreover remarkably incurved. These

* See folio 1239.

differences are expressed at the bottom of the accompanying plate, where fig. 1. represents *D. Pierardi*, and fig. 2. *D. cucullatum*. Whether or not the latter is really a distinct species may possibly be doubted; but it is certainly so remarkable a form, as to have amply justified Dr. Brown in separating it from *Pierardi*.

Allied to these plants are some new species, which we avail ourselves of the present opportunity of making known to Botanists.

1. *D. ochreatum* (Lindl. in Wall. Cat. no. 7410); caulibus brevibus e squamis membranaceis ventricosis erumpentibus, foliis ovato-lanceolatis acutis, vaginis ventricosis, floribus geminatis sepalis petalisque majoribus lanceolatis acuminatis, labello rhomboideo-ovato unguiculato cucullato intus tomentosus.

Hab. in Chittagong, *Wallich* (*hab. s. sp. e Museo Anglo-Indico*).

Species vaginis laxis, alioquin facie *D. Pierardi*, primo intuitu distinguenda. Flores expansi ferè 3 uncias lati, pallidi. Labellum purpureo-maculatum; ungue intus supra medium calloso.

A beautiful plant.

2. *D. Cunninghamii*; caulibus gracilibus pendulis ramosis vaginis foliorum corneis transversim corrugatis squamatis, foliis ovato-linearibus obtusiusculis, pedunculis oppositifoliis bifloris foliis multò brevioribus, sepalis ovatis acutis, petalis oblongis acutis latioribus, labelli trilobi lobo intermedio subrotundo undulato basi 5-lamellato lateralibus nanis acutis.

Hab. supra truncos *Callistemonis elliptici* (*A. C.*) in insula septentrionali *Novæ Zelandiæ*, juxta mare, incolis *Raumangha*; *R. Cunningham* (*hab. s. sp. comm. cel. A. Cunningham*).

This species is nearly allied to *D. biflorum*, of which I had examined no specimens at the time the *Gen. et Sp. Orch.* was published. Having since received that plant in a good state from Mr. Mathews, who found it in Otaheite hanging from the branches of trees, I am enabled to offer a correct definition of that very rare plant.

- D. biflorum* Swartz; caule pendulo gracili tereti, foliis lineari-lanceolatis acuminatis planis, pedunculis bifloris lateralibus e paleis corneis erumpentibus, sepalis petalisque acuminatissimis, labello rhomboideo medio bilamellato trilobo lobis lateralibus acutis nanis intermedio deltoideo acuminato margine fimbriato.

3. *D. Griffithianum*; caulibus erectis elongatis clavatis sursum tetragonis apice diphyllis, foliis lanceolato-oblongis obtusis, racemis laxis flexuosis multifloris pendulis e latere caulium natis, bracteis oblongis membranaceis pedicellis 4-plò brevioribus, petalis oblongis ciliolatis sepalis duplò latioribus, labello ovato pubescente denticulato subrepando, capsulis pyriformibus angulatis.

Hab. in Regno *Burmano* supra arbores, *W. Griffith*. (*hab. s. sp. comm. cel. Griffith*.)

A beautiful species with much the appearance of *D. aggregatum*.

4. *D. extincorium*; pseudobulbis depressis orbiculatis aggregatis, foliis, pedunculis terminalibus unifloris, sepalis lateralibus posticis basi longè in extincorii forma productis, labelli ungue longissimo tenui, limbo trilobo: lobis lateralibus erectis acutis intermedio oblongo rotundato, capsula pyriformi angulata.

Hab. in Regno *Burmano* locis humidis supra truncos *Careyæ* arboreæ, *W. Griffith* (*hab. s. sp. comm. cel. Griffith*.)

This is next *D. pusillum* Blume, and has very much the appearance of a *Bolbophyllum*. Its leaves have not been seen.



* *BEGÓNIA* petalódes.*Petaled Begonia.*

MONŒCIA POLYANDRIA.

Nat. ord. BEGONIACEÆ, Juss. (Introduction to the Natural System of Botany, p. 169.)

BEGONIA.—*Supra, vol. 4. fol. 284.*

B. petalodes; caulescens, foliis æquilateris orbiculatis 5-9-lobis incisiss serratis cucullatis, floribus masculis disepalis dipetalis, fœmineis tetrasepalis tetrapetalis, fructus alis subæqualibus acuminatis, cymis 2-3 floris.

Stipulæ ovatæ, serratæ. Foliorum lamina petiolo brevior. Cymæ longè pedunculatæ 2-3-floræ. Flores masculi sepalis 2 subrotundis roseis, petalis 2 conformibus sed minoribus albis. Flores fœminei sepalis 4, parvis, subrotundis roseis, petalis totidem conformibus albis quorum 2 minora sunt. Alæ fructus immaturi æquales, acuminatæ.

A pretty little species sent us by Mr. Richard Harrison, in April 1833, and we presume a native of Brazil. It adds another to the beautiful genus *Begonia*, which cultivators scarcely appreciate in this country. The species are all so very easily cultivated, and have so neat an appearance that they are exceedingly well adapted for ornamenting a damp stove. One of the prettiest things in the gardens of Schönbrunn is a *Begonia* house, filled with this genus and Ferns growing upon decayed wood and old tan.

To Botanists this is peculiarly instructive, because it shews better than any species we have previously seen what the regular form of the genus may be considered. To us it

* See fol. 1252.

is the more interesting, because its petals, distinct from the sepals, shew we were right in considering the tendency of *Begonia* to be towards the production of a corolla, and that it is consequently correctly associated in the *Nixus plantarum* with Polypetalous orders; and secondly, that two or four being the number of the floral envelopes, when distinctly formed, the relationship of *Begoniaceæ* to *Onagrariæ* is almost demonstrated.

1758.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly May. 1. 1835.

J. Watts. sc.

* ONCIDIUM citrinum.

Lemon-coloured Oncidium.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆE § VANDEÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.)

ONCIDIUM.—Supra, vol. 13. fol. 1050.

O. citrinum; pseudobulbis oblongis compressis, foliis ensiformibus rigidis scapo simplici brevioribus, sepalis petalisque labelli longitudine lineari-oblongis undulatis, labello cordato utrinque introrsum arcuato apice dilatato subreniformi, cristâ 8-tuberculatâ pubescente, alis minimis, stigmatе orbiculari.

Planta O. altissimo (fol. 1651.) valdè affinis et fortè mera varietas. Diversa tamen videtur scapo non ramoso, floribus parum maculatis, sepalis petalisque minùs acuminatis, cristâ tuberculatâ potius quam digitatâ, demum alis minimis, et stigmatе orbiculari nec angusto compressoque.

A native of Trinidad, whence it was introduced by Messrs. Loddiges, in whose collection our drawing was made last November. Unfortunately the plant soon after sickened and died; so that it is for the present lost to the country.

It approaches very nearly to *O. altissimum*, figured at fol. 1651 of this work; and is principally distinguished by the following characters. Its flowering stem is simple and not branched; its flowers are of a pale lemon colour, very distant from each other, and by no means so much spotted; the crest of the lip consists of about eight warts, which are slightly downy, and not of nine smooth finger-like processes; its stigma is nearly orbicular, and not long and narrow, and the wings of the column are exceedingly small; and finally, both the pseudo-bulb and the leaves have a singularly yellow tint, as we are informed by Mr. George Loddiges.

Like the rest of the genus it requires a hot and damp stove.

* See folio 1542.



* *PHYSIANTHUS* albens.*White Bladderbloom.*

PENTANDRIA DIGYNIA.

Nat. ord. ASCLEPIADEÆ, R. Br. (*Introduction to the Natural System of Botany*, p. 210.)

PHYSIANTHUS Martius. Corolla campanulata, tubo inflato-ventricoso, limbo 5-fido connivente. Columna fructificationis inclusa. Corona staminea inclusa, 5-phylla, foliolis tubo stamineo insertis, deinde corollæ adnatis, sursum liberis cucullatis. Antheræ membranâ terminatæ. Pollinis massæ 10, ceraceæ, compresso-clavatæ, in cruribus retinaculi deflexis pendulæ. Stigma bipiculatum. Semina comosa. Martius nov. g. et sp. 1. p. 53.

Ph. *albens*; herbacea, volubilis, foliis oppositis integerrimis acutis basi cordato-truncatis subtùs albo pruinosis, floribus subdichotomo-cymosis. Martius l. c. t. 32. Graham in Edinb. New Phil. Journ. Oct. 1832. Bot. Mag. 3201.

Corolla alba, coriacea, campanulata; limbo patulo, laciniis ovatis undulatis medio roseo penicillatis; tubo basi ventricoso. Corona staminea imâ basi corollæ inclusa; appendiculis 5, ovatis, carnosis, extrorsùm cucullatis, inflexis, staminibus oppositis. Stamina connata in conum pistillum circumdantem; antherarum loculis sagittatis, staminei coloris. Pollinia pendula, glandulâ nigrâ. Stigma globosum, apice bicornæ. Squamæ hypogynæ 0.

We received a specimen of this rare plant in August last from Mr. Blair, gardener to Miss Martineau, as a native of Mexico. Upon comparing it, however, with the figure of *Physianthus albens*, given by von Martius from specimens collected by him in the woods of Ypanema in the Province of St. Pauls, and with the plate in the Botanical Magazine of the same plant from Buenos Ayres, we are obliged to conclude either that there has been some mistake with regard

* Literally bladder-flower, in allusion to the inflated corolla.

to its introduction to the London gardens, or that it stretches over a very great extent of the American Continent.

It is a climbing plant, which runs rapidly over the rafters of a hothouse, or greenhouse, and may be easily propagated by cuttings.



* *BLÉTIA* refléxa.*Reflexed Bletia.*

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆÆ § EPIDENDRÆÆ, Lindl. (*Introduction to the Natural System of Botany*, p. 262.)

BLETIA.—*Supra*, vol. 17. fol. 1401.

B. reflexa; sepalis lineari-lanceolatis lateralibus reflexis, petalis cuneato-lanceolatis supra columnam conniventibus, labelli trilobi lobis lateralibus rotundatis planis intermedio angusto undulato lamellis 5 altis parallelis indivisis ad pedem columnæ decurrentibus, foliis angustis ensiformibus plicatis.

Folia angusta, ensiformia, plicata. Scapus $1\frac{1}{2}$ -2-pedalis, teres, distanter vaginatus, 2-4-florus. Bracteæ oblongo-lineares, membranaceæ, ovario duplò breviores. Sepala angusto-lanceolata, ex viridi rubescentia basi alba; lateralibus reflexa, supremum paulò latius magis erectum, apice tantùm reflexum. Petala ex viridi rubescentia, cuneato-lanceolata, erecta, acuta, collateralibus columnam a tergo tegentibus. Labellum cucullatum, parte inferiore album, cum basi columnæ articulatum, oblongum, trilobum; lobis lateralibus erectis rotundatis, albis, margine roseis, intermedio elongato pariter rotundato, crispo, atropurpureo, patente, lamellis 5 altis integris subundulatis totum axim labelli percurrentibus. Columna purpurea, clavata, apice alata, dente unico incurvo post cardinem antheræ. Pollinia 8, quorum 4 cæteris minora.

Drawn in the hothouse of the Messrs. Loddiges, in November 1834. It is a terrestrial species, and a native of Mexico. The greenish colour of its flowers, and the remarkably narrow reflexed sepals, distinguish it at once from the remainder of the genus.

The specimen figured was less brilliant in its colours, and altogether smaller than is to be expected hereafter; for wild specimens now before us, collected by Baron de Karwinsky, and forming part of the Royal Bavarian Museum, have a rich purple cast upon every part of the flowers, and the scapes each bear from two to four blossoms.

Requires the same treatment as other Bletias.

* See folio 1401.



* CAPRIFOLIUM hispidulum.

Bristly Honeysuckle.

PENTANDRIA MONOGYNIA.

*Nat. ord. CAPRIFOLIACEÆ, Juss. (Introduction to the Natural System of Botany, p. 206.)**CAPRIFOLIUM.—Supra, vol. 15. fol. 1232.*

C. hispidulum; tota hispido-pilosa, umbellis pedunculatis, corollis glabris bilabatis tubo limbo duplò longiore, staminibus exsertis, foliis petiolatis cordato-ovatis obtusis subtus glaucis summis sessilibus liberis, caule filiformi. *Lonicera hispidula. Douglas.*

Caulis in genere debilis, filiformis, volubilis, vel prostratus, pilis rectis distantibus ut ferè omnes aliæ partes, hispidus. Folia parva. Flores parvi, rosei, pedunculis foliorum ferè longitudine, glomerulis bibracteatis.

A very rare hardy shrub, discovered by Mr. Douglas in the woods of North West America. It is quite different from all the other honeysuckles, and is nearly scentless.

It will not readily grow more than two or three feet high, and seems as if it preferred lying prostrate to twining round other plants. In common soil it can scarcely be kept alive, but in peat and loam it grows as readily as any other hardy American plant.

Our drawing was made in the garden of the Horticultural Society in July last.

* See fol. 1232.



* LEPÁNTHES tridentáta.

Three-toothed Lepadthes.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDEÆ § MALAXIDEÆ, Lindl. (*Introduction to the Natural System of Botany*, p. 262.)

LEPANTHES Swz. *Sepala* patula, basi, præsertim lateralia, connata. *Petala* 2, libera, nana. *Labellum* apice sub-bilobum, cum columna connatum. *Columna* teres, nana, aut elongata. *Pollinia* 2.—Herbæ epiphytæ, caulibus filiformibus, monophyllis, vaginatis, spicis terminalibus, Pleurothallis omnino habitu. Flores parvi, lutescentes vel rubescentes.

L. tridentata; folio ovato acuto apice tridentato, pedunculis subbifloris folio longioribus, sepalis ovatis lateralibus semiconnatis, petalis semisagittatis tomentosis, labelli ciliati lobis lateralibus rotundatis, intermedio brevissimo acutiusculo incurvo, columnâ nanâ.

L. tridentata. Swartz *Prodr.* p. 125. *fl. Ind. occ.* p. 1561.

Caules cæspitosi, vaginis lævibus laxis cincti, staturâ a lineis 2 usque ad uncias totidem variantes. Folia ovato-oblonga, $1\frac{1}{2}$ unciam longa, crassa, apice acutè tridentata, subtus pallidiora. Pedunculi terminales, bracteati, subbiflori, foliis longiores. Flores parvi, bilabiati, oculo nudo purpureo-lutei. Sepala ovata, acuta, æqualia; supremum purpureum luteo-marginatum; lateralia semiconnata lutescentia, parum purpurata. Petala atropurpurea, tomentosa, semisagittata: lobo postico rotundato, sepalis dimidio breviora. Labellum minus, lutescens, basi concavum, purpurascens, cum basi columnæ connatum eoque parallelum, imò longius; trilobum, lobis lateralibus ascendentibus ciliatis columnam amplectantibus, intermedio incurvo minimo, minùs manifestè ciliato. Columna teres, minima, bicornis, basi tantum labello adnata; stigmatè depresso subcuneiformi. Pollinia 2, parallela, conica, caudiculis apice incrassato aurantiaco coherentibus.

This which is one of the most pigmy of Orchideæ, not much exceeding in size the moss amongst which it grows, will nevertheless amply repay the observer for a careful exami-

* So called from λεπός bark, or λεπτός small, and ανθος a flower; because the plants of this genus have very small flowers and grow upon the bark of trees.

nation of it. In order to explain its singular structure we have given magnified figures of the more important parts of its fructification.

The first thing that strikes us is the singular termination of its leaves, which do not end in a sharp point as it seems to the naked eye; but have three little delicate teeth (fig. 1.). The flowers are covered externally by three sepals, the uppermost of which is purple with a yellowish border, and the lateral ones yellow with a little stain of purple; these are frosted over with brilliant little tubercles, and glitter in the sunshine as if encrusted with emeralds. The petals are two deep purple downy parts (fig. 2.), standing upright in the centre of the blossom, and forming a sort of back for the column to rest against; the column itself (fig. 5.) is a minute two-horned body, which buries its head between the lobes of the labellum (fig. 3.); and the latter is a yellowish three-lobed downy organ, whose two lateral divisions stand up on each side of the column, in the form of tiny bucklers ready to defend it from aggression, while the centre lobe projects in the fashion of a little horn (fig. 3. and 4.).

This is the first species of the genus which has been seen alive in Europe. It is a native of the highest parts of the mountains of Jamaica, where it is found growing on the bark of trees, among mosses. It can only be preserved alive with great care by being kept under a bell glass among damp moss in a cool part of a stove. We received it from Messrs. Loddiges in January last.

1763.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly May. 1. 1835.

J. Watts. sc.

* **ÓROBUS** atropurpúreus.*Dark Purple Orobus.*

DIADELPHIA DECANDRIA.

Nat. ord. LEGUMINOSÆ, *Juss.* (*Introduction to the Natural System of Botany*, p. 87.)

OROBUS L. *Calyx* campanulatus, 5-fidus, lobis 2 superioribus brevioribus. *Corolla* papilionacea. *Stamina* diadelpa. *Stylus* gracilis, linearis, apice villosus. *Legumen* cylindræum, oblongum, 1-loculare, bivalve, polyspermum. *Semina* hilo lineari.—Herbæ erectæ. *Stipulæ* semisagittatæ. *Petioli* in setum brevem simplicem desinentes. *Folia* abruptè pinnata, paucijuga. *Racemi* axillares, pedunculati. *De Cand. prodr.* 2. 376.

O. atropurpureus; caule sub-simplici striato, foliis 1-plurijugis, foliolis linearibus acuminatis glabris, stipulis semisagittatis subunidentatis, pedunculis foliis longioribus, racemis densis secundis multifloris, corollis elongatis.

O. atropurpureus. *Desf. fl. atl.* 2. 157. t. 196. *De Cand. prodr.* 2. 386.

O. siculus. *Rafin. caratt.* 72.

O. Rafinesquii. *Presl. del. Prag.* 41.

A native of wild places near Algiers, where it was first noticed by Desfontaines. It is also met with in Sicily, and in the loamy meadows of eastern Calabria, near Cotrone and Cassano. In the gardens it is a hardy perennial, flowering in May; our drawing was made in the Garden of the Horticultural Society.

It varies in the number of its leaflets from one to several pair; Desfontaines' figure and ours represent extreme states.

* Said to be so named from *ορω* to excite and *βους* a bullock; that is to say, exciting to cattle. The Orobus of the Greeks was some sort of vetch.



S. F. Hook. del.

Pub by J. Ridgway 169 Piccadilly

June 1. 1835.

S. Walp. sc.

* CLAVÍJA ornáta.

Elegant Clavija.

PENTANDRIA MONOGYNIA.

Nat. ord. ARDISIACEÆ § C. THEOPHRASTEÆ, Bartl. ord. nat. p. 165.
THEOPHRASTEACEÆ, D. Don in G. Don. gen. syst. 4. p. 24.

CLAVIJA.—Ruiz et Pavon. Calyx 5-partitus. Corolla rotata, 5-loba. Coronæ appendices 5, carnosæ, adnatæ. Filamenta in tubum connata! Antheræ cohærentes. Ovarium 1-loculare, multi-ovulatum. Bacca crustacea, mono-polysperma.—Arbores (Amer. Æquin.) sempervirentes, trunco simplicissimo, recto, more Palmarum apice frondoso. Folia alternatim conferta, undique patentia, oblonga, coriacea, glabra, reticulato-venosa, margine dentato-spinosa v. integerrima; petioli basi callosi. Flores racemosi, albi aut aurantiaci. D. Don. MSS.

C. ornata, foliis elongato-lanceolatis mucronatis dentato-spinosis, petiolis unguicularibus, racemis glabris, baccis subdispermis. D. Don.

Clavija ornata. D. Don in edinb. phil. journ. January 1831. in G. Don gen. syst. 4. p. 25.

Theophrasta longifolia. Jacq. coll. 4. p. 136. hort. schænbr. 1. t. 116. Spreng. syst. 1. p. 670.

Arbor (in caldario culta) vix ultrà 6-pedalis, tamen in loco natali sæpè altitudinem 20 pedum attingens, trunco recto, simplicissimo, apice frondoso, cortice fusco rugoso-rimoso induto. Folia conferta, subverticillata, patentia, oblongo-spathulata v. lanceolata, mucronata, dentato-spinosa, lætè viridia, glaberrima, cartilaginea, suprà nitidula, pedalia v. sesquipedalia, spithamam lata, costá validá pallidiori subtùs elevatá, rumis primariis transversis parallelis, venulisque ramosissimis reticulata, basi attenuata, integerrima. Petioli unguiculares, robusti, fusco-purpurei, suprà planiusculi. Racemi supra-axillares, simplices, multiflori, palmares v. spithamæi, plerumque penduli, axi tereti, glabro, viridi. Flores odore Primulæ veris fragrantissimi, crocei, undique sparsi, inferioribus sæpè abortivis. Pedicelli brevissimi, robusti, teretes, glabri, pallidè aurantiaci. Calyx 5-partitus, persistens: laciniis rotundatis, coriaceis, impressè punctatis, margine membranaceis, leviter erosis, æstivatione imbricatis. Corolla rotata, substantiá crassá coriaceá; tubo brevissimo, subobconico, intùs sulcato; limbo 5-lobò, patulo; lobis subrotundis, integerrimis, æstivatione imbricatis, disco convexiusculis. Coronæ appendices 5, flavæ, carnosæ, deltoideæ, adnatæ, tuberculiformes, suprà bisulcatæ, lobis corollæ alternæ. Stamina 5, corollæ lobis opposita; filamenta in tubum membranaceum infernè ventricosum connata; antheræ cohærentes, extrorsæ, biloculares, appendiculá brevissimá rotundatá membranaceá coronatæ; loculis oblongis, distinctis, parallelis. Pollen pulvercum. Ovarium liberum, pyramidatum, uniloculare, aurantiacum; ovulis numerosis,

* Named by the authors of the Flora Peruviana in compliment to Jose Clavijo Faxardo, a Spanish Naturalist, who translated the works of Buffon into his own language.

placenta centrali insertis. Stylus vix ullus. Stigma peltatum, 5-angulum pruinatum. Bacca magnitudine Cerasi minoris, subglobosa, crustacea, lateribus parùm compressa, unilocularis, evalvis, plerumque disperma, rarò mono v. trisperma, basi calyce et apice stylo persistentibus instructa, crustâ intùs virenti nervis venisque plurimis peragrata. Semina subrotunda, substantiâ pulposâ copiosè tecta, subinde drupacea! aquâ tepidâ madida pulpa exit voluminosa, basi umbilico amplo orbiculato concaviusculo instructa: testa cartilaginea; albumen magnum, corneum, pelliculâ membranaceâ arcuè adhärenti vestitum. Embryo erectus, axilis, lacteus: cotyledones ovales, subfoliaceæ; radícula cylindræa, obtusa, cotyledonibus subæqualis, umbilico obversa. D. Don. MSS.

This very elegant tree is a native of Caraccas, whence it was introduced to the Imperial Gardens at Schönbrunn, where it flowered for the first time in November 1789, and a description and figure have been given of it by the elder Jacquin in his magnificent work on the plants of that garden. A number of young plants were raised at the late Mr. Colvill's nursery, in 1829, from seeds collected in Caraccas by Mr. Fanning, to whom I am indebted for specimens of the ripe fruit. The young plants were observed to vary a good deal in the breadth of their leaves, and as some of them produced flowers in the spring of 1834, they were found to agree precisely with the plant here figured, which blossomed about the same time in the stove of the Chelsea Botanic Garden, and which had been previously introduced from the continent under the name of *Theophrasta latifolia*.

The plant loves a high temperature, and should be grown in a soil composed of equal parts of loam and vegetable earth, and plentifully supplied with water while in a growing state.

The late Sir James Edward Smith has expressed, in Rees's Cyclopædia, a suspicion of this species being identical with the *Eresia* of Plumier, the *Theophrasta americana* of Linnæus; but the latter having a large polyspermous fruit, and the former having seldom more than two seeds, completely sets the question at rest as to their specific difference at least.

The *Theophrasteaceæ*, consisting of *Theophrasta*, *Clavija*, *Jacquinia*, and *Leonia*, constitute a small group intermediate between *Myrsinæ* and *Sapotæ*, being distinguished from the former by their polyspermous fruit, foliaceous embryo, extrorse anthers, and by the presence of coronary appendages alternating with the stamina, and which are to be regarded as the rudiments of a second series of those organs. *Theophrasta* agrees with the present genus in having extrorse anthers, but differs in its campanulate corolla, annular inclosed crown, and free stamina.

The dried flowers and seeds of *Clavija* and *Jacquinia* when immersed in warm water emit an agreeable fragrance resembling that of the Primrose. *D. Don.*

For these valuable notes we are indebted to Mr. Don, who had an opportunity of examining the specimen from which our drawing was made. We are only acquainted with the plant from the accounts of Jacquin to which Mr. Don has referred.



* EPIDÉNDRUM grácile.

Graceful Epidendrum.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆ, Juss. (Introduction to the Natural System of Botany, p. 262.)

EPIDENDRUM.—Supra, vol. 1. fol. 17.

E. gracile ; foliis in pseudobulbos ovatos corrugatos pluribus levato-ensiformibus, racemo simplici longissimo, sepalis oblongis petalisque cuneatis patentibus, labelli ferè liberi trilobi lobis lateralibus semiovatis intermedio oblongo crispo obtusissimo duplò minoribus disco bicostato.

Labellum tripartitum, cum basi columnæ leviter connatum ; laciniis lateralibus erectis, semiovatis, obtusis, subundulatis, flavis sanguineo venosis, intermediâ subrotundâ crispâ ejusdem coloris brevioribus ; in medio crassum, album venis purpureis, bicostatum ; costis utrinque infra medium obsolete unidentatis. Ovarium brevè cuniculatum. Columna semiteres, marginata apice utrinque auriculata. Gynizus deorsum trilobus ; rostillum crassum, breve, rotundatum. Anthera complete 4-loculares ; dissepimentis membranâ marginatis. Pollinia 4, compressa basi unidentata ; caudiculis totidem pulvereis replicatis.

A native of the Bahamas, whence it was sent to the Horticultural Society by John Campbell Lees, Esq. in 1833. From a plant which flowered last September the accompanying drawing has been taken.

The plant is very nearly allied to *E. odoratissimum*, but its leaves are three times as long, and the flowering stem shoots up to the length of three feet, without a branch or a leaf beyond the little scales that appear upon it here and there ; the flowers moreover are larger, and the form of the segments of the lip materially different, while the

* See folio 1415.

callosities upon the disk of the latter distinguish it positively. In *E. odoratissimum* the disk is simply thickened and flattened and gradually thins away towards the base; in *E. gracile* it is distinctly two-ribbed, and has two obsolete teeth a little below the middle.

The want of beauty in this plant is amply compensated for by its delicious fragrance, early in the morning and at night. It seems as if it were more terrestrial than epiphytal, and grows freely in any light well drained soil. The patch sent home by Mr. Lees consisted of a mass of pseudobulbs nearly two feet in diameter.



Miss Drake. del.

Pub. by J. Ridgway 169 Piccadilly June 1. 1835.

J. Walpole. sc.

* **VERBÉNA** multífida; *var. contracta*.*Dwarf Purple Vervain.*

DIDYNAMIA ANGIOSPERMIA.

*Nat. ord. VERBENACEÆ, Juss. (Introduction to the Natural System of Botany, p. 238.)**VERBENA.—Supra, vol. 4. fol. 294.**V. multifida*; diffusa pilosa, foliis 3-5-partitis laciniatis, corymbis subcapitatis pedunculatis, calycibus hispidè pilosis, corollæ laciniis patentibus emarginatis.*V. multifida. Flora Peruv. vol. 1. p. 21. tab. 33. f. c. Syst. p. 338. (1798.)**V. erinoides. Willd. enum. p. 634. no. 12. (1809!)—Hooker Bot. Misc. 1. 168.**Erinus laciniatus. Linn. Sp. Pl. 879.*"Lychnidea Verbenæ tenuifoliæ folio, vulgò Sandia Laguen. *Feuill. Per. v. 3. t. 25.*"

One of the commonest of all plants on the alps of Chile and Mendoza, growing according to Dr. Gillies to the elevation of 8000 feet above the level of the sea; and varying extremely in the colour of the flowers, in stature, and in the degree in which the leaves are cut. In some individuals the flowers are said to be scarlet, in others blue, in the case before us purplish, and we can scarcely doubt that the *V. sulphurea* of fol. 1748 is in reality another form.

It is a hardy perennial; and one of its more robust varieties is now a common ornament of the flower garden, over which it runs during the summer, forming patches several feet in diameter.

What we now figure is a dwarfer and more short-jointed kind, our drawing of which was made in the Garden of

* See folio 1184.

the Horticultural Society last July. It looks almost like a species of scentless Thyme, and grows into a very dense patch which has but little disposition to extend itself.

As there is no reason why the original name of *V. multifida* should be made to give way to Willdenow's alteration to *V. erinaides*, we retain the oldest designation.



* *GESNÉRA allagophýlla.**Shifting-leaved Gesnera.*

DIDYNAMIA ANGIOSPERMIA.

Nat. ord. GESNEREÆ, Richard. (Introduction to the Natural System of Botany, p. 227.)

GESNERA.—Supra, vol. 4. fol. 329.

G. allagophylla; caule erecto piloso sæpè trigono, foliis villosis subsessilibus ternis oppositis vel sparsis lineari oblongis spatulatisve obtusis crenatis, floribus sessilibus in verticillis spicâve longâ terminali, corollæ tubo piloso cylindrico: limbo æquali patenti glabro, glandulis hypogynis duabus posticis.
De Martius Nov. Gen. et Sp. pl. 3. 36.

A stove plant, our drawing of which was made last July from a specimen belonging to Mr. Young of Epsom. Unfortunately we had no opportunity of making any notes concerning its structure, and consequently we are uncertain whether it is rightly referred to the above named species. It agrees very well with von Martius' careful description, except that we did not observe the upper part of the stem to be at all triangular.

If we are right in the species to which we have referred it, this is a native of the auriferous plains of Brazil in the Province of the Mines, in various places, especially about Villa Rica, the town of S. João d'El Rey, in the Serras da Pires and da Piedade, and elsewhere. It has also been met with by Sellow, beyond the Tropic, in the Province of St. Paul's. (*Martius.*)

A very pretty neat species, and an interesting addition to the showy and easily cultivable genus to which it belongs.

* See folio 1158.



* CAMPÁNULA Gargánica.

The Harebell of St. Angelo.

PENTANDRIA MONOGYNIA.

*Nat. ord. CAMPANULACEÆ, Juss. (Introduction to the Natural System of Botany, p. 185.)**CAMPANULA.—Supra, vol. 1. fol. 56.*

C. garganica; foliis radicalibus reniformibus longissimè petiolatis, caulinis cordatis omnibus crenato-dentatis pubescentibus, floribus axillaribus subfasciculatis (subsolitariis) corollis 5-partitis. *Alph. Decand. Monogr. des Campan. p. 299.*

C. garganica. Tenore in *Floræ neap. add. cum app. sem. h. r. neap. 1827. Sylloge, p. 95.* Sweet's *Brit. Fl. Gard. t. 252.*

For the opportunity of figuring this rare and beautiful species of harebell we are indebted to the kindness of Mrs. Marryat of Wimbledon, in whose rich collection it flowered in great perfection in July last. It had been raised from seeds presented to Mrs. Pallisser by Professor Tenore.

The following are the observations of that Botanist concerning this little known species. It is a perennial, flowering in June, and grows wild on Mount Gargano, among the ruins of the ancient Monastery, and near fort St. Angelo. It agrees in the form of its leaves, and in the branching of its peduncles with *C. Portenschlagiana*; but differs in the lobes of the calyx being twice not thrice as short as the corolla, and in the deep divisions of the latter. From *C. diffusa*, which it resembles in many respects, it may be distinguished by its deeply 5-parted corolla, with narrow lanceolate, revolute segments, and not half 5 cleft with broad ovate spreading lobes; the flowers moreover are racemose and not corymbose. From *C. Elatine* it differs in its stem

* See fol. 1738.

being altogether diffuse and by no means ascending, tolerably firm, and from 3 to 6 inches long, in its root-leaves being nearly round and kidney-shaped, with double wavy crenellings, and not ovate, or between ovate and heart-shaped, with taper pointed pectinate teeth, in its equal corolla, which is pale blue and not bluish purple.

In many respects this account is somewhat at variance with the plant we have figured, and accords better with that published by Mr. Don in Sweet's Flower Garden; but our wild specimens from the shady rocks of Mount Gargano agree exactly with the individual that flowered in Mrs. Marryat's garden, and we conclude that the species is liable to variations in the crenelling of its leaves and in the colour and depth of the lobes of the corolla.

It is presumed that like most of the plants of the South of Italy this will require protection in winter from the heavy rains of this climate, otherwise it will probably prove hardy in the clefts of rockwork, of which it is a charming ornament.



Miss Drake, del.

Pub by J. Ridgway 109 Piccadilly June. 1. 1835.

J. Watts, sc.

* **PSORÁLEA macrostácha.***Long-spiked Psoralea.*

DIADELPHIA DECANDRIA.

Nat. ord. LEGUMINOSÆ, Juss. (Introduction to the Natural System of Botany, p. 87.)

PSORALEA.—*Supra, vol. 12. fol. 968.*

P. macrostachya; foliis pinnatim 3-foliolatis pubescentibus, foliolis ovatis mucronatis, petiolis glanduloso-scabris, pedunculis axillaribus foliis quadruplò longioribus, spicis cylindræis rachi bracteis calycibusque hirsutissimis. *Decand. prodr. v. 2. 220.*

Herba perennis, erecta, ramosa, 3-pedalis, undique pubescens. Foliola integerrima, utrinque præsertim supra glandulosa; lateralia petiolulata. Stipulæ membranacæ, ovatæ, acuminatæ. Pili racheos, bractearum, calycumque nigri densissimi; pedunculorum ferè nulli. Flores decidui, rachin hirsutum rectum linguentes. Calycis labium superius 4-dentatum. Corolla purpurea calyce parum longior; alis obtusissimis. Legumen ovatum, mucronatum, pilosum, calycis longitudine. Semen oblongum, aterrimum, glabrum.

This species is described by De Candolle, from specimens communicated by La Gasca, and purporting to come from Nutka. Our drawing was however made from plants obtained from California through Mr. Douglas; and we therefore suspect some error in the former statement.

It is a handsome hardy perennial, yielding seed in tolerably plenty, and readily increasing by division. Our drawing was made in the Garden of the Horticultural Society in July of last year.

It is remarkable for the singular hairiness of the rachis and green parts of the flower, beginning abruptly with the

* So named, from ψωραλεος warty, in allusion to the very general presence of little tubercular secretions upon the surface of different species.

origin of the first flower, but which is scarcely at all visible on the flower-stalk itself. After the flowers have fallen off this produces a singular appearance, the naked rachis remaining at the end of each flower-stalk in the form of a long hairy tail.

Fig. 1. represents a side view of an entire flower ; 2. is a front view of the same ; 3. a calyx with the corolla pulled off to shew the stamens ; 4. a pod, and 5. a ripe seed.



* **PENTSTÉMON** staticifólius.*Sea Lavender-leaved Pentstemon.*

DIDYNAMIA ANGIOSPERMIA.

Nat. ord. SCROPHULARINEÆ, Juss. (Introduction to the Natural System of Botany, p. 228.)

PENTSTEMON.—*Supra, vol. 13. fol. 1121.*

P. staticifolius; caule ascendente pubescente, foliis radicalibus oblongo-lanceolatis in petiolum longum angustatis integerrimis glabris: caulinis sessilibus cordato-ovatis dentatis pubescentibus, cymis subsessilibus calycibusque tomentosis, corollis ventricosis pubescentibus labiorum laciniis ovatis obtusis subæqualibus.

Caulis subbipedalis, apice tomentosus et cymis subsessilibus verticillastros Labiatarum referentibus onustus. Folia radicalia cum petiolo 7-poll. longa. Corolla violacea, speciosa, fere pollicem et dimidium longa.

A new hardy species of this beautiful genus, sent to the Horticultural Society by the late Mr. Douglas from California. It is most nearly related to *P. diffusus*, figured at fol. 1132 of this work, from which it differs in its much larger and more lilac flowers, in the form of its leaves, and in those next the root being perfectly entire.

Our drawing was made in the Garden of the Society in June last. At present the species is extremely rare; only one plant having been originally raised. It grows and flowers freely in a peat border.

* See fol. 1245.

We avail ourselves of a vacant leaf or two in this number to present our Botanical friends with a revision of the whole of the genera of the natural order to which *Pentstemon* belongs. For this valuable contribution we are indebted to George Bentham, Esq. the learned author of the *Genera and Species Labiatarum*.

Tribus I. *VERBASCEÆ*, *Bartl.* Corolla tubo abbreviato vel subgloboso, limbo explanato 4-5-fido vel bilabiato non ringente. Stamina fertilia 2-5, sæpius declinata. Antheræ approximatae vel coherentes nunc uniloculares nunc subbiloculares loculis divaricatis confluentibus. Capsula bivalvis septicide dehiscens, valvulis sæpe bifidis, dissepimento e marginibus valvulorum inflexis duplicato solubili. Semina nuda. Genera Solanaceis rotatis affinia sed æstivatione corollæ abunde distincta.

* *Corolla subrotata 5-loba.*

1. *VERBASCUM*, *Linn.* Stamina fertilia 5.
2. *CELSIA*, *Linn.* Stamina fertilia 4. Capsula globosa usque ad basin dehiscens.
3. *ALONSOA*, *Ruiz et Pav.* Stamina fertilia 4. Capsula ovato-oblonga vix ad medium dehiscens. Flores torsione pedicelli resupinati. (*Hemimeris Auct. quoad species Americana*).

** *Corolla bilabiata labiis concavis.*

4. *JOVELLANA*, *Cav.* Corollæ lobi concavi subæquales patentés.
5. *CALCEOLARIA*, *Linn.* Corollæ lobus inferior incurvus calceolatus.

*** *Corolla tubo subgloboso limbo valde inæquali.*

6. *SCROPHULARIA*, *Linn.*

Tribus II. *HEMIMERIDEÆ*. Corolla tubo abbreviato, fauce concava, limbo explanato lacinia infima (sæpe bifida) basi concava 1-2-saccata vel calcarata. Stamina fertilia 2 vel 4 didynama adscendentia. Antheræ approximatae uniloculares vel biloculares loculis divaricatis. Capsula bivalvis, loculicide dehiscens valvulis integris bifidisve vel septicide dehiscens valvulis bifidis, rarius indehiscens, dissepimento e marginibus valvularum inflexis duplicato solubili vel rarius concreto.

* *Corolla ecalcarata breviter bisaccata. Capsula globosa.*

7. *ANGELONIA*, *Humb. et Kunth.* Capsula loculicide dehiscens valvulis integris.
8. *PHYLACANTHUS*, *Nees et Mart.* Capsula indehiscens.
9. *HEMIMERIS*, *Linn.* Capsula loculicide dehiscens septicide bipartibilis vel dehiscens. (*Diascia, Link.*)

** *Corolla calcarata. Capsula compressa apice truncata.*

10. *NEMESIA*, *Vent.*

Tribus III. *ANTIRRHINEÆ*, *Chavannes.* Corolla tubulosa limbo personato vel ringente bilabiato rarius subæquali. Stamina fertilia 4 didynama. Antheræ per paria approximatae biloculares. Capsula dentibus seu valvulis pluribus operculisve circumscissis dehiscens vel irregulariter rumpens. Semina nuda vel testa arilliformi inclusa.

* *Corolla basi calcarata.*

11. ANARRHINUM, *Desf.* Corolla bilabiata. Capsula operculis 2 valviformibus dehiscens.

12. LINARIA, *Tourn.* Corolla personata. Capsula operculis circumscissis vel plurimis dentiformibus vel valviformibus dehiscens.

* *Corolla basi saccata gibba vel æqualis.*

13. ANTIRRHINUM, *Linn.* Corolla personata. Capsula apice poris vel foraminibus 2-3 dehiscens.

14. MAURANDIA, *Jacq.* Corolla personata. Capsula apice dentibus 10 dehiscens (*Usteria, Cav.*).

15. GALVESIA, *Juss.* Corolla bilabiata. Capsula sub apice irregulariter rumpens; tubus corollæ intus nudus (*Agassizia, Chavannes*).

16. LOPHOSPERMUM, *Don.* Corolla bilabiata. Capsula sub apice irregulariter rumpens; tubus corollæ intus pilorum seriebus 2 percursus.

17. RHODOCHITON, *Zucc.* Calyx amplus coloratus. Corollæ limbus subæqualiter 5-fidus. Capsula sub apice irregulariter rumpens.

Tribus IV. SALPIGLOSSIDÆ. Corolla tubo elongato rarius abbreviato, limbo obliquo nunc irregulariter bilabiato nunc concavo vel subplano lobis 5 sæpe incisus. Stamina fertilia 2 vel 4 didynama declinata. Antheræ biloculares loculis apice demum sæpius confluentibus. Capsula 2-4-valvis valvulis septicide dehiscentibus bifidisve. Genera plura a Solanaceis capsularibus (præsertim a *Petunia*) nonnisi æstivatione corollæ imbricativa et staminibus 4 nec 5 differrunt.

* *Corolla basi supra gibba limbo valde irregulari.*

18. COLLINSIA, *Nutt.* Corolla 5-fida. Capsula globosa 4-valvis.

19. SCHIZANTHUS, *Ruiz et Pav.* Corolla multifida. Capsula oblonga.

** *Corolla infundibuliformis limbo parum inæquali.*

20. SALPIGLOSSIS, *Ruiz et Pav.* Corollæ tubus ampliatus subrectus, limbi explanati laciniæ emarginatæ vel bifidæ.

21. BROWALLIA, *Linn.* Corollæ tubus tenuis rectus apice parum dilatatus, limbi lobis integerrimis emarginatisve.

22. FRANCISCEA, *Pohl.* Corollæ tubus tenuis apice incurvus, limbi lobis integerrimis planis.

23. ? BRUNSFELSIA, *Linn.* Corollæ tubus elongatus rectus æqualis, limbi lobis integerrimis planis. Capsula baccata.

24. ANTHOCERCIS, *Labill.* Corollæ tubus ampliatus limbo subrecto.

Tribus V. DIGITALEÆ. Corolla tubulosa sæpius ventricosa limbo bilabiato. Stamina basi declinata apice sæpissime adscendentia, 4 fertilia didynama, quinto summo sterili vel nullo. Antheræ biloculares loculis demum sæpissime divaricatis confluentibus. Stigma simplex, vel in lobis brevibus styli 2 vix incrassata. Capsula dura bivalvis, valvulis septicide dehiscentibus sæpius bifidis vel bipartitis, dissepimento duplicato e marginibus valvularum inflexis demum bipartibili. Placentæ a dissepimento demum sæpius solutæ. Genera priora Bignoniaceis affinia sed abunde distincta. Tribus habitu magis quam characteribus a Gratioleis differt.

* *Stamen quintum præsens sterile.*

25. CHELONE, *Linn.* Corollæ labium superius amplum concavum. Semina membranaceo-alata.

26. PENTSTEMON, *Lhér.* Corollæ laciniæ subplanæ. Semina nuda.

** *Staminis quinti vestigium nullum.*

27. RUSSELIA, *Jacq.* Corollæ laciniæ vix inæquales.

28. *DIGITALIS*, Linn. Corolla declinata labio inferiore longiore.
 29. *ISOPLEXIS*, Lindl. Corolla incurva labio superiore longiore.

Tribus VI. *GRATIOLEÆ*. Corolla tubulosa limbo bilabiato. Stamina fertilia 2 vel 4 adscendentia. Antheræ biloculares muticæ. Capsula 2-4-valvis septicide vel loculicide dehiscens valvulis integris bifidisve, placentæ dissepimento non bipartibili demum libero adnatæ. Semina nuda.

§ 1. *Antherarum loculi disjuncti.*

30. *PTEROSTIGMA*, Benth. Stamina superiora fertilia antherarum loculis disjunctis, inferiora sterilia vel dimidiata. Calyx sub 5-partitus lacinia suprema majore. (*Adenosma villosum*, *A. macrophyllum* et *Stemodia capitata*. Benth. in Wall. Cat. n. 3851, 3853 et 3926, cum specia quarta nova ex China.)

31. *LINDENBERGIA*, Link. et Otto. Stamina 4 fertilia loculis disjunctis. Calyx semi 5-fidus (*Brachycoris*, Schrader, Bovea, Decaisne).

32. *STEMODIA*, Linn. Stamina 4 fertilia loculis disjunctis. Calyx 5-partitus (*Modestia* et *Diamoste*, Cham. et Schlecht. *Cybbanthera*, Hamilt.)

§ 2. *Antherarum loculi contigui. Capsulæ valvulæ 4 vel si 2 loculicide dehiscentes vel fissæ.*

* *Stamina fertilia 4. Calyx semi-5-fidus. Capsulæ valvularum margines septiferæ inflexæ.*

33. *DODARTIA*, Linn. Calyx tubuloso-campanulatus dentibus rectis. Corollæ labium inferius basi papillosum bigibbosum. Herba rigida paucifoliata.

34. *MAZUS*, Lour. Calyx campanulatus dentibus patentibus. Corollæ labium inferius basi papillosum bigibbosum. Herbæ humiles scapis subnudis (*Hornemannia*, Willd.).

35. *LIMNOPHILA*, Br. Calyx tubuloso-campanulatus labio inferiore glabro laciniis planis. Herbæ uliginosæ.

** *Stamina fertilia 4. Calyx 5-partitus. Capsulæ valvularum margines septiferæ inflexæ.*

36. ? *CAPRARIA*, Linn. Corolla vix bilabiata. Stigma obtusum vix incrassatum (*C. biflora*, Linn.).

37. *MORGANIA*, Br. Corolla bilabiata. Stigma bilamellatum.

*** *Stamina fertilia 4. Calyx tubulosus 5-dentatus. Capsulæ valvularum margines subplanæ.*

38. *MIMULUS*, Linn. Capsula 2-valvis.

39. *UVEDALIA*, Br. Capsula 4-valvis.

**** *Stamina fertilia 4. Calyx 5-partitus. Capsulæ valvularum margines planæ.*

40. *HERPESTIS*, Gaertn. Capsulæ valvulæ bipartitæ. Calyces laciniæ valde inæquales (*Monniera Mich.* *Bramia Lam.*).

41. *SPHÆROTHECA*, Cham. et Schlecht. Capsulæ valvulæ bipartitæ. Calycis laciniæ subæquales.

42. *MATOUREA*, Aubl. Capsulæ valvulæ subintegre loculicide dehiscentes. Calycis laciniæ parum inæquales (*Species americanæ 4-5 inclusa Gratiola acuminata*, Elliott.).

***** *Stamina fertilia 2. Calyx 5-partitus.*

43. *GRATIOLA*, Linn. Stamina superiora fertilia.

44. *BEYRICHIA*, Cham. et Schlecht. Stamina inferiora fertilia. Capsulæ valvulæ vix fissæ (Sp. 4, Brasilienses).

45. ? ACHETARIA, Cham. et Schlecht. Stamina inferiora fertilia. Capsulæ valvulæ bipartibiles.

***** *Stamina fertilia 2. Calyx 3-5-dentatus vel 5-fidus.*

46. DOPATRIUM, Hamilt. Corolla calyce multo longior fauce ampliata (Gratiola juncea, lobelioides et nudicaulis).

47. MICROCARPÆA, Br. Corolla calycem minutum vix excedens.

§ 3. *Antherarum loculi contigui. Capsulæ valvulæ 2 integræ marginibus planis, dissepimento membranaceo parallelo.*

* *Stamina fertilia 2.*

48. PEPLIDIUM, Delil. Stamina sterilia nulla. Capsula sæpius irregulariter rupta valvulis vix solutis.

49. BONNAYA, Link. et Ott. Stamina sterilia 2. Capsula constanter valvatim dehiscens.

** *Stamina fertilia 4, inferiorum filamenta basi appendiculata vel gibba.*

50. VANDELLIA, Linn. Calyx 5-partitus vel regulariter 5-dentatus non plicatus. Corollæ tubus intus nudus (Tittmannia, Reichb. Torenia, Cham. et Schlecht.).

51. ARTANEMA, Don. Calyx 5-partitus. Corollæ tubus intus 4-squamatus (Diceros, Pers. non Lour.).

52. TORENIA, Linn. Calyx tubulosus plicatus oblique 5-dentatus (Nortenia, Dup. Thou.).

*** *Stamina fertilia 4, filamentis nudis integris.*

53. ? HETERANTHIA, Nees et Mart. Antheræ filamentorum inferiorum magnæ stylum amplectentes (Vrolikia, Spreng.).

54. LINDERNIA, Linn. Antheræ conformes. Calyx 5-partitus.

55. LIMOSELLA, Linn. Antheræ conformes. Calyx 5-dentatus vel 5-fidus.

Tribus VII. BUCHNEREÆ. Corolla tubo tenui, limbo subplano 4-5-fido, laciniis sæpe bifidis. Stamina fertilia 4 didynama adscendentia. Antheræ uniloculares vel loculis demum divaricatis confluentibus. Capsula 2-valvis valvulis integris bifidisve. Genera nonnulla Selagineis habitu affines.

* *Capsulæ valvulæ loculicido-dehiscentes elasticæ integræ.*

56. BUCHNERA, Linn. (Campuleia, Dup. Thou. tubo apice incurvo. Piripea, Aubl. tubo apice recto.)

** *Capsulæ valvulæ bifidæ marginibus septiferis inflexis.*

57. NYCTERINIA, Don. Calyx oblongo-tubulosus 5-angulatus 5-dentatus. Corollæ laciniæ emarginatæ vel bifidæ (Erini Capenses).

58. ERINUS, Linn. Calyx 5-partitus. Corollæ laciniæ emarginatæ (E. alpinus).

59. MANULEA, Linn. Calyx 5-partitus. Corollæ laciniæ integerrimæ inæquales (Nemia, Berg.).

60. SUTERA, Roth. Calyx 5-partitus. Corollæ laciniæ 5 brevissimæ integræ æquales vix patentes (S. glandulosa, Roth. et Capraria multifida, Linn.?).

Tribus VIII. BUDDLEIÆ. Corolla tubo recto, limbo plano patente æqualiter 4-fido. Stamina fertilia 4 subæqualia. Antheræ distantes biloculares. Capsula 2-valvis.

61. BUDDLEIA, Linn.

Tribus IX. TEEDIEÆ. Corolla tubulosa bilabiata vel subinfundibuliformis, limbo subæqualiter 5-fido. Stamina fertilia 4 tubo inclusa didynama vel subæqualia distantia. Antheræ biloculares. Fructus baccatus.

* *Bacca indehiscens.*

62. LEUCOCARPUS, *Don.* Corolla bilabiata. Calyx 5-dentatus.

63. TEEDIA, *Rud.* Corollæ limbus subæqualis. Calyx 5-partitus.

** *Capsula baccata 4-valvis.*

64. HEMIPHRAGMA, *Wall.*

Tribus X. VERONICEÆ. Corolla rotata infundibuliformis vel rarius irregulariter bilabiata. Stamina fertilia nunc 4 (vel 6-7?) subæqualia exserta distantia nunc 2 (superiora) inferiorum vestigiis nullis. Stylus simplex stigmatē tenui vel rarius incrassato subcapitato. Capsula 2-valvis loculicide dehiscens valvulis sæpissime septicide bifidis bipartitisve. Semina nuda vel testa laxa arilliformi inclusa.

* *Stamina 4 (vel 6-7?). Corolla 5 (vel 6-7?) fida.*

65. OURISIA, *Comm.* Corolla campanulata 5-fida. Stamina 4.

66. ? DISANDRA, *Linn.* Corolla rotata 6-8-fida. Stamina 5-7. (*An hujus ordinis?*)

67. SIBTHORPIA, *Linn.* Corolla rotata 5-fida. Stamina 4.

** *Stamina 4. Corolla 4-fida.*

68. SCOPARIA, *Linn.* Corolla rotata.

69. GEOCHORDA, *Cham. et Schlecht.* Corolla infundibuliformis.

70. PICRORHIZA, *Royle.* Corolla campanulata calyce brevior. (*Veronica Lindleyana, Wall.*)

*** *Stamina 2.*

71. VERONICA, *Linn.* Corolla rotata vel infundibuliformis (*Pæderota, Linn. Leptandra, Nutt. Diplophyllum, Lehm. Cochlidiospermum, Reichb. Omphalospora, Bess. Hebe, Juss. Aidelus, Spreng.*).

72. CALORHABDOS, *Benth.* Corolla tubulosa incurva ore oblique 2-3-fido (*Veronica Brunoniana, Wall.*).

73. WULFENIA, *Jacq.* Corolla tubulosa recta limbo subbilabiato. Calyx 4-5-partitus.

74. GYMNANDRA, *Pall.* Corolla tubulosa recta limbo subbilabiato. Calyx 2-partitus (*Lagotis, Gærtn.*)

Tribus XI. GERARDIEÆ. Corolla campanulata infundibuliformis vel tubulosa, limbo 5-fido laciniis rotundatis planis. Stamina fertilia 4 adscendentia. Antheræ approximatae 2-loculares loculis discretis parallelis sæpe acuminatis. Stylus simplex, stigmatē obtuso sæpius elongato. Capsula 2-valvis loculicide dehiscens valvulis integris bifidisve. Semina sæpissime testa membranacea laxa arilliformi inclusa.

* *Calyx 5-dentatus. Corolla infundibuliformis.*

75. ESCOBEDIA, *Ruiz et Pav.* Corolla tubus elongatus limbo amplo 5-fido.

76. PHYSOCALYX, *Pohl.* Calyx inflatus. Corollæ limbus parvus æqualiter 5-fidus.

** *Calyx 5-dentatus vel 5-fidus. Corolla campanulata vel tubulosa.*

77. VIRGULARIA, *Ruiz et Pav.* Calyx 5-dentatus. Corolla ample tubulosa.

78. MACRANTHERA, *Nutt.* Calyx profunde 5-fidus. Corolla tubulosa vel incurva. Capsula acuminata.

79. GERARDIA, *Linn.* Calyx 5-dentatus vel semi-5-fidus. Corolla campanulata. Capsula obtusa (*Melasma, Berg.? Sopubia, Hamilt.*).

80. ? SEYMERIA, *Pursh.* Calyx 5-partitus. Corolla rotato-campanulata. Capsula acuta (*Afzelia, Gmel.*)

81. GLOSSOSTYLIS, *Cham. et Schlecht.* Calyx 2-5-dentatus. Corolla oblique campanulata. Semina intra membranam lineari-cuneatam minima. (Species observatæ tres: 1. *G. aspera Cham. et Schlecht. Brasiliensis*, 2. *Rhinanthus scaber Thunb. ? Capensis*, 3. *Hymenospermum dentatum Benth. Avensis*, et forsàn quarta inter plantas Senegalenses Perrottetianas).

*** *Calyx compressus hinc fissus.*

82. CENTRANTHERA, *Br.* (*Razumovia, Spreng. ?*)

Tribus XII. RHINANTHÆ. Corollæ limbus bilabiatus labio superiore concavo galeato vel lineari, integro vel emarginato, inferiore patente trifido. Stamina fertilia 4 didynama (rarius 2) adscendentia. Antheræ biloculares loculis discretis parallelis sæpe acuminatis. Stylus simplex. Capsula 2-valvis loculicide dehiscens valvulis sæpissime integris. Seminum nunc testa laxa membranacea inclusa, nunc nuda.

* *Antherarum loculi tenues mutici.*

83. ORTHOCARPUS, *Nutt.* Antherarum loculi inæquales. Calyx tubulosus vel campanulatus 4-fidus.

84. CASTILLEJA, *Mut.* Antherarum loculi inæquales. Calyx compressus bifidus vel quadrifidus hinc fissus (*Euchroma, Nutt.*).

85. SCHWALBEA, *Linn.* Antherarum loculi æquales. Calyx compressus oblique 5-fidus.

** *Antherarum loculi crassi sæpius acuti vel aristati.*

86. LAMOUROUXIA, *Humb. et Kunth.* Corolla tubulosa compressa ventricosa labio inferiore brevi. Calyx 4-5-fidus.

87. CYMBARIA, *Linn.* Corolla tubulosa ventricosa. Calyx 10-partitus.

88. ODONTITES, *Stev.* Calyx campanulatus 4-fidus. Corolla profunde bilabiata labio superiore laciniisque inferioris subæquilongis integerrimis vel intermedia retusa. Antheræ oblongæ galeam superantes.

89. EUPHRASIA, *Linn.* Calyx campanulatus 4-fidus. Corollæ labium superius galeatum emarginatum inferius magnum patens laciniis emarginatis (*Parentucellia, Viv. ?*)

90. SIPHONOSTEGIA, *Benth.* Calyx infundibuliformis tubo elongato limbo 4-5-partito. Corollæ labium superius incurvum inferius brevius (Species unica *Chinensis*).

91. BARTSIA, *Linn.* Calyx campanulatus vel tubulosus 4-fidus. Corollæ labium superius galeatum compressum inferioris laciniæ integerrimæ. Antheræ ovatæ galea breviores (*Trixago, Stev.*).

92. PEDICULARIS, *Linn.* Calyx 5-fidus vel inæqualiter 2-3-fidus. Corollæ galea elongata sæpius incurva.

93. RHINANTHUS, *Linn.* Calyx compressus membranaceo-inflatus. Corollæ labium superius galeatum. Semina pauca ala membranacea cincta (*Alectrolophus, Stev.*).

94. ELEPHAS, *Tourn.* Corollæ labium superius lineare proboscideum. Semina pauca striata (*Rhinanthus, Stev.*).

95. MELAMPYRUM, *Linn.* Corollæ galea compressa lobis replicatis. Semina in loculo solitaria magna exalata.

Genus incertæ sedis.

96. TOZZIA, *Linn.* Corolla Gratiolearum, stamina Gerardiearum, vel Rhinanthearum. Capsula abortu monosperma.

Genera mihi non satis nota. Conobea, *Aubl.* Gomara, Sanchesia, Xuarezia et Calytriplex, *Ruiz et Pav.* Leucophyllum, *Humb. et Kunth.* Dichroma, *Cav.* Curtia, *Cham. et Schlecht.* Lafuentia, *Lag.* Hemianthus, *Nutt.* Willichia, *Mut.* (Hydrantheium, *Humb. et Kunth.*). Aptosimum, *Burch.* Hydrotriche, *Zucc.* Diceros, Picria et Tripinna, *Lour.* Bœa, *Commers.* Palmstruckia, *Retz, fil.*



PÆÓNIA. Moutan; *lacera*.

Double-red curled Tree Pæony.

POLYANDRIA MONOGYNIA.

Nat. ord. RANUNCULACEÆ, Juss. (Introduction to the Natural System of Botany, p. 6.)

PÆONIA.—Supra, vol. 1. fol. 42.

GARDEN VARIETY.

Raised from the seed of *Pæonia Moutan* by Mr. William Hyland, Gardener to the Earl of Sandwich, at Hinchingsbrook, near Huntingdon, and communicated to us by Michael Foster, Esq.

This very beautiful variety is strikingly different from the other Moutans in the bright rosy red of the petals, the innermost of which are very much cut and gashed, curled up, and distinctly bordered with a narrow edge of light carmine, which sets them off to great advantage, and gives the whole flower a peculiarly rich and finished appearance. Mr. Hyland informs us that the seed was sown in September, and did not germinate for eighteen months afterwards; and that he finds such to be the case with all the seeds he has sown of this kind of *Pæony*.

It flowered for the first time in April 1834, when the plant was only three years old. It may therefore be expected that the future blossoms will be still finer than what we have figured.



campylocentrum

* ANGRÆCUM micranthum.

Small-flowered Angræcum.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆE § VANDEÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.)

ANGRÆCUM.—*Supra, vol. 18. fol. 1522.*

A. micranthum; caule brevi, foliis oblongis trinerviis apice obliquis, spicis secundis multifloris horizontalibus congestis foliis brevioribus, sepalis petalisque subæqualibus angustè ovatis apice patulis, labello conformi basi bilobo bidentato medio pubescente, calcare obtuso incurvo postico.

Caulis 2-pollicaris. Folia $1\frac{1}{2}$ -p. longa, $1\frac{1}{2}$ -p. lata, plana, crassa. Flores minuti, albi.

A very small and unattractive species, lately introduced by Messrs. Loddiges from Sierra Leone. It is quite distinct from any previously described, but retains the peculiar characters of the genus without deviation. It flowered in February.

It is curious that Angræcum should be so exclusively an African genus, that no certain species has yet been found beyond its continent or immediately dependent islands, although it evidently stretches across the whole of the tropical part of Africa itself. In addition to this from Sierra Leone, three or four other species are now in our gardens from the same place, but all of them equally inconspicuous with the present. Of these *A. clandestinum*, *teretifolium* and *distichum* have already flowered with Messrs. Loddiges. Fig. 1. represents a lip with the spur attached; 2. an entire flower; and 3, a pair of pollen masses with their gland.

* See folio 1522.



Miss Drake. del.

Pub by F. Ridgway 169 Piccadilly July 1. 1835.

J. Walts. sc.

* **RUSSELLIA juncea.***Rushy Russelia.*

DIDYNAMIA ANGIOSPERMIA.

Nat. ord. SCROPHULARINEÆ, Juss. Trib. 5. DIGITALEÆ, Bentham.
(Introduction to the Natural System of Botany, p. 228.)

RUSSELLIA, Jacq.—Corolla tubulosa recta limbo patente subbilabiata labio superiore emarginato vel bifido inferiore trifido laciniis omnibus subæqualibus planis. Stamina fertilia 4, basi declinata apice adscendentia, sterile nullum. Antheræ sub labio superiore approximatae glabræ biloculares loculis demum divaricatis confluentibus. Stylus simplex stigmatate integro parum incrassato. Semina nuda.—Frutices, ramis sæpius pendulis glabris. Folia opposita dentata vel integra. Flores nunc in racemis cymisve axillaribus dense glomerati nunc laxè racemosi pedunculis elongatis subunifloris. Corollæ sæpius coccineæ. Bentham MSS.

R. juncea; ramis tetragonis erectis junceis, foliis minimis petiolatis ovatis sub-integris, pedunculis filiformibus subbifloris.

R. juncea. Zuccarini in litt.

Caulis plantæ 3-pedalis erectus, angulatus, viridis, ramis onustus gracillimis, tetragonis, pallidè viridibus, junioribus subaphyllis, nutantibus. Folia sæpe minima, semper parva, ramulis debilioribus majora, petiolata, ovata, acuta, subtiliter serrulato-ciliata, aliquando dente uno alterove munita. Inflorescentia primo adspectu paniculata, sed revera e ramulis racemosim dispositis, gracilibus, distantibus constans, quorum apice insident flos unus duove pedicellati et cernui. Calyx parvus sepalis 5 ovatis acutis imbricantibus. Corolla tubulosa, coccinea, unciam longa, sursum dilatata, limbo subæqualiter 5-fido; laciniis obtusis duabus superioribus magis approximatis. Stamina 4, didynama. Antheræ glabræ oblongæ, apiculatæ, lobis parallelis. Stigma bilobum; lobis oblongis planis.

A very pretty Greenhouse or Stove plant found in Mexico (probably near Oaxaca), by Count Karwinski, and lately introduced to the gardens of this country from

* Named after Dr. Alexander Russel, an English Physician, who resided for some years at Aleppo, and published an account of his observations upon the Natural History of that place in the year 1756.

Berlin and Munich. We have specimens from Dr. von Martius under the name we have adopted, but we know not in what publication Professor Zuccarini has described the species.

It forms a tufted light green bush, about three feet high, almost destitute of leaves, but having a graceful appearance in consequence of the large number of its drooping slender twigs, which look like those of an Equisetum or a Casuarina, and which form a singular contrast with the handsome scarlet flowers that appear in abundance in the months of July and August.

Propagated very easily by cuttings.

We are indebted to Mr. Bentham for the amended character of the genus. Fig. 1. is a corolla cut open by dividing the lip through the middle; 2. is a stigma with its style; 3. a stamen.



Miss Drake. del.

Pub. by J. Ridgway, 169 Piccadilly July 1. 1835.

J. Walts. sc.

* **ERIÓGONUM** compósitum.*Compound Eriogonum.*

ENNEANDRIA TRIGYNIA.

Nat. ord. POLYGONEÆ, Juss. Trib. ERIOGONEÆ, Benth. in Linn. Trans. ined. (Introduction to the Natural System of Botany, p. 169.)

ERIOGONUM Michx.—*Involucrum* tubulosum campanulatum vel cyathi-forme vix angulatum subæqualiter 6-dentatum, multiflorum. *Receptaculum* bracteolis intra pedicellos sæpe instructum. *Flores* hermaphroditæ vel dioici. *Perianthia* exserta profunde 6-fida laciniis 3 exterioribus, 3 interioribus sæpe dissimilibus. *Stamina* 9 basi subcoalita. *Ovarium* triquetrum. *Styli* 3 filiformes. *Stigmata* terminalia tenuia vel parum incrassata capitata. *Achenium* triquetrum vel vix trilobatum.—Herbæ suffruticesve sæpissime lanuginosæ. *Folia caulina alterna ad basin caulis approximata cæspitosa vel secus ramos fasciculata exstipulata*, petiolo basi dilatato sæpius amplexicauli vel vaginante. *Pedunculi vel rami floriferi terminales nunc simplices involucri solitario terminali sæpius 2-3-chotome vel umbellatim ramosi, involucri solitarii glomeratisve, intra ramificationes, secus ramos vel ad apices sessilibus pedicellatisve.* *Bractææ vel folia ramorum floralium sub ramificationibus orta tot quot rami vel (ramis abortientibus) numerosiora nunc minuta squamæformia, nunc herbacea et foliis caulinis difformia, sæpissime sessilia, exstipulata vel stipulis scariosis aucta.* Bentham in Linn. Trans. ined.

§ 3. *Umbellata*, Benth. *Folia omnia exstipulata. Genitalia pilosa, filamenta prope basin ovarii. Inflorescentia umbellata pedunculo (sæpe scapiformi) apice umbellam simplicem vel duplicem pluriradiatam nonnunquam in capitulum globosum contractam vel ad involucrum solitarium reductam gerente. Bentham l. c.*

E. compositum (Dougl. MSS.) foliis ad basin caulis approximatis longe petiolatis ovatis basi rotundatis cordatisve supra demum glabratis subtus dense albolanatis, pedunculo longissimo nudo apice breviter biumbellato, involucri breviter pedicellatis campanulatis multifloris. *Bentham l. c.*

Habitu et foliis *E. latifolio* affine, inflorescentia et floribus ad hanc sectionem referendum. *Petioli basi vaginantes, extus villosissimi, 2-4-pollicares, limbo 1-1½-pollicari. Pedunculus scapiformis ultrapedalis, in exemplaribus cultis folio sæpe instructis. Bractææ sub radiis umbellæ lineares, oblongæ, vel (in exempl. cultis) dilatato-ovatae. Radii umbellæ 1½-3-pollicares, umbellularum vix semipollicares, villosuli. Flores majores quam in E. latifolio. Genitalia minus pilosa quam in præcedentibus. Perianthia post anthesin aucta, laciniæ interiores elongatæ obovatæ, exteriores breviores latiores, marginibus membranaceis crispis.* Bentham MSS.

* "Eriov wool, and γόνυ a joint; in allusion to the woolliness of the stems.

We are indebted to Mr. Bentham for the foregoing extracts from an unpublished paper read before the Linnean Society in May of the present year, and comprising the characters of several new genera, and species of this curious section of Polygoneæ.

The subject of the present article was found by Douglas on the rocky gravelly banks of rivers in New Albion.

It is a hardy herbaceous plant, forming a patch of neat dark green leaves which throw up in abundance, in May and June, their woolly scapes terminated by masses of yellowish white flowers. It thrives equally in peat earth, or common soil, best in a damp situation, and may be increased by cuttings of its well ripened shoots, struck in peat and sand in an almost exhausted hotbed.

We have illustrated the structure of this rare genus in more detail than can usually be afforded in a work so cheap as this. 1. Is a magnified view of a flower as seen externally. 2. Is a section to show the connection of the stamens at their base, and how they are placed by threes in face of the inner segments of the calyx. 3. An ovary with its recurved styles. 4. A fruit enveloped in the permanent calyx. 5. A ripe nut. 6. A seed taken from out of the latter. 7. A section of a seed to shew the position of the embryo in the albumen.



* **CLÍANTHUS** puniceus.*Crimson Glory-pea.*

DIADELPHIA DECANDRIA.

Nat. ord. LEGUMINOSÆ § PAPILIONACEÆ, Juss. (*Introduction to the Natural System of Botany*, p. 86.)

CLIANTHUS, Soland. *Mss. in mus. Britt.*—Calyx latè campanulatus, subæqualis, 5-dentatus. Vexillum acuminatum, reflexum, alis parallelis longius; carina scaphiformis, vexillo alisque multò longior, omninò monopetala. Stamina manifestè perigyna, diadelpa, omnia fertilia. Stylus staminibus duplò longior, versus apicem hinc leviter barbatus, stigmatè simplicissimo. Legumen pedicellatum, coriaceum, acuminatum, ventricosum, polyspermum, intus lanulosum, suturâ dorsali rectâ ventrali convexâ. Semina reniformia, funiculis longiusculis affixa. —Suffrutices, Herbæve foliis impari-pinnatis stipulatis, floribus speciosissimis racemosis.

C. puniceus; suffruticosus diffusus glaber, foliolis alternis oblongis subemarginatis, racemis pendulis multifloris, calyce 5-dentato, legumine glabro.

C. puniceus, Soland. *MSS. in mus. Britt. All. Cunn. in Hort. Trans. vol. 1. n. s. p. 521. t. 22.*

Donia punicea. *Don's gen. syst. of gard. 2. p. 468.*

Suffrutex diffusus, ramosus, 2-3-pedalis, omni pube orbatus nisi levissimâ in paginam inferiorem foliolorum juniorum et inflorescentiæ partes virides; ramis viridibus, parum lignosis. Folia 8-juga cum impare; foliolis oblongis obtusis subemarginatis alternis; stipulis viridibus ovatis, reflexis, foliolis multò minoribus. Racemi penduli, multiflori; axi flexuosâ; bracteis angustè ovatis reflexis pedicellis bracteolatis filiformibus multò brevioribus. Calyx semi-quinquefidus, dentibus acuminatis. Vexillum ovato-lanceolatum, acuminatum reflexum 2-p. longum, atro-sanguineum versus basin albo interrupte vittatum, posticè roseum. Alæ atro-sanguineæ, obtusæ 1½-p. longæ. Carina omninò monopetala, acuminata, fere 3-p. longa, sanguineo-aurantiaca, basi pallida. Legumen ferè 3-p. longum, atro-fuscum, venosum. Semina reniformia, fusca, atro-nebulosa.

The only account that has yet been given of this most beautiful plant is in the last part of the Horticultural Transactions; where a figure is published from a specimen furnished by Mr. Levison Gower from his garden at Titsey, near Godstow. We are indebted to the same gentleman for the specimen from which our plate has been prepared.

It is a native of New Zealand, and, in the opinion of Mr. Allan Cunningham, probably to the southward of the Bay of Islands; and

* From κλειος *glory*, and ανθος *a flower*, in reference to the noble aspect of the species of this beautiful genus.

perhaps, on the shores of the River Thames, or at Mercury Bay. Its seeds were sent by the Missionaries under the name of *Kowaingutukaka*, or *Parrot's-bill*; and were stated to be the produce of a large tree; but this was no doubt a mistake, as there is no reason to believe that the plant will grow more than three or four feet high.

We extract the following account of it from the Horticultural Transactions.

“When planted in a peat border in the open air, where it succeeds the best, it forms a half-herbaceous evergreen shrub, not very unlike an evergreen Vetch, or more correctly speaking, a scarlet *Colutea*, (*Sutherlandia frutescens*). Its leaves are smooth, pinnated, and of rather a succulent texture, consisting of about eight pairs and an odd one. The stem is entirely free from furrows or angles, and is about as thick as a goose's quill. The flowers grow in oval clusters, hanging down from the axils of the leaves upon the lateral branches; each flower is rather more than three inches from the tip of the standard to the tip of the keel; the petals are of a light bright rich crimson, without any mottling or marking; the standard, which is of an ovate-lanceolate figure, and much tapered to the point, is reflexed so as almost to lie back upon the calyx; the wings are very much shorter than the keel, the point of which is so much prolonged as to look like the beak of some bird, although it must be confessed not much like that of a parrot. The flowers are succeeded by brownish black pods, two inches and a half long, seated on a slender stipe, and convex on the upper instead of the lower edge: so that unless attention be paid to their manner of growth, it would seem as if the seeds grew to the lower instead of the upper edge. They are covered all over inside with a delicate cottony down, in which lie the small kidney-shaped seeds, of a dull yellowish-ochre colour, mottled with small dark brown blotches and speckles.

“From the trials that have been made of the proper mode of managing it, both by Mr. Gower and the Rev. John Coleman, by whom it was given to the former gentleman, it would appear that it succeeds best when treated as a hardy plant, and turned out into a peat border; for in such a situation it has now been two years in Mr. Gower's garden, and the plants continue to look very healthy, with a profusion of blossoms forming for next year. Kept in a greenhouse it was sickly, and did not flower in the hands of Mr. Gower's gardener; but Mr. Coleman succeeded in blossoming it in a large pot in the greenhouse, and in inducing it to ripen its pods, one of which is that here figured.

“Considering that the climate of New Zealand is in some places so much like that of England, that some species, such as *Edwardsia microphylla*, will bear the rigour of our winters, it is not improbable that this may also prove a hardy plant. If so its extraordinary beauty will render it one of the most valuable species that has been introduced of late years; and even if it should be no hardier than *Sutherlandia frutescens*, it will still form one of the most important and welcome of all the modern additions to our flower gardens.”

We refer our readers to the observations upon the other species made by Mr. Allan Cunningham in the place whence the foregoing passage has been extracted, for further information concerning these remarkable plants; adding only, that in the first place, the supposed attachment of the seeds to the lower suture of the pod, as described by Mr. Don, is a mistake, their attachment being in *Clianthus* as in all other Leguminosæ, namely, to the upper suture; and in the second, that as we are not aware of any reason why Dr. Solander's well-known name, *Clianthus*, should have been changed to that of *Donia*, we have not felt authorized to sanction the innovation.



* COLLETIA horrida.

Bristling Colletia.

PENTANDRIA MONOGYNIA.

Nat. ord. RHAMNEÆ, Juss. (*Introduction to the Natural System of Botany*, p. 113.)

COLLETIA, Kunth.—*Calyx* campanulatus, membranaceus, coloratus. *Petala* O, v. minutissima, linearia. *Stamina* antheris ovatis bilocularibus v. reniformibus unilocularibus. *Discus* brevis, cupulæformis, fundo calycis adnatus. *Ovarium* liberum, triloculare. *Stylus* simplex, elongatus. *Fructus* basi calycis tubo persistente cinctus, tricoccus, dehiscens. *Semina* sessilia. *Ad. Brong. in Ann. sc. nat.* x. 365.

C. horrida; spinis rigidis simplicibus vel tri-multipartitis, pedunculis sub geminatis, calycibus ovato-oblongis, staminibus sessilibus.

C. horrida. *Ad. Brong. l. c. an. Vent.?*

C. ferox. *Gill. & Hook. in Bot. Misc.* 1. 154. t. 44. f. B.

Frutex uliginosus, atroviridis, rigidus, nudus aut leviter pubescens, aphyllus, vel in ramis vegetioribus parcè foliatus. Rami spinescentes nunc omnino simplices, nunc tripartiti, sæpe multipartiti, validi et rigidi. Folia dum adsunt, minuta, ovata, serrata, atroviridia, glabra. Pedunculi sæpius geminati, nunc aggregati, sub anthesi nutantes, postea rigidi, ascendentes. Calyx ovato-oblongus, laciniis erectis, demum supra discum annularem revolutum transverse separabilis et deciduus. Stamina omninò sessilia. Ovarium superum, triloculare, ovulo unico erecto cuique loculo. Stylus 1. Stigmata 3, subcapitata.

A hardy evergreen shrub, with much the habit of a furze bush, like which it consists chiefly of strong spines of a deep green colour. The flowers, which are of a greenish white, stained with dull purple, are produced in some abundance at the ends of the shoots, and appear in May and June.

Leaves the plant has none when old; but its young branches are furnished with them in the form of opposite

* Named by Commerson after one Mons. Collet, a French Botanist, who is said by Jussieu to have been a fierce opponent of the method of Tournefort.

bright green sawed scales, at the base of each of which is a small stipule. They however speedily fall off, leaving the branches to act as leaves by the aid of their soft parenchyma with which they are clothed in the form of bark.

A native of Chili and Mendoza, where it is probably common. It is by no means so handsome a species as the *C. spinosa* of the same country, whose branches are loaded with masses of white bloom which must have an exceedingly pretty effect.

It grows in common garden soil, and prefers a hot exposed dry situation, such as the foot of a south wall, without any kind of shade. We have not as yet heard of its having been propagated; but it is often raised from Chilian seeds, under the name of *Retanilla*.



* PHOLIDÓTA imbricáta.

Imbricated Pholidota.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆ § MALAXIDÆ. (*Introduction to the Natural System of Botany*, p. 262.)

PHOLIDOTA.—*Supra*, vol. 14. fol. 1213.

P. imbricata; pseudobulbis oblongis corrugato-sulcatis obtusis, foliis solitariis oblongo-lanceolatis plicatis acutis, spicis pendulis foliorum longitudine, bracteis membranaceis concavis acutis, sepalis lateralibus ovatis carinatis, labello subgloboso cucullato: lobis lateralibus parvis erectis intermedio bilobo: laciniis rotundatis.

P. imbricata. *Lindl. in Hook. Exot. Fl. t. 138. Loddiges Bot. Cab. t. 1934.*

It appears that there are two different species confounded under the name of *Ph. imbricata*; namely, this one and that which is figured at fol. 1213 of this work.

So long a time has elapsed since we saw the latter in flower that we are scarcely able to point out any very satisfactory characters to distinguish them, although we have no difficulty in acceding to the opinion of cultivators that they are in fact entirely distinct. We therefore propose to call *Ph. pallida* the plant before figured by the name of *Ph. imbricata*, and to retain the latter designation for this, which was originally figured by Dr. Hooker in the Exotic Flora, and afterwards by Messrs. Loddiges in their Botanical Cabinet.

They are both natives of India, and possibly occupy

* See fol. 1213.

similar regions; they principally differ in the following particulars. *P. pallida* has very round blunt bracts, white flowers, smaller leaves, and grows very weakly and unwillingly under ordinary circumstances. *P. imbricata* has pointed bracts, yellowish flowers with a dash of violet, very long strong leaves, and grows and flowers most freely.

Our specimens were communicated by Richard Harrison, Esq., in May, 1834.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly Aug. 1. 1835.

J. Watts. sc.

* **MYÁNTHUS** barbátus.*Bearded Flywort.*

GYNANDRIA MONANDRIA.

*Nat. ord. ORCHIDÆÆ §. VANDEÆ. (Introduction to the Natural System of Botany, p. 262.)**MYANTHUS.—Supra, vol. 1. fol. 1538.**M. barbatus*; labello in pilis succulentis barbæformibus dissoluto basi supra unicorni.

Caules vetusti fusiformi-cylindræci, 4-5-poll. longi, parum induviati, luteo-olivacei. Folia atroviridia, undulata, Cataseti, oblongo-lanceolata, versus basin angustata. Racemus radicalis, prostratus vel pendulus, flores 9-10 inversos, i. e. labello quoad axin superiore gerens. Bractææ lineari-lanceolatæ membranææ, pedicellis purpurascens breviores. Perianthium bilabiatum; sepalis petalisque lineari-lanceolatis, herbaceis, fusco-sanguineo maculatis. Sepalum supremum, cum petalis in galeam agglutinatum; lateralia patentissima. Labellum cum columna subarticulatum, lineare, medio infractum et saccatum; margine in fila tenuia alba numerosissima dissoluto, cristam aviculæ cujusdam simulantia; cornu deorsum falcatum, candidum, basi superiore tridentatum gerens. Columna herbacea, fusco-sanguineo maculata, cornu apicis galeæ longitudine, cirrhis duabus rectis purpurascens apice pellucidis cornu labelli amplexentibus. Glandula maxima, cartilaginea, oblonga, stigmatæ cum vi elasticâ dissiliens, et convoluta.

This curious new species is a native of Demerara, near the falls of Wapopekai on the Massarony river, some hundreds of miles from its mouth, where it was found by Mr. John Henschman growing in the clefts of trees in shady situations. It was first sent us in February last by Mr. Lowe, with whom it first flowered, but almost immediately after we received it from Chatsworth, and from the garden of Mr. Willmore, of Oldford, near Birmingham. The

* See folio 1721.

first specimens were pallid, as if they had suffered from want of light, but Mr. Willmore's were much darker. In the latter the sepals were of a deep green spotted with purple, the petals were a deep rich brownish red; and the lip itself, with its innumerable thread-like feelers, was tinged with red in all parts, except a strong white horn, which stands upon it near its base, curving downwards towards the little pit which occupies the centre of the lip.

This plant has altogether the habit of *Catasetum*, and no doubt requires the same sort of management.



Miss Drake, del.

Bot by J. Purdyway 169 Piccadilly Aug. 1. 1835.

J. Walto. sc.

* **DENDRÓBIUM** cúpreum.*Copper-coloured Dendrobium.*

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆE §. MALAXIDÆE, Lindl. (Introduction to the Natural System of Botany, p. 262.)

DENDROBIUM.—Supra, vol. 7. fol. 548.

D. cupreum; caulibus teretibus pendulis, foliis oblongis obtusis emarginatis, racemis lateralibus laxis multifloris, bracteis ovato-linearibus canaliculatis obtusis, sepalis ovatis obtusiusculis petalis oblongis explanatis obtusis minoribus, labello unguiculato cochleariformi obtusissimo extus villosus intus nudo.
D. cupreum. Herb. MSS.

A native of the East Indies, whence it was sent by Dr. Wallich to the Hon. and Rev. W. Herbert, about the year 1825. It flowered at Spofforth for the first time at Midsummer 1834, when the drawing was made by Mr. Herbert, which we have now the opportunity of publishing.

It has been suspected that there are two *Dendrobia* in the East Indies, which agree with each other in a sort of tessellated appearance of the flowers, and a remarkably inflated slipper-like hairy labellum; one of these, called *D. moschatum*, has pale orange coloured blossoms, and a lip with 5 crested veins near its base; the other, to which the name of *D. Calceolaria* has been applied by Dr. Hooker, has rose coloured flowers with a little yellow in the centre, and is said to be destitute of the crested veins of the lip. There is, moreover, a *D. clavatum* of Dr. Wallich's Indian Catalogue, the specimens that were distributed of which

* See folio 1239.

were too imperfect to be described in the Genera and Species of Orchideous plants ; but whether this plant is one of the preceding, or something altogether distinct is unknown. Furthermore, the plant now represented belongs to the same little cluster, and appears as if it might form a fourth species, unless indeed it should prove to be this *D. clavatum*. We have had no opportunity of examining it, but Mr. Herbert describes it to us in such a manner as to shew that it must be different from *D. Calceolaria* at least. “ Its flowers are of a pale copper colour, veined with a redder tinge, and have two brown red blotches inside the lip. The spike of ten large flowers all expanded simultaneously, and the progress was so rapid that only about a week or ten days elapsed between its shewing bud and bursting into bloom. Independently of the colour of the flowers this species differs from *D. Calceolaria* in not having such long shoots, and in having the leaves less attenuated and shorter. *D. Calceolaria* under the same treatment makes shoots above four feet long ; this plant under three feet. It is curious that these Dendrobiums if they miss flowering put forth a young plant instead of a spike of flowers at the point of inflorescence, as *D. Calceolaria* is doing here now.”



* *LASTHÉNIA* glabrata.*Smooth Lasthenia.*

SYNGENESIA POLYGAMIA SUPERFLUA.

Nat. ord. COMPOSITÆ. (Introduction to the Natural System of Botany, p. 197.)

LASTHENIA, De Cand. MSS.—Involucrum monophyllum, multidentatum, ebracteatum. *Receptaculum* conicum, scrobiculatum. *Flosculi* radii fœminei ligulati; disci hermaphroditi infundibulares. *Antheræ* apice appendiculatæ, basi muticæ. *Stigmata* rhombeolanceolata ad angulos barbata. *Pappus* O. *Achenia* compressa, lævia, apice disco brevi semicylindræo mucronata.

L. glabrata, glaberrima; foliis subdentatis, involucris turbinatis.



Herba annua, insipida, diffusa, glabrata, imò lucida, foliis oppositis a basi sub-amplexicauli acuminatis, subsucculentis, nunc integerrimis, nunc dentatis. Capitula majora quam in *L. californica*, flosculis magis vitellinis. Ligulæ oblongæ bidentatæ. Flosculi disci valde glandulosi. Achenia atra lævigata.

A new hardy annual, native of California, whence it was introduced by the Horticultural Society in 1834. It flowers in May and June, if sown very early in the spring or in the previous autumn, and forms a pretty gay mass of yellow in the beds of the flower garden. It seeds profusely.

The characters of the genus *Lasthenia* have not yet been published; the name is employed in the manuscripts, of Professor De Candolle for another species,† which is also in our gardens, but which is rather less ornamental than the present.

* An unexplained manuscript name of Professor De Candolle.

† *L. californica* (De Cand.) subpubescens; foliis integerrimis, capitulis basi umbonatis.



* ANGRÆCUM distichum.

Two-rowed Angræcum.

GYNANDRIA MONANDRIA.

*Nat. ord. ORCHIDÆE § VANDEÆ. (Introduction to the Natural System of Botany, p. 262.)**ANGRÆCUM.—Supra, vol. 18. fol. 1522.*

A. distichum ; caule imbricato, foliis distichis compressis recurvis obtusis supra canaliculatis, floribus solitariis axillaribus pedunculis foliis subæqualibus, sepalis ovatis petalisque angustioribus secundis obtusis, labello postico oblongo concavo apice tridentato calcare tereti horizontali pedunculo brevior.

Caules 3-4 pollices longi, atrovirides, foliis densissimè imbricati. Flores parvi, lactei, carnosì, inodori. Labelli dentes laterales rotundati intermedio recto acuto.

A remarkably neat and pretty looking species, on account of its small closely packed deep bright green leaves and tufted stems, but having no beauty of a striking nature in its flowers.

It is a native of Sierra Leone, where it is found on the bark of trees. Imported by Messrs. Loddiges, in whose Nursery our drawing was made in October last. It appears to grow freely in the damp hot atmosphere which so many similar plants delight in.

* See folio 1522.



Miss Drake del.

Pub by J. Ridgway 169 Piccadilly Aug. 1. 1835.

J. Watts sc.

* *DYCKIA rariflora*.*Scattered-flowered Dyckia.*

HEXANDRIA TRIGYNIA.

Nat. ord. BROMELIACEÆ, Juss. (Introduction to the Natural System of Botany, p. 256.)

DYCKIA.—*Schultes fil. Calyx* tripartitus; laciniis erectis concavis. *Corolla* urceolato-campanulata; petalis erectis rhomboideis carnosis basi nudis. *Stamina* basi monadelpha; *antheris* erectis. *Ovarium* superum, tripartibile, polyspermum; *stylis* tribus *stigmatibusque* totidem dilatatis complicatis fimbriatis; *ovula* disticha.—*Folia* carnosâ, epidermide *crassâ corned paululum lepidotâ vestita* margine spinosa.

D. rariflora; foliis lineari-lanceolatis recurvo-patentibus, spicâ rariflorâ, spathis scapi ciliolato-serrulatis, floralibus calycem acutiusculum subæquantibus. *Schultes fil. in Rom. et Sch. Syst. veg. v. 7. p. 1195. Graham in Jameson's Journal, July, 1835, p. 202.*

Introduced from the Berlin Garden by the Horticultural Society in 1833. This plant is a native of the Serra of Villa Rica, in Brazil, where it and two more species were discovered by the indefatigable travellers Spix and Martius. It flowers in June and propagates very slowly by offsets, after the manner of an Aloe, with which it agrees in many of its habits.

* Named in compliment to his Highness the Prince of Salm—Reifferscheid-Dyck, a great lover of Gardening, and one of the most liberal and intelligent of the noble patrons of science of the present day.

The dry stove seems to suit it, for there it produces its rich orange flowers in great perfection, and retains them in all their freshness and beauty for several weeks.

We do not understand upon what principle this genus is referred to the Linnæan *Hexandria Monogynia*, instead of *Trigynia*, for it unquestionably has 3 distinct styles; unless it is to be considered a new case of the necessity of understanding the natural affinities of plants in order to use the sexual system.

Few persons, when they look at the leaves of a plant, ever think of the curious internal mechanism by which all its vital actions are put and maintained in motion; and yet there is not in the whole range of the creation a more singular object than a leaf, nor one whose structure is a more admirable instance of design and forethought. The internal anatomy in this species is highly curious and very easily examined. It consists as usual of a quantity of cellular matter enclosed in a cuticle, but the arrangement of the parts, which is most uncommon, is probably connected with the habits of life of the species in its native wildernesses. The cuticle is hard and composed on the upper surface of three, and on the lower of five layers of extremely minute compact cubical cells. The leaf itself is plane above and convex below; corresponding with the convexity is a stratum of equal thickness of dodecahedral cells, which are green, and pierced towards their upper side by the parallel veins of the leaf; above this structure is a very thick plano-convex bed of hard prismatical cells, which are planted nearly perpendicularly below the cuticle; so that when the section of the leaf is viewed by the naked eye it appears, as is

indicated in our figure, as if it consisted of a common thin channelled leaf, whose concavity is filled up by cartilaginous matter. It is also worthy of remark, that in the lower stratum the tissue below each vein is much more lax and cavernous than that which is interspersed between the veins. The stomates are small and imperfect, and occupy a double or triple line in each furrow, lying concealed among the scurfiness.



* **EMPÉTRUM** rúbrum.*Red Crowberry.*

DICECIA TRIANDRIA.

Nat. ord. EMPETREÆ, Nuttall. (Introduction to the Natural System of Botany, p. 109.)

EMPETRUM, L. ♂ *Squamæ imbricatæ, quarum tres interiores petaloideæ. Stamina 3. ♀ Squamæ maris. Ovarium 6-9-loculare; ovulis solitariis ascendentibus. Stigma sessile, radiis plicatis tot quot loculi. Pericarpium baccatum, seminibus osseis.*

E. rubrum; ramulis foliisque margine lanulosis.

E. rubrum. Willd. sp. pl. 4. 713.

Suffrutex sempervirens, virgatus, procumbens, visu subcinereo. Rami purpurascens, graciles, ramosi, laná laxá intricatá vestiti. Folia linearia, 3 lineas longa, petiolata, marginibus veris reflexis in medio dorsi attingentibus, spuriis laná ciliatis. Flores axillares, solitarii; masculi e squamis constantes imbricatis, circiter novem, quarum 3 interiores corollæ formam induunt et sub aspectu petalorum trium linearium cuneatorum, fusco-purpureorum, apice reflexorum se expandunt. Stamina 3; antheris oblongis 2-locularibus in filamenta debilia vacillantibus. Rudimentum ovarii. Fæminei maribus sunt similes, sed omninò anandri. Ovarium sphericum, 9-loculare. Stigma sessile, fusco-purpureum, radiis novem plicatis stellam referentibus.

A hardy evergreen shrub, lately introduced by Mr. Low of the Clapton Nursery, under the name of "the Cranberry of Staten Island."

It is a plant that is found all over the Southern point of South America, growing with *Pernetia mucronata* along the sandy coast, spreading over the stones, but especially thriving at the back of the low sandhills, by which the shore

* From *εν* in, and *πετρος* a stone, in allusion to the rocky places it grows among.

is often skirted. It is, however, by no means confined to this locality, for we have specimens of it collected at Concepcion by Macrae. According to Gaudichaud the red berries are pleasant to eat.

It grows freely in peat among other plants of a similar kind, and appears perfectly hardy. We have hitherto seen only fertile individuals alive, but in all probability sterile ones exist somewhere in collections, and if so we may soon expect to see its fruit.

Although not striking in its appearance, it forms an interesting addition to the hardy shrubs of this country.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly Aug. 7. 1835.

J. Watts. sc.

* EUTÓCA diváricata.

Stragglng Eutoca.

PENTANDRIA MONOGYNIA.

Nat. ord. HYDROPHYLLÆ, R. Br. (*Introduction to the Natural System of Botany*, p. 244.)

EUTOCA.—*Supra*, fol. 1180.

E. divaricata; caulibus dichotomo-divaricatis, foliis omnibus ovatis indivisis, placentis 12-20-ovulatis. *Bentham in Trans. Linn. soc. vol. 17. p. 278.*

Annua. Caulis decumbens, fragilis, pilosus. Folia indivisa, pilosa, petiolata, oblonga, obtusa. Pedunculi axillares, foliis longiores. Corolla calyce parum longior, subcampanulata limbo plano; plicis decem tubi squamularum loco. Stamina exserta. Ovarium ovatum, hispidum; stylo filiformi hispido, apice furcato. Capsula calyce acuto inclusa, ovata, acuta, compressa, pilis rigidis hispida; unilocularis, bivalvis; valvulis intus nitidis medio placentam corrugatam polyspermam gerentibus. Semina plurima abortiva; matura ovalia, compressa, subtriangularia, scrobiculata; embryo tereti in axi albuminis cornei.

A Californian annual of no great beauty, flowering in May and June. Its seeds should be sown in August in a situation neither overshadowed, nor too much exposed to the sun; the young plants should not be allowed to stand nearer than four or five inches, and they should be kept carefully cleared of weeds; if this is not attended to, the branches will become so weak and brittle that they will not bear the water. Taking these precautions the flowers will be obtained in the spring, before the summer heats set in, and the seeds will ripen just when the season has become too dry to sustain the vegetation of the species.

The light violet blossoms are pretty in bouquets, but are hardly of more than Botanical interest.

* See folio 1180.



* *GESNÉRA* fauciális.*Wide-mouthed Gesnera.*

DIDYNAMIA ANGIOSPERMIA.

Nat. ord. GESNEREÆ, Richard. (Introduction to the Natural System of Botany, p. 227.)

GESNERA.—Suprà, fol. 4. fol. 329.

G. faucialis; herbacea, foliis sub-sessilibus cordatis oblongis acutis crenatis rugosis tomentosis, racemo terminali, bracteis ovatis acutis reflexis, corollis tomentosis labio superiore oblongo bilobo basi angustato inferiore minimo revoluto fauce latissimâ truncatâ.

Communicated in July 1834, by the Hon. and Rev. W. Herbert, who received it from Mr. Tate. It is no doubt a native of Brazil, and is nearly related to *G. Selloi*, from which, however, it differs in many respects, especially in the great depth of the throat of the corolla from back to front, and in the narrow base of the upper lip.

Mr. Herbert considers it by far the finest of the genus, but he can only distinguish it by the size, brilliancy, and paucity of its flowers, from *G. bulbosa*, though it eclipses it. It is equally hardy, and thrives in the greenhouse, if not damp in the winter.

* See folio 1158.



* **ERYTHRONIUM** grandiflorum.*Large American Dog's-tooth Violet.*

HEXANDRIA MONOGYNIA.

*Nat. ord. LILIACEÆ, Juss. (Introduction to the Natural System of Botany, p. 279.)**ERYTHRONIUM, L.—Perianthium hexaphyllum, campanulatum, foliis reflexis: petalis basi bituberculatis. Stamina hypogyna, tria longiora. Stylus trigonus. Stigma 3-lobum v. 3-sulcatum. Capsula polysperma, loculicidotrivalvis. Semina ovata.—Folia radicalia, gemina; flores speciosi; scapi sæpius uniflori, nunc ramosi.**E. grandiflorum; foliis oblongo-lanceolatis subcomplicatis obtusis, perianthii laciniis ovato-lanceolatis acuminatis fere ab imâ basi reflexis, stigmate tripartito.**E. grandiflorum. Pursh. fl. am. sept. 1. 231.**Species Erythroniis alteris americanis affinis sed valdè distincta. Ab americano vulgari et Nuttalliano differt foliis concavis impunctatis et stigmate tripartito; ab albido formâ foliolorum perianthii diversissimâ et stigmate altiùs trifido; ab omnibus floribus duplò majoribus intensiùs luteis, foliolis ab imâ basi ferè reflexis.*

Of this extremely rare plant, a single bulb was received by the Horticultural Society from North West America eight or nine years ago; it has continued to grow slowly in a peat border, and at last put forth its beautiful flowers last May. It has not however produced any seed, and it will probably be many years before it can possibly be distributed.

Mr. Douglas, who discovered it, considered it the *Erythronium grandiflorum* of Pursh; and we adopt his opinion, although we confess ourselves unable to reconcile with it the statement by that writer, that his plant has linear-lanceolate leaves.

* The common Dog's-tooth violet was the *σατύριον ἐρυθρόνιον*, or *red Satyrion*, of Dioscorides, and hence the latter word has been adopted by the moderns.

Its very large flowers, with their segments bent back almost to their base, distinguish this at first sight from the other American species.

Beautiful as it is this cannot be compared with an unpublished species from the same country, which we possess from Mr. Douglas; and which is most remarkable for having *an irregularly branched scape*. This may be named and characterised as follows:—

Erythronium giganteum; foliis oblongis lanceolatisve, scapo irregulariter ramoso 5-floro foliis duplò longiore, foliolis perianthii acuminatis medio reflexis, stigmatè 3-lobò.



* ONCIDIUM pulchellum.

Pretty Oncidium.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆ, Juss. § VANDEÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.)

ONCIDIUM.—Suprà, vol. 13. fol. 1050.

§. 2. Folia triquetra v. teretia.

- O. pulchellum*; foliis acutè triquetris carinatis subfalcatis integris, scapo confertissime multifloro subcernuo, sepalis cymbiformibus acuminatis lateralibus connatis, petalis ovatis undulatis acutis, labelli lobis lateralibus rotundatis intermedio subrotundo sessili retuso v. emarginato subæqualibus, cristâ trilobâ anticè 1-callosâ, alis columnæ acinaciformibus denticulatis.
- O. pulchellum. Hooker in Bot. Mag. t. 2773. Lindl. Gen. & Sp. Orch. p. 206. Lodd. Bot. Cab. t. 1984.*

A beautiful species found in several parts of the West Indies, and probably mistaken for *O. variegatum* till Dr. Hooker distinguished it. When in flower its panicle is so loaded with white blossoms tinged with yellow and pink as to be weighed down with their profusion.

It requires the treatment usually given to its tribe.

* See folio 1542.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly Sept. 1. 1835.

J. Walp. sc.

* *AZARA* dentáta.*Toothed Azara.*

POLYANDRIA MONOGYNIA.

Nat. ord. BIXINEÆ, Kunth. (Introduction to the Natural System of Botany, p. 152.)

AZARA, Ruiz et Pav. Cal. 4-6-partitus. Petala O. Stamina plurima filamentis filiformibus liberis. Antheræ biloculares. Ovarium 1-loculare, placentis tribus polyspermis parietalibus; stigma indivisum. Bacca stylo apiculata, 1-locularis, polysperma. — Frutices Chilenses. Stipulæ nullæ, v. aliquando solitariæ folium parvum mentientes. Flores fragrantés.

A. dentata; ramis pubescentibus, foliis oblongis glabris nitidis crenato-serratis, stipulis rotundis, racemis corymbosis foliis brevioribus, staminibus calyce parum longioribus.

*A. dentata. Ruiz et Pavon. Fl. Peruv. syst. p. 138.**

A common shrub in the woods near Conception in Chile, also frequent near Valparaiso.

In this country it forms a very handsome evergreen bush, with remarkably glossy deep bright green leaves. It appears quite hardy if nailed to a wall and protected from wet in winter, but it will not bear the winters near London under other circumstances. No drought seems to affect it; for at the time we are writing, after nearly two months of the hottest and driest weather known in England, on the face of a south wall, the leaves of this plant are perfectly fresh and green.

It may be increased by cuttings, but layers are preferable.

* Named after Joseph Nicolas Azara, a Spanish Gentleman, of whom nothing further is known, except that he was a patron of science.



* ONCIDIUM Lemonianum.

Sir C. Lemon's Oncidium.

GYNANDRIA MONANDRIA.

*Nat. Ord. ORCHIDÆ, Juss. § VANDEÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.)**ONCIDIUM.—Suprà, vol. 13. fol. 1050.*

§ 2. Folia triquetra v. teretia.

O. Lemonianum ; foliis compressis acuminatis suprà sulcatis, scapo stricto paucifloro, sepalis parvis spatulatis apiculatis omnibus liberis, petalis oblongis undulatis, labelli laciniis lateralibus linearibus abbreviatis intermediâ maximâ reniformi bilobâ : lobis dentatis ungue brevi margine denticulato, columnæ alis subquadratis truncatis oblique unidentatis.

“ This curious little epiphyte was among a collection imported from the Havannah in March last, by Capt. Sutton of Flushing near Falmouth, and by him presented to Sir Charles Lemon, Bart., M. P., in whose garden at Carclew it flowered in May. It appears to us to be totally different from any species hitherto noticed, and is readily distinguished by its peculiar foliage.”

“ *Pseudobulbs* very small, scarcely discernible, roundish, one-leaved. *Leaves* from one to three inches long, and about a quarter of an inch broad, dagger-shaped, two-sided or edged, and sharp at the point ; pale green, slightly striated, and covered with a profusion of minute, roughish, pale dots or warts. *Scape* about 9 inches high, round and slender, nearly erect, and in our specimen 5-flowered. Its colour is a pale brownish green, unless at the joints where it is a

* See folio 1542.

dark brown. *Sepals* yellowish, marked along the back with reddish spots. The upper one which rises erect is spathulate, and appears to have generally 2 spots on the inside behind the column. The two lower ones are small and acuminate. *Labellum* bright yellow, beautifully spotted. The lower division is crescent-shaped, two-lobed, slightly unguiculate and fringed at the outer edge. The upper divisions are panduriform and of the same colour as the lower, but without the spots. The centre is tuberculated and ciliated on each side. *Column* deep yellow, with comparatively large spreading wings."

For the previous note we are indebted to Mr. Booth, under whose successful management this and several other curious new Orchideous plants have been raised at Carclew. The species is nearly allied to *O. Cebolleta*, from which, however, its characters abundantly distinguish it.





* *KENNÉDYA* *Marryattæ*.*Mrs. Marryat's Kennedyya.*

DIADELPHIA DECANDRIA.

*Nat. Ord. LEGUMINOSÆ, Juss. (Introduction to the Natural System of Botany, p. 86.)**KENNÉDYA.*—*Suprà, vol. 11. fol. 944.*§ 1. *Foliis trifoliolatis, carinâ rectâ vexillo sublongiore, De Cand.**K. Marryattæ*; foliolis tribus oblongis obtusis undulatis petiolo brevioribus, junioribus caulibusque villosissimis, stipulis bracteisque cordatis apiculatis, pedunculis 4-floris.

A beautiful greenhouse climber, obtained from Swan River seeds sent home by Sir James Stirling. Our first specimens were communicated by Mrs. Marryat, we afterwards received others from the garden of Mr. Robert Mangles. It flowers abundantly from April to July, producing a striking appearance with its numerous scarlet blossoms.

As it is easily propagated by cuttings it will soon become a common plant.

Different as this is from *K. prostrata* it is not easy to fix upon any very decisive mark of distinction. It is much larger in all its parts, and very much more hairy; in fact its young leaves and stems are perfectly shaggy. The leaflets are larger, far more wavy, and shorter than their stalk. The flowers grow about four in a cluster, instead of two, or being altogether solitary. The fruit we have not seen.

* See folio 1421.



* ARCTOSTÁPHYLOS tomentósa.

Downy Bearberry.

DECANDRIA MONOGYNIA.

Nat. ord. ERICEÆ, Juss. (*Introduction to the Natural System of Botany*, p. 182.)

ARCTOSTAPHYLOS, Adanson. Omnia Arbuti, sed *Drupa lævis*, composita, putamine pluriloculari, loculis monospermis.—Frutices.

A. *tomentosa*; ramis tomentosis sæpius setoso-hispidis, foliis ovatis acutis coriaceis junioribus tomentosis, racemis brevibus compositis, bracteis inferioribus herbaceis linearibus racemosis, fructu depresso.

Arbutus tomentosa. *Pursh. fl. amer. sept.* 1. 282. *Hooker & Arnott in Beech. Voy.* 1. 144.

a. *hispida*; ramis setoso-hispidis tomentosis. *Hooker Fl. Bor. Am.* 2. t. 130. f. 1. *Bot. Mag. t.* 3320.

β. *nuda*; ramis tomentosis. *Hooker l. c. f.* 4.

A curious and very rare hardy evergreen shrub, native of rocky places on the west side of North America, from Puget's Sound in the north to California and the Mexican mountains in the south.

It requires to be cultivated in peat and loam, and in a sheltered situation, when it flowers in March; Dr. Hooker informs us that at Glasgow it is kept in the greenhouse. Our specimens were obligingly communicated from the very select collection of William Harrison, Esq., of Cheshunt, where it has been kept in the open air for about four years, and where alone it has as yet flowered in England.

We quite agree with Dr. Hooker in considering the plant which occurs without any bristly hairs upon its branches as a mere variety of this. But we possess two others, like it in habit, which it is important to distinguish. These are

1. *Arctostaphylos cordifolia*; ramis tomentosis, foliis oblongis obtusis subcordatis coriaceis subtus tomentosis supra nitidis, racemis brevibus compositis, bracteis inferioribus foliaceis oblongis coriaceis racemi longitudine, fructu . . .

On the north-west coast of America, Mr. Menzies.

2. *Arctostaphylos glauca*; glabra, glauca, foliis ovato-oblongis acutis coriaceis basi obtusissimis, racemis brevibus compositis, bracteis inferioribus squamæformibus, fructu ovato.

California, Douglas.

There are no plants more general favourites among Collectors than the species of this Natural Order Ericææ, beginning with Heaths at one end of the series and passing through Andromedas, Vacciniums and Azaleas to Kalmias and Rhododendrons at the other. It is, however, very remarkable, that notwithstanding the extensive commerce of England, the zeal of her merchants, and the enterprize of

* The English name Bear-berry, or Bear-grape, is a literal translation of the Greek words of which this name is composed.

individuals, some of the most magnificent of these plants are still known to Europeans only from the dried specimens in the herbaria of Botanists. We allude to the noble genus *Befaria*, which contains many species more beautiful than even *Rhododendron* and *Azalea*, to the *Thibaudias* with their long tubular crimson blossoms, and to many species of *Gaylussaccia*. The finest of these plants inhabit the Cordilleras of Peru, in the country of the Cinchonas; and certainly if one half the sum that has been sometimes wasted in ill considered undertakings were applied judiciously to an expedition into this region, there would be no reasonable doubt of success, and the results would be indescribably important. We should be glad to explain our views more in detail to any one who might be disposed to engage in such an undertaking, and we cannot help anticipating that the project will be seriously entertained by some of those liberal and wealthy nobles who are indeed becoming the Mæcenates of Natural History in Great Britain. We have been led to these remarks by the examination of a collection of dried specimens lately sent from the Cinchona country by Mr. Mathews, in which we find a larger number of very beautiful plants than we ever have had the good fortune to examine in any collection of similar extent. Among other things there is a most lovely plant which constitutes a new genus related to *Thibaudia*, and which, as it cannot fail before many years to find its way in a living state to England, we have named in compliment to one of the most liberal of those patrons of science whose noble actions render them the pride of their country as well as the surest pillars of their own high hereditary order. The name of Cavendish has long since been enrolled even in the ranks of practical science, but it shines forth in these later days with renewed lustre in the person of his Grace the present Duke of Devonshire, whose noble residence at Chatsworth is rapidly becoming, by the good taste and munificence of its present proprietor, as remarkable for richness in objects of Botanical importance as it has long been admirable for its extent and princely magnificence.

In order that this plant may be fixed in the records of science we add its technical characters, with a brief description of its general appearance.

CAVENDISHIA.

Calyx superus, campanulatus, truncatus, obsolete quadridentatus. Corolla tubuloso-cylindracea, 4-dentata. Stamina 8, inclusa, basi imâ corollæ inserta; filamentis carnosis monadelphis; connectivis alternis hastatis auriculis rotundatis, alternis rhomboidalibus. Antheræ muticæ; loculis sulco alto exaratis, apice liberis et rimâ dehiscentibus. Ovarium quadrangulare, 4-loculare, polyspermum. Stigma simplex—*Frutex Peruvianus, sempervirens; gemmis magnis imbricatis; foliis laurinis; floribus speciosis, purpureis, capitatis, involucreatis.*

1. *Cavendishia nobilis.*

This is apparently an evergreen shrub, with broad laurel-like leaves and buds, which are covered over with imbricated scales, very much like those of a *Camellia*. The flowers are arranged in very close terminal capitate racemes, enclosed within a sort of involucre composed of broad ovate bracts, which are downy on the outside, and coloured red at the edge and in the inside. The corolla is bright crimson and tubular, about an inch long. Sometimes the flowers grow from the side of the old wood, but generally from within involucre at the ends of the branches.



* CALOTRÓPIS procéra.

Tall Calotropis.

PENTANDRIA DIGYNIA.

*Nat. Ord. ASCLEPIADEÆ, R. Br. (Introduction to the Natural System of Botany, p. 210.)**CALOTROPIS.—Suprà, vol. 1. fol. 58.*

C. procera; corollæ laciniis patulis. *R. Br. in Hort. Kew. ed. 2. 2. 78.*
Asclepias procera. Hort. Kew. 1. 305. Willd. sp. pl. 1. 1263.
Asclepias gigantea. Andr. Reposit. t. 271.
Beidelsar. Prosp. Alp. ægypt. cap. 25. c. iv.

“This plant was raised in the garden of Sir Charles Lemon, Bart., M. P., at Carclew, in June 1832, from seeds collected by Lieut. James Sullivan, R. N., at Porto Praya, St. Jago. In the note which accompanied the seed it is described as being “a shrub or small tree growing from 10 to 20 feet high, and flowering in clusters at the ends of the branches. Leaves six inches long. Pod about the size of the leaf, called by the natives *calmady*.” With us it appears to be a tender, lactescent, upright growing, deciduous shrub, requiring the constant heat of the stove, and thriving in a soil composed of sandy loam, and decayed vegetable earth. It flowered in April.

“*Stem* round, pale green, and together with the leaves thickly covered with hoary pubescence which rubs off on being touched. *Leaves* opposite, decussate, spreading, those of the stem obovate, acuminate, about five inches long and two and a half inches broad. At the extremity of the shoots

* Literally “beautifully twisted,” apparently in reference to the corolla of *C. gigantea*.

they are much smaller and more pointed. *Petioles* short, with a deep groove on the upper side. *Flowers* stalked, loosely paniced and spirally arranged, 7, 10, or more in each panicle. *Calyx* 5-parted. *Sepals* very small, stellate, adpressed. *Corolla* divided rather more than half its length, slightly campanulate, and nearly an inch in diameter. When fully expanded it is stellate, and measures about $1\frac{1}{4}$ inch from point to point of the opposite segments, which are somewhat cordate, acuminate, and exceed a quarter of an inch in breadth. Outwardly they are of a pale silvery colour, but inwardly of a deep purplish red, at the point becoming paler and spotted towards the centre of the flower, in which as in others of this Natural Order there is a curious formation, angular at the top, secreting the parts of fructification, and supported by five pale, purplish-tinged, rounded, bracket-like processes. These latter and the divisions of the calyx point in one direction, being placed alternate with the petals and angular part of the centre which point in another. At the extremity of each of the angles is a small aperture, which on being enlarged and having the film removed exhibits two obovate, transparent, greenish yellow bodies, like the pollen masses of Orchideous plants.

“The flowers are produced in succession for several weeks. In the shade, or when the plant is in a room, they are scentless, but in sunshine, or a warm atmosphere, they are highly fragrant.”

Mr. Booth has favoured us with the foregoing account and the drawing that illustrates it. The plant is no doubt the *Beid el sar* of Prosper Alpinus, and appears to extend across the whole of the north of Africa, even into Persia, following it may be supposed the desert region till it stops to the westward of Delhi. Alpinus says its juice, which is extremely acrid, was administered successfully as a remedy for ringworm and other cutaneous affections; and that it is also a powerful agent in removing hair from skin.



Miss Drake del.

Pub by J. Ridgway 163 Piccadilly Oct. 1. 1835.

J. Walter sc.

* **CORYANTHES maculáta.***Spotted Coryanthes.*

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆ, § VANDEÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.)

CORYANTHES, Hooker. Perianthium patentissimum. Sepala dilatata, flexuosa, conduplicata; lateralibus maximis, basi distinctis. Petala multò minora, erecta. Labellum unguiculatum, maximum, galeatum, cum basi columnæ continuum, nullo modo articulatum, tridentatum, in medio unguis appendice poculiformi circumdatum. Columna teres, basi bicornis, elongata, apice recurva, bialata. Stigma rima transversa. Anthera bilocularis. Pollinia 2, compressa, posticè sulcata, caudiculâ lineari arcuatâ, glandulâ lunatâ apicibus approximato-recurvis.—Herbæ epiphytæ, pseudo-bulbosæ. Folia striata. Racemi penduli. Flores maximi. Lindl. Gen. et Sp. Orch. 159.

C. maculata; foliis lato-lanceolatis, sepalis lateralibus semicordatis rectis, labello intùs purpureo maculato cornubus elongatis.

C. maculata. Hooker Bot. Mag. t. 3102. Lindl. l. c.

Sepala membranacea, tenuia, citò flaccida, supremum lanceolatum acuminatum undulatum, lateralia maxima (2½-pollicaria) libera, basi angusta, semicordata, primùm explanata vespertilionis alarum more, dein flaccida. Petala lanceolata undulata flexuosa in labelli galeam prona. Labellum carnosum, 2½ poll. longum, unguiculatum, luteo purpureoque variegatum; hypochilio dilatato lateribus inflexis galeato extus rotundato, in mesochilio angusto canaliculato transeunta; epichilio maximo minus carnosum galeato apice ovato inflexo. Columna libera, basi cornubus duobus elongatis succum stillantibus instructa, apice clavata et recurva marginibus in alasproductis. Stigma rima transversa.

A native of the woods in Demerara, where it is not uncommon hanging from the branches of trees, and suspending in the air the singular lips of its flowers like fairy buckets, as if for the use of the birds and insects that inhabit the surrounding foliage.

* Literally, Helmet-flower; in allusion to the peculiar form of both the hypochilium and epichilium.

There certainly is not a more singular genus than this in the whole vegetable kingdom, nor one whose flowers are less like flowers to the eye of the ordinary observer. The sepals are of the most delicate texture; when young they spread equally round the centre, but after a few hours they collapse, and assume the appearance of a bat's wing half closed. The lip is furnished near its base with a yellow cup, over which hang two horns constantly distilling water into it, and in such abundance as to fill it several times; this cup communicates by a narrow channel formed of the inflated margin of the lip, with the upper end of the latter, and this also is a capacious vessel very much like an old helmet, into which the honey that the cup cannot contain may run over.

C. maculata differs from *C. speciosa*, not only in the colour of the flowers and the greater breadth of its leaves, but also in its lateral sepals being almost exactly half-cordate, without any bend in the middle, and in the horns of the base of its lip being much longer; the middle lobe of its lip too is less distinctly trifid.

C. macrantha, which has lately flowered at Mr. Knight's, surpasses both the others in the richness and magnitude of its flower, and is decidedly distinguished by the mesochilium being irregularly plaited and ridged.

Our drawing of this species was made from a specimen communicated by the Horticultural Society in July 1833.



* IPOMŒA Aitóni.

Mr. Aiton's Ipomœa.

PENTANDRIA MONOGYNIA.

Nat. ord. CONVULVACEÆ, Juss. (Introduction to the Natural System of Botany, p. 218.)

IPOMŒA.—Suprà, vol. 1. fol. 9.

I. *Aitoni*; villosa, foliis cordato-subrotundis, trilobis: lobis acutis lateralibus abbreviatis, pedunculis multifloris, bracteis sepalisque divergentibus acuminatissimis, corollâ subcampanulatâ tubo incrassato, staminibus basi glandulosis.

I. *Aitoni. Hort.*

Radix perennis. Caulis villosissimus. Folia villosovelutina; lobo intermedio subrhombico, acuminato; lamina petioli longitudine. Pedunculi petiolis longiores, multiflori. Flores densè aggregati; bracteis angustis, canaliculatis, acuminatis, recurvis, villosis, sepalorum longitudine. Sepala conformia, corollæ tubo longiora. Corolla violacea, lobis brevibus acutiusculis, tubo incrassato sub staminibus glanduloso. Stamina 5, ultra tubum exserta, quorum duo longiora. Ovarium disco cyathiformi insertum, biloculare; ovulis duobus cuique loculo; stylus simplex; stigma incrassatum bilobum.

A pretty perennial stove climber, not uncommon in collections under the name we have adopted. It flowers from April to October. Its native country is unknown to us.

This strictly belongs to the genus *Ipomœa*, as it has at length been limited by Professor Choisy. The only points in its structure which particularly deserve mention, are the thickened tube of its corolla, and the collection of glands at the base of each stamen, imitating as it were in *Ipomœa* the scale of the genus *Lepistemon*.

Increased by seeds and cuttings, the former of which are produced in some abundance.

The flowers open in the morning.

* From ψ a climbing plant, and $\sigma\mu\omicron\iota\varsigma$ similar; in allusion to the resemblance of this genus to *Convolvulus*.



Miss Drake del.

Pub-try J. Ridgway 169

Piccadilly Oct. 1. 1835.

J. Watts. sc.

* GOVÉNIA supérba.

Superb Govenia.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDEÆ, § VANDEÆ, Lindl. (*Introduction to the Natural System of Botany*, p. 262.)

GOVENIA, Lindl. *Perianthium* bilabiatum. *Sepala* lateralia falcata, labello supposita, basi paululum connata, supremo paulò majora. *Petala* sub sepalo supremo conniventia, breviora, obliqua. *Labellum* integerrimum, ecalcaratum, concavum, cum basi parum productâ columnæ articulatum, sessile. *Columna* basi paulò producta, teres, subfusiformis, apice utrinque marginata. *Anthera* calyptriformis, 1-ocularis. *Pollinia* 4, solida, incumbentia, caudiculâ brevi, glandulâ minore triangulari.—Terrestres. *Folia plicata*. *Spicæ radicales, multifloræ*. *Flores speciosi*. Lindl. gen. et sp. Orch. 153.

G. superba; labello ovato-cordato, spicâ cylindræâ, bracteis acuminatis, foliis oblongis acuminatis basi angustatis scapo æqualibus.

G. superba, Lindl. in *Lodd. Bot. Cab. t. 1709. Gen. & Sp. Orch. l. c.*
Maxillaria superba. *Lexarsa et Llave Orch. Mex. 2. 13.*

Caulis cum foliis 5-pedalis. Folia 3-pedalia, et ultra, basi vaginis purpurascens vestita. Racemus 1-1½-pedalis, cylindræus, floribus onustus. *Sepala* 7 lineas longa, lateralia semi-oblonga. *Petala* obliquè rhomboidea acuminata. *Labellum* 3 lineas tantum longum, lividum, unguiculatum, cordatum, ovatum, obtusum, suprâ canaliculatum.

This noble species was originally discovered by Messrs. Lexarza and La Llave on the mountains near Valladolid, a town of Mexico, flowering in August. The native name is *Azuzena amarilla*. It is one of the handsomest plants of the whole order of Orchideæ, and is especially remarkable for its stately appearance, the rich orange of its flowers, and the long time they continue to open. The specimen in Messrs. Loddiges' collection grows to the height of a man; that from which our drawing was made was furnished by Mr. Bateman, in February 1834.

* So called in compliment to James Robert Gowen, Esq. under whose care were conducted many of the curious experiments upon cross fertilization at Highclere, the seat of the Earl of Carnarvon.

It is found to bear the hottest and dampest stove, but like all the terrestrial species requires a season of repose.

Beautiful as is this plant it must be equalled in the showiness of its blossoms by the *Maxillaria liliacea*, or *Azuzena del Monte* of the people of Valladolid, in the neighbourhood of which town it grows beneath the shade of trees. This plant is described as having a scape about a foot high, terminated by a thryse of snow-white flowers, whose segments are variegated with purple lines, and whose lip is spotted at the base with yellow. It is also in all probability a *Govenia*, and may be named *G. liliacea*.

Of a similar noble aspect is a species collected in Mexico by Count Karwinski, near Atotovilco el Chico, in a cool climate, flowering in June and July. Specimens of this plant, belonging to the Royal Herbarium of Munich, are now before us. The leaves are oblong-lanceolate, tapering to each end, especially to the point, a foot long and about 5 inches wide, and very like those of *Govenia superba*, except that they are smaller. The stem is stated to be five feet high. It is terminated by a short and very dense headed raceme of large flowers, to each of which is an oblong obtuse bract, not quite so long as the ovary. The raceme is 4 inches long, and as much broad at the base. In the form of the flowers it resembles *Govenia superba* very much, but they are three times as large; the sepals are an inch and half long and the other parts in proportion. The lip has an ovate form, but is not cordate at its base, it is not quite half so long as the sepals, and is marked towards its end with largish round very dark dots. The colour of the flowers cannot be ascertained from the dried specimen, but seems to have been pale and uniform; probably pale yellow. This plant may be thus distinguished —

G. capitata; labello ovato obtuso, racemo capitato, bracteis oblongis obtusiusculis, foliis oblongis acuminatis basi angustatis scapo multò brevioribus.



* **MÍMULUS** lúteus ; *var. variegatus*.*Variegated Monkey-flower.*

DIDYNAMIA ANGIOSPERMIA.

Nat. ord. SCROPHULARINEÆ, Juss. (Introduction to the Natural System of Botany, p. 228.)

MIMULUS.—*Supra, vol. 11. fol. 874.*

§. Caule basi sæpius procumbente vel repente, rarius erecto ; foliis 3-5-nerviis dentatis ; calycibus ovatis tubulosive sæpiùs inflatis. *Bentham Scroph. Ind. p. 28.*

M. luteus ; glaber, foliis eroso-dentatis inferioribus longè petiolatis ovatis sublyratisve superioribus rotundatis cordato-amplexicaulibus, calycibus ovatis fructiferis campanulatis dentibus ovatis acutis supremo maximo. *Benth. l. c.*

M. luteus. *Linn. Sp. pl. 884. aliorumque.*

Var. variegatus ; corollæ tubo stramineo intus maculato, limbo purpureo.

M. variegatus. *Lodd. Bot. Cab. t. 1872.*

M. luteus variegatus. *Hooker in Bot. Mag. 3336.*

The first plants of this species which appeared in this country were reared from seeds received from Paris ; but we believe without any intimation of its native country. Since that time Chile has been stated to be the origin of the species, but we know not upon what certain authority ; not a specimen have we ever seen of it from that country. We should almost be inclined to suspect it to be some garden production, if it did not remain true from seed ; at all events we agree with Dr. Hooker in referring it to *M. luteus* as a variety.

It is the prettiest of the genus, and is raised with the greatest facility, provided its seed is sown in a damp place under the shade of a wall, or where it is not fully exposed to

* See folio 1330.

the sun's rays. So treated it grows vigorously and flowers beautifully in May and June, and will come up again in the same place year after year from its self-sown seeds. But if exposed to too bright a sun, and to dryness, it loses its vigorous habit, and becomes a poor dwindling thing not worth cultivation.

It will also strike out freely from cuttings.



* PLEUROTHÁLLIS Gróbyi.

The Groby Pleurothallis.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆ, §. MALAXIDÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.

PLEUROTHALLIS.—Suprà, vol. 9. fol. 759.

P. Grobyi; folio obovato emarginato caule triplo longiore racemo laxo erecto multoties brevior, bracteis minimis membranaceis, sepalis costatis oblongis acutis lateralibus apice tantùm sejunctis, petalis lanceolatis acutis, labello lineari obtuso carnosio superne unisulcato.

P. Grobyi. Bateman in litt.

Folia cum caulibus brevibus secundariis cæspitem efficientibus, atroviridia. Racemi laxi flexuosi. Sepala intus vitellina extus sanguineo vittata. Petala flava medio sanguinea. Labellum petalis paulò longius, sanguineum apice luteum. Columna semiteres, marginata, clinandrio anticè bidentato. Pollinia 2, parva, in unum ferè confluentia, basi (apice) pellucida sed glandulæ nullæ se affigentia.

A native of Demerara, whence it was imported by Mr. Bateman, who named it in compliment to Lord Grey of Groby, of whose skill and ardour in the cultivation of these curious plants we have already offered our own feeble acknowledgment. (See *Grobya Amherstiæ*, fol. 1740.)

It is nearly related to *P. picta* from the same country, and with it belongs to a very pretty set of species, all of which are characterized by having their leaves so much longer than the secondary stems as to conceal them and to form dense tufts.

It is readily cultivated in the Orchideous house; but where the atmosphere is not very damp, will require to be kept under a bell-glass.

Our drawing was made at Messrs. Loddiges in April last.

WE take this opportunity of filling a spare leaf with the characters of some unpublished species of *Pleurothallis*, together with a new arrangement of the genus, and a few remarks upon *Specklinia* and *Physosiphon*.

PLEUROTHALLIS.

§. 1. *Acaules. Flores aggregati v. solitarii.*

Sp. 1. *ruscifolia*; 2. *laurifolia*; 3. *succosa*; 4. *tribuloides*; 5. *cordata*.

6. *P. grandiflora*; folio perfoliato oblongo-lanceolato caudato apice tridentato, pedunculis unifloris fasciculatis, sepalo supremo ventricoso acuminato lateralibus in unum connatis recurvis triplo latiore, petalis lineari-lanceolatis hispido-ciliatis, labello brevioris unguiculato subrotundo carnosio ciliato supra unguem excavato.—In *Peruvia*; supra arbores in convalle Lloæ, alt 8000-ped, *Hall.* (*hab. s. sp. comm. cel. Hooker.*)

7. *P. angustifolia*; caulibus ascendentibus folio lineari-lanceolato longioribus, floribus pedicellatis subsolitariis, sepalis carinatis, petalis lanceolatis denticulatis, labello lineari acuto marginibus pone basin incrassatis.—In *Mexico*, prope Xalapam, *Deppe.* (*h. s. sp.*)

8. *P. bidentata*; caulibus ascendentibus folio oblongo-lanceolato apice bidentato longioribus, floribus pedicellatis subsolitariis, sepalis ecarinatis, labello lineari obtuso apice ciliato.—Prope *Rio Janeiro*, *Forbes.* (*h. s. sp.*)

9. *P. papillosa*; caulibus strictis folio angustè oblongo apice tridentato paulò longioribus, pedunculis subaggregatis unifloris flore vix longioribus, sepalis campanulatis lateralibus semiconnatis supremo oblongo brevioribus, labello subhastato obtusissimo apice vittisq. duabus elevatis carnosis papillosis.—In *Brazilia* prov. *Minarum* in arboribus ad *S. João Baptista*, *Martius* (*ex. s. sp. in hb. M.*) Flores majusculi.

10. *P. discoidea*; caulibus strictis folio ovato-lanceolato brevioribus, pedunculis solitariis unifloris, sepalis oblongis acutis lateralibus omnino connatis, petalis linearibus acuminatis decurvis, labello cordato-ovato undulato obtusiusculo sepalis subæquali.—In insula *Trinitatis* (*v. ic. pict. in hb. Hooker.*) Facies *P. ruscifoliae*. Flores flavi maculâ sanguineâ oblongâ anticè bidentatâ in medio labelli.

11. *P. punctata*; caulibus strictis folio crasso ovali plano-convexo marginato longioribus, floribus solitariis breviter pedicellatis, petalis lanceolatis apice pubescentibus, labello unguiculato cordato-sagittato obtuso, columna elongata.—In *Brazilia* (*exam. v. c.*). Folia subtus purpureo punctata. Flores atropurpurei. Petala herbacea. Tota planta vix $1\frac{1}{4}$ poll. alta.

§. 2. *Acaules. Flores racemosi.*

a. *Folia caulibus secundariis longiora (Cæspitosæ).*

12. *alpestris*; 13? *laxa*; 14. *Grobyi*.

15. *P. picta*; folio spathulato marginato retuso racemis laxis duplò brevioribus, bracteis minimis, sepalis acuminatis lateralibus apice tantum sejunctis, petalis lineari-lanceolatis acutis, labello lineari obtuso carnosissimo supra 1-sulcato.—In *Demerara, Loddiges (exam. v. c.)*.

16. *P. hymenantha*; folio lineari-lanceolato apice obsolete 3-dentato racemis flexuosis capillaribus subæquali, bracteis minimis, sepalis acutis lateralibus semi-sejunctis, petalis acuminatis parum brevioribus, labello oblongo obtuso subundulato complicato.—In *Brazilia*; in arboribus vetustis prope Retiro provinciæ Sebastianop. *Martius. (exam. s. sp. in hb. M.)*

17. *P. ochreata*; folio crasso lanceolato complicato mucrone acuto racemo denso flexuoso nutante duplò longiore, bracteis cucullatis imbricatis acutis, sepalis carinatis acuminatis lateralibus semiconnatis, petalis lineari-oblongis nanis mucronatis, labello —In *Brazilia*; in rupibus Serra do Monte Santo provinciæ Bahiensis, *Martius. (exam. s. sp. in hb. M.)* Calyx ruber v. miniatus. Petala duplo minora flava, omnia subæquali. *Mart. mss.*

18. *P. rupestris*; folio tereti mucronato anticè sulcato racemo stricto rigido paulò brevioribus, bracteis parvis rigidis concavis, sepalis acutis lateralibus semiconnatis, petalis lanceolatis ciliatis, labello subtrilobo membranaceo obtuso appendice magno carnosissimo bifurco: cruribus parallelis porrectis acutis.—In *Brazilia*, inter rupes M. Itacolumi et Moro de V^a. R^a. provinciæ Minarum, *Martius. (exam. s. sp. in hb. M.)* Scapus rubenti-virens. Sepala purpurascens. Petala purpurea. *Mart. mss.*

19. *P. teres*; folio tereti racemo gracili brevioribus, bracteis membranaceis ochreatis, floribus pendulis, sepalis acuminatis lateralibus apice tantum liberis, petalis minimis oblongis acutis, labello lineari canaliculato acuto marginibus incrassatis, columnâ basi pubescente.—In *Brazilia, Loddiges (exam. v. c.)*. Flores cinnamomei. Folia omnino *Leptotis bicoloris*.

20. *P. hians*; caule brevi, folio oblongo basi angustato obtuso racemo flexuoso 6-9-floro duplò brevioribus, bracteis minutis ochreatis, floribus bilabiatis: labio postico 2-dentato antico oblongo obtuso, petalis cuneatis, labello ovato acuto complicato.—In *Brazilia*, in montibus Organ (*v. ic. pict. et exam. fl. s. in herb. Hooker.*) Caulis pollicaris. Folium 3-pollicare. Scapus fusco purpureus. Flores ejusdem coloris, aciculatim maculati, extus pallidè cinnamomei. Sepalum supremum anticum, lateralibus connatis posticis æquilatum. Petala labellumq. nana atropurpurea. Columna ejusdem coloris clinandrii margine membranaceo latissimo.

21. *spiralis*. 22. *racemiflora*.

b. *Folia caulibus secundariis breviora v. æqualia. (Macropodæ.)*

23. *prolifera*; 24. *saurocephala*; 25. *quadrifida* (= *Dendrobium quadrifidum. La Llave*); 26. *pulchella*; 27. *Lanceana*.

28. *P. Jamiesoni*; folio lineari-oblongo carnosissimo submucronato racemo secundo stricto duplò brevioribus, bracteis densis brevibus ochreatis cucullatis, sepalis

oblongis subæqualibus leviter carinatis lateralibus basi tantum connatis, petalis duplo brevioribus apice rotundatis, labello brevioris membranaceo rhombeo-trilobo plicâ unguis transversâ.—In *Peruvia*, in præcipitiis Pichinchæ haud procul a Quito, *Jamieson*. (*exam. s. sp. in hb. Hooker.*) *P. pulchellæ* valdè similis, sed minor.

29. *P. capillaris*; folio lineari-lanceolato racemis capillaribus fasciculatis subæquali, sepalis acuminatis lateralibus basi connatis, petalis ferè æquilongis serratis acuminatis, labello lanceolato basi angustato acuto membranaceo 3-venio.—In *Brazilia*, in arborum truncis provinciæ Sebastianop. *Martius*. (*exam. s. sp. in hb. M.*) Facies *P. ruscifoliæ*.

30. lanceolata.

31. *P. sclerophylla*; folio oblongo petiolato obtuso carnosissimo racemis multò brevioribus, bracteis ovatis cucullatis, sepalis acuminatissimis omnibus ferè liberis, petalis nanis obtusis, labello oblongo conduplicato apice piloso-glanduloso.—In *Brazilia*; in Serra de Piedade, prov. Minarum, etiam in rupibus campestribus prov. S. Pauli, *Martius*. (*exam. s. sp. in hb. M.*)

32. *P. rubens*; folio oblongo-lanceolato obtuso, racemis longissimis, bracteis oblongis ochreatis, sepalis lanceolatis acuminatis lateralibus basi tantum connatis petalis obtusis, labello obovato repando obtuso lineis 2 elevatis flexuosis carnosissimis.—In *Brazilia*; in rupibus ad Opp. S. Joao d' El Rey, *Martius*. (*exam. s. sp. in hb. M.*)

§. 3. *Caulescentes*; i. e. rhizomate nullo sed caule folioso.

33. macrorhiza. 34. caulescens.

Under the name of *Stelis tubata*, Messrs. Loddiges have published a very curious plant with the habit of *Pleurothallis*, but with its sepals all united into a tube, which is inflated at the base and contracted at the mouth. Otherwise its fructification has the structure of *Stelis*. I propose to distinguish this as a peculiar genus, with the following name and character.

PHYSOSIPHON.

Calyx tubulosus basi ventricosus, apice 3-fidus. Petala in fundo calycis carnosissima, nana. Labellum et columna Stelidis. Pollinia 2, spherica.—Herbæ epiphytæ habitu *Pleurothallidis*.

1. *P. Loddigesii*; (= *Stelis tubata*, *Lodd. Bot. Cab. t. 1601.*)

2. *P. emarginata*; (= *Pleurothallis emarginata*, *Lindl. Gen. et. Sp. Orch. Pl. p. 6.*)

3. *P. spiralis*; folio angustè oblongo subemarginato carnosissimo spicâ spirali confertâ duplò brevioribus, bracteis minimis ovatis acutis cucullatis, calyce semitripartito laciniis erectis acutis, petalis cuneatis carnosissimis, labello ovato-subrotundo membranaceo apiculato trivenio.—In *Brazilia*, ad Portum Sæ. Catharinæ, *Tweedie*. (*hab. s. sp. comm. Cel. Hooker.*)

SPECKLINIA.

To this genus is to be added the following:—

S. atropurpurea; vaginis caulis 2-3 ventricosis, folio oblongo plano 7-9-venio, flore solitario subsessili, petalis 3-cuspidatis, labelli laminâ sagittatâ.—
In *Jamaica*. (*Exam. s. cult. in herb. Hooker. ex horto Liverp.*) Folium 2½ poll. longum, basi angustatum. Vaginæ infundibulares, pollicem longæ, in caulem 2-pollicarem. Flores atropurpurei. Ovarium lineam longum, hexapterum; alis apice truncatis petalinis minoribus. Alabastrus quem tantum vidi semunciam longus, angustus, falcatus. Sepala anteriora basi gibbosa, oblongo-linearia. Petala nana, vix 2 lineas longa, ovata, 3-cuspidata, obliqua, sæpiùs venâ alterâ excurrente, alterâ in dimidio majore petali abbreviatâ. Labellum unguiculatum, in columnam pronum, cum ejus basi articulatum; ungue lineari, 3-venio axi cristato, laminâ sagittatâ apicibus posticis obtusis. Columna petaloideo-alata, apice tridentata, basi paulò producta; pèdis marginibus carnosus rotundatis medio unidentatis. Anthera parva 1-ocularis. Pollinia 2, pyriformia, atropurpurea, apice materie diaphanâ granulosâ stigmati adhærentia.

And probably several spurious *Dendrobia* of authors, especially—

<i>S. retusa</i>	=	<i>Dendr. retusum</i> , <i>La Llave</i> .
<i>S. scariosa</i>	=	<i>Dendr. scariosum</i> , <i>Do.</i>
<i>S. pusilla</i>	=	<i>Dendr. pusillum</i> , <i>Humb. et Kunth.</i>
<i>S. acuminata</i>	=	<i>Dendr. acuminatum</i> , <i>Do.</i>
<i>S. elegans</i>	=	<i>Dendr. elegans</i> , <i>Do.</i>



* EDWÁRDSIA chilénsis.

Chilian Mayu Tree.

DECANDRIA MONOGYNIA.

Nat. ord. LEGUMINOSÆ, Juss. § PAPILIONACEÆ. (Introduction to the Natural System of Botany, p. 87.)

EDWARDSIA.—Suprà, vol. 9. fol. 738.

E. chilensis; foliolis 13-19 elliptico-oblongis obtusis rigidis subtus sericeis, petalo supremo lateralium longitudine, legumine binodoso aptero.

Edwardsia chilensis. Miers Trav. in Chili; the name only.

Sophora macrocarpa. Smith in Rees. De Cand. Prodr. 2. 96. Hooker et Arnott in Bot. Misc. 3. 177.

A fine tree, native of Chile, where the inhabitants call it *Mayu*. It was introduced by Messrs. Loddiges in 1822, having been sent them by General Paroissien under the name of "*Sophora Myrospermum*."

It proves to be a hardy handsome tree, flowering in May in great profusion. We believe that it at present exists in no other collection.

This is plainly an *Edwardsia*, notwithstanding the opinion of the late Sir James Smith to the contrary, and notwithstanding the alleged want of wings to its pods. Supposing that fact to be as stated, it only shews that the winged pod is unessential to the genus *Edwardsia*, and not that the genus itself is a bad one; for surely if there be any one genus on which Nature has set her mark more distinctly than another it is this. Mistakes like those of Smith are the certain accompaniments of the meagre system of artificial Botany.

* See folio 738.

It is the short loose axillary racemes, the hard cup-shaped calyx, slit on the upper side, and the broad vexillum which overlaps the other petals and is usually absolutely parallel with them, that distinguish *Edwardsia* from *Sophora*, whose racemes are terminal, calyx soft and regularly toothed, and vexillum narrow and recurved, or reflexed.



* MAXILLÁRIA crócea.

Saffron-coloured Maxillaria.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆ, § VANDEÆ, Lindl. (*Introduction to the Natural System of Botany*, p. 262.)

MAXILLARIA.—*Suprà*, vol. 11. fol. 897.

§. 2. *Scapigeræ*. Pedunculi radicales. Labellum anticum. Scapi uniflori.

M. crocea; pseudobulbis oblongis compressis foliatis, foliis oblongis undulatis obtusis emarginatis latè vaginantibus, scapis erectis unifloris laxè vaginatis foliis duplo brevioribus, sepalis petalisque angustè triangularibus apice elongatis abruptè acutis, labello obovato obtuso levissimè trilobo apice carnosocrispo supra medium unituberculato.

We have received the drawing and following account of this plant from Mr. Booth.

“ We owe the introduction of this curious plant to Capt. Sutton, of his Majesty’s Packet establishment at Falmouth, who imported it from Rio Janeiro in the spring of 1833, and presented it, with others, to Sir C. Lemon, Bart., M.P., in whose collection at Carclew, Cornwall, the subject of the present notice produced its flowers in August 1834.

“ *Root* fibrous. *Pseudo-bulbs* small, oblong, compressed, nearly covered by the three sheathing, strap-shaped leaves, which are each thick and rigid, from four to six inches long, and an inch in width, a little twisted, with an oblique sharp point. The leaf on the bulb is similar to the others but much longer. All of them are of a deep green above, and a pale silvery green beneath. *Scapes* four inches high, one-flowered, round and slender, rising from the base of the leaves, and covered with three or four, thin, pale green,

* See folio 1428.

acuminate bracts, forming a joint at their junction with the stem. *Flowers* before expansion having some resemblance to the beak of a bird, pale green with a brownish tinge at the spur. *Perianth* of a greenish yellow, becoming of an orange colour when expanded. *Sepals* narrow and pointed, the three exterior ones being each about the same size and length, with their edges turned back. The two interior ones are rather shorter than the others, somewhat curved, and crossing so as almost to conceal the column. *Labellum* about half the length of the sepals, thick and fleshy, connected to the lengthened base of the column, which forms the spur; cucullate and crisp at the margin, which is rather paler than the sepals, otherwise it is of a brownish red colour, together with the column.

“It is obviously allied to *M. picta* and *punctata*, from both which the character assigned to it, abundantly distinguish it.

“Like most of the terrestrial Orchideous plants from Tropical countries, it requires the constant heat of the stove, and to be kept rather moist. It seems to flourish in the soil we find in the hollow decayed trunks of trees, mixed with some finely chopped moss; and promises to increase freely by off-sets.”



* STANHÓPEA oculáta.

Eyed Stanhopea.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆ, § VANDEÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.)

STANHOPEA.—*Suprà, vol. 18. fol. 1529.*

S. oculata; labello medio constricto, hypochilii elongati lateribus cornutis erectis basi bimaculatis, epichilio ovato acuminato cornubus duobus baseos incurvis.

Lindl. gen. et sp. Orch. Pl. p. 158.

Ceratochilus oculatus. Lodd. Bot. Cab. t. 1764.

Imported by Messrs. Loddiges. It is the most interesting of this splendid genus, on account of the extremely delicate waxy appearance of its surface, the softness of its ground colour, and the richness of the deep purple spots, which lying upon a bright yellow field so very conspicuously ornament the base of the lip and the petals.

Our drawing was taken from a most noble specimen produced in July 1834, in the stove of Mr. Bateman. The leaves, including the petioles, were more than two feet long, and being of the darkest green had a most imposing appearance. Mr. Bateman informs us that he finds it requires eight such leaves to form a flowering plant; and that those who wish to succeed in blossoming it must suffer it to remain a long time undisturbed, since a good many pseudo-bulbs are required to furnish one really fine specimen.

The singular form of the labellum of this and some other Orchideous plants has rendered it necessary to distinguish it into three parts for the convenience of description.

* See folio 1529.

The lowest part of the lip is the *hypochilium*, the upper end the *epichilium*, and the intermediate portion the *mesochilium*. All these parts are, however, nothing more than divisions in a very fleshy lip, produced by contractions of its sides, and by the lobes, so common in the order, variously arranged and combined.

The species is at once known by the narrowness and length of the hypochilium, independently of all other circumstances.

It is said to be a native of Brazil; but we suspect some mistake in this statement. It is certainly wild in Mexico, for it exists among Count Karwinski's plants in the Royal Munich Herbarium.



* **PRUNUS japonica.***The Single Chinese Plum.*

 ICOSANDRIA MONOGYNIA.

Nat. ord. ROSACEÆ, § AMYGDALÆÆ, Juss. (Introduction to the Natural System of Botany, p. 81.)

PRUNUS.—*Suprà, vol. 2. fol. 136.*

P. japonica. Suprà, vol. 1. fol. 27. The double state.

It is always interesting to procure the wild forms of cultivated species, and so see from what humble originals Nature produces some of the most striking of her works.

The double Chinese Plum, or Almond, as it is often incorrectly called, is beyond all comparison the handsomest plant of its season; there is nothing to vie with its crowded clusters of most delicate blushing flowers, the petals of which are loosely, but symmetrically, arranged into the most perfect of vegetable beauties. The simple shrub now represented is its origin, and is one of the many examples of the creation by the patient Chinese of the fairest ornaments of the garden, from the most inconspicuous plants of the woods.

For its introduction the public is indebted to John Reeves, Esq., by whom so large a proportion of all the fine Chinese flowers now common all over Europe have been procured for this country.

It appears to be a hardy shrub; our specimen, however, was taken in January last, from a plant which had flowered in a greenhouse.

The double state of the species is represented in the first volume of this work.



* *MAXILLÁRIA picta.**Painted Maxillaria.*

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆ, § VANDEÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.)

MAXILLARIA.—Suprà, vol. 11. fol. 897.

- §. 2. *Scapigeræ.* Pedunculi radicales. Labellum anticum. Scapi uniflori.
M. picta; pseudobulbis ovatis subsulcatis 1-2-phyllis, foliis oblongo-lanceolatis planis, scapis erectis unifloris distanter vaginatis foliis duplò brevioribus, sepalis petalisque lanceolatis (abruptè acutis) sub-conniventibus, labello trilobo medio unituberculato lobis oblongis obtusis lateralibus brevioribus.
M. picta. *Hooker in Bot. Mag. t. 3154. Lindl. gen. & sp. Orch. p. 146.*

A truly beautiful plant found wild in the Organ mountains of Brazil, and originally introduced by the late lamented Mrs. Arnold Harrison. When well managed it throws up a profusion of its gay speckled flowers, which remain perfect for some time.

Our drawing was made in the stove of the Messrs. Loddiges in November 1834.

There are two other unpublished species of *Maxillaria*, belonging to this section, which we take this opportunity of naming.

1. *M. rufescens*; pseudobulbis ovatis subtetragonis monophyllis, foliis lanceolatis utrinque acuminatis, scapis unifloris prostratis vaginis minimis distantibus, sepalis petalisque oblongis conformibus obtusis, labello oblongo trilobo etuberculato laciniis lateralibus minimis acutis intermediâ elongatâ emarginatâ.

A native of Trinidad, whence it was imported by Mr. Lowe of Clapton. Our description is taken from a drawing and flower transmitted from Chatsworth in December 1834.

* See folio 1428.

t. 1802. Vitch refers to this number as an illustration of *M. rufescens*.

It is a species of no beauty. The sepals are a dull greenish purple, the petals and lip yellow, the latter speckled with dull purplish red. Possibly the specimen was bleached, and the colours would become brighter if the plant had flowered in brighter weather. Stands next *M. picta*.

2. *M. graminea* ; pseudobulbis nullis, foliis gramineis recurvis, perianthio campanulato, sepalis oblongis obtusis lateralibus paululum connatis, petalis angustioribus, labello cum pede longè producto columnæ articulato trilobo : lobis lateralibus semi-ovatis acutis ascendentibus intermedio majore reniformi, tuberculo disci magno carnosio truncato posticè bilobo.

A native of Demerara; imported by Mr. Lowe; flowered in Messrs. Loddiges' collection, January 1835. A small and inconspicuous species. The blossoms are pale yellow, with the segments a little banded with purple near the base. Not very nearly related to any known species.

1803.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly Oct. 1. 1835.

J. Watts. sc.

* **CHIRÓNIA** pedunculáris.*Long-stalked Chironia.*

PENTANDRIA MONOGYNIA.

*Nat. Ord. GENTIANEÆ, Juss. (Introduction to the Natural System of Botany, p. 215.)**CHIRONIA.—Suprà, vol. 3. fol. 197.*

C. peduncularis; glabra, foliis ovato-lanceolatis acuminatis sessilibus 3-5-veniis internodiis subbrevioribus, pedunculis unifloris foliis longioribus, calyce tubo corollæ brevioris, caule tereti.

C. trinervis. Hort. nec Linn.

Frutex diffusus, 3-4-pedalis, atroviridis, glaber, succo amarissimo sublacteo scatens. Internodia basi purpurascens. Folia 3-5-venia, 2-poll. longa, sæpe internodiis breviora. Pedunculi axillares et terminales, foliis duplò longiores, uniflori. Calyx 5-partitus, laciniis linearibus acuminatis, tubo corollæ brevioribus. Corolla persistens, tubo gracili viridi semunciali, limbo patente 5-partito, purpureo, ejusdem longitudinis; laciniis ovatis acutis post anthesin convolutis. Stamina 5, fauce tubi inserta, a stylo aversa, exserta; antheris linearibus poro duplici apicis dehiscentibus, defloratis rectis. Ovarium teres, acuminatum, uniloculare marginibus carpellorum 4 inflexis ovuliferis; stylo continuo; stigmatibus dilatato bilobo. Fructus semimaturus carnosus, amarissimus.

A very pretty greenhouse shrub, covered with a succession of purple flowers from July to October, and propagated freely by cuttings.

Nothing can be easier than its management, as it grows in any kind of soil, will thrive out of doors in summer, and will survive the winter without injury in a very indifferent greenhouse.

* Named in honour of Chiron the Centaur, one of the earliest medical practitioners whose names have been preserved. The intense bitterness of the genus indicates its powerful tonic properties.

We are unacquainted with its history; it does not seem to have been before described; and we have not been able to learn even its native country. In gardens it is called *Chironia trinervis*, but it is a totally different plant from the Ceylon species so named by Linnæus, which appears to be a genuine *Exacum*, as that genus is limited by Dr. Brown in his *Prodromus*.

The bitterness of the species is most remarkable, even among its bitter neighbours.



* MAXILLARIA densa.

Dense-flowered Maxillaria.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆ, § VANDEÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.)

MAXILLARIA.—Suprà, vol. 11. fol. 897.

§. 1. *Axillifloræ.* Pedunculi axillares.

M. densa; pseudobulbis oblongis compressis axillaribus monophyllis, foliis oblongo-lanceolatis obtusis emarginatis, racemis axillaribus densissimè aggregatis, bracteis cucullatis, perianthiis bilabiatis, sepalis lineari-lanceolatis acuminatis carinatis, petalis paulo minoribus, labello oblongo indiviso apice recurvo et canaliculato, medio lineâ transversâ elevatâ.

A native of Mexico, whence it was imported by the Messrs. Loddiges, in whose stove it flowered in January last.

It belongs to that set of Maxillarias on which the genus was founded by the authors of the Flora Peruviana, and which in some respects appear to differ from those with which we are but acquainted in Gardens. The stems rise some height above the ground, and are closely invested with brown withered scales, from the axils of which spring the pseudo-bulbs and flowers; in most Maxillarias on the contrary, there seems to be no stem, but the pseudo-bulbs sit close upon the ground, as in *M. picta* and *ochroleuca*. In truth, however, there is a stem in both kinds; only it is erect and above ground in the one, and prostrate or underground in the other. And so it is with all the pseudo-bulbous Orchideæ; their true stem is in most cases a rhizoma, like that of Iris, rooting on the side that touches the ground, and pushing up leaves and pseudo-bulbs on the other.

* See folio 1428.



* FÚCHSIA díscolor.

Port Famine Fuchsia.

OCTANDRIA MONOGYNIA.

*Nat. ord. ONAGRACEÆ, Juss. (Introduction to the Natural System of Botany, p. 56.)**FUCHSIA.—Suprà, vol. 10. fol. 847.*

F. discolor; dumosa, ramosissima, vinosa, ramulis levissimè pubescentibus, foliis ovatis denticulatis undulatis petiolatis oppositis ternatisque, pedunculis foliis longioribus, petalis obtusis convolutis calycis laciniis acuminatis brevioribus, staminibus longè exsertis.

A native of Port Famine in the Falkland Islands, whence it was some years since introduced by Mr. Lowe. Our drawing was made from a plant in the Garden of the Horticultural Society in August last. It has the same season of flowering as the other Fuchsias, and may be increased readily by cuttings in like manner.

If we are asked to state in what respects this differs botanically from *F. gracilis* and *tenella*, we should find it very difficult to answer the question. And yet it is a decidedly different plant; remarkable for its compact, bushy manner of growth, its deep purple branches, its small very undulated leaves, and also for its being apparently more hardy than any other Fuchsia yet in the Gardens. For the latter reason we attach especial importance to it, for by a judicious intermixture of its pollen with such beautiful plants as *F. conica*, *globosa*, and its other more tender relatives, the whole race may probably be rendered capable of bearing

* See folio 1269.

the climate of Great Britain, and may thus become far more generally valuable than they yet are.

It is probable that those are right who consider the greater part of the Chilean Fuchsias mere varieties of *F. macrostema*, and if so this will have to be added to the list as a dwarf contracted kind. *F. globosa* and *conica* are, however, to all appearance distinct species.



1.

2.

FERNANDÉZIA acúta.

* *Sharp-leaved Fernandezia.*

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDÆ, §. VANDEÆ, Lindl. (Introduction to the Natural System of Botany. p. 262.)

FERNANDEZIA, Fl. Peruv. Perianthium patens. Sepala libera. Petala conformia, sub sepalo supremo conniventia. Labellum anticum v. posticum, ecalcaratum, liberum, trilobum, disco tuberculatum. Columna brevis, utrinque aurita. Anthera bilocularis. Pollinia 2, solida, obovata, glandulá caudiculá —Epiphytæ caulescentes. Folia disticha, equitantia, imbricata. Racemi pauciflori, terminales v. laterales. Flores parvi lutei. Lindl. Gen. et Sp. Orch. p. 207.

F. acuta; foliis acuminatis carinatis, corymbo laxo multifloro, bracteis obtusis membranaceis, labello lineari 3-lobo laciniis lateralibus rectis abbreviatis intermediâ emarginatâ obtusâ, disco pulvinateo tuberculo minimo pone basin.

Omnia fere F. elegantis nisi folia et labellum. Hoc pallidè luteum disco prominulo pulvinateo anticè bilobo sanguineo-marginato; laciniis lateralibus nullo modo hastæ effigiem referentibus sed laciniâ intermediâ parallelis.

A native of Trinidad, whence it was imported by Mr. Knight, in whose collection it flowered in June last.

In habit is very like *F. elegans*, from which it is principally distinguished by the tapering form of its leaves, and the shape of its lip, which, instead of having a sort of hastate form, in consequence of the divergence of its lateral lobes, and a number of irregular unequal tubercles upon its disk, has the lateral lobes parallel with the middle one, only a single small tubercle at its base, and a slightly elevated disk which is 2-lobed, and bordered with crimson in front.

A hot damp stove is necessary to keep this rare species in health.

* So called after George Garcias Fernandez, a Spanish Botanist, who is unknown, except through the medium of this genus.



Miss Drake del.

Publ. by J. Peasway 769 Broadway Nov. 1. 1835.

J. Wallis sc.

* *CÉREUS* triangúlaris. = *HYLOCEREUS*
undatus

Triangular Torch-thistle.

ICOSANDRIA MONOGYNIA.

Nat. ord. CACTACEÆ. (Introduction to the Natural System of Botany, p. 54.)

CEREUS, D. C. Sepala numerosissima, imbricata, basi ovario adnata, in tubum elongatum concreta, exteriora breviora calycinalia, media longiora colorata, intima petaliformia. Stamina numerosissima cum tubo concreta. Stylus filiformis apice multifidus. Bacca sepalorum reliquiis areolata, tuberculata aut squamata. Cotyledones nullæ? (depauperatæ).—Frutices carnosì, elongati, axi ligneo internè medullifero donati, angulis verticalibus spinarum fasciculos gerentibus regulariter sulcati. Anguli seu alæ nunc plurimæ, nunc paucissimæ, rariùs duæ tantum et tunc rami compresso-alati. Flores ampli è spinarum fasciculis aut crenis angulorum orti. De Cand. Prodr. 3. 463.

§. 2. *Serpentini.* Caulis articulatus prostratus radicans vel volubilis. D. C.
 *** 3-4-angulares.

C. triangularis; repens trigonus, aculeis brevibus quaternis decussatis, sepalis exterioribus spatulatis apice foliaceis, petalis lanceolatis cuspidatis.

Cactus triangularis. *Linn. sp. pl.* 666. *Jacq. amer.* 152.

Cereus triangularis. *Haworth. syn.* 180. *De Cand. prodr. l. c.*

A native of Mexico and the West India Islands, whence it was introduced long since, but it flowers so rarely that it has never yet been represented from an European specimen. Bradley, who has given it in his work on succulent plants, has only figured its stem.

For the specimen from which our drawing was made we are indebted to Sir George Staunton, Bart., in whose garden at Leigh Park, near Havant, it flowered in Sep-

* Literally, a torch or taper; a name translated by the English Torch-thistle; and given to these plants in consequence of the upright kinds having something the appearance of the tapers used in the ceremonies of the Roman Catholic religion.

September 1834, under the good management of Mr. Robert S. Wilson the gardener. The plant had been in the collection at Leigh Park upwards of fifteen years without blossoming ; between March and September it produced shoots upwards of seven feet long.

Two flowers were perfected, of which one opened about six o'clock in the afternoon of the 22nd of September, and faded about eleven A.M. on the following day ; the other was despatched to London by coach on the 24th of the same month, and reached London in perfect condition.

It was indeed a beautiful object, its petals were of the most dazzling whiteness, the effect of which was greatly heightened by the dense mass of yellow stamens occupying the centre, and by the border of olive green sepals, on which the petals reposed. This is said to have the largest flowers of all the species, not even excepting the common night-blowing *Cereus* ; its fruit is described as being quite smooth, of a rich scarlet, and with the size and form of a goose's egg.



* EÚTOCA víscida.

Clammy Eutoca.

PENTANDRIA MONOGYNIA.

Nat. ord. HYDROPHYLLACEÆ. (Introduction to the Natural System of Botany. p. 244.)

EUTOCA.—Suprà, vol. 14. fol. 1180.

E. viscida; glanduloso-pilosa viscosa, caule erecto ramoso, foliis cordato-ovatis subangulatis serratis, racemis elongatis furcatis simplicibusque, placentis multi-ovulatis.

E. viscida. Bentham MSS.

Annua, undique pilis nigro-capitatis glandulosis viscosa; caule tereti ramoso bipedali. Folia tactu mollia et oleosa, 2 p. lata, paulò longiora, sensim decrescentia, superiora grosse simpliciter dentata. Racemi gyratim elongantes, multiflori, ebracteati. Sepala linearia, obtusa, capsulæ longitudine. Corolla generis, amæne cærulea tubo roseo. Filamenta setacea basi villosa; antheræ flavæ. Capsula ovata, mucronata, apice bivalvis, semibilocularis, placentis parietalibus polyspermis. Semina minuta, fusca, scrobiculata, marginibus areolarum acutis centro foveola excavata.

A new hardy annual, found in California by Mr. Douglas, and raised by the Horticultural Society, in whose garden it flowered this summer for the first time. It is perfectly hardy and will grow in any common soil. Our drawing was made last July.

The blue of the flowers is remarkably deep and brilliant, so that the plant has a handsome appearance; but its leaves are rather coarse and weedy. We know, however, of no plant better adapted for bouquets; for it will go on growing and flowering in water for two or three weeks after being gathered.

The whole of its surface is covered with hairs, having little black heads filled with a viscid secretion, and sticking to the fingers like those of the Henbane; these are so crowded about the pedicels as to give them quite a sooty aspect.

* See folio 1180.

1809.



S. Watts. sc.

Pub by J. Ridgway 169 Piccadilly Nov. 1. 1835.

S. Watts. sc.

* VÁNDA téres.

Taper-leaved Vanda.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDACEÆ, § VANDEÆ. (Introduction to the Natural System of Botany, p. 262.)

VANDA.—*Suprà, vol. 6. fol. 506.*

V. teres; foliis teretibus, racemis ascendentibus subbifloris foliis æqualibus, sepalis oblongis obtusis: supremo erecto lateralibus semitortis labello suppositis, petalis majoribus suborbiculatis undulatis, labello basi conico: laciniis lateralibus ascendentibus subtruncatis intermediâ pubescente apice dilatâtâ truncatâ emarginatâ. *Lindl. gen. et sp. Orch. p. 217.*

This superb epiphyte was originally discovered upon trees in Sylhet by Dr. Wallich, and it has more recently been met with by Mr. William Griffith abundantly near Medown in the Burmese Empire, also growing on trees in the woods.

When Dr. Wallich came to England, this species formed one of the numerous living plants brought home under his care, and blossomed during the voyage; the very flowers which were then produced we are so fortunate as to possess in spirits, through the kindness of their liberal discoverer; they measure nearly four inches and a half from the tip of one petal to that of the other, but only two were formed upon the spike. The specimens from Mr. Griffith are not

* Vandá is the Sanscrit name of the original species of this genus.

quite so large in the flower, but there are three blossoms to each spike.

That from which our drawing was taken was produced at Syon, in the collection of his Grace the Duke of Northumberland, by whose permission we are enabled to publish it. It was not quite so fine as what we have represented, but considering what the native specimens are we have felt quite justified in forming our figure between the two. The flowers in the accompanying plate, though larger than those on the plant at Syon, are considerably smaller than those we possess in spirits.

Nothing can exceed the flowers of this plant in delicacy of texture or softness of colour, the deep purple of the petals softens away to the margin, and seems to melt as it were into the purer white of the sepals, while the rich crimson and yellow of the lip renders the brilliancy of the other parts still more conspicuous.

We found the plant had been managed by having its stem covered with moss, which if kept damp would maintain the surface in a uniform state. We do not, however, perceive any trace of the roots in our wild specimens having been growing among moss, and we consider it probable that it will succeed as well if the roots are exposed freely to the air.



* **CRATÆGUS** Douglásii.*The Douglas Thorn.*

ICOSANDRIA MONO-PENTAGYNIA.

*Nat. ord. ROSACEÆ, § POMEÆ. (Introduction to the Natural System of Botany, p. 83.)**CRATÆGUS.—Suprà, vol. 83. fol. 1128.**C. Douglasii*; foliis obovatis ovalibusque inciso-serratis acutis basi cuneatis glabris demum subcoriaceis sublucidis, laciniis calycis ovatis glanduloso-serratis pedunculisque glabris, spinis rigidis rectiusculis.*C. punctata. β. brevispina. Douglas in Hook. Fl. Bor. Am. 1. 201.*Rami *ascendentes, cinereo-purpurei, glabri, spinis rigidis lævigatis, nunc brevibus nunc longissimis. Folia semper glabra, petiolis omninò eglandulosi, inciso-serrata nullo modo angulata vel plicata, sub fine anni coriacea et sublucida, citò tamen decidua, paulò post C. glandulosam et punctatam. Inflorescentia cymosa glabra, floribus magnitudine mediocri subtetragynis. Sepala ovata glanduloso-serrata. Fructus parvus, atropurpureus.*

A native of North West America, where it was collected by Mr. Douglas. A hardy tree of small size flowering in May.

We believe this to be essentially different from all the published species of this genus, but the whole of the synonymy is in so deplorable a state of confusion, from want of some monographer, who at the same time is well acquainted with the living trees, and has access to authentic sources of information, that we are by no means certain that it has not already appeared in print as a species.

Mr. Douglas considered it a variety of *C. punctata*, but the leaves of that species are far thinner, and have an evident tendency to be furrowed as if plaited; its branches are

* See folio 1161.

horizontal, and its fruit large and either red or yellow, in all which respects it differs from this. Its inflorescence, moreover, is downy. It is possible that what is called *C. macracantha* in the gardens may not be specifically distinct, but as its fruit is red it requires further examination.

The leaves of this species are remarkably leathery in the autumn, and then acquire a purplish cast and are shining. They fall off about the same time as those of *C. punctata* and *pyrifolia*.

The plant from which our drawing was taken stands in the Arboretum of the Horticultural Society.



* **MAXILLÁRIA** cristáta.*Crested Maxillaria.*

GYNANDRIA MONANDRIA.

*Nat. ord. ORCHIDACEÆ, § VANDEÆ. (Introduction to the Natural System of Botany, p. 262.)**MAXILLARIA.—Suprà, vol. 11. fol. 897.*

M. cristata; pseudobulbis ovatis sulcatis monophyllis, foliis oblongo-lanceolatis plicatis, scapo pendulo bifloro squamis laxiusculis vaginato, floribus explanatis, sepalis petalisque lanceolatis acutis æqualibus, labello multo minore tripartito carnosio: laciniis lateralibus falcatis intermediâ rotundatâ cristato-fimbriatâ utrinque unidentatâ, ungue subcristato disco bidentato.

Sepala $1\frac{3}{4}$ unciam longa, alba, sanguineo interruptè striata et fasciata. Petala æqualia et conformia, apice sanguinea basi maculata dorso alba. Labellum purpureum, ungue viridi cristâ dentibusque albis; crista laciniæ intermediæ fimbriata pilis submoniliformibus: unguis e cirrhulis 4-6 rectis submoniliformibus constans; dentes disci compressi divaricati margine crenati posteriore duplò majore. Columna basi viridis, apice lutea, et utrinque subalata, rostello longissimo subulato. Caudicula polliniorum longissima, glandulâ parvâ subtriangulari.

For the opportunity of publishing this remarkably beautiful epiphyte, we are obliged to Mr. Knight, of the King's Road Nursery, with whom it flowered in the stove in July last.

It is a native of Trinidad, growing on old decayed branches of trees near the Mud Lake. We have seen a drawing of it taken on the spot, and it was in no degree superior in beauty to the present specimen.

The lip is a most curious organ. It is of a firm fleshy texture, and is deeply divided into three parts; of these the

* See folio 1206.

two lateral ones are falcate and smooth, while the middle one is rounded, has a strong tooth on each side, and is bordered by a deep white fringe of necklace-shaped hairs ; then the short stalk of the lip has four or more such hairs growing straight from its border, and the disk is furnished with a large white two-lobed flat tooth, the divisions of which diverge from each other.

The striping, banding, and painting of the delicate white flowers with rich crimson, produces a very rich and striking effect.



* GARDOQUÍA Gilliésii.

The Gillies Gardoquia.

DIDYNAMIA GYMNOSPERMIA.

*Nat. ord. LABIACEÆ. (Introduction to the Natural System of Botany, p. 239.)**GARDOQUIA.—Suprà, vol. 21. fol. 1747.*

G. Gilliesii; suffruticosa, ramis divaricatis pubescentibus, foliis oblongo-linearibus cuneatisve obtusis integerrimis basi angustatis utrinque viridibus planis, floralibus conformibus bracteisque calyce subbrevioribus, verticillastris paucis multifloris laxiusculis irregularibus, calycibus elongatis erectis, dentibus lanceolato-subulatis subæqualibus, fauce intùs nudâ. *Bentham Gen. & Sp. Labiat. p. 403.*

*G. Gilliesii. Graham in Edinb. Phil. Journ. 1831, Sept.**G. chilensis. Bentham in Hooker & Arn. Beech. Voy. 58.*

A neat little half-shrubby herbaceous plant, growing not more than 6 or 8 inches high, flowering from June to September in the open border, and readily multiplied by cuttings which root freely in peat and sand. It requires a little protection in winter.

It appears to be a common Chilean plant, for it occurs in almost every collection that has yet reached England from the neighbourhood of Valparaiso.

Our drawing was made in the Garden of the Horticultural Society.

Although this is far less showy than *G. Hookeri*, figured at fol. 1747, yet as it is a hardier plant it will be to many the more interesting of the two.

Our figure represents the analysis of the parts of fructification: 1. is a magnified entire flower; 2. is a calyx; 3. a corolla split open; and, 4. one of the anthers with a part of the filament.



Miss Drake del.

Pub by J. Ridgway 16j Piccadilly Nov. 1. 1835.

J. Walp. sc.

* **DAUBÉNYA** aúrea.*Golden Daubenyia.*

HEXANDRIA MONOGYNIA.

Nat. ord. ASPHODELACEÆ. (Introduction to the Natural System of Botany, p. 273.)

DAUBENYA.—*Inflorescentia* umbellata, sessilis, epigæa. *Perianthium* tubulosum, ovario arcte appressum, limbo bilabiato, labio superiore nano tridentato, inferiore majore tripartito : in floribus radii maximo, disci depauperato. *Stamina* 6 inæqualia basi laciniarum adnata, declinata, in floribus disci sæpè submonadelphæ. *Ovarium* elongatum, subtriangulare angulis rotundatis, in stylo angustatum ; 3-loculare, loculis polyspermis apice vacuis. *Stigma* simplex.—*Bulbi* capenses habitu omnino *Massoniæ*.

D. aurea.

Massonia lutea. Hort.

Folia 2, oblonga, carnosa, sulcata, prostrata, umbellam sessilem multifloram e sinu suo emittentia. Flores lutei, densè aggregati ; radii labio exteriori (inferiore) 3-partito ; laciniis obovatis rotundatis lateralibus brevioribus ; disci laciniis omnibus angustis labii majoris acutis paulò elongatis. *Stamina* 6, basi laciniarum adnata, declinata, laciniæ inferiori intermediæ oppositum cum eâ altiùs connatum. *Antheræ* dorso affixæ. *Ovarium* laciniis nullis inter loculos. *Ovula* rotunda funiculo brevi placentæ adnata.

A greenhouse bulb, native of the Cape of Good Hope, whence it was obtained by Messrs. Young of Epsom, under the name of *Massonia lutea*. It flowers in June, and is very pretty as well as singular, so long as its blossoms remain expanded, which is for about three weeks ; afterwards its flaccid green leaves have little to attract attention.

No trace of this plant is to be found in books, so that it is probably some recent discovery made in the interior of the Cape Colony. From *Massonia*, with which it remark-

* We name this new and curious genus after Dr Charles Daubeny, Professor of Botany at Oxford, whose interesting researches in Vegetable Chemistry have materially conduced to improve our knowledge of the physiology of plants.

ably agrees in habit, it differs essentially in its tubular not campanulate, very irregular, perianth, and in the absence of the honey-pores which form so remarkable a part of the character of *Massonia*. With the fruit we are unacquainted.

Fig. 1. represents a flower of the ray with the larger lip; fig. 2. is one of the flowers of the disk, in which both lips are very small and imperfect; fig. 3. is a vertical section of the ovary, with the empty upper portion of the cells; fig. 4. is a transverse section of the ovary, shewing how closely it is invested by the tube of the perianth; this takes place to such a degree as to give the ovary the appearance of being inferior.



* **ELICHRYSUM** bicolor.*Two-coloured Elichrysum.*

SYNGENESIA POLYANDRIA.

*Nat. ord. ASTERACEÆ. (COMPOSITÆ.) Introduction to the Natural System of Botany, p. 197.)**ELICHRYSUM.—Suprà, vol. 1. fol. 21.*

E. bicolor; annuum; foliis lineari-lanceolatis acuminatis basi obtusis scabro-ciliatis, superioribus subulatis, caule glabro ramoso ramulis monocephalis esquamatis, bracteis involucri fulvis aureisque acutis.

Caulis 2-pedalis, sulcatus, fastigio ramoso. Folia basi aliquando subcordata, margine scabra, supra scabriuscula. Capitula facie omnia *E. bracteati*.

A beautiful new hardy annual introduced by Mr. Low of the Clapton Nursery. In appearance it resembles the now common *E. bracteatum*, but is much handsomer.

It is a native of Van Diemen's Land, whence our excellent correspondent Mr. Gunn has sent beautiful specimens (No. 111), agreeing in all respects with the cultivated plant.

It is one of the prettiest new species of the season that has just past by.

* This is an old Greek name, meaning a golden spiral, and consequently should be written Helichrysum. It is supposed to have been the plant now called Gnaphalium Stæchas.



Miss Drake del.

Pub by J. Ridgway 169 Piccadilly Decr. 1. 1835.

J. Watts sc.

* **MACRADÉNIA** triándra.*Triandrous Long-gland.*

GYNANDRIA MONANDRIA.

*Nat. ord. ORCHIDACEÆ, § VANDEÆ, Lindl. (Introduction to the Natural System of Botany, p. 262.)**MACRADENIA.—Suprà, vol. 8. fol. 612.*

M. triandra; foliis coriaceis lineari-oblongis acuminatis, racemo prostrato, lahello in medio trilamellato, clinandrio serrato antheris duabus sterilibus.

Pseudobulbi cylindranei paulo attenuati. Folia plurivenia. Racemi foliis multò breviores. Sepala et petala intus sanguinea herbaceo limbata. Labelli lamellæ rotundatæ, costæformes, antrorsum crassiores. Antheræ steriles ovales, integræ, sanguineo marginatæ.

The *Macradenia lutescens* of this work (folio 612) we have never seen; but as it was described by Dr. Brown we have no difficulty in deciding, upon the evidence that the description affords, that it is specifically different from this.

No notice is taken by Dr. Brown of the existence of any lamellæ in the middle of the lip, nor of the abortive anthers which are always present; besides which the clinandrium here is regularly and strongly serrated, the leaves much narrower, and the raceme pendulous or prostrate instead of erect.

A native of Surinam, whence it was brought by Mr. Lance. It flowered in the possession of the Horticultural Society in May last. A strong damp stove heat is required to keep it in health.

* So named in allusion to the long caudicula of the pollen masses, fig. 5. from μακρός long, and αδην a gland.



Miss Drake, del.

Pub by J. Ridgway 169 Piccadilly Dec. 1. 1835.

J. Watts, sc.

* COCCOLÓBA virens.

Green Sea-side Grape.

OCTANDRIA TRIGYNIA.

Nat. ord. POLYGONACEÆ. (Introduction to the Natural System of Botany, p. 221.)

COCCOLOBA, L. *Calyx* 5-partitus, semi-inferus, laciniis imbricatis; in fructu incrassatus baccans. *Stamina* 7-10 serie simplici fauce inserta, filamentis basi in urceolum brevissimum coalitis. *Ovarium* semi-superum, trigynum; stigmatibus spathulatis nunc lobulatis. *Achenium* basi lobatum, osseum, calyce baccato tectum.—Frutices arboresve *minores, occidentales*. *Flores racemosi*. *Fructus subacidi, edules*.

C. virens; foliis ovato-lanceolatis obtusis basi in petiolum angustatis, racemis nutantibus, floribus decandris.

Folia omninò depilata, ovato-lanceolata, semper basi angustata nec ullo modo basi obtusa vel cordata; omnia conformia. Racemi virides, nutantes, foliis minoribus æquales.

The species of *Coccoloba* are so ill defined in books that it is by no means easy in the absence of fruit to determine them. This in some respects resembles *C. obtusifolia* of Jacquin, in others *C. microstachya* of Willdenow. From the former it differs in the form of its leaves, from the latter in their size and proportion to the racemes.

A hothouse plant, communicated from Wormleybury by Sir Abraham Hume, with whom it flowered in August 1833. It was called *C. excoriata*, which is a very different plant.

We are unacquainted with its native country, but presume it to be the West Indies.

* From *κοκκος* a fruit (in this case seed) and *λοβος* a lobe, in allusion to the lobed seeds.



* *OXALIS* Piottæ.*Piotta's Oxalis.*

DECANDRIA PENTAGYNIA.

*Nat. ord. OXALIDACEÆ. (Introduction to the Natural System of Botany, p. 140.)**OXALIS.—Suprà, vol. 15. fol. 1249.*

-
- O. Piottæ*; acaulis, petiolo planiusculo, foliolis 3 obcordatis glabris, scapo pubescente unifloro foliis longiore, sepalis integris, stylis longissimis, stigmatibus plumosis.
- O. Piottæ. Colla Hort. Rip. p. 98. t. 1.*
-

A truly beautiful little half hardy, or frame perennial, flowering most copiously during the months of July and August. A little tuft does not indeed produce much appearance, but a pot filled with its dense green leaves and covered with the large salmon coloured flowers is a lovely object.

Said to be a native of the Cape of Good Hope. Our specimens were obligingly communicated by Mrs. Marryat, to whom they were sent by Mrs. Palliser. That lady obtained them both from Professor Savi of Pisa and from the direction of other Italian Botanic Gardens.

We believe the plant is at present in the possession of no one in this country except Mrs. Marryat, and those of her friends to whom she has given it.

The species seems nearly allied to *O. compressa*.

* See folio 1249.



* GALATÉLLA punctáta.

Dotted Galatella.

SYNGENESIA POLYGAMIA FRUSTRANEA.

Nat. ord. ASTERACEÆ. (COMPOSITÆ.) (*Introduction to the Natural System of Botany*, p. 197.)

GALATELLA, Cassini. *Capitulum* radiatum; radio uniseriali, ligulis fœmineis abortivis; disci flosculis hermaphroditis fertilibus tubulosis; stigmatibus apice spathulatis obtusis. *Involucrum* disco brevius; foliolis pluriserialibus imbricatis inappendiculatis; interioribus membranaceis exterioribus sæpè trinerviis. *Receptaculum* alveolatum, marginibus alveolorum paleaceo-laceris. *Pappus* simplex, pluriserialis, radiis filiformibus rigidis serrulato-scabris. *Achenia* densè hirsuta v. strigoso-setacea. *Nees Genera et Sp. Asterearum*, p. 158. (terminis mutatis.)

G. punctata; foliis lineari-lanceolatis acuminatis integerrimis trinerviibus punctatis scabris, ramulorum linearibus, caule apice corymboso ramis elongatis paucifloris contractis, involucris brevibus turbinatis, radio longo. *Nees l. c.*

Aster punctatus. *W. & Kit. Pl. Hung.* 2. 113. t. 109.

Galatella intermedia. *Cass. in dict. des sc. nat. xviii.* p. 58.

Aster desertorum. *Fisch. ined.*

A hardy herbaceous plant, native of salt marshes in Hungary, Podolia, and elsewhere in the east of Europe. It also occurs in Siberia. In this country it flowers in July and August, growing to the height of about two feet, and forming a thick compact bush.

We greatly doubt whether it is really a distinct species from *G. hyssopifolia* and *acris*, or they from each other, so very difficult is it to distinguish them with absolute certainty when one has a long series of specimens under examination. From the former it may, however, be generally known by its compact inflorescence, and larger flower heads, which are

quite regularly corymbose, instead of being of different heights as in *hyssopifolia*; its leaves too are more uniformly three-ribbed.

In *G. acris* the leaves are pretty uniformly one-ribbed, and the inflorescence much smaller and more contracted.

All the species of this genus are well adapted for borders of shrubberies, and for places where shade-loving plants alone will grow.



* **OCHRANTHE** *arguta*.*Fine-toothed Palebloom.*

PENTANDRIA TRIGYNIA.

Nat. ord. HYPERICACEÆ ANOMALÆ.

OCHRANTHE. *Calyx* membranaceus, 5-phyllus, imbricatus, *corollæ* pentapetalæ simillimus. *Stamina* 5 hypogyna. *Discus* urceolatus pentagonus. *Carpella* 3, basi juncta, stylis disjunctis. *Ovula* cuique carpello 6, placentæ centrali affixa.—*Folia* opposita, serrata. *Stipulæ* interpetiolares serrata. *Flores* terminales, pallidi.

Ochranthe arguta.

Caulis fruticosus, glaber, adultus cinereus, junior viridis cinereo maculatus. *Folia* opposita, glabra, petiolata, obovato-lanceolata, acuminata, serrata, basi integra. *Stipulæ* intrapetiolares, ovatae, serrulatae, pallidae. *Thyrus* terminalis, congestus, pauciflorus. *Flores* inodori, albidii, demum flavescentes. *Pedicelli* basi 1-, medio 2-bracteati, glabri. *Calyx* erecto-patens, irregularis, 5-phyllus; sepalis concavis, obtusis, inæqualibus, exterioribus corolla brevioribus, margine subciliatis, æstivatione imbricatis $\frac{3}{4}$. *Petala* 5, hypogyna, subconvoluta, rarò aliquot patentia, unguiculata, oblonga, obtusa, versus medium 3-venia. *Stamina* 5, hypogyna, erecta, rigida, pistilli longitudine, petalis alterna; antheræ medio affixæ, introrsæ, erectæ, biloculares, longitudinaliter dehiscentes; pollen rotundo-triquetrum, angulis (inflatis?) globosis pellucidis, nunc subrotundum angulis nullis. *Discus* cyathiformis, pentagonus, carnosus, angulis planatis. *Ovarium* superum, ovatum, obtusè trigonum, stylis 3 subulatis erectis, versus basin pilis raris patentibus; 3-loculare, ovulis circiter 6 in utroque loculo, placentæ versus apicem axeos appensis.

A native of China, and consequently a Greenhouse plant. It flowered in the Garden of the Horticultural Society so long since as March 1826; but shortly after died, and has never again made its appearance. We have kept the drawing ever since in our portfolio, in the hope of the species being

* From ὀχρὸς pale, and ανθος a flower.

again discovered, but we now find it necessary to publish some account of the plant.

The fruit being unknown, we are unable to obtain even an approximation to its true station in the system. In some respects it resembles Cunoniaceæ, especially in its remarkable stipulation, and apocarpous many-seeded pistil, combined with opposite leaves; but it differs in having only five hypogynous stamens, three instead of two carpels, and a remarkably imbricated calyx.

Eucryphia and Carpodontos among Hypericaceæ, are polyandrous and have no stipules, besides other points of discrepancy.

Hugoniaceæ have alternate leaves, a different habit, and a larger number of stamens.

Anisadenia, which seems to be a shrubby Elatinaceous plant, has monadelphous stamens, and no hypogynous cup, &c. Its leaves are also alternate without stipules.

In all probability Ochrante forms a part of some unknown order, the station of which will be in the Calycese group of polypetalous Dicotyledons, either in the Cistal or Guttal alliance. (*See Key to Structural, Physiological and Systematic Botany, p. 51.*)



* RHODODÉNDRON pulcherrimum.

The lovely Rhododendron.

DECANDRIA MONOGYNIA.

*Nat. ord. ERICACEÆ. (Introduction to the Natural System of Botany, p. 182.)**RHODODENDRON.—Suprà, vol. 1. fol. 37.*

GARDEN VARIETY.

A most beautiful plant obtained by Mr. Waterer of Knap-hill between *R. arboreum* and *caucasicum*. It is of rather delicate appearance, but we are informed that it is quite hardy, and an abundant flowerer. Our drawing was made in March last.

Fig. 1 is a kind called *Rhododendron Nobleanum*, which is very much like the other in all respects, except that its flowers are of a deep and brilliant rose colour.

Both are among the handsomest hardy shrubs in cultivation.

* See folio 1240.



* EULÓPHIA lúrida.

Lurid Eulophia.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDACEÆ, § VANDEÆ, Lindl. (*Introduction to the Natural System of Botany*, p. 262.)

EULOPHIA.—*Suprà*, vol. 17. fol. 1433.

E. lurida; foliis (lineari-) lanceolatis scapo ramoso multò brevioribus, bracteis minimis subulatis, sepalis lineari-spatulatis obtusis, petalis paulò latioribus, labelli tripartiti basi callosi lobis lateralibus obtusis recurvis: intermedio obcordato, calcare cylindraceo inflexo obtuso. *Lindl. gen. et sp. Orch.* p. 182.

One of the easiest of all stove Orchideous plants to cultivate, growing upon a damp wall, or a rough stone, or under almost any kind of condition, provided the air be hot and damp, and uniform. It then flowers profusely at intervals throughout the whole year. Our drawing was made in the noble collection of the Messrs. Loddiges in January last.

A native of Sierra Leone, whence great quantities are occasionally brought. It appears to be extremely common upon the trunks of trees in some parts of that colony.

* See folio 1433.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly Jan. 1. 1836.

J. Watts. sc.

* COSMÉLIA rúbra.

Red Cosmelia.

PENTANDRIA MONOGYNIA.

Nat. ord. EPACRIDACEÆ.

COSMELIA. Calyx foliaceus. Corolla tubulosa. Stamina epipetala; antheræ apicibus ciliatis filamentorum adnatæ. Squamulæ 5, hypogynæ. Capsula placentis columnæ centrali adnatis.—Frutex paludosus, erectus, ramosus, ramis dum denudatis non cicatrisatis. Folia basibus cucullatis semivaginantibus. Flores ramos laterales breves terminantes, solitarii, cernui. Calyx foliis minoribus imbricatus. Corolla ruberrima. Stamina inclusa, antheris basi solutis. Placentæ utraque extremitate liberæ. Brown prodr. 553.


C. rubra. Brown l. c.

According to Dr. Brown this is a marsh plant, found upon the South Coast of New Holland, and remarkable for its leaves, although so thick and rigid, not leaving conspicuous scars upon the branches when they fall off.

In this country it is a very pretty greenhouse plant with the habit of an Epacris. Our drawing was made in Messrs. Loddiges collection in May last.

The dissected parts of the flower in the accompanying plate afford a good illustration of the Natural order Epacridaceæ, which are only known from true Ericaceæ by their anthers having single cells opening by a pore. 1. Is the top of a corolla cut open, with the stamens sticking to it; 2. an anther with a portion of the upper end of a filament; 3. a pistil, with the five hypogynous scales.

* Apparently from κοσμεω to ornament, in allusion to its beauty.



Miss Drake. del.

Pub by J. Ridgway 109 Piccadilly Jan. 1. 1836.

J. Walts. sc.

* *LASTHENIA californica*.*Downy Lasthenia.*

SYNGENESIA POLYGAMIA SUPERFLUA.

Nat. ord. ASTERACEÆ (COMPOSITÆ).

LASTHENIA.—*Suprà*, vol. 21. fol. 1780.*L. californica*; subpubescens; foliis integerrimis, capitulis basi umbonatis.*Supra l. c.**L. californica*. *De Cand. prodr. ined.*

This species differs from the one already published in the present work in little except its smaller size, and differently formed flower heads, which are impressed at the base with a cavity for the peduncle, and like the peduncles themselves are slightly downy.

It is an annual plant, flowering for about six weeks at different periods of the year, according to the season at which its seeds are sown.

Professor De Candolle, in a manuscript list of the genera of Compositæ, which we have just received from him, includes it in his first series *Tubulifloræ*, fourth tribe *Senecionideæ*, fifth sub-tribe *Heleniæ*, first division *Gaillardieæ*, second sub-division *Euheleniæ*, along with *Cephalophora* and *Helenium*, both of which are known in our gardens. We observe that Professor De Candolle attributes the genus to Cassini, but we have not succeeded in discovering it in any of the writings of that most unmethodical of Botanists.

* See folio 1780.



Miss Drake. del.

Pub by J. Ridgway 169 Piccadilly Som. 1. 1836.

J. Watts. sc.

* ARISTOLOCHÍA fœtens.

Stinking Birthwort.

GYNANDRIA HEXANDRIA.

Nat. ord. ARISTOLOCHIACEÆ.

ARISTOLOCHIA.—*Suprà*, vol. 8. fol. 689.

A. fœtens; foliis lato-cordatis acutis, caule volubili, pedunculis solitariis, bracteâ perfoliatâ, limbo calycis maximo integro cordato labio longissimè caudato, tubo extus glabro.

Nauseosa, fœtida, demum semiputrida ramis scandit longis volubilibus glabris. Folia cordata sinu aperto, integerrima, subrotunda, acuminata, subtus levissimè pubescentia, glaucescentia. Bracteæ solitariæ, orbiculatæ, perfoliatæ. Flores maximi, tubo extus glabro, basi inflato, sursum angustato costato arcuato, limbo subrotundo cordato basi clauso apice in appendicem longum linearem subspiralem acuminatum producto, luteo purpureoque livido variegato maculato et punctato.

A native of the West Indies, whence it has been obtained by Mrs. Marryat, in whose stove at Wimbledon it flowered in June last. Like all the other hothouse species it strikes readily from cuttings.

It is chiefly remarkable for the large size, and singular colour of its flowers, which are beautifully variegated with purple and dirty yellow; they have a most disagreeable disgusting smell, which will prevent the plant from becoming a favourite.

Nearly allied to *A. grandiflora*, from which it is distinguished by the smooth tube of its calyx, the colour of the same part, &c. &c.

* See folio 1399.



3.



1.



5.

4.



2.

* PLEUROTHÁLLIS picta.

Painted Pleurothallis.

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDACEÆ, § MALAXIDEÆ.

PLEUROTHALLIS.—*Suprà*, vol. 9. fol. 759.

P. picta; folio spathulato marginato retuso racemis laxis duplò brevioribus, bracteis minimis, sepalis acuminatis lateralibus apice tantùm sejunctis, petalis linearilanceolatis acutis, labello lineari obtuso carnosissimo supra 1-sulcato. *Suprà* fol. 1797.

Very near the *P. Grobyi* already figured in this work (fol. 1797), from which it principally differs in its spathula-shaped retuse leaves, and in the narrower form of all the parts of the flower; as will be seen by comparing the dissections in the two plates.

This also is a native of Demerara, and was sent us by Mr. Loddiges first in October 1834, and next in March 1835, when our drawing was made. It is a graceful pretty species, and well deserves an attentive examination.

Its tufts are rapidly formed under good management, but it seems to require the close atmosphere of a bell-glass.

Fig. 1. is a view of the column, petals, and lip. 2. The column by itself. 3. The upper side of the lip. 4. The pollen mass in situ. 5. The same separated from each other.

* See folio 1298.



* HIBISCUS *Rosa sinensis*.*Single-flowered Chinese Rose Mallow.*

MONADELPHIA POLYANDRIA.

Nat. ord. MALVACEÆ.

HIBISCUS—*Suprà*, vol. 1. fol. 29.

- § IV. *KETMIA*. Carpella seu capsulæ loculamenta polysperma. Semina glabra. Corollæ expansæ. Involucella 5-7-phylla. Calyx 5-lobus non longitudinaliter ruptus. *D. C. prodr.* 1. 448.
- H. *Rosa-sinensis*; caule inermi arboreo, foliis ovatis acuminatis glabris basi integerrimis apice grossè dentatis subincisis, pedicellis folii longitudine, involucello 7-phylo. *D. C. l. c.*
- H. *Rosa Sinensis*. *Linn. sp. pl.* 977. *Lour. fl. cochinch.* 2. 419. *Cav. diss.* 3. 1. 69. f. 2.
- Flos Festalis. *Rumph. Amb. book* 6. chap. 11. t. 8.

The double varieties of this species, crimson, yellow, buff, and even white, are not uncommon in collections, but the single state which is now represented, although much handsomer, is comparatively rare.

Instead of a crowd of ragged, unequal, ill-arranged petals, destitute of all symmetry, occupying the centre of the flower, we have a long graceful curved crimson tube, terminated by a brush of bright yellow anthers, which surround five little crimson velvet cushions of stigmas.

It is easily propagated by cuttings, but requires a good stove to flower in perfection. Our drawing was made in that of the Horticultural Society in September 1834.

According to Loureiro this species is spontaneous both in China and CochinChina; where it is employed for many

* See folio 1463.

purposes as an emollient. The flowers tinge spirits of wine red, or by the addition of a little alum a beautiful violet. Shoes or other similar things are blackened, by rubbing the flowers upon them. In CochinChina the plant is so common that garden hedges are often made of it.

Rumf says that in Amboyna it was in his days employed as a common ornament on occasions of festivity and even at funerals. From its constant use for blacking shoes it had acquired the barbarous Portuguese name of Fula Sapato.



* *PIMELÉA* ligustrîna.*Privet-leaved Pimelea.*

DIANDRIA MONOGYNIA.

Nat. ord. THYMELACEÆ.

PIMELEA.—*Suprà*, vol. 15. fol. 1268.

1. *Folia opposita. Capitulum terminale. Involucrum foliis rameis dissimile.* Br. prodr. 359.

P. ligustrina; involucris tetraphyllis, foliolis ovatis extùs venosis intus pubescentibus, perianthiis sericeis, receptaculis fructiferis ovatis pube brevissima, foliis ovalibus oblongo-lanceolatisve venosis. *Br. l. c.*

P. ligustrina. *Lab. Fl. Nov. Holl.* 1. p. 9. t. 3.

A neat species of the extensive genus *Pimelea*, found from Van Diemen's Land to Port Jackson, and growing in its native places as much as ten feet high. We have specimens from Mr. Gunn of Launceston, and others.

It is a hardy greenhouse plant, readily multiplied by cuttings. It only requires good ventilation during winter, so as to prevent its being destroyed by damp. Our drawing was made in Mr. Lowe's Nursery in March 1834.

We are not aware of the particular part of the island of Van Diemen in which this shrub occurs; but if it is from the south side it will probably prove hardy in the southern parts of England.

* See folio 1268.



Miss Drake del.

Pub. by J. Ridgway 169 Piccadilly Jan. 1820.

J. Walp. sc.

* **DENDRÓBIUM densiflórum.***Dense-flowered Dendrobium.*

GYNANDRIA MONANDRIA.

Nat. ord. ORCHIDACEÆ, § MALAXIDEÆ.

DENDROBIUM.—*Suprà, vol. 7. fol. 548.*

D. densiflorum ; caulibus articulatis clavatis pendulis apice foliosis, foliis oblongis acutis nervosis, racemis lateralibus multifloris foliis longioribus ; junioribus strobiliformibus, bracteis oblongis plicatis recurvis pedicellis longioribus, sepalis patentissimis ovatis obtusis, petalis conformibus majoribus, labello majore rhomboideo unguiculato serrulato retuso. *Lindl. Gen. & Sp. Orch. p. 90. Wall. Pl. as. rar. no. 40.*

Planta pendula, lævis, cæspitosa, arborum parasitica, pedalis-2-pedalis. Radix constans fibris numerosis, fasciculatis, teretibus, indivisis. Caules plures, rigidi, cylindræo-clavati, lucidi, ætate flavicantes, maximâ parte nudi, sulcis aliquot profundis totidemque angulis obtusissimis notati, vestigiis vaginalium remotis annulati, extrorsum digitum crassi, apice foliosi. Folia valdè approximata, patentia, alterna, ovato-oblonga, acuta, 5-pollicaria, firma, subcarnosa, lucida, plana, nervosa, basi contractâ obsoletissimè petiolatâ abientia in vaginas breves, adpressas, deciduas. Racemus magnus, solitarius, laxè pendulus, versus apicem caulis infra folia lateralis, pedunculatus, 6-8-pollicaris, ovato-oblongus, obtusus, multiflorus ; novellus, longè ante expansionem, sessilis, ovatus, strobiliformis, 2-pollicaris, valdè regulariter imbricatus bracteis adpressis, planis, alternantibus. Pedunculus crassus, cylindricus, 2-3-pollicaris, bracteis sparsis, oblongis, recurvis, nervosis, membranaceis, ferè pollicaribus, persistentibus. Flores valdè numerosi, majusculi, flavi, approximati, undique sparsi, patentissimi, insidentes pedicellis teretibus, pollicaribus, suffultis bracteâ magnâ, obovatâ, nervosâ, persistente. Sepala oblonga, obtusa, semipollicaria ; lateralia parùm majora, basique dilatâ infra basin columnæ in calcar brevissimum, subcylindricum, obtusum connata, uti petala ovata, subæquantia, patentissima. Labellum planum, patens, retusum, ciliato-denticulatum, suprâ pubescens, basi desinens in unguem brevem, canaliculatum. Columna brevis, conica, crassa, basi leviter producta. Anthera ovata. Wallich.

“ This lovely *Orchidea* comes so near to Roxburgh's *Dendrobium clavatum* (Hort. Beng. p. 63), that I should consider them as identical, if he had not ascribed bulbs to his plant, of which ours is entirely destitute. In the Calcutta garden the two plants maintain that character, although in other respects they are alike. It is possible that under particular circumstances, the bulbs or pseudo-bulbs may become elongated into clavate stems, and if so the Silhet plant would be the same species with ours from Nipal.”

The above we borrow from Dr. Wallich's splendid *Plantæ Asiaticæ*.

Beautiful as is the specimen of this plant, represented in the accompanying figure, it is still inferior to what is produced in India, so that cultivators have still a point to gain in respect to this charming species. Our drawing was made in the stove of Messrs. Loddiges in May last.

Since the last addition to the published species of *Dendrobium*, we have received from Mr. Allan Cunningham specimens of a most remarkable new kind, with the following memorandum.

“*Dendrobium cassythoides*; aphyllum, caule glabro stolonifero, floribus racemosis, racemis (stolonibus oppositis) subtribrachiatis, bracteis lanceolato-ovatis appressis, labello convoluto indiviso apice incurvo crispato, [callo lineari in medio et altero ovato acuminato sub apice.”]

“A leafless stoloniferous plant, with tribrachiate racemes of flowers, of which the perianths are of a brownish or golden yellow colour, and the labellum white, and elegantly pencilled within, as in *D. Pierardi*.”

“ Found growing from the crevices of sandstone rocks, on the eastern side of Sydney Cove, Port Jackson, where it was observed in flower in October 1834.”—*R. C.*

“ The above is extracted from a letter from my brother, Mr. Richard Cunningham, dated August 2, 1834. He adds,

“ I send you in a bottle a Dendrobium, which, not finding noticed in either printed or manuscript enumerations in my possession I have called Cassythoides, from the *primâ facie* resemblance it has to the genus Cassytha, not only in its leafless character and short racemes of flowers, but in its peculiar chocolate bronze or japanned papulose stems—it may be found, that it also resembles it in its climbing propensities.”

It is not a little remarkable that so highly curious a plant as this should so long have been overlooked, although a native of a locality which, as Mr. Allan Cunningham remarks, has doubtless been traversed by Botanists of many countries in Europe, who have visited Port Jackson in ships of discovery since the Colony was founded in January 1788, viz., French, Spanish, German, Swedish, Russian, &c., besides many of our own countrymen.