

CON
2116

2282

HARVARD UNIVERSITY.



LIBRARY

OF THE

MUSEUM OF COMPARATIVE ZOOLOGY.

No 4340
 GIFT OF *the Connecticut*
Academy

January 25 - July 16. 1878.

TRANSACTIONS
OF THE
CONNECTICUT ACADEMY
OF
ARTS AND SCIENCES.

VOLUME III.



NEW HAVEN :
PUBLISHED BY THE ACADEMY.
1874 1874 to 1878.

Tuttle, Morehouse & Taylor, Printers, New Haven.

CONTENTS.

	PAGE
LIST OF ADDITIONS TO THE LIBRARY,	v
ART. I.—REPORT ON THE DREDGINGS IN THE REGION OF ST. GEORGE'S BANKS, IN 1872. BY S. I. SMITH AND O. HARGER. PLATES 1-8,	1
ART. II.—DESCRIPTIONS OF NEW AND RARE SPECIES OF HYDROIDS FROM THE NEW ENGLAND COAST. BY S. F. CLARK. PLATES 9-10,	58
ART. III.—ON THE CHONDRODITE FROM THE TILLY-FOSTER IRON MINE, BREWSTER, N. Y. BY E. S. DANA. PLATES 11-13,	67
ART. IV.—ON THE TRANSCENDENTAL CURVES $\sin y \sin my = a \sin x \sin nx + b$. BY H. A. NEWTON AND A. W. PHILIPS. PLATES 14-37,	97
ART. V.—ON THE EQUILIBRIUM OF HETEROGENEOUS SUBSTANCES. BY J. W. GIBBS. FIRST PART,	108
ART. VI.—THE HYDROIDS OF THE PACIFIC COAST OF THE UNITED STATES SOUTH OF VANCOUVER ISLAND, WITH A REPORT UPON THOSE IN THE MUSEUM OF YALE COLLEGE. BY S. F. CLARK. PLATES 38-41,	249
ART. VII.—ON THE ANATOMY AND HABITS OF NEREIS VIRENS. BY F. M. TURNBULL. PLATES 42-44,	265
ART. VIII.—MEDIAN AND PAIRED FINS, A CONTRIBUTION TO THE HISTORY OF VERTEBRATE LIMBS. BY J. K. THACHER. PLATES 49-60,	281
ART. IX.—EARLY STAGES OF HIPPA TALPOIDA, WITH A NOTE ON THE STRUCTURE OF THE MANDIBLES AND MAXILLÆ IN HIPPA AND REMIPES. BY S. I. SMITH. PLATES 45-48,	311
ART. X.—ON THE EQUILIBRIUM OF HETEROGENEOUS SUBSTANCES (CONCLUDED). BY J. W. GIBBS,	343

OFFICERS OF THE ACADEMY.

President,

ELIAS LOOMIS.

Vice-President,

HUBERT A. NEWTON.

Secretary,

OSCAR HARGER.

Librarian,

ADDISON VAN NAME.

Treasurer,

HENRY C. KINGSLEY.

Publishing Committee,

HUBERT A. NEWTON,

ELIAS LOOMIS,

GEORGE J. BRUSH,

ADDISON E. VERRILL,

CHESTER S. LYMAN,

WILLIAM D. WHITNEY.

Auditing Committee,

HUBERT A. NEWTON,

DANIEL C. EATON,

HENRY T. BLAKE.

ADDITIONS TO THE LIBRARY,

FROM DECEMBER 1, 1873, TO JUNE 1, 1876.

- American Association for the Advancement of Science.* Proceedings. Twenty-second, twenty-third Meetings. 8°. Memoirs. I. 4. Salem, 1874-5.
- ALBANY.—*Institute.* Transactions. Vol. VII, VIII. 1872-6. 8°. Proceedings. Vol. I. 3-4, II. 1. 1871-4. 8°.
- *New York State Cabinet of Natural History.* Twenty-third, twenty-fifth Annual Reports. 1870-2. 8°.
- *New York State Library.* Fifty-sixth Annual Report. 1874. 8°.
- BOSTON.—*American Academy of Arts and Sciences.* Proceedings. Vol. VIII-X. 1868-75. 8°.
- *Society of Natural History.* Memoirs. Vol. II. i. 1, ii. 4, iii. 1-5, iv. 1-4. 1871-6. 4°. Proceedings. Vol. XV. 4, XVI, XVII, XVIII. 1-2. 1873-6. 8°. Henz, N. M. The Spiders of the United States. (Occasional Papers. II). 1875. 8°.
- BUFFALO.—*Society of Natural Sciences.* Bulletin. Vol. I. 4, II. III. 1-2. 1874-6. 8°.
- CAMBRIDGE.—*Museum of Comparative Zoology.* Illustrated Catalogue. No. IV-VIII. 1871-4. 4°. Bulletin. Vol. II. 3-5, III. 1-14. 1871-6. 8°. Annual Report. 1870-1875. 8°.
- MINNEAPOLIS.—*Minnesota Academy of Natural Sciences.* Bulletin. 1874. 8°.
- NEW YORK.—*Lyceum of Natural History.* Annals. Vol. XI. 3-6. 1875. 8°.
- POUGHKEEPSIE.—*Society of Natural Science.* Proceedings. Vol. I. 1-2. 1876. 8°.
- ST. LOUIS.—*Academy of Science.* Transactions. Vol. III. 2. 1875. 8°.
- SALEM.—*Essex Institute.* Bulletin. Vol. V. 6-12, VI, VII. 1-7. 1873-5. 8°.
- *Peabody Academy of Science.* Memoirs. Vol. I. 4. 1875. 4°.
- SAN FRANCISCO.—*California Academy of Sciences.* Proceedings. Vol. I. 1854-7 (2d ed. 1873), III. 2. 1873. 8°.
- SAVANNAH.—*Georgia Historical Society.* Jones, C. C., Jr. Sergeant Wm. Jasper. An Address before the Georgia Historical Society, Jan. 3, 1876. 8°.
- WASHINGTON.—*Surgeon General's Office.* Annual Report of the Surgeon General, U. S. Army. 1874. 8°. Circular No. 8. Report on the Hygiene of the U. S. Army. 1875. 4°.
- *United States Naval Observatory.* Astronomical and Meteorological Observations. 1871, 1873. 4°.
- WORCESTER.—*American Antiquarian Society.* Proceedings. No. 62-65. 1874-5. 8°.

- AMSTERDAM.—*Koninklijke Akademie van Wetenschappen*. Verslagen en Mededeelingen. Afdeel. Natuurkunde. Tweede Reeks. Deel VII. 1873. 8°. Jaarboek. 1872. 8°.
- AUGSBURG.—*Naturhistorischer Verein*. Bericht XXII, XXIII. 1873-5. 8°.
- BASEL.—*Naturforschende Gesellschaft*. Bericht über die Verhandlungen. I-VIII. 1835-1849. 8°. Verhandlungen. Theil VI. 1-2. 1874-5. 8°.
- BATAVIA.—*Natuurkundige Vereeniging*. Natuurkundig Tijdschrift voor Nederlandsch Indie. Deel XXII, XXIII. 1871-3. 8°.
- *Société des Arts et des Sciences*. Tijdschrift. Vol. XX. 4-6, XXI. 1-2. 1872-4. 8°. Notulen. X. 4. XI. 1-4. 1873-4. 8°. Codicum Arabicorum (Catalogus. 1873. 8°. Alphabetische Lijst van Kaarten. 1873. 8°.
- BELFAST.—*Natural History and Philosophical Society*. Proceedings. Session 1872-3, 1873-4, 1874-5. 8°.
- BERLIN.—*Königliche Akademie der Wissenschaften*. Physikalische Abhandlungen. 1838, 1841, 1842, 1845, 1849. 4°. Mathematische Abhandlungen. 1845. 4°. Bericht über die Verhandlungen. 1854-5. 8°. Monatsbericht. 1856-9. 8°.
- BOLOGNA.—*Accademia delle Scienze dell' Istituto di Bologna*. Rendiconto. 1873-4. 8°.
- BONN.—*Naturhistorischer Verein der preussischen Rheinlande und Westphalens*. Verhandlungen. Jahrg. XXIX. 2, XXX, XXXI, XXXII. 1. 1872-5. 8°.
- BORDEAUX.—*Société des Sciences Physiques et Naturelles*. Mémoires. T. IX, X, II. Sér. I. 1. 1873-5. 8°.
- *Société Linnéenne*. Actes. T. XXVII. 2, XXVIII. 1872. 8°.
- BREMEN.—*Naturwissenschaftlicher Verein*. Abhandlungen. Bd. III. 4, IV. 1. 1873-4. 8°. Beilage. No. 3. 1873. 4°.
- BRÜNN.—*Naturforscher Verein*. Verhandlungen. Bd. V, VI, XI, XII. 1866-74. 8°.
- BRUSSELS.—*Académie Royale des Sciences, des Lettres et des Beaux-Arts de Belgique*, Mémoires. T. XL. 1873. 4°. Mémoires Couronnés et Mémoires des Savants Étrangers. T. XXXVII, XXXVIII. 1873-4. 4°. Mémoires Couronnés et Autres Mémoires. T. XXIII. 1873. 8°. Bulletins. II. Sér. T. XXXV-XXXVII. 1872-4. 8°. Annuaire. 1874. 8°. Centième anniversaire de fondation. 1872. 2 vols. 8°.
- *Observatoire Royal*. Annales. T. XXI, XXII. 1872-3. 4°.
- Observations des phénomènes périodiques. 1872. 4°. Notices extraits de l' Annuaire pour 1874. 16°. Quetelet, A. Congrès international de statistique. 1873. 4°. Quetelet, E. La comète de Coggia. 8°. pp. 10. Quetelet, E. Rapport sur l'aréographie de M. Terby. 8°. p. 6.
- *Société Entomologique de Belgique*. Annales. T. I-XIV, XVI. 1857-1873. 8°. Compte-rendu. Sér. II. No. 18. 1875. 8°.
- *Société Géologique de Belgique*. Annales. T. I. 1874. 8°.
- BUENOS AYRES.—*Academia Nacional de Ciencias Exactas*. Boletín. Entrega I. 1874. 8°.
- CALCUTTA.—*Asiatic Society of Bengal*. Journal. 1873. I. 2-4, II. 3-4; 1874; 1875, I., II. 1-3. 8°. Proceedings. 1873. v-x, 1874, 1875. 8°.
- CARLSRUHE.—*Polytechnische Schule*. Programm. 1875-6. 8°. Riffel, A. Ueber die anatomischen und physiologischen Eigenschaften der äusseren Haut. Tübingen, 1875. 8°.

- CATANIA.—*Accademia Gioenia di Scienze Naturali*. Atti. Ser. III. T. VII, VIII. 1872-3. 4°. Carta geologica della città di Catania e dintorni. Per Carmelo Sciuto-Patti. 8 Tavole.
- CHEMNITZ.—*Naturwissenschaftliche Gesellschaft*. Bericht IV. 1873. 8°.
- CHERBOURG.—*Société Nationale des Sciences Naturelles*. Mémoires. T. XVII, XVIII. 1873. 8°. Catalogue de la bibliothèque. 2^e partie, 1^{er} livr. 1873. 8°.
- CHUR.—*Naturforschende Gesellschaft Graubündens*. Jahresbericht. Neue Folge. XV-XVII. 1869-72. 8°.
- COPENHAGEN.—*Kongelige Danske Videnskabernes Selskab*. Oversigt over Forhandlinger. 1873, 1874, 1875, i. 8°.
- DANZIG.—*Naturforschende Gesellschaft* Schriften. Neue Folge. Bd. III. 2-3. 1873-4. 8°.
- DIJON.—*Académie des Sciences, Arts et Belles Lettres*. Mémoires. III. Ser. T. I. 1871-3. 8°.
- DORPAT.—*Gelehrte Estnische Gesellschaft*. Verhandlungen. Bd. VIII. 2. 1875. 8°. Sitzungsbericht. 1874. 8°.
- *Naturforscher Gesellschaft*. Sitzungsberichte. Bd. III. 1-6, IV. 1. 1869-75. 8°. Archiv für die Naturkunde Liv-, Ehst- und Kurlands. I. Ser. Bd. V. 1-4, VI, VII, 1-4. 1870-5. II. Ser. Bd. V, VII. 1-2. 1867-75. 8°.
- DRESDEN.—*Kais. Leopold.-Carolin. Deutsche Akademie der Naturforscher*. Leopoldina. Heft VII-X. 1871-4. 4°.
- *Naturwissenschaftliche Gesellschaft Isis*. Sitzungsberichte. 1873. Apr.-Dec., 1874. Jan.-Sept., 1875. Jan.-Dec. 8°.
- *Verein für Erdkunde*. Jahresbericht. X, XI, XII. 1874-5. 8°.
- DUBLIN.—*Royal Irish Academy*. Transactions. Vol. XXIV; Antiquities, Pt. ix; Vol. XXV; Science, Pt. i-xx. 1872-5. 4°. Proceedings. Vol. X. 4; Series II. Vol. II. 1-3. 1870-5. 8°.
- EDINBURGH.—*Geological Society*. Transactions. Vol. II. 3. 1874. 8°.
- EMDEN.—*Naturforschende Gesellschaft*. Kleine Schriften. XVII. Hannover, 1875. 4°. Jahresbericht. LIX, LX. 1873-4. 8°.
- ERFURT.—*Königl. Akademie gemeinnütziger Wissenschaften*. Jahrbuch. Neue Folge. Heft. VII. 1873. 8°.
- FALMOUTH.—*Royal Cornwall Polytechnic Society*. Forty-second Annual Report. 1874. 8°.
- FIRENZE.—*R. Comitato Geologico d'Italia*. Bolletino. 1873, 1874, 1875. i-iv. 8°.
- FRANKFURT A. M.—*Neue Zoologische Gesellschaft*. Der Zoologische Garten. Jahrg. XIV. 7-12, XV, XVI. 1-6. 1873-5. 8°.
- FREIBURG I. B.—*Naturforschende Gesellschaft*. Berichte. Bd. VI. 1-2. 1873. 8°.
- GENÈVE.—*Institut National Genevois*. Bulletin. T. XX. 1875. 8°.
- *Société de Physique et d'Histoire Naturelle*. Mémoires. T. XXIII, XXIV. 1. 1873-5. 4°.
- GLASGOW.—*Philosophical Society*. Proceedings. Vol. IX. 1873-5. 8°.
- GÖRLITZ.—*Naturforschende Gesellschaft*. Abhandlungen. Bd. XV. 1875. 8°.
- GÖTEBORG.—*Kongl. Vetenskaps och Vitterhets-Samhalle*. Handlingar. Ny Tidsföljd. Häftet XII-XIV. 1873-4. 8°.
- HALLE.—*Naturforschende Gesellschaft*. Abhandlungen. Bd. XII. 3-4, XIII. 2. 4°. Bericht. 1873, 1874. 4°.
- *Naturwissenschaftlicher Verein für Sachsen und Thüringen*. Zeitschrift der gesammten Naturwissenschaften. Bd. VII-X. Berlin, 1873-4. 8°.

- HAMBURG.—*Naturwissenschaftlicher Verein*. Abhandlungen. Bd. V. 4, VI. 1. 1873. 4°.
- HANNOVER.—*Naturhistorische Gesellschaft*. Jahresbericht. XXII–XXIV. 1872–4. 8°.
- HARLEM.—*Musée Teyler*. Archives. Vol. III. 3. 1873. 8°.
- HEIDELBERG.—*Naturhistorisch-Medicinischer Verein*. Verhandlungen. Neue Folge. Bd. I. 1, 3. 1874–6. 8°.
- HELSINGFORS.—*Societas Scientiarum Fennica*. Öfversigt af Förhandlingar. XIV–XVI. 1871–4. 8°. Bidrag till Kännedom af Finlands Natur och Folk. Häft. XVIII, XIX, XXI, XXII, XXIII. 1871–3. 8°.
- HERMANNSTADT.—*Siebenbürgischer Verein*. Verhandlungen. Jahrg. XXV. 1875. 8°.
- HOBART TOWN.—*Royal Society of Tasmania*. Monthly Notices for 1872. 8°.
- KÖNIGSBERG.—*Königl. physikalisch-ökonomische Gesellschaft*. Schriften. Jahrg. XIV–XV. 1873–4. 4°.
- KRAKAU.—*K. k. Sternwarte*. Materialy do Klimatografii Galicyi. Rok 1872–1874. 8°.
- LAUSANNE.—*Société Vaudoise des Sciences Naturelles*. Bulletin. II. Ser. No. 64–65, 71–75. 1870–1876. 8°.
- LEIDEN.—*Sternwarte*. Annalen. Bd. IV. Haag, 1875. 4°.
- LEIPZIG.—*Astronomische Gesellschaft*. Vierteljahrsschrift. Jahrg. VIII. 3–4, IX, X. 1–3. 1873–5. 8°. Publication XIII. 1874. 4°.
- LIÉGE.—*Société Royale des Sciences*. Mémoires. II. Ser. T. IV, V. 1873–4. 8°.
- LINZ.—*Handels- und Gewerbekammer*. Bericht. 1870, 1871, 1872. 8°. Bericht über die Lage und Bedürfnisse des Kleingewerbes in Oberösterreich. 1872. 8°.
- LIVERPOOL.—*Literary and Philosophical Society*. Proceedings. No. XXVII–XXIX. 1872–5. 8°.
- LONDON.—*Mathematical Society*. Proceedings. No. 62–86. 1873–6. 8°.
- LUXEMBOURG.—*Institut Royal Grand-Ducal*. Publications. T. XIII. Section des Sci. Nat. et Mathémat. T. XIV, XV. 1873–5. 8°. Observations Météorologiques faites à Luxembourg par F. Reuter. Vol. II. 1874. 8°.
- LYON.—*Académie des Sciences, Belles-Lettres et Arts*. Mémoires. Classe des Sciences. T. XX. 1873–4. 8°.
- MANCHESTER.—*Literary and Philosophical Society*. Memoirs. III. Ser. Vol. IV. 1871. 8°. Proceedings. Vol. VIII–XII. 1869–73. 8°.
- *Scientific Students Association*. Annual Report. 1872, 1873. 8°.
- MELBOURNE.—*Royal Society of Victoria*. Transactions and Proceedings. Vol. X, XI. 1874. 8°.
- METZ.—*Académie*. Mémoires. Année L–LV. 1868–74. 8°. Tables générales. 1819–1871. 8°.
- MEXICO.—*Sociedad de Geografía y Estadística*. Boletín. III Epoque. T. II. 5–6. 1875. 8°.
- MILANO.—*Reale Istituto Lombardo*. Rendiconti. Serie II. Vol. VI. 6–20. VII. 1–16. 1873–4. 8°.
- *Reale Osservatorio di Brera*. Pubblicazione. No. II, IV, V, VII–X. 1873–5. 4°.
- *Società Italiana di Scienze Naturali*. Atti. Vol. XV. 3–5. XVI, XVII. 1–3. 1872–5. 8°.
- MONTPELLIER.—*Académie des Sciences et des Lettres*. Mémoires. Lettres, T. V. 4; Sciences, T. VIII. 2; Médecine, T. IV. 6. 1870–2. 4°.
- MOSCOW.—*Société Impériale des Naturalistes*. Nouveaux Mémoires, T. XIII. 4, 1874. 4°. Bulletin. 1873. ii–iv, 1874, i–iv. 8°.

- MÜNCHEN.—*Königl. Bayerische Akademie der Wissenschaften*. Sitzungsberichte der philosoph.- philolog. und histor. Classe. 1872. iv-v, 1873, 1874. i-iii. 8°. Sitzungsberichte der mathemat.-physikal. Classe. 1872. iii, 1873, 1874. i-ii. 8°.
- Bietz, W. Der Antheil der königl. Bayer. Akademie an der Entwicklung der Electricitätslehre. München, 1873. 4°.
- Bischoff, T. L. W. von. Ueber den Einfluss des Freih. Justus v. Liebig auf die Entwicklung der Physiologie. München, 1874. 4°.
- Döllinger, J. von. Rede, 25 Juli, 1873. München, 1874. 8°.
- Pettenkofer, Max von. Dr. Justus Freih. von Liebig. zum Gedächtniss. München, 1874. 4°.
- Prantl, K. von. Gedächtnissrede auf F. A. Trendelenburg. München, 1873. 4°.
- Vogel, A. Justus Freih. von Liebig als Begründer der Agrikultur-Chemie. München, 1874. 4.
- *Sternwarte*. Annalen. Bd. XIX. 1873. 8°. Supplementbd. XIII. 1874. 8°.
- *Landeswirthschaftlicher Verein in Bayern*. Haus- und Landwirthschafts Kalender. 1874. 4°.
- NAPOLI.—*Società Reale di Napoli*. Accademia delle Scienze Fisiche e Matematiche. Atti. Vol. V. 1873. 4° Rendiconto. Anno IX-XI. 1870-72. 4°.
- NEU-BRANDENBURG.—*Verein der Freunde der Naturgeschichte in Mecklenburg*. Archiv. Jahrg. XXVII-XXIX. 1873-5. 8°.
- NEUCHATEL.—*Société des Sciences Naturelles*. Bulletin. T. IX, X. 1-2. 1871-5. 8°.
- OFFENBACH A. M.—*Verein für Naturkunde*. Bericht XIII, XIV. 1871-3. 8°.
- PARIS. — *Société d'Acclimatation*. Bulletin Mensuel. II Ser. T. X. 6-11, III Ser. T. I. 2-12, II, III. 1. 2. 1873-6. 8°.
- *Société Géologique de France*. Bulletin. III Ser. T. I. 1-5, II. 1-5, 7, and Tables, III. 1-2, 4-5, 8, IV. 1. 1872-6. 8°.
- *Société Américaine*. Annuaire. 1873. 8°.
- PRAG.—*Königl. böhmische Akademie der Wissenschaften*. Abhandlungen. Sechste Folge. Bd. VI, VII. 1873-4. 4°. Sitzungsberichte. 1872. ii, 1873, 1874. 8°.
- *K. k. Sternwarte*. Astronomische, magnetische und meteorologische Beobachtungen. 1873, 1874. 4°.
- PULKOWA.—*Nicolai Hauptsternwarte*. Jahresbericht. 1871-2, 1872-3. 8°.
- Dollen, W. Die Zeitbestimmung vermittelst des tragbaren Durchgangsinstruments im Verticale des Polarsterns. Zweite Abhandlung. St. Petersburg, 1874. 4°.
- QUEBEC.—*Literary and Historical Society*. Transactions. New Series. Part X, XI. 1873-5. 8°.
- REGENSBURG.—*Zoologisch-mineralogischer Verein*. Abhandlungen. Heft X. München. 1875. 8°. Correspondenz-Blatt. Jahrg. XXVII, XXVIII. 1873-4. 8°.
- *Historischer Verein vom Oberpfalz und Regensburg*. Verhandlungen. Bd. XXIX, XXX. Verzeichniss über Bd. I-XXX. 1874. 8°.
- RIGA.—*Naturforscher Verein*. Correspondenzblatt. Jahrg. XX. 1874. 8°.
- ST. GALLEN.—*Naturwissenschaftliche Gesellschaft*. Bericht. 1872-3. 8°.

- SANTIAGO.—*Universidad de Chile*. Anales. T. XXVIII–XLIV. 1866–73. 8°. Memorias presentados al Congreso Nacional de 1874, viz.: Memoria de Relaciones Exteriores e de Colonizacion; del Interior; de Justicia, Culto e Instruccion Publica; de Guerra; de Marina. Santiago and Valparaiso. 1874. 5 vols. 8°. Anuario de la Oficina Central Meteorolojica de Santiago de Chile. 1871–2. 8°. Briseño, R. Estadistica Bibliografica de la Literatura Chiliena. Santiago. 1862. 4°. Domeyko, D. Ignacio. Quarto Apendice al Reino Mineral de Chle. Santiago, 1874. 8°. Varas, J. A. Colonizacion de Llanquihue, Valdivia, Arauco. Santiago, 1872. 8°.
- ST. PETERSBURG.—*Jardin Impérial de Botanique*. Trudi I, 2, II, III. 1872–4. 8°. *Schweizerische Naturforschende Gesellschaft*. Verhandlungen in Schaffhausen, 1873. Jahresversammlung LVI. 8°.
- STOCKHOLM.—*Konigl. Svenska Vetenskaps Akademien*. Handlingar. Ny Följd. Bd. IX, 2, X, XII. 1870–3. 4°. Bibang till Handlingar. Bd. I, II, 1872–5. 8°. Öfversigt. Arg. XXVIII–XXXI. 1871–4. 8°. Meteorologiska Jagttagelser. Bd. XII–XIV. 1870–2. 4°. Minnesteckning öfver J. A. von Hartmansdorff. 1872. 8°. Minnesteckning öfver Hans Järta. 1874. 8°.
- STUTT GART.—*Verein für vaterländische Naturkunde in Württemberg*. Jahreshfte. Jahrg. XXX, XXXI. 1874–5. 8°.
- SYDNEY.—*Government Observatory*. Results of Meteorological Observations. 1872, 1873. 8°.
- TORONTO. *Magnetical Observatory*. Monthly Meteorological Register. 1873–4. 8°. General Meteorological Register. 1873–5. 8°. Abstracts and Results of Meteorological Observations, 1841–71. 1875. 8°. Third Report of the Meteorological Office of the Dominion of Canada. 1873. 8°.
- UPSALA.—*Regia Societas Scientiarum*. Nova Acta. Ser. III, T. VIII. 2. 1873. 4°. Bulletin Météorologique Mensuel. IV, V. 1–6. 1872–3. 4°.
- WIEN.—*Kaiserliche Akademie der Wissenschaften*. Sitzungsberichte. Math.-naturwiss. Classe. Abtheil. i, ii, Bd. LXIII, LXIV. Abtheil. i, Bd. LXV–LXXI. 1871–5. 8°. Wex, G. Ueber die Wasserabnahme in den Quellen, Flüssen und Strömen. Wien, 1873. 4°.
- *K. k. geologische Reichsanstalt*. Abhandlungen. Bd. V, VI, VII, 1–3, VIII. 1. 1871–5. 4°. Jahrbuch. Bd. XIX. 4, XX–XXV. 1869–75. 8°. Verhandlungen. Jahrg. 1869. No. 14–18, 1870–3, 1874. No. 1–13, 16–18, 1875. 8°. General Register zu Jahrbuch XI–XX und Verhandlungen 1860–1870. 8°. Hauer, F. v. Zur Erinnerung an Wilhelm Haidinger. 8°.
- *K. k. zoologisch-botanische Gesellschaft*. Verhandlungen. Bd. XXIII, XXIV. 1873–4. 8°.
- WIESBADEN.—*Nassauischer Verein für Naturkunde*. Jahrbücher. Jahrg. XXV–XXVIII. 1871–4. 8°.

WÜRZBURG.—*Physikalisch-medicinische Gesellschaft.* Sitzungsberichte. 1868–74. 8°. Festschrift zur Feier des fünfundzwanzigjährigen Bestehens der Gesellschaft. 1875. 4°.

Kölliker, A. Festsrede, 8. December, 1874. 8°.

ZÜRICH.—*Naturforschende Gesellschaft.* Vierteljahrsschrift. Jahrg. XIV–XVII. 1869–72. 8°.

Agardh, J. G. Till Algernes Systematik. Nya Bidrag. Lund, 1872. 4°.

From the Author.

Galle, J. G. Ueber eine Bestimmung der Sonnen-Parallaxe aus correspondirenden Beobachtungen des Planeten Flora, 1873. Breslau, 1875. 8°.

From the Author.

Haughton, S. Principles of Animal Mechanics. Second Edition. Lond., 1873. 8°.

From the Author.

Macedo, J. M. de. Notions de chorographie du Brésil. Leipzig, 1873. 8°.

Schiner, J. R. Diptera Austriaca. I. Wien, 1854.

Separatabdruck naturwiss. Abhandlungen aus den Schriften des zoolog.-botanischen Vereins, in Wien. Wien, 1856. 8°.

From Dr. V. Plöson.

Meunier, S. Cours de géologie comparée. Paris, 1874. 8°.

From the Author.

Morren, É. L'horticulture a l'Exposition Universelle de Paris de 1867. Bruxelles, 1870. 8°.

—————Rapport séculaire sur les travaux de botanique et de physiologie végétale. Bruxelles, 8°.

—————Éloge de Jean-Theodore Lacordaire. Liège, 1870. 8°.

From the Author.

Newberry, J. S. The Surface Geology of Ohio. Columbus, 1874. 8°.

—————The Structure and Relations of the Dinichthys. With two charts. Columbus, 1875. 8°.

From the Author.

Saussure, H. de. Mémoires pour servir à l'histoire naturelle de Mexique, des Antilles et des États Unis. Première livraison. Crustacées. Genève, 1858. 8°.

From the Author.

Winchell, N. H. Geological and Natural History Survey of Minnesota. Second and third Annual Reports, 1873, 1874. St. Paul, 1874–5. 8°.

From the Author.

I. REPORT ON THE DREDGINGS IN THE REGION OF ST. GEORGE'S
BANKS, IN 1872.* BY S. I. SMITH AND O. HARGER.

[Published by permission of the Superintendent of the U. S. Coast Survey.]

DURING the summer of 1872, a series of dredgings was carried on by the authors in the neighborhood of St. George's Banks. The work was undertaken at the instance of Professor Baird, United States Commissioner of Fish and Fisheries, and carried on, through the coöperation of the Coast Survey, from the steamer *Bache*, on board of which accommodations were furnished for two persons, with the necessary apparatus. On board the steamer we were received and treated throughout with the utmost courtesy by Commander J. A. Howell, and the other officers of the vessel, Lieutenants Jaques, Hagerman, Jacob and Rush; and although the dredging was carried on in connection with the special hydrographic work of the Coast Survey, all these gentlemen manifested a degree of interest in our work equal to that which they felt in their own.

On account of the lateness of the season at which operations were begun, the weather was most of the time quite unfavorable for dredging, so that the number of hauls made with the dredge was much smaller than had been expected, and no opportunities were afforded for using the large trawl or the rake dredges which had been provided, with the rest of the outfit for the natural history department of the expedition, by the United States Fish Commissioner. Still, the collections which were made from these comparatively few dredgings have proved rich and very important, giving nearly the only information which we possess of the character of the fauna of the fishing banks, and adding very largely to the knowledge of the distribution, both geographical and bathymetrical, of the marine animals of our northern coast.

* The text of this report was written and presented to the Superintendent of the Coast Survey, very nearly in its present form, in December, 1872, but its publication has been unavoidably delayed until the present. The figures illustrating some of the species mentioned have been added since the report was first prepared.

After we were obliged, late in September, to leave the expedition, Prof. A. S. Packard, Jr., and Mr. Caleb Cooke, of Salem, Mass., went in the steamer on another trip, which was mainly devoted to dredging. On this trip a number of successful hauls were made at different localities along the northern extremity of George's Bank, in 40 to 150 fathoms. The region visited on this trip was quite distant from any of the localities examined by us, and the bottom, in the deeper dredgings, was of an entirely different character, so that the collections made by Prof. Packard and Mr. Cooke contain many species not found by us, and add very greatly to the value of the results.

We wish specially to acknowledge the assistance rendered us in the preparation of this report by Professor Verrill, who has identified all the worms mentioned, and the more difficult mollusks and radiates.

The following table will facilitate references to the localities at which the dredgings were made. The letters in the first column are the same as those used by Professor Verrill in his papers in the *American Journal of Science*. When more than one haul of the dredge was made at any of the localities, the number of hauls is indicated in parenthesis.

Station.	N. Lat.	W. Long.	Depth in fath.	Nature of bottom.	Temperature.		
					Air.	Surface.	Bottom.*
<i>a</i>	41° 40'	68° 10'	25	Soft sand.			
<i>b</i>			30	" "			
<i>c</i>	41 25	66 45	28	Coarse sand.			
<i>d</i>	" "	66 24.8	50	Sand and shells.	66°	62°	45°
<i>e</i>	" "	65 58.3	60	Shells and sand.	61	58	58
<i>f</i>	" "	65 50.3	65	Dead shells.	64	60	55
<i>g</i>	" "	65 42.3	430	Sand, gravel and stones.	66	65	51
<i>h</i>	42 56.5	64 51.3	45	Gravel and stones.	64	61	36
<i>i</i>	42 44	64 36	60	Gravel, stones and sponges.	62	62	
<i>j</i>			20	Mud and fine sand.			
<i>o</i> (3)	42 5	67 49	110	Soft mud and sand.		56	49
<i>p</i>	42 3	" "	85	" " " "		56	49
<i>q</i> (2)	42 0	67 42	45	Coarse sand.			
<i>r</i>	42 3	67 31	40	" "			
<i>s</i> (2)	42 11	67 17	150	Soft sandy mud.		52	52

The first dredgings were made on the evening of August 29, to the west of George's Shoal, about latitude 41° 40' north, longitude 68° 10'

* Very little confidence can be placed in these bottom temperatures, as the Miller-Casella thermometers used did not give uniform results. Most of the temperatures are manifestly much too high.

west. The first haul, (*a*), in 25 fathoms, soft sandy bottom, gave the following species:

CRUSTACEA.

Eupagurus Bernhardus Brandt; abundant.

Crangon vulgaris Fabricius; abundant.

Conilera polita Harger (Stimpson).

Epelys montosus Smith (Stimpson).

Balanus porcatus Costa; common.

ANNELIDA.

Cistenides granulata Malmgren.

TURBELLARIA.

Meckelia lurida (?) Verrill.

MOLLUSCA.

Bela turricula (Montagu).

B. harpularia H. and A. Adams (Couthouy).

Buccinum undatum Linné; very large and abundant.

Neptunea pygmaea H. and A. Adams (Gould); abundant.

Tritia trivittata H. and A. Adams (Say); abundant.

Lunatia heros H. and A. Adams (Say).

L. immaculata H. and A. Adams (Totten).

Crepidula plana Say (*unquiformis* Stimpson); several, living.

Ensatella Americana Verrill (Gould).

Modiolaria nigra Lovén (Gray).

RADIATA.

Echinarachnius parma Gray; very abundant.

Hydractinia polyclina Agassiz.

At the second haul (*b*), in 30 fathoms, the bottom was of the same character, but a greater variety of species was obtained.

CRUSTACEA.

Cancer irroratus Say, young; common.

Eupagurus Bernhardus Brandt; abundant.

E. pubescens Brandt; common.

Crangon vulgaris Fabricius; abundant.

Pandalus annulicornis Leach; common.

Stenothoë peltata Smith, sp. nov. [Plate III, figures 5-8.]

Photis (?) sp.

Ampelisca sp.

Xenoclea megachir Smith, sp. nov. [Plate III, figures 1-4.]

Unciola irrorata Say.

Dulichia sp.

ANNELIDA.

Aphrodita aculeata Linné.

Clymenella torquata Verrill; tubes only.

TURBELLARIA.

Meckelia turida (?) Verrill.

M. ingens (?) Leidy.

MOLLUSCA.

Bela turricula (Montagu).

B. harpularia H. and A. Adams (Couthouy).

Admete viridula (O. Fabricius).

Buccinum undatum Linné; large and abundant.

Neptunea pygmaea H. and A. Adams (Gould); large and common.

Lunatia heros H. and A. Adams (Say).

“ “ var. *triseriata* (Say).

Crepidula plana Say (*unguiformis* Stimpson).

Siliqua costata H. and A. Adams (Say).

Yoldia limatula Stimpson (Say).

Modiolaria nigra Lovén (Gray).

Farella familiaris Smitt (Gros); abundant.

Gemellaria loricata Busk (Linné).

RADIATA.

Echinarachnius parma Gray; very abundant.

Asterias vulgaris Stimpson.

Campanularia verticillata Lamarck (Linné).

Sertularia cupressina Linné.

S. latiuscula? Stimpson.

Hydrallmania falcata Hincks (Linné); abundant.

Five successful hauls were made on the line of soundings running east from George's Bank, on the parallel of 41° 25' north latitude, to 63° 20' west longitude. The first of these hauls (*e*), beginning at the western end of the line, was in about longitude 66° 45' west, from 28 fathoms, coarse sandy bottom, September 16. Here the following species occurred.

CRUSTACEA.

Cancer irroratus Say, young; abundant.

Eupagurus Bernhardus Brandt.

Crangon vulgaris Fabricius; common.

Pandalus annulicornis Leach.

Ampelisca sp.

ANNELIDA.

Nereis pelagica Linné.

TURBELLARIA.

Meckelia lurida (?) Verrill.

MOLLUSCA.

Bela harpularia H. and A. Adams (Couthouy).

Tritia trivittata H. and A. Adams (Say); common.

Lanatia heros H. and A. Adams (Say), variety *triseriata*; common.

L. immaculata H. and A. Adams (Totten).

Crepidula fornicata Lamarek (Linné); one dead specimen.

Scalaria Grœnlandica Sowerby.

Clidiophora trilineata Carpenter.

Maetra solidissima Chemnitz (Gray).

Astarte castanea Say.

Crenella glandula H. and A. Adams (Totten).

Ostrea Virginica Lister; only dead specimens.

Glandula arenicola Verrill; abundant.

RADIATA.

Strongylocentrotus Dröbachiensis A. Agassiz.

Echinarachnius parma Gray.

At the second haul (*d*), longitude 66° 24'8" west, 50 fathoms, sandy and shelly bottom, August 31, the following species occurred.

PYCNOGONIDA.

Nymphon grossipes Kroyer.

CRUSTACEA.

Cancer irroratus Say; young.

Hyas coarctatus Leach; abundant.

Eupagurus Bernhardus Brandt; common.

E. Kroyerii Stimpson; common.

E. pubescens Brandt; common.

Pandalus annulicornis Leach; common.

Vetumnus serratus Goës.

Melita dentata Boeck (*Gammarus purpuratus* Stimpson).

Mæra Danæ Bate (Stimpson); common.

Cerapus rubricornis Stimpson; common.

Podocerus nitidus Stimpson.

Unciola irrorata Say; common.

Balanus porcatus Costa; common.

ANNELIDA.

Aphrodita aculeata Linné.

Harmothoë imbricata Malmgren (Linné).

Phyllodoce catenula Verrill. [Plate IV, figure 3.]

Nereis pelagica Linné; abundant.

Cistenides granulata Malmgren.

Thelepus cincinnatus Malmgren (Fabricius).

Potamilla oculifera Verrill (Leidy).

P. neglecta Malmgren.

Spirorbis nautiloides? Lamarck. [Plate IV, figure 4.]

GEPHYREA.

Phascolosoma cementarium Verrill.

MOLLUSCA.

Bela turricula (Montagu).

B. harpularia H. and A. Adams (Couthouy).

B. pleurotomaria H. and A. Adams (Couthouy).

B. decussata (Couthouy).

Neptunea curta Verrill (Jeffreys sp.; *Fusus Islandicus* Gould).

N. decemcostata (Say).

N. pygmaea H. and A. Adams (Gould).

Lunatia immaculata H. and A. Adams (Totten).

Natica clausa Broderip and Sowerby; common.

Amawopsis helicoides Stimpson; rare.

Crepidula plana Say (*unquiformis* Stimpson).

Acirsa borealis Möreh (Beck).

Margarita obscura Gould (Couthouy).

Hanleya mendicaria Carpenter (Mighels and Adams).

Entalis striolata Stimpson. [Plate I, figure 3.]

Æolis sp.

Thracia truncata Mighels and Adams.

Cyprina Islandica Lamarck (Linné).

Cardium pinnulatum Conrad; abundant.

Cyclocardia borealis Conrad; common.

Astarte castanea Say.

A. quadrans Gould.

Modiolaria discors Beck (Linné).

Pecten Islandicus Chemnitz (Müller).

Anomia aculeata Gmélin; abundant.

“ “ smooth variety.

Boltenia clavata Stimpson.

Pera crystallina Verrill (Möller); young. [Plate VIII, figure 1.]

Amarœcium sp.

Cellularia ternata Johnston (Busk).

“ “ var. *duplex* Smitt.

Caberea Ellisii Smitt (Fleming).

Bugula Murrayana Busk (Bean); abundant.

Cellaria fistulosa Linné.

Discopora Skenei Smitt.

Cellepora scabra Smitt (Fabricius).

C. ramulosa Linné, var.; with the two last species abundant on hydroid stems.

RADIATA.

Strongylocentrotus Dröbachiensis A. Agassiz.

Cribrella sanguinolenta Lütken.

Leptasterias compta Verrill.

Ophiopholis aculeata Gray; common.

Amphipholis elegans Ljungman.

Ophioglypha robusta Lyman.

Hydractinia polyclina Agassiz.

Eudendrium ramosum Ehrenberg.

E. capillare Alder (?)

Tubularia indivisa Linné.

Gonothyraea Lovéni Allman.

Campanularia verticillata Lamarek (Linné); common.

C. Hincksii Alder.

C. volubilis Alder (Linné).

Laföæa dumosa Sars (Fleming); abundant on Bryozoa.

L. gracillima G. O. Sars (Alder); with last, common.

Calycella syringa Hincks (Linné); common.

Cuspidella humilis Hincks.

Halecium Beanii Johnston.

H. tenellum Hincks.

Sertularella polyzonias Gray, var. *gigantea* Hincks; common.

S. tricuspidata Hincks (Alder); common.

Diphasia fallax Agassiz (Johnston); abundant.

Sertularia cupressina Linné; common.

S. latiuscula Stimpson.

S. abietina Linné.

Hydrallmania falcata Hincks; very abundant.

Urticina crassicornis Ehrenberg.

SPONGES.

Thecophora ibla Wyville Thompson. [Plate VII, figure 1.]

Other undetermined species.

At the third haul (*e*), longitude 65° 58'3", 60 fathoms, shelly and sandy bottom, September 16, the following were found:

CRUSTACEA.

- Hyas coarctatus* Leach; abundant.
Eupagurus Kroyerii Stimpson; abundant.
E. pubescens Brandt; abundant.
Sabinea septemcarinata Owen; one specimen.
Pandalus annulicornis Leach; common.
Paramphithoë pulchella Bruzelius (Kroyer).
Melita dentata Boeck (*Gammarus purpuratus* Stimpson).
Podocerus nitidus Stimpson.
Caprella sp.
Balanus porcatus Costa; common.

ANNELIDA.

- Harmothoë imbricata* Malmgren.
Rhynchobolus capitatus Verrill (Ørsted sp., not of Claparède).
Thelepus cincinnatus Malmgren (Fabricius).
Spirorbis nautiloides? Lamarck. [Plate IV, figure 4.]

GEPHYREA.

- Phascolosoma cementarium* Verrill.

MOLLUSCA.

- Bela violacea* (Mighels and Adams).
B. harpularia H. and A. Adams (Couthouy).
Neptunea curta Verrill (Jeffreys sp.; *Fusus Islandicus* Gould).
N. decemcostata (Say).
N. pygmæa H. and A. Adams (Gould); common.
Lunatia heros H. and A. Adams (Say), variety *triseriata*.
Natica clausa Broderip and Sowerby.
Amautropsis helicoides Stimpson; rare.
Stylifer Stimpsonii Verrill; parasitic on *Strongylocentrotus Drö-*
bachiensis A. Agassiz. [Plate I, figure 1.]
Aporrhais occidentalis Sowerby.
Acirsa borealis Mörch (Beck).
Margarita obscura Gould (Couthouy); common.
Diodora noachina Gray (Linné).
Hanleya mendicaria Carpenter (Mighels and Adams).
Cylichna alba Lovén (Brown).
Entalis striolata Stimpson; common. [Plate I, figure 3.]
Dendronotus arborescens Alder and Hancock.
Cyprina Islandica Lamarck (Linné); very abundant.
Cardium pinnulatum Conrad; common.
Astarte castanea Say.
A. quadrans Gould.

Astarte undata Gould. [Plate I, figures 6–9.]

Modiolaria corrugata Mörch (Stimpson).

Pecten tenuicostatus Mighels.

P. Islandicus Chemnitz (Müller).

Anomia aculeata Gmëlin, smooth variety; common.

Cellularia ternata Johnston (Busk); common.

Bugula Murrayana Busk (Bean).

Cellaria fistulosa Linné.

Discopora Skenei Smitt, variety.

Cellepora scabra Smitt (Fabricius); with the last on hydroïd stems.

RADIATA.

Lophothuria Fabricii Verrill.

Psolus phantapus Oken.

Strongylocentrotus Dröbachiensis A. Agassiz; abundant.

Echinarachnius parma Gray; abundant.

Crossaster papposus Müller and Troschel.

Cribrella sanguinolenta Lütken.

Hydractinia polyclina Agassiz; common.

Eudendrium capillare Alder.

E. ramosum Ehrenberg.

Tubularia indivisa Linné; common.

Campanularia verticillata Lamarek (Linné); common.

C. Hincksi Alder.

C. volubilis Alder.

Gonothyrea hyalina Hincks.

Lafoëa dumosa Sars (Fleming); common.

L. gracillima G. O. Sars (Alder).

Grammaria abietina Sars.

Coppinia areta Hincks (Dalyell); on hydroïd stems.

Sertularella tricuspidata Hincks (Alder); abundant.

Diphasia mirabilis Verrill.

D. fallax Agassiz.

Sertularia latiuscula Stimpson.

S. cupressina Linné.

Hydrallmania falcata Hincks (Linné); common.

Epizoanthus Americanus Verrill; coating shells inhabited by *Eupagurus pubescens*, and also on hydroïd stems. [Plate VIII, fig. 2.]

Urticina crassicornis Ehrenberg.

At the fourth haul (*f*), longitude 65° 50' 3", 65 fathoms, the bottom composed of dead shells, September 15, midnight, the following occurred:

CRUSTACEA.

Eupagurus Kroyerii Stimpson.

E. pubescens Brandt.

Crangon vulgaris Fabricius.

MOLLUSCA.

Bela decussata (Couthouy).

Natica clausa Broderip and Sowerby.

Stylifer Stimpsonii Verrill; parasitic on *Strongylocentrotus Dröbachiensis* A. Agassiz. [Plate I, figure 1.]

Acirsa borealis Mörch (Beck).

Margarita obscura Gould (Couthouy), variety.

Mactra solidissima Chemnitz (Gray); abundant.

Cyprina Islandica Lamarek (Linné); common.

Cyclocardia borealis Conrad.

Astarte undata Gould. [Plate I, figures 6-9.]

Crenella glandula H. and A. Adams (Totten).

RADIATA.

Strongylocentrotus Dröbachiensis A. Agassiz; common.

Echinarachnius parma Gray; abundant.

Hydractinia polyclina Agassiz.

Tubularia indivisa Linné.

Campanularia verticillata Lamarek (Linné).

Sertularella tricuspida Hincks (Alder); common.

Sertularia cupressina Linné.

Epizoanthus Americanus Verrill; coating shells inhabited by *Eupagurus pubescens*. [Plate VIII, figure 2.]

Urticina crassicornis Ehrenberg.

The fifth haul (*g*) on this line was made on the evening of September 15, to the east of the bank, in longitude $65^{\circ} 42' 3''$ west, at a depth of about 430 fathoms, on a bottom of sand, gravel, small and large stones. Here the following species occurred:

PYCNOGONIDA.

Pycnogonum littorale Müller (*pelagicum* Stimpson); common.

CRUSTACEA.

Eupagurus Kroyerii Stimpson; common.

Pandalus annulicornis Leach; several specimens.

Thysanopoda sp.; several specimens, perhaps not from the bottom.

Uciola irrorata Say; several specimens.

Melita dentata Boeck.

Scalpellum Stroemi Sars; on hydroid stems. [Plate III, fig. 9.]

ANNELIDA.

Nothria conchylega Malmgren (Sars); abundant. [Plate VII, fig. 3.]

Nephtys circumata Verrill, sp. nov.

Lumbriconereis fragilis (Ersted) (Müller).

Leodice vivida Verrill (*Eunice vivida* Stimpson). [Plate V, fig. 5.]

Rhyuchobolus capitatus Verrill (Ersted sp., not of Claparède).

GEPHYREA.

Phascolosoma tubicola Verrill.

MOLLUSCA.

Bela cancellata (Mighels and Adams).

Neptunea pygmaea H. and A. Adams (Gould).

Lunatia Groenlandica (Möller).

L. immaculata H. and A. Adams (Totten).

Natica clausa Broderip and Sowerby.

Margarita obscura Gould (Couthouy).

Diodora noachina Gray (Linné), variety *princeps* (Mighels and Adams).

Entalis striolata Stimpson. [Plate I, figure 3.]

Astarte lens Stimpson; dwarf variety.

Cryptodon obesus Verrill. [Plate I, figure 11.]

Pecten pustulosus Verrill.

Vesicularia armata Verrill; on *Sertularia argentea*.

Several other species of Bryozoa.

RADIATA.

Lophothuria squamata Verrill.

Pentacta assimilis (Duben and Koren).

Schizaster fragilis Agassiz (Duben and Koren).

Strongylocentrotus Dröbachiensis A. Agassiz; several.

Echinarachnius parma Gray; common.

Ophioglypha Sarsi Lyman; common.

Eudendrium ramosum Ehrenberg (Linné).

Tubularia indivisa Linné.

Campanularia verticillata Lamarck (Linné).

Lafoëa gracillima G. O. Sars (Alder).

Calycella producta G. O. Sars.

Halecium robustum Verrill.

Sertularella Gayi Hincks (Lamoroux).

S. tricuspidata Hincks (Alder); with reproductive capsules.

Sertularia argentea Linné, slender variety; with reproductive capsules.

Epizoanthus Americanus Verrill; upon small stones.

Urticina crassicornis Ehrenberg; young specimens.

U. nodosa Verrill (Fabricius); two large specimens.

On the line of soundings from near Cape Sable, Nova Scotia, to latitude $41^{\circ} 25'$ north, longitude $63^{\circ} 20'$ west, two successful hauls were made September 12, on Le Have Bank. The first (*h*), latitude $42^{\circ} 56.5'$ north, longitude $64^{\circ} 51.3'$ west, 45 fathoms, gravelly and stony bottom, gave the following species :

CRUSTACEA.

- Hyas coarctatus* Leach ; very abundant.
Eupagurus Kroyeri Stimpson ; abundant.
Hippolyte spina Leach ; several specimens.
H. pusiola Kroyer.
Syrrhoë crenulata Goës ; a single specimen.
Tiron acanthurus Lilljeborg ; one specimen.
Paramphithoë cataphracta Smith (Stimpson).
Tritopsis aculeata Boeck, and several undetermined Amphipods.

ANNELIDA AND GEPHYREA.

- Nychia cirrosa* Malmgren (Pallas).
Eunoë nodosa Malmgren (Sars).
Harmothoë imbricata Malmgren (Linné).
Nereis pelagica Linné ; abundant.
Nothria conchylega Malmgren (Sars) ; very abundant. [Plate VII, figure 3.]
Spiochaetopterus (?) ; tubes only.
Cistenides granulata Malmgren.
Thelepus cincinnatus Malmgren (Fabricius).
Potamilla oculifera Verrill (Leidy).
P. neglecta Malmgren ; very abundant.
Spirorbis valida Verrill, sp. nov.
S. lucidus Mörch.
Phascolosoma camentarium Verrill.

TURBELLARIA.

- Leptoplana ellipsoides* Girard.

MOLLUSCA.

- Bela violacea* (Mighels and Adams).
Admete viridula Stimpson (O. Fabricius).
Buccinum undatum Linné ; common.
Neptunea decemcostata (Say).
N. pygmaea H. and A. Adams (Gould) ; common.
Trophon Gunneri Lovén ; three specimens.
Lunatia Grœnlandica (Möller).
Natica clausa Broderip and Sowerby ; abundant.
Crepidula plana Say (*unquiformis* Stimpson) ; one alive.
Trichotropis borealis Sowerby ; abundant.

- Aporrhais occidentalis* Sowerby.
Turritella erosa Couthouy; common.
Scalaria Greenlandica Sowerby; abundant.
Margarita cinerea Gould; common.
M. Greenlandica Möller (*M. undulata* Gould).
Trachydermon album Carpenter (Montagu).
Entalis striolata Stimpson; abundant. [Plate I, figure 3.]
Dendronotus arboreseens Alder and Hancock.
Mya truncata Linné.
Cardium pinnulatum Conrad; common.
Cyclocardia Novangliæ Morse; common.
Astarte elliptica (Brown); very abundant. [Plate I, figure 10.]
A. Banksii Leach; common. [Plate I, figure 12.]
A. undata Gould; common. [Plate I, figures 6–9.]
Pecten Islandicus Chemnitz (Müller); abundant.
Boltenia Bolteni.
Cynthia carnea Verrill.
Terebratulina septentrionalis (Couthouy); common.
Rhynchonella psittacea (Gmélin).
Myriozoum coarctatum Smitt (Sars); common.
Eschara papposa Packard.
Escharoides rosacea Smitt.
Cellepora avicularis Hincks.

RADIATA.

- Lophothuria Fabricii* Verrill.
Strongylocentrotus Dröbachiensis A. Agassiz; common.
Crossaster papposus Müller and Troschel; young.
Pteraster militaris Müller and Troschel.
Ophiopholis aculeata Gray; common.
Ophioglypha Sarsii Lyman.
O. robusta Lyman.
Clytia Johnstoni Hincks (Alder).
Hydractinia polyelina Agassiz; abundant.
Eudendrium capillare Alder.
Tubularia indivisa Linné; common.
Campanularia verticillata Lamarck (Linné).
C. Hincksii Alder.
Lafoëa gracillima G. O. Sars.
Calycella syringa Hincks (Linné).
Sertularella tricuspidata Hincks (Alder); common.
S. polyzonias Gray, variety *gigantea* Hincks.
Thuiaria articulata Fleming (Pallas).
Urticina crassicornis Ehrenberg; abundant.

At the second haul (*i*), latitude 41° 44' north, longitude 64° 36' west, 60 fathoms, coarse gravel, stones, and sponges, the following occurred:

HYDROZOÏDA.

Nymphon grossipes Kroyer.

CRUSTACEA.

Hyas coarctatus Leach.

H. araneus Leach.

Eupagurus Kroyerii Stimpson; abundant.

Sabinea septemcarinata Owen; two specimens.

Tritopsis aculeata Boeck.

Acanthozone cuspidata Boeck.

ANNELIDA.

Eunoë nodosa Malmgren (Sars).

Harmothoë imbricata Malmgren (Linné).

Lagisca rarispina Malmgren (Sars).

Nothria conchylega Malmgren (Sars); abundant. [Plate VII, fig. 3.]

Thelepus cincinnatus Malmgren (Fabricius).

Potamilla neglecta Malmgren; abundant.

Spirorbis valida Verrill.

MOLLUSCA.

Admete viridula Stimpson (O. Fabricius); common.

Trophon Gunneri Lovén.

Natica clausa Broderip and Sowerby.

Aporrhais occidentalis Sowerby.

Turritella reticulata Mighels and Adams.

Margarita cinerea Gould.

Hanleia mendicaria Carpenter (Mighels and Adams).

Trachydermon album Carpenter (Montagu).

Entalis striolata Stimpson; abundant.

Æolis rufibranchialis Alder and Hancock (?).

Cardium pinnulatum Conrad.

Terebratulina septentrionalis (Couthouy); common.

Amarœcium glabrum Verrill.

Discopora Skenei Smitt.

RADIATA.

Crëbrella sanguinolenta Lütken.

Ophiopholis aculeata Gray; abundant.

Ophioglypha Sarsii Lyman.

Lafoëa gracillima G. O. Sars.

Sertularella tricuspida Hincks.

Sertularella polyzonias Gray, variety *gigantea* Hincks.
Hydrallmania falcata Hincks, var. *tenerrima* (Stimpson).
Aglaophenia myriophyllum Lamoroux (Linné).
Grammaria abietina Sars.

Many species of sponges were also obtained, but most of them are as yet undetermined. Among them is *Thecophora ibla* W. Thompson.

On leaving Halifax, N. S., September 11, one haul (*j*) was made just off Chebucto Head, in 20 fathoms, soft mud and fine sand with decaying sea weed. Here the following were found:

CRUSTACEA.

Hyas araneus Leach; common.
Eupagurus pubescens Brandt.
Diastylis quadrispinosa G. O. Sars; common.
D. sculpta G. O. Sars.
Halirages fulvocinctus Boeck (Sars).
Gammarus ornatus Edwards; perhaps from floating sea-weed.
Ampelisca sp.; common.
Monoculodes borealis Boeck.

ANNELIDA.

Harmothoë imbricata Malmgren (Linné).
Goniada maculata Ersted.
Brada sp.
Cistenides granulata Malmgren.

MOLLUSCA.

Aporrhais occidentalis Sowerby; common.
Turritella reticulata Mighels and Adams; common.
Margarita varicosa Mighels and Adams; common.
M. obscura Gould (Couthouy), variety.
Thracia myopsis Beek.
Macoma proxima (Gray).
Astarte elliptica (Brown). [Plate I, figure 10.]
Anomia aculeata Gmélin.
Terebratulina septentrionalis (Couthouy).
Rhynchonella psittocæa (Gmélin).
Crisia eburnea Lamoroux (Linné). [Plate II, figures 3-4.]
Flustra papyrea (Pallas).

RADIATA.

Ophiopholis aculeata Gray.
Ophioglypha robusta Lyman.
Manania auricula Clark (?)
Hydrallmania falcata Hincks (Linné).

The dredging while Dr. Packard and Mr. Cooke were on board the Bache was at five localities. First (*o*), just on the northwestern border of George's Bank, latitude 42° 5' north, longitude 67° 49' west, in 110 fathoms, three hauls were made from a bottom of soft sandy mud with a few stones, and the following collected :

CRUSTACEA.

- Caridion Gordoni* Goës; one small specimen.
Harpina fusiformis (Stimpson); common.
Stegocephalus ampulla Bell; one large specimen.
Unciola irrorata Say; common.
Anthura brachiata Stimpson.

ANNELIDA.

- Lænilla?* *mollis* G. O. Sars.
Pholoë minuta Malmgren.
Nephtys ciliata Malmgren (Müller). [Plate V, figure 1.]
N. ingens Stimpson.
Phyllodoce sp.
Eteone depressa Malmgren (?).
Nereis pelagica Linné; common.
Lumbriconereis fragilis Örsted (Müller). [Plate V, figure 2.]
Ninoë nigripes Verrill. [Plate V, figure 3.]
Leodice vivida Verrill (Stimpson). [Plate V, figure 5.]
Nothria conchylega Malmgren (Sars). [Plate VII, figure 3.]
N. opalina Verrill; common. [Plate VII, figure 4.]
Goniada maculata Örsted.
Rhynchobolus capitatus Verrill (Örsted sp., not of Claparède).
Ammotrypane fimbriata Verrill.
Eumenia crassa Örsted.
Trophonia aspera Verrill (Stimpson).
Sternaspis fossor Stimpson.
Scolecolepis cirrata Malmgren (Sars).
Notomastus latericius Sars.
Ancistria capillaris Verrill.
Maldane Sarsii Malmgren.
Rhodine Lovéni Malmgren.
Nicomache lunbriculis Malmgren (Fabricius).
Aziothea catenata Malmgren (?).
Praxilla prætermissa Malmgren.
P. gracilis Malmgren.
P. species undetermined.
Ammochares assimilis Sars. [Plate V, figure 4.]
Ampharete arctica Malmgren.

- Ampharete Finnarchica* (?) Malmgren (Sars).
A. gracilis Malmgren.
Amphicteis Gunneri Malmgren (Sars); abundant.
Samytha sexcirrata Malmgren (Sars).
Samythella elongata Verrill.
Melinna cristata Malmgren (Sars.)
Amphitrite cirrata Müller.
Pista cristata Malmgren (Müller). [Plate IV, figure 2.]
Grymæa spiralis Verrill. [Plate IV, figure 1.]
Terebellides Stroemi Sars.
Polycirrus sp.
Sabella pavonia (?) Malmgren.
Potamilla neglecta Malmgren.
Protula media Stimpson; tubes only. [Plate VI.]
P. borealis (?) Sars; tubes only.

GEPHYREA.

- Phascolosoma cæmentarium* Verrill.
P. boreale Keferstein (?).
P. tubicola Verrill.
Chætoderma nitidulum Lovén. [Plate VIII, figures 3-4.]

TURBELLARIA.

- Meckelia lurida* (?) Verrill.

MOLLUSCA.

- Pleurotomella Puckardii* Verrill; one living.
Admete viridula Stimpson (O. Fabricius).
Neptunea pygmaea H. and A. Adams (Gould); common.
Ringicula nitida Verrill. [Plate I, figure 2.]
Lunatia Greenlandica (Möller).
Natica clausa Broderip and Sowerby.
Velutina laevigata (Linné).
Margarita cinerea Gould.
Lepeta cæca Müller.
Trachydermon album Carpenter (Montagu).
Cylichna alba Lovén (Brown).
Philine sp.
Entalis striolata Stimpson; common.
Neera arctica Sars.
Thracia myopsis Beck.
Macoma proxima (Gray).
Cardium pinnulatum Conrad.
Astarte lens Stimpson, dwarf var.; common.

- Cryptodon Gouldii* H. and A. Adams (Philippi).
Yoldia obesa Stimpson.
Leda tenuisulcata Stimpson (Couthony); common.
Crenella glandula H. and A. Adams (Totten).
Pecten Islandicus Chemnitz (Müller).
P. tenuicostatus Mighels.
Anomia aculeata Gmélin.
Ascidiopsis complanata Verrill (Fabricius).
Terebratulina septentrionalis (Couthony).
Discofascigera lucernaria Sars.
Cellularia sp.
Bugula avicularia Busk, variety *fastigiata*.

RADIATA.

- Lophothuria Fabricii* Verrill.
Pentacta assimilis (Duben and Koren).
Thyone scabra Verrill.
Schizaster fragilis Agassiz (Duben and Koren); several.
Ctenodiscus crispatus Duben and Koren.
Ophioglypha affinis Lyman.
O. Sarsii Lyman.
Ophiopholis aculeata Gray.
Ophiacantha spinulosa Müller and Troschel.
Archaster arcticus Sars.
Pennatula aculeata Danielsen.
Cerianthus borealis Verrill. [Plate II, figure 5.]

Second (*p*), a little to the southeast of the first, latitude 42° 3', longitude 67° 49', 85 fathoms, one haul from a bottom of the same character as at first locality:

CRUSTACEA.

- Harpina fusiformis* Smith (Stimpson); common.

ANNELIDA.

- Antinoë Sarsii* Kinberg.
Nephtlys ingens Stimpson.
N. circinata Verrill, sp. nov.
Lumbriconereis fragilis (Ersted) (Müller). [Plate V, figure 2.]
Nothria conchylega Malmgren (Sars). [Plate VII, figure 3.]
Ammotrypane fimbriata Verrill.
Eumenia crassa (Ersted).
Trophonia aspera Verrill (Stimpson).
Sternaspis fossor Stimpson.
Chaetozone setosa Malmgren.

Ancistria capillaris Verrill.
Maldane Sarsii Malmgren.
Rhodine Lovéni Malmgren.
Præcilla prætermissa Malmgren.
Ammochares assimilis Sars.
Ampharete arctica Malmgren.
Amphicteis Sundevalli Malmgren.
Terebellides Stroemi Sars.
Polycirrus sp.

GEPHYREA.

Phascolosoma cementarium Verrill.

MOLLUSCA.

Natica clausa Broderip and Sowerby.
Scalaria Greenlandica Sowerby.
Yoldia obesa Stimpson.
Y. thraciformis Stimpson (Storer).

RADIATA.

Schizaster fragilis Agassiz (Duben and Koren).
Edwardsia sp.

Third (*q*), still farther to the southeast, latitude 42°, longitude 67° 42', two hauls in 45 fathoms, coarse sandy bottom :

CRUSTACEA.

Hyas coarctatus Leach ; very abundant.
Cancer irroratus Say ; one young specimen.
Eupagurus Bernhardus Brandt.
E. Kroyeri Stimpson ; common.
E. pubescens Brandt ; common.
Crangon vulgaris Fabricius ; abundant.
Hippolyte pusiola Kroyer.
Pandalus annulicornis Leach ; common.
Vertumnus serratus Goës.
Parumphithoë cataphracta Smith (Stimpson).
P. pulchella Bruzelius.
Phoxus Kroyeri Stimpson.
Melita dentata Boeck.
Pontogeneia inermis Boeck ; one specimen.
Cerapus rubricornis Stimpson.
Xenoclea megachir Smith, sp. nov. [Plate III, figures 1-4.]
Unciola irrorata Say.
Balanus porcatus Costa.

ANNELIDA.

- Harmothoë imbricata* Malmgren.
Lagisca propinqua Malmgren.
Eusyllis phosphorea Verrill, sp. nov. [Plate VII, figure 2.]
Nereis pelagica Linné.
Leodice vivida Verrill (Stimpson). [Plate V, figure 5.]
Nothria conchylega Malmgren (Sars). [Plate VII, figure 3.]
Ammochares assimilis Sars. [Plate V, figure 4.]
Amphitrite Greenlandica Malmgren.
Thelepus cincinnatus Malmgren (Fabricius).
Chone infundibuliformis Kroyer.
Spirorbis nautiloides Lamarek?

MOLLUSCA.

- Bela harpularia* H. and A. Adams (Couthouy).
B. pleurotomaria H. and A. Adams (Couthouy).
B. turricula (Montagu).
Buccinum undatum Linné.
Neptunea curta Verrill (Jeffreys sp., *Fusus Islandicus* Gould).
N. pygmaea H. and A. Adams (Gould).
Lunatia heros H. and A. Adams, variety *triseriata* (Say).
L. immaculata H. and A. Adams (Totten).
Turritella erosa Couthouy.
T. acicula Stimpson.
Margarita obscura Gould (Couthouy); common.
Diodora noachina Gray (Linné).
Hanleyia mendicaria Carpenter; large specimens.
Maetra solidissima Chemnitz (Gray).
Cyprina Islandica Lamarek (Linné).
Cardium pinnulatum Conrad; common.
Astarte quadrans Gould.
Leda tenuisulcata Stimpson (Couthouy); common.
Crenella glandula H. and A. Adams (Totten).
Modiola modiolus Turton (Linné).
Modioluria laevigata (Gray).
Pecten tenuicostatus Mighels.
P. Islandicus Chemnitz (Müller).
Anomia aculeata Gmélin.
Glandula arenicola Verrill.
Cellularia ternata Johnston (Busk).
Gemellaria loricata Busk (Linné).
Bugula Murrayana Busk (Bean).
Cellepora tuberosa D'Orbigny.

RADIATA.

- Lophothuria Fabricii* Verrill.
Strongylocentrotus Dröbachiensis A. Agassiz; common.
Echinarachnius parma Gray; very abundant.
Solaster endeca Forbes.
Crossaster papposus Müller and Troschel.
Cribrella sanguinolenta Lütken.
Asterias vulgaris Stimpson.
Leptasterias Stimpsoni Verrill.
Ophiopholis aculeata Gray; common.
Ophioglypha robusta Lyman.
Eudendrium ramosum Ehrenberg (Linné).
E. capillare Alder.
Campanularia verticillata Lamarek; common.
C. Hincksii Alder.
Laföëa dumosa Sars (Fleming); common.
Calycella syringa Hincks (Linné).
Grammaria abietina Sars (*G. robusta* Stimpson).
Coppinia arcta Hincks; on hydroid stems.
Halecium labrosum Alder.
Sertularella polyzonias Gray (Linné); common.
 " " variety *gigantea* Hincks.
S. tricuspida Hincks (Alder); abundant.
Sertularia abietina Linné; one, very large.
S. latiuscula Stimpson.
S. cypressina Linné; common.
S. argentea Ellis and Solander. [Plate III, figure 2.]
Hydrallmania fulcata Hincks; abundant.

Third (*r*), north and a little east of the last, latitude $42^{\circ} 3'$, longitude $67^{\circ} 31'$, in 40 fathoms, coarse sandy bottom:

CRUSTACEA.

- Eupagurus Bernhardus* Brandt.

ANNELIDA.

- Dodecaceria concharum* (Ersted).
Spirorbis quadrangularis Stimpson.

MOLLUSCA.

- Bela harpularia* H. and A. Adams (Couthouy).
Natica clausa Broderip and Sowerby.
Scalaria Greenlandica Sowerby.
Margarita obscura Gould (Couthouy).
Mactra solidissima Chemnitz (Gray).

Cyprina Islandica Lamarek (Linné).

Cardium pinnulatum Conrad.

Astarte castanea Say.

Pecten tenuicostatus Mighels.

Cellularia sp.

Bugula Murrayana Busk (Bean).

RADIATA.

Echinarachnius parma Gray.

Hydractinia polyclina Agassiz.

Sertularella polyzonias Gray, var. *gigantea* Hincks.

SPONGES.

Chalina oculata Bowerbank, and a massive siliceous sponge.

Fifth (s) a little to northeast of the bank, latitude $42^{\circ} 11'$, longitude $67^{\circ} 17'$, two hauls in 150 fathoms, soft sandy mud with a few pebbles:

CRUSTACEA.

Hyas coarctatus Leach.

Eupagurus Bernhardus Brandt.

E. Kroyeri Stimpson.

E. pubescens Brandt.

Ptilocheirus pinguis Stimpson.

Æga psora Bate and Westwood.

Conilera polita Harger (Stimpson).

Balanus porcatus Costa.

ANNELIDA.

Lætmonice flicornis Kinberg.

Harmothoë imbricata Malmgren (Linné).

Antinoë angusta Verrill, sp. nov.

Eucranta villosa Malmgren.

Nephtys ingens Stimpson.

Lumbriconereis fragilis Ørsted (Müller). [Plate V, figure 2.]

Nothria conchylega Malmgren (Sars). [Plate VII, figure 3.]

N. opalina Verrill. [Plate VII, figure 4.]

Goniada maculata Ørsted.

Scalibregma inflatum Rathke.

Spiochaetopterus (?); tubes exactly like those of this genus.

Scolecopsis cirrata Malmgren (Sars).

Ancistria capillaris Verrill.

Maldane Sarsii Malmgren; abundant.

- Cistenides granulata* Malmgren.
Amphicteis Gunneri Malmgren (Sars).
Amage auricula Malmgren.
Smythella elongata Verrill.
Melinna cristata Malmgren (Sars).
Pista cristata Malmgren (Sars). [Plate IV, figure 2.]
Grymæa spiralis Verrill. [Plate IV, figure 1.]
Terebellides Stroemi Sars.
Protula borealis Sars?; tubes only.

GEPHYREA.

- Phascolosoma cæmentarium* Verrill.
P. boreale Keferstein (?)
P. tubicola Verrill.

MOLLUSCA.

- Admete viridula* Stimpson (Müller).
Neptunea pygmea H. and A. Adams (Gould); abundant.
Ringicula nitida Verrill; one living. [Plate I, figure 2.]
Lunatia Grœnlandica (Möller).
Natica clausa Broderip and Sowerby.
Volutina zonata Gould.
Torrellia vestita Jeffreys; one specimen.
Aporrhais occidentalis Sowerby.
Margarita cinerea Gould.
M. obscura Gould (Couthouy).
Trachydermon album Carpenter.
Scaphander puncto-striatus Stimpson; one very large.
Cylichna alba Lovén (Brown).
Entalis striolata Stimpson; abundant. [Plate I, figure 3.]
Dentalium occidentale Stimpson; one specimen.
Nuxa arctica Sars.
Thracia myopsis Beck.
Cardium pinnulatum Conrad.
Astarte quadrans Gould.
A. lens Stimpson. dwarf variety; common.
Yoldia obesa Stimpson.
Leda tenuisulcata Stimpson (Couthouy).
Arca pectunculoides Scacchi; several.
Pecten pustulosus Verrill; one living.
Anomia aculeata Gmélin.
Glandula arenicola Verrill; common.
Terebratulina septentrionalis (Couthouy); abundant.

- Hornerea lichenoides* Smitt (Linné).
Discoporella verrucaria Smitt.
Cellularia ternata Johnston, var. *gracilis* (*arctica* Busk sp.).
C. Peachii Busk.
Caberea Ellisii Smitt (Fleming).
Bugula Murrayana Busk (Bean); abundant.
B. avicularia Busk, var. *fastigiata*.
Cellepora scabra Smitt (Fabricius).
C. ramulosa (Linné).

RADIATA.

- Thyone scabra* Verrill.
Schizaster fragilis Agassiz (Duben and Koren); several.
Echinarachnius parma Gray.
Solaster fureifer Duben and Koren; one specimen.
Archaster arcticus Sars; one specimen.
Ophioglypha Sarsii Lyman.
O. affinis Lyman.
Ophiacantha spinulosa Müller and Troschel.
Clytia Johnstoni Hincks.
Eudendrium capillare Alder.
Sertularella tricuspida Hincks (Alder).
S. Gayi Hincks (Lamoroux).
Sertularia epressina Linné.
Pennatula aculeata Danielsen.
Virgularia Lyungmanii Kölliker.
Bolocera Tuediæ Gosse; tentacles only.
Urticina crassicornis Ehrenberg.
Cerianthus borealis Verrill. [Plate II, figure 5.]

The lists of species from all the localities (*a, b, c, d, e, f, q, r,*) on George's Bank itself, show that the fauna of that region is almost exactly the same as in the Bay of Fundy, at the same depths and on similar bottom. To be sure, on the one hand, several arctic species, not yet found in the Bay of Fundy, occurred upon the Bank; but on the other hand, several apparently more southern forms were found, as the species of *Crepidula* and *Stylifer*. The two dredgings (*h* and *i*) upon LeHave Bank seem to indicate, as we might expect, a somewhat more arctic fauna than that upon George's Bank, since several arctic species, not known from George's Bank or the Bay of Fundy, occurred there, though *Crepidula plana* was also found.

The dredgings in deep water near the Banks indicate a fauna quite different from that upon the Banks themselves. This is undoubtedly

partially owing to the difference in the character of the bottom as well as to the difference in depth. Of the species occurring in deep water, a much larger proportion than in the shallower waters are the same as those of northern Europe. At the greatest depth reached by the dredge, about 430 fathoms, at the locality (*g*) east of George's Bank, almost all the species which were not also found in shallow water are European. Some of these species, however, were dredged in 1872, by Prof. Verrill, in the central part of the Bay of Fundy, east of Grand Menan Island, in 95 to 106 fathoms, where the character of the bottom was quite similar to that of our deepest dredging.

At each of the three deepest of Dr. Packard's dredgings, (*o*) 110 fathoms, (*p*) 85 fathoms, and (*s*) 150 fathoms, the bottom was composed of soft sandy mud, very different in character from that at any of the localities examined by us. The fauna of the bottom at these three places was essentially the same, and, although many of the species, on account of the character of the bottom, were different from those at the locality in 430 fathoms, about the same proportion are identical with European species.

Although the dredgings in deep water were so few, the facts presented in the foregoing lists with reference to the bathymetrical distribution of species, are important and very interesting. Of the species enumerated from 430 fathoms, considerably more than half are well known shallow water forms, many of them even occurring between tides in the Bay of Fundy and at other points on the coast, while nearly all the species mentioned are also found at less than 50 fathoms depth. The same remarks will apply to the deeper dredgings of Dr. Packard and Mr. Cooke. The species from the deepest dredging belong apparently to as highly organized groups of animals as do those from shallow water. We were not able to detect any decrease in the intensity of the colors in individuals from this depth. The colors of *Pandanus annulicornis*, *Eupagurus Kroyeri*, *Unciola irrorata*, and *Urticina crassicornis*, all brightly-colored species, seemed to have lost none of their intensity at the depth of 430 fathoms.

Besides the investigation of the fauna of the bottom by means of the dredge, every opportunity was employed for collecting those animals which live in part or wholly at the surface of the water. Notwithstanding the unfavorable character of the weather during most of the time we were at sea, towing nets were used, whenever soundings were being made, and usually with very good results. Nets of small size were several times successfully used even when the

steamer was under weigh. In this way a great number of surface species were collected, and a large proportion of them are additions to the fauna of our coast. Many of these species belong to genera previously known only from much farther south, or from the eastern or southern Atlantic, while quite a number are undescribed.

August 29, on and near Cultivator Shoal (*k*), where the surface temperature of the water was 62°, the following were taken: *Trachynema digitale* A. Agassiz, *Pleurobrachia rhododactyla* Agassiz, species of *Sagitta* and *Autolytus*, several species of Copepod crustacea, *Calliopius leviusculus* Boeck (among floating rock-weed), the young of some Brachyuran in the zoëa and megalops stages of growth, and a species of *Motella* (?).

East of George's Bank, in latitude 41° 20' to 30', longitude 63° to 63° 30', September 14, during the day, many species were taken, but as they all occurred, with many additional species, on the following day, it is not necessary to enumerate them separately.

On the evening of September 14, from nine to ten o'clock, still east of the Bank (*m*), in latitude 41° 25', longitude 63° 55', while the surface temperature was 65°, the following forms occurred: *Pleurobrachia* sp.; a species of *Salpa* in abundance; several species of Heteropods and Pteropods, among the latter *Spirialis Gouldii* Stimpson, and species of *Styliola*; a species of *Sagitta*; a species of *Sapphirina* and a great many other Copepods; species of *Hyperia*, *Phrosina*, and of another allied genus; a species of *Thysanopoda*, which was beautifully phosphorescent; young Brachyura in the zoëa and megalops stages, and the young of some Macrouran.

September 15, on the same line of soundings, in latitude 41° 25', longitude 65° 5' to 30', the temperature of the water varying from 66° to 70°, but most of the time at the latter point, very many species occurred, and among them the following: *Physalia pelagica* Lamarck (Portuguese man-of-war), *Cestum Veneris* Lesueur (?), *Stomolophus meleagris* Agassiz, *Charybdea periphylla* Péron and Lesueur, *Pelagia cyanella* Péron and Lesueur; species of *Salpa* and *Sagitta* in great abundance; *Lepas pectinata* Spengler and *L. fascicularis* Ellis and Solander; two species of *Sapphirina* and many other genera of Copepoda; species of *Oxycephalus*, *Platyscelus*, *Pronoë*, *Anchylomera*, *Thyropus*, *Phronima* (?), and *Hyperia*; *Calliopius leviusculus* Boeck, common among floating rock-weed; species of *Lucifer* and *Mysis*; *Latreutes ensiferus* Stimpson, *Nautilograpsus minutus* Edwards, and *Neptunus Sayi* Stimpson among gulf-weed, and the latter frequently seen swimming at some distance from the sea-weed; three species of Heteropods and ten species of Pteropods, all new to our coast.

Among the Pteropods are *Styliola acus* (Eschscholtz sp.), and four other species of the same genus, two of *Pleuropus*, *Spirialis Gouldii*, etc. Many of these species and genera are quite new to the fauna of the United States, and nearly all of them to the coast of New England. They are nearly all, as far as known, characteristic Gulf Stream forms.

*Notes on some of the Species enumerated; by S. I. SMITH.**

CRUSTACEA.

Eupagurus Bernhardus Brandt.

Pagurus Bernhardus (Linné sp.) Fabricius, Entomologia systematica, ii, p. 469, 1793, and Supplementum, p. 411, 1798.

Pagurus (subgenus *Eupagurus*, section *Streptodactylus*) *Bernhardus* Brandt, Middendorff's Sibirische Reise, Krebse, p. 106, 1851.

Eupagurus Bernhardus Stimpson, Crust. Pacific Shores of North America, Journal Boston Soc. Nat. Hist., vi, p. 483 (separate copies, p. 43), 1857.

I have recently † wrongly given Stimpson as authority for this and the next species, not having at the time access to Brandt's work, and not being able to comprehend his absurdly complex nomenclature from the quotation of his names by other authors.

Eupagurus pubescens Brandt.

Pagurus pubescens Kroyer (in part), Grönlands Amphipoder, p. 68, 1838, and Naturhistorisk Tidsskrift, ii, p. 251, 1839.

Pagurus (subgenus *Eupagurus*, section *Orthodactylus*) *pubescens* Brandt, op. cit., p. 111, 1851.

Eupagurus pubescens Stimpson, Prodrômus descriptionis Animalium evertibratorum, etc., Proceedings Acad. Nat. Sci., Philadelphia, 1858, p. 237 (separate copies, p. 75), 1859, and Notes on North American Crustacea, Annals Lyceum Nat. Hist., New York, vii, p. 89 (separate copies, p. 43), 1859.

This species is common on our eastern coast north of Cape Cod, but is not quite as abundant as the last species and is seldom found at low water. South of Cape Cod it is apparently confined to the deeper and colder waters.

* With the exception of the portion relating to the Crustacea, these notes have had the benefit of Professor Verrill's revision, and the descriptions of all the new species have been copied from his published papers, or, in the case of those here for the first time described, have been prepared by him specially for these pages, and are marked by his initials.

† Report upon the Invertebrate Animals of Vineyard Sound, in Report of the U. S. Commissioner of Fish and Fisheries, Part I, 1873 (published in 1874).

Eupagurus Kroyeri Stimpson.

Notes on North American Crustacea, *Annals Lyceum Nat. Hist.*, vii, p. 89 (43)^r 1859.

This species is very closely allied to the last and is very easily confounded with it, especially when young. The differences in the relative proportions of the chelipeds and ambulatory legs, given by Stimpson, will not hold for distinguishing the two species, but the differences in the amount of pubescence and especially in the form and armature of the chelipeds seem to be constant characters, sufficient for distinguishing them.

The *Kroyeri* has about the same range, on our coast, as the last species, although I have never seen it south of Cape Cod, but is apparently less abundant and more confined to the deeper waters.

Sabinea septemcarinata Owen (*Sabine* sp.)

This species was dredged in 68 fathoms off Casco Bay in the summer of 1873. It has also been found by Mr. Whiteaves in the Gulf of St. Lawrence and by Dr. Packard on the coast of Labrador. It is an exceedingly arctic and circumpolar species.

Caridon Gordoni Goës (*Bate* sp.?)

Goës, *Crustacea decapoda podophthalma marina Sueciæ* (from *Æfversight af Kongl. Vetenskaps-Akad. Förhandlingar*, Stockholm, 1863), p. 10.

We have dredged this species in 50 fathoms in the Bay of Fundy, and Dr. Packard and Mr. Cooke obtained it on Cashe's Ledge in 1873.

Our specimens agree well with the detailed description given by Goës, except that they have a well developed epipodus ("flagellum") upon the second, third and fourth cephalothoracic legs, as in some species of *Hippolyte*, while Goës says of the second legs, "nec palpo nec (quoad viderim) flagello ullo instructis," and of the third to fifth, "flagellum basale nullum inspicere potui." From the guarded manner in which Goës mentions these wholly negative characters, I am inclined to regard them as doubtful. Our specimens agree so completely in all other respects that it seems highly improbable that they should be distinct from the European species.

Diastylis quadrispinosa G. O. Sars.

Æfversight af Kongl. Vetenskaps-Akademiens Förhandlingar, 1871, Stockholm, p. 27; and *Beskrivelse af de Paa Fregatten Josephiens Expedition Fundne Cumaceer*, in *Kongl. Svenska Vetenskaps-Akademiens Handlingar*, ix, p. 28, plates 10, 11, figs. 51-61, 1871.

This is the most abundant species of the genus from off Buzzard's Bay and Vineyard Sound to Nova Scotia. It ranges north at least as far as the Gulf of St. Lawrence.

Diastylis sculpta G. O. Sars.

Loc. cit., *Øfversigt*, p. 71; *Handlingar*, p. 24, pls. 1-9, figs. 1-49.

This species is not uncommon in Casco Bay and the Bay of Fundy.

Phoxus Kroyeri Stimpson.

Marine Invertebrates of Grand Manan, p. 58, 1853.

We have dredged this species in 10 to 29 fathoms in and off Vineyard Sound, on sandy and muddy bottoms in shallow water in Casco Bay, and have found it from low water to 20 fathoms in the Bay of Fundy. Mr. Whiteaves has dredged it in the Gulf of St. Lawrence in 200 fathoms, muddy bottom.

Our species is very closely allied to, and probably identical with, the *P. Holbollii* Kroyer which is found in Greenland, Iceland and northern Scandinavia.

Harpina fusiformis Smith.

Phoxus fusiformis Stimpson, Marine Invertebrates of Grand Manan, p. 57, 1853.

This species is very likely identical with the *H. plumosa* Boeck (*Phoxus plumosus* Kroyer), which has very nearly the same range as *Phoxus Holbollii*.

We have dredged our species in 20 to 60 fathoms, muddy bottom, in the Bay of Fundy. Mr. Whiteaves has dredged it frequently, in the Gulf of St. Lawrence.

Stenothoe peltata Smith, sp. nov.

Plate IV, figures 5 to 8.

Female. Eyes round and nearly white in alcoholic specimens. Antennulæ considerably shorter than the epimera of the fourth segment; first segment of the peduncle stout, fully as long as the head, the second shorter, and the third very short and like the segments of the flagellum; flagellum scarcely longer than the peduncle, composed of about eight segments. Antennæ slightly longer than the antennulæ; the ultimate and penultimate segments of the peduncle about equal in length; flagellum about as long as the flagellum of the antennulæ. Second epimeron (figure 5) rudely ovate, twice as high as broad; third somewhat rectangular, no wider than the second but considerably deeper; fourth (figure 6) very large, slightly deeper than the third and a third or a fourth longer than deep, being about as long as the first five segments of the thorax, the inferior margin regularly curved and the posterior convex in outline. First legs (figure 7) small and slender; merus triangular and broader distally than the carpus, which is not quite twice as long as broad and has the lateral margins parallel; propodus narrower but slightly longer than the carpus and narrowed distally; dactylus about half as long as the pro-

podus. Second legs (figure 5) stouter; merus short triangular; carpus much broader than long and only slightly produced beneath the propodus; propodus about as long as the breadth of the epimeron, nearly twice as long as broad; palmary margin (figure 8) convex in outline, slightly oblique, with an acute lobe and a spine at the posterior angle, within which the tip of the dactylus closes. Third and fourth legs slender and nearly naked. Basal segment in the fifth legs slender, four times as long as broad, not wider than the merus. Sixth and seventh legs slightly shorter than the fifth, the basal segments posteriorly dilated and squamiform in both pairs, but broader in the seventh than in the sixth. Posterior caudal stylets with the ramus slightly longer than the peduncle.

Length of largest specimen, from front of head to tip of telson, about 6^{mm}.

The mandibles are without palpi or molar tubercles, and in all other characters the species agrees with the genus *Stenothoë* as restricted by Boeck, but it seems to be very distinct from either of the European species.

Near Cultivator Shoal (haul *b*), 30 fathoms, soft, sandy bottom, August 29.

Syrrhoë crenulata Goës.

Crustacea amphipoda maris Spetsbergiam alluentis, Cefversight af Kongl. Vetenskaps-Akad. Förhandlingar, Stockholm, 1865, p. 527, pl. xl, fig. 25; Boeck, Crustacea amphipoda borealia et arctica (Vidensk.-Selskabs Forhandlingar, Christiania, 1870), p. 67, 1870.

We have also dredged this species, in 1872, in 12 fathoms in Johnson's Bay, near Eastport, Maine, and in 90 to 100 fathoms off Grand Menan, and have examined specimens dredged, in 1873, in 30 fathoms, in Gaspé Bay, Gulf of St. Lawrence. Our specimens have all been considerably larger than the one figured by Goës, but otherwise agree perfectly. It seems to be an exceedingly arctic form, being found in Europe from Spitzbergen to the western coast of Norway.

Tiron acanthurus Lilljeborg.

Boeck, op. cit., p. 69. *Syrrhoë bicuspis* Goës, loc. cit., p. 528, pl. xl, fig. 26. ? *Thessarops hastata* Norman, Annals and Magazine Nat. Hist., IV, ii, p. 412, pl. xxii, figs. 4-7, 1868.

This species has apparently not been noticed on our coast before. It has been found in Greenland, Finmark, and on the western coast of Norway, while Norman's *Thessarops* was from the English coast.

Ædiceros lynceus Sars.

Oversigt over nordsk-arct. Krebsdyr. Forhandl. i Vidensk.-Selsk. i Christiania, 1858, p. 143 (teste Boeck); Boeck, op. cit., p. 82. *Ædiceros propinquus* Goës, loc. cit.,

p. 526, 1865, pl. xxxix, fig. 19. *Monoculodes nubilius* Packard, Memoirs Boston Soc. Nat. Hist., i, p. 398, 1867.

We dredged this species in the Bay of Fundy in 1868 and 1872, the latter year in 60 to 80 fathoms; in Casco Bay, in 27 fathoms, in 1873, and Dr. Packard and Mr. Cooke obtained it at several localities, in the "Gulf of Maine," from 50 to 90 fathoms, on the expedition of the Bache in 1873. I have also examined specimens dredged in the Gulf of St. Lawrence by Mr. Whiteaves and on the coast of Labrador by Dr. Packard. It extends to Greenland, Iceland, Spitzbergen and Finmark.

Monoculodes borealis Boeck.

Op. cit., p. 88, 1870. *Ediceros affinis* Goës, loc. cit., p. 527, pl. xxxix, fig. 21', 1865 (non Bruzelius).

This species is recorded from Spitzbergen and northern Norway by Goës and Boeck, but seems not to have been noticed on this side of the Atlantic before.

Paramphithoe pulchella Bruzelius (Kroyer sp.)

We have dredged this species off Casco Bay and in the Bay of Fundy, on hard bottoms, in from 40 to 90 fathoms, and it was dredged on Cashe's Ledge and Stellwagen's Bank, in 1873, by Dr. Packard and Mr. Cooke. It extends north to the Gulf of St. Lawrence, and, according to Boeck, to Greenland, Iceland, Spitzbergen, and the western coast of Norway.

Paramphithoe cataphracta Smith.

Amphithonotus cataphractus Stimpson, Synopsis of the Marine Invertebrata of Grand Manan, p. 52, 1853 (description copied in Bate, Catalogue of Amphipodus Crustacea in the British Museum, p. 152, 1862.)

This species is apparently a true *Paramphithoë*, as restricted by Boeck, and closely allied to, if not identical with, *P. panopla* Bruzelius (*Amphithoë panopla* Kroyer). Boeck places *Pleustes tuberculatus* Bate as a synonym of Kroyer's species, and if he is correct in this, our species is undoubtedly distinct. The *cataphracta* appears to be an inhabitant of hard or coarse sandy and shelly bottoms from 5 to 50 fathoms. We have dredged it sparingly in Casco Bay and the Bay of Fundy, and Dr. Packard has dredged it on the coast of Labrador.

Vertumnus serratus? Goës (Fabricius sp.)

Acanthonotus serratus Stimpson, Synopsis of the Marine Invertebrata of Grand Manan, p. 52, 1853.

Our specimens all differ from the descriptions and figures given by Boeck and Kroyer in the armature of the posterior margin of the

third segment of the abdomen. In our specimens the upper process from this margin is armed with four or five teeth above and at the tip, while the lower process is armed with five or six teeth similarly situated, but with no teeth on the lower margin except just at the tip. In Kroyer's figure (Grönlands Ampfipoder, plate II, figure 8) the upper process is represented as terminating in a single tooth and the lower process as toothed along both sides; Boeck's description agrees with this except that he says there are two teeth at the tip of the upper process.

It is not uncommon on hard bottoms in from 5 to 50 fathoms in the Bay of Fundy. We have also dredged it in Casco Bay and have received it from the Gulf of St. Lawrence, where it was dredged by Mr. Whiteaves.

Acanthozone cuspidata Boeck.

This species is quite common on hard, and especially on spongy bottoms in 5 to 40 fathoms in the Bay of Fundy, although it is not mentioned by Stimpson in his work on Grand Menan. We have also dredged it in Casco Bay, and Mr. Whiteaves has obtained it in the Gulf of St. Lawrence. It ranges to Greenland, Spitzbergen and Finmark.

Byblis Gaimardi Boeck (Kroyer sp.)

We have frequently dredged this species in Casco Bay and the Bay of Fundy, on muddy bottoms in 10 to 60 fathoms. It extends north to the Gulf of St. Lawrence (Whiteaves), Labrador (Packard), and, according to Boeck, to Greenland, Iceland, Spitzbergen and Norway. The *Ampelisca Gaimardi* of Bate, and Bate and Westwood, is not this species but a true *Ampelisca*.

All the species of this sub-family are undoubtedly tube dwellers. Lilljeborg noticed the habit in *Haploöps*; it has been observed in species of *Ampelisca* by Professor Verrill and myself. In this species, the glands which secrete the cementing fluid are situated principally in the meral and basal segments of the third and fourth pairs of thoracic legs.

Xenoclea megachir Smith, sp. nov.

Plate IV, figures 1 to 4.

Male. Eyes large, black, very slightly elongated, and approaching closely the edges of the triangular prominence of the inferior angle of the front margin of the head. Peduncle of the antennulæ about as long as the head and the first two segments of the thorax, the second segment longest, the first and third about equal in length,

flagellum about as long as the peduncle and composed of twelve to sixteen segments. Antennæ a little longer than the antennulæ; ultimate and penultimate segments of the peduncle sub-equal in length; flagellum slightly shorter than the peduncle and composed of eleven to fifteen segments. First epimeron (figure 1) as broad as high; second (figure 2) broader than high; third (figure 3) and fourth not broader than high and successively deeper than the first and second; fifth (figure 4) slightly deeper than the fourth and its terminal portion as broad. In the first legs (figure 1), the carpus longer and broader than the propodus, which is somewhat oval and twice as long as broad; the dactylus slender, slightly curved and fully as long as the propodus. The inferior distal margin of the propodus is regularly curved to a short distance from the extremity, where there is a small but deep emargination, beyond which and round upon the short distal margin the edge is serrate with minutely crenulated teeth; the posterior margin is furnished with numerous slender setæ and with a single stout spine at the emargination near the distal end. The inner edge of the dactylus is armed with a series of acute teeth directed obliquely toward the tip. In the second pair of legs (figure 2) the propodus is very stout, about twice as long as the epimeron and scarcely one-half longer than broad; the palmary margin oblique and armed near the middle with two stout obtuse teeth; the dactylus stout and its inner edge sinuous. Third (figure 3) and fourth pair of legs alike; ischium and carpus short, each nearly or quite as broad as long; merus fully as long as the epimeron and half as broad as long; propodus slender, not more than half as broad as the carpus but twice as long; dactylus slender, about half as long as the propodus. Basal segment in the fifth legs (figure 4) squamiform, oval, nearly as broad as long and with a marked angular emargination at the inferior posterior angle; carpus only slightly longer than the breadth of the merus; dactylus slightly curved and acute. Second and third segments of the abdomen with the inferior portion of the posterior margin sinuous, and the inferior angle prominent, but scarcely less than right-angled. The outer rami in all the caudal stylets slightly shorter than the inner, and all the rami armed with short spines above and more slender spines at the tips. Telson stout, about as broad as long and scarcely more than half as long as the peduncle of the posterior caudal stylets, the posterior margin with a few setiform hairs each side.

In the female the hands in the second pair of limbs are proportionally much smaller and more abundantly provided with hairs, while the teeth, or lobes of the palmary margin, are further apart and

separated by a broad and deep, rounded sinus; the dactylus is not so stout, and has the inner margin evenly curved and serrated.

Length, from front of head to tip of telson, 5.5 to 7.5^{mm}.

I refer this species with some hesitation to Boeck's genus *Xenoclea*, which is known to me only from the very short diagnosis of the genus and of the single species *X. Batei*, given in his *Crustacea Amphipoda Borealia et Arctica*, p. 155. "Pedes 3tii et 4ti paris articulo 1mo latissimo" of the generic diagnosis would scarcely apply to our species, but in all the other generic characters it agrees perfectly, as it does also with the diagnosis of the sub-family Photinæ, except that the mandibles each bear six serrated spines instead of the usual number, four.

Near Cultivator Shoal (haul *b*), 30 fathoms, soft, sandy bottom, August 29; and on the northern side of George's Bank (haul *q*), north latitude 42°, west longitude 67° 42', 45 fathoms, coarse sandy bottom. Also, in 18 fathoms, off Watch Hill, Rhode Island.

When first examining the alcoholic specimens of this species, I noticed a peculiar opaque glandular structure filling a large portion of the third and fourth pairs of thoracic legs, which in most, if not all, the non-tube-building Amphipoda are wholly occupied by muscles. A further examination shows that the terminal segment (dactylus) in these legs is not acute and claw-like, but truncated at the tip and apparently tubular. In this species, a large cylindrical portion of the gland lies along each side of the long basal segment, and these two portions uniting at the distal end pass through the ischial and along the posterior side of the meral and carpal segments and doubtless connect with the tubular dactylus. (See Plate III, figure 3.) There can be no doubt that these are the glands which secrete the cement with which the tubes are built, and that these two pairs of legs are specialized for that purpose. A hasty examination revealed a similar structure of the corresponding legs in *Amphithoë maculata*, *Ptilocheirus pinguis*, *Cerapus rubricornis*, *Byblis Gaimardi*, and a species of *Ampelisca*. In all these except the last two a very large proportion of the gland is in the basal segment. In the *Amphithoë* this segment is thickened and the gland is in the middle. In the *Cerapus* it is very broad and almost entirely filled by the gland, with only very slender muscles through the middle, and the orifice in the dactylus is not at the very tip but sub-terminal on the posterior side. In the *Ptilocheirus* the gland forms three longitudinal masses in the basal segment and is also largely developed in the meral and carpal segments. The dactylus is long and slender and the orifice sub-terminal. In *Ampelisca* and *Byblis* (which, like *Haploöps*, are tube-building genera) the meral segments of the specialized legs are nearly

as large as the basal and contain a proportionally large part of the gland.

Scapellum Stroemi Sars.

Plate III, figure 9.

I am not aware that a description of this species has yet been published, although the name was used by Prof. Michael Sars in his list of animals living at great depths in the sea, published in 1869,* and the species has since been incidentally figured, without any detail, on the stems of *Mopsea borealis*, by Dr. G. O. Sars in his recent work on "Some Remarkable Forms of Animal Life from Great Depths off the Norwegian Coast" (Plate V, figure 2). Dr. G. O. Sars has, however, very kindly compared a drawing of one of our specimens, and he writes me that it agrees in every detail with the Norwegian form. It is very distinct from any of the species described in Darwin's great work, and also from the species recently described from the Challenger Expedition.

Since our specimens were obtained from 430 fathoms, Dr. Packard and Mr. Cooke have dredged in 50 to 70 fathoms near Cashe's Ledge, and in 142 fathoms, 20 miles east of Cape Race (both localities within the "Gulf of Maine"). All the specimens were attached to stems of hydroids. On the Norwegian coast the species has the same habit and has been found by Dr. G. O. Sars in from 80 to 300 fathoms.

ANNELIDA.

Lænilla (?) *mollis* G. O. Sars.

Bidrag til Kundskaben om Christianiafjordens Fauna, iii, p. 7, plate xiv, figs. 1-12, 1873.

Body large, rather stout, medially convex. Head short and broad, narrowed posteriorly, prominently rounded laterally, and produced into two very small conical points anteriorly. The anterior eyes are larger than the others, situated on the outer and upper surface of the lateral prominences, and look outward and upward; the posterior pair are nearer together, on the lateral slopes of the narrowed part of the head. The median tentacle is wanting in our specimen, but its basal segment is of moderate size and cylindro-conical; the antennæ are slender, and nearly three times the length of the head, banded with brown; the palpi are rather slender and regularly tapered, smooth, or nearly so, four or five times the length of the head. The dorsal and tentacular cirri and the scales are wanting in the single specimen obtained. The lateral appendages are large and

* Forhandlinger i Videnskabs-Selskabet i Christiania, 1868, p. 259, 1869.

prominent, with large fascicles of long, slender setæ in the lower rami, and much shorter and stouter ones in the upper rami. The appendages, including setæ, equal or exceed the breadth of the body. Breadth of body, exclusive of appendages, 7^{mm}; length of the latter, without setæ, 3·5^{mm}; with setæ, 10^{mm}; length of body to the 15th segment, 25^{mm}. The setæ of the upper ramus are very stout, and all of nearly the same form, the upper ones being merely smaller and stouter than the rest; they are nearly straight or slightly recurved, with rather conspicuous, moderately close transverse series of denticles, which extend nearly to the ends, leaving only stout, naked, straight tips. The setæ of the lower ramus are much longer and far more slender, with a long, slender shaft, and a slightly expanded terminal portion, which is conspicuously, but not closely, spinulated on both sides to the tips; many of these are nearly straight, but most are slightly curved; the upper ones are most slender, and mostly have the tips only very slightly bidentate, and the spinules exceed the diameter of the setæ and increase toward the end, the last ones projecting considerably beyond the tip; the middle ones are about twice as stout, having the terminal part more expanded; their spinulation is similar, but the tips are more distinctly, though slightly, bidentate, the denticles being partially obscured by the terminal spinules that project beyond them; the lower ones are more slender and like the upper ones in form and character.

Near St. George's Bank, 110 fathoms, mud. Coast of Norway, 40-200 fathoms (G. O. Sars).

Our specimen is imperfect, but the head and setæ are quite peculiar. The latter are remarkable for the length of the spinules, and for the minuteness of the denticles at the tips.—A. E. V.

Antinoe angusta Verrill, sp. nov.

Body narrow, rather slender, elongated, tapering gradually posteriorly. Head small, short, rounded, broader than long, the lateral lobes short, not prolonged into points anteriorly, but obtusely rounded; the lateral borders also well rounded. Eyes small, nearly equal; the posterior pair situated on the dorsal side of the vertex; the anterior pair farther apart on the outer and upper surface of the lateral prominences. Tentacle long and very slender, about three times the length of the head; antennæ small and short, scarcely one-third as long as the head; palpi moderately large, glabrous, considerably longer than the tentacle. Dorsal cirri slender, pretty regularly but not closely covered with slender papillæ. The lateral appendages, except anteriorly, bear large fascicles of long, fine capillary setæ, which gives a

villous appearance to the sides. The elytra, in our specimen, are wanting. The color, in alcohol, is light brown, crossed by lighter transverse lines. Length, 15^{mm}; breadth, without appendages, 2^{mm}; breadth, including setæ, 4^{mm}.

On the middle segments the setæ of the upper ramus are quite unequal in size and length; the upper ones are stout, with the ends more or less recurved; the middle ones are still larger and more than twice as long, slightly curved, and, like the former, conspicuously transversely serrulate almost to the extreme tips; the lower ones are shorter, less stout, and slightly curved. The setæ of the lower ramus are longer and extremely slender; the upper ones are mostly but slightly expanded in the middle, with very long, flexible capillary tips, finely tapered to the end, and very minutely serrulate or nearly smooth; the median ones are stouter, more expanded in the middle, with long, acuminate, slender, sharp tips, and with conspicuous, rather distant spinules on one or both sides, which become very fine and more crowded distally; the lower ones are much shorter, and have shorter but still very slender tips, and fewer and more distant spinules. The ventral cirri are slender, tapered, with few, distantly scattered, small papillæ.—A. E. V.

Near Saint George's Bank, 150 fathoms, mud (locality *s*).

Antinoe Sarsi Kinberg.

Malmgren, Nordiska Hafs-Annulater, Öfversigt Kongl. Vetenskaps-Akad. Förhandlingar, Stockholm, 1865, p. 75, pl. 9, fig. 6; *Annulata Polychæta*, p. 13, 1867.

Our specimen of this species agrees very well with Malmgren's figures and description. It is much larger and stouter than the preceding, and the head is longer and quite different in form, the lateral lobes extending forward into acute conical points.

The setæ are similar to those of the former, but the median and inferior setæ of the lower ramus are relatively somewhat stouter and have the tips less attenuated and elongated, while the spinules are larger and more conspicuous, especially on the upper setæ of the lower ramus.

Near Saint George's Bank, 85 fathoms, mud. Gulf of Saint Lawrence (Whiteaves, t. McIntosh).—A. E. V.

Eucranta villosa Malmgren.

Eucranta villosa Malmgren, Nordiska Hafs-Annulater, Öfversigt af Kongl. Vetenskaps-Akad. Förhandlingar, Stockholm, 1865, p. 80, pl. 10, fig. 9; *Annulata Polychæta*, p. 14, 1867.

? *Eupolynoë occidentalis* McIntosh, *Annals and Magazine Nat. Hist.*, IV, vol. xiii, p. 264, pl. 9, figs. 8-13, 1874.

This large species is easily distinguished, even when destitute of

its scales, by the short, stout, strongly curved setæ of the upper ramus, and much longer, slender, fasciated setæ of the lower ramus, among which the upper ones have a strongly spinulose, slender, acuminate, terminal portion, with a nearly straight, split, or forceps-like, slender tip, while the middle and lower ones have a short, cuspidate terminal portion, with few large spinules, and naked acute tips.

I am unable to find anything in the figures and description of the species recently described by McIntosh to indicate that it is distinct from the present species, with which, however, he has not compared it.

Near Saint George's Bank, 150 fathoms, mud (locality *s*). Gulf of Saint Lawrence, 110 fathoms (Whiteaves, t. McIntosh).—A. E. V.

Nephtys circinata Verrill, sp. nov.

Body slender, elongated, rather depressed, tapering gradually posteriorly. Head sub-pentagonal, rather broader than long; a pair of short, tapering antennæ at the anterior angles, about one-fourth as long as the width of the anterior border of the head; another pair of longer, slender, tapering antennæ at the lateral angles; tentacular cirri long and tapering. Proboscis smooth toward the base; its distal portion with rows of slender acute papillæ, which increase rapidly in length toward the end, where they become very prominent.

The lateral appendages, including the setæ, are as long as the breadth of the body; the setæ are very numerous, long and slender. The caudal cirrus is long and slender, tapering to a slender tip. Length of body, 50^{mm}; diameter, 2.5^{mm}; diameter, including appendages, 5^{mm}.

The lateral appendages of the middle region are moderately long, the rami separated by a space scarcely equal to half their height. Superior ramus, with a short, broad ovate, obtuse, or slightly acuminate upper lamella, directed outward, and considerably exceeding the setigerous lobe, and a much smaller ovate median lamella; branchial cirrus long, rather slender, tapered, curved downward and inward (circinate), forming rather more than a complete whorl; the appendage at its base, on the anterior segments, is short and broad, subtruncate distally, and with a small papilliform process projecting downward from its lower angle, nearly in contact with the branchial cirrus; on the median segments it is broad and long-ovate, unequally acuminate, leaf-like. The lower ramus has a very long and wide ligulate lamella, directed obliquely upward and outward, usually more than twice as long as the setigerous lobe, and about equal to it in width; its lower edge at about the middle is sometimes incurved, and its tip is acuminate and blunt-pointed; the ventral cirrus is slender and tapered.

The capillary setæ form large fascicles and are very long and slender, nearly smooth, and with very attenuated tips; their length is about three times that of the appendages themselves; the transversely marked setæ are scarcely one-fourth as long, and about the same in diameter, with very slender tips.—A. E. V.

East of Saint George's Bank, 430 fathoms (locality *g*); north of Saint George's Bank, 85 fathoms, mud (locality *p*).

Nephtys ingens Stimpson.

Synopsis of the Marine Invertebrata of Grand Manan, p. 33, 1853; Verrill, Report on the Invertebrate Animals of Vineyard Sound and Adjacent Waters, in Report of U. S. Commissioner of Fish and Fisheries, part I, 1873, p. 583 (separate copies, p. 289), plate xii, figs. 59, 60, 1874.

? *Nephtys incisa* Malmgren, Öfversigt af Kongl. Vet.-Akad. Förhandlingar, 1865, p. 105, plate xii, fig. 21.

This is the most common and abundant species on muddy bottoms in the deep water along the whole New England coast. It occurs at all depths from 2 to 430 fathoms.

It is easily distinguished by the stout quadrangular body, deeply incised posteriorly; by the blackish setæ, and by the remarkably elongated and widely separated rami of the posterior appendages. There is a long, odd, median papilla on the dorsal side of the proboscis, and a smaller one beneath; the papillæ in the longitudinal rows are rather small.—A. E. V.

Phyllodoce catenula Verrill.

Report on the Invertebrate Animals of Vineyard Sound, in Report of U. S. Commissioner of Fish and Fisheries, part I, 1873, p. 587, 1874; Exploration of Casco Bay by the U. S. Fish Commission, Proceedings American Association for the Advancement of Science, 1873, p. 380, pl. 3, fig. 1, 1874.

Plate IV, figure 3.

George's Bank, 50 fathoms (locality *d*). It also occurs at Watch Hill, Rhode Island, in 4 to 6 fathoms, among rocks and algæ, and in tide-pools; at Wood's Hole, at surface, evening, July 3; in Casco Bay, 8 to 30 fathoms; and is very common in the Bay of Fundy, from low-water to 50 fathoms.

This species is closely allied to *P. pulchella* Malmgren, from northern Europe, but differs somewhat in the form of the head, which is shorter and rounder in the latter; the branchiæ also differ in form.

Eusyllis phosphorea Verrill, sp. nov.

Plate VII, figure 3.

Body slender, elongated, tapering gradually posteriorly. Head, in alcoholic specimens, broader than long, well-rounded in front, the posterior margin incurved; but in living specimens the head is longer

than broad and slightly narrowed posteriorly. Eyes small, but conspicuous, wide apart, the anterior considerably farther apart than the posterior ones. Palpi large in preserved specimens, broad ovate, and well rounded anteriorly, in contact at their bases; but in living specimens more elongated and oblong, exceeding the length of the head. Antennæ (or tentacles) long and slender, distinctly and rather regularly annulated, but not moniliform. Tentacular cirri, in preserved specimens, similar to the antennæ; the upper ones are of about the same length, but the lower are little more than half as long. Dorsal cirrus of the second segment is as long as, or even longer than, the antennæ. The dorsal cirri on the 3d, 4th, and 5th segments are shorter, about equal, longer than the lower tentacular cirrus, and about half as long as the dorsal cirri of the succeeding segments, which are alternately longer and shorter, the longer ones about half as long as the breadth of the body. While living, the alternate dorsal cirri are usually held extended and curled up over the back. The two anal cirri are long and slender; in one preserved specimen they are more than twice the breadth of the body, while in the same specimen the dorsal cirri on the second and third segments preceding the anal one are considerably longer than those on the segments farther forward.

The setæ are all compound, rather long, mostly considerably bent, with a short, acute-triangular terminal piece, which is very distinctly bidentate at the tip.

Color of body, when living, deep salmon, or light yellowish orange, with dark brown intestinal line, darker posteriorly; eyes dark brown.

Length, when living, about 25^{mm}; breadth, 1.5^{mm}.

Saint George's Bank, 45 fathoms, among hydroids; Bay of Fundy, off Grand Menan, 52 fathoms, among hydroids.

This species, when living, was most brilliantly phosphorescent, with a bright green light, so intense as to be distinctly visible in daylight, or close to a large kerosene lamp.—A. E. V.

Ninoe nigripes Verrill.

Report on the Invertebrate Animals of Vineyard Sound, in Report of U. S. Commissioner of Fish and Fisheries, part I, 1873, p. 595, 1874; Proceedings American Association for Advancement of Science, 1873, p. 382, pl. 3, fig. 5, 1874.

Plate V, figure 3.

Locality *o*, 110 fathoms. Also Fisher's Island Sound, Vineyard Sound, and Buzzard's Bay, and waters outside, in 8 to 29 fathoms, mud; Casco Bay, 10 to 68 fathoms; off the coast of Maine, at various depths to 107 fathoms.

Leodice vivida Verrill.

Eunice vivida Stimpson, Marine Invertebrata of Grand Manan, p. 35, 1853.

Leodice vivida Verrill, American Journal of Science, III, vol. v, p. 9, January, 1873.

Plate V, figure 5.

Nothria conchylega Malmgren.

Onuphis conchylega Sars, Beskrivelsir og Iagttagelser, p. 61, pl. 10, fig. 28 (*teste* Malmgren), 1835.

Onuphis Eschrichti (Ersted, Grönlands Annulata Dorsibranchiata, p. 20, pl. 3, figs. 33-41, 45, 1843.

Northia conchylega Johnston, Catalogue of British Worms, p. 138, 1865.

Nothria conchylega Malmgren, Annulata Polychæta, p. 66, 1867.

Plate VII, figure 3.

This species is abundant in the deeper waters, especially upon hard bottoms, on the whole northern coast of New England, and in the Gulf of St. Lawrence. Malmgren records it, in 30 to 250 fathoms, from Greenland, Spitzbergen, Finnmark, and the coast of Norway.

The name "Nothria" was substituted for *Northia* (Johnston) by Malmgren for reasons that are scarcely sufficient. The latter name was, however, previously in use for a genus of shells (Gray, 1847), and must be rejected on that account.

Nothria opalina Verrill.

American Journal of Science, III, vol. v, p. 102, 1873.

Plate VII, figure 4.

Body long and slender, narrowed anteriorly, much depressed and of nearly uniform width throughout most of its length; the five anterior segments much longer than the others. Palpi inferior, rather large, hemispherical; antennæ small, ovate, close together, on the front of head. Three central tentacles very long and slender, tapering, acute, the basal portion regularly annulated and thickened for a considerable distance, beyond which the surface is smooth, with an occasional distant annulation; the central odd one is somewhat shorter and more slender than the two adjacent ones, which reach to or beyond the 10th segment; outer pair much shorter, being less than half the length of the central ones. Tentacular cirri small and very slender. Lateral appendages or "feet" of the first six setigerous segments similar in structure but more prominent than the following ones, from which they also differ in having the ventral cirrus well developed, long and tapering, but shorter and thicker on the first segment than on the five following. Those of the first pair have a stout stalk, which terminates in a small, bluntly rounded setigerous lobe, with a long, slender, subterminal cirrus-like lobe above,

longer than the stalk; dorsal cirrus arising from near the base, longer and more slender than the terminal cirrus; branchial filament simple, long and very slender, about equalling the dorsal cirrus and united to it above its base; ventral cirrus ovate, tapering, blunt, arising from near the base. The second pair of feet are similar to those of the first, except that in the largest specimens there are two branchial filaments, and the ventral cirrus is longer and more slender. The 3d, 4th, 5th, and 6th pairs have essentially the same structure, but the ventral cirrus becomes gradually longer to the 6th, where it is longer than the stalk and nearly equal to the terminal cirrus. The succeeding feet are much shorter; the ventral cirrus is a mere conical papilla, which soon disappears; the terminal cirriform lobe becomes smaller and disappears after the 10th pair; the branchial filament becomes larger and longer to the middle region, where it exceeds in length half the diameter of the body, while the dorsal cirrus at the same time becomes smaller and shorter, until it is less than one-fourth the length of the branchia.

The setæ of the anterior feet consist of slender, acutely pointed, curved ones, mixed with much stouter, blunt pointed compound ones; farther back there are two fascicles of more slender acute setæ, and in the lower bundles a few long, stout, bidentate hooks, with a thin, rounded, terminal expansion.

Color, in alcohol, pale yellowish white, but everywhere very brilliantly iridescent, with opaline lustre and colors.

Length, 75 to 125^{mm}; diameter, 2.5 to 4^{mm}.

Common in 110 and 150 fathoms, hauls *s* and *o*. It was also dredged in 1873, off Casco Bay, in 30 to 94 fathoms, and on Jeffrey's Bank, in 79 to 105 fathoms. It was also abundant, on muddy bottoms in deep water, at all the localities in the Gulf of Maine examined by Dr. Packard and Mr. Cooke in 1873.

Goniada maculata Ersted.

Ann. Dan. consp., p. 33, figs. 16, 23, 91, 95, 97, 98 (t. Malmgren). *Glyceria viridescens* Stimpson, Marine Invertebrata of Grand Manan, p. 33, 1853.

North of Saint George's Bank, 110 and 150 fathoms, mud (localities *o* and *s*); Saint George's Bank, 20 fathoms (locality *j*). Off Casco Bay, 30 to 90 fathoms, mud. Bay of Fundy, 20 to 70 fathoms. Common in the Gulf of Maine, 60 to 100 fathoms. Northern coasts of Europe, from Finmark to Scotland, 10 to 130 fathoms (Malmgren).—A. E. V.

Rhynchobolus capitatus Verrill.

Glycera capitata Ersted, Grönl. Ann. Dorsibranchiata, p. 44, plate VII, figs. 87, 88, 90-94, 96, 99; Malmgren, Annulata Polychæta, p. 70, 1867 (*non* Claparède).

This species is furnished with four well-developed jaws, and therefore belongs to the genus *Rhynchobolus*, as constituted by Claparède. The species without jaws, which he refers to *Glycera*, must be distinct.

Saint George's Bank, 60 fathoms (locality *e*); 20 fathoms (locality *j*); 110 fathoms (locality *o*); east of Saint George's, 430 fathoms (locality *g*). Greenland, Iceland, Spitzbergen, and northern coasts of Europe to Great Britain.—A. E. V.

Samythella Verrill.

Body elongated, composed of about 50 segments, 15 of which bear fascicles of setæ; and posteriorly about 35 bear uncini only, but have a small conical papilla above the uncigerous lobe, as in *Melinna*; the uncini commence on the 4th setigerous ring. Branchiæ six, placed side by side in a continuous transverse row. Cephalic lobe oblique, somewhat shield-shape, with a narrowed prominent front. Buccal lobe shorter. Tentacles numerous, smooth and slender.

This genus is closely allied to *Samytha* of Malmgren, in the structure of the head and number of branchiæ, but differs in having a much larger number of segments (in this respect approaching *Melinna*), and in having only 15 setigerous segments, instead of 17.

Samythella elongata Verrill.

American Journal of Science, III, vol. v, p. 99, 1873.

Body slender, composed of 54 segments in the specimens examined, tapering regularly to the posterior end. Cephalic lobe about as broad as long, broadly rounded posteriorly, with the postero-lateral corners prominent and well rounded, the sides slightly incurved and rapidly narrowing to the front, which is about half the width of the back, and subtruncate, projecting forward; the middle region is a raised and convex oblong area as wide as the front edge, into which it runs. Buccal lobe a little shorter. Tentacles numerous, slender, tapering. Branchiæ subequal, slender, tapering, about twice the length of the cephalic lobe. Setæ numerous and long in all the fascicles except the first three, the longest nearly one-third the diameter of the body. The posterior end of the body is surrounded by about eight small papillæ, of which the two upper ones are largest.

Length of largest specimen, in alcohol, 46^{mm}; diameter, 2.5 to 3^{mm}.

The tubes consist of a thin and tough lining, to which a close layer of sand, in grains of moderate and nearly uniform size, is firmly cemented.

Grymæa spiralis Verrill.

Am. Journal of Science, III, vol. vii, p. 407, fig. 2, and plate V, fig. 4, April, 1874.

Plate IV, figure 1.

Body long and slender, spirally coiled, composed of over 150 segments, of which about 120 bear fascicles of slender setæ. Branchiæ long filiform, two or three times the diameter of body, arising in three clusters on each side, easily detached and often partially absent. Setæ on the first six or seven segments a little longer than the following ones. General color dark red. Tube composed of firmly cemented mud and sand, coiled in a double spiral, the two halves revolving in opposite directions.

Also dredged, in 1872, off Grand Menan Island, Bay of Fundy, in 60 fathoms; and in 1873, off Casco Bay, in 90 fathoms, mud; and in 80 fathoms on Jeffrey's Bank.

Fig. 1.*

? *Potamilla neglecta* Malmgren.

Öfversigt af Kongl. Vet.-Akad. Förhandlingar, 1865, p. 401, plate 27, fig. 84.

Subella neglecta Sars, Reise i Lofot. og Finn., p. 83 (t. Malmgren).

This species was very abundant at localities *d*, *h*, *i*, *q*, and also occurred in 110 fathoms (locality *o*).

The tubes are long and tough, covered externally with sand. One specimen from Le Have Bank, 45 fathoms (locality *h*), had a large number of young ones within the tube, adhering to its inner surface.—A. E. V.

Spirorbis valida Verrill, sp. nov.

Tubes much larger than usual in the genus, round, strong, thick, opaque, white, transversely wrinkled, rather rapidly enlarging, sinistral, or coiling in the same direction with the hands of a watch; in some specimens, found attached to flat shells, the tubes form low, rapidly enlarging spirals of several turns, the last whorl enveloping and concealing the others externally, except near its termination, where it rises obliquely upon the preceding one, but leaving a broad, shallow umbilicus in which the previous whorls are visible; in other specimens, attached to convex univalve shells (*Tarritella erosa*, etc.), the whorls rest upon the upper side of each of the preceding ones, forming an elevated and often somewhat irregular spiral, increasing in size upward, with a small umbilicus, and usually with the last part of the upper whorl slightly free from the preceding one and ascending

* Tube of *Grymæa spiralis*, natural size.

obliquely. Diameter of the larger tubes, at end, 1.75^{mm} to 2^{mm}; height of the more elevated spirals, 3^{mm} to 5^{mm}.

There are 15 large branchiæ in the adult specimens: 8 on the left side, 7 on the right, with the operculum; the pinnae are long, slender, extending to near the ends of the branchiæ, which have slender and short, naked tips. Operculum large, white, calcareous, irregularly obconic, obliquely truncated, with the outer surface concave, the dorsal side gibbous, the margin slightly sinuous but entire, except for a small notch, or emargination, in the dorsal edge; the dorsal portion is translucent, while the ventral portion is opaque and contains small, round, ova-like bodies; the peduncle is rather short and stout, gradually expanding into the base of the operculum, but swollen in the middle, on the dorsal side. Collar, in the specimens examined, considerably mutilated, apparently with a sinuous but not revolute anterior margin, and with a long posterior dorsal point. The region covered by the collar bears, at least on the left side, three large fascicles of slender, acute, yellowish setæ, both above and below; the anterior fascicles are directed forward, and the upper anterior one is larger than the other fascicles.

Le Have Bank, 45 and 60 fathoms (localities *h* and *i*).

The size of this species is exceptionally large, and the branchiæ are unusually numerous for the genus *Spirorbis*, to which I refer it with some hesitation. When living specimens can be studied it may prove to be a new genus. It has, like *Vermilia*, a calcareous operculum, but in form and structure this organ resembles that of some species of *Spirorbis*.—A. E. V.

? *Spirorbis nautiloides* Lamarck.

Anim. sans Vert., ed. I, vol. v, p. 359, 1818. ? *Spirorbis communis* Quatrefages, Histoire naturelle des Annelés, vol. ii, p. 489.

Plate IV, figure 4.

The species figured agrees pretty well with that described by Quatrefages, but may not be the same as that of Lamarck, which is regarded by several writers as synonymous with it, and by others with *S. borealis*, the species so abundant on *Fucus* at low-water mark, on our shores.

The present species is seldom, if ever, found at low-water mark, and occurs chiefly on stones and shells in deep water. The tubes are opaque, white, cylindrical, rather closely coiled, the terminal portion not erect, and the surface is more or less conspicuously marked with lines of growth.

Abundant on the hard bottoms at Saint George's Bank; Casco

Bay; Cashe's Ledge; and in the Bay of Fundy, 10 to 106 fathoms.—A. E. V.

Protula media Stimpson.

Marine Invertebrata of Grand Manan, p. 30, 1853.

Plate VI.

This species usually forms much contorted and irregularly bent tubes, which are cylindrical and nearly smooth, but with irregular lines of growth.

North of Saint George's Bank, 110 fathoms (locality *o*). Often brought up by fishermen on Saint George's Bank, attached to shells and stones. Abundant on Cashe's Ledge, 50 to 70 fathoms; off Grand Manan, 30 to 50 fathoms; off Casco Bay.—A. E. V.

? *Protula borealis* Sars.

Vidensk. Selsk. Forhandlinger, 1871, p. 417 (separate copies, p. 14).

Numerous empty tubes from the muddy bottoms in 110 and 150 fathoms (localities *o* and *s*) differ considerably in form from those of the *P. media*, ordinarily met with, and may be this species, if distinct. But they may, very likely, prove to be only a variation of the former, due to the muddy character of the bottom. The tubes are much less bent and contorted, often but slightly curved, or nearly straight, nearly smooth, but with occasional ridges or folds, indicating periods of growth.—A. E. V.

GEPHYREA.

Phascolosoma cæmentarium Verrill (Quatrefages sp.).

American Journal of Science, III, vol. v, p. 99, 1873; and Report upon the Invertebrate Animals of Vineyard Sound, in Report of U. S. Commissioner of Fish and Fisheries, part I, 1873, p. 627, pl. xviii, fig. 92, 1874.

Very common on the coast of New England, from Long Island Sound northward, in 5 to 430 fathoms, in dead univalve shells.

Phascolosoma tubicola Verrill.

American Journal of Science, III, vol. v, p. 99, 1873; Proceedings American Association for Advancement of Science, 1873, p. 388, 1874.

Body versatile in form; in contraction short, cylindrical, oval or fusiform, 12 to 25^{mm} long, 2.5 to 4^{mm} in diameter; in full extension the body is more or less fusiform, gradually tapering anteriorly into the long, slender, nearly cylindrical retractile portion, which is longer than the rest of the body, and bears, near the end, a circle of about ten to sixteen simple, slender tentacles, beyond which the terminal

portion is often extended into a short proboscis, with the mouth at the end; below the tentacles there is sometimes a dilation, but this is without special spines or granules, and like the rest of the retractile portion in texture. The posterior end of the body is bluntly rounded, and the skin is transversely wrinkled and rough, and covered with small, round, somewhat raised verrucæ or suckers, to which dirt adheres, and at the end nearly always bears from 3 to 8 small, but prominent, peculiar bodies, having a slender pedicle and a clavate or globular head; their nature is doubtful. (They may be sense-organs, but should be examined on living specimens.) At about the posterior third of the proper body is an irregular zone of numerous, dark brown, hard chitinous hooks, arranged in several rows, broad triangular in form, with acute points directed forward; among the hooks are also a few suckers; the middle region is covered with small, round, slightly raised suckers, which become much more prominent and crowded at the anterior end toward the base of the retractile portion, and have here the form of small, subconical, elevated warts, to which dirt usually adheres firmly; the retractile portion is covered throughout with minute conical verrucæ or papillæ, most prominent toward the base.

In many respects *P. cæmentarium* agrees very closely with this, but it has the posterior end much smoother, and with less conspicuous suckers; the hooks are not so numerous, less acute, and lighter colored; the anterior part of the body has smaller and less prominent suckers or verrucæ; the skin is lighter colored, thinner, and more translucent, and there is a zone bearing several rows of minute, slender, acute, chitinous spinules, a little below the tentacles.

Hauls *p*, *o*, and *s*, 85 to 110 fathoms. It has also been dredged, in 60 to 94 fathoms, off Casco Bay.

? *Phascolosoma boreale* Keferstein.

Beiträge zur Anat. und syst. Kenntniss der Sipunculiden, p. 206.

This species is rather short and thick, obtuse posteriorly, nearly smooth to the naked eye, and destitute of both hooks and distinct suckers, but the skin is minutely wrinkled transversely, and covered with almost microscopic slender papillæ, and is minutely specked with dirty yellowish brown; the retractile portion is more distinctly granulated anteriorly. The tentacles are rather numerous, small, and simple.

Dredged also off Casco Bay, 64 fathoms; Cashe's Ledge, 50 to 72 fathoms; and in the Gulf of St. Lawrence (Whiteaves).

MOLLUSCA.

Pleurotomella Packardii Verrill.

American Journal of Science, III, vol. v, 1873, p. 15 (December, 1872).

Shell thin, fragile, translucent, pale flesh-colored, moderately stout, with an acute, somewhat turreted spire. Whorls nine; the apical whorls, for about two and one-half turns, are nearly smooth, regular, convex, chestnut-colored; below this the whorls are shouldered, strongly convex in the middle, but with a smooth concave band below the suture, corresponding to the posterior notch in the outer lip; the whorls are crossed below the sub-sutural band by about 16 strong, prominent, rounded, somewhat oblique ribs, most prominent on the middle of the whorl, but not angulated; on the last whorl these ribs become very oblique below the middle, and follow the curve of the edge of the lip, nearly fading out anteriorly; the surface between the ribs is marked by faint lines of growth and by fine, unequal, slightly raised revolving lines, which pass over the ribs without interruption. They become more evident on the lower part of the last whorl, and are very faint on the sub-sutural band, which is more decidedly marked by receding, strongly curved lines of growth. The aperture is rather broad above, elongated below, sub-oval, outer lip very thin, sharp, prominent above, separated from the preceding whorl by a wide and very deep sinus, extending back for about one-fifth of the circumference of the whorl; the anterior border of the lip is incurved near the end, and obliquely truncate, forming a short, straight canal. Columella simple, nearly straight, its inner edge toward the end sharp, and obliquely excurved. No operculum. Length, 21.2^{mm}; breadth, 11.2^{mm}; length of aperture, 12.0^{mm}; breadth of same, 5.0^{mm}. The absence of eyes and operculum, great size of the posterior sinus, and character of the apex, indicate that this shell represents a new genus.

One living specimen from (*o*) 110 fathoms.

Ringicula nitida Verrill.

American Journal of Science, III, vol. v, 1873, p. 16 (December, 1872).

Plate I, figure 2.

Shell small, white, smooth, broad oval, with five whorls, spire rapidly and regularly tapered, sub-acute, shorter than the aperture. Whorls very convex, regularly rounded, the sutures well impressed; a well marked, impressed, revolving line just below the suture; the surface otherwise nearly smooth, but with more or less distinct, distant, microscopic revolving lines, most distinct anteriorly. Aperture somewhat crescent-shaped. Outer lip evenly rounded, forming

the segment of a circle, the border regularly thickened, receding a little posteriorly, near the suture. Callus on the body whorl narrow, nearly even, but a little swollen in the middle and slightly raised. Columella stout, recurved at the end, with two strong, very prominent, equal, spiral folds—the anterior one projecting beyond the canal, with the end rounded. Length, 4.2^{mm}; breadth, 3.1^{mm}; length of aperture, 2.5^{mm}; breadth of aperture, .11^{mm}.

From 110 and 150 fathoms (localities *s* and *o*).

Torellia vestita Jeffreys.

This shell in form and size somewhat resembles large specimens of *Margarita helicina*, but it has a ciliated epidermis resembling that of *Velutina levigata*. The spire is small and low; whorls four, the last large, well rounded, forming the bulk of the shell. Suture deep. Umbilicus small and deep, somewhat concealed by the reflected outer edge of the columella, which recedes in front and joins the outer lip at an obtuse angle, forming a broad, shallow, anterior emargination; inner border of the columella a little excavated near the body whorl, slightly swollen in the middle. Outer lip sharp, regularly rounded. Epidermis thick, greenish, with conspicuous lines of growth, finely reticulated by raised revolving lines, along which arise numerous slender, but short, hair-like processes. Shell beneath the epidermis white, nearly smooth. Length, 7.5^{mm}; breadth, 10^{mm}; length of aperture, 6^{mm}; breadth, 4.5^{mm}.

Only one specimen, dead and inhabited by a *Phascolosoma*, was found in 1872. Since this, however, during the explorations of 1873, it was dredged by Dr. Packard and Mr. Cooke, in 52 to 90 fathoms, on Cashe's Ledge, off the coast of Maine.

Stylifer Stimpsonii Verrill.

American Journal of Science, III, vol. iii, p. 283, 1872.

Plate I, figure 1.

Shell white, short, swollen, broad oval; spire short, rapidly enlarging. Whorls four or five, the last one forming a large part of the shell; convex, rounded, with the suture impressed; surface smooth, or with faint striæ of growth. Color, when living, pale orange yellow. Length, about 4^{mm}; breadth, 3^{mm}.

Parasitic on *Strongylocentrotus Dröbachiensis*. In 32 fathoms off the coast of New Jersey (Capt. Gedney); 60 and 65 fathoms (*e* and *f*), George's Banks; 8 fathoms off Fisher's Island, mouth of Long Island Sound.

Astarte undata Gould.

Verrill, American Journal of Science, III, vol. iii, pp. 213, 287, 1872; and Report on the Invertebrate Animals of Vineyard Sound, in Report of U. S. Commissioner of Fish and Fisheries, part I, 1873, p. 384, pl. 29, fig. 203, 1874.

Plate I, figures 6 to 9.

The figures given in Gould's works are scarcely characteristic of this, the most abundant species of the northern coast of New England, and we here publish several figures, prepared by Professor Verrill, which more fully illustrate the different forms of the species. The name *undata* was proposed by Gould for a form of his *Astarte sulcata*.

Astarte lens Stimpson.

Astarte crebricostata Gould, Invertebrata of Massachusetts, 2d edition, edited by Binney, p. 126, fig. 440, 1870 (not of Forbes, *teste* Verrill).

Astarte lens Stimpson, MS., Gould, op. cit., p. 127; Verrill, American Journal of Science, III, vol. iii, pp. 213, 287, 1872.

Plate I, figures 4 and 5.

This species seems to be more exclusively a deep-water form than the last, although the specimens dredged by us at the localities (*g*, *o*, and *s*) mentioned are all much smaller than the common form of the species in the Bay of Fundy, and may well be regarded as a dwarf variety.

Pecten pustulosus Verrill.

American Journal of Science, III, vol. v, 1873, p. 14 (December, 1872).

Upper valve more convex than the lower, a little swollen toward the umbo; length and breadth nearly equal, the margin diverging nearly at right angles from the beak to the middle of the anterior and posterior borders, on each of which there is an obtuse angle, from which the outline of the ventral margin forms a regular curve, nearly semicircular, but a little produced ventrally; the surface with about 14 radiating rows of relatively large, prominent, round, hollow vesicles, those in the middle rows nearly hemispherical, while part of those of the lateral ones are subconical and smaller; seven or eight of the rows are first developed, at a short distance from the apex of the shell, the other ones afterward coming in between the primary ones; the rows are distant in the middle and more crowded together toward the borders; between the rows of vesicles the surface is marked by distant, fine, impressed grooves, which pass between and separate the vesicles; on the umbos, above the origin of the vesicles, the border of the groove rises into a thin, slightly elevated lamella. Lower valve with fine, close, slightly raised, concentric lamellæ, be-

coming faint toward the beak. Auricles unequal, that of the upper valve small, and a little projecting posteriorly, much longer and more prominent, with a deep, curved emargination anteriorly, its surface with concentric lamellæ and radiating rows of small, conical vesicles; that of the lower valve with a deep, angular byssal notch anteriorly, its surface with concentric lamellæ and faint radiating ridges. Color yellowish white. Length, 7.5^{mm}; height, 8.0^{mm}; thickness, 2.5^{mm}.

East of St. George's Banks (*g*), in 430 fathoms, dead but fresh valves; and north of the Banks, locality (*s*), 150 fathoms, living.

Pera crystallina Verrill.

Clavelina crystallina Möller, Naturhistorisk Tidsskrift, vol. iv, p. 95, 1842.

Pera pellucida Stimpson, Proceedings Boston Soc. Nat. Hist., vol. iv, p. 232, 1852.

Pera crystallina Verrill, American Journal of Science, III, vol. iii, p. 213, pl. 8, fig. 9, 1872.

Plate VIII, figure 1.

This species was described by Stimpson from specimens, adhering to stems *Sertularella polyzonias*, variety *gigantea*, taken in 30 fathoms on St. George's Banks. Professor Verrill records it from Murray Bay, Gulf of St. Lawrence.

Glandula arenicola Verrill.

American Journal of Science, III, vol. iii, pp. 211, 288, 1872; Report on the Invertebrate Animals of Vineyard Sound, in Report of U. S. Commissioner of Fish and Fisheries, 1873, p. 701, 1874.

This species, which was dredged by us in immense numbers in 28 fathoms (haul *c*), has also been dredged, by Dr. Dawson, at Murray Bay, Gulf of St. Lawrence, by Mr. T. M. Prudden, in Buzzard's Bay, and off New London, Conn., by A. E. Verrill.

Thyone scabra Verrill.

American Journal of Science, III, vol. v, p. 100, 1873.

Thyone fusus? Verrill, American Journal of Science, III, vol. v, p. 14, 1873 (*non* Koren).

Body fusiform, gradually tapered behind, with a long, slender, posterior portion, covered throughout with very numerous, rather rigid, slender, scabrous papillæ; skin rather rigid, scabrous with small, rough points, which project from the plates. Tentacles ten; eight large ones much elongated and arborescently divided from near the base; the two small ones are very short, nearly sessile, subdivided from the base. The calcareous plates of the skin are very flat, somewhat imbricated, irregularly oval, triangular, or subpolygonal, with an undulated or crenulated margin, pierced by about 20 to 24 unequal round openings, two or three central ones larger than the rest, the

interspaces mostly as wide as the pores; from the center of the upper side arises an open, slender, flat, acute spinous process, composed of two anastomosing pieces. The plates of the papillæ or suckers are narrow, elongated, bent into a bow-shape, the middle expanded and usually pierced by about four pores, two of which are larger; the ends are also usually dilated and pierced with small pores; from the middle arises a flat, spinous process, similar to that of the skin-plates, but smaller.

Length, in alcohol, about 50^{mm}; greatest diameter, 6 to 9^{mm}; length of longest tentacles, 7.5^{mm}. Color of preserved specimens, yellowish brown.

Localities *o* and *s*, 110 and 150 fathoms. Also dredged, in 1873, off Caseo Bay.

This species resembles *T. raphanus* Duben and Koren (Troschel sp.) in form, but the latter has long-stalked tentacles, branching only near the ends, and the plates of the skin are different in form, and in the perforations, and lack the spinous processes which give the species its rough, scabrous surface.

? *Charybdea periphylla* Péron and Lesueur.

Verrill, Report upon the Invertebrate Animals of Vineyard Sound, p. 724, 1874.

This species, originally described and figured by Péron and Lesueur from mutilated specimens taken under the equator in the Atlantic Ocean, is doubtfully identified by Professor Verrill with a specimen obtained by us east of George's Banks.

The body in the alcoholic specimen is elevated, bell-shaped, rounded above, with a marked constriction toward the border; transparent, the inner cavity showing through as a large, conical, dark reddish brown spot, with the apex slightly truncated. Border deeply divided into sixteen long, flat lobes, which are of nearly uniform breadth throughout, and slightly rounded, or sub-truncate, at the end; the edges and end thin and more or less frilled; the inner side with two sub-marginal carinæ. Eyes inconspicuous, but small bright red specks are scattered over the marginal lobes. The intervals between the lobes are narrow and generally smoothly rounded, without distinct evidence of the existence of tentacles, except that, in one of these intervals, there is a small and short papilliform process, with brown pigment at the base. The ovaries are mostly wanting, but portions are to be seen as slightly convoluted organs in the marginal region, opposite the intervals between the lobes.

Lafoea gracillima G. O. Sars (Alder sp.).

Lafoëa fruticosa Hincks, History of British Hydroid Zoophytes, p. 202, pl. 41, fig. 2, 1868; and Annals and Magazine Nat. Hist., IV, vol. xiii, pp. 132, 148, pl. 6, figs. 6-10, pl. 7, fig. 16. 1874.

Lafoëa gracillima G. O. Sars (Alder sp.), Bidrag til Kundskaben om Norges Hydroider, in Vidensk.-Selskabs Forhandling, Christiania, for 1873, p. 115 (27), pl. 4, figs. 19-21.

Hincks reports this species from 100 fathoms off the coast of Iceland, and G. O. Sars from a depth of 150 fathoms off the Norwegian coast. It has been dredged by Professor Verrill in the Bay of Fundy and in Casco Bay.

Halecium robustum Verrill.

American Journal of Science, III, vol. v, 1873, p. 9, December, 1872.

Stem stout and coarse, composed of many tubes; branches stout, tapering, compound except at tips, pinnately or bipinnately branched, the branchlets spreading at an angle of about 45° ; yellowish white and translucent, about $\frac{1}{5}$ of an inch long, divided by simple distant constrictions, the long internodes usually bearing from two to four hydroids. Hydrothecæ alternate, large, deep, somewhat vase-shaped, with an even, slightly everted rim, below which there is a slight constriction; the middle region is slightly smaller, gradually narrowed toward the base, with a simple diaphragm near the base within. The hydrothecæ are articulated upon slightly prominent projections from the stem, in an oblique and excentric position, so as to produce a decidedly geniculated appearance. Most of the hydrothecæ are simple, but some have one or two slightly prominent secondary rims near the margin. Height about 100^{mm} .

East of St. George's Bank, 430 fathoms (haul *g*).

Sertularella polyzonias Gray, var. **gigantea** Hincks.

Annals and Magazine Nat. Hist., IV, vol. xiii, p. 151, pl. 7, figs. 11, 12, 1874.

Diphasia mirabilis Verrill.

American Journal of Science, III, vol. v, 1873, p. 9, December, 1872.

Stem stout, rather rigid, narrowed at base, pinnately branched, somewhat flexuous between the branches, which are alternate, stout, rigid, straight, constricted at base, spreading at an angle of about 45° . Hydrothecæ on the main stem in two rows, nearly opposite; on the branches mostly in six regular rows, occupying all sides of the branches, those in the adjacent rows alternating. The hydrothecæ have large, appressed, somewhat swollen bases, but the upper portion is rapidly narrowed, prominent and curved outward; aperture strongly bilabiate, operculated. Reproductive capsules not observed.

Le Have Bank, 60 fathoms (haul *e*).

Pennatula aculeata Danielsen.

Pennatula aculeata Danielsen, Forhandling i Vedenskabs-Selskabet i Christiania, 1858, p. 25 (teste Kölliker); Verrill, loc. cit., p. 100, 1873.

Pennatula phosphorea, var. *aculeata*, Kölliker, Anatomisch-systematische Beschreibung der Alcyonarien, 1 Abtheilung, 1 Hälfte, p. 134, pl. 9, fig. 73, 1870 (from Ahandlungen d. Senckenberg. Naturf. Gesellschaft, Frankfurt, Bd. vii).

Pennatula Canadensis Whiteaves, Annals and Magazine of Natural History, IV, vol. x, p. 346, November, 1872.

Pennatula, near *P. phosphorea* Verrill, Am. Journal of Science, III, vol. v, p. 5, 1873.

Localities *o* and *s*, 110 and 150 fathoms. Also dredged by Mr. Whiteaves in 200 fathoms in the Gulf of St. Lawrence.

Virgularia Lyungmanii Kölliker.

Op. cit., 2 Hälfte, 1 Heft, p. 196, pl. 13, figs. 133, 134, 1871; Verrill, American Journal of Science, III, vol. v, p. 100, 1873; Whiteaves, Report on a Second Deep-sea Dredging Expedition to the Gulf of St. Lawrence, p. 13, 1873.

This species was described by Kölliker from specimens obtained in 30 to 80 fathoms, among the Azores, by the Josephine Expedition sent out by the Swedish government. It was also dredged in 1872, in the Gulf of St. Lawrence, at a depth of 200 fathoms, by Mr. Whiteaves.

Urticina nodosa Verrill.

Actinia nodosa Fabricius, Fauna Grœnlandica, p. 350, 1780.

Urticina digitata Verrill, Am. Jour. of Science, III, vol. v, p. 5, 1873 (not of Müller?).

This species has been dredged also in deep water off Casco Bay (Professor Verrill), and in the Gulf of St. Lawrence (Mr. Whiteaves).

Cerianthus borealis Verrill.

American Journal of Science, III, vol. v, 1873, p. 5, December, 1872.

Plate II, figure 5.

Body much elongated, tapering gradually to the abactinal opening, the surface smooth but more or less sulcated longitudinally. Marginal tentacles very numerous and unequal, the inner ones longest, in the largest specimens 56^{mm} long, and 3^{mm} in diameter at base, gradually tapering, acute; the outer ones 25^{mm} and less in length. Oral tentacles numerous, crowded in several rows, in the largest specimens about 25^{mm} long, slender, acute. Color of body olive-brown or dark chestnut-brown, sometimes pale bluish or purplish just below the tentacles; disk pale yellowish-brown; space within the oral tentacles, around the mouth, deep brown, with lighter radiating lines; oral tentacles pale chestnut-brown; marginal ones deep salmon or yellowish-brown, the longest usually barred transversely with six to eight dark reddish-brown spots, each spot partially divided along the median line into two lateral ones; part of the tentacles often have flake white spots on each side, at the base.

The two largest specimens, dredged in 1872, in 28 fathoms, east of Grand Menan, by Professor Verrill, measured 125^{mm} across the disk and tentacles, but their bodies were mutilated. Entire ones of much smaller size were dredged by Dr. Packard and Mr. Cook in 110 and 150 fathoms, soft muddy bottom, hauls *s* and *o*. The largest of these was 200^{mm} long, and like other species of the genus, inhabited a thick, tough, felt-like, muddy tube. It was also dredged, in 1873, in Casco Bay, from 7 to 94 fathoms. One of these specimens, dredged off Seguin Island, in 70 fathoms, was 450^{mm} long, 40^{mm} in diameter, and 175^{mm} across the tentacles. A small specimen has been dredged in 18 fathoms off Watch Hill, R. I.

Epizoanthus Americanus Verrill.

Plate VIII, figure 2.

This species lives upon stones as well as upon shells inhabited by *Eupagurus*. The specimens from 430 fathoms (*g*) were on stones, while those from 60 and 65 fathoms (*s* and *f*) were on shells. It ranges from off the coast of New Jersey to the Gulf of St. Lawrence.

Sponges.

Most of the sponges obtained have not yet been sufficiently studied to be reported upon, but the two following species are of special interest.

Hyalonema longissimum Sars.

G. O. Sars, on some Remarkable Forms of Animal Life from the Great Depths off the Norwegian Coast, p. 70, pl. 6, figs. 35–45, 1872.

Only a single and somewhat abnormal specimen of this remarkable species was dredged by us in 430 fathoms, but it has since been dredged in considerable abundance by Professor Verrill, in 95 fathoms, off Casco Bay, and by Dr. Packard and Mr. Cooke on Cashe's Ledge.* Mr. Whiteaves reports it also from deep water in the Gulf of St. Lawrence.

Thecophora ibla Wyville Thompson.

Depths of the Sea, p. 147, fig. 24, 1873; Verrill, American Journal of Science, III, vol. vii, p. 500, pl. 8, fig. 8, 1874.

Plate VII, figure 1.

This species, first described by Wyville Thompson, from specimens dredged in 344 fathoms, off the Shetland Islands, by the Porcupine expedition, and dredged by us in 50 and 60 fathoms (hauls *e* and *d*), has since been dredged by Dr. Packard and Mr. Cook on Cashe's Ledge and Jeffrey's Ledge in the Gulf of Maine.

* American Journal of Science, III, vol. vi, p. 440, 1873.

EXPLANATION OF PLATES.

PLATE I.

- Figure 1.—*Stylifer Stimpsonii* Verrill; specimen from 60 fathoms, George's Bank (haul *i*); enlarged 10 diameters.
- Figure 2.—*Ringicula nitida* Verrill; specimen from 110 fathoms; enlarged 14 diameters.
- Figure 3.—*Entalis striolata* Stimpson; several views of animal, with the foot in different states of expansion; enlarged about $1\frac{1}{2}$ diameters.
- Figure 4.—*Astarte lens* Stimpson; adult; natural size.
- Figure 5.—The same; young specimen; natural size.
- Figure 6.—*Astarte undata* Gould; inside of valves, showing the hinge; natural size.
- Figure 7.—The same; young specimen; natural size.
- Figure 8.—The same; adult specimen; natural size.
- Figure 9.—Variety of the same; adult specimen; natural size.
- Figure 10.—*Astarte elliptica* (Brown); natural size.
- Figure 11.—*Cryptodon obesus* Verrill; inside of valve; enlarged 3 diameters.
- Figure 12.—*Astarte Banksii* Leach; natural size.

Figure 1 was drawn from nature by S. I. Smith; 2, 5, 6, 7, 8, 9, 11, by Professor Verrill; 3, by J. H. Emerton; the rest from Binney's Gould.

PLATE II.

- Figure 1.—*Sertularia argentea* Ellis and Solander; a branch bearing reproductive capsules (gonothecæ) with the soft parts removed; much enlarged.
- Figure 2.—*Acyonium carneum* Agassiz; three of the polyps fully expanded; much enlarged.
- Figure 3.—*Crisia eburnea* Lamouroux; a cluster of branches, enlarged.
- Figure 4.—The same; a branch bearing ovicells, more highly magnified.
- Figure 5.—*Cerianthus borealis* Verrill; entire animal removed from its tube and fully expanded; about one-third natural size.

Figures 1 and 2 were drawn from nature by Professor Verrill; 3 and 4 by Professor A. Hyatt; 5 by J. H. Emerton.

PLATE III.

- Figure 1.—*Xenoclea megachir* Smith, male; one of the first pair of legs with its epimeron, seen from the outside; enlarged 20 diameters.
- Figure 2.—The same; one of the second pair of legs, seen in the same position and enlarged the same amount.
- Figure 3.—The same; one of the third pair of legs, with its epimeron and gill, seen from the outside, and showing the glandular organ within; enlarged 20 diameters; *a*, the tip of the dactylus, showing the perforation; enlarged 100 diameters.
- Figure 4.—The same; one of the fifth pair of legs, with its epimeron and gill, seen from the outside; enlarged 20 diameters.
- Figure 5.—*Stenothoe pellata* Smith, female; one of the second pair of legs, with its epimeron, seen from the outside; enlarged 16 diameters.
- Figure 6.—The same; one of the fourth pair of legs, with its epimeron, seen from the outside; enlarged 16 diameters.
- Figure 7.—The same; one of the first pair of legs, seen from the outside; enlarged 50 diameters.
- Figure 8.—The same; distal portion of the propodus, with the dactylus, of one of the second pair of legs, seen from the outside; enlarged 125 diameters.
- Figure 9.—*Scalpelum Stroemi* Sars; side view; enlarged 5 diameters.

All the figures were drawn on wood, from alcoholic specimens, by S. I. Smith.

PLATE IV.

Figure 1.—*Grymæa spiralis* Verrill; head and anterior part of body; enlarged.

Figure 2.—*Pista cristata* Malmgren; head and anterior part of body; enlarged.

Figure 3.—*Phyllodoce catenula* Verrill; dorsal view of anterior part of body and head, and extended proboscis; enlarged about 4 diameters.

Figure 4.—? *Spirorbis nautiloides* Lamarck; entire animal; much enlarged.

All the figures were drawn from life by J. H. Emerton.

PLATE V.

Figure 1.—*Nephtys ciliata* Rathke; one of the lateral appendages; enlarged 10 diameters.

Figure 2.—*Lumbriconereis fragilis* (Ersted); anterior part of body and head, dorsal view; enlarged about 6 diameters.

Figure 3.—*Ninoë nigripes* Verrill; one of the lateral appendages from the middle part of the body; greatly enlarged.

Figure 4.—*Ammochares assimilis* Sars; entire animal; enlarged about 4 diameters.

Figure 5.—*Leodice vivida* Verrill; head and anterior part of the body and 12th segment; dorsal view; enlarged about 4 diameters.

Figure 1 was copied from Ehlers; all the others were drawn from nature by J. H. Emerton.

PLATE VI.

Protula media Stimpson; animal removed from the tube; enlarged 4 diameters.

Drawn from life by J. H. Emerton, from a specimen dredged near Grand Menan, Bay of Fundy, by Professor Verrill, in 1872.

PLATE VII.

Figure 1.—*Thecophora ibla* W. Thompson; specimen from 60 fathoms, Le Have Bank (haul *i*); natural size.

Figure 2.—*Eusyllis phosphorea* Verrill; anterior and posterior portions of the animal; dorsal view; much enlarged.

Figure 3.—*Nothria conchylega* Malmgren; anterior portion; enlarged.

Figure 4.—*Nothria opalina* Verrill; anterior portion; enlarged.

Figure 1 was drawn from nature by Sherman; the others from life by J. H. Emerton.

PLATE VIII.

Figure 1.—*Pera crystallina* Verrill; enlarged 3 diameters.

Figure 2.—*Epizoanthus Americanus* Verrill; a single polyp expanded; enlarged about 6 diameters.

Figure 3.—*Chectoderma nitidulum* Lovén; entire animal; enlarged 4 diameters.

Figure 4.—The same; posterior portion with the gills expanded; enlarged 24 diameters.

Figure 1 was drawn from nature by Professor Verrill; the others were drawn from life by J. H. Emerton.

ERRATA.

Page 1, line 13, for Hagerman, read Hagenman.

" 13, " 34, " *cappilare*, read *capillare*.

" 28, " 19, " **Caridon**, read **Caridion**.

" 35, " 3, " **Scapellum**, read **Scalpellum**.

" 58, " 14, " branches, read branchlets.

" 60, " 12, " Plate X, read Plate IX.

EXPLANATION OF PLATES.

PLATE I.

- Figure 1.—*Stylifer Stimpsonii* Verrill; specimen from 60 fathoms, George's Bank (haul ϵ); enlarged 10 diameters.
- Figure 2.—*Ringicula nitida* Verrill; specimen from 110 fathoms; enlarged 14 diameters.
- Figure 3.—*Entalis striolata* Stimpson; several views of animal, with the foot in different states of expansion; enlarged about $1\frac{1}{2}$ diameters.
- Figure 4.—*Astarte lens* Stimpson; adult; natural size.
- Figure 5.—The same; young specimen; natural size.
- Figure 6.—*Astarte undata* Gould; inside of valves, showing the hinge; natural size.
- Figure 7.—The same; young specimen; natural size.
- Figure 8.—The same; adult specimen; natural size.
- Figure 9.—Variety of the same; adult specimen; natural size.
- Figure 10.—*Astarte elliptica* (Brown); natural size.
- Figure 11.—*Cryptodon obesus* Verrill; inside of valve; enlarged 3 diameters.
- Figure 12.—*Astarte Banksii* Leach; natural size.

Figure 1 was drawn from nature by S. I. Smith; 2, 5, 6, 7, 8, 9, 11, by Professor Verrill; 3, by J. H. Emerton; the rest from Binney's Gould.

PLATE II.

- Figure 1.—*Sertularia argentea* Ellis and Solander; a branch bearing reproductive capsules (gonothecæ) with the soft parts removed; much enlarged.
- Figure 2.—*Alcyonium carneum* Agassiz; three of the polyps fully expanded; much enlarged.
- Figure 3.—*Crisia eburnea* Lamouroux; a cluster of branches, enlarged.
- Figure 4.—The same; a branch bearing ovicells, more highly magnified.
- Figure 5.—*Cerianthus borealis* Verrill; entire animal removed from its tube and fully expanded; about one-third natural size.

Figures 1 and 2 were drawn from nature by Professor Verrill; 3 and 4 by Professor A. Hyatt; 5 by J. H. Emerton.

PLATE III.

- Figure 1.—*Xenoclea megachir* Smith, male; one of the first pair of legs with its epimeron, seen from the outside; enlarged 20 diameters.
- Figure 2.—The same; one of the second pair of legs, seen in the same position and enlarged the same amount.
- Figure 3.—The same; one of the third pair of legs, with its epimeron and gill, seen from the outside, and showing the glandular organ within; enlarged 20 diameters; *a*, the tip of the dactylus, showing the perforation; enlarged 100 diameters.
- Figure 4.—The same; one of the fifth pair of legs, with its epimeron and gill, seen from the outside; enlarged 20 diameters.
- Figure 5.—*Stenothoë peltata* Smith, female; one of the second pair of legs, with its epimeron, seen from the outside; enlarged 16 diameters.

PLATE IV.

- Figure 1.—*Grymæa spiralis* Verrill; head and anterior part of body; enlarged.
 Figure 2.—*Pista cristata* Malmgren; head and anterior part of body; enlarged.
 Figure 3.—*Phyllodoce catenula* Verrill; dorsal view of anterior part of body and head, and extended proboscis; enlarged about 4 diameters.
 Figure 4.—? *Spirorbis nautiloides* Lamarek; entire animal; much enlarged.
 All the figures were drawn from life by J. H. Emerton.

PLATE V.

- Figure 1.—*Nephtys ciliata* Rathke; one of the lateral appendages; enlarged 10 diameters.
 Figure 2.—*Lumbriconereis fragilis* Ersted; anterior part of body and head, dorsal view; enlarged about 6 diameters.
 Figure 3.—*Ninoë nigripes* Verrill; one of the lateral appendages from the middle part of the body; greatly enlarged.
 Figure 4.—*Ammochares assimilis* Sars; entire animal; enlarged about 4 diameters.
 Figure 5.—*Leodice vivida* Verrill; head and anterior part of the body and 12th segment; dorsal view; enlarged about 4 diameters.
 Figure 1 was copied from Ehlers; all the others were drawn from nature by J. H. Emerton.

PLATE VI.

- Protula media* Stimpson; animal removed from the tube; enlarged 4 diameters.
 Drawn from life by J. H. Emerton, from a specimen dredged near Grand Menan, Bay of Fundy, by Professor Verrill, in 1872.

PLATE VII.

- Figure 1.—*Thecophora ibla* W. Thompson; specimen from 60 fathoms, Le Have Bank (haul *i*); natural size.
 Figure 2.—*Eusyllis phosphorea* Verrill; anterior and posterior portions of the animal; dorsal view; much enlarged.
 Figure 3.—*Nothria conchylega* Malmgren; anterior portion; enlarged.
 Figure 4.—*Nothria opalina* Verrill; anterior portion; enlarged.
 Figure 1 was drawn from nature by Sherman; the others from life by J. H. Emerton.

PLATE VIII.

- Figure 1.—*Pera crystallina* Verrill; enlarged 3 diameters.
 Figure 2.—*Epizoanthus Americanus* Verrill; a single polyp expanded; enlarged about 6 diameters.
 Figure 3.—*Chatoderma nitidulum* Lovén; entire animal; enlarged 4 diameters.
 Figure 4.—The same; posterior portion with the gills expanded; enlarged 24 diameters.
 Figure 1 was drawn from nature by Professor Verrill; the others were drawn from life by J. H. Emerton.

ERRATA.

- Page 5, line 30, for *Vetumnus*, read *Vertumnus*.
 " 9, last line, for 1873, read 1874.
 " 11, line 31, for *virticillata*, read *verticillata*.

2
1
III.—DESCRIPTIONS OF NEW AND RARE SPECIES OF HYDROIDS FROM
THE NEW ENGLAND COAST. BY S. F. CLARK.

THE material for this paper was gathered while at work on the Hydroids in the Museum of Yale College. This collection is mostly from the New England Coast, and is very large and complete.

Obelia bicuspidata, sp. nov.

Plate IX, fig. 1.

The stem is erect, slender, straight or nearly so, compound, consisting of many united tubes which gradually diminish in number toward the top, varying in color from a light horn, to a light whitish brown, sparingly branched, and with three or four annulations just above the origin of each branch; branches short, ascending, slender and irregularly arranged, sometimes one, and often two branches starting from a node; branches few, very short, slender and ascending.

Hydrothecæ very deeply campanulate, narrow, tapering slightly toward the base, very hyaline, and with eight to ten longitudinal lines extending from the distal extremity nearly to the base; the rim is armed with very acute teeth, varying in number from sixteen to twenty-two, and arranged in pairs, the spaces in which the longitudinal lines terminate being a trifle wider and deeper than the alternate spaces; the pedicels supporting the hydrothecæ are long and tapering, consisting of about fifteen annulations. Gonothecæ unknown.

Height of largest specimens, about three inches (80^{mm}).

The specimens from which this species is described were taken in 3-5 fathoms, on the reefs near Thimble Islands, Long Island Sound, September 23, 1874.

This species is closely allied to *O. bidentata*, but is readily distinguished from the latter by its entirely different habit, the narrower and deeper calyces, and by the long tapering pedicels upon which the calyces are supported.

Obelia bidentata, sp. nov.

Plate IX, fig. 2.

Stems clustered, straight or slightly flexuous, compound, composed, at the base, of eight or ten slender, united tubes, varying in color from a light horn to a dingy white, densely branched, and with three

or four annulations just above the origin of each branch; mode of branching irregular, two branches often starting from a node, sometimes an alternate arrangement of branches on opposite sides of the stem, one branch at each node. A few of the lowest branches sometimes attain a considerable length and resemble the main stem, the upper ones are short, sparingly branched and with the pinnae diverging at a slight angle; the branchlets and ends of the branches are simple, slender, translucent, and very graceful. Hydrothecæ very deeply campanulate, tapering slightly toward the base, and with nine to twelve longitudinal lines extending from the distal extremity nearly to the base; the rim is ornamented with from eighteen to twenty-four very acute teeth, arranged in pairs, the spaces in which the longitudinal markings terminate being a trifle wider and deeper than the alternate spaces; the pedicels supporting the hydrothecæ are usually short and stout, consisting of three to six strong rings, but some of the hydrothecæ near the base of the stem have the pedicel slightly tapering, and composed of from ten to twelve annulations. Gonothecæ unknown.

The largest specimen has a height of about 6 inches (150^{mm}).

We have had this species from but one locality, Greenport, Long Island, where it was collected August 5th, 1874, in considerable abundance, on the piles of the wharves at low water,—U. S. Fish Commission.

O. bidentata resembles *O. gelatinosa* in the delicacy and grace of its habit, in the flexibility of the compound stem and branches, and in the pellucid whiteness of the upper portion of its branches and branchlets.

Campanularia pygmæa, sp. nov.

Plate IX, fig. 9.

Stem often creeping, with short, stout, coarsely annulated, upright pedicels, sometimes with one or two short, annulated branchlets, each bearing a single calycle. Hydrothecæ large, deep campanulate, tapering slightly toward the base, and with longitudinal lines at regular intervals, extending down about one-fourth the length of the calycle; the rim is ornamented with from ten to fourteen square-cut denticles, which are more or less hollowed out above, and separated by rather shallow evenly rounded notches, of about the same breadth. Gonothecæ unknown. Height about 1^{mm}.

Found growing on a specimen of *Sertularia latiuscula*, from Casco Bay, Maine,—U. S. Fish Commission.

Campanularia noliformis McCrady.

Plate X, fig. 5.

A few specimens of this species were collected at low-water, near Savin Rock, in the latter part of September, 1874, attached to *Zostera marina*.

The hydrarium agrees very well with McCrady's description. We were not fortunate enough to find the gonothecæ.

This is the first time this species has been recorded since McCrady described it from the harbor of Charleston, S. Carolina, in 1857.

Campanularia calceolifera Hincks.

Ann. and Mag. of Nat. Hist., vol. viii, Aug., 1871, page 78, pl. vi.

Plate X, figs. 7, 8.

Stem filiform, slender, flexuous, sometimes slightly branched, ringed at the base and above the origin of the branches, light horn-color, with the upper portions pellucid white; branches short, curving outward, undivided, and bearing but two or three calyces. Hydrothecæ alternate, broadly campanulate, deep, with a slightly everted entire rim, and borne on annulated pedicels of variable length, those on the upper portion of the stem consisting of five to eight rings, those near the base, of twelve to twenty. At each bend of the stem a single hydrotheca is given off, and these all curving outward give to this species a very graceful habit. Gonothecæ axillary, borne on pedicels consisting of three or four rings, largest at the distal extremity and tapering gradually toward the base, with a peculiar incurved coil or twist at the distal end near the opening; the aperture is shield-shaped and placed in a depression on one side of the distal end. An internal membrane extends inward from the shield-shaped opening and terminates in a circular orifice near the distal extremity. Height about one inch (25^{mm}).

Noank, Conn., from the bottom of an old scow, Sept. 9, 1874, with gonothecæ; piles of wharves at Woods Hole, Mass., Aug., 1871, with gonothecæ—U. S. Fish Commission.

Hincks' figures represent the hydrothecæ as being more everted than they are in the American specimens; otherwise they exactly correspond. This is the first time this species has been recorded from the American coast.

Gonothyræa tenuis, sp. nov.

Plate X, fig. 8.

Stem simple, somewhat flexuous, slender, and annulated above the origin of each of the numerous branches, which are arranged alternately, some simple and some compound, the latter bearing numerous branchlets, the lower branches sometimes half the length of the main stem; base of the stem and of the lower branches light horn-color, the upper portions of the same and all of the branchlets pellucid white; branches and branchlets spreading, giving quite a bushy appearance to a well developed colony. Hydrothecæ variable in size, deeply campanulate, tapering quite rapidly from a little below the middle to the base; the rim is ornamented with teeth which show considerable variation, both in number and in shape; in some cases they are quite sharp and shallow, while on other calyces upon the same stem they are of a castellated form and sometimes slightly emarginate; in number they vary from ten to sixteen; the pedicels which support the hydrothecæ also vary greatly, some being composed of but three or four annulations, others of as many as fourteen.

Gonothecæ axillary, very much elongated, narrow, obconic, tapering gradually from the distal to the proximal end, borne on short pedicels of but three to five rings; the number of medusæ holding planulæ contained in each reproductive capsule varies from two to five and the number of planulæ in each medusa varies to the same extent; the tentacles of the medusæ vary considerably in length and in number, some of them being over half as long as the diameter of the medusa, while others are not more than one-third that length, in number they vary from eight to fourteen. The planulæ at the time of liberation are regularly cylindrical, and their length is equal to nearly four times the width. Height usually 1 to 1.25 inches (25 to 38^{mm}).

New Haven, Conn., on piles of Long Wharf, June 2nd, 1875, — S. F. Clark. Found in considerable abundance at low-water, loaded with reproductive capsules. The large size of the latter, together with the clusters of extracapsular medusæ, make this quite a showy species for one of such humble growth.

Opercularella pumila, sp. nov.

Plate IX, figs. 3, 4, 5.

Stem rather stout, erect or creeping, slightly flexuous, annulated throughout, sparingly branched; branches erect, undivided, some-

times attaining a considerable length. Hydrothecæ largest in the middle, tapering very slightly toward the base, rapidly converging at the distal end, and supported on short annulated pedicels, consisting of three to five rings. Gonothecæ fusiform, with the tapering neck often somewhat elongated; length about twice that of the hydrothecæ. They contain one to three small globular or ovate immature medusoids. The pedicels consist of three to six annulations. In the creeping form the hydrothecæ appear at intervals borne on short ringed stalks consisting of about three to six rings.

Portland, Maine, August, 1873, with gonothecæ, on piles of wharves; off Montauk Pt., Long Island, 5-15 fathoms, August, 1874,—U. S. Fish Commission.

This species closely resembles *O. lacerata* of Hincks, from which it is distinguished by the forms both of the hydrothecæ and gonothecæ. The hydrothecæ are also smaller than those of *O. lacerata*. As the reproduction has not been traced in this species, it can only be referred to the genus provisionally. I am inclined to think, from the shape of the gonothecæ and from the fact that they often contain two or three distinct reproductive bodies, that it may not belong under *Opercularella*.

Opercularella lacerata Hincks.

Campanularia lacerata Johnston, Brit. Zoöph., p. 111, Pl. xxviii, fig. 3.

Opercularella lacerata Hincks, Brit. Hydr. Zoöph., p. 194, Pl. xxxix, fig. 1.

Plate IX, fig. 6.

Stem erect, simple, slightly flexuous, more or less annulated throughout, sparingly branched; branches short undivided. Hydrothecæ ovato-fusiform and borne on short pedicels of but two or three annulations; operculum composed of six to eight segments. Gonothecæ, of the female colonies, a trifle wider at the distal end, and tapering very slightly toward the base, supported on short ringed pedicels; the medusoid (sporosac) containing the planulæ is quite large, the diameter being about equal to the length of the gonotheca; from two to five planulæ in each medusoid.

New Haven, Conn., on piles of Long Wharf, May 13th, 1875, with extracapsular medusæ.—S. F. Clark.

The hydrothecæ in this species average about one third larger than those of *O. pumila*; the segments of the opercula are more deeply cleft; and there are differences in the form of the gonothecæ.

Calycella syringa Hincks. Peculiar variety.

Plate X, figs. 1, 2, 3.

Sertularia syringa Linn., Syst. 1311.

Calicella syringa Hincks, Cat. Devon Zooph., 23; Ann. N. H. (3d ser.), VIII, 294.

Calycella syringa Hincks, British Hydroid Zoophytes, Vol. I, p. 206, Plate xxxix, figs. 2, 2a.

Stem simple, creeping, nearly smooth. Hydrothecæ hyaline, colorless or tinged with a light horn-color, cylindrical, rounded off below, with an everted rim, to which is attached an operculum consisting of from five to eight segments and supported on twisted pedicels of considerable length, with eight to fifteen twists; some of the hydrothecæ have an addition in the shape of a wide ring, ornamented with from ten to fourteen longitudinal markings, which rises for some distance above the rim and on the summit of which there is borne either the operculum or another ring; in some cases there are as many as four of these rings with an operculum at the summit. The opercula usually point upward, but are occasionally deflected into the calycele.

From Casco Bay, Me., 9 fathoms,—U. S. Fish Commission.

Halecium articulatum, sp. nov.

Plate X, fig. 6.

Stem dark brown and tapering gradually, very stout, sparingly branched, compound, consisting of many, slender, anastomosing, serpentine tubes; branches short and irregularly arranged on all sides of the main stem; branchlets few and very short; both branches and branchlets are divided into very short, stout internodes by distinct joints placed at right angles to the stem; branches and branchlets simple, whitish, delicate, becoming more numerous toward the top of the stem. The internodes become shorter very gradually toward the ends of the branches and branchlets. Hydrothecæ alternate, short and wide, one to each segment; some of them have a cup within a cup, as is so often the case in the species of *Halecium*. Gonothecæ borne in rows on the upper side of the pinnae. The female gonothecæ are large, obovate, and have the opening on one side and nearer the distal than the proximal end; the male gonothecæ are oblong, subcylindrical, and, like the female, are sessile. Height of the largest specimens, 5 inches (125^{mm}).

Eastern end of Long Island Sound, 8–12 fathoms; Cox's Ledge, S. E. of Block Island, 17–21 fathoms; Casco Bay, Maine; Eastport, Me.,—U. S. Fish Commission.

H. articulatum resembles *H. plumosum*, but has a stouter habit; the internodes are shorter and stouter; and it may also be distinguished by the direction of the joints, at right angles to the stem, and by the very wide-mouthed calyces. The female gonothecæ somewhat resemble those of *H. Beanii*, but are of a stouter build; they are relatively larger at the distal extremity, the orifice is differently shaped and is differently situated, being nearer the distal extremity.

Sertularia argentea Ellis and Solander, var. *divaricata* nov.

Plate X, fig. 7.

Stem simple, stout, erect, straight or slightly flexuous, of a deep horn color, regularly jointed, each joint having two or three branches; branches alternate, sparingly branched, diverging at right angles from the main stem and all in the same plane, divided quite regularly by joints, each bearing two pairs of hydrothecæ, much resembling a young shoot of the usual form of *S. argentea*. Hydrothecæ nearly opposite, curving strongly outwards, with a bilabiate mouth, the upper lip being considerably smaller than the lower; hydrothecæ are also scattered along the stem in pairs. Gonothecæ unknown.

Collected at Casco Bay, Me., 1873,—U. S. Fish Commission.

I at first thought that this was a distinct species, but I have since had intermediate forms which prove quite conclusively that it is only a variety of *S. argentea*. Considerable variation is shown in the hydrothecæ of this variety; some of them, on the same stem, are more directly opposite and curve outwards more than others.

Plumularia Verrillii, sp. nov.

Plate X, fig. 9.

Stems erect, simple, straight or slightly curved, slender, two to four inches high, of a bright horn-color, branched and regularly jointed by transverse divisions; the branches have their origin near the base of the stem, are erect and resemble the main stem in all particulars; pinnæ occasionally branched, regularly arranged on two sides of the main stem and branches, separated by an angle of ninety degrees, composed of long similar joints, each bearing a hydrotheca and a number of nematophores; occasionally there is an odd, intermediate joint bearing only one or two nematophores and no hydrothecæ; a single pinna to each joint. Nematophores sessile, compound, large, tapering to the base, with a round cup-like opening: there are four to six on each hydrotheca-bearing joint, one on each side of the upper

edge of the hydrotheca, two or three between the hydrotheca and the proximal end of the joint, and occasionally one at the distal end; on the main stem there are usually two in each axil and two or three on each joint. Hydrothecæ small, shallow, about equal in depth to the length of the nematophores, attached to the stem by their entire length; rim entire. Gonothecæ borne in the axils of the stem and of the branched pinnæ, sessile, tapering at the base, the remaining portion either regularly cylindrical or slightly swollen in the middle; aperture large, terminal. Height, 2.5 inches (64^{mm}).

Eastport, Maine, 10–20 fathoms, 1868,—A. E. Verrill and S. I. Smith.

P. Verrillii is a beautiful little, pellucid, white species, with a delicate, graceful habit which readily distinguishes it from any of the forms now known upon our coast. It is the second genuine *Plumularia* from the New England coast, both having been discovered by Professor Verrill. The previously described species (*P. tenella* Verrill) was dredged in 1871, off Gay Head, Martha's Vineyard, in 10 fathoms. It has since been dredged in 4–5 fathoms off the Thimble Islands, near New Haven, Conn., and it was also found on the piles at Greenport, Long Island, August 5th, 1874, with gonothecæ. It differs greatly from the present species in the form of the gonothecæ, which are in the shape of an elegantly curved cornucopia, slender at the base and gradually enlarged to the end, and with a cluster of nematophores at the base. The hydrothecæ are also different in form.

EXPLANATION OF PLATES.

PLATE IX.

- Figure 1. *Obelia bicuspidata*; from Thimble Islands.
 Figure 2. *Obelia bidentata*; from Greenport, Long Island.
 Figure 3. *Opercularella pumila*; creeping form.
 Figure 4. The same, young, with stem erect; from off Montauk.
 Figure 5. The same, from Portland, Me., showing a more luxuriant growth; *a* and *c*, the hydrarium; *b*, gonotheca, enlarged 32 diameters.
 Figure 6. *Opercularella lacerata*; *a*, hydrarium; *b*, gonotheca; *c*, medusoid; *d*, undeveloped planulæ.
 Figure 7. *Campanularia calceolifera*; from Noank, Conn.
 Figure 8. The same; *a*, hydrotheca; *b*, gonotheca; from Noank, Conn.
 Figure 9. *Campanularia pygmæa*; from Casco Bay.

PLATE X.

Figure 1. *Calycella syringa*; from Casco Bay, showing a peculiar variation in the operculum.

Figure 2. The same, showing the variation in the size and shape of the hydrothecæ and in the length of the pedicel.

Figure 3. The same, with one secondary ring.

Figure 4. *Clytia Johnstoni*, from Noank, Conn.

Figure 5. *Campanularia notiformis*, showing variations in the pedicels.

Figure 6. *Halecium articulatum*; from Coxe's Ledge. A, a branch bearing both hydrothecæ and gonothecæ; a, gonothecæ; b, hydrothecæ. B, a branch with hydrothecæ only; b, a branchlet; c, hydrothecæ.

Figure 7. *Sertularia argentea*, var. *divaricata*; from Casco Bay.

Figure 8. *Gonothyræa tenuis*; a, branch with hydrothecæ; b, gonotheca with extracapsular medusæ; c, medusa with radiating tubes and tentacles; d, planula.

Figure 9. *Plumularia Verrillii*; a, branch showing hydrothecæ and nematophores and the arrangement of the joints; b, gonotheca; c, a single joint.

III. ON THE CHONDRODITE FROM THE TILLY-FOSTER IRON MINE,
BREWSTER, NEW YORK. BY EDWARD S. DANA.

WITH THREE PLATES.

THE interesting discovery by Scacchi,* of the existence of three types in the crystals of the Vesuvian humite, gives especial interest to the study of chondrodite—a mineral identical with humite in chemical composition, and yet very different in appearance, as well as in origin and method of occurrence. The same subject of humite has since been more exhaustively investigated by vom Rath,† with the entire confirmation of Scacchi's views. These authors have shown that the crystals of humite are to be divided into three groups, all bearing the same relation to each other in respect to their lateral axes, while the vertical axis has a distinct value for each type. In other words, the planes occurring upon a given crystal bear simple relations to each other, whereas only very complex symbols result when the planes of one type are referred to the axes of another. For a full explanation of this subject reference must be made to the valuable memoirs above alluded to. It will be sufficient to give as an example the symbols of the occurring pyramids of the ν series on the second and third types of chondrodite (see beyond); (1) as referred to their own axes; (2) as referred to the axes of the second type.

(1.)		(2.)	
II.	III.	II.	III.
$\frac{4}{7}\sqrt{2} (\nu^1)$	$\frac{8}{11}\sqrt{2} (\nu^3)$	$\frac{4}{7}\sqrt{2}$	$\frac{36}{55}\sqrt{2}$
$\frac{4}{5}\sqrt{2} (\nu^2)$	$\frac{8}{9}\sqrt{2} (\nu^4)$	$\frac{4}{5}\sqrt{2}$	$\frac{36}{45}\sqrt{2}$
$\frac{4}{3}\sqrt{2} (\nu^3)$	$\frac{8}{7}\sqrt{2} (\nu^5)$	$\frac{4}{3}\sqrt{2}$	$\frac{36}{35}\sqrt{2}$
$4\sqrt{2} (\nu^4)$	$\frac{8}{5}\sqrt{2} (\nu^6)$	$4\sqrt{2}$	$\frac{36}{25}\sqrt{2}$
	$\frac{8}{3}\sqrt{2} (\nu^7)$		$\frac{36}{15}\sqrt{2}$
	$8\sqrt{2} (\nu^8)$		$\frac{36}{5}\sqrt{2}$

* Pogg. Ann., Erg. Bd. iii, 161, 1851.

† Pogg. Ann., Erg. Bd. v, 321, 1871; vi, 385, 1873.

The chemical composition of the three types of humite has been most recently investigated by vom Rath,* and although analyses lead to somewhat different results in the three cases, he concludes that in composition they are still essentially the same, and that the cause of the variation in crystalline form is *not* to be found in the relative amount of fluorine present, as has been often assumed.

A further remarkable peculiarity true of two of the three types is their hemihedral character, which is clearly set forth in the memoirs referred to. These points are alluded to here because of their direct bearing on the crystallization of chondrodite, which forms the subject of this paper.

Chondrodite was first shown by Rammelsberg to be identical with humite in chemical composition, but its crystallographic relation to it was not brought out until the investigations of Kokscharow. He showed, in his "Materialien zur Mineralogie Russlands," vol. vi, p. 73, 1870, that the crystals from Pargas, Finland, were identical in form and angles with type II of humite. Vom Rath has followed with the description of crystals from Nya-Kopparberg, Sweden, and proved that the same fact is true of them.

The study of the chondrodite from the Tilly-Foster iron mine, Brewster, Putnam Co., New York, which I have been able to make during the past season, has shown that it, too, is for the most part identical in crystalline form with type II of humite, but that at the same time crystals exist belonging to *type I*, and others which belong to *type III*. Further than this, the chemical composition of the second type crystals, as shown by an analysis by Mr. G. W. Hawes (p. 21), agrees with great exactness with that of the Swedish mineral analyzed by vom Rath. Moreover, the detailed study of these crystals has shown that while they agree with humite in the character of their hemihedrism, as well as in angles, they surpass it in the multiplicity of secondary planes. Thus a single solid angle has been observed which was modified by fifteen distinct and well-defined, though very minute, planes. This, as will be seen when the facts are described in detail, implies a delicacy in the action of the crystallogenic forces at work which is unparalleled, and sustains the opinion that chondrodite, or humite, is unique among mineral species.

The method of occurrence at the Tilly-Foster iron mine has been fully described by Prof. Dana in a memoir entitled, "Serpentine pseudomorphs and other kinds, etc.," *Journal of Science*, viii, pp. 371,

* Pogg. Ann., cxlvii, 246, 1872.

447, 1874. It may be of interest, however, to review the subject again so far as the chondrodite itself is immediately concerned.

The chondrodite forms the gangue of the magnetite, being everywhere disseminated through it in varying proportions. In the parts of the mine where the ore is purest and perfectly firm and solid,—the so-called “blue ore,”—the associated chondrodite is sparsely sprinkled through it in small yellow grains, showing no trace of crystalline form. Occasionally, however, the firmer ore contains the chondrodite in very large but imperfect crystals, or crystalline masses, associated directly with enstatite and enveloped with dolomite, which have a dark, rich brown color, and a brilliant luster on the fracture. A distinctly laminated structure is uniformly present, which is perhaps due to cleavage (?) (See page 21.) Isolated grains imbedded in dolomite often show traces of crystalline faces, though nothing that admits of even approximate determination. An analysis of this variety of the mineral gave Mr. Breidenbach (*Am. J. Sci.*, III, vi, 209),

Si 35.42, Fe 5.72, Mg 54.22, Fl 9.00=104.36; equivalent of oxygen replaced by fluorine, 3.79.

In the larger portion of the mine as now opened the soft “yellow ore” predominates: the chondrodite is present in it in much larger quantities, and, like the other minerals present, it has almost universally suffered extensive alteration. A long list of these products of alteration has been fully described by Prof. Dana in the memoir already alluded to. The chondrodite forms the main portion of the material taken out, and many tons of this refuse matter are yearly thrown away. It varies much in color, but is generally of a light yellow; it usually has more or less of a soapy feel and shows a variety of transition-products between the semi-altered material and serpentine. The chondrodite in this “yellow ore” is generally massive; but occasionally fragments of large coarse crystals have been found, some of which measure five or six inches in length. These are always more or less altered; moreover, the material of which they are formed is far from homogeneous, masses of magnetite, and also chlorite, being often enclosed. Dolomite is the most constantly associated mineral and occurs in rhombohedrons of considerable size; these, as well as the crystals of chondrodite, are often coated with magnetite.

Better crystals of chondrodite than those just mentioned are sometimes found in what were once cavities in this massive material. Unfortunately these have all suffered from the general alteration and now have little or no luster, and often are not even smooth. These cavities are almost invariably filled with a soft mealy serpentine,

which can be cut out with a knife. These crystals vary in size, being sometimes an inch or two in length. A crystallographic examination of them is seldom possible, but a few of the crystals found allow of it, and the results are described beyond. The form is usually very simple, and the color varies from a deep red to a light yellow. This may be said to be the common method of occurrence at the locality.

Fortunately, material much better adapted for crystallographic study also occurs, though this is very rarely true. Narrow veins are sometimes met with, two or three inches across, which were originally lined with more or less perfectly crystallized chondrodite and also with dodecahedrons of magnetite, crystals of ripidolite, and rarely apatite, and then subsequently filled in with dolomite. Where this has been the case and the dolomite has remained intact the chondrodite has been protected and the crystals have retained perfectly their luster and color. Only in a few instances were the crystals polished when covered simply by a soft serpentine. The chemical composition of this chondrodite is given beyond, after the description of the crystals. It has a deep, garnet-red color, and a luster equal to that of the finest Binneenthal blende.

1. *Description of Crystals belonging to Type II.*

The remark of vom Rath in regard to the irregularity of form of the Swedish chondrodite is eminently true of the Brewster crystals. For in the same little group no two are alike; so that each one deserves and requires an especial study.

The first point to be determined was the values of the fundamental angles. Some difficulty was found in obtaining these from the fact that many crystals, though faultless in luster, yet gave uncertain measurements. This was due to the fractured condition of many of the planes, which, though often not very apparent at first sight, yet gave rise to a variety of reflected images in the goniometer, of which no one could be accepted as trustworthy. All the larger crystals show a multitude of internal fractures; and, where such crystals have been subjected to altering influences, this circumstance has hastened their destruction, and in all cases the external condition of the planes has been more or less affected. The direction of the fracture lines was in most cases entirely irregular, though in a number of cases they were distinctly parallel to $e^2(2-\bar{i})$. The presence of these cracks gave the crystals the appearance of having suffered sudden contraction, by which the planes had been irregularly drawn inward, forming re-entrant angles; in fact, in this respect, as in general appearance, this chondrodite might be aptly compared to a resin.

The smallest crystals proved to be free from this cause of irregularity, and one of them, on which the faces were exceptionally brilliant, was chosen for careful measurement. It may be added that all the measurements were made with an Oertling goniometer, provided with two telescopes.

The mean of 30 measurements of A ($O=001$) on $r^1 \left(\frac{4}{7} \bar{2} = \bar{2}47 \right)$ gave: $135^\circ 18' 50''$. The maximum variation from the mean given was $\pm 45''$. The mean of 30 measurements of A on $e^a \left(\frac{2}{5} \bar{3} = 205 \right)$ gave: $149^\circ 55' 48''$. Maximum variation $\pm 45''$.

These were accepted as the fundamental angles, and as the agreement between the other angles measured and the calculations made

TABLE I.

Chondrodite.

Humite.

			$C=i\bar{i}$ (010) Calculated.	$A=O$ (001) Measured.	Calculated.	A (v. Rath) Calculated.
i	$1\bar{i}$	011	147° 32' 39"	122° 44' (ap.)	122° 27' 21"	122° 27' 49"
e^a	$\frac{2}{5}\bar{3}$	205	90	*149 55 48	149 58 48	
e^1	$\frac{2}{3}\bar{3}$	203	90	135 59	136 1 17	135 52 15
e^2	$2\bar{3}$	201	90	109 4	109 3 24	108 57 50
r^1	$\frac{4}{7}\bar{2}$	$\bar{2}47$	129 42 9	*135 18 50	135 18 50	135 17 40
r^2	$\frac{4}{5}\bar{2}$	245	137 25 45	125 52	125 50 6	125 49 0
r^3	$\frac{4}{3}\bar{2}$	$\bar{2}43$	146 27 42	113 25½	113 25 36	113 24 45
r^4	$4\bar{2}$	241	154 2 9	98 14	98 13 6	98 12 47
m^2	$6\frac{3}{2}\bar{3}$	$\bar{6}41$	125 43 56	95 22	95 19 40	95 17 59
n^1	$\frac{2}{3}$	$\bar{2}23$	127 1 31		125 3 49	125 2 47
n^2	2	221	135 45 24	103 11	103 10 4	103 9 35

upon the above basis proved to be as close as could be desired, having regard to the nature of the plane in each individual case, no

attempt to correct them was deemed desirable. Calculated from these angles the parameters are :

$$a \text{ (vert.)} = 1.57236 ; b = 1 ; c = 1.08630 ;$$

and the angle for the fundamental prism is

$$I \wedge I (110 \wedge \bar{1}\bar{1}0) = 85^\circ 15' 46'' \text{ or } 94^\circ 44' 14''.$$

It may be added that the angle of A on r' behind ($\bar{2}\bar{1}7$) was measured with equal care, and found to be $135^\circ 18' 40''$.

The preceding table* (I) includes the principal angles measured on the same crystal, and also those calculated from the above parameters; in addition, the corresponding angles for humite, type II, are also given, as calculated by vom Rath. The angles of the macrodomes agree very closely, it will be observed, in chondrodite and humite; in the brachydomes, on the other hand, there is a divergence of 6 or 7 minutes.

The angles given in table I, and also in tables XII, XIII, XIV, for types I and III, are the actual angles. In all the other tables, however, the supplement (normal) angles, as measured and calculated, are uniformly given. The reason for this was that these angles having special reference to the sphere of projection, and being chiefly of value in calculating with it, it did not seem worth while to change them from the form in which they had been used.

It is necessary to explain, also, the system of symbols and letters here and subsequently employed. The fundamental form adopted is the same as that used in Dana's "Mineralogy," and first suggested by the author of that work in *Am. J. Sci.* II, xiv, 175, 185. It is to be remembered that Scacchi made the prism of the r series (i. e., $i\bar{2}$) the fundamental prism, and gave to the vertical axes lengths, in the ratio of 7 : 5 : 9 to each other, for types I, II and III respectively. Vom Rath followed him in this latter respect, but for the vertical prism took that of the n series (i. e., I , or 110 of Dana). Prof. Dana took a modified view of the relations of the three forms, and chose for the fundamental macrodome in each type the plane making with $A(O)$ an angle of 122° to 125° : thus on humite (vom Rath), type I, $124^\circ 17'$, II, $122^\circ 28'$, III, $125^\circ 15'$. In this view the vertical axes have one-third (I), one-half (II) and one-quarter (III) of the lengths assumed by Scacchi; in other words, their relation to each other are

* Both the symbols of Naumann (in the form used in Dana's "Mineralogy") and also of Miller are given; the signs belonging to each plane omitted here, as the relations of the planes are shown with sufficient clearness on the spherical projection, Plate xiii.

as $\frac{7}{3} : \frac{5}{2} : \frac{9}{4}$ or 28 : 30 : 27. The method adopted by Scacchi and vom Rath has the advantage of expressing the simplest possible numerical ratios between the three types. It is the view of the author above referred to, however, that the variation in the angle of $\Delta \wedge 1\text{-}\bar{7}(011)$ is to be regarded in the same light as the variation in the vertical axes of the rhombohedral carbonates, or in the orthorhombic sulphates; or in other words, the three types form an isomorphous series, and the variation observed is no greater than is constantly seen in analogous isomorphous groups. This view seems to find confirmation in the crystallographic relations of humite and *chrysolite*, a subject already discussed by Scacchi, Rammelsberg and vom Rath. Taking the fundamental form, as in Dana's "Mineralogy" (here $i\text{-}\bar{2} = I$ of other authors), the lateral axes are nearly identical with those of humite, while the vertical axis (1.25928 Kokscharow) has exactly the ratio of $\frac{8}{15}$ to that of humite, type II, and $\frac{8}{9}$ to that of humite, type III; in other words, we have the ratios:

	Humite.		
Chrysolite.	-----		
	III.	I.	II.
24	: 27	: 28	: 30.

If we adopt the vertical axes of Scacchi and quadruple that of chrysolite, we obtain

II.	I. Chrysolite.	III.
5	: 7	: 8 : 9

These relations were in effect brought out by Scacchi when he showed that what he called the common fundamental form of humite, obtained by dividing the vertical axes by 7 (I), 5 (II), and 9 (III) respectively, was nearly identical with that accepted by him for chrysolite. This fact seemed to Rammelsberg of so much importance that he proposed to refer all the planes of humite to this common fundamental form; and in this he has been followed by Kokscharow. The result of this will be seen in the following table, which gives the symbols thus obtained for the two common macrodomes in each type.

I.	II.	III.
$\frac{7}{10} \left(\frac{3}{5} \text{-}\bar{7} \right)$	$\frac{5}{8} \left(\frac{1}{2} \text{-}\bar{7} \right)$	$\frac{3}{4} \left(\frac{2}{3} \text{-}\bar{7} \right)$
$\frac{7}{6} (1\text{-}\bar{7})$	$\frac{5}{4} (1\text{-}\bar{7})$	$\frac{9}{8} (1\text{-}\bar{7})$
$\frac{7}{2} (3\text{-}\bar{7})$		$\frac{9}{4} (2\text{-}\bar{7})$

It will be seen from these few examples that the plan proposed introduces a set of common molecular axes at the expense of all crystallographic simplicity in the relations of the several planes of each type. Moreover, the view of Rammelsberg loses some of its plausibility, if, as shown by vom Rath, the vertical axes do *not* stand in direct relation to the amount of fluorine present. The view of Prof. Dana here advocated seems to have the advantage of presenting all the relations in their most natural light.

It may be added, as completing the history of the subject, that DesCloizeaux refers all the humite planes to one, and that the second, type.

In regard to the letters employed, it seemed to offer the simplest solution of an obvious difficulty to retain all the letters of Scaechi for the second type, and for the third to use simply the corresponding Greek letters in the same order, and for the first type to use the corresponding capital letters. It was not deemed advisable to use the same letter for two planes, on different types, which bore no immediate relation to each other.

TABLE II.
A = O(001).

			Calculated.	II.	III.	IV.
i^1	1- $\bar{7}$	011	57° 33'	57° 28'		
e^a	$\frac{2}{5}\bar{5}$	205 30 4				
e^1	$\frac{2}{3}\bar{5}$	203 43 58 $\frac{1}{2}$				
e^2	2- $\bar{5}$	201 70 57				
r^1	$\frac{4}{7}\bar{2}$	$\bar{2}47$ 44 41	$\left\{ \begin{array}{l} 44 \quad 40\frac{1}{2} \\ 44 \quad 40 \end{array} \right.$	44° 37'	44° 41'	
r^2	$\frac{4}{5}\bar{2}$	$\bar{2}45$ 54 10		54 12		
r^3	$\frac{4}{3}\bar{2}$	$\bar{2}43$ 66 34	$\left\{ \begin{array}{l} 66 \quad 39 \\ 66 \quad 32 \end{array} \right.$	66 25	66 35	
r^4	4- $\bar{2}$	$\bar{2}41$ 81 47		81 52		
n^1	$\frac{2}{3}$	$\bar{2}23$ 54 56				
n^2	2	$\bar{2}21$ 76 50				
m^2	6- $\frac{3}{2}$	$\bar{6}41$ 84 40				

TABLE III.
C = $i\bar{7}$ (010).

Calculated.	II.	III.	IV.
32° 27'	$\left\{ \begin{array}{l} 32^\circ \quad 28' \\ 32 \quad 32 \end{array} \right.$		
90			
90			
90			
50 18	$\left\{ \begin{array}{l} 50 \quad 7 \\ 50 \quad 30 \end{array} \right.$	$\left\{ \begin{array}{l} 50^\circ \quad 13' \\ 50 \quad 30 \end{array} \right.$	50° 12'
42 34	42 39	42 32	42 35
33 32	$\left\{ \begin{array}{l} 33 \quad 31 \\ 33 \quad 32 \end{array} \right.$	33 30	33 27
25 58		$\left\{ \begin{array}{l} 26 \quad 0 \\ 25 \quad 57\frac{1}{2} \end{array} \right.$	
44 14 $\frac{1}{2}$		44 11	
52 58			
54 16		54 18	

In the tables, II, III, IV, are given the angles as measured in a series of crystals (each crystal is numbered). They are important as showing how far the angles are constant. Some considerable variations from calculated angles in a few instances are to be explained by the cause of irregularity already mentioned—the occurrence of irregular fractures across the planes. In table V are given the supplement angles calculated for all the planes on e^2 and also the angles measured on the several crystals (as before numbered). The angles are in pairs corresponding to 201 and $\bar{2}01$, or $\bar{2}01$ and 201 in each case. (Compare the spherical projection, plate xiii.)

TABLE IV.
Angles measured on $C = i\bar{i}$ (010).

		V.	VI.	VII.	VIII.	IX.	X.	XI.	XII.
r^1	$\frac{4}{7}\bar{2}$ 247			50° 11'	$\left\{ \begin{array}{l} 50^\circ 16' \\ 50 \quad 16 \end{array} \right.$	$\left\{ \begin{array}{l} 50^\circ 17' \\ 50 \quad 17 \end{array} \right.$	50° 12'		50° 24'
r^2	$\frac{4}{5}\bar{2}$ 245	42° 34'	$\left\{ \begin{array}{l} 42^\circ 20' \\ 42 \quad 12 \end{array} \right.$	$\left\{ \begin{array}{l} 42 \quad 30 \\ 42 \quad 23 \end{array} \right.$	42 28	$\left\{ \begin{array}{l} 42 \quad 36 \\ 42 \quad 29 \end{array} \right.$		42° 34'	$\left\{ \begin{array}{l} 42 \quad 38 \\ 42 \quad 36 \end{array} \right.$
r^3	$\frac{4}{3}\bar{2}$ 243			33 26	33 32½	$\left\{ \begin{array}{l} 33 \quad 33 \\ 33 \quad 34 \end{array} \right.$	33 27		33 32
r^4	$4\bar{2}$ 241	25 59	$\left\{ \begin{array}{l} 26 \quad 1 \\ 26 \quad 0 \end{array} \right.$	25 55		25 57	25 56½	26 0	
n^1	$\frac{2}{3}$ 223	$\left\{ \begin{array}{l} 44 \quad 20 \\ 44 \quad 10 \end{array} \right.$	$\left\{ \begin{array}{l} 44 \quad 26 \\ 44 \quad 14 \end{array} \right.$	44 14		44 15			
n^2	2 221	52 57	52 55			52 52			
m^2	$6\frac{3}{2}$ 641			54 31					

Of the planes which occur, according to Scacchi and vom Rath, on type II of humite, I have identified all but m of Scacchi ($\frac{6}{5}\bar{2}$) and $\frac{2}{5}e$ of vom Rath ($\frac{6}{5}\bar{i}$). Of new planes I have found the following, which fall in the old vertical or horizontal zones, and many others to be described later; o ($i\bar{2}=210$), i^2 ($2\bar{i}=021$), ia ($\frac{4}{3}\bar{i}=047$), ib ($\frac{2}{5}\bar{i}=025$), ea ($\frac{2}{5}\bar{i}=205$), ra ($\frac{2}{5}\bar{2}=\bar{4}89$); of these the most interesting is the prism $i\bar{2}$, as hitherto no prism has been found on either the 2d or 3d types.

Hemihedrism—The peculiar hemihedral character of humite has been alluded to, and it is a little striking that the crystals of chondrodite should show so entire a correspondence to it. Taking the same position for the crystals as vom Rath, r^2 and r^4 appear uniformly in the positive (or upper) quadrants, r^1 and r^3 always in the

TABLE V.

Angles measured on $e^2=2-\bar{i}$ (201 and $\bar{2}01$).

			Calculated.	V.	VII.	IX.	I.	X.	VI.
e^1	$\frac{2}{3}\bar{i}$	203	26° 57' 53"	26° 55'			26° 51'		26° 59'
r^1	$\frac{4}{7}\bar{2}$	$\bar{2}47$	{ 59 19 58 87 22 38				59 20	59° 20'	
r^2	$\frac{4}{5}\bar{2}$	245	{ 59 14 0 82 34 28	59 14	59° 33' (approx)	{ 59° 30' 59 30	82 34		
r^3	$\frac{4}{3}\bar{2}$	$\bar{2}43$	{ 60 30 0 76 32 14		76 26	76 15		76 35	XI.
r^4	$4\bar{2}$	241	{ 64 2 9 69 51 3		64 0	63 57			64° 0'
n^1	$\frac{2}{3}$	$\bar{2}23$	{ 44 38 20 70 20 31	44 34					
n^2	2	221	{ 45 45 24 56 42 18	{ 45 40 45 50	45 46 56 42		45 42 56 42		
m	$6\frac{3}{2}$	$\bar{6}41$	{ 37 34 34 42 57 12			{ 42 43 42 32	42 51		

negative (or lower) and n^2 is both + and -, but where occurring alone is generally negative; n^1 is generally, and m^2 always, negative. Of the brachydomes it may be said that they are often holohedral, but this is not always the case. The various figures on the two plates will show the true relations better than words. It is to be said, however, that when the brachydomes are \pm they are still distinguished from each other physically. Thus the + series may be largely developed and rough, destitute of any semblance to polish, when the negative series is as lustrous as the pyramidal planes. When $e^2(2-\bar{i}=201)$ is only once present it is uniformly positive. The macrodomes are always holohedral on humite; here this is sometimes the case, but there is a good deal of irregularity (as will be seen in the figures), and this is conspicuous in figure 7 where i and $i\frac{1}{2}$ occur together and also ia , $i\beta$, and i .

Habit.—With regard to the general habit of the crystals, it is interesting to note the wide variation which is shown. Figures 1, 2, 6, 9, 10, are intended to give some idea of the crystals as drawn symmetrically, and figures 7, 10, 14, 15, 16, 17, 18, 19, , of their actual appearance. As will be seen, the figures are drawn with $C(i\bar{i}, 010)$ in front: this was necessary in order to give a true idea of their real appearance. The prism $i\bar{2}$ is so acute ($49\frac{1}{2}^\circ$) that when directed

toward the eye the projection gives it but little width. As a matter of fact the crystals have C uniformly well developed and are generally attached approximately by the extremity of the brachy-diagonal axis. This having been explained, it will be clear that while fig. 3 is an almost exact reproduction of an actual crystal, fig. 5, by the other method of projection, gives an entirely false idea of its appearance. It is certainly true that the latter method shows the hemihedrism in its true light, but this should not weigh against the other more important consideration.

The crystals from which the partial figures, 7, 10, 16, 17, were drawn were united, along with others quite as diverse, in one small fragment only half an inch in length. It is to be noted that figures 14 and 15 are really more different than would appear at first glance. The crystals drawn in figures 16 and 19 also occurred closely conjoined in the same group; and other examples might be mentioned. One crystal of a very prismatic appearance (when placed in an inverted position) is shown in figure 19.

Presence of minute planes.—The most remarkable feature of the mineral from this locality is yet to be mentioned. I refer to the multitude of minute planes which modify many of the solid angles. One single case will be discussed in detail, as the planes admitted of more than usually exact determination; it serves well to illustrate the subject. A horizontal projection of a portion of the crystal is shown in fig. 14. The crystal itself was small, and unfortunately so imbedded in dolomite that it was for the most part rough and beyond even approximate measurements. The part available showed C ($i\bar{i}$) faultless; also r^3 good; and less satisfactory r^1 , r^2 and r^4 . On the solid angle between C , r^3 and r^2 a large number of minute planes were observed; they were so extremely small (all covering a surface not $\cdot 03$ of an inch in breadth) that any exact measurements seemed at first hopeless. They were sharply defined, however, and brilliant, and when the attempt was made it was found that they gave perfectly distinct though faint reflections. It may be remarked here that measurements in this case were only made possible by the substitution of a cross, cut in tin foil and illuminated very brilliantly by a gas burner, for the ordinary spider lines in the second telescope—a device for which I am indebted to Prof. Schrauf of Vienna.*

The measurements were all made with the greatest care, though, as will be readily understood, the exact adjustment of planes so small

* Ber. Ak. Wien, lxvii, 1873.

was not an easy task. The symbols were calculated from the angles thus obtained. But as even then some doubt remained as to the degree of dependence which was to be placed in them, the measurements were all repeated with the same care as before. The result was perfectly satisfactory, as the variation in no case exceeded the probable error of observation given to each angle when measured for the first time. This variation in most cases did not exceed $\pm 2'$. The following supplement angles were obtained for the more prominent planes: $C \wedge r^3$ gave $33^\circ 33'$ (required $33^\circ 32\frac{1}{4}'$); and $C \wedge r^4$ gave $25^\circ 53'$ (required $25^\circ 58'$); and $r^3 \wedge r^4$ gave $31^\circ 31'$ (required $31^\circ 38'$). The following table contains the angles for each of these minute planes as measured on C , and r^3 and also on i^2 , itself one of this group.

TABLE VI.

			$C = i\bar{i}(010).$		$r^3 = \bar{2}43.$		$i^2 = 201.$	
			Meas.	Calc.	Meas.	Calc.	Meas.	Calc.
i^2	$2\bar{1}$	021	$17^\circ 37'$	$17^\circ 38\frac{1}{2}'$	$23^\circ 58'$	$23^\circ 49'$		
$+x^1$	$\frac{12}{7}\bar{6}$	2·12·7	22 9	21 53	31 4	31 12	$8^\circ 21'$	$8^\circ 37'$
$+x^2$	$\frac{26}{9}\bar{13}$	2·26·9	12 55	12 58	29 32	29 27	6 36	6 35
$+x^3$	$\frac{26}{7}\bar{13}$	2·26·7	10 34	10 30	30 37	30 42	8 50	8 54
$+x^4$	$\frac{34}{7}\bar{17}$	$\frac{17}{2}$ 4·34·7	9 42	9 39	33 41	33 41	12 0	11 53
$-x^6$	$\frac{12}{7}\bar{24}$	$\bar{1}$ ·24·14	20 21	20 28	21 11	21 6	3 25	3 24
$-x^5$	$\frac{13}{12}\bar{13}$	$\bar{5}$ ·13·12	34 32	34 27	7 32	7 15	21 15	21 8
$-x^7$	$\frac{13}{7}\bar{13}$	$\bar{1}$ ·13·7	18 50	19 17	19 40	19 47	4 5	4 3
$-x^8$	$\frac{24}{13}\bar{4}$	$\bar{6}$ ·24·13	22 24	22 31	12 15	11 59	12 22	12 22
$-x^9$	$\frac{5}{2}\bar{15}$	$\bar{2}$ ·15·6	16 5	15 47	18 51	19 7	7 39	7 34
$-x^{10}$	$\frac{13}{7}\bar{13}$	$\bar{4}$ ·13·7	24 22	23 59	9 26	9 47	15 25	15 3
$-y^2$	$\frac{25}{9}\bar{25}$	$\bar{7}$ ·25·9	18 54	19 2	14 45	14 38	14 40	14 52
$-y^1$	$3\bar{8}$	$\bar{9}$ ·24·8	21 55	22 4	13 18	13 14	19 32	19 29
$-y^4$	$9\bar{9}$	$\bar{2}$ 91	11 43	12 13	23 31	23 14		
$-y^3$	$24\bar{24}$	$\bar{9}$ ·24·1	18 45	19 5	25 34	25 27		

The calculated symbols are also given with the angles which belong to them. It will be noticed that $i^2(2\bar{7}=021)$ is itself one of the minute planes of the same character as those surrounding it, and its presence gives a reality to them which they would not otherwise have, and shows what degree of reliance is to be placed on the angles. Cases of a similar character will be noted hereafter.

The symbols* calculated for this series of planes are certainly not simple; and yet a moment's consideration will show that this was exactly what was to be expected. Crowded together so closely, they would be abnormal if occurring on crystals of any species, while this becomes still more true for a mineral like chondrodite. The constantly recurring, common planes have ratios which in any other species would be considered next to impossible: thus, in type II, $1 : \frac{1}{3} : \frac{1}{5} : \frac{1}{7}$; and in type III, $1, \frac{1}{3}, \frac{1}{5}, \frac{1}{7}, \frac{1}{9}, \frac{1}{11}$. It is to be noticed that these are the true ratios of the r series of pyramids, which exist no matter what change be made in the assumed axes. It is not surprising, then, that these secondary planes should themselves have symbols totally at variance with the accepted law of simplicity of the indices. Many cases of planes with what may be called abnormal indices have been described, but frequently they are to be explained as has been done by Brezina in the case of the $\frac{2\bar{2}}{4}\cdot\bar{7}$ ($25\cdot0\cdot24$), which he proved to exist in wiserine, as a tendency to a plane with a simple index ($1\bar{7}=101$), which has resulted in a plane with approximately the given index. The case in hand, however, is quite different.

It will be noticed, however, that, lawless as they appear at first, there is an attempt at system in the symbols given. Thus in the ratio of the brachydiagonal to the vertical axis, we have:

x^1	12 : 7	x^7	13 : 7
x^5	12 : 7	x^{10}	13 : 7
y^2	24 : 13	x^3	26 : 7
y^1	24 : 8	x^2	26 : 9
y^3	24 : 1	x^5	13 : 12

and so on. The ratios for the other axes might be drawn out in the same way, but they are already contained in the symbols given in the table. A little surprise is felt at first that the uniformity in ratio is not greater, that, for instance, x^1 is not $2\cdot13\cdot7$, instead of $2\cdot12\cdot7$; but the measurements are too good and reliable to allow of such a

* In the symbols given in the tables (i. e., those according to Naumann) the dash over the second figure, or fraction, has been omitted (in order to simplify the work of the printer). This has also been done in all the following tables, being made possible by the fact that all these planes, with one or two exceptions, belong to the *macrodiagonal* series.

supposition. It is to be noticed that when ratios of this character are allowed, a slight change in the measured angle will alter entirely the calculated index; the liberty in this respect is not, however, quite so great as it would seem at first sight to be. For example, the ratios $\frac{5}{6}(\frac{3}{6}\frac{5}{6})$ and $\frac{4}{7}(\frac{3}{6}\frac{6}{6})$ approach pretty closely to each other, and it might be a question which was to be accepted as the true ratio of the two axes for a certain plane; and yet if the ratio of one of these axes with the third be unquestionably expressed in sevenths, e. g., $\frac{8}{7}$, then there seems little doubt but that the ratio $\frac{4}{7}$ is to be accepted, for that would give $8\cdot4\cdot7$ or $\frac{8}{7}\cdot 2$, while the other supposition would give $35\cdot72\cdot63$ or $\frac{8}{9}\cdot\frac{7}{3}\frac{2}{5}$. This principle has been accepted in obtaining all the indices given in the following tables.

A remarkable fact connected with these planes,—in fact, implied in what has already been said,—is that there is so little tendency among them to lie in zones. For example, x^1 , x^9 , y^2 and y^1 lie very nearly in a zone with each other and i^2 , and yet the reflections in the goniometer deny that this is exactly true, while no satisfactory indices can be obtained on this supposition. (x^1 , i^2 and y^2 are, however, in a zone.)

In regard to these planes two points are to be noticed. In the first place, the question suggests itself whether, if referred to a common fundamental form (see above), or to that of either of the other types, the relations of the planes would be at all more simple. This is answered in the negative, as will be seen to be necessary if the trial is made, and also evidently because planes whose normals make angles of a few degrees only with one another can never bear simple relations to each other, no matter what axes be assumed.

In the second place, it might be urged that such ratios as those above given being accepted, there is no reason why we should attempt to express the relations of the prominent planes—those of humite, type II, for example, with simple numbers (see above, page 7). But, as has just been stated, the attempt to refer these planes themselves to other axes leads to disastrous results, while further, as has been shown, these planes are truly secondary and subordinate and bear no relations to other types of the species.

This case has been dwelt upon at considerable length, because it was believed that theoretically the existence of such planes was a matter of some interest and importance, and because this single crystal offered opportunities for exact determination which did not exist to the same degree in any other case. Almost all of the twenty and more smaller crystals examined showed some of these secondary planes. In some cases, however, there was a tendency to rounded edges

without the formation of distinct planes, as is so common in many species; and then nothing of course could be done.

The following tables, VII, VIII, IX, X, include the measured angles, with the symbols obtained, and the corresponding angles belonging to them, for a considerable number of these minute planes. Those occurring on each individual crystal are arranged together, being expressed by the same letter, and where the crystal has been figured, this is also indicated. Upwards of one-hundred of these planes were measured, and an attempt was made in every case to obtain a satisfactory index. It was concluded, however, to discard the

TABLE VII.

			$C = i\bar{i} (010).$		$+e^2 = 2\bar{i} (2\bar{0}1).$		
			Meas.	Calc.	Meas.	Calc.	
Crystal VII (figure 16).	-a	8-4	281	14° 6'	13° 41'	79° 2'	79° 16'
	-v ₁	$\frac{11}{4} - \frac{11}{5}$	$\bar{5} \cdot 11 \cdot 4$	25 14	25 34	73 29	73 7
	-v ₂	$\frac{8}{3} - \frac{16}{11}$	$\bar{11} \cdot 16 \cdot 6$	33 50	34 5	64 40	64 28
	-v ₃	$\frac{13}{4} - \frac{5}{4}$	$\bar{5} 2 \cdot 65 \cdot 20$	37 7	37 19	60 0	59 49
	-s ₁	$\frac{3}{2} - \frac{15}{16}$	$\bar{1} 6 \cdot 15 \cdot 10$	46 43	46 56	57 38	57 22
	-t ₁	$\frac{5}{7} - \frac{5}{3}$	$\bar{3} 57$	46 4	46 21	81 35	80 49
	-t ₂	$\frac{9}{14} - \frac{9}{4}$	$\bar{4} 9 \cdot 14$	46 57	46 58	87 7	87 31
	-s ₂	$\frac{7}{9} - \frac{7}{8}$	$\bar{8} 79$	53 6	53 7	64 9	64 7
	-t ₃	$\frac{1}{2} - \frac{3}{2}$	$\bar{2} 36$	54 31	54 42	85 0	84 33
	+t ₄	$\frac{7}{11} - \frac{7}{2}$	$\bar{2} 7 \cdot 11$	45 47	45 57	92 46	93 6
Crystal X (figure 15).	+t ₅	$\frac{9}{10} - \frac{9}{2}$	$\bar{2} 9 \cdot 10$	36 16	36 21	91 20	91 44
	-n ²	2	$\bar{2} 21$	44 13	44 $14\frac{1}{2}$	56 45	56 42
	-ζ	$\frac{9}{5} - \frac{9}{7}$	$\bar{7} 95$	38 45	38 37	63 44	63 58
	-θ	$\frac{15}{4} - \frac{3}{4}$	$\bar{2} 0 \cdot 15 \cdot 4$	50 58	51 6	46 13	46 4
	i ¹	1- $\bar{2}$	011	32 34	32 27	79 55	79 54
	+p ¹	$\frac{5}{7} - 5$	157	42 0	42 17	109 36	110 7
	-p ²	$\frac{20}{17} - 5$	$\bar{4} 20 \cdot 17$	29 41	29 $44\frac{1}{2}$	90 0	89 58

larger portion as untrustworthy, retaining those which had given the best angles. It is not pretended that the symbols deduced are, even in the majority of cases, correct beyond question; for the angles, while perfectly reliable in some cases, are in others somewhat uncertain, and for reasons already explained this throws still greater doubt over the indices which calculation may produce.

TABLE VIII.

			$C = i\bar{i} (010).$		$A = O (001).$	
			Meas.	Calc.	Meas.	Calc.
Crystal III (figure 17).	$-z^1$	$\frac{7}{6} - \frac{7}{5}$	$\bar{6}75$	40° 0'	39° 52'	65° 25' 65° 51'
	$-z^2$	$\frac{11}{6} - \frac{11}{5}$	$\bar{5} \cdot 11 \cdot 6$	28 44	28 32	72 13 72 15
	$-z^3$	$\frac{15}{4} - \frac{15}{14}$	$\bar{14} \cdot 15 \cdot 4$	41 0	41 14	82 45 82 40
	$-z^4$	$\frac{17}{3} - \frac{17}{4}$	$\bar{4} \cdot 17 \cdot 3$	14 6	13 43	83 55 83 44
	$-z^5$	$\frac{10}{11} - 5$	$\bar{2} \cdot 10 \cdot 11$	35 42	35 54	55 44 55 28
	$-m$	$6 - \frac{3}{2}$	$\bar{6}41$	54 18	54 17	84 37 84 40
Crystal IV (figure 10).	$-\beta$	$\frac{5}{9} - \frac{5}{2}$	$\bar{2}59$	50 32	50 16	42 45 42 56
	$-\xi_1$	$\frac{7}{9} - \frac{7}{5}$	$\bar{5}79$	46 45	46 10	55 48 55 29
	$-\xi_2$	$\frac{8}{9} - \frac{4}{3}$	$\bar{6}89$	45 18	44 51	59 59 30

The group of planes clustered about i^2 has already been described. It is interesting to note that in two instances analogous groups were observed, of which $i(1\bar{i} = 011)$ was a member (see fig. 15), and in two other cases the common and prominent planes n^2 (fig. 15) and m^2 (fig. 17) were found in the same relations. The angles obtained for these planes show conclusively the degree of dependence to be placed on those measured for the other planes. It will be remembered that in all cases these planes were exceedingly minute.

The fact already mentioned, that all of these planes belong to the macrodiagonal series, may possibly be explained in part by the fact that it is uniformly that portion of the crystal (i. e., near $C, i\bar{i}$) which is exposed and well developed. There still remains the fact, which will be noticed by a glance at the spherical projection (plate xiii) and which does not allow of an analogous explanation, that

TABLE IX.

			$e^2 = 2\bar{i}$ (201).				$A = O$ (001).			
			Meas.		Calc.		Meas.		Calc.	
Crystal II (figure 7).	i^2	$\frac{1}{2} - \bar{i}$	012		51° 50'	51° 50'				
	ia	$\frac{4}{7} - \bar{i}$	047		47½-48	48 4				
	$i\beta$	$\frac{2}{5} - \bar{i}$	025		58 18	57 51				
	$-ra$	$\frac{8}{9} - \bar{2}$	489		40 6	40 24	56° 42'	56° 59'		
	$-\tau$	$\frac{1}{2} - \frac{5}{4}$	4·5·10		55 45	55 47	45 7	44 18		
	$-q$	$\frac{11}{10} - \frac{11}{7}$	7·11·10		39 33	39 32	63 23	63 26		
	$+\pi$	$\frac{1}{2} - \frac{5}{3}$	3·5·10		54 2	54 13	41 58	41 55		
Crystal I.	$-ny$	$\frac{5}{2}$	552		43 40	43 42	79 33	79 22		
	$-nx$	30	30·30·1		42 40	42 39	89 7	89 6		

these minute planes almost always lie in the negative (lower) quadrants; this has necessitated the drawing of some of the crystals in an inverted position. The figures show, in addition to planes mentioned in the tables, some others for which no symbols were obtained.

TABLE X.

			$C = i - \bar{i}$ (010).				$-e^1 = \frac{2}{3} - \bar{i}$ (203).			
			32° 20'		32° 27'		67° 15'		67° 16'	
Crystal VIII.	i	$1 - \bar{i}$	011							
	$-w_1$	$1 - \frac{15}{4}$	4·15·15		34 41	34 18	58 21	58 43		
	$-w_2$	$\frac{17}{10} - \frac{17}{6}$	6·17·10		26 33	26 25	63 29	63 37		
	$-w_3$	$\frac{3}{4} - 3$	134		41 34	42 3	52 45	52 18		
	$-\tau$	$\frac{1}{2} - \frac{5}{4}$	4·5·10		55 27	55 47	36 40	36 37		
	$-\delta$	$\frac{8}{9} - 4$	289		37 10	36 57	56 27	57 21		
	$-u_1$	$\frac{15}{13} - 5$	3·15·13		29 22	30 10	63 45	63 2		
$-u_2$	$\frac{12}{11} - 6$	2·12·11		31 8	31 6	62 55	63 13			

Crystals of simpler habit.—The crystals thus far described have all been of a more or less complicated character. But allusion has been made to some very simple crystals, which also deserve description. The distinction is in most cases probably unimportant, though believed to be of interest at first. The simple crystals are uniformly large; they are so generally altered, and appear so differently from their small brilliant relatives, that it was supposed that they differed from them at least in the purity of the original material, if not more essentially. One brilliant exception, however, to the general rule in regard to the altered condition of these crystals, in the form of an isolated crystal of faultless luster, and deep red color, about $\frac{3}{4}$ of an inch in length, as well as numerous examples of transition products between the altered and unaltered material, made it probable that all the crystals in question were originally of the same character. Some examples are given in figures 3, 4 and 5. The angles could be measured approximately only with a hand goniometer, but there is no question that they, as well as others, belong to type II. On one of them $\pm r$ was observed.

In what has been said exception must be made in regard to the large coarse crystals, and crystalline masses, mentioned in the early part of this article, and which are made up of a more or less heterogeneous mass of chondrodite, magnetite and sometimes ripidolite; some, at least, of these last, belong to type I. (See p. 25.)

Twins.—The humite crystals of Vesuvius, as well as the Swedish chondrodite, has been shown by vom Rath to possess so great a tendency to twinning that it is a little remarkable that the contrary should be true of the mineral from Brewster. Figs. 20, 21, show the only method of twinning which has been found, as well as the only distinct twin-crystal. The axis of revolution here is the vertical axis of the crystal, and the composition-face the basal plane *A*. Unfortunately the crystal in question was quite imperfect, and all that was available is shown in the figure. The plane *B* ($i\bar{z}=\bar{1}00$) gave no reflections, so that all measurements were made on e^2 (that is 201 and 20 $\bar{1}$); in this case these planes were similar in luster as a result of the twinning. A revolution of the kind mentioned (in a perfectly symmetrical crystal) would, so far as this half of the crystal goes, have the effect only of making it holohedral, giving no re-entrant angles; but, in case of any irregularity, it might give, as here, a re-entrant angle in the planes which are hemihedral in their occurrence.

The measurement of the re-entrant angle here observed gave for $m^2 \wedge \bar{m}^2$, $10^\circ 38'$ and $10^\circ 40'$; required $10^\circ 39'$. The other angles measured on the same crystal are given in the following table.

TABLE XI.

		$e^2 = 2\bar{1}$ (201).		$e^2 = 2\bar{1}$ (201).	
		Meas.	Calc.	Meas.	Calc.
\bar{m}^2	641	37° 39'	37° 34½'	42° 54'	42° 57'
m^2	641	42 57	42 57	37 28	37 34½
\bar{m}^2	641	37 36	37 34½	42 55	42 57
m^2	641	42 57	42 57	37 33	37 34½
n^2	221	45 45	45 45	56 42	56 42
n^2	221	56 44	56 42	45 45	45 45
n^2	221	45 34	45 45	56 41	56 42
n^2	221	56 15	56 42	45 34	45 45
n^1	223	44 45	44 38	70 16	70 20
	223	44 48	44 38		
r^2	245	59 13	59 14	82 34	82 34½
r^1	241	64 3	64 2		
o	210	33 52	33 50½	33 48	33 50½
ϕ	60·38·1	38 14	38 17	39 24	39 27½
"	"	641		641	
		4 31	4 25	6 31	6 36½

Besides the interest of the twin, a noteworthy fact in the crystal is the occurrence of the prism ($i\bar{2} = 210$) the first time that any one of the vertical prisms has been observed in the 2d type of either humite or chondrodite. It lies in both the zones $e^2(201)$, $n^2(2\bar{2}1)$, and $e^2(22\bar{1})$, $n^2(2\bar{2}1)$ which answers sufficiently to determine what it is; and the result thus obtained is fully sustained by good measurements, as will be seen in the table. This plane is distinctly present on one side only of B ; on the other side its presence is barely indicated. Its place here is taken by a well polished and conspicuous plane ϕ , which is another striking instance of the peculiar nature of this species; its position is indicated in the sphere of projection,* and the angles on e^2 are given in the table. The index was calculated for each pair of measurements 38° 14', 39° 24 and 4° 31, 6° 31 (see in the table), and the results obtained were identical. From the first pair of measured angles $\phi \wedge B$ was found to be 34° 30½' and from the second $\phi \wedge B = 34° 31'$ (required 34° 32' 38"). The index obtained was 60· $\frac{38}{9}$ or 60·38·1, and abnormal as it certainly is, it expresses the exact posi-

* In the projection ϕ is placed incorrectly in the negative quadrants; it should be in the positive with $+n$, $+n^2$, $+r$, etc.

tion of the plane. It will be noticed that the four planes upon which the inclination of Φ was measured are so situated, that any variation from the true position in the index would show itself very conspicuously. The fundamental angles for Φ are as follows :

$$\Phi \wedge A(001) = 88^{\circ} 51' 41''$$

$$\Phi \wedge B(100) = 34 \quad 32 \quad 38$$

$$\Phi \wedge C(010) = 55 \quad 28 \quad 49$$

In figure 16 a crystal, or portion of one, is exhibited which is holohedral. It is irregular in this respect, however, that r^2 forms a re-entrant angle with r^3 . This is not a point of special importance, as an irregularity such as this is often observed; but, in view of the crystal which has just been described, it is possible that here also there has been a semi-revolution parallel to the basal plane. A more interesting crystal, already once alluded to, is shown in fig. 4. It is conspicuously hemimorphic, as far as the form goes. It is large, and admits only of approximate measurements, but there is no doubt but that the planes as given have been determined correctly. In view of the fact that a revolution parallel to O would produce just the effect we have here, and as such a twinning law has been shown to exist in another conspicuous case, it is altogether probable that this forms an ample explanation of what is observed. Another exactly parallel case is noted under the description of two crystals of the 3d type. The above described crystal was somewhat altered, and so far imbedded in the matrix that any experiments as to its pyro-electrical character were out of the question.

Chemical composition.—I am glad to be able to add here the results of a chemical examination of the chondrodite of the 2d type from this locality, by Mr. G. W. Hawes of the Sheffield Scientific School. It obviously increases much the value of this memoir. An analysis by Mr. Breidenbaugh has already been quoted (p. 3).

The material analyzed by Mr. Hawes consisted of fragments of crystals of the 2d type, selected with great care to avoid the presence of any altered material. It had a deep garnet-red color and a brilliant vitreous luster. Its specific gravity as determined by Mr. Hawes was 3.22.

	Analysis I.	Analysis II.
Silica	34.10	34.05
Magnesia	53.17	53.72
Ferrous oxide	7.17	7.28
Alumina	.48	.41
Fluorine	4.14	3.88
	<hr/>	<hr/>
	99.06	99.34

Following the view of Rammelsberg, that the higher values of each constituent are nearest to the truth, Mr. Hawes's analysis becomes as follows. For comparison the results obtained by vom Rath for 2d type crystals from Vesuvius and from Sweden are added.

	CHONDRODITE.		HUMITE.
	Brewster, N. Y., <i>Hawes.</i>	Sweden, <i>v. Rath.</i>	Vesuvius, <i>v. Rath.</i>
Silica	34.10	33.96	34.02
Magnesia	53.72	53.51	59.23
Ferrous oxide	7.28	6.83	1.78
Alumina	0.48	0.72	0.99
Fluorine	4.14	4.24	2.74
	<hr/>	<hr/>	<hr/>
	99.72	99.26	98.76
Silicon	15.91	15.85	15.88
Magnesium	32.23	32.11	35.54
Iron	5.66	5.31	1.38
Aluminum	0.26	0.38	0.53
Fluorine	4.14	4.24	2.74
Oxygen	39.78	39.58	41.54
	<hr/>	<hr/>	<hr/>
	97.98	97.47	97.61

Transforming the iron into an equivalent of magnesium, as also the alumina ($2Al=3Mg$), Mr. Hawes obtains further:

Silicon 15.91, Magnesium 35.00, Fluorine 4.14, Oxygen 39.78.

From these values a formula is deduced, which is essentially that of the Swedish mineral according to *v. Rath*, $20(Mg_5Si_2O_9)+Mg_5Si_2F_{18}$. The close correspondence between the three analyses in the above table is certainly very remarkable. It would have been extremely interesting to have added an analysis also of crystals of the 1st and 3d types; but, as will be apparent from what follows, the material was not to be obtained.

In completing the description of this variety of the mineral, in general it may be repeated that it occurs usually in narrow veins, and when free from alteration has uniformly a deep *garnet-red* color. A *cleavage* such as exists in humite (parallel to the basal plane) and has been observed by *Koksharov* on chondrodite from Pargas, could in no case be discovered. The fracture is always conchoidal, and the only thing that suggested cleavage was the laminated structure of the massive specimens described, and the fractures parallel to e^2 which were quite conspicuous on two or three specimens. The material in hand was not well adapted for optical determinations;

but some interesting results have been obtained which are described in the closing pages of this paper.

2. *Description of Crystals of Type III.*

Since it was shown by Kokscharow that the Pargas chondrodite was identical with the second type of humite, it was natural to expect that further investigation would prove the existence also of the two remaining types. Up to the present time that expectation has been unfulfilled, and it has been reserved for the Brewster locality to give this confirmation of Scacchi's interesting discovery. The crystals of the 3d type are exceedingly rare, three or four specimens being all that have thus far been found, and from these only two individual crystals could be obtained which allowed of measurement. Fortunately these two crystals are very satisfactory, being small and brilliant, and establish the fact as well as a hundred could do. Figures 11 and 12 show one of the crystals, and figure 13 the other. The appearance of the first crystal is best shown in the second of these figures. As will be seen, the planes are the same as in humite, and they are for the most part hemihedral and situated in the same way; i. e., they include $+\rho^4, +\rho^6, +\rho^8$, and $-\rho^3, -\rho^5, -\rho^7$, and $-\nu^2, -\nu^2, -\nu^3, -\nu^4$ as also i^2, i^3, i^4 . In the $n(\nu)$ series the planes are distinct in the negative half, but the crystal is incomplete, and it is a little uncertain whether the $+$ series should not in part be added in the symmetrical drawing fig. 11; on humite, III, these planes are both positive and negative. No brachydomes are visible, the edge being rounded and rough. m^2 of Scacchi may also be present, but that is a little doubtful.

The second crystal is of very different form, and while the first was affixed to the rock so that only one-half was developed, this one was imbedded in brucite, and entirely free in it. It was perfectly formed on all sides, being almost as perfect as the projected drawing, with the exception, however, of the acute (brachydiagonal) edge, which was mostly broken. When only the upper part of the crystal is considered, it will be seen that the hemihedrism is like that in the other case, except that ρ^6 is holohedral. For macrodomes there are $i^1(\frac{2}{3}\bar{1}=023)$, $i^2(1\bar{7}=011)$, $i^3(2\bar{7}=021)$, $i^4(4\bar{7}=041)$; the last has not been observed on humite. On measuring the planes below it was found that they were not distributed as was expected in accordance with the monoclinic character of the crystal; instead, either extremity of the brachydiagonal axis was differently developed. This is clear in the figure, it being but a more complicated repetition of what was observed in one

of the very simple crystals of the 2d type (see fig. 4 and page 18). There are present also at one extremity $\pm \epsilon' (\frac{1}{4}\bar{1}=407)$, though the plane could only be approximately measured. This is probably also to be explained as having resulted from a revolution parallel to the basal plane. The crystal was very small and not at all adapted to experiments having in view the discovery of any proper hemimorphic development. The angles measured on both these crystals are contained in the following tables.

TABLE XII.

		Chondrodite.			Humite.				
		Calc.	$A = O(001)$. Meas (XX). Meas (XXI)		v. Rath. Calculated.				
ϵ	$\frac{1}{7}\bar{1}$	407	143°	20'	9''		143°	11'	29''
ρ^1	$4\bar{1}$	041	100	1	7				
ρ^3	$2\bar{1}$	021	109	27	35		109	27	54
ρ^2	$1\bar{1}$	011	125	14	49		125	15	18
ρ^1	$\frac{2}{3}\bar{1}$	023	136	40	4		136	40	34
ρ^3	$\frac{8}{11}\bar{2}$	2811	131	25	57	131° 46'	131° 24'	131	24 49
ρ^4	$\frac{8}{9}\bar{2}$	289	125	50	6	{ 125 37 125 47	125 48	125 49	0
ρ^5	$\frac{8}{7}\bar{2}$	287	119	19	18	{ 119 35 119 15	118 36	119 18	19
ρ^6	$\frac{8}{5}\bar{2}$	285	111	51	38	{ 111 44 111 49	112 0	111 50	50
ρ^7	$\frac{8}{3}\bar{2}$	283	103	32	4	103 41	103 38	103 31	33
ρ^8	$8\bar{2}$	281	94	35	15	{ 94 31 94 13	94 48	94 35	4
ν^1	$\frac{4}{7}$	447	132	17	48		132 16	132 16	43
ν^2	$\frac{4}{5}$	445	123	1	8		122 32	123 0	8
ν^3	$\frac{4}{3}$	443	111	18	7		111 5	111 17	23
ν^4	4	441	97	24	20		97 29	97 24	3

Unfortunately the inclination to *C* on no one of the pyramidal planes could be measured with perfect accuracy; the measurements are good, yet not entirely trustworthy. These planes, though brilliant, are uniformly fractured in the manner already explained, and this made all

TABLE XIII.

Chondrodite.

			<i>C</i> = $i\bar{i}$ (010).			
			Calc.	Meas (XX).		Meas (XXI)
i^4	$4\bar{i}$	041	169°	58'	53"	170° 2'
i^3	$2\bar{i}$	021	160	32	25	160 35
i^2	$1\bar{i}$	011	144	45	11	144 47
i^1	$\frac{2}{3}\bar{i}$	023	133	19	56	133 (ap.)
ρ^3	$\frac{8}{11}\bar{2}$	$\bar{2}811$	132	56	12	{ 133 13 132 38 } 132° 46'
ρ^4	$\frac{8}{9}\bar{2}$	289	137	26	13	{ 137 36 137 27 } 137 21
ρ^5	$\frac{8}{7}\bar{2}$	$\bar{2}87$	142	22	50	{ 142 35 142 20 } 142 38
ρ^6	$\frac{8}{5}\bar{2}$	285	147	28	24	147 26
ρ^7	$\frac{8}{3}\bar{2}$	$\bar{2}83$	152	1	44	152 2
ρ^8	$8\bar{2}$	281	154	53	20	155 0 154 58
ν^1	$\frac{4}{7}$	447	123	31	12	
ν^2	$\frac{4}{5}$	445	128	5	52	
ν^3	$\frac{4}{3}$	443	133	16	51	133 57
ν^4	4	441	136	51	9	136 42

these angles a little uncertain. The macrodomes in one case gave good measurements; and making use of the best of them, $C \wedge i^2(1\bar{i}) = 144^\circ 47'$, and also the same prism as in type II, after the analogy of humite, the angles were calculated throughout. It was found, however, that wherever trustworthy they corresponded so closely to what was required to satisfy the ratio of 10 : 9 for the vertical axes (asserted as approximately true by Scacchi, i. e., 5 : 9 for his axes, and finally proved rigidly by vom Rath), that the calculations were made on this supposition. The calculated angles, as now given, consequently have as their basis the prismatic angle $I \wedge I = 94^\circ 44' 14''$ and the macrodome angle $C \wedge i^2 = 144^\circ 45' 11''$.

The corresponding parameters are :

$$a \text{ (vert.)} = 1.41512; \quad b = 1; \quad c = 1.08630.$$

Very little further can be said in regard to the crystals of the 3d type. Those observed had a somewhat different color from those of type II; that is, the color was more yellowish, less of a pure garnet-red—though this may be accidental. No analysis was possible of course; and even the specific gravity was out of the question also, for the one loose crystal, in addition to its small size, had imbedded in it a still smaller crystal of ripidolite, making any gravity determinations obviously unreliable.

The method of occurrence was much like that of the brilliant crystals of the second type; that is, they were found implanted on the massive rock adjoining small veins. The associated minerals were magnetite, ripidolite in clear transparent crystals, and, probably as a later formation, brucite.

3. *Description of Crystals of Type I.*

The occurrence of large, coarse crystals of quite impure chondrodite, imbedded in the massive material, has already been described; these belong, at least in part, to the *first* of Seacchi's types. As has been remarked, the crystals of this character do not often admit of exact determination, but in two cases they were so good as to allow of their crystallographic relations being accurately made out. The accompanying wood-cuts, figures 22 and 23, give faithful representations of their appearance and size.




Fig. 22.




Fig. 23.

It will be seen that they are both quite imperfect, and it was on this account that no attempt was made to make a symmetrical drawing of either of them. In each case the crystals were so distorted as to give a sharp edge between the diagonally situated pyramidal planes; this furnished an opportunity for reliable measurements. The planes

were, of course, destitute of all luster, but they were mostly smooth and large enough to allow of the convenient use of the hand goniometer.

R^2 on R^2 (behind) gave measurements varying, in a series of trials, from 78° to 79° , required $79^\circ 4'$.

R^3 on R^3 (behind) gave $62\frac{1}{2}^\circ$, required $63^\circ 1'$.

R^2 on R^3 (behind) gave 72 , required $71 17\frac{1}{2}'$.

R^3 on R^2 (behind) gave 72 , required $71 17\frac{1}{2}'$.

These angles on both crystals were identical within the allowed error of observation (say $30'$). The above are the best angles afforded by any of the planes.

It is entirely impossible to refer these angles to any of the forms of the second type. When compared with the third type, it is seen that on making the supposition that R^2 and R^3 (front) are ρ^3 and ρ^4 respectively, and R^2 and R^3 (behind) are ρ^4 and ρ^5 , we obtain for:

$\rho^3 \wedge \rho^4$ (behind) = $77^\circ 12'$; $\rho^4 \wedge \rho^5$ (behind) = $65^\circ 6'$;

$\rho^3 \wedge \rho^5$ (behind) = $70^\circ 32'$; $\rho^4 \wedge \rho^4$ (behind) = $71^\circ 37'$.

It will be seen, by comparing these with the previously given angles, that the measured angles correspond much better with the first type, and my confidence in them is so great that this would alone be regarded as sufficient to establish the point; and that, without reference to the fact that the supposed method of occurrence of the third type planes is contrary to all the laws of the species.

The decisive proof is derived from the fact that both crystals are certainly holohedral, the planes on both sides being similar with the exception of R^1 , and there is nothing of the obliquity which is observed in the hemihedral forms.

The measured angles of C on R^5 , right and left, were identical, though not obtainable with exactness; the measurements gave $152\frac{1}{2}^\circ$ – 154° : this is also true for C on R^3 , right and left, = $140\frac{1}{2}^\circ$ – $142\frac{1}{2}^\circ$.

In the first crystal $J^1 = \frac{3}{5} - i$ {035} occurs, and in the other J^2 ($1\bar{7} = 011$). The occurrence of C is also to be noticed, as it is rare on humite; in fig. 23, the oscillatory combination of R^4 and R^5 will be also observed.

The following table includes the most important angles for the occurring planes, calculated from the fundamental form of the second type on the assumption that the lateral axes are equal, and the vertical axes have the ratio of 14 : 15. The measured angles are also added though only approximate; in the form given they were obtained immediately from the measurements over the top of the crystals (see above).

TABLE XIV.

			Chondrodite.			Humite.						
			$C = i\bar{i} (010)$. Calculated.			$A = O (001)$. Meas. Cale.			v. Rath. Calculated.			
J ²	$1\bar{i}$	011	145° 43' 44"			124° 16' 16"			124° 16' 45"			
J ¹	$\frac{3}{5}\bar{i}$	035							138 38 38			
R ¹	$\frac{3}{5}\bar{2}$	3·6·10	129	12	57	135°	135	53	35	135	52	23
R ²	$\frac{3}{4}\bar{2}$	368	134	28	38	129 $\frac{1}{4}$	129	32	3	129	30	52
R ³	$1\bar{2}$	122	140	30	10	121 $\frac{1}{4}$	121	45	28	121	44	23
R ⁴	$\frac{3}{2}\bar{2}$	364	147	6	34		112	25	28	112	24	37
R ⁵	$3\bar{2}$	362	152	49	49	101	101	39	30	101 ¹	39	2

The two crystals described are the only ones which could be positively identified. It is very probable, however, that of those found others also belong here, as they have much the same appearance and habit. These crystals are all considerably altered, being generally soft enough to be cut with a knife, and for this reason a chemical analysis would be of little value. The color of the crystals is gray to grayish-yellow, and the material of which they are composed is never pure, and often quite heterogeneous. In this respect they recall the brown crystals described by v. Rath as occurring at Nya-Kopparberg.

Whether brilliant crystals of the first type exist, as they do of the other types, must be left for the present undecided.

4. *On the Optical Properties of Chondrodite.*

In the preceding pages the question of the orthorhombic or clinorhombic crystallization of the chondrodite has not been discussed. In fact, nothing was detected by the measurements sustaining any other conclusion than that of Scacchi and vom Rath, that the crystals were fundamentally orthometric. Still the hemihedral character of the second and third types seem to point to a clinometric form, and this is apparently supported by the optical characters obtained. The material available for optical investigations was very scanty, and, with the exception of one crystal, poorly adapted for the purpose.

The crystal referred to was, properly, but the fragment of what was originally a specimen of considerable size and beauty;

when unbroken it must have been nearly an inch in length. In the condition in which it was found it showed only the brachydomes e^1 and e^2 , with the pyramids n^1 , n^2 , and m^2 ; it had the deep garnet-red color of crystals of the second type, and with the exception of the universally present fractures was perfectly clear and transparent.

On the optical properties of the mineral in question, we have, as far as I have been able to find, no information except what has been given by DesCloizeaux, *Manuel de Mineralogie*, 1862, p. 141. He says: "Double refraction energetic; positive bisectrix normal to g^1 ; optic-axes situated in the plane parallel to the base; divergence in oil for red and yellow rays, $82^\circ 14'$. Dispersion almost nothing; $\rho < v$?"

Guided by the above, a section was cut from the crystal described, which was parallel to C , i. e., perpendicular to the brachydomes present. The examination of this section showed: 1st, that the acute bisectrix is normal to C ($i\bar{7}, 010$); 2d, that this bisectrix is positive; 3d, that the optic-axial angle is large, the axes being seen only when oil is used; but 4th, that *the axes do not lie in the basal plane*, but in a plane making an angle of about 154° with it. This last point was so unexpected and anomalous that every effort was made to explain the measurements in some other way, but with no success. The planes on the crystal had been carefully measured, before the slicing, and the angles agreed perfectly with those of type II for the planes mentioned, so that it was impossible to assume that the crystal had not been correctly put into position. By means of a stauroscope, made by Fuess in Berlin, after the excellent pattern of Groth,* the position of the two axes of polarization, as referred to e^1 , and also to e^2 in plane C , were carefully determined. The measurements were repeated twenty times, the error arising from an imperfect adjustment of the Nicols being eliminated in the usual manner. The result was as follows:

Supplement angle made by the plane of the axes—

with e^1 ($\frac{2}{3}\bar{i}=203$), $18^\circ 9'$; hence with the basal plane, C , $25^\circ 50'$.

with e^2 ($2\bar{i}=201$), $45^\circ 9'$; " " " " $25^\circ 46'$.

In order to confirm these results, other crystals were sought, which would admit of like determinations. None could be found which would serve for measuring the axial angle; but two small ones, on which the plane C was naturally developed, proved to be clear enough to allow of measurements with the stauroscope. The first

*See Pogg. Ann., cxliv, 34, 1870.

alone gave accurate results; on it the angle of the same plane with e^a ($i\gamma=205$) was determined with equal care. The results were:

$4^\circ 55'$ for the angle with e^a ; and hence $25^\circ 59'$ with C .

The agreement with the angles given above is as close as could be desired. In the other case, the rather rare plane B ($i\gamma=100$) was present; the crystal was minute, however, and the determination only approximate. It was found that the normal to the axial plane made with B an angle of $65^\circ-70^\circ$, and hence with the normal to the basal plane $20^\circ-25^\circ$.

With so ample confirmation the point made cannot be even questioned, and it remains to reconcile it with the crystallographic properties of the species. It will be seen at once that the position of the optic axes is totally at variance with the accepted orthorhombic character of the crystals; but it conforms to the rule for monoclinic crystals, as one axis of polarization is normal to the plane of symmetry C , and the others lie in it, or in other words, the optic-axes lie in a plane perpendicular to the axis of symmetry. The angles measured and calculated, given in the various tables, show that the variation from the rectangular type, if it really exist, must be very slight, as the agreement between the angles measured and those calculated on the assumed prismatic basis is very close—it being remarked that some considerable variations in the angles given in the tables are amply explained by the imperfection of the crystals. Note the angles measured for $m^2 \wedge \overline{m}^2$ on the twin crystal described on page 18. It was not to be expected that the variation in the optical character of the crystals would be so decided in view of the slight divergence which is possible in the crystalline form. I reserve for the future the careful revision of the angles of this species, when I shall hope to be able to command a more abundant supply of satisfactory material. It may be added that the hemihedral character of the second and third types of humite long ago suggested the idea that they were oblique in form; but all the crystallographic investigations thus far have seemed to deny this. In the Mineralogy of Brook and Miller, the form is made oblique, but this seems to be due to a misunderstanding of the planes occurring on the crystals.

It would have been interesting to extend these observations to the two remaining types, but the material did not allow of it. It was also desired to investigate the same subject for humite, but, though some good specimens are to be found in the Yale cabinet, there were no satisfactory crystals to be had, and the matter is left for others, who have a larger choice of specimens. The axes as already mentioned do not appear distinctly except in oil; in the first-mentioned

section they admitted of good measurements. The mean of thirty determinations of the angle for red rays gave—

$$2H\alpha = 88^{\circ} 48' : \text{the extremes being } 88^{\circ} 36' \text{ and } 89^{\circ} 0'.$$

With a yellow light (sodium) the angle was essentially the same, but the mean was 10' or 15' smaller, which would indicate that the dispersion is $\rho > v$, but the matter cannot be considered to be beyond doubt. No other dispersion was observed, that is, none parallel or perpendicular to the plane of polarization.

The index of refraction of the oil employed, as determined by Professor Wright and myself, was 1.466.

In conclusion, I have to express my very great obligations to Prof. Allen for his kindness in giving me free use of all the specimens in his valuable cabinet. Both of the crystals of the third type, as well as several others mentioned, came from his collection; in fact it was Prof. Allen who first made known the special interest connected with the locality. To Mr. Cosgriff, the superintendent of the Tilly-Foster Iron Mine, I am also much indebted for his uniform kindness and courtesy to me at the several occasions when I have visited the mine; as also for the gift of several fine specimens.

IV.—ON THE TRANSCENDENTAL CURVES $\sin y \sin my = a \sin x \sin nx + b$.
 With Plates XIV—XXXVII. BY H. A. NEWTON AND A. W. PHILLIPS.

1. ALGEBRAIC curves have been studied hitherto more than transcendental. A few of the latter have been given in the text books, but attempts to classify the numerous varieties of transcendental curves have been rare.

From the form of a transcendental curve it is not easy to state an equation that can represent it. The simpler inverse problem of describing the curve from the equation is naturally the first to be undertaken. The forms that result may, when compared, suggest the solution of the direct problem. We have thought it worth while, therefore, to select for study a single one of the numberless transcendental equations, and to exhibit a few of the very many plane curves which that one equation furnishes. The equation selected is,

$$\sin y \sin my = a \sin x \sin nx + b, \quad (1)$$

in which there are four arbitrary constants a , b , m , and n , with two coördinates, x and y .

2. We assume that m and n are each less than unity. If either, for example m , is greater than unity, we may change the unit for y in the ratio of $1 : m$ by writing $y' = my$. The first member of Eq. (1) then becomes $\sin y' \sin \frac{1}{m} y'$, where the coefficient of y' is $\frac{1}{m}$, which is less than unity.

In the equation thus changed, we have assumed in our figures the units for x and y equal, and the axes rectangular. The effect of a different supposition in either particular can be readily understood.

3. *Curves whose equations are $y = a \sin x \sin mx$.* It was found convenient to draw several auxiliary curves whose equations are of the form,

$$y = a \sin x \sin mx. \quad (2)$$

A convenient arbitrary value being assumed for a , to m was given in turn all the values of the proper fractions, which, reduced to their lowest terms, have denominators less than 12. The forms of these curves are shown on plates XIV and XV, excepting a few in which m has 11 for denominator. In fig. 37 is shown the beginning of the curve when m has the irrational value $\sqrt{\frac{1}{3}}$. The axis of x is drawn

in the figures. The origin is the point at the left of each figure where the curve touches the axis of x .

It will be convenient at times to put $m = \frac{p}{q}$, and $n = \frac{p'}{q'}$, where p, q, p' , and q' , are integers, and the fractions are reduced to their lowest terms.

4. *Properties of the curves of Eq. (2).* By inspection of the curves on plates XIV and XV, and of their equations, we readily deduce the following properties :

- a. The value of y is not greater than a .
- b. When either x or mx is a multiple of π , $y=0$.
- c. There are maxima or minima values of y when $m \tan x = -\tan mx$.
- d. When m is rational the values of y repeat ; after $q\pi$ if p and q are both *odd* ; after $2q\pi$ if either p or q is even.
- e. When m is irrational the curve does not repeat its form.
- f. The curve is symmetrical about the axis of y , and about an axis through the middle point of each cycle.
- g. If p or q is *even*, the curve is symmetrical about the point $y=0$,

$$x = \frac{q\pi}{2}.$$
- h. There are, in each distance $2q\pi$ along x , $p+q$ maxima, and an equal number of minima values of y .

5. The value of y in Eq. (2) may be regarded as made up of two parts, since $y = a \sin x \sin mx = \frac{a}{2} \cos(1-m)x - \frac{a}{2} \cos(1+m)x$. In fig. 22, where $m = \frac{7}{8}$, these parts are separately shown. The continuous line represents the curve $y = \frac{a}{2} \cos(1 - \frac{7}{8})x$, having one complete oscillation in a distance of 16π along x . By laying off below and above this curve the second part of y , that is $-\frac{a}{2} \cos(1 + \frac{7}{8})x$, we have the curve $y = a \sin x \sin \frac{7}{8}x$.

6. *Use of the auxiliary curves, Eq. (2).* To draw the curves from equation (1), even after all the usual devices for saving labor have been employed, requires the frequent solution of equations of the form $\sin x \sin mx = c$. This equation gives a set of values of x for each cycle of the curve. To find each value of x requires a solution by trial and error, a very simple process, but when often repeated quite tedious. By the curves figs. 1-37 carefully traced on cross-section paper we may by merely running the eye along the line $y=c$

obtain by inspection all the values of x to a sufficient degree of accuracy.

7. *Equation (1) when $a=b=0$.* The equation

$$\sin y \sin my=0, \tag{3}$$

consists of the two equations $\sin y=0$, and $\sin my=0$, and is satisfied by the values $y=l\pi$, and $my=l\pi$, where l is 0, or any integer. In fig. 66 the horizontal lines belong to the equation $\sin y \sin \frac{4}{5}y=0$. They consist of two series, one at intervals of π , the other at intervals of $2\frac{1}{4}\pi$. If through the intersections of the curve in fig. 25 with the axis of x there be drawn lines perpendicular to that axis, the lines for $\sin x \sin \frac{4}{5}x=0$ would be obtained. The heavy lines of fig. 66 represent double lines, corresponding to points of tangency in fig. 25.

8. *Equation (1) where $a=0$.* The equation (1) becomes by making $a=0$, and for convenience changing the axes,

$$\sin x \sin mx=b. \tag{4}$$

This does not contain y , and therefore represents straight lines parallel to the axis of y . If the straight line $y=b$ be drawn parallel to the axis of x to cut the curve $y=\sin x \sin mx$, and through the several points of intersection straight lines be drawn parallel to the axis of y , these lines will evidently be those represented by the equation $\sin x \sin mx=b$.

In fig. 66 the vertical lines represent the equation $\sin x \sin \frac{5}{3}x=\frac{2}{5}$. If the curve in fig. 26 be cut by a line parallel to the axis of x and distant from it two-fifths of the largest ordinate, the intersections will correspond with the intersections of any horizontal line in fig. 66 by the several vertical lines.*

9. *Equation (1) where $a=1, b=0, m=n=1$.* The equation

$$\sin y \sin y=\sin x \sin x \tag{5}$$

becomes $\sin y=\pm\sin x$, or $y=l\pi \pm x$, l being 0, or an integer. The curve consists of two series of parallel equidistant straight lines, the one parallel to $y=x$, the other to $y=-x$, and both cutting the axes at intervals of π . The locus is represented in fig. 38, where the origin is any point of intersection.

10. *Equation (1) where $a=1, b=0, m=n=\frac{1}{q}$.* The equation

$$\sin y \sin \frac{1}{q}y=\sin x \sin \frac{1}{q}x \tag{6}$$

is one of the simpler examples of equation (1).

*The unit of abscissas in plates XIV and XV is smaller than in the other plates.

- a.* It is satisfied if $y=x$, or if $y=-x$. Hence the two straight lines $y=\pm x$ form part of the locus of equation (6).
- b.* If $2lq\pi+x$ be put for x , l being an integer, the equation is unchanged, whether q be *odd* or *even*.
- c.* If q be *odd* the equation will be unchanged if $lq\pi+x$ be put for x .
- d.* The curve repeats itself to the right and left, and also above and below, at intervals of $q\pi$ if q is *odd*, and at intervals of $2q\pi$ if q is *even*.
- e.* Straight lines parallel to $y=\pm x$, and cutting the axes at intervals of $q\pi$, or $2q\pi$, according as q is *odd* or *even*, belong to the locus of equation (6).
- f.* These straight lines divide the infinite plane of coördinates into equal squares for a given value of q . Each square contains a similar and equal portion of the locus. If q is *odd*, that portion is not always similarly placed, for it may have two positions with respect to an axis.
- g.* If q is *even*, isolated points at the centers of the squares (*f*) belong to the locus.
- h.* The equation (6) is satisfied if $\sin x=0$, and $\sin y=0$. Hence the locus of (6) passes through the angular points of all the squares formed by the lines $\sin x=0$, and $\sin y=0$ (Art. 7.)
- i.* A few curves representing equation (6) are shown in figs. 40-47. The axes are not drawn. Any point of intersection of straight lines that is surrounded by an oval may be taken for the origin. The several propositions of this article will be more easily understood by inspection of the curves.

11. Equation (1) when $a=1$, $b=0$, $m=n=\frac{p}{q}$. In this case the equation becomes,

$$\sin y \sin \frac{p}{q} y = \sin x \sin \frac{p}{q} x. \quad (7)$$

The properties of the curves of equation (7) are in many respects like those of equation (6).

- a.* The two straight lines $y=\pm x$ belong to the locus.
- b.* If p and q are both *odd*, the equation is unchanged, if y or x is increased or diminished by multiples of $q\pi$.
- c.* If either p or q is *even*, the equation is unchanged if y or x is increased or diminished by multiples of $2q\pi$.
- d.* The curve repeats in the direction of either axis; at intervals of $q\pi$ if p and q are both *odd*, at intervals of $2q\pi$ if either p or q is *even*.

- e.* Straight lines parallel to $y = \pm x$ and cutting the axes at intervals of $q\pi$, or $2q\pi$, according as $p+q$ is *even* or *odd*, belong to the locus of equation (7).
- f.* These straight lines divide the plane of coördinates into equal squares for any given value of m . Each square contains a similar and equal portion of the locus, though not always similarly placed.
- g.* Equation (7) is satisfied if $\sin x \sin mx = 0$, and $\sin y \sin my = 0$. Hence the locus passes through all the angular points of the rectangles formed by these two series of parallel straight lines (Art. 7).
- h.* If $p+q$ is *odd* isolated points appear, belonging to the locus, at the centers of the squares.
- i.* The maxima and minima values of y are determined by the equation $\frac{p}{q} \tan x = -\tan \frac{p}{q} x$ (Art. 4, *c*). This equation represents straight lines parallel to the axis of y . There are $2(p+q)$ of the lines (Art. 4, *h*) in an interval of $2q\pi$.
- j.* The same equation in y gives the maxima and minima values of x .
- k.* These equations are also the conditions of the isolated and double points. Hence there can be isolated or double points only at the intersections of the lines $\frac{p}{q} \tan x = -\tan \frac{p}{q} x$ with the lines $\frac{p}{q} \tan y = -\tan \frac{p}{q} y$.
- l.* The propositions (*i*), (*j*), and (*k*) hold equally true for any values of a and b in equation (1), and there are similar properties if m is not equal to n .
- m.* The figs. 48-65, 68, and 70, represent curves belonging to equation (7). Any point where two straight lines meet, and that is surrounded by an oval may be the origin.
- n.* If through the double points on the line $y=x$ vertical and horizontal lines be drawn, these lines will pass through all the points of maxima and minima ordinates and abscissas. By their intersections they will mark all the possible positions of double points for any values of a and b .

12. Equation (1) when $a=1$, $b=0$, $m=n=$ an irrational number.

The equation

$$\sin y \sin \sqrt{\frac{1}{5}} y = \sin x \sin \sqrt{\frac{1}{5}} x, \quad (8)$$

represents a class of curves that do not repeat their forms but change

continually throughout the plane of coördinates. The curve is symmetrical about either axis, and also about either line $y = \pm x$. These two lines belong to the curve.

The origin and a portion of the curve, principally in the first quadrant, are given in fig. 67, plate XXII.

13. *Equation (1) when $a = -1, b = 0, m = n = \frac{1}{q}$.* If q is *even* the equation

$$\sin y \sin \frac{1}{q} y = - \sin x \sin \frac{1}{q} x \quad (9)$$

merely changes the sign of the second member if we substitute $q\pi + x$ for x . Hence the curves in figures 40, 42, 44, and 47 represent equation (9), when q is *even*, the origin being at an isolated point.

But if q is *odd* we obtain new forms which have these properties.

a. The origin is an isolated point.

b. If $q = 1$, the locus consists solely of points (fig. 39).

c. If $q = 3$, each point is surrounded by one closed curve (fig. 93).

d. If $q = 5$, each point is surrounded by two closed curves (fig. 74).

e. The resemblance of these figures to parts of figs. 40, 42, and 44, and the law of their formation makes it unnecessary to give further examples.

f. A dot and four surrounding closed curves in fig. 47, would fairly represent the element for equation (9), when $q = 9$.

14. *Equation (1) when $a = -1, b = 0, m = n = \frac{p}{q}$.* Curves whose equations are of the form

$$\sin y \sin \frac{p}{q} y = - \sin x \sin \frac{p}{q} x \quad (10)$$

are shown in figures 69, 71, 99, and 108. There are no straight lines belonging to the locus. The origin is at any one of the isolated points. The first two are placed beside figures 68 and 70 for ease in comparing.

The following propositions of Art. 11, for equation (7) apply also to equation (10), without change of terms, viz: $b, c, d, g, i, j,$ and k .

15. *Equation (1) when $a = 1, b = 0, m = 1,$ and $n = \frac{p'}{q'}$.* The figures 76-79, and 81, represent curves whose equations are

$$\sin y \sin y = \sin x \sin \frac{p'}{q'} x. \quad (11)$$

In the direction of y they repeat at intervals of π . In the direction of x they repeat at intervals of $q'\pi$, or $2q'\pi$, according as $p'+q'$ is even or odd.

Fig. 80 gives a similar curve except that $a = -1$.

16. Equation (1) when $a=1, b=0, m = \frac{p}{q}$, and $n = \frac{p'}{q}$. The equation (11) is a special case of the equation

$$\sin y \sin \frac{p}{q} y = \sin x \sin \frac{p'}{q} x. \quad (12)$$

Examples of curves from equation (12) are given in figures 82–91, 123, and 141. The number of different curves that this equation gives us is quite large, even if q and q' are limited to small numbers. If 11 is the maximum value of q and q' , the number of independent curves belonging to the equation is nearly a thousand. Equations (5), (6), (7) and (11) are special cases of (12).

17. *Farther consideration of the curves of equation (12).*

- a. If the parallel straight lines $\sin x \sin \frac{p'}{q} x = 0$ be drawn (Art. 7) the plane of coördinates is divided by those lines into portions. When two lines coincide the portion between them may be regarded as real but infinitesimal. In crossing any of these lines the sign of the second member of (12) changes from plus to minus, or *vice-versa*.
- b. In like manner in crossing any of the parallel lines $\sin y \sin \frac{p}{q} y = 0$, the sign of the first member changes.
- c. The lines $\sin y \sin \frac{p}{q} y = 0$, and $\sin x \sin \frac{p'}{q} x = 0$, divide the plane into rectangles (some of which are infinitesimal). The curve of equation (12) passes through each of the angular points of these rectangles.
- d. Since the signs of the two members of (12) must be alike the curve passes at any angle of a rectangle into the rectangle vertically opposite. It passes from a rectangle only at the angles.
- e. If, however, any rectangle is of infinitesimal breadth and finite length, the curve at its extremity becomes tangent to the line that limits the infinitesimal parallelogram.
- f. If a rectangle becomes infinitesimal in both directions, the curve has at that point an isolated or a double point.
- g. The horizontal and vertical lines of fig. 148, and the rectangles formed by them, illustrate the above propositions. The con-

tinuous closed line represents the curve of equation (12) in this case. The axis of y is the heavy vertical line, and the axis of x the upper heavy horizontal line. These heavy lines are double lines.

- h.* Several of the propositions of Art. 11 apply to equation (12) with evident modifications. If $\frac{p}{q}$ is not equal to $\frac{p'}{q'}$, there are no straight lines belonging to the locus.
- i.* We may regard the plane of the curve as divided into equal rectangles by lines parallel to the axes, the altitudes of the rectangles being $q\pi$, or $2q\pi$, according as $p+q$ is *even* or *odd*, and whose bases are $q'\pi$, or $2q'\pi$, according as $p'+q'$ is *even* or *odd*. The curve (12) repeats itself in each of these rectangles without any variation, through the whole extent of the plane.
- j.* The origin of (12) is a real double point.

18. *Effect of a change of the value of a in equation (1), when $b=0$.* The effect of a change in the value of the coefficient of the second member may be observed by comparing some of the figures: for example, figs. 38 with 39; figs. 41 with 93; figs. 45 with 72 and 73; figs. 77 with 80; figs. 43 with 74 and 75; figs. 123 with 131-135; figs. 136 with 141 and 145.

19. The effect of the change of this factor can be better observed in the simpler equation

$$\sin y = k \sin x, \quad (13)$$

where k represents a as assuming several values. Figure 130 represents a *faisceau* of curves for equation (13). The origin is the nodal point near the lower left hand corner of the figure. Let k change from $-\infty$ to $+\infty$.

- a.* If $k = \infty$, we have the vertical equidistant straight lines.
- b.* If $k = -2$, we have the curved lines represented by uniform fine dots. At the origin it is tangent to $y = -2x$.
- c.* If $k = -1$, we have the straight lines in which dots and strokes alternate.
- d.* If $k = -\frac{1}{2}$, we have the continuous curved lines.
- e.* If $k = 0$, we have horizontal straight lines.
- f.* If $k = \frac{1}{2}$, we have the heavy dotted curved lines.
- g.* If $k = 1$, we have straight lines of which $y = x$ is one, and the others are similarly marked.
- h.* If $k = 2$, we have the curved lines consisting of a stroke and three dots alternating.

i. If $k = +\infty$, we have vertical lines again. The curve is at the origin always tangent to $y = kx$. The faisceau has nodal points wherever x and y are both multiples of π .

20. If we consider in like manner the faisceau of curves

$$\sin y \sin \frac{2}{3}y = k \sin x \sin \frac{2}{3}x, \quad (14)$$

for various values of k (fig. 148), we shall find similar but more complicated changes. The origin is the intersection of the heavy lines near the top of the figure. The figure represents the loci for six values of k , viz: ∞ , -1 , $-\frac{1}{2}$, 0 , $+1$, and $+2$. Each of the six loci passes through each nodal point, if isolated points be counted as branches of a locus.

- a.* For $k = \infty$, we have the vertical straight lines. The heavy line is a double line.
- b.* For $k = -1$, we have the uniformly dotted curves.
- c.* For $k = -\frac{1}{2}$, we have the curves represented by strokes and four dots alternating.
- d.* For $k = 0$, we have horizontal straight lines, the heavy lines being double.
- e.* For $k = 1$, we have the continuous curves (compare fig. 147).
- f.* For $k = 2$, we have the curved lines consisting of a long stroke and a short stroke alternating.

By removal upward or downward a distance of 5π , the curve (*b*) coincides with (*e*). In general any one of the curves by such change coincides with that one for which k has an equal value with opposite sign.

21. We may in like manner obtain a faisceau of curves from the equation

$$\sin y \sin my = k \sin x \sin nx + b, \quad (15)$$

by giving to k different values.

The curve will be the horizontal lines $\sin y \sin my = b$ (Art. 8), if $k = 0$. It will be the vertical lines $\sin x \sin nx = 0$, if $k = 0$. For other values of k , the curve will pass through all the points of intersections of these series of straight lines. Figure 66 represents (with the axes interchanged) the vertical and horizontal lines in a special case.

The lines of maxima and minima values of x and y , and the possible positions of double points (Art. 11, *i, j, k*), are independent of k and b . The origin is not upon the curve if k and b are finite.

22. *Change of value of b in equation (1).* It remains to consider the effect of a change in the constant b in equation (1). That it may change entirely the appearance of the locus will be seen by a glance at figures 92, 93, and 94. The same curves are superposed in fig. 95. Though each locus may have its own double points, they cannot when superposed cut each other.

23. In the figures 96-103, the curves of the equation

$$\sin y \sin \frac{2}{7}y = -\sin x \sin \frac{2}{7}x + k \quad (16)$$

are shown for certain specified values of k . The origin is the place of the isolated point in fig. 99. The several curves if superposed will not intersect. The values of k were selected so as to furnish curves with double points.

24. A series of twelve curves from the equation

$$\sin y \sin \frac{3}{11}y = -\sin x \sin \frac{3}{11}x + k \quad (17)$$

is given in the figures 104-115. By tracing any selected portions of the figure through the series the effect of the change in k will be seen. As in equation (16) values of k were chosen which give (except fig. 108) real double points. In each case other curves of the series with real double points might have been given.

25. Another series of fourteen curves is given in figures 116-129 from the equation

$$\sin y \sin \frac{1}{2}y = \sin x \sin \frac{3}{4}x + k. \quad (18)$$

The complete series would give 18 curves with double or isolated points. The omitted curves are those having isolated points, one at the beginning and one at the end of the series, one between figs. 127 and 128, and one between figs. 129 and 130.

26. Similar partial series can be seen in figs. 136-138, in figs. 139-143, and in figs. 144-146.

27. The superposition of the several curves of a series is shown in figure 147 where the curves represent the equation

$$\sin y \sin \frac{2}{5}y = \sin x \sin \frac{2}{5}x + k.$$

A little more than one complete figure of the curves is shown. The origin is at the double point near the top of the figure. The value of k varies from curve to curve by intervals of $\frac{1}{5}$, and it cannot numerically exceed 2. The full line corresponds to $k=0$.

The multiple that k is of $+\frac{1}{5}$ is denoted by the number of *dots* between the long strokes of the lines.

The multiple that k is of $-\frac{1}{5}$ is denoted by the number of *short strokes* between the long strokes in the lines.

28. The resemblance of fig. 147 to a series of contour lines in surveying, suggests a corresponding treatment of the equation. Let

$$z = \sin y \sin my - a \sin x \sin nx - b \tag{19}$$

be the equation of a surface, and let it be intersected by planes parallel to the plane of xy , and we may obtain the groups of curves described in Arts. 22-27.

The surface of equation (19) may be described by continuous motion, as follows: Let $z = \sin y \sin my$ be a plane curve (figs. 1-37), and let it move parallel to itself so that each point of it shall describe a straight line parallel to the axis of x . The curve shall then describe a cylindrical surface whose equation is

$$z = \sin y \sin my. \tag{20}$$

Let $z = -a \sin x \sin nx - b$ be the equation of a second plane curve, and let this curve move parallel to the plane xz , in such a manner that the axis of x of the curve shall always lie in the cylindrical surface (20). The curve will describe by its motion the surface of equation (19).

The surface will consist of one continuous sheet lying between the two parallel planes $z = \pm (1 + a + b)$, the positive numerical values of a and b being here taken.

29. By means of the two arbitrary constants, a and b , in equation (1) the curve may be made to pass through any two points of the plane.

In a rectangle whose base is $2q'\pi$, and altitude $2q\pi$, there are $4(p+q)(p'+q')$ possible positions of double points (Art. 11, k .) If the curve passes through such a point it must have there two branches real or imaginary.

Hence we may assign to a and b such values that the curve will have double points, in general, at any two of the $4(p+q)(p'+q')$ possible positions.

ERRATUM IN PLATE XVI.

In figure 40, plate XVI, there is a series of ovals about one-half of the real double points. There should be added to the curve, as represented, a like series of ovals around each of the remaining real double points.

V. ON THE EQUILIBRIUM OF HETEROGENEOUS SUBSTANCES.

BY J. WILLARD GIBBS.

“Die Energie der Welt ist constant.
Die Entropie der Welt strebt einem Maximum zu.”

CLAUSIUS.*

THE comprehension of the laws which govern any material system is greatly facilitated by considering the energy and entropy of the system in the various states of which it is capable. As the difference of the values of the energy for any two states represents the combined amount of work and heat received or yielded by the system when it is brought from one state to the other, and the difference of entropy is the limit of all the possible values of the integral $\int \frac{dQ}{t}$, (dQ denoting the element of the heat received from external sources, and t the temperature of the part of the system receiving it,) the varying values of the energy and entropy characterize in all that is essential the effects producible by the system in passing from one state to another. For by mechanical and thermodynamic contrivances, supposed theoretically perfect, any supply of work and heat may be transformed into any other which does not differ from it either in the amount of work and heat taken together or in the value of the integral $\int \frac{dQ}{t}$. But it is not only in respect to the external relations of a system that its energy and entropy are of predominant importance. As in the case of simply mechanical systems, (such as are discussed in theoretical mechanics,) which are capable of only one kind of action upon external systems, viz., the performance of mechanical work, the function which expresses the capability of the system for this kind of action also plays the leading part in the theory of equilibrium, the condition of equilibrium being that the variation of this function shall vanish, so in a thermodynamic system, (such as all material systems actually are,) which is capable of two different kinds of action upon external systems, the two functions which express the twofold capabilities of the system afford an almost equally simple criterion of equilibrium.

*Pogg. Ann. Bd. cxxv (1865), S. 400; or Mechanische Wärmetheorie, Abhand. ix., S. 44.

CRITERIA OF EQUILIBRIUM AND STABILITY.

The criterion of equilibrium for a material system which is isolated from all external influences may be expressed in either of the following entirely equivalent forms:

I. *For the equilibrium of any isolated system it is necessary and sufficient that in all possible variations of the state of the system which do not alter its energy, the variation of its entropy shall either vanish or be negative.* If ϵ denote the energy, and η the entropy of the system, and we use a subscript letter after a variation to indicate a quantity of which the value is not to be varied, the condition of equilibrium may be written

$$(\delta\eta)_{\epsilon} \leq 0. \tag{1}$$

II. *For the equilibrium of any isolated system it is necessary and sufficient that in all possible variations in the state of the system which do not alter its entropy, the variation of its energy shall either vanish or be positive.* This condition may be written

$$(\delta\epsilon)_{\eta} \geq 0. \tag{2}$$

That these two theorems are equivalent will appear from the consideration that it is always possible to increase both the energy and the entropy of the system, or to decrease both together, viz., by imparting heat to any part of the system or by taking it away. For, if condition (1) is not satisfied, there must be some variation in the state of the system for which

$$\delta\eta > 0 \text{ and } \delta\epsilon = 0;$$

therefore, by diminishing both the energy and the entropy of the system *in its varied state*, we shall obtain a state for which (considered as a variation from the original state)

$$\delta\eta = 0 \text{ and } \delta\epsilon < 0;$$

therefore condition (2) is not satisfied. Conversely, if condition (2) is not satisfied, there must be a variation in the state of the system for which

$$\delta\epsilon < 0 \text{ and } \delta\eta = 0;$$

hence there must also be one for which

$$\delta\epsilon = 0 \text{ and } \delta\eta > 0;$$

therefore condition (1) is not satisfied.

The equations which express the condition of equilibrium, as also its statement in words, are to be interpreted in accordance with the general usage in respect to differential equations, that is, infinitesimals

of higher orders than the first relatively to those which express the amount of change of the system are to be neglected. But to distinguish the different kinds of equilibrium in respect to stability, we must have regard to the absolute values of the variations. We will use Δ as the sign of variation in those equations which are to be construed *strictly*, i. e., in which infinitesimals of the higher orders are not to be neglected. With this understanding, we may express the necessary and sufficient conditions of the different kinds of equilibrium as follows;—for stable equilibrium

$$(\Delta\eta)_\varepsilon < 0, \text{ i. e., } (\Delta\varepsilon)_\eta > 0: \quad (3)$$

for neutral equilibrium there must be some variations in the state of the system for which

$$(\Delta\eta)_\varepsilon = 0, \text{ i. e., } (\Delta\varepsilon)_\eta = 0, \quad (4)$$

while in general

$$(\Delta\eta)_\varepsilon \leq 0, \text{ i. e., } (\Delta\varepsilon)_\eta \geq 0; \quad (5)$$

and for unstable equilibrium there must be some variations for which

$$(\Delta\eta)_\varepsilon > 0, \quad (6)$$

i. e., there must be some for which

$$(\Delta\varepsilon)_\eta < 0, \quad (7)$$

while in general

$$(\delta\eta)_\varepsilon \leq 0, \text{ i. e., } (\delta\varepsilon)_\eta \geq 0. \quad (8)$$

In these criteria of equilibrium and stability, account is taken only of *possible* variations. It is necessary to explain in what sense this is to be understood. In the first place, all variations in the state of the system which involve the transportation of any matter through any finite distance are of course to be excluded from consideration, although they may be capable of expression by infinitesimal variations of quantities which perfectly determine the state of the system. For example, if the system contains two masses of the same substance, not in contact, nor connected by other masses consisting of or containing the same substance or its components, an infinitesimal increase of the one mass with an equal decrease of the other is not to be considered as a possible variation in the state of the system. In addition to such cases of essential impossibility, if heat can pass by conduction or radiation from every part of the system to every other, only those variations are to be rejected as impossible, which involve changes which are prevented by passive forces or analogous resistances to change. But, if the system consist of parts between which there is supposed to be no thermal communication, it will be necessary to regard as impossible any diminution of the entropy of any of

these parts, as such a change can not take place without the passage of heat. This limitation may most conveniently be applied to the second of the above forms of the condition of equilibrium, which will then become

$$(\delta\varepsilon)_{\eta', \eta'', \text{ etc.}} \cong 0, \quad (9)$$

$\eta', \eta'', \text{ etc.}$, denoting the entropies of the various parts between which there is no communication of heat. When the condition of equilibrium is thus expressed, the limitation in respect to the conduction of heat will need no farther consideration.

In order to apply to any system the criteria of equilibrium which have been given, a knowledge is requisite of its passive forces or resistances to change, in so far, at least, as they are capable of *preventing* change. (Those passive forces which only retard change, like viscosity, need not be considered.) Such properties of a system are in general easily recognized upon the most superficial knowledge of its nature. As examples, we may instance the passive force of friction which prevents sliding when two surfaces of solids are pressed together,—that which prevents the different components of a solid, and sometimes of a fluid, from having different motions one from another,—that resistance to change which sometimes prevents either of two forms of the same substance (simple or compound), which are capable of existing, from passing into the other,—that which prevents the changes in solids which imply plasticity, (in other words, changes of the form to which the solid tends to return,) when the deformation does not exceed certain limits.

It is a characteristic of all these passive resistances that they prevent a certain kind of motion or change, however the initial state of the system may be modified, and to whatever external agencies of force and heat it may be subjected, within limits, it may be, but yet within limits which allow finite variations in the values of all the quantities which express the initial state of the system or the mechanical or thermal influences acting on it, without producing the change in question. The equilibrium which is due to such passive properties is thus widely distinguished from that caused by the balance of the active tendencies of the system, where an external influence, or a change in the initial state, infinitesimal in amount, is sufficient to produce change either in the positive or negative direction. Hence the ease with which these passive resistances are recognized. Only in the case that the state of the system lies so near the limit at which the resistances cease to be operative to prevent change, as to create a

doubt whether the case falls within or without the limit, will a more accurate knowledge of these resistances be necessary.

To establish the validity of the criterion of equilibrium, we will consider first the sufficiency, and afterwards the necessity, of the condition as expressed in either of the two equivalent forms.

In the first place, if the system is in a state in which its entropy is greater than in any other state of the same energy, it is evidently in equilibrium, as any change of state must involve either a decrease of entropy or an increase of energy, which are alike impossible for an isolated system. We may add that this is a case of *stable* equilibrium, as no infinitely small cause (whether relating to a variation of the initial state or to the action of any external bodies) can produce a finite change of state, as this would involve a finite decrease of entropy or increase of energy.

We will next suppose that the system has the greatest entropy consistent with its energy, and therefore the least energy consistent with its entropy, but that there are other states of the same energy and entropy as its actual state. In this case, it is impossible that any motion of masses should take place; for if any of the energy of the system should come to consist of *vis viva* (of sensible motions), a state of the system identical in other respects but without the motion would have less energy and not less entropy, which would be contrary to the supposition. (But we cannot apply this reasoning to the motion within any mass of its different components in different directions, as in diffusion, when the momenta of the components balance one another.) Nor, in the case supposed, can any conduction of heat take place, for this involves an increase of entropy, as heat is only conducted from bodies of higher to those of lower temperature. It is equally impossible that any changes should be produced by the transfer of heat by radiation. The condition which we have supposed is therefore sufficient for equilibrium, so far as the motion of masses and the transfer of heat are concerned, but to show that the same is true in regard to the motions of diffusion and chemical or molecular changes, when these can occur without being accompanied or followed by the motions of masses or the transfer of heat, we must have recourse to considerations of a more general nature. The following considerations seem to justify the belief that the condition is sufficient for equilibrium in every respect.

Let us suppose, in order to test the tenability of such a hypothesis, that a system may have the greatest entropy consistent with its energy without being in equilibrium. In such a case, changes in the

state of the system must take place, but these will necessarily be such that the energy and the entropy will remain unchanged and the system will continue to satisfy the same condition, as initially, of having the greatest entropy consistent with its energy. Let us consider the change which takes place in any time so short that the change may be regarded as uniform in nature throughout that time. This time must be so chosen that the change does not take place in it infinitely slowly, which is always easy, as the change which we suppose to take place cannot be infinitely slow except at particular moments. Now no change whatever in the state of the system, which does not alter the value of the energy, and which commences with the same state in which the system was supposed at the commencement of the short time considered, will cause an increase of entropy. Hence, it will generally be possible by some slight variation in the circumstances of the case to make all changes in the state of the system like or nearly like that which is supposed actually to occur, and not involving a change of energy, to involve a necessary decrease of entropy, which would render any such change impossible. This variation may be in the values of the variables which determine the state of the system, or in the values of the constants which determine the nature of the system, or in the form of the functions which express its laws,—only there must be nothing in the system as modified which is thermodynamically impossible. For example, we might suppose temperature or pressure to be varied, or the composition of the different bodies in the system, or, if no small variations which could be actually realized would produce the required result, we might suppose the properties themselves of the substances to undergo variation, subject to the general laws of matter. If, then, there is any tendency toward change in the system as first supposed, it is a tendency which can be entirely checked by an infinitesimal variation in the circumstances of the case. As this supposition cannot be allowed, we must believe that a system is always in equilibrium when it has the greatest entropy consistent with its energy, or, in other words, when it has the least energy consistent with its entropy.

The same considerations will evidently apply to any case in which a system is in such a state that $\Delta\eta \leq 0$ for any possible infinitesimal variation of the state for which $\Delta\varepsilon = 0$, even if the entropy is not the least of which the system is capable with the same energy. (The term *possible* has here the meaning previously defined, and the character Δ is used, as before, to denote that the equations are to be

construed strictly, i. e., without neglect of the infinitesimals of the higher orders.)

The only case in which the sufficiency of the condition of equilibrium which has been given remains to be proved is that in which in our notation $\delta\eta \leq 0$ for all possible variations not affecting the energy, but for some of these variations $\Delta\eta > 0$, that is, when the entropy has in some respects the characteristics of a minimum. In this case the considerations adduced in the last paragraph will not apply without modification, as the change of state may be infinitely slow at first, and it is only in the initial state that the condition $\delta\eta_e \leq 0$ holds true. But the differential coefficients of all orders of the quantities which determine the state of the system, taken with respect of the time, must be functions of these same quantities. None of these differential coefficients can have any value other than 0, for the state of the system for which $\delta\eta_e \leq 0$. For otherwise, as it would generally be possible, as before, by some infinitely small modification of the case, to render impossible any change like or nearly like that which might be supposed to occur, this infinitely small modification of the case would make a finite difference in the value of the differential coefficients which had before the finite values, or in some of lower orders, which is contrary to that continuity which we have reason to expect. Such considerations seem to justify us in regarding such a state as we are discussing as one of theoretical equilibrium; although as the equilibrium is evidently unstable, it cannot be realized.

We have still to prove that the condition enunciated is in every case necessary for equilibrium. It is evidently so in all cases in which the active tendencies of the system are so balanced that changes of every kind, except those excluded in the statement of the condition of equilibrium, can take place *reversibly*, (i. e., both in the positive and the negative direction,) in states of the system differing infinitely little from the state in question. In this case, we may omit the sign of inequality and write as the condition of such a state of equilibrium

$$(\delta\eta)_e = 0, \quad \text{i. e.,} \quad (\delta\varepsilon)_\eta = 0 \quad (10)$$

But to prove that the condition previously enunciated is in every case necessary, it must be shown that whenever an isolated system remains without change, if there is any infinitesimal variation in its state, not involving a finite change of position of any (even an infinitesimal part) of its matter, which would diminish its energy by a

quantity which is not infinitely small relatively to the variations of the quantities which determine the state of the system, without altering its entropy,—or, if the system has thermally isolated parts, without altering the entropy of any such part,—this variation involves changes in the system which are prevented by its passive forces or analogous resistances to change. Now, as the described variation in the state of the system diminishes its energy without altering its entropy, it must be regarded as theoretically possible to produce that variation by some process, perhaps a very indirect one, so as to gain a certain amount of work (above all expended on the system). Hence we may conclude that the active forces or tendencies of the system favor the variation in question, and that equilibrium cannot subsist unless the variation is prevented by passive forces.

The preceding considerations will suffice, it is believed, to establish the validity of the criterion of equilibrium which has been given. The criteria of stability may readily be deduced from that of equilibrium. We will now proceed to apply these principles to systems consisting of heterogeneous substances and deduce the special laws which apply to different classes of phenomena. For this purpose we shall use the second form of the criterion of equilibrium, both because it admits more readily the introduction of the condition that there shall be no thermal communication between the different parts of the system, and because it is more convenient, as respects the form of the general equations relating to equilibrium, to make the entropy one of the independent variables which determine the state of the system, than to make the energy one of these variables.

THE CONDITIONS OF EQUILIBRIUM FOR HETEROGENEOUS MASSES IN CONTACT WHEN UNINFLUENCED BY GRAVITY, ELECTRICITY, DISTORTION OF THE SOLID MASSES, OR CAPILLARY TENSIONS.

In order to arrive as directly as possible at the most characteristic and essential laws of chemical equilibrium, we will first give our attention to a case of the simplest kind. We will examine the conditions of equilibrium of a mass of matter of various kinds enclosed in a rigid and fixed envelop, which is impermeable to and unalterable by any of the substances enclosed, and perfectly non-conducting to heat. We will suppose that the case is not complicated by the action of gravity, or by any electrical influences, and that in the solid portions of the mass the pressure is the same in every direction.

We will farther simplify the problem by supposing that the variations of the parts of the energy and entropy which depend upon the surfaces separating heterogeneous masses are so small in comparison with the variations of the parts of the energy and entropy which depend upon the quantities of these masses, that the former may be neglected by the side of the latter; in other words, we will exclude the considerations which belong to the theory of capillarity.

It will be observed that the supposition of a rigid and non-conducting envelop enclosing the mass under discussion involves no real loss of generality, for if any mass of matter is in equilibrium, it would also be so, if the whole or any part of it were enclosed in an envelop as supposed; therefore the conditions of equilibrium for a mass thus enclosed are the general conditions which must always be satisfied in case of equilibrium. As for the other suppositions which have been made, all the circumstances and considerations which are here excluded will afterward be made the subject of special discussion.

Conditions relating to the Equilibrium between the initially existing Homogeneous Parts of the given Mass.

Let us first consider the energy of any homogeneous part of the given mass, and its variation for any possible variation in the composition and state of this part. (By *homogeneous* is meant that the part in question is uniform throughout, not only in chemical composition, but also in physical state.) If we consider the amount and kind of matter in this homogeneous mass as fixed, its energy ε is a function of its entropy η , and its volume v , and the differentials of these quantities are subject to the relation

$$d\varepsilon = t d\eta - p dv, \quad (11)$$

t denoting the (absolute) temperature of the mass, and p its pressure. For $t d\eta$ is the heat received, and $p dv$ the work done, by the mass during its change of state. But if we consider the matter in the mass as variable, and write m_1, m_2, \dots, m_n for the quantities of the various substances S_1, S_2, \dots, S_n of which the mass is composed, ε will evidently be a function of $\eta, v, m_1, m_2, \dots, m_n$, and we shall have for the complete value of the differential of ε

$$d\varepsilon = t d\eta - p dv + \mu_1 dm_1 + \mu_2 dm_2 \dots + \mu_n dm_n, \quad (12)$$

$\mu_1, \mu_2, \dots, \mu_n$ denoting the differential coefficients of ε taken with respect to m_1, m_2, \dots, m_n .

The substances S_1, S_2, \dots, S_n , of which we consider the mass composed, must of course be such that the values of the differen-

tials dm_1, dm_2, \dots, dm_n shall be independent, and shall express every possible variation in the composition of the homogeneous mass considered, including those produced by the absorption of substances different from any initially present. It may therefore be necessary to have terms in the equation relating to component substances which do not initially occur in the homogeneous mass considered, provided, of course, that these substances, or their components, are to be found in some part of the whole given mass.

If the conditions mentioned are satisfied, the choice of the substances which we are to regard as the components of the mass considered, may be determined entirely by convenience, and independently of any theory in regard to the internal constitution of the mass. The number of components will sometimes be greater, and sometimes less, than the number of chemical elements present. For example, in considering the equilibrium in a vessel containing water and free hydrogen and oxygen, we should be obliged to recognize three components in the gaseous part. But in considering the equilibrium of dilute sulphuric acid with the vapor which it yields, we should have only two components to consider in the liquid mass, sulphuric acid (anhydrous, or of any particular degree of concentration) and (additional) water. If, however, we are considering sulphuric acid in a state of maximum concentration in connection with substances which might possibly afford water to the acid, it must be noticed that the condition of the independence of the differentials will require that we consider the acid in the state of maximum concentration as one of the components. The quantity of this component will then be capable of variation both in the positive and in the negative sense, while the quantity of the other component can increase but cannot decrease below the value 0.

For brevity's sake, we may call a substance S_a an *actual component* of any homogeneous mass, to denote that the quantity m_a of that substance in the given mass may be either increased or diminished (although we may have so chosen the other component substances that $m_a = 0$); and we may call a substance S_b a *possible component* to denote that it may be combined with, but cannot be subtracted from the homogeneous mass in question. In this case, as we have seen in the above example, we must so choose the component substances that $m_b = 0$.

The units by which we measure the substances of which we regard the given mass as composed may each be chosen independently. To fix our ideas for the purpose of a general discussion, we may suppose

all substances measured by weight or mass. Yet in special cases, it may be more convenient to adopt chemical equivalents as the units of the component substances.

It may be observed that it is not necessary for the validity of equation (12) that the variations of nature and state of the mass to which the equation refers should be such as do not disturb its homogeneity, provided that in all parts of the mass the variations of nature and state are infinitely small. For, if this last condition be not violated, an equation like (12) is certainly valid for all the infinitesimal parts of the (initially) homogeneous mass; i. e., if we write $D\varepsilon$, $D\eta$, etc., for the energy, entropy, etc., of any infinitesimal part, $dD\varepsilon = t dD\eta - p dDv + \mu_1 dDm_1 + \mu_2 dDm_2 \dots + \mu_n dDm_n$, (13) whence we may derive equation (12) by integrating for the whole initially homogeneous mass.

We will now suppose that the whole mass is divided into parts so that each part is homogeneous, and consider such variations in the energy of the system as are due to variations in the composition and state of the several parts remaining (at least approximately) homogeneous, and together occupying the whole space within the envelop. We will at first suppose the case to be such that the component substances are the same for each of the parts, each of the substances $S_1, S_2, \dots S_n$ being an actual component of each part. If we distinguish the letters referring to the different parts by accents, the variation in the energy of the system may be expressed by $\delta\varepsilon' + \delta\varepsilon'' + \text{etc.}$, and the general condition of equilibrium requires that

$$\delta\varepsilon' + \delta\varepsilon'' + \text{etc.} \geq 0 \quad (14)$$

for all variations which do not conflict with the *equations of condition*. These equations must express that the entropy of the whole given mass does not vary, nor its volume, nor the total quantities of any of the substances $S_1, S_2, \dots S_n$. We will suppose that there are no other equations of condition. It will then be necessary for equilibrium that

$$\begin{aligned} & t' \delta\eta' - p' \delta v' + \mu_1' \delta m_1' + \mu_2' \delta m_2' \dots + \mu_n' \delta m_n' \\ & + t'' \delta\eta'' - p'' \delta v'' + \mu_1'' \delta m_1'' + \mu_2'' \delta m_2'' \dots + \mu_n'' \delta m_n'' \\ & + \text{etc.} \geq 0 \end{aligned} \quad (15)$$

for any values of the variations for which

$$\delta\eta' + \delta\eta'' + \delta\eta''' + \text{etc.} = 0, \quad (16)$$

$$\delta v' + \delta v'' + \delta v''' + \text{etc.} = 0, \quad (17)$$

$$\left. \begin{aligned} \delta m_1' + \delta m_1'' + \delta m_1''' + \text{etc.} &= 0, \\ \delta m_2' + \delta m_2'' + \delta m_2''' + \text{etc.} &= 0, \\ \text{and } \delta m_n' + \delta m_n'' + \delta m_n''' + \text{etc.} &= 0. \end{aligned} \right\} \quad (18)$$

For this it is evidently necessary and sufficient that

$$t' = t'' = t''' = \text{etc.} \quad (19)$$

$$p' = p'' = p''' = \text{etc.} \quad (20)$$

$$\left. \begin{aligned} \mu_1' &= \mu_1'' = \mu_1''' = \text{etc.} \\ \mu_2' &= \mu_2'' = \mu_2''' = \text{etc.} \\ \mu_n' &= \mu_n'' = \mu_n''' = \text{etc.} \end{aligned} \right\} \quad (21)$$

Equations (19) and (20) express the conditions of thermal and mechanical equilibrium, viz., that the temperature and the pressure must be constant throughout the whole mass. In equations (21) we have the conditions characteristic of chemical equilibrium. If we call a quantity μ_x , as defined by such an equation as (12), the *potential* for the substance S_x in the homogeneous mass considered, these conditions may be expressed as follows:

The potential for each component substance must be constant throughout the whole mass.

It will be remembered that we have supposed that there is no restriction upon the freedom of motion or combination of the component substances, and that each is an actual component of all parts of the given mass.

The state of the whole mass will be completely determined (if we regard as immaterial the position and form of the various homogeneous parts of which it is composed), when the values are determined of the quantities of which the variations occur in (15). The number of these quantities, which we may call the independent variables, is evidently $(n + 2)\nu$, ν denoting the number of homogeneous parts into which the whole mass is divided. All the quantities which occur in (19), (20), (21), are functions of these variables, and may be regarded as known functions, if the energy of each part is known as a function of its entropy, volume, and the quantities of its components. (See eq. (12).) Therefore, equations (19), (20), (21), may be regarded as $(\nu - 1)(n + 2)$ independent equations between the independent variables. The volume of the whole mass and the total quantities of the various substances being known afford $n + 1$ additional equations. If we also know the total energy of the given mass, or its total entropy, we will have as many equations as there are independent variables.

But if any of the substances $S_1, S_2 \dots S_n$ are only possible components of some parts of the given mass, the variation δm of the quantity of such a substance in such a part cannot have a negative value, so that the general condition of equilibrium (15) does not require that the potential for that substance in that part should be equal to the potential for the same substance in the parts of which it is an actual component, but only that it shall not be less. In this case instead of (21) we may write

$$\left. \begin{array}{l}
 \mu_1 = M_1 \\
 \text{for all parts of which } S_1 \text{ is an actual component, and} \\
 \mu_1 \geq M_1 \\
 \text{for all parts of which } S_1 \text{ is a possible (but not actual) component,} \\
 \mu_2 = M_2 \\
 \text{for all parts of which } S_2 \text{ is an actual component, and} \\
 \mu_2 \geq M_2 \\
 \text{for all parts of which } S_2 \text{ is a possible (but not actual) component,} \\
 \text{etc.,}
 \end{array} \right\} \quad (22)$$

M_1, M_2 , etc., denoting constants of which the value is only determined by these equations.

If we now suppose that the components (actual or possible) of the various homogeneous parts of the given mass are not the same, the result will be of the same character as before, provided that all the different components are *independent*, (i. e., that no one can be made out of the others,) so that the total quantity of each component is fixed. The general condition of equilibrium (15) and the equations of condition (16), (17), (18) will require no change, except that, if any of the substances $S_1, S_2 \dots S_n$ is not a component (actual or possible) of any part, the term $\mu \delta m$ for that substance and part will be wanting in the former, and the δm in the latter. This will require no change in the form of the particular conditions of equilibrium as expressed by (19), (20), (22); but the number of single conditions contained in (22) is of course less than if all the component substances were components of all the parts. Whenever, therefore, each of the different homogeneous parts of the given mass may be regarded as composed of some or of all of the same set of substances, no one of which can be formed out of the others, the condition which (with equality of temperature and pressure) is necessary and sufficient for equilibrium between the different parts of the given mass may be expressed as follows:

The potential for each of the component substances must have a constant value in all parts of the given mass of which that substance is an actual component, and have a value not less than this in all parts of which it is a possible component.

The number of equations afforded by these conditions, after elimination of $M_1, M_2, \dots M_n$, will be less than $(n + 2)(\nu - 1)$ by the number of terms in (15) in which the variation of the form δm is either necessarily nothing or incapable of a negative value. The number of variables to be determined is diminished by the same number, or, if we choose, we may write an equation of the form $m = 0$ for each of these terms. But when the substance is a possible component of the part concerned, there will also be a condition (expressed by \geq) to show whether the supposition that the substance is not an actual component is consistent with equilibrium.

We will now suppose that the substances $S_1, S_2, \dots S_n$ are not all independent of each other, i. e., that some of them can be formed out of others. We will first consider a very simple case. Let S_3 be composed of S_1 and S_2 combined in the ratio of a to b , S_1 and S_2 occurring as actual components in some parts of the given mass, and S_3 in other parts, which do not contain S_1 and S_2 as separately variable components. The general condition of equilibrium will still have the form of (15) with certain of the terms of the form $\mu \delta m$ omitted. It may be written more briefly

$$\sum (t \delta \eta) - \sum (p \delta v) + \sum (\mu_1 \delta m_1) + \sum (\mu_2 \delta m_2) \dots + \sum (\mu_n \delta m_n) \geq 0,$$

the sign \sum denoting summation in regard to the different parts of the given mass. But instead of the three equations of condition,

$$\sum \delta m_1 = 0, \quad \sum \delta m_2 = 0, \quad \sum \delta m_3 = 0, \quad (24)$$

we shall have the two,

$$\left. \begin{aligned} \sum \delta m_1 + \frac{a}{a+b} \sum \delta m_3 &= 0, \\ \sum \delta m_2 + \frac{b}{a+b} \sum \delta m_3 &= 0. \end{aligned} \right\} (25)$$

The other equations of condition,

$$\sum \delta \eta = 0, \quad \sum \delta v = 0, \quad \sum \delta m_4 = 0, \quad \text{etc.}, \quad (26)$$

will remain unchanged. Now as all values of the variations which satisfy equations (24) will also satisfy equations (25), it is evident that all the particular conditions of equilibrium which we have already deduced, (19), (20), (22), are necessary in this case also. When these are satisfied, the general condition (23) reduces to

$$M_1 \sum \delta m_1 + M_2 \sum \delta m_2 + M_3 \sum \delta m_3 \geq 0. \quad (27)$$

For, although it may be that μ_1' , for example, is greater than M_1 , yet it can only be so when the following $\delta m_1'$ is incapable of a negative value. Hence, if (27) is satisfied, (23) must also be. Again, if (23) is satisfied, (27) must also be satisfied, so long as the variation of the quantity of every substance has the value 0 in all the parts of which it is not an actual component. But as this limitation does not affect the range of the possible values of $\sum \delta m_1$, $\sum \delta m_2$, and $\sum \delta m_3$, it may be disregarded. Therefore the conditions (23) and (27) are entirely equivalent, when (19), (20), (22) are satisfied. Now, by means of the equations of condition (25), we may eliminate $\sum \delta m_1$ and $\sum \delta m_2$ from (27), which becomes

$$- a M_1 \sum \delta m_3 - b M_2 \sum \delta m_3 + (a + b) M_3 \sum \delta m_3 \geq 0, \quad (28)$$

i. e., as the value of $\sum \delta m_3$ may be either positive or negative,

$$a M_1 + b M_2 = (a + b) M_3, \quad (29)$$

which is the additional condition of equilibrium which is necessary in this case.

The relations between the component substances may be less simple than in this case, but in any case they will only affect the equations of condition, and these may always be found without difficulty, and will enable us to eliminate from the general condition of equilibrium as many variations as there are equations of condition, after which the coefficients of the remaining variations may be set equal to zero, except the coefficients of variations which are incapable of negative values, which coefficients must be equal to or greater than zero. It will be easy to perform these operations in each particular case, but it may be interesting to see the form of the resultant equations in general.

We will suppose that the various homogeneous parts are considered as having in all n components, S_1, S_2, \dots, S_n , and that there is no restriction upon their freedom of motion and combination. But we will so far limit the generality of the problem as to suppose that each of these components is an actual component of some part of the given mass.* If some of these components can be formed out of others, all such relations can be expressed by equations such as

$$\alpha \mathcal{E}_a + \beta \mathcal{E}_b + \text{etc.} = \kappa \mathcal{E}_k + \lambda \mathcal{E}_l + \text{etc.} \quad (30)$$

where $\mathcal{E}_a, \mathcal{E}_b, \mathcal{E}_k$, etc. denote the units of the substances S_a, S_b, S_k , etc.,

* When we come to seek the conditions of equilibrium relating to the formation of masses unlike any previously existing, we shall take up *de novo* the whole problem of the equilibrium of heterogeneous masses enclosed in a non-conducting envelop, and give it a more general treatment, which will be free from this limitation.

(that is, of certain of the substances $S_1, S_2, \dots S_n$) and α, β, κ , etc. denote numbers. These are not, it will be observed, equations between abstract quantities, but the sign $=$ denotes qualitative as well as quantitative equivalence. We will suppose that there are r independent equations of this character. The equations of condition relating to the component substances may easily be derived from these equations, but it will not be necessary to consider them particularly. It is evident that they will be satisfied by any values of the variations which satisfy equations (18); hence, the particular conditions of equilibrium (19), (20), (22) must be necessary in this case, and, if these are satisfied, the general equation of equilibrium (15) or (23) will reduce to

$$M_1 \sum \delta m_1 + M_2 \sum \delta m_2 \dots + M_n \sum \delta m_n \geq 0. \quad (31)$$

This will appear from the same considerations which were used in regard to equations (23) and (27). Now it is evidently possible to give to $\sum \delta m_a, \sum \delta m_b, \sum \delta m_k$, etc. values proportional to $\alpha, \beta, -\kappa$, etc. in equation (30), and also the same values taken negatively, making $\sum \delta m = 0$ in each of the other terms; therefore

$$\alpha M_a + \beta M_b + \text{etc.} \dots - \kappa M_k - \lambda M_l - \text{etc.} = 0, \quad (32)$$

or,

$$\alpha M_a + \beta M_b + \text{etc.} = \kappa M_k + \lambda M_l + \text{etc.} \quad (33)$$

It will be observed that this equation has the same form and coefficients as equation (30), M taking the place of \mathfrak{S} . It is evident that there must be a similar condition of equilibrium for every one of the r equations of which (30) is an example, which may be obtained simply by changing \mathfrak{S} in these equations into M . When these conditions are satisfied, (31) will be satisfied with any possible values of $\sum \delta m_1, \sum \delta m_2, \dots \sum \delta m_n$. For no values of these quantities are possible, except such that the equation

$$(\sum \delta m_1) \mathfrak{S}_1 + (\sum \delta m_2) \mathfrak{S}_2 \dots + (\sum \delta m_n) \mathfrak{S}_n = 0 \quad (34)$$

after the substitution of these values, can be derived from the r equations like (30), by the ordinary processes of the reduction of linear equations. Therefore, on account of the correspondence between (31) and (34), and between the r equations like (33) and the r equations like (30), the conditions obtained by giving any possible values to the variations in (31) may also be derived from the r equations like (33); that is, the condition (31) is satisfied, if the r equations like (33) are satisfied. Therefore the r equations like (33) are with (19), (20), and (22) the equivalent of the general condition (15) or (23).

For determining the state of a given mass when in equilibrium and having a given volume and given energy or entropy, the condition of equilibrium affords an additional equation corresponding to each of the r independent relations between the n component substances. But the equations which express our knowledge of the matter in the given mass will be correspondingly diminished, being $n - r$ in number, like the equations of condition relating to the quantities of the component substances, which may be derived from the former by differentiation.

Conditions relating to the possible Formation of Masses Unlike any Previously Existing.

The variations which we have hitherto considered do not embrace every possible infinitesimal variation in the state of the given mass, so that the particular conditions already formed, although always necessary for equilibrium (when there are no other equations of condition than such as we have supposed), are not always sufficient. For, besides the infinitesimal variations in the state and composition of different parts of the given mass, infinitesimal masses may be formed entirely different in state and composition from any initially existing. Such parts of the whole mass in its varied state as cannot be regarded as parts of the initially existing mass which have been infinitesimally varied in state and composition, we will call *new parts*. These will necessarily be infinitely small. As it is more convenient to regard a vacuum as a limiting case of extreme rarefaction than to give a special consideration to the possible formation of empty spaces within the given mass, the term *new parts* will be used to include any empty spaces which may be formed, when such have not existed initially. We will use $D\varepsilon$, $D\eta$, Dv , Dm_1 , Dm_2 , . . . Dm_n to denote the infinitesimal energy, entropy, and volume of any one of these new parts, and the infinitesimal quantities of its components. The component substances S_1 , S_2 , . . . S_n must now be taken to include not only the independently variable components (actual or possible) of all parts of the given mass as initially existing, but also the components of all the new parts, the possible formation of which we have to consider. The character δ will be used as before to express the infinitesimal variations of the quantities relating to those parts which are only infinitesimally varied in state and composition, and which for distinction we will call *original parts*, including under this term the empty spaces, if such exist initially, within the envelop bounding the system. As we may divide the given mass into as many parts as we choose, and as not only the

Now, using Lagrange's "method of multipliers,"* we will subtract $T (\sum \delta \eta + \sum D\eta) - P (\sum \delta v + \sum Dv)$ from the first member of the general condition of equilibrium (37), T and P being constants of which the value is as yet arbitrary. We might proceed in the same way with the remaining equations of condition, but we may obtain the same result more simply in another way. We will first observe that

$$(\sum \delta m_1 + \sum Dm_1) \xi_1 + (\sum \delta m_2 + \sum Dm_2) \xi_2 \dots + (\sum \delta m_n + \sum Dm_n) \xi_n = 0, \quad (42)$$

which equation would hold identically for any possible values of the quantities in the parentheses, if for r of the letters $\xi_1, \xi_2, \dots, \xi_n$ were substituted their values in terms of the others as derived from equations (38). (Although $\xi_1, \xi_2, \dots, \xi_n$ do not represent abstract quantities, yet the operations necessary for the reduction of linear equations are evidently applicable to equations (38).) Therefore, equation (42) will hold true if for $\xi_1, \xi_2, \dots, \xi_n$ we substitute n numbers which satisfy equations (38). Let M_1, M_2, \dots, M_n be such numbers, i. e., let

$$\left. \begin{aligned} a_1 M_1 + a_2 M_2 \dots + a_n M_n &= 0, \\ b_1 M_1 + b_2 M_2 \dots + b_n M_n &= 0, \\ \text{etc.} \end{aligned} \right\} r \text{ equations, } (43)$$

then

$$M_1 (\sum \delta m_1 + \sum Dm_1) + M_2 (\sum \delta m_2 + \sum Dm_2) \dots + M_n (\sum \delta m_n + \sum Dm_n) = 0. \quad (44)$$

This expression, in which the values of $n - r$ of the constants M_1, M_2, \dots, M_n are still arbitrary, we will also subtract from the first member of the general condition of equilibrium (37), which will then become

$$\begin{aligned} \sum D\varepsilon + \sum (t \delta \eta) - \sum (p \delta v) + \sum (\mu_1 \delta m_1) \dots + \sum (\mu_n \delta m_n) \\ - T \sum \delta \eta + P \sum \delta v - M_1 \sum \delta m_1 \dots + M_n \sum \delta m_n \\ - T \sum D\eta + P \sum Dv - M_1 \sum Dm_1 \dots - M_n \sum Dm_n \geq 0. \end{aligned} \quad (45)$$

That is, having assigned to $T, P, M_1, M_2, \dots, M_n$ any values consistent with (43), we may assert that it is necessary and sufficient for equilibrium that (45) shall hold true for any variations in the state of the system consistent with the equations of condition (39), (40), (41). But it will always be possible, in case of equilibrium, to assign such values to $T, P, M_1, M_2, \dots, M_n$, without violating equations (43).

* On account of the sign \geq in (37), and because some of the variations are incapable of negative values, the successive steps in the reasoning will be developed at greater length than would be otherwise necessary.

that (45) shall hold true for all variations in the state of the system and in the quantities of the various substances composing it, even though these variations are not consistent with the equations of condition (39), (40), (41). For, when it is not possible to do this, it must be possible by applying (45) to variations in the system not necessarily restricted by the equations of condition (39), (40), (41) to obtain conditions in regard to $T, P, M_1, M_2, \dots M_n$, some of which will be inconsistent with others or with equations (43). These conditions we will represent by

$$A \geq 0, \quad B \geq 0, \quad \text{etc.}, \tag{46}$$

A, B , etc. being linear functions of $T, P, M_1, M_2, \dots M_n$. Then it will be possible to deduce from these conditions a single condition of the form

$$\alpha A + \beta B + \text{etc.} \geq 0, \tag{47}$$

α, β , etc. being positive constants, which cannot hold true consistently with equations (43). But it is evident from the form of (47) that, like any of the conditions (46), it could have been obtained directly from (45) by applying this formula to a certain change in the system (perhaps not restricted by the equations of condition (39), (40), (41)). Now as (47) cannot hold true consistently with eqs. (43), it is evident, in the first place, that it cannot contain T or P , therefore in the change in the system just mentioned (for which (45) reduces to (47))

$$\sum \delta \eta + \sum D\eta = 0, \quad \text{and} \quad \sum \delta v + \sum Dv = 0,$$

so that the equations of condition (39) and (40) are satisfied. Again, for the same reason, the homogeneous function of the first degree of $M_1, M_2, \dots M_n$ in (47) must be one of which the value is fixed by eqs. (43). But the value thus fixed can only be zero, as is evident from the form of these equations. Therefore

$$\begin{aligned} (\sum \delta m_1 + \sum Dm_1) M_1 + (\sum \delta m_2 + \sum Dm_2) M_2 \dots \\ + (\sum \delta m_n + \sum Dm_n) M_n = 0 \end{aligned} \tag{48}$$

for any values of $M_1, M_2 \dots M_n$ which satisfy eqs. (43), and therefore

$$\begin{aligned} (\sum \delta m_1 + \sum Dm_1) \xi_1 + (\sum \delta m_2 + \sum Dm_2) \xi_2 \dots \\ + (\sum \delta m_n + \sum Dm_n) \xi_n = 0 \end{aligned} \tag{49}$$

for any numerical values of $\xi_1, \xi_2, \dots \xi_n$ which satisfy eqs. (38). This equation (49) will therefore hold true, if for r of the letters $\xi, \xi_2, \dots \xi_n$ we substitute their values in terms of the others taken from eqs. (38), and therefore it will hold true when we use

$\mathfrak{S}_1, \mathfrak{S}_2, \dots \mathfrak{S}_n$, as before, to denote the units of the various components. Thus understood, the equation expresses that the values of the quantities in the parentheses are such as are consistent with the equations of condition (41). The change in the system, therefore, which we are considering, is not one which violates any of the equations of condition, and as (45) does not hold true for this change, and for all values of $T, P, M_1, M_2, \dots M_n$ which are consistent with eqs. (43), the state of the system cannot be one of equilibrium. Therefore it is necessary, and it is evidently sufficient for equilibrium, that it shall be possible to assign to $T, P, M_1, M_2, \dots M_n$ such values, consistent with eqs. (43), that the condition (45) shall hold true for any change in the system irrespective of the equations of condition (39), (40), (41).

For this it is necessary and sufficient that

$$t = T, \quad p = P, \quad (50)$$

$$\mu_1 \delta m_1 \geq M_1 \delta m_1, \quad \mu_2 \delta m_2 \geq M_2 \delta m_2, \quad \dots \quad \mu_n \delta m_n \geq M_n \delta m_n \quad (51)$$

for each of the *original parts* as previously defined, and that

$$D\varepsilon - T D\eta + P Dv - M_1 Dm_1 - M_2 Dm_2 \dots - M_n Dm_n \geq 0, \quad (52)$$

for each of the *new parts* as previously defined. If to these conditions we add equations (43), we may treat $T, P, M_1, M_2, \dots M_n$ simply as unknown quantities to be eliminated.

In regard to conditions (51), it will be observed that if a substance, \mathcal{S}_1 , is an actual component of the part of the given mass distinguished by a single accent, $\delta m_1'$ may be either positive or negative, and we shall have $\mu_1' = M_1$; but if \mathcal{S}_1 is only a possible component of that part, $\delta m_1'$ will be incapable of a negative value, and we will have $\mu_1' \geq M_1$.

The formulæ (50), (51), and (43) express the same particular conditions of equilibrium which we have before obtained by a less general process. It remains to discuss (52). This formula must hold true of any infinitesimal mass in the system in its varied state which is not approximately homogeneous with any of the surrounding masses, the expressions $D\varepsilon, D\eta, Dv, Dm_1, Dm_2, \dots Dm_n$ denoting the energy, entropy, and volume of this infinitesimal mass, and the quantities of the substances $\mathcal{S}_1, \mathcal{S}_2, \dots \mathcal{S}_n$ which we regard as composing it, (not necessarily as *independently* variable components). If there is more than one way in which this mass may be considered as composed of these substances, we may choose whichever is most convenient. Indeed it follows directly from the relations existing between M_1, M_2, \dots and M_n that the result would be the same in

any case. Now, if we assume that the values of $D\varepsilon$, $D\eta$, Dv , Dm_1 , Dm_2 , . . . Dm_n are proportional to the values of ε , η , v , m_1 , m_2 , . . . m_n for any large homogeneous mass of similar composition, and of the same temperature and pressure, the condition is equivalent to this, that

$$\varepsilon - T\eta + Pv - M_1 m_1 - M_2 m_2 . . . - M_n m_n \geq 0 \quad (53)$$

for any large homogeneous body which can be formed out of the substances S_1 , S_2 . . . S_n .

But the validity of this last transformation cannot be admitted without considerable limitation. It is assumed that the relation between the energy, entropy, volume, and the quantities of the different components of a very small mass surrounded by substances of different composition and state is the same as if the mass in question formed a part of a large homogeneous body. We started, indeed, with the assumption that we might neglect the part of the energy, etc., depending upon the surfaces separating heterogeneous masses. Now, in many cases, and for many purposes, as, in general, when the masses are large, such an assumption is quite legitimate, but in the case of these masses which are formed within or among substances of different nature or state, and which at their first formation must be infinitely small, the same assumption is evidently entirely inadmissible, as the surfaces must be regarded as infinitely large in proportion to the masses. We shall see hereafter what modifications are necessary in our formulæ in order to include the parts of the energy, etc., which are due to the surfaces, but this will be on the assumption, which is usual in the theory of capillarity, that the radius of curvature of the surfaces is large in proportion to the radius of sensible molecular action, and also to the thickness of the lamina of matter at the surface which is not (sensibly) homogeneous in all respects with either of the masses which it separates. But although the formulæ thus modified will apply with sensible accuracy to masses (occurring within masses of a different nature) much smaller than if the terms relating to the surfaces were omitted, yet their failure when applied to masses infinitely small in all their dimensions is not less absolute.

Considerations like the foregoing might render doubtful the validity even of (52) as the necessary and sufficient condition of equilibrium in regard to the formation of masses not approximately homogeneous with those previously existing, when the conditions of equilibrium between the latter are satisfied, unless it is shown that in establishing this formula there have been no quantities neglected relating to the

mutual action of the new and the original parts, which can affect the result. It will be easy to give such a meaning to the expressions $D\varepsilon$, $D\eta$, Dv , Dm_1 , Dm_2 , . . . Dm_n that this shall be evidently the case. It will be observed that the quantities represented by these expressions have not been perfectly defined. In the first place, we have no right to assume the existence of any surface of absolute discontinuity to divide the new parts from the original, so that the position given to the dividing surface is to a certain extent arbitrary. Even if the surface separating the masses were determined, the energy to be attributed to the masses separated would be partly arbitrary, since a part of the total energy depends upon the mutual action of the two masses. We ought perhaps to consider the case the same in regard to the entropy, although the entropy of a system never depends upon the mutual relations of parts at sensible distances from one another. Now the condition (52) will be valid if the quantities $D\varepsilon$, $D\eta$, Dv , Dm_1 , Dm_2 . . . Dm_n are so defined that none of the assumptions which have been made, tacitly or otherwise, relating to the formation of these new parts, shall be violated. These assumptions are the following:—that the relation between the variations of the energy, entropy, volume, etc., of any of the original parts is not affected by the vicinity of the new parts; and that the energy, entropy, volume, etc., of the system in its varied state are correctly represented by the sums of the energies, entropies, volumes, etc., of the various parts (original and new), so far at least as any of these quantities are determined or affected by the formation of the new parts. We will suppose $D\varepsilon$, $D\eta$, Dv , Dm_1 , Dm_2 . . . Dm_n to be so defined that these conditions shall not be violated. This may be done in various ways. We may suppose that the position of the surfaces separating the new and the original parts has been fixed in any suitable way. This will determine the space and the matter belonging to the parts separated. If this does not determine the division of the entropy, we may suppose this determined in any suitable arbitrary way. Thus we may suppose the total energy in and about any new part to be so distributed that equation (12) as applied to the original parts shall not be violated by the formation of the new parts. Or, it may seem more simple to suppose that the imaginary surface which divides any new part from the original is so placed as to include all the matter which is affected by the vicinity of the new formation, so that the part or parts which we regard as original may be left homogeneous in the strictest sense, including uniform *densities of energy and entropy*, up to the very

bounding surface. The homogeneity of the new parts is of no consequence, as we have made no assumption in that respect. It may be doubtful whether we can consider the new parts, *as thus bounded*, to be infinitely small even in their earliest stages of development. But if they are not infinitely small, the only way in which this can affect the validity of our formulæ will be that in virtue of the equations of condition, i. e., in virtue of the evident necessities of the case, finite variations of the energy, entropy, volume, etc., of the original parts will be caused, to which it might seem that equation (12) would not apply. But if the nature and state of the mass be not varied, equation (12) will hold true of finite differences. (This appears at once, if we integrate the equation under the above limitation.) Hence, the equation will hold true for finite differences, provided that the nature and state of the mass be infinitely little varied. For the differences may be considered as made up of two parts, of which the first are for a constant nature and state of the mass, and the second are infinitely small. We may therefore regard the new parts to be bounded as supposed without prejudice to the validity of any of our results.

The condition (52) understood in either of these ways (or in others which will suggest themselves to the reader) will have a perfectly definite meaning, and will be valid as the necessary and sufficient condition of equilibrium in regard to the formation of new parts, when the conditions of equilibrium in regard to the original parts, (50), (51), and (43), are satisfied.

In regard to the condition (53), it may be shown that with (50), (51), and (43) it is always sufficient for equilibrium. To prove this, it is only necessary to show that when (50), (51), and (43) are satisfied, and (52) is not, (53) will also not be satisfied.

We will first observe that an expression of the form

$$-\varepsilon + T\eta - Pv + M_1 m_1 + M_2 m_2 \dots + M_n m_n \quad (54)$$

denotes the work obtainable by the formation (by a reversible process) of a body of which $\varepsilon, \eta, v, m_1, m_2, \dots, m_n$ are the energy, entropy, volume, and the quantities of the components, within a medium having the pressure P , the temperature T , and the potentials M_1, M_2, \dots, M_n . (The medium is supposed so large that its properties are not sensibly altered in any part by the formation of the body.) For ε is the energy of the body formed, and the remaining terms represent (as may be seen by applying equation (12) to the medium) the decrease of the energy of the medium, if, after the

formation of the body, the joint entropy of the medium and the body, their joint volumes and joint quantities of matter, were the same as the entropy, etc., of the medium before the formation of the body. This consideration may convince us that for any given finite values of v and of T, P, M_1 , etc. this expression cannot be infinite when ε, η, m_1 , etc. are determined by any real body, whether homogeneous or not, (but of the given volume), even when T, P, M_1 , etc. do not represent the values of the temperature, pressure, and potentials of any real substance. (If the substances $S_1, S_2, \dots S_n$ are all actual components of any homogeneous part of the system of which the equilibrium is discussed, that part will afford an example of a body having the temperature, pressure, and potentials of the medium supposed.)

Now by integrating equation (12) on the supposition that the nature and state of the mass considered remain unchanged, we obtain the equation

$$\varepsilon = t\eta - pv + \mu_1 m_1 + \mu_2 m_2 \dots + \mu_n m_n \quad (55)$$

which will hold true of any homogeneous mass whatever. Therefore for any one of the original parts, by (50) and (51),

$$\varepsilon - T\eta + Pv - M_1 m_1 - M_2 m_2 \dots - M_n m_n = 0. \quad (56)$$

If the condition (52) is not satisfied in regard to all possible new parts, let N be a new part occurring in an original part O , for which the condition is not satisfied. It is evident that the value of the expression

$$\varepsilon - T\eta + Pv - M_1 m_1 - M_2 m_2 \dots - M_n m_n \quad (57)$$

applied to a mass like O including some very small masses like N , will be negative, and will decrease if the number of these masses like N is increased, until there remains within the whole mass no portion of any sensible size without these masses like N , which, it will be remembered, have no sensible size. But it cannot decrease without limit, as the value of (54) cannot become infinite. Now we need not inquire whether the least value of (57) (for constant values of $T, P, M_1, M_2, \dots M_n$) would be obtained by excluding entirely the mass like O , and filling the whole space considered with masses like N , or whether a certain mixture would give a smaller value,—it is certain that the least possible value of (57) per unit of volume, and that a negative value, will be realized by a mass having a certain homogeneity. If the new part N for which the condition (52) is not satisfied occurs between two different original parts O' and O'' , the argument need not be essentially varied. We may consider the

value of (57), for a body consisting of masses like O' and O'' separated by a lamina N . This value may be decreased by increasing the extent of this lamina, which may be done within a given volume by giving it a convoluted form; and it will be evident, as before, that the least possible value of (57) will be for a homogeneous mass, and that the value will be negative. And such a mass will be not merely an ideal combination, but a body capable of existing, for as the expression (57) has for this mass in the state considered its least possible value per unit of volume, the energy of the mass included in a unit of volume is the least possible for the same matter with the same entropy and volume,—hence, if confined in a non-conducting vessel, it will be in a state of not unstable equilibrium. Therefore when (50), (51), and (43) are satisfied, if the condition (52) is not satisfied in regard to all possible new parts, there will be some homogeneous body which can be formed out of the substances $S_1, S_2, \dots S_n$ which will not satisfy condition (53).

Therefore, if the initially existing masses satisfy the conditions (50), (51), and (43), and condition (53) is satisfied by every homogeneous body which can be formed out of the given matter, there will be equilibrium.

On the other hand, (53) is not a necessary condition of equilibrium. For we may easily conceive that the condition (52) shall hold true (for any very small formations within or between any of the given masses), while the condition (53) is not satisfied (for all large masses formed of the given matter), and experience shows that this is very often the case. Supersaturated solutions, superheated water, etc., are familiar examples. Such an equilibrium will, however, be *practically* unstable. By this is meant that, although, strictly speaking, an infinitely small disturbance or change may not be sufficient to destroy the equilibrium, yet a very small change in the initial state, perhaps a circumstance which entirely escapes our powers of perception, will be sufficient to do so. The presence of a small portion of the substance for which the condition (53) does not hold true, is sufficient to produce this result, when this substance forms a variable component of the original homogeneous masses. In other cases, when, if the new substances are formed at all, different kinds must be formed simultaneously, the initial presence of the different kinds, and that in immediate proximity, may be necessary.

It will be observed, that from (56) and (53) we can at once obtain (50) and (51), viz., by applying (53) to bodies differing infinitely little from the various homogeneous parts of the given mass. There-

fore, the condition (56) (relating to the various homogeneous parts of the given mass) and (53) (relating to any bodies which can be formed of the given matter) with (43) are always sufficient for equilibrium, and always necessary for an equilibrium which shall be practically stable. And, if we choose, we may get rid of limitation in regard to equations (43). For, if we compare these equations with (38), it is easy to see that it is always immaterial, in applying the tests (56) and (53) to any body, how we consider it to be composed. Hence, in applying these tests, we may consider all bodies to be composed of the *ultimate* components of the given mass. Then the terms in (56) and (53) which relate to other components than these will vanish, and we need not regard the equations (43). Such of the constants M_1, M_2, \dots, M_n as relate to the ultimate components, may be regarded, like T and P , as unknown quantities subject only to the conditions (56) and (53).

These two conditions, which are sufficient for equilibrium and necessary for a practically stable equilibrium, may be united in one, viz., (if we choose the ultimate components of the given mass for the component substances to which m_1, m_2, \dots, m_n relate) that it shall be possible to give such values to the constants $T, P, M_1, M_2, \dots, M_n$ in the expression (57) that the value of the expression for each of the homogeneous parts of the mass in question shall be as small as for any body whatever made of the same components.

Effect of Solidity of any Part of the given Mass.

If any of the homogeneous masses of which the equilibrium is in question are solid, it will evidently be proper to treat the proportion of their components as invariable in the application of the criterion of equilibrium, even in the case of *compounds of variable proportions*, i. e., even when bodies can exist which are compounded in proportions infinitesimally varied from those of the solids considered. (Those solids which are capable of absorbing fluids form of course an exception, so far as their fluid components are concerned.) It is true that a solid may be increased by the formation of new solid matter on the surface where it meets a fluid, which is not homogeneous with the previously existing solid, but such a deposit will properly be treated as a distinct part of the system, (viz., as one of the parts which we have called *new*). Yet it is worthy of notice that if a homogeneous solid which is a compound of variable proportions is in contact and equilibrium with a fluid, and the actual components of the solid (considered as of variable composition) are also actual com-

ponents of the fluid, and the condition (53) is satisfied in regard to all bodies which can be formed out of the actual components of the fluid, (which will always be the case unless the fluid is practically unstable,) all the conditions will hold true of the solid, which would be necessary for equilibrium if it were fluid.

This follows directly from the principles stated on the preceding pages. For in this case the value of (57) will be zero as determined either for the solid or for the fluid considered with reference to their ultimate components, and will not be negative for any body whatever which can be formed of these components; and these conditions are sufficient for equilibrium independently of the solidity of one of the masses. Yet the point is perhaps of sufficient importance to demand a more detailed consideration.

Let $S_a, \dots S_g$ be the actual components of the solid, and $S_h, \dots S_k$ its possible components (which occur as actual components in the fluid); then, considering the proportion of the components of the solid as variable, we shall have for this body by equation (12)

$$d\varepsilon' = t d\eta' - p' dv' + \mu'_a dm'_a \dots + \mu'_g dm'_g + \mu'_h dm'_h \dots + \mu'_k dm'_k. \tag{58}$$

By this equation the potentials $\mu'_a \dots \mu'_k$ are perfectly defined. But the differentials $dm'_a \dots dm'_k$, considered as independent, evidently express variations which are not *possible* in the sense required in the criterion of equilibrium. We might, however, introduce them into the general condition of equilibrium, if we should express the dependence between them by the proper equations of condition. But it will be more in accordance with our method hitherto, if we consider the solid to have only a single independently variable component S_x , of which the nature is represented by the solid itself. We may then write

$$\delta\varepsilon' = t' \delta\eta' - p' \delta v' + \mu'_x \delta m'_x. \tag{59}$$

In regard to the relation of the potential μ'_x to the potentials occurring in equation (58) it will be observed, that as we have by integration of (58) and (59)

$$\varepsilon' = t' \eta' - p' v' + \mu'_a m'_a \dots + \mu'_g m'_g, \tag{60}$$

and
$$\varepsilon' = t' \eta' - p' v' + \mu'_x m'_x; \tag{61}$$

therefore
$$\mu'_x m'_x = \mu'_a m'_a \dots + \mu'_g m'_g. \tag{62}$$

Now, if the fluid has besides $S_a \dots S_g$ and $S_h \dots S_k$ the actual components $S_l \dots S_n$, we may write for the fluid

$$\begin{aligned} \delta \varepsilon'' = t'' \delta \eta'' - p'' \delta v'' + \mu_a'' \delta m_a'' \dots + \mu_g'' \delta m_g'' \\ + \mu_h'' \delta m_h'' \dots + \mu_k'' \delta m_k'' + \mu_l'' \delta m_l'' \dots + \mu_n'' \delta m_n'', \end{aligned} \quad (63)$$

and as by supposition

$$m_x' \mathcal{E}_x = m_a' \mathcal{E}_a \dots + m_g' \mathcal{E}_g \quad (64)$$

equations (43), (50), and (51) will give in this case on elimination of the constants T, P , etc.,

$$t' = t'', \quad p' = p'', \quad (65)$$

and

$$m_x' \mu_x' = m_a' \mu_a'' \dots + m_g' \mu_g''. \quad (66)$$

Equations (65) and (66) may be regarded as expressing the conditions of equilibrium between the solid and the fluid. The last condition may also, in virtue of (62), be expressed by the equation

$$m_a' \mu_a' \dots + m_g' \mu_g' = m_a' \mu_a'' \dots + m_g' \mu_g''. \quad (67)$$

But if condition (53) holds true of all bodies which can be formed of $S_a \dots S_g, S_h, \dots S_k, S_l \dots S_n$, we may write for all such bodies

$$\begin{aligned} \varepsilon - t'' \eta + p'' v - \mu_a'' m_a \dots - \mu_g'' m_g - \mu_h'' m_h \\ \dots - \mu_k'' m_k - \mu_l'' m_l \dots - \mu_n'' m_n \geq 0. \end{aligned} \quad (68)$$

(In applying this formula to various bodies, it is to be observed that only the values of the unaccented letters are to be determined by the different bodies to which it is applied, the values of the accented letters being already determined by the given fluid.) Now, by (60), (65), and (67), the value of the first member of this condition is zero when applied to the solid in its given state. As the condition must hold true of a body differing infinitesimally from the solid, we shall have

$$\begin{aligned} d\varepsilon' - t'' d\eta' + p'' dv' - \mu_a'' dm_a' \dots - \mu_g'' dm_g' \\ - \mu_h'' dm_h' \dots - \mu_k'' dm_k' \geq 0, \end{aligned} \quad (69)$$

or, by equations (58) and (65),

$$\begin{aligned} (\mu_a' - \mu_a'') dm_a' \dots + (\mu_g' - \mu_g'') dm_g' \\ + (\mu_h' - \mu_h'') dm_h' \dots + (\mu_k' - \mu_k'') dm_k' \geq 0. \end{aligned} \quad (70)$$

Therefore, as these differentials are all independent,

$$\mu_a' = \mu_a'', \dots \mu_g' = \mu_g'', \quad \mu_h' \geq \mu_h'', \dots \mu_k' \geq \mu_k''; \quad (71)$$

which with (65) are evidently the same conditions which we would have obtained if we had neglected the fact of the solidity of one of the masses.

We have supposed the solid to be homogeneous. But it is evident that in any case the above conditions must hold for every separate point where the solid meets the fluid. Hence, the temperature and pressure and the potentials for all the actual components of the solid must have a constant value in the solid at the surface where it meets the fluid. Now, these quantities are determined by the nature and state of the solid, and exceed in number the independent variations of which its nature and state are capable. Hence, if we reject as improbable the supposition that the nature or state of a body can vary without affecting the value of any of these quantities, we may conclude that a solid which varies (continuously) in nature or state at its surface cannot be in equilibrium with a stable fluid which contains, as independently variable components, the variable components of the solid. (There may be, however, in equilibrium with the same stable fluid, a *finite* number of different solid bodies, composed of the variable components of the fluid, and having their nature and state completely determined by the fluid.)*

Effect of Additional Equations of Condition.

As the equations of condition, of which we have made use, are such as always apply to matter enclosed in a rigid, impermeable, and non-conducting envelop, the particular conditions of equilibrium which we have found will always be sufficient for equilibrium. But the number of conditions necessary for equilibrium, will be diminished, in a case otherwise the same, as the number of equations of condition is increased. Yet the problem of equilibrium which has been treated will sufficiently indicate the method to be pursued in all cases and the general nature of the results.

It will be observed that the position of the various homogeneous parts of the given mass, which is otherwise immaterial, may determine the existence of certain equations of condition. Thus, when different parts of the system in which a certain substance is a variable component are entirely separated from one another by parts of which this substance is not a component, the quantity of this substance will be invariable for each of the parts of the system which are thus separated, which will be easily expressed by equations of condition. Other equations of condition may arise from the passive forces (or resistances to change) inherent in the given masses. In the prob-

* The solid has been considered as subject only to isotropic stresses. The effect of other stresses will be considered hereafter.

lem which we are next to consider there are equations of condition due to a cause of a different nature.

Effect of a Diaphragm (Equilibrium of Osmotic Forces).

If the given mass, enclosed as before, is divided into two parts, each of which is homogeneous and fluid, by a diaphragm which is capable of supporting an excess of pressure on either side, and is permeable to some of the components and impermeable to others, we shall have the equations of condition

$$\delta\eta' + \delta\eta'' = 0, \quad (72)$$

$$\delta v' = 0, \quad \delta v'' = 0, \quad (73)$$

and for the components which cannot pass the diaphragm

$$\delta m_a' = 0, \quad \delta m_a'' = 0, \quad \delta m_b' = 0, \quad \delta m_b'' = 0, \quad \text{etc.}, \quad (74)$$

and for those which can

$$\delta m_h' + \delta m_h'' = 0, \quad \delta m_i' + \delta m_i'' = 0, \quad \text{etc.} \quad (75)$$

With these equations of condition, the general condition of equilibrium (see (15)) will give the following particular conditions:

$$t' = t'', \quad (76)$$

and for the components which can pass the diaphragm, if actual components of both masses,

$$\mu_h' = \mu_h'', \quad \mu_i' = \mu_i'', \quad \text{etc.}, \quad (77)$$

but not

$$p' = p'',$$

nor

$$\mu_a' = \mu_a'', \quad \mu_b' = \mu_b'', \quad \text{etc.}$$

Again, if the diaphragm is permeable to the components in certain proportions only, or in proportions not entirely determined yet subject to certain conditions, these conditions may be expressed by equations of condition, which will be linear equations between $\delta m_1'$, $\delta m_2'$, etc., and if these be known the deduction of the particular conditions of equilibrium will present no difficulties. We will however observe that if the components S_1 , S_2 , etc. (being actual components on each side) can pass the diaphragm simultaneously in the proportions a_1 , a_2 , etc. (without other resistances than such as vanish with the velocity of the current), values proportional to a_1 , a_2 , etc. are possible for $\delta m_1'$, $\delta m_2'$, etc. in the general condition of equilibrium, $\delta m_1''$, $\delta m_2''$, etc. having the same values taken negatively, so that we shall have for one particular condition of equilibrium

$$a_1 \mu_1' + a_2 \mu_2' + \text{etc.} = a_1 \mu_1'' + a_2 \mu_2'' + \text{etc.} \quad (78)$$

There will evidently be as many independent equations of this form

as there are independent combinations of the elements which can pass the diaphragm.

These conditions of equilibrium do not of course depend in any way upon the supposition that the volume of each fluid mass is kept constant, if the diaphragm is in any case supposed immovable. In fact, we may easily obtain the same conditions of equilibrium, if we suppose the volumes variable. In this case, as the equilibrium must be preserved by forces acting upon the external surfaces of the fluids, the variation of the energy of the sources of these forces must appear in the general condition of equilibrium, which will be

$$\delta \epsilon' + \delta \epsilon'' + P' \delta v' + P'' \delta v'' \geq 0, \tag{79}$$

P' and P'' denoting the external forces per unit of area. (Compare (14).) From this condition we may evidently derive the same internal conditions of equilibrium as before, and in addition the external conditions

$$p' = P', \quad p'' = P''. \tag{80}$$

In the preceding paragraphs it is assumed that the permeability of the diaphragm is perfect, and its impermeability absolute, i. e., that it offers no resistance to the passage of the components of the fluids in certain proportions, except such as vanishes with the velocity, and that in other proportions the components cannot pass at all. How far these conditions are satisfied in any particular case is of course to be determined by experiment.

If the diaphragm is permeable to all the n components without restriction, the temperature and the potentials for all the components must be the same on both sides. Now, as one may easily convince himself, a mass having n components is capable of only $n+1$ independent variations in nature and state. Hence, if the fluid on one side of the diaphragm remains without change, that on the other side cannot (in general) vary in nature or state. Yet the pressure will not necessarily be the same on both sides. For, although the pressure is a function of the temperature and the n potentials, it may be a many-valued function (or any one of several functions) of these variables. But when the pressures are different on the two sides, the fluid which has the less pressure will be *practically unstable*, in the sense in which the term has been used on page 133. For

$$\epsilon'' - \epsilon' + p'' v'' - \mu_1'' m_1'' - \mu_2'' m_2'' \dots - \mu_n'' m_n'' = 0, \tag{81}$$

as appears from equation (12) if integrated on the supposition that the nature and state of the mass remain unchanged. Therefore, if $p' < p''$ while $\epsilon' = \epsilon''$, $\mu_1' = \mu_1''$, etc.,

$$\varepsilon'' - t' \eta'' + p' v'' - \mu_1' m_1'' - \mu_2' m_2'' \dots - \mu_n' m_n'' < 0. \quad (82)$$

This relation indicates the instability of the fluid to which the single accents refer. (See page 133.)

But independently of any assumption in regard to the permeability of the diaphragm, the following relation will hold true in any case in which each of the two fluid masses may be regarded as uniform throughout in nature and state. Let the character D be used with the variables which express the nature, state, and quantity of the fluids to denote the increments of the values of these quantities actually occurring in a time either finite or infinitesimal. Then, as the heat received by the two masses cannot exceed $t' D\eta' + t'' D\eta''$, and as the increase of their energy is equal to the difference of the heat they receive and the work they do,

$$D\varepsilon' + D\varepsilon'' \leq t' D\eta' + t'' D\eta'' - p' Dv' - p'' Dv'', \quad (83)$$

i. e., by (12),

$$\mu_1' Dm_1' + \mu_1'' Dm_1'' + \mu_2' Dm_2' + \mu_2'' Dm_2'' + \text{etc.} \leq 0, \quad (84)$$

or

$$(\mu_1'' - \mu_1') Dm_1'' + (\mu_2'' - \mu_2') Dm_2'' + \text{etc.} \leq 0. \quad (85)$$

It is evident that the sign $=$ holds true only in the limiting case in which no motion takes place.

DEFINITION AND PROPERTIES OF FUNDAMENTAL EQUATIONS.

The solution of the problems of equilibrium which we have been considering has been made to depend upon the equations which express the relations between the energy, entropy, volume, and the quantities of the various components, for homogeneous combinations of the substances which are found in the given mass. The nature of such equations must be determined by experiment. As, however, it is only *differences* of energy and of entropy that can be measured, or indeed, that have a physical meaning, the values of these quantities are so far arbitrary, that we may choose independently for each simple substance the state in which its energy and its entropy are both zero. The values of the energy and the entropy of any compound body in any particular state will then be fixed. Its energy will be the sum of the work and heat expended in bringing its components from the states in which their energies and their entropies are zero into combination and to the state in question; and its entropy is the value of the integral $\int \frac{dQ}{t}$ for any *reversible* process

by which that change is effected (dQ denoting an element of the heat communicated to the matter thus treated, and t the temperature of the matter receiving it). In the determination both of the energy and of the entropy, it is understood that at the close of the process, all bodies which have been used, other than those to which the determinations relate, have been restored to their original state, with the exception of the sources of the work and heat expended, which must be used only as such sources.

We know, however, *a priori*, that if the quantity of any homogeneous mass containing n independently variable components varies and not its nature or state, the quantities $\varepsilon, \eta, v, m_1, m_2, \dots, m_n$ will all vary in the same proportion; therefore it is sufficient if we learn from experiment the relation between all but any one of these quantities for a given constant value of that one. Or, we may consider that we have to learn from experiment the relation subsisting between the $n + 2$ ratios of the $n + 3$ quantities $\varepsilon, \eta, v, m_1, m_2, \dots, m_n$. To fix our ideas we may take for these ratios $\frac{\varepsilon}{v}, \frac{\eta}{v}, \frac{m_1}{v}, \frac{m_2}{v}$, etc., that is, the separate densities of the components, and the ratios $\frac{\varepsilon}{v}$ and $\frac{\eta}{v}$, which may be called the *densities of energy and entropy*. But when there is but one component, it may be more convenient to choose $\frac{\varepsilon}{m}, \frac{\eta}{m}, \frac{v}{m}$ as the three variables. In any case, it is only a function of $n + 1$ independent variables, of which the form is to be determined by experiment.

Now if ε is a known function of $\eta, v, m_1, m_2, \dots, m_n$, as by equation (12)

$$d\varepsilon = t d\eta - p dv + \mu_1 dm_1 + \mu_2 dm_2 \dots + \mu_n dm_n, \quad (86)$$

$t, p, \mu_1, \mu_2, \dots, \mu_n$ are functions of the same variables, which may be derived from the original function by differentiation, and may therefore be considered as known functions. This will make $n + 3$ independent known relations between the $2n + 5$ variables, $\varepsilon, \eta, v, m_1, m_2, \dots, m_n, t, p, \mu_1, \mu_2, \dots, \mu_n$. These are all that exist, for of these variables, $n + 2$ are evidently independent. Now upon these relations depend a very large class of the properties of the compound considered,—we may say in general, all its thermal, mechanical, and chemical properties, so far as *active tendencies* are concerned, in cases in which the form of the mass does not require consideration. A single equation from which all these relations may

be deduced we will call a *fundamental equation* for the substance in question. We shall hereafter consider a more general form of the fundamental equation for solids, in which the pressure at any point is not supposed to be the same in all directions. But for masses subject only to isotropic stresses an equation between $\varepsilon, \eta, v, m_1, m_2, \dots, m_n$ is a fundamental equation. There are other equations which possess this same property.*

Let

$$\psi = \varepsilon - t \eta, \quad (87)$$

then by differentiation and comparison with (86) we obtain

$$d\psi = -\eta dt - p dv + \mu_1 dm_1 + \mu_2 dm_2 \dots + \mu_n dm_n. \quad (88)$$

If, then, ψ is known as a function of $t, v, m_1, m_2, \dots, m_n$, we can find $\eta, p, \mu_1, \mu_2, \dots, \mu_n$ in terms of the same variables. If we then substitute for ψ in our original equation its value taken from eq. (87), we shall have again $n + 3$ independent relations between the same $2n + 5$ variables as before.

Let

$$\chi = \varepsilon + p v, \quad (89)$$

then by (86),

$$d\chi = t d\eta + v dp + \mu_1 dm_1 + \mu_2 dm_2 \dots + \mu_n dm_n. \quad (90)$$

If, then, χ be known as a function of $\eta, p, m_1, m_2, \dots, m_n$, we can find $t, v, \mu_1, \mu_2, \dots, \mu_n$ in terms of the same variables. By eliminating χ , we may obtain again $n + 3$ independent relations between the same $2n + 5$ variables as at first.

Let

$$\zeta = \varepsilon - t \eta + p v, \quad (91)$$

then, by (86)

$$d\zeta = -\eta dt + v dp + \mu_1 dm_1 + \mu_2 dm_2 \dots + \mu_n dm_n. \quad (92)$$

If, then, ζ is known as a function of $t, p, m_1, m_2, \dots, m_n$, we can

* M. Massieu (Comptes Rendus, T. lxxix, 1869, p. 858 and p. 1057) has shown how all the properties of a fluid "which are considered in thermodynamics" may be deduced from a single function, which he calls a characteristic function of the fluid considered. In the papers cited, he introduces two different functions of this kind; viz., a function of the temperature and volume, which he denotes by ψ , the value of which in our notation would be $\frac{-\varepsilon + t\eta}{t}$ or $\frac{-\psi}{t}$; and a function of the temperature and pressure, which he denotes by ψ' , the value of which in our notation would be $\frac{-\varepsilon + t\eta - pv}{t}$ or $\frac{-\zeta}{t}$. In both cases he considers a constant quantity (one kilogram) of the fluid, which is regarded as invariable in composition.

find $\eta, v, \mu_1, \mu_2, \dots, \mu_n$ in terms of the same variables. By eliminating ζ , we may obtain again $n + 3$ independent relations between the same $2n + 5$ variables as at first.

If we integrate (86), supposing the quantity of the compound substance considered to vary from zero to any finite value, its nature and state remaining unchanged, we obtain

$$\varepsilon = t\eta - pv + \mu_1 m_1 + \mu_2 m_2 \dots + \mu_n m_n, \quad (93)$$

and by (87), (89), (91)

$$\psi = -pv + \mu_1 m_1 + \mu_2 m_2 \dots + \mu_n m_n, \quad (94)$$

$$\chi = t\eta + \mu_1 m_1 + \mu_2 m_2 \dots + \mu_n m_n, \quad (95)$$

$$\zeta = \mu_1 m_1 + \mu_2 m_2 \dots + \mu_n m_n. \quad (96)$$

The last three equations may also be obtained directly by integrating (88), (90), and (92).

If we differentiate (93) in the most general manner, and compare the result with (86), we obtain

$$-v dp + \eta dt + m_1 d\mu_1 + m_2 d\mu_2 \dots + m_n d\mu_n = 0, \quad (97)$$

or

$$dp = \frac{\eta}{v} dt + \frac{m_1}{v} d\mu_1 + \frac{m_2}{v} d\mu_2 \dots + \frac{m_n}{v} d\mu_n = 0. \quad (98)$$

Hence, there is a relation between the $n + 2$ quantities $t, p, \mu_1, \mu_2, \dots, \mu_n$, which, if known, will enable us to find in terms of these quantities all the ratios of the $n + 2$ quantities $\eta, v, m_1, m_2 \dots m_n$. With (93), this will make $n + 3$ independent relations between the same $2n + 5$ variables as at first.

Any equation, therefore, between the quantities

$$\varepsilon, \quad \eta, \quad v, \quad m_1, \quad m_2, \dots, m_n, \quad (99)$$

$$\text{or} \quad \psi, \quad t, \quad v, \quad m_1, \quad m_2, \dots, m_n, \quad (100)$$

$$\text{or} \quad \chi, \quad \eta, \quad p, \quad m_1, \quad m_2, \dots, m_n, \quad (101)$$

$$\text{or} \quad \zeta, \quad t, \quad p, \quad m_1, \quad m_2, \dots, m_n, \quad (102)$$

$$\text{or} \quad t, \quad p, \quad \mu_1, \quad \mu_2, \dots, \mu_n, \quad (103)$$

is a fundamental equation, and any such is entirely equivalent to any other.* For any homogeneous mass whatever, considered (in general) as variable in composition, in quantity, and in thermodynamic state, and having n independently variable components, to which

* The distinction between equations which are, and which are not, *fundamental*, in the sense in which the word is here used, may be illustrated by comparing an equation

the subscript numerals refer, (but not excluding the case in which $n = 1$ and the composition of the body is invariable,) there is a relation between the quantities enumerated in any one of the above sets, from which, if known, with the aid only of *general* principles and relations, we may deduce all the relations subsisting for such a mass between the quantities $\varepsilon, \psi, \chi, \zeta, \eta, v, m_1, m_2, \dots, m_n, t, p, \mu_1, \mu_2, \dots, \mu_n$. It will be observed that, besides the equations which define ψ, χ , and ζ , there is one finite equation, (93), which subsists between these quantities independently of the form of the fundamental equation.

Other sets of quantities might of course be added which possess the same property. The sets (100), (101), (102) are mentioned on account of the important properties of the quantities ψ, χ, ζ , and because the equations (88), (90), (92), like (86), afford convenient definitions of the potentials, viz.,

$$\mu_1 = \left(\frac{d\varepsilon}{dm_1} \right)_{\eta, v, m} = \left(\frac{d\psi}{dm_1} \right)_{t, v, m} = \left(\frac{d\chi}{dm_1} \right)_{\eta, p, m} = \left(\frac{d\zeta}{dm_1} \right)_{t, p, m} \quad (104)$$

etc., where the subscript letters denote the quantities which remain constant in the differentiation, m being written for brevity for all the letters m_1, m_2, \dots, m_n except the one occurring in the denominator. It will be observed that the quantities in (103) are all independent of the quantity of the mass considered, and are those which must, in general, have the same value in contiguous masses in equilibrium.

On the quantities ψ, χ, ζ .

The quantity ψ has been defined for any homogeneous mass by the equation

$$\psi = \varepsilon - t\eta. \quad (105)$$

between

$$\varepsilon, \eta, v, m_1, m_2, \dots, m_n$$

with one between

$$\varepsilon, t, v, m_1, m_2, \dots, m_n.$$

As, by (86),

$$t = \left(\frac{d\varepsilon}{d\eta} \right)_{vm},$$

the second equation may evidently be derived from the first. But the first equation cannot be derived from the second; for an equation between

$$\varepsilon, \left(\frac{d\varepsilon}{d\eta} \right)_{vm}, v, m_1, m_2, \dots, m_n$$

is equivalent to one between

$$\left(\frac{d\eta}{d\varepsilon} \right)_{vm}, \varepsilon, v, m_1, m_2, \dots, m_n.$$

which is evidently not sufficient to determine the value of η in terms of the other variables.

We may extend this definition to any material system whatever which has a uniform temperature throughout.

If we compare two states of the system of the same temperature, we have

$$\psi' - \psi'' = \varepsilon' - \varepsilon'' - t(\eta' - \eta''). \quad (106)$$

If we suppose the system brought from the first to the second of these states without change of temperature and by a reversible process in which W is the work done and Q the heat received by the system, then

$$\varepsilon' - \varepsilon'' = W - Q, \quad (107)$$

and
$$t(\eta' - \eta'') = Q. \quad (108)$$

Hence

$$\psi' - \psi'' = W; \quad (109)$$

and for an infinitely small reversible change in the state of the system, in which the temperature remains constant, we may write

$$-d\psi = dW. \quad (110)$$

Therefore, $-\psi$ is the force function of the system for constant temperature, just as $-\varepsilon$ is the force function for constant entropy. That is, if we consider ψ as a function of the temperature and the variables which express the distribution of the matter in space, for every different value of the temperature $-\psi$ is the different force function required by the system if maintained at that special temperature.

From this we may conclude that when a system has a uniform temperature throughout, the additional conditions which are necessary and sufficient for equilibrium may be expressed by

$$(\delta\psi)_t \geq 0. * \quad (111)$$

* This general condition of equilibrium might be used instead of (2) in such problems of equilibrium as we have considered and others which we shall consider hereafter with evident advantage in respect to the brevity of the formulæ, as the limitation expressed by the subscript t in (111) applies to every part of the system taken separately, and diminishes by one the number of independent variations in the state of these parts which we have to consider. The more cumbersome course adopted in this paper has been chosen, among other reasons, for the sake of deducing *all* the particular conditions of equilibrium from one general condition, and of having the quantities mentioned in this general condition such as are most generally used and most simply defined; and because in the longer formulæ as given, the reader will easily see in each case the form which they would take if we should adopt (111) as the general condition of equilibrium, which would be in effect to take the thermal condition of equilibrium for granted, and to seek only the remaining conditions. For example, in the problem treated on pages 116 ff., we would obtain from (111) by (88) a condition precisely like (15), except that the terms $t\delta\eta'$, $t\delta\eta''$, etc. would be wanting.

When it is not possible to bring the system from one to the other of the states to which ψ' and ψ'' relate by a reversible process without altering the temperature, it will be observed that it is not necessary for the validity of (107)–(109) that the temperature of the system should remain constant during the reversible process to which W and Q relate, provided that the only source of heat or cold used has the same temperature as the system in its initial or final state. Any external bodies may be used in the process in any way not affecting the condition of reversibility, if restored to their original condition at the close of the process; nor does the limitation in regard to the use of heat apply to such heat as may be restored to the source from which it has been taken.

It may be interesting to show directly the equivalence of the conditions (111) and (2) when applied to a system of which the temperature in the given state is uniform throughout.

If there are any variations in the state of such a system which do not satisfy (2), then for these variations

$$\delta\varepsilon < 0 \quad \text{and} \quad \delta\eta = 0.$$

If the temperature of the system in its varied state is not uniform, we may evidently increase its entropy without altering its energy by supposing heat to pass from the warmer to the cooler parts. And the state having the greatest entropy for the energy $\varepsilon + \delta\varepsilon$ will necessarily be a state of uniform temperature. For this state (regarded as a variation from the original state)

$$\delta\varepsilon < 0 \quad \text{and} \quad \delta\eta > 0.$$

Hence, as we may diminish both the energy and the entropy by cooling the system, there must be a state of uniform temperature for which (regarded as a variation of the original state)

$$\delta\varepsilon < 0 \quad \text{and} \quad \delta\eta = 0.$$

From this we may conclude that for systems of initially uniform temperature condition (2) will not be altered if we limit the variations to such as do not disturb the uniformity of temperature.

Confining our attention, then, to states of uniform temperature, we have by differentiation of (105)

$$\delta\varepsilon - t \delta\eta = \delta\psi + \eta \delta t. \tag{112}$$

Now there are evidently changes in the system (produced by heating or cooling) for which

$$\delta\varepsilon - t \delta\eta = 0 \quad \text{and therefore} \quad \delta\psi + \eta \delta t = 0, \tag{113}$$

neither $\delta\eta$ nor δt having the value zero. This consideration is sufficient to show that the condition (2) is equivalent to

$$\delta\varepsilon - t\delta\eta \geq 0. \tag{114}$$

and that the condition (111) is equivalent to

$$\delta\psi + \eta\delta t \geq 0 \tag{115}$$

and by (112) the two last conditions are equivalent.

In such cases as we have considered on pages 115–137, in which the form and position of the masses of which the system is composed is immaterial, uniformity of temperature and pressure are always necessary for equilibrium, and the remaining conditions, when these are satisfied, may be conveniently expressed by means of the function ζ , which has been defined for a homogeneous mass on page 142, and which we will here define for any mass of uniform temperature and pressure by the same equation

$$\zeta = \varepsilon - t\eta + pv. \tag{116}$$

For such a mass, the condition of (internal) equilibrium is

$$(\delta\zeta)_{t,p} \geq 0. \tag{117}$$

That this condition is equivalent to (2) will easily appear from considerations like those used in respect to (111).

Hence, it is necessary for the equilibrium of two contiguous masses identical in composition that the values of ζ as determined for equal quantities of the two masses should be equal. Or, when one of three contiguous masses can be formed out of the other two, it is necessary for equilibrium that the value of ζ for any quantity of the first mass should be equal to the sum of the values of ζ for such quantities of the second and third masses as together contain the same matter. Thus, for the equilibrium of a solution composed of a parts of water and b parts of a salt which is in contact with vapor of water and crystals of the salt, it is necessary that the value of ζ for the quantity $a + b$ of the solution should be equal to the sum of the values of ζ for the quantities a of the vapor and b of the salt. Similar propositions will hold true in more complicated cases. The reader will easily deduce these conditions from the particular conditions of equilibrium given on page 128.

In like manner we may extend the definition of χ to any mass or combination of masses in which the pressure is everywhere the same, using ε for the energy and v for the volume of the whole and setting as before

$$\chi = \varepsilon + pv. \tag{118}$$

If we denote by Q the heat received by the combined masses from external sources in any process in which the pressure is not varied, and distinguish the initial and final states of the system by accents we have

$$\chi'' - \chi' = \varepsilon'' - \varepsilon' + p(v'' - v') = Q. \quad (119)$$

This function may therefore be called the *heat function for constant pressure* (just as the energy might be called the heat function for constant volume), the diminution of the function representing in all cases in which the pressure is not varied the heat given out by the system. In all cases of chemical action in which no heat is allowed to escape the value of χ remains unchanged.

POTENTIALS.

In the definition of the potentials μ_1, μ_2 , etc., the energy of a homogeneous mass was considered as a function of its entropy, its volume, and the quantities of the various substances composing it. Then the potential for one of these substances was defined as the differential coefficient of the energy taken with respect to the variable expressing the quantity of that substance. Now, as the manner in which we consider the given mass as composed of various substances is in some degree arbitrary, so that the energy may be considered as a function of various different sets of variables expressing quantities of component substances, it might seem that the above definition does not fix the value of the potential of any substance in the given mass, until we have fixed the manner in which the mass is to be considered as composed. For example, if we have a solution obtained by dissolving in water a certain salt containing water of crystallization, we may consider the liquid as composed of m_s weight-units of the hydrate and m_w of water, or as composed of m_s of the anhydrous salt and m_w of water. It will be observed that the values of m_s and m_w are not the same, nor those of m_w and m_w , and hence it might seem that the potential for water in the given liquid considered as composed of the hydrate and water, viz.,

$$\left(\frac{d\varepsilon}{dm_w} \right)_{\eta, v, m_s}$$

would be different from the potential for water in the same liquid considered as composed of anhydrous salt and water, viz.,

$$\left(\frac{d\varepsilon}{dm_w} \right)_{\eta, v, m_s}$$

The value of the two expressions is, however, the same, for, although m_w is not equal to m_w , we may of course suppose dm_w to be equal to dm_w , and then the numerators in the two fractions will also be equal, as they each denote the increase of energy of the liquid, when the quantity dm_w or dm_w of water is added without altering the entropy and volume of the liquid. Precisely the same considerations will apply to any other case.

In fact, we may give a definition of a potential which shall not presuppose any choice of a particular set of substances as the components of the homogeneous mass considered.

Definition.—If to any homogeneous mass we suppose an infinitesimal quantity of any substance to be added, the mass remaining homogeneous and its entropy and volume remaining unchanged, the increase of the energy of the mass divided by the quantity of the substance added is the *potential* for that substance in the mass considered. (For the purposes of this definition, any chemical element or combination of elements in given proportions may be considered a substance, whether capable or not of existing by itself as a homogeneous body.)

In the above definition we may evidently substitute for entropy, volume, and energy, respectively, either temperature, volume, and the function ψ ; or entropy, pressure, and the function χ ; or temperature, pressure, and the function ζ . (Compare equation (104).)

In the same homogeneous mass, therefore, we may distinguish the potentials for an indefinite number of substances, each of which has a perfectly determined value.

Between the potentials for different substances in the same homogeneous mass the same equations will subsist as between the units of these substances. That is, if the substances, S_a, S_b , etc., S_k, S_l , etc., are components of any given homogeneous mass, and are such that

$$\alpha \mathfrak{S}_a + \beta \mathfrak{S}_b + \text{etc.} = \kappa \mathfrak{S}_k + \lambda \mathfrak{S}_l + \text{etc.}, \quad (120)$$

$\mathfrak{S}_a, \mathfrak{S}_b$, etc., $\mathfrak{S}_k, \mathfrak{S}_l$, etc. denoting the units of the several substances, and α, β , etc., κ, λ , etc. denoting numbers, then if μ_a, μ_b , etc., μ_k, μ_l , etc. denote the potentials for these substances in the homogeneous mass,

$$\alpha \mu_a + \beta \mu_b + \text{etc.} = \kappa \mu_k + \lambda \mu_l + \text{etc.} \quad (121)$$

To show this, we will suppose the mass considered to be very large. Then, the first number of (121) denotes the increase of the energy of the mass produced by the addition of the matter represented by the first member of (120), and the second member of (121) denotes the

increase of energy of the same mass produced by the addition of the matter represented by the second member of (120), the entropy and volume of the mass remaining in each case unchanged. Therefore, as the two members of (120) represent the same matter in kind and quantity, the two members of (121) must be equal.

But it must be understood that equation (120) is intended to denote equivalence of the substances represented *in the mass considered*, and not merely chemical identity; in other words, it is supposed that there are no passive resistances to change in the mass considered which prevent the substances represented by one member of (120) from passing into those represented by the other. For example, in respect to a mixture of vapor of water and free hydrogen and oxygen (at ordinary temperatures), we may not write

$$9 \mathfrak{S}_{A_9} = 1 \mathfrak{S}_H + 8 \mathfrak{S}_O,$$

but water is to be treated as an independent substance, and no necessary relation will subsist between the potential for water and the potentials for hydrogen and oxygen.

The reader will observe that the relations expressed by equations (43) and (51) (which are essentially relations between the potentials for actual components in different parts of a mass in a state of equilibrium) are simply those which by (121) would necessary subsist between the same potentials in any homogeneous mass containing as variable components all the substances to which the potentials relate.

In the case of a body of invariable composition, the potential for the single component is equal to the value of ζ for one unit of the body, as appears from the equation

$$\zeta = \mu m \tag{122}$$

to which (96) reduces in this case. Therefore, when $n = 1$, the fundamental equation between the quantities in the set (102) (see page 143) and that between the quantities in (103) may be derived either from the other by simple substitution. But, with this single exception, an equation between the quantities in one of the sets (99)–(103) cannot be derived from the equation between the quantities in another of these sets without differentiation.

Also in the case of a body of variable composition, when all the quantities of the components except one vanish, the potential for that one will be equal to the value of ζ for one unit of the body. We may make this occur for any given composition of the body by

choosing as one of the components the matter constituting the body itself, so that the value of ζ for one unit of a body may always be considered as a potential. Hence the relations between the values of ζ for contiguous masses given on page 147 may be regarded as relations between potentials.

The two following propositions afford definitions of a potential which may sometimes be convenient.

The potential for any substance in any homogeneous mass is equal to the amount of mechanical work required to bring a unit of the substance by a reversible process from the state in which its energy and entropy are both zero into combination with the homogeneous mass, which at the close of the process must have its original volume, and which is supposed so large as not to be sensibly altered in any part. All other bodies used in the process must by its close be restored to their original state, except those used to supply the work, which must be used only as the source of the work. For, in a reversible process, when the entropies of other bodies are not altered, the entropy of the substance and mass taken together will not be altered. But the original entropy of the substance is zero; therefore the entropy of the mass is not altered by the addition of the substance. Again, the work expended will be equal to the increment of the energy of the mass and substance taken together, and therefore equal, as the original energy of the substance is zero, to the increment of energy of the mass due to the addition of the substance, which by the definition on page 149 is equal to the potential in question.

The potential for any substance in any homogeneous mass is equal to the work required to bring a unit of the substance by a reversible process from a state in which $\psi = 0$ and the temperature is the same as that of the given mass into combination with this mass, which at the close of the process must have the same volume and temperature as at first, and which is supposed so large as not to be sensibly altered in any part. A source of heat or cold of the temperature of the given mass is allowed, with this exception, other bodies are to be used only on the same conditions as before. This may be shown by applying equation (109) to the mass and substance taken together.

The last proposition enables us to see very easily, how the value of the potential is affected by the arbitrary constants involved in the definition of the energy and the entropy of each elementary substance. For we may imagine the substance brought from the state in which $\psi = 0$ and the temperature is the same as that of the given

mass, first to any specified state of the same temperature, and then into combination with the given mass. In the first part of the process the work expended is evidently represented by the value of ψ for the unit of the substance in the state specified. Let this be denoted by ψ' , and let μ denote the potential in question, and W the work expended in bringing a unit of the substance from the specified state into combination with the given mass as aforesaid; then

$$\mu = \psi' + W. \quad (123)$$

Now as the state of the substance for which $\varepsilon = 0$ and $\eta = 0$ is arbitrary, we may simultaneously increase the energies of the unit of the substance in all possible states by any constant C , and the entropies of the substance in all possible states by any constant K . The value of ψ , or $\varepsilon - t\eta$, for any state would then be increased by $C - tK$, t denoting the temperature of the state. Applying this to ψ' in (123) and observing that the last term in this equation is independent of the values of these constants, we see that the potential would be increased by the same quantity $C - tK$, t being the temperature of the mass in which the potential is to be determined.

ON COEXISTENT PHASES OF MATTER.

In considering the different homogeneous bodies which can be formed out of any set of component substances, it will be convenient to have a term which shall refer solely to the composition and thermodynamic state of any such body without regard to its quantity or form. We may call such bodies as differ in composition or state different *phases* of the matter considered, regarding all bodies which differ only in quantity and form as different examples of the same phase. Phases which can exist together, the dividing surfaces being plane, in an equilibrium which does not depend upon passive resistances to change, we shall call *coexistent*.

If a homogeneous body has n independently variable components, the phase of the body is evidently capable of $n + 1$ independent variations. A system of r coexistent phases, each of which has the same n independently variable components is capable of $n + 2 - r$ variations of phase. For the temperature, the pressure, and the potentials for the actual components have the same values in the different phases, and the variations of these quantities are by (97) subject to as many conditions as there are different phases. Therefore, the number of independent variations in the values of these quantities, i. e., the number of independent variations of phase of the system, will be $n + 2 - r$.

Or, when the r bodies considered have not the same independently variable components, if we still denote by n the number of independently variable components of the r bodies taken as a whole, the number of independent variations of phase of which the system is capable will still be $n + 2 - r$. In this case, it will be necessary to consider the potentials for more than n component substances. Let the number of these potentials be $n + h$. We shall have by (97), as before, r relations between the variations of the temperature, of the pressure, and of these $n + h$ potentials, and we shall also have by (43) and (51) h relations between these potentials, of the same form as the relations which subsist between the units of the different component substances.

Hence, if $r = n + 2$, no variation in the phases (remaining coexistent) is possible. It does not seem probable that r can ever exceed $n + 2$. An example of $n = 1$ and $r = 3$ is seen in the coexistent solid, liquid, and gaseous forms of any substance of invariable composition. It seems not improbable that in the case of sulphur and some other simple substances there is more than one triad of coexistent phases; but it is entirely improbable that there are four coexistent phases of any simple substance. An example of $n = 2$ and $r = 4$ is seen in a solution of a salt in water in contact with vapor of water and two different kinds of crystals of the salt.

Concerning $n + 1$ Coexistent Phases.

We will now seek the differential equation which expresses the relation between the variations of the temperature and the pressure in a system of $n + 1$ coexistent phases (n denoting, as before, the number of independently variable components in the system taken as a whole).

In this case we have $n + 1$ equations of the general form of (97) (one for each of the coexistent phases), in which we may distinguish the quantities η , v , m_1 , m_2 , etc. relating to the different phases by accents. But t and p will each have the same value throughout, and the same is true of μ_1 , μ_2 , etc., so far as each of these occurs in the different equations. If the total number of these potentials is $n + h$, there will be h independent relations between them, corresponding to the h independent relations between the units of the component substances to which the potentials relate, by means of which we may eliminate the variations of h of the potentials from the equations of the form of (97) in which they occur.

Let one of these equations be

$$v' dp = \eta' dt + m_a' d\mu_a + m_b' d\mu_b + \text{etc.}, \tag{124}$$

and by the proposed elimination let it become

$$v' dp = \eta' dt + A_1' d\mu_1 + A_2' d\mu_2 \dots + A_n' d\mu_n \tag{125}$$

It will be observed that μ_a , for example, in (124) denotes the potential in the mass considered for a substance S_a which may or may not be identical with any of the substances S_1, S_2 , etc. to which the potentials in (125) relate. Now as the equations between the potentials by means of which the elimination is performed are similar to those which subsist between the units of the corresponding substances, (compare equations (38), (43), and (51),) if we denote these units by $\mathfrak{S}_a, \mathfrak{S}_b$, etc., $\mathfrak{S}_1, \mathfrak{S}_2$, etc., we must also have

$$m_a' \mathfrak{S}_a + m_b' \mathfrak{S}_b + \text{etc.} = A_1' \mathfrak{S}_1 + A_2' \mathfrak{S}_2 \dots + A_n' \mathfrak{S}_n \tag{126}$$

But the first member of this equation denotes (in kind and quantity) the matter in the body to which equations (124) and (125) relate. As the same must be true of the second member, we may regard this same body as composed of the quantity A_1' of the substance S_1 , with the quantity A_2' of the substance S_2 , etc. We will therefore, in accordance with our general usage, write m_1', m_2' , etc. for A_1', A_2' , etc. in (125), which will then become

$$v' dp = \eta' dt + m_1' d\mu_1 + m_2' d\mu_2 \dots + m_n' d\mu_n \tag{127}$$

But we must remember that the components to which the m_1', m_2' , etc. of this equation relate are not necessarily independently variable, as are the components to which the similar expressions in (97) and (124) relate. The rest of the $n + 1$ equations may be reduced to a similar form, viz.,

$$v'' dp = \eta'' dt + m_1'' d\mu_1 + m_2'' d\mu_2 \dots + m_n'' d\mu_n, \tag{128}$$

etc.

By elimination of $d\mu_1, d\mu_2, \dots, d\mu_n$ from these equations we obtain

$$\begin{vmatrix} v' & m_1' & m_2' & \dots & m_n' \\ v'' & m_1'' & m_2'' & \dots & m_n'' \\ v''' & m_1''' & m_2''' & \dots & m_n''' \\ \cdot & \cdot & \cdot & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & \cdot \end{vmatrix} dp = \begin{vmatrix} \eta' & m_1' & m_2' & \dots & m_n' \\ \eta'' & m_1'' & m_2'' & \dots & m_n'' \\ \eta''' & m_1''' & m_2''' & \dots & m_n''' \\ \cdot & \cdot & \cdot & \dots & \cdot \\ \cdot & \cdot & \cdot & \dots & \cdot \end{vmatrix} dt. \tag{129}$$

In this equation we may make v', v'' , etc. equal to unity. Then m_1', m_2', m_1'' , etc. will denote the separate densities of the components in the different phases, and η', η'' , etc. the densities of entropy.

When $n = 1$,

$$(m'' v' - m' v'') dp = (m'' \eta' - m' \eta'') dt, \quad (130)$$

or, if we make $m' = 1$ and $m'' = 1$, we have the usual formula

$$\frac{dp}{dt} = \frac{\eta' - \eta''}{v' - v''} = \frac{Q}{t(v'' - v')}, \quad (131)$$

in which Q denotes the heat absorbed by a unit of the substance in passing from one state to the other without change of temperature or pressure.

Concerning Cases in which the Number of Coexistent Phases is less than $n + 1$.

When $n > 1$, if the quantities of all the components S_1, S_2, \dots, S_n are proportional in two coexistent phases, the two equations of the form of (127) and (128) relating to these phases will be sufficient for the elimination of the variations of all the potentials. In fact, the condition of the coexistence of the two phases together with the condition of the equality of the $n - 1$ ratios of m_1', m_2', \dots, m_n' with the $n - 1$ ratios of $m_1'', m_2'', \dots, m_n''$ is sufficient to determine p as a function of t if the fundamental equation is known for each of the phases. The differential equation in this case may be expressed in the form of (130), m' and m'' denoting either the quantities of any one of the components or the total quantities of matter in the bodies to which they relate. Equation (131) will also hold true in this case, if the total quantity of matter in each of the bodies is unity. But this case differs from the preceding in that the matter which absorbs the heat Q in passing from one state to another, and to which the other letters in the formula relate, although the same in quantity, is not in general the same in kind at different temperatures and pressures. Yet the case will often occur that one of the phases is essentially invariable in composition, especially when it is a crystalline body, and in this case the matter to which the letters in (131) relate will not vary with the temperature and pressure.

When $n = 2$, two coexistent phases are capable, when the temperature is constant, of a single variation in phase. But as (130) will hold true in this case when $m_1' : m_2' :: m_1'' : m_2''$, it follows that for constant temperature the pressure is in general a maximum or a minimum when the composition of the two phases is identical. In like manner, the temperature of the two coexistent phases is in general a maximum or a minimum, for constant pressure, when the composition of the two phases is identical. Hence, the series of simultaneous values of t and p for which the composition of two coexistent phases

is identical separates those simultaneous values of t and p for which no coexistent phases are possible from those for which there are two pair of coexistent phases. This may be applied to a liquid having two independently variable components in connection with the vapor which it yields, or in connection with any solid which may be formed in it.

When $n = 3$, we have for three coexistent phases three equations of the form of (127), from which we may obtain the following,

$$\begin{vmatrix} v' & m_1' & m_2' \\ v'' & m_1'' & m_2'' \\ v''' & m_1''' & m_2''' \end{vmatrix} dp = \begin{vmatrix} \eta' & m_1' & m_2' \\ \eta'' & m_1'' & m_2'' \\ \eta''' & m_1''' & m_2''' \end{vmatrix} dt + \begin{vmatrix} m_1' & m_2' & m_3' \\ m_1'' & m_2'' & m_3'' \\ m_1''' & m_2''' & m_3''' \end{vmatrix} d\mu_3. \quad (132)$$

Now the value of the last of these determinants will be zero, when the composition of one of the three phases is such as can be produced by combining the other two. Hence, the pressure of three coexistent phases will in general be a maximum or minimum for constant temperature, and the temperature a maximum or minimum for constant pressure, when the above condition in regard to the composition of the coexistent phases is satisfied. The series of simultaneous values of t and p for which the condition is satisfied separates those simultaneous values of t and p for which three coexistent phases are not possible, from those for which there are two triads of coexistent phases. These propositions may be extended to higher values of n , and illustrated by the boiling temperatures and pressures of saturated solutions of $n - 2$ different solids in solvents having two independently variable components.

INTERNAL STABILITY OF HOMOGENEOUS FLUIDS AS INDICATED BY FUNDAMENTAL EQUATIONS.

We will now consider the stability of a fluid enclosed in a rigid envelop which is non-conducting to heat and impermeable to all the components of the fluid. The fluid is supposed initially homogeneous in the sense in which we have before used the word, i. e., uniform in every respect throughout its whole extent. Let $S_1, S_2, \dots S_n$ be the *ultimate* components of the fluid; we may then consider every body which can be formed out of the fluid to be composed of $S_1, S_2, \dots S_n$, and that in only one way. Let $m_1, m_2, \dots m_n$ denote the quantities of these substances in any such body, and let ϵ, η, v , denote its energy, entropy, and volume. The fundamental equation for compounds of $S_1, S_2, \dots S_n$, if completely determined, will give us all possible sets of simultaneous values of these variables for homogeneous bodies.

Now, if it is possible to assign such values to the constants $T, P, M_1, M_2, \dots, M_n$ that the value of the expression

$$\epsilon - T\eta + Pv - M_1 m_1 - M_2 m_2 \dots - M_n m_n \quad (133)$$

shall be zero for the given fluid, and shall be positive for every other phase of the same components, i. e., for every homogeneous body* not identical in nature and state with the given fluid (but composed entirely of S_1, S_2, \dots, S_n), the condition of the given fluid will be stable.

For, in any condition whatever of the given mass, whether or not homogeneous, or fluid, if the value of the expression (133) is not negative for any homogeneous part of the mass, its value for the whole mass cannot be negative; and if its value cannot be zero for any homogeneous part which is not identical in phase with the mass in its given condition, its value cannot be zero for the whole except when the whole is in the given condition. Therefore, in the case supposed, the value of this expression for any other than the given condition of the mass is positive. (That this conclusion cannot be invalidated by the fact that it is not entirely correct to regard a composite mass as made up of homogeneous parts having the same properties in respect to energy, entropy, etc., as if they were parts of larger homogeneous masses, will easily appear from considerations similar to those adduced on pages 131-133.) If, then, the value of the expression (133) for the mass considered is less when it is in the given condition than when it is in any other, the energy of the mass in its given condition must be less than in any other condition in which it has the same entropy and volume. The given condition is therefore stable. (See page 110.)

Again, if it is possible to assign such values to the constants in (133) that the value of the expression shall be zero for the given fluid mass, and shall not be negative for any phase of the same components, the given condition will be evidently not unstable. (See page 110.) It will be stable unless it is possible for the given matter in the given volume and with the given entropy to consist of homogeneous parts for all of which the value of the expression (133) is zero, but which are not all identical in phase with the mass in its given condition. (A mass consisting of such parts would be in equilibrium, as we have already seen on pages 133, 134.) In this case, if we disregard the quantities connected with the surfaces which divide the

* A vacuum is throughout this discussion to be regarded as a limiting case of an extremely rarified body. We may thus avoid the necessity of the specific mention of a vacuum in propositions of this kind.

homogeneous parts, we must regard the given condition as one of neutral equilibrium. But in regard to these homogeneous parts, which we may evidently consider to be all different phases, the following conditions must be satisfied. (The accents distinguish the letters referring to the different parts, and the unaccented letters refer to the whole mass.)

$$\left. \begin{aligned} \eta' + \eta'' + \text{etc.} &= \eta, \\ v' + v'' + \text{etc.} &= v, \\ m_1' + m_1'' + \text{etc.} &= m_1, \\ m_2' + m_2'' + \text{etc.} &= m_2, \\ \text{etc.} & \end{aligned} \right\} \quad (134)$$

Now the values of η , v , m_1 , m_2 , etc. are determined by the whole fluid mass in its given state, and the values of $\frac{\eta'}{v'}$, $\frac{\eta''}{v''}$, etc., $\frac{m_1'}{v'}$, $\frac{m_1''}{v''}$, etc., $\frac{m_2'}{v'}$, $\frac{m_2''}{v''}$, etc., etc., are determined by the phases of the various parts. But the phases of these parts are evidently determined by the phase of the fluid as given. They form, in fact, the whole set of coexistent phases of which the latter is one. Hence, we may regard (134) as $n + 2$ linear equations between v' , v'' , etc. (The values of v' , v'' , etc. are also subject to the condition that none of them can be negative.) Now one solution of these equations must give us the given condition of the fluid; and it is not to be expected that they will be capable of any other solution, unless the number of different homogeneous parts, that is, the number of different coexistent phases, is greater than $n + 2$. We have already seen (page 153) that it is not probable that this is ever the case.

We may, however, remark that in a certain sense an infinitely large fluid mass will be in neutral equilibrium in regard to the formation of the substances, if such there are, other than the given fluid, for which the value of (133) is zero (when the constants are so determined that the value of the expression is zero for the given fluid, and not negative for any substance); for the tendency of such a formation to be reabsorbed will diminish indefinitely as the mass out of which it is formed increases.

When the substances S_1 , S_2 , . . . S_n are all independently variable components of the given mass, it is evident from (86) that the conditions that the value of (133) shall be zero for the mass as given, and shall not be negative for any phase of the same components, can only be fulfilled when the constants T , P , M_1 , M_2 , . . . M_n are equal to the temperature, the pressure, and the several potentials in the given

mass. If we give these values to the constants, the expression (133) will necessarily have the value zero for the given mass and we shall only have to inquire whether its value is positive for all other phases. But when $S_1, S_2, \dots S_n$ are not all independently variable components of the given mass, the values which it will be necessary to give to the constants in (133) cannot be determined entirely from the properties of the given mass; but T and P must be equal to its temperature and pressure, and it will be easy to obtain as many equations connecting $M_1, M_2, \dots M_n$ with the potentials in the given mass as it contains independently variable components.

When it is not possible to assign such values to the constants in (133) that the value of the expression shall be zero for the given fluid, and either zero or positive for any phase of the same components, we have already seen (pages 129–134) that if equilibrium subsists without passive resistances to change, it must be in virtue of properties which are peculiar to small masses surrounded by masses of different nature, and which are not indicated by fundamental equations. In this case, the fluid will necessarily be unstable, if we extend this term to embrace all cases in which an initial disturbance confined to a small part of an indefinitely large fluid mass will cause an ultimate change of state not indefinitely small in degree throughout the whole mass. In the discussion of stability as indicated by fundamental equations it will be convenient to use the term in this sense.*

* If we wish to know the stability of the given fluid when exposed to a constant temperature, or to a constant pressure, or to both, we have only to suppose that there is enclosed in the same envelop with the given fluid another body (which cannot combine with the fluid) of which the fundamental equation is $\epsilon = T\eta$, or $\epsilon = -Pv$, or $\epsilon = T\eta - Pv$, as the case may be, (T and P denoting the constant temperature and pressure, which of course must be those of the given fluid,) and to apply the criteria of page 110 to the whole system. When it is possible to assign such values to the constants in (133) that the value of the expression shall be zero for the given fluid and positive for every other phase of the same components, the value of (133) for the whole system will be less when the system is in its given condition than when it is in any other. (Changes of form and position of the given fluid are of course regarded as immaterial.) Hence the fluid is stable. When it is not possible to assign such values to the constants that the value of (133) shall be zero for the given fluid and zero or positive for any other phase, the fluid is of course unstable. In the remaining case, when it is possible to assign such values to the constants that the value of (133) shall be zero for the given fluid and zero or positive for every other phase, but not without the value zero for some other phase, the state of equilibrium of the fluid as stable or neutral will be determined by the possibility of satisfying, for any other than the given condition of the fluid, equations like (134), in which, however, the first or the second or both are to be stricken out, according as we are considering the

In determining for any given positive values of T and P and any given values whatever of $M_1, M_2, \dots M_n$ whether the expression (133) is capable of a negative value for any phase of the components $S_1, S_2, \dots S_n$, and if not, whether it is capable of the value zero for any other phase than that of which the stability is in question, it is only necessary to consider phases having the temperature T and pressure P . For we may assume that a mass of matter represented by any values of $m_1, m_2, \dots m_n$ is capable of at least one state of not unstable equilibrium (which may or may not be a homogeneous state) at this temperature and pressure. It may easily be shown that for such a state the value of $\varepsilon - T\eta + Pv$ must be as small as for any other state of the same matter. The same will therefore be true of the value of (133). Therefore if this expression is capable of a negative value for any mass whatever, it will have a negative value for that mass at the temperature T and pressure P . And if this mass is not homogeneous, the value of (133) must be negative for at least one of its homogeneous parts. So also, if the expression (133) is not capable of a negative value for any phase of the components, any phase for which it has the value zero must have the temperature T and the pressure P .

It may easily be shown that the same must be true in the limiting cases in which $T=0$ and $P=0$. For negative values of P , (133) is always capable of negative values, as its value for a vacuum is Pv .

For any body of the temperature T and pressure P , the expression (133) may by (91) be reduced to the form

$$\zeta - M_1 m_1 - M_2 m_2 \dots - M_n m_n. \quad (135)$$

We have already seen (pages 131, 132) that an expression like (133), when $T, P, M_1, M_2, \dots M_n$ and v have any given finite values, cannot have an infinite negative value as applied to any real body. Hence, in determining whether (133) is capable of a negative value for any phase of the components $S_1, S_2, \dots S_n$, and if not, whether it is capable of the value zero for any other phase than that of which the stability is in question, we have only to consider the least value of which it is capable for a constant value of v . Any body giving this value must satisfy the condition that for constant volume

$$d\varepsilon - T d\eta - M_1 dm_1 - M_2 dm_2 \dots - M_n dm_n \geq 0, \quad (136)$$

stability of the fluid for constant temperature, or for constant pressure, or for both. The number of coexistent phases will sometimes exceed by one or two the number of the remaining equations, and then the equilibrium of the fluid will be neutral in respect to one or two independent changes.

or, if we substitute the value of $d\varepsilon$ taken from equation (86), using subscript $a \dots g$ for the quantities relating to the actual components of the body, and subscript $h \dots k$ for those relating to the possible,

$$t d\eta + \mu_a dm_a \dots + \mu_g dm_g + \mu_h dm_h \dots + \mu_k dm_k - T d\eta - M_1 dm_1 - M_2 dm_2 \dots - M_n dm_n \geq 0. \quad (137)$$

That is, the temperature of the body must be equal to T , and the potentials of its components must satisfy the same conditions as if it were in contact and in equilibrium with a body having potentials $M_1, M_2, \dots M_n$. Therefore the same relations must subsist between $\mu_a \dots \mu_g$, and $M_1 \dots M_n$ as between the units of the corresponding substances, so that

$$m_a \mu_a \dots + m_g \mu_g = m_1 M_1 \dots + m_n M_n; \quad (138)$$

and as we have by (93)

$$\varepsilon = t\eta - p v + \mu_a m_a \dots + \mu_g m_g, \quad (139)$$

the expression (133) will reduce (for the body or bodies for which it has the least value per unit of volume) to

$$(P - p) v, \quad (140)$$

the value of which will be positive, null, or negative, according as the value of

$$P - p \quad (141)$$

is positive, null, or negative.

Hence, the conditions in regard to the stability of a fluid of which all the ultimate components are independently variable admit a very simple expression. If the pressure of the fluid is greater than that of any other phase of the same components which has the same temperature and the same values of the potentials for its actual components, the fluid is stable without coexistent phases; if its pressure is not as great as that of some other such phase, it will be unstable; if its pressure is as great as that of any other such phase, but not greater than that of every other, the fluid will certainly not be unstable, and in all probability it will be stable (when enclosed in a rigid envelop which is impermeable to heat and to all kinds of matter), but it will be one of a set of coexistent phases of which the others are the phases which have the same pressure.

The considerations of the last two pages, by which the tests relating to the stability of a fluid are simplified, apply to such bodies as actually exist. But if we should form arbitrarily any equation as a fundamental equation, and ask whether a fluid of which the proper-

ties were given by that equation would be stable, the tests of stability last given would be insufficient, as some of our assumptions might not be fulfilled by the equation. The test, however, as first given (pages 156–159) would in all cases be sufficient.

Stability in respect to Continuous Changes of Phase.

In considering the changes which may take place in any mass, we have already had occasion to distinguish between infinitesimal changes in existing phases, and the formation of entirely new phases. A phase of a fluid may be stable in regard to the former kind of change, and unstable in regard to the latter. In this case it may be capable of continued existence in virtue of properties which prevent the commencement of discontinuous changes. But a phase which is unstable in regard to continuous changes is evidently incapable of permanent existence on a large scale except in consequence of passive resistances to change. We will now consider the conditions of stability in respect to continuous changes of phase, or, as it may also be called, stability in respect to adjacent phases. We may use the same general test as before, except that the expression (133) is to be applied only to phases which differ infinitely little from the phase of which the stability is in question. In this case the component substances to be considered will be limited to the independently variable components of the fluid, and the constants M_1, M_2 , etc. must have the values of the potentials for these components in the given fluid. The constants in (133) are thus entirely determined and the value of the expression for the given phase is necessarily zero. If for any infinitely small variation of the phase, the value of (133) can become negative, the fluid will be unstable; but if for every infinitely small variation of the phase the value of (133) becomes positive, the fluid will be stable. The only remaining case, in which the phase can be varied without altering the value of (133) can hardly be expected to occur. The phase concerned would in such a case have coexistent adjacent phases. It will be sufficient to discuss the condition of stability (in respect to continuous changes) without coexistent adjacent phases.

This condition, which for brevity's sake we will call the condition of stability, may be written in the form

$$\varepsilon'' - t' \eta'' + p' v'' - \mu_1' m_1'' \dots - \mu_n' m_n'' > 0, \quad (142)$$

in which the quantities relating to the phase of which the stability is in question are distinguished by single accents, and those relating to

the other phase by double accents. This condition is by (93) equivalent to

$$\begin{aligned} & \varepsilon'' - t' \eta'' + p' v'' - \mu_1' m_1'' \dots - \mu_n' m_n'' \\ & - \varepsilon' + t' \eta' - p' v' + \mu_1' m_1' \dots + \mu_n' m_n' > 0, \end{aligned} \quad (143)$$

and to

$$\begin{aligned} & -t' \eta'' + p' v'' - \mu_1' m_1'' \dots - \mu_n' m_n'' \\ & + t'' \eta'' - p'' v'' + \mu_1'' m_1'' \dots + \mu_n'' m_n'' > 0. \end{aligned} \quad (144)$$

The condition (143) may be expressed more briefly in the form

$$\Delta \varepsilon > t \Delta \eta - p \Delta v + \mu_1 \Delta m_1 \dots + \mu_n \Delta m_n, \quad (145)$$

if we use the character Δ to signify that the condition, although relating to infinitesimal differences, is not to be interpreted in accordance with the usual convention in respect to differential equations with neglect of infinitesimals of higher orders than the first, but is to be interpreted *strictly*, like an equation between finite differences. In fact, when a condition like (145) (interpreted strictly) is satisfied for infinitesimal differences, it must be possible to assign limits within which it shall hold true of finite differences. But it is to be remembered that the condition is not to be applied to any arbitrary values of $\Delta \eta$, Δv , Δm_1 , . . . Δm_n , but only to such as are determined by a change of phase. (If only the quantity of the body which determines the value of the variables should vary and not its phase, the value of the first member of (145) would evidently be zero.) We may free ourselves from this limitation by making v constant, which will cause the term $-p \Delta v$ to disappear. If we then divide by the constant v , the condition will become

$$\Delta \frac{\varepsilon}{v} > t \Delta \frac{\eta}{v} + \mu_1 \Delta \frac{m_1}{v} \dots + \mu_n \Delta \frac{m_n}{v}, \quad (146)$$

in which form it will not be necessary to regard v as constant. As we may obtain from (86)

$$d \frac{\varepsilon}{v} = t d \frac{\eta}{v} + \mu_1 d \frac{m_1}{v} \dots + \mu_n d \frac{m_n}{v}, \quad (147)$$

we see that *the stability of any phase in regard to continuous changes depends upon the same conditions in regard to the second and higher differential coefficients of the density of energy regarded as a function of the density of entropy and the densities of the several components, which would make the density of energy a minimum, if the necessary conditions in regard to the first differential coefficients were fulfilled.* When $n = 1$, it may be more convenient to regard m as constant

in (145) than v . Regarding m a constant, it appears that the stability of a phase depends upon the same conditions in regard to the second and higher differential coefficients of the energy of a unit of mass regarded as a function of its entropy and volume, which would make the energy a minimum, if the necessary conditions in regard to the first differential coefficients were fulfilled.

The formula (144) expresses the condition of stability for the phase to which t' , p' , etc. relate. But it is evidently the necessary and sufficient condition of the stability of all phases of certain kinds of matter, or of all phases within given limits, that (144) shall hold true of any two infinitesimally differing phases within the same limits, or, as the case may be, in general. For the purpose, therefore, of such *collective* determinations of stability, we may neglect the distinction between the two states compared, and write the condition in the form

$$- \eta \Delta t + v \Delta p - m_1 \Delta \mu_1 \dots - m_n \Delta \mu_n > 0, \quad (148)$$

or

$$\Delta p > \frac{\eta}{v} \Delta t + \frac{m_1}{v} \Delta \mu_1 \dots + \frac{m_n}{v} \Delta \mu_n. \quad (149)$$

Comparing (98), we see that it is necessary and sufficient for the stability in regard to continuous changes of all the phases within any given limits, that within those limits the same conditions should be fulfilled in respect to the second and higher differential coefficients of the pressure regarded as a function of the temperature and the several potentials, which would make the pressure a minimum, if the necessary conditions with respect to the first differential coefficients were fulfilled.

By equations (87) and (94), the condition (142) may be brought to the form

$$\begin{aligned} & \psi'' + t'' \eta'' + p' v'' - \mu_1' m_1'' \dots - \mu_n' m_n'' \\ & - \psi' - t' \eta' - p' v' + \mu_1' m_1' \dots + \mu_n' m_n' > 0. \end{aligned} \quad (150)$$

For the stability of all phases within any given limits it is necessary and sufficient that within the same limits this condition shall hold true of any two phases which differ infinitely little. This evidently requires that when $v' = v''$, $m_1' = m_1''$, \dots , $m_n' = m_n''$,

$$\psi'' - \psi' + (t'' - t') \eta' > 0; \quad (151)$$

and that when $t' = t''$

$$\begin{aligned} & \psi'' + p' v'' - \mu_1' m_1'' \dots + \mu_n' m_n'' \\ & - \psi' - p' v' + \mu_1' m_1' \dots + \mu_n' m_n' > 0. \end{aligned} \quad (152)$$

These conditions may be written in the form

$$[\Delta\psi + \eta \Delta t]_{v,m} < 0, \tag{153}$$

$$[\Delta\psi + p \Delta v - \mu_1 \Delta m_1 \dots - \mu_n \Delta m_n]_t > 0, \tag{154}$$

in which the subscript letters indicate the quantities which are to be regarded as constant, m standing for all the quantities $m_1 \dots m_n$. If these conditions hold true within any given limits, (150) will also hold true of any two infinitesimally differing phases within the same limits. To prove this, we will consider a third phase, determined by the equations

$$t''' = t', \tag{155}$$

and

$$v''' = v'', \quad m_1''' = m_1'', \quad \dots \quad m_n''' = m_n''. \tag{156}$$

Now by (153),

$$\psi''' - \psi'' + (t''' - t'') \eta'' < 0; \tag{157}$$

and by (154),

$$\begin{aligned} & \psi''' + p' v''' - \mu_1' m_1''' \dots - \mu_n' m_n''' \\ & - \psi'' - p' v'' + \mu_1' m_1'' \dots + \mu_n' m_n'' > 0. \end{aligned} \tag{158}$$

Hence,

$$\begin{aligned} & \psi'' + t'' \eta'' + p' v''' - \mu_1' m_1''' \dots - \mu_n' m_n''' \\ & - \psi' - t''' \eta'' - p' v'' + \mu_1' m_1'' \dots + \mu_n' m_n'' > 0, \end{aligned} \tag{159}$$

which by (155) and (156) is equivalent to (150). Therefore, the conditions (153) and (154) in respect to the phases within any given limits are necessary and sufficient for the stability of all the phases within those limits. It will be observed that in (153) we have the condition of thermal stability of a body considered as unchangeable in composition and in volume, and in (154), the condition of mechanical and chemical stability of the body considered as maintained at a constant temperature. Comparing equation (88), we see that the condition (153) will be satisfied, if $\frac{d^2\psi}{dt^2} < 0$, i. e., if $\frac{d\eta}{dt}$ or $t \frac{d\eta}{dt}$ (the specific heat for constant volume) is positive. When $n = 1$, i. e., when the composition of the body is invariable, the condition (154) will evidently not be altered, if we regard m as constant, by which the condition will be reduced to

$$[\Delta\psi + p \Delta v]_{t,m} > 0. \tag{160}$$

This condition will evidently be satisfied if $\frac{d^2\psi}{dv^2} > 0$, i. e., if $-\frac{dp}{dv}$ or $-v \frac{dp}{dv}$ (the elasticity for constant temperature) is positive. But when $n > 1$, (154) may be abbreviated more symmetrically by making v constant.

Again, by (91) and (96), the condition (142) may be brought to the form

$$\begin{aligned} & \zeta'' + t'' \eta'' - p'' v'' - \mu_1' m_1'' \dots - \mu_n' m_n'' \\ & - \zeta' - t' \eta' + p' v' + \mu_1' m_1' \dots + \mu_n' m_n' > 0. \end{aligned} \quad (161)$$

Therefore, for the stability of all phases within any given limits it is necessary and sufficient that within the same limits

$$[\Delta\zeta + \eta \Delta t - v \Delta p]_m < 0, \quad (162)$$

and

$$[\Delta\zeta - \mu_1 \Delta m_1 \dots - \mu_n \Delta m_n]_{t,p} > 0, \quad (163)$$

as may easily be proved by the method used with (153) and (154). The first of these formulæ expresses the thermal and mechanical conditions of stability for a body considered as unchangeable in composition, and the second the conditions of chemical stability for a body considered as maintained at a constant temperature and pressure. If $n = 1$, the second condition falls away, and as in this case $\zeta = m\mu$, condition (162) becomes identical with (148).

The foregoing discussion will serve to illustrate the relation of the general condition of stability in regard to continuous changes to some of the principal forms of fundamental equations. It is evident that each of the conditions (146), (149), (154), (162), (163) involve in general several particular conditions of stability. We will now give our attention to the latter. Let

$$\Phi = \varepsilon - t' \eta + p' v - \mu_1' m_1 \dots - \mu_n' m_n, \quad (164)$$

the accented letters referring to one phase and the unaccented to another. It is by (142) the necessary and sufficient condition of the stability of the first phase that, for constant values of the quantities relating to that phase and of v , the value of Φ shall be a minimum when the second phase is identical with the first. Differentiating (164), we have by (86)

$$\begin{aligned} d\Phi = (t - t') d\eta - (p - p') dv + (\mu_1 - \mu_1') dm_1 \\ \dots - (\mu_n - \mu_n') dm_n. \end{aligned} \quad (165)$$

Therefore, the above condition requires that if we regard v, m_1, \dots, m_n as having the constant values indicated by accenting these letters, t shall be an increasing function of η , when the variable phase differs sufficiently little from the fixed. But as the fixed phase may be any one within the limits of stability, t must be an increasing function of η (within these limits) for any constant values of v, m_1, \dots, m_n . This condition may be written

$$\left(\frac{\Delta t}{\Delta \eta} \right)_{v, m_1, \dots, m_n} > 0. \quad (166)$$

When this condition is satisfied, the value of Φ , for any given values of v, m_1, \dots, m_n , will be a minimum when $t = t'$. And therefore, in applying the general condition of stability relating to the value of Φ , we need only consider the phases for which $t = t'$.

We see again by (165) that the general condition requires that if we regard t, v, m_2, \dots, m_n as having the constant values indicated by accenting these letters, μ_1 shall be an increasing function of m_1 , when the variable phase differs sufficiently little from the fixed. But as the fixed phase may be any one within the limits of stability, μ_1 must be an increasing function of m_1 (within these limits) for any constant values of t, v, m_2, \dots, m_n . That is,

$$\left(\frac{\Delta\mu_1}{\Delta m_1}\right)_{t, v, m_2, \dots, m_n} > 0. \tag{167}$$

When this condition is satisfied, as well as (166), Φ will have a minimum value, for any constant values of v, m_2, \dots, m_n , when $t = t'$ and $\mu_1 = \mu_1'$; so that in applying the general condition of stability we need only consider the phases for which $t = t'$ and $\mu_1 = \mu_1'$.

In this way we may also obtain the following particular conditions of stability:

$$\left(\frac{\Delta\mu_2}{\Delta m_2}\right)_{t, v, m_1, m_3, \dots, m_n} > 0, \tag{168}$$

.

$$\left(\frac{\Delta\mu_n}{\Delta m_n}\right)_{t, v, m_1, \dots, m_{n-1}} > 0. \tag{169}$$

When the $n + 1$ conditions (166)–(169) are all satisfied, the value of Φ , for any constant value of v , will be a minimum when the temperature and the potentials of the variable phase are equal to those of the fixed. The pressures will then also be equal and the phases will be entirely identical. Hence, the general condition of stability will be completely satisfied, when the above particular conditions are satisfied.

From the manner in which these particular conditions have been derived, it is evident that we may interchange in them ν, m_1, \dots, m_n in any way, provided that we also interchange in the same way t, μ_1, \dots, μ_n . In this way we may obtain different sets of $n + 1$ conditions which are necessary and sufficient for stability. The quantity v might be included in the first of these lists, and $-p$ in the second, except in cases when, in some of the phases considered, the entropy or the quantity of one of the components has the value zero.

Let us write R_{n+1} for the determinant of the order $n + 1$

$$\begin{vmatrix} \frac{d^2 \varepsilon}{d\eta^2} & \frac{d^2 \varepsilon}{dm_1 d\eta} & \cdots & \frac{d^2 \varepsilon}{dm_n d\eta} \\ \frac{d^2 \varepsilon}{d\eta dm_1} & \frac{d^2 \varepsilon}{dm_1^2} & \cdots & \frac{d^2 \varepsilon}{dm_n dm_1} \\ \cdot & \cdot & \cdots & \cdot \\ \frac{d^2 \varepsilon}{d\eta dm_n} & \frac{d^2 \varepsilon}{dm_1 dm_n} & \cdots & \frac{d^2 \varepsilon}{dm_n^2} \end{vmatrix} \quad (173)$$

of which the constituents are by (86) the same as the coefficients in equations (172), and $R_n, R_{n-1},$ etc. for the minors obtained by erasing the last column and row in the original determinant and in the minors successively obtained, and R_1 for the last remaining constituent. Then if $dt, d\mu_1, \dots, d\mu_{n-1},$ and dv all have the value zero, we have by (172)

$$R_n d\mu_n = R_{n+1} dm_n, \quad (174)$$

that is,

$$\left(\frac{d\mu_n}{dm_n}\right)_{t, v, \mu_1, \dots, \mu_{n-1}} = \frac{R_{n+1}}{R_n}. \quad (175)$$

In like manner we obtain

$$\left(\frac{d\mu_{n-1}}{dm_{n-1}}\right)_{t, v, \mu_1, \dots, \mu_{n-2}, m_n} = \frac{R_n}{R_{n-1}}, \left. \begin{matrix} \\ \text{etc.} \end{matrix} \right\} \quad (176)$$

Therefore, the conditions obtained by writing d for Δ in (166)–(169) are equivalent to this, that the determinant given above with the n minors obtained from it as above mentioned and the last remaining constituent $\frac{d^2 \varepsilon}{d\eta^2}$ shall all be positive. Any phase for which this condition is satisfied will be stable, and no phase will be stable for which any of these quantities has a negative value. But the conditions (166)–(169) will remain valid, if we interchange in any way η, m_1, \dots, m_n (with corresponding interchange of t, μ_1, \dots, μ_n). Hence the order in which we erase successive columns with the corresponding rows in the determinant is immaterial. Therefore none of the minors of the determinant (173) which are formed by erasing corresponding rows and columns, and none of the constituents of the principal diagonal, can be negative for a stable phase.

We will now consider the conditions which characterize the *limits of stability* (i. e., the limits which divide stable from unstable phases)

with respect to continuous changes.* Here, evidently, one of the conditions (166)–(169) must cease to hold true. Therefore, one of the differential coefficients formed by changing \mathcal{J} into d in the first members of these conditions must have the value zero. (That it is the numerator and not the denominator in the differential coefficient which vanishes at the limit appears from the consideration that the denominator is in each case the differential of a quantity which is necessarily capable of progressive variation, so long at least as the phase is capable of variation at all under the conditions expressed by the subscript letters.) The same will hold true of the set of differential coefficients obtained from these by interchanging in any way η , m_1, \dots, m_n , and simultaneously interchanging t , μ_1, \dots, μ_n in the same way. But we may obtain a more definite result than this.

Let us give to η or t , to m_1 or μ_1, \dots to m_{n-1} or μ_{n-1} , and to v , the constant values indicated by these letters when accented. Then by (165)

$$d\Phi = (\mu_n - \mu_n') dm_n \quad (177)$$

Now

$$\mu_n - \mu_n' = \left(\frac{d\mu_n}{dm_n} \right)' (m_n - m_n') \quad (178)$$

approximately, the differential coefficient being interpreted in accordance with the above assignment of constant values to certain variables, and its value being determined for the phase to which the accented letters refer. Therefore,

$$d\Phi = \left(\frac{d\mu_n}{dm_n} \right)' (m_n - m_n') dm_n, \quad (179)$$

and

$$\Phi = \frac{1}{2} \left(\frac{d\mu_n}{dm_n} \right)' (m_n - m_n')^2. \quad (180)$$

The quantities neglected in the last equation are evidently of the same order as $(m_n - m_n')^3$. Now this value of Φ will of course be different (the differential coefficient having a different meaning) according as we have made η or t constant, and according as we have made m_1 or μ_1 constant, etc.; but since, within the limits of stability, the value of Φ , for any constant values of m_n and v , will be the least when $t, p, \mu_1, \dots, \mu_{n-1}$ have the values indicated by accenting these letters, the value of the differential coefficient will be at least as small

* The limits of stability with respect to discontinuous changes are formed by phases which are coexistent with other phases. Some of the properties of such phases have already been considered. See pages 152–156.

when we give these variables these constant values, as when we adopt any other of the suppositions mentioned above in regard to the quantities remaining constant. And in all these relations we may interchange in any way η, m_1, \dots, m_n , if we interchange in the same way t, μ_1, \dots, μ_n . It follows that, within the limits of stability, when we choose for any one of the differential coefficients

$$\frac{dt}{d\eta}, \frac{d\mu_1}{dm_1}, \dots, \frac{d\mu_n}{dm_n} \tag{181}$$

the quantities following the sign d in the numerators of the others together with v as those which are to remain constant in differentiation, the value of the differential coefficient as thus determined will be at least as small as when one or more of the constants in differentiation are taken from the denominators, one being still taken from each fraction, and v as before being constant.

Now we have seen that none of these differential coefficients, as determined in any of these ways, can have a negative value within the limit of stability, and that some of them must have the value zero at that limit. Therefore, in virtue of the relations just established, one at least of these differential coefficients determined by considering constant the quantities occurring in the numerators of the others together with v , will have the value zero. But if one such has the value zero, all such will in general have the same value. For if

$$\left(\frac{d\mu_n}{dm_n}\right)_{t, v, \mu_1, \dots, \mu_{n-1}} \tag{182}$$

for example, has the value zero, we may change the density of the component S_n without altering (if we disregard infinitesimals of higher orders than the first) the temperature or the potentials, and therefore, by (98), without altering the pressure. That is, we may change the phase without altering any of the quantities $t, p, \mu_1, \dots, \mu_n$. (In other words, the phases adjacent to the limits of stability exhibit *approximately* the relations characteristic of neutral equilibrium.) Now this change of phase, which changes the density of one of the components, will in general change the density of the others and the density of entropy. Therefore, all the other differential coefficients formed after the analogy of (182), i. e., formed from the fractions in (181) by taking as constants for each the quantities in the numerators of the others together with v , will in general have the value zero at the limit of stability. And the relation which characterizes the limit of stability may be expressed, in general, by setting any one of these differential coefficients equal to zero. Such

an equation, when the fundamental equation is known, may be reduced to the form of an equation between the independent variables of the fundamental equation.

Again, as the determinant (173) is equal to the product of the differential coefficients obtained by writing d for Δ in the first members of (166)–(169), the equation of the limit of stability may be expressed by setting this determinant equal to zero. The form of the differential equation as thus expressed will not be altered by the interchange of the expressions η, m_1, \dots, m_n , but it will be altered by the substitution of v for any one of these expressions, which will be allowable whenever the quantity for which it is substituted has not the value zero in any of the phases to which the formula is to be applied.

The condition formed by setting the expression (182) equal to zero is evidently equivalent to this, that

$$\left[\frac{d\mu_n}{d\frac{m_n}{v}} \right]_{t, \mu_1, \dots, \mu_{n-1}} = 0, \quad (183)$$

that is, that

$$\left[\frac{d\frac{m_n}{v}}{d\mu_n} \right]_{t, \mu_1, \dots, \mu_{n-1}} = \infty, \quad (184)$$

or by (98), if we regard t, μ_1, \dots, μ_n as the independent variables,

$$\left(\frac{d^2 p}{d\mu_n^2} \right) = \infty. \quad (185)$$

In like manner we may obtain

$$\frac{d^2 p}{dt^2} = \infty, \quad \frac{d^2 p}{d\mu_1^2} = \infty, \quad \dots \quad \frac{d^2 p}{d\mu_{n-1}^2} = \infty. \quad (186)$$

Any one of these equations, (185), (186), may be regarded, in general, as the equation of the limit of stability. We may be certain that at every phase at that limit one at least of these equations will hold true.

GEOMETRICAL ILLUSTRATIONS.

Surfaces in which the Composition of the Body represented is Constant.

In vol. ii, p. 382, of the *Trans. Conn. Acad.*, a method is described of representing the thermodynamic properties of substances of invariable composition by means of surfaces. The volume, entropy, and energy

of a constant quantity of a substance are represented by rectangular co-ordinates. This method corresponds to the first kind of fundamental equation described on pages 140–144. Any other kind of fundamental equation for a substance of invariable composition will suggest an analogous geometrical method. Thus, if we make m constant, the variables in any one of the sets (99)–(103) are reduced to three, which may be represented by rectangular co-ordinates. This will, however, afford but four different methods, for, as has already (page 150) been observed, the two last sets are essentially equivalent when $n = 1$.

The method described in the preceding volume has certain advantages, especially for the purposes of theoretical discussion, but it may often be more advantageous to select a method in which the properties represented by two of the co-ordinates shall be such as best serve to identify and describe the different states of the substance. This condition is satisfied by temperature and pressure as well, perhaps, as by any other properties. We may represent these by two of the co-ordinates and the potential by the third. (See page 143.) It will not be overlooked that there is the closest analogy between these three quantities in respect to their parts in the general theory of equilibrium. (A similar analogy exists between volume, entropy, and energy.) If we give m the constant value unity, the third co-ordinate will also represent ζ , which then becomes equal to μ .

Comparing the two methods, we observe that in one

$$v = x, \quad \eta = y, \quad \varepsilon = z, \tag{187}$$

$$p = -\frac{dz}{dx}, \quad t = \frac{dz}{dy}, \quad \mu = \zeta = z - \frac{dz}{dx}x - \frac{dz}{dy}y; \tag{188}$$

and in the other

$$t = x, \quad p = y, \quad \mu = \zeta = z, \tag{189}$$

$$\eta = -\frac{dz}{dx}, \quad v = \frac{dz}{dy}, \quad \varepsilon = z - \frac{dz}{dx}x - \frac{dz}{dy}y. \tag{190}$$

Now $\frac{dz}{dx}$ and $\frac{dz}{dy}$ are evidently determined by the inclination of the tangent plane, and $z - \frac{dz}{dx}x - \frac{dz}{dy}y$ is the segment which it cuts off on the axis of Z . The two methods, therefore, have this reciprocal relation, that the quantities represented in one by the position of a point in a surface are represented in the other by the position of a tangent plane.

The surfaces defined by equations (187) and (189) may be distinguished as the v - η - ε surface, and the t - p - ζ surface, of the substance to which they relate.

In the t - p - ζ surface a line in which one part of the surface cuts another represents a series of pairs of coexistent states. A point through which pass three different parts of the surface represents a triad of coexistent states. Through such a point will evidently pass the three lines formed by the intersection of these sheets taken two by two. The perpendicular projection of these lines upon the p - t plane will give the curves which have recently been discussed by Professor J. Thomson.* These curves divide the space about the projection of the triple point into six parts which may be distinguished as follows: Let $\zeta^{(V)}$, $\zeta^{(L)}$, $\zeta^{(S)}$ denote the three ordinates determined for the same values of p and t by the three sheets passing through the triple point, then in one of the six spaces

$$\zeta^{(V)} < \zeta^{(L)} < \zeta^{(S)}, \quad (191)$$

in the next space, separated from the former by the line for which $\zeta^{(L)} = \zeta^{(S)}$,

$$\zeta^{(V)} < \zeta^{(S)} < \zeta^{(L)}, \quad (192)$$

in the third space, separated from the last by the line for which $\zeta^{(V)} = \zeta^{(S)}$,

$$\zeta^{(S)} < \zeta^{(V)} < \zeta^{(L)}, \quad (193)$$

in the fourth

$$\zeta^{(S)} < \zeta^{(L)} < \zeta^{(V)}, \quad (194)$$

in the fifth

$$\zeta^{(L)} < \zeta^{(S)} < \zeta^{(V)}, \quad (195)$$

in the sixth

$$\zeta^{(L)} < \zeta^{(V)} < \zeta^{(S)}. \quad (196)$$

The sheet which gives the least values of ζ is in each case that which represents the stable states of the substance. From this it is evident that in passing around the projection of the triple point we pass through lines representing alternately coexistent stable and coexistent unstable states. But the states represented by the intermediate values of ζ may be called stable *relatively* to the states represented by the highest. The differences $\zeta^{(L)} - \zeta^{(V)}$, etc. represent the amount of work obtained in bringing the substance by a reversible process from one to the other of the states to which these quantities relate, in a medium having the temperature and pressure common to the two states. To illustrate such a process, we may suppose a plane perpendicular to the axis of temperature to pass through the points

* See the Reports of the British Association for 1871 and 1872; and Philosophical Magazine, vol. xlvii. (1874), p. 447.

representing the two states. This will in general cut the double line formed by the two sheets to which the symbols (L) and (V) refer. The intersections of the plane with the two sheets will connect the double point thus determined with the points representing the initial and final states of the process, and thus form a *reversible path* for the body between those states.

The geometrical relations which indicate the stability of any state may be easily obtained by applying the principles stated on pp. 156 ff. to the case in which there is but a single component. The expression (133) as a test of stability will reduce to

$$\varepsilon - t' \eta + p' v - \mu' m, \quad (197)$$

the accented letters referring to the state of which the stability is in question, and the unaccented letters to any other state. If we consider the quantity of matter in each state to be unity, this expression may be reduced by equations (91) and (96) to the form

$$\zeta - \zeta' + (t - t') \eta - (p - p') v, \quad (198)$$

which evidently denotes the distance of the point (t', p', ζ') below the tangent plane for the point (t, p, ζ), measured parallel to the axis of ζ . Hence if the tangent plane for every other state passes above the point representing any given state, the latter will be stable. If any of the tangent planes pass below the point representing the given state, that state will be unstable. Yet it is not always necessary to consider these tangent planes. For, as has been observed on page-160, we may assume that (in the case of any real substance) there will be at least one not unstable state for any given temperature and pressure, except when the latter is negative. Therefore the state represented by a point in the surface on the positive side of the plane $p = 0$ will be unstable only when there is a point in the surface for which t and p have the same values and ζ a less value. It follows from what has been stated, that where the surface is doubly convex upwards (in the direction in which ζ is measured) the states represented will be stable in respect to adjacent states. This also appears directly from (162). But where the surface is concave upwards in either of its principal curvatures the states represented will be unstable in respect to adjacent states.

When the number of component substances is greater than unity, it is not possible to represent the fundamental equation by a single surface. We have therefore to consider how it may be represented by an infinite number of surfaces. A natural extension of either of the methods already described will give us a series of surfaces in

which every one is the v - η - ε surface, or every one the t - p - ζ surface for a body of constant composition, the proportion of the components varying as we pass from one surface to another. But for a simultaneous view of the properties which are exhibited by compounds of two or three components without change of temperature or pressure, we may more advantageously make one or both of the quantities t or p constant in each surface.

Surfaces and Curves in which the Composition of the Body represented is Variable and its Temperature and Pressure are Constant.

When there are three components, the position of a point in the X - Y plane may indicate the composition of a body most simply, perhaps, as follows. The body is supposed to be composed of the quantities m_1, m_2, m_3 of the substances S_1, S_2, S_3 , the value of $m_1 + m_2 + m_3$ being unity. Let P_1, P_2, P_3 be any three points in the plane, which are not in the same straight line. If we suppose masses equal to m_1, m_2, m_3 to be placed at these three points, the center of gravity of these masses will determine a point which will indicate the value of these quantities. If the triangle is equiangular and has the height unity, the distances of the point from the three sides will be equal numerically to m_1, m_2, m_3 . Now if for every possible phase of the components, of a given temperature and pressure, we lay off from the point in the X - Y plane which represents the composition of the phase a distance measured parallel to the axis of Z and representing the value of ζ (when $m_1 + m_2 + m_3 = 1$), the points thus determined will form a surface, which may be designated as the m_1 - m_2 - m_3 - ζ surface of the substances considered, or simply as their m - ζ surface, for the given temperature and pressure. In like manner, when there are but two component substances, we may obtain a curve, which we will suppose in the X - Z plane. The coordinate y may then represent temperature or pressure. But we will limit ourselves to the consideration of the properties of the m - ζ surface for $n = 3$, or the m - ζ curve for $n = 2$, regarded as a surface, or curve, which varies with the temperature and pressure.

As by (96) and (92)

$$\zeta = \mu_1 m_1 + \mu_2 m_2 + \mu_3 m_3,$$

and (for constant temperature and pressure)

$$d\zeta = \mu_1 dm_1 + \mu_2 dm_2 + \mu_3 dm_3,$$

if we imagine a tangent plane for the point to which these letters relate, and denote by ζ' the ordinate for any point in the plane,

and by m_1', m_2', m_3' , the distances of the foot of this ordinate from the three sides of the triangle $P_1 P_2 P_3$, we may easily obtain

$$\zeta' = \mu_1 m_1' + \mu_2 m_2' + \mu_3 m_3', \quad (199)$$

which we may regard as the equation of the tangent plane. Therefore the ordinates for this plane at P_1, P_2 , and P_3 are equal respectively to the potentials μ_1, μ_2 , and μ_3 . And in general, the ordinate for any point in the tangent plane is equal to the potential (in the phase represented by the point of contact) for a substance of which the composition is indicated by the position of the ordinate. (See page 149.) Among the bodies which may be formed of S_1, S_2 , and S_3 , there may be some which are incapable of variation in composition, or which are capable only of a single kind of variation. These will be represented by single points and curves in vertical planes. Of the tangent plane to one of these curves only a single line will be fixed, which will determine a series of potentials of which only two will be independent. The phase represented by a separate point will determine only a single potential, viz., the potential for the substance of the body itself, which will be equal to ζ .

The points representing a set of coexistent phases have in general a common tangent plane. But when one of these points is situated on the edge where a sheet of the surface terminates, it is sufficient if the plane is tangent to the edge and passes below the surface. Or, when the point is at the end of a separate line belonging to the surface, or at an angle in the edge of a sheet, it is sufficient if the plane pass through the point and below the line or sheet. If no part of the surface lies below the tangent plane, the points where it meets the plane will represent a stable (or at least not unstable) set of co-existent phases.

The surface which we have considered represents the relation between ζ and m_1, m_2, m_3 for homogeneous bodies when t and p have any constant values and $m_1 + m_2 + m_3 = 1$. It will often be useful to consider the surface which represents the relation between the same variables for bodies which consist of parts in different but coexistent phases. We may suppose that these are stable, at least in regard to adjacent phases, as otherwise the case would be devoid of interest. The point which represents the state of the composite body will evidently be at the center of gravity of masses equal to the parts of the body placed at the points representing the phases of these parts. Hence from the surface representing the properties of homogeneous bodies, which may be called the primitive surface, we

may easily construct the surface representing the properties of bodies which are in equilibrium but not homogeneous. This may be called the secondary or derived surface. It will consist, in general, of various portions or sheets. The sheets which represent a combination of two phases may be formed by rolling a double tangent plane upon the primitive surface: the part of the envelop of its successive positions which lies between the curves traced by the points of contact will belong to the derived surface. When the primitive surface has a triple tangent plane or one of higher order, the triangle in the tangent plane formed by joining the points of contact, or the smallest polygon without re-entrant angles which includes all the points of contact, will belong to the derived surface, and will represent masses consisting in general of three or more phases.

Of the whole thermodynamic surface as thus constructed for any temperature and any positive pressure, that part is especially important which gives the least value of ζ for any given values of m_1 , m_2 , m_3 . The state of a mass represented by a point in this part of the surface is one in which no dissipation of energy would be possible if the mass were enclosed in a rigid envelop impermeable both to matter and to heat; and the state of any mass composed of S_1 , S_2 , S_3 in any proportions, in which the dissipation of energy has been completed, so far as internal processes are concerned, (i. e., under the limitations imposed by such an envelop as above supposed,) would be represented by a point in the part which we are considering of the m - ζ surface for the temperature and pressure of the mass. We may therefore briefly distinguish this part of the surface as the *surface of dissipated energy*. It is evident that it forms a continuous sheet, the projection of which upon the X - Y plane coincides with the triangle $P_1 P_2 P_3$, (except when the pressure for which the m - ζ surface is constructed is negative, in which case there is no surface of dissipated energy,) that it nowhere has any convexity upward, and that the states which it represents are in no case unstable.

The general properties of the m - ζ lines for two component substances are so similar as not to require separate consideration. We now proceed to illustrate the use of both the surfaces and the lines by the discussion of several particular cases.

Three coexistent phases of two component substances may be represented by the points A, B, and C, in figure 1, in which ζ is measured toward the top of the page from $P_1 P_2$, m_1 toward the left from $P_2 Q_2$, and m_2 toward the right from $P_1 Q_1$. It is supposed that $P_1 P_2 = 1$. Portions of the curves to which these points belong

are seen in the figure, and will be denoted by the symbols (A), (B), (C). We may, for convenience, speak of these as separate curves, without implying anything in regard to their possible continuity in parts of the diagram remote from their common tangent AC. The *line of dissipated energy* includes the straight line AC and portions of the primitive curves (A) and (C). Let us first consider how the




FIG. 1.

diagram will be altered, if the temperature is varied while the pressure remains constant. If the temperature receives the increment dt , an ordinate of which the position is fixed will receive the increment $\left(\frac{d\zeta}{dt}\right)_{p, m} dt$, or $-\eta dt$. (The reader will easily convince himself that this is true of the

ordinates for the secondary line AC, as well as of the ordinates for the primitive curves.) Now if we denote by η' the entropy of the phase represented by the point B considered as belonging to the curve (B), and by η'' the entropy of the composite state of the same matter represented by the point B considered as belonging to the tangent to the curves (A) and (C), $t(\eta' - \eta'')$ will denote the heat yielded by a unit of matter in passing from the first to the second of these states. If this quantity is positive, an elevation of temperature will evidently cause a part of the curve (B) to protrude below the tangent to (A) and (C), which will no longer form a part of the line of dissipated energy. This line will then include portions of the three curves (A), (B), and (C), and of the tangents to (A) and (B) and to (B) and (C). On the other hand, a lowering of the temperature will cause the curve (B) to lie entirely above the tangent to (A) and (C), so that all the phases of the sort represented by (B) will be unstable. If $t(\eta' - \eta'')$ is negative, these effects will be produced by the opposite changes of temperature.

The effect of a change of pressure while the temperature remains constant may be found in a manner entirely analogous. The variation of any ordinate will be $\left(\frac{d\zeta}{dp}\right)_{t, m} dp$ or $v dp$. Therefore, if the volume of the homogeneous phase represented by the point B is a greater than the volume of the same matter divided between the the phases represented by A and C, an increase of pressure will give diagram indicating that all phases of the sort represented by curve (B) are unstable, and a decrease of pressure will give a diagram indi-

ating two stable pairs of coexistent phases, in each of which one of the phases is of the sort represented by the curve (B). When the relation of the volumes is the reverse of that supposed, these results will be produced by the opposite changes of pressure.

When we have four coexistent phases of three component substances, there are two cases which must be distinguished. In the first, one of the points of contact of the primitive surface with the quadruple tangent plane lies within the triangle formed by joining the other three; in the second, the four points may be joined so as to form a quadrilateral without re-entrant angles. Figure 2 represents the projection upon the X - Y plane (in which m_1, m_2, m_3 are measured) of a part of the surface of dissipated energy, when one of the points of contact D falls within the triangle formed by the other three A, B, C . This surface includes the triangle ABC in the quadruple tangent plane, portions of the three sheets of the primitive surface which touch the triangle at its vertices, EAF, GBH, ICK , and portions of the three developable surfaces formed by a tangent plane rolling upon each pair of these sheets. These developable surfaces are repre-




FIG. 2.




FIG. 3.

sented in the figure by ruled surfaces, the lines indicating the direction of their rectilinear elements. A point within the triangle ABC represents a mass of which the matter is divided, in general, between three or four different phases, in a manner not entirely determined by the position of a point. (The quantities of matter in these phases are such that if placed at the corresponding points, A, B, C, D , their center of gravity would be at the point representing the total mass.)

Such a mass, if exposed to constant temperature and pressure, would be in neutral equilibrium. A point in the developable surfaces represents a mass of which the matter is divided between two coexisting phases, which are represented by the extremities of the line in the figure passing through that point. A point in the primitive surface represents of course a homogeneous mass.

To determine the effect of a change of temperature without change of pressure upon the general features of the surface of dissipated energy, we must know whether heat is absorbed or yielded by a mass in passing from the phase represented by the point *D* in the primitive surface to the composite state consisting of the phases *A*, *B*, and *C* which is represented by the same point. If the first is the case, an increase of temperature will cause the sheet (*D*) (i. e., the sheet of the primitive surface to which the point *D* belongs) to separate from the plane tangent to the three other sheets, so as to be situated entirely above it, and a decrease of temperature, will cause a part of the sheet (*D*) to protrude through the plane tangent to the other sheets. These effects will be produced by the opposite changes of temperature, when heat is yielded by a mass passing from the homogeneous to the composite state above mentioned.

In like manner, to determine the effect of a variation of pressure without change of temperature, we must know whether the volume for the homogeneous phase represented by *D* is greater or less than the volume of the same matter divided between the phases *A*, *B*, and *C*. If the homogeneous phase has the greater volume, an increase of pressure will cause the sheet (*D*) to separate from the plane tangent to the other sheets, and a diminution of pressure will cause a part of the sheet (*D*) to protrude below that tangent plane. And these effects will be produced by the opposite changes of pressure, if the homogeneous phase has the less volume. All this appears from precisely the same considerations which were used in the analogous case for two component substances.

Now when the sheet (*D*) rises above the plane tangent to the other sheets, the general features of the surface of dissipated energy are not altered, except by the disappearance of the point *D*. But when the sheet (*D*) protrudes below the plane tangent to the other sheets, the surface of dissipated energy will take the form indicated in figure 3. It will include portions of the four sheets of the primitive surface, portions of the six developable surfaces formed by a double tangent plane rolling upon these sheets taken two by two, and portions of three triple tangent planes for these sheets taken by threes, the sheet (*D*) being always one of the three.

But when the points of contact with the quadruple tangent plane which represent the four coexistent phases can be joined so as to form a quadrilateral ABCD (fig. 4) without reentrant angles, the surface of dissipated energy will include this plane quadrilateral, portions of the four sheets of the primitive surface which are tangent to it, and portions of the four developable surfaces formed by double




FIG. 4.




FIG. 5.

tangent planes rolling upon the four pairs of these sheets which correspond to the four sides of the quadrilateral. To determine the general effect of a variation of temperature upon the surface of dissipated energy, let us consider the composite states represented by the point I at the intersection of the diagonals of the quadrilateral. Among these states (which all relate to the same kind and quantity of matter) there is one which is composed of the phases A and C, and another which is composed of the phases B and D. Now if the entropy of the first of these states is greater than that of the second, (i. e., if heat is given out by a body in passing from the first to the second state at constant temperature and pressure,) which we may suppose without loss of generality, an elevation of temperature while the pressure remains constant will cause the triple tangent planes to (B), (D), and (A), and to (B), (D), and (C), to rise above the triple tangent planes to (A), (C), and (B), and to (A), (C), and (D), in the vicinity of the point I. The surface of dissipated energy will therefore take the form indicated in figure 5, in which there are two plane triangles and five developable surfaces besides portions of the four primitive sheets. A diminution of temperature will give a different but entirely analogous form to the surface of dissipated energy. The quadrilateral ABCD will in this case break into two triangles along the diameter BD. The effects produced by

variation of the pressure while the temperature remains constant will of course be similar to those described. By considering the difference of volume instead of the difference of entropy of the two states represented by the point I in the quadruple tangent plane, we may distinguish between the effects of increase and diminution of pressure.

It should be observed that the points of contact of the quadruple tangent plane with the primitive surface may be at isolated points or curves belonging to the latter. So also, in the case of two component substances, the points of contact of the triple tangent line may be at isolated points belonging to the primitive curve. Such cases need not be separately treated, as the necessary modifications in the preceding statements, when applied to such cases, are quite evident. And in the remaining discussion of this geometrical method, it will generally be left to the reader to make the necessary limitations or modifications in analogous cases.

The necessary condition in regard to simultaneous variations of temperature and pressure, in order that four coexistent phases of three components, or three coexistent phases of two components, shall remain possible, has already been deduced by purely analytical processes. (See equation (129).)

We will next consider the case of two coexistent phases of identical composition, and first, when the number of components is two. The coexistent phases, if each is variable in composition, will be represented by the point of contact of two curves. One of the curves will in general lie above the other except at the point of contact; therefore, when the temperature and pressure remain constant, one phase cannot be varied in composition without becoming unstable, while the other phase will be stable if the proportion of either component is increased. By varying the temperature or pressure, we may cause the upper curve to protrude below the other, or to rise (relatively) entirely above it. (By comparing the volumes or the entropies of the two coexistent phases, we may easily determine which result would be produced by an increase of temperature or of pressure.) Hence, the temperatures and pressures for which two coexistent phases have the same composition form the limit to the temperatures and pressures for which such coexistent phases are possible. It will be observed that as we pass this limit of temperature and pressure, the pair of coexistent phases does not simply become unstable, like pairs and triads of coexistent phases which we have considered before, but there ceases to be any such pair of coexistent phases. The same result has already been obtained analytically on

page 155. But on that side of the limit on which the coexistent phases are possible, there will be two pairs of coexistent phases for the same values of t and p , as seen in figure 6. If the curve AA' represents vapor, and the curve BB' liquid, a liquid (represented by) B may exist in contact with a vapor A , and (at the same temperature and pressure) a liquid B' in contact with a vapor A' . If we compare these phases in respect to their composition, we see that in one case the vapor is richer than the liquid in a certain




FIG. 6.

component, and in the other case poorer. Therefore, if these liquids are made to boil, the effect on their composition will be opposite. If the boiling is continued under constant pressure, the temperature will rise as the liquids approach each other in composition, and the curve BB' will rise *relatively* to the curve AA' , until the curves are tangent to each other, when the two liquids become identical in nature, as also the vapors which they yield. In composition, and in the value of ζ per unit of mass, the vapor will then agree with the liquid. But if the curve BB' (which has the greater curvature) represents vapor, and AA' represents liquid, the effect of boiling will make the liquids A and A' differ more in composition. In this case, the relations indicated in the figure will hold for a temperature higher than that for which (with the same pressure) the curves are tangent to one another.

When two coexistent phases of three component substances have the same composition, they are represented by the point of contact of two sheets of the primitive surface. If these sheets do not intersect at the point of contact, the case is very similar to that which we have just considered. The upper sheet except at the point of contact represents unstable phases. If the temperature or pressure are so varied that a part of the upper sheet protrudes through the lower, the points of contact of a double tangent plane rolling upon the two sheets will describe a closed curve on each, and the surface of dissipated energy will include a portion of each sheet of the primitive surface united by a ring-shaped developable surface.

If the sheet having the greater curvatures represents liquid, and the other sheet vapor, the boiling temperature for any given pressure will be a maximum, and the pressure of saturated vapor for any given temperature will be a minimum, when the coexistent liquid and vapor have the same composition.

But if the two sheets, constructed for the temperature and pressure of the coexistent phases which have the same composition, intersect

at the point of contact, the whole primitive surface as seen from below will in general present four re-entrant furrows, radiating from the point of contact, for each of which a developable surface may be formed by a rolling double tangent plane. The different parts of the surface of dissipated energy in the vicinity of the point of contact are represented in figure 7. ATB, ETF are parts of one sheet of the primitive surface, and CTD, GTH are parts of the other. These are united by the developable surfaces BTC, DTE, FTG, HTA. Now we may make either sheet of the primitive surface sink relatively to the other by the proper variation of temperature or pressure. If the sheet to which ATB, ETF belong is that which sinks relatively, these




FIG. 7.

parts of the surface of dissipated energy will be merged in one, as well as the developable surfaces BTC, DTE, and also FTG, HTA. (The lines CTD, BTE, ATF, HTG will separate from one another at T, each forming a continuous curve.) But if the sheet of the primitive surface which sinks relatively is that to which CTD and GTH belong, then these parts will be merged in one in the surface of dissipated energy, as will be the developable surfaces BTC, ATH, and also DTE, FTG.

It is evident that this is not a case of maximum or minimum temperature for coexistent phases under constant pressure, or of maximum or minimum pressure for coexistent phases at constant temperature.

Another case of interest is when the composition of one of three coexistent phases is such as can be produced by combining the other two. In this case, the primitive surface must touch the same plane in three points in the same straight line. Let us distinguish the parts of the primitive surface to which these points belong as the sheets (A), (B), and (C), (C) denoting that which is intermediate in position. The sheet (C) is evidently tangent to the developable surface formed upon (A) and (B). It may or it may not intersect it at the point of contact. If it does not, it must lie above the developable surface, (unless it represents states which are unstable in regard to continuous changes,) and the surface of dissipated energy will include parts of the primitive sheets (A) and (B), the developable surface joining them, and the single point of the sheet (C) in which it meets this developable surface. Now, if the temperature or pressure is varied so as to make the sheet (C) rise above the

developable surface formed on the sheets (A) and (B), the surface of dissipated energy will be altered in its general features only by the removal of the single point of the sheet (C). But if the temperature or pressure is altered so as to make a part of the sheet (C) protrude through the developable surface formed on (A) and (B), the surface




FIG. 8.

of dissipated energy will have the form indicated in figure 8. It will include two plane triangles ABC and A'B'C', a part of each of the sheets (A) and (B), represented in the figure by the spaces on the left of the line aAA'a' and on the right of the line bBB'b', a small part CC' of the sheet (C), and developable surfaces formed upon these sheets taken by pairs ACC'A', BCC'B', aABb, a'A'B'b', the last two being different portions of the same developable surface.

But if, when the primitive surface is constructed for such a temperature and pressure that it has three points of contact with the same plane in the same straight line, the sheet (C) (which has the middle position) at its point of contact with the triple tangent plane intersects the developable surface formed upon the other sheets (A) and (B), the surface of dissipated energy will not include this developable surface, but will consist of portions of the three primitive sheets with two developable surfaces formed on (A) and (C) and on (B) and (C). These developable surfaces meet one another at the point of contact of (C) with the triple tangent plane, dividing the




FIG. 9.

portion of this sheet which belongs to the surface of dissipated energy into two parts. If now the temperature or pressure are varied so as to make the sheet (C) sink relatively to the developable surface formed on (A) and (B), the only alteration in the general features of the surface of dissipated energy will be that the developable surfaces

formed on (A) and (C) and on (B) and (C) will separate from one another, and the two parts of the sheet (C) will be merged in one. But a contrary variation of temperature or pressure will give a

surface of dissipated energy such as is represented in figure (9), containing two plane triangles ABC , $A'B'C'$ belonging to triple tangent planes, a portion of the sheet (A) on the left of the line $aAA'a'$, a portion of the sheet (B) on the right of the line $bBB'b'$, two separate portions $cC\gamma$ and $c'C'\gamma'$ of the sheet (C), two separate portions $aACc$ and $a'A'C'e'$ of the developable surface formed on (A) and (C), two separate portions $bBC\gamma$ and $b'B'C'\gamma'$ of the developable surface formed on (B) and (C), and the portion $A'ABB'$ of the developable surface formed on (A) and (B).

From these geometrical relations it appears that (in general) the temperature of three coexistent phases is a maximum or minimum for constant pressure, and the pressure of three coexistent phases a maximum or minimum for constant temperature, when the composition of the three coexistent phases is such that one can be formed by combining the other two. This result has been obtained analytically on page 156.

The preceding examples are amply sufficient to illustrate the use of the m - ζ surfaces and curves. The physical properties indicated by the nature of the surface of dissipated energy have been only occasionally mentioned, as they are often far more distinctly indicated by the diagrams than they could be in words. It will be observed that a knowledge of the lines which divide the various different portions of the surface of dissipated energy and of the direction of the rectilinear elements of the developable surfaces, *as projected upon the X - Y plane*, without a knowledge of the form of the m - ζ surface in space, is sufficient for the determination (in respect to the quantity and composition of the resulting masses) of the combinations and separations of the substances, and of the changes in their states of aggregation, which take place when the substances are exposed to the temperature and pressure to which the projected lines relate, except so far as such transformations are prevented by passive resistances to change.

CRITICAL PHASES.

It has been ascertained by experiment that the variations of two coexistent states of the same substance are in some cases limited in one direction by a terminal state at which the distinction of the coexistent states vanishes.* This state has been called the *critical state*. Analogous properties may doubtless be exhibited by compounds of variable composition without change of temperature or

* See Dr. Andrews "On the continuity of the gaseous and liquid states of matter." *Phil. Trans.*, vol. 159, p. 575.

pressure. For if, at any given temperature and pressure, two liquids are capable of forming a stable mixture in any ratio $m_1 : m_2$ less than a , and in any greater than b , a and b being the values of that ratio for two coexistent phases, while either can form a stable mixture with a third liquid in all proportions, and any small quantities of the first and second can unite at once with a great quantity of the third to form a stable mixture, it may easily be seen that two coexistent mixtures of the three liquids may be varied in composition, the temperature and pressure remaining the same, from initial phases in each of which the quantity of the third liquid is nothing, to a terminal phase in which the distinction of the two phases vanishes.

In general, we may define a *critical phase* as one at which the distinction between coexistent phases vanishes. We may suppose the coexistent phases to be stable in respect to continuous changes, for although relations in some respects analogous might be imagined to hold true in regard to phases which are unstable in respect to continuous changes, the discussion of such cases would be devoid of interest. But if the coexistent phases and the critical phase are unstable only in respect to the possible formation of phases entirely different from the critical and adjacent phases, the liability to such changes will in no respect affect the relations between the critical and adjacent phases, and need not be considered in a theoretical discussion of these relations, although it may prevent an experimental realization of the phases considered. For the sake of brevity, in the following discussion, phases in the vicinity of the critical phase will generally be called stable, if they are unstable only in respect to the formation of phases entirely different from any in the vicinity of the critical phase.

Let us first consider the number of independent variations of which a critical phase (while remaining such) is capable. If we denote by n the number of independently variable components, a pair of coexistent phases will be capable of n independent variations, which may be expressed by the variations of n of the quantities $t, p, \mu_1, \mu_2, \dots, \mu_n$. If we limit these variations by giving to $n-1$ of the quantities the constant values which they have for a certain critical phase, we obtain a linear* series of pairs of coexistent phases terminated by the critical phase. If we now vary infinitesimally the values of these $n-1$ quantities, we shall have for the new set of values considered constant a new linear series of pairs of coexistent phases. Now for every pair of phases in the first series, there must be pairs of phases in the

* This term is used to characterize a series having a *single* degree of extension.

second series differing infinitely little from the pair in the first, and *vice versa*, therefore the second series of coexistent phases must be terminated by a critical phase which differs, but differs infinitely little, from the first. We see, therefore, that if we vary arbitrarily the values of any $n - 1$ of the quantities $t, p, \mu_1, \mu_2, \dots, \mu_n$, as determined by a critical phase, we obtain one and only one critical phase for each set of varied values; i. e., a critical phase is capable of $n - 1$ independent variations.

The quantities $t, p, \mu_1, \mu_2, \dots, \mu_n$ have the same values in two coexistent phases, but the ratios of the quantities $\eta, v, m_1, m_2, \dots, m_n$ are in general different in the two phases. Or, if for convenience we compare equal volumes of the two phases (which involves no loss of generality), the quantities $\eta, m_1, m_2, \dots, m_n$ will in general have different values in two coexistent phases. Applying this to coexistent phases indefinitely near to a critical phase, we see that in the immediate vicinity of a critical phase, if the values of n of the quantities $t, p, \mu_1, \mu_2, \dots, \mu_n$, are regarded as constant (as well as v), the variations of either of the others will be infinitely small compared with the variations of the quantities $\eta, m_1, m_2, \dots, m_n$. This condition, which we may write in the form

$$\left(\frac{d\mu_n}{dm_n} \right)_{t, v, \mu_1, \dots, \mu_{n-1}} = 0, \tag{200}$$

characterizes, as we have seen on page 171, the limits which divide stable from unstable phases in respect to continuous changes.

In fact, if we give to the quantities $t, \mu_1, \mu_2, \dots, \mu_{n-1}$ constant values determined by a pair of coexistent phases, and to $\frac{m_n}{v}$ a series of values increasing from the less to the greater of the values which it has in these coexistent phases, we determine a linear series of phases connecting the coexistent phases, in some part of which μ_n —since it has the same value in the two coexistent phases, but not a uniform value throughout the series (for if it had, which is theoretically improbable, all these phases would be coexistent)—must be a decreasing function of $\frac{m_n}{v}$, or of m_n , if v also is supposed constant. Therefore, the series must contain phases which are unstable in respect to continuous changes. (See page 168.) And as such a pair of coexistent phases may be taken indefinitely near to any critical phase, the unstable phases (with respect to continuous changes) must approach indefinitely near to this phase.

Critical phases have similar properties with reference to stability as determined with regard to discontinuous changes. For as every stable phase which has a coexistent phase lies upon the limit which separates stable from unstable phases, the same must be true of any stable critical phase. (The same may be said of critical phases which are unstable in regard to discontinuous changes if we leave out of account the liability to the particular kind of discontinuous change in respect to which the critical phase is unstable.)

The linear series of phases determined by giving to n of the quantities $t, p, \mu_1, \mu_2, \dots, \mu_n$ the constant values which they have in any pair of coexistent phases consists of unstable phases in the part between the coexistent phases, but in the part beyond these phases in either direction it consists of stable phases. Hence, if a critical phase is varied in such a manner that n of the quantities $t, p, \mu_1, \mu_2, \dots, \mu_n$ remain constant, it will remain stable in respect both to continuous and to discontinuous changes. Therefore, μ_n is an increasing function of m_n when $t, v, \mu_1, \mu_2, \dots, \mu_{n-1}$ have constant values determined by any critical phase. But as equation (200) holds true at the critical phase, the following conditions must also hold true at that phase:

$$\left(\frac{d^2 \mu_n}{dm_n^2}\right)_{t, v, \mu_1, \dots, \mu_{n-1}} = 0, \quad (201)$$

$$\left(\frac{d^3 \mu_n}{dm_n^3}\right)_{t, v, \mu_1, \dots, \mu_{n-1}} \geq 0. \quad (202)$$

If the sign of equality holds in the last condition, additional conditions, concerning the differential coefficients of higher orders, must be satisfied.

Equations (200) and (201) may in general be called the equations of critical phases. It is evident that there are only two independent equations of this character, as a critical phase is capable of $n-1$ independent variations.

We are not, however, absolutely certain that equation (200) will always be satisfied by a critical phase. For it is possible that the denominator in the fraction may vanish as well as the numerator for an infinitesimal change of phase in which the quantities indicated are constant. In such a case, we may suppose the subscript n to refer to some different component substance, or use another differential coefficient of the same general form (such as are described on page 171 as characterizing the limits of stability in respect to continuous changes), making the corresponding changes in (201) and (202). We may be certain that some of the formulæ thus formed will not fail. But for a perfectly rigorous method there is an advan-

tage in the use of $\eta, v, m_1, m_2, \dots, m_n$ as independent variables. The condition that the phase may be varied without altering any of the quantities $t, \mu_1, \mu_2, \dots, \mu_n$ will then be expressed by the equation

$$R_{n+1} = 0, \tag{203}$$

in which R_{n+1} denotes the same determinant as on page 169. To obtain the second equation characteristic of critical phases, we observe that as a phase which is critical cannot become unstable when varied so that n of the quantities $t, p, \mu_1, \mu_2, \dots, \mu_n$ remain constant, the differential of R_{n+1} for constant volume, viz.,

$$\frac{dR_{n+1}}{d\eta} d\eta + \frac{dR_{n+1}}{dm_1} dm_1 \dots + \frac{dR_{n+1}}{dm_n} dm_n \tag{204}$$

cannot become negative when n of the equations (172) are satisfied. Neither can it have a positive value, for then its value might become negative by a change of sign of $d\eta, dm_1$, etc. Therefore the expression (204) has the value zero, if n of the equations (172) are satisfied. This may be expressed by an equation

$$S = 0, \tag{205}$$

in which S denotes a determinant in which the constituents are the same as in R_{n+1} , except in a single horizontal line, in which the differential coefficients in (204) are to be substituted. In whatever line this substitution is made, the equation (205), as well as (203), will hold true of every critical phase without exception.

If we choose $t, p, m_1, m_2, \dots, m_n$ as independent variables, and write U for the determinant

$$\begin{vmatrix} \frac{d^2\zeta}{dm_1^2} & \frac{d^2\zeta}{dm_2 dm_1} & \dots & \frac{d^2\zeta}{dm_{n-1} dm_1} \\ \frac{d^2\zeta}{dm_1 dm_2} & \frac{d^2\zeta}{dm_2^2} & \dots & \frac{d^2\zeta}{dm_{n-1} dm_2} \\ \dots & \dots & \dots & \dots \\ \frac{d^2\zeta}{dm_1 dm_{n-1}} & \frac{d^2\zeta}{dm_2 dm_{n-1}} & \dots & \frac{d^2\zeta}{dm_{n-1}^2} \end{vmatrix} \tag{206}$$

and V for the determinant formed from this by substituting for the constituents in any horizontal line the expressions

$$\frac{dU}{dm_1}, \quad \frac{dU}{dm^2}, \quad \dots, \quad \frac{dU}{dm_{n-1}}, \tag{207}$$

the equations of critical phases will be

$$U = 0, \quad V = 0. \tag{208}$$

It results immediately from the definition of a critical phase, that an infinitesimal change in the condition of a mass in such a phase

may cause the mass, if it remains in a state of dissipated energy (i. e., in a state in which the dissipation of energy by internal processes is complete), to cease to be homogeneous. In this respect a critical phase resembles any phase which has a coexistent phase, but differs from such phases in that the two parts into which the mass divides when it ceases to be homogeneous differ infinitely little from each other and from the original phase, and that neither of these parts is in general infinitely small. If we consider a change in the mass to be determined by the values of $d\eta$, dv , dm_1 , dm_2 , . . . dm_n , it is evident that the change in question will cause the mass to cease to be homogeneous whenever the expression

$$\frac{dR_{n+1}}{d\eta} d\eta + \frac{dR_{n+1}}{dv} dv + \frac{dR_{n+1}}{dm_1} dm_1, \dots + \frac{dR_{n+1}}{dm_n} dm_n \quad (209)$$

has a negative value. For if the mass should remain homogeneous, it would become unstable, as R_{n+1} would become negative. Hence, in general, any change thus determined, or its reverse (determined by giving to $d\eta$, dv , dm_1 , dm_2 , . . . dm_n the same values taken negatively), will cause the mass to cease to be homogeneous. The condition which must be satisfied with reference to $d\eta$, dv , dm_1 , dm_2 , . . . dm_n , in order that neither the change indicated, nor the reverse, shall destroy the homogeneity of the mass, is expressed by equating the above expression to zero.

But if we consider the change in the state of the mass (supposed to remain in a state of dissipated energy) to be determined by arbitrary values of $n+1$ of the differentials dt , dp , $d\mu_1$, $d\mu_2$, . . . $d\mu_n$, the case will be entirely different. For, if the mass ceases to be homogeneous, it will consist of two coexistent phases, and as applied to these only n of the quantities t , p , μ_1 , μ_2 , . . . μ_n will be independent. Therefore, for arbitrary variations of $n+1$ of these quantities, the mass must in general remain homogeneous.

But if, instead of supposing the mass to remain in a state of dissipated energy, we suppose that it remains homogeneous, it may easily be shown that to certain values of $n+1$ of the above differentials there will correspond three different phases, of which one is stable with respect both to continuous and to discontinuous changes, another is stable with respect to the former and unstable with respect to the latter, and the third is unstable with respect to both.

In general, however, if n of the quantities p , t , μ_1 , μ_2 , . . . μ_n , or n arbitrary functions of these quantities, have the same constant values as at a critical phase, the linear series of phases thus determined will be stable, in the vicinity of the critical phase. But if less

than n of these quantities or functions of the same together with certain of the quantities $\eta, v, m_1, m_2, \dots, m_n$, or arbitrary functions of the latter quantities, have the same values as at a critical phase, so as to determine a linear series of phases, the differential of R_{n+1} in such a series of phases will not in general vanish at the critical phase, so that in general a part of the series will be unstable.

We may illustrate these relations by considering separately the cases in which $n=1$ and $n=2$. If a mass of invariable composition is in a critical state, we may keep its volume constant, and destroy its homogeneity by changing its entropy (i. e., by adding or subtracting heat—probably the latter), or we may keep its entropy constant and destroy its homogeneity by changing its volume; but if we keep its pressure constant we cannot destroy its homogeneity by any thermal action, nor if we keep its temperature constant can we destroy its homogeneity by any mechanical action.

When a mass having two independently variable components is in a critical phase, and either its volume or its pressure is maintained constant, its homogeneity may be destroyed by a change of entropy or temperature. Or, if either its entropy or its temperature is maintained constant, its homogeneity may be destroyed by a change of volume or pressure. In both these cases it is supposed that the quantities of the components remain unchanged. But if we suppose both the temperature and the pressure to be maintained constant, the mass will remain homogeneous, however the proportion of the components be changed. Or, if a mass consists of two coexistent phases, one of which is a critical phase having two independently variable components, and either the temperature or the pressure of the mass is maintained constant, it will not be possible by mechanical or thermal means, or by changing the quantities of the components, to cause the critical phase to change into a pair of coexistent phases, so as to give three coexistent phases in the whole mass. The statements of this paragraph and of the preceding have reference only to infinitesimal changes.*

* A brief abstract (which came to the author's notice after the above was in type) of a memoir by M. Duclaux, "Sur la séparation des liquides mélangés, etc." will be found in *Comptes Rendus*, vol. lxxxi. (1875), p. 815.

ON THE VALUES OF THE POTENTIALS WHEN THE QUANTITY OF ONE OF THE COMPONENTS IS VERY SMALL.

If we apply equation (97) to a homogeneous mass having two independently variable components S_1 and S_2 , and make t , p , and m_1 constant, we obtain

$$m_1 \left(\frac{d\mu_1}{dm_2} \right)_{t, p, m_1} + m_2 \left(\frac{d\mu_2}{dm_2} \right)_{t, p, m_1} = 0. \quad (210)$$

Therefore, for $m_2=0$, either

$$\left(\frac{d\mu_1}{dm_2} \right)_{t, p, m_1} = 0, \quad (211)$$

or

$$\left(\frac{d\mu_2}{dm_2} \right)_{t, p, m_1} = \infty. \quad (212)$$

Now, whatever may be the composition of the mass considered, we may always so choose the substance S_1 that the mass shall consist solely of that substance, and in respect to any other variable component S_2 , we shall have $m_2=0$. But equation (212) cannot hold true *in general* as thus applied. For it may easily be shown (as has been done with regard to the potential on pages 148, 149) that the value of a differential coefficient like that in (212) for any given mass, when the substance S_2 (to which m_2 and μ_2 relate) is determined, is independent of the particular substance which we may regard as the other component of the mass; so that, if equation (212) holds true when the substance denoted by S_1 has been so chosen that $m_2=0$, it must hold true without such a restriction, which cannot generally be the case.

In fact, it is easy to prove directly that equation (211) will hold true of any phase which is stable in regard to continuous changes and in which $m_2=0$, *if m_2 is capable of negative as well as positive values.* For by (171), in any phase having that kind of stability, μ_1 is an increasing function of m_1 when t , p , and m_2 are regarded as constant. Hence, μ_1 will have its greatest value when the mass consists wholly of S_1 , i. e., when $m_2=0$. Therefore, if m_2 is capable of negative as well as positive values, equation (211) must hold true for $m_2=0$. (This appears also from the geometrical representation of potentials in the m - ζ curve. See page 177.)

But if m_2 is capable only of positive values, we can only conclude from the preceding considerations that the value of the differential coefficient in (211) cannot be positive. Nor, if we consider the physical significance of this case, viz., that an increase of m_2 denotes an

addition to the mass in question of a substance not before contained in it, does any reason appear for supposing that this differential coefficient has generally the value zero. To fix our ideas, let us suppose that S_1 denotes water, and S_2 a salt (either anhydrous or any particular hydrate). The addition of the salt to water, previously in a state capable of equilibrium with vapor or with ice, will destroy the possibility of such equilibrium at the same temperature and pressure. The liquid will dissolve the ice, or condense the vapor, which is brought in contact with it under such circumstances, which shows that μ_1 (the potential for water in the liquid mass) is diminished by the addition of the salt, when the temperature and pressure are maintained constant. Now there seems to be no *a priori* reason for supposing that the ratio of this diminution of the potential for water to the quantity of the salt which is added vanishes with this quantity. We should rather expect that, for small quantities of the salt, an effect of this kind would be proportional to its cause, i. e., that the differential coefficient in (211) would have a finite negative value for an infinitesimal value of m_2 . That this is the case with respect to numerous watery solutions of salts is distinctly indicated by the experiments of Wüllner* on the tension of the vapor yielded by such solutions, and of Rüdorff† on the temperature at which ice is formed in them; and unless we have experimental evidence that cases are numerous in which the contrary is true, it seems not unreasonable to assume, as a *general* law, that when m_2 has the value zero and is incapable of negative values, the differential coefficient in (211) will have a finite negative value, and that equation (212) will therefore hold true. But this case must be carefully distinguished from that in which m_2 is capable of negative values, which also may be illustrated by a solution of a salt in water. For this purpose let S_1 denote a hydrate of the salt which can be crystallized, and let S_2 denote water, and let us consider a liquid consisting entirely of S_1 and of such temperature and pressure as to be in equilibrium with crystals of S_1 . In such a liquid, an increase or a diminution of the quantity of water would alike cause crystals of S_1 to dissolve, which requires that the differential coefficient in (211) shall vanish at the particular phase of the liquid for which $m_2 = 0$.

Let us return to the case in which m_2 is incapable of negative values, and examine, without other restriction in regard to the substances

* Pogg. Ann., vol. ciii. (1858), p. 529; vol. cv. (1858), p. 85; vol. cx. (1860), p. 564.

† Pogg. Ann., vol. cxiv. (1861), p. 63.

denoted by S_1 and S_2 , the relation between μ_2 and $\frac{m_2}{m_1}$ for any constant temperature and pressure and for such small values of $\frac{m_2}{m_1}$ that the differential coefficient in (211) may be regarded as having the same constant value as when $m_2 = 0$, the values of t , p , and m_1 being unchanged. If we denote this value of the differential coefficient by $\frac{-A}{m_1}$, the value of A will be positive, and will be independent of m_1 .

Then for small values of $\frac{m_2}{m_1}$, we have by (210), approximately,

$$m_2 \left(\frac{d\mu_2}{dm_2} \right)_{t, p, m_1} = A, \quad (213)$$

i. e.,

$$\left(\frac{d\mu_2}{d \log m_2} \right)_{t, p, m_1} = A. \quad (214)$$

If we write the integral of this equation in the form

$$\mu_2 = A \log \frac{Bm_2}{m_1}, \quad (215)$$

B like A will have a positive value depending only upon the temperature and pressure. As this equation is to be applied only to cases in which the value of m_2 is very small compared with m_1 , we may regard $\frac{m_1}{v}$ as constant, when temperature and pressure are constant, and write

$$\mu_2 = A \log \frac{Cm_2}{v}, \quad (216)$$

C denoting a positive quantity, dependent only upon the temperature and pressure.

We have so far considered the composition of the body as varying only in regard to the proportion of two components. But the argument will be in no respect invalidated, if we suppose the composition of the body to be capable of other variations. In this case, the quantities A and C will be functions not only of the temperature and pressure but also of the quantities which express the composition of the substance of which together with S_2 the body is composed. If the quantities of any of the components besides S_2 are very small (relatively to the quantities of others), it seems reasonable to assume that the value of μ_2 , and therefore the values of A and C , will be nearly the same as if these components were absent.

Hence, if the independently variable components of any body are $S_a, \dots S_g$, and $S_h, \dots S_k$, the quantities of the latter being very small as compared with the quantities of the former, and are incapable of negative values, we may express approximately the values of the potentials for $S_h, \dots S_k$ by equations (subject of course to the uncertainties of the assumptions which have been made) of the form

$$\mu_h = A_h \log \frac{C_h m_h}{v}, \tag{217}$$

$$\dots \dots \dots$$

$$\mu_k = A_k \log \frac{C_k m_k}{v}, \tag{218}$$

in which $A_h, C_h, \dots A_k, C_k$ denote functions of the temperature, the pressure, and the ratios of the quantities $m_a, \dots m_g$.

We shall see hereafter, when we come to consider the properties of gases, that these equations may be verified experimentally in a very large class of cases, so that we have considerable reason for believing that they express a general law in regard to the limiting values of potentials.*

ON CERTAIN POINTS RELATING TO THE MOLECULAR CONSTITUTION OF BODIES.

It not unfrequently occurs that the number of proximate components which it is necessary to recognize as independently variable in a body exceeds the number of components which would be sufficient to express its ultimate composition. Such is the case, for example, as has been remarked on page 117, in regard to a mixture at ordinary temperatures of vapor of water and free hydrogen and oxygen. This case is explained by the existence of three sorts of molecules in the gaseous mass, viz., molecules of hydrogen, of oxygen, and of hydrogen and oxygen combined. In other cases, which are essentially the same in principle, we suppose a greater number of different sorts of molecules, which differ in composition, and the relations between

* The reader will not fail to remark that, if we could assume the universality of this law, the statement of the conditions necessary for equilibrium between different masses in contact would be much simplified. For, as the potential for a substance which is only a *possible* component (see page 117) would always have the value $-\infty$, the case could not occur that the potential for any substance should have a greater value in a mass in which that substance is only a possible component, than in another mass in which it is an actual component; and the conditions (22) and (51) might be expressed with the sign of equality without exception for the case of possible components.

these may be more complicated. Other cases are explained by molecules which differ in the quantity of matter which they contain, but not in the kind of matter, nor in the proportion of the different kinds. In still other cases, there appear to be different sorts of molecules, which differ neither in the kind nor in the quantity of matter which they contain, but only in the manner in which they are constituted. What is essential in the cases referred to is that a certain number of some sort or sorts of molecules shall be equivalent to a certain number of some other sort or sorts in respect to the kinds and quantities of matter which they collectively contain, and yet the former shall never be transformed into the latter within the body considered, nor the latter into the former, however the proportion of the numbers of the different sorts of molecules may be varied, or the composition of the body in other respects, or its thermodynamic state as represented by temperature and pressure or any other two suitable variables, provided, it may be, that these variations do not exceed certain limits. Thus, in the example given above, the temperature must not be raised beyond a certain limit, or molecules of hydrogen and of oxygen may be transformed into molecules of water.

The differences in bodies resulting from such differences in the constitution of their molecules are capable of continuous variation, in bodies containing the same matter and in the same thermodynamic state as determined, for example, by pressure and temperature, as the numbers of the molecules of the different sorts are varied. These differences are thus distinguished from those which depend upon the manner in which the molecules are combined to form sensible masses. The latter do not cause an increase in the number of variables in the fundamental equation; but they may be the cause of different values of which the function is sometimes capable for one set of values of the independent variables, as, for example, when we have several different values of ξ for the same values of $t, p, m_1, m_2, \dots, m_n$, one perhaps being for a gaseous body, one for a liquid, one for an amorphous solid, and others for different kinds of crystals, and all being invariable for constant values of the above mentioned independent variables.

But it must be observed that when the differences in the constitution of the molecules are entirely determined by the quantities of the different kinds of matter in a body with the two variables which express its thermodynamic state, these differences will not involve any increase in the number of variables in the fundamental equation. For example, if we should raise the temperature of the mixture of

vapor of water and free hydrogen and oxygen, which we have just considered, to a point at which the numbers of the different sorts of molecules are entirely determined by the temperature and pressure and the total quantities of hydrogen and of oxygen which are present, the fundamental equation of such a mass would involve but four independent variables, which might be the four quantities just mentioned. The fact of a certain part of the matter present existing in the form of vapor of water would, of course, be one of the facts which determine the nature of the relation between ζ and the independent variables, which is expressed by the fundamental equation.

But in the case first considered, in which the quantities of the different sorts of molecules are *not* determined by the temperature and pressure and the quantities of the different kinds of matter in the body as determined by its ultimate analysis, the components of which the quantities or the potentials appear in the fundamental equation must be those which are determined by the proximate analysis of the body, so that the variations in their quantities, with two variations relating to the thermodynamic state of the body, shall include all the variations of which the body is capable.* Such cases present no especial difficulty; there is indeed nothing in the physical and chemical properties of such bodies, so far as a certain range of experiments is concerned, which is different from what might be, if the proximate components were incapable of farther reduction or transformation. Yet among the various phases of the kinds of matter concerned, represented by the different sets of values of the variables which satisfy the fundamental equation, there is a certain class which merit especial attention. These are the phases for which the entropy has a maximum value for the same matter, as determined by the ultimate analysis of the body, with the same energy and volume. To fix our ideas let us call the proximate components S_1, \dots, S_n , and the ultimate components S_a, \dots, S_h ; and let m_1, \dots, m_n denote the quantities of the former, and m_a, \dots, m_h , the quantities of the latter. It is evident that $m_a \dots m_h$ are homogeneous functions of the first degree of m_1, \dots, m_n ; and that the relations between the substances S_1, \dots, S_n might be expressed by homogeneous equations of the first degree between the units of these substances, equal in number to the difference of the numbers of the proximate and of the ultimate com-

* The terms proximate or ultimate are not necessarily to be understood in an absolute sense. All that is said here and in the following paragraphs will apply to many cases in which components may conveniently be regarded as proximate or ultimate, which are such only in a relative sense.

ponents. The phases in question are those for which η is a maximum for constant values of ε , v , $m_a, \dots m_h$; or, as they may also be described, those for which ε is a minimum for constant values of η , v , $m_a, \dots m_h$; or for which ζ is a minimum for constant values of t , p , $m_a, \dots m_h$. The phases which satisfy this condition may be readily determined when the fundamental equation (which will contain the quantities $m_1, \dots m_n$ or $\mu_1, \dots \mu_n$) is known. Indeed it is easy to see that we may express the conditions which determine these phases by substituting $\mu_1, \dots \mu_n$ for the letters denoting the units of the corresponding substances in the equations which express the equivalence in ultimate analysis between these units.

These phases may be called, with reference to the kind of change which we are considering, phases of dissipated energy. That we have used a similar term before, with reference to a different kind of changes, yet in a sense entirely analogous, need not create confusion.

It is characteristic of these phases that we cannot alter the values of $m_1, \dots m_n$ in any real mass in such a phase, while the volume of the mass as well as its matter remain unchanged, without diminishing the energy or increasing the entropy of some other system. Hence, if the mass is large, its equilibrium can be but slightly disturbed by the action of any small body, or by a single electric spark, or by any cause which is not in some way proportioned to the effect to be produced. But when the proportion of the proximate components of a mass taken in connection with its temperature and pressure is not such as to constitute a phase of dissipated energy, it may be possible to cause great changes in the mass by the contact of a very small body. Indeed it is possible that the changes produced by such contact may only be limited by the attainment of a phase of dissipated energy. Such a result will probably be produced in a fluid mass by contact with another fluid which contains molecules of all the kinds which occur in the first fluid (or at least all those which contain the same kinds of matter which also occur in other sorts of molecules), but which differs from the first fluid in that the quantities of the various kinds of molecules are entirely determined by the ultimate composition of the fluid and its temperature and pressure. Or, to speak without reference to the molecular state of the fluid, the result considered would doubtless be brought about by contact with another fluid, which absorbs all the proximate components of the first, $S_1, \dots S_n$, (or all those between which there exist relations of equivalence in respect to their ultimate analysis), independently, and without passive resistances, but for which the phase is completely deter-

mined by its temperature and pressure and its ultimate composition (in respect at least to the particular substances just mentioned). By the absorption of the substances $S_1, \dots S_n$ independently and without passive resistances, it is meant that when the absorbing body is in equilibrium with another containing these substances, it shall be possible by *infinitesimal* changes in these bodies to produce the exchange of all these substances in either direction and independently. An exception to the preceding statement may of course be made for cases in which the result in question is prevented by the occurrence of some other kinds of change; in other words, it is assumed that the two bodies can remain in contact preserving the properties which have been mentioned.

The term *catalysis* has been applied to such action as we are considering. When a body has the property of reducing another, without limitation with respect to the proportion of the two bodies, to a phase of dissipated energy, in regard to a certain kind of molecular change, it may be called a *perfect catalytic agent* with respect to the second body and the kind of molecular change considered.

It seems not improbable that in some cases in which molecular changes take place slowly in homogeneous bodies, a mass of which the temperature and pressure are maintained constant will be finally brought to a state of equilibrium which is entirely determined by its temperature and pressure and the quantities of its ultimate components, while the various transitory states through which the mass passes, (which are evidently not completely defined by the quantities just mentioned,) may be completely defined by the quantities of certain proximate components with the temperature and pressure, and the matter of the mass may be brought by processes approximately reversible from permanent states to these various transitory states. In such cases, we may form a fundamental equation with reference to all possible phases, whether transitory or permanent; and we may also form a fundamental equation of different import and containing a smaller number of independent variables, which has reference solely to the final phases of equilibrium. The latter are the phases of dissipated energy (with reference to molecular changes), and when the more general form of the fundamental equation is known, it will be easy to derive from it the fundamental equation for these permanent phases alone.

Now, as these relations, theoretically considered, are independent of the rapidity of the molecular changes, the question naturally arises, whether in cases in which we are not able to distinguish such transi-

tory phases, they may not still have a theoretical significance. If so, the consideration of the subject from this point of view, may assist us, in such cases, in discovering the form of the fundamental equation with reference to the ultimate components, which is the only equation required to express all the properties of the bodies which are capable of experimental demonstration. Thus, when the phase of a body is completely determined by the quantities of n independently variable components, with the temperature and pressure, and we have reason to suppose that the body is composed of a greater number n' of proximate components, which are therefore not independently variable (while the temperature and pressure remain constant), it seems quite possible that the fundamental equation of the body may be of the same form as the equation for the phases of dissipated energy of analogous compounds of n' proximate and n ultimate components, in which the proximate components are capable of independent variation (without variation of temperature or pressure). And if such is found to be the case, the fact will be of interest as affording an indication concerning the proximate constitution of the body.

Such considerations seem to be especially applicable to the very common case in which at certain temperatures and pressures, regarded as constant, the quantities of certain proximate components of a mass are capable of independent variations, and all the phases produced by these variations are permanent in their nature, while at other temperatures and pressures, likewise regarded as constant, the quantities of these proximate components are not capable of independent variation, and the phase may be completely defined by the quantities of the ultimate components with the temperature and pressure. There may be, at certain intermediate temperatures and pressures, a condition with respect to the independence of the proximate components intermediate in character, in which the quantities of the proximate components are independently variable when we consider all phases, the essentially transitory as well as the permanent, but in which these quantities are not independently variable when we consider the permanent phases alone. Now we have no reason to believe that the passing of a body in a state of dissipated energy from one to another of the three conditions mentioned has any necessary connection with any discontinuous change of state. Passing the limit which separates one of these states from another will not therefore involve any discontinuous change in the values of any of the quantities enumerated in (99)–(103) on page 143, if $m_1, m_2, \dots, m_n, \mu_1, \mu_2, \dots, \mu_n$ are

understood as always relating to the ultimate components of the body. Therefore, if we regard masses in the different conditions mentioned above as having different fundamental equations, (which we may suppose to be of any one of the five kinds described on page 143,) these equations will agree at the limits dividing these conditions not only in the values of all the variables which appear in the equations, but also in all the differential coefficients of the first order involving these variables. We may illustrate these relations by supposing the values of t , p , and ζ for a mass in which the quantities of the ultimate components are constant to be represented by rectilinear coordinates. Where the proximate composition of such a mass is not determined by t and p , the value of ζ will not be determined by these variables, and the points representing connected values of t , p , and ζ will form a solid. This solid will be bounded in the direction opposite to that in which ζ is measured, by a surface which represents the phases of dissipated energy. In a part of the figure, all the phases thus represented may be permanent, in another part only the phases in the bounding surface, and in a third part there may be no such solid figure (for any phases of which the existence is experimentally demonstrable), but only a surface. This surface together with the bounding surfaces representing phases of dissipated energy in the parts of the figure mentioned above forms a continuous sheet, without discontinuity in regard to the direction of its normal at the limits dividing the different parts of the figure which have been mentioned. (There may, indeed, be different sheets representing liquid and gaseous states, etc., but if we limit our consideration to states of one of these sorts, the case will be as has been stated.)

We shall hereafter, in the discussion of the fundamental equations of gases, have an example of the derivation of the fundamental equation for phases of dissipated energy (with respect to the molecular changes on which the proximate composition of the body depends) from the more general form of the fundamental equation.

THE CONDITIONS OF EQUILIBRIUM FOR HETEROGENEOUS MASSES UNDER THE INFLUENCE OF GRAVITY.

Let us now seek the conditions of equilibrium for a mass of various kinds of matter subject to the influence of gravity. It will be convenient to suppose the mass enclosed in an immovable envelop which is impermeable to matter and to heat, and in other respects, except in regard to gravity, to make the same suppositions as on pages 115, 116. The energy of the mass will now consist of two parts, one of

which depends upon its intrinsic nature and state, and the other upon its position in space. Let Dm denote an element of the mass, $D\varepsilon$ the intrinsic energy of this element, h its height above a fixed horizontal plane, and g the force of gravity; then the total energy of the mass (when without sensible motions) will be expressed by the formula

$$\int D\varepsilon + \int fgh Dm, \quad (219)$$

in which the integrations include all the elements of the mass; and the general condition of equilibrium will be

$$\delta \int D\varepsilon + \delta \int fgh Dm \geq 0, \quad (220)$$

the variations being subject to certain equations of condition. These must express that the entropy of the whole mass is constant, that the surface bounding the whole mass is fixed, and that the total quantities of each of the component substances is constant. We shall suppose that there are no other equations of condition, and that the independently variable components are the same throughout the whole mass; and we shall at first limit ourselves to the consideration of the conditions of equilibrium with respect to the changes which may be expressed by infinitesimal variations of the quantities which define the initial state of the mass, without regarding the possibility of the formation at any place of infinitesimal masses entirely different from any initially existing in the same vicinity.

Let $D\eta$, Dv , Dm_1, \dots, Dm_n denote the entropy of the element Dm , its volume, and the quantities which it contains of the various components. Then

$$Dm = Dm_1 \dots + Dm_n, \quad (221)$$

and

$$\delta Dm = \delta Dm_1 \dots + \delta Dm_n. \quad (222)$$

Also, by equation (12),

$$\delta D\varepsilon = t \delta D\eta - p \delta Dv + \mu_1 \delta Dm_1 \dots + \mu_n \delta Dm_n. \quad (223)$$

By these equations the general condition of equilibrium may be reduced to the form

$$\begin{aligned} \int t \delta D\eta - \int p \delta Dv + \int f \mu_1 \delta Dm_1 \dots + \int f \mu_n \delta Dm_n \\ + \int f g \delta h Dm + \int f g h \delta Dm_1 \dots + \int f g h \delta Dm_n \geq 0. \end{aligned} \quad (224)$$

Now it will be observed that the different equations of condition affect different parts of this condition, so that we must have, separately,

$$\int t \delta D\eta \geq 0, \quad \text{if} \quad \int \delta D\eta = 0; \quad (225)$$

$$-f p \delta Dv + f g \delta h Dm \geq 0, \tag{226}$$

if the bounding surface is unvaried ;

$$\left. \begin{aligned} f \mu_1 \delta Dm_1 + f g h \delta Dm_1 &\geq 0, & \text{if } f \delta Dm_1 = 0; \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ f \mu_n \delta Dm_n + f g h \delta Dm_n &\geq 0, & \text{if } f \delta Dm_n = 0. \end{aligned} \right\} \tag{227}$$

From (225) we may derive the condition of thermal equilibrium,

$$t = \text{Const.} \tag{228}$$

Condition (226) is evidently the ordinary mechanical condition of equilibrium, and may be transformed by any of the usual methods. We may, for example, apply the formula to such motions as might take place longitudinally within an infinitely narrow tube, terminated at both ends by the external surface of the mass, but otherwise of indeterminate form. If we denote by m the mass, and by v the volume, included in the part of the tube between one end and a transverse section of variable position, the condition will take the form

$$-f p \delta dv + f g \delta h dm \geq 0, \tag{229}$$

in which the integrations include the whole contents of the tube. Since no motion is possible at the ends of the tube,

$$f p \delta dv + f \delta v dp = f d(p \delta v) = 0. \tag{230}$$

Again, if we denote by γ the density of the fluid,

$$f g \delta h dm = f g \frac{dh}{dv} \delta v \gamma dv = f g \gamma \delta v dh. \tag{231}$$

By these equations condition (229) may be reduced to the form

$$f \delta v (dp + g \gamma dh) \geq 0. \tag{232}$$

Therefore, since δv is arbitrary in value,

$$dp = - g \gamma dh, \tag{233}$$

which will hold true at any point in the tube, the differentials being taken with respect to the direction of the tube at that point. Therefore, as the form of the tube is indeterminate, this equation must hold true, without restriction, throughout the whole mass. It evidently requires that the pressure shall be a function of the height alone, and that the density shall be equal to the first derivative of this function, divided by $-g$.

Conditions (227) contain all that is characteristic of chemical equilibrium. To satisfy these conditions it is necessary and sufficient that

$$\left. \begin{aligned} \mu_1 + g h &= \text{Const.} \\ \cdot & \cdot \cdot \cdot \\ \mu_n + g h &= \text{Const.} \end{aligned} \right\} \tag{234}$$

The expressions μ_1, \dots, μ_n denote quantities which we have called the potentials for the several components, and which are entirely determined at any point in a mass by the nature and state of the mass about that point. We may avoid all confusion between these quantities and the potential of the force of gravity, if we distinguish the former, when necessary, as *intrinsic potentials*. The relations indicated by equations (234) may then be expressed as follows:

When a fluid mass is in equilibrium under the influence of gravity, and has the same independently variable components throughout, the intrinsic potentials for the several components are constant in any given level, and diminish uniformly as the height increases, the difference of the values of the intrinsic potential for any component at two different levels, being equal to the work done by the force of gravity when a unit of matter falls from the higher to the lower level.

The conditions expressed by equations (228), (233), (234) are necessary and sufficient for equilibrium, except with respect to the possible formation of masses which are not approximately identical in phase with any previously existing about the points where they may be formed. The possibility of such formations at any point is evidently independent of the action of gravity, and is determined entirely by the phase or phases of the matter about that point. The conditions of equilibrium in this respect have been discussed on pages 128–134.

But equations (228), (233), and (234) are not entirely independent. For with respect to any mass in which there are no surfaces of discontinuity (i. e., surfaces where adjacent elements of mass have finite differences of phase), one of these equations will be a consequence of the others. Thus by (228) and (234), we may obtain from (97), which will hold true of any continuous variations of phase, the equation

$$v dp = -g (m_1 \dots + m_n) dh; \quad (235)$$

or
$$dp = -g \gamma dh; \quad (236)$$

which will therefore hold true in any mass in which equations (228) and (234) are satisfied, and in which there are no surfaces of discontinuity. But the condition of equilibrium expressed by equation (233) has no exception with respect to surfaces of discontinuity; therefore in any mass in which such surfaces occur, it will be necessary for equilibrium, in addition to the relations expressed by equations (228) and (234), that there shall be no discontinuous change of pressure at these surfaces.

This superfluity in the particular conditions of equilibrium which we have found, as applied to a mass which is everywhere continuous

in phase, is due to the fact that we have made the elements of volume variable in position and size, while the matter initially contained in these elements is not supposed to be confined to them. Now, as the different components may move in different directions when the state of the system varies, it is evidently impossible to define the elements of volume so as always to include the same matter; we must, therefore, suppose the matter contained in the elements of volume to vary; and therefore it would be allowable to make these elements fixed in space. If the given mass has no surfaces of discontinuity, this would be much the simplest plan. But if there are any surfaces of discontinuity, it will be possible for the state of the given mass to vary, not only by infinitesimal changes of phase in the fixed elements of volume, but also by movements of the surfaces of discontinuity. It would therefore be necessary to add to our general condition of equilibrium terms relating to discontinuous changes in the elements of volume about these surfaces,—a necessity which is avoided if we consider these elements movable, as we can then suppose that each element remains always on the same side of the surface of discontinuity.

Method of treating the preceding problem, in which the elements of volume are regarded as fixed.

It may be interesting to see in detail how the particular conditions of equilibrium may be obtained if we regard the elements of volume as fixed in position and size, and consider the possibility of finite as well as infinitesimal changes of phase in each element of volume. If we use the character Δ to denote the differences determined by such finite differences of phase, we may express the variation of the intrinsic energy of the whole mass in the form

$$\int \delta D\varepsilon + \int \Delta D\varepsilon, \quad (237)$$

in which the first integral extends over all the elements which are infinitesimally varied, and the second over all those which experience a finite variation. We may regard both integrals as extending throughout the whole mass, but their values will be zero except for the parts mentioned.

If we do not wish to limit ourselves to the consideration of masses so small that the force of gravity can be regarded as constant in direction and in intensity, we may use Γ to denote the potential of the force of gravity, and express the variation of the part of the energy which is due to gravity in the form

$$-\int \Gamma \delta Dm - \int \Gamma \Delta Dm. \quad (238)$$

We shall then have, for the general condition of equilibrium,

$$f \delta D\varepsilon + f' \Delta D\varepsilon - f' \Gamma \delta Dm - f' \Gamma \Delta Dm \geq 0; \quad (239)$$

and the equations of condition will be

$$f \delta D\eta + f' \Delta D\eta = 0, \quad (240)$$

$$\left. \begin{aligned} f \delta Dm_1 + f' \Delta Dm_1 &= 0, \\ \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ f \delta Dm_n + f' \Delta Dm_n &= 0. \end{aligned} \right\} \quad (241)$$

We may obtain a condition of equilibrium independent of these equations of condition, by subtracting these equations, multiplied each by an indeterminate constant, from condition (239). If we denote these indeterminate constants by T, M_1, \dots, M_n , we shall obtain after arranging the terms

$$\begin{aligned} & \int \overline{\delta D\varepsilon - \Gamma \delta Dm - T \delta D\eta - M_1 \delta Dm_1 \dots - M_n \delta Dm_n} \\ & + \int \overline{\Delta D\varepsilon - \Gamma \Delta Dm - T \Delta D\eta - M_1 \Delta Dm_1 \dots - M_n \Delta Dm_n} \geq 0. \end{aligned} \quad (242)$$

The variations, both infinitesimal and finite, in this condition are independent of the equations of condition (240) and (241), and are only subject to the condition that the varied values of $D\varepsilon, D\eta, Dm_1, \dots, Dm_n$ for each element are determined by a certain change of phase. But as we do not suppose the same element to experience both a finite and an infinitesimal change of phase, we must have

$$\delta D\varepsilon - \Gamma \delta Dm - T \delta D\eta - M_1 \delta Dm_1 \dots - M_n \delta Dm_n \geq 0, \quad (243)$$

and

$$\Delta D\varepsilon - \Gamma \Delta Dm - T \Delta D\eta - M_1 \Delta Dm_1 \dots - M_n \Delta Dm_n \geq 0. \quad (244)$$

By equation (12), and in virtue of the necessary relation (222), the first of these conditions reduces to

$$\begin{aligned} (t - T) \delta D\eta + (\mu_1 - \Gamma - M_1) \delta Dm_1 \dots \\ + (\mu_n - \Gamma - M_n) \delta Dm_n \geq 0; \end{aligned} \quad (245)$$

for which it is necessary and sufficient that

$$t = T, \quad (246)$$

$$\left. \begin{aligned} \mu_1 - \Gamma &= M_1, \\ \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \\ \mu_n - \Gamma &= M_n. \end{aligned} \right\} * \quad (247)$$

* The gravitation potential is here supposed to be defined in the usual way. But if it were defined so as to *decrease* when a body falls, we would have the sign + instead of - in these equations; i. e., for each component, the sum of the gravitation and intrinsic potentials would be constant throughout the whole mass.

Condition (244) may be reduced to the form

$$\Delta D\varepsilon - T \Delta D\eta - (\Gamma + M_1) \Delta Dm_1 \dots - (\Gamma + M_n) \Delta Dm_n \geq 0; \quad (248)$$

and by (246) and (247) to

$$\Delta D\varepsilon - t \Delta D\eta - \mu_1 \Delta Dm_1 \dots - \mu_n \Delta Dm_n \geq 0. \quad (249)$$

If values determined subsequently to the change of phase are distinguished by accents, this condition may be written

$$D\varepsilon' - t D\eta' - \mu_1 Dm_1' \dots - \mu_n Dm_n' - D\varepsilon + t D\eta + \mu_1 Dm_1 \dots + \mu_n Dm_n \geq 0, \quad (250)$$

which may be reduced by (93) to

$$D\varepsilon' - t D\eta' - \mu_1 Dm_1', \dots - \mu_n Dm_n' + p Dv \geq 0. \quad (251)$$

Now if the element of volume Dv is adjacent to a surface of discontinuity, let us suppose $D\varepsilon'$, $D\eta'$, Dm_1' , \dots , Dm_n' to be determined (for the same element of volume) by the phase existing on the other side of the surface of discontinuity. As t , μ_1 , \dots , μ_n have the same values on both sides of this surface, the condition may be reduced by (93) to

$$-p' Dv + p Dv \geq 0. \quad (252)$$

That is, the pressure must not be greater on one side of a surface of discontinuity than on the other.

Applied more generally, (251) expresses the condition of equilibrium with respect to the possibility of discontinuous changes of phases at any point. As $Dv' = Dv$, the condition may also be written

$$D\varepsilon' - t D\eta' + p Dv' - \mu_1 Dm_1', \dots - \mu_n Dm_n' \geq 0, \quad (253)$$

which must hold true when t , p , μ_1 , \dots , μ_n have values determined by any point in the mass, and $D\varepsilon'$, $D\eta'$, Dv' , Dm_1' , \dots , Dm_n' have values determined by any possible phase of the substances of which the mass is composed. The application of the condition is, however, subject to the limitations considered on pages 128-134. It may easily be shown (see pages 160, 161) that for constant values of t , μ_1 , \dots , μ_n , and of Dv' , the first member of (253) will have the least possible value when $D\varepsilon'$, $D\eta'$, Dm_1' , \dots , Dm_n' are determined by a phase for which the temperature has the value t , and the potentials the values μ_1 , \dots , μ_n . It will be sufficient, therefore, to consider the condition as applied to such phases, in which case it may be reduced by (93) to

$$p - p' \geq 0. \quad (254)$$

That is, the pressure at any point must be as great as that of any phase of the same components, for which the temperature and the

potentials have the same values as at that point. We may also express this condition by saying that the pressure must be as great as is consistent with equations (246), (247). This condition with the equations mentioned will always be sufficient for equilibrium; when the condition is not satisfied, if equilibrium subsists, it will be at least practically unstable.

Hence, the phase at any point of a fluid mass, which is in stable equilibrium under the influence of gravity (whether this force is due to external bodies or to the mass itself), and which has throughout the same independently variable components, is completely determined by the phase at any other point and the difference of the values of the gravitation potential for the two points.

FUNDAMENTAL EQUATIONS OF IDEAL GASES AND GAS-MIXTURES.

For a constant quantity of a perfect or ideal gas, the product of the volume and pressure is proportional to the temperature, and the variations of energy are proportional to the variations of temperature. For a unit of such a gas we may write

$$\begin{aligned} p v &= a t, \\ d\varepsilon &= c dt, \end{aligned}$$

a and c denoting constants. By integration, we obtain the equation

$$\varepsilon = c t + E,$$

in which E also denotes a constant. If by these equations we eliminate t and p from (11), we obtain

$$d\varepsilon = \frac{\varepsilon - E}{c} d\eta - \frac{a}{v} \frac{\varepsilon - E}{c} dv,$$

or

$$c \frac{d\varepsilon}{\varepsilon - E} = d\eta - a \frac{dv}{v}.$$

The integral of this equation may be written in the form

$$c \log \frac{\varepsilon - E}{c} = \eta - a \log v - H,$$

where H denotes a fourth constant. We may regard E as denoting the energy of a unit of the gas for $t=0$; H its entropy for $t=1$ and $v=1$; a its pressure in the latter state, or its volume for $t=1$ and $p=1$; c its specific heat at constant volume. We may extend the application of the equation to any quantity of the gas, without altering the values of the constants, if we substitute $\frac{\varepsilon}{m}$, $\frac{\eta}{m}$, $\frac{v}{m}$ for ε , η , v , respectively. This will give

$$c \log \frac{\varepsilon - Em}{cm} = \frac{\eta}{m} - H + a \log \frac{m}{v}. \quad (255)$$

This is a fundamental equation (see pages 140–144) for an ideal gas of invariable composition. It will be observed that if we do not have to consider the properties of the matter which forms the gas as appearing in any other form or combination, but solely as constituting the gas in question (in a state of purity), we may without loss of generality give to E and H the value zero, or any other arbitrary values. But when the scope of our investigations is not thus limited, we may have determined the states of the substance of the gas for which $\varepsilon=0$ and $\eta=0$ with reference to some other form in which the substance appears, or, if the substance is compound, the states of its components for which $\varepsilon=0$ and $\eta=0$ may be already determined; so that the constants E and H cannot in general be treated as arbitrary.

We obtain from (255) by differentiation

$$\frac{c}{\varepsilon - Em} d\varepsilon = \frac{1}{m} d\eta - \frac{a}{v} dv + \left(\frac{cE}{\varepsilon - Em} + \frac{c+a}{m} - \frac{\eta}{m^2} \right) dm, \quad (256)$$

whence, in virtue of the general relation expressed by (86),

$$t = \frac{\varepsilon - Em}{cm}, \quad (257)$$

$$p = a \frac{\varepsilon - Em}{cv}, \quad (258)$$

$$\mu = E + \frac{\varepsilon - Em}{cm^2} (cm + am - \eta). \quad (259)$$

We may obtain the fundamental equation between ψ , t , v , and m from equations (87), (255), and (257). Eliminating ε we have

$$\psi = Em + cm t - t \eta,$$

and
$$c \log t = \frac{\eta}{m} - H + a \log \frac{m}{v};$$

and eliminating η , we have the fundamental equation

$$\psi = Em + m t \left(c - H - c \log t + a \log \frac{m}{v} \right). \quad (260)$$

Differentiating this equation, we obtain

$$d\psi = -m \left(H + c \log t + a \log \frac{v}{m} \right) dt - \frac{amt}{v} dv + \left(E + t \left(c + a - H - c \log t + a \log \frac{m}{v} \right) \right) dm; \quad (261)$$

whence, by the general equation (88),

$$\eta = m \left(H + c \log t + a \log \frac{v}{m} \right), \quad (262)$$

$$p = \frac{a m t}{v}, \quad (263)$$

$$\mu = E + t \left(c + a - H - c \log t + a \log \frac{m}{v} \right). \quad (264)$$

From (260), by (87) and (91), we obtain

$$\zeta = Em + m t \left(c - H - c \log t + a \log \frac{m}{v} \right) + p v,$$

and eliminating v by means of (263), we obtain the fundamental equation

$$\zeta = Em + m t \left(c + a - H - (c + a) \log t + a \log \frac{p}{a} \right). \quad (265)$$

From this, by differentiation and comparison with (92), we may obtain the equations

$$\eta = m \left(H + (c + a) \log t - a \log \frac{p}{a} \right), \quad (266)$$

$$v = \frac{a m t}{p}, \quad (267)$$

$$\mu = E + t \left(c + a - H - (c + a) \log t + a \log \frac{p}{a} \right). \quad (268)$$

The last is also a fundamental equation. It may be written in the form

$$\log \frac{p}{a} = \frac{H - c - a}{a} + \frac{c + a}{a} \log t + \frac{\mu - E}{a t}, \quad (269)$$

or, if we denote by e the base of the Naperian system of logarithms,

$$p = a e^{\frac{H - c - a}{a} + \frac{c + a}{a} \log t + \frac{\mu - E}{a t}} \quad (270)$$

The fundamental equation between χ , η , p , and m may also be easily obtained; it is

$$(c + a) \log \frac{\chi - Em}{(c + a)m} = \frac{\eta}{m} - H + a \log \frac{p}{a}, \quad (271)$$

which can be solved with respect to χ .

Any one of the fundamental equations (255), (260), (265), (270), and (271), which are entirely equivalent to one another, may be

regarded as defining an ideal gas. It will be observed that most of these equations might be abbreviated by the use of different constants. In (270), for example, a single constant might be used for

$$\alpha e^{\frac{H-c-a}{a}}, \text{ and another for } \frac{c+a}{a}.$$

The equations have been given in the above form, in order that the relations between the constants occurring in the different equations might be most clearly exhibited. The sum $c+a$ is the specific heat for constant pressure, as appears if we differentiate (266) regarding p and m as constant.*

* We may easily obtain the equation between the temperature and pressure of a saturated vapor, if we know the fundamental equations of the substance both in the gaseous, and in the liquid or solid state. If we suppose that the density and the specific heat at constant pressure of the liquid may be regarded as constant quantities (for such moderate pressures as the liquid experiences while in contact with the vapor), and denote this specific heat by k , and the volume of a unit of the liquid by V , we shall have for a unit of the liquid

$$t \, d\eta = k \, dt,$$

whence

$$\eta = k \log t + H',$$

where H' denotes a constant. Also, from this equation and (97),

$$d\mu = -(k \log t + H') \, dt + V \, dp,$$

whence

$$\mu = kt - kt \log t - H' t + Vp + E', \tag{A}$$

where E' denotes another constant. This is a fundamental equation for the substance in the liquid state. If (268) represents the fundamental equation for the same substance in the gaseous state, the two equations will both hold true of coexistent liquid and gas. Eliminating μ we obtain

$$\log \frac{p}{a} = \frac{H-H'+k-c-a}{a} - \frac{k-c-a}{a} \log t - \frac{E-E'}{at} + \frac{V}{a} \frac{p}{t}.$$

If we neglect the last term, which is evidently equal to the density of the vapor divided by the density of the liquid, we may write

$$\log p = A - B \log t - \frac{C}{t},$$

A , B , and C denoting constants. If we make similar suppositions in regard to the substance in the solid state, the equation between the pressure and temperature of coexistent solid and gaseous phases will of course have the same form.

A similar equation will also apply to the phases of an ideal gas which are coexistent with two different kinds of solids, one of which can be formed by the combination of the gas with the other, each being of invariable composition and of constant specific heat and density. In this case we may write for one solid

$$\mu_1 = k't - k't \log t - H't + V'p + E',$$

and for the other

$$\mu_2 = k''t - k''t \log t - H''t + V''p + E'',$$

and for the gas

$$\mu_3 = E + t \left(c+a-H - (c+a) \log t + a \log \frac{p}{a} \right).$$

The preceding fundamental equations all apply to gases of *constant composition*, for which the matter is entirely determined by a single

Now if a unit of the gas unites with the quantity λ of the first solid to form the quantity $1 + \lambda$ of the second it will be necessary for equilibrium (see pages 121, 122) that

$$\mu_3 + \lambda \mu_1 = (1 + \lambda) \mu_2.$$

Substituting the values of μ_1, μ_2, μ_3 , given above, we obtain after arranging the terms and dividing by at

$$\log \frac{p}{a} = A - B \log t - \frac{C}{t} + D \frac{p}{t},$$

when

$$A = \frac{H + \lambda H' - (1 + \lambda) H'' - c - a - \lambda k' + (1 + \lambda) k''}{a},$$

$$B = \frac{(1 + \lambda) k'' - \lambda k' - c - a}{a},$$

$$C = \frac{E + \lambda E' - (1 + \lambda) E''}{a}, \quad D = \frac{(1 + \lambda) V'' - \lambda V'}{a}.$$

We may conclude from this that an equation of the same form may be applied to an ideal gas in equilibrium with a liquid of which it forms an independently variable component, when the specific heat and density of the liquid are entirely determined by its composition, except that the letters A, B, C , and D must in this case be understood to denote quantities which vary with the composition of the liquid. But to consider the case more in detail, we have for the liquid by (A)

$$\frac{\zeta}{m} = \mu = kt - kt \log t - H't + V\rho + E',$$

where k, H', V, E' denote quantities which depend only upon the composition of the liquid. Hence, we may write

$$\zeta = \mathbf{k}t - \mathbf{k}t \log t - \mathbf{H}t + \mathbf{V}\rho + \mathbf{E},$$

where $\mathbf{k}, \mathbf{H}, \mathbf{V}$, and \mathbf{E} denote functions of m_1, m_2 , etc. (the quantities of the several components of the liquid). Hence, by (92),

$$\mu_1 = \frac{d\mathbf{k}}{dm_1} t - \frac{d\mathbf{k}}{dm_1} t \log t - \frac{d\mathbf{H}}{dm_1} t + \frac{d\mathbf{V}}{dm_1} \rho + \frac{d\mathbf{E}}{dm_1}.$$

If the component to which this potential relates is that which also forms the gas, we shall have by (269)

$$\log \frac{p}{a} = \frac{H - c - a}{a} + \frac{c + a}{a} \log t + \frac{\mu_1 - E}{at}.$$

Eliminating μ_1 , we obtain the equation

$$\log \frac{p}{a} = A - B \log t - \frac{C}{t} + D \frac{p}{t},$$

in which A, B, C , and D denote quantities which depend only upon the composition of the liquid, viz:

$$A = \frac{1}{a} \left(H - \frac{d\mathbf{H}}{dm_1} - c - a + \frac{d\mathbf{k}}{dm_1} \right),$$

$$B = \frac{1}{a} \left(\frac{d\mathbf{k}}{dm_1} - c - a \right),$$

variable (m). We may obtain corresponding fundamental equations for a mixture of gases, in which the proportion of the components shall be variable, from the following considerations.

It is a rule which admits of a very general and in many cases very exact experimental verification, that if several liquid or solid substances which yield different gases or vapors are simultaneously in equilibrium with a mixture of these gases (cases of chemical action between the gases being excluded,) the pressure in the gas-mixture is equal to the sum of the pressures of the gases yielded at the same temperature by the various liquid or solid substances taken separately. Now the potential in any of the liquids or solids for the substance which it yields in the form of gas has very nearly the same value when the liquid or solid is in equilibrium with the gas-mixture as when it is in equilibrium with its own gas alone. The difference of the pressure in the two cases will cause a certain difference in the values of the potential, but that this difference will be small, we may infer from the equation

$$\left(\frac{d\mu_1}{dp}\right)_{t, m} = \left(\frac{dv}{dm_1}\right)_{t, p, m}, \quad (272)$$

which may be derived from equation (92). In most cases, there will be a certain absorption by each liquid of the gases yielded by the others, but as it is well known that the above rule does not apply to cases in which such absorption takes place to any great extent, we may conclude that the effect of this circumstance in the cases with which we have to do is of secondary importance. If we neglect the slight differences in the values of the potentials due to these circumstances, the rule may be expressed as follows:

The pressure in a mixture of different gases is equal to the sum of the pressures of the different gases as existing each by itself at the same temperature and with the same value of its potential.

To form a precise idea of the practical significance of the law as thus stated with reference to the equilibrium of two liquids with a mixture of the gases which they emit, when neither liquid absorbs the gas emitted by the other, we may imagine a long tube closed at each end and bent in the form of a W to contain in each of the descending

$$C = \frac{1}{a} \left(E - \frac{d\mathbf{E}}{dm_1} \right) \quad D = \frac{1}{a} \frac{d\mathbf{V}}{dm_1}$$

With respect to some of the equations which have here been deduced, the reader may compare Professor Kirchhoff "Ueber die Spannung des Dampfes von Mischungen aus Wasser und Schwefelsäure," Pogg. Ann., vol. civ. (1858), p. 612; and Dr. Rankine "On Saturated Vapors," Phil. Mag., vol. xxxi. (1866), p. 199.

loops one of the liquids, and above these liquids the gases which they emit, viz., the separate gases at the ends of the tube, and the mixed gases in the middle. We may suppose the whole to be in equilibrium, the difference of the pressures of the gases being balanced by the proper heights of the liquid columns. Now it is evident from the principles established on pages 203–210 that the potential for either gas will have the same value in the mixed and in the separate gas *at the same level*, and therefore according to the rule in the form which we have given, the pressure in the gas-mixture is equal to the sum of the pressures in the separate gases, *all these pressures being measured at the same level*. Now the experiments by which the rule has been established relate rather to the gases in the vicinity of the surfaces of the liquids. Yet, although the differences of level in these surfaces may be considerable, the corresponding differences of pressure in the columns of gas will certainly be very small in all cases which can be regarded as falling under the laws of ideal gases, for which very great pressures are not admitted.

If we apply the above law to a mixture of ideal gases and distinguish by subscript numerals the quantities relating to the different gases, and denote by Σ_1 the sum of all similar terms obtained by changing the subscript numerals, we shall have by (270)

$$p = \Sigma_1 \left(a_1 e^{\frac{H_1 - c_1 - a_1}{a_1}} t^{\frac{c_1 + a_1}{a_1}} e^{\frac{\mu_1 - E_1}{a_1 t}} \right), \quad (273)$$

It will be legitimate to assume this equation provisionally as the fundamental equation defining an ideal gas-mixture, and afterwards to justify the suitability of such a definition by the properties which may be deduced from it. In particular, it will be necessary to show that an ideal gas-mixture as thus defined, when the proportion of its components remains constant, has all the properties which have already been assumed for an ideal gas of invariable composition; it will also be desirable to consider more rigorously and more in detail the equilibrium of such a gas-mixture with solids and liquids, with respect to the above rule.

By differentiation and comparison with (98) we obtain

$$\frac{\eta}{v} = \Sigma_1 \left((c_1 + a_1 - \frac{\mu_1 - E_1}{t}) e^{\frac{H_1 - c_1 - a_1}{a_1}} t^{\frac{c_1 + a_1}{a_1}} e^{\frac{\mu_1 - E_1}{a_1 t}} \right), \quad (274)$$

$$\left. \begin{aligned} \frac{m_1}{v} &= e^{\frac{H_1 - c_1 - a_1}{a_1} t} \frac{c_1}{a_1} e^{\frac{\mu_1 - E_1}{a_1 t}}, \\ \frac{m_2}{v} &= e^{\frac{H_2 - c_2 - a_2}{a_2} t} \frac{c_2}{a_2} e^{\frac{\mu_2 - E_2}{a_2 t}}, \\ \text{etc.} \end{aligned} \right\} \quad (275)$$

Equations (275) indicate that the relation between the temperature, the density of any component, and the potential for that component, is not affected by the presence of the other components. They may also be written

$$\left. \begin{aligned} \mu_1 &= E_1 + t \left(c_1 + a_1 - H_1 - c_1 \log t + a_1 \log \frac{m_1}{v} \right), \\ \text{etc.} \end{aligned} \right\} \quad (276)$$

Eliminating $\mu_1, \mu_2, \text{etc.}$ from (273) and (274) by means of (275) and (276), we obtain

$$p = \sum_1 \frac{a_1 m_1 t}{v}, \quad (277)$$

$$\eta = \sum_1 \left(m_1 H_1 + m_1 c_1 \log t + m_1 a_1 \log \frac{v}{m_1} \right). \quad (278)$$

Equation (277) expresses the familiar principle that the pressure in a gas-mixture is equal to the sum of the pressures which the component gases would possess if existing separately with the same volume at the same temperature. Equation (278) expresses a similar principle in regard to the entropy of the gas-mixture.

From (276) and (277) we may easily obtain the fundamental equation between $\psi, t, v, m_1, m_2, \text{etc.}$ For by substituting in (94) the values of $p, \mu_1, \mu_2, \text{etc.}$ taken from these equations, we obtain

$$\psi = \sum_1 \left(E_1 m_1 + m_1 t \left(c_1 - H_1 - c_1 \log t + a_1 \log \frac{m_1}{v} \right) \right). \quad (279)$$

If we regard the proportion of the various components as constant, this equation may be simplified by writing

$$\begin{aligned} m &\text{ for } \sum_1 m_1, \\ c m &\text{ for } \sum_1 (c_1 m_1), \\ a m &\text{ for } \sum_1 (a_1 m_1), \\ E m &\text{ for } \sum_1 (E_1 m_1), \end{aligned}$$

and $H m - a m \log m$ for $\sum_1 (H_1 m_1 - a_1 m_1 \log m_1)$.

The values of $c, a, E,$ and $H,$ will then be constant and m will denote the total quantity of gas. As the equation will thus be reduced to the

form of (260), it is evident that an ideal gas-mixture, as defined by (273) or (279), when the proportion of its components remains unchanged, will have all the properties which we have assumed for an ideal gas of invariable composition. The relations between the specific heats of the gas-mixture at constant volume and at constant pressure and the specific heats of its components are expressed by the equations

$$c = \sum_1 \frac{m_1 c_1}{m}, \quad (280)$$

and

$$c + a = \sum_1 \frac{m_1 (c_1 + a_1)}{m}. \quad (281)$$

We have already seen that the values of t , v , m_1 , μ_1 in a gas-mixture are such as are possible for the component G_1 (to which m_1 and μ_1 relate) existing separately. If we denote by p_1 , η_1 , ψ_1 , ε_1 , χ_1 , ζ_1 the connected values of the several quantities which the letters indicate determined for the gas G_1 as thus existing separately, and extend this notation to the other components, we shall have by (273), (274), and (279)

$$p = \sum_1 p_1, \quad \eta = \sum_1 \eta_1, \quad \psi = \sum_1 \psi_1; \quad (282)$$

whence by (87), (89), and (91)

$$\varepsilon = \sum_1 \varepsilon_1, \quad \chi = \sum_1 \chi_1, \quad \zeta = \sum_1 \zeta_1. \quad (283)$$

The quantities p , η , ψ , ε , χ , ζ relating to the gas-mixture may therefore be regarded as consisting of parts which may be attributed to the several components in such a manner that between the parts of these quantities which are assigned to any component, the quantity of that component, the potential for that component, the temperature, and the volume, the same relations shall subsist as if that component existed separately. It is in this sense that we should understand the law of Dalton, that every gas is as a vacuum to every other gas.

It is to be remarked that these relations are consistent and possible for a mixture of gases which are not ideal gases, and indeed without any limitation in regard to the thermodynamic properties of the individual gases. They are all consequences of the law that the pressure in a mixture of different gases is equal to the sum of the pressures of the different gases as existing each by itself at the same temperature and with the same value of its potential. For let p_1 , η_1 , ε_1 , ψ_1 , χ_1 , ζ_1 ; p_2 , etc.; etc. be defined as relating to the different gases existing each by itself with the same volume, temperature, and potential as in the gas-mixture; if

$$p = \sum_1 p_1,$$

then
$$\left(\frac{d\rho}{d\mu_1}\right)_{t, \mu_2 \dots \mu_n} = \left(\frac{d\rho_1}{d\mu_1}\right)_t;$$

and therefore, by (98), the quantity of any component gas G_1 in the gas-mixture, and in the separate gas to which p_1, η_1 , etc. relate, is the same and may be denoted by the same symbol m_1 . Also

$$\eta = v \left(\frac{d\rho}{dt}\right)_{\mu_1, \dots, \mu_n} = v \sum_1 \left(\frac{d\rho}{dt}\right)_{\mu_1} = \sum_1 \eta_1;$$

whence also, by (93)-(96),

$$\varepsilon = \sum_1 \varepsilon_1, \quad \psi = \sum_1 \psi_1, \quad \chi = \sum_1 \chi_1, \quad \zeta = \sum_1 \zeta_1.$$

All the same relations will also hold true whenever the value of ψ for the gas-mixture is equal to the sum of the values of this function for the several component gases existing each by itself in the same quantity as in the gas-mixture and with the temperature and volume of the gas-mixture. For if $p_1, \eta_1, \varepsilon_1, \psi_1, \chi_1, \zeta_1; p_2$, etc.; etc. are defined as relating to the components existing thus by themselves, we shall have

$$\psi = \sum_1 \psi_1,$$

whence

$$\left(\frac{d\psi}{dm_1}\right)_{t, v, m} = \left(\frac{d\psi_1}{dm_1}\right)_{t, v}.*$$

Therefore, by (88), the potential μ_1 has the same value in the gas-mixture and in the gas G_1 existing separately as supposed. Moreover,

$$\eta = - \left(\frac{d\psi}{dt}\right)_{v, m} = - \sum_1 \left(\frac{d\psi_1}{dt}\right)_{v, m} = \sum_1 \eta_1,$$

and

$$p = - \left(\frac{d\psi}{dv}\right)_{t, m} = - \sum_1 \left(\frac{d\psi_1}{dv}\right)_{t, m} = \sum_1 p_1,$$

whence

$$\varepsilon = \sum_1 \varepsilon_1, \quad \chi = \sum_1 \chi_1, \quad \zeta = \sum_1 \zeta_1.$$

Whenever different bodies are combined without communication of work or heat between them and external bodies, the energy of the body formed by the combination is necessarily equal to the sum of the energies of the bodies combined. In the case of ideal gas-mixtures, when the initial temperatures of the gas-masses which are combined

* A subscript m after a differential coefficient relating to a body having several independently variable components is used here and elsewhere in this paper to indicate that each of the quantities m_1, m_2 , etc., unless its differential occurs in the expression to which the suffix is applied, is to be regarded as constant in the differentiation.

are the same, (whether these gas-masses are entirely different gases, or gas-mixtures differing only in the proportion of their components,) the condition just mentioned can only be satisfied when the temperature of the resultant gas-mixture is also the same. In such combinations, therefore, the final temperature will be the same as the initial.

If we consider a vertical column of an ideal gas-mixture which is in equilibrium, and denote the densities of one of its components at two different points by γ_1 and γ_1' , we shall have by (275) and (234)

$$\frac{\gamma_1}{\gamma_1'} = e^{\frac{\mu_1 - \mu_1'}{a_1 t}} = e^{\frac{g(h' - h)}{a_1 t}}. \quad (284)$$

From this equation, in which we may regard the quantities distinguished by accents as constant, it appears that the relation between the density of any one of the components and the height is not affected by the presence of the other components.

The work obtained or expended in any reversible process of combination or separation of ideal gas-mixtures at constant temperature, or when the temperatures of the initial and final gas-masses and of the only external source of heat or cold which is used are all the same, will be found by taking the difference of the sums of the values of ψ for the initial, and for the final gas-masses. (See pages 145, 146). It is evident from the form of equation (279) that this work is equal to the sum of the quantities of work which would be obtained or expended in producing in each different component existing separately the same changes of density which that component experiences in the actual process for which the work is sought.*

We will now return to the consideration of the equilibrium of a liquid with the gas which it emits as affected by the presence of different gases, when the gaseous mass in contact with the liquid may be regarded as an ideal gas-mixture.

It may first be observed, that the density of the gas which is emitted by the liquid will not be affected by the presence of other gases which are not absorbed by the liquid, when the liquid is protected in any way from the pressure due to these additional gases. This may be accomplished by separating the liquid and gaseous

* This result has been given by Lord Rayleigh, (Phil. Mag., vol. xlix, 1875, p. 311). It will be observed that equation (279) might be deduced immediately from this principle in connection with equation (260) which expresses the properties ordinarily assumed for perfect gases.

masses by a diaphragm which is permeable to the liquid. It will then be easy to maintain the liquid at any constant pressure which is not greater than that in the gas. The potential in the liquid for the substance which it yields as gas will then remain constant, and therefore the potential for the same substance in the gas and the density of this substance in the gas and the part of the gaseous pressure due to it will not be affected by the other components of the gas.

But when the gas and liquid meet under ordinary circumstances, i. e., in a free plane surface, the pressure in both is necessarily the same, as also the value of the potential for any common component S_1 . Let us suppose the density of an insoluble component of the gas to vary, while the composition of the liquid and the temperature remain unchanged. If we denote the increments of pressure and of the potential for S_1 by dp and $d\mu_1$, we shall have by (272)

$$d\mu_1 = \left(\frac{d\mu_1}{dp}\right)_{t, m}^{(L)} dp = \left(\frac{dv}{dm_1}\right)_{t, p, m}^{(L)} dp,$$

the index (L) denoting that the expressions to which it is affixed refer to the liquid. (Expressions without such an index will refer to the gas alone or to the gas and liquid in common.) Again, since the gas is an ideal gas-mixture, the relation between p_1 and μ_1 is the same as if the component S_1 existed by itself at the same temperature, and therefore by (268)

$$d\mu_1 = a_1 t d \log p_1.$$

Therefore

$$a_1 t d \log p_1 = \left(\frac{dv}{dm_1}\right)_{t, p, m}^{(L)} dp. \quad (265)$$

This may be integrated at once if we regard the differential coefficient in the second member as constant, which will be a very close approximation. We may obtain a result more simple, but not quite so accurate, if we write the equation in the form

$$dp_1 = \gamma_1 \left(\frac{dv}{dm_1}\right)_{t, p, m}^{(L)} dp, \quad (266)$$

where γ_1 denotes the density of the component S_1 in the gas, and integrate regarding this quantity also as constant. This will give

$$p_1 - p_1' = \gamma_1 \left(\frac{dv}{dm_1}\right)_{t, p, m}^{(L)} (p - p'). \quad (267)$$

where p_1' and p' denote the values of p_1 and p when the insoluble component of the gas is entirely wanting. It will be observed that $p - p'$ is nearly equal to the pressure of the insoluble component, in the phase of the gas-mixture to which p_1 relates. S_1 is not neces-

sarily the only common component of the gas and liquid. If there are others, we may find the increase of the part of the pressure in the gas-mixture belonging to any one of them by equations differing from the last only in the subscript numerals.

Let us next consider the effect of a gas which is absorbed to some extent, and which must therefore in strictness be regarded as a component of the liquid. We may commence by considering in general the equilibrium of a gas-mixture of two components S_1 and S_2 with a liquid formed of the same components. Using a notation like the previous, we shall have by (98) for constant temperature,

$$dp = \gamma_1 d\mu_1 + \gamma_2 d\mu_2,$$

and

$$dp = \gamma_1^{(L)} d\mu_1 + \gamma_2^{(L)} d\mu_2;$$

whence

$$(\gamma_1^{(L)} - \gamma_1) d\mu_1 = (\gamma_2 - \gamma_2^{(L)}) d\mu_2.$$

Now if the gas is an ideal gas-mixture,

$$d\mu_1 = \frac{a_1 t}{p_1} dp_1 = \frac{dp_1}{\gamma_1}, \quad \text{and} \quad d\mu_2 = \frac{a_2 t}{p_2} dp_2 = \frac{dp_2}{\gamma_2},$$

therefore

$$\left(\frac{\gamma_1^{(L)}}{\gamma_1} - 1 \right) dp_1 = \left(1 - \frac{\gamma_2^{(L)}}{\gamma_2} \right) dp_2. \quad (288)$$

We may now suppose that S_1 is the principal component of the liquid, and S_2 is a gas which is absorbed in the liquid to a slight extent. In such cases it is well known that the ratio of the densities of the substance S_2 in the liquid and in the gas is for a given temperature approximately constant. If we denote this constant by A , we shall have

$$\left(\frac{\gamma_1^{(L)}}{\gamma_1} - 1 \right) dp_1 = (1 - A) dp_2. \quad (289)$$

It would be easy to integrate this equation regarding γ_1 as variable, but as the variation in the value of p_1 is necessarily very small we shall obtain sufficient accuracy if we regard γ_1 as well as $\gamma_1^{(L)}$ as constant. We shall thus obtain

$$\left(\frac{\gamma_1^{(L)}}{\gamma_1} - 1 \right) (p_1 - p_1') = (1 - A) p_2, \quad (290)$$

where p_1' denotes the pressure of the saturated vapor of the pure liquid consisting of S_1 . It will be observed that when $A = 1$, the presence of the gas S_2 will not affect the pressure or density of the gas S_1 . When $A < 1$, the pressure and density of the gas S_1 are greater than if S_2 were absent, and when $A > 1$, the reverse is true.

The properties of an ideal gas-mixture (according to the definition which we have assumed) when in equilibrium with liquids or solids have been developed at length, because it is only in respect to these properties that there is any variation from the properties usually attributed to perfect gases. As the pressure of a gas saturated with vapor is usually given as a little less than the sum of the pressure of the gas calculated from its density and that of saturated vapor in a space otherwise empty, while our formulæ would make it a little more, when the gas is insoluble, it would appear that in this respect our formulæ are less accurate than the rule which would make the pressure of the gas saturated with vapor equal to the sum of the two pressures mentioned. Yet the reader will observe that the magnitude of the quantities concerned is not such that any stress can be laid upon this circumstance.

It will also be observed that the statement of Dalton's law which we have adopted, while it serves to complete the theory of gas-mixtures (with respect to a certain class of properties), asserts nothing with reference to any solid or liquid bodies. But the common rule that the density of a gas necessary for equilibrium with a solid or liquid is not altered by the presence of a different gas which is not absorbed by the solid or liquid, if construed *strictly*, will involve consequences in regard to solids and liquids which are entirely inadmissible. To show this, we will assume the correctness of the rule mentioned. Let S_1 denote the common component of the gaseous and liquid or solid masses, and S_2 the insoluble gas, and let quantities relating to the gaseous mass be distinguished when necessary by the index (G), and those relating to the liquid or solid by the index (L). Now while the gas is in equilibrium with the liquid or solid, let the quantity which it contains of S_2 receive the increment dm_2 , its volume and the quantity which it contains of the other component, as well as the temperature, remaining constant. The potential for S_1 in the gaseous mass will receive the increment

$$\left(\frac{d\mu_1}{dm_2}\right)_{t, v, m}^{(G)} dm_2$$

and the pressure will receive the increment

$$\left(\frac{dp}{dm_2}\right)_{t, v, m}^{(G)} dm_2.$$

Now the liquid or solid remaining in equilibrium with the gas must experience the same variations in the values of μ_1 and p . But by (272)

$$\left(\frac{d\mu_1}{dp}\right)_{t, m}^{(L)} = \left(\frac{dr}{dm_1}\right)_{t, p, m}^{(L)}.$$

Therefore,

$$\left(\frac{dv}{dm_1}\right)_{t, p, m}^{(L)} = \frac{\left(\frac{d\mu_1}{dm_2}\right)_{t, v, m}^{(G)}}{\left(\frac{dp}{dm_2}\right)_{t, v, m}^{(G)}}.$$

It will be observed that the first member of this equation relates solely to the liquid or solid, and the second member solely to the gas. Now we may suppose the same gaseous mass to be capable of equilibrium with several different liquids or solids, and the first member of this equation must therefore have the same value for all such liquids or solids; which is quite inadmissible. In the simplest case, in which the liquid or solid is identical in substance with the vapor which it yields, it is evident that the expression in question denotes the reciprocal of the density of the solid or liquid. Hence, when a gas is in equilibrium with one of its components both in the solid and liquid states (as when a moist gas is in equilibrium with ice and water), it would be necessary that the solid and liquid should have the same density.

The foregoing considerations appear sufficient to justify the definition of an ideal gas-mixture which we have chosen. It is of course immaterial whether we regard the definition as expressed by equation (273), or by (279), or by any other fundamental equation which can be derived from these.

The fundamental equations for an ideal gas-mixture corresponding to (255), (265), and (271) may easily be derived from these equations by using inversely the substitutions given on page 217. They are

$$\sum_1 (c_1 m_1) \log \frac{\varepsilon - \sum_1 (E_1 m_1)}{\sum_1 (c_1 m_1)} = \eta + \sum_1 (a_1 m_1 \log \frac{m_1}{v} - H_1 m_1), \quad (291)$$

$$\begin{aligned} \sum_1 (c_1 m_1 + a_1 m_1) \log \frac{\chi - \sum_1 (E_1 m_1)}{\sum_1 (c_1 m_1 + a_1 m_1)} \\ = \eta + \sum_1 \left(a_1 m_1 \log \frac{p m_1}{\sum_1 (a_1 m_1)} - H_1 m_1 \right), \end{aligned} \quad (292)$$

$$\begin{aligned} \zeta = \sum_1 \left(E_1 m_1 + m_1 t (c_1 + a_1 - H_1) \right) \\ - \sum_1 (c_1 m_1 + a_1 m_1) t \log t + \sum_1 \left(a_1 m_1 t \log \frac{p m_1}{\sum_1 (a_1 m_1)} \right). \end{aligned} \quad (293)$$

The components to which the fundamental equations (273), (279), (291), (292), 293) refer, may themselves be gas-mixtures. We may for example apply the fundamental equations of a binary gas-mixture

to a mixture of hydrogen and air, or to any ternary gas-mixture in which the proportion of two of the components is fixed. In fact, the form of equation (279) which applies to a gas-mixture of any particular number of components may easily be reduced, when the proportions of some of these components are fixed, to the form which applies to a gas-mixture of a smaller number of components. The necessary substitutions will be analogous to those given on page 217. But the components must be entirely different from one another with respect to the gases of which they are formed by mixture. We cannot, for example, apply equation (279) to a gas-mixture in which the components are oxygen and air. It would indeed be easy to form a fundamental equation for such a gas-mixture with reference to the designated gases as components. Such an equation might be derived from (279) by the proper substitutions. But the result would be an equation of more complexity than (279). A *chemical* compound, however, with respect to Dalton's law, and with respect to all the equations which have been given, is to be regarded as entirely different from its components. Thus, a mixture of hydrogen, oxygen, and vapor of water is to be regarded as a ternary gas-mixture, having the three components mentioned. This is certainly true when the quantities of the compound gas and of its components are all independently variable in the gas-mixture, without change of temperature or pressure. Cases in which these quantities are not thus independently variable will be considered hereafter.

Inferences in regard to Potentials in Liquids and Solids.

Such equations as (264), (268), (276), by which the values of potentials in pure or mixed gases may be derived from quantities capable of direct measurement, have an interest which is not confined to the theory of gases. For as the potentials of the independently variable components which are common to coexistent liquid and gaseous masses have the same values in each, these expressions will generally afford the means of determining for liquids, at least approximately, the potential for any independently variable component which is capable of existing in the gaseous state. For although every state of a liquid is not such as can exist in contact with a gaseous mass, it will always be possible, when any of the components of the liquid are volatile, to bring it by a change of pressure alone, its temperature and composition remaining unchanged, to a state for which there is a coexistent phase of vapor, in which

the values of the potentials of the volatile components of the liquid may be estimated from the density of these substances in the vapor. The variations of the potentials in the liquid due to the change of pressure will in general be quite trifling as compared with the variations which are connected with changes of temperature or of composition, and may moreover be readily estimated by means of equation (272). The same considerations will apply to volatile solids with respect to the determination of the potential for the substance of the solid.

As an application of this method of determining the potentials in liquids, let us make use of the law of Henry in regard to the absorption of gases by liquids to determine the relation between the quantity of the gas contained in any liquid mass and its potential. Let us consider the liquid as in equilibrium with the gas, and let $m_1^{(G)}$ denote the quantity of the gas existing as such, $m_1^{(L)}$ the quantity of the same substance contained in the liquid mass, μ_1 the potential for this substance common to the gas and liquid, $v^{(G)}$ and $v^{(L)}$ the volumes of the gas and liquid. When the absorbed gas forms but a very small part of the liquid mass, we have by Henry's law

$$\frac{m_1^{(L)}}{v^{(L)}} = A \frac{m_1^{(G)}}{v^{(G)}}, \quad (294)$$

where A is a function of the temperature; and by (276)

$$\mu_1 = B + C \log \frac{m_1^{(G)}}{v^{(G)}}, \quad (295)$$

B and C also denoting functions of the temperature. Therefore

$$\mu_1 = B + C \log \frac{m_1^{(L)}}{A v^{(L)}}. \quad (296)$$

It will be seen (if we disregard the difference of notation) that this equation is equivalent in form to (216), which was deduced from *a priori* considerations as a probable relation between the quantity and the potential of a small component. When a liquid absorbs several gases at once, there will be several equations of the form of (296), which will hold true simultaneously, and which we may regard as equivalent to equations (217), (218). The quantities A and C in (216), with the corresponding quantities in (217), (218), were regarded as functions of the temperature and pressure, but since the potentials in liquids are but little affected by the pressure, we might anticipate that these quantities in the case of liquids might be regarded as functions of the temperature alone.

In regard to equations (216), (217), (218), we may now observe that by (264) and (276) they are shown to hold true in ideal gases or gas-mixtures, not only for components which form only a small part of the whole gas-mixture, but without any such limitation, and not only approximately but absolutely. It is noticeable that in this case quantities A and C are functions of the temperature alone, and do not even depend upon the nature of the gaseous mass, except upon the particular component to which they relate. As all gaseous bodies are generally supposed to approximate to the laws of ideal gases when sufficiently rarefied, we may regard these equations as approximately valid for gaseous bodies in general when the density is sufficiently small. When the density of the gaseous mass is very great, but the separate density of the component in question is small, the equations will probably hold true, but the values of A and C may not be entirely independent of the pressure, or of the composition of the mass in respect to its principal components. These equations will also apply, as we have just seen, to the potentials in liquid bodies for components of which the density in the liquid is very small, whenever these components exist also in the gaseous state, and conform to the law of Henry. This seems to indicate that the law expressed by these equations has a very general application.

Considerations relating to the Increase of Entropy due to the Mixture of Gases by Diffusion.

From equation (278) we may easily calculate the increase of entropy which takes place when two different gases are mixed by diffusion, at a constant temperature and pressure. Let us suppose that the quantities of the gases are such that each occupies initially one half of the total volume. If we denote this volume by V , the increase of entropy will be

$$m_1 a_1 \log V + m_2 a_2 \log V - m_1 a_1 \log \frac{V}{2} - m_2 a_2 \log \frac{V}{2},$$

or $(m_1 a_1 + m_2 a_2) \log 2.$

Now $m_1 a_1 = \frac{pV}{2t},$ and $m_2 a_2 = \frac{pV}{2t}.$

Therefore the increase of entropy may be represented by the expression

$$\frac{pV}{t} \log 2. \tag{297}$$

It is noticeable that the value of this expression does not depend upon the kinds of gas which are concerned, if the quantities are such as has been supposed, except that the gases which are mixed must be

of different kinds. If we should bring into contact two masses of the same kind of gas, they would also mix, but there would be no increase of entropy. But in regard to the relation which this case bears to the preceding, we must bear in mind the following considerations. When we say that when two different gases mix by diffusion, as we have supposed, the energy of the whole remains constant, and the entropy receives a certain increase, we mean that the gases could be separated and brought to the same volume and temperature which they had at first by means of certain changes in external bodies, for example, by the passage of a certain amount of heat from a warmer to a colder body. But when we say that when two gas-masses of the same kind are mixed under similar circumstances there is no change of energy or entropy, we do not mean that the gases which have been mixed can be separated without change to external bodies. On the contrary, the separation of the gases is entirely impossible. We call the energy and entropy of the gas-masses when mixed the same as when they were unmixed, because we do not recognize any difference in the substance of the two masses. So when gases of different kinds are mixed, if we ask what changes in external bodies are necessary to bring the system to its original state, we do not mean a state in which each particle shall occupy more or less exactly the same position as at some previous epoch, but only a state which shall be undistinguishable from the previous one in its sensible properties. It is to states of systems thus incompletely defined that the problems of thermodynamics relate.

But if such considerations explain why the mixture of gas-masses of the same kind stands on a different footing from the mixture of gas-masses of different kinds, the fact is not less significant that the increase of entropy due to the mixture of gases of different kinds, in such a case as we have supposed, is independent of the nature of the gases.

Now we may without violence to the general laws of gases which are embodied in our equations suppose other gases to exist than such as actually do exist, and there does not appear to be any limit to the resemblance which there might be between two such kinds of gas. But the increase of entropy due to the mixing of given volumes of the gases at a given temperature and pressure would be independent of the degree of similarity or dissimilarity between them. We might also imagine the case of two gases which should be absolutely identical in all the properties (sensible and molecular) which come into play while they exist as gases either pure or mixed with each other,

but which should differ in respect to the attractions between their atoms and the atoms of some other substances, and therefore in their tendency to combine with such substances. In the mixture of such gases by diffusion an increase of entropy would take place, although the process of mixture, dynamically considered, might be absolutely identical in its minutest details (even with respect to the precise path of each atom) with processes which might take place without any increase of entropy. In such respects, entropy stands strongly contrasted with energy. Again, when such gases have been mixed, there is no more impossibility of the separation of the two kinds of molecules in virtue of their ordinary motions in the gaseous mass without any special external influence, than there is of the separation of a homogeneous gas into the same two parts into which it has once been divided, after these have once been mixed. In other words, the impossibility of an uncompensated decrease of entropy seems to be reduced to improbability.

There is perhaps no fact in the molecular theory of gases so well established as that the number of molecules in a given volume at a given temperature and pressure is the same for every kind of gas when in a state to which the laws of ideal gases apply. Hence the quantity $\frac{pV}{t}$ in (297) must be entirely determined by the number of molecules which are mixed. And the increase of entropy is therefore determined by the number of these molecules and is independent of their dynamical condition and of the degree of difference between them.

The result is of the same nature when the volumes of the gases which are mixed are not equal, and when more than two kinds of gas are mixed. If we denote by v_1, v_2 , etc., the initial volumes of the different kinds of gas, and by V as before the total volume, the increase of entropy may be written in the form

$$\sum_1 (m_1 a_1) \log V - \sum_1 (m_1 a_1 \log v_1).$$

And if we denote by r_1, r_2 , etc., the numbers of the molecules of the several different kinds of gas, we shall have

$$r_1 = C m_1 a_1, \quad r_2 = C m_2 a_2, \quad \text{etc.},$$

where C denotes a constant. Hence

$$v_1 : V :: m_1 a_1 : \sum_1 (m_1 a_1) :: r_1 : \sum_1 r_1;$$

and the increase of entropy may be written

$$\frac{\sum_1 r_1 \log \sum_1 r_1 - \sum_1 (r_1 \log r_1)}{C}. \quad (298)$$

The Phases of Dissipated Energy of an Ideal Gas-mixture with Components which are Chemically Related.

We will now pass to the consideration of the phases of dissipated energy (see page 200) of an ideal gas-mixture, in which the number of the proximate components exceeds that of the ultimate.

Let us first suppose that an ideal gas-mixture has for proximate components the gases G_1 , G_2 , and G_3 , the units of which are denoted by \mathfrak{G}_1 , \mathfrak{G}_2 , \mathfrak{G}_3 , and that in ultimate analysis

$$\mathfrak{G}_3 = \lambda_1 \mathfrak{G}_1 + \lambda_2 \mathfrak{G}_2, \quad (299)$$

λ_1 and λ_2 denoting positive constants, such that $\lambda_1 + \lambda_2 = 1$. The phases which we are to consider are those for which the energy of the gas-mixture is a minimum for constant entropy and volume and constant quantities of G_1 and G_2 , as determined in ultimate analysis. For such phases, by (86),

$$\mu_1 \delta m_1 + \mu_2 \delta m_2 + \mu_3 \delta m_3 \geq 0. \quad (300)$$

for such values of the variations as do not affect the quantities of G_1 and G_2 as determined in ultimate analysis. Values of δm_1 , δm_2 , δm_3 proportional to λ_1 , λ_2 , -1 , and only such, are evidently consistent with this restriction: therefore

$$\lambda_1 \mu_1 + \lambda_2 \mu_2 = \mu_3. \quad (301)$$

If we substitute in this equation values of μ_1 , μ_2 , μ_3 taken from (276), we obtain, after arranging the terms and dividing by t ,

$$\lambda_1 a_1 \log \frac{m_1}{v} + \lambda_2 a_2 \log \frac{m_2}{v} - a_3 \log \frac{m_3}{v} = A + B \log t - \frac{C}{t}, \quad (302)$$

where

$$A = \lambda_1 H_1 + \lambda_2 H_2 - H_3 - \lambda_1 c_1 - \lambda_2 c_2 + c_3 - \lambda_1 a_1 - \lambda_2 a_2 + a_3, \quad (303)$$

$$B = \lambda_1 c_1 + \lambda_2 c_2 - c_3, \quad (304)$$

$$C = \lambda_1 E_1 + \lambda_2 E_2 - E_3. \quad (305)$$

If we denote by β_1 and β_2 the volumes (determined under standard conditions of temperature and pressure) of the quantities of the gases G_1 and G_2 which are contained in a unit of volume of the gas G_3 , we shall have

$$\beta_1 = \frac{\lambda_1 a_1}{a_3}, \text{ and } \beta_2 = \frac{\lambda_2 a_2}{a_3}, \quad (306)$$

and (302) will reduce to the form

$$\log \frac{m_1^{\beta_1} m_2^{\beta_2}}{m_3 v^{\beta_1 + \beta_2 - 1}} = \frac{A}{a_3} + \frac{B}{a_3} \log t - \frac{C}{a_3 t}. \quad (307)$$

Moreover, as by (277)

$$p v = (a_1 m_1 + a_2 m_2 + a_3 m_3) t, \quad (308)$$

we have on eliminating v

$$\log \frac{m_1^{\beta_1} m_2^{\beta_2} p^{\beta_1 + \beta_2 - 1}}{m_3 (a_1 m_1 + a_2 m_2 + a_3 m_3)^{\beta_1 + \beta_2 - 1}} = \frac{A}{a_3} + \frac{B'}{a_3} \log t - \frac{C}{a_3 t}, \quad (309)$$

where

$$B' = \lambda_1 c_1 + \lambda_2 c_2 - c_3 + \lambda_1 a_1 + \lambda_2 a_2 - a_3. \quad (310)$$

It will be observed that the quantities β_1, β_2 will always be positive and have a simple relation to unity, and that the value of $\beta_1 + \beta_2 - 1$ will be positive or zero, according as gas G_3 is formed of G_1 and G_2 with or without condensation. If we should assume, according to the rule often given for the specific heat of compound gases, that the thermal capacity at constant volume of any quantity of the gas G_3 is equal to the sum of the thermal capacities of the quantities which it contains of the gases G_1 and G_2 , the value of B would be zero. The heat evolved in the formation of a unit of the gas G_3 out of the gases G_1 and G_2 , without mechanical action, is by (283) and (257)

$$\lambda_1 (c_1 t + E_1) + \lambda_2 (c_2 t + E_2) - (c_3 t + E_3),$$

or

$$B t + C,$$

which will reduce to C when the above relation in regard to the specific heats is satisfied. In any case the quantity of heat thus evolved divided by $a_3 t^2$ will be equal to the differential coefficient of the second member of equation (307) with respect to t . Moreover, the heat evolved in the formation of a unit of the gas G_3 out of the gases G_1 and G_2 under constant pressure is

$$B t + C + \lambda_1 a_1 t + \lambda_2 a_2 t - a_3 t = B' t + C,$$

which is equal to the differential coefficient of the second member of (309) with respect to t , multiplied by $a_3 t^2$.

It appears by (307) that, except in the case when $\beta_1 + \beta_2 = 1$, for any given finite values of m_1, m_2, m_3 , and t (infinitesimal values being excluded as well as infinite), it will always be possible to assign such a finite value to v that the mixture shall be in a state of dissipated energy. Thus, if we regard a mixture of hydrogen, oxygen, and vapor of water as an ideal gas-mixture, for a mixture containing any given quantities of these three gases at any given tem-

perature there will be a certain volume at which the mixture will be in a state of dissipated energy. In such a state no such phenomenon as explosion will be possible, and no formation of water by the action of platinum. (If the mass should be expanded beyond this volume, the only possible action of a catalytic agent would be to resolve the water into its components.) It may indeed be true that at ordinary temperatures, except when the quantity either of hydrogen or of oxygen is very small compared with the quantity of water, the state of dissipated energy is one of such extreme rarefaction as to lie entirely beyond our power of experimental verification. It is also to be noticed that a state of great rarefaction is so unfavorable to any condensation of the gases, that it is quite probable that the catalytic action of platinum may cease entirely at a degree of rarefaction far short of what is necessary for a state of dissipated energy. But with respect to the theoretical demonstration, such states of great rarefaction are precisely those to which we should suppose that the laws of ideal gas-mixtures would apply most perfectly.

But when the compound gas G_3 is formed of G_1 and G_2 without condensation, (i. e., when $\beta_1 + \beta_2 = 1$), it appears from equation (307) that the relation between m_1 , m_2 , and m_3 which is necessary for a phase of dissipated energy is determined by the temperature alone.

In any case, if we regard the total quantities of the gases G_1 and G_2 (as determined by the ultimate analysis of the gas-mixture), and also the volume, as constant, the quantities of these gases which appear uncombined in a phase of dissipated energy will increase with the temperature, if the formation of the compound G_3 *without change of volume* is attended with evolution of heat. Also, if we regard the total quantities of the gases G_1 and G_2 , and also the pressure, as constant, the quantities of these gases which appear uncombined in a phase of dissipated energy, will increase with the temperature, if the formation of the compound G_3 *under constant pressure* is attended with evolution of heat. If $B = 0$, (a case, as has been seen, of especial importance), the heat obtained by the formation of a unit of G_3 out of G_1 and G_2 without change of volume or of temperature will be equal to C . If this quantity is positive, and the total quantities of the gases G_1 and G_2 and also the volume have given finite values, for an infinitesimal value of t we shall have (for a phase of dissipated energy) an infinitesimal value either of m_1 or of m_2 , and for an infinite value of t we shall have finite (neither infinitesimal nor infinite) values of m_1 , m_2 , and m_3 . But if we suppose the pressure instead of the volume to have a given finite value (with

suppositions otherwise the same), we shall have for infinitesimal values of t an infinitesimal value either of m_1 or m_2 , and for infinite values of t finite or infinitesimal values of m_3 according as $\beta_1 + \beta_2$ is equal to or greater than unity.

The case which we have considered is that of a ternary gas-mixture, but our results may easily be generalized in this respect. In fact, whatever the number of component gases in a gas-mixture, if there are relations of equivalence in ultimate analysis between these components, such relations may be expressed by one or more equations of the form

$$\lambda_1 \mathfrak{G}_1 + \lambda_2 \mathfrak{G}_2 + \lambda_3 \mathfrak{G}_3 + \text{etc.} = 0, \quad (311)$$

where $\mathfrak{G}_1, \mathfrak{G}_2, \text{etc.}$ denote the units of the various component gases, and $\lambda_1, \lambda_2, \text{etc.}$ denote positive or negative constants such that $\sum_1 \lambda_1 = 0$. From (311) with (86) we may derive for phases of dissipated energy,

$$\lambda_1 \mu_1 + \lambda_2 \mu_2 + \lambda_3 \mu_3 + \text{etc.} = 0, \\ \text{or} \quad \sum_1 (\lambda_1 \mu_1) = 0. \quad (312)$$

Hence, by (276),

$$\sum_1 \left(\lambda_1 a_1 \log \frac{m_1}{v} \right) = A + B \log t - \frac{C}{t}, \quad (313)$$

where A, B and C are constants determined by the equations

$$A = \sum_1 (\lambda_1 H_1 - \lambda_1 e_1 - \lambda_1 a_1), \quad (314)$$

$$B = \sum_1 (\lambda_1 e_1), \quad (315)$$

$$C = \sum_1 (\lambda_1 E_1). \quad (316)$$

Also, since $p v = \sum_1 (a_1 m_1) t$,

$$\sum_1 (\lambda_1 a_1 \log m_1) - \sum (\lambda_1 a_1) \log \sum_1 (a_1 m_1) \\ + \sum (\lambda_1 a_1) \log p = A + B' \log t - \frac{C}{t}, \quad (317)$$

where

$$B' = \sum_1 (\lambda_1 e_1 + \lambda_1 a_1). \quad (318)$$

If there is more than one equation of the form (311), we shall have more than one of each of the forms (313) and (317), which will hold true simultaneously for phases of dissipated energy.

It will be observed that the relations necessary for a phase of dissipated energy between the volume and temperature of an ideal gas-mixture, and the quantities of the components which take part in the chemical processes, and the pressure due to these components, are not affected by the presence of neutral gases in the gas-mixture.

From equations (312) and (234) it follows that if there is a phase of dissipated energy at any point in an ideal gas-mixture in equilibrium under the influence of gravity, the whole gas-mixture must consist of such phases.

The equations of the phases of dissipated energy of a binary gas-mixture, the components of which are identical in substance, are comparatively simple in form. In this case the two components have the same potential, and if we write β for $\frac{a_1}{a_2}$ (the ratio of the volumes of equal quantities of the two components under the same conditions of temperature and pressure), we shall have

$$\log \frac{m_1^\beta}{m_2 v^{\beta-1}} = \frac{A}{a_2} + \frac{B}{a_2} \log t - \frac{C}{a_2 t}, \quad (319)$$

$$\log \frac{m_1^\beta p^{\beta-1}}{m_2 (a_1 m_1 + a_2 m_2)^{\beta-1}} = \frac{A}{a_2} + \frac{B'}{a_2} \log t - \frac{C}{a_2 t}; \quad (320)$$

where

$$A = H_1 - H_2 - c_1 + c_2 - a_1 + a_2, \quad (321)$$

$$B = c_1 - c_2, \quad B' = c_1 - c_2 + a_1 - a_2, \quad (322)$$

$$C = E_1 - E_2. \quad (323)$$

Gas-mixtures with Convertible Components.

The equations of the phases of dissipated energy of ideal gas-mixtures which have components of which some are identical in ultimate analysis to others have an especial interest in relation to the theory of gas-mixtures in which the components are not only thus equivalent, but are actually transformed into each other within the gas-mixture on variations of temperature and pressure, so that quantities of these (proximate) components are entirely determined, at least in any permanent phase of the gas-mixture, by the quantities of a smaller number of ultimate components, with the temperature and pressure. Such gas-mixtures may be distinguished as having *convertible components*. The very general considerations adduced on pages 197–203, which are not limited in their application to gaseous bodies, suggest the hypothesis that the equations of the phases of dissipated energy of ideal gas-mixtures may apply to such gas-mixtures as have been described. It will, however, be desirable to consider the matter more in detail.

In the first place, if we consider the case of a gas-mixture which only differs from an ordinary ideal gas-mixture for which some of the components are equivalent in that there is perfect freedom in regard to the transformation of these components, it follows at once from the general formula of equilibrium (1) or (2) that equilibrium is only possible for such phases as we have called phases of dissipated energy, for which some of the characteristic equations have been deduced in the preceding pages.

If it should be urged, that regarding a gas-mixture which has convertible components as an ideal gas-mixture of which, for some reason, only a part of the phases are actually capable of existing, we might still suppose the particular phases which alone can exist to be determined by some other principle than that of the free convertibility of the components (as if, perhaps, the case were analogous to one of *constraint* in mechanics), it may easily be shown that such a hypothesis is entirely untenable, when the quantities of the proximate components may be varied independently by suitable variations of the temperature and pressure, and of the quantities of the ultimate components, and it is admitted that the relations between the energy, entropy, volume, temperature, pressure, and the quantities of the several proximate components in the gas-mixture are the same as for an ordinary ideal gas-mixture, in which the components are not convertible. Let us denote the quantities of the n' proximate components of a gas-mixture A by m_1, m_2 , etc., and the quantities of its n ultimate components by $\mathbf{m}_1, \mathbf{m}_2$, etc. (n denoting a number less than n'), and let us suppose that for this gas-mixture the quantities $\varepsilon, \eta, v, t, p, m_1, m_2$, etc. satisfy the relations characteristic of an ideal gas-mixture, while the phase of the gas-mixture is entirely determined by the values of $\mathbf{m}_1, \mathbf{m}_2$, etc., with two of the quantities $\varepsilon, \eta, v, t, p$. We may evidently imagine such an ideal gas-mixture B having n' components (not convertible), that every phase of A shall correspond with one of B in the values of $\varepsilon, \eta, v, t, p, m_1, m_2$, etc. Now let us give to the quantities $\mathbf{m}_1, \mathbf{m}_2$, etc. in the gas-mixture A any fixed values, and for the body thus defined let us imagine the $v\text{-}\eta\text{-}\varepsilon$ surface (see page 174) constructed; likewise for the ideal gas-mixture B let us imagine the $v\text{-}\eta\text{-}\varepsilon$ surface constructed for every set of values of m_1, m_2 , etc. which is consistent with the given values of $\mathbf{m}_1, \mathbf{m}_2$, etc., i. e., for every body of which the *ultimate* composition would be expressed by the given values of $\mathbf{m}_1, \mathbf{m}_2$, etc. It follows immediately from our supposition, that every point in the $v\text{-}\eta\text{-}\varepsilon$ surface relating to A must coincide with some point of one of the $v\text{-}\eta\text{-}\varepsilon$ surfaces relating

to B not only in respect to position but also in respect to its tangent plane (which represents temperature and pressure); therefore the v - η - ε surface relating to A must be tangent to the various v - η - ε surfaces relating to B , and therefore must be an envelop of these surfaces. From this it follows that the points which represent phases common to both gas-mixtures must represent the phases of dissipated energy of the gas-mixture B .

The properties of an ideal gas-mixture which are assumed in regard to the gas-mixture of convertible components in the above demonstration are expressed by equations (277) and (278) with the equation

$$\varepsilon = \sum_1 (c_1 m_1 t + m_1 E_1). \quad (324)$$

It is usual to assume in regard to gas-mixtures having convertible components that the convertibility of the components does not affect the relations (277) and (324). The same cannot be said of the equation (278). But in a very important class of cases it will be sufficient if the applicability of (277) and (324) is admitted. The cases referred to are those in which in certain phases of a gas-mixture the components are convertible, and in other phases of the same proximate composition the components are not convertible, and the equations of an ideal gas-mixture hold true.

If there is only a single degree of convertibility between the components, (i. e., if only a single kind of conversion, with its reverse, can take place among the components,) it will be sufficient to assume, in regard to the phases in which conversion takes place, the validity of equation (277) and of the following, which can be derived from (324) by differentiation, and comparison with equation (11), which expresses a necessary relation,

$$[t d\eta - p dv - \sum_1 (c_1 m_1) dt]_m = 0.* \quad (325)$$

We shall confine our demonstration to this case. It will be observed that the physical signification of (325) is that if the gas-mixture is subjected to such changes of volume and temperature as do not alter its proximate composition, the heat absorbed or yielded may be calculated by the same formula as if the components were not convertible.

Let us suppose the thermodynamic state of a gaseous mass M , of such a kind as has just been described, to be varied while within the limits within which the components are not convertible. (The quantities of the proximate components, therefore, as well as of the ulti-

* This notation is intended to indicate that m_1, m_2 , etc. are regarded as constant

mate, are supposed constant). If we use the same method of geometrical representation as before, the point representing the volume, entropy, and energy of the mass will describe a line in the v - η - ε surface of an ideal gas-mixture of inconvertible components, the form and position of this surface being determined by the proximate composition of M . Let us now suppose the same mass to be carried beyond the limit of inconvertibility, the variations of state after passing the limit being such as not to alter its proximate composition. It is evident that this will in general be possible. Exceptions can only occur when the limit is formed by phases in which the proximate composition is uniform. The line traced in the region of convertibility must belong to the same v - η - ε surface of an ideal gas-mixture of inconvertible components as before, continued beyond the limit of inconvertibility for the components of M , since the variations of volume, entropy and energy are the same as would be possible if the components were not convertible. But it must also belong to the v - η - ε surface of the body M , which is here a gas-mixture of convertible components. Moreover, as the inclination of each of these surfaces must indicate the temperature and pressure of the phases through which the body passes, these two surfaces must be tangent to each other along the line which has been traced. As the v - η - ε surface of the body M in the region of convertibility must thus be tangent to all the surfaces representing ideal gas-mixtures of every possible proximate composition consistent with the ultimate composition of M , continued beyond the region of inconvertibility, in which alone their form and position may be capable of experimental demonstration, the former surface must be an envelop of the latter surfaces, and therefore a continuation of the surface of the phases of dissipated energy in the region of inconvertibility.

The foregoing considerations may give a measure of *a priori* probability to the results which are obtained by applying the ordinary laws of ideal gas-mixtures to cases in which the components are convertible. It is only by experiments upon gases in phases in which their components are convertible that the validity of any of these results can be established.

The very accurate determinations of density which have been made for the peroxide of nitrogen enable us to subject some of our equations to a very critical test. That this substance in the gaseous state is properly regarded as a mixture of different gases can hardly be doubted, as the proportion of the components derived from its density on the supposition that one component has the molecular formula

NO_2 and the other the formula N_2O_4 is the same as that derived from the depth of the color on the supposition that the absorption of light is due to one of the components alone, and is proportioned to the separate density of that component.*

MM. Sainte-Claire Deville and Troost† have given a series of determinations of what we shall call the *relative densities* of peroxide of nitrogen at various temperatures under atmospheric pressure. We use the term *relative density* to denote what it is usual in treatises on chemistry to denote by the term *density*, viz., the actual density of a gas divided by the density of a standard perfect gas at the same pressure and temperature, the standard gas being air, or more strictly, an ideal gas which has the same density as air at the zero of the centigrade scale and the pressure of one atmosphere. In order to test our equations by these determinations, it will be convenient to transform equation (320), so as to give directly the relation between the relative density, the pressure, and the temperature.

As the density of the standard gas at any given temperature and pressure may by (263) be expressed by the formula $\frac{p}{a_s t}$, the relative density of a binary gas-mixture may be expressed by

$$D = (m_1 + m_2) \frac{a_s t}{p v}. \quad (326)$$

Now by (263)

$$a_1 m_1 + a_2 m_2 = \frac{p v}{t}. \quad (327)$$

By giving to m_2 and m_1 successively the value zero in these equations, we obtain

$$D_1 = \frac{a_s}{a_1}, \quad D_2 = \frac{a_s}{a_2}, \quad (328)$$

where D_1 and D_2 denote the values of D when the gas consists wholly of one or of the other component. If we assume that

$$D_2 = 2 D_1, \quad (329)$$

we shall have

$$a_1 = 2 a_2. \quad (330)$$

From (326) we have

$$m_1 + m_2 = D \frac{p v}{a_s t},$$

* Salet, "Sur la coloration du peroxyde d'azote," Comptes Rendus, vol. lxxvii, p. 488.

† Comptes Rendus, vol. lxxiv, p. 237.

and from (327), by (328) and (330),

$$2 m_1 + m_2 = D_2 \frac{p v}{a_s t} = 2 D_1 \frac{p v}{a_s t},$$

whence

$$m_1 = (D_2 - D) \frac{p v}{a_s t}, \tag{331}$$

$$m_2 = 2 (D - D_1) \frac{p v}{a_s t}. \tag{332}$$

By (327), (331), and (332) we obtain from (320)

$$\log \frac{(D_2 - D)^2 p}{2 (D - D_1) a_s} = \frac{A}{a_2} + \frac{B'}{a_2} \log t - \frac{C}{t}. \tag{333}$$

This formula will be more convenient for purposes of calculation if we introduce common logarithms (denoted by \log_{10}) instead of hyperbolic, the temperature of the ordinary centigrade scale t_c instead of the absolute temperature t , and the pressure in atmospheres p_{at} instead of p the pressure in a rational system of units. If we also add the logarithm of a_s to both sides of the equation, we obtain

$$\log_{10} \frac{(D_2 - D)^2 p_{at}}{2 (D - D_1)} = \mathbf{A} + \frac{B'}{a_2} \log_{10} (t_c + 273) - \frac{\mathbf{C}}{t_c + 273}, \tag{334}$$

where \mathbf{A} and \mathbf{C} denote constants, the values of which are closely connected with those of A and C .

From the molecular formulæ of peroxide of nitrogen NO_2 and N_2O_4 , we may calculate the relative densities

$$D_1 = \frac{14 + 32}{2} .0691 = 1.589, \text{ and } D_2 = \frac{28 + 64}{2} .0691 = 3.178. \tag{335}$$

The determinations of MM. Deville and Troost are satisfactorily represented by the equation

$$\log_{10} \frac{(3.178 - D)^2 p_{at}}{2 (D - 1.589)} = 9.47056 - \frac{3118.6}{t_c + 273}, \tag{336}$$

which gives

$$D = 3.178 + \Theta - \sqrt{\Theta (3.178 + \Theta)}$$

where
$$\log_{10} \Theta = 9.47056 - \frac{3118.6}{t_c + 273} - \log_{10} p_{at}$$

In the first part of the following table are given in successive columns the temperature and pressure of the gas in the several experiments of MM. Deville and Troost, the relative densities calculated from these numbers by equation (336), the relative densities as observed, and the difference of the observed and calculated relative

densities. It will be observed that these differences are quite small, in no case reaching .03, and on the average scarcely exceeding .01. The significance of such correspondence in favor of the hypothesis by means of which equation (336) has been established is of course diminished by the fact that two constants in the equation have been determined from these experiments. If the same equation can be shown to give correctly the relative densities at other pressures than that for which the constants have been determined, such correspondence will be much more decisive.

t_c	p_{at}	D calculated by eq. (336).	D observed.	diff.	Observers.
26.7	1	2.676	2.65	-.026	D. & T.
35.4	1	2.524	2.53	+.006	D. & T.
39.8	1	2.443	2.46	+.017	D. & T.
49.6	1	2.256	2.27	+.014	D. & T.
60.2	1	2.067	2.08	+.013	D. & T.
70.0	1	1.920	1.92	.000	D. & T.
80.6	1	1.801	1.80	-.001	D. & T.
90.0	1	1.728	1.72	-.008	D. & T.
100.1	1	1.676	1.68	+.004	D. & T.
111.3	1	1.641	1.65	+.009	D. & T.
121.5	1	1.622	1.62	-.002	D. & T.
135.0	1	1.607	1.60	-.007	D. & T.
154.0	1	1.597	1.58	-.017	D. & T.
183.2	1	1.592	1.57	-.022	D. & T.
97.5	1	1.687			
97.5	$\frac{19450}{26397}$	1.631	1.783	+.152	P. & W.
24.5	1	2.711			
24.5	$\frac{18090}{42529}$	2.524	2.52	-.004	P. & W.
11.3	1	2.891			
11.3	$\frac{4265}{44205}$	2.620	2.645	+.025	P. & W.
4.2	1	2.964			
4.2	$\frac{6023}{35438}$	2.708	2.588	-.120	P. & W.

Messrs. Playfair and Wanklyn have published* four determinations of the relative density of peroxide of nitrogen at various temperatures when diluted with nitrogen. Since the relations expressed by equations (319) and (320) are not affected by the presence of a third gas which is different from the gases G_1 and G_2 (to which m_1 and m_2 relate) and neutral to them, (see the remark at the foot of page 233),—provided that we take p to denote the pressure which we attribute to the gases G_1 and G_2 , i. e., the total pressure diminished by the pressure which the third gas would exert if occupying alone the same space at the same temperature,—it follows that the relations expressed for

*Transactions of the Royal Society of Edinburg, vol. xxii, p. 441.

peroxide of nitrogen by (333), (334), and (336) will not be affected by the presence of free nitrogen, if the pressure expressed by p or p_{at} and contained implicitly in the symbol D (see equation (326) by which D is defined) is understood to denote the total pressure diminished by the pressure due to the free nitrogen. The determinations of Playfair and Wanklyn are given in the latter part of the above table. The pressures given are those obtained by subtracting the pressure due to the free nitrogen from the total pressure. We may suppose such reduced pressures to have been used in the reduction of the observations by which the numbers in the column of observed relative densities were obtained. Besides the relative densities calculated by equation (336) for the temperatures and (reduced) pressures of the observations, the table contains the relative densities calculated for the same temperatures and the pressure of one atmosphere.

The reader will observe that in the second and third experiments of Playfair and Wanklyn there is a very close accordance between the calculated and observed values of D , while in the second and fourth experiments there is a considerable difference. Now the weight to be attributed to the several determinations is very different. The quantities of peroxide of nitrogen which were used in the several experiments were respectively .2410, .5893, .3166, and .2016 grammes. For a rough approximation, we may assume that the probable errors of the relative densities are inversely proportional to these numbers. This would make the probable error of the first and fourth observations two or three times as great as that of the second and considerably greater than that of the third. We must also observe that in the first of these experiments, the observed relative density 1.783 is greater than 1.687, the relative density calculated by equation (336) for the temperature of the experiment and the pressure of one atmosphere. Now the number 1.687 we may regard as established directly by the experiments of Deville and Troost. For in seven successive experiments in this part of the series the calculated relative densities differ from the observed by less than .01. If then we accept the numbers given by experiment, the effect of diluting the gas with nitrogen is to increase its relative density. As this result is entirely at variance with the facts observed in the case of other gases, and in the case of this gas at lower temperatures, as appears from the three other determinations of Playfair and Wanklyn, it cannot possibly be admitted on the strength of a single

observation. The first experiment of this series cannot therefore properly be used as a test of our equations. Similar considerations apply with somewhat less force to the last experiment. By comparing the temperatures and pressures of the three last experiments with the observed relative densities, the reader may easily convince himself that if we admit the substantial accuracy of the determinations in the two first of these experiments (the second and third of the series, which have the greatest weight), the last determination of relative density 2.588 must be too small. In fact, it should evidently be greater than the number in the preceding experiment 2.645.

If we confine our attention to the second and third experiments of the series, the agreement is as good as could be desired. Nor will the admission of errors of .152 and .120 (certainly not large in determinations of this kind) in the first and fourth experiments involve any serious doubt of the substantial accuracy of the second and third, when the difference of weight of the determinations is considered. Yet it is much to be desired that the relation expressed by (336), or with more generality by (334), should be tested by more numerous experiments.

It should be stated that the numbers in the column of pressures are not quite accurate. In the experiments of Deville and Troost the gas was subject to the actual atmospheric pressure at the time of the experiment. This varied from 747 to 764 millimeters of mercury. The precise pressure for each experiment is not given. In the experiments of Playfair and Wanklyn the mixture of nitrogen and peroxide of nitrogen was subject to the actual atmospheric pressure at the time of the experiment. The numbers in the column of pressures express the fraction of the whole pressure which remains after subtracting the part due to the free nitrogen. But no indication is given in the published account of the experiments in regard to the height of the barometer. Now it may easily be shown that a variation of $\frac{1}{760}$ in the value of p can in no case cause a variation of more than .005 in the value of D as calculated by equation (336). In any of the experiments of Playfair and Wanklyn a variation of more than 30^{mm} in the height of the barometer would be necessary to produce a variation of .01 in the value of D . The errors due to this source cannot therefore be very serious. They might have been avoided altogether in the discussion of the experiments of Deville and Troost by using instead of (336) a formula expressing the relation between the relative density, the temperature, and the actual density, as the reciprocal of the latter quantity is given for each ex-

periment of this series. It seemed best, however, to make a trifling sacrifice of accuracy for the sake of simplicity.

It might be thought that the experiments under discussion would be better represented by a formula in which the term containing $\log t$ (see equation (333)) was retained. But an examination of the figures in the table will show that nothing important can be gained in this respect, and there is hardly sufficient motive for adding another term to the formula of calculation. Any attempt to determine the *real* values of A , B' , and C in equation (333), (assuming the absolute validity of such an equation for peroxide of nitrogen,) from the experiments under discussion would be entirely misleading, as the reader may easily convince himself.

From equation (336), however, the following conclusions may be deduced. By comparison with (334) we obtain

$$A + \frac{B'}{a_2} \log_{10} t - \frac{C}{t} = 9.47056 - \frac{3118.6}{t},$$

which must hold true approximately between the temperatures 11° and 90° . (At higher temperatures the relative densities vary too slowly with the temperatures to afford a critical test of the accuracy of this relation.) By differentiation we obtain

$$\frac{MB'}{a_2 t} + \frac{C}{t^2} = \frac{3118.6}{t^2},$$

where M denotes the modulus of the common system of logarithms. Now by comparing equations (333) and (334) we see that

$$C = \frac{MC}{a_2} = .43429 \frac{C}{a_2}.$$

Hence

$$B' t + C = 7181 a_2 = 3590 a_1,$$

which may be regarded as a close approximation at 40° or 50° , and a tolerable approximation between the limits of temperature above mentioned. Now $B' t + C$ represents the heat evolved by the conversion of a unit of NO_2 into N_2O_4 under constant pressure. Such conversion cannot take place at constant pressure without change of temperature, which renders the experimental verification of the last equation less simple. But since by equations (322)

$$B' = B + a_1 - a_2 = B + \frac{1}{2} a_1,$$

we shall have for the temperature of 40°

$$B t + C = 3434 a_1.$$

Now $B t + C$ represents the decrease of energy when a unit of NO_2 is transformed into N_2O_4 without change of temperature. It therefore

represents the excess of the heat evolved over the work done by external forces when a mass of the gas is compressed at constant temperature until a unit of NO_2 has been converted into N_2O_4 . This quantity will be constant if $B = 0$, i. e., if the specific heats at constant volume of NO_2 and N_2O_4 are the same. This assumption would be more simple from a theoretical stand-point and perhaps safer than the assumption that $B' = 0$. If $B = 0$, $B' = a_2$. If we wish to embody this assumption in the equation between D , p , and t , we may substitute

$$6.5228 + \log_{10} (t_c + 273) - \frac{2977.4}{t_c + 273}$$

for the second member of equation (336). The relative densities calculated by the equation thus modified from the temperatures and pressures of the experiments under discussion will not differ from those calculated from the unmodified equation by more than .002 in any case, or by more than .001 in the first series of experiments.

It is to be noticed that if we admit the validity of the volumetrical relation expressed by equation (333), which is evidently equivalent to an equation between p , t , v , and m (this letter denoting the quantity of the gas without reference to its molecular condition), or if we admit the validity of the equation only between certain limits of temperature and for densities less than a certain limit of density, and also admit that between the given limits of temperature the specific heat of the gas at constant volume may be regarded as a constant quantity when the gas is sufficiently rarefied to be regarded as consisting wholly of NO_2 ,—or, to speak without reference to the molecular state of the gas, when it is rarefied until its relative density D approximates to its limiting value D_1 ,—we must also admit the validity (within the same limits of temperature and density) of all the calorimetrical relations which belong to ideal gas-mixtures with convertible components. The premises are evidently equivalent to this,—that we may imagine an ideal gas with convertible components such that between certain limits of temperature and above a certain limit of density the relation between p , t , and v shall be the same for a unit of this ideal gas as for a unit of peroxide of nitrogen, and for a very great value of v (within the given limits of temperature) the thermal capacity at constant volume of the ideal and actual gases shall be the same. Let us regard t and v as independent variables; we may let these letters and p refer alike to the ideal and real gases, but we must distinguish the entropy η' of the ideal gas from the entropy η of the real gas. Now by (88)

$$\frac{d\eta}{dv} = \frac{dp}{dt}, \tag{337}$$

therefore

$$\frac{d}{dv} \frac{d\eta}{dt} = \frac{d}{dt} \frac{d\eta}{dv} = \frac{d}{dt} \frac{dp}{dt} = \frac{d^2p}{dt^2}. \tag{338}$$

Since a similar relation will hold true for η' , we obtain

$$\frac{d}{dv} \frac{d\eta}{dt} = \frac{d}{dv} \frac{d\eta'}{dt}, \tag{339}$$

which must hold true within the given limits of temperature and density. Now it is granted that

$$\frac{d\eta}{dt} = \frac{d\eta'}{dt} \tag{340}$$

for very great values of v at any temperature within the given limits, (for the two members of the equation represent the thermal capacities at constant volume of the real and ideal gases divided by t .) hence, in virtue of (339), this equation must hold true in general within the given limits of temperature and density. Again, as an equation like (337) will hold true of η' , we shall have

$$\frac{d\eta}{dv} = \frac{d\eta'}{dv}. \tag{341}$$

From the two last equations it is evident that in all calorimetrical relations the ideal and real gases are identical. Moreover the energy and entropy of the ideal gas are evidently so far arbitrary that we may suppose them to have the same values as in the real gas for any given values of t and v . Hence the entropies of the two gases are the same within the given limits; and on account of the necessary relation

$$d\varepsilon = t d\eta - p dv,$$

the energies of the two gases are in like manner identical. Hence the fundamental equation between the energy, entropy, volume, and quantity of matter must be the same for the ideal gas as for the actual.

We may easily form a fundamental equation for an ideal gas-mixture with convertible components, which shall relate only to the phases of equilibrium. For this purpose, we may use the equations of the form (312) to eliminate from the equation of the form (273), which expresses the relation between the pressure, the temperature, and the potentials for the proximate components, as many of the potentials as there are equations of the former kind, leaving the

potentials for those components which it is convenient to regard as the ultimate components of the gas-mixture.

In the case of a binary gas-mixture with convertible components, the components will have the same potential, which may be denoted by μ , and the fundamental equation will be

$$p = a_1 L_1 t^{\frac{c_1 + a_1}{a_1}} e^{\frac{\mu - E_1}{a_1 t}} + a_2 L_2 t^{\frac{c_2 + a_2}{a_2}} e^{\frac{\mu - E_2}{a_2 t}}, \quad (342)$$

where

$$L_1 = e^{\frac{H_1 - c_1 - a_1}{a_1}}, \quad L_2 = e^{\frac{H_2 - c_2 - a_2}{a_2}}. \quad (343)$$

From this equation, by differentiation and comparison with (98), we obtain

$$\begin{aligned} \frac{\eta}{v} = & L_1 \left(c_1 + a_1 - \frac{\mu - E_1}{t} \right) t^{\frac{c_1}{a_1}} e^{\frac{\mu - E_1}{a_1 t}} \\ & + L_2 \left(c_2 + a_2 - \frac{\mu - E_2}{t} \right) t^{\frac{c_2}{a_2}} e^{\frac{\mu - E_2}{a_2 t}}, \end{aligned} \quad (244)$$

$$\frac{m}{v} = L_1 t^{\frac{c_1}{a_1}} e^{\frac{\mu - E_1}{a_1 t}} + L_2 t^{\frac{c_2}{a_2}} e^{\frac{\mu - E_2}{a_2 t}}. \quad (345)$$

From the general equation (93) with the preceding equations the following is easily obtained,—

$$\frac{\varepsilon}{v} = L_1 (c_1 t + E_1) t^{\frac{c_1}{a_1}} e^{\frac{\mu - E_1}{a_1 t}} + L_2 (c_2 t + E_2) t^{\frac{c_2}{a_2}} e^{\frac{\mu - E_2}{a_2 t}}. \quad (346)$$

We may obtain the relation between p , t , v , and m by eliminating μ from (342) and (345). For this purpose we may proceed as follows. From (342) and (345) we obtain

$$p - a_2 t^{\frac{m}{v}} = (a_1 - a_2) L_1 t^{\frac{c_1 + a_1}{a_1}} e^{\frac{\mu - E_1}{a_1 t}} \quad (347)$$

$$a_1 t^{\frac{m}{v}} - p = (a_1 - a_2) L_2 t^{\frac{c_2 + a_2}{a_2}} e^{\frac{\mu - E_2}{a_2 t}}; \quad (348)$$

and from these equations we obtain

$$\begin{aligned} a_1 \log \left(p - a_2 t^{\frac{m}{v}} \right) - a_2 \log \left(a_1 t^{\frac{m}{v}} - p \right) = & (a_1 - a_2) \log (a_1 - a_2) \\ & + a_1 \log L_1 - a_2 \log L_2 + (c_2 - c_1 + a_1 - a_2) \log t - \frac{E_1 - E_2}{t}. \end{aligned} \quad (349)$$

(In the particular case when $a_1 = 2 a_2$ this equation will be equivalent to (333)). By (347) and (348) we may easily eliminate μ from (346).

The reader will observe that the relations thus deduced from the fundamental equation (342) without any reference to the different components of the gaseous mass are equivalent to those which relate to the phases of dissipated energy of a binary gas-mixture with components which are equivalent in substance but not convertible, except that the equations derived from (342) do not give the quantities of the proximate components, but relate solely to those properties which are capable of direct experimental verification without the aid of any theory of the constitution of the gaseous mass.

The practical application of these equations is rendered more simple by the fact that the ratio $a_1 : a_2$ will always bear a simple relation to unity. When a_1 and a_2 are equal, if we write a for their common value, we shall have by (342) and (345)

$$p v = a m t, \tag{350}$$

and by (345) and (346)

$$\frac{\varepsilon}{m} = \frac{L_1(c_1 t + E_1) + L_2(c_2 t + E_2)t}{L_1 + L_2 t} \frac{\frac{c_2 - c_1}{a} \frac{E_1 - E_2}{e^{at}}}{\frac{c_2 - c_1}{a} \frac{E_1 - E_2}{e^{at}}}. \tag{351}$$

By this equation we may calculate directly the amount of heat required to raise a given quantity of the gas from one given temperature to another at constant volume. The equation shows that the amount of heat will be independent of the volume of the gas. The heat necessary to produce a given change of temperature in the gas at constant pressure, may be found by taking the difference of the values of χ , as defined by equation (89), for the initial and final states of the gas. From (89), (350), and (351) we obtain

$$\frac{\chi}{m} = \frac{L_1(c_1 t + at + E_1) + L_2(c_2 t + at + E_2)t}{L_1 + L_2 t} \frac{\frac{c_2 - c_1}{a} \frac{E_1 - E_2}{e^{at}}}{\frac{c_2 - c_1}{a} \frac{E_1 - E_2}{e^{at}}}. \tag{352}$$

By differentiation of the two last equations we may obtain directly the specific heats of the gas at constant volume and at constant pressure.

The fundamental equation of an ideal ternary gas-mixture with a single relation of convertibility between its components is

$$\begin{aligned}
 p = & a_1 e^{\frac{H_1 - c_1 - a_1}{a_1}} \frac{c_1 + a_1}{t} \frac{\mu_1 - E_1}{e^{a_1 t}} \\
 & + a_2 e^{\frac{H_2 - c_2 - a_2}{a_2}} \frac{c_2 + a_2}{t} \frac{\mu_2 - E_2}{e^{a_2 t}} \\
 & + a_3 e^{\frac{H_3 - c_3 - a_3}{a_3}} \frac{c_3 + a_3}{t} \frac{\lambda_1 \mu_1 + \lambda_2 \mu_2 - E_3}{e^{a_3 t}}, \quad (353)
 \end{aligned}$$

where λ_1 and λ_2 have the same meaning as on page 230.

(To be continued.)

ERRATA.

Page 167, formula (168), for m_1 read μ_1 .

“ formula (169), for m_1, \dots, m_{n-1} read μ_1, \dots, μ_{n-1} .

Page 239, formula (333), for $\frac{C}{t}$ read $\frac{C}{a_2 t}$.

VI. THE HYDROIDS OF THE PACIFIC COAST OF THE UNITED STATES,
SOUTH OF VANCOUVER ISLAND. WITH A REPORT UPON THOSE
IN THE MUSEUM OF YALE COLLEGE. BY S. F. CLARK.

Read Jan. 19. 1876.

THE Museum is indebted for its collection of Californian Hydroids chiefly to Prof. D. C. Eaton, who has presented during the last two or three years, a large number of specimens, that were received by him with dried algæ from that coast. They were collected and sent to him by Dr. C. W. Anderson, Santa Cruz, Cal.; Dr. L. N. Dimmick, Santa Barbara, Cal.; Mrs. Ellwood Cooper, Santa Barbara, Cal.; and Miss Mitchell of Vancouver Island. All the specimens received from these sources were collected in tide-pools along the shore or attached to algæ, washed in from deeper water. A few alcoholic specimens have also been received from San Diego, Cal., collected on the piles of the wharves and along the shore, by Dr. E. Palmer, and a fine specimen of *Plumularia setacea* was dredged in six to eight fathoms, off San Diego, by Mr. Henry Hemphill.

Some of the species, including most of the *Sertularidæ*, do not seem to be injured by being dried, but others, as the *Campanularidæ*, are usually rendered useless for description. The specimens of the two species of *Campanularia* described below are unusually well preserved, both hydrothecæ and gonothecæ being in good condition.

There has been very little published on the hydroids of the western coast of North America, up to the present time. In 1857 Dr. Trask* described and figured nine new species of Zoöphytes from the Bay of San Francisco and adjacent localities. Five of these are Bryozoa; the remaining four represent three genera of the family Sertulariæ, as follows: *Sertularia anguina*, *S. falcata*, *Sertularella turgida* and *Hydrallmania Franciscana*, all of which, so far as I am aware, are peculiar to that coast; unless indeed the last named species prove to be identical with *H. falcata* of Europe,

* Proceedings of the California Academy of Natural Sciences, vol. i, March, 1857.
Dr. J. B. Trask.

Africa and New England. In 1860 Andrew Murray* described and figured five species from the Californian coast, of which three are new, and the other two, *Sertularia labrata* and *Plumularia gracilis*, are synonymous with two of Trask's species, viz: *Sertularia anguina* and *Plumularia Franciscana*. Mr. Alexander Agassiz† in 1865 described seven species and recorded seven others from the Bay of San Francisco; and he had three of the same from the Gulf of Georgia, W. T. Five species were also mentioned by him from the North Pacific. Two of these five northern species, *Bougainvillia Mertensii* Ag. and *Cotulina Greenei* A. Ag., are also found at San Francisco. The latter species having also been collected at Santa Barbara, Cal., has the wide range of nearly three thousand miles upon our western coast. Professor Allman mentions having found sixteen species in a collection from the Californian coast, submitted to him for examination; two of them, *Lafoëa dumosa* and *Sertularia pumila*, are common on the European and New England coasts, and the former species is also recorded from South Africa. The collection in the Museum of Yale College contains twelve species and one variety. Of these four are new; nine are recorded only from the Pacific coast of North America, as yet; and three, *Halecium tenellum*, *Sertularia argentea* and *Plumularia setacea*, are also common on the European shores; the first two of these have also been found on the New England coast, from Maine to Long Island Sound. The most common form on the Californian coast is the showy *Aglaophenia struthionides*, which is apparently as abundant there as *Sertularia argentea* and *S. pumila* are upon our eastern shores, for it forms the bulk of every package sent to us from the western coast. The following table gives a list of all the Hydroids known on the western coast of the United States, from Vancouver's Island to San Diego, with the range of the different species and the names of some of the collectors.

List of Hydroida known to occur between San Diego and Vancouver Island.

<i>Coryne rosaria</i> A. Ag.	Bay of San Francisco, Cal. (A. Agassiz).
<i>Tubularia elegans</i> Clark.	San Diego, Cal. (Dr. E. Palmer).
<i>Thamnocnidia tubularoides</i> A. Ag.	Bay of San Francisco, Cal. (A. Agassiz).

* The Annals and Magazine of Natural History, Series 3, No. XXVIII, April, 1860. Descriptions of new species of Hydroids from the Californian Coast. By Andrew Murray.

† Illustrated Catalogue of the Museum of Comparative Zoölogy. No. II. North American Aclephæ. By Alexander Agassiz. 1865.

<i>Parypha microcephala</i> A. Ag.	Bay of San Francisco, Cal. (A. Agassiz).
<i>Bimeria gracilis</i> Clark.	San Diego, Cal. (E. Palmer).
<i>Bougainvillia Mertensii</i> Agassiz.	Bay of San Francisco, Cal. (A. Agassiz).
<i>Eudendrium</i> , sp.	Santa Cruz, Cal. (C. W. Anderson).
<i>Campanularia everta</i> Clark.	San Diego, Cal. (H. Hemphill), to Vancouver Island (J. M. Dawson).
<i>Campanularia fusiformis</i> Clark.	Vancouver Island (J. M. Dawson).
<i>Campanularia cylindrica</i> Clark.	Santa Cruz, Cal. (C. W. Anderson).
<i>Laomedea rigida</i> A. Ag.	Bay of San Francisco, Cal. (A. Agassiz).
<i>Laomedea Pacifica</i> A. Ag.	Gulf of Georgia (A. Ag.) to Bay of San Francisco (A. Agassiz).
<i>Lafoëa dumosa</i> Sars.	
<i>Halecium tenellum</i> Hincks.	San Diego, Cal. (Dr. E. Palmer).
<i>Sertularia anguina</i> Trask.	Santa Cruz, Cal. (C. W. Anderson), to Vancouver Island (J. M. Dawson).
<i>Sertularia anguina</i> , var <i>robusta</i> Clark.	San Diego, Cal. (H. Hemphill), to Vancouver Island (J. M. Dawson).
<i>Sertularia argentea</i> E. and S.	Santa Barbara, Cal. (Mrs. Ellwood Cooper).
<i>Sertularia pumila</i> Linn.	
<i>Sertularia Greenei</i> Murray.	Santa Barbara, Cal. (Mrs. Ellwood Cooper), to Vancouver Island (J. M. Dawson).
<i>Sertularia furcata</i> Trask.	San Diego (Dr. E. Palmer), to Bay of San Francisco (J. B. Trask).
<i>Sertularia corniculata</i> Murray.	Bay of San Francisco (A. Murray).
<i>Sertularella turgida</i> Clark (Trask).	San Diego, Cal., to Vancouver Island (J. M. Dawson).
<i>Plumularia setacea</i> Lamarck.	San Diego, Cal. (Dr. E. Palmer), to Vancouver Island (J. M. Dawson).
<i>Aglaophenia struthionides</i> Clark (Murray).	San Diego, Cal. (D. C. Cleveland), to Vancouver Island (Miss Mitchell).

This list of twenty-four species is very small compared with that of the eastern coast, from Maine to New York, the fauna of the latter region containing five times as many species as that of the former, notwithstanding that the region included on the western coast is over thirteen hundred miles in length, while that of the New England coast is only about eight hundred. It should be borne in mind however that most of the collecting on the Pacific coast has been done along the shore, the dredge having been little used, and there is little doubt that when the fauna has been more thoroughly investigated the number of Hydroids may be at least doubled. Such a variety as exists on the New England coast can hardly be expected from our Pacific shores south of Vancouver Island, for the waters there do not afford the same diversity in temperature.

Bimeria (?) gracilis, sp. nov.

Plate XXXVIII, figure 3.

Stems clustered, rooted by a creeping stolon, erect, simple, delicate, not divided by distinct joints, thickly branched; branches suberect, the larger ones reaching to the end of the stem and resembling the main stalk, the smaller ones bear but one or two hydranths and are also unjointed; perisarc extending over the hydranths and partially covering the tentacles, annulated at the base of each branch and branchlet. Sporosacs developed from the hydrophyton, a single one at the base of each hydranth-bearing branchlet, oval or ovate, supported by a short peduncle consisting of one or two annulations. Hydranths large, tapering uniformly from the distal end to the base, provided with about ten or twelve tentacles and with a large, rounded or slightly conical proboscis. Height of best specimen, 55^{mm}.

Collected on the piles of wharves at San Diego, Cal., by Dr. E. Palmer, 1875.

Our specimens were not in a good condition when they arrived, having been crowded in a tin can with many other things, which pressed them all out of shape, and the quantity of alcohol not being sufficient to preserve so much animal matter, the hydroids suffered considerably; the hydranths and sporosacs especially were in a very worn and mutilated state. It is not easy to determine just how far the perisarc extends upon the hydranth, but it certainly covers the body of the latter, and it must, I think, be developed over a portion of the tentacles, for after soaking them in a dilute solution of caustic potash for forty-eight hours the tentacles still retained their normal position, nor did they show any decrease in size. The potash seemed to act very slowly, for after being in the warm solution forty-eight hours the hydranths were not entirely dissolved out. The fact of the tentacles being unaffected would seem to indicate that they are entirely protected by chitin, but tentacles so protected would be of little or no use to the animal, and I think it more probable that the distal portions are free and may be contracted into the basal covering. It is impossible to determine from our specimens how the tentacles are held, whether in a single erect verticil as in *Garveia* or with each alternate tentacle depressed, as in *Bimeria vestita* of Wright. With such imperfect data I feel some doubt about placing this species in the genus *Bimeria*, and only do so provisionally.

Tubularia elegans, sp. nov.

Plate XXXVIII, figure 2.

Stems clustered, rooted by a creeping stolon, erect, unbranched, more or less annulated at intervals toward the base. Hydranths large, with about thirty tentacles in the proximal set and twenty to twenty-four in the distal. Gonophores borne in clusters just inside the proximal tentacles, twelve to twenty in a cluster, each of the larger ones crowned with four conical tubercles. Height of finest specimen, 75^{mm}.

Collected on the piles of the wharf at San Diego, by Dr. E. Palmer, 1875. Intermingled with it and often attached to it were numerous shoots of *Bimeria*. Many of the young had attached themselves to the parent stalk, giving at first sight the appearance of branching stems.

The specimens from which this species is described were crowded in the same can with the *Bimeria* described above, and are in the same dilapidated condition. There is a Tubularian, *Thamnocnidia tubularoides*, from the Bay of San Francisco, described by A. Agassiz (Cat. of N. A. Aclephæ, p. 196), which he says "is readily distinguished from its eastern congeners by the stoutness of the stem and large size of the head." The description is a very meagre one, but from these two characters I conclude that it must be distinct from *T. elegans*, for the latter species has neither a stouter stem nor larger head than *Thamnocnidia spectabilis* of the New England coast.

Eudendrium, sp.

Plate XXXVIII, figure 1.

We have also received from the California coast the perisarc or chitinous portion of what I take to be a species of *Eudendrium*.

Stems stout, erect, dark horn color, strongly annulated throughout, rather sparingly branched; branches sub-erect, springing from all sides of the stem and much divided. Hydranths borne at the extremity of the short ramuli. The entire perisarc is strongly ringed, giving it a close resemblance to the tracheæ of an insect. Height of largest specimen, 80^{mm}.

Santa Cruz, Bay of Monterey, Cal.,—Dr. C. W. Anderson.

Campanularia everta, sp. nov.

Plate XXXIX, figure 4.

Stems rather stout, arising at intervals from the creeping stolon, with two annulations at the base of the hydrothecæ, the lower one

smaller than the upper; the remainder of the stem has a wavy outline or is slightly annulated. Hydrothecæ broadly campanulate, not deep, tapering more or less gradually from the distal end to the base, the rim strongly everted and bearing about fifteen rather shallow teeth. Gonothecæ, large, turgid, nearly cylindrical, tapering a little at the base, borne on short, stout peduncles and with the aperture terminal, small and cylindrical.

Found creeping on an Alga from San Diego, Cal.,—H. Hemphill.

This is a very pretty form and may readily be distinguished by the broad hydrothecæ with their strongly everted, toothed rims. The peculiar shape of the gonothecæ is also very characteristic.

Campanularia cylindrica, sp. nov.

Plate XXXIX, figures 1-1^d.

Stems are simple, unbranched pedicels, of very variable length, more or less annulated over the entire length and with a single well-marked ring at the base of the hydrothecæ, rooted by a creeping, twisted stolon. Hydrothecæ campanulate, nearly cylindrical, tapering but very slightly toward the base, varying greatly in depth, rim armed with about fifteen very shallow, sharply pointed teeth. The gonothecæ also show considerable variation in size, there being occasionally one or two which are at least twice the size of the ordinary form; they are subfusiform, tapering slightly more toward the proximal than the distal end, supported on short pedicels with one or two annulations.

Campanularia fusiformis, sp. nov.

Plate XXXIX, figures 2-2^g.

Hydrocaulus simple, creeping, bearing the pedicels at irregular intervals; pedicels of variable length, usually two or three times the length of the hydrothecæ, never more than six times their length, with a more or less wavy outline. Hydrothecæ small, deeply campanulate, tapering at the base, rim ornamented with about twelve stout, shallow, acute teeth, a single distinct annulation at the base. Gonothecæ small, fusiform, constricted at both ends, sessile, aperture small, terminal.

Vancouver Island,—J. M. Dawson. Found growing on *Sertularia anguina* var. *robusta*.

This species is closely allied to *C. cylindrica* of the Californian coast from which it may be distinguished by the size of the hydro-

thecae and by their shape, not being rounded at the base; by the form of the gonothecae, which are sessile and have a circular terminal aperture.

Found creeping on the old stems of a *Eudendrium*-like form, taken at Santa Cruz, Cal., Bay of Monterey, by Dr. C. W. Anderson.

The variation in the length of the stems is very great; sometimes they are about equal to the length of the hydrothecae, and again they will be five or six times that length. The stolon is quite uniformly twisted and is at least twice the diameter of the stems.

Halecium tenellum Hincks.

Halecium tenellum Hincks, Annals and Mag. of Nat. Hist., 3, VIII, 252, pl. VI.

Plate XXXIX, figure 5.

Some very good specimens of this delicate species have been received from San Diego. There were no gonothecae but the hydrosome is so exactly similar to our New England specimens and to the figure and description of Hincks that I do not hesitate to call it the same. A glance at our figure will show how exactly it corresponds.

Found parasitic on a species of *Bimeria*, collected on the piles of wharves, San Diego, Cal.,—Dr. E. Palmer, 1875.

Sertularia anguina Trask.

Sertularia anguina Trask, Proc. Cal. Acad. Nat. Sci., 112, Plate V, fig. 1. March 30, 1857.

Sertularia labrata Murray, Ann. and Mag. for April, 1860, 250, Plate XI, fig. 2.

Plate XL, figures 1, 1^a, 2.

Stems clustered, simple, erect, straight from the proximal end to the first branch, above the first branch flexuous, becoming more and more so toward the distal end, sparingly branched, divided by transverse joints into short internodes, those below the first branch bearing a single pair of nearly opposite hydrothecae, while those above the first branch have three hydrothecae and give origin to a single branch; branches regularly alternate, ascending, slightly curved, mostly short, a few have a much larger growth and exactly imitate the main stems in every particular; color corneous. Hydrothecae nearly opposite, somewhat flask-shaped or tapering evenly to the distal end without any constriction or flask-shaped neck; aperture usually entire, slightly oblique, facing toward the stem, or with the outer margin much more produced than the inner and in some cases showing a dis-

tinctly sinuous outline. Gonothecæ unknown. Height of largest specimen, 75^{mm}. Plentiful on a large species of *Mytilus* from Monterey, Tomales Pt., Punta Reyes, and on old shells, Bay of San Francisco (Dr. Trask); Santa Cruz,—Dr. Anderson; Vancouver Island,—J. M. Dawson.

Our specimens of *S. anguina* agree so closely with Hinck's description and figures of *S. filicula* that I cannot separate the trophosomes, but the gonosomes being unknown, I prefer to let the species remain distinct rather than to unite them on such incomplete data. Murray also noticed the similarity to *S. filicula* although he only possessed "a minute portion without vesicles."

Dr. Trask says of his specimens, "Their affinity is with that of *S. fallax* of Johnston more nearly than with any other species with which I am acquainted." He could not have known of *S. filicula* at that time or he would at once have noticed the much closer resemblance to that form. He also says that his specimens have four hydrothecæ between each pair of branchlets, while ours have but three: one pair opposite each other and one odd cell in the axil of the branch. His description and figure agree so well, however, in every other respect that I am inclined to regard this as an error of observation.

This description has been made from specimens which were dried before they were sent to us and have since been soaked out in warm water. The perisarc being very stout and durable I do not think that it can have changed to any great extent.

Sertularia anguina, variety *robusta* Clark.

Plate XL, figures 3, 4, 5.

The variety *robusta* differs from the ordinary form in having a stouter stem, larger hydrothecæ, longer pinnae, and in being in every way a much larger and stouter form. The mode of growth, the branching, the shape and arrangement of the hydrothecæ are the same as in the normal form. Gonothecæ borne on the pinnae, more or less fusiform, arising from just below the hydrothecæ, distal extremity slightly curved to one side, the terminal aperture, large, circular. Length of largest specimens, 100^{mm}.

San Diego, Cal.,—Henry Hemphill; taken from kelp roots washed ashore during a storm.

San Diego, Cal.,—D. C. Cleveland, 1875; Santa Cruz, Cal., Bay of Monterey,—Dr. C. W. Anderson; Vancouver Island,—J. M. Dawson.

This variety is very similar to *S. abietina* of Linnæus in many

respects. It has the same mode of growth, the same robust habit, the same style and arrangement of hydrothecæ. There is quite a difference in the natural size figures of *S. abietina* given by Hincks and by Johnston. Our specimens from the New England coast agree very well with Johnston's figure, which is just about twice the size of Hincks' figure.

Sertularia argentea Ellis and Solander (Linn.)

Three fine specimens of this widely distributed species have been found at Santa Barbara, California; they are in good condition and loaded with gonothecæ. The only characteristic which shows any variation is the gonothecæ, which are proportionally a trifle longer and also have the orifice a little larger than in our east coast specimens. This slight difference being the only one and this character varying considerably in the same specimen, I should not think of separating them.

Santa Barbara, Cal.,—Mrs. Ellwood Cooper. Height of largest specimen, 160^{mm}.

Sertularia Greenei Murray.

Sertularia tricuspidata Murray, Ann. and Mag. for April, 1860, p. 250, Pl. XI, fig. 1.

Sertularia Greenei Murray, Ann. and Mag., v, p. 504, 1860.

Cotulina Greenei A. Agassiz, Cat. of N. Amer. Acalophæ, 1865, p. 147.

Plate XXXVIII, figure 6.

Stems erect, slender, densely clustered, simple, thickly branched, basal portion straight, above the first branch becoming slightly flexuous, forming a graceful arc between each two branches, color corneous, usually darkest at the base; joints placed at right angles to the stem and very irregularly distributed, forming internodes which bear from one to eight pairs of hydrothecæ; branches alternate, erect, many of them short; some of the lower ones equal in length to the main stem and closely resembling it in every respect; others from the middle portion of the stem are of a medium length and, like the longest branches from the lower part of the stem, reach to the extremity of the main stem forming a corymb-like structure. The branchlets, like the branches, are mostly short, but a few are of considerable length, extending to the ends of the branches. Hydrothecæ sub-alternate, tapering uniformly to the distal end, with oblique, toothed apertures, which face toward the stem; on the outer edge of the aperture are two large, prominent teeth separated by a deep notch.

Gonothecæ borne in rows on the upper sides of the branchlets; the upper portion cylindrical, the proximal half tapering toward the base, aperture terminal, in a small cylindrical process elevated from the center of the distal end. Height of largest specimens, 90^{mm}.

Bay of San Francisco, Cal.,—Murray; Santa Cruz, Cal.,—Dr. C. W. Anderson; Santa Barbara, Cal.,—Mrs. Ellwood Cooper; Vancouver Island,—J. M. Dawson.

This is an interesting form as it is the only member of the *Sertulariidae* on the American coast having the peculiar aperture to the gonothecæ, by Agassiz called bottle-shaped, though it is by no means an uncommon form among our *Campanulariidae*,—*Obelia gelatinosa*, *O. geniculata* and *O. dichotoma* having the same general form of gonothecæ.

A peculiar discrepancy occurs in the descriptions of Murray and A. Agassiz in regard to the number of teeth on the rims of the hydrothecæ; the former describes them with three teeth, the central one being larger than the two lateral, while Agassiz describes them with four, two prominent exterior points and two smaller ones near the stem. We have quite a large supply of specimens in a good state of preservation and after having carefully examined them all, I cannot find a single hydrotheca that would afford any reason for changing the above description of *two* teeth upon the rim of each cell.

I should judge from Murray's figure that his specimens were not well preserved and by contracting had thrown out the inner margin of the rim, giving it, in some views, the appearance of a tooth. But how he made out one tooth to be much larger than the other two, I am at a loss to understand. And the fact of Agassiz having seen four teeth I am at present unable to account for.

Sertularia furcata Trask.

Sertularia furcata Trask, Proc. Cal. Acad. Nat. Sci., Mar. 30, 1857, 112, Plate V, figs. 2, a, b, c, d, e.

Plate XXXIX, figure 3.

Stems short, unbranched, rooted by a creeping stolon, simple, spreading in every direction forming dense verticillated clusters around the pieces of fucus on which it is usually found, attached to the stolon by a short, slender, twisted process about the length of an internode, divided by transverse joints into short regular internodes each bearing a single pair of hydrothecæ, color corneous. Hydrothecæ opposite, deeply immersed in the stem, with two large, sharp teeth on

the outer margin and a large aperture generally reaching to the stem. Gonothecæ large, sessile, generally borne near the base of the stems though occasionally found scattered over the entire length, of an elongated oval form, sometimes slightly compressed, with a large circular, terminal aperture. Height of largest specimen, 45^{mm}.

Bay of San Francisco and Farallone Islands,—Trask; Santa Cruz, Cal., Bay of Monterey,—C. W. Anderson; San Diego, Cal.,—Dr. E. Palmer; Santa Barbara, Cal.,—Mrs. Ellwood Cooper; Santa Barbara, Cal.,—Dr. L. N. Dimmick.

S. furcata seems to be more nearly allied to *S. Greenei* than to any other Sertularian of the west coast known to us, both having the same style of hydrothecæ, arranged in about the same manner and with similarly toothed apertures. It is the same style seen in *S. operculata* of Linnæus.

S. furcata may be readily distinguished from *S. Greenei* by its entirely different gonothecæ, by the different size of the aperture in the hydrothecæ, by the extent to which the latter are immersed in the stem and by the habit or mode of growth. In general appearance it strongly reminds one of the *Sertularia pumila* of the New England coast.

Sertularella turgida Clark (Trask).

Sertularia turgida Trask, Proc. Cal. Acad. Nat. Sci., Mar. 30, 1857, 113, Plate IV, fig. 1.

Sertularia turgida A. Agassiz, Cat. N. Am. Aculephæ, p. 145, 1865.

Plate XXXVIII, figures 4, 5.

Stems attached by a creeping stolon, sparingly branched, attached to the stolon by a pedicel consisting of three or four rings, short, stout, simple, spreading in every direction from the branches of fucæ and pieces of laminaria which seem to be the favorite stations of this species, divided by oblique joints into short, stout internodes each bearing a single hydrotheca, color light corneous; branches stout, erect, usually about half the length of the main stem and very irregularly arranged, in some cases alternately, in others all the branches spring from one side of the stem and sometimes there seems to be no regular arrangement. Hydrothecæ large, full, alternate, deeply immersed in the stem, the inner angle of the proximal end extending more deeply into the stem than the outer, aperture large, armed with three stout teeth, two of which are larger than the other and are situated on the outer side of the rim, facing the stem, the third tooth

is on the inner margin of the rim midway between the other two. Gonothecæ, arising in the axils of the hydrothecæ, are large and of two forms; the larger form, similar to the gonotheca of *Sertularella polyzonias* Gray (Linn.), is obovate, sessile, armed with a few stout, blunt spines around the distal end, aperture, terminal and at the outer end of a small cylindrical process formed by a constriction or a very sudden tapering near the extremity and surrounded by a number of the largest spines; the smaller form is supplied with about twice as many spines as the larger form and is shorter and proportionally broader, the broadest portion being nearer to the distal end; this variation in form undoubtedly indicates a sexual difference, the smaller form probably being the male and the larger the female gonothecæ; both forms have the surface more or less roughened by transverse wrinkles. Length of largest specimen, 38^{mm}.

Bay of San Francisco, Monterey, Tomales Point, Cal., on mollusca and algæ.—Trask; Santa Cruz, Bay of Monterey, Cal.,—Dr. C. W. Anderson; San Diego, Cal.,—D. C. Cleveland; Vancouver Island, —J. M. Dawson.

The nearest ally of *S. turgida* is the *S. polyzonias* of Gray, to which in many respects it bears a striking resemblance.

Hydrallmania Franciscana Clark (Trask).

Plumularia Franciscana Trask, Proc. Cal. Acad. of Nat. Sci., vol. i, p. 113, Pl. IV, fig. 3.

Plumularia gracilis Murray, Ann. and Mag. of Nat. Hist. for April, 1860, p. 251, Pl. XII, fig. 1.

Trask and Murray both had representatives of a species which they referred to the genus *Plumularia* and which, from their descriptions and figures, undoubtedly belongs to Hinck's genus *Hydrallmania*, though at the time their descriptions were published, this genus had not been recognized. This species is certainly very close to *H. fulcata* of Hincks, but Murray, who had an opportunity of comparing the two, says they are distinct.

Unfortunately we have had no specimens.

Bay of San Francisco, Cal., among rejectamenta on the beach, —Trask.

Plumularia setacea Lamarek.*Sertularia pinnata*, β , Linn., Syst. Nat., 1312.*Sertularia setacea* Pallas, Elench., p. 148.*Plumularia setacea* Lamk., An. s. Vert. (2d ed.), ii, 165.

Plate XLI, figures 1, 2.

Stems simple, slender, erect, rooted by a creeping stolon and divided by transverse joints into short internodes of uniform size, regularly branched; pinnae alternate, regularly arranged, one from each internode, arising from the stem by a prominent process produced from the outer and upper side of each internode, divided by transverse or slightly oblique joints into internodes of two sizes arranged alternately. In large specimens 150^{mm}. to 300^{mm}. long, the main stems are considerably branched, the branches alternately arranged and clustered, extending quite or nearly to the distal end of the stem; the branches give off branchlets, which like themselves resemble the main stems in every particular. Hydrothecae with an even rim, small, borne on the larger sized internodes of the pinnae. Nematophores compound, those on the pinnae not quite equal in length to the hydrothecae, those on the main stems a little longer than the hydrothecae; three on each internode of the stem, two in the axil of each pinna, one on the opposite side of the internode near the base, one only on the upper side of the smaller joints of the pinnae and three on the larger joints, one just below the hydrotheca, and one on each side of the aperture. Gonothecae sessile, borne in the axils of the pinnae; female elongate oval, produced at the distal end into a tubular neck with a discoidal, terminal orifice; male, smaller than the female, fusiform, and with a much smaller aperture.

Santa Cruz, Cal.,—C. W. Anderson; San Diego, Cal.,—Dr. E. Palmer; San Diego, Cal.,—H. Hemphill; Vancouver Island,—Dawson.

Most of our specimens consist of dense clusters of the delicate shoots, about 50^{mm}. to 80^{mm}. long, and usually attached to some large, coarse alga. Ellis' old name of "*Sea Bristles*" was well chosen, for it conveys quite an accurate idea of the appearance of these smaller forms. The larger forms are more branched, usually of a darker color and have a closer resemblance to hair than to bristles; one of our largest specimens from San Diego consists of a tuft 200^{mm}. in length composed of about three hundred branched shoots; this had been washed ashore and was found by Dr. Palmer; a still larger specimen was dredged off San Diego in six to ten fathoms by Mr. Hemphill, which measures 300^{mm}. in length and forms a thick cluster of about a thousand shoots.

Aglaophenia struthionides Clark (Murray).

Plumularia struthionides Murray, Ann. and Mag. of Nat. Hist. for April, 1870, 251, Plate XII, fig. 2.

Aglaophenia franciscana A. Agassiz, Cat. N. A. Acalephæ, p. 140, 1865.

Plate XLI, figures 3-3^d.

Stems rooted by a creeping stolon, simple, erect or spreading in every direction, divided by slightly oblique joints into very short internodes of equal length, each bearing a single pinna, varying from the lightest to the darkest horn-color; shoots tall, stout, plumose, tapering slightly toward the base, the distal end abruptly pointed; pinnae slightly curved, sub-erect, unbranched, not in the same plane, the sides bearing the hydrothecæ curving toward each other, divided into short internodes by slightly oblique joints, each internode bearing a single hydrotheca. Hydrothecæ large, cup-shaped, expanding toward the distal end, aperture large, patulous, rim denticulated, armed usually with eleven, sometimes nine, sharp, uneven teeth. Nematophores tubular, the lateral ones of medium size, projecting ear-like from the sides of the hydrothecæ, the anterior one long, adnate for the greater part of its length, free near the distal end, extending nearly or quite to the edge of the toothed rim, aperture small, discoidal, terminal; those upon the corbulæ are a trifle larger than the lateral ones and are arranged in transverse rows, the ends of which do not meet. Corbulæ large, cylindrical, with numerous ridges (ten to sixteen) composed of oblique rows of nematophores; usually from two to six hydrothecæ at the base of the corbula. In luxurious specimens the corbulæ are very abundant, there being between seventy and eighty on a single shoot. Length of largest specimen, 150^{mm}.

Bay of San Francisco,—Trask; San Francisco, Cal.,—A. Agassiz; Santa Barbara, Cal.,—L. F. Dimmick; Santa Barbara, Cal.,—Mrs. Ellwood Cooper; Santa Cruz, Cal., Bay of Monterey,—C. W. Anderson; San Diego, Cal.,—D. C. Cleveland; Vancouver Island,—Miss Mitchell; Vancouver Island,—J. M. Dawson.

In the various lots of Hydroids which we have received from the western coast, this species has always been the most abundant. It seems to be as common and as widely distributed on the western coast of the United States as *Sertularia pumila* is upon the eastern coast. Both are also very often found parasitic on algæ, but *A. struthionides* is often found in deeper water than *S. pumila*.

As Murray remarks, this species is nearly allied to *Plumularia cristata*, the *A. pluma* of Linnaeus, but is much closer in habit, the hydrothecæ are wider-mouthed and shallower and the teeth upon the rim are unequal. An important error occurs in the synonymy of this species given by Mr. A. Agassiz. He has, under the name *Aglaophenia Franciscana* the following synonymy:

Plumularia Franciscana Trask.

Plumularia struthionides Murray.

The *Plumularia Franciscana* of Trask belongs to the genus *Hydrallmania* of Hincks, as a glance at the figures and description of Trask will show; and it is synonymous with the *Plumularia gracilis* of Murray. Hence the name of the above described species should not be *A. Franciscana*, but *A. struthionides*.

EXPLANATION OF PLATES.

PLATE XXXVIII.

- Figure 1.—*Eudendrium*, sp.; from Santa Cruz, Cal.
 Figure 2.—*Tubularia elegans*; *a*, cluster of medusæ buds; *a*, *a'*, and *a''*, buds in different stages of development; *a'''*, an actinula escaping.
 Figure 3.—*Bimeria gracilis*; *a*, *a'* and *a''*. sporosacs.
 Figure 4.—*Sertularella turgida*; *a*, the gonangium or gonotheca.
 Figure 5.—*Sertularella turgida*; another form of gonotheca.
 Figure 6.—*Sertularia Greenei*; *a*, and *a'*, the gonangia (gonothecæ); *h*, hydrotheca.

PLATE XXXIX.

- Figure 1.—*Campanularia cylindrica*; *1a*, the same showing the full length of the pedicel; *1b*, the female gonotheca; *1c*, an abnormally developed hydrotheca; *1d*, the male gonotheca.
 Figure 2.—*Campanularia fusiformis*; *2a*, *2b*, and *2d*, the same showing the amount of variation in the width and depth of the hydrothecæ; *2e*, *2f*, and *2g*, the gonothecæ; *r*, the rootstock or creeping stem.
 Figure 3.—*Sertularia furcata*; *a*, and *a'*, the gonothecæ.
 Figure 4.—*Campanularia everta*; *a*, and *a'*, the gonothecæ; *r*, the creeping stem.
 Figure 5.—*Hylecium tenellum*; from San Diego, Cal.; *r*, the creeping stem.

PLATE XL.

- Figure 1.—*Sertularia anguina*; *1a*, a single hydrotheca showing the outline of the outer margin of the rim.
 Figure 2.—*Sertularia anguina*; a portion of the main stem.
 Figure 3.—*Sertularia anguina*, var. *robusta*; a portion of the main stem.
 Figure 4.—The same; portion of a branch: *a*, gonotheca.
 Figure 5.—The same; with a monstrosity, *a*, at the extremity of the branch.

PLATE XII.

- Figure 1.—*Plumularia setacea*; portion of a branch with nematophores and female gonotheca, *a*.
- Figure 2.—The same; a portion of the main stem; *n*, nematophores; *h*, hydrotheca; *2a*, male gonotheca.
- Figure 3.—*Aglaophenia struthionides*; a portion of a pinna; *3a*, *3b*, and *3c*, different views of the same; *n* and *n'*, nematophores; *h*, hydrotheca; *3d*, corbula; *a*, the body of the corbula; *b*, the wing-like processes at the base; *n*, and *n'*, the nematophores; *h*, the hydrotheca.

VII. ON THE ANATOMY AND HABITS OF NEREIS VIRENS. BY
FREDERICK M. TURNBULL.

Read January 19, 1876.*

THE *Nereis* (*Alitta*) *virens*, which is one of the largest and most common of our marine annellids, is found under stones or burrowing in the sand and mud of sheltered shores, both at low-water mark, and at a considerable distance farther up. It grows to the length of eighteen inches or more, and is quite stout in its proportions.

It is very active and voracious, feeding on other worms and various kinds of marine animals which it finds when burrowing in the sand.† It will even devour its own immediate relatives, if hungry when it meets them. It suddenly thrusts out its proboscis and seizes its prey with the two powerful jaws, then withdraws the proboscis, the jaws closing at the same time. In this way it will tear large pieces from the body of its victim, being able, at one bite, to cut in two a worm of its own size. One which I had confined in a small dish of water, bit its own body in two pieces at the middle. As the proboscis is turned inside out, when it is protruded, whatever has been seized by the jaws will be drawn by them inside the proboscis as soon as the latter resumes its natural position, the proboscis then acting as a sort of gizzard.

These worms, by secreting a viscid fluid, will surround themselves in a few minutes with a translucent sheath which binds the grains of sand together, forming a loose and flexible tube. They remain most of the time in these tubes, which are nearly always situated in sand and mud or under flat stones, and they move in them with consider-

* Abstract of a graduation thesis presented at the Sheffield Scientific School, June, 1875.

† Later observations show that this species does not restrict itself to an animal diet. Several large specimens, taken by me in October, 1875, had the intestine completely filled with algæ of several species, among which *Ulva latissima* was most abundant. The algæ were torn into fragments and large shreds and rolled together into long pellets, but even after passing through the intestine their nature could be easily recognized.—A. E. VERRILL.

able freedom and rapidity, pushing themselves along by means of their aciculæ, setæ and ligulæ, the aciculæ being controlled by special muscles. They hold their tentacular cirri straight out in front of them, as they move, in order to have warning of anything that they may approach.

The tantog, scup and other fishes dig them out of the sand and devour them eagerly. But at certain times, especially at night, they leave their burrows, and swim about like eels or snakes, in large numbers, and at such times fall an easy prey to many kinds of fishes. This habit seems to be connected with the season of reproduction. They were thus observed swimming at the surface in the day time, near Newport, in April, 1872, by Mr. T. M. Prudden and Mr. T. H. Russell, and several times by Professor Verrill, later in the season. At Watch Hill, R. I., April 12th, 1873, Professor Verrill found great numbers of the males swimming in the pools among the rocks at low-water, and discharging their milt. The males were also seen swimming in the tide-pools and shallow waters at Savin Rock, April, 1875, by Professor D. C. Eaton and Mr. Kleeberger. The *Nereis virens* is abundant at all seasons of the year, in most places along the sandy and muddy shores, both of the sounds and estuaries, burrowing near low-water mark. It occurs all along the coast, from New York to the Arctic Ocean, and is also found on the northern coasts of Europe.*

The body consists of a large number of rings or segments. This number varies with the size and age of the worm. It may be less than one hundred, or as many as two hundred. The increase in length takes place by the addition of new posterior segments, in advance of the caudal segment. New segments may also be formed when a part of the body is broken off, and in this way a considerable part of the posterior portion of the body can be completely reproduced. The head (figs. 1 and 17) is very fully developed, being provided with two pairs of eyes and two pairs of antennæ. It is attached to the dorsal side of the first segment, which is called the buccal or mouth-ring (figs. 1 and 15, *d*), because it contains the month (fig. 15, *m*). There are also four feelers, called tentacular cirri (fig. 1, *ee*, *ee'*, *e* and *e'*), arising from the buccal ring on each side of the head.

The abdominal rings (fig. 1, *g*) follow the buccal ring. Each one of these has a pair of lateral lamelliform appendages (figs. 1 and 17, *h*), used as paddles in swimming and also serving the purpose of gills. The last segment or ring of the body bears a pair of cirri, similar to the tentacular cirri, and also contains the anal orifice.

* See, for habits of this and allied species, Professor Verrill's report in First Report of U. S. Commission of Fish and Fisheries, p. 318. 1873.

The males are easily distinguished from the females by differences in color and in the form of the side appendages. The color of the body in the male is an intense steel-blue, which blends into green at the base of the lateral appendages. These have a rich and brilliant sea-green color which is heightened by the complimentary effect of the numerous red blood-vessels they contain. The latter are especially noticeable toward the posterior end of the body where the skin is thinner and less opaque, so that the appendages, with their network of capillaries, appear bright red.

In the female the body is of a dull greenish color, with a slight tinge of orange and red. The appendages are orange-green at the base, and become bright orange toward their extremities; but sometimes they are greenish throughout. The whole surface of the skin, in both male and female, is iridescent, reflecting bright hues when placed in the light.

The head (figs. 1 and 15, *a*) is small, and flattened on the dorsal and ventral sides. From the position of the anterior eyes the sides taper toward the anterior extremity, where it is rounded off and terminated by a pair of small antennæ (figs. 1 and 15, *b, b*). There are two pairs of eyes on the upper surface of the head, one pair near its base and another pair more anterior and farther apart. The anterior eyes are situated near the middle and on the broadest portion of the head. On each side of the head, attached to its anterior half and also to the buccal ring, is a large antenna or palpus, as it is sometimes called (figs. 1 and 15, *c*). These are stout, fleshy and somewhat contractile organs. Each has a small rounded lobe at the tip (figs. 1 and 15, *c'*).

The buccal segment and the head constitute the cephalic or head region. The tegument about the mouth is wrinkled and folded longitudinally, presenting an appearance like the mouth of a purse, when drawn together by strings. The tentacular cirri are long, slender and quite flexible. They receive their nerves from the first abdominal nerve-ganglion, whereas the antennæ receive theirs from the head-ganglion.

There are four pairs of tentacular cirri, two dorsal (fig. 1, *ee, ee'*) and two ventral (fig. 1, *e, e'*). The ventral ones are situated nearer to the palpi than the dorsal, and hence the two are called respectively the internal and external tentacular cirri. The relative lengths of the tentacular cirri are shown in fig. 1. The two large dorsal tentacular cirri (*ee, ee'*) are longer in the male than in the female. In the male they will reach to about the middle of the ninth segment, when laid along the back. In the female they will reach to about the middle of the

fifth segment. The other tentacular cirri have more nearly the same length in both sexes.

The abdominal segments increase in size to about the eighth, and then remain nearly the same for some distance along the middle region, but the posterior rings decrease in size, causing the body to taper until it becomes quite slender. The appendages, also, are longer and broader along the middle region than toward either extremity. On the ventral side of the segments the part continuous with the feet is smooth, but the other parts show the strong transverse muscles (fig. 26), which, by their contraction, lessen the size of the body cavity.

The lateral appendages or feet of *Nereis virens* are quite complicated and wonderful organs. They are biramous (fig. 12), having two rami, one dorsal (A) and the other ventral (B). The aciculæ of each foot arise from a crypt which is attached by shroud-like muscles to the base of the foot. The setæ arise from the interior of the two rami. The feet are complicated by the addition of other organs, serving for locomotion, sensation and respiration.

The respiratory organs, often called ligulæ, are more or less flattened lobes with their teguments very thin and filled with a rich vascular network (figs. 22, *s*, 25, 26). The upper ramus has two ligulæ, a superior one (fig. 12, *b*) on its upper, and an inferior one (fig. 12, *d*) on its lower side. At the base of the superior ligula, on a sort of shoulder of its upper edge, is the dorsal cirrus (fig. 12, *a*). There are also two setigerous lobes smaller than the ligulæ, one on each side of the opening through which the setæ protrude (fig. 12, *c* and *k*). The anterior (*c*) is longer than the other (*k*) and is connected with the inferior ligula (*d*) of the upper ramus. The acicula is attached to the inner walls of the ramus forming a partition, which terminates with the end of the acicula between the lobes *c* and *d* (fig. 12) and generally forms, in the middle and posterior parts of the body, a third and smaller lobe (fig. 12, *c'*).

The lower ramus lacks a superior ligula, but it has an inferior one (fig. 12, *g*) more rounded and not so broad as the others. In the lower ramus, as in the upper, there is normally only one fascicle of setæ, but in the lower ramus it is divided into two clusters by the acicula, which, by its attachment to the inner walls of the ramus, forms a partition. Here, as in the upper ramus, there are two flattened setigerous lobes, about equal in length, one on each side of the opening for the setæ (fig. 12, *f* and *e*), and the partition formed by the acicula extends to the extremity of the anterior one, dividing it into two small lobes (fig. 12, *e* and *e'*).

The inferior ligula of the upper ramus, with its two lobes (fig. 12, *c, d*) corresponds, apparently to the lobe (*e*) and its divisions, while the lobe (*k*) of the upper ramus corresponds to the lobe (*f*) of the lower ramus. In the upper ramus, the acicula and its partition do not divide the bunch of setæ, as it comes out above the acicula. The ventral cirrus arises from a slight protuberance of the inferior side of the lower ramus (fig. 12, *h*).

The feet are not all alike from one end of the body to the other; they change in form most along the anterior region, and in the first five the variation is considerable; along the middle and posterior regions, it is slight in comparison.

I have figured the first five, the forty-fifth, and the one hundred-and-ninth feet of a female worm, having one hundred and twenty-one segments; also the first, fifth, forty-fifth, and one hundred-and-ninth in a male worm of nearly equal size, but having one hundred and sixty-one segments. These figures show the posterior view of the feet. Those of the female will be first described, and then compared with those of the male.

In the first foot of the female (fig. 3) the upper ramus has only the dorsal cirrus and the superior ligula developed. The former is about one-fifth longer and a little less than one-third as broad as the latter, being very well developed, while the ligula is somewhat rudimentary. The ligula is rounded and simple in shape, having a constriction at its base. Of the lower ramus, all the parts are present. The posterior setigerous lobe (*f*) is longer than the anterior setigerous lobe (*e*), and is oval and flattened. The anterior setigerous lobe (*e*) is seen to be divided by the acicula into two smaller terminal lobes (*e* and *e'*), indicated by dotted lines where they are covered by the posterior setigerous lobe (*f*). The inferior ligula (*g*), of the lower ramus, is of about the same size and shape as the superior ligula (*b*), and the inferior cirrus (*h*), is like the superior cirrus (*a*). The setæ of the lower ramus, as shown in the figure, extend just beyond the posterior setigerous lobe (*f*). They are somewhat rudimentary and are for the most part blunt. The acicula is also rather small.

The second foot (fig. 4) has the anterior setigerous lobe (*e*), and the acicula a little larger and better developed, and the dorsal cirrus is seen to be a little shorter than that of the first foot; otherwise the second foot is scarcely different from the first.

In the third foot (fig. 5) the upper ramus has developed an inferior ligula, setæ and acicula. It also has an anterior setigerous lobe (*e*). This lobe is a part of the inferior ligula (*d*). It shows more

distinctly in some of my figures of the feet of the male. The superior ligula is of about the same size and shape as the inferior ligula. The lower ramus is a little more developed than it is in either the first or second feet, but the inferior cirrus (*h*) is shorter.

The fourth foot (fig. 6) is more highly developed than the third; but the inferior cirrus is shorter in comparison with the lower ramus.

The fifth foot (fig. 7) is still more highly developed than the fourth, and the superior ligula is larger than it is in the preceding feet. The dorsal and ventral cirri are both smaller. If we now compare the figure of the fifth with the figures of the first and second feet, the difference is seen to be considerable, particularly in the length of the cirri.

The superior ligula (*b*) begins to increase in size at the fifth foot, and continues to do so, until in the forty-fifth (fig. 8, *b*) it is larger than any other part of the foot. It has also become flat and pointed. The shoulder (*i*) is much larger. The inferior ligula (*d*) is also flat and pointed. A posterior setigerous lobe (*k*) is now large enough to be easily seen. The posterior setigerous lobe (*f*) of the lower ramus is somewhat pointed, and its lower edge is oblique. The anterior setigerous lobe (*e*) is now as long as the posterior setigerous lobe, and its two divisions are nearly equally developed. The inferior ligula (*g*) is rounded and somewhat tapering toward its extremity. The dorsal and ventral cirri are now quite small, particularly the latter, while otherwise the foot is much larger. The setæ have become longer and more perfect, from the anterior toward the posterior, attaining their maximum in the middle region of the body. The aciculæ are also larger here than in either the anterior or posterior regions; although, in the posterior region they are longer in comparison with the size of the foot than anywhere else.

The one hundred and ninth foot (fig. 9), as shown by the figures, is nearly the same in form as the forty-fifth, but smaller.

On comparing the first, fifth, forty-fifth and hundred and ninth feet of the female with the same in the male, we find that there is a marked difference in the cirri. The dorsal cirrus in the first foot of the male (fig. 10) is one-half longer than it is in the female, and it is larger in proportion. In the fifth foot (fig. 11) the difference is the same. In the forty-fifth (fig. 12) the dorsal cirrus of the male is longer and also slenderer than it is in the female. In the one hundred and ninth foot (fig. 13) the dorsal cirri are about the same in length, but those of the male are more slender.

The difference between the dorsal cirri decreases toward the posterior end of the body, being greatest in the anterior segments.

The *ventral* cirrus of the first foot of the male (fig. 10) is not quite so long as the *dorsal* cirrus, but it is a little longer than the ventral cirrus of the female. In the fifth foot (fig. 11) it is a little longer and much more slender than it is in the female. In the forty-fifth (fig. 12) it is twice as long and of the same diameter as it is in the female. In the one hundred and ninth (fig. 13) there is the same difference which we see in the forty-fifth.

The difference between the ventral cirri of the male and female, increases from anterior to posterior, being greatest in the posterior segments.

Beginning with the forty-fifth foot (fig. 12) a shoulder (*x*) appears on the dorsal side of the lower ramus in the male. It increases in size and definiteness farther back, but finally disappears. This shoulder is never seen in the female, and is a good character for distinguishing the sexes, but it requires microscopic examination. The difference between the cirri of the male and female is sufficient to distinguish them without the use of a lens.

The setæ under the microscope are very delicate and beautiful (figs. 2, 2*a*). They consist of two parts, the shaft (*a*) and the blade (*b*). The shaft has a transversely striated appearance, which is exceedingly regular. The blade is held in a sort of socket in the end of the shaft, and one edge is toothed like a saw. There are two forms: one in which the blade is short, having its extremity blunt and slightly hooked (fig. 2), and one with the blade tapering to a fine point, the blade being long and slender (fig. 2*a*). The length varies and with it the delicacy of the point. In most of the setæ the latter is so sharp that it seems to vanish, and can be seen only with a high power. Those setæ which are hooked have the hooks and toothed edge turned upward; and these are always confined to the lower ramus, and to the lower side of the latter in both bunches. In the middle and posterior regions these hooked setæ disappear, their place being taken by the other kind; but the latter are shorter than those in the upper part of the bunch. The same forms of setæ are found in both male and female. Those of the anterior feet are shorter than those in the middle region. When viewed by transmitted light the bunches present all the colors of the spectrum.

The aciculæ (figs. 12 and 22) are simple thorns, in the form of an elongated cone, generally a little bent. They are black, except at the base, showing through the translucent integuments of the foot. At the base they are hollow and therefore lighter colored.

Muscular System.

The walls of the body contain two muscular layers, which are well defined. The first is thick on the ventral side of the body (fig. 22, *k*), but is thinner elsewhere. It forms in each ring a muscular plane of transverse fibres. The second is placed within the preceding and is formed of great fasciæ which are attached to the interannular partitions (fig. 16, *b*). This layer does not exist on the ventral side, between the bases of the feet. The interannular partitions (fig. 16, *a*) are attached to the intestine, which they hold in place, dividing the general cavity of the body into a series of chambers (fig. 16); but as the inner subcutaneous layer does not extend over the ventral floor of the cavity, the chambers open into one another beneath the intestine. I have already mentioned the museles attached to the fleshy knobs, which hold the aciculæ. These museles (fig. 22), when they contract all together, force the aciculæ outward. When different ones contract they move the aciculæ to one side.

The *Nereis virens* in swimming moves its body laterally, like a snake. It sometimes has also an undulatory movement, up and down. These motions are all produced by the subcutaneous museles. The lateral appendages are used as paddles, but their principal use is to push the worm along in its tube, and for crawling. They are greatly aided by the stiff aciculæ, controlled by their special museles (fig. 22), and by the setæ. When burrowing, the proboscis is used to push away the sand in front and is then withdrawn, while the body is moved forward partly by a vermicular motion and partly by the side appendages.

Alimentary System.

The proboscis is a very remarkable organ and constitutes a formidable weapon. It is divided by M. DeQuatrefages into three regions: the pharyngeal, the dental, and the œsophageal (figs. 16–18). The mouth has already been partially described. The walls of its cavity are connected by several small, delicate museles, with the walls of the body cavity, as I have shown in figure 16, *m* and *n*. The pharyngeal region commences immediately back of the buccal cavity (figs. 16 and 20, *x*), and has two muscular partitions (fig. 16, *e*).

The dental region (20 and 16, *p*) is very muscular, and is provided with a considerable number of small teeth, or denticles, which are arranged in groups on the anterior, inner surface of its walls. It also has two large and powerful jaws attached by their hollow bases to the museles of the posterior inner surface. The worm has the power of turning this dental region inside out.

When the proboscis is inside the body, it takes the position shown in figures 16 and 20, the œsophagus being curved and pushed back against the intestine; but when it is forced out, the œsophagus straightens, and that part of the proboscis which is protruded takes the position shown in figures 17 and 18, these figures showing the arrangement of the jaws and denticles, figure 17 giving the dorsal view and figure 18 the ventral view; the buccal ring is very much stretched; and the head, antennæ and tentacular cirri are forced over on the back (fig. 17). The jaws are imbedded in and attached to the special muscles, in such a manner that the more the proboscis is protruded the farther apart their points move, and when the proboscis is withdrawn they close like a pair of scissors, their points crossing. The jaws (fig. 19) are curved like hooks, and have their inner concave edge denticulated with about ten teeth. They and the denticles are composed of a black chitinous material. The denticles, which are conical and pointed, are not attached to muscles, but are simply imbedded in the surface. The number and size of the denticles, and even their positions, vary considerably in the different specimens examined. I think it would be hard to find two specimens exactly alike in this respect. Among seven worms, I found five with one denticle, one with two denticles, and one with seven denticles on the median anterior area of the dorsal side (fig. 17, *o*).

Among six worms, I found two with two denticles and four with none at all on the posterior median area of the dorsal side (fig. 17, *t*). On the left submedian anterior area of the dorsal side (fig. 17, *l*), the number of denticles varied from three to eight; and on the right submedian anterior area of the dorsal side (fig. 17, *l'*) from two to eight. In only one case did the two last areas have the same number of denticles. On the left submedian posterior area of the dorsal side (fig. 17, *s*) the number of denticles varied from none to five; and on the right submedian posterior area of the dorsal side (fig. 17, *s'*) from one to four.

The denticles on the lateral and ventral posterior areas (fig. 17 and 18, *r*) vary considerably in number, position and size. Among four worms, the number of denticles on the right lateral anterior area (figs. 17 and 18, *n*) varied from twelve to twenty-five; and on the left lateral anterior area (figs. 17 and 18, *n'*) from eleven to thirty.

Among five worms the number of denticles on the anterior median area of the ventral side (fig. 18, *y*) varied from four to twelve. On each of the two submedian areas of the ventral side (fig. 18, *x'*) there was one denticle.

In some of the large worms the denticles are as small as those of smaller worms, while in other cases they are much larger. The large denticles probably are knocked off in some way and new ones grow in their place. This would account in part for the large number of very small denticles, and also for the variations in number.

Attached to the anterior end of the œsophagus, one on each side, are two salivary glands (figs. 16 and 20, *j*). These are free except at one end, and are ciliated on their outer surface. The intestine proper (figs. 16 and 20, *r*) is straight and is constricted somewhat by the muscular partitions of each segment through which it passes. It is brilliant greenish yellow in color and is surrounded by a regular capillary network of blood vessels (fig. 20).

The internal surface of the œsophagus is tessellated with low, rounded papillæ or tubercles. These are regular in shape and equal in size. Their sides are diagonal to the length of the œsophagus. They show through the walls of the œsophagus, so that its outside appears tessellated with dark squares, and as the œsophagus is stretched or contracted they become diamond shaped. The tubercles are of a dull color, between brown and yellowish green.

The interior surface of the intestine is also covered with regular longitudinal rows of low, rounded tubercles, which are much smaller than those of the œsophagus. They are greenish-yellow like the outside of the intestine. The end of the œsophagus projects into the cavity of the intestine, and its opening, which has sphincter and also longitudinal muscles, can be enlarged or contracted to a considerable extent. The outer surface of this end of the œsophagus is continuous with and like the internal surface of the intestine. The latter secretes a brown fluid in its interior and probably acts as a hepatic organ. The dental portion of the proboscis acts like a gizzard, and the œsophagus is probably a sort of stomach.

Circulation and Respiration.

The circulatory system is highly developed and complicated. The blood is red, and the vascular system is complete and closed. The principal vessels have a longitudinal course, occupying the whole length of the median line of the body, one as a dorsal (figs. 20–24, *a*), and the other as a ventral vessel (*b*). They are contractile, and by a sort of peristaltic motion the blood is pushed forward in the dorsal vessel, and in the opposite direction by the ventral vessel. The dorsal vessel is visible for nearly its whole length through the more or less transparent walls of the body, and its blood can be seen

moving in a series of waves toward the head. The ventral vessel sends off, in each segment of the body, except a few in the region of the proboscis, two smaller vessels, one on each side. These two vessels fork, each sending a branch to the inferior ramus of the foot of the next segment to the rear (figs. 20-24, *f'*), and another larger branch (*e*) around the intestine, by the side of the transverse partition, to the dorsal vessel, receiving, also, on its way, a vessel from the upper ramus of the foot of its own segment (*d*). Besides these principal lateral vessels, there are five other vessels on each side in each segment, coming from the ventral vessel (fig. 20). These form a loose but regular network that surrounds the intestine and is connected with five other convoluted vessels, which join the dorsal vessel. This network on the intestine probably supplies the hepatic organ with material for its secretion, and very likely may receive nutritive material from the digested food. The blood moves in waves, at regular intervals, through the peripheral vessels (figs. 20-24, *e*) to the dorsal vessel, but I could not see in which direction the blood moved in the network. The blood is forced into it at each pulsation of the dorsal vessel, but the normal flow may be in the opposite direction. The peripheral vessels are also connected with this network (fig. 20). The dorsal and ventral vessels are connected at the posterior extremity of the body by a simple peripheral vascular ring (fig. 23, *e*), in which the blood flows from the ventral to the dorsal vessel. In the region of the proboscis, the ventral vessel sends lateral branches directly to all the feet but the first three (figs. 20 and 21). It then sends a pair of vessels to the œsophagus (figs. 20 and 21, *e*), which pass back along the œsophagus, one on each side, as far as the intestine, being connected with smaller vessels on the surface of the œsophagus. The ventral vessel next sends off a pair of vessels which expand into capillary networks, one on each side (figs. 20, 21, *s* and *g*). Each of these networks sends small branches to the first three feet on its own side, and then merges into a vessel (figs. 20 and 21, *k*), which goes to the base of the tentacular cirri. The ventral vessel now goes upward to the under surface of the proboscis, and there divides into three branches (figs. 20, 21, *t*, *t* and *l*). The middle branch (fig. 21, *l*) passes under a muscle and along the median line of the ventral surface, as far as the pharynx, where it divides into two, forming a small vascular ring (fig. 21, *n*) about the latter. The two lateral branches pass upward and backward on the proboscis, each expanding into a remarkably rich and delicate network on its own side of the proboscis (figs. 20 and 21, *u*). From each of these networks a vessel

(*v*) passes to the base of the head, where it joins the dorsal vessel (*a*), thus completing the circulation. From this junction small vessels probably go into the head and antennæ. The lateral vessel passing to the lower ramus divides into branches ramifying on that portion which is continuous with the foot on the ventral side of the segment, and also over the lower ramus (figs. 22 and 26). There a connection is made with vessels of the upper ramus, and I think this is done by the vessel marked *x* in figures 22 and 24, because it is quite large at the base of the inferior ligula of the upper ramus, and grows smaller at first and then swells out again before joining the vessels of the upper ramus, in the superior ligula. The branch *d* (figs. 20, 22 and 24), coming from the dorsal ramus, receives blood from that organ and also from a peculiar and beautiful arrangement of capillaries on the dorsal side of the body (fig. 25).

In the first four segments, in the region of the proboscis, the dorsal vessel has no branches, but in the remaining segments, commencing with the fifth, it has five pairs of long peripheral branches (fig. 20, *c, c'*), corresponding to the peripheral vessels of the posterior part of the body. They are not attached to the proboscis, but are simply connected with the dorsal and ventral vessels by their ends. The one coming from the dorsal vessel in the fifth segment is connected with the ventral vessel in the fourth segment (fig. 20.) The first three feet probably do not act as gills, as very little blood is sent to them. The two networks (figs. 20 and 21, *u*) on the proboscis are probably for carrying on the exchange between the blood and the liquid of the body cavity.

The respiration is carried on by the red fluid in the beautiful arrangement of capillaries on the body and feet, especially the latter. The flat ligulæ of the feet are exceedingly delicate in structure and take the place of gills, absorbing the oxygen from the water to purify the blood received from the ventral vessel, which then returns to the dorsal vessel.

The disposition of this respiratory arrangement is shown in figures 22, 24, 25 and 26.

The Nervous System.

The nervous system of *Nereis virens* (figs. 27 and 28) is complicated and well developed, being composed of a series of ganglia, sending out branches and connected by nervous cords. It lies mainly on the ventral floor of the body beneath the large ventral vessel. The first and largest ganglion (figs. 27 and 28, *a*), analogous to the brain of higher animals, is situated in the head. It is composed of several

smaller ganglia joined together. It bears the eyes, on four short nervous peduncles, on its dorsal side. In front it sends four nerves to the antennæ (*b*, *b* and *c*, *c*, figs. 27 and 28). Laterally it sends out two branches called the connectives (*d*, *d*), which pass around the mouth and proboscis to join the first of the abdominal ganglia (*h*, fig. 27). Near the junction of the connective with the head ganglion, is a small ganglion sending nerves to the internal tentacular cirri (*e*, *e*, figs. 27 and 28). The connectives, near their lower extremity, send two nerves (*g*, *g*, fig. 27) to a series of ganglia and nerves on the ventral side of the proboscis (*w*, *w*, fig. 27). There is also an accessory connective (figs. 27 and 28, *d'*) on each side, passing from the first abdominal ganglion to the ganglion supplying the external tentacular cirri (*e'*, *e'*, figs. 27 and 28). This accessory connective also has a ganglion (*n'*, fig. 27) at the middle, sending nerves to the muscular partitions of the proboscis.

Each of the first three abdominal ganglia sends, from its anterior portion, on each side, a nerve that forks, one branch (fig. 27, *n*) going to the muscular partition and the other (*o*) passing through the partition to the preceding segment. In the remaining abdominal ganglia, beginning with the fourth, the branches *n* and *o* become separate nerves (fig. 27). From the posterior portion of these ganglia a nervous trunk on each side (*m*), goes to each foot, where there is a small ganglion (*k*) sending off a cutaneous branch and a branch (*i*), supplying nerves to the foot.

The ganglia (fig. 28, *e*, *e*) and the head-ganglion (*a*) send some very slender nerves (fig. 28, *z*, *z*) to a series of ganglia on the dorsal side of the proboscis (*y*, *y*). In figure 27 the series of ganglia (*w*, *w*) are drawn as if the proboscis had been revolved about a line drawn through its anterior end, so that the ventral surface would be uppermost. In figure 28 the ganglia (*y*, *y*) are in their natural position. The dorsal ganglia (fig. 28, *x*, *x*) are connected with the ventral ganglia (fig. 27, *v*, *v*) by means of nervous cords; the dorsal ganglia (*s*, *s*, fig. 28) with the ventral ganglia (fig. 27, *l*, *l*), by means of nerves passing around the proboscis outside the points of the retracted jaws (*f*, *f'*); and the dorsal ganglia (*t*, *t*, fig. 28) with the ventral ganglia (*p*, *p*, fig. 27) by means of two short, thick nervous commissures which send off the nerves (*u*, *u*, figs. 27 and 28). These two nerves (*u*, *u*) terminate in the ganglia (*r*, *r*, figs. 27 and 28).

These ganglia and nerves of the proboscis lie on its walls, underneath the muscles.

The sense-organs are the four eyes, the four antennæ, the tentacular cirri, and the dorsal and ventral cirri of the feet; also the long slender cirri of the posterior extremity. The antennæ and cirri are organs of touch.

Organs of Reproduction.

The sexes are separate, and the genital organs appear as simple glandular bodies, ovaries or spermaries, which project from the ventral surface into the cavity of the body, between the transverse muscular partitions. At the sexual period they are filled with eggs or spermatic particles, although at other times they can scarcely be seen. Neither the spermaries nor the ovaries have special excretory ducts, which open upon the surface of the body. The sperm and ova are discharged into the cavity of the body, which at this period is often filled with them. At the base of the lower ramus of each foot (fig. 22, *g*) is a glandular body, called the segmental organs. Some of these are normally kidneys, as urea has been found in them, but some are usually modified to act as oviducts, having a trumpet-shaped mouth opening into the body cavity and communicating with the exterior. I found the segmental organs all along the body beyond the region of the proboscis, but was unable to find the trumpet-shaped tubes. These are probably situated in the posterior segments, as Professor Verrill has seen the male worms discharging their milt from that portion of the body. The fecundation takes place in the water.

EXPLANATION OF PLATES.

PLATE XLII.

Figure 1.—*Nereis virens*, female; dorsal view of the anterior portion of the body; *a*, head, with four eyes; *b, b*, antennæ; *c, c*, palpi; *c', c'*, lobes of palpi; *d*, buccal ring; *ee, ee*, longer dorsal pair of tentacular cirri; *ee', ee'*, shorter dorsal pair of tentacular cirri; *e, e*, longer ventral pair of tentacular cirri; *e', e'*, shorter ventral pair of tentacular cirri; *h*, lateral appendages; *g*, abdominal rings.

Fig. 2 and 2*a*.—Two forms of setæ; *a*, shaft; *b*, blade.

Fig. 3.—First lateral appendage of female, posterior view; lettering the same as in fig. 8.

Fig. 4.—Second lateral appendage of female, posterior view; lettering the same as in fig. 8.

Fig. 5.—Third lateral appendage of female, posterior view; lettering the same as in fig. 8.

Fig. 6.—Fourth lateral appendage of female, posterior view; lettering the same as in fig. 8.

Fig. 7.—Fifth lateral appendage of female, posterior view; lettering the same as in fig. 8.

- Fig. 8.—Forty-fifth lateral appendage of female, posterior view; A, upper ramus; B, lower ramus; *a*, dorsal cirrus; *b*, superior ligula of upper ramus; *c*, anterior setigerous lobe; *k*, posterior setigerous lobe; *d*, inferior ligula of upper ramus; *e*, *e'*, divisions of anterior setigerous lobe of lower ramus; *f*, posterior setigerous lobe; *g*, inferior ligula; *h*, ventral cirrus; *i*, shoulder of upper ramus; *s*, *s'*, setæ; *y*, *y'*, aciculæ.
- Fig. 9.—One hundred and ninth lateral appendage of female, posterior view; lettering the same as in fig. 8.
- Fig. 10.—First lateral appendage of male, posterior view; lettering the same as in fig. 8.
- Fig. 11.—Fifth lateral appendage of male, posterior view; lettering the same as in fig. 8.
- Fig. 12.—Forty-fifth lateral appendage of male, posterior view; *c*, extra division of anterior setigerous lobe; *x*, shoulder, peculiar to the male, on the dorsal edge of the lower ramus; otherwise the lettering is the same as in fig. 8.
- Fig. 13.—One hundred and ninth lateral appendage of male, posterior view; lettering the same as in fig. 8.

PLATE XLIII.

- Fig. 15.—*Nereis virens*; ventral view of the head and mouth, the proboscis withdrawn; *a*, head; *b*, *b*, antennæ; *c*, *c*, palpi; *c'*, *c'*, lobes of palpi; *d*, buccal ring; *ee'*, *ee'*, shorter dorsal pair of tentacular cirri; *e*, *e*, longer ventral pair of tentacular cirri; *e'*, *e'*, shorter ventral pair of tentacular cirri; *m*, mouth; *g*, abdominal rings; *h*, *h*, lateral appendages.
- Fig. 16.—*Nereis virens*; the walls of the body are cut through longitudinally on the dorsal side, so as to show the perivisceral cavity with the alimentary canal; *m*, mouth; *n*, muscles of mouth; *x*, pharyngeal region of the proboscis; *c*, muscular partitions of proboscis; *b*, perivisceral cavity; *p*, dental region of proboscis; *o*, oesophageal region of proboscis; *j*, salivary glands; *r*, intestine proper; *a*, muscular partitions.
- Fig. 17.—Head of *Nereis virens*, with the proboscis protruded, dorsal view; *a*, head; *b*, *b*, antennæ; *c*, *c*, palpi; *c'*, *c'*, lobes of palpi; *d*, buccal ring; *ee*, longer dorsal pair of tentacular cirri; *ee'*, *ee'*, shorter dorsal pair of tentacular cirri; *e*, *e*, longer ventral pair of tentacular cirri; *e'*, *e'*, shorter ventral pair of tentacular cirri; *f*, *f*, jaws; *o*, anterior median area of dorsal side; *l*, *l'*, left and right anterior sub-median areas of dorsal side; *n*, *n'*, anterior lateral areas; *t*, posterior median area of dorsal side; *s*, *s'*, left and right posterior sub-median areas of dorsal side; *r*, posterior lateral and ventral areas.
- Fig. 18.—Protruded proboscis, ventral side; *f*, *f*, jaws; *n*, *n'*, anterior lateral areas; *y*, anterior median area of ventral side; *x*, *x'*, left and right anterior sub-median areas of ventral side; *r*, posterior lateral and ventral areas.
- Fig. 19.—Jaw of *Nereis virens*, much enlarged.
- Figs. 27 and 28.—Nervous system of *Nereis virens*; *h*, abdominal ganglia; *n*, *n*, nerves to muscular partitions; *o*, *o*, nerves passing through partition to preceding segment; *m*, *m*, nervous trunks to feet; *k*, *k*, ganglia sending off a cutaneous branch and a branch (*i*) supplying nerves to the feet; *d*, *d*, connectives; *d'*, *d'*, accessory connectives; *g*, *g*, nerves communicating with the ganglia of the proboscis; *e*, *e*, ganglia sending branches to the internal or ventral tentacular cirri; *e'*, *e'*, ganglia sending

branches to the external or dorsal tentacular cirri; *a*, head-ganglion with four eyes; *b, b*, nerves to antennæ; *c, c*, nerves to palpi; *w, w*, series of ganglia and nerves on the ventral side of proboscis; *f, f*, jaws, ventral side; *y, y*, the series of ganglia and nerves on the dorsal side of the proboscis; *f, f*, jaws, dorsal side.

In figure 27 the proboscis has been revolved about a line passing through *g, g*, so that the ventral side is uppermost. The head and abdominal ganglia are in their natural position. The position of figure 28 is reversed so as to show the relations of the dorsal ganglia *y, y*, to the ventral ganglia *w, w* (fig. 27).

PLATE XLIV.

Fig. 20.—Circulation of blood in *Nereis virens*, and also the alimentary canal in its natural position; *m*, mouth; *x*, pharyngeal region of proboscis; *p*, dental region of proboscis; *o*, œsophageal region of proboscis; *r*, intestine, covered with a vascular network, which is connected in each segment with the large dorsal and ventral vessels by short branches; *j*, salivary glands; *a*, large dorsal vessel; *b*, large ventral vessel; *c, c', c''*, peripheral vessels; *d*, branches from the dorsal side of the feet; *e*, branch to œsophagus; *f, f''*, branches to the ventral side of the feet; *s*, lateral branch, supplying the vascular network (*g*) and first three lateral appendages; *k*, vessel from network (*g*) to base of tentacular cirri; *l*, branch from the vascular network (*u*) on the proboscis, to the large ventral vessel (*b*); *v*, branch from the large dorsal vessel (*a*), at base of head, to the network (*u*) on the proboscis.

Fig. 21.—Diagram, showing the disposition of the large ventral vessel, and its branches on both sides, in the anterior portion of the body; *l*, continuation of the large ventral vessel along the median ventral line beneath the muscles of the proboscis; *n*, vascular ring surrounding the pharyngeal region of proboscis; otherwise the lettering is the same as in fig. 20. The arrows indicate the direction in which the blood flows.

Fig. 22.—Diagram to show the circulation of the blood, and also the relative position of the parts, in one segment of the body; *i*, intestine; *p*, perivisceral cavity; *h*, crypt from which aciculæ grow; *l*, muscles of crypt, which are attached to the base of the foot; *g, g*, segmental organs; *k*, walls of body; *n*, a ganglion of the abdominal chain; *a*, large dorsal vessel; *b*, large ventral vessel; *c*, peripheral vessel; *f*, branch to ventral side of foot; *d*, branch from dorsal side of foot.

Fig. 23.—Circulation of the blood in the last posterior segment; *a*, large dorsal vessel; *b*, large ventral vessel; *c*, vascular ring, with no branches.

Fig. 24.—Lateral view of the circulation in one segment; *d*, branch from the dorsal side of foot; *f*, branch to the ventral side of foot in the adjacent posterior segment.

In the last two figures arrows indicate the direction in which the blood flows.

Fig. 25.—Dorsal view of two segments showing the vascular network in the lateral appendages and beneath the skin of the back; lettering the same as in fig. 22; the large dorsal vessel, *a*, and the peripheral vessels *c, c*, show through the translucent walls of the body.

Fig. 26.—Segment showing the vascular network beneath the skin of the ventral side and in the lateral appendages; lettering as in fig. 22.

VII. MEDIAN AND PAIRED FINS, A CONTRIBUTION TO THE HISTORY OF VERTEBRATE LIMBS. BY JAMES K. THACHER.

Median Fins in Amphioxus.

THE quadrate markings seen at the base of the median fin in *Amphioxus* extend on the dorsal side from one extremity of the animal to the other, or nearly so, and on the ventral side from the porus abdominalis aborad toward the extremity of the tail. They are largest and most distinct in the middle of the body, and become smaller and less clearly marked (as seen from the outside), toward the head and tail until they seem to fade out entirely as they closely approach these extremities.

As Stieda* has shown these are but the external marks of a series of cavities, containing what is described as "a transparent, wholly structureless mass, resembling a coagulation."

Thus the relation of these bodies to the somewhat similarly placed primordial fin rays, or "interneural spines," of the Craniote fishes is not so simple and direct as indicated in the mistaken representations of Rathke† and of Müller.‡ Still they occupy a position similar to that of the primordial fin rays of other fishes, and the fact that they do not agree with the segmentation of the lateral muscles, seems to have some pertinency here, and to this alone I wish to call attention.

In the middle of the back there are about five of these bodies to a single segment, and on the ventral side just aborad of the abdominal pore there are about four to each. As we shall see hereafter, the structures of the median line (genuine fin-rays except in *Amphioxus*) exhibit quite generally throughout the fishes, a total disregard of the segmentation of the lateral muscles, and are more numerous than those segments.

* Studien über den *Amphioxus lanceolatus* von Dr. Ludwig Stieda, Mem. de l'Acad. Imp. des Sciences de St. Petersburg, VII^e Serie, Tome xix. No. 7.

† Rathke, Bemerkungen über den Bau des *Amphioxus lanceolatus*. Königsberg, 1841.

‡ Johannes Müller, Ueber den Bau und die Lebenserscheinungen des *Branchiostoma tubricum*, Abhandl. der Berliner Academie, 1842.

Median Fins in Myxine.

In *Myxine glutinosa* the median fin extends but a short distance forward. In a specimen 24 centimeters long, from the Bay of Fundy, the fin reaches 4 cm. from the extremity of the tail on the dorsal side, and 2.5 on the ventral.

The fin-rays, now unquestionable homologues of the primordial fin-rays of Gnathostomes, though not yet having assumed the histological structure of true cartilage, support the thin fold of skin which forms the fin. They are simple tapering rods, extending distally to the edge of the fin, and proximally scarcely dipping below the general body contours.

The only deviation from simple rods which I have been able to find is the dichotomous splitting of some of the rods where the fin rounds the extremity of the tail.

The numerical relation between these rays and the corresponding muscular segments is as three to one on the dorsal side, and as two and a half to one on the ventral.

I have been unable to detect any muscular fibers in the composition of the fin.

Median Fins in Petromyzon.

Here the median fins are much better developed. In a specimen (*Petromyzon marinus*, from the Connecticut River), 77 cm. long, the caudal fin extends forward along the dorsal side 7.5 cm., sloping downward nearly to the body, then the second dorsal rises abruptly and runs orad 16 cm., where it reaches by a gentle slope the general outline of the body. There follows a finless space 3.5 cm. in length which is succeeded by the first dorsal, whose extent is 9 cm., being therefore shorter as it is lower than the second dorsal. The anus is opposite the orad part of the second dorsal.

The fins, therefore, take up almost the whole of the hinder half of the mid-dorsal line. In *Myxine* only one-sixth was thus occupied. On the ventral side we have only the caudal, extending about as far here as it does above.

These fins are supported by a series of chondroid rays, lying quite close to one another in the median plane. They are straight and slope aborad from the fatty-fibrous ridge-pole of the myelonal canal (Pl. XLIX, fig. 1, a,) to the very edge of the fin. They are found in all the fins. Their form is represented in Pl. XLIX, fig. 1, where only one ray is drawn complete. As shown, it bifurcates twice and thus ends distally in four fine branches. This figure is from the central part of

the large second dorsal, and shows one of the longest rays. Where they are shorter we may have only one bifurcation, or one of the two primary branches, that toward the longer rays, may again divide, while the other toward the shorter remains unclift. Farther forward at the beginning of the second dorsal, where the rays are still shorter, they do not divide at all, but end somewhat bluntly though compressed from side to side.

Each ray is largest in the middle and here lies quite close to its adjacent rays; below they grow more slender, and therefore are somewhat spaced, but expand somewhat to a foot resting on the myelonal canal.

I have seen no cases of conerescence between adjacent rays. With the exception of the variation in the branching and length in different parts of the fin, before alluded to, the rays are all similar and parallel one to another.

This branching is plainly a true dichotomy and not the product of conerescence, as is evidenced by the total absence of anything else resembling conerescence, by the similarity in size between two adjacent differently branched rays, and by the regularity of the branching.

On each side of the row of skeletal elements are muscular bundles of a somewhat blacker color than the two great masses of lateral muscles. The muscles of the median fin wedge themselves into the angle between the lateral muscles of the two sides along the median line. They are sharply distinguished from these. The fibers of the lateral muscles run longitudinally, while those of the fin muscles are parallel to the primordial median fin-rays. There is absolutely no continuity between the two in any part. Moreover the bundles of the fin-muscles show no relations to the segments of the lateral muscles. A cross section, Pl. XLIX, fig. 3, shows the relation between the fin-muscle bundles and fin-rays.

The numerical relation between the fin-rays and the segments of the lateral muscle is shown in Pl. XLIX, fig. 2, where we have a little less than four of the former to one of the latter.

The relation between the fin-rays and the neural arches is shown in figure 1, where we have 35 rods and 23 arches. These neural arches rise from the sheath of the notochord, to stiffen the fibrous sides of the myelonal canal and to apply themselves to its fatty-fibrous ridge-pole. The fin-rays abut on the same ridge-pole in the mid-dorsal line, but they are in no way connected with the neural arches. I have met with no cases even of conerescence between the two.

Figure 4, Pl. XLIX, shows the relation between the neural arches and the segments of the lateral muscles. There are two arches to one segment. In figure 1, then, there must have been $11\frac{1}{2}$ segments, which gives us a very little more than *three* as the ratio between the fin-rays and the segments. From figure 2 we obtained a little under *four*. Both results are necessarily correct. There is considerable variation in the relation between the fin-rays and the muscular segments and the neural arches. This is exhibited in the following table of observations on a single specimen.

	Fin-rays.	Arches.	Ratio of fin-rays to arches.	Ratio of fin-rays to segments.
In the orad half of 1D.	19	8	2.4	4.8
In the aborad half of 1D.	20	8	2.5	5.
In the orad part of 2D.	20	11	1.8	3.6
In the next 35 rays of 2D.	35	23	1.5	3.
In the next 27 rays of 2D.	27	17	1.6	3.2
In the next 12 rays of 2D.	12	7	1.7	3.4

The Lampreys then have advanced beyond the Myxines toward the Gnathostomes as regards the structure of the median fins, in the greater development and efficiency of those organs, and in the addition of special fin-muscles which seem to be wanting in the lower group. And while in Myxine the original independence of the median fin skeleton and the axial skeleton is shown by the existence of the median fin-rays before any neural arches have appeared, and by their want of agreement with the muscular segments to which the axial skeleton will conform when it does arise, in *Petromyzon* we have the same independence more strikingly reaffirmed by the simultaneous existence of neural arches and median fin-rays, and their perfect indifference to one another.

*Median Fins of Elasmobranchii.**

We shall here consider the skeleton of the dorsal and anal fins alone. That of the caudal fin has on the ventral side undergone peculiar modifications by the union of fin-rays with hæmal spines.

* The term Elasmobranchii includes the Chimeroids with the Sharks and Rays. I have had no opportunity of examining the fin skeletons of the former. But *Chimæra* seems to me to be but a divergent form of Sharks and to have its nearest living relative in *Cestracion*. They are quite specialized forms, *Cestracion* the less so.

I will state here that the Elasmobranchs described came from Wood's Hole, Mass. The names used are those given by Dr. Gill in the U. S. Fish Comm. Rep. for 1871-72.

In general the structure of the median fin resembles what we have seen in *Petromyzon*, but there are important differences. The rays are of hyaline cartilage and they do not usually reach down to the ridge-pole of the myelonal canal. In *Eulamia*, for example, in the smaller of the specimens figured, the rays approach within a centimeter and a half at the oral extremity of the first dorsal fin, but are three and a half centimeters distant at the other extremity. In *Squalus* (*Acanthias*) they come closer.

This ridge-pole consists of a cord of rather peculiar white longitudinal fibers, constituting now a "*ligamentum longitudinale*." It appears to me undoubtedly homologous with the fatty-fibrous body in *Petromyzon*. The cartilaginous arches unite under, and do not extend around over it, though they clasp it somewhat.

The rays are segmented, usually twice. Dichotomy is rare if not altogether absent. Concrecence of adjacent rays is by no means uncommon. The reduction of rays in size is exhibited in all degrees.

Calcification presents itself in a thin superficial layer on each side of the somewhat flattened ray, but it fails on the edges, i. e. as we come close to the median plane.

The muscles of the fin, as in *Petromyzon*, are in total independence of the large masses of segmented lateral muscles, but they are in more definite relation with the skeleton of the fin. This is accurately represented in Pl. LIX, fig. 66, though that is a section of a pectoral and not of a median fin. We see that each ray has on each side a special muscle, separated from its fellows by the fibrous sheet which runs from between the rays to the integument. Each little muscle developes in its median line a flat tendon, which, parallel to the surface of the fin, inserts itself in the fascia covering the extremities of the fin-rays and the proximal ends of the well known horny fibers, which here supplement the primordial skeleton, as the secondary fin-rays of Ganoids and Teleosts do.

The relation between the number of fin-rays and that of the vertebræ opposite to them is similar to what we have seen in the lower forms. In the Nictitantes, for example, there are on the average about 2.5 rays to one vertebra. But there is considerable variation, even in individuals of the same species. The extreme numbers, so far as I have observed, are 3.5 in an anal of *Sphyrna*, and 2 in a first dorsal of *Eulamia*.

We turn now to the more minute examination of several species.

First Dorsal of Mustelus canis, Pl. XLIX and L, figs. 5-10.

In Pl. XLIX, fig. 5, we have 24 separate rods, unless 3 be the terminal piece of 2; but its conformation seems to testify to its independent but reduced character. The number here then is 24 or 23.

In Pl. XLIX, fig. 6, we have again the same alternative, without quite so strong a case for 24, but still quite strong.

In Pl. L, fig. 7, we have the choice between 24 and 25, but in favor of the latter.

In Pl. L, fig. 8, we have 23 or 24, but the former has the greater probability.

Pl. L, fig. 9 gives us 22, 23 or 24, 23 being more probable.

Pl. L, fig. 10 exhibits 23 or 24, dependent on the view taken of rays 5, 6, and 7. The probability seems in favor of 24.

I think we may sum up then with regard to the number of rays constituting the first dorsal of *Mustelus canis* thus: it has usually 24 rays but this may vary to 23 or 25.

Nearly all the rods are segmented twice. The distal line of segmentations fails in the one or two orad; and the proximal, in the two or three aborad ones. Additional segmentations are very rare. What might be reckoned as such are seen in fig. 7, ray 4; fig. 6, rays 5 and 6. This makes an average of $\cdot 0 +$.*

The union of adjacent rays is rare. I estimate it at $\cdot 04$ of the total possible concrescence.†

Shortening or reduction in size is likewise rare. We have first those questionable cases of which fig. 5, rod 3 seems the least questionable; and then plainer but less extreme instances in fig. 7, ray 5; fig. 9, ray 18; fig. 10, ray 7, then we have the usual shortening of the rays at the extremities of the fin. When these aborad rays shorten up, those next in front of them have a remarkable tendency to grow up under them. This is well shown in fig. 9.

Moreover when in the aborad rays the proximal joint becomes very short, it is sometimes divided into two lateral halves. This is the case for example in rod 22 of fig. 10.

Occasionally we have a minute piece or pieces of cartilage forming a tip to a ray. It cannot act as an extra joint, by giving increased flexibility to the ray. And it seems doubtful whether the origin of the two is to be referred to the same causes. Yet intermediate forms occur so as to raise the question whether they are to be referred to one or the other category. These tips seem to be exhibited in fig. 7, ray 4; fig. 8, rays 2 and 3; fig. 9, ray 3. This gives $\frac{4}{6 \times 24} = \cdot 0 +$.

I now find the ratio of the proximal piece of the middle ray of each fin to the middle piece of the same. The average of these ratios is $\cdot 6$. The method gives $\cdot 3$ as the ratio between the terminal and middle joints.

We have then for the first dorsal of *Mustelus canis*:

Number of rays 24. Extra segmentations $\cdot 0 +$.

Concrescence $\cdot 04$. Betipping $\cdot 0 +$. Ratio of proximal to middle piece of middle ray $\cdot 6$. Ratio of distal to middle piece $\cdot 3$.

* The decimal is obtained by dividing the number of additional segments by the number of rays.

† The amount of concrescence between two adjacent rays is the ratio between the length of the union and the whole distance through which they are adjacent and might have united. The sum of these fractions divided by the number of rays less the number of fins, gives the estimate of the concrescence.

Second Dorsal of Mustelus canis, Pl. L and LI, figs. 11–15.

Fig. 11 has 24 rays, fig. 12 has 22, fig. 13 has 23, fig. 14 has 24, fig. 15 has 24. Thus of the five examined three have 24, one 23 and one 22 rays. It seems probable that a wider examination would give us forms having 25 or more rarely 26. Thus we have for the second dorsal 24 rays with some slight variation.

Extra segmentations appear only in fig. 11, ray 5; and fig. 14, ray 2. This gives $\cdot 0+$.

I estimate the concrescence at $\cdot 06$. The concrescence is mostly confined to the proximal row, and is more frequent at the two ends than in the middle of the series.

Betipping is seen only in fig. 14, ray 23. This gives $\cdot 0+$.

Ratio of proximal piece of middle ray to middle piece $\cdot 6$.

Ratio of distal piece of middle ray to middle piece $\cdot 4$.

The downward prolongation of the proximal parts of one or two of the oral rays is noticeable, being quite pronounced in all the cases except that represented in fig. 11.

The reduction of the rays is rare, but shown to an extreme extent in fig. 11, ray 1; and fig. 15, ray 1.

Anal of Mustelus canis, Pl. LI and LII, figs. 16–19.

Figs. 16 and 17 have each 18 rays. Fig. 18 has 17 or 18 according as the last ray is double or not. Fig. 19 has 18 or 19 under the same conditions. The great width of the last ray in the last two cases makes the larger number probable. Thus we have 18 as the normal number, with probably slight variations.

Extra segmentations are seen in fig. 17, ray 5; and fig. 19, ray 3. This gives $\cdot 0+$.

The concrescence I estimate at $\cdot 09$. Betipping occurs in fig. 16, ray 3. This gives $\cdot 0+$.

Ratio of proximal piece of middle ray to middle piece is $\cdot 7$.

Ratio of distal piece of middle ray to middle piece is $\cdot 6$.

First Dorsal of Galeocerdo tigrinus, Pl. LII, fig. 20.

In this sole specimen there are 25 rays.

Extra segmentation occurs in 10, 20, 21, 22, 23, which gives $\cdot 2$. It should be noticed that this extra segmentation is in each case here a doubling of the proximal line of segmentations.

Concrescence is estimated at $\cdot 06$. Betipping is seen in 18, 19 and 25. This gives $\cdot 1$.

Ratio of proximal piece of middle ray to middle piece $1\cdot 1$.

Ratio of distal piece of middle ray to middle piece $\cdot 6$.

Shortening is seen in 6 and 22. In the latter the proximal piece is excluded from the edge of the fin by *a*, the proximal piece of 23, and by the proximal piece of 21. The piece *a* consists of two lateral halves.

Second Dorsal of Galeocerdo tigrinus, Pl. LII, fig. 21.

Number of rays 13. Extra segmentation in 3, giving $\cdot 1$.

Concrescence is estimated at $\cdot 01$. Betipping, in 5, 8 and 13, gives $\cdot 2$.

Ratio of proximal piece of middle ray to middle piece $1\cdot 3$.

Ratio of distal piece of middle ray to middle piece $\cdot 6$.

Anal of Galeocerdo tigrinus, Pl. LII, fig. 22.

Number of rays 12. Extra segmentation in 6, 7, 8 and 10, gives $\cdot 3$.

Concrescence is estimated at $\cdot 05$. Betipping in 4 gives $\cdot 1$.

Ratio of proximal piece of middle ray to middle piece $1\cdot 1$.

Ratio of distal piece of middle ray to middle piece $\cdot 7$.

First Dorsal of Eulamia Milberti, Pl. LII and LIII, figs. 23 and 24.

Number of rays 28 or 29.

As indicated by the numbering of the rays, I take number 21 in each figure to be a single ray, which has widened at the top, and been segmented in the way figured.

Extra segmentations in fig. 23, rays 20 and 21 (2 extra segmentations in the latter) in fig. 24, rays 4 and 21. This gives $\cdot 1$.

Concrescence is estimated at $\cdot 09$. Betipping is absent.

Ratio of proximal to middle piece $\cdot 9$.

Ratio of distal to middle piece $\cdot 9$.

Second Dorsal of Eulamia Milberti, Pl. LIII, figs. 25 and 26.

The number of rays differs remarkably in the two specimens, being 12 in the one and 16 in the other. It must, however, be remembered that the second dorsal has become very small and of very little physiological importance. Organs which have thus become functionless are peculiarly prone to vary. They thus secure more easily some other and new function. We will take the average number 14 as the normal one for the rays of this fin.

Extra segmentation occurs in fig. 25, ray 6 (twice), and in fig. 26, ray 3. This gives $\cdot 1$.

Concrescence is estimated at $\cdot 09$.

Betipping is absent.

Ratio of proximal to middle piece of middle ray $\cdot 8$.

Ratio of distal to middle piece of middle ray $\cdot 5$.

Anal of Eulamia Milberti, Pl. LIII, figs. 27 and 28.

Number of rays 17 or 18.

Extra segmentations in fig. 27, rays 7, 10 and 12 (twice in the latter); in fig. 28, twice in 12, once in 16, give $\cdot 2$.

This implies a certain interpretation of the ambiguous rays 11 and 12 in figure 27. In fig. 28 we seem to have a plain case. Here the ray 12 is broadened at the top, and its distal piece divided by two intersecting cuts into four pieces. Ray 12 in fig. 27 is explained in the same way. Ray 11 is a little shortened, and excluded from the edge by the tips of 10 and of 12. This appears to me the most probable view of the case.

Concrescence is estimated at $\cdot 12$. Betipping absent.

Ratio of proximal to middle piece of middle ray $\cdot 7$.

Ratio of distal to middle piece of middle ray $\cdot 4$.

First Dorsal of Sphyrna zygaena, Pl. LIII and LIV, figs. 29 and 30.

Number of rays 33 and 34.

I regard the three pieces at the extremity of 28 as belonging to that ray. It has been widened and divided like the instances in *Eulamia*.

Extra segmentations, fig. 29, rays 2 and 3, twice; rays 4 and 5; ray 28, twice; fig. 30, ray 28, twice, give $\cdot 2$.

Concrescence is estimated at $\cdot 07$. Betipping absent.

Ratio of proximal piece of middle ray to middle piece $\cdot 7$.

Ratio of distal to middle piece of middle ray $2\cdot 4$.

In fig. 29 the proximal line of segmentation fails in rays 9–17, except in the joined rays 11 and 12 where it is present. In fig. 30 it fails in rays 8–16.

Second Dorsal of Sphyrna zygaena, Pl. LIV, fig. 31.

Number of rays 14. Extra segmentations amount to 1.0.

Concrescence is estimated at .07.

Betipping is absent.

Ratio* of proximal to middle piece of middle ray .3.

Ratio of distal to middle piece of middle ray .3.

The last ray, both in the second dorsal and the anal, is large and round.

Anal of Sphyrna zygaena, Pl. LIV, fig. 32.

Number of rays, 27.

Extra segmentations 8 (1), 9 (1), 10 (1), 11 (1), 12 (1), 13 (1), 14 (1), 15 (1), 16 (1), 17 (1), 18 (2), 19 (1), 20 (1). This gives .5.

Concrescence is estimated at .03. Betipping none.

Ratio of proximal to middle piece of middle ray .4.

Ratio of distal to middle piece of middle ray .3.

First Dorsal of Eugomphodus litoralis, Pl. LIV, and LV, figs. 33-39.

Specimens figured in figs. 33, 34 and 36 have plainly 16 rays. Those in figs. 37 and 39 have plainly 17. Those in figs. 35 and 38 have 16 separate rays, but the last is quite broad. Where we have plainly 17 rays, figs. 37 and 39, the last two rays have united with the exception of their distal joints. We may fairly conclude that figs. 35 and 36 present a more complete concrescence of those rays. We have then as the number of rays 16 or 17, the former in three cases, the latter in four.

Extra segmentation is estimated at .7. Concrescence is estimated to be .05.

Betipping reaches .4, each separate piece being counted. These small nodules of cartilage sometimes seem very evidently to be a continuation of a ray upon the following ray. But not infrequently they seem to be scattered rather irregularly along the edge of the fin. It will be noticed that they are most frequent in the oral part of the fin, though not on the first two or three rays. It is very probable that the estimate of their frequency should be higher than given, for they are easily lost in the preparation of the specimen.

Ratio of proximal to middle piece of middle ray .5. Ratio of distal to middle piece of middle ray .4.

Second Dorsal of Eugomphodus litoralis, Pl. LV and LVI, figs. 40-46.

In fig. 42 we have 16; in 45, 17; in 40 and 46 we have 18 rays. These are all plain cases. Fig. 43 exhibits 17 rays, but raises a suspicion of 18 by the breadth of the last ray. Fig. 41 gives 16 or 17, probably the latter. Fig. 44 leaves us in doubt between 14, 15 and 16, with, as it seems to me 15, the most probable. We may take 17 as the normal number. As far as the evidence here goes the second dorsal is more liable to vary than the first. We see that in each the greater the number of rays, the greater is the amount of concrescence.

Extra segmentation amounts to .6.

Concrescence amounts to .10. Betipping amounts to .4.

Ratio of proximal to middle piece of middle ray is .3.

Ratio of distal to middle piece of middle ray is .3.

* Where a segmentation is double the point half way between the joints is taken as the limit between the middle and extreme piece. Where it is triple the middle segmentation is taken.

Anal of Eugomphodus litoralis, Pl. LV1 and LVII, figs. 47-50.

The number of rays is 21 in figs. 47 and 49. In 48 we have 22, and in 50, 20 rays. We may take then 21 as the normal number.

Extra segmentations amount to .4.

Concrescence is estimated at .10. Betipping amounts to .3.

Ratio of proximal to middle piece of middle ray 4.

Ratio of distal to middle piece of middle ray .5.

We may sum up the results of this investigation of the resemblances and differences of the forms so far examined in the following table.

	No. of rays.	Extra segmentations.	Concrescence.	Betipping.	Ratio of prox. to mid. piece of mid. ray.	Ratio of distal to mid. piece of mid. ray.
First Dorsal.	<i>Eugomphodus</i> .. 17	.7	.05	.4	.5	.4
	<i>Mustelus</i> 24	.0	.04	.0	.6	.3
	<i>Galeocerdo</i> 25	.2	.06	.1	1.1	.6
	<i>Eulamia</i> 29	.1	.09	.0	.9	.9
	<i>Sphyrna</i> 34	.2	.07	.0	.7	2.4
Second Dorsal.	<i>Eugomphodus</i> .. 17	.6	.10	.4	.3	.3
	<i>Mustelus</i> 24	.0	.06	.0	.6	.4
	<i>Galeocerdo</i> 13	.1	.01	.2	1.3	.6
	<i>Eulamia</i> 14	.1	.09	.0	.8	.5
	<i>Sphyrna</i> 14	1.0	.07	.0	.3	.3
Anal.	<i>Eugomphodus</i> .. 21	.4	.10	.3	.4	.5
	<i>Mustelus</i> 18	.0	.09	.0	.7	.6
	<i>Galeocerdo</i> 12	.3	.05	.1	1.1	.7
	<i>Eulamia</i> 18	.2	.12	.0	.7	.4
	<i>Sphyrna</i> 27	.5	.03	.0	.4	.3

The changes which are presented consist chiefly in concrescence. This takes place in various ways. The proximal portions more frequently unite than the distal, but we may have concrescence of the distal points while the proximal portions are separate. The reduction of rays is exhibited in all degrees from the slightest shortening to the extreme degree shown in Pl. LVII, fig. 50, ray 9.

It is perhaps noteworthy that the changes which would have sufficed to differentiate the fin of one species from another, if they had continuously advanced for a few generations, are changes which in no slight degree are now taking place between parents and children. I mean changes of number, segmentation, concrescence, and relative lengths of parts of rays.

I exhibit some figures of the dorsal fins of *Squalus*, Pl. LVII, figs. 51-56, and Pl. LVIII, fig. 57, *Raia*, Pl. LVIII, figs. 58, 59, and *Myliobatis*, Pl. LVIII, fig. 60. It is perfectly certain that they have been derived from a series of parallel rays, and that the principal process of change has been in the way of concrescence. They tell their own story as far as it can be told without the investigation of other closely related forms.

The last ray in the first dorsal of *Raia*, Pl. LVIII, fig. 58, is remarkable, and it is represented in the second dorsal, fig. 59, by a series of separate nodules.

I also add figures of the dorsal, Pl. LVIII, fig. 61, and anal, Pl. LIX, fig. 62, of *Acipenser*. These are very similar to the simpler shark dorsals. It will be noticed that we have the predominant division into three pieces, but the terminal piece is very short.

Conclusions regarding Median Fins.

The primordial median fin-rays in whatever form they occur are derivatives from a series of simple parallel chondroid rods, which grew up in the median fold in total independence of the cartilaginous arches above and below the notochord. These earliest representatives of these parts were from two to four times as numerous as the vertebrae opposite them. In the Gnathostomi true hyaline cartilage replaced the lower form of tissue seen in *Myxine* and *Petromyzon*.

Segmentation and conerescence, as well as reduction in size, were common changes in the Gnathostomes, and here a division into three parts is the usual though not invariable rule.

Hence it is seen that Gegenbaur's* statement that, in their simplest form, the primordial fin-rays are mere prolongations of the neural spines is incorrect. It has been demonstrated that this was not the *earliest* form. The Dipnoans, however, seem to offer an example where the primordial median fin-rays are mere prolongations of the neural spines. They demand a moment's consideration.

True neural spines are first found in the Ganoids. They are absent in the Elasmobranchs and Agnathostomi. Pl. LIX, fig. 63 represents the projection of a section of a vertebral segment of *Acipenser* cut through the middle line of the arch and neural spine. As the latter slopes backward, it is considerably foreshortened in the figure.

Now the cartilaginous arches springing from the sheath of the notochord pass upwards to lay themselves on each side of the fibrous cord *b*, and here they spread inward to meet one another on the ventral side of *b*, and also prolong themselves above to almost or quite meet, and then they are followed by the dorsal spine *a*, from which they are separated by a segmentation.

Now the cord *b*, which is the same as the *ligamentum longitudi-*

* Grundriss der Vergleichenden Anatomie, 1873, p. 488. Gegenbaur's assertion in the same place that they usually correspond in number to the vertebrae opposite, is very strange. We have already seen that they do not do this in the earlier and more significant forms; and the statement of Gegenbaur would decidedly misrepresent what we find, for example, in the figures of fish skeletons in Agassiz's *Poissons Fossiles*.

nale in *Ceratodus*,* is also, without the slightest doubt, homologous with the chord which lies entirely above the neural arch in the Elasmobranchs. The peculiar fibrous character is almost exactly the same in each. We have seen that this cord in the sharks is in all probability homologous with the fibrous fatty ridge-pole of the neural canal in *Petromyzon*. Thus while in the latter the arches of one side and the other are entirely separate, in the sharks they have spread beneath the ligament so as to meet, and in the higher Gnathostomes they have also joined above it, or nearly joined, for the origin of *a* is still to be discussed.

There are two possibilities with regard to the neural spine *a*. Either it is formed by the union of a median fin-ray with the neural arches, the ray thus constituting the keystone of the arch, or else by the union of the neural rods from each side and their prolongation dorsad.

But the junction between *a* and *b* is quite close; the neural spines correspond in number and position with the lateral parts of the arch; while fig. 61 shows conclusively the absolute independence of neural spines and primordial median fin-rays.

The second of the two possibilities is then the true one. Thus neither are median fin-rays derived from neural spines, nor neural spines, where they occur, from primordial fin-rays.

But the cartilaginous supports of the median fold in the Dipnoans are very long and segmented. They are simply elongated neural spines and are not primordial fin-rays in any homological sense. If they were formed by the reduction in number of the primordial fin-rays and their coalescence with the neural spines it is impossible that we should not have here and there an extra one, and some evidence in the case of others of such a junction. But there is nothing of the kind, either in the descriptions of Günther in the case of *Ceratodus*,† or in those of Owen‡ and Peters§ in that of *Protopterus annectens*, or in those of Bischoff|| in that of *Lepidosiren paradoxa*. Gün-

* Günther's Description of *Ceratodus*, Phil. Trans., vol. clxi, pt. ii, Pl. XXXVIII, Figs. 3-9, é.

† Phil. Trans., vol. clxi, pt. ii, 1871. Günther, Description of *Ceratodus*. In Günther's fig. 2, Pl. XXX, the proximal joint of the 14th neural spine seems to bear two "interneurals," one orad of the other. But as no notice is taken in the text of this, which would be a very noteworthy fact, if it were fact, and as the description of these parts there given is such as would demand a notice of this exception, it is evident that it must be an inaccuracy in the figure.

‡ Trans. Linnean Soc., vol. xviii, pt. iii. Owen, Description of *Lepidosiren annectens*.

§ Müller's Archiv., 1845. Peters, Ueber einen dem *Lepidosiren annectens* verwandten, Fisch von Quellimane.

|| Ann. Sc. Nat., xiv, 1840. Bischoff, Sur le *Lepidosiren paradoxa*.

ther's denomination, then, of the ultimate and penultimate joints of the neural spines of *Ceratodus* as "interneural first" and "interneural second," is ill chosen, and rests on a mistake in homology. That great genetic group, then, consisting of Dipnoi, Amphibia and Amniota, seems to have entirely lost those primordial median fin-rays which appeared so early and are found even in *Myxine*.

Limb-skeleton of Air-breathing Vertebrata.

In 1864, Gegenbaur* set forth the splendid results of a widely extended investigation of the limb-skeleton of the air-breathing vertebrates. Herein was established the typical form of these parts for this large group, consisting of Amphibia and Amniota. Inasmuch as there is no doubt of the natural, that is the genetic, character of this group, and inasmuch as it is marked out from all other vertebrates by the development of a *fenestra ovalis* and the modification of the proximal part of the second post-oral, or hyoid, arch into a stapes in connection therewith, I venture to use the name Stapedifera in place of the circumloutory air-breathing Vertebrates. For the Stapedifera, then, the typical limb-skeleton was established; typical in the sense of the older anatomists, as that ideal form from which we could in our minds easily derive the various actual forms now living; but typical also in the newer sense, as that actual form, the limb-skeleton of the latest common ancestors of all Stapedifera, from which have been developed the corresponding parts in all living Stapedifera.

The same form belongs to both fore and hind limbs. Using the names applicable to the former, we have, as is well known, humerus, radius and ulna, radiale, intermedium and ulnare, a centrale, and then set around these, five carpalia followed by their metacarpals and phalanges. Moreover, the strong suspicion of the double nature of the centrale, as evidenced in the descriptions of *Cryptobranchus Japonicus*,† by Schmidt, Goddard, and J. Van der Hoeven, is later, 1865, confirmed by the careful observations and clear presentation of the anatomy of that animal by Hyrtl.‡ The Ichthyosaurs§ and

* Gegenbaur, Untersuchungen zur vergleichenden Anatomie der Wirbelthiere, Hft. 1, Carpus und Tarsus.

† Gegenbaur, Unters., Hft. 1, p. 57.

‡ Hyrtl, Schediasma anatomicum. 1865. Gegenbaur, Unters., Hft. 2, p. 165.

§ Gegenb., Unters., Hft. 2, p. 165, and Jena Zeitschr., Bd. v, Hft. 2, 1870. Gegenbaur, Ueber das Gliedmaassenskelet der Enaliosaurier. In this last a further modification is made in the recognition of the pisiform as the remains of a sixth row, and as being an essential part of the carpus and not merely a sesamoid bone.

Plesiosaurs give too their unambiguous evidence in the same direction.

This limb of the latest common ancestors of the Stapedifera, this typical limb of that group, has been named by Huxley, *chiropterygium*,* and the term will be found convenient.

The Chiropterygium and the Fins of Fishes.

The homology between the paired fins of fishes and the limbs of Stapedifera has long been recognized; but the special homologies of the skeletal parts of each has been the subject of much controversy. For a historical sketch of the various divergent opinions endorsed by the highest authorities I must refer to the second volume of Gegenbaur's *Untersuchungen*.

Two pairs of limbs are found throughout the great genetic group of the Gnathostomi. The chiropterygium having been established, the determination of that earlier form typical for all the Gnathostomi became a more pressing question.

To an answer to this question the investigations of Gegenbaur now begin to lead.

The first part of the second volume of the *Untersuchungen* discusses the shoulder girdle, and the result is that we are now able to trace clearly and surely the primordial shoulder girdle, the scapulo-coracoid, throughout the Vertebrata. We are no longer in doubt as to what part of the fish fin and girdle corresponds to limb and what to girdle of the Stapedifera. The results of Gegenbaur's work with which we are here concerned were confirmed by the later but independent researches of Parker.†

Development of the Archipterygium Theory.

This preliminary question having been satisfactorily answered, the derivation of the chiropterygium is next attempted.

The second half of the second volume of the *Untersuchungen* (1865) takes up the pectoral fin of fishes. Quite a number of very excellent figures of numerous Elasmobranchs, Ganoids and Teleosts are given, and the limb skeleton of *Protopterus* is discussed. The conclusions drawn are as follows:

We may take as the most generalized form of limb that of the Elasmobranchs, where its various parts are most plainly presented in

* Proc. Zool. Soc. London, 1876, pt. i, p. 56. T. H. Huxley, On *Ceratodus Forsteri*.

† Parker, Shoulder-girdle and Sternum. Ray Soc., 1868.

the rays. It is divided into three parts, Propterygium, Mesopterygium and Metapterygium. Each of these consists of a *basale*, which articulates with the shoulder girdle, and a number of rays set on its edge.

The fin of *Protopterus* is derived from this by the destruction of the pro- and mesopterygium. The metapterygium is here represented by the long articulated rod, which alone remains in *Lepidosiren paradoxa*. The row of cartilages along its sides are the metapterygial rays.

In the Ganoids, *Polypterus* alone has the three divisions represented. In this, neither metapterygium nor propterygium bears rays. These are confined to the mesopterygium, which is excluded from the articulation with the shoulder girdle.

In the other Ganoids the propterygium fails. Between the mesopterygium and metapterygium a number of rays are brought into articulation with the shoulder girdle, resembling what is seen in some of the Rays.

The Teleosts in the main resemble this second group of Ganoids.

The chiropterygium is derived from the metapterygium alone, and thus resembles the limb of *Protopterus*. The fore limb will serve as an example. The *Stammreihe* or *basale metapterygii* is presented by the humerus, radius, radiale, carpale radii, the metacarpal and phalanges of the thumb. The other bones are the rays belonging to this, and their arrangement will be best understood by looking at the *Ichthyosaurus* limb, fig. 70. In 1870,* Gegenbaur published his explanation of the limb of the Enaliosaurs. The unbroken lines in fig. 70 of *Ichthyosaurus* exhibit his view of the relations of the fin with that of fishes. This may be regarded as closing the first stage of the development of the theory in Gegenbaur's publications.

The second immediately opens. For in the next number† of the *Jena Zeitschrift* there appears an extended article on the ventral fins of Elasmobranchs. The previous view is modified as follows. The fin-skeleton of the latest common ancestors of all Gnathostomes, is represented pure and simple in the fore limb of *Protopterus amneetens*, and with only slight modification in the ventral fins of Elasmobranchs. It now has a name given to it. It is called *archipterygium*. There is a limb girdle, complete ventrad. On each side is articulated to this

* *Jenaische Zeitschr.*, Bd. v, Hft. 3. Gegenbaur, Ueber das Gliedmaassenskelet der Enaliosaurier, Feb., 1870.

† *Jen. Zeitschr.*, Bd. v, Hft. 4. Gegenbaur, Ueber der Gliedmaassen der Wirbelthiere im Allgemeinen und der Hintergliedmaassen der Selachier insbesondere, May, 1870.

the stem-row (*Stamm-reihe*), a long tapering many-jointed cartilaginous rod which bears on the outer side a series of rays. This evidently calls for no change of view regarding the Enaliosaurs or Stapedifera. But the fin-skeleton of fishes exhibits everywhere, except in *Protopterus* and *Scymnus*, a slipping off of the rays from the stem-row and their articulation with the girdle, and very commonly their articulation with one another and considerable fusion (concrecence).

Still another change awaits the primordial limb, even the named archipterygium. In 1871, Günther* published his description of *Ceratodus*. Here the stem-row has a series of rays down each side. The archipterygium is modified to accord with this in the Jena Zeitschrift published April 22, 1872,† where Gegenbaur adopts the "*Biseriale Archipterygium*" as the parent form, and attempts to show that there are some traces of the median row of rays in the pectoral fins of some Elasmobranchs. With the exception of *Ceratodus* and the questionable exception of these Elasmobranch pectorals, the biserial has been everywhere reduced to the uniserial form, and still further reduced as heretofore explained.

In the third volume of the *Untersuchungen*, dated May, 1872, a suggestion‡ is made of the possible origin of the Archipterygium and the limb-girdles. They are assimilated to the branchial arches and their diverging rays, where rays move up upon, and articulate with, the longest middle ray. It is but justice to say that the suggestion is a little vaguely and hesitatingly made.

In confirmation of Gegenbaur's views, Bunge,§ in 1874, published a further investigation of the pectoral fin of Elasmobranchs, showing a number of rays which might be regarded as median, in several species not examined by Gegenbaur. Finally, in 1876, Huxley|| took up the question, and, while he accepted the archipterygium, he modified the interpretation of a large number of the forms.

* Proc. Roy. Soc., 1871, p. 378, and more fully, with a figure of the fin-skeleton, in Ann. and Mag. of Nat. Hist., March, 1871. To these Gegenbaur refers, Jen. Zeitschr., Bd. vii, Hft. 2, p. 132, note. But a much fuller description is given by Günther, Phil. Trans., vol. clxi, pt. ii, pp. 511-572. This was published early in 1872.

† Jen. Zeitschr., Bd. vii, Hft. 2, pp. 131-141. Gegenbaur, Ueber das Archipterygium.

‡ Gegenbaur, Unters., Hft. III. Kopfskelet der Selachier, p. 181, note. 1872.

§ A. Bunge, Jena. Zeitschr., Bd. 8, Hft. 2, 1874. Ueber die Nachweisbarkeit eines biserialen Archipterygium bei Selachiern und Dipnoern. Bunge also calls attention to the fact that the fringing rays in *Protopterus unnectens* are on the median (i. e., ventral,) side of the axis, and not, as in Elasmobranchs, on the lateral (i. e., dorsal,) side.

|| T. H. Huxley, Proc. Zool. Soc. Lon. for 1876, Pt. 1. On *Ceratodus Forsteri*.

Most of the modifications introduced by Huxley, though perhaps not all, spring from a question which is independent of any theory with regard to the skeleton, archipterygium or other, but which, superior to them, must determine the application of them to the passage from the fish limb to that of Stapedifera.

If an Elasmobranch pectoral fin, for example, of *Mustelus*, be removed and laid on the corresponding hand, with the propterygial edge toward the thumb, and the metapterygial edge toward the little finger, then the ventral surface of the fin will look in the same direction as the palmar surface of the hand. But if it be turned over so that the metapterygial edge corresponds to the thumb and the propterygial to the little finger, then the dorsal surface of the fin will correspond to the palmar surface of the hand.

One or the other of these views must be taken. There is no third possible. Huxley takes the first, Gegenbaur the second. This, however is no new question and no new difference of opinion. Cuvier, following Bakker, named the two ossifications of the scapulo-coracoid which are so generally found in osseous fishes, radius and ulna. Owen simply reversed this nomenclature and Mettenheimer followed him. The question was the same as now respecting the homologies of faces and edges of fin and limb. On the one side, then, we have Bakker, Cuvier and Huxley; on the other, Owen, Mettenheimer and Gegenbaur. The weight of evidence seems to me to be in favor of the view entertained by the latter group, namely, that the metapterygial edge of the fish fin corresponds with the radial or thumb side of the hand, and consequently that the dorsal surface of the fish fin is the palmar (or plantar) surface. But I have no new facts.

By reviewing Gegenbaur's work it will be seen that this theory of his rests upon the form of the limbs in the Elasmobranchii and Dipnoi. In the former group it is the hind limbs which furnish nearly all the evidence. The fore limbs (pectorals) are brought in merely to testify to the *biserial* character of the archipterygium, of which no Elasmobranch *ventral* gives a sign. That is to say, the ventrals having testified to the archipterygium, and that having been accepted, the pectorals find use for themselves in showing that it was fringed down the median as well as the lateral side. If then the same form of limb is found in Elasmobranch and Dipnoan, the same form was undoubtedly possessed by their common ancestors. But as their common ancestors were also undoubtedly common ancestors of all Gnathostomes, therefore all Gnathostome limbs must have been derived from this form.

The testimony of the Ganoids and Teleosts seems to me to be somewhat adverse to the theory. Again it is impossible to think that that of the Stapedifera can be very clearly in its favor, when Huxley, while accepting the archipterygium as the parent form, gives an explanation of the chiropterygium entirely distinct from and utterly inconsistent with that of Gegenbaur.

Any opinion adverse to the archipterygium theory will have difficulty in maintaining itself, so long as it does not show that the resemblance between the fins of sharks, and those of Dipnoi is a merely superficial one, and is not able to suggest how a certain show of resemblance might have arisen in two entirely distinct and different series of developments.

Another View of the Origin of Vertebrate Limbs.

Into competition with this theory, which sees in the fin of *Ceratodus* that from which all other limbs have been derived, I bring a second which sees in the same only a special development peculiar to the Dipnoi. It is this.

As the dorsal and anal fins were specializations of the median folds of Amphioxus, so the paired fins were specializations of the two lateral folds which are supplementary to the median in completing the circuit of the body. These lateral folds, then, are the homologues of the Wolffian ridges, in embryos of higher forms. Here, as in the median fins, there were formed chondroid and finally cartilaginous rods. These became at least twice segmented. The oral ones, with more or less concrecence proximally, were prolonged inwards. The cartilages spreading met in the middle line, and a later extension of the cartilages dorsal completed the limb girdle.

If now we seek to determine the form of limb for the *Protognathostomi*, that is to say, for that time for which the archipterygium in its entirety is proposed, we should propose this.

The limbs of the Protognathostomi consisted of a series of parallel articulated cartilaginous rays. They may have coalesced somewhat proximally and oral. In the ventral pair they had extended themselves mesial until they had nearly or quite met and formed the hip girdle. They had not here extended themselves dorsal. In the pectoral limb the same state of things prevailed, but was carried a step further, namely, by the dorsal extension of the cartilage constituting the scapular portion, thus more nearly forming a ring or girdle.

This theory naturally divides itself into two parts, namely, the derivation of the Gnathostome limb from a series of simple parallel

cartilages, and the derivation of the latter from the lateral folds of *Amphioxus*.

Though the last mentioned portion of the theory would derive considerable strength from the establishment of the first, it is not a necessary consequence of it, and the first might be true even if the last were false. If the last be true, of course the first must be true.

The establishment of the derivation from the lateral folds of *Amphioxus* is made difficult from the absence of limbs or anything representing them in the two groups which (in a sense) stand between *Amphioxus* and the Elasmobranchs, namely, the Myxinoids and Lampreys.

As will be seen, it assumes the essential correctness of Huxley's suggestion with regard to the relation between the folds which grow down to inclose the atrial cavity of *Amphioxus* and the body walls of higher vertebrates. But it is equally consistent with Huxley's* entire suggestion, as put forth by him, or with Ray Lankester's† modification of it.

On the other hand, it is inconsistent with Kowalewsky's‡ view of the homology between these and opercular folds. This must perhaps be considered still an open question, though Rolph's§ arguments on the other side seem to me of much less weight than they do to Semper.||

Yet even if this homology with the lateral folds should have to be given up (the embryology of the Marsipobranchs will throw considerable light on it), the very frequent occurrence of the formation of external lateral folds parallel to the axis of the body in the bilateral animals in general and in the Vertebrates in particular, renders it quite possible that the paired fins may have had a *similar* origin. At present, however, I am strongly of the opinion that they are

* Journ. of Linn. Soc., vol. xii, No. 59, May, 1875. Huxley, Classification of the Animal Kingdom.

† Quarterly Journ. of Micr. Sc., New series, No. 59, July, 1875. Ray Lankester, New Points in the Structure of *Amphioxus*.

‡ Mem. St. Petersburg Acad., VII Series, tome xi, No. 4, 1867. A. Kowalewsky, Entwicklungsgeschichte des *Amphioxus lanceolatus*.

§ Sitzungsberichte der Naturforschenden Gesellschaft zu Leipzig. Jahrg. II, No. 1, Jan. 29, 1875. Rolph, Unters. über den Bau des *Amphioxus lanceolatus*. See also for a complete account of his investigations, under the same title, Morph. Jahrb., Bd., ii, Hft. 1, 1876.

|| C. Semper, Die Verwandtschaftsbeziehungen der gegliederten Thiere, 1875, p. 317. (Sep. Abdr. aus Semper: Arbeiten a. d. Zoolog-zootom Institut zu Würzburg, Bd. II).

derived directly from the lateral folds of *Amphioxus*. These in their turn may be referred to a reduplication of the process which has already formed the atrial space, but which is not carried so far here, in the lateral folds.

Ventral Fin of Acipenser.

The ventral fin of a young specimen of *Acipenser brevirostris* is exhibited in Pl. LIX, figs. 64 and 65. The fin of one side is separate from that of the other, no synchondrosis uniting the two halves of the girdle. But the part *b* approaches closely its fellow. The same separateness of the two sides obtains in the shoulder girdle. The composition of the fin is perfectly evident. Beginning at the aboral end of the row, we have first three separate and parallel rays. The proximal joints increase in length from the first to the last of the three. In the remaining rays these basal joints, increasing still more in length, have united with each other to form the large pelvic piece *ab*. The composite nature of this is confirmed by the groovings of the surface, which extend about a centimeter before they finally fade out.

The iliac process, *a*, is half a centimeter high.

In fig. 64 the proximal joint of the penultimate ray grows up a little under the last ray, in the manner familiar in the median fins. It does not happen to occur in the fin of the other side, fig. 65.

The predominant three-fold division obtains. But the penultimate ray in fig. 64 has a tip or an extra segmentation, and *c* is without the distal segmentation.

The breadth and the outline of *c* raises the suspicion of its double character.

I have had no opportunity of examining other Ganoid fins, and this one of *Acipenser* seems, on the whole, that which most nearly approaches the parent form of the Gnathostomes. But while in the independence of the two sides, in the separateness of the rays, and the simple segmentation, it gives us the early form more complete than is elsewhere found, in the number of rays and in the absence of the iliac process the shark ventrals are less advanced.

Elasmobranch Ventrals.

We now turn to the derivation of the ventral fins of sharks, one of the two abutments of the broad span of the archipterygium theory.

The series of steps by which I conceive them to have been derived


from the row of parallel rays is presented in woodcuts A, B, C and D.




The kind of 'change invoked is simply coneresence, with some spreading of the cartilage. The former of these processes is abundantly shown in the case of the median fins, while something of the latter process is seen in Pl. LI, figs. 12, 13, 14, 15; Pl. LIII, fig. 27; Pl. LVI, fig. 46; Pl. LVII, fig. 49. And it is noteworthy that here the rays which prolong themselves proximally are the oral ones, just as they are in the ventral fins. As for the coneresence, this has been carried much farther in the dorsal fins of *Squalus Americanus*, *Myliobatis* and *Raia levis* than it has in the shark ventrals.

It is barely possible that the definiteness and constancy of the coneresence in the latter may be in whole or in part determined by the copulatory function of the last part of the fin in male Elasmobranchs.

While the derivation of the ventral fins is thus easy from a series of parallel cartilages, we find much greater difficulty in the case of some of the median fins, in *Raia levis*, for example, which is, unless my own preconceptions deceive me, a far better case of a biserial archipterygium than any furnished in the paired fins, aside from *Ceratodus*.



Indeed I may state that the origin of this paper lay in an observation of a fin of another species of *Raia*, not however well enough preserved



for drawing. The very striking similarity to the uniserial archipterygium raised the question whether the median fins, at any rate the dorsal and anal fins, might not have arisen from the same archipterygium. The result of my investigations was a decided negative. It has been absolutely proved that they did not so originate, and the way in which they did originate has been clearly shown.

While then, on the one hand it has been shown that the development of a pair of fins, whose skeleton consists of a series of parallel rays clothed on each side with a layer of muscle, as a specialization of the lateral folds (metapleura of Ray Lankester) of *Amphioxus*, contains no steps which have not been taken in the same animals in the case of the median fins, so also it has been shown that the development, of the ventral fins and the pelvic girdle of sharks from such a series exhibits no processes or kinds of change which are not also exhibited in the median fins of those same fishes. When we contrast the changes from a series of parallel rays to the completed ventral fin of the shark, as it has been given above, with the changes which Gegenbaur supposes to have made it out of the archipterygium, namely, the stripping off of every one of the median rays, for no sign of them is ever found in the ventral fins aside from the Dipnoi, and the slipping off of the oral portion of the rays to immediately articulate with the shoulder girdle, I hardly think that those changes of his will appear so well evidenced as these changes which I believe to have taken place. And when the utter darkness that covers the development of the archipterygium itself (for it does not seem fair to the archipterygium to make much account of the suggestion of Gegenbaur respecting the branchial arches, *Unters.*, Hft. iii, p. 181, note) is contrasted with the familiar changes which would have brought these Selaechian fins out of the lateral folds *Amphioxus*, I hardly think the advantage can lie with the archipterygium.

*Homodynamism of Median and Paired Fins.**

Let us compare the ventral with the dorsal fins, say in *Mustelus canis*.

* Since this paper was written, I have found a paper of Humphrey's on the Homological Relations of Mesial and Lateral Fins of Osseous Fishes, *Journ. of Anat. and Phys.*, Nov., 1870. Here a comparison between the fins in question is made in the case of the Pike, and the "Iliac" or "Pubic" bones in osseous fishes are assimilated to the interneural spines or to the proximal part of them. Goodsir had made some earlier comparisons without valuable result. See *Anatomical Memoirs*, vol. ii, p. 106.

In each there is a layer of muscle on each side of the cartilaginous skeleton; this flat mass is in each divided into separate muscles by septa running from between the skeletal rods straight to the integument, in the way exhibited in Pl. LIX, fig. 66; in each these muscles develop a tendon in their middle plane parallel to the sides of the fin; this is inserted in the fascia over the terminal cartilages where the horny fibers begin. These last are the same in nature and arrangement in each.

The skeletal elements remain (see Pl. LIX, fig. 67). We have a short terminal piece in each, then comes a longer middle piece. There is left in the median fin a proximal row of cartilages, for the most part separate, which are again much shorter than the middle pieces. In the ventral fin the solid basale metapterygii and a half of the girdle correspond to these in every particular, except in not being of separate rays. The similarity between the two fins is complete except in a single particular. And that exception would be removed by a process which is familiar in both, namely conerescence. Even as they stand, I think that a ventral fin with one half the girdle resembles the dorsal more than it does the pectoral.

A certain amount of similarity warrants us in inferring an earlier state when the similarity was greater. It has been proved that at that earlier time the median fins were composed of separate rays. The greater similarity, then, can only be attained by the resolution of the basale metapterygii into its component parts.

In the Elasmobranchs, as is well known, the primordial fin-skeleton is supplemented in both median and paired fins by the well known horny fibers. In the higher fishes these are replaced by the dermal rays. The presence of horny fibers in the adipose fin of the salmon, shows that the horny fibers were the earlier form. Therefore the same changes have been taking place in the median and paired fins at the same time. The same general result, i. e., of concomitant variation in median and paired fins is confirmed in the sub-groups of Elasmobranchs, with regard to minor changes in the primordial skeleton.

The homodynamism of median and paired fins comes out strikingly in *Centrino Salviani*. Here, according to descriptions, a fold of skin is raised along the median line of the back, recalling the early continuous fold of skin along the back in *Amphioxus*. But similarly there appear two folds of skin along the sides, recalling the continuous lateral folds of *Amphioxus*. It is at once a proof of the homodynamism of the two, and a confirmation of the views here presented of their origin.

The Dipnoan Fin.

In the Teleosts and Ganoids, which are in a sense intermediate between the Dipnoi and Elasmobranchii, the limb skeleton has been comparatively little modified from the form in this paper set forth as the typical limb skeleton of Gnathostomi. Scarcely any other processes than reduction of the number of rays and concrescence are concerned.

Now in the fin of *Ceratodus* the archipterygium form has certainly been developed, and if the previous views be correct, it has been developed from this series of parallel rays.

Günther* has suggested one way, and Gegenbaur† another, in which a row of parallel rays might transform themselves into an archipterygium form. But it is possible that these fringing rays are new developments. They look very much like it in *Protopterus*. Peters‡ has called attention, in this connection, to the similar structure of the dorsal finlets of *Polypterus*, and these might throw some light on the subject. For myself, I am strongly inclined to suspect that the three portions of the second piece of the limb of *Ceratodus*, which Günther describes, indicate three fin-rays, and that the feathering of one of these is a later development. The fact that Huxley could find no sign of division in his specimens seems of little weight in view of the complete fusion which we know takes place here and there in median fins.

The Limb of Stapedifera.

Pl. LX, fig. 70 (*Ichthyosaurus*), exhibits my view of the composition of the limb of air-breathing vertebrates. The dotted lines indicate the separate rays of which it is composed. But there are other ways in which it may have been derived from a series of parallel rays, and I offer this merely as the most probable interpretation so far as I can now see. Fig. 71 does the same for the hind limb of *Cryptobranchus Japonicus*. The curvature of the rays has been exhibited in a marked degree quite frequently in Elasmobranch median fins.

The Innervation of the Paired Fins.

I have made complete and definite observations of the innervation only in a single case, namely in the pectoral fin of *Mustelus canis*. This fin is supplied by the first 15 myelonal nerves together with a very small branch from the vagus. The simplest condition is seen in the aborad nerves.

* Phil. Trans., vol. clxi, Pt. ii, p. 534.

† Unters., Hft. iii, p. 181, note.

‡ Muller's Archiv, 1845, p. 3.

The aborad four (12-15) branches, coming directly from the myelon, advance, each by itself, close to the metapterygium, where they each bifurcate, sending one branch to the dorsal and the other to the ventral side of the fin. The next four (8-11) unite to form a rather loose plexus, which separates again into four nerves, which then sub-divide in the same way as the last mentioned four, except that the orad of these behaves a trifle differently in a manner hereafter to be described.

Now the first seven nerves unite with one another and with the minute branch of the vagus in the following way. The vagal branch emerges from the skull with that nerve, but already rolled up as a separate branch and easily to be separated from it; this joins the first myelonal nerve and this the second, and their sum the third, and so on, until we have a cord formed of the vagal and first seven myelonal branches. This sends off a branch to the muscles and integument in front of the shoulder girdle, but the main part of it proceeds on its way to enter the foramen called by Gegenbaur, *Eintrittsöffnung*, and then divides within the cartilage of the girdle in the way which he has described, and similarly to the aborad nerves, which he has left unnoticed, sending one branch to the dorsal and the other to the ventral muscles of the fin. Now the eighth nerve sends off its ventral branch like those aborad of it, but the dorsal branch enters the entrance-opening with the cord of the vagus and 1-7 spinal nerves; but it does not unite with this cord till after the latter has divided, and then unites with its dorsal branch and emerges with that from the cartilage on the dorsal side of the fin. In another specimen, this dorsal branch of the eighth nerve enters the cartilage by a minute separate foramen, but unites with the dorsal branch of the anterior cord, as in this case.

As stated, my observations in the other cases have not been as thorough, and I cannot give the number of nerves, but in the ventral fin the arrangement is as follows. A number of nerves are gathered together to form the orad cord. This, on coming opposite the foramen in the pelvic girdle, divides and sends its branch to the ventral side of the fin through that. Then the other aborad nerves coming out, each by itself, to the metapterygium divide into two branches for the two sides of the fin, just as in the case of the pectoral fin. This is in *Mustelus canis*.

In *Eugomphodus litoralis*, see Pl. LX, fig. 60, from the articulation of two or three rays with the girdle, aborad of those which by their concrescence mark themselves out as the propterygium, it appears

that the articulation (or failure of concrecence) of the metapterygium and girdle has taken place farther aborad, and consequently a greater number of rays devote their basal parts to the formation of the pelvic girdle. Expectedly then, we find that the cartilage does in fact spread around the branches of the next two nerves. Through the foramina the ventral branches of these nerves pass, while the branches to the dorsal side pass along to that side above the cartilage.

The difference in respect to the relation between the cartilage and the nerves in the pectoral and ventral limb is, that in the former the cartilage thickens so as to include the branching place of the first nerve or bundle of nerves, while in the pelvic limb, it is thinner and merely transmits the ventral branch. In his Memoir on the Shoulder Girdle, Gegenbaur has called attention to the two branching canals or two pairs of openings in the shoulder girdle of the Batoidei. He states that he has not observed whether the aborad one is traversed by a nerve. I have examined this in the case of *Raia erinaceus*, and found that both fore and after openings transmit nerves in the same way. This is evidently what would be anticipated from what has been herein said. We have here what we had in the ventral fin (and girdle) of *Eugomphodus*; a greater number of rays are devoted to girdle building, and another bundle of nerves is included in the spreading cartilage.

The observations of Rolph on the innervation of *Amphioxus** are in complete agreement with what would be required by the view here advocated. He says "Der ventrale Ast verläuft herab bis in die Seitenfalten. Beim Eintritt in dieselben theilt er sich in zwei Aeste, deren einer (n_2) an der Aussenwand der Seitenfalte hinzieht; der andere durchläuft die Seitenfalte in querer Richtung, um in die Bauchmuskulatur über zu treten, in der er sich nicht weiter verfolgen lassen. Zuvor jedoch gibt er noch einen Zweig ab (n_3), welcher, n_2 parallel, an der inneren Wand der Seitenfalte verläuft."

The manner of innervation, then, seems to me as totally inconsistent with the Archipterygium theory as it is in thorough and telling harmony with the view which I have here presented.

ADDENDUM.

Since the views expressed in the foregoing pages were complete in my own mind six or eight months ago, I had looked for confirmation of them in the brilliant investigations of Balfour on the development

* Morph. Jahrb., Bd. ii, Hft. 1, p. 107, 1876.

of Elasmobranchs. The preliminary account, however, in the Journal of Microscopical Science, contained nothing bearing on the point, and the papers in the Journal of Anatomy and Physiology I have been able to obtain only irregularly. Immediately after the last proof of the preceding pages had been received, the number of that Journal for October, 1876, came into my hands. Here Balfour devotes three or four pages to the limbs. He says: "If the account just given of the development of the limb is an accurate record of what really takes place, it is not possible to deny that some light is thrown by it upon the first origin of the vertebrate limbs. The fact can only bear one interpretation, viz: *that the limbs are the remnants of continuous lateral fins.*"

"The development of the limbs is almost identically similar to that of the dorsal fins." He goes on to state that while none of his researches throw any light on the nature of the skeletal parts of the limb, they certainly lend no support to Gegenbaur's view of their derivation from the branchial skeleton. Thus these results have not only been reached independently, but from two different classes of facts. To the belief in the original continuity of the lateral fins and the homodynamism of median and paired fins I was led by observations on adult forms, and particularly on the skeleton. Balfour comes to the same results from embryological investigations, in that group from which on general grounds an answer was most to be expected; nor do these investigations regard the skeleton.

I have also just received the last number of the Morph. Jahrb. It contains a paper by Wiedersheim* confirming Gegenbaur's view respecting the double nature of the centrale. This had previously been shown only in the tarsus of *Cryptobranchus Japonicus*, (and in the Enaliosaurs). Wiedersheim shows its double character in three Siberian species of Urodela, in both carpus and tarsus. This is a very important confirmation of the chiropterygium, and relieves us of suspicions with regard to its correctness when we push our inquiries into earlier history and more simple forms.

In the same number of the Jahrbuch is a paper by Gegenbaur† on the archipterygium theory. He modifies his explanation of the Stapediferal limb to accord with Huxley's view of the homology of edges and faces of limb and fin. He says that while he does not

* Morph. Jahrb., Bd. ii, Hft. 3. R. Wiedersheim, Die ältesten Formen des Carpus und Tarsus der heutigen Amphibien.

† C. Gegenbaur, Zur Morphologie der Gliedmaassen der Wirbelthiere.

think the correctness of this view fully demonstrated, still he thinks there is a decided balance of probability in its favor. Therefore the ulnar side of the arm now appears as the *Stammreihe*. In other particulars Gegenbaur reaffirms his previous views. He proceeds to devote considerable space to the discussion of the origin of the archipterygium, and again proposes to assimilate the limb and limb-girdles to the gill-arches with their rays. He supports this suggestion with considerable argumentation. To this position the archipterygium theory leads him.

I take this opportunity for expressing my sense of the great advantages furnished by the U. S. Fish Commission for the study of marine life on our coasts, and in particular by the biological laboratory at Wood's Hole, established in connection with that Commission, and also for acknowledging my personal indebtedness to Professor Baird, through whose courtesy I have enjoyed these facilities for a number of summers.

EXPLANATION OF PLATES.

The figures are all drawn with a camera, and photo-lithographed. They are all three-fourths of the size of the originals, except figs. 1, 2, 3, 4, 20, 21, 22, 60; figs. 20, 21 and 22 are three-eighths, and 60 is a little less than twice natural size.

PLATE XLIX.

- Figure 1.—*Petromyzon marinus*. *a*, Ligamentum longitudinale, or ridge-pole of myelonal canal; *b*, notochord; *c*, neural arches.
 Figure 2.—*Petromyzon marinus*. *a*, fin-rays; *b*, intermuscular septa.
 Figure 3.—*Petromyzon marinus*. *a*, fin-ray; *b*, fin-muscles.
 Figure 4.—Section of *Petromyzon marinus*, to show the relation of the neural arches to the muscular segments; *a*, intermuscular septum; *b*, neural arch; *c*, blood-vessel; *d*, muscular segment; *e*, fatty-fibrous ridge-pole of neural canal.
 Figures 5, 6.—*Mustelus canis*. First dorsal.

PLATE L.

- Figures 7-10.—*Mustelus canis*. First dorsal.
 Figure 11.—*Mustelus canis*. Second dorsal.

PLATE LI.

- Figures 12-15.—*Mustelus canis*. Second dorsal.
 Figures 16, 17.—*Mustelus canis*. Anal.

PLATE LII.

- Figures 18, 19.—*Mustelus canis*. Anal.
 Figure 20.—*Galeocerdo tigrinus*. First dorsal.
 Figure 21.—*Galeocerdo tigrinus*. Second dorsal.
 Figure 22.—*Galeocerdo tigrinus*. Anal.
 Figure 23.—*Eulamia Milberti*. First dorsal.

PLATE LIII.

- Figure 24.—*Eulamia Milberti*. First dorsal.
 Figures 25, 26.— “ “ Second dorsal.
 Figures 27, 28.— “ “ Anal.
 Figure 29.—*Sphyrna zygaena*. First dorsal.

PLATE LIV.

- Figure 30.—*Sphyrna zygaena*. First dorsal.
 Figure 31.— “ “ Second dorsal.
 Figure 32.— “ “ Anal.
 Figures 33-35.—*Eugomphodus (Odontaspis) litoralis*. First dorsal.

PLATE LV.

- Figures 36-39.—*Eugomphodus litoralis*. First dorsal.
 Figures 40, 41.— “ “ Second dorsal.

PLATE LVI.

- Figures 42-46.—*Eugomphodus litoralis*. Second dorsal.
 Figures 47, 48.— “ “ Anal.

PLATE LVII.

Figures 49, 50 — *Eugomphodus litoralis*. Anal.

Figures 51–54.—*Squalus (Acanthias) Americanus*. First dorsal.

Figures 55, 56.— “ “ “ Second dorsal. The spine is removed in figs. 53, 55, 57; but not in figs. 51, 52, 54, 56.

PLATE LVIII.

Figure 57.—*Squalus Americanus*. Second dorsal.

Figure 58.—*Raia levis*. First dorsal.

Figure 59.— “ “ Second dorsal.

Figure 60.—*Myliobatis Fremenvillei*. Sole dorsal.

Figure 61.—*Acipenser brevirostris*. Sole dorsal; *a*, neural arch; *b*, intercalary cartilages; *c*, neural spine; *d*, foramen for ventral branch of spinal nerve; *e*, foramen for dorsal branch of spinal nerve.

PLATE LIX.

Figure 62.—*Acipenser brevirostris*. Anal. Opposite $2\frac{1}{2}$ vertebræ.

Figure 63.—Section of vertebrarium of *Acipenser brevirostris*; *a*, neural spine; *b*, ligamentum longitudinale; *c*, neural arch.

Figure 64.—*Acipenser brevirostris*, left ventral, from above; *a*, iliac process.

Figure 65.—*Acipenser brevirostris*, portion of right ventral, from below.

Figure 66.—Section of pectoral fin of *Mustelus canis*.

Figure 67.—*Mustelus canis*, ventral.

Figure 68.—*Mustelus canis*, pectoral detached from girdle.

PLATE LX.

Figure 69.—*Eugomphodus litoralis*. Ventrals; *a*, *b* and *c* nerve-foramens.

Figure 70.—Pectoral limb of *Ichthyosaurus*, after Cuvier. *Car.*, carpalia; *Cen.*, centralia; *u*, ulnare; *i*, intermedium; *r*, radiale; *U*, ulna; *R*, radius; *H*, humerus.

I take, though with some hesitation, Gegenbaur's identification of the radial and ulnar sides in this limb of *Ichthyosaurus*.

Figure 71.—*Cryptobranchus Japonicus*. Hind foot, after Hyrtl, Schediasma anatomicum, 1865.

VIII.--THE EARLY STAGES OF *HIPPA TALPOIDA*, WITH A NOTE ON THE STRUCTURE OF THE MANDIBLES AND MAXILLÆ IN *HIPPA* AND REMIPES. BY SIDNEY I. SMITH.


THE biological station, established under the auspices of the United States Commissioner of Fish and Fisheries, at Wood's Hole, Massachusetts, during the summer of 1875, afforded several naturalists, and among them the writer, excellent facilities for studying the marine animals of Vineyard Sound and the adjacent waters. The locality is very favorable for obtaining in abundance the free-swimming larvæ of a great variety of marine animals. Among the young of numerous species of crustacea, the zocæ of *Hippa* were particularly interesting, and I succeeded in obtaining a nearly complete series of the post-embryonal stages of that peculiar genus.

Since almost nothing has been published in regard to the habits of any of the species of Hippidæ or Albulidæ, a few words in regard to the habits of the only species, of either of these groups, living upon the coast of New England may not be out of place here.

Hippa talpoida inhabits the entire eastern coast of the United States from Cape Cod southward to the west coast of Florida; Egmont Key being its most southern and western habitat known to me. At what point it is met or replaced by the Brazilian *H. emerita*, I am unable to determine, never having seen specimens of either species from, or the record of their occurrence in, the West Indies or Central America, although some species of the genus probably inhabits both these regions.

On the sandy coasts of the southern United States the *H. talpoida* is apparently very abundant, while on the coast of New England it is much less common, being found only in special localities, although,

Fig. 1.*



* *Hippa talpoida*, adult female with the antennæ extruded, dorsal view, enlarged about two diameters.

from its gregarious habits, it is usually found in abundance in such localities. The northern range of this, as well as of numerous other, southern species is undoubtedly restricted by the extreme cold of the winters; and exceptionally cold seasons probably destroy a large part of the individuals over considerable portions of the coast. During the summer of 1870 not a specimen of the adult or half grown *Hippa* could be found at Fire Island Beach, Long Island, although the extensive sandy beaches of that region offer specially favorable localities, which were thoroughly searched; but during the last of August and early September, the young just changed from the zoea, and also in a little later stage, appeared abundantly upon the beaches. During the following summer no fully grown specimens were found on the shores of Vineyard Sound, though half grown specimens (perhaps from the young of the previous season) were common. During the summer of 1875, fully grown specimens of both sexes were found in great abundance at a single, very restricted locality near Nobska Point, on the shore of Vineyard Sound, although at this time none could be found at the particular locality where they were common in 1871.

Upon our shores, as far as I have observed, the *Hippa* inhabits sandy beaches which are somewhat exposed to the action of the waves. It seems to prefer only a narrow zone of the shore, at or very near low water mark, where it lives gregariously, burrowing in the loose and changing sands. At the locality near Nobska Point above referred to, it was obtained in great abundance by digging over the sand just at the edge of the receding waves. Several individuals were often thrown out at a single stroke of the spade, but the wonderful rapidity with which these animals burrow made it extremely difficult to secure more than one or two of them at a time. The smooth, oval form of the animal, with the peculiar structure of the short and stout second, third, and fourth pairs of thoracic legs, enables them to burrow with far greater rapidity than any other crustacean I have observed. Like many other sand-dwelling crustaceans, they burrow only backwards; and the wedge-shaped posterior extremity of the animal, formed by the abrupt bend in the abdomen, adapts them admirably for movement in this direction. When thrown upon the wet beach, they push themselves backward with the burrowing thoracic legs and, by digging with the appendages of the sixth segment of the abdomen slightly into the surface, direct the posterior extremity of the body downward into the sand. Upon the beaches, at least where there are any waves, they seem usually to be buried completely

beneath the surface. Occasionally, however, they are found swimming about in pools left by the tide, and they undoubtedly, when undisturbed, sometimes come out and swim in the same way along the shore, though they probably never venture far from the bottom.

When first placed in an aquarium with a few inches of sand at the bottom, they invariably plunged at once entirely beneath the sand, but, after a few moments of quiet, usually worked themselves gradually towards the surface, resting in a nearly perpendicular position with just the tips of the antennulæ and eyes at the surface, while the ex-current water from the branchiæ formed a small opening and a slightly boiling motion in the sand. Occasionally, when entirely undisturbed, they would suddenly leave the sand and swim rapidly round the top of the aquarium for a moment and then dive suddenly to the bottom and bury themselves in the sand. In swimming, as well as in burrowing, the telson was carried appressed to the sternum and they invariably moved backward, the motion being apparently produced by the appendages of the sixth abdominal segment and the anterior thoracic legs, while the latter served also as steering organs.

During all the ordinary motions of swimming and burrowing, I have never seen the antennæ extruded, although the peculiar arrangement of the peduncular segments and their complex system of muscles are apparently specially adapted for extending and withdrawing these beautiful organs. When the animals are thrown into alcohol however, the antennæ are sometimes thrown out convulsively and then immediately retracted. In life the antennæ are most of the time held in the position in which they are usually found in alcoholic specimens, that is, between the second and external maxillipeds, with the peduncles crossed in front, and the flagella curved down and entirely round the mouth so that their dense armament of setæ all project inward. When extruded, the distal segments of the peduncle are revolved half way round on the proximal ones, so as to carry the whole appendage to its own side of the animal and throw the curve of the flagellum into a reversed position. Judging from the peculiarly armed setæ of the flagella, one of the principal offices of the antennæ is the removal of parasitic growths and all other foreign substances from the appendages of the anterior portion of the animal.

The mouth parts of the adult are not adapted for ordinary prehension or mastication, but I am unable to make any positive statement in regard to the food of these animals. In all specimens examined the alimentary canal was filled with fine sand which seemed to be nearly free from animal or vegetable matter. The material from the

stomach, however, shew, under the microscope, a small quantity of vegetable matter, and it seems probable that the sand is swallowed for the nutritive matter it may contain.

Upon the beaches of Vineyard Sound the two sexes appeared to occur in about equal numbers, although in museum collections the males are often rare. This is probably due to the great inequality in size between the male and female, the length of the carapax in the larger females from Vineyard Sound being 20 to 22^{mm}, while in the largest males it does not exceed 14^{mm}. The sexes differ also in the form of the telson (Plate XLVIII, figs. 7, 8) which is narrower and more triangular in the male than in the female.

Females carrying eggs were found during the entire month of August, and during that period the embryos within the eggs were nearly fully developed in many of them. Undoubtedly, however, the term of carrying eggs extends over a much longer period than this. The eggs are nearly spherical, .40 to .45^{mm} in diameter, and the yolk mass is orange yellow while the formed tissues of the embryo are nearly colorless. Numerous attempts to obtain newly hatched young, by keeping egg-carrying females in aquaria, failed from the parent invariably casting off the eggs before they were fully matured. Consequently I failed to secure the earliest stage of the zoea, for the youngest individuals taken in the towing net were apparently in the second stage.

Very nearly fully developed embryos, when removed from the egg, were found to possess all the normal articulated appendages of the fully formed zoæ, but there was no appearance of lateral spines upon the carapax and the rostrum was broad and obtuse. In this stage the embryo agrees almost perfectly with the figure of the zoea of *Hippa emerita* from the coast of Brazil, given by Fritz Müller in his work entitled "Für Darwin."* The difference between the embryo in this stage and the second zoea-stage (Plate XLV, fig. 1), in which

* English translation, London, 1869, p. 54, fig. 25. The figure is accompanied by the following paragraph: "The Zoëa of the Tatuira [*Hippa*] also appears to differ but little from those of the true Crabs, which it likewise resembles in its mode of locomotion. The carapax possesses only a short, broad frontal process; the posterior margin of the tail is edged with numerous short setæ." This, as far as I am aware, is the only published account of the development of any species of Hippidae, except a note by myself (in an article on "The Metamorphoses of the Lobster and other Crustacea," in the Report of the U. S. Commissioner of Fish and Fisheries, Part I, 1873, p. 530) recording the occurrence, at the surface in Vineyard Sound, of the young in what is described further on in these pages as the megalops-stage.

the rostrum and lateral spines are enormously developed, suggests the possibility that Müller had observed only imperfectly developed young zoeæ in which the rostrum and lateral spines were not expanded. It seems scarcely probable that such a difference could exist between the first stage of the zoea, when the veiling membrane, in which, on first escaping from the egg, the young are usually enveloped, has been entirely cast off and the lateral spines and the rostrum are fully expanded, and the second zoea-stage about to be described. The three later, true zoea-stages obtained are evidently contiguous steps in the development and are here designated the second, third, and last stages of the zoea. From this last stage the zoea passes at once into a stage closely resembling the adult in general form, but with the eyes still very large and the abdomen furnished with powerful swimming legs. This condition of the animal corresponds perfectly to the Brachyuran megalops and may properly be designated as the megalops-stage.

Second zoea-stage.

In this stage the young (Plate XLV, fig. 1, ventral view) are a little over 3^{mm} in length, from tip of rostrum to the posterior margin of the carapax, and a little over 2^{mm} between the tips of the lateral spines. In general form the carapax is oval, with the smaller end forward, and its surface is very smooth and regularly rounded. The dorsal surface of the carapax is strongly convex but very regularly rounded and wholly devoid of any rudiment of a dorsal spine, which is so generally characteristic of the zoeæ of Brachyura. At the bases of the ocular peduncles the carapax is sharply contracted laterally into an exceedingly long, very slender, and slightly tapering rostrum curved regularly downward until, toward the tip, it becomes nearly parallel with the posterior margin of the carapax. The lateral spines are nearly as long as the diameter of the carapax, are situated far back and low down on the sides of the carapax, and are directed downward and obliquely outward, but are not strongly curved. Beneath, the carapax curves inward on all sides, leaving a comparatively small opening which is wholly inferior, with its anterior portion about as broad as the telson, but posteriorly contracted into a narrow abdominal sinus, of which the rounded posterior margin is nearly on a line between the lateral spines. This shortening of the inferior opening, carrying the abdomen forward and wholly beneath the carapax, together with the absence of the dorsal spine, gives the animal an appearance unlike ordinary Brachyuran zoeæ.

The ocular peduncles are stout, regularly tapering to near the bases, and are usually carried perpendicular to the mesial plain, though they admit of considerable motion in all directions. The cornea is considerably larger than the diameter of the peduncle, its diameter being nearly a third of the horizontal diameter of the carapax, and, when the peduncle is held straight out, reaches slightly beyond the lateral margin of the carapax.

The antennulæ (Plate XLVI, fig. 1) are still rudimentary, simple, sack-like, unarticulated appendages, tapering toward the tip, which is furnished, as usual in this stage of development, with three stout, filiform, obtuse setæ, differing slightly in length, diameter, and amount of curvature, and of which the longest is about half the length of the antennula itself.

The antennæ (Plate XLVI, fig. 2) are of about the same length as the antennulæ, but of nearly the same diameter throughout, and are armed distally, at the outer edge, with an acute, dentiform process (*a*, fig. 2) directed straight forward and itself armed with a minute, setiform spine on the inner edge near the tip. Between the base of this process and a slight, rounded prominence (*c*, fig. 2), situated at the extremity of the inner margin, and which represents the rudimentary flagellum, there is a similar, but slightly more slender, process (*b*) attached at its base by an oblique articulation and armed, near the tip, with a minute, setiform spine like that upon the outer process.

The oral appendages differ very little from their condition in the last zoea-stage, under which they are fully described. The labrum and labium differ scarcely at all, except in size, in the three zoea-stages here described. The labrum, as seen from beneath, is a broad, somewhat triangular prominence between the bases of the antennulæ and the tips of the mandibles. The labium is deeply bilobed, though far less deeply than in the adult, with the lobes broadly rounded and the entire margin clothed with microscopic hairs.

The mandibles are nearly as in the last zoea-stage. They are stout at the bases, but taper to very slender tips, which are only slightly different on the right and left side. There is no molar area, but the crown of the mandible is longest in a vertical direction and is armed inferiorly with four long, but blunt, teeth which decrease rapidly in size as they approach the middle of the crown, where they are met by a series of six or seven long, slender, seta-like processes which occupy the superior half of the coronal margin.

The first pair of maxillæ (Plate XLVI, fig. 11) are symmetrical and composed of the same parts as in the adult. The inner lobe

(*a*, fig. 11) is small and tipped with three long setae. The outer lobe (*b*, fig. 11) is broader than the inner and armed at the extremity with three nearly equal, long and slender teeth, of which the distal one appears like a process from the margin, showing no line of articulation at its base. The palpus (*c*, fig. 11) is very small, composed of a single segment and tipped with a long, plumose seta.

The second pair of maxillae (Plate XLVII, fig. 1) are very imperfect. The protognath (*a*, fig. 1) is a small, obtuse lobe tipped with three short setae. The scaphognath (*c*, fig. 1) projects beyond it anteriorly as a slightly larger lobe, while posteriorly it is broad but short and truncated, and the anterior lobe and the outer edge are, as yet, alone furnished with setae.

The first and second pairs of maxillipeds, or natatory legs, (Plate XLV, fig. 1, second pair) are similar in structure to those of most zoeae and differ only slightly from each other. In both pairs the basal portion, or protognath, is alike stout, about as long as the exognath, and unarmed, except by three or four minute setae on the distal portion of the inner margin. The exognaths, or natatory branches, alike in both pairs, are nearly cylindrical, but flattened at the tips, where they each bear a series of eight slender, plumose setae, which, in the middle, are as long as the exognath itself but decrease in length to the outer ones, which are scarcely more than two-thirds as long. The inner branch, or endognath, in both pairs, is composed of four cylindrical segments subequal in length. In the first pair, however, the inner branch is shorter than the exognath, while in the second pair it is considerably longer. In both pairs the three proximal segments of the endognath are each armed with two or three small setae on the inner side, and the distal segment, which is much more slender than the others, is tipped with four setae, of which two are nearly as long as the segment itself and pectenated with minute, setiform spinules along one side, and the two others shorter and apparently unarmed.

The third pair of maxillipeds and the four anterior pairs of thoracic legs are, even at this early state, represented by a series of clearly defined, though entirely unsegmented, processes situated just above and back of the bases of the second maxillipeds and entirely within the carapax, but visible through it, in a lateral view of the animal, in a line nearly parallel with the posterior margin of the carapax. In the single specimen examined, no lobes representing the slender, posterior thoracic legs of the adult could be discovered. Above each of the processes representing the first four pairs of thoracic legs there is a minute, papilliform process apparently representing one of the branchial appendages belonging to these legs in the later stages.

The abdomen is smaller and has much less freedom of motion in the mesial plane than in most Brachyuran zoeae. The first segment is not clearly differentiated from the thorax. The second, third, fourth, and fifth segments are entirely without appendages; the second and third are nearly equal in length and sub-cylindrical; the fourth is slightly shorter and is expanded considerably at the posterior extremity; and the fifth is about as long as the fourth, compressed vertically, and broadly expanded at the postero-lateral angles so that it is about twice as broad as long. The sixth segment is consolidated with the telson, forming a broad, lamelliform, caudal appendage about as long as the middle breadth of the carapax. The appendages of the sixth segment (Plate XLVIII, fig. 10) are small, rudimentary, appressed to the under side of the telson so as to be hidden from above, and are each composed of a stout basal segment and a single narrow lamella (the outer) tipped with two slender setae, of which the outer is about as long as the lamella itself and the inner much longer.

The lateral margins of the telson are slightly curved outward and unarmed, but project posteriorly into a stout tooth each side of the strongly arcuate posterior margin. This posterior margin has, in all the zoea-stages here described, a remarkably complex armament of ciliated spines and minute teeth (Plate XLVIII, figs. 13, 14, 15). In a considerable number of specimens in the third and the last stages, the number of these ciliated spines is usually twenty-six, of which the eighth, counting from either side, is the largest, and the sixth and tenth usually the next in size. One of the specimens in the second stage (Plate XLVIII, fig. 13) conforms strictly with this: there are two sub-median spines (*a, a*, fig. 13) separated by a single denticle, then each side a slightly larger spine (*b, b*) separated from the sub-median ones by a single denticle, then four alternately smaller and larger spines (*c, d, e, f*) separated from each other by two denticles at each interspace, the outer (*f*) of these four spines being the eighth, counting from either side, and the largest. Outside this large spine there are, each side, seven smaller spines separated by interspaces which increase toward the outer margin and are armed with from two to nine denticles. The space between the outer spine each side and the tooth of the lateral margin is greater than any of the interspaces between the spines and is armed with twelve or thirteen denticles. In the other specimen in this stage there are only twenty-five spines, a single median spine (Plate XLVIII, fig. 14, *a*) taking the place of the two sub-median spines and the denticle between them; otherwise the spines and denticles are essentially as in

the first specimen and as in the succeeding stages. This is apparently an abnormal variation in the armament of the telson. The arrangement of these spines, and especially whether they be odd or even in number, I have usually found a constant character for distinguishing the larval forms in different groups of Podophthalmia.

Of this stage only two specimens were secured, one taken on the evening of September 4, the other in the day-time the next day. In coloration and habits they agreed essentially with the young in the succeeding zoea-stages.

Third zoea-stage.

In general form and appearance the zoeæ in this stage very closely resemble those in the second, although they have increased considerably in size, and especially in the length of the rostrum, which is relatively longer and more slender. They are about 4.5^{mm} in length, from the tip of the rostrum to the posterior margin of the carapax, and nearly 3^{mm} between the tips of the lateral spines.

The ocular peduncles and eyes have increased only slightly in absolute size and are relatively smaller than in the second stage.

The antennulæ have changed very little. There is, as yet, apparently no distinction of peduncle and flagellum, although the two or three distal segments of the latter are faintly indicated, and, on the inner side, there are two or three filiform setæ on the penultimate segment in addition to the three on the terminal segment.

The antennæ (Plate XLVI, fig. 3) have increased in size but show no indication of segmentation. The two dentiform processes (*a*, *b*, fig. 3) have each two or three minute spirules at the tip, but are otherwise unchanged; the flagellum (*c*, fig. 3), however, has increased so as to project beyond the tips of the dentiform processes and show plainly its true character.

The labrum, labium, mandibles, and first maxillæ, except in size, do not differ appreciably from their condition in the second stage. The second maxillæ differ but little, the scaphognath being a little more elongated posteriorly, so as to approach slightly its form in the next stage.

The first and second pairs of maxillipeds differ from those of the first stage only in the exognaths, which are each furnished with ten instead of eight terminal setæ.

The lobes representing the third pair of maxillipeds and the four anterior pairs of thoracic legs have increased much in size, are

curled up closely beneath the sternum, and, in all the specimens examined, show the segments of the succeeding stage faintly indicated within. The posterior thoracic legs appear to be represented by a small process each side, just back of, and nearly hidden by, the rudimentary fourth pair. Four pairs of gills on each side are represented by two slender processes, one above the other, at the bases of each of the four anterior pairs of legs.

The proximal segments of the abdomen are almost exactly as in the previous stage, except there are very slight elevations beneath the second, third, fourth, and fifth segments, where the rudimentary legs are to appear in the succeeding stage. The sixth segment is still consolidated with the telson. Its appendages (Plate XLVIII, fig. 11) have increased much in size and the inner lamella (*c*, fig. 11) has appeared as a small, sack-like appendage at the base of the outer lamella (*b*, fig. 11), which is twice as long as in the previous stage, very narrow, only slightly expanded in the middle, and sub-truncate at the extremity, where it is furnished with four slender and curved setæ, of which the median ones are longer than the lamella itself, while the outer are little more than half as long. The telson is of the same form as in the previous stage and has the same number of ciliated spines in the armament of the posterior border, while the number of denticles in the interspaces has considerably increased, though they are not as numerous as in the succeeding stage. In one specimen there is the same abnormal arrangement of the spines described under the second stage, that is, there are only twenty-five spines in all, one median spine taking the place of the two sub-median spines and the denticles separating them.

The young in this stage were taken on several occasions, both in the day-time and evening, from August 28 to September 8. Their habits and coloration in life were the same as in the last zoea-stage.

Last zoea-stage.

The length from the tip of the rostrum to the posterior margin of the carapax and the breadth between the tips of the lateral spines are nearly twice as great as in the second stage, while the rostrum is relatively considerably longer than in either the second or the third stage, its entire length being nearly twice that of the carapax proper. The general form and appearance of the young in this stage are shown upon Plate XLV, figs. 2, 3, 4.

The eyes and ocular peduncles are very little larger absolutely than in the last stage. The diameter of the cornea is scarcely a

fourth the horizontal diameter of the carapax and the ocular peduncles are slightly shorter, proportionally, than in the earlier stages.

In the antennulae (Plate XLVI, fig. 5) the segmentation of the flagellum is carried nearly or quite to the peduncle, which, however, shows no division into segments and no clear separation from the flagellum. The flagellum itself is composed of six or seven segments which are a little broader than long and of which the terminal one is furnished with three filiform setae, the penultimate and antepenultimate with two or three each which are situated upon the inner side at the distal articulations, while there are two similarly situated, but small and rudimentary setae, upon the fourth segment from the tip.

The antennae (Plate XLVI, fig. 4) have increased very much in size, and the flagellum is much longer than the peduncle. The peduncle shows but one distinct articulation, which is near the bases of the dentiform processes and apparently represents the articulation between the second and third segments of the fully developed appendage. The dentiform processes (*a*, *b*, fig. 4) are much more slender and proportionally smaller than in the third stage, but are armed with the same number of spines at the tips. The flagellum (*c*, fig. 4) externally shows no indication of segmentation, but, in all the specimens examined, the articulations of the flagellum of the succeeding megalops-stage is distinctly visible beneath the integument, as shown in the figure.

The labrum (Plate XLVI, fig. 5, *b*), as seen from beneath, is a conspicuous, somewhat triangular prominence between the bases of the antennulae and the mandibles, with the margins and the inferior surface regularly rounded and without emarginations at any point. The labium (*d*, fig. 5) is deeply bilobed, the regularly rounded lobes projecting each side of the oral opening nearly to the tips of the mandibles and having the edges clothed with microscopic hairs throughout.

The mandibles (Plate XLVI, fig. 5, *c*; and fig. 6) are almost exactly as in the earlier stages. They are wholly without molar areas and terminate in narrow crowns, which are only slightly different on the right and left sides. In each mandible the inferior half of the coronal margin is armed with four stout teeth, the two most inferior of which are stout and obtusely pointed, the inferior being straight and much longer than the next, which slightly overlaps it at base, as seen from before or behind, and curves upward at the tip; while the two succeeding ones are short, triangular, more acute, and separated from each other by a considerable sinus. On the right side the inferior

tooth is a little more anterior than the others, while on the left side it is more posterior, so that these inferior marginal teeth overlap each other when the mandibles are brought together. The posterior side of the crown of the right mandible is slightly convex while the same side of the left is a little concave. On the superior half of the crown of each mandible there is a series of six slender processes, or teeth, of which the five inferior are very slender, setæ-like, and twice as long as the two triangular teeth just mentioned, while the superior, or marginal, one is stouter, acutely triangular, and divided at tip. In the specimen figured, there is also a small supplementary tooth on the right mandible, between the bases of the third and fourth processes of the superior half of the margin.

The first pair of maxillæ (Plate XLVI, fig. 5, *e*; and fig. 12) have changed scarcely at all, except in size, from their simple form in the second stage. The inner lobe has a single rudimentary seta, on the inner edge near the tip, in addition to the three terminal ones, the three teeth of the outer lobe are relatively a little shorter and very slightly stouter, and the palpus is considerably larger.

The second pair of maxillæ (Plate XLVII, fig. 2) have advanced considerably in their development and have apparently partially assumed their adult function. The protognath shows a slight indication of division into two lobes (*a*, *b*, fig. 2), of which the outer projects as far forward as the scaphognath, from which it is much more deeply separated than in the earlier stages. The scaphognath is twice as long as the protognath, and its posterior portion is elongated and narrowed at the extremity as in the later stages, but its inner edge is not yet margined with the characteristic plumose setæ.

The first and second pairs of maxillipeds (Plate XLV, figs. 2 and 4, and Plate XLVII, fig. 5) have the same form and structure as in the preceding stages, the only noticeable difference being the addition of still another pair of setæ at the tip of each exognath, making twelve in all, of which the outer are scarcely more than half as long as the middle ones, which are not quite equal in length to the exognath itself.

The third pair of maxillipeds and all the thoracic limbs (Plate XLV, fig. 4, and Plate XLVIII, fig. 9) are still curved inward beneath the sternum and entirely inclosed within the posterior portion of the carapax, although the four anterior pairs of legs have begun to show clearly the form which they assume in the megalops and later stages. The third pair of maxillipeds (*b*, Plate XLVIII, fig. 9) are still sub-cylindrical and in each the meral segment is only

a little stouter than the palpus, of which only the elongated dactylus and the stout propodus are clearly distinguishable. The first four pairs of thoracic legs (*c, d, e, f*, fig. 9) show clearly only the distal segments, but, of these, the propodus and dactylus have already assumed the peculiar form which characterizes them in the megalops-stage and in the adult, the propodus in the first, second, and third pairs being elongated into a prominent lobe on the inside at the distal extremity, while in the fourth pair (*f*, fig. 9) this segment is truncated at the distal end. The posterior thoracic legs (*g*, fig. 9) are slender, cylindrical, and the three distal segments are nearly equal in length, but there is no indication of the cheliform character which they assume in the megalops-stage. The branchial processes (*h*, fig. 9) above the bases of the legs have become more conspicuous than in the earlier stages and represent the full number of branchiæ in the adult, there being one above the base of each external maxilliped and two each upon the four anterior thoracic legs, making nine pairs in all.

The first segment of the abdomen is still in the same condition as in the second stage, not distinctly differentiated from the thorax, and the remaining segments themselves retain very nearly the same form as before. The abdominal legs of the second, third, fourth, and fifth segments (Plate XLV, fig. 4) are as long as the segments to which they belong, but are still sack-like, the base separated from the terminal portion by an obscure articulation, but with no indication of the separation of the outer from the inner lamella, and the appendages are evidently of no functional importance. The sixth segment is still closely united with the telson, although some of the specimens show a slight indication of the approaching articulation. The appendages of the sixth segment (Plate XLVIII, fig. 12) have increased very much in size, and the outer lamella (*b*) has become narrow-oval in outline, about a third as broad as long, with the tip rounded and furnished with six strongly curved setæ, of which the third from the outside is much longer than the lamella itself, while all the others are much shorter and decrease in length each way from the longest. The inner lamella (*c*, fig. 12) is two-thirds as long as the outer, nearly as broad, and regularly oval in outline, without hairs or setæ.

The telson (Plate XLV, figs. 3 and 4) has the same form as in the previous stages. Its posterior margin (Plate XLVIII, fig. 15) is armed with twenty-six ciliated spines having the same arrangement as in the earlier stages. The number of denticles in the interspaces between the spines is very much increased, there being three between

sub-median spines, two each side between these and the next outside of them, from three to twelve in each interspace between the other spines, and from thirty to forty between the outer spine each side and the tooth of the lateral margin.

The zoeæ in this stage were frequently taken at the surface, both in the day-time and evening, from August 7th to September 10th. In life, the entire animal is translucent with a slight greenish tint, except a brilliant spot of orange pigment with metallic lustre at the base of the rostrum and at the base of each of the lateral spines. These bright spots would often catch the eye in looking into the water when the rest of the animal was nearly or quite invisible. Their motions in the water are similar to those of zoeæ in general, except that the movements are much less rapid, and they seem comparatively sluggish in habit. In aquaria they were usually seen swimming slowly about the surface, seldom whirling off with the mad, gyratory motion so characteristic of many Brachyuran zoeæ.

Individuals in this last zoea-stage, when kept in confinement, often changed, at a single molt, to the megalops-stage described beyond, although many died during the process of molting.

The following table shows the relative measurements of specimens in each of the zoea-stages described :

	Second Stage.	Third Stage.	Last Stage.
Length from tip of rostrum to posterior margin of carapax,--	3·2mm.	4·4mm.	6·2mm.
Breadth between tips of lateral spines, -----	2·2	2·8	4·3
Length of rostrum from tip to front of ocular peduncles,----	1·8	2·8	4·7
“ “ lateral spines, about,-----	·8	1·1	1·3
“ “ carapax from front of ocular peduncles to posterior margin, -----	1·3	1·8	2·5
Breadth of carapax in the middle,-----	·9	1·2	1·7
“ “ telson, -----	·7	1·0	1·4

Megalops-stage.

In specimens recently changed from the last zoea-stage, the length of the carapax is slightly over 3^{mm} and that of the abdomen, when fully extended, a little less, while the breadth of the carapax is about 2^{mm}. In general form, the young in this stage resemble the adult, but differ essentially in much the same way that the Brachyuran megalops differs from its adult; the eyes still being relatively large, with short and thick peduncles, as in the last stage of the zoea, while the second, third, fourth, and fifth abdominal segments, as well as the sixth, are provided with strong swimming appendages.

The form of the carapax resembles that of the adult very closely, but is broader in proportion, being a half longer than broad; the lobes of the frontal margin (Plate XLV, fig. 5) are more obtuse and much broader; and the lateral lobes form the antero-lateral angles of the carapax, instead of being separated from the angles by a transverse portion, each side, nearly as long as the thickness of the peduncles of the antennæ, while the lateral margins of the carapax are more regular in outline and do not project downward so much between the first and fourth pairs of legs.

The ocular peduncles are still very short and stout, though the eyes have increased only a very little in absolute size since the last zoea-stage. The cornea is a little elongated and occupies the lateral portion of the tip of the peduncle; its greater diameter is still about a fourth of the horizontal diameter of the carapax and more than half the length of the peduncle.

The antennulæ (Plate XLVIII, fig. 1) reach to the extremities of the peduncles of the antennæ and show a marked advance over the previous stages. The segments of the peduncle are fully differentiated and the second segment (*b*, fig. 1) has already a marked prominence, tipped with a few plumose hairs, in place of the elongation so conspicuous upon the inferior side of the distal extremity of the same segment in the adult (Plate XLVIII, fig. 3). The flagellum (*d*, fig. 1) is composed of eight or nine short and stout segments, all, except one or two of the most proximal, armed inferiorly with plumose, or pectinate, setæ. The secondary flagellum (*e*, fig. 1) is represented by a single, minute segment tipped with a plumose seta.

The antennæ (Plate XLV, fig. 5, and Plate XLVI, figs. 7, 7*a*) are slightly longer than the carapax and have assumed all the important features of the adult. The segments of the peduncle (Plate XLVI, figs. 7, 7*a*) are fully differentiated and essentially the same as in the adult; and they have the same peculiar structure adapted to folding the antennæ across in front of the mouth and within the external maxillipeds. To aid in accomplishing this, there is a supplemental segment or rod (*f*, figs. 7, 7*a*) on the outer side of the third segment—and apparently a separately calcified part of it—which articulates proximally just within the lateral spine of the second segment and distally with the outer edge of the fourth segment, and is so separated, by non-calcified, flexible integument, from the third segment itself, as to move independently of it. When the antenna is extruded, this supplemental rod lies parallel with the outer margin of the third segment, from which it is then separated by a considerable space of

the non-calcified integument; but when the antenna is folded away within the external maxillipeds, the rod is transverse to its former position and is almost or quite hidden between the second and fourth segments, the outer, calcified portion of the latter segment folding over the previously exposed area of thin integument. The flagellum is stout, tapers very gradually to an obtuse and rather thick tip, and is composed of about twenty-five segments, which are very short proximally, but increase regularly in length distally until, near the tip, they are longer than broad. Each segment is armed upon one side with a fascicle of long, pectinated setæ. As seen in a transverse section of the flagellum (Plate XLV, fig. 6), the outer one of these setæ, on each side of the fascicle, is strongly curved inward at the extremity and much longer than the inner ones, which are nearly straight and armed, for only a part of their length, with much shorter, stronger, flattened, and blade-shaped teeth. In this stage there are usually from five to seven setæ to each segment, the median one or two often being simply acicular, or even very small and rudimentary.

This structure of the flagellum of the antenna is essentially the same as in the adult, where, however, the number and size of the setæ, as well as the extent of their armament, is enormously increased upon each segment, while the number of segments is several times greater than in the megalops-stage. In the ordinary adult specimens, there are one hundred to one hundred and fifty segments in the flagellum and eight to twelve setæ to each segment. There are two forms of these setæ in each fascicle, as in the megalops-stage. The outer one, each side of every fascicle, is very long and convolutedly curved inward at the extremity; while all the others are shorter, though varying much in length among themselves, and nearly straight. The long, outer setæ are armed, for nearly their whole length, with very long, almost filiform, secondary setæ, which are arranged in a double series along the inner side of the curve. These secondary setæ are exceedingly slender, very slightly tapering, from $\cdot 2$ to $\cdot 25^{\text{mm}}$ long,—the longer ones being on the distal half though not at the extremity,—and are placed so thickly that, in the middle portion of the seta, there are two hundred in the space of a millimeter. The extremity of the shaft of the seta itself is unarmed for a very short distance, curved sharply so as to be nearly parallel with the last of the secondary setæ, and tapers to a very slender and acute tip. The remaining setæ of each fascicle are nearly straight, or very slightly bent, and armed upon the outside of the curve. The teeth, as in the outer setæ, are arranged in two series, which here, however,

approach so closely that the bases are nearly in the same line, those of one series alternating with those of the other. The teeth themselves are much shorter than the secondary setæ of the outer setæ, being $\cdot 05$ to $\cdot 07^{\text{mm}}$ long, much stouter and less crowded, so that there are about an hundred in the space of a millimeter in the middle portion of the setæ. Near the base of the seta, a few of the teeth are very slender and hair-like, but the rest are stout, truncated at the tips, lamelliform, and placed with their bases transverse to the shaft of the seta, but with the blades twisted nearly half way round, so that the surface at the tip is nearly at a right angle to the base. The shaft of the seta itself extends a little way beyond the teeth in an acutely cultriform tip.

The oral appendages (Plate XLVI, figs. 8, 9, 13; Plate XLVII, figs. 3, 4, 6, 7) have undergone a transformation even more wonderful than the usual change from the zoea to the megalops. The mandibles have lost, almost entirely, the structure and function usual to them in all the Thoracostraca, and, together with the other oral appendages, have assumed very nearly the adult form.

The labrum, as seen from beneath (Plate XLVI, fig. 8, *a*), is much more elongated than in the zoea-stages, being about as broad as long, but with the sides still arcuate outward and not at all incurved as in the adult. The labium (*e*, fig. 8) is divided very deeply into two long and obtuse lobes, fringed with microscopic hairs along the oral margins, and projecting forward, each side of the mouth, to the coronal portions of the mandibles.

The mandibles (Plate XLVI, fig. 8, *b*, *c*, *d*) have become thin and foliaceous and completely consolidated with the walls of the oral opening. They have become differentiated, however, into two portions, apparently corresponding to protognath and palpus, or endognath, though these parts are not separated by distinct articulations. The protognathal portion (*b*, fig. 8), corresponding to the entire non-palpigerous mandible of the zoea-stages, is coalesced with the lateral walls of the mouth, except at the broad foliaceous tip, which scarcely projects into the oral opening. The distal margin of this is, however, obscurely dentate, the denticulation varying somewhat in different individuals in the same stage (fig. 8, *b*, and fig. 9), although the mandibles have apparently ceased to perform any of the usual mandibular functions. The endognathal portion (*c*, *d*, fig. 8) evidently represents a palpus with its segments completely coalesced, although the fold between the terminal (*d*) and the outer, spinous portion (*e*) apparently marks the union of the two distal segments.

The distal extremity of the palpus is foliaceous, the margins obtusely rounded and armed with long, plumose setæ, which extend over the oral edge of the labrum and the anterior edge of the mouth itself. The outer margin (*c*), which is thickly spinous in the adult, projects considerably laterally, but is, as yet, armed with only three or four spines.

The first pair of maxillæ (Plate XLVI, fig. 13) have assumed the general form and arrangement of parts which they present in the adult. The inner lobe (*a*, fig. 13) is relatively larger than in the zoea-stages, being considerably broader than the outer lobe; and is armed at the tip, and a little way down the inner margin, with long and stout setæ, most of which are slightly spinulose or plumose distally. The outer lobe (*b*) is long, rather narrow, and the terminal margin is obtusely rounded and armed with setiform spines, which are short and stout toward the inner margin but increase in length outward, and, at the outer margin, grade suddenly into very long and slender, plumose setæ, of which the outermost one is nearly three times as long as the width of the lobe itself. On the inner, lateral margin there are several plumose hairs and on the outer margin, a single short spinule. The palpus (*c*) projects laterally from near the base of the outer lobe as an irregular, sack-like appendage. The articulation at the base of the inner lobe is much less distinct than in the adult, although the articulation at the base of the outer lobe is conspicuous.

The peculiar structure of the mandibles and maxillæ of the adult are more fully discussed in a special note at the end of this article.

The second pair of maxillæ (Plate XLVII, fig. 3) have assumed nearly the adult form. The lobes of the protognath (*a*, *b*, fig. 3) are deeply separated, and a small, supplementary, papiliform lobe (*a'*), tipped with a long, plumose seta, has appeared between the inner and outer lobe, arising near the base of the inner. The terminal margin of the inner lobe is armed with plumose setæ, arranged in three series of varying lengths, the longest, situated upon the very edge, being nearly as long as the lobe itself. The outer lobe is similarly armed, but the setæ are all much shorter and less plumose, and some of them are stout and serrate. Between the outer lobe and the anterior projection of the scaphognath, there is a small, triangular lobe (*d*) apparently representing the endognath. The characteristic fringing of plumose hairs has extended round upon the inner margin of the broad, posterior portion of the scaphognath (*e*), which has changed comparatively little since the last zoea-stage, but still is of

nearly the same form as in the adult, and has apparently fully acquired the adult function.

The three pairs of maxillipeds (Plate XLVII, fig. 4, 6, 7) have assumed so nearly the adult form that detailed descriptions of them are unnecessary.

As in the adult, the anterior lobe of the protognath of the first maxillipeds (Plate XLVII, fig. 4, *a*) is very much elongated; the straight inner margin is thickly beset with plumose setæ throughout, and with an additional series of much longer setæ on the posterior half, and also at the extremity. The endognath seems to be represented only by the long, slender, and soft appendage (*b*), apparently arising from near the base of the inner side of the stout, two jointed exognath (*c*). The basal segment of the exognath is nearly as long as the distal lobe of the protognath, while the terminal segment is somewhat shorter, but fully as broad, and thickly marginal throughout with plumose setæ, which become very long at its extremity. In the adult, the terminal segment is more triangular in outline and the tip is more acute.

The endognath in the second pair of maxillipeds (Plate XLVII, fig. 6, *a*) differs in form slightly from that of the adult, being proportionally stouter and less flattened; the terminal segment, in particular, is proportionally, considerably shorter and consequently more tapering. The basal segment of the exognath (*b*, fig. 6) is not so much narrowed distally, and the oval, terminal segment is a little narrower than in the adult.

The broad, opercular, external maxilliped (Plate XLVII, fig. 7) is transversely truncated at the distal extremity where the palpus articulates, wanting almost wholly the conspicuous, rounded prominence of the anterior angle, just within the articulation, in the adult; and the posterior angle of the inner margin is less prominent and more broadly rounded. The palpus is much less slender, less compressed, and the terminal segment is proportionally shorter than in the adult.

The thoracic legs, like the maxillipeds, are so much like those of the adult, both in form and function, that detailed descriptions of them seem needless here. The anterior pair (Plate XLV, fig. 5) are a little more slender than in the adult, and the terminal segments, or dactyli, are not quite as thickly margined with plumose setæ. The second and third pairs are almost exactly alike and, together with the fourth pair, are specially adapted for burrowing. The fourth pair differ from the second and third principally in having the two

terminal segments more simple in form, the propodus being nearly square, while the dactylus is small and acutely triangular, with the tip obtuse.

The posterior thoracic legs are exceedingly slender and, as in the adult, are in life, usually at least, held concealed within the branchial cavities. They are armed with comparatively few of the very slender spinulose setæ with which they are so abundantly furnished, especially toward the distal extremity, in the adult; and the series of spinulose teeth upon the prehensile edges of the short fingers of the chelæ are very short and composed of only a few individuals upon each finger. There is no doubt that the peculiar position, structure, and armament, of these slender posterior thoracic legs, in *Hippa* and allied genera, specially adapt them for cleaning the branchiæ and branchial chambers of parasites and other foreign bodies. It is worthy of note, in connection with this, that none of the maxillipeds or thoracic legs possess any traces of either exipodal or epipodal branches, some of the last of which perform the office of branchiæ-cleaners in the majority of the *Brachyura* and *Macrura*.

The abdomen (Plate XLVIII, fig. 4), when fully extended, is about as long as the carapax and resembles that of the adult in the form and proportions of the segments, but differs essentially in the structure of the appendages, especially those of the second to the fifth segment, as is usual in the megalops-stage. The first segment,* as seen from above (*a*, fig. 4), is a small plate, with the outline of the

* Latreille and Milne-Edwards have mistaken this small first segment of the abdomen of *Remipes* and *Hippa* for the last segment of the thorax, and I am not aware that the mistake has been corrected by any subsequent author. Latreille (*Genera Crustaceorum et Insectorum*, i, p. 45) simply says of *Remipes testudinarius*, "Cauda segmento baseos aliis multo latiore." But Milne-Edwards (*Histoire naturelle des Crustacés*, ii, pp. 203, 206, 208), after saying in the description of the genus *Albunea*, "Le premier anneau de l'abdomen est petit, et reçu dans une échancrure de la carapace; le second est au contraire très-grand et présente de chaque côté un grand prolongement lamelleux," goes on to say, under the genus *Remipes*, "Le dernier anneau thoracique, qui porte ces appendices [les petites postérieures], est complet en dessus, mobile, et pas recouvert par la carapace, de manière qu'on pourrait facilement le prendre pour le premier segment de l'abdomen. Celui-ci est très-grand, et présente de chaque côté un prolongement lamelleux ovalaire qui chevauche sur la carapace; son bord postérieur est échancré pour loger le second anneau abdominal, qui est ovalaire; * * * * le cinquième et la sixième sont également petits, mais sont sondés entre eux. * * * * Les trois premiers anneaux portent, chez la femelle, des filits ovifères simples." Under *Hippa*, he further says, "Le dernier anneau thoracique n'est pas libre et à découvert comme chez les Remipèdes; mais le premier article de l'abdomen est à peu près de

segment of a circle, filling an arcuate sinus in the posterior margin of the carapax; with which, however, it is not yet consolidated, as in the adult, for it is slightly movable upon the carapax and usually separates from it and remains attached to the second segment when the abdomen is torn away from the cephalo-thorax. As usual in the megalops-stage, and in the larval forms of *Podophthalmia* generally, it is without appendages. The second-segment is about five times as broad as long, three-fourths as broad as the carapax, and nearly twice as broad as the third segment; its great breadth being a result of a broad, lamellar expansion each side. The anterior margin is nearly straight, and the lamellar portion each side slips slightly over the posterior margin of the carapax, when the abdomen is folded beneath the sternum. The lateral margins are very oblique and converge rapidly to the posterior margin, which is no longer than the breadth of the third segment, for the reception of which it is excavated throughout nearly its whole extent; but the lateral expansions do not project so far posteriorly each side of the third segment as they do in the adult. The margins each side are armed with a few stiff setæ. The third segment is about as long as the second, the lateral margins only slightly projecting and rounded and, together with the anterior margin, armed with a few setæ. The fourth and fifth segments are nearly as long as the third, and each successively, very slightly narrower than the one in front of it; they both project very little laterally and are sparsely armed on the lateral and anterior margins with stiff setæ. The sixth segment is nearly as wide as the fifth and only a little wider than long, both the anterior and posterior margins

même forme, et les anneaux suivans présentent aussi la disposition que nous avons déjà remarquée chez ces Crustacés."

Without any reference to the homology of the parts, so clearly shown by their structure and appendages in the megalops-stage, there need be no question in regard to the broad, winged segment in both *Hippa* and *Remipes* being the second, from the fact that it bears the first pair of large ovigerous appendages in the female, and that, if it be the first, that somewhere, two of the succeeding segments must have become completely consolidated into one, although there is no indication, in the segments themselves, of any such consolidation. Quite as conclusive also is the fact that, in the allied genus *Albunea*, while all the segments of the abdomen are perfectly distinct and movable one upon the other, and the anterior ones have nearly the same form as in *Hippa* and *Remipes*, the small first segment is entirely free from the carapax, though fitting closely in a deep sinus in its posterior margin, as described by Milne-Edwards in the passage just quoted. Although this consolidation dorsally of the first segment of the abdomen with the mandibular segment is of rare occurrence in the Malacostraca, it certainly seems no more abnormal than the appearance of the last thoracic segment in the same position, would be.

being considerably curved outward in the middle. The telson is proportionally a little broader than in the adult,—the breadth being rather more than half the length, while in the adult it is usually less,—and much less acute posteriorly and at the antero-lateral angles. The margin is sparsely fringed, nearly all round, with slender setæ or hairs.

The appendages of the second, third, fourth, and fifth segments are similar, though varying considerably in the length of the terminal lamellæ. The peduncles, or protopods, of the appendages of the second segment (Plate XLVIII, fig. 5) are stout and nearly as long as the breadth of the segment itself. The outer lamella, or exopod, (*a*, fig. 5) is nearly as broad as the peduncle, more than half as long, and margined with about twelve slender, plumose setæ (fig. 5*a*), of which the terminal ones are nearly as long as the peduncle, while the others decrease in length as they recede from the tip of the lamella. The inner lamella, or endopod, (*b*, fig. 5) is a small papilli-form appendage, not as long as the diameter of the peduncle, and armed at the tip with a number of very minute appendages, which are apparently only extremely modified spines or setæ. These appendages, as shown in the accompanying illustrations (Plate XLVIII, fig. 6*a*, 6*b*), are composed of a cap, shaped like a concavo-convex lens, supported, on the less convex side, by a short peduncle inserted at the margin and nearly vertical to the under surface of the cap. The diameter of the cap is from .007 to .009^{mm} and the height from the surface of the lamella about the same. In alcoholic specimens at least, these appendages seem to be hard and chitinous, and I am at a loss to understand what purpose they serve in the economy of the animal, unless it be as scrapers for the removal of parasites and other foreign substances from some part of the under surface of the body, in which case it is difficult to see just how they could be applied.

The appendages of the third, fourth, and fifth segments differ from those of the second in the successive shortening of the peduncle, but, more particularly, in the successive increase in the size of the inner lamella, which in the appendages of the fifth segment (Plate XLVIII, fig. 6) have become more than half as long as the outer; the inner lamellæ are similarly armed at the tips, and the outer are all alike margined with long plumose setæ, of which there are usually a very few more on the posterior than on the anterior pairs.

The appendages of the sixth segment (Plate XLVIII, fig. 4) are of nearly the same form and proportion as in the adult. The peduncle, or protopod, is stout, broadest distally, and a little longer than the

segment itself. The terminal lamellæ are nearly alike, each being almost as long as the peduncle, less than half as broad, oval, broadly rounded at the tip, and fringed with slender, plumose setæ, which are very long at the tips of the lamellæ but shorter along the sides. The outer lamella, or exopod, is connected with the peduncle by a very short basal segment.

The young in the megalops-stage were frequently taken at the surface in the towing-net from the 25th of August to the 10th of September. Most of them were taken in the evening, but single individuals were once or twice taken in the day-time. They were also several times obtained directly from specimens in the last zoea-stage kept in aquaria, though the greater part of the zoeæ so kept died during the process of molting. The color of the megalops during life is very much like that of the zoeæ, except that the megalops wants the spots of bright metallic pigment so conspicuous in the zoea.

While in aquaria, the young in the megalops-stage swam rapidly about the walls, at the surface of the water, but not quite as actively as in the zoea-stages. When placed in a shallow dish of sea-water with a little sand heaped up upon one side, they at once, on approaching it, darted for the edge of the miniature sand-beach and backed themselves into it, with an evident satisfaction and with an ease and agility that could not have been excelled by their sand-loving parents after months or years of practice. One, only a few hours after changing from the zoea-stage, evinced the same anxiety to be settled in a home of changing sand. They seemed perfectly content to remain thus buried just at the edge of the water and were never noticed to leave their tiny sand-beach, unless disturbed. If pushed out of the sand and placed on the opposite side of the dish, they swam uneasily about until they found their former retreat and at once plunged themselves backward quite out of sight into it.

On the southern side of Fire Island Beach, Long Island, in September, 1870, I noticed the alacrity with which the young hippas buried themselves in the beach, when they were brought in and left on the wet sand by the waves, and then suspected they were just relinquishing their early pelagic habits, but unfortunately I preserved no specimens at this time. A few days later, the young were found in considerable numbers burrowing in the damp sand in company with species of *Orchestidæ*, just above the edge of the waves; a situation in which I have never since observed either the young or adults.

Three of the specimens, found upon the beach and preserved at this time, prove interesting as they apparently represent a stage next succeeding the megalops. In these specimens the carapax is a little over 4^{mm} in length and the whole integument is of a much firmer consistency than in the megalops-stage. The cephalo-thoracic appendages have all advanced slightly toward the fully mature form. The ocular peduncles have increased in length but are more slender, the diameter at the tips being apparently, absolutely very slightly less than in the megalops-stage. In the antennulæ (Plate XLVIII, fig. 2), the distal prominence upon the under side of the second segment of the peduncle, projects as far as the distal end of the third segment and is more densely fringed with setæ than before. The flagellum has increased only slightly in length and has three or four additional segments, but the secondary flagellum is a third as long as the primary and is composed of four segments, of which the ultimate is as long as the penultimate and antepenultimate together. The oral appendages and thoracic legs have changed comparatively little, although the slender posterior pair of thoracic legs are much more completely armed with setæ and spinules than before.

The first segment of the abdomen has become pretty firmly consolidated with the carapax, though the suture marking the union is still conspicuous and much deeper than in the adult. The expansions on the sides of the second segment are a little broader longitudinally, and the tip and the antero-lateral angles of the telson are more acute than in the megalops. The most marked change, however, has taken place in the appendages of the anterior segments. In all three of the specimens examined, there are no appendages whatever upon either one of the five anterior segments. The sudden and total disappearance of these appendages raises the question, whether the ovigerous appendages of the female are derived directly from the swimming legs of the megalops, or are developed specially, at a later period. It is of course very difficult, if not impossible, to distinguish the sexes, at so early a period, by any of the primary sexual characters; so that the three specimens examined may very likely all be young males; in which case the early disappearance of the appendages might, naturally enough, be expected, as there are no appendages on any of the five anterior segments in the adults of that sex. I have, however, examined several additional specimens of the young, in which the carapax was less than 7^{mm} in length, without, in any case, finding even rudiments of appendages on any of the anterior segments of the abdomen. The smallest individual in which I have been able to detect ovigerous appendages is a specimen, with the

carapax about 7^{mm} in length, from Egmont Key, Florida. In this specimen, the appendages upon the second segment are very small and slender, about .8^{mm} long, unbranched, and composed of three segments—as in the adult female (Plate XLVIII, fig. 16)—of which the terminal one is very short and tipped with a sparsely plumose seta, similar to a single other seta upon the side of the proximal segment; there are no other setæ or hairs discoverable. The appendages of the third segment are similar, but still smaller, being scarcely .5^{mm} long, and composed of two segments only; although the newly formed appendages, as seen through the integument, show indication of division into three segments. No appendages whatever were discoverable upon the fourth segment. As the specimen was about to molt, there is a mere possibility that there may have been very rudimentary appendages upon this segment. In the adult female there are three pairs of unbranched appendages, composed of three segments each, and borne upon the second, third, and fourth segments; no appendages whatever being borne upon the fifth segment. In mature specimens of ordinary size (in which the carapax is about 20^{mm} long), the appendages of the second segment (Plate XLVIII, fig. 16) are about 10^{mm} long; those upon the fourth, about 3.5^{mm}, while those upon the third are intermediate between the two other pairs. Though these observations are by no means conclusive or satisfactory, it seems probable that the egg-bearing appendages of the adult *Hippa* are special developments, and not metamorphosed from the swimming legs of the megalops-stage.

The following table gives, in millimeters, the results of measurements, of the megalops and later stages, side by side for comparison. The measurements in the second column are from one of the three specimens, previously referred to, from Fire Island Beach, Long Island; the others are from specimens from Vineyard Sound.

	Megalops.	Fire I.	Male.	Male.	Female.	Female.
Length of carapax,-----	3.2	4.3	8.3	14.0	13.0	20.6
Breadth of "-----	2.0	2.8	6.2	10.5	9.0	15.4
Length of abdomen,-----	3.0	4.5	9.0	16.0	14.5	25.0
" ocular peduncle,-----	.6	.7	2.4	3.8	3.6	4.6
Diameter of ocular peduncle at tip,-----	.3	.25	.4	.5	.5	.6
Length of antennula,-----	1.0	1.3	3.8	5.6	5.4	7.0
" antenna,-----	3.4	4.8	10.0	19.0	16.0	25.0
" ultimate segment of anterior legs,-----	.8	1.3	3.0	4.1	3.8	5.8
" peduncle of appendage of sixth segment of abdomen,-----	.65	1.0	2.0	3.0	2.8	4.6
" lamellæ of the same,-----	.65	1.0	2.0	2.9	2.8	4.5
" telson,-----	1.7	2.5	5.3	9.2	8.5	14.0
Breadth of "-----	.9	1.3	2.4	4.0	3.9	7.0

A zoea from Zanzibar described briefly by Claus, in his recent work on the development of Crustacea,* and by him doubtfully regarded as one of the early stages of some species of *Albunea*, is certainly the larva of some near ally of *Hippa*, though the figures and description are scarcely sufficiently detailed for a close comparison with the here described zoea-stages of that genus. The zoea described by Claus is apparently in a stage corresponding to that which I have called the third zoea-stage of *Hippa*, but it is already much larger than the zoeæ of the last stage in that genus. As far as can be judged from Claus' imperfect figures and description, the most important character, distinguishing his zoea from that of *Hippa*, is the presence of the exognaths of the external maxillipeds, which form a third pair of swimming appendages somewhat smaller than the two anterior pairs; all the other distinctions seem to be of minor importance. The exognaths of the external maxillipeds are entirely wanting in the adult Hippidæ but are present in the Albanidæ, and their presence in the supposed *Albunea* zoea, coupled with their absence in the latest zoea-stages of *Hippa*, tends strongly to confirm Claus' supposition that his zoea really belongs to the Albanidæ, and most likely to the genus *Albunea*.

From his examination of this so-called zoea of *Albunea*, together with Fritz Müller's note upon the zoea of *Hippa emerita*, previously referred to, Claus concludes that the larval stages of the *Hippa* and *Albunea* group of crustaceans agree much more closely with the zoea-stages of the Brachyura than with the larval stages of the Paguroids or any of the Macrura. The additional evidence, derived from the nearly complete post-embryonal history of *Hippa*, here presented, abundantly confirms this view. There is nothing in the zoea- or megalops-stages of *Hippa* to recall particularly any of the known early stages of the Paguroids, while there is much that forcibly recalls the Brachyuran zoea and megalops. The only conspicuous character distinguishing the *Hippa* and *Albunea* zoea from the ordinary zoeæ of Brachyura, is the absence of a dorsal spine upon the carapax, and this is apparently a distinction of minor importance. Claus has suggested a resemblance between these Hippoid zoeæ and the *Erichthus* larval stage; the motions and color of the living zoeæ of *Hippa* do certainly remind one of the early stages of some of the Stomatopods; but these distant resemblances do not appear to be

* Untersuchungen zur Erforschung der Genealogischen Grundlage des Crustaceen-Systems, Wein, 1876, p. 59, plate ix, figs. 1-10.

the result of any structural similarity. In fact the full series of the early stages of *Hippa* shows conclusively that the young undergo an essentially Brachyural, zoea and megalops, post-embryonal development, and it furnishes an important addition to the evidence tending to confirm the view that the Anomoura are a heterogeneous group, made up, probably, of outlying Brachyural and Macrural families.

Note on the Structure of the Mandibles and Maxilla in Hippa and Remipes.

In the description of the megalops-stage, I have already alluded to the remarkable structure of the mandibles and the first pair of maxillæ in the adult *Hippa*, but these appendages in *Hippa* and the allied genera seem worthy of more special attention, since they furnish important characters for distinguishing the Hippidæ and Alburnidæ. Although the oral appendages of *Alburnea*, and apparently also of the allied genera, *Blepharopoda* Randall and *Lepidopa* Stimpson, have been correctly described by all the later authorities, the structure and homology of the parts of the mandibles and the first pair of maxillæ in *Hippa* and *Remipes*, though very important in a taxonomic point of view, seem to have been misunderstood by all carcinologists who have written upon these genera.

Milne-Edwards, in his great general work on crustaceans, appears to have been the first to describe, with care, the oral appendages of any of the Hippidæ or Alburnidæ, and the errors, into which he fell in describing *Hippa* and *Remipes*, have been perpetuated by succeeding writers. To be sure, J. C. Fabricius had before described the parts about the mouth, in his original description of the genus *Hippa* (Supplementum Entomologiæ Systematicæ, p. 329), but his account of the inner appendages is so confused that I am quite unable to understand what parts he had in view as the mandible and its palpus. He says, "*Mandibula* brevis, cornea, fornicata, obtusa, dorso palpigera. *Palpus* setaceus, tenuis, uti videtur multiarticulatus," which could scarcely be made to apply to the parts of the mandible as they really exist, either in *Hippa* or *Remipes*. Milne-Edwards, in his description of the genus *Remipes* (Histoire naturelle des Crustacés, ii, p. 205), says, "Les mâchoires de la seconde paire ne présentent rien de bien remarquable; celles de la première paire sont très-petites, et refoulées en avant, entre la mandibule et la lèvre supérieure, qui est très-grande et fort saillante. Enfin la mandibule, qui est fortement dentelée, porte un palpe composé de deux petits articles lamelleux, séparés du corps de la mandibule par un grand sillon membraneux." This strongly dentate "mandible"

is evidently the terminal portion of the inner lobe of the first maxilla (Plate XLVII, fig. 10, *a*); but its "palpus composed of two small, lamellar segments" is not so easily interpreted, though it may possibly be the real mandible itself. In the figures of *Remipes testudinarius* given in Milne-Edwards' volume on the Crustacea in the third edition of *Le Règne animal de Cuvier*, figure 1^d, plate 42, is simply the inner lobe of the first maxilla, which is entirely wanting in the otherwise correct figure of the maxilla itself; but the only allusion to the mandibular palpus is in the explanation of figure, "1^d, Mandibules (ces organes sont dépourvus d'appendice palpiforme)." In Milne-Edwards' description of the genus *Hippa* (*Histoire naturelle des Crustacés*, ii, p. 207) no reference is made to the maxillæ or mandibles, and no part of either of these appendages is given in the figures of *H. emerita* in *Le Règne animal de Cuvier*.

De Haan's figures of mouth appendages of *Remipes* (*Fauna Japonica*, Plate Q) are like those of Milne-Edwards just referred to, except that he has left the basal portion of the inner lobe of the first maxilla properly attached in place to answer for the inner lobe which he describes as the "lacina interna brevissima" (op. cit., p. 201). De Haan however failed to find the mandibular palpus, for he distinctly says in the description of these maxillary mandibles, "palpis nullis." In regard to *Hippa*, De Haan seems to have been more fortunate, for, in his description of the genus (op. cit., p. 201), after correctly describing the maxillæ, he says, "Mandibulæ minutissimæ, cartilagineæ, corona integra, palpis membranaceis triarticularis." His figure of the mandible of *Hippa Asiatica* (op. cit., plate Q; copied as figure 14, Plate XLVI, of the accompanying illustrations), although evidently drawn from the real mandible, is apparently very incorrect.

The following descriptions and figures of the mandibles and maxillæ in *Hippa* and *Remipes* are taken from *H. talpoida* Say, and *R. Pacificus* Dana.

Hippa.—The form and position of the labrum, labium, and mandibles, as seen from beneath, are shown in figure 10 on Plate XLVI. The mouth is a narrow, longitudinal opening bounded, each side and posteriorly, by a margin raised above the general surface of the buccal area, and in front, by the labrum (*a*, figure 10), which projects anteriorly as a triangular, lamelliform plate above the epistome, from which, however, it is separated by a considerable space. Beneath the labrum, the margin of the epistome is reflected and projects forward each side into a slender, lamelliform process (*b*) tipped with

very long, plumose setae extending along each side of the labrum. At the sides of the oral opening are the mandibles (*c*, *d*, *e*), the tips of their palpi (*e*) approaching each other just in front of it. The mesial side of the basal portion of the mandible is consolidated with and forms a part of the raised lateral margin of the oral opening, while the outer side is continuous with and consolidated to the buccal walls each side. The palpus arises from this basal portion of the mandible and inwraps the posterior part of the margin of the labrum by means of an oblique fold, which distinctly separates it into an outer (*d*) and an inner (*e*) portion. This oblique fold is most marked at the anterior extremity of the palpus, where it presents a sigmoid curve, the thickening of the palpus posteriorly rendering it less distinct though well seen in a transverse section, as in figure 10*a*. The free margin (*d*, figure 10) of this outer portion is thickly armed with a series of stout spinules, while the inner portion projects above the posterior part of the labrum, as a broad, oval-pointed, lamelliform lobe, the edges of which are thickly beset with slender setae. The coronal portion (*c*) of each mandible is a broad, obtusely rounded, thin lamella, the free terminal part of which lies just above the palpus. Each lamella of the deeply bifid labium (*f*) lies just above the mandible each side and closely infolds the margin of the oral opening.

The protognathal portion of the first maxilla (Plate XLVII, fig. 8) is divided into two deeply separated lobes and fits closely upon the labium and mandible. The somewhat oval, terminal portion of the posterior, or inner, lobe (*a*, fig. 8) lies just over the lobes of the labium each side, and the whole length of its nearly straight, inner margin is bent downward into the posterior part of the oral opening, and is edged with a double series of long, serrate setae; while the outer convex surface is armed with a scattered, double line of minute spinules. This terminal portion of the lobe is free all round, except near the outer, posterior angle where it is connected with the buccal walls, and where a thickened, rod-like portion connects it, by an imperfect articulation, with the base of the anterior lobe. The anterior lobe (*b*) is very long and narrow, and its oblique, terminal margin lies just over the inner portion of the mandibular palpus, and is thickly armed with short, setiform spinules, while the lateral margins are thickly clothed with slender, plumose setae, of which those at the anterior extremity of the outer margin are very much the longest. The palpus (*c*) is soft and sack-like, arises from near the base of the anterior lobe, and projects laterally into the efferent passages of the branchial chamber.

The second pair of maxillæ offer no marked peculiarity and are essentially of the same form as in the megalops-stage (Plate XLVII, fig. 3). The two principal lobes of the protognath are very similar in form and structure to, lie directly over, and doubtless act in direct conjunction with, the two corresponding lobes of the first maxilla.

Remipes.—The general structure and arrangement of the parts of the mandibles and both pairs of maxillæ is the same as in *Hippa*, but there are important differences in the details of the form and structure of special parts, particularly in the posterior lobe of the protognath in the first pair of maxillæ.

The mandibles (Plate XLVII, fig. 9) are consolidated with and closely invest the walls of the oral opening as in *Hippa*, but the protognathal portion (*a*) is farther removed from the opening, is even more rudimentary, its margins are unarmed, and it apparently serves only as a base for the attachment of the palpus. The free margin (*b*) of the outer portion of the palpus projects in an obtuse angle in front and is armed with a few, very stout, short, and obtuse spines. The inner portion (*c*), lying just above the labrum, is somewhat orbicular, and its terminal margin is armed with numerous, exceedingly long setæ, of which the longest are plumose.

The terminal portion of the posterior lobe in the first pair of maxillæ (Plate XLVII, fig. 10, *a*) is a thick, heavily calcified, rectilinear plate united, by a broad articulation, with the basal, rod-like portion and at the posterior angle, directly with the buccal walls. Its terminal margin is armed externally with a series of five, very large, short, stout, conical, dentiform spines, and at the angles at each end of the series with a similar, but very much smaller spine. Upon the short posterior margin there are several still smaller spines of the same character. The terminal margin beneath is thickly beset with stout, plumose setæ projecting inward. The anterior lobe (*b*) is long, very narrow to near the expanded, spatulate extremity, the convex terminal margin of which is densely armed with setæ which are stout and spiniform, except anteriorly where they are long and more slender. The palpus (*c*) is nearly the same as in *Hippa*, but a little shorter and thicker.

The different parts of the second pair of maxillæ are arranged in the same way as in the *Hippa*, but the posterior lobe of the protognath is proportionally much smaller and its edge is armed with very long, plumose setæ, which are of nearly uniform thickness to very near the tips; while the anterior lobe is spatulate in form and thickly armed with similar, but somewhat shorter, setæ.

EXPLANATION OF PLATES.

PLATE XLV.

- Figure 1.—Zoea in the second stage, ventral view, with the abdomen bent forward beneath the cephalo-thorax so as to show the dorsal side of the terminal portion of the abdomen, enlarged twenty-four diameters.
- Figure 2.—Zoea in the last stage, dorsal view, enlarged twelve diameters.
- Figure 3.—The same, in the same position as in figure 1, ventral view, the natatory limbs omitted, enlarged twelve diameters.
- Figure 4.—The same, lateral view, enlarged twelve diameters.
- Figure 5.—Megalops just changed from the last zoea-stage; dorsal view of anterior portion of the animal showing the eyes, antennulæ, antennæ, and the extremities of the first pair of thoracic legs, enlarged twenty-four diameters. The antennæ are shown fully extruded and the whole animal under slight compression.
- Figure 6.—One of the segments of the flagellum of an antenna, megalops-stage, seen in transverse section to show the arrangement of the setæ, enlarged forty diameters.

PLATE XLVI.

- Figure 1.—Antennula, second zoea-stage, enlarged seventy-five diameters.
- Figure 2.—Antenna of the right side seen from beneath, second zoea-stage, enlarged seventy-five diameters; *a*, *b*, dentiform processes; *c*, flagellum.
- Figure 3.—The same, third zoea-stage, enlarged seventy-five diameters; *a*, *b*, dentiform processes; *c*, rudimentary flagellum.
- Figure 4.—The same, last zoea-stage, enlarged forty diameters; *a*, *b*, dentiform processes; *c*, rudimentary flagellum showing the multiarticulate flagellum of the megalops-stage already formed within.
- Figure 5.—Antennule and inner mouth appendages seen from beneath, last zoea-stage, enlarged thirty-six diameters; *a*, *a*, antennulæ; *b*, labrum; *c*, mandible; *d*, labium; *e*, first maxilla.
- Figure 6.—Tips of the mandibles, last zoea-stage, enlarged 150 diameters.
- Figure 7.—Peduncle of the antenna seen from beneath, megalops-stage, enlarged twelve diameters; *a*, *b*, *c*, *d*, *e*, segments of peduncets; *f*, supplementary rod or plate.
- Figure 8.—Labrum, labium and mandibles seen in place from beneath, megalops-stage, enlarged seventy-five diameters; *a*, labrum; *b*, coronal portion of mandible; *c*, *d*, mandibular palpus; *e*, labium.
- Figure 9.—Coronal portion of the mandible of another specimen in the same stage, enlarged 150 diameters.
- Figure 10.—Labrum, labium, and mandibles seen in place from beneath, adult female, enlarged twenty-four diameters; *a*, labrum; *b*, setæ-bearing tubercle above the margin of the labrum; *c*, coronal portion of the mandible; *d*, lateral spinous margin of the basal portion of the mandibular palpus; *e*, terminal portion of the palpus; *f*, labium; *g*, *h*, line of the section shown in figure 10*a*. The letter *h*, is directly over the anterior extremity of the elongated mouth opening. 10*a*, outline of a transverse section of the mandible through the line *gh* in figure 10; the dotted lines representing the anterior edges of the appendage projected upon the section; *c*, tip of the coronal portion of the mandible; *g*, lateral spinous margin of the palpus; *h*, inner extremity of the terminal portion of the palpus.
- Figure 11.—First maxilla of the left side seen from beneath, second zoea-stage, enlarged seventy-two diameters; *a*, inner lobe; *b*, outer lobe; *c*, palpus.
- Figure 12.—Same, last zoea-stage, enlarged seventy-two diameters.
- Figure 13.—Same, megalops-stage, enlarged seventy-five diameters.
- Figure 14.—Mandible of *Hippa Asiatica*, copied from DeHaan, Fauna Japonica, Crustacea, Plate Q.

PLATE XLVII.

- Figure 1.—Second maxilla of the left side seen from beneath, second zoea-stage, enlarged thirty-six diameters; *a* protognath; *b*, scaphognath.
- Figure 2.—Same, last zoea-stage, enlarged thirty-six diameters; *a*, *b*, protognath; *c*, scaphognath.

- Figure 3.—Same, megalops-stage, enlarged forty diameters; *a*, *a'*, *b*, lobes of the protognath; *d*, endognath; *c*, scaphognath.
- Figure 4.—First maxilliped of the left side seen from beneath, megalops-stage, enlarged forty diameters; *a*, protognath; *b*, endognath; *c*, exognath.
- Figure 5.—Second maxilliped of the left side seen from behind, last zoea-stage, enlarged forty diameters; *a*, protognath; *b*, endognath, *c*, natatory exognath.
- Figure 6.—Second maxilliped of the left side seen from beneath, megalops-stage, enlarged forty diameters; *a*, endognath; *b*, exognath.
- Figure 7.—Third (external) maxilliped of the left side seen from beneath, megalops-stage, enlarged twenty-four diameters.
- Figure 8.—First maxilla of the right side seen from beneath, adult female *Hippa*, enlarged twelve diameters; *a*, inner lobe; *b*, outer lobe; *c*, palpus.
- Figure 9.—*Remipes Pacificus*. Mandible of the right side seen from beneath, adult female, enlarged twenty-four diameters; *a*, basal part of the mandible itself (protognath); *b*, lateral spinous margin of the basal portion of the palpus; *c*, terminal portion of the palpus.
- Figure 10.—First maxilla of the right side, from the same specimen as the last figure, enlarged twelve diameters; *a*, inner lobe; *b*, outer lobe; *c*, palpus.

PLATE XLVIII.

- Figure 1.—Antennula, seen from the side, megalops-stage, enlarged twenty-four diameters; *a*, *b*, *c*, first, second, and third segments of the peduncle; *d*, primary flagellum; *e*, rudimentary secondary flagellum.
- Figure 2.—Same, from a specimen about 5^{mm} in length, enlarged twenty-four diameters.
- Figure 3.—Same, adult female, enlarged six diameters.
- Figure 4.—Abdomen, megalops-stage, dorsal view, enlarged twelve diameters; *a*, first segment.
- Figure 5.—One of the swimming legs of the second segment of the abdomen, megalops-stage, enlarged twenty-four diameters; *a*, outer lamella, or exopod; *b*, inner lamella, or endopod. *5a*, portion of one of the plumose setæ, more enlarged.
- Figure 6.—One of the swimming legs of the fifth segment of the abdomen of the same specimen, enlarged twenty-four diameters; *a*, outer lamella; *b*, inner lamella. *6a*, *6b*, side and front view of one of the peculiar appendages with which the tip of the inner lamella is armed, enlarged 500 diameters.
- Figure 7.—Telson of adult male, dorsal view, enlarged two diameters.
- Figure 8.—Same, adult female, enlarged two diameters.
- Figure 9.—Rudimentary thoracic legs and third maxilliped of the left side, last zoea-stage, enlarged twelve diameters; *a*, base of second maxillipeds; *b*, third maxilliped; *c*, *d*, *e*, *f*, *g*, thoracic legs; *h*, branchiæ. The appendages are represented as seen from the inner (mesial) side and under slight compression so as to show the extremities which are curled in and hidden from the outside; in consequence, the branchiæ are represented diagrammatically.
- Figure 10.—Appendage of the right side of the sixth segment of the abdomen, second zoea-stage, dorsal view, enlarged forty diameters; *a*, basal portion; *b*, outer lamella. *10a*, tip of lamella with the bases of the terminal setæ, enlarged 175 diameters.
- Figure 11.—Corresponding appendage, third zoea-stage, enlarged forty diameters; *a*, base; *b*, outer lamella; *c*, inner lamella.
- Figure 12.—Corresponding appendage, last zoea-stage, enlarged forty diameters.
- Figure 13.—Posterior margin of telson, second zoea-stage, enlarged 100 diameters. In this and the next two figures the arrow indicates the median line of the telson, the margin on the left of it being exactly the reverse of that shown upon the right.
- Figure 14.—The same, from another individual in the same stage, showing a single median spine in place of the two median spines and denticle in other specimens.
- Figure 15.—Posterior margin of telson, last zoea-stage, enlarged 100 diameters. *15a*, two of the denticles from the margin between the spines, enlarged 300 diameters.
- Figure 16.—Ovigerous appendage of the second abdominal segment of an adult female, enlarged four diameters.

IX. ON THE EQUILIBRIUM OF HETEROGENEOUS SUBSTANCES.

BY J. WILLARD GIBBS.

(Continued from page 248).

THE CONDITIONS OF INTERNAL AND EXTERNAL EQUILIBRIUM FOR SOLIDS IN CONTACT WITH FLUIDS WITH REGARD TO ALL POSSIBLE STATES OF STRAIN OF THE SOLIDS.

IN treating of the physical properties of a solid, it is necessary to consider its *state of strain*. A body is said to be *strained* when the relative position of its parts is altered, and by its *state of strain* is meant its state in respect to the relative position of its parts. We have hitherto considered the equilibrium of solids only in the case in which their state of strain is determined by pressures having the same values in all directions about any point. Let us now consider the subject without this limitation.

If x', y', z' are the rectangular co-ordinates of a point of a solid body in any completely determined state of strain, which we shall call the *state of reference*, and x, y, z , the rectangular co-ordinates of the same point of the body in the state in which its properties are the subject of discussion, we may regard x, y, z as functions of x', y', z' , the form of the functions determining the second state of strain. For brevity, we may sometimes distinguish the variable state, to which x, y, z relate, and the constant state (state of reference), to which x', y', z' relate, as the *strained* and the *unstrained* states; but it must be remembered that these terms have reference merely to the change of form or *strain* determined by the functions which express the relations of x, y, z and x', y', z' , and do not imply any particular physical properties in either of the two states, nor prevent their possible coincidence. The axes to which the co-ordinates x, y, z , and x', y', z' relate will be distinguished as the axes of X, Y, Z , and X', Y', Z' . It is not necessary, nor always convenient, to regard these systems of axes as identical, but they should be similar, i. e., capable of superposition.

The state of strain of any element of the body is determined by the values of the differential coefficients of x, y , and z with respect to x', y' , and z' ; for changes in the values of x, y, z , when the differential coefficients remain the same, only cause motions of translation of the

body. When the differential coefficients of the first order do not vary sensibly except for distances greater than the radius of sensible molecular action, we may regard them as completely determining the state of strain of any element. There are nine of these differential coefficients, viz.,

$$\left. \begin{array}{ccc} \frac{dx}{dx'} & \frac{dx}{dy'} & \frac{dx}{dz'} \\ \frac{dy}{dx'} & \frac{dy}{dy'} & \frac{dy}{dz'} \\ \frac{dz}{dx'} & \frac{dz}{dy'} & \frac{dz}{dz'} \end{array} \right\} \quad (354)$$

It will be observed that these quantities determine the orientation of the element as well as its strain, and both these particulars must be given in order to determine the nine differential coefficients. Therefore, since the orientation is capable of three independent variations, which do not affect the strain, the strain of the element, considered without regard to directions in space, must be capable of six independent variations.

The physical state of any given element of a solid in any unvarying state of strain is capable of one variation, which is produced by addition or subtraction of heat. If we write ε_v , and η_v , for the energy and entropy of the element divided by its volume in the state of reference, we shall have for any constant state of strain

$$\delta \varepsilon_v = t \delta \eta_v.$$

But if the strain varies, we may consider ε_v , as a function of η_v , and the nine quantities in (354), and may write

$$\begin{aligned} \delta \varepsilon_v = & t \delta \eta_v + X_{x'} \delta \frac{dx}{dx'} + X_{y'} \delta \frac{dx}{dy'} + X_{z'} \delta \frac{dx}{dz'} \\ & + Y_{x'} \delta \frac{dy}{dx'} + Y_{y'} \delta \frac{dy}{dy'} + Y_{z'} \delta \frac{dy}{dz'} \\ & + Z_{x'} \delta \frac{dz}{dx'} + Z_{y'} \delta \frac{dz}{dy'} + Z_{z'} \delta \frac{dz}{dz'}, \end{aligned} \quad (355)$$

where $X_{x'}$, . . . $Z_{z'}$, denote the differential coefficients of ε_v , taken with respect to $\frac{dx}{dx'}$, . . . $\frac{dz}{dz'}$. The physical signification of these quantities will be apparent, if we apply the formula to an element which in the state of reference is a right parallelepiped having the edges dx' , dy' , dz' , and suppose that in the strained state the face in which x' has the smaller constant value remains fixed, while the opposite face is moved parallel to the axis of X . If we also suppose

no heat to be imparted to the element, we shall have, on multiplying by $dx' dy' dz'$,

$$\delta \varepsilon_v, dx' dy' dz' = X_x, \delta \frac{dx}{dx'} dx' dy' dz'.$$

Now the first member of this equation evidently represents the work done upon the element by the surrounding elements; the second member must therefore have the same value. Since we must regard the forces acting on opposite faces of the elementary parallelepiped as equal and opposite, the whole work done will be zero except for the face which moves parallel to X . And since $\delta \frac{dx}{dx'}$ represents the distance moved by this face, $X_x, dy' dz'$ must be equal to the component parallel to X of the force acting upon this face. In general, therefore, if by the positive side of a surface for which x' is constant we understand the side on which x' has the greater value, we may say that $X_x,$ denotes the component parallel to X of the force exerted by the matter on the positive side of a surface for which x' is constant upon the matter on the negative side of that surface per unit of the surface measured in the state of reference. The same may be said, *mutatis mutandis*, of the other symbols of the same type.

It will be convenient to use Σ and Σ' to denote summation with respect to quantities relating to the axes X, Y, Z , and to the axes X', Y', Z' , respectively. With this understanding we may write

$$\delta \varepsilon_v = t \delta \eta_v + \Sigma \Sigma' \left(X_x, \delta \frac{dx}{dx'} \right). \tag{356}$$

This is the complete value of the variation of ε_v for a given element of the solid. If we multiply by $dx' dy' dz'$, and take the integral for the whole body, we shall obtain the value of the variation of the total energy of the body, when this is supposed invariable in substance. But if we suppose the body to be increased or diminished in substance at its surface (the increment being continuous in nature and state with the part of the body to which it is joined), to obtain the complete value of the variation of the energy of the body, we must add the integral

$$\int \varepsilon_v, \delta N' Ds'$$

in which Ds' denotes an element of the surface measured in the state of reference, and $\delta N'$ the change in position of this surface (due to the substance added or taken away) measured normally and outward in the state of reference. The complete value of the variation of the intrinsic energy of the solid is therefore

$$\int \int \int t \delta \eta_v dx' dy' dz' + \int \int \int \Sigma \Sigma' \left(X_x \delta \frac{dx'}{dx'} \right) dx' dy' dz' + \int \epsilon_v \delta N' Ds'. \quad (357)$$

This is entirely independent of any supposition in regard to the homogeneity of the solid.

To obtain the conditions of equilibrium for solid and fluid masses in contact, we should make the variation of the energy of the whole equal to or greater than zero. But since we have already examined the conditions of equilibrium for fluids, we need here only seek the conditions of equilibrium for the interior of a solid mass and for the surfaces where it comes in contact with fluids. For this it will be necessary to consider the variations of the energy of the fluids only so far as they are immediately connected with the changes in the solid. We may suppose the solid with so much of the fluid as is in close proximity to it to be enclosed in a fixed envelop, which is impermeable to matter and to heat, and to which the solid is firmly attached wherever they meet. We may also suppose that in the narrow space or spaces between the solid and the envelop, which are filled with fluid, there is no motion of matter or transmission of heat across any surfaces which can be generated by moving normals to the surface of the solid, since the terms in the condition of equilibrium relating to such processes may be cancelled on account of the internal equilibrium of the fluids. It will be observed that this method is perfectly applicable to the case in which a fluid mass is entirely enclosed in a solid. A detached portion of the envelop will then be necessary to separate the great mass of the fluid from the small portion adjacent to the solid, which alone we have to consider. Now the variation of the energy of the fluid mass will be, by equation (13),

$$\int^F t \delta D\eta - \int^F p \delta Dv + \Sigma_1 \int^F \mu_1 \delta Dm_1, \quad (358)$$

where \int^F denotes an integration extending over all the elements of the fluid (within the envelop), and Σ_1 denotes a summation with regard to those independently variable components of the fluid of which the solid is composed. Where the solid does not consist of substances which are components, actual or possible (see page 117), of the fluid, this term is of course to be cancelled.

If we wish to take account of gravity, we may suppose that it acts in the negative direction of the axis of Z . It is evident that the variation of the energy due to gravity for the whole mass considered is simply

$$\int \int \int g I' \delta z dx' dy' dz', \quad (359)$$

where g denotes the force of gravity, and I' the density of the

element in the state of reference, and the triple integration, as before, extends throughout the solid.

We have, then, for the general condition of equilibrium,

$$\begin{aligned} f'f't \delta \eta_v, dx' dy' dz' + f'f'f' \sum' \left(X_{x'} \delta \frac{dx'}{dx'} \right) dx' dy' dz' \\ + f'f'f'g I' \delta z dx' dy' dz' + f' \varepsilon_v, \delta N' Ds' \\ + f'^F t \delta D\eta - f'^F p' \delta Dv + \sum_1 f'^F \mu_1 \delta Dm_1 \geq 0. \end{aligned} \quad (360)$$

The equations of condition to which these variations are subject are:

(1) that which expresses the constancy of the total entropy,

$$f'f'f' \delta \eta_v, dx' dy' dz' + f' \eta_v, \delta N' Ds' + f'^F \delta D\eta = 0; \quad (361)$$

(2) that which expresses how the value of δDv for any element of the fluid is determined by changes in the solid,

$$\delta Dv = -(\alpha \delta x + \beta \delta y + \gamma \delta z) Ds - v_v, \delta N' Ds', \quad (362)$$

where α, β, γ denote the direction cosines of the normal to the surface of the body in the state to which x, y, z relate, Ds the element of the surface in this state corresponding to Ds' in the state of reference, and v_v , the volume of an element of the solid divided by its volume in the state of reference;

(3) those which express how the values of $\delta Dm_1, \delta Dm_2$, etc. for any element of the fluid are determined by the changes in the solid,

$$\left. \begin{aligned} \delta Dm_1 &= -I'_1 \delta N' Ds', \\ \delta Dm_2 &= -I'_2 \delta N' Ds', \\ \text{etc.,} \end{aligned} \right\} \quad (363)$$

where I'_1, I'_2 , etc. denote the separate densities of the several components in the solid in the state of reference.

Now, since the variations of entropy are independent of all the other variations, the condition of equilibrium (360), considered with regard to the equation of condition (361), evidently requires that throughout the whole system

$$t = \text{const.} \quad (364)$$

We may therefore use (361) to eliminate the first and fifth integrals from (360). If we multiply (362) by p , and take the integrals for the whole surface of the solid and for the fluid in contact with it, we obtain the equation

$$f'^F p \delta Dv = -f'p (\alpha \delta x + \beta \delta y + \gamma \delta z) Ds - f'p v_v, \delta N' Ds', \quad (365)$$

by means of which we may eliminate the sixth integral from (360).

If we add equations (363) multiplied respectively by μ_1, μ_2 , etc., and take the integrals, we obtain the equation

$$\sum_1 f^F \mu_1 \delta Dm_1 = -f \sum_1 (\mu_1 \Gamma_1') \delta N' Ds', \quad (366)$$

by means of which we may eliminate the last integral from (360).

The condition of equilibrium is thus reduced to the form

$$\begin{aligned} f' f' f' \sum \sum' \left(X_{x'} \delta \frac{dx'}{dx'} \right) dx' dy' dz' + f' f' f' g \Gamma' \delta z dx' dy' dz' \\ + f' \varepsilon_{v'} \delta_2 N' Ds' - f' t \eta_{v'} \delta N' Ds' + f' p (\alpha \delta x + \beta \delta y + \gamma \delta z) Ds \\ + f' p v_{v'} \delta N' Ds' - f' \sum_1 (\mu_1 \Gamma_1') \delta N' Ds' \geq 0, \quad (367) \end{aligned}$$

in which the variations are independent of the equations of condition, and in which the only quantities relating to the fluids are p and μ_1 , μ_2 , etc.

Now by the ordinary method of the calculus of variations, if we write α' , β' , γ' for the direction-cosines of the normal to the surface of the solid in the state of reference, we have

$$\begin{aligned} f' f' f' X_{x'} \delta \frac{dx'}{dx'} dx' dy' dz' \\ = f' \alpha' X_{x'} \delta x Ds' - f' f' f' \frac{dX_{x'}}{dx'} \delta x dx' dy' dz', \quad (368) \end{aligned}$$

with similar expressions for the other parts into which the first integral in (367) may be divided. The condition of equilibrium is thus reduced to the form

$$\begin{aligned} -f' f' f' \sum \sum' \left(\frac{dX_{x'}}{dx'} \delta x \right) dx' dy' dz' + f' f' f' g \Gamma' \delta z dx' dy' dz' \\ + f' \sum \sum' (\alpha' X_{x'} \delta x) Ds' + f' p \sum (\alpha \delta x) Ds \\ + f' [\varepsilon_{v'} - t \eta_{v'} + p v_{v'} - \sum_1 (\mu_1 \Gamma_1')] \delta N' Ds' \geq 0. \quad (369) \end{aligned}$$

It must be observed that if the solid mass is not continuous throughout in nature and state, the surface-integral in (368), and therefore the first surface-integral in (369), must be taken to apply not only to the external surface of the solid, but also to every surface of discontinuity within it, and that with reference to each of the two masses separated by the surface. To satisfy the condition of equilibrium, as thus understood, it is necessary and sufficient that throughout the solid mass

$$\sum \sum' \left(\frac{dX_{x'}}{dx'} \delta x \right) - g \Gamma' \delta z = 0; \quad (370)$$

that throughout the surfaces where the solid meets the fluid

$$Ds' \sum \sum' (\alpha' X_{x'} \delta x) + Ds p \sum (\alpha \delta x) = 0, \quad (371)$$

and

$$[\varepsilon_{v'} - t \eta_{v'} + p v_{v'} - \sum_1 (\mu_1 \Gamma_1')] \delta N' \geq 0; \quad (372)$$

and that throughout the internal surfaces of discontinuity

$$\sum \sum' (\alpha' X_x, \delta x)_1 + \sum \sum' (\alpha' X_x, \delta x)_2 = 0, \tag{373}$$

where the suffixed numerals distinguish the expressions relating to the masses on opposite sides of a surface of discontinuity.

Equation (370) expresses the mechanical conditions of internal equilibrium for a continuous solid under the influence of gravity. If we expand the first term, and set the coefficients of δx , δy , and δz separately equal to zero, we obtain

$$\left. \begin{aligned} \frac{dX_{x'}}{dx'} + \frac{dX_{y'}}{dy'} + \frac{dX_{z'}}{dz'} &= 0, \\ \frac{dY_{x'}}{dx'} + \frac{dY_{y'}}{dy'} + \frac{dY_{z'}}{dz'} &= 0, \\ \frac{dZ_{x'}}{dx'} + \frac{dZ_{y'}}{dy'} + \frac{dZ_{z'}}{dz'} &= g I'. \end{aligned} \right\} \tag{374}$$

The first member of any one of these equations multiplied by $dx' dy' dz'$ evidently represents the sum of the components parallel to one of the axes X, Y, Z of the forces exerted on the six faces of the element $dx' dy' dz'$ by the neighboring elements.

As the state which we have called the state of reference is arbitrary, it may be convenient for some purposes to make it coincide with the state to which x, y, z relate, and the axes X', Y', Z' with the axes X, Y, Z . The values of X_x, \dots, Z_z , on this particular supposition may be represented by the symbols X_x, \dots, Z_z . Since

$$X_{y'} = \frac{d\varepsilon_{y'}}{d\frac{dx'}{dy'}}, \quad \text{and} \quad Y_{x'} = \frac{d\varepsilon_{x'}}{d\frac{dy'}{dx'}}$$

and since, when the states x, y, z and x', y', z' coincide, and the axes X, Y, Z , and X', Y', Z' , $d\frac{dx'}{dy'}$ and $d\frac{dy'}{dx'}$ represent displacements which differ only by a rotation, we must have

$$X_y = Y_x, \tag{375}$$

and for similar reasons,

$$Y_z = Z_y, \quad Z_x = X_z. \tag{376}$$

The six quantities X_x, Y_y, Z_z, X_y or Y_x, Y_z or Z_y , and Z_x or X_z are called the *rectangular components of stress*, the three first being the *longitudinal stresses* and the three last the *shearing stresses*. The mechanical conditions of internal equilibrium for a solid under the influence of gravity may therefore be expressed by the equations

$$\left. \begin{aligned} \frac{dX_x}{dx} + \frac{dX_y}{dy} + \frac{dX_z}{dz} &= 0, \\ \frac{dY_x}{dx} + \frac{dY_y}{dy} + \frac{dY_z}{dz} &= 0, \\ \frac{dZ_x}{dx} + \frac{dZ_y}{dy} + \frac{dZ_z}{dz} &= gI, \end{aligned} \right\} \quad (377)$$

where I denotes the density of the element to which the other symbols relate. Equations (375), (376) are rather to be regarded as expressing necessary relations (when X_x, \dots, Z_z are regarded as internal forces determined by the state of strain of the solid) than as expressing conditions of equilibrium. They will hold true of a solid which is not in equilibrium,—of one, for example, through which vibrations are propagated,—which is not the case with equations (377).

Equation (373) expresses the mechanical conditions of equilibrium for a surface of discontinuity within the solid. If we set the coefficients of $\delta x, \delta y, \delta z$, separately equal to zero we obtain

$$\left. \begin{aligned} (\alpha' X_x + \beta' X_y + \gamma' X_z)_1 + (\alpha' X_x + \beta' X_y + \gamma' X_z)_2 &= 0, \\ (\alpha' Y_x + \beta' Y_y + \gamma' Y_z)_1 + (\alpha' Y_x + \beta' Y_y + \gamma' Y_z)_2 &= 0, \\ (\alpha' Z_x + \beta' Z_y + \gamma' Z_z)_1 + (\alpha' Z_x + \beta' Z_y + \gamma' Z_z)_2 &= 0. \end{aligned} \right\} \quad (378)$$

Now when the α', β', γ' represent the direction-cosines of the normal in the state of reference on the positive side of any surface within the solid, an expression of the form

$$\alpha' X_x + \beta' X_y + \gamma' X_z, \quad (379)$$

represents the component parallel to X of the force exerted upon the surface in the strained state by the matter on the positive side per unit of area measured in the state of reference. This is evident from the consideration that in estimating the force upon any surface we may substitute for the given surface a broken one consisting of elements for each of which either x' or y' or z' is constant. Applied to a surface bounding a solid, or any portion of a solid which may not be continuous with the rest, when the normal is drawn outward as usual, the same expression taken negatively represents the component parallel to X of the force exerted upon the surface (per unit of surface measured in the state of reference) by the interior of the solid, or of the portion considered. Equations (378) therefore express the condition that the force exerted upon the surface of discontinuity by the matter on one side and determined by its state of strain shall be equal and opposite to that exerted by the matter on the other side. Since

$$(\alpha')_1 = -(\alpha')_2, \quad (\beta')_1 = -(\beta')_2, \quad (\gamma')_1 = -(\gamma')_2,$$

we may also write

$$\left. \begin{aligned} \alpha'(X_{x'})_1 + \beta'(X_{y'})_1 + \gamma'(X_{z'})_1 &= \alpha'(X_{x'})_2 + \beta'(X_{y'})_2 + \gamma'(X_{z'})_2, \\ \text{etc.,} \end{aligned} \right\} (380)$$

where the signs of α' , β' , γ' may be determined by the normal on either side of the surface of discontinuity.

Equation (371) expresses the mechanical condition of equilibrium for a surface where the solid meets a fluid. It involves the separate equations

$$\left. \begin{aligned} \alpha' X_{x'} + \beta' X_{y'} + \gamma' X_{z'} &= -\alpha p \frac{Ds}{Ds'}, \\ \alpha' Y_{x'} + \beta' Y_{y'} + \gamma' Y_{z'} &= -\beta p \frac{Ds}{Ds'}, \\ \alpha' Z_{x'} + \beta' Z_{y'} + \gamma' Z_{z'} &= -\gamma p \frac{Ds}{Ds'}, \end{aligned} \right\} (381)$$

the fraction $\frac{Ds}{Ds'}$ denoting the ratio of the areas of the same element of the surface in the strained and unstrained states of the solid. These equations evidently express that the force exerted by the interior of the solid upon an element of its surface, and determined by the strain of the solid, must be normal to the surface and equal (but acting in the opposite direction) to the pressure exerted by the fluid upon the same element of surface.

If we wish to replace α and Ds by α' , β' , γ' , and the quantities which express the strain of the element, we may make use of the following considerations. The product αDs is the projection of the element Ds on the Y - Z plane. Now since the ratio $\frac{Ds}{Ds'}$ is independent of the form of the element, we may suppose that it has any convenient form. Let it be bounded by the three surfaces $x' = \text{const.}$, $y' = \text{const.}$, $z' = \text{const.}$, and let the parts of each of these surfaces included by the two others with the surface of the body be denoted by L , M , and N , or by L' , M' and N' , according as we have reference to the strained or unstrained state of the body. The areas of L' , M' , and N' are evidently $\alpha' Ds'$, $\beta' Ds'$, and $\gamma' Ds'$; and the sum of the projections of L , M and N upon any plane is equal to the projection of Ds upon that plane, since L , M , and N with Ds include a solid figure. (In propositions of this kind the *sides* of surfaces must be distinguished. If the normal to Ds falls outward from the small solid figure, the normals to L , M , and N must fall inward, and *vice*

versa). Now L' is a right-angled triangle of which the perpendicular sides may be called dy' and dz' . The projection of L on the Y - Z plane will be a triangle, the angular points of which are determined by the co-ordinates

$$y, z; \quad y + \frac{dy}{dy'} dy', \quad z + \frac{dz}{dz'} dz'; \quad y + \frac{dy}{dz'} dz', \quad z + \frac{dz}{dy'} dy';$$

the area of such a triangle is

$$\frac{1}{2} \left(\frac{dy}{dy'} \frac{dz}{dz'} - \frac{dz}{dy'} \frac{dy}{dz'} \right) dy' dz',$$

or, since $\frac{1}{2} dy' dz'$ represents the area of L' ,

$$\left(\frac{dy}{dy'} \frac{dz}{dz'} - \frac{dz}{dy'} \frac{dy}{dz'} \right) \alpha' Ds'.$$

(That this expression has the proper sign will appear if we suppose for the moment that the strain vanishes.) The areas of the projections of M and N upon the same plane will be obtained by changing y', z' and α' in this expression into z', x' , and β' , and into x', y' , and γ' . The sum of the three expressions may be substituted for $\alpha' Ds$ in (381).

We shall hereafter use Σ' to denote the sum of the three terms obtained by rotary substitutions of quantities relating to the axes X', Y', Z' , (i. e., by changing x', y', z' into y', z', x' , and into z', x', y' , with similar changes in regard to α', β', γ' , and other quantities relating to these axes,) and Σ to denote the sum of the three terms obtained by similar rotary changes of quantities relating to the axes X, Y, Z . This is only an extension of our previous use of these symbols.

With this understanding, equations (381) may be reduced to the form

$$\Sigma' (\alpha' X_{xi}) + p \Sigma' \left\{ \alpha' \left(\frac{dy}{dy'} \frac{dz}{dz'} - \frac{dz}{dy'} \frac{dy}{dz'} \right) \right\} = 0, \quad \left. \begin{array}{l} \\ \text{etc.} \end{array} \right\} \quad (382)$$

The formula (372) expresses the additional condition of equilibrium which relates to the dissolving of the solid, or its growth without discontinuity. If the solid consists entirely of substances which are actual components of the fluid, and there are no passive resistances which impede the formation or dissolving of the solid, $\delta N'$ may have either positive or negative values, and we must have

$$\varepsilon_{v_i} - t \eta_{v_i} + p v_{v_i} = \Sigma_1 (\mu_1 \Gamma_1'). \quad (383)$$

But if some of the components of the solid are only possible com-

ponents (see page 117) of the fluid, $\delta\mathcal{N}'$ is incapable of positive values, as the quantity of the solid cannot be increased, and it is sufficient for equilibrium that

$$\varepsilon_v - t\eta_v + p v_v \leq \Sigma_1(\mu_1 \Gamma_1'). \quad (384)$$

To express condition (383) in a form independent of the state of reference, we may use ε_v , η_v , Γ_1' , etc., to denote the densities of energy, of entropy, and of the several component substances in the *variable* state of the solid. We shall obtain, on dividing the equation by v_v ,

$$\varepsilon_v - t\eta_v + p = \Sigma_1(\mu_1 \Gamma_1'). \quad (385)$$

It will be remembered that the summation relates to the several components of the solid. If the solid is of uniform composition throughout, or if we only care to consider the contact of the solid and the fluid at a single point, we may treat the solid as composed of a single substance. If we use μ_1 to denote the potential for this substance in the fluid, and Γ to denote the density of the solid in the variable state, (Γ' , as before denoting its density in the state of reference,) we shall have

$$\varepsilon_v - t\eta_v + p v_v = \mu_1 \Gamma', \quad (386)$$

and

$$\varepsilon_v - t\eta_v + p = \mu_1 \Gamma. \quad (387)$$

To fix our ideas in discussing this condition, let us apply it to the case of a solid body which is homogeneous in nature and in state of strain. If we denote by ε , η , v , and m , its energy, entropy, volume, and mass, we have

$$\varepsilon - t\eta + p v = \mu_1 m. \quad (388)$$

Now the mechanical conditions of equilibrium for the surface where a solid meets a fluid require that the traction upon the surface determined by the state of strain of the solid shall be normal to the surface. This condition is always satisfied with respect to three surfaces at right angles to one another. In proving this well known proposition, we shall lose nothing in generality, if we make the state of reference, which is arbitrary, coincident with the state under discussion, the axes to which these states are referred being also coincident. We shall then have, for the normal component of the traction per unit of surface across any surface for which the direction-cosines of the normal are α , β , γ , [compare (379), and for the notation \mathcal{X}_x , etc., page 349,]

$$\begin{aligned}
 S &= \alpha (\alpha X_x + \beta X_y + \gamma X_z) \\
 &\quad + \beta (\alpha Y_x + \beta Y_y + \gamma Y_z) \\
 &\quad + \gamma (\alpha Z_x + \beta Z_y + \gamma Z_z),
 \end{aligned}$$

or, by (375), (376),

$$\begin{aligned}
 S &= \alpha^2 X_x + \beta^2 Y_y + \gamma^2 Z_z \\
 &\quad + 2 \alpha \beta X_y + 2 \beta \gamma Y_z + 2 \gamma \alpha Z_x. \quad (389)
 \end{aligned}$$

We may also choose any convenient directions for the co-ordinate axes. Let us suppose that the direction of the axis of X is so chosen that the value of S for the surface perpendicular to this axis is as great as for any other surface, and that the direction of the axis of Y (supposed at right angles to X) is such that the value of S for the surface perpendicular to it is as great as for any other surface passing through the axis of X . Then, if we write $\frac{dS}{d\alpha}$, $\frac{dS}{d\beta}$, $\frac{dS}{d\gamma}$ for the differential coefficients derived from the last equation by treating α , β , and γ as *independent* variables,

$$\frac{dS}{d\alpha} d\alpha + \frac{dS}{d\beta} d\beta + \frac{dS}{d\gamma} d\gamma = 0,$$

when

$$\alpha d\alpha + \beta d\beta + \gamma d\gamma = 0,$$

and

$$\alpha = 1, \quad \beta = 0, \quad \gamma = 0.$$

That is,

$$\frac{dS}{d\beta} = 0, \quad \text{and} \quad \frac{dS}{d\gamma} = 0,$$

when

$$\alpha = 1, \quad \beta = 0, \quad \gamma = 0.$$

Hence,

$$X_y = 0, \quad \text{and} \quad Z_x = 0. \quad (390)$$

Moreover,

$$\frac{dS}{d\beta} d\beta + \frac{dS}{d\gamma} d\gamma = 0,$$

when

$$\alpha = 0, \quad d\alpha = 0,$$

$$\beta d\beta + \gamma d\gamma = 0,$$

and

$$\beta = 1, \quad \gamma = 0.$$

Hence

$$Y_z = 0. \quad (391)$$

Therefore, when the co-ordinate axes have the supposed directions, which are called the *principal axes of stress*, the rectangular components of the traction across any surface (α , β , γ) are by (379)

$$\alpha X_x, \quad \beta Y_y, \quad \gamma Z_z. \quad (392)$$

Hence, the traction across any surface will be normal to that surface,—

(1), when the surface is perpendicular to a principal axis of stress ;

(2), if two of the *principal tractions* X_x, Y_y, Z_z are equal, when the surface is perpendicular to the plane containing the two corresponding axes, (in this case the traction across any such surface is equal to the common value of the two principal tractions);

(3), if the principal tractions are all equal, the traction is normal and constant for all surfaces.

It will be observed that in the second and third cases the position of the principal axes of stress are partially or wholly indeterminate, (so that these cases may be regarded as included in the first,) but the values of the principal tractions are always determinate, although not always different.

If, therefore, a solid which is homogeneous in nature and in state of strain is bounded by six surfaces perpendicular to the principal axes of strain, the mechanical conditions of equilibrium for these surfaces may be satisfied by the contact of fluids having the proper pressures, [see (381),] which will in general be different for the different pairs of opposite sides, and may be denoted by p', p'', p''' . (These pressures are equal to the principal tractions of the solid taken negatively.) It will then be necessary for equilibrium with respect to the tendency of the solid to dissolve that the potential for the substance of the solid in the fluids shall have values $\mu_1', \mu_1'', \mu_1'''$ determined by the equations

$$\varepsilon - t \eta + p' v = \mu_1' m, \tag{393}$$

$$\varepsilon - t \eta + p'' v = \mu_1'' m, \tag{394}$$

$$\varepsilon - t \eta + p''' v = \mu_1''' m. \tag{395}$$

These values, it will be observed, are entirely determined by the nature and state of the solid, and their differences are equal to the differences of the corresponding pressures divided by the density of the solid.

It may be interesting to compare one of these potentials, as μ_1' , with the potential (for the same substance) in a fluid of the same temperature t and pressure p' which would be in equilibrium with the same solid subjected on all sides to the uniform pressure p' . If we write $[\varepsilon]_{p'}$, $[\eta]_{p'}$, $[v]_{p'}$, and $[\mu_1]_{p'}$ for the values which ε, η, v , and μ_1 would receive on this supposition, we shall have

$$[\varepsilon]_{p'} - t [\eta]_{p'} + p' [v]_{p'} = [\mu_1]_{p'} m. \tag{396}$$

Subtracting this from (393), we obtain

$$\varepsilon - [\varepsilon]_{p'} - t \eta + t [\eta]_{p'} + p' v - p' [v]_{p'} = \mu_1 m - [\mu_1]_{p'} m. \tag{397}$$

Now it follows immediately from the definitions of energy and entropy

that the first four terms of this equation represent the work spent upon the solid in bringing it from the state of hydrostatic stress to the other state without change of temperature, and $p'v - p'[v]_{p'}$, evidently denotes the work done in displacing a fluid of pressure p' surrounding the solid during the operation. Therefore, the first number of the equation represents the total work done in bringing the solid *when surrounded by a fluid of pressure p'* from the state of hydrostatic stress p' to the state of stress p', p'', p''' . This quantity is necessarily positive, except of course in the limiting case when $p' = p'' = p'''$. If the quantity of matter of the solid body be unity, the increase of the potential in the fluid on the side of the solid on which the pressure remains constant, which will be necessary to maintain equilibrium, is equal to the work done as above described. Hence, μ_1' is greater than $[\mu_1]_{p'}$, and for similar reasons, μ_1'' is greater than the value of the potential which would be necessary for equilibrium if the solid were subjected to the uniform pressure p'' , and μ_1''' greater than that which would be necessary for equilibrium if the solid were subjected to the uniform pressure p''' . That is, (if we adapt our language to what we may regard as the most general case, viz., that in which the fluids contain the substance of the solid but are not wholly composed of that substance,) the fluids in equilibrium with the solid are all supersaturated with respect to the substance of the solid, except when the solid is in a state of hydrostatic stress; so that if there were present in any one of these fluids any small fragment of the same kind of solid subject to the hydrostatic pressure of the fluid, such a fragment would tend to increase. Even when no such fragment is present, although there must be perfect equilibrium so far as concerns the tendency of the solid to dissolve or to increase by the accretion of similarly strained matter, yet the presence of the solid which is subject to the distorting stresses, will doubtless facilitate the commencement of the formation of a solid of hydrostatic stress upon its surface, to the same extent, perhaps, in the case of an amorphous body, as if it were itself subject only to hydrostatic stress. This may sometimes, or perhaps generally, make it a necessary condition of equilibrium in cases of contact between a fluid and an amorphous solid which can be formed out of it that the solid at the surface where it meets the fluid shall be sensibly in a state of hydrostatic stress.

But in the case of a crystalline solid, subjected to distorting stresses and in contact with solutions satisfying the conditions deduced above, although crystals of hydrostatic stress would doubtless commence to

form upon its surface (if the distorting stresses and consequent supersaturation of the fluid should be carried too far), before they would commence to be formed within the fluid or on the surface of most other bodies, yet within certain limits the relations expressed by equations (393)–(395) must admit of realization, especially when the solutions are such as can be easily supersaturated.*

It may be interesting to compare the variations of p , the pressure in the fluid which determines in part the stresses and the state of strain of the solid, with other variations of the stresses or strains in the solid, with respect to the relation expressed by equation (388). To examine this point with complete generality, we may proceed in the following manner.

Let us consider so much of the solid as has in the state of reference the form of a cube, the edges of which are equal to unity, and parallel to the co-ordinate axes. We may suppose this body to be homogeneous in nature and in state of strain both in its state of reference and in its variable state. (This involves no loss of generality, since we may make the unit of length as small as we choose.) Let the fluid meet the solid on one or both of the surfaces for which Z' is constant. We may suppose these surfaces to remain perpendicular to the axis of Z in the variable state of the solid, and the edges in which y' and z' are both constant to remain parallel to the axis of X . It will be observed that these suppositions only fix the position of the strained body relatively to the co-ordinate axes, and do not in any way limit its state of strain.

It follows from the suppositions which we have made that

$$\frac{dz}{dx'} = \text{const.} = 0, \quad \frac{dz}{dy'} = \text{const.} = 0, \quad \frac{dy}{dx'} = \text{const.} = 0; \quad (398)$$

and

$$X_{z'} = 0, \quad Y_{z'} = 0, \quad Z_{z'} = -p \frac{dx}{dx'} \frac{dy}{dy'}. \quad (399)$$

Hence, by (355),

$$d\varepsilon_v = t d\eta_v + X_{x'} d\frac{dx}{dx'} + X_{y'} d\frac{dx}{dy'} + Y_{y'} d\frac{dy}{dy'} - p \frac{dx}{dx'} \frac{dy}{dy'} d\frac{dz}{dz'}. \quad (400)$$

Again, by (388),

* The effect of distorting stresses in a solid on the phenomena of crystallization and liquefaction, as well as the effect of change of hydrostatic pressure common to the solid and liquid, was first described by Professor James Thomson. See *Trans. R. S. Edin.*, vol. xvi, p. 575; and *Proc. Roy. Soc.*, vol. xi, p. 473, or *Phil. Mag.*, S. 4, vol. xxiv, p. 395.

$$d\varepsilon = t d\eta + \eta dt - p dx - v dp + m d\mu_1, \quad (401)$$

Now the suppositions which have been made require that

$$v = \frac{dx}{dx'} \frac{dy}{dy'} \frac{dz}{dz'}, \quad (402)$$

and

$$dv = \frac{dy}{dy'} \frac{dz}{dz'} d\frac{dx}{dx'} + \frac{dz}{dz'} \frac{dx}{dx'} d\frac{dy}{dy'} + \frac{dx}{dx'} \frac{dy}{dy'} d\frac{dz}{dz'}. \quad (403)$$

Combining equations (400), (401), and (403), and observing that ε_v , and η_v , are equivalent to ε and η , we obtain

$$\begin{aligned} & \eta dt - v dp + m d\mu_1 \\ & = \left(X_{x'} + p \frac{dy}{dy'} \frac{dz}{dz'} \right) d\frac{dx}{dx'} + X_{y'} d\frac{dx}{dy'} + \left(Y_{y'} + p \frac{dz}{dz'} \frac{dx}{dx'} \right) d\frac{dy}{dy'}. \end{aligned} \quad (404)$$

The reader will observe that when the solid is subjected on all sides to the uniform normal pressure p , the coefficients of the differentials in the second member of this equation will vanish. For the expres-

sion $\frac{dy}{dy'} \frac{dz}{dz'}$ represents the projection on the Y - Z plane of a side of the parallelepiped for which x' is constant, and multiplied by p it will be equal to the component parallel to the axis of X of the total pressure across this side, i. e., it will be equal to $X_{x'}$, taken negatively.

The case is similar with respect to the coefficient of $d\frac{dy}{dy'}$; and $X_{y'}$, evidently denotes a force tangential to the surface on which it acts. It will also be observed, that if we regard the forces acting upon the sides of the solid parallelepiped as composed of the hydrostatic pressure p together with addition forces, the work done in any infinitesimal variation of the state of strain of the solid by these additional forces will be represented by the second member of the equation.

We will first consider the case in which the fluid is identical in substance with the solid. We have then, by equation (97), for a mass of the fluid equal to that of the solid,

$$\eta_F dt + v_F dp + m d\mu_1 = 0, \quad (405)$$

η_F and v_F denoting the entropy and volume of the fluid. By subtraction we obtain

$$\begin{aligned} & -(\eta_F - \eta) dt + (v_F - v) dp \\ & = \left(X_{x'} + p \frac{dy}{dy'} \frac{dz}{dz'} \right) d\frac{dx}{dx'} + X_{y'} d\frac{dx}{dy'} + \left(Y_{y'} + p \frac{dz}{dz'} \frac{dx}{dx'} \right) d\frac{dy}{dy'}. \end{aligned} \quad (406)$$

Now if the quantities $\frac{dx}{dx'}$, $\frac{dx}{dy'}$, $\frac{dy}{dy'}$ remain constant, we shall have for the relation between the variations of temperature and pressure which is necessary for the preservation of equilibrium

$$\frac{dt}{dp} = \frac{v_F - v}{\eta_F - \eta} = t \frac{v_F - v}{Q}, \tag{407}$$

where Q denotes the heat which would be absorbed if the solid body should pass into the fluid state without change of temperature or pressure. This equation is similar to (131), which applies to bodies subject to hydrostatic pressure. But the value of $\frac{dt}{dp}$ will not generally be the same as if the solid were subject on all sides to the uniform normal pressure p ; for the quantities v and η (and therefore Q) will in general have different values. But when the pressures on all sides are normal and equal, the value of $\frac{dt}{dp}$ will be the same, whether we consider the pressure when varied as still normal and equal on all sides, or consider the quantities $\frac{dx}{dx'}, \frac{dy}{dy'}, \frac{dz}{dz'}$ as constant.

But if we wish to know how the temperature is affected if the pressure between the solid and fluid remains constant, but the strain of the solid is varied in any way consistent with this supposition, the differential coefficients of t with respect to the quantities which express the strain are indicated by equation (406). These differential coefficients all vanish, when the pressures on all sides are normal and equal, but the differential coefficient $\frac{dt}{dp}$, when $\frac{dx}{dx'}, \frac{dy}{dy'}, \frac{dz}{dz'}$ are constant, or when the pressures on all sides are normal and equal, vanishes only when the density of the fluid is equal to that of the solid.

The case is nearly the same when the fluid is not identical in substance with the solid, if we suppose the composition of the fluid to remain unchanged. We have necessarily with respect to the fluid

$$d\mu_1 = \left(\frac{d\mu_1}{dt}\right)_{p, m}^{(F)} dt + \left(\frac{d\mu_1}{dp}\right)_{t, m}^{(F)} dp,^* \tag{408}$$

where the index (F) is used to indicate that the expression to which it is affixed relates to the fluid. But by equation (92)

$$\left(\frac{d\mu_1}{dt}\right)_{p, m}^{(F)} = -\left(\frac{d\eta}{dm_1}\right)_{t, p, m}^{(F)}, \text{ and } \left(\frac{d\mu_1}{dp}\right)_{t, m}^{(F)} = \left(\frac{dv}{dm_1}\right)_{t, p, m}^{(F)}. \tag{409}$$

Substituting these values in the preceding equation, transposing terms, and multiplying by m , we obtain

* A suffixed m stands here, as elsewhere in this paper, for all the symbols m_1, m_2 , etc., except such as may occur in the differential coefficient.

$$m \left(\frac{d\eta}{dm_1} \right)_{t, p, m}^{(F)} dt - m \left(\frac{dv}{dm_1} \right)_{t, p, m}^{(F)} dp + m d\mu_1 = 0. \quad (410)$$

By subtracting this equation from (404) we may obtain an equation similar to (406), except that in place of η_F and v_F we shall have the expressions

$$m \left(\frac{d\eta}{dm_1} \right)_{t, p, m}^{(F)} \quad \text{and} \quad m \left(\frac{dv}{dm_1} \right)_{t, p, m}^{(F)}$$

The discussion of equation (406) will therefore apply *mutatis mutandis* to this case.

We may also wish to find the variations in the composition of the fluid which will be necessary for equilibrium when the pressure p or the quantities $\frac{dx}{dx'}$, $\frac{dy}{dy'}$, $\frac{dz}{dz'}$ are varied, the temperature remaining constant. If we know the value for the fluid of the quantity represented by ζ on page 142 in terms of t, p , and the quantities of the several components m_1, m_2, m_3 , etc., the first of which relates to the substance of which the solid is formed, we can easily find the value of μ_1 in terms of the same variables. Now in considering variations in the composition of the solid, it will be sufficient if we make all but one of the components variable. We may therefore give to m_1 a constant value, and making t also constant, we shall have

$$d\mu_1 = \left(\frac{d\mu_1}{dp} \right)_{t, m}^{(F)} dp + \left(\frac{d\mu_1}{dm_2} \right)_{t, p, m}^{(F)} dm_2 + \left(\frac{d\mu_1}{dm_3} \right)_{t, p, m}^{(F)} dm_3 + \text{etc.}$$

Substituting this value in equation (404), and cancelling the term containing dt , we obtain

$$\begin{aligned} & \left\{ m \left(\frac{d\mu_1}{dp} \right)_{t, m}^{(F)} - v \right\} dp + m \left(\frac{d\mu_1}{dm_2} \right)_{t, p, m}^{(F)} dm_2 \\ & + m \left(\frac{d\mu_1}{dm_3} \right)_{t, p, m}^{(F)} dm_3 + \text{etc.} = \left(X_x + p \frac{dy}{dy'} \frac{dz}{dz'} \right) d \frac{dx}{dx'} \\ & + X_y d \frac{dx'}{dy'} + \left(Y_y + p \frac{dz}{dz'} \frac{dx}{dx'} \right) d \frac{dy}{dy'}. \end{aligned} \quad (411)$$

This equation shows the variation in the quantity of any one of the components of the fluid (other than the substance which forms the solid) which will balance a variation of p , or of $\frac{dx}{dx'}$, $\frac{dy}{dy'}$, $\frac{dz}{dz'}$, with respect to the tendency of the solid to dissolve.

Fundamental Equations for Solids.

The principles developed in the preceding pages show that the solution of problems relating to the equilibrium of a solid, or at least their reduction to purely analytical processes, may be made to depend upon our knowledge of the composition and density of the solid at every point in some particular state, which we have called the state of reference, and of the relation existing between the quantities which have been represented by ε_{v_1} , η_{v_1} , $\frac{dx}{dx'}$, $\frac{dy}{dy'}$, \dots , $\frac{dz}{dz'}$, x' , y' , and z' . When the solid is in contact with fluids, a certain knowledge of the properties of the fluids is also requisite, but only such as is necessary for the solution of problems relating to the equilibrium of fluids among themselves.

If in any state of which a solid is capable, it is homogeneous in its nature and in its state of strain, we may choose this state as the state of reference, and the relation between ε_{v_1} , η_{v_1} , $\frac{dx}{dx'}$, \dots , $\frac{dz}{dz'}$, will be independent of x' , y' , z' . But it is not always possible, even in the case of bodies which are homogeneous in nature, to bring all the elements simultaneously into the same state of strain. It would not be possible, for example, in the case of a Prince Rupert's drop.

If, however, we know the relation between ε_{v_1} , η_{v_1} , $\frac{dx}{dx'}$, \dots , $\frac{dz}{dz'}$ for any kind of homogeneous solid, with respect to any given state of reference, we may derive from it a similar relation with respect to any other state as a state of reference. For if x' , y' , z' denote the co-ordinates of points of the solid in the first state of reference, and x'' , y'' , z'' the co-ordinates of the same points in the second state of reference, we shall have necessarily

$$\frac{dx}{dx'} = \frac{dx}{dx''} \frac{dx''}{dx'} + \frac{dy}{dy''} \frac{dy''}{dx'} + \frac{dz}{dz''} \frac{dz''}{dx'}, \text{ etc. (nine equations),} \quad (412)$$

and if we write R for the volume of an element in the state (x'', y'', z'') divided by its volume in the state (x', y', z') , we shall have

$$R = \begin{vmatrix} \frac{dx''}{dx'} & \frac{dy''}{dy'} & \frac{dz''}{dz'} \\ \frac{dy''}{dx'} & \frac{dy''}{dy'} & \frac{dy''}{dz'} \\ \frac{dz''}{dx'} & \frac{dz''}{dy'} & \frac{dz''}{dz'} \end{vmatrix}, \quad (413)$$

$$\varepsilon_{v_i} = R \varepsilon_{v_{ii}}, \quad \eta_{v_i} = R \eta_{v_{ii}}. \quad (414)$$

If, then, we have ascertained by experiment the value of ε_{v_i} in terms of $\eta_{v_i}, \frac{dx}{dx'}, \dots \frac{dz}{dz'}$, and the quantities which express the composition of the body, by the substitution of the values given in (412)–(414), we shall obtain $\varepsilon_{v_{ii}}$ in terms of $\eta_{v_{ii}}, \frac{dx}{dx''}, \dots \frac{dz}{dz''}, \frac{dx''}{dx'}, \dots \frac{dz''}{dz'}$, and the quantities which express the composition of the body.

We may apply this to the elements of a body which may be variable from point to point in composition and state of strain in a given state of reference (x'', y'', z''), and if the body is fully described in that state of reference, both in respect to its composition and to the displacement which it would be necessary to give to a homogeneous solid of the same composition, for which ε_{v_i} is known in terms of $\eta_{v_i}, \frac{dx}{dx'}, \dots \frac{dz}{dz'}$, and the quantities which express its composition, to bring it from the state of reference (x', y', z') into a similar and similarly situated state of strain with that of the element of the non-homogeneous body, we may evidently regard $\frac{dx''}{dx'}, \dots \frac{dz''}{dz'}$ as known for each element of the body, that is, as known in terms of x'', y'', z'' . We shall then have $\varepsilon_{v_{ii}}$ in terms of $\eta_{v_{ii}}, \frac{dx}{dx''}, \dots \frac{dz}{dz''}, x'', y'', z''$; and since the composition of the body is known in terms of x'', y'', z'' , and the density, if not given directly, can be determined from the density of the homogeneous body in its state of reference (x', y', z'), this is sufficient for determining the equilibrium of any given state of the non-homogeneous solid.

An equation, therefore, which expresses for any kind of solid, and with reference to any determined state of reference, the relation between the quantities denoted by $\varepsilon_{v_i}, \eta_{v_i}, \frac{dx}{dx'}, \dots \frac{dz}{dz'}$, involving also the quantities which express the composition of the body, when that is capable of continuous variation, or any other equation from which the same relations may be deduced, may be called a *fundamental equation* for that kind of solid. It will be observed that the sense in which this term is here used, is entirely analogous to that in which we have already applied the term to fluids and solids which are subject only to hydrostatic pressure.

When the fundamental equation between $\varepsilon_{v_i}, \eta_{v_i}, \frac{dx}{dx'}, \dots \frac{dz}{dz'}$ is

known, we may obtain by differentiation the values of $t, X_{x_i}, \dots, Z_{z_i}$ in terms of the former quantities, which will give eleven independent relations between the twenty-one quantities

$$\varepsilon_{v_i}, \eta_{v_i}, \frac{dx}{dx'}, \dots, \frac{dz}{dz'}, t, X_{x_i}, \dots, Z_{z_i}, \quad (415)$$

which are all that exist, since ten of these quantities are independent. All these equations may also involve variables which express the composition of the body, when that is capable of continuous variation.

If we use the symbol ψ_{v_i} to denote the value of ψ (as defined on pages 144, 145) for any element of a solid divided by the volume of the element in the state of reference, we shall have

$$\psi_{v_i} = \varepsilon_{v_i} - t \eta_{v_i}. \quad (416)$$

The equation (356) may be reduced to the form

$$\delta \psi_{v_i} = - \eta_{v_i} \delta t + \sum \sum' \left(X_{x_i} \delta \frac{dx}{dx'} \right). \quad (417)$$

Therefore, if we know the value of ψ_{v_i} in terms of the variables $t, \frac{dx}{dx'}, \dots, \frac{dz}{dz'}$, together with those which express the composition of the body, we may obtain by differentiation the values of $\eta_{v_i}, X_{x_i}, \dots, Z_{z_i}$ in terms of the same variables. This will make eleven independent relations between the same quantities as before, except that we shall have ψ_{v_i} instead of ε_{v_i} . Or if we eliminate ψ_{v_i} by means of equation (416), we shall obtain eleven independent equations between the quantities in (415) and those which express the composition of the body. An equation, therefore, which determines the value of ψ_{v_i} , as a function of the quantities $t, \frac{dx}{dx'}, \dots, \frac{dz}{dz'}$, and the quantities which express the composition of the body when it is capable of continuous variation, is a fundamental equation for the kind of solid to which it relates.

In the discussion of the conditions of equilibrium of a solid, we might have started with the principle that it is necessary and sufficient for equilibrium that the temperature shall be uniform throughout the whole mass in question, and that the variation of the force-function ($-\psi$) of the same mass shall be null or negative for any variation in the state of the mass not affecting its temperature. We might have assumed that the value of ψ for any same element of the solid is a

function of the temperature and the state of strain, so that for constant temperature we might write

$$\delta\psi_v = \sum \sum' \left(X_{x_i} \delta \frac{dx}{dx'} \right),$$

the quantities X_{x_i}, \dots, Z_{z_i} being defined by this equation. This would be only a formal change in the definition of X_{x_i}, \dots, Z_{z_i} , and would not affect their values, for this equation holds true of X_{x_i}, \dots, Z_{z_i} as defined by equation (355). With such data, by transformations similar to those which we have employed, we might obtain similar results.* It is evident that the only difference in the equations would be that ψ_v would take the place of ϵ_v , and that the terms relating to entropy would be wanting. Such a method is evidently preferable with respect to the directness with which the results are obtained. The method of this paper shows more distinctly the rôle of *energy* and *entropy* in the theory of equilibrium, and can be extended more naturally to those dynamical problems in which motions take place under the condition of constancy of entropy of the elements of a solid (as when vibrations are propagated through a solid), just as the other method can be more naturally extended to dynamical problems in which the temperature is constant. (See note on page 145.)

We have already had occasion to remark that the state of strain of any element considered without reference to directions in space is capable of only six independent variations. Hence, it must be possible to express the state of strain of an element by six functions of $\frac{dx}{dx'}, \dots, \frac{dz}{dz'}$, which are independent of the position of the element.

For these quantities we may choose the squares of the ratios of elongation of lines parallel to the three co-ordinate axes in the state of reference, and the products of the ratios of elongation for each pair of these lines multiplied by the cosine of the angle which they include in the variable state of the solid. If we denote these quantities by A, B, C, a, b, c , we shall have

* For an example of this method, see Thomson and Tait's *Natural Philosophy*, vol. i, p. 705. With regard to the general theory of elastic solids, compare also Thomson's Memoir "On the Thermo-elastic and Thermo-magnetic Properties of Matter" in the *Quarterly Journal of Mathematics*, vol. i, p. 57 (1855), and Green's memoirs on the propagation, reflection, and refraction of light in the *Transactions of the Cambridge Philosophical Society*, vol. vii.

$$A = \sum \left(\frac{dx}{dx'} \right)^2, \quad B = \sum \left(\frac{dy}{dy'} \right)^2, \quad C = \sum \left(\frac{dz}{dz'} \right)^2, \quad (418)$$

$$a = \sum \left(\frac{dx}{dy'} \frac{dx'}{dz'} \right), \quad b = \sum \left(\frac{dx}{dz'} \frac{dx'}{dx'} \right), \quad c = \sum \left(\frac{dx}{dx'} \frac{dx'}{dy'} \right). \quad (419)$$

The determination of the fundamental equation for a solid is thus reduced to the determination of the relation between ϵ_{v_i} , η_{v_i} , A , B , C , a , b , c , or of the relation between ψ_{v_i} , t , A , B , C , a , b , c .

In the case of isotropic solids, the state of strain of an element, so far as it can affect the relation of ϵ_{v_i} and η_{v_i} , or of ψ_{v_i} and t , is capable of only three independent variations. This appears most distinctly as a consequence of the proposition that for any given strain of an element there are three lines in the element which are at right angles to one another both in its unstrained and in its strained state. If the unstrained element is isotropic, the ratios of elongation for these three lines must with η_{v_i} determine the value ϵ_{v_i} , or with t determine the value of ψ_{v_i} .

To demonstrate the existence of such lines, which are called the *principal axes of strain*, and to find the relations of the elongations of such lines to the quantities $\frac{dx}{dx'}$, . . . $\frac{dz}{dz'}$, we may proceed as follows. The ratio of elongation r of any line of which α' , β' , γ' are the direction-cosines in the state of reference is evidently given by the equation

$$\begin{aligned} r^2 = & \left(\frac{dx}{dx'} \alpha' + \frac{dy}{dy'} \beta' + \frac{dz}{dz'} \gamma' \right)^2 \\ & + \left(\frac{dy}{dx'} \alpha' + \frac{dy}{dy'} \beta' + \frac{dy}{dz'} \gamma' \right)^2 \\ & + \left(\frac{dz}{dx'} \alpha' + \frac{dz}{dy'} \beta' + \frac{dz}{dz'} \gamma' \right)^2. \end{aligned} \quad (420)$$

Now the proposition to be established is evidently equivalent to this—that it is always possible to give such directions to the two systems of rectangular axes X' , Y' , Z' , and X , Y , Z , that

$$\left. \begin{aligned} \frac{dx}{dy'} = 0, \quad \frac{dx}{dz'} = 0, \quad \frac{dy}{dz'} = 0, \\ \frac{dy}{dx'} = 0, \quad \frac{dz}{dx'} = 0, \quad \frac{dz}{dy'} = 0. \end{aligned} \right\} \quad (421)$$

We may choose a line in the element for which the value of r is at least as great as for any other, and make the axes of X and X' parallel to this line in the strained and unstrained states respectively.

Then
$$\frac{dy}{dx'} = 0, \quad \frac{dz}{dx'} = 0. \quad (422)$$

Moreover, if we write $\frac{d(r^2)}{d\alpha'}, \frac{d(r^2)}{d\beta'}, \frac{d(r^2)}{d\gamma'}$ for the differential coefficients obtained from (420) by treating α', β', γ' as *independent* variables,

$$\frac{d(r^2)}{d\alpha'} d\alpha' + \frac{d(r^2)}{d\beta'} d\beta' + \frac{d(r^2)}{d\gamma'} d\gamma' = 0,$$

when $\alpha' d\alpha' + \beta' d\beta' + \gamma' d\gamma' = 0,$

and $\alpha' = 1, \beta' = 0, \gamma' = 0.$

That is, $\frac{d(r^2)}{d\beta'} = 0,$ and $\frac{d(r^2)}{d\gamma'} = 0,$

when $\alpha' = 1, \beta' = 0, \gamma' = 0.$

Hence,
$$\frac{dx}{dy'} = 0, \quad \frac{dx}{dz'} = 0. \quad (423)$$

Therefore a line of the element which in the unstrained state is perpendicular to X' is perpendicular to X in the strained state. Of all such lines we may choose one for which the value of r is at least as great as for any other, and make the axes of Y' and Z' parallel to this line in the unstrained and in the strained state respectively. Then

$$\frac{dz}{dy'} = 0; \quad (424)$$

and it may easily be shown by reasoning similar to that which has just been employed that

$$\frac{dy}{dz'} = 0. \quad (425)$$

Lines parallel to the axes of $X', Y',$ and Z' in the unstrained body will therefore be parallel to $X, Y,$ and Z in the strained body, and the ratios of elongation for such lines will be

$$\frac{dx}{dx'}, \quad \frac{dy}{dy'}, \quad \frac{dz}{dz'}.$$

These lines have the common property of a stationary value of the ratio of elongation for varying directions of the line. This appears from the form to which the general value of r^2 is reduced by the positions of the co-ordinate axes, viz.,

$$r^2 = \left(\frac{dx}{dx'}\right)^2 \alpha'^2 + \left(\frac{dy}{dy'}\right)^2 \beta'^2 + \left(\frac{dz}{dz'}\right)^2 \gamma'^2.$$

Having thus proved the existence of lines, with reference to any particular strain, which have the properties mentioned, let us proceed to find the relations between the ratios of elongation for these lines (the *principal axes of strain*) and the quantities $\frac{dx}{dx'}$, . . . $\frac{dz}{dz'}$ under the most general supposition with respect to the position of the co-ordinate axes.

For any principal axis of strain we have

$$\frac{d(r^2)}{d\alpha'} d\alpha' + \frac{d(r^2)}{d\beta'} d\beta' + \frac{d(r^2)}{d\gamma'} d\gamma' = 0,$$

when
$$\alpha' d\alpha' + \beta' d\beta' + \gamma' d\gamma' = 0,$$

the differential coefficients in the first of these equations being determined from (420) as before. Therefore,

$$\frac{1}{\alpha'} \frac{d(r^2)}{d\alpha'} = \frac{1}{\beta'} \frac{d(r^2)}{d\beta'} = \frac{1}{\gamma'} \frac{d(r^2)}{d\gamma'}. \tag{426}$$

From (420) we obtain directly

$$\frac{\alpha' d(r^2)}{2 d\alpha'} + \frac{\beta' d(r^2)}{2 d\beta'} + \frac{\gamma' d(r^2)}{2 d\gamma'} = r^2. \tag{427}$$

From the two last equations, in virtue of the necessary relation $\alpha'^2 + \beta'^2 + \gamma'^2 = 1$, we obtain

$$\frac{1}{2} \frac{d(r^2)}{d\alpha'} = \alpha' r^2, \quad \frac{1}{2} \frac{d(r^2)}{d\beta'} = \beta' r^2, \quad \frac{1}{2} \frac{d(r^2)}{d\gamma'} = \gamma' r^2, \tag{428}$$

or, if we substitute the values of the differential coefficients taken from (420),

$$\left. \begin{aligned} \alpha' &\geq \left(\frac{dx}{dx'}\right)^2 + \beta' \geq \left(\frac{dx}{dx'} \frac{dx}{dy'}\right) + \gamma' \geq \left(\frac{dx}{dx'} \frac{dx}{dz'}\right) = \alpha' r^2, \\ \alpha' &\geq \left(\frac{dx}{dy'} \frac{dx}{dx'}\right) + \beta' \geq \left(\frac{dx}{dy'}\right)^2 + \gamma' \geq \left(\frac{dx}{dy'} \frac{dx}{dz'}\right) = \beta' r^2, \\ \alpha' &\geq \left(\frac{dx}{dz'} \frac{dx}{dx'}\right) + \beta' \geq \left(\frac{dx}{dz'} \frac{dx}{dy'}\right) + \gamma' \geq \left(\frac{dx}{dz'}\right)^2 = \gamma' r^2, \end{aligned} \right\} \tag{429}$$

If we eliminate α' , β' , γ' from these equations, we may write the result in the form,

$$\left| \begin{array}{ccc} \geq \left(\frac{dx}{dx'}\right)^2 - r^2 & \geq \left(\frac{dx}{dx'} \frac{dx}{dy'}\right) & \geq \left(\frac{dx}{dx'} \frac{dx}{dz'}\right) \\ \geq \left(\frac{dx}{dy'} \frac{dx}{dx'}\right) & \geq \left(\frac{dx}{dy'}\right)^2 - r^2 & \geq \left(\frac{dx}{dy'} \frac{dx}{dz'}\right) \\ \geq \left(\frac{dx}{dz'} \frac{dx}{dx'}\right) & \geq \left(\frac{dx}{dz'} \frac{dx}{dy'}\right) & \geq \left(\frac{dx}{dz'}\right)^2 - r^2 \end{array} \right| = 0. \tag{430}$$

We may write

$$-r^6 + Er^4 - Fr^2 + G = 0. \tag{431}$$

Then

$$E = \sum' \left(\frac{dx}{dx'} \right)^2. \tag{432}$$

Also*

$$\begin{aligned} F &= \sum' \left\{ \sum \left(\frac{dx}{dx'} \right)^2 \sum \left(\frac{dx}{dy'} \right)^2 - \sum \left(\frac{dx}{dx'} \frac{dx}{dy'} \right) \sum \left(\frac{dx}{dx'} \frac{dx}{dy'} \right) \right\} \\ &= \sum' \sum \left\{ \left(\frac{dx}{dx'} \right)^2 \sum \left(\frac{dx}{dy'} \right)^2 - \frac{dx}{dx'} \frac{dx}{dy'} \sum \left(\frac{dx}{dx'} \frac{dx}{dy'} \right) \right\} = \\ \sum' \sum &\left\{ \left(\frac{dx}{dx'} \right)^2 \left(\frac{dy}{dy'} \right)^2 + \left(\frac{dx}{dx'} \right)^2 \left(\frac{dz}{dy'} \right)^2 - \frac{dx}{dx'} \frac{dx}{dy'} \frac{dy}{dx'} \frac{dy}{dy'} - \frac{dx}{dx'} \frac{dx}{dy'} \frac{dz}{dx'} \frac{dz}{dy'} \right\} \\ &= \sum' \sum \left\{ \left(\frac{dx}{dx'} \right)^2 \left(\frac{dy}{dy'} \right)^2 + \left(\frac{dy}{dx'} \right)^2 \left(\frac{dx}{dy'} \right)^2 - 2 \frac{dx}{dx'} \frac{dx}{dy'} \frac{dy}{dx'} \frac{dy}{dy'} \right\} \\ &= \sum' \sum \left(\frac{dx}{dx'} \frac{dy}{dy'} - \frac{dy}{dx'} \frac{dx}{dy'} \right)^2. \tag{433} \end{aligned}$$

This may also be written

$$F = \sum' \sum \left| \begin{array}{cc} \frac{dx}{dx'} & \frac{dx}{dy'} \\ \frac{dy}{dx'} & \frac{dy}{dy'} \end{array} \right|^2. \tag{434}$$

In the reduction of the value of G , it will be convenient to use the symbol \sum_{3+3} to denote the sum of the *six* terms formed by changing x, y, z , into y, z, x ; z, x, y ; x, z, y ; y, x, z ; and z, y, x ; and the symbol \sum_{3-3} in the same sense except that the last three terms are to be taken negatively; also to use \sum'_{3-3} in a similar sense with respect to x', y', z' ; and to use x', y', z' as equivalent to x', y', z' , except that they are not to be affected by the sign of summation. With this understanding we may write

$$G = \sum'_{3-3} \left\{ \sum \left(\frac{dx}{dx'} \frac{dx}{dx'} \right) \sum \left(\frac{dx}{dy'} \frac{dx}{dy'} \right) \sum \left(\frac{dx}{dz'} \frac{dx}{dz'} \right) \right\}. \tag{435}$$

* The values of F and G given in equations (434) and (438), which are here deduced at length, may be derived from inspection of equation (430) by means of the usual theorems relating to the multiplication of determinants. See Salmon's *Lessons Introductory to the Modern Higher Algebra*, 2d Ed., Lesson III; or Baltzer's *Theorie und Anwendung der Determinanten*, § 5.

In expanding the product of the three sums, we may cancel on account of the sign \sum'_{3-3} the terms which do not contain all the three expressions dx , dy , and dz . Hence we may write

$$\begin{aligned} G &= \sum'_{3-3} \sum_{3+3} \left(\frac{dx}{dx'} \frac{dx}{dx'} \frac{dy}{dy'} \frac{dy}{dy'} \frac{dz}{dz'} \frac{dz}{dz'} \right) \\ &= \sum_{3+3} \left\{ \frac{dx}{dx'} \frac{dy}{dy'} \frac{dz}{dz'} \sum'_{3-3} \left(\frac{dx}{dx'} \frac{dy}{dy'} \frac{dz}{dz'} \right) \right\} \\ &= \sum_{3-3} \left(\frac{dx}{dx'} \frac{dy}{dy'} \frac{dz}{dz'} \right) \sum'_{3-3} \left(\frac{dx}{dx'} \frac{dy}{dy'} \frac{dz}{dz'} \right). \end{aligned} \tag{436}$$

Or, if we set

$$H = \begin{vmatrix} \frac{dx}{dx'} & \frac{dx}{dy'} & \frac{dx}{dz'} \\ \frac{dy}{dx'} & \frac{dy}{dy'} & \frac{dy}{dz'} \\ \frac{dz}{dx'} & \frac{dz}{dy'} & \frac{dz}{dz'} \end{vmatrix}, \tag{437}$$

we shall have

$$G = H^2. \tag{438}$$

It will be observed that F represents the sum of the squares of the nine minors which can be formed from the determinant in (437), and that E represents the sum of the squares of the nine constituents of the same determinant.

Now we know by the theory of equations that equation (431) will be satisfied in general by three different values of r^2 , which we may denote by r_1^2 , r_2^2 , r_3^2 , and which must represent the squares of the ratios of elongation for the three principal axes of strain; also that E , F , G , are symmetrical functions of r_1^2 , r_2^2 , r_3^2 , viz.,

$$\left. \begin{aligned} E &= r_1^2 + r_2^2 + r_3^2, & F &= r_1^2 r_2^2 + r_2^2 r_3^2 + r_3^2 r_1^2, \\ G &= r_1^2 r_2^2 r_3^2. \end{aligned} \right\} \tag{439}$$

Hence, although it is possible to solve equation (431) by the use of trigonometrical functions, it will be more simple to regard ϵ_v as a function of η_v , and the quantities E , F , G (or H), which we have expressed in terms of $\frac{dx}{dx'}$, . . . $\frac{dz}{dz'}$. Since ϵ_v is a single-valued function of η_v , and r_1^2 , r_2^2 , r_3^2 (with respect to all the changes of which the body is capable), and a symmetrical function with respect to r_1^2 , r_2^2 , r_3^2 , and since r_1^2 , r_2^2 , r_3^2 are *collectively* determined without ambiguity by the values of E , F , and H , the quantity ϵ_v must be a

single-valued function of η_v , E , F , and H . The determination of the fundamental equation for isotropic bodies is therefore reduced to the determination of this function, or (as appears from similar considerations) the determination of ψ_v , as a function of t , E , F , and H .

It appears from equations (439) that E represents the sum of the squares of the ratios of elongation for the principal axes of strain, that F represents the sum of the squares of the ratios of enlargement for the three surfaces determined by these axes, and that G represents the square of the ratio of enlargement of volume. Again, equation (432) shows that E represents the sum of the squares of the ratios of elongation for lines parallel to X' , Y' , and Z' ; equation (434) shows that F represents the sum of the squares of the ratios of enlargement for surfaces parallel to the planes $X'-Y'$, $Y'-Z'$, $Z'-X'$; and equation (438), like (439), shows that G represents the square of the ratio of enlargement of volume. Since the position of the co-ordinate axes is arbitrary, it follows that the sum of the squares of the ratios of elongation or enlargement of three lines or surfaces which in the unstrained state are at right angles to one another, is otherwise independent of the direction of the lines or surfaces. Hence, $\frac{1}{3}E$ and $\frac{1}{3}F$ are the mean squares of the ratios of linear elongation and of superficial enlargement, for all possible directions in the unstrained solid.

There is not only a practical advantage in regarding the strain as determined by E , F , and H , instead of E , F , and G , because H is more simply expressed in terms of $\frac{dx}{dx'}$, . . . $\frac{dz}{dz'}$, but there is also a certain theoretical advantage on the side of E , F , H . If the systems of co-ordinate axes X , Y , Z , and X' , Y' , Z' , are either identical or such as are capable of superposition, which it will always be convenient to suppose, the determinant H will always have a positive value for any strain of which a body can be capable. But it is possible to give to x , y , z such values as functions of x' , y' , z' that H shall have a negative value. For example, we may make

$$x = x', \quad y = y', \quad z = -z'. \quad (440)$$

This will give $H = -1$, while

$$x = x', \quad y = y', \quad z = z' \quad (441)$$

will give $H = 1$. Both (440) and (441) give $G = 1$. Now although such a change in the position of the particles of a body as is represented by (440) cannot take place while the body remains solid, yet

a method of representing strains may be considered incomplete, which confuses the cases represented by (440) and (441).

We may avoid all such confusion by using E , F , and H to represent a strain. Let us consider an element of the body strained which in the state (x', y', z') is a cube with its edges parallel to the axes of X' , Y' , Z' , and call the edges dx' , dy' , dz' according to the axes to which they are parallel, and consider the ends of the edges as positive for which the values of x' , y' , or z' are the greater. Whatever may be the nature of the parallelepiped in the state (x, y, z) which corresponds to the cube dx' , dy' , dz' and is determined by the quantities $\frac{dx}{dx'}$, . . . $\frac{dz}{dz'}$, it may always be brought by continuous changes to the form of a cube and to a position in which the edges dx' , dy' shall be parallel to the axes of X and Y , the positive ends of the edges toward the positive directions of the axes, and this may be done without giving the volume of the parallelepiped the value zero, and therefore without changing the sign of H . Now two cases are possible;—the positive end of the edge dz' may be turned toward the positive or toward the negative direction of the axis of Z . In the first case, H is evidently positive; in the second, negative. The determinant H will therefore be positive or negative,—we may say, if we choose, that the volume will be positive or negative,—according as the element can or cannot be brought from the state (x, y, z) to the state (x', y', z') by continuous changes without giving its volume the value zero.

If we now recur to the consideration of the principal axes of strain and the principal ratios of elongation r_1, r_2, r_3 , and denote by U_1, U_2, U_3 and U_1', U_2', U_3' the principal axes of strain in the strained and unstrained element respectively, it is evident that the sign of r_1 , for example, depends upon the direction in U_1 which we regard as corresponding to a given direction in U_1' . If we choose to associate directions in these axes so that r_1, r_2, r_3 shall all be positive, the positive or negative value of H will determine whether the system of axes U_1, U_2, U_3 is or is not capable of superposition upon the system U_1', U_2', U_3' so that corresponding directions in the axes shall coincide. Or, if we prefer to associate directions in the two systems of axes, so that they shall be capable of superposition, corresponding directions coinciding, the positive or negative value of H will determine whether an even or an odd number of the quantities r_1, r_2, r_3 are negative. In this case we may write

$$r_1 r_2 r_3 = H = \begin{vmatrix} \frac{dx}{dx'} & \frac{dx}{dy'} & \frac{dx}{dz'} \\ \frac{dy}{dx'} & \frac{dy}{dy'} & \frac{dy}{dz'} \\ \frac{dz}{dx'} & \frac{dz}{dy'} & \frac{dz}{dz'} \end{vmatrix}. \quad (442)$$

It will be observed that to change the signs of two of the quantities r_1, r_2, r_3 is simply to give a certain rotation to the body without changing its state of strain.

Whichever supposition we make with respect to the axes U_1, U_2, U_3 , it is evident that the state of strain is completely determined by the values E, F , and H , not only when we limit ourselves to the consideration of such strains as are consistent with the idea of solidity, but also when we regard any values of $\frac{dx}{dx'}, \dots, \frac{dz}{dz'}$ as possible.

Approximative Formulæ.—For many purposes the value of ε_v , for an isotropic solid may be represented with sufficient accuracy by the formula

$$\varepsilon_v = i' + e' E + f' F + h' H, \quad (443)$$

where i', e', f' , and h' denote functions of η_v ; or the value of ψ_v , by the formula

$$\psi_v = i + e E + f F + h H, \quad (444)$$

where i, e, f , and h denote functions of t . Let us first consider the second of these formulæ. Since E, F , and H are symmetrical functions of r_1, r_2, r_3 , if ψ_v is any function of t, E, F, H , we must have

$$\left. \begin{aligned} \frac{d\psi_v}{dr_1} &= \frac{d\psi_v}{dr_2} = \frac{d\psi_v}{dr_3}, \\ \frac{d^2\psi_v}{dr_1^2} &= \frac{d^2\psi_v}{dr_2^2} = \frac{d^2\psi_v}{dr_3^2}, \\ \frac{d^2\psi_v}{dr_1 dr_2} &= \frac{d^2\psi_v}{dr_2 dr_3} = \frac{d^2\psi_v}{dr_3 dr_1} \end{aligned} \right\} \quad (445)$$

whenever $r_1 = r_2 = r_3$. Now i, e, f , and h may be determined (as functions of t) so as to give to

$$\psi_v, \quad \frac{d\psi_v}{dr_1}, \quad \frac{d^2\psi_v}{dr_1^2}, \quad \frac{d^2\psi_v}{dr_1 dr_2}$$

their proper values at every temperature for some isotropic state of strain, which may be determined by any desired condition. We shall suppose that they are determined so as to give the proper

values to ψ_v , etc., when the stresses in the solid vanish. If we denote by r_0 the common value of r_1, r_2, r_3 which will make the stresses vanish at any given temperature, and imagine the true value of ψ_v , and also the value given by equation (444) to be expressed in terms of the ascending powers of

$$r_1 - r_0, \quad r_2 - r_0, \quad r_3 - r_0, \tag{446}$$

it is evident that the expressions will coincide as far as the terms of the second degree *inclusive*. That is, the errors of the values of ψ_v , given by equation (444) are of the same order of magnitude as the cubes of the above differences. The errors of the values of

$$\frac{d\psi_v}{dr_1}, \quad \frac{d\psi_v}{dr_2}, \quad \frac{d\psi_v}{dr_3}$$

will be of the same order of magnitude as the squares of the same differences. Therefore, since

$$\frac{d\psi_v}{d\bar{x}} = \frac{d\psi_v}{dr_1} \frac{dr_1}{d\bar{x}} + \frac{d\psi_v}{dr_2} \frac{dr_2}{d\bar{x}} + \frac{d\psi_v}{dr_3} \frac{dr_3}{d\bar{x}} \tag{447}$$

whether we regard the true value of ψ_v , or the value given by equation (444), and since the error in (444) does not affect the values of

$$\frac{dr_1}{d\bar{x}}, \quad \frac{dr_2}{d\bar{x}}, \quad \frac{dr_3}{d\bar{x}},$$

which we may regard as determined by equations (431), (432), (434), (437) and (438), the errors in the values of \bar{X}_v , derived from (444) will be of the same order of magnitude as the squares of the differences in (446). The same will be true with respect to $\bar{X}_v, \bar{X}_z, \bar{Y}_v$, etc., etc.

It will be interesting to see how the quantities e, f , and h are related to those which most simply represent the elastic properties of isotropic solids. If we denote by V and R the *elasticity of volume* and the *rigidity** (both determined under the condition of constant temperature and for states of vanishing stress), we shall have as definitions:

$$V = -v \left(\frac{dp}{dv} \right)_t, \quad \text{when} \quad v = r_0^3 v', \tag{448}$$

* See Thomson and Tait's *Natural Philosophy*, vol. i, p. 711.

where p denotes a uniform pressure to which the solid is subjected, v its volume, and v' its volume in the state of reference; and

$$\left. \begin{aligned} R &= \frac{dX_{v'}}{d\frac{dx}{dy'}} = \frac{d^2\psi_{v'}}{\left(d\frac{dx}{dy'}\right)^2}, \\ \text{when} \quad \frac{dx}{dx'} &= \frac{dy}{dy'} = \frac{dz}{dz'} = r_0, \\ \text{and} \quad \frac{dx}{dy'} &= \frac{dx}{dz'} = \frac{dy}{dz'} = \frac{dy}{dx'} = \frac{dz}{dx'} = \frac{dz}{dy'} = 0. \end{aligned} \right\} \quad (449)$$

Now when the solid is subject to uniform pressure on all sides, if we consider so much of it as has the volume unity in the state of reference, we shall have

$$r_1 = r_2 = r_3 = v^{\frac{1}{3}}, \quad (450)$$

and by (444) and (439),

$$\psi_{v'} = i + 3e v^{\frac{2}{3}} + 3f' v^{\frac{4}{3}} + h v. \quad (451)$$

Hence, by equation (88), since $\psi_{v'}$ is equivalent to ψ ,

$$-p = \left(\frac{d\psi}{dv}\right)_t = 2e v^{-\frac{1}{3}} + 4f' v^{\frac{1}{3}} + h, \quad (452)$$

$$-v \left(\frac{dp}{dv}\right)_t = -\frac{2}{3}e v^{-\frac{4}{3}} + \frac{4}{3}f' v^{-\frac{2}{3}}; \quad (453)$$

and by (448),

$$V = -\frac{2}{3}\frac{e}{r_0} + \frac{4}{3}f'r_0. \quad (454)$$

To obtain the value of R in accordance with the definition (449), we may suppose the values of E , F , and H given by equations (432), (434), and (437) to be substituted in equation (444). This will give for the value of R

$$R = 2e + 4f'r_0^2. \quad (455)$$

Moreover, since p must vanish in (452) when $v = r_0^3$, we have

$$2e + 4f'r_0^2 + hr_0 = 0. \quad (456)$$

From the three last equations may be obtained the values of e , f' , h , in terms of r_0 , V , and R ; viz.,

$$e = \frac{R - 3r_0 V}{4}, \quad f' = \frac{R + 3r_0 V}{8r_0^2}, \quad h = -\frac{R}{r_0}. \quad (457)$$

The quantity r_0 , like R and V , is a function of the temperature, the differential coefficient $\frac{d \log r_0}{dt}$ representing the rate of linear expansion of the solid when without stress.

It will not be necessary to discuss equation (443) at length, as the case is entirely analogous to that which has just been treated. [It must be remembered that η_{v_i} , in the discussion of (443) will take the place everywhere of the temperature in the discussion of (444).] If we denote by V' and R' the *elasticity of volume* and the *rigidity*, both determined under the condition of *constant entropy*, (i. e., of *no transmission of heat*), and for states of vanishing stress, we shall have the equations:

$$V' = -\frac{2e'}{3r_0} + \frac{4}{3}f'r_0, \tag{458}$$

$$R' = 2e' + 4f'r_0^2, \tag{459}$$

$$2e' + 4f'r_0^2 + h'r_0 = 0. \tag{460}$$

Whence

$$e' = \frac{R' - 3r_0V'}{4}, \quad f' = \frac{R' + 3r_0V'}{8r_0^2}, \quad h' = -\frac{R'}{r_0}. \tag{461}$$

In these equations r_0 , R' , and V' are to be regarded as functions of the quantity η_{v_i} .

If we wish to change from one state of reference to another (also isotropic), the changes required in the fundamental equation are easily made. If a denotes the length of any line of the solid in the second state of reference divided by its length in the first, it is evident that when we change from the first state of reference to the second the values of the symbols ε_{v_i} , η_{v_i} , ψ_{v_i} , H are divided by a^3 , that of E by a^2 , and that of F by a^4 . In making the change of the state of reference, we must therefore substitute in the fundamental equation of the form (444) $a^3\psi_{v_i}$, a^2E , a^4F , a^3H for ψ_{v_i} , E , F , and H , respectively. In the fundamental equation of the form (443), we must make the analogous substitutions, and also substitute $a^3\eta_{v_i}$ for η_{v_i} . [It will be remembered that i' , e' , f' , and h' represent functions of η_{v_i} , and that it is only when their values in terms of η_{v_i} are substituted, that equation (443) becomes a fundamental equation.]

Concerning Solids which absorb Fluids.

There are certain bodies which are solid with respect to some of their components, while they have other components which are fluid. In the following discussion, we shall suppose both the solidity and the fluidity to be perfect, so far as any properties are concerned which can affect the conditions of equilibrium,—i. e., we shall suppose that the solid matter of the body is entirely free from plasticity, and that there are no passive resistances to the motion of the fluid

components except such as vanish with the velocity of the motion,—leaving it to be determined by experiment how far and in what cases these suppositions are realized.

It is evident that equation (356) must hold true with regard to such a body, when the quantities of the fluid components contained in a given element of the solid remain constant. Let $\Gamma'_a, \Gamma'_b,$ etc., denote the quantities of the several fluid components contained in an element of the body divided by the volume of the element in the state of reference, or, in other words, let these symbols denote the densities which the several fluid components would have, if the body should be brought to the state of reference while the matter contained in each element remained unchanged. We may then say that equation (356) will hold true, when $\Gamma'_a, \Gamma'_b,$ etc., are constant. The complete value of the differential of $\varepsilon_v,$ will therefore be given by an equation of the form

$$d\varepsilon_v = t d\eta_v + \sum \sum' \left(X_x, d\frac{dx}{dx'} \right) + L_a d\Gamma'_a + L_b d\Gamma'_b + \text{etc.} \quad (462)$$

Now when the body is in a state of hydrostatic stress, the term in this equation containing the signs of summation will reduce to $-p dv_v,$ ($v_v,$ denoting, as elsewhere, the volume of the element divided by its volume in the state of reference). For in this case

$$X_x = -p \left(\frac{dy}{dy'} \frac{dz}{dz'} - \frac{dz}{dy'} \frac{dy}{dz'} \right), \quad (463)$$

$$\sum \sum' \left(X_x, d\frac{dx}{dx'} \right) = -p \sum \sum' \left\{ \left(\frac{dy}{dy'} \frac{dz}{dz'} - \frac{dz}{dy'} \frac{dy}{dz'} \right) d\frac{dx}{dx'} \right\}$$

$$= -p d \begin{vmatrix} \frac{dx}{dx'} & \frac{dx}{dy'} & \frac{dx}{dz'} \\ \frac{dy}{dx'} & \frac{dy}{dy'} & \frac{dy}{dz'} \\ \frac{dz}{dx'} & \frac{dz}{dy'} & \frac{dz}{dz'} \end{vmatrix}$$

$$= -p dv_v. \quad (464)$$

We have, therefore, for a state of hydrostatic stress,

$$d\varepsilon_v = t d\eta_v - p dv_v + L_a d\Gamma'_a + L_b d\Gamma'_b + \text{etc.}, \quad (465)$$

and multiplying by the volume of the element in the state of reference, which we may regard as constant,

$$d\varepsilon = t d\eta - p dv + L_a dm_a + L_b dm_b + \text{etc.}, \quad (466)$$

where ε , η , v , m_a , m_b , etc., denote the energy, entropy, and volume of the element, and the quantities of its several fluid components. It is evident that the equation will also hold true, if these symbols are understood as relating to a homogeneous body of finite size. The only limitation with respect to the variations is that the element or body to which the symbols relate shall always contain the same solid matter. The varied state may be one of hydrostatic stress or otherwise.

But when the body is in a state of hydrostatic stress, and the solid matter is considered invariable, we have by equation (12)

$$d\varepsilon = t d\eta - p dv + \mu_a dm_a + \mu_b dm_b + \text{etc.} \quad (467)$$

It should be remembered that the equation cited occurs in a discussion which relates only to bodies of hydrostatic stress, so that the varied state as well as the initial is there regarded as one of hydrostatic stress. But a comparison of the two last equations shows that the last will hold true without any such limitation, and moreover, that the quantities L_a , L_b , etc., when determined for a state of hydrostatic stress, are equal to the potentials μ_a , μ_b , etc.

Since we have hitherto used the term *potential* solely with reference to bodies of hydrostatic stress, we may apply this term as we choose with regard to other bodies. We may therefore call the quantities L_a , L_b , etc., the *potentials* for the several fluid components in the body considered, whether the state of the body is one of hydrostatic stress or not, since this use of the term involves only an extension of its former definition. It will also be convenient to use our ordinary symbol for a potential to represent these quantities. Equation (462) may then be written

$$d\varepsilon_v = t d\eta_v + \sum \sum' \left(\Lambda_{x'} d\frac{dx'}{x'} \right) + \mu_a d\Gamma'_a + \mu_b d\Gamma'_b + \text{etc.} \quad (468)$$

This equation holds true of solids having fluid components without any limitation with respect to the initial state or to the variations, except that the solid matter to which the symbols relate shall remain the same.

In regard to the conditions of equilibrium for a body of this kind, it is evident in the first place that if we make Γ'_a , Γ'_b , etc., constant, we shall obtain from the general criterion of equilibrium all the conditions which we have obtained for ordinary solids, and which are expressed by the formulæ (364), (374), (380), (382)–(384). The quantities Γ'_1 , Γ'_2 , etc., in the last two formulæ include of course

those which have just been represented by Γ'_a, Γ'_b , etc., and which relate to the fluid components of the body, as well as the corresponding quantities relating to its solid components. Again, if we suppose the solid matter of the body to remain without variation in quantity or position, it will easily appear that the potentials for the substances which form the fluid components of the solid body must satisfy the same conditions in the solid body and in the fluids in contact with it, as in the case of entirely fluid masses. See eqs. (22).

The above conditions must however be slightly modified in order to make them sufficient for equilibrium. It is evident that if the solid is dissolved at its surface, the fluid components which are set free may be absorbed by the solid as well as by the fluid mass, and in like manner if the quantity of the solid is increased, the fluid components of the new portion may be taken from the previously existing solid mass. Hence, whenever the *solid* components of the solid body are actual components of the fluid mass, (whether the case is the same with the *fluid* components of the solid body or not,) an equation of the form (383) must be satisfied, in which the potentials μ_a, μ_b , etc., contained implicitly in the second member of the equation are determined from the solid body. Also if the *solid* components of the solid body are all possible but not all actual components of the fluid mass, a condition of the form (384) must be satisfied, the values of the potentials in the second member being determined as in the preceding case.

The quantities

$$t, X_{x_1}, \dots Z_{z_1}, \mu_a, \mu_b, \text{ etc.}, \quad (469)$$

being differential coefficients of ε_v , with respect to the variables

$$\eta_v, \frac{dx}{dx'}, \dots \frac{dz}{dz'}, \Gamma'_a, \Gamma'_b, \text{ etc.}, \quad (470)$$

will of course satisfy the necessary relations

$$\frac{dt}{d\frac{dx}{dx'}} = \frac{dX_{x_1}}{d\eta_v}, \text{ etc.} \quad (471)$$

This result may be generalized as follows. Not only is the second member of equation (468) a complete differential in its present form, but it will remain such if we transfer the sign of differentiation (d) from one factor to the other of any term (the sum indicated by the symbol $\sum \Sigma'$ is here supposed to be expanded into nine terms), and at the same time change the sign of the term from $+$ to $-$. For to

substitute $-\eta_{V_i} dt$ for $t d\eta_{V_i}$, for example, is equivalent to subtracting the complete differential $d(t\eta_{V_i})$. Therefore, if we consider the quantities in (469) and (470) which occur in any same term in equation (468) as forming a pair, we may choose as independent variables either quantity of each pair, and the differential coefficient of the remaining quantity of any pair with respect to the independent variable of another pair will be equal to the differential coefficient of the remaining quantity of the second pair with respect to the independent variable of the first, taken positively, if the independent variables of these pairs are both affected by the sign d in equation (468), or are neither thus affected, but otherwise taken negatively. Thus

$$\left(\frac{dX_{X_i}}{dI'_a}\right)_{dx} = \left(\frac{d\mu_a}{d\frac{dx}{dx'}}\right)_{\Gamma_a'}, \quad \left(\frac{dX_{X_i}}{d\mu_a}\right)_{dx} = -\left(\frac{dI'_a}{d\frac{dx}{dx'}}\right)_{\mu_a}, \quad (472)$$

$$\left(\frac{d\frac{dx}{dx'}}{d\mu_a}\right)_{X_{X_i}} = \left(\frac{dI'_a}{dX_{X_i}}\right)_{\mu_a}, \quad \left(\frac{d\frac{dx}{dx'}}{dI'_a}\right)_{X_{X_i}} = -\left(\frac{d\mu_a}{dX_{X_i}}\right)_{\Gamma_a'}, \quad (473)$$

where in addition to the quantities indicated by the suffixes, the following are to be considered as constant: either t or η_{V_i} , either X_{Y_i} , or $\frac{dx}{dy'}$, . . . either Z_{Z_i} , or $\frac{dz}{dz'}$, either μ_b or I'_b , etc.

It will be observed that when the temperature is constant the conditions $\mu_a = \text{const.}$, $\mu_b = \text{const.}$ represent the physical condition of a body in contact with a fluid of which the phase does not vary, and which contains the components to which the potentials relate. Also that when I'_a , I'_b , etc., are constant, the heat absorbed by the body in any infinitesimal change of condition per unit of volume measured in the state of reference is represented by $t d\eta_{V_i}$. If we denote this quantity by dQ_{V_i} , and use the suffix Q to denote the condition of no transmission of heat, we may write

$$\left(\frac{d \log t}{d\frac{dx}{dx'}}\right)_Q = \left(\frac{dX_{X_i}}{dQ_{V_i}}\right)_{dx}, \quad \left(\frac{d \log t}{dX_{X_i}}\right)_Q = -\left(\frac{d\frac{dx}{dx'}}{dQ_{V_i}}\right)_{X_{X_i}}, \quad (474)$$

$$\left(\frac{dQ_{V_i}}{dX_{X_i}}\right)_t = \left(\frac{d\frac{dx}{dx'}}{d \log t}\right)_{X_{X_i}}, \quad \left(\frac{dQ_{V_i}}{d\frac{dx}{dx'}}\right)_t = -\left(\frac{dX_{X_i}}{d \log t}\right)_{dx}, \quad (475)$$

where I'_a , I'_b , etc., must be regarded as constant in all the equations, and either X_{Y_i} , or $\frac{dx}{dy'}$, . . . either Z_{Z_i} , or $\frac{dz}{dz'}$, in each equation.

INFLUENCE OF SURFACES OF DISCONTINUITY UPON THE EQUILIBRIUM OF HETEROGENEOUS MASSES.—THEORY OF CAPILLARITY.

We have hitherto supposed, in treating of heterogeneous masses in contact, that they might be considered as separated by mathematical surfaces, each mass being unaffected by the vicinity of the others, so that it might be homogeneous quite up to the separating surfaces both with respect to the density of each of its various components and also with respect to the densities of energy and entropy. That such is not rigorously the case is evident from the consideration that if it were so with respect to the densities of the components it could not be so in general with respect to the density of energy, as the sphere of molecular action is not infinitely small. But we know from observation that it is only within very small distances of such a surface that any mass is sensibly affected by its vicinity,—a natural consequence of the exceedingly small sphere of sensible molecular action,—and this fact renders possible a simple method of taking account of the variations in the densities of the component substances and of energy and entropy, which occur in the vicinity of surfaces of discontinuity. We may use this term, for the sake of brevity, without implying that the discontinuity is absolute, or that the term distinguishes any surface with mathematical precision. It may be taken to denote the non-homogeneous film which separates homogeneous or nearly homogeneous masses.

Let us consider such a surface of discontinuity in a fluid mass which is in equilibrium and uninfluenced by gravity. For the precise measurement of the quantities with which we have to do, it will be convenient to be able to refer to a geometrical surface, which shall be sensibly coincident with the physical surface of discontinuity, but shall have a precisely determined position. For this end, let us take some point in or very near to the physical surface of discontinuity, and imagine a geometrical surface to pass through this point and all other points which are similarly situated with respect to the condition of the adjacent matter. Let this geometrical surface be called the *dividing surface*, and designated by the symbol S . It will be observed that the position of this surface is as yet to a certain extent arbitrary, but that the directions of its normals are already everywhere determined, since all the surfaces which can be formed in the manner described are evidently parallel to one another. Let us also imagine a closed surface cutting the surface S and including a part of the homogeneous mass on each side. We will so far limit the

form of this closed surface as to suppose that on each side of S , as far as there is any want of perfect homogeneity in the fluid masses, the closed surface is such as may be generated by a moving normal to S . Let the portion of S which is included by the closed surface be denoted by \mathfrak{s} , and the area of this portion by s . Moreover, let the mass contained within the closed surface be divided into three parts by two surfaces, one on each side of S , and very near to that surface, although at such distance as to lie entirely beyond the influence of the discontinuity in its vicinity. Let us call the part which contains the surface \mathfrak{s} (with the physical surface of discontinuity) M , and the homogeneous parts M' and M'' , and distinguish by $\varepsilon, \varepsilon', \varepsilon'', \eta, \eta', \eta'', m_1, m_1', m_1'', m_2, m_2', m_2'',$ etc., the energies and entropies of these masses, and the quantities which they contain of their various components.

It is necessary, however, to define more precisely what is to be understood in cases like the present by the energy of masses which are only separated from other masses by imaginary surfaces. A part of the total energy which belongs to the matter in the vicinity of the separating surface, relates to pairs of particles which are on different sides of the surface, and such energy is not in the nature of things referable to either mass by itself. Yet, to avoid the necessity of taking separate account of such energy, it will often be convenient to include it in the energies which we refer to the separate masses. When there is no break in the homogeneity at the surface, it is natural to treat the energy as distributed with a uniform density. This is essentially the case with the initial state of the system which we are considering, for it has been divided by surfaces passing in general through homogeneous masses. The only exception—that of the surface which cuts at right angles the non-homogeneous film—(apart from the consideration that without any important loss of generality we may regard the part of this surface within the film as very small compared with the other surfaces) is rather apparent than real, as there is no change in the state of the matter *in the direction perpendicular to this surface*. But in the variations to be considered in the state of the system, it will not be convenient to limit ourselves to such as do not create any discontinuity at the surfaces bounding the masses M, M', M'' : we must therefore determine how we will estimate the energies of the masses in case of such infinitesimal discontinuities as may be supposed to arise. Now the energy of each mass will be most easily estimated by neglecting the discontinuity, i. e., if we estimate the energy on the supposition that

beyond the bounding surface the phase is identical with that within the surface. This will evidently be allowable, if it does not affect the total amount of energy. To show that it does not affect this quantity, we have only to observe that, if the energy of the mass on one side of a surface where there is an infinitesimal discontinuity of phase is greater as determined by this rule than if determined by any other (suitable) rule, the energy of the mass on the other side must be less by the same amount when determined by the first rule than when determined by the second, since the discontinuity relative to the second mass is equal but opposite in character to the discontinuity relative to the first.

If the entropy of the mass which occupies any one of the spaces considered is not in the nature of things determined without reference to the surrounding masses, we may suppose a similar method to be applied to the estimation of entropy.

With this understanding, let us return to the consideration of the equilibrium of the three masses M , M' , and M'' . We shall suppose that there are no limitations to the possible variations of the system due to any want of perfect mobility of the components by means of which we express the composition of the masses, and that these components are independent, i. e., that no one of them can be formed out of the others.

With regard to the mass M , which includes the surface of discontinuity, it is necessary for its internal equilibrium that when its boundaries are considered constant, and when we consider only *reversible* variations (i. e., those of which the opposite are also possible), the variation of its energy should vanish with the variations of its entropy and of the quantities of its various components. For changes within this mass will not affect the energy or the entropy of the surrounding masses (when these quantities are estimated on the principle which we have adopted), and it may therefore be treated as an isolated system. For fixed boundaries of the mass M , and for reversible variations, we may therefore write

$$\delta\varepsilon = A_0 \delta\eta + A_1 \delta m_1 + A_2 \delta m_2 + \text{etc.}, \quad (476)$$

where A_0 , A_1 , A_2 , etc., are quantities determined by the initial (unvaried) condition of the system. It is evident that A_0 is the temperature of the lamelliform mass to which the equation relates, or the *temperature at the surface of discontinuity*. By comparison of this equation with (12) it will be seen that the definition of A_1 , A_2 , etc., is entirely analogous to that of the potentials in homo-

genous masses, although the mass to which the former quantities relate is not homogeneous, while in our previous definition of potentials, only homogeneous masses were considered. By a natural extension of the term *potential*, we may call the quantities A_1, A_2 , etc., the *potentials at the surface of discontinuity*. This designation will be farther justified by the fact, which will appear hereafter, that the value of these quantities is independent of the thickness of the lamina (M) to which they relate. If we employ our ordinary symbols for temperature and potentials, we may write

$$\delta\varepsilon = t \delta\eta + \mu_1 \delta m_1 + \mu_2 \delta m_2 + \text{etc.} \quad (477)$$

If we substitute \geq for $=$ in this equation, the formula will hold true of all variations whether reversible or not;* for if the variation of energy could have a value less than that of the second member of the equation, there must be variation in the condition of M in which its energy is diminished without change of its entropy or of the quantities of its various components.

It is important, however, to observe that for any given values of $\delta\eta, \delta m_1, \delta m_2$, etc., while there *may* be possible variations of the nature and state of M for which the value of $\delta\varepsilon$ is greater than that of the second member of (477), there *must* always be possible variations for which the value of $\delta\varepsilon$ is equal to that of the second member.

* To illustrate the difference between variations which are reversible, and those which are not, we may conceive of two entirely different substances meeting in equilibrium at a mathematical surface without being at all mixed. We may also conceive of them as mixed in a thin film about the surface where they meet, and then the amount of mixture is capable of variation both by increase and by diminution. But when they are absolutely unmixed, the amount of mixture can be increased, but is incapable of diminution, and it is then consistent with equilibrium that the value of $\delta\varepsilon$ (for a variation of the system in which the substances commence to mix) should be greater than the second member of (477). It is not necessary to determine whether precisely such cases actually occur; but it would not be legitimate to overlook the possible occurrence of cases in which variations may be possible while the opposite variations are not.

It will be observed that the sense in which the term *reversible* is here used is entirely different from that in which it is frequently used in treatises on thermodynamics, where a process by which a system is brought from a state A to a state B is called reversible, to signify that the system may also be brought from the state B to the state A through the same series of intermediate states taken in the reverse order by means of external agencies of the opposite character. The variation of a system from a state A to a state B (supposed to differ infinitely little from the first) is here called reversible when the system is capable of another state B' which bears the same relation to the state A that A bears to B.

It will be convenient to have a notation which will enable us to express this by an equation. Let $\delta\varepsilon$ denote the smallest value (i. e., the value nearest to $-\infty$) of $\delta\varepsilon$ consistent with given values of the other variations, then

$$\delta\varepsilon = t \delta\eta + \mu_1 \delta m_1 + \mu_2 \delta m_2 + \text{etc.} \quad (478)$$

For the internal equilibrium of the whole mass which consists of the parts M, M', M'', it is necessary that

$$\delta\varepsilon + \delta\varepsilon' + \delta\varepsilon'' \geq 0 \quad (479)$$

for all variations which do not affect the enclosing surface or the total entropy or the total quantity of any of the various components. If we also regard the surfaces separating M, M', and M'' as invariable, we may derive from this condition, by equations (478) and (12), the following as a *necessary* condition of equilibrium:

$$\begin{aligned} & t \delta\eta + \mu_1 \delta m_1 + \mu_2 \delta m_2 + \text{etc.} \\ & + t' \delta\eta' + \mu_1' \delta m_1' + \mu_2' \delta m_2' + \text{etc.} \\ & + t'' \delta\eta'' + \mu_1'' \delta m_1'' + \mu_2'' \delta m_2'' + \text{etc.} \geq 0, \end{aligned} \quad (480)$$

the variations being subject to the equations of conditions

$$\left. \begin{aligned} \delta\eta + \delta\eta' + \delta\eta'' &= 0, \\ \delta m_1 + \delta m_1' + \delta m_1'' &= 0, \\ \delta m_2 + \delta m_2' + \delta m_2'' &= 0, \\ \text{etc.} \end{aligned} \right\} \quad (481)$$

It may also be the case that some of the quantities $\delta m_1'$, $\delta m_1''$, $\delta m_2'$, $\delta m_2''$, etc., are incapable of negative values or can only have the value zero. This will be the case when the substances to which these quantities relate are not actual or possible components of M' or M''. (See page 117.) To satisfy the above condition it is necessary and sufficient that

$$t = t' = t'', \quad (482)$$

$$\mu_1' \delta m_1' \geq \mu_1 \delta m_1', \quad \mu_2' \delta m_2' \geq \mu_2 \delta m_2', \quad \text{etc.}, \quad (483)$$

$$\mu_1'' \delta m_1'' \geq \mu_1 \delta m_1'', \quad \mu_2'' \delta m_2'' \geq \mu_2 \delta m_2'', \quad \text{etc.} \quad (484)$$

It will be observed that, if the substance to which μ_1 , for instance, relates is an actual component of each of the homogeneous masses, we shall have $\mu_1 = \mu_1' = \mu_1''$. If it is an actual component of the first only of these masses, we shall have $\mu_1 = \mu_1'$. If it is also a possible component of the second homogeneous mass, we shall also have $\mu_1 \geq \mu_1''$. If this substance occurs only at the surface of dis-

continuity, the value of the potential μ_1 will not be determined by any equation, but cannot be greater than the potential for the same substance in either of the homogeneous masses in which it may be a possible component.

It appears, therefore, that the particular conditions of equilibrium relating to temperature and the potentials which we have before obtained by neglecting the influence of the surfaces of discontinuity (pp. 119, 120, 128) are not invalidated by the influence of such discontinuity in their application to homogeneous parts of the system bounded like M' and M'' by imaginary surfaces lying within the limits of homogeneity,—a condition which may be fulfilled by surfaces very near to the surfaces of discontinuity. It appears also that similar conditions will apply to the non-homogeneous films like M' , which separate such homogeneous masses. The properties of such films, which are of course different from those of homogeneous masses, require our farther attention.

The volume occupied by the mass M is divided by the surface s into two parts, which we will call v''' and v'''' , v''' lying next to M' , and v'''' to M'' . Let us imagine these volumes filled by masses having throughout the same temperature, pressure and potentials, and the same densities of energy and entropy, and of the various components, as the masses M' and M'' respectively. We shall then have, by equation (12), if we regard the volumes as constant,

$$\delta\varepsilon''' = t' \delta\eta''' + \mu_1' \delta m_1''' + \mu_2' \delta m_2''' + \text{etc.}, \quad (485)$$

$$\delta\varepsilon'''' = t'' \delta\eta'''' + \mu_1'' \delta m_1'''' + \mu_2'' \delta m_2'''' + \text{etc.}; \quad (486)$$

whence, by (482)–(484), we have for reversible variations

$$\delta\varepsilon''' = t \delta\eta''' + \mu_1 \delta m_1''' + \mu_2 \delta m_2''' + \text{etc.}, \quad (487)$$

$$\delta\varepsilon'''' = t \delta\eta'''' + \mu_1 \delta m_1'''' + \mu_2 \delta m_2'''' + \text{etc.} \quad (488)$$

From these equations and (477), we have for reversible variations

$$\begin{aligned} \delta(\varepsilon - \varepsilon''' - \varepsilon''') &= t \delta(\eta - \eta''' - \eta''') \\ + \mu_1 \delta(m_1 - m_1''' - m_1''') &+ \mu_2 \delta(m_2 - m_2''' - m_2''') + \text{etc.} \end{aligned} \quad (489)$$

Or, if we set*

$$\varepsilon^s = \varepsilon - \varepsilon''' - \varepsilon''', \quad \eta^s = \eta - \eta''' - \eta''', \quad (490)$$

$$m_1^s = m_1 - m_1''' - m_1''', \quad m_2^s = m_2 - m_2''' - m_2''', \quad \text{etc.}, \quad (491)$$

* It will be understood that the s here used is not an algebraic exponent, but is only intended as a distinguishing mark. The Roman letter S has not been used to denote any quantity.

we may write

$$\delta \epsilon^s = t \delta \eta^s + \mu_1 \delta m_1^s + \mu_2 \delta m_2^s + \text{etc.} \quad (492)$$

This is true of reversible variations in which the surfaces which have been considered are fixed. It will be observed that ϵ^s denotes the excess of the energy of the actual mass which occupies the total volume which we have considered over that energy which it would have, if on each side of the surface S the density of energy had the same uniform value quite up to that surface which it has at a sensible distance from it; and that η^s , m_1^s , m_2^s , etc., have analogous significations. It will be convenient, and need not be a source of any misconception, to call ϵ^s and η^s the energy and entropy of the surface (or the *superficial* energy and entropy), $\frac{\epsilon^s}{s}$ and $\frac{\eta^s}{s}$ the *superficial densities* of energy and entropy, $\frac{m_1^s}{s}$, $\frac{m_2^s}{s}$, etc., the *superficial densities* of the several components.

Now these quantities (ϵ^s , η^s , m_1^s , etc.) are determined partly by the state of the physical system which we are considering, and partly by the various imaginary surfaces by means of which these quantities have been defined. The position of these surfaces, it will be remembered, has been regarded as fixed in the variation of the system. It is evident, however, that the form of that portion of these surfaces, which lies in the region of homogeneity on either side of the surface of discontinuity cannot affect the values of these quantities. To obtain the complete value of $\delta \epsilon^s$ for reversible variations, we have therefore only to regard variations in the position and form of the limited surface s , as this determines all of the surfaces in question lying within the region of non-homogeneity. Let us first suppose the form of s to remain unvaried and only its position in space to vary, either by translation or rotation. No change in (492) will be necessary to make it valid in this case. For the equation is valid if s remains fixed and the material system is varied in position; also, if the material system and s are both varied in position, while their relative position remains unchanged. Therefore, it will be valid if the surface alone varies its position.

But if the form of s be varied, we must add to the second member (492) terms which shall represent the value of

$$\delta \epsilon^s - t \delta \eta^s - \mu_1 \delta m_1^s - \mu_2 \delta m_2^s - \text{etc.}$$

due to such variation in the form of s . If we suppose s to be suffi-

ciently small to be considered uniform throughout in its curvatures and in respect to the state of the surrounding matter, the value of the above expression will be determined by the variation of its area δs and the variations of its principal curvatures δc_1 and δc_2 , and we may write

$$\delta \varepsilon^s = t \delta \eta^s + \mu_1 \delta m_1^s + \mu_2 \delta m_2^s + \text{etc.} \\ + \sigma \delta s + C_1 \delta c_1 + C_2 \delta c_2, \quad (493)$$

or

$$\delta \varepsilon^s = t \delta \eta^s + \mu_1 \delta m_1^s + \mu_2 \delta m_2^s + \text{etc.} \\ + \sigma \delta s + \frac{1}{2} (C_1 + C_2) \delta (c_1 + c_2) + \frac{1}{2} (C_1 - C_2) \delta (c_1 - c_2), \quad (494)$$

σ , C_1 , and C_2 denoting quantities which are determined by the initial state of the system and position and form of \mathfrak{s} . The above is the complete value of the variation of ε^s for reversible variations of the system. But it is always possible to give such a position to the surface \mathfrak{s} that $C_1 + C_2$ shall vanish.

To show this, it will be convenient to write the equation in the longer form [see (490), (491)]

$$\delta \varepsilon - t \delta \eta - \mu_1 \delta m_1 - \mu_2 \delta m_2 - \text{etc.} \\ - \delta \varepsilon''' + t \delta \eta''' + \mu_1 \delta m_1''' + \mu_2 \delta m_2''' + \text{etc.} \\ - \delta \varepsilon'''' + t \delta \eta'''' + \mu_1 \delta m_1'''' + \mu_2 \delta m_2'''' + \text{etc.} \\ = \sigma \delta s + \frac{1}{2} (C_1 + C_2) \delta (c_1 + c_2) + \frac{1}{2} (C_1 - C_2) \delta (c_1 - c_2), \quad (495)$$

i. e., by (482)-(484) and (12),

$$\delta \varepsilon - t \delta \eta - \mu_1 \delta m_1 - \mu_2 \delta m_2 - \text{etc.} + p' \delta v''' + p'' \delta v'''' \\ = \sigma \delta s + \frac{1}{2} (C_1 + C_2) \delta (c_1 + c_2) + \frac{1}{2} (C_3 - C_2) \delta (c_1 - c_2). \quad (496)$$

From this equation it appears in the first place that the pressure is the same in the two homogeneous masses separated by a plane surface of discontinuity. For let us imagine the material system to remain unchanged, while the plane surface \mathfrak{s} without change of area or of form moves in the direction of its normal. As this does not affect the boundaries of the mass M ,

$$\delta \varepsilon - t \delta \eta - \mu_1 \delta m_1 - \mu_2 \delta m_2 - \text{etc.} = 0.$$

Also $\delta s = 0$, $\delta (c_1 + c_2) = 0$, $\delta (c_1 - c_2) = 0$, and $\delta v''' = -\delta v''''$. Hence $p' = p''$, when the surface of discontinuity is plane.

Let us now examine the effect of different positions of the surface \mathfrak{s} in the same material system upon the value of $C_1 + C_2$, supposing at first that in the initial state of the system the surface of discontinuity is plane. Let us give the surface \mathfrak{s} some particular position. In the

initial state of the system this surface will of course be plane like the physical surface of discontinuity, to which it is parallel. In the varied state of the system, let it become a portion of a spherical surface having positive curvature; and at sensible distances from this surface let the matter be homogeneous and with the same phases as in the initial state of the system; also at and about the surface let the state of the matter so far as possible be the same as at and about the plane surface in the initial state of the system. (Such a variation in the system may evidently take place negatively as well as positively, as the surface may be curved toward either side. But whether such a variation is consistent with the maintenance of equilibrium is of no consequence, since in the preceding equations only the initial state is supposed to be one of equilibrium.) Let the surface \mathbf{s} , placed as supposed, whether in the initial or the varied state of the surface, be distinguished by the symbol \mathbf{s}' . Without changing either the initial or the varied state of the material system, let us make another supposition with respect to the imaginary surface \mathbf{s} . In the unvaried system let it be parallel to its former position but removed from it a distance λ on the side on which lie the centers of positive curvature. In the varied state of the system, let it be spherical and concentric with \mathbf{s}' , and separated from it by the same distance λ . It will of course lie on the same side of \mathbf{s}' as in the unvaried system. Let the surface \mathbf{s} , placed in accordance with this second supposition, be distinguished by the symbol \mathbf{s}'' . Both in the initial and the varied state, let the perimeters of \mathbf{s}' and \mathbf{s}'' be traced by a common normal. Now the value of

$$\delta\varepsilon - t\delta\eta - \mu_1\delta m_1 - \mu_2\delta m_2 - \text{etc.}$$

in equation (496) is not affected by the position of \mathbf{s} , being determined simply by the body \mathbf{M} : the same is true $p'\delta v''' + p''\delta v''''$ or $p'\delta(v''' + v''''), v''' + v''''$ being the volume of \mathbf{M} . Therefore the second member of (496) will have the same value whether the expressions relate to \mathbf{s}' or \mathbf{s}'' . Moreover, $\delta(c_1 - c_2) = 0$ both for \mathbf{s}' and \mathbf{s}'' . If we distinguish the quantities determined for \mathbf{s}' and for \mathbf{s}'' by the marks ' and '' , we may therefore write

$$\sigma'\delta s' + \frac{1}{2}(C_1' + C_2')\delta(c_1' + c_2') = \sigma''\delta s'' + \frac{1}{2}(C_1'' + C_2'')\delta(c_1'' + c_2'').$$

Now if we make $\delta s'' = 0$,

we shall have by geometrical necessity

$$\delta s' = s\lambda\delta(c_1'' + c_2'').$$

Hence

$$\sigma' s \lambda \delta(c_1'' + c_2'') + \frac{1}{2} (C_1' + C_2') \delta(c_1' + c_2') = \frac{1}{2} (C_2'' + C_2'') \delta(c_1'' + c_2'').$$

But
$$\delta(c_1' + c_2') = \delta(c_1'' + c_2'').$$

Therefore,
$$C_1' + C_2' + 2 \sigma' s \lambda = C_1'' + C_2''.$$

This equation shows that we may give a positive or negative value to $C_1'' + C_2''$ by placing \mathbf{s}'' a sufficient distance on one or on the other side of \mathbf{s}' . Since this is true when the (unvaried) surface is plane, it must also be true when the surface is nearly plane. And for this purpose a surface may be regarded as nearly plane, when the radii of curvature are very large in proportion to the thickness of the non-homogeneous film. This is the case when the radii of curvature have any sensible size. In general, therefore, whether the surface of discontinuity is plane or curved it is possible to place the surface \mathbf{s} so that $C_1 + C_2$ in equation (494) shall vanish.

Now we may easily convince ourselves by equation (493) that if \mathbf{s} is placed within the non-homogeneous film, and $s = 1$, the quantity σ is of the same order of magnitude as the values of ε^s , η^s , m_1^s , m_2^s , etc., while the values of C_1 and C_2 are of the same order of magnitude as the changes in the values of the former quantities caused by increasing the curvature of \mathbf{s} by unity. Hence, on account of the thinness of the non-homogeneous film, since it can be very little affected by such a change of curvature in \mathbf{s} , the values of C_1 and C_2 must in general be very small relatively to σ . And hence, if \mathbf{s}' be placed within the non-homogeneous film, the value of λ which will make $C_1'' + C_2''$ vanish must be very small (of the same order of magnitude as the thickness of the non-homogeneous film). The position of \mathbf{s} , therefore, which will make $C_1 + C_2$ in (494) vanish, will in general be sensibly coincident with the physical surface of discontinuity.

We shall hereafter suppose, when the contrary is not distinctly indicated that the surface \mathbf{s} , in the unvaried state of the system, has such a position as to make $C_1 + C_2 = 0$. It will be remembered that the surface \mathbf{s} is a part of a larger surface S , which we have called the *dividing surface*, and which is coextensive with the physical surface of discontinuity. We may suppose that the position of the dividing surface is everywhere determined by similar considerations. This is evidently consistent with the suppositions made on page 380 with regard to this surface.

We may therefore cancel the term

$$\frac{1}{2} (C_1 + C_2) \delta (c_1 + c_2)$$

in (494). In regard to the following term, it will be observed that C_1 must necessarily be equal to C_2 , when $c_1 = c_2$, which is the case when the surface of discontinuity is plane. Now on account of the thinness of the non-homogeneous film, we may always regard it as composed of parts which are approximately plane. Therefore, without danger of sensible error, we may also cancel the term

$$\frac{1}{2} (C_1 - C_2) \delta (c_1 - c_2).$$

Equation (494) is thus reduced to the form

$$\delta \varepsilon^s = t \delta \eta^s + \sigma \delta s + \mu_1 \delta m_1^s + \mu_2 \delta m_2^s + \text{etc.} \quad (497)$$

We may regard this as the complete value of $\delta \varepsilon^s$, for all reversible variations in the state of the system supposed initially in equilibrium, when the dividing surface has its initial position determined in the manner described.

The above equation is of fundamental importance in the theory of capillarity. It expresses a relation with regard to surfaces of discontinuity analogous to that expressed by equation (12) with regard to homogeneous masses. From the two equations may be directly deduced the conditions of equilibrium of heterogeneous masses in contact, subject or not to the action of gravity, without disregard of the influence of the surfaces of discontinuity. The general problem, including the action of gravity, we shall take up hereafter: at present we shall only consider, as hitherto, a small part of a surface of discontinuity with a part of the homogeneous mass on either side, in order to deduce the additional condition which may be found when we take account of the motion of the dividing surface.

We suppose as before that the mass especially considered is bounded by a surface of which all that lies in the region of non-homogeneity is such as may be traced by a moving normal to the dividing surface. But instead of dividing the mass as before into four parts, it will be sufficient to regard it as divided into two parts by the dividing surface. The energy, entropy, etc., of these parts, estimated on the supposition that its nature (including density of energy, etc.) is uniform quite up to the dividing surface, will be denoted by ε' , η' , etc., ε'' , η'' , etc. Then the total energy will be $\varepsilon^s + \varepsilon' + \varepsilon''$, and the general condition of internal equilibrium will be that

$$\delta \varepsilon^s + \delta \varepsilon' + \delta \varepsilon'' \geq 0, \quad (498)$$

when the bounding surface is fixed, and the total entropy and total quantities of the various components are constant. We may suppose $\eta^s, \eta', \eta'', m_1^s, m_1', m_1'', m_2^s, m_2', m_2'',$ etc., to be all constant. Then by (497) and (12) the condition reduces to

$$\sigma \delta s - p' \delta v' - p'' \delta v'' = 0. \tag{499}$$

(We may set $=$ for \cong , since changes in the position of the dividing surface can evidently take place in either of two opposite directions.) This equation has evidently the same form as if a membrane without rigidity and having a tension σ , uniform in all directions, existed at the dividing surface. Hence, the particular position which we have chosen for this surface may be called the surface of tension, and σ the superficial tension. If all parts of the dividing surface move a uniform normal distance δN , we shall have

$$\delta s = (c_1 + c_2) s \delta N, \quad \delta v' = s \delta N, \quad \delta v'' = -s \delta N;$$

whence
$$\sigma (c_1 + c_2) = p' - p'', \tag{500}$$

the curvatures being positive when their centers lie on the side to which p' relates. This is the condition which takes the place of that of equality of pressure (see pp. 119, 128) for heterogeneous fluid masses in contact, when we take account of the influence of the surfaces of discontinuity. We have already seen that the conditions relating to temperature and the potentials are not affected by these surfaces.

Fundamental Equations for Surfaces of Discontinuity.

In equation (497) the initial state of the system is supposed to be one of equilibrium. The only limitation with respect to the varied state is that the variation shall be reversible, i. e., that an opposite variation shall be possible. Let us now confine our attention to variations in which the system remains in equilibrium. To distinguish this case, we may use the character d instead δ , and write

$$d\varepsilon^s = t d\eta^s + \sigma ds + \mu_1 dm_1^s + \mu_2 dm_2^s + \text{etc.} \tag{501}$$

Both the states considered being states of equilibrium, the limitation with respect to the reversibility of the variations may be neglected, since the variations will always be reversible in at least one of the states considered.

If we integrate this equation, supposing the area s to increase from zero to any finite value s , while the material system to a part of which the equation relates remains without change, we obtain

$$\varepsilon^s = t \eta^s + \sigma s + \mu_1 m_1^s + \mu_2 m_2^s + \text{etc.}, \tag{502}$$

which may be applied to any portion of any surface of discontinuity (in equilibrium) which is of the same nature throughout, or throughout which the values of t , σ , μ_1 , μ_2 , etc. are constant.

If we differentiate this equation, regarding all the quantities as variable, and compare the result with (501), we obtain

$$\eta^s dt + s d\sigma + m_1^s d\mu_1 + m_2^s d\mu_2 + \text{etc.} = 0. \quad (503)$$

If we denote the *superficial densities* of energy, of entropy, and of the several component substances (see page 386) by ϵ_s , η_s , Γ_1 , Γ_2 , etc., we have

$$\epsilon_s = \frac{\epsilon^s}{s}, \quad \eta_s = \frac{\eta^s}{s}, \quad (504)$$

$$\Gamma_1 = \frac{m_1^s}{s}, \quad \Gamma_2 = \frac{m_2^s}{s}, \quad \text{etc.}, \quad (505)$$

and the preceding equations may be reduced to the form:—

$$d\epsilon_s = t d\eta_s + \mu_1 d\Gamma_1 + \mu_2 d\Gamma_2 + \text{etc.}, \quad (506)$$

$$\epsilon_s = t \eta_s + \sigma + \mu_1 \Gamma_1 + \mu_2 \Gamma_2 + \text{etc.}, \quad (507)$$

$$d\sigma = -\eta_s dt - \Gamma_1 d\mu_1 - \Gamma_2 d\mu_2 - \text{etc.} \quad (508)$$

Now the contact of the two homogeneous masses does not impose any restriction upon the variations of phase of either, except that the temperature and the potentials for actual components shall have the same value in both. [See (482)–(484) and (500).] For however the values of the pressures in the homogeneous masses may vary (on account of arbitrary variations of the temperature and potentials), and however the superficial tension may vary, equation (500) may always be satisfied by giving the proper curvature to the surface of tension, so long, at least, as the difference of pressures is not great. Moreover, if any of the potentials μ_1 , μ_2 , etc. relate to substances which are found only at the surface of discontinuity, their values may be varied by varying the superficial densities of those substances. The values of t , μ_1 , μ_2 , etc. are therefore independently variable, and it appears from equation (508) that σ is a function of these quantities. If the form of this function is known, we may derive from it by differentiation $n+1$ equations (n denoting the total number of component substances) giving the values of η_s , Γ_1 , Γ_2 , etc. in terms of the variables just mentioned. This will give us, with (507), $n+3$ independent equations between the $2n+4$ quantities which occur in that equation. These are all that exist, since $n+1$

of these quantities are independently variable. Or, we may consider that we have $n+3$ independent equations between the $2n+5$ quantities occurring in equation (502), of which $n+2$ are independently variable.

An equation, therefore, between

$$\sigma, t, \mu_1, \mu_2, \text{ etc.}, \tag{509}$$

may be called a fundamental equation for the surface of discontinuity. An equation between

$$\varepsilon^s, \eta^s, s, m_1^s, m_2^s, \text{ etc.}, \tag{510}$$

or between $\varepsilon_s, \eta_s, \Gamma_1, \Gamma_2, \text{ etc.},$ (511)

may also be called a fundamental equation in the same sense. For it is evident from (501) that an equation may be regarded as subsisting between the variables (510), and if this equation be known, since $n+2$ of the variables may be regarded as independent (viz., $n+1$ for the $n+1$ variations in the nature of the surface of discontinuity, and one for the area of the surface considered), we may obtain by differentiation and comparison with (501), $n+2$ additional equations between the $2n+5$ quantities occurring in (502). Equation (506) shows that equivalent relations can be deduced from an equation between the variables (511). It is moreover quite evident that an equation between the variables (510) must be reducible to the form of an equation between the ratios of these variables, and therefore to an equation between the variables (511).

The same designation may be applied to any equation from which, by differentiation and the aid only of general principles and relations, $n+3$ independent relations between the same $2n+5$ quantities may be obtained.

If we set
$$\psi^s = \varepsilon^s - t \eta^s, \tag{512}$$

we obtain by differentiation and comparison with (501)

$$d\psi^s = -\eta^s dt + \sigma ds + \mu_1 dm_1^s + \mu_2 dm_2^s + \text{etc.} \tag{513}$$

An equation, therefore, between $\psi^s, t, s, m_1^s, m_2^s, \text{ etc.},$ is a fundamental equation, and is to be regarded as entirely equivalent to either of the other fundamental equations which have been mentioned.

The reader will not fail to notice the analogy between these fundamental equations, which relate to surfaces of discontinuity, and those relating to homogeneous masses, which have been described on pages 140-144.

On the Experimental Determination of Fundamental Equations for Surfaces of Discontinuity.

When all the substances which are found at a surface of discontinuity are components of one or the other of the homogeneous masses, the potentials μ_1, μ_2 , etc., as well as the temperature, may be determined from these homogeneous masses.* The tension σ may be determined by means of the relation (500). But our measurements are practically confined to cases in which the difference of the pressures in the homogeneous masses is small; for with increasing differences of pressure the radii of curvature soon become too small for measurement. Therefore, although the equation $p' = p''$ (which is equivalent to an equation between t, μ_1, μ_2 , etc., since p' and p'' are both functions of these variables) may not be exactly satisfied in cases in which it is convenient to measure the tension, yet this equation is so nearly satisfied in all the measurements of tension which we can make, that we must regard such measurements as simply establishing the values of σ for values of t, μ_1, μ_2 , etc., which satisfy the equation $p' = p''$, but not as sufficient to establish the rate of change in the value of σ for variations of t, μ_1, μ_2 , etc., which are inconsistent with the equation $p' = p''$.

To show this more distinctly, let t, μ_2, μ_3 , etc. remain constant, then by (508) and (98)

$$d\sigma = -\Gamma_1 d\mu_1,$$

$$dp' = \gamma_1' d\mu_1,$$

$$dp'' = \gamma_1'' d\mu_1,$$

γ_1' and γ_1'' denoting the densities $\frac{m_1'}{v'}$ and $\frac{m_1''}{v''}$. Hence,

$$dp' - dp'' = (\gamma_1' - \gamma_1'') d\mu_1,$$

and $\Gamma_1 d(p' - p'') = (\gamma_1'' - \gamma_1') d\sigma$.

But by (500)

$$(c_1 + c_2) d\sigma + \sigma d(c_1 + c_2) = d(p' - p'').$$

Therefore,

$$\Gamma_1 (c_1 + c_2) d\sigma + \Gamma_1 \sigma d(c_1 + c_2) = (\gamma_1'' - \gamma_1') d\sigma,$$

or $\{\gamma_1'' - \gamma_1' - \Gamma_1 (c_1 + c_2)\} d\sigma = \Gamma_1 \sigma d(c_1 + c_2)$.

* It is here supposed that the thermodynamic properties of the homogeneous masses have already been investigated, and that the fundamental equations of these masses may be regarded as known.

Now $\Gamma_1(c_1 + c_2)$ will generally be very small compared with $\gamma_1'' - \gamma_1'$. Neglecting the former term, we have

$$\frac{d\sigma}{\sigma} = \frac{\Gamma_1}{\gamma_1'' - \gamma_1'} d(c_1 + c_2).$$

To integrate this equation, we may regard $\Gamma_1, \gamma_1', \gamma_1''$ as constant. This will give, as an approximate value,

$$\log \frac{\sigma}{\sigma'} = -\frac{\Gamma_1}{\gamma_1'' - \gamma_1'} (c_1 + c_2),$$

σ' denoting the value of σ when the surface is plane. From this it appears that when the radii of curvature have any sensible magnitude, the value of σ will be sensibly the same as when the surface is plane and the temperature and all the potentials except one have the same values, unless the component for which the potential has not the same value has very nearly the same density in the two homogeneous masses, in which case, the condition under which the variations take place is nearly equivalent to the condition that the pressures shall remain equal.

Accordingly, we cannot in general expect to determine the superficial density Γ_1 from its value $-\left(\frac{d\sigma}{d\mu_1}\right)_{t,\mu}$ * by measurements of superficial tensions. The case will be the same with Γ_2, Γ_3 , etc., and also with η_s , the superficial density of entropy.

The quantities $\epsilon_s, \eta_s, \Gamma_1, \Gamma_2$, etc. are evidently too small in general to admit of direct measurement. When one of the components, however, is found only at the surface of discontinuity, it may be more easy to measure its superficial density than its potential. But except in this case, which is of secondary interest, it will generally be easy to determine σ in terms of t, μ_1, μ_2 , etc., with considerable accuracy for plane surfaces, and extremely difficult or impossible to determine the fundamental equation more completely.

Fundamental Equations for Plane Surfaces of Discontinuity.

An equation giving σ in terms of t, μ_1, μ_2 , etc., which will hold true only so long as the surface of discontinuity is plane, may be called a fundamental equation for a plane surface of discontinuity. It will be interesting to see precisely what results can be obtained from such an equation, especially with respect to the energy and entropy

* The suffixed μ is used to denote that all the potentials except that occurring in the denominator of the differential coefficient are to be regarded as constant.

and the quantities of the component substances in the vicinity of the surface of discontinuity.

These results can be exhibited in a more simple form, if we deviate to a certain extent from the method which we have been following. The particular position adopted for the dividing surface (which determines the superficial densities) was chosen in order to make the term $\frac{1}{2}(C_1 + C_2) \delta(c_1 + c_2)$ in (494) vanish. But when the curvature of the surface is not supposed to vary, such a position of the dividing surface is not necessary for the simplification of the formula. It is evident that equation (501) will hold true for plane surfaces (supposed to remain such) without reference to the position of the dividing surfaces, except that it shall be parallel to the surface of discontinuity. We are therefore at liberty to choose such a position for the dividing surface as may for any purpose be convenient.

None of the equations (502)–(513), which are either derived from (501), or serve to define new symbols, will be affected by such a change in the position of the dividing surface. But the expressions $\varepsilon^s, \eta^s, m_1^s, m_2^s$, etc., as also $\varepsilon_s, \eta_s, \Gamma_1, \Gamma_2$, etc. and ψ^s , will of course have different values when the position of that surface is changed. The quantity σ , however, which we may regard as defined by equations (501), or, if we choose, by (502) or (507), will not be affected in value by such a change. For if the dividing surface be moved a distance λ measured normally and toward the side to which v'' relates, the quantities

$$\varepsilon_s, \eta_s, \Gamma_1, \Gamma_2, \text{ etc.},$$

will evidently receive the respective increments

$\lambda(\varepsilon_v'' - \varepsilon_v'), \lambda(\eta_v'' - \eta_v'), \lambda(\gamma_1'' - \gamma_1'), \lambda(\gamma_2'' - \gamma_2')$, etc., $\varepsilon_v', \varepsilon_v'', \eta_v', \eta_v''$ denoting the densities of energy and entropy in the two homogeneous masses. Hence, by equation (507), σ will receive the increment

$$\lambda(\varepsilon_v'' - \varepsilon_v') - t\lambda(\eta_v'' - \eta_v') - \mu_1\lambda(\gamma_1'' - \gamma_1') - \mu_2\lambda(\gamma_2'' - \gamma_2') - \text{etc.}$$

But by (93)

$$\begin{aligned} -p'' &= \varepsilon_v'' - t\eta_v'' - \mu_1\gamma_1'' - \mu_2\gamma_2'' - \text{etc.}, \\ -p' &= \varepsilon_v' - t\eta_v' - \mu_1\gamma_1' - \mu_2\gamma_2' - \text{etc.} \end{aligned}$$

Therefore, since $p' = p''$, the increment in the value of σ is zero. The value of σ is therefore independent of the position of the dividing surface, when this surface is plane. But when we call this quantity the superficial tension, we must remember that it will not have

its characteristic properties as a tension with reference to any arbitrary surface. Considered as a tension, its position is in the surface which we have called the surface of tension, and, strictly speaking, nowhere else. The positions of the dividing surface, however, which we shall consider, will not vary from the surface of tension sufficiently to make this distinction of any practical importance.

It is generally possible to place the dividing surface so that the total quantity of any desired component in the vicinity of the surface of discontinuity shall be the same as if the density of that component were uniform on each side quite up to the dividing surface. In other words, we may place the dividing surface so as to make any one of the quantities Γ_1, Γ_2 , etc., vanish. The only exception is with regard to a component which has the same density in the two homogeneous masses. With regard to a component which has very nearly the same density in the two masses such a location of the dividing surface might be objectionable, as the dividing surface might fail to coincide sensibly with the physical surface of discontinuity. Let us suppose that γ_1' is not equal (nor very nearly equal) to γ_1'' , and that the dividing surface is so placed as to make $\Gamma_1 = 0$. Then equation (508) reduces to

$$d\sigma = -\eta_{s(1)}dt - \Gamma'_{2(1)}d\mu_2 - \Gamma'_{3(1)}d\mu_3 - \text{etc.}, \quad (514)$$

where the symbols $\eta_{s(1)}, \Gamma'_{2(1)}$, etc., are used for greater distinctness to denote the values of η_s, Γ_2 , etc., as determined by a dividing surface placed so that $\Gamma_1 = 0$. Now we may consider all the differentials in the second member of this equation as independent, without violating the condition that the surface shall remain plane, i. e., that $dp' = dp''$. This appears at once from the values of dp' and dp'' given by equation (98). Moreover, as has already been observed, when the fundamental equations of the two homogeneous masses are known, the equation $p' = p''$ affords a relation between the quantities t, μ_1, μ_2 , etc. Hence, when the value of σ is also known for plane surfaces in terms of t, μ_1, μ_2 , etc., we can eliminate μ_1 from this expression by means of the relation derived from the equality of pressures, and obtain the value of σ for plane surfaces in terms of t, μ_2, μ_3 , etc. From this, by differentiation, we may obtain directly the values of $\eta_{s(1)}, \Gamma'_{2(1)}, \Gamma'_{3(1)}$, etc., in terms of t, μ_2, μ_3 , etc. This would be a convenient form of the fundamental equation. But, if the elimination of p', p'' , and μ_1 from the finite equations presents algebraic difficulties, we can in all cases easily eliminate $dp', dp'', d\mu_1$ from the corresponding differential equations and thus obtain a

differential equation from which the values of $\eta_{s(1)}$, $\Gamma_{2(1)}$, $\Gamma_{3(1)}$, etc. in terms of t , μ_1 , μ_2 , etc., may be at once obtained by comparison with (514).*

* If liquid mercury meets the mixed vapors of water and mercury in a plane surface, and we use μ_1 and μ_2 to denote the potentials of mercury and water respectively, and place the dividing surface so that $\Gamma_1=0$, i. e., so that the total quantity of mercury is the same as if the liquid mercury reached this surface on one side and the mercury vapor on the other without change of density on either side, then $\Gamma_{2(1)}$ will represent the amount of water in the vicinity of this surface, per unit of surface, above that which there would be, if the water-vapor just reached the surface without change of density, and this quantity (which we may call the quantity of water condensed upon the surface of the mercury) will be determined by the equation

$$\Gamma_{2(1)} = - \frac{d\sigma}{d\mu_2}.$$

(In this differential coefficient as well as the following, the temperature is supposed to remain constant and the surface of discontinuity plane. Practically, the latter condition may be regarded as fulfilled in the case of any ordinary curvatures.)

If the pressure in the mixed vapors conforms to the law of Dalton (see pp. 215, 218), we shall have for constant temperature

$$dp_2 = \gamma_2 d\mu_2,$$

where p_2 denotes the part of the pressure in the vapor due to the water-vapor, and γ_2 the density of the water-vapor. Hence we obtain

$$\Gamma_{2(1)} = - \gamma_2 \frac{d\sigma}{dp_2}.$$

For temperatures below 100° centigrade, this will certainly be accurate, since the pressure due to the vapor of mercury may be neglected.

The value of σ for $p_2=0$ and the temperature of 20° centigrade must be nearly the same as the superficial tension of mercury in contact with air, or 55.03 grammes per linear metre according to Quincke (Pogg. Ann., Bd. 139, p. 27). The value of σ at the same temperature, when the condensed water begins to have the properties of water in mass, will be equal to the sum of the superficial tensions of mercury in contact with water and of water in contact with its own vapor. This will be, according to the same authority, 42.58 + 8.25, or 50.83 grammes per metre, if we neglect the difference of the tensions of water with its vapor and water with air. As p_2 , therefore, increases from zero to 236400 grammes per square metre (when water begins to be condensed *in mass*), σ diminishes from about 55.03 to about 50.83 grammes per linear metre. If the general course of the values of σ for intermediate values of p_2 were determined by experiment, we could easily form an approximate estimate of the values of the superficial density $\Gamma_{2(1)}$ for different pressures less than that of saturated vapor. It will be observed that the determination of the superficial density does not by any means depend upon inappreciable differences of superficial tension. The greatest difficulty in the determination would doubtless be that of distinguishing between the diminution of superficial tension due to the water and that due to other substances which might accidentally be present. Such determinations are of considerable practical importance on account of the use of mercury in measurements of the specific gravity of vapors.

The same physical relations may of course be deduced without giving up the use of the surface of tension as a dividing surface, but the formulæ which express them will be less simple. If we make t, μ_3, μ_4 , etc. constant, we have by (98) and (508)

$$\begin{aligned} dp' &= \gamma_1' d\mu_1 + \gamma_2' d\mu_2, \\ dp'' &= \gamma_1'' d\mu_1 + \gamma_2'' d\mu_2, \\ d\sigma &= -\Gamma_1 d\mu_1 - \Gamma_2 d\mu_2, \end{aligned}$$

where we may suppose Γ_1 and Γ_2 to be determined with reference to the surface of tension. Then, if $dp' = dp''$,

$$(\gamma_1' - \gamma_1'') d\mu_1 + (\gamma_2' - \gamma_2'') d\mu_2 = 0,$$

and

$$d\sigma = \Gamma_1 \frac{\gamma_2' - \gamma_2''}{\gamma_1' - \gamma_1''} d\mu_2 - \Gamma_2 d\mu_2.$$

That is,

$$\left(\frac{d\sigma}{d\mu_2}\right)_{p'-p'', t, \mu_3, \mu_4, \text{etc.}} = -\Gamma_2 + \Gamma_1 \frac{\gamma_2' - \gamma_2''}{\gamma_1' - \gamma_1''}. \quad (515)$$

The reader will observe that $\frac{\Gamma_1}{\gamma_1' - \gamma_1''}$ represents the distance between the surface of tension and that dividing surface which would make $\Gamma_1 = 0$; the second number of the last equation is therefore equivalent to $-\Gamma_{2(x)}$.

If any component substance has the same density in the two homogeneous masses separated by a plane surface of discontinuity, the value of the superficial density for that component is independent of the position of the dividing surface. In this case alone we may derive the value of the superficial density of a component with reference to the surface of tension from the fundamental equation for plane surfaces alone. Thus in the last equation, when $\gamma_2' = \gamma_2''$, the second member will reduce to $-\Gamma_2$. It will be observed that to make $p' - p'', t, \mu_3, \mu_4$, etc. constant is in this case equivalent to making t, μ_1, μ_3, μ_4 , etc. constant.

Substantially the same is true of the superficial density of entropy or of energy, when either of these has the same density in the two homogeneous masses.*

* With respect to questions which concern only the *form* of surfaces of discontinuity, such precision as we have employed in regard to the position of the dividing surface is evidently quite unnecessary. This precision has not been used for the sake of the mechanical part of the problem, which does not require the surface to be defined with greater nicety than we can employ in our observations, but in order to give

Concerning the Stability of Surfaces of Discontinuity.

We shall first consider the stability of a film separating homogeneous masses with respect to changes in its nature, while its position and the nature of the homogeneous masses are not altered. For this purpose, it will be convenient to suppose that the homogeneous masses are very large, and thoroughly stable with respect to the possible formation of any different homogeneous masses out of their components, and that the surface of discontinuity is plane and uniform.

Let us distinguish the quantities which relate to the actual components of one or both of the homogeneous masses by the suffixes a, b , etc., and those which relate to components which are found only at the surface of discontinuity by the suffixes g, h , etc., and consider the variation of the energy of the whole system in consequence of a given change in the nature of a small part of the surface of discontinuity, while the entropy of the whole system and the total quantities of the several components remain constant, as well as the volume of each of the homogeneous masses, as determined by the surface of tension. This small part of the surface of discontinuity in its changed state is supposed to be still uniform in nature, and such as may subsist in equilibrium between the given homogeneous masses, which will evidently not be sensibly altered in nature or thermodynamic state. The remainder of the surface of discontinuity is also supposed to remain uniform, and on account of its infinitely greater size to be infinitely less altered in its nature than the first part. Let $\Delta\varepsilon^s$ denote the increment of the superficial energy of this first part, $\Delta\eta^s, \Delta m_a^s, \Delta m_b^s$, etc., $\Delta m_g^s, \Delta m_h^s$, etc., the increments of its superficial

determinate values to the superficial densities of energy, entropy, and the component substances, which quantities, as has been seen, play an important part in the relations between the tension of a surface of discontinuity, and the composition of the masses which it separates.

The product σs of the superficial tension and the area of the surface, may be regarded as the *available energy* due to the surface in a system in which the temperature and the potentials μ_1, μ_2 , etc.—or the differences of these potentials and the gravitational potential (see page 208) when the system is subject to gravity—are maintained sensibly constant. The value of σ , as well as that of s , is sensibly independent of the precise position which we may assign to the dividing surface (so long as this is sensibly coincident with the surface of discontinuity), but ε_s , the *superficial density of energy*, as the term is used in this paper, like the superficial densities of entropy and of the component substances, requires a more precise localization of the dividing surface.

entropy and of the quantities of the components which we regard as belonging to the surface. The increments of entropy and of the various components which the rest of the system receive will be expressed by

$$-\Delta\eta^s, \quad -\Delta m_a^s, \quad -\Delta m_b^s, \quad \text{etc.}, \quad -\Delta m_g^s, \quad -\Delta m_h^s, \quad \text{etc.},$$

and the consequent increment of energy will be by (12) and (501)

$$-t\Delta\eta^s - \mu_a\Delta m_a^s - \mu_b\Delta m_b^s - \text{etc.} - \mu_g\Delta m_g^s - \mu_h\Delta m_h^s - \text{etc.}$$

Hence the total increment of energy in the whole system will be

$$\left. \begin{aligned} \Delta\varepsilon^s - t\Delta\eta^s - \mu_a\Delta m_a^s - \mu_b\Delta m_b^s - \text{etc.} \\ - \mu_g\Delta m_g^s - \mu_h\Delta m_h^s - \text{etc.} \end{aligned} \right\} \quad (516)$$

If the value of this expression is necessarily positive, for finite changes as well as infinitesimal in the nature of the part of the film to which $\Delta\varepsilon^s$, etc. relate,* the increment of energy of the whole system will be positive for any possible changes in the nature of the film, and the film will be stable, at least with respect to changes in its nature, as distinguished from its position. For, if we write

$$D\varepsilon^s, \quad D\eta^s, \quad Dm_a^s, \quad Dm_b^s, \quad \text{etc.}, \quad Dm_g^s, \quad Dm_h^s, \quad \text{etc.}$$

for the energy, etc. of any element of the surface of discontinuity, we have from the supposition just made

$$\begin{aligned} \Delta D\varepsilon^s - t\Delta D\eta^s - \mu_a\Delta Dm_a^s - \mu_b\Delta Dm_b^s - \text{etc.} \\ - \mu_g\Delta Dm_g^s - \mu_h\Delta Dm_h^s - \text{etc.} > 0; \end{aligned} \quad (517)$$

and integrating for the whole surface, since

$$\Delta f Dm_g^s = 0, \quad \Delta f Dm_h^s = 0, \quad \text{etc.},$$

we have

$$\Delta f D\varepsilon^s - t\Delta f D\eta^s - \mu_a\Delta f Dm_a^s - \mu_b\Delta f Dm_b^s - \text{etc.} > 0. \quad (518)$$

Now $\Delta f D\eta^s$ is the increment of the entropy of the whole surface, and $-\Delta f D\eta^s$ is therefore the increment of the entropy of the two homogeneous masses. In like manner, $-\Delta f Dm_a^s$, $-\Delta f Dm_b^s$, etc. are the increments of the quantities of the components in these masses. The expression

$$-t\Delta f D\eta^s - \mu_a\Delta f Dm_a^s - \mu_b\Delta f Dm_b^s - \text{etc.}$$

* In the case of infinitesimal changes in the nature of the film, the sign Δ must be interpreted, as elsewhere in this paper, without neglect of infinitesimals of the higher orders. Otherwise, by equation (501), the above expression would have the value zero.

denotes therefore, according to equation (12), the increment of energy of the two homogeneous masses, and since $\Delta f D\varepsilon^s$ denotes the increment of energy of the surface, the above condition expresses that the increment of the total energy of the system is positive. That we have only considered the possible formation of such films as are capable of existing in equilibrium between the given homogeneous masses can not invalidate the conclusion in regard to the stability of the film, for in considering whether any state of the system will have less energy than the given state, we need only consider the state of least energy, which is necessarily one of equilibrium.

If the expression (516) is capable of a negative value for an infinitesimal change in the nature of the part of the film to which the symbols relate, the film is obviously unstable.

If the expression is capable of a negative value, but only for finite and not for infinitesimal changes in the nature of this part of the film, the film is *practically unstable*,* i. e., if such a change were made in a small part of the film, the disturbance would tend to increase. But it might be necessary that the initial disturbance should also have a finite magnitude in respect to the extent of surface in which it occurs; for we cannot suppose that the thermodynamic relations of an infinitesimal part of a surface of discontinuity are independent of the adjacent parts. On the other hand, the changes which we have been considering are such that every part of the film remains in equilibrium with the homogeneous masses on each side; and if the energy of the system can be diminished by a finite change satisfying this condition, it may perhaps be capable of diminution by an infinitesimal change which does not satisfy the same condition. We must therefore leave it undetermined whether the film, which in this case is practically unstable, is or is not unstable in the strict mathematical sense of the term.

Let us consider more particularly the condition of practical stability, in which we need not distinguish between finite and infinitesimal changes. To determine whether the expression (516) is capable of a negative value, we need only consider the least value of which it is capable. Let us write it in the fuller form

$$\left. \begin{aligned} \varepsilon^{S''} - \varepsilon^{S'} - t(\eta^{S''} - \eta^{S'}) - \mu_a(m_a^{S''} - m_a^{S'}) - \mu_b(m_b^{S''} - m_b^{S'}) - \text{etc.} \\ - \mu'_g(m_g^{S''} - m_g^{S'}) - \mu'_h(m_h^{S''} - m_h^{S'}) - \text{etc.,} \end{aligned} \right\} (519)$$

where the single and double accents distinguish the quantities which

* With respect to the sense in which this term is used, compare page 133.

relate to the first and second states of the film, the letters without accents denoting those quantities which have the same value in both states. The differential of this expression when the quantities distinguished by double accents are alone considered variable, and the area of the surface is constant, will reduce by (501) to the form

$$(\mu_g'' - \mu_g') dm_g^{s''} + (\mu_h'' - \mu_h') dm_h^{s''} + \text{etc.}$$

To make this incapable of a negative value, we must have

$$\mu_g'' = \mu_g', \quad \text{unless } m_g^{s''} = 0,$$

$$\mu_h'' = \mu_h', \quad \text{unless } m_h^{s''} = 0.$$

In virtue of these relations and by equation (502), the expression (519), i. e., (516), will reduce to

$$\sigma'' s - \sigma' s,$$

which will be positive or negative according as

$$\sigma'' - \sigma' \tag{520}$$

is positive or negative.

That is, if the tension of the film is less than that of any other film which can exist between the same homogeneous masses (which has therefore the same values of t , μ_a , μ_b , etc.), and which moreover has the same values of the potentials μ_g , μ_h , etc., so far as it contains the substances to which these relate, then the first film will be stable. But the film will be practically unstable, if any other such film has a less tension. [Compare the expression (141), by which the practical stability of homogeneous masses is tested.]

It is, however, evidently necessary for the stability of the surface of discontinuity with respect to *deformation*, that the value of the superficial tension should be positive. Moreover, since we have by (502) for the surface of discontinuity

$$\varepsilon^s - t \eta^s - \mu_a m_a^s - \mu_b m_b^s - \text{etc.} - \mu_g m_g^s - \mu_h m_h^s - \text{etc.} = \sigma s,$$

and by (93) for the two homogeneous masses

$$\varepsilon' - t \eta' + p v' - \mu_a m_a' - \mu_b m_b' - \text{etc.} = 0,$$

$$\varepsilon'' - t \eta'' + p v'' - \mu_a m_a'' - \mu_b m_b'' - \text{etc.} = 0,$$

if we denote by

$$\varepsilon, \eta, v, m_a, m_b, \text{ etc.}, m_g, m_h, \text{ etc.},$$

the total energy, etc. of a composite mass consisting of two such homogeneous masses divided by such a surface of discontinuity, we shall have by addition of these equations

$$\varepsilon - t\eta + pv - \mu_a m_a - \mu_b m_b - \text{etc.} - \mu_g m_g - \mu_h m_h - \text{etc.} = \sigma s.$$

Now if the value of σ is negative, the value of the first member of this equation will decrease as s increases, and may therefore be decreased by making the mass to consist of thin alternate strata of the two kinds of homogeneous masses which we are considering. There will be no limit to the decrease which is thus possible with a given value of v , so long as the equation is applicable, i. e., so long as the strata have the properties of similar bodies in mass. But it may easily be shown (as in a similar case on pages 131, 132) that when the values of

$$t, p, \mu_a, \mu_b, \text{etc.}, \mu_g, \mu_h, \text{etc.}$$

are regarded as fixed, being determined by the surface of discontinuity in question, and the values of

$$\varepsilon, \eta, m_a, m_b, \text{etc.}, m_g, m_h, \text{etc.}$$

are variable and may be determined by any body having the given volume v , the first member of this equation cannot have an infinite negative value, and must therefore have a least possible value, which will be negative, if any value is negative, that is, if σ is negative.

The body determining ε, η , etc. which will give this least value to this expression will evidently be sensibly homogeneous. With respect to the formation of such a body, the system consisting of the two homogeneous masses and the surface of discontinuity with the negative tension is by (53) (see also page 133) at least practically unstable, if the surface of discontinuity is very large, so that it can afford the requisite material without sensible alteration of the values of the potentials. (This limitation disappears, if all the component substances are found in the homogeneous masses.) Therefore, in a system satisfying the conditions of practical stability with respect to the possible formation of all kinds of homogeneous masses, negative tensions of the surfaces of discontinuity are necessarily excluded.

Let us now consider the condition which we obtain by applying (516) to infinitesimal changes. The expression may be expanded as before to the form (519), and then reduced by equation (502) to the form

$$s(\sigma'' - \sigma') + m_g^{s''}(\mu_g'' - \mu_g') + m_h^{s''}(\mu_h'' - \mu_h') + \text{etc.}$$

That the value of this expression shall be positive when the quantities are determined by two films which differ infinitely little is a necessary condition of the stability of the film to which the single

accents relate. But if one film is stable, the other will in general be so too, and the distinction between the films with respect to stability is of importance only at the limits of stability. If all films for all values of μ_g, μ_h , etc. are stable, or all within certain limits, it is evident that the value of the expression must be positive when the quantities are determined by any two infinitesimally different films within the same limits. For such collective determinations of stability the condition may be written

$$-s \Delta\sigma - m_g^s \Delta\mu_g - m_h^s \Delta\mu_h - \text{etc.} > 0,$$

or

$$\Delta\sigma < -\Gamma_g \Delta\mu_g - \Gamma_h \Delta\mu_h - \text{etc.} \quad (521)$$

On comparison of this formula with (508), it appears that within the limits of stability the second and higher differential coefficients of the tension considered as a function of the potentials for the substances which are found only at the surface of discontinuity (the potentials for the substances found in the homogeneous masses and the temperature being regarded as constant) satisfy the conditions which would make the tension a maximum if the necessary conditions relative to the first differential coefficients were fulfilled.

In the foregoing discussion of stability, the surface of discontinuity is supposed plane. In this case, as the tension is supposed positive, there can be no tendency to a change of form of the surface. We now pass to the consideration of changes consisting in or connected with motion and change of form of the surface of tension, which we shall at first suppose to be and to remain spherical and uniform throughout.

In order that the equilibrium of a spherical mass entirely surrounded by an indefinitely large mass of different nature shall be neutral with respect to changes in the value of r , the radius of the sphere, it is evidently necessary that equation (500), which in this may be written

$$2\sigma = r(p' - p''), \quad (522)$$

as well as the other conditions of equilibrium, shall continue to hold true for varying values of r . Hence, for a state of equilibrium which is on the limit between stability and instability, it is necessary that the equation

$$2d\sigma = (p' - p'')dr + rdp'$$

shall be satisfied, when the relations between $d\sigma$, dp' , and dr are determined from the fundamental equations on the supposition that

the conditions of equilibrium relating to temperature and the potentials remain satisfied. (The differential coefficients in the equations which follow are to be determined on this supposition.) Moreover, if

$$r \frac{dp'}{dr} < 2 \frac{d\sigma}{dr} - p' + p'', \quad (523)$$

i. e., if the pressure of the interior mass increases less rapidly (or decreases more rapidly) with increasing radius than is necessary to preserve neutral equilibrium, the equilibrium is stable. But if

$$r \frac{dp'}{dr} > 2 \frac{d\sigma}{dr} - p' + p'', \quad (524)$$

the equilibrium is unstable. In the remaining case, when

$$r \frac{dp'}{dr} = 2 \frac{d\sigma}{dr} - p' + p'', \quad (525)$$

farther conditions are of course necessary to determine absolutely whether the equilibrium is stable or unstable, but in general the equilibrium will be stable in respect to change in one direction and unstable in respect to change in the opposite direction, and is therefore to be considered unstable. In general, therefore, we may call (523) the condition of stability.

When the interior mass and the surface of discontinuity are formed entirely of substances which are components of the external mass, p' and σ cannot vary and condition (524) being satisfied the equilibrium is unstable.

But if either the interior homogeneous mass or the surface of discontinuity contains substances which are not components of the enveloping mass, the equilibrium may be stable. If there is but one such substance, and we denote its densities and potential by γ'_1 , Γ_1 , and μ_1 , the condition of stability (523) will reduce to the form

$$\left(r \frac{dp'}{d\mu_1} - 2 \frac{d\sigma}{d\mu_1} \right) \frac{d\mu_1}{dr} < p'' - p',$$

or, by (98) and (508),

$$(r \gamma'_1 + 2 \Gamma_1) \frac{d\mu_1}{dr} < p'' - p'. \quad (526)$$

In these equations and in all which follow in the discussion of this case, the temperature and the potentials μ_2 , μ_3 , etc. are to be regarded as constant. But

$$\gamma'_1 v' + \Gamma_1 s,$$

which represents the total quantity of the component specified by the suffix, must be constant. It is evidently equal to

$$\frac{4}{3} \pi r^3 \gamma_1' + 4 \pi r^2 \Gamma_1.$$

Dividing by 4π and differentiating, we obtain

$$(r^2 \gamma_1' + 2 r \Gamma_1) dr + \frac{1}{3} r^3 d\gamma_1' + r^2 d\Gamma_1 = 0,$$

or, since γ_1' and Γ_1 are functions of μ_1 ,

$$(r \gamma_1' + 2 \Gamma_1) dr + \left(\frac{r^2}{3} \frac{d\gamma_1'}{d\mu_1} + r \frac{d\Gamma_1}{d\mu_1} \right) d\mu_1 = 0. \quad (527)$$

By means of this equation, the condition of stability is brought to the form

$$\frac{(r \gamma_1' + 2 \Gamma_1)^2}{\frac{r^2}{3} \frac{d\gamma_1'}{d\mu_1} + r \frac{d\Gamma_1}{d\mu_1}} > p' - p''. \quad (528)$$

If we eliminate r by equation (522), we have

$$\frac{\left(\frac{\gamma_1'}{p' - p''} + \frac{\Gamma_1}{\sigma} \right)^2}{\frac{1}{3(p' - p'')} \frac{d\gamma_1'}{d\mu_1} + \frac{1}{2\sigma} \frac{d\Gamma_1}{d\mu_1}} > 1. \quad (529)$$

If p' and σ are known in terms of t, μ_1, μ_2 , etc., we may express the first member of this condition in terms of the same variables and p'' . This will enable us to determine, for any given state of the external mass, the values of μ_1 which will make the equilibrium stable or unstable.

If the component to which γ_1' and Γ_1 relate is found only at the surface of discontinuity, the condition of stability reduces to

$$\frac{\Gamma_1^2}{\sigma} \frac{d\mu_1}{d\Gamma_1} > \frac{1}{2}. \quad (530)$$

Since

$$\Gamma_1 = - \frac{d\sigma}{d\mu_1},$$

we may also write

$$\frac{\Gamma_1}{\sigma} \frac{d\sigma}{d\Gamma_1} < - \frac{1}{2}, \quad \text{or} \quad \frac{d \log \sigma}{d \log \Gamma_1} < - \frac{1}{2}. \quad (531)$$

Again, if $\Gamma_1 = 0$ and $\frac{d\Gamma_1}{d\mu_1} = 0$, the condition of stability reduces to

$$\frac{3 \gamma_1'^2}{p' - p''} \frac{d\mu_1}{d\gamma_1'} > 1. \quad (532)$$

Since

$$\gamma_1' = \frac{dp'}{d\mu_1},$$

we may also write

$$\frac{\gamma_1'}{p' - p''} \frac{dp'}{d\gamma_1'} > \frac{1}{3}, \quad \text{or} \quad \frac{d \log (p' - p'')}{d \log \gamma_1'} > \frac{1}{3}. \quad (533)$$

When r is large, this will be a close approximation for any values of I'_1 , unless γ_1' is very small. The two special conditions (531) and (533) might be derived from very elementary considerations.

Similar conditions of stability may be found when there are more substances than one in the inner mass or the surface of discontinuity, which are not components of the enveloping mass. In this case, we have instead of (526) a condition of the form

$$(r \gamma_1' + 2 I'_1) \frac{d\mu_1}{dr} + (r \gamma_2' + 2 I'_2) \frac{d\mu_2}{dr} + \text{etc.} < p'' - p', \quad (534)$$

from which $\frac{d\mu_1}{dr}$, $\frac{d\mu_2}{dr}$, etc. may be eliminated by means of equations derived from the conditions that

$$\gamma_1' v' + I'_1 s, \quad \gamma_2' v' + I'_2 s, \quad \text{etc.}$$

must be constant.

Nearly the same method may be applied to the following problem. Two different homogeneous fluids are separated by a diaphragm having a circular orifice, their volumes being invariable except by the motion of the surface of discontinuity, which adheres to the edge of the orifice:—to determine the stability or instability of this surface when in equilibrium.

The condition of stability derived from (522) may in this case be written

$$r \frac{d(p' - p'')}{dv'} < 2 \frac{d\sigma}{dv'} - (p' - p'') \frac{dr}{dv'}, \quad (535)$$

where the quantities relating to the concave side of the surface of tension are distinguished by a single accent.

If both the masses are infinitely large, or if one which contains all the components of the system is infinitely large, $p' - p''$ and σ will be constant, and the condition reduces to

$$\frac{dr}{dv'} < 0.$$

The equilibrium will therefore be stable or unstable according as the surface of tension is less or greater than a hemisphere.

To return to the general problem:—if we denote by x the part of the axis of the circular orifice intercepted between the center of the orifice and the surface of tension, by R the radius of the orifice, and by V' the value of v' when the surface of tension is plane, we shall have the geometrical relations

$$R^2 = 2 r x - x^2,$$

and

$$\begin{aligned} v' &= V' + \frac{2}{3} \pi r^2 x - \frac{1}{3} \pi R^2 (r - x) \\ &= V' + \pi r x^2 - \frac{1}{3} \pi x^3. \end{aligned}$$

By differentiation we obtain

$$(r - x) dx + x dr = 0,$$

and $dv' = \pi x^2 dr + (2 \pi r x - \pi x^2) dv;$

whence $(r - x) dv' = - \pi r x^2 dr.$ (536)

By means of this relation, the condition of stability may be reduced to the form

$$\frac{dp'}{dr'} - \frac{dp''}{dr''} - \frac{2}{r} \frac{d\sigma}{dv'} < (p' - p'') \frac{r - x}{\pi r^2 x^2}. \quad (537)$$

Let us now suppose that the temperature and all the potentials except one, μ_1 , are to be regarded as constant. This will be the case when one of the homogeneous masses is very large and contains all the components of the system except one, or when both these masses are very large and there is a single substance at the surface of discontinuity which is not a component of either; also when the whole system contains but a single component, and is exposed to a constant temperature at its surface. Condition (537) will reduce by (98) and (508) to the form

$$\left(\gamma_1' - \gamma_1'' + \frac{2 \Gamma_1}{r} \right) \frac{d\mu_1}{dv'} < (p' - p'') \frac{r - x}{\pi r^2 x^2}. \quad (538)$$

But $\gamma_1' v' + \gamma_1'' v'' + \Gamma_1 s$

(the total quantity of the component specified by the suffix) must be constant; therefore, since

$$dv'' = - dv', \quad \text{and} \quad ds = \frac{2}{r} dv',$$

$$\left(v' \frac{d\gamma_1'}{d\mu_1} + v'' \frac{d\gamma_1''}{d\mu_1} + s \frac{d\Gamma_1}{d\mu_1} \right) d\mu_1 + \left(\gamma_1' - \gamma_1'' + \frac{2 \Gamma_1}{r} \right) dv' = 0. \quad (539)$$

By this equation, the condition of stability is brought to the form

$$\frac{\left(\gamma_1' - \gamma_1'' + \frac{2 \Gamma_1}{r} \right)^2}{v' \frac{d\gamma_1'}{d\mu_1} + v'' \frac{d\gamma_1''}{d\mu_1} + s \frac{d\Gamma_1}{d\mu_1}} > (p' - p'') \frac{x - r}{\pi x^2 r^2}. \quad (540)$$

When the substance specified by the suffix is a component of either of the homogeneous masses, the terms $\frac{2 \Gamma_1}{r}$ and $s \frac{d\Gamma_1}{d\mu_1}$ may generally be neglected. When it is not a component of either, the terms γ_1' , γ_1'' , $v' \frac{d\gamma_1'}{d\mu_1}$, $v'' \frac{d\gamma_1''}{d\mu_1}$ may of course be cancelled, but we must not apply the formula to cases in which the substance spreads over the diaphragm separating the homogeneous masses.

In the cases just discussed, the problem of the stability of certain surfaces of tension has been solved by considering the case of neutral equilibrium,—a condition of neutral equilibrium affording the equation of the limit of stability. This method probably leads as directly as any to the result, when that consists in the determination of the value of a certain quantity at the limit of stability, or of the relation which exists at that limit between certain quantities specifying the state of the system. But problems of a more general character may require a more general treatment.

Let it be required to ascertain the stability or instability of a fluid system in a given state of equilibrium with respect to motion of the surfaces of tension and accompanying changes. It is supposed that the conditions of internal stability for the separate homogeneous masses are satisfied, as well as those conditions of stability for the surfaces of discontinuity which relate to small portions of these surfaces with the adjacent masses. (The conditions of stability which are here supposed to be satisfied have been already discussed in part and will be farther discussed hereafter.) The fundamental equations for all the masses and surfaces occurring in the system are supposed to be known. In applying the general criteria of stability which are given on page 110, we encounter the following difficulty.

The question of the stability of the system is to be determined by the consideration of states of the system which are slightly varied from that of which the stability is in question. These varied states of the system are not in general states of equilibrium, and the relations expressed by the fundamental equations may not hold true of them. More than this,—if we attempt to describe a varied state of the system by varied values of the quantities which describe the initial state, if these varied values are such as are inconsistent with equilibrium, they may fail to determine with precision any state of the system. Thus, when the phases of two contiguous homogeneous masses are specified, if these phases are such as satisfy all the conditions of equilibrium, the nature of the surface of discontinuity (if without additional components) is entirely determined; but if the phases do not satisfy all the conditions of equilibrium, the nature of the surface of discontinuity is not only undetermined, but incapable of determination by specified values of such quantities as we have employed to express the nature of surfaces of discontinuity in equilibrium. For example, if the temperatures in contiguous homogeneous masses are different, we cannot specify the thermal state of the surface of discontinuity by assigning to it any particular temperature. It would be

necessary to give the law by which the temperature passes over from one value to the other. And if this were given, we could make no use of it in the determination of other quantities, unless the rate of change of the temperature were so gradual, that at every point we could regard the thermodynamic state as unaffected by the change of temperature in its vicinity. It is true that we are also ignorant in respect to surfaces of discontinuity *in equilibrium* of the law of change of those quantities which are different in the two phases in contact, such as the densities of the components, but this, although unknown to us, is entirely determined by the nature of the phases in contact, so that no vagueness is occasioned in the definition of any of the quantities which we have occasion to use with reference to such surfaces of discontinuity.

It may be observed that we have established certain differential equations, especially (497), in which only the initial state is necessarily one of equilibrium. Such equations may be regarded as establishing certain properties of states bordering upon those of equilibrium. But these are properties which hold true only when we disregard quantities proportional to the square of those which express the degree of variation of the system from equilibrium. Such equations are therefore sufficient for the determination of the conditions of equilibrium, but not sufficient for the determination of the conditions of stability

We may, however, use the following method to decide the question of stability in such a case as has been described.

Beside the real system of which the stability is in question, it will be convenient to conceive of another system, to which we shall attribute in its initial state the same homogeneous masses and surfaces of discontinuity which belong to the real system. We shall also suppose that the homogeneous masses and surfaces of discontinuity of this system, which we may call the imaginary system, have the same fundamental equations as those of the real system. But the imaginary system is to differ from the real in that the variations of its state are limited to such as do not violate the conditions of equilibrium relating to temperature and the potentials, and that the fundamental equations of the surfaces of discontinuity hold true for these varied states, although the condition of equilibrium expressed by equation (500) may not be satisfied.

Before proceeding farther, we must decide whether we are to examine the question of stability under the condition of a constant external temperature, or under the condition of no transmission of

heat to or from external bodies, and in general, to what external influences we are to regard the system as subject. It will be convenient to suppose that the exterior of the system is fixed, and that neither matter nor heat can be transmitted through it. Other cases may easily be reduced to this, or treated in a manner entirely analogous.

Now if the real system in the given state is unstable, there must be some slightly varied state in which the energy is less, but the entropy and the quantities of the components the same as in the given state, and the exterior of the system unvaried. But it may easily be shown that the given state of the system may be made stable by constraining the surfaces of discontinuity to pass through certain fixed lines situated in the unvaried surfaces. Hence, if the surfaces of discontinuity are constrained to pass through corresponding fixed lines in the surfaces of discontinuity belonging to the varied state just mentioned, there must be a state of stable equilibrium for the system thus constrained which will differ infinitely little from the given state of the system, the stability of which is in question, and will have the same entropy, quantities of components and exterior, but less energy. The imaginary system will have a similar state, since the real and imaginary systems do not differ in respect to those states which satisfy all the conditions of equilibrium for each surface of discontinuity. That is, the imaginary system has a state, differing infinitely little from the given state, and with the same entropy, quantities of components, and exterior, but with less energy.

Conversely, if the imaginary system has such a state as that just described, the real system will also have such a state. This may be shown by fixing certain lines in the surfaces of discontinuity of the imaginary system in its state of less energy and then making the energy a minimum under the conditions. The state thus determined will satisfy all the conditions of equilibrium for each surface of discontinuity, and the real system will therefore have a corresponding state, in which the entropy, quantities of components, and exterior will be the same as in the given state, but the energy less.

We may therefore determine whether the given system is or is not unstable, by applying the general criterion of instability (7) to the imaginary system.

If the system is not unstable, the equilibrium is either neutral or stable. Of course we can determine which of these is the case by reference to the imaginary system, since this determination depends upon states of equilibrium, in regard to which the real and imaginary

systems do not differ. We may therefore determine whether the equilibrium of the given system is stable, neutral, or unstable, by applying the criteria (3)–(7) to the imaginary system.

The result which we have obtained may be expressed as follows:— In applying to a fluid system which is in equilibrium, and of which all the small parts taken separately are stable, the criteria of stable, neutral, and unstable equilibrium, we may regard the system as under constraint to satisfy the conditions of equilibrium relating to temperature and the potentials, and as satisfying the relations expressed by the fundamental equations for masses and surfaces, even when the condition of equilibrium relating to pressure [equation (500)] is not satisfied.

It follows immediately from this principle, in connection with equations (501) and (86), that in a stable system each surface of tension must be a surface of minimum area for constant values of the volumes which it divides, when the other surfaces bounding these volumes and the perimeter of the surface of tension are regarded as fixed; that in a system in neutral equilibrium each surface of tension will have as small an area as it can receive by any slight variations under the same limitations; and that in seeking the remaining conditions of stable or neutral equilibrium, when these are satisfied, it is only necessary to consider such varied surfaces of tension as have similar properties with reference to the varied volumes and perimeters.

We may illustrate the method which has been described by applying it to a problem but slightly different from one already (pp. 408, 409) discussed by a different method. It is required to determine the conditions of stability for a system in equilibrium, consisting of two different homogeneous masses meeting at a surface of discontinuity, the perimeter of which is invariable, as well as the exterior of the whole system, which is also impermeable to heat.

To determine what is necessary for stability in addition to the condition of minimum area for the surface of tension, we need only consider those varied surfaces of tension which satisfy the same condition. We may therefore regard the surface of tension as determined by v' , the volume of one of the homogeneous masses. But the state of the system would evidently be completely determined by the position of the surface of tension and the temperature and potentials, if the entropy and the quantities of the components were variable; and therefore, since the entropy and the quantities of the components are constant, the state of the system must be completely determined by the position of the surface of tension. We may therefore regard

all the quantities relating to the system as functions of v' , and the condition of stability may be written

$$\frac{d\varepsilon}{dv'} dv' + \frac{1}{2} \frac{d^2\varepsilon}{dv'^2} dv'^2 + \text{etc.} > 0,$$

where ε denotes the total energy of the system. Now the conditions of equilibrium require that

$$\frac{d\varepsilon}{dv'} = 0,$$

Hence, the general condition of stability is that

$$\frac{d^2\varepsilon}{dv'^2} > 0. \quad (541)$$

Now if we write ε' , ε'' , ε^s for the energies of the two masses and of the surface, we have by (86) and (501), since the total entropy and the total quantities of the several components are constant,

$$d\varepsilon = d\varepsilon' + d\varepsilon'' + d\varepsilon^s = -p' dv' - p'' dv'' + \sigma ds,$$

or, since $dv'' = -dv'$,

$$\frac{d\varepsilon}{dv'} = -p' + p'' + \sigma \frac{ds}{dv'}. \quad (542)$$

Hence,

$$\frac{d^2\varepsilon}{dv'^2} = -\frac{dp'}{dv'} + \frac{dp''}{dv'} + \frac{d\sigma}{dv'} \frac{ds}{dv'} + \sigma \frac{d^2s}{dv'^2}, \quad (543)$$

and the condition of stability may be written

$$\sigma \frac{d^2s}{dv'^2} > \frac{dp'}{dv'} - \frac{dp''}{dv'} - \frac{d\sigma}{dv'} \frac{ds}{dv'}. \quad (544)$$

If we now simplify the problem by supposing, as in the similar case on page 409, that we may disregard the variations of the temperature and of all the potentials except one, the condition will reduce to

$$\sigma \frac{d^2s}{dv'^2} > \left(\gamma_1' - \gamma_1'' + \Gamma_1 \frac{ds}{dv'} \right) \frac{d\mu_1}{dv'}. \quad (545)$$

The total quantity of the substance indicated by the suffix 1 is

$$\gamma_1' v' + \gamma_1'' v'' + \Gamma_1 s.$$

Making this constant, we have

$$\left(\gamma_1' - \gamma_1'' + \Gamma_1 \frac{ds}{dv'} \right) dv' + \left(v' \frac{d\gamma_1'}{d\mu_1} + v'' \frac{d\gamma_1''}{d\mu_1} + s \frac{d\Gamma_1}{d\mu_1} \right) d\mu_1 = 0. \quad (546)$$

The condition of equilibrium is thus reduced to the form

$$\sigma \frac{d^2s}{dv'^2} > - \frac{\left(\gamma_1' - \gamma_1'' + \Gamma_1 \frac{ds}{dv'} \right)^2}{v' \frac{d\gamma_1'}{d\mu_1} + v'' \frac{d\gamma_1''}{d\mu_1} + s \frac{d\Gamma_1}{d\mu_1}}, \quad (547)$$

where $\frac{ds}{dv'}$ and $\frac{d^2s}{dv'^2}$ are to be determined from the form of the surface of tension by purely geometrical considerations, and the other differential coefficients are to be determined from the fundamental equations of the homogeneous masses and the surface of discontinuity. Condition (540) may be easily deduced from this as a particular case.

The condition of stability with reference to motion of surfaces of discontinuity admits of a very simple expression when we can treat the temperature and potentials as constant. This will be the case when one or more of the homogeneous masses, containing together all the component substances, may be considered as indefinitely large, the surfaces of discontinuity being finite. For if we write $\sum \Delta \varepsilon$ for the sum of the variations of the energies of the several homogeneous masses, and $\sum \Delta \varepsilon^s$ for the sum of the variations of the energies of the several surfaces of discontinuity, the condition of stability may be written

$$\sum \Delta \varepsilon + \sum \Delta \varepsilon^s > 0, \tag{548}$$

the total entropy and the total quantities of the several components being constant. The variations to be considered are infinitesimal, but the character Δ signifies, as elsewhere in this paper, that the expression is to be interpreted without neglect of infinitesimals of the higher orders. Since the temperature and potentials are sensibly constant, the same will be true of the pressures and surface-tensions, and by integration of (86) and (501) we may obtain for any homogeneous mass

$$\Delta \varepsilon = t \Delta \eta - p \Delta v + \mu_1 \Delta m_1 + \mu_2 \Delta m_2 + \text{etc.},$$

and for any surface of discontinuity

$$\Delta \varepsilon^s = t \Delta \eta^s + \sigma \Delta s + \mu_1 \Delta m_1^s + \mu_2 \Delta m_2^s + \text{etc.}$$

These equations will hold true of finite differences, when $t, p, \sigma, \mu_1, \mu_2$, etc. are constant, and will therefore hold true of infinitesimal differences, under the same limitations, without neglect of the infinitesimals of the higher orders. By substitution of these values, the condition of stability will reduce to the form

$$- \sum (p \Delta v) + \sum (\sigma \Delta s) > 0,$$

or
$$\sum (p \Delta v) - \sum (\sigma \Delta s) < 0. \tag{549}$$

That is, the sum of the products of the volumes of the masses by their pressures diminished by the sum of the products of the areas of the surfaces of discontinuity by their tensions must be a maximum. This is a purely geometrical condition, since the pressures and ten-

sions are constant. This condition is of interest, because it is always sufficient for stability with reference to motion of surfaces of discontinuity. For any system may be reduced to the kind described by putting certain parts of the system in communication (by means of fine tubes if necessary) with large masses of the proper temperatures and potentials. This may be done without introducing any new movable surfaces of discontinuity. The condition (549) when applied to the altered system is therefore the same as when applied to the original system. But it is sufficient for the stability of the altered system, and therefore sufficient for its stability if we diminish its freedom by breaking the connection between the original system and the additional parts, and therefore sufficient for the stability of the original system.

On the Possibility of the Formation of a Fluid of different Phase within any Homogeneous Fluid.

The study of surfaces of discontinuity throws considerable light upon the subject of the stability of such homogeneous fluid masses as have a less pressure than others formed of the same components (or some of them) and having the same temperature and the same potentials for their actual components.*

In considering this subject, we must first of all inquire how far our method of treating surfaces of discontinuity is applicable to cases in which the radii of curvature of the surfaces are of insensible magnitude. That it should not be applied to such cases without limitation is evident from the consideration that we have neglected the term $\frac{1}{2}(C_1 - C_2)\delta(c_1 - c_2)$ in equation (494) on account of the magnitude of the radii of curvature compared with the thickness of the non-homogeneous film. (See page 390). When, however, only spherical masses are considered, this term will always disappear, since C_1 and C_2 will necessarily be equal.

Again, the surfaces of discontinuity have been regarded as separating homogeneous masses. But we may easily conceive that a globular mass (surrounded by a large homogeneous mass of different nature) may be so small that no part of it will be homogeneous, and that even at its center the matter cannot be regarded as having any phase of matter *in mass*. This, however, will cause no difficulty, if we regard the phase of the interior mass as determined by the same

* See page 161, where the term stable is used (as indicated on page 159) in a less strict sense than in the discussion which here follows.

relations to the exterior mass as in other cases. Beside the phase of the exterior mass, there will always be another phase having the same temperature and potentials, but of the general nature of the small globule which is surrounded by that mass and in equilibrium with it. This phase is completely determined by the system considered, and in general entirely stable and perfectly capable of realization in mass, although not such that the exterior mass could exist in contact with it at a plane surface. This is the phase which we are to attribute to the mass which we conceive as existing within the dividing surface.*

With this understanding with regard to the phase of the fictitious interior mass, there will be no ambiguity in the meaning of any of the symbols which we have employed, when applied to cases in which the surface of discontinuity is spherical, however small the radius may be. Nor will the demonstration of the general theorems require any material modification. The dividing surface, which determines the value of ε^s , η^s , \bar{m}_1^s , m_2^s , etc., is as in other cases to be placed so as to make the term $\frac{1}{2}(C_1 + C_2)\delta(c_1 + c_2)$ in equation (494) vanish, i. e., so as to make equation (497) valid. It has been shown on pages 387–389 that when thus placed it will sensibly coincide with the physical surface of discontinuity, when this consists of a non-homogeneous film separating homogeneous masses, and having radii of curvature which are large compared with its thickness. But in regard to globular masses too small for this theorem to have any application, it will be worth while to examine how far we may be certain that the radius of the dividing surface will have a real and positive value, since it is only then that our method will have any natural application.

The value of the radius of the dividing surface, supposed spherical, of any globule in equilibrium with a surrounding homogeneous fluid may be most easily obtained by eliminating σ from equations (500) and (502), which have been derived from (497), and contain the radius implicitly. If we write r for this radius, equation (500) may be written

$$2\sigma = (p' - p'')r, \quad (550)$$

the single and double accents referring respectively to the interior and exterior masses. If we write $[\varepsilon]$, $[\eta]$, $[m_1]$, $[m_2]$, etc. for the

* For example, in applying our formulæ to a microscopic globule of water in steam, by the density or pressure of the interior mass we should understand, not the actual density or pressure at the center of the globule, but the density of liquid water (in large quantities) which has the temperature and potential of the steam.

excess of the total energy, entropy, etc. in and about the globular mass above what would be in the same space if it were uniformly filled with matter of the phase of the exterior mass, we shall have necessarily with reference to the whole dividing surface

$$\varepsilon^S = [\varepsilon] - v'(\varepsilon_v' - \varepsilon_v''), \quad \eta^S = [\eta] - v'(\eta_v' - \eta_v''),$$

$$m_1^S = [m_1] - v'(\gamma_1' - \gamma_1''), \quad m_2^S = [m_2] - v'(\gamma_2' - \gamma_2''), \text{ etc.,}$$

where ε_v' , ε_v'' , η_v' , η_v'' , γ_1' , γ_1'' , etc. denote, in accordance with our usage elsewhere, the volume-densities of energy, of entropy, and of the various components, in the two homogeneous masses. We may thus obtain from equation (502)

$$\sigma s = [\varepsilon] - v'(\varepsilon_v' - \varepsilon_v'') - t[\eta] + t v'(\eta_v' - \eta_v'')$$

$$- \mu_1[m_1] + \mu_1 v'(\gamma_1' - \gamma_1'') - \mu_2[m_2] + \mu_2 v'(\gamma_2' - \gamma_2'') - \text{etc.} \quad (551)$$

But by (93),

$$p' = -\varepsilon_v' + t\eta_v' + \mu_1\gamma_1' + \mu_2\gamma_2' + \text{etc.},$$

$$p'' = -\varepsilon_v'' + t\eta_v'' + \mu_1\gamma_1'' + \mu_2\gamma_2'' + \text{etc.}$$

Let us also write for brevity

$$W = [\varepsilon] - t[\eta] - \mu_1[m_1] - \mu_2[m_2] - \text{etc.} \quad (552)$$

(It will be observed that the value of W is entirely determined by the nature of the physical system considered, and that the notion of the dividing surface does not in any way enter into its definition.)

We shall then have

$$\sigma s = W + v'(p' - p''), \quad (553)$$

or, substituting for s and v' their values in terms of r ,

$$4\pi r^2 \sigma = W + \frac{4}{3}\pi r^3(p' - p''), \quad (554)$$

and eliminating σ by (550),

$$\frac{2}{3}\pi r^3(p' - p'') = W, \quad (555)$$

$$r = \left(\frac{3W}{2\pi(p' - p'')} \right)^{\frac{1}{3}}. \quad (556)$$

If we eliminate r instead σ , we have

$$\frac{16\pi\sigma^3}{3(p' - p'')^2} = W, \quad (557)$$

$$\sigma = \left(\frac{3W(p' - p'')^2}{16\pi} \right)^{\frac{1}{3}}. \quad (558)$$

Now, if we first suppose the difference of the pressures in the homogeneous masses to be very small, so that the surface of discontinuity is nearly plane, since without any important loss of generality

we may regard σ as positive (for if σ is not positive when $p' = p''$, the surface when plane would not be stable in regard to position, as it certainly is, in every actual case, when the proper conditions are fulfilled with respect to its perimeter), we see by (550) that the pressure in the interior mass must be the greater; i. e., we may regard σ , $p' - p''$, and r as all positive. By (555), the value of W will also be positive. But it is evident from equation (552), which defines W , that the value of this quantity is necessarily real, in any possible case of equilibrium, and can only become infinite when r becomes infinite and $p' = p''$. Hence, by (556) and (558), as $p' - p''$ increases from very small values, W , r , and σ have single, real, and positive values until they simultaneously reach the value zero. Within this limit, our method is evidently applicable; beyond this limit, if such exist, it will hardly be profitable to seek to interpret the equations. But it must be remembered that the vanishing of the radius of the somewhat arbitrarily determined *dividing surface* may not necessarily involve the vanishing of the physical heterogeneity. It is evident, however, (see pp. 387-389,) that the globule must become insensible in magnitude before r can vanish.

It may easily be shown that the quantity denoted by W is the work which would be required to form (by a reversible process) the heterogeneous globule in the interior of a very large mass having initially the uniform phase of the exterior mass. For this work is equal to the increment of energy of the system when the globule is formed without change of the entropy or volume of the whole system or of the quantities of the several components. Now $[\eta]$, $[m_1]$, $[m_2]$, etc. denote the increments of entropy and of the components in the space where the globule is formed. Hence these quantities with the negative sign will be equal to the increments of entropy and of the components in the rest of the system. And hence, by equation (86),

$$-t[\eta] - \mu_1[m_1] - \mu_2[m_2] - \text{etc.}$$

will denote the increment of energy in all the system except where the globule is formed. But $[\varepsilon]$ denotes the increment of energy in that part of the system. Therefore, by (552), W denotes the total increment of energy in the circumstances supposed, or the work required for the formation of the globule.

The conclusions which may be drawn from these considerations with respect to the stability of the homogeneous mass of the pressure p'' (supposed less than p' , the pressure belonging to a different phase of the same temperature and potentials) are very obvious.

Within those limits within which the method used has been justified, the mass in question must be regarded as in strictness stable with respect to the growth of a globule of the kind considered, since W , the work required for the formation of such a globule of a certain size (viz., that which would be in equilibrium with the surrounding mass), will always be positive. Nor can smaller globules be formed, for they can neither be in equilibrium with the surrounding mass, being too small, nor grow to the size of that to which W relates. If, however, by any external agency such a globular mass (of the size necessary for equilibrium) were formed, the equilibrium has already (page 406) been shown to be unstable, and with the least excess in size, the interior mass would tend to increase without limit except that depending on the magnitude of the exterior mass. We may therefore regard the quantity W as affording a kind of measure of the *stability* of the phase to which p'' relates. In equation (557) the value of W is given in terms of σ and $p' - p''$. If the three fundamental equations which give σ , p' , and p'' in terms of the temperature and the potentials were known, we might regard the stability (W) as known in terms of the same variables. It will be observed that when $p' = p''$ the value of W is infinite. If $p' - p''$ increases without greater changes of the phases than are necessary for such increase, W will vary at first very nearly inversely as the square of $p' - p''$. If $p' - p''$ continues to increase, it may perhaps occur that W reaches the value zero; but until this occurs the phase is certainly stable with respect to the kind of change considered. Another kind of change is conceivable, which initially is small in degree but may be great in its extent in space. Stability in this respect or *stability in respect to continuous changes of phase* has already been discussed (see page 162), and its limits determined. These limits depend entirely upon the fundamental equation of the homogeneous mass of which the stability is in question. But with respect to the kind of changes here considered, which are initially small in extent but great in degree, it does not appear how we can fix the limits of stability with the same precision. But it is safe to say that if there is such a limit it must be at or beyond the limit at which σ vanishes. This latter limit is determined entirely by the fundamental equation of the surface of discontinuity between the phase of which the stability is in question and that of which the possible formation is in question. We have already seen that when σ vanishes, the radius of the dividing surface and the work W vanish with it. If the fault in the homogeneity of the mass vanishes at the same time, (it evidently

cannot vanish sooner,) the phase becomes unstable at this limit. But if the fault in the homogeneity of the physical mass does not vanish with r , σ and W ,—and no sufficient reason appears why this should not be considered as the general case,—although the amount of work necessary to upset the equilibrium of the phase is infinitesimal, this is not enough to make the phase unstable. It appears therefore that W is a somewhat one-sided measure of stability.

It must be remembered in this connection that the fundamental equation of a surface of discontinuity can hardly be regarded as capable of experimental determination, except for plane surfaces, (see pp. 394, 395,) although the relation for spherical surfaces is in the nature of things entirely determined, at least so far as the phases are separately capable of existence. Yet the foregoing discussion yields the following practical results. It has been shown that the real stability of a phase extends in general beyond that limit (discussed on pages 160, 161), which may be called the limit of practical stability, at which the phase can exist in contact with another at a plane surface, and a formula has been deduced to express the degree of stability in such cases as measured by the amount of work necessary to upset the equilibrium of the phase when supposed to extend indefinitely in space. It has also been shown to be entirely consistent with the principles established that this stability should have limits, and the manner in which the general equations would accommodate themselves to this case has been pointed out.

By equation (553), which may be written

$$W = \sigma s - (p' - p'') v', \quad (559)$$

we see that the work W consists of two parts, of which one is always positive, and is expressed by the product of the superficial tension and the area of the surface of tension, and the other is always negative, and is numerically equal to the product of the difference of pressure by the volume of the interior mass. We may regard the first part as expressing the work spent in forming the surface of tension, and the second part the work gained in forming the interior mass.*

* To make the physical significance of the above more clear, we may suppose the two processes to be performed separately in the following manner. We may suppose a large mass of the same phase as that which has the volume v' to exist initially in the interior of the other. Of course, it must be surrounded by a resisting envelop, on account of the difference of the pressures. We may, however, suppose this envelop permeable to all the component substances, although not of such properties that a mass can form on the exterior like that within. We may allow the

Moreover, the second of these quantities, if we neglect its sign, is always equal to two-thirds of the first, as appears from equation (550) and the geometrical relation $v' = \frac{2}{3}rs$. We may therefore write

$$W = \frac{1}{3} \sigma s = \frac{1}{2} (p' - p'') v'. \quad (560)$$

On the Possible Formation at the Surface where two different Homogeneous Fluids meet of a Fluid of different Phase from either.

Let A, B, and C be three different fluid phases of matter, which satisfy all the conditions necessary for equilibrium when they meet at plane surfaces. The components of A and B may be the same or different, but C must have no components except such as belong to A or B. Let us suppose masses of the phases A and B to be separated by a very thin sheet of the phase C. This sheet will not necessarily be plane, but the sum of its principal curvatures must be zero. We may treat such a system as consisting simply of masses of the phases A and B with a certain surface of discontinuity, for in our previous discussion there has been nothing to limit the thickness or the nature of the film separating homogeneous masses, except that its thickness has generally been supposed to be small in comparison with its radii of curvature. The value of the superficial tension for such a film will be $\sigma_{AC} + \sigma_{BC}$, if we denote by these symbols the tensions of the surfaces of contact of the phases A and C, and B and C, respectively. This not only appears from evident mechanical considerations, but may also be easily verified by equations (502) and (93), the first of which may be regarded as defining the quantity σ . This value will not be affected by diminishing the thickness of the film, until the

envelop to yield to the internal pressure until its contents are increased by v' without materially affecting its superficial area. If this be done sufficiently slowly, the phase of the mass within will remain constant. (See page 139.) A homogeneous mass of the volume v' and of the desired phase has thus been produced, and the work gained is evidently $(p' - p'')v'$.

Let us suppose that a small aperture is now opened and closed in the envelop so as to let out exactly the volume v' of the mass within, the envelop being pressed inwards in another place so as to diminish its contents by this amount. During the extrusion of the drop and until the orifice is entirely closed, the surface of the drop must adhere to the edge of the orifice, but not elsewhere to the outside surface of the envelop. The work done in forming the surface of the drop will evidently be σs or $\frac{2}{3}(p' - p'')v'$. Of this work, the amount $(p' - p'')v'$ will be expended in pressing the envelop inward, and the rest in opening and closing the orifice. Both the opening and the closing will be resisted by the capillary tension. If the orifice is circular, it must have, when widest open, the radius determined by equation (550).

limit is reached at which the interior of the film ceases to have the properties of matter in mass. Now if $\sigma_{AC} + \sigma_{BC}$ is greater than σ_{AB} , the tension of the ordinary surface between A and B, such a film will be at least practically unstable. (See page 403.) We cannot suppose that $\sigma_{AB} > \sigma_{AC} + \sigma_{BC}$, for this would make the ordinary surface between A and B unstable and difficult to realize. If $\sigma_{AB} = \sigma_{AC} + \sigma_{BC}$, we may assume, in general, that this relation is not accidental, and that the ordinary surface of contact for A and B is of the kind which we have described.

Let us now suppose the phases A and B to vary, so as still to satisfy the conditions of equilibrium at plane contact, but so that the pressure of the phase C determined by the temperature and potentials of A and B shall become less than the pressure of A and B. A system consisting of the phases A and B will be entirely stable with respect to the formation of any phase like C. (The case is not quite identical with that considered on page 161, since the system in question contains two different phases, but the principles involved are entirely the same.)

With respect to variations of the phases A and B in the opposite direction we must consider two cases separately. It will be convenient to denote the pressures of the three phases by p_A, p_B, p_C , and to regard these quantities as functions of the temperature and potentials.

If $\sigma_{AB} = \sigma_{AC} + \sigma_{BC}$ for values of the temperature and potentials which make $p_A = p_B = p_C$, it will not be possible to alter the temperature and potentials at the surface of contact of the phases A and B so that $p_A = p_B$, and $p_C > p_A$, for the relation of the temperature and potentials necessary for the equality of the three pressures will be preserved by the increase of the mass of the phase C. Such variations of the phases A and B might be brought about in separate masses, but if these were brought into contact, there would be an immediate formation of a mass of the phase C, with reduction of the phases of the adjacent masses to such as satisfy the conditions of equilibrium with that phase.

But if $\sigma_{AB} < \sigma_{AC} + \sigma_{BC}$, we can vary the temperature and potentials so that $p_A = p_B$, and $p_C > p_A$, and it will not be possible for a sheet of the phase of C to form *immediately*, i. e., while the pressure of C is sensibly equal to that of A and B; for mechanical work equal to $\sigma_{AC} + \sigma_{BC} - \sigma_{AB}$ per unit of surface might be obtained by bringing the system into its original condition, and therefore produced without any external expenditure, unless it be that of heat at the temperature of the system, which is evidently incapable of producing the work.

The stability of the system in respect to such a change must therefore extend beyond the point where the pressure of C commences to be less than that of A and B. We arrive at the same result if we use the expression (520) as a test of stability. Since this expression has a finite positive value when the pressures of the phases are all equal, the ordinary surface of discontinuity must be stable, and it must require a finite change in the circumstances of the case to make it become unstable.*

In the preceding paragraph it is shown that the surface of contact of phases A and B is stable under certain circumstances, with respect to the formation of a thin sheet of the phase C. To complete the demonstration of the stability of the surface with respect to the formation of the phase C, it is necessary to show that this phase cannot be formed at the surface in lentiform masses. This is the more necessary, since it is in this manner, if at all, that the phase is likely to be formed, for an incipient sheet of phase C would evidently be unstable when $\sigma_{AB} < \sigma_{AC} + \sigma_{BC}$, and would immediately break up into lentiform masses.

It will be convenient to consider first a lentiform mass of phase C




FIG. 10.

in equilibrium between masses of phases A and B which meet in a plane surface. Let figure 10 represent a section of such a system through the centers of the spherical surfaces, the mass of phase A lying on the left of DEH'FG, and that of phase B on the right of DEH''FG. Let the line joining the centers cut the spherical surfaces in H' and H'', and the plane of the surface of contact of A and B in I. Let the radii of EH'F and EH''F be denoted by r' , r'' , and the segments I I', I I'' by x' , x'' . Also let IE, the radius of the circle in which the spherical surfaces intersect, be denoted by R . By a suitable application of the general condition of equilibrium we may easily obtain the equation

$$\sigma_{AC} \frac{r' - x'}{r'} + \sigma_{BC} \frac{r'' - x''}{r''} = \sigma_{AB}, \quad (561)$$

* It is true that such a case as we are now considering is formally excluded in the discussion referred to, which relates to a plane surface, and in which the system is supposed thoroughly stable with respect to the possible formation of any different homogeneous masses. Yet the reader will easily convince himself that the criterion (520) is perfectly valid in this case with respect to the possible formation of a thin sheet of the phase C, which, as we have seen, may be treated simply as a different kind of surface of discontinuity.

which signifies that the components parallel to EF of the tension σ_{AC} and σ_{BC} are together equal to σ_{AB} . If we denote by W the amount of work which must be expended in order to form such a lentiform mass as we are considering between masses of indefinite extent having the phases A and B, we may write

$$W = M - N, \tag{562}$$

where M denotes the work expended in replacing the surface between A and B by the surfaces between A and C and B and C, and N denotes the work gained in replacing the masses of phases A and B by the mass of phase C. Then

$$M = \sigma_{AC} s_{AC} + \sigma_{BC} s_{BC} - \sigma_{AB} s_{AB}, \tag{563}$$

where s_{AC} , s_{BC} , s_{AB} denote the areas of the three surfaces concerned; and

$$N = V' (p_C - p_A) + V'' (p_C - p_B), \tag{564}$$

where V' and V'' denote the volumes of the masses of the phases A and B which are replaced. Now by (500),

$$p_C - p_A = \frac{2\sigma_{AC}}{r'}, \quad \text{and} \quad p_C - p_B = \frac{2\sigma_{BC}}{r''}. \tag{565}$$

We have also the geometrical relations

$$\left. \begin{aligned} V' &= \frac{2}{3} \pi r'^2 x' - \frac{1}{3} \pi R^2 (r' - x'), \\ V'' &= \frac{2}{3} \pi r''^2 x'' - \frac{1}{3} \pi R^2 (r'' - x''). \end{aligned} \right\} \tag{566}$$

By substitution we obtain

$$\begin{aligned} N &= \frac{4}{3} \pi \sigma_{AC} r' x' - \frac{2}{3} \pi R^2 \sigma_{AC} \frac{r' - x'}{r'} \\ &\quad + \frac{4}{3} \pi \sigma_{BC} r'' x'' - \frac{2}{3} \pi R^2 \sigma_{BC} \frac{r'' - x''}{r''}, \end{aligned} \tag{567}$$

and by (561),

$$N = \frac{4}{3} \pi \sigma_{AC} r' x' + \frac{4}{3} \pi \sigma_{BC} r'' x'' - \frac{2}{3} \pi R^2 \sigma_{AB}. \tag{568}$$

Since

$$2 \pi r' x' = s_{AC}, \quad 2 \pi r'' x'' = s_{BC}, \quad \pi R^2 = s_{AB},$$

we may write

$$N = \frac{2}{3} (\sigma_{AC} s_{AC} + \sigma_{BC} s_{BC} - \sigma_{AB} s_{AB}). \tag{569}$$

(The reader will observe that the ratio of M and N is the same as that of the corresponding quantities in the case of the spherical mass treated on pages 416-422.) We have therefore

$$W = \frac{1}{3} (\sigma_{AC} s_{AC} + \sigma_{BC} s_{BC} - \sigma_{AB} s_{AB}). \tag{570}$$

This value is positive so long as

$$\sigma_{AC} + \sigma_{BC} > \sigma_{AB},$$

since

$$s_{AC} > s_{AB}, \quad \text{and} \quad s_{BC} > s_{AB}.$$

But at the limit, when

$$\sigma_{AC} + \sigma_{BC} = \sigma_{AB},$$

we see by (561) that

$$s_{AC} = s_{AB}, \quad \text{and} \quad s_{BC} = s_{AB},$$

and therefore

$$W = 0.$$

It should however be observed that in the immediate vicinity of the circle in which the three surfaces of discontinuity intersect, the physical state of each of these surfaces must be affected by the vicinity of the others. We cannot, therefore, rely upon the formula (570) except when the dimensions of the lentiform mass are of sensible magnitude.

We may conclude that after we pass the limit at which p_C becomes greater than p_A and p_B (supposed equal) lentiform masses of phase C will not be formed until either $\sigma_{AB} = \sigma_{AC} + \sigma_{BC}$, or $p_C - p_A$ becomes so great that the lentiform mass which would be in equilibrium is one of insensible magnitude. [The diminution of the radii with increasing values of $p_C - p_A$ is indicated by equation (565).] Hence, no mass of phase C will be formed until one of these limits is reached. Although the demonstration relates to a *plane* surface between A and B, the result must be applicable whenever the radii of curvature have a sensible magnitude, since the effect of such curvature may be disregarded when the lentiform mass is of sufficiently small.

The equilibrium of the lentiform mass of phase C is easily proved to be unstable, so that the quantity W affords a kind of measure of the stability of plane surfaces of contact of the phases A and B.*

* If we represent phases by the position of points in such a manner that coexistent phases (in the sense in which the term is used on page 152) are represented by the same point, and allow ourselves, for brevity, to speak of the phases as having the positions of the points by which they are represented, we may say that three coexistent phases are situated where three series of pairs of coexistent phases meet or intersect. If the three phases are all fluid, or when the effects of solidity may be disregarded, two cases are to be distinguished. Either the three series of coexistent phases all intersect,—this is when each of the three surface-tensions is less than the sum of the two others,—or one of the series terminates where the two others intersect,—this is where one surface tension is equal to the sum of the others. The series of coexistent phases will be represented by lines or surfaces, according as the phases have one or two independently variable components. Similar relations exist when the number of components is greater, except that they are not capable of geometrical representation without some limitation, as that of constant temperature or pressure or certain constant potentials.

Essentially the same principles apply to the more general problem in which the phases A and B have moderately different pressures, so that their surfaces of contact must be curved, but the radii of curvature have a sensible magnitude.

In order that a thin film of the phase C may be in equilibrium between masses of the phases A and B, the following equations must be satisfied—

$$\sigma_{AC}(c_1 + c_2) = p_A - p_C,$$

$$\sigma_{BC}(c_1 + c_2) = p_C - p_B,$$

where c_1 and c_2 denote the principal curvatures of the film, the centers of positive curvature lying in the mass having the phase A. Eliminating $c_1 + c_2$, we have

$$\sigma_{BC}(p_A - p_C) = \sigma_{AC}(p_C - p_B),$$

or

$$p_C = \frac{\sigma_{BC} p_A + \sigma_{AC} p_B}{\sigma_{BC} + \sigma_{AC}}. \tag{571}$$

It is evident that if p_C has a value greater than that determined by this equation, such a film will develop into a larger mass; if p_C has a less value, such a film will tend to diminish. Hence, when

$$p_C < \frac{\sigma_{BC} p_A + \sigma_{AC} p_B}{\sigma_{BC} + \sigma_{AC}}, \tag{572}$$

the phases A and B have a stable surface of contact.

Again, if more than one kind of surface of discontinuity is possible between A and B, for any given values of the temperature and potentials, it will be impossible for that having the greater tension to displace the other, at the temperature and with the potentials considered. Hence, when p_C has the value determined by equation (571), and consequently $\sigma_{AC} + \sigma_{BC}$ is one value of the tension for the surface between A and B, it is impossible that the ordinary tension of the surface σ_{AB} should be greater than this. If $\sigma_{AB} = \sigma_{AC} + \sigma_{BC}$, when equation (571) is satisfied, we may presume that a thin film of the phase C actually exists at the surface between A and B, and that a variation of the phases such as would make p_C greater than the second number of (571) cannot be brought about at that surface, as it would be prevented by the formation of a larger mass of the phase C. But if $\sigma_{AB} < \sigma_{AC} + \sigma_{BC}$ when equation (571) is satisfied, this equation does not mark the limit of the stability of the surface between A and B, for the temperature or potentials must receive a finite

change before the film of phase C, or (as we shall see in the following paragraph) a lentiform mass of that phase, can be formed.

The work which must be expended in order to form on the surface between indefinitely large masses of phases A and B a lentiform mass of phase C in equilibrium, may evidently be represented by the formula

$$W = \sigma_{AC} S_{AC} + \sigma_{BC} S_{BC} - \sigma_{AB} S_{AB} \\ - p_C V_C + p_A V_A + p_B V_B, \quad (573)$$

where S_{AC} , S_{BC} denote the areas of the surfaces formed between A and C, and B and C, S_{AB} the diminution of the area of the surface between A and B, V_C the volume formed of the phase C, and V_A , V_B the diminution of the volumes of the phases A and B. Let us now suppose σ_{AC} , σ_{BC} , σ_{AB} , p_A , p_B to remain constant and the external boundary of the surface between A and B to remain fixed, while p_C increases and the surfaces of tension receive such alterations as are necessary for equilibrium. It is not necessary that this should be physically possible in the actual system; we may suppose the changes to take place, for the sake of argument, although involving changes in the fundamental equations of the masses and surfaces considered. Then, regarding W simply as an abbreviation for the second member of the preceding equation, we have

$$dW = \sigma_{AC} dS_{AC} + \sigma_{BC} dS_{BC} - \sigma_{AB} dS_{AB} \\ - p_C dV_C + p_A dV_A + p_B dV_B - V_C dp_C. \quad (574)$$

But the conditions of equilibrium require that

$$\sigma_{AC} dS_{AC} + \sigma_{BC} dS_{BC} - \sigma_{AB} dS_{AB} \\ - p_C dV_C + p_A dV_A + p_B dV_B = 0. \quad (575)$$

Hence,

$$dW = -V_C dp_C. \quad (576)$$

Now it is evident that V_C will diminish as p_C increases. Let us integrate the last equation supposing p_C to increase from its original value until V_C vanishes. This will give

$$W'' - W' = \text{a negative quantity}, \quad (577)$$

where W' and W'' denote the initial and final values of W . But $W'' = 0$. Hence W' is positive. But this is the value of W in the original system containing the lentiform mass, and expresses the work necessary to form the mass between the phases A and B. It is therefore impossible that such a mass should form on a surface be-

tween these phases. We must however observe the same limitation as in the less general case already discussed,—that $p_C - p_A$, $p_C - p_B$ must not be so great that the dimensions of the lentiform mass are of insensible magnitude. It may also be observed that the value of these differences may be so small that there will not be room on the surface between the masses of phases A and B for a mass of phase C sufficiently large for equilibrium. In this case we may consider a mass of phase C which is in equilibrium upon the surface between A and B in virtue of a *constraint* applied to the line in which the three surfaces of discontinuity intersect, which will not allow this line to become longer, although not preventing it from becoming shorter. We may prove that the value of W is positive by such an integration as we have used before.

Substitution of Pressures for Potentials in Fundamental Equations for Surfaces.

The fundamental equation of a surface which gives the value of the tension in terms of the temperature and potentials seems best adapted to the purposes of theoretical discussion, especially when the number of components is large or undetermined. But the experimental determination of the fundamental equations, or the application of any result indicated by theory to actual cases, will be facilitated by the use of other quantities in place of the potentials, which shall be capable of more direct measurement, and of which the numerical expression (when the necessary measurements have been made) shall depend upon less complex considerations. The numerical value of a potential depends not only upon the system of units employed, but also upon the arbitrary constants involved in the definition of the energy and entropy of the substance to which the potential relates, or, it may be, of the elementary substances of which that substance is formed. (See page 152.) This fact and the want of means of direct measurement may give a certain vagueness to the idea of the potentials, and render the equations which involve them less fitted to give a clear idea of physical relations.

Now the fundamental equation of each of the homogeneous masses which are separated by any surface of discontinuity affords a relation between the pressure in that mass and the temperature and potentials. We are therefore able to eliminate one or two potentials from the fundamental equation of a surface by introducing the pressures in the adjacent masses. Again, when one of these masses is a gas-

mixture which satisfies Dalton's law as given on page 215, the potential for each simple gas may be expressed in terms of the temperature and the partial pressure belonging to that gas. By the introduction of these partial pressures we may eliminate as many potentials from the fundamental equation of the surface as there are simple gases in the gas-mixture.

An equation obtained by such substitutions may be regarded as a fundamental equation for the surface of discontinuity to which it relates, for when the fundamental equations of the adjacent masses are known, the equation in question is evidently equivalent to an equation between the tension, temperature, and potentials, and we must regard the knowledge of the properties of the adjacent masses as an indispensable preliminary, or an essential part, of a complete knowledge of any surface of discontinuity. It is evident, however, that from these fundamental equations involving pressures instead of potentials we cannot obtain by differentiation (without the use of the fundamental equations of the homogeneous masses) precisely the same relations as by the differentiation of the equations between the tensions, temperatures, and potentials. It will be interesting to inquire, at least in the more important cases, what relations may be obtained by differentiation from the fundamental equations just described alone.

If there is but one component, the fundamental equations of the two homogeneous masses afford one relation more than is necessary for the elimination of the potential. It may be convenient to regard the tension as a function of the temperature and the difference of the pressures. Now we have by (508) and (98)

$$d\sigma = -\eta_s dt - \Gamma d\mu_1,$$

$$d(p' - p'') = (\eta_{v'} - \eta_{v''}) dt + (\gamma' - \gamma'') d\mu_1.$$

Hence we derive the equation

$$d\sigma = -\left(\eta_s - \frac{\Gamma}{\gamma' - \gamma''} (\eta_{v'} - \eta_{v''})\right) dt - \frac{\Gamma}{\gamma' - \gamma''} d(p' - p''), \quad (578)$$

which indicates the differential coefficients of σ with respect to t and $p' - p''$. For surfaces which may be regarded as nearly plane, it is evident that $\frac{\Gamma}{\gamma' - \gamma''}$ represents the distance from the surface of tension to a dividing surface located so as to make the superficial density of the single component vanish, (being positive, when the

latter surface is on the side specified by the double accents,) and that the coefficient of dt (without the negative sign) represents the superficial density of entropy as determined by the latter dividing surface, i. e., the quantity denoted by $\eta_{s(1)}$ on page 397.

When there are two components, neither of which is confined to the surface of discontinuity, we may regard the tension as a function of the temperature and the pressures in the two homogeneous masses. The values of the differential coefficients of the tension with respect to these variables may be represented in a simple form if we choose such substances for the components that in the particular state considered each mass shall consist of a single component. This will always be possible when the composition of the two masses is not identical, and will evidently not affect the values of the differential coefficients. We then have

$$\begin{aligned} d\sigma &= -\eta_s dt - \Gamma'_i d\mu_i - \Gamma''_u d\mu_u, \\ dp' &= \eta_{v'} dt + \gamma' d\mu_i, \\ dp'' &= \eta_{v''} dt + \gamma'' d\mu_u, \end{aligned}$$

where the marks $'$ and $''$ are used instead of the usual $_1$ and $_2$ to indicate the identity of the component specified with the substance of the homogeneous masses specified by $'$ and $''$. Eliminating $d\mu_i$ and $d\mu_u$ we obtain

$$d\sigma = -\left(\eta_s - \frac{\Gamma'_i}{\gamma'} \eta_{v'} - \frac{\Gamma''_u}{\gamma''} \eta_{v''}\right) dt - \frac{\Gamma'_i}{\gamma'} dp' - \frac{\Gamma''_u}{\gamma''} dp''. \quad (579)$$

We may generally neglect the difference of p' and p'' , and write

$$d\sigma = -\left(\eta_s - \frac{\Gamma'_i}{\gamma'} \eta_{v'} - \frac{\Gamma''_u}{\gamma''} \eta_{v''}\right) dt - \left(\frac{\Gamma'_i}{\gamma'} + \frac{\Gamma''_u}{\gamma''}\right) dp. \quad (580)$$

The equation thus modified is strictly to be regarded as the equation for a plane surface. It is evident that $\frac{\Gamma'_i}{\gamma'}$ and $\frac{\Gamma''_u}{\gamma''}$ represent the distances from the surface of tension of the two surfaces of which one would make Γ'_i vanish, and the other Γ''_u , that $\frac{\Gamma'_i}{\gamma'} + \frac{\Gamma''_u}{\gamma''}$ represents the distance between these two surfaces, or the *diminution of volume* due to a unit of the surface of discontinuity, and that the coefficient of dt (without the negative sign) represents the excess of entropy in a system consisting of a unit of the surface of discontinuity with a part of each of the adjacent masses above that which the same matter would have if it existed in two homogeneous masses of the same phases but without any surface of discontinuity.

(A mass thus existing without any surface of discontinuity must of course be entirely surrounded by matter of the same phase.)*

The form in which the values of $\left(\frac{d\sigma}{dt}\right)_p$ and $\left(\frac{d\sigma}{dp}\right)_t$ are given in equation (580) is adapted to give a clear idea of the relations of these quantities to the particular state of the system for which they are to be determined, but not to show how they vary with the state of the system. For this purpose it will be convenient to have the values of these differential coefficients expressed with reference to ordinary components. Let these be specified as usual by $_1$ and $_2$. If we eliminate $d\mu_1$ and $d\mu_2$ from the equations

$$\begin{aligned} -d\sigma &= \eta_s dt + I'_1 d\mu_1 + I'_2 d\mu_2, \\ dp &= \eta_v' dt + \gamma_1' d\mu_1 + \gamma_2' d\mu_2, \\ dp &= \eta_v'' dt + \gamma_1'' d\mu_1 + \gamma_2'' d\mu_2, \end{aligned}$$

* If we set

$$V = -\frac{\Gamma'}{\gamma'} - \frac{\Gamma''}{\gamma''}, \quad (a)$$

$$H_s = \eta_s - \frac{\Gamma'}{\gamma'} \eta_v' - \frac{\Gamma''}{\gamma''} \eta_v'', \quad (b)$$

and in like manner

$$E_s = \varepsilon_s - \frac{\Gamma'}{\gamma'} \varepsilon_v' - \frac{\Gamma''}{\gamma''} \varepsilon_v''. \quad (c)$$

we may easily obtain, by means of equations (93) and (507),

$$E_s = t H_s + \sigma - p V. \quad (d)$$

Now equation (580) may be written

$$d\sigma = -H_s dt + V dp. \quad (e)$$

Differentiating (d), and comparing the result with (e), we obtain

$$dE_s = t dH_s - p dV. \quad (f)$$

The quantities E_s and H_s might be called the superficial densities of energy and entropy quite as properly as those which we denote by ε_s and η_s . In fact, when the composition of both of the homogeneous masses is invariable, the quantities E_s and H_s are much more simple in their definition than ε_s and η_s , and would probably be more naturally suggested by the terms *superficial density of energy* and *of entropy*. It would also be natural in this case to regard the quantities of the homogeneous masses as determined by the total quantities of matter, and not by the surface of tension or any other dividing surface. But such a nomenclature and method could not readily be extended so as to treat cases of more than two components with entire generality.

In the treatment of surfaces of discontinuity in this paper, the definitions and nomenclature which have been adopted will be strictly adhered to. The object of this note is to suggest to the reader how a different method might be used in some cases with advantage, and to show the precise relations between the quantities which are used in this paper and others which might be confounded with them, and which may be made more prominent when the subject is treated differently.

we obtain

$$d\sigma = \frac{B}{A} dt + \frac{C}{A} dp, \tag{581}$$

where

$$A = \gamma_1'' \gamma_2' - \gamma_1' \gamma_2'', \tag{582}$$

$$B = \begin{vmatrix} \eta_s & \Gamma_1 & \Gamma_2 \\ \eta_v' & \gamma_1' & \gamma_2' \\ \eta_v'' & \gamma_1'' & \gamma_2'' \end{vmatrix}, \tag{583}$$

$$C = \Gamma_1 (\gamma_2'' - \gamma_2') + \Gamma_2 (\gamma_1' - \gamma_1''). \tag{584}$$

It will be observed that A vanishes when the composition of the two homogeneous masses is identical, while B and C do not, in general, and that the value of A is negative or positive according as the mass specified by $'$ contains the component specified by $_1$ in a greater or less proportion than the other mass. Hence, the values both of $\left(\frac{d\sigma}{dt}\right)_p$ and of $\left(\frac{d\sigma}{dp}\right)_t$ become infinite when the difference in the composition of the masses vanishes, and change sign when the greater proportion of a component passes from one mass to the other. This might be inferred from the statements on page 155 respecting coexistent phases which are identical in composition, from which it appears that when two coexistent phases have nearly the same composition, a small variation of the temperature or pressure of the coexistent phases will cause a relatively very great variation in the composition of the phases. The same relations are indicated by the graphical method represented in figure 6 on page 184.

With regard to gas-mixtures which conform to Dalton's law, we shall only consider the fundamental equation for plane surfaces, and shall suppose that there is not more than one component in the liquid which does not appear in the gas-mixture. We have already seen that in limiting the fundamental equation to plane surfaces we can get rid of one potential by choosing such a dividing surface that the superficial density of one of the components vanishes. Let this be done with respect to the component peculiar to the liquid, if such there is; if there is no such component, let it be done with respect to one of the gaseous components. Let the remaining potentials be eliminated by means of the fundamental equations of the simple gases. We may thus obtain an equation between the superficial tension, the temperature, and the several pressures of the simple gases in the gas-mixture or all but one of these pressures. Now, if we eliminate $d\mu_2, d\mu_3,$ etc. from the equations

$$\begin{aligned}d\sigma &= -\eta_{s(1)} dt - \Gamma_{2(1)} d\mu_2 + \Gamma_{3(1)} d\mu_3 + \text{etc.}, \\dp_2 &= \eta_{v2} dt + \gamma_2 d\mu_2, \\dp_3 &= \eta_{v3} dt + \gamma_3 d\mu_3, \\&\text{etc.},\end{aligned}$$

where the suffix ₁ relates to the component of which the surface-density has been made to vanish, and γ_2, γ_3 , etc. denote the densities of the gases specified in the gas mixture, and p_2, p_3 , etc., η_{v2}, η_{v3} , etc. the pressures and the densities of entropy due to these several gases, we obtain

$$\begin{aligned}d\sigma &= -\left(\eta_{s(1)} - \frac{\Gamma_{2(1)}}{\gamma_2} \eta_{v2} - \frac{\Gamma_{3(1)}}{\gamma_3} \eta_{v3} - \text{etc.}\right) dt \\&\quad - \frac{\Gamma_{2(1)}}{\gamma_2} dp_2 - \frac{\Gamma_{3(1)}}{\gamma_3} dp_3 - \text{etc.} \quad (585)\end{aligned}$$

This equation affords values of the differential coefficients of σ with respect to t, p_2, p_3 , etc., which may be set equal to those obtained by differentiating the equation between these variables.

Thermal and Mechanical Relations pertaining to the Extension of a Surface of Discontinuity.

The fundamental equation of a surface of discontinuity with one or two component substances, beside its statical applications, is of use to determine the heat absorbed when the surface is extended under certain conditions.

Let us first consider the case in which there is only a single component substance. We may treat the surface as plane, and place the dividing surface so that the surface density of the single component vanishes. (See page 397.) If we suppose the area of the surface to be increased by unity without change of temperature or of the quantities of liquid and vapor, the entropy of the whole will be increased by $\eta_{s(1)}$. Therefore, if we denote by Q the quantity of heat which must be added to satisfy the conditions, we shall have

$$Q = t \eta_{s(1)}, \quad (586)$$

and by (514),

$$Q = -t \frac{d\sigma}{dt} = -\frac{d\sigma}{d \log t}, \quad (587)$$

It will be observed that the condition of constant quantities of liquid and vapor as determined by the dividing surface which we have adopted is equivalent to the condition that the total volume shall remain constant.

Again, if the surface is extended without application of heat, while the pressure in the liquid and vapor remains constant, the temperature will evidently be maintained constant by condensation of the vapor. If we denote by M the mass of vapor condensed per unit of surface formed, and by η_M' and η_M'' the entropies of the liquid and vapor per unit of mass, the condition of no addition of heat will require that

$$M (\eta_M'' - \eta_M') = \eta_{s(1)}. \tag{588}$$

The increase of the volume of liquid will be

$$\frac{\eta_{s(1)}}{\gamma' (\eta_M'' - \eta_M')}, \tag{589}$$

and the diminution of the volume of vapor

$$\frac{\eta_{s(1)}}{\gamma'' (\eta_M'' - \eta_M')}. \tag{590}$$

Hence, for the work done (per unit of surface formed) by the external bodies which maintain the pressure, we shall have

$$W = \frac{p \eta_{s(1)}}{\eta_M'' - \eta_M'} \left(\frac{1}{\gamma''} - \frac{1}{\gamma'} \right). \tag{591}$$

and, by (514) and (131),

$$W = -p \frac{d\sigma}{dt} \frac{dt}{dp} = -p \frac{d\sigma}{dp} = -\frac{d\sigma}{d \log p}. \tag{592}$$

The work expended directly in extending the film will of course be equal to σ .

Let us now consider the case in which there are two component substances, neither of which is confined to the surface. Since we cannot make the superficial density of both these substances vanish by any dividing surface, it will be best to regard the surface of tension as the dividing surface. We may, however, simplify the formula by choosing such substances for components that each homogeneous mass shall consist of a single component. Quantities relating to these components will be distinguished as on page 431. If the surface is extended until its area is increased by unity, while heat is added at the surface so as to keep the temperature constant, and the pressure of the homogeneous masses is also kept constant, the phase of these masses will necessarily remain unchanged, but the quantity of one will be diminished by Γ_a , and that of the other by Γ_b . Their entropies will therefore be diminished by $\frac{\Gamma_a}{\gamma'} \eta_V'$ and $\frac{\Gamma_b}{\gamma''} \eta_V''$, respect-

ively. Hence, since the surface receives the increment of entropy η_s , the total quantity of entropy will be increased by

$$\eta_s - \frac{\Gamma'_i}{\gamma'} \eta_{V'} - \frac{\Gamma''_i}{\gamma''} \eta_{V''},$$

which by equation (580) is equal to

$$- \left(\frac{d\sigma}{dt} \right)_p.$$

Therefore, for the quantity of heat Q imparted to the surface, we shall have

$$Q = -t \left(\frac{d\sigma}{dt} \right)_p = - \left(\frac{d\sigma}{d \log t} \right)_p. \quad (593)$$

We must notice the difference between this formula and (587). In (593) the quantity of heat Q is determined by the condition that the temperature and pressures shall remain constant. In (587) these conditions are equivalent and insufficient to determine the quantity of heat. The additional condition by which Q is determined may be most simply expressed by saying that the total volume must remain constant. Again, the differential coefficient in (593) is defined by considering p as constant; in the differential coefficient in (587) p cannot be considered as constant, and no condition is necessary to give the expression a definite value. Yet, notwithstanding the difference of the two cases, it is quite possible to give a single demonstration which shall be applicable to both. This may be done by considering a cycle of operations after the method employed by Sir William Thomson, who first pointed out these relations.*

The diminution of volume (per unit of surface formed) will be

$$V = \frac{\Gamma'_i}{\gamma'} + \frac{\Gamma''_i}{\gamma''} = - \left(\frac{d\sigma}{dp} \right)_t; \quad (594)$$

and the work done (per unit of surface formed) by the external bodies which maintain the pressure constant will be

$$W = -p \left(\frac{d\sigma}{dp} \right)_t = - \left(\frac{d\sigma}{d \log p} \right)_t. \quad (595)$$

Compare equation (592).

The values of Q and W may also be expressed in terms of quantities relating to the ordinary components. By substitution in (593) and (595) of the values of the differential coefficients which are given by (581), we obtain

* See *Proc. Roy. Soc.*, vol. ix, p. 255, (June, 1858); or *Phil. Mag.*, Ser. 4, vol. xvii, p. 61.

$$Q = -t \frac{B}{A}, \quad W = -p \frac{C}{A}. \quad (596)$$

where A , B , and C represent the expressions indicated by (582)–(584). It will be observed that the values of Q and W are in general infinite for the surface of discontinuity between coexistent phases which differ infinitesimally in composition, and change sign with the quantity A . When the phases are absolutely identical in composition, it is not in general possible to counteract the effect of extension of the surface of discontinuity by any supply of heat. For the matter at the surface will not in general have the same composition as the homogeneous masses, and the matter required for the increased surface cannot be obtained from these masses without altering their phase. The infinite values of Q and W are explained by the fact that when the phases are nearly identical in composition, the extension of the surface of discontinuity is accompanied by the vaporization or condensation of a very large mass, according as the liquid or the vapor is the richer in that component which is necessary for the formation of the surface of discontinuity.

If, instead of considering the amount of heat necessary to keep the phases from altering while the surface of discontinuity is extended, we consider the variation of temperature caused by the extension of the surface while the pressures remain constant, it appears that this variation of temperature changes sign with $\gamma_1''\gamma_2' - \gamma_1'\gamma_2''$, but vanishes with this quantity, i. e., vanishes when the composition of the phases becomes the same. This may be inferred from the statements on page 155, or from a consideration of the figure on page 184. When the composition of the homogeneous masses is initially absolutely identical, the effect on the temperature of a finite extension or contraction of the surface of discontinuity will be the same,—either of the two will lower or raise the temperature according as the temperature is a maximum or minimum for constant pressure.

The effect of the extension of a surface of discontinuity which is most easily verified by experiment is the effect upon the tension before complete equilibrium has been reestablished throughout the adjacent masses. A fresh surface between coexistent phases may be regarded in this connection as an extreme case of a recently extended surface. When sufficient time has elapsed after the extension of a surface originally in equilibrium between coexistent phases, the superficial tension will evidently have sensibly its original value, unless there are substances at the surface which are either not found

at all in the adjacent masses, or are found only in quantities comparable to those in which they exist at the surface. But a surface newly formed or extended may have a very different tension.

This will not be the case, however, when there is only a single component substance, since all the processes necessary for equilibrium are confined to a film of insensible thickness, and will require no appreciable time for their completion.

When there are two components, neither of which is confined to the surface of discontinuity, the reestablishment of equilibrium after the extension of the surface does not necessitate any processes reaching into the interior of the masses except the transmission of heat between the surface of discontinuity and the interior of the masses. It appears from equation (593) that if the tension of the surface diminishes with a rise of temperature, heat must be supplied to the surface to maintain the temperature uniform when the surface is extended, i. e., the effect of extending the surface is to cool it; but if the tension of any surface increases with the temperature, the effect of extending the surface will be to raise its temperature. In either case, it will be observed, the immediate effect of extending the surface is to increase its tension. A contraction of the surface will of course have the opposite effect. But the time necessary for the reestablishment of sensible thermal equilibrium after extension or contraction of the surface must in most cases be very short.

In regard to the formation or extension of a surface between two coexistent phases of more than two components, there are two extreme cases which it is desirable to notice. When the superficial density of each of the components is exceeding small compared with its density in either of the homogeneous masses, the matter (as well as the heat) necessary for the formation or extension of the normal surface can be taken from the immediate vicinity of the surface without sensibly changing the properties of the masses from which it is taken. But if any one of these superficial densities has a considerable value, while the density of the same component is very small in each of the homogeneous masses, both absolutely and relatively to the densities of the other components, the matter necessary for the formation or extension of the normal surface must come from a considerable distance. Especially if we consider that a small difference of density of such a component in one of the homogeneous masses will probably make a considerable difference in the value of the corresponding potential [see eq. (217)], and that a small difference in the value of the potential will make a considerable difference in the ten-

sion [see eq. (508)], it will be evident that in this case a considerable time will be necessary after the formation of a fresh surface or the extension of an old one for the reëstablishment of the normal value of the superficial tension. In intermediate cases, the reëstablishment of the normal tension will take place with different degrees of rapidity.

But whatever the number of component substances, provided that it is greater than one, and whether the reëstablishment of equilibrium is slow or rapid, extension of the surface will generally produce increase and contraction decrease of the tension. It would evidently be inconsistent with stability that the opposite effects should be produced. In general, therefore, a fresh surface between coexistent phases has a greater tension than an old one.* By the use of fresh surfaces, in experiments in capillarity, we may sometimes avoid the effect of minute quantities of foreign substances, which may be present without our knowledge or desire, in the fluids which meet at the surface investigated.

When the establishment of equilibrium is rapid, the variation of the tension from its normal value will be manifested especially during the extension or contraction of the surface, the phenomenon resembling that of viscosity, except that the variations of tension arising from variations in the densities at and about the surface will be the same in all directions, while the variations of tension due to any property of the surface really analogous to viscosity would be greatest in the direction of the most rapid extension.

We may here notice the different action of traces in the homogeneous masses of those substances which increase the tension and of those which diminish it. When the volume-densities of a component are very small, its surface-density may have a considerable positive value, but can only have a very minute negative one.† For the value when negative cannot exceed (numerically) the product of the greater volume-density by the thickness of the non-homogeneous

* When, however, homogeneous masses which have not coexistent phases are brought into contact, the superficial tension may increase with the course of time. The superficial tension of a drop of alcohol and water placed in a large room will increase as the potential for alcohol is equalized throughout the room, and is diminished in the vicinity of the surface of discontinuity.

† It is here supposed that we have chosen for components such substances as are incapable of resolution into other components which are independently variable in the homogeneous masses. In a mixture of alcohol and water, for example, the components must be pure alcohol and pure water.

film. Each of these quantities is exceedingly small. The surface-density when positive is of the same order of magnitude as the thickness of the non-homogeneous film, but is not necessarily small compared with other surface-densities because the volume-densities of the same substance in the adjacent masses are small. Now the potential of a substance which forms a very small part of a homogeneous mass certainly increases, and probably very rapidly, as the proportion of that component is increased. [See (171) and (217).] The pressure, temperature, and the other potentials, will not be sensibly affected. [See (98).] But the effect on the tension of this increase of the potential will be proportional to the surface-density, and will be to diminish the tension when the surface-density is positive. [See (508).] It is therefore quite possible that a very small trace of a substance in the homogeneous masses should greatly diminish the tension, but not possible that such a trace should greatly increase it.*

Impermeable Films.

We have so far supposed, in treating of surfaces of discontinuity, that they afford no obstacle to the passage of any of the component substances from either of the homogeneous masses to the other. The case, however, must be considered, in which there is a film of matter at the surface of discontinuity which is impermeable to some or all of

* From the experiments of M. E. Duclaux, (*Annales de Chimie et de Physique*, Ser. 4, vol. xxi, p. 383,) it appears that one per cent. of alcohol in water will diminish the superficial tension to .933, the value for pure water being unity. The experiments do not extend to pure alcohol, but the difference of the tensions for mixtures of alcohol and water containing 10 and 20 per cent. water is comparatively small, the tensions being .322 and .336 respectively.

According to the same authority (page 427 of the volume cited), one 3200th part of Castile soap will reduce the superficial tension of water by one-fourth; one 800th part of soap by one-half. These determinations, as well as those relating to alcohol and water, are made by the method of drops, the weight of the drops of different liquids (from the same pipette) being regarded as proportional to their superficial tensions.

M. Athanase Dupré has determined the superficial tensions of solutions of soap by different methods. A statical method gives for one part of common soap in 5000 of water a superficial tension about one-half as great as for pure water, but if the tension be measured on a jet close to the orifice, the value (for the same solution) is sensibly identical with that of pure water. He explains these different values of the superficial tension of the same solution as well as the great effect on the superficial tension which a very small quantity of soap or other trifling impurity may produce, by the tendency of the soap or other substance to form a film on the surface of the liquid. (See *Annales de Chimie et de Physique*, Ser. 4, vol. vii, p. 409, and vol. ix, p. 379.)

the components of the contiguous masses. Such may be the case, for example, when a film of oil is spread on a surface of water, even when the film is too thin to exhibit the properties of the oil in mass. In such cases, if there is communication between the contiguous masses through other parts of the system to which they belong, such that the components in question can pass freely from one mass to the other, the impossibility of a direct passage through the film may be regarded as an immaterial circumstance, so far as states of equilibrium are concerned, and our formulæ will require no change. But when there is no such indirect communication, the potential for any component for which the film is impermeable may have entirely different values on opposite sides of the film, and the case evidently requires a modification of our usual method.

A single consideration will suggest the proper treatment of such cases. If a certain component which is found on both sides of a film cannot pass from either side to the other, the fact that the part of the component which is on one side is the same kind of matter with the part on the other side may be disregarded. All the general relations must hold true, which would hold if they were really different substances. We may therefore write μ_1 for the potential of the component on one side of the film, and μ_2 for the potential of the same substance (to be treated as if it were a different substance) on the other side; m_1^s for the excess of the quantity of the substance on the first side of the film above the quantity which would be on that side of the dividing surface (whether this is determined by the surface of tension or otherwise) if the density of the substance were the same near the dividing surface as at a distance, and m_2^s for a similar quantity relating to the other side of the film and dividing surface. On the same principle, we may use Γ_1 and Γ_2 to denote the values of m_1^s and m_2^s per unit of surface, and $m_1', m_2'', \gamma_1', \gamma_2''$ to denote the quantities of the substance and its densities in the two homogeneous masses.

With such a notation, which may be extended to cases in which the film is impermeable to any number of components, the equations relating to the surface and the contiguous masses will evidently have the same form as if the substances specified by the different suffixes were all really different. The superficial tension will be a function of μ_1 and μ_2 , with the temperature and the potentials for the other components, and $-\Gamma_1, -\Gamma_2$ will be equal to its differential coefficients with respect to μ_1 and μ_2 . In a word, all the general relations which have been demonstrated may be applied to this case, if

we remember always to treat the component as a different substance according as it is found on one side or the other of the impermeable film.

When there is free passage for the component specified by the suffixes $_1$ and $_2$ through other parts of the system, (or through any flaws in the film,) we shall have in case of equilibrium $\mu_1 = \mu_2$. If we wish to obtain the fundamental equation for the surface when satisfying this condition, without reference to other possible states of the surface, we may set a single symbol for μ_1 and μ_2 in the more general form of the fundamental equation. Cases may occur of an impermeability which is not absolute, but which renders the transmission of some of the components exceedingly slow. In such cases, it may be necessary to distinguish at least two different fundamental equations, one relating to a state of approximate equilibrium which may be quickly established, and another relating to the ultimate state of complete equilibrium. The former may be derived from the latter by such substitutions as that just indicated.

The Conditions of Internal Equilibrium for a System of Heterogeneous Fluid Masses without neglect of the Influence of the Surfaces of Discontinuity or of Gravity.

Let us now seek the complete value of the variation of the energy of a system of heterogeneous fluid masses, in which the influence of gravity and of the surfaces of discontinuity shall be included, and deduce from it the conditions of internal equilibrium for such a system. In accordance with the method which has been developed, the intrinsic energy, (*i. e.*, the part of the energy which is independent of gravity,) the entropy, and the quantities of the several components must each be divided into two parts, one of which we regard as belonging to the surfaces which divide approximately homogeneous masses, and the other as belonging to these masses. The elements of intrinsic energy, entropy, etc., relating to an element of surface Ds will be denoted by $D\varepsilon^s$, $D\eta^s$, Dm_1^s , Dm_2^s , etc., and those relating to an element of volume Dv , by $D\varepsilon^v$, $D\eta^v$, Dm_1^v , Dm_2^v , etc. We shall also use Dm^s or ΓDs and Dm^v or γDv to denote the total quantities of matter relating to the elements Ds and Dv respectively. That is,

$$Dm^s = \Gamma Ds = Dm_1^s + Dm_2^s + \text{etc.}, \quad (597)$$

$$Dm^v = \gamma Dv = Dm_1^v + Dm_2^v + \text{etc.} \quad (598)$$

The part of the energy which is due to gravity must also be divided

into two parts, one of which relates to the elements Dm^s , and the other to the elements Dm^v . The complete value of the variation of the energy of the system will be represented by the expression

$$\delta f D\varepsilon^v + \delta f D\varepsilon^s + \delta f g z Dm^v + \delta f g z Dm^s, \quad (599)$$

in which g denotes the force of gravity, and z the height of the element above a fixed horizontal plane.

It will be convenient to limit ourselves at first to the consideration of reversible variations. This will exclude the formation of new masses or surfaces. We may therefore regard any infinitesimal variation in the state of the system as consisting of infinitesimal variations of the quantities relating to its several elements, and bring the sign of variation in the preceding formula after the sign of integration. If we then substitute for $\delta D\varepsilon^v$, $\delta D\varepsilon^s$, δDm^v , δDm^s , the values given by equations (13), (497), (597), (598), we shall have for the condition of equilibrium with respect to reversible variations of the internal state of the system

$$\begin{aligned} & f^t \delta D\eta^v - f^p \delta Dv + f^{\mu_1} \delta Dm_1^v + f^{\mu_2} \delta Dm_2^v + \text{etc.} \\ & + f^t \delta D\eta^s + f^\sigma \delta Ds + f^{\mu_1} \delta Dm_1^s + f^{\mu_2} \delta Dm_2^s + \text{etc.} \\ & + f g \delta z Dm^v + f g z \delta Dm_1^v + f g z \delta Dm_2^v + \text{etc.} \\ & + f g \delta z Dm^s + f g z \delta Dm_1^s + f g z \delta Dm_2^s + \text{etc.} = 0, \quad (600) \end{aligned}$$

Since equation (497) relates to surfaces of discontinuity which are initially in equilibrium, it might seem that this condition, although always necessary for equilibrium, may not always be sufficient. It is evident, however, from the form of the condition, that it includes the particular conditions of equilibrium relating to every possible deformation of the system, or reversible variation in the distribution of entropy or of the several components. It therefore includes all the relations between the different parts of the system which are necessary for equilibrium, so far as reversible variations are concerned. (The necessary relations between the various quantities relating to each element of the masses and surfaces are expressed by the fundamental equation of the mass or surface concerned, or may be immediately derived from it. See pp. 140-144 and 391-393.)

The variations in (600) are subject to the conditions which arise from the nature of the system and from the supposition that the changes in the system are not such as to affect external bodies. This supposition is necessary, unless we are to consider the variations in the state of the external bodies, and is evidently allowable in seeking the conditions of equilibrium which relate to the interior of the sys-

tem.* But before we consider the equations of condition in detail, we may divide the condition of equilibrium (600) into the three conditions

$$f't \delta D\eta^v + f't \delta D\eta^s = 0, \quad (601)$$

$$-f'p \delta Dv + f'\sigma \delta Ds + f'g \delta z Dm^v + f'g \delta z Dm^s = 0, \quad (602)$$

$$\begin{aligned} & f'\mu_1 \delta Dm_1^v + f'\mu_1 \delta Dm_1^s + f'gz \delta Dm_1^v + f'gz \delta Dm_1^s \\ & + f'\mu_2 \delta Dm_2^v + f'\mu_2 \delta Dm_2^s + f'gz \delta Dm_2^v + f'gz \delta Dm_2^s \\ & + \text{etc.} = 0. \end{aligned} \quad (603)$$

For the variations which occur in any one of the three are evidently independent of those which occur in the other two, and the equations of condition will relate to one or another of these conditions separately.

The variations in condition (601) are subject to the condition that the entropy of the whole system shall remain constant. This may be expressed by the equation

$$f' \delta D\eta^v + f' \delta D\eta^s = 0. \quad (604)$$

To satisfy the condition thus limited it is necessary and sufficient that

$$t = \text{const.} \quad (605)$$

throughout the whole system, which is the condition of thermal equilibrium.

The conditions of mechanical equilibrium, or those that relate to the possible deformation of the system, are contained in (602), which may also be written

$$-f'p \delta Dv + f'\sigma \delta Ds + f'g \gamma \delta z Dv + f'g \Gamma \delta z Ds = 0. \quad (606)$$

It will be observed that this condition has the same form as if the different fluids were separated by heavy and elastic membranes without rigidity and having at every point a tension uniform in all directions in the plane of the surface. The variations in this formula,

* We have sometimes given a physical expression to a supposition of this kind, in problems in which the peculiar condition of matter in the vicinity of surfaces of discontinuity was to be neglected, by regarding the system as surrounded by a rigid and impermeable envelop. But the more exact treatment which we are now to give the problem of equilibrium would require us to take account of the influence of the envelop on the immediately adjacent matter. Since this involves the consideration of surfaces of discontinuity between solids and fluids, and we wish to limit ourselves at present to the consideration of the equilibrium of fluid masses, we shall give up the conception of an impermeable envelop, and regard the system as bounded simply by a imaginary surface, which is not a surface of discontinuity. The variations of the system must be such as do not deform the surface, nor affect the matter external to it.

beside their necessary geometrical relations, are subject to the conditions that the external surface of the system, and the lines in which the surfaces of discontinuity meet it, are fixed. The formula may be reduced by any of the usual methods, so as to give the particular conditions of mechanical equilibrium. Perhaps the following method will lead as directly as any to the desired result.

It will be observed the quantities affected by δ in (606) relate exclusively to the position and size of the elements of volume and surface into which the system is divided, and that the variations δp and $\delta\sigma$ do not enter into the formula either explicitly or implicitly. The equations of condition which concern this formula also relate exclusively to the variations of the system of geometrical elements, and do not contain either δp or $\delta\sigma$. Hence, in determining whether the first member of the formula has the value zero for every possible variation of the system of geometrical elements, we may assign to δp and $\delta\sigma$ any values whatever, which may simplify the solution of the problem, without inquiring whether such values are physically possible.

Now when the system is in its initial state, the pressure p , in each of the parts into which the system is divided by the surfaces of tension, is a function of the co-ordinates which determine the position of the element Dv , to which the pressure relates. In the varied state of the system, the element Dv will in general have a different position. Let the variation δp be determined solely by the change in position of the element Dv . This may be expressed by the equation

$$\delta p = \frac{dp}{dx} \delta x + \frac{dp}{dy} \delta y + \frac{dp}{dz} \delta z, \tag{607}$$

in which $\frac{dp}{dx}$, $\frac{dp}{dy}$, $\frac{dp}{dz}$ are determined by the function mentioned, and δx , δy , δz by the variation of the position of the element Dv .

Again, in the initial state of the system the tension σ , in each of the different surfaces of discontinuity, is a function of two co-ordinates, ω_1 , ω_2 , which determine the position of the element Ds . In the varied state of the system, this element will in general have a different position. The change of position may be resolved into a component lying in the surface and another normal to it. Let the variation $\delta\sigma$ be determined solely by the first of these components of the motion of Ds . This may be expressed by the equation

$$\delta\sigma = \frac{d\sigma}{d\omega_1} \delta\omega_1 + \frac{d\sigma}{d\omega_2} \delta\omega_2, \tag{608}$$

in which $\frac{d\sigma}{d\omega_1}$, $\frac{d\sigma}{d\omega_2}$ are determined by the function mentioned, and $\delta\omega_1$, $\delta\omega_2$, by the component of the motion of Ds which lies in the plane of the surface.

With this understanding, which is also to apply to δp and $\delta\sigma$ when contained implicitly in any expression, we shall proceed to the reduction of the condition (606).

With respect to any one of the volumes into which the system is divided by the surfaces of discontinuity, we may write

$$\int p \delta Dv = \delta \int p Dv - \int \delta p Dv.$$

But it is evident that

$$\delta \int p Dv = \int p \delta N Ds,$$

where the second integral relates to the surfaces of discontinuity bounding the volume considered, and δN denotes the normal component of the motion of an element of the surface, measured outward. Hence,

$$\int p \delta Dv = \int p \delta N Ds - \int \delta p Dv.$$

Since this equation is true of each separate volume into which the system is divided, we may write for the whole system

$$\int p \delta Dv = \int (p' - p'') \delta N Ds - \int \delta p Dv, \quad (609)$$

where p' and p'' denote the pressures on opposite sides of the element Ds , and δN is measured toward the side specified by double accents.

Again, for each of the surfaces of discontinuity, taken separately,

$$\int \sigma \delta Ds = \delta \int \sigma Ds - \int \delta \sigma Ds,$$

and

$$\delta \int \sigma Ds = \int \sigma (c_1 + c_2) \delta N Ds + \int \sigma \delta T Dl,$$

where c_1 and c_2 denote the principal curvatures of the surface, (positive, when the centers are on the side opposite to that toward which δN is measured,) Dl an element of the perimeter of the surface, and δT the component of the motion of this element which lies in the plane of the surface and is perpendicular to the perimeter, (positive, when it extends the surface). Hence we have for the whole system

$$\int \sigma \delta Ds = \int \sigma (c_1 + c_2) \delta N Ds + \int \Sigma (\sigma \delta T) Dl - \int \delta \sigma Ds, \quad (610)$$

where the integration of the elements Dl extends to all the lines in which the surfaces of discontinuity meet, and the symbol Σ denotes a summation with respect to the several surfaces which meet in such a line.

By equations (609) and (610), the general condition of mechanical equilibrium is reduced to the form

$$-f(p' - p'') \delta N Ds + f \delta p Dv + f \sigma (c_1 + c_2) \delta N Ds + f \Sigma(\sigma \delta T) Dl - f \delta \sigma Ds + f g \gamma \delta z Dv + f g \Gamma \delta z Ds = 0.$$

Arranging and combining terms, we have

$$f(g \gamma \delta z + \delta p) Dv + f[(p'' - p') \delta N + \sigma (c_1 + c_2) \delta N + g \Gamma \delta z - \delta \sigma] Ds + f \Sigma(\sigma \delta T) Dl = 0. \quad (611)$$

To satisfy this condition, it is evidently necessary that the coefficients of Dv , Ds , and Dl shall vanish throughout the system.

In order that the coefficient of Dv shall vanish, it is necessary and sufficient that, in each of the masses into which the system is divided by the surfaces of tension, p shall be a function of z alone, such that

$$\frac{dp}{dz} = -g \gamma. \quad (612)$$

In order that the coefficient of Ds shall vanish in all cases, it is necessary and sufficient that it shall vanish for normal and for tangential movements of the surface. For normal movements we may write

$$\delta \sigma = 0, \quad \text{and} \quad \delta z = \cos \mathcal{S} \delta N,$$

where \mathcal{S} denotes the angle which the normal makes with a vertical line. The first condition therefore gives the equation

$$p' - p'' = \sigma (c_1 + c_2) + g \Gamma \cos \mathcal{S}, \quad (613)$$

which must hold true at every point in every surface of discontinuity. The condition with respect to tangential movements shows that in each surface of tension σ is a function of z alone, such that

$$\frac{d\sigma}{dz} = g \Gamma. \quad (614)$$

In order that the coefficient of Dl in (611) shall vanish, we must have, for every point in every line in which surfaces of discontinuity meet, and for any infinitesimal displacement of the line,

$$\Sigma(\sigma \delta T) = 0. \quad (615)$$

This condition evidently expresses the same relations between the tensions of the surfaces meeting in the line and the directions of perpendiculars to the line drawn in the planes of the various surfaces, which hold for the magnitudes and directions of forces in equilibrium in a plane.

In condition (603), the variations which relate to any component are to be regarded as having the value zero in any part of the system in

which that substance is not an actual component.* The same is true with respect to the equations of condition, which are of the form

$$\left. \begin{aligned} f \delta Dm_1^y + f \delta Dm_1^s &= 0, \\ f \delta Dm_2^y + f \delta Dm_2^s &= 0, \\ \text{etc.} \end{aligned} \right\} \quad (616)$$

(It is here supposed that the various components are independent, *i. e.*, that none can be formed out of others, and that the parts of the system in which any component actually occurs are not entirely separated by parts in which it does not occur.) To satisfy the condition (603), subject to these equations of condition, it is necessary and sufficient that the conditions

$$\left. \begin{aligned} \mu_1 + gz &= M_1, \\ \mu_2 + gz &= M_2, \\ \text{etc.,} \end{aligned} \right\} \quad (617)$$

(M_1, M_2 , etc. denoting constants,) shall each hold true in those parts of the system in which the substance specified is an actual component. We may here add the condition of equilibrium relative to the possible absorption of any substance (to be specified by the suffix a) by parts of the system of which it is not an actual component, *viz.*, that the expression $\mu_a + gz$ must not have a less value in such parts of the system than in a contiguous part in which the substance is an actual component.

From equation (613) with (605) and (617) we may easily obtain the differential equation of a surface of tension (in the geometrical sense of the term), when p' , p'' , and σ are known in terms of the temperature and potentials. For $e_1 + e_2$ and \mathcal{S} may be expressed in terms of the first and second differential coefficients of z with respect to the horizontal co-ordinates, and p' , p'' , σ , and l in terms of the temperature and potentials. But the temperature is constant, and for each of the potentials we may substitute $-gz$ increased by a constant. We thus obtain an equation in which the only variables are z and its first and second differential coefficients with respect to the horizontal co-ordinates. But it will rarely be necessary to use so exact a method. Within moderate differences of level, we may regard γ' , γ'' , and σ as constant. We may then integrate the equation [derived from (612)]

$$d(p' - p'') = g(\gamma'' - \gamma') dz,$$

* The term *actual component* has been defined for homogeneous masses on page 117, and the definition may be extended to surfaces of discontinuity. It will be observed that if a substance is an actual component of either of the masses separated by a surface of discontinuity, it must be regarded as an actual component for that surface, as well as when it occurs at the surface but not in either of the contiguous masses.

which will give

$$\rho' - \rho'' = g(\gamma'' - \gamma')z, \tag{618}$$

where z is to be measured from the horizontal plane for which $\rho' = \rho''$. Substituting this value in (613), and neglecting the term containing l' , we have

$$c_1 + c_2 = \frac{g(\gamma'' - \gamma')}{\sigma}z, \tag{619}$$

where the coefficient of z is to be regarded as constant. Now the value of z cannot be very large, in any surface of sensible dimensions, unless $\gamma'' - \gamma'$ is very small. We may therefore consider this equation as practically exact, unless the densities of the contiguous masses are very nearly equal. If we substitute for the sum of the curvatures its value in terms of the differential coefficients of z with respect to the horizontal rectangular co-ordinates, x and y , we have

$$\frac{\left(1 + \frac{dz^2}{dy^2}\right) \frac{d^2z}{dx^2} - 2 \frac{dz}{dx} \frac{dz}{dy} \frac{d^2z}{dx dy} + \left(1 + \frac{dz^2}{dx^2}\right) \frac{d^2z}{dy^2}}{\left(1 + \frac{dz^2}{dx^2} + \frac{dz^2}{dy^2}\right)^{\frac{3}{2}}} = \frac{g(\gamma'' - \gamma')}{\sigma}z. \tag{620}$$

With regard to the sign of the root in the denominator of the fraction, it is to be observed that, if we always take the positive value of the root, the value of the whole fraction will be positive or negative according as the greater concavity is turned upward or downward. But we wish the value of the fraction to be positive when the greater concavity is turned toward the mass specified by a single accent. We should therefore take the positive or negative value of the root according as this mass is above or below the surface.

The particular conditions of equilibrium which are given in the last paragraph but one may be regarded in general as the conditions of chemical equilibrium between the different parts of the system, since they relate to the separate components.* But such a designation is not entirely appropriate unless the number of components is greater than one. In no case are the conditions of mechanical equilibrium entirely independent of those which relate to temperature and the potentials. For the conditions (612) and (614) may be regarded as consequences of (605) and (617) in virtue of the necessary relations (98) and (508).†

* Concerning another kind of conditions of chemical equilibrium, which relate to the molecular arrangement of the components, and not to their sensible distribution in space, see pages 197-203.

† Compare page 206, where a similar problem is treated without regard to the influence of the surfaces of discontinuity.

The mechanical conditions of equilibrium, however, have an especial importance, since we may always regard them as satisfied in any liquid (and not decidedly viscous) mass in which no sensible motions are observable. In such a mass, when isolated, the attainment of mechanical equilibrium will take place very soon; thermal and chemical equilibrium will follow more slowly. The thermal equilibrium will generally require less time for its approximate attainment than the chemical; but the processes by which the latter is produced will generally cause certain inequalities of temperature until a state of complete equilibrium is reached.

When a surface of discontinuity has more components than one which do not occur in the contiguous masses, the adjustment of the potentials for these components in accordance with equations (617) may take place very slowly, or not at all, for want of sufficient mobility in the components of the surface. But when this surface has only one component which does not occur in the contiguous masses, and the temperature and potentials in these masses satisfy the conditions of equilibrium, the potential for the component peculiar to the surface will very quickly conform to the law expressed in (617), since this is a necessary consequence of the condition of mechanical equilibrium (614) in connection with the conditions relating to temperature and the potentials which we have supposed to be satisfied. The necessary distribution of the substance peculiar to the surface will be brought about by expansions and contractions of the surface. If the surface meets a third mass containing this component and no other which is foreign to the masses divided by the surface, the potential for this component in the surface will of course be determined by that in the mass which it meets.

The particular conditions of mechanical equilibrium (612)–(615), which may be regarded as expressing the relations which must subsist between contiguous portions of a fluid system in a state of mechanical equilibrium, are serviceable in determining whether a given system is or is not in such a state. But the mechanical theorems which relate to finite parts of the system, although they may be deduced from these conditions by integration, may generally be more easily obtained by a suitable application of the general condition of mechanical equilibrium (606), or by the application of ordinary mechanical principles to the system regarded as subject to the forces indicated by this equation.

It will be observed that the conditions of equilibrium relating to temperature and the potentials are not affected by the surfaces of

discontinuity. [Compare (228) and (234).] * Since a phase cannot vary continuously without variations of the temperature or the potentials, it follows from these conditions that the phase at any point in a fluid system which has the same independently variable components throughout, and is in equilibrium under the influence of gravity, must be one of a certain number of phases which are completely determined by the phase at any given point and the difference of level of the two points considered. If the phases throughout the fluid system satisfy the general condition of practical stability for phases existing in large masses (viz., that the pressure shall be the least consistent with the temperature and potentials), they will be entirely determined by the phase at any given point and the differences of level. (Compare page 210, where the subject is treated without regard to the influence of the surfaces of discontinuity.)

Conditions of equilibrium relating to irreversible changes.—The conditions of equilibrium relating to the absorption by any part of the system of substances which are not actual components of that part have been given on page 448. Those relating to the formation of new masses and surfaces are included in the conditions of stability relating to such changes, and are not always distinguishable from them. They are evidently independent of the action of gravity. We have already discussed the conditions of stability with respect to the formation of new fluid masses within a homogeneous fluid and at the surface when two such masses meet (see pages 416–429), as well as the condition relating to the possibility of a change in the nature of a surface of discontinuity. (See pages 400–403, where the surface considered is plane, but the result may easily be extended to curved surfaces.) We shall hereafter consider, in some of the more important cases, the conditions of stability with respect to the formation of new masses and surfaces which are peculiar to lines in which several surfaces of discontinuity meet, and points in which several such lines meet.

Conditions of stability relating to the whole system.—Beside the conditions of stability relating to very small parts of a system, which are substantially independent of the action of gravity, and are discussed elsewhere, there are other conditions, which relate to the

* If the fluid system is divided into separate masses by solid diaphragms which are permeable to all the components of the fluids independently, the conditions of equilibrium of the fluids relating to temperature and the potentials will not be affected. (Compare page 139.) The propositions which follow in the above paragraph may be extended to this case.

whole system or to considerable parts of it. To determine the question of the stability of a given fluid system under the influence of gravity, when all the conditions of equilibrium are satisfied as well as those conditions of stability which relate to small parts of the system taken separately, we may use the method described on page 413, the demonstration of which (pages 411, 412) will not require any essential modification on account of gravity.

When the variations of temperature and of the quantities M_1, M_2 , etc. [see (617)] involved in the changes considered are so small that they may be neglected, the condition of stability takes a very simple form, as we have already seen to be the case with respect to a system uninfluenced by gravity. (See page 415.)

We have to consider a varied state of the system in which the total entropy and the total quantities of the various components are unchanged, and all variations vanish at the exterior of the system,—in which, moreover, the conditions of equilibrium relating to temperature and the potentials are satisfied, and the relations expressed by the fundamental equations of the masses and surfaces are to be regarded as satisfied, although the state of the system is not one of complete equilibrium. Let us imagine the state of the system to vary continuously in the course of time in accordance with these conditions and use the symbol d to denote the simultaneous changes which take place at any instant. If we denote the total energy of the system by E , the value of dE may be expanded like that of δE in (599) and (600), and then reduced (since the values of $t, \mu_1 + gz, \mu_2 + gz$, etc. are uniform throughout the system, and the total entropy and total quantities of the several components are constant) to the form

$$\begin{aligned} dE &= -j'p dDv + j'g dz Dm^v + j'\sigma dDs + j'g dz Dm^s \\ &= -j'p dDv + j'g \gamma dz Dv + j'\sigma dDs + j'g \Gamma dz Ds, \end{aligned} \quad (621)$$

where the integrations relate to the elements expressed by the symbol D . The value of p at any point in any of the various masses, and that of σ at any point in any of the various surfaces of discontinuity are entirely determined by the temperature and potentials at the point considered. If the variations of t and M_1, M_2 , etc. are to be neglected, the variations of p and σ will be determined solely by the change in position of the point considered. Therefore, by (612) and (614),

$$dp = -g \gamma dz, \quad d\sigma = g \Gamma dz;$$

and

$$\begin{aligned} dE &= -f p dDv - f dp Dv + f \sigma dDs + f d\sigma Ds \\ &= -df p Dv + df \sigma Ds. \end{aligned} \tag{622}$$

If we now integrate with respect to d , commencing at the given state of the system, we obtain

$$\Delta E = -\Delta f p Dv + \Delta f \sigma Ds, \tag{623}$$

where Δ denotes the value of a quantity in a varied state of the system diminished by its value in the given state. This is true for finite variations, and is therefore true for infinitesimal variations without neglect of the infinitesimals of the higher orders. The condition of stability is therefore that

$$\Delta f p Dv - \Delta f \sigma Ds < 0, \tag{624}$$

or that the quantity

$$f p Dv - f \sigma Ds \tag{625}$$

has a maximum value, the values of p and σ , for each different mass or surface, being regarded as determined functions of z . (In ordinary cases σ may be regarded as constant in each surface of discontinuity, and p as a linear function of z in each different mass.) It may easily be shown (compare page 416) that this condition is always *sufficient* for stability with reference to motion of surfaces of discontinuity, even when the variations of t , M_1 , M_2 , etc. cannot be neglected in the determination of the *necessary* condition of stability with respect to such changes.

On the Possibility of the Formation of a New Surface of Discontinuity where several Surfaces of Discontinuity meet.

When more than three surfaces of discontinuity between homogeneous masses meet along a line, we may conceive of a new surface being formed between any two of the masses which do not meet in a surface in the original state of the system. The condition of stability with respect to the formation of such a surface may be easily obtained by the consideration of the limit between stability and instability, as exemplified by a system which is in equilibrium when a very small surface of the kind is formed.

To fix our ideas, let us suppose that there are four homogeneous masses A, B, C, and D, which meet one another in four surfaces, which we may call A-B, B-C, C-D, and D-A, these surfaces all meeting along a line L. This is indicated in figure 11 by a section of the sur-

faces cutting the line L at right angles at a point O . In an infinitesimal variation of the state of the system, we may conceive of a small surface being formed between A and C (to be called $A-C$), so that the section of the surfaces of discontinuity by the same plane takes the form indicated in figure 12. Let us suppose that




FIG. 11.




FIG. 12.




FIG. 13.

the condition of equilibrium (615) is satisfied both for the line L in which the surfaces of discontinuity meet in the original state of the system, and for the two such lines (which we may call L' and L'') in the varied state of the system, at least at the points O' and O'' where they are cut by the plane of section. We may therefore form a quadrilateral of which the sides $\alpha\beta$, $\beta\gamma$, $\gamma\delta$, $\delta\alpha$ are equal in numerical value to the tensions of the several surfaces $A-B$, $B-C$, $C-D$, $D-A$, and are parallel to the normals to these surfaces at the point O in the original state of the system. In like manner, for the varied state of the system we can construct two triangles having similar relations to the surfaces of discontinuity meeting at O' and O'' . But the directions of the normals to the surfaces $A-B$ and $B-C$ at O' and to $C-D$ and $D-A$ at O'' in the varied state of the system differ infinitely little from the directions of the corresponding normals at O in the initial state. We may therefore regard $\alpha\beta$, $\beta\gamma$ as two sides of the triangle representing the surfaces meeting at O' , and $\gamma\delta$, $\delta\alpha$ as two sides of the triangle representing the surfaces meeting at O'' . Therefore, if we join $\alpha\gamma$, this line will represent the direction of the normal to the surface $A-C$, and the value of its tension. If the tension of a surface between such masses as A and C had been greater than that represented by $\alpha\gamma$, it is evident that the initial state of the system of surfaces (represented in figure 11) would have been stable with respect to the possible formation of any such surface. If the tension had been less, the state of the system would have been at least practically unstable. To determine whether it is unstable in the strict sense of the term, or whether or not it is prop-

erly to be regarded as in equilibrium, would require a more refined analysis than we have used.*

The result which we have obtained may be generalized as follows. When more than three surfaces of discontinuity in a fluid system meet in equilibrium along a line, with respect to the surfaces and masses immediately adjacent to any point of this line we may form a polygon of which the angular points shall correspond in order to the different masses separated by the surfaces of discontinuity, and the sides to these surfaces, each side being perpendicular to the corresponding surface, and equal to its tension. With respect to the formation of new surfaces of discontinuity in the vicinity of the point especially considered, the system is stable, if every diagonal of the polygon is less, and practically unstable, if any diagonal is greater, than the tension which would belong to the surface of discontinuity between the corresponding masses. In the limiting case, when the diagonal is exactly equal to the tension of the corresponding surface, the system may often be determined to be unstable by the application of the principle enunciated to an adjacent point of the line in which the surfaces of discontinuity meet. But when, in the polygons constructed for all points of the line, no diagonal is in any case greater

* We may here remark that a nearer approximation in the theory of equilibrium and stability might be attained, by taking special account, in our general equations, of the lines in which surfaces of discontinuity meet. These lines might be treated in a manner entirely analogous to that in which we have treated surfaces of discontinuity. We might recognize linear densities of energy, of entropy, and of the several substances which occur about the line, also a certain linear tension. With respect to these quantities and the temperature and potentials, relations would hold analogous to those which have been demonstrated for surfaces of discontinuity. (See pp. 391-393.) If the sum of the tensions of the lines L' and L'' , mentioned above, is greater than the tension of the line L , this line will be in strictness stable (although practically unstable) with respect to the formation of a surface between A and C , when the tension of such a surface is a little less than that represented by the diagonal ay .

The different use of the term *practically unstable* in different parts of this paper need not create confusion, since the general meaning of the term is in all cases the same. A system is called practically unstable when a very small (not necessarily indefinitely small) disturbance or variation in its condition will produce a considerable change. In the former part of this paper, in which the influence of surfaces of discontinuity was neglected, a system was regarded as practically unstable when such a result would be produced by a disturbance of the same order of magnitude as the quantities relating to surfaces of discontinuity which were neglected. But where surfaces of discontinuity are considered, a system is not regarded as practically unstable, unless the disturbance which will produce such a result is very small compared with the quantities relating to surfaces of discontinuity of any appreciable magnitude.

than the tension of the corresponding surface, but a certain diagonal is equal to the tension in the polygons constructed for a finite portion of the line, farther investigations are necessary to determine the stability of the system. For this purpose, the method described on page 413 is evidently applicable.

A similar proposition may be enunciated in many cases with respect to a point about which the angular space is divided into solid angles by surfaces of discontinuity. If these surfaces are in equilibrium, we can always form a closed solid figure without re-entrant angles of which the angular points shall correspond to the several masses, the edges to the surfaces of discontinuity, and the sides to the lines in which these edges meet, the edges being perpendicular to the corresponding surfaces, and equal to their tensions, and the sides being perpendicular to the corresponding lines. Now if the solid angles in the physical system are such as may be subtended by the sides and bases of a triangular prism enclosing the vertical point, or can be derived from such by deformation, the figure representing the tensions will have the form of two triangular pyramids on opposite sides of the same base, and the system will be stable or practically unstable with respect to the formation of a surface between the masses which only meet in a point, according as the tension of a surface between such masses is greater or less than the diagonal joining the corresponding angular points of the solid representing the tensions. This will easily appear on consideration of the case in which a very small surface between the masses would be in equilibrium.

The Conditions of Stability for Fluids relating to the Formation of a New Phase at a Line in which Three Surfaces of Discontinuity meet.

With regard to the formation of new phases there are particular conditions of stability which relate to lines in which several surfaces of discontinuity meet. We may limit ourselves to the case in which there are three such surfaces, this being the only one of frequent occurrence, and may treat them as meeting in a straight line. It will be convenient to commence by considering the equilibrium of a system in which such a line is replaced by a filament of a different phase.

Let us suppose that three homogeneous fluid masses, A, B, and C, are separated by cylindrical (or plane) surfaces, A-B, B-C, C-A, which at first meet in a straight line, each of the surface-tensions σ_{AB} , σ_{BC} , σ_{CA} being less than the sum of the other two. Let us suppose that the

system is then modified by the introduction of a fourth fluid mass D, which is placed between A, B, and C, and is separated from them by cylindrical surfaces D-A, D-B, D-C meeting A-B, B-C, and C-A in straight lines. The general form of the surfaces is shown by figure 14, in which the full lines represent a section perpendicular to all the surfaces. The system thus modified is to be in equilibrium, as well as the original system, the position of the surfaces A-B, B-C, C-A being unchanged. That the last condition is consistent with equilibrium will appear from the following mechanical considerations.




FIG. 14.




FIG. 15.




FIG. 16.

Let v_D denote the volume of the mass D per unit of length or the area of the curvilinear triangle abc . Equilibrium is evidently possible for any values of the surface-tensions (if only σ_{AB} , σ_{BC} , σ_{CA} satisfy the condition mentioned above, and the tensions of the three surfaces meeting at each of the edges of D satisfy a similar condition) with any value (not too large) of v_D , if the edges of D are constrained to remain in the original surfaces A-B, B-C, and C-A, or these surfaces extended, if necessary, without change of curvature. (In certain cases one of the surfaces D-A, D-B, D-C may disappear and D will be bounded by only two cylindrical surfaces.) We may therefore regard the system as maintained in equilibrium by forces applied to the edges of D and acting at right angles to A-B, B-C, C-A. The same forces would keep the system in equilibrium if D were rigid. They must therefore have a zero resultant, since the nature of the mass D is immaterial when it is rigid, and no forces external to the system would be required to keep a corresponding part of the original system in equilibrium. But it is evident from the points of application and directions of these forces that they cannot have a zero resultant unless each force is zero. We may therefore introduce a fourth mass D without disturbing the parts which remain of the surfaces A-B, B-C, C-D.

It will be observed that all the angles at a , b , c , and d in figure 14 are entirely determined by the six surface-tensions σ_{AB} , σ_{BC} , σ_{CA} , σ_{DA} , σ_{DB} , σ_{DC} . [See (615).] The angles may be derived from the tensions

by the following construction, which will also indicate some important properties. If we form a triangle $\alpha\beta\gamma$ (figure 15 or 16) having sides equal to σ_{AB} , σ_{BC} , σ_{CA} , the angles of the triangle will be supplements of the angles at d . To fix our ideas, we may suppose the sides of the triangle to be perpendicular to the surfaces at d . Upon $\beta\gamma$ we may then construct (as in figure 16) a triangle $\beta\gamma\delta'$ having sides equal to σ_{BC} , σ_{DC} , σ_{DB} , upon $\gamma\alpha$ a triangle $\gamma\alpha\delta''$ having sides equal to σ_{CA} , σ_{DA} , σ_{DC} , and upon $\alpha\beta$ a triangle $\alpha\beta\delta'''$ having sides equal to σ_{AB} , σ_{DB} , σ_{DA} . These triangles are to be on the same sides of the lines $\beta\gamma$, $\gamma\alpha$, $\alpha\beta$, respectively, as the triangle $\alpha\beta\gamma$. The angles of these triangles will be supplements of the angles of the surfaces of discontinuity at a , b , and c . Thus $\beta\gamma\delta' = dab$, and $\alpha\gamma\delta'' = dba$. Now if δ' and δ'' fall together in a single point δ within the triangle $\alpha\beta\gamma$, δ''' will fall in the same point, as in figure 15. In this case we shall have $\beta\gamma\delta + \alpha\gamma\delta = \alpha\gamma\beta$, and the three angles of the curvilinear triangle adb will be together equal to two right angles. The same will be true of the three angles of each of the triangles bdc , cda , and hence of the three angles of the triangle abc . But if δ' , δ'' , δ''' do not fall together in the same point within the triangle $\alpha\beta\gamma$, it is either possible to bring these points to coincide within the triangle by increasing some or all of the tensions σ_{DA} , σ_{DB} , σ_{DC} , or to effect the same result by diminishing some or all of these tensions. (This will easily appear when one of the points δ' , δ'' , δ''' falls within the triangle, if we let the two tensions which determine this point remain constant, and the third tension vary. When all the points δ' , δ'' , δ''' fall without the triangle $\alpha\beta\gamma$, we may suppose the greatest of the tensions σ_{DA} , σ_{DB} , σ_{DC} —the two greatest, when these are equal, and all three when they all are equal—to diminish until one of the points δ' , δ'' , δ''' is brought within the triangle $\alpha\beta\gamma$.) In the first case we may say that the tensions of the new surfaces are too small to be represented by the distances of an internal point from the vertices of the triangle representing the tensions of the original surfaces (or, for brevity, that they are too small to be represented as in figure 15); in the second case we may say that they are too great to be thus represented. In the first case, the sum of the angles in each of the triangles adb , bdc , cda is less than two right angles (compare figures 14 and 16): in the second case, each pair of the triangles $\alpha\beta\delta'''$, $\beta\gamma\delta'$, $\gamma\alpha\delta''$ will overlap, at least when the tensions σ_{DA} , σ_{DB} , σ_{DC} are only a little too great to be represented as in figure 15, and the sum of the angles of each of the triangles adb , bdc , cda will be greater than two right angles.

Let us denote by v_A, v_B, v_C the portions of v_D which were originally occupied by the masses A, B, C, respectively, by s_{DA}, s_{DB}, s_{DC} , the areas of the surfaces specified per unit of length of the mass D, and by s_{AB}, s_{BC}, s_{CA} , the areas of the surfaces specified which were replaced by the mass D per unit of its length. In numerical value, v_A, v_B, v_C will be equal to the areas of the curvilinear triangles bcd, cad, abd ; and $s_{DA}, s_{DB}, s_{DC}, s_{AB}, s_{BC}, s_{CA}$ to the lengths of the lines bc, ca, ab, cd, ad, bd . Also let

$$W_S = \sigma_{DA} s_{DA} + \sigma_{DB} s_{DB} + \sigma_{DC} s_{DC} - \sigma_{AB} s_{AB} - \sigma_{BC} s_{BC} - \sigma_{CA} s_{CA}, \quad (626)$$

and
$$W_V = p_D v_D - p_A v_A - p_B v_B - p_C v_C. \quad (627)$$

The general condition of mechanical equilibrium for a system of homogeneous masses not influenced by gravity, when the exterior of the whole system is fixed, may be written

$$\Sigma (\sigma \delta s) - \Sigma (p \delta v) = 0. \quad (628)$$

[See (606).] If we apply this both to the original system consisting of the masses A, B, and C, and to the system modified by the introduction of the mass D, and take the difference of the results, supposing the deformation of the system to be the same in each case, we shall have

$$\begin{aligned} &\sigma_{DA} \delta s_{DA} + \sigma_{DB} \delta s_{DB} + \sigma_{DC} \delta s_{DC} - \sigma_{AB} \delta s_{AB} - \sigma_{BC} \delta s_{BC} \\ &- \sigma_{CA} \delta s_{CA} - p_D \delta v_D + p_A \delta v_A + p_B \delta v_B + p_C \delta v_C = 0. \end{aligned} \quad (629)$$

In view of this relation, if we differentiate (626) and (627) regarding all quantities except the pressures as variable, we obtain

$$\begin{aligned} dW_S - dW_V = &s_{DA} d\sigma_{DA} + s_{DB} d\sigma_{DB} + s_{DC} d\sigma_{DC} \\ &- s_{AB} d\sigma_{AB} - s_{BC} d\sigma_{BC} - s_{CA} d\sigma_{CA}. \end{aligned} \quad (630)$$

Let us now suppose the system to vary in size, remaining always similar to itself in form, and that the tensions diminish in the same ratio as lines, while the pressures remain constant. Such changes will evidently not impair the equilibrium. Since all the quantities $s_{DA}, \sigma_{DA}, s_{DB}, \sigma_{DB}$, etc. vary in the same ratio,

$$s_{DA} d\sigma_{DA} = \frac{1}{2} d(\sigma_{DA} s_{DA}), \quad s_{DB} d\sigma_{DB} = \frac{1}{2} d(\sigma_{DB} s_{DB}), \quad \text{etc.} \quad (631)$$

We have therefore by integration of (630)

$$W_S - W_V = \frac{1}{2} (\sigma_{DA} s_{DA} + \sigma_{DB} s_{DB} + \sigma_{DC} s_{DC} - \sigma_{AB} s_{AB} - \sigma_{BC} s_{BC} - \sigma_{CA} s_{CA}), \quad (632)$$

whence, by (626),

$$W_S = 2 W_V, \quad (633)$$

The condition of stability for the system when the pressures and tensions are regarded as constant, and the position of the surfaces

A-B, B-C, C-A as fixed, is that $W_s - W_v$ shall be a minimum under the same conditions. [See (549).] Now for any constant values of the tensions and of p_A, p_B, p_C , we may make v_D so small that when it varies, the system remaining in equilibrium, (which will in general require a variation of p_D), we may neglect the curvatures of the lines da, db, dc , and regard the figure $abcd$ as remaining similar to itself. For the *total curvature* (*i. e.*, the curvature measured in degrees) of each of the lines ab, bc, ca may be regarded as constant, being equal to the constant difference of the sum of the angles of one of the curvilinear triangles adb, bdc, cda and two right angles. Therefore, when v_D is very small, and the system is so deformed that equilibrium would be preserved if p_D had the proper variation, but this pressure as well as the others and all the tensions remain constant, W_s will vary as the lines in the figure $abcd$, and W_v as the square of these lines. Therefore, for such deformations,

$$W_v \propto W_s^2.$$

This shows that the system cannot be stable for constant pressures and tensions when v_D is small and W_v is positive, since $W_s - W_v$ will not be a minimum. It also shows that the system is stable when W_v is negative. For, to determine whether $W_s - W_v$ is a minimum for constant values of the pressures and tensions, it will evidently be sufficient to consider such varied forms of the system as give the least value to $W_s - W_v$ for any value of v_D in connection with the constant pressures and tensions. And it may easily be shown that such forms of the system are those which would preserve equilibrium if p_D had the proper value.

These results will enable us to determine the most important questions relating to the stability of a line along which three homogeneous fluids A, B, C meet, with respect to the formation of a different fluid D. The components of D must of course be such as are found in the surrounding bodies. We shall regard p_D and $\sigma_{DA}, \sigma_{DB}, \sigma_{DC}$ as determined by that phase of D which satisfies the conditions of equilibrium with the other bodies relating to temperature and the potentials. These quantities are therefore determinable, by means of the fundamental equations of the mass D and of the surfaces D-A, D-B, D-C, from the temperature and potentials of the given system.

Let us first consider the case in which the tensions, thus determined, can be represented as in figure 15, and p_D has a value consistent with the equilibrium of a small mass such as we have been considering. It appears from the preceding discussion that when v_D is

sufficiently small the figure $abcd$ may be regarded as rectilinear, and that its angles are entirely determined by its tensions. Hence the ratios of v_A, v_B, v_C, v_D , for sufficiently small values of v_D , are determined by the tensions alone, and for convenience in calculating these ratios, we may suppose p_A, p_B, p_C to be equal, which will make the figure $abcd$ absolutely rectilinear, and make p_D equal to the other pressures, since it is supposed that this quantity has the value necessary for equilibrium. We may obtain a simple expression for the ratios of v_A, v_B, v_C, v_D in terms of the tensions in the following manner. We shall write $[D B C]$, $[D C A]$, etc., to denote the areas of triangles having sides equal to the tensions of the surfaces between the masses specified.

$$\begin{aligned}
 v_A : v_B &:: \text{triangle } b d c : \text{triangle } a d c \\
 &:: bc \sin bed : ac \sin acd \\
 &:: \sin bac \sin bed : \sin abc \sin acd \\
 &:: \sin \gamma \delta \beta \sin \delta \alpha \beta : \sin \gamma \delta \alpha \sin \delta \beta \alpha \\
 &:: \sin \gamma \delta \beta \delta \beta : \sin \gamma \delta \alpha \delta \alpha \\
 &:: \text{triangle } \gamma \delta \beta : \text{triangle } \gamma \delta \alpha \\
 &:: [D B C] : [D C A].
 \end{aligned}$$

Hence,

$$v_A : v_B : v_C : v_D :: [D B C] : [D C A] : [D A B] : [A B C], \quad (634)$$

where

$$\frac{1}{4} \sqrt{[(\sigma_{AB} + \sigma_{BC} + \sigma_{CA})(\sigma_{AB} + \sigma_{BC} - \sigma_{CA})(\sigma_{BC} + \sigma_{CA} - \sigma_{AB})(\sigma_{CA} + \sigma_{AB} - \sigma_{BC})]}$$

may be written for $[A B C]$, and analogous expressions for the other symbols, the sign $\sqrt{}$ denoting the positive root of the necessarily positive expression which follows. This proportion will hold true in any case of equilibrium, when the tensions satisfy the condition mentioned and v_D is sufficiently small. Now if $p_A = p_B = p_C$, p_D will have the same value, and we shall have by (627) $W_V = 0$, and by (633) $W_S = 0$. But when v_D is very small, the value of W_S is entirely determined by the tensions and v_D . Therefore, whenever the tensions satisfy the condition supposed, and v_D is very small (whether p_A, p_B, p_C are equal or unequal),

$$0 = W_S = W_V = p_D v_D - p_A v_A - p_B v_B - p_C v_C, \quad (635)$$

which with (634) gives

$$p_D = \frac{[D B C] p_A + [D C A] p_B + [D A B] p_C}{[D B C] + [D C A] + [D A B]}. \quad (636)$$

Since this is the only value of p_D for which equilibrium is possible when

the tensions satisfy the condition supposed and v_D is small, it follows that when p_D has a less value, the line where the fluids A, B, C meet is stable with respect to the formation of the fluid D. When p_D has a greater value, if such a line can exist at all, it must be at least practically unstable, *i. e.*, if only a very small mass of the fluid D should be formed it would tend to increase.

Let us next consider the case in which the tensions of the new surfaces are too small to be represented as in figure 15. If the pressures and tensions are consistent with equilibrium for any very small value of v_D , the angles of each of the curvilinear triangles adb , bdc , $cd a$ will be together less than two right angles, and the lines ab , bc , ca , will be convex toward the mass D. For given values of the pressures and tensions, it will be easy to determine the magnitude of v_D . For the tensions will give the total curvatures (in degrees) of the lines ab , bc , ca ; and the pressures will give the radii of curvature. These lines are thus completely determined. In order that v_D shall be very small it is evidently necessary that p_D shall be less than the other pressures. Yet if the tensions of the new surfaces are only a very little too small to be represented as in figure 15, v_D may be quite small when the value of p_D is only a little less than that given by equation (636). In any case, when the tensions of the new surfaces are too small to be represented as in figure 15, and v_D is small, W_v is negative, and the equilibrium of the mass D is stable. Moreover, $W_s - W_v$, which represents the work necessary to form the mass D with its surfaces in place of the other masses and surfaces, is negative.

With respect to the stability of a line in which the surfaces A-B, B-C, C-A meet, when the tensions of the new surfaces are too small to be represented as in figure 15, we first observe that when the pressures and tensions are such as to make v_D moderately small but not so small as to be neglected, [this will be when p_D is somewhat smaller than the second member of (636),—more or less smaller according as the tensions differ more or less from such as are represented in figure 15,] the equilibrium of such a line as that supposed (if it is capable of existing at all) is at least practically unstable. For greater values of p_D (with the same values of the other pressures and the tensions) the same will be true. For somewhat smaller values of p_D , the mass of the phase D which will be formed will be so small, that we may neglect this mass and regard the surfaces A-B, B-C, C-A as meeting in a line in stable equilibrium. For still smaller values of p_D , we may likewise regard the surfaces A-B, B-C, C-A as capable

of meeting in stable equilibrium. It may be observed that when v_D , as determined by our equations, becomes quite insensible, the conception of a small mass D having the properties deducible from our equations ceases to be accurate, since the matter in the vicinity of a line where these surfaces of discontinuity meet must be in a peculiar state of equilibrium not recognized by our equations.* But this cannot affect the validity of our conclusion with respect to the stability of the line in question.

The case remains to be considered in which the tensions of the new surfaces are too great to be represented as in figure 15. Let us suppose that they are not very much too great to be thus represented. When the pressures are such as to make v_D moderately small (in case of equilibrium) but not so small that the mass D to which it relates ceases to have the properties of matter in mass, [this will be when p_D is somewhat greater than the second member of (636),—more or less greater according as the tensions differ more or less from such as are represented in figure 15,] the line where the surfaces A-B, B-C, C-A meet will be in stable equilibrium with respect to the formation of such a mass as we have considered, since $W_s - W_v$ will be positive. The same will be true for less values of p_D . For greater values of p_D , the value of $W_s - W_v$, which measures the stability with respect to the kind of change considered, diminishes. It does not vanish, according to our equations, for finite values of p_D . But these equations are not to be trusted beyond the limit at which the mass D ceases to be of sensible magnitude.

But when the tensions are such as we now suppose, we must also consider the possible formation of a mass D within a closed figure in which the surfaces D-A, D-B, D-C meet together (with the surfaces A-B, B-C, C-A) in two opposite points. If such a figure is to be in equilibrium, the six tensions must be such as can be represented by

* See note on page 455. We may here add that the linear tension there mentioned may have a negative value. This would be the case with respect to a line in which three surfaces of discontinuity are regarded as meeting, but where nevertheless there really exists in stable equilibrium a filament of different phase from the three surrounding masses. The value of the linear tension for the supposed line, would be nearly equal to the value of $W_s - W_v$ for the actually existing filament. (For the exact value of the linear tension it would be necessary to add the sum of the linear tensions of the three edges of the filament.) We may regard two soap-bubbles adhering together as an example of this case. The reader will easily convince himself that in an exact treatment of the equilibrium of such a double bubble we must recognize a certain negative tension in the line of intersection of the three surfaces of discontinuity.

the six distances of four points in space (see page 455),—a condition which evidently agrees with the supposition which we have made. If we denote by w_v the work gained in forming the mass D (of such size and form as to be in equilibrium) in place of the other masses, and by w_s the work expended in forming the new surfaces in place of the old, it may easily be shown by a method similar to that used on page 459 that $w_s = \frac{3}{2}w_v$. From this we obtain $w_s - w_v = \frac{1}{2}w_v$. This is evidently positive when p_D is greater than the other pressures. But it diminishes with increase of p_D , as easily appears from the equivalent expression $\frac{1}{3}w_s$. Hence the line of intersection of the surfaces of discontinuity A-B, B-C, C-A is stable for values of p_D greater than the other pressures (and therefore for all values of p_D) so long as our method is to be regarded as accurate, which will be so long as the mass D which would be in equilibrium has a sensible size.

In certain cases in which the tensions of the new surfaces are much too large to be represented as in figure 15, the reasoning of the two last paragraphs will cease to be applicable. These are cases in which the six tensions cannot be represented by the sides of a tetrahedron. It is not necessary to discuss these cases, which are distinguished by the different shape which the mass D would take if it should be formed, since it is evident that they can constitute no exception to the results which we have obtained. For an increase of the values of σ_{DA} , σ_{DB} , σ_{DC} cannot favor the formation of D, and hence cannot impair the stability of the line considered, as deduced from our equations. Nor can an increase of these tensions essentially affect the fact that the stability thus demonstrated may fail to be realized when p_D is considerably greater than the other pressures, since the *a priori* demonstration of the stability of any one of the surfaces A-B, B-C, C-A, taken singly, is subject to the limitation mentioned. (See page 426.)

*The Condition of Stability for Fluids relating to the Formation
of a New Phase at a Point where the Vertices of
Four Different Masses meet.*

Let four different fluid masses A, B, C, D meet about a point, so as to form the six surfaces of discontinuity A-B, B-C, C-A, D-A, D-B, D-C, which meet in the four lines A-B-C, B-C-D, C-D-A, D-A-B, these lines meeting in the vertical point. Let us suppose the system stable in other respects, and consider the conditions of stability for the vertical point with respect to the possible formation of a different fluid mass E.

If the system can be in equilibrium when the vertical point has been replaced by a mass E against which the four masses A, B, C, D abut, being truncated at their vertices, it is evident that E will have four vertices, at each of which six surfaces of discontinuity meet. (Thus at one vertex there will be the surfaces formed by A, B, C, and E.) The tensions of each set of six surfaces (like those of the six surfaces formed by A, B, C, and D) must therefore be such that they can be represented by the six edges of a tetrahedron. When the tensions do not satisfy these relations, there will be no particular condition of stability for the point about which A, B, C, and D meet, since if a mass E should be formed, it would distribute itself along some of the lines or surfaces which meet at the vertical point, and it is therefore sufficient to consider the stability of these lines and surfaces. We shall suppose that the relations mentioned are satisfied.

If we denote by W_V the work gained in forming the mass E (of such size and form as to be in equilibrium) in place of the portions of the other masses which are suppressed, and by W_S the work expended in forming the new surfaces in place of the old, it may easily be shown by a method similar to that used on page 459 that

$$W_S = \frac{3}{2} W_V, \tag{637}$$

whence
$$W_S - W_V = \frac{1}{2} W_V; \tag{638}$$

also, that when the volume E is small, the equilibrium of E will be stable or unstable according as W_S and W_V are negative or positive.

A critical relation for the tensions is that which makes equilibrium possible for the system of the five masses A, B, C, D, E, when all the surfaces are plane. The ten tensions may then be represented in magnitude and direction by the ten distances of five points in space $\alpha, \beta, \gamma, \delta, \varepsilon$, viz., the tension of A-B and the direction of its normal by the line $\alpha\beta$, etc. The point ε will lie within the tetrahedron formed by the other points. If we write v_E for the volume of E, and v_A, v_B, v_C, v_D for the volumes of the parts of the other masses which are suppressed to make room for E, we have evidently

$$W_V = p_E v_E - p_A v_A - p_B v_B - p_C v_C - p_D v_D. \tag{639}$$

Hence, when all the surfaces are plane, $W_V = 0$, and $W_S = 0$. Now equilibrium is always possible for a given small value of v_E with any given values of the tensions and of p_A, p_B, p_C, p_D . When the tensions satisfy the critical relation, $W_S = 0$, if $p_A = p_B = p_C = p_D$. But when v_E is small and constant, the value of W_S must be independent of p_A, p_B, p_C, p_D , since the angles of the surfaces are determined by the tensions and their curvatures may be neglected. Hence, $W_S = 0$, and

$W_v = 0$, when the critical relation is satisfied and v_E small. This gives

$$p_E = \frac{v_A p_A + v_B p_B + v_C p_C + v_D p_D}{v_E}. \tag{640}$$

In calculating the ratios of v_A, v_B, v_C, v_D, v_E , we may suppose all the surfaces to be plane. Then E will have the form of a tetrahedron, the vertices of which may be called a, b, c, d, (each vertex being named after the mass which is not found there,) and v_A, v_B, v_C, v_D will be the volumes of the tetrahedra into which it may be divided by planes passing through its edges and an interior point e. The volumes of these tetrahedra are proportional to those of the five tetrahedra of the figure $\alpha \beta \gamma \delta \epsilon$, as will easily appear if we recollect that the line a b is common to the surfaces C-D, D-E, E-C, and therefore perpendicular to the surface common to the lines $\gamma \delta, \delta \epsilon, \epsilon \gamma$, *i. e.*, to the surface $\gamma \delta \epsilon$, and so in other cases, (it will be observed that γ, δ , and ϵ are the letters which do not correspond to a or b); also that the surface a b c is the surface D-E and therefore perpendicular to $\delta \epsilon$, etc. Let tetr abcd, trian abc, etc. denote the volume of the tetrahedron or the area of the triangle specified, $\sin(ab, bc)$, $\sin(abc, dbc)$, $\sin(abc, ad)$, etc. the sines of the angles made by the lines and surfaces specified, and [B C D E], [C D E A], etc., the volumes of tetrahedra having edges equal to the tensions of the surfaces between the masses specified. Then, since we may express the volume of a tetrahedron either by $\frac{1}{3}$ of the product of one side, an edge leading to the opposite vertex, and the sine of the angle which these make, or by $\frac{2}{3}$ of the product of two sides divided by the common edge and multiplied by the sine of the included angle,

$$\begin{aligned} v_A : v_B &:: \text{tetr bcde} : \text{tetr acde} \\ &:: bc \sin(bc, cde) : ac \sin(ac, cde) \\ &:: \sin(ba, ac) \sin(bc, cde) : \sin(ab, bc) \sin(ac, cde) \\ &:: \sin(\gamma \delta \epsilon, \beta \delta \epsilon) \sin(\alpha \delta \epsilon, \alpha \beta) : \sin(\gamma \delta \epsilon, \alpha \delta \epsilon) \sin(\beta \delta \epsilon, \alpha \beta) \\ &:: \frac{\text{tetr } \gamma \beta \delta \epsilon \text{ tetr } \beta \alpha \delta \epsilon}{\text{trian } \beta \delta \epsilon \text{ trian } \alpha \delta \epsilon} : \frac{\text{tetr } \gamma \alpha \delta \epsilon \text{ tetr } \alpha \beta \delta \epsilon}{\text{trian } \alpha \delta \epsilon \text{ trian } \beta \delta \epsilon} \\ &:: \text{tetr } \gamma \beta \delta \epsilon : \text{tetr } \gamma \alpha \delta \epsilon \\ &:: [\text{B C D E}] : [\text{C D E A}]. \end{aligned}$$

Hence,

$$v_A : v_B : v_C : v_D :: [\text{B C D E}] : [\text{C D E A}] : [\text{D E A B}] : [\text{E A B C}], \tag{641}$$

and (640) may be written

$$p_E = \frac{[\text{B C D E}]p_A + [\text{C D E A}]p_B + [\text{D E A B}]p_C + [\text{E A B C}]p_D}{[\text{B C D E}] + [\text{C D E A}] + [\text{D E A B}] + [\text{E A B C}]} \tag{642}$$

If the value of p_E is less than this, when the tensions satisfy the critical relation, the point where vertices of the masses A, B, C, D meet is stable with respect to the formation of any mass of the nature of E. But if the value of p_E is greater, either the masses A, B, C, D cannot meet at a point in equilibrium, or the equilibrium will be at least practically unstable.

When the tensions of the new surfaces are too small to satisfy the critical relation with the other tensions, these surfaces will be convex toward E; when their tensions are too great for that relation, the surfaces will be concave toward E. In the first case, W_V is negative, and the equilibrium of the five masses A, B, C, D, E is stable, but the equilibrium of the four masses A, B, C, D meeting at a point is impossible or at least practically unstable. This is subject to the limitation that when p_E is sufficiently small the mass E which will form will be so small that it may be neglected. This will only be the case when p_E is smaller—in general considerably smaller—than the second number of (642). In the second case, the equilibrium of the five masses A, B, C, D, E will be unstable, but the equilibrium of the four masses A, B, C, D will be stable unless v_E (calculated for the case of the five masses) is of insensible magnitude. This will only be the case when p_E is greater—in general considerably greater—than the second member of (642).

Liquid Films.

When a fluid exists in the form of a thin film between other fluids, the great inequality of its extension in different directions will give rise to certain peculiar properties, even when its thickness is sufficient for its interior to have the properties of matter in mass. The frequent occurrence of such films, and the remarkable properties which they exhibit, entitle them to particular consideration. To fix our ideas, we shall suppose that the film is liquid and that the contiguous fluids are gaseous. The reader will observe our results are not dependent, so far as their general character is concerned, upon this supposition.

Let us imagine the film to be divided by surfaces perpendicular to its sides into small portions of which all the dimensions are of the same order of magnitude as the thickness of the film,—such portions to be called *elements of the film*,—it is evident that far less time will in general be required for the attainment of approximate equilibrium between the different parts of any such element and the other fluids which are immediately contiguous, than for the attainment of equi-

librium between all the different elements of the film. There will accordingly be a time, commencing shortly after the formation of the film, in which its separate elements may be regarded as satisfying the conditions of internal equilibrium, and of equilibrium with the contiguous gases, while they may not satisfy all the conditions of equilibrium with each other. It is when the changes due to this want of complete equilibrium take place so slowly that the film appears to be at rest, except so far as it accommodates itself to any change in the external conditions to which it is subjected, that the characteristic properties of the film are most striking and most sharply defined.

Let us therefore consider the properties which will belong to a film sufficiently thick for its interior to have the properties of matter in mass, in virtue of the approximate equilibrium of all its elements taken separately, when the matter contained in each element is regarded as invariable, with the exception of certain substances which are components of the contiguous gas-masses and have their potentials thereby determined. The occurrence of a film which precisely satisfies these conditions may be exceptional, but the discussion of this somewhat ideal case will enable us to understand the principal laws which determine the behavior of liquid films in general.

Let us first consider the properties which will belong to each element of the film under the conditions mentioned. Let us suppose the element extended, while the temperature and the potentials which are determined by the contiguous gas-masses are unchanged. If the film has no components except those of which the potentials are maintained constant, there will be no variation of tension in its surfaces. The same will be true when the film has only one component of which the potential is not maintained constant, provided that this is a component of the interior of the film and not of its surface alone. If we regard the thickness of the film as determined by *dividing surfaces* which make the surface-density of this component vanish, the thickness will vary inversely as the area of the element of the film, but no change will be produced in the nature or the tension of its surfaces. If, however, the single component of which the potential is not maintained constant is confined to the surfaces of the film, an extension of the element will generally produce a decrease in the potential of this component, and an increase of tension. This will certainly be true in those cases in which the component shows a tendency to distribute itself with a uniform superficial density.

When the film has two or more components of which the potentials are not maintained constant by the contiguous gas masses, they will not in general exist in the same proportion in the interior of the film as on its surfaces, but those components which diminish the tensions will be found in greater proportion on the surfaces. When the film is extended, there will therefore not be enough of these substances to keep up the same volume- and surface-densities as before, and the deficiency will cause a certain increase of tension. The value of the *elasticity of the film*, (*i. e.*, the infinitesimal increase of the united tensions of its surfaces divided by the infinitesimal increase of area in a unit of surface), may be calculated from the quantities which specify the nature of the film, when the fundamental equations of the interior mass, of the contiguous gas-masses, and of the two surfaces of discontinuity are known. We may illustrate this by a simple example.

Let us suppose that the two surfaces of a plane film are entirely alike, that the contiguous gas-masses are identical in phase, and that they determine the potentials of all the components of the film except two. Let us call these components S_1 and S_2 , the latter denoting that which occurs in greater proportion on the surface than in the interior of the film. Let us denote by γ_1 and γ_2 the densities of these components in the interior of the film, by λ the thickness of the film determined by such dividing surfaces as make the surface-density of S_1 vanish (see page 397), by $\Gamma_{2(1)}$ the surface-density of the other component as determined by the same surfaces, by σ and s the tension and area of one of these surfaces, and by E the elasticity of the film when extended under the supposition that the total quantities of S_1 and S_2 in the part of the film extended are invariable, as also the temperature and the potentials of the other components. From the definition of E we have

$$2 d\sigma = E \frac{ds}{s}, \tag{643}$$

and from the conditions of the extension of the film

$$\frac{ds}{s} = -\frac{d(\lambda \gamma_1)}{\lambda \gamma_1} = -\frac{d(\lambda \gamma_2 + 2 \Gamma_{2(1)})}{\lambda \gamma_2 + 2 \Gamma_{2(1)}}. \tag{644}$$

Hence we obtain

$$\lambda \gamma_1 \frac{ds}{s} = -\gamma_1 d\lambda - \lambda d\gamma_1,$$

$$(\lambda \gamma_2 + 2 \Gamma_{2(1)}) \frac{ds}{s} = -\gamma_2 d\lambda - \lambda d\gamma_2 - 2 d\Gamma_{2(1)};$$

and eliminating $d\lambda$,

$$2\gamma_1\Gamma_{2(1)}\frac{ds}{s} = -\lambda\gamma_1 d\gamma_2 + \lambda\gamma_2 d\gamma_1 - 2\gamma_1 d\Gamma_{2(1)}, \quad (645)$$

If we set
$$r = \frac{\gamma_2}{\gamma_1}, \quad (646)$$

we have
$$dr = \frac{\gamma_1 d\gamma_2 - \gamma_2 d\gamma_1}{\gamma_1^2}, \quad (647)$$

and
$$2\Gamma_{2(1)}\frac{ds}{s} = -\lambda\gamma_1 dr - 2d\Gamma_{2(1)}. \quad (648)$$

With this equation we may eliminate ds from (643). We may also eliminate $d\sigma$ by the necessary relation [see (514)]

$$d\sigma = -\Gamma_{2(1)}d\mu_2.$$

This will give

$$4\Gamma_{2(1)}^2 d\mu_2 = E(\lambda\gamma_1 dr + 2d\Gamma_{2(1)}), \quad (649)$$

or

$$\frac{4\Gamma_{2(1)}^2}{E} = \lambda\gamma_1 \frac{dr}{d\mu_2} + 2\frac{d\Gamma_{2(1)}}{d\mu_2}, \quad (650)$$

where the differential coefficients are to be determined on the conditions that the temperature and all the potentials except μ_1 and μ_2 are constant, and that the pressure in the interior of the film shall remain equal to that in the contiguous gas-masses. The latter condition may be expressed by the equation

$$(\gamma_1 - \gamma_1') d\mu_1 + (\gamma_2 - \gamma_2') d\mu_2 = 0, \quad (651)$$

in which γ_1' and γ_2' denote the densities of S_1 and S_2 in the contiguous gas-masses. [See (98).] When the tension of the surfaces of the film and the pressures in its interior and in the contiguous gas-masses are known in terms of the temperature and potentials, equation (650) will give the value of E in terms of the same variables together with λ .

If we write G_1 and G_2 for the total quantities of S_1 and S_2 per unit of area of the film, we have

$$G_1 = \lambda\gamma_1, \quad (652)$$

$$G_2 = \lambda\gamma_2 + 2\Gamma_{2(1)}. \quad (653)$$

Therefore,

$$G_2 = G_1 r + 2\Gamma_{2(1)},$$

$$\left(\frac{dG_2}{d\mu_2}\right)_{G_1} = \lambda\gamma_1 \frac{dr}{d\mu_2} + 2\frac{d\Gamma_{2(1)}}{d\mu_2}, \quad (654)$$

where the differential coefficients in the second member are to be determined as in (650), and that in the first member with the additional condition that G_1 is constant. Therefore,

$$\frac{4 \Gamma_{2(1)}^2}{E} = \left(\frac{dG_2}{d\mu_2} \right)_{G_1},$$

and

$$E = 4 \Gamma_{2(1)}^2 \left(\frac{d\mu_2}{dG_2} \right)_{G_1}, \tag{655}$$

the last differential coefficient being determined by the same conditions as that in the preceding equation. It will be observed that the value of E will be positive in any ordinary case.

These equations give the elasticity of any element of the film when the temperature and the potentials for the substances which are found in the contiguous gas-masses are regarded as constant, and the potentials for the other components, μ_1 and μ_2 , have had time to equalize themselves throughout the element considered. The increase of tension immediately after a rapid extension will be greater than that given by these equations.

The existence of this elasticity, which has thus been established from *a priori* considerations, is clearly indicated by the phenomena which liquid films present. Yet it is not to be demonstrated simply by comparing the tensions of films of different thickness, even when they are made from the same liquid, for difference of thickness does not necessarily involve any difference of tension. When the phases within the films as well as without are the same, and the surfaces of the films are also the same, there will be no difference of tension. Nor will the tension of the same film be altered, if a part of the interior drains away in the course of time, without affecting the surfaces. In case the thickness of the film is reduced by evaporation, the tension may be either increased or diminished. (The evaporation of the substance S_1 , in the case we have just considered, would diminish the tension.) Yet it may easily be shown that extension increases the tension of a film and contraction diminishes it. When a plane film is held vertically, the tension of the upper portions must evidently be greater than that of the lower. The tensions in every part of the film may be reduced to equality by turning it into a horizontal position. By restoring the original position we may restore the original tensions, or nearly so. It is evident that the same element of the film is capable of supporting very unequal tensions. Nor can this be always attributed to viscosity of the film. For in many cases, if we hold the film nearly horizontal, and elevate first one side and then another, the lighter portions of the film will dart from one side to the other, so as to show a very striking mobility in the film. The differences of tension which cause these rapid movements are only a very

small fraction of the difference of tension in the upper and lower portions of the film when held vertically.

If we account for the power of an element of the film to support an increase of tension by viscosity, it will be necessary to suppose that the viscosity offers a resistance to a deformation of the film in which its surface is enlarged and its thickness diminished, which is enormously great in comparison with the resistance to a deformation in which the film is extended in the direction of one tangent and contracted in the direction of another, while its thickness and the areas of its surfaces remain constant. This is not to be readily admitted as a physical explanation, although to a certain extent the phenomena resemble those which would be caused by such a singular viscosity. (See page 439.) The only natural explanation of the phenomena is that the extension of an element of the film, which is the immediate result of an increase of external force applied to its perimeter, causes an increase of its tension, by which it is brought into true equilibrium with the external forces.

The phenomena to which we have referred are such as are apparent to a very cursory observation. In the following experiment, which is described by M. Plateau,* an increased tension is manifested in a film while contracting after a previous extension. The warmth of a finger brought near to a bubble of soap-water with glycerine, which is thin enough to show colors, causes a spot to appear indicating a diminution of thickness. When the finger is removed, the spot returns to its original color. This indicates a contraction, which would be resisted by any viscosity of the film, and can only be due to an excess of tension in the portion stretched on the return of its original temperature.

We have so far supposed that the film is thick enough for its interior to have the properties of matter in mass. Its properties are then entirely determined by those of the three phases and the two surfaces of discontinuity. From these we can also determine, in part at least, the properties of a film at the limit at which the interior ceases to have the properties of matter in mass. The elasticity of the film, which increases with its thinness, cannot of course vanish at that limit, so that the film cannot become unstable with respect to extension and contraction of its elements immediately after passing that limit.

Yet a certain kind of instability will probably arise, which we may

* "Statique expérimentale et théorique des liquides soumis aux seules forces moléculaires," vol. i, p. 294.

here notice, although it relates to changes in which the condition of the invariability of the quantities of certain components in an element of the film is not satisfied. With respect to variations in the distribution of its components, a film will in general be stable, when its interior has the properties of matter in mass, with the single exception of variations affecting its thickness without any change of phase or of the nature of the surfaces. With respect to this kind of change, which may be brought about by a current in the interior of the film, the equilibrium is neutral. But when the interior ceases to have the properties of matter in mass, it is to be supposed that the equilibrium will generally become unstable in this respect. For it is not likely that the neutral equilibrium will be unaffected by such a change of circumstances, and since the film certainly becomes unstable when it is sufficiently reduced in thickness, it is most natural to suppose that the first effect of diminishing the thickness will be in the direction of instability rather than in that of stability. (We are here considering liquid films between gaseous masses. In certain other cases, the opposite supposition might be more natural, as in respect to a film of water between mercury and air, which would certainly become stable when sufficiently reduced in thickness.)

Let us now return to our former suppositions—that the film is thick enough for the interior to have the properties of matter in mass, and that the matter in each element is invariable, except with respect to those substances which have their potentials determined by the contiguous gas-masses—and consider what conditions are necessary for equilibrium in such a case.

In consequence of the supposed equilibrium of its several elements, such a film may be treated as a simple surface of discontinuity between the contiguous gas-masses (which may be similar or different), whenever its radius of curvature is very large in comparison with its thickness,—a condition which we shall always suppose to be fulfilled. With respect to the film considered in this light, the mechanical conditions of equilibrium will always be satisfied, or very nearly so, as soon as a state of approximate rest is attained, except in those cases in which the film exhibits a decided viscosity. That is, the relations (613), (614), (615) will hold true, when by σ we understand the tension of the film regarded as a simple surface of discontinuity (this is equivalent to the sum of the tensions of the two surfaces of the film), and by Γ its mass per unit of area diminished by the mass of gas which would occupy the same space if the film should be suppressed and the gases should meet at its surface of tension. This

surface of tension of the film will evidently divide the distance between the surfaces of tension for the two surfaces of the film taken separately, in the inverse ratio of their tensions. For practical purposes, we may regard Γ simply as the mass of the film per unit of area. It will be observed that the terms containing Γ in (613) and (614) are not to be neglected in our present application of these equations.

But the mechanical conditions of equilibrium for the film regarded as an approximately homogeneous mass in the form of a thin sheet bounded by two surfaces of discontinuity are not necessarily satisfied when the film is in a state of apparent rest. In fact, these conditions cannot be satisfied (in any place where the force of gravity has an appreciable intensity) unless the film is horizontal. For the pressure in the interior of the film cannot satisfy simultaneously condition (612), which requires it to vary rapidly with the height z , and condition (613) applied separately to the different surfaces, which makes it a certain mean between the pressures in the adjacent gas-masses. Nor can these conditions be deduced from the general condition of mechanical equilibrium (606) or (611), without supposing that the interior of the film is free to move independently of the surfaces, which is contrary to what we have supposed.

Moreover, the potentials of the various components of the film will not in general satisfy conditions (617), and cannot (when the temperature is uniform) unless the film is horizontal. For if these conditions were satisfied, equation (612) would follow as a consequence. (See page 449.)

We may here remark that such a film as we are considering cannot form any exception to the principle indicated on page 450,—that when a surface of discontinuity which satisfies the conditions of mechanical equilibrium has only one component which is not found in the contiguous masses, and these masses satisfy all the conditions of equilibrium, the potential for the component mentioned must satisfy the law expressed in (617), as a consequence of the condition of mechanical equilibrium (614). Therefore, as we have just seen that it is impossible that all the potentials in a liquid film which is not horizontal should conform to (617) when the temperature is uniform, it follows that if a liquid film exhibits any persistence which is not due to viscosity, or to a horizontal position, or to differences of temperature, it must have more than one component of which the potential is not determined by the contiguous gas-masses in accordance with (617).

The difficulties of the quantitative experimental verification of the properties which have been described would be very great, even in cases in which the conditions we have imagined were entirely fulfilled. Yet the general effect of any divergence from these conditions will be easily perceived, and when allowance is made for such divergence, the general behavior of liquid films will be seen to agree with the requirements of theory.

The formation of a liquid film takes place most symmetrically when a bubble of air rises to the top of a mass of the liquid. The motion of the liquid, as it is displaced by the bubble, is evidently such as to stretch the two surfaces in which the liquid meets the air, where these surfaces approach one another. This will cause an increase of tension, which will tend to restrain the extension of the surfaces. The extent to which this effect is produced will vary with the nature of the liquid. Let us suppose that the case is one in which the liquid contains one or more components which, although constituting but a very small part of its mass, greatly reduce its tension. Such components will exist in excess on the surfaces of the liquid. In this case the restraint upon the extension of the surfaces will be considerable, and as the bubble of air rises above the general level of the liquid, the motion of the latter will consist largely of a running out from between the two surfaces. But this running out of the liquid will be greatly retarded by its viscosity as soon as it is reduced to the thickness of a film, and the effect of the extension of the surfaces in increasing their tension will become greater and more permanent as the quantity of liquid diminishes which is available for supplying the substances which go to form the increased surfaces.

We may form a rough estimate of the amount of motion which is possible for the interior of a liquid film, relatively to its exterior, by calculating the descent of water between parallel vertical planes at which the motion of the water is reduced to zero. If we use the coefficient of viscosity as determined by Helmholtz and Piotrowski,* we obtain

$$V = 581 D^2, \tag{656}$$

where V denotes the mean velocity of the water (*i. e.*, that velocity

* Sitzungsberichte der Wiener Akademie, (mathemat.-naturwiss. Classe), B. xl, S. 607. The calculation of formula (656) and that of the factor ($\frac{5}{3}$) applied to the formula of Poiseuille, to adapt it to a current between plane surfaces, have been made by means of the general equations of the motion of a viscous liquid as given in the memoir referred to.

which, if it were uniform throughout the whole space between the fixed planes, would give the same discharge of water as the actual variable velocity) expressed in millimetres per second, and D denotes the distance in millimetres between the fixed planes, which is supposed to be very small in proportion to their other dimensions. This is for the temperature of 24.5° C. For the same temperature, the experiments of Poiseuille* give

$$V = 337 D^2$$

for the descent of water in long capillary tubes, which is equivalent to

$$V = 899 D^2 \quad (657)$$

for descent between parallel planes. The numerical coefficient in this equation differs considerably from that in (656), which is derived from experiments of an entirely different nature, but we may at least conclude that in a film of a liquid which has a viscosity and specific gravity not very different from those of water at the temperature mentioned the mean velocity of the interior relatively to the surfaces will not probably exceed $1000 D^2$. This is a velocity of $.1^{\text{mm}}$ per second for a thickness of $.01^{\text{mm}}$, $.06^{\text{mm}}$ per *minute* for a thickness of $.001$ (which corresponds to the red of the fifth order in a film of water), and $.036^{\text{mm}}$ per *hour* for a thickness of $.0001^{\text{mm}}$ (which corresponds to the white of the first order). Such an internal current is evidently consistent with great persistence of the film, especially in those cases in which the film can exist in a state of the greatest tenuity. On the other hand, the above equations give so large a value of V for thicknesses of 1^{mm} or $.1^{\text{mm}}$, that the film can evidently be formed without carrying up any great weight of liquid, and any such thicknesses as these can have only a momentary existence.

A little consideration will show that the phenomenon is essentially of the same nature when films are formed in any other way, as by dipping a ring or the mouth of a cup in the liquid and then withdrawing it. When the film is formed in the mouth of a pipe, it may sometimes be extended so as to form a large bubble. Since the elasticity (*i. e.*, the increase of the tension with extension) is greater in the thinner parts, the thicker parts will be most extended, and the effect of this process (so far as it is not modified by gravity) will be to diminish the ratio of the greatest to the least thickness of the film. During this extension, as well as at other times, the increased elasticity due to imperfect communication of heat, etc., will serve to protect the bubble from fracture by shocks received from the air or the

* *Ibid.*, p. 653; or *Mémoires des Savants Étrangers*, vol. ix, p. 532.

pipe. If the bubble is now laid upon a suitable support, the condition (613) will be realized almost instantly. The bubble will then tend toward conformity with condition (614), the lighter portions rising to the top, more or less slowly, according to the viscosity of the film. The resulting difference of thickness between the upper and the lower parts of the bubble is due partly to the greater tension to which the upper parts are subject, and partly to a difference in the matter of which they are composed. When the film has only two components of which the potentials are not determined by the contiguous atmosphere, the laws which govern the arrangement of the elements of the film may be very simply expressed. If we call these components S_1 and S_2 , the latter denoting (as on page 469) that which exists in excess at the surface, one element of the film will tend toward the same level with another, or a higher, or a lower level, according as the quantity of S_2 bears the same ratio to the quantity of S_1 in the first element as in the second, or a greater, or a less ratio.

When a film, however formed, satisfies both the conditions (613) and (614), its thickness being sufficient for its interior to have the properties of matter in mass, the interior will still be subject to the slow current which we have already described, if it is truly fluid, however great its viscosity may be. It seems probable, however, that this process is often totally arrested by a certain gelatinous consistency of the mass in question, in virtue of which, although practically fluid in its behavior with reference to ordinary stresses, it may have the properties of a solid with respect to such very small stresses as those which are caused by gravity in the interior of a very thin film which satisfies the conditions (613) and (614).

However this may be, there is another cause which is often more potent in producing changes in a film, when the conditions just mentioned are approximately satisfied, than the action of gravity on its interior. This will be seen if we turn our attention to the edge where the film is terminated. At such an edge we generally find a liquid mass, continuous in phase with the interior of the film, which is bounded by concave surfaces, and in which the pressure is therefore less than in the interior of the film. This liquid mass therefore exerts a strong suction upon the interior of the film, by which its thickness is rapidly reduced. This effect is best seen when a film which has been formed in a ring is held in a vertical position. Unless the film is very viscous, its diminished thickness near the edge causes a rapid upward current on each side, while the central portion slowly

descends. Also at the bottom of the film, where the edge is nearly horizontal, portions which have become thinned escape from their position of unstable equilibrium beneath heavier portions, and pass upwards, traversing the central portion of the film until they find a position of stable equilibrium. By these processes, the whole film is rapidly reduced in thickness.

The energy of the suction which produces these effects may be inferred from the following considerations. The pressure in the slender liquid mass which encircles the film is of course variable, being greater in the lower portions than in the upper, but it is everywhere less than the pressure of the atmosphere. Let us take a point where the pressure is less than that of the atmosphere by an amount represented by a column of the liquid one centimetre in height. (It is probable that much greater differences of pressure occur.) At a point near by in the interior of the film the pressure is that of the atmosphere. Now if the difference of pressure of these two points were distributed uniformly through the space of one centimetre, the intensity of its action would be exactly equal to that of gravity. But since the change of pressure must take place very suddenly (in a small fraction of a millimetre), its effect in producing a current in a limited space must be enormously great compared with that of gravity.

Since the process just described is connected with the descent of the liquid in the mass encircling the film, we may regard it as another example of the downward tendency of the interior of the film. There is a third way in which this descent may take place, when the principal component of the interior is volatile, viz., through the air. Thus, in the case of a film of soap-water, if we suppose the atmosphere to be of such humidity that the potential for water at a level mid-way between the top and bottom of the film has the same value in the atmosphere as in the film, it may easily be shown that evaporation will take place in the upper portions and condensation in the lower. These processes, if the atmosphere were otherwise undisturbed, would occasion currents of diffusion and other currents, the general effect of which would be to carry the moisture downward. Such a precise adjustment would be hardly attainable, and the processes described would not be so rapid as to have a practical importance.

But when the potential for water in the atmosphere differs considerably from that in the film, as in the case of a film of soap-water in a dry atmosphere, or a film of soap-water with glycerine in a moist

atmosphere, the effect of evaporation or condensation is not to be neglected. In the first case, the diminution of the thickness of the film will be accelerated, in the second, retarded. In the case of the film containing glycerine, it should be observed that the water condensed cannot in all respects replace the fluid carried down by the internal current but that the two processes together will tend to wash out the glycerine from the film.

But when a component which greatly diminishes the tension of the film, although forming but a small fraction of its mass, (therefore existing in excess at the surface,) is volatile, the effect of evaporation and condensation may be considerable, even when the mean value of the potential for that component is the same in the film as in the surrounding atmosphere. To illustrate this, let us take the simple case of two components S_1 and S_2 , as before. (See page 469.) It appears from equation (508) that the potentials must vary in the film with the height z , since the tension does, and from (98) that these variations must (very nearly) satisfy the relation

$$\gamma_1 \frac{d\mu_1}{dz} + \gamma_2 \frac{d\mu_2}{dz} = 0, \quad (658)$$

γ_1 and γ_2 denoting the densities of S_1 and S_2 in the interior of the film. The variation of the potential of S_2 as we pass from one level to another is therefore as much more rapid than that of S_1 , as its density in the interior of the film is less. If then the resistances restraining the evaporation, transmission through the atmosphere, and condensation of the two substances are the same, these processes will go on much more rapidly with respect to S_2 . It will be

observed that the values of $\frac{d\mu_1}{dz}$ and $\frac{d\mu_2}{dz}$ will have opposite signs, the tendency of S_1 being to pass down through the atmosphere, and that of S_2 to pass up. Moreover, it may easily be shown that the evaporation or condensation of S_2 will produce a very much greater effect than the evaporation or condensation of the same quantity of S_1 . These effects are really of the same kind. For if condensation of S_2 takes place at the top of the film, it will cause a diminution of tension, and thus occasion an extension of this part of the film, by which its thickness will be reduced, as it would be by evaporation of S_1 . We may infer that it is a general condition of the persistence of liquid films, that the substance which causes the diminution of tension in the upper parts of the film must not be volatile.

But apart from any action of the atmosphere, we have seen that a

film which is truly fluid in its interior is in general subject to a continual diminution of thickness by the internal currents due to gravity and the suction at its edge. Sooner or later, the interior will somewhere cease to have the properties of matter in mass. The film will then probably become unstable with respect to a flux of the interior (see page 473), the thinnest parts tending to become still more thin (apart from any external cause) very much as if there were an attraction between the surfaces of the film, insensible at greater distances, but becoming sensible when the thickness of the film is sufficiently reduced. We should expect this to determine the rupture of the film, and such is doubtless the case with most liquids. In a film of soap-water, however, the rupture does not take place, and the processes which go on can be watched. It is apparent even to a very superficial observation that a film of which the tint is approaching the black exhibits a remarkable instability. The continuous change of tint is interrupted by the breaking out and rapid extension of black spots. That in the formation of these bright spots a separation of different substances takes place, and not simply an extension of a part of the film, is shown by the fact that the film is made thicker at the edge of these spots.

This is very distinctly seen in a plane vertical film, when a single black spot breaks out and spreads rapidly over a considerable area which was before of a nearly uniform tint approaching the black. The edge of the black spot as it spreads is marked as it were by a string of bright beads, which unite together on touching, and thus becoming larger, glide down across the bands of color below. Under favorable circumstances, there is often quite a shower of these bright spots. They are evidently small spots very much thicker—apparently many times thicker—than the part of the film out of which they are formed. Now if the formation of the black spots were due to a simple extension of the film, it is evident that no such appearance would be presented. The thickening of the edge of the film cannot be accounted for by *contraction*. For an extension of the upper portion of the film and contraction of the lower and thicker portion, with descent of the intervening portions, would be far less resisted by viscosity, and far more favored by gravity than such extensions and contractions as would produce the appearances described. But the rapid formation of a thin spot by an internal current would cause an accumulation at the edge of the spot of the material forming the interior of the film, and necessitate a thickening of the film in that place.

That which is most difficult to account for in the formation of the black spots is the arrest of the process by which the film grows thinner. It seems most natural to account for this, *if possible*, by passive resistance to motion due to a very viscous or gelatinous condition of the film. For it does not seem likely that the film, after becoming unstable by the flux of matter from its interior, would become stable (without the support of such resistance) by a continuance of the same process. On the other hand, gelatinous properties are very marked in soap-water which contains somewhat more soap than is best for the formation of films, and it is entirely natural that, even when such properties are wanting in the interior of a mass or thick film of a liquid, they may still exist in the immediate vicinity of the surface (where we know that the soap or some of its components exists in excess), or throughout a film which is so thin that the interior has ceased to have the properties of matter in mass.* But these considerations do not amount to any *a priori* probability of an arrest of the tendency toward an internal current between adjacent elements of a black spot which may differ slightly in thickness, in time to prevent rupture of the film. For, in a thick film, the increase of the tension with the extension, which is necessary for its stability with respect to extension, is connected with an excess of the soap (or of some of its components) at the surface as compared with the interior of the film. With respect to the black spots, although the interior has ceased to have the properties of matter in mass, and any quantitative determinations derived from the surfaces of a mass of the liquid will not be applicable, it is natural to account for the stability with reference to extension by supposing that the same general difference of composition still exists. If therefore we account for the arrest of internal currents by the increasing density of soap or some of its components in the interior of the film, we must still suppose that the characteristic difference of composition in the interior and surface of the film has not been obliterated.

The preceding discussion relates to liquid films between masses of gas. Similar considerations will apply to liquid films between other liquids or between a liquid and a gas, and to films of gas between

* The experiments of M. Plateau (chapter VII of the work already cited) show that this is the case to a very remarkable degree with respect to a solution of saponine. With respect to soap-water, however, they do not indicate any greater superficial viscosity than belongs to pure water. But the resistance to an internal current, such as we are considering, is not necessarily measured by the resistance to such motions as those of the experiments referred to.

masses of liquid. The latter may be formed by gently depositing a liquid drop upon the surface of a mass of the same or a different liquid. This may be done (with suitable liquids) so that the continuity of the air separating the liquid drop and mass is not broken, but a film of air is formed, which, if the liquids are similar, is a counterpart of the liquid film which is formed by a bubble of air rising to the top of a mass of the liquid. (If the bubble has the same volume as the drop, the films will have precisely the same form, as well as the rest of the surfaces which bound the bubble and the drop.) Sometimes, when the weight and momentum of the drop carry it through the surface of the mass on which it falls, it appears surrounded by a complete spherical film of air, which is the counterpart on a small scale of a soap-bubble hovering in air.* Since, however, the substance to which the necessary differences of tension in the film are mainly due is a component of the liquid masses on each side of the air film, the necessary differences of the potential of this substance cannot be permanently maintained, and these films have little persistence compared with films of soap-water in air. In this respect, the case of these air-films is analogous to that of liquid films in an atmosphere containing substances by which their tension is greatly reduced. Compare page 479.

Surfaces of Discontinuity between Solids and Fluids.

We have hitherto treated of surfaces of discontinuity on the supposition that the contiguous masses are fluid. This is by far the most simple case for any rigorous treatment, since the masses are necessarily isotropic both in nature and in their state of strain. In this case, moreover, the mobility of the masses allows a satisfactory experimental verification of the mechanical conditions of equilibrium. On the other hand, the rigidity of solids is in general so great, that any tendency of the surfaces of discontinuity to variation in area or form may be neglected in comparison with the forces which are produced in the interior of the solids by any sensible strains, so that it is not generally necessary to take account of the surfaces of discontinuity in determining the state of strain of solid masses. But we must take account of the nature of the surfaces of discontinuity

* These spherical air-films are easily formed in soap-water. They are distinguishable from ordinary air-bubbles by their general behavior and by their appearance. The two concentric spherical surfaces are distinctly seen, the diameter of one appearing to be about three-quarters as large as that of the other. This is of course an optical illusion, depending upon the index of refraction of the liquid.

between solids and fluids with reference to the tendency toward solidification or dissolution at such surfaces, and also with reference to the tendencies of different fluids to spread over the surfaces of solids.

Let us therefore consider a surface of discontinuity between a fluid and a solid, the latter being either isotropic or of a continuous crystalline structure, and subject to any kind of stress compatible with a state of mechanical equilibrium with the fluid. We shall not exclude the case in which substances foreign to the contiguous masses are present in small quantities at the surface of discontinuity, but we shall suppose that the nature of this surface (*i. e.*, of the non-homogeneous film between the approximately homogeneous masses), is entirely determined by the nature and state of the masses which it separates, and the quantities of the foreign substances which may be present. The notions of the *dividing surface*, and of the *superficial densities* of energy, entropy, and the several components, which we have used with respect to surfaces of discontinuity between fluids (see pages 380 and 386), will evidently apply without modification to the present case. We shall use the suffix ₁ with reference to the substance of the solid, and shall suppose the dividing surface to be determined so as to make the superficial density of this substance vanish. The superficial densities of energy, of entropy, and of the other component substances may then be denoted by our usual symbols (see page 397),

$$\varepsilon_{s(1)}, \quad \eta_{s(1)}, \quad I'_{2(1)}, \quad I'_{3(1)}, \quad \text{etc.}$$

Let the quantity σ be defined by the equation

$$\sigma = \varepsilon_{s(1)} - t \eta_{s(1)} - \mu_2 I'_{2(1)} - \mu_3 I'_{3(1)} - \text{etc.}, \quad (659)$$

in which t denotes the temperature, and μ_2, μ_3 , etc. the potentials for the substances specified at the surface of discontinuity.

As in the case of two fluid masses, (see page 421,) we may regard σ as expressing the work spent in forming a unit of the surface of discontinuity—under certain conditions, which we need not here specify—but it cannot properly be regarded as expressing the tension of the surface. The latter quantity depends upon the work spent in *stretching* the surface, while the quantity σ depends upon the work spent in *forming* the surface. With respect to perfectly fluid masses, these processes are not distinguishable, unless the surface of discontinuity has components which are not found in the contiguous masses, and even in this case, (since the surface must be supposed to be formed out of matter supplied at the same potentials which belong to the matter in the surface,) the work spent in increasing the surface infinitesi-

mally by stretching is identical with that which must be spent in forming an equal infinitesimal amount of new surface. But when one of the masses is solid, and its states of strain are to be distinguished, there is no such equivalence between the stretching of the surface and the forming of new surface.*

With these preliminary notions, we now proceed to discuss the condition of equilibrium which relates to the dissolving of a solid at the surface where it meets a fluid, when the thermal and mechanical conditions of equilibrium are satisfied. It will be necessary for us to consider the case of isotropic and of crystallized bodies separately, since in the former the value of σ is independent of the direction of the surface, except so far as it may be influenced by the state of strain of the solid, while in the latter the value of σ varies greatly with the direction of the surface with respect to the axes of crystallization, and in such a manner as to have a large number of sharply defined minima.† This may be inferred from the phenomena which crystalline bodies present, as will appear more distinctly in the following discussion. Accordingly, while a variation in the direction of an

* This will appear more distinctly if we consider a particular case. Let us consider a thin plane sheet of a crystal in a vacuum (which may be regarded as a limiting case of a very attenuated fluid), and let us suppose that the two surfaces of the sheet are alike. By applying the proper forces to the edges of the sheet, we can make all stress vanish in its interior. The *tensions* of the two surfaces, are in equilibrium with these forces, and are measured by them. But the tensions of the surfaces, thus determined, may evidently have different values in different directions, and are entirely different from the quantity which we denote by σ , which represents the work required to form a unit of the surface by any reversible process, and is not connected with any idea of direction.

In certain cases, however, it appears probable that the values of σ and of the superficial tension will not greatly differ. This is especially true of the numerous bodies which, although generally (and for many purposes properly) regarded as solids, are really very viscous fluids. Even when a body exhibits no fluid properties at its actual temperature, if its surface has been formed at a higher temperature, at which the body was fluid, and the change from the fluid to the solid state has been by insensible gradations, we may suppose that the value of σ coincided with the superficial tension until the body was decidedly solid, and that they will only differ so far as they may be differently affected by subsequent variations of temperature and of the stresses applied to the solid. Moreover, when an amorphous solid is in a state of equilibrium with a solvent, although it may have no fluid properties in its interior, it seems not improbable that the particles at its surface, which have a greater degree of mobility, may so arrange themselves that the value of σ will coincide with the superficial tension, as in the case of fluids.

† The differential coefficients of σ with respect to the direction-cosines of the surface appear to be discontinuous functions of the latter quantities.

element of the surface may be neglected (with respect to its effect on the value of σ) in the case of isotropic solids, it is quite otherwise with crystals. Also, while the surfaces of equilibrium between fluids and soluble isotropic solids are without discontinuities of direction, being in general curved, a crystal in a state of equilibrium with a fluid in which it can dissolve is bounded in general by a broken surface consisting of sensibly plane portions.

For isotropic solids, the conditions of equilibrium may be deduced as follows. If we suppose that the solid is unchanged, except that an infinitesimal portion is dissolved at the surface where it meets the fluid, and that the fluid is considerable in quantity and remains homogeneous, the increment of energy in the vicinity of the surface will be represented by the expression

$$J'[\varepsilon_V' - \varepsilon_V'' + (c_1 + c_2) \varepsilon_{s(1)}] \delta N Ds$$

where Ds denotes an element of the surface, δN the variation in its position (measured normally, and regarded as *negative* when the solid is dissolved), c_1 and c_2 its principal curvatures (positive when their centers lie on the same side as the solid), $\varepsilon_{s(1)}$ the surface-density of energy, ε_V' and ε_V'' the volume-densities of energy in the solid and fluid respectively, and the sign of integration relates to the elements Ds . In like manner, the increments of entropy and of the quantities of the several components in the vicinity of the surface will be

$$J'[\eta_V' - \eta_V'' + (c_1 + c_2) \eta_{s(1)}] \delta N Ds,$$

$$J'[\gamma_1' - \gamma_1''] \delta N Ds,$$

$$J'[-\gamma_2'' + (c_1 + c_2) \Gamma_{2(1)}] \delta N Ds,$$

etc.

The entropy and the matter of different kinds represented by these expressions we may suppose to be derived from the fluid mass. These expressions, therefore, with a change of sign, will represent the increments of entropy and of the quantities of the components in the whole space occupied by the fluid except that which is immediately contiguous to the solid. Since this space may be regarded as constant, the increment of energy in this space may be obtained [according to equation (12)] by multiplying the above expression relating to entropy by $-t$, and those relating to the components by $-\mu_1''$, $-\mu_2$, etc.,* and taking the sum. If to this

* The potential μ_1'' is marked by double accents in order to indicate that its value is to be determined in the fluid mass, and to distinguish it from the potential μ_1'

we add the above expression for the increment of energy near the surface, we obtain the increment of energy for the whole system. Now by (93) we have

$$p'' = -\varepsilon_v'' + t\eta_v'' + \mu_1''\gamma_1'' + \mu_2\gamma_2'' + \text{etc.}$$

By this equation and (659), our expression for the total increment of energy in the system may be reduced to the form

$$J[\varepsilon_v' - t\eta_v' - \mu_1''\gamma_1' + p'' + (c_1 + c_2)\sigma] \delta N Ds. \quad (660)$$

In order that this shall vanish for any values of δN , it is necessary that the coefficient of $\delta N Ds$ shall vanish. This gives for the condition of equilibrium

$$\mu_1'' = \frac{\varepsilon_v' - t\eta_v' + p'' + (c_1 + c_2)\sigma}{\gamma_1'}. \quad (661)$$

This equation is identical with (387), with the exception of the term containing σ , which vanishes when the surface is plane.*

We may also observe that when the solid has no stresses except an isotropic pressure, if the quantity represented by σ is equal to the true tension of the surface, $p'' + (c_1 + c_2)\sigma$ will represent the pressure in the interior of the solid, and the second member of the equation will represent [see equation (93)] the value of the potential in the solid for the substance of which it consists. In this case, therefore, the equation reduces to

$$\mu_1'' = \mu_1',$$

that is, it expresses the equality of the potentials for the substance of the solid in the two masses—the same condition which would subsist if both masses were fluid.

Moreover, the compressibility of all solids is so small that, although σ may not represent the true tension of the surface, nor $p'' + (c_1 + c_2)\sigma$ the true pressure in the solid when its stresses are isotropic, the quantities ε_v' and η_v' if calculated for the pressure $p'' + (c_1 + c_2)\sigma$ with the actual temperature will have sensibly the same values as if calculated for the true pressure of the solid. Hence, the second member

relating to the solid mass (when this is in a state of isotropic stress), which, as we shall see, may not always have the same value. The other potentials μ_2 , etc., have the same values as in (659), and consist of two classes, one of which relates to substances which are components of the fluid mass, (these might be marked by the double accents,) and the other relates to substances found only at the surface of discontinuity. The expressions to be multiplied by the potentials of this latter class all have the value zero.

* In equation (387), the density of the solid is denoted by Γ , which is therefore equivalent to γ_1' in (661).

of equation (661), when the stresses of the solid are sensibly isotropic, is sensibly equal to the potential of the same body at the same temperature but with the pressure $p'' + (c_1 + c_2) \sigma$, and the condition of equilibrium with respect to dissolving for a solid of isotropic stresses may be expressed with sufficient accuracy by saying that the potential for the substance of the solid in the fluid must have this value. In like manner, when the solid is not in a state of isotropic stress, the difference of the two pressures in question will not sensibly affect the values of ε_v' and η_v' , and the value of the second member of the equation may be calculated as if $p'' + (c_1 + c_2) \sigma$ represented the true pressure in the solid in the direction of the normal to the surface. Therefore, if we had taken for granted that the quantity σ represents the tension of a surface between a solid and a fluid, as it does when both masses are fluid, this assumption would not have led us into any practical error in determining the value of the potential μ_1'' which is necessary for equilibrium. On the other hand, if in the case of any amorphous body the value of σ differs notably from the true surface-tension, the latter quantity substituted for σ in (661) will make the second member of the equation equal to the true value of μ_1' , when the stresses are isotropic, but this will not be equal to the value of μ_1'' in case of equilibrium, unless $c_1 + c_2 = 0$.

When the stresses in the solid are not isotropic, equation (661) may be regarded as expressing the condition of equilibrium with respect to the dissolving of the solid, and is to be distinguished from the condition of equilibrium with respect to an increase of solid matter, since the new matter would doubtless be deposited in a state of isotropic stress. (The case would of course be different with crystalline bodies, which are not considered here.) The value of μ_1'' necessary for equilibrium with respect to the formation of new matter is a little less than that necessary for equilibrium with respect to the dissolving of the solid. In regard to the actual behavior of the solid and fluid, all that the theory enables us to predict with certainty is that the solid will not dissolve if the value of the potential μ_1'' is greater than that given by the equation for the solid with its distorting stresses, and that new matter will not be formed if the value of μ_1'' is less than the same equation would give for the case of the solid with isotropic stresses.* It seems probable, however, that

* The possibility that the new solid matter might differ in composition from the original solid is here left out of account. This point has been discussed on pages 134-137, but without reference to the state of strain of the solid or the influence of the curvature of the surface of discontinuity. The statement made above may be

if the fluid in contact with the solid is not renewed, the system will generally find a state of equilibrium in which the outermost portion of the solid will be in a state of isotropic stress. If at first the solid should dissolve, this would supersaturate the fluid, perhaps until a state is reached satisfying the condition of equilibrium with the stressed solid, and then, if not before, a deposition of solid matter in a state of isotropic stress would be likely to commence and go on until the fluid is reduced to a state of equilibrium with this new solid matter.

The action of gravity will not affect the nature of the condition of equilibrium for any single point at which the fluid meets the solid, but it will cause the values of p'' and μ_1'' in (661) to vary according to the laws expressed by (612) and (617). If we suppose that the outer part of the solid is in a state of isotropic stress, which is the most important case, since it is the only one in which the equilibrium is in every sense stable, we have seen that the condition (661) is at least sensibly equivalent to this:—that the potential for the substance of the solid which would belong to the solid mass at the temperature t and the pressure $p'' + (c_1 + c_2) \sigma$ must be equal to μ_1'' . Or, if we denote by (p') the pressure belonging to solid with the temperature t and the potential equal to μ_1'' , the condition may be expressed in the form

$$(p') = p'' + (c_1 + c_2) \sigma. \quad (662)$$

Now if we write γ'' for the total density of the fluid, we have by (612)

$$dp'' = -g \gamma'' dz.$$

By (98)

$$d(p') = \gamma_1' d\mu_1'',$$

and by (617)

$$d\mu_1'' = -g dz;$$

whence

$$d(p') = -g \gamma_1' dz.$$

Accordingly we have

$$d(p') - dp'' = g (\gamma'' - \gamma_1') dz,$$

and

$$(p') - p'' = g (\gamma'' - \gamma_1') z,$$

z being measured from the horizontal plane for which $(p') = p''$. Substituting this value in (662), we obtain

$$c_1 + c_2 = \frac{g (\gamma'' - \gamma_1')}{\sigma} z, \quad (663)$$

generalized so as to hold true of the formation of new solid matter of any kind on the surface as follows:—that new solid matter of any kind will not be formed upon the surface (with more than insensible thickness), if the second member of (661) calculated for such new matter is greater than the potential in the fluid for such matter.

precisely as if both masses were fluid, and σ denoted the tension of their common surface, and (p') the true pressure in the mass specified. [Compare (619).]

The obstacles to an exact experimental realization of these relations are very great, principally from the want of absolute uniformity in the internal structure of amorphous solids, and on account of the passive resistances to the processes which are necessary to bring about a state satisfying the conditions of theoretical equilibrium, but it may be easy to verify the general tendency toward diminution of surface, which is implied in the foregoing equations.*

Let us apply the same method to the case in which the solid is a crystal. The surface between the solid and fluid will now consist of plane portions, the directions of which may be regarded as invari-

* It seems probable that a tendency of this kind plays an important part in some of the phenomena which have been observed with respect to the freezing together of pieces of ice. (See especially Professor Faraday's "Note on Regelation" in the *Proceedings of the Royal Society*, vol. x, p. 440; or in the *Philosophical Magazine*, 4th ser., vol. xxi, p. 146.) Although this is a body of crystalline structure, and the action which takes place is doubtless influenced to a certain extent by the directions of the axes of crystallization, yet, since the phenomena have not been observed to depend upon the orientation of the pieces of ice, we may conclude that the effect, so far as its general character is concerned, is such as might take place with an isotropic body. In other words, for the purposes of a general explanation of the phenomena we may neglect the differences in the values of σ_{IW} (the suffixes are used to indicate that the symbol relates to the surface between ice and water) for different orientations of the axes of crystallization, and also neglect the influence of the surface of discontinuity with respect to crystalline structure, which must be formed by the freezing together of the two masses of ice when the axes of crystallization in the two masses are not similarly directed. In reality, this surface—or the necessity of the formation of such a surface if the pieces of ice freeze together—must exert an influence adverse to their union, measured by a quantity σ_{II} , which is determined for this surface by the same principles as when one of two contiguous masses is fluid, and varies with the orientations of the two systems of crystallographic axes relatively to each other and to the surface. But under the circumstances of the experiment, since we may neglect the possibility of the two systems of axes having precisely the same directions, this influence is probably of a tolerably constant character, and is evidently not sufficient to alter the general nature of the result. In order wholly to prevent the tendency of pieces of ice to freeze together, when meeting in water with curved surfaces and without pressure, it would be necessary that $\sigma_{II} \geq 2\sigma_{IW}$, except so far as the case is modified by passive resistances to change, and by the inequality in the values of σ_{II} and σ_{IW} for different directions of the axes of crystallization.

It will be observed that this view of the phenomena is in harmony with the opinion of Professor Faraday. With respect to the union of pieces of ice as an indirect consequence of pressure, see page 198 of volume xi of the *Proceedings of the Royal Society*; or the *Philosophical Magazine*, 4th ser., vol. xxiii, p. 407.

able. If the crystal grows on one side a distance δN , without other change, the increment of energy in the vicinity of the surface will be

$$(\varepsilon_v' - \varepsilon_v'') s \delta N + \sum' (\varepsilon_{s(1)'} l' \operatorname{cosec} \omega' - \varepsilon_{s(1)} l' \cot \omega') \delta N,$$

where ε_v' and ε_v'' denote the volume-densities of energy in the crystal and fluid respectively, s the area of the side on which the crystal grows, $\varepsilon_{s(1)}$ the surface-density of energy on that side, $\varepsilon_{s(1)'}$ the surface-density of energy on an adjacent side, ω' the external angle of these two sides, l' their common edge, and the symbol \sum' a summation with respect to the different sides adjacent to the first. The increments of entropy and of the quantities of the several components will be represented by analogous formulæ, and if we deduce as on pages 485, 486 the expression for the increase of energy in the whole system due to the growth of the crystal without change of the total entropy or volume, and set this expression equal to zero, we shall obtain for the condition of equilibrium

$$(\varepsilon_v' - t \eta_v' - \mu_1'' \gamma_1' + p'') s \delta N + \sum' (\sigma' l' \operatorname{cosec} \omega' - \sigma l' \cot \omega') \delta N = 0, \quad (664)$$

where σ and σ' relate respectively to the same sides as $\varepsilon_{s(1)}$ and $\varepsilon_{s(1)'}$ in the preceding formula. This gives

$$\mu_1'' = \frac{\varepsilon_v' - t \eta_v' + p''}{\gamma_1'} + \frac{\sum' (\sigma' l' \operatorname{cosec} \omega' - \sigma l' \cot \omega')}{s \gamma_1'}. \quad (665)$$

It will be observed that unless the side especially considered is small or narrow, we may neglect the second fraction in this equation, which will then give the same value of μ_1'' as equation (387), or as equation (661) applied to a plane surface.

Since a similar equation must hold true with respect to every other side of the crystal of which the equilibrium is not affected by meeting some other body, the condition of equilibrium for the crystalline form (when unaffected by gravity) is that the expression

$$\frac{\sum' (\sigma' l' \operatorname{cosec} \omega' - \sigma l' \cot \omega')}{s} \quad (666)$$

shall have the same value for each side of the crystal. (By the value of this expression for any side of the crystal is meant its value when σ and s are determined by that side and the other quantities by the surrounding sides in succession in connection with the first side.) This condition will not be affected by a change in the size of a crystal while its proportions remain the same. But the tendencies of similar crystals toward the form required by this condition, as measured by the inequalities in the composition or the temperature of

the surrounding fluid which would counterbalance them, will be inversely as the linear dimensions of the crystals, as appears from the preceding equation.

If we write v for the volume of a crystal, and $\Sigma(\sigma s)$ for the sum of the areas of all its sides multiplied each by the corresponding value of σ , the numerator and denominator of the fraction (666), multiplied each by δN , may be represented by $\delta \Sigma(\sigma s)$ and δv respectively. The value of the fraction is therefore equal to that of the differential coefficient

$$\frac{d\Sigma(\sigma s)}{dv}$$

as determined by the displacement of a particular side while the other sides are fixed. The condition of equilibrium for the form of a crystal (when the influence of gravity may be neglected) is that the value of this differential coefficient must be independent of the particular side which is supposed to be displaced. For a constant volume of the crystal, $\Sigma(\sigma s)$ has therefore a minimum value when the condition of equilibrium is satisfied, as may easily be proved more directly.

When there are no foreign substances at the surfaces of the crystal, and the surrounding fluid is indefinitely extended, the quantity $\Sigma(\sigma s)$ represents the work required to form the surfaces of the crystal, and the coefficient of $s \delta N$ in (664) with its sign reversed represents the work gained in forming a mass of volume unity like the crystal but regarded as without surfaces. We may denote the work required to form the crystal by

$$W_s - W_v,$$

W_s denoting the work required to form the surfaces [*i. e.*, $\Sigma(\sigma s)$], and W_v the work gained in forming the mass as distinguished from the surfaces. Equation (664) may then be written

$$-\delta W_v + \Sigma(\sigma \delta s) = 0. \tag{667}$$

Now (664) would evidently continue to hold true if the crystal were diminished in size, remaining similar to itself in form and in nature, if the values of σ in all the sides were supposed to diminish in the same ratio as the linear dimensions of the crystal. The variation of W_s would then be determined by the relation

$$dW_s = d\Sigma(\sigma s) = \frac{3}{2} \Sigma(\sigma ds),$$

and that of W_v by (667). Hence,

$$dW_s = \frac{3}{2} dW_v,$$

and, since W_s and W_v vanish together,

$$\begin{aligned} W_s &= \frac{2}{3} W_v, \\ W_s - W_v &= \frac{1}{3} W_s = \frac{1}{2} W_v, \end{aligned} \quad (668)$$

—the same relation which we have before seen to subsist with respect to a spherical mass of fluid as well as in other cases. (See pages 421, 425, 465.)

The equilibrium of the crystal is unstable with respect to variations in size when the surrounding fluid is indefinitely extended, but it may be made stable by limiting the quantity of the fluid.

To take account of the influence of gravity, we must give to μ_1'' and p'' in (665) their average values in the side considered. These coincide (when the fluid is in a state of internal equilibrium) with their values at the center of gravity of the side. The values of γ_1' , ϵ_v' , η_v' may be regarded as constant, so far as the influence of gravity is concerned. Now since by (612) and (617)

$$dp'' = -g \gamma'' dz,$$

and

$$d\mu_1'' = -g dz,$$

we have

$$d(\gamma_1' \mu_1'' - p'') = g(\gamma'' - \gamma_1') dz.$$

Comparing (664), we see that the upper or the lower faces of the crystal will have the greater tendency to grow, (other things being equal,) according as the crystal is lighter or heavier than the fluid. When the densities of the two masses are equal, the effect of gravity on the form of the crystal may be neglected.

In the preceding paragraph the fluid is regarded as in a state of internal equilibrium. If we suppose the composition and temperature of the fluid to be uniform, the condition which will make the effect of gravity vanish will be that

$$\frac{d(\gamma_1' \mu_1'' - p'')}{dz} = 0,$$

when the value of the differential coefficient is determined in accordance with this supposition. This condition reduces to

$$\left(\frac{d\mu_1}{dp}\right)_{t,m}'' = \frac{1}{\gamma_1'},^*$$

which, by equation (92), is equivalent to

$$\left(\frac{dv}{dm_1}\right)_{t,p,m}'' = \frac{1}{\gamma_1'}. \quad (669)$$

* A suffixed m is used to represent all the symbols m_1 , m_2 , etc., except such as may occur in the differential coefficient.

The tendency of a crystal to grow will be greater in the upper or lower parts of the fluid, according as the growth of a crystal at constant temperature and pressure will produce expansion or contraction.

Again, we may suppose the composition of the fluid and its entropy per unit of mass to be uniform. The temperature will then vary with the pressure, that is, with z . We may also suppose the temperature of different crystals or different parts of the same crystal to be determined by the fluid in contact with them. These conditions express a state which may perhaps be realized when the fluid is gently stirred. Owing to the differences of temperature we cannot regard ε_V' and η_V' in (664) as constant, but we may regard their variations as subject to the relation $d\varepsilon_V' = t d\eta_V'$. Therefore, if we make $\eta_V' = 0$ for the mean temperature of the fluid, (which involves no real loss of generality,) we may treat $\varepsilon_V' - t \eta_V'$ as constant. We shall then have for the condition that the effect of gravity shall vanish—

$$\frac{d(\gamma_1' \mu_1'' - \gamma'')}{dz} = 0,$$

which signifies in the present case that

$$\left(\frac{d\mu_1}{dp}\right)_{\eta, m}'' = \frac{1}{\gamma_1'},$$

or, by (90),

$$\left(\frac{dv}{dm_1}\right)_{\eta, p, m}'' = \frac{1}{\gamma_1'} \tag{670}$$

Since the entropy of the crystal is zero, this equation expresses that the dissolving of a small crystal in a considerable quantity of the fluid will produce neither expansion nor contraction, when the pressure is maintained constant and no heat is supplied or taken away.

The manner in which crystals actually grow or dissolve is often principally determined by other differences of phase in the surrounding fluid than those which have been considered in the preceding paragraph. This is especially the case when the crystal is growing or dissolving rapidly. When the great mass of the fluid is considerably supersaturated, the action of the crystal keeps the part immediately contiguous to it nearer the state of exact saturation. The farthest projecting parts of the crystal will therefore be most exposed to the action of the supersaturated fluid, and will grow most rapidly. The same parts of a crystal will dissolve most rapidly in a fluid considerably below saturation.*

* See O. Lehmann "Ueber das Wachstum der Krystalle," *Zeitschrift für Kristallographie und Mineralogie*, Bd. i, S. 453; or the review of the paper in *Wiedemann's Beiblätter*, Bd. ii, S. 1.

But even when the fluid is supersaturated only so much as is necessary in order that the crystal shall grow at all, it is not to be expected that the form in which $\Sigma(\sigma s)$ has a minimum value (or such a modification of that form as may be due to gravity or to the influence of the body supporting the crystal) will always be the ultimate result. For we cannot imagine a body of the internal structure and external form of a crystal to grow or dissolve by an entirely continuous process, or by a process in the same sense continuous as condensation or evaporation between a liquid and gas, or the corresponding processes between an amorphous solid and a fluid. The process is rather to be regarded as periodic, and the formula (664) cannot properly represent the true value of the quantities intended unless δN is equal to the distance between two successive layers of molecules in the crystal, or a multiple of that distance. Since this can hardly be treated as an infinitesimal, we can only conclude with certainty that sensible changes cannot take place for which the expression (664) would have a positive value.*

* That it is necessary that certain relations shall be precisely satisfied in order that equilibrium may subsist between a liquid and gas with respect to evaporation, is explained (see Clausius "Ueber die Art der Bewegung, welche wir Wärme nennen," *Pogg. Ann.*, Bd. c, S. 353; or *Abhand. über die mech. Wärmetheorie*, XIV,) by supposing that a passage of individual molecules from the one mass to the other is continually taking place, so that the slightest circumstance may give the preponderance to the passage of matter in either direction. The same supposition may be applied, at least in many cases, to the equilibrium between amorphous solids and fluids. Also in the case of crystals in equilibrium with fluids, there may be a passage of individual molecules from one mass to the other, so as to cause insensible fluctuations in the mass of the solid. If these fluctuations are such as to cause the occasional deposit or removal of a whole layer of particles, the least cause would be sufficient to make the probability of one kind of change prevail over that of the other, and it would be necessary for equilibrium that the theoretical conditions deduced above should be precisely satisfied. But this supposition seems quite improbable, except with respect to a very small side.

The following view of the molecular state of a crystal when in equilibrium with respect to growth or dissolution appears as probable as any. Since the molecules at the corners and edges of a perfect crystal would be less firmly held in their places than those in the middle of a side, we may suppose that when the condition of theoretical equilibrium (665) is satisfied several of the outermost layers of molecules on each side of the crystal are incomplete toward the edges. The boundaries of these imperfect layers probably fluctuate, as individual molecules attach themselves to the crystal or detach themselves, but not so that a layer is entirely removed (on any side of considerable size), to be restored again simply by the irregularities of the motions of the individual molecules. Single molecules or small groups of molecules may indeed attach themselves to the side of the crystal but they will speedily be dislodged, and if any molecules are thrown out from the middle of a surface, these deficiencies

Let us now examine the special condition of equilibrium which relates to a line at which three different masses meet, when one or more of these masses is solid. If we apply the method of page 685 to a system containing such a line, it is evident that we shall obtain in the expression corresponding to (660), beside the integral relating to the surfaces, a term of the form

$$\int \Sigma(\sigma \delta T) Dl$$

to be interpreted as the similar term in (611), except so far as the definition of σ has been modified in its extension to solid masses. In order that this term shall be incapable of a negative value it is neces-

will also soon be made good; nor will the frequency of these occurrences be such as greatly to affect the general smoothness of the surfaces, except near the edges where the surfaces fall off somewhat, as before described. Now a continued growth on any side of a crystal is impossible unless new layers can be formed. This will require a value of μ_1'' which may exceed that given by equation (665) by a finite quantity. Since the difficulty in the formation of a new layer is at or near the commencement of the formation, the necessary value of μ_1'' may be independent of the area of the side, except when the side is very small. The value of μ_1'' which is necessary for the growth of the crystal will however be different for different kinds of surfaces, and probably will generally be greatest for the surfaces for which σ is least.

On the whole, it seems not improbable that the form of very minute crystals in equilibrium with solvents is principally determined by equation (665), (*i. e.*, by the condition that $\Sigma(\sigma s)$ shall be a minimum for the volume of the crystal except so far as the case is modified by gravity or the contact of other bodies,) but as they grow larger (in a solvent no more supersaturated than is necessary to make them grow at all), the deposition of new matter on the different surfaces will be determined more by the nature (orientation) of the surfaces and less by their size and relations to the surrounding surfaces. As a final result, a large crystal, thus formed, will generally be bounded by those surfaces alone on which the deposit of new matter takes place least readily, with small, perhaps insensible truncations. If one kind of surfaces satisfying this condition cannot form a closed figure, the crystal will be bounded by two or three kinds of surfaces determined by the same condition. The kinds of surface thus determined will probably generally be those for which σ has the least values. But the relative development of the different kinds of sides, even if unmodified by gravity or the contact of other bodies, will not be such as to make $\Sigma(\sigma s)$ a minimum. The growth of the crystal will finally be confined to sides of a single kind.

It does not appear that any part of the operation of removing a layer of molecules presents any especial difficulty so marked as that of commencing a new layer; yet the values of μ_1'' which will just allow the different stages of the process to go on must be slightly different, and therefore, for the continued dissolving of the crystal the value of μ_1'' must be less (by a finite quantity) than that given by equation (665). It seems probable that this would be especially true of those sides for which σ has the least values. The effect of dissolving a crystal (even when it is done as slowly as possible) is therefore to produce a form which probably differs from that of theoretical equilibrium in a direction opposite to that of a growing crystal.

sary that at every point of the line

$$\Sigma(\sigma \delta T) \geq 0 \quad (671)$$

for any *possible* displacement of the line. Those displacements are to be regarded as possible which are not prevented by the solidity of the masses, when the interior of every solid mass is regarded as incapable of motion. At the surfaces between solid and fluid masses, the processes of solidification and dissolution will be possible in some cases, and impossible in others.

The simplest case is when two masses are fluid and the third is solid and insoluble. Let us denote the solid by S, the fluids by A and B, and the angles filled by these fluids by α and β respectively. If the surface of the solid is continuous at the line where it meets the two fluids, the condition of equilibrium reduces to

$$\sigma_{AB} \cos \alpha = \sigma_{BS} - \sigma_{AS}. \quad (672)$$

If the line where these masses meet is at an edge of the solid, the condition of equilibrium is that

$$\text{and } \left. \begin{aligned} \sigma_{AB} \cos \alpha &\leq \sigma_{BS} - \sigma_{AS}, \\ \sigma_{AB} \cos \beta &\leq \sigma_{AS} - \sigma_{BS}; \end{aligned} \right\} \quad (673)$$

which reduces to the preceding when $\alpha + \beta = \pi$. Since the displacement of the line can take place by a purely mechanical process, this condition is capable of a more satisfactory experimental verification than those conditions which relate to processes of solidification and dissolution. Yet the frictional resistance to a displacement of the line is enormously greater than in the case of three fluids, since the relative displacements of contiguous portions of matter are enormously greater. Moreover, foreign substances adhering to the solid are not easily displaced, and cannot be distributed by extensions and contractions of the surface of discontinuity, as in the case of fluid masses. Hence, the distribution of such substances is arbitrary to a greater extent than in the case of fluid masses, (in which a single foreign substance in any surface of discontinuity is uniformly distributed, and a greater number are at least so distributed as to make the tension of the surface uniform,) and the presence of these substances will modify the conditions of equilibrium in a more irregular manner.

If one or more of three surfaces of discontinuity which meet in a line divides an amorphous solid from a fluid in which it is soluble, such a surface is to be regarded as movable, and the particular conditions involved in (671) will be accordingly modified. If the soluble solid is a crystal, the case will properly be treated by the method used on page 490. The condition of equilibrium relating to the line

will not in this case be entirely separable from those relating to the adjacent surfaces, since a displacement of the line will involve a displacement of the whole side of the crystal which is terminated at this line. But the expression for the total increment of energy in the system due to any internal changes not involving any variation in the total entropy or volume will consist of two parts, of which one relates to the properties of the masses of the system, and the other may be expressed in the form

$$\delta \Sigma(\sigma s),$$

the summation relating to all the surfaces of discontinuity. This indicates the same tendency toward changes diminishing the value of $\Sigma(\sigma s)$, which appears in other cases.*

General Relations.—For any constant state of strain of the surface of the solid, we may write

$$d\varepsilon_{s(1)} = t d\eta_{s(1)} + \mu_2 dI_{2(1)} + \mu_3 dI_{3(1)} + \text{etc.}, \quad (674)$$

since this relation is implied in the definition of the quantities involved. From this and (659) we obtain

$$d\sigma = -\eta_{s(1)} dt - I_{2(1)} d\mu_2 - I_{3(1)} d\mu_3 - \text{etc.}, \quad (675)$$

which is subject, in strictness, to the same limitation—that the state of strain of the surface of the solid remains the same. But this limitation may in most cases be neglected. (If the quantity σ represented the true tension of the surface, as in the case of a surface between fluids, the limitation would be wholly unnecessary.)

Another method and notation.—We have so far supposed that we have to do with a non-homogeneous film of matter between two homogeneous (or very nearly homogeneous) masses, and that the nature and state of this film is in all respects determined by the

* The freezing together of wool and ice may be mentioned here. The fact that a fiber of wool which remains in contact with a block of ice under water will become attached to it seems to be strictly analogous to the fact that if a solid body be brought into such a position that it just touches the free surface of water, the water will generally rise up about the point of contact so as to touch the solid over a surface of some extent. The condition of the latter phenomenon is

$$\sigma_{sA} + \sigma_{wA} > \sigma_{sw},$$

where the suffixes s , A , and w refer to the solid, to air, and to water, respectively. In like manner, the condition for the freezing of the ice to the wool, if we neglect the æolotropic properties of the ice, is

$$\sigma_{sw} + \sigma_{1w} > \sigma_{s1},$$

where the suffixes s , w , and 1 relate to wool, to water, and to ice, respectively. See *Proc. Roy. Soc.*, vol. x, p. 447; or *Phil. Mag.*, 4th ser., vol. xxi, p. 151.

nature and state of these masses together with the quantities of the foreign substances which may be present in the film. (See page 483.) Problems relating to processes of solidification and dissolution seem hardly capable of a satisfactory solution, except on this supposition, which appears in general allowable with respect to the surfaces produced by these processes. But in considering the equilibrium of fluids at the surface of an unchangeable solid, such a limitation is neither necessary nor convenient. The following method of treating the subject will be found more simple and at the same time more general.

Let us suppose the superficial density of energy to be determined by the excess of energy in the vicinity of the surface over that which would belong to the solid, if (with the same temperature and state of strain) it were bounded by a vacuum in place of the fluid, and to the fluid, if it extended with a uniform volume-density of energy just up to the surface of the solid, or, if in any case this does not sufficiently define a surface, to a surface determined in some definite way by the exterior particles of the solid. Let us use the symbol (ϵ_s) to denote the superficial energy *thus defined*. Let us suppose a superficial density of entropy to be determined in a manner entirely analogous, and be denoted by (η_s). In like manner also, for all the components of the fluid, and for all foreign fluid substances which may be present at the surface, let the superficial densities be determined, and denoted by (Γ_2), (Γ_3), etc. These *superficial densities of the fluid components* relate solely to the matter which is fluid or movable. All matter which is immovably attached to the solid mass is to be regarded as a part of the same. Moreover, let ς be defined by the equation

$$\varsigma = (\epsilon_s) - t(\eta_s) - \mu_2(\Gamma_2) - \mu_3(\Gamma_3) - \text{etc.} \quad (676)$$

These quantities will satisfy the following general relations—

$$d(\epsilon_s) = t d(\eta_s) + \mu_2 d(\Gamma_2) + \mu_3 d(\Gamma_3) + \text{etc.}, \quad (677)$$

$$d\varsigma = -(\eta_s) dt - (\Gamma_2) d\mu_2 - (\Gamma_3) d\mu_3 - \text{etc.} \quad (678)$$

In strictness, these relations are subject to the same limitation as (674) and (675). But this limitation may generally be neglected. In fact, the values of ς , (ϵ_s), etc. must in general be much less affected by variations in the state of strain of the surface of the solid than those of σ , $\epsilon_{s(1)}$, etc.

The quantity ς evidently represents the tendency to contraction in that portion of the surface of the fluid which is in contact with the solid. It may be called *the superficial tension of the fluid in contact with the solid*. Its value may be either positive or negative.

It will be observed that for the same solid surface and for the same temperature but for different fluids the values of σ (in all cases to which the definition of this quantity is applicable) will differ from those of ς by a constant, viz., the value of σ for the solid surface in a vacuum.

For the condition of equilibrium of two different fluids at a line on the surface of the solid, we may easily obtain

$$\sigma_{AB} \cos \alpha = \varsigma_{BS} - \varsigma_{AS}, \tag{679}$$

the suffixes, etc., being used as in (672), and the condition being subject to the same modification when the fluids meet at an edge of the solid.

It must also be regarded as a condition of theoretical equilibrium at the line considered, [subject, like (679), to limitation on account of passive resistances to motion,] that if there are any foreign substances in the surfaces A-S and B-S, the potentials for these substances shall have the same value on both sides of the line; or, if any such substance is found only on one side of the line, that the potential for that substance must not have a less value on the other side; and that the potentials for the components of the mass A, for example, must have the same values in the surface B-C as in the mass A, or, if they are not actual components of the surface B-C, a value not less than in A. Hence, we cannot determine the difference of the surface-tensions of two fluids in contact with the same solid, by bringing them together upon the surface of the solid, unless these conditions are satisfied, as well as those which are necessary to prevent the mixing of the fluid masses.

The investigation on pages 442-448 of the conditions of equilibrium for a fluid system under the influence of gravity may easily be extended to the case in which the system is bounded by or includes solid masses, when these can be treated as rigid and incapable of dissolution. The general condition of mechanical equilibrium would be of the form

$$-f p \delta Dv + f' g \gamma \delta z Dv + f \sigma \delta Ds + f' g \Gamma \delta z Ds + f' g \delta z Dm + f' \varsigma \delta Ds + f' g (\Gamma) \delta z Ds = 0, \tag{680}$$

where the first four integrals relate to the fluid masses and the surfaces which divide them, and have the same signification as in equation (606), the fifth integral relates to the movable solid masses, and the sixth and seventh to the surfaces between the solids and fluids, (Γ) denoting the sum of the quantities (Γ_2), (Γ_3), etc. It should be observed that at the surface where a fluid meets a solid

δz and δz , which indicate respectively the displacements of the solid and the fluid, may have different values, but the components of these displacements which are normal to the surface must be equal.

From this equation, among other particular conditions of equilibrium, we may derive the following—

$$d\zeta = g(L') dz, \quad (681)$$

[compare (614),] which expresses the law governing the distribution of a thin fluid film on the surface of a solid, when there are no passive resistances to its motion.

By applying equation (680) to the case of a vertical cylindrical tube containing two different fluids, we may easily obtain the well-known theorem that the product of the perimeter of the internal surface by the difference $\zeta' - \zeta''$ of the superficial tensions of the upper and lower fluids in contact with the tube is equal to the excess of weight of the matter in the tube above that which would be there, if the boundary between the fluids were in the horizontal plane at which their pressures would be equal. In this theorem, we may either include or exclude the weight of a film of fluid matter adhering to the tube. The proposition is usually applied to the column of fluid *in mass* between the horizontal plane for which $p' = p''$ and the actual boundary between the two fluids. The superficial tensions ζ' and ζ'' are then to be measured in the vicinity of this column. But we may also include the weight of a film adhering to the internal surface of the tube. For example, in the case of water in equilibrium with its own vapor in a tube, the weight of all the water-substance in the tube above the plane $p' = p''$, diminished by that of the water-vapor which would fill the same space, is equal to the perimeter multiplied by the difference in the values of ζ at the top of the tube and at the plane $p' = p''$. If the height of the tube is infinite, the value of ζ at the top vanishes, and the weight of the film of water adhering to the tube and of the mass of liquid water above the plane $p' = p''$ diminished by the weight of vapor which would fill the same space is equal in numerical value but of opposite sign to the product of the perimeter of the internal surface of the tube multiplied by ζ'' , the superficial tension of liquid water in contact with the tube at the pressure at which the water and its vapor would be in equilibrium at a plane surface. In this sense, the total weight of water which can be supported by the tube per unit of the perimeter of its surface is directly measured by the value of $-\zeta$ for water in contact with the tube.

MODIFICATION OF THE CONDITIONS OF EQUILIBRIUM BY ELECTRO-
MOTIVE FORCE.—THEORY OF A PERFECT ELECTRO-CHEMICAL
APPARATUS.

We know by experience that in certain fluids (electrolytic conductors) there is a connection between the fluxes of the component substances and that of electricity. The quantitative relation between these fluxes may be expressed by an equation of the form

$$De = \frac{Dm_a}{\alpha_a} + \frac{Dm_b}{\alpha_b} + \text{etc.} - \frac{Dm_g}{\alpha_g} - \frac{Dm_h}{\alpha_h} - \text{etc.}, \quad (682)$$

where De , Dm_a , etc. denote the infinitesimal quantities of electricity and of the components of the fluid which pass simultaneously through any same surface, which may be either at rest or in motion, and α_a , α_b , etc., α_g , α_h , etc. denote positive constants. We may evidently regard Dm_a , Dm_b , etc., Dm_g , Dm_h , etc., as independent of one another. For, if they were not so, one or more could be expressed in terms of the others, and we could reduce the equation to a shorter form in which all the terms of this kind would be independent.

Since the motion of the fluid as a whole will not involve any electrical current, the densities of the components specified by the suffixes must satisfy the relation

$$\frac{\gamma_a}{\alpha_a} + \frac{\gamma_b}{\alpha_b} + \text{etc.} = \frac{\gamma_g}{\alpha_g} + \frac{\gamma_h}{\alpha_h} + \text{etc.} \quad (683)$$

These densities, therefore, are not independently variable, like the densities of the components which we have employed in other cases.

We may account for the relation (682) by supposing that electricity (positive or negative) is inseparably attached to the different kinds of molecules, so long as they remain in the interior of the fluid, in such a way that the quantities α_a , α_b , etc. of the substances specified are each charged with a unit of positive electricity, and the quantities α_g , α_h , etc. of the substances specified by these suffixes are each charged with a unit of negative electricity. The relation (683) is accounted for by the fact that the constants α_a , α_g , etc. are so small that the electrical charge of any sensible portion of the fluid varying sensibly from the law expressed in (683) would be enormously great, so that the formation of such a mass would be resisted by a very great force.

It will be observed that the choice of the substances which we regard as the components of the fluid is to some extent arbitrary, and that the same physical relations may be expressed by different equa-

tions of the form (682), in which the fluxes are expressed with reference to different sets of components. If the components chosen are such as represent what we believe to be the actual molecular constitution of the fluid, those of which the fluxes appear in the equation of the form (682) are called the *ions*, and the constants of the equation are called their *electro-chemical equivalents*. For our present purpose, which has nothing to do with any theories of molecular constitution, we may choose such a set of components as may be convenient, and call those *ions*, of which the fluxes appear in the equation of the form (682), without farther limitation.

Now, since the fluxes of the independently variable components of an electrolytic fluid do not necessitate any electrical currents, all the conditions of equilibrium which relate to the movements of these components will be the same as if the fluid were incapable of the electrolytic process. Therefore all the conditions of equilibrium which we have found without reference to electrical considerations, will apply to an electrolytic fluid and its independently variable components. But we have still to seek the remaining conditions of equilibrium, which relate to the possibility of electrolytic conduction.

For simplicity, we shall suppose that the fluid is without internal surfaces of discontinuity (and therefore homogeneous except so far as it may be slightly affected by gravity), and that it meets metallic conductors (*electrodes*) in different parts of its surface, being otherwise bounded by non-conductors. The only electrical currents which it is necessary to consider are those which enter the electrolyte at one electrode and leave it at another.

If all the conditions of equilibrium are fulfilled in a given state of the system, except those which relate to changes involving a flux of electricity, and we imagine the state of the system to be varied by the passage from one electrode to another of the quantity of electricity δe accompanied by the quantity δm_a of the component specified, without any flux of the other components or any variation in the total entropy, the total variation of energy in the system will be represented by the expression

$$(V'' - V') \delta e + (\mu_a'' - \mu_a') \delta m_a + (F' - F'') \delta m_a,$$

in which V' , V'' denote the electrical potentials in pieces of the same kind of metal connected with the two electrodes, F' , F'' , the gravitational potentials at the two electrodes, and μ_a' , μ_a'' , the intrinsic potentials for the substance specified. The first term represents the increment of the potential energy of electricity, the second the incre-

ment of the intrinsic energy of the ponderable matter, and the third the increment of the energy due to gravitation.* But by (682)

$$\delta m_a = \alpha_a \delta e$$

It is therefore necessary for equilibrium that

$$V'' - V' + \alpha_a (\mu_a'' - \mu_a' - F'' + F') = 0. \quad (684)$$

To extend this relation to all the electrodes we may write

$$\begin{aligned} V' + \alpha_a (\mu_a' - F') &= V'' + \alpha_a (\mu_a'' - F'') \\ &= V''' + \alpha_a (\mu_a''' - F''') = \text{etc.} \end{aligned} \quad (685)$$

For each of the other cations (specified by ν etc.) there will be a similar condition, and for each of the anions a condition of the form

$$\begin{aligned} V' - \alpha_g (\mu_g' - F') &= V'' - \alpha_g (\mu_g'' - F'') \\ &= V''' - \alpha_g (\mu_g''' - F''') = \text{etc.} \end{aligned} \quad (686)$$

When the effect of gravity may be neglected, and there are but two electrodes, as in a galvanic or electrolytic cell, we have for any cation

$$V'' - V' = \alpha_a (\mu_a' - \mu_a''), \quad (687)$$

and for any anion

$$V'' - V' = \alpha_g (\mu_g'' - \mu_g'), \quad (688)$$

where $V'' - V'$ denotes the electromotive force of the combination. That is:—

When all the conditions of equilibrium are fulfilled in a galvanic or electrolytic cell, the electromotive force is equal to the difference in the values of the potential for any ion or apparent ion at the surfaces of the electrodes multiplied by the electro-chemical equivalent of that ion, the greater potential of an anion being at the same electrode as the greater electrical potential, and the reverse being true of a cation.

Let us apply this principle to different cases.

(I.) If the ion is an independently variable component of an electrode, or by itself constitutes an electrode, the potential for the ion (in any case of equilibrium which does not depend upon passive resistances to change) will have the same value within the electrode as on its surface, and will be determined by the composition of the electrode with its temperature and pressure. This might be illustrated by a cell with electrodes of mercury containing certain quantities of zinc in solution (or with one such electrode and the other of pure

* It is here supposed that the gravitational potential may be regarded as constant for each electrode. When this is not the case, the expression may be applied to small parts of the electrodes taken separately.

zinc) and an electrolytic fluid containing a salt of zinc, but not capable of dissolving the mercury.* We may regard a cell in which hydrogen acts as an ion between electrodes of palladium charged with hydrogen as another illustration of the same principle, but the solidity of the electrodes and the consequent resistance to the diffusion of the hydrogen within them (a process which cannot be assisted by convective currents as in a liquid mass) present considerable obstacles to the experimental verification of the relation.

(II.) Sometimes the ion is soluble (as an independently variable component) in the electrolytic fluid. Of course its condition in the fluid when thus dissolved must be entirely different from its condition when acting on an ion, in which case its quantity is not independently variable, as we have already seen. Its diffusion in the fluid in this state of solution is not necessarily connected with any electrical current, and in other relations its properties may be entirely changed. In any discussion of the internal properties of the fluid (with respect to its fundamental equation, for example,) it would be necessary to treat it as a different substance. (See page 117.) But if the process by which the charge of electricity passes into the electrode, and the ion is dissolved in the electrolyte is *reversible*, we may evidently regard the potentials for the substance of the ion in (687) or (688) as relating to the substance thus dissolved in the electrolyte. In case of absolute equilibrium, the density of the substance thus dissolved would of course be uniform throughout the fluid, (since it can move independently of any electrical current,) so that by the strict application of our principle we only obtain the somewhat barren result, that if any of the ions are soluble in the fluid without their electrical charges, the electromotive force must vanish in any case of absolute equilibrium not dependent upon passive resistances. Nevertheless, cases in which the ion is thus dissolved in the electrolytic fluid only to a very small extent, and its passage from one electrode to the other by ordinary diffusion is extremely slow, may be regarded as approximating to the case in which it is incapable of diffusion. In such cases, we may regard the relations (687), (688) as approximately valid, although the condition of equilibrium

* If the electrolytic fluid dissolved the mercury as well as the zinc, equilibrium could only subsist when the electromotive force is zero, and the composition of the electrodes identical. For when the electrodes are formed of the two metals in different proportions, that which has the greater potential for zinc will have the less potential for mercury. [See equation (98).] This is inconsistent with equilibrium, according to the principle mentioned above, if both metals can act as cations.

relating to the diffusion of the dissolved ion is not satisfied. This may be the case with hydrogen and oxygen as ions (or apparent ions) between electrodes of platinum in some of its forms.

(III.) The ion may appear in mass at the electrode. If it be a conductor of electricity, it may be regarded as forming an electrode, as soon as the deposit has become thick enough to have the properties of matter in mass. The case therefore will not be different from that first considered. When the ion is a non-conductor, a continuous thick deposit on the electrode would of course prevent the possibility of an electrical current. But the case in which the ion being a non-conductor is disengaged in masses contiguous to the electrode but not entirely covering it, is an important one. It may be illustrated by hydrogen appearing in bubbles at a cathode. In case of perfect equilibrium, independent of passive resistances, the potential of the ion in (687) or (688) may be determined in such a mass. Yet the circumstances are quite unfavorable for the establishment of perfect equilibrium, unless the ion is to some extent absorbed by the electrode or electrolytic fluid, or the electrode is fluid. For if the ion must pass *immediately* into the non-conducting mass, while the electricity passes into the electrode, it is evident that the only possible terminus of an electrolytic current is at the line where the electrode, the non-conducting mass, and the electrolytic fluid meet, so that the electrolytic process is necessarily greatly retarded, and an approximate ceasing of the current cannot be regarded as evidence that a state of approximate equilibrium has been reached. But even a slight degree of solubility of the ion in the electrolytic fluid or in the electrode may greatly diminish the resistance to the electrolytic process, and help toward producing that state of complete equilibrium which is supposed in the theorem we are discussing. And the mobility of the surface of a liquid electrode may act in the same way. When the ion is absorbed by the electrode, or by the electrolytic fluid, the case of course comes under the heads which we have already considered, yet the fact that the ion is set free in mass is important, since it is in such a mass that the determination of the value of the potential will generally be most easily made.

(IV.) When the ion is not absorbed either (IV) by the electrode or by the electrolytic fluid, and is not set free in mass, it may still be deposited on the surface of the electrode. Although this can take place only to a limited extent (without forming a body having the properties of matter in mass), yet the electro-chemical equivalents of all substances are so small that a very considerable flux of electricity

may take place before the deposit will have the properties of matter in mass. Even when the ion appears in mass, or is absorbed by the electrode or electrolytic fluid, the non-homogeneous film between the electrolytic fluid and the electrode may contain an additional portion of it. Whether the ion is confined to the surface of the electrode or not, we may regard this as one of the cases in which we have to recognize a certain superficial density of substances at surfaces of discontinuity, the general theory of which we have already considered.

The deposit of the ion will affect the superficial tension of the electrode if it is liquid, or the closely related quantity which we have denoted by the same symbol σ (see pages 482–500) if the electrode is solid. The effect can of course be best observed in the case of a liquid electrode. But whether the electrodes are liquid or solid, if the external electromotive force $V' - V''$ applied to an electrolytic combination is varied, when it is too weak to produce a lasting current, and the electrodes are thereby brought into a new state of polarization, in which they make equilibrium with the altered value of the electromotive force, without change in the nature of the electrodes or of the electrolytic fluid, then by (508) or (675)

$$\begin{aligned} d\sigma' &= -\Gamma'_a d\mu'_a, \\ d\sigma'' &= -\Gamma''_a d\mu''_a; \end{aligned}$$

and by (687),

$$d(V' - V'') = -\alpha_a (d\mu'_a - d\mu''_a).$$

Hence

$$d(V' - V'') = \frac{\alpha_a}{\Gamma'_a} d\sigma' - \frac{\alpha_a}{\Gamma''_a} d\sigma''. \quad (689)$$

If we suppose that the state of polarization of only one of the electrodes is affected (as will be the case when its surface is very small compared with that of the other), we have

$$d\sigma' = \frac{\Gamma'_a}{\alpha_a} d(V' - V''). \quad (690)$$

The superficial tension of one of the electrodes is then a function of the electromotive force.

This principle has been applied by M. Lippmann to the construction of the electrometer which bears his name.* In applying equations (689) and (690) to dilute sulphuric acid between electrodes of mercury, as in a Lippmann's electrometer, we may suppose that the

* See his memoir: "Relations entre les phénomènes électriques et capillaires," *Annales de Chimie et de Physique*, 5^e série, t. v, p. 494.

suffix refers to hydrogen. It will be most convenient to suppose the *dividing surface* to be so placed as to make the surface-density of mercury zero. (See page 397.) The matter which exists in excess or deficiency at the surface may then be expressed by the surface-densities of sulphuric acid, of water, and of hydrogen. The value of the last may be determined from equation (690). According to M. Lippmann's determinations, it is negative when the surface is in its natural state (i. e., the state to which it tends when no external electromotive force is applied), since σ' increases with $V'' - V'$. When $V'' - V'$ is equal to nine-tenths of the electromotive force of a Daniell's cell, the electrode to which V'' relates remaining in its natural state, the tension σ' of the surface of the other electrode has a maximum value, and there is no excess or deficiency of hydrogen at that surface. This is the condition toward which a surface tends when it is extended while no flux of electricity takes place. The flux of electricity per unit of new surface formed, which will maintain a surface in a constant condition while it is extended, is represented by $\frac{I'_a}{\alpha_a}$ in numerical value, and its direction, when I''_a is negative, is from the mercury into the acid.

We have so far supposed, in the main, that there are no passive resistances to change, except such as vanish with the rapidity of the processes which they resist. The actual condition of things with respect to passive resistances appears to be nearly as follows. There does not appear to be any passive resistance to the electrolytic process by which an ion is transferred from one electrode to another, except such as vanishes with the rapidity of the process. For, in any case of equilibrium, the smallest variation of the externally applied electromotive force appears to be sufficient to cause a (temporary) electrolytic current. But the case is not the same with respect to the molecular changes by which the ion passes into new combinations or relations, as when it enters into the mass of the electrodes, or separates itself in mass, or is dissolved (no longer with the properties of an ion) in the electrolytic fluid. In virtue of the passive resistance to these processes, the external electromotive force may often vary within wide limits, without creating any current by which the ion is transferred from one of the masses considered to the other. In other words, the value of $V' - V''$ may often differ greatly from that obtained from (687) or (688) when we determine the values of the potentials for the ion as in cases I, II, and III. We may, however, regard these equations as entirely valid, when the potentials for the

ions are determined at the surface of the electrodes with reference to the ion in the condition in which it is brought there or taken away by an electrolytic current, without any attendant irreversible processes. But in a complete discussion of the properties of the surface of an electrode it may be necessary to distinguish (both in respect to surface-densities and to potentials) between the substance of the ion in this condition and the same substance in other conditions into which it cannot pass (directly) without irreversible processes. No such distinction, however, is necessary when the substance of the ion can pass at the surface of the electrode by reversible processes from any one of the conditions in which it appears to any other.

The formulæ (687), (688) afford as many equations as there are ions. These, however, amount to only one independent equation additional to those which relate to the independently variable components of the electrolytic fluid. This appears from the consideration that a flux of any cation may be combined with a flux of any anion in the same direction so as to involve no electrical current, and that this may be regarded as the flux of an independently variable component of the electrolytic fluid.

General Properties of a Perfect Electro-chemical Apparatus.

When an electrical current passes through a galvanic or electrolytic cell, the state of the cell is altered. If no changes take place in the cell except during the passage of the current, and all changes which accompany the current can be reversed by reversing the current, the cell may be called a perfect electro-chemical apparatus. The electromotive force of the cell may be determined by the equations which have just been given. But some of the general relations to which such an apparatus is subject may be conveniently stated in a form in which the ions are not explicitly mentioned.

In the most general case, we may regard the cell as subject to external action of four different kinds. (1) The supply of electricity at one electrode and the withdrawal of the same quantity at the other. (2) The supply or withdrawal of a certain quantity of heat. (3) The action of gravity. (4) The motion of the surfaces enclosing the apparatus, as when its volume is increased by the liberation of gases.

The increase of the energy in the cell is necessarily equal to that which it receives from external sources. We may express this by the equation

$$d\varepsilon = (V' - V'') de + dQ + dW_G + dW_P, \quad (691)$$

in which $d\varepsilon$ denotes the increment of the intrinsic energy of the cell, de the quantity of electricity which passes through it, V' and V'' the electrical potentials in masses of the same kind of metal connected with the anode and cathode respectively, dQ the heat received from external bodies, dW_g the work done by gravity, and dW_p the work done by the pressures which act on the external surface of the apparatus.

The conditions under which we suppose the processes to take place are such that the increase of the entropy of the apparatus is equal to the entropy which it receives from external sources. The only external source of entropy is the heat which is communicated to the cell by the surrounding bodies. If we write $d\eta$ for the increment of entropy in the cell, and t for the temperature, we have

$$d\eta = \frac{dQ}{t}. \tag{692}$$

Eliminating dQ , we obtain

$$d\varepsilon = (V' - V'') de + t d\eta + dW_g + dW_p, \tag{693}$$

or

$$V'' - V' = -\frac{d\varepsilon}{de} + t \frac{d\eta}{de} + \frac{dW_g}{de} + \frac{dW_p}{de}. \tag{694}$$

It is worth while to notice that if we give up the condition of the reversibility of the processes, so that the cell is no longer supposed to be a perfect electro-chemical apparatus, the relation (691) will still subsist. But, if we still suppose, for simplicity, that all parts of the cell have the same temperature, which is *necessarily* the case with a perfect electro-chemical apparatus, we shall have, instead of (692),

$$d\eta \geq \frac{dQ}{t}, \tag{695}$$

and instead of (693), (694)

$$(V'' - V') de \leq -d\varepsilon + t d\eta + dW_g + dW_p. \tag{696}$$

The values of the several terms of the second member of (694), for a given cell, will vary with the external influences to which the cell is subjected. If the cell is enclosed (with the products of electrolysis) in a rigid envelop, the last term will vanish. The term relating to gravity is generally to be neglected. If no heat is supplied or withdrawn, the term containing $d\eta$ will vanish. But in the calculation of the electromotive force, which is the most important application of the equation, it is generally more convenient to suppose that the temperature remains constant.

The quantities expressed by the terms containing dQ and $d\eta$ in (691), (693), (694), and (696) are frequently neglected in the consideration of cells of which the temperature is supposed to remain constant. In other words, it is frequently assumed that neither heat nor cold is produced by the passage of an electrical current through a perfect electro-chemical combination (except that heat which may be indefinitely diminished by increasing the time in which a given quantity of electricity passes), and that only heat can be produced in any cell, unless it be by processes of a secondary nature, which are not immediately or necessarily connected with the process of electrolysis.

It does not appear that this assumption is justified by any sufficient reason. In fact, it is easy to find a case in which the electromotive force is determined entirely by the term $t \frac{d\eta}{de}$ in (694), all the other terms in the second member of the equation vanishing. This is true of a Grove's gas battery charged with hydrogen and nitrogen. In this case, the hydrogen passes over to the nitrogen,—a process which does not alter the energy of the cell, when maintained at a constant temperature. The work done by external pressures is evidently nothing, and that done by gravity is (or may be) nothing. Yet an electrical current is produced. The work done (or which may be done) by the current outside of the cell is the equivalent of the work (or of a part of the work) which might be gained by allowing the gases to mix in other ways. This is equal, as has been shown by Lord Rayleigh,* to the work which may be gained by allowing each gas separately to expand at constant temperature from its initial volume to the volume occupied by the two gases together. The same work is equal, as appears from equations (278), (279) on page 217, (see also page 220,) to the increase of the entropy of the system multiplied by the temperature.

It is possible to vary the construction of the cell in such a way that nitrogen or other neutral gas will not be necessary. Let the cell consist of a U-shaped tube of sufficient height, and have pure hydrogen at each pole under very unequal pressures (as of one and two atmospheres respectively) which are maintained constant by properly weighted pistons, sliding in the arms of the tube. The difference of the pressures in the gas-masses at the two electrodes must of course be balanced by the difference in the height of the two columns of acidulated water. It will hardly be doubted that such an apparatus

* *Philosophical Magazine*, vol. xlix, p. 311.

would have an electromotive force acting in the direction of a current which would carry the hydrogen from the denser to the rarer mass. Certainly the gas could not be carried in the opposite direction by an external electromotive force without the expenditure of as much (electromotive) work as is equal to the mechanical work necessary to pump the gas from the one arm of the tube to the other. And if by any modification of the metallic electrodes (which remain unchanged by the passage of electricity) we could reduce the passive resistances to zero, so that the hydrogen could be carried reversibly from one mass to the other without finite variation of the electromotive force, the only possible value of the electromotive force would be represented by the expression $t \frac{d\eta}{de}$, as a very close approximation. It will be observed that, although gravity plays an essential part in a cell of this kind by maintaining the difference of pressure in the masses of hydrogen, the electromotive force cannot possibly be ascribed to gravity, since the work done by gravity, when hydrogen passes from the denser to the rarer mass, is negative.

Again, it is entirely improbable that the electrical currents caused by differences in the concentration of solutions of salts, (as in a cell containing sulphate of zinc between zinc electrodes, or sulphate of copper between copper electrodes, the solution of the salt being of unequal strength at the two electrodes,) which have recently been investigated theoretically and experimentally by MM. Helmholtz and Moser,* are confined to cases in which the mixture of solutions of different degrees of concentration will produce heat. Yet in cases in which the mixture of more and less concentrated solutions is not attended with evolution or absorption of heat, the electromotive force must vanish in a cell of the kind considered, if it is determined simply by the diminution of energy in the cell. And when the mixture produces cold, the same rule would make any electromotive force impossible except in the direction which would tend to increase the difference of concentration. Such conclusions as would be quite irreconcilable with the theory of the phenomena given by Professor Helmholtz.

A more striking example of the necessity of taking account of the variations of entropy in the cell in *a priori* determinations of electromotive force is afforded by electrodes of zinc and mercury in a solution of sulphate of zinc. Since heat is absorbed when zinc is dissolved

* *Annalen der Physik und Chemie*, Neue Folge, Band iii, February, 1878.

in mercury,* the energy of the cell is increased by a transfer of zinc to the mercury, when the temperature is maintained constant. Yet in this combination, the electromotive force acts in the direction of the current producing such a transfer.† The couple presents certain anomalies when a considerable quantity of zinc is united with the mercury. The electromotive force changes its direction, so that this case is usually cited as an illustration of the principle that the electromotive force is in the direction of the current which diminishes the energy of the cell, *i. e.*, which produces or allows those changes which are accompanied by evolution of heat when they take place directly. But whatever may be the cause of the electromotive force which has been observed acting in the direction from the amalgam through the electrolyte to the zinc (a force which according to the determinations of M. Gaugain is only one twenty-fifth part of that which acts in the reverse direction when pure mercury takes the place of the amalgam), these anomalies can hardly affect the general conclusions with which alone we are here concerned. If the electrodes of a cell are pure zinc and an amalgam containing zinc not in excess of the amount which the mercury will dissolve at the temperature of the experiment without losing its fluidity, and if the only change (other than thermal) accompanying a current is a transfer of zinc from one electrode to the other,—conditions which may not have been satisfied in all the experiments recorded, but which it is allowable to suppose in a theoretical discussion, and which certainly will not be regarded as inconsistent with the fact that heat is absorbed when zinc is dissolved in mercury,—it is impossible that the electromotive force should be in the direction of a current transferring zinc from the amalgam to the electrode of pure zinc. For, since the zinc eliminated from the amalgam by the electrolytic process might be re-dissolved directly, such a direction of the electromotive force would involve the possibility of obtaining an indefinite amount of electromotive work, and therefore of mechanical work, without other expenditure than that of heat at the constant temperature of the cell.

None of the cases which we have been considering involve combinations by definite proportions, and, except in the case of the cell with electrodes of mercury and zinc, the electromotive forces are quite small. It may perhaps be thought that with respect to those cells in which combinations take place by definite proportions the electromotive force may be calculated with substantial accuracy from

* J. Regnauld, *Comptes Rendus*, t. li, p. 778.

† Gaugain, *Comptes Rendus*, t. xlii, p. 430.

the diminution of the energy, without regarding the variation of entropy. But the phenomena of chemical combination do not in general seem to indicate any possibility of obtaining from the combination of substances by any process whatever an amount of mechanical work which is equivalent to the heat produced by the direct union of the substances.

A kilogramme of hydrogen, for example, combining by combustion under the pressure of the atmosphere with eight kilogrammes of oxygen to form liquid water, yields an amount of heat which may be represented in round numbers by 34000 calories.* We may suppose that the gases are taken at the temperature of 0° C., and that the water is reduced to the same temperature. *But this heat cannot be obtained at any temperature desired.* A very high temperature has the effect of preventing to a greater or less extent, the combination of the elements. Thus, according to M. Sainte-Claire Deville,† the temperature obtained by the combustion of hydrogen and oxygen cannot much if at all exceed 2500° C., which implies that less than one-half of the hydrogen and oxygen present combine at that temperature. This relates to combustion under the pressure of the atmosphere. According to the determinations of Professor Bunsen‡ in regard to combustion in a confined space, only one-third of a mixture of hydrogen and oxygen will form a chemical compound at the temperature of 2850° C. and a pressure of ten atmospheres, and only a little more than one-half when the temperature is reduced by the addition of nitrogen to 2024° C., and the pressure to about three atmospheres *exclusive* of the part due to the nitrogen.

Now 10 calories at 2500° C. are to be regarded as reversibly convertible into one calorie at 4° C. together with the mechanical work representing the energy of 9 calories. If, therefore, all the 34000 calories obtainable from the union of hydrogen and oxygen under atmospheric pressure could be obtained at the temperature of 2500° C., and no higher, we should estimate the electromotive work performed in a perfect electro-chemical apparatus in which these elements are combined or separated at ordinary temperatures and under atmospheric pressure as representing nine-tenths of the 34000 calories, and the heat evolved or absorbed in the apparatus as representing one-tenth of the 34000 calories.§ This, of course, would give an electromotive

* See Rühlmann's *Handbuch der mechanischen Wärmetheorie*, Bd. ii, p. 290.

† *Comptes Rendus*, t. lvi, p. 199; and t. lxiv, 67.

‡ *Pogg. Ann.*, Bd. cxxxi (1867), p. 161.

§ These numbers are not subject to correction for the pressure of the atmosphere, since the 34000 calories relate to combustion under the same pressure.

force exactly nine-tenths as great as is obtained on the supposition that all the 34000 calories are convertible into electromotive or mechanical work. But, according to all indications, the estimate 2500° C. (for the temperature at which we may regard all the heat of combustion as obtainable) is far too high,* and we must regard the theoretical value of the electromotive force necessary to electrolyze water as considerably less than nine-tenths of the value obtained on the supposition that it is necessary for the electromotive agent to supply all the energy necessary for the process.

The case is essentially the same with respect to the electrolysis of hydrochloric acid, which is probably a more typical example of the process than the electrolysis of water. The phenomenon of dissociation is equally marked, and occurs at a much lower temperature, more than half of the gas being dissociated at 1400° C.† And the heat which is obtained by the combination of hydrochloric acid gas with water, especially with water which already contains a considerable quantity of the acid, is probably only to be obtained at temperatures comparatively low. This indicates that the theoretical value of the electromotive force necessary to electrolyze this acid (*i. e.*, the electromotive force which would be necessary in a reversible electrochemical apparatus), must be very much less than that which could perform in electromotive work the equivalent of all the heat evolved in the combination of hydrogen, chlorine and water to form the liquid submitted to electrolysis. This presumption, based upon the phenomena exhibited in the direct combination of the substances, is corroborated by the experiments of M. Favre, who has observed an absorption of heat in the cell in which this acid was electrolyzed.‡ The

* Unless the received ideas concerning the behavior of gases at high temperatures are quite erroneous, it is possible to indicate the general character of a process (involving at most only such difficulties as are neglected in theoretical discussions) by which water may be converted into separate masses of hydrogen and oxygen without other expenditure than that of an amount of heat equal to the difference of energy of the matter in the two states and supplied at a temperature far below 2500° C. The essential parts of the process would be (1) vaporizing the water and heating it to a temperature at which a considerable part will be dissociated, (2) the partial separation of the hydrogen and oxygen by filtration, and (3) the cooling of both gaseous masses until the vapor they contain is condensed. A little calculation will show that in a continuous process all the heat obtained in the operation of cooling the products of filtration could be utilized in heating fresh water.

† Sainte-Claire Deville, *Comptes Rendus*, t. lxiv, p. 67.

‡ See *Mémoires des Savants Étrangers*, Sér. 2, t. xxv, No. 1, p. 142; or *Comptes Rendus*, t. lxxiii, p. 973. The figures obtained by M. Favre will be given hereafter, in connection with others of the same nature.

electromotive work expended must therefore have been less than the increase of energy in the cell.

In both cases of composition in definite proportions which we have considered, the compound has more entropy than its elements, and the difference is by no means inconsiderable. This appears to be the rule rather than the exception with respect to compounds which have less energy than their elements. Yet it would be rash to assert that it is an invariable rule. And when one substance is substituted for another in a compound, we may expect great diversity in the relations of energy and entropy.

In some cases, there is a striking correspondence between the electromotive force of a cell and the rate of diminution of its energy per unit of electricity transmitted, the temperature remaining constant. A Daniell's cell is a notable example of this correspondence. It may perhaps be regarded as a very significant case, since of all cells in common use, it has the most constant electromotive force, and most nearly approaches the condition of reversibility. If we apply our previous notation [compare (691)] with the substitution of finite for infinitesimal differences to the determinations of M. Favre,* estimating energy in calories, we have for each equivalent (32.6 kilogrammes) of zinc dissolved

$$(V'' - V') \Delta e = 24327^{\text{cal.}}, \quad \Delta \varepsilon = -25394^{\text{cal.}}, \quad \Delta Q = -1067^{\text{cal.}}$$

It will be observed that the electromotive work performed by the cell is about four per cent. less than the diminution of energy in the cell.† The value of ΔQ , which, when negative, represents the heat evolved in the cell when the external resistance of the circuit is very great, was determined by direct measurement, and does not appear to have been corrected for the resistance of the cell. This correction would diminish the value of $-\Delta Q$, and increase that of $(V'' - V') \Delta e$, which was obtained by subtracting $-\Delta Q$ from $-\Delta \varepsilon$.

It appears that under certain conditions neither heat nor cold is produced in a Grove's cell. For M. Favre has found that with different degrees of concentration of the nitric acid sometimes heat and sometimes cold is produced.‡ When neither is produced, of course

* See *Mém. Savants Étrang.*, loc. cit., p. 90; or *Comptes Rendus*, vol. lxix, p. 35, where the numbers are slightly different.

† A comparison of the experiments of different physicists has in some cases given a much closer correspondence. See Wiedemann's *Galvanismus*, etc., 2^{te} Auflage, Bd. ii, §§ 1117, 1118.

‡ *Mém. Savants Étrang.*, loc. cit., p. 93; or *Comptes Rendus*, t. lxix, p. 37, and t. lxxiii, p. 893.

the electromotive force of the cell is exactly equal to its diminution of energy per unit of electricity transmitted. But such a coincidence is far less significant than the fact that an absorption of heat has been observed. With acid containing about seven equivalents of water ($\text{HNO}_6 + 7\text{HO}$), M. Favre has found

$$(V'' - V') \Delta e = 46781 \text{ cal.}, \quad \Delta \varepsilon = -41824 \text{ cal.}, \quad \Delta Q = 4957 \text{ cal.};$$

and with acid containing about one equivalent of water ($\text{HNO}_6 + \text{HO}$),

$$(V'' - V') \Delta e = 49847 \text{ cal.}, \quad \Delta \varepsilon = -52714 \text{ cal.}, \quad \Delta Q = -2867 \text{ cal.}$$

In the first example, it will be observed that the quantity of heat absorbed in the cell is not small, and that the electromotive force is nearly one-eighth greater than can be accounted for by the diminution of energy in the cell.

This absorption of heat in the cell he has observed in other cases, in which the chemical processes are much more simple.

For electrodes of cadmium and platinum in hydrochloric acid his experiments give*

$$\begin{aligned} (V'' - V') \Delta e &= 9256 \text{ cal.}, & \Delta \varepsilon &= -8258 \text{ cal.}, \\ \Delta W_P &= -290 \text{ cal.}, & \Delta Q &= 1288 \text{ cal.} \end{aligned}$$

In this case the electromotive force is nearly one-sixth greater than can be accounted for by the diminution of energy in the cell with the work done against the pressure of the atmosphere.

For electrodes of zinc and platinum in the same acid one series of experiments gives†

$$\begin{aligned} (V'' - V') \Delta e &= 16950 \text{ cal.}, & \Delta \varepsilon &= -16189 \text{ cal.}, \\ \Delta W_P &= -290 \text{ cal.}, & \Delta Q &= 1051 \text{ cal.}; \end{aligned}$$

and a later series,‡

$$\begin{aligned} (V'' - V') \Delta e &= 16738 \text{ cal.}, & \Delta \varepsilon &= -17702 \text{ cal.}, \\ \Delta W_P &= -290 \text{ cal.}, & \Delta Q &= -674 \text{ cal.} \end{aligned}$$

In the electrolysis of hydrochloric acid in a cell with a porous partition, he has found§

* *Comptes Rendus*, t. lxxviii, p. 1305. The total heat obtained in the whole circuit (including the cell) when all the electromotive work is turned into heat, was ascertained by direct experiment. This quantity, 7968 calories, is evidently represented by $(V'' - V') \Delta e - \Delta Q$, also by $-\Delta \varepsilon + \Delta W_P$. [See (691).] The value of $(V'' - V') \Delta e$ is obtained by adding ΔQ , and that of $-\Delta \varepsilon$ by adding $-\Delta W_P$, which is easily estimated, being determined by the evolution of one kilogramme of hydrogen.

† *Ibid.*

‡ *Mém. Savants Étrang.*, loc. cit., p. 145.

§ *Ibid.*, p. 142.

$$(V' - V'') \Delta e = 34825^{\text{cal.}} \quad \Delta Q = 2113^{\text{cal.}},$$

whence

$$\Delta \varepsilon - \Delta W_p = 36938.$$

We cannot assign a precise value to ΔW_p , since the quantity of chlorine which was evolved in the form of gas is not stated. But the value of $-\Delta W_p$ must lie between $290^{\text{cal.}}$ and $580^{\text{cal.}}$, probably nearer to the former.

The great difference in the results of the two series of experiments relating to electrodes of zinc and platinum in hydrochloric acid is most naturally explained by supposing some difference in the conditions of the experiment, as in the concentration of the acid, or in the extent to which the substitution of zinc for hydrogen took place.* That which it is important for us to observe in all these cases is that there are conditions under which heat is absorbed in a galvanic or electrolytic cell, so that the galvanic cell has a greater electromotive force than can be accounted for by the diminution of its energy, and the operation of electrolysis requires a less electromotive force than would be calculated from the increase of energy in the cell,—especially when the work done against the pressure of the atmosphere is taken into account.

It should be noticed that in all these experiments the quantity represented by ΔQ (which is the critical quantity with respect to the point at issue) was determined by direct measurement of the heat absorbed or evolved by the cell when placed alone in a calorimeter. The resistance of the circuit was made so great by a rheostat placed outside of the calorimeter that the resistance of the cell was regarded as insignificant in comparison, and no correction appears to have been made in any case for this resistance. With exception of the error due to this circumstance, which would in all cases diminish the heat absorbed in the cell (or increase the heat evolved), the probable error of ΔQ must be very small in comparison with that of $(V' - V'') \Delta e$, or with that of $\Delta \varepsilon$, which were in general determined by the compar-

* It should perhaps be stated that in his extended memoir published in 1877 in the *Mémoires des Savants Étrangers*, in which he has presumably collected those results of his experiments which he regards as most important and most accurate, M. Favre does not mention the absorption of heat in a cell of this kind, or in the similar cell in which cadmium takes the place of zinc. This may be taken to indicate a decided preference for the later experiments which showed an evolution of heat. Whatever the ground of this preference may have been, it can hardly destroy the significance of the absorption of heat, which was a matter of direct observation in repeated experiments. See *Comptes Rendus*, t. lxxviii, p. 1305.

ison of different calorimetric measurements, involving very much greater quantities of heat.

In considering the numbers which have been cited, we should remember that when hydrogen is evolved as gas the process is in general very far from reversible. In a perfect electrochemical apparatus, the same changes in the cell would yield a much greater amount of electromotive work, or absorb a much less amount. In either case, the value of ΔQ would be much greater than in the imperfect apparatus, the difference being measured perhaps by thousands of calories.*

It often occurs in a galvanic or electrolytic cell that an ion which is set free at one of the electrodes appears in part as gas, and is in part absorbed by the electrolytic fluid, and in part absorbed by the electrode. In such cases, a slight variation in the circumstances, which would not sensibly affect the electromotive force, would cause all of the ion to be disposed of in one of the three ways mentioned, if the current were sufficiently weak. This would make a considerable

* Except in the case of the Grove's cell, in which the reactions are quite complicated, the absorption of heat is most marked in the electrolysis of hydrochloric acid. The latter case is interesting, since the experiments confirm the presumption afforded by the behavior of the substances in other circumstances. (See page 514.) In addition to the circumstances mentioned above tending to diminish the observed absorption of heat, the following, which are peculiar to this case, should be noticed.

The electrolysis was performed in a cell with a porous partition, in order to prevent the chlorine and hydrogen dissolved in the liquid from coming in contact with each other. It had appeared in a previous series of experiments (*Mém. Savants Étrang.*, loc. cit., p. 131; or *Comptes Rendus*, t. lxvi, p. 1231,) that a very considerable amount of heat might be produced by the chemical union of the gases in solution. In a cell without partition, instead of an absorption, an evolution of heat took place, which sometimes exceeded 5000 calories. If, therefore, the partition did not perfectly perform its office, this could only cause a diminution in the value of ΔQ .

A large part at least of the chlorine appears to have been absorbed by the electrolytic fluid. It is probable that a slight difference in the circumstances of the experiment—a diminution of pressure, for example,—might have caused the greater part of the chlorine to be evolved as gas, without essentially affecting the electromotive force. The solution of chlorine in water presents some anomalies, and may be attended with complex reactions, but it appears to be always attended with a very considerable evolution of heat. (See Berthelot, *Comptes Rendus*, t. lxxvi, p. 1514.) If we regard the evolution of the chlorine in the form of gas as the normal process, we may suppose that the absorption of heat in the cell was greatly diminished by the retention of the chlorine in solution.

Under certain circumstances, oxygen is evolved in the electrolysis of dilute hydrochloric acid. It does not appear that this took place to any considerable extent in the experiments which we are considering. But so far as it may have occurred, we may regard it as a case of the electrolysis of water. The significance of the fact of the absorption of heat is not thereby affected.

difference in the variation of energy in the cell, and the electromotive force cannot certainly be calculated from the variation of energy alone in all these cases. The correction due to the work performed against the pressure of the atmosphere when the ion is set free as gas will not help us in reconciling these differences. It will appear on consideration that this correction will in general increase the discordance in the values of the electromotive force. Nor does it distinctly appear which of these cases is to be regarded as normal and which are to be rejected as involving secondary processes.*

If in any case secondary processes are excluded, we should expect it to be when the ion is identical in substance with the electrode upon which it is deposited, or from which it passes into the electrolyte. But even in this case we do not escape the difficulty of the different forms in which the substance may appear. If the temperature of the experiment is at the melting point of a metal which forms the ion and the electrode, a slight variation of temperature will cause the ion to be deposited in the solid or in the liquid state, or, if the current is in the opposite direction, to be taken up from a solid or from a liquid body. Since this will make a considerable difference in the variation of energy, we obtain different values for the electromotive force above and below the melting point of the metal, unless we also take account of the variations of entropy. Experiment does not indicate the existence of any such difference,† and when we take account of variations of entropy, as in equation (694), it is apparent

that there ought not to be any, the terms $\frac{d\varepsilon}{de}$ and $t\frac{d\eta}{de}$ being both

* It will be observed that in using the formulae (694) and (696) we do not have to make any distinction between *primary* and *secondary* processes. The only limitation to the generality of these formulae depends upon the *reversibility* of the processes, and this limitation does not apply to (696).

† M. Raoult has experimented with a galvanic element having an electrode of bismuth in contact with phosphoric acid containing phosphate of bismuth in solution. (See *Comptes Rendus*, t. lxxviii, p. 643.) Since this metal absorbs in melting 12.64 calories per kilogramme or 885 calories per equivalent (70^{kil.}), while a Daniell's cell yields about 24000 calories of electromotive work per equivalent of metal, the solid or liquid state of the bismuth ought to make a difference of electromotive force represented by .037 of a Daniell's cell, if the electromotive force depended simply upon the energy of the cell. But in M. Raoult's experiments no sudden change of electromotive force was manifested at the moment when the bismuth changed its state of aggregation. In fact, a change of temperature in the electrode from about fifteen degrees above to about fifteen degrees below the temperature of fusion only occasioned a variation of electromotive force equal to .002 of a Daniell's cell.

Experiments upon lead and tin gave similar results.

affected by the same difference, viz., the heat of fusion of an electrochemical equivalent of the metal. In fact, if such a difference existed, it would be easy to devise arrangements by which the heat yielded by a metal in passing from the liquid to the solid state could be transformed into electromotive work (and therefore into mechanical work) without other expenditure.

The foregoing examples will be sufficient, it is believed, to show the necessity of regarding other considerations in determining the electromotive force of a galvanic or electrolytic cell than the variation of its energy alone (when its temperature is supposed to remain constant), or corrected only for the work which may be done by external pressures or by gravity. But the relations expressed by (693), (694), and (696) may be put in a briefer form.

If we set, as on page 144,

$$\psi = \varepsilon - t \eta,$$

we have, for any constant temperature,

$$d\psi = d\varepsilon - t d\eta;$$

and for any perfect electrochemical apparatus, the temperature of which is maintained constant,

$$V'' - V' = -\frac{d\psi}{de} + \frac{dW_g}{de} + \frac{dW_p}{de}; \quad (697)$$

and for any cell whatever, when the temperature is maintained uniform and constant,

$$(V'' - V') de \leq -d\psi + dW_g + dW_p. \quad (698)$$

In a cell of any ordinary dimensions, the work done by gravity, as well as the inequalities of pressure in different parts of the cell may be neglected. If the pressure as well as the temperature is maintained uniform and constant, and we set, as on page 147,

$$\zeta = \varepsilon - t \eta + p v,$$

where p denotes the pressure in the cell, and v its total volume (including the products of electrolysis), we have

$$d\zeta = d\varepsilon - t d\eta + p dv,$$

and for a perfect electro-chemical apparatus,

$$V'' - V' = -\frac{d\zeta}{de}, \quad (699)$$

or for any cell,

$$(V'' - V') de \leq -d\zeta. \quad (700)$$

SYNOPSIS OF SUBJECTS TREATED.

	Page
PRELIMINARY REMARK on the rôle of energy and entropy in the theory of thermodynamic systems.....	108
CRITERIA OF EQUILIBRIUM AND STABILITY.	
Criteria enunciated.....	109
Meaning of the term <i>possible variations</i> ,.....	110
Passive resistances.....	111
Validity of the criteria.....	112
THE CONDITIONS OF EQUILIBRIUM FOR HETEROGENEOUS MASSES IN CONTACT, WHEN UNINFLUENCED BY GRAVITY, ELECTRICITY, DISTORTION OF THE SOLID MASSES, OR CAPILLARY TENSIONS.	
Statement of the problem.....	115
Conditions relating to equilibrium between the initially existing homogeneous parts of the system.....	116
Meaning of the term <i>homogeneous</i> ,.....	116
Variation of the energy of a homogeneous mass,.....	116
Choice of substances to be regarded as components.—Actual and possible components,.....	116
Deduction of the particular conditions of equilibrium when all parts of the system have the same components,.....	118
Definition of the potentials for the component substances in the various homogeneous masses,.....	119
Case in which certain substances are only possible components in a part of the system,.....	120
Form of the particular conditions of equilibrium when there are relations of convertibility between the substances which are regarded as the components of the different masses,.....	121
Conditions relating to the possible formation of masses unlike any previously existing,.....	124
Very small masses cannot be treated by the same method as those of considerable size,.....	129
Sense in which formula (52) may be regarded as expressing the condition sought,.....	129
Condition (53) is always sufficient for equilibrium, but not always necessary,.....	131
A mass in which this condition is not satisfied, is at least <i>practically</i> unstable,.....	133
(This condition is farther discussed under the head of <i>Stability</i> . See p. 156).	
Effect of solidity of any part of the system.....	134
Effect of additional equations of condition,.....	137
Effect of a diaphragm,—equilibrium of osmotic forces,.....	138
FUNDAMENTAL EQUATIONS.	
Definition and properties,.....	140
Concerning the quantities ψ , χ , ζ ,.....	144
Expression of the criterion of equilibrium by means of the quantity ψ ,.....	145
Expression of the criterion of equilibrium in certain cases by means of the quantity ζ ,.....	147
POTENTIALS.	
The value of a potential for a substance in a given mass is not dependent on the other substances which may be chosen to represent the composition of the mass,.....	148
Potentials defined so as to render this property evident,.....	149

	Page
In the same homogeneous mass we may distinguish the potentials for an indefinite number of substances, each of which has a perfectly determined value. Between the potentials for different substances in the same homogeneous mass the same equations will subsist as between the units of these substances,	149
The values of potentials depend upon the arbitrary constants involved in the definition of the energy and entropy of each elementary substance.....	151
COEXISTENT PHASES.	
Definition of <i>phases</i> —of <i>coexistent phases</i> ,	152
Number of the independent variations which are possible in a system of <i>coexistent phases</i> ,.....	152
Case of $n + 1$ <i>coexistent phases</i> ,.....	153
Cases of a less number of <i>coexistent phases</i> ,.....	155
INTERNAL STABILITY OF HOMOGENEOUS FLUIDS AS INDICATED BY FUNDAMENTAL EQUATIONS.	
General condition of absolute stability,	156
Other forms of the condition,	160
Stability in respect to continuous changes of phase,.....	162
Conditions which characterize the limits of stability in this respect,.....	169
GEOMETRICAL ILLUSTRATIONS.	
Surfaces in which the composition of the body represented is constant,	172
Surfaces and curves in which the composition of the body represented is variable and its temperature and pressure are constant,.....	176
CRITICAL PHASES.	
Definition,	188
Number of independent variations which are possible for a critical phase while remaining such,	188
Analytical expression of the conditions which characterize critical phases.—Situation of critical phases with respect to the limits of stability,.....	189
Variations which are possible under different circumstances in the condition of a mass initially in a critical phase,.....	191
ON THE VALUES OF THE POTENTIALS WHEN THE QUANTITY OF ONE OF THE COMPONENTS IS VERY SMALL.....	
	194
ON CERTAIN POINTS RELATING TO THE MOLECULAR CONSTITUTION OF BODIES.	
Proximate and ultimate components,.....	197
Phases of dissipated energy,.....	200
Catalysis.—perfect catalytic agent,	201
A fundamental equation for phases of dissipated energy may be formed from the more general form of the fundamental equation,.....	201
The phases of dissipated energy may sometimes be the only phases the existence of which can be experimentally verified,.....	201
THE CONDITIONS OF EQUILIBRIUM FOR HETEROGENEOUS MASSES UNDER THE INFLUENCE OF GRAVITY.	
The problem is treated by two different methods:	
The elements of volume are regarded as variable,.....	203
The elements of volume are regarded as fixed,	207
FUNDAMENTAL EQUATIONS OF IDEAL GASES AND GAS-MIXTURES.	
Ideal gas,.....	210
Ideal gas-mixture—Dalton's Law,	215
Inferences in regard to potentials in liquids and solids,	225
Considerations relating to the increase of entropy due to the mixture of gases by diffusion,	227
The phases of dissipated energy of an ideal gas-mixture with components which are chemically related,.....	230

	Page
Gas-mixtures with convertible components,	234
Case of peroxide of nitrogen,	237
Fundamental equations for the phases of equilibrium,	245

SOLIDS.

The conditions of internal and external equilibrium for solids in contact with fluids with regard to all possible states of strain,	343
Strains expressed by nine differential coefficients,	344
Variation of energy in an element of a solid,	344
Deduction of the conditions of equilibrium,	346
Discussion of the condition which relates to the dissolving of the solid,	352
Fundamental equations for solids,	361
Concerning solids which absorb fluids,	375

THEORY OF CAPILLARITY.

SURFACES OF DISCONTINUITY BETWEEN FLUID MASSES.

Preliminary notions.—Surfaces of discontinuity.—Dividing surface,	380
Discussion of the problem.—The particular conditions of equilibrium for contiguous masses relating to temperature and the potentials which have already been obtained are not invalidated by the influence of the surface of discontinuity.—Superficial energy and entropy.—Superficial densities of the component substances.—General expression for the variation of the superficial energy.—Condition of equilibrium relating to the pressures in the contiguous masses,	380
Fundamental equations for surfaces of discontinuity between fluid masses,	391
Experimental determination of the same,	394
Fundamental equations for plane surfaces,	395
Stability of surfaces of discontinuity—	
(1) with respect to changes in the nature of the surface,	400
(2) with respect to changes in which the form of the surface is varied,	405
On the possibility of the formation of a fluid of different phase within any homogeneous fluid,	416
On the possible formation at the surface where two different homogeneous fluids meet of a fluid of different phase from either,	422
Substitution of pressures for potentials in fundamental equations for surfaces,	429
Thermal and mechanical relations pertaining to the extension of surfaces of discontinuity,	434
Impermeable films,	440
The conditions of internal equilibrium for a system of heterogeneous fluid masses without neglect of the influence of the surfaces of discontinuity or of gravity,	442
Conditions of stability,	451
On the possibility of the formation of a new surface of discontinuity where several surfaces of discontinuity meet,	453
The conditions of stability for fluids relating to the formation of a new phase at a line in which three surfaces of discontinuity meet,	456
The conditions of stability for fluids relating to the formation of a new phase at a point where the vertices of four different masses meet,	464
Liquid films,	467
Definition of an element of the film,	467
Each element may generally be regarded as in a state of equilibrium.—Properties of an element in such a state and sufficiently thick for its interior to have the properties of matter in mass.—Conditions under which an extension of the film will not cause an increase of tension.—When the film has more than one component which does not belong to the contiguous masses, extension will in general cause an increase of tension.—Value of the elasticity of the film deduced from the fundamental equations of the surfaces and masses.—Elasticity manifest to observation,	468
The elasticity of a film does not vanish at the limit at which its interior ceases to have the properties of matter in mass, but a certain kind of instability is developed,	472
Application of the conditions of equilibrium already deduced for a system under the influence of gravity (pages 447, 448) to the case of a liquid film,	473
Concerning the formation of liquid films and the processes which lead to their destruction.—Black spots in films of soap-water,	475

SURFACES OF DISCONTINUITY BETWEEN SOLIDS AND FLUIDS.

	Page
Preliminary notions,	482
Conditions of equilibrium for isotropic solids,	485
Effect of gravity,	488
Conditions of equilibrium in the case of crystals,	489
Effect of gravity,	492
Limitations,	493
Conditions of equilibrium for a line at which three different masses meet, one of which is solid,	495
General relations,	497
Another method and notation,	497

ELECTROMOTIVE FORCE.

Modification of the conditions of equilibrium by electromotive force,	501
Equation of fluxes.—Ions.—Electro-chemical equivalents,	501
Conditions of equilibrium,	502
Four cases,	503
Lippmann's electrometer,	506
Limitations due to passive resistances,	507
General properties of a perfect electro-chemical apparatus,	508
Reversibility the test of perfection,	508
Determination of the electromotive force from the changes which take place in the cell.—Modification of the formula for the case of an imperfect apparatus,	509
When the temperature of the cell is regarded as constant, it is not allowable to neglect the variation of entropy due to heat absorbed or evolved.—This is shown by a Grove's gas battery charged with hydrogen and nitrogen,	510
by the currents caused by differences in the concentration of the electrolyte,	511
and by electrodes of zinc and mercury in a solution of sulphate of zinc,	511
That the same is true when the chemical processes take place by definite proportions is shown by <i>a priori</i> considerations based on the phenomena exhibited in the direct combination of the elements of water or of hydrochloric acid,	513
and by the absorption of heat which M. Favre has in many cases observed in a galvanic or electrolytic cell,	516
The different physical states in which the ion is deposited do not affect the value of the electromotive force, if the phases are coexistent.—Experiments of M. Raoult,	518
Other formulæ for the electromotive force,	520

I N D E X .

- Acanthouotus serratus*, 31.
Acanthozone cuspidata, 32, 14.
Acipenser, ventral fin of, 300.
Acirsa borealis, 6, 8, 10.
Actinia nodosa, 54.
Admete viridula, 4, 12, 14, 17, 23.
Ega psora, 22.
Eolis rubibranchialis, 14.
 species, 6.
Aglaophenia Franciscana, 262, 263.
 myriophyllum, 15.
 struthionides, 262, 250, 251, 263.
Amage auricula, 23.
Amarœcium glabrum, 14.
 species, 6.
Amauropsis helicoides, 6, 8.
Ammodontes assimilis, 16, 19, 20.
Ammotrypane fimbriata, 16, 18.
Ampeliscia Gaimardi, 32.
 species, 3, 4, 15.
Ampharete arctica, 16, 19.
 Finnarchica, 17.
 gracilis, 17.
Amphicteis Gunneri, 17, 19, 23.
Amphipholis elegans, 7.
Amphioxus, median fins in, 281.
Amphithoë maculata, 34.
 panopla, 31.
Amphithonotus cataphractus, 31.
Amphitrite cirrata, 17.
 Greenlandica, 20.
Anchylomera, 26.
Ancistria capillaris, 16, 19, 22.
Anomia aculeata, 6, 9, 15, 18, 20, 23.
Anthura brachiata, 16.
Antinœ angusta, 36, 22.
 Sarsi, 37, 18.
Aphrodita aculeata, 4, 5.
Aporrhais occidentalis, 8, 13, 14, 15, 23.
Arca pectunculoides, 23.
Archaster arcticus, 18, 24.
Ascidioopsis complanata, 18.
Astarte Banksii, 13.
 castanea, 5, 6, 8, 22.
 crebricostata, 50.
 elliptica, 13, 15.
 lens, 50, 11, 17, 23.
 quadrans, 6, 8, 20, 23.
 sulcata, 50.
 undata, 50, 9, 10, 13.
Asterias vulgaris, 4, 21.
Autolytus, 26.
Axiotea catenata, 16.

Balanus porcatus, 3, 5, 8, 19, 22.
Bela cancellata, 11.
 decussata, 6, 10.
 harpularia, 3, 4, 5, 6, 8, 20, 21.
 pleurotomaria, 6, 20.
 turricula, 3, 4, 6, 20.
 violacea, 8, 12.
Bimeria gracilis, 252, 251.
Bolocera Tuediæ, 24.
Boltenia Bolteni, 13.
 clavata, 6.
Bougainvillia Mertensii, 250, 251.
Brada species, 15.
Buccinum undatum, 3, 4, 12, 20.
Bugula avicularia, 18, 24.
 Murrayana, 7, 9, 20, 22, 23.
Byblis Gaimardi, 32, 34.

Caberea Ellisii, 6, 24.
Calliopius leviusculus, 26.
Calycella producta, 11.
 syringa, 7, 13, 21, 63.
Campanularia calceolifera, 60.
 cylindrica, 254, 251.
 everta, 253, 251.
 fusiformis, 254, 251.
 Huicksii, 7, 9, 13, 21.
 noliformis, 60.
 pygmæa, 59.
 verticillata, 4, 7, 9, 10, 11, 13, 21.
 volubilis, 7, 9.
Cancer irroratus, 3, 4, 5, 19.
Caprella species, 8.
Cardium pinnulatum, 6, 8, 13, 14, 17, 20,
 22, 23.
Caridion Gordoni, 28, 16.
Cellaria fistulosa, 7, 9.
Cellepora avicularis, 13.
 ramulosa, 7, 24.
 scabra, 7, 9, 24.
 tuberosa, 20.
Cellularia Peachii, 24.
 species, 18, 22.
 ternata, 6, 9, 20, 24.
Cerapus rubricornis, 5, 19, 34.
Cerianthus borealis, 54, 18, 24.

- Cestum Veneris*, 26.
Chaetozone setosa, 18.
Chalina oculata, 22.
Charybdea periphylla, 52, 26.
 Chondrodite from the Tilly Foster iron mine, Brewster, New York, by E. S. Dana, 67.
Chone infundibuliformis, 20.
Cibrella sanguinolenta, 7, 9, 14, 21.
Cistenides granulata, 3, 6, 12, 15, 23.
 Clark, S. F., new and rare species of Hydroids from the New England coast, 58.
 hydroids of the Pacific Coast of the United States south of Vancouver Island. With a report upon those in the museum of Yale College, 249.
Clavelina crystallina, 51.
Clidiophora trilineata, 5.
Clymenella torquata, 4.
Clytia Johnstoni, 13, 24.
Conilera polita, 3, 22.
Coppinia arcta, 9, 21.
Coryne rosaria, 250.
Cotulina Greenei, 250, 257.
Crangon vulgaris, 3, 4, 10, 19.
Crenella glandula, 5, 10, 18, 20.
Crepidula fornicata, 5.
 plana, 3, 4, 6, 12, 24.
 unguiformis, 3, 4, 6, 12.
Crisea eburnea, 15.
Crossaster papposus, 9, 13, 21.
Cryptodon Gouldii, 18.
 obesus, 11.
Ctenodiscus crispatus, 18.
Cuspidella humilis, 7.
Cyclocardia borealis, 6, 10.
 Novanglicæ, 13.
Cylichna alba, 8, 17, 23.
Cynthia carnea, 13.
Cyprina Islandica, 6, 8, 10, 20, 22.
 Dana E. S., chondrodite from the Tilly-Foster iron mine, Brewster, New York, 67.
Dentalium occidentale, 23.
Dendronotus arborescens, 8, 13.
Diastylis quadrispinosa, 28, 15.
 sculpta, 29, 15.
Diodora noachiana, 8, 11, 20.
Diphasia fallax, 7, 9.
 mirabilis, 53, 9.
Discofascigera lucernaria, 18.
Discopora Skenei, 7, 9, 14.
Discoporella verrucaria, 24.
Dodecaceria concharum, 21.
 Dredgings in the region of St. George's Banks in 1872, report on, by S. I. Smith and O. Harger, 1.
Dulichia species, 3.
Echinarachnius parma, 3, 4, 5, 9, 10, 11, 21, 22, 24.
Edwardsia species, 19.
Elasmobranchii, median fins of, 284.
Ensatella Americana, 3.
Entalis striolata, 6, 8, 11, 13, 14, 17, 23.
Epelys montosus, 3.
Epizoanthus Americanus, 55, 9, 10, 11.
Equilibrium of heterogeneous substances, by J. W. Gibbs, 108, 343.
Eschara papposa, 13.
Escharoides rosacea, 13.
Eteone depressa, 16.
Eueranta villosa, 37, 22.
Eudendrium capillare, 7, 9, 13, 21, 24.
 ramosum, 7, 9, 11, 21.
 species, 253, 251.
Eumenia crassa, 16, 18.
Eunice vivida, 11, 41.
Eunoë nodosa, 12, 14.
Eupagurus Bernhardus, 27, 3, 4, 5, 19, 21, 22.
 Kroyeri, 28, 5, 8, 10, 12, 14, 22, 25.
 pubescens, 27, 3, 5, 8, 9, 10, 15, 19, 22.
Eupolynoë occidentalis, 37.
Eusyllis phosphorea, 39, 20.
Farrella familiaris, 4.
 Fins, median and paired, a contribution to the history of vertebrate limbs, by J. K. Thacher, 281.
Flustra papyrea, 15.
Fusus Islandicus, 6, 8, 20.
Gammarus ornatus, 15.
 purpuratus, 5, 8.
Gemellaria loricata, 4, 20.
 Gibbs, J. W., equilibrium of heterogeneous substances, 108, 343.
Glandula arenicola, 51, 5, 20, 23.
Glycera capitata, 43.
 viridescens, 42.
Goniada maculata, 42, 15, 16, 22.
Gonothyræa hyalina, 9.
 Lovéni, 7.
 tenuis, 61.
Grammaria abietina, 9, 15, 21.
 robusta, 21.
Grymæa spiralis, 44, 17, 23.
Halecium articulatum, 63.
 Beanii, 7, 64.
 labrosum, 21.
 plumosum, 64.
 robustum, 53, 11.
 teuellum, 7, 250, 251.
Halirages fulvocinctus, 15.
Hanleia mendicaria, 6, 8, 14, 20.
 Harger, O., report on the dredgings in the region of St. George's Banks, in 1872, 1.
Harmothoë imbricata, 5, 8, 12, 14, 15, 20, 22.
Harpina fusiformis, 29, 16, 18.
 plumosa, 29.

- Hippa, structure of the mandibles and maxillæ in. 337.
 Hippa talpoida, early stages of, with a note on the structure of the mandibles and maxillæ in Hippa and Remipes, by S. I. Smith, 311.
 Hippolyte pusiola, 12, 19.
 spina, 12.
 Hornerea lichenoides, 24.
 Hyalonema longissimum. 55.
 Hyas araneus, 14, 15.
 coarctatus, 5, 8, 12, 14, 19, 22.
 Hydractinia polyclina, 3, 7, 9, 10, 13, 22.
 Hydrallmania falcata, 4, 7, 9, 15, 21, 249, 260.
 Franciscana, 260, 249.
 Hydroids, new and rare species from the New England coast, by S. F. Clark, 58.
 of the Pacific Coast of the United States south of Vancouver Island. With a report upon those in the museum of Yale College. by S. F. Clark, 249.
 Hyperia, 26.
 Lænilla mollis, 35, 16.
 Lættonice flicornis, 22.
 Lafoëa dumosa, 7, 9, 21, 250, 251.
 fruticosa, 53.
 gracillima, 53, 7, 9, 11, 13, 14.
 Lagisca propinqua, 20.
 rarispinia, 14.
 Laomedea Pacifica, 251.
 rigida, 251.
 Latreutes ensiferus, 26.
 Leda tenuisulcata, 18, 20, 23.
 Lepas fascicularis, 26.
 pectinata, 26.
 Lepeta cæca, 17.
 Leptasterias compta, 7.
 Stimpsoni, 21.
 Leptoplana ellipsoides, 12.
 Leodice vivida, 41, 11, 16, 20.
 Library, additions to the, v.
 Lophothuria Fabricii, 9, 13, 18, 21.
 squamata, 11.
 Lucifer, 26.
 Lumbriconereis fragilis, 11, 16, 18, 22.
 Lunatia Grœnlandica, 11, 12, 17, 23.
 heros, 3, 4, 5, 8, 20.
 immaculata, 3, 5, 6, 11, 20.
 Macoma proxima, 15, 17.
 Mactra solidissima, 5, 10, 20, 21.
 Mæra Danae, 5.
 Maldane Sarsii, 16, 19, 22.
 Manania auricula, 15.
 Margarita cinerea, 13, 14, 17, 23.
 Grœnlandica, 13.
 helicina, 49.
 obscura, 6, 8, 10, 11, 20, 21, 23.
 undulata, 13.
 varicosa, 15.
 Meckelia ingens, 4.
 lurida, 3, 4, 5, 17.
 Melinna eristata, 17, 23.
 Melita dentata, 5, 8, 10, 19.
 Modiolaria modiolus, 20.
 Modiolaria corrugata, 9.
 discors, 6.
 lævigata, 20.
 nigra, 3.
 Monoculodes borealis, 31, 15.
 nubilus, 31.
 Mopsea borealis, 35.
 Mya truncata, 13.
 Myrriozoum coarctatum, 13.
 Mysis, 26.
 Myxine, median fins in, 282.
 Natica clausa, 6, 8, 10, 11, 12, 14, 17, 19, 21, 23.
 Nautilograpsus minutus, 26.
 Næra arctica, 17, 23.
 Nephthys ciliata, 16.
 circinata, 38, 11, 18.
 incisa, 39.
 ingens, 39, 16, 18, 22.
 Neptunea curta, 6, 8, 20.
 decemcostata, 6, 8, 12.
 pygmæa, 3, 4, 6, 8, 11, 12, 17, 20, 23.
 Neptunus Sayi, 26.
 Nereis pelagica, 4, 6, 12, 16, 20.
 virens, anatomy and habits of, by F. M. Turnbull, 265.
 Newton, H. A., on the transcendental curves, $\sin y \sin my = a \sin x \sin nx + b$, 97.
 Nicomache lumbricalis, 16.
 Ninoe nigripes, 40, 16.
 Nothria conchylega, 41, 10, 12, 14, 16, 18, 20, 22.
 opalina, 41, 16, 22.
 Notomastus latericius, 16.
 Nychia cirrosa, 12.
 Nymphon grossipes, 5, 14.
 Obelia bicuspidata, 58.
 bidentata, 58.
 dichotoma, 258.
 gelatinosa, 59, 258.
 geniculata, 258.
 Oedicerus affinis, 31.
 lynceus, 30.
 propinquus, 30.
 Onuphis conchylega, 41.
 Eschrichti, 41.
 Opercularella lacerata, 62.
 pumila, 61.
 Ophiacantha spinulosa, 18, 24.
 Ophioglypha affinis, 18, 24.
 robusta, 7, 13, 15, 21.
 Sarsii, 11, 13, 14, 18, 24.
 Ophiopholis aculeata, 7, 13, 14, 15, 18, 21.
 Ostrea Virginica, 5.
 Oxycephalus, 26.

- Pagurus Bernhardus, 27.
 pubescens, 27.
 Pandalus annulicornis, 3, 4, 5, 8, 10, 19, 25.
 Paramphithoë cataphracta, 31, 12, 19.
 panopla, 31.
 pulchella, 31, 8, 19.
 Parypha microcephala, 251.
 Pecten Islandicus, 6, 9, 13, 18, 20.
 pustulosus, 50, 23.
 tenuicostatus, 9, 18, 20, 22.
 Pelagia cyanella, 26.
 Pennatula aculeata, 54, 18, 24.
 Canadensis, 54.
 phosphorea, 54.
 Pentacta assimilis, 11, 18.
 Pera crystallina, 51, 6, 11.
 pellucida, 51.
 Petromyzon, median fins in, 282.
 Phascolosoma boreale, 47, 17, 23.
 cæmentarium, 46, 6, 8, 12, 17, 19, 23.
 tubicola, 46, 11, 17, 23.
 Philine species, 17.
 Phillips, A. W., on the transcendental curves, $\sin y \sin my = a \sin x \sin nx + b$, 97.
 Pholoë minuta, 16.
 Photis species, 3.
 Phoxus fusiformis, 29.
 Holbollii, 29.
 Kroyeri, 29, 19.
 plumosus, 29.
 Phronima, 26.
 Phrosina, 26.
 Phyllodoce catenula, 39, 5.
 pulchella, 39.
 species, 16.
 Physalia pelagica, 26.
 Pisa cristata, 17, 23.
 Platyscelus, 26.
 Pleurobrachia rhododactyla, 26.
 species, 26.
 Pleuropus, 27.
 Pleurotomella Packardii, 48, 17.
 Pleustes tuberculatus, 31.
 Plumularia cristata, 263.
 Franciscana, 260, 263.
 gracilis, 250, 260, 263.
 setacea, 261, 249, 250, 251.
 struthionides, 262, 263.
 tenella, 65.
 Verrillii, 64.
 Podocerus nitidus, 5, 8.
 Polycirrus species, 17, 19.
 Pontogeneia inermis, 19.
 Potamilla neglecta, 44, 6, 12, 14, 17.
 oculifera, 6, 12.
 Praxilla, 16.
 gracilis, 16.
 prætermisssa, 16, 19.
 Tronoë, 26.
 Protula borealis, 46, 17, 23.
 media, 46, 17.
 Psolus phantapus, 9.
 Pteraster militaris, 13.
 Ptilocheirus pinguis, 22, 34.
 Pycnogonum littorale, 10.
 pelagicum, 10.
 Remipes, structure of the mandibles and maxillæ in, 337.
 Ringicula nitida, 48, 17, 23.
 Rhodine Lovéni, 16, 19.
 Rhynchobolus capitatus, 43, 8, 11, 16.
 Rhynchonella psittacea, 13, 15.
 Sabella neglecta, 44.
 pavonia, 17.
 Sabinea septemcarinata, 28, 8, 14.
 Sagitta, 26.
 Salpa, 26.
 Samytha sexcirrata, 17.
 Samythella, 43.
 elongata, 43, 17, 23.
 Sapphirina, 26.
 Scalaria Grœnlandica, 5, 13, 19, 21.
 Scalibregma inflatum, 22.
 Scalpellum Strœmi, 35, 10.
 Scaphander puncto-striatus, 23.
 Schizaster fragilis, 11, 18, 19, 24.
 Scolecolepis cirrata, 16, 22.
 Sertularella Gayi, 11, 24.
 polyzonias, 7, 13, 15, 21, 22, 51, 53, 260.
 tricuspidata, 7, 9, 10, 11, 13, 14, 21, 24.
 turgida, 259, 249, 251, 260.
 Sertularia abietina, 7, 21.
 anguina, 255, 256, 249, 250, 251, 254.
 argentea, 64, 11, 21, 257, 250, 251.
 corniculata, 251.
 cupressina, 4, 7, 9, 10, 21, 24.
 Franciscana, 250.
 furcata, 258, 249, 251.
 Greenii, 257, 251.
 labrata, 250, 255.
 latiuscula, 4, 7, 9, 21, 59.
 pinnata, 261.
 pumila, 250, 251, 262.
 setacea, 261.
 tricuspidata, 257.
 Siliqua costata, 4.
 Smith, S. I., report on the dredgings in the region of St. George's Banks, in 1872, 1.
 the early stages of Hippa talpoida, with a note on the structure of the mandibles and maxillæ in Hippa and Remipes, 311.
 Solaster endeca, 21.
 furcifer, 24.
 Spiocætopterus, 12, 22.
 Spiralis Gouldii, 26.
 Spirorbis borealis, 45.
 communis, 45.
 incidus, 12.
 nautiloides, 45, 6, 8, 20.
 quadrangularis, 21.

- Spirorbis valida*, 44, 12, 14.
Stegocephalus ampulla, 16.
Stenothoë peltata, 29, 3.
Sternaspis fossor, 16, 18.
Stromolophus meleagris, 26.
Strongylocentrotus Dröbachiensis, 5, 7, 8,
 9, 10, 11, 13, 21, 49.
Stylifer, 24.
 Stimpsonii, 49, 8, 10.
Styliola, 26.
 acus, 27.
Syrrhoë bicuspis, 30.
 crenulata, 30, 12.

Terebellides Stroemi, 17, 19, 23.
Terebratulina septentrionalis, 13, 14, 15,
 18, 23.
 Thacher, J. K., median and paired fins, a
 contribution to the history of vetebrate
 limbs, 281.
Thamnocnidia tubularoides, 250.
Thracia myopsis, 15, 17, 23.
 truncata, 6.
Thecophora ibla, 55, 7.
Thelepus cincinnatus, 6, 8, 12, 14, 20.
Thessarops hastata, 30.
Thuiaria articulata, 13.
Thyone fusus, 51.
 raphanus, 52.
 scabra, 51, 18, 24.
Thyropus, 26.
 species, 10, 26.
Tiron acanthurus, 30, 12.
Torellia vestita, 49, 23.
Trachydermon album, 13, 14, 17, 23.

Trachynema digitale, 26.
 Transcendental curves, by H. A. Newton
 and A. W. Phillips, 97.
Trichotropis borealis, 12.
Tritia trivittata, 3, 5.
Tritopsis aculeata, 12, 14.
Trophon Gunneri, 12, 14.
Trophonia aspera, 16, 18.
Tubularia elegans, 253, 250.
 indivisa, 7, 9, 10, 11, 13.
 Turnbull, F. M., anatomy and habits of
Nereis virens, 265.
Turritella acicula, 20.
 erosa, 13, 20, 44.
 reticulata, 14, 15.

Unciola irrorata, 3, 5, 10, 16, 19, 25.
Urticina crassicornis, 7, 9, 10, 11, 13, 24,
 25.
 digitata, 54.
 nodosa, 54, 11.

Velutina lævigata, 17, 49.
 zonata, 23.
Vesicularia armata, 11.
Vetumnus serratus, 31, 5, 19.
Virgularia Lyungmanii, 54, 24.

Xenoclea Batei, 34.
 megachir, 32, 3, 19.

Yoldia limatula, 4.
 obesa, 18, 19, 23.
 thraciformis, 19.

ERRATA TO VOL. III.

Page 1, line 13, for Hagerman, read Hageuman.

Page 5, line 30, for *Vetumnus*, read *Vertumnus*.

Page 9, last line, for 1873, read 1874.

Page 11, line 31, for *virticillata*, read *verticillata*.

Page 13, line 34, for *cappilare*, read *capillare*.

Page 23, line 4, for *Smythella*, read *Samythella*.

Page 28, line 19, for **Caridon**, read **Caridion**.

Page 35, line 3, for **Scapellum**, read **Scalpellum**.

Page 58, line 14, for branches, read branchlets.

Page 60, line 12, for Plate X, read IX.

Page 167, formula (168), for m_1 , read μ_1 .

Page 167, formula (169), for m_1, \dots, m_{n-1} , read μ_1, \dots, μ_{n-1} .

Page 239, formula (333), for $\frac{C}{t}$ read $\frac{C}{a_2 t}$.

Page 295, note †, 1st line, for Ueber de Gliedmassen, read Ueber das Skelett der Gliedmassen.

Page 304, note, for prominal, read proximal.

Page 356, last line but two, for crystalline solid, read solid of continuous crystalline structure.

Page 385, line 13, for M' , read M .








Pages 391, 394, 395, 400, in headings, after *Discontinuity*, add *between Fluid Masses*.

Page 403, line 16, after any other film, add of the same components.


Page 405, line 29, after this, add case.

Page 432, line 15 of foot-note, for II, read II₁.


In figure 40, plate XVI, there is a series of ovals around one-half of the real double points. There should be added to the curve, as represented, a like series of ovals around each of the remaining real double points.




1.




2.




4.



5.




3.



1.

2.


3.



4.

5.


6.



7.


8.

9.






1.



2.



3.



4.







1.



2.




3.



4.




5.











1.




3.



2.



4.






1.




2.




3.



4.



1



2



3



4



7



8



5



9



6









Fig. 1.




Fig. 2.




Fig. 5.




Fig. 3.




Fig. 4.




Fig. 10.




Fig. 6.




Fig. 7.




Fig. 8.





Fig. 9.






Fig. 11.




Fig. 12.




Fig. 13.




Fig. 14.




Fig. 15.




Fig. 16.




Fig. 18.




Fig. 17.




Fig. 19.




Fig. 20.






Fig. 21.

Curves whose equations are $y = a \sin x \sin mx$.




1. $m = 1$.




2. $m = \frac{1}{2}$.




3. $m = \frac{1}{3}$.



4. $m = \frac{2}{5}$.




5. $m = \frac{1}{4}$.



6. $m = \frac{3}{4}$.




7. $m = \frac{1}{5}$.



8. $m = \frac{2}{5}$.




9. $m = \frac{3}{5}$.




10. $m = \frac{1}{6}$.




11. $m = \frac{1}{6}$.




12. $m = \frac{5}{6}$.




13. $m = \frac{1}{7}$.




14. $m = \frac{2}{7}$.




15. $m = \frac{3}{7}$.




16. $m = \frac{4}{7}$.




17. $m = \frac{5}{7}$.




18. $m = \frac{6}{7}$.



19. $m = \frac{1}{8}$.




20. $m = \frac{3}{8}$.



21. $m = \frac{5}{8}$.

Curves whose equations are $y = a \sin x \sin nx$.




22. $m = \frac{1}{6}$.

23. $m = \frac{1}{9}$.

24. $m = \frac{2}{9}$.

25. $m = \frac{4}{9}$.




26. $m = \frac{1}{3}$.

27. $m = \frac{7}{9}$.

28. $m = \frac{8}{9}$.


29. $m = \frac{1}{10}$.



30. $m = \frac{2}{10}$.

31. $m = \frac{7}{10}$.

32. $m = \frac{9}{10}$.




33. $m = \frac{1}{11}$.


34. $m = \frac{3}{11}$.

35. $m = \frac{5}{11}$.


36. $m = \frac{7}{11}$.




37. $m = \sqrt{\frac{1}{5}}$.




38. $\sin y \sin y = \sin x \sin x.$




39. $\sin y \sin y = -\sin x \sin x.$




40. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{1}{2} x.$




41. $\sin y \sin \frac{1}{3} y = \sin x \sin \frac{1}{3} x.$




42. $\sin y \sin \frac{1}{4} y = \sin x \sin \frac{1}{4} x.$




44. $\sin y \sin \frac{1}{6} y = \sin x \sin \frac{1}{6} x.$




43. $\sin y \sin \frac{1}{5} y = \sin x \sin \frac{1}{5} x.$




45. $\sin y \sin \frac{1}{4} y = \sin x \sin \frac{1}{4} x.$




46. $\sin y \sin \frac{1}{5} y = \sin x \sin \frac{1}{5} x.$




47. $\sin y \sin \frac{1}{10} y = \sin x \sin \frac{1}{10} x.$




48. $\sin y \sin \frac{2}{3} y = \sin x \sin \frac{2}{3} x.$




49. $\sin y \sin \frac{3}{4} y = \sin x \sin \frac{3}{4} x.$




50. $\sin y \sin \frac{2}{3} y = \sin x \sin \frac{2}{3} x.$




51. $\sin y \sin \frac{3}{4} y = \sin x \sin \frac{3}{4} x.$




53. $\sin y \sin \frac{3}{4} y = \sin x \sin \frac{3}{4} x.$




52. $\sin y \sin \frac{2}{3} y = \sin x \sin \frac{2}{3} x.$




54. $\sin y \sin \frac{4}{5} y = \sin x \sin \frac{4}{5} x.$




55. $\sin y \sin \frac{3}{8} y = \sin x \sin \frac{3}{8} x.$




56. $\sin y \sin \frac{5}{8} y = \sin x \sin \frac{5}{8} x.$




57. $\sin y \sin \frac{2}{3} y = \sin x \sin \frac{2}{3} x.$




58. $\sin y \sin \frac{1}{3} y = \sin x \sin \frac{1}{3} x.$




59. $\sin y \sin \frac{3}{10} y = \sin x \sin \frac{3}{10} x.$




60. $\sin y \sin \frac{2}{11} y = \sin x \sin \frac{2}{11} x.$




61. $\sin y \sin \frac{3}{11} y = \sin x \sin \frac{3}{11} x.$




62. $\sin y \sin \frac{4}{11} y = \sin x \sin \frac{4}{11} x.$




63. $\sin y \sin \frac{5}{11} y = \sin x \sin \frac{5}{11} x.$




64. $\sin y \sin \frac{9}{11} y = \sin x \sin \frac{9}{11} x.$




65. $\sin y \sin \frac{7}{11} y = \sin x \sin \frac{7}{11} x.$




66. $\sin y \sin \frac{4}{5} y (\sin x \sin \frac{5}{8} x - .40) = 0.$




67. $\sin y \sin \sqrt{\frac{2}{3}} y = \sin x \sin \sqrt{\frac{2}{3}} x.$




68. $\sin y \sin \frac{5}{7} y = \sin x \sin \frac{5}{7} x.$




69. $\sin y \sin \frac{5}{7} y = -\sin x \sin \frac{5}{7} x.$




70. $\sin y \sin \frac{5}{8} y = \sin x \sin \frac{5}{8} x.$




71. $\sin y \sin \frac{5}{8} y = -\sin x \sin \frac{5}{8} x.$




72. $\sin y \sin \frac{1}{7} y = .28 \sin x \sin \frac{1}{7} x.$




73. $\sin y \sin \frac{1}{7} y = .63 \sin x \sin \frac{1}{7} x.$




74. $\sin y \sin \frac{1}{5} y = -\sin x \sin \frac{1}{5} x.$




75. $\sin y \sin \frac{1}{5} y = .35 \sin x \sin \frac{1}{5} x.$




76. $\sin y \sin y = \sin x \sin \frac{1}{2} x.$




77. $\sin y \sin y = \sin x \sin \frac{1}{3} x.$




78. $\sin y \sin y = \sin x \sin \frac{1}{4} x.$




79. $\sin y \sin y = \sin x \sin \frac{1}{5} x.$




80. $\sin y \sin y = -\sin x \sin \frac{1}{3} x.$




82. $\sin y \sin \frac{4}{7} y = \sin x \sin \frac{3}{7} x.$




81. $\sin y \sin y = \sin x \sin \frac{3}{7} x.$




83. $\sin y \sin \frac{4}{7} y = \sin x \sin \frac{3}{11} x.$




84. $\sin y \sin \frac{3}{8} y = \sin x \sin \frac{2}{7} x.$




85. $\sin y \sin \frac{1}{4} y = \sin x \sin \frac{7}{11} x.$




86. $\sin y \sin \frac{1}{6} y = \sin x \sin \frac{1}{2} x.$




91. $\sin y \sin \frac{2}{11} y = \sin x \sin \frac{1}{11} x.$




92. $\sin y \sin \frac{1}{3} y = -\sin x \sin \frac{1}{3} x + .56.$




93. $\sin y \sin \frac{1}{3} y = -\sin x \sin \frac{1}{3} x.$




94. $\sin y \sin \frac{1}{3} y = -\sin x \sin \frac{1}{3} x - .44.$




95. $\sin y \sin \frac{1}{3} y = -\sin x \sin \frac{1}{3} x + k.$




96. $\sin y \sin \frac{3}{7} y = -\sin x \sin \frac{3}{7} x - .80.$




97. $\sin y \sin \frac{3}{7} y = -\sin x \sin \frac{3}{7} x - .38.$




99. $\sin y \sin \frac{3}{7} y = -\sin x \sin \frac{3}{7} x$




98. $\sin y \sin \frac{3}{7} y = -\sin x \sin \frac{3}{7} x - .22$




100. $\sin y \sin \frac{3}{7} y = -\sin x \sin \frac{3}{7} x + .11.$




101. $\sin y \sin \frac{3}{7} y = -\sin x \sin \frac{3}{7} x + .31.$




102. $\sin y \sin \frac{3}{7} y = -\sin x \sin \frac{3}{7} x + .69.$



103. $\sin y \sin \frac{3}{7} y = -\sin x \sin \frac{3}{7} x + 1.00.$




104. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x - .49.$




105. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x - .21.$




106. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x - .18.$




107. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x - .11.$



108. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x$.



109. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x + .04$.




110. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x + .07$.



111. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x + .17$.




112. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x + .22.$



113. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x + .32.$



114. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x + .47.$



115. $\sin y \sin \frac{3}{11} y = -\sin x \sin \frac{3}{11} x + .60.$

116. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{3}{7} x - 1.46.$

117. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{3}{7} x - 1.00.$

118. $\widehat{\sin y} \widehat{\sin \frac{1}{2} y} = \widehat{\sin x} \widehat{\sin \frac{3}{7} x} - .77.$


119. $\widehat{\sin y} \widehat{\sin \frac{1}{2} y} = \widehat{\sin x} \widehat{\sin \frac{3}{7} x} - .69.$

120. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{3}{7} x - .39.$


121. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{3}{7} x - .23.$

122. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{3}{7} x - .11.$


123. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{3}{7} x.$




124. $\sin y \sin \frac{1}{2}y = \sin x \sin \frac{3}{7}x + .08.$




125. $\sin y \sin \frac{1}{2}y = \sin x \sin \frac{3}{7}x + .15.$




126. $\sin y \sin \frac{1}{2}y = \sin x \sin \frac{3}{7}x + .38.$




127. $\sin y \sin \frac{1}{2}y = \sin x \sin \frac{3}{7}x + .66.$




128. $\sin y \sin \frac{1}{2}y = \sin x \sin \frac{3}{7}x + .92.$




129. $\sin y \sin \frac{1}{2}y = \sin x \sin \frac{3}{7}x + 1.15.$




130. $\sin y = k \sin x.$




131. $\sin y \sin \frac{1}{2} y = .77 \sin x \sin \frac{3}{7} x.$




132. $\sin y \sin \frac{1}{2} y = .84 \sin x \sin \frac{3}{7} x.$




133. $\sin y \sin \frac{1}{2} y = 1.11 \sin x \sin \frac{3}{7} x.$




134. $\sin y \sin \frac{1}{2} y = 2.0 \sin x \sin \frac{3}{7} x.$




135. $\sin y \sin \frac{1}{2} y = 6.9 \sin x \sin \frac{3}{7} x.$




136. $\sin y \sin \frac{1}{2} y = .77 \sin x \sin \frac{1}{3} x.$




137. $\sin y \sin \frac{1}{2} y = .77 \sin x \sin \frac{1}{3} x + .34.$




138. $\sin y \sin \frac{1}{2} y = .77 \sin x \sin \frac{1}{3} x + .77.$




139. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{1}{3} x + .23.$



140. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{1}{3} x + .21.$




141. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{1}{3} x.$




142. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{1}{3} x - .56.$




143. $\sin y \sin \frac{1}{2} y = \sin x \sin \frac{1}{3} x - .77.$




144. $\sin y \sin \frac{1}{2} y = 1.37 \sin x \sin \frac{1}{3} x + .60.$




145. $\sin y \sin \frac{1}{2} y = 1.37 \sin x \sin \frac{1}{3} x.$




146. $\sin y \sin \frac{1}{2} y = 1.37 \sin x \sin \frac{1}{3} x - .77.$



















$$147. \quad \text{Sin } y \text{ Sin } \frac{2}{5} y = \text{sin } x \text{ sin } \frac{3}{5} x + \kappa.$$














148. $\sin y \sin \frac{2}{7} y = k \sin x \sin \frac{3}{7} x.$





























Fig 11




Fig 7.




Fig. 9.




Fig 8.






Fig 10








Fig. 12.



Fig. 16



Fig. 13




Fig. 14




Fig. 15.



























Fig. 17.



















Fig. 61.




Fig. 60




Fig. 59




Fig. 58




Fig. 57.




Fig 68




Fig 63




Fig 64




Fig 65



Fig 67




Fig 66



Fig 62



Fig. 69




Fig. 71







Fig. 70









3 2044 106 253 149

DATE DUE

~~DEC 31 1983~~

MAY 03 2003

