
This is a reproduction of a library book that was digitized by Google as part of an ongoing effort to preserve the information in books and make it universally accessible.

Google™ books

<https://books.google.com>





Division. 658 Shelf.

American Geographical Society.

Acquired June 10 1914


by purchase

from B. F. Stevens & Brown


No. 19465 Cost £ 25-10-0

(For 17 vols.)

Rebarded
by A.G.S.



LIBRARY of the
OHIO STATE
UNIVERSITY



1. The first part of the document discusses the importance of maintaining accurate records of all transactions and activities. It emphasizes the need for transparency and accountability in financial reporting.

2. The second part of the document outlines the various methods and techniques used to collect and analyze data. It highlights the importance of using reliable sources and ensuring the accuracy of the information gathered.

3. The third part of the document focuses on the interpretation and analysis of the collected data. It discusses the various statistical and analytical tools used to identify trends and patterns in the data.

4. The fourth part of the document provides a detailed overview of the findings and conclusions drawn from the analysis. It discusses the implications of the results and offers recommendations for future research and action.

[No. 28.]

JOURNAL
OF THE
STRAITS BRANCH
OF THE
ROYAL ASIATIC SOCIETY.

JUNE, 1891.

AMERICAN
GEOGRAPHICAL
SOCIETY

SINGAPORE:

PRINTED AT THE GOVERNMENT PRINTING OFFICE.

AGENTS OF THE SOCIETY:

London and America. ... TRÜBNER & Co.
Paris, ... ERNEST LEROUX & C^{IE}.
Germany, ... K. F. KOEHLER'S ANTIQUARIUM, Leipzig.

100-1000
1000-1000
1000-1000

STATE OIL

1000-1000

Digitized by Google

TABLE OF CONTENTS.

The Grasses and Sedges of the Malay Peninsula— <i>by H. N. Ridley, M.A., F.L.S.,</i>	1
Outline of the History of the Dindings from the 17th Century to the present time— <i>by E. M. Merwether,</i> ...	35
Native Names of Streets in Singapore— <i>by H. T. Haughton,</i> ...	49
Notes on a Trip to Bukit Etam, Selangor— <i>by Lieut. H. J. Kelsall, B.E.,</i>	67
Anatomical Notes on Malay Apes— <i>by Dr. A. Keith,</i> ...	77
The Fishing Industry of Krian and Kurau, Perak— <i>by A. T. Dew,</i>	95
A Day at Christmas Island— <i>by H. N. Ridley, M.A., F.L.S.,</i> ...	123
Occasional Notes:—	
Discovery of a Stone Implement in Singapore, ...	141
Pantang Larang of Negri Sembilan,	142
On the Occurrence of a Rhododendron in Singapore,...	144
Pogonia punctata, Bl., in Singapore,	146
The Keringga,	147
Eudromias veredus in Singapore,	147
Frugivorous Habits of the Tupaia,	148

THE GRASSES AND SEDGES OF THE MALAY PENINSULA.

BY

H. N. RIDLEY, M.A., F.L.S.



IN publishing this list of the Grasses and Sedges of the Peninsula (*Gramineæ* and *Cyperaceæ*), I am well aware that it is by no means complete. From many parts of the Peninsula we have no specimens of these families, and especially is this the case as regards the great mountain region which traverses the Peninsula. From the uplands we shall doubtless procure many interesting kinds when the country is more opened up, and from the northern parts of the Peninsula bordering on Siam we may expect to obtain many Assam and Burmese species not yet met with. This list rather represents the lowland glumaceous flora of the South.

A country which like this is covered with dense forest for the greater part is, as a rule, poorly provided with grasses and sedges, for few of these plants occur in high forest. Yet on the whole there is a considerable variety, and among them not a few very interesting and curious plants are to be met with.

I am much indebted to Professor HACKEL of St. Polten for identifying many of the grasses, and to Mr. C. B. CLARKE for much assistance in the matter of *Cyperaceæ*.

Habitats.—The most productive localities for grasses and sedges are the damp low swamps and rice-fields, the banks of streams and the sandy shores of rivers and seas. The open country where the jungle has been cleared and secondary forest is returning is very barren of plants of interest, although large tracts are covered with glumaceous plants. *Imperata*, *Ischæmum*, *Paspalum*, *Panicum* of the *Digitaria*

section form the bulk of the grass flora here, while the *Cyperaceæ* are represented by a few common *Sclerias*, *Rhynchospora Wallichii* and *Fimbristylis*. The low swampy ground and rice-fields are rich in the genera *Panicum*, *Isachne*, *Lcersia*, *Scirpus*, *Rhynchospora*, *Scleria*, *Heleocharis* and *Cyperus*. Along the streams and rivers some fine species may be met with, among which *Scleria oryzoides*, *Scirpus grossus* and *Sc. mucronatus*, *Lepironia*, many *Cyperi*, the common reed *Phragmites Roxburghii*, and the wild sugar-cane *Saccharum arundinaceum* and *S. Ridleyi* are very conspicuous; the last three grasses forming huge thickets in many places. The *Sclerias*, too—*Scl. malaccensis* and *Scl. oryzoides*—often cover large tracts of wet open land, forming jungles about six feet high, very laborious to traverse. In the dense forests the glumaceous plants almost disappear, yet there are several very interesting genera to be met with. The large group of *Hypolytrææ* is well represented here by many species of *Mapania* and *Pandanophyllum*. The latter of which, as the name denotes, so much resemble the smaller species of screw-pines that, unless the inflorescence is found, they may be easily mistaken for them. Grasses are almost unrepresented in the deep jungle. *Centotheca lappacea* and *Lophatherum*, however, sometimes find their way far in, being transported widely by their very adherent spikelets attaching themselves to wild beasts. In many places, too, there are large tracts of bamboos, but at present specimens of these in flower have rarely been met with, so that of what species they are is not yet known.

On the sandy districts by the sea and along the river banks, a large number of species are to be met with, and among them several very striking plants. Such are the Porcupine grass (*Spinifex*), *Thouarea*, a curious creeping grass, *Schænus calostachyus*, Poir, and two species of *Eriachne*, *Gahnia*, *Lepturus*, *Tricostularia* and many other species worthy of cultivation.

Some of our native grasses are worthy of cultivation as ornamental plants, among these the *Saccharums* and *Phragmites* form large tufts like those of Pampas grass. The *Pandanophylla* and *Mapaniæ* are fine foliage plants suited for pot culture, and *Pogonatherum* and *Fimbristylis monophylla* with

their fine leaves and stems also make elegant pot plants for mingling with those of larger foliage.

The larger *Andropogons*—*A. citratus*, *A. nardus* and *A. muricatus*—are cultivated for their sweet scent; the first producing the well known lemon grass oil, the second citronella. Of other grasses of economic value, it is only necessary to mention rice (*Oryza sativa*), sugar-cane (*Saccharum officinarum*), millet (*Sorghum vulgare* and *S. saccharatum*), and the numerous species of bamboos cultivated here.

The plumed seeds of *Imperata arundinacea* are used by the Malays to stuff pillows; of the stem of *Scirpus grossus* and *Lepironia mucronata* they make mats, and walking sticks are made from *Thysanotana* in Sungei Ujong. A kind of beer was at one time made from the rhizomes of Lalang, which contain a good deal of sugar. The manufacture was not remunerative and was soon given up.

The grasses and sedges which form the greater part of the turf here are *Andropogon acicularis*, *Cynodon dactylon*, *Ischæmum ciliare*, besides which, in some places *Fimbristylis monophylla*, *F. tenera* and *F. communis* form a considerable portion. I have seen a lawn almost entirely composed of *Zoysia pungens*, a soft inland form. It makes a very deep soft turf, but is unsuitable for tennis lawns, except in dry places, where it becomes short and compact. For sandy places *Andropogon aciculare* and *Cynodon dactylon* are best. Where the soil is better the *Ischæmums* form a better class of turf. The best native fodder grass here is unquestionably *Ischæmum muticum*; *I. ciliare* and *I. timorense* are also very good. *Panicum jumentorum*, the Guinea grass, has been introduced as a fodder plant, but must be used with caution, as horses have been killed by overfeeding with it.

Distribution.—A large number of the glumaceous plants of the Peninsula are very widely distributed throughout the Indo-Malayan region. A few are cosmopolitan, occurring in all parts of the world that are warm enough for them; such are *Cyperus polystachyus*, *C. rotundus*, *C. distans*, and *Fimbristylis communis* among *Cyperaceæ*; *Cynodon dactylon*, *Digitaria sanguinalis*, *Eleusine indica* among grasses. A

small number of sea-shore plants have an Australian facies; such are *Schænus calostachyus*, *Timbristylis sericea*, *Spinifex squarrosus*, *Lepturus repens*. With them occur also such Australian forms as *Casuarina equisetifolia*, *Leucopogon*, *Dianella*, *Melaleuca* and *Philydrum* in the lowlands near the sea; and on the mountains of the interior we also meet with *Bocckia*, *Leptospermum*, *Cryptostylis*, *Corysanthes* and *Dacrydium*. All these are to be found also in the Malayan Archipelago, and a few, e. g., *Dianella*, *Cryptostylis*, *Spinifex* and the *Lepturus* as far West as Ceylon. This seems to indicate a wave of Australian immigrants westwards, some of which, such as *Casuarina* and the grasses, at least were brought by sea-currents along the island shores.

There are two grasses which are certainly of American origin, viz., *Chloris barbata* and *Paspalum conjugatum*. The latter is very common in the cultivated districts, but seems to disappear in the interior. In Penang, are two Indian grasses which do not appear in the South of the Peninsula, viz., *Thysanolanæ acarifera*, Nees, which is typically Burmese, and *Eragrostis Wightiana*, a rare grass only known in the mouth of the Hooghly and in one place in Ceylon.

Native names.—Such native names as I have been able to find I have inserted. The derivations or translations are only suggested. It is often very difficult to get at the idea of a Malay when he names a plant, so many of the names are absurd and apparently meaningless.

The common name for grass is "Rumput," but this means also almost any small plant, such as Rumput Kra-Nas, Ape's rice grass, *Vandellia crustacea*. "Senderaian" is a general name for *Cyperis* and *Sclerias*.

CYPERUS.

C. pumilus, Linn. Sandy places. Not common.

Singapore—Changi, Selitar.

Malacca—Ayer Panas. Also collected by GRIFFITH.

Penang—Dato Kramat.

Native name, Rumput Taman (Malacca).

C. polystachyus, Rottb. Very common in waste ground, sea shores, etc. Singapore—Very common everywhere.

Penang—Waterfall, Government Hill, Telok Bahang, etc.
 Sungei Ujong—Seremban.
 Pahang—Pekan, Sungei Meang.
 Kelantan—Kamposa.

Native name, Rumput Parah Betina.

Var. *laxiflora*. Usually in damper spots.

Singapore—Changi, Tanglin.

Penang—Telok Bahang.

Selangor—Bukit Kudah.

Var. *ferruginea*.

Singapore—KUBZ.

C. alopecuroides, Rottb. Damp spots by streams. Rare.

Penang—Bagan Jermal.

Kelantan—Kamposa.

C. castaneus, Willd.

Perak—Dr. KING's Collector.

These are the only specimens I have seen from the Peninsula.

C. cuspidatus, H. B. K.

Penang—Dato Kramat. This is the form *angustifolia*.

C. compressus, L. Common in waste ground.

Singapore—Changi, Tanglin, etc.

Penang—Sepoy Lines.

Malacca—Ayer Panas.

Sungei Ujong—Port Dickson.

Native name, Rumput Tiga Sari (Malacca), "three angled grass."

C. Zollingeri, Steud. Common in kampongs, orchards, etc.

Singapore—Tanglin, Changi, Fresh Water Isle.

Penang—Waterfall Hill, Government Hill, Telok Bahang.

Pahang—Pekan.

Malacca—Pulau Besar.

C. platystylis, R. Br. Was collected in Penang by WALLICH, and distributed with the number 3359D. I have not met with it.

C. Griffithii, Steud. Common on sandy shores on both coasts. This is a very variable plant, the most striking form of which is one with swollen thickened spikelets, which was described as *C. radians* by NEES, but which is certainly not specifically distinct. It is much less common than the typical *C. Griffithii* with flat spikelets, but occurs mixed with it, at Pekan. There is also a dwarf form with very short stem, having the lower spikelets almost buried in the sand, and I met also with a curious flaccid form with long grassy leaves, and the glumes almost, or indeed in some specimens quite, foliace-

6 GRASSES AND SEDGES OF THE MALAY PENINSULA.

ous. This was growing in a very damp spot where a stream entered the sand, at Sungei Meang in Pahang.

Singapore—Changi. Plentiful.

Malacca—Pulau Besar.

Penang—Telok Bahang, Batu Feringgi.

Pahang—Rumpin River, Cherating, Sungei Meang, Pekan, Kwala Pahang. Very common along the coast.

Kelantan—Kamposa.

C. haspan, L. Common in swampy spots.

Singapore—Tanglin, Tivoli, etc.

Penang—Tanjong Bunga.

Selangor—Kwala Lumpur.

Malacca—Chabau, Ayer Panas.

Pahang—Kwala Pahang.

Native names, Rumpit Sumbo (? “wick grass”), R. Bilis Jantan (Bilis is a kind of fish, “male fish grass,” probably because it grows by streams).

C. flavidus, Retz. With the preceding, but less common.

Singapore—Tanglin.

Penang—Tanjong Bunga.

C. elegans, L. (*C. diffusus*, Vahl). Woods. Fairly common.

Singapore—Bukit Timah.

Selangor—Pataling Caves, Kwala Lumpur.

Penang—Pulau Betong. Also collected here by WALLICH (No. 3474).

Pahang—Katapong near Pekan.

C. turgidulus, C. B. Clarke. Common, especially near the sea in sandy kampongs and open woods.

Singapore—Changi.

Penang—Tanjong Bunga, Waterfall.

Sungei Ujong—Gunong Brembun.

Pahang—Rumpin River, Praman, Pekan.

Siam—Bangtaphan (Dr. KEITH).

Native name, Rumpit Chukor Kerbau, “buffalo razor-grass.”

C. iria, L. A common weed in gardens and plantations.

Singapore—Tanglin, Jurong, Chan Chu Kang, Changi.

Penang—Bagan Jermal, Waterfall, Dato Kramat.

Kelantan—Kamposa.

Malacca—Ayer Panas.

Native name, Rumpit Suloh Bulahlang (Malacca). Literally, “dragon-flies torch.”

Var. micriria. This is a very small form, only an inch or so tall,

with narrow leaves, which not rarely occurs in damp spots rather speedily dried. I met with it at Toas in Singapore.

C. pulcherrimus, Willd. Damp spots.

Penang—Pulau Betong.

Selangor—Kwala Lumpur.

Pahang—Katapong.

Siam—Bangtaphan (Dr. KEITH).

C. distans, Linn. fil. A common roadside and waste ground weed here, as it is almost all over the tropics.

Singapore—Ang Mo Kio, Tanglin, and many other places.

Penang—Sepoy Lines, Dato Kramat.

Selangor—Caves, Kwala Lumpur.

Kelantan—Kamposa.

Siam—Bangtaphan (Dr. KEITH).

Malacca—Pulau Undan.

Native name, Rumput Vanggi (Malacca). Perhaps “Wangi” “scented grass.”

C. malaccensis, Lam. Muddy streams near the sea, and on the beach. Not very common.

Penang—Waterfall.

Pahang—Pekan.

Tringganu—Pulau Ketam.

Kelantan—Kamposa.

C. pilosus, Vahl. One of the commonest species, and somewhat variable in height, colouring and compactness of umbel.

Singapore—Changi, Ang Mo Kio, Tanglin, Jurong, Chan Chu Kang.

Johor—Tanah Merah Road.

Malacca—Common.

Penang—Waterfall, Tanjong Bunga.

Selangor—Kwala Lumpur near the Club, Bukit Kuda.

Sungei Ujong—Port DICKSON, Kwala Sawar.

Pahang—Pekan, Katapong.

C. procerus, Rottb. Not common.

Penang—Waterfall.

Kelantan—Kamposa.

Malacca—Chabau, Ching.

Native name, Rumput Munsiang.

C. rotundus, L. A common weed.

Singapore—Tanglin, Chan Chu Kang, &c.

Penang—Bagan Jermal, Sepoy Lines.

Pahang—Pekan.

8 GRASSES AND SEDGES OF THE MALAY PENINSULA.

- Malacca—Common in the town and elsewhere.
Siam—Bangtaphan.
- C. stoloniferus*, Retz. Muddy shores, and grassy spots near the sea. Common.
Singapore—Changi, Tanjong Ru; also Pulau Buru and Raffles Lighthouse Island.
Penang—Bagan Jermal.
Pahang—Kwala Pahang.
Kedah—Langkawi Islands.
Malacca—Shores near the town.
- C. auricomus*, Sieber. Damp spots near stream. Local.
Penang—Dato Kramat.
Selangor—Bukit Kudah, Kwala Lumpur.
Pahang—Katapong near Pekan.
Sungei Ujong—Kwala Sawar.
- C. elatus*, L. Was collected in Penang by WALLICH (No. 3341a).
I have not seen any other specimens.
- C.* (§ *Diclidium*) *ferax*, A Rich. Damp spots. Rare.
Selangor—Kwala Lumpur.
Pahang—Katapong.

MARISCUS.

- M. dilutus*, C. B. Clarke. Damp spots. Not rare.
Singapore—Bukit Timah, Selitar.
Selangor—Kwala Lumpur.
Perak—Gopeng (Calcutta Herbarium, C. B. CLARKE).
Pahang—Pekan.
Kelantan—Kamposa.
- M. pennatus*, Lam. Sea shores common, more rarely inland.
Singapore—Selitar, Bajau, Tanjong Ru.
Johor—Tana Runto.
Malacca—Ayer Panas, Alor Gajah.
Kelantan—Kamposa.
Pahang—Sungei Meang.
- Native names, Rumput Bumbot, "twisted grass," R. Surai, R. Sulengsen.
- M. biglumis*, Gaertn. Local.
Var. *cylindrostachys*.
Penang—Pulau Betong.
Sungei Ujong—Pantai.
Pahang—Pulau Tioman.
Singapore—Common.

Var. cyperina.

Penang.

Pekan—Katapong.

M. dubius, Rottb. Very common on sandy shores by the sea.

Singapore—Cathedral compound, Changi.

Penang—Tanjong Bunga.

Province Welleslev—(Dr. KING).

Malacca—Pulau Besar.

Pahang—Pekan, Rumpin River, Sungei Meang.

Kelantan—Kamposa.

Siam—Bangtaphan (Dr. KERTH).

M. umbellatus, Vahl. Very common in waste grounds, woods, etc.*Var. typica.*

Sungei Ujong—Gunong Bumbur.

Pahang—Katapong.

Var. cylindrostachys. The commonest form.

Singapore—Ang Mo Kio, Selitar, Tanglin.

Johor—Tanah Runto, Pengerang.

Malacca—Alor Gajah.

Selangor—Kwala Lumpur.

Penang—Government Hill.

Pahang—Rumpin River.

Siam—Bangtaphan (Dr. KERTH).

Var. picta. A very compact headed form with almost sessile squarrose spikes.

Sungei Ujong—Gunong Bumbur.

Siam—Bangtaphan (Dr. KERTH).

The native names are Rumput Tangot Baong "bearded umbrella grass," and Rumput Pinang "palm grass."

KYLINGA.

K. monocephala, Rottb.

Singapore—Botanic Gardens, etc.

Malacca—Ayer Panas.

Pahang—Pekan.

Native Names, Rumput Tuki, Rumput Butang, "button grass," a name also applied to *Eriocaulons* and *Rhynchospora Wallichii*.*K. brevifolia*, Rottb. A common weed.

Singapore—Almost everywhere, Selitar, Bajau, etc.

Penang—Government Hill.

Johor—Scudai River.

10 GRASSES AND SEDGES OF THE MALAY PENINSULA.

Pahang—Kwala Pahang, Pekan.
Malacca—Common.

- K. cylindrica*, Nees. Local. Not at all common.
Singapore—Raffles Institute (HULLETT).
Siam—Bangtaphan (Dr. KEITH).

FIMBRISTYLIS.

- F. nutans*, Vahl. Damp marshy spots on sandy heaths. Local.
Singapore—Chan-i.
Malacca—Pulau Besar.
Penang—Telok Bahang, Tanjong Bunga.
Pahang—Rumpin River, Kwala Pahang.

- F. schoenoides*, Vahl.

Penang—Telok Bahang, Tanjong Bunga.

- F. argentea*, Vahl. Sandy banks of rivers. Rare.
Kelantan—Kamposa.

- F. æstivalis*, Vahl. Damp sandy spots.

Singapore—Tanglin.

Pahang—Katapong, Kwala Pahang.

Native name, Rumput Surai.

- F. diphylla*, Kth. Very common.

Singapore—Changi, Tanglin, Selitar, etc.

Malacca—Pulau Besar near the town.

Sungei Ujong—Seremban.

Penang—Waterfall.

Johor—Tanah Merah Besar.

Pahang—Pekan.

Native names, Rumput Parah (Parah is a measure), R. Pürun Batu, “mat grass,” which grows in stony places.

- F. sericea*, R. Br. Sandy shores. Rare. East coast only.

Pahang—Cherating, Kwala Pahang, Pekan, Sungei Meang.

Kelantan—Kamposa.

- F. pauciflora*. (*F. malaccana*, Boeck. *F. filiformis*, Kth). Very common.

Singapore—Abundant—Tanglin, Changi, Chan Chu Kang,

Government Hill, Rochor, Pulau Tekong.

Johor—Tanah Merah Road.

Sungei Ujong.

Penang—Waterfall, etc.

Pahang—Pekan, Praman.

Native name, Rumput Girah.

- F. tenera*. Var. *obtusata*, C. B. Clarke. Not rare. In sandy open

places. Mr. CLARKE says "a very remarkable plant, which I have had several times before. It has the leaves of *F. disticha*, but the glumes are not distichous. It is perhaps a species."

Singapore—Botanic Gardens in the turf, Jurong sandy spots, Bukit Mandai roadside, Bukit Timah.

Penang—Telok Bahang.

Malacca—Sungei Baru Ulu.

Pahang—Kwala Pahang.

F. asperima, Beckler. Forms large tufts in woods. Not very common.

Singapore—Bukit Mandai (base of the hill), Botanic Gardens a weed in the flower beds).

Penang—Government Hill, Tanjong Bunga, Waterfall, Telok Bahang.

Malacca—Tanjong Kling, Bukit Bruang.

Sungei Ujong—Gunong Burumban.

Native names, Rumput Siamet. Perhaps from "Siya," "useless."

R. Bavang (Sungei Ujong). Perhaps "Bawang," "onion grass," from the resemblance of its leaves to those of an onion. R. Pulot (Sungei Ujong), "rice grass."

F. miliacea, Vahl. Common in muddy damp spots.

Singapore—Tanglin, Ang Mo Kio, Selitar, etc.

Penang—Waterfall, Tanjong Bunga.

Malacca—Ayer Panas, Chiung, etc.

Johor—Tanah Runto.

Pahang—Pekan, Kwala Pahang.

Kelantan—Kamposa.

Native name, Rumput 'tai Kerbau "buffalo dung grass" (Malacca).

This is probably so called from its coming up where buffaloes have been feeding, these animals eating it, and passing the seed which eventually germinates.

F. globulosa, Kth. Damp spots. Local.

Malacca—Bukit Sabukor, Ayer Panas.

Penang—Waterfall.

Pahang—Pekan, Kwala Pahang.

Var. *foliata*, Beckler. Malacca (Jagor).

Native name, Rumput Sandang.

F. retusa, Thwaites. Stated by BOECKLER (Cyperaceæ, p. 582) to have been gathered by GRIFFITH in Malacca. I have not met with.

F. glomerata, Nees. Sandy places usually near the sea.

12 GRASSES AND SEDGES OF THE MALAY PENINSULA.

- Singapore—Tampenis Road near Bukit Mandai, Saranggong.
Johor—Tanah Merah Road.
Pahang—Cherating, Kwala Pahang.
Malacca—Tanjong Kling.
- F. setacea*, Benth. Wet spots. Local.
Singapore—Bank of the lake in the Gardens.
Pahang—Pekan.
- F. longispica*, Steud. Sandy spots, East coast. Rare.
Pahang—Kwala Pahang, Rumpin River, Pramau.
- F. tetragona*, Br. Sandy heaths.
Pahang—Kwala Pahang.
- F. leptoclada* Benth. Sandy places.
Singapore—Bukit Timah, Changi.
Pahang—Kwala Pahang.

SCIRPUS.

- Sc. grossus*, Linn. fil. Not rare in rice-fields. Used in making mats and baskets, under the name of Rumput Musiang and R. Murong (Penang).
Malacca and Penang—In the paddy-fields. Not rare.
Pahang—Near Pekan.
Kelantan—Kamposa.
- Sc. mucronatus*, L. Common in ponds and ditches.
Singapore—Swamp near Thompson Road, Reservoir, Ang Mo Kio, Tivoli.
Malacca—Ayer Panas.
Selangor—Kwala Lumpur, Klang.
Penang—Pulau Betong.
Pahang—Ayer Etam, Pekan.
Sungei Ujong—Kwala Sawar.
- Sc. supinus*, L. Local. In swamps and sawars.
Penang—Waterfall.
Malacca—Selanda.
It is called Rumput Prut Tikus in Malacca. This is literally "mouse intestine grass."
- S. juncooides*, Roxb. In similar localities to the preceding.
Malacca—Bukit Sabukor, Bukit Tunggal.
Penang—Waterfall.

BULBOSTYLIS.

- B. barbatus*, Rottb. Common on sandy paths, sea-shores, etc. A curious proliferous form occurred at Pekan, with long slen-

der branches one inch long springing from the unusually large capitulum of flowers.

Singapore—Changi, Tanglin, etc.

Penang—Batu Feringgi.

Perak.

Pahang—Kwala Pahang, Pekan, Katapong.

Tringganu—Pulau Ketam.

Malacca—Tanjong Kling, etc.

B. puberula, Poir. Sandy places. Rare.

Singapore—No locality (CANTLEY).

Karimoon Isles.

Malacca—(GRIFFITH).

TRICOSTULARIA.

T. borneensis, Benth. Sandy heaths. Rare, but plentiful where found.

Pahang—Pekan, Praman.

FUIRENA.

F. glomerata, Lam. Common in swamps, rice-fields and ditches.

Singapore and Malacca—Common everywhere.

Penang—Waterfall, Sungei Ujong, Bukit Sulu.

Pahang—Pekan.

It is called Rumput Buku Buloh (Buku Buluh, *i.e.*, grass with nodes like a bamboo), R. Kululot (Malacca), Trigonia grass. (The Kululot is a small species of bee, genus *Trigonia*, which collects pollen from this plant. R. Lidah Munkerang (Sungei Ujong).

LIPOCARPHA.

L. argentea, R. Br. Common in damp spots, by streams in open country.

Singapore—Very common, Tanglin, Chan Chu Kang, &c.

Malacca—Common.

Sungei Ujong.

Johor—Tanah Merah Road.

It is called Rumput Rotan, "rattan grass," in Sungei Ujong.

L. microcephala, Steud. Rare. A new record for the Peninsula.

Singapore—Chan Chu Kang in dry turf, Bukit Mandai ditch by roadside.

14 GRASSES AND SEDGES OF THE MALAY PENINSULA.

LEPIRONIA.

- L. mucronata*, Rich. Local in muddy spots near mangrove swamps.
Singapore—Jurong. Tampenis Road near Changi.
Malacca—(GRIFFITH).

ELEOCHARIS.

- E. chætaria*, R. & S. Common in shallow ditches and damp spots.
An elongate floating form occurs not rarely in deeper water.
Singapore—Tanglin, Chan Chu Kang, etc.
Selangor—Kwala Lumpur.
Pahang—Pekan, Kwala Pahang.
Malacca—Tanjong Kling, Ayer Molek.
- E. capitata*, R. Br. Rare.
Singapore—Changi, Tampenis. Plentiful in ditches by the roadside.
- E. variegata*, Var. *laxiflora*. Common in ditches, ponds, etc.
Singapore—Tanglin, Reservoir.
Penang—Telok Bahang, Waterfall.
Pahang—Kwala Pahang.
Malacca—Near Ayer Molek.
- E. ochrostachys*, Steud. Fairly common. Very closely resembling the preceding one.
Singapore—Near the Garden lake.
Malacca—Merlimau.
Pahang—Katapong near Pekan.
- E. equisetina*, Presl. Rare.
Penang—Stone quarry near the Waterfall.

MAPANIA.

- M. longa*. Benth. Rare. A large tufted plant growing in streams, resembling a *Sparganium* in habit.
Singapore—Chan Chu Kang.
- M. humilis*, Naves and Villar. An elegant plant for pot cultivation, the leaves being somewhat like those of a *Pandanus*, and purple. Common in jungle.
Singapore—Kranji, Bukit Timah, Bajau, Chan Chu Kang.
Perak—Maxwell's Hill.
Malacca—Selandar.
- Called Poko Seak Seak Rimbah. Perhaps this should be "Poko Siya Siya Rimbah," *i.e.*, useless jungle plant. It is used for fever.

- M. bancana*, Benth and Hook. A common woodland plant growing in damp spots in the jungles.
Singapore—Changi, Ang Mo Kio, Kranji, Toas, Selitar.
Malacca.
Penang—Waterfall Hill.
Sungei Ujong—Gunong Brumbun.
Selangor—Pataling.
Pahang—Pekan.
It is known as Rumpit Supidang, R. Giring Giring, "rattle grass," R. Susat Balukar (Sungei Ujong).
- M. palustris*, Benth and Hook. Forms large tufts in thick jungle.
Singapore—Pulau Ubin, Chan Chu Kang.
It is called Poko Mengkuang Tudong and P. Menkuang Lubo, from its resemblance to the Mengkuang (*Pandanus furcatus*).
- M. multispicata*, C. B. Clarke.
Singapore—Bukit Timah.

SCIRPODENDRON.

- Sc. costatum*, Kurz.
Singapore—North Selitar and Changi. I believe this very curious plant is not at all uncommon, but it resembles so much a young *Pandanus* that it is easily overlooked. It is a native of Ceylon and Java, and this new locality forms a connecting link between these two regions.

HYPOLYTRUM.

- H. proliferum*, Boeck. Local. In wet jungle.
Singapore—Selitar, Chan Chu Kang.
New to British India.
- H. latifolium*, Rich. In wet jungle.
Singapore—Chan Chu Kang, Jurong.
Pahang—Ayer Etam.

REMIREA.

- R. maritima*, L. Common on most of the sandy shores.
Singapore—Changi near the Police Station.
Malacca—Tanjong Kling.
Penang—Batu Feringgi.
Pahang—Cherating, Sungei Meang, Kwala Pahang.
Kedah—Langkawi Islands.
Siam—Bangtaphan (Dr. KEITH).

RHYNCHOSPORA.

- R. aurea*, Vahl. Very common in damp spots.
Singapore and Malacca—Everywhere.
Selangor—Batu Tiga, Kwala Lumpur.
Pahang—Pekan.
Sungei Ujong—Common, Kuala Sawar.
- R. Wallichiana*, Kunth. Not rare. In dry open country among
bracken andalang.
Singapore—Jurong, Selitar, Bajau, etc., Sungei Murai. Fresh
Water Island.
Malacca—Pulau Besar, Bukit Tunggal. Also collected by
GRIFFITH.
Pahang—Kwala Pahang.
It is called Rumput Butang ("button grass") in Singapore.
Forma sylvestris.
Leaves very slender and grassy; heads very small and com-
pact, usually green and not red. Damp woods.
Karimon Isles.
Malacca—Mount Ophir (R. DERRY).
Penang—Government Hill.
- R. glauca*, Vahl. Var. *chinensis*. Damp sandy spots. Rare.
Singapore—Changi near Tanah Merah, abundant.
- R. malasica*, C. B. Clarke. Very local.
Singapore—Bukit Mandai, growing in streams. I have only
found this curious plant in this one spot, but it is very plenti-
ful here. Mr. CLARKE says he has also seen it from Malacca
and Borneo.

CLADIUM.

- Cl. Maingayii*, C. B. Clarke.
Malacca—Mount Ophir (P. W. HULLETT).
- Cl. glomeratum*, Nees. Rare. In water.
Singapore—Bukit Mandai, Changi towards Tanah Merah.
"The first specimens I have seen from Malaya," says Mr. CLARKE.
"It is a common Australian plant, also known from China
and Japan."

SCHÆNUS.

- Sch. calostachyus*, Poir. Sandy places near the sea. Rare.
Singapore—Changi near Tanah Merah. Abundant.
Pahang—Praman near Pekan.
Mr. CLARKE says:—"An interesting though apparently wide-

spread plant. I have it from various parts of Australia, Louisiade Archipelago, Waigiou and Borneo, but not before from British India."

GAHNIA.

- G. tristis*, Nees. Sea shores, more rarely from inland localities. Rather common, but not to be met with everywhere. Usually in rocky spots above the sea. There is a weaker form which occurs in woods in the interior, with very much slenderer leaves.
- Singapore—Forts Siloso and Serapong, Upper Mandai, Tampenis Road near Changi. Pulau Battam.
- Karimon Isles.
- Selangor—Seppan.
- Johor—Scudai River, Tanah Runto.
- G. javanica*, Zoll. Mountains at a considerable altitude.
- Perak—Gunong Batu Puteh at 6,700 feet altitude (Wray).
- G. sp.*
- Perak—Maxwell's Hill (Curtis).

SCLERIA.

This genus, which is rather a difficult one, is well represented here, but I am sure that a number more will be found when carefully sought.

- Scl. caricina*. *Diplacrum caricinum*, R. Br. Common in waste places, patis and damp spots.
- Singapore—Tanglin, Tivoli, Changi. Common.
- Malacca—Merlimau, Pulau Besar.
- Pahang—Kwala Pahang, Pekan.
- Scl. lithosperma*, Nees. Common in dry country.
- Malacca—Alor Gajah.
- Penang—Waterfall Garden.
- Pahang—Rumpin River, Kwala Pahang.
- Sungei Ujong—Linggi River.
- Scl. lateriflora*, Bekler. Not common.
- Singapore—Changi.
- Penang—Telok Bahang.
- Scl. stuedeliana*, Miq. Rare.
- Penang—Waterfall.
- Scl. hebecarpa*, Nees. Common in open country.
- Singapore—Jurong.
- Penang—Tanjong Bunga.

- Malacca—(GRIFFITH).
 Pahang—Pekan.
 Siam—Bangtaphan (Dr. KEITH).
- ScL. oryzoides*, Presl. River banks. Local, but plentiful where it occurs.
 Singapore—Changi towards Tanah Merah.
 Malacca—Banks of Merlimau River, and also near Malacca, Ching.
- Native name, Rumput Leeku Dana. Possibly an error for "Lichu Daun," polishing leaves, *i.e.*, used for polishing wood.
- ScL. malaccensis*, Bekler. Very common, often covering large tracts of wet ground.
 Singapore—Everywhere, Tanglin, Jurong, Bukit Timah.
 Malacca—Common.
 Selangor—Kwala Lumpur.
 Pahang—Pekan.
- ScL. multifoliata*, Bekler. In thickets, scrambling through bushes.
 Malacca—Pulau Besar.
 Pahang—Pekan.
- ScL. laevis*, Willd.
 Johor—Tanah Merah Road.
 Pahang—Kwala Pahang.
- ScL. sumatrensis*, Retz. Common in dry open places.
 Singapore—Tanglin, Bukit Timah, Jurong.
 Penang—Government Hill, Tanjong Bunga.
 Malacca—Alor Gajah, and other places.
 Johor—Tanah Merah Road.
 Sungei Ujong—Gunong Brumbun.
 Selangor—Bukit Kuda, Klang.
 Perak.
 Pahang—Kwala Pahang.
 Siam—Bangtaphan (Dr. KEITH).
- It is called Rumput Kumba in Sungei Ujong. "Kumba" is an ornament made of coco-nut leaves, which is perhaps the origin of the name.
- ScL. chinensis*, Kth. New to this country.
 Pahang—Kwala Brawas, Pekan.
- ScL. androgyna*, Nees. Collected by GRIFFITH in Malacca. I have not met with.

CAREX.

- C. cryptostachys*, Brngn. Wet rocky places.

Singapore—Bukit Timah.
 Penang—Government Hill.
 Perak—Taiping.
 Sungei Ujong.

Native name, Rumput Ringgin (Sungei Ujong).

C. sp.

Penang Hill, rocky places on the summit.

GRAMINEÆ.

PASPALUM.

P. scrobiculatum, Linn. Common everywhere, and apparently native.

Singapore—All over the island.
 Johor—Tanah Merah.
 Malacca—Ayer Panas.
 Selangor—Kwala Lumpur.
 Penang—Waterfall Garden.
 Sungei Ujong.
 Pahang—Pekan, Kwala Pahang, etc.

Native names, Rumput Tulo Sintadok, "caterpillar grass," R. Liku, "smooth grass" (Malacca), R. Ijok, "green grass" (Sungei Ujong).

P. distichum, Burm. In sandy mud near the seashore. Common.

Singapore—Tanjong Karong, Bajau, Sungei Murai.
 Johor—Tanah Merah Road, Tanjong Bunga.
 Penang—Bagan Jermal.
 Pahang—Sungei Meang.

P. conjugatum, Berg. Native of South America, but now widely distributed. It chiefly occurs along paths and in waste places near villages, being transported by its very adherent fruits, but in the wilder parts of the country does not occur. It forms a good fodder grass.

Singapore—Common everywhere.
 Johor—Shores of the strait near Johor.
 Malacca—Common.
 Penang—Waterfall Garden.
 Pahang—Pekan.
 Siam—Bangtaphan (Dr. KEITH).
 Sungei Ujong—Common.

ISACHNE.

I. australis, R. Br. Common in damp swampy spots.

- Singapore—Tanglin, Changi.
 Johor—Summit of Mount Ophir (HULLETT), Pengerang.
 Pahang—Pekan, Katapong.
 Penang—Waterfall Garden.
 Malacca—Ayer Panas, common.
- I. pulchella*, Roth. Common in rice fields and wet spots.
 Singapore.—Chan Chu Kang, Ang Mo Kio, Kranji.
 Malacca—Bukit Tunggal.
 Penang.
- Ophismenus burmanni*, Retz. Not common.
 Penang—Sepoy Lines.
- O. compositus*, Wight. Shady woods. Widely distributed, but not very common.
 Malacca—Bukit Tunggal.
 Pahang—Kwala Pahang.
 Pulau Buru, near Karimon Islands.

PANICUM.

- P. (Digitaria) sanguinale*, Trin. Common in waste ground.
 Singapore—Botanic Gardens, Changi.
 Penang—Waterfall.
 Pahang—Rumpin River, Sungei Meang.
- Var. *australe*. Occurs in Singapore, and Penang near the Waterfall.
- P. parvulum*, Nees. Common weed.
 Singapore—Botanic Gardens, Changi, Jurong, Raffles Light-house.
 Penang—Waterfall.
 Pahang—Sungei Meang.
- P. volascens*, Nees. Rare.
 Singapore—Bajau. A form with glabrous spikelets.
- P. heteranthum*, Nees. Rare. Sandy shores, beneath the *casuarinas*.
 Pahang—Rumpin River, Sungei Meang.
- P. cimicinum*, Retz. Weed near villages and old cultivations.
 Singapore—Tanglin, Tivoli, Changi, Chan Chu Kang.
 Penang—Waterfall, Tanjung Bunga.
- P. (Brachiuria) nodosum*, Kth. *P. multinode* Presl. Common in thickets and woods.
 Singapore—Tanglin, Dalvey Road, etc., Chan Chu Kang.
 Freshwater Isle.
 Penang—Waterfall, Government Hill, Pulau Betong.

Pahang—Pekan.

Kelantan—Kamposa.

Native name, Sarong Buaya (Penang). Perhaps this should be "Sarang," *i.e.*, crocodile's nest.

P. subquadriparum, Trin. Rare.

Singapore—Changi.

P. luzonense, Presl. Not very rare. Chiefly in new cleared ground and roadsides.

Penang—Waterfall.

Malacca—Alor Gajah.

Pahang—Katapong.

Perak.

P. trigonum. Shady woods. Common.

Malacca—Pulau Besar, Sungei Udang.

Penang—Sepoy Lines.

Pahang—Pekan, Kwala Pahang.

Sungei Ujong—Bukit Sulu.

Native name, Rumput Mutubong, R. Kurubong Padi (Sungei Ujong). Perhaps grass which surrounds the rice, *i.e.*, weed in the rice fields.

P. radicans, Retz. Shady woods. Common.

Singapore—Bukit Timah, Changi, Tanglin.

Penang—Government Hill.

Pahang—Pekan.

Selangor—Klang.

Malacca—Alor Gajah, Ayer Molek, etc.

Perak.

Sungei Ujong—Port Dickson.

Native name, Rumput Telor Ikan ("fish-egg grass"), from the resemblance of the small black spikelets to fish eggs.

P. ovalifolium, Poir. Rare. Shady woods.

Pahang—Kwala Pahang.

Penang—Government Hill.

Malacca—Bukit Bruang.

P. ischæmoides, Retz. Sandy spots. Local.

Singapore—Changi.

Pahang—Pekan.

Kelantan—Kamposa.

Penang—Waterfall.

Malacca—Ching.

P. concinnum, Nees. A very handsome grass not rare in woods.

Singapore—Sumbawang, Tanglin.

22 GRASSES AND SEDGES OF THE MALAY PENINSULA.

Malacca—Selandor. Bukit Sadanau.

Penang—Moniot's Road.

Perak.

Selangor—Klang.

Called Rumput Jangot Ali and Akar Tongkat Ali. *i.e.*, "Ali's beard grass," "Ali's walking-stick root." The roots are eaten by Malays in Malacca along with betel-nuts. It is also called Rumput Poko Kulubong.

P. incomptum, Trin.

Perak—Gunong Hijau (WRAY).

P. jumentorum. Is cultivated here and there for fodder.

P. (Hymenachne) auritum, Presl. Pools and ditches. Not rare.

Singapore—In the Reservoir, Changi, Ang Mo Kio.

Penang—Waterfall, Tanjong Bunga.

Pahang—Pekan, Mahang.

Siam—Bangtaphan (Dr. KEITH).

Malacca—Stream near Alor Gajah.

P. myurus, Meyer. Ditches. Not very common.

Malacca—Ayer Panas, Ching.

Penang—Near the coast.

Sungei Ujong—Burunang, Kwala Sawa, where it is called Rumput Kumpai, and used for making lamp-wicks.

P. myosuroides, R. Br. Common in ditches.

Singapore—Ang Mo Kio, Kranji.

Malacca—Chabau.

Pahang—Pekan.

Penang—Pulau Betong.

Native name, Rumput Kumani.

P. indicum, L. Very common. Dry waste places.

Singapore—Bukit Timah, Changi, etc.

Johor—Tanah Merah.

Pahang—Praman.

Penang—Waterfall.

Malacca—Selandar, etc.

Native names, Rumput Bidis, R. Bonto Darat.

P. Ridleyi, Hackel. Dry thickets.

Malacca—Pulau Besar, Bukit Bruang.

Pahang—Pekan.

Near *P. foliosum* and *zizanioides*; with the latter General MONRO classed it. It was also collected by GRIFFITH in Malacca.

P. plicatum. Rare.

Selangorv—Caes, Kwala Lumpur.

Sungei Ujong—Pantai.

P. (Echinochloa) crus-galli, L. Not common.

Singapore.

Penang—Waterfall, Bagan Jermal.

Kelantan—Kamposa.

Selangor—Bukit Kudah.

P. colonum, L. Common in waste grounds.

Singapore—Tanglin, Changi.

Penang—Near the coast.

Malacca—Near the town.

SETARIA.

S. glauca, Beauv. Open country. Not very common.

Singapore—Ang Mo Kio, Tanglin.

Penang—Abundant in paddy-fields.

Malacca—Chenana Puteh.

Pahang—Pekan.

Native name, Rumput Julong-Julong. Perhaps "julung-julung," waving grass.

S. dasyura. Waste places, open country.

Singapore—Chan Chu Kang.

Pahang—Pekan.

PENNISSETUM.

P. italicum, L. Millet, a plant rarely cultivated here, has turned up now and again as a waif of cultivation, Malacca.

THUAREA.

Th. sarmentosum, Thouars. In sand on the sea shore. Rare.

Singapore—Changi near the Police Station.

Pahang—Common under the casuarinas on the shore at Rumpin River, Cherating and Sungei Meang.

SPINIFEX.

S. squarrosus, Labill. Porcupine grass. Sandy shores of the East Coast. Plentiful in places.

Siam—Bangtaphan (Dr. KEITH).

Pahang—Rumpin River, Cherating, Sungei Meang, Kwala Pahang.

It is very curious to see the detached bristling heads of this grass whirled along the sands by the wind.

LEPTASPIS.

L. urceolata. Damp woods. Local, but very widely distributed.

24 GRASSES AND SEDGES OF THE MALAY PENINSULA.

The spikelets are very adhesive, and by adhering to animals get carried about the jungle.

Singapore—Pulau Ubin, Chan Chu Kang.

Johor—Summit of Mount Ophir (HULLETT).

Malacca—Telandar, Ayer Panas, Bukit Sadanen, etc.

Selangor—Kwala Lumpur, Bukit Kudah, Bukit Etam (KEL-SALL).

Perak—Tapa (WRAY).

Sungei Ujong—Gaong Jalan.

It is called Tapa Kulang or Rulang; Tampo Glang and Getah Puyuh (Malacca), "pigeon birdlime," because the spikelets adhere to pigeons.

COIX.

C. lachryma-Jobi, L. Job's tears. Is apparently not native. It occurs commonly near villages, as in Singapore at Selitar; near Johor town; Penang, Bukit Tumujang; Sungei Ujong, etc.

The native name is Mulai Tikus.

ZEA.

Z. mays, L. Maize is cultivated, but to a small extent, and chiefly used as a vegetable. It is more extensively grown in Kelantan.

ORYZA.

O. sativa, L. Commonly cultivated, and often occurring as an escape.

Native name, Padi Pulot.

O. sativa, Var.? A very curious plant, with broad green leaves and thin flower spikes, the spikelets falling off very readily. Grows in damp thickets and muddy spots by the stream at Ayer Etam in Pekan, and also nearer to the town, in the open country at the back. It is perhaps a form of the common rice, gone wild, or possibly a distinct species. It is very distinct and dissimilar to the common form.

LEERSIA.

L. hexandra, Sw. Swampy ground, ditches, etc. Common.

Singapore—Tanglin, Changi, etc.

Penang—Waterfall.

Pahang—Pekan.

Malacca—Rice-fields.

THYSANOLÆNA.

Th. acarifera, Nees. Woods. Rare.

Penang Hill.

Sungei Ujong.

It is called Buluh Tubarau, and used for making walking canes in Sungei Ujong. It is an elegant bamboo-like grass, forming thick tufts about six feet tall.

PEROTIS.

P. latifolia, Br. Sandy places usually near the sea. Not rare.

Singapore—Changi, Cathedral compound.

Karimon Islands.

Penang—Tanjong Bunga, Telok Bahang.

Pahang—Kwala Pahang, Pekan. Common.

Siam—Bangtaphan (Dr. KEITH).

ZOYSIA.

Z. pungens, L. Sandy ground. Common, especially near the sea.

On the sea-shore it becomes very stiff and wiry. In damper ground it forms a very deep soft turf, but is not suited for lawns, except in sandy spots, where it makes a short compact turf.

Singapore—Common. Changi, Tanglin, Tanjong Ru, etc.

Malacca—Common, in and around the town.

Pahang—Pekan.

Tringganu—Sea shore.

IMPERATA.

I. exaltata, Retz. Rather a handsome lalang, larger and with a more branching spike than the common one. Rare, in thickets.

Singapore—Chan Chu Kang, Toas, Bajau.

Selangor—Pataling near Kwala Lumpur, and Langat.

Sungei Ujong—Along the rail banks between Port Dickson and Kwala Sawar.

It is called Lalang Jawa, *i.e.*, Javanese lalang.

I. cylindrica, Cyr. Far too common in the Peninsula wherever cultivation has spread, and often covering large tracts of country. Its growth after burning of the forests is remarkably rapid, as it propagates itself by its feathery seeds, as well as by its

rhizome. A large tract covered with the plant in fruit looks from a distance as if it were covered with snow. It seems to prefer hot and dry places, but dislikes wet or very sandy soil. In the first, its place is taken by *Scleria malaccensis* and other such plants, in the latter by *Bracken* (*Pteris aquilina*, L.) or *Gleichenia*. It is a grass of very little use, as cattle and horses do not care for it on account of its hardness. Pigs, however, are very fond of the young shoots. Attempts have been made to use it for paper making, but without much success, and a kind of beer is sometimes made from the rhizomes. The plumed seeds are sometimes used to stuff pillows. It is very common in Singapore, Johor, Penang, Malacca, Province Wellesley and Selangor, but seems to be rarer in the less cultivated districts, and notably in Tringganu, Pahang and Kelantan.

SACCHARUM.

S. arundinaceum, Retz. River banks. Local. A very ornamental grass. In good wet localities it grows to a height of 12 feet or more, with very large panicles of flowers.

Selangor—Kwala Lumpur near the Caves.

Pahang—Ayer Etam near Pekan, forming large thickets.

S. officinarum, L. Sugar-cane is cultivated chiefly in Province Wellesley and Perak.

S. Ridleyi, Hackel. Banks of the river at Pekan and for some way up above, forming large tufts and covering a large tract of flat country on the right bank. It has narrow leaves and tall spikes of purple flowers. Professor HACKEL remarks that it is a very interesting plant belonging to the section *Miscanthus*, of which only one species (*S. fuscum*, Griff.), an Indian plant, was previously known.

Pollinia ciliata, Linn. Forming large clumps on the river bank. Rare.

Pahang—At Renchong above Pekan.

P. Ridleyi, Hackel. On sandy shores very close to the sea. Rare, but plentiful where it occurs.

Pahang—Rumpin River, Sungei Meang.

Allied to *P. Cumingiana*, Steud., a Philippines plant.

POGONATHERUM.

P. polystachya, R. Br. Often cultivated in Singapore as a pot plant. Not common.

Penang—Rocks by the Waterfall.

Pahang—Banks of the river at **Benchong**.
 Siam—Klingtoi River, **Bangtaphan** (Dr. **KEITH**).

DIMERIA.

- D. ornithopoda**, Trin. Paths and dry open ground. Not rare in Singapore.
 Paths in the Garden jungle, and in the grass plots, **Tivoli**, **Selitar**.
 Var. *subramosa*, sub-var. *imperfecta*, **Changi**.

ROTTBOELLIA.

- R. glandulosa**, Trin. Not rare on the edges of woods, and in thickets.
 Singapore.
 Pulau Buru near **Karimon Isles**.
 Selangor—Caves, **Kwala Lumpur**, **Bukit Kudah**.
 Pahang—**Pekan**, **Rumpin River**.
R. geminata, Hackel. Open country in **Pekan** and **Rumpin River**.
 Rare.
 This is a new and curious species, which has not as yet been found elsewhere. It was described from these specimens by Professor **HACKEL**.
R. sp. **Changi**.

ISCHEMUM.

- I. muticum**, L. One of our commonest grasses, forming a good turf and a good fodder grass. Very variable in height and form. In long grass or fern, it becomes drawn up and tall, in open dry spots, more prostrate, shorter and stiffer, creeping widely. Singapore, **Johor**, **Penang**, **Malacca**, **Sungei Ujong**. Very common.
 Pahang—**Pekan**, **Sungei Meang**.
 Perak—**Changkat Jerim** (**WRAY**).
 Siam—**Bangtaphan** (Dr. **KEITH**).
 The natives call it **Rumput Trutoos**, **R. Tamaga**, and consider its leaves a cure for headache.
I. ciliare, Retz. Common on roadsides in Singapore, and probably elsewhere in the Peninsula. I have gathered it at **Kranji**, **Bukit Timah** and **Ang Mo Kio**.
I. timorense, Kunth. Roadsides.
 Singapore—**Kranji**, **Tanglin**.
 Penang.
I. aristatum, L. Dry open country. Not common.

Singapore—Blakang Mati.
 Sungei Ujong—Without locality.

Malacca—Near Ayer Molek.

Var. *submuticum*, Hack. A variety with the very small awn enclosed in the glumes. Fresh Water Island.

This plant is called Rumput Ekor Chari.

ANDROPOGON.

A. contortus, L. Sandy turf near the sea shore.

Singapore—Changi near the Police Station.

Malacca—Pulau Besar.

Penang—Waterfall Gardens, Batu Feringgi.

Pahang—Near the Sultan's tombs, Kwala Pahang.

A. aristulatus, Hochst. Sandy shores. Rare.

Siam—Bangtaphan (Dr. KEITH).

Pahang—Kwala Pahang, Pekan.

A. aciculatus, Retz. Very common and known in the Straits as "love grass." It forms a good turf, and in Pahang covers large tracts of country.

Singapore, Johor, Malacca.—Abundant.

Penang—Waterfall Gardens.

Pahang—Pekan, etc.

A. intermedius, R. Br. The Straits form is, Professor HACKEL says, a laxer form than the Australian type. Not common.

Malacca—Ayer Panas.

Pahang—Pekan.

The natives call it Rumput Pipit, "sparrow grass."

A. Schoenanthus, L. and *A. nardus*, L. Often occur as escapes from cultivation, but are not wild here.

A. squarrosus, L. Possibly native, but often cultivated. I have seen it half wild at Bajau in Singapore, and Mr. CURTIS sends it from Tanjong Tokong in Penang.

SORGHUM.

S. vulgare, L., and the variety *saccharatum* are sometimes cultivated.

THEMEDA.

Th. arguens, Hack. Roadsides. Not common.

Malacca—Selandar.

Penang—Telok Bahang, Waterfall Gardens.

It is called in Malacca Rumput Sarang Pipit, "bird's nest

- grass," as the birds use its spikelets for building their nests.
Th. gigantea, var. *villosa*. Common. A very tall showy grass forming big tufts, and throwing up very tall panicles.
 Singapore—Fort Siloso (said to have been introduced).
 Johor—Tanah Merah Road near Johor Bharu.
 Malacca—Pulau Besar.
 Selangor—Kwala Lumpur, by the railway and near the Caves.
 Penang—Bukit Mertajam.
 Pahang—Pekan.
 Sungei Ujong—Near Kwala Sawar.

APLUDA.

- A. varia*, Hack. Not common. Pastures and river banks.
 Pahang—Katapong, Renchong.

SPOROBOLUS.

- S. diander*, Beauv. Dry spots.
 Singapore—Raffles Lighthouse.
 Malacca—Ayer Panas.
 Johor—Tana Runto.
 Native name, Rumpul Tulo Bulalang, *i.e.*, the grass on which the dragon-fly balances itself.
S. elongatus, Beauv. Dry heaths.
 Johor—Tanah Runto. Fresh Water Island.
 Penang—Sepoy Lines.
 Pahang—Pekan.
 Karimon Islands.

ERIACHNE.

- E. chinensis*, Benth. Dry, open, sandy spots. Common where it occurs, but local.
 Singapore—Fort Siloso.
 Pahang—Pekan.
E. trisetia, Nees. Dry, sandy heaths. Rare.
 Pahang—Pekan and Kwala Pahang, abundant.
 Malacca—Pangkalan Balau.

CYNODON.

- C. dactylon*, L. Rather common in sandy and dry spots.
 Singapore—Common.
 Penang—Sepoy Lines, Waterfall.
 Malacca—In the town.

CHLORIS.

- C. barbata*, Probably introduced from the West Indies.
 Johor—Near the Sawmills.
 Province Wellesley—Batu Kawan.

ELEUSINE.

- E. indica*, L. Very common in waste grounds near villages.
 Singapore—Tanglin, Changi.
 Johor—Tanah Runto, Johor Bharu.
 Penang—Waterfall Gardens.
 Perak—Tapa (WRAY).
 Sungei Ujong—Port Dickson, with a form approaching the cultivated *E. coracana*.
E. ægyptiaca, L. Not very common. On the sea shore.
 Singapore—Changi.
 Pahang—Kwala Pahang, Sungei Meang.

LEPTOCHLOA.

- L. sp.*
 Penang—Bagan Jemal.
L. chinensis, Nees.
 Kelantan—Banks of River Kamposa.

ARUNDO.

- A. donax*, L. This, a native of Southern Europe, is often cultivated here, especially a form with variegated leaves.

PHRAGMITES.

- P. Roxburghii*, Kth. The common reed. Banks of streams and damp spots. Not rare.
 Singapore—Fresh Water Isle.
 Selangor—Kwala Lumpur, Pataling, all along the river.
 Malacca—Merlimau, Selandar.
 Penang—Western Ayer Etam.
 Pahang—Pekan River bank.
 Native name, Rumput Mata Burong Puding, "variegated bird's-eye grass."

CENTOTHECA.

- C. lappacea*, Beauv. Very common in woods and along paths.
 Carried about by its adhesive spikelets.
 Singapore—Both the hispid and glabrous varieties common

over the whole island.

Johor—Tanjong Bunga, etc.

Penang—Government Hill, Tanjong Bunga.

Malacca—Ayer Panas, Pulau Besar, etc.

Pahang—Kwala Pahang.

Nagri Sembilan—Bukit Tumiang.

Native names, Rumput Silit Kain. Perhaps the grass that damages the clothes, *i.e.*, by adhering to them. R. Darah, "blood grass."

LOPHOTHERUM.

L. gracile, Brngn. Woods. Rather local.

Singapore—Chan Chu Kang.

Malacca—Chabau.

Sungei Ujong.

Perak—Birch's Hill, Larut (WRAY), Taiping (HERVEY).

Native names, Rumput Kururut, R. Jarang, "spreading grass,"
R. Klurat.

ERAGROSTIS.

E. pilosa, L. Weed in cultivated ground.

Singapore.

Penang.

E. plumosa, Retz. A garden weed. Very common.

Singapore—Abundant.

Penang, Dato Kramat, the Fort.

Malacca—Ayer Panas, etc.

Pahang—Pekan, Katapong.

Kelantan—Kamposa.

E. Brownii, Kth. Dry spots. Widely scattered.

Karimon Islands.

Penang—Government Hill, Waterfall.

Kelantan—Kamposa.

Pahang—Kwala Pahang.

Malacca—Common.

E. zeylanica, Br. Damp, sandy spots.

Singapore—(N. CANTLEY).

Pahang—Kwala Pahang.

E. ferruginea, Thunb. Sandy spots.

Singapore—Changi, Bukit Timah.

Pahang—Pramau, Pekan, Katapong, common.

E. unioloides, Nees. Very common in dry places.

Singapore—Tanglin, Changi, Jurong.

32 GRASSES AND SEDGES OF THE MALAY PENINSULA.

Johor—Scudai River, Johor Bharu.
Malacca—Tanjong Kling, Ayer Panas.
Penang—Waterfall.
Pahang—Pekan.
Kelantan—Kamposa.

- E. Wightiana*, Nees. Very rare.
Penang—Near the coast.
E. nutans, Retz. Dry, sandy heaths. Not common.
Pahang—Pekan.
Siam—Bangtaphan (Dr. KEITH).

LEPTURUS.

- L. repens*, R. Br. Sands of the sea shore. Rather rare.
Pahang—Sungei Meang, Rumpin River.

BAMBUSACEÆ.

There are a good number of species of bamboos to be met with in a wild state in the Peninsula, but very few of them have been collected. It is nearly impossible to identify a bamboo out of flower, and, as is well known, it is by no means common to find plants in flower. So that it is almost a rule among tropical botanists that whenever a bamboo is seen to be in flower specimens must be collected. There are large extents of bamboos in many parts of the Peninsula, but of what species is not yet known. I met with a large patch of a species of *Dendrocalamus* in Bukit Kuda near Klang, of which every joint was full of most excellent water, even the twigs being full though the ground beneath was remarkably dry. Several species of bamboo are cultivated, some for ordinary use, others as hedges, or for the edible shoots, especially *Bambusa nana*, the hedge bamboo.

GIGANTOCHLOA.

- G. verticillata*, Monr. Common and often cultivated. I found it in flower at Tivoli in Singapore.
G. heterostachya, Murr.
Malacca—Ayer Panas (GRIFFITH).
I have not seen this species.

MELOCANNA.

- M. Blumei*, Nees. Singapore—Roadside at Selitar. The flowering glumes, HACKEL says, are larger than the type. It is perhaps

a variety. It is not a large species, about 12 or 14 feet tall, with rather broad dark green leaves.

M. gracilis, Kurz.

Singapore—(WALLICH 5,032).

I have not seen this. It appears to be a small plant about 10 feet tall.

DENDROCALAMUS.

D. strictus, Nees. MUNRO, in the Monograph of Bamboos, mentions having received this from Singapore from MONTGOMERIE, and says that it is a variety with very large whorls of the inflorescence. I do not think the species is anywhere wild in Singapore, but perhaps his specimens came from a garden. It is the well-known male bamboo so much in request for spear handles.

D. flagellifer, Munro. Common and probably native, but it is most frequently seen near villages, in Singapore at least.

Singapore—Jurong.

Malacca—Bukit Sabukor.

Siam—Bangtaphan, extremely common (Dr. KEITH).

D. giganteus, Munro. *Bambusa gigantea*, Wallich. WALLICH obtained his specimens from Penang, but whether it was wild there, or cultivated, does not appear. I have not seen or heard of it from there. It is stated to be indigenous to Malacca, and Mr. DERRY told me he thinks he has found the plant in Bukit Sadanen, whence he sent cuttings to the Botanic Gardens. The magnificent plants in Peradeniya Gardens, Ceylon, are well known to every visitor there. It has been often introduced into Singapore, but does not grow at all to its normal size, and indeed appears not to thrive at much less than 1,000 feet altitude.

BAMBUSA.

B. nana, Thw.

The hedge bamboo most cultivated, supposed to be a native of China.

OUTLINE OF THE HISTORY OF THE DINDINGS
FROM THE 17TH CENTURY TO THE
PRESENT TIME.

BY

E. M. MEREWETHER.



IN the following brief account of the Dindings, I do not pretend to place before the readers of this Journal much that is new. I have merely endeavoured to put together in a connected form as much of the History of the Dindings as I have been able to gather from the sources at my disposal, in the hope that it may be of some interest to those who only know the Dindings by name. For the History of the Dutch Occupation of Pangkor, I am indebted to Mr. W. E. MAXWELL'S article on the Dutch in Perak, published in Vol. No. 10 of this Journal, and to his Note on the same subject in Vol. No. 11.

The earliest mention of the Dindings is found in a letter from the Governor-General and Board of Administration of the United East India Company, dated 1st October, 1661, discovered among the Dutch records in Malacca, in which an order is given to cut 200 pieces of a certain red wood at Pulau Dinding (Pangkor) to be sent to "Patria" (Holland). From this it may be assumed that the Island became known to the Dutch before that date, probably when they established their factory in Perak in 1650, which was cut off in 1651. 1661.

Two years later, on the 29th November, 1663, an old Dutch navigator named WOUTER SCHOUTEN visited Pulau Dinding on his way to Bengal, and wrote an account of his voyage, from which I take the following extracts:— 1663.

"On the 25th November in the evening sighted Malacka for the second time, and advanced four miles with the land

wind; then anchored, waiting for daybreak; weather now lovely. Sunshine and a temperate sky with a following breeze with which we set sail and passed the green, rocky mountains of Cape Ressados (Rachado?), steering now South-West for some hours, and then North to fetch above the dangerous reef of Poelo Parselar. Coming under the green coast of the kingdom of Pera, we sighted the Poelo Sambilan or Nine Islands, which having passed, we headed for the island of Dinding, and arrived on the 29th November in the Roadstead between that Island and the mainland of Pera, close to the Watering-place.....

“ Having reached the neighbourhood of the Watering-place on the inner side of the Island Dinding above-mentioned, we immediately sent a good party of sailors to the coast of Pera opposite to procure firewood for our further voyage to Bengal. The others went to Poelo Dinding to fetch fresh water from one of the principal Rivers of the island, and we, not to be idle, went also on shore with a line of 80 fathoms and brought up fish out of the Gulfs and Bays of the Island Dinding, going on board in the evening with a good haul of all sorts of well-flavoured, delicate fish.

“ In the same way, on the next day, the 30th November, our people still being engaged in fetching water and firewood, we roamed all about and visited all parts of the Island Dinding, taking at last a good haul; we remained on shore all night with our Sub-Merchant ABRAHAM DE WIJS and others in the same way inclined and there we enjoyed our catch.....

“ The Island Dinding, about 30 miles to the North-West of Malacka, is uninhabited, full of high Mountains, vast Forests and very dreadful Wildernesses. The sea coast is here and there covered with terrible large Rocks and overhanging Cliffs, which are overgrown in a wonderful way with Verdure and Underwood, and some with very high Trees, so that one cannot very well walk round the Island along the beach.....The sweet water flowed down from the high woody Mountains between great ravines, making its way down to the Sea in numerous little Rivers, and we found it lovely, agreeable and clear. It is said that in Amboyna and on this Island Dinding

the best fresh water of the whole East Indies is found, and this I believe to be the fact, for I myself (in my own opinion) have never in any other country in India drunk better water than in these two places."

He goes on to relate how he and his friends "heard in the wildernesses many Rattle-snakes," but they did not see any although they "made search for these monsters." The Pangkor of to-day is very well provided in the matter of reptiles, but the rattlesnakes must have left the island when the Dutch did, for I have never heard of one being seen. What SCHOUTEN heard was probably the hamadryad (*Ophiophagus elaps*), which makes a peculiar noise, but not with its tail. He also mentions that they "plucked the Oysters of the Trees," and gravely proceeds to explain how they got there, as "this might seem to some people incredible."

After a stay of five days, WOUTER SCHOUTEN left Pangkor and proceeded on his voyage to Bengal on the 3rd December, but, meeting with bad weather immediately after leaving, he had to put back to refit, and left again finally the next day.

Turning again to the Malacca records, we find a letter dated 5th August, 1670, in which orders are given to take possession of Pulau Dinding, and to build there a stronghold of wood; and another letter dated 31st October in the same year laying down that the garrison shall consist of 1 sergeant, 3 soldiers and 3 sailors. There is nothing to show when this order was carried out, but DAMPIER, who visited Pangkor 19 years later, found there "a Governour and about 20 or 30 soldiers" and a fort of stone (brick?). Besides the fort, the Governor had a house about a hundred yards away, where he used to spend the day, but he never trusted himself outside the fort at night on account of the hostility of the Malays, who, as VALENTYN says of Perak Malays generally, were, "very foul and murderous." The Perak Malays seem to have had a bad reputation from the earliest times. HAMILTON, alluding to the "cutting off" of the Dutch factory in Perak in 1651, remarks that "the inhabitants are so treacherous, faithless and bloody that no European nation can keep factories there;" and, speaking of "Selangore" and "Parsalore," he observes that

the inhabitants "have too many of the *Perak qualities* to be trusted with honest men's lives and money."

DAMPIER gives an interesting description of the fort and the Governor's house as he found them. The ruins of the former still exist, and, though they are in a very tumble-down condition, it is easy to see how accurate DAMPIER'S description was; but no trace remains of the house, which was built of timber.

He also gives a most amusing account of an entertainment given by the Governor to Captain MINCHIN and a Mr. and Mrs. RICHARDS who were on board his ship. They went ashore in the afternoon to call on the Governor, and Captain MINCHIN treated him to brandy-punch, in return for which the Governor sent out a boat to catch some fresh fish, as an addition to the other fare of the fort, and invited them to "supper." The boat came back about 5 o'clock, the fish was cooked at once, a fresh bowl of punch was brewed, and the Governor sat down with his guests and the officers from the fort prepared to do ample justice to this good cheer. But hardly had they fallen to, when one of the soldiers raised an alarm of "Malayans," which produced a most remarkable effect on the Dutch section of the party. The Governor, without saying a single word, leapt through a window and was off like a hare in the direction of the fort; the officers and servants followed suit with great promptitude; and the three guests were left behind in amazement "at this sudden Consternation of the Governour and his people." They followed, however, to the fort, where they found the Governor (possibly rather ashamed of his precipitate flight now that he was in safe quarters) waiting to receive them. The door was then shut, and the soldiers proceeded to fire the guns to show the "Malayans" that they were ready for them, but no attack was made, and their alarm must have given way to a feeling of regret at the loss of the supper and the punch which they had left behind. It is noticeable that DAMPIER, in common with other early voyagers, speaks of the excellence of the harbour at Pangkor, and also of the timber. Of the latter he says "the Trees are of divers sorts, many of which are good

timber, and large enough for any Use. Here are also some good for Masts and Yards, they being naturally light yet tough and serviceable."

In 1690, one year after DAMPIER'S visit, the Dutch garrison at Pangkor was "cut off" by the Malays under one PANGLIMA KULUP; and in 1693 an order was given that, in consequence of this massacre, no garrison should be posted again at Pulau Dinding, but that a stone pillar should be erected there, having on one side the arms of the United East India Company, and on the other those of the United Provinces. 1690. 1693.

The pillar above-mentioned seems to have been erected, and either to have fallen down or to have been pulled down by the Malays, probably the latter, for in a letter dated 8th August, 1695, an order is given to re-erect it and to clean it yearly and keep it in repair. 1695.

Having thus set up a monument of their occupation—and abandonment—of Pangkor, the Dutch seem to have left the place alone for a period of 50 years. In 1745, however, an order was given to build again a small fort at Pulau Dinding, and to put there a garrison of 30 European and the same number of native soldiers, but *no Bugis*. This order was apparently carried out, but the occupation did not last long, for in 1748 an order was given to remove the garrison on account of the insalubrity of the place, and to send them to Perak. Thus ended the Dutch occupation of Pangkor. 1745. 1748.

How long the stone pillar referred to above was maintained, it is impossible to say; but it may be inferred that some difficulty was found in doing so, for, before leaving Pangkor, the Dutch carved the arms of the United Provinces on a large granite boulder near the fort, which may be seen to this day. There are also traces of an inscription below the arms, but, with the exception of a stray letter here and there, it has been entirely effaced, partly by the action of time and weather, and partly by well-meant but injudicious attempts to clean the face of the rock.

In 1822 Pangkor was visited by CRAWFURD, who found it uninhabited. Having read DAMPIER'S description of the Dutch fort, he searched for it, and found it exactly as described. In 1822.

his account of Pangkor, CRAWFURD specially notices the "beautiful and safe harbour running North and South, and seemingly sheltered from every wind."

We now come to the British occupation of Pangkor. In 1826 that island, together with the Sembilan Islands, was ceded to the British by Perak, with a view to the suppression of piracy; but it does not appear that any use was made of the concession either at the time or for some years after. 1826.

In 1867, a Kedah Malay named Haji MAHOMED AKIB conceived the idea of settling at Pangkor, and applied to the Lieut.-Governor of Penang (Col. MAN) for permission to do so. After some consultation, the authorities being at first doubtful whether it was British territory, he was allowed to settle; a concession of 100 orlongs of land was given to him; 1867.

and in 1870 he was appointed Penghulu of Pangkor by Colonel (now Sir ARCHIBALD) ANSON. He induced a few other people to join him, but most of them went away again, as Pangkor and Pulau Sembilan were in those days the favourite haunts of pirates, and frequent descents were made by them on the settlers, so that for the first few years Haji MAHOMED AKIB lived almost alone. How he arranged matters with the pirates I do not know, and I never succeeded in getting a satisfactory answer to this question from him; but they do not seem to have visited him as they did other people. I may add, however, that when the Government undertook the suppression of piracy in these waters in earnest, Haji MAHOMED AKIB rendered valuable assistance on several occasions. 1870.

In January, 1874, the Pangkor Treaty was signed, by which, in addition to Pangkor, a strip of territory on the mainland containing an area of about 200 square miles was ceded to the British Government, and the whole territory was called the Dindings. It was made a dependency of the Settlement of Penang, but it was arranged, as a matter of convenience, that it should be administered by the Resident of Perak, the Revenue and Expenditure being, of course, included in the Budget of the Colony. 1874.

After the Pangkor Treaty, some Police were placed on the

island under a European Sergeant to assist the Penghulu in keeping order, and in March, 1875, Inspector WARNE was sent to take charge of the Station. After remaining there six months, he was relieved by Inspector COOPER in September of the same year, and the latter was in charge till April, 1877, when Captain LLOYD, who was appointed Superintendent of the Dindings on the 29th March, took charge.

On the 26th October, 1878, after only eighteen months' service, Captain LLOYD was brutally murdered by some of the Chinese coolies from the Lumut Estate, who, being thrown out of employment, naturally turned to their favourite pastime of gang-robbery. The details of this crime are familiar to most people in the Straits, and I will not dwell on them here.

After Captain LLOYD'S murder, Mr. R. R. BRUCE, then an Inspector in the Perak Police, was placed temporarily in charge of the Dindings. He was appointed Superintendent at the beginning of 1879, and continued to hold that post till 15th September, 1882, when he nearly lost his life under the following circumstances. One Haji OSMAN BIN HASAN had gone "amok" at Telok Sera and wounded four boys, and had then been killed by the Penghulu and Assistant Penghulu. On going into the case, Mr. BRUCE found that one Haji HASAN, who was then at Taiping, was required as a witness, and had him brought to Pangkor. On the 15th September, Mr. BRUCE examined Haji HASAN in the old "balei" or court-house at Pangkor, Captain LLOYD'S former residence. Haji HASAN knew nothing of the "amok" at Telok Sera, and at first Mr. BRUCE did not tell him about it, but subsequently he did so. One of the boys who had been wounded was related to Haji HASAN, and the latter seemed much disturbed and put out by the news. He asked leave to go at once to Telok Sera and see the boy, but as he was not reported to be dangerously wounded, Mr. BRUCE told Haji HASAN he could go as soon as the enquiry was finished. He said nothing further and seemed satisfied. Shortly afterwards Mr. BRUCE and Inspector MCKEON of the Perak Police adjourned to an inner room for breakfast. They had not been there long, when Inspector MCKEON happened to look up,

and saw Haji HASAN behind Mr. BRUCE, who was sitting with his back to the door, with a "golok" in his hand, which he was just about to plunge into Mr. BRUCE'S left shoulder. Inspector MCKEON sprang to his feet and seized Haji HASAN by the throat with one hand, while with the other he attempted to seize Haji HASAN'S right hand in which he had the "golok." He was only able, however, to alter the direction of the blow, and Mr. BRUCE received a serious wound in the back, but he undoubtedly saved Mr. BRUCE'S life, for, had he been stabbed on the left shoulder as Haji HASAN intended to do, the wound must have been fatal. Inspector MCKEON then had a severe struggle with Haji HASAN, in the course of which he received an ugly wound in the left forearm, but he eventually succeeded in wresting the "golok" from him, and getting him down in a corner, when the Police came to his assistance. This sudden attack on Mr. BRUCE appears to have been quite unpremeditated. It is probable that the news of the "amok" at Telok Sera upset Haji HASAN'S mental equilibrium and caused him to "amok" himself.

Mr. BRUCE suffered a great deal from the effects of his wound, which obliged him to go to Europe for a change, and ultimately to resign his appointment.

In June, 1883, MR. URBAN BRUCE was appointed Superin- 1883.
tendent of the Dindings, and held that post till 23rd Novem-
ber, 1884, when he met his death through a most unfortunate 1884.
accident in Penang, where he had gone to recruit his health
after an attack of fever.

On his death Mr. A. T. DEW, of the Perak Government
service, was appointed Acting Superintendent of the Dindings,
and was confirmed in the appointment in May, 1886. He re- 1886.
mained in charge of the district till July, 1886, when it was
decided by the Government of the Colony to take the admin-
istration into its own hands, and an Officer of the Colonial
service was placed in charge. By Ordinance XI of 1886, the
territory and islands of the Dindings ceased to be a part
of the Settlement of Penang and were made a dependency of
the Settlement of Singapore. They have since been re-

transferred to the Settlement of Penang by Ordinance IX of 1890.

Although the contemporary history of the Dindings only extends over a period of 17 years, it cannot be said to have been an uneventful one, as two out of the first four Superintendents met their deaths while holding the appointment, and a third was seriously wounded. It is to be hoped, however, that the tragic era has passed, and that a peaceful and prosperous future is in store for a district which, when developed, should be a valuable accession to the Colony.

In concluding this account, it may not be out of place to give a brief description of the district under notice. It consists of Pulau Pangkor, Pulau Sembilan (the Nine Islands), lying about 10 miles to the South of Pangkor, and the strip of territory on the mainland already alluded to, which is bounded on the North, East and South by Perak, and on the West by the sea. The northern boundary is 8 miles in length, the eastern 22 miles, and the southern 5 miles, while the length of coast line, roughly speaking, is between 25 and 30 miles. The area of this portion of the district is about 200 square miles. The island of Pulau Jarra, called by the Malays Pulau Temurah, is also a dependency of the Dindings. It is about half a mile long forming a rounded hill 500 feet high, in the middle of the Straits of Malacca, with 25 to 40 fathoms all round.

It has two rivers—Sungei Bruas at the northern end, a narrow winding stream, which is a good deal used, nevertheless, for bringing down timber and jungle produce from the interior; and Sungei Dinding opposite Pangkor, which for about 15 miles from its mouth is a fine deep river, navigable for the largest vessels. At a place called Kota Siam it divides itself into three branches—Sungei Kota Siam, Sungei Sigari and Sungei Raja Itam—the latter of which is navigable for launches for some 5 or 6 miles above Kota Siam, or about 20 miles in all from the mouth. The late Penghulu, Haji MAHOMED AKIB, is fond of relating how he once piloted H. M. S. *Iron Duke* up to Kota Siam and back.

At Kota Siam there is an old earthwork which, the Malays

say, was once occupied by Siamese, who carried on a continual and lively warfare with the Malays under Raja ITAM, whose fort is a little higher up the river, and has given the name to one of the branches.

Two causes have hitherto operated to prevent the river becoming the port of Perak, as it seems intended by nature to be—(1) the fact that it is in British territory, and that the Government of Perak naturally wanted to have a port in their own State; and (2) the entire absence of communication between the river and the interior of Perak. The question of railway communication between the Dindings and Kinta recently engaged the attention of the Government, but has been shelved for the time being owing to the construction of a line from Telok Anson to Tapa.

The coast is bold, and in many places rocky, with fine, sandy beaches, presenting a marked contrast to the Perak coast on either side, of which mud-banks and mangrove swamps are the chief characteristics. A range of hills runs along the coast for nearly its entire length, and also for some distance along the North bank of the Dinding River, the highest point being at Telok Sera. I believe that the height of this point has never been ascertained, as it is not given in the Admiralty chart, which gives the heights of other points, but it cannot be less than 1,500 feet, and I should say is probably more. On the banks of the Dinding River, opposite Pangkor, there are three hills over 1,000 feet in height and several smaller ones. The soil is fertile and well suited to the cultivation of such products as tapioca, pepper, gambier, Liberian coffee, spices, &c. North of Telok Sera there is a considerable tract of fine, low, swampy land suitable for the cultivation of padi, and, I should say, of sugar also, though the latter has not yet been tried. Padi grows excellently there, and if all this land were to be brought under cultivation, a valuable rice crop might be produced.

The chief product of the district has, however, hitherto been timber, of which a considerable quantity must have been exported during the last ten or twelve years. The quantity exported in 1887 was, in round numbers, 2,200 tons. I know

of at least one timber merchant in Penang who started on a very small scale on the Dindings some years ago, and who has made a large fortune out of timber. Steps are now being taken to establish forest reserves before all the best kinds of timber are exhausted, for although the district is a small one, it is the only part of the Colony in which there is any extent of forest left.

Tin is the only metal found in any quantity in the district. There are considerable deposits of alluvial tin in three or four places, and experts judge from the appearance of the ore, that it has not travelled far from the lodes. There being no hills outside the district nearer than the main Perak range, it is reasonable to conclude that the lodes are located in the hills at the foot of which the alluvial deposits are found.

The tin has not yet been worked by Europeans, nor, except on a very small scale, by natives, notwithstanding the great facilities that exist in the way of water-carriage. Before the collapse of the tin syndicate, there was every prospect of land being taken up in the Dindings by Chinese capitalists from Taiping (Perak), but the fall in the price of tin caused the scheme to fall through. Round about Telok Sera there are traces in many places of old workings (chiefly *Lampau*), which are said by the Malays to be of Siamese origin.

Gold has been found in small quantities at one or two places, and in Pangkor Island there is place which is still called Telok Klian Mas.

The island of Pangkor is about 12 miles long and 4 or 5 miles wide at the broadest part. It is very hilly, there being three peaks over 1,100 feet high and three over 700 feet. The highest point is 1,313 feet. These figures are taken from the Admiralty chart. There is not much level land, but the soil is good, and many of the smaller hills are well adapted for planting. The headquarters of the district were, till last year, situated on the East side of the island, where there is an excellent harbour, with deep water and well sheltered from all winds. It is well known for the abundance and excellence of the fish caught there, a large quantity of which is cured and exported. A few small steamers call occasion-

ally at Pangkor, but the bulk of the trade is carried on by native craft, chiefly Chinese.

The one drawback to this beautiful island is its unhealthiness. From the appointment of the first Superintendent up to the present time, the journals of the officers in charge have been one long record of sickness caused by the malarial fever which seems to have made Pangkor its home, and which spares no one. Judging from appearances, one would say that the island ought to be very healthy, and it is difficult to say what causes the fever. The late Rev. J. E. TENNISON-WOODS, who spent some weeks at Pangkor in 1884, attributed the unhealthiness of the place to the fish-curing which is carried on at Raja Bayang, the principal village of the island. He says: "The effluvium around the village of Raja Byong was unbearable to those who had not been inured to it by previous education. Whenever the wind blew from the village towards our quarters, we had to leave the house. Our Chinese servants, in spite of every prohibition, spread their mats under the bungalow at night, and exposed themselves to the full force of these mephitic breezes. The consequence was that they were all stricken down with fever, and some nearly died. Pankore has the name of being a very unhealthy place, *but the marvel is how anybody lives there at all.* The inhabitants suffer much from what is called malarial fever, *but the malaria here is undoubtedly mephitism from putrid fish offal.*" * This may be true as far as the village of Raja Bayang is concerned, but other parts of the island, where there is no fish-curing, are just as unhealthy, and the cause must, therefore, be sought elsewhere. It is probable that nothing but extensive clearing of the jungle will effect any improvement. The headquarters of the district were moved last year to Lumut at the mouth of the Dindings River, which it was hoped would be a healthier position, but the results of the move have not been very encouraging so far.

The group of islands called Pulau Sembilan lie about 10 miles to the South of Pangkor, opposite the mouth of the

* Essay on the Fisheries of the Oriental Region, p. 27. The italics are mine.—E. M. M.

Perak River. They are uninhabited, except during the season when the turtles lay their eggs, when the person to whom the right of collecting the eggs is farmed places a few Malays there to see that no poaching takes place, and to collect the eggs. There is excellent water on one of the islands, and in dry weather the Malays from the mainland are very glad to avail themselves of this supply in spite of the distance they have to go for it. I have been told by Malays that there is a cave on one of the islands where edible birds' nests are to be found, but I could never find any one who could show me the entrance. One man said that when the pirates left the neighbourhood they closed up the entrance to the cave with a big stone. If the cave ever existed outside the imagination of the natives, it is probable that this statement is true.



NATIVE NAMES OF STREETS IN SINGAPORE.

—:0:—



ACTING under the powers conferred by section 143 of the Municipal Ordinance 1887, the Commissioners have affixed at the corner of every street in the town of Singapore a board on which is inscribed "the name by which such street shall be known."

But, as most of the residents are aware, the names given by the Municipality to the various streets are only used by the European portion of the population, and the Chinese, Tamils and Malays have names for the streets very different from their Municipal titles. In the selection of names for the streets, Chinese names were very properly selected for the Chinese part of the town, and Malay names for many of the streets in Kampong Glam, and one would naturally have supposed that such names as "Hongkong Street" and "Macao Street" would have been adopted by the Chinese, and that the Malays would have had no hesitation in appropriating a name like "Jalan Sultan."

But, whatever the reason may be, the fact remains that the Municipal names are ignored by the natives, with the exception of the police, who are, of course, compelled to learn them. I think, therefore, it will be of interest to give a list of some of the principal Chinese and Tamil names, as the names illustrate, to a certain extent, the characteristics of each nation. For example, what can be more characteristic of the Singapore Tamil than the names he has given to the Esplanade, to Bain Court, and to Tank Road? With regard to the Chinese names, they may, I think, be divided into three classes, *i.e.*, purely Malay names pronounced in Chinese fashion, Malay names translated into Chinese, and lastly descriptive names.

The descriptive names are mostly very good, and while the appropriateness of the name for Church Street may only strike those who are acquainted with police work in Singapore, yet we can all admire the picturesque names for Thomson and Alexandra Roads. I have not attempted to include the Malay names in the list, as Malays take but little notice of streets, and, as a rule, only describe places by *Kampongs*, which are of course well known in the case of Singapore.

I am indebted to my friend Mr. J. O. ANTHONISZ for the list of Tamil names, and I have to thank Messrs. LI A SIK and TAN TIANG SIONG for preparing the list of Chinese names. The Chinese names are given in the list as pronounced by the Hokkien Chinese.

H. T. HAUGHTON.

CHINESE NAMES OF STREETS AND ROADS IN SINGAPORE.

Albert Street. 監公望舊魯 *or* (2) 磨蔴油街

Kam Kong Mang Ku Lu (Kampong Bencoolen). *Or* (2) Boh Mua lu Koi—"The street where sesamum oil is expressed."

Alexandra Road. 水磨內 *or* (2) 老只脚

Chwi Bo Lai—"The water rice-mill." *Or* (2) Lau Chi Kha—"The foot of the betel-nuts."

Almeida Street. 牛車水戲園後

Gu Chia Chwi Hi Hng Au—"The back of the Kreta Ayer Theatre." "Gu Chia Chwi," literally "water cart," a translation of the Malay name "Kreta Ayer."

Amoy Street. 媽祖宮後 *or* (2) 義學口

Ma Cho Keng Au—"The back of the Ma Cho Temple." *Or* (2) Ghi Oh Khau—"The front of the school."

Angus Street. 監公嗎呔呷

Kam Kong Ma La Kak (Kampong Malaka). The same name is also given to Cumming Street, Fisher Street, Kerr Street, Keng Cheow Street, Omar Road and Solomon Street.

Arab Street. 爪亞街

Jiawa Koi—"Javanese Street."

Armenian Street. 成寶新厝後

Seng Poh Sin Chu Au—"The back of Seng Poh's new building."

Bain Street and Holloway Lane. 色仔厘禮拜對面巷

Sek Kia Ni Loi Pai Twi Bin Hang—"The lane opposite the Portuguese Church." "Sek Kia Ni"=Serani.

Balestier Road. * 鵝曹大伯公

Go Cho Tua Pek Kong—"Rochor Temple." "Go Cho" =Rochor.

Bali Lane. 嘛哩內

Bali Lai. "Lai"=inside, off the road. Compare Haji Lane and Shaikh Madarsah Lane.

Battery Road. 土庫後

Tho Kho Au—"Back of the godowns." "Tho Kho" = Godown, literally "earth treasury."

Beach Lane. 鐵巴虱巷

Thi Pa Sat Hang—"Iron market lane."

Beach Road. 小坡海墘 *or* (2) 鐵巴虱口

Sio Po Hai Ki—"Small Singapore's seaside." "Sio Po" = "Small Singapore" (Kampong Glam), in contradistinc-

* Balestier Road called after JOSEPH BALESTIER, Consul for the United States of America who died about 1848.

tion to **大坡** "Twa Po"="Big Singapore" across the river. *Or* (2) Thi Pa Sat Khau—"Iron market mouth." The iron market is of course Clyde Terrace Market.

Boat Quay. **淡邦路頭** (2) **十三行** (3) **十八間**
(4) **水厝尾** *or* (5) **溪墘**

Tiam Pang Lo Thau—"The place to go for sampans." "Tiam Pang"=sampans. "Lo Thau"=the beginning of the way. *Or* (2) Chap Sa Hang—"The thirteen shops." There is a street in Canton of the same name. *Or* (3) Chap Poet Keng—"The eighteen houses." *Or* (4) Chwi Chu Boi—"Bathing house end." *Or* (5) Khoi Ki—"Stream bank."

Bras Basah Road. **老脚舊間口**

Lau Kha Ku Keng Khau—"Mouth of the old jail." "Kha Ku" literally "ankle chains."

Buffalo Road. **監公加路武**

Kam Kong Ka La Bu (Kampong Kerbau)—"Kandang Kerbau."

Bukit Timah Road. **馬車路尾** *or* (2) **新山路**

Be Chia Lo Boi—"End of the horse-carriage street." Meaning that the road ends at Kranji. *Or* (2) Sin Swa Lo—"Johor Road." "Sin Swa"=Johor, literally "New mountain."

Canal Road. **監公嗎呖呷港**

Kam Kong Ma La Kak Kang—"The Kampong Malaka stream."

Canton Street. **橫街仔**

Whe Koi Kia—"The small road that crosses," *i. e.*, Boat Quay and Circular Road.

Carpenter Street. **戲館街** *or* (2) **義福街**

Hi Kwan Koi—"Wayang-house street." *Or* (2) Gli Hok Koi—"The street of the Ghi Hoks."

Cheang Hong Lim Street. **八間仔**

Poeh Keng A—"The eight small houses."

Chin Chew Street. **荳腐街**

Tan Hu Hoi—"Beancake street," so called from the beancake sellers who live there. Beancake is called by Babas "Kweh Koyak."

Chin Hin Street. **大坡新街**

Twa Po Sin Koi—"The new street of big Singapore."

China Street. **繖間前** *or* (2) **福建義興公司前**

Kiau Keng Cheng—"The front of the gambling houses."
Or (2) Hok Kien Ghi Hin Kong Si Cheng—"The front of the Hokkien Ghi Hin Kongsì."

Church Street **繖間口** *or* (2) **五代天福宮口**

Kiau Keng Khau—"The mouth of the gambling houses."
Or (2) Ngo Tai Tian Hok Kiong Khau—"The mouth of the five generations the Tian Hok Temple."

Circular Road. **十三行後** *or* (2) **十八間後**

Chap Sa Hang Au—"The back of the thirteen shops," *i. e.*, the back of Boat Quay. *Or* (2) Chap Poet Keng Au—"The back of the eighteen houses," *i. e.*, Boat Quay.

Clyde Street. **鐵巴虱馬打厝對面巷**

Thi Pa Sat Ma Ta Chu Twi Bin Hang—"The lane opposite the iron-market Police Station."

Clyde Terrace. **碗店口**

Ua Tiam Khau—"The mouth of the crockery shops."
There are many crockery shops in this street much patronized by the Bugis traders. The Malay name is "Jalan Kedei Pinggan Mangko."

Coleman Street. **勐尼新厝後**

Hiok Ni Sin Chu Au—"At the back of (Tan) Hiok Ni's new building."

Collyer Quay. **土庫後**

Tho Kho Au—"At the back of the godowns."

Commercial Square. **土庫口** *or* (2) **花園角**

Tho Kho Khau—"The mouth of the godowns." *Or* (2)
Hua Hooi Kak—"Flower garden corner."

Crawford Street. **鵝曹尾** *or* (2) **小坡老巴剎**

Go Cho Bue—"The end of Rochor." *Or* (2) Sio Po Lau
Pa Sat—"The old market of small Singapore."

Cross Street. **咕嚕仔街**

Kiat Leng Kia Koi—"The Klingmen's street," so called on account of the numerous Klings, boatmen principally, who live there.

Ellenborough Street. **新巴剎口**

Sin Pa Sat Khau—"The mouth of the new market."

Esplanade. **大葛前草坡**

Twa Kok Cheng Chau Po—"The grass field in front of the great Court."

Fish Court. **咯嗎班囊巷**

Lo Ma Pan Jiang Hang (Rumah Panjang Lane). The "Rumah Panjang" was the old atap building in Rochor Road where the *demi-monde* lived before Fraser Street was built.

Fish Street. **新巴剎邊**

Sin Pa Sat Pi—"At the side of the new market."

Flint Street. **土庫尾**

Tho Kho Bue—"The end of the godowns."

Fraser Street. **小坡新街**

Sio Po Sin Koi—"The new street of small Singapore."

Garden Street and Jedda Street. **鐵巴虱對面巷**

Thi Pa Sat Twi Bin Hang—"The lane opposite the iron market."

George Street. **玻璃後順豐街**

Po Le Au Sun Hong Koi—"The Sun Hong Street at the back of the Police," *i. e.*, Central Station. "Sun Hong" was a shop formerly in that street and it gave its name to one of the five divisions of the annual Chin Ge procession.

Synagogue Street bears the same name.

Haji Lane. **舊峇厘**

Ku Ba Li—"Old Bali (lane)."

Havelock Road. **槓石街 or (2) 酒廊路**

Kong Chioh Koi—"Break-stone street." *Or (2)* Chiu Long Lo—"Spirits-shed street," so called from an arrack manufactory there.

High Street. **大葛街 or (2) 雙井路**

Twa Kok Koi—"Supreme Court street." *Or (2)* Siang Che Lo—"Two-wells street," *i. e.*, the two wells at the foot of Fort Canning.

Hill Street. **王家山脚 or (2) 吊橋頭**

Ong Ke Swa Kha—"The foot of the Governor's hill." Compare the Malay name of Fort Canning—"Bukit Tuan Bonham." *Or (2)* Tiau Kio Thau—"At the end of the suspension bridge," *i. e.*, Coleman Bridge.

Hokkien Street. **造馬車街**

Cho Be Chia Koi—"The street where horse-carriages are made."

Hong Lim Quay. **監公嗎吹呷海墘**

Kam Kong Ma La Kak Hai Ki—"Kampong Malaka sea beach."

Hongkong Street. 馬交街
Ma Kau Koi—"Macao street,"

Hylam Street. 海南會館後
Hai Lam Hoi Kuan Au—"The back of the Hailam meeting house"

Jalan Sultan. 二十間
Ji Chap Keng—"The twenty houses."

Japan Street. 媽祖宮邊
Ma Cho Keng Pi—"The side of the Ma Cho temple."

Java Road. 小坡打石街
Sio Po Phah Chioh Koi—"Small Singapore cut-stone road."

Kalang Road. 呀喼橋
Ga Lang Kio—"Kalang bridge." The vulgarism of "Gallang Road" evidently comes through the Chinese pronunciation.

Kampong Bharu. 新監公
Sin Kam Kong—"New kampong."

Kampong Glam Beach. 大井脚
Twa Che Kha—"The foot of the big well." There used to be an old well in the middle of the road at Sultan's Gate.

Kampong Java Road. 紅毛塚
Ang Mo Phun—"European tombs."

Kandang Kerbau and Selegie Road. 竹仔脚
Tek Kia Kha—"The foot of the small bamboos." This

name is probably given on account of the bamboo clumps in the neighbourhood.

Killenev Road. 東陵巴虱對面巷

Tang Leng Pa Sat Twi Bin Hang—"The lane in front of the Tanglin market."

Kim Seng Road. 豐興橋

Hong Hin Kio—"Hong Hin bridge." "Hong Hin"= chop of Tan Kim Seng.

Kling Street. 山仔頂 *or* (2) 鐵條脚

Swa Kia Teng—"Small hill top." *Or* (2) Thi Thiau Kha "The foot of the iron pillars." Many iron pillars were used in the construction of houses in this street.

Lavender Street 鵝曹大公司 *or* (2) 菜園內

Go Cho Twa Kong Si—"The big Rochor kongsi." The chief meeting house of the Thien Thi Hoi was in this street. * *Or* (2) Chai Hng Lai, meaning that vegetable gardens are alongside the road. "Chai Hng"=Vegetable gardens; and "Lai"=inside.

Little Cross Street. 縐亞打銅街

Jiowa Phak Tang Koi—"The street of the Javanese copper-smiths."

Lorong Telok. 篾籠街

Bi Lang Koi—"Bamboo basket street."

Macao Street. 玻璃邊

Po Li Pi—"The side of the Police," *i.e.*, Central Station.

Magazine Road. 陳聖王街

Tan Seng Ong Koi—"Tan Seng Ong street." Tan Seng Ong was the ancestor of the Seh "Tan."

* See "Manners and Customs of the Chinese of the Straits Settlements," by J. D. VAUGHAN, p. 110.

- Malabar Street. 舊海南會館
Ku Hai Lam Hoi Kwan—"The old Hailam meeting-house."
- Malacca Street. 老巴虱口 *or* (2) 萬興街
Lau Pa Sat Khau—"Mouth of old market." *Or* (2) Ban Hin Koi—"Ban Hin street," so called from the firm of that name.
- Manila Street. 色仔呢街
Sek Kia Ni Koi—"Serani street." Queen Street bears the same name.
- Market Street. 中街 *or* (2) 老巴虱口
Tiong Koi—"Centre street." This is one of the divisions of the Chin Ge procession. *Or* (2) Lau Pa Sat Khau—one of the names for Malacca Street.
- Merchant Road. 新街口戲園街
Sin Koi Khau Hi Hng Hoi—"The theatre street at the mouth of the new street," *i.e.*, Chin Hin Street.
- Middle Road. 小坡紅毛打鐵
Sio Po Ang Mo Pah Thi—"European foundry of small Singapore." The foundry referred to formerly belonged to Messrs. CAZALAS. It now belongs to a Chinese firm, the Ban Hap Kongsì.
- Nankin Street. 松柏館街
Siong Pek Kwan Koi—"Siong Pek Kwan street." "Siong Pek Kwan" was the Keh branch of the Thien Thi Hoi now suppressed along with the other dangerous societies.
- Neil Road. 牛車水斜路
Gu Chia Chwi Kia Lo—"The steep street of Kreta Ayer."
- New Bridge Road. 新巴虱馬打厝前 *or* (2) 二馬路

Sin Pa Sat Ma Ta Chu Cheng—"The front of new market Police Station. *Or* (2) Ji Ma Lo—"Number two horse way." South Bridge Road being Tai Ma Lo—"The great horse way."

New Harbour. 新監公船澳 *or* (2) 石叻門

Sin Kam Kong Chun U—"Kampong Bharu Dock." *Or* (2) Sek Lat Moi—"Selat passage." Sek Lat=Malay "Selat." Telok Blanga Road bears the same name.

New Market Road. 新巴虱水棚脚

Sin Pa Sat Chwi Pe Kha—"The foot of the bathing house at the new market."

North Boat Quay. 掬霜前

Kek S'ng Cheng—"The front of the ice factory."

North Bridge Road. 水仙門火車路

Chwi Sian Moi Hui Chia Lo—Literally "Water fairy door fire cart road." Hui Chia=steam tram.

North Canal Road. 溝仔墘 *or* (2) 單邊街

Kau Kia Ki—"The side of the little drain," *i. e.*, the old Singapore canal. *Or* (2) Tan Pin Kai—"One side street," so called because there are houses only in one side of the street.

Orchard Road. 東陵巴虱街

Tang Leng Pa Sat Koi—"Tanglin market street."

Outram Road. 時牌坡

Si Pai Po—"The Sepoy's field."

Pagoda Street. 咕啞仔禮拜後

Kek Leng Kia Loi Pai Au—"The back of the Kling place of worship."

Pekin Street. 衣箱街

I Sio Koi—"Clothing box street," so called from the cabinetmakers living there.

Phillip Street. 老爺宮口

Lau Ia Keng Khau—"Mouth of the Lau Ia temple," *i. e.*, the Teo Chiu temple. "Lau Ia"=grandfather.

River Valley Road. 王家山脚 *or* (2) 龍頭井

Ong Ke Swa Kha—"The foot of the Governor's hill" *i. e.*, Fort Canning. *Or* (2) Leng Thau Che—"Dragon's head well," *i. e.*, the fountain by the side of the road opposite the Ice Works.

Rochor Road. 略嗎班襄

Lo Ma Pan Jiang—"Rumah Panjang." See Fish Court.

Sago Street. 牛車水戲園前

Gu Chia Chwi Hi Hng Cheng—"The front of Kreta Ayer theatre."

Saranggong Road. 後港路

Au Kang Lo—"The Au Kang road." "Au Kang" is the Kangka at the end of the road.

Selegie Road.

See Kandang Kerbau. Is Selegie the Malay word "Sëligi," a "wooden-dart"?

Shaikh Madarsah Lane. 新巴厘

Sin Ba Li—"New Bali." Compare Haji Lane, "Old Bali."

Smith Street. 牛車水戲園街

Gu Chia Chwi Hi Hng Koi—"Kreta Ayer theatre street."

South Bridge Road. 大馬路 *or* (2) 漆木街

Tai Ma Lo—"The great horse-way." *Or* (2) Chat Bok Koi "Paint wood street." This last name is however only given to that portion of South Bridge Road lying between Elgin

Bridge and the corner of North Canal Road, and it is so called on account of the numerous painters who live there.

South Canal Road. **港仔墘**

Kang A Ki—"The side of the small river," *i. e.*, the old Singapore Canal.

Stamford Road. **流水澗**

Lau Chwi Khe—"Flowing water stream."

Sultan Gate. **王府口** *or* (2) **打鐵街**

Ong Hu Khau—"The mouth of the palace." *Or* (2) Phah Thi Koi—"The street of the iron-smiths."

Sultan Road. **小坡打鐵街**

Sio Po Phah Thi Koi—"Small Singapore's iron-smiths' street."

Tanglin Road. **大東陵**

Twa Tang Leng—"Great Tanglin."

Tank Road. **王家山後**

Ong Ke Swa Au—"The back of the Governor's hill," *i. e.*, Fort Canning.

Telok Ayer Street. **源順街**

Gwan Sun Koi—"Gwan Sun street." "Gwan Sun" was the chop of a firm in the street which gave its name to one of the five divisions of the Chin Ge procession.

Thomson Road. **井水港**

Chia Chwi Kang—"Fresh water stream," *i. e.*, the Kalang River.

Tringganu Street. **牛車水橫街**

Gu Chia Chwi Wha Koi—"The cross street of Kreta Ayer," *i. e.*, crossing Smith Street and Sago Street.

Upper Circular Road. 馬車街

Be Chia Koi—"Horse carriage street," so called from the gharry stand there.

Upper Cross Street. 海山公司街

Hai San Kong Si Koi—"Hai San kongsi street," so called because the meeting-house of that kongsi used to be in the street.

Upper Hokkien Street. 叻巴街 or (2) 長泰街

Lam Pa Koi—"Swamp street." Or (2) Chiang Tai Koi—"The Chiang Tai temple street."

Victoria Street. 後馬車路

Au Be Chia Lo—"The back of horse-carriage street," *i. e.*, at the back of North Bridge Road.

TAMIL NAMES OF STREETS AND ROADS IN SINGAPORE.

Albert Street. தீமிரித் திடல்

Thimiri Thidal—"Place where people tread on fire." The ceremony of walking through the fire in October-November was held in the street.

Arab Street. பூக்கடைச் சடக்கு

Pūkadei Sadakku—"Street of the flower shops."

Bain Court. புருவன் துரை சாயப்பட்டரை

Brown Thurei Sāyapattare—"Brown's carriage painting establishment."

Bain and Holloway Streets. பாக்கு தோப்பு

Pākku Thoppu—"Betel-nut garden."

Balestier Road. தண்ணீர் கம்பம்

Thannir Kampam—"The water kampong," so named as water was conveyed from this place to town for sale.

Beach Road. கடல்கரை சடக்கு

Kadal Karei Sadakku—"Road by the seaside."

Chin Chew Street. ஆரம்பிள்ளை சடக்கு

Arapillei Sadakku—"Arapillei's road."

Clyde Terrace. இரும்பு பாசா

Irumpu Pasar—"Iron market."

Commercial Square. கிட்டங்கு தெருவு

Kidangu Theruvu—"Street of the godowns."

Cross Street. பால்கடைச் சடக்கு

Palkadei Sadakku—"Street of the milk shops." [The Malay name is "Kampong Susu."]

Dhoby Ghaut. வண்ணன் தெருவு

Vannan Theruvu—"Street of the dhobies."

Esplanade. ஜனவெரி கிடல்

January Thidal—"January place" so named from the sports held on 1st January.

Havelock Road. மாசா ஆரா சடக்கு

Masak Arak Sadakku—"Arrack distilling street." ["Jalan Masak Arak" is the Malay name.]

Kalang Road. காலாங் விளக்கு கூடு

Kalang Villakukhûdu—"Kalang light cage." (Gasometer).

Kampong Kapor. சுண்ணாம்பு கம்பம்

Sunnambu Kampam—"The lime kampong."

Kling Street. கத்திக்கடைச் சடக்கு

Kathi Kadei Sadakku—"Knife shop street." [The Malay

name is "Jalan Kedei Pisau."]

Lavender Street. குசுத் தெருவு
Kosa Theruvu—"Potters' street."

Mackenzie Road. தண்ணீர் ஆலை
Thannir Âlei—"Water engine."

Market Street. செட்டித் தெருவு
Chetty Theruvu—"Chetties' street."

Orchard Road (town limit). வயிராசி மடம்
Vairakimadam—"Fakir's place."

Orchard Road (from Koek's Market to Upper Tang-
lin). மேட்டு தான்
Mattu Than—"It is high ground."

Pagoda Street. மாரியம்மன் கோயில் பக்கத்து சடக்கு
Mariammen Kovil Pakkathu Sadakku—"The side street
by temple of Mariammen."

Queen Street. டோபி கம்பம்
Dhoby Kampam—"Street of the dhobies." [The Malay
name is "Kampong Dhobi."]

River Valley Road. கல் ஆலை
Kal Âlei—"Stone crusher" from the steam-crusher which
is kept at the corner of Tank Road and River Valley Road.

Rochor Road. கமமங்காலா பழைய கூத்துமேடை சடக்கு
Kammangala Paleia Kuthû Mâdei Sadakku—"Street of
Kampong Glam old Hindu theatre."

Selegie Road (near Mackenzie Road). நாகப்பன் தான்
Nagappen Thân—"It is Nagappen's."

South Bridge Road. கலப்பித்தி கடை சடக்கு
Kalapithi Kadei Sadakku—"Cawker's shop street."

Sultan Gate. இராசா கோட்டை

- Raja Kôttei—"Rajah's palace."
- Tanglin, Upper. வம்பு மலை
Vampumalei—"Whampoa's hill."
- Tank Road. கல்லப்பன் தான்
Nallappen Thán—"It is Nallappen's."
- Thomson Road. தண்ணீர் பீலி சடக்கு
Thanîr Pîlei Sadakku—"Water pipe Street."
- Victoria Street. கமமங்காலா புதுகூத்து மேடை சடக்கு
Kammangala Puthu Kûthu Madei Sadakku—"Street of
Kampong Glam new Hindu theatre."
- Victoria Street (lower). போயான் கம்பம்
Boyan Kampam—"Boyan kampong." [The Malay name
is "Kampong Boyan Lama."]
- Victoria Street (a lane off). பால் கம்பம்
Pal Kampam—"Milk village."
- Waterloo Street. கிருஷ்ணன் கோவில் சடக்கு
Krishnen Kovil Sadakku—"Street of Krishnen temple."

H. T. HAUGHTON.

NOTES ON A TRIP TO BUKIT ETAM, SELANGOR.

BY

LIEUT. H. J. KELSALL, R. E.



ON the 1st of January, 1891, I started in company with Mr. LAWDER, District Officer at Kajang, for a five days' trip in the jungle. Our destination was Bukit Etam, situated about 25 miles due East of Kwala Lumpur. By road the distance is about 30 miles.

We reached as far as Ulu Langat, a native village about 13 miles from Kwala Lumpur, and slept there that night in the Police Station, a room of which is set apart for the use of travellers. Here we got eight or nine coolies to carry our baggage.

At 7.20 A.M. on the 2nd, we left Ulu Langat. We rode the first 12 or 13 miles to the foot of the hill, there being a good bridle-path, which had been made by Mr. LAWDER for the use of the Resident a few weeks before. Previous to this there had only been a foot-path through the jungle. The path crosses the Langat River at Ulu Langat, and then runs more or less parallel to the left bank of the river, passing for the first two miles through second growth jungle, and now and then crossing a stream by means of a rough bridge. The jungle then becomes more dense, the ordinary lowland jungle, with thick undergrowth, the most striking tree being the beautiful and orange *Ixora*, the "Jarum Jarum" of the Malays, which here grows to a height of 25 or 30 feet. Three miles from Ulu Langat a path branches off to the hot springs of Dusun Tuah. There are five or six of these springs altogether.

all near the river. One, I am told, actually rises in the bed of the stream. The principal stream rises out of a mass of granitic rock about 20 feet high and 30 or 40 in circumference. Sulphuretted hydrogen is given off in considerable quantities by the springs. The temperature of the water is, I believe, about 180° Fahr., but as I had no thermometer I could not test it myself. Butterflies of several species appeared to like the hot water, for they hovered about and settled on the moist rocks and sucked up the water. Here I got three or four specimens of that prince of butterflies, the beautiful *Ornithoptera Brookeana*, resplendent in black and green velvet. The water of the river here is quite cold and clear like an English stream, and orchids are pretty abundant on the trees. No beasts were seen in the jungle, and only a few birds. One or two red woodpeckers, a few tailor-birds (*Orthotomus ruficeps*) and bulbuls, a solitary pair of crows, and now and then a wagtail were the only representatives of the feathered tribes that were seen.

The Sungei Lui, an affluent of the Sungei Langat, had to be forded three or four times in the course of our ride, but was nowhere much more than kneedeep. On the sandy banks of the river, especially at one place where there were a lot of durian skins, and at moist spots on the paths, numbers of butterflies, principally belonging to the family *Pieridæ*, but including some *Papilios* and others, were seen, and every now and then one of the blues would cross the path with a flash of metallic light, or a big swallow-tail zigzag down the road, or a glorious black and gold *Ornithopteron* come sailing lazily out of the jungle, at one side only, to disappear a moment or two later on the other. Leeches were, as usual, plentiful in this low damp jungle. A beautiful orange-flowered globba (*Globba aurantiaca*) was plentiful along the sides of the path, and I found a single plant of a small-leaved begonia.

At the foot of the hill, where the path to Ginting Peris branches off, we dismounted, and sent our ponies back to Ulu Langat. After a rest of some hours, and tiffin, we started to walk the remaining four or five miles, all of which is uphill and pretty steep. After ascending a short way, a belt of

giant bamboos is passed through, and the character of the jungle begins to change. The undergrowth is not so dense. There is also a perceptible difference in the air—there is a freshness in it that that of the low jungle does not possess. The path here zigzags a good deal, owing to the steepness of the hill. At a height of 2,000 to 2,500 feet the jungle becomes more stunted: fewer large trees are seen and magnificent tree ferns reaching a height of 20 or 30 feet are fairly common. The path follows the crest of a spur of the main ridge, till it reaches a small Sakei *ladang*, or clearing, where a bamboo hut had been erected for the use of the Resident when he visited the hill some weeks before. Here there were also one or two deserted Sakei huts—mere attap sheds raised on poles some ten feet or so from the ground—and a small plantation of sugarcane. Here we spent the night very comfortably, although it was rather cold, and were lulled to sleep by the sweet wild music of the Sakei wind-organs. These are made of long bamboos, in each joint of which is made a small slit. They are lashed upright to the top branches of a tree, and when the wind blows act something like gigantic flutes. The sound they make is weird yet soothing.

On the morning of the 3rd January, we started at 7 A.M., and under the guidance of the only Sakei we found in the place (who informed us that he had been with CAMERON, the well-known Malayan explorer) went to the top of the ridge. All the valleys below were full of white mist, which marked out the courses of the streams, but as this cleared away under the influence of the morning sun, we had a magnificent view—Kwala Lumpur due West, and beyond that Klang Straits, and a little more to the northward Pulau Angsa. By climbing a tree we got a view all round, and found we were at the highest point of the ridge. The hill consists of a hog-backed ridge four or five miles long, running North and South and rising from South to North, with several spurs lying at right angles to it. The top of the ridge is only a few yards wide and the sides slope steeply down. An animal track runs along the crest, and the rocks and roots of trees between which it passes are in many places worn smooth by

animals brushing past. There were merely pools of rain water all along the path, and in the mud the fresh track of a rhinoceros. The jungle on and near the summit is entirely different from that lower down. The trees are stunted and gnarled, and they as well as the ground are thickly clothed with luxuriant moss nearly a foot thick and saturated with moisture. From the top of one of the trees I was able to get a view of the twin peaks of Gunong Hantu and Gunong Besar to the North-East, and setting the coolies to work we made a small clearing which enabled us to get a good view of the country in the direction of Kwala Lumpur and Klang to the West, and Ulu Selangor to the North-West, with all the intervening country. The ridge of limestone rocks near Batu stands out distinctly like a great wall or dyke running nearly East and West. While the men were at work clearing, I spent the time collecting plants. Amongst those I obtained, were a *Rhododendron*, a fine *Medinilla* with a pink flower and red undersides to the leaves, *Burmanicea longifolia* with beautiful pale bluish-white flowers hanging in clusters from the top of a stem a foot or eighteen inches high, the glorious *Nepenthes sanguinea* with pouches eight or nine inches in length, a *Sotterila*, and a large number of orchids—23 or more species, including a fine *Cælogyne* (probably new), one or two specimens of *Spathoglottis aurea*, and several species of *Anectochilus*, one a fine red one with a broad yellow band down the centre, and a green variety of the same, another of a beautiful velvety bronze colour with crimped edges, and another of a deep red-brown with pink edges and undersides. Few birds were seen on the top of the hill. A species of thrush of a uniform dark brown colour, and a few small flower-peckers were almost the only ones. Not having a gun I could not obtain specimens. Insects were not numerous. There were some flies and bees, and I got a single specimen of *Vanessa perakana* and one of *Atella sinha*.

Having brought a good supply of *kajangs*, we made a smart hut about 500 feet higher up than that where we spent the first night, and about half a mile from the crest of the ridge.

The next morning we set the coolies to work again at the clearings on the summit, while we followed up the course of the stream which ran near our camp. The bed was full of large boulders of granite, amongst which the stream eventually disappeared about a quarter of a mile from the crest of the ridge. I got some more orchids and two species of begonia growing on the rocks in the stream. In the gully in which the stream ran grew a very large species of *Amomum*, the fronds of which were 25 or 30 feet long, but I obtained no flowers of it.

By the evening the coolies had made a sufficient clearing to enable us to get a good view of Gunong Besar and Gunong Hantu. The view from the summit was really magnificent, extending over the whole of Selangor, a great part of Jelevu, and part of Sungei Ujong. That night we slept at our new camp, and were glad of a log fire in the evening. Early the next morning we started for Kwala Lumpur, walking the first six miles, and then, meeting our ponies, riding to within nine miles of Kwala Lumpur, where a dog-cart was waiting to take us on.

On the 15th January I started for a second visit to Bukit Etam. This time with only a Chinese boy and a few coolies. I slept, as before, at Ulu Langat for the first night. Having a minimum thermometer with me I was able to take some readings. At 8 p. m. on the night of the 15th, the temperature in the verandah of the Police Station was 68° Fahr. It had been raining all the afternoon since 4 p. m. The following morning the thermometer registered a temperature of 65°. At 7 a. m. it was 72°. This time I walked the whole way, stopping for an hour at the hot springs at Dusun Tuah, in the hopes of taking some more specimens of *Ornithoptera Brookeana*, but I only saw one, which I failed to secure.

At the bank of one of the streams I had to cross, I got a fair number of insects, including two specimens of *Clerome fannula*, several *Papilios*, and *Pieridæ*.

I reached the big hut at the Sakei clearings about 4 p. m.; my coolies about two hours later. They appeared not to be accustomed to hill work, and were quite done up, although

they had light loads. The temperature inside the hut at 8.30 p. m was 64° Fahr.

Next morning I started at 6 a. m. and moved up to the upper camp, and started collecting plants. I got a good number, including 25 species of fir and some good orchids, amongst which were a lot of *Anectochlis* similar to those I got on the previous trip; also a good *Bulbophyllum*. I gathered a sack full of orchids for Mr. RIDLEY of the Botanic Gardens, Singapore.

The temperature at 6 p. m. outside the hut was 63° Fahr. The minimum reading on the morning of the 18th was 56°, and at 11 a. m. the temperature was 64°. The thermometer was hung outside the north end of the hut, the hut being on the western slope of the hill. I collected another sack full of orchids, all from the top of the hill. I also found three specimens of a saprophyte—*Thismia aseroe*—and one of my men brought me a small land crab of a reddish-brown colour and about half an inch across. *

Finding my men had run out of rice, and having collected as much as I could carry, I decided to return to Kwala Lumpur that afternoon, and started at 2 p. m. in pouring rain, leaving my men to follow with the baggage. It rained all the way to Kwala Lumpur, which I reached at 9.45 p. m. My men came in the next afternoon.

THE FOLLOWING PLANTS WERE COLLECTED BY LIEUT. KELSALL ON BUKIT ETAM, SELANGOR :—

Dillenia aurea, Sm.

Illicium cambodianum, Kurz.

Pyrenaria Kunstleri, King.

Schima Noronhæ, Var. ? In fruit only. The leaves thick, rounded and blunt, with entire edges. Possibly a new species.

Rourea acuminata, Hook.

Allomorpha exigua, Grif.

Sonerila tenuifolia, Bl. A Javanese plant, not hitherto recorded from the Peninsula. [There were also two

* Since returning to Singapore, I have found a similar one on Bukit Timah.

- or three other species of *Sonerila* unknown to me.]
Bauhinia emarginata, Jack.
Argostemma inæquale, Benn. A good find, as it has not previously been found in the Peninsula. It is a native of Java.
Argostemma. Three other species apparently undescribed.
Ophiorrhiza sp. An unusually large species.
Lasianthus sp. nr. *cyanocarpus*.
Cephaelis Griffithii, Hook.
Ardisia villosa, Jack.
Rhododendron malayanum, Jack.
Diplycosia microphylla, Becc.
Tabernæmontana javanica, Miq.? A plant much resembling *T. corymbosa*, but with sessile cymes.
Didymocarpus marginata, C. B. Clarke.
D. platypus, C. B. Clarke. I suppose at least that this plant, which I have also from Kwala Lumpur, is this species. It fits the description well, but I have a specimen of what appears to be *D. crinitus* from Bukit Timah, Singapore, where it is common, labelled "*D. platypus*" from Kew. Mr. CLARKE states that the two species are very nearly allied, and as *D. crinitus* is rather variable in form of leaf, these two species may be the same.
D. sp. nov. A most remarkable plant, which is evidently new, but the specimens were all flowerless. The leaves were crowded and resembled those of such a fern as *Polypodium decorum*, being most elegantly lobed and cut, and bright red. The flowers were white. It would be a charming plant for cultivation.
Eschynanthus speciosus, Hook. This plant was also brought by Lieut. KELSALL, having been collected by Mr. SYERS in Ulu Selangor. It is a fine new addition to the Straits Flora, being hitherto only known from Borneo and Java.
Strobilanthes near *Maingayi*. Just the same as a plant so named from Kew, which was collected on Penang Hill by Mr. CURTIS.

Nepenthes sanguinea, Hook.

Elatostemma acuminatum, Benn.

Ficus hispida, Miq.

F. diversifolia, Bl. Two forms were collected—one almost typically the common form, the other var. *lutescens*.

Chloranthus officinalis, Bl.

Podocarpus neriifolius.

P. cupressina.

Spathoglottis aurea, Lindl.

Calanthe veratrifolia, R. Br.

Microstylis acutangula, Hook. The flowers of this were green, not white as described in Flor. Brit. India.

Otherwise the plant was similar.

Ærides odoratum, Lour.

Phaius pauciflorus, Bl. Apparently this rare Javanese plant, but the spur of the flower is somewhat larger than in BLUME'S figure. It is the first record of the plant from the Peninsula.

Dendrobium. New species of the *Cadetia* section. Will be described later.

Cælogyne. New species. A very lovely plant of the *Flaccidæ* group.

C. tomentosa, Lindl.

Cryptostylis arachnites, Bl.

Stenochasma urceolare, Griff.

Alpinia Rafflesiana, Wall. This charming orange-flowered *Alpinia* is not at all common. I only know one other locality for it, viz., at Toas, in Western Singapore.

Globba leucantha, Miq.

G. aurantiaca, Miq.

Susum malayanum.

Scleria malaccensis.

Heptaspis urceolata, R. Br.

Smilax hypoleuca, Mig.

Dianella revoluta, Br.

Burmannia longifolia, Becc.

Thismia aseroe, Becc.

Trichomanes pluma.

Hymenophyllum Neesii.

H. javanicum.

Prosaptia Emersonii.

Thamnopteris nidus-avis var. *phyllitidis.*

Lecanopteris carnosa, Bl.

Oleandra neriiformis.

Calostoma Junghuhnii, Schl. This curious fungus I also met with in the jungle at Pataling near Kwala Lumpur. It is known from Sikkim and Java and Sumatra.

Besides these, there were a number of plants of more difficult orders, which I have not yet identified, and a large quantity of living orchids, which have not as yet shewn signs of flower, some of which will certainly prove novelties.

This list gives a fair idea of the style of the Flora of this high region. There is a large Bornean element in it, but many of the plants are common also to the Perak mountains. The Flora is evidently every rich and interesting, and the collection shows what can be done in a very short and hurried journey by any one possessed of sufficient energy to collect as he goes along.

H. N. R.

ANATOMICAL NOTES ON MALAY APES.

BY

DR. A. KEITH.

—:O:O:—

I.—PRELIMINARY.



IN the autumn of 1889 my thoughts were directed towards the Pathology of Fever and Ague. Some years ago, research exposed in the blood an influx of a particular micro-organism during the attack. These microbes may be the actual fever-raising virus; or they may be the producers of the fever poison; or they may only be a concomitant of that poison. Be that as it may, the next important step in the evolution of our knowledge of the disease is to determine on what particular part of the body this poison immediately acts. Symptoms, remedies and alkaloids whose specific actions are known, point to a powerful affection of the great visceral sympathetic system of nerves. This system is probably the butt of the fever poison.

II.—OBJECT OF RESEARCH.

- (a) — It seemed well worth determining in a more exact manner than hitherto had been done, the anatomical relations of this nervous system, and I selected the quadrumana of this Siamese Province—Bangtaphan—lying at the eastern basis of the Peninsula.
- (b) — And as about 80% of the natives, along the banks of the river of that Province, suffer to a greater or less degree from enlargement of the spleen, it was of interest to see if the quadrumana kept them company in this. But by confining my attention within so narrow limits, I should have committed great havoc among them for a small result.

- (c)—So, after I had learned what might be called their normal structure, I looked for variations in their anatomy, and these variations—their amount and kind—signify a great deal.
- (d)—In the young and almost adult members I observed the later acquired stages of development, to which DARWIN'S Law of Heredity can be applied.
- (e)—And again, between the species of the same genus there are differences—the result of physiological processes. These are of great interest.

III.—THE QUADRUMANA OF BANGTAPHAN.

A.—*Hylobates lar* (gibbon). (Wa-Wa).

Dissected 2 Adult Females.
1 Adult Male.
1 Fœtus.

I take the white circumfacial ring with the white dorsal aspects of the hands and feet as the distinctive mark of the species. In the three specimens the amount and distribution of the interdigital web varied; so it does in all the species of monkeys I have examined; the colour of hair is of little value, it varies with the sex, it varies with the year, and it varies with the animal's age.

B.—*Papio nemestrinus* (pig-tailed baboon).

Dissected 3 Adult Males.
3 Adult Females.
3 Juveniles.
1 Fœtus.

A steadfast species. It does not answer exactly to the description in MASON'S "Burmah," having the ringed hair, and also a remarkable leonine appearance—in the male only. The baby is of a greyish yellow colour with dark brown hairs along the spine. There is a remarkable resemblance to the young of the stumpy-tailed monkey.

C.—*Papio arctoides* (the stumpy-tailed baboon).

Not nearly so abundant as the last species. Of these, as yet I have made no dissections, but at present I have a couple of tame ones. One is quite a baby with a white face, hair of

a dirty yellow hue, and a black stripe down its back, which is spreading laterally.

D.—*Papio cynomolgus* (crab-eating baboon).

Abundant near the sea side. No dissections.

I have had tame specimens of all three species. Mentally they differ immensely. A young female *P. cynomolgus* was full of mischief, fun and fight. *P. nemestrinus* was quiet and retiring, but given to lasting fits of anger; while *P. arctoides* was a most staid individual with a far-away gaze and his whole mind seemed steadily occupied sexually.

E.—*Semnopithecus albocinereus* (Schinz), (steel-grey monkey).

Dissected 13 Adults.

2 Fœtuses.

F.—*Semnopithecus femoralis* (Horsfield), (black monkey).

Dissected 1 Adult.

1 Fœtus.

The fœtus and babies of both are of a dull buff yellow. *S. albocinereus* is anatomically very variable, as one would expect from the immense number of nearly allied species. In Bangtaphan both of these species are very abundant. They have very distinctive cries; that of *S. albocinereus* is a deep guttural ejaculation, while that of *S. femoralis*, though evidently produced in a similar manner, has a distinctive metallic timbre, which medical men would describe as a "cracked pot" sound.

G.—*Nycticebus tardigradus*. (Slow loris).

Rare. No dissections.

Elucidation of some of the details has required dissections of other animals, amongst these:—

1. A young Bornean mias (orang-utan).
2. *Galeopithecus volans* (2 specimens) (flying lemur).
3. *Pteropus marginatus* (2 specimens) (bat).

IV.—VARIATION OF THE SPLEEN.

Within the bounds of health, the spleen varies in weight with the phase of digestion and period of the day. But I take it, that an increase of one-third in weight, beyond the average

relative weight of an organ, constitutes an abnormal condition of that organ. The accompanying table gives the actual and relative weight in most of the specimens I have examined, and chart I is a more lucid representation of the same. The spleen appears to be comparatively large in the mature fœtus, and enlargements are more common in adult females than adult males. In *Papio* and *Hylobates* the spleen seems to be wonderfully steadfast, while in the genus *Scmnopithecus* the variability is great—from .0010 to .0055. But in none could I say there was a resemblance to the human ague spleen. The nearest approach to this condition I have yet come across was the spleen of a young female orang-utan. I have a young stumpy-tailed baboon who suffered at one time greatly from anæmia, his ears and face having the almost characteristic hue of malarious cachexia, but the fluctuations in his temperature were slight— 102° to 102.75° F. Another very young stumpy-tailed monkey's temperature is 101.2° F., while the crab-eating monkey has a temperature of 101° F. Thus the results of the enlargement of the spleen, as an indication of fever in the monkeys, are negative.

V.—SOME OF THE VARIATIONS IN STRUCTURE.

In the jungle, remote from reference, it is impossible to give an adequate account of these. I will broadly note the chief ones.

Of the three species I have examined, *Scmnopithecus albocinereus* is the most variable; next *Hylobates lar*; while has *Papio nemestrinus* has a wonderfully steadfast structure. One would expect to find variation in those parts or organs:—

- 1st, which differ in nearly allied species and genera; or
- 2nd, which have undergone an increase or change of function.

The assumption of the semi-upright position by the lower quadrumana, and the downright position (using downright in distinction to upright as applied to man) by the higher quadrumana (*Hylobates* and *Simia*), has led to a change of function, in degree and kind, of many of the organs of the body. The rough diagrams *a*, *b*, *c* and *d* show partly what I mean. From these it can be seen that the line of support is continually changing. Thus the spinal column, especially about

the lumbar region, ought to show variations. This position further leads to a change in the mode of respiration, and we ought again to find variations in the bony structure of the chest, as in the sternum cartilages and ribs. And those parts are the most variable in the monkey organism.

(a)—*Variations of the Sternum.* (Breast-bone).

In the cartilaginous fetal sternum, centres of ossification appear, and it is the almost universal condition in the mammalian kingdom, for the first centre to spread and form the first piece or manubrium, while the second, third, fourth and fifth may or may not unite to form the sternum. But in the genus *Hylobates* (gibbons) the rule seems to be for the first and second pieces to unite and form the manubrium. Of the four specimens I dissected, two united in this manner (diag. IV), and two did not (diag. III). Professor DWIGHT called attention to this point (Journal of Anatomy, 1890) and gave the following cases, which I tabulate:—

Specimens in which the 1st and 2nd pieces united to form manubrium.	Specimens in which the 1st and 2nd pieces did not unite.
<i>Hylobates leuciscus</i> , 2 specimens, (Dwight). <i>Hylobates varié</i> , 2 Specimens, (Knox). <i>Hylobates syndactylus</i> , 1 Specimen. <i>Hylobates lar</i> , 2 specimens.	<i>Hylobates lar</i> , 1 specimen, (Dwight). <i>Hylobates lar</i> , 2 specimens.
Total, 7 specimens.	Total, 3 specimens.

Professor DWIGHT gives diag. I shewing the same condition in man, and diag. II shows it in *Hylobates leuciscus*. I found a very close approximation to the same form in a specimen of *Semnopithecus albocinereus* (see diag. VI). Professor DWIGHT is of the opinion that it means nothing more than that nature is on one of her accidental excursions, but the condition seems rather too wide spread for that. The sternum in the quadru-

mana is in a plastic condition, and the great development of the pectoral muscles—especially the manubrial portion of them—may determine the union of the first and second parts.

The sixth centre of sternal ossification appears late in juvenile life in the *Scmnopithec*i, and in one adult it had failed to put in an appearance (see diag. VIII).

(b)—*Variations in Ribs and Chest Walls.*

In the same genus the eighth pair of ribs seem undecided whether they should terminate in the fibro-cartilage in front of the xiphoid or not come within an inch of it (see diags. VI, VII, VIII, IX, X). In *Papio nemestrinus* the eighth pair of ribs always reach the sternum (see diag. VIII). In a specimen of *Scmnopithecus* the fifth rib terminated in a piece of cartilage to which some fibres of the diaphragm were attached (see diag. VI). In three specimens of the same genus, I came across rib-like developments in the fifth, sixth and seventh costal cartilages (see diag. VIII), and in all the adults, these cartilages were more or less ossified.

There are three floating ribs as a rule in *Scmnopithecus*, but in two specimens the tenth rib had joined those above and become a false rib.

(c)—*Variations in the Lumbar Region.*

In a fœtus of *Papio nemestrinus*, I found the transverse process of the first lumbar vertebra enlarged and representing a thirteenth rib. It was costal in dimensions and appearance (see diag. XI). There was a similar condition in the second lumbar vertebra of *Hylobates lar*, representing a fourteenth rib (see diag. XII).

In the gibbon the twenty-sixth vertebra affects the chief support of the pelvic bones, while in the monkeys and baboons it is the twenty-seventh. Diag. XV shows the peculiar condition found in one of the *Scmnopithec*i, on one side the twenty-seventh as usual is supported by the pelvic bones, but on the other side it is the twenty-eighth, the twenty-seventh on that side having a plain transverse process. Professor G. B. HOWES (Journal of Anatomy, July, 1890) shows a similar condition in a frog (diag. XIV). The above-mentioned specimen shows considerable aggregations of cartilage on its twenty-fifth and twenty-sixth transverse process as if at one time they had

had intentions of having connection with the pelvic bones. Such growth of cartilage on the tips of transverse processes is not of uncommon occurrence (diag. XVI). In a fœtus of *Hylobates lar* the transverse process of the twenty-seventh vertebra was in a much further state of development than the twenty-sixth (diags. XVIII and XVII).

In one specimen of *Semnopithecus* the abdominal aorta divided on the twenty-fourth vertebra instead of the twenty-sixth, and smaller degrees of variation are common.

The arrangement of blood vessels in the pelvis is very variable. The insertion of the diaphragm may shift down a vertebra and the muscles arising from the anterior aspect of the lumbar region are also liable to variation.

(d)—*Variations in Muscles of the Limbs.*

With the great increase and change of functions in the arms of the *Hylobates*, one would not be surprised to come across variations in the muscles of these limbs. The biceps becomes the great muscle of locomotion, and in diag. XIX we have what might be called the normal condition—one head coming from the coracoid process, the other coming from the top of the glenoid cavity; while in diag. XX we have the glenoid head coming as usual, while the coracoid comes from the edge of the biceps groove. A curious thing, in connection with this, is the transference of the triceps head of the latissimus dorsi to the biceps, thus from an extensor in the monkeys and baboons, it becomes a flexor in the gibbons (see diag. XX).

There are also numerous variations amongst the extensor muscles of the forearm and leg, while there are numerous forms of arrangement in the arteries and veins of the lower limbs.

(e)—*Variations in the Viscera.*

Diag. XXI shows the appendix vermiformis of the gibbons—long, narrow and worm-like; that of the orang is almost similar; diag. XXII shows that of the pig-tailed baboon, only a contraction of the cæcum; diag. XXV gives the usual form of the cæcum of the *S. albocinereus*, with hardly an appendix at all, but sometimes it resembles that of the pig-tailed baboon as in diag. XXIV.

Sometimes in dissections of the human body we find the kidneys united by an isthmus of renal matter, such an arrangement being called the horse-shoe kidney. Diag. XXVI shows such a variation in *Semnopithecus*. In *Hylobates* (see diag. XXXVII) the thyroid gland has a large isthmus lying in front of the trachea; in *Papio* this isthmus is very attenuated; in *Semnopithecus* it is non-existent, although sometimes a rudimentary one may be met with. The thymus gland sometimes exists in adults, and sometimes is absent in infants.

This is but an introduction to the more superficial variations in the quadrumana. We can never know their correct systematic position, nor the direction of their evolutionary tendency, nor their relations to man and to their brother quadrumana, until these be known. And as the all-pervading tropical jungle is gradually being replaced by a vegetation necessary for the sustenance of increasing humanity, the quadrumana will be replaced by man. The century that will hold the apes in its conservatories of rarities, is not far distant, if coming centuries are at all like the great nineteenth.

VI.—A FEW POINTS IN DEVELOPMENT.

(a)—*Ear.*

In each species of quadrumana the ear steadily maintains its characteristics. It attains its full growth long before maturity is reached, and still some of its transformations take place late in adult life. Diags. XXVIII and XXIX are from DARWIN'S well-known illustrations showing the occurrence in the human helix of a slight protuberance probably likely representing the aural tip of the lower animals. In *Papio cynomolgus* this tip is large and remains unfolded (diag. XXX). In *Papio nemestrinus* and *Semnopithecus albocinereus* one can see this tip folding in late in foetal life (diags. XXI and XXXII). In a specimen of *Hylobates lar* this tip was unfolded (diag. XXXII). The aural edge of the orang is folded to a greater extent than that of man and has no lobe, but on the other hand that of the *Semnopithecus* has the lobe and resembles closely the human ear. Man seems to have picked many of his parts in common with several of the quadrumana. The teeth of the *Semnopithecus* are more human than those of the orang.

(b)—*The Intertemporal Space and the Temporal Ridges.*

(See diags. XXXVI, XXXVII, XXXVIII, XXXIX).

The intertemporal space is bounded laterally by the temporal ridges from which rise the temporal muscles; in front by the frontal ridges, behind by the occipital ridges, from which rise the occipital muscles. In foetal life and youth this space covers nearly the whole skull, but as the animal grows older the temporal muscles climb up the sides of the skull, while the occipital scramble up behind and the intertemporal space melts away before them. Thus the temporal ridges may crush it out and meet, as in the adult orang, and form a ridge. As a general rule, it may be taken that *the larger this space the greater the relative size of the brain to the body.* A small intertemporal space means large temporal and occipital muscles; large temporal muscles mean large canine teeth (fighting teeth) or coarse food; coarse food means a big stomach, and a big stomach, as I shall presently show, means a small brain. Keep away the effect of the big canine teeth and this intertemporal space can be formulated so as to give a wonderfully exact indication of the relative amount of brain matter.

But the proper signification of these temporal ridges has an important value in classification. They certainly cannot have any generic value, and little stress can be laid on them as specific distinctions, seeing the ease with which they could be varied and the degree in which they do vary with age and sex. They do not reach their final position until well into adult life.

(c)—*Epiphysis of the Bones.*

Take the scapula (shoulder blade) of the young monkeys for instance. Along the base we find a border of cartilage. This border foreshadows the further development the bone is to undergo (diags. XL, XLI, XLII.) According to DARWIN'S Law of Heredity, these borders represent the latest evolutionary acquisitions of the scapula. Thus XL, XLI, XLII represent the latest scapular additions made to *Semnopithecus*, *Papio*, and *Hylobates*. *Semnopithecus* and *Hylobates* have had their infra-spinatus space augmented, giving them increased arm

climbing power, while *Papio* has had both supra and infra-spinatus enlarged equally, which is probably connected with their all-fours locomotion on the ground.

The lateral extension of the sternum seems to have a similar meaning. Indeed from the epiphysis we can read the late history of the animal.

Many of the tendinous insertions of muscles in adult life get assimilated with the periosteum of the underlying bones and thus there appears a difference in the adult and foetal insertion of a muscle. (See diags. XLIII and XLIV).

VII.—A FEW PHYSIOLOGICAL CONSIDERATIONS.

These problems are of even more interest than those of variation and development already given. But to understand their real meaning one must break through the conventional idea of human time. While the clock that marks the progress of things human has moved a century, the pendulum that registers the progress of the things of evolution has but swung a second. The failure to grasp a wider than vulgar view of time has kept many of the conservative naturalists from appreciating the final problems of evolution.

How the Stomach, Brain and Muscles are correlated.

The stomach of the three gibbons weighed, on an average, 903 grains, while on an average their contents (mostly green acid figs) weighed 4,100 grains. Their proportion to the weight of the body was respectively .0110 and .0462. Taking half-a-dozen of the *Semnopithec*i, their stomachs weigh 3,216 grains (relative weight=.0400) and contents 25,000 grains (.3200 of body weight). The table shows this more clearly.

	Body Weight.	Stomach Weight.	Contents Weight.	Relative Weight, Stomach	Relative Weight, Contents
Hylobates lar, 3 Specimens, ...	12.5 lbs.	903 grs.	4,100 grs.	.0110	.0462
Semno. albocinereus, 6 Specimens,	15.4 lbs.	3,216 grs.	25,000 grs.	.0400	.3200

That is, the gibbon carries a diet about one-twenty-fifth part of his weight and spends one-hundredth of his internal economy in carrying it and doing the first part of his digestion. On the other hand, this white-eyed, steel-grey monkey has to perform his locomotion with a diet load nearly one-third of his weight and expends as much as one-twenty-fifth part of his bodily substance on his stomach, as much as he does on his liver.

And it is from the stomach that the brain receives its impulse for work. And the progress is from a plentiful food supply that is difficult of digestion, such as the steel-grey monkey's staple and abundant diet amongst the bamboo leaves, to a scarcer and more inaccessible but more easily digestible food, as the green acid figs, and other fruit foods of the gibbon. On the one hand an intricate, large and expensive apparatus is used; on the other a simpler and smaller organ does the work. The extensive foraging excursions of the gibbon entail much more brain labour than the 'barn-door' feeding of the grey monkeys. The gibbon has added to his brain and subtracted from his stomach. And it was this simple mathematical calculation that brought us to where we are.

Such a statement of the facts of the case bear out. We tabulate the brain statistics of the above group of monkeys, and this is how the matter stands:—

	Average Bodily Weight.	Actual Brain Weight.	Relative Brain Weight.
Hylobates lar, ...	12.5 lbs.	1,607 grs.	.0187
S. albocinereus, ...	15.5 lbs.	1,113 grs.	.0120

The relative weight of the brain is in inverse ratio to that of the stomach, the gibbons spending $\frac{1}{50}$ of their tissue on brain + $\frac{1}{100}$ on stomach, while the grey monkey spends about $\frac{1}{50}$ on his brain + $\frac{1}{25}$ on his stomach. Man has reached the furthest point in this direction; he spends about $\frac{1}{100}$ of his economy upon a stomach, he gives about $\frac{1}{50}$ to his brain. In a young rang-utan (7 months), I found the brain weighing one-four-

teenth of its bodily weight, while its stomach weighed a little over one-hundredth.

Chart II looks as if it did not bear out the statement that increase of brain means decrease of stomach. The orang is quite out of line, and in connection with this I would point out that the development of the brain seems not to be governed by the law that regulates the growth of other organs. The relative amount of brain matter is greatest in foetal life; from that point onwards to full growth it diminishes. I tabulate the statistics derived from mother and child in the following species:—

Species.	Weight of Adult Brain.		Juvenile or Foetal Brain.	
	Actual.	Relative.	Actual.	Relative.
Female Hylobates lar,	1,620 grs.	.017	85 grs.	.12
Semno. albocinereus,	1,160 „	.016	460 „	.060
Papio nemestrinus, ...	1,340 „	.019	1,300 „	.216
Bitch and Pup, ...	1,180 „	.0056	595 „	.0218

Contrast this with the spinal cord development:—

Species.	Weight of Spinal Cord in Adult.		Weight of Spinal Cord in Baby and Pup.	
	Actual.	Relative.	Actual.	Relative.
Papio nemestrinus, ...	175 grs.	.0025	26 grs.	.0040
Bitch and Pup, ...	290 grs.	.0014	38 grs.	.00138


The other organs of the body increase much in ratio with the rest; it is so with the spinal cord, but the brain grows with no relation to the state of the other organs, save the stomach.

And the greater the cerebral development the greater is its disproportion to the other organs in foetal life.

I should have liked to give some statements as to the correlation of brain and stomach, or what is the same thing, internal economy with external surroundings, by means of the muscular system, but as my paper has crept to considerable length, I will defer these and some other curious details which I have observed in the quadrumana of this region.


1.—Chart showing Degree of Variation of Weight of Spleen in *Quadrumania*.

Sp. I. II. III. IV. V. VI. VII. VIII. IX. X. XI. XII. XIII. XIV. = Specimens of each Species.




— = Semnopithecus.
 = Papio.
 - - - - = Hylobates.


II.—Chart showing Correlation of Stomach and Brain.




* Relative Weight of the Brain to Weight of the Body.




I




II




III




IV




V




VI




VII




VIII




IX




X



XI



XII




XIII




XIV



XV



XVI



XVII




VII



VIII


XVIII




XIX



XX




XXII




4 3/4


XXI




XXIV




XXV




XXVI



H. lar



P. nemestrinus



XXVII


S. abbocinereus



XXVIII



XXIX



XXX^a



XXX




XXXI



XXXII



XXXIII



XXXIV




XXXV


PHOTO-ZINCOGRAPHED AT THE COL: ENG S. OFFICE S. S.




XXXVI



XXXVII




XXXVIII




XXXIX




XL




XLI



XLII



XLIII



XLIV



XLVI



THE FISHING INDUSTRY

OF
KRIAN AND KURAU, PERAK.

BY

A. T. DEW,

*Collector and Magistrate and Harbour Master,
Matang and Port Weld, Perak.*




THE Chinese fishing village situated on the South bank of the entrance to the Kurau River is said to have been in existence for more than thirty years. ONG GAN, one of the oldest inhabitants, has lived here for twenty-six years, and there were a good many fishermen then—about 15 houses—and the village has gradually grown to its present size. It now consists of from 70 to 80 houses, with between 450 and 500 Hokkien Chinese, all fishermen, working about 75 nets (*pukat*) and about 140 boats. Besides nets, they at present work fishing stakes (*blat*) as under:—

- 1 *Blat Jerumal.*
- 1 *Blat Telok.*
- 4 *Blat Kedah;* and about
- 30 *Blat Langei.*

Large fishing stakes will not stand here, as the mud is too soft; they require firmer ground.


The nets used are of two kinds—*pukat hanyut* (drifting net) and *pukat lengkong* (seine net). Besides these there are the hand nets used by prawn-catchers (*siring udang* and *sungkar udang*). A brief description of these stakes and nets may perhaps be of interest.

Blat Ferumal (an out-shore *blat*).




This is made facing the ebb tide. *ABCD*, enclosure of stakes, with a net-work of bamboo or rattan *sega* round it. *CF*, *DE*, fences of ditto guiding in the fish. *CD* are high nibong posts with pulleys on top. At the bottom of the enclosure is a rattan net called the "*daun*" (leaf); ropes pass through the pulleys on *C* and *D* and make fast to the outside corners of the "*daun*." Men watch in the hut at ebb tide, and, when they see enough fish are in, haul up the outer edge of the "*daun*" first, and the fish are caught. The inside end is then raised, and the fish thrown out into the boat, and the "*daun*" again set. This trap is fixed in about $3\frac{1}{2}$ fathoms of water, low spring tides.

Blat Telok (an in-shore *blat*).



Set facing and close to the shore. The water runs out at ebb tide, and the fish all run into the trap. In this trap are caught *Ikan Sembilan*, *Ketrak*, *Sablak*, and sometimes prawns. The construction is similar to the *Jerumal*, but the mesh of the net-work is much smaller, and the whole thing is made of much lighter material.


Blat Kedah (an in-shore *blat*). This is a smaller edition of the *Blat Kombang*, and is an in-shore fishing stake.



Made of bakau-wood stakes, the net-work of bamboo or rattan *sega*. Fish enter with the ebb tide, and are then hustled from the first and second *kurongs* into the third with a rattan arrangement called a *sisir*.

It is worked by two or three men.


Blat Langei (an in-shore *blat*).



Made of bakau (mangrove) wood stakes. The trap is a string net in the shape of a bag. There are six rattan rings on it, and the net travels up and down on stakes passed through the rings.


A man watches in a boat outside the net, and pulls it up to empty out the fish, and re-sets it; the operation is repeated as long as the ebb tide lasts.

Pukat Hanyut (drift net).




This net is from 80 to 100 fathoms long and 6 fathoms deep, of a mesh (*mata serat*) of about $1\frac{1}{2}$ inches square.

At one end it is made fast to a large float *F* (*polompong*) about $1\frac{1}{2}$ feet in diameter. The other end is made fast to the boat. The net is kept up by floats along the top. It has no weights at the bottom. It drifts along dragging the boat after it, and can be worked in both tides. Only one or two men are required to work it, and large fish are caught in it. The fish strike against it and get entangled.

Pukat Lengkong (seine net).

This net is from 80 to 100 fathoms long and 2 fathoms deep; it has floats along the top, and one large one attached to the centre of the top edge *F* by a two-fathom line; the bottom is weighted with tin. The big float (*polompong*) is first thrown overboard. The net is put out at ebb tide; one end is held by a large boat, and a small boat takes the other end round to the big boat, encircling the fish. The leads must be on the ground, or the fish escape underneath. When hauling the net into the boat, a man goes overboard and gets inside the net to tend the leads, see they do not get foul of sticks, &c., and let the fish out. The mesh is about $\frac{1}{2}$ inch, and all kinds of fish and prawns are caught, though the large kinds are thus caught but seldom, as they frequent the deeper waters. Five men are required to work it.

Siring Udang (hand prawn net).



This is made of coarse cloth (*siring*), has a shallow mouth and a deep bag at the end. It is attached to two sticks crossed as above, and the man, going into the water, places the fork *S* against his shoulder, and pushes it along in front of him on the ground, and the prawns are shovelled into the bag.


The *Sungkar Udang* is similar but made of string.

These are all the methods, at present in vogue, of taking fish at Kwala Kurau.

I am, however, acquainted with some other methods of taking fish followed in this country, and, as they may be any day adopted here, it may be as well to enumerate them.

The *Blat Kombang* is made on the same principle as the *Blat Kedah*, but is longer, and is an out-shore fishing stake, and will take up to eight men to work it, according to its size. The *Blat Batawie* (out-shore *blat*) is the same as the *Blat Kombang*, only the rattan netting is of a larger, coarser mesh, and is for catching large fish. It is set in channels in about 2 fathoms of water, low spring tides. The *Blat Lengkong* (in-shore *blat*) is the same as the latter, only the netting round it is much finer and closer, being made of *bertam*. With it is caught the *Ikan bunga ayer*, about 1 to 2 inches long, which is very good eating, and fetches, when salted, about 5 cents per catty.

The *Rawei*.




104 FISHING INDUSTRY OF KRIAN AND KURAU, PERAK.

A B is a long rope, running up to 200 fathoms in length.

At intervals of about $1\frac{1}{2}$ fathoms are placed bamboos to act as floats, *C D E F G H I J K L M*. From the rope *A B* are suspended, at the end of lines about 1 fathom long, hooks of iron, of a fairly large size; about 10 hang from between each float.

The hooks are unbaited, and the contrivance is either allowed to float about or is anchored at one end. Any *scaleless* fish of a moderate size trying to pass get hooked about the body. Fish with scales, such as the *Selangin*, *Sinahong*, &c., are protected by their scales and do not get hooked. It is set at ebb tide and taken up at flood; two or three men work it in a boat, generally Chinese. With it are caught *Ikan Yu* (shark), *Pari* (skate, three kinds, the black with white spots is the best), *Yu parang* (sword fish), *Duri*, *Goh*, *Bargu*, *Seludu*, *Pelotan*, &c. The last five are all of one family, and are much relished by the Malays. No *Raweis* are used at Kurau, but it is said that at Sungei Magat Aris there are one or two, and I am told they are used in the neighbourhood of Pasir Itam, Larut.

Pompong.



A B C D are two pieces of bamboo fixed to one another at right angles. Hanging from them is a net-work *C E D*, the *Penaju*, guiding the fish, entering by the mouth *C D* into the net trap *F G H I*. It is anchored to the ground, and set at ebb tide. It is an in-shore fish trap.

With the out-shore *blats* are caught *Ikan Selangin*, *Sinahong*, *Barwei* (three kinds, viz., *Barwei chermin*, *Barwei tambar* and *Barwei kedaywas*), *Duri*, *Pupus*, *Pelatar*, *Chincharu*, *Kurau*, *Temeras*, *Siakup*, *Pitcha priok*, &c. With the in-shore *blats* are caught *Ikan Glammar*, *Selangin*, *Chinkarong*, *Pupus*, *Selangat*, *Membeli*, *Udang kertas*, *Kiki*, *Yu*, *Pari*, crabs, &c.

The large *blats*, i.e., *Blat Jerumal*, *Blat Kombang*, and *Blat Batawie*, are owned by and worked by Malays; as is also the *Blat Kedah*. They make them themselves, and sell the fish to the Chinamen at the fishing village. Sometimes they get an advance from one of these Chinamen to assist them in putting up the *blat*, they agreeing in return to sell their fish to or through the Chinaman, the latter taking 10 per cent. of the value as interest, the Malay repaying the original debt out of the price fetched by the fish. The smaller *blats*, i.e., *Blat Telok*, *Blat Langei*, &c., are owned by and worked by the Chinese generally, and they make them themselves. The services of a Malay *Pawang* (sorcerer) are not called into requisition in choosing the sites for these *blats*, but in the case of *Blat Jerumal*, *Blat Kombang*, *Blat Batawie* and *Blat Kedah*, no one would think of commencing one until the spot had been fixed on by the driving of the first stake by a *Pawang Laut* (sea sorcerer) accompanied by the usual ceremonies (*Jamu*). The first is called "*Bacha doa*," in which, after the first stake has been fixed in by the *Pawang*, certain prayers are read from a book by the *Pawang*, the *Pawang* looking towards the prepared yellow rice (*nasi kunyit*) brought for the occasion, the others taking part in the ceremony holding their hands up in praying attitude, and casting their eyes to heaven.

The second is the "*Ratap*," or prayer, or lament, given in chorus, which every one knows by heart.

The third is more "*Bacha doa*."

The fourth is "*Minta jauh bala, akan datang datang rahamat yang kabajikan*" (asking that bad luck may be far from them and that good luck may befall them).

The *Pawang* then scatters some of the yellow rice on the waters, and the participants in the ceremony then eat the remainder of the rice, and the fowls, which have generally been killed and cooked on shore beforehand.

ONG GAN states that he has never, since he has been here, known the fish to get so scarce at Kuala Kurau as during the last two or three years. AH LIEW and TAN SIANG, also very old inhabitants (the former, the farm agent, has been here twenty-eight years, and is *the* oldest inhabitant), all tell the same story. They cannot account in any way for the fish getting short, but think perhaps it is because there are too many people fishing. Four years ago the place increased very much in fishermen, but during the last two years some have gone away. None of them seem to have any idea when or where the fish breed, or whence or whither they come and go, and when I suggested that if we could find out we should, perhaps, order a close time to give the fish a chance of breeding, they said that would be very hard on them, as what were they to do if not allowed to fish? In fact they want to eat their cake and have it as well. I told them it was done in Europe with good results. ONG GAN says he has only known the fishing get bad like this once before, about twenty years ago. I do not believe this, however, as at Telok Rubiah, about 6 miles further off, I was told it had happened often before, the fish coming and going. I then enquired into the worldly position of the fishermen. They are all Hokkiens, and mostly related to one another. They pay their own passages over from China to join their friends, and then go into partnership with some others in the ownership of a net (*Pukat Lengkong*). Each net is divided into $7\frac{1}{2}$ shares, and there are six men to a net. The owner has $1\frac{1}{2}$ share extra for owning the net, and he works with the other five, and they all, including the owner, get one share each. Thus the owner has altogether $2\frac{1}{2}$ shares. One

of the six men stops in the house and salts and dries the fish, cooks the food, &c. The cost of the house, net and boats amounts to about \$350, and the owner mentioned above owns the whole concern, including all furniture, tubs, &c.

In good seasons one net (*Pukat Lengkong*) can get fifty or sixty pikuls of salt fish in a month, but in bad seasons as little as eight or ten pikuls.

The fish is classed for export in three classes:—

No. 1.—Dried prawns, which fetch from	\$7.50 to \$10.00 per pikl.
„ 2.—Large dried salt fish fetch from	\$3.00 to \$ 7.00 „
„ 3.—Small dried salt fish (formanure)	\$0.85 to \$ 1.30 „
Blachan,	\$1.50 to \$ 2.50 „
Prawn shell manure,	about 50 cents per pikul.

The fish manure is sold to sugar planters for manuring the sugar-cane, and the prawn shell manure is chiefly used for manuring nutmeg trees, &c.

The men are nearly all opium-smokers, but are extremely healthy and of fine physique, as men employed on the sea generally are; and there is a total absence of those nasty sores and ulcers on the legs, &c., so common among the agricultural and mining coolies.

The fishing is at its best during the 10th, 11th and 12th and 1st Chinese moons, roughly, from November 3rd to March 12th, *i.e.*, during the North-East Monsoon, when it is fine, dry weather with but little wind or storms and waves to interrupt operations. The fishing is at its worst during the 5th, 6th and 7th moons, roughly, from June 9th to September 5th, *i.e.*, during the South-West Monsoon, when it is very wet and windy. Every month no fishing with *pukat lengkongs* can be carried on between the 13th and 21st of the Chinese moon, and between the 28th and 5th, *i.e.*, during the high spring tides; but between the 12th, 13th, 14th, 15th and 16th plenty of fish are to be got in the *jerumals* and *blats*, but not in the *pukats* (both kinds). There is thus a period varying from sixteen to eighteen days every month during which these nets cannot be used.

Generally speaking, the higher the tides get the less fish are to be found. During these periods of enforced idleness the

men make and mend their nets, cut their supply of firewood, &c.

They say that the *Ikan kurau* and *Ikan tambrai* have got scarce. It is a big fish, which, salted, fetches eight cents per catty, and in the wet season 12 cents. *Ikan selangin* and *Ikan sinahong* are also getting scarce. I suggested that perhaps the cause of this was that they caught all the young ones, but was told that this was not the case. The parent fish are only, as a rule, caught in deep waters in the out-shore *blats* and *jerumals*. They do not come into the shallow waters, where *pukat lengkong*, &c. are worked, and the young which accompany the big fish can go in and out of these *blats* without being caught.

The people complain of the heavy taxes imposed on their industry. They state that in the old days, under the Sultan of Perak, the first tax imposed was one of \$12 per annum on each *pukat lengkong*, paid to the farmer. After two or three years this was found to be too heavy and was reduced to \$8. The *jerumals*, *blats*, *pukat hanyuts* and other contrivances for taking fish were all free. No other payments whatever were imposed. The Opium Farmer at Kuala Kurau paid about \$100 per annum to the Sultan, and imported and sold as much opium as he liked. He cooked his own opium. There were, however, but few fishermen there then. They now complain that they have to pay one-tenth of the value of their fish to the farmer, pay \$10 per annum for every fishing stake, whether large or small, out-shore or in-shore, and that each person using any other means of catching fish must pay for a personal license of \$1.20 per annum. Thus the six men manning a *pukat lengkong* each pay \$1.20 per annum, total \$7.20 per annum, and the two men working a *pukat hanyut* pay \$2.40 between them. The *jerumal* and *blat* men, besides having to pay \$10 per annum for their *blat* licenses, also pay \$1.20 per annum each for a personal license. A *blat* or *jerumal*, requiring, say, six men, will thus cost \$7.20 per annum extra. This was levied by mistake by the clerk misunderstanding the rule, and will be refunded to them. Besides this they have to pay 25 cents

for a port clearance, whenever their *tongkangs*, laden with fish for export, leave the port, and they have to pay annually for the numbering of each large *pukat* boat 80 cents, for a small ditto 45 cents, and the large and small *sampan kotas* are charged the same. For the numbering of *tongkangs* \$1.30 is charged. They also have to pay \$1 for the cost of numbering of each *blat* or *jerumat*. They are also charged 10 cents per foot frontage (Land Revenue) for their houses. I should think this, on an average, would come to about \$2 per annum per house. This was imposed from January 1st, 1886, and they think it very hard that they should be charged town allotment fees for occupying a part of the useless mud bank of the river.

As regards the payment of one-tenth of the value of the fish to the farmer, they state that they have come to an arrangement with the farmer, by which they pay him \$1.40 per month for each *pukat lengkong* instead. If the farmer did this from pure philanthropy it was very good of him; perhaps he came to the conclusion that, if he claimed his full pound of flesh, he would kill the goose that was laying the golden eggs.

It appears that, previous to December, 1885, no licenses were imposed on nets and fishing stakes in the district, though in Larut fishing-stake licenses seem to have been imposed since 1880, but no fishing-net licenses until last year. In that month, however, the Assistant Resident (Mr. CREAGH) issued the order, which I have marked *A*, annexed, and which was issued as a notice by the Magistrate and Collector, Dr. LEECH. The fishermen of this district (and I believe of other districts as well) petitioned against it, with the result that the order was suspended for further consideration, and, after much deliberation, extending over nearly a year, during which period no payments for licenses at all were imposed, the Assistant Resident issued the new order dated 1st October, 1887, which I have marked *B*.

The following table will show the difference between the old and new rules:—

110 FISHING INDUSTRY OF KRIAN AND KURAU, PERAK.

Old Rule dated December, 1885.		New Rule dated October, 1887.	
Large fishing stakes per annum, ...	\$ 10 00	Large fishing stakes per annum, ...	\$ 10 00
Small fishing stakes per annum, ...	8 00	Small fishing stakes per annum, ...	10 00
Large Pukat (<i>Lengkong</i>),	10 00	Say, six men @ \$1.20, ...	7 20
Small Pukat (<i>Hanyul</i>),	8 00	„ two men @ \$1.20, ...	2 40
Prawn (<i>Siring</i> or <i>Sungkar Udang</i>), ...	4 80	„ one man @ \$1.20, ...	1 20
Total, ...	40 80	Total, ...	\$ 30 80

By the new rules the fees are payable half-yearly.

It will be seen from the above table that the new rule makes a total reduction of 25 per cent. on the old rule.

As regards the statement that the fish has got scarce during the last two or three years, the following totals of exports from Kwala Kurau and Telok Rubiah, taken from the Customs Clerk's book at Kwala Kurau, will show :—

Description.	1885.		1886.		1887.	
	Weight.		Weight.		Weight.	
	Pikuls.	Catties.	Pikuls.	Catties.	Pikuls.	Catties.
Dried prawns, ...	2,771 00	32,116 50	4,414 00	37,993 60	971 00	10,362 00
Salt fish, ...	8,865 00	20,555 30	3,214 00	8,573 40	8,997 75	22,151 40
Blachan, ...	2,383 00	5,527 20	354 50	689 00	455 00	1,242 00
Fish and prawn shell manure,	17,640 00	15,740 10	1,530 00	2,172 00	9,317 00	9,995 00
Totals, ...	31,659 00	73,939 10	9,512 50	49,428 00	19,740 75	43,750 40

From this it will be seen that the export of *salt fish* was less than half in 1886 of what it was in 1885, but in 1887 it had risen again to more than it was in 1885.

Dried prawns were nearly double in quantity in 1886 of what it was in 1885, but in 1887 it had fallen to about a third of what it was in 1885.

Blachan was only about one-seventh in 1886 of what it was in 1885, and the quantity had only slightly risen again in 1887, though, from its better quality, it was worth nearly double that of 1886, though it is still less than a quarter the value of that of 1885.

Fish and prawns shell manure was in 1886 only about one-tenth of that in 1885, but again rose in 1887 to about one-half of what it was in 1885.

Comparing the two years, 1886 and 1887, there seems to have been an improvement in 1887 over 1886 under every heading except prawns, and the fish about which the people specially complained is more than double what it was in 1886, and more even than it was in 1885. It would, therefore, seem, from this return, that the prawn-fishing is really all they have just cause to complain about at present. I am inclined to think it is only a temporary scarcity; but, if it turns out, after watching the returns for a few more years, to be permanent, I do not think it will be a matter for great surprise, considering the constant shovelling up of these creatures by the million in the *Sirings* and *Sungkars*, besides the large number that are caught in the *Pukat Lengkongs* and small *Blats*.

It is only in 1887 that they fell off; had the quantity remained anything like that of 1886, then the total value of the fishery export for 1887, instead of being only \$43,750, would have been about \$63,000, only \$10,000 short of 1885, an increase of about \$14,000 on 1886. As it is the grand total for 1887 only shows a decrease on 1886 of less than \$6,000.

Prawns are the most valuable product of the fishery, fetching, as I have stated in paragraph 28, from \$7.50 to \$10 per pikul, and, consequently, a bad prawn year makes an enormous deficit in the fishermen's profits.

Telok Rubiah.

Telok Rubiah, situated on the coast about 6 or 8 miles to the South of Kwala Kurau, is a small fishing village of about eighteen houses, some of which are, however, unoccupied at present, with about thirty-five or forty fishermen—Hokkiens. The oldest inhabitant, KOW CHUNG, the farmer, has lived there for twenty-four years. He states that the fish has been scarce for about two and-a-half years, but he has known this happen before several times, the fish coming and going. Some of the people have, in consequence, gone to Pangkor (Dindings), Bernam, and Lower Perak, until good times return. Before, there were thirty or thirty-five *Pukat Hanyut* nets here, but now there are only seventeen or eighteen. The people here only use *Pukat Hanyut* and *Blat Langei*, no other *Blats* or *Ferumals*; they also use the *Siring Udang* a good deal.

The sea here is constantly encroaching, washing away the land. During KOW CHUNG'S time, the site of the village has had to be shifted four times, being gradually washed away, and they will have soon to move again for the fifth time.

There is no river or stream here, and, after their supply of rain water, kept in tubs, has given out in the dry season, they bring water either from Matang, Larut, or Pulau Jerejak, Penang. The health of the fishermen, like that of those at Kwala Kurau, is very good, the abominable smells proceeding from the fish and prawns in process of being cured, and more especially from the manufacture of *Blachan*, and in the thick of which they live and sleep, having no neutralising effect on the really healthy out-at-sea part of their lives. I think Chinamen like smells, and that they do them good. I have felt before now considerable inconvenience when three miles to leeward of one of these fishing villages, and I unpleasantly remembered my two hours' sojourn in the close atmosphere of KOW CHUNG'S house, surrounded with packages of *blachan*, &c., for days afterwards.

The people here pay no rent to Government for the land on which their houses are built; it would be hard if they did, seeing that the sites of their houses are washed away every few years. Neither do they pay one-tenth of the value of their

fish to the farmer. I suppose the farmer could claim it if he liked, but has never yet asked for it. The people here, with the exception of grumbling a little at the scarcity of fish, and at having to pay for licenses for nets, boats, &c., had little to complain of. The farmer, old KOW CHUNG, however, stated that, in the old days of the Sultan, he only had to pay \$1 per ball duty on opium and was allowed to cook it himself. Now he has to pay \$7 per ball to the coast farmer. There were no other taxes at all formerly, and now the proceeds of the fishing only produce about \$8 per month profit for each man—a large fall from the \$20 and more sometimes gained by each in good times, but nevertheless sufficient, I think, to live on comfortably with a little to spare. The farmer has to pay to the farmer at Kwala Kurau \$25 per month for his license to sell chandoo. He buys it at \$1.35 and sells it at \$1.50 per tael. He says he is now losing money, as he only sells about 110 taels per month.

Three years ago the crocodiles were very bad here. One man had his leg bitten very badly and was sent to Penang, where it was amputated, and he is still living and has gone to Trong, Larut, and lives with Malays, attap-making. Three men were afterwards bitten in one day. The Malay paddy-planters at the place (only about seven families) then arranged to call a *Pawang* from Perak to charm away the crocodiles and to bring back the former prosperity to their crops. Each Malay house subscribed 25 cents, and every Chinaman in the fishing village subscribed 30 cents. This was to pay for the buffalo for the sacrifice, the *Pawang's* expenses being paid by the Malays. The son-in-law of the former Datoh ALI of Kurau called the *Pawang*. The ceremony took place, and there have been no crocodiles there ever since and the Malays' crops improved.

There is no separate account kept of the export of fish from Telok Rubiah. They declare their cargoes at Kwala Kurau, running in there to take out port clearances, and the declared amount of fish is entered in the Kwala Kurau book as if it was Kwala Kurau fish.

Tanjong Piandang.

At Tanjong Piandang there is a fishing village of about twenty houses with thirty or forty fishermen. They only catch prawns. About fifteen years ago, in Datoh ALI'S time, there were about two hundred men there, and they used nets also; and people from Nibong Tebal used to come and put up *Blats* there. The people, who had always had the reputation of being a lawless, turbulent lot, and from whose numbers it was notorious that gang robbers, thieves and other criminals were recruited, at last brought matters to a crisis about four years ago by attacking the Police when they came there to make an arrest. As a punishment the Government sent a force of Police there to burn the village down, which was accordingly done. At that time there were about one hundred fishermen. The village has been partially rebuilt, but now, as I have said above, there are only about thirty or forty men, all (but one) Te-Chius. The exception is an old Hokkien, named OOI IN, who has been living there for thirty-nine years and is seventy years old—the oldest inhabitant.

In Datoh ALI'S time the farmer paid \$10 per month and was allowed to cook his own opium and spirits. The fishermen then paid no taxes at all.

The people complain that the prawns have got very short in quantity. They have got scarce before, but never to such an extent as during the last four years. They say that, before the last four years, each man could get \$6, \$7 or \$8 profit per month, and that now often they do not get enough to buy sufficient rice to eat. They cannot account in any way for the prawns getting short, but do not think the big fish eat them and cause the scarcity.

There are a good many crocodiles, and perhaps they eat them, they say. A man was killed by one about eight years ago, but no further harm seems to have been done up to about three years ago, since which time two men have been bitten, but no one killed.

Sometimes they get a *Pawang* to come and pray and charm

them, and the reptiles get good for a time, but it does not last long.

Like Telok Rubiah, the village has had to be moved a great many times, owing to the encroachments of the sea. The people in this fishing village are not charged house rent by Government, nor does the farmer claim one-tenth the value of the fish taken, as I suppose he might do if he liked.

These fishermen were all originally agricultural *Sinkheh* coolies, and, on their agreements expiring, took to this means of making a living, not being strong men nor fitted for work requiring much strength nor for carrying heavy weights.

(NOTE.—For export returns see table at end.)

Sungei Bharu.

At Sungei Bharu there are only about twenty Chinese fishermen, all prawn-catchers. There were many more formerly, and the people complain of the prawns having got scarce lately. The export return is included in that of Tanjong Piandang.

Bagan Tiang.

At Bagan Tiang is a very small and miserable Chinese fishing village of about fifteen houses. They are all Te-Chius, and work six *Pukat Lengkong* and *Siring Udang*, and carry fish to Parit Buntar for sale. There are only about thirty of these fishermen. There are, however, 19 *Blat Jerumal*, 3 *Blat Kombang*, 1 *Blat Telok*, worked by Malays under advances chiefly to Chinamen in Penang. There must thus be a very considerable amount of fish taken here, and the reason that the export return shows so little is that it is only cured fish that is entered in the export book, and nearly all the fish taken here is taken direct from the *Blats*, &c., over to Penang, and sold fresh, it fetching a higher price than salt fish. Boats also come over from Penang to the fishing stakes, and buy up all they can get and run straight away with it.

Some time ago a small steamer tried its luck in running over here for fish, but I believe it was a losing venture. Some

of the fish is also taken up to Simpang Ampat, Nibong Tebal and Parit Buntar and sold fresh.

Many of the people here declare the fish is much scarcer than it used to be. Unfortunately the export return for this part of the district throws no light whatever on the subject, for, as I have said, nearly all the fish is taken away (fresh) without ever being reported.

The export account, such as it is, however, is quite untrustworthy, being kept by careless Malay clerks over whom no supervision seems to have ever been exercised.

For instance, in glancing through the export account book for 1886, I noticed an entry "4,000 catties of dried prawns, value \$40." Knowing that dried prawns are worth about \$10 per pikul, I at once brought this to the clerk's notice, and he agreed with me, that it was evidently meant for \$400. Thus, in this single instance alone, the return shows only one-tenth the value it should have, and, if this is a fair sample of the way the accounts are kept, they are not much to be depended on. The clerk, however, says this mistake was made by his predecessor.

The farmer here does not exact one-tenth of the value of the fish taken, as I suppose he is entitled to do. He would need to be a sharp man to collect it from the *Jerumal* and *Blat* men, who run their fish over to Penang.

Penghulu LEBBY LAH, who has lived there thirty years, says it is always the same, the fish come and go with the seasons (monsoons); that many people declare the fish has got scarcer altogether, but that he cannot tell. If they are scarcer, perhaps it is because there are too many people fishing, he says. People come over from Penang and fish here, and there are all our own coast people at work too.

The Krani tells me that only nineteen licenses to catch fish with hook and line (*panching*) have been taken out. Now, according to Mr. CREAGH'S new regulation (marked *B*, paragraph 2) each person using any other means (besides stakes) of catching fish or prawns is to take out a license, for which he will be charged 60 cents half-yearly. A number of Malays came to me about this when the new regulations were

issued, with the usual story about being very poor—*baniak miskin*, &c.,—and that it was very hard lines, &c. I could only explain to them that Mr. CREAGH had taken a whole year to decide about this matter, and that it must, therefore, have been well thought out, and that paragraph 2 was so clear that *every one taking fish by any other means than blats* was to pay 60 cents half-yearly, that they undoubtedly must take out licenses for line fishing (*panching ikan*).

Now, it is ridiculous to suppose that there are only nineteen people line fishing on all this coast, and the inference is that just as many people fish as before, and that the order is simply ignored. The Malays say, with truth, that it is hard that they cannot go out and catch a fish for their dinners without taking out a license; but how is one to draw the line between such cases and those of the people who go out three or four in a boat, say, daily, and who all day long keep three or four lines over the side and take perhaps a pikul of the best fish, *Selangin* or *Sinahong*, &c.? The license only comes to 10 cents per month per man, and the sale of one catty of this fish about pays for it. Besides, the chances are that the man who only goes out, honestly enough perhaps, with the intention of catching a fish for his dinner, is not likely to resist the temptation of catching as many as he can if he finds them biting well. There are, I know, a great many men always employed in this way, and they sell their fish out at sea to the *jerumal* or *blat* men or the Penang boats, and make a good living out of it.

The Malays and Chinese have both conjectured that the alleged scarcity is caused by too many people fishing. If so, to repeal this light tax would act as a further incentive to every man, woman and child to renew their efforts to make matters worse; whereas a little judicious raiding by the Police now and then, leading to the occasional fining of one of the offenders, will, I have no doubt, stop a good deal of it, to the improvement of the fishery for those who invest capital in it and make a regular business of it, and to the increase of the revenue. I do not see any reason why the trade of the regular professional fishermen, Chinese and Malays, who pay

their taxes and other licenses honestly for their privileges of using nets and fishing stakes, should be ruined, or, let us say, damaged, by men paying nothing.

Final Remarks.

If any reliance can be placed on the Kwala Kurau export account above, it would seem that the actual quantity of *fish* has not much, if at all, fallen off in this district, but that the *prawns*, the most remunerative part of the fishery, have decreased very much during the last two years. but that there is nothing to show whether the decrease is of a permanent or only of a temporary nature.

If, in the course of two or three years more, the decrease is found to be of a permanent nature, the nature and habits of the prawn in these seas will have to be studied, in order to protect it and encourage its breeding, if necessary by establishing a close season and the same will apply to the fish, if it also should get alarmingly scarce.

However, notwithstanding the favourable aspect of the salt fish return in the Kwala Kurau book, there is no doubt that there is a general complaint of the scarcity of fish throughout the district, whether wholly true, partially true, or false, I cannot say, but I am inclined to think that there is a good deal in it, as I hear the same report from Penang; and, when on a recent visit to Pangkor, I was told that there had been a great scarcity of fish ever since I gave up charge of the Dindings, in the middle of 1886, and there are now not half the number of fishermen there that there used to be.

I would suggest that *Outshore Stakes*, *i.e.*, *Blat Jerumal*, *Blat Kombang*, and *Blat Batawei*, should not be fixed in depth of more than $3\frac{1}{2}$ fathoms (low water spring tides), nor should each fishing stake be situated at less than 400 fathoms from any other fishing stake, nor should it exceed 300 fathoms in length.

Inshore Stakes, *i.e.*, every other kind of stakes, also *Pompons*, should not be fixed in a depth of more than $2\frac{1}{2}$ fathoms (low water spring tides), nor should each fishing stake be situated less than 200 fathoms from any other fishing stake, and should not exceed 120 fathoms in length.

It will be seen that an outshore *blat* is more than twice the length of an inshore *blat*, and it is, therefore, only fair that the same charge should not be made for both.

I would further suggest that the use or possession of the murderous *Rawel*, with its 1,000 or 2,000 hooks, should be made illegal.

ARTHUR T. DEW.

KRIAN, PERAK,
21st April, 1888.



1204 & 1205

85.

A.

Notice is hereby given that all fishing stakes and nets on the Coast of the Kran and Kurau Districts will be licensed from the 1st January, 1886, at the following rates:—

Fishing stakes (large) and seine nets,	\$10.00	a year.
„ (small)	\$ 8.00	„
Pukat nets (small)	\$ 8.00	„
Prawn nets	\$ 4.80	„

The boundaries of the district are as follows:—

On the North a line in continuation of the boundary ditch, and on the South a line running due East from the most Southern point of Tanjong Blanga.

Any person wishing to erect a permanent fishing stake must obtain permission of the Chief Officer of the District before doing so. Any person found fishing within the above limits without a license will be prosecuted.

C. LEECH,

Collector and Magistrate, Kran.

Kran, 16th December, 1885.

B.

Government Notification.

1. Owners of fishing stakes, as at present, to pay \$5 half-yearly for each set of stakes, without regard to the number of men employed.
2. Each person using any other means of catching fish or prawns to take out a license, for which he will be charged 60 cents half-yearly.
3. This license to be not transferable, and to entitle the holder to fish with any kind of net he pleases, fishing stakes excepted.
4. Any breach of the above rules to be punishable with a fine not exceeding \$25 or three months' rigorous imprisonment.
5. Every fisherman to be required to have his license on his person while fishing or in the boat he is using.
6. Any breach of rules to be punishable with a fine not exceeding \$5 or 14 days' rigorous imprisonment.
7. Every one in a fishing *kongsi* except the cook to be held a fisherman and required to have a license.

C. V. CREAGH,

Assistant Resident, Perak.

ASSISTANT RESIDENT'S OFFICE,
Perak, 1st October, 1887.

Comparative Statement of Export of the undermentioned Articles for the years 1885, 1886 and 1887.

District.	Description.	1885.			1886.			1887.					
		Weight.		Amount. \$ c.	Weight.		Amount. \$ c.	Weight.		Amount. \$ c.			
		Pikuls.	Cattles.		Pikuls.	Cattles.		Pikuls.	Cattles.				
B. Tiang, ...	Blachan, ...	727	00	985	50	13	00		
T. Piandang, ...	Do., ...	47	00	118	00	11	00	16	50	76	00	228	40
B. Tiang, ...	Dried Prawns, ...	87	50	914	60	113	06	114	00	63	08	617	00
T. Piandang, ...	Do., ...	277	64	3,172	50	53	00	1,397	92	229	50	3,106	60
B. Tiang, ...	Salt Fish, ...	9	00	6	00	41	00	89	75	123	50	895	50
T. Piandang, ...	Do., ...	27	00	66	00	44	00	288	75	118	30	589	10
B. Tiang, ...	{ Fish & Prawns { Manure,	38	40
T. Piandang, ...	Do.,
Total...		1,166	14	5,262	60	275	06	1,945	32	610	38	5,436	60

A DAY AT CHRISTMAS ISLAND

BY

H. N. RIDLEY, M. A., F. L. S.



In a naturalist's eyes there is always a peculiar interest attaching to an oceanic island. For, owing to its isolation, we are able to obtain many clues to obscure points in the distribution and development of species, by investigating its fauna and flora.

In speaking of an oceanic island, I mean one which has, as far as we know, arisen by volcanic or other action from the sea, and is not merely a detached portion of an adjacent continent, or of a continent which has in lapse of time been destroyed. Under this category come most, if not all, of the chain of islands which lie scattered throughout the Atlantic Ocean far away from land, such as the Azores, Canaries, Madeira, Cape Verde, Fernando de Noronha, St. Paul's Rocks, Tristan d'Acunha, St. Helena, South Trinidad and Martin Vaz, the Crozets, Diego Garcia, and, farther East, Cocos and Christmas Islands. The plants and animals inhabiting nearly all these islands have now been pretty well explored, and good accounts, especially of the plants, have been published by various travellers and naturalists. Mr. HEMSLEY has collected all the work done on the botany of these islands, and added much thereto in the great work of "The Voyage of the *Challenger*." One or two, however, remain to be more thoroughly examined, as at present but little is known of them. These are South Trinidad and its companion Martin Vaz lying off the coast of Brazil, and Christmas Island, now a British Colony attached to the Straits Settlements.

Christmas Island was visited by H. M. S. *Flying Fish* in 1887, and by H. M. S. *Egeria* in the year 1888, and Mr. J. J. LISTER, who remained on the island for a week, made considerable collections of the plants and animals there, accounts of which were published in the Journals of the Linnean and Zoological Society. Much, however, remains to be done, and it was with great pleasure that I found myself last August in H. M. S. *Redpole* bound for Christmas Island. Unfortunately the vessel was only able to remain there for the inside of one day, so that I could not get more than a glimpse of its flora and fauna. Many of the trees and shrubs were out of flower at that time too, so that I was unable to get specimens of them. I managed, however, to make several additions to the list of plants published by Mr. HEMSLEY from Mr. LISTER'S collections. Among the most interesting of which was a very pretty new orchid, *Saccolabium archytas*.

Viewed from the sea, near Flying Fish Cove, the island appeared as a long straight ridge rising abruptly from the sea, and densely covered with high forest, the deep green of which is dotted with white specks—the innumerable nesting sea-fowl—while here and there are orange-red patches of the bloom of a handsome coral-tree (*Erythrina*).

The whole island is about nine miles across, and its highest point is nearly 1,200 feet above sea level. The cliffs are mostly abrupt, but in some places are patches of beach, composed of broken coral, shells and stones. The surrounding seas are very deep, and there are but few spots shallow enough for anchorage of a large ship.

On the beach at Flying Fish Cove, Mr. ANDREW ROSS has established his little colony, at present consisting of 15 persons, and is cultivating coffee, coco-nuts and vegetables, for which purpose the brown soil saturated with guano is well suited.

The core of the island is said to be composed of volcanic rock, and I saw a few pebbles on the beach consisting of some kind of trap, but though I ascended to the top of the ridge, I saw none of the rock *in situ*. The whole island is covered with coral-limestone, a hard yellowish rock emitting a ringing sound when struck. In many places it occurs in the form of

cliffs rising one above the other, shewing that the island has been slowly and gradually elevated from the sea bottom. Where the upper surface is exposed it weathers into pinnacles and curiously shaped angular projections, and though weathering externally into holes, the inner portions are usually homogeneous. I did not see any fossils in it, but the distinguishable remains of shells and coral would probably be found, if sought for.

Exactly similar rock occurs in Fernando de Noronha, an island lying off the East coast of Brazil, which I visited in 1887, and with which Christmas Island has many points in common. This coral rock has been described by Mr. BRANNER in some notes on the petrology of Fernando de Noronha, published lately in the American Journal of Science as being derived from sand-dunes hardened by carbonate of lime. He shows, however, by analysis that it consists almost entirely of carbonate of lime with a very small proportion of silica which would at once negative that suggestion. Furthermore, it was quite easy to find at Fernando de Noronha spots where the coral-reef, which was still growing at the outer edge in the sea, passed into the compact ringing rock with no distinguishable organic remains in it. It is true that on Fernando de Noronha there are sand-dunes which might, and indeed had in one place, become hardened more or less into rock, not much resembling, however, the limestone, but in Christmas Island there is no level place on which could develop sand-dunes sufficiently extensive to form the immense mass of rock of which the greater portion of the island consists.

On our return from Christmas Island to Singapore we stopped for an hour or two at Anjer Point in Java, opposite the volcanic island Krakatau, of which so large a portion was destroyed by an eruption a few years ago. Anjer Point was then struck by an enormous wave which overthrew the light-house and caused a great loss of life. On the shore near the remains of the old light-house are some very large blocks of stone which, I was informed, were thrown up there from Krakatau, when in eruption. However, on examining them, I found they were masses of ordinary coral reef, which were probably thrown up from the sea by the large wave. They were

quite of the same nature as the Christmas Island rock as far as I could make out, but the organic remains were plainly visible and the rock less compact.

The soil which covers the rock beds in the forest is powdery, and dark brown, strongly impregnated with guano from the droppings of the numerous sea birds nesting upon the trees very similar to that of Fernando de Noronha. The forest is composed of a large variety of trees, many of very large size and producing good timbers, among them was the Gayam (*Inocarpus edulis*), species of *Eugenia* and *Picus*, *Sideroxylon sunduicum*, an *Erythrina* with very small but brilliant orange flowers, and several other kinds of which I was unable to procure enough material to identify. The trees are clothed with ferns of several species, orchids and a wax-plant (*Hoya Aldrichii*, Hemsl.) which is called by the colonists the Flower of Paradise, from its beauty. It is very abundant, but I saw no trace of fruit or flowers, and had to content myself with bringing home some living plants. The orchids were abundant, too, the most common being a new species to which I have given the name of *Saccolabium archytas*. It has thick bright green leaves and long racemes of small white flowers spotted with pink. Besides these there was a *Dendrobium* of the *Cadetia* section, *D. crumenatum*, the well-known pigeon-orchid, a curious new species of *Sarcochilus* and what appeared to be one of the *Phreatias* described from LISTER'S collection by Mr. ROLFE. None of these latter were in flower at the time of my visit, but I secured live plants, and the *Sarcochilus* has since flowered in Singapore.

The pigeon-orchid is one of the most widely distributed of tree orchids in the East Indian region, and I was much surprised to pass a large number of plants apparently of this species quite fresh and green floating in the sea between Singapore and Batavia. One does not imagine that so delicate a plant as an orchid could survive prolonged immersion in the sea.

Terrestrial orchids were represented by *Corymbis veratrifolia*, Bl., also a very widely distributed species occurring in Western Africa, Assam, all over the Malay Peninsula and on the Malay Archipelago. It is true that BLUME in his "Or-

chids of the Indian Archipelago" distinguishes the African, Assamese and Malayan plants as three distinct plants, but the distinguishing characters appear to me too slight, especially as the plant is evidently a very variable one. I recently found specimens in Pulau Ubin near Singapore which were quite indistinguishable from some of the African forms.

Ferns are very abundant. Mr. LISTER collected fifteen species; two of which, viz., an *Asplenium* and an *Acrostichum* were new to science. The ground beneath the bigger trees in the forest produced many shrubs and small plants, among which were *Anisomeles ovata*, a large straggling herb with purple flowers like a large dead nettle; and another horehound-like Labiate with white flowers, a shrubby species of *Achyranthes*, *Ehretia buxifolia*, *Laportea crenulata*, Gaud., a tree nettle with stinging petioles, clumps of *Pandani*, *Oplismenus compositus*, *Fleurya æstuans*, and other plants. A *Randia*, perhaps of an undescribed species, formed coffee-like bushes with small white flowers and orange berries. It is evidently allied to *R. densiflora* which is also stated to occur here, but entirely different from that species in its smaller size of parts and habit. *Ochrosia Ackeringæ*, Miq., is a small *Apocynaceous* tree with white flowers and twin yellow fruits. In more open places near the shore were a tall Mallow (*Abutilon* sp.), with fairly large buff flowers, one of the common tree vines (*Leea sambucina*, L.) resembling an elder bush, and a scrambling bryony with small flowers (*Zehneria mucronata*, Miq.) and on the sandy spots by the shore were the common Waroo (*Hibiscus tiliaceus*), the goat's foot convolvulus (*Ipomea pes-capræ*), *Scævola Koenigii*, L., and *Ischoemum foliosum Hack var. leiophyllum*. This grass was identified for me by Professor HACKEL, who says, that this is a new variety differing from the type only in the glabrous leaves. This type is only known from New Caledonia.

The native palm (*Arenga Listeri*, Becc.) is plentiful, especially on the lower ground near the sea. It is a beautiful species about twenty feet in height with a stout green stem about four inches through, smooth except for the rings. The leaves are large, pinnate, light green above and glaucous beneath. The fruits resemble coffee-berries and are

bright red, each containing three seeds. They are peculiarly irritating even when handled like those of *Caryota urens* and *Kentia Macarthurii*. The tree produces a very excellent sago which is used as food by Mr. Ross' party. Indeed Christmas Island produces, for an island of its size, a remarkably large supply of natural vegetable and animal food, for besides the sago, there are plenty of Gayam (*Inocarpus edulis*) and Indian almonds (*Terminalia catappa*) both eatable and excellent fruits; while land crabs, pigeons and frigate birds supply plenty of animal food.

The birds of the island are somewhat interesting. A pretty brown ground thrush with a red abdomen was very abundant and tame, hopping about everywhere, and small green warblers (*Zosterops natalis*, Sharpe) flew about in little flocks and were very bold and fearless. A small brown hawk like a kestrel (*Urospizias natalis*, Sharpe) appeared on one occasion and fell to the gun. But the attraction to the party who landed from the *Redpole* lay in the pigeons, of which there were two species, one resembling the common green pigeon, the other a large blue dove, known as *Carphophaga Whartoni*, peculiar to this island. This is a very beautiful bird allied to the common Pergam (*C. ænea*), but of a rich deep blue colour. It is very abundant especially on the top of the hill, and towards sunset the woods resounded with its loud note. It sits high up in the lofty trees and is by no means easy to see. It is a very strong bird, several receiving four complete charges of shot before dropping and, like the common Pergam, is tough and not worth eating, but as an ornamental bird it has been successfully introduced into the Cocos Islands by Mr. Ross.

Of the marine birds, the Frigate Bird (*Fregata aquila*), tropic birds (*Phaethon flavirostris*) and boobies (*Sula piscatrix*) were very abundant nesting on the trees and filling the air with their fishy odour. The Frigate Bird is, we found, very good eating, having dark brown flesh with a good and not fishy flavour. Towards evening the great fox bats (*Pteropus natalis*, Thom.), began to appear and clung squeaking to the branches of the trees which were in fruit. They are quite different from the common fox bat, being entirely black. I saw

also a small insectivorous bat flying about, but could not catch it.

The native rats, *Mus Maclearii* and *M. nativitatis*, Thom., do not appear till dark, and as we left before sunset we could not secure any specimens. There are several kinds of lizards here, two of which I caught; one was a very dark, coloured gecko inhabiting the old stumps and rotten trees lying on the ground. As is constantly the case in islands and especially in oceanic islands there are no snakes here, though Mr. ROSS has seen, he tells me, their skeletons washed up on shore. But the most conspicuous denizens of the forest are the large blue and red land crabs (*Birgus latro*) which are exceedingly abundant living in holes in the ground, but constantly wandering about in the woods. Insects are not numerous. I noticed three species of butterfly, a yellow Terias, (*T. amplexa*) a Hypolimnias and what was probably the endemic species *Vadebra Maclearii*, but I could only get a single specimen of the first named species. Mr. ROSS presented me with a pair of very fine maroon-coloured hawkmoths, and told me that there was another and much finer kind to be met with of which he hoped to secure specimens. Of other insects I saw an abundant species of small red wasp, and a fine large *Buprestid beetle*, of a light but brilliant green colour glittering all over, and a large green larval mantis, too young to identify.

I append a list of all the animals and plants recorded, as far as I know, from the island, and hope that Mr. ROSS, who takes much interest in his insular home, will be enabled to assist us in getting a more complete idea of the fauna and flora of Christmas Island.

LIST OF ANIMALS AND PLANTS RECORDED FROM
CHRISTMAS ISLAND.

[NOTE:—In this list all endemic plants and animals are printed in italics. The (!) denotes that specimens were seen or gathered by myself, and the asterisk (*) marks those that have not previously been recorded.]

MAMMALS.

- * *Pteropus natalis*, Thomas. !
Small insectivorous bat. !
- Crocidura fuliginosa* var, trichura, Dobs.
- Mus Maclearii*, Thomas.
- Mus nativitatis*, Thomas.

BIRDS.

- * *Merula erythropleura*, Sharpe. !
Zosterops natalis, Sharpe. !
Collocalia natalis, Sharpe.
- * *Carpophaga Whartoni*, Sharpe. !
Chalcophaps natalis, Sharpe.
- * *Urospizias natalis*, Sharpe. !
Minox natalis, Sharpe.
- * *Ardea jugularis*, Forster. !
Charadrius Geoffroyi, Wagler.
Tringoides hypoleucus, L.
Phæthon phœnicurus, G. M.
Phæthon flavirostris, Brandt. !
- * *Fregata aquila*, L.
Sula piscatrix, L. !

REPTILES.

- Gymnodactylus marmoratus*, Kahl.
- Gecko Listeri*, Blgr.
- Lygosoma nativitatis*, Blgr.
- Ablepharus egerix*, Blgr.
- Typhlops exocæti*, Blgr.
- Chelonia virgata*.

LEPIDOPTERA.

- Valebra Maclearii*, Butl.
Hypolimnias Listeri, Butl.
Nacaduba aluta, Butl.
 * *Terias amplexa*, Butl.
Terias patruelis, Moore.
Porthesia irrorata, Butl.
Hydrillodes, sp.
Endragana limbata, Butl.
Sphingida, sp.
Pyralis Listeri, Butl.
Boarmia compactaria, Walk.

COLEOPTERA.

- Morio orientalis*, Dejean.
Hololepta, sp.
Poederus, sp.
Paroegus Listeri, Gahan.
Leptaulax, sp.
Chrysodema simplex, Waterh.
Stigmatium, sp.
Muephilus, sp.
Hopatrum, sp.
Sessinia, 2 sp.
Cercesium nigrum, Gahan.

HEMIPTERA.

- Lygocus subrufescens*, Kirby.

HOMOPTERA.

- Oxypleura calipso*, Kirby.
Ricania flavicostalis, Kirby.
R. affinis, Kirby.
R. hyalina, Kirby.

DIPTERA.

- Laphria nigrocoerulea*, Kirby.
Stilbomyia jucunda, Kirby.

MYRIAPODA.

- Cryptops hortensis*, Leach.
Cryptops inermipes, Pocock.
Mecistocephalus castaneiceps, Haase.
Cylindrodesmus hirsutus, Pocock.
Spirostreptus exocoeti, Pocock.

ORTHOPTERA.

- Labidura nigricornis*, Kirby.
Blatta livida, Fabr.
Panesthia javanica, Serv.
Clitumnus stilpnoides.
 A large larval Mantis.
Phisis Listeri, Kirby.
Gryllacris rufocaria, Kirby.
Primnia orientalis, Kirby.
Cyrtacanthacris fusilinea, Walk.
C. disparilis, Kirby.
Epacromia rufostriata, Kirby.

HYMENOPTERA.

- Camponotus melichlorus*, Kirby, (Formicidæ).
Lobopelta diminuta, Kirby, (Poneridæ).
Odynerus polypemus, Kirby, (Eumenidæ).
Polistes belder, Kirby, (Vespidæ).

ARACHNIDA.

- Nephila nigritarsis*, Koch.
Homalattus auratus, Koch.
Heteropoda venataria, L.

CRUSTACEA.

- Hyllocarcinus natalis*, Pocock.
Birgus latro, L.
Monchammus naticciatis, Gahan.
Praonethra perplexa, Gahan.
Micracantha, sp.

CRUSTACEA,—*Continued.*

Epilachum, sp.
Piezonotus discoidalis, Waterh.

MOLLUSCA.

Ariophanta Normani, Smith.
A. Mabelæ, Smith.
A. Mildredæ, Smith.
Succinea solidula, Pfeiffer.
S. solitaria, Smith.
S. Listeri, Smith.
Pythia scaraboeus, L.
Melampús luteus, Quoy.
Melampus fasciatus, Deshayes.
Truncatella valida, Pfeiffer.
Leptopoma mouhoti, Pfeiffer.

PLANTS.

DICOTYLEDONES.

- Abutilon indicum* var. ?
 * *A.* sp. !
 * *Hibiscus abelmoschus*, L. ! Near the Settlement.
 * *H. tiliaceus*. ! Common on the shores.
Vitis pedata. !
Leea horrida, Teysm.
 * *L. sambucina*, L. ! Near the shore.
Erythrina, sp. ! Tall tree. Flowers small, scarlet orange.
Inocarpus edulis, Forst.
Terminalia catappa, L.
Eugenia, sp.
Barringtonia racemosa, Bl.
Pemphis acidula, Forst.
Zehneria mucronata, Miq. Near the shores.
Heptapleurum ellipticum, Seem. !
Randia densiflora, Benth.
 * *R.* sp. ! Bushes common near the Settlement.
 * *Psychotria*, sp. ! On the top of the ridge, small dark green bushes.
Blumea spectabilis, Dec.

PLANTS,— *Continued.*

- * *Spilanthes acmella*, L. ! Near the huts.
- Scævola Koenigii*, !
- Ardisia complanata*, Wall.
- Sideroxylon sundaicum*, Miq. ! A lofty tree near shore.
- Ochrosia Ackeri*gæ, Miq. ! Small tree near shore.
- Hoya Aldrichii*, Hems! Covering all the trees luxuriantly.
- Cordia subcordata*, Lam.
- Ehretia buxifolia*, Roxb.
- Tournefortia argentea*, Linn.
- * *Ipomea pescaprae*, ! Linn. Sea shore.
- Solanum biflorum*, ! Lour. !
- Datura alba*.
- Dicliptera Maclearii*, Hems.
- Anisomeles ovata*, R. B. L. In the woods, common.
- Callicarpa longifolia*, Lam.
- Tectona grandis*, Linn.
- Boerhaavia repanda*, Willd.
- Pisonia excelsa*, Bl.
- Achyranthes aspera*, L.
- * *A. sp.* ! A shrubby species near the shore in the woods.
- Deeringia celosioides*, R. Br.
- Peperomia*, sp.
- Hernandia ovigera*, L.
- Euphorbia hypericifolia*, L.
- Cleidion javanicum*, Bl.
- Macaranga tanarius*, Muell. Arg.
- Cudrania javanica*, Trec.
- Laportea crenulata*, Gaud. !
- Fleurya ruderalis*, Gaud. !

MONOCOTYLEDONES.

- * *Dendrobium crumenatum*, Lind. !
- * *D. (§ cadetia)* sp.
- Phreatia Listeri*, Rolfe.
- Ph. congesta*, Rolfe.
- * *Sarcochilus carinatifolius* sp. One plant only found.
- * *Saccolabium Archytas* sp. ! Very abundant.

MONOCOTYLEDONES,—*Continued.*

- * *Corymbis veratrifolia*, Bl. ! Top of the ridge.
Arenga Listeri, Becc. !
Pandanus, sp. Bushes near shore.
Fimbristylis cymosa, R. Br.
- * *Digitaria sanguinalis*, L. Near the Settlements.
- * *Oplismenus compositus*, Beauv. ! In the woods.
- * *Ischæmum muticum*, L.
- * *I. foliosum* var. *leiophyllum*, Hack. ! On the shore,
forming big tufts.
- * *Eleusine indica*, L. ! Near the Settlement.
Eragrostis plumosa, Lk.

FERNS.

Davallia solida, Sw.
D. dissecta, J. Sm.
Asplenium Nidus, L. !
A. falcatum, Lam. !
A. centrifugale, Bak.
Nephrodium truncatum, Presl.
N. syrmatium, Bak.
Aspidium membranaceum, Hook.
Nephrolepis acuta, Presl.
N. ramosa, Moore.
Polypodium adnascens, Sw.
P. irioides, Lam.
Vittaria elongata, Sw. !
Acrostichum flagelliferum, Wall.
A. Listeri, Bak.

LYCOPODIACEÆ.

Lycopodium phlegmaria, L.

MOSESSES.

Neckera Lepineana, Mont.
Thyridium fasciculatum, Nutt.

HENATICÆ.

Ptychanthus squarrosus, Mont.

Lejeunia serpyllifolia, Lib.

LICHENS.

Usnea trichodea, Ach.

FUNGI.

Polyporus australis, Fr.

P. conchatus, Fr.

Stereum lobatum, Kze.

NEW SPECIES DESCRIBED.

Sarcophilus carinatifolius.

Stem six inches long, flattened. Leaves fleshy oblong-elliptic slightly unequally bilobed blunt at the lip, bright green very strongly keeled especially at the base, two inches long one inch across; the base flattened laterally so as to form a broad flat petiole; sheath very deeply cleft on side opposite to lamina, when dry strongly ribbed. Scapes very graceful erect slender pale green about six inches long with one or two small sheathing bracts. Raceme of few flowers, opening one or two at a time, rachis thickened terete; bracts lanceate acute; green $\frac{1}{8}$ of an inch in length; flowers small thin textured, white withering buff; ovary and pedicel $\frac{1}{2}$ inch long, straight bright green. Posticous sepal lanceolate acuminate with recurved apex, laterals similar but the base prolonged below into an angle, keeled outside. Petals shorter linear lanceolate. Lip white base prolonged to form a blunt spur, lateral lobes very long narrow linear acuminate, midlobe shorter thick ovate blunt; a lorate raised ridge on the disc ending in a blunt rounded callus on the midlobe; a yellow spot on the base of the lip outside. Column straight thick with a distinct foot, clinandrium nearly flat, wings of column bent over the face,

foot linear upcurved. Another quadrate rounded at the top front edge truncate straight. Pollinia four unequal oval flattened with a small oval disc. Stigma deep oval; rostellum small.

A curious little plant, which bears no very distinct relation to any other species I know, but is probably an ally of *S. leopardinus*, Par., or at all events belongs to the section which includes this plant.

Saccolabium Archytas, nsp.

Stems short usually crowded in large masses with very many roots. Leaves lorate few, bright green, apices unequally bilobed, with blunt lobes, six inches long, by one broad. Racemes pendulous covered with small flowers, four to six inches long with a rather slender angled green rachis. Bracts short ovate acute. Flowers hardly $\frac{1}{2}$ inch across, opening a few at a time, ovary and pedicel $\frac{1}{2}$ inch long white, terete. Dorsal sepal narrow oblong obtuse boat-shaped, laterals obtusely subtriangular. Petals spatulate with a broad obtuse apex, shorter than the sepals. All white. Lip small with side-lobes large and rounded white with violet spots, midlobe absent. Spur large straight dependent blunt. Column base narrow, above dilated, white with a violet face. Anther ovate acute in front, yellow. Pollen masses two globose pale yellow, pedicel linear rather broad, disc ovate. Capsule slender elongate three inches long, ribs not much elevated.

This is a pretty little plant although the flowers are so small. It is remarkable for the subtriangular sepals and the lip being reduced almost to a spur with the lateral lobes flanking the mouth.

SUMMARY.

The fauna and flora is typically Malayan as might be expected from the position of the island, and is evidently primarily derived from the coasts of Java which is the nearest land. Perhaps when the southern regions of Java are better worked out we shall find some of the peculiar Christmas Island species occurring there.

The greater part of the plants fall under one of the two heads, those whose seeds or fruits are carried about unharmed by sea currents, and those which having edible fruits are dispersed by birds. These in fact are the two methods by which most of the oceanic islands are populated. To the first class belong *Hibiscus tiliaceus*, *Inocarpus edulis*, *Terminalia Cattappa*, *Barringtonia racemosa*, *Pemphis acidula*, *Ochrosia*, *Ackeringa*, *Cordia subcordata*, *Tournefortia argentea*, *Ipomea pes capræ*, *Boerhaavia repanda*, *Hernandia ovigera*, *Pandanus*, sp., *Fimbristylis cymosa*, and probably also *Erythrina*, *Abutilon*, *Ehretia buxifolia* and *Tectona*. *Dendrobium crumenatum* and the other orchids may have been derived from plants drifted ashore on fallen trees.

As fruit-eating birds and bats are very numerous it is not to be wondered at that there are many species with edible fruits, which have been brought to the island by them. To this class belong *Vitis*, *Leea*, *Zehneria*, *Heptapleurum*, *Randia*, *Psychotria*, *Ardisia*, *Solanum*, *Collicarpa*, *Deeringia*, *Ficus*, *Didymosperma* and probably also *Eugenia*, *Sideroxylon*, *Euphorbia*, and *Cleidion*. The fruits of *Didymosperma* are as stated above very irritating but this is no bar to their being devoured by birds. The fruits of *Kentia Macarthuri*, a New Guinea palm, are nearly as irritating, but they are greedily eaten by the blue starlings in the Botanic Gardens, at Singapore and I found, in Fernando de Noronha, a species of *Sapium* which was so poisonous that the fruit falling on a horse would blister the skin, yet it was often eaten by small birds.

A few species have been introduced accidentally by man into Christmas Island, and this class will doubtless increase largely. Nearly all of these occur close to the Settlements. They include *Hibiscus Abelmoschus*, *Spilanthes*, *Datura*, *Achyranthes aspera*, *Eleusine*, *Digitaria* and perhaps *Abutilon indicum*. Such plants as *Pisonia excelsa* with its sticky fruits, *Achyranthes* sp. and *Oplismenus* may have been borne to the island by birds, as their fruits are more or less adhesive.

It might be imagined that winged or plumed seeds would be easily carried to distant islands and consequently form an important element in the flora. This is not so. In the majo-

urity of cases these seeds are carried a short way only and even if they were widely scattered by prolonged and violent winds, they would have but little chance of accidentally striking an island far off in the sea. To the class with plumed seeds belong *Hoya Aldrichii*, and *Blumea spectabilis*. Both may have been introduced by wind-currents.

Orchids are acknowledged to be rare in Oceanic Islands although their minute seeds are blown from the split capsules for a considerable distance. Perhaps that the Christmas Island species were in this manner drifted to their present home, but it is also possible that the epiphytic species were floated there on logs of fallen trees.

Ferns, the spores of which are produced in enormous quantities and are exceedingly light, are certainly widely scattered by the blowing of their spores to long distances, and the same remark applies to Fungi, Mosses and other Cellular Cryptogams.

————:0:————

LIST OF PAPERS PUBLISHED ON CHRISTMAS ISLAND.

Proc. Zool. Soc. 1887.

MACLEAN, Capt., General Report, on the Collections of H. M. S. <i>Flying Fish</i> , ...	p.	508
GUNTHER, A., Zoology,	"	507
THOMAS, O., Report on Mammals,	"	511
SHARPE, R. B., Report on Birds,	"	515
BOULENGER, G. A., Report on Reptiles,	"	516
SMITH, E. A., Report on Molluscs,	"	517
WATERHOUSE, C. O., Report on Coleoptera,	"	520
POCOCK, R. J., Report on Crustacea,	"	520
BELL, F. J., Report on Echinoderms,	"	523
DENDY, A., Report on Porifera,	"	524

Proc. Zool. Soc., 1888.

LISTER, J. J., Natural History,	p.	512
THOMAS, O., Mammals,	"	532
BOULENGER, Reptiles,	"	534
GAHAN, C. S., Coleoptera,	"	538
BUTLER, A. G., Lepidoptera,	"	542
SMITH, E. A., Terrestrial Molluscs,	"	536
KIRBY, W. F., Various Insects,	"	546
POCOCK, R. J., Arachnids, Myriapods, and Crustacea,	"	556
HEMSLEY, Report on Botanic Collections, Journ. Linn. Soc., 1890, p. 351.		
MACLEAN, J. P., Note on Christmas Island. <i>Nature</i> , xxxvi. p. 13.		
LISTER, J. J., <i>Nature</i> , xxxvii, p. 203.		
WHARTON, J. L., <i>Proc. Geogr. Soc.</i> , 1888, p. 613-624.		

OCCASIONAL NOTES.

DISCOVERY OF A STONE IMPLEMENT IN SINGAPORE.

A short time ago, Lieutenant A. D. COX, while walking on a road at the barracks at Tanglin, picked up from among the laterite which was being put on the road a large stone implement, which he has presented to the Museum. The weapon is five inches in length and nearly four inches across and about one and a half thick in the thickest part. It is oblong, with one end abruptly truncate, the other ground off to a rounded cutting edge. It is a good deal worn, and at one spot bears a small coral, showing that it had recently come from the sea. It is of a dark chocolate-brown externally, but by dipping it a little at one corner it was found to be composed of a very hard compact granite containing very small flakes of mica. On enquiry I found that the contractor who was laying down the laterite had obtained it from Tanjong Karang on the West Coast of Singapore. This spot I have since visited, and found that the stone was being taken from below high water mark, which would account for the presence of the coral upon the specimen. Tanjong Karang is a small promontory, consisting of a core of rather hard iron-stone, covered with about two feet of humus and gravelly soil. There is what is called a *Kramat* at the corner nearest to the spot whence the specimen must have come, but this *Kramat* merely consists of a detached block of iron-stone, which in shape more or less resembles a tomb. The overlying soil on the promontory has so shifted from denudation that it is impossible to get any idea of its age; fragments of modern pottery occurring even at the part where it rests on the iron-stone. I sought carefully for any more weapons, but could find none, and indeed it was hardly to be expected, as they are almost always found singly here.

Hitherto, as far as I am aware, no worked stone of this class has ever been found in Singapore, though stone implements have been obtained in some numbers in Perak and Pahang, where they are known as *Batu Linta*. These latter are generally made of a black igneous rock, and of a narrower form than the present one. The form, however, of a stone axe must always depend on the texture and hardness of the stone and its cleavage. And the exceedingly hard texture of this granite is, no doubt, the reason for its broad shape. The edge has evidently been ground very carefully, but not equally, one side being flatter than the other. From this I gather that the weapon was used as an adze or *billong*.

Of what race were the makers of these weapons we are entirely ignorant, none of the wild tribes use stone weapons, nor do they know anything about those that are found in the Peninsula. Nor is the stone of which the Pahang specimens are made, known at all from this region. It is probable that these were brought from farther East, but the Singapore specimen is made of granite, and of such granite as is found at no great distance from Tanjong Karang, namely, at Pulau Ubin, and I also found an outcrop of granite in one of the hills between Tanjong Karang and Toas. It is probable, therefore, that it was made on or near the spot where it was found, possibly from a sea-shore pebble of granite from the neighbourhood of Pulau Ubin.

H. N. R.

PANTANG LARANG OF NEGRI SEMBILAN.

The Yam Tuan Besar of Negri Sembilan has recently caused to be published certain rules for the conduct of the Malays of that State. These rules—*Pantang Larang*—are not new, but are some of the old rules of the State. They are now published to show the masses that the Malay customs are in full force as hitherto. The following is a translation :—

It is forbidden—

1. To fire guns or cannons during the three days previous to either of the great feasts called *Hari Raia* or *Hari Raia Haji*, *i. e.*, on the 28th, 29th or 30th of Jal Haija.

2. To disobey a lawful order of the Raja or Penghulu.

3. For any one to wear clothes of the colours which are set apart as the special colour attached to the Rajaship, or to any State office, such as yellow, which is the Raja's colour: or to carry the hair-ornamented spear, or to wear entirely black clothing, which may only be the dress of officers such as the *Laksamana*.

4. For any person to make use of and display weapons or articles similar to the insignia of a Chief or State officer, as such insignia.

5. To recite the form of prayer called *Hothbah*, except at the Istana, Penghulu's or Lembaga's Court-house, Mosque, or at any other place specially set apart by the Raja or Penghulu.

6. For any person to wear weapons having gold or silver ornaments covering either the end of the handle or of the sheath, except by permission or rank.

7. To arrange his house or premises similarly to a royal hall, called *Balei jariga*, which is fenced round with split *runyong* (the Kabong tree) according to the ancient custom in the country of Menangkabau, with gates roofed (*i. e.*, with an arch over them), with the exception of persons who are permitted by the Raja or Penghulu.

8. The big drum of the Mosque is not to be beaten, except on feast days or on the occasion of any public calamity, or at a calling together of the people.

Dahga Dahgi. This means that the people of low rank must never disobey the orders of those of higher rank than themselves.

Sumbang Salah. This means that the children of two women who are sisters cannot marry each other. It is against the law of the country.

Chelaka Derahka means that whosoever disobeys these orders shall be considered guilty of treason.

Whoever disobeys the aforesaid rules is liable to a fine not exceeding twenty-five dollars.

M. L.

ON THE OCCURRENCE OF A RHODODENDRON IN SINGAPORE.

In January of this year, while walking near the bungalow on the top of Bukit Timah with Lieutenant KELSALL, I observed in the upper branches of a lofty Dipterocarpous tree, probably a species of *Shorea*, an epiphytic plant with red flowers. On going to the foot of the tree we picked up several fallen blossoms, from which it was evident that the plant was a species of *Rhododendron*, and that, one not known to occur in the Malay Peninsula. The tree was about 150 feet high, and about 6 feet through, without a branch for fully 80 feet, so that it was by no means easy to reach it. The following plan was then adopted. A number of large spike nails were driven into the tree, and a pole lashed to them so as to form a ladder, for about a quarter of a distance. A Malay then ascended this with a further supply of nails, and a pulley block being fixed to the tree another pole was hauled up and made fast to the lower one. This was done again and again till the lowest branch was reached, and eventually the plant was lowered to the ground. There were two plants on the tree, which were transferred to the Botanic Gardens, where one remained a long time in flower. The plant is an exceedingly beautiful one, with clusters of nodding flowers about two inches long, of a beautiful crimson-red colour. It appears to be identical with one figured in the Botanical Magazine under the name of *Rhododendron javanicum* var. *tubiflorum* (Tab. 9850), differing only in the colour, which in the picture is a very dull red. The plant from which this drawing was made was ob-

tained by Mr. CURTIS in Solok in Sumatra, and sent home to Mr. VEITCH, who cultivated it in England. It seems, to me however, that this so-called variety is specifically quite distinct from *Rhododendron javanicum*, which is also a native of the Peninsula.

Rhododendron javanicum, Benn., is based on a plant found in Java by Dr. HORSFIELD, and is figured and described in BENNETT & HORSFIELD'S *Plantæ Javanicæ Rariores* (Tab. xix), and I have what is evidently the same plant from Gunong Hijau in Perak, collected by Mr. CURTIS, but the tube of the flowers is rather shorter and more infundibuliform. The Singapore plant differs from the true *javanicum* in the leaves, which are shorter, thicker and blunter and more conspicuously dotted with glands, and the base of the blade is not narrowed into the petiole, but ends abruptly. The flowers are nodding on rather long pedicels. The tube is very long in proportion to the limb, nearly $1\frac{1}{2}$ inch, the lobes of the limb are short, broad and rounded, about half an inch long. The tube is cylindrical, slightly curved, and when alive grooved at the base, where, as in the Perak plant, it is almost funnel shaped, with very large lobes to the limb. The stamens again are much thinner, the anther only half the size. The style is thick and the stigma large and somewhat trilobed in the Perak plant and in BENNETT'S figure, whereas in our plant the style is thin, with a much smaller more simple stigma. These points are so marked, that it appears to me that the two plants should be specifically separated. It is very possible that the Singapore plant may be identical with *Rh. longiflorum*, Lindl., a plant which has been found in Borneo at sea level by Professor BECCARI, whose description fits it fairly well, but I have not here LINDLEY'S original description of that plant.

Besides this species, there are four other kinds of *Rhododendron* in the Malay Peninsula.

Rh. malayanum, Jack, which is the commonest, occurs on Mount Ophir, Gunong Hijau, and Maxwell's Hill, and in other mountains of the central range, and also in Sumatra, Java and Celebes. It is a straggling plant with small crimson flowers.

Rh. jasminiflorum, Hook., with clusters of white jessamine-like flowers, only known from Mount Ophir.

Rh. Teysmanni, Miq., a yellow flowered plant, grows on Penang Hill. It is an ally of *Rh. javanicum*, Benn., which is the fourth species.

All these grow on the trees as epiphytes, usually at considerable altitudes, *i. e.*, from 2,000 feet upwards.

The only other plant of the order *Ericaceæ* yet recorded from the Peninsula is a *Diplycosia microphylla*, Becc., a remarkable little creeping plant, which appears to be common at high altitudes. It has more of the appearance of one of the bilberry family, with tough, wiry branches, and small oval leaves.

The allied orders of *Vacciniaceæ* and *Epacridæ* are represented respectively, in Singapore, by *Vaccinium malaccense* a bush of considerable size, with pink or white sweet-scented heather bell-shaped flowers, and small eatable black bilberries; and the remarkable *Leucopogon malayanus*, Jack, a large bush with hard, sharp-pointed leaves, very small white flowers, and sweet but very small orange berries. Both grow on sandy ground near Changi towards Tanah Merah, *Leucopogon* forming here dense thickets. Another species of *Vaccinium* also grows in Pekan, with longer and more crimson flowers.

H. N. R.

POGONIA PUNCTATA, BL., IN SINGAPORE.

The genus of orchids *Pogonia* has not hitherto been recorded from the Malay Peninsula, but as the surrounding countries produce a number of species, it is not to be wondered at that at least one kind should occur here. Some months ago, I found in a stream on the western slope of Bukit Timah a small plant of some species of this genus, and by carefully transferring the small tuber with its solitary leaf to the Botanic Gardens, I was able to induce it at last to flower, when it proved to be *P. punctata*, Bl.—a plant obtained by BLUME from the slopes of Mount Pangerang in Java, and figured by

him in the "Orchids of the Malay Archipelago" (Tab. 42).

This plant belongs to the section of the genus in which the umbrella-shaped leaf appears alone, and, after remaining for a short time, suddenly withers and is followed by a stem bearing one or more flowers. In this species the leaf is about three inches tall, the blade being orbicular, cordate dark green and more or less purple beneath. The flower stem is about the same height, covered with sheaths, which are thickly dotted with purple, whence the trivial name. It bears a single nodding flower about an inch and a half long, with a purple ovary and a pale glower thickly spotted all over outside with purple. The petals and sepals are very narrow linear acute. The lip is white with violet spots. The flower has a faint scent of cucumbers. It never seems fully to open. Probably it is not so rare as it would appear, for, without doubt, it is an exceedingly difficult plant to find, as it is very inconspicuous and occurs apparently quite sporadically. I have several times since hunted in the spot where I found this little plant, and never seen any more.

H. N. R.

THE KERINGGA.

Since writing the account of the *Caringa* (more correctly *Keringga*) in the last number of the Journal, I have received a letter from Mr. WATERHOUSE, of the British Museum, in which he tells me that the specimens I sent him are *Æcophylla smaragdina* and not *Formica gracilipes*. It apparently takes its name of *smaragdina* (emerald green) from the curious dusk-green colour of the winged female. It is identical with the insect of Ceylon and Southern India.

H. N. R.

EUDROMIAS VEREDUS IN SINGAPORE.

ON the 13th of May, 1891, the Museum Taxidermist, Mr. L. A. FERNANDEZ, saw a Malay boy capturing birds by

means of hair nooses on the race course. On inspecting his captures, he found a pair of plovers ♂ and ♀ with which he was not acquainted; these he purchased for the very moderate sum of 10 cents for the pair, and they are now mounted. They prove to be *Eudromias veredus* of Gould., originally described P. Z. S., 1848, p. 38. The species is rare in this part of the world. It has been recorded once from the Andamans (S. F. I., 83) and once from the Sunderbunds. It however seems to occur not unfrequently in Java and the islands farther East.

W. D.

I may also note that when at Penang in May last I noticed a flight of about 20 birds of the rare tern *Sterna leucoptera*.

W. D.

FRUGIVOROUS HABITS OF THE TUPAIA.

This little insectivore (*Tupaia javanensis*) is very common in Singapore, and especially in the Botanic Gardens, where it may be often seen running about among the trees. It is easily mistaken for the common little squirrel (*Sciurus hippurus*), of which it has much the appearance. When alarmed it quickly darts up the trunk of the nearest tree, but is a poor climber, and never seems to go high up like the squirrel. But besides these points of resemblance, it appears also to be largely frugivorous. It was found that the seeds sown in boxes were constantly being dug up and devoured by some animal, and traps baited with pieces of coco-nut or banana were set, and a number of tupaia were caught. These being put into a cage appear to live very comfortably upon bananas, pineapple, rice and other such things; refusing meat. The Rev. T. G. WOOD, in his Natural History, states that *T. ferruginea* is said to feed on beetles, but to vary its diet with certain fruits. The common species here seems to be almost entirely frugivorous, though its teeth are those of a typical insectivore.

H. N. R.

[No. 24.]

JOURNAL
OF THE
STRAITS BRANCH
OF THE
ROYAL ASIATIC SOCIETY.

DECEMBER, 1891.

SINGAPORE:

PRINTED AT THE GOVERNMENT PRINTING OFFICE.

AGENTS OF THE SOCIETY:

London and America, ... TRÜBNER & Co.

Paris, ... ERNEST LEROUX & CIE.

Germany, ... K. F. KOEHLER'S ANTIQUARIUM. Leipzig.

169

TABLE OF CONTENTS.

Council for 1892,	v
List of Members for 1892,	vi
Proceedings of the General Meeting,	xv
Council's Annual Report for 1891,	xvii
Treasurer's Account for 1891,	xix

An Old Minute by Sir Stamford Raffles— <i>by H. A. O'Brien,</i>	1
Some Notes on the Sakai Dialects of the Malay Peninsula— <i>by Hugh Clifford,</i>	13
An Account of a Journey across the Malay Peninsula from Koh Lak to Mergui— <i>by Arthur Keith, M.B., C.M.,</i> ...	81
The Thermal Springs of Selangor and Malacca— <i>by W.</i> <i>Bott, F.C.S., F.G.C.S., &c.,</i>	43
Notes on the Siamese Provinces of Koowi, Bangtaphan, Pateeo and Champoon— <i>by Arthur Keith, M.B., C.M.,</i> ...	63
The Alleged Discovery of Mercury in Malacca— <i>by W.</i> <i>Bott, F.C.S., F.G.C.S., &c.,</i>	79
A Note on Rengas Poisoning— <i>by W. C. Brown, M.D.,</i> ...	83
A New Collection of Malay Proverbs— <i>by Hugh Clifford,</i> ...	87

TABLE OF CONTENTS,—*Continued.*

A Bibliography of Malaya from July, 1890, to June, 1891,— <i>by C. Davies Sherborn, F.G.S., F.Z.S.,</i>	121
Occasional Notes :—	
The Putri of Mount Ophir,	165
Diamonds in the Malay Peninsula,	166
Description of a new Species of Jungle Fowl, said to come from Borneo,	167
A large Mias in Singapore,	168
Note on the Nest and Eggs of <i>Nyctiornis Amicta</i> ,	169
Nest and Eggs of <i>Henicurus Ruficapillus</i> , Temm.,	170
On the Occurrence of <i>Petrosavia</i> in Perak.	170

THE
STRAITS BRANCH
OF THE
ROYAL ASIATIC SOCIETY.

PATRON:

His Excellency Sir CECIL CLEMENTI SMITH, G.C.M.G.

COUNCIL FOR 1892.

His Excellency Major-General Sir CHARLES WARREN, G.C.M.G.,
K.C.B., *President.*

The Right Revd. Bishop G. F. HOSE, *Vice-President, Singapore.*

D. LOGAN, Esquire, *Vice-President, Penang.*

H. N. RIDLEY, Esquire, *Honorary Secretary.*

H. T. HAUGHTON Esquire, *Honorary Treasurer.*

A. KNIGHT, Esquire,

H. L. NOBONHA, Esquire,

C. W. S. KYNERSLEY, Esquire. } *Councillors.*

Lieut. H. J. KELSALL, R.A.,

The Hon'ble J. W. BONSER.

LIST OF MEMBERS

FOR

1892.

Nos.	Names.	Addresses.
1	ABRAHAMSON, E. E.	China North Borneo Co. Ltd., Sandakan, B. N. B.
2	ALLINGHAM, S.	25, Grosvenor Street, Grosvenor Square, Lon- don, W.
3	ANTHONISZ, J. O.	Singapore (Europe).
4	AYRE, A. F.	Singapore (Europe).
5	BAMPFYLDE, C. A.	Kuching, Sarawak.
6	BECHER, H. M.	Singapore.
7	BEESTON, Capt. R. D., B.A.	Sandakan, B. N. B.
8	BELLAMY, G. C., B.A.	Selangor.
9	BERNARD, F. G.	Woodleigh, Saranggong Road, Singapore.
10	BICKNELL, W. A.	Audit Department, Penang.
11	BIRCH, J. K.	First Magistrate, Penang.
12	BLAND, R. N.	Land Department, Singa- pore.
13	BLAGDEN, C. O.	Malacca.
14	BONSER, The Hon'ble J. W.	Singapore.
15	BOTT, Dr. W.	The Grange, Grange Road, Singapore.
16	BRADDON, Dr. W. L.	Jelebu.
17	BRANDT, D.	Singapore.

MEMBERS FOR 1892,—*Continued.*

Nos.	Names.	Addresses.
18	BROWN, L. C.	Penang.
19	BROWN, Dr. W. C.	Beach Street, Penang.
20	BRYANT, A. T.	The Dindings.
21	BUCKLEY, C. B.	Orchard Road, Singapore.
22	BURBIDGE, W.	Sophia Road, Singapore.
23	BURKINSHAW, J.	Mount Elizabeth, Orchard Road, Singapore (England).
24	CAMERON, Capt. M. A., R.E.	Dover, England.
25	CAMUS, M. DE	Singapore.
26	CERRUTI, G. B.
27	CLAINE, J.	21, Northam Road, Penang.
28	CLIFFORD, H.	Kwala Lipis, Ulu Pahang.
29	COPLEY, GEORGE	Municipality, Malacca.
30	CREAGH, His Excellency C. VANDELEUR, C.M.G.	British North Borneo.
31	CROIX, J. ERRINGTON DE LA	France.
32	CURRIE, ANDREW	28, Fenchurch Street, London.
33	DANE Dr. R.	Singapore.
34	DENT, Sir ALFRED, K.C.M.G.	11, Old Broad Street, London, E. C.
35	DEW, A. T.	Matang, Perak.
36	DIETHELM, W. H.	Hooglandt & Co., Singapore.
37	DOWN, St. V. B.	Singapore (England).
38	DUNLOP, C.	Powell & Co., Singapore.
39	EBHARDT, HANS	Hilty & Co., Singapore.
40	EGERTON, WALTER	Butterworth, Province Wellesley.
41	ELCUM, J. B.	Penang (Europe).

MEMBERS FOR 1892,—*Continued.*

Nos.	Names.	Addresses.
42	ESCHKE, H. H.	German Consulate, Singapore.
43	EVERETT, H. H.	Labuan.
44	EVERETT, A. HART	Labuan.
45	FERGUSON, A. M., Jr.	18 & 19, Baillie Street, Fort, Colombo.
46	GAGGINO, G.	Gaggino & Co., Singapore.
47	GAYNOR, Lieut. H. F., R.E.	Tanglin Barracks, Singapore.
48	GENTLE, ALEX.	Grange Road, Singapore.
49	GOLDNEY, Mr. Justice J. T.	Penang.
50	GOSLING, T. L.	River Valley Road, Singapore.
51	GOTTLIEB, G. S. H.	Penang.
52	GRAHAM, JAMES	London.
53	GULLAND, W. G.	Paterson, Simons & Co., London.
54	HALE, A.	Negri Sembilan.
55	HAUGHTON, H. T., B.A.	Singapore.
56	HAVILAND, DF.	Kuching, Sarawak.
57	HERVEY, The Hon'ble D. F. A., C.M.G.	Resident Councillor, Malacca.
58	HILL, E. C. H.	Inspector of Schools, Singapore.
59	HILL, FENTON WALTER	Kuala Lumpur, Selangor.
60	HOSE, Right Revd. Bishop G. F., M.A., D.D. (Honorary Member)	Sarawak.
61	HOSE, C.	Baram, Sarawak.
62	HOUTHUYSEN, C. L.	Netherlands Trading Society, Singapore.
63	HOYNCK, P. C. VAN PAPENDRECHT	Singapore.

MEMBERS FOR 1892,—*Continued.*

Nos.	Names.	Addresses.
64	HULLETT, R. W., M.A., F.L.S.,	Singapore.
65	HUTCHINSON, E. R.
66	IBRAHIM BIN ABDULLAH, Dato Dalam	Johor Baharu.
67	IRVING, C. J., C.M.G.	Hillands Tiverton, Devon- shire, England.
68	JOAQUIM, J. P.	Buitenzorg, Paterson Road, Singapore.
69	JOHOR, H. H. the Sultan of the State and Territory of, G.C.M.G., G.C.S.I. (Honorary Member)	Johor.
70	KEHDING, F.	Labuan, Deli.
71	KEITH, Dr. A.	Bangkok, Siam.
72	KELSALL, Lieut. H. J., R.A.	Fort Canning, Singapore.
73	KENNEDY, ARCHIBALD	Batu Gajah, Kinta, Perak.
74	KER, T. RAWSON	Johor Baharu, Johor.
75	KNIGHT, ARTHUR	Grassdale, River Valley Road, Singapore.
76	KROM MUN DEWAWONGSE VARO- PRAKAR, H. R. II. Prince	Bangkok.
77	KYNERSLEY, C. W. S.	Singapore.
78	LAUGHER, H.	Singapore.
79	LAWES, Revd. W. G. (Honorary Member).	Port Moresby, New Guinea.
80	LAVINO, G.	Dutch Consulate, Singa- pore.
81	LEASK, Dr. J. T.	Malacca.
82	LEES, F. BALFOUR	Singapore Insurance Com- pany Limited, Singapore.

MEMBERS FOR 1892,—*Continued.*

Nos.	Names.	Addresses.
83	LEWIS, JOHN E. A., B.A.	Government Printing Office, Sarawak.
84	LISTER, The Hon. MARTIN	Negri Sembilan.
85	LITTLE, R. M.	Gaya, Sandakan.
86	LOGAN, D.	Solicitor-General, Penang.
87	LOW, Sir HUGH, G.C.M.G.	London.
88	MACBEAN, W.	Straits Insurance Office, Hongkong.
89	McKILLOP, J.	Pulau Brani, Singapore.
90	MACHADO, A. D.	Temoh Gold Mines, Ke- lantan.
91	MARTIN, Dr. L.	Mabar Estate, Deli.
92	MAXWELL, The Hon'ble W. E., C.M.G.	Colonial Secretary's House.
93	MAXWELL, R. W.	Inspector-General's Office, Singapore.
94	MEREWETHER, E. M.	Singapore.
95	MILLER, JAMES	Gilfillan, Wood & Co., Singapore.
96	NANSON, WM., B.A., F.S.A.	Craigton, Tanglin, Singa- pore.
97	NAPIER, W. J., M.A.	Paterson Road, Singapore.
98	NEAVE, D. C.	Cluny Road, Tanglin, Singapore.
99	NEWTON, HOWARD	Grasslands, St. Thomas Walk, Singapore.
100	NORONHA, H. L.	Devonshire Road, Singa- pore.
101	O'BRIEN, The Hon'ble H. A.	Singapore.
102	O'SULLIVAN, A. W. S., B.A.	Penang.
103	OWEN, J. F.	Kwala Pahang.

MEMBERS FOR 1892,—*Continued.*

Nos.	Names.	Addresses.
104	PATERSON, D. W.	Guthrie & Co., Singapore.
105	PAUL, W. F. B.	Sungei Ujong.
106	PERAK, GOVERNMENT MUSEUM.	Larut.
107	PERHAM, The Ven. Archdeacon J. (Honorary Member)	Singapore.
108	PICKERING, W. A., C.M.G.	England.
109	READ, W. H. M., C.M.G.	9, Petersham Terrace, Queen's Gate, London.
110	REITH, Revd. G. M., M.A.	Mount Elizabeth, Singa- pore.
111	RICKETT, C. B.	Hongkong and Shanghai Bank, Foochow.
112	RIDLEY, H. N., M.A., F.L.S.	Botanic Gardens, Singa- pore.
113	RODGER, J. P.	Pekan, Pahang.
114	ROST, Dr. REINHOLD (Honorary Member).	India Office Library, Lon- don, S. W.
115	SARAWAK, H. H. The Raja of, K.C.M.G. (Honorary Member)	Kuching, Sarawak.
116	SATOW, E. M., C.M.G. (Honorary Member)	Monte Video, S. America.
117	SCHAALJE, M.	Medan, Deli.
118	SCOTT, Dr. DUNCAN	Batu Gajah, Kinta, Perak.
119	SEAH LIANG SEAH.	Chop "Ilin Hin," Singa- pore.
120	SEAH SONG SEAH	Chop "Chin Hin," Singa- pore.
121	SHELFORD, The Hon'ble T., C.M.G.	Broadfields, Paterson Road, Singapore.
122	SKINNER, The Hon'ble A. M., C.M.G.	Resident Councillor, Pe- nang.

MEMBERS FOR 1892,—*Continued.*

Nos.	Names.	Addresses.
123	SMITH, H. E. Sir CECIL CLEMENTI, M.A., G.C.M.G.	Government House.
124	SOHST, T.	Mount Rosie, Chancery Lane, Singapore.
125	SOUBINDRO MOHUN TAGORE, Raja, Mus. Doc.	Pathuria Ghatha Raj Bati, Calcutta, India.
126	ST. CLAIR, W. G.	Singapore Free Press Office, Singapore.
127	STRINGER, C.	Singapore (Europe).
128	SWETTENHAM, F. A., C.M.G.	Resident, Perak (Europe).
129	SYED MOHAMED BIN AHMED AL SAGOFF	Singapore.
130	SYED ABUBAKER BIN OMAR AL JUNIED	Arabia.
131	SYERS, H. C.	Kwala Lumpur, Selangor.
132	TALBOT, A. P.	Singapore (Europe).
133	THOMPSON, A. B.	Deli, Sumatra.
134	THOROLD, F. THOROLD	Perak (Europe).
135	TREACHER, W. H., C.M.G.	Secretary to Government, Perak.
136	TRÜBNER & Co.	Ludgate Hill, London, E.C.
137	VAN BENINGEN VAN HELSDIN- GEN, Dr. R.	Deli, Sumatra.
138	VERMONT, The Hon'ble J. M. B.	Batu Kawan Estate, Pro- vince Wellesley.
139	WALKER, Lieut.-Col. R. S. F., C.M.G.	Perak.

MEMBERS FOR 1892,—*Continued.*

Nos.	Names.	Addresses.
140	WALKER, H.	Land and Survey Department, Sandakan, B. N. B.
141	WARREN, H. E. Major-General Sir CHARLES, G.C.M.G., K.C.B.,	Singapore.
142	WATSON, E. A.	Bentong, Pahang.
143	WEST, F. G.	Selangor.
144	WHEATLEY, J. J. L.	Muar.
145	WILDMAN, R.	Singapore.
146	WISE, E. A.	Pahang.
147	WRAY, L., Jr.	Perak Museum, Perak.
148	WBENCH, W. T.	Raffles Institution, Singapore.
149	YULE, Colonel HENRY, R.E., C.B. (Honorary Member)	Penywern Road, London, S. W.

Members are requested to inform the Secretary of any change of address or decease of members in order that the list may be as complete as possible.

All communications concerning the publications of the Society should be addressed to the Secretary ; all subscriptions to the Treasurer.

Members may have on application forms authorising their Bankers or Agents to pay their subscription to the Society regularly each year.

11

PROCEEDINGS
OF THE
ANNUAL GENERAL MEETING
OF THE
STRAITS BRANCH
OF THE
ROYAL ASIATIC SOCIETY,
HELD AT THE
RAFFLES MUSEUM
ON
THURSDAY, 28TH JANUARY, 1892.

PRESENT :

The Right Rev. Bishop G. F. HOSE, D.D., *Vice-President*,
H. T. HAUGHTON, Esq., *Honorary Treasurer*, the Hon'ble
J. W. BONSER, C. W. SNEYD KYNMERSLEY, Esq., A. KNIGHT,
Esq., H. L. NORONHA, Esq., Lieut. KELSALL, R.A., *Councillors* ;
E. M. MEREWETHER, Esq., Capt. SMITH, C. B. BUCKLEY, Esq.,
Mr. Justice J. T. GOLDNEY, and H. N. RIDLEY, Esq., *Honorary
Secretary*.

The minutes of the last general meeting were read and confirmed.

The Vice-President opened the proceedings with a short address, in which he referred to the decease of Sir J. FREDERICK DICKSON, K.C.M.G., and to the resolution agreed to by the Council expressing their regret at his death.

The resolution was then read by the Secretary, and Mr. Justice GOLDNEY proposed and Mr. KYNNEERSLEY seconded, that it should be adopted as an expression of the sentiments of the whole Society. This was agreed to unanimously.

The Secretary and Treasurer then read their respective Reports, and Mr. Justice GOLDNEY proposed and Mr. MEREWETHER seconded their adoption. The Reports were adopted.

The Hon'ble J. W. BONSER proposed that the new Council should revise the Rules and have them correctly printed. This was seconded by Mr. H. T. HAUGHTON and carried.

The following Officers were then elected for the ensuing year:—

President,—His Excellency Major-General Sir CHARLES WARREN, G.C.M.G., K.C.B.

Vice-President,—Singapore: The Right Rev. Bishop G. F. HOSE, D.D.; Penang: D. LOGAN, Esq.

Honorary Secretary,—H. N. RIDLEY, Esq.

Honorary Treasurer,—H. T. HAUGHTON, Esq.

Councillors,—A. KNIGHT, Esq., H. L. NORONHA, Esq., C. W. SNEYD KYNNEERSLEY, Esq., Lieut. H. KELSALL, R.A., and the Hon'ble J. W. BONSER.

The Hon'ble J. W. BONSER moved a vote of thanks to the Chairman which was carried unanimously.

—:O:O:—

ANNUAL REPORT
OF THE
COUNCIL
OF THE
STRAITS BRANCH
OF THE
ROYAL ASIATIC SOCIETY,
FOR THE YEAR 1891.

THE Council desires to record its sense of the loss which the Society has sustained in the death of Sir J. FREDERICK DICKSON, K.C.M.G., who for five years has been President of the Society, and, by the great interest he has taken in the affairs and his own wide knowledge of Oriental literature, has contributed so much to the Society's success and progress

During the year, the following new members have been elected:—

<p>Mr. R. DANE. The Hon'ble H. A. O'BRIEN. Mr. H. VON PAPENDRECHT. Mr. Justice GOLDNEY. Mr. E. A. WISE. Mr. F. J. OWEN. Mr. E. B. HUTCHINSON.</p>	<p>His Excellency Major-General Sir CHARLES WARREN, G.C.M.G., K.C.B., F.R.S. Dr. HAVILAND. Mr. ROUNSEVELLE WILDMAN. Mr. A. D. MACHADO.</p>
---	--

We have lost by death Sir FREDERICK A. WELD, G.C.M.G., and Sir J. FREDERICK DICKSON, K.C.M.G.; and the following members have resigned:—

<p>Mr. V. SERGEL. Mr. H. W. C. LEECH. Mr. E. KOEK. Col. S. DUNLOP, C.M.G.</p>	<p>Mr. F. H. GOTTLIEB. Dr. T. I. ROWELL. Mr. O. MUHRY.</p>
---	--

The Council have examined the Rules, and found that various alterations which had been already sanctioned by the Society had not been embodied in the printed Rules, and they recommend that the Rules should be reprinted correctly.

The usual number of Journals was published, containing papers on Ethnological and Natural History subjects. A new edition of the Map of the Malay Peninsula was published, which found a ready sale.

A large number of publications of kindred Societies were received in exchange for those of the Society, and correspondence and exchange has been opened up with the following additional ones:—

Royal Academy of Modena.
Society for the Exploration of Amurland.
Royal University of Upsala.
Natur-Historisches Museum, Hamburg.

The financial position of the Society is satisfactory, the balance at the bank being now \$857.

STRAITS BRANCH OF THE ROYAL ASIATIC SOCIETY.

Dr. Honorary Treasurer's Cash Account from 1st January to 31st December, 1891. Cr.

	\$	c.		\$	c.
1891.			To paid for KOH YEW HEAN Press for Paper, ...	180	00
			" " for setting up Journal No. 22, ...	138	00
			" " for KOH YEW HEAN Press for Paper, ...	164	00
			" " for setting up Journal No. 23, &c., ...	270	00
			" " for Mr. C. D. SHERBORN for Bibliography of Malaya (draft for £17 at 3/1½), ...	109	90
			" " for revising Proof-sheets of Bibliography of Malaya, ...	10	00
			" " for Advertising in local papers, ...	14	00
			" " for Clerk's Salary from 1st January to 30th November, 1891, ...	110	00
			<i>Carried forward, ...</i>	995	90
	\$	c.			
	1,115	77			
	509	47			
	85	00			
	160	00			
	19	50			
	0	15			
	11	00			
	13	45			
			<i>Carried forward, ...</i>	1,914	34

H
Honorary Treasurer's Cash Account from 1st January to 31st December, 1891,—Continued.

1891.		Brought forward, ..	\$ 1,914 34
		Brought forward,...	\$ 995 90
		To paid for Stationery, ...	8 16
		" " for Postages, Cooly-hire and Petty Expenses,	46 45
		" " Discount on Cheques and Bank Notes, ...	0 40
		" " Balance in Chartered Mercantile Bank, ...	857 43
		" " Balance with Honorary Treasurer, ..	6 00

AN OLD MINUTE BY SIR STAMFORD RAFFLES.



THE following interesting record was recently found amongst some old documents in the Singapore Treasury. It is signed by Sir STAMFORD RAFFLES, and the concluding portion of the minute as well as the final signature are in the handwriting of that officer. The date is June, 1823, still legible as when it was written, but the day of the date has been eaten out of the paper.

The Proclamation and the Minute which follows it may accordingly be accepted as one of the first official utterances of the founder of Singapore, after the transfer of the island from the Government of Bencoolen to that of Bengal some three years before its amalgamation with Penang and Malacca.

Apart from its intrinsic value as a state paper, this document is interesting when we compare and contrast the present state of our laws with the provisions there indicated and foreshadowed nearly seventy years ago.

As the Minute is reprinted *in extenso*, I need only draw the reader's special attention to the author's views upon gambling, prostitution, registration of deeds, adulteration, the sanctity of oaths, and municipal regulation, amongst many other points touched upon. The doctrine of the liability of publicans may raise a smile, but it is a theory which still finds support amongst the apostles of temperance in England, where the legal sanction of the publican's errors lies only in the hands of the exponents of the licensing laws.

The Lieutenant-Governor's views upon deportation will, no doubt, be read with interest at the present time: while his brief remarks upon the characteristics of the Malay race, which he knew so well, are as deserving of study and acceptance to-day as when they were penned.

H. A. O'BRIEN.

Singapore, 7th August, 1891.

PROCLAMATION.

Provision having been made by Regulations Nos. III and IV of 1823 for the establishment of an efficient Magistracy at Singapore and for the mode in which local Regulations having the force of Law should be enacted, and by whom such Laws should be administered, it now becomes necessary to state the principles and objects which should be kept in view in framing such Regulations, and, as far as circumstances may admit, to apprise all parties of their respective rights and duties, in order that ignorance thereof may not hereafter be pleaded on the part of any individual or class of people.

The Lieutenant-Governor is, in consequence, induced to give publicity to the following Minute containing the leading principles and objects to be attended to:—

MINUTE BY THE LIEUTENANT-GOVERNOR.

As the population of Singapore will necessarily consist of a mixture, in various proportions, of strangers from all parts of the world having commercial concerns at this Port, though chiefly of Chinese and Malays, it would be impracticable for any Judicial Authority to become perfectly acquainted with the Laws and Customs having the force of Law which are acknowledged in their own countries respectively by the varied classes of so mixed a population, and to administer them in such a manner as to preserve them inviolate even in the mutual intercourse of those classes severally amongst themselves, far more so when justice is to be done between

the Englishman and Chinese, the Bugguese and Hindoo, and the like. On the other hand, to apply the law of Europe direct, with all its accumulated processes and penalties, to a people of whom more than nine-tenths will probably be natives of China and the Malay Archipelago, would be as repugnant to universal and natural justice as it would be inconsistent with the benevolence and liberality which has ever marked the British rule in India.

Under these circumstances, nothing seems to be left but to have recourse to first principles, to use every precaution against the existence of temptation to crime that is found consistent with the perfect liberty of those who have no evil intentions, and when these precautions fail, to secure redress to the injured party, when possible, and such punishment as will be most likely to prevent a repetition of the crime, either by the party himself offending, or by those who may be inclined to follow his example. Nothing should be endured in the Settlement, however sanctioned by the local usage of particular tribes who resort to it, that has either a direct effect, or notoriously strong tendency to endanger the safety or liberty of person or the security of property, and in the same manner no want of what are considered legal formalities in any country should debar a person from having substantial justice rendered to him, *so that legal and moral obligation may never be at variance.*

Taking this as the fundamental principle for the Laws of the Settlement, it may be presumed that no local Regulation would be enacted that the society if left to themselves would not desire to see carried into effect; no public institution or source of expense would exist of which the benefit was not obvious to the enlightened part at least, if not to the whole body of the community, who would therefore soon feel that the Government was not made to tyrannize over the people, but for their protection and happiness.

Under such a system of administration, it is not unreasonable to expect that every facility would be afforded by the mass of the population to the Executive in carrying the Laws into effect, for even the midnight robber and swindler have no

desire that their own persons or property should be liable to those evils which they inflict on the rest of the community, and will readily join in their suppression when other delinquents are the objects of the terrors of the Law.

In carrying such a system into effect, it ought to be fully understood and maintained on all occasions, that while individuals are allowed to *protect* themselves as far as possible against wrongs, the *redress* of wrongs cannot be left to the resentment or the revenge of the parties conceiving themselves injured. That must be done solely by Government through the instrumentality of the Judicial and Executive Officers whom it appoints for that purpose.

No one therefore being allowed to be a judge in his own case, or to revenge his own quarrel, arms or weapons capable of inflicting instant death as habitually worn by the Malays become unnecessary, and, by dispensing with them, the greatest temptation to and power of doing to others the greatest and irremediable wrong in depriving them of life is in a great measure removed. If a man takes another's horse or cow by robbery or theft or under a mistaken idea that he has a right to the property in question, redress can be afforded to him as soon as he is convicted of his crime or discovers his error, but if from revenge or under false impressions a man is suddenly excited to take the life of a fellow creature, it is in vain that he afterwards discovers that he was misled by passion or had been deceived by appearances. It often happens too in these countries that a man who considers himself aggrieved by a particular individual and finding himself in possession of a sharp weapon, attempts the life of every one he meets indiscriminately, and without having any wrong at their hands to complain of. It is impossible to see who may or may not be guilty of such acts of inhuman cruelty, and therefore all should agree to lay aside the use of the weapon that is commonly employed by persons who then transform themselves to wild beasts by giving way to brutal passion.

On the same principle, it has been found by experience that those who indulge frequently in gaming and cockfighting, are not only liable to engage in quarrels with those who

have won their money, but also that they are incited to acts of fraud and robbery in order to obtain the means of amusement or of attempting to retrieve their losses; it is therefore the duty of Government to suppress both gaming and cock-fighting as far as possible without trespassing on the free will of private conduct. No man should be allowed to receive any money either directly or indirectly for conducting a gaming table or cock-pit, and winners of money at such places should be compelled to restore the amount to the losers, and should on no account be permitted to enforce payment from those with whom they have gambled on credit.

Intoxication being a source of personal danger to the community, and the indulgence in that vice being a frequent cause of betraying those who are addicted to it to the commission of acts of dishonesty, it is the duty of a good Magistrate to throw every obstacle in the way. In the first place the Officers of Police should be required to place in constraint any person seen in public in a state of intoxication until he becomes sober, and in the next place the vender of intoxicating articles who supplied him with the means of inebriety, should be visited with proof* and fined, and be liable to make good the amount of any loss which the person so intoxicated can prove he suffered during his inebriety from being unable to take care of himself; the extent of this fine must necessarily be discretionary on the part of the Magistrate, depending principally on the degree of inebriety produced; it should always be of such an amount that the fear of being subject to it may be sufficient to outweigh in the mind of the vender the temptation of profit in the sale of his goods; of course if it should appear in evidence that the individual was supplied with the means of intoxication for the purpose of taking advantage of him in that state, the object converts the simple misdemeanour into a crime according to the particular purpose contemplated, and further punishment to the guilty as well as redress to the individual injured must be awarded accordingly. The use of spirituous liquors, though innocent in moderation, becomes vicious when indulged in to

* *Sic*, probably "reproof."

excess; the consumption may be diminished by the enhancement of price, and in this way the indulgence may be made so expensive as to be only attainable beyond the bounds of moderation by those whose means give them a station in society that induces them to be guarded in their conduct for the sake of preserving the respect of those whose eyes are turned upon them; thus, while gaming as practised by the Chinese and cock-fighting by the Malays are absolutely pernicious in every degree in which they come under public cognizance, the use of opium and spirituous liquors may be repressed by exacting a heavy tax in the way of License from the venders.

There are many important considerations that stand in the way of enacting laws against prostitution, indeed it would, in a country where concubinage is not forbidden, be difficult to draw a line between the concubine and the common prostitute; it is practicable however in some degree, and highly desirable, that the temptation to profit should not exist to induce the seduction of women into this course of life by others of their own sex; the unfortunate prostitute should be treated with compassion, but every obstacle should be thrown in the way of her service being a source of profit to any one but herself. It should therefore be declared unlawful for any person whatever to share the hire or wages of prostitution or to derive any profit or emolument either directly or indirectly by maintaining or procuring prostitutes, as for any parent or guardian of a female or any other person to ask or receive directly or indirectly any reward for bestowing a female in prostitution, any custom, law or usage of the country in which such female or her parents or her guardians were born notwithstanding, reserving only for a jury to advise what constitutes a legal obligation on the man to support the woman thus bestowed, or in other words a contract of marriage by local usage and what a connection of prostitution; the penalty must here also be modified by circumstances. It is much more criminal to induct a girl into prostitution than to facilitate her pursuit of vice after she has entered upon it as a profession.

It may be necessary to make specific Regulations for the

protection of the community generally against fire, both with regard to the construction of buildings, the storing of gunpowder and combustibles, the manufacture of arrack, &c., &c., the power of infringing on a neighbour's property after a fire has broken out either for the purpose of access to the means of extinguishing it or to prevent its spreading to a greater distance.

Boatmen and parties offering themselves publicly for hire may also be subjected to regulation with the view of facilitating the attainment of redress when they are guilty of fraud and negligence.

Weights and measures of the acknowledged standard should be accessible to all, and those used in purchases and sales ought to be in strict conformity with such standards. Certain Magisterial Officers, therefore, should be employed to examine those used by persons who openly keep goods exposed for sale. When found defective the person in whose behalf they are used should be liable to fine proportioned to his supposed means and the apparent degree of fraud resorted to.

Fraud with respect to the quality of articles is a crime more readily detected, and may be left to private prosecution. In giving redress to the individual, punishment ought to be annexed in proportion as the fraud is of an injurious nature.

As a great check to fraud and falsehood, a general Registry Office for all written agreements or engagements which are liable to be made the ground of dispute before a Court of Justice, should be opened for the public. Regulation should be made for the authenticity of the document in the first instance, and either party or any party interested should be entitled to a copy, paying for the same a moderate fee as a compensation for the trouble given to the Registrar and his Establishment. Precaution must of course be taken against the falsification or abstraction of such documents from the Registrar's Office. All deeds which may be so registered should have an avowed preference over one that is not so registered, unless the holder of the latter can shew a clear, distinct and satisfactory cause why he has not been able to have *his* deed registered and the *onus* of establishing this ought decidedly to rest on him.

Nuisances generally speaking may be safely left to the complaint of individuals in each particular instance where the cause of nuisance is not obvious to all, or directly injurious to particular individuals, as crowding the river with vessels, &c., when it may be made subject of special regulation.

All house-holders should be registered and all houses numbered; auctioneers and pawnbrokers should be placed under specific regulations, and none allowed to act as such without giving security for complying with the same and taking out a license for the purpose.

With respect to the employment of informers, it may be observed that Magistrates must have information, but no bad passion should be elicited in the procuring of it. No temptation to lead others to vice for the sake of reward for informing, no inducement to betray confidence, and the act of giving information should be treated as a public and honourable duty.

Precautionary measures being taken on the above principles for preserving the peace and good order of society and removing as far as practicable the immediate temptations to crime and violence, it next becomes necessary to define what shall be considered Crimes, what lawful punishments and how injuries shall be redressed.

By the constitution of England, the absolute rights of the subject are defined as follows:—

1st. "The right of personal security; which consists in a person's legal uninterrupted enjoyment of his life, his limbs, his body, his health and his reputation."

2nd. "The right of personal liberty; which consists in the power of locomotion, of changing situation or removing one's person to whatever place one's own inclination may direct, without imprisonment or restraint, unless by due course of Law."

3rdly. "The right of property; which consists in the use, enjoyment and disposal of all acquisitions without any control or diminution save only by the Laws of the Land."

There seems no reason for denying corresponding rights to all classes of people residing under the protection of the British Flag at Singapore, the Laws of the Land being such

as are or may be enacted under the provisions of Regulation No. III of 1823, dated the 20th January last, with such others of a more general nature as may be directed by a higher Authority or which may necessarily accrue under the provisions of the Legislature and the political circumstances of the Settlement as a Dependence of Great Britain. Admitting these rights to exist, it follows that all acts by which they are invaded are wrongs, that is to say, crimes or injuries.

In the enactment of Laws for securing these rights, legal obligation must never supercede or take place of or be inconsistent with or more or less onerous than moral obligation. The English practice of teaching prisoners to plead not guilty, that they may thus have a chance of escaping from punishment, is inconsistent with this and consequently objectionable. It is indeed right and proper that the Court should inform itself of all the circumstances of a crime from witnesses as well as from the declaration of the prisoner himself. Denial is in fact an aggravation of a crime according to every idea of common sense. It disarms punishment of one of its most beneficial objects by casting a shade of doubt over its justice.

The sanctity of oaths should also be more upheld than in the English Courts. This may be done by never administering them except as a *dernier resort*. If they are not frequently administered, not only will their sanction be more regarded and in this way their breach be less proportionately frequent, but of necessity much more *absolutely* uncommon and consequently much more certainly visited with due punishment in all cases of evidence given before a Court of Justice.

The imprisonment of an unfortunate debtor at the pleasure of the creditor, by which the services of the individual are lost to all parties, seems objectionable in this Settlement, and it is considered that the rights of property may be sufficiently protected by giving to the creditor a right to the value of the debtor's services for a limited period in no case exceeding 5 years, and that the debtor should only be liable to imprisonment in case of fraud, and as far as may be necessary for the security of his person in the event of his not being able to find bail during the process of the Court and for the performance

of the decree after judgment may be passed.

It is well known that the Malay race are sensibly alive to shame, and that in many instances they would prefer death to ignominy. That is a high and honourable feeling and ought to be cherished; let great care be taken to avoid all punishments which are unnecessarily degrading. Both the Malays and Chinese are a reasoning people, and though each may reason in a way peculiar to themselves and different in some respects from our own way of reasoning. This germ of civilization should not be checked. Let no man be punished without a reason assigned. Let the principles of British Law be applied not only with mildness but with a patriarchal kindness and indulgent consideration for the prejudices of each tribe as far as natural justice will allow, but also with reference to their reasoning powers however weak, and that moral principle which, however often disregarded, still exists in the consciences of all men. Let the native institutions as far as regards religious ceremonies, marriage and inheritance be respected when they may not be inconsistent with justice and humanity or injurious to the peace and morals of society.

Let all men be considered equal in the eye of the Law.

Let no man be banished the country without a trial by his peers or by due course of Law.

Let no man be deprived of his liberty without a cause, and no man detained in confinement beyond 48 hours without a right to demand a hearing and trial according to due course of Law.

Let the people have a voice through the Magistracy by which their sentiments may at all times be freely expressed.

In fixing a scale of punishments, the first principle to be attended to is that they should be so graduated as to attach to each particular crime its due and relative punishment according to its enormity, and with regard to the nature of the punishments they should be as mild and humane as the general security of person and property admits of. Severity of punishment defeats its own end, and the laws should in all cases be so mild that no one may be deterred from prosecuting a criminal by considerations of humanity. No feeling

interferes with justice in behalf of a murderer, let this crime be punished by death, and no other. Banishment is the next in order. Solitary confinement proportioned to the degree of the offence or pertinacity of the offender in his criminal course seems the least objectionable of all sorts of punishment. Disgrace may also be a form of punishment, but much caution is required in this respect lest a too frequent enforcement of the punishment destroy the feeling which can alone make it a punishment. Personal chastisement is only for the lower orders who are incapable of feeling the shame of disgrace and may probably be had recourse to in cases of wilful perjury where the falsehood of the witness is palpable and his object particularly mischievous. In all cases let it be considered as no less an object of the Law to afford redress to the party injured than to punish the offender. Compensation should in all cases, where it is possible, be made to the injured party to the extent of the means of the offender, as in the case of the Malay *Bangoon* where when the father is murdered the family are entitled to pecuniary compensation for his loss.

(Signed) T. S. RAFFLES.

With these views and principles the Lieutenant-Governor has this day transmitted to the Acting Magistrates such a graduated Scale of Crimes and Punishments as appears to him sufficient to meet the existing circumstances of the Settlement and to answer the end of substantial justice, with instructions that they will duly deliberate on the subject and after such revision as their local knowledge and experience may suggest, submit the same to the Chief Local Authority with their opinion, and in the form of a Code of Laws to be established for the Settlement and to be in force after publication by the Resident until rescinded by a higher Authority, or altered under the provisions laid down for the enactment of local Laws and Regulations.

The Magistrates have further been required to frame in the form of a Police Regulation, to be approved and published by Government, such further Regulations as may be advisable in that Department.

It is to be hoped that the provisions that will be thus made will be found sufficient for the public peace and the protection of person and property until circumstances may admit of the establishment of a more regular Court of Judicature, every arrangement that can be now made being necessarily of a provisional nature.

Dated at Singapore this day of June, 1823.

(Signed) T. S. RAFFLES.



SOME NOTES ON THE SAKAI DIALECTS

OF THE
MALAY PENINSULA.

BY
HUGH CLIFFORD.

"For the purpose of disclosing to us the real character of language left to itself to follow its own laws, without let or hindrance, a study of Chinese and the Turanian dialects, a study even of the jargons of the savages of Africa, Polynesia and Melanesia is far more instructive than the most minute analysis of Sanskrit or Hebrew."

"On the Stratification of Languages."—MAX MÜLLER.



THE present paper deals with some of the jargons referred to by Professor MAX MÜLLER in the extract from the lecture above quoted, and as the dialects spoken by the largest Sakai tribes of the Peninsula have hitherto practically escaped observation, I trust that even the scanty data in my possession may prove of interest to the readers of this Journal. I do not propose to publish at the present time an exhaustive vocabulary of any of the dialects in question, as the material in my possession is not, in my opinion, sufficiently complete to render any publication that I could now make, of permanent value. I venture to think, however, that as during the last seven years I have visited many aboriginal tribes, and have collected vocabularies of their dialects in several parts of the Peninsula, many of the facts which I have ascertained, and the conclusions to which, in my opinion, these facts point, may be new and worthy of consideration by those who care for philological study. At some future date, when I have had further

opportunities of perfecting and completing my grammar and vocabularies of Sakai dialects, I hope to be able to publish them in a form which will be more thorough and satisfactory than any which I could now attempt, and I shall, therefore, restrict myself in the present paper to a few notes on these dialects, and their connection with Malay.

In the same way, and for the same reasons, I shall not attempt to embody in this article any facts concerning the Semang or Pangan, as the Negritos proper are variously called by the Malays, and even in the Sakai dialects, I shall deal chiefly with the Sēn-oi dialect, which is that with which I have the more intimate acquaintance, and which appears to be the purest form of Sakai extant.

Before going any further, however, it is necessary to clearly state that the aborigines of the Peninsula consist of people of two distinct races. The first, or Sakai, are a light-coloured, slenderly built people with the wavy, abundant hair, and in many cases the drooping nose of the Polynesian. The second, or Semang, who are true Negrito, are short, dark and thickset, with woolly hair, flat features, thick lips and general Negro characteristics. The former have attained a degree of civilization which is far superior to anything which the Negrito have reached. Sakai live in houses, and plant as well as hunt. The Negrito lives by his bow and blow-pipe alone, and lives in a temporary lean-to shed in spots where game is most plentiful. The Sakai affects to look down upon the Negrito, while the latter is a happy-go-lucky, cheery, little hunter who looks down on nobody.

The Sakai tribe is now split up into innumerable clans, each consisting of a few families, living in places surrounded by the Malays, and thus cut off from intercommunication with one another. These small clans, as might be anticipated, show many signs of the influence exerted over them by their Malay neighbours in their language and customs, and though it is comparatively rare to find them embracing the Muhammadan faith, still their civilization is more advanced, and they themselves are more degenerate than their brothers the Sakai of the far interior. There is a clan of Sakai in Kuantan (Pahang)

which actually keeps a Malay *Pēndekar* to teach the young idea the art of Malay fencing! This is an exceptional case, but, as a rule, when a Sakai has come to wear clothes he is morally a ruined man.

Among these clans the dialect spoken is so interspersed with Malay words and phrases, as to be merely a bastard form of the original tongue, and it is not from them that we can learn what the pure Sakai language is. A study of such mutilated dialects, except as a means of watching the rapid decadence of a barbarous tongue, would not repay the labour. Fortunately there is one large district in the Peninsula which is still purely Sakai country, the only permanent inhabitants being all of the Sakai race. This district is situated in almost the exact centre of the Peninsula, and comprises the head waters of the Jelai, Telom, and Serau Rivers in Pahang, the *ulu* of the Batang Padang, Bidor, Kampar, and Plus Rivers in Perak, and of the Galas and Nenggiri Rivers in Kelantan. This large tract of country having from time immemorial been almost exclusively peopled by Sakai, it is here alone that the pure Sakai is met with, and then only in the interior of the district, in places where the Malay language is still unknown. Here we find the Sakai tongue spoken as it has been spoken for generations, and containing comparatively few Malay words, or signs of the influence of the Malay language. Outside influence, as is well known, is a great modifier of all languages, and more especially of the dialects of a wholly unlettered people, and it is, therefore, a matter of some interest to find the Sakai in those places where such influence has scarcely had an opportunity of being brought to bear upon him, or where such disturbing elements have been reduced to an absolute minimum. Comparison with the dialects spoken by the scattered tribes, who have long been surrounded by Malays, shews that in the pure Sakai dialects many words are found which have been replaced by Malay words among the former tribes, and that the general purity of the language has been much destroyed by the introduction of foreign words or phrases.

For these reasons, I shall deal chiefly with the two

dialects of Sakai spoken in the tract of jungle, or "Sakai country" above referred to, and shall only use the vocabularies collected in other parts of the Peninsula for purposes of comparison and illustration.

The two dialects here mentioned are called by the Sakai of the lower half of this district "Sĕn-oi" and "Tĕm-be'", respectively, though the term "Sĕn-oi" is applied to itself by each of the tribes in question, neither of which will allow that the other has any right to the appellation. The term "Gob" (a stranger) is applied to all mankind other than the hill-tribesmen, the term "Gob Mălĕyu" being used for the Malays, "Gob Chĕ-na'" for a Chinaman, and "Gob Bĕ-ŭg'" (the white stranger) for a European. A line drawn from Blanja on the Perak River to the Bidor mountains, and thence to Kuala Nenggiri in the State of Kelantan, will roughly divide these two tribes from one another, the country South of this line being inhabited by Sĕn-oi, and the northern division by Tĕm-be'. Trade and other intercourse is carried on between the Tĕm-be' and Sĕn-oi, but though the dialects spoken are, in my opinion, undoubtedly mere variations one of another, still they differ sufficiently to render it impossible for a Sĕn-oi to understand a Tĕm-be' and *vice versâ*, unless either is familiar with the dialect of the other.

The grounds on which I base my statement as to the near connection between these dialects, and their intimate relation with one another, are the curiously close resemblance which exists between many of the more elementary words in both Sĕn-oi and Tĕm-be'. The following examples, taken almost hap-hazard from my Sakai vocabularies, will amply illustrate the above fact, and, in my opinion, are sufficient to justify the belief that both tongues are sprung from the same source, even if one is not a mere corruption of the other:—

<i>English.</i>	<i>Sĕn-oi.</i>	<i>Tĕm-be'.</i>
To eat	Cha'	Cha'
To go	Chĕp	Chĕp
To give	Ōk	Ōk
To take	Kod	Kod
To laugh	Lĕk	Lĕp

<i>English.</i>	<i>Sěn-oi.</i>	<i>Těm-be'.</i>
Cooked rice	Cha'-na'	Cha'-na'
To be; to have, &c.	Müng	Moh
Fire	Ois	Ois
A bird	Chêp	Chêp
Earth-hand	Tê	Tê
A husband	Ĕn-sîrr	Sîrr
Down stream	Rêh	Rêh
A path	Nong	Nong
The head	Kû-i	Kû-i
The stomach	Kut	Kut
Jungle	Sě-râk ^a	Sě-rok
High	Chě-rûk ^a	Jě-rôk
Tired; fatigued	Gě-hêl	Gě-hêl
To; towards, &c.	Ma'	Ma'

The above examples can easily be multiplied, but they are, in my opinion, sufficient to establish the intimate connection which exists between the Sěn-oi and Těm-be' dialects: nevertheless it must not be supposed that all or even the majority of the words used by these two clans are equally similar to one another. The following specimens of common and elementary words will amply prove that this is not the case, although even in some of them I maintain that a near connection can also be traced:—

<i>English.</i>	<i>Sěn-oi.</i>	<i>Těm-be'.</i>
I, me, we, us	Ĕng	Yêh
You, he, she, it	Hêh	Hâh
Come	Da	Bê
To do; to make	Ū-i	Ĕl
Smoke	Bi-chûl	Jêt-jôt
A lie	Pa'-ho'	Li-bus
A tiger	Rak ^a	Ma'-nu
Flooring	Ching-karr	Nîs
A pillow	Tě-nû-i	Chěng-kol
A mother	Mê	Boh
A father	Bî	Běr
A child	Kě-nod	Kwod
A blow-pipe	Bě-lau	Blâ-hû

<i>English.</i>	<i>Sěn-oi</i>	<i>Tēm-be'.</i>
To hear	Gěr-ték	Kě-yok
A house	Děrk ^a	Dêh
Good; pretty	Bor	Mêng
Here	Dih	Doh
Great; large	Ēn-toi	Mě-nû

The numerals, which are a curious feature in Sakai dialects, also differ in Sěn-oi and Tēm-be'. No Sakai can count, in his own dialect, above three, and among the pure Sakai tribes of the interior no words are in use to express four, five, etc., a word which means "many, etc." being used for any number greater than three. The Sakai who are met with near Malay settlements have, for the most part, adopted the Malay numerals up to ten, but this is merely another instance of the influence which the Malays have exerted on the manner of thought, and consequently on the language of these savages. The numerals as known to the pure Sakai are as follows:—

<i>English.</i>	<i>Sěn-oi.</i>	<i>Tēm-be'.</i>
One	Na'-nu	Nêh
Two	Nar	Nar
Three	Nê	Ne'
Many, etc.	Kěrp ^a	Cha'-tuk ^a

In only one case have I encountered a higher numeral than three among any of the aboriginal dialects of the Peninsula. The tribe in whose vocabulary a word for the numeral "four" was found lived, in 1885, in the jungles near Sadang on the geographical right bank of the Perak River, and consisted at that time of only seven members. I was told by the father of this family that he, his wife and his children were all who survived of a comparatively large clan, which, I make little doubt, will shortly be wholly extinct. Professor MAX MÜLLER tells of a traveller in South America who saw a parrot which was the only living creature that could speak some words of a forgotten tongue. A vocabulary which I compiled of this dialect will act the part of the parrot, and will shortly be the only record that such a dialect was ever spoken by man. This tribe called itself Semang as opposed to the Sakai tribes of

Lěngkûas near Blanja, but its representatives differed very considerably in physical characteristics from the Semang of Ulu Perak, and the Pangan of Ulu Kelantan and Ulu Pahang. The following are the numerals in use among these Semang:—

<i>English.</i>	<i>Semang of Sadang.</i>
One	Nai
Two	Bêh
Three	Pat
Four	Sa-bêh

In other respects, however, this dialect was even more primitive than any Sakai language with which I am acquainted, no word being used for "male," "female" and many other elementary expressions.

As I trust that I have now been able to establish the near connection between the Sên-oi and Têm-be' dialects, I propose, in the following table, to give examples of twenty words found in four Sakai dialects collected in Perak and Pahang, which will, in my opinion, support my contention that the isolated or the scattered Sakai clans all speak languages which have a common origin, and which are closely connected one with another. Of the Sakai of the sea-coast, if indeed these people are really Sakai, and not some other race, I do not profess to treat, as I have had few opportunities of gaining any sufficient knowledge concerning them. The Sakai who speak the dialects, to which the table printed below refers, are all hill-tribes, or land Sakai, and all present the same physical characteristics which I have described in an earlier paragraph of this paper:—

<i>English.</i>	<i>Sên-oi.</i>	<i>Têm-be'.</i>	<i>Blanja dialect.</i>	<i>Slim dialect.</i>
1 To sit	Gû-i	Gûl	Gû-i	Gěr-i
2 That	Jih	Nah	Jih	Jih
3 Recently; just	Pai	Pai	Pai	Pai
4 To throw away	Wêh	Gas	Gas	Gas
5 A pig	Gau	Gaur	Gau	Gau
6 To bring	Ên	Ên	Ên	An
7 A rhinoceros	Â-gâp	Â-gap	Â-gâp	Sě-jâp

<i>English.</i>	<i>Sěn-oi.</i>	<i>Tēm-be'.</i>	<i>Blanja dialect.</i>	<i>Slim dialect.</i>
8 A dog	Cho'	Chû-or	Cho'	Cho'
9 A cloud; the sky	Râ-hu	Râ-hu	Sû-i	Râ-hu
10 To sleep	Bêt	Sě-log	Bêt	Bêt
11 Clothes	Â-bat	Â-bat ²	Â-bat	Â-bau
12 A wife	Kě-nah	Kě-nah	Kě-nah	Kě-nah
13 Water	Tê-u	Org	Tê-u	Tê-u
14 A river	Tê-u	Tê-u	Tê-u	Tê-u
15 Wood	Jě-hu	Jě-hu	Jě-hu	Jě-hu
16 A fowl	Puk	Ma'-nuk	Puk	Puk
17 The moon	Gě-che'	Gě-che'	Gě-che'	Bi-che'
18 To see	Neng	Neh	Neng	Nen
19 To fly	Gi	Guh	Gi	Gi
20 Female	Krě-dol	Ba'-bo'	Krě-dol	Krě-dol

The above examples will suffice for the purposes of illustration, and I may add that every dialect with which I have yet come in contact shews an equal resemblance to one or other of the two principal variations of the Sakai language—the Sěn-oi and Tēm-be' dialects. Leaving the question of the connection which I maintain exists between the various Sakai dialects and sub-dialects, I shall now turn to an examination of some of the characteristics of the Sěn-oi language, which I have selected as being the form of Sakai with which I have the more intimate acquaintance.

This dialect is spoken by about 6,000 people who, as already stated, inhabit the lower portion of the Sakai country, and many of the isolated clans speak dialects which are merely modifications of it. In its grammar, and in its range of sound, it is merely a type of all Sakai dialects, and the vowels and consonants necessary to transliterate it are common to all these jargons.

A fact which strikes the observer as curious in the Sakai dialects is that in so primitive a tongue so vast a variety of sounds should be found. In Sěn-oi there are the vowels *a*, *e*, *i*, *o* and *u*, each of which has four variations, viz., medium, long (ˆ), short (˘), and abrupt ('), with the exception of *u* which has no abrupt tone.

a is pronounced like the vowel sound in *calm*, ex., Da=to come.

á is pronounced like the vowel sound in *blast*, ex., A-gap=a rhinoceros.

ǎ very short *a* sound, ex., Am-cha'=to eat; Dǎk=to bear, to support.

a' similar to sound of the final *ak* in the Malay word *Banyak* in the dialects in which the *k* is silent, ex., Pa'-ho'=to lie; Ma'=to, towards.

e as in the English word *get*, ex., Jel-jol=to hang.

é as in the French word *est*, ex., Jék=an axe; Nê=three.

ě a very short *e* sound, ex., Rěn-tak=the tongue; Mě-nang=a younger brother or sister.

e' similar to the *e* sound in the Malay word *Chapek*, the *k* being silent, ex., Nya'-ne'=a ghost, a spirit.

i as in the English word *hit*, ex., Kě-mit=a mosquito.

í similar to the vowel sound in *cheat*, ex., Ku-f=language, speech; Í-ok^a=to fall.

ĩ a very short *i* sound, shorter than in the English word *hit*, ex., Bĩ-lut=to extinguish; Bi-jog=wet.

i' as in the Malay word *badik*, the *k* being silent, ex., Si'=to fell.

o as in *ox*, ex., Sok=hair.

ó as in *broken*, ex., Shók=the navel; Ók=to give.

õ shorter than the *o* in *ox*, ex., Děng-dõk^a=a branch.

o' similar to the final *ok* in *puchok*, the *k* being silent, ex., Da'-do'=sufficient, enough.

u similar to the vowel sound in *moon*, ex., Nyun=near; Ku-rul=the knee.

ú similar to the *u* sound in *acute*, ex., Kû-ish=a porcupine; Dúl=the handle of a weapon.

ũ rather shorter than the *u* sound in *hug*, ex., Krě-rüg=to pull or pluck out.

In addition to the above vowel sounds, there are the semi-vowels *w* and *y*, pronounced as in English, ex., Wêk=to shoot with a bow; Wih=do not!, desist!; Yatⁿ=a grandfather; Yap=to speak.

There are also the diphthongs *ai*, pronounced *eye*, ex., Pai=recently, just, only just; Laish=an ant.

au pronounced like the vowel sound in *how*, ex., Pě-lau-i=a star; Bě-lau=a blow-pipe.

oi pronounced like the vowel sound in *boy*, ex., Sěn-oi=a man; Chě-noi=a post.

The consonants are as follows, and are similar to those in English, in so far as their value is concerned:—

b, ch, d, g, h, j, k, l, m, n, p, r, s, t, and z.

These consonants combine as follows:—

ng similar to the Malay ڠ , ex., Ngěn-tap=the testicles; Mêng=the cheeks.

ny similar to the Malay ڤ , or the Spanish \tilde{n} , ex., Nyûh=heavy; Nyun=near.

sh, which is equivalent to the Malay ش , ex., Kě-lěsh=a mountain; Jîsh=daylight; Shôk=the navel.

There are two distinct *r*'s in Sěn-oi—one, which is guttural, pronounced very deep down in the throat, I have rendered *r*, ex., Râ-hu=the sky, the heavens; Rěj-ă-roj=lost, to lose one's way: and the other, which I have rendered *rr*, is a rolled *r* more pronounced than the rolled *r* in French.

Another peculiarity of Sěn-oi is the existence of what, for want of a better name, I have termed "nasal finals." These are final *kⁿ*, final *pⁿ*, and final *tⁿ*. They are pronounced by closing the lips and emitting a very slight nasal *n* sound in a descending tone after the final *k*, *p* or *t* has been articulated. A similar sound is found, I believe, in the Hottentot dialects, but I have as yet had no opportunity of comparing Sakai with the African tongues.

Every syllable in Sakai is pronounced separately and distinctly, with a kind of catch between each, and when Malay words are adopted into the dialect, they too are always split up into their component syllables.

The grammar of the language appears to be simple. The following are the principal rules:—

I.—The nominative precedes the verb it governs.

II.—The accusative follows the verb which governs it.

III.—Adjectives and demonstrative pronouns follow the noun they qualify.

IV.—The cases are formed by the use of prepositions, with the exception of the genitive case, which is formed by placing the possessor immediately after the thing possessed. The idea *at, to, towards* is rendered by the preposition *Ma'*. The idea *on* and *from* by the preposition *Kên*. These prepositions (with the additional particle *Pă=ai, ow*) are found to exist in the same or in slightly modified forms in all the dialects of Sakai with which I am acquainted.

V.—There are no inflections of nouns or conjugation of verbs, but the cases are indicated by the personal pronouns, and the words and tenses by means of the auxiliaries *Hôt=to wish, to want, and Tě-lăs=to be done, over, done, finished, enough.*

The following sentences will illustrate all the above rules:—

- | | | | | | |
|-------|-------------------|--------------|--------------|--------------|--------------|
| (i) | Eng | cha' | cha'-na'. | | |
| | ₁ | ₂ | ₃ | | |
| | I | eat | rice. | | |
| | ₁ | ₂ | ₃ | | |
| (ii) | Héh | tě-lăs | kuh | ka' | jih. |
| | ₁ | ₂ | ₃ | ₄ | ₅ |
| | He | has | killed | that | fish. |
| | ₁ | ₂ | ₃ | ₅ | ₄ |
| (iii) | Derk ^a | êng | bê | ma'-chut. | |
| | ₁ | ₂ | ₃ | ₄ | |
| | My | house | [is] | very | small. |
| | ₂ | ₁ | | ₃ | ₄ |
| (iv) | Ôk | i-ôdz | êng | ma' | hê. |
| | ₁ | ₂ | ₃ | ₄ | ₅ |
| | Give | my | chopper | to | him. |
| | ₁ | ₃ | ₂ | ₄ | ₅ |

(v) Hê chíp ma' lor.
 1 2 3 4

Where [are] you going (*Lit.*, You go to where?).

(vi) Êng hôt chíp ma' ta'.
 1 2 3 4 5

I [am] going upstream (*Lit.*, I wish to go to the interior).

(vii) Hê gû-i kěn tê nyun derk^a jh.
 1 2 3 4 5 6 7

Sit you upon the ground near this house.

(viii) Êng pai hól kěn rê.
 1 2 3 4 5

I [have] just arrived from downstream.

(ix) Ia-lok mẽ-nang êng hôt chíp ma' sě-rak^a bort chęp.
 1 2 3 4 5 6 7 8 9

To-morrow [my] brother will go to [the] jungle [to] catch birds.

(x) Bi-chól â-bat^a sěn-oi.
 1 2 3

Smoke [is] the garment [of the] Sakai. (a Sěn-oi proverb).

The above will, I hope, give a sufficiently clear idea of the manner in which Sěn-oi sentences are formed, but before passing on to other matters, there are one or two peculiarities of the Sěn-oi dialect which I should like to mention. One is the extraordinarily close resemblance which exists between many words in this dialect, and which, at first, is very puzzling to one who has to compile his own grammar and vocabulary of these aboriginal tongues. The following examples will

illustrate this peculiarity with sufficient clearness :—

Cho'	=	A dog
Choh	=	To void
Chók	=	To stab
Chók ⁿ	=	A root, a rattan
Dol	=	To place, to set down
Dúl	=	The handle of a knife
Êng	=	I, we
Ên	=	To bring
Hol	=	A small black ape—the <i>siamang</i> of the Malays
Hól	=	To arrive
Ka'	=	A fish
Kah	=	To cut, to gash
Kol	=	Pith
Kól	=	To fall down
Kû-i	=	The head
Ku-f	=	Language, speech

The colours which have names in the Sakai language are as follows :—

Rě-ngah	=	Black
Bi-úg	=	White
Chěng-ul	=	Red

All dark colours are included under "Black;" green, blue, mauve, &c. under "White;" and crimson, orange, yellow and brown under "Red."

Hitherto all efforts to connect the Sěn-oi dialects with any other tongue have failed, so far as my attempts are concerned, but I have now arranged to procure some further vocabularies for purposes of comparison, and hope to be able to discover a connection either with the Papuan dialects or with some of the jargons spoken by the Dyaks or other aboriginal tribes of the Malay Archipelago. The statement made by Mr. VAUGHAN STEVENS that the Sakai dialects were closely connected with the language of Thibet is totally incorrect. After a careful comparison of the grammars and vocabularies of the two tongues, I am in a position to state that they have neither a root nor a grammatical form in common. Thibetan is a polysyllabic language with an elaborate grammar,

which in no point is similar with anything found in Sakai or Semang. The language spoken by the latter people I hope to be able to connect with the inhabitants of the Andaman Islands. The physical characteristics of the two people are strikingly similar, and a comparison of their dialects will be full of interest. It would be somewhat premature, however, to do more than study the Sakai dialects themselves, and, as already stated, I have not hitherto succeeded in finding a single root in common in any of the vocabularies (including that of the Veddahs of Ceylon, which would appear to be a bastard Indian dialect) which I have as yet been able to examine.

There is yet another point on which I should wish to touch before concluding these notes. I refer to the connection between Sakai and Malay. In the introduction to his *Malay Manual*, Mr. W. E. MAXWELL has propounded a theory to the effect that some Malay roots may possibly be derived from Sakai. He says:—"Another characteristic list of words might "be made compounded with the monosyllable *Tang* (which in "Sakai and Semang means 'hand'), and conveying an idea "of seizing or holding. *Tang-an*=the hand; *Tang-kap*=to "seize," &c., &c.

Now, in Sën-oi the word for "hand" is *Terk* and in Tēm-be' it is *Pih*. Semang dialects are absolutely distinct from Sakai, having but few roots in common, and in no dialect that I know does the word *Tang* occur meaning a "hand." This being so, I should be disinclined to accept Mr. MAXWELL'S theory, the more so as there is much evidence to prove that at the time the Malays first penetrated into the Peninsula and other Malay countries, they spoke a language which, both in its words and in the rules by which its substantives were formed, did not differ appreciably from the Malay of to-day.

The names *duri-an*, the thorny fruit, *rambut-an*, the hairy fruit, and *pulas-an*, the twisted fruit, were all given to the fruits in question (which are indigenous in the Malay countries alone) by a people in whose language the words *duri*, a thorn, *rambut*, hair, and *pulas*, to twist, were all accepted terms, and at a period when the inseparable affix *an* had come to be employed for the formation of substantives as it is to the

present day. The language being thus formed when the Malays first arrived in the Peninsula would preclude the possibility of the Malay language deriving elementary roots from Sakai dialects. Among the Sakai tribes, too, *sem-pa*,¹ a durian, has no connection with *tër-lákⁿ*, a thorn, and the names for the other fruits are equally distinct, and seeing that even to the present day the durian groves of the far interior are one of the principal factors in the Sakai's annual food supply, it is only to be expected that the name of so important an article of food should be one of the first elementary words to be embodied in the language of a primitive people dwelling in the Malay Peninsula.

But the evidence leads us further yet. Most people who have travelled in the interior of the Peninsula have seen the stone implements which are frequently discovered by the natives. These implements, called *batu halilintar* or thunder bolts, by the Malays, who believe them to be the bolts hurled from the heavens during storms, are of three kinds—stone axes, shaped somewhat like the Malay *bliong*; spear-heads; and choppers. At the present day similar tools wrought in metal are sold to the Sakai by the Malays, but it is a significant fact that they are all called by Sakai names by the aborigines. The following are the names in question:—

<i>English.</i>	<i>Malay.</i>	<i>Sěn-oi.</i>	<i>Tëm-be.¹</i>
Axe	Bliong	Jêk	Jêk
Spear	Lembang	Ta-rok	Bě-lush
Chopper	Parang	I-ôdz	I-ôdz

Now, in spite of VOLTAIRE'S famous saying that "*pour Messieurs les étymologistes les consonnes ne lui coûtent rien et les voyelles bien peu de chose,*" I doubt whether any one will maintain that any connection exists between the Malay and aboriginal words for these implements. But the Malays also sell hatchets (*kapak*) to the Sakai, and this implement, which has no equivalent among the stone implements of the Peninsula, is called by the Malay name, *ku'-pak* being the Sakai modified form. Now these facts, I contend, point to the conclusion that at one time the tools made of stone were

used by the Sakai and bore the names which are now given to the metal tools, of a similar shape, introduced by the Malays. That the metal weapons were introduced by a foreign race is proved by the fact that even to this day the pure Sakai have no knowledge of the art of fusing metals. That the Malays were the race which introduced these metal tools to the Sakai is rendered more probable by the fact that the weapons for which equivalents do not occur among the stone implements bear Malay names as already stated. If this point is allowed, there remains no alternative but to accept, what in my opinion is an undoubted fact, viz., that the Malays invaded the Peninsula at a period when they had attained to a considerable degree of knowledge in the useful arts, and their language formed in its essential characteristics, and that, therefore, the Malay language does not, and could not possibly, owe anything (saving perhaps the names of a few plants and beasts) to Sakai roots.

In the above remarks I have had occasion to state that the word *Kapak*, a hatchet, is adopted by the Sakai and becomes *Ka'-pak* as pronounced by the aborigines. Now this needs a word of comment, as it has frequently been remarked with surprise that the Sakai in adopting Malay words ending with *k*, which letter is silent in the dialects of the Peninsula, always give the silent *k* its full *written* value. Among other edifying deductions which have been drawn from this simple fact, it has been gravely argued that the peculiarity has arisen from the fact that the Malays of the Peninsula at some former period spoke as do their neighbours of Borneo and Sumatra, who pronounce all final *k*'s. That the Sakai learned it at that period, and have never abandoned the practice, though why they should have retained a peculiarity of pronunciation which the Malays of the Peninsula have relinquished, was not explained.

Now, the true explanation of this matter really is that in Sakai there are certain phonetic laws, of which the Sakai themselves are unconscious, but which, so far as I yet know, are employed without exception in all cases where Malay words are adopted into Sĕn-oi. These rules can be stated,

but not explained, any more than it can be told why in Malay words with an initial *s* form the verb by dropping that letter and substituting *meny* မှ်. In both Sakai and Malay it is probable that euphony to the native ear is alone responsible for such rules. In Sakai the following are the phonetic changes which words adopted by the aborigines from Malay undergo, according to the original termination of the Malay word:—

- (I) All Malay words split up into component syllables.
- (II) A final vowel becomes that vowel in its abrupt (') form. Thus, *luka*, a wound, becomes in Sên-oi *lu-ka'*.
- (III) Final *m* becomes final nasal *p*ⁿ. The vowel in the last syllable sometimes changing from *a* to *u*. Thus, *jeram*, a rapid, becomes *jě-rupⁿ*.
- (IV) Final *ng* becomes final nasal *k*ⁿ. Thus, *kuching*, a cat, becomes *ku-chikⁿ*; *cherang*, a clearing, becomes *chě-rakⁿ*.

(V) Final *ak*, *ek*, *ik*, *ok* and *uk* in Malay, though silent in that language, are pronounced as they are written and not as they are pronounced by the Malays of the Peninsula.

The reason for rule (V) is apparent. The Malay sound of the finals written *ak*, *ek*, *ik*, *ok* and *uk* are already in use by the Sakai for another class of words adopted from the Malay, as will be seen by rule (II) above. As stated in rule (IV), *k*ⁿ, the only other appropriate sound, is employed for words adopted from the Malay ending in *ng*. The only resource left to the Sakai is, therefore, to pronounce the *k* in order to distinguish between the final *a*, *e*, *i*, *o* or *u*, and the final *ak*, *ek*, *ik*, *ok* and *uk*, and this is accordingly done.

I have now written all that I think it advisable to publish at the present time, and finally I will briefly recapitulate the conclusions which, I think, are shewn to be probable, if not certain, from the evidence which I have submitted. Firstly, then, I hold that the Sakai all speak various dialects of a common tongue; secondly, that they are more ancient inhabitants of the Peninsula than are the Malays; and lastly, that the former have not derived any elementary roots from the Sakai.

15th September, 1891.

AN ACCOUNT OF A JOURNEY
ACROSS THE MALAY PENINSULA
FROM KOH LAK TO MERGUI.

BY

ARTHUR KEITH, M. B., C. M.



ULLY a century ago much of the traffic between Siam and the West passed over the Malay Peninsula between Mergui and Koh Lak, and in the month of June of this year, having occasion to visit Mergui, I chose this old route. In those olden days carriages with ladies riding in them and driven by cockaded coachmen were wont to pass to and fro by this route, but the remains of the old road that can be seen to-day lead one to suspect that such a statement was the gloss some old writer put upon the rustic, squeaking bullock-carts with their native drivers that wore their hair cut in a shoe-brush fashion by way of a cockade. In those days Tenasserim stood at the terminus on the Burmese side, then the capital of a kingdom and often spoken of as a Venice of the East, for she possessed a large fleet of vessels that carried her commerce all over the Bay of Bengal, while at the Siamese terminus, Koh Lak sheltered many junks and big boats that carried the overland traffic to Ayuthia or to Bangkok and further if necessary. But the rapid transit brought about by steam has killed this trade, Tenasserim is little better than a memory, a white man is a rare sight in Koh Lak, and the jungle has seized and made much of the road its own again.

Koh Lak itself is a small island rising from the sea like a rudely splintered church tower, but on the adjacent mainland is a small town bearing the same name. Along the sea-board here, are numerous small patches of paddy fields with rude bamboo houses, raised high above the ground, scattered amongst them, and numerous water-buffaloes feeding in the deserted clearings. Looking westwards, some ten miles distant, one can see Khow Maun (known on the Burmese side as Khow Mordaun) rising from the ridge that divides Siam from Burma, over the shoulder of which the path we were to take turned. It was then the 7th of June, and the South-West Monsoon had set in, and although upon the East side of the Peninsula not a drop of rain was falling, we could see heavy masses of vapour lying upon the West side of the hills, and evidently rain falling plentifully on there.

My guide, who had been known to me for over two years as a miner and hunter at Bangtaphan, was a Siamese with a dash of Burmese blood in him and had relations living on both sides of the Peninsula whom he visited frequently, and known to those of the Siamese side as *Nai Yeet*, to those in Burma as *Moung See*. He would assent gravely to the most improbable statements, and the real state of matters never stood a moment in his way from putting them as he thought they were wanted to be. He had announced this road as eminently suited for elephants, and that a pony might go along, so that I was tempted to travel at ease. Luckily I had with me another man, NUAN by name, a most faithful Siamese and quite an ornament to the skin he wears.

Having followed the coast line from Bangtaphan northwards to Koh Lak, we there turned inland and made for the pass across the hills at Khow Maun. For the first four miles, a rough dray-path led us over a shallow, level, damp soil carrying a stunted jungle clear overhead, so that we made a swinging pace. But at the end of the four miles the path ended in a clearing in the jungle covered with tall lalang grass and full of shallow pools, and for the following five miles we followed an irregular footpath that wound through the jungle. The soil was here deeper and drier, and the trees grew larger and

offered great obstruction to our passage. Before entering this footpath we noticed by the wayside a small clearing covered with green grass and where probably had stood a rest-house for travellers. After leaving the houses on the coast we passed but one human habitation, some six miles inland—the house of one of those men that live by the produce of the jungle. Darkness was beginning to fall when we emerged from the ill-marked footpath on to a deeply rutted dray-path that wound up the slopes of Khow Maun, and by the side of this path we spread our *kajangs*, and spent the night.

MOUNG SEE said the path before us was now free from all obstruction, but as he had made the same statement every morning for the last four days, and as we had found it utterly false—for it had cost us an immense amount of labour to bring the elephant that distance, owing to the obstruction offered by the numerous low branches across the road—I went in front to inspect the path, and at the shoulder of the hill found that it was completely obliterated by the jungle. On the slope of the hill the rains kept the road scoured, so that vegetation could not spring up on it, but on the shoulder of the hill, the soil being left at peace, was soon seized and grown over. Sending the elephant and all superfluous baggage back, and taking a waterproof coat, a blanket, a gun, rice and fish for five days, as well as other things necessary for my business at Mergui, NUAN, MOUNG SEE and I set out together.

On the side of Khow Maun, along this dray-path, are many shallow trenches running round the hill as if at one time an army had encamped there. Just on the shoulder of the hill, a few large spreading trees shelter a considerable expanse of sward, whereon at one time stood a temple—a rest-house for the overland travellers. Near by, in a ravine a little lower down, is a well with excellent clear water, for all the streams were dry, and we had been drinking from the buffalo pools, which the natives held drinkable if there was no marked odour, so that this water was very welcome. The country here is granitic, full of deep narrow ravines, and here and there we saw the deep cuttings that had been made to carry the old road through them.

There is a steep ascent for the last two or three hundred yards before gaining the top of the pass, which is about 750 feet above the sea level, and where there is a mound covered with all sorts of idols, but the prevailing type was a small, very fat clay elephant set upon exceedingly stunted limbs. This was an altar to the spirit of the hills, and NUAN and MOUNG SEE coming up, stooped down and made their obeisance, and NUAN in a prayer informed the spirit of my state and station, and begged he would not think hardly of me for having but two followers, and told him of the more numerous retinue I had brought away with me, and finally concluded his prayer with this promise "and now if you will give us a safe conduct to Muang Meerit (Mergui), and keep us from robbers, tigers and "all the other ills that beset travellers, we will make a great "deal of *merit* when we safely reach our journey's end."

On the Burmese side everything was wet with the constant drizzle. The rain clouds must hang very low, for the range of hills here are very little over one thousand feet in height, yet they are high enough to determine the rainy season on either side. From a glimpse through the trees, which we got descending from the pass, we could see that the country in front of us was entirely covered with forest jungle, with hills in the distance. The foot-path leads down a gentle descent, and at the foot of the hills we found the country uneven, cut up by ravines, with a deep alluvial soil covered by a lofty evergreen forest. We reached a stream about seven miles from the pass, and, beneath a large tree with very thick foliage, upon its bank we spent the night.

About ten o'clock a downpour of rain commenced. I drew my water-proof coat over my blanket, while NUAN and MOUNG SEE crouched over the fire; but the wet began to steal in all round and the rain drowned the fire, so that we waited anxiously for morning to break. As soon as we could see, we packed up our things and set out, the rain coming heavier than ever. For five miles we went through a luxuriant bamboo jungle, where we started a herd of wild buffaloes, and jungle fowl were abundant, but as we had only a Colt's repeating rifle with us we were unable to procure any for our pot. By the

side of the path were the ruins of a brick temple, which at one time must have been rather an elaborate structure, and there the men again made their obeisance.

When we emerged from the bamboo jungle we struck a stream coming from the S. E., shallow and about thirty feet wide. On its left bank were three dilapidated huts, where probably Chinamen or natives had resided while prospecting the stream which is reported to carry gold. We followed this stream—the *Klong Pan Peng*—for some distance, then crossed and recrossed it several times, until we reached an open space in which there was a rude hut called by MOUNG SEE “the house of the father of the buffaloes.” The rain cleared away, and we had time to dry our clothes and cook some food before it set in again, and continued all night.

The soil here is deep and carries lofty trees with a rich and thick undergrowth of trailing and twining plants. Ever since leaving the bamboo jungle leeches lay thickly in wait for us, but next day they surrounded us like besetting sins. On the path in front one could see a perfect little forest of miniature elephant trunks nodding on the ground, and no railway guard catches the moving foot-board so cleverly as they catch the traveller. In a clearing an attempt was made to rid ourselves of the enemy, but they boarded us quite as quickly as we could throw them over. Few travellers pass this way, and how these crowds get a living I do not know.

Here and there, through this part of the jungle we came on small pieces of green sward surrounded by large trees, and sometimes we passed larger clearings. Evidently people had at one time a habitation here, and even now, during the dry season, herdsmen drive their cattle up from the lowland flats of Burma to graze in these patches and in the bamboo jungle.

About seven miles from the field of the father of the buffaloes, this stream from the S. E., which we had followed and crossed, joins a large stream from the N. E., and just at the junction of the streams our path ran into the river. After half a day's fruitless search for it on the further side of the river amidst a constant downpour, we recrossed to a sand bank, on which we threw up a hut. We were fortunate enough, during

our search for the path, to find a turtle weighing some 45 lbs. feeding in a marsh by the river. He seemed sadly astonished when the two men started to remove his breast plate, but he ought to have congratulated himself on escaping the sad fate of the small land tortoise which is invariably cooked by being hung over a fire and roasted alive.

The rain continued all night, and next morning the river was so flooded that we could not cross it. So a council was held and we determined to build a raft of bamboos with which to proceed down the river, although NUAN objected sadly to trust himself to the mercy of the river spirit. By four o'clock over forty bamboos had been cut, lashed together in three tiers, so that the raft had rather the appearance of a rather broad ladder. We were anxious to test its capabilities, and foolishly started that night. It went swiftly and smoothly along the flooded stream until a difference of opinion between NUAN and MOUNG SEE landed it broadside on a small island in the middle of the stream, and in a moment we were over and in the water. Luckily I had tied everything to the raft with the exception of my only pair of shoes, and when the raft was righted we found they were gone and everything soaking wet. Darkness came down, so we moored. The rain was constant; our clothes, blankets and matches wet; and the best piece of ground we could get was damp and marshy, so that we longed all night for the light of morning.

The country now became covered with jungle-clad ranges of hills, set closely together, separated only by the narrowest ravines, and rising up some 500 feet or more. These hills ran right across the course of the river, so that it seemed to dash against the first range, then eddying along its flanks burst through the first vulnerable point it reached and dashed against the next range, where it again searched for and found an exit, and so bounding and turning, rushing and eddying, it at last burst through this hilly country and sailed out on the flatter country beyond. It took us a day and an half to get through this tortuous channel, sixteen hours punting at $2\frac{1}{2}$ miles an hour, so that to reach a point some fifteen miles distant as the crow flies we had to cover about forty.

After spending such an uncomfortable night, we found ourselves just at the entrance to this hilly country. The river was now in big flood, and the rapids were full of great boulders, against which the river dashed itself. Below those rapids came great silent whirlpools in the shadow of the hills, so deep that our long bamboo punt poles failed to reach the bottom, and our raft rotated again and again. It required all our energies and wits to shoot those rapids, and often we bounced against the snags and jutting boulders, coming off as a rule with but slight damage to the raft, but more than once we were on the eve of complete disaster, when we got righted almost by miracle.

We had just shot a rapid, and were taking matters easily after our exertions, when our attention was suddenly attracted by a thundering tearing noise as of some large animal bursting through the narrow coppice between the hill and the river. Presently a great male banteng (*Bos frontalis*) appeared on the bank about fifty yards behind us, plunged into the flood and made swiftly for the other side. At one time he looked threateningly towards us, so we were glad to get rid of the brute, but just as we were turning the next bend we again heard the thunder of his hoofs, and he instantly appeared on the bank quite opposite us. As he plunged into the river the men threw their punt poles from them and covered on the raft, while I hastily undid the fastenings of my gun and loaded it. But by that time the bull had thought better of it and had made across the river just in front of us and was ascending the opposite bank as I took a flying shot at him, but he disappeared evidently none the worse. The beast, I believe, had not the slightest intention of molesting us; but the hills threw their steepest sides so close to the river, that he had to cross and recross to get along the narrow bank that was sometimes on one side sometimes on the other.

A little after midday the rain ceased, and in a blink of sunshine that followed, we dried our matches and shortly afterwards moored our raft and prepared to spend the night. We made a great fire, dried our clothes and blankets, cooked the

the last of our rice, and threw up a rude roof of leaves, so that when the rain fell with the darkness we were prepared for it.

It required some cajoling to get NUAN to again trust his life on the raft, for said he "Last night the river spirit came"—here NUAN held out his hands and shook his head quite despondently. But the river was the only way out of the hills; we knew dwellings could not be far off, and the river had abated somewhat; these and other inducements allowed us to again resume our journey in the morning. Early in the forenoon we came upon tracks of honey and gum-dammar collectors, and at midday we sailed out between two hills that stood as sentinels to a flat, jungle-covered country with small rounded hills scattered over it. The soil was deep and eminently suited for agricultural uses, but we saw no vestige of cultivation until the day was well spent, when we suddenly came upon a small settlement of houses. These bamboo houses were evidently of recent erection, and stood amongst plantains and maize, which grew amongst the fresh trunks of lately felled trees. The inhabitants were Siamese, most of whom had recently come over from the Provinces on the East coast, and many of them had worked as miners at Bangtaphan. Siamese priests had come up from Wat Kew, an old Siamese settlement a day's journey down the river, and had taken advantage of the flood which had made the river navigable for big boats to visit this outlying Colony of their parishioners and collect their tithes. All night long the old priest with his two young brethren were fêted, so that we were allowed to spend the night there in comparative peace.

Next morning the priests set out for their 'Wat,' and we obtained permission to occupy part of the boat. A little further down we reached a larger village, also a Siamese Colony—for the Siamese occupy the valley of this river down to Tenasserim. There the priests stopped to take their midday meal. The headman told me he had been settled there for upwards of eight years, and that he came from the Province of Koowi. The soil, he also informed me, was much richer than any on the East Coast, and that they cultivate only a little rice, devoting their time more to the raising of

bananas and maize, which they sell down the river, and besides they keep a good few buffaloes. I have no doubt also they enjoy the security of property and freedom from oppression under British protection.

After they had finished their meal we set out again, the young priests '*chowing*' the boat while the old man reclined within. Round about him, built in great piles were the worthy man's presents from his parishioners, consisting of great heaps of half roasted fish, baskets full of sweetmeats and fancy cakes, bags of rice and bunches of bananas, betel-nuts and coco-nuts '*galore*,' in fact stores large enough to hold a great priesthood eating for weeks. In fact, so full was the boat of good things (and I believe the priests were quite as full) that, although the boat was a fairly large one, there was neither room to sit or stand, and it was quite a treat to see the old priest's little boy scrambling amongst those things to supply the betel-nut wants of his superior.

The river got wider, deeper and flowed more slowly, but the surrounding country was still of the same character—flat and jungle-covered, with rounded hills here and there. By landing some distance above Wat Keo and walking over a neck of land, we were able to cut a good few miles of the river and reach the '*Wat*' that night, where we slept. A beautiful park surrounds the abode of the priests, and the Wat is built after the Burmese design, but all the priests are Siamese, some of them coming from the Provinces on the East Coast. The village of Wat Keo (or Takay as it is called on the maps) has about 400 inhabitants, mostly Siamese, and is surrounded by large paddy fields and extensive pasture land. NUAN had now an opportunity of fulfilling his vow, for in his hour of danger he had promised the water spirit, at the first Wat he reached, to make merit to the extent of ten ticals, but now was quite convinced it was a rash promise and considered three ticals quite enough.

Next morning we left Wat Keo and proceeded further down the river. Only two house-boats were available, both belonging to local trading Chinamen, and these, practising their usual policy on the stranger, extracted three times the

customary amount, and no doubt divided the profits. Moreover the boat was small, one had to crawl into it, and when there lie on one's back. The Chinaman was an old Perak miner who had come up to Mergui, married a woman of the country, started trading on the river, and now had two daughters to assist him in 'chowing' the boat.

The following morning we arrived at Tenasserim, once the proud capital of a kingdom of that name. Its ruler, its power, its people and its fleet are all gone, and all that remains to mark its ancient claims of a Venice of the East is the dilapidated remains of a terrace leading up from the river. The inhabitants are mostly Burmese and do not number more than 600, and there seems to be no life in the place. Yet it occupies a situation surpassingly fine. The foot of the high bank on which it sits is swept by the smaller Tenasserim River, which a little below the town bends to meet the larger river coming circling from the North and enclosing between them an expansive 'haugh' of green sward. The 'haugh', the circling rivers and the town are enclosed in a circuit of forest-clad hills, so that Tenasserim sits on the South side of an amphitheatre.

Just as we were leaving Tenasserim, two little Burmese boys came running down to the boat and asked in good English "Where are you going, Sir?" They were wonderfully intelligent lads; they attended school in Mergui, but were then home for holidays, their father being a merchant in Tenasserim.

The river between Mergui and Tenasserim is broad and deep, in many parts it really looks like a series of lakes surrounded by hills; and here and there along its banks are fishing villages. During the rainy season steam launches, drawing four or five feet of water, can go to Tenasserim, and formerly one of the British India steamships used to pass up the river to within eight miles of Tenasserim, where her further passage was prevented by a bar of rocks that cross the river there.

The mouth of the river is surrounded by numerous swampy mangrove-covered islands, and sailing along these we came

in sight of Mergui next forenoon. The town contains about 15,000 inhabitants, and is the capital of the southern part of Lower Burma. The Government houses are situated on a small plateau that rises from the town with the native houses—mostly built of wood—round it reaching down to the water's edge. The inhabitants are mostly Burmese, but Chinamen are the active traders.

NUAN went ashore and procured me shoes and other necessaries, and fitted me out more as a Chinaman than as a European, to face civilization. I was most hospitably entertained by Mr. BATTEN and Mr. HUGHES, and after a few days left on board the *S. S. Mergui* to return to Bangtaphan *viâ* the isthmus of Kra.



THE THERMAL SPRINGS OF SELANGOR AND MALACCA.

BY

D R. W. B O T T, F.C.S., F.G.C.S., &c.



THE following account represents a brief summary of the results of two official visits to Selangor and Malacca undertaken in August and November, 1890, respectively. Many of the results have already appeared in the report published by Government.

A great many details—such as description of apparatus employed, methods of collecting gases, and analytical methods used, which really form the most valuable part of the paper from a strictly scientific point of view—have been omitted, as being obviously not suitable for publication in this Journal; these will be found in the complete paper which will shortly be published in England and Germany (Proc. Royal Soc. and Berlin Chem. Soc.). In reference to the analytical data given, I wish to point out, that these only represent the approximate composition of the water of the springs, owing to unavoidable admixture of ordinary surface water in most cases. I hope at a future date to be able to visit these springs again, provided with such apparatus and appliances as will enable me to collect the water absolutely free from foreign admixtures. The numbers given, however, if not absolutely true, will, I believe, be found to require but a trifling correction in each case.

GENERAL NATURE OF THE SPRINGS.

All the springs visited by me and described more fully later on, belong to the class of "Simple Thermal Waters," viz., they are characterised by a high temperature, but do not contain any considerable amount of dissolved mineral matter. This circumstance, and the small amount of lime in the water, seems to indicate that all these springs rise from silicious and not from calcareous tracts, although no strictly reliable relation can ordinarily be established in such cases between the chemical nature of a spring and its ultimate source. The immediate and obvious surroundings of the springs are old granitic formations. This also applies to those springs which are situated in padi swamps, and where granite, though not apparent, is found below the mud surface. In addition to common and porphyritic granite, quartz, as rock and pebbles, diorite, gneiss and greissen (a kind of granite practically free from felspar which often forms the matrix of tin lodes) are found, but limestone is not apparent. The latter remark, however, does not apply to the springs of Ulu Selangor, which I was unable to visit in person.

The springs themselves appear as pools or puddles of hot water, varying in area from 1 to 3 or more square yards, and generally fed by two or more distinct streams or jets issuing from holes and crevices in the bottom of the basin. Bubbles of gas are observed continually to emerge with these jets and ascend through the supernatant hot water. These consist essentially of nitrogen and carbon dioxide with a small quantity of hydrogen sulphide and traces of hydrogen and marsh gas. The presence of hydrogen sulphide is apparent from the sulphurous smell in the vicinity of the springs and the water itself owes to this gas a slight smell and taste. The beds of the springs consist of granitic rock more or less decomposed on the surface and coated with silicious sinter. The floor of these beds is covered with all kinds of loose vegetable and mineral debris in the form of granular sediment or soft fibrous and gelatinous masses. The latter accumulate in parts and often form a thick continuous layer on

the surface of the water having a green and in some cases a bright orange red colour. The consistency and general appearance of this substance varies from a loose, fibrous structure resembling soft, dense moss, to a gelatinous, homogeneous texture not unlike India-rubber. A careful microscopical examination of these masses showed that they consist of decayed organic matter derived from leaves and other vegetable substances which find their way into the water and undergo decomposition at a comparatively high temperature. They represent a dense magma of cells of most variable shape and appearance, representative of almost every kind of vegetable cell in existence. How far this growth may be looked upon as a whole, as an individual, as a living thing increasing actively, or a dead product growing passively by mechanical addition,—these are questions of considerable interest, to be settled by a separate investigation, and I hope to discuss this subject fully in a future paper. In several specimens of the vegetable matter from Ulu Klang and Setapak, a distinct form of algæ was found, and I have since been informed, that similar specimens had been found in Ayer Panas water, Malacca, and sent to the British Museum for examination. It appears, however, that this has led to no further results. The temperature of the springs ranges from about 100° to 185° F. in Selangor, and 95° to 130° F. in Malacca. It varies moreover in each spring according to the distance from the feeders, being highest at the point where they enter.

A number of complete analyses were made of the air in the vicinity of the springs. The air was found to be of normal composition though free from active oxygen or ozone. Air collected close to the springs was found to contain traces of hydrogen sulphide and rather more than the normal amount of carbonic acid and ammonia. The details of these analyses will be given subsequently.

All the springs contain a considerable amount of dissolved silica. On keeping samples of the water part of this is deposited in a pulverulent state, sometimes, however, very thin, soft, transparent, glistening films separate from the water and these on analysis are found to consist of silica, organic matter and traces of sulphur.

SPRINGS IN SELANGOR.

SETAPAK SPRING.

Temperature, ...	47-50° Centigrade (118-122° Fahrenheit).
Total solids dissolved in water,	18.35 grains per gallon.
Chlorine,	0.4 " "
Hardness,	3.6 " "
Free Ammonia,	0.036 parts per million.
Albuminoid Ammonia,	0.10 " " "

The quantity of dissolved constituents amounts to 3.10108 parts in 10,000 of the water, viz. :—

Calcium Carbonate,	0.23	parts in 10,000.
Calcium Sulphate,	0.25	" "
Magnesium "	0.014	" "
Sodium "	0.21	" "
Potassium "	0.08	" "
Sodium Carbonate,	0.43	" "
Ammonium "	0.00108	" "
Sodium Chloride,	0.043	" "
Potassium "	0.005	" "
Lithium "	trace	" "
Sodium Sulphide,	0.025	" "
Boric Acid,	trace	" "
Carbonic Acid,	0.71	" "
Nitrogen,	0.09	" "
Hydrogen Sulphide,	0.034	" "
Organic Matter,	0.120	" "
Silica,	0.859	" "
	<u>3.10108</u>	" "

The gases escaping from the spring were collected and analysed, giving the following percentage composition :—

Nitrogen,	97 %.
Carbonic Acid,	2.5 %.
Hydrogen Sulphide,	0.5 %.
Hydrogen,	trace.
Marsh Gas,	"

Total, ... 100.00.

A number of analyses were also made of samples of air collected close to the spring, and the following mean results were obtained :—

Oxygen, 20.885%	by volume.
Nitrogen, 79.115	„ „
Carbonic Acid, 0.06	„ „
	<u>100.060</u>	

The ammonia was determined separately and amounted to 0.055 parts by weight in 1,000 of air. The air near the spring differs consequently from ordinary air slightly by containing a little more carbonic acid (0.06%, as against 0.03 to 0.04 in normal air).

GOMBAK SPRING.

Temperature, ... 50-54° Centigrade (122-129° Fahrenheit).

Total solids dissolved in the water, 21.333 grains per gallon.

Hardness, 3.6	„ „
Chlorine, 0.55	„ „
Free Ammonia, 0.04	parts per million.
Albuminoid Ammonia, 0.05	„ „ „

Total constituents dissolved in 10,000 parts of the water, 3.9942, viz. :—

Calcium Carbonate, 0.26	parts in 10,000.
Sodium „ 0.22	„ „
Ammonium „ 0.0012	„ „
Calcium Sulphate, 0.285	„ „
Magnesium „ 0.016	„ „
Sodium „ 0.199	„ „
Potassium „ 0.093	„ „
Sodium Chloride, 0.065	„ „
Potassium „ 0.012	„ „
Lithium „	
Sodium Sulphide, 0.018	„ „

Hydrogen Sulphide,	0.030	parts in 10,000.
Boric Acid,		
Carbonic Acid,	0.81	,,
Nitrogen ,,	0.08	,,
Organic Matter,	0.09	,,
Silica,	1.765	,,
		<hr/>	
Total, ...		3.9942	,,

Composition of Gases escaping from Spring.

Nitrogen,	98.1%	by volume
Carbonic Acid,	1.9%	,,
Hydrogen Sulphide,	0.3%	,,
		<hr/>	
		100.3	

Composition of Air close to Spring.

Oxygen,	20.90%	by volume.
Nitrogen,	79.10%	,,
Carbonic Acid,	0.06%	,,
		<hr/>	
		100.06%	

SEMUNIAH SPRING.

Temperature,	45-50°C. (113-122° F.).
Hardness,	3.4 grains per gallon.
Chlorine,	0.5 ,,
Total solids dissolved in the water,	20.5 ,,
Free Ammonia,	0.04 parts per million.
Albuminoid Ammonia,	0.05 ,,

Constituents contained in 10,000 parts of the water:—

Calcium Carbonate,	0.21	parts in 10,000.
Sodium ,,	0.1900	,,
Ammonium ,,	0.0015	,,

DUSUN TUA SPRING.

Temperature, ...	45-50° C. (122° F.).
Total solids dissolved in water, 15.4 grains per gallon.	
Hardness,	2.3 " "
Chlorine,	0.25 " "
Free Ammonia,	0.04 parts per million.
Albuminoid Ammonia,	0.03 " "

Constituents contained in 10,000 parts of the water:—

Calcium Carbonate,	0.1357 parts in 10,000.
Sodium "	0.08000 " "
Ammonium "	0.00053 " "
Calcium Sulphate,	0.11400 " "
Sodium "	0.09900 " "
Magnesium "	0.01200 " "
Potassium "	0.06300 " "
Sodium Chloride,	0.05100 " "
Potassium "	0.00400 " "
Lithium "	trace " "
Sodium Sulphide,	0.00900 " "
Hydrogen "	0.03200 " "
Boric Acid,	trace " "
Carbonic "	0.92000 " "
Nitrogen "	0.07000 " "
Organic Matter,	0.04031 " "
Silica,	0.61200 " "
Total,	2.20141 " "

Gases escaping from Spring.

Nitrogen,	96.5% by volume.
Carbonic Acid,	2.9 " "
Hydrogen Sulphide,	0.4 " "
Hydrogen and Marsh Gas,	traces.

Composition of Air close to Spring.

Nitrogen,	79.04
Oxygen,	20.91
Carbonic Acid,	0.05
Total,	100.00

ULU SELANGOR SPRING.

Temperature,	...	100.2% F. (38 C.).
Hardness,	...	3.5 grains per gallon.
Total solids dissolved in the water,	19.5	" "
Chlorine,	...	0.5 " "
Free Ammonia,	...	0.16 parts per million.
Albuminoid Ammonia,	...	0.12 " "

Constituents dissolved in 10,000 parts of the water:—

Calcium Carbonate,	...	0.2100 parts in 10,000
Sodium "	...	0.1800 " "
Ammonium "	...	0.0025 " "
Calcium Sulphate,	...	0.1900 " "
Magnesium "	...	0.0180 " "
Sodium "	...	0.1850 " "
Potassium "	...	0.0600 " "
Sodium Chloride,	...	0.0700 " "
Potassium "	...	0.0090 " "
Lithium "	...	trace " "
Potassium Iodide,	...	trace " "
Sodium Sulphide,	...	0.0090.
Hydrogen* "	...	not determined.
Boric Acid,	...	trace.
Carbonic Acid,*	...	not determined.
Nitrogen,*	...	" "
Organic Matter,	...	0.1100 parts in 10,000
Silica,	...	1.2000 " "

Total, ... 1.2135 parts in 10,000

Composition of Gases from Ulu Selangor Spring and surrounding Air.

Not determined, as I did not visit the spring in person.

* These constituents were not determined, as I did not personally visit this spring, and they can only be correctly estimated in freshly collected water. Judging by analogy they would differ but slightly from the amounts under "Gombah."

ULU KLANG SPRING.

Temperature,	83-84° C. (181-183° F.).
Hardness,	2.0 grains per gallon.
Chlorine,	0.45 " "
Free Ammonia,	0.03 parts per million.
Albuminoid Ammonia,	0.04 " "
Total solids dissolved in the water,	21.95 grains per gallon.

Constituents dissolved in 10,000 parts of the water:—

Calcium Carbonate,	0.1600 parts in 10,000
Sodium	"	...	0.1950 " "
Ammonium	"	...	0.0050 " "
Calcium Sulphate,	0.1250 " "
Magnesium	"	...	0.0201 " "
Sodium	"	...	0.2200 " "
Potassium	"	...	0.0955 " "
Sodium Chloride,	0.0500 " "
Potassium	"	...	0.0150 " "
Lithium	"	...	trace " "
Sodium Sulphide,	0.0200 " "
Hydrogen	"	...	0.0350 " "
Boric Acid,	" "
Carbonic Acid,	0.8050 " "
Nitrogen,	0.0700 " "
Potassium Iodide,	trace " "
Organic Matter,	0.0950 " "
Silica,	1.5000 " "
Total,...			3.4606 " "

Composition of Gases escaping from Spring.

Nitrogen,	95.85
Carbonic Acid,	3.50
Hydrogen Sulphide,	0.65
Hydrogen and Marsh Gas,	trace

100.00

Composition of Air close to Spring.

Oxygen,	20.870
Nitrogen,	79.060
Carbonic Acid,	0.070
Hydrogen Sulphide,	trace
				100.00

(Ozone or active Oxygen absent).

Ammonia in air near spring, 0.11 parts in 1,000 parts by weight.

A sample of air was collected about 1½ miles from the spring, and the ozone in it determined.

The mean result of the separate experiments was 0.000,0011 milligrams per litre of air.

Composition of Rocks from the Spring.

It has already been stated that the formations in the immediate vicinity of the springs are granitic, the principal component minerals being accordingly:—Quartz, Felspar, Mica (Muscovite and Zinwaldite) and Hornblende.

The average composition of the rock (from 6 analysis) is shown below:—


Silica,	74.30%
Alumina,	14.20,,
Alkalis,	5.10,,
Lime,	3.00,,
Magnesia,	0.60,,
Iron (taken as Ferric Oxide),	1.10,,
Moisture, Organic Matter, Manganese, &c.,	1.70,,
				100.00%

Tin ore in varying though small quantities is found disseminated through the mass of some of the specimens. Gold was absent in all the samples I took, with the exception of a piece

of Dicrite found at Ulu Klang, which contained a quantity corresponding to $\frac{3}{4}$ oz. to the ton. I do not for the present attach any importance to this circumstance, as it may be a purely sporadic occurrence; yet it might be worth while to obtain a proper sample taken from a large quantity of the rock near Ulu Klang and have it tested for gold. I am not aware whether or not gold has ever been found or been supposed to exist at Ulu Klang.

SPRINGS IN MALACCA.

The springs found in Malacca in all respects closely resemble the Selangor Springs (*see* under General Nature of the Springs). One spring at Cherana Puteh is situated in the jungle, those at Ayer Panas and Alor Gajah occur in padi swamps. The latter two have been properly set for bathing purpose, so I can say nothing about their original condition and appearance. The spring at Cherana Puteh appears as a pool or puddle of hot water in a bed of dark grey mud from which bubbles of gas are continually ascending. At the time of my visit a little brook was flowing close to the spring and mixing with it to a certain extent. No line of demarcation between the brook and the water from the spring was visible, but could be readily detected with a thermometer; the differences observed are worthy of note (*see* sketch below).



It will be noticed that within small distances of a foot or even 6 inches the temperature varies very considerably. The points marked \times indicate places where the temperature was taken, those marked \oplus mark the entrance of feeders or "eyes" of the spring, and naturally possess the highest temperature.

SPRING AT AYER PANAS.

The temperature of this water varies considerably in the different tanks or wells in which the spring has been collected.

a.—Tanks inside house.

- No. 1 = 45° Centigrade (113° Fahrenheit).
- „ 2 = 35° „ (95° „).
- „ 3 = 45° „ (113° „).
- „ 4 = 52° „ (123° „).

b.—Tank outside house on ground immediately adjoining.

- No. 5 = 55° Centigrade (131° Fahrenheit).

c.—Tank in padi swamp behind house

- No. 6 = 52° Centigrade (125° Fahrenheit).

d.—Tanks in front of house in the field.

- (Left) No. 7 = 33° Centigrade (91° Fahrenheit).
- (Right) No. 8 = 48° „ (118° „).

Analysis of water taken from hottest tank, No. 5 :—

Total solids dissolved in the water,	18.40 grains per gallon.
Hardness, 2.5 „ „
Chlorine, 0.7 „ „
Free Ammonia, 0.05 parts per million.
Albuminoid Ammonia, 0.05 „ „

Constituents dissolved in 10,000 parts of the water :—

Calcium Carbonate, 0.160 parts in 10,000
Calcium Sulphate, 0.180 „ „
Magnesium „ 0.015 „ „
Sodium „ 0.190 „ „
Potassium „ 0.085 „ „
Sodium Carbonate, 0.450 „ „
Ammonium „ 0.00018,, „

Sodium Chloride	0.095	parts in 10,000
Potassium "	0.006	" "
Sodium Sulphide,	0.020	" "
Hydrogen "	0.025	" "
Carbonic Acid,	0.585	" "
Nitrogen,	0.085	" "
Silica,	0.780	" "
Organic Matter,	0.250	" "
Total,	2.92618	" "

SPRING NEAR ALOR GAJAH.

Part of this spring has been collected in a tank inside a little house, part of it rises in the bed of a little brook close to this house, and a third outlet has been set and received in a square bricked trough or well. The latter evidently (from its temperature, etc.) corresponds to the main feeder.

Temperature of water inside house,	35° C.	(95° F.).
Temperature of water in the brook,	40° "	(104° F.).
Temperature of water in bricked well,	55° "	(131° F.).
Hardness,	...	2.5
Total solids dissolved in the water,	...	1.7 grains per gallon.
Chlorine,	...	0.5 " "
Free Ammonia,	...	0.04 parts per million.
Albuminoid Ammonia,	...	0.05 " "

Constituents contained in 10,000 parts of the water:—

Calcium Carbonate,	0.200	parts in 10,000.
Calcium Sulphate,	0.140	" "
Magnesium "	0.025	" "
Sodium "	0.150	" "
Potassium "	0.095	" "
Sodium Carbonate,	0.550	" "
Ammonium "	0.00015	" "
Sodium Chloride,	0.075	" "
Potassium "	0.005	" "
Lithium "	trace	" "

Sodium Sulphide,	0.019	parts in 10,000.
Hydrogen „	0.020	„ „
Carbonic Acid,	0.480	„ „
Nitrogen,	0.075	„ „
Silica,	0.590	„ „
Organic Matter,	0.295	„ „
			<u>2.71915</u>	„ „

SPRING AT CHERANA PUTEH.

Situated in a piece of jungle with bed of soft mud mixed with particles of quartz and granite. A little brook of cold water flows close to the spring and partly mixes with it.

Maximum temperature, 55° C. (131° F.).

Analysis of a sample of the water after simple filtration to remove suspended matter:—

Total solids dissolved in water, 22.5 grains per gallon.

Hardness,	4	
Chloride,	0.7	„ „
Free Ammonia,	0.04	parts per million.
Albuminoid Ammonia,	0.06	„ „

Constituents dissolved in 10,000 parts of the water:—

Calcium Carbonate,	0.250	parts in 10,000.
Calcium Sulphate,	0.270	„ „
Magnesium „	0.015	„ „
Sodium „	0.185	„ „
Potassium,	0.095	„ „
Sodium Carbonate,	0.320	„ „
Ammonium Carbonate,	0.000180	„ „
Sodium Chloride,	0.100	„ „
Potassium „	0.090	„ „
Lithium, „	trace	„ „
Boric Acid,	„ „
Sodium Sulphide,	0.025	„ „

Iron, Alumina and Manganese,	0.125	parts in 10,000.
Hydrogen Sulphide,	0.019	” ”
Carbonic Acid,	0.450	” ”
Nitrogen,	0.089	” ”
Silica,	0.595	” ”
Organic Matter,	0.195	” ”
	<hr/>	
	2.82318	parts in 10,000

Composition of Gases escaping from the Springs.

A number of analyses were made of the gases which are given off by these hot springs, and they were found to consist principally of Nitrogen and Carbonic Acid with small quantities of Hydrogen Sulphide and Marsh Gas.

THERAPEUTIC VALUE OF THE SPRINGS.

In my remarks upon the general nature of the springs, I have already pointed out that they must be classed with simple thermal waters, as they contain but an insignificant amount of solid matter excepting Silica which has but little physiological action. All the springs, however, contain Hydrogen Sulphide, and although the quantity present is sufficient to warrant their being classed with “Sulphurous Waters” in the strict sense of the word, yet it suffices to impart to them a distinct therapeutic value. In addition to this, their temperature is an important item, considered medicinally. There are a number of well-known springs in Europe, which owe their virtues apparently to their temperature only and contain as little or less mineral matter than the Selangor springs and less Hydrogen Sulphide or none at all (Matlock; Buxton; Wildbad; Aix en Provence).

Among the Selangor springs Ulu Klang ranks first as possessing the highest temperature and containing most Hydrogen Sulphide. The remainder may be placed in order of merit thus:—Setapak; Dusun Tua; Semunieh; Gombah. Ulu Selangor would probably be the last in the list, although I cannot say so definitely, as I did not personally inspect and examine this spring. In any case it would come after Setapak

On reference to the data of analysis of the Ulu Klang and Setapak waters, it will be seen that they contain the following amounts of Hydrogen Sulphide dissolved in every 10,000 parts of the water :—

Ulu Klang,	0.035 (Temp. 151° F.).
Setapak,	0.034 („ 122° F.).

Now the average amount of Hydrogen Sulphide in *cold* sulphurous springs may be taken as 0.090 parts in 10,000; hence the Ulu Klang and Setapak contain a very fair fraction of this amount and possess a high temperature at the same time. Both of them may, therefore, be recommended for bathing or drinking purposes. Their action would be stimulating and diaphoretic, and they would be found valuable in :— Chronic Skin Diseases and Rheumatism; Chlorosis; Amenorrhœa; Secondary Syphilis; Dyspepsia due to disordered action of the liver. I do not think it practicable to bottle and sell the water, as this would result in the loss of almost all its volatile and valuable constituents by evaporation and subsequent decomposition. The water should be used on the spot and as nearly at the temperature of the spring as may be found endurable.

The suggestions arising from the above are :—

I.—That the hot springs in Selangor, more particularly those at Ulu Klang and Setapak, are by virtue of their temperature and chemical composition therapeutically valuable.

II.—The springs at Ulu Klang and Setapak should be properly set and made available for bathing and drinking purposes. It is almost needless to say that every care should be taken to prevent any communication between the water used for bathing and drinking purposes respectively.

It should be mentioned here that the Malays and Chinese seem to be well aware of the virtues of these springs, and particularly of their specific action in skin diseases. They bathe in them freely, and I am informed that the Chinese more or less monopolise some springs, the Malays others. As far as I could learn they never drink the water.

Ultimate Origin of the Springs.

The geology of the springs is a subject of great interest, but unfortunately one not to be approached or settled in an easy, off-hand manner. Much patient study of actual sections in the immediate vicinity of the springs, examination of deeper strata by boring, in short a complete geological survey alone, can throw light upon this question. One is accustomed to associate hot springs with volcanic action, yet these springs occur in parts far distant from any known volcanic belt.

No basalt and no minerals indicative of direct volcanic action are to be found in the neighbourhood. The composition of the water is, in some respects, similar to the water ejected by the geysers of Iceland, but here the analogy ends. Beyond conjectures pure and simple, I can at present pronounce no opinion as to the final source of these springs, viz., the question of direct volcanic action or simple intra-terrestrial heat. I sincerely hope, however, to have an opportunity to follow up this subject later on.

In conclusion, I wish to refer to a paper by Mons. STANISLAS MEUNIER published in J. cx, p. 1085 of the *Compt. Rend.* 'Examen chimique d'eaux minérales provenant de Malairé. Mineral d'étain de formation actuelle.' The paper describes the analysis of two bottles of water brought by Mons. J. ERRINGTON DE LA CROIX from Ayer Panas and Cheras, and contains the following statements among others:—

- 1.—The water had a bad smell on opening bottle.
- 2.—On heating it gave off gas bubbles.
- 3.—On evaporation it left a syrupy residue which was incinerated and gave two milligrams of ash per 100 cc. water taken.
- 4.—Exposed to the air for a few hours the water fills itself with white, viscous vegetations not botanically determined.
- 5.—The Ayer Panas water had no taste; the Cheras water no smell, but a slightly saline taste, and on heating gave off myriads of gas, probably pure Nitrogen amounting to 10 cc. per litre.

6.—The water contained Chlorine, but no Sulphates, and about 1.4 grams Sodium Chloride per litre.

7.—A mineral found in the spring is described, having a spec. gravity 2.1, cavernous and tuberculous structure, whitish grey colour with small black dendritic particles in it. The composition of this is given as :—

Silica,	91.8 %
Water,	7.5 "
Tin Oxide,	0.5 "
Iron,	0.2 "
Aluminium,	traces
			100.0 "

From this the author draws the remarkable conclusion that this substance is a kind of opal similar to geyselite, but containing tin as a peculiar and characteristic constituent.

He goes on to say that this is the first time that tin has been found in *statu nascenti* as it were, viz., in the act of deposition from its mother liquor, and looks upon this as an important contribution to the theory of the formation of tin ore.

It needs but a glance at the statements above cited to show that they are of no value at all, in fact mostly absurd, but I append a few remarks upon these points, lest a casual reader of the paper should be deceived by them and attach credit to the grand but unfortunately erroneous deduction (No. 7) in which that paper culminates. It seems strange that matter so devoid of point and substance should have found its way into the Compt. Rend. :—

1. *Bad Smell.*—Most waters would have a bad smell after being carried from Selangor to France.
2. *Bubbles on heating.*—All water, unless specially treated, gives off bubbles on heating.
3. *Syrupy Residue on evaporation.*—The water leaves a *solid* residue. The ash is more than 4 milligrams per 100 cc.
4. *White, viscous vegetations.*—These are probably the

siliceous deposits mentioned in my paper. They do not form in a few hours as stated, and the water does not *fill itself* with them.

6. *No Sulphates, but about 1.4 grams Sodium Chloride per litre.*—The water naturally contains Sulphates, and the Chloride amounts to 0.015 grams at most.
7. The description given is that of common sinter met with in all hot springs, almost without exception. As for the tin in it, it would be difficult in these parts to pick up a mineral which did *not* contain a trace of tin, as is well enough known. There is absolutely nothing remarkable in the presence of tin in this sinter, the very granite upon which it grows carries traces of tin. The bold theory of *tin oxide* being deposited from water containing *Sulphuretted Hydrogen* is not compatible with chemical laws, is a chemical impossibility.

I desire to add that I have appended the above remarks purely by way of correcting erroneous statements, and nothing is further from my thoughts than "sitting on" the author of them. It must be borne in mind that he could only work with a small sample sent to him in France, and which probably arrived in a more or less altered condition. He never saw the springs nor the water fresh from them.

W. BOTT.



NOTES ON THE SIAMESE PROVINCES OF KOOWI, BANGTAPHAN, PATEEO AND CHAMPOON.

BY

ARTHUR KEITH, M. B., C. M.

Their Position and Outstanding Features.



IN that part of the Malay Peninsula lying between $10^{\circ} 20'$ and 12° N. Latitude, the backbone range of hills almost skirts the East coast and thus between the hills and the sea lies a narrow strip of country, the northernmost part forming the province of Koowi; Bangtaphan, Pateeo and Champoon following in order towards the South. This backbone range, covered evenly over with deep jungle like the plain that lies between it and the sea, averages in height some 1,600 feet above the sea level, but at Khov Pra it reaches some 3,500 feet, while it attains 4,326 feet—its highest point—in Khov Luang, standing as the boundary wall between Burma and Siam. It is broken here and there by gaps, honoured by the name of passes, the best known of which perhaps is that of Kra, 250 feet above sea level, lying at the southern extremity of the piece of country of which I write, and frequented by the Siamese living in the Pak-chan valley when visiting their relatives in Champoon; while better known in former times and lying quite at the other extremity, the pass of Koowi, some 750 feet above sea level, affords the colony of Siamese living in the valley of the smaller Tenasserim River, a rude path to their native provinces on the East Coast. Between

these two lie ill-marked crossings at Krat, Paron and Bangtaphanoi, the last being used by the Siamese living in the Lenya valley; but the truth of the matter is that the numerous bands of dacoits that infest this region cross anywhere (*vide* map). To the seaward of the main ridge, and for the greater part running parallel to it, are numerous small ranges, while scattered hills rise with their burden of forest jungle over the plain and thickly skirt the coast, often standing into the sea as promontories offering their bluff side to the waves; and one can see that the same features characterise the seabottom—the hills rising from the water as numerous jungle-covered islands.

Prominent Geological Features.

The basis of the backbone ridge is a rough-grained granite, rapidly decomposing when exposed to the atmosphere, and the streams flowing from it (as indeed is the case almost everywhere in the Peninsula) often carry *tin* and more frequently and more abundantly *titanic oxide of iron*. Lying on the granite and rising as the basis of the small ranges of hills is a semi-metamorphosed clay-slate much broken up, non-fossiliferous and probably of the Cambrian period, which forms the "country rock." To the seaward of this slate and lying on it, runs a great bed of conglomerate rock, composed of flat, water-worn slate pebbles and roughly rounded pieces of quartz welded together by a red clay containing much iron. Its stratification is much disturbed, and it sometimes rises into low hills. Where the slate and this conglomerate meet, one frequently finds great outcrops of a porphyritic rock evidently that of the Mergui series, and here also one finds large outcrops of gold-bearing quartz as in the concession of the Goldfields of Siam Company. Again it can be seen further South in Champoon where another mine has been opened out. (*Vide* Map.) In the beds of the streams running through these quartz-bearing parts, one finds gold, and sometimes for a mile or more inland from either bank alluvial gold occurs along its sides for long stretches, but from its comparatively even distribution and level deposition one has to look for

some means other than the river to account for its presence here and thus. Still proceeding seawards, and having crossed this land of conglomerate, one again comes to the "country rock"—metamorphosed clay slate. On the coast, laterite is abundant, often forming low cliffs, but the hills skirting the sea and most of the islands are of a limestone that gives forth a metallic ring when struck.

Limestone Hills and Caves.

About a mile to the West of the village of Bangtaphan, raising its somewhat flattish top some five hundred feet above the surrounding alluvial plain with its old sea beaches, is a rather typical limestone hill with a sloping side to the land, and like others of its kind, a steepish side to the sea sheltering the mouths of numerous caves of various sorts and sizes. To gain the principal cave, one ascends some fifty feet to find the mouth as sharply cut as, and somewhat after the manner of, a cathedral door, and as the cave retreats from its mouth it increases in every dimension. The floor, running backwards for some 80 feet, mounts in three great tiers some 60 feet wide, the last and highest tier being sacred to a gilt Bhudda with a perfect myriad of leaden apostles round about him, for the cave is used as an occasional place of worship, as these caves commonly are in Siam. The roof rises in vaults, the highest of all rising to some forty feet and is pierced by what the Scottish people familiarly know as a "Hell's Lum." From the shape of the caves, from those "Hell's Lums," and from their situation on the steep side facing the sea, one naturally concludes they are the result of sea-action. One can see lines, more or less parallel to the surface slope of the hill running along the walls of those caves, as if they demarcated concentric layers which had been deposited by some spring that rose in the centre of the hill, overflowed, and laved its sides, and the not uncommon nearly concave top lends some support to such a guess. It seems highly probable that out of the solidified debris in the floors of those caves, animal remains, recent or otherwise, might be obtained, although a search made in those of Borneo (see the Society's Journal for

1879)—was practically negative, but the religious purposes to which these caves are put deters one from actively pursuing operations.

These limestone hills and caves seem fairly well distributed along either coast of the Peninsula—seldom I believe rising many hundred feet above sea level. On the West coast away as far North as Moulmein, one finds similar hills with famous caves—also used as temples; in the Lenya valley again there are said to be some remarkable specimens, and there are others in Selangor, Kedah and Pahang, besides in many of the islands lying along that coast. In the four Siamese provinces they are common, and at Pateeo a group of these limestone hills rises from the sea like a great set of decapitated sugar cones, while away much further North beyond these provinces, at Petcheburee, there is a famous cave containing a gigantic wooden image of Bhudda. Among the islands and shores along the Eastern side of the Gulf of Siam one comes across them, while they abound in North Borneo.

The Rising of the Land.

Assuming then that these caves are the result of sea-action, as in all probability they are, and seeing that some of them are now high and dry some hundred feet or more, we may conclude that the land has at any rate risen that amount in recent times. A writer in this Journal for 1879, discussing those limestone hills and caves in North Borneo, concludes that that part of the island had risen about 500 feet in recent times, and before I saw that article I had concluded that at any rate the land in these four provinces lying three hundred feet above the sea level had emerged from the water in a recent era. MASON, in his work on Burma, gives it as his belief that the Burmese coast is rising, and states that the land on the other side of the Bay of Bengal is sinking. Probably the whole great tract of country—almost the whole of the Indo-China Peninsula—over which this limestone formation occurs is rising; at any rate that small part of which I write, I believe to be in course of elevation. In the great flat plain,

sloping from the hills to the sea, one finds extensive beds of clay almost of one level and sloping towards the sea, and over this again great beds of evidently sea-rolled gravel also comparatively at one level and of an even thickness, lying some 450 feet above the present sea level. The hills have their sloping sides to the mainland, and their steep sides—often sea-marked cliffs—to the sea. Then again (see vertical section across Province of Bangtaphan at lower right-hand corner of map) going towards and within two miles of the sea, one crosses undulating and wavy old sea beaches of sand following each other in rapid succession. Lem Tong Lan (*vide* section) is a hill standing out to the sea with the usual characteristic shape and joined to the land by a muddy isthmus, over which old inhabitants say it used to be possible to sail at high water with a boat, which is now impossible. The section running through Koh Yeu shows that island with its sloping side in very shallow water towards the land, while that towards the surf is steep and faces deep water. The shape of those islands and hills, this little tongue of land joining Lem Tong Lan to the mainland, these old sea beaches, the limestone caves, and the other recent geological formations even in the absence of recent marine remains, seem to point to the land having risen and still to be rising, but of course in this extension seawards of the land deposition has played a considerable part.

The River System and the Effects of Deposition.

In this narrow strip of country the rivulets from the main range and subsidiary hills meet on the plain below to form considerable streams which, running over beds of sand and gravel, make on the whole a straight course across the plain to the sea. One of those streams has seldom more than 150 square miles of a drainage area, but the river of Champoon, like the large streams draining the other side, and like the rivers of the Peninsula in general, runs parallel to the main range of hills for the greater part of its course until, near its termination, it turns outwards to end in the sea, and thus drains an area of about 450 square miles—three times that of

any of the other streams. On the West side, on the other hand, with an area almost three times as great to be drained, the place of those babbling small streams is taken by comparatively large rivers; the Pak Chan has a drainage area of 600 square miles, the Linya 800 square miles, and the smaller Tenasserim River has over 2,000 square miles of a drainage area. These rivers on the West side, with their large drainage areas, receiving a rainfall fully double that of those smaller ones on the East coast, and carrying an infinitely larger volume of water with its inherent properties of disintegration, denudation, transportation and deposition, represent powerful factors at work on the Burmese side almost absent in these four Siamese Provinces. Sandy beaches form the seaboard of these, with sand bars at the mouths of the streams, and with patches of mangrove in the narrow marshes and pools between the more recent of the old sea beaches. The Champoon River, however, partaking of the rainfall as well as of the character of these rivers on the West side, is surrounded at the mouth by great mud-banks, large mangrove swamps and tracts of rich alluvial soil. On the West coast these evidences of deposition are extremely extensive, for the islands lend a calm to the water round the mouths of the rivers favourable to the settling of suspended matter, and the mangrove trees stepping further and further into the water as subsidence goes on highly favour further deposition among the roots; so that deposition and extension of the mangrove swamps proceed *pari passu*. Knowing somewhat of the drainage area of a river one may roughly guess, from the extent of mangrove swamp surrounding its mouth, the rainfall of the district.

Lately, at the mouth of the Krat River, on the eastern shore of the Gulf of Siam, I came across some remarkable evidence of the rate of deposition. The Krat River, rising at the western side of the Battambang Hills, drains an area of some 250 square miles, with a rainfall of probably 200 inches per annum, and runs through a clay country, a considerable extent of which is cultivated, so that it carries a great deal of matter in suspension. In 1859 H. M. *Saracen* laid down the

islands at the mouth of this river showing wide channels between them and the mainland. Now those islands are almost in contact with the mainland; roughly guessing I should say some 500 acres have come above high-water mark since then.

Meteorological.

The main range of hills, although only 1,600 feet in height, determines the rainy season on the West side with the S.W. Monsoon, while in these four provinces on the East side with the N.E. Monsoon. But with the exception of some heavy downfalls in November and December, there is little of a rainy season, rain falling more or less all the year through.

Rainfall at Paron, Bangtaphan, for 1890.

Jan.	Feb.	Mar.	April.	May.	June.	July.	Aug.	Sept.	Oct.	Nov.	Dec.
1.5	3.1	1.5	5.7	4.5	1.2	3.5	4.6	3.4	8.2	5.2	20.6

The total rainfall for 1890 at Bangtaphan was 63 inches, but at Mergui, on the other side of the Peninsula, it was over 200 inches. The change of the Monsoon always came gradually. Towards the end of April the wind veered towards the South and kept changing, so that May was half spent before it finally settled in the S.W. So again in October; for several days it may have blown from the N.E. and again returned to the S.W., so that November had well begun before we enjoyed those steady cold breezes from the N.E. that gave a piquancy to life.

In May, the hottest month of the year, the thermometer in the shade registered on an average 84.5° at 9 a.m. and 91° at midday, while at night it never sunk below 75°. But in December, I have seen the minimum thermometer register outside 59°, and in the shade 65° at night.

The Jungle Covering.

Jungle forest sweeps evenly over the Provinces of Koowi and Bangtaphan, but is interrupted in Pateeo by a large tract of grass country—of most interesting origin—but occurs again in Champoon although of a much less primitive nature than that in the two northern provinces. Rather stunted trees, surrounded by and supporting numerous twining and climbing shrubs, cover the hills with a thick, almost impenetrable undergrowth. In the valleys, near the backbone ridge, on the other hand, grow lofty trees affording valuable timber, shading a damp green undergrowth of sapling-like shrubs. In the plain, at the foot of the smaller hills, are large tracts of bamboo forest, while the jungle extending over the plain is much the same as that occurring in the valleys. Various species of figs are extremely common, and fine gum-dammar trees occurs all over these provinces and are farmed out for their oil. The oil exudes into small wells dug out of the trunk some four feet from the ground, and the exudation is at times stimulated by setting fire to the oil in the well. It is mostly used for making torches; hast is dipped in the oil, wrapt into a banana leaf, the whole tied up with pieces of rattan constituting a torch—one of the principal articles of export.

In the month of December, the leaves begin to fall, so that by the beginning of March, just before many of the trees burst into bloom, large tracts of the drier jungle are comparatively leafless. In many of these trees this shedding of the leaf is likely due to some physiological adaptation; in others that bloom while in this leafless condition, it may be to increase the display and attractiveness or facilitate the means for fertilization; and in some possibly it may be a bequest from some distant ancestor that grew in a temperate climate.

Competition for life is keen in the jungle, and although I have no statistics to offer, I believe there is a high rate of mortality amongst the jungle trees. Tottering trees infested with white ants abound, the fig trees are everywhere making victims, and after a slight gale many trunks lie prone with tons of soil upturned with their roots. When the trunk has

rotted away, this soil remains as if it were a broken piece of an earth-dyke by the side of a shallow trench—very puzzling until one knows their origin. In time the rain blots them out, but from their abundance one can see that this is one of nature's methods of ploughing the jungle ground and must play not an unimportant part among the more important factors in altering the face of the land.

The sappan-wood tree is abundant, so are many species of rattan, but these with the dammar oil are the only jungle products exported from those provinces.

The Grass Country of Pateeo.

The jungle suddenly ceases at Bangtaphannoi, and from there southwards to beyond the village of Pateeo where the jungle again appears in patches, stretches an undulating country waving with lalang grass some sixty square miles in extent. Some twenty-seven years ago, so the people say—people that saw what they tell—a great typhoon crossed the Peninsula here, levelling nearly every tree as it came and included villages in the ruin, so that not a few human lives were lost. A fire following completed the work and left an open, blackened country that speedily became covered with grass that took the place of the former thick jungle forest. Here and there still stand charred stumps, while heaps of ashes covered with grass and half buried fallen boles of trees through which one's foot sinks when walking, abound everywhere. Towards the confines of this open country, the hills have their slopes facing the S.W., denuded of jungle and grass-covered, while their slopes towards the N.E. still retain their thick covering of trees. Every year the grass is set on fire and burns its border line a little further into the jungle, so that steadily it increases. The edge of the jungle, like the border of an unhealthy wound, shows no robustness nor vitality, and falls an easy pray to the all-devouring annual flames.

The result of the action of these warring elements has been to totally alter the climatic conditions, the fauna and the flora of this small locality. A few species of grass have supplanted the numerous forms of jungle growth; the small

barking deer and the black leopard abound on it, with a few wild buffaloes. But there is not a human habitation upon it, notwithstanding the abundant pasture, for the Siamese keep cattle only as a means of transport and labour.

Old Paddy Fields.

With the exception of the grass country of Pateeo all the other open spaces in the jungle, which although numerous seldom exceed forty acres in extent, are old paddy-fields cleared at one time for cultivation by man. When a piece of jungle has been cleared for hill-paddy and the crop reaped and carried away, the old tree stumps sprout, young saplings spring up, and the jungle soon regains its own; but in the low-lying level alluvial ground prepared for 'wet' paddy, and which is usually put under a course of crops, thick grass springs up, and being set on fire every dry season drives the jungle quite as far back as it is able to regain during the rains, so that it becomes a permanent opening in the jungle.

The Rice Crop.

Although small patches of tobacco are grown, and one may see cotton trees, plantains, coco-nut and betel-nut palms in gardens surrounding the houses, yet rice is by far the chief crop of those four provinces. The rice-fields lie upon the flat, alluvial soil surrounding the villages, which are invariably situated upon the banks of a stream towards its mouth. In the end of July, low earthen walls are thrown up, dividing this land into plots containing about one square rood, and the soil turned over by means of a rude wooden plough. As soon as rain falls buffaloes are turned out to trample and soften the soil and to further prepare it for sowing, they are yoked to a log of wood set with wooden teeth and the ground thus harrowed. A small seed bed is prepared, and rice sown extremely thickly, and when the crowded plants have got their heads some six inches above the ground they are transplanted and set one by one some eight inches apart in the plots already prepared by ploughing and harrowing. By the end of November, the crop is ripe, the heads are cut and gathered by

the sickle, and the grain trampled out on the threshing-floor by buffaloes. The paddy is stored away in small raised bamboo houses specially made for the purpose.

An extremely small proportion of the land is under cultivation—less than 2%. Those four provinces include 2,200 square miles of the King of Siam's kingdom, and out of these 2,200 square miles about $3\frac{3}{4}$ square miles only are under cultivation. An average crop produces about 1,000 lbs. of clean rice, so that in an average year the total production in these provinces would be about 2,300,000 lbs of rice. This has to feed a population of some 16,500 souls—rather less than one half pound per head per diem—for as far as I know, there is neither export nor import in this commodity. A man and his wife could with ease cultivate four acres of rice and produce 4,000 lbs of rice, but nearly everyone grows his own supply only, and very few cultivate that amount.

From passing through those provinces and residing in them, I believe their size, acreage under cultivation, and population are approximately what I here tabulate:—

<i>Province.</i>	<i>Total square mileage.</i>	<i>Acres cultivated.</i>	<i>Population.</i>
Champooon, ...	740	800	6,000
Pateeo, ...	450	500	3,000
Bangtaphan, ...	480	300	2,500
Koowi, ...	560	700	5,000

Fishing Industry.

In the dietary of the Siamese, fish occupies almost as important a place as rice, and although every Siamese is partly an agriculturist, yet there are many that devote their lives entirely to either calling. All the villages have easy access

to the sea; along the shore are many small collections of fishermen's houses, and here and there are Chinese *kongsis*, from all of which at the break of day boats issue to prosecute the fishing close inshore. They use nets of every shape and size almost entirely, and return about 10 o'clock, commonly with good catches, which they dispose of at something like three cents per lb., and what is not bought for immediate consumption is salted and dried in the sun. The Chinamen living in those *kongsis* on the beach are engaged in catching prawns, which they do by dredging a very fine meshed net along the sands, for the production of "blachang." At night, too, one can see torches flitting about on the water, the fisherman spearing the fish that are attracted to the light. In the streams one comes across dams in which are set all sorts of ingenious bamboo traps for fish.

Means of Communication.

The sea is the highway between these provincial towns and Bangkok the capital. Some twenty years ago Chinese junks and large boats of Siamese build crowded the Menam at Bangkok bringing in the produce from the coasts all round the gulf and carrying back products of civilization in return, but now, although two or three are always riding at anchor in the river at every village and town, they have been almost completely supplanted by small steamers flying at their stern the white elephant, having long since passed that condition of evidence required to carry the Union Jack.

On land, the towns and villages lying along the coast are connected by a path often winding along the beach, always rough and uneven, wending its way across morasses and soft miry paddy-fields, full of holes, crossing streams with neither bridge nor ford, so that it is laborious travelling indeed, and in the three southern Provinces fit for elephants only. Elephants abound in these Provinces and are used for transport, but in Koowi, where an elephant cannot be found, the roads are much better, being well adapted for buffalo drays, which are the sole means of transport.

Four years ago a telegraph line from Bangkok was carried through these provinces, and a station was opened at Bangtaphan, but so frequently was the wire broken and instruments out of order, that it was seldom of any practical use. Already there is scarcely a sound pole, and certainly not a perpendicular one, and in many places the wire has forsaken the insulators and takes its support directly from the ground.

Location of Towns and Villages.

Dacoity, or rather midnight robbery, is so rife in the district that the people, afraid to live in isolated and remote houses, congregate in towns and villages. These, surrounded by their paddy-fields, are situated upon the banks of a stream just so far from the mouth as a junk or big Siamese boat can reach at high water. In a small stream as that of Krat, the village is but a quarter of a mile from the sea; Bangtaphan, which stands upon a much larger stream, is two miles up the river, while Champoon, upon a much larger stream still, is nearly ten miles from the sea. The town occupying such a situation has the advantages of an ample supply of fresh water except at full tide, easy access to the sea for trade and fishing, and still in the midst of its paddy-fields. At the mouth of the river is commonly a small fishing hamlet known as the *Paknam*, but which as in Taiyang, at the mouth of the Champoon River, may wax bigger than the principal town situated higher up the river. The village of Paron upon the concession of "The Gold Fields of Siam Company," is a product of the mining industry there and is the only exception I know of in these provinces to the general situation upon the flat alluvial soil near the mouth of a river.

Composition of a Village or Town.

These villages or towns resemble each other as much as peas do; they agree in being an irregular row of bamboo houses covered with attaps, raised upon posts some five feet above the ground and usually about one hundred of them huddled upon one or both banks of the stream. Gardens surround the houses usually containing chilies, papaya trees,

cotton plants, and other vegetables, while round about are numerous groves of coco-nut and betel-net palms, plantains and jack fruit trees. A house rather larger than the others, but quite as dilapidated, proclaims the residence of the Governor, and an open shed near his Excellency's house serves as a court of justice by day and a *sala* or rest-house by night. At a little distance from the village, commonly in the healthiest and prettiest situation, always shaded by trees, stands the *wat* or pagoda, the centre of education and religion and which shelters a yellow-robed priest to every forty inhabitants in the town. Five or six houses of rather large dimensions, but, unlike all the other houses, not raised on posts, with numerous pigs feeding in front of them and with the doors pasted over with red posters, are the shops of the village, invariably owned and conducted by the Chinese. Here are sold dyes and calico prints, Manchester and Birmingham goods of very inferior quality, while buffalo hides and horns, dried fish, coco-nuts and betel-nuts are bought or exchanged. The opium farm, the gambling farm, and the spirit farm are always in the hands of the Chinese, and while waiting sometimes in the "grog-shop" I have been surprised to see men and women come straggling in, tendering their two cents and tossing off their *arrack* much as one sees in England.

The Inhabitants.

Settling down at one of those villages, and taking a random hundred of the inhabitants, one would find them something like this:—6 Chinamen wedded to 6 Siamese women and having 13 children between them; 16 would be found Siamese-Chinese of a former generation. Of pure Siamese 10 men have wedded 10 Siamese women and 18 children have been the outcome, while 11 would be unmarried Siamese—male and female. Forty-one per cent. is a low estimate of the proportion of Chinese and Chinese descendants that still wear a queue, for in some towns such as Taiyong they constitute more than 70% of the population. Into the numerous villages lying along the 3,000 miles of coast between

Rangoon and Tonquin, Chinamen have been continually dropping for centuries, mixing and marrying with the natives, so that the wonder is not that they have affected the race along the shore, but that they have not affected it more. The children of a Chinese-Siamese marriage—unlike the weaklings produced by such a wide divergence as the Siamese and European—are robust, of larger stature, and more muscular than the native, comparatively dark in complexion, and wanting the business smartness and mental activity of the father, as well as lacking that peculiar mental subtlety commonly found in the Siamese. Very curious to state, while the female offspring of a Chinese-Burmese marriage is as a rule of very prepossessing appearance, yet her Siamese cousin is invariably very ugly. The Chinese-Siamese is often a man of great ability, as can be seen by the large numbers that have raised themselves to affluence by trade, and in the last century, when the Burmese had overrun Siam and were grinding her down, one PHAYA FAK (a Chinese-Siamese) arose, rallied his countrymen round him, led them from victory to victory, until Siam was again free. Those that know the dallying nature of the Siamese must recognize the Herculean task this man performed.

Each of these provinces has its Governor appointed by the King, that of Champoon occupying the highest rank, but without exception they are greedy, ignorant, narrow-minded men, who hate the foreigner, his trade and his habits. They form a marked contrast to the discreet Chinamen that preside over the destinies of the provinces lying South of these, who in every way offer inducements to the capitalist to invest and open out the resources of the country, either agricultural or mining. But these old Siamese Governors are conservative of all their rights, and desire nothing more urgently than to keep the bustle and activity of civilization as far away as possible, and to be allowed to tread their own way, and I, for one, do not blame them.


What the future of those provinces may be, I cannot say; no doubt they contain much latent mineral and agricultural wealth, but as long as the Siamese labourer can gain his

pound of rice for $2\frac{1}{2}$ cents and his pound of fish at the same rate, he will wait a long time ere he will sweat himself to uncover their wealth. Besides the struggle of the native of these provinces is not for bread, it is for health, the high mortality leaving those that survive room enough and to spare to make an ample living.




SIAMSE PROVINCES KHOH LAK TO MERGUI

KHOH LAK TO MERGUI



10°



1000 ft. to inch
700 ft. to inch

Cross-section across Bangtaphan Province

THE ALLEGED DISCOVERY OF MERCURY IN MALACCA.

BY

DR. W. BOTT, F.C.S. F.G.C.S., F.P.S., &C.



IN November last year, I was informed that a quantity of mercury had been discovered during excavations on St. Paul's Hill, Malacca, on the site of the new water reservoir for the town. The discovery, I believe, produced some little excitement at the time, and may, in certain quarters, even have conjured up pleasant visions of a new-born revenue of Malacca from a rich mercury mine. Now the existence of such a treasure would doubtless have resulted in a transfer of the proposed reservoir to a new place and the giving up of the venerable hill of St. Paul to the rude attacks of the pick and spade of the miner, and, as the work of excavation was then progressing, and the very spot of the discovery was about to be covered up, it seemed desirable to enquire into the matter without delay. In due course, I received two large bags of the supposed ore—one from the exact place where the mercury had been collected, the other from the immediate vicinity. Both samples consisted of bright red earth, and the colour might or might not have been due to cinnabar (Sulphide of Mercury) as far as mere appearance went. On closer examination, visible globules of metallic mercury were found throughout the whole of the first sample, while the second contained none. Unfortunately the character of the soil in which the mercury was found, did not accord with its occurrence, for further examination shewed that the colour of the soil was entirely due to red, argillaceous ferric hydroxide. and that besides

metallic mercury it contained no combined mercury, no cinnabar, or in fact a sulphide of any kind. Now the presence of mercury in such a matrix of laterite pure and simple is a mineralogical impossibility, and my first attempt to solve the problem was by sharply questioning the Laboratory Steward whether he had lately spilled any mercury about the place. This led to indignant protests and no result, and thus there was left but one possible hope of explanation, viz., the existence of a cinnabar vein in St. Paul's Hill, from which the mercury had been derived, although it seemed strange that none of the cinnabar had found its way into the sample with it. A report was sent to the effect that the mercury was there, that it had no business to be there, and its presence could only be accounted for by a somewhat improbable hypothesis. At the instance of the Acting Resident Councillor, I was then asked to visit the place in person, but was unable to do so until the end of January, when I found a great part of the site covered up with brickwork. No difficulty was, however, experienced in ascertaining the nature of the entire formation, and proving the absence of a cinnabar vein and of sulphides generally. The existence of metallic mercury, however, was confirmed; appreciable quantities of it were found in several places, in fact 2 oz. were collected in about ten minutes. Having proved that this metal could not primarily have been present in the soil, it follows that it must have found its way into it by accident, and this opens up an unlimited field for speculation, unfortunately, however, without the redeeming feature of being able to tell the correct guess from the rest.

As several pounds of the metal had been obtained with little trouble in one place, and more might have been collected in other parts, it is evident, that a considerable quantity of mercury must, at one time or another, have been placed into the ground intentionally or accidentally; this suggests the following possibilities and new questions:—

1. The metal was intentionally poured into the ground. It is hard to conceive a reason for this; "salting" mercury mines would be a new idea entirely.

2. The metal was hidden in the ground. This is possible, and may have been done for an unlimited number of reasons, but then suitable vessels would have been used, and no trace of such has been unearthed.

3. The mercury was spilled by accident. This too is possible; it must have been a pretty big and expensive spill, and the operator does not seem to have tried very hard to pick up the precious fluid.

4. Who buried or spilled the mercury? There is no record of mercury being stored on St. Paul's Hill during recent years; the Dutch or Portuguese may, therefore, have been the guilty party.

5. What was the mercury intended for? Gold-mining?

6. Why was it buried in the ground if we discard the spilling theory?

The above and a legion more questions and conjectures environ the mercurial puzzle of St. Paul's Hill, Malacca. Whether any light will ever be thrown upon the dark problem of the bright metal, whether one of the many contributors of this Journal familiar with side issues and odd details of Malacca history will be able to produce facts bearing upon the matter, remains to be seen. Until then, to Malacca will belong the credit of an inexplicable conjuring trick in the mineral line, unless the reader traces the mercury back to spiritualistic agencies, or concludes, with TOPSY, that it "growed."

W. BOTT.

NOTE.—Since writing the above I have found the possible explanation of the discovery of mercury in St. Paul's Hill. CAMERON in his book "Malayan India," chapter XIII, states, that about 1864 Captain PLAYFAIR discovered at the base of the hill an old cellar or store-room which had formed part of the old Portuguese Government buildings. In it—in a small recess—were found forty or fifty earthenware pots mostly crumbled to pieces, but each of those which were whole contained a small quantity of mercury. Only about four pounds

altogether were collected, but had the pots been full, as they probably were when first stowed away, they must have contained more than a ton of mercury. The metal had been brought there in connection with gold-mining on Mount Ophir. As only a small quantity of the metal was recovered in 1864 and recently, a ton or so of the metal, representing a value of about \$2,000 would still remain buried in St. Paul's Hill. It remains, however, still to be proved that the mercury recently found is part of that formerly discovered by PLAYFAIR. The latter distinctly states that the mercury was discovered at the base of the hill, whereas the recent finds were made some distance above the base. It is difficult to conceive that this mercury should have worked its way up the hill, and it is more likely to be traced to another hiding place or store-room made on a higher level.

W. BOTT.



A NOTE ON RENGAS POISONING.

BY

W. C. BROWN, M. D.



THE commonest of the various species of Rengas is *Melanorrhæa Curtisii Oliv*, a tall and handsomely foliated jungle tree which flourishes luxuriantly on many of the hill slopes of the Malay Peninsula, and extends northwards as far as the Southern Provinces of Burma. All the trees of this family contain in abundance an acrid sap, which is well known to native woodmen as a substance that, coming in contact with the skin, produces disagreeable and even dangerous results.

If the healthy skin is rubbed lightly with the juice from a freshly cut twig, violent inflammation, with smarting and burning pain, follows within twenty-four hours, and results in a characteristic pustular eruption—an eruption of blebs filled with matter. If the injured surface be of any extent, fever and other constitutional disturbances follow the local injury. When a large extent of skin has been affected, as happens when a native with unprotected body struggles through broken branches, this fever is said to be so irritant and septic that it not infrequently ends fatally. I have not, however, seen a case of such gravity, but from the peculiarly severe symptoms produced by the sap on a small surface there can be little doubt but that where a large extent of skin is involved, the consequences might be most serious. An extract or tincture of the twigs made by soaking them in proof spirit has active blistering properties, and might be of value in medicine as a counter-irritant, did we not possess in croton oil and tartarated antimony remedies

whose action in producing a pustular eruption is identical with Rengas sap, and which have the additional advantage that their constitutional action is never dangerous. Internally, the juice acts as a violent irritant, causing vomiting and purging, and its administration is in the highest degree dangerous.

Malays assert that simple contact with the leaves or unbroken branches is sufficient to give rise to equally evil effects, but experiment points to the fact that it is the sap of the tree that possesses the noxious qualities, and that simple contact with unbroken twigs is not usually hurtful. It is a matter of some importance, however, to note that the wood of the Rengas tree, which, being closely grained and capable of taking a high polish, would otherwise be of great value for cabinet-making, retains its irritant properties long after the sap has apparently dried. I am informed by Messrs. KNIGHT & Co. of Singapore that, after years of seasoning, when the wood is cut up it gives rise to painful and intractable eruptions on the hands and bodies of the workmen, and that, for this reason alone, its value as an economic wood is seriously impaired.

The immediate treatment of the poison is generally successful; it should be to wrap the injured part in bandages with some dry alkaline powder, such as bicarbonate of soda, the object being to counteract the acid of the poison, and to absorb the exuded secretion from the skin. Vaseline or lard smeared on the hands and bodies of the woodcutters prevents the sap from setting up inflammation by affording a protective covering to the skin. Many cases, however, especially those that have been neglected, are refractory to treatment of any sort, and the ulcers that form between the fingers and toes as a result of Rengas poison are extremely untractable and sluggish in taking on a healthy and healing action; the eruption in emaciated and unhealthy subjects being very apt to develop into a condition of chronic ulceration.

If we pass from the region of observed fact to the more dubious realm of Malay belief and narrative, we are told that when a Rengas tree is felled, the exhalations from the cut

stem are so noxious that on the woodmen inhaling them, their faces become swollen as if they had been stung by hornets, and that, although none of the sap has touched the skin, fever and blindness are usual consequences. It is impossible to accept these statements, but it is quite possible that some inflammation of the outer membrane of the eyes might be caused by the irritating vapour of Rengas sap.

With such distinctive characteristics as these, it is not surprising to find that Malays have invested the Rengas tree with supernatural properties. Before they venture to fell the tree to obtain the highly prized *tiangs* of this wood, elaborate exorcisms have to be undertaken to counteract the influences of evil-disposed *djins* and *langsuyar*. It is a matter of popular belief that, if the incantations are efficiently performed, the woodcutter is rendered proof against the poison of the sap; and three Malays living at Ayer Etam in Penang are believed to be such adepts in the art that their bodies have become impervious to the influence of the juice. I have experienced great difficulty in getting other natives to collect branches, as they insist that it is essential to have some one to do it who thoroughly understands the spells proper for the occasion. On investigation, it was found that all that these Malays claimed to achieve by their incantations was to attain exemption from the effects of the noxious vapour of the cut trees, and that they are careful to augment the protective powers of their spells by rubbing their hands and bodies with coco-nut oil.

The various species of the Rengas family must be frequently met with in the Peninsula in clearing jungle for plantations or railways, and it is obviously a matter of importance to be able to identify trees that may expose workmen to serious danger.

Penang, 29th October, 1891.

A
NEW COLLECTION
OF
MALAY PROVERBS.

—:o:—



ANY of the following Proverbs have been known to me for a considerable period, but though familiar with their use in colloquial Malay, I never reduced them to writing until lately when, travelling in the interior of Pahang, I was driven to do so for the sake of the occupation. At the same time I considerably enlarged my original stock of hitherto unpublished Proverbs, and, to the best of my belief, the present collection contains no Proverb which has already appeared in print.

The Malay language is one which lends itself to that terseness of expression and felicity of diction which is so essential to an aphorism, and which is often the despair of the translator, who attempts to compress the meaning conveyed by a few pithy Malay words, into as many lines of English. The Malay Proverb is also interesting from a philological point of view, because in these old-time sayings many words are found which do not occur in any Malay writings, though such terms are, for the most part, common enough in the colloquial dialects.

By the Malay, who is conservative to the backbone, and a *laudator temporis acti* by nature and education, all the wisdom of the ancients (*per-kata-an orang tua-tua*) whether preserved in Proverb, Charm, or Precept, is accepted with perfect faith as incontrovertible truth. As a race, too, the Ma-

lays delight in obscure hints and darksome metaphors. An educated Malay will ask for his neighbour's daughter in marriage to his son in a letter which is simply a string of *doubles entendres*, and in familiar colloquial discourse a native will quote the first line of a verse (which in its entirety contains four lines, the first two of which are wholly irrelevant and are only introduced for the sake of rhyming with the two last lines), leaving his bewildered hearer to infer his meaning from a knowledge of the lines which form the rest of the verse, the first line of which has been given. In discussions among Malays, too, it is the man who can quote, and not he who can reason, that bears away the palm. I need hardly add that a Proverb which is both ancient and obscurely metaphorical, is immensely popular with all classes of Malays.

In preparing these Proverbs for the press, I have endeavoured to combine, as far as possible, an absolutely literal translation with a correct rendering into English of the Malay meaning.

The Proverbs are arranged in order according to the sequence of the Malay alphabet.

HUGH CLIFFORD.

1 اداكه هملغ بيسا اولر مپومف دباوه اكر

Ada-kah hilang bisa ular meny-usop di-bawah akar.

Will a snake lose its venom through creeping under a root?
i., e., will a Chief lose his rank through being civil to his inferiors?

2 اصل انكوتوغل اداله بيس

Asal angkau tugal ada-lah benis.

If you make holes in the ground, seed shall be forthcoming.
Used to signify that if an offence is committed retaliation will follow.

Tugal توغل = the holes in the ground prepared for the reception of seed; to sew seed in holes prepared for it; to make holes in the ground for the reception of seed.

FAVRE renders this word "*Culture du riz dans des terres élevées et sèches*," but omits to mention the particular manner of planting to which alone this word (so far as my experience goes) is applied.

3 انكوق بوكن كيليف يا

Anggok bukan géling ya.

Not a nod but a shake of the head.

To feign a desire for one thing while really desiring something different.

4 التن فانه لسونغ تا بر بوي

Antan patah lesong hilang.

If the pestle be broken, the mortar will be lost.

If the husband be impotent, the wife will prove unfaithful.

5

الجميع كالعنق باني براني
Anjing galak babi brani.

The dogs are ferocious and the pigs are daring
 Used to signify that both sides are 'spoiling for a fight.'

6

انق بايق مننتو موليق
Anak baik menantu molek.

A good son and a pretty daughter-in-law.
 Used in speaking of people who are entirely of one mind.

7

اولر بوكن ايكن فون بوكن
Ular bukan ikan pun bukan.

Neither snake nor fish.
 'Neither fish, flesh, nor fowl, nor good red-herring.'

8

ايكن بياردافت سرمشغ جاغن فوكه
Ikan biar dapat serampang jangan pokah.

Let the fish be caught, but let not the fish spear be bent.
 To act with due discretion.

Compare Prov. 27 in No. 11 of this Journal, and the two Proverbs with which Mr. MAXWELL there compares it. The familiar version of Prov. 27 with which I am best acquainted, runs as follows:—

اولر بيمر ماتي تانه جاغن لكوق بولوه جاغن فاته

Ular biar mati tanah jangan lekok buloh jangan patah.

Let the snake die, but let not the ground be dented, nor the bamboo broken.

Pokah فوكه = to bend ; to become bent ; to be bent ; to

be out of the straight ; bent ; crooked.

This term, though a common colloquial expression, is not, I believe, to be found in any Malay Dictionary now published.

9

ايکن ٲرکيرف جال ٲيب

Ikan ter-kirap jala tiba.

As the fish shake themselves clear, the casting net arrives.
Too late.

10

باتو کچيل برگوليف نايک باتو بسر برگوليف ٲورن

Batu kechil ber-guling naik, batu besar ber-guling turun.

Little stones roll up, and big stones roll down.

The end of all things, when the common people shall be greater than the nobles.

11

باتو هيٲتم ٲا' برمنديغ

Batu hitam ta' ber-sanding.

A black stone without projections.
Difficult to injure.

12

باغوم منٲا' اكو لهير بادق ٲ منٲا اكو داغيغ

*Bangau ! Bangau ! minta aku lehir. Badak ! Badak !
minta aku daging.*

Oh stork ! I beg from thee thy neck ! Oh rhinoceros ! I beg from thee thy flesh !

Compare :—

Prov. 64 of Mr. MAXWELL'S collection published in No. 2 of this Journal.

دمتعا كقد يغار

Di-minta ka-pada iang ada.

Ask from those who possess, &c.

13

باكي الجيغ ترمشيت دفاكر

Bagai anjing ter-sepit di-pagar.

Like a dog squeezed in a fence.

Viz., Making a hideous clamour.

The form :—

باكي الجيغ ترمشيت ايكور

Bagai anjing ter-sepit ekor.

Like a dog when its tail is squeezed.

And also met with.

14

باكي الجيغ ملنتغ دلي

Bagai anjing me-lentang denai.

Like a dog hunting wide.

(*Lit.*, crossing a jungle path.)

Denai دلي = a small foot-path in deep jungle. This word, though often met with in colloquial Malay, is not to be found in any Dictionary.

15

باكي الجيغ مهالق دفتتس غاجه

Bagai anjing meny-alak di-pantat gajah.

Like a dog barking at an elephant's stern.

Impotent rage.

16

باغي النى دارا مابق الدم

Bagai anak dara mabok andam.

Like a maiden overcome by the fringe upon her forehead.

Self-conscious ; shy.

Andam الدم = the fringe of short hair cut upon the forehead of newly married girls, or girls who are about to be married. FAVRE mistranslates this word—'Arranged, adjusted, &c.'

17

باغي اور گنتوغ کتبيغ باغي تببيغ گنتوغ کا اور

Bagai aur gantong ka-tebing, bagai tebing gantong ka-aur.

Like a bamboo hanging from a river bank, and like a river bank depending from a bamboo.

To lean on a broken reed.

18

باغي اير تيتيق كباتو

Bagai ayer titek ka-batu.

Like water dripping on to a stone.

19

باغي ايكن كنا توب

*Bagai ikan kena tuba.*Like a fish poisoned by the *tuba* (*Derris elliptica*) root.

In difficulties. 'Up a tree.'

The above Proverb is generally used with several others:—

باغي ايكن كنا توب اودغ ددالم تفكوق

باغي اولس نغا ترهمفس كباتو

باغي روان ددالم توار

باغي انق مسمم كتهور
باغي دکن دهاوه ففکل بوله

Bagai ikan kena tuba.

Udang di-dalam tanggak.

Bagai ulat nangka ter-hempas ka-batu.

Bagai ruan di-dalam tuar.

Bagai anak sepat ka-tohor.

Bagai dekan di-bawah pangkal buloh.

Like a fish poisoned with the *tuba* root.

Like a prawn in a prawn-basket.

Like jack-fruit grub dashed against a stone.

Like a *ruan* fish in a rattan fish-trap.

Like a little *sepat* fish in the shallows.

Like the bamboo-rat under the roots of the bamboo.

Ruan روان and *Sepat* مسمم are both small fish found in the *padi* swamps, and are usually caught by means of a *serkap*.

Tuar توار = a fish-trap of a conical shape, made from the *onak* thorn, and so constructed that a fish can enter without difficulty, but cannot get out without being hooked by the thorns.

20

باغي ايم دماکن توغو

Bagai ayam di-makan tungau.

Like a fowl devoured by parasites.

Poorly; seedy-looking; in poor condition.

Tungau توغو = a small red parasite found on the sand-sprits which line the river-banks in the Peninsula. It breeds in the dung of buffaloes.

21

باغي بودق مافو هيفومس
Bagai budak sapu hingus.

Like a child which has its nose wiped for it.
 A muff.

22

باغي تاگر دڤولو سمبيلان
Bagai tagar di-pulau sembilanan.

Like a thunder-clap at Pulau Sembilan.
 Used to describe a very loud noise.

This Proverb is in use in Perak. The place mentioned is a group of islands at the mouth of the Perak River.

23

باغي تومن ماكن انق
Bagai toman makan anak.

Like the *toman* fish which eats its own young.

Applied to Chiefs who misuse their powers, oppressing those whom they are bound to protect.

Toman تومن = a fresh water fish which attains a considerable size. Its colour is a reddish yellow, and its flesh is of a saffron hue. It subsists on other fish.

24

باغي رابق دغن ابي اصل بردكف مپاللاه اي
Bagai rabok dengan api asal ber-dekat me-nyala-lah ia.

Like tinder and fire, which if they come near one another burst into a blaze.

Applied to people who are deadly enemies.

25

باغي فليدق ددالم چراغ.

Bagai pelandok di-dalam cherang.

Like a mouse-deer in a clearing, *i. e.*, stupid, bewildered, timid.

Cherang چراغ = a clearing ; an opening in the forest made for planting purposes. This word is much used in the States on the Eastern slope of the Peninsula. It is probably derived from *Trang* تراغ light, open, clear, &c. The form *chě-rakn* (a clearing) met with in the Sĕn-oi dialect of Sakai, has the appearance of being formed from the Malay word and not *vice versâ*.

26

باغي کرا دافس چفگوغ

Bagai kra dapat changgong.

Like an ape which has caught hold of a snag.
Clinging like grim death ; as a fool clings to bad advice.

27

باغي کليف ۲ تريغ مالم

Bagai klip-klip terbang malam.

Like a fire-fly flying by night. Said of a supposed secret which is widely known.

Compare :—

ايم فوئيه تريغ مالم

Ayam puteh terbang malam.

A white fowl flying by night, in Mr. MAXWELL'S collection of Malay Proverbs.

28

باغي کوچيغ دغن ففگغ

Bagai kuching dengan panggang.

Like the cat with the roast.

Impossible to keep apart. Said of the mutual attraction between young people of opposite sexes.

29

باگي گاله دتغه اروس

Bagai galah di-tengah arus.

Like a boat-pole in mid-stream, *i. e.*, shaking from head to foot.

30

باگي مامندول بهرو برانق

Bagai ma' mandul bharu ber-anak.

Like a hitherto barren woman who has just given birth to a child.

"As pleased as Punch."

Compare :—

باگي اورغ بوت بهرو چليق

Bagai orang buta bharu chelék.

Like a blind man who has newly recovered his sight.

31

باگي مناريق رمبوت ددالم تفوغ

Bagai men-arek rambut di-dalam tepung.

Like pulling a hair out of flour.

"As easy as lying."

32

باگي هريمو برانق مودا

Bagai harimau ber-anak muda.

Like a tiger with young cubs.

Irritable; angry without provocation.

33

بايك فوطيه تولغ جاغن فوطيه مات

Baik puteh tulang jangan puteh mata.

Better white bones than white eyes.

This Proverb contains an idiom ; *puteh mata* (white eyes) being a common Malay expression signifying "ill-feeling," "black blood," &c. The meaning of the above Proverb thus is, that it is better that a quarrel should come to a head, even though it ends in the death of one of the parties concerned, rather than that ill-feeling should exist endlessly.

34 بيراڤكه تاجم فيسو فارغ تاجم لاڤي مولوت مانسي
Be-ber-apa-kah tajam pisau parang tajam lagi mulut manusia.

However sharp is a knife or a chopper, sharper yet is the human tongue.

35 بتول ۲ ايڤوراڤيغ كالو بڤيماڤ فون اد جوڤ بيفكوڤن
Betul-betul ekor anjing kalau bagai-mana pun ada juga bengkok-nya.

However straight a dog's tail may be, it will still be crooked.
 A rogue will never be altogether honest.

36 بچاراكن رومفت دهلامن اورغ دهلامن سنديري رومفت
 مسفي ككاكي تڤاڤا
Bechara-kan rumput di-halaman orang di-halaman sendiri rumput sampai ka-kaki tangga.

To take thought of the grass on one's neighbour's lawn while that on one's own house-plot spreads even unto the foot of the door-step.

The beam and the mote.

37 بيراڤ فون انڤيغ مپالڤ بوكيم بولهكه رونته
Be-ber-apa pun anjing meny-alak bukit bulih-kah runtoh.

However much dogs may bark, will a hill fall to pieces? *i. e.*, Does a man of assured position regard the slander of his inferiors?

38 برتليڠكوه انتن دلڠسوڠ ايم جوڠ يڠكنيڠڠ

Ber-telingkōh antan di-lesong ayam juga yang kenyang.

When the pestles fall together into the mortar, the fowls have their bellies-full.

When men of rank quarrel, the common people reap the benefit.

Telingkōh تليڠكوه = to collide; to clash. So far as I know this word is not to be found in any Malay Dictionary.

39 برچري تيدق برطلق برنيكج تيدق برقضي

Ber-cherai tidak ber-talak ber-nikah tidak ber-kathi.

Divorced without the ceremony of divorce, and married without the assistance of the clergy.

Said of the parents of a bastard.

40

بري بتيس هندق فها

Bri betis handak paha.

Give the leg and the thigh is wanted.

"Give an inch they take an ell."

Compare Proverb 99 in this collection.

41 بلوم توارڠ فنڠڠ بواه سڠكواڠ مبسر بتيس

Belum tuarang panjang buah sengkuaḡ s'besar betis.

Before a drought has lasted for a long time the *sengkuaḡ* fruit are grown to be as big as the calf of one's leg.

Said when false reports are spread without any foundation for them.

Tuarang = a drought which is even of longer duration than the ordinary drought called *Kamarau* کامارو.

Sengkuang = the name of a creeping plant which has an edible root. The fruit is said only to grow large during very dry weather.

42

بیرم نایک کامات

Biar-biar naik ka-mata.

Intestinal worms ascend even unto the eyes.

“Do not put off till to-morrow what may be done to-day.”

This Proverb contains a play upon the words *biar-biar* بیرم (intestinal worms), and *biar* بیر (to leave alone; to let be); but it is impossible to render this pun in an English translation. This Proverb is generally used in answer to any one who says *Biar-lah dahulu* بیرله دهول “let it be for the present;” in reply to an order bidding him do anything.

43

تا بوله دبايكي جاشن دفيچهكن

Ta' bulih di-baik-i jangan di-pichah-kan.

If you cannot mend it, do not break it more.

44

تا موغله گنه داون دشن مبنر گنه باتغ جوگ

Ta' sunggoh getah daun dengan s'benar getah batang juga.

Not really resin from the leaf, but in truth resin from the trunk of the tree.

Said of one who has authority which is the result of the influence of some one greater than he, with whom he is connected.

45

تانه لمبه کندوشن اير کايو بيغکوق ليتين کرا

Tanah lembah kandong-an ayer, kayu bengkok titi-an kra.

Low-lying land holds water ; a crooked bough is used by monkies as a bridge.

The rich prevail, the poor go to the wall.

46 تاومسهله اجر النى بوای برنغ دي مده فندي مدي

Ta' usah-lah ajar anak buaya bernang, dia sudah pandai sedia.

It is not necessary to teach a young crocodile how to swim, as it is already expert.

"To teach one's grandmother to suck eggs."

Compare :—

ايتيق داجر برنغ

Itek di-ajar bernang.

The duck is taught to swim.

(No. 77 of KLINKERT'S Collection of Proverbs).

47 تاومسه بيمبغ گولا ددالم مولف بيل هندق تان تان

Ta' usah bimbang gula di-dalam mulut bila handak telan telan.

Be not solicitous concerning the sugar which is in your mouth, when you desire to swallow it, do so !

Do not be anxious about trifling undertakings.

48 تاهن جرم سوروغ كفلا

Tahan jerat surong kapala.

Set a snare and thrust your head into it.

Applied to those who through excess of cunning overreach themselves.

49

تراص تراولجم گوبل ملايغ
بر تينتيغ بر تلمشي تراص
گوبل ملايغ تراص تينگل

Tras ter-unjam gubal me-layang.
Ber-tinting ber-tampi tras.
Gubal me-layang tras tinggal.

The core of the tree is fixed firmly in the ground, and the light wood which surrounds it is blown away. When the core of the tree is sifted, the light wood is blown away, but the core of the tree remains.

If common people compete with nobles, they come to an evil end, while the nobles remain unharmed.

Unjam اولجم = to stick upright in the ground. This word is exactly similar in meaning to *Chachak*, چاچق.

50

ترکنا فد ايکن بر مسورق
ترکنا فد باتغ مامم

Ter-kena pada ikan ber-sorak.
Ter-kena pada batang masam.

To shout with triumph if one strikes a fish, and to be cross if one strikes a log.

Unduly elated by success and unduly depressed by adversity.

51

چاغن دمغکاکن ايکن لایس ۲ تیدق میغف

Jangan di-sangka-kan ikan lais-lais tidak meny-engat.

Do not fancy that the *lais-lais* fish will not sting.

“Despise not thy enemy.”

Lais-lais لايس ۲ = a small fresh water fish which is armed with sharp stings in its fins.

52 جاشن داچر انى هريمو ماكن داكيغ
Jangan di-ajar anak harimau makan daging.

Do not teach a tiger cub to eat flesh.

Said to those who give bad advice to a young *Raja* or Chief who unassisted will learn evil ways quite quickly enough.

53 چكلو برانق ايكوف كامف بيدن
Fikalau ber-anak ikut kata bidan.

If thou art confined of a child, do as the midwife bids thee.
Follow the advice of experts.

54 چاريق ۲ بولو ايم لام ۲ اي برچنتوم فول
Charek-charek bulu ayam lama-lama ia ber-chantum bula.

Though a fowl's feathers may be torn to tatters they will eventually grow again thickly.

Though relatives may quarrel, they will eventually make friends again.

Chantum چنتوم is translated by FAVRE, “Piqué, piqué à l'aiguille,” but that I believe to be only a secondary meaning, the term being principally used to express a thick growth of any kind.

In Pahang, *Chantum* چنتوم in this Proverb is often replaced by *Tampun* تمشون, which has the same signification as *Chan-*

tum, but is not as universally understood among the Malays of the Peninsula.

55 دتوب مهاجكه ايكن دجال جاربع بوككه ايكن
Di-tuba sahaja-kah ikan di-jala jaring bukan-kah ikan.

Are those alone fish which are poisoned with the *tuba* root?
 Are those not also fish which are caught in the nets?

Usually applied by a plain woman in the sense "are accomplishments confined to those who are well favoured?"

56 دمفكاكن لاغيت ايه رنده
 دندغ دكف دچافي تا بوله
Di-sangka-kan langit itu rendah.
Di-pandang dekat di-chapai ta'bulih.

To think that the heavens are low, but though they look as though they were close they cannot be touched with the hand.

Said of the relationship which exists between a *Raja* or Chief and his followers.

57 دمان نمف كوتو هندق ماكن چكلو ليدق داتس كثلا
Di-mana tempat kutu handak makan jikalau tidak di-atas
kapala.

Where shall lice feed if not upon the head?
i. e., Where shall the poor find relief if not from the bounty of the great?

58 دمانكه برتراس كايو مهغ
Di-mana-kah ber-tras kayu mahang.

Where is there found a hard wood core in a *mahang* tree?
 Do not expect impossibilities.

Mahang مهغ = the name of a forest tree of very soft wood (*Macaranga*). Native tradition says that the name of the State of Pahang is derived from this word. The tiny brook which debouches at Kuala Tembeling and gives its name to the great stream formed by the Jelai and Tembeling Rivers, taking its source, it is said, at a spot where a giant *mahang* tree once stood.

59

دمثيب تيدق برسگيب

دتميب تيدق برتالي

*Dempit tidak ber-sanggit.**Di-tambat tidak ber-tali.*

Pressed together but not rubbing, tied without twine.

Said of persons who live together, but who are not bound by any real tie.

Sanggit شگيب = to rub against; to touch against anything. To come in contact, or to be in contact with anything.

The root, which is found in *Dempit* دمثيب, is one which has retained its form and its meaning in a more marked degree than almost any other Malay root. This is exemplified by the following words:—

Apit افيب = to press, &c.

Kapit كافيب = to press, pressed, &c. (See *Pulut kapit* فولوب كافيب = pressed olenaceous rice.)

Kepit كثيب = to carry under the armpit, *viz.*, pressed between the arm and the body.

Sepit سثيب = to squeeze, &c. (See Proverb 13 of this Collection.)

Hempit همثيب = to get close to anything; to press against it, &c.

Sempit شَمْفَيْت = narrow, confined.

Sumpit سَمُومْفَيْت or *Kempit* كَمْفَيْت = a narrow rice bag (a place in which rice is confined).

Chubit چُوبَيْت = to pinch.

The word *Apit* اْفَيْت has been traced to a Sanskrit root; but either the resemblance is a chance coincidence, or else this root which runs through so many common words in Malay is itself derived from Sanskrit, which would seem improbable.

60 جاغن دمسل گولوغ برلاري هيلغ كابوت تمفتله دي
Jangan di-sesal gunong ber-lari, hilang kabut tampak-lah dia.

Fret not through fear that a mountain will run away, when the mist clears it will be still in view.

Do not be anxious about remote contingencies.

61 سام كاین بامه مهلي تيمه مسوكو كالوتا تيمه فرغي مراسم
منداف

*Sama kain basah s'hele timah s'suku, kalau ta' timah
pergi s'rasa men-dapat.*

A bathing cloth and a *suku* of money are of equal value.

If only one did not pay the money away, one would feel as though one had gained by the bargain.

Said of one who wishes to "have his cake and eat it too."

62 مسرف انجیح دافة فاسير
Seperti anjing dapat pasir.

Like a dog on a sand-bank.

Running heedlessly hither and thither through sheer rude health and robust spirits.

- 63 مسفرم ددالو افي هيغكف كفوهن كا يو
 هيغكف كا ابو ايون ماتي
 هيغكف كرنتيغ رنتيغ فاته
Seperti dedalu api hinggap ka-pohon kayu.
Hinggap ka-ibu ibu-nya mati.
Hinggap ka-ranting ranting-nya patah.

Like a spark of fire falling upon a tree.

If it falls upon the trunk the trunk will die.

If it falls upon a branch the branch will be broken.

“ Evil communication corrupts good manners.”

- 64 مسفرم اورغ ماتي جكلو تباد اورغ مفاشكف بيلاكه برغري
 Seperti orang mati jikalau tiada orang meng-angkat
 bila-kah ber-grak.

Like a dead man. If no one lifts him up, when will he be able to move?

Helpless. Impotent.

- 65 مسفرم برگنتوغ فد رمبوم سهلي
 Seperti ber-gantong pada rambut s'hele.

Like hanging on a single hair.

A precarious position.

- 66 مسفرم تبولفس كدالم مولم غاجه فاه اكن كلوان
 Seperti tebu lepas ka-dalam mulut gajah payah akan ka-
 luar-nya.

Like sugar-cane which has entered an elephant's mouth, difficult to extricate.

Applied to missed opportunities.

67

مفردة لمفردت جاچه لالو

Seperti tempat gajah lalu.

Like a place through which elephants have passed.
Said of places through which a royal progress has been made.

68

مفردت ساك دانس تالم

Seperti saga di-atas talam.

Like a single pea on a large tray.
Said of one who is close-fisted and inhospitable. The Bar-
mecide's feast.

69

مفردة فيسو راوة بيغكوق اورغ بوله بتول بيغكوق

مسنديري تا بوله بتول

*Seperti pisau raut bengkok orang bulih betul bengkok
kita ta' bulih betul.*

Like a (bent) knife used for scraping *rotan*. The twists in
others can be made straight, but one's own kinks can never be
straightened.

Applied to himself by one who considers that his offences
are more severely punished than the offences committed by
others.

"One man may steal a horse, and another may not look over
the fence."

A *pisau raut* has a slightly curved blade which is about
the size of an ordinary razor. This blade is attached to a
handle, which is usually double or treble the length of the
blade. As its name implies it is used to "scrape" *rotan*, etc.

70

مفردة كربو مسندوق انق دغن فامر ليدق دغن هوجغ

*Seperti kerbau men-andok anak dengan papar tidak dengan
hujong.*

Like a buffalo butting its young with the flat and not with the points of the horn.

Parents' anger to their children is always tempered with love.

71

مفثرة كلي دوا ملوبڠ

Seperti kli dua s'lobang.

Like two *kli* fish in one hole.

Said of people who are in one mind in all their enterprises. *Kli* كلي = a long scaleless fish found in the *padi* swamps.

In Ulu Pahang the two Chiefs who live in the far *Ulu* of the Lipis District bear the titles of TO' KLI and TO' BAKAR, respectively. Neither of them has ever visited the Sultan of Pahang, which fact caused the Sultan to declare that TO' KLI was too slippery to hold, and TO' BAKAR (Bakar باكر = to burn) too hot to handle. This royal witticism is a good specimen of the Malay pun.

72

مگن برتان مسيب جان

Segan ber-tanya sesat jalan.

If you are too shy to make enquiries, you will lose your way.

73

مسمليه ايم دغن فيسو مسمليه اورڠ دغن كاس

Semleh ayam dengan pisau, semleh orang dengan kapas.

Cut a fowl's throat with a knife, and a man's throat with cotton.

Men are deceived by soft words.

74

موسو ددادا تا'دافت دايلثكن

Susu di-dada ta'dapat di-elak-kan.

Milk at the breast cannot be shirked.

Responsibilities cannot be evaded.

Elak ايلق = to shirk ; to avoid ; to escape from. Though this word is of daily occurrence in colloquial Malay, I have not met it in any Malay Dictionary now published.

75

مياث مفاكو بيرق دلفه جان

Si-apa meng-aku berak di-tengah jalan.

Who will confess to having committed a nuisance ?

i. e., Who will own to a disgraceful act ?

76

ميتكور كاتق دباوه تمفورونغ دمفكانك تيا دليا يغلان

S'ekor katak di-bawah tempurong di-sangka-kan tiada dunia iang lain.

A frog beneath a coco-nut shell believes that there is no other world.

Applied to those who are unduly impressed with the importance of themselves and their own immediate surrounding.

77

ميتكور كربو ممباوا لومفر مسموان ترفاليت

S'ekor kerbau mem-bawa' lumpor samoa ter-palit.

One buffalo brings mud and all the herd are smeared with it.

Evil example is quickly followed.

78

عبارت بورغ ماة لفس بادلي تركورونغ

Ibarat burong, mata lepas badan-nya ter-kurong.

Like a bird whose eyes are at liberty to wander, but whose body is in confinement.

- 79 عبارة بورغ مولت مالنسي جاغن دفاكي
Ībarat burong mulut manis jangan di-pakai.

The metaphor of a bird—Do not place trust in sweet words.
 Said of chance travellers who “love and ride away.”

- 80 فادم مپالا تاريق فوننتوغ
Padam me-nyala tarek puntong.

When the blaze has gone out, you can draw the logs from
 the fire-place.

This Proverb is used as a boast, meaning that the speaker,
 like Mr. AUGUSTUS MODEL, “will never be taken alive.”

- 81 فاته توغكت ترجرامغ
Patah tongkat ber-jeremang.

The staff being broken, to go supporting oneself upon all
 fours.

Said of one who will never give in and confess himself to
 be beaten.

- 82 فاته تومبه هيلغ برغنتي
Patah tumbuh hilang ber-ganti.

That which is broken off shoots afresh. That which is lost
 is replaced.

All things have their compensation.

“When I have seen the hungry ocean gain

“Encroachment on the kingdom of the shore,

“And the firm ground win of the watery main

“Increasing store with loss and loss with store, &c.”

83

كالو تا' سير بردگف سير ملايغ

Kalau ta'sir ber-degap sir me-layang.

If not head over ears in love, at least slightly smitten.

Sir سير = to care, to desire, to want, to wish for, to be anxious to obtain. To condescend to.

In his collection of Malay Proverbs (Prov. 272) printed in No. 3 of this Journal, Mr. MAXWELL gives this word as *Sior* which, however, is only the local pronunciation of the word in the Perak dialect, where a final *ir* is always rendered *ior* in the colloquial pronunciation, e. g., *Kikir* كيكير, a file, is pronounced *Kikior* in the Perak dialect. This word is in common colloquial use in Perak, Pahang, Kelantan, Trengganu and Patani. Another word *Sualak* سوالق, which is not included in lists of words yet published, is also frequently met with in the colloquial dialects, especially in Perak. Mr. MAXWELL states that *Sir* is identical in meaning with *Ingin* اينجين, but in the negative form *Ta' sir*, and *Ta' sualak* also, are perhaps better rendered into polite Malay by the phrase *Ta' sudi* تا' سودي.

Degap دگف which is also a slang word not found in any Dictionary, signifies 'thoroughly, altogether,' &c. *Kena degap* كنا دگف has much the same signification as *Kena tepat* كنا تفت to be hit full, to be hit fair.

84

كالو غاجه هندق هندق گاديغن
كالو هريمو هندق هندق بلغن

Kalau gajah handak pandang gading-nya.

Kalau harimau handak pandang blang-nya.

If an elephant, he wants to examine its tusks. If a tiger, he wants to see its stripes.

Said of one who is ready for "anything from pitch and toss to manslaughter."

Blang بلغ has often been incorrectly translated 'piebald' whereas its true meaning is *striped*, 'piebald' being exactly rendered by the Malay work *Tompok* تومفق. FAVRE would seem to have missed the meaning of this latter word in this connection, as he translates it 'Pile, monceau, tas, amas, 'être en pile, en groupe' and renders *pie* (piebald) by *plang* بلغ, which is merely a corrupted form of *blang* بلغ

85 كاله منغ تا'تاهو برمورق بوله
Kalah menang ta'tahu ber-sorak bulih.

We know not whether we shall win or lose, but anyhow we can shout !

Said by those who anticipate a victory.

86 كچيل تا'بوله دمفكاكن اتق
بسر تا'بوله دمفكاكن باف
Kechil ta'bulih di-sangka-kan anak.
Besar ta'bulih di-sangka-kan bapa.

Small beasts should not be accounted young, nor large beasts be accounted parents.

Do not judge by appearances.

87 كسيهنكن راج دانس اوموغ
Kasih-an-kan raja di-atas usong.

To feel pity for a prince upon his litter.

Misplaced pity.

88

كفلا سام هيتم هائي ماميغ ۲

Kapala sama hitam hati masing-masing.

Our heads alike are black, but our hearts are different in each one of us.

89

كومبغ تامة ئيكور بوغا تامة سكاكي

Kumbang ta' s'ekor, bunga ta' s'kaki.

There is not only one beetle, and but a single flower.

"There are as good fish in the sea as ever came out of it."

90

كيم بهرو چافي فغايرة اورغ موده ليب كسمرغ

Kita baru chapai peng-ayuh orang sudah tiba ka-sebrang.

We have just taken up our paddles when others have already crossed the stream.

Said of one who is too late.

"To be asked for the wedding and arrive for the christening."

91

كيم مموا مائي تئافي قبور ماميغ ۲

Kita samoa mati tetapi kubor masing-masing.

We all alike die, but our graves are separate.

Men's lots are divided.

92

كيجيقي انق ملاك
 بوال انق منغكابو
 تيفوانق وميو
 بيدعه انق ترغكالو
 مومبوغ انق فهغ

Kéchék anak Malaka.

Bual anak Menangkabau.

Tipu anak Rembau.

Bidaäh anak Trengganu.

Sombong anak Pahang.

The men of Malacca wheedle; the men of Menangkabau draw the long bow; the men of Rembau cheat; the men of Trengganu lie; and arrogant are the men of Pahang!

The above proverbial saying hits off the most prominent characteristic of each nation of Malays mentioned.

93

کورغ نبق رڭکوه یغلبه

Kurang krat rengkoh yang lebeh.

The less we sever by cutting the more we break by pulling and twisting.

We cannot shirk our responsibilities, and if we put them aside in one form they will start up afresh in another.

Rengkoh رڭکوه = to break by pulling, twisting or bending. It has much the same meaning as that more common word *Lentor* لنتور.

94 غاجه همدق بیرق بسر کنچیل همدق بیرق بسر ایسق کببغ

*Gajah handak berak besar, kanchil handak berak besar
esok ka-bebang.*

The elephant passes huge excrement; the mouse-deer desires to do the same: in the end the latter will have a stoppage of the anus.

The frog and the bull.

Bebang بڭ = a stoppage of the anus, or of the womb in child-birth.

95

گھرو دباکر کمون بر باؤ

Gharu di-bakar kemenyan ber-bau.

Ligom aloes is burned and gum benzoin gives forth a sweet odour.

One man does the work and another reaps the benefit.

96

لمه لیبے کا یو اکر دلنتور بولہ دفالہ تا دافے

Lemah liat kayu akar di-lentok bulih di-patah ta'dapat.

Yielding yet tough like a root which can be bent but not broken.

Said of one who knows when to resist and when to give way.

97

لار لار ماکن داوفیہ لای فوئیہ لای دکروه

Lar-lar makan di-upih lagi puteh lagi di-kroh.

Insects eating the betel-palm sheath the whiter it grows the more turbid is the water which cleanses it.

98

لیدہ بر چابغ بائی بیاق

Lidah ber-chabang bagai biawak.

A tongue forked like that of an iguana.

Said of one who breaks faith with another.

99

ماکین مورہ ماکین منارار

Makin murah makin men-awar.

The lower the price, the more you cheapen it.

"Give an inch you take an ell."

See Proverb 40 of this collection.

100

مالو ماكن فروة لافر
مالو برکايوه فراهو تالاجو

Malu makan prot lapar.

Malu ber-kayuh prau ta'laju.

If one is shy to eat, one's belly will feel hunger. If one is ashamed to paddle, one's boat will not travel swiftly.

Indolence leads to poverty.

Compare Proverbs 168 and 173 in Mr. MAXWELL'S collection published in No. 2 of this Journal.

101

مغات دولغ فاکو مرفيه

Meng-ata dulang paku serpih.

The chipped nail abuses the tray.

This Proverb is best explained by the line which is usually tacked on to it:—

مغات اورغ اوق يغ لبه

Meng-ata orang awak yang lebeh.

When we abuse others we infer that we are better than they are.

102

منتاداره فد داغ

Minta darah pada daing.

To ask a dried fish for its blood.

"You cannot get blood from a stone."

103

مسيبون تانه يغ تيفكي مشگالي تانه يغ لكوق

Men-ambun tanah iang tinggi, meng-gali tanah iang lekok.

To heap up earth on a place where the ground is already high, and to dig where the earth is already indented.

Useless labour.

104

مننتي ارا تا برگنه

Me-nanti ara ta' ber-getah.

To wait for a fig which is devoid of milk (latex).

To wait endlessly.

105

مننتي فليدو كمبيغ ترفوتس

Me-nanti pelir kambing ter-putus.

To wait the severing of the goat's penis.

To wait for something which although it appears to be imminent, never really comes to pass.

106

مولوت ما نيسي مماتهنك تولغ

Mulut manis mem-atah-kan tulang.

Gentle words (*lit.*, a sweet mouth) break bones.

Men are deceived by flattery.

Compare Proverb 72 in this collection.

107

نامسي تا داينجين فينگن تا رتق

Nasi ta' dingin pinggan ta' retak.

If the rice be not cold the plate will not be cracked.

This expression is best explained by the following line which is often added to it:—

اديق فون تا ايفين ابغ فون تا هندق

Adek pun ta' ingin abang pun ta' handak.

If thou, oh sister mine, dost not desire it, I, thy brother, desire it not at all.

The words 'brother' and 'sister' when used in this manner of course really mean 'lover' and 'mistress.'

108 نايق ملومفت تورن ترجون
Naik me-lompat turun terjun.

To ascend leaping and come down with a plunge.
"Pride comes before a fall."

109 هارفكن توغكت ميباوا ربه
Harap-kan tongkat tongkat mem-bawa rebah.

To trust in a staff, and the staff helps one to a fall.
"To lean on a reed which breaks and pierces the hand."
This Proverb is usually applied in speaking of a *Raja* or Chief who has been deceived by his followers.

Compare:—

فاگر ماکن فالي
Pagar makan padi.

The fence eating the crop, &c.

110 هارفكن تينتيق امبون نغه هاري
Harap-kan titik ambun tengah hari.

To hope for drops of dew in the middle of the day.
To expect impossibilities.

111 هریمو مغاوم تا' منغكت كالو تا' مغاوم راجين منغكت
Harimau meng-aum ta' men-angkap kalau ta' meng-aum rajin men-angkap.

A tiger when it roars does not make a kill, when it is silent it often finds prey.

A boastful fellow is not usually brave, and *vice versâ*.

112

هندق کتینگى دفرلدهکن
هندق کرلده دفرلینگیکن

Handak ka-tinggi di-pe-rendah-kan.

Handak ka-rendah di-per-tinggi-kan.

When wanting height to make a thing low, and desiring lowness to heighten.

Foolish action which defeats its own ends.

113

هندق کشفخ دکرمه
هندق کفندی دهوبغ

Handak ka-panjang di-krat-krat.

Handak ka-pendek di-hubong-hubong.

Desiring length, to cut a thing short. Desiring shortness, to lengthen a thing by joining.

Compare the last preceding Proverb.

114

هیلغ ایکن ددالم کرابو

Hilang ikan di-dalam krabu.

To lose a fish in the preserved fish.

"Not to be able to see the town for the houses."

A BIBLIOGRAPHY OF MALAYA,*

FROM JULY, 1890, TO JUNE, 1891.

BY

C. DAVIES SHERBORN, F.G.S., F.Z.S.

In compiling this Bibliography, all sources of information have been utilized. In inserting, therefore, every publication that has come under his notice, the compiler hopes that the entries will prove of considerable assistance; but, as a large proportion of the literature of this district, either never reaches England at all, or else arrives so long after as to be too late for examination for this purpose, he begs the reader's indulgence for any error that may be present. His thanks are due to M. Martinus Nijhoff of The Hague for information as to some of the more recent books.

ABDUL KARIM TJIAT.—Sjair atau Panton akan djoedoh Perak deri Bangsawan Ambon, 1865-14 Maart-1890. 8vo. *Ambon*, 1891, 70 pp.

ACTA van het Zending-Congres, gehouden te Amsterdam op den 28, 29 en 30 Januari, 1890, waarin zijn opgenomen eene Bijdrage van W. van der Bergh: de Referaten van Kuijper, Schaafsma, Wijmalen, Cachet, Wagenaar, Fortman; en de Zendingrede van J. C. Sikkel. 8vo. *Amsterdam*, 1890, iv, 122 pp.

* By "Malaya" is here meant that part of the Archipelago enclosed in a line drawn round the North of Siam and the Philippines, though Macassar Strait between Lombok and Bali, round the outlying Islands of Java and Sumatra and to the East of Nicobar and Andaman Islands.

- AGASSIS.—L'île de Java. *Bull. Soc. Geogr., Lyon*, vol. ix, 1890, pp. 285-310.
- AGIUS, J. J.—Población de Filipinas. *Rev. Españ.*, vol. 133, 1891, pp. 75-93, 161-173.
- ALBRECHT, J. E.—Klapper op de Wetboeken in het Staatsblad van Nederlandsch-Indië, benevens op het Bijblad op dat Staatsblad. Jaarg. 1884-1889. 8vo. *Leiden*, 1890. viii, 195 pp.
- ALGEMEEN Verslag van den staat van het middelbaar en lager onderwijs voor Europeanen en met dezen gelijkgestelden in Nederlandsch-Indië over 1888. 8vo. *Batavia*, 1889, viii, 125 pp.
- 1890. 8vo. *Batavia*, 1890, viii, 164 pp.
- ALGEMEEN vijfjarig Verslag van het Inlandsch onderwijs in Nederlandsch-Indië loopende over de jaren 1883 t/m 1887. 4to. *Batavia*, 1890, x, 394 pp.
- ALMANAK.—Regeeringsalmanak voor Nederlandsch-Indië, 1890. 3 pts. 8vo. *Batavia*, 1890, xvi, 443: vii, 665: xxxii, 919 pp.
- Regeeringsalmanak voor Nederlandsch-Indië, 1891. 2 pts. 8vo. *Batavia*, 1891 (I,—xvi, 452, 668 pp.; II,—xxvi, 928 pp.).
- Javaansche Almanak voor 1891. 8vo. *Semarang* 1890, vi, 200, 160, 26 pp., pts.
- Javaansche Almanak (7 Jaargang). Met het portret van den Kroonprins van Djokjakarta. 8vo. *Djokja*, 1891, iv, 242, 112 pp., pts. [In Javaansche Karakteren.]
- Dengan tafel pematjaan elkitab jang tjoendoek sasaharj doewa nats, bage tahun 1890. 8vo. *Batavia*, 1889, 12 pp.
- 1891. 8vo. *Batavia*, 1890, 12 pp.
- Maleische Almanak (14^e Jaargang). 8vo. *Djokjakarta*, 1890, iv, 230, 32, 42 pp., portrait.
- Maleische Almanak (15^e Jaargang). Met het portret van den Kroonprins van Djokjakarta. 8vo. *Djokjakarta*, 1891, iv, 283, vi pp.

- ALMANAK.—(Takwin) atau hitoengan Hari, Boelan, Tahoen Orang Mesehi 1891, Tahoen Orang Islam 1308-1309, Tahoen Orang Tjina, Kong-si, 16-17. 8vo. *Batavia*, 1890, 40 pp.
- ANDERSON, J.—The Selungs of the Mergui Archipelago. 8vo. *London*, 1890, 47 pp.
- ANON.—Blokade van *Atjeh*. *Ind. Gids*, vol. xiii, 1891, pp. 864, 865.
- De *Atjeh*-zaak in verband met de laatste gebeurtenissen. *Ind. Gids*, vol. xii, 18—, pp. 1-27.
- Resultaten van proeven met een *ballon* captif in *Atjeh*. *Ind. Gids*, vol. xiii, 1891, pp. 150-152.
- De productie van *Billiton-tin* in de laatste jaren. *Ind. Gids*, vol. xiii, 1891, pp. 691, 692.
- Een Engelsch geestelijke over *Bornei's* Westkust. *Tijdschr. Ned. Ind.*, 1891, pp. 174-179.
- Ceram. *Ned. Zendingsts*, vol. ii, 1890, pp. 305-320.
- Petitie van *Chineezzen* te Singapore tegen den opiumverkoop. *Ind. Gids*, vol. xiii, 1891, p. 954.
- Chineesche begrafenisfonds merk Hong San Sie Boekoe peratoeran mengoempoelken oewang boeat menoeloeng atas kematian orang-orang Tjina di *Batawi*. 8vo. *Batavia*, 1890, 22 pp.
- De Chineesche geheime eedgenootschappen. *Tijdschr. Ned.-Ind.*, 1890, pp. 57-63.
- Een inlandsch *Christen* in Moeite. *Ind. Gids*, vol. xiii, 1891, pp. 856, 857.
- Chronicle and Directory for China, Japan, Straits Settlements, Indo-China, Philippines, etc., 1890. 28th year. *Hongkong*, 1890.
- De werking der Indische *Comptabiliteitswet*. *Ind. Gids*, vol. xiii, 1891, pp. 857-863.
- Voorschriften betreffende de *Comptabiliteit* in Nederlandsch-Indië. 8vo. *Batavia*, 1889, iv. 145 pp.

- ANON.—De heer Cremer over *Decentralisatie* in Indië. *Ind. Gids*, vol. xiii, 1891, pp. 237-243.
- Een discussie in de Eerste Kamer over het *Defensieplan* voor Ned.-Indië. *Ind. Gids*, vol. xiii, 1891, pp. 595-599.
- Een Keizerlijke optocht in *Djokjokarta*. *Eigen Haard*, 1890, p. 278.
- Zedekundige *Fabelen* naar't Maleisch Boenga Rampaj. [Javaansch.] 8vo. *Soerabaia*, 1890, 66 pp.
- Aanteekening over eene Javaansche welriekende *Grassoort*. *Teysmannia*, Jaarg. 1, 1890, pt. 9.
- Statistiek van den *Handel*, de scheepvaart en de in en uitvoerrechten in Nederlandsch-Indië over het jaar 1889. 8vo. *Batavia*, 1891, 8, lii, 149 pp.
- Handel* van Ned.-Indië met Genua en China. *Ind. Gids*, vol. xiii, 1891, pp. 683, 684.
- Handel* tusschen Ned.-Indië en Groot-Britannië. *Ind. Gids*, vol. xiii, 1891, pp. 480-482.
- Handel* van Ned.-Indië met Boston. *Ind. Gids*, vol. xiii, 1891, p. 1112.
- De *Handel* in Suiker te Glasgow in 1890. *Ind. Gids*, vol. xiii, 1891, pp. 940-951.
- Het eigendunkelijke van onze *behandeling* der inlandsche vorsten. *Ind. Gids*, vol. xiii, 1891, pp. 73-77.
- Hygiënische* Maatregelen. *Ind. Gids*, vol. xiii, 1891, pp. 875-878.
- “Jasa” in Westen in Midden-en Oost-*Java*. *Ind. Gids*, vol. xiii, 1891, pp. 324, 325.
- Een deputatus synodi voor Midden-*Java*. *Ind. Gids*, vol. xiii, 1891, p. 419, 420.
- De sterkte van den veestapel op *Java* en Madoera. *Ind. Gids*, vol. xiii, 1891, pp. 601, 602.
- De verdediging van *Java*. *Ind. Gids*, vol. xiii, 1891, pp. 82-87.

- ANON.—Waardeert de *Javaan* ons bestuur? *Ind. Gids*, vol. xiii, 1891, pp. 249, 250.
- [Geslachtslijst van den eersten Sultan van *Jogja* tot den zesden door Raden Mas Aria Nata Asmara II.] 8vo. *Jogjakarta*, 1890, 60 pp. [In Javanese.]
- In de Kamer over heeren-en désadiensten. *Ind. Gids*, vol. xiii, 1891, pp. 87-95.
- Toestand van de *Kerk* in Oost-Indië in 1644. *Ned. Zendingsts*, vol. ii, 1890, pp. 208-253.
- De ordonnante betreffende het verkoopen van gemalen *Koffie*. *Ind. Gids*, vol. xiii, 1891, pp. 79-81.
- De oud-Hoofdinspecteur Heyting contra de Staatscommissie voor de *Koffiecultuur*. *Ind. Gids*, vol. xii, 1890, pp. 1942-1975.
- Officiële Mededeelingen omtrent de Gouvernements-*Koffiecultuur* op Jaav. *Ind. Gids*, vol. xiii, 1891, pp. 580-593.
- Uit het *Koloniaal* Verslag van 1890. I. Christelijke Godsdienst. II. Mohammedaansche Godsdienst. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 252-274.
- De opkomst van het Nederlandsch gezag in Oost-Indië. Verzameling van onuitgegeven stukken uit het *oud-Koloniaal* Archief. 2 Bks. Uitg. door Dr P. A. Tiele. Deel 2, bew-door J. E. Heeres. 8vo. *s'Gravenhage*, 1890, xii, lxxxiv, 396, xix pp.
- Lichtenlijst van het Koninkrijk der Nederlanden en de *Koloniën*. 4to. *s'Gravenhage*, 1891, 88 pp.
- Uit de *Koloniale* begrooting voor 1891. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 441-444.
- Landbouw* op verhuurde Landen in Solo en Djokja. *Ind. Gids*, vol. xiii, 1891, pp. 688-691.
- Tegen een *landrente-kadaster*. *Ind. Gids*, vol. xiii, 1891, pp. 78, 79.
- Inlichtingen van den Minister betreffende de *Ma-doereesche* Grooten. *Ind. Gids*, vol. xiii, 1891, pp. 609-611.

- ANON.—Verordeningen en keuren voor de residentie *Madioen*, verzameld en in 't Maleisch overgebracht door *T. A. Winátá*. 8vo. *Batavia*, 1889, iv, 56 pp.
- Observations made at the *Magnetical* and Meteorological Observatory at Batavia. vol. xii, 1889. 4to. *Batavia*, 1890, 10, 244 pp.
- De debatten in de Kamer over de *Mailcontracten*. *Ind. Gids*, vol. xiii, 1891, pp. 403-419.
- Verslag van het *Mijnwezen* in Nederlandsch-Indië over het jaar 1888-1889. *Jaarb. Mijnw. Ned. O. I.*, vol. xix, 1890, Techn. Ged. pp. 143-191.
- De generale staf en de voorbereiding van *Militaire Expeditiën* in Indië. *Ind. Gids*, vol. xiii, 1891, pp. 503-523.
- Een bezoek aan *Modjo-Warno*. *Ned. Zendingsts*, vol. ii, 1890, pp. 262-265.
- Mohamedaansch-godsdiensstige* broederschappen. Door een oud-Oost-Indisch ambtenaar. 8vo. *Batavia*, 1890.
- Staatsblad van *Nederlandsch-Indië* over het jaar 1889. 8vo. *Batavia*, 1890.
- Het Reglement op het *Notarisambt* in Nederlandsch Oost-Indië benevens het tarief van het honorarium toegelicht door P. Vellema. 8vo. *s'Gravenhage*, 1890 217 pp.
- De financiëele resultaten der *Opiumrègie* in Cochin China van 1882 tot 1890. *Ind. Gids*, vol. xiii, 1891, p. 952.
- Opiumdebiet* en *Opiumprijzen* op Java en Madoera *Ind. Gids*, vol. xiii, 1891, pp. 1112-1114.
- Opiumconsumptie* en *Opiumprijzen* in Cochin-China en op Java. *Ind. Gids*, vol. xiii, 1891, pp. 952, 953.
- Uit het rapport van den heer Groeneveldt over het *Opiummonopolie* in Fransch Indo-China. *Ind. Gids* vol. xiii, 1891, pp. 694-724. [Deals with Ned.-Indië.]

- ANON.—*Overeenkomsten* met Sintang, Tajan-Meliouw en Boeloengan. *Ind. Gids*, vol. xiii, 1891, pp. 846-853.
- Overeenkomst* met Pasir. *Ind. Gids*, vol. xiii, 1891, pp. 1053-1064.
- Overeenkomst* met Assahan en verklaringen, afgelegd door bestuurders in de Residentie Oostkust van Sumatra met betrekking tot het haven beheer, enz. *Ind. Gids*, vol. xiii, 1891, pp. 570-580.
- Overeenkomsten* met Malaboet, Tandjoeng Kassin, Siantar en Rokan Ampat Kotta. *Ind. Gids*, vol. xiii, 1891, pp. 395-401.
- Pacht Apioen Atoeran* Baroe reperti di temtoeken dingen Oendang Oendang tanggal 14 Juli 1890, Staatsblad No. 149. 8vo. *Soerabaia*, 1890, 24 pp.
- Het onderwijs op de *Philippijnen*. *Ind. Gids*, vol. xiii, 1891, pp. 325-327.
- De begrooting voor openbare werken in de practijk. *Ind. Gids*, vol. xiii, 1891, pp. 867-871.
- Het eiland *Randjoewa*. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 332-334.
- Geen *Repeteergeweer* voor Indië. *Ind. Gids*, vol. xiii, 1891, pp. 246-247.
- Een nieuwe bewerking van de *Sěrat Aji Saka*. *Tijdschr. taal-land-volk. N. I.*, vol. xxxiv, 1890, p. 104.
- Een Amerikaansche gedachte over een *Spoorweg* van Djember naar Banjoewangi. *Ind. Gids*, vol. xiii, 1891, pp. 81, 82.
- Een nieuwe lezing omtrent de quaestie der *Spoorwegtarieven*. *Ind. Gids*, vol. xiii, 1891, pp. 865-867.
- Een plan voor Rentegarantie ten behoeve van vier nieuwe *Spoorwegen*. *Ind. Gids*, vol. xiii, 1891, pp. 593-595.
- Verslag over den aanleg en de exploitatie van de *Staatsspoorwegen* in Nederlandsch-Indië over het jaar 1889. 4to. *Batavia*, 1891, 300 pp. 3 pts.

- ANON.—*Suiker* productie per bouw in de Gouvernementslanden van Java. *Ind. Gids*, vol. xiii, 1891, pp. 954-956.
- Over het gewicht en de uitgebreidheid van het wortelstelsel bij het *Suikerriet*. *Ind. Gids*, vol. xiii, 1891, pp. 148-150.
- Bilder aus *Sumatra*. *Schorer's Familienblatt*, Heft 10, 1891.
- Tocht naar het rijk van Poeloe-Lawan (Sumatra's Oostkust). Onderzoek van het stroomgebied der Bottang-Nila. *Tijdschr. Binnenl. Bestuur*, vol. iii, 1890, p. 129.
- Wat de particuliere industrie op *Sumatra's* Oostkust aan de Schatkist kost en wat zij voor de Schatkist opbrengt. *Ind. Gids*, vol. xiii, 1891, p. 77.
- Tabaks* ondernemingen op de Oostkust van Sumatra, Langkat, Deli, Serdang, Bedagei, Padang, Batoe-Bahara, Asahan, Kwaloe, Bilah, Panei en Laboean-Batoe. 4to. Amsterdam, 1891, 18 pp.
- Eindelijk een parc *vaccinogène* op Java. *Ind. Gids*, vol. xiii, 1891, pp. 871-875.
- Reglement voor het *Weduwen* en *Wezenfonds* voor de Officiëren der landmacht in Nederlandsch-Indië. Bygewerkt tot ultimo 1890 van toelichtende nooten voorzien door C. H. W. Boers. 8vo. *s'Gravenhage*, 1891, 34 pp.
- Gezondheidstoestand bij de *Zeemacht* in Indië. *Ind. Gids*, vol. xiii, 1891, p. 72.
- De *Zending* in ons Parlement. *Ned. Zendingsts*, vol. ii, 1890, pp. 278-311; 335-379.
- Maatregelen in het belang der *Zending*. *Ind. Gids*, vol. xiii, 1891, pp. 854-856.
- Een gewraakte uitgave in zake de serch. *Ind. Gids*, vol. xiii, 1891, pp. 69-71.
- Archief voor de geschiedenis der oud-Hollandsche *Zending*. V. de Molukken 1603-1624. 8vo. *Utrecht*, 1890, viii, 248 pp.

- ARNTZENIUS, G.—Cultuur en volk. Beschouwingen over de Gouvernements koffiecultuur op Java. 8vo. *Gravenhage*, 1891, 4, 159 pp.
- ATOERAN tanah Mardika disebelah koelen Kali Tjimanoeck ja'itoe tanah-tanah jang ada didalem Residentie Banten, Betawi, Krawang dan Priangan. Staatsblad tahun 1836 No. 19... 1854, No. 1.....1827, No. III. 8vo. *Batavia*, 1890, 96 pp.
- BABAD—Peladjarran, No. 4. 8vo. *Samarang*, 1890.
- BAESSLER, A.—Reisen im Malayischen Archipel. *Zeitschr. Ethnol.*, vol. xxii, 1891, pp. 493-500.
- Ethnographische Beiträge zur Kenntniss des Oostindischen Archipels, I-V. *Intern. Arch. Ethnogr.*, vol. iv, 1891, pp. 66-83, pls. iv-ix.
- Vocabular von Negrito-Worten. *Zeitschr. Ethnol.*, vol. xxii, 1891, p. 500.
- Ethnographische Beiträge zur Kenntniss des Ost-Indischen Archipels. 8vo. *Leiden*, 1891, 6 pls. and figs.
- BAHWA INI HIKAJAT DJAHIDIN.—Terkarang pada bahasa Wolanda oleh J. A. Uitkens dan dikarangken pada bahasa Melajoe oleh A. F. von de Wall. 8vo. *Betawi*, 1890, 80 pp.
- BALEN, J. H. VAN.—De Nederlanders in Oost en West, te water en te land. Het geheim van mijnheer Tukker. De opstand van het Kakihan-verbond op Ceram in 1860. Indische zeeroovers. 8vo. *Amsterdam*, 1891.
- De kroon van Mataram. 8vo. *Amsterdam*, 1890, 199 pp. 12 pls.
- BARTELS, M.—Das javanische Modell einis Wajang-spiel. *Zeitschr. Ethno.*, vol. xxii, 1890, pp. 266-270.
- BEAUREGARD, G. M. O.—Proverbs et dictons malays, III *Rev. tradit. pop.*, vol. v, 1890, pp. 722-724.
- BECCARI, O.—Phoenix, etc. *Malesia*, vol. iii. pp. 353-384 (Feb.); 385-432 (Mar.) 1890 [completing the publication].

- BENKO, J. V.—Das Datum der Philippinen. *Mith. Geogr. Ges. Wien*, vol. xxxiii, 1890, pp. 560-576.
- "BENEDICTUS."—Les musiques bizarres à l'Exposition recueillies et transcrites par Benedictus. Dessins de F. H. Gorguet. 8vo. *Paris*, 1890, iv, 87 pp. [Le Gamelang Javanais.]
- BIBÉLI, O.—Ma bi dëdemmo. Bijbelsche geschiedenissen in de Galillareesche taal. 8vo. *Utrecht*, 1889, iv 416 pp.
- BIEGER, P.—Staat der gemeenten in Kediri-Madioen op ultimo 1889. *Med. Ned. Zendel*, vol. xxxiv, 1890, p. 390.
- Bezoeken op Soemba. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 1-30.
- Een Doodenfeest te Rendé, op Soemba, *ibid.*, pp. 151-161.
- BIET.—Vocabulaire Iysson. *Mém. Soc. Acad. Indo-Chin.* vol. i, 1879 (1890), pp. 21-42.
- BINK, G. L.—Lijst van telwoorden en eeinge zelfstandige naamwoorden enz. [Wandammen Baäi]. *Tijdschr. Ind. taal-land-volk.*, vol. xxxiv, 1890, pp. 62-71.
- BLANCHARD, E.—Les preuves de la dislocation de l'extrémité sud-est du continent asiatique pendant l'âge moderne de la terre. *Comptes Rendus Acad. Sci. Paris*, vol. cx, 1890, pp. 369-373.
- BLOMMEN [?VAN BLOEMEN WAANDERS].—Indische Schetsen, No. vii, viii. 8vo. *Gravenhage*, 1890, pp. 149-193.
- BLUMENTRITT, F.—Vademecum etnográfico de Filipinas. *Bol. Soc. Geogr. Madrid*, vol. xxvii, pp. 246-271.
- Uber de Eingeborenen der Insel Palawan und der Inselgruppe der Calamianen. *Globus*, vol. lix, 1891, pp. 166-184.
- Die Kianganen (Luzon). *Ausland*, 1891, pp. 118-129.
- Die Calingas. *Ausland*, 1891, p. 328.
- Beiträge zur Kenntniss der Mindayas (Mindanao) *Mith. Geogr. Ges. Wien*, vol. xxxiii, pp. 232-243.

- BLUMENTRITT, F.—Die Tiruray der Insel Mindanao. *Globus*, vol. lviii, 1890, pp. 129-131.
- Alphabetisches Verzeichnis der eingebornen Hämme der Philippinen. *Zeitschr, Ges. Grok. Berlin*, vol. xxv, 1890, pp. 127-146.
- Las razas del archipélago filipino. *Bol. Soc. Geogr. Madrid*, vol. xxviii, 1890, pp. 7-42, map.
- Die Subanos, Mindanao. *Ausland*, 1890, pp. 392-395.
- Das Stromgebiet des Rio Grande de Mindanao. *Petermann's Mitth.*, vol. xxxvii, 1891, pp. 108-113, map, 1:1,000,000.
- BOEKOE OBAT dan Katoeronggo Koeda ija itoe menerangkan tjatjat atawa baiknja koeda, kentara darie boeloe atawa tandanja bernama Mathie, atawa oeijeng oeijengan, bagimana tjaritanja orang doeloe kala. 8vo. *Djokja*, 1890, 56 pp.
- BOEKOE PANTOEN dari Tjeritanja Nonah Asschepoetster tersalin bahasa Melajoe Terkarang oleh H. 8vo. *Soerabaia*, 1890, 32 pp.
- BOEKOE SAIRAN dari Tjeritanja Samphek, Eng Taij tersalin darie Boekoe Tjina. Terkarang oleh The. T. L. Aken goenanja bekin ingetan anak moeda djaman sekarang. 8vo. *Soerabaia*, 1890, 116 pp.
- BOEKOE TJERITA-AN Hikajatuja Baron van Munchausen tersalin bahasa Melajoe. 8vo. *Soerabaia*, 1890, 86 pp.
- Tanah Djawa. Bagian ka I. 8vo. *Soerabaia*, 1890. 91 pp.
- BOEKOE TJERITA Doeloe kala di negari Tjiena, Menjeritakan Keiser Lie Sie Bient-Ijoe Tee Hoe. Tersalin dari boekoe Tjina "See Ijoe" oleh Boen Sing Hoo. II. 8vo. *Samarang*, 1890, 102 pp.
- BOCK, C.—De Kontei à Bandjirmasin. *Tour de Monde*, 1890, pp. 337-368.

- BOEKOE ELMOE PERITOENGAN Tersalin dariê boekoe tjina doeloe kala Yak Kieng Ho Thian Ik So. Njang amat bergoena sekalie aken memboewat segala pertanjakan. 4to. *Soerabaja*, 1890, IV, 128 pp.
- BOEKOE Ka-adilan hoekoeman atas bangsa djawa dan sebrang di Hindia Nederland. Menoeroet salinan jang telah di oendang-oendang ken oleh Kandjeng Goebnernement, dengan di pindahken kapada hoeroef Arab. 8vo. *Batavia*, 1890, 184 pp.
- BOEKOE KITAB TJAP Dji YA ILMOE MELIATIN dari Radja Seitan Thio Tian Soe. Tersalin dari boekoe Tjina oleh O. H. T. & Y. P. S., Soekaboemi. 8vo. *Batavia*, 1890, 36 pp.
- BOEKOE (KOA H MIEA) petang-petangan orang Tjina tersalin dari boekoe TJAP DJIE SHIO terkarang oleh L. S. Y. & T. H. L. 8vo. *Batavia*, 1890, 80 pp.
- BOER, H. B. DE.—Beschrijving van het Stroomgebied van de Rivieren in het Rijk van Indragiri en de naburige kust landen. *Tijdschr. Aardr. Genootsch.* ser. 2, vol. vii, 1890, pp. 557-576, 3 maps.
- B [OERLAGE], J. G.—Ternataansche plantennamen. *Teysmannia*, Jaarg 2, 1891.
- BORNEO.—Zum Klima von Nord Borneo, Sandakan, 1888. *Met. Zeitschr.*, vol. vii, 1890, p. 119.
- BOSSE, VAN.—De hervorming van het rechtswezen op Sumatra's West-kust. *Batav. Handelsblad*, 1890.
- BOINLLEVAUX,—Ma visite anx ruines cambodgiennes en 1850, *Mem. Sor. acad. indo-chin.*, vol. i, 1879, [1890], pp. 1-18.
- BON KON SIN TJIONG,—Satoe Pladjaran Baik bagei Orang Lelaki dan Prampoean tersalin dari Roepa-roepa kitab tjina dan ti tambain dengen swatoe pringetan. Di KARANG OLEH OEY TJOEN BIN. 8vo *Batavia*, 1890, 66 pp.
- BOULENGER, G. A.—Remarks on the Herpetological Fauna of Mount Kina Baloo, North Borneo. *Annals Mag. Nat. Hist.* [6] vol. vii, 1891, pp. 341-345.

- BOULENGER G. A.—List of Reptiles, Batrachians and fresh-water fish collected by Prof. Moesch and Mr. Iverson, in the district of Deli, Sumatra *Proc.Zool. Soc.* 1891, pp. 31-40
- BOUMAN AND ROESSINGH VAN ITERSON.—De Indische comptabiliteitswet. *Ind-Gids*, vol. xiii, 1891, pp. 40-68.
- BRAAM-MORRIS, D. F. VAN.—Nota van toelichting behoorende bij het contract gesloten met het Landschap Bima. *Tijdschr. taal-land-volk*. N. I. vol. xxxiv, 1890., pp. 176-233.
- BRAU DE SAINT-POL-LIAS, H.—La Côte de poivre. Voyage à Sumatra. 8vo. *Paris*, 1891, vi, 240, pp. gigs.
-
- Les Orangs-Atché. *Nouv. Revue*, 15 Juli 1890.
-
- La Côte des douze colonies. Sumatra. *Revue Franç.*, 1890, xi, pp. 129-137.
- BRENNER-FELSACH, F. S.—Reise durch die unobhändigen Battahland und auf der Insel Nias. *Mitt. k.-k. Geogr. Soc. Wien*, 1890, xxxiii, pp. 276-306 map 1 : 200,000.
- BRONS MIDDEL, R.—Losse schetsen over Nederlandsch Oost-Indië. 8vo. *Groningen*, 1890, 496 pp.
- BRUINSMA, J. F. D.—Onze versterking in het landschap Edi en de krijgsverrichtingen aldaar gedurende de maanden April en Mei 1889. 8vo. *Sneek*, 1890, 52 pp. 3 kaarten.
-
- De verovering van Atjeh's groote Missigit. 8vo. *Sneek*, 1889, 4 pl.; 3 portr. 1 kaart.
- BULÉE.—Van kwaad tot erger. *Ind. Gids*, vol. xiii, 1891, pp. 258-262.
- BURCK, W.—Opmerkingen over de onder den naam van *Erythroxylon coca*, in Nederlandsch-Indië gecultiveerde gewassen. *Teysmannia*, Jaarg. 1, 1890, pp.
- BUYS, M.—Depok. *Ind. Gids*, vol. xii, 1890, pp. 1239-1248.
- CASTILLO, R. DE.—Gran Diccionario Geográfico, estadístico é histórico de España e sus provincias de Cuba Puerto Rico, Filipinas y posesiones de Afrika vol. i. 8vo. *Barcelona*, 1890, 720 pp.

- ALPHABETISCHE CATALOGUS van de Bibliotheek der Soekaboemische Landbouw-Vereeniging. Jaarg. 1. 8vo. *Batavia*, 1890, 51 pp.
- CATALOGUS van hetgeen verkrijgbaar is bij's Landsdepôt van leermiddelen te Weltevreden of voor Gouvernements inlandsche scholen aangevraagd kan worden ter verstreking van Batavia. 8vo. *Batavia*, 1890, 44 pp.
- van de Bibliotheek van het Hoofdbureau van den Generalen Staf. 8vo. *Batavia*, 1890, iv, 77 pp.
- CLERCO, F. S. A. DE.—Bijdragen tot de kennis der Residentie Ternate. 8vo. *Leiden*, 1890, xx, 359 pp.
- Wederzijdsche bezoeken van den Sultan en den Resident van Ternate. *Ind. Gids*, vol. xii, 1890, pp. 2611-2616.
- Moet Nieuw-Guinea een tweede Atjeh worden. *Ind. Gids*, vol. xii, 1890, pp. 1682-1694.
- COBLIJN, W. A.—De verdediging van Java. 8vo. *s' Gravenhage*, 1890, ii, 185 pp.
- COOL, W.—De opleiding en hogere vorming van den officier der landmacht hier te lande en voor Indië. 8vo.'s. *Gravhage*, 1890, 84 pp.
- COOTS, A.—Notes on the recent developement, explorations, and commercial Geography of British North Borneo. *Four. Manchester Geogr.* 8vo. 1890, pp. 63-76.
- CORDES, J. W. H.—In Pasoeroeans bergen. *Ind. Gids*, vol. xii, 1890, pp. 216-228.
- In Preanger Regentschappen, I. *Ind. Gids*, vol. xii, 1890, pp. 2106-2124.
- In de Preanger Regentschappen, II. III. *Ind. Gids*, vol. xii, 1890, pp. 2372-2378; 2630-2648.
- CREMER, J. T.—Oelische schetsen. *Eigen Haard*, 1890, pp. 564, 587, 675.
- CROIZIER, DE.—Le Cambodge d'après les historiens chinois. *Mém. Soc. Acad. Indo-Chin.*, vol. i, 1879 [1890], pp. 163-174.

- CROIZIER, DE.—Historie de l'architecture Cambodgienne. *Mém. Sor. Acad. Indo-Chin.*, vol. i, 1879 [1890], pp. 85-108.
- Notice des manuscrits siamois de la Bibliothèque nationale: *Mém. Soc. Acad. Indo-Chin.*, vol. i, 1879, [1890], pp. 213-270.
- Les monumens de l'ancien Cambodge. *Mém. Soc. Acad. Indo-Chin.*, vol. i, 1879 [1890], pp. 273-289.
- CROZET'S, voyage to Tasmania, New Zealand, the Ladrone Islands, and the Philippines in the years 1771-1772. Transl. and ed. by H. L. Roth. 8vo. *London*, 1891.
- DAUM, P. A.—"Waarom de Minister Keuchenins viel." 8vo. *Batavia*, 1890, ii, 16 pp.
- DAVISON, W.—Journal of a trip to Pahang. *Jour. Straits Br. R. A. S.*; 1890, No. 20, pp. 83-91.
- DEDEM, W. K. VAN.—De agrarische kwestie of Java. *Ind. Gids*, vol. xii, 1890, pp. 2125-2170.
- DEKKER, V.—*Multatuli*.
- DE LOS REYES Y FLORENTINO, J.—Folk-Lore Filipino, vol. ii, 16mo. *Manila*, 1890, 300 pp.
- DE SAIR MADI KENTJANA.—Vertaald door C. den Hamer. 8vo. *Batavia*, 1890, 34 pp. [Reprinted from *Tijdschr. taal-land-volk.*, vol. xxxiii.]
- DEVENTER, M. L. VAN.—Daendels--Raffles. *Ind. Gids*, vol. xiii, 1891, pp. 341, 524 and 818.
- Het Nederlandsch gezag over Java en onderhoorigheden sedert 1811. Deel I. 1811—1820. 8vo. *s'Gravenhage*, 1891, ccx, 349 pp.
- DHARMA,—De Koffiekultuur als Volkscultuur. *Ind. Gids*, vol. xii, 1890, pp. 889-893.
- DIJK, P. VAN.—Omsmelting en zuivering van drie partijen Banka-Tin aan het Koloniaal Etablissement te Amsterdam. *Jaarb. Mijnw. Ned O. I.* vol. xix, 1890, Techn. Geo., pp. v-xxiii.
- DOBERCK, W.—Mittlerer Luftdruck zu Iloilo, Philippineir. *Met. Zeitshi.*, vol. vi, 1889, p. 156.

- DOEDES, N. D.—De Nederlanders. Een schets onzer vaderlandsche en koloniale geschiedenis. 8vo. *Leeuwarden*, 1890, iv, 339 pp.
- DONGENG—Dongeng noc Aranch, Sawareh na di karang, Saraweh deni di pindahkeun tina basa Malajoe kana basa Soenda koe Raden Hadji Moehamad Moesa, Hoofd Panghoeloe Kabupaten Limbangan. 8vo. *Betawi*, 1890, 71 pp.
- ELOUT VAN SOETERWOUDE, W.—De opiumvloek op Java. 4to. *s'Gravenhage*, 1890, 85, 8, 13, 4 pp.
- EMANTS, G.—Het Indische conversie vraagstuk. *Vrag. tijds.*, vol. xvii, 1891, pp. 1-15.
- ENGELBRECHT, W. A.—De Nederlandsch-Indische Wetboeken. De Grondwet van het Koninkrijk der Nederlanden, het Reglement op het beleid der regeering van Nederlandsch-Indië en andere wetten, wettelijke verordeningen en besluiten, met verwijzing naar de op elk artikel betrekking hebbende Nederlandsch-Indische, Nenderlandsche en Fransche wetsbepalingen. 8vo. *Semarang*, 1890, xxxiii, 1788 pp.
- ERNST,—Afschaffing of afkoopbaarstelling der heerendiensten op Java. *Ind. Gids*, vol. xii, 1890, pp. 1434-1462.
- ERNST, A.—Das Gold-und Silbererz-Vorkommen von Tambang-Salida auf Sumatras westküste. 8vo. *Freiburg i/s.*, 1890, 15 pp., map.
- ESCHE, J. M.—Een en ander naar aanleiding der onttrekking van koffie aan de verplichte levering. *Ind. Gids*, vol. xiii, 1891, pp. 184-219.
- ESSER, J. P.—Bijdrage tot de geschiedenis der zending in den N. I. archipel 1816-1875. *Ned. Zendingsts*, vol. ii, 1890, pp. 138-164.
- "EXCAVATOR"—De generale staf en the voorbereiding van militaire expeditiën in Indië. *Ind. Gids*, vol. xiii, 1891, pp. 770-791.
- FÉER.—Le bouddhisme à Siam. *Mém. Soc. Acad. Indo-Chin.*, vol. i, 1879 [1890], pp. 146-163.

FÉER.—Notice des manuscrits birmans et des manuscrits cambodgiens. *Mém. Soc. Acad. Indo-Chin.*, vol. i, 1879 [1890], pp. 189-198.

———Les nouveaux caractères cambodgiens de l'Imprimerie nationale. *Mém. Soc. Acad. Indo-Chin.*, vol. i, 1879 [1890], pp. 270-273.

FENNEMA, R.—Rapport omtrent het voorkomen van petroleum in Beneden-Langkat. *Zaarb. mijnw. ned. O. I.*, vol. xix, 1890, Verh. Techn. Gil., pp. 10-91.

FIGIE, S.—Uitkomsten van Meteorologische Waarnemingen in Nederlandsch-Indië gedurende Januari, Februari, Maart, April, Mei en Juni van 1889. *Nat. Tijdschr. Ned.-Ind.*, vol. xlix, 1890, pp. 443-500.

———and H. ONNEN.—Vulkanische verschijnselen en aardbevingen in den O. I. Archipel waargenomen gedurende het jaar 1888. *Nat. Tijdschr. Ned. Ind.*, yearly.

FINANCE.—Particulars of the following will be found in *Ind. Gids.* month by month:—Financiëel Overzicht; Prijzen van Indische Fondsen; Wisselkoersen; Javasche Bank; Koffie Statistiek; Specerijen: Tin, &c., by *E. Rittner Bos*.

FLANMARION, C.—L'éruption der Krakatoa. 16mo. *Paris*, 1890.

FOKKER, A. A.—Volledige leercursus om de Javaansche taal te leeren. Brief III-V. 8vo. *Zutphen*, 1890 (pp. 33-80).

FOORE, ANNIE—Bogoriana. Roman uit Indie. 8vo. *Haarlem*, 1890, viii, 414 pp., portrait.

FORTINI-SANTARELLI, S.—Un viaggio a Nias. *Rass. Nazion.*, 1890, Feb. 16.

FRANCIS, W.—Is het noodig, dat er in N. I. een afzonderlijk strafwetboek voor Europeanen en een afzonderlijk strafwetboek voor inlanders bestaan. *Ind.-Gids.*, vol. xii, 1890, pp. 421-436.

GAHAN, C. J.—Notes on Longicorn Coleoptera of the group *Cerambycinæ* with descriptions of new Genera and Species. *Annals Mag. Nat. Hist.*, [6]. vol. vi, 1890, pp. 247-261, and vol. vii, 1891, pp. 19-35.

- GARCIN DE TASSY.—La langue hindoustanie en Indo-Chine, *Mém. Soc. Acad. Indo-Chin.*, vol. i, 1875 [1890], pp. 18.
- GELDER, W. VAN.—Uit Indië de aarde rond. Aardrijkskundig leerboekje. 1^o. d. Nederlandsch-Oost-Indië. Ed. 3. 8vo. *Groningen*, 1890.
- Mangle, nja eta roeparoepa tjarita rendeung tjonto pikeun pangkat kadøewa di sakola Soenda. 8vo. *Leiden*, 1890, iv, 66, ii pp.
- Uit Indië de aarde rond. (Aadr. leerboek.) I. Ned. Oost-Indië. Ed. 3, 8vo. *Groningen*, 1890, vii, 78 pp.
- GERTH VAN WIJK, D.—Spraa kleer der Maleische taal. 8vo. *Batavia*, 1890, xi, 304, vi pp.
- GODWIN-AUSTEN, H. H.—On supposed new Species of Land-Mollusca from Borneo belonging to the Genera, *Opisthostoma* and *Diplommatina*. *Annals Mag. Nat. Hist.* [6] vol. vi, 1890, pp. 244-246, pl. vii.
- On a Collection of Land Shells made in Borneo by Mr. A. Everett, with description of supposed new species. Pt. 2. Zonitidæ and Helicidæ. *Proc. Zoot-Soc.*, 1891, pp. 22-47, pls. ii-iv.
- HOEBELK.—eber Javanische Lebermoose. *Ann. Jard. Botan. Buitenzorg*, vol. ix, pt. 1, 1890, pp. 1-40.
- GOLTSTEIN, W. VAN.—Het doodvonnis der Gouvernementskoffiecultuur. *Ind. Gids*, vol. xii, 1890, pp. 1249-1268.
- GORKUM, K. W. VAN.—De Oost-Indische Cultures, in betrekking tot Handel en Nijverheid. 8vo. *Amsterdam*, 1890.
- GORKOM, K. W.—De Oost-Indische Cultuurs in betrekking tot Handel en Nijverheid. Supplement. 8vo. *Amsterdam*, 1890, viii, 304 pp.
- GRAAFLAND, N.—De Ambtenaar bij het Binnenlandsch Bestuur en het Inlandsch onderwijs. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 133-151.
- Het inlandsch onderwijs. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 223-252.

- GRAAFLAND, N.—Depok. Ene ethnographische Studie. *Med. Ned. Zendel*, vol. xxxv, 1891, pp. 1-43.
- GRABOWSKY, F.—Neber Aenoseringen geistigen Lebens bei den olo ngadju in Sued-Ost-Borneo. *Bijdr. taal-landvolkenk. Ned. Ind.*
- GRASHIUS, G. J.—Bijdrage tot de kennis van het Soendaneesch. 8vo. *Leiden*, 1891.
- Soendanesche Bloemlezing. Legenden en Moslimsche leerboekjes. 8vo. *Leiden*, 1891, viii, 136 pp.
- GRESHOFF, M.—Eene welriekende varen. *Teysmannia*, Jaarg. 1, 1890.
- Aperçu du premier rapport du Laboratoire Chimico-Pharmacologique du Jardin Botanique de l'Etat de Buitenzorg. *Ann. Jard. Botan. Buitenzorg*, vol. ix, pt. 2, 1891, pp. 247-260.
- Boeloe Ongko (*Artocarpus venenosa*, Zoll. et Mor.) *Teysmannia*, Jaarg. 1, 1890, pt. 9.
- Indische producten in de nieuwe Nederlandsche en duitsche pharmacopeën. *Teysmannia*, Jaarg. 1, 1890, pt. 9.
- Een boek over Sirih. *Teysmannia*, Jaarg. 2, 1891.
- Eenste verslag van het onderzoek naar de plantenstoffen van Nederlandsch-Indië. Mededeelingen uit 'sLands Placentuin. vii. Chemisch-pharmacologisch Laboratorium. 8vo. *Batavia*, 1890, iv, 128 pp.
- GRONEMAN, J.—Onjuiste voorstellingen. *Intern. Arch. Ethnogr*, vol. iii, 1890, p. 154.
- En avond bij den rijksbestuurder van Jogjakartã. *Intern. Arch. Ethnogr*, vol. iii, 1890, pp. 175-180, pl. xiv.
- "H."—Nog eens *Gloriosa Superba*. *Teysmannia*, Jaarg. 1, 1890.
- HAAK, J.—Opiumregie met normaal Tjandoe. 8vo. *Samarang*, 1889.

- HAARSMA, G. E.—Der Tabaksbau auf Deli. 8vo. *Amsterdam*, 1890, 2, vi, 240, pp. figs. and sketch maps.
- HARTHOORN, S. E.—De opperheerschappij van China over den Maleischen Archipel volgens Chineesche bescheiden. *Ind. Gids*, vol. xii, 1890, pp. 2170-2185.
- De Bantamshe staat en handel. *Ind. Gids*, vol. xii, 1890, pp. 1306-1322.
- “Onze uiterlijk beste en eerste vrienden” in Oost-Indië. *Ind. Gids*, vol. xii, 1890, pp. 1895-1913.
- HASSELT, A. W. M. VAN.—Araneae ex Archipelago Malayano, Weber. *Zool. Ergebn. Reise Ned. O. I.*, Heft 2, 1891, pp. 193-211.
- HASSELT, L. J. VAN.—Wat is voor een Papoea noodig om op het tooneel der hoogste beschaving eene plaats te kunnen innemen? *Ned. Zendingst.*, vol. ii, 1890, pp. 129-138.
- HAGEN, B.—Anthropologische Studien aus Insulinde Veröffentlicht durch die K. Akad. wiss zu Amsterdam. 4to *Amsterdam*, 1890, ii, 151, 16 pp.
- Die Pflanzen-und Thierwelt von Deli auf der Ostküste Sumatra's. *Tijdschr. Kon. Ned. aardrijksk. Gen.*, vol. vii, 1890, pp. 1-240.
- HALL J. N. VAN—Uit Multatuli's Brieven. *De Gids*, Marz, 1890, pp. 1-15.
- HAMER, C. DEN—De Sair Madi Kentjana. *Tijdschr. Ind. Taal-land volk.*, vol. xxxiii, 1890, pp. 531-564.
- HANDBOEK voor cultuur-en handelsondernemingen in Nederlandsch-Indië. 3e. Jaarg. 8vo. *Amsterdam*, 1890, viii, 601. pp.
- HAUGHTON, H. T.—Notes on Names of Places in the Island of Singapore. *Journ. Straits Br. R. A. S.*, vol. xx, 1890, pp. 75-83.
- HAVE, J. J. TEN.—Beknopte Aardrijkskunde van Nederlandsch Oost-en West-Indië. Ed. 2, 8vo. s'. *Hage*, 1891.

- HEERES, J. E.—De Gouverneurs-en Commissarissen-Generaal van Nederlandsch-Indië. *Ned. Spectator*, 1891, No. 9.
- HEIN, A. R.—De bildenden Künste bei den Dayaks auf Borneo. 8vo. *Wien*, 1890, 228 pp., 10 pls., 90 figs.
- HELFRICH, O. L.—Verzameling Lampongsche Teksten, getranscribeerd onder toezicht van *Vert. Batav. Gen. Kunst.*, vol. xlv, 1891.
- Proeve van een Lampongsch-Hollandsche woordenlijst, bepaaldelijk voor het dialect van Kroë. 8vo., *Batavia*, 1891, 2, 116 pp.
- HELDERMAN, W. D.—De tijger en het bijgeloof der Batahs. *Tijdschr. taal-land-volk.*, N. I., vol. xxxiv, 1890, pp. 170-176.
- HENNEMANN, J. P.—De opleiding van inlandsche arbeiders *Ned. Zendingsts.*, vol. ii, 1890, pp. 15-27.
- Het eerste tiental jaren van het Seminarie te Depok, 1878-1888. *Ned. Zendingsts.*, vol. ii, 1890, pp. 212-221.
- HIGGISON, S. J.—Java, the Pearl of the East. 12 mo., *Boston*, 1890, ii, 204 pp.
- HIKAJAT SOELTAN IBRAHIM.—Tertjitak pada Kantor pengelœwaren mèrèk Regina Orientis di Betawi. 4to., *Betawi*, 1890, 42 pp.
- Ibnoe Adaham Walijoe'llah. Tlahiran baharoe. 8vo., *Regensburg*, 1891, 52 pp.
- HIKAJAT SRI PADOEKA jang di Pertoewan Besar Jan Pieterzoon Cœn Jang memarintahi tanah Hindia Olanda deri pada tahun 1619 sampei tahun 1623 dan kadoewa kali deri pada tahun 1627 sampei tahun 1629 terkarang oleh J. Diepenhorst. 4to., *Betawi*, 1890, 64 pp.
- HINDORF, R.—Leitfaden zur Erlernung der Malayischen Umgangssprache (Küsten Malayisch). 8vo., *Betawi*, 1890, 72 pp.
- “HIPPOFIEL.”—De remonteering van het Nederlandsch-Indische leger en de verbetering der Indische paardenrassen. *Ind. Gids.*, vol. xii, 1890, pp. 1864-1895.

- HOËVELL, G. W. W. C. VAN.—Bijdrage tot de Ethnographie van den Indischen Archipel. *Intern. Arch. Ethnogr.*, vol. iii, 1890, pp. 186-188, pl. xvi.
- De Assistent-Residentie Gorontalo voor zoover die onder rechtstreeksch. Bestuur is gebracht. *Tijdschr. Ned. Aard. Genootsch.* [2], vol. viii, 1891, pp. 26-44.
- HOGENDORP, W.—De Sangir en Talanteilanden. *Ned. Zendingsts.*, vol. ii, 1890, pp. 113-123.
- HOLB, K. F.—Heerendienstregeling in Kedoe. *Ind. Gids.*, vol. xii, 1890, pp. 1132-1142.
- HOLLANDER, J. J. DE.—Handleiding bij de behandeling van't Maleisch letterschrift. Ed. 3, 8vo., *Utrecht*, 1890, ii, 32 pp.
- HOLLE, K. F.—Nota over de rijstcultuur voor Sumatra bewerkt door E. Th. van Delden vertaald door D. Gerth van Wijk. 8vo., *Batavia*, 1890, 40 pp.
- HORST, D. W.—De eed der Chineezzen. *Ind. Gids*, vol. xii, 1890, pp. 1502-1510.
- ILMOE OBAT obattan bergoena pada sekalian Orang Lelaki dan Prampoean jang dapat roepa-roepa penjakit terpetik dari boekoe tjina: Tat Seng Pian, No. 1. 8vo., *Batavia*, 1890, xvi, 95 pp.
- No. 2, *Ibid*, 1890, pp. 97-194.
- IMMINK, A. J.—Over de rechterlijke macht in Nederlandsch-Indië. *De Gids*, April, 1890, pp. 102-140.
- INILAH KITAB EDJA ja itoe pada menjatakan permoelaän beladjar membatja, terkarang oleh L. K. Harmsen (Tjitakan jang k-anam kalinja.) 8vo. Gouvernements besluit do 15 Juli, 1890, No. 28. Tertjitak di bandar Batawi pada perteraän Gouvernemen: tahoen 1890. 62 pp.
- ISNARD, L. C.—Gemencheh (District of Johol) Negri Sembilan. *Journ. Straits Br. R. A. S.*, 1890, No. 21.
- JAARBOEK, VAN DE K.—Nederlandsche Zeemacht, 1888-1889. 8vo. *'sGravenhage*, 1890, iii, 350 pp. [pp. 303-349. Verrichtingen in Oost-Indië.]

- JACOBS, J. R.—Neerlands driekleur in Neerlandsch-Indië, door Neerlands dapperen bewaakt. Schetsen uit het Indische Krijgsleven, met portretten, opgedragen aan de ridders der Militaire Willemsorde, beneden den rang van Officier. 8vo. *Rotterdam*, 1890, iv, 117 pp.
- JACOBS, JUL.—Een Javaansch volksfeest. *Eigen. Haard*, 1890, pp. 538 & 550.
- And J. J. MEIJER.—De Badoej's. 8vo. 's*Gravenhage*, 1891, 4 bl., 175 pp.
- JACOBSEN, A.—Reisen in Ostindischen Archipel. *Petermann's Mitth.*, 1890, pp. 103-105, map.
- JAEGER, W.—Van ginds (Atjeh). Herinneringen van een jong invalide. 8vo. *Amsterdam*, 1890, iv, 200 pp.
- JANSE, J. M.—Mededeelingen uit 's Lands Plantentuin. viii. Proeve eener verklaring van Serehverschijnselen door J. M. Janse. 8vo. *Batavia*, 1891, 40 pp.
- Chlorose van cultuurplanten en hare genezing. *Teysmannia*, Jaarg I, 1890.
- Druppelen. *Teysmannia*, Jaarg, 2, 1891.
- JURIEN DE LA GRAVIÈRE.—Les ouvriers de la onzième heure. Histoire des premières navigations des Anglais et des Hollandais dans les mers polaires et dans la mer des Indes. 2 vols. 8vo. *Paris*, 1890, iv, 338 and iv, 362 pp.
- "V. K."—Cijfers en feiten uit het kolonial verslag van 1889. *Ind. Gids*, vol. xii, 1890, pp. 692-727.
- KAN, C. M.—De Litteratuur over Nederlandsch-Indië sedert Juli, 1889. *Tijdschr. Aardr. Genootsch.*, ser. 2, vol. vii, 1890, pp. 543-557.
- De belangrijkste reizen der Nederlanders, in de 19e eeuw ondernomen. De voornaamste werken in dat tijdperk op geographisch gebied verschenen. *Tijdschr. Ned. Aard. Gen.*, vol. vi, 1890, pp. 510-581.
- KATECHISMUS, atou manga kakiwalo dingangu sasimbahe bou tatintirongu agama Mesehi ko Susi. Nisalin su bahasang sangie bou i. f. Kelling. 8vo. *Batavia*, 1890, 49. pp.

- KEANE, A. H.—Malays. *Chamber's Encycl.*, vol. vi, 1890.
- KERCKHOFF, C. E. P. VAN.—Eenige mededeelingen en opmerkingen betreffende de slavernij in Nederlandsch-Indië en hare afschaffing. *Ind. Gids.*, vol. xiii, 1891, pp. 743-769.
- Eenige opmerkingen betreffende de zoogenaamde "orang loeboe" op Sumatra's westkust. *Tijdschr. Ned. Aard. Genootsch.*, ser. 2, vol. vii, 1890, p. 576.
- Rechtspraak volgens de adat. *Ind. Gids.*, vol. xiii, 1891, pp. 1-11, 422.
- KERN, H.—Taalkundig gegevens ter bepaling van het stamland der maleisch-polynesische volken. *Versl. Med. ak. Amsterdam* (Letterk.), 2k. 3, vol. vi, 1890; and separately 8vo. 18 pp., 3 pls.
- KIELSTRA, E. B.—De ondergang van het Bandjermasinsche Rijk. *Ind. Gids.*, vol. xii, 1890, pp. 2389-2419; vol. xiii, 1891, pp. 12-39-364-387-819-845.
- Bijdragen tot de geschiedenis van Borneo's Westerafdeeling. *Ind. Gids.*, vol. xii, 1890, pp. 450-475; 682-692; 857-879; 1085-1113-1482-1502, 1694-1724, 1922-1942, 2185-2227.
- Het eiland Madoera. *De Gids*, 1890, pp. 517-546.
- Sumatra's Westkust van 1836-1840. *Bijdr. taal-land-volk.*, N. I., ser. 5, vol. v, 1890, pp. 263-348.
- KING, G.—Materials for a Flora of the Malayan Peninsula, No. 2. *Calcutta*, 1890.
- KITAB EDJA dan Badja bagi anak-anak jang baroe moelai beladjar oleh Toewan van Duijn, Goeroe jang kadoea pada midarsah beladjar goeroe di negri Probolinggo. 8vo. Gouvernements besluit dd^o 15 Juli, 1890, No. 28. Tertara di bandar Batawi pada pertaraän Goewernemen, 1890. 24 pp.
- KITAB jang bergoena kapada kanak-kanak jang hendak beladjar menbilang. Bahagian jang ka-ampat and kalima. 8vo. *Betawi*, 1889, 59, 87 pp.

- KLERK DE REUS, G. C.—De expeditie naar Bengalen in 1759 ii, iii. *Ind. Gids*, vol. xii, 1890, pp. 27-91; 247-279.
- KLINKERT, H. C.—Bloemlezing uit de Maleishche geschriften, ten behoeve van de gouvernements inlandsche scholen en van hen die het Maleische met Latijnsch karakter beoefenen. 8vo. *Leiden*, 1890, x, 413 pp.
- KOCK, W. H. W. DE.—Handboek voor den Algemeenen Ontvanger. 8vo. *Semarang*, 1860, lxxix, 416, lxvii pp.
- KOHLBRÜGGE, J. H. F.—Versuch einer anatomie des Genus. *Hylobates*. Weber *Zool., Ergebn. Reise Ned. O. I.*, Heft 2, 1891, pp. 211-354, pls. xvii-xix, 24 figs.
- KREEMER, J.—Regenafweerders. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 320-325: and *Ind. Gids*, vol. xii, 1890, pp. 2082-2086.
- Is uitbreiding van het Christendom langs vreedzamen weg op Java en Sumatra mogelijk. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 301-312.
- KRUIJF, E. F.—Missie, Antimissie en Omissie. *Ned. Zendingsts*, vol. 11, 1890, pp. 321-325.
- KRUIJT, A.—Staat van de gemeenten, de scholen en de ziekenpractijk, in den werkkring van Môdjô-warnô, over 1889. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 293-301.
- Spaarbank voor Inlanders te Môdjô-warnô. *Med. Ned. Gendel*. vol. xxxiv, 1890, pp. 312-314.
- Verslag aangaande den werkkring Môdjô-warnô over het jaar 1889, *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 274-293.
- KRUIJT, H. C.—Berichten uit Deli. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 325-332.
- Berichten (Deli tuwa), *Med. Ned. Zendel*, vol. xxxv, 1891, pp. 43-75.
- KUNEMAN, JULIUS.—De Gouvernements koffiecultuur op Java. Onderzoek en advies. 8vo. *s'Gravenhage*, 1891, 4, 201 pp
- “V. L”.—De nieuwe zending onder de Bataks. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 210-222.

- LALOY, L.—Présentation d'un Javanais et d'un Malais par Deje et Martin. *L'Aurore*, vol. 1, 1890, p. 628.
- LANGE, G. A. DE.—Mijn ontginnig van staatswege op Billiton. 8vo. *s'Gravenhage*, 1891, 64 pp.
- LANGENDOORF, M. M.—Een en ander over mijn arbeidsveld (Letti). *Ned. Zending's*, vol. ii, 1890, pp. 203-212.
- LENS, C., J. P. G. KRUIJT, and J. A. VAN DEN BROEK.—Maleische brieven van Sumatra: Noord-Oost en Oost Kust. Ed. 2, 4to. *Delft*, 1891.
- LEVYSSOHN NORMAN, H. D.—De Kedoese heerdienstregeling. *Ind. Gids.*, vol. xii, 1890, pp. 597-634.
- Misverstand. *Ind. Gids.*, vol. xii, 1890, pp. 1913-1922.
- LEWIN, L.—Neber das Babelkanen. *Intern. Arch. Ethnogr.*, vol. iii, 1890, pp. 61-65.
- LIDTH DE JENDE, TH. W. VAN.—Reptilia from the Malay Archipelago, ii, Ophidia, Weber. *Zool. Ergebn. Reise Ned. O. I.*, Heft. 2, 1891, pp. 178-192, pls. xv, xvi.
- LION, J.—Verzameling van de voornaamste publiek-en privaatrechtelijke algemeene verordeningen voor Ned.-Indië. 8vo. *Samarang*, 1890.
- LORGEAN,—Le Chant de l'Éléphant traduit du Siamois. *Mém. Soc. Acad. Indo-Chin.*, vol. 1, 1879 [1890], pp. 67-85.
- LONTER, J. DE.—Een British-Indische gemeentewet. *Ind. Gids.*, vol. xii, 1890, pp. 1041-1085.
- LOUWERIER, J.—Het huwelijk der Inlandsche Christenen in de Minabassa. *Ned. Zending's*, vol. ii, 1890, pp. 311-315.
- LUBBERS, A.—Eene bijdrage tot de Anthropologie der Atjehers. *Geneesk. Tijdschr. Ned.-Ind.*, vol. xxx, 1890.
- LUITJES, F. A.—Kitab edja dan batja. Bagian jang kadoea. 8vo. *Batawi*, 1889, 67 pp.
- "M".—Kleinleed. Schets uit het leven in Indië. *Eigen Haard*, 1891, pp. 70-72.
- MACMAHON, A. R.—The Ancient Shan Kingdom of Pong. *Asiat. Quart. Rev.*, July, 1890.

- MADRAZO, J. DE.—Filipinas. *Rev. Españ.*, vol. xxvii, pp. 438, 556; vol. xxviii, pp. 61, 391, 461; vol. xxix, pp. 64, 153.
- MARGADANT, W. F.—Voor drie jaren naar de Oost. 8vo. *Zutphen*, 1890.
- MARRE, ARISTIDE.—Notice des manuscrits Malaïs de la Bibliothèque Nationale. *Mém. Soc. Acad. Indo-Chin.*, vol. i, 1879 [1890], pp. 174-189.
- MARRE,—Index des mots Malaïs Français. *Mém. Soc. Acad. Indo-Chin.*, vol. I, 1879 [1890], pp. 108-146.
- MARTIN, K.—Versteinerungen der sogen. Alten Schieferformation von West Borneo. *Samml. Geol. Reichs, Mus. Leiden*, vol. iv (.), 1890, pp. 198-207.
- Neber nene *Sigodon*, Reste aus Java. 4to. *Amsterdam*, 1890, 3 pls. [Veröff. d. d. k. ak. wiss. amst.]
- MATTHES, B. F.—Bijbelsche geschiedverhalen in tweemaal twee en vijftig lessen. In het Makas-aarsch vertaald. 8vo. *Amsterdam*, 1890, iv, 437 pp.
- MAURITS.—Indische menschen in Holland. Oorspronkelijke roman. 8vo. *Leiden*, 1890, iv, 290 pp.
- MENDELS, J.—Herman Willem Daendels, voor zijne benoeming tot Gouverneur-Generaal van Oost-Indië (1762-1807). 8vo. *s'Gravenhage*, 1890, xvi, 394, 210 pp.
- METEOROLOGISCHE—Waarnemingen in Nederlandsch-Indië. *Nat. Tijdschr. Ned. Ind.*, yearly.
- METZGER, E.—Grenzlinien zwischen Asien und Australien. *Ausland*, 1890, pp. 609 & 629.
- Vierzig Jahre Niederländischer Kolonialherrschaft in Ostindien. 8vo. *Hamburg*, 1890, iv, 52 pp.
- Ein dest der Dajak. *Ausland*, 1890, pp. 474-476.
- Java. *Chamber's Encycl.*, vol. vi, 1890.
- MEYER, A. B. & A. SCHADENBERG.—Die Philippinen. I. Nord-Luzon. 4to. *Dresden*, 1890, 26 pp., 18 pls.

- MEYER, A. B. & A. SCHADENBERG.—Album von Philippinen-typen (Nord-Luzon). 4to. *Dresden*, 1891, 19 pp. 50 pls.
- MEYER, J. J.—Les fixations légales relatives à la religion musulmane aux Indes néerlandaises. *Ann. Extr. Orient*, vol. xiv, 1890, pp. 257-267, 289-298.
- L'Administration de la justice parmi les indigènes dans les îles de Java et de Madoura. *Ann. Extr. Orient*, vol. xiv, 1890, pp. 129-142.
- MEYNEERS D'ESTREY, H.—Notre des manuscrits malais de la Bibliothèque de la Société des Arts et Sciences de Batavia. *Mém. Soc. Acad. Indo-Chin.*, vol. 1, 1879, [1890], pp. 198-213.
- Peuplement excessif de Java. *Rev. Géogr.*, Novr., 1890, xxvii, pp. 340-349.
- Les Kongsis ou républiques d'émigrants chinois dans l'ouest de Bornéo, i-x. *Ann. Extr. Orient*, vol. xiv, 1890, pp. 65-74, 97-106, 165-175.
- MILNE.—Building in the Philippines. *Min. Proc. Inst. Civ. Engineers*, vol. 100, 1890.
- MOREN, H. H. & J. M. ROSSKOPF.—Eenige mededeelingen omtrent het Protestantsch Weeshuis te Semarang in de laatste jaren. *Ind. Gids*, vol. xiii, 1891, pp. 252-257.
- MORRIS, D. F. VAN BRAAM.—Nota van toelichting behoorende bij het contact gesloten met het Landschap Bima. *Tijdschr. Ind. taal-land-volk.*, vol. xxxiv, 1890, pp. 176-232.
- MUHAMMED, *Arsad B. Abdallah el Bungari*, Sebil el-muhtadin. [Transl. into Javanese from the Arabic.] 4to. *Bulog*, 1, 307, 265, 279 pp.
- MÜLLER, A.—Orientalische Bibliographie meter mit wirkung der Herren Bezenberger, Gleiniger, Gottheil, J. Müller, Strack, Vollers, und Wijnmalen. 8vo. *Berlin*, Jahrg., iv, (1890), 1890 and 1891: Jahrg. v, (1891), 1891.
- MULTATULI [E. D. DEKKER].—Nogeens: Vrije-arbeid in Nederlandsch-Indië, Multatuli en zijne werken geschetst door Aart-Admiral. Ed. 3, 8vo. *Arnheim*, 1890, 192 pp.

MULTATULI [E. D. DEKKER].—Brieven. Bijdragen tot de kennis van zijn leven. Gerangschikt en toegel. Door Douwes Dekker. D. I (Wordingsgeschiedenis van den Havelaar.) 8vo. *Amsterdam*, 1890, 155 pp.

-----Twee fragmenten uit een onafgewerkt blijspel. 8vo. *Amsterdam*, 1891.

-----Brieven. Bijdragen tot de kennis van zijn leven. Gerangschikt en toegelicht door Mevrouw Douwes Dekker, geb. Hamminck Schepel (Deel ii, De Havelaar verschenen 1860). 8vo. *Amsterdam*. 1890, iv, 240 pp.

"H. N."—Singapore. *Pall Mall Gazette*, Aug., 21, 1890, p. 3.

-----Old and new Siam. An Interview with H.R.H. Devawongse Varoprakar, the Siamese Minister of Foreign Affairs. *Pall Mall Gazette*, August 30, 1890.

NAAMLIJST der Europeesche inwoners van het mannelijk geslacht in Nederlandsch-Indië en opgaven omtrent hun burgerlijken stand, 1890. 8vo. *Batavia*, 1890, 4, 412 pp.

-----der Europeesche bewoners van het mannelijk geslacht in Nederlandsch-Indië en opgaven omtrent hun burgerlijken stand, 1891. 8vo. *Batavia*, 1891, 4, 420 pp.

NAAM-EN RANGLIJST der officieren van het Koninklijke leger der Nederlanden en van de Nederlandsche Indiën voor 1891, Jaarg, 60. 8vo. *Gorinchem*, 1891, xvi, 396 pp.

NEDERLANDSCHE MAIL.—Wekelijksche dienst op Nederlandsch Oost-Indië. 8vo. *Amsterdam*, 1891.

NETHERLAND INDIES.—Finances for the year 1890-1891. *Parliamentary Papers*, [6205-156].

NEURDENBURG, J. C.—De opleiding van onze Zendelingen. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 416-437.

-----Overzicht van het Zendingveld in Nederlandsch-Indië. *Ned. Zendingsts*, vol. ii, 1890, pp. 90-102.

-----Statistiek uit de Minahassa. *Ned. Zending*, vol. iii, 1891, pp. 60-62.

- NEURDENBURG, J. C.—Uit de binnenlanden van Borneo. *Med. Ned. Zendel.* vol. xxxiv, 1890, pp. 437-441.
- NEVE, E. K. A. DE.—De grondslagen der organisatie van het Indische leger, i, ii. *Ind. Gids.*, vol. xii, 1890, pp. 833-857; 1113-1132.
- De grondslagen der organisatie van het Indische leger. iii, iv. *Ind. Gids.*, vol. xii, 1890, pp. 1268-1275, 1462-1482.
- NIEMANN, G. K.—Nog iets over het eiland Rotti of Rote. *Med. Ned. Zendel.*, vol. xxxiv, 1890, pp. 43-45.
- NIJPELS, G.—Het wetsontwerp tot regeling van de opleiding en de hogere vorming van den officier van het Indische leger in Nederland. *Ind. Gids.*, vol. xii, 1890, pp. 298-313.
- OHL, A. E. U. W.—Vier brieven aan een lid van de Tweede Kamer over de muntdepêche van den Minister van Kolonien. (Overged: uit de Soerabaya-Courant). 8vo. *Soerabaya*, 1890, 42 pp.
- OOMEN, J. H. B.—Een nieuw hulpmiddel bij de inning van belastingen. *Ind. Gids.*, vol. xiii, 1891, pp. 161-183.
- De Gouvernements-koffiecultuur en art. 564, van het Regeerings reglement. *Ind. Gids.*, vol. xii, 1890, pp. 279-298.
- OOST INDISCH AMBTENAAR.—Mahomedaansch-godsdienstige broederschappen. 8vo. *Batavia*, 1890.
- OPHUIJZEN, C. A. VAN.—Bataksche spreekwoorden en spreekwijzen. *Tijdschr. taal-land-volk.*, N. I., vol. xxxiv, 1890, pp. 7²-104.
- OTTOW, W. M.—Over vervalschingen van kajoepoti olie. *Geneesk. Tijdschr. Ned.-Ind.*, vol. xxx, 1890.
- OUDEMANS, A. C.—Over verontreiniging van Banka tin. *Jaarb. Mijnw. Ned. O. I.*, vol. xix, 1890, *Techn. Ged.*, pp. xxiv-xxxvi.
- PAGENSTECHEER, A.—Beiträge zur lepidopteren fauna des Malayischen Archipels. v. Neber Schmetterlinge von Ost-Java. 8vo. *Wiesbaden*, 1891, 18 pp.

- PERATOERAN hoekoeman policie boeat sekalian orang bangsa Djawa dan sebrang di tanah Hindia Nederland. Tjitakan jang kadoewa. 8vo. *Batavia*, 1890, 24 pp.
- PERELAER, M. T. H.—Baboe dalima. Opuim-roman. 2 vols., Ed. 2. 8vo. *Amsterdam*, 1889, xii, 326; vi, 317 pp.
- PETIKAN dari Staatsblad, No. 149 en 158, 1890, atoeran dari pacht amfioen serta perdjandjian goenanja di tanah Djawa dan Madoera. 8vo. *Soerabaia*, 1890, 34 pp.
- PETIT, W. L. DE—La conquête de la vallée d'Atchin par les Hollandais. Une page d'histoire coloniale contemporaine. 8vo. *La Haye*, 1891, viii, 371 pp., map.
- PLEYTE, C. M.—Indonesisches Feuerzeng. *Globus*, lix, 1890, pp. 52-56.
- Image en bois: est-elle de Bornéo? *Intern. Arch. Ethnogr.*, vol. iv, 1891, p. 93.
- PLISCHKE, KARL.—Kurze Mittheilungen über zwei Malayische Spiele. *Intern. Arch. Ethnogr.*, vol. iii, 1890, pp. 189-194.
- POENSEN, C.—De organisatie der door de zending ontstane christen-gemeenten in Ned. Indië. *Ned. Zendingsts*, vol. ii, 1890, pp. 27-46.
- De Evangelisatie op Java. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 391-416.
- POSEWITZ, TH.—Mittheilungen über Nord-Borneo. *Petermann's Mitth.*, vol. xxxvi, 1890, pp. 203, 204.
- Diè neueste Durchkrenzung von British Nord-Borneo. *Petermann's Mitth.*, vol. xxxvii, 1891, p. 47.
- Vulkane in Borneo. *Ausland*, 1890, p. 219.
- POTTER, F. J.—De kultuur van suikerriet op Java. 8vo. *Arnheim*, 1890, 70 pp.
- POURTALÈS, DE.—Représentation de l'industrie française aux Indes Néerlandaises. Ed. 2, 8vo. *Paris*, 1890, ii, 23 pp.
- RADEN MAS ARIA NATA ASMARA II.—Geslachtslijst van den eersten Sultan van Jogja tot den Zesden. 8vo. *Jogjakarta*, 1890, 60 pp.

- RAJAL Y LARRE, J.—Memoria acerca de la provincia de Nueva Ecija, en Filipinas. *Bol. Soc. Geogr. Madrid*, vol. xxvii, pp. 290-340.
- REINBOLD, TH.—Sargassen voor Indischen Archipel. *Ann. Jaard. Botan. Buitenzorg*, vol. x, pt. 1, 1891, pp. 67-74.
- RENAN, D. G. A. H.—Verslag van een onderzoek naar Petroleum in Langkat. *Jaarb. Mijnw. Ned. O. I.*, vol. xix, 1890, verb. zedur. Gid., pp. 1-9.
- RHEDE VAN DER KLOOT, M. A. VAN.—De Gouverneurs-Generaal en Commissarissen-Generaal van Nederlandsch-Indië. 1610-1888. Historisch-geneologisch beschreven. Met wapenafbeeldingen door M. J. Lion. 8vo. 's *Gravenhage*, 1891, xii, 355 pp.
- RHO, F.—A traverso l'arcipelago Malese. Dalle note di un viaggio di circumnavigazione sulla R. Corvetta "Caracciolo." 8vo. *Firenza*, 1890, 127 pp.
- RIEDEL, J. G. F.—Les idées spécifiques du droit de propriété foncière chez les Indonésiens. *T'oung Pao*, vol. i, 1891, pp. 277-296.
- RIEMENS, J.—Overzicht van de zesde Zendings-conferentie, gehouden te Batavia. *Ned Zendingsts*, vol. ii, 1890, pp. 257-262.
- RIDLEY, H. N.—Destruction of Coconut Palms by Beetles. *Journ. Straits Br. R. A. S.* No. 20 (1889), 1890.
- RINNOOY, N.—Het buiten-christelijk huwelijk. *Ned Zendingsts*, vol. ii, 1890, pp. 193-203.
- ROCHEMONT, J. I. DE.—Rudolf van Meerkerke. 8vo. 's *Gravenhage*, 1890, 10, 340 pp.
- ROGGE, C.—Eene dienstreis van Amboina naar de Noordkust van Ceram. *Tijdschr. Ned. Aard. Genootsch*, ser. 2, vol. vii, 1890, pp. 844-866.
- ROHLFS, G.—Gustav Nachtigal's Abschied vor seiner Reise nach Bornu. *Deutsch. Rundsch. Geogr.*, xiii, 1891, pp. 289-298.

- ROOKER, H.—Bericht nopens den voortgang der Evangelisatie in de Minahassa over 1888. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 60-87; 174-210.
- Staat der Genootschapsscholen in de Minhassa, ult. 1888. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 54-60.
- Statistieke opgaven van het aantal gedoopten, enz. in de Minahassa over 1888. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 45-54.
- ROOS, K. H. F.—Aanteekeningen over de afdeeling Kaoer in de residentie Bengkoelen. *Tijdschr. Binnenl. Bestuur*, vol. v, 1890, p. 122.
- ROOY, DE.—De positie der volkshoofden in een gedeelte der Padangsche Bovenlanden; hunne ethnographische en hunne politieke beteekenis. *Ind. Gids*, vol. xii, 1890, pp. 634-682.
- De koffiecultuur ter Sumatra's Westkust en het rapport der Staats commissie. *Ind. Gids*, vol. xii, pp. 1409-1434.
- ROUVEROY van NIEUWAAL, M. C. VAN.—Het Maleisch in de kazerne. Maleische woorden en uitdrukkingen, die in de kazerne tegenover niet-Europeesche militairen te pas komen. Gesprekken en taalregels, gevolgd door eene lijst der meest voorkomende woorden en twee aanhangsels. 8vo. *Breda*, 1891, 4, 97 pp.
- RUGE, G.—Anatomisches über den Rumpf der Hylobatiden Ein Beitrag zur Bestimmung der stellung dieses Genus in systeme. Weber, *Zool. Ergebn. Reise Ned O. I.* Heft 2, 1891, pp. 366-461, pls. xxi-xxv.
- SALINAN.—Staatsblad dan Bijblad dari jangbaroe kaloear teroes keblakang sampe tahun 1816 aken grenanja anak negri dan sesamanja. 8vo. *Batavia*, 1890, 172 pp.
- SAMBOENGAN.—Bockoe tjerita-an tanah djawa. Bagian ka 2. 8vo. *Soerabaia*, 1890, 172 pp.
- Bockoe tjerita-an tanah djawa. Bagian ka 3. 8vo. *Soerabaia*, 1890, 55 pp.

- SANDWICK, R. A. VAN.—In het rijk van Vulcaan. De uitbarsting van Krakatau en hare gevolgen. 8vo. *Zutphen*, 1891, 195 pp., map.
- SCHEIDENAGEL, M.—El archipélago do Legaspi. 8vo. *Madrid*, 1890, 320 pp.
- SCHERER, G. A.—Hoe moet Atjeh gepacificeerd worden. Voordracht gehouden in het Indisch Genootschap. 8vo. *'sGravenhage*, 1891, 46 pp.
- Nog altoos het Atjehvraagstuk. *Tijdschr. Ned. I.*, 1891, pp. 283-316.
- SCHLEGEL, G.—Chinese loanwords in the Malay language. *T'oung Pao*, vol. 1, 1891, pp. 391-405.
- SCHMELTZ, J. D. E.—Indonesische Prunkwaffen. *Intern. Arch. Ethnol.*, vol. iii, 1890, pp. 85-118.
- Poignards de Bornéo. *Intern. Arch. Ethnogr.* vol. iii, 1890, p. 196.
- Beiträge zur Ethnographie von Borneo. *Intern. Arch. Ethnogr.*, vol. iii, 1891, pp. 238-242, pl. xix.
- SCHOEMAKER, J. P.—Schetsen uit den Atjeh-oorlog. ii. 8vo. *'sGravenhage*, 1890, vi, 248 pp.
- De contra-guerilla in de N. O. linie te Atjeh. *Haagsche Stemmen*, 1890, No. 29.
- De verovering van Kota-Pasir (Mela-boeh). *Haagsche Stemmen*, 1890, No. 30.
- SCHOT, J. G.—Moslemen en Christendom. 8vo. *Leiden*, 1890, iv, 91 pp.
- Exploitatiegeest of zedelijk bewustzijn? *Ind. Gids*, vol. xiii, 1891, pp. 547-569.
- Kolonisatie in onze Oost. Sociaal-economische studie. 8vo. *Amsterdam*, 1891, 51 pp.
- Eenige Nederlandsch-Indische belangen. I. De inmenging der Staten-General. II. Het concubinaat der Europeesche Ambtenaren. III. De honding der Regeering tegenoner Islam en Christendom. IV. De opium quaestie. *Ind. Gids*, vol. xii, 1890, pp. 1625-1682.

- SCHUCHARDT, H.—Kreolische Studien. IX. Ueber das Malaiio-Portugiesische von Batavia und Tugu. *Sitz. k. Ak. Wiss Wien*, vol. cxxii.
- SCHUICK, A.—Die Mangarai-Strasse. *Petermann's Mitth.*, 1890, pp. 180, 181.
- SCHULZE, L. F. M.—Führer auf Java. 8vo. *Leipzig*, 1890. viii, 480 pp. Map of railways in Dutch. Gids voor Java. 8vo. *Leipzig*, 1890, viii, 480 pp.
- SCHUURMANS, N. D.—De Tarigah Nagsjibendijah op Java. *Ned. Zendingsts*, vol. ii, 1890, pp. 265-278.
- Mededeelingen van Javanen over Javaansch heidendom vestaald en toegelicht. *Ned. Zendingsts*, vol. ii, 1890, pp. 65-90.
- Javaansche middelen tegen de Cholera. *De Macedonier*, 1891, (22 pp).
- De "Twee Woorden" van den Islam, volgens de Tariqah asj-sjattarijjah. *Ned. Zending*, vol. iii, 1891, pp. 23-60.
- SEMBOER-SOEROEK, djopo-mentro. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 335-360.
- SERAT KANTJIL; het boek van den Kantjil, Javaansch dierenepos. 8vo. 'sGravenhage, 1889, 167 pp.
- SERRURIER, L.—De Pionier, handleiding voor het verzamelen en waarnemen op natuurkundig gebied, inzonderheid in tropische gewesten.
- SIE-ANOE.—Beknopte handleiding om de Maleische taal te verstaan en te spreken. 8vo. *Haarlem*, 1890, 93 pp.
- SJA'IR, djalanan krëta api ja'itoe Bataviasche Oosterspoorweg dengan personeelnja bij gelegenheid van de opening der lijn Tjikarang-Kedoeng-Gedé bezongen oleh Tan Teng Kie. 8vo. *Betawi*, 1891, 24 pp.
- Midah Tjoemboewan tersalin dari pada hoeroef 'Arab serta di sahken karangannja. 8vo. *Betawi*, 1890, 104 pp.
- SKERTCHLY, S. B. J.—On fire-making in North Borneo. *Fourn. Anthropol. Inst.*, vol. xix, 1891, pp. 445-452, pl. xi.

- SKERTCHLY, S. B. J.—On some Bornean Traps. *Journ. Anthropol. Inst.*, vol. xx, 1890, pp. 211-219, pls. iv-x.
- SOERAT Kabar, Indiëa Nederland. 4to. *Amsterdam*, 1890, 8 pp.
- Kandoengan boeat goenanja segala priajie-priajie, jang memegang pakardjân di tanah gouvernemennan di poelo Djawa dan Madoera.....terkoempoel oleh L. T. M [eyer]. i-vi. 8vo. *Semarang*, 1889-90, 562 pp.
- Ketrangan dari pada hal kaadaän bangsa Tjina di negri Hindia Olanda. terkarang oleh Padoeka Toean J. E. Albrecht. 8vo. *Batavia*, 1890, 61 pp.
- SOERAT Peratoeran jang bahroe atas Pacht Apioen di tanah Djawa dan Madoera (Staatsblad 1890, No. 149), dan di Pertja Barat, Palembang, Bangkahoela, Tanah Lampong dan di.....Djambi dan taäloknja dan di Residentie Borneo salatan dan timoer sehingga jang masok pada koewasa Goebnernemen Olanda (Staatsblad 1890, No. 155. 8vo. *Batavia*, 1890, 90 pp.
- SOLMS-LAUBACH, H. Z.—Ueber die species in der gattung. *Rafflesia*, insonderheit ueber die auf den Philippinen sich findenden Arten. *Ann. Jard. Botan. Buitenzorg*, vol. ix, pt. 2, 1891, pp. 184-246.
- STELLWAGEN, A. W.—De Gouverneur-Generaal Jan Pieterzoon Coen. *Elsevier's Geill. Maandsch*, 1891, pp. 292-307, Portrait.
- TABAI DE TOOVENAAR.—Schets van Dajaksche zeden. *Tijdschr. Ned. Indië*, 1890, ii, pp. 57-70.
- TATINTIRO MANIMBU, Mangalene: Manga Kakiwalo dingangu sasimbahe bou Hikajetu Elkitab. Nisalim su bahasang Sangie von i F. Kelling. 8vo. *Batavia*, 1890, 43 pp.
- TERPETIK dari Burgerlijke Wetboek di salin bahasa Melaijoe goenanja toean Tanah serta toean Tanah di Hindia Nederland. 8vo. *Soerabaia*, 1890, 34 pp.
- TIELE, P. A.—Bouwstoffen voor de geschiedenis der Nederlanden in den Maleischen Archipel. 8vo. *s'Hage*, 1889-1890.

- TIELE, P. A.—Bouwstoffen voor de geschiedenis der Nederlanders in den Maleischen Archipel, pt. 2, by J. E. Heeres. *Haag*, 1890.
- TORAT OEMMAT ISJRAËL.—Perhimpoeanan akan melebehken pembatja-an soerat-soeratan Meseli di Betawi. 8vo. *Batavia*, 1890, 54 pp.
- TREACHER.—Sketches of Brunai, Sarawak, Labuan and North Borneo. *Four. Straits Br. R. A. S.*, No. 20 (1889), 1890.
- TREUB, M.—Verslag omtrent den staat van s'Lands Plantentuin te Buitenzorg en de daarbij behoorende inrichtingen over het jaar, 1889. 8vo. *Batavia*, 1890, 88 pp.
- J. E. Teysmann. Eene Schets. *Teysmannia*, vol. i.
- TROMP, S. W.—Herinneringen van een Oost-Indisch Ambtenaar. *Eigen Haard*, 1890, pp. 86, 134, 154, etc.
- Mededeelingen uit Borneo. *Tijdschr. Ned. Aard. Genootsch.*, ser. 2, vol. vii, 1890, pp. 728-763.
- TRUONG-VINH-KY, P. J. B.—Kim, Vân, Kiêu Truyên. Poèmes populaires annamites transcr. en gôt-ngu, précédées d'un résumé analytique. 16mo. *Saigon*, 1889, 136 pp.
- VAN DER BERG, L. W. C.—De toekomst van Atjeh. *Tijdschr. Ned. Ind.*, 1890, pt. i, pp. 209-219.
- VAN DER BERG, N. P.—De Bataviasche Decemberbeweging van 1795. *De Gids*, Feb., 1890, pp. 369-413.
- Regeling der verhouding tusschen werkgevers en arbeiders in Nederlandsch-Indië. Voor- dracht gehouden in de algemeene vergadering van de Nederlandsche afdeeling der Nederlandsch-Indische Maatschappij van Nijverheid en Landbouw op 9 Decem- ber, 1890. 8vo. *Amsterdam*, 1891, 31 pp.
- VAN DER CHIJS, J. A.—Philip Cartaret in Nederlandsch-Indië 1767-1768. *Tijdschr. taal-land-volk.*, vol. xxxiv, 1890, pp. 1-62.

- VAN DER CHIJS, J. A.—Nederlandsch-Indisch Plakaatboek, 1602-1811, vi, Deel. 1750-1754. 8vo. *Batavia*, 1889, ii, 873 pp.
-
- Nederlandsch-Indisch Plakaatboek, 1602-1811, Deel. vii, 1755-1764. 8vo. *Batavia*, 1890, ii, 880 p2.
-
- Dagh-Register gehouden in't Casteel Batavia. Anno 1661. Uitgegeven.....van der Chijs. 8vo. *Batavia*, 1889, 2, 557 pp.
- VAN DER ENT, M.—Hollandsch-Soendasche spreekwijzen, brieven en korte opstellen. 8vo. *s'Gravenhage*, 1891, 106 pp.
- VAN DER KELLEN.—Brief over Gambos en zijne bewoners. Medeged door P. J. Veth. *Tijdschr. Kon. Ned. Aard. Gen.*, vol. vi, 1890, pp. 91-104.
- VAN DEN KEMP, H. P.—Het Rapport van den Hoofdinspecteur van Financien H. J. van de Graaff, dat de aanleiding is geweest tot de intrekking der landverhuur in de Vorstenlanden. *Tijdschr. Nijverh. Landb. N. I.*, Dec., 1890.
- VAN DER KEMP, P. A.—De voorschriften op het Europeesch lager onderwijs in Nederlandsch-Indië verzameld en toegelicht. 8vo. *Batavia*, 1890, xxiv, 431 pp.
- VAN DER STOK, J. P.—Regenwaarnemingen in Nederlandsch-Indië, xi, Jaarg, 1889. 8vo. *Batavia*, 1890, x, 418 pp.
- VAN DER TOORN, J. L. and L. K. HARMSSEN.—Pada menjatakan sagala halahwal ilmoe bilangan jang boleh habis dibahagi dan lain sabalsainja. 8vo. *Betawi*, 1889, 43, ii, pp.
- VAN DER VALK, T. A. F.—Beschrijving van Nederlandsch-Indië. Ed. 2, 8vo. *Batavia*, 1890, v, 192 pp.
- VERBEEK, R. D. M.—Minéraux utiles que l'on trouve dans l'archipel Indo-Néerlandais. *Cochinchine Franç. Excursions*, 1890, xiv, No. 32, pp. 227.235.
- [VERNE, JULES].—Hikaijat Fileas Fogg atawa tjariteranja orang bertarohan sanggoep mengoelilingi boemi dalem 80 hari lamanja oleh toewan L. Th. M. Menoeroet. 8vo. *Semarang*, 1890, iv, 148 pp.

- VERSTIJNEN, F.—Een ontwerp-ordonnantie voor de regeling der landrente op Java en Madoera met toelichtingen en een ontwerp-instructie voor die uitvoering. *Ind. Gids*, vol. xiii, 1891, pp. 979-1052.
- VETH, P. J.—Hasan-Hosein of Taboetfeest. *Intern. Arch. Ethnogr.*, vol. iii, 1890, p. 194.
- VILLEMEREUIL, DE.—Dondart de Lagrée et ses manuscrits. *Mém. Soc. Acad. Indo-Chin.*, vol. i, 1879 [1890], pp. 42-67.
- VIS, M. H.—Gunther von Bülzingslöwen bij de 2^e Expeditie tegen Atjeh. 8vo. *Soerabaya*, 1890.
- VORDERMAN, A. G.—*Euplocamus sumatranus* (Dub.) en *Euplocamus Vieilloti* (G. R. Gray). Derde bijdrage tot de kennis der ornithologie van Sumatra. *Nat. Tijdschr. Ned. Indië*, vol. xlix, 1890, pp. 97-104.
- Eurycoma longifolia*, Jack., de moederplant van "Kajoe bidara laout." *Teysmannia*, Jaarg. 1, 1890.
- De Chineesche behandelingswijze van Keeldiphtheritis. 8vo. *Batavia*, 1890, iv, 92, ii pp.
- VORSTMAN, R.—Een vraag van den dag. 8vo. *Macassar*, 1890, ii, 49 pp.
- VOUTE, E. J.—Chevelures. • *Teysmannia*, Jaarg. 1, 1890.
- Araucaria excelsa*, R. Br. *Teysmannia*, Jaarg. 1, 1890, pt. 9.
- Nieuwigheden en indische tuinbouw. *Teysmannia*, Jaarg. 2, 1891.
- Bladgroenten. *Teysmannia*, Jaarg. 2, 1891.
- VRIES, S. J. DE.—De invloed van den Islâm in Indië. *Med. Ned. Zendel*, vol. xxxiv, 1890, pp. 30-43.
- WAAL, G. DE.—Transscriptie van het Manangkabausch Maleisch, eene proeve tot beëindiging der Quaestie. *Tijdschr. Ind. taal-land-volk.*, vol. xxxiv, 1890, pp. 100-103.

- WALLACE, A. R.—The Malay Archipelago, the land of the Orang-utan and the Bird of Paradise. New Ed. 8vo. London, 1891, xviii, 516 pp., maps and figs.
- WEBER, M.—Eigenthümliche Lagerung der Leber und Niere bei Siluroiden. *Weber, Zool. Ergebn. Reise Ned. O. I., Heft 2*, 1891, pp. 355-365, pl. xx.
- Reptilia from the Malay Archipelago. I. Sauria, Crocodylidae, Chelonia. *Weber's Zool. Ergebn. Reise N. O.-I., Heft 2*, 1890.
- Eenige mededeelingen over zijne reizen in Indie in het vorige jaar. *Tijdschr. Ned. Aard. Genootsch.*, Ser. 2, vol. vii, 1890, pp. 454-462.
- WERUMEUS-BUNING, A.—Indische marineleven. *Eigen Haard*, 1890, pp. 42, 58, 74, 689, 705.
- Een Battaksche kampong. *Tijdschr. Ned. Ind.* 1890, pp. 382-390.
- WESLY, L. F.—De onderafdeeling Rawas (Residentie Palembang). *Tijdschr. Binnenl. Bestuur*, vol. iii, 1890, p. 275.
- WESSELS, L.—De opheffing van het monopolie en de vervanging van de gedwongen Koffiecultuur op Java door een Staatscultuur in vrijen arbeid. 8vo. *s'Gravenhage*, 1890, ii, 72 pp.
- De landrente op Java in verband met het kadaster. *Ind. Gids*, vol. xii, 1890, pp. 91-125.
- De Gouvernements—Koffiecultuur in de Minahassa. *Tijdschr. Ned. I.*, 1891, pp. 50 and 123.
- WESTENBERG, C. J.—Nota over de onafhankelijke Bataklanden. *Tijdschr. taal-land-volk., N. I.*, vol. xxxiv, 1890, pp. 105-117.
- WESTERHOUT, A. R.—De doodenakkers van Indië, Geschiedkundig verhaal nit den Atjch-oorlog, Afl. 1. 8vo. *Amsterdam*, 1891, pp. 1-32.
- WICHMANN, A.—Die aufnahme der Molo-strasse. *Petermann's Mitth.*, 1890, pp. 153, map.

- WICHMANN, A.—Bericht über eine in Jahre 1888-1889 ausgeführte Reise nach dem Indischen Archipel. *Tijdschr. Ned. Aard Genootsch.* [2], vol. vii, 1891, pp. 907-994.
- WIGMAN, H. J.—Cacao. *Teysmannia*, Jaarg. 1, 1890.
- Aanleg van tuinen en erven. *Teysmannia*, jaarg. 1, 1890.
- Stamrozen. *Teysmannia*, Jaarg. 1, 1890, pt. 9.
- Water in tuinen. *Teysmannia*, Jaarg. 2, 1891.
- Over tuin-aanleg, hoofdzakelijk aan de hand van Leonard A. Springer. *Teysmannia*, Jaarg. 2, 1891.
- WIJNGAARDEN, J. K.—Eerste Verslag. Savoe. *Med. Ned. Zendel.* vol. xxxiv, 1890, pp. 366-390.
- WILDEMAN, E. DE—Les Trentepohlia des Indes Néerlandaises. *Ann. Jard. Botan. Buitenzorg*, vol. ix, pt. 2, 1891, pp. 127-142.
- WILKEN, G. A.—Over het huwelijks-en erfrecht bij de volken van Zuid-Sumatra. Ene bijdrage tot de ontwikkelingsgeschiedenis van het familierecht. 8vo. *s'Gravenhage*, 1891, 87 pp.
- Het Animisme bij de Volken van den Oost-Indischen Archipel. *Tijdschr. Binnenl. Bestuur*, vol. v, 1890, pp. 34-38.
- Struma en cretinisme in den Inischen Archipel. *Bijdr. taal-land-volk.*, N. I., ser. 5, vol. v, 1890, pp. 349-425.
- WINKLER, F. C.—Kitab iboe dan anak Aken goenanja estri njang soeda kawin. Tersalin bahasa melaijoe. 8vo. *Soerabaija*, 1890, ii, 116 pp.
- WRAY, L.—Journal of a collecting expedition to the Mountain of Batang Padang, Perak. *Journ. Straits Br. R. A. S.*, 1890, No. 21.
- "X".—Segli in 1885-1887. *Ind. Gids*, vol. xii, 1890, pp. 1849-1864.

- “X”—De tarieven voor het vervoer van suiker op de lijn Semarang-Vorstenlanden en het overnemen van dien spoorweg door den Staat. *Ind. Gids*, vol. xiii, 1891, pp. 388-394.
- Irrigatie werken of Java. *Ind. Gids*, vol. xiii, 1891, pp. 321, 673, 948.
- YOUNG, J. W.—Eenige aantellingen betreffende de Chineezzen in Nederlandsch-Indië. *Tijdschr. Ned. Indië*, 1890, pp. 362-375.
- De wetgeving ten aanzien van geheime genootschappen of broederschappen onder de Chineezzen in de Straits Settlements en in Nederlandsch-Indië. *Tijdschr. Ned. Indië*, 1890, pp. 179-209: 214-292.
- ZEGERS, J. L.—Het opuinvraagstuk in Nederlandsch Oost-Indië. 8vo. *Nijmegen*, 1890.
- ZORAB, G.—De publiekrechtelijke toestand der vreemde Oosterlingen in Nederlandsch Oost-Indië. Proefschrift. 8vo. *Leiden*, 1890, viii, 182 pp.

MAPS.

- BAWEAN.—Reede Sangkapoera. Trigonometrisch opgenomen door A. W. F. C. van Woerden 1: 20,000. *Batavia*, (Hydrogr. Bureau), 1888.
- WESTKUST BELITOENG.—Reede Tanjoeng Pandan en vaarwaters naar de reede. 1: 50,000. *Batavia*, 1890.
- JAVA.—Approaches to *Batavia*. 1: 73,000. *Washington*, (Hydrogr. Bureau), 1890.
- Blad iii Gecompileerd op het Hydrographisch Bureau te *Batavia*. 1: 500,000, 1890. *Batavia*.
- MADOERA Zuidkust. Reede Soemenep. 1: 75,000. *Batavia*, 1890.
- KAART VAN JAVA EN MADOERA.—By Dornseiffen, J. 1: 950,000 4 Bl. *Amsterdam*, 1890.

NOORDKUST JAVA.—Blad VII. Eiland Ketapang tot Duiveneiland. Trigonometrisch opgenomen door A. W. F. C. van Woerden, A. H. Hoekwater en J. W. A. F. van Maren Bentz van den Berg, 1886-1890. 1 : 200,000. *Batavia*, 1890.

—————Blad VIII. Trigonometrisch opgenomen door Z. M. opnemings vaartuigen Hydrograaf, Melvill van Carnbee en Blommendal, Com. de Luit. ter zec der 1^e kl., van Woerden, Hoekwater, en Bentz v. d. Berg, 1886-1889. 1 : 200,000. 1890, *Batavia*.

MADOERA, ZUIDKUST.—Reede Soemenep. Trigon. opgenomen door Z. M. Opnem. Blommendal, Com. Luit. 1^e kl., Bentz v. d. Berg, 1887. 1 : 75,000. 1890, *Batavia*.

MALAY PENINSULA, East Coast. Entrance to Kuantan River. 1 : 14,600. Entrance to Patani River. 1 : 18,250. Gulf of Siam, Koh Sichang harbour 1 : 13,500. *London* (Admiralty), 1889-90.

MOLUCCA.—Plannen van ankerplaatsen in den *Molukschen Archipel*. Bl. iv. *Batavia*, (Hydrogr. Bureau), 1889.

INDEX van de Kaarten van den *Nederlandsch-Indischen Archipel*. *Batavia* (Hydrogr. Bureau) 1890.

SCHOOL ATLAS van *Nederlandsch-Oost-Indië*, by W. van Gelder. Ed. 2. *Groningen*, 1890, 4to. 9 Bl.

ATLAS van *Nederlandsch-Indië*, by W. van Gelder. Ed. 2, 15 maps. *Groningen*, 1890.

WANDKAART van *Nederlandsch-Oost-Indië* by P. R. Bos, R. R. Rijkens and W. van Gelder. 6 Bl. Ed. 2. *Groningen*, 1890, with "Handleiding" in 8vo. 35 pp.

ATLAS van *Nederland* en zijne overzeesche bezittingen, by L. A. E. van der Lei. 4to. *Amsterdam*, 1890, 13 kaarten.

KAART der concessiën in het Landschap *Sambas*, 1 : 200,000. *Amsterdam* 1890.

SERMATA en Tanimber-eilanden. Plannen van Ankerplaatsen. *Batavia* (Hydrogr. Bureau), 1890.

SINGAPORE AND RHIO STRAITS.—1: 146,000: New Harbour, Singapore Straits, 1:12,170: Singapore Roads, 1: 12,170. *Washington*, (Hydrogr. Office), 1890.

NOORDOOSTKUST SUMATRA van Diamantpunt tot Hoek Tamiang. 1: 250,000. *Batavia* (Hydrogr. Bureau), 1889.

—van Hoek Tamiang tot de Broeders. 1,250,000. *Batavia* (Hydrogr. Bureau), 1889.

PLANNEN van Ankerplaatsen en Mondingen van Rivieren op de *Noord en Oostkust van Sumatra*, 1 Bl. *Batavia* (Hydrogr. Bureau), 1890.

OOSTKUST SUMATRA. Geul der Palembang of Moesi River. 1:40,000 *Batavia* (Hydrogr. Bureau), 1889.

ATLAS des Missions de la Societé des Missions Étrangères, by A. LAUNAY. Fo. 27 Maps with text. *Lille*, 1890. [Siam, 1:5,000,000. Malaka, 1:1,574,000.]



OCCASIONAL NOTES.



THE PUTRI OF MOUNT OPHIR.

The Dato' of Johol states. Putri GANDARÍA lives on Mount Ophir, but she often visits other territories, such as the watershed of the Gemencheh River.

She is seen by mankind under the following circumstances. Boatmen going down the stream will see a very ugly old woman on the bank. She has with her a cat, and slung over her shoulder is a small bag containing saffron.

The old woman shouts to the boatmen to stop and take her with them down stream. The boatmen seeing that she is old and ugly take no notice of her. In a moment the boat grounds on a sand-bank and the boatmen cannot get off again.

The old woman then cries "Let me get into the boat and all will be well." The boatmen agree, and the old woman gets into the boat. No sooner has she got in than the boat glides off down stream. They travel together for some time when the old woman says she wishes to land. The boatmen draw up to the bank and she lands, at the same time giving them each a piece of the saffron from her bag, which the instant it touches their hands turns into gold. At the same time the ugly old woman and her cat disappear.

This is how Putri GANDARÍA is seen by men, and to meet her does not forbode misfortune, but on the contrary is the signal of rare good luck.

She can transform her cat into a tiger, and people are afraid to hunt her out and molest her. The tradition is that if she is sought for or disturbed, she will certainly order her cat to assume the shape of a tiger, who will avenge her by devouring such obtrusive persons. Putri GANDARÍA does not live on the very summit of Gunong Lédang. I never heard of any one living there.

In Monsieur LEÓN JAUSSEN'S translation of GODINHO DE EREDIA'S "Account of Malacca," the Putri of Gunong Lédang is mentioned:—

" * * * * Gunoledam, where lived the Enchantress 'Putry Raynha' a female Magician who in like manner to the Sorceress ERICHTHO the Thessalonian transformed, by virtue of certain plants, women into tigers, into birds or into other kinds of animals." And again:—

"According to a fable common amongst the Malays, the Queen Putri, wife of PERMICURI, the founder of Malacca, retired to the mountain and she lives there still having become immortal by means of magic art. Her dwelling place is on the summit of the mountain in a cavern, where she reclines on a couch formed of the bones of the dead and she shows herself in the shape of a beautiful young girl dressed in silk * * * * *"

The Dato' of Johol's version is probably the aboriginal version representing the Putri as an ugly old woman with a cat. GODINHO DE EREDIA'S version is the civilized Malay version treating the lady as young and beautiful and dressed in silk in accordance probably with the ideas in similar Arabic stories.

M. L.

DIAMONDS IN THE MALAY PENINSULA.

GARCIA DA ORTA in his *Aromatum Historia*, Lib. I, p. 171, treating of diamonds says:—"Alia est rupes ad fretum Tanjam in Malacæ tractu, quæ etiam de Rupe veteri cognominatos profert. Exigui quidem sunt sed laudati: unum tamen habent vitium quod ponderosi sint, quo nomine gratiores sunt venditoribus quam emptoribus."

"There is another rock at the Strait of Tanjam in the territory of Malacca, which produces [diamonds] of the class called old-rock diamond. They are few but valuable: however they have one fault, that they are heavy, wherefore they are more valued by the sellers than the buyers."

We have not any record of late of the discovery of diamonds in the Peninsula, but where is the Strait of Tanjam? Is it a perversion of Tanjong, and if so which Cape is it?

H. N. R.

DESCRIPTION OF A NEW SPECIES OF JUNGLE FOWL, SAID TO COME FROM BORNEO.

The specimen from which this description is taken was in confinement in the Botanic Gardens, Singapore, and was obtained in 1890 from a Malay dealer who had two of them (both males) for sale, and who said they came from Borneo. I have named this species *Gallus violaceus* on account of the conspicuous violet gloss of the hackles and tail feathers, which is the most distinguishing feature of the bird. This beautiful jungle fowl resembles *Gallus varius* of Java in having only a single throat wattle and in the hackles being round ended instead of lance shaped as in the other members of the genus. It however differs from *G. varius* in having a serrated comb, and in its colouring.

Description of Adult Male.

Comb large and serrated. Throat wattle single. A yellowish patch edged with purplish blue at each side of throat wattle. Neck hackles rounded at the ends, not lance shaped, and of a beautiful metallic violaceous purple, each feather very narrowly edged with black. Lower feathers of the neck shot with metallic green. Wings.—Primaries black slightly glossed with invisible green. Secondaries black, broadly edged with dull chestnut on the outer webs. Greater wing coverts black, glossed with invisible green, the uppermost row broadly edged on both webs with deep red brown, those of the secondaries slightly edged outwardly with chestnut. Median coverts and scapularies bright chestnut shaded with black. Lesser coverts dark green edged with black and some of the feathers glossed with metallic purple. Feathers of the back

and rump bright golden chestnut broadly shaded with metallic purple and black. Tail black glossed with metallic purple, metallic green at base. Under tail-coverts olive black glossed with metallic green and purple. Feathers of the throat black streaked with chestnut. Abdomen blackish slightly glossed with purple. Feathers of and vent black tipped with chestnut. Under side of wings dark brown with a silvery sheen. Irides yellow ochre. Bill light brown, darker at base. Legs pinkish white. Claws whitish.

In the specimen described, there are one or two irregular white blotches on the quills of the wings.

Total length (living bird) about 28 inches. Wing 9 inches. Bill from gape 1.0 inch. Tarsus 3.5 inches.

H. J. KELSALL.

A LARGE MIAS IN SINGAPORE.

There was in Singapore in November last one of the largest, if not the largest, Mias that has ever been captured. It was a male and probably of the species known as *Simia satyrus*, Linn., or the Mias Pappan of the Dyaks. He was captured in Borneo, and bought by a native dealer in Singapore, who eventually sold him to a German ship's captain, by whom he has been, it is believed, taken to Germany.

As far as I could judge, his height must have been close on 4 feet 5 inches. The cage in which he was confined was 4 feet 2 inches or thereabouts in height, and he could easily touch the top of it with his head without standing erect. His face was immensely broad, the cheeks being flattened out sideways into a sort of disc. The hair was long (about 4 inches) and thick and of a bright red colour and he had a distinct short pointed beard. The eyes dark brown.

WALLACE in the *Malay Archipelago*, chap. 4, discusses the size of the Mias at some length, and concludes by saying "on the whole, therefore, I think it will be allowed that up to this

time we have not the least reliable evidence of the existence of Orangs in Borneo more than 4 feet 2 inches high." This specimen, however, was distinctly bigger than this.

The method said to have been adopted by the natives in catching this huge animal was ingenious. They pounded up a quantity of chillies into a paste, of which they made pellets. These pellets they blew by means of their *sumpitans* into the eyes of the unfortunate Mias, or Orang Utan, who rubbed its eyes to try and ease the pain, and thus made them worse and soon became quite blinded for the time being, and was of course in frightful agony. The natives then proceeded to cut down the tree in which the wretched beast was, and when it was on the ground, not knowing which way to turn from its tormentors, they seized it by the neck by means of two long poles fastened together like a gigantic pair of scissors, and thus placed it in the cage prepared for it, which was then firmly closed.

H. J. K.

NOTE ON THE NEST AND EGGS OF *NYCTIORNIS AMICTA*.

On the 22nd August, 1891, when walking along a jungle track close to the limestone cave at Kota Glanggi (near Pulau Tawar, Pahang) a Malay who was just a short distance in front of me collecting plants for Mr. RIDLEY, called my attention to a hole in the ground, which he said was a "*Sarang Burong*," or bird's nest, and he said he was sure there were eggs in it, as he had seen a green bird about the size of a ground dove (*Chalcophaps indica*) come out of it. The hole was close to the path and in the side of a slight mound. The mouth was just large enough for me to get my hand into, and oval in shape. I could not reach the end of the tunnel, so I made the Malay cut away the ground, following the tunnel, and when he had got about 30 inches from the mouth he took out two white almost globular eggs. There

appeared to be no nest proper, the eggs being simply laid at the end of the tunnel, which was nowhere more than 6 or 8 inches below ground and as nearly as possible horizontal. Mr. DAVISON, to whom I gave the eggs as soon as I got to camp, identified them as those of a bee-eater and undoubtedly of *Nyctiornis amicta*.

The dimensions of the eggs are as follows:—

1.25 × 1.15 inches.

1.22 × 1.10 inches.

The eggs are now in the Singapore Museum.

H. J. K.

NEST AND EGGS OF *HENICURUS*
RUFICAPILLUS, TEMM.

On the 23rd July, 1891, while out collecting in the neighbourhood of our camp at Kuala Tahan (Pahang Ulu) and looking out for birds up a small streamlet flowing through the thick jungle, I found a nest of *Henicurus ruficapillus* with three eggs in it. The hen bird was sitting on the nest, and flew off on my approach. I did not get a shot at her. The nest, which is shallow, cup-shaped, and 6 or 7 inches in diameter, is made entirely of moss and lined with the skeletons of leaves. It was built on to the side of an almost perpendicular rock, very much after the fashion of that of the water ouzel. Nest and eggs are now in the Singapore Museum.

The eggs are white spotted and blotched with light red and pale purplish red, the spots being larger and concentrated into a ring round the larger end. One egg was broken while blowing, the dimensions of the other two are as follows:—

0.97 × 0.66 inches.

0.90 × 0.68 inches.

H. J. K.

ON THE OCCURRENCE OF *PETROSAVIA*
IN PERAK.

During a short trip to Perak in February last, I collected a quantity of a small yellow saprophytic plant growing sporadically among the vegetation along the banks of the road

from the tea-gardens to Maxwell's Hill, and also on the Hermitage Hill. On examining it at leisure, I found it to agree very well with the description of a rare plant known as *Petrosavia stellata*, Becc., described and figured in the *Nuove Giornale Botanico Italiano*, iii 7, t. 1. BECCARI collected his specimens in Borneo, on Mount Poe, near Sarawak, at 3,000 feet altitude, and, as far as I am aware, it has not been collected there since, and it has not hitherto been recorded from the Malay Peninsula. It, therefore, forms another addition to the Bornean types to be met with in the Malay Peninsula.

Petrosavia is a slender, wiry, yellow herb with an underground rhizome, over three inches long, covered with small papery scale leaves. The stems rise from the end of the rhizome, one or more together, from three to seven inches tall, often slightly zigzag, and thickest at the base, where they are covered with numerous crowded sheathing leaves, lanceolate acuminate in shape, about $\frac{1}{4}$ inch long. The stems are slenderer above, and the leaves fewer and smaller. The flowers are arranged in a corymbose raceme, they are small, one-eighth of an inch across, yellow, and on rather long ($\frac{3}{8}$ inch) pedicels. The bracts resemble the upper leaves, and are about $\frac{1}{8}$ inch long. There are two to each flower. The sepals are lanceolate acuminate, short and small. The petals much larger, ovate, blunt, alternating with the sepals. The stamens are six in number, three opposite the sepals, and three opposite the petals. Their filaments are shorter than the petals, thickened at the base and tapering upwards. The anthers are oblong, rounded, dorsifixed, the cells divergent at the base, and splitting along the edge. The pistils are three, connate at the base, widely spreading above. The ovaries are conical follicular, tapering to the small round stigmas. The carpels are three in number, and split on the inner and upper face nearly as far as the stigmas; they each contain about twelve elliptic, oblong, brown, nodulose seeds, which are full of oil and proteids, but contain no starch. Mr. PERCY GROOM, during his residence in Singapore, has examined them carefully under the microscope, and has discovered the embryo, which had escaped all other observers. It is very minute, and resembles

that of the parasitic *Monotropa*, or Indian pipe, of the English woods.

The plant is probably saprophytic, taking its nourishment from the decaying leaves in which it grows, but it is possible that it is parasitic on some other plant or tree. It frequents the drier parts of the wood, at from 2,000 to 3,000 feet elevation.

Petrosavia seems certainly to belong to the order *Liliaceæ*, an order but scantily represented in the Malayan region, but its affinity to any other liliaceous plant is very obscure. The whole order is most plentifully represented in the North temperate region, and in Africa and Australia, but species occur in almost every part of the globe. In the Malay Peninsula we have very few genera—*Dracæna*, with a number of species; *Smilax*, four or five; *Dianella*, one species; and a very curious plant, the *Tupistra* (?) *singaporiæna*, *Bak.* found in Singapore many years ago by Dr. WALLICH, and never seen since, unless a remarkable plant, of which one specimen in fruit was found by myself and Mr. CURTIS in the *Cypripedium* valley on Penang Hill this spring, is this species. All these have berried fruit, but *Petrosavia* belongs to the capsule-bearing section of the order.

Mr. BAKER, in the Journal of the Linnean Society, vol. XVII, p. 492, and HOOKER in the *Genera Plantarum*, put it in the neighbourhood of *Tofieldia*, which in the latter work is classed with the section *Narthecix*, but it seems to me to be more nearly allied to the *Anthericææ*. It is, however, very distinct from any genus in either of these groups, and, owing to its peculiar habit (for it is the only known saprophytic liliacea), it is so modified that many of its characters throw no light on its relationship. Besides its saprophytic habit, it is almost unique in having the carpels of the capsule free from each other and spreading widely, and in the peculiar structure of the seeds. Our further researches into the Malayan flora may, we may hope, eventuate in discovering other liliaceous plants allied to *Petrosavia* which will enable us to clear up doubts as to its position.

H. N. R.

[No. 25.]

JOURNAL
OF THE
STRAITS BRANCH
OF THE
ROYAL ASIATIC SOCIETY.

JANUARY, 1894.

SINGAPORE:

PRINTED AT THE GOVERNMENT PRINTING OFFICE

AGENTS OF THE SOCIETY:

London and America, ... TRÜBNER & Co.

Paris, ... ERNEST LEROUX & CIE.

Germany, ... K. F. KOEHLER'S ANTIQUARIUM, Leipzig

TABLE OF CONTENTS.

Council for 1893, 	v
List of Members for 1893, 	vi
Proceedings of the General Meeting, 	xv
Council's Annual Report for 1892,	xvii
Treasurer's Account for 1892, 	xix
Rules of the Straits Asiatic Society, 	xxi

A Journey to the Source of the Indau— <i>by H. W. Lake</i> , ...	1
On the Dispersal of Seeds by Mammals— <i>by H. N. Ridley</i> , M.A., F.L.S., 	11
Account of a trip up the Pahang, Tembeling and Tahan Rivers, and an Attempt to reach Gunong Tahan— <i>by</i> <i>Lieut. H. J. Kelsall</i> , R.A., 	33
List of Mammals recorded from Pahang— <i>by H. N. Ridley</i> , M.A., F.L.S., 	57
A Catalogue of the the Flowering Plants and Ferns found growing wild in the Island of Penang— <i>by C. Curtis</i> , F.L.S., 	67

TABLE OF CONTENTS,—*Continued.*

Occasional Notes:—

Earthquake in the Malay Peninsula,	169
On the Occurrence of the rare Bat-Hawk in Johor, ...	171
A large Beetle caught in a Pitcher of <i>Nepenthes</i> , ...	172
The Bird-dropping Spider (<i>Ornithoscatoides</i>) in Johor,	172
Notes on <i>Gallus Violaceus</i> , *.. ..	173
A Malay Lullaby,	174

THE
STRAITS BRANCH
OF THE
ROYAL ASIATIC SOCIETY.

PATRON:

His Excellency Sir CECIL CLEMENTI SMITH, G.C.M.G.

COUNCIL FOR 1893.

His Excellency Major-General Sir CHARLES WARREN, G.C.M.G.,
K.C.B., *President.*

The Hon'ble W. E. MAXWELL, C.M.G., *Vice-President, Singapore.*

D. LOGAN, Esquire, *Vice-President, Penang.*

H. N. RIDLEY, Esquire, *Honorary Secretary.*

H. T. HAUGHTON, Esquire, *Honorary Treasurer.*

A. KNIGHT, Esquire,

H. L. NORONHA, Esquire,

C. W. S. KYNNESELEY, Esquire, } *Councillors.*

Lieut. H. J. KELSALL, R.A.,

The Hon'ble J. W. BONSER,

LIST OF MEMBERS

FOR

1893.

Nos.	Names.	Addresses.
1	ABRAHAMSON, E. E.	Sandakan, B. N. B.
2	ALLINGHAM, S.	25, Grosvenor Street, Grosvenor Square, Lon- don, W.
3	ANTHONISZ, J. O.	Singapore.
4	ATKINSON, H. S.	Singapore.
5	BAMPFYLDE, C. A.	Kuching, Sarawak.
6	BARNES, W. D.	Penang.
7	BECHER, H. M.	Singapore.
8	BEESTON, Capt. R. D.	Sandakan, B. N. B.
9	BELLAMY, G. C.	Kuala Lumpur, Selangor.
10	BERNARD, F. G.	Woodleigh, Serangong Road, Singapore.
11	BERRINGTON, A. T. D.	Selangor.
12	BICKNELL, W. A.	Audit Department, Penang.
13	BIRCH, J. K.	First Magistrate, Penang.
14	BLAGDEN, C. O.	Malacca.
15	BLAND, R. N.	Land Department, Singa- pore.
16	BONSER, The Hon'ble J. W.	Singapore.
17	BOTT, Dr. W.	The Grange, Grange Road, Singapore.
18	BRADDON, Dr. W. L.	Sungei Ujong.
19	BRANDT, D.	Europe.

MEMBERS FOR 1893,—*Continued.*

Nos.	Names.	Addresses.
20	BROWN, Dr. W. C.	Beach Street, Penang.
21	BRYANT, A. T.	District Office, Dindings.
22	BUCKLEY, C. B.	Orchard Road, Singapore.
23	BURKINSHAW, J.
24	CAMERON, Capt. M. A., R.E.	Dover, England.
25	CAMUS, M. DE	Singapore.
26	CLAIN, J.	87, Rue du Cherche Midi, Paris.
27	CLIFFORD, H. C.	Kuala Lipis, Ulu Pahang.
28	COPLEY, GEORGE	Municipality, Malacca.
29	CREAGH, His Excellency C. VANDELEUR, C.M.G.	Sandakan, B. N. B.
30	CROIX, J. ERRINGTON DE LA	Paris.
31	CURRIE, ANDREW	28, Fenchurch Street, Lon- don.
32	DANE, Dr. R.	Singapore.
33	DENT, Sir ALFRED, K.C.M.G.	11, Old Broad Street, Lon- don, E. C.
34	DEW, A. T.	Perak.
35	DIETHELM, W. H.	Hooglandt & Co., Singa- pore.
36	DOWN, St. V. B.	Singapore.
37	DUNLOP, C.	Powell & Co., Singapore.
38	EBHARDT, HANS	Hilty & Co., Singapore.
39	EGERTON, WALTER	Penang.
40	ELCUM, J. B.
41	ESCHKE, H. H.	German Consulate, Sin- gapore.
42	EVERETT, A. HART	Labuan.
43	EVERETT, H. H.	Sarawak.

MEMBERS FOR 1893,—*Continued.*

Nos.	Names.	Addresses.
44	FERGUSON, A. M., Jr.	18 & 19, Baillie Street, Fort, Colombo.
45	FEILDING, J.
46	GAGGINO, G.	Gaggino & Co., Singapore.
47	GENTLE, ALEX.	Grange Road, Singapore.
48	GOLDNEY, Sir JOHN T.	Trinidad.
49	GOSLING, T. L.	River Valley Road, Singa- pore.
50	GOTTLIEB, G. S. H.	Penang.
51	GRAHAM, JAMES	Glasgow.
52	GULLAND, W. G.	Paterson, Simons & Co., England.
53	HALE, A.	Negri Sembilan.
54	HAUGHTON, H. T.	Singapore.
55	HAVILAND, Dr.	Kuching, Sarawak.
56	HERVEY, The Hon'ble D. F. A., C.M.G.	Resident Councillor, Ma- lacca.
57	HILL, E. C. H.	Inspector of Schools, Singapore.
58	HILL, F. W.	Selangor.
59	HOSE, Right Revd. Bishop G. F., M.A., D.D. (Honorary Member)	Singapore.
60	HOSE, C.	Baram, Sarawak.
61	HOUTHUYSEN, C. L.	Batavia.
62	HOYNCK VAN PAPENDRECHT, P. C.	Singapore.
63	HUDSON, H. H.	Singapore.
64	HULLETT, R. W., M.A., F.L.S.,	Singapore.
65	IBRAHIM BIN ABDULLAH, Dato Dalam	Johor Bharu.
66	IRVING, C. J., C.M.G.	Hillands, Tiverton, Devon- shire, England.

MEMBERS FOR 1893,—*Continued.*

Nos.	Names.	Addresses.
67	JOAQUIM, J. P.	Singapore.
68	JOHOR, H. H. the Sultan of the State and Territory of, G.C.M.G., G.C.S.I. (Honorary Member)	Johor.
69	KEGAN, PAUL TRENCH, TRÜBNER & Co.	London.
70	KEHDING, F.	Laboean, Deli.
71	KEITH, Dr. A.	Bangkok.
72	KELSALL, Lieut. H. J., R.A.	Fort Canning, Singapore.
73	KENNEDY, ARCHIBALD	Batu Gajah, Kinta, Perak.
74	KEE, T. RAWSON	Johor Baharu, Johor.
75	KNIGHT, ARTHUR	Grassdale, River Valley Road, Singapore.
76	KROM MUN DEWAWONGSE VAROPRAKAR, H. R. H. Prince	Bangkok, Siam
77	KYNNEERSLEY, C. W. S.	Singapore.
78	LAKE, H.
79	LAUGHER, H.	Singapore.
80	LAVINO, G.	Dutch Consulate, Singapore.
81	LAWES, Revd. W. G. (Honorary Member).	Port Moresby, New Guinea.
82	LEASK, Dr. J. T.	Malacca.
83	LEES, F. BALFOUR
84	LEWIS, JOHN E. A.	Government Printing Office, Sarawak.
85	LISTER, Hon. MARTIN	Negri Sembilan.
86	LITTLE, R. M.	Gaya, B. N. B.
87	LITTON, G. B.	Canton.
88	LOGAN, DANIEL	Penang.

MEMBERS FOR 1893,—*Continued.*

Nos.	Names.	Addresses.
89	LOW, Sir HUGH, G.C.M.G.	England.
90	MACBEAN, W.	Straits Insurance Office, Hongkong.
91	MACHADO, A. D.	Temoh Gold Mines, Pa- tani.
92	MARTIN, Dr. L.	Mabar Estate, Deli, Suma- tra.
93	MAXWELL, The Hon'ble W. E., C.M.G.	Colonial Secretary's House, Singapore.
94	MAXWELL, R. W.	Inspector-General's Office, Singapore.
95	MCKILLOP, J.	Pulau Brani, Singapore.
96	MELDRUM, Dato'	Johor.
97	MEREWETHER, E. M.	Malacca.
98	MILLER, JAMES	Gilfillan, Wood & Co., Singapore.
99	NANSON, Wm., F.S.A.	Craigton, Tanglin, Singa- pore.
100	NAPIER, W. J.	Paterson Road, Singapore.
101	NEAVE, D. C.	Cluny Road, Tanglin, Singapore.
102	NEWTON, HOWARD	Grasslands, St. Thomas Walk, Singapore.
103	NORONHA, H. L.	Killeney Road, Singapore.
104	O'BRIEN, H. A.	Singapore.
105	O'SULLIVAN, A. W. S.	Penang.
106	OWEN, J. F.	Kuala Pahang.
107	PARKES, D. G.	Thaiping, Perak.
108	PAUL, W. F. B.	Sungei Ujong.

MEMBERS FOR 1893,—*Continued.*

Nos.	Names.	Addresses.
109	PERAK GOVERNMENT MUSEUM (Curator of)	Taiping, Perak.
110	PERHAM, The Ven. Archdeacon J. (Honorary Member)	Singapore.
111	READ, W. H. M., C.M.G.	9, Petersham Terrace, Queen's Gate, London.
112	REITH, Revd. GEO. M.	Mount Elizabeth, Singa- pore.
113	RICKETT, C. B.	Hongkong and Shanghai Bank, Foochow.
114	RIDLEY, H. N., F.L.S.	Botanic Gardens, Singa- pore.
115	RODGER, J. P.	Pekan, Pahang.
116	ROST, Dr. REINHOLD (Honorary Member).	India Office Library, Lon- don, S. W.
117	SALMON, J.
118	SARAWAK, H. H. The Raja of,	Kuching, Sarawak.
119	G.C.M.G. (Honorary Member)	
120	SATOW, E. M., C.M.G. (Honorary Member)	Monte Video, S. America Medan, Deli.
121	SCHAALJE, M.	Europe.
122	SCOTT, Dr. DUNCAN	Chop "Chin Hin," Singa- pore.
123	SEAH LIANG SEAH.	Chop "Chin Hin," Singa- pore.
124	SEAH SONG SEAH	Broadfields, Paterson Road, Singapore.
125	SHELFORD, The Hon'ble T., C.M.G.	Singapore.
126	SHELFORD, W. H.	Resident Councillor, Pe- nang.
127	SKINNER, The Hon'ble A. M., C.M.G.	

MEMBERS FOR 1893,—*Continued.*

Nos.	Names.	Addresses.
128	SMITH, H. E. Sir CECIL CLEMENTI, G.C.M.G.	Government House.
129	SOHST, T.	Mount Rosie, Chancery Lane, Singapore.
130	SOUBINDRO MOHUN TAGORE, Raja, Mus. Doc.	Pathuria Ghata Raj Bati, Calcutta, India.
131	ST. CLAIR, W. G.	<i>Singapore Free Press</i> Office, Singapore.
132	STRINGER, C.	Singapore.
133	SWETTENHAM, F. A., C.M.G.	Resident, Perak.
134	SYED MOHAMED BIN AHMED AL SAGOFF	Singapore.
135	SYED ABUBAKAR BIN OMAR AL JUNIED	Arabia.
136	SYERS, H. C.	Kuala Lumpur, Selangor.
137	TALBOT, A. P.	Government Hill, Singa- pore.
138	THOROLD, F. THOROLD	England.
139	TREACHER, W. H., C.M.G.	Resident, Selangor
140	VAN BENNINGEN VAN HELSDIN- GEN, Dr. R.	Deli, Sumatra.
141	VERMONT, The Hon'ble J. M. B.	Batu Kawan Estate, Pro- vince Wellesley.
142	WALKER, Lieut.-Col. R. S. F., C.M.G.	Thaiping, Perak.
143	WALKER, H.	Land and Survey Depart- ment, Sandakan, B. N. B.
144	WARREN, H. E. Major-General Sir CHARLES, G.C.M.G., K.C.B.,	Singapore.

MEMBERS FOR 1893,—*Continued.*

Nos.	Names.	Addresses.
145	WATSON, E. A.	Pahang.
146	WEST, F. G.	Selangor.
147	WHEATLEY, J. J. L.	Muar.
148	WILDMAN, R.	America, U. S. A.
149	WISE, E. A.	Pekan, Pahang.
150	WRAY, L., Jr.	Perak Museum, Perak.
151	WRENCH, D. T.	Singapore.

Members are requested to inform the Secretary of any change of address or decease of members in order that the list may be as complete as possible.

All communications concerning the publications of the Society should be addressed to the Secretary ; all subscriptions to the Treasurer.

Members may have on application forms authorising their Bankers or Agents to pay their subscription to the Society regularly each year.

PROCEEDINGS
OF THE
ANNUAL GENERAL MEETING
OF THE
STRAITS BRANCH
OF THE
ROYAL ASIATIC SOCIETY,
HELD AT THE
RAFFLES MUSEUM
ON
MONDAY, 23RD JANUARY, 1893.

PRESENT :

His Excellency Major-General Sir CHARLES WARREN, G.C.M.G., K.C.B., *President*; The Hon'ble W. E. MAXWELL, C.M.G., H. T. HAUGHTON, Esq., W. J. NAPIER, Esq., A. KNIGHT, Esq., Lieut. H. J. KELSALL, R.A., H. L. NORONHA, Esq., Dr. W. BOTT, C. W. S. KYNNEERSLEY, Esq., W. G. ST. CLAIR, Esq., and H. N. RIDLEY, Esq., *Honorary Secretary*.

The minutes of the last general meeting were read and confirmed.

The Annual Report was read and accepted.

The Accounts of the Honorary Treasurer were passed.

The following Officers were then elected for the ensuing year:—

President,—His Excellency Major-General Sir CHARLES WARREN, G.C.M.G., K.C.B.

Vice-Presidents,—Singapore: The Hon'ble W. E. MAXWELL, C.M.G.; Penang: D. LOGAN, Esq.

Honorary Secretary,—H. N. RIDLEY, Esq.

Honorary Treasurer,—H. T. HAUGHTON, Esq.

Councillors.—A. KNIGHT, Esq., H. L. NORONHA, Esq., C. W. S. KYNNEERSLEY, Esq., Lieut. H. J. KELSALL, R.A., and the Hon'ble J. W. BONSER.

The following new members were elected :—

Dato MELDRUM, Sir E. C. BOVILL, Kt., H. H. HUDSON, Esq., and H. S. ATKINSON, Esq.

The Secretary stated that, in accordance with a minute, he had written to the chief Printing Presses to enquire what would be the cost of printing the Journal, and that he had received two answers—one from the American Mission Press at 75 cents. a page for 300 copies, and one from the Singapore and Straits Printing Office at 2 dollars a page for 200 copies.

It was agreed to put the work in the hands of the former.



ANNUAL REPORT
OF THE
COUNCIL
OF THE
STRAITS BRANCH
OF THE
ROYAL ASIATIC SOCIETY,
FOR THE YEAR 1892.

The Council are happy to state that the affairs of the Society continue to be financially in a satisfactory condition.

The following members were elected by the Council during the past year :—

Mr. A. T. D. BERRINGTON.	Mr. J. SALMON.
Mr. G. B. LITTON.	Mr. W. H. SHELFORD.
Mr. W. D. BARNES.	Mr. D. G. PARKES.
Mr. H. LAKE.	Mr. J. FEILDING.

Surg.-Capt. F. SMITH and Mr. W. A. PICKERING, C.M.G., tendered their resignation.

During the year, Journal No. 24 was published, and materials for the next number are in the hands of the printer. The Council regret that the publication of the Journal is still in arrear, but hope that steps now being taken will result in their being able in future to publish at least two numbers annually, as was formerly done.

As the old edition of the map of the Malay Peninsula is now out of print, and there are no more copies to be had, it was

decided to obtain materials for the compilation of a new edition, and for this a Committee was formed. Examination shows that in the earlier edition there were many inaccuracies in the coast line, and a more correct one is being prepared from the Admiralty charts.

In answer to a circular asking for copies of new survey corrections, etc., the Committee have received a number of additional maps of various parts of the Peninsula, and more are promised shortly. These include maps of several portions of Pahang, hitherto blanks in the map; a map of Malacca; and the extreme North of the Peninsula on the borders of Siam; and a complete survey of Johor; so that it is hoped that the new edition may be a great improvement on the earlier ones.

A number of publications of kindred Societies were received and added to the Society's Library, and a copy of MASON'S "Burma" was presented by Mr. BLAGDEN.

Honorary Treasurer's Cash Account from 1st January to 31st December, 1892,—Continued.

		\$	c.
1892.	<i>Brought forward, ..</i>	\$	2,015 01
		\$	2,015 01
	<i>Brought forward,...</i>	\$	886 51
	To paid for setting up Journal No. 24, ...	182	00
	" " Hire of Chronometer for one month, ...	12	00
	Balance in Chartered Bank on 31st Decem- ber, 1892, ...	1,075	89
	Balance in Chartered Mercantile Bank on 31st December, 1892, Balance with Honorary Treasurer, ..	897	03
		11	58
		\$	2,015 01

SINGAPORE,
25th January, 1893

H. T. HAUGHTON,
Honorary Treasurer.

R U L E S
OF THE
S T R A I T S A S I A T I C S O C I E T Y .

—:O:—

I.—Name and Objects.

1. The name of the Society shall be "THE STRAITS ASIATIC SOCIETY."
2. The objects of the Society shall be—
 - a. The investigation of subjects connected with the Straits of Malacca and the neighbouring Countries.
 - b. The publication of papers in a Journal.
 - c. The formation of a Library of books bearing on the objects of the Society.

II.—Membership.

3. Members shall be classed as Ordinary and Honorary.
4. Ordinary Members shall pay an annual subscription of \$5, payable in advance on the 1st January of each year.
5. Honorary Members shall pay no subscription.
6. On or about the 30th June of every year, the Honorary Treasurer shall prepare a list of those Members whose subscriptions for the current year remain unpaid, and such persons shall be deemed to have resigned their membership. But the operation of this rule, in any particular case, may be suspended by a vote of the Council of the Society.
7. Candidates for admission as Members shall be proposed by one and seconded by another Member of the Society, and if agreed to by a majority of the Council shall be deemed to be duly elected.

8. Honorary Members must be proposed for election by the Council at a general meeting of the Society.

III.—Officers.

9. The Officers of the Society shall be :—
A President ;
Two Vice-Presidents, one of whom shall be selected from amongst the members resident in Penang ;
An Honorary Secretary and Librarian ;
An Honorary Treasurer ; and
Five Councillors.

Those Officers shall hold office until their successors are chosen.

10. Vacancies in the above offices shall be filled for the current year by a vote of the remaining Officers.

IV.—Council.

11. The Council of the Society shall be composed of the Officers for the current year, and its duties shall be :—

- a.* To administer the affairs, property and trusts of the Society.
- b.* To elect Ordinary Members, and recommend Honorary Members for election by the Society.
- c.* To decide on the eligibility of papers to be read before general meetings.
- d.* To select papers for publication in the Journal, and to supervise the printing and distribution of the said Journal.
- e.* To select and purchase books for the Library.
- f.* To accept or decline donations on behalf of the Society.
- g.* To present to the Annual Meeting at the expiration of their term of office a Report of the proceedings and condition of the Society.

12. The Council shall meet for the transaction of business once a month, or oftener if necessary. At Council meetings, three Officers shall constitute a quorum.

13. The Council shall have authority, subject to confirmation by a general meeting, to make and enforce such by-laws and regulations for the proper conduct of the Society's affairs as may, from time to time, be expedient.

V.—Meetings.

14. The Annual General Meeting shall be held in January of each year.

15. General Meetings shall be held, when practicable, once in every month, and oftener if expedient, at such hour as the Council may appoint.

16. At Meetings of the Society, eleven members shall form a quorum for the transaction of business.

17. At all Meetings, the Chairman shall, in case of an equality of votes, be entitled to a casting vote in addition to his own.

18. At the Annual General Meeting, the Council shall present a Report for the preceding year, and the Treasurer shall render an account of the financial condition of the Society. Officers for the current year shall also be chosen.

19. The work of Ordinary General Meetings shall be the transaction of routine business, the reading of papers approved by the Council, and the discussion of topics connected with the general objects of the Society.

20. Notice of the subjects intended to be introduced for discussion by any Member of the Society should be handed in to the Secretary before the Meeting.

Visitors may be admitted to the Meetings of the Society, but no one who is not a Member shall be allowed to address the Meeting, except by invitation or permission of the Chairman.

VI.—Publications of the Society.

21. A Journal shall be published, when practicable, every six months, under the supervision of the Council. It shall comprise a selection of the papers read before the Society, the Report of the Council and Treasurer, and such other matter as the Council may deem it expedient to publish.

22. Every member of the Society shall be entitled to one copy of the Journal, deliverable at the place of publication. The Council shall have power to present copies to other Societies and to distinguished individuals, and the remaining copies shall be sold at such prices as the Council shall, from time to time, direct.

23. Twenty-four copies of each paper published in the Journal shall be placed at the disposal of the Author.

24. The Council shall have power to sanction the publication in a separate form, of papers or documents laid before the Society, if in their opinion practicable and expedient.

VII.--Popular Lectures.

25. Occasional Popular Lectures upon literary or scientific subject may be delivered, under the sanction of the Council, on evenings other than those appointed for General Meetings of the Society.

VIII.—Amendments.

26. Amendments to these Rules must be proposed in writing to the Council, who shall, after notice given, lay them before a General Meeting of the Society. A Committee of Resident Members shall thereupon be appointed, in conjunction with the Council, to report on the proposed Amendments to the General Meeting next ensuing, when a decision may be taken.

A JOURNEY TO THE SOURCE OF THE INDAU.

BY

H. W. LAKE.



ON August 2nd, 1891, in pursuance of instructions received from H. H. the Sultan of Johor, I left Johor Baharu on the Government Steamer *Pulai*, bound for Kuala Indau. One of the objects of the party of which I was in charge was to determine more exactly, if possible, the source of the Indau River.

Existing maps, notably that published under the auspices of the Straits Branch of the Royal Asiatic Society, show the Indau as a comparatively small river, having its source a little beyond Mount Janing, in Johor Territory, and emptying itself into the China Sea in Lat. $2^{\circ} 40' 0''$ and Long. $103^{\circ} 36' 10''$.

In 1879, Mr. D. F. A. HERVEY explored the Indau Sembrong, a large tributary of the Indau proper, and afterwards published a most interesting account of his journey in the *Journal of the Asiatic Society*.

In 1875, VON MIKLUCHO MACLAY went up the Indau to Mount Janing, from thence, however, he appears to have turned seawards to Pekan.

Some few other Europeans have, I believe, penetrated as far as Mount Janing: beyond this point, however, nothing definite seems to have been known either of the course of the Indau, or the topography of the hill country where it takes its rise.

Owing to special work, which necessitated our calling at the Sedili Rivers, the *Pulai* did not arrive off Kuala Indau until August 11th.

On the North bank at the Kuala, there is a Police Station flying the Sultan of Pahang's flag, whilst on the opposite bank is Kampong Padang, the residence of the Sultan of Johor's *Naib*, or officer in charge. The coast just here is low and sandy, and a bar at the Kuala prevents steamers of any tonnage from entering the inner harbour. The village of Padang numbers about 500 Malay inhabitants, with a few Chinese merchants and shop-keepers. The trade with Singapore, which is small but growing, consists chiefly in the export of damar, rotans, getah and kapur baru (camphor wood).

On August 13th, I proceeded up-stream with a small party of Malays in *jalors* (dug-out canoes). For the first few miles the Indau averages $\frac{1}{3}$ of a mile in width, the banks are low, and the country somewhat swampy. There are numerous small clearings on either bank, but very little cultivated land. Steamers of light draught can easily run up as far as Kuala Sembrong, a distance of 30 miles from Padang.

The junction of the Indau and Indau Sembrong was reached early on the morning of the 14th. Near this point is a small group of hills, the principal summits of which, are known as Bukit Tanah Abang and Bukit Langkap. The former hill is composed of granite, with, on the lower slopes, an overburden of white and yellow clays. Here I found a little cassiterite (tin ore) in small waterworn grains.

At the Station at Kuala Sembrong I obtained the assistance of the Malay *Batin* of Jakuns, and, accompanied by him, proceeded in a North-West direction along the Upper Indau to Batu Gajah, a Jakun village on the Johor bank, about 22 miles above the Sembrong. Here a camp was established, in order to enable the *Batin* to collect a sufficient number of aborigines to take myself and party up to the source. The river at this point is shallow and winding with a pebbly bed and high banks covered with fine timber.

But little is known of the Jakuns of the Upper Indau. They are in most cases darker and smaller than the ordinary Malay, and appear to present anthropological characteristics similar to those of the Orang Sakai described by MACLAY. A certain amount of crossing has probably, rendered the Indau type

of aborigine less distinct, for instance, their hair does not curl so closely as that of the true Sakai and in some cases it is quite straight.

In the neighbourhood of Batu Gajah, about one hundred men, women and children live, three or four families sharing the same one-roomed hut. They subsist mainly on *ubi kayu* (tapioca root), fruits and fish, with a little rice which they obtain, together with salt and tobacco, from Malay traders, in exchange for rotan, damar, getah and camphor wood. I was informed by the *Batin* that during the greater portion of the year when dispersed in search of jungle products, these people live entirely on *ubi kayu* and fruit.

Most of the men carry long spears, but none of them appear to use the *sumpitan* or blowpipe, which with its poisoned darts is so common amongst the aborigines of the Keratong and Jekati Rivers.

The increasing intercourse with the Malays is most strikingly exemplified in the dying out of the aboriginal tongue, which on the Indau has become a mere dialect, two-thirds at least of the words being apparently either Malay or of Malay origin. They have not yet, however, adopted the Mohamadan religion.

On the Sungei Mas, which flows from the West and enters the Indau about nine miles above Kuala Sembrong, is another Jakun settlement consisting of about 30 men with nearly 100 women and children; they cultivate a little paddy, but in other respects are similar to those previously described.

On arriving at Batu Gajah I had some difficulty in getting the Malay boatmen to camp on the river bank, as they assured me that there were many tigers in the neighbourhood. At first I paid no attention to this and was anything but impressed when a cry of *rimau! rimau!!* roused everybody at midnight. The alarm proved a false one, having apparently originated in the fertile brain of a boatman who had supped recklessly on underdone *ubi kayu*.

Next day, however, two Jakuns were brought in very badly mauled by a tiger. They stated that they had been attacked whilst asleep on a sandbank some distance up the river; one

man's scalp wounds I judged to be of a fatal nature, the other, a youngster, was badly bitten in the forearm. They both refused to be treated by an European, and later in the day I saw them lying in the blazing sun with their wounds well smeared with wood ashes and wrapped in leaves. After this occurrence we found the tiger traps, consisting of a bamboo spear set across the paths, after the manner of a spring gun, a great deal more alarming than the idea of the tigers themselves, and that same evening a man was fatally wounded in the thigh by carelessly stepping across a trap of this kind.

I took advantage of the delay at Batu Gajah to ascend Gunong Janing. This mountain is situated on the left bank of the Indau, and is consequently in Johor Territory. The ascent commences from the river bank and is at first fairly easy. The last few hundred feet, however, is as steep as it well can be without being absolutely perpendicular, and in many places the Jakuns had to build ladders of poles lashed with rotan.

The height of Janing, determined by aneroid barometer, is 1,950 feet. As far as I was enabled to judge, the mountain is largely if not entirely composed of a coarse whitish sandstone which crops out here and there in wall-like masses. The summit is densely wooded, and it was only by building a ladder to the top of a tall tree that a complete view of the surrounding country could be obtained.

Janing appears to be the chief of a small group of hills which rise on either bank of the Indau; Kendok, a long ridge-like hill, lies opposite Janing on the Pahang bank.

Away to the South on the verge of the horizon are the Belumut hills with the two chief summits—Gunong Belumut and Gunong Chemundong—distinctly visible.

To the North and North-East lies the jungle-covered plain of the Rumpin River, with the sea beyond, whilst to the North-West a confused mass of hills and mountains stretch away as far as the eye can reach, amongst these is the source of the Indau.

With a field glass I could make out the long spit of sand at Kuala Indau with Tiunan Island in the offing.

On August 21st, I left Batu Gajah and proceeded up-stream, accompanied by 4 Malays and 15 Jakuns in a dozen small *jalors*. Personal baggage and equipment was cut down to a very few pounds in order to enable us to carry sufficient rice to last a month at least. Mount Janing was soon left behind, and at noon on the same day the first *jeram*, or rapid, was reached. Progress now became very slow, as *jeram* followed *jeram* in rapid succession; at each of these the canoes had to be unloaded and dragged over the rocks, whilst the stores were carried along the shore or borne on the heads of the natives, who were at times breast deep in the water. Many of the *jerams* are really small waterfalls over which the river rushes with considerable force. Several times the canoes were torn from the hauling ropes of rotan, and swamped, and in one or two cases stove in, I believe, so that before the close of the first day of this sort of work, there was not a single package of stores that had not been under water at least two or three times. We had now entered a hilly country and the river became rockier and more winding every hour.

The geology of this region appears to be very simple.

A granite bed rock overlaid by a series of clays and clay shales, traversed by many dykes of quartz felsite, and quartz porphyry, with here and there irregularly intruded masses of felsite, diorite, trachyte and other felspathic rocks.

The *jerams* are in most cases formed by outcropping masses of quartz felsite, some few, however, are of granite and granite porphyry. Many of the hills are distinctly conical, notably Gunong Berumbun and Bukit Tenegon, which rise abruptly from the Pahang bank.

From Mount Janing onwards the country on either side of the river, presents one mass of hilly uninhabited jungle.

On the afternoon of the 23rd, we reached the highest point navigable by small canoes. Here the Indau is simply a broad shallow stream heaped up with boulders of granite felsite and diorite, we, therefore, abandoned the boats and continued the journey on foot, cutting a path along the bank or more frequently wading in the bed of the stream; heavy loads and torrents of rain made matters somewhat trying for the next

few days. The hills now rapidly closed in on every side and the course of the stream lay in a narrow rocky ravine. At a point about 30 miles above Mount Janing, an old jungle path leads in a Westerly direction across the hills to the plain of Tenang, in the Segamat district, one long day's march for men without loads. Two days' journey towards the South-West brought us to the base of Gunong Besar and Gunong Chabang-tiga. At this point there is an extremely picturesque little waterfall, the stream falling over a mass of granite rocks nearly 40 feet high; here also are two huge boulders of granite which may serve future travellers as a landmark. Above the waterfall the Indau becomes a mere mountain rivulet with numerous tiny tributaries. The actual source is on the Eastern slope of a lofty ridge (1,800 feet above sea level) which connects Gunong Besar on the North with Gunong Chabang-tiga on the South. The formation here is a stiff yellow clay overlying the granitic country rock. On the Western slope of the ridge, or saddle back, is the source of the Segamat, which eventually empties itself into the Muar River. I spent several days in mapping this hill country, which does not appear to have been previously explored.

Gunong Besar is composed of a fine-grained grey granite traversed here and there by eruptive dykes and overlaid on the lower slopes with clays and clay shales. Height determined by aneroid 2,600 feet above sea level.

South of Gunong Besar and connected by the forementioned ridge is Chabang-tiga with its three granite peaks, the highest of which I estimate to be over 3,000 feet above sea level. To the North, East and South-East a small series of mountains stretch away to the horizon in one confused mass of jungle-covered peaks, ridges and chains. To the South-West lies the plain of Tenang, watered by the rivers Juassih, Tenang and Segamat, all tributaries of the Muar.

The range in which Gunong Besar and Gunong Chabang-tiga constitute the chief features, runs approximately North and South and is apparently separated from the hills to the far North and North-East by a flat tract of jungle country.

For topographical purposes, I have called this small system

of hills the Tenang range, the larger portion of which lies within the Johor frontier.

On September 1st, we crossed the Tenang Hills into Ulu Segamat, from this point my work took me northwards towards Keratong and the Jekati River. Five weeks later I returned to Johor Baharu by way of Muar and the West coast.

Itinerary from Kuala Sembrong to Ulu Indau.

I have compiled the following from my journal and sketch map in the hope that it may prove useful to future travellers.

The mileage given is, of course, only approximate, still I think that the error is well under 10%. I have not thought it necessary to give any of the more precise topographical data, as it would be of little value here unless accompanied by a map of Johor. In every case when the mileage is given, it refers to the distance above Kuala Sembrong and not Kuala Indau.

Mas River.—Left bank of Indau (ascending) at 8 miles above Kuala Sembrong; mouth about 50 feet wide; source in Johor Territory. Jakun village about 2 hours' journey up.

Lemakau River.—Left bank at 18 miles; mouth about 20 feet wide; source in Johor Territory.

Batu Gajah.—Jakun village on left bank at 22½ miles.

Mount Fasing.—On left bank at 24¾ miles; height 1,950 feet; path leads up South slope to summit. Kendok hills on right bank directly opposite, highest point navigable by large canoes.

Fasing River.—Left bank at 29 miles; source in the neighbourhood of Mount Janing; Kuala about 50 feet wide, very rocky.

The first *jeram* or rapid occurs here.

Bukit Tenegon.—A conical hill near the right bank.

Keng Kim River.—Right bank at 30¼ miles; source in low hills, one day's journey distant towards the Rumpin River.

Kanu Stream.—Left bank at 32¾ miles; Gunong Berumbun on right bank some little distance inland.

Temapan River.—Right bank at $34\frac{1}{2}$ miles; mouth about 45 feet broad, very rocky; source unknown.

Danoy River.—Left bank at $35\frac{1}{2}$ miles.

Lawing River.—Left bank at $36\frac{1}{2}$ miles.

Highest navigable point of Indau at $37\frac{1}{2}$ miles; river very rocky. Bukit Salokris on left bank; from the Jasing River to here there are 15 distinct series of rapids of granite and porphyry.

Kemupoy River.—Left bank at $38\frac{1}{2}$ miles; source in Bukit Batu Dandan; mouth about 40 feet wide, very shallow and rocky.

Kidir Stream.—Left bank at $40\frac{3}{4}$ miles; source probably in Batu Dandan.

Granite Rocks.—Right bank at $42\frac{3}{4}$ miles, $\frac{1}{2}$ mile further up a stream comes in from the S.E. (slopes of Chabang-tiga).

Source of Indau at $44\frac{3}{4}$ miles on a ridge about 2,800 feet above sea level.

The Tenang Hills.—The following are the chief features of this system.

The bearings and approximate distances are taken from the summit of Gunong Besar.

Gunong Besar.—Point of observation 2,500 feet high.

Bukit Salokris.—Bearing E. distance 5 miles from Gunong Besar.

Bukit Batu Dandan.—Bearing S.S.E. distance $2\frac{3}{4}$ miles, height 1,830 feet.

Gunong Selai.—Bearing South, distance 2 miles; source of Selai River a tributary of the Indau Sembrong.

Gunong Tiang.—Bearing South, distance 3 miles; height 2,230 feet.

Gunong Tenang.—Bearing South, distance $4\frac{1}{2}$ miles; source of the Tenang River, a tributary of the Muar.

Gunong Chabang-tiga.—Bearing S.S.W.,

distance $1\frac{1}{2}$ miles; 3 distinct peaks,
height about 3,000 feet.

Gunong Pukin.—Bearing N.N.W., distance
 $1\frac{1}{2}$ miles; source of Pukin River, a tri-
butary of the Keratong.

Length of the Upper Indau from the source to Kuala Sem-
brong, $44\frac{1}{2}$ miles, general course Easterly.

Length of the Indau from Kuala Sembrong to the sea, 30
miles, course Southerly.

Total length of river about 75 miles.

ON THE DISPERSAL OF SEEDS BY MAMMALS.

BY

H. N. RIDLEY, M.A., F.L.S.



THE relations of animals to plants in the matter of fertilization has been the subject of many hundred papers and books written by various observers, especially since DARWIN published his well-known researches. But the various modifications and adaptations of the seed and fruit for distribution by animals, although of almost equal importance in the evolution of new forms, has been very much neglected. No one can avoid being struck by the observation that there are a very large number of plants in some orders, which closely resemble each other in the form and colouring of the flowers and yet differ very materially in the fruit. In many of these cases it is the necessity of special adaptation for dispersal of the seed that is the cause of the various modifications of the fruit or seed. Seeds are, as is well known, dispersed by the aid of animals, either by being swallowed by them and afterwards passed from the body at some distance from the parent plant, or by adhering to their fur or feathers and so being borne away, or by being thrown to a distance by them, as will be explained later on. Or again they may be dispersed by the aid of wind or water, being in the first instance blown far from the tree, and in the latter case being drifted away by sea or river currents; and lastly they may be scattered by merely mechanical means, as in the explosive capsules of the Castor-oil (*Ricinus communis*), and other Euphorbiaceous plants, or by merely rolling by their own weight when falling from the top of a lofty tree.

I may here call attention to a fact which has been overlooked by some naturalists in dealing with this subject, namely, that it is of no advantage to a plant to have its seeds borne to great distances but that on the contrary it may be positively injurious. In the first place the plants may be partially or wholly unisexual, the male and female flowers being on different trees. This is especially common among East Indian jungle trees, notably in the orders Euphorbiaceæ, Sapindaceæ, and Myristicaceæ.

Now if we suppose that a flock of pigeons have swooped down upon a nutmeg tree in fruit and swallowed a quantity of the seeds, and then flown away in all directions for, say, a hundred miles before passing the seed, the plants will eventually be at so great a distance apart, that it will be impossible for the flowers of the female trees to be fertilized, and such isolated trees can never reproduce themselves.

The same observation applies to plants which require a special fertilizing insect. If the seed is borne to a district where the fertilizer does not exist, it is to all intents and purposes destroyed. In this case, however, it is possible that another fertilizer may be found who can do the work fairly well, and indeed it is rare that any plant depends on a single species of fertilizer. And lastly there is always a risk of the seeds being deposited in an unsuitable locality, if borne too far away. This, however, is obviated by the conservative habits of the agents, thus fruit pigeons which always inhabit thick jungle, even if they did travel a long way with the seeds, inasmuch as they naturally fly to thick jungle, would almost certainly drop the seed in a locality similar to the one they took it from, and in the same way wading birds carrying seeds adhering to their feathers, would in their long migratory flights pass over jungles and deserts and only stop at pools or swamps where the seeds might get planted.

In the matter of wind-dispersed seeds also it is not difficult to show, especially in the flora of the big jungles, that travelling for a long distance is of no object and indeed does not seem to be aimed at. The object aimed at is rather to sprinkle the seeds at such a distance from the parent tree that the roots of

the latter may not interfere with the growth of the young plant, but not too widely to be absolutely isolated, nor on ground unsuited to the requirements of the species. It must be pointed out also that in the thick jungle where the foliage is very dense there is always a risk of the fruit when fallen resting in the branches, or in the tangled mass of creepers that mats together the branches of the big trees. It is, therefore, important that the fruits or seeds should be in many cases modified so as to avoid this accident. Besides in the case of such palms as *Pholidocarpus* and *Elaeis* where the stem is roughened by the projecting bases of the fallen leaves, there is always danger of the seeds resting in the spaces between these leaf bases and the trunk. This is usually obviated by the fruits being made edible and sought by birds, or mammals, as in the date-palms (*Phoenix*) or by the inflorescence being elongate so that the fruit is held out from the tree on long branches as in *Pholidocarpus*. The oil-palm (*Elaeis guineensis*) is not a native of Malaya, though often cultivated, and the spadix of fruit is short-stalked and hidden among the leaves, so that, unless some animal or bird devours the fruit, there is a great risk of the seed lodging among the leaf stalks, and as no bird here seems to care for the fruit, this is what often happens in trees in the Malay Peninsula. I have seen trees in the Botanic Gardens in which the spaces between the old leaf bases and the stem at the top of the trees were full of seedling oil-palms which, of course, would soon perish from want of nourishment.

In the Malay Peninsula, as elsewhere, the birds play the most important part in the dispersal of seed, but the mammals are agents also of considerable importance, and in this paper I intend to speak of their action in this work, and to compare it with that of the birds.

In the dense jungles that cover the hills of the Malay Peninsula one very soon notices that bird life gets scantier and scantier the further one penetrates them. Pigeons, hornbills and finches become scarcer, and almost entirely disappear, and even monkeys are not so abundant as one would expect.

In the woods of the Tahan valley in Pahang I have seen the

ground in some places strewn with fruits of various big trees absolutely untouched by animals or birds. In such spots the struggle for dispersal must be very great, and there must be a very large number of seeds wasted. Many trees and climbers in the dense jungle fruit very heavily and one frequently finds (especially where monkeys are absent) the ground beneath these plants almost carpeted with seedlings a short time after the fall of the fruit, but on visiting the same spot a few weeks later only a few of these young plants are to be seen. Nearly all have perished, partly from overcrowding and partly from absence of light. Nor does the waste of young plants end here, for a very large proportion of those that do become trees can never push their branches through the dense mass of older trees so as to be fully exposed to the light, when alone they can flower.

The contrast between the woods of the colder climates and the jungles of the tropics is most strongly brought out by the wonderful disproportion of species in a given area. In cold climates one frequently sees woods consisting almost exclusively of one species of tree such as the beechwoods of Southern England, and the firwoods of Norway, but in a Malay forest all the trees appear to be of different species.

Indeed it is only in exceptional places which are suited to a limited number of species (such a spot, for instance, as a mangrove swamp) that one sees a large number of individuals of one species together. In the jungles, which are suited to the requirements of a great variety of species, the different individuals are isolated, for here the ground is already so thickly covered with older trees and shrubs, that there are but few vacancies to be filled up. And thus of the immense number of seeds which fall from the trees, but few can find vacant spots on which they can develop into trees.

The assistance of the wind or of the mammals or birds which dwell in the forests is used to fill up these vacancies.

The plants which make use of animals to disperse their seeds either possess juicy or fleshy eatable fruits of which the seeds are passed through the bodies of the animals unhurt, or dry nuts, or again the fruits may be adhesive either by some viscid

material or by hooks or recurved bristles by which they adhere to the fur and are thus borne away.

The following are the mammalia of the Malay Peninsula which eat fruit, and probably all act to a certain extent, as some do to a very large extent, as seed dispersers :—

Quadrumana:—*Hylobates* (Wau-Wau), two species; *Semnopithecus* (Lotong), one or more; *Macacus*, two or three species; Cheiroptera (bats); *Pteropus* (Fruit-bats), two or more species. Insectivora:—*Tupaia*, several. Carnivora:—*Viverra* (civetcats), several species; *Arctictis* (Binturong) and probably *Hemigale*; the bear *Helarcios malayanus*; Rodentia *Sciurus* (Squirrels), several; *Sciuropterus* (Flying Squirrels), *Mus* (Mice and Rats), probably several. In addition to these must be added the deer, the wild pig (*Sus indicus*), the elephant, and probably the wild ox (*Bos gaurus*) and the buffalo (*Bubalus aruco*), which disperse seeds of grasses and other herbaceous plants both by swallowing seeds in the herbage they eat, and by bearing, attached to their skins, the adhesive fruits and seeds.

Quadrumana.—The monkeys haunt the big jungles wandering about in small flocks and avoiding the open country. They live chiefly on fruit, and of these they generally attack the juicy or succulent kinds, although they will occasionally eat the chestnuts and other dry fruits. The commonest species in Singapore is the Kra (*Macacus cynomolgus*) and this is one of the most important of the seed-dispersing mammals. The fruits I have chiefly seen it devouring are those of various species of *Eugenia*, *Baccaurea*, *Mangifera*, *Willughbeia*, *Dialium*, *Trichosanthes*, *Nephelium*, *Careya*, *Strychnos*, *Elæocarpus*, *Randia*, *Calophyllum*, *Gardenia*. And here I will point out that the fruits eaten by mammals are usually plain green in colour and inconspicuous, and frequently of considerable size, whereas those which are intended for dispersal by birds are often scarlet, orange or yellow, more rarely blue or white and small. The reasons for this are, that the mammals move comparatively slowly through the jungle searching the fruit close at hand, for it is impossible for them to see far into the thick mass of foliage, whereas the birds moving more

rapidly and at a higher elevation can detect fruit, if coloured, at a great distance and can speedily make their way to it, and that besides many of the fruit-eating mammals are nocturnal and, therefore, colour would be useless to them in the dark.

Berries and drupes are eaten whole (except for the skin) by the monkeys, if they are small, like *Nephelium*, *Zizyphus*. In these fruits the sweet pulp often adheres strongly to the stone making it so slippery that it is almost impossible to avoid swallowing the latter. *Zizyphus calophyllus*, a common creeper with small globose fruit, and the Mata-kuching (*Nephelium malaiense*) are good instances of this form of drupe. In *Baccaurea motleyana*, Hook., the Rambai, the seeds, of which there are three in a fruit inclosed in sweet pulp, are very thin, and are quite troublesome to eject.

The Malays and the Sakais in eating these fruits generally swallow the seeds even of such large drupes as the Rambutan, and I have seen in the deserted encampments of the Sakais in Pahang germinating seeds of the Rambutan which had been swallowed and had passed through their bodies. The Malays indeed say that this is the most wholesome and pleasant way of eating these fruits.

There are two forms of the fruit of the Polessan (*Nephelium mutabile*), in one of which the flesh adheres tightly to the stone, and in the other it is firmer, and readily breaks away, and can be nibbled off easily. If a monkey ate the cling-stone variety the seed would slip down its throat, while from the firmer-fleshed free-stone variety it would nibble the flesh and throw the stone away. In drupes of this kind it is essential that they should not be too large for an animal to swallow, and there is a decided advantage in the sweet pulp being very thin as it is thus more slippery and cannot be detached by biting. Many large and heavy fruits like those of the wild mangoes (*Mangifera cæsia*, *lagenifera*, etc.) and *Careya* are carried by the monkeys who gather them to a convenient perch to be eaten, and in doing so they frequently drop them, so that one finds large fruits partially eaten often at considerable distances from the parent tree. But the weight of these fruits has also another advantage, by preventing their lodging in the tangled

mass of creepers and thick foliage which often mats the tops of the highest trees together, and causing them to fall and roll to some distance.

The *Willughbeias* are lofty climbers in the jungles. They have large dull green or yellowish inconspicuous fruits, pear-shaped or globose. The seeds are soft and enclosed in a sweet and eatable flesh, of which the monkeys are very fond. There are a number of seeds in a single fruit, and if, as sometimes happens, the fruit falls whole on the ground the seeds all germinate in the fruit, so that one finds a small cluster of plants where a fruit has fallen and decayed. After these have grown a few inches, all or nearly all perish from overcrowding, and in localities where monkeys are scarce I have seen the ground covered in places with seedlings, of which in a few weeks none are left. If, however, a flock of monkeys visits the locality when the plant is in fruit, all the best fruits are speedily devoured. Monkeys like other mammals are very greedy eaters, and when there is plenty of fruit on a *Willughbeia*, they do not entirely finish each fruit, but tear it to bits and scatter the seeds in different directions, only eating bits of it. The seeds of the *Willughbeia* are soft and comparatively tasteless, or if they have a flavour it is bitter and unpleasant, but a great many do get bitten up and destroyed by the monkeys, even if a considerable number are detached and scattered about, unharmed.

It has been pointed out to me that *Willughbeia* seed to be satisfactorily planted must be thrown at the base of or near a big tree so that the creeper may have a support to climb on, and this is brought about in the following way. A monkey seizes a fine fruit to eat, the others of the flock immediately rush at him to take it away, he scrambles into the nearest big tree and getting into the fork or behind a big branch devours it with hurried bites for fear of being robbed, throwing the seeds at the foot of the tree up which the young plant can eventually climb.

Nearly all the *Willughbeias* and the plants of the allied genera *Melodinus* and *Leuconotis* are climbers in thick jungle, and in this case the fruits are, as above said, green, or at most

tinted a little with yellow or orange. Those, however, that grow in more open country or on the exposed edges of jungles have bright yellow or orange coloured fruits.

In the heathy country bordering the Pahang River, I found a species of *Willughbeia* which bore exceedingly pleasant, small, oval fruits of a bright apricot-yellow colour and very conspicuous. It grew in low thickets in open sandy country, where monkeys do not go, as they have an objection to travelling far on the ground on account of the risks from tigers, wild cats, dogs and other enemies. This conspicuous fruited *Willughbeia* had probably developed its showy colour to attract birds, of which there were many large fruit-eating kinds, and the reduction in size of this fruit is also an assistance in dispersal as even the hornbill can hardly manage to carry a globose fruit as large as that of *Willughbeia edulis*.

The various species of *Dialium*, known to the natives as Kranji, are big trees with ovoid black pods, each containing one hard seed which is enclosed in a somewhat acid but pleasantly flavoured pithy substance. The monkeys are very fond of these and one often sees the remains of the fruit on the ground. The fruit is unfortunately relished by the monkeys before it is ripe so that very often the whole crop is gathered green by them and so destroyed, and here I may call attention to the value of acidity of unripe fruits in preventing animals from eating them too soon, which would soon exterminate the trees by destroying the seeds.

Though many of the fruits eaten by animals are sweet or pleasant to our taste, a large proportion of those very popular with monkeys are either tasteless or nauseous—often astringent in flavour to us. Some may even be poisonous as *Strychnos*.

Cheiroptera.—There are several kinds of fruit-eating bats in the Malay Peninsula, but of their habits little is known. The largest kind, *Pteropus edulis*, is very irregular in its appearance. In some years there are hardly any to be seen in Singapore, but some years ago there were enormous numbers roosting every day in the Garden jungle. They fly great distances and may be seen far out at sea. They eat great quantities of fruit of different kinds. *Cynopterus marginatus*

is a small and very common species which during the day hides in the leaves of plantains, or palms, or, when it can, in caves. It eats fruits voraciously, especially figs (*Ficus Miquelii* and *Benjamina*), and I have seen it, or allied species, in great numbers at the Tembusu tree (*Fagrea fragrans*), *Livistona australis*, the *Nepheliums*, the Chiko (*Achras sapota*) and other trees.

It is difficult to see how these animals detect the fruit on the trees in the night, but perhaps they use the powers of smell. In any case it must be easier for them to find the fruits than for the insectivorous bats to see and catch insects at night. It appears to me that fruit-eating bats are much less intelligent than the insectivorous kinds. They are slower in flight and more constantly caught in house at night being utterly confused by the light, whereas insectivorous bats fly readily in and out.

The Malays keep these animals off from the fruit trees by attaching to the boughs the thorny *flagella* of the rattans which are collected and sold for this purpose, so that the bats in flying to the fruit get their wings entangled and torn by the thorns.

Viverridæ.—Of the greater number of the species of this group little or nothing is known as to their habits. All appear to be omnivorous, and certainly eat a very large quantity of fruit. They are nocturnal and find the fruit probably by the scent. The common civet cat or Musang (*Viverra malaccensis*) is a most destructive eater of cultivated fruit.

I have seen in its excreta the seeds of coffee, *Gnetum scandens*, *Caryota Cumingii*, *Mimusops elengi*, and it is also very partial to the fruits of *Artocarpus rigida* and *integrifolia* (the Jack), *Diospyros discolor*, *Achras sapota*, and many other fruits.

Its habits of selecting the best coffee berries to eat is well known to planters, who often collect the seeds passed by the Musangs for cultivation, as giving stronger plants. It has a habit of dropping its excreta on open spaces, especially paths, so that the seeds passed by it can very readily grow.

The Binturong (*Arctictis binturong*) lives much on fruit.

•

One kept in captivity ate papayas (*Carica papaya*) and Rambais (*Baccaurea mottleyana*, Hook.) swallowing the seeds and passing them apparently uninjured.

Ursidæ.—The common bear (*Helarctos malayanus*) is a nocturnal fruit-eater, and is particularly fond of Durians. The fruit of the common Durian (*Durio zibethinus*) when ripe falls entire upon the ground, and when found by the bear is torn to pieces and the seeds scattered about. The bear, however, also ascends the tree as well and helps itself to the Durians, which grow, as is well known, on the thick branches, where it can get at them.

In the wild Durian (*D. Oxleyanus*) the fruit splits on the tree and lets the seeds fall. In one wild Durian which I saw at Pekan, the flesh of the seed was pink and the capsule split so as to show the bright colour. It grew in a part of the country where there were no bears and was probably dispersed by some such bird as the hornbill.

I gave a bear a wild Durian (*Durio oblongus*) to eat. It tore it to bits with its paws and ate the aril of the seed and a good deal of the placentas, but would not eat the seed and spit it out so that it fell some way off. Another bear seeing the seed fall bit it, but did not like the taste and would not eat it. There was no particular taste to the part that the bear ate, though it was very eager to eat it, nor had the rejected seeds any taste that seemed objectionable to me.

I will here digress a little to compare the several forms of fruit of the *Durionæ* showing their various modifications and their meaning with respect to dispersal.

In the genus *Durio*, the fruit is a thorny capsule, dull brown or green, and the seeds have a white or pink edible aril.

Durio zibethinus, L. The fruits are borne on the strong branches, on short stalks, and are strongly scented so as to be easily found at night, and are dispersed by the bear, a heavy nocturnal animal.

D. oblongus, Mast. The fruits resemble those of *zibethinus*, but are not scented. They open on the tree, and eventually drop the seed. They are probably dispersed by birds, as the

plant is a native of Singapore where there are no bears.

D. sp. has a scentless fruit with a conspicuous red aril on the seeds. There were no bears in the locality, so that it was probably dispersed by hornbills, which were abundant.

D. testudinarum, Becc., has the fruit at the base of tree. This is called Durian Karkura (Tortoise Durian) in Borneo by the natives, according to BECCARI, and is perhaps eaten by these animals.

Neesia synandra, Mast. This has a very hard woody capsule, bluish grey, not or hardly armed. Seed black, with a small yellow waxy aril. The fruit drops whole, and the seeds are protected till ripe by a quantity of pungent irritating yellow bristles lining the inner walls of the woody capsule. The aril and sometimes the whole seed is eaten by mice. (The seed falls out of the capsule when the capsule falls.)

Cælostegia Griffithii, Mast., has a very large rather round capsule covered with thorns, and of a showy orange colour. The seeds are chesnut colour and conspicuous, with an orange waxy aril. They do not fall out of the fruit, but fall with it. I imagine that they are dispersed by the agency of birds, but I found rats very partial to them.

Boschia Griffithii, Mast. A very small Durian with a conspicuous scarlet capsule, splitting on the tree, and exposing the black conspicuous seeds. The form and colouring of this remind one of the fruits of *Sterculia lævis* which is dispersed by birds. The fruits are borne on small twigs, and I have little doubt but that the seeds are scattered by birds. BECCARI gives Durian Tupai as one of the native names of the plant in Borneo, this would mean Squirrel Durian, but does not imply that squirrels eat it, *tupai* being really equivalent to *tikus* (mouse) meaning of small size, as opposed to *gajah* (elephant) which means in speaking of fruit, &c. large. Thus *Commersonia platyphylla* anders is called in Singapore Durian Tupai because the fruit looks more or less like a very small Durian.

Insectivora.—*Tupaia ferruginea* and other allied species although belonging to the order Insectivora are chiefly frugivorous. They appear to eat the smaller fruits such as those of *Marlea nobilis* and *Eugenias* which have a firm texture and

are not very hard.

Ungulata.—The Ungulates of the Malay Peninsula include the elephant; rhinoceros, one or two species; tapir, wild ox (*Bos gaurus*); deer, one or two species; mouse deer, two or more species; and the wild pig. To which must be added as a seed disperser the buffalo (*Bubalus arnee*). These animals act more as scatterers of seed attached to their hair or hides, but probably also, to a certain extent, by swallowing grass-seeds in the herbage. The first four are inhabitants of the densest jungles, especially of the hill regions, and feed chiefly on the bushes and leaves of trees. They make long tracks through the dense forests, and wander often to great distances. I have seen many seedlings, apparently of some small herb, springing up in dung of elephants dropped in their tracks. The wild ox lives, to a small extent, on fruit. One brought down to Singapore ate greedily the fruits of the Sentol (*Sandoricum indicum*).

Scoparia dulcis L., is a small herb introduced accidentally from South America which has been widely scattered by the water buffalo. In Pahang, I traced it up the Pahang and Tembeling Rivers as far as the buffalo went. On sandbanks in the river where for some reason buffaloes had not gone this plant was absent, and I saw it and also *Cleome viscosa* springing from masses of buffalo-dung, in several places. Many of the smaller herbs and especially grasses and sedges must be distributed by this animal in this way, and *Fimbristylis miliacea*, a sedge very abundant in marshes where these animal go, is called by the Malays *Rumput Tahi Kerbau* (buffalo's dung grass) for this reason.

Rodentia.—The important seed distributors in the family are the rats and the squirrels.

The rats and mice of the Malay Peninsula are as yet very little known. I have seen at the foot of Mount Ophir, in dense jungle by a stream, a large reddish rat eating the fallen fruit of a wild species of mango, of which it might easily have borne off fruits to its holes under the boulders to some distance from the tree.

There are a great number of herbaceous plants, the fruits

of which are more or less concealed among the leaves or in the ground. Such is *Curculigo sumatrana* which has small inconspicuous sweet fruits with very small seeds. These disappear as soon as ripe, and are certainly eaten by some rodent. The *Scitaminceæ* again have fruits which are much sought by these little animals. Most of the jungle loving species have the fruits at the base of the stems as *Amomum*, *Zingiber*.

The fruits are inconspicuous, but in many cases the bracts which enclose them are red. This colouring, however, bears rather a relation to the floral stage of development and is intended to make the flowers more conspicuous to the insect fertilizer. When the plant is in fruit, the bracts have usually become shabby and inconspicuous.

Nicolaia hemisphærica and *Amomum laterale* are two species which have plain green fruits, in the former in a head on a short, stout stem, in the latter in a stout, cylindric, lateral spike about a foot above the ground. These fruits are devoured by some rodent (probably a squirrel) as soon as they are ripe.

There is a great contrast between the fruits of these jungle gingers where the inflorescence is a compact head and radical or low down, and those which live in more open country and possess terminal inflorescence. In the former the fruits are inconspicuous and often green, while in the latter they are either showy and orange as in the *Alpinias* of the river banks, or they are white in the plants of the open jungle as in the case of *Clinogyne* and *Alpinia galanga*.

The squirrels (*Sciurus*) probably disperse more seeds than the *Muridæ*, and being diurnal can more easily be seen at work. They do not, as a rule, eat sweet or juicy fruits, but those of firmer texture, as those of the Daroo (*Sideroxylon sundaicum*), *Marlea nobilis*, and *Pyrenaria acuminata*. All these are inconspicuous, small, green fruits containing hard seeds, and it is very common to find gnawn fruits lying some way off from the trees, usually with the seeds uninjured. In many cases a tree is completely denuded of fruit as fast as it is ripe, and the squirrels carry it so far that it is impossible to find any. To some of the introduced fruits they are very

destructive especially to cocoa (*Theobroma cacao*) and to coco-nuts, destroying the latter by biting round holes into the fruit and eating the interior, so that trees near jungle, if unprotected, lose all their fruits.

But it is on the fruits of oaks and chestnuts that these animals chiefly live. These trees fruit very heavily, more so than any class of tree here, and the ground beneath an oak in fruit is often covered with acorns. The chestnuts (*Castanopsis*) nearly all have their fruit arranged in close spikes and usually covered with a prickly involucre. The whole spike readily breaks off the tree, but it is difficult to separate the individual chestnuts. A squirrel seizes a spike and breaks it off, and holding it in its paws attempts to nibble through the prickly husk to eat the fruit and it often happens that owing to the prickles being too sharp for it, it drops the whole spike before it has succeeded in eating more than one nut.

The squirrels invariably, if possible, when they have gathered one of these fruits run to a short distance to eat it conveniently, so that the nut or acorn may be carried to some distance before it is dropped. The big *Sciurus bicolor* is an entirely arboreal squirrel living in very dense jungle and very rarely if ever coming down to the ground, and when it takes a fruit it runs to a suitable spot to devour it. It sits transversely on the bough, holding on with its hind feet, its head and fore-arms hanging down over the bough on one side and its tail on the other. In this position it is very likely to drop a nut either too prickly or too smooth for it to hold fast. The smaller squirrels (*Sc. notatus* and *Sc. tenuis*) when they descend the trees to pick up the fallen acorns or chestnuts, which *Sc. bicolor* never does, always run up an adjacent tree to eat them, and I have frequently seen one carry an acorn in its mouth for some distance before eating it. I recently saw a small red-bellied squirrel (*Sc. notatus*) eating the fruits of an *Elæocarpus*. When it took a fruit, it hung head downward from a bough by its hind feet only. *Sc. tenuis* too usually hangs from the trunk of a tree by its hind feet head downwards when eating acorns. As there is no season here when a squirrel cannot get food, it never stores up

seeds in holes as the English squirrel does for the winter. Among the oaks, which are more abundant than the chestnuts, there seems at first sight even less protection for the seed or means for its dispersal than for the latter, but there are certain slight modifications which have a most important effect in these matters. The acorn is nearly always quite exposed, and the cup, which corresponds to the involucre of the chestnut, is not armed, although in some species it is roughened with soft hooks (*Quercus hystrix*).

If one examines the acorns which have fallen from a tree where there are many squirrels, one notices that they are all nibbled at the base, and there are often marks of teeth as of ineffectual bites on the sides. I gave a *Sciurus bicolor* some acorns of *Quercus lucida*, a large rounded acorn with a thick but shallow cup. Taking them between its paws, it made an attempt to bite into the side of the acorn, but the outer coat was so smooth that its teeth slipped and it could not get a hold. It then turned the acorn round and bit the cup, and the acorn immediately fell out of the cup and rolled away. Had it been up in a tree when it tried to eat the acorn, the fruit would have fallen down and rolled perhaps far from the parent tree. On giving it the acorn again it began to bite the rim at its base, but it was clear that the smooth polished surface of the fruit was too slippery for its paws, and even on the floor of its cage it had some trouble in holding it.

Many of the acorns have a fairly firm outer coat thus polished, and fall very readily from the cup when ripe, but some such as *Q. encleisocarpa*, and *Q. Cantleyi*, two of our commonest species, have an improvement on this. The acorn is coated with a very fine silk, which has almost a greasy feel. It is not at all easy for a squirrel to hold these in its paws to eat, and it is very common to see the acorns of the former scattered all over a wood in which there is a tree in fruit, and nearly all of these bear the marks of squirrels' teeth, but for all practical purposes are unhurt. *Q. encleisocarpa* has the cup in the form of a thin brown covering, from which, though it is much cracked and split when ripe, the acorn never falls. In *Q. Cantleyi* the acorn is readily detached

from its cup and is most difficult for a squirrel to hold. It can take it in its mouth by the aid of the raised circular rim at the bottom and thus carry it off, but to eat it it must hold the slippery conical portion in its paws so as to nibble at the base, the only place where its teeth can get a purchase, and it naturally lets many of these silky-coated acorns slip unhurt from its grasp.

This method of dispersal is a very expensive one, a large proportion of seeds being destroyed by the squirrels, compared with those that are deposited by them in suitable positions for development into trees, but so large is the crop produced at one time that the number safely planted is quite sufficient to keep up the stock. It must be remembered too that it is necessary for the trees to supply enough nuts to tempt the squirrels. If the squirrels did not get enough off the trees to eat or these seeds were so well protected that they could not get at them, they would be less likely to visit the trees at all and indeed where there are few or no squirrels, as in the hill forests, oaks and chestnuts are much more scanty than in the low country where they abound.

WALLACE (*Tropical Nature*, ed. 1891, p. 400,) says of most of the plants whose large seeded nuts cannot be eaten without destroying their germinating power:—"It is a suggestive fact that they are among the most ancient of known dicotyledonous plants—oaks and beeches going back to the cretaceous period with little change of type so that it is not improbable that they are older than any fruit-eating mammal adapted to feed upon their fruits."

Still in the prickly husk of the chestnut, and in the smooth polished or silky exterior of the acorns, we see that these fruits have undergone modifications by which the little enemies have been prevented from exterminating the trees, and have been utilized as dispersers of the seeds.

Very much remains to be observed still as to the action of fruit-eating mammalia as seed-dispersers. Many of them are difficult to watch in a wild state on account of their shyness and nocturnal habits, and even in Malaya there are several such as the Loris (*Nycticebus tardigradus*), the Galago

(*Galeopithecus volans*) (which, however, certainly lives very largely and possibly altogether on leaves), the Binturong (*Arctictis*), the larger *Viverras*, *Hemigale*, *Paradoxurus*, and the fruit-bats, of the habits of which in the wild state little or nothing is known. But I think it may be noticed that the fruits especially sought by mammals are inconspicuous and dull-coloured and not brilliant. This, however, must not be held to imply that bright colours are not visible to the diurnal mammals such as monkeys. Sir JOHN LUBBOCK has used the argument of the coloured fruits forming the food of monkeys against the theory of MAGNUS, GEIGER and GLADSTONE that the ancients were colour blind, saying that "if monkeys and apes could distinguish colours surely we may infer that even the most savage of men could do so too." (Flowers, Fruits and Leaves, p. 74.) There is, however, no doubt on other grounds, notably the brilliant colours of the males of some monkeys themselves, that monkeys are not colour blind, while the fact that the Malays have fewer names for colours than they can see, and the Pahang Sakais, as Mr. CLIFFORD has shown in the last number of the Journal, have only three names for colours—black, white and red—although they can apparently distinguish others, entirely negatives the colour-blind theories, which are but another example of the worthlessness of ethnological deductions based on the study of ancient literature only, unsupplemented by comparison with the characteristics of modern savages.

List of plants, the seeds of which are certainly dispersed by mammals.

Adinandra dumosa. Fruit white, with small seeds. Eaten by bats.

Durio sibethinus, L. By bears.

Neesia synandra, Mast. Seeds with eatable yellow aril. Mice.

Canarium rufum and other species. Fruits green or dull purple with a turpentiney or sweet taste. Eaten by monkeys. A single large seed protected by an excessively hard shell.

Calophyllum pulcherrimum and other small fruited kinds

Fruits green with a hard seed. Bats and monkeys.

Elæocarpus. Fruit green or blue. Stone hard. Monkeys.

Pyrenaria. Fruit green. Seeds hard. Monkeys, squirrels.

Parinarium Griffithii, Hook. Fruits firm dull purplish. A large hard seed. Monkeys.

Strombosia javanica, Bl. Fruit green. Stone hard. Squirrels.

Mangifera. Fruit green, yellowish or grey, large. Stone hard. Rats.

Nephelium. Mostly dull coloured. Civets, monkeys, bats.

Xerospermum. Fruit yellow. do.

Lansium domesticum, Jack. *Langsat*. Fruit whitish. Seed covered with sweet pulp. Civets.

Dialium. Fruit black. Seeds hard. Monkeys.

Eugenia grandis, Willd. Fruit green. Stone hard. Monkeys, squirrels.

E. scoparia, Wall. Fruit blue. Stone hard. Monkeys, squirrels.

E. malaccensis, L. Pink or white. Stone hard. Monkeys, civets.

Careya arborea. Large green fruits. Monkeys.

Psidium guava, L. Fruit green. Seeds small and hard. Civets.

Barringtonia racemosa. Fruits modified for sea-travelling, but I have seen a squirrel run off with one to some distance. They are dull green with a large stone.

Melothria, *Mukia* and other small terrestrial *Cucurbitaceæ*. Fruits usually dull green. With many hard seeds. Eaten by rats.

Marlea nobilis, C. B. Clarke. Fruits green, hard stone. Squirrel, tupaia.

Citrus aurantium, L. Fruit orange, usually green when ripe in the tropics. Monkeys, civets (Opossum in Brazil).

Coffea. Berries red. Seeds hard enclosed in a thin sweet pulp. Civets, monkeys.

Gardenia Griffithii, Hook. Capsule green, full of flat seeds in a sweet pulp. Monkeys.

Randia anisophylla, Do. do.

Dichopsis obovata, C. B. Clarke. Fruit green passing into orange. Seed hard. Monkeys.

Sideroxylon sundaicum. Fruit green. Seed hard, polished. Squirrels.

Mimusops elengi, L. Fruit orange. Civets (also birds).

Achras sapota, L. Fruit brown. Seed hard, polished. Civets, bats.

Diospyros discolor. Fruit pink, dull and inconspicuous. Scented Civets.

Willughbeia. Fruit green. Monkeys.

Strychnos Tiente. Fruit green. Seeds poisonous, imbedded in a sweet pulp. Monkeys.

Fagraea fragrans. Fruit orange. Usually eaten by birds, but also by bats.

Baccaurea Motleyana, Hook. Fruit white. Monkeys, squirrels.

Laurinea sp. Small inconspicuous, green fruit. Hard seed. Monkeys.

Artocarpus rigida, Bl. Yellow. Fruits large yellow, inconspicuous. Seeds small enclosed in a sweet orange pulp. Civets.

Ficus (Figs). Inconspicuous fruited kinds, e.g., *F. Miquelii* *F. Benjaminia*. Bats, also more rarely monkeys and also birds.

Quercus, Castanopsis. Fruit inconspicuous. Squirrels, more rarely monkeys.

Gnetum scandens, Bl. Fruit orange. Hard seed. Civets.

Amomum. Fruit usually inconspicuous, Squirrels, rats.

Zingiber. Do. do.

Musa. Wild plantain. Fruit green, inconspicuous. Seeds small, hard. Monkeys.

Curculigo. Fruit inconspicuous, hidden. Probably eaten by mice.

Livistona. Fruit black. Seed large hard. Bats.

Caryota. Fruit dark red. Seed hard. Civets.

Zalacca. Fruit brown, acid. Seed large hard. Rats or squirrels.

Calami. Fruit brown or yellow. Seed hard, covered with thin pulp. Apparently eaten by squirrels, very speedily taken, the nibbled skin only left.

Scirpodendron. This aberrant *Sedge* has its inconspicuous fruits always nibbled by some small mammal.

ADHESIVE FRUITS DISTRIBUTED BY MAMMALS.

In the Malay Peninsula there are fewer plants furnished with means of adhesion to fur or feather than in many parts of the world. This is owing to the limited amount of open country, the greater part of this region being covered thickly with a dense jungle of lofty trees. For the greater part of the adhesive fruits belong to herbs, or half-shrubby plants living in flat, open country. Of such as we do find here, a number are aliens more or less accidentally introduced, such are the white *Plumbago* (*Plumbago zeylanica*, L.), *Urena lobata*, the sensitive plant (*Mimosa pudica*, L.), *Triumfetta*, and *Paspalum conjugatum*, L. Nearly all of these are carried about by man or domestic animals.

In *Plumbago zeylanica*, L. the calyx which encloses the capsule is provided with sticky hairs, which readily adhere to clothes or fur. The plant is common in villages, but I never saw it at any distance from cultivation. In *Triumfetta*, a roadside weed, the capsule is provided with hooks.

Paspalum conjugatum is a common grass, the very small spikelets of which are rounded and edged with short, bristly hairs. They are very easily detached from the rachis on which they are arranged, and attach themselves readily especially in wet weather to clothes, &c. This grass has travelled further than any of the introduced weeds throughout the Peninsula. I have found it growing in crevices of rocks in the Tahan River as far as I have been, and on Padang Batu on Mount Ophir, I saw a plant growing at the stream close to the camping ground, at the spot where the natives who visit the spot are accustomed to bathe and wash their clothes. A very long way from the flat country where it is abundant.

In *Chrysopogon aciculatus*, Beauv., commonly known here as love-grass, the spikelets are arranged in an erect panicle with slender, wiry branches each of which bears one fertile spikelet the base and one or more barren spikelets. From the

bottom of the lowest spikelet projects a spur covered except on one side with stiff yellow hairs pointing upwards. When the fruit is ripe this fertile spikelet readily breaks off and adheres by its spur to cloth or the fur of an animal and is borne away. This grass is very abundant in dry open country, and forms an extensive turf in many places.

Besides these grasses, there are three species which inhabit the dense jungles, and excepting bamboos, are the only jungle grasses here. They are *Leptaspis urceolata*, Br., *Lophatherum gracile*, Brngn., and *Centotheca lappacea*, Beauv.

The first of these has a loose spreading panicle bearing curious oval spikelets, of which one of the outer glumes is, in the female flower, swollen up and entirely encloses the fruit, this outer glume is covered thickly with short but strong, abruptly hooked hairs, by which it clings very tightly to a passing animal. So adhesive is it that in brushing past it it often happens that the whole inflorescence is torn off.

In an allied species *L. cochleata*, a native of Ceylon, the spikelet is smaller and kidney-shaped with five ridges and covered in like manner with very short hooked hairs.

In *Lophatherum* the spikelets have several glumes, of which the eight terminal ones bear awns covered thickly with minute processes pointing downwards. When the fruit is ripe the awns become hooked by drying, and by this and the minute processes the spikelets can attach themselves to any animal.

In *Centotheca* not only are the branches of the panicle provided with short processes (pointing upwards in this plant) but from each side of one of the upper glumes which encloses the fruit, project a double row of long, white processes by which means the whole panicle readily adheres to the clothes of man or to the fur of an animal. These two latter grasses are especially common along paths and animal tracks in the thickest jungles, but where it is too thick for animals to go easily, one does not find them.

It is probable that more plants will be found which possess these adhesive fruits in the Malay Peninsula, but these will, I think, be chiefly introduced weeds. In any case the number will be very much smaller than that of plants dispersed in any

other of the ways mentioned above, viz., by being swallowed by animals or birds, or by being drifted by wind or water.

The part played by mammals is insignificant compared with that played by birds in the dispersal of seeds, but as has been shewn it is too important to be overlooked.

Of the great waves of evolution which in past times have altered the whole character of the Flora of the world, the first and greatest was probably that due to the appearance of pollen- and honey-seeking insects through whose agency the brilliant colours and elaborate forms of flowers were evolved. The next was due to the evolution of the frugivorous birds through whom we have obtained much of the colouring and sweetness of the fruits. Through the evolution of mammals, we have also obtained many modifications of fruits, and the development of some groups of plant, notably the grasses and some of the other herbaceous plants will, I believe, be eventually shown to have borne a close relation to the evolution of the graminivorous mammals so abundant at one period of the world's history.

—:O:O:—

ACCOUNT OF A TRIP UP THE PAHANG, TEMBELING, AND TAHAN RIVERS, AND AN ATTEMPT TO REACH GUNONG TAHAN.

— : o : —



OUR party, composed of Messrs. H. N. RIDLEY, Director of the Botanic Gardens, Singapore, W. DAVISON, Superintendent of the Raffles Museum, and Lieut. H. J. KELSALL, R.A., with a staff of native plant-collectors and bird-skinners, left Singapore at 3 p.m. on 23rd June, 1891, in the s.s. *Glanggi*, and arrived at Kuala Pahang about 10 a.m. on the 24th.

Leaving our men to look after the baggage we proceeded to Pekan in a steam-launch belonging to Mr. HOLE, of Pekan, that gentleman having kindly offered to take us with him, as he was returning to Pekan at once.

Arrived there, we proceeded at once to the Residency; Mr. HUGH CLIFFORD, the Acting Resident having made arrangements for our accommodation.

We found that Mr. HOLE had, in accordance with previous arrangement, despatched a boat-load of rice and other stores, with 25 coolies, to await our arrival at Kuala Tembeling. So far all was satisfactory, but we found that there was some difficulty in getting a supply of small boats, which would be necessary above Temerloh (Kuala Semantan), owing to the shallowness of the river—the season being an exceptionally dry one.

We went all round Pekan endeavouring to buy three or four boats suitable to our needs, but in vain, there were none

34 ACCOUNT OF A TRIP UP THE PAHANG, AND OTHER RIVERS.

for sale. In this difficulty Mr. HUGH CLIFFORD came to our assistance, and placed at our disposal a large native built boat. His head boatman, after some little difficulty, succeeded in getting together a gang of 25 coolies, amongst whom were two Dyak lads, one of whom turned out a most useful hand in the jungle, and set them to work to caulk and clean the boats which had not been in use for some time, and consequently required overhauling.

All this delayed us in Pekan till the 28th, but the time was not altogether wasted, as we did some collecting in the immediate neighbourhood.

On the North side of river, which is here over half-a-mile wide, are extensive sandy plains interspersed with patches of heavy jungle. On the sandy portions, the trees are all much stunted and grow in small clumps or coppices, giving quite a distinctive appearance to the landscape.

Portions of these sandy tracts were riddled with the burrows of the bee-eaters (*Merops philippinus*), at the extremity of which they lay their eggs. These burrows are excavated in the hard ground and slope downwards for the first foot or so, then run more or less horizontally. Among other birds noticed were the Brahmany Kite (*Haliastur indus*), which appears to be the common kite of the southern portion of Malaya, the "Burra Burra" (*Trachycomus ochrocephalus*), one of the best songsters amongst Malayan birds; most of the common Kingfishers (the Raja Udang of the Malays), and some rarer ones (*Halcyon coromanda* and *Alcedo meningting*), a few woodpeckers and several swifts. Butterflies were not plentiful, the most noteworthy being perhaps the beautiful *Parthenes gambrisius*.

Of mammals, the only ones noticed were the common mangrove monkey, Kra of the Malays (*Mucacus cynomolgus*), and several of the common squirrels (*Sciurus notatus* and *Sciurus tenuis*). Of reptiles, the large Monitor (*Hydrosaurus salvator*) and the beautiful many-coloured sand lizard.

On the 28th, everything being at last ready and being provided with letters from Mr. CLIFFORD to the Sultan asking him to render us any assistance in his power in the way of

providing guides, we started up-stream about 11 a.m., being obliged to wait for night tide to get out into the main stream, Pekan being situated on a sort of loop. Mr. TOWNSON, a young Australian, accompanied us. By 5 p.m. we had reached Tanjong Bedang, and went ashore for a short time. Here we saw some peafowl, which are very plentiful all along this part of the river, and large numbers of the ground dove "Terkuku" (*Turtur tigrina*), which are exceedingly good eating, also specimens of the Burmese lapwing (*Lobivanellus atronuchalis*), and the small Malayan hornbill (*Authracoceros convexus*). At dusk we started again meaning to travel all night, but owing to our constantly running on sand-banks, with which the river abounds, we made but little progress. Our means of progression was by means of poles or *gallahs*, eight men poling at a time, four at each side.

29th June. We went ashore for an hour or two in the morning to collect but did not get anything of note; a couple of specimens of the Eastern little Tern (*Sterna sinensis*) were shot.

The general character of the country bordering the river up as far as Kuala Tembeling is pretty much the same. Along the river bank, is a strip of jungle, 100 to 200 yards wide, with frequent small villages and plantations of coco-nut, banana, and mangosteen trees. Inside of this lies a tract of open grass or swampy land, varying from 100 yards to half-a-mile in width, and beyond this comes the jungle proper. The tract of open land is in places ploughed and cultivated by the natives, who use rude wooden ploughs, which, however, do the work required of them fairly well. The draught animal is the water-buffalo (*Bos bubalus*), of which there are large herds in a semi-domesticated condition. We continued travelling by night, and made better progress than before.

30th June. We heard a Wau Wau in the early morning, the first we had noticed. We made good progress during the day, and got as far as Kuala Luit, about 50 miles from Pekan. Here we went ashore for a short time, and got specimens of *Tupaia javanica*; *Cymborrhynchus macrorrhynchus* the Rouge-et-noir Broadbill, and the Malayan Falconet

(*Microhierax fringillarius*), in addition to which were noted an osprey (*Pandion haliaëtus*), a night jar (*Lyncornis Temminckii*) and a jay (*Platysmurus leucopterus*). We started again at dark, and travelled during the night.

1st July. We continued steadily up-stream all day, and towards evening went ashore to collect, but got very little, we however shot a couple of hornbills (*Anthracocrus convexus*), which were very plentiful, and which are excellent eating. They are best stewed and have rich and pleasant though rather strong flavour. We always looked upon them as a great addition to our bill of fare. We travelled all night as usual, and next day, 2nd July, reached Kuala Berar about noon. Just below Kuala Berar the river was divided into two by a long sandbank, and unfortunately we took the wrong channel, and when we got to the upper end we found that the water was too shallow to allow of our boat, which did not draw more than about two feet of water, passing, so we were obliged to return down stream and follow up the other bank of the river, this time with success. This mistake delayed us more than an hour. At Kuala Berar we stopped several hours, as we wished, if possible, to engage a few more men, but in this we were unsuccessful. Here we noticed in addition to birds already mentioned, the small red Kingfisher (*Ceyx rufidorsa*), the green Broadbill (*Calyptomena viridis*), the little brown Barbet (*Calorhamphus Hayi*) and the not common *Anthothreptes hypogrammica*. At 3 p.m. we started again, and at 5 p.m. had another halt at Kampong Pamum to bathe and let the men have their evening meal. At 6 p.m. we again started and got along well for some time, the river being much narrower and consequently less encumbered with sandbanks. The little tern was still plentiful, flying up and down the river. About 7 o'clock a very heavy thunderstorm came on and it became so dark that we were obliged to tie up to the bank, as we could not see to proceed. The lightning was magnificent, like rivers of liquid fire running right across the sky. There was very little thunder and scarcely any rain. It was some hours before we could proceed and we did not pass Kuala Triang till dawn.

3rd July. As we knew that we should be unable to get much beyond Temerloh in our large boat, we began to look out for smaller ones and landing at Pulau Guai, where there is a straggling village, we found, after some search, one sampan which the owners, after some bargaining, let us have for \$11. We shot a few birds here but nothing unusual. Taking our newly acquired boat in tow, we proceeded up-stream still looking out for boats, and presently we passed another which looked suitable but it was some time before we could find the owner who was at Mosque, it being Friday, but eventually, after some hours' delay, we agreed to give \$15 for the boat and went away with it in tow. A nest of the Racket-tailed Drongo was noticed in a tree near the bank of the river. An unsuccessful attempt, owing to its being at the end of a thin branch, was made to get it. The nest is a very frail structure being a cup of open basket-work of grass stems about as large as an ordinary large breakfast cup suspended below the branch. It contained young birds and the parents resented the attempted robbery most pluckily, flying round the head of the intruder and uttering shrill screams. We halted for the night at Pulau Jelam where we saw some teal (*Dendrocygna javanica*). The night was so dark that we could not travel. About 10 a.m. on the 4th we reached Temerloh where, finding that it was impossible to get the big boat any further, we stayed the whole of that day and the next trying to get boats which, owing to the demand for them by the miners in the Ulu, were difficult to obtain. At last, however, and after a great deal of bargaining we got three of a suitable size at a fairly reasonable price.

The night before our arrival a buffalo calf had been killed by a tiger about three miles from Temerloh, and the following night Mr. OWEN, the District Officer, accompanied by one of our party, sat up for some hours over the hill in the hopes that the tiger would return, but without success. The pretty striped squirrel called Tupai B'lang (*Sciurus Rafflesii*) was here very plentiful. On the 6th July, having transferred all our baggage to the small boats, seven in number, including the two small sampans we had brought from Pekan, we got under

way about 10.30 a.m. and went on steadily till 6 p.m. when we halted for the night at Tanjong Doyang, camping on a sandbank with no shelter but our mosquito curtains which we found sufficient to keep out the heavy dew. A specimen of *Sterna javanica* was seen here. Next morning (7th July) we spent an hour or two collecting before starting and got the following birds amongst others, *Dryocotaphus nigricapitatus*; a brown Babbler (*Timalea nigricollis*); one of the beautiful yellow crested Woodpeckers (*Gecinus chlorolophus*); and the Malayan Falconet (*Microhierax fringillarum*). Starting at 9.30 a.m. we went on steadily till 6 p.m. when we halted and camped on a large sandbank at Pulau Changai. Here we found great tracks of elephant and crocodile on the sand and heard the peculiar cry of the Argus Pheasant, *Burong Kuang* of the Malays.

On the morning of the 8th, we went out collecting, but saw nothing of note but the large Horn-bill (*Buceros rhinoceros*), of which, however, we could not obtain specimens as they kept in the tops of the tallest trees.

During the day, the Semantan Hills were conspicuous to the West and also some limestone hills to the East and in some places the river became much narrower and deeper, flowing between rocky banks.

At 4.30 p.m., we halted at Tanjong Antan where there was a good sandbank. Here we decided to camp as it was doubtful if we would find another sandbank, it being very difficult to camp elsewhere especially where the river banks are steep. From our camp we had a fine view of the Semantan Hills over which the sun set with a fine after-glow of purple and gold.

Next morning we found tracks of elephant and obtained a few birds including a specimen of *Halietus plumbeus*, the rare lesser sea-eagle. Several Wauwaus were also heard. About 4.30 p.m. we reached Pulau Tawar, the place of residence of the Sultan of Pahang, and camped on a sandbank, a short distance above the village. The name is rather misleading as there is no island, the village being built along the left (East) bank of the river. The banks are here about 30 feet high and steep.

The Sultan informed us that very little was known of Gunong Tahan and was not sure if there was any one who could act as a guide. However he gave us a letter to Panglima Garang YUSUF at Kuala Tembeling in whose district he thought there was a man who had once been to Gunong Tahan, but of this he was not sure. He questioned us as to our objects in going into the jungle and took much interest in an 8-bore gun we had with us, being himself a bit of a sportsman and possessing several guns.

Next day (11th July), we got to within a few miles of Kuala Tembeling which we reached early on the 12th. There we found our 25 coolies and our stores safely deposited in the Police Station.

Being told here that Kuala Tahan was only one day's journey from this place, it was arranged that Mr. KELSALL and Mr. TOWNSON should go ahead with 16 men in three boats and as much of the stores as they could carry in addition, as far as Kuala Tahan, where they would form a camp. They started at 2 p.m. and at 4.30 p.m. reached the first rapids where all hands had to be turned on to each boat in turn to drag them up. At 6 p.m., they reached Kuala Kuang where they camped for the night.

The next day they got as far as Kampong Pulau Manis and on the 14th reached Kuala Tahan, after passing several bad rapids. The largest boat was stopped about $1\frac{1}{2}$ miles below Kuala Tahan by the river becoming very shallow. The other two, smaller ones, had therefore to make several trips to and fro, and by 7.30 p.m. all the baggage was safely deposited on a shingle bank about half-a-mile up the Tahan.

The Tahan is deep and still at its mouth, and is much impeded with fallen trees, and is dark and dismal looking. It, however, gets shallow very soon and opens out somewhat.

On the 16th, Mr. KELSALL returned with the three boats to Kuala Tembeling passing on the way Mr. RIDLEY who, having obtained some more small boats, and found Panglima Garang YUSUF had started on the 14th.

On the 16th, Messrs. DAVISON and KELSALL started with the remainder of the baggage, on the way they dynamited

several pools in the hopes of getting some fresh fish. Nothing was got out of any of them but one, but that made up for all the rest as 35 large fish were killed in it. On the 17th, at 3 p.m., they reached Kuala Tahan. A very curious fish was here obtained, about 3 feet long with a curious humped back and two rows of teeth on its tongue. Mr. RIDLEY, accompanied by Panglima Garang YUSUF, visited the Penghulu Raja near Kuala Tahan in the hopes of obtaining a guide or at least reliable information as to the route to Gunong Tahan. He, however, was either unwilling or unable to supply guides nor could any reliable information be obtained as to the district. Indeed it appeared doubtful whether any Malays had ever been nearer to the mountain than was sufficient to see it. A number go from time to time up the Tahan valley to collect guttah and rotans, but as they go in boats the distance that they travelled is very short and probably such information as they are possessed of with reference to the Gunong Tahan is derived from the Sakais.

Gunong Tahan, we were told, could be approached by two routes, one following the river Tahan, the other the river Ketchau. By the route the mountain can probably be approached nearer but it was said that on the W. or S.W. side, from which the Ketchau takes its rise, it is exceedingly precipitous and unpracticable, while by the Tahan route it is possible to ascend. The natives further state that there is a lake or swamp from which flow the rivers Tahan and Kelantan.

On the 19th, Mr. RIDLEY, accompanied by Mr. TOWNSON, started off with four boats and fourteen men up-stream and two men were sent back to Pekan in one of the boats with an order for more rice as we found we would require more, and owing to the previous bad season, rice was hardly procurable and very expensive in Pahang. Panglima Garang YUSUF tried to get us a guide but without success. He produced an old man who was supposed to know the way to Gunong Tahan and offered to guide us there for the modest sum of \$60. However as we discovered after some conversation with him that he had never been up to the mountain but had only seen

it from the river we dispensed with his services. Panglima Garang YUSUF also left us at this point.

At this camp we heard for the first time a most peculiar noise which we at first attributed to horn-bills but subsequently found to be produced by a large monkey, of what species we were unable to determine, as although many attempts were made to obtain specimens, we were entirely unsuccessful owing to the extreme wariness of the monkeys. The cry commenced with a series of deep sonorous barks which gradually becoming quicker and shriller ended in a sort of unearthly laugh and then commenced again. We frequently heard these monkeys again up the Tahan valley. Here we obtained our first specimens of the pretty little chestnut-backed Forktail (*Hydrocichla ruficapillus*), the bird which frequents all the small rocky mountain streams and of which a nest was subsequently obtained. The nest which is cup-shaped is constructed of moss and lined with dead leaves and built against the almost perpendicular face of a large rock overhanging the water. For three nights in succession a beautiful specimen of that glorious butterfly (*Zeuxidia aureola*) crossed the river just below our camp at dusk, but unfortunately we did not succeed the capturing it.

On the evening of the 21st, three of the boats which Mr. RIDLEY had taken up-stream returned and we found that he had only been able to reach a point about three miles up-stream at the mouth of the Sungei Tenok beyond which point the boats could not proceed owing to the bed of the stream being choked up with boulders. In fact it was with great difficulty that Mr. RIDLEY had succeeded in getting the boats up thus far as the stream was exceedingly shallow and much encumbered with fallen trees, and owing to having to stop to make a channel for the boats, moving stones to either side and cutting through fallen logs, it took the better part of two days to accomplish this distance.

On the 22nd the three boats were again started up-stream with a second load of rice and other stores and fifteen men. There was heavy rain during the evening and the river rose some six or eight feet during the night, nearly washing away

some of our stores which had been left on the shingle bank instead of being moved up to our store-house on the bank, we woke, however, just in time to save them.

On the 23rd Mr. KELSALL with three men to cut a path ascended a hill about six or seven hundred feet high about two miles N.E. of Kuala Tahan and from the top of a tree got a view of a high ranged mountain about 20 to 30 miles distant to the N. W. This was doubtless the Tahan range. It consisted of a long range running apparently N.E. and S.W. and culminating in several peaks, the highest of which was towards the N.E. end of the range and appeared to be from 8,000 to 10,000 feet high.

Mr. RIDLEY also ascended some of the hills in the neighbourhood of Kuala Tenok in the hopes of getting a view and saw a high range to the S.E. which he could not identify but owing to the density of the jungle he could not see any distance in other direction.

On the 24th, Messrs. DAVISON and KELSALL with the remaining stores proceeded to Kuala Tenok, the boats having returned the previous evening. Two men were left at Kuala Tahan in charge of stores we left behind and six coolies for whom there was no room in the boats. Owing to Mr. RIDLEY having cleared a passage for the boats and to the greater depth of water due to the heavy rain that had fallen they were able to accomplish in $4\frac{1}{2}$ hours what had taken Mr. RIDLEY almost two days in spite of the fact that the boats had to be dragged our shoals almost half the distance traversed.

Finding that it was impossible to get the boats any higher up the river, Mr. RIDLEY had commenced cutting a path along the right bank.

The jungle is very thick for the most part and contains a considerable number of valuable timber trees such as *Tampanis*, *Kayu Minyak* and other dipterocarpaceous trees but the river is too shallow and full of rapids to allow of their being floated down in the usual way. There is also still a considerable quantity of getah percha (*Dichopsis gutta*), getah grip (*Willughbeia edulis*), besides jelutong and other gutta-producing trees of less value. Rotans of various species are also very abundant and there is much dammar.

On the 25th July, Mr. RIDLEY started off early with a few men to clear the path ahead. The rest of the men were employed most of the day making "*ambongs*" or back baskets in which to carry loads.

On the 26th, we all started early with as much baggage as the men could carry and marched about 2½ miles along the path cleared by Mr. RIDLEY. This path was in parts very bad, being steep, narrow and along the face of a hill which made the progress of the coolies with the baggage very slow, and it was well on in the afternoon before the distance was accomplished.

The river here is very beautiful flowing in a narrow valley between steep jungle-covered hills rising to a height of 800 to 1,000 feet. The stream itself is full of large boulders.

Camp was formed in a small, clear space in a small ravine near the river which was the only few yards of level ground we could find. The six men we had left at Kwala Tahan to follow overtook us at this camp (Camp No. 3) soon after dark. About 7.30 p.m. it commenced to rain heavily and continued for three or four hours. We had with us three large sheets of strong linen to serve as tents and a small tent of Mr. TOWNSON. Two of these we used, the native servants had one and one was used to cover up the stores at night. These tents we found were anything but waterproof and being open at the ends the rain used to blow in and we all frequently got very wet. The coolies made *pondoks* of palm leaves for themselves each night.

On the 27th, Messrs. RIDLEY, KELSALL and TOWNSON went forward with a few men cutting a path, while Mr. DAVISON spent the day exploring the neighbourhood of the camps for birds and insects but without much success. The bulk of the men were sent back to Kuala Tenok in charge of a Mandor to bring up more stores. During the day about two miles of path were cut along the river bank and Mr. RIDLEY and his party returned to camp at 5 p.m.

The comparative absence of bird life in these jungles is most remarkable. Sometimes one would spend the whole day in search of specimens and only obtain half a dozen or so.

44 ACCOUNT OF A TRIP UP THE PAHANG, AND OTHER RIVERS.

This is partly owing to the denseness of the foliage which renders it very difficult to see any birds that are about, and also to the fact that except quite early in the morning and for a short time in the late afternoon most of the birds are quite silent.

On the 28th July, Mr. RIDLEY accompanied by Mr. TOWNSON with a small party of men went on cutting the path, while Messrs. DAVISON and KELSALL superintended the transport of the stores from the third camp to the fourth about a mile further on. This necessitated two journeys and occupied nearly the whole day, as the path was very bad. Mr. KELSALL saw a specimen of the rare striped civet cat (*Hemigale Hardwickii*) but not having a gun could not add it to our collections.

Our rice was made up in 50-lb. bags which is the very outside a man can carry in the jungle. The direction followed was North-westerly along the bank of the river and the path was a constant series of ascents and descents over the spurs of the hills between which the river flows, and was most difficult for men carrying loads. This was especially the case when small streams running into the Tahan had to be crossed. The banks of these streams are usually steep and after rain become muddy and exceedingly slippery and in many places it was only with the assistance of branches and roots that the men could get up and down with the loads. In spite of every effort being made any thing like rapid progress was impossible and it was only by the exercise of a very large amount of patience and tact that the men were got to do the amount of work they did. On one or two occasions some of the men threatened to run away and one mandor gave a great deal of trouble, not only would he not do any work himself but he did all in his power to make the other men discontented.

On the 29th, Mr. RIDLEY and Mr. TOWNSON went on as before with three or four men cutting a path while Mr. DAVIDSON and Mr. KELSALL looked after the remainder of the men, some of whom were sent back to Kuala Tenok to bring up more rice, and the rest were sent on after Mr.

RIDLEY with stores for several days so that he might form an advance camp and thus be saved the journey back to the main camp every day. A specimen of the handsome squirrel (*Sciurus insignis*) was obtained at this camp.

On the 30th of July, Mr. RIDLEY'S best plant collector who had been complaining of fever for some days was so bad that he had to be sent back to the camp at Kuala Tenok. He was so weak that he required the assistance of two men to enable him to walk. Mr. KELSALL also went back to Kuala Tenok to get some things that had been left locked up and to see this man safely back. Sakai and wild beast tracks occur on both sides of the river and these were often made use of, being enlarged for the baggage carriers. In some places trees had to be cut down to form bridges over the river and the ravines that had to be crossed in the jungle. During the whole time we were in the Tahan jungles we did not meet a single Sakai although we constantly met with evidence of their presence in the shape of rough leantos or "*pondoks*" consisting of a long pole supported at each end on a forked stick stuck in the ground against which palm leaves are leant. Other evidences were small platforms in the forks of trees, 15 or 20 feet from the ground, dead fires and on one or two occasions newly cut branches of trees. At one place Mr. RIDLEY found on the banks of the river a group of seven small huts or booths made of palm leaves tied together at the top in the shape of a bee hive and with a thick bed of palm leaves in each of them.

The wild men are evidently very shy, as they never let us catch a glimpse of them although they evidently watched us all along, and on our moving from one camp were quick to clear off all old tins and other waste which had been thrown away.

On the 31st, Mr. RIDLEY continued cutting the path and Mr. KELSALL followed him to make a rough compass sketch of the route, overtaking him not far from the 6th camp, while Mr. DAVISON superintended the coolies moving the stores on to the 5th camp. This part of the path was very bad. Mr. RIDLEY and his party during this day ascended a hill of considerable altitude on the left bank of the river and

got a good view of the Tahan range—a broken ridge densely covered with trees. On one side could be seen a vertical cliff white in colour and possibly of limestone more or less clothed with vegetation and to the North rose the big peak of Gunong Tahan. It did not appear to be of the estimated height of 14,000 feet, and is probably not more than 8,000 to 10,000 feet in altitude.

On the 1st of August, Mr. RIDLEY sent back all his men to assist in bringing up stores, and the whole were brought up to the 6th camp. The next morning twelve men in charge of a mandor were sent back to Kuala Tenok to bring up all available rice and fish, the rest of the men moved everything on to the 7th camp. The road between the 6th and 7th camps was very bad being mostly along the side of a steep hill. Mr. RIDLEY had also been obliged to cut down a tree of considerable size to form a bridge over the river.

On the 3rd, Mr. RIDLEY and Mr. TOWNSON with some men went on cutting the path, and formed a fresh camp about a mile and-a-half ahead, while Mr. DAVISON and Mr. KELSALL did some collecting.

On the 4th, Mr. RIDLEY and party went on about one and-a-half or two miles and formed the 9th and what proved to be the last camp. Mr. KELSALL and Mr. DAVISON went on collecting, Mr. KELSALL going as far as a little beyond the 9th camp. The twelve men who had been sent back for rice returned this day.

The 5th and 6th were spent by Mr. RIDLEY'S party trying various routes as it appeared impossible to continue along the banks of the Tahan beyond the 9th camp owing to the steepness of the hills between which the river here flows and at last they took a well-worn Sakai track which went away in a South-westerly direction and then turned North again. This track followed roughly the direction of a small stream which falls into the Tahan from the West and which appeared to come from the slopes of Gunong Tahan. After following this track for some distance they left it and ascended a high ridge which barred further progress up the valley of the Tahan, at its lowest point.

While Mr. RIDLEY and Mr. TOWNSON were on this hill, a message came from Mr. DAVISON'S camp saying that Mr. KELSALL was ill, having been attacked with fever on the night of the 4th and would probably be unable to move for some days. Mr. RIDLEY and Mr. TOWNSON returned to the 7th camp on the 7th to obtain more provisions and consult as to the next step. On looking over the stores it was found that there was only enough food for five days, and as the supply that had been sent for to Pekan had not arrived at Kuala Tenok and in all probability would not do so for five or six days a retreat was decided upon.

The expedition started back on the 9th (Mr. KELSALL being carried on a raft or stretcher for two days) and reached Kuala Tenok on the 11th; the 12th was spent re-packing stores and waiting for the last of the baggage to arrive. The whole of the 13th was occupied moving down to Kuala Tahan, the river was a good deal swollen owing to the recent heavy rains, and one small boat was upset, fortunately there was nothing in it of much value. One day had to be spent at Kuala Tahan repairing the boats and constructing rafts of bamboo to convey some of the men and the collections of live plants down stream.

Pulau Tawar was reached on the 19th of August and a stay of one day was made in order to visit the limestone caves at Kota Glanggi. Some good things were obtained in the neighbourhood of the caves including a specimen of BOSCH'S beautiful ground thrush (*Pitta Boschi*) and in one of the caves, Kota Burong, a pair of long-tailed porcupines (*Hystrix longicauda*) were taken. Here also was found a nest of the forest Bee-eater (*Nyctiornis amicta*). It consists, like those of the other Bee-eaters, of a hole in the ground some two or three feet long. The eggs are pure white. Leeches were also more plentiful in these woods than we found them anywhere else.

These caves appear to be frequented by the wild men, for in all of them we found remains of fires and beds of palm leaves but of the people themselves we saw nothing.

The failure of the expedition to reach its goal was due to a

variety of causes of which the chief were the difficulties of transport owing to the low state of the river and bad coolies. Food for the expedition for two months—the time allotted by the Government—was taken but owing to the want of rain the rivers were so shallow that it was with the greatest difficulty that the boats were taken up and this delayed progress greatly. Added to this were the difficulties of making our way through the Tahan jungles which were far greater than had been anticipated. The coolies who were chiefly Kelantan men proved, with a few exceptions, a very worthless lot and many of them suffered from beri-beri, fever, diarrhœa, dysentery and a most loathsome form of skin disease, known among the Malays as "*kurap*."

Judging from recent information about this part of Pahang, the route adopted was probably not the best, as it gave us much more actual jungle work than would have been necessary had the Tembeling River been followed up to its source and a path cut from thence to Gunong Tahan.

One man can carry in the jungle but little more provisions than suffice for fifteen men for one day, so that for ten days in the jungle away from the base of supplies ten men out of fifteen are required at the start to carry provisions and every additional day away from the base increases the difficulties of transport.

The Tahan jungles appear to be very unhealthy especially at the further points reached. Heavy rain fell here every night, and the ground was covered with decaying vegetable remains. At night frequently the whole ground round the tents was illuminated by phosphorescence, (probably bacterial) on the decaying leaves.

Unfortunately though the jungles contain plenty of game, it is not easily procurable and cannot be caught without trapping, and this, time did not permit. All the streams, however, contain plenty of fish and they can generally be obtained with the aid of dynamite.

It was hoped that some observations might have been taken that would materially increase the topographical knowledge of the Tahan region, but owing to the close nature of the

country and the impossibility of getting any view, only the roughest observations could be made.

Although two out of three plant collectors were ill—one with fever, the other from an injury to his foot—for the greater part of the time in the Tahan River valley, and were, therefore, useless, the collection of herbarium specimens was very successful, over two thousand being obtained.

H. J. K.

VEGETATION OF PAHANG.

During the expedition no opportunity was lost of collecting plants, and these with the collections made in earlier visits to Pahang (in 1889 and 1890) give a very fair idea of the flora of the low country here, which was hitherto unknown.

The notes and descriptions of these plants being somewhat voluminous are published elsewhere, but a short account of the flora of Pahang, as far as we have seen it, will not, I think, be out of place.

The sea-shore at the mouth of the Pahang River and the adjoining heaths country was explored in 1890. This district, so different from any other part of the Malay Peninsula as far as we have seen, produced a number of interesting and new plants. Along the sea-shore is a single row of *Casuarinas* on whose branches grew besides lichens and fungi a curious mistletoe (*Loranthus*) with leaves and flowers of a bright orange colour. Beneath the trees the ground is covered with pink and white *Ipomeas*, the Porcupine Grass (*Spinifex*), a very small species of *Premna*, and a very pretty violet flowered *Vitex* apparently a prostrate maritime form of *V. Lagundi* which in land forms a small tree. Here and there along the coasts are patches of mangrove, but owing to the sandy nature of this district they are by no means extensive. From the shore inland and along the river on the left bank to some distance above Pekan extends the sandy heath district interspersed with woods, a very interesting region. Here are bushes of *Vacciniums* of two species, with rosy or white sweet-scented flowers and small eatable berries. *Eugenias*, *Rhodomyrtus*, the Sea Olive, (*Olea maritima*) *Ilex*, and many other shrubs, upon whose branches

grow many small orchids, among which were *Eria acervata*, a new *Bulbophyllum*, *Dendrobiums*, and the beautiful climbing white spider orchid (*Renanthera alba* Ridl.). The ground is covered with grasses and sedges, and many small herbs, *Mitrasacme*, with small buff and white flowers; *Utricularias*, pink, yellow and white; the creeping blue flowered *Cyanotis*; and many others. Among the larger trees here are the Tembusu (*Fagraea fragrans*), many figs and *Eugénias* and a magnificent specimen of the big flowered *Fagraea imperialis* was found at one spot in full bloom.

On the right bank of the river, the country is more swampy and wooded, with pools full of the pink Lotus (*Nelumbium speciosum*), and other aquatics. Here are great clumps of *Licuala* palms (commonly called Penang Lawyers). Two new species of grass (*Rottboellia geminata* Hack, and *Saccharum Ridleyi* Huck) were obtained here. The latter forms dense thickets somewhat resembling clumps of Pampas grass with erect panicles of purple flowers.

Up to Kuala Tembeling, the country on the banks of the river is open and flat. For the most part covered with low woods and grassy pastures. Further inland are bigger and denser forests, while the edge of the river is often covered with dense thickets of *Cucurbitaceæ*, *Bauhinias* with brilliant orange flowers, clumps of *Clinogyne*, with white blossoms, big *Zingiberaceæ* and many other beautiful plants. Some of the trees along this part of the river are very striking, *Cassia siamea* and *Cæsalpinia sappan* are bright with yellow flowers, *Lagerstræmias* of two species, *Cassia nodosa* with innumerable rosy blossoms, and *Millettia atropurpurea*, with its deep purple blossom give a brilliancy of colour to the banks. *Bignonia indica*, with its huge sword-like pods is very abundant especially on some of the sandy islets in the river.

The woods in the neighbourhood of Pulau Tawar proved, botanically, exceedingly rich, and many novelties were collected here, including a remarkable *Scitamineous* plant allied to *Lowia*, with five stamens, some very curious *Rubiaceous* herbs, and here the lovely *Didymocarpus quinquevulnerus*, a new species with large white flowers tipped beautifully with

carmine was abundant. The limestone region of Kota Glanggi, was also a field of great interest. The rocks and adjoining woods abounded in remarkable and curious plants. *Trichopus zeylanicus*, a small herbaceous plant allied to the yams was abundant. This plant has not hitherto been collected in the Malay Peninsula, being only known from Ceylon and Southern India; *Begonias*, *Elatostemmas* and ferns, clothed the rocks, and on the higher parts were many orchids, including several new species of *Sarcochilus* and *Saccolabium*. The curious *Arisæma fimbriatum*, and several species of *Amorphophallus*, *Peperomia portulacoides* (a dwarf succulent plant not hitherto known except in Southern India), a very fine violet flowered *Calanthe* and many other plants of interest were collected here. At Kuala Tembeling a good lot of rare and curious plants were met with, both on the river banks and in the woods a little way inland, of which the most interesting were the yellow dead nettle, *Gomphostemma*, and the parasitic *Brugmansia*, one of the *Rafflesiaceæ* (the first of this order recorded from the Malay Peninsula, although Mr. WRAY tells me he has long known of the occurrence of the *Rafflesia* itself in Perak). The *Brugmansia*, which is a native also of Borneo, was found growing on the prostrate stem of a vine, in a dense thicket of tall *Scitamineæ* on the borders of a wood, about two miles from the river.

In the more open woods here and elsewhere in this part of Pahang, grows a very beautiful yellow flowered *Dillenia* well worthy of cultivation. Another interesting tree which occurred here was the "Kapayung" or "Payung," (*Pangium edule*). The fruit of this tree produces a rather coarse oil used by the natives in medicine and also for attracting fish. An old Malay fisherman, whom we met here, had a bamboo full of the pounded seeds, in the form of a dark brown oily mess. He put a little into the water of the river near a deep hole, where, after waiting a few minutes, we fired a charge of dynamite and took a fairly large number of fish.

Along the Tembeling River, the forests came down in many places closer to the water's edge, but the collections made here were more scanty as the expedition was hurrying on.

At one spot, among other plants, a new wild plantain was collected, with the flower bracts of a bright yellow instead of either violet or brown as in the two common Peninsular species.

The entrance to the Tahan River is marked by dense jungle coming down to the water's edge, and the flora here is very different to that of the Tembeling and Pahang Rivers, though some characteristic plants of this region had found their way down the former river for some way. The greater number of novelties found during the expedition were collected in the Tahan valley. Among the more interesting of which were a very fine new species of *Dipterocarpus* (*D. pulcherrimus*) a big tree with hard red timber. At the time of our visit it was covered with its pink two-winged fruits. The flowers, which like those of most *Dipterocarpi* are large and creamy pink with a delicious fragrancly were also collected. The well-known "Champedak," (*Artocarpus polyphemia*) is truly wild in these woods. "Pulawan," (*Tristania Whitiana*), with corymbs of evil-smelling white flowers was another abundant tree. Oaks and nutmegs were comparatively rare. Of palms, "Ebol" (*Orania macrocladus* Bl.), "Langkap" (*Arenga obtusiloba* Miq.), *Caryota mitis* Lour, and "Bayas" (*Oncosperma horrida*) were common, and the grand "Daun Payoh" (*Teysmannia altifrons* Miq.) was met with in several places. The paddle-shaped blades of the leaf of this palm were over six feet long, and form excellent roofs for huts. Of the smaller palms there were a number of Pinangas of all sizes, Licualas, and Iguanuras, two or three Zalaccas, and a number of kinds of rattans. Along the banks of the river many of the herbs and shrubs growing nearest to the water were noticed to have peculiarly long and narrow leaves, among these, were a *Calophyllum* (new), an *Ixora*, an *Antidesma*, a small fig, (*Ficus pyrifolia*), a little purple flowered *Hygrophila*, and a curious new genus of *Asclepiads*, a *Podochilus* and some aroids. These all grew on the rocks close to the water's edge, and must have been subject to constant rushes of water when the river, as it often does, suddenly rises. In such a case as this, the broader leaved plants would suffer heavily, as the rush of the stream would tear their

foliage to pieces, while narrow-leaved plants offering a less resistance would be comparatively uninjured.

Among the herbaceous plants of this region the *Didymocarpus* are most attractive, growing on banks and open spaces in the woods; a number of different kinds were collected including the *D. quinquevulnerus* mentioned above, but the form here had the carmine spots replaced by an exquisite purple violet colour; there was also a very similar species of a dark crimson red with a yellow spot in the tube, and some white and yellow kinds. There were also a number of the smaller flowered species, violet, pink and white. Nearly all the plants of this charming group were new to science. It is remarkable how very local the species of this genus are. None of the numerous species of the Perak Hills occurred here, nor were those of the Malacca, Penang or Johor Hills seen.

Another curious little plant first found here was *Neckia*, one of the *Violaceæ*, but in no way recalling a violet. It is a little erect herb, the stem covered with brown bristles, the leaves narrow, and the flowers very fugacious, pink, on long, slender stalks. It has since been collected on Batu Pahat and Gunong Panti in Johor, but hitherto was only known from Borneo.

Among the aroids of this region, the most interesting is *Rhynchosyris*, a singular little plant abundant on rocks in the streams and watercourses. It was not previously known outside Borneo, but one and perhaps more species are common in similar places in Perak and Johor.

The Ginger family, (*Scitamineæ*) is very well represented, and several novelties were collected. At one place were tall thickets of a curious new *Phrynium* with white flowers. A pretty little turmeric (*Curcuma*) was abundant, *Elettariopsis*, with its long runners almost hidden in wet rotten sticks and throwing up solitary white flowers and globular white fruits, grew in the damp watercourses. *Zingibers*, of which the finest was *Z. spectabile* with its tall scarlet cones from which protrude curious black-veined, yellow flowers; *Amomums* of several kinds, and a *Globba* with yellow flowers and yellow or scarlet bracts, were all plentiful.

A curious plant which requires investigation is *an epiphytic Pandanus* which grows in the form of a tuft often in tall trees. Plants were seen far up the Tahan River but they bore no flowers nor fruit. Similar plants have been seen in Perak and in Johor where it is called "Pandau Lari." It is evidently a true *Pandanus* and not a *Freycinetia*. Orchids are abundant on the trees overhanging the river. Most conspicuous is *Grammatophyllum*, in grand flower at the time of our visit. There were also an abundance of *Dendrobiums*, *Erias*, *Bulbophylla*, *Thecostele*, *Podochili*, *Appendiculas*, *Cælogynes*, including the lovely *C. Forstermanni*. Among the prettiest novelties were a beautiful orange *Dendrobium*, and a *Phalænopsis* in the way of *Ph. sumatrana* but with smaller flowers. The gravelly islets of the river were bright with *Arundina speciosa*, the best form with the very dark lip.

Ferns produced no novelties, but among the most conspicuous were *Dipteris Horsfieldii*, and *D. Lobbii*, *Adiantum æthiopicum*, many *Lindsayas* and *Trichomaues* and *Hymenophylla*.

The common *Selaginellas* and *Lycopodiums* were plentiful, and a number collected. A few mosses were also obtained, among which Mr. BROTHERUS (of Helsingfors) has found two novelties.

Jungle Produce.

The Pahang gutta still holds its own as one of the best class guttas in the market. The tree (*Dichopsis gutta*) is abundant in many of the upper forests and was fairly plentiful in the Tahan valley though here as elsewhere the trees are scattered often at some distance apart in the jungle. *Willughbeias*, (Getah Grip) are abundant, and we had often to cut through big lianes of them in the Tahan valley. A new species of this genus with eatable apricot-coloured fruit grew in the open heath country near Pekan. Jelutong, (*Dyera costulata*, Hook.) is also plentiful.

Gaharu Wood (*Aquilaria malaccensis*) is procured from the Tahan valley as elsewhere in Pahang and specimens of the tree in fruit were collected.

Many *Dipterocarpaceae* trees, such as *Dipterocarpus pterygocalyx* which produces the wood-oil, (Minyak Krueng), and the *Shorea*s and *Hopeas* which produce Dammars were seen throughout the jungles.

Rattans are abundant, but near the villages we noticed comparatively few of any value. The commonest near the Pahang River is Rotan Chin-Chin, the *Dæmonorops crinitus* which appears to be little valued by the Malays. In the wilder parts, such as the Taban valley, we met with many other and more valuable species, which had escaped the search of the natives.

All these products are collected by the Sakais and sold to the Malays who float them down the river on rafts of bamboo to Pekan where the bamboos themselves fetch a certain price as they are scarce in the lower part of the river.

Cultivation.

The Pahang Malay does but little in cultivation and the Chinese have not yet established themselves here as planters, although the soil in many places is very suitable. In Pekan a small quantity of copra is made, but the natives seem to prefer to make coco-nut sugar rather than to attempt to grow the coco-nuts for the fruit. Paddy is cultivated to a small extent, and very carelessly, and here and there are patches of Indian corn and Italian millett (*Panicum italicum*). Some plants of Arabian coffee very strong and free from disease were seen at Temerloh, in the garden of a Malay, who, however, was quite ignorant of the use of the berries, only using the leaves to make tea of.

The ordinary Malay vegetables and fruits were to be seen in the village gardens, but as the Pahang Malay does not care to grow more than he actually requires at the time, they were neither abundant nor very good.

Fruit trees such as Durians, Rambutans, and Pulassan were often found in dense jungle, but I doubt that any were truly wild. They seemed to be either relics of deserted villages or derived from seeds dropped in the jungles by wandering Malays and Sakais.

Of truly wild fruits, the following were seen :—

Champedak (*Artocarpus polyphemia*) ; Rambutan Pachat (*Xerospermum Noronhianum*) ; and a remarkable climber with large fruits, resembling apples in form and flavour, called Akar Panti, in the Tahan woods ; Monkey-jack (*Artocarpus rigidus*), and Tampoi (*Baccaurea malayana*) in the Pulau Tawar woods, where also were trees of apparently a species of *Canarium* called Drija, of which the kernel produced an oil, apparently much sought after by the natives. And here were also fruiting trees of the Kumbang Samangko (*Sterculia scaphigera*) the seed of which when immersed in water produces a mucilage used medicinally by the Malays ; Rumania (*Bouea microphylla*) ; various species of *Garcinia* producing the fruits known as Kandis. Asam Gelugur (*Garcinia atroviridis*) was met with in a wild or half wild state in many spots along the route. The red fruited Mangosteen (*G. hombroniana*) grew abundantly in the sandy country near Kuala Pahang, but was not in fruit at the time of our visit.

Of eatable smaller berried fruit, the Kamuning (*Rhodomyrta tomentosus*) ; Nasi-nasi (*Eugenia zeylanica*) ; Mata Pelandok (*Ardisia crenulata*) ; the wild raspberry (*Rubus moluccanus*) were all abundant in the open country.

H. N. R.



LIST OF MAMMALS RECORDED FROM PAHANG.
[Specimens were obtained of those marked with a †.]

1. *Hylobates albimanus*, Vig. and Horsf.
The black Wau Wau, "Ungka" of the Malays, is common in the Pahang jungles.
2. *Macacus nemestrinus*, Linn.
The Coco-nut Monkey or Berok.
3. *Macacus cynomolgus*, Schreb.
The Mangrove Monkey or Kra is common along the Coast and for some distance up the rivers.
- †4. *Semnopithecus obscurus*, Reid.
The grey Lotong. This Monkey, which is common in Pahang, varies very much in colour. Here it is of a dark grey, while further South, in Johor, it is almost black.
5. *Nycticebus tardigradus*, Waterhouse.
The slow Loris.
6. *Felis tigris*, Linn.
The tiger, "Harimau" of the Malays, appears to be widely distributed, especially in the low country, disappearing in the hills.
7. *Felis pardus*, Linn.
The leopard or panther. There is considerable difference of opinion as to whether the leopard and so-called panther are distinct species. There are two distinct forms, one, which includes the black panther being much more thickset and heavy looking than the typical leopard and the spots are usually rings and not rosettes as in that animal. But different specimens vary so much in shade of colour and the form and arrangement of the spots that it is by no means easy to decide.

- The black variety appears to be by far the most common form in the Malay Peninsula, the yellow one being comparatively seldom met with.
8. *Felis Temminckii*, Vig. and Horsf.
The red tiger-cat. A living specimen from Pahang was for some time in the Botanical Gardens, Singapore.
There are probably several other species of wild cats found in Pahang, but there are no records of them. Tracks were often seen of small species in sandy spots.
9. *Viverricula malaccensis*, Gm.
The common Musang or small civet cat.
10. *Hemigale Hardwickii*, Less. Seen in Tahan woods.
11. *Cuon rutilans*.
The wild dog of the Malay Peninsula, called by the Malays "Anjing Utan" is said by them to hunt in packs. Living specimens have been sent down to Singapore from Pahang.
12. *Lutra leptonyx*, Gray.
The Otter. Malay "Anjing Ayer." Seen on the Tahan River.
13. *Helarctos Malayanus*.
The Malayan Honey bear. Tracks seen by the Tahan River. This little bear is easily kept in captivity and becomes exceedingly tame.
- †14. *Tupaia ferruginea*, Raffl.
Malay "Tupaia tanah."
- †15. *Tupaia javanica*, Horsf.
- †16. *Galeopithecus volans*, Linn.
The flying lemur. Kuala Tahan.
17. *Pteropus edulis*, Gray.
The large Fruit Bat, or, as it is sometimes called, the flying Fox. Common at Pekan.
There are many other bats, but few have as yet been collected and identified.
- †18. *Sciurus bicolor*, Sparr.
This very variable Squirrel is in Pahang usually of a

- dirty yellowish white colour, almost tawny on the back. Seen far up the Tahan River, and at Pulau Tawar.
- †19. *Sciurus Rafflesii*, Vig. and Horsf,
The Coco-nut Squirrel, "Tupai B'lang" of the Malays. This squirrel is most destructive amongst coco-nuts. It gnaws a round hole in the nuts to get at the inside. Common at Kuala Semantan.
- †20. *Sciurus laticaudatus*, Diard.
This appears to be a rare species. Kota Glanggi.
- †21. *Sciurus griseimanus*, A. M. Edwards.
- †22. *Sciurus insignis*, F. Cerv. This is a pretty squirrel with three black stripes down the back. It appears to live almost entirely on the ground. Tahan River.
- †23. *Sciurus notatus*, Bodd. This is perhaps the commonest of all the Malayan squirrels. Common everywhere.
- †24. *Hystrix longicauda*, Marsden.
The long-tailed porcupine. Two caught in the Kota Glanggi caves.
25. *Elephas maximus*, L.
The elephant, though common all through Pahang, is never caught and tamed. Abundant in the Tahan woods.
26. *Rhinoceros sp.* Tahan River woods. Tracks seen and animal heard at night.
27. *Tapirus malayanus*. Tracks have been seen of this animal at near Temerloh.
Bos gaurus, Ham. Smith.
The S'ladang is to be met with all through the Pahang jungles, but, owing to its shy and retiring habits, is difficult of approach. Tahan woods and all down Pahang River.
28. *Bos bubalus*, Ham. Smith.
The water buffalo is common in a semi-domesticated state along the rivers. It is very doubtful if it occurs in a wild state.
29. *Menorrhædus sumatrensis*, Shaw.

- The "Kambing Utan" of the Malays. This wild goat is supposed to occur in the high mountains of the interior, but there is no record of its having been obtained.
30. *Cervulus mountjac*, Brooke.
The "Kijang" of the Malays. Kuala Tahan.
31. *Cervus equinus*. Cuv. "Rusa" or "Rusa Hitam."
The Malayan Sambur, apparently the same as the Indian species, but the horns do not attain the same length. Heard at night along the Pahang River.
32. *Tragulus napu*, A. M. Edward.
The large mouse-deer.
33. *Tragulus javanicus*, A. M. Edward.
The small mouse-deer.
- †34. *Sus cristatus*, Wagner.
The wild pig, common in low country; seen also far up the Tahan.
35. *Manis javanica*, Desm. The Pangolin or "Tingiling."
The Scaly Armadillo. Pramau, Pekan.

LIST OF BIRDS OBSERVED OR COLLECTED
DURING TRIP IN PAHANG.

1. *Accipiter virgatus*, Temm.
This sparrow-hawk is fairly common.
2. *Haliaetus leucogaster*, Gm.
The white-bellied sea eagle is common along the coast and for a considerable distance up the river.
- 2a. *Haliaetus plumbeus*. Tembeling River.
3. *Haliastur indus*, Bodd.
The Brahmany kite or maroon kite. This, the common kite of Singapore, is also abundant in Pahang for a considerable distance up the river.
- †4. *Microhierax fringillarius*, Drap.
The smallest falcon in the world, scarcely as large as a common starling. Fairly plentiful. Kuala Berar.

5. *Ketupa javanensis*, Less.
The fishing owl is found in the low lands near the Coast. Kuala Pahang.
Several other species of owl were heard, but, owing to their nocturnal habits, it is difficult to obtain specimens of these birds.
6. *Corone macrorhyncha*, Wagl.
The common crow of the coasts and open country.
7. *Corone enca*, Sharpe.
This crow is only found in the jungles and singly or in pairs.
8. *Platysmurus leucopterus*, Temm.
The white winged jay. Found in the higher jungles. Tahan woods.
9. *Oriolus xanthonotus*, Horsf.
10. *Dissemurus platurus*, Vieill.
The racket-tailed drongo is plentiful all through the Pahang jungles. This bird is a wonderful mocking bird, and imitates the notes of many other birds.
11. *Artamides sumatrensis*, S. Müll. Pulau Tawar.
12. *Campophaga minor*, Davison.
A new species taken at Pulau Tawar.
13. *Lalage orientalis*, Bodd. Common in open country.
14. *Pericrocotus igneus*, Blyth.
15. *Gerygone pectoralis*, Davison. A new bird shot among Casuarinas at Kwala Pahang.
16. *Rhipidura javanica*, Blyth. Very common in open country.
17. *Rhipidura perlata*, S. Müll.
18. *Terpsiphone affinis*, Hay.
19. *Terpsiphone incii*, Gould.
20. *Philentoma relatum*, Temm. Kwala Tahan.
21. *Philentoma pyrrhopterum*, Temm.
22. *Siphia elegans*, Temm. Kwala Tahan.
23. *Aegithina tiphia*, Linn. Pulau Tawar.
24. *Chloropsis zosterops*, Vig. Kota Glanggi.
25. *Pinarocichla lunulata*, Davison. A new bulbul.
26. *Criniger phaeocephalus*, Hartl. Tahan woods.

27. *Trachycomus ochrocephalus*, G. M. "Barabara" common near Pekan.
28. *Pycnonotus analis*, Horsf. The common bulbul.
29. *Pycnonotus plumosus*, Blyth. Common.
30. *Irena cyanea*, Begbie. Kwala Tahan.
31. *Copsychus musicus*, Raffl. Common in the cultivated country.
32. *Cittocincla tricolor*, Levaill. Pulau Chengei.
33. *Orthotomus ruficeps*, Lesson. Open country.
34. *Hydrocichla ruficapilla*, Temm. Kwala Tahan in streams.
35. *Pomatorhinus bornöensis*, Cab. Kwala Tahan.
36. *Stachyris nigricollis*, Temm. Pahang River. Common.
37. *Turdinus abotti*, Blyth. Common.
38. *Turdinus magnirostris*, Moore. Common.
39. *Turdinus macrodactylis*, Strickl. Kwala Tahan.
40. *Dryocalaphus nigricapitatus*, Eyton. Tanjong Loyang.
41. *Malacopterum magnum*, Eyton. Kwala Tahan.
42. *Malacopterum melanoccephalum*, Davison.
43. *Miscornis gularis*, Raffl. Common.
44. *Miscornis erythropterum*, Blyth. River banks etc. Common.
45. *Macrorhis ptilosus*, Lard & Selb. Common. Pahang river.
46. *Ptilocichla leucogastra*, Davison. River Tahan.
47. *Lanius cristatus*, Linn. The common shrike inhabits open country.
48. *Calchostetha insignis*, Lard. Open country.
49. *Aethopyga siparaja*, Raffl. Open country.
50. *Cinnyris pectoralis*, Horsf. Open country.
51. *Anthothreptes hypogrammica*, S. Müll. Kwala Berar, etc.
52. *Anthothreptes malaccensis*, Scop. Open country.
53. *Dicaeum cruentatum*, Linn. Open country common.
54. *Dicaeum chrysorrhoeum*, Temm. Open country common.
55. *Hirundo javanica*, Sparrm. Common swallow.

56. *Hirundo badia*, Cassin.
57. *Anthus rufulus*. Vieill. Meadow pipit. Open country.
58. *Passer montanus*. Linn. Common sparrow.
59. *Acridotheres torquatus*, Davison. Pulau Tawar.
60. *Mainatus javanensis*, Osb. Pulau Tawar. The "Tiong."
61. *Calornis chalybea*, Horsf. The green starling, common in open country near Pekan.
62. *Munia maja*, Linn. Common. A nest was taken at Pekan.
63. *Munia atricapilla*, Vieill. Common.
64. *Uroloncha acuticauda*, Hodgs. Common.
65. *Ploceus baya* Blyth. The weaver bird. Nesting at Pekan and Pulau Tawar.
66. *Pitta tucullata*, Hartt. Bentong.
67. *Pitta cyanoptera*, Temm. Bentong.
68. *Pitta megarhyncha*, Schl. Kwala Pahang.
69. *Pitta boschi*, Müll. Caves Kota Glanggi.
70. *Calypdomena viridis*, Raffl. Very common.
71. *Eurylaemus javanicus*, Horsf.
72. *Eurylaemus ochromelas*, Raffl. Black and yellow broadbill. Common on the Tembeling river.
73. *Corydon sumatranus*, Raffl. Malay "Tiong batu."
74. *Cymborhynchus macrorhynchus*, G.M. Very common all up the river.
75. *Chaetura gigantea*, Temm. Open country near Pekan.
76. *Macropteryx comatus*, Temm. Tahan river.
77. *Caprimulgus macrurus*, Horsf. Open country.
78. *Lyncornis temminckii*, Gould.
79. *Eurystomus orientalis*, Linn. The Eastern roller.
80. *Merops sumatranus*, Raffl.
81. *Merops philippinus*, Linn. Very common near Pekan.
82. *Nyctiornis amicta*, Temm. The forest bee eater.
83. *Pelargopsis malaccensis*, Sharpe. Common all up the Pahang river.
84. *Alcedo ispida*, Linn. Common small king-fisher.
85. *Alcedo meninting*, Horsf. Pulau Chengai, &c.
86. *Halcyon coromandus*, Lath. Kuala Pahang.
87. *Halcyon smyrnensis*, Linn. Open country along

Pahang river.

88. *Halcyon pileatus*, Bodd.
 89. *Halcyon humii*, Sharpe. Very common at Kuala Pahang.
 90. *Buceros rhinoceros*, Linn. Common up the river.
 91. *Anthracoceros convexus*, Temm. Very common along lower reaches of Pahang river.
 92. *Rhinoplax vigil*, Forst. River Tembeling.
 93. *Harpactes kasumba*, Raffl. The common Trogon Kuala Tahan &c.
 94. *Harpactes diardi*, Temm. do. do.
 95. *Harpactes duvaucelli*, Temm. do. do.
 †96. *Gecinus puniceus*, Horsf.
 †97. *Chrysophlegma malaccense*, Lath.
 †98. *Chrysophlegma humii*, Hargitt.
 †99. *Iyungipicus auritus*, Eyt.
 †100. *Lepocistes porphyromelas*, Boie.
 †101. *Miglyptes grammithorax*, Mach.
 †102. *Miglyptes tukki*, Less.
 †103. *Micropternus brachyurus*, Vieill.
 †104. *Tiga javanensis*, Ljung.
 †105. *Chrysocolaptes rallidus*, Temm. Kuala Tembeling.
 †106. *Humilophus pulverulentus*, Temm.
 †107. *Thriponax javensis*, Horsf.
 †108. *Calorhamphus hayi*, Gray. Very common in open country.
 †109. *Cyanops henrici*, Temm. Tahan jungles.
 110. *Cacomantis passerinus*, Vahl. Open country.
 111. *Centropus bengalensis*, G.M. Open country secondary jungle.
 112. *Rhinortha chlorophoea*, Raffl. The cat bird, common everywhere.
 113. *Psittinus incertus*, Salv. Small paroquet.
 113a. *Pavo muticus*. Common peacock.
 113b. *Gallus ferrugineus*. Common. Kuala Pahang.
 114. *Osmotreron olax*, Temm.
 115. *Osmotreron vernans*, Linn.
 116. *Ptilonopus jambu*, G. M.

- 117. *Chalcophaps indica*, Linn.
- 118. *Lobivanellus atronuchalis*, Blyth.
- 119. *Charadrius fulvus*, G. M.
- 120. *Dendrocygna javanica*, Horsf.
- 121a. *Aegialites godefroyi*. Kuala Pahang.
- 121b. *Gallinago sthenura*, Pin tailed snipe. Pulau Tawar.
- 122. *Butorides javanica*, Horsf.
- 123. *Herodias garzetta*, The common Egret. Pekan.
- 124. *Leptoptilus javanicus*, Adjutant or "Burong Babi"
common all along river.



A CATALOGUE OF THE FLOWERING PLANTS AND FERNS FOUND GROWING WILD IN THE ISLAND OF PENANG.

COMPILED BY

C. CURTIS, F.L.S.

Assistant Superintendent of Forests, Penang.



THE Island of Penang, situated in Lat. $5^{\circ} 24'$ North, and Long. $100^{\circ} 20'$ East, has an area of about 106 square miles.

The greater portion is hilly, the highest point being West Hill, about the centre of the Island, which is 2,750 feet above sea level. Government Hill, on which many of the plants recorded have been collected, is about 250 feet lower. From an altitude of about 1,000 feet, and in some places much lower, these hills are covered with a varied and luxuriant vegetation, conspicuous among the larger trees being Dipterocarps, *Agathis loranthifolius*, *Dacrydium elatum*, Oaks, Eugenias, &c. Underneath these is found a heterogeneous mass consisting of seedlings of the larger trees intermixed with a variety of shrubs and smaller trees, all struggling for supremacy.

Deep ravines are numerous, the bottoms of which are a mass of more or less rounded granite boulders piled one on top of the other in great confusion, so that to cross from one side to the other is often a matter of difficulty. In many of these ravines, up to 1,500-2,000 feet, there are small but permanent streams of water that can only be detected by the

sound of their rippling several feet below the surface, where they have found a passage among the boulders.

It is in such places, where the direct rays of the sun never penetrate, growing on these boulders, that the great wealth of Ferns, Aroids, *Medinillas*, *Rhododendron*, *Cypripedium barbatum*, *Didymocarps*, *Sonerilas*, small Orchids and such like things are most abundant.

The soil of all the hills is very similar, being of a reddish or yellow colour, and composed mainly of partially decomposed granite. In the lower lands where the soil is of a different nature, and where one might naturally expect to find a somewhat different flora, forests have long disappeared, and with them probably many plants that were once common. When forests have once been felled and burnt off there is an end to the most interesting vegetation, at least for a long period. Even though no cultivation be attempted most of the trees shrubs, &c. that spring up on the cleared land are different from the original ones, and of as little value commercially as they are botanically uninteresting. Among the first to assume possession, often to the exclusion of every other plant, is the "lalang" (*Imperata arundinacea*), and in places where this is absent "resam" (*Gleichenia* spp.). Of woody plants, *Rhodomyrtus tomentosa*, *Eurya acuminata*, *Trema amboinense*, and *Adinandra dumosa* are among the first and most numerous in a new clearing. The present site of the Botanic Garden had at some time in the earlier days of the Settlement of Penang been planted with cloves and nutmegs, but at the time the formation of a garden was commenced these trees had long been dead and the land was covered with secondary jungle, in many places as dense as that of the surrounding hills that apparently have never been cleared. Notwithstanding the fact that these hills rise abruptly on three sides and are covered to the summit with large *Dipterocarps*, *Sterculias*, *Eugenias*, *Swintonia*, &c., so that in a good seed-bearing season thousands of seeds must be washed down into the valley by the heavy rains, there was scarcely any trace of these re-occupying their original position on the land that had been cultivated.

What actually occupied the land was thousands of *Eurya*

acuminata, *Rhodomyrtus tomentosa*, *Melastoma malabathrica*; and in lesser numbers *Glochidium obscurum*, *Crypteronia pubescens*, *Morinda tinctoria*, and a few others. Where there were no trees the "lalang" had taken absolute possession.

The mean temperature of the Island is a little above 80° F., with a range of 14°-15°, and the average rainfall for the past five years 120 inches, spread more or less over the whole year, though most falls during the latter half.

There is no well-defined flowering season; some things can be collected at almost any time, while others appear to flower only at intervals of several years. During the past seven years there has been but one really good flowering season, which I attribute to the unusually long drought of the preceding period.

Several large collections of plants have, at various times, been made in Penang, the most important being those of WALLICH, MAINGAY, PORTER and PHILLIPS. A great many of the plants found by these, and other of the earlier collectors, I have been able to identify and hope yet to add largely to the number, but owing to the increased area under cultivation since their time, and the complete change that follows clearing, as already pointed out, it is probable that some of the plants recorded by them will not be again collected on the Island.

So far as I am aware no separate catalogue of plants collected in Penang has yet been published, and thus it is hoped that the present, though incomplete, may be of use to those interested in the botany and forestry of the region until material is available for a better.

Not having access to many books in which scattered records of Penang plants occur, I have included (in cases where I have not myself collected a specimen or seen the plant growing wild) only those for which Penang is given as a locality in the Flora of British India, Annals of the Royal Botanic Gardens Calcutta, and Materials for a Flora of the Malayan Peninsula, so far as these works are published; HOOKER'S Synopsis Filicum, and BEDDOME'S Ferns of British India. No doubt many plants, whose distribution is so general throughout the Malayan

Islands and the Peninsula that their occurrence in Penang is not mentioned in either of the above works, have been collected here, but if I have not seen them they are omitted from this list.

The earlier orders to the end of Tiliaceæ, and also Myristicaceæ and Cupuliferæ, have been carefully compared and determined by Dr. KING in the course of preparing the Annals and Materials, so that the determination in these orders may be accepted with greater confidence than others on which the same critical examination has not been brought to bear.

The most incomplete orders are those not yet taken up in the Flora of British India, as I have no means of ascertaining what species were collected by the earlier collectors, or of identifying those collected by myself. Mr. RIDLEY'S intimate knowledge of Orchids and Grasses has enabled me to identify nearly all the species collected.

Numbers bracketed thus (C. 241-259) are those under which specimens of the plant have been distributed, and the letter (C.) in brackets implies that though not distributed I have collected it, or it is well known to me in a living state. Malay names, where given, are printed in inverted commas, but these have been sparingly used, for, except in the case of plants used either as food or medicine, native names in Penang are unreliable. Even in the case of timbers in general use the same name is often applied to several species, and sometimes to trees belonging to different orders, so that their practical value as an aid to identification is not great. I am aware that in the other Settlement where there is still a considerable Malay agricultural population this does not apply with the same force.

I cannot sufficiently express my obligation to Dr. KING, the Kew authorities, Mr. H. N. RIDLEY and Bishop HOSE for assistance in determining my collections, and I take this opportunity of requesting those who have received sets to kindly point out any errors of determination that may come under their notice, and to supply the names when wanting.

Penang, 12th November, 1892.

RANUNCULACEÆ.

Clematis smilacifolia, Wall.; strong growing climber. Penara Bukit, rare. (C. 1048).

Naravelia laurifolia, Wall.; long slender climber. Waterfall, rare. (C. 1709).

DILLENiaceÆ.

Delima sarmentosa, L.; woody climber with white sweet-scented flowers. Not uncommon. (C. 294).

Tetracera assa, DC.; climbing shrub; the commonest plant of this order. (C. 69).

Tetracera macrophylla, Wall.; large woody climber. Government Hill. (C. 3008).

Tetracera, sp.; a larger and more robust plant than the preceding. Government Hill. (C. 1495).

Acrotrema costatum, Jack.; stemless, flower yellow; leaves silvery grey. Waterfall, common. (C. 456).

Wormia oblonga, Wall.; medium sized tree. Telok Bahang, not common. (C. 257).

Wormia Kunstlerii, King; tree 60-70 feet; fruit about 1 in. through, flowers not seen. Nalm Pass. (C. 2465).

Dillenia indica, L.; "Simpoh;" tree 40-60 feet; fl. white, 5-6 in. across. Not common. (C.).

Dillenia ovata, Wall.; small tree 20-30 feet; fl. yellow, 4 in. across, fruit globular 3-3½ in. through. Waterfall (C. 228).

Dillenia grandifolia, Wall.; I have not seen in flower, but have leaf specimens that agree with the description in Fl. B. I. (C.).

MAGNOLIACEÆ.

Magnolia Maingayi, King; tree 30-40 feet; bark grey, fl. white, fragrant. Government Hill at from 2,000-2,500 feet. (C. 292).

Michelia champaca, L.; largely cultivated and often found on abandoned land, but not truly wild in Penang. (C.).

Talauma mutabilis, Bl.; small tree, fl. white, sweet-scented. Not uncommon. (C. 1022).

Talauma lanigera, Hook. fil.; Small tree. Government Hill, rare. (C. 3008).

Kadsura scandens, Bl.; woody climber, fl. almost white, about $\frac{1}{2}$ in. in diameter. Penara Bukit. (C. 1529).

Kadsura lanceolata, King; slender woody climber. Not uncommon. (C. 2821).

Kadsura cauliflora, Bl.; stem stouter and leaves larger than in the preceding; carpel red, the size of a pea. (C. 2440).

ANONACEÆ.

Stelechocarpus, sp.; a tree 30-40 feet high, fl. greenish, carpel globular, $\frac{3}{4}$ in. Waterfall. (C. 2277).

Sageraca elliptica, Hook. fil.; a large tree. Not seen.

Cyathostemma viridiflorum, Griff. Not seen.

Cyathostemma Hookeri, King; long woody climber, fl. small, yellow. Waterfall and Government Hill. (C. 1213).

Uvaria dulcis, Dunal; woody climber. Government Hill and Muka Head, not common. (C. 729-1414).

Uvaria Lobbiana, Hk. fil.; large climber; not common; at low elevation. (C. 841-1234)

Uvaria excelsa, Wall.; large woody climber. Abundant. (C. 50).

Uvaria macrophylla, Roxb.; a large scandent or climbing shrub. This and the preceding are the two most common Uvarias in Penang. (C. 263, 842, 1312).

Uvaria purpurea, Bl.; large woody climber; found at low elevations only. (C. 234).

Uvaria hirsuta, Jack; long climber, petals red. Waterfall and Western Hill. (C. 1212).

Uvaria Curtisii, King; a long climber, fl. yellowish. Government Hill 2,000 feet, rare. (C. 1415).

Uvaria pauciovulata, Hk. fil. Government Hill. (C. 825).

Uvaria micrantha, Hk. fil. Not identified.

Uvaria sub-repanda, Wall.; scandent shrub. (C. 1408).

(*Ellipeia nervosa*, Hk. fil.; for which Penang is given as a locality in the Materials for a Flora of the Malayan Peninsula was collected in Perak.)

Cyathocalyx Maingayi, Hk. fil.; a large tree, flowers greenish yellow, Telok Bahang. (C. 1035).

Artobotrys suaveolens, Bl.; long climber. Common. (C. 26-2234).

Drepananthus pruniferus, Maingay; small tree, 25-40 feet high; branches long, drooping. (C. 1417).

Canangium odoratum, Baill. Not truly wild in Penang. (C).

Unona Dunalii, Wall.; tree. Muka Head. (C. 727).

Unona desmos, Dunal; a climber. West Hill and Batu Feringgi. (C. 807-1413).

Unona stenopetala, Hook. fil. King's collector, Scortechini.

Polyalthia clavigera, King; a tree 30-40 feet high; carpels yellow. Penara Bukit. (C. 2444).

Polyalthia macrophylla, Hook. fil.; a tree 30-40 feet high. Pulau Betong. (C. 2140).

Polyalthia cinnamomea, Hook. fil.; a tree 60 feet high; flowers dull red, principally on leafless branches. Not uncommon in the south of the island. (C. 2470).

Polyalthia oblonga, King; small tree. Common. (C. 1409-1277-1281).

Anaxagorea Scortechinii, King; a tree 30-40 feet high. Government Hill. (C. 1054).

Goniothalamus Kunstleri, King; an erect growing shrub 6-8 feet high. West Hill. (C. 1540).

Var. *macrantha*. Waterfall. (C. 1596).

Goniothalamus giganteus, Hook. fil.; a tree 40-50 feet, petals golden yellow. Waterfall. (C. 2162).

Goniothalamus macrophyllus, Hook. fil.; a shrub 6-8 feet high, scarcely at all branched. (C. 1577).

Goniothalamus tapis, Miq.; a large shrub. Government Hill 2,000-2,500 feet. (C. 302).

Mitrephora macrophylla, Oliver; a large shrub. (C. 813-1279).

Mitrephora Maingayi, Hook. fil.; small tree. Not uncommon. (C. 157-1553).

Popowia ramosissima, Hook. fil.; a small tree. Collected Pulau Betong. Not distributed. (C.).

Popowia tomentosa, Maingay; a small tree. Government Hill 2,000-2,500 feet (C. 648).

Popowia nervifolia, Maingay; a small tree. (C. 893).

Oxymitra calycina, King; a woody climber. Government Hill 2,000 feet, rare. (C. 767).

- Oxymitra glauca*, Hook. fil. Not identified.
Melodorum manubriatum, Hook. fil. Collected by Maingay.
Melodorum latifolium, Hook. fil.; a large climber reaching to the top of tall trees. Common. (C. 1411-1196).
Melodorum lanuginosum, Hook. fil.; large woody climber. Government Hill 1,000-2,000 feet. (C. 797-1195-1383).
Melodorum Maingayi, Hook. fil.; large climber. (C. 1046).
Melodorum prismaticum, Hook. fil. Government Hill. (C. 730).
Melodorum rubiginosum var. *oblongum*, King. Waterfall (C. 843).
Melodorum elegans, Hook. fil. Not seen. Wallich 6474a.
Melodorum pisocarpum, Hook. fil. Not seen.
Xylopia Curtisii, King; a tree 30 feet high. (C. 1569).
Xylopia elliptica, Maingay; a tree 25-30 feet high; petals yellowish white. Sungei Ujong. (C. 2482).
Xylopia stenopetala, Oliver; a tree 59-60 feet high. Government Hill 690 feet. (C. 857-880).
Phæanthus nutans, Hook. fil.; a small tree. Government Hill 2,000 feet. (C. 1416).
Phæanthus lucidus, Oliver; a small tree. Waterfall. Not uncommon. (C. 839-1407).
Alphonsea Curtisii, King; a scandent shrub. Muka Head. (C. 1410).
Mezzettia Curtisii, King; a tree 30-40 feet high; carpels about 2 in. in diam. (2266).
 Undetermined sps. of Anonaceæ in Herb. Penang. (C. 2744, 2745, 2775, 2768).

MENISPERMACEÆ.

- Tinospora crispa*, Miers; climbing shrub. Waterfall. Not common. (C. 1231).
Tinomisium petiolare, Miers; climber. Penara Bukit and Pulau Betong. (C. 208).
Fibraurea chloroleuca, Miers; large woody climber. Waterfall and West Hill. (C. 208).
Cosciniun Blumeianum, Miers; large climber; ripe drupes round, tomentose, $\frac{3}{4}$ in. (C. 8).

Limacia triandra, Miers; climbing shrub. Tanjong Bunga, near the beach. (C. 447).

Limacia oblonga, Miers; a climber. Government Hill. (C. 671).

Limacia velutina, Miers; a woody climber. At from 1,000-2,000 feet elevation. (C. 2296).

Hyserpa triflora, Miers, "Kelintek Nyamok." A slender climbing shrub. (C. 949-1558).

Pericampylus incanus, Miers; a slender climber. The most common plant of this order in Penang. (C. 122).

Stephania hernandifolia, Walp.; slender climber. Penara Bukit 1,000 feet. (C. 1260).

NYPHEACEÆ.

Nymphea stellata, Willd. "Klipoh." Common in ditches and paddy fields. (C. 518).

CRUCIFERÆ.

Cardamine hirsuta, L.; small annual herb found sparingly on the top of Government Hill. (C. 2156).

CAPPARIDÆ.

Cleome viscosa, L.; annual herb 1-2 feet. Coast. (C. 1859).

Capparis Scortechinii, King; scandent shrub. I know of but one plant at Batu Feringgi. (C. 239).

Capparis micrantha, DC.; shrubby. Not common. (C. 1762).

VIOLACEÆ.

Viola serpens, Wall.; occurs sparingly on the top of Government Hill. (C.).

Alsodeia Wallichiana, Hk. fil.; erect shrub 3-5 feet; seen only at the Waterfall. (C. 1532-1899).

Alsodeia lanceolata, Wall.; small shrub, the only plant of the genus that is at all common. (C. 94-179).

Alsodeia lanceolata var. *foliis latioribus* is a much larger plant than the type. (C. 75).

Alsodeia Kunstleriana, King; shrub about the same size as *A. Wallichianus*. Waterfall, Not common. (C. 1898).

Alsodeia echinocarpa, Korth, is mentioned in Dr. King's

Materials for the Flora of Malayan Peninsula as occurring in Penang but I have not seen it.

BIXINEÆ.

Scolopia rhinantha, Clos.; small tree. Government Hill. Not common. (C. 991).

Scolopia Roxburghii, Clos.; tree 20-30 feet. Government Hill. (C. 778).

Scolopia crenata, Clos.; large shrub, sometimes small tree. Not uncommon near the coast. (C. 706).

Flacourtia inermis, Rox., is said to have been collected in Penang by Jack. It is known to me as a cultivated tree only. (C.).

Flacourtia rukam, Zoll. & Moritz; a tree attaining a height of about 30 feet. Not uncommon. (C. 1566).

Flacourtia cataphracta, Rox., was collected on Government Hill by Kunstler. Not seen.

Hydnocarpus Curtisii, King; small tree. Batu Feringgi and Waterfall. (C. 800-1535).

Hydnocarpus nana, King; small tree. Telok Kumbar, rare. (C. 854).

Ryparosa Scortechinii, King; medium-sized tree, flowers from the stem. (C. 1741).

PITTOSPOREÆ.

Pittosporum ferrugineum, Ait., "Chabe hantu"; small tree, common about Batu Feringgi. (C. 453-1533).

POLYGALEÆ.

Polygala venenosa, Hassk.; shrub 2-3 feet. Not uncommon in damp shady places at about 2,000 feet. (C. 291).

Polygala brachystachya, Bl. Flowers yellow in open grassy places. (C.).

Trigonostrium hypoleucum, Miq.; tree 30-40 feet with long slender branches. Government Hill. (C. 387).

Salomonina oblongifolia, DC.; dwarf herb, common in open grassy places. (C.).

Salomonina cantoniensis, Lour. Waterfall. (C. 1599).

Xanthophyllum affine, Korth.; tree not large; flowers open white, soon change to yellow. (C. 1188-2263).

Xanthophyllum Maingayii, Hook. fil.; small tree; flowers white. Government Hill, rare. (C. 1439).

Xanthophyllum eurhynchum, Miq.; tree. Telok Bahang near the coast. (C. 1552).

Xanthophyllum Wrayii, King; small tree. Balik Pulau. (C. 677).

Xanthophyllum pulchrum, King. Penara Bukit, rare. (C. 1167).

Xanthophyllum Kunstleri, King; medium-sized leafy tree 40 feet high. Government Hill. (C. 1590).

Xanthophyllum Curtisii, King; small tree; flowers yellowish white. Moniot's Road. (C. 1591).

Xanthophyllum Scortechinii, King; tree 25-30 feet, flowers pink; fruit globose 2½ in. Government Hill. (C. 2407).

Xanthophyllum, sps. undetermined. (C. 474 and 1486).

PORTULACÆ.

Portulaca oleracea, L.; annual herb, sometimes used as a salad. Common. (C. 1650-1875).

Portulaca quadrifida, L.; small succulent herb. Collected in the Fort. (C. 2148).

HYPERICINÆ.

Hypericum japonicum, Thunb.; dwarf herb with yellow flowers. Rice fields, south of the Island. Common. (C. 1946).

Cratoxylon arborescens, Bl. var. *Miquelii*; large tree, 60 ft. Government Hill 1,500-2,000 feet. Common. (C. 285).

Cratoxylon polyanthum, Korth. "Drum"; medium-sized tree. Not common. (C. 1168).

Cratoxylon polyanthum var. *ligustrinum*, Bl. (C. 824).

Cratoxylon formosum, Bth. & Hk.; tree about 30 feet; the most common of the genus. (C. 172-665).

Cratoxylon Maingayii, Dyer, is probably the plant in Herb. Penang (C. 2484).

GUTTIFERÆ.

Garcinia merguensis, Hook. fil.; small tree. Telok Bahang. Not uncommon. (C. 900-2408).

Garcinia eugeniæfolia, Wall.; small tree. Government Hill 1,500 feet. (C. 669).

Garcinia Scortechinii, King; tree 50-40 feet. Telok Bahang. Not common. (C. 1249).

Garcinia Hombroniana, Pierre; fruit much like a mangosteen but smaller. A common tree. (C. 690-2453).

Garcinia cornea, L. (C. 690 perhaps belongs to this species according to Dr. King).

Garcinia fascicularis, Wall.: small tree. Waterfall. (C. 840-1421).

Garcinia Penangiana, Pierre; small tree. Waterfall and Government Hill up to 1000 feet. (C. 1430-1549-1514).

Garcinia bancana, Miq.; small tree, seen only near the Coast. (C. 240).

Garcinia atro-viridis, Griff. "Asam Gelugur"; tall handsome tree; fruit yellow grooved, larger than mangosteen. Often cultivated. (C. 855).

Garcinia Griffithii, T. And.; small straight tree. (C. 1609).

Garcinia nigro-lineata, Pl.; tree 30-40 feet, branches slender; fruit 1-1½ inch yellow. (C. 1420-2478).

Garcinia Xanthochymus, Hook. fil. Not seen.

Garcinia, sps. undetermined and apparently distinct from any of the above. (C. 2293) (899-1424) 2412-1080 (691-1423) (2246) (1550).

Calophyllum spectabile, Willd. All the sps. of *Calophyllum* produce valuable timber known locally as "Bintangor." (C. 1153-1154).

Calophyllum canum, Hook. fil.; medium-sized tree. Government Hill. (C. 1543).

Calophyllum pulcherrimum, Wall.; tall straight tree. (C. 418).

Calophyllum Kunstlerii, King; medium-sized tree. Common along the coast. (C. 1425).

Calophyllum Curtisii, King; 40-50 feet; fruit the size of a pea. Government Hill. Rare. (C. 523).

Calophyllum molle, King; small tree. Government Hill 1,000 feet. Not common. (C. 1426).

Calophyllum inophyllum, L. "Penaga"; medium-sized

tree; occurs sparingly along the sea shore. (C. 133).

Calophyllum sp. near *floribundum*; medium-sized tree. Government Hill 1,000 feet. (C. 830).

Kayea racemosa, Pl. and Trian.; tree 40-50 feet Government Hill, rare. (C. 1441).

Kayea Kunstlerii, King; small much branched tree Water-fall, &c., common. (C. 805, 1418, 1419).

Kayea near *neuvosa*, T. And.; West Hill. (C. 748).

Mesua ferrea, L. "Matopus"; large tree, valuable timber. Muka Head and Government Hill. (C. 420).

TERNSTRÆMIACEÆ.

Ternstræmia penangiana, Choisy; tree 30-50 feet; flowers yellowish white. Government Hill. (C. 905).

Ternstræmia coriacea, Scheff.; medium-sized tree; bark grey, smooth. Top of Government Hill. (C. 1055).

Adinandra dumosa, Jack. "Kayu Gula"; tree 20-30 feet, Common on abandoned land in secondary jungle. (C. 35, 1725).

Adinandra acuminata, Korth.; small tree, does not occur below 1,006 feet. (C. 482).

Adinandra Miquelii, King. (C. 1612).

Adinandra integerrima, T. And. Not identified. Wallich Nos. 3663 and 7070.

Adinandra maculosa, T. And. Not identified.

Adinandra Hulletti, King. (C. 275, in part).

Adinandra villosa, Choisy; small tree 25 feet. Government Hill 2,000 feet, rare. (C. 2241).

Eurya acuminata, DC.; small tree 25-30 feet. Very common in places. (C. 299).

Eurya japonica var nitida, Penang, Lobb in fil. Brit. Ind. is probably an error.

Saurauja tristyla, DC.; small tree. Government Hill, damp shady ravines. (C. 786).

Schima Noronhæ, Reinw. "Medang Bequoi"; medium-sized tree; wood soft. Moderately common all over the island. (C. 161, 166).

Pyrenaria acuminata, Planch. is said to occur in Penang. Not seen.

Gordonia excelsa, Bl.; tall tree; flowers large yellow. Government Hill, rare. (C. 834).

Gordonia sp.; tree 20-30 feet. Top of Government Hill. (C. 2281).

Gordonia anomala, Spring, is a doubtful Penang plant stated to have been sent to Calcutta from Penang. It is a native of China.

Archytæa Vahlîi, Choisy; shrub 6-10 feet; flowers pink. Batu Feringgi, not seen elsewhere. (C. 1069).

DIPTEROCARPEÆ.

Dipterocarpus grandiflorus, Blanco; "Kayu minyak"; a large tree which yields an useful oil. (C. 424).

Dipterocarpus cornutus, Dyer; large tree 70-80 feet high. Waterfall, not common. (C. 1402).

Dipterocarpus fagincus, Vesque; "Kruen"; a large tall tree yields oil; moderately common on Government Hill up to 1,000 feet. (C. 1401).

Dipterocarpus, sp.; much resembling the preceding and scarcely distinguishable when not in fruit. Government Hill. (C. 1560).

Dipterocarpus Kerrii, King; a large tall tree closely resembling the species named after Dr. KERR, but in the absence of flowers there is some doubt. Waterfall, not common. (C. 1653). Typical *D. Kerrii* was collected on Gunong Tunggal in the Dindings and the local name is "Kruen chaia." (C. 1561).

Dipterocarpus Skinnerii, King; an exceedingly rare species named in honour of the Resident Councillor of Penang. A tree about 70 feet high, bark rough, warty. West Hill 1,500 feet, only one tree seen. (C. 1403).

Ancistrocladus extensus, Wall. var. *pinangianus*; a large woody climber, abundant in places. (C. 774).

Anisoptera Curtisii, Dyer. "Rengkon"; large tree, pretty common on Government Hill up to 2,000 feet. (C. 428).

Vatica pallida, Dyer; small tree, branches slender, drooping. Common on the lower slopes of the hills and one of the few Dipterocarps that flowers annually. (C. 117).

Vatica Curtisii, King: medium-sized tree, fruit in young state brick red. Waterfall, not common. (C. 1579).

Vatica nitens, King; middling-sized tree, not common. Telok Bahang. (C. 1404).

Vatica Wallichii, Dyer, "Ressak" small tree, bark smooth grey. Abundant. (C. 1218 1161 1391).

Shorea glauca, King, "Damar laut daun besar"; large tree wood hard and durable, one of the best forms of timber that are locally known as "Damar laut." (C. 372).

Shorea sericea, Dyer, "Seraya"; large tree, wood easily worked and used for a variety of purposes. Not common in Penang. (C. 431).

Shorea Curtisii, Dyer; "Meranti tai"; tree 60-80 feet high, leaves silvery grey, wood soft but much used for indoor work. More or less common all over the island. (C. 427-1394-1395).

Shorea ciliata, King; medium-sized tree, wood hard, durable. Government Hill, rare. (C. 1578).

Shorea utilis, King, "Damar laut No. satu"; large tree, wood hard and close grained. Undoubtedly the best timber in the island. Waterfall and Muka Head, too much in demand to be common. (C. 423).

Shorea pauciflora, King; medium-sized tree. Government Hill. (C. 1527).

Shorea parvifolia, Dyer, "Meranti daun kecil"; tall straight tree, bark rough deeply grooved. Abundant. C. 201).

Shorea, sp. near *parvifolia* but with smaller fruit and probably a distinct species. West Hill 2,000 feet. (C. 437).

Shorea costata, King, "Damar laut"; large tree. Government Hill, rare. (C. 199).

Shorea bracteata, Dyer, "Semah" tall straight tree. Waterfall, not common. (C. 1405).

Shorea macroptera, Dyer, tree 50-60 feet high. Government Hill. (C. 1392).

Hopea intermedia, King. "Jankang"; medium-sized tree, not uncommon. (C. 425-1397).

Hopea micrantha, Hook. fil.; medium-sized rather tall tree much like the preceding. Government Hill (C. 266).

Hopea Curtisii, King; large straight tree 60-80 feet high; immature fruit greenish yellow. Waterfall. (C. 1562).

Balanocarpus pinangianus, King. "Damar itam"; large tree. Government Hill, &c., common. (C. 1, 429, 1393).

Balanocarpus Curtisii, King; small tree 15-30 feet high, bark smooth, polished. Waterfall, abundant in two places. (C. 1406).

MALVACEÆ.

Sida carpinifolia, L. "Katombe." A shrub 1-2 feet high flowers yellow. Common. (C. 946, 1894).

Sida rhombæfolia, L. Not uncommon. (C.).

Urena lobata, L. "Pulut"; very common. (C. 3).

Abutilon indicum, G. Don; dwarf shrub. Open places near the coast. (C. 127).

Hibiscus surattensis, L.; climber. Not common. (C. 70).

Hibiscus abelmoschus, L.; under shrub with large yellow conspicuous flowers. Open places not common. (C. 1925).

Hibiscus macrophyllus, Rox. Recorded from Penang with no collector's name in Fl. Brit. Ind. probably an error. Not collected by me.

Hibiscus tiliaceus, L.; littoral tree, 30-40 feet high. Common. (C. 274).

Thespesia populnea, Corr. "Baru"; a tree 20-30 feet. Not common. (C. 1715).

Eriodendron anfractuosum, D.C. A tall tree found in many native gardens, but not truly wild in Penang. (C.)

Durio zibethinus, L. "Durian"; cultivated, and often found in a half wild state on abandoned land. (C.).

Durio testudinarum, var. *pinangianus*, Beccari; a tall straight tree; flowers and fruit from the stem, generally within a few inches of the ground. Fruit round, the size of a large orange. West Hill 2,500 feet. (C. 293).

Boschia Griffithii, Mast.; a tree 30-40 feet; fruit bright red. Telok Bahang. (C. 381).

Neesia synandra, Mast. Penang, Maingay. Not since collected here.

STERCULIACEÆ.

Sterculia parvifolia, Wall.; small tree 15-20 feet. Government Hill. (C. 526).

Sterculia lævis, Wall.; small tree. Government Hill. (C. 1624).

Sterculia rubiginosa, Vent.; tree 15-20 feet. Penara Bukit. Not common. (C. 1195-2471).

Sterculia ensifolia, Mast.; shrub 6-10 feet flowers dull red. Waterfall. (C. 272, 1428, 2299).

Sterculia parviflora, Roxb.; medium-sized tree 20-30 feet, conspicuous when in fruit. Not common. (C. 770-1429).

Sterculia colorata, Roxb.; small tree. Government Hill, rare. (C. 144).

Sterculia hyposticta, Miq.; large shrub. Government Hill, (C.)

Sterculia macrophylla, Vent.; tree 30-40 feet. Waterfall, rare. (C. 3008).

Sterculia, sp. large tree. Waterfall. (C. 2762).

Sterculia campanulata, Wall. large tree, deciduous; fruit white. Bukit Pulai. (C. 2783).

Tarrietia perakensis, King; tree 40-60 feet high. Government Hill 2,400 feet. (C. 2229).

Tarrietia Curtisii, King; leafy spreading tree 30-40 feet; leaves digitate; samara almost black 1,000-2,000 feet. (C. 1427).

Heriteria littoralis, Dryand, "Dungun"; medium sized tree. Tidal swamps, common. (C. 517).

Helicteres isora, L.; small tree. Government Hill. (C. 1012).

Helicteres hirsuta, Lour. var. *oblonga*. Wallich No. 1183. George Porter. (C.)

Pterospermum Jackianum, Wall.; tall straight tree. Waterfall, not uncommon, (C. 783).

Pterospermum Blumeianum, Korth.; small tree. Balik Pulau. (C. 2772).

Melochia corchorifolia, L.; common undershrub. (C. 413).

Buettneria Jackiana, Wall.; climbing shrub. (C. 86).

Buettneria Curtisii, Oliv.; climbing shrub, flowers white,

Batu Feringgi and Telok Bahang .(C. 817-1166).

Commersonia platyphylla, And.; tree 30-40 feet; fl. white. Waterfall, not common. (C. 776).

Leptonychia glabra, Turcz.; erect shrub 6-10 feet. Common. (C. 38, 407).

TILIACEÆ.

Corchorus olitorius, L. Telok Bahang. (C. 2483).

Pentace Curtisii, King; large tree; flowers white. Waterfall. (C. 1573).

Pentace, sp.; small tree, flowers much smaller than those of *P. Curtisii*. Penara Bukit. (C. 3006).

Schoutenia accrescens, Mast.; tree 50-60 feet; calyx papery, yellowish white. Waterfall. (C. 1520).

Grewia umbellata, Rox.; scandent shrub. Common. (C. 444-714).

Grewia fibrocarpa, Mast.; small tree; drupe pyriform, yellow. Government Hill. (C. 522-1712).

Grewia globulifera, Mast.; medium-sized tree. Penara Bukit. (C. 2414).

Grewia laurifolia, Hk. fil.; small tree. Government Hill 500 feet (C. 183, 1,488).

Grewia paniculata, Roxb. "Chindrie"; small tree 15-30 feet high; ripe drupes the size of a pea, black edible. (C. 108).

Triumfetta rhomboidea, Jacq.; undershrub 2-3 feet high Coast. (C. 484, 2,141).

Elæocarpus Ganitrus, Roxb.; tree 30-40 feet; Penara Bukit, not common. (C. 775).

Elæocarpus parvifolius, Wall.; tree 25-30 feet not uncommon at 2,000-2,500 feet on Government Hill. (C. 310, 376).

Elæocarpus stipularis, Bl.; not seen.

Elæocarpus nitidus, Jack; small tree. Waterfall. (C. 282-463).

Elæocarpus robustus, Roxb.; large tree. Telok Bahang. (C. 1,049).

Elæocarpus glabrescens, Mast.; middling sized tree. Government Hill. (C. 1,092).

Elæocarpus Griffithii, Mast.; not seen. King's collector.

Elæocarpus pedunculata, Wall. ; small tree, rather common and conspicuous when in flower. (C. 256).

Elæocarpus apiculatus, Mast. Collected by Wallich No. 3969.

Elæocarpus petiolatus, Wall. ; tree 25-30 feet. Waterfall, not uncommon. (C. 383, 464).

Elæocarpus Hullettii, King ; small tree. (C. 1,091).

Elæocarpus Jackiana, Wall. ; tree. (C. 465).

LINEÆ.

Roucheria Griffithiana, P. C. ; large climbing shrub, flower yellow. (C. 165, 207).

Erythroxyton burmanicum, Griff. "Chinta mula" ; tree 25-30 feet, fruit red. Waterfall. (C.).

Ixonanthes icosandra, Jack. Wallich No. 4409 and 4802. (C. 717).

Ixonanthes obovata, Hook. fil. ; not seen.

Ixonanthes reticulata, Jack ; medium-sized tree. Government Hill. (C. 978).

MALPIGHIACEÆ.

Brachylophon Curtisii, Oliv. ; erect shrub 3-5 feet, fl. yellow. Not uncommon along the Coast. (C. 231).

Hiptage madablota, Gærtn. ; climbing shrub. (C. 1159-1561).

Hiptage sericea, Hk. fil. Wallich 1814. Balik Pulau. (C. 1525).

Aspidopterys concava, A. Juss. ; flexuous climber. Government Hill 1,000 feet. (C. 138-798).

Aspidopterys, sp. ; climber. Telok Bahang. (C. 195).

GERANIACEÆ.

Oxalis corniculata, L. ; common creeping weed with yellow flowers. (C. 1836).

Oxalis Martiana, Zucc. ; herb common on Government Hill. Introduced. (C. 1726).

Biophytum sensitivum, D.C. ; herb 6-10 in. petals yellow ; not common. (C. 2144).

Hydrocera triflora, W. & A. Paddy fields and ditches. (C. 1190).

RUTACEÆ.

Evodia Roxburghiana, Benth. Not seen. Collected by Wallich No. 8065.

Evodia robusta, Hook. fil.; large shrub. Government Hill. (C. 1485).

Evodia latifolia, D.C.; small tree. Pulau Betong. (C. 731-939).

Evodia glabra, Bl. Tree 40-50 feet. Waterfall. (C. 2428).

Tetractomia Roxburghii, Hk. F.; tree 20-30 feet; fl. white. Government Hill, not uncommon. (C. 281).

Xanthoxylum myriacanthum, Wall.; tree 25-30 feet; stem covered with prickles, rare. (C. 1,076).

Acronychia laurifolia, Bl.; tree 20-30 feet (C. 315, 374, 694).

Acronychia Porteri, Hk. fil.; a smaller and much less common tree than the preceding. (C. 1,140, 1,162).

Glycosmis pentaphylla, Correa; small shrub, fruit round white $\frac{1}{4}$ inch in diameter, a very variable plant. (C. 89).

Glycosmis sapindoides, Lind.; small shrub. Government Hill, not uncommon. (C. 722).

Glycosmis puberula, Lind.; small tree. Government Hill, not uncommon. (C. 88, 1,223).

Micromelum pubescens, Bl. (C. 1,760).

Micromelum hirsutum, Oliv.; shrub. Ayer Hitam. (C. 1436).

Clauseria excavata, Burm.; "Chenamah" small tree 10-20 feet. Common near the Coast. (C. 150).

Paramignya Griffithii, Hk. fil.; Wallich No. 6,358. Penang. G. Porter is the only authority for this, which was probably not collected here.

Paramignya monophylla, Wight.; large climbing shrub with axillary spines 1 inch long; fruit (unripe) $\frac{1}{2}$ - $\frac{3}{4}$ inch long. Waterfall. (C. 2,207).

Atalantia Roxburghiana, Hk. fil.; "A native of Pulo Pinang" Roxburgh. A very little known plant. Not identified.

Atalantia monophylla, Corr.; small tree; not uncommon. (C. 467).

SIMARUBEÆ.

Ailantus malabarica, D. C.; tall tree, conspicuous when in

young fruit. Waterfall. (C. 710).

Eurycoma longifolia, Jack.; "Tongkat barinda" small tree, common. (C. 141).

Eurycoma apiculata, A. W. B.; small tree 6-10 feet high. Waterfall. (C. 2,763).

OCHNACEÆ.

Gomphia sumatrana, Jack.; "Jangot Klee," small tree; petals yellow. Not uncommon near the Court. (C. 221).

Gomphia Hookerii, Planch, larger than the preceding; fruit bright red. Government Hill. (C. 1147, 2154).

Euthemis leucocarpa, Jack.; small erect shrub; berries white (C. 1768).

Euthemis minor, Jack. Not seen.

BURSERACEÆ.

Canarium commune, L. The only tree known to me in Penang is cultivated. (C.).

Canarium nitidum. Benn.; small tree. Ayer Hitam (C. 495).

Canarium grandiflorum, Benn.; small tree. Government Hill (C. 803, 959).

Canarium, sp.; tree 30 ft.; fruit ovate, 1 in long. Muka Head and Government Hill. (C. 1432, 1433).

Canarium purpurascens, Benn.; small tree. Government Hill 1200 ft. (C. 862).

Canarium hirtellum, Benn.; small tree. Government Hill (C. 656).

Canarium laxum, Benn.; tree 30-40 ft. Government Hill (C. 1431).

Canarium, sp.; small tree. West Hill. (C. 1544)

Canarium pilosum, Benn. ?; tree 30-40 ft. (2251).

Canarium rugosum, Miq.; tree 20-30 ft. Muka Head. (C. 1434).

MELIACEÆ.

Melia tomentosa, Rox. Not identified. A drawing of this at Kew and Roxburgh's description are all that are known of it. Roxburgh says "a native of Pulo Pinang where it

grows into a large tree" and gives Malay name Barang bahee possibly an error for Berangan babi.

Melia excelsa, Jack. Not identified, collected by Jack only here.

Dysoxylum cauliflorum, Hiern; tree 30-40 feet, flowers white, often from old wood. Government Hill. (C. 204).

Dysoxylum cuneatum, Hiern; small tree. West Hill. (C. 1045).

Dysoxylum, sp.; near *binectariferum*, tree 30-40 feet; fruit pink, 2-3 in. Government Hill. (C. 2437).

Chisocheton spicatus, Hiern; small tree. Not common. (C. 655).

Chisocheton penduliflorus, Bl.; small tree 10-15 feet with a stem 2 in. diam. Government Hill. (C. 460).

Chisocheton divergens, Bl.; small tree. Pulau Betong, rare. (C. 892).

Chisocheton glomeratus, Hiern. Collected by Porter (Wall Cat 9040).

Chisocheton, sp.: small tree, racemes 20-24 in., pendulous; capsules 1 in., bright pink. (C. 2295).

Chisocheton princeps, Hemsley; tree 40 feet. Waterfall. (C. 1519).

Chisocheton, sp.; tree 60 feet; leaves 5-6 feet; panicles as long as the leaves; fl. dirty white. Pulau Betong. (C. 2469).

Chisocheton, sp.; tree 60-70 feet; capsule $1\frac{1}{2}$ - $1\frac{3}{4}$ in.; flesh colored, 1-2 seeded; arillus bright red. (C. 2467-2468).

Sandoricum indicum, Cov.; "Sentol" large tree, cultivated; doubtful if indigenous in Penang. (C.).

Aglaia odorata, Lour. is not wild in Penang. (C.).

Aglaia tenuicaulis, Hiern; small tree with a stem 2-3 in. diam. Government Hill. (C. 747).

Aglaia minutiflora, Bedd.; small tree; branches slender, drooping. Not uncommon. (C. 894-961).

Aglaia minutiflora, var. *Griffithii*. (C. 2003).

Aglaia palembanica, Miq. Government Hill. (C. 768).

Aglaia, sp. aff. *paniculata*, King. Small tree. (C. 895-896).

Aglaia, sp.; spreading tree 30-40 feet; fruit pyriform, silvery grey. Waterfall. (C. 2287).

Aglaia, sp. small tree; fruit ovate, brown. (2448).

Lansium domesticum, Jack. is cultivated, not wild. (C.)

Amoora rohituka, W. & A.; small tree. Not common. (C. 969).

Heynea trijuga, Rox.; large tree. Penara Bukit. (C. 676).

Carapa moluccensis, Lam.; "Nireh" small tree. Common in swamps. (C. 515, 742).

Cedrela Toona, Rox.; one very large tree on the top of Government Hill, probably planted there. (C. 826).

Meliacea? small tree. Moniot's Road 2,000 feet (C. 1,690).

CHAILLETIACEÆ.

Chailletia Laurocerasus, Pl.; scandent or climbing shrub, common on Government Hill. (C. 152).

Chailletia, sp. Wall. Cat., 7443 referred to in Fl. B. Ind. I have not seen; there appears but the one species in Penang.

OLACINEÆ.

Ximenea americana, Will.; large shrub. Bata Feringgi, in damp places. (C. 1572).

Erythropalum scandens, Bl.; climbing shrub. Waterfall. (C. 2777),

Ochanostachys amentacea, Mast.; tree 50 feet, fruit globular 2-2½ inch, Government Hill. (C. 1500).

Ctenolophon grandifolius, Oliv.; large tree; flower reddish, fragrant. Muka Head, rare. (C. 721).

Strombosia javanica, Bl. not met with.

Strombosia, sp.; not the preceding. (C. 859).

Lasianthera malaccensis, Mast.; small tree (C. 912, 957).

Gomphandra penangiana, Wall.; small tree 10-15 feet. (C. 1225).

Gomphandra, sp. shrub 2-4 feet; leaves ¼ + 4-5 inch. West Hill 2,500 feet. (C. 1,265, 737).

Pteleocarpa malaccensis, Oliv.; tall tree; flower yellow, very showy when in bloom, not uncommon. (C. 835, 1,494).

Phytocrene bracteata, Wall.; large climber. Balik Pulau. (C. 1754).

Phytocrene oblonga, Wall. Not seen. Porter collected it.

Phytocrene palmata, Wall. Much more common than *Ph. bracteata*. Government Hill, &c. (C. 2327).

Iodes oblonga, Pl.; long slender flexuous climber; flower white. Waterfall. Not common. (C. 2438).

Platea latifolia, Bl.? tree 30-40 feet; fruit pear-shaped, yellow, 1 inch long. Near the waterfall. (C. 2,421).

ILICINEÆ.

Ilex macrophylla, Wall. "Medang tulok"; tree 25-30 feet. Common in all parts of the Island. (C. 290, 379, 304).

Ilex Maingayii, Hk. fil.; tree 20-25 feet. Top of Government Hill, rare. (C. 2152).

Ilex cymosa, Bl.; small tall tree. Batu Feringgi. (C. 1036).

CELASTRINEÆ.

Eunonymus javanicus, Bl.; small tree. Not uncommon. (C. 736).

Micropteris bivalvis, Wall.; small tree. Government Hill 2,000-2,500 feet. (C. 651, 652, 1025, 1727).

Micropteris, sp.; large shrub. Moniot's Road. (C. 345, 1531).

Micropteris, sp.; small tree. West Hill. (C. 968).

Lophopetalum pallidum, Laws.; medium sized tree (C. 1577).

Lophopetalum reflexum, Laws.?; tree 40-50 feet. (C. 1502).

Lophopetalum, sp.; tree. Government Hill. (C. 1501).

Lophopetalum, sp.; tree about 40 feet. Government Hill 2,000 feet. (C. 1576).

Celastrus monosperma, Wall.; large climbing shrub. Government Hill. (C. 404).

Kurrimia pulcherrima, Wall.; tree 30-40 ft. (C. 1064, 1106).

Kurrimia paniculata, Wall.; medium-sized tree 30-40 ft. Government Hill. (C. 307).

Hippocratea, sp.; small tree. West Hill, rare. (C. 175).

Salacia flavescens, Kurz; large scandent shrub. Government Hill. (C. 12).

Salacia longifolia, Hook. f.; small tree 20 ft. Government Hill, not uncommon. (C. 134, 1266).

Salacia grandiflora, Kurz; large scandent shrub. Government Hill. (C. 151).

Salacia grandiflora, variety?; a much smaller plant. (C. 249, 1737).

Salacia oblonga, Wall.? scandent shrub; fruit green, the size of a small orange. (C. 693).

Salacia Griffithii, Laws.? (C. 692).

Salacia sp.; scandent shrub. West Hill. (C. 1704).

Siphonodon celastrineus, Griff. not identified.

RHAMNEÆ.

Ventilago leiocarpa, Benth.; large climbing shrub. (C. 311, 1751).

Zizyphus Ænopia, Mill.; large shrub. (C. 107).

Zizyphus catophylla, Wall.; large climbing shrub. Not uncommon. (C. 119).

Colubrina asiatica, Brongn. "Prea Pantie;" a common coast plant. (C. 236).

AMPELIDEÆ.

Cissus quadrangularis, L.; is cultivated, the leaves and shoots being used in curries. Not seen in a wild state. (C.)

Cissus hastata, Planch. Not seen. Porter.

Cissus glaberrima, Wall. Not common. (C. 1138).

Cissus repens, Lamk. Penara Bukit, not common. (1135).

Cissus adnata, Roxb. Not seen.

Cissus carnos, Lam. Common. (C. 187).

Cissus angustifolia, Wall.; fl. sweet scented, retaining their perfume long after being dried. (C. 1137).

Cissus moltissima, Wall. Muka Head. (C. 1435).

Ampelocissus thyrsoiflora, Planch. (C. 3122).

Ampelocissus cinnamomea, Wall. (C. 47).

Ampelocissus compositifolia, Planch. Not identified. Penang, collected by Philips.

Ampelocissus nitida, Planch. Not identified. Wallich.

Ampelocissus polystachya, Planch. Collected by Walker.

Tetrastigma peduncularis, Wall. Not seen. Wallich No. 6024.

- Tetrastigma lanceolarium*, Planch. Government Hill 685.
Tetrastigma pedata, Vahl. Not identified. Lawson gives it for Penang in Flor. Brit. Ind. but with no collector's name.
Cissus, sp.; undetermined in herb. Penang. (C. 1051, 1241, 1245).
Pterisanthes araneosa, Miq.; leaves simple, cordate toothed. (C. 168).
Pterisanthes, sp.; leaves trifoliate, leaflets lanceolate, $\frac{1}{2}$ in. broad. Government Hill. (C. 761).
Leea rubra, Bl.; small erect shrub. Penara Bukit, rare. (C. 1107).
Leea sambucina, Willd.; erect shrub 6-10 feet. Very common. (C. 96).

SAPINDACEÆ.

- Cardiospermum halicacabum*, L.; annual climbing weed. Not common. (C. 349).
Erioglossum edule, Bl.; tree 20-30 feet; ripe fruit almost black. (C. 701).
Allophyllus Cobbe, Bl.; small tree. Government Hill, not uncommon. (C. 340, 771, 1264).
Mischocarpus fuscescens, Bl.; small tree. Government Hill. (C. 1243).
Cupania glabrata, Kurz., tree 30-40 feet. Top of Government Hill. (C. 1041).
Cupania pleuropteris, Bl.; small tree. (C. 646).
Var. bijuga, tree 20-40 feet (C. 789, 1158).
Cupania Lessertiana, Camb.; tree, small on the coast, larger on the Hill. (C. 227, 288).
Lepisanthes cuneata, Hiern. Not identified. Penang. Porter.
Otophora paucijuga, Hiern.; small tree. Government Hill. (C. 1164).
Xerospermum Noronhianum, Bl.; small tree. (C. 846, 1232, 2285).
Nephelium, sp.; large tree. Government Hill. (C. 1575).
Pometia tomentosa, Kurz.; large spreading tree, 50 feet high. (C. 888).

Dodonæa viscosa, L. Telok Bahang. (C. 2847).

Turpinia sphærocarpa, Hassk. ; tree 40-50 feet. Govern-
ment Hill 2000 feet. (C. 1033, 1717).

Sapindaceæ; small tree; leaves pinnate. (C. 1600).

SABIACEÆ.

Sabia limoniacea, Wall.; climbing shrub. Sungei Penang.
(C. 1221).

Meliosma lancifolia, Hook. fil.; small tree. Government Hill,
rare. (C. 1215).

ANACARDIACEÆ.

Mangifera quadrifida, Jack. Not identified.

Mangifera fætida, Lour. "Bachang"; large tree common
in orchards, often on abandoned land, but not truly wild.
(C. 1747).

Mangifera, sp.; large tree; leaves 12-15 in. by 3-4 in.;
fruit round, about 3 in. Government Hill. (C. 1598).

Mangifera, sp.; small tree; fruit not seen. (C. 1496).

Mangifera, sp.; tree 25-30 ft. fruit ovate, brown, 3 × 2½
in. long. Telok Bahang, rare. (C.)

Mangifera, sp.; tree 30-40 ft.; fruit globular 1½-2 in.
Telok Bahang. (C. 2306).

Anacardium occidentale, L. quite common in sandy places
near the coast. (Introduced) (C.)

Bouea microphylla, Griff. "Rumania;" compact growing
tree 20-40 ft. Waterfall and Government Hill. (C. 223).

Bouea macrophylla, Griff. is cultivated not wild in Pen-
ang. (C.)

Gluta elegans, Wall.; small tree; drupes 1-1½ in long;
calyx red, petals white. Common. (C. 153, 1062).

Buchanania lucida, Bl.; tree 20-40 ft. Government Hill.
(C. 319, 696).

Buchanania acuminata, Turcz.; medium sized tree.
(C. 154).

Melanorrhæa Curtisii, Oliv. "Rengas;" large tree, heart-
wood dark red, juice poisonous. (C. 242. 433).

Swintonia Griffithii, Kurz; large tall tree. Waterfall.
(C. 1579).

Swintonia spicifera, Hk. f. "Moopoo;" large tree, abundant in one or two places. (C. 371).

Odina wodier, Roxb.; I have seen but a single tree which may have been introduced. (C. 1499).

Parishia Maingayii, Hk. f.; small tree; young fruit full of white milky juice. Waterfall. (C. 2264).

Camptosperma Griffithii, March.; tree 50-60 ft.; leaves of young plants 2-3 times larger than in adult trees. Government Hill. (C. 1037).

Spondias mangifera, Willd. "Kadongdong"; often met with on abandoned land: not truly wild. (C. 752).

Dracontomelum mangiferum, Bl. Not met with. Collected by Maingay perhaps from a cultivated tree.

Anacardiaceæ, L.; tree 60 feet; bark rough, juice black, resinous. (C. 1567).

Anacardiaceæ, L.; tree 40-50 feet; drupe 1-1½ in. Government Hill. (C. 2475).

CONNARACEÆ.

Agelæa vestita, Hk. F.; large scandent shrub. Telok Bahang. Not common. (C. 1556).

Rourea rugosa, Pl.; scandent shrub. Government Hill. (C, 191).

Rourea similis, Bl. Government Hill, near the chalet. (C. 473).

Roureopsis pubinervis, Hook. fil.; collected by Porter. (Wall Cat 9050). Batu Ferengy. (C. 2749).

Connarus ferrugineus, Jack. Not seen.

Connarus semidecandrus, Jack. Not seen. Philips, Wallich.

Connarus gibbosus, Wall.; tree. Telok Bahang. (C. 1151 2899).

Connarus grandis, Jack; small tree. Telok Bahang. (C. 27).

Connarus oligophyllus, Pl. Not identified. Porter.

Cnestis ramiflora, Griff.; climbing shrub. (C. 145-1157).

Ellipanthus Griffithii, Hk. F.; small tree. Government Hill near the chalet. (1014-1097).

LEGUMINOSEÆ.

- Crotolaria retusa*, L.; undershrub. coast. (C. 109).
Crotolaria striata, D.C. is very common. (C.)
Millettia sericea, W. & A.; large climbing shrub; pod almost black, velvety. Penara Bukit. (C. 844).
Millettia eriantha, Benth.; climbing shrub. Telok Bahang, not common. (C. 338).
Millettia atropurpurea, Benth.; tree 30-40 ft. attaining a moderate size. Not uncommon. (C. 182).
Millettia, sp.; climbing shrub. Penara Bukit. (C. 1190).
Millettia, sp.; tree. Telok Bahang. (C. 1160).
Tephrosia Hookeriana, W. and A.; undershrub 4-6 ft. (C. 1878).
Tephrosia purpurea, Pers.; Undershrub 2-4 ft. (C. 1895).
Uraria crinita, Desv.; shrubby perennial; flowers lavender. (C. 916).
Alysicarpus vaginalis, D.C. is common in open grassy places, especially at the Waterfall. (C. 1893).
Desmodium umbellatum, D.C.; shrub 6-10 ft.; corolla white. Telok Bahang on the beach. (C. 1082).
Desmodium polycarpum, D.C.; shrub 2-3 ft. Ayer Hitam. (C. 392, 459).
Desmodium triflorum, D.C. Waterfall, common. (C. 1842).
Desmodium heterophyllum, D.C. not seen.
Abrus precatorius, L.; slender climber, very common. (C. 41).
Mucuna acuminata, Graham. not seen.
Erythrina indica, Lam. "Dadap;" is commonly planted, but not truly wild in Penang. (C.)
Spatholobus Roxburghii, Benth. var. *denudatus*. Not seen.
Spatholobus gyrocarpus, Benth.; large woody climber. Government Hill. (C. 271).
Spatholobus acuminatus, Benth. Probably not a native.
Spatholobus crassifolius, Benth. "Penang 1822. Wallich." Doubtless an error. It is a native of North India.
Dioclea reflexa, Hook. fil. not seen. "Penang or Singapore" Wallich. Probably not native.

- Pueraria phaseoloides*, Benth. Not identified.
Pachyrhizus angulatus, Rich. is cultivated. (C.).
Cajanus indicus, Spreng. Introduced.? (C. 118).
Flemingia congesta, Roxb.; shrub 2-4 feet. Not uncommon. (C. 87, 1926).
Dalbergia Championi, Thwaites?; large woody climber. Government Hill. (C. 1572).
Dalbergia tamarindifolia, Roxb.; large woody climber. Waterfall. (C. 1492).
Dalbergia monosperma, Dalz.; climbing shrub. Telok Bahang. (C. 220).
Dalbergia, sp.; climbing shrub; pod 2-3 inches long. Ayer Hitam. (C. 1437).
Pterocarpus indicus, Willd.; is the principal shade tree planted in Penang. Not wild. (C.).
Pongamia glabra, Vent.; tree 30-40 feet. Not uncommon along the Coast. (C. 382).
Derris scandens, Benth.; climbing shrub. Common on the banks of tidal rivers. (C. 279).
Derris uliginosa, Benth.; climbing shrub. (C. 918).
Derris elliptica, Benth. "Akar tubah"; cultivated; root used as an insecticide and for poisoning fish. Not seen in a wild state. (C. 1438).
Derris thyrsoiflora, Benth.; robust climber. (C. 248, 802).
Derris, sp.; large climbing shrub, not referable to any of the above. Waterfall. (C.)
Cæsalpinia Bonducella, Flem.; straggling shrub. Tanjong Bunga. (C. 99).
Cæsalpinia sepiaria, Roxb.; climbing prickly shrub. Top of Government Hill. (C. 385).
Cæsalpinia digyna, Rottb. Tanjong Bunga. (C. 448).
Peltophorum ferrugineum, Benth.; tree 25-30 feet; corolla yellow. A coast plant. (C. 370).
Cassia occidentalis, L.; dwarf undershrub. common. (C. 131).
Cassia sophora, L.; not identified.
Cassia siamea, Lam.; small tree. (C. 92).
Cassia mimosoides, L.; perennial 6-18 in.; corolla yellow. common on Government Hill. (C. 820).

Cassia Kleinii, W. & A.; shrubby 2 feet. Government Hill. (C.).

Cassia javanica, L.; tree 30-40 feet. Waterfall. (C. 1576).

Cynometra polyandra, Roxb. is quoted from Penang and Malacca in the Flora of British India without any collector's name. Wallich and Roxburgh only got it in Khasiya and Silhet.

Cynometra cauliflora, L. "Nam-nam"; is cultivated; not seen wild. (C.)

Sindora Wallichii, Benth.; large tree; timber much valued. Not common. (C. 430).

Dialium Maingayii, Baker. "Kranji"; large tree, fruit sometimes eaten. (C. 440).

Saraca triandra, Baker; small tree. Government Hill, not common. (C. 163-647).

Bauhinia tomentosa, L. Penang, in the Fl. Brit. Ind. with no collector's name. It is certainly not wild in the peninsula.

Bauhinia cornifolia, Baker; climbing shrub. Government Hill. (C. 295).

Bauhinia emarginata, Jack; collected by Porter.

Bauhinia integrifolia, Roxb.; large woody climber; far more common than any of the other sps. (C. 300-500).

Bauhinia bidentata, Jack; petals orange; handsome plant when in flower. Government Hill. (C. 136).

Bauhinia, sp. near *bidentata*. Top of Government Hill. (C. 488).

Bauhinia elongata, Korth. Not identified. Collected by Porter. (Wallich 5782).

Bauhinia glauca, Wall. Not identified.

Bauhinia ferruginea, Roxb. var. *Griffithiana*. (C. 784). var. *excelsa*. (C. 211).

Bauhinia purpurea, L. is cultivated, not wild I think. (C.).

Bauhinia lucida, Wall. is only known from leaf specimens of a plant collected at Calcutta said to have come from Penang. (No. 5779b).

Bauhinia, sps. undetermined. (C. 801, 1541).

Neptunia oleracea, Lour.; floating herb; fl. yellow. Not common. (C.).

Entada scandens, Benth.; large climber. Government Hill. (C. 115).

98 A CATALOGUE OF THE FLOWERING PLANTS AND FERNS, &C.

Adenanthera pavonina, L. is often found growing near villages, but never in the forest. (C.).

Parkia biglandulosa, W. & A. is commonly cultivated. (C.).

Leucæna glauca, Benth.; large shrub. Waterfall. (C. 49).

Mimosa pudica, L. is one of the pests of the island. (C.).

Acacia Farnesiana, Willd. Not uncommon, but a doubtful native (C.).

Acacia pernata, Willd. var. *pluricapitata*; not seen.

Albizzia lebbek, Benth.; tall tree. Telok Bahang. (C. 206).

Albizzia myriophylla, Benth.; small tree. (C. 788-1702).

Albizzia, sp.; tree 30-40 feet; pod $6 \times 2\frac{1}{2}$ in. Waterfall. (C. 1921).

Calliandra umbrosa, Benth.; quoted without authority for Penang in Flora of British India was really collected in Silhet.

Pithecolobium dulce, Benth.; largely used as a fence plant. (C.)

Pithecolobium buhalinum, Benth. Not seen.

Pithecolobium microcarpum, Benth.; small tree. (C. 1093).

Pithecolobium fasciculatum, Benth.; small tree, rare. (C. 720).

Pithecolobium lobatum, Benth. "Jering"; tree 20-30 feet. (C. 105-711).

Pithecolobium contortum, Mart.; shrub 4-10 feet. (C. 19-264),

Pithecolobium clypcaria, Benth.; small tree. Ayer Hitam. (C. 209-189).

Pithecolobium angulatum, Benth.: spreading tree. (C. 489).

ROSACEÆ.

Parinarium costatum, Bl. "Poko Obie;" tree 30-40 ft. (C. 259).

Parinarium oblongifolium, Hook. f. ?; tree about 40 ft.; drupe $1\frac{1}{4}$ - $1\frac{1}{2}$ in. (C. 2163).

Parinarium asperulum, Miq.; tree 30-40 ft.; petals white (C. 203).

Parinarium nitidum, Hook. fil.; small tree 20-25 ft.; drupe round the size of a large pea. (C. 147, 853).

Parinarium Griffithianum, Benth.; large tree; drupe 1 in.

long of a dark colour. Muka Head, not common. (C. 1514, 2416).

Parastemon urophyllum, A. D. C.; small tree, branches slender. (C. 1149, 2480).

Prunus martabanica, Hook. f.; tree 40-50 ft. Government Hill. (C. 1512).

Pygeum lanceolatum. Hook. f.; tree 20-30 ft.; branches slender, drooping. (C. 216. 735).

Pygeum parviflorum, Teys. and Binnend.; tree 20-30 ft.; fruit black. Government Hill &c., common. (C. 162).

Pygeum, sp., tree 30-40 ft.; fruit $\frac{3}{4}$ in. Nalm. (C. 938).

Rubus glomeratus, Bl.; common in all parts of the Island. (C. 74).

Rubus moluccanus, L. not seen.

Fragaria indica, Andr.; does not occur other than as a cultivated plant. (C.)

SAXIFRAGACEÆ.

Polyosma integrifolia, Bl.; tree 30-40 ft. Penara Bukit. (C. 1081).

Polyosma mutabilis, Bl.; tree 20-30 ft. Not uncommon. (C. 377, 758).

Polyosma, sp.; tree larger in all its parts than either of the preceding; may be a form of *P. integrifolia*. (C. 1165).

CRASSULACEÆ.

Bryophyllum calycinum, Salisb.; herb 2-3 feet. Common. (C.)

HAMAMELIDEÆ.

Maingaya malayana, Oliv.; tree about 25 feet tall with a stem 6-9 inches diameter. Government Hill. (C. 659).

RHIZOPHOREÆ.

Rhizophora conjugata, L.; tree 20-40 feet, largely used for firewood. Common in tidal swamps. (C. 516).

Brugueria eriopetala, W. & A.; tree, not uncommon. (C. 514).

Carallia integerrima, D. C.; tree 30 feet. Government Hill 1000 2000 feet, not common. (C. 1078).

Pellacalyx axillaris, Korth.; medium-sized tree. (C. 937).

Pellacalyx Saccardianus, Scort.; small tree. Penara Bukit. (C. 1031).

Gynotroches axillaris, Miq; tree 25-30 feet, very common. (C. 324, 865).

Anisophyllea zeylanica, Benth. var.; tree 25-30 feet; branches slender, drooping. West Hill. (C. 746).

Anisophyllea grandifolia, Henslow; tree 30-40 feet. Government Hill, not uncommon. (C. 521).

Anisophyllea, sp.; tree 30-40 feet; drupe globular, 2-2½ inches. (This may be *A. Griffithii*). Waterfall. (C. 148, 1511).

COMBRETACEÆ.

Calycopteris floribunda, Lamk.; large shrub. Waterfall (C. 184).

Combretum squamosum, Roxb.; large scandent shrub. (C. 258 1077).

Combretum chinense, Roxb. Not identified. Collected by Porter.

Combretum acuminatum, Roxb. Large climber, petals yellowish white. Balik Pulau. (C. 2474).

Quisqualis densiflora, Wall.; scandent shrub. (C. 190, 889).

MYRTACEÆ.

Tristania Whitiana, Griff. Not seen.

Tristania Maingavii, Duthie; "Palawan" tree 20-40 feet; bark smooth, grey. Government Hill, &c., common. (C. 238).

Rhodamnia trinervia, Bl.; "Monkoyan" tree 25 feet or more. Common. (C. 51).

Rhodamnia? sp.; tree 40 feet. Penara Bukit 1000 feet. (C. 1446).

Rhodomyrtus tomentosa, Wight; "Kamunting"; shrub 4-10 feet; abundant in places, fruit eaten. (C. 200).

Decaspermum paniculatum, Kurz; small tree 15-20 feet; not uncommon. (C. 33, 177).

Eugenia malaccensis, L. is cultivated. (C).

Eugenia Jambos, L. is not wild. (C).

Eugenia javanica, Lamk.;? small tree, not common. (C. 1114).

Eugenia grandis, Wight; "Kiean batu"; large tree, valuable timber. Common near the coast. (C. 751).

Eugenia filiformis, Wall; small tree; branches slender, drooping. Government Hill. (C. 744, 1443).

Eugenia ramo issima, Wall.; tree. Pulau Betong. (C. 973).

Eugenia inophylla, Rox. (C. 974 ?)

Eugenia Thumra, Rox.; medium-sized tree. Waterfall. (C. 2410).

Eugenia acuminatissima, Kurz; tree. Government Hill. (C. 179).

Eugenia claviflora, Rox. in Flora of British India, is queried for Penang by Wallich in his catalogue which is apparently the only authority for its being a Penang plant.

Eugenia leptantha, Wight; medium-sized tree; fruit black, edible. (C. 697, 749).

Eugenia zeylanica, Wight; tree 20-30 feet, common. (C. 317, 511, 2245).

Eugenia grata, Wall. "Gelam Tikus"; small tree; bark used for tanning purposes. (C. 756).

Eugenia penangiana, Duthie; medium-sized tree. Government Hill 500 feet. (C. 193).

Eugenia skiophila, Duthie. Government Hill in deep shaded ravines. Maingay. Not met with since.

Eugenia chlorantha, Duthie.; tree. West Hill 1,500 ft. (C. 180, 1448).

Eugenia lineata, Bl.; small tree. Muka Head &c., (C. 728 976 ? 2417).

Eugenia bracteolata, Wight; small tree. Government Hill. (C. 247, 1089).

Eugenia valdivenosa, Duthie; small tree. Government Hill at 2,000-2,500 feet. (C. 55).

Eugenia expansa, Wall. Not identified. (Wall. cat. 3,567) collected by himself.

Eugenia subdecussata, Wall.; small tree. West Hill 2,500 ft. (C. 864).

Eugenia oblata, Wall?; large branching tree. Government Hill. (C. 212).

Eugenia lævicaulis, Duthie; small tree. (C. 2246).

Eugenia brachiata, Rox. "Krean batu;" tree 25-30 feet. Waterfall, common. (C. 32).

Eugenia verecunda, Wall?; tree. Government Hill. (C. 654).

Eugenia, sps. undetermined in Herb. Penang. (C. 194, 653, 666, 759, 1090, 1152, 1442, 1604, 2224, 2228).

Pseudeugenia perakensis, Scort.; large shrub; not uncommon on Government Hill. (C. 461, 886).

Barringtonia speciosa, Forst.; medium-sized tree. (C. 909).

Barringtonia acutangula, Gaertn.; tree 25-30 ft.; not common. (C. 397).

MELASTOMACEÆ.

Melastoma malabathricum, L.; shrub 3-6 ft. (C. 71).

Melastoma imbricatum, Wall.; not identified. Wall. 4047.

Melastoma molle, Wall.; shrub 3-5 feet. Government Hill 1000-2000 ft., not common. (C. 683).

Melastoma appressum, Wall.; Penang Hill, Hullett.

Melastoma longifolium, Naud. Government Hill Road to Penara Bukit at 1800 feet. (C. 2789).

Allomorpha exigua, Bl.; ? shrub 1-3 feet, common. (C. 73,399).

Sonerila erecta, Jack; stem 3-12 inches; common from 1000-2000 feet (C. 1238).

Sonerila affinis, Arn.; stem 3-6 inches. Government Hill 2000-2500 feet; not common. (C.)

Sonerila moluccana, Roxb.; stem 1-2 inches. Not uncommon in damp shady places from sea-level up to 2000 feet. (C. 412).

Marumia nemorosa, Bl.; large climbing shrub; flower large, pink. Penara Bukit, rare. (C. 1008).

Dissochæta annulata, Houk. fil. West Hill. (C. 740).

Dissochæta punctulata, Hook. fil. Said to have been collected here by Walker. Wants confirmation.

Dissochæta bractenta, Bl.; large twining shrub. Government Hill 500-1000 feet. (C. 2298, 80?)

Dissochæta pallida, Bl. Government Hill. (C. 2297).

Dissochæta intermedia, Bl. Not identified. Wallich 4052.

Dissochæta celebica, Bl. Not identified. Wallich No. 4050 4052 pars. These two numbers are labelled Singapore and

Penang 1822 and were probably obtained only in the former locality.

Dissochæta gracilis, Bl.; slender twining shrub. (C. 398).

Dissochæta? sp. undetermined. (C. 1078).

Anplectrum glaucum, Triana. Not seen. Collected by McNair, Jack, Wallich, etc.

Anplectrum pallens, Bl.; twining shrub, branches slender. Government Hill 1000-2500 feet. (C. 471).

Anplectrum divaricatum, Triana. Not identified. Wallich (4051), and Walker.

Anplectrum annulatum, Triana. Not identified. Wallich Cat. No. 4056 obtained by Porter.

Anplectrum polyanthum, C. B. C. Waterfall (C. 61).

Medinilla rubicunda, Bl. Not seen. The plant numbered in Wallich's catalogue 4086 was collected at Cape Rachado in Sungei Ujong not in Penang.

Medinilla speciosa, Bl.; shrub 4-6 feet., panicle long, pink, pendulous. (C. 874).

Pogonanthera pulverulenta Bl.; shrub 6-10 feet. Batu Ferengy. (C. 1880).

Pachycentria macrorrhiza Becc., compact growing shrub 2-3 feet; generally on rocks or in the forks of large trees. (C. 347).

Pachycentria, sp.; straggling shrub. Damp ravines at 2,000 feet. (C. 2225).

Astronia smilacifolia, Triana; small tree 25 feet. Top of Government Hill. (C. 743).

Pternandra carulescens, Jack; tree 25-30 feet. (C. 879).

Var. *fackiana*. (C. 2220).

Pternandra capitellata, Jack. "Kulit nipis"; tree 20-30 feet. Government Hill, &c., common. (C. 67).

Pternandra paniculata, Benth. Not identified.

Kibessia echinata, Cogn., tree, branches rather slender. (C. 270).

Kibessia pubescens, Dcne. Not identified.

Kibessia, sp.; small tree apparently different to either of the preceding. Nalm. (C. 953).

Memecylon myrsinoides, Bl.; small compact growing tree; fl. blue. (C. 100-2219).

Memecylon heteropleurum, Bl.; small tree. West Hill. (C. 457-814).

Memecylon microstomum, C. B. C.; small tree. Government Hill 2,000 feet, not common. (C. 766).

Memecylon cœruleum, Jack; small erect shrub; common along the coast, not seen elsewhere. (C. 54).

Memecylon amplexicaule, Roxb.; shrub 8-10 feet tall Government Hill 2,000 feet. (C. 695).

Memecylon garcinoides, Bl.; small tree. West Hill. (C. 816).

Memecylon acuminatum, Smith, var. *flavescens*; tree 25-30 feet. West Hill 1,000 feet. (C. 815).

Memecylon edule, Roxb.; tree 25 feet high or sometimes much less, fl. blue. Common along the coast. (C. 723).

Memecylon oligoneurum, Bl.? small tree. Ayer Hitam. (C. 1065).

Memecylon, sp.; erect shrub. West Hill. (C. 1773).

Memecylon, sp.; tree 20-30 ft. Government Hill 2000 ft. (C. 2243).

LYTHRACEÆ.

Crypteronia pubescens, Bl. "Bequoie;" tree 20-40 ft. very common at the Waterfall. (C. 110).

Crypteronia glabra, Bl.; large tree 40-50 ft. Government Hill 2500 ft., rare. (C. 739).

Crypteronia Griffithii, C. B. C.; tree 25-30 ft. (C. 1739, 660?)

Duabanga sonneratioides, Ham.; large tree. Balik Pulau, not common. (C. 678).

Sonneratia acida, L.; small tree. Tidal swamps. (C. 1084).

Sonneratia Griffithii, Kurz; small tree. (C. 1108).

ONAGRACEÆ.

Fussia repens, L. Common in ditches. (C. 1936).

Fussia suffruticosa, L. Rice fields, and generally in damp places. (C. 123, 1972?)

Ludwigia prostrata, Roxb. is not uncommon. (C. 57).

SAMYDACEÆ.

Casearia glomerata, Rox. ; small tree. Pulau Betong. (C. 982, 2466?)

Casearia macrocarpa, C. B. C. ; tree 20-30 ft. fl. small white. (C. 220, 960).

Casearia albicans, Wall. not seen. (Wallich no 7197).

Casearia, sp. ; small tree ; fl. minute, white. Government Hill 300-500 ft. (C. 2143).

Casearia sp. ; small tree. West Hill. (C. 1574).

Homalium longifolium, Benth. ; tree 30-40 feet ; conspicuous when in flower. (C. 301, 929).

Homalium sp. ; tree 30-40 feet ; leaves 6-8 × 3-4 inches. Waterfall, only one tree seen. (C. 1592).

Samyda esculenta, Roxb. ? (C. 1019).

PASSIFLOREÆ.

Passiflora fœtida, L. is naturalized and much more common than any other plant of this order. (C. 1236)

Modecca singaporeiana, Mast. ; twining shrub. Penara Bukit, rare. (C. 869).

Modecca nicobarica, Kurz. ; very slender twiner ; fruit 1½-2 inches, bright red. Not uncommon. (C. 1521).

CUCURBITACEÆ.

Hodgsonia heteroclita, Hook. fil. Not seen. Collected by Porter (Wall cat 6685) and Phillips.

Trichosanthes palmata, Roxb. is not uncommon. (C.)

Gynopetalum quinquelobium, Miq. Not identified. Collected by Porter.

Mormodica cochinchinensis, Spreng. (C. 890).

Cerasiocrarpum? penangense. C. B. C. not identified. A doubtful plant collected by Wallich. (No. 6704).

BEGONIACEÆ.

Begonia Evansiana, Andr. In Andrews' Botanist's Repository where this plant was first described it is said that Mr. Evan's collector found it growing in the clefts of rocks in the Island of Pulo Penang in 1808. It has never been since met with here and could hardly have been overlooked. It is a Chinese plant.

Begonia guttata, Wall. Not identified. (Wall Cat 3671a).

Begonia prolifera, A. D. C. Erroneously attributed to Penang on the strength of a plant collected by Finlayson which really came from one of the Siamese islands.

Begonia sinuata, Wall. Government Hill, &c., common. (C. 390, 481).

Begonia malabarica, Lam. ? (C. 1262, 1738).

Begonia, sp. Pulau Betong, rare. (C. 1023).

UMBELLIFERÆ.

Hydrocotyle asiatica, L. "Pegaga;" a common herb; leaves used for curries &c. (C.)

Hydrocotyle rotundifolia, Roxb. Government Hill. (C. 1752).

ARALIACEÆ.

Aralia, sp. aff. *A. Thompsonii*, Seem.; small prickly tree 10-15 ft. Government Hill. (C. 462).

Heptapleurum venulosum, Seem.; shrub. (C. 972).

Heptapleurum subulatum. Seem.; shrub. (C. 971, 1246).

Heptapleurum cephalotes, C. B. C. ? tree 20-30 ft, not large, Government Hill. (C. 837).

Heptapleurum heterophyllum, Seem. Moniot's Road. (C. 2301).

Heptapleurum, sp. collected at Pulau Betong (C. 1950).

Trevesia palmata, Vis. var.; said to have been collected in Penang by Porter, was probably from Province Wellesley. (C.)

Arthrophyllum diversifolium, Bl.; tree 20-30 ft. not large. Waterfall. (C. 781).

Arthrophyllum pinnatum, C. B. C.; shrub. West Hill 2000 ft. (C. 334).

CORNACEÆ.

Marlea begoniæfolia, Rox.; small tree. Pulau Betong. (C. 940).

Marlea nobilis, C. B. C, Government Hill. (C. 1505).

Mastixia Maingayii, C. B. C.; tall tree. Government Hill. (C. 1564).

Mastixia, sp.; tree 30-40 ft. Pulau Betong. (C. 919).

CAPRIFOLIACEÆ.

Viburnum sambucinum, Reinw; large shrub, sometimes a small tree; flowers white, sweet scented. Top of Government Hill, common. (C. 278).

RUBIACEÆ.

Sarcocophalus Funghuhnii, Miq.; large tree. Government Hill 500 ft. (C. 303, 2751.?)

Adina rubescens, Hemsley var. *acuminata*; medium-sized tree. Government Hill (C. 369).

Nauclea peduncularis, Wall.; tree 25-30 ft. Sungei Penang, not common (C. 1056).

Uncaria sclerophylla, Roxb.; large climbing shrub. Govern-Hill. (C. 6, 185).

Uncaria attenuata, Korth. Government Hill. (C. 135).

Uncaria canescens, Korth. Government Hill. (C. 331).

Uncaria pteropoda, Miq. Government Hill. (C. 332, 1247?)

Uncaria ovalifolium, Roxb. not identified collected by George Porter.

Uncaria dasyneura, Korth. Government Hill. (C. 1070),

Uncaria gambier, Rox. is not wild in Penang.

Uncaria lanosa, Wall. Waterfall. (C. 917).

Uncaria acida, Roxb. is probably one of the above but the description is too meagre to guess which and no type is known to exist.

Coptosapelta flavescens, Korth.; climbing shrub; flower open white, gradually turning yellow. (C. 312, 1029).

Dentella repens, Forst.; prostrate herb; fl. white, $\frac{1}{4}$ in. (C. 1850).

Grœnia Jackii, W. and A.; erect shrub, 6-10 ft.; a common plant near the Coast. (C. 106).

Argostemma pictum, Wall.; small herb; common on damp rocks at 1,000-2,000 feet. (C. 408).

Argostemma verticillatum, Wall. An error. Wallich's specimens came from Nepal, not Penang as stated in Flora of British India

- Argostemma humile*, Wall. Not met with. It was collected by Jack.
- Argostemma unifolium*, Benn. Damp rocks, common. (C. 990).
- Argostemma elatostemma*, Hook. fil. West Hill. (C. 962).
- Argostemma*, sps. undetermined. (C. 342, 955).
- Hedyotis macrophylla*, Wall. Not identified. (Wallich No. 842).
- Hedyotis capitellata*, Wall; climber; the most common of the genus. (C. 23).
- Hedyotis mollis*, Wall. Penara Bukit. (C. 1096).
- Hedyotis vestita*, Br. Not seen.
- Hedyotis glabra*, Br. Government Hill. (C. 39).
- Hedyotis pinifolia*, Wall. Not seen.
- Hedyotis tenelliflora*, Bl. (C. 2236).
- Hedyotis hispida*, Retz. Waterfall. (C. 1847).
- Hedyotis congesta*, Br. Penang Hill (C. 2846).
- Hedyotis scabra*, Wall. Bagan Jennal. (C.)
- Hedyotis*, sp. probably new. (C. 977).
- Oldenlandia corymbosa*, L.; a common weed. (C. 1849, 1988).
- Oldenlandia diffusa*, Roxb. Waterfall; not very common. (C. 1846, 1845).
- Oldenlandia Heynii*, Br. Common. (C. 1848).
- Oldenlandia trinervia*, Retz.; fl. minute, white. (C. 1844).
- Ophiorrhiza munges*, L.; small herb. West Hill. (C. 979).
- Ophiorrhiza discolor*, Bl. Pulau Betong. (C. 1761),
- Ophiorrhiza tomentosa*, Jack. West Hill. (C. 907).
- Mussænda glabra*, Wall.; climbing shrub; common. (C. 116).
- Mussænda villosa*, Wall. (C. 289, 1934)?
- Mussænda cordifolia*, Wall. Not identified. (Wallich 6260). Imperfectly known.
- Lucinæa morinda*, D. C. Not seen here. "Penang and Singapore" No. 3487 Wallich.
- Trisciadia truncata*, Hook. fil. Not identified. On the Hill, Wallich in 1822. Not collected since.
- Adenosacme longifolia*, Wall.; shrubby, 1-2 feet; fruit white. (C. 845).

Aulacodiscus premnoides, Hook. fil.; small tree. Penara Bukit, not common. (C. 987).

Urophyllum Griffithianum, Wight; small tree. Government Hill, rare. (C. 1189, 2294).?

Urophyllum glabrum, Wall.; common at 1000-2000 feet on Government Hill. (C. 260, 1759).

Urophyllum streptopodium, Wall.; small tree. West Hill. (C. 1770).

Urophyllum Blumeanum, Wight; Government Hill. (C. 261).

Urophyllum villosum, Wall.; large shrub; common. (C. 178).

Urophyllum? sp.; shrub, smaller in all its parts than either of the preceding. (C.)

Webera odorata, Roxb. Not identified. Collected by Wallich.

Webera fragrans, Bl.; small tree. (C. 480).

Webera, sp. Near *Maingayii*, Hook. fil.; large shrub. (C. 1060).

Webera mollis, Wall.? small tree. West Hill. (C. 745).

Webera stellulata, Hook. fil. Not identified. Wallich, Griffith.

Webera longifolia, Hook. fil. shrub 6-10 feet; cymes 6-10 inches, pendulous; flower white. Waterfall. (C. 947, 1144).

Webera Wallichii, Hook. fil. Not identified. Wallich 8401. I.

Webera, sp.; large shrub. Waterfall. (C. 2217).

Anomanthodia auriculata, Hook. fil.; small tree. Moniot's Road, not common. (G. 1075).

Randia, sp. near *longiflora*, Lam.; slender tall tree. (C. 818).

Randia densiflora, Benth.; small tree; common. (C. 128-923).

Randia, sp. near *densiflora*, Benth. (C. 796).

Randia anisophylla, Lamk. "Randa"; tree 25-30 feet; fl. white fugacious. The commonest of all the *Randias*. (C. 164).

Randia exaltata, Griff.; spreading tree; corolla white with purple markings; fruit the size of a small orange, black. (C. 793).

Randia macrophylla, Br.; shrub; rare in Penang, (C. 966).

Randia, sp.; large climbing shrub; fl. yellowish white; fruit 1 in., globular. Government Hill. (C. 927).

Randia, sp. small erect spinous shrub. Waterfall. (C. 1498).

Gardenia carinata, Wall.; tree 34-40 feet, spreading. (C. 525).

110 A CATALOGUE OF THE FLOWERING PLANTS AND FERNS, &C.

Gardenia tubifera, Wall. ; tree 20-30 feet, compact, leafy. Government Hill. (C. 686).

Petunga longifolia, D C. Not identified. Penang. Phillips, Wallich.

Petunga, sp. ; small slender tree 15-20 feet. Government Hill. (C. 726).

Di lospora? sp ; small tree. Penara Bukit. (C. 1449).

Timonius jambosella, Thw. ; small close-growing tree ; not uncommon. (C. 95-263).

Timonius Rumphii, D. C. Government Hill. (C. 2844).

Canthium didymum, Roxb. ; tree 25 feet. Waterfall. (C. 219).

Canthium glabrum, Bl. ; small tree. Penara Bukit. (C. 938).

Canthium confertum, Korth. Government Hill. (C. 1694-1119).

Canthium horridum, Bl. Not seen. Cantley, Wallich.

Canthium, sp. ; probably *oliganthum*, Miq. (C. 1756).

Ixora Brunonis, Wall. Not identified. Wallich. Phillips.

Ixora pendula, Jack ; large shrub. Waterfall (C. 2240).

Ixora grandifolia, Zoll and Moritz. Not seen.

Ixora villosa, Roxb. Not identified. Wallich.

Ixora coccinea, L. ; shrub ; cultivated, wild as an escape. (C.)

Ixora amœna, Wall. Government Hill. (C. 79).

Ixora fu'gens, Roxb. ; tall shrub ; common (C. 479, 1730? 2257? 2255?)

Ixora congesta, Roxb. ; shrub 6-10 ft. (C. 667, 729).

Ixora, sp. near *congesta* ; shrub, not exceeding 12 in. (C. 400).

Ixora opaca, Br. ; small tree. Muka Head. (C. 719).

Ixora lucida, Br. Not identified. Wallich 6135, very little known.

Ixora nigricans, Br. ; large shrub. Pulau Betong. (C. 986, 2451).

Ixora, sp. ; small tree 20 ft. (C. 1112).

Pavetta indica, L. Pulau Betong. (C. 908).

Coffea? sp. ; small tree, fl. not seen. (C. 757, 1774).

Morinda tinctoria, Roxb. ; tree 15-25 feet ; very common in places. (C. 679).

- Morinda umbellata*, L.; large climbing shrub. (C. 149).
Prismatomeris albidiflora, Thw.; shrub 3-5 ft.; fl. white; common near the Coast. (C. 330).
Gynochthoides sublancoolata, Miq. Wallich No. 8385 "Penang" is the only authority for this here and the specimens have neither flower nor fruit, so are doubtful.
Psychotria malayana, Jack. Not seen.
Psychotria polycarpa, Miq.; climber. Penara Bukit. (C. 1095).
Psychotria sarmentosa, Bl.; common Government Hill. (C. 951).
Psychotria morindæfolia, Wall.; climber. (C. 478).
Psychotria stipulacea, Wall.; erect shrub, 4-6 ft. collected by Jack. (C. 732).
Psychotria Jackii, Hook. fil.; not seen in Penang, common at Pangkor. (C.)
Psychotria tortilis, Bl.; 6-18 in.; stem fleshy. Not uncommon. (C. 405, 1269).
Psychotria angulata, Korth.; erect shrub, 2-4 feet. (C. 726, 14)?
Psychotria Helferiana, Kurz. not met with. Recorded from "Penang Wallich" in Flor. Brit. Ind. but Wallich's No. 8368 was collected in Singapore where this plant is common.
Psychotria perangensis, Hook. fil., Not identified, insufficiently known. Top of Government Hill, Maingay.
Psychotria sp.; sarmentose shrub, berries white. (C. 2215).
Psychotria? sps. undetermined. (C. 751, 981, 160).
Chasalia curviflora, Thw.; shrub 2-3 feet; stem rather fleshy; common C. 414, 680).
Var longifolia. Top of Government Hill.
Geophila reniformis, Dcn.; creeping herb; corolla ivory white; fruit bright red; common. (C. 1930).
Lasianthus cynnocarpus, Jack; shrub 4-6 feet; fruit blue $\frac{1}{4}$ - $\frac{1}{2}$ inch. Not uncommon. (C. 650, 1116).
Lasianthus sp. near *Wallichii*, Wight; shrub 6-7 feet. Government Hill in damp ravines. (C. 284).
Lasianthus adpressus, Hook. fil. Not identified. Wallich 8443, 8442 in part.

Lasianthus densifolius, Miq. Not identified.

Lasianthus crinitus, Hook. L. Not identified. Collected by Wallich, locality doubtful.

Lasianthus Kurzii, Hook. fil. Not identified; locality doubtful more likely from Martaban.

Lasianthus chinensis, Benth.; tall shrub. West Hill 2000 feet. (C. 1594).

Lasianthus, sps. undetermined. (C. 760, 785).

Hydnophytum montanum, Bl. (C. 2164).

Pædiria fœtida, L.; slender twining shrub. (C. 24).

Spermacoce hispida, L.; prostrate herb; flower pink; common in open grassy places. (C. 1986, 1987).

COMPOSITÆ.

Vernonia chinensis, Less.; shrubby 1-2 feet; flower blue; abundant in orchards and waste places. (C. 931).

Vernonia arborea, Ham.; tree 25-30 feet. (C. 2-926).

Vernonia cinerea, Less.; annual herb, 1-3 feet. Waterfall. (C. 1698).

Elephantopus scaber, L.; dwarf rigid herb; abundant in coco-nut plantations. (C. 130).

Adenostemma viscosa, Forst; erect annual herb. (C. 985).

Ageratum conyzoides, L.; fl. blue or white, abundant. (C.).

Mikania scandens, Willd.; slender twining herb. Penara Bukit. (C. 1042).

Microglossa volubilis, D.C. Not seen. Collected by Jack and Wallich.

Blumea chinensis, D.C. Not seen.

Blumea balsamifera, D.C.; 6-10 feet. Tanjong Bunga. (C. 29).

Blumea lacera, D.C.; annual, 1-3 feet. Waterfall. (C. 1699).

Laggera flava, Benth. Not identified. Quoted as occurring in Penang in Flora British India but with no collector's name.

Pluchea indica, Less.; "Bluntas" shrubby 4-5 feet; common in marshy places. (C. 509).

Sphæranthus africanus, L.; annual herb 1-2 feet; not uncommon in rice fields. (C. 1942).

Eclipta alba, Hassk.; much branched, sometimes erect. (C. 1057).

Wedelia biflora, D.C.; climbing herb, fl. yellow. (C. 1568-40).

Spilanthes acmella, L.; common annual herb. (C. 1929).

Synadrella nodiflora, Gærtn.; branching herb; common in open places. (C. 913).

Bidens pilosa, L.; one of the commonest weeds. (C. 1669).

Gynura sarmentosa, D.C.; twining herb. Penara Bukit. (C. 1205).

Gynura Pseudo-china, D.C. Government Hill. (C. 2995).

Emilia sonchifolia, D.C.; annual herb. Penara Bukit. (C. 37).

Launæa pinnatifida, Cass. Government Hill.

GOODENOVIEÆ.

Scævola Kænigiï, Vahl.; shrub 6-10 feet; branches stout, soft. Common along the coast. (C. 320).

CAMPANULACEÆ.

Pentaphragma begoniæfolium, Wall.; herb 6-12 in.; common 1,000-2,000 feet. (C. 329).

ERFIACEÆ.

Rhododendron Teysmanii, Miq.; shrub 2-4 feet; flowers yellow. Government Hill 2,000 feet. (C. 530).

PLUMBAGINEÆ.

Plumbago zeylanica, L.; is not uncommon, but not indigenous. (C.).

Plumbago rosea, L.; found in one or two places, not indigenous. (C.).

MYRSINEÆ.

Mæsa ramentacea A. D.C.; large rambling shrub; never a tree in Penang. Common. (C. 48-251).

Mæsa ramentacea var ovata, not identified.

Myrsine capitellata, Wall.; small tree. Government Hill. (C. 1013).

Embelia ribes, Burm.; large climbing shrub, common. (C. 18).

Embelia ribes var. pinangiana, Oliv.; fruit larger than the type, of a reddish colour. (C. 306).

- Embelia coriacea*, A. D.C.; climber, rare. (C. 1047).
Embelia careacens, Jack. Not identified. Only collected by Jack.
Embelia Limpanii, Schiff. Government Hill. (C. 305).
Labisia pothoina, Lindl.; shrubby 6-10 inches, common. (C. 1192).
Ardisia lancrolata, Rox. Not identified. Roxburgh.
Ardisia colorata, Rox.; small tree. Penara Bukit. (C. 1052).
Ardisia divergens, Rox. Not identified. Jack and Maingay.
Ardisia tuberculata, Wall. var. *Ophirensis*; small shrub. West Hill 2500 feet. (C. 2249).
Ardisia porosa, Clarke; straggling shrub. West Hill 2000 feet. (C. 1769).
Ardisia crenata, Roxb.; shrub 2-4 feet; very common. (C. 337).
Ardisia villosa, Roxb.; shrub 1-2 feet. Government Hill. 2000 feet. (C. 597).
Ardisia oxyphylla, Wall.; small tree. Government Hill. (C. 668).
Ardisia humilis, Vahl; small tree 20-25 feet. (C. 217).
Ardisia, sp. in herb. Penang; leaves $10 \times 4\frac{1}{2}$ inches; berry $\frac{1}{4}$ inch. (C. 2755).
Pimelandra Wallichii, A. D. C.; small tree 10-15 feet. Government Hill, not uncommon. (C. 1156).
Aegiceras majus, Gaertn.; tree 30 feet; common in swamps near the coast. (C. 62, 513).

SAPOTACEÆ.

- Chrysophyllum* sp.; tree 25-30 feet. Batu Feringgi. (C. 1072).
Sideroxylon malaccense, Clarke; tree 40 feet; fruit 1-1 $\frac{1}{2}$ inches, rusty tomentose. (C. 1452).
Sideroxylon ferruginum, Hk. & A.; tree 40 feet, coast. (C. 273, 703).
Sideroxylon, sp. near *firmum*, Burck; small tree. West Hill 2,500 feet, rare. (C. 1575).
Sideroxylon, sp.; small tree. Batu Feringgi. (C. 2254).
Sideroxylon, sp.; small tree. Penara Bukit. (C. 1450).

Dichopsis gutta, Benth.; tree 50-60 feet. Government Hill, &c. (C. 780).

Bussia Mo'leyana, Clarke; tree 50 feet. Government Hill, not common. (C. 1451).

Payena Maingayii, Clarke; small tree. West Hill 2,000 feet. (C. 1565).

Payena maluccensis, Clarke. Telok Bahang. (C. 159).

Payena lucida, A. D.C. Not identified. (Wallich 4147).

Payena costata, tree. Waterfall (C. 1516)

Payena sp.; near *Leerii*, Hook. fil.; tree 50-60 feet. Government Hill. (C. 777).

EBENACEÆ.

Maba buxifolia, Pers.; low close growing bush. Telok Bahang, not common. (C. 708).

Maba merguensis, Hiern? large shrub. (C. 705).

Maba sp.; large tree. Muka Head. (C. 718).

Maba sp.; small tree. Penara Bukit. (C. 448).

Diospyros apiculata, Hiern; shrub 6-8 feet. Government Hill 1,000 feet, rare. (C. 1584).

Diospyros microphylla, Bedd.; small tree. Government Hill 300 feet. (C. 1513).

Diospyros sapotoides, Kurz; small tree. Telok Bahang. (C. 1564).

Diospyros decipiens, Clarke; small tree about 20-30 feet. Telok Bahang. (C. 1453-1454).

Diospyros sp.; near *oleifolia*, Wall.; tall tree. Government Hill 1,200 feet. (C. 1573)

Diospyros undulata, Wall.; tree medium-size, 40 feet or more. Telok Bahang, &c. (C. 1085).

Diospyros lucida, Wall.; tree 25-30 feet. Government Hill, near the chalet. (C. 250).

Diospyros oblonga Wall. "Koomoi"; tree 30-40 feet high. Sungei Penang. (C. 2487).

Diospyros venosa, Wall. Not identified.

Diospyros sp.; tree 20-30 feet; leaves 6-8 x 2½ in.; fruit conical ¾-1 in. Waterfall. (C. 2221).

Diospyros sp. near *ebenum* small tree. Batu Feringgi. (C. 1602).

Tabernaemontana malaccensis, Clarke; small shrub 2 feet Waterfall, not common. (C. 1775).

Parsonsia spiralis, Wall; twining shrub. Batu Feringig. (C. 450).

Parsonsia ovata, Wall.; Old road to Batu Feringgi. (C. 820).

Wrightia lævis, Wall.; tree 40-50 ft.; fl. greenish white, 1 in. in diam. Government Hill. (C. 1767).

Wrightia? sp.; small tree. Pulau Belong. (C. 848).

Strobhanthus Jackianus, Wall. "Bunga hantu"; shrub 3-4 feet; flower brick red, fugacious. (C. 188).

Strobhanthus dichotomus, D. C. not seen. Common in Singapore. The drawing labelled Penang in Herb. Kew (the only authority for its occurring here) was probably wrongly localised.

Urceola elastica, Roxb.; large climbing shrub. Government Hill. (C. 823).

Urceola lucida, Benth.; woody climber. Muka Head. (C. 661, 941).

Urceola brachysepala, Hook fil. Panti Achie. (C. 403).

Parameria glandulifera, Benth.; climbing shrub. Government Hill, rare. (C. 1142).

Parameria polyneura, Hook fil? Waterfall. (C. 1455, 1456).

Parameria densiflora, Oliv.; large climbing shrub; flower dull red. Government Hill. (C. 158).

Chonemorpha macrophylla, Don; large climber. (C. 832).

Aganosma marginata, Don; large shrub. (C. 664).

Aganosma? sp.? climbing shrub; flower white. Penara Bukit. (C. 2441).

Rhynchodia Wallichii, Benth.; flower white, fragrant. (C. 838).

Anodendron pauciflorum, Hook fil; climbing shrub. Government Hill. (C. 663, 1457).

Anodendron paniculatum, A. D. C.; large climbing shrub. (C. 1740, 2258).

Ichnocarpus ovatifolium, A.D.C. Tanjong Bunga. (C. 810).

Microchites polyantha, Miq.; climber, covering rocks and trees; roots emitted the whole length of the stem. (C. 850).

ASCLEPIADEÆ.

Antherandra acutifolia, Dcne. Not identified. Wallich 445^N).

Streptocaulon Wallichii, W. and A.; twining shrub; very common. (C. 4, 472, 2122).

Secamone emetica, Br.; slender twining shrub. (C. 1896, 225).

Secamone Finlaysonii, Wight? (C. 922).

Toxocarpus Griffithii, Dcne. Not identified. Collected by Porter.

Toxocarpus acuminatus, Benth.?; glabrous twining shrub; fl. white. Government Hill. (C. 244).

Calotropis gigantea, Br. I have seen only in gardens. (C.)

Asclepias curassavica, L. is met with rarely. (C. 344).

Cynanchum ovalifolium, Wight; twiner. Penara Bukit. (C. 871, 1270).

Cynanchum corymbosum, Wight. Not identified. 8231 F, Wallich.

Gougronema, sp.; West Hill. (C. 1545).

Pergularia puberula, Miq.; twining shrub, rare. (C. 932).

Tylophora hirsuta, Wight. Not identified.

Tylophora Wallichii, Hook. fil.; slender twiner. Pulau Betong, very rare. (C. 933).

Heterosiemma, sp.; twiner. Telok Bahang. (C. 1707).

Dischidia nummularia, Br.; slender herb; often completely covering trunks of trees or rocks. (C.)

Dischidia benghalensis, Colebr. Not common. West Hill at from 2000-2500 ft. (C. 2160).

Dischidia rhombifolia, Bl. Batu Feringgi. (C. 1897).

Hoya multiflora, Bl. The only evidence for this here is a drawing labelled Penang preserved at Kew.

Hoya parviflora, Wight. (C. 689).

Hoya latifolia, Don. Not seen.

Hoya obtusifolia, Wight. Not identified. "Penang? Wallich" (No 8167) probably from elsewhere.

Hoya coronaria, Bl. Telok Bahang. (C. 236).

Hoya diversifolia, Bl. Not uncommon on rocks close to the sea. (C. 928).

Diospyros sp. aff. *decipiens*, Clarke; small tree, 20 feet. Bukit Padre. (C. 1276).

Diospyros sp.; small tree. Moniot's Road. (C. 1693).

Diospyros sp.; small tree; leaves 10-12 inches; flower small, white. (C. 898, 1932).

STYRACACEÆ.

Symplocos ferruginea, Roxb.; tree 25 feet; common. (C. 206).

Symplocos sp. aff. *fasciculata*, Zoll.; tree 30-40 feet. Penara Bukit. (C. 1039).

Symplocos adenophylla, Wall; small tree; not uncommon. (C. 402).

Symplocos racemosa, Roxb.; tree 40-50 feet; flower white. Muka Head and Government Hill. (C. 715).

Symplocos cerasifolia, Wall.; tree 30-40 feet. Government Hill 1200 feet. (C. 1503).

Symplocos rubiginosa, Wall.; small tree. Penara Bukit and Government Hill. (C. 528, 1692).

Symplocos sp. near *rosea*, Benth.; small tree; fruit $\frac{3}{4}$ inch long. Penara Bukit, but one tree seen. (C. 1099).

Symplocos sp.; small tree. West Hill. (C. 325).

Symplocos sp.; tree. Muka Head. (C. 958).

Styrax serrulatum, Roxb.; tree 25-30 feet; branches slender, drooping; flower white. (C. 1181, 1538).

OLEACEÆ.

Jasminum pubescens var. *bracteata*, Rox. Not identified. Recorded from Penang, Wallich in Fl. Brit. Ind. Wallich's specimens (No. 2867) came from the Calcutta gardens, and were originally obtained from Bencoolen in Sumatra.

Jasminum Maingayii, Clarke. Government Hill. (C. 643).

Jasminum b farium, Wall. (C. 83)?

Jasminum scandens, Vahl. (C. 1920. 42)?

Jasminum subtriplinerve, Bl. (C. 643).

Jasminum sp. near *dispermum*, Wall. (C. 1227).

Linociera pauciflora, Clarke; tree 30 feet; not uncommon. (C. 277, 695).

Olea maritima, Wall.; small tree. Coast. (C. 222).

Olea dentata, Wall.; small tree. Telok Bahang. (C. 950).

Myxopyrum nervosum, Telok Bahang. (C. 2891).

APOCYNACÆ.

Willughbeia coriæa, Wall.; large climbing shrub. Waterfall. (C. 1500, 847).

Willughbeia sp.; large shrub. Penara Bukit. (C. 1121).

Chilocarpus Maingayii, Dyer? (C. 475).

Leuconotis eugenifolius, A. D. C. "Getah Gharu"; scandent shrub; not uncommon. (C. 1051).

Melodinus monogynus, Roxb. Not identified; It is said to have been collected by Maingay, otherwise only known from the Himalaya and Assam.

Melodinus orientalis, Bl.; large rambling shrub; fruit the size of a small orange. (C. 687).

Melodinus coriaceus, Oliv.; large climber reaching to the top of tall trees; fruit round, the size of an orange. (C. 1040).

Alyxia coriacea, Wall. Not identified. Collected by Wallich.

Alyxia pilosa, Miq. is a doubtful Penang plant. Not seen. Mount Ophir is the correct locality for this plant.

Alyxia stellata, Roem. and Sch. Not identified. Collected by Wallich.

Hunteria corymbosa, Rox.; small tree, common. (C. 237, 707).

Cerbera odollam, Gærtn.; tree 20-30 ft.; common. (C.)

Vincetoxicum rosea, L. is quite common in open sandy places. (C.)

Alstonia scholaris, Br. "Getah Paule;" tall straight tree; wood white, light. (C.)

Alstonia macrophylla, Wall. "Chengah Petrie;" tree tall but not large. Common. (C. 378).

Alstonia angustifolia, Wall. Not identified. Collected by Maingay.

Dyera costulata, Hook. fil.; large tree; wood very similar to that of *Alstonia scholaris*, not common. (C.)

Tabernæmontana corymbosa, Rox.; shrub 6-8 ft. Pulau Betong, not common. (C. 1763).

Tabernæmontana peduncularis, Wall.; small shrub. (C. 790).

Hoya coriacea, Bl. Not identified. Penang? Wallich 8163. (Probably wrong locality).

Ceropegia lucida, Wall.; slender twining herb. Penara Bukit, rare. (C. 1007).

LOGANIACEÆ.

Buddleia, sp. Top of Government Hill, probably introduced. (C. 1058).

Fagraea obovata, Wall.; small tree. Government Hill, not common. (C. 346, 2432).

Fagraea racemosa, Jack; small tree 10-15 ft. Common at the Waterfall. (C. 280).

Fagraea fragrans, Rox. "Temusu;" tree 20-30 ft.; not common. (C. 314, 1366).

Fagraea Wallichiana, Benth. "Temusu Bukit;" tree 40-50 ft.; fl. yellow. Common at 2000-2500 ft., and a valuable timber. (C. 375).

Strychnos laurina, Wall.; climbing shrub. (C. 1021).

Strychnos malaccensis, Benth; Muka Head. (C. 1515).

Strychnos pubescens, Clarke. (C. 970).

Strychnos Tieute, Lesch.? (C. 709, 1490).

BORAGINEÆ.

Cordia subcordata, Lam.; large shrub. Coast. (C. 303).

Ehretia Wallichiana, Hk. f. & T.; tree 40-50 ft. (C. 1034).

Tournefortia Wallichii, DC; large rambling shrub. (C. 1011).

Heliotropium indicum, L.; common annual weed. (C. 345).

CONVOLVULACEÆ.

Erycibe Princei, Wall. Recorded from Penang as collected by Wallich in Flor. Brit. India is an error. Wallich's plant came from Singapore where it is common.

Erycibe Griffithii, Clarke;? large climbing shrub; fruit $1\frac{1}{2}$ inch. long, brown. Government Hill. (C. 772).

Erycibe,?; large scandent shrub. (C. 2837).

Erycibe coriacea, Wall. scandent shrub; berry $\frac{1}{4}$ inch. Government Hill. (C. 911, 1273).

Erycibe, sp.; large climber; named *E. Griffithii* in herb. Penang but is not the same as 772. (C. 181).

Argyrea tiliaefolia, Wight. Pulau i etong. (C. 1970).

Lettsomia strigosa, Roxb. Waterfall. (C. 498).

Lettsomia adpressa, Miq.; twiner; corolla rosy pink. A rather common plant. (C. 318).

Lettsomia penangiana, Miq.; large climber; flower not seen. (C. 1586).

Ipomen angustifolia, Jacq.; slender twiner; corolla yellow (C. 1068).

Ipomea denticulata, Choisy. (C. 1974).

Ipomea staphylinn, Roem. & Sch.; large climber. (C. 470).

Ipomea cymosa, Roem. & Sch.; twiner; corolla pink. Bukit Padre. (C. 1083).

Ipomea pes-tigridis, L. Open sunny places, (C. 3033).

Ipomia Turpethum, Br. (C. 1703).

Calystegia hedericea, Wall. "Penang, Porter."

Evolvulus alsinoides, L.; prostrate herb; corolla pale blue. Open sandy places. (C. 1601).

Breweria cordata, Bl.; large climber; corolla white, not uncommon. (C. 1066).

SOLANACEÆ.

Solanum nigrum, L., herb 12-18 in.; fruit black. (C.).

Solanum torvum, Swartz; shrub 4-6 feet. (C. 90).

Solanum indicum, L.; shrubby 3-4 feet. (C. 10).

Solanum sarmentosum, Nees. 2628 F. Penang 1822 Wallich. (locality probably wrong).

Solanum trilobatum, L.; climber, rare. (C. 851).

Solanum aculeatissimum, Jacq.; procumbent; berry globose 1-1½ in., orange. (C.).

SCROPHULARINEÆ.

Adenosma capitatum, Benth.; herb 1-2 feet; fl. blue. Waterfall (C. 1886).

Adenosma hirsutum, Kz. (C. 22381).

Limnophila villosa, Bl. Waterfall. (C. 1839).

Herpestis Monniera, H. B. & K.; small herb; common in swamps. (C. 1939).

Artanema sesamoides, Benth.; large herb; fl. lilac. Not uncommon in the south of the island. (C. 945).

Torenia polygonoides, Benth. Not seen.

Torenia mucronulata, Benth.; prostrate herb. Waterfall. (C. 1837).

Torenia ciliata, Smith. Waterfall, not common. (C.).

Vandellin crustacea, Benth.; fl. lilac or blue. (C. 1854).

Vandellia pedunculata, Benth. "Kra Nasi." Paddy fields' common. (C. 1973).

Bonnaya veronicaefolia, Spreng.; fl. white and pink, or lavender and white; common. (C. 1852).

Bonnaya brachiata, Link. & Otto. (C. 1851).

Scoparia dulcis, L. "Cha Padang." Very common. (C. 688).

Striga lutea, Lour.; herb 4-6 in.; fl. yellow. Common in open places among long grass, (C. 1840).

LENTIBULARIÆ.

Utricularia flexuosa, Vahl.; fl. golden yellow; common in ditches and rice fields. (C. 519).

Utricularia bifida, L.; not uncommon in marshy land among grass. (C. 1177).

Utricularia racemosa, Wall.; fl. white; grows with *U. bifida* at the Waterfall. (C.).

Utricularia orbiculata, is said to have been collected in Penang near the Waterfall by Prof. Goebel. Not seen.

GESNERIACEÆ.

Æschynanthus Lobbianna, Hook. fil.; Government Hill. (C.).

Æschynanthus marmorata, N.E. Brown; fl. green and yellow. Government Hill 2,000-2,500 feet; generally on large trees. (C. 1700-2142).

Didymocarpus cordata, Wall.; 1-2 feet or more high; corolla white. Damp shady ravines. (C. 452).

Didymocarpus corchorifolia, Wall. Government Hill. (C. 1239).

Didymocarpus crinita, Jack; stem 6-12 in.; corolla white (pale purple form in Perak) common. (C. 410).

Didymocarpus reptans, Jack; creeping; corolla violet or white. Waterfall, not common. (C. 1706).

Didymocarpus incana, Benth.; leaves silvery grey; corolla blue. Common on damp rocks (C. 297).

Didissandra frutescens, Jack; shrubby, 2-3 feet. Damp ravines at 1,600-2,000 feet. (C. 828).

Cyrtandromæa acuminata, Benth.; stem 1-2 feet; corolla white. Penara Bukit, rare. (C. 1016).

Stauranthera grandiflora, Benth.; herb 3-6 in.; fl. lavender and yellow. Balik Pulau, very rare. (C. 1753).

Rhynchotechum parviflorum, Bl. var. *penangensis*, not identified. A very doubtful plant based on imperfect material.

Cyrtandra dispar, D. C.; shrub 3-5 ft. Moniot's Rd. (C. 1204).

Cyrtandra decurrens, De Vriese; stem 12-18 in, fleshy; corolla greenish white. (C. 1018).

Cyrtandra, sp. Pulau Betong. (C. 3035).

BIGNONIACEÆ.

Oroxylum indicum, Vent. tree 30-40 ft. Pulau Betong, not common. (C. 983).

Dolichandrone Rheedii, Seem.; small tree; corolla large white, fugacious. Tidal Swamps. (C. 1184).

Stereospermum chelonioides, D. C. tree 50-60 ft; corolla pale yellow; Capsule 12-24 in long. (C. 2431).

Stereospermum glandulosum, Miq.; small tree. (C. 140).

Diplanthera bancana, Scheff.; tree 30-40 ft.; corolla yellow. Not common. (C. 870, 1569).

PEDALINEÆ.

Sesamum indicum, D. C. annual herb 1-2 ft. (C. 1582).

ACANTHACEÆ.

Thunbergia alata, Bojer. Government Hill. Introduced. (C.)

Nelsonia campestris, Br.; prostrate herb. Balik Pulau. (C. 1750).

Ebermaiera lanceolata, Hassk. Not identified. Penang Porter, Wall. No. 7489.

Ebermaiera angustifolia, T. And.; dwarf herb. Pulau Betong. (C. 1931).

Ebermaiera axillaris, Nees. Not identified. Is stated by Anderson to have been collected by Wallich, No. 2414.

Ebermaiera elongata, Nees. "Wallich in Penang." Nees makes no mention of it in the *Plantæ Asiaticæ Rariores* which he doubtless would have done had he found it in Wallich's collections. It is a native of Java.

Ebermaiera lasiobotrys, Nees.; Pulau Betong. (C. 1350).

Ebermaiera racemosa, Miq.; herb 6-10 in. Common on West and Government Hills 2,000-25,000 ft. (C. 406).

Hypographila salicifolia, Nees. Pulau Betong. (C. 1757).

Ruellia repens, L.; stem 9-12 in.; corolla rose colour fugacious. (C. 1876).

Strobilanthes Maingayii, Clarke. Not identified. One of Maingay's plants, locality quite uncertain.

Strobilanthes collinus, Nees. Not identified. Collected by Porter. Wall Cat 2343.

Strobilanthes, sp.; shrubby; corolla white. Government Hill. (C. 298).

Strobilanthes, sp. Muka Head. (C. 954).

Strobilanthes, sp.; prostrate herb. Government Hill 2,000 feet. (C. 2761).

Acanthus ebracteatus, Vahl. Common in tidal swamps. The seeds are reputed cure for boils. (C. 510).

Barleria conspicua, Nees. Not identified.

Asystasia intrusa, Br.; stem 3-4 ft. Waterfall. (C. 1853).

Asystasia coromandeliana, Nees; fl. white, purple, or yellow. Common. (C. 2419).

Asystasia Kunthiana, Nees. Not identified. A doubtful plant collected by Wallich.

Eranthemum album, Nees. Not seen. Collected by Porter.

Eranthemum malaccensis, Clarke; small shrub. Waterfall, not uncommon. (C. 499, 348).

Eranthemum graciliflorum, Nees. Not identified. A very doubtful plant. Wall. Cat 2427.

Andrographis paniculata, Nees; stem 6-18 in. Not uncommon in open spots. (C. 1877).

Justicia inconspicua, Wall. Cat 2475 is absolutely unknown.

Justicia gandarussa, L. Balik Pulau. (C. 791).

Justicia decussata, Roxb. Pulau Betong. (C. 1765).

Justicia ptychostoma, Wall. Not identified. Collected by Porter.

Justicia Maingayii, Clarke. Not identified. Collected by Maingay.

Adhatoda vasica, Nees; shrub 4-5 ft. Ayer Hitam. (C. 1017).

Peristrophe acuminata, Nees. Balik Pulau. (C. 681).

VERBENACEÆ.

Lantana indica, L.; shrub 1-2 feet; fl. pink. Balik Pulau, not common. (C. 121).

Lantana camara, L.; has run wild all over the Island.

Stachytarpheta indica, Vahl; 2-3 feet. Common. (C. 867).

Callicarpa arborea, Rox.; small tree 25-30 feet. Telok Bahang, not common. (C. 699).

Callicarpa cana, L. apparently Wallich's No. 1834. 1 is taken as the authority for this as a Penang plant in the Flora of British India but that was cultivated in the Calcutta gardens.

Callicarpa pedunculata, Br. Wallich 1834, 2. Collected by Geo. Porter. Not seen.

Callicarpa longifolia, Lamk.; shrub 6-10 feet; berry white. Government Hill. (C. 309).

Premna cordifolia, Rox. shrub. (C. 681.)?

Premna divaricata, Wall; scandent shrub. (C. 215).

Gmelina asiatica, L. tree 25-30 feet. Common. (C. 454).

Vitex trifolia, L.; small tree. (C. 98).

Vitex pubescens, Vahl. "Halban"; a very common tree. (C. 64).

Vitex coriacea, Clarke; tree compact, 20-25 feet; flower purple. Government Hill. (C. 323).

Vitex vestita, Wall.; small tree; corolla yellow. Government Hill. (C. 265).

Vitex glabrata, Br.; tree, larger than either of the preceding. Not common. (C. 1071).

Clerodendron Lobbii, Clarke. Not identified. Collected by Lobb.

Clerodendron calamitosum, L.; small shrub. Introduced? (C).

Clerodendron deflexum, Wall.; shrub 1-3 feet. Government Hill. (C. 81).

Clerodendron paniculatum, Rox.; shrub 6-10 feet; corolla red. Not seen except near villages. (C. 313).

Clerodendron neriifolium, Wall.; common on the sea shore. (C. 2782).

Clerodendron villosum, Bl.; shrub 4-6 ft. A far more common plant than either of the others. (C. 101).

Clerodendron acuminatum, Wall.; shrub. (C. 1539).

Peronema canescens, Jack; tree 25-30 ft. Common near the race course; not seen elsewhere. (C. 2427).

Sphenodesma triflora, Wight; scandent shrub. Waterfall. (C. 210).

Sphenodesma barbata, Schauer. Collected by Geo. Porter.

Sphenodesma pentandra, Jack; large rambling shrub (C. 269).

Avicennia officinalis, L.; small tree common in Mangrove swamps. (C.).

LABIATÆ.

Ocimum basilicum, L.; herb 2 ft. Tanjong Tokong. (C. 2492).

Mochosma polystachyum, Benth; annual herb. (C. 34).

Coleus atropurpureus, Benth. Waste places, common. (C.)

Hyptis brevipes, Poit.; is a very common plant. (C. 1111).

Hyptis suaveolens, Poit.; is not nearly so common as *brevipes*. (C. 1250).

Pogostemon patchouli, Pelletier; found as an escape, is not in my opinion indigenous (C.)

Diosphylla auricularia, Bl. "Ekor Kuching." Common in damp places. (C. 868).

Anisomeles ovata, Br. Waterfall &c.; common. (C. 65).

Anisomeles malabarica, Br. Not identified. Collected by Wallich no 2037, 2.

Leucas zeylanica, Br. Annual herb 1-2 ft.; corolla white. Very common. (C. 409).

Leucas tinifolia, Spreng. is a much smaller plant. (C. 1115).

Gomphostemma parviflorum, Wall. not seen. There is some error in the Flora of British India and in Bentham's paper on the Labiatae in the Plantae Asiaticae Rariores ii 12. *G. parviflorum* no 2158 was collected at Silhet but *G. crinitum*. Wall. Cat no 2159 was collected in Penang by Wallich in 1822. It has not been gathered there since.

PLANTAGINEÆ.

Plantago major, L. Government Hill 2,500 feet. (C. 2157).

NYCTAGINEÆ.

Boerhaavia repens, L. Not uncommon near George Town. (C. 902).

AMARANTACEÆ.

Amaranthus spinosus, L. "Bayam." A good vegetable (C. 245).

Cyathula prostrata, Bl. Common in coco-nut gardens. (C. 17-944).

Pupalia atropurpurea, Miq. The authority for this is Wallich's Catalogue 6933-2. Penang and Singapore, as the plant never since been seen in the Malay Peninsula, the locality is probably wrong.

Achyranthes aspera, L. Common in sandy soils. (C. 1835).

Alternanthera sessilis, Bl. Common everywhere. (C. 1043).

POLYGONACEÆ.

Polygonum tomentosum, Willd.; tall herb. Damp places. (C. 16).

Polygonum barbatum, L. Pulau Betong. (C. 943).

Polygonum strigosum, Br. Not seen. Collected by Wallich.

Polygonum pedunculare, Wall. Commoner than either of the others. (C. 921).

NEPENTHACEÆ.

Nepenthes ampullaria, Jack. Penara Bukit, rare. (C.).

Nepenthes phyllamphora, Willd. Common among low bushes in swampy places. (C. 1202).

Nepenthes albo-marginata, Lobb. Government Hill, common. (C.).

ARISTOLOCHIACEÆ.

Thottea dependens, Klotzsch; shrubby 1-2 feet. Waterfall. (C. 1170-1507).

Aristolochia Roxburghii, Klotzsch; twiner. Waterfall. (C. 2464).

Aristolochia, sp. leaves deeply lobed. Government Hill. (C. 330),

Bragantia tomentosa, Bl. Waterfall. Common. (C. 2754).

PIPERACEÆ.

Piper ribesioides, Wall. West Hill, rare. (C. 2256).

Piper sumatranum, Cas. Not identified. Wallich 6646B. Not only is the identification of this specimen doubtful but the locality Penang is marked doubtful in the catalogue.

Piper caninum, Bl. Balik Pulau. (C.).

Piper longum, L. "Kudak." Common. (C. 887).

Piper pectuloides, Rox. Government Hill 2,000 ft. (C. 2291).

Piper miniatum, Bl. Batu Feringgi. (C. 821, 2290).

Piper rostratum, Rox.? West Hill. (C. 967).

Piper penangensis, Cas. Not identified. Wallich 6644 B.

Piper lonchitis, Roem. and Schultes. Government Hill. (C. 734).

Piper leptonema, Hook. fil. (C. 1771).

Piper attenuatum, Ham. Not identified. Locality very doubtful.

Piper subpeltatum, Willd. Wallich 6638a. Not seen.

Piper sulcatum, Bl. Government Hill. (C. 443).

Piper porphyrophyllum, N. E. Brown. Common on Government Hill. (C.)

Piper, sp.; near *Blumii*. Government Hill. (C. 792).

Peperomia pellucida, Kunth.; herb, succulent, 6-18 in. Common about George town. (C. 1968).

CHLORANTHACEÆ.

Chloranthus officinalis, Bl.; compact undershrub; berries white. (C. 1180, 1708).

Chloranthus brachystachys, Bl.; shrubby, less compact than the preceding, berries red. (C. 1050).

MYRISTICACEÆ.

Myristica fragrans, Houtt. Cultivated, often found on land that has reverted to forest. (C.)

Myristica bracteata; A. D. C.; large tree 50 ft. high. Government Hill, rare. (C. 1497).

Myristica Maingayii, Hook. fil.; a tree 50-60 ft. high and 18-24 in. in diam.; fruit oblong 3 × 2 in. larger than *M. fragrans* Government Hill, rare. (C. 2455).

Myristica elliptica, Wall.; tree 30-40 ft.; fruit larger than *M. fragrans*; and mace blood-red, lacinate as stated by Maingay. (C. 1122).

Myristica superba, Hook. fil.; collected by Phillips. (C. 2966).

Myristica tomentosa, Hook. fil.; tree 25-30 ft. Waterfall and other places (C. 1748).

Myristica, sp. closely resembling *M. Canarioides*, King. (C. 943).

Myristica Griffithii, Hook. fil.; a tree 40-50 ft. Government Hill 1500 ft. (C. 2406).

Myristica Farquhariana, Wall.; medium-sized tree, not uncommon. (C. 487, 804).

Myristica Forbesii, King. Collected by Maingay. Not seen.

Myristica Irya, Gærtn.; large spreading tree 60-70 ft. high. South of the Island, not common. (C. 936).

Myristica oblongifolia, King. Not seen.

Myristica Scortechinii, King; Government Hill. (C. 2769).

Myristica intermedia, Bl.; medium sized tree. West Hill 1500 ft. (C. 1044).

Myristica furfuracea, Hook. fil.; Government Hill 2000 feet. (C. 1459).

Myristica furfuracea var. *major*, King. (C. 1459, 2456).

Myristica laurina, Bl.; small tree. West Hill, not uncommon. (C. 1191, 2457).

Myristica glauca, Bl. (C. 935).

Myristica Missionis, Wall.; tree 20-30 feet. Pulau Betong. (C. 700, 935).

Myristica glaucescens, Hook. fil.; tree 25-30 feet, common. (C. 1559).

Myristica Hookeriana, Wall.; a tall but not very large tree. Government Hill. (C. 202).

Myristica Curtisii, King; a tree with slender branches. Waterfall. (C. 1024, 1301).

Myristica racemosa, King; tree 40-50 feet high. Nalm pass, rare. (C. 934).

Myristica Ridlevana, King; tree 40-50 feet high. Government Hill. (C. 2458.)?

Myristica eugeniaefolia, A. D. C. is said to have been collected in Penang. Not identified. It is little known plant and probably identical with one of the preceding.

MONIMIACEÆ.

Kibara coriacea, Endl.; small tree 20 feet. Government Hill, not common. (C. 1242, 2255).

LAURINEÆ.

Apollonias sp.; tree 40-50 feet. Penara Bukit. (C. 1098).

Beilschmiedia sp.; large tree; fruit ovate, 1 inch long. Government Hill 2000 feet, rare. (C. 1015).

Dehaasia Kurzii, King? compact growing tree. West Hill. (C. 1193).

Dehaasia sp.; tree 30-40 feet; fruit with fleshy red pedicel $\frac{3}{4}$ inch long. Government Hill. (C. 1183).

Endiandra sp.; tall tree, branches slender; fruit oblong, $1\frac{1}{4}$ inches. Moniot Road. (C. 1230).

Ediandra, sp.; tree 40-50 feet high, fruit 2 in. long black-purple. Batu Feringy. (C. 3068).

Endiandra, tree 30-40 tall, leaves obovate 7×4 in. fruit oblong dark purple. Pantie Achie (C. 3067).

Micropora Curtisii, Hook fil; tree 40 feet leafy. Government Hill 1000-2000 feet. (C. 524, 1214).

Cinnamomum javanicum, Bl. very doubtful, some leaves collected by Walker are the only authority for this here.

Cinnamomum iners, Reinw.; tree medium-size, very common. (C. 156).

Cinnamomum nitidum, Bl.; recorded hence in Flor. Brit. Ind. is an error. No. 2583a Wall. Cat. is *C. iners* Reinw. and

was collected in Penang No. 2583 B.C.D. which are referred to *C. nitidum* were not from Penang.

Cinnamomum vimineum, Nees; small tree; leaves when bruised emit a strong aromatic scent. Government and West Hills. (C. 1571).

Cinnamomum mollissimum, Hook fil; small tree 20-25 feet. West Hill 2500 feet. (C. 308)

Cinnamomum Parthenoxylon, Meissn. "Medang"; large tree, timber valuable. (C. 512, 1038).

Cinnamomum Lampongum, Miq.; Medium-sized tree. (C. 2252).

Machilus rimosa, Bl.; large spreading tree. Penara Bukit, not uncommon. (C. 942, 1030).

Phæbe opaca, Bl. Not seen. Collected by Porter.

Phæbe multiflora, Bl.; large tree. Pulau Betong. (C. 2739).

Alseodaphne peduncularis, Hook fil. Not identified, Wallich 2596.

Alseodaphne sp.; middling sized tree. Waterfall. (C. 1536).

Actinodaphne macrophylla, Nees; tree 30-40 feet. Waterfall and Pulau Betong. (C. 2304, 2473).

Actinodaphne pruinosa, Nees; tree 40-50 feet. Moniot's Road, rare. (C. 1020).

Litsæa sebifera, Pers.; tree 20-30 feet. West Hill. (C. 2305).

Litsæa sp.; near *sebifera*, but with fasciculate not racemose inflorescence (C. 1182).

Litsæa sessiliflora, Hook fil; large shrub. Government Hill. (C. 649).

Litsæa grandis, Wall.; tree 40-50 feet. Top of Government Hill. (C. 1104).

Litsæa polyantha, Juss.; small tree. Government Hill 2500 feet (C. 283).

Litsæa amara, Bl. Not identified. Maingay.

Litsæa salicifolia, Rox.; small tree. Batu Hitam. (C. 787).

Litsæa læta, Wall.; small tree. Moniot's Road. (C. 2159).

Litsæa penangiana, Hook fil; small tree. (C. 1587) ?

Litsæa myrsiticifolia, Wall.; tree 30-40 feet. (C. 769, 1079).

Litsæa nitida, Bl.; tree 40 feet. Moniot's Road. (C. 795).

Litsæa zeylanica, C. & Fr. small tree. Tanjong Bunga, &c., common. (C. 468, 446).

Litsæa rugosa, Kurz; Not identified. an obscure plant probably not a *Litsea* at all.

Litsæa Noronhiana, Bl.; large tree 50-60 feet. Government Hill. 1200 feet. (C. 2449).

Litsæa sp.; tree 30-40 feet. Top of Government Hill. (C. 1103).

Litsæa sp.; small tree; fruit the size of a pea. (C. 733, 1073).

Lindera oxyphylla, Benth. Not identified. Porter. (Wall cat 2547).

Cassytha filiformis, L.; twiner, common on rocks and low bushes near the coast. (C. 1203).

PROTEACEÆ.

Helicia attenuata, Bl.; small tree. Telok Bahang. (C. 1181).

Helicia, sp.; small tree quite distinct from *H attenuata*. Penara Bukit. (C. 3020).

Helicia petiolaris, Bennett; not seen. Formerly cultivated in Penang according to Jack. It is not rare in Singapore.

THYMELEACEÆ.

Wikstræmia indica, C. A. Mey.; small shrub. Open places, often in Nutmeg plantations. (C. 30).

Linostoma pauciflorum, Griff.; climbing shrub. Muka Head. (C. 713).

Aquilaria Malaccensis, Lam.; tree 40-60 feet bark smooth, grey; flower white, scented. (C. 1534).

This tree produces the *Gaharu* Wood.

ELEAGNACEÆ.

Elæagnus latifolia, L.; large climbing shrub. Top of Government Hill. (C. 904).

LORANTHACEÆ.

Loranthus Lobbii, Hook fil. Government Hill. (C. 85, 389).

Loranthus pulcher, D. C. (C. 394).

Loranthus pentapetalus, Roxb. Government Hill and Penara Bukit, common. (C. 205).

Loranthus coccineus, Jack; Government Hill. (C. 243).

Loranthus ferrugineus, Rox. Government Hill; common. (C.)

Loranthus pentandrus, L. Ayer Hitam. (C. 476).

Loranthus ampullaceus, Rox. Government Hill. (C. 66).

Loranthus globosus, Rox. Not identified. Apparently only differs from the preceding in having fleshier narrower leaves and globose fruit, but *L. ampullaceus* is very variable in the form of the leaf.

Loranthus albidus, Bl. Government Hill and Penara Bukit, not uncommon. (C. 1113).

Viscum ovalifolium, Wall. Not seen.

Viscum articulatum, Burm. Government Hill. (C.).

Notothixos malayanus, Oliv.; parasitic, much branched shrub; fruit white. Coast. (C. 233).

SANTALACEÆ.

Henslowia lobbiana, A. D. C.; long twining shrub. Government Hill, common. (C. 21).

Henslowia buxifolia, Bl.; parasitic on the roots of trees in damp shady places. (C. 477).

Champereia Griffithiana, Bl.; small tree. Telok Bahang, not common. (C. 155, 1163, 2262).

EUPHORBIACEÆ.

Euphoria atoto, Forst.; shrubby 1-3 ft. Open sandy places near the coast. (C. 333, 1883).

Euphorbia pilulifera, L.; erect annual herb 1-2 ft. (C. 1838).

Euphorbia thymifolia, Burm.; much branched prostrate herb. Government Hill. (C. 1945).

Euphorbia, sp. near *neruifolia*, L. erect shrub, 2-5 ft. Common at the Waterfall. (C. 1567).

Bridelia stipularis, Bl. Not identified. Collected by Wallich. (C. 7878 N).

Bridelia tomentosa, Bl.; small tree. Common. (C. 112).

Bridelia penangiana, Hook. fil.; small tree. Government Hill. (C. 521).

Bridelia rufa, Hook. fil.; small tree. Waterfall. (C. 1480).

Bridelia Curtisii, Hook. fil.; large shrub. Telok Bahang. (C. 97).

Cleistanthus hirsutululus, Hook. fil.; small tree. Waterfall. (C. 1464).

Cleistanthus membranaceus, Hook. fil. (C. 1481).

Cleistanthus nitidus, Hk. f.; small tree, Government Hill. (C. 146).

Cleistanthus pedicellatus, Hook. fil.; small compact tree. Government Hill, rare. (C. 169).

Actephila excelsa var. *javanica*, Miq.; shrub, not common. (C. 674).

Phyllanthus emblica, L. "Buah Malakka;" spreading tree. Not common. (C.).

Phyllanthus frondosa, Wall.; shrub 3-5 ft.; fruit the size of a pea. Batu Feringgi. (C. 819, 1263).

Phyllanthus Kunstlerii, Hook. fil. Penang at 800-1000 feet, Kunstler Not seen.

Phyllanthus urinaria, L. Not seen.

Phyllanthus niruri, L.; herb 6-18 in. Common. (C. 1841).

Phyllanthus acutus, Wall. Collected by Finlayson (Wall. Cat. No. 7931) was probably not from Penang at all.

Phyllanthus pulcher, Wall.; shrub 2-4 ft.; common near villages, doubtfully wild. (C. 78).

Phyllanthus distichus, Muell. Arg. Cultivated.

Glochidion insulare, Hook. fil.; small tree. Government Hill. (C. 1728).

Glochidion hirsutum, Muell. Arg. Not identified.

Glochidion macrostigma, Hook. fil. Penara Bukit. (C. 852).

Glochidion Wallichianum, Muell. Arg. Not identified. Wallich 7873).

Glochidion obscurum, Bl.; tree 30-40 feet. Waterfall, common. (C. 46).

Glochidion nanogynum, Hook. fil. Not identified. Wallich 8003 B.

Glochidion desmocarpum, Hook. fil. Not identified. Penang, herb. Hooker perhaps from elsewhere.

Glochidion lævigatum, Hook. fil.; small tree. (C. 833, 1593).

Glochidion superbum, Baill.; small tree, common. (C. 111).

Glochidion coronatum, Hook fil.; large shrub sometimes a tree. (C. 213)

Glochidion Curtisii, Hook fil.; large shrub. Government Hill. (C. 670).

Glochidion perakense, Hook fil.; shrub. West Hill. (C. 1120).

Breynia discigera, Muell. Arg.; large shrub. West Hill. (C. 196).

Breynia coronata, Hook fil.; large shrub or small tree; common. (C.).

Sauropus albicans, Bl.; small undershrub. West Hill. (C. 963)

Cyclostemon longifolius, Bl.; tall tree. Government Hill.

Cyclostemon Curtisii, Hook fil.; small tree. Government Hill. (C.).

Cyclostemon sp.; small tree 12-15 feet. Waterfall. (C. 1776).

Choriophyllum malayanum, Benth.; tree 40 feet or more. Government Hill, &c.; common. (C. 1148, 881, 1198).

Aporosa ficifolia, Baill.; small tree. West Hill. (C. 1483, 1583).

Aporosa nigricans, Hook fil.; large shrub. West Hill. (C. 1570, 1574).

Aporosa microsphaera, Hook fil.; small tree. (C. 1482, 1466, 1470, 1467).

Aporosa aurea, Hook fil.; large shrub. Waterfall. (C. 1460).

Aporosa lunata, Benth; tree 20-30 feet. Moniots Road. (C. 1468).

Aporosa Benthamiana, Hook fil.; small tree 25-30 feet. (C. 1469).

Aporosa stellifera, Hook fil.; small tree. Penara Bukit. (C. 1472).

Aporosa Prainana, King mss.; small tree. Waterfall and Government Hill. (C. 1476, 1477).

Antidesma pachystachys, Hook fil. Not identified. Wallich 8569.

Antidesma velutinosum, Bl.; large shrub or small tree. Government Hill. (C. 672).

Antidesma velutinosum, var. *lancifolia*; shrub 6-10 feet. (C. 863).

Antidesma Ghæsembilla, Gærtn.; low spreading tree. Waterfall, &c. Common. (C. 782-1554).

Antidesma leucocladon, Hk. f.; shrub 5-6 feet high; ripe fruit black, smooth fleshy. (C. 2280).

Antidesma Bunius, Spreng. Moniot's Road 2278.

Antidesma fallax, Muell. Arg.; tree 30-40 feet. Penara Bukit. (C. 1473, 984?).

Antidesma Moritzii, Muell. Arg. small tree. Tanjong Bonga. (C. 230).

Antidesma sp.; small tree. Telok Bahang. (C. 1150).

Baccaurea parviflora, Muell. Arg.; small tree, common. (C. 166, 1169).

Baccaurea macrophylla, Muell. Arg. Not identified. Penang. Phillips.

Baccaurea Griffithii, Hook fil; tree 30-40 feet. Waterfall. (C. 1474).

Baccaurea brevipes, Hook fil; small tree, not uncommon. (C. 1568, 1478).

Baccaurea Wallichii, Hook fil. Not identified. Wallich 8073.

Baccaurea, sp. A. of Fl. Brit. Ind.; tree 30-40 feet; bark almost black. Government Hill. (C. 1463).

Baccaurea, sp. tree 20-30 feet high; berries white, globular, pubescent. (C. 3021).

Galearia fulva, Muell. Arg. Not identified. Wallich 8585c. Phillips.

Galearia pedicellata, Br. Not identified. Wallich 8585. Phillips.

Galearia Jackiana, Br. Not identified. Penang, Jack.

Galearia subulata, Muell. Arg. In Sonder's herbarium from Lindley's herbarium. Not identified, probably from elsewhere.

Galearia Lindleyana, Muell. Arg. shrub. West Hill. (C. 176).

Microdesmis caseariæfolia, Pl.; erect shrub 4-6 feet. Waterfall. (C. 1491).

Microdesmis sp. small shrub. Waterfall. (C. 1489).

Aleurites moluccana, Willd.; tree 30-40 feet or more. Not seen in any place where it might not easily have been introduced. (C. 1210).

Croton argyratus, Bl.; small tree 20-25 ft., common. (C. 232)
Trigonostemon longifolius, Baill.; shrub 8-10 ft. Waterfall.
 (C. 1475).

Trigonostemon indicus, Muell. Arg.; small tree 15-25 ft.;
 back smooth, grey. Government Hill. (C. 218).

Ostodes muricata, Hook. fil.; shrub 6-10 ft. Waterfall.
 (C. 1484).

Dimorphocalyx malayanus, Hook. fil.; tree 20-25 feet;
 flower white, conspicuous. Waterfall. (C. 806).

Dimorphocalyx Kunstlerii, King; small tree. Waterfall.
 (C. 1517).

Erismanthus obliqua, Wall.; small tree. 25 ft. Waterfall,
 common. (C. 1471).

Agrostistachys Gaudichaudi, Muell. Arg. collected by Gau-
 dichaud. Not identified.

Agrostistachys longifolia, Benth. var. *malayana*. Not iden-
 tified. Wallich.

Claoxylon indicum, Hassk.; small tree. Pulau Betong. (C.
 897).

Claoxylon longifolium, Muell. Arg.; small tree. (C. 675).

Claoxylon longifolium var. *brachystachys*. (C. 1101).

Claoxylon Wallichianum, Muell. Arg.; shrub. Government
 Hill. (C. 644).

Acalypha indica, L.; annual herb, common. (C. 891, 2147).

Cœlodepas Wallichianum, Benth.; small tree with very hard
 wood. Penara Bukit. (C. 1271).

Alchornea villosa, Muell. Arg., var. *glabrata*; tall shrub.
 Government Hill. (C. 214).

Alchornia discolor, Hook. fil.; shrub. Government Hill. (C.
 822).

Cœlodiscus montanus, Muell. Arg.; shrub. Penara Bukit.
 (C. 1261).

Mallotus barbatus, Muell. Arg. Not identified. Wallich
 7822.

Mallotus cochinchinensis, Lour.; small tree; common. (C.
 43).

Mallotus Porterianus, Muell. Arg.; large shrub or small
 tree. (C. 1461, 1462).

Mallotus Griffithianus, Hook. fil.; tree 25-30 feet. Waterfall and Government Hill. (C. 2260-2286).

Mallotus lancifolius, Hook. fil.; shrub 8-10 feet. Government Hill 2,000-2 500 feet. (C. 684).

Mallotus sub-peltatus, Muell. Arg. Government Hill. (C. 157).

Mallotus penangensis, Muell. Arg. Government Hill. (C. 755).

Mallotus repandus, Muell. Arg. Not seen.

Cleidon javanicum, Bl.; not seen.

Macaranga pustulata, King; small tree. Government Hill. (C.).

Macaranga Tanarius, Muell. Arg.; spreading tree 30-40 feet. Pulau Butong, not common. (C. 1949).

Macaranga Curtisii, Hook. fil.; medium-sized tree. West Hill. (C. 286).

Macaranga hypoleuca, Muell. Arg.; small straight tree 25-30 feet. Government Hill. (C).

Macaranga megalophylla, Muell. Arg.; tree 30-40 feet. Government Hill and Waterfall. (C. 1588).

Macaranga populifolia, Muell. Arg. Not identified.

Macaranga javanica, Muell. Arg.; small tree, common. (C. 76).

Macaranga triloba, Muell. Arg.; small tree. Government Hill. (C. 1479).

Macaranga Lowii, King; small tree 15-20 feet. Government Hill, not uncommon. (C. 276, 799).

Endospermum malaccense, Muell. Arg. medium-sized spreading tree. Government Hill. (C. 316).

Gelonium multiflorum, A. Juss.; small tree. Pulau Betong. (C. 924).

Gelonium bifarium, Rox.; small tree. Government Hill. (C. 53).

Gelonium glomerulatum, Hassk.; shrub. Telok Bahang. (C. 224).

Chaetocarpus castanocarpus, Thwaites; small tree. Waterfall Government Hill &c.; common. (C. 725, 449, 658).

Baliospermum axillare, Bl.; shrub. Government Hill. (C. 77).

Epiprinus malayanus, Griff.; small tree 15-20 ft. Nalm, rare. (C. 2472).

Cnesmone javanica, Bl. "Jelatang Rusa"; climbing shrub with stinging hairs. Government Hill. (C. 384).

Sapium baccatum, Rox.; tree 25-30 feet. Not uncommon. (C. 836).

Sapium indicum, Willd.; tree 30 feet. Coast. (C. 1605).

Excæcaria agallocha, L.; tree; common in Mangrove swamps. (C. 11, 1026).

Excæcaria quadrangularis, Muell. Arg.; shrub 2-4 feet. Waterfall and Pulau Betong (C. 989).

Sebastiana chamælea Muell. Waterfall-gardens in grass plots.

URTICACEÆ.

Trema timorensis, Bl.; small tree; branches slender. Bagian Jermal, rare. (C. 1935).

Trema angustifolia, Bl.; small tree; branches rigid. (C. 1893).

Trema amboinensis, Bl.; tree 20-30 feet; very quick growing. Common. (C. 102, 268).

Gironniera nervosa, Pl.; medium-sized tree, 30-40 feet. Government Hill, rare. (C. 192).

Gironniera subæqualis, Pl.; tree 30-40 feet. Telok Bahang, rare. (C. 1074).

Gironniera parvifolia, Pl.; tree 20-30 feet. Government Hill, common. (C. 529, 1506).

Phyllochlamys Wallichii, King; thorny shrub 6-10 feet. Waterfall, not uncommon. (C. 174).

Streblus asper, Lour. Not seen.

Sloetia Sideroxylon, Teysm and Binnend. "Tampinis"; tree medium size, wood durable, in great demand. (C. 198).

Sloetia penangiana, Oliv. "Tampinis daun besar"; tree, branches less rigid and leaves larger than the preceding. (C. 380).

Sloetia Wallichii, King; not identified. Wallich No. 5090.

Ficus pisifera, Wall.; small tree. Sungei Penang. (C. 1132).

Ficus gibbosa, Bl.; small tree. Coast. (C. 948).

Ficus parietalis, Bl.; Not identified.

- Ficus pilosa*, Reinw. Not identified. Very doubtful.
Ficus bracteata, Wall. Not seen.
Ficus globosa, Bl. (C. 2508).
Ficus xylophylla, Wall. Government Hill 2500 feet. (C. 2216).
Ficus rigida, Miq.; is common. (C).
Ficus dubia, Wall.; tree 30-40 feet. Penara Bukit. (C. 1128).
Ficus glabella, Bl.; small tree. Tanjung Bunga. (C. 1126).
Ficus retusa, L. Not seen.
Ficus vasculosa, Wall.; tree, Penara Bukit. (C. 1133).
Ficus punctata, Thunb. Not seen. Wallich.
Ficus rostrata, Lamk. Penara Bukit, common. (C. 1124, 1127).
Ficus hispida. L. f.; small tree 15-20 feet, common. (C).
Ficus Miquelii, King; tree. Waterfall. (C. 1749, 1889).
Ficus Ribes, Reinw. Not identified.
Ficus fistulosa, Reinw. Not identified.
Ficus lepicarpa, Bl.; small tree. Waterfall. (C. 1130).
Ficus recurva, Bl.; creeping shrub. (C. 171, 1131).
Ficus annulata, Bl. Sungei Penang. (C. 2509).
Ficus Curtisii, King. A large tree. Glugor and Telok Bahang. (C. 1982).
Ficus villosa, Bl.; woody creeper; generally on rocks. West Hill, &c., common. (C. 173).
Ficus diversifolia, Bl.; small erect shrub 2-6 feet; not uncommon. (C. 873).
Ficus alba, Reinw.; small tree 10-12 feet, common. (C. 1129).
Ficus chrysocarpa, Reinw. Not seen. Wallich.
Ficus glandulifera, Wall.; small tree; not uncommon up to 2500 feet. (C. 1134).
Ficus variegata, Bl. Not seen. King's collector.
Ficus chartacea, Wall. var. *torulosa*. (C. 1711).
Artocarpus rigida, Bl. "Iamponey"; large tree, sometimes cultivated for its fruit. (C. 1984).
Artocarpus integrifolia, L.; The Jack, cultivated in almost every village, often found in a half-wild state. (C).

Artocarpus polyphema, Persoon. The Champedak. Cultivated.

Artocarpus lanceifolia, Rox. "Nangka pipit"; tree. Penara Bukit. (C.).

Artocarpus chaplasha, Rox. Government Hill. (C.).

Artocarpus Lakoocha, Rox.: tree 30-40 feet, deciduous. Government Hill 2,000 feet. (C. 1222, 006, 1251),

Artocarpus Gomeziana, Wall. var. *Griffithii*, King. "Tampong"; compact leafy tree 30-40 feet. (C. 657).

Balanostreblus ilicifolia, Kurz. Not identified. On Penang Island. King's collector.

Conocephalus suaveolens, Bl. "Ara jankang"; large climbing shrub, Pulau Betong. (C. 1745).

Conocephalus amœnus, King; large climber. Pulau Betong (C. 1980).

Conocephalus Scortechinii, King. Government Hill. (C.)

Conocephalus subtrinervius, Miq. Penara Bukit, by the side of streams. (C. 1009)

Fleurya interrupta, Gaud.; annual herb 1-2 feet; common in waste places. (C. 915).

Laportea sp.; shrub 6-10 feet. Wang Seraya and Nalm Pass, not common. (C. 1100).

Pilea muscosa, L. Abundant at the Waterfall. (C. 1233).

Pellionia javanica, Wedd.; dwarf herb. Balik Pulau. (C. 682).

Pellionia acaulis, Hook. fil. Not identified. On damp rocks, King's collector.

Elatostema sessile, Forst.; herb 12-18 inches. Damp shady places, Balik Pulau. (C. 922).

Elatostema molle, Wedd. Not identified. Phillips.

Procris frutescens, Bl.; stem woody 1-2 feet. Government Hill. (C. 486).

Pouzolzia mdica, Gaud.; prostrate herb. (C. 1938, 2145).

Pouzolzia pentandra, Benn.; 2-3 feet, common. (C. 1879).

Pipturus mollissimus, Wedd.; large climbing shrub, Penara Bukit. (C. 673, 1032).

JUGLANDEÆ.

Engelhardtia Wallichiana, Lindl.; medium-sized tree. Government Hill 2000-2500 feet. (C. 56).

Engelhardtia serrata, Bl.; tree 30-40 feet. Muka Head. (C. 1716).

Engelhardtia nudiflora, Hook. fil. Not seen. Government Hill. Maingay.

MYRICACEÆ.

Myrica sp.; middling sized tree. West Hill 2500 feet. (C. 903).

CASUARINEÆ.

Casuarina equisetifolia, Forst. "Kayu Ru." Cultivated; indigenous. (C.)

CUPULIFERÆ.

Quercus odocarpa, Korth. "Berangan Babi"; large tree. Telok Bahang, rare. (C. 434).

Quercus Wallichiana, Lindl.; medium-sized tree, 20-40 feet. Government Hill, common. (C. 252, 253).

Quercus sundaica, Bl.; low spreading tree. Government Hill 2000 feet, rare. (C. 367, 442).

Quercus Curtisi, King; medium-sized tall tree. Waterfall. (C. 1253).

Quercus Wenzigiana, King; tree 20-30 feet. Government Hill 2,000 feet. (C. 360, 368).

Quercus Rassa, Miq.; tree, branches slender, drooping. Government Hill 2000 feet. (C. 362, 363, 255).

Quercus lucida, Rox.; tall tree, not very large. West Hill 2500 feet. (C. 436, 930).

Quercus Omalkos, Korth.; tall tree. Government Hill 2000 feet. rare. (C. 435).

Quercus cyclophora, Endl.; large tree, not uncommon. Government Hill 1000-2000 feet. (C. 361).

Quercus discocarpa, Hance; tree about 25 feet. Top of Government Hill. (C. 1155).

Quercus encleisocarpa, Korth.; small tree. Government Hill 1000 feet. (C. 254).

Quercus Maingayii, Benth.; large tree. Government Hill and Bukit Laksamana, rare. (C. 267).

Quercus sp. (*nitida*, Bl. ?); small tree. West Hill 2500 feet. (C. 366).

Castanopsis javanica, A.D.C.; large tree. Government Hill. 1000 feet. (C. 419).

Castanopsis sumatrana, A.D.C.; medium-sized tree. Waterfall. (C. 1563).

Castanopsis Wallichii, King; tree tall, not large. Government Hill 1000 feet, (C. 416).

Castanopsis Curtisii, King; medium-sized tree. Penara Bukit, rare; only one tree seen. (C. 1691).

SALICICNEÆ.

Salix tetrasperma, Rox.; tree 20-30 feet; not uncommon in damp places. (C. 698).

GNETACEÆ.

Gnetum Gnemon, L. "Buah Maningo"; erect pyramidal tree 30-40 feet, cultivated for its fruit. (C. 856),

Gnetum Brunonianum, Griff.; erect shrub 3-5 feet; fruit smaller than *G. Gnemon*. (C. 878).

Gnetum neglectum, Bl.; extensive climber completely covering low trees; common. (C. 877).

Gnetum macrostachyum, Hook. fil; large climber; stem 2-3 inches diam. Government Hill 2000-2500 feet. (C. 1109).

CONIFERÆ.

Dacrydium elatum, Wall.; large tree. Government Hill; common. (C.)

Podocarpus neriifolia, Don; tree 30 feet or more Government Hill. (C.)

Podocarpus cupressina, Br.; small graceful tree (C.).

Agathis loranthifolia, Salis.; tree 60-100 feet with a straight clean stem, common. (C.).

HYDROCHARIDÆ.

Blyxa malavana, Ridl.; aquatic herb. Telok Bahang, common. (C. 1888).

Ottelia alismoides, Pers.; aquatic herb. Common in ditches near the town. (C. 851).

BURMANNIACÆ.

Burmannia cælestis, Don. Not uncommon in open grassy places. (C.).

ORCHIDÆ.

Microstylis congesta, Reich. f. Pulau Betong 500 feet, rare. (C.).

Microstylis Maingayii, Hook. fil. Bukit Laksamana 1800 feet; not common. (C. 1005).

Oberonia anceps, Lindl. Waterfall. (C.).

Liparis elegans, Lindl. Moniot's Road on rocks, not uncommon. (C. 507).

Liparis longipes, Lindl. (C.).

Dendrobium pumilum, Rox. Near the Coast. (C.).

Dendrobium lonchophyllum, Hook. fil. Government Hill 2,000 feet. (C.).

Dendrobium serra, Lindl. Collected by Porter and Maingay.

Dendrobium grande, Hook. fil. A figure in Kew Herbarium is the only record for this plant here.

Dendrobium Leonis, Rchb. fil. On mangosteen trees. (C.).

Dendrobium sinuatum, Lindl. Obtained by Maingay.

Dendrobium subulatum, Hook. fil. Western Road, growing on the shade trees. (C. 2408).

Dendrobium villosulum, Wall. Collected by Porter.

Dendrobium crumenatum, Swartz. Pigeon-orchid. Common on coco-nut trees; the most abundant of any Orchid. (C.).

Dendrobium secundum, Wall. Not common. (C.).

Dendrobium sanguinolentum, Lindl. Government Hill 2000 feet. (C.).

Dendrobium bifarium, Lindl. Government Hill 2000 feet. (C.).

Bulbophyllum pileatum, Lindl. Government Hill. (C. 1141).

Bulbophyllum Medusæ, Rchb. fil, Government Hill. (C.).

Bulbophyllum capitatum, Lindl. Government Hill on *Dacrydium elatum*, not uncommon. (C. 2883).

Bulbophyllum leptosepalum, Hook. fil. Government Hill. (C. 415).

Bulbophyllum bisetosum, Lindl. Telok Tikus, rare. (C. 1979).

Cirrhopetalum vaginatum, Lindl. West Hill. (C. 355, 1777).

Cirrhopetalum concinnum, Hook. fil. var. Top of Government Hill.

Cirrhopetalum longescapum Teysm. Penang, Lobb. The specimen in the Buitenzorg Herbarium consists only of a portion of a stout scape without flowers or leaves. The plant has not since been seen.

Eria Maingayii, Hook. fil. Government Hill. Maingay.

Eria nutans, Lindl. Government Hill 2000 feet. (C.).

Eria neglecta, Ridl. Government Hill. (C.).

Eria pulchella, Lindl. On rocks near the coast. (C. 1735).

Eria pellicipes, Rchb. fil. Maingay.

Eria ferox, Bl. Lobb. Maingay.

Eria oligantha, Hook. fil. Government Hill.

Eria velutina, G. Loddiges, Government Hill. (C. 1696).

Eria leiophylla, Lindl. Government Hill. (C. 1,139, 1,240).

Claderia viridiflora, Hook. West Hill. (Government Hill 2,000 feet).

Spathoglottis plicata, Bl. Common from sea level up to 2,000 feet. (G. 72).

Tainia penangiana, Hook. fil. Government Hill. (C.).

Tainia Maingayii, Hook. fil. Maingay.

Agrostophyllum majus, Hook. fil. Government Hill 2,000-2,500 feet, on tall trees. (C.).

Cælogyne macrobulbon, Hook. fil. Wallich 1969-2.

Cælogyne tomentosa, Lindl. Government Hill. (C.).

Cælogyne tomentosa var. *penangensis*. Maingay.

Cælogyne Cumingi, Lindl. West Hill. (C.).

Cælogyne speciosa, Lindl. Government Hill 2,000 feet. (C.).

Calanthe curculigoides, Wall. Government and West Hill 2,000-2,500 feet, common. (C.).

Eulophia squalida, Lindl. Government Hill. (C. 1064).

Cymbidium Finlaysonianum, Lindl. Common on trees and rocks on the sea coast. (C. 353).

Geodorum purpureum, var. Telok Bahang, rare. (C. 354).

Geodorum citrinum, Jackson. Drawing in Kew Herbarium.

Grammatohyllum speciosum, Bl. Not common. (C.).

Bromheadia palustris, Lindl. Common on sloping banks of red soil up to 2,500 feet. (C. 359).

Polystachya penangensis, Ridl. Government Hill 1,800 feet (C. 1006).

Luisia tristis, Hook. fil. Penara Bukit, on Durian trees. (C. 1176).

Sarcochilus trichoglottis, Hook. fil. Orchards, on Mangosteen trees. (C.).

Sarcochilus (Cuculla) lilacinus, Griff. Telok Bahang, among long grass. (C.).

Sarcochilus (Cuculla) brachystachys, Hook. fil. Maingay.

Sarcochilus (Cuculla) arachnites, Reich. fil. Telok Bahang, on trees by the river. (C.).

Ærides suavissimum, Lindl. A very rare plant in Penang, I have collected it on two occasions only. (C.).

Renanthera elongata, Lindl. Ayer Hitam, rare. (C. 485).

(*Saccolabium penangianum*, Hook. fil.; is not a Penang plant. Sungkei River where my specimens were collected is in Perak. Fl. B. I. Vol. VI p. 57.)

Saccolabium tenuicaule, Hook. fil. Waterfall and West Hill. (C. 994).

Saccolabium cornigerum, Ridl. ined. West Hill. (C.).

Acampe penangiana, Ridl. ined. Government Hill. (C. 1963).

Sarcanthus secundus, Griff. Near the coast. (C.).

Sarcanthus Scortechinii, Hook. fil. Coast. (C. 2310).

Cleisostoma spicatum, Lindl. Tanjong Bunga, common. (C. 1834).

Teniophyllum serrula, Hook. fil. Western Road, on shade trees. (C.).

Acriopsis indica, Wight, was collected by Maingay.

Acriopsis javanica, Reinw. Common all over the Island. (C. 2472).

Podochilus acicularis, Hook. fil. Government Hill 2,000-2,500 feet, on damp rocks, common. (C. 1733).

Appendicula callosa, Bl. Government Hill. (C. 358).

Appendicula Lewisii, Griff. West Hill 2,000 feet (C. 993).
Appendicula Maingayii, Hook. fil. Government Hill 2,000 feet. (C. 2822).

Oxyanthera decurva, Hook. fil. Not seen. Government Hill, Maingay.

Galeola altissima, Rech. fil. Not common. (C. 352).

Galeola Hydra, Rech. fil. (C. 351).

Vanilla Griffithii, Rechb. fil. (*V. albida* Hook. fil. non Bl.)
 Waterfall, not uncommon. (C. 2271).

Corymbis longiflora, Hook. fil. Balik Pulau, rare. (C.).

Anæctochilus brevilabris, Lindl. Telok Bahang. (C.).

Anæctochilus Reinwardtii, Bl. Government Hill. (C. 1063).

Hetæria obliqua, Bl. Pulau Betong. (C.).

Odontochilus calcaratus, Hook. fil. Government Hill 2,000 feet. (C. 2823).

Hæmaria discolor, Bl. Wallich No. 7390 (C.).

Zeuxine affinis, Benth. Government Hill.

Cryptostylis arachnites, Bl. Government Hill 2,500 feet. (C.).

Pogonia punctata, Bl. Waterfall, rare. (C.)

Habenaria lacertifera, var. *robusta*, Abundant. (C. 357).

Cypripedium barbatum, Lindl. Government Hill. (C.)

Apostasia Wallichii, Br. West Hill 2,500 feet. (C. 925).

Neuwiedia Lindleyii, Rolf. Sungei Penang. (C. 469)

Neuwiedia Curtisii, Rolf. Government Hill, rare. (C. 1186).

SCITAMINEÆ.

Globba Wallichii, Baker; herb. Government Hill. (C. 455).

Globba uliginosa, Miquel. Government Hill. King.

Globba pendula, Rox. Roxburgh.

Globba versicolor, Smith, (Fl. Brit. Ind.).

Globba, sps. undetermined. (C. 956-2851).

Curcuma petiolata, Roscoe, bracts violet coloured. Common. (C. 1744).

Gastrochilus pulcherrima, Wall. Maingay.

Kæmpferia parvula, King. Government Hill 2,000 feet. Common. (C.).

Kæmpferia rotunda, L. Government Hill. (C. 189.)

- Amomum sphærocephalum*, Baker. Collected by Maingay.
Amomum megalocheilos, Baker. Cooly lines Government Hill. (C. 2419).
Amomum metrocheilos, Baker. Government Hill. (C. 1530).
Amomum biflorum, Jack. No specimen of this is known to exist. Can it be an *Elettariopsis*?
Zingiber gracile, Jack. Waterfall. Not common. (C.).
Zingiber spectabile, Griff. (C. 1200, 1978, 2161).
Costus speciosus, Smith. Common. (C.).
Costus Kingii, Baker. Pulau Betong. (C. 1976).
Elettariopsis Curtisii, Baker. West Hill 2,500 feet. (C. 1570).
Elettariopsis serpentina, Baker. Waterfall. (C. 2276).
Alpinia galanga, Sw. Penara Bukit. (C.).
Alpinia Wrayii, King. Not identified.
Alpinia mutica, Rox. Sent to the Calcutta Gardens from Penang.
Alpinia Rafflesiana, Wall. Not seen.
Phrynium parviflorum, Rox. Government Hill. (C. 2420).
Canna indica, L. Common. (C.).
Canna indica, var., *flava*. Not common (C.).
Undetermined *Scitamineæ*. (C. 327, 2275, 2884).

HÆMODORACEÆ.

- Peliosanthes Teta*, Andr.; dwarf herb, seeds blue. Waterfall, common. (C. 139).
Peliosanthes humilis, Andrews. Maingay.
Peliosanthes albida, Baker; leaves long; fl. greenish. Government Hill 2,000 feet. (C. 753).
Peliosanthes, sp.; leaves long, fl. purple. (C. 2757).
Ophiopogon prolifera, Lindl. T. Lewis.
Ophiopogon sp. West Hill. Not *prolifera*. (C.).

AMARYLLIDÆÆ.

- Curculigo latifolia*, Dryand. Common in dense jungle. (C.)
Crinum asiaticum, L. Common on the sea coast. (C. 1274).

TACCACEÆ.

Tacca cristata, Jack. Common at low elevations. (C. 335).

DIOSCOREACEÆ.

Dioscorea demonum, Rox. Climber. Muka Head. (C. 2415).

Dioscorea orbiculata, Hook. Government Hill. (C. 1900?)

Dioscorea oppositifolia, L. climber. Government Hill. (C. 2232).

Dioscorea laurifolia, Wall. (C. 60. 63 ?.)

Dioscorea gibbiflora, Hook. fil. Wallich 5106 B in part.

Dioscorea bulbifera, L.? Pulau Betong. (C. 1565. 1943).

ROXBURGHACEÆ.

Stemona Curtisii, Hook. fil.; twining, fl. pink. Waterfall. (C. 1522).

LILIACEÆ.

Smilax myosotiflora, A. De C. Porter.

Smilax lævis, Wall.; climbing shrub. Government Hill. (C. 1172).

Smilax extensa, Wall. Moniot's Road. (C. 1244).

Smilax polyacantha, Wall. Porter. Wall. Cat. 5127.

Smilax leucophylla, Bl. leaves large; glaucous beneath. Government Hill 1,000-2,000 feet. (C. 1145).

Dracæna brachystachys, Hook. fil.; small tree 10-12 feet high, 2-3 in. diam.; fl. white. (C. 2302).

Dracæna Porteri, Baker. Waterfall. (C. C.).

Dracæna terniflora, var. *Curtisii*, Hook. fil. Pulau Betong. (C. 901).

Dracæna gracilis, Wall.; small shrub. West Hill 2,500 feet. (C. 1186).

Dracæna Cantleyi, Baker; a large tree; fruit bright red the size of a cherry. Not common. (C.).

Dracæna, sp.; shrubby; panicle branched, 6-10 in long. (C. 2269).

Dracæna Jackiana, Wall cat 5145. A. B. Collected by Porter.

Dianella ensifolia, Redoute. West Hill 2,000 feet. (C. 1701).

Gloriosa superba, L. Not common in Penang. (C.).

PONTEDERiaceÆ.

Monochoria hastifolia Presl; herb. Ditches. (C.)

Monochoria vaginalis, Presl; fl. blue. Common. (C. 1235).

XYRIDeÆ.

Xyris indica, L. "Jerangoo Padang." Paddy fields south of the Island. (C. 1927).

COMMELINACEÆ.

Commelina nudiflora, L.; creeping herb; fl. blue. (C. 1224).

Commelina benghalensis. L.; Common. (C. 1843).

Aneilema nudiflorum, Bl.; small herb; fl blue; common in open damp places. (C. 1856).

Aneilema conspicuum, Kunth. Damp dense jungle. (C. 980, 1977).

Forrestia marginata, Hassk.; herb; stem creeping, rooting from the nodes. Pulau Betong. (C. 1948).

Cyanotis barbata, Don. Wallich.

Floscopa scandens, Lour. prostrate herb. Waterfall. (C.).

FLAGELLARIEÆ.

Flagellaria indica, L. "Rotan binni" Common. (C. 25).

Susum anthelminticum, Bl. Government Hill 2,000 feet. (C. 335).

PALMEÆ.

Areca catechu, L. "Pinang." Cultivated and common on abandoned land. (C.).

Pinanga polymorpha, Becc.; stem 2-3 feet. (C.).

Pinanga disticha, Bl.; stem 2-6 feet. Pantie Achie. (C. 391).

Pinanga subruminata, Becc. about 1,500-2,000 King.

Pinanga malaiana, Scheff.; stem 8-12 feet. (C.).

Oncosperma horrida, (Griff) "Nibong." Common. (C.).

Iguanura Wallichiana, Hook. fil. Collected by Porter.

Arenga saccharifera, Labill. Common. (C.).

Arenga obtusifolia, Mart. Waterfall Hill.

Caryota mitis, Lour.; stem 15-25 feet. One of the commonest palms in the Island. (C. 2149).

- Nipa fruticans*, Wurm. Common in tidal swamps. (C.)
Phoenix paludosa, Rox. ; stem 10-25 feet. Swamps. (C.)
Licuala spinosa, Wurm. (C.)
Licuala acutifida, Mart. Common. (C.)
Livistona cochinchinensis, Mart. (Lewis,)
Calamus Draco, Griff. Penang. Lewis.
Calamus javensis, Bl. West 1 ill. (C. 2268).
Calamus javensis subvar. *purpurascens*, Becc. Not identified.
Calamus javensis subvar. *penangiana*, Becc. Not identified.
Calamus melanacanthus, Mart. Penang according to Martius.
Calamus martianus, Becc. Penang, Gaudichaud.
Calamus verticillaris, Griff. (C. 2222. ? 2232. ?)
Calamus hystrix, Griff. Government Hill.
Calamus geniculatus, Griff. (C. 712, 1476. ?)
Calamus riminalis, Willd. Wallich.
Calamus Lewisianus, Griff. "Kichum" Penang. Lewis.
Calamus monticolus, Griff. Penang Hill half way up. Lewis.
Zalacca affinis, Griff. ; leaves 12-15 feet ; fruit. pear shaped densely bristly. (C. 2435).
Zalacca glabrescens, Griff. ; leaves 15-20 feet. Government Hill in damp shady jungle. (C.)
Zalacca edulis, Reinw. "Salak Kumbang" sent by Lewis to Griffith. (I do not think this is wild anywhere in the Malay Peninsula).
Zalacca Wallichiana, Griff. ; is said to occur in Penang. Not seen.
Zalacca conferta, Griff. Not seen. Lewis sent it to Griffith under the name of Asam Kumbang.
Plectocomia elongata, Bl. ; an immense climber. Government Hill 2,000-2,500 feet. (C. 2436).
Engessonia triste, Griff. ; " Bertam ; " stemless ; leaves 15-20 feet. The most common of all the palms. (C. 2218).
- PANDANÆ.
- Pandanus*, sp. ; stem 3-5 feet high ; fruit about 2½ in. (C. 1821).
Pandanus sp. several undetermined.

Pandanus helicopus, Miq. 6-10 feet high; fruit 4-5 in long. West Hill 2,500 feet. (C. 2272).

Freycinetia angustifolia, Bl.; stem slender. Court. (C. 1171).

Freycinetia scandens, Gaud.? Government Hill. Resembles *F. insignis* but has narrower leaves and bracts not red.

ARACEÆ.

Cryptocoryne ciliata, Fischer, Common in tidal swamps. (C. 1940).

Arisæma cuspidatum? Waterfall. (C. 1882).

Arisæma filiforme, Bl. Waterfall, not common. (C.)

Arisæma Kunstleri, Hook. fil.; spathe green. Waterfall. (C. 2887).

Arisæma Scortechinii, Hook. fil. Government Hill 2,000 feet. (C. 143).

Arisæma Roxburghii, Kunth is said to have been collected by Curtis on Government Hill in Flor. Brit. Ind.

Typhonium Roxburghii, Schott. Common in open spots. (C. 1858).

Typhonium Motleyanum, Schott. Pulau Tikus. (C. 2888).

Amorphophallus campanulatus, Bl. Common. (C.)

Amorphophallus Ptainii, Hook. fil. Waterfall. (C.)

Pothos Curtisii, King. Batu Feringgi. (C. 808).

Pothos, sp. West Hill. Only one specimen collected. (C.)

Lasia heterophylla, Endl. Tidal swamps, common. (C. 1941).

Homalomena angustifolium, Hook. fil. Telok Bahang on stones in mid-stream, common. (C. 2886).

Homalomena ovatum, Hook. fil. Wallich Waterfall. (C. 1884).

Homalomena humilis, Hook. fil. Government Hill.

Homalomena sagittifolia, Jung. Waterfall, common. (C. 1857).

Homalomena aromatica, Schott. Moniot's Road.

Homalomena Miqueliana, Schott; stem 2-3 ft. high. (C. 1881).

Homalomena obliquata, Hook. fil. Collected by Phillips.

Schismatoglottis longipes, Miq. Waterfall, damp ravines (2828).

Amydrium humile, Schott. Government Hill 2,500 ft. (C. 1855).

Raphidophora, sp. Batu Feringgi. (C. 1923).

Epipremnum giganteum, Schott. Collected by Roxburgh.

Epipremnum Maingayii, Pulau Betong. (C. 2885).

Epipremnum humile, Scott. Government Hill 2,500 feet. (C. 1855).

Epipremnum mirabile, Scott. Penang. Roxburgh not seen.

Anadendrum montanum, Scott. Collected by Wallich.

Anadendrum marginatum, Schott. collected by Porter.

Aglaonema nitidum, Kth. was collected here by Jack.

Aglaonema simplex, Bl. (C. 1721).

Colocasia antiquorum, L. Pulau Betong (C. 1933).

ERIOCAULONEÆ.

Eriocaulon Wallichianum, Common.

CYPERACEÆ.

Cyperus polystachyus, Rottbl. Waterfall. (C. 1784, 1864).
var *laxiflorus*, Benth. (C. 1862)

Cyperus pumilus, L. Dato Kramat. (C. 1782).

Cyperus cuspidatus var. *angustifolia*. (C. 1831).

Cyperus compressus, L. Common. (C. 1779).

Cyperus Haspan, L. Very common. (C. 1788).

Cyperus pulcherrimum Wight. Common in Rice fields.

Cyperus tria, L. (C. 1780, 1871, 1952).

Cyperus distans, L. common anywhere. (C. 875, 1785, 1832).

Cyperus rotundus, L. moderately common. (C. 1781, 1953).

Cyperus pilosus, Vahl. Tanjong Bunga. (C. 1830).

Cyperus procerus, Rottbl. Waterfall. (C. 1787).

Cyperus auricomus, Sieb. Ditches; not common. (C. 872).

Cyperus pennatus, Lour. Common. (C. 104).

Cyperus umbellatus, Benth. Waterfall, not common. (C. 491, 1783).

Cyperus turgidulus, Clarke. Tanjong Bunga. (C. 883).

Cyperus Griffithii, Steud. Open places in pure sand. (C. 884).

Cyperus Zollingerii, Steud. Waterfall, rare. (C. 1833).

- Cyperus radiatus*, Vahl. Bagian Jermal. (C. 1956).
Cyperus, sp. may be *flavidus*. Waterfall. (C. 1863).
Cyperus, sp. in way of *turgidulus*, Clarke. (C. 2173).
Mariscus Dregianus, Kunth. Tanjong Tokong. (C.).
Mariscus sieberianus, Nees. (C. 1955).
Mariscus cyperinus, Vahl. (C. 1786).
Mariscus microcephalus, Presl. Waterfall, common. (C. 1873, 1874).
Kyllinga brevifolia, Rottbl. Common all over the Island. (C. 1808).
Kyllinga monocephala, Rottbl. (C. 1823).
Fimbristylis nutans, Vahl. (C. 1868, 1869).
Fimbristylis schænoides, Vahl. (C. 1822).
Fimbristylis miliacca, Vahl. (C. 1792).
Fimbristylis asperrima, Bæck. (C. 1595, 1793).
Fimbristylis filiformis, Kunth. (C. 1789).
Fimbristylis diphylla, Kunth. (C. 1790).
Fimbristylis globulosa, Vahl. (C. 1791).
Bulbostylis barbata, Kunth. (C. 885).
Scirpus grossus, L. (C. 350).
Scirpus debilis, Pursh. (C. 1904, 2178).
Scirpus mucronatus, L. (C. 1962).
Eleocharis equisetina, Presl. (C. 1905, 2176).
Eleocharis variegata var. *laxiflora*. (C. 1865, 2177).
Fuirena glomerata, Lam. (C. 497).
Lipocarpa argentea, R. Br. (C. 1807).
Hypolytrum giganteum, Wall. (C. 15. 490).
Mapania, sp. (C. 1820).
Remirea maritima, Aubl. (C. 1861).
Rhynchospora aurea, Vahl. (C. 1256).
Scleria lithosperma, Willd. (C. 1794).
Scleria hebecarpa, Nees. (C. 1828).
Scleria sumatrensis, Retz. (C. 22).
Scleria Steudeliana, Bæck. (C. 1907).
Carex indica, L. (C. 1207).
Carex cryptostachys, Brongn. (C. 1910).

GRAMINEÆ.

- Paspalum scrobiculatum*, L. (C. 492).
Paspalum conjugatum, Berg. (C. 493).
Paspalum distichum, Burm. (C. 1957).
Isachne pulchella, Roth. (C. 496).
Isachne australis, Br. (C. 1813).
Panicum sanguinale var. *australe*, (C. 1801, 1914).
Panicum parvulum, Trin. (C. 1919).
Panicum colonum, L. (C. 68).
Panicum myosuroides, R. Br. (C. 1960).
Panicum Crus-galli, L. (C. 1958).
Panicum indicum, L. (C. 1802).
Panicum miliare, Lam. (C. 1803).
Panicum cimicinum, Retz. (C. 1829).
Panicum incomptum, Trin. (C. 132, 1809, 1908).
Panicum nodosum, Kunth. (C. 1917, 1918).
Panicum auritum, Presl. (C. 1818).
Panicum trigonum, Retz. (C. 1824).
Panicum Myurus, Lam. (C. 1903).
Panicum luzonense, Presl. (C. 1915).
Panicum multinode F. (C. 1961).
Panicum radicans, Retz. (C. 1916).
Panicum parvulum, Nees. (C. 1814).
Panicum fimbriatum, Nees. (C. 1257).
Oplismenus compositus, R. and S.
Oplismenus Burmanni, Beauv. (C. 1825).
Setaria glauca, Beauv. (C. 1259, 1959).
Leptaspis urceolata, Br. and Benn. (C. 1758).
Coix lachryma, Fobi; L. (C. 58).
Leersia hexandra, Swartz. (C. 1902).
Perotis latifolia, Ait. (C. 9, 1817).
Zoysia pungens, Willd. (C. 1812).
Imperata arundinacea, Cyr. "Lalang" (C. 1816).
Pogonatherum polystachyum, R. and S. (C. 877).
Ischæmum muticum, L. (C. 1810).
Ischæmum ciliare, Retz. (C. 1806).
Ischæmum timorense, Kunth. (C. 2167).
Rottboellia glandulosa, Trin. (C. 1913).

- Andropogon contortus*, L. (C. 1906, 1912).
Andropogon nardus, L. (C. 91).
Andropogon squarrosus, L. (C. 1819).
Chrysopogon aciculare, Trin. (C. 1799).
Themeda gigantea var. *villosa*. (C. 1971).
Themeda arguens, Hack. (C. 1911);
Sporobolus elongatus, R. Br. (C. 876).
Sporobolus diander, Beauv. (C. 1826).
Cynodon dactylon, Pers. (C. 1811).
Eleusine indica, Gærtn. (C. 1796).
Eleusine coracana, Gærtn. (C. 1118).
Eleusine ægyptiaca, Pers. (C. 1797).
Leptochloa chinensis, Nees. (C. 1951).
Phragmites Roxburghii, Kunth. (C. 125).
Eragrostis unioloides, R. and S. (C. 1800).
Eragrostis Brownii, Kunth. (C. 1805).
Eragrostis Wightiana, Bl. (C. 882).
Eragrostis plumosa, Retz. (C. 1827, 2169).
Eragrostis pilosa, Beauv. (C. 1804).
Cenotheca lappacea, Desv. (C. 36, 1815).
Thysanolæna acarifera, Nees. Government Hill. (C. 1909).
Arundo donax, cultivated. (C. 1981).
Bambusa nana, Roxb. The Hedge bamboo is cultivated.
(C. 1720).

FILICES.

- Gleichenia longissima*, Bl.; fronds 8-10 feet long. Government Hill 2,000-2,500 feet. (C. 531).
Gleichenia flagellaris, Spreng. "Resam." Common up to 1,000-1,500 feet. (C. 533).
Gleichenia hirta, Bl. Government Hill, Hullett.
Gleichenia dichotoma, Willd. Abundant. (C. 532).
Gleichenia, sp.; appears to be distinct from either of the preceding (C. 534).
Cyathea Brunonis, Wall.; stem 3-4 feet; frond 2-3 feet. Government Hill 2,000-2,500 feet. (C. 535).
Amphicosmia alterans, Hook. fil., collected by Wallich, Sir William Norris and Cantley.

- Alsophila latebrosa*, Hook. fil. ; stem tall. Government Hill. (C.)
- Alsophila glauca*, Smith ; stem stout. Not uncommon. (C.).
- Alsophila comosa*, Hook. Government Hill. (537).
- Alsophila glabra*, Hook. Government Hill. (538).
- Cibotium Barometz*, Link. Government at 1,000 (Cantley).
- Hymenophyllum tenellum*, Klein. On damp rocks at 1,500-2,000 feet. (C. 540).
- Hymenophyllum rarum*, Br. Government Hill. (Bishop Hose).
- Hymenophyllum polyanthos*, Sw. var. *Blumeanum* ; fronds 12-18 in. long. Damp shady ravines (C. 1724).
- Hymenophyllum australe*, Willd. (*Javanicum Spreng*). Government Hill. (C. 539). var. *Badium*. Government Hill (Hullett).
- Hymenophyllum Smithii*, Hook. West Hill 2,500 feet. (C.)
- Hymenophyllum aculeatum*, V. D. B. at 3,000 feet. King.
- Trichomanes digitatum*, Sw. West Hill 2,000 feet. (C. 1174).
- Trichomanes bipunctatum*, Poir. Government Hill, on rocks (C. 542).
- Trichomanes javanicum*, Bl. West Hill, damp ravines. (C. 541).
- Trichomanes radicans* var. *Kunzeanum* Government Hill. (Hullett).
- Trichomanes pallidum*, Bl. Not seen. Collected by Lady Dalhousie.
- Humata heterophylla*, Smith. Not seen.
- H. angustata*, Wall. West Hill, on trees. (C. 543).
- H. pedata*, Smith. Abundant in many places. (C. 544).
- Leucostegia hymenophylla*, Not identified.
- Leucostegia affinis*, Hook. Ayer Hitam. (C. 545).
- Prosaptia Emersonii*, Hook. fil. Moniot's Road, not uncommon. (C. 546).
- Davallia solida*, Sw. ; caudex stout ; fronds large. Common on rocks at low elevations. (C. 547).
- Davallia elegans*, Sw. Penang Hill.
- Davallia Lorrainii*, Hance ; fronds 6-8 in. Government Hill 2,000-2,500 feet. (C.).

Microlepia pinnata, Cav.; fronds 12-18 in. Government Hill 2,000-2,500 feet, abundant (C. 548).

Microlepia speluncæ L. (C. 629).

Stenoloma chinensis, Sw. var. *Veitchii*. Government Hill 1,500-2,000 feet. (C. 549).

Lindsaya cultrata, Sw.; fronds 6-7 in. Waterfall, rare (C. 1710).

Lindsaya scandens, Hook. Government Hill. (C. 550).

Lindsaya flabellulata, Dry.; a very variable plant. (C. 553).

Lindsaya trapeziformis, Dry. Government Hill. (C. 551, 554).

Lindsaya divergens, Wall. Government Hill 2,000 feet. (C. 552).

Lindsaya lanuginosa, Wall. Not seen.

Schizoloma davallioides, Bl. Government Hill. (Hullett).

Schizoloma lobata, Poir. Government Hill.

Adiantum lunulatum, Burm. Balik Pulau. (C.).

Adiantum flabellulatum, L.; frond 8-12 in long. Abundant in one or two places. (C. 555).

Cheilanthes tenuifolia, Sw. Steep banks, Ayer Hitam. (C. 556).

Pteris longifolia, L. Not uncommon on old brick walls. (C.).

Pteris cretica, L. Government Hill.

Pteris pellucida, Presl. West Hill on boulders in mid-stream. (C. 998).

Pteris crenata, Sw. Common in hedges. (C. 557).

Pteris semipinnata, L. Batu Hitam 1,000 feet, rare. (C. 635).

Pteris Dalhousiæ, Hook. Not seen. Apparently only collected by Lady Dalhousie.

Pteris quadriaurita, Retz. Balik Pulau road. (C. 1,001).

Pteris longipinnula, Wall. Not seen.

Pteris aquilina, L. Not uncommon at 1,000-2,000 feet and occasionally near sea-level. (C.).

Campteria biaurita, L. Pulau Betong. (C. 538).

Litobrochia incisa, Thunb. Common. (C.).

- Litobrochia marginata*, Bory. Balik Pulau. (C.).
Ceratopteris thalictroides, L. Ditches and swampy places. (C. 113).
Blechnum orientale, L. Abundant, up to 2,000 feet. (C.).
Blechnum Finlaysonianum, Wall. Government Hill 2,000 feet, not common. (C. 1002).
Thamnopteris nidus, L. Common on trees and rocks, not far from the coast. (C. 559).
Thamnopteris nidus, var. *phyllitidis*.
Asplenium Griffithianum, Hook; frond simple 6-10 in long. (C. 656).
Asplenium subavenium, Hook. Collected by Mactier.
Asplenium unilaterale, Lam. Damp ravines, rare. (C. 567).
Asplenium longissimum, Bl. Penara Bukit on steep banks, abundant. (C. 561).
Asplenium Wightianum, Wall. Not common. (C. 561).
Asplenium tenerum, Forst. Not uncommon on trees, at about 2,000 feet elevation. (C. 560).
Asplenium hirtum, Kaulf. Batu Hitam, rare. (C. 564).
Asplenium macrophyllum, Sw. Common. (C. 562, 566).
Asplenium paradoxum, Bl. Not identified.
Asplenium nitidum, Sw. Not common. (C. 568).
Asplenium Mactieri, Bedd. Collected by Mactier.
Asplenium laserpitijifolium, Lam. Government Hill. (C. 563).
Asplenium bulbiferum, Forst. not seen.
Asplenium subserratum, Bl. Moniot's Road. (C. 570).
Asplenium pallidum, Bl. Penara Bukit. (C. 1268).
Diplazium porrectum, Wall. West Hill 2500 H. (C. 1000).
Diplazium tomentosum, Hook. fil. Penara Bukit. (C. 1267).
Diplazium chlorophyllum, Bak. (C.).
Diplazium Bantamense, Bl. not seen.
Diplazium speciosum, Bl. ? (C. 571-999) Government Hill.
Diplazium sylvaticum, Presl. var. *Prescottianum*; Wall. Government Hill.
Diplazium Sorzogonense, Presl. fronds 18-24 in (C. 1723).
Anisogonium cordifolium, Mett. Penara Bukit, rare. (C. 1194).

- Anisogonium esculentum*, Presl. Common in damp places (C.).
Anisogonium lineolatum, Mett. Moniot's Road. (C. 573).
Hemidictyon Finlaysonianum, Wall. Not seen.
Mesochloena polycarpa, Baker. not uncommon (C. 574).
Aspidium aculeatum, Sw. Government Hill 1000-2000 feet,
 common. (C. 575.)
Aspidium semibipinnatum, Wall. Not seen.
Aspidium variolosum, Wall. Waterfall. (C. 1608). Bishop
 Hose.
Aspidium vastum, Bl. Penang.
Aspidium Singaporianum, Baker. Waterfall. (C. 576).
Aspidium polymorpha, Baker. Waterfall. common (C. 577).
Lastrea immersa, Bl. Open places, common. (C. 580).
Lastrea calcarata, Hook. fil. Government Hill (Hullett).
Lastrea crassifolia, Bl. Government Hill. (C. 579).
Lastrea intermedia, Bl. and var. *Blumei*. (C.).
Nephrodium intermedium, Baker. Government Hill. (C.
 636).
Nephrodium unitum, R. Br. Common. (C. 583).
Nephrodium molle and the var. *procurrens*, Baker. Govern-
 ment Hill. (C. 585).
Nephrodium pennigerum, Bl. Government Hill Hullett.
Nephrodium aridum, Baker. Pulau Betong. (C. 581).
Nephrodium moulemeinense? (C. 584).
Nephrodium urophyllum, Bedd. (C. 582).
Nephrodium multilineatum, Wall. Penang Hill, Wallich.
Nephrolepis exaltata, Schott. Common. (C.).
Nephrolepis volubilis, J. Smith. (C. 587).
Nephrolepis acuta, Presl. (C.).
Oleandra neriiformis, Cav. Government Hill, rare. (C.
 1607).
Polypodium subvenosum, Baker; small plant; fronds 3-4
 in. Government Hill on damp rocks. (C. 588).
Polypodium univarse, Bak. (C.).
Polypodium decorum, Brack. Government Hill, on trees
 and rocks. (C. 589).
Dictyopteris Barberi, Hook. fil. Not seen.
Goniophlebium verrucosum, Wall. Not seen.

- Niphobolus adnascens*, Sw. Very common. (C. 590, 591).
Niphobolus penangianus, Hook; fronds 12-18 in. Government Hill, towards Ayer Hitam. (C. 592).
Pleopeltis sinuosa, Wall.; frond simple 6-9 in. Ayer Hitam, on trees. (C. 595).
Pleopeltis stenophylla, Bl.; fronds about 6 in. (C. 602).
Pleopeltis angustata, Sw. Top of Government Hill. (C. 596).
Pleopeltis nigrescens, Bl.; stipe 6-18 in.; frond 12-18 in. Damp places, on rocks. (C. 600).
Pleopeltis membranacea, Don. Government Hill. (C. 597).
Pleopeltis musæfolia, Bl.; frond 2-3 feet. West Hill at about 2,000 feet, rare. (C. 598).
Pleopeltis phymatodes, L. Sea coast, abundant. (C. 599).
Drynaria quercifolia, L. Waterfall, common. (C. 594).
Drynaria rigidula, Sw. Government Hill. (C. 593).
Drynaria palmata, Bl. Government Hill, on damp rocks. (C.).
Dipteris Horsfieldii, Br.; stipe 3-7 feet; frond 1-3 feet. West Hill 2,000-2,500 feet, abundant. (C. 632).
Gymnogramma alismæfolia, Hook. fil.; stipe 10-12 in.; frond 12-18 in. West Hill 2,000 feet. (C. 603).
Gymnogramma avenia, Baker; frond 6-18 in. Damp shady places at 1,000-2,000 feet. (C. 605).
Selliguea Feeii, Hook. fil.; stipe 4-10 in.; frond 3-5 in. long. Government Hill, common. (C. 604).
Selliguea involuta, Don. Not identified.
Drymoglossum piloselloides, Presl. Abundant. (C. 1003).
Meniscium salicifolium, Wall. Ayer Hitam. (C. 616).
Antrophyum plantagineum, Kaulf. (C. 606).
Antrophyum reticulatum, var. *parvum* of Beddome. Not identified.
Vittaria elongata, Sw. Common. (C.).
Vittaria scolopendrina, Presl.; fronds 18-24 in Government Hill, not common. (C. 608).
Tænitis blechnoides, Sw. Government Hill, common. (C. 610).

Stenochlæna Norrisii, Hook. Government Hill 2,000 feet. (C. 1606.?)

Stenochlæna palustris, L. Waterfall &c. abundant. (C. 611).

Polybotrya appendiculata, Willd. Common. (C. 612).

Gymnopteris sub-repanda, Hook. fil.; stipe of barren frond 6-9 in.; frond 6-12 in. Balik Pulau, rare. (C. 628).

Gymnopteris virens, Wall. Government Hill. (C. 618.?)

Gymnopteris flagellifera, Wall. Government Hill. (C. 615).

Gymnopteris spicatum, L. West Hill, common. (C. 613, 614).

Acrostichum aureum, L. Tidal swamps, abundant. (C.).

Photinopteris rigida, Wall. West Hill, rare. (C. 619).

Photinopteris drynarioides, Hook. fil. Government Hill near the Bungalow, not common. (C.).

Platyserium bifforme, Bl. More or less common all over the Island (C. 639).

Schizæa digitata, Sw. West Hill (C.).

Lygodium circinatum, Sw. (C. 622).

Lygodium scandens, var. *microphylla*. Br. (C. 623).

Lygodium pinnatifidum, Sw. (C. 623).

Lygodium polystachyum, Wall. Waterfall, not common (C. 625).

Angiopteris evecta, Hoffm. Not uncommon (C.).

LYCOPODIACEÆ.

Lycopodium cernuum, L. Common in Penang.

Lycopodium phlegmaria, L. collected by Wallich.

Lycopodium Dalhousieanum, Spring. collected by Lady Dalhousie.

Lycopodium nummularifolium, Bl. collected by Lady Dalhousie.

Selaginella pinangensis, Spring. collected by Gaudichaud.

Selaginella trichobasis, Baker. collected by Wallich.

Selaginella alutacea, Spring. Damp banks on Penang Hill Maingay.

Selaginella atroviridis, Spring. Wallich, Gaudichaud.

Selaginella plumosa, Bak. Wallich No. 122.

Selaginella Wallichii, Spring. Wallich, Lady Dalhousie.

Selaginella Willdenovii, Spring. (C).

Selaginella caulescens, Spring. Wallich, Gaudichaud.

Selaginella caulescens, var. *argentea*, Wallich 127. Lady Dalhousie.

Selaginella chrysocaulos, Spring. Wallich 127 bis.

Psilotum triquetrum Sw. collected by Wallich.

THE BOTANISTS OF PENANG.

The number of students of botany in Penang in past years has not been great, and of some of those that are here mentioned I can get little or no information. Some whose names appear associated with plants seem merely to have collected a few specimens and transmitted them to Europe but as they are sometimes the only authority for the occurrence of certain plants in Penang, I have thought it as well to collect what information I can about them.

WILLIAM ROXBURGH was born in 1759 and took charge of the Calcutta Gardens in 1793. He seems to have never visited Penang, but received a certain number of living plants thence which he cultivated in the Gardens, and described in the *Flora Indica* published after his death in 1820.

Some of these such as *Melia tomentosa* and *Alpinia mutica* have not since been met with in Penang, and it is very likely that they were either cultivated in Penang and sent to him as if native there, or were wrongly labelled in the Calcutta Gardens.

In 1807 WILLIAM HUNTER, of the Bengal Medical Establishment, published a paper in the Linnean Society's Transactions on Gambier, as cultivated in Penang.

NATHANIEL WALLICH was born in Copenhagen in 1786, and went to India in 1807, taking charge of the Calcutta Gardens then belonging to the East India Company in 1815. He made his first great exploring expedition into Nepal in 1820, and returning thence ill went for a voyage to Penang and Singapore and visited several other parts of the Peninsula

(1822). He remained here for five months residing in Singapore in a house called Botany Hall on what is now Mount Wallich near the New Harbour Docks. While here he met Sir STAMFORD RAFFLES and WILLIAM JACK and also G. FINLAYSON, all of whom contributed plants to his herbarium.

WILLIAM JACK accompanied RAFFLES as botanist to Bencoolen in Sumatra in December, 1818, and later went to Penang. In both of these localities he made most extensive collections and published two short papers in the Malayan Miscellanies which have been reprinted in Hooker's Botanical Magazine and later in the Indo-Malaysian Essays of this Society, JACK'S health broke down in 1822, and he died at Bencoolen as he was starting for the Cape of Good Hope to recover his health. A number of his plants were sent to WALLICH and distributed with the East Indian Company's herbarium. The rest with all his manuscripts and a large series of over two thousand drawings were on board the illfated vessel, the "Fame" which was burnt at sea on the way to England, the whole collection being destroyed. Most of the plants described by JACK from Penang and those that were sent by him to WALLICH have been re-discovered of late years, but a few have escaped recent collectors, and it is possible that some of those which were distributed by WALLICH as from Penang were really collected in Sumatra. JACK'S name has been associated with several Malayan plants including the beautiful Rubiaceous tree, *Jackia ornata*.

GEORGE FINLAYSON was a native of Thurso, who became Assistant Surgeon in the 8th Regiment of Light Dragoons and was attached to the Mission from the Bengal Government to Siam and Cochin China in 1821. He returned to Singapore in 1822 seriously ill with consumption and died shortly after reaching England.

He made good collections in the various parts of the peninsula he visited, chiefly in Siam and Cochin China, and was the first botanist to visit the Dindings where he collected *Dracæna Finlaysoniana* which grows there to this day.

Many of his plants were sent to the East India Company's Herbarium and were finally distributed when that Herbarium

was broken up. Unfortunately many specimens appear not to have been adequately labelled, and some of those cited in books as coming from Penang may have come from Cochin China. WALLICH named the rare Asclepiad *Finlaysonia obovata* after him and his name has been perpetuated by two of our commonest and most charming orchids *Bromheadia Finlaysoniana* and *Cymbidium Finlaysonianum*.

After his return to Calcutta Wallich received many plants from collectors employed in Penang and chief among these was GEORGE PORTER, whose name is associated with the common dwarf *Dracæna Dracæna Porteri*. Dr. Prain has kindly examined the Wallichian correspondence at Calcutta to find out something about this collector, and Mr. KYNNEERSLEY has also given me some additional information about him. The latter says that in August 1822 the Headmaster of the Penang Free School resigned and Mr. PORTER who was recommended by Dr. WALLICH was appointed at a salary of \$100.

The same year however a Government Botanic Garden was started near Ayer Hitam and Porter was put in charge, though a judge, Mr. LEYCESTER, was nominally the curator. He remained there till 1834, at least, when the gardens were sold by Governor Murchison for 1250 rupees. He did not apparently have a very happy time as Superintendent of the gardens for Governor IBBETSON or rather his wife made trouble because Porter did not supply enough vegetables for their table. In 1834 he sent the first plants of Patchouli to Calcutta Gardens, and this is the last I have heard of him, Dr. PRAIN thinks that he got appointed Schoolmaster again through the interests of WALLICH'S friends,

The East India Herbarium was eventually dispersed, the study set being preserved at the Linnean Society's rooms in London. WALLICH compiled and published a catalogue of it, and the plants distributed are quoted in this paper as (Wall. cat.) After this period botany seems to have faded away in Penang, and nothing was done in it for many years.

Mention must be made however of Governor W. E. PHILLIPS who sent some plants to Kew which are mentioned in the Flora of British India. Mr. KYNNEERSLEY is my authority for the fol-

lowing notes. Mr. PHILLIPS was appointed Secretary to Government in 1800, and became Collector of Customs and Land Revenues in 1805, and eventually Governor in 1820. He retired in 1824, being presented then by the inhabitants of Penang with a gold cup. He was an able man and gave special attention to the land question which he seems thoroughly to have understood. He resided at Suffolk House where Crawford visited him in 1819, and describes the place as an English gentleman's mansion and park where cloves and nutmegs in full bearing were substituted for oaks, elms, and ashes. The grounds contained two to three hundred spotted deer. It was he who started the Ayer Hitam gardens.

The next botanist who appears at Penang was a Colonel Walker who in or about 1837 collected a number of plants which were distributed to various European Museums.

In 1842 or thereabouts GEORGE GRIFFITH came as Government Botanist to Malacca. He was well known for his explorations in Assam, where he had made extensive collections. He seems never to have visited Penang but received a few plants thence from T. LEWIS, Assistant Resident Councillor, after whom he named *Appendicula Lewisii* and *Calamus Lewisianus*.

Amongst other collectors who sent plants to England about this time may be mentioned Sir WILLIAM NORRIS (after whom *Norrisia* was named) who sent specimens to Sir WILLIAM HOOKER, and Lady DALHOUSIE, who is better known for her Botanical work in the Himalayas but who sent home also a collection of plants from Penang.

WILLIAM LOBB, an Orchid collector for Messrs. VEITCH visited this region in 1845 in search of ornamental plants for cultivation and besides sending home many plants alive, made a collection of dried specimens, which have been distributed to various Herbaria. Unfortunately many of these were either not at all or wrongly localised and as he collected not only in Penang and Singapore but also in Borneo and the Philippine islands, some of the plants quoted in books as, "Penang LOBB," were really collected in the further islands of the Malay Archipelago. Among the well-

known plants which bear his name are *Aeschynanthus Lobbii* and *Dipteris Lobbiana*.

Surgeon-General Maingay resided in Malacca from 1863 to 1868 and thence visited Penang and made extensive collections of plants, in both these places as well as in Singapore. After his death in the Andaman islands in 1870, his collections were bequeathed to Kew and the plants described in the Flora of British India and other publications. Most of the species said to have been collected by him in Penang have been since rediscovered, but perhaps some of those which have not been met with again have been wrongly localised as his plant-tickets often had no localities marked on them.

The flora of Penang has probably altered but little since Wallich's time. There has not been here the extensive denudation of forests which has occurred in Singapore, but without doubt the flora of the lowlying country especially near the town has under gone much change. Large as the number of known species in so limited an area is, without doubt many more remain undiscovered and with them perhaps we shall get again most of those formerly found only by the earlier collectors. In this catalogue the earlier authorities are only quoted for species which have not been seen of late years.

H. N. R.



OCCASIONAL NOTES.

EARTHQUAKE IN THE MALAY PENINSULA.

The rarity of earthquakes in the Malay Peninsula is somewhat remarkable, as though the volcanic belt of the Archipelago is absolutely outside this region, it approaches so near that one would imagine that disturbances would constantly make themselves felt here.

From time to time tremors more or less faint have been experienced in Singapore, but no record seems to have been kept of these. It is stated by residents that a shock equalling in intensity the one which shook Singapore and a large portion of the Peninsula on the night of May 17th, 1892, was felt in 1861, but no details of this earlier occurrence were preserved. The recent shock occurred at 8.10 p.m., and at Tanglin it commenced comparatively lightly and increased rapidly in violence till the whole house was violently shaken, so that glasses and furniture rattled and doors kept banging to and fro, and then it gradually died away.

The duration of the tremors was very variously reported by observers as from six seconds to three minutes, but no one seems to have taken an accurate record. At Tanglin it seemed, as nearly as I could judge (for I did not notice its commencement), to be nearly four minutes before the vibration of the house had entirely died away, but the violent period I estimated at about a minute's duration. One observer, Mr. T. A. WANDALE, residing at Pasir Panjang, noticed two distinct oscillations, the first lasting apparently thirty seconds, the second (which was more violent) with very distinct undulations lasting for about twenty-five seconds, there being an almost complete cessation of movement for ten seconds between the two waves. In Deli (Sumatra) "the shocks were more severe and had a slow, rolling and tremulous motion culminating in a heavy

shock which occurred three times in succession." (*Straits Times*.) No distinct shocks were felt in any part of the Peninsula, nor was any sound heard during the tremors, except that of the moving timbers, glasses, etc. No damage is reported from any part of the Peninsula, nor any absolute displacement of furniture, except that at Telok Ayer (Singapore) where a lamp glass and ruler were shaken off a table upon the floor. Much damage, however, is reported at Padang Sidempuan in Tapanuli district, Sumatra, and also to a less extent in Deli, and Rantan Perapat. In Singapore much alarm was caused to the natives, who ran out of their houses, and one Chinaman was so frightened that he leaped out of a window and broke his leg. Mr. Justice GOLDNEY reports that just before the shock was felt a number of black and white robins (*Copsycus musicus*) flew into the house, and some were caught by the cat.

No tidal wave was observed on the Singapore coasts, but at Muar, a steamer lying at the wharf was moved repeatedly, and off Singapore and Johor ships and boats were rocked about.

The earthquake was felt all over Singapore, in Johor, Muar, Malacca, Jelebu, Penang, Province Wellesley and at Pekan, besides the places mentioned in Sumatra, but not in Borneo nor Java. In certain spots in Singapore nothing was noticed, such were Government Hill and Fort Canning. It is well known that earthquakes have a habit of skipping over certain places, which spots have been termed "earthquake bridges." MILNE (Earthquakes, p. 141) says: "When an elastic wave passes from one bed of rock to another of a different character a certain portion of the wave is transmitted and refracted and bridges we may conceive of as occurring where the phenomenon of total refraction occurs." It is possible that this may account for the absence of the tremors in these spots, but in some cases where nothing was noticed, the observers were walking or standing on the ground, and the shock being comparatively weak was not perceived as it was by persons in wooden houses on piles which naturally were more unstable and thus would move with the slightest shock. The tremors

were all horizontal, and in the Peninsula ran from West to East. From the nature of the vibrations it may be suggested that the shock itself was at a considerable distance from Singapore. The exact direction in which the wave ran does not appear to be quite certain, as no one seems to have attempted to settle it by experiment at the time. From Malacca the vibrations are reported to have come from the N.N.W. and this is probable as they were more violent at Deli which is N.N.W. of Malacca. At Pasir Panjang they apparently travelled S.S.W. to N.N.E. or from S.W. to N. E. At Tanglin they appeared to travel from S.W. to N.E., but from the movements of a certain door, I believe they were really from the North-West.

In Medan, on the other hand they were felt as travelling from East to West. If this is correct, the starting point of the shocks must have been somewhere between South of Deli and North of Malacca, and a volcano called Sarek Berapi is said to have been the one from which the vibrations started. In conclusion, it is, I think, worth remarking that the weather for some time before the earthquake occurred was remarkably hot and oppressive in Singapore, as unusually hot weather has in other cases of earthquake been observed as preceding the shock.

H. N R.

ON THE OCCURRENCE OF THE RARE BAT-HAWK IN JOHOR.

In December last, the Bird Collector of the Raffles Museum shot in Johor a fine specimen of the very rare Hawk, *Machærhamphus alcinus* (Westerm.).

Of the genus *Machærhamphus*, only two species are known, viz., this one and *M. Andersoni*, whose habitat is Damara Land in South-West Africa, and Madagascar.

With reference to *M. alcinus*, Mr. E. W. OATES in his "Birds of British Burmah" says:—

"The slender-billed Pern is a very rare species, about which little is known. Mr. HOUGH procured one specimen at

“ Malewoon in Tenasserim, and Captain BINGHAM informs me that he thinks he once saw it in the Thoungyeen Valley.

‘ It has been known to occur at Malacca, in Borneo, and in New Guinea.

“ This species is probably crepuscular in its habits; and if this is the case its apparent scarcity is accounted for. An allied species in Africa feeds on bats.

“ This Hawk has a remarkably narrow carinated bill, large eyes, a very wide gape and an elongated occipital crest ”

The Johor specimen is about 18 inches in length, and its general colour is dark amber brown, almost black in parts; the throat and upper breast are white.

This is the only specimen of this bird in the Raffles Museum.

H. J. K.

A LARGE BEETLE CAUGHT IN A PITCHER OF NEPENTHES.

The greater number of the insects which find their death in the pitchers of the pitcher plant (*Nepenthes*) are very small, such as ants, small cockroaches and flies, and I do not think that any insect has been found fairly entrapped as big as a beetle which I found recently in a pitcher of the beautiful *Nepenthes sanguinea* on the very summit of Gunong Ledang, commonly known as Mount Ophir. This was a female of the brown stag-beetle, *Odontolabris gazella*, 2 inches in length and 1 across the body, exclusive of the spread of its legs. It was quite dead and floating flat in the water contained in the pitcher, which was one of very large size.

THE BIRD-DROPPING SPIDER (ORNITHOS- CATOIDES) IN JOHOR.

Among the large number of curious and interesting spider in the Malayan region, few are more remarkable than the *Ornithoscatoides*, which is so coloured as to exactly mimic a piece

of bird's-dung fallen upon a leaf. This spider was described by H. O. FORBES in his "Wanderings of a Naturalist" and by the Rev. O. PICKARD CAMBRIDGE, in the same work. During a visit to Gunong Panti in Johor, I was pleased to meet with a species apparently of this genus, which was engaged in sucking the juice of a red bug, which it had captured. FORBES describes and figures his species as spinning a thin web upon a leaf to represent the watery portion of the excreta and then lying upon its back on the web to which it holds by some strong spines on the back of its legs, it waits for some incautious butterfly to alight on the supposed bird's dropping, when it immediately secures the prey. The specimen I found had left its web, a thin circular white film on the leaf of a wild plantain, to devour its capture. On comparing it with the descriptions above quoted, I find that, in the colouring of the body and several structural points, the Gunong Panti spider differs from FORBES' *Ornithoscatoides decipiens*, collected in Java and Sumatra, and it is probably a distinct species.

H. N. R.

NOTES ON GALLUS VIOLACEUS.

In the description of *Gallus violaceus* in No. 24 of this Journal page 167 3rd line from foot of page for 'shaded' read 'shafted'
 „ 168 1st line „ do. „ do.
 „ 168 8th line „ 'brown' „ 'horn.'

Two more specimens (both males) of this bird have recently come under my notice. They were in the possession of a native animal dealer in Singapore but he could not give me any definite information as to where they came from. He said he thought they came from Java but was not certain and it is far more probable that they come from further east.

H. J. K.

A MALAY LULLABY.

The following may be of interest as representing the "Baby Bunting" style of rhyme of the Malay.

This one is sung when putting children to sleep, and is generally used in Naniŋ and also in the Negri Sembilan.

It is hence called "Lagu Buai." The tune is pretty though monotonous and suits the words well from what might be termed an onomatopœic view, as it is evident that the words of the lines have no actual meaning, and I cannot discover any special origin for them.

"Chapah menggulai chapah lah sayang."

"Chapah menggulai chapah lah sayang."

"Chapah didalam kelong."

"Chapah didalam kelong."


"Má mana bapa di mana lah sayang."


"Má mana bapa di mana lah sayang."

"Má ada di pintu kelong."

"Má ada di pintu kelong."

M. L.





The Ohio State University



3 2435 05425200 2

THE OHIO STATE UNIVERSITY BOOK DEPOSITORY



D	AISLE	SECT	SHLF	SIDE	POS	ITEM	C
8	03	39	13	8	04	010	5