

SERVICIO NACIONAL DE SANIDAD, INOCUIDAD Y CALIDAD AGROALIMENTARIA

DIRECCION GENERAL DE SANIDAD VEGETAL

ПЕРЕЧЕНЬ РЕГУЛИРУЕМЫХ ВРЕДНЫХ ОРГАНИЗМОВ МЕКСИКИ, 2011 г.

<https://www.ippc.int/content/lista-de-plagas-reglamentadas-de-mexico-2011> по состоянию на 1 октября 2014 г.

| Вредный организм | Кто описал |
|--|--|
| Вирус кольцевой пятнистости акации/ <i>Acacia ringspot virus</i> | |
| <i>Acalitus gossypii</i> | Banks |
| <i>Acanthospermum australe</i> | (Loefl.) Kuntze |
| <i>Acanthospermum hispidum</i> | DeCandolle |
| <i>Acarus siro</i> | Linnaeus |
| <i>Aceria sheldoni</i> | (Erwing) |
| <i>Aceria tulipae</i> | (Keifer, 1938) |
| <i>Acidovorax avenae</i> subsp. <i>avenae</i> | (Manns, 1909) Willems et al., 1992 |
| <i>Acidovorax avenae</i> subsp. <i>cattleyae</i> | (Pavarino, 1911) Willems et al., 1992 |
| <i>Acidovorax avenae</i> subsp. <i>citrulli</i> | (Schaad et al., 1978) Willems et al., 1992 |
| <i>Acremonium maydis</i> | |
| <i>Acrobasis nuxvorella</i> | Neunzig, 1970 |
| <i>Acrobasis pyrivorella</i> | (Matsumura) |
| <i>Acrolepiopsis assectella</i> | Zeller |
| <i>Acrophialophora fuispora</i> | (Saksena, 1953) Samson, 1970 |
| <i>Acrospermum maxoni</i> | |
| <i>Aculops fuchsiae</i> | Keifer |
| <i>Aculus fockeui</i> | Nalepa y Trouessart |
| <i>Adoxophyes orana</i> | Fischer von Roeslerstamm, 1834 |
| <i>Aegilops cylindrica</i> | Host |
| <i>Aeolothrips collaris</i> | Priesner, 1919 |
| <i>Aeolothrips intermedius</i> | Bagnall, 1934 |
| <i>Aeolothrips scabiosatibia</i> | Moulton, 1930 |
| <i>Aeolothrips tenuicornis</i> | Bagnall, 1926 |
| <i>Agasphaerops nigra</i> | Horn, 1876 |
| <i>Agriotes lineatus</i> | |
| <i>Agrostemma githago</i> | Linnaeus |
| <i>Alcidodes dentipes</i> | (Olivier, 1790) |
| <i>Aleurocanthus spiniferus</i> | (Quaintance, 1903) |
| <i>Aleurocanthus woglumi</i> | |
| <i>Aleurodicus destructor</i> | Mackie, 1912 |
| <i>Aleurodicus dispersus</i> | Russell, 1965 |
| <i>Aleurothrixus floccosus</i> | (Maskell, 1896) |
| Alfalfa mosaic virus | |
| <i>Alopecurus myosuroides</i> | Huds. (1762) |
| <i>Alsophila pometaria</i> | Harris, 1841 |
| <i>Alternaria brassicicola</i> | (Schwein., 1832) Wiltshire, 1947 |
| <i>Alternaria cucumerina</i> | (Ellis y Everh., 1895) Elliott, 1917 |
| <i>Alternaria dauci</i> | (Kühn, 1855) Groves y Skolko, 1944 |

| | |
|------------------------------------|----------------------------------|
| Alternaria dianthi | Stevens y Hall, 1909 |
| Alternaria gaisen | Nagano, 1928 |
| Alternaria helianthi | (Hansf.) Tubaki y Nishih. |
| Alternaria japonica | Yoshii, 1941 |
| Alternaria macrospora | Zimmerm., 1931 |
| Alternaria mali | Roberts, 1914 |
| Alternaria radicina | Meier et al., 1922 |
| <i>Alternaria ricini</i> | (Yoshii, 1929) Hansf., 1943 |
| Alternaria stanphylioides | |
| Alternaria steviae | Ishiba et al., 1982 |
| Alternaria tenuissima | (Kunze, 1818) Wiltshire, 1933 |
| Alternaria triticina | Prasada y Prabhu, 1963 |
| Alternaria zinniae | Ellis, 1972 |
| <i>Amaranthus blitum</i> | Linnaeus, 1753 |
| <i>Amblypelta cocophaga</i> | China |
| <i>Amblypelta lutescens</i> | (Distant) |
| <i>Amblyptilia pica</i> | Walsingham, 1880 |
| American plum line pattern virus | |
| Amerrhynus ynca | Sahlberg, 1823 |
| <i>Amitermes desertorum</i> | (Desneaux) |
| <i>Ammannia baccifera</i> | Linnaeus, 1753 |
| <i>Amphitetranychus viennensis</i> | (Zacher) |
| <i>Amrasca biguttula biguttula</i> | Ishida, 1912 |
| <i>Amrineus cocofolius</i> | Flechtmann, 1994 |
| <i>Anarsia lineatella</i> | Zeller, 1839 |
| <i>Anastrepha fraterculus</i> | Wiedemann, 1830 |
| <i>Anastrepha grandis</i> | Macquart, 1846 |
| <i>Anastrepha ludens</i> | Loew, 1873 |
| <i>Anastrepha obliqua</i> | Macquart, 1835 |
| <i>Anastrepha serpentina</i> | Wiedemann, 1830 |
| <i>Anastrepha striata</i> | Schiner, 1868 |
| <i>Anastrepha suspensa</i> | Loew, 1862 |
| Andean potato latent virus | |
| Andean potato mottle virus | |
| Anemone ringspot virus | |
| <i>Anguina agrostis</i> | (Steinbuch, 1799) Filipjev, 1936 |
| <i>Anguina tritici</i> | (Steinbuch, 1799) Chitwood, 1935 |
| <i>Anomis flava</i> | Fabricius 1775 |
| <i>Anoplophora chinensis</i> | (Forster, 1771) |
| <i>Anthonomus eugenii</i> | Cano , 1894 |
| <i>Anthonomus grandis</i> | Boheman , 1843 |
| <i>Anthonomus pomorum</i> | Linnaeus |
| <i>Anthonomus signatus</i> | Say , 1831 |
| <i>Anthonomus vestitus</i> | Boheman, 1859 |
| <i>Anthoxanthum odoratum</i> | Linnaeus |
| <i>Anthoxanthum puellii</i> | Lecoq y Lamotte |
| <i>Apera spica-venti</i> | (Linnaeus) Beauv. |
| Aphanomyces euteiches | Drechsler, 1925 |

| | |
|--|--|
| <i>Aphelenchoides besseyi</i> | Christie, 1942 |
| <i>Aphelenchoides fragariae</i> | (Ritzema-Bos, 1891) Christie, 1932 |
| <i>Aphelenchoides parietinus</i> | (Bastian, 1865) Steiner, 1932 |
| <i>Aphelenchoides ritzemabosi</i> | (Schwartz, 1911) Steiner y Buhner, 1932 |
| <i>Aphelenchoides tagetae</i> | Steiner, 1941 |
| <i>Aphis nasturtii</i> | Kaltenbach, 1843 |
| <i>Apion soleatum</i> | Wagner |
| <i>Apiospora camptospora</i> | Penz. y Sacc., 1897 |
| <i>Apomyelois ceratoniae</i> | Zeller, 1839 |
| Apple chlorotic leaf spot virus | |
| Apple mosaic virus | |
| Apple ringspot disease | |
| Apple rubbery wood phytoplasma | |
| Apple stem grooving virus | |
| Apricot pseudo-chlorotic leaf spot virus | |
| Arabis mosaic virus | |
| <i>Archips argyrospilus</i> | Walker, 1863 |
| <i>Archips rosana</i> | Linnaeus |
| <i>Argyrotaenia citrana</i> | (Fernald, 1889) |
| <i>Argyrotaenia ljunghiana</i> | (Thunberg) |
| <i>Argyrotaenia velutinana</i> | Walker, 1863 |
| <i>Armillaria tabescens</i> | (Scop., 1772) Emel, 1921 |
| <i>Arracacha virus A</i> | |
| Artichoke latent virus | |
| <i>Artona catoxantha</i> | (Hampson, 1892) |
| <i>Asclepias syriaca</i> | Linnaeus |
| <i>Ascochyta cycadina</i> | Scalia |
| <i>Ascochyta hyalospora</i> | (Cooke & Ellis, 1878) Boerema et al., 1977 |
| <i>Ascochyta plumbaginicola</i> | Henn. |
| Asparagus virus 1 | |
| Asparagus virus 2 | |
| <i>Aspergillus terreus</i> | Thom, 1918 |
| <i>Asperisporium pongamiae</i> | (Syd., 1913) Deighton, 1976 |
| <i>Asphondylia ilicicola</i> | Foote, 1953 |
| <i>Aspidiotus nerii</i> | Bouché, 1833 |
| Aster yellows phytoplasma group | |
| <i>Asteromella pongamiae</i> | (Sydow y Sydow) Aa, 2002 |
| <i>Athelia rolfsii</i> | (Curzi, 1932) Tu y Kimbr., 1978 |
| <i>Athelia rolfsii</i> var. <i>delphinii</i> | (Curzi) Tu y Kimbrough |
| <i>Aulacaspis rosae</i> | (Bouché, 1833) |
| <i>Aulacaspis tegalensis</i> | (Zehntner, 1898) |
| <i>Aulacorthum solani</i> | (Kaltenbach, 1843) |
| <i>Autographa gamma</i> | (Linnaeus, 1758) |
| <i>Avena sterilis</i> | Linnaeus |
| Avocado sunblotch avsunviroid | |
| <i>Bactericera cockerelli</i> | (Sulc, 1909) |
| <i>Bactrocera carambolae</i> | Drew y Hancock, 1994 |
| <i>Bactrocera cucurbitae</i> | Coquillett, 1899 |

| | |
|--|---|
| <i>Bactrocera dorsalis</i> | (Hendel, 1912) |
| <i>Bactrocera latifrons</i> | (Hendel, 1915) |
| <i>Bactrocera minax</i> | (Enderlein, 1920) |
| <i>Bactrocera musae</i> | (Tryon, 1927) |
| <i>Bactrocera oleae</i> | Gmelin, 1790 |
| <i>Bactrocera passiflorae</i> | (Froggatt, 1911) |
| <i>Bactrocera pedestris</i> | (Bezzi, 1914) |
| <i>Bactrocera psidii</i> | Froggatt, 1899 |
| <i>Bactrocera tryoni</i> | (Froggatt, 1897) |
| <i>Bactrocera tsuneonis</i> | (Miyake, 1919) |
| <i>Bactrocera zonata</i> | (Saunders, 1841) |
| Balansia oryzae-sativae | Hashioka, 1971 |
| Banana bunchy top virus | |
| Barley yellow dwarf viruses | |
| <i>Batrachedra amydraula</i> | Meyrick, 1916 |
| Bean leafroll virus | |
| Beet curly top virus | |
| Beet western yellows virus | |
| <i>Bemisia tabaci</i> | (Gennadius, 1889) |
| <i>Bemisia tabaci biot. B</i> | (Gennadius, 1889) |
| Bidens mottle virus | |
| Blueberry leaf mottle virus | |
| <i>Boeremia foveata</i> | (Foister, 1940) Aveskamp et al., 2010 |
| <i>Boerhavia diffusa</i> | Linnaeus, 1753 |
| <i>Bondia comonana</i> | (Kearfott, 1907) |
| Botryosphaeria berengeriana f.sp. pyricola | (Nose, 1933) Koganezawa y Sakuma, 1984 |
| <i>Botryosphaeria cocogena</i> | Subileau et al., 1994 |
| <i>Botryosphaeria obtusa</i> | (Schweinitz, 1822) Shoem. |
| Botryosphaeria protearum | Denman & Crous, 2003 |
| Botryosphaeria ribis | Grossenb. y Duggar, 1911 |
| Botryotinia convoluta | (Drayton, 1937) Whetzel, 1945 |
| Botryotinia fuckeliana | (de Bary) Whetzel, 1945 |
| Botryotinia porri | (Beyma, 1927) Whetzel, 1945 |
| <i>Botryotinia ricini</i> | (Godfrey, 1919) Whetzel, 1945 |
| Botryotinia squamosa | Vienn., 1953 |
| Botrytis aclada | Fresen., 1850 |
| Botrytis paeoniae | Oudem., 1897 |
| Botrytis tulipae | (Lib., 1830) Lind, 1913 |
| <i>Brachycaudus helichrysi</i> | Kaltenbach 1843 |
| <i>Brachycerus albidentatus</i> | Gyllenhal , 1840 |
| <i>Brachystola sp.</i> | |
| <i>Bradysia sp.</i> | |
| <i>Bregmatothrips iridis</i> | Watson, 1924 |
| Bremia lactucae | Regel, 1843 |
| Brenneria rubrifaciens | (Wilson et al., 1967) Hauben et al., 1999 |
| <i>Brevipalpus californicus</i> | (Banks, 1904) |
| <i>Brevipalpus chilensis</i> | Baker, 1949 |
| <i>Brevipalpus phoenicis</i> | (Geijskes, 1936) |

| | |
|--------------------------------------|---|
| Broad bean mottle virus | |
| Broad bean wilt virus | |
| <i>Bromus sterilis</i> | Linnaeus |
| <i>Brontispa longissima</i> | Gestro, 1885 |
| <i>Buglossoides arvensis</i> | (Linnaeus) Johnston |
| Burkholderia andropogonis | (Smith, 1911) Gillis et al., 1995 |
| Burkholderia caryophylli | (Burkholder, 1942) Yabuuchi et al., 1993 |
| Burkholderia cepacia | (ex Burkholder) Yabuuchi et al., 1993 |
| Burkholderia gladioli pv. alliicola | (Burkholder 1942) comb. nov. |
| Burkholderia gladioli pv. gladioli | (Severini, 1913) Yabuuchi et al., 1993 |
| Burkholderia glumae | (Kurita y Tabei, 1967) Urakami et al., 1994 |
| <i>Busseola fusca</i> | Fuller, 1901 |
| <i>Byturus fumatus</i> | Schreiber, 1910 |
| Cacao swollen shoot virus | |
| Cacao yellow mosaic virus | |
| <i>Cacoecimorpha pronubana</i> | Hübner 1800 |
| <i>Cactoblastis cactorum</i> | Berg, 1885 |
| Cactus virus X | |
| <i>Cacyreus marshalli</i> | Butler, 1898 |
| <i>Callopistria floridensis</i> | (Guenée, 1852) |
| <i>Callosobruchus analis</i> | Fabricius |
| Calonectria kyotensis | Terashita, 1968 |
| <i>Calystegia sepium</i> | (Linnaeus) Br. |
| Candidatus Liberibacter asiaticus | Jagoueix, Bové y Garnier |
| Candidatus Liberibacter solanacearum | |
| Candidatus Phytoplasma aurantifolia | Zreik et al., 1995 |
| Candidatus Phytoplasma mali | Seemüller y Schneider, 2004 |
| Candidatus Phytoplasma prunorum | Seemüller y Schneider, 2004 |
| Candidatus Phytoplasma pyri | Seemüller y Schneider, 2004 |
| Carnation etched ring virus | |
| Carnation necrotic fleck virus | Inouye & Mitsuata, 1973 |
| <i>Carposina sasakii</i> | Matsumura, 1900 |
| <i>Carthamus lanatus</i> | Linnaeus |
| <i>Carthamus oxyacantha</i> | Bieb. |
| <i>Castniomera licus</i> | (Drury) |
| Caulotops distanti | (Reuter, 1905) |
| Celery latent virus | |
| Celery mosaic virus | |
| <i>Cephaleuros virescens</i> | Künze |
| <i>Cerataphis brasiliensis</i> | (Hempel) |
| <i>Cerataphis orchidearum</i> | (Westwood) |
| <i>Ceratitis capitata</i> | (Wiedemann, 1824) |
| <i>Ceratitis cosyra</i> | (Walker, 1849) |
| <i>Ceratitis quinaria</i> | (Bezzi, 1918) |
| <i>Ceratitis rosa</i> | Karsch, 1887 |
| <i>Ceratitis rubivora</i> | Coquillett, 1901 |
| Ceratocystis paradoxa | (Dade, 1928) Moreau, 1952 |
| Ceratorhiza decidua | (Davis, 1919) Roberts, 1999 |

| | |
|--|---|
| <i>Ceratothrips discolor</i> | (Karny, 1907) |
| <i>Cercospora amaryllidis</i> | Ellis y Everh. |
| <i>Cercospora beticola</i> | Sacc., 1876 |
| <i>Cercospora callae</i> | Peck y Clinton, 1876 |
| <i>Cercospora callicarpae</i> | Cooke |
| <i>Cercospora dendrobii</i> | Burnett |
| <i>Cercospora kikuchii</i> | (Matsumoto y Tomoy., 1925) Gardner, 1927 |
| <i>Cercospora steviae</i> | |
| <i>Ceriporiopsis carnegiae</i> | (Baxter, 1941) Gilb. y Ryvarden, 1985 |
| <i>Ceroplastes destructor</i> | Newstead, 1917 |
| <i>Ceroplastes rubens</i> | Maskell, 1893 |
| <i>Ceruraphis viburnicola</i> | (Gillette, 1909) |
| <i>Ceutorhynchus pleurostigma</i> | (Marsham) Stephens , 1829 |
| <i>Chaetanaphothrips orchidii</i> | (Moulton, 1907) |
| <i>Chalara elegans</i> | Raj y Kendrick, 1975 |
| <i>Chenopodium petiolare</i> | Kunth |
| Cherry leaf roll virus | |
| Cherry necrotic rusty mottle virus | |
| Cherry rasp leaf virus | |
| Cherry rusty mottle disease | |
| <i>Chilo partellus</i> | (Swinhoe, 1885) |
| <i>Chloroclystis rectangulata</i> | Linnaeus, 1758 |
| <i>Choanephora cucurbitarum</i> | (Berk. y Ravenel, 1875) Thaxt., 1903 |
| <i>Choristoneura rosaceana</i> | (Harris, 1841) |
| <i>Chromatomyia syngenesiae</i> | Hardy, 1849 |
| Chrysanthemum chlorotic mottle pelamoviroid | |
| Chrysanthemum flower distortion virus | |
| Chrysanthemum stunt pospiviroid | |
| Chrysanthemum virus B | |
| <i>Chrysodeixis includens</i> | (Walker, 1858) |
| <i>Chrysomyxa rhododendri</i> | (DC., 1815) De Bary, 1879 |
| <i>Chrysopogon aciculatus</i> | (Retz.) Trin. |
| <i>Cirsium arvense</i> | (Linnaeus) Scop., 1772 |
| Citrus blight disease | |
| Citrus cachexia hostuviroid | |
| Citrus exocortis pospiviroid | |
| Citrus impietratura disease | Ruggieri, 1955 |
| Citrus infectious variegation virus | |
| Citrus leprosis virus | |
| Citrus psorosis B | |
| Citrus tristeza virus | |
| Citrus vein enation disease | |
| <i>Cladosporium iridis</i> | (Fautrey y Roumeguère) Vries, 1952 |
| <i>Cladosporium tenuissimum</i> | Cooke, 1878 |
| <i>Clavibacter michiganensis</i> subsp. <i>insidiosus</i> | (McCulloch, 1925) Davis et al., 1984 |
| <i>Clavibacter michiganensis</i> subsp. <i>michiganensis</i> | (Smith, 1910) Davis et al., 1984 |
| <i>Clavibacter michiganensis</i> subsp. <i>nebraskensis</i> | (Vidaver y Mandel, 1974) Davis et al., 1984 |
| <i>Clavibacter michiganensis</i> subsp. <i>sepedonicus</i> | (Spieckermann y Kotthoff, 1914) Dye y Kemp |

| | |
|--|--|
| Clavibacter michiganensis subsp. tessellarius | (Carlson y Vidaver, 1982) Davis et al., 1984 |
| Claviceps africana | Frederickson et al., 1991 |
| Claviceps sorghi | Kulk. et al., 1976 |
| <i>Cleome viscosa</i> | Linnaeus |
| Coccomyces coronatus | (Schumacher, 1803) De Notaris, 1859 |
| <i>Coccus viridis</i> | (Green, 1889) |
| <i>Cochliobolus carbonum</i> | Nelson, 1959 |
| Cocoa necrosis virus | |
| Coconut cadang-cadang cocadviroid | |
| <i>Coelaenomenodera elaeidis</i> | Maulik |
| <i>Coelogyne pulverula</i> | Teijsm. y Binn., 1862 |
| Coffee ringspot virus | |
| Coleosporium asterum | (Dietel, 1899) Syd. y Syd., 1914 |
| Coleosporium tussilaginis | (Pers., 1801) Lév., 1849 |
| Coleosporium tussilaginis f.sp. campanulae-trachel | |
| Colletotrichum acutatum | Simmonds, 1968 |
| Colletotrichum boninense | Moriwaki et al., 2003 |
| <i>Colletotrichum capsici</i> | (Sydow & Sydow, 1913) Butler & Bisby, 1931 |
| Colletotrichum coccodes | (Wallr., 1833) Hughes, 1958 |
| Colletotrichum coffeanum var. virulans | Noack |
| Colletotrichum dematium | (Pers., 1801) Grove, 1918 |
| Colletotrichum sublineolum | Henn., 1913 |
| Colletotrichum truncatum | (Schwein., 1832) Andrus y Moore, 1935 |
| <i>Commelina benghalensis</i> | Linnaeus, 1753 |
| Coniothyrium callicarpae | Cooke |
| Coniothyrium wernsdorffiae | Laubert, 1905 |
| Conoderus falli | Lane, 1956 |
| <i>Conogethes punctiferalis</i> | (Guenée) |
| <i>Conotrachelus aguacatae</i> | Barber, 1923 |
| <i>Conotrachelus nenuphar</i> | (Herbst) Harris , 1841 |
| <i>Conotrachelus perseae</i> | Barber, 1919 |
| <i>Conringia orientalis</i> | (Linnaeus) Dumort. |
| <i>Copitarsia consueta</i> | (Walker, 1857) |
| <i>Copturus aguacatae</i> | Kissinger |
| Corynespora cassiicola | (Berk. y Curtis, 1868) Wei, 1950 |
| <i>Corythucha marmorata</i> | (Uhler, 1878) |
| Cotton anthocyanosis virus | |
| Cotton leaf curl virus | |
| <i>Crioceris asparagi</i> | (Linnaeus, 1758) |
| Crocus mosaic virus | |
| <i>Crupina vulgaris</i> | Pers. ex Cass. |
| Cryptosporella umbrina | (Jenkins, 1918) Jenkins y Wehmeyer, 1935 |
| Ctnicera pruinina | Horn, 1871 |
| Cucumber green mottle mosaic virus | |
| Cucumber leaf spot virus | |
| Cucumber mosaic virus | |
| <i>Curculio caryae</i> | Horn, 1873 |
| <i>Curculio elephas</i> | Gyllenhal, 1836 |

| | |
|--|--|
| <i>Curtobacterium flaccumfaciens</i> pv. <i>flaccumfaciens</i> | (Hedges, 1922) Collins y Jones, 1983 |
| <i>Curtobacterium flaccumfaciens</i> pv. <i>poinsettiae</i> | (Starr y Pirone, 1942) Collins y Jones, 1983 |
| <i>Curvularia</i> spp. | |
| <i>Curvularia trifolii</i> | (Kauffman, 1920) Boedijn, 1933 |
| <i>Cuscuta campestris</i> | |
| <i>Cuscuta corymbosa</i> | |
| <i>Cuscuta corymbosa</i> var. <i>stylosa</i> | |
| <i>Cuscuta europaea</i> | Linnaeus, 1753 |
| <i>Cuscuta indecora</i> | |
| <i>Cuscuta potosina</i> var. <i>globifera</i> | |
| <i>Cuscuta squamata</i> | |
| <i>Cuscuta</i> spp. | |
| <i>Cuscuta umbellata</i> | |
| <i>Cydia caryana</i> | Fitch, 1856 |
| <i>Cydia latiferreana</i> | (Walsingham, 1879) |
| <i>Cydia splendana</i> | (Hübner, 1796) Hübner, 1799 |
| <i>Cylindrocladium circumscriptum</i> | |
| <i>Cylindrocladium pteridis</i> | Wolf, 1926 |
| Cymbidium mosaic virus | |
| Cymbidium ringspot virus | |
| <i>Cynthia cardui</i> | Linnaeus, 1758 |
| <i>Dacus ciliatus</i> | Loew, 1862 |
| <i>Daedalea ambigua</i> | Berk., 1845 |
| Dahlia mosaic stunt virus | |
| Dahlia mosaic virus | |
| Dasheen mosaic virus | |
| <i>Datana major</i> | Grote & Robinson |
| <i>Davidiella macrospora</i> | (Kleb., 1924) Crous y Braun, 2003 |
| <i>Davidiella tassiana</i> | (De Notaris, 1863) Crous y Braun, 2003 |
| <i>Delia antiqua</i> | Meigen, 1826 |
| <i>Delia floralis</i> | (Fallén, 1824) |
| <i>Delia florilega</i> | Zetterstedt, 1945 |
| <i>Delia platura</i> | (Meigen, 1826) |
| Dendrobium mosaic virus | |
| <i>Dendrothrips ornatus</i> | (Jablonowski, 1894) |
| <i>Dendryphiella vinosa</i> | (Berkeley y Curtis) Reisinger, 1968 |
| <i>Deroceras</i> spp. | |
| <i>Desmella superficialis</i> | Sydow y Sydow, 1919 |
| <i>Diaphorina citri</i> | Kuwayama, 1908 |
| <i>Diaporthe phaseolorum</i> var. <i>meridionalis</i> | Fernández, 1996 |
| <i>Diaporthe phaseolorum</i> var. <i>sojae</i> | (Lehman, 1922) Wehm., 1933 |
| <i>Diaporthe vaccinii</i> | Shear, 1931 |
| <i>Diaporthe vexans</i> | Gratz, 1942 |
| <i>Diaprepes abbreviatus</i> | (Linnaeus, 1758) Hustache, 1929 |
| <i>Diaspis bromeliae</i> | (Kerner, 1778) Kerner |
| <i>Diaspis echinocacti</i> | (Bouché, 1833) |
| <i>Diatraea indigenella</i> | Dyar y Heinrich, 1927 |
| <i>Dichromothrips corbeti</i> | (Priesner, 1936) |

| | |
|-----------------------------------|---|
| Dickeya chrysanthemi | (Burkholder et al., 1953) Samson et al., 2005 |
| Dickeya dianthicola | Samson et al., 2005 |
| Dickeya paradisiaca | (Fernández y López, 1970) Samson et al., 2005 |
| Dickeya zeae | Samson et al., 2005 |
| <i>Dicladispa armigera</i> | (Olivier, 1808) |
| Didymella bryoniae | (Auersw., 1869) Rehm, 1881 |
| Didymella fuchsiae | Cooke Y Harkn. |
| Didymella holci | (Tehon, 1937) Arx, 1987 |
| <i>Didymella ligulicola</i> | (Baker et al., 1949) Arx, 1962 |
| Didymella rabiei | (Kovatschevski, 1936) Arx, 1962 |
| <i>Digitaria abyssinica</i> | (Rich.) Stapf, 1907 |
| <i>Digitaria velutina</i> | (Forssk.) Beauv. |
| <i>Dinoderus minutus</i> | (Fabricius, 1775) |
| <i>Diparopsis tephrogramma</i> | Baker, 1911 |
| <i>Diplotaxis erucoides</i> | (Linnaeus) DeCandolle |
| Diplothecha tunae | (Spreng., 1820) Starbäck, 1893 |
| <i>Ditylenchus angustus</i> | (Butler, 1913) Filipjev, 1936 |
| <i>Ditylenchus destructor</i> | Thorne, 1945 |
| <i>Ditylenchus dipsaci</i> | (Kühn, 1857) Filip'ev, 1936 |
| Dothiorella rot | Especie inexistente |
| Dothiorella spp. | |
| Dracaena mottle virus | |
| <i>Draeculacephala minerva</i> | Ball, 1927 |
| <i>Drosophila suzukii</i> | (Matsumura, 1931) |
| <i>Dysaphis tulipae</i> | Boyer de Fonscolombe, 1841 |
| <i>Dysdercus cingulatus</i> | (Fabricius, 1775) |
| <i>Dysdercus fasciatus</i> | Signoret |
| Dysmicoccus neobrevipes | Beardsley |
| <i>Earias biplaga</i> | Walker, 1866 |
| <i>Earias cupreoviridis</i> | Walker, 1862 |
| <i>Earias insulana</i> | Boisduval, 1833 |
| <i>Earias vittella</i> | (Fabricius, 1794) |
| Echidnodes bromeliacearum | (Rehm) Theiss. y Sydow, 1918 |
| <i>Echinochloa frumentacea</i> | (Roxb.) Link |
| <i>Echinochloa oryzoides</i> | (Ard.) Fritsch |
| <i>Echium vulgare</i> | Linnaeus |
| Eggplant latent elaviroid | |
| Eggplant mottled dwarf virus | |
| <i>Eldana saccharina</i> | Walker, 1865 |
| <i>Elsinoë brasiliensis</i> | Bitancourt & Jenkins, 1942 |
| <i>Elymus repens</i> | (Linnaeus) Gould, 1947 |
| Emericella nidulans | (Eidam, 1883) Vuillemin, 1927 |
| <i>Emex australis</i> | Steinh. |
| <i>Emex spinosa</i> | (Linnaeus) Campd. |
| <i>Endothenia hebesana</i> | Walker, 1863 |
| <i>Enneadesmus obtusedentatus</i> | Lesne |
| <i>Enneadesmus trispinosus</i> | (Olivier, 1795) |
| Entopeltis interrupta | (Winter, 1884) Höhnel, 1910 |

| | |
|---|---------------------------------------|
| <i>Entyloma eryngii</i> | (Corda, 1839) de Bary, 1874 |
| <i>Entyloma lobeliae</i> | Farlow, 1883 |
| <i>Entyloma veronicae</i> | (Halsted, 1890) Lagerheim, 1891 |
| <i>Epicaerus cognatus</i> | Sharp, 1891 |
| <i>Epicauta pensylvanica</i> | (De Geer, 1775) |
| <i>Epichoristodes acerbella</i> | (Walker, 1864) |
| <i>Epiphyas postvittana</i> | Walker, 1863 |
| <i>Epitrix tuberis</i> | Gentner, 1944 |
| <i>Eriophyes eucricotesi</i> | |
| <i>Eriophyes hibisci</i> | Nalepa |
| <i>Eriophyes pyri</i> | (Pagenstecher) |
| <i>Erwinia amylovora</i> pv. <i>crenatoserrata</i> | |
| <i>Erwinia chrysanthemi</i> pv. <i>chrysanthemi</i> | (Burkholder et al., 1953) Dye, 1969 |
| <i>Erwinia herbicola</i> pv. <i>gypsophilae</i> | (Brown, 1934) Miller et al., 1981 |
| <i>Erwinia</i> sp. | |
| <i>Erwinia tracheiphila</i> | (Smith, 1895) |
| <i>Erynnis lucilius</i> | Scudder y Burgess, 1870 |
| <i>Erysiphe australiana</i> | (McAlpine, 1899) Braun y Takam., 2000 |
| <i>Eudocima fullonia</i> | (Clerck, 1764) |
| <i>Eulophonotus myrmeleon</i> | Felder, 1874 |
| <i>Eumerus funeralis</i> | Meigen, 1822 |
| <i>Eupalamides cyparissias</i> | Fabricius, 1777 |
| <i>Euphorbia esula</i> | Linnaeus |
| <i>Euphorbia helioscopia</i> | Linnaeus, 1753 |
| <i>Euphranta canadensis</i> | (Loew, 1873) |
| <i>Euphydryas viburnii</i> | |
| <i>Euplatypus segnis</i> | Bright & Skidmore, 2002 |
| <i>Euplexia benesimilis</i> | McDunnough, 1922 |
| <i>Euplexia similis</i> | |
| <i>Eupoecilia ambiguella</i> | Hübner, 1796 |
| <i>Euproctis chrysorrhoea</i> | (Linnaeus, 1758) |
| <i>Euscepes postfasciatus</i> | (Fairmaire, 1849) |
| <i>Eutetranychus buxi</i> | |
| <i>Eutetranychus orientalis</i> | Klein |
| <i>Eutinobothrus brasiliensis</i> | (Hambleton, 1937) Lima, 1938 |
| <i>Euwallacea fornicatus</i> | (Eichhoff, 1868) Wood y Bright, 1992 |
| <i>Exartema ferriferanum</i> | (Walker, 1863) Clemens y Zeller |
| Fiji disease virus | |
| <i>Frankliniella cestrum</i> | Moulton |
| <i>Frankliniella intonsa</i> | (Trybom, 1895) |
| <i>Frankliniella occidentalis</i> | (Pergande, 1895) |
| <i>Frankliniella schultzei</i> | (Trybom, 1910) |
| Freesia mosaic virus | |
| <i>Fumaria officinalis</i> | Linnaeus, 1753 |
| <i>Fusarium cerealis</i> | (Cooke, 1878) Saccardo, 1886 |
| <i>Fusarium incarnatum</i> | (Roberge, 1849) Saccardo, 1886 |
| <i>Fusarium oxysporum</i> f.sp. <i>aechmeae</i> | Gordon, 1965 |
| <i>Fusarium oxysporum</i> f.sp. <i>asparagi</i> | Cohen, 1946 |

| | |
|--|---|
| <i>Fusarium oxysporum</i> f.sp. <i>cattleyae</i> | Foster |
| <i>Fusarium oxysporum</i> f.sp. <i>cepae</i> | (Hanzawa, 1914) Snyder & Hansen, 1940 |
| <i>Fusarium oxysporum</i> f.sp. <i>conglutinans</i> | (Wollenweber, 1913) Snyder & Hansen, 1940 |
| <i>Fusarium oxysporum</i> f.sp. <i>cubense</i> | (Smith, 1910) Snyder y Hansen, 1940 |
| <i>Fusarium oxysporum</i> f.sp. <i>cubense</i> raza 4 | (Smith, 1910) |
| <i>Fusarium oxysporum</i> f.sp. <i>gladioli</i> | (Massey, 1926) Snyder y Hansen, 1940 |
| <i>Fusarium oxysporum</i> f.sp. <i>melongenae</i> | Matuo y Ishigami, 1958 |
| <i>Fusarium oxysporum</i> f.sp. <i>radicis-cucumerinum</i> | Vakalounakis, 1996 |
| <i>Fusarium oxysporum</i> f.sp. <i>ricini</i> | (Wollenweber, 1940) Gordon, 1965 |
| <i>Fusarium oxysporum</i> f.sp. <i>sesami</i> | Jacz. |
| <i>Fusarium oxysporum</i> f.sp. <i>vasinfectum</i> | (Atkinson, 1892) Snyder y Hansen, 1940 |
| <i>Fusarium solani</i> f.sp. <i>cucurbitae</i> | Snyder y Hansen, 1941 |
| <i>Fusarium solani</i> f.sp. <i>pisi</i> | (Jones, 1923) Snyder y Hansen, 1941 |
| <i>Fusarium</i> sp. | |
| <i>Gaeumannomyces graminis</i> var. <i>Graminis</i> | (Saccardo) Arx y Olivier |
| <i>Galega officinalis</i> | Linnaeus |
| <i>Galeopsis tetrahit</i> | Linnaeus |
| <i>Gangara thyrsis</i> | (Fabricius, 1775) |
| <i>Ganoderma lucidum</i> | (Curtis, 1781) Karsten, 1881 |
| <i>Gastrolobium grandiflorum</i> | Muell., 1862 |
| <i>Gibberella acuminata</i> | Wollenweber |
| <i>Gibberella avenacea</i> | Cook |
| <i>Gibberella fujikuroi</i> | (Sawada, 1919) Wollenw., 1931 |
| <i>Gibberella intricans</i> | Wollenw., 1930 |
| <i>Gibberella stilboides</i> | Gordon ex Booth |
| <i>Gladiolus grassines</i> (MLO) | |
| <i>Globodera pallida</i> | (Stone, 1973) Behrens, 1975 |
| <i>Globodera rostochiensis</i> | (Wollenweber, 1923) Behrens, 1975 |
| <i>Gloeosporium convallariae</i> | Allescher, 1895 |
| <i>Glomerella tucumanensis</i> | (Spegazzini) Arx y Müller, 1954 |
| <i>Golovinomyces orontii</i> | (Castagne) Heluta |
| Grapevine enation disease | |
| Grapevine flavescence doree phytoplasma | |
| Grapevine leafroll-associated viruses | |
| Grapevine vein necrosis virus | |
| Grapevine virus A | |
| <i>Grapholita funebrana</i> | Treitschke, 1835 |
| <i>Grapholita inopinata</i> | Heinrich, 1928 |
| <i>Grapholita molesta</i> | (Busck, 1916) |
| <i>Grapholita packardi</i> | Zeller |
| <i>Grapholita prunivora</i> | (Walsh) |
| Green mosaic virus | |
| <i>Guignardia musae</i> | Raciborsky, 1909 |
| <i>Gymnaspis aechmeae</i> | Newstead, 1898 |
| <i>Gymnosporangium clavipes</i> | Cooke y Peck, 1873 |
| <i>Gymnosporangium globosum</i> | Farlow, 1880 |
| <i>Gymnosporangium juniperi-virginianae</i> | Schweinitz, 1822 |
| <i>Hadula trifolii</i> | (Hufnagel, 1766) |

| | |
|---------------------------------------|--------------------------------------|
| <i>Halticus bractatus</i> | (Say, 1832) |
| Haplobasidion musae | Ellis, 1957 |
| <i>Haplothrips kurdjumovi</i> | Karny, 1913 |
| <i>Harrisina brillians</i> | |
| <i>Heilipus lauri</i> | Boheman , 1845 |
| <i>Helicotylenchus pseudorobustus</i> | (Steiner, 1914) Golden, 1956 |
| <i>Helicoverpa armigera</i> | (Hübner, 1809) |
| <i>Helicoverpa zea</i> | (Boddie, 1850) |
| <i>Heliopsis virescens</i> | (Fabricius, 1777) |
| <i>Heliophris haemorrhoidalis</i> | (Bouché, 1833) |
| Helminthosporium solani | Durieu y Montagne, 1848 |
| <i>Hemiberlesia rapax</i> | (Comstock, 1881) |
| Hemileia coffeicola | Maublanc y Roger, 1934 |
| Hemileia vastatrix | Berkeley y Broome, 1869 |
| Henderinia cerei | |
| <i>Heracleum mantegazzianum</i> | Sommier y Levier, 1895 |
| <i>Hercinothrips bicinctus</i> | (Bagnall, 1919) |
| <i>Hercinothrips femoralis</i> | (Reuter, 1891) |
| <i>Heterodera avenae</i> | Wollenweber, 1924 |
| <i>Heterodera glycines</i> | Ichinohe, 1952 |
| <i>Heterodera oryzae</i> | Luc y Brizuela, 1961 |
| <i>Heterodera schachtii</i> | Schmidt, 1871 |
| <i>Heterodera trifolii</i> | Goffart, 1932 |
| <i>Heterodera zea</i> | Koshy et al., 1971 |
| Heterosporium fulvarosae | |
| <i>Hibiscus trionum</i> | Linnaeus |
| Hollyhock yellow-vein mosaic disease | |
| <i>Homalodisca vitripennis</i> | Germar, 1821 |
| <i>Homoeosoma heinrichi</i> | Pastrana, 1961 |
| Hop stunt hostuviroid | |
| <i>Horcias nobilellus</i> | (Berg, 1883) |
| Hydrangea ringspot virus | |
| Hydrangea virescense MLO | |
| <i>Hylemya sp.</i> | |
| <i>Hypera zoilus</i> | (Fabricius, 1763) |
| <i>Hypothenemus hampei</i> | (Ferrari, 1867) Wood y Bright , 1992 |
| <i>Icerya aegyptiaca</i> | (Douglas, 1890) |
| <i>Idiopterus nephrolepidis</i> | Davis, 1909 |
| Impatiens necrotic spot virus | |
| <i>Imperata cylindrica</i> | (Linnaeus) Raeschel, 1797 |
| Indian citrus ringspot virus | |
| <i>Ipomoea carnea</i> | Jacq. |
| <i>Ipomoea polymorpha</i> | Roem. & Schult. |
| Irenopsis cupaniae | (Stevens, 1916) Stevens, 1927 |
| Iris fulva mosaic virus | |
| Iris mild mosaic virus | |
| Iris severe mosaic virus | |
| <i>Ischaemum rugosum</i> | Salisb., 1791 |

| | |
|---|--|
| Jasmine infectious variegation disease | |
| <i>Kabatiella lini</i> | (Lafferty, 1921) Karakulin, 1957 |
| <i>Kabatiella microsticta</i> | Bubák, 1907 |
| <i>Kellermania anomala</i> | (Cooke) Höhnel, 1915 |
| <i>Khuskia oryzae</i> | Hudson, 1963 |
| <i>Lasiodiplodia theobromae</i> | (Pat., 1892) Griffiths y Maubl., 1909 |
| Lavender yellow decline (MLO) | |
| <i>Leifsonia xyli</i> subsp. <i>xyli</i> | (Davis et al., 1984) Evtushenko et al., 2000 |
| <i>Lepidosaphes ulmi</i> | (Linnaeus, 1758) |
| <i>Leptinotarsa decemlineata</i> | Say, 1824 |
| <i>Leptochloa chinensis</i> | (Linnaeus) Nees, 1824 |
| <i>Leptopharsa gibbicarina</i> | Froeschner |
| <i>Leptothyrium rosarum</i> | Cooke |
| Lettuce big-vein virus | |
| Lettuce mosaic virus | |
| <i>Ligniera vasculorum</i> | (Matz) Cook |
| <i>Ligyryus ebenus</i> | (De Geer, 1774) |
| <i>Ligyryus gibbosus</i> | (Degeer, 1774) |
| <i>Lilioceris lili</i> | (Scopoli, 1763) |
| Lily mottle virus | |
| Lily rosette virus | |
| <i>Limonius californicus</i> | Mannerheim, 1843 |
| <i>Linaria vulgaris</i> | Miller |
| <i>Liriomyza bryoniae</i> | (Kaltenbach, 1858) |
| <i>Liriomyza huidobrensis</i> | (Blanchard, 1926) |
| <i>Liriomyza pusilla</i> | (Meigen, 1830) |
| <i>Liriomyza sativae</i> | Blanchard, 1938 |
| <i>Liriomyza trifolii</i> | (Burgess in Comstock, 1880) Burgess, 1880 |
| <i>Lissachatina fulica</i> | (Bowdich, 1822) |
| <i>Listroderes costirostris</i> | Schönherr, 1826 |
| Little cherry virus | |
| <i>Lobesia botrana</i> | Denis y Schiffermüller, 1776 |
| <i>Lolium rigidum</i> | Gaudin, 1811 |
| <i>Longicaudus trirhodus</i> subsp. <i>trirhodus</i> | |
| <i>Longidorus elongatus</i> | (de Man, 1876) Micoletzky, 1922 |
| <i>Lophodermium exaridum</i> | (Cooke y Peck, 1878) Saccardo, 1883 |
| <i>Lophodermium schweinitzii</i> | Wilson & Robertson, 1947 |
| <i>Lopholeucaspis japonica</i> | (Cockerell, 1897) |
| <i>Lopidea davisii</i> | Knight, 1917 |
| <i>Ludwigia hyssopifolia</i> | (Don) Exell |
| <i>Lycium ferocissimum</i> | Miers |
| <i>Lyctus africanus</i> | Lesne |
| <i>Lymantria dispar</i> | Linnaeus, 1758 |
| <i>Maconellicoccus hirsutus</i> | (Green, 1908) |
| <i>Macronoctua onusta</i> | Grote, 1874 |
| Macrophoma pandani | (Léveillé, 1846) Berlese y Voglino, 1886 |
| <i>Macrophya simillima</i> | Rohwer, 1917 |
| <i>Macrosiphoniella artemisiae</i> subsp. <i>artemisiae</i> | (Harrington, 1985) |

| | |
|---------------------------------------|---------------------------------------|
| <i>Macrosiphoniella sanborni</i> | (Gillette, 1908) |
| <i>Macrosiphum euphorbiae</i> | Thomas, 1878 |
| <i>Malacosoma americanum</i> | Fabricius, 1793 |
| <i>Malacosoma disstria</i> | Hübner, 1819 |
| <i>Maladera castanea</i> | (Arrow, 1913) |
| Maple mosaic disease | |
| <i>Marasmia trapezalis</i> | (Guenée) |
| Marasmius palmivorus | Sharples, 1928 |
| <i>Marmara gulosa</i> | Guillén y Davis, 2001 |
| <i>Mayetiola destructor</i> | (Say, 1817) |
| <i>Megalopyge opercularis</i> | Smith y Abbot, 1797 |
| <i>Melaleuca quinquenervia</i> | (Cav.) Blake |
| Melanconium pandani | Léveillé, 1845 |
| Melanomma dothideasporum | |
| <i>Melanoplus sp.</i> | |
| <i>Melanotus communis</i> | Gyllenhal , 1817 |
| <i>Melanthrips fuscus</i> | (Stutzer, 1776) |
| <i>Melastoma malabathricum</i> | Linnaeus |
| Meliola dracaena | Stevens, 1925 |
| Meliola praetervisa | Gaillard, 1892 |
| <i>Melochia corchorifolia</i> | Linnaeus |
| <i>Meloidogyne arenaria</i> | (Neal, 1889) Chitwood, 1949 |
| <i>Meloidogyne chitwoodi</i> | Golden et al., 1980 |
| <i>Meloidogyne exigua</i> | Goeldi, 1892 |
| <i>Meloidogyne hapla</i> | Chitwood, 1949 |
| <i>Meloidogyne javanica</i> | (Treub, 1885) Chitwood, 1949 |
| <i>Meloidogyne marioni</i> | (Cornu, 1879) Chitwood y Oteifa, 1952 |
| <i>Meloidogyne spp.</i> | |
| <i>Melolontha melolontha</i> | (Linnaeus, 1758) |
| Melon necrotic spot virus | |
| <i>Merodon equestris</i> | Fabricius, 1794 |
| <i>Metcalfa pruinosa</i> | (Say, 1830) |
| Microbotryum violaceum | (Persoon) Deml y Oberwinkler, 1982 |
| <i>Microcephalothrips abdominalis</i> | (Crawford, 1910) |
| Microdochium dimerum | (Penzig, 1882) Arx, 1984 |
| <i>Microlophium carnosum</i> | (Buckton, 1876) |
| Microsphaeropsis hellebori | (Cooke y Massee, 1887) Aa, 2002 |
| <i>Mikania micrantha</i> | Kunth |
| Milesia insularis | Faull, 1932 |
| <i>Monarthropalpus buxi</i> | (Laboulbène) |
| Monilinia fructigena | (Aderh. y Ruhland) Honey, 1945 |
| Moniliophthora perniciosa | (Stahel) Aime y Mora, 2006 |
| Moniliophthora roreri | (Ciferri, 1933) Evans et al., 1978 |
| <i>Monochoria vaginalis</i> | (Burm.) Presl, 1827 |
| Monographella albescens | (Thümen, 1889) Parkinson et al., 1981 |
| <i>Mononychus vulpeculus</i> | Schoenherr , 1825 |
| Mycoleptodiscus indicus | (Sahni) Sutton, 1973 |
| Mycosphaerella anthurii | Miles, 1917 |

| | |
|--------------------------------------|---|
| <i>Mycosphaerella fijiensis</i> | Morelet, 1969 |
| <i>Mycosphaerella opuntia</i> | (Ellis y Everhart, 1888) Dearness, 1919 |
| <i>Mycosphaerella pinodes</i> | (Berk. & Bloxam) Vestergren |
| <i>Myndus crudus</i> | Van Duzee, 1907 |
| <i>Myrothecium roridum</i> | Tode, 1790 |
| <i>Mythimna unipuncta</i> | Haworth |
| <i>Myzaphis rosarum</i> | (Kaltenbach, 1843) |
| <i>Nacoleia octasema</i> | Meyrick (1886) |
| <i>Nandina mosaic virus</i> | |
| <i>Naohidemyces vacciniorum</i> | (Link) Spooner, 1999 |
| <i>Narcissus mosaic virus</i> | |
| <i>Narcissus yellow stripe virus</i> | |
| <i>Nassella trichotoma</i> | (Nees) Hackel ex Arech., 1896 |
| <i>Naupactus leucoloma</i> | Boheman , 1840 |
| <i>Naupactus xanthographus</i> | (Germar) Boheman in Schoenh. , 1833 |
| <i>Nectria magnoliae</i> | Lohman y Hepting, 1943 |
| <i>Nectriella pironii</i> | Alfieri y Samuels, 1980 |
| <i>Nemorimyza maculosa</i> | (Malloch, 1913) |
| <i>Neomyzus circumflexum</i> | (Buckton, 1876) |
| <i>Neonectria radicola</i> | (Gerlach y Nilsson) Mantiri y Samuels, 2001 |
| <i>Neoscytalidium dimidiatum</i> | (Penzig) Crous y Slippers, 2006 |
| <i>Neosilba pendula</i> | (Bezzi, 1919) |
| <i>Neslia paniculata</i> | (Linnaeus) Desv. |
| <i>Nodonota puncticollis</i> | (Say) |
| <i>Oberea bimaculata</i> | (Olivier, 1795) |
| <i>Odontoglossum ringspot virus</i> | |
| <i>Oligonychus coffeae</i> | (Nietner) |
| <i>Oligonychus ilicis</i> | |
| <i>Olpidium brassicae</i> | (Woronin, 1878) Dangeard, 1886 |
| <i>Omophlus lepturoides</i> | (Fabricius, 1787) |
| <i>Omophoita aequinoctialis</i> | (Linnaeus, 1758) |
| <i>Oncidium ringspot virus</i> | |
| <i>Onion yellow dwarf virus</i> | |
| <i>Operophtera brumata</i> | Linnaeus, 1758 |
| <i>Opogona sacchari</i> | Bojer, 1856 |
| <i>Orchamoplatus mammaeferus</i> | (Quaintancer y Baker, 1917) |
| <i>Orchidophilus aterrimus</i> | (Waterhouse) |
| <i>Orgyia leucostigma</i> | (Smith, 1797) |
| <i>Orgyia postica</i> | (Walker, 1855) |
| <i>Orobanche cernua</i> | Loefl., 1758 |
| <i>Orobanche ramosa</i> | Linnaeus (1753) |
| <i>Orobanche spp.</i> | |
| <i>Orseolia oryzivora</i> | Harris y Gagné, 1982 |
| <i>Orthosia hibisci</i> | (Guenée, 1852) |
| <i>Oryza longistaminata</i> | Chev. y Roehr., 1914 |
| <i>Oryza punctata</i> | Kotschy ex Steud. |
| <i>Oryza rufipogon</i> | Griffiths |
| <i>Ostrinia nubilalis</i> | (Hübner) |

| | |
|---|--|
| <i>Otala lactea</i> | (Muller, 1774) |
| <i>Otiorhynchus sulcatus</i> | (Fabricius, 1775) Germar |
| <i>Ovalisia virgata</i> | (Motschulsky) |
| <i>Pachnaeus azureus</i> | Gyllenhal , 1834 |
| <i>Pachymerus nucleorum</i> | (Fabricius) |
| <i>Pachysandra terminalis</i> | Siebold y Zucc. |
| <i>Paecilomyces variotii</i> | Bainier, 1907 |
| <i>Paeonia virus 1</i> | |
| <i>Paleacrita vernata</i> | (Peck, 1795) |
| Palm lethal yellowing phytoplasma | |
| <i>Pandemis heparana</i> | Denis y Schiffermüller |
| <i>Pantoea stewartii</i> | (Smith, 1898) Mergaert et al., 1993 |
| <i>Pantomorus cervinus</i> | (Boheman) |
| <i>Papaipema cataphracta</i> | Grote, 1864 |
| <i>Papaipema nebris</i> | Guenée, 1852 |
| <i>Papaipema purpurifasciata</i> | Grote y Robinson, 1868 |
| Papaya meleira virus | |
| <i>Parabemisia myricae</i> | (Kuwana, 1927) |
| <i>Paratachardina lobata lobata</i> | (Chamberlin, 1923) |
| <i>Paratrichodorus minor</i> | (Colbran, 1956) Siddiqi, 1974 |
| <i>Paratrichodorus porosus</i> | (Allen, 1957) Siddiqi, 1974 |
| <i>Parlatoria blanchardi</i> | (Targioni Tozzetti) Leonardi, 1899 |
| <i>Parlatoria ziziphi</i> | (Lucas, 1853) |
| <i>Paspalum scrobiculatum</i> | Linnaeus |
| <i>Passalora bataticola</i> | (Cif. y Bruner, 1931) Braun y Crous, 2003 |
| <i>Passalora crotonis</i> | (Ellis y Everhart, 1893) Crous y Braun, 2003 |
| <i>Passalora sojina</i> | (Hara, 1915) Shin y Braun, 1996 |
| Pea enation mosaic virus-1 | |
| Pea seed-borne mosaic virus | |
| Pea streak virus | |
| Peach latent mosaic pelamoviroid | |
| Peach rosette mosaic virus | |
| Peach rosette phytoplasma | |
| Peach X disease phytoplasma | |
| Peach yellows phytoplasma | |
| Peanut stunt virus | |
| Pear blister canker apscaviroid | |
| Pear bud drop disease | |
| <i>Pectinophora gossypiella</i> | Saunders, 1843 |
| <i>Pectinophora scutigera</i> | Holdaway, 1926 |
| <i>Pectobacterium atrosepticum</i> | (van Hall, 1902) Gardan et al., 2003 |
| <i>Pectobacterium carnegieanum</i> | (Standring, 1942) Brenner et al., 1973 |
| <i>Pectobacterium carotovorum</i> subsp. <i>carotovorum</i> | (Jones, 1901) Hauben et al., 1999 |
| <i>Pectobacterium cypripedii</i> | (Hori, 1911) Brenner et al., 1973 |
| Pelargonium leaf curl virus | |
| <i>Penicillium expansum</i> | Link, 1809 |
| <i>Pennisetum macrourum</i> | Trin. |
| <i>Pennisetum pedicellatum</i> | Trin. |

| | |
|---|--|
| <i>Pennisetum polystachion</i> | (Linnaeus, 1759) Schultes (1824) |
| Pepper mottle virus | |
| <i>Periconia circinata</i> | (Mangin, 1899) Saccardo, 1905 |
| <i>Peridroma saucia</i> | Hübner, 1803/808 |
| <i>Perkinsiella saccharicida</i> | Kirkaldy, 1903 |
| <i>Perofascia lepidii</i> | (McAlpine, 1895) Constant., 2002 |
| <i>Peronosclerospora maydis</i> | (Raciborski, 1897) Shaw, 1978 |
| <i>Peronosclerospora philippinensis</i> | (Weston) Shaw, 1978 |
| <i>Peronosclerospora sacchari</i> | (Miyake) Shirai y Hara, 1927 |
| <i>Peronospora farinosa</i> f.sp. <i>chenopodii</i> | Byford |
| <i>Peronospora farinosa</i> f.sp. <i>spinaciae</i> | Byford |
| <i>Pestalotiopsis palmarum</i> | (Cooke, 1875) Steyaert, 1949 |
| <i>Petrobia latens</i> | (Müller, 1776) |
| <i>Peyronellaea curtisii</i> | (Berkeley, 1878) Aveskamp et al., 2010 |
| <i>Phaeochoropsis palmicola</i> | (Stevens, 1920) Hyde y Cannon, 1999 |
| <i>Phakopsora cherimoliae</i> | Cummins, 1941 |
| <i>Phakopsora pachyrhizi</i> | Sydow y Sydow, 1914 |
| <i>Phialophora cinerescens</i> | (Wollenweber, 1930) Beyma, 1940 |
| <i>Phoenicococcus marlatti</i> | Cockerell, 1899 |
| <i>Phoma anemones</i> | Kauffman, 1936 |
| <i>Phoma destructiva</i> | Plowright, 1881 |
| <i>Phoma exigua</i> var. <i>exigua</i> | Saccardo, 1879 |
| <i>Phoma iowana</i> | Saccardo |
| <i>Phoma</i> sp. | |
| <i>Phoma tracheiphila</i> | (Petri) Kantachveli y Gikachvili, 1948 |
| <i>Phomopsis longicolla</i> | Hobbs, 1985 |
| <i>Phthorimaea operculella</i> | (Zeller 1873) |
| <i>Phyllachora pomigena</i> | (Schweinitz) Saccardo, 1883 |
| <i>Phyllachora torrendiella</i> | (Batista, 1948) Subileau et al., 1993 |
| <i>Phyllachora yapensis</i> subsp. <i>pongamiae</i> | (Berkeley y Broome, 1875) Cannon, 1991 |
| <i>Phyllocnistis citrella</i> | Stainton, 1856 |
| <i>Phyllognathus excavatus</i> | (Forster, 1771) |
| <i>Phyllonorycter blancardella</i> | (Fabricius, 1781) |
| <i>Phyllosticta ghaesembillae</i> | Koorders, 1907 |
| <i>Phyllosticta maculicola</i> | Halsted, 1894 |
| <i>Phyllosticta solitaria</i> | Ellis y Everhart, 1895 |
| <i>Phyllosticta telopeae</i> | Yip, 1989 |
| <i>Phyllotreta cruciferae</i> | (Goeze, 1777) |
| <i>Phymatotrichopsis omnivora</i> | (Duggar, 1916) Hennebert, 1973 |
| <i>Phytomonas staheli</i> | McGhee y McGhee, 1979 |
| <i>Phytomyza aconiti</i> | Hendel, 1920 |
| <i>Phytomyza ilicis</i> | Curtis, 1846 |
| <i>Phytomyza minuscula</i> | Goureau, 1851 |
| <i>Phytomyza rufipes</i> | Meigen, 1830 |
| <i>Phytonemus pallidus</i> | (Banks) |
| <i>Phytophthora cactorum</i> | (Lebert y Cohn) Schröter, 1886 |
| <i>Phytophthora capsici</i> | Leonian, 1922 |
| <i>Phytophthora cinnamomi</i> | Rands, 1922 |

| | |
|---|---|
| Phytophthora cryptogea | Pethybridge y Lafferty, 1919 |
| Phytophthora erythroseptica var. erythroseptica | (Waterhouse, 1963) |
| Phytophthora fragariae | Hickman, 1940 |
| Phytophthora infestans | (Montagne) de Bary, 1876 |
| Phytophthora megakarya | Brasier y Griffin, 1979 |
| Phytophthora ramorum | Werres et al., 2001 |
| Phytophthora richardiae | Buisman, 1927 |
| Phytophthora rubi | (Wilcox y Duncan, 1993) Man in't Veld, 2007 |
| Phytophthora sojae | Kaufmann y Gerdemann, 1958 |
| Pinnaspis buxi | (Bouché) |
| <i>Pinnaspis dysoxyl</i> | (Maskell, 1885) |
| <i>Planococcus citri</i> | (Risso, 1813) |
| <i>Planococcus ficus</i> | Signoret, 1875 |
| <i>Planococcus kenya</i> | (Le Pelley, 1935) |
| <i>Planococcus lilacinus</i> | Cockerell, 1905 |
| <i>Planococcus minor</i> | (Maskell, 1897) |
| Plasmopara halstedii | (Farlow) Berlese y de Toni, 1888 |
| Platycarpa scaly bark disease | |
| <i>Platynota idaeusalis</i> | Walker, 1859 |
| Pleospora betae | (Berlese) Nevodovsky, 1915 |
| Plowrightia circumscissa | Tracy y Earle, 1901 |
| Plum pox virus | |
| <i>Plutella xylostella</i> | Linnaeus, 1767 |
| <i>Pococera atramentalis</i> | (Lederer) |
| <i>Podischnus agenor</i> | (Olivier, 1789) |
| <i>Poecilopsus lineatus</i> | (Fabricius, 1798) |
| <i>Polygonum barbatum</i> | Linnaeus |
| <i>Polygonum convolvulus</i> | Linnaeus |
| <i>Polygonum lapathifolium</i> | Linnaeus |
| <i>Polyphagotarsonemus latus</i> | Banks |
| Polyscytalum pustulans | (Owen y Wakefield, 1919) Ellis, 1919 |
| <i>Popillia japonica</i> | Newman |
| Porroca | Ferrand, 1960 |
| Potato black ringspot virus | |
| Potato deforming mosaic virus | |
| Potato latent virus | |
| Potato leaflet stunt disease | |
| Potato leafroll virus | |
| Potato mop-top virus | |
| Potato spindle tuber pospiviroid | |
| Potato stolbur phytoplasma | |
| Potato virus A | |
| Potato virus M | |
| Potato virus S | |
| Potato virus T | |
| Potato virus X | |
| Potato virus Y | |
| Potato virus Yn | |

| | |
|--|--|
| Potato virus Yntn | |
| Potato witches' broom phytoplasma | |
| Potato yellow dwarf virus | |
| Potato yellow vein virus | |
| <i>Prathoda longissima</i> | (Deighton & MacGarvie, 1968) Simmons, 2007 |
| <i>Pratylenchus coffeae</i> | (Zimmermann, 1898) Filipjev y Steckhoven, 1941 |
| <i>Pratylenchus penetrans</i> | (Cobb, 1917) Filipjev y Steckhoven, 1941 |
| <i>Pratylenchus pratensis</i> | (de Man, 1880) Filipjev, 1936 |
| <i>Pratylenchus vulnus</i> | Allen y Jensen, 1951 |
| <i>Pratylenchus zae</i> | Graham, 1951 |
| <i>Prays citri</i> | Millière, 1873 |
| <i>Premnotrypes latithorax</i> | (Pierce) |
| <i>Premnotrypes sanfordi</i> | (Pierce) |
| <i>Premnotrypes solani</i> | Pierce , 1913 |
| <i>Premnotrypes vorax</i> | (Hustache) |
| Privet infectious variegation disease | |
| <i>Proeulia auraria</i> | (Clarke, 1949) |
| <i>Proeulia chrysopteris</i> | (Butler, 1883) |
| Prune dwarf virus | Saponari Maria |
| Prunus necrotic ringspot virus | |
| <i>Psammotermes hypostoma</i> | Desmeux |
| <i>Pseudaulacaspis cockerelli</i> | (Cooley, 1897) |
| <i>Pseudaulacaspis pentagona</i> | (Targioni Tozzetti, 1886) MacGillivray, 1921 |
| <i>Pseudischnaspis bowreyi</i> | (Cockerell, 1893) |
| <i>Pseudocercospora pongamiae-pinnatae</i> | Ram y Mallaiah, 1993 |
| <i>Pseudocneorhinus bifasciatus</i> | (Roelofs) |
| <i>Pseudococcus comstocki</i> | (Kuwana, 1902) |
| <i>Pseudococcus longispinus</i> | (Targioni Tozzetti, 1867) Targioni Tozzetti |
| <i>Pseudococcus viburni</i> | (Signoret, 1875) Signoret |
| <i>Pseudocochliobolus pallescens</i> | Tsuda y Ueyama, 1983 |
| <i>Pseudomonas cichorii</i> | (Swingle, 1925) Stapp, 1928 |
| <i>Pseudomonas erodii</i> | Lewis |
| <i>Pseudomonas fluorescens</i> | (Trevisan, 1889) Migula, 1895 |
| <i>Pseudomonas gardeniae</i> | |
| <i>Pseudomonas marginalis</i> | (Brown, 1918) Stevens, 1925 |
| <i>Pseudomonas marginalis</i> pv. <i>marginalis</i> | (Brown, 1918) Stevens, 1925 |
| <i>Pseudomonas savastanoi</i> pv. <i>glycinea</i> | (Coerper, 1919) Gardan et al., 1992 |
| <i>Pseudomonas savastanoi</i> pv. <i>savastanoi</i> | (ex Smith, 1908) Gardan et al., 1992 |
| <i>Pseudomonas syringae</i> pv. <i>aptata</i> | (Brown y Jamieson, 1913) Young et al., 1978 |
| <i>Pseudomonas syringae</i> pv. <i>atropaciens</i> | (McCulloch, 1920) Young et al., 1978 |
| <i>Pseudomonas syringae</i> pv. <i>coronafaciens</i> | (Elliott, 1920) Young et al., 1978 |
| <i>Pseudomonas syringae</i> pv. <i>delphinii</i> | (Smith, 1904) Young et al., 1978 |
| <i>Pseudomonas syringae</i> pv. <i>garcae</i> | (Amaral et al., 1956) Young et al., 1978 |
| <i>Pseudomonas syringae</i> pv. <i>lachrymans</i> | (Smith y Bryan, 1915) Young et al., 1978 |
| <i>Pseudomonas syringae</i> pv. <i>lapsa</i> | (Ark, 1940) Young et al., 1978 |
| <i>Pseudomonas syringae</i> pv. <i>maculicola</i> | (McCulloch, 1911) Young et al., 1978 |
| <i>Pseudomonas syringae</i> pv. <i>morsprunorum</i> | (Wormald, 1931) Young et al., 1978 |

| | |
|---|--|
| <i>Pseudomonas syringae</i> pv. <i>primulae</i> | (Ark y Gardner, 1936) Young et al., 1978 |
| <i>Pseudomonas syringae</i> pv. <i>sesami</i> | (Malkoff, 1906) Young et al., 1978 |
| <i>Pseudomonas syringae</i> pv. <i>syringae</i> | van Hall, 1902 |
| <i>Pseudomonas syringae</i> pv. <i>tomato</i> | (Okabe) Young et al. |
| <i>Pseudomonas syringae</i> pv. <i>zizaniae</i> | (ex Bowden y Percich, 1983) Young et al., 1991 |
| <i>Pseudomonas viridiflava</i> | (Burkholder, 1930) Dowson, 1939 |
| <i>Pseudomonas xanthochlora</i> | (Schuster) Stapp, 1928 |
| <i>Pseudospiropes indicus</i> | Rai y Rai, 1995 |
| <i>Psylla buxi</i> | (Linné, 1758) |
| <i>Puccinia albulensis</i> | Magnus, 1890 |
| <i>Puccinia allii</i> | (DeCandolle, 1815) Rudolphi, 1829 |
| <i>Puccinia andina</i> | Dietel y Neger, 1899 |
| <i>Puccinia andropogonis</i> | Schweinitz, 1832 |
| <i>Puccinia anisacanthi</i> | Dietel y Holw., 1901 |
| <i>Puccinia arenariae</i> | (Schumacher, 1803) Winter, 1880 |
| <i>Puccinia aristidae</i> | Tracy, 1893 |
| <i>Puccinia arundiariae</i> | Schweinitz, 1822 |
| <i>Puccinia bartholomaei</i> | Dietel, 1892 |
| <i>Puccinia calcitrapae</i> | DeCandolle, 1805 |
| <i>Puccinia campanulae</i> | Carmichael, 1836 |
| <i>Puccinia cannae</i> | Hennings, 1902 |
| <i>Puccinia caricina</i> var. <i>caricina</i> | DeCandolle, 1815 |
| <i>Puccinia chelonis</i> | Dietel y Holway, 1897 |
| <i>Puccinia chrysanthemi</i> | Roze, 1900 |
| <i>Puccinia cirsii-lanceolati</i> | Schröter, 1887 |
| <i>Puccinia cnici-oleracei</i> | Persoon, 1823 |
| <i>Puccinia concrescens</i> | Ellis y Everhart, 1915 |
| <i>Puccinia conoclinii</i> | Seymour, 1885 |
| <i>Puccinia cooperiae</i> | Long, 1902 |
| <i>Puccinia cyani</i> | Passerini, 1874 |
| <i>Puccinia cyperi</i> | Arthur, 1891 |
| <i>Puccinia delphinii</i> | Dietel y Holway, 1893 |
| <i>Puccinia dichondrae</i> | Montagne, 1852 |
| <i>Puccinia dioicae</i> var. <i>extensicola</i> | (Plowright, 1889) Henderson, 1961 |
| <i>Puccinia douglasii</i> | Ellis y Everhart, 1893 |
| <i>Puccinia dulichii</i> | Sydow y Sydow, 1904 |
| <i>Puccinia eleocharidis</i> | Arthur, 1884 |
| <i>Puccinia ellisiana</i> | Thümen, 1878 |
| <i>Puccinia enceliae</i> | Dietel y Holway, 1897 |
| <i>Puccinia extensicola</i> var. <i>asteris</i> | (Thümen, 1877) Arthur |
| <i>Puccinia gentianae</i> | (Strauss) Martius, 1824 |
| <i>Puccinia graminis</i> f.sp. <i>tritici</i> raza TTKS | Erikssoon y Henning |
| <i>Puccinia grindeliae</i> | Peck, 1879 |
| <i>Puccinia heliconiae</i> | (Dietel, 1897) Arthur, 1918 |
| <i>Puccinia horiana</i> | Hennings, 1901 |
| <i>Puccinia iridis</i> | Wallroth, 1844 |
| <i>Puccinia iridis</i> f.sp. <i>floridana</i> | Sobers, 1966 |

| | |
|--|--|
| <i>Puccinia irrequisita</i> | Jackson, 1921 |
| <i>Puccinia kuehnii</i> | (Krüger, 1890) Butler, 1914 |
| <i>Puccinia lateritia</i> | Schweinitz, 1853 |
| <i>Puccinia leonotidicola</i> | Hennings, 1903 |
| <i>Puccinia liatridis</i> | (Bethel ex Arthur y Fromme, 1920) Arthur, 1934 |
| <i>Puccinia limosae</i> | Magnus, 1877 |
| <i>Puccinia lobeliae</i> | Gerard, 1873 |
| <i>Puccinia mccleanii</i> | |
| <i>Puccinia melampodii</i> | Dietel y Holway, 1897 |
| <i>Puccinia obtecta</i> | Peck, 1873 |
| <i>Puccinia panici</i> | Dietel y Holway, 1895 |
| <i>Puccinia pelargonii-zonalis</i> | Doidge, 1926 |
| <i>Puccinia peridermiospora</i> | (Ellis y Tracy, 1890) Arthur, 1899 |
| <i>Puccinia physostegiae</i> | Peck y Clinton, 1878 |
| <i>Puccinia pimpinellae</i> | (Strauss, 1810) Link, 1824 |
| <i>Puccinia poarum</i> | Nielsen, 1877 |
| <i>Puccinia probabilis</i> | Arthur y Cummins, 1934 |
| <i>Puccinia psidii</i> | Winter, 1884 |
| <i>Puccinia punctiformis</i> | (Strauss, 1810) Röhling, 1813 |
| <i>Puccinia rhaetica</i> | Fischer, 1899 |
| <i>Puccinia rubigo-vera</i> var. <i>agropyri</i> | (Erikss.) Arthur |
| <i>Puccinia stipae</i> | (Opiz, 1852) Arthur, 1884 |
| <i>Puccinia tenuis</i> | Bonorden |
| <i>Puccinia tillandsiae</i> | Cummins y Pollack, 1974 |
| <i>Puccinia triticina</i> | Eriksson, 1899 |
| <i>Puccinia vagans</i> | (DeCandolle) Arthur, 1934 |
| <i>Puccinia venustula</i> | (Arthur, 1916) Arthur, 1918 |
| <i>Puccinia verbesinae</i> | Schweinitz, 1822 |
| <i>Pucciniastrum epilobii</i> | Otth, 1861 |
| <i>Pyrenophora chaetomioides</i> | Spegazzini, 1899 |
| <i>Pyricularia setariae</i> | Nisikado, 1927 |
| <i>Pyrilla perpusilla</i> | (Walker, 1851) Walker |
| <i>Pyrria umbra</i> | Hüfnagel, 1766 |
| <i>Pythium myriotylum</i> | Drechsler, 1930 |
| <i>Pythium splendens</i> | Braun, 1925 |
| <i>Radopholus similis</i> | (Cobb, 1893) Thorne, 1949 |
| <i>Ralstonia solanacearum</i> | (Smith, 1896) Yabuuchi et al., 1996 |
| <i>Ralstonia solanacearum</i> raza 2 | (Smith, 1896) Yabuuchi et al., 1996 |
| <i>Ralstonia solanacearum</i> raza 3 | (Smith, 1896) Yabuuchi et al., 1996 |
| <i>Ranunculus repens</i> | Linnaeus |
| <i>Raoiella indica</i> | Hirst |
| Raspberry bushy dwarf virus | |
| Raspberry leaf curl virus | |
| Raspberry ringspot virus | |
| <i>Ravenelia hobsonii</i> | Cooke, 1880 |
| <i>Ravenelia humphreyana</i> | Hennings, 1898 |
| <i>Ravenelia spegazziniana</i> | Lindquist, 1946 |

| | |
|-------------------------------------|--|
| Ravenelia texensis | Ellis y Galloway, 1894 |
| Red clover vein mosaic virus | |
| <i>Retithrips syriacus</i> | (Mayet, 1890) |
| <i>Retracrus elaeis</i> | Keifer |
| <i>Rhabdoscelus obscurus</i> | (Boisduval) |
| Rhabdospora oudemansii | Hennings, 1900 |
| <i>Rhadinaphelenchus cocophilus</i> | (Cobb, 1919) Goodey, 1960 |
| <i>Rhagoletis basiola</i> | (Sacken, 1877) |
| <i>Rhagoletis cerasi</i> | Linnaeus, 1758 |
| <i>Rhagoletis indifferens</i> | Curran, 1932 |
| <i>Rhagoletis pomonella</i> | Walsh, 1867 |
| <i>Rhagoletis suavis</i> | Loew, 1862 |
| <i>Rhagoletis tomatis</i> | Foote, 1981 |
| <i>Rhigopsidius tucumanus</i> | Heller, 1906 |
| <i>Rhinostomus barbirostris</i> | (Fabricius) Rafinesque |
| Rhizobium radiobacter | (Beijerinck y van Delden, 1902) Young et al., 2001 |
| Rhizobium rhizogenes | (Riker et al., 1930) Young et al., 2001 |
| Rhizoctonia crocorum | (Persoon) DeCandolle, 1815 |
| <i>Rhizoecus falcifer</i> | Herculais, 1878 |
| <i>Rhizoecus floridanus</i> | Hambleton, 1973 |
| Rhodococcus fascians | (Tilford, 1936) Goodfellow, 1984 |
| <i>Rhopalosiphoninus latysiphon</i> | (Davidson, 1912) |
| <i>Rhopobota ilicifoliana</i> | Kearfott, 1907 |
| Rhopoglyphus filicinus | (Fries, 1823) Nitschke ex Fuckel, 1870 |
| <i>Rhynchites foveipennis</i> | Fairm. , 1888 |
| <i>Rhynchophorus ferrugineus</i> | (Olivier, 1790) |
| <i>Rhyssomatus subtilis</i> | Fiedler , 1937 |
| Rice black streaked dwarf virus | |
| Rice hoja blanca virus | |
| Rice stripe virus | |
| <i>Richardia brasiliensis</i> | Gomes, 1801 |
| <i>Richardia grandiflora</i> | (Cham. & Schltdl.) Steud. |
| <i>Richardia stellaris</i> | (Cham. & Schltdl.) Steud. |
| Rose wilt disease | |
| Rose X disease | |
| <i>Rottboellia cochinchinensis</i> | (Lour.) Clayton (1981) |
| <i>Rotylenchulus parvus</i> | (Williams, 1960) Sher, 1961 |
| <i>Rubus fruticosus</i> | Linnaeus |
| <i>Rubus moluccanus</i> | Linnaeus |
| Rubus stunt phytoplasma | Seemüller et al., 1998 |
| <i>Sabulodes aegrotata</i> | (Guenée, 1858) |
| <i>Sacadodes pyralis</i> | Dyar, 1912 |
| Saccharata proteae | (Wakefield, 1922) Denman & Crous, 2004 |
| <i>Saccharum spontaneum</i> | Linnaeus, 1771 |
| <i>Salsola vermiculata</i> | Linnaeus |
| Sandal spike phytoplasma | |
| Sarocladium oryzae | (Sawada, 1922) Gams y Hawksworth, 1975 |

| | |
|--|---|
| <i>Schistocerca piceifrons piceifrons</i> | (Walker, 1870) |
| <i>Schoenoplectiella juncooides</i> | (Roxb.) Lye |
| <i>Scirtothrips aurantii</i> | Faure, 1929 |
| <i>Scirtothrips dorsalis</i> | Hood, 1919 |
| <i>Sclerophoma eustomonis</i> | Taubenhaus y Ezekiel |
| <i>Sclerophthora rayssiae</i> var. <i>zeae</i> | Payak y Renfro, 1967 |
| <i>Sclerotinia denigriicans</i> | |
| <i>Sclerotinia minor</i> | Jagger, 1920 |
| <i>Sclerotinia trifoliorum</i> | Eriksson, 1880 |
| <i>Sclerotium denigrans</i> | Pape, 1943 |
| <i>Scolytus rugulosus</i> | (Müller) |
| <i>Scolytus schevyrewi</i> | Semenov, 1902 |
| <i>Scutellonema brachyurus</i> | (Steiner, 1938) Andrassy, 1958 |
| <i>Senecio vulgaris</i> | Linnaeus, 1753 |
| <i>Septoria atriplicis</i> | (Westendorp, 1851) Fuckel, 1870 |
| <i>Septoria helianthi</i> | Ellis y Kellerman, 1883 |
| <i>Septoria steviae</i> | Ishiba et al., 1982 |
| <i>Sesamia calamistis</i> | Hampson, 1910 |
| <i>Sesamia cretica</i> | Lederer, 1857 |
| <i>Sesamia inferens</i> | (Walker, 1856) |
| <i>Setaria pumila</i> | (Poir.) Roemer y Schultes, 1817 |
| <i>Setaria spp.</i> | |
| <i>Sigmodon hispidus</i> | Say y Ord, 1825 |
| <i>Silene noctiflora</i> | Linnaeus |
| <i>Silybum marianum</i> | (Linnaeus) Gaertn. |
| <i>Sinapis arvensis</i> | Linnaeus |
| <i>Sitodiplosis mosellana</i> | (Gehin, 1857) |
| <i>Solanum carolinense</i> | Linnaeus (1753) |
| <i>Solanum peruvianum</i> | Linnaeus |
| <i>Solanum ptycanthum</i> | Dunal |
| <i>Solanum viarum</i> | Dunal |
| <i>Sonchus arvensis</i> | Linnaeus, 1753 |
| Sonchus yellow net virus | |
| Sowbane mosaic virus | |
| <i>Sparganothis sulfureana</i> | Clemens, 1860 |
| <i>Sphaceloma poinsettiae</i> | Jenkins y Rühle, 1942 |
| <i>Sphaceloma venezuelanum</i> | Bitancourt y Jenkins |
| <i>Sphacelotheca erianthi</i> | (Sydow y Sydow) Mundkur, 1942 |
| <i>Sphacelotheca macrospora</i> | Yen y Wang, 1955 |
| <i>Sphacelotheca reiliana</i> | (Kühn, 1875) Clinton, 1902 |
| <i>Sphaeropsis vitis</i> | Jakubski, 1965 |
| <i>Sphaeropsis hedericola</i> | (Spegazzini) Saccardo, 1884 |
| <i>Sphenarium sp.</i> | |
| <i>Sphenoclea zeylanica</i> | Gaertn., 1788 |
| <i>Sphenophorus venatus vestitus</i> | Chittendon |
| <i>Spilonota albicana</i> | Motschulsky |
| <i>Spilonota ocellana</i> | (Denis y Schiffermüller) Schiffermüller |
| <i>Spilosoma virginica</i> | Fabricius, 1798 |

| | |
|--|--|
| <i>Spiroplasma citri</i> | Saglio et al., 1973 |
| <i>Spodoptera exigua</i> | (Hübner- 1803-1808) |
| <i>Spodoptera littoralis</i> | (Boisduval, 1833) |
| <i>Spodoptera litura</i> | (Fabricius, 1775) |
| <i>Spodoptera mauritia</i> | Boisduval, 1833 |
| <i>Spoladea recurvalis</i> | (Fabricius, 1775) |
| <i>Spongospora subterranea</i> | (Wallr.) Lagerh., 1892 |
| Squash mosaic virus | |
| <i>Steneotarsonemus concavuscutum</i> | Lofego & Gondim |
| <i>Steneotarsonemus laticeps</i> | (Halbert) |
| <i>Steneotarsonemus spinki</i> | Smiley, 1967 |
| <i>Stenoma catenifer</i> | Walsingham, 1912 |
| <i>Stigmina palmivora</i> | (Saccardo, 1898) Hughes, 1952 |
| Strawberry crinkle virus | |
| Strawberry latent C virus | |
| Strawberry latent ringspot virus | |
| Strawberry mild yellow edge virus | |
| Strawberry mottle virus | |
| Strawberry pseudo mild yellow edge virus | |
| Strawberry vein banding virus | |
| Strawberry witches' broom phytoplasma | |
| <i>Streptomyces scabiei</i> | (ex Thaxter, 1892) Lambert y Loria, 1989 |
| <i>Striga angustifolia</i> | (Don) Saldanha, 1963 |
| <i>Striga asiatica</i> | (Linnaeus, 1753) Kuntze, 1891 |
| <i>Striga densiflora</i> | (Benth., 1835) Benth., 1863 |
| <i>Striga hermonthica</i> | (Del., 1813) Benth., 1836 |
| <i>Striga spp.</i> | |
| <i>Stromatinia cepivora</i> | (Berkeley, 1841) Whetzel, 1945 |
| Sugarcane bacilliform virus | |
| Sugarcane sereh disease | |
| Sugarcane streak virus | |
| Sugarcane yellow leaf virus | |
| <i>Symmetrischema tangolia</i> | Turner, 1919 |
| <i>Synchytrium aureum</i> | Schröter, 1870 |
| <i>Synchytrium edgertonii</i> | Cook, 1945 |
| <i>Synchytrium endobioticum</i> | (Schilberszky) Percival, 1909 |
| <i>Synchytrium fulgens</i> | Schröter, 1873 |
| <i>Synchytrium nigrescens</i> | Davis, 1924 |
| <i>Synchytrium vaccinii</i> | Thomas, 1889 |
| <i>Taeniothrips eucharii</i> | (Whetzel, 1923) |
| <i>Taphrina bullata</i> | (Berkeley y Broome, 1855) Tulasne, 1866 |
| <i>Teratosphaeria jonkershoekensis</i> | (Wyk ET AL., 1975) Crous y Braun, 2007 |
| <i>Tetraleurodes ursorum</i> | (Cockerell, 1910) |
| <i>Thanatephorus cucumeris</i> | (Frank) Donk, 1956 |
| <i>Thanatephorus theobromae</i> | (Talbot y Keane) Roberts, 1999 |
| <i>Thaumatotibia leucotreta</i> | Meyrick |
| <i>Thecaphora solani</i> | Barrus, 1944 |
| <i>Thekopsora areolata</i> | (Fries, 1818) Magnus, 1875 |

| | |
|-----------------------------------|----------------------------------|
| <i>Themeda quadrivalvis</i> | (Linnaeus) Kuntze |
| <i>Theretra oldenlandiae</i> | (Fabricius, 1775) |
| <i>Thlaspi arvense</i> | Linnaeus (1753) |
| <i>Thrips angusticeps</i> | Uzel, 1895 |
| <i>Thrips flavus</i> | Schrank |
| <i>Thrips hawaiiensis</i> | (Morgan, 1913) |
| <i>Thrips meridionalis</i> | (Priesner, 1926) |
| <i>Thrips nigropilosus</i> | Uzel |
| <i>Thrips palmi</i> | Karny, 1925 |
| <i>Thrips simplex</i> | (Morison, 1930) |
| <i>Thrips sumatrensis</i> | (Priesner, 1934) |
| <i>Tilletia barclayana</i> | (Brefeld) Saccardo y Sydow, 1899 |
| <i>Tilletia controversa</i> | Kühn, 1874 |
| <i>Tilletia indica</i> | Mitra, 1931 |
| <i>Tilletia tritici</i> | (Bjerkander, 1775) Winter, 1874 |
| <i>Tipula paludosa</i> | Meigen, 1830 |
| Tobacco etch virus | |
| Tobacco mosaic virus | |
| Tobacco necrosis virus | |
| Tobacco rattle virus | |
| Tobacco ringspot virus | |
| Tobacco streak virus | |
| Tomato aspermy virus | |
| Tomato black ring virus | |
| Tomato bushy stunt virus | |
| Tomato mottle virus | |
| Tomato ringspot virus | |
| Tomato spotted wilt virus | |
| <i>Toxoptera citricida</i> | (Kirkaldy) |
| <i>Toxotrypana sp.</i> | |
| <i>Tremex fuscicornis</i> | (Fabricius, 1787) |
| <i>Tretendophragmia palmivora</i> | Subramanian, 1994 |
| <i>Trialeurodes vaporariorum</i> | (Westwood, 1856) |
| <i>Trichoconiella padwickii</i> | (Ganguly, 1948) Jain, 1976 |
| <i>Trichodorus spp.</i> | |
| <i>Trichodorus viruliferus</i> | Hooper, 1963 |
| <i>Trichothecium roseum</i> | (Persoon, 1794) Link, 1809 |
| <i>Trichromothrips xanthius</i> | (Williams, 1917) |
| <i>Trioza alacris</i> | Flor, 1861 |
| <i>Tripleurospermum inodorum</i> | (Linnaeus) Sch. |
| <i>Tripleurospermum maritimum</i> | (Linnaeus) Koch |
| <i>Trirhithrum coffeae</i> | Bezzi, 1918 |
| <i>Trogoderma granarium</i> | Everts, 1899 |
| Tulip breaking virus | |
| <i>Tuta absoluta</i> | (Meyrick, 1917) Povolny (1994) |
| <i>Tylenchorhynchus claytoni</i> | Steiner, 1937 |
| <i>Typophorus nigrilus</i> | (Fabricius, 1801) |
| <i>Udea rubigalis</i> | (Guenée) Guenée, 1854 |

| | |
|---|---|
| <i>Uleodothis pteridis</i> | Stevens, 1920 |
| <i>Ulex europaeus</i> | Linnaeus |
| <i>Unaspis citri</i> | (Comstock, 1883) |
| <i>Unaspis yanonensis</i> | (Kuwana, 1923) |
| <i>Uredo behnickiana</i> | Hennings, 1905 |
| <i>Uredo gladioli-buettneri</i> | Bubák, 1911 |
| <i>Urocarpidium peruvianum</i> | (Linnaeus, 1753) Krapov. |
| <i>Urochloa panicoides</i> | Beauv. |
| <i>Urocystis sorosporioides</i> | Körnicker ex von Waldheim, 1877 |
| <i>Uromyces asclepiadis</i> | Cooke, 1877 |
| <i>Uromyces compactus</i> | Peck, 1882 |
| <i>Uromyces dianthi</i> | (Persoon) Niessl 1872 |
| <i>Uromyces euphorbiae</i> | Cooke y Peck, 1873 |
| <i>Uromyces funci</i> | |
| <i>Uromyces gentianae</i> | Arthur, 1891 |
| <i>Uromyces holwayi</i> | Lagerheim, 1889 |
| <i>Uromyces houstoniatus</i> | Sheldon, 1909 |
| <i>Uromyces junci</i> | (Desmazières) Tulasne y Tulasne, 1854 |
| <i>Uromyces limonii</i> | (DeCandolle, 1805) Léveillé, 1849 |
| <i>Uromyces lupini</i> | Berkeley y Curtis, 1860 |
| <i>Uromyces occidentalis</i> | Dietel, 1903 |
| <i>Uromyces pisi-sativi</i> | (Persoon) Liro, 1908 |
| <i>Uromyces probus</i> | Arthur 1911 |
| <i>Uromyces sparganii</i> | Cooke y Peck, 1874 |
| <i>Uromyces triquetrus</i> | Cooke, 1862 |
| <i>Uromycladium tepperianum</i> | (Saccardo, 1889) McAlpine, 1906 |
| <i>Ustilago cissi</i> | Tulasne y Tulasne |
| <i>Ustilago nuda</i> f.sp. <i>tritici</i> | Schaffnit, 1928 |
| <i>Ustilago scitaminea</i> | Sydow, 1924 |
| <i>Vaccaria hispanica</i> | (Miller) Rauschert |
| <i>Vankya vaillantii</i> | <i>Vankya vaillantii</i> (Tul. & C. Tul.) Ershad 2000 |
| <i>Verticillium albo-atrum</i> | Reinke y Berthold, 1879 |
| <i>Verticillium dahliae</i> | Klebahn, 1913 |
| <i>Verticillium heterocladium</i> | Penzig, 1882 |
| <i>Villosiclava virens</i> | (Sakurai ex Nakata) Tanaka y Tanaka, 2009 |
| <i>Vinsonia stellifera</i> | (Westwood, 1871) |
| <i>Virachola bimaculata</i> | Hewitson, 1874 |
| <i>Virus marchitez variegada</i> | |
| <i>Viteus vitifoliae</i> | (Fitch, 1851) |
| <i>Volutella albido-pila</i> | Boudier, 1891 |
| Watermelon mosaic virus-II | Wang et al., 1993 |
| Wheat streak mosaic virus | |
| <i>Xanthomonas albilineans</i> | (Ashby, 1929) Dowson, 1943 |
| <i>Xanthomonas arboricola</i> pv. <i>pruni</i> | (Smith, 1903) Vauterin et al. |
| <i>Xanthomonas axonopodis</i> pv. <i>alfalfae</i> | (Riker et al., 1935) Vauterin et al., 1995 |
| <i>Xanthomonas axonopodis</i> pv. <i>begoniae</i> | (Takimoto 1934) Vauterin et al. 1995 |
| <i>Xanthomonas axonopodis</i> pv. <i>citri</i> | (Hasse, 1915) Vauterin et al., 1995 |
| <i>Xanthomonas axonopodis</i> pv. <i>dieffenbachiae</i> | (McCulloch y Pirone, 1939) Vauterin et al., |

| | |
|---|---|
| | 1995 |
| <i>Xanthomonas axonopodis</i> pv. <i>malvacearum</i> | (Smith, 1901) Vauterin et al., 1995 |
| <i>Xanthomonas axonopodis</i> pv. <i>poinsettiicola</i> | (Patel et al., 1951) Vauterin et al., 1995 |
| <i>Xanthomonas axonopodis</i> pv. <i>ricini</i> | (Yoshii & Takimoto, 1928) Vauterin et al., 1995 |
| <i>Xanthomonas axonopodis</i> pv. <i>vasculorum</i> | (Cobb, 1893) Vauterin et al., 1995 |
| <i>Xanthomonas campestris</i> | (Pammel, 1895) Dowson, 1939 |
| <i>Xanthomonas campestris</i> pv. <i>aberrans</i> | (Knösel, 1961) Dye, 1978 |
| <i>Xanthomonas campestris</i> pv. <i>armoraciae</i> | (McCulloch, 1929) Dye, 1978 |
| <i>Xanthomonas campestris</i> pv. <i>campestris</i> | (Pammel, 1895) Dowson, 1939 |
| <i>Xanthomonas campestris</i> pv. <i>sesami</i> | (Sabet y Dowson, 1960) Dye, 1978 |
| <i>Xanthomonas cucurbitae</i> | (Bryan, 1926) Vauterin et al., 1995 |
| <i>Xanthomonas fragariae</i> | Kennedy y King, 1962 |
| <i>Xanthomonas hortorum</i> pv. <i>carotae</i> | (Kendrick, 1934) Vauterin et al., 1995 |
| <i>Xanthomonas hortorum</i> pv. <i>hederae</i> | (Arnaud, 1920) Vauterin et al., 1995 |
| <i>Xanthomonas hortorum</i> pv. <i>pelargonii</i> | (Brown, 1923) Vauterin et al., 1995 |
| <i>Xanthomonas hyacinthi</i> | (Wakker, 1883) Vauterin et al., 1995 |
| <i>Xanthomonas oryzae</i> pv. <i>oryzae</i> | (Ishiyama, 1922) Swings et al., 1990 |
| <i>Xanthomonas oryzae</i> pv. <i>oryzicola</i> | (Fang et al., 1957) Swings et al., 1990 |
| <i>Xanthomonas translucens</i> pv. <i>cerealis</i> | (Hagborg, 1942) Vauterin et al., 1995 |
| <i>Xanthomonas vasicola</i> pv. <i>holcicola</i> | (Elliott, 1930) Vauterin et al., 1995 |
| <i>Xanthomonas vesicatoria</i> | (Doidge, 1920) Dowson, 1939 |
| <i>Xanthopastis timais</i> | (Cramer, 1782) |
| <i>Xestia c-nigrum</i> | (Linnaeus, 1758) |
| <i>Xiphinema coxi</i> | Tarjan, 1964 |
| <i>Xiphinema diversicaudatum</i> | (Micoletzky, 1927) Thorne, 1939 |
| <i>Xiphinema index</i> | Thorne y Allen, 1950 |
| <i>Xiphinema rivesi</i> | Dalmasso, 1969 |
| <i>Xylella fastidiosa</i> subsp. <i>fastidiosa</i> | (Wells et al., 1987) |
| <i>Xylella fastidiosa</i> subsp. <i>multiplex</i> | (Wells et al., 1987) |
| <i>Xylella fastidiosa</i> subsp. <i>pauca</i> | (Wells et al., 1987) |
| <i>Xylomyges curialis</i> | (Grote, 1873) |
| <i>Xylophilus ampelinus</i> | (Panagopoulos, 1969) Willems et al., 1987 |
| <i>Xyphon fulgida</i> | (Nottingham, 1932) |
| <i>Yucca bacilliform virus</i> | |
| <i>Yuccaborus frontalis</i> | LeConte, 1876 |
| <i>Zeuzera pyrina</i> | Linnaeus, 1761 |
| <i>Zonosemata electa</i> | (Say, 1830) |
| <i>Zygotylenchus guevarai</i> | (Jiménez, 1963) Braun y Loof, 1966 |