



Adansonnia-Consulting



Vegetation survey of Mlele Beekeeping Zone in 2018



Dr. Urs Bloesch

Evilard, 1 May 2019

Acknowledgements

I sincerely thank ADAP Geneva for entrusting to us the vegetation survey of Mlele Beekeeping Zone. I am deeply grateful to Inyonga Beekeepers Association (IBA) and to the ADAP Project Supervisor Shabani Halfani and his team for their generous hospitality and support throughout the mission. I would like to express my warmest thanks to the Village Game Scout (VGS) team including Dicksoni Malembeka, Johni Gervasi, Felesiano Futakamba.

My special thanks go to Lucile Daudet who later finished the two vegetation transects together with the VGS team and to Frank Mbago for his field assistance and the plant identification at the Herbarium of the University of Dar es Salaam. Finally, I am grateful to the Project driver Yahya Ally for careful driving and tolerance during field work. The field work would not have been possible without the tireless commitment of all team members.

Abstract

A vegetation survey of Mlele Beekeeping Zone (BKZ) in Inyonga District has been conducted in 2018. The objective of the survey was a) to detect possible vegetation changes within the Mlele BKZ and b) to assess the current stage of the vegetation of the whole Mlele BKZ and its current threats (baseline).

13 plots out of the 30 vegetation plots from the 2004 survey have been resurveyed during this assessment. The species richness of the Mlele BKZ comprises 83 species for the 2018 vegetation survey. The 13 vegetation plots resurveyed will serve as permanent plots for monitoring and evaluating any vegetation changes at site level.

Moreover, two vegetation transects of about 10 km length each have been outlined and 800 vegetation points have been sampled with the Point-Centered Quarter Method (PCQM). Tree density, basal area, dominant tree height, and estimated standing volume are significantly higher on upper land compared to the miombo woodlands below the escarpment.

The most frequent trees/shrubs in Mlele BKZ in the lower diameter class ($DBH < 20\text{cm}$) are *Diplorhynchus condylocarpon*, *Pseudolachnostylis maprouneifolia*, and *Pterocarpus angolensis*. These species are quite fire-tolerant what may explain their prevalence in the annually burnt Mlele BKZ. In the upper diameter class ($DBH \geq 20\text{cm}$) *Julbernardia globiflora* and *Brachystegia manga/utilis* are the most frequent tree species in Mlele BKZ. Young *Pterocarpus angolensis* from the lower diameter class ($DBH < 20\text{cm}$) are widespread in Mlele BKZ and are amongst the five most frequent and dominant species.

The vegetation surveys in 2018 allowed completing substantially the former tree/shrub list for Mlele BKZ from Mwanguango (2004). The checklist now includes in total 123 trees/shrubs with their scientific and Konongo names including also species identified from opportunistic sampling.

In total 28 human site disturbances have been assessed including 34 trees cut (many of them *Pterocarpus angolensis*) along the two transects covering roughly an area of 160 ha. About 80% of the disturbances happened on the lower land what may be explained by the proximity of the principal road Inyonga to Mpanda.

The use of the PCQM along the vegetation transects is an appropriate and quite simple method to assess the vegetation baseline and in addition also all human disturbances along the transects. It is suggested to use this method also for the neighbouring protected areas what would allow to compare ecosystem (vegetation) health of protected areas under different managements.

The Miombo habitats of Mlele BKZ are well protected. No sign of recent poaching was detected and no logging happened in the last five years in the two belt transects as a result of the participatory management of the beekeeping zone with regular patrolling, support of beekeeping activities and awareness raising campaign by the Project.

List of abbreviations

| | |
|------|--|
| ADAP | Association for the Development of Protected Areas |
| ba | Basal area |
| BKZ | Beekeeping Zone |
| DBH | Diameter at Breast Height |
| FR | Forest Reserve |
| GCA | Game Controlled Area |
| IBA | Inyonga Beekeepers Association |
| MNRT | Ministry of Natural Resources and Tourism |
| MoU | Memorandum of Understanding |
| NGO | Non Governmental Organisation |
| PA | Protected Area |
| PCQM | Point-Centered Quarter Method |
| TAWA | Tanzania Wildlife Management Authority |
| TFS | Tanzania Forest Services Agency |
| VGS | Village Game Scout |

Content

| | |
|---|----|
| Acknowledgements..... | 2 |
| Abstract..... | 3 |
| List of abbreviations..... | 4 |
| 1) Introduction | 6 |
| 2) Study area | 7 |
| 3) Methods..... | 9 |
| 3.1 Data collection | 9 |
| 3.2 Data analysis | 12 |
| 4) Results..... | 13 |
| 5) Discussions | 17 |
| 6) Conclusions | 19 |
| 7) Recommendations | 19 |
| 8) Bibliography | 20 |
| Appendix A: Permanent vegetation plots..... | 22 |
| Appendix B: Vegetation transects..... | 36 |
| Appendix C: Tree and shrub checklist Mlele BKZ (27/3/2019) | 70 |
| Appendix D: Vegetation survey 2018 of Mlele Beekeeping Zone | 73 |

1) Introduction

Following a transfer of management rights negotiated with the government in 2010, the Mlele Beekeeping Zone (BKZ) is managed by the Inyonga Beekeepers Association (IBA) in collaboration with the District authorities and the villages. Since 2002, ADAP (Association for the Development of Protected Areas) is supporting IBA in beekeeping activities and conservation of natural resources in general of the BKZ. ADAP is a Swiss NGO based in Geneva founded in 1997. Its objective is to generate additional incomes for local communities in order to gain their support for environmental conservation.

The current forest resources of the Mlele BKZ and their threats are not well known. According to the National Forest Policy 1998 as well as the Forest Act, No. 14 of 2002, a forest management plan, including a harvesting plan, needs to be established for all forest reserves. The current harvesting plan for Mulele Hills FR was established in 2014 relying heavily on the former forest inventory conducted by the Ministry of Natural Resources and Tourism (MNRT) through the Forestry and Beekeeping Division in 2005 and on other secondary data from 2006-2014 such as annual district forest harvesting reports, tree felling licences, transit passes and illegal harvesting reports (TFS 2014). No field survey has been carried out to verify and to update these secondary data. Therefore, the current harvesting plan of the Mulele Hills FR might not be realistic.

In 2004, ADAP has mandated a tree survey of Mlele BKZ in order to know the melliferous trees and their abundance, the use of forest products and the human disturbances (Mwangulango 2004). A discussion with Oscar Malembeka, a retired Village Game Scout (VGS) who accompanied the late Nathan Mwangulango in the vegetation survey allowed clarifying their methodology used. A main objective of the 2004 survey was to establish 30 permanent vegetation plots for monitoring vegetation changes over time. Unfortunately, this objective was not respected in the second vegetation survey carried out in 2013 (Kayombo et al. 2013) where new 42 new plots were established following a different methodology. Therefore, any direct comparison of the results of the tree survey of 2013 with 2004 is hampered and any possible vegetation change cannot be detected.

The **objective** of the new vegetation survey in 2018 was twofold:
a) to detect possible vegetation changes within the Mlele BKZ and
b) to assess the current stage of the vegetation of the whole Mlele BKZ and its currents threats (baseline).

To establish any possible vegetation changes we decided to resurvey only vegetation plots from the 2004 survey. This choice was based on the following reasons:

- Longer time period for detecting vegetation changes (2004-2018);
- The regeneration potential of the plots (trees/shrubs < 20 cm have only been recorded in 2004 (Mwangulango 2004);
- Exact location of plots (coordinates) not clear in the 2013 vegetation survey (Kayombo et al. 2013);
- Unfortunately we could not meet the authors of the 2013 vegetation survey to clarify their methodology applied.

The vegetation plots from 2004 have been established mostly along roads often nearby the boundary where access is easier. Possible vegetation changes in individual plots are barely representative for the whole area of Mlele BKZ.

Therefore, we decided in addition to the survey of permanent vegetation plots from 2004, to outline two vegetation transects of about 10 km length each for improving the data set and

having a sound baseline for 2018. Thereby, also all human disturbances could be assessed along the transects.

This third vegetation survey has been assessed by two field trips. Dr. Urs Bloesch led the team composed of Frank Mbago (22 June to 26 June 2018) and the VGSs Dicksoni Malembeka, Johni Gervasi, and Felesiano Futakamba in the first field trip from 22 June to 2 July 2018. Lucile Daudet together with the VGS completed in a second field trip from 9 to 14 September 2018 the two vegetation transects.

The rainfall was exceptionally high in the preceding rainy season resulting in very tall and dense grass layer. Some of the larger Mbugas¹ were still flooded but most of the few existing tracks were passable by car during the first field trip while during the second field trip towards the end of the dry season in September the grass layer and the soil were dry along the transects.

2) Study area

Miombo woodland is the dominating vegetation type of unimodal rainfall areas in southeastern Africa (Smith & Allen 2004). Miombo woodlands are the most extensive vegetation type in Tanzania (Shirima et al. 2014) thereby playing a key role as carbon source and sink. Miombo woodlands are characterized by a distinct and often continuous grass layer and an open to closed tree canopy layer and often annually burnt. Miombo are dominated by the genera *Brachystegia* and *Julbernadia* with *Brachystegia spiciformis* and *Julbernadia globiflora* as the most common tree species (Frost 1996).

Usually we distinguish between floristically rich “wetter miombo” of the higher rainfall areas (>1000 mm per annum) and floristically more poor “drier miombo” (<1000 mm per annum). Miombo woodlands have a low soil nutrient content, are well drained, highly leached, acidic and low in organic matter (Frost 1996). The timber and non-timber products from the miombo woodlands are essential for the livelihoods of millions of people living inside and outside the miombo woodlands (Malaisse 1997).

The BKZ in Mlele District is located northwest of Inyonga township, along the main road to Mpanda and the road to Rukwa Game Reserve Headquarters covering an area of about 850,000 ha. Mlele BKZ is surrounded by a dense network of protected areas in Katavi Region (see Fig. 1). The altitude of Mlele BKZ ranges between 1200 to 1500 m a.s.l. from the lower land in the northeast to the upper land (hilltop, plateau) in the southwest separated by the escarpment.

Miombo woodlands are the predominating vegetation type in Mlele BKZ (Mwangulango 2004, Kayombo et al. 2013) covering more than 90% of the area. The average annual rainfall is estimated to oscillate between 1000 and 1200 mm. The miombo type is transitional between “drier” and “wetter” miombo. The seasonally waterlogged mbugas are mainly covered with grasslands and occasionally with few trees and shrubs (wooded grasslands). Narrow riverine forests occur along the permanent and seasonal streams and some evergreen forest patches exist on deeper soils in depressions (south of Ngaramira area and at Masigo according to Kayombo et al. 2013).

Mlele BKZ is part of the Mulele Hills Forest Reserve of 519,295 ha which was gazetted in 1953 (TFS 2014). Based on the memorandum of understanding (MoU) with the Ministry of

¹ Seasonally waterlogged grasslands and grass savannas; term frequently used in Tanzania.

Natural Resources and Tourism (MNRT) in 2010, IBA is managing the beekeeping zone on behalf of the MNRT. The MoU is currently under revision as requested by the Tanzania Forest Services Agency (TFS). TFS was established in 2011 as an executive agency with the mandate of managing national forests (natural and plantations) and bee resources in a sustainable manner by ensuring sustainable supply of various forest and bee products and services, stable ecosystem and maintaining biological diversity. Mlele FR is also a Game Controlled Area (GCA) under the auspice of the Tanzania Wildlife Management Authority (TAWA) from the MNRT. Moreover, hunting companies which have also anti-poaching and road maintenance tasks are active in Mlele FR.




Fig. 1. Beekeeping zone and other protected areas of Mlele District, in Katavi Region (Mermod 2016).

The Wakonongo are the original tribe of Mlele District. Traditionally they were hunter-gatherer practising slash-and-burn agriculture having an intimate relation with the miombo woodlands (Mermod 2016).

3) Methods

3.1 Data collection

Vegetation plots

13 plots out of the 30 vegetation plots from 2004 (Mwangulango 2004) have been resurveyed from 23 to 28 June (see Appendix A and D). For time reasons, facility in access was considered in the selection of the plots for this survey.

As far as possible the methodology from 2004 was respected. Note that Mwangulango (2004) did not explicitly specify the minimal Diameter at Breast Height (DBH²) recorded for the lower diameter < 20 cm class (probably DBH \geq 7 cm); we recorded all trees with a DBH \geq 3 cm. Unfortunately, Mwangulango (2004) gives only the total number of tree species of the upper and lower diameter class without indicating the exact DBH of each tree recorded. Therefore, the basal area and the class diameter distribution for each tree species could not be established for the 2004 vegetation survey.

For time and practical reasons the size of the vegetation plots from 2004 was reduced to a circular sample plot of 0.1 ha (radius of 17.85m). Recording of all trees and shrubs in a rectangular plot of 100 x100 m (plot numbers 1 - 15) or 50 x 50 m (plot numbers 16 – 30) as in the 2004 survey seemed to us very difficult to realise and is also very time-consuming. Therefore, we used smaller and circular plots which are easier to survey using a measure tape from the centre of the plot. Finally, the 2018 survey recorded the following parameters:

- Location and general site description: we use a handheld GPS (Garmin 12 XL) to find the coordinates of the centre of the 2004 plot and to record the elevation; the nearest tree was selected and marked with white colour (in 2004 blue colour was used to mark the trees of the plot); slope and slope exposure were determined with a Büchi compass (P3252).
- Systematically, each plot centre was photographed in each cardinal direction (Canon powershot S95) resulting in 52 photographs in total (see example in Fig.2).
- Each tree/shrub species with a DBH \geq 3 cm has been identified and its diameter recorded. This allows establishing for each species the diameter class distribution. The diameter of multi-stemmed trees has been taken for each single stem (DBH \geq 3 cm) for calculating the basal area (ba). For calculating the tree density per ha, multi-stemmed trees are considered as one tree.
- The height of the dominant trees has been estimated visually (ocular estimation) using a levelling rod of 2 m height as a reference at the base of the trunk.
- The canopy cover³ of the dominant trees and of the understorey has been recorded assuming that all trees/shrubs are in full foliage; in addition the vegetation cover of the herbaceous layer has been estimated and the dominant species recorded according to the cover-abundance scale from Braun-Blanquet (1964).

² Diameter at breast height, or DBH, is a standard method of expressing the diameter of the trunk or bole of a standing tree at 1.3m height.

³ Percentage of the soil surface covered by vertically projected tree and shrubs crowns.




Fig. 2. Example of visual illustration through photographs from the centre of the plot in each cardinal direction (here vegetation plot 3).

Transects

Google Earth has been used for locating the transects. The main vegetation types (canopy cover) within Mlele BKZ and possible access constraints for sampling along the transects have been quickly identified. Transects should be outlined perpendicular to the main mountain ridges thereby following a variable slope gradient with different vegetation types. Finally, two transects have been established with an azimuth of 47° or 227° considering possible access by existing tracks (see Appendix D). The first transect is located on the upper land starting at the boundary road about 300 m southeast from the IBA camp (azimuth of 47°). A nearby track east of the transect was facilitating the access for the first five kilometres. A second transect was located on the lower land starting from the main Mpanda road (azimuth of 227°). The nearby German road has similar direction thereby facilitating access the sampling on transect for about ten kilometers.

Along the transects the Point-Centered Quarter Method (PCQM) has been used (Mueller-Dombois, & Ellenberg 1974). Four quarters are established at the sampling point through a cross formed by two lines. One line is the compass direction (47° or 227°) and the second a line running perpendicular to the compass direction through the sampling point. The distance

to the mid-point of the nearest tree from the sampling point is measured in each quarter (see Fig. 3). The interval between two sampling points has been set at 300 steps systematically.

50 points have been recorded along each transect resulting in a transect length of about 10 km. The first 27 points of the first transect on upper land have been recorded from 29 June to 1 July 2018, while the remaining 23 points have been recorded from 13 to 14 September 2018. The second transect on lower land nearby German road has been recorded on 2 July 2018 (first 8 points) and completed for the remaining points (9 – 50) from 9 to 12 September 2018. This plotless method has the advantage in that they do not require laying out plot boundaries what saves considerable time.

We used a handheld GPS (Garmin 12 XL) to record the coordinates of each sampling point. All tree/shrub species and their DBH were recorded separately for two size categories: larger trees ($DBH \geq 20$ cm) and smaller trees/shrubs ($3 \geq DBH < 20$ cm). The height of the dominant trees has been estimated visually (ocular estimation) using a levelling rod of 2 m height as a reference at the base of the trunk. Slope and slope exposure were determined with a Büchi compass (P3252). Systematically, each direction of the cross marking the four quarters was photographed from the sampling point (Canon powershot S95) resulting in 200 photographs for each transect (see example in Fig. 2). The data for the two transects are given in Appendix B.

The parameters obtained with the PCQM method are (Mueller-Dombois, & Ellenberg 1974):

- Species;
- Density (from mean distance);
- For each tree its diameter class diameter distribution;
- Basal area (ba) and dominance for each species;
- Absolute frequency (as the occurrence of a species at a sampling point).

The tree basal area⁴ of the surrounding stand was also estimated additionally in a simple but efficient way using the Bitterlich method (1948). Please note that the Bitterlich method was only applied in the vegetation survey from 22 June to 2 July 2018.

The Bitterlich method counts trees (shrubs) in a circle from the sampling point with an angle-gauge (wooden stick of 50 cm length with a fixed small metal sheet with four angular widths at its end). Only trees (shrubs) that are larger in diameter than the smallest angular width (counting factor $k = 1$) are included in the count.

⁴ Tree basal area is the cross-sectional area (over the bark) at breast height (1.3 metres above the ground) measured in square meters.



Fig. 3. PCQM method: Distance to the mid-point of the nearest tree from the sampling point is measured in each quarter and tree species and diameter is recorded.

All human ecosystem disturbances have been recorded on both sides along the transects by two VGSs (total sampling width of 80 m). Each VGS surveyed one half (i.e. 40 m) of the total width of the belt transect. Disturbances could include logging (saw pits, stumps), tree barking (for bee hives, ropes), trapping, infrastructures (camps), encroachments, and grazing by cattle. Please note that fire signs were not recorded since most of the miombo woodlands are burnt annually.

3.2 Data analysis

If plants could not be identified in the field, voucher specimens were identified by Frank Mbago at the herbarium of the University of Dar es Salaam. A plant list of all recorded specimen has been elaborated (see Appendix C).

For the permanent vegetation plots, mean α -diversity (mean number of species of subunits), γ -diversity (total species diversity for the landscape and β -diversity ($\beta = \gamma / \alpha$) were defined (see Whittaker 1970).

For the vegetation transects, absolute frequency i.e., the occurrence of a species at the sampling points has been calculated as follows:

(N points with species/total points) x 100.

Moreover, the species dominance as product of mean ba and number of species per ha has been calculated.

As an example the diameter class distribution of the highly searched Mninga (*Pterocarpus angolensis*) timber species has been established in a diagram.

4) Results

Vegetation plots 2004 / 2018

The vegetation data of the 13 permanent vegetation plots resurveyed in 2018 are presented in Appendix A (data 2004 and 2018). Most of the marked trees of the resurveyed vegetation plots from 2004 had still their blue marks.

As outlined a direct comparision between the vegetation surveys of the permanent plots from 2018 and 2004 is difficult and not possible for some parameters due to:

- a) Different size of the plots in 2004 and 2018;
- b) The 2004 survey distinguishes only between DBH smaller or larger (equal) 20 cm and do not indicate the exact DBH of a tree/shrub (basal area cannot be defined);
- c) Minimal diameter for the lower diameter class not specified in the 2004 survey.




Fig. 4. Large *Brachystegia spiciformis* tree on the upper land.

Table 1 below shows different diversity indices of the species richness for the vegetation surveys in 2004 and 2018.

Table 1: Species diversity indices

| Species diversity indices | Survey 2004 (plot size 1 ha and 0.25 ha) | Survey 2018 (plot size 0.1 ha) |
|---------------------------|---|-----------------------------------|
| Mean α-diversity | 34.2 | 19.4 |
| β-diversity | 3.2 | 4.3 |
| γ-diversity | 108 | 83 |

Table 2 below lists the tree/shrub density for the vegetation surveys in 2004 and 2018.

Table 2: Tree and shrub density

| Tree/shrub density | Survey 2004 (plot size 1 ha and 0.25 ha) | Survey 2018 (plot size 0.1 ha) |
|---|---|-----------------------------------|
| Mean tree/shrub density (DBH < 20 cm) per ha | - ⁵ | 508 (317 ⁶) |
| Mean tree density (DBH ≥ 20 cm) per ha | 141 (30) | 118 (42 ⁶) |

Vegetation baseline using transects

Table 3 below includes the tree/shrub density, the mean basal area, estimated mean tree height, and estimated standing volume per ha for the miombo landscape on upper land (transect 1) and lower land (transect 2).

Table 3: Forest parameters

| | Transect 1 upper land | | | Transect 2 lower land | | |
|---|-----------------------|----------|-------|-----------------------|----------|-------|
| | Ø < 20cm | Ø ≥ 20cm | Total | Ø < 20cm | Ø ≥ 20cm | Total |
| Tree/shrub density/ha | 414.77 | 84.3 | | 324.2 | 50.4 | |
| Mean basal area/ha (m ²) | 2.69 | 8.91 | 11.60 | 2.43 | 4.95 | 7.38 |
| Estimated mean tree height (m) ⁷ | 6 | 11 | | 5 | 8 | |
| Estimated standing volume/ha (m ³) ⁸ | 8.1 | 49.0 | 57.1 | 6.1 | 19.8 | |

⁵ Minimal diameter recorded not specified.

⁶ Standard deviation

⁷ Mean tree height for the canopy and the understorey on upper land estimated at 11 m (from mean dominant tree height: 14.6 m) and 6 m, respectively; mean tree height for the canopy and the understorey on lower land estimated at 8 m (from mean dominant tree height: 9.6 m) and 5 m, respectively.

⁸ The standing volume was estimated using a mean tree coefficient of 0.5.

The absolute frequency of the first five tree/shrub species for the lower (DBH < 20cm) and the upper diameter class (DBH ≥ 20cm) is presented for both transects in tables 4 and 5 below.

Table 4: Absolute frequency for transect 1 on upper land

| $\emptyset < 20\text{cm}$ | | $\emptyset \geq 20\text{cm}$ | |
|------------------------------------|--------------------|---|--------------------|
| Species | Absolute frequency | Species | Absolute frequency |
| <i>Brachystegia glaucescens</i> | 28 | <i>Julbernardia globiflora</i> | 48 |
| <i>Diplorhynchus condylocarpon</i> | 28 | <i>Brachystegia spiciformis</i> | 46 |
| <i>Pseudolachnostylis maproun.</i> | 26 | <i>Parinari curatellifolia</i> | 28 |
| <i>Hexalobus monopetalus</i> | 24 | <i>Brachystegia manga/utilis</i> ⁹ | 18 |
| <i>Pterocarpus angolensis</i> | 22 | <i>Anisophyllea pomifera</i> | 18 |

Table 5: Absolute frequency for transect 2 on lower land

| $\emptyset < 20\text{cm}$ | | $\emptyset \geq 20\text{cm}$ | |
|------------------------------------|--------------------|----------------------------------|--------------------|
| Species | Absolute frequency | Species | Absolute frequency |
| <i>Diplorhynchus condylocarpon</i> | 32 | <i>Brachystegia manga/utilis</i> | 38 |
| <i>Pseudolachnostylis maproun.</i> | 32 | <i>Brachystegia glaucescens</i> | 32 |
| <i>Brachystegia manga/utilis</i> | 22 | <i>Julbernardia globiflora</i> | 32 |
| <i>Pterocarpus angolensis</i> | 22 | <i>Pterocarpus angolensis</i> | 22 |
| <i>Combretum fragrans</i> | 20 | <i>Pericopsis angolensis</i> | 16 |
| <i>Hymenocardia acida</i> | 20 | | |

The dominance values for the first five tree/shrub species for the lower (DBH < 20cm) and the upper diameter class (DBH ≥ 20cm) is presented for both transects in tables 6 and 7 below.

Table 6: Species dominance for transect 1 on upper land

| $\emptyset < 20\text{cm}$ | | $\emptyset \geq 20\text{cm}$ | |
|------------------------------------|---------|----------------------------------|---------|
| Species | Ranking | Species | Ranking |
| <i>Pterocarpus angolensis</i> | 1 | <i>Brachystegia spiciformis</i> | 1 |
| <i>Pseudolachnostylis maproun.</i> | 2 | <i>Julbernardia globiflora</i> | 2 |
| <i>Brachystegia glaucescens</i> | 3 | <i>Brachystegia manga/utilis</i> | 3 |
| <i>Phyllocosmus leimareanus</i> | 4 | <i>Parinari curatellifolia</i> | 4 |
| <i>Brachystegia spiciformis</i> | 5 | <i>Erythrophleum africanum</i> | 5 |

Table 7: Species dominance for transect 2 on lower land

| $\emptyset < 20\text{cm}$ | | $\emptyset \geq 20\text{cm}$ | |
|------------------------------------|---------|----------------------------------|---------|
| Species | Ranking | Species | Ranking |
| <i>Diplorhynchus condylocarpon</i> | 1 | <i>Brachystegia glaucescens</i> | 1 |
| <i>Pseudolachnostylis maproun.</i> | 2 | <i>Brachystegia manga/utilis</i> | 2 |
| <i>Brachystegia manga/utilis</i> | 3 | <i>Julbernardia globiflora</i> | 3 |
| <i>Oldfieldia dactylophylla</i> | 4 | <i>Pterocarpus angolensis</i> | 4 |
| <i>Pterocarpus angolensis</i> | 5 | <i>Pericopsis angolensis</i> | 5 |

⁹ The species is either *Brachystegia manga* or *Brachystegia utilis*.

The diameter class distribution of Mnenga (*Pterocarpus angolensis*) is shown in Fig. 5 below.




Fig. 5. Diameter class distribution of *Pterocarpus angolensis*.

The vegetation surveys in 2018 allowed completing substantially the former tree/shrub list for Mlele BKZ from Mwanguulango (2004). The checklist now includes in total 123 trees/shrubs with their scientific and Konongo names including also species identified from opportunistic sampling (see Appendix C). This quite exhaustive plant list including for most species the vernacular name will be very useful for any vegetation work in future.

All human disturbances recorded on a width of 80 m along transect 1 on upper land and transect 2 on lower land are summarised in tables 8 and 9 below.

Table 8: Human disturbances along transect 1 on upper land

| Type of disturbances | Magnitude (quantitiy) | Position along the transect |
|--|---|-----------------------------|
| Logging | 1 <i>Pterocarpus a.</i> cut (>10 yrs) | Point 3, +75 steps |
| | 1 <i>Monotes katangensis</i> recorded cut for pole | Point 10 |
| | 1 <i>Pterocarpus a.</i> cut (>10 yrs) (10 steps) | Point 12, +10 steps |
| Building constructions | Beekeepers' camp | Point 3, +30 steps |
| Tree cutting for harvesting wild honey | 1 <i>Pterocarpus angolensis</i> cut for wild honey harvesting (4 yrs) | Point 25, +130 steps |
| | 1 tree cut for honey | Point 46, +20 steps |
| Total disturbances | 6 site disturbances, 5 trees cut | |

Table 9: Human disturbances along transect 2 on lower land

| Type of disturbances | Magnitude (quantitiy) | Position along the transect |
|--|--|--|
| Logging | 3 <i>Pseudolachnostylis m.</i> cut for poles (\varnothing 12-15 cm; >5 yrs) 5 <i>Pterocarpus angolensis</i> cut for timber (>5 yrs) (90 steps) 1 <i>Julbernardia g.</i> tree barking for bee hives (\varnothing 30 cm; >5 yrs) 1 <i>Pterocarpus angolensis</i> cut for timber (\varnothing 35 cm; >5 yrs) 1 <i>Julbernardia globiflora</i> cut for timber? (\varnothing 25 cm; >5 yrs) 1 tree cut for timber (>10 years) 1 tree cut for timber (>5 years) | Point 6, +30-50 steps Point 6, +90 steps Point 6, +145 steps Point 7, +190 steps Point 9, +280 steps Point 10, +20 steps Point 11, +10 steps Point 11, +64 steps Point 11, +130 steps Point 11, +175 steps Point 11, +240 steps Point 14, +28 steps |
| Tree cutting for harvesting wild honey | 1 <i>Brachystegia utilis</i> cut for wild honey harvesting 1 tree cut for wild honey harvesting (> 5 years) 1 tree cut for wild honey harvesting (> 5 years) | Point 7, +50 steps Point 10, +80 steps Point 10, +290 steps |
| Tree barking | 1 <i>Julbernardia g.</i> tree barking for bee hives (\varnothing 22 cm; >5 yrs) 1 tree debarked for beehives (>3 years) | Point 6, +70 steps Point 11, +280 steps |
| Tree cutting for harvesting wild honey | 2 trees cut for honey 1 tree cut for honey 1 tree cut for honey 1 tree cut for honey (>3 years) | Point 20, +280 steps Point 32, +10 steps Point 33, +170 steps Point 45, +180 steps |
| Total disturbances | 22 site disturbances; 29 trees cut or debarked | |

5) Discussions

Vegetation plots 2004 / 2018

The species composition of the Mlele BKZ is rather typical for the wetter Miombo type. *Combretum* and *Acacia* species typical for the drier Miombo are not very abundant. The species richness of the Mlele BKZ comprises 83 species for the 2018 vegetation survey (108 in 2004). This γ -diversity is quite similar with the 86 trees and shrubs recorded by Backéus et al. (2006) in their survey of miombo woodlands in Kilosa District having similar rainfall.

The higher mean α -diversity and γ -diversity of the 2004 vegetation plots is certainly codetermined by the considerably larger plot size of 1 ha (plots 1 – 15) respectively 0.25 ha (plots 16 – 30) what results also in a higher total species richness (γ -diversity). The mean tree density per ha for the upper diameter class (DBH \geq 20 cm) is quite similar in the 2004 and 2018 vegetation surveys with 141 and 118 trees, respectively.

The 13 vegetation plots resurveyed will serve as permanent plots for monitoring and evaluating any vegetation changes at site level.

Vegetation baseline using transects

In total 800 trees and shrubs have been recorded by the PCQM. The two transects have been recorded within ten days, i.e. in average 10 points were sampled per day. The VGSSs have been trained on the spot and they mastered the different steps of the methods quite quickly.

Tree density, basal area, dominant tree height, and estimated standing volume are significantly higher on upper land compared to the miombo woodlands below the escarpment (see table 3). This may point to higher soil fertility on upper land and to more and longer inundated wetlands (Mbugas) at the foothill of the escarpment. The higher tree density in the vegetation plots compared to the area covered by the transects might be explained by the selection of the vegetation plots which were placed in 2004 in rather closed and undisturbed miombo vegetation (see table 2 and 3).

The most frequent trees/shrubs in Mlele BKZ in the lower diameter class (DBH < 20cm) are *Diplorhynchus condylocarpon*, *Pseudolachnostylis maprouneifolia*, and *Pterocarpus angolensis* (see tables 4-5). According to Backéus et al. (2006) these three species are quite fire-tolerant what may explain their prevalence in the annually burnt Mlele BKZ. In the upper diameter class (DBH ≥ 20cm) *Julbernardia globiflora* and *Brachystegia manga/utilis* are the most frequent tree species in Mlele BKZ.

The diameter class distribution of *Pterocarpus angolensis* in Fig. 5 shows a distinct underrepresentation for DBH > 18 cm. This IUCN near-threatened species is highly sought for timber (see Jew et al. 2016 and Schwartz et al. 2002). On the other hand, the regeneration of *Pterocarpus angolensis* is quite abundant apart from the 10.5 cm diameter class. Young *Pterocarpus angolensis* from the lower diameter class (DBH < 20cm) are widespread in Mlele BKZ and are amongst the five most frequent and dominant species (see tables 4-7). This is line with Backéus et al. (2006) who reported a good natural regeneration capacity of Mnenga, possibly because individuals below logging size have a good seed set. By contrast Boaler (1966) and Schwartz et al. (2002) reported very low recruitment levels for Mnenga.

Ecosystem health

The vegetation surveys revealed that the Miombo habitats of Mlele BKZ are intact. Natural regeneration of key species is quite abundant. In addition, the survey along the belt transects of about 20 km length and 80 m width allowed to assess all human disturbances within an area covering roughly 160 ha. We believe that the location and the total length of transects may give quite representative results for Mlele BKZ.

Fires are the prevailing disturbance in Mlele BKZ but their signs have not been recorded during the survey since most of Mlele BKZ is burnt annually including early and late dry season fires. It seems that there is no clear prescribed burning management by IBA for Mlele BKZ. The VGSSs are applying both early and late dry season burning although their impact on the tree:grass ratio is adverse thereby either favouring grazers or browsers (see Bloesch 1999, 2008). Moreover, hot late dry season fires may affect melliferous trees and their natural regeneration.

In total 28 human site disturbances has been assessed including 32 trees cut (many of them *Pterocarpus angolensis*) along the two transects. About 80% of the disturbances happened on the lower land what may be explained by the proximity of the principal road Inyonga to Mpanda.

It is remarkable that all recorded human disturbances occurred several years back. No sign of recent poaching was detected and no logging happened in the last five years in the two belt transects. This is quite amazing considering the greatly increased human pressure in recent years on protected areas of the Katavi/Rukwa ecosystem. Illegal activities such as poaching, logging of high value hardwood timber trees, encroachment of farmers's fields, and grazing by Sukuma cattle are more and more threatening the ecosystems (Mermod 2016). As a result, most of these protected areas managed by the central government are increasingly degraded.

On the other hand, the joint management of Mlele BKZ between IBA (ADAP) and the MNRT based on a strong participatory approach with the local communities has allowed an effective protection of the ecosystems in the beekeeping zone. It would be interested to further analyse to what extent this is the result of the efficient protection system in place with regular patrolling of Mlele BKZ by the VGSs or the Project's support of income generation activities (mainly modern beekeeping) and awareness raising.

6) Conclusions

The use of the PCQM along the vegetation transects is an appropriate and quite simple method to assess the vegetation baseline and in addition also all human disturbances along the transects. This method can now also be applied by the trained VGSs to monitor and evaluate changes in the vegetation and in human disturbances. It is suggested to use this method also for the neighbouring protected areas what would allow to compare ecosystem (vegetation) health of protected areas under different managements.

The Miombo habitats of Mlele BKZ are well protected. No sign of recent poaching was detected and no logging happened in the last five years in the two belt transects as a result of the participatory management of the beekeeping zone with regular patrolling, support of beekeeping activities and awareness raising campaign by the Project.

Regarding the management of Mlele BKZ it is suggested to elaborate a vegetation map with the main vegetation units. The vegetation map is an important management tool and will also serve to elaborate a fire management plan for the zone. This map could be also an important planning tool for the district authorities.

7) Recommendations

Based on the observations of the field mission the following recommendations to IBA / ADAP are made:

- 1) Use the transect methodology for monitoring and evaluating any vegetation changes at site level.
- 2) Put the coordinates of the PCQM points on a map to localise the two transects.

- 3) Elaborate a vegetation map of Mlele BKZ with its main vegetation units. The vegetation data from the 2018 field campaign could be used as ground truth for elaborating the map.
- 4) IBA in consultation with TFS should elaborate a fire management plan for Mlele BKZ including a seasonal prescribed burning plan.
- 5) We suggest using the transect methodology with the PCQM for the planned vegetation assessment of Rungwa FR. Appropriate transects should be located using google earth. The trained VGSs from Mlele BKZ will participate at the beginning of the survey until the VGSs from Rungwa FR have been trained on the spot in the methodology.
- 6) Amend the ecological information of the Miombo species and assess the use of the Miombo trees/shrub species for the adjacent local communities (importance for their livelihoods).
- 7) It is suggested to include Tanzanian students in joint BSc/MSc studies with Swiss students.
- 8) Elaborate a paper for a research journal with a focus on the transect methodology using PCQM and its potential for rapidly establishing the vegetation baseline for monitoring and evaluating the vegetation.

8) Bibliography

- Backéus, I., Petterson, B., Strömquist, L. & Ruffo, C. (2006) Tree communities and structural dynamics in miombo (Brachystegia-Julbernardia) woodland, Tanzania. *Forest Ecology and Management*, 230, 171-178.
- Bitterlich, W. (1948) Die Winkelzählprobe. Allg. Forst- und Holzwirtsch. Ztg. 59, 4-5.
- Bloesch, U. (1999) Fire as a tool in the management of a savanna / dry forest reserve in Madagascar. *Applied Vegetation Science*, 2, 117-124.
- Bloesch, U. (2002) The dynamics of thicket clumps in the Kagera savanna landscape, East Africa. PhD thesis N° 14386, Swiss Federal Institute of Technology Zurich (ETH). Shaker, Aachen.
- Bloesch, U. (2008) Thicket clumps: A typical feature of the Kagera savanna landscape, East Africa. *Journal of Vegetation Science*, 19, 31-44.
- Boaler, S.B. (1966). Ecology of a miombo site, Lupa north forest reserve, Tanzania: II. Plant communities and seasonal variation in the vegetation. *Journal of Ecology*, 54, 465–479.
- Braun-Blanquet, J. (1964) Pflanzensoziologie, Grundzüge der Vegetationskunde, 3rd edition. Springer, Wien, New York.
- Frost, P. G. H. (1996). The ecology of miombo woodlands. In B. Campbell (Ed.), *The Miombo in transition: Woodlands and welfare in Africa* (pp. 11–57). Bogor, Indonesia: CFIOR.

Jew, K.K., Dougill, A.J., Sallu, S.M., O'Connell, J. & Benton, T.G. (2016) Miombo woodland under threat: Consequences for tree diversity and carbon storage. *Forest Ecology and Management*, 361, 144-153.

Kayombo, C.J., Mpinga, I. & Natai, H. (2013) Melliferous Status and Activities Endangering Tree Species. Composition and Diversity Survey of Mlele Bee Keeping Zone [Mbkz], in Mlele District, Katavi Region-Tanzania. Consultancy carried out by the Forestry Training Institute-Olmotonyi, Arusha-Tanzania.

Malaisse, F. (1997) Se nourrir en forêt claire africaine. Approche écologique et nutritionnelle. Centre Technique de Coopération Agricole et Rurale, Wageningen / Les presses agronomiques de Gembloux.

Mermod, S. (2016) What does the future hold for the Mlele District Forest Reserves? The interaction of the social, political, and ecological spheres in Tanzania's Katavi Region. MSc in geography. Faculty of Geosciences and Environment (GSE), University of Lausanne.

Mueller-Dombois, D. & Ellenberg, H. (1974) Aims and methods of vegetation ecology. John Wiley & Sons, New York, London, Sydney and Toronto.

Mwanguango, N. (2004) Vegetation survey in Mlele beekeeping zone. April 2004 - July 2004.

Schwartz, M.W., Caro, T.M. & Banda-Sakala, T. (2002) Assessing the sustainability of harvest of Pterocarpus angolensis in Rukwa Region, Tanzania. *Forest Ecology and Management*, 170, 259-269.

Shirima, D.D., Totland, O., Munishi, P.K.T. and Moe, S.R. (2014) Relationships between tree species richness, evenness and aboveground carbon storage in montane forests and miombo woodlands of Tanzania. *Basic and Applied Ecology*, 16, 239–249.

Smith, P. & Allen, Q. (2004) Field guide to the trees and shrubs of the Miombo woodlands. Royal Botanic Gardens, Kew.

Strømgaard, P. (1992). Immediate and long-term effects of fire and ash-fertilization on a Zambian miombo woodland soil. *Agriculture, Ecosystems & Environment*, 41, 19–37.

TFS (2014) Forest Harvesting Plan for Mulele Forest Reserve. Mlele District – Katavi Region. Five Years Plan for 2014/2015-2018/2019. Zonal Manager TFS Western Zone Office, Ministry of Natural Resources and Tourism, Tabora.

Whittaker, R.H. (1970) Communities and ecosystems. Macmillan Co., Collier-Macmillan Ltd., London.

Appendix A: Permanent vegetation plots

Vegetation plot 1 (recorded 25/6/2018)

Coordinates: S 6.64310° / E 31.73194° (white-marked *Brachystegia glaucescens*)

Elevation: 1365 m.a.s.l.; slope: +/- flat

Dominant tree height: 16 m;

Vegetation cover: canopy (8-16m): 35%; understorey (2-8m): 10%; herbaceous layer (0-2m): 75%

Herbaceous layer: *Hyparrhenia newtonii* 2, *Urelytrum digitatum* 2, *Margaritaria discoidea* 1, *Smilax anceps* 1, *Zonetrache inamoena* 1

Photos: 5972, 5973, 5974, 5975

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|------------------------------------|--|------------------|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Pseudolachnostylis maproun.</i> | 4,11,15,11,11,12,3,7/7,9,5, 5/14/15,4,11,8,3=15 | 25/5=1 | 37 | 16 |
| <i>Diplorhynchus condylocarpon</i> | 4,4/9,3,3,10,10,7,4,3, 5/7/10/17,8,7/10,22=13 | 22=1 | 48 | 6 |
| <i>Julbernardia globiflora</i> | 15,8,10,16,12,11=6 | 28,36/31,20,30=4 | 18 | 64 |
| <i>Brachystegia spiciformis</i> | 4,10,9,4,10,5,5,4=8 | - | 23 | 5 |
| <i>Dichrostachys cinerea</i> | 3/5/7,4/8,5/3,3,3,3/3,4/3=7 | - | 24 | - |
| <i>Erythrophleum africanum</i> | 7,12,7,4,4,9=6 | - | 33 | 5 |
| <i>Brachystegia glaucescens</i> | 10,5,9,10,15=5 | - | 38 | 3 |
| <i>Flacourtie indica</i> | 3/3,5,3=3 | - | 8 | - |
| <i>Isoberlinia angolensis</i> | 14,5,4=3 | - | - | 11 |
| <i>Vitex doniana</i> | 8,3=2 | 26=1 | 1 | 3 |
| <i>Annona senegalensis</i> | 4/3=1 | - | 5 | - |
| <i>Azanza garckeana</i> | 3/3=1 | - | - | - |
| <i>Brachystegia utilis/manga</i> | - | 21=1 | - | - |
| <i>Burkea africana</i> | 5=1 | - | - | - |
| <i>Cassipourea mollis</i> | 3=1 | - | - | 6 |
| <i>Lannea schimperi</i> | - | 26=1 | 2 | 11 |
| <i>Pterocarpus angolensis</i> | 9=1 | - | 3 | 2 |
| <i>Kigelia africana</i> | 15=1 | - | - | - |
| <i>Anisophyllea pomifera</i> | - | - | - | 3 |
| <i>Bobgunnia madagascar.</i> | - | - | 12 | - |
| <i>Brachystegia boehmii</i> | - | - | 12 | 3 |
| <i>Combretum collinum</i> | - | - | 1 | 4 |
| <i>Combretum fragrans</i> | - | - | 1 | - |
| <i>Combretum psidiodoides</i> | - | - | 1 | - |
| <i>Diospyros cornii</i> | - | - | - | 1 |
| <i>Garcinia huillensis</i> | - | - | - | 1 |
| <i>Hexalobus monopetalus</i> | - | - | 2 | |
| <i>Hymenocardia acida</i> | - | - | 9 | - |
| <i>Lonchocarpus capassa</i> | - | - | - | 2 |
| <i>Mallotus oppositifolius?</i> | - | - | 14 | - |
| <i>Monotes engleri</i> | - | - | - | 1 |
| <i>Ozoroa insignis</i> | - | - | 1 | - |
| <i>Parinari curatellifolia</i> | - | - | - | 4 |
| <i>Pericopsis angolensis</i> | - | - | - | 1 |
| <i>Psorospermum febrifugum</i> | - | - | 1 | - |
| <i>Rothmannia engleriana</i> | - | - | 1 | - |
| <i>Schrebera trichoclada</i> | - | - | 3 | - |
| <i>Securidaca longepeduncul.</i> | - | - | 1 | - |
| <i>Strychnos cocculoides</i> | - | - | 2 | 1 |
| <i>Strychnos spinosa</i> | - | - | - | 2 |

| | | | | |
|------------------------------|------------|-----------|------------|------------|
| <i>Terminalia kaiseriana</i> | - | - | 5 | 2 |
| <i>Ximenia caffra</i> | - | - | 2 | - |
| <i>Msinde</i> | - | - | - | 1 |
| Tree density (per ha) | 740 | 90 | 308 | 158 |
| Species richness | | 18 | | 39 |

Vegetation plot 2 (recorded 25/6/2018)

Coordinates: S 6.60126° / E 31.74842° (white-marked *Julbernardia globiflora*)

Elevation: 1380 m.a.s.l.; slope: +/- flat

Dominant tree height: 22(24) m;

Vegetation cover: canopy (8-22{24}m): 55%; understorey (2-8m): 5%; herbaceous layer (0-2m): 85%

Herbaceous layer: *Anthephora elongata* 2, *Hyparrhenia newtonii* 2, *Urelytrum digitatum* 2, *Indigofera rhynchosarpa* 1, *Tetracera masuiana* 1

Photos: 5976, 5977, 5978, 5979

Remark: Several dead trees

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|-----------------------------------|---------------|--|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Julbernardia globiflora</i> | - | 33,62,27/28,32/39,25,21, 29,37,40=9 | 1 | 56 |
| <i>Ximenia caffra</i> | 10,2,9,15=4 | - | 9 | - |
| <i>Brachystegia spiciformis</i> | - | 39,31,34=3 | 54 | - |
| <i>Diplorhynchus condylocarp.</i> | 8,10/11/16=2 | - | 18 | - |
| <i>Pseudolachnostylis mapr.</i> | 14/18,13=2 | - | 19 | 2 |
| <i>Cassipourea mollis</i> | 14=1 | - | 7 | 3 |
| <i>Combretum zeyheri</i> | 5=1 | - | 1 | - |
| <i>Erythrophleum africanum</i> | - | 23=1 | 2 | 12 |
| <i>Hexalobus monopetalus</i> | 7=1 | - | 34 | 1 |
| <i>Lannea schimperi</i> | 7=1 | - | 3 | 2 |
| <i>Terminalia kaiseriana</i> | 5=1 | - | 2 | - |
| <i>Brachystegia glaucescens</i> | - | 32=1 | 10 | 7 |
| <i>Annona senegalensis</i> | - | - | 1 | - |
| <i>Bobgunnia madagascar.</i> | - | - | 15 | - |
| <i>Cassia abbreviata</i> | - | - | 1 | - |
| <i>Combretum collinum</i> | - | - | 4 | 10 |
| <i>Combretum psidiooides</i> | - | - | 1 | 1 |
| <i>Commiphora africana</i> | - | - | 1 | - |
| <i>Dichrostachys cinerea</i> | - | - | 3 | - |
| <i>Garcinia huillensis</i> | - | - | - | 1 |
| <i>Lonchocarpus capassa</i> | - | - | 4 | - |
| <i>Oldfieldia dactylophylla</i> | - | - | 11 | - |
| <i>Pericopsis angolensis</i> | - | - | 1 | 9 |
| <i>Pterocarpus angolensis</i> | - | - | 11 | 11 |
| <i>Rothmannia engleriana</i> | - | - | 1 | - |
| <i>Strychnos cocculoides</i> | - | - | - | 5 |
| <i>Kaguaguva</i> | - | - | 2 | 1 |
| Tree density (per ha) | 130 | 140 | 216 | 121 |
| Species richness | | 12 | | 27 |

Vegetation plot 3 (recorded 26/6/2018)

Coordinates: S 6.50590° / E 31.72323° (white-marked *Julbernardia globiflora*)

Elevation: 1220 m.a.s.l.; slope: 1-2%, slope exposure: 300°

Dominant tree height: 24 m;

Vegetation cover: canopy (8-24m): 55%; understorey (2-8m): 30%; herbaceous layer (0-2m): 80%

Herbaceous layer: *Margaritaria discoidea* 2, *Hyparrhenia newtonii* 2, *Friesodielsia obovata* 1

Photos: 5984, 5985, 5986, 5987

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|-----------------------------------|--------------------|------------------------|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Julbernardia globiflora</i> | - | 32,42,36,36,50,33,41=7 | - | 46 |
| <i>Diplorhynchus condylocarp.</i> | 8,9,5,9/6,10=5 | 24=1 | 16 | 16 |
| <i>Hexalobus monopetalus</i> | 4,4,8/12/15,11,6=5 | - | 25 | 2 |
| <i>Burkea africana</i> | 13,11,11,15/12=4 | - | 1 | 1 |
| <i>Monanthotaxis discolor</i> | 12,8,8,7=4 | - | 8 | - |
| <i>Pseudolachnostylis mapr.</i> | 10,15,10/10=3 | 27=1 | 13 | 7 |
| <i>Pavetta stuhlmannii</i> | 3,4,3=3 | - | - | - |
| <i>Hymenocardia acida</i> | 7,3=2 | - | 18 | 6 |
| <i>Phyllanthus engleri</i> | 4,3=2 | - | 5 | - |
| <i>Schrebera trichoclada</i> | 3,5=2 | - | 3 | 4 |
| <i>Brachystegia utilis/manga</i> | - | 33=1 | 4 | 16 |
| <i>Bobgunnia madagascar.</i> | 11=1 | - | 2 | - |
| <i>Dalbergia nitidula</i> | 9/14=1 | - | - | - |
| <i>Erythrophleum africanum</i> | 7=1 | - | 8 | 15 |
| <i>Margaritaria discoidea</i> | 4=1 | - | - | - |
| <i>Lannea schimperi</i> | - | 24/14=1 | 5 | 7 |
| <i>Pterocarpus angolensis</i> | - | 31=1 | 6 | 8 |
| <i>Pterocarpus tinctorius</i> | - | 9/53=1 | - | 8 |
| <i>Pericopsis angolensis</i> | - | 24/26=1 | 6 | 5 |
| <i>Terminalia cf. mollis</i> | 4=1 | - | - | - |
| <i>Afzelia quanzensis</i> | - | - | - | 3 |
| <i>Annona senegalensis</i> | - | - | 5 | 1 |
| <i>Borassus aethiopium</i> | - | - | 1 | - |
| <i>Brachystegia spiciformis</i> | - | - | 7 | 3 |
| <i>Cassia abbreviata</i> | - | - | - | 2 |
| <i>Catunaregam spinosa</i> | - | - | 2 | - |
| <i>Crossopteryx febrifuga</i> | - | - | 1 | - |
| <i>Combretum collinum</i> | - | - | - | 1 |
| <i>Combretum molle</i> | - | - | - | 1 |
| <i>Combretum psidiooides</i> | - | - | 1 | - |
| <i>Flacourtie indica</i> | - | - | 3 | - |
| <i>Friesodielsia obovata</i> | - | - | 21 | - |
| <i>Garcinia huillensis</i> | - | - | 1 | - |
| <i>Kigelia africana</i> | - | - | - | 1 |
| <i>Lonchocarpus capassa</i> | - | - | 2 | 2 |
| <i>Mallotus oppositifolius?</i> | - | - | 44 | 11 |
| <i>Markhamia obtusifolia</i> | - | - | 8 | 2 |
| <i>Parinari curatellifolia</i> | - | - | 1 | - |
| <i>Rothmannia engleriana</i> | - | - | 3 | - |
| <i>Sclerocarya birrea</i> | - | - | - | 1 |
| <i>Stereospermum kunthianum</i> | - | - | - | 2 |
| <i>Terminalia kaiseriana</i> | - | - | 2 | 4 |
| <i>Vangueria madagascariens.</i> | - | - | 3 | - |
| <i>Vitex payos</i> | - | - | 2 | - |
| Tree density (per ha) | 350 | 140 | 227 | 175 |
| Species richness | | 20 | | 40 |

Vegetation plot 5 (recorded 26/6/2018)

Coordinates: S 6.58370° / E 31.88180° (white-marked *Erythrophleum africanum*)

Location: Mtalazia

Elevation: 1180 m.a.s.l.; slope: +/- flat

Dominant tree height: 17-18m m;

Vegetation cover: canopy (10-17/18m): 40%; understorey (2-10m): 15%; herbaceous layer (0-2m): 75%

Herbaceous layer: *Hyparrhenia newtonii* 2, *Julbernardia globiflora* 2, *Pennisetum polystachyon*

Photos: 5988, 5989, 5990, 5991

Remark: 1 old termitaria

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|------------------------------------|--------------------------|---------------|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Pseudolachnostylis mapr.</i> | 15,8,13,11,4,9,18,16/5=8 | 22,/24/24=1 | 22 | 21 |
| <i>Brachystegia utilis/manga</i> | 4=1 | 25,36,42,26=4 | 10 | 41 |
| <i>Terminalia kaiseriana</i> | 5,6,11,17=4 | 24=1 | 11 | - |
| <i>Diplorhynchus condylocarp.</i> | 4,18,11/12,4,4=5 | - | 10 | 3 |
| <i>Julbernardia globiflora</i> | 6=1 | 27,22,33=3 | 8 | 14 |
| <i>Hymenocardia acida</i> | 5,10,13/17=3 | - | 13 | - |
| <i>Lannea schimperi</i> | - | 20/8,20,23=3 | - | 8 |
| <i>Brachystegia boehmii?</i> | - | 42,38=2 | - | - |
| <i>Cassia abbreviata</i> | 6,13=2 | - | - | - |
| <i>Combretum molle</i> | 11,6=2 | - | - | - |
| <i>Erythrophleum africanum</i> | 3,17=2 | - | 21 | 9 |
| <i>Schrebera trichoclada</i> | 4,5=2 | - | 1 | - |
| <i>Bridelia duvigneaudii</i> | 3=1 | - | - | - |
| <i>Burkea africana</i> | 3=1 | - | - | 1 |
| <i>Dalbergia nitidula</i> | 13=1 | - | - | - |
| <i>Hexalobus monopetalus</i> | - | 21=1 | 1 | 4 |
| <i>Pericopsis angolensis</i> | - | 36=1 | - | 1 |
| <i>Vitex doniana</i> | - | 21=1 | 1 | - |
| <i>Vitex fischeri</i> | 6=1 | - | - | - |
| <i>Ziziphus mucronata</i> | - | 12/26/6=3 | 1 | 1 |
| <i>Borassus aethiopium</i> | - | - | 1 | - |
| <i>Brachystegia glaucescens</i> | - | - | 8 | 15 |
| <i>Bobgunnia madagascar.</i> | - | - | 9 | - |
| <i>Cassipourea mollis</i> | - | - | 4 | - |
| <i>Combretum collinum</i> | - | - | 3 | 6 |
| <i>Combretum zeyheri</i> | - | - | 4 | - |
| <i>Crossopteryx febrifuga</i> | - | - | 1 | - |
| <i>Diospyros cornii</i> | - | - | 2 | - |
| <i>Flacourtie indica</i> | - | - | 3 | - |
| <i>Isoberlinia angolensis</i> | - | - | 1 | 7 |
| <i>Mallotus oppositifolius?</i> | - | - | 12 | - |
| <i>Monanthotaxis discolor</i> | - | - | 6 | - |
| <i>Monotes engleri?</i> | - | - | 5 | 6 |
| <i>Oldfieldia dactylophylla</i> | - | - | 1 | - |
| <i>Piliostigma thonningii</i> | - | - | 1 | 1 |
| <i>Pterocarpus angolensis</i> | - | - | 1 | 5 |
| <i>Pterocarpus tinctorius</i> | - | - | - | 3 |
| <i>Rothmannia engleriana</i> | - | - | 4 | - |
| <i>Strychnos cocculoides</i> | - | - | 6 | 6 |
| <i>Strychnos potatorum</i> | - | - | - | 1 |
| <i>Tamarindus indica</i> | - | - | 1 | - |
| <i>Vangueria madagascariensis.</i> | - | - | 2 | - |
| <i>Vitex payos</i> | - | - | 1 | - |

| | | | | |
|------------------------------|------------|------------|------------|------------|
| Tree density (per ha) | 340 | 200 | 175 | 153 |
| Species richness | 20 | | 37 | |

Vegetation plot 10 (recorded 23/6/2018)

Coordinates: S 6.82835° / E 31.70690° (white-marked *Julbernardia globiflora*)

Elevation: 1420 m.a.s.l.; slope: 3-4%, slope exposure: 100°

Dominant tree height: 20 m;

Vegetation cover: canopy (8-20m): 50%; understorey (2-10m): 25%; herbaceous layer (0-2m): 65%

Herbaceous layer: *Anthephora elongata* 2, *Hyparrhenia newtonii* 2, *Hexalobus monopetalus* 2,

Tetracera masuiana 1

Photos: 5926, 5928, 5929, 5930

Remark: 1 large termitaria

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|-----------------------------------|--------------------|---------------|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Diplorhynchus condylocarp.</i> | 15,17,15,5,3,3,7=7 | | 34 | 3 |
| <i>Brachystegia glaucescens</i> | 3,3=2 | 27,24=2 | 9 | 2 |
| <i>Julbernardia globiflora</i> | - | 47,38,39,51=4 | 21 | 49 |
| <i>Lonchocarpus capassa</i> | 4,4,3=3 | - | 2 | - |
| <i>Hexalobus monopetalus</i> | 3,15=2 | | 57 | 1 |
| <i>Uapaca nitida</i> | 9,7=2 | - | - | - |
| <i>Rhus vulgaris</i> | 4,7=2 | - | - | - |
| <i>Anisophyllea pomifera</i> | 4=1 | - | 12 | 8 |
| <i>Azanza garckeana</i> | 4=1 | - | - | - |
| <i>Brachystegia microphylla</i> | 3=1 | - | - | - |
| <i>Cassia singueana</i> | 5=1 | - | - | - |
| <i>Combreutia molle</i> | - | 26=1 | 5 | - |
| <i>Lannea schimperi</i> | 5=1 | - | 1 | - |
| <i>Memecylon flavovires</i> | 5=1 | - | - | - |
| <i>Pericopsis angolensis</i> | 3=1 | - | 2 | 6 |
| <i>Pseudolachnostylis mapr.</i> | 9=1 | - | 2 | 2 |
| <i>Uapaca kirkiana</i> | 7=1 | - | 10 | 1 |
| <i>Vitex doniana</i> | - | 34=1 | 2 | - |
| <i>Albizia versicolor</i> | - | - | - | 1 |
| <i>Annona senegalensis</i> | - | - | 1 | - |
| <i>Bobgunnia madagascar.</i> | - | - | 1 | - |
| <i>Brachystegia spiciformis</i> | - | - | 9 | 2 |
| <i>Brachystegia longifolia?</i> | - | - | 7 | 2 |
| <i>Brachystegia taxifolia?</i> | - | - | 39 | 16 |
| <i>Burkea africana</i> | - | - | 3 | - |
| <i>Cassia abbreviata</i> | - | - | - | 1 |
| <i>Cassipourea mollis</i> | - | - | 5 | 10 |
| <i>Dalbergia nitidula</i> | - | - | 4 | 3 |
| <i>Erythrophleum africanum</i> | - | - | 3 | 5 |
| <i>Flacourtie indica</i> | - | - | - | 1 |
| <i>Markhamia obtusifolia</i> | - | - | 1 | 1 |
| <i>Monanthotaxis discolor</i> | - | - | 2 | - |
| <i>Monotes engleri?</i> | - | - | - | 1 |
| <i>Ochna holstii</i> | - | - | 2 | - |
| <i>Oldfieldia dactylophylla</i> | - | - | 1 | - |
| <i>Parinari curatellifolia</i> | - | - | 63 | 7 |
| <i>Pterocarpus angolensis</i> | - | - | 10 | 1 |
| <i>Pterocarpus tinctorius</i> | - | - | 1 | - |
| <i>Rothmannia engleriana</i> | - | - | 2 | - |

| | | | | |
|------------------------------|------------|-----------|------------|------------|
| <i>Schrebera trichoclada</i> | - | - | 1 | - |
| <i>Vitex payos</i> | - | - | 2 | - |
| <i>Ximenia caffra</i> | - | - | 3 | - |
| Tree density (per ha) | 270 | 80 | 317 | 123 |
| Species richness | | 18 | | 36 |

Vegetation plot 11 (recorded 23/6/2018)

Coordinates: S 6.79391° / E 31.65827° (white-marked *Julbernardia globiflora*)

Elevation: 1475 m.a.s.l.; slope: 3%, slope exposure: 70°

Dominant tree height: 18-20 m;

Vegetation cover: canopy (6-18/20m): 35%; understorey (2-6m): 10%; herbaceous layer (0-2m): 45%

Herbaceous layer: *Hyparrhenia newtonii* 2-3, *Zonetrichie inamoena* 1-2, *Psychotria eminiana* 1

Photos: 5922, 5923, 5924, 5925

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|-----------------------------------|---------------|----------------------------------|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Julbernardia globiflora</i> | - | 29,37,39,33,35,31,24,25, 50=9 | - | 62 |
| <i>Hexalobus monopetalus</i> | 4,5,4,4,4=5 | - | 30 | - |
| <i>Phyllocosmus leimareanus</i> | 18,8,10=3 | 27=1 | - | - |
| <i>Brachystegia spiciformis</i> | 3=1 | 77,28,47=3 | 25 | 11 |
| <i>Anisophyllea pomifera</i> | 18,9=2 | 22=1 | 10 | 7 |
| <i>Erythrophleum africanum</i> | 4,8=2 | - | 2 | 10 |
| <i>Dichrostachys cinerea</i> | 3,5=2 | - | 13 | - |
| <i>Pseudolachnostylis mapr.</i> | 3,4=2 | - | 13 | - |
| <i>Diplorhynchus condylocarp.</i> | 3=1 | - | 30 | - |
| <i>Brachystegia glaucescens</i> | 3=1 | - | 35 | 1 |
| <i>Burkea africana</i> | - | 22=1 | 2 | 1 |
| <i>Lannea schimperi</i> | 11=1 | - | 3 | 1 |
| <i>Pterocarpus angolensis</i> | 8=1 | - | 2 | 1 |
| <i>Terminalia kaiseriana</i> | 6=1 | - | 7 | - |
| <i>Vitex doniana</i> | 7=1 | - | 7 | 2 |
| <i>Albizia versicolor</i> | - | - | 7 | 1 |
| <i>Annona senegalensis</i> | - | - | 1 | - |
| <i>Bobgunnia madagascar.</i> | - | - | 8 | - |
| <i>Brachystegia</i> sp. | - | - | - | 1 |
| <i>Combretum molle</i> | - | - | 2 | - |
| <i>Catunaregam spinosa</i> | - | - | 1 | - |
| <i>Dalbergia nitidula</i> | - | - | 2 | - |
| <i>Garcinia huillensis</i> | - | - | - | 1 |
| <i>Lonchocarpus capassa</i> | - | - | 3 | - |
| <i>Ochna holstii</i> | - | - | 8 | 12 |
| <i>Parinari curatellifolia</i> | - | - | 2 | 7 |
| <i>Protea</i> sp. | - | - | 2 | - |
| <i>Rothmannia engleriana</i> | - | - | 6 | - |
| <i>Strychnos cocculoides</i> | - | - | 3 | 4 |
| <i>Uapaca kirkiana</i> | | | 2 | - |
| <i>Vitex payos</i> | - | - | 1 | - |
| <i>Ximenia caffra</i> | - | - | 11 | - |
| <i>Mtonga</i> | - | - | 1 | - |
| Tree density (per ha) | 230 | 150 | 239 | 122 |
| Species richness | | 15 | | 32 |

Vegetation plot 12 (recorded 23/6/2018)

Coordinates: S 6.74754° / E 31.62633° (white-marked *Bobgunnia madagascariensis*)

Elevation: 1360 m.a.s.l.; slope: 1-2%, slope exposure: 50°

Dominant tree height: 20-22 m;

Vegetation cover: canopy (10-20/22m): 55%; understorey (2-10m): 10%; herbaceous layer (0-2m): 65%

Herbaceous layer: *Hyparrhenia newtonii* 2, *Hexalobus monopetalus* 1, *Hyparrhenia rufa* 1, *Zonetrichia inamoena* 1

Photos: 5934, 5935, 5936, 5937

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|-----------------------------------|----------------------|-------------|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Diplorhynchus condylocarp.</i> | 10,7,6,3,5,4,7,7,3=9 | - | 19 | -- |
| <i>Phyllocosmus leimareanus</i> | 12,24,11,11,13/11=5 | - | - | - |
| <i>Brachystegia spiciformis</i> | 4,7,16=3 | 29/46,34=2 | 5 | 15 |
| <i>Julbernardia globiflora</i> | - | 32,61,34=3 | 2 | 44 |
| <i>Hexalobus monopetalus</i> | 3,6,7=3 | - | 12 | 2 |
| <i>Anisophyllea pomifera</i> | 10,19,15=2 | - | 10 | 10 |
| <i>Bobgunnia madagascar.</i> | 13,14=2 | - | 4 | - |
| <i>Brachystegia microphylla</i> | 3,5=2 | - | | |
| <i>Erythrophleum africanum</i> | 3/3=1 | 21=1 | 8 | 8 |
| <i>Lannea schimperi</i> | - | 20,21=2 | 1 | 3 |
| <i>Albizia antunesiana</i> | 14=1 | - | - | - |
| <i>Brachystegia floribunda?</i> | 4=1 | - | - | - |
| <i>Bridelia scleroneura</i> | 5=1 | - | - | - |
| <i>Cassia abbreviata</i> | 10=1 | - | 1 | - |
| <i>Dichrostachys cinerea</i> | 5=1 | - | 1 | - |
| <i>Lonchocarpus capassa</i> | 3=1 | - | 6 | - |
| <i>Parinari curatellifolia</i> | 3=1 | | 2 | 3 |
| <i>Chrysophyllum bangweolen.</i> | 4=1 | | - | - |
| <i>Strychnos innocua</i> | 4=1 | - | 1 | 2 |
| <i>Vitex fischeri</i> | 4=1 | | - | - |
| <i>Ximenia caffra</i> | 3=1 | - | 3 | - |
| <i>Albizia versicolor</i> | - | - | 2 | 1 |
| <i>Afzelia quanzensis</i> | - | - | - | 1 |
| <i>Annona senegalensis</i> | - | - | 2 | - |
| <i>Brachystegia bohmii</i> | - | - | 1 | 22 |
| <i>Cassipourea mollis</i> | - | - | 2 | - |
| <i>Dalbergia nitidula</i> | - | - | 5 | - |
| <i>Flacourtie indica</i> | - | - | 2 | - |
| <i>Garcinia huillensis</i> | - | - | 3 | 1 |
| <i>Kigelia africana</i> | - | - | 1 | - |
| <i>Mallotus oppositifolius?</i> | - | - | 7 | - |
| <i>Monanthotaxis discolor</i> | - | - | 2 | - |
| <i>Monotes engleri?</i> | - | - | 1 | 2 |
| <i>Ochna holstii</i> | - | - | 16 | 8 |
| <i>Pericopsis angolensis</i> | - | - | - | 3 |
| <i>Pseudolachnostylis mapr.</i> | - | - | 8 | 7 |
| <i>Pterocarpus angolensis</i> | - | - | 2 | - |
| <i>Rothmannia engleriana</i> | - | - | 3 | - |
| <i>Schrebera trichoclada</i> | - | - | 1 | - |
| <i>Securidaca longependuncul.</i> | - | - | 1 | - |
| <i>Strychnos cocculoides</i> | - | - | 1 | - |
| <i>Terminalia kaiseriana</i> | - | - | 2 | 1 |
| <i>Trichilia emetica</i> | - | - | 1 | - |
| <i>Uapaca kirkiana</i> | - | - | 7 | - |

| | | | | |
|------------------------------|------------|-----------|------------|------------|
| <i>Uapaca nitida</i> | - | - | 3 | - |
| <i>Vitex doniana</i> | - | - | 7 | - |
| <i>Vitex payos</i> | - | - | 1 | - |
| <i>Mwasya</i> | - | - | - | 13 |
| <i>Mtonga</i> | - | - | 1 | - |
| Tree density (per ha) | 380 | 80 | 157 | 146 |
| Species richness | | 21 | | 42 |

Vegetation plot 13 (recorded 25/6/2018)

Coordinates: S 6.69570° / E 31.60895° (white-marked *Erythrophleum africanum*)

Elevation: 1310 m.a.s.l.; slope: 1%, slope exposure: 245°

Dominant tree height: 18-20 m;

Vegetation cover: canopy (8-18/20m): 55%; understorey (2-8m): 20%; herbaceous layer (0-2m): 15%

Herbaceous layer: *Adenodolichos punctatus* 1, *Hyparrhenia newtonii* 1, *Bothriochloa bladhii* 1,

Psychotria eminiana 1, *Urelytrum digitatum* 1, *Tetracera masuiana* 1, *Zonetriche inamoena* 1

Photos: 5964, 5965, 5966, 5967

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|-----------------------------------|--------------------------------------|--|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Hexalobus monopetalus</i> | 4,13,3,7,7,6,7,4,4,5,3,3,3,7 = 14 | 33,62,27/28,32/39,25,21, 29,37,40=9 | 20 | 8 |
| <i>Diplorhynchus condylocarp.</i> | 11,3,6/8,3/3,8/8,10/6,12,5/8 = 8 | - | 57 | 1 |
| <i>Hymenocardia acida</i> | 3,3,3,3,4,3,5,3=8 | - | 43 | 1 |
| <i>Julbernardia globiflora</i> | 10,4,8,6,5,3,3=7 | 53=1 | 13 | 35 |
| <i>Brachystegia spiciformis</i> | 13,13,9,18=4 | 32,20,98=3 | 24 | 18 |
| <i>Maprounea africana</i> | 9,6,4,10,6,14,9=7 | | - | - |
| <i>Pseudolachnostylis mapr</i> | 3,9,7,3,7,6,4=7 | - | 6 | 3 |
| <i>Brachystegia utilis/manga</i> | 14=1 | 24,21,51,32/10,14=5 | - | - |
| <i>Terminalia kaiseriana</i> | 19,15,9,4=4 | - | 23 | 12 |
| <i>Uapaca nitida</i> | 17/13,9,13=3 | - | - | - |
| <i>Cassipourea mollis</i> | 4,3,3=3 | - | 8 | - |
| <i>Erythrophleum africanum</i> | 3,9,15=3 | - | 46 | 9 |
| <i>Maranthes floribunda</i> | 12/9,3,7=3 | - | - | - |
| <i>Parinari curatellifolia</i> | 4/4,5,4=3 | - | 9 | 7 |
| <i>Strychnos innocua</i> | 4,5=2 | - | 14 | - |
| <i>Albizia antunesiana</i> | 14=1 | - | - | - |
| <i>Bridelia duvigneaudii</i> | 4=1 | - | - | - |
| <i>Bobgunnia madagascar.</i> | 12=1 | - | 9 | - |
| <i>Combretum zeyheri</i> | 4=1 | | 3 | - |
| <i>Dalbergia nitidula</i> | 4=1 | - | 8 | - |
| <i>Lannea schimperi</i> | 4=1 | - | 15 | 3 |
| <i>Margaritaria discoidea</i> | 9=1 | - | - | - |
| <i>Monanthotaxis discolor</i> | 3=1 | - | - | - |
| <i>Oldfieldia dactylophylla</i> | 5=1 | - | 10 | 1 |
| <i>Securidaca longependuncul.</i> | 5=1 | - | | |
| <i>Vitex doniana</i> | 3=1 | - | 4 | 14 |
| <i>Afzelia quanzensis</i> | - | - | 1 | - |
| <i>Albizia versicolor</i> | - | - | 1 | - |
| <i>Anisophyllea pomifera</i> | - | - | 1 | - |
| <i>Annona senegalensis</i> | - | - | 4 | - |
| <i>Borassus aethiopum</i> | - | - | 1 | - |
| <i>Brachystegia boehmii</i> | - | - | 26 | 8 |
| <i>Cassia abbreviata</i> | - | - | 1 | - |

| | | | | |
|-----------------------------------|------------|------------|------------|------------|
| <i>Catunaregam spinosa</i> | - | - | 1 | - |
| <i>Combretum collinum</i> | - | - | - | 6 |
| <i>Combretum fragrans</i> | - | - | 1 | - |
| <i>Combretum molle</i> | - | - | 6 | - |
| <i>Mallotus oppositifolius?</i> | - | - | 85 | - |
| <i>Markhamia obtusifolia</i> | - | - | 1 | - |
| <i>Monanthotaxis discolor</i> | - | - | 15 | - |
| <i>Garcinia huillensis</i> | - | - | 1 | - |
| <i>Pericopsis angolensis</i> | - | - | 19 | 6 |
| <i>Phyllanthus engleri</i> | - | - | 3 | - |
| <i>Pterocarpus angolensis</i> | - | - | 1 | 8 |
| <i>Pterocarpus tinctorius</i> | - | - | 2 | 1 |
| <i>Rothmannia engleriana</i> | - | - | 4 | - |
| <i>Schrebera trichoclada</i> | - | - | 9 | - |
| <i>Strychnos cocculoides</i> | - | - | 2 | - |
| <i>Strychnos</i> sp. | - | - | - | 1 |
| <i>Uapaca kirkiana</i> | - | - | 4 | - |
| <i>Vangueria madagascariensis</i> | - | - | 1 | - |
| <i>Vitex payos</i> | - | - | 3 | 2 |
| <i>Ximenia caffra</i> | - | - | 1 | - |
| <i>Ziziphus mucronata</i> | - | - | 3 | 1 |
| <i>Mtonga</i> | - | - | 2 | 1 |
| Tree density (per ha) | 880 | 180 | 511 | 146 |
| Species richness | | 26 | | 46 |

Vegetation plot 14 (recorded 25/6/2018)

Coordinates: S 6.66932° / E 31.64857° (white-marked *Julbernardia globiflora*)

Elevation: 1375 m.a.s.l.; slope: +/- flat

Dominant tree height: 22 m;

Vegetation cover: canopy (8-22m): 40%; understorey (2-8m): 15%; herbaceous layer (0-2m): 65%

Herbaceous layer: *Hyparrhenia newtonii* 2, *Indigofera hirsuta* 1, *eminiana* 1, *Tetracera masuiana* 1

Photos: 5968, 5969, 5970, 5971

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|-----------------------------------|---------------------|----------------------------------|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Julbernardia globiflora</i> | - | 35,31,31,26,30,24/22,25, 36=8 | 32 | 58 |
| <i>Hexalobus monopetalus</i> | 4/4,5,5,4,5,3,3=7 | - | 1 | - |
| <i>Pseudolachnostylis mapr.</i> | 12,14,15,8/10/4,8=5 | - | 20 | 11 |
| <i>Erythrophleum africanum</i> | 16/10,11,3=3 | - | 15 | 12 |
| <i>Albizia gummiifera</i> | 14=1 | 41=1 | - | - |
| <i>Anisophyllea pomifera</i> | 3/5/9=1 | 41=1 | 1 | 10 |
| <i>Brachystegia spiciformis</i> | - | 26,23=2 | 68 | 53 |
| <i>Diplorhynchus condylocarp.</i> | 3,14/13=2 | - | 29 | - |
| <i>Combretum zeyheri</i> | 4/11,5=2 | - | - | - |
| <i>Brachystegia glaucescens</i> | 15/5=1 | 22=1 | 8 | 7 |
| <i>Combretum collinum</i> | 13=1 | - | 2 | 2 |
| <i>Dalbergia nitidula</i> | 12=1 | - | 10 | - |
| <i>Elaeodendron schweinfurth.</i> | 8=1 | - | - | - |
| <i>Lonchocarpus capassa</i> | 3=1 | - | - | - |
| <i>Terminalia kaiseriana</i> | 16=1 | - | 1 | - |
| <i>Albizia versicolor</i> | - | - | 2 | - |
| <i>Bobgunnia madagascar.</i> | - | - | 1 | - |
| <i>Burkea africana</i> | - | - | 1 | - |

| | | | | |
|---------------------------------|------------|------------|------------|------------|
| <i>Cassipourea mollis</i> | - | - | 1 | - |
| <i>Garcinia huillensis</i> | - | - | 1 | - |
| <i>Isoberlinia angolensis</i> | - | - | 4 | - |
| <i>Lannea schimperi</i> | - | - | 1 | 5 |
| <i>Mallotus oppositifolius?</i> | - | - | 6 | - |
| <i>Monanthotaxis discolor</i> | - | - | 1 | - |
| <i>Ochna holstii</i> | - | - | 30 | 16 |
| <i>Oldfieldia dactylophylla</i> | - | - | 2 | - |
| <i>Parinari curatellifolia</i> | - | - | 1 | 3 |
| <i>Phyllanthus engleri</i> | - | - | 4 | - |
| <i>Phyllocosmus leimareanus</i> | - | - | 1 | - |
| <i>Protea madiensis</i> | - | - | 11 | 2 |
| <i>Pterocarpus angolensis</i> | - | - | 5 | 4 |
| <i>Pterocarpus tinctorius</i> | - | - | 5 | - |
| <i>Schrebera trichoclada</i> | - | - | 4 | - |
| <i>Strychnos cocculoides</i> | - | - | 8 | - |
| <i>Vitex doniana</i> | - | - | - | 1 |
| <i>Vitex payos</i> | - | - | 8 | - |
| <i>Ximenia caffra</i> | - | - | 15 | 3 |
| <i>Mkalya</i> | - | - | 2 | - |
| <i>Mtonga</i> | - | - | 2 | - |
| Tree density (per ha) | 270 | 130 | 303 | 187 |
| Species richness | 15 | | 35 | |

Vegetation plot 15 (recorded 25/6/2018)

Coordinates: S 6.63017° / E 31.95254° (white-marked *Pterocarpus angolensis*, *Brachystegia spiciformis*?)

Elevation: 1245 m.a.s.l.; slope: 1-2%, slope exposure: 10°

Dominant tree height: 18(20) m;

Vegetation cover: canopy (10-18/20m): 35%; understorey (2-10m): 25%; herbaceous layer (0-2m): 65%

Herbaceous layer: *Hyparrhenia newtonii* 2-3, *Tristachya superba* 1-2, *Combretum molle* 1

Photos: 5992, 5993, 5994, 5995

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|-----------------------------------|-------------------------|---------------|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Julbernardia globiflora</i> | 5,5,10,10,6,4,19=7 | 36,38,29,41=4 | 59 | 35 |
| <i>Diplorhynchus condylocarp.</i> | 11/15,6,10,17,7,5,3,7=8 | - | 14 | 5 |
| <i>Brachystegia spiciformis</i> | 11,11,11,11,5,10,4=7 | 39,31,34=3 | 45 | 10 |
| <i>Rothmannia engleriana</i> | 4,5,5,4=4 | - | 1 | - |
| <i>Combretum molle</i> | 18,7,11=3 | - | - | 1 |
| <i>Crosopteryx febrifuga</i> | 11, 8,17=3 | - | 8 | 1 |
| <i>Pterocarpus angolensis</i> | 7,17=2 | 20=1 | 14 | 12 |
| <i>Ximenia americana</i> | 9,9,12=3 | - | - | - |
| <i>Combretum zeyheri</i> | 8,11=2 | - | 3 | 2 |
| <i>Dalbergia nitidula</i> | 9,6=2 | - | 2 | 12 |
| <i>Oldfieldia dactylophylla</i> | 4,10=2 | - | 1 | - |
| <i>Pericopsis angolensis</i> | 15,11=2 | - | 8 | 4 |
| <i>Pseudolachnostylis mapr.</i> | 10,3=2 | - | 16 | - |
| <i>Strychnos spinosa</i> | 5,6=2 | - | 1 | - |
| <i>Monotes katangensis</i> | 28/27=1 | - | 7 | 3 |
| <i>Brachystegia glaucescens</i> | 8=1 | - | 11 | 20 |
| <i>Burkea africana</i> | 8=1 | - | - | - |

| | | | | |
|-----------------------------------|------------|-----------|------------|------------|
| <i>Cassipourea mollis</i> | 6=1 | - | 1 | - |
| <i>Combretum collinum</i> | 6=1 | - | 3 | 1 |
| <i>Dichrostachys cinerea</i> | 5/4=1 | - | 4 | - |
| <i>Hexalobus monopetalus</i> | 9=1 | - | 11 | 3 |
| <i>Catunaregam spinosa</i> | 3=1 | - | 1 | - |
| <i>Erythrophleum africanum</i> | 10=1 | - | 4 | 1 |
| <i>Lannea schimperi</i> | 13=1 | - | 5 | 2 |
| <i>Parinari curatellifolia</i> | 5=1 | - | 10 | 7 |
| <i>Pterocarpus tinctorius</i> | 5=1 | - | 1 | 2 |
| <i>Terminalia kaiseriana</i> | 8=1 | - | 14 | 2 |
| <i>Afzelia quanzensis</i> | - | - | - | 1 |
| <i>Albizia versicolor</i> | - | - | 1 | - |
| <i>Bobgunnia madagascariensis</i> | - | - | 6 | 1 |
| <i>Commiphora africana</i> | - | - | 1 | - |
| <i>Flacourtie indica</i> | - | - | 1 | - |
| <i>Mallotus oppositifolius?</i> | - | - | 12 | - |
| <i>Schrebera trichoclada</i> | - | - | 1 | - |
| <i>Strychnos cocculoides</i> | - | - | 1 | - |
| <i>Strychnos innocua</i> | - | - | 9 | 1 |
| <i>Monotes engleri?</i> | - | - | - | 4 |
| <i>Pericopsis angolensis</i> | - | - | 8 | 4 |
| <i>Phyllanthus engleri</i> | - | - | 1 | - |
| <i>Tamarindus indica</i> | - | - | 1 | - |
| <i>Ximenia caffra</i> | - | - | 4 | - |
| <i>Mkalya</i> | - | - | 1 | - |
| <i>Monga</i> | - | - | 7 | 1 |
| Tree density (per ha) | 640 | 80 | 298 | 135 |
| Species richness | | 27 | | 38 |

Vegetation plot 16 (recorded 28/6/2018)

Coordinates: S 6.67304° / E 31.94294° (white-marked *Bobgunnia madagascar.*)

Elevation: 1205 m.a.s.l.; slope: 0-1%, slope exposure: 30%

Dominant tree height: 15 m;

Vegetation cover: canopy (8-15m): 30%; understorey (2-8m): 25%; herbaceous layer (0-2m): 55%

Herbaceous layer: *Hyparrhenia newtonii* 2, *Julbernardia globiflora* 1-2, *Mundulea sericea* 1-2

Photos: 5996, 5997, 5998, 5999

| Species | 2018 (0.1 ha) | | 2004 (0.25 ha) | |
|-----------------------------------|---|-------------|----------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Julbernardia globiflora</i> | 8,4,10,4,10,5/7,4,4,9,9,4, 4,5,10,6=15 | 22=1 | 6 | 5 |
| <i>Diplorhynchus condylocarp.</i> | 2,10/4,4,7,12,5,14,7,15,5= 10 | 6/3/28=1 | 51 | 3 |
| <i>Erythrophleum africanum</i> | 8,3,6,5,3,4,5,4,3,6,6=11 | - | 8 | 2 |
| <i>Strychnos pungens</i> | 10,17,10,11=4 | 24,22,23=3 | - | - |
| <i>Brachystegia glaucescens</i> | 17,10,10/15,9,14/13=5 | 25=1 | 1 | 6 |
| <i>Burkea africana</i> | 16,9,14,5=4 | - | 5 | - |
| <i>Pterocarpus angolensis</i> | 16,6,17,7=4 | 22=1 | - | 1 |
| <i>Bobgunnia madagascar.</i> | 5,17,17=3 | - | 5 | - |
| <i>Brachystegia spiciformis</i> | 13,4=2 | - | - | 3 |
| <i>Flacourtie indica</i> | 5,4=2 | - | 8 | - |
| <i>Maprounea africana</i> | 10,10=2 | - | - | - |
| <i>Monanthotaxis discolor</i> | 4,4=2 | - | - | - |
| <i>Ochna oxyphylla</i> | 7,10=2 | - | - | - |

| | | | | |
|-----------------------------------|------------|-------------|------------|------------|
| <i>Terminalia kaiseriana</i> | - | 28/7,22/6=2 | 1 | - |
| <i>Brachystegia stipulata?</i> | 3=1 | - | - | - |
| <i>Bridelia scleroneura</i> | 9=1 | - | - | - |
| <i>Afzelia quanzensis</i> | - | 24/23=1 | - | - |
| <i>Dalbergia nitidula</i> | 13=1 | - | 1 | - |
| <i>Dichrostachys cinerea</i> | 7=1 | - | - | - |
| <i>Hymenocardia acida</i> | 3=1 | - | 6 | - |
| <i>Monotes kantangensi</i> | 3=1 | - | - | - |
| <i>Pseudolachnostylis mapr.</i> | 15=1 | - | 3 | 3 |
| <i>Terminalia mollis</i> | 7=1 | - | - | - |
| <i>Vitex payos</i> | 10=1 | - | 2 | - |
| <i>Brachystegia boehmii</i> | - | - | 3 | 1 |
| <i>Bauhinia thonningii</i> | - | - | - | 1 |
| <i>Combretum collinum</i> | - | - | 1 | - |
| <i>Combretum fragrans</i> | - | - | 3 | 2 |
| <i>Combretum zeyheri</i> | - | - | 1 | - |
| <i>Commiphora africana</i> | - | - | - | 2 |
| <i>Diospyros cornii</i> | - | - | 2 | - |
| <i>Hexalobus monopetalus</i> | - | - | 1 | - |
| <i>Holarrhena pubescens</i> | - | - | 3 | - |
| <i>Isoberlinia angolensis</i> | - | - | - | 1 |
| <i>Kigelia africana</i> | - | - | - | 1 |
| <i>Lannea schimperi</i> | - | - | - | 1 |
| <i>Mallotus oppositifolius?</i> | - | - | 2 | - |
| <i>Monotes engleri?</i> | - | - | - | 2 |
| <i>Oldfieldia dactylophylla</i> | - | - | 3 | - |
| <i>Ozoroa insignis</i> | - | - | 1 | - |
| <i>Parinari curatellifolia</i> | - | - | 2 | 3 |
| <i>Protea madiensis</i> | - | - | 3 | - |
| <i>Rothmannia engleriana</i> | - | - | 4 | - |
| <i>Securidaca longependuncul.</i> | - | - | 1 | - |
| <i>Strychnos cocculoides</i> | - | - | 3 | 6 |
| <i>Strychnos spinosa</i> | - | - | 3 | - |
| <i>Kaulwampako</i> | - | - | - | 1 |
| Tree density/0.25 ha | | | 133 | 44 |
| Tree density/ha | 750 | 100 | 532 | 176 |
| Species richness | | 24 | | 37 |

Vegetation plot 17 (recorded 28/6/2018)

Coordinates: S 6.70926° / E 31.93098° (white-marked *Pterocarpus angolensis*);

Location: Ulaya

Elevation: 1205 m.a.s.l.; slope: 0-2%, slope exposure: 220°

Dominant tree height: 16 m;

Vegetation cover: canopy (8-16 m): 30%; understorey (2-8m): 15%; herbaceous layer (0-2m): 75%

Herbaceous layer: *Hyparrhenia newtonii* 3, *Combretum molle* 1

Photos: 6000, 6001, 6002, 6003

Remark: Along the border plot *Combretum molle* (5) and *Pseudolachnostylis maprouneifolia* (1) cut for poles; 1 large termitaria (*Macrotermes*)

| Species | 2018 (0.1 ha) | | 2004 (0.25 ha) | |
|--------------------------------|------------------------------------|-------------|----------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Terminalia kaiseriana</i> | 3,5,4/10,8,3,5,6,11,3,3/5,3 =11 | - | 6 | - |
| <i>Brachystegia stipulata?</i> | 12,9,14,8,7=5 | 22=1 | - | 1 |

| | | | | |
|-----------------------------------|-------------------|------------|------------|-----------|
| <i>Pseudolachnostylis mapr.</i> | 12,3,19,8/13,10=5 | 23,23,23=3 | - | 2 |
| <i>Maranthes floribunda</i> | 8=1 | 29/18,31=2 | - | - |
| <i>Combretum fragrans</i> | 3,18=2 | 9/23=1 | 6 | 8 |
| <i>Julbernardia globiflora</i> | - | 30,30,35=3 | - | 1 |
| <i>Bauhinia thonningii</i> | 8/5,11=2 | - | - | - |
| <i>Brachystegia glaucescens</i> | 8/12,9=2 | - | - | 1 |
| <i>Maytenus senegalensis</i> | 4/3,3=2 | - | - | - |
| <i>Bauhinia petersiana</i> | 10=1 | - | - | - |
| <i>Catunaregam spinosa</i> | 3/3=1 | - | 2 | - |
| <i>Combretum molle</i> | 15=1 | - | - | - |
| <i>Dichrostachys cinerea</i> | 4/3=1 | - | - | - |
| <i>Diplorhynchus condylocarp.</i> | 10=1 | - | 4 | 2 |
| <i>Ekebergia capensis</i> | 4=1 | - | - | - |
| <i>Pterocarpus tinctorius</i> | - | 31=1 | - | - |
| <i>Terminalia mollis</i> | 10=1 | - | - | - |
| <i>Uapaca kirkii</i> | 8/14=1 | - | - | - |
| <i>Vangueria madagascariensis</i> | 14=1 | - | - | - |
| <i>Ziziphus mucronata</i> | 14/3=1 | - | - | - |
| <i>Acacia polyacantha</i> | - | - | - | 2 |
| <i>Combretum psidiooides</i> | - | - | 5 | 3 |
| <i>Commiphora africana</i> | - | - | 1 | - |
| <i>Lannea schimperi</i> | - | - | - | 1 |
| <i>Parinari curatellifolia</i> | - | - | 8 | - |
| <i>Pericopsis angolensis</i> | - | - | - | 1 |
| <i>Phyllanthus engleri</i> | - | - | 4 | - |
| <i>Schrebera trichoclada</i> | - | - | 3 | - |
| <i>Stereospermum kunthianum</i> | - | - | 1 | - |
| <i>Ximenia caffra</i> | - | - | 2 | - |
| Tree density/0.25 ha | | | 42 | 22 |
| Tree density/ha | 400 | 110 | 168 | 88 |
| Species richness | | 20 | | 18 |

Vegetation plot 21 (recorded 24/6/2018)

Coordinates: S 6.72350° / E 31.74830° (white-marked *Pterocarpus angolensis*)

Location: Kikusi area

Elevation: 1500 m.a.s.l.; slope: +/- flat

Dominant tree height: 10(12) m;

Vegetation cover: canopy (6-10{12}m): 30%; understorey (2-6m): 20%; herbaceous layer (0-2m): 15%

Herbaceous layer: *Anthephora elongata* 1, *Andropogon schirensis* 1, *Zonetriche inamoena* 1

Photos: 5943, 5944, 5945, 5946

Remark: Several *Pterocarpus angolensi* debarked, two of them in vegetation plot

| Species | 2018 (0.1 ha) | | 2004 (1 ha) | |
|--------------------------------|---|---------------------|-------------|-------------|
| | DBH < 20 cm | DBH ≥ 20 cm | DBH < 20 cm | DBH ≥ 20 cm |
| <i>Julbernardia globiflora</i> | 4,9,5,5,5,11,10,10/13,5,12,3,12,12/13,5,9,8,7,5,10,9,8,11,3,10,4,6,3,7,3,12,7,7,9,11,3,9,7,6,5,4,7,3,10,9/5,4,14,12,4,10,10,3,10,13/7,5,10,3,7,6/7=58 | 43,32=2 | 19 | 3 |
| <i>Pterocarpus angolensis</i> | 12/15/13,13/12,10/7,1010/7,8,8/7,13,313,7,9,14,8,8,11,610/10/17,19=18 | 6/25/7,15/26/9,44=3 | 27 | - |
| <i>Monotes katangensi</i> | 5,6,3,3,3,3,6/6,6,4,3,9,3, | - | - | - |

| | | | | |
|--|---|-------------|------------|------------|
| | 3/5,4,5,11,16/17,4,4,7,11, 10=12 | | | |
| <i>Parinari curatellifolia</i> | 6,6,3,5,3,10=6 | - | 14 | 1 |
| <i>Rothmannia engleriana</i> | 4,3,5,4,3,3=6 | - | 3 | - |
| <i>Albizia antunesiana</i> | 5,3,3,6,4=5 | - | 4 | - |
| <i>Ochna afzelii</i> ssp. <i>afzelii</i> | 4,5,8/7/7,3,6=5 | | 2 | - |
| <i>Diplorhynchus condylocarp.</i> | 3,10,7=3 | - | - | - |
| <i>Erythrophleum africanum</i> | 8,17/7=2 | 26=1 | - | - |
| <i>Lannea schimperi</i> | 3,9=2 | - | 1 | - |
| <i>Anisophyllea pomifera</i> | 4=1 | - | 2 | - |
| <i>Brachystegia spiciformis</i> | 7=1 | - | 1 | 1 |
| <i>Combretum molle</i> | 7=1 | - | 7 | 1 |
| <i>Hexalobus monopetalus</i> | 8=1 | - | 8 | 3 |
| <i>Strychnos pungens</i> | 5=1 | - | - | - |
| <i>Multidentia crassa</i> | 9/6=1 | - | - | - |
| <i>Cassipourea mollis</i> | - | - | 3 | - |
| <i>Combretum psidoides</i> | - | - | 2 | - |
| <i>Dalbergia nitidula</i> | - | - | 3 | - |
| <i>Garcinia huillensis</i> | - | - | - | 1 |
| <i>Monanthotaxis discolor</i> | - | - | 4 | - |
| <i>Monotes engleri?</i> | - | - | 19 | 15 |
| <i>Ximenia caffra</i> | - | - | 5 | - |
| Tree density/0.25 ha | | | 124 | 25 |
| Tree density/ha | 1230 | 60 | 496 | 100 |
| Species richness | | 16 | | 18 |

Appendix B: Vegetation transects

Transect 1: Upper land (above escarpment)

Starting point road: S 6°80616 / E 31°66380; azimuth: 47° (towards vegetation plot 21)

Sampling point 1:

Coordinates: S 6°80492 / E 31°66522

Slope: 5%, slope exposure: 110°

Dominant tree height: 11 m

Basal area (Bitterlich, k=1): 10 m²

Photos: 6004, 6005, 6006, 6007

Disturbances/notes: stony blocks (boulders)

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia microphylla</i> | 22 | 6.80 |
| | <i>Pterocarpus angolensis</i> | 24 | 20.90 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 26 | 26.50 |
| | <i>Combretum molle</i> | 25 | 26.20 |
| DBH < 20 cm | <i>Vitex payos</i> | 12 | 4.05 |
| | <i>Parinari curatellifolia</i> | 5 | 4.00 |
| | <i>Combretum collinum</i> | 8 | 2.00 |
| | <i>Strychnos innocua</i> | 5 | 2.24 |

Sampling point 2:

Coordinates: S 6°80370 / E 31°66665

Slope: 8%, slope exposure: 80°

Dominant tree height: 22 m

Basal area (Bitterlich, k=1): 16 m²

Photos: 6008, 6009, 6010, 6011

Disturbances/notes (2-3): old traditional bee hive on the ground (290 steps)

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Parinari curatellifolia</i> | 23 | 6.84 |
| | <i>Brachystegia microphylla</i> | 21 | 6.00 |
| | <i>Julbernardia globiflora</i> | 24 | 9.11 |
| | <i>Parinari curatellifolia</i> | 37 | 13.11 |
| DBH < 20 cm | <i>Parinari curatellifolia</i> | 6 | 2.41 |
| | <i>Uapaca kirkiana</i> | 14 | 3.60 |
| | <i>Julbernardia globiflora</i> | 7 | 4.45 |
| | <i>Brachystegia microphylla</i> | 15 | 2.65 |

Sampling point 3:

Coordinates: S 6°80225 / E 31°66787

Slope: 3-4%, slope exposure: 340°

Dominant tree height: 20 m

Basal area (Bitterlich, k=1): 14 m²

Photos: 6012, 6013, 6014, 6015

Disturbances/notes (3-4): beekeepers camp (30 steps); 1 *Pterocarpus a.* cut (>10 yrs) (75 steps)

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Pterocarpus angolensis</i> | 38 | 27.16 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 21 | 2.60 |
| | <i>Pericopsis angolensis</i> | 38 | 5.58 |
| | <i>Brachystegia microphylla</i> | 30 | 6.51 |
| DBH < 20 cm | <i>Hexalobus monopetalus</i> | 5 | 5.50 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 13 | 2.16 |
| | <i>Hexalobus monopetalus</i> | 5 | 3.15 |
| | <i>Hexalobus monopetalus</i> | 3 | 13.88 |

Sampling point 4:

Coordinates: S 6°80085 / E 31°66928

Slope: +/- flat, slope exposure: 335°

Dominant tree height: 18 m

Basal area (Bitterlich, k=1): 10.5 m²

Photos: 6016, 6017, 6018, 6019

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Parinari curatellifolia</i> | 21 | 13.56 |
| | <i>Brachystegia utilis</i> | 37 | 5.55 |
| | <i>Brachystegia utilis</i> | 33 | 3.15 |
| | <i>Brachystegia glaucescens</i> | 21 | 16.65 |
| DBH < 20 cm | <i>Pterocarpus angolensis</i> | 6 | 7.91 |
| | <i>Parinari curatellifolia</i> | 19 | 7.76 |
| | <i>Uapaca kirkiana</i> | 9 | 8.77 |
| | <i>Uapaca kirkiana</i> | 9 | 6.96 |

Sampling point 5:

Coordinates: S 6°79929 / E 31°67060

Slope: 3-4%, slope exposure: 320°

Dominant tree height: 18 m

Basal area (Bitterlich, k=1): 9 m²

Photos: 6020, 6021, 6022, 6023

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 46 | 8.45 |
| | <i>Brachystegia utilis</i> | 36 | 16.24 |
| | <i>Parinari curatellifolia</i> | 41 | 11.73 |
| | <i>Parinari curatellifolia</i> | 62 | 8.56 |
| DBH < 20 cm | <i>Phyllocosmus leimareanus</i> | 12 | 5.08 |
| | <i>Brachystegia utilis</i> | 5 | 4.70 |
| | <i>Pterocarpus angolensis</i> | 16 | 3.55 |
| | <i>Uapaca kirkiana</i> | 5 | 2.08 |

Sampling point 6:

Coordinates: S 6°79778 / E 31°67186

Slope: 3-4%, slope exposure: 360°

Dominant tree height: 18 m

Basal area (Bitterlich, k=1): 10 m²

Photos: 6024, 6025, 6026, 6027

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Pseudolachnostylis maprouneifolia</i> | 20 | 15.85 |
| | <i>Brachystegia utilis</i> | 59 | 12.95 |
| | <i>Vitex doniana</i> | 35 | 8.00 |
| | <i>Brachystegia utilis</i> | 32 | 13.40 |
| DBH < 20 cm | <i>Bobgunnia madagascariensis</i> | 15 | 10.10 |
| | <i>Pterocarpus angolensis</i> | 3 | 3.51 |
| | <i>Diplorhynchus condylocarpon</i> | 5 | 10.15 |
| | <i>Brachystegia utilis</i> | 4 | 2.35 |

Sampling point 7:

Coordinates: S 6°79645 / E 31°67337

Slope: 3-4%, slope exposure: 360°

Dominant tree height: 18 m

Basal area (Bitterlich, k=1): 13 m²

Photos: 6028, 6029, 6030, 6031

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 33 | 9.21 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 32 | 4.70 |
| | <i>Julbernardia globiflora</i> | 25 | 8.85 |
| | <i>Julbernardia globiflora</i> | 29 | 7.70 |
| DBH < 20 cm | <i>Brachystegia glaucescens</i> | 5 | 5.30 |
| | <i>Brachystegia glaucescens</i> | 15 | 8.88 |
| | <i>Burkea africana</i> | 3 | 5.61 |
| | <i>Diplorhynchus condylocarpon</i> | 5 | 1.04 |

Sampling point 8:

Coordinates: S 6°79527 / E 31°67500

Slope: 2-3%, slope exposure: 360°

Dominant tree height: 15 m

Basal area (Bitterlich, k=1): 13 m²

Photos: 6032, 6033, 6034, 6035

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia utilis</i> | 24 | 11.57 |
| | <i>Uapaca nitida</i> | 21 | 13.80 |
| | <i>Bobgunnia madagascariensis</i> | 22 | 13.51 |
| | <i>Julbernardia globiflora</i> | 43 | 9.82 |
| DBH < 20 cm | <i>Uapaca nitida</i> | 8 | 2.03 |
| | <i>Julbernardia globiflora</i> | 3 | 4.21 |
| | <i>Bobgunnia madagascariensis</i> | 10 | 1.86 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 3 | 6.06 |

Sampling point 9:

Coordinates: S 6°79413 / E 31°67652

Slope: 1-2%, slope exposure: 340°

Dominant tree height: 15 m

Basal area (Bitterlich, k=1): 7 m²

Photos: 6036, 6037, 6038, 6039

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Monotes katangensis</i> | 22 | 13.89 |
| | <i>Brachystegia glaucescens</i> | 22 | 20.15 |
| | <i>Uapaca kirkiana</i> | 27 | 11.40 |
| | <i>Brachystegia utilis</i> | 21 | 25.75 |
| DBH < 20 cm | <i>Albizia antunesiana</i> | 11 | 1.50 |
| | <i>Pterocarpus angolensis</i> | 14 | 3.31 |
| | <i>Brachystegia glaucescens</i> | 12 | 2.38 |
| | <i>Pterocarpus angolensis</i> | 5 | 5.32 |

Sampling point 10:

Coordinates: S 6°79270 / E 31°67792

Slope: 3-4%, slope exposure: 330°

Dominant tree height: 16 m

Basal area (Bitterlich, k=1): 13.5 m²

Photos: 6040, 6041, 6042, 6043

Disturbances/notes: * 1 *Monotes katangensis* recorded cut for pole

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Albizia antunesiana</i> | 52 | 15.30 |
| | <i>Brachystegia glaucescens</i> | 27 | 6.50 |
| | <i>Pterocarpus angolensis</i> | 20 | 13.98 |
| | <i>Pterocarpus angolensis</i> | 30 | 8.97 |
| DBH < 20 cm | <i>Monotes katangensis</i> | 9 | 0.70 |
| | <i>Brachystegia glaucescens</i> | 5 | 4.21 |
| | * <i>Monotes katangensis</i> | 10 | 2.35 |
| | <i>Monotes katangensis</i> | 18 | 5.75 |

Sampling point 11:

Coordinates: S 6°79131 / E 31°67926

Slope: +/- 1%, slope exposure: 320°

Dominant tree height: 14 m

Basal area (Bitterlich, k=1): 9 m²

Photos: 6044, 6045, 6046, 6047

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia utilis</i> | 70 | 9.38 |
| | <i>Combretum molle</i> | 24 | 10.00 |
| | <i>Brachystegia utilis</i> | 41 | 4.91 |
| | <i>Brachystegia utilis</i> | 24 | 11.42 |
| DBH < 20 cm | <i>Terminalia mollis</i> | 14 | 10.99 |
| | <i>Terminalia kaiseriana</i> | 8 | 9.52 |
| | <i>Diplorhynchus condylocarpon</i> | 3 | 1.05 |
| | <i>Terminalia kaiseriana</i> | 10 | 5.21 |

Sampling point 12:

Coordinates: S 6°78970 / E 31°68052

Slope: 1-2%, slope exposure: 310°

Dominant tree height: 16 m

Basal area (Bitterlich, k=1): 6 m²

Photos: 6048, 6049, 6050, 6051

Disturbances/notes (12 – 13): 1 *Pterocarpus angolensis* cut (>10 yrs) (10 steps)

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia spiciformis</i> | 21 | 10.76 |
| | <i>Erythrina abyssinica</i> | 75 | 8.93 |
| | <i>Combretum molle</i> | 23 | 11.20 |
| | <i>Ziziphus mucronata</i> | 24 | 12.82 |
| DBH < 20 cm | <i>Brachystegia glaucescens</i> | 8 | 10.49 |
| | <i>Annona senegalensis</i> | 8 | 8.23 |
| | <i>Combretum molle</i> | 7 | 1.02 |
| | <i>Azanza garckeana</i> | 7 | 3.70 |

Sampling point 13:

Coordinates: S 6°79801 / E 31°68150

Slope: 3%, slope exposure: 280°

Dominant tree height: 17 m

Basal area (Bitterlich, k=1): 15 m²

Photos: 6052, 6053, 6054, 6055

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Erythrophleum africanum</i> | 21 | 10.59 |
| | <i>Brachystegia spiciformis</i> | 20 | 5.13 |
| | <i>Erythrophleum africanum</i> | 33 | 5.49 |
| | <i>Brachystegia spiciformis</i> | 40 | 8.65 |
| DBH < 20 cm | <i>Pseudolachnostylis maprouneifolia</i> | 5 | 3.01 |
| | <i>Brachystegia glaucescens</i> | 3 | 4.62 |
| | <i>Terminalia mollis</i> | 8 | 5.16 |
| | <i>Terminalia kaiseriana</i> | 7 | 4.81 |

Sampling point 14:

Coordinates: S 6°78626 / E 31°68255

Slope: 1-2%, slope exposure: 330°

Dominant tree height: 17 m

Basal area (Bitterlich, k=1): 15 m²

Photos: 6056, 6057, 6058, 6059

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia spiciformis</i> | 36 | 9.60 |
| | <i>Brachystegia spiciformis</i> | 75 | 16.90 |
| | <i>Brachystegia spiciformis</i> | 22 | 15.85 |
| | <i>Monanthotaxis discolor</i> | 27 | 2.47 |
| DBH < 20 cm | <i>Vitex doniana</i> | 5 | 4.40 |
| | <i>Combretum collinum</i> | 7 | 7.68 |
| | <i>Brachystegia spiciformis</i> | 6 | 2.31 |
| | <i>Brachystegia glaucescens</i> | 5 | 1.92 |

Sampling point 15:

Coordinates: S 6°78486 / E 31°68393

Slope: 0-1%, slope exposure: 290°

Dominant tree height: 18 m

Basal area (Bitterlich, k=1): 8 m²

Photos: 6060, 6061, 6062, 6063

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Vitex doniana</i> | 46 | 3.60 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 20 | 8.84 |
| | <i>Julbernardia globiflora</i> | 42 | 4.35 |
| | <i>Brachystegia spiciformis</i> | 38 | 10.65 |
| DBH < 20 cm | <i>Oldfieldia dactylophylla</i> | 10 | 3.95 |
| | <i>Oldfieldia dactylophylla</i> | 4 | 11.92 |
| | <i>Vitex doniana</i> | 4 | 10.30 |
| | <i>Oldfieldia dactylophylla</i> | 17 | 7.07 |

Sampling point 16:

Coordinates: S 6°78355 / E 31°68534

Slope: +/- 1%, slope exposure: 320°

Dominant tree height: 12 m

Basal area (Bitterlich, k=1): 7 m²

Photos: 6064, 6065, 6066, 6067

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Pterocarpus angolensis</i> | 21 | 10.93 |
| | <i>Pericopsis angolensis</i> | 21 | 7.85 |
| | <i>Kigelia africana</i> | 35 | 18.10 |
| | <i>Phyllanthus engleri</i> | 45 | 19.40 |
| DBH < 20 cm | <i>Lonchocarpus capassa</i> | 3 | 2.00 |
| | <i>Pterocarpus angolensis</i> | 8 | 1.56 |
| | <i>Pterocarpus angolensis</i> | 8 | 2.18 |
| | <i>Lonchocarpus capassa</i> | 5 | 1.12 |

Sampling point 17:

Coordinates: S 6°78227 / E 31°68687

Slope: +/- 1%, slope exposure: 320°

Dominant tree height: 18 m

Basal area (Bitterlich, k=1): 9.5 m²

Photos: 6068, 6069, 6070, 6071

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Ochna holsti</i> | 22 | 3.44 |
| | <i>Combretum collinum</i> | 21 | 11.04 |
| | <i>Erythrophleum africanum</i> | 24 | 18.59 |
| | <i>Pericopsis angolensis</i> | 33 | 15.20 |
| DBH < 20 cm | <i>Diplorhynchus condylocarpon</i> | 15 | 9.05 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 4 | 8.10 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 5 | 8.63 |
| | <i>Terminalia kaiseriana</i> | 11 | 5.49 |

Sampling point 18:

Coordinates: S 6°78082 / E 31°68813

Slope: +/- 1%, slope exposure: 300°

Dominant tree height: 18 m

Basal area (Bitterlich, k=1): 18 m²

Photos: 6072, 6073, 6074, 6075

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia spiciformis</i> | 55 | 8.80 |
| | <i>Brachystegia spiciformis</i> | 21 | 6.50 |
| | <i>Strychnos innocua</i> | 23 | 8.00 |
| | <i>Pterocarpus angolensis</i> | 23 | 3.97 |
| DBH < 20 cm | <i>Oldfieldia dactylophylla</i> | 7 | 2.08 |
| | <i>Oldfieldia dactylophylla</i> | 10 | 3.59 |
| | <i>Oldfieldia dactylophylla</i> | 13 | 3.50 |
| | <i>Combretum collinum</i> | 14 | 2.01 |

Sampling point 19:

Coordinates: S 6°77945 / E 31°68953

Slope: +/- 1%, slope exposure: 260°

Dominant tree height: 24 m

Basal area (Bitterlich, k=1): 12 m²

Photos: 6076, 6077, 6078, 6079

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia spiciformis</i> | 53 | 3.85 |
| | <i>Mystroxylon aethiopicum</i> | 21 | 9.75 |
| | <i>Strychnos innocua</i> | 26 | 11.89 |
| | <i>Combretum molle</i> | 30 | 14.48 |
| DBH < 20 cm | <i>Monanthotaxis discolor</i> | 3 | 1.72 |
| | <i>Pterocarpus angolensis</i> | 4 | 1.08 |
| | <i>Pterocarpus angolensis</i> | 13 | 2.41 |
| | <i>Pterocarpus angolensis</i> | 3 | 2.06 |

Sampling point 20:

Coordinates: S 6°77808 / E 31°69095

Slope: 1-2%, slope exposure: 280°

Dominant tree height: 22 m

Basal area (Bitterlich, k=1): 11 m²

Photos: 6080, 6081, 6082, 6083

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Erythrophleum africanum</i> | 33 | 11.43 |
| | <i>Erythrophleum africanum</i> | 37 | 14.14 |
| | <i>Brachystegia spiciformis</i> | 46 | 3.85 |
| | <i>Erythrophleum africanum</i> | 40 | 11.17 |
| DBH < 20 cm | <i>Markhamia obtusifolia</i> | 7 | 1.80 |
| | <i>Brachystegia spiciformis</i> | 3 | 2.66 |
| | <i>Albizia antunesiana</i> | 8 | 2.03 |
| | <i>Hexalobus monopetalus</i> | 8 | 7.62 |

Sampling point 21:

Coordinates: S 6°77674 / E 31°69236

Slope: 1-2%, slope exposure: 280°

Dominant tree height: 18 m

Basal area (Bitterlich, k=1): 13 m²

Photos: no photographs

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Maranthes floribunda</i> | 56 | 3.43 |
| | <i>Olax obtusifolia</i> | 25 | 12.22 |
| | <i>Brachystegia spiciformis</i> | 37 | 6.32 |
| | <i>Brachystegia spiciformis</i> | 39 | 3.91 |
| DBH < 20 cm | <i>Lannea schimperi</i> | 17 | 4.15 |
| | <i>Hexalobus monopetalus</i> | 5 | 2.69 |
| | <i>Julbernardia globiflora</i> | 12 | 2.80 |
| | <i>Albizia antunesiana</i> | 6 | 11.28 |

Sampling point 22:

Coordinates: S 6°77529 / E 31°69374

Slope: +/- flat

Dominant tree height: 12 m

Basal area (Bitterlich, k=1): 8 m²

Photos: 6085, 6086, 6087, 6088

Disturbances/notes: –

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Parinari curatellifolia</i> | 40 | 15.45 |
| | <i>Anisophyllea pomifera</i> | 41 | 11.80 |
| | <i>Phyllocosmus leimareanus</i> | 21 | 8.45 |
| | <i>Brachystegia utilis</i> | 43 | 9.90 |
| DBH < 20 cm | <i>Brachystegia spiciformis</i> | 5 | 3.58 |
| | <i>Erythrophleum africanum</i> | 3 | 5.56 |
| | <i>Erythrophleum africanum</i> | 7 | 4.40 |
| | <i>Brachystegia spiciformis</i> | 6 | 3.96 |

Sampling point 23:

Coordinates: S 6°77376 / E 31°69503

Slope: +/- flat

Dominant tree height: 18 m

Basal area (Bitterlich, k=1): 13.5 m²

Photos: 6089, 6090, 6091, 6092

Disturbances/notes: –

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 39 | 10.62 |
| | <i>Parinari curatellifolia</i> | 31 | 12.34 |
| | <i>Anisophyllea pomifera</i> | 25 | 8.57 |
| | <i>Brachystegia spiciformis</i> | 31 | 3.92 |
| DBH < 20 cm | <i>Phyllocosmus leimareanus</i> | 4 | 12.02 |
| | <i>Phyllocosmus leimareanus</i> | 16 | 7.01 |
| | <i>Hexalobus monopetalus</i> | 3 | 2.90 |
| | <i>Combretum molle</i> | 3 | 9.29 |

Sampling point 24:

Coordinates: S 6°77249 / E 31°69654

Slope: +/- flat

Dominant tree height: 15 m

Basal area (Bitterlich, k=1): 7 m²

Photos: 6093, 6094, 6095, 6096

Disturbances/notes: –

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Maranthes floribunda</i> | 34 | 10.10 |
| | <i>Erythrophleum africanum</i> | 25 | 4.40 |
| | <i>Erythrophleum africanum</i> | 22 | 7.78 |
| | <i>Brachystegia spiciformis</i> | 47 | 9.49 |
| DBH < 20 cm | <i>Diplorhynchus condylocarpon</i> | 3 | 2.63 |
| | <i>Albizia antunesiana</i> | 3 | 2.05 |
| | <i>Multidentia crassa</i> | 3 | 12.71 |
| | <i>Diplorhynchus condylocarpon</i> | 4 | 3.29 |

Sampling point 25:

Coordinates: S 6°77094 / E 31°69788

Slope: 0-1%, slope exposure: 300°

Dominant tree height: 17 m

Basal area (Bitterlich, k=1): 5.5 m²

Photos: 6097, 6098, 6099, 6100

Disturbances/notes: 1 *Pterocarpus angolensis* cut for wild honey harvesting (4 yrs) (130 steps)

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|-----------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Musa</i> | 47 | 14.35 |
| | <i>Maranthes floribunda</i> | 43 | 3.98 |
| | <i>Julbernardia globiflora</i> | 34 | 16.69 |
| | <i>Parinari curatellifolia</i> | 23 | 11.40 |
| DBH < 20 cm | <i>Chrysophyllum bangweolense</i> | 4 | 4.29 |
| | <i>Ximenia americana</i> | 10 | 10.40 |
| | <i>Memecylon flavovirens</i> | 3 | 5.53 |
| | <i>Hexalobus monopetalus</i> | 5 | 8.42 |

Sampling point 26:

Coordinates: S 6°76946 / E 31°69907

Slope: +/- 1%, slope exposure: 290°

Dominant tree height: 20 m

Basal area (Bitterlich, k=1): 9 m²

Photos: 6101, 6102, 6103, 6104

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|-----------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Anisophyllea pomifera</i> | 21 | 13.10 |
| | <i>Parinari curatellifolia</i> | 29 | 4.09 |
| | <i>Maranthes floribunda</i> | 36 | 6.12 |
| | <i>Phyllocosmus leimareanus</i> | 35 | 3.66 |
| DBH < 20 cm | <i>Bobgunnia madagascariensis</i> | 3 | 7.09 |
| | <i>Julbernardia globiflora</i> | 8 | 5.68 |
| | <i>Brachystegia glaucescens</i> | 6 | 2.07 |
| | <i>Erythrophleum africanum</i> | 3 | 9.09 |

Sampling point 27:

Coordinates: S 6°76817 / E 31°70052

Slope: +/- flat

Dominant tree height: 15 m

Basal area (Bitterlich, k=1): 14 m²

Photos: 6105, 6106, 6107, 6108

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 40 | 5.53 |
| | <i>Parinari curatellifolia</i> | 23 | 11.59 |
| | <i>Julbernardia globiflora</i> | 36 | 15.58 |
| | <i>Brachystegia spiciformis</i> | 44 | 17.30 |
| DBH < 20 cm | <i>Brachystegia spiciformis</i> | 8 | 1.32 |
| | <i>Strychnos spinosa</i> | 3 | 2.11 |
| | <i>Brachystegia spiciformis</i> | 5 | 2.42 |
| | <i>Brachystegia spiciformis</i> | 8 | 3.16 |

Sampling point 28 (28 – 50 Lucile)

Coordinates: S 6°76632 / E 31°70142
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 8-10m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4112 / 4113 / 4114 / 4115
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Anisopyllea pomifera</i> | 27 | 22.6 |
| | <i>Brachystegia manga</i> | 45 | 13.32 |
| | <i>Brachystegia spiciformis</i> | 42 | 18.10 |
| | <i>Brachystegia spiciformis</i> | 76 | 10.00 |
| DBH < 20 cm | <i>Parinari curatellifolia</i> | 12 | 4.35 |
| | <i>Julbernardia globiflora</i> | 3 | 1.78 |
| | <i>Brachystegia manga</i> | 10 | 2.23 |
| | <i>Parinari curatellifolia</i> | 4 | 3.15 |

Sampling point 29:

Coordinates: S 6°76474 / E 31°70244
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 8-10 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4116 / 4117 / 4118 / 4119
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Anisopylea pomifera</i> | 25 | 12.00 |
| | <i>Anisopylea pomifera</i> | 31 | 16.96 |
| | <i>Pericopsis angolensis</i> | 35 | 3.63 |
| | <i>Pericopsis angolensis</i> | 32 | 16.39 |
| DBH < 20 cm | <i>Julbernardia globiflora</i> | 6 | 0.59 |
| | <i>Brachystegia spiciformis</i> | 8 | 1.28 |
| | <i>Brachystegia glaucescens</i> | 6 | 6.39 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 6 | 3.75 |

Sampling point 30:

Coordinates: S 6°76314 / E 31°70400
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 10 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4120 / 4121 / 4122 / 4123
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia spiciformis</i> | 46 | 2.51 |
| | <i>Parinari curatellifolia</i> | 36 | 10.50 |
| | <i>Brachystegia spiciformis</i> | 48 | 7.05 |
| | <i>Brachystegia spiciformis</i> | 39 | 11.10 |
| DBH < 20 cm | <i>Memecylon flavovirens</i> | 4 | 2.00 |
| | <i>Parinari curatellifolia</i> | 4 | 3.24 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 13 | 5.75 |
| | <i>Garcinia huillensis</i> | 18 | 5.25 |

Sampling point 31:

Coordinates: S 6°76177 / E 31°70546
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 10 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4124 / 4125 / 4126 / 4127
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia spiciformis</i> | 60 | 19 |
| | <i>Brachystegia spiciformis</i> | 35 | 19.50 |
| | <i>Brachystegia spiciformis</i> | 48 | 11.35 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 33 | 8.85 |
| DBH < 20 cm | <i>Albizia antunesiana</i> | 13 | 6.35 |
| | <i>Pterocarpus angolensis</i> | 11 | 0.83 |
| | <i>Garcinia huillensis</i> | 4 | 3.4 |
| | <i>Monanthotaxis discolor</i> | 4 | 1.22 |

Sampling point 32:

Coordinates: S 6°76314 / E 31°70400
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 10 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4128 / 4129 / 4130 / 4131
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Phyllocosmus leimareanus</i> | 31 | 11.34 |
| | <i>Erythrophleum africanum</i> | 27 | 10.52 |
| | <i>Erythrophleum africanum</i> | 28 | 8.45 |
| | <i>Brachystegia manga</i> | 23 | 13.75 |
| DBH < 20 cm | <i>Brachystegia spiciformis</i> | 4 | 1.40 |
| | <i>Brachystegia spiciformis</i> | 8 | 1.27 |
| | <i>Erythrophleum africanum</i> | 10 | 2.20 |
| | <i>Brachystegia spiciformis</i> | 6 | 1.60 |

Sampling point 33:

Coordinates: S 6°75852 / E 31°70751
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 10 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4132 / 4133 / 4134 / 4135
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 37 | 9.84 |
| | <i>Brachystegia spiciformis</i> | 41 | 16.95 |
| | <i>Hexalobus monopetalus</i> | 39 | 14.70 |
| | <i>Parinari curatellifolia</i> | 36 | 9.42 |
| DBH < 20 cm | <i>Erythrophleum africanum</i> | 16 | 3.30 |
| | <i>Brachystegia glaucescens</i> | 8 | 1.93 |
| | <i>Erythrophleum africanum</i> | 4 | 3.25 |
| | <i>Diplorhynchus condylocarpon</i> | 11 | 2.83 |

Sampling point 34:

Coordinates: S 6°75658 / E 31°70879
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 12-14m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4136 / 4137 / 4138 / 4139
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Pseudolachnostylis maprouneifolia</i> | 25 | 3.44 |
| | <i>Albizia antunesiana</i> | 31 | 2.50 |
| | <i>Julbernardia globiflora</i> | 39 | 35.10 |
| | <i>Phyllocosmus leimareanus</i> | 24 | 5.50 |
| DBH < 20 cm | <i>Hexalobus monopetalus</i> | 7 | 3.84 |
| | <i>Combretum molle</i> | 8 | 4.10 |
| | <i>Diplorhynchus condylocarpon</i> | 8 | 8.60 |
| | <i>Bobgunnia madagascariensis</i> | 7 | 2.45 |

Sampling point 35:

Coordinates: S 6°75501 / E 31°71039
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 14 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4141 / 4142 / 4143 / 4144
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 48 | 2.10 |
| | <i>Julbernardia globiflora</i> | 48 | 1.90 |
| | <i>Julbernardia globiflora</i> | 44 | 13 |
| | <i>Brachystegia spiciformis</i> | 24 | 2.15 |
| DBH < 20 cm | <i>Pseudolachnostylis maprouneifolia</i> | 12 | 4.95 |
| | <i>Hexalobus monopetalus</i> | 5 | 2.90 |
| | <i>Lannea schimperi</i> | 14 | 13.50 |
| | <i>Diplorhynchus condylocarpon</i> | 6 | 9.00 |

Sampling point 36:

Coordinates: S 6°75334 / E 31°71138
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 11 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4147 / 4148 / 4149 / 4150
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia spiciformis</i> | 32 | 5.50 |
| | <i>Julbernardia globiflora</i> | 36 | 5.25 |
| | <i>Phyllocosmus leimareanus</i> | 25 | 11.04 |
| | <i>Julbernardia globiflora</i> | 40 | 10.80 |
| DBH < 20 cm | <i>Brachystegia glaucescens</i> | 15 | 8.00 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 12 | 4.95 |
| | <i>Pterocarpus angolensis</i> | 5 | 14.27 |
| | <i>Albizia antunesiana</i> | 4 | 8.73 |

Sampling point 37:

Coordinates: S 6°75182 / E 31°71277
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 14 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4151 / 4152 / 4153 / 4154
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 32 | 1.10 |
| | <i>Julbernardia globiflora</i> | 36 | 3.13 |
| | <i>Julbernardia globiflora</i> | 37 | 1.00 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 21 | 19.50 |
| DBH < 20 cm | <i>Pseudolachnostylis maprouneifolia</i> | 10 | 11.90 |
| | <i>Maranthes floribunda</i> | 16 | 8.20 |
| | <i>Pterocarpus angolensis</i> | 19 | 8.82 |
| | <i>Monanthotaxis discolor</i> | 4 | 6.20 |

Sampling point 38:

Coordinates: S 6°75022 / E 31°71382
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 8 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4156 / 4157 / 4158 / 4159
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 40 | 9.45 |
| | <i>Brachystegia glaucescens</i> | 25 | 6.00 |
| | <i>Anisophyllea pomifera</i> | 26 | 5.82 |
| | <i>Albizia antunesiana</i> | 40 | 10.77 |
| DBH < 20 cm | <i>Phyllocosmus leimareanus</i> | 11 | 5.37 |
| | <i>Phyllocosmus leimareanus</i> | 4 | 1.22 |
| | <i>Brachystegia glaucescens</i> | 11 | 3.35 |
| | <i>Phyllocosmus leimareanus</i> | 13 | 5.70 |

Sampling point 39:

Coordinates: S 6°74841 / E 31°71442
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 10 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4160 / 4161 / 4162 / 4163
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Monotes africanus</i> | 36 | 22.85 |
| | <i>Vitex doniana</i> | 27 | 16.75 |
| | <i>Brachystegia spiciformis</i> | 37 | 13.20 |
| | <i>Lannea schimperi</i> | 28 | 13.15 |
| DBH < 20 cm | <i>Brachystegia spiciformis</i> | 15 | 4.20 |
| | <i>Erythrophleum africanum</i> | 4 | 4.72 |
| | <i>Erythrophleum africanum</i> | 4 | 2.74 |
| | <i>Diplorhynchus condylocarpon</i> | 7 | 1.55 |

Sampling point 40:

Coordinates: S 6°74684 / E 31°71562
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 8-10 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4164 / 4165 / 4166 / 4167
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 35 | 23.94 |
| | <i>Erythrophleum africanum</i> | 23 | 17.60 |
| | <i>Julbernardia globiflora</i> | 40 | 12.85 |
| | <i>Brachystegia manga</i> | 22 | 6.25 |
| DBH < 20 cm | <i>Diplorhynchus condylocarpon</i> | 7 | 4.62 |
| | <i>Strychnos innocua</i> | 10 | 4.30 |
| | <i>Monotes africanus</i> | 14 | 9.00 |
| | <i>Strychnos innocua</i> | 6 | 1.35 |

Sampling point 41:

Coordinates: S 6°74531 / E 31°71693
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 12 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4168 / 4169 / 4170 / 4171
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 30 | 8.10 |
| | <i>Monotes africanus</i> | 38 | 13.40 |
| | <i>Erythrophleum africanum</i> | 35 | 9.40 |
| | <i>Parinari curatellifolia</i> | 53 | 29.15 |
| DBH < 20 cm | <i>Diplorhynchus condylocarpon</i> | 12 | 2.43 |
| | <i>Monotes africanus</i> | 8 | 1.16 |
| | <i>Brachystegia glaucescens</i> | 4 | 4.89 |
| | <i>Uapaca nitida</i> | 6 | 2.93 |

Sampling point 42:

Coordinates: S 6°74379 / E 31°71832
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 12 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4172 / 4173 / 4174 / 4175
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Pterocarpus angolensis</i> | 23 | 9.86 |
| | <i>Maranthes floribunda</i> | 45 | 17.10 |
| | <i>Parinari curatellifolia</i> | 27 | 6.10 |
| | <i>Julbernardia globiflora</i> | 44 | 6.65 |
| DBH < 20 cm | <i>Julbernardia globiflora</i> | 4 | 2.10 |
| | <i>Maytenus senegalensis</i> | 4 | 1.45 |
| | <i>Brachystegia glaucescens</i> | 4 | 2.80 |
| | <i>Julbernardia globiflora</i> | 4 | 3.40 |

Sampling point 43:

Coordinates: S 6°74219 / E 31°71948
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 12-14 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: ???
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 39 | 7.20 |
| | <i>Parinari curatellifolia</i> | 25 | 7.02 |
| | <i>Parinari curatellifolia</i> | 23 | 7.50 |
| | <i>Julbernardia globiflora</i> | 51 | 6.10 |
| DBH < 20 cm | <i>Monanthotaxis discolor</i> | 9 | 3.06 |
| | <i>Ochna inermis</i> | 5 | 3.23 |
| | <i>Diplorhynchus condylocarpon</i> | 12 | 4.65 |
| | <i>Hexalobus monopetalus</i> | 6 | 3.90 |

Sampling point 44:

Coordinates: S 6°74048 / 31°72050
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 10-12 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4210 / 4211 / 4212 / 4213
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Anisophyllea pomifera</i> | 28 | 15.16 |
| | <i>Julbernardia globiflora</i> | 47 | 4.72 |
| | <i>Julbernardia globiflora</i> | 31 | 10.93 |
| | <i>Parinari curatellifolia</i> | 28 | 10.80 |
| DBH < 20 cm | <i>Brachystegia spiciformis</i> | 18 | 2.64 |
| | <i>Diplorhynchus condylocarpon</i> | 6 | 2.45 |
| | <i>Hexalobus monopetalus</i> | 5 | 8.90 |
| | <i>Cassipourea mollis</i> | 6 | 6.20 |

Sampling point 45:

Coordinates: S 6°73917/ E 31°72186
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 12-13 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4214 / 4215 / 4216 / 4217
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Lannea schimperi</i> | 35 | 10.35 |
| | <i>Brachystegia spiciformis</i> | 102 | 12.50 |
| | <i>Brachystegia spiciformis</i> | 86 | 17.60 |
| | <i>Lannea schimperi</i> | 35 | 17.70 |
| DBH < 20 cm | <i>Hexalobus monopetalus</i> | 4 | 3.53 |
| | <i>Hexalobus monopetalus</i> | 7 | 6.30 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 3 | 2.34 |
| | <i>Brachystegia glaucescens</i> | 11 | 4.55 |

Sampling point 46:

Coordinates: S 6°73760 / E 31°72293
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 12-14 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4218 / 4219 / 4220 / 4221
 Disturbances/notes: 20 step (1) tree cut for honey

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 34 | 4 |
| | <i>Brachystegia spiciformis</i> | 45 | 8.50 |
| | <i>Julbernardia globiflora</i> | 33 | 14.30 |
| | <i>Anisophyllea pomifera</i> | 29 | 6.56 |
| DBH < 20 cm | <i>Rhus longipes</i> | 6 | 4.32 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 15 | 13.65 |
| | <i>Phyllocosmus leimareanus</i> | 19 | 4.40 |
| | <i>Diplorhynchus condylocarpon</i> | 8 | 1.80 |

Sampling point 47:

Coordinates: S 6°73570 / E 31°72468
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 14-16 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4222 / 4223 / 4224 / 4225
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 29 | 8.81 |
| | <i>Brachystegia spiciformis</i> | 84 | 9.35 |
| | <i>Julbernardia globiflora</i> | 37 | 9.00 |
| | <i>Pterocarpus angolensis</i> | 44 | 24.80 |
| DBH < 20 cm | <i>Albizia antunesiana</i> | 9 | 6.25 |
| | <i>Memecylon flavovirens</i> | 4 | 9.90 |
| | <i>Anisophyllea pomifera</i> | 19 | 11.25 |
| | <i>Olax obtusifolia</i> | 4 | 5.15 |

Sampling point 48:

Coordinates: S 6°73429 / 31°72614
 Slope: not recorded, slope exposure: not recorded
 Dominant tree height: 16-18 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4226 / 4227 / 4228 / 4229
 Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Maranthes floribunda</i> | 32 | 13.40 |
| | <i>Julbernardia globiflora</i> | 34 | 13.20 |
| | <i>Julbernardia globiflora</i> | 45 | 15 |
| | <i>Olax obtusifolia</i> | 36 | 5.37 |
| DBH < 20 cm | <i>Schrebera trichoclada</i> | 4 | 10.17 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 8 | 10.90 |
| | <i>Pterocarpus angolensis</i> | 6 | 7.85 |
| | <i>Erythrophleum africanum</i> | 18 | 10.85 |

Sampling point 49:

Coordinates: S 6°73270 / 31°72739

Slope: not recorded, slope exposure: not recorded

Dominant tree height: 10-12 m

Basal area (Bitterlich, k=1): not recorded

Photos: 4231 / 4232 / 4233 / 4234

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 47 | 15.30 |
| | <i>Brachystegia spiciformis</i> | 41 | 9.05 |
| | <i>Julbernardia globiflora</i> | 35 | 15 |
| | <i>Pterocarpus angolensis</i> | 24 | 16.20 |
| DBH < 20 cm | <i>Albizia antunesiana</i> | 6 | 6.45 |
| | <i>Lannea schimperi</i> | 19 | 9.30 |
| | <i>Combretum collinum</i> | 14 | 4.75 |
| | <i>Albizia antunesiana</i> | 6 | 5.50 |

Sampling point 50:

Coordinates: S 6°73103 / E 31°72838

Slope: not recorded, slope exposure: not recorded

Dominant tree height: 14 m

Basal area (Bitterlich, k=1): not recorded

Photos: 4235 / 4236 / 4237 / 4238

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Anisophyllea pomifera</i> | 29 | 7.06 |
| | <i>Phyllocosmus leimareanus</i> | 27 | 20.25 |
| | <i>Erythrophleum africanum</i> | 29 | 7.50 |
| | <i>Julbernardia globiflora</i> | 39 | 6.35 |
| DBH < 20 cm | <i>Erythrophleum africanum</i> | 4 | 1.50 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 11 | 6.90 |
| | <i>Pterocarpus angolensis</i> | 13 | 11.76 |
| | <i>Brachystegia manga</i> | 7 | 5.40 |

Transect 2: Lower land near German road (below escarpment)

Starting point main road: S 6°55010 / E 31°83249; azimuth: 227°

Sampling point 1:

Coordinates: S 6°55138 / E 31°83089

Slope: 1-2%, slope exposure: 105°

Dominant tree height: 12 m

Basal area (Bitterlich, k=1): 12 m²

Photos: 6110, 6111, 6112, 6113

Disturbances/notes: -

| Diameter class | Nearest species in the each quarter | DBH (cm) | Distance (m) |
|----------------|-------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Erythrophleum africanum</i> | 20 | 9.18 |
| | <i>Pterocarpus angolensis</i> | 25 | 9.51 |
| | <i>Pterocarpus angolensis</i> | 21 | 0.88 |
| | <i>Strychnos innocua</i> | 21 | 13.47 |
| DBH < 20 cm | <i>Combretum molle</i> | 12 | 2.06 |
| | <i>Bobgunnia madagascariensis</i> | 10 | 8.00 |
| | <i>Brachystegia utilis</i> | 18 | 2.19 |
| | <i>Terminalia kaiseriana</i> | 4 | 1.24 |

Sampling point 2:

Coordinates: S 6°55242 / E 31°82926

Slope: +/- flat

Dominant tree height: 16 m

Basal area (Bitterlich, k=1): 13 m²

Photos: 6114, 6115, 6116, 6117

Disturbances/notes: -

| Diameter class | Nearest species in the each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Combretum fragrans</i> | 25 | 2.69 |
| | <i>Combretum fragrans</i> | 29 | 6.36 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 20 | 4.56 |
| | <i>Julbernardia globiflora</i> | 35 | 9.99 |
| DBH < 20 cm | <i>Pseudolachnostylis maprouneifolia</i> | 10 | 6.62 |
| | <i>Lonchocarpus capassa</i> | 5 | 12.36 |
| | <i>Diplorhynchus condylocarpon</i> | 5 | 9.21 |
| | <i>Hymenocardia acida</i> | 4 | 5.95 |

Sampling point 3:

Coordinates: S 6°55346 / E 31°82764

Slope: - 1%; slope exposure: 90°

Dominant tree height: 16 m

Basal area (Bitterlich, k=1): 7 m²

Photos: 6118, 6119, 6120, 6121

Disturbances/notes: -

| Diameter class | Nearest species in the each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Isoberlinia angolensis</i> | 20 | 3.63 |
| | <i>Julbernardia globiflora</i> | 27 | 7.92 |
| | <i>Julbernardia globiflora</i> | 34 | 19.25 |
| | <i>Isoberlinia angolensis</i> | 22 | 8.70 |
| DBH < 20 cm | <i>Pseudolachnostylis maprouneifolia</i> | 14 | 6.00 |
| | <i>Brachystegia utilis</i> | 13 | 6.87 |
| | <i>Hymenocardia acida</i> | 4 | 4.90 |
| | <i>Diplorhynchus condylocarpon</i> | 7 | 4.99 |

Sampling point 4:

Coordinates: S 6°55468 / E 31°82611

Slope: - 1%; slope exposure: 105°

Dominant tree height: 15 m

Basal area (Bitterlich, k=1): 11.5 m²

Photos: 6122, 6123, 6124, 6125

Disturbances/notes: -

| Diameter class | Nearest species in the each quarter | DBH (cm) | Distance (m) |
|----------------|-------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Isoberlinia angolensis</i> | 31 | 15.45 |
| | <i>Erythrophleum africanum</i> | 20 | 8.11 |
| | <i>Julbernardia globiflora</i> | 32 | 3.23 |
| | <i>Brachystegia spiciformis</i> | 25 | 17.57 |
| DBH < 20 cm | <i>Memecylon flavovirens</i> | 5 | 2.06 |
| | <i>Rothmannia engleriana</i> | 3 | 3.64 |
| | <i>Bobgunnia madagascariensis</i> | 7 | 2.93 |
| | <i>Bobgunnia madagascariensis</i> | 4 | 1.30 |

Sampling point 5:

Coordinates: S 6°55580 / E 31°82459

Slope: +/- flat

Dominant tree height: 16 m

Basal area (Bitterlich, k=1): 11 m²

Photos: 6126, 6127, 6128, 6129

Disturbances/notes: -

| Diameter class | Nearest species in the each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia utilis</i> | 35 | 12.89 |
| | <i>Julbernardia globiflora</i> | 31 | 6.88 |
| | <i>Brachystegia utilis</i> | 36 | 6.35 |
| | <i>Brachystegia utilis</i> | 25 | 13.60 |
| DBH < 20 cm | <i>Burkea africana</i> | 16 | 1.02 |
| | <i>Burkea africana</i> | 15 | 3.21 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 11 | 5.96 |
| | <i>Erythrophleum africanum</i> | 7 | 5.06 |

Sampling point 6:

Coordinates: S 6°55694 / E 31°82294

Slope: - 1%; slope exposure: 90°

Dominant tree height: 16 m

Basal area (Bitterlich, k=1): 6 m²

Photos: 6130, 6131, 6132, 6133

Disturbances/notes (6 – 7):

3 *Pseudolachnostylis* m. cut for poles (Ø 12-15 cm; >5 yrs) (30 – 50 steps)1 *Julbernardia globiflora* tree barking for bee hives (Ø 22 cm; >5 yrs) (70 steps)5 *Pterocarpus angolensis* cut for timber (>5 yrs) (90 steps)1 *Julbernardia globiflora* tree barking for bee hives (Ø 30 cm; >5 yrs) (145 steps)

| Diameter class | Nearest species in the each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 37 | 8.40 |
| | <i>Pericopsis angolensis</i> | 41 | 19.88 |
| | <i>Pterocarpus angolensis</i> | 35 | 13.50 |
| | <i>Isoberlinia angolensis</i> | 35 | 14.95 |
| DBH < 20 cm | <i>Crossopteryx febrifuga</i> | 3 | 5.26 |
| | <i>Brachystegia utilis</i> | 3 | 10.93 |
| | <i>Bobgunnia madagascariensis</i> | 5 | 6.95 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 13 | 12.70 |

Sampling point 7:

Coordinates: S 6°55796 / E 31°82143

Slope: +/- flat

Dominant tree height: 18 m

Basal area (Bitterlich, k=1): 8.5 m²

Photos: 6134, 6135, 6136, 6137

Disturbances/notes (7 – 8):

1 *Brachystegia utilis* cut wild honey harvesting (50 steps)

1 *Pterocarpus angolensis* cut for timber (Ø 35 cm; >5 yrs) (190 steps)

1 *Julbernardia globiflora* cut for timber? (Ø 25 cm; >5 yrs) (190 steps)

| Diameter class | Nearest species in the each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Pericopsis angolensis</i> | 21 | 9.28 |
| | <i>Brachystegia utilis</i> | 30 | 4.04 |
| | <i>Brachystegia utilis</i> | 24 | 12.63 |
| | <i>Brachystegia utilis</i> | 31 | 6.11 |
| DBH < 20 cm | <i>Diplorhynchus condylocarpon</i> | 19 | 6.40 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 9 | 7.30 |
| | <i>Julbernardia globiflora</i> | 3 | 8.53 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 13 | 7.30 |

Sampling point 8:

Coordinates: S 6°55909 / E 31°81994

Slope: +/- flat

Dominant tree height: 17 m

Basal area (Bitterlich, k=1): 16 m²

Photos: 6138, 6139, 6140, 6141

Disturbances/notes: -

| Diameter class | Nearest species in the each quarter | DBH (cm) | Distance (m) |
|----------------|-------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 25 | 3.20 |
| | <i>Diplorhynchus condylocarpon</i> | 24 | 7.47 |
| | <i>Brachystegia utilis</i> | 25 | 8.06 |
| | <i>Borassus aethiopum</i> | 47 | 9.85 |
| DBH < 20 cm | <i>Uapaca nitida</i> | 8 | 6.05 |
| | <i>Brachystegia utilis</i> | 6 | 3.60 |
| | <i>Bobgunnia madagascariensis</i> | 11 | 2.93 |
| | <i>Uapaca nitida</i> | 8 | 3.80 |

Sampling point 9 (9 – 50 Lucile)

Coordinates: S 6°56054 / E 31°81868

Slope: +/- flat

Dominant tree height: 9 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3844 / 3845 / 3846 / 3847

Disturbances/notes: + 280 steps: (1) tree cut for timber (>10 years)

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|-----------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 23 | 13.50 |
| | <i>Brachystegia manga</i> | 35 | 11 |
| | <i>Brachystegia manga</i> | 36 | 10.70 |
| | <i>Julbernardia globiflora</i> | 35 | 19.50 |
| DBH < 20 cm | <i>Bobgunnia madagascariensis</i> | 10 | 4.20 |
| | <i>Julbernardia globiflora</i> | 12 | 2.35 |
| | <i>Brachystegia spiciformis</i> | 9 | 3 |
| | <i>Maprounea africana</i> | 9 | 3 |

Sampling point 10

Coordinates: S 6°56202 / E 31°81752

Slope: +/- flat

Dominant tree height: 6-7 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3848 / 3849 / 3850 / 3851

Disturbances/notes: + 20 steps (1) tree cut for timber (>10 years)

+ 80 steps (1) cutting down tree for honey (> 5 years)

+290 steps (1) cutting down tree for honey (> 5 years)

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia spiciformis</i> | 28 | 8.30 |
| | <i>Pterocarpus angolensis</i> | 25 | 7.00 |
| | <i>Brachystegia spiciformis</i> | 29 | 9.30 |
| | <i>Afzelia quanzensis</i> | 27 | 4.70 |
| DBH < 20 cm | <i>Strychnos pungens</i> | 14 | 7.70 |
| | <i>Bobgunnia madagascariensis</i> | 11 | 8.40 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 14 | 6.55 |
| | <i>Hymenocardia acida</i> | 5 | 4.50 |

Sampling point 11

Coordinates: S 6°56198 / E 31°81753

Slope: +/- flat

Dominant tree height: 8 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3855 / 3856 / 3857 / 3858

Disturbances/notes: + 280 steps (1) debarking tree for beehives (>3 years) / +10 steps (1) timbering (>10 years) / +64 steps (1) timbering (>10 years) /+130 steps (1) timbering (>10 years) /+175 steps (1) timbering (>10 years) / +240 steps (1) timbering (>10 years)

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 28 | 7 |
| | <i>Strychnos pungens</i> | 22 | 3.90 |
| | <i>Brachystegia manga</i> | 24 | 3.50 |
| | <i>Julbernardia globiflora</i> | 27 | 8.30 |
| DBH < 20 cm | <i>Ziziphus</i> sp. | 9 | 15.70 |
| | <i>Erythrophleum africanum</i> | 6 | 4 |
| | <i>Diplorhynchus condylocarpon</i> | 12 | 7.60 |
| | <i>Brachystegia manga</i> | 18 | 4.70 |

Sampling point 12

Coordinates: S 6°56526 / E 31°81499

Slope: +/- flat

Dominant tree height: 10 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3859 / 3860 / 3861 / 3862

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 21 | 11.50 |
| | <i>Julbernardia globiflora</i> | 26 | 8.10 |
| | <i>Julbernardia globiflora</i> | 32 | 8.80 |
| | <i>Brachystegia spiciformis</i> | 34 | 10 |
| DBH < 20 cm | <i>Pterocarpus angolensis</i> | 17 | 8.20 |
| | <i>Erythrophleum africanum</i> | 14 | 5.50 |
| | <i>Erythrophleum africanum</i> | 12 | 5.80 |
| | <i>Diplorhynchus condylocarpon</i> | 11 | 1.40 |

Sampling point 13

Coordinates: S 6°56687 / E 31°81383

Slope: +/- flat

Dominant tree height: 6-8 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3868 / 3869 / 3870 / 3871

Disturbances/notes: -

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 38 | 9.60 |
| | <i>Lannea schimperi</i> | 29 | 17.60 |
| | <i>Julbernardia globiflora</i> | 35 | 16.10 |
| | <i>Julbernardia globiflora</i> | 38 | 10.90 |
| DBH < 20 cm | <i>Pseudolachnostylis maprouneifolia</i> | 7 | 3.80 |
| | <i>Euphorbia matabensis</i> | 12 | 0.70 |
| | <i>Flacourtie indica</i> | 5 | 2.90 |
| | <i>Pterocarpus angolensis</i> | 7 | 3.20 |

Sampling point 14

Coordinates: S 6°56875 / E 31°81290

Slope: +/- flat

Dominant tree height: 8-10m

Basal area (Bitterlich, k=1): not recorded

Photos: 3891 / 3892 / 3893 / 3894

Disturbances/notes: +28 steps (1) timbering (> 5 years)

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 27 | 6.80 |
| | <i>Brachystegia manga</i> | 36 | 9.30 |
| | <i>Brachystegia manga</i> | 51 | 6.45 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 23 | 4.50 |
| DBH < 20 cm | <i>Brachystegia glaucescens</i> | 12 | 2.90 |
| | <i>Diplorhynchus condylocarpon</i> | 16 | 3.45 |
| | <i>Monanthotaxis discolor</i> | 9 | 10.95 |
| | <i>Hymenocardia acida</i> | 6 | 15.10 |

Sampling point 15

Coordinates: S 6°57085/ E 31°81202

Slope: +/- flat

Dominant tree height: 10-12m

Basal area (Bitterlich, k=1): not recorded

Photos: 3895 / 3896 / 3897 / 3898

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Lannea schimperi</i> | 25 | 7.80 |
| | <i>Brachystegia manga</i> | 20 | 10.60 |
| | <i>Brachystegia manga</i> | 27 | 9.60 |
| | <i>Isoberlinia angolensis</i> | 32 | 13.53 |
| DBH < 20 cm | <i>Isoberlinia angolensis</i> | 6 | 2.65 |
| | <i>Hymenocardia acida</i> | 5 | 3.07 |
| | <i>Brachystegia spiciformis</i> | 14 | 8.65 |
| | <i>Oldfieldia dactylophylla</i> | 3 | 5.15 |

Sampling point 16

Coordinates: S 6°57246 / E 31°81025

Slope: +/- flat

Dominant tree height: 6-8m

Basal area (Bitterlich, k=1): not recorded

Photos: 3903 / 3904 / 3905 / 3906

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Pericopsis angolensis</i> | 37 | 7.35 |
| | <i>Pterocarpus angolensis</i> | 26 | 2.35 |
| | <i>Combretum molle</i> | 26 | 15.37 |
| | <i>Brachystegia glaucescens</i> | 25 | 8.45 |
| DBH < 20 cm | <i>Oldfieldia dactylophylla</i> | 7 | 3.45 |
| | <i>Oldfieldia dactylophylla</i> | 17 | 7.12 |
| | <i>Pterocarpus angolensis</i> | 15 | 2.65 |
| | <i>Pterocarpus angolensis</i> | 8 | 8.80 |

Sampling point 17

Coordinates: S 6°57419/ E 31°80906

Slope: +/- flat

Dominant tree height: 6-7 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3907 / 3908 / 3909 / 3910

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 22 | 10.50 |
| | <i>Combretum fragrans</i> | 26 | 11.40 |
| | <i>Pterocarpus tinctorius</i> | 25 | 14.40 |
| | <i>Dalbergia nitidula</i> | 37 | 3.90 |
| DBH < 20 cm | <i>Brachystegia glaucescens</i> | 10 | 8.55 |
| | <i>Bauhinia thonningii</i> | 7 | 7.30 |
| | <i>Annona senegalensis</i> | 6 | 12.60 |
| | <i>Bauhinia thonningii</i> | 7 | 5.30 |

Sampling point 18

Coordinates: S 6°57592 / E 31°80795

Slope: +/- flat

Dominant tree height: 8 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3911 / 3912 / 3913 / 3914

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Combretum fragrans</i> | 25 | 10.40 |
| | <i>Diplorhynchus condylocarpon</i> | 23 | 8.15 |
| | <i>Brachystegia glaucescens</i> | 56 | 17.40 |
| | <i>Dalbergia nitidula</i> | 32 | 8.05 |
| DBH < 20 cm | <i>Combretum fragrans</i> | 9 | 6.60 |
| | <i>Diplorhynchus condylocarpon</i> | 9 | 7.95 |
| | <i>Lannea schimperi</i> | 8 | 5.90 |
| | <i>Combretum molle</i> | 7 | 6.25 |

Sampling point 19

Coordinates: S 6°57777 / E 31°80689

Slope: +/- flat

Dominant tree height: 6-8 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3915 / 3916 / 3917 / 3918

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Dalbergia nitidula</i> | 36 | 25.56 |
| | <i>Combretum molle</i> | 26 | 8.35 |
| | <i>Brachystegia glaucescens</i> | 42 | 14.15 |
| | <i>Brachystegia glaucescens</i> | 35 | 25.40 |
| DBH < 20 cm | <i>Diplorhynchus condylocarpon</i> | 10 | 5 |
| | <i>Combretum molle</i> | 13 | 8.07 |
| | <i>Terminalia kaiseriana</i> | 11 | 3.20 |
| | <i>Pterocarpus tinctorius</i> | 13 | 3.30 |

Sampling point 20

Coordinates: S 6°57956 / E 31°80575

Slope: +/- flat

Dominant tree height: 6-8m

Basal area (Bitterlich, k=1): not recorded

Photos: ??? / 3920 / 3921 / 3922

Disturbances/notes: +280 steps (2) tree cut for honey

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Pterocarpus tinctorius</i> | 24 | 4.40 |
| | <i>Brachystegia glaucescens</i> | 43 | 9.20 |
| | <i>Brachystegia glaucescens</i> | 25 | 27.50 |
| | <i>Brachystegia glaucescens</i> | 51 | 25.20 |
| DBH < 20 cm | <i>Pterocarpus tinctorius</i> | 11 | 6.50 |
| | <i>Acacia gerrardii</i> | 5 | 8.23 |
| | <i>Brachystegia glaucescens</i> | 10 | 18.70 |
| | <i>Combretum fragrans</i> | 6 | 14.10 |

Sampling point 21

Coordinates: S 6°58118 / E 31°80454

Slope: +/- flat

Dominant tree height: 6-7m

Basal area (Bitterlich, k=1): not recorded

Photos: 3924 / 3925 / 3926 / 3927

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia glaucescens</i> | 43 | 10.40 |
| | <i>Sterculia quinqueloba</i> | 24 | 16.10 |
| | <i>Combretum fragrans</i> | 32 | 31.00 |
| | <i>Combretum molle</i> | 25 | 6.10 |
| DBH < 20 cm | <i>Combretum fragrans</i> | 8 | 1.40 |
| | <i>Sterculia quinqueloba</i> | 4 | 5.12 |
| | <i>Combretum fragrans</i> | 12 | 7.30 |
| | <i>Diplorhynchus condylocarpon</i> | 13 | 6.00 |

Sampling point 22

Coordinates: S 6°58294 / E 31°80357

Slope: +/- flat

Dominant tree height: 10-12 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3928 / 3929 / 3930 / 3931

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia glaucescens</i> | 35 | 12.45 |
| | <i>Brachystegia glaucescens</i> | 31 | 10.75 |
| | <i>Brachystegia glaucescens</i> | 56 | 30.80 |
| | <i>Brachystegia glaucescens</i> | 37 | 9.10 |
| DBH < 20 cm | <i>Acacia gerrardii</i> | 6 | 3.74 |
| | <i>Combretum fragrans</i> | 8 | 12.15 |
| | <i>Lannea schimperi</i> | 6 | 16 |
| | <i>Diplorhynchus condylocarpon</i> | 6 | 12.08 |

Sampling point 23

Coordinates: S 6°58455 / E 31°80247

Slope: +/- flat

Dominant tree height: 6-8m

Basal area (Bitterlich, k=1): not recorded

Photos: 3932 / 3933 / 3934 / 3935

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia glaucescens</i> | 31 | 9.70 |
| | <i>Brachystegia glaucescens</i> | 77 | 48.65 |
| | <i>Brachystegia glaucescens</i> | 62 | 16.70 |
| | <i>Pterocarpus tinctorius</i> | 43 | 6.80 |
| DBH < 20 cm | <i>Brachystegia glaucescens</i> | 13 | 9.50 |
| | <i>Annona senegalensis</i> | 7 | 5.32 |
| | <i>Pterocarpus angolensis</i> | 17 | 11.40 |
| | <i>Dichrostachys cinerea</i> | 4 | 11.95 |

Sampling point 24

Coordinates: S 6°58594 / E 31°80065

Slope: +/- flat

Dominant tree height: 5-6 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3938 / 3939 / 3940 / 3941

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Dalbergia nitidula</i> | 42 | 20 |
| | <i>Euphorbia matabensis</i> | 27 | 17 |
| | <i>Terminalia mollis</i> | 26 | 46.45 |
| | <i>Ozoroa insignis subsp. reticulata</i> | 53 | 50.40 |
| DBH < 20 cm | <i>Bauhinia thonningii</i> | 4 | 2.88 |
| | <i>Acacia gerrardii</i> | 5 | 1.15 |
| | <i>Acacia gerrardii</i> | 5 | 6.15 |
| | <i>Combretum fragrans</i> | 8 | 3.60 |

Sampling point 25

Coordinates: S 6°58765 / E 31°79945

Slope: +/- flat

Dominant tree height: 4-6m

Basal area (Bitterlich, k=1): not recorded

Photos: 3946 / 3947 / 3948 / 3949

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Pericopsis angolensis</i> | 26 | 48.6 |
| | <i>Combretum molle</i> | 33 | 42.10 |
| | <i>Combretum fragrans</i> | 23 | 25.10 |
| | <i>Combretum fragrans</i> | 22 | 57 |
| DBH < 20 cm | <i>Combretum fragrans</i> | 4 | 7.26 |
| | <i>Combretum fragrans</i> | 4 | 13.35 |
| | <i>Bauhinia thonningii</i> | 11 | 4.35 |
| | <i>Acacia gerrardii</i> | 8 | 6.00 |

Sampling point 26

Coordinates: S 6°58816 / E 31°79709

Slope: +/- flat

Dominant tree height: 4-6 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3950 / 3951 / 3952 / 3953

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia glaucescens</i> | 23 | 14.80 |
| | <i>Combretum collinum</i> | 33 | 7.60 |
| | <i>Oldfieldia dactylophylla</i> | 32 | 17.70 |
| | <i>Erythrophleum africanum</i> | 36 | 7.55 |
| DBH < 20 cm | <i>Acacia gerrardii</i> | 4 | 9.90 |
| | <i>Combretum collinum</i> | 5 | 7.72 |
| | <i>Brachystegia glaucescens</i> | 18 | 1.15 |
| | <i>Diplorhynchus condylocarpon</i> | 16 | 3.55 |

Sampling point 27

Coordinates: S 6°58901 / E 31°79709

Slope: +/- flat

Dominant tree height: 4-6 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3954 / 3955 / 3956 / 3957

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Combretum molle</i> | 27 | 6.70 |
| | <i>Brachystegia glaucescens</i> | 66 | 35.90 |
| | <i>Pericopsis angolensis</i> | 30 | 14.60 |
| | <i>Combretum molle</i> | 25 | 26.86 |
| DBH < 20 cm | <i>Combretum fragrans</i> | 11 | 2.90 |
| | <i>Terminalia kaiseriana</i> | 9 | 5.40 |
| | <i>Pterocarpus tinctorius</i> | 18 | 6.35 |
| | <i>Combretum fragrans</i> | 8 | 1.64 |

Sampling point 28

Coordinates: S 6°59042 / E 31°79450

Slope: +/- flat

Dominant tree height: 8-10 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3963 / 3964 / 3965 / 3966

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia glaucescens</i> | 44 | 3.50 |
| | <i>Brachystegia glaucescens</i> | 46 | 4.07 |
| | <i>Xeroderris stuhlmannii</i> | 34 | 44.66 |
| | <i>Pericopsis angolensis</i> | 39 | 21.60 |
| DBH < 20 cm | <i>Combretum molle</i> | 6 | 1.53 |
| | <i>Pterocarpus tinctorius</i> | 8 | 2.67 |
| | <i>Diplorhynchus condylocarpon</i> | 7 | 1.90 |
| | <i>Combretum molle</i> | 7 | 1.75 |

Sampling point 29

Coordinates: S 6°56054 / E 31°81868

Slope: +/- flat

Dominant tree height: 9 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3967 / 3968 / 3969 / 3970

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Diospyros cornii</i> | 31 | 39.36 |
| | <i>Diospyros cornii</i> | 22 | 17.05 |
| | <i>Diospyros cornii</i> | 31 | 12.13 |
| | <i>Diospyros cornii</i> | 31 | 25.05 |
| DBH < 20 cm | <i>Combretum fragrans</i> | 5 | 12.95 |
| | <i>Acacia gerrardii</i> | 5 | 7.55 |
| | <i>Acacia gerrardii</i> | 6 | 6.44 |
| | <i>Acacia gerrardii</i> | 4 | 10.10 |

Sampling point 30

Coordinates: S 6°59351 / E 31°79137

Slope: +/- flat

Dominant tree height: 9 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3971 / 3972 / 3973 / 3974

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia glaucescens</i> | 31 | 27.90 |
| | <i>Brachystegia glaucescens</i> | 50 | 28.10 |
| | <i>Combretum molle</i> | 24 | 7.84 |
| | <i>Brachystegia glaucescens</i> | 30 | 10.15 |
| DBH < 20 cm | <i>Pterocarpus angolensis</i> | 7 | 7.88 |
| | <i>Combretum molle</i> | 18 | 3.35 |
| | <i>Acacia polyacantha</i> | 6 | 5.55 |
| | <i>Pterocarpus tinctorius</i> | 5 | 11.10 |

Sampling point 31

Coordinates: S 6°59473 / E 31°78987
 Slope: +/- flat
 Dominant tree height: 10-12 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 3975 / 3976 / 3977 / 3978
 Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia glaucescens</i> | 45 | 12.82 |
| | <i>Acacia stuhlmannii</i> | 24 | 1190 |
| | <i>Pterocarpus angolensis</i> | 43 | 3.34 |
| | <i>Lannea schimperi</i> | 28 | 19.80 |
| DBH < 20 cm | <i>Eriosema</i> | 7 | 10.90 |
| | <i>Sterculia africana</i> | 6 | 8.45 |
| | <i>Diplorhynchus condylocarpon</i> | 10 | 1.42 |
| | <i>Diplorhynchus condylocarpon</i> | 17 | 5.55 |

Sampling point 32

Coordinates: S 6°59624 / E 31°78880
 Slope: +/- flat
 Dominant tree height: 9 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 3979 / 3980 / 3981 / 3982
 Disturbances/notes: + 10 steps (1) tree cut for honey

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia glaucescens</i> | 44 | 6.85 |
| | <i>Pterocarpus angolensis</i> | 37 | 12.00 |
| | <i>Brachystegia glaucescens</i> | 39 | 4.82 |
| | <i>Erythrophleum africanum</i> | 24 | 15.15 |
| DBH < 20 cm | <i>Pterocarpus angolensis</i> | 4 | 1.60 |
| | <i>Oldfieldia dactylophylla</i> | 7 | 5 |
| | <i>Terminalia kaiseriana</i> | 10 | 2.45 |
| | <i>Combretum collinum</i> | 6 | 7.55 |

Sampling point 33

Coordinates: S 6°59788 / E 31°78773
 Slope: +/- flat
 Dominant tree height: 9 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 3983 / 3984 / 3985 / 3986
 Disturbances/notes: +170 steps (1) tree cut for honey

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Pterocarpus angolensis</i> | 32 | 6.15 |
| | <i>Brachystegia glaucescens</i> | 42 | 14.80 |
| | <i>Pterocarpus angolensis</i> | 26 | 18.20 |
| | <i>Brachystegia glaucescens</i> | 40 | 9.70 |
| DBH < 20 cm | <i>Pseudolachnostylis maprouneifolia</i> | 8 | 11.60 |
| | <i>Combretum collinum</i> | 17 | 3.90 |
| | <i>Pterocarpus angolensis</i> | 5 | 16.08 |
| | <i>Combretum collinum</i> | 8 | 4.00 |

Sampling point 34

Coordinates: S 6°59917 / E 31°78625

Slope: +/- flat

Dominant tree height: m

Basal area (Bitterlich, k=1): not recorded

Photos: 3988 / 3989 / 3990 / 3991

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|------------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Strychnos potatorum</i> | 25 | 6.70 |
| | <i>Combretum fragrans</i> | 22 | 7.85 |
| | <i>Lannea schimperi</i> | 32 | 18.70 |
| | <i>Monotes africanus</i> | 35 | 8.15 |
| DBH < 20 cm | <i>Diplorhynchus condylocarpon</i> | 7 | 2.95 |
| | <i>Brachystegia glaucescens</i> | 6 | 3.12 |
| | <i>Brachystegia glaucescens</i> | 5 | 3.04 |
| | <i>Julbernardia globiflora</i> | 7 | 1.75 |

Sampling point 35

Coordinates: S 6°60099 / E 31°78449

Slope: +/- flat

Dominant tree height: 10-12 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3992 / 3993 / 3994 / 3995

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Terminalia kaiseriana</i> | 23 | 19.15 |
| | <i>Lannea schimperi</i> | 27 | 4.75 |
| | <i>Brachystegia manga</i> | 46 | 6.00 |
| | <i>Pterocarpus angolensis</i> | 35 | 18.35 |
| DBH < 20 cm | <i>Pterocarpus angolensis</i> | 4 | 4.50 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 4 | 4.10 |
| | <i>Flacourtie indica</i> | 6 | 5.70 |
| | <i>Oldfieldia dactylophylla</i> | 4 | 3.50 |

Sampling point 36

Coordinates: S 6°60270 / E 31°78331

Slope: +/- flat

Dominant tree height: 6 m

Basal area (Bitterlich, k=1): not recorded

Photos: 3996 / 3997 / 3998 / 3999

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Burkea africana</i> | 28 | 23.80 |
| | <i>Cassia singueana</i> | 27 | 14.90 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 27 | 11.00 |
| | <i>Lannea schimperi</i> | 26 | 14.95 |
| DBH < 20 cm | <i>Brachystegia manga</i> | 7 | 3.30 |
| | <i>Pterocarpus angolensis</i> | 4 | 7.75 |
| | <i>Combretum fragrans</i> | 4 | 5.68 |
| | <i>Erythrophleum africanum</i> | 4 | 8.00 |

Sampling point 37

Coordinates: S 6°60428 / E 31°78217

Slope: +/- flat

Dominant tree height: 6-8m

Basal area (Bitterlich, k=1): not recorded

Photos: 4000 / 4001 / 4002 / 4003

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 35 | 5.65 |
| | <i>Brachystegia manga</i> | 24 | 4.05 |
| | <i>Brachystegia manga</i> | 29 | 37.50 |
| | <i>Brachystegia manga</i> | 29 | 20.10 |
| DBH < 20 cm | <i>Oldfieldia dactylophylla</i> | 6 | 8.20 |
| | <i>Dalbergia nitidula</i> | 5 | 6.20 |
| | <i>Oldfieldia dactylophylla</i> | 4 | 4.15 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 5 | 4.40 |

Sampling point 38

Coordinates: S 6°60569 / E 31°78066

Slope: +/- flat

Dominant tree height: 8-10 m

Basal area (Bitterlich, k=1): not recorded

Photos: 4004 / 4005 / 4006 / 4007

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 31 | 3.72 |
| | <i>Terminalia kaiseriana</i> | 27 | 13.30 |
| | <i>Pterocarpus angolensis</i> | 39 | 11.33 |
| | <i>Brachystegia glaucescens</i> | 57 | 20.62 |
| DBH < 20 cm | <i>Phyllocosmus leimareanus</i> | 9 | 4.00 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 5 | 1.65 |
| | <i>Oldfieldia dactylophylla</i> | 17 | 7.00 |
| | <i>Terminalia kaiseriana</i> | 6 | 1.45 |

Sampling point 39

Coordinates: S 6°60706 / E 31°77926

Slope: +/- flat

Dominant tree height: 8 m

Basal area (Bitterlich, k=1): not recorded

Photos: 4009 / 4010 / 4011 / 4012

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 25 | 8.06 |
| | <i>Brachystegia manga</i> | 24 | 10.01 |
| | <i>Brachystegia manga</i> | 23 | 6.05 |
| | <i>Pterocarpus angolensis</i> | 30 | 9.10 |
| DBH < 20 cm | <i>Hymenocardia acida</i> | 6 | 13.35 |
| | <i>Oldfieldia dactylophylla</i> | 10 | 2.08 |
| | <i>Brachystegia manga</i> | 19 | 3.05 |
| | <i>Monotes africanus</i> | 17 | 7.55 |

Sampling point 40

Coordinates: S 6°60842 / E 31°77768
 Slope: +/- flat
 Dominant tree height: 8-10 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4013 / 4014 / 4015 / 4016
 Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Pericopsis angolensis</i> | 28 | 15.83 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 58 | 16.75 |
| | <i>Burkea africana</i> | 37 | 5.02 |
| | <i>Julbernardia globiflora</i> | 36 | 10.72 |
| DBH < 20 cm | <i>Hymenocardia acida</i> | 11 | 4.55 |
| | <i>Pterocarpus angolensis</i> | 13 | 1.45 |
| | <i>Bobgunnia madagascariensis</i> | 13 | 10.70 |
| | <i>Oldfieldia dactylophylla</i> | 11 | 1.28 |

Sampling point 41

Coordinates: S 6°60977/ E 31°77606
 Slope: +/- flat
 Dominant tree height: 6-8 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4018 / 4019 / 4020 / 4021
 Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 28 | 12.78 |
| | <i>Brachystegia manga</i> | 50 | 22.65 |
| | <i>Brachystegia manga</i> | 27 | 19.15 |
| | <i>Terminalia kaiseriana</i> | 29 | 5.86 |
| DBH < 20 cm | <i>Phyllocosmus leimareanus</i> | 19 | 0.86 |
| | <i>Brachystegia manga</i> | 3 | 0.90 |
| | <i>Hymenocardia acida</i> | 4 | 5.36 |
| | <i>Brachystegia manga</i> | 9 | 2.87 |

Sampling point 42

Coordinates: S 6°60888 / E 31°77792
 Slope: +/- flat
 Dominant tree height: 6-8 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4022 / 4023 / 4024 / 4025
 Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 70 | 12.87 |
| | <i>Diplorhynchus condylocarpon</i> | 28 | 15.34 |
| | <i>Brachystegia manga</i> | 38 | 10.85 |
| | <i>Brachystegia manga</i> | 49 | 3.75 |
| DBH < 20 cm | <i>Oldfieldia dactylophylla</i> | 18 | 8.40 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 14 | 4.28 |
| | <i>Monotes africanus</i> | 6 | 4.16 |
| | <i>Vitex mombassae</i> | 13 | 3.40 |

Sampling point 43

Coordinates: S 6°61250 / E 31°77320

Slope: +/- flat

Dominant tree height: 10-12 m

Basal area (Bitterlich, k=1): not recorded

Photos: 4026 / 4027 / 4028 / 4029

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 34 | 16.80 |
| | <i>Brachystegia manga</i> | 35 | 10.65 |
| | <i>Lannea schimperi</i> | 28 | 28.00 |
| | <i>Pterocarpus tinctorius</i> | 57 | 13.10 |
| DBH < 20 cm | <i>Pterocarpus angolensis</i> | 6 | 4.60 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 19 | 3.88 |
| | <i>Julbernardia globiflora</i> | 5 | 3.66 |
| | <i>Brachystegia manga</i> | 4 | 1.90 |

Sampling point 44

Coordinates: S 6°61356 / E 31°77159

Slope: +/- flat

Dominant tree height: 6-8 m

Basal area (Bitterlich, k=1): not recorded

Photos: 4031 / 4032 / 4033 / 4034

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 30 | 20.47 |
| | <i>Brachystegia manga</i> | 31 | 36.30 |
| | <i>Brachystegia glaucescens</i> | 48 | 14.38 |
| | <i>Combretum molle</i> | 25 | 42.30 |
| DBH < 20 cm | <i>Julbernardia globiflora</i> | 10 | 2.86 |
| | <i>Flacourtie indica</i> | 4 | 2.00 |
| | <i>Julbernardia globiflora</i> | 4 | 2.90 |
| | <i>Julbernardia globiflora</i> | 4 | 4.68 |

Sampling point 45

Coordinates: S 6°61569 / E 31°76971

Slope: +/- flat

Dominant tree height: 10-12 m

Basal area (Bitterlich, k=1): not recorded

Photos: 4042 / 4043 / 4044 / 4045

Disturbances/notes: + 180 steps (1) tree cut for honey (>3 years)

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Julbernardia globiflora</i> | 22 | 23.05 |
| | <i>Pericopsis angolensis</i> | 54 | 5.70 |
| | <i>Julbernardia globiflora</i> | 82 | 15.63 |
| | <i>Julbernardia globiflora</i> | 61 | 8.50 |
| DBH < 20 cm | <i>Combretum fragrans</i> | 3 | 4.81 |
| | <i>Diplorhynchus condylocarpon</i> | 12 | 3.72 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 13 | 4.96 |
| | <i>Diplorhynchus condylocarpon</i> | 9 | 3.75 |

Sampling point 46

Coordinates: S 6°61701 / E 31°76800

Slope: +/- flat

Dominant tree height: 10-12 m

Basal area (Bitterlich, k=1): not recorded

Photos: 4046 / 4047 / 4048 / 4049

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Monotes africanus</i> | 29 | 21.70 |
| | <i>Monotes africanus</i> | 25 | 7.34 |
| | <i>Brachystegia manga</i> | 37 | 15.05 |
| | <i>Monotes africanus</i> | 38 | 8.25 |
| DBH < 20 cm | <i>Julbernardia globiflora</i> | 6 | 1.63 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 10 | 6.04 |
| | <i>Brachystegia manga</i> | 4 | 6.59 |
| | <i>Julbernardia globiflora</i> | 4 | 3.45 |

Sampling point 47

Coordinates: S 6°61835 / E 31°76674

Slope: +/- flat

Dominant tree height: 10-12 m

Basal area (Bitterlich, k=1): not recorded

Photos: 4053 / 4054 / 4055 / 4056

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Erythrophleum africanum</i> | 23 | 12.75 |
| | <i>Borassus aethiopum</i> | 55 | 17 |
| | <i>Brachystegia manga</i> | 80 | 17.80 |
| | <i>Brachystegia spiciformis</i> | 50 | 17.35 |
| DBH < 20 cm | <i>Flacourtie indica</i> | 13 | 3.65 |
| | <i>Phyllocosmus leimareanus</i> | 5 | 5.40 |
| | <i>Lannea schimperi</i> | 16 | 2.47 |
| | <i>Hymenocardia acida</i> | 4 | 1.80 |

Sampling point 48

Coordinates: S 6°61998 / E 31°76550

Slope exposure: SW avec azimuth 227 (N)

Dominant tree height: 10 m

Basal area (Bitterlich, k=1): not recorded

Photos: 4064 / 4065 / 4066 / 4067

Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Brachystegia manga</i> | 25 | 16.20 |
| | <i>Brachystegia manga</i> | 23 | 13.05 |
| | <i>Brachystegia manga</i> | 44 | 14.60 |
| | <i>Brachystegia manga</i> | 53 | 9.90 |
| DBH < 20 cm | <i>Flacourtie indica</i> | 6 | 8.00 |
| | <i>Ximenia caffra</i> | 4 | 10.33 |
| | <i>Brachystegia manga</i> | 4 | 2.10 |
| | <i>Combretum molle</i> | 18 | 2.48 |

Sampling point 49

Coordinates: S 6°62115 / E 31°76402
 Slope exposure: SW avec azimuth 227 (N)
 Dominant tree height: 12-14 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4068 / 4069 / 4070 / 4071
 Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|--|----------|--------------|
| DBH ≥ 20 cm | <i>Combretum collinum</i> | 26 | 10.47 |
| | <i>Monotes africanus</i> | 45 | 10.77 |
| | <i>Julbernardia globiflora</i> | 29 | 3.80 |
| | <i>Kigelia africana</i> | 30 | 22.71 |
| DBH < 20 cm | <i>Diplorhynchus condylocarpon</i> | 13 | 3.40 |
| | <i>Diplorhynchus condylocarpon</i> | 6 | 2 |
| | <i>Ximenia caffra</i> | 3 | 3.90 |
| | <i>Pseudolachnostylis maprouneifolia</i> | 15 | 7.30 |

Sampling point 50

Coordinates: S 6°62278 / E 31°76323
 Slope exposure: NORD
 Dominant tree height: 10 m
 Basal area (Bitterlich, k=1): not recorded
 Photos: 4072 / 4073 / 4074 / 4075
 Disturbances/notes:

| Diameter class | Nearest species in each quarter | DBH (cm) | Distance (m) |
|----------------|---------------------------------|----------|--------------|
| DBH ≥ 20 cm | <i>Pterocarpus angolensis</i> | 32 | 6.70 |
| | <i>Brachystegia microphylla</i> | 39 | 20.00 |
| | <i>Brachystegia microphylla</i> | 25 | 17.05 |
| | <i>Julbernardia globiflora</i> | 29 | 16.65 |
| DBH < 20 cm | <i>Hymenocardia acida</i> | 5 | 3.70 |
| | <i>Brachystegia microphylla</i> | 7 | 0.86 |
| | <i>Brachystegia microphylla</i> | 5 | 2.04 |
| | <i>Terminalia kaiseriana</i> | 15 | 3.20 |

Appendix C: Tree and shrub checklist Mlele BKZ (27/3/2019)

Plant list revised and amended from Mwangulango 2004

| Scientific name | Kikonongo |
|--|---------------------------|
| <i>Acacia gerrardii</i> | Ulula |
| <i>Acacia polyacantha</i> | Muwombwe |
| <i>Acacia stuhlmannii</i> | Nunga? |
| <i>Afzelia quanzensis</i> | Mkola |
| <i>Albizia antunesiana</i> | Mphilipili |
| <i>Albizia glaberrima</i> | |
| <i>Albizia grandibracteata</i> | |
| <i>Albizia gummosa</i> | |
| <i>Albizia harveyi</i> | |
| <i>Albizia versicolor</i> | Masako |
| <i>Anisophyllea pomifera</i> | Msindwi |
| <i>Annona senegalensis</i> | Mfilafila |
| <i>Antidesma membranaceum</i> | Msekela |
| <i>Azanza garckeana</i> | Mtowo |
| <i>Bauhinia thonningii</i> | Mfundwa mbogo |
| <i>Bobgunnia madagascariensis</i> | Kasanda |
| <i>Borassus aethiopum</i> | Sandala |
| <i>Brachystegia boehmii</i> | |
| <i>Brachystegia floribunda?</i> | Msilanga |
| <i>Brachystegia glaucescens</i> | Myombo |
| <i>Brachystegia microphylla</i> | Mkongolo |
| <i>Brachystegia spiciformis</i> | Mtundu / Umtundu |
| <i>Brachystegia stipulata</i> | |
| <i>Brachystegia utilis/floribunda/manga?</i> | Msilanga |
| <i>Brachystegia taxifolia</i> | Kapepe |
| <i>Bridelia duvigneaudii</i> | |
| <i>Bridelia scleroneura</i> | |
| <i>Burkea africana</i> | Mgandosinsi |
| <i>Cassia abbreviata</i> | Mzoka |
| <i>Cassia singueana</i> | Mzoka |
| <i>Cassipourea mollis</i> | Mlugala |
| <i>Catunaregam spinosa</i> | Mpongole |
| <i>Chrysophyllum bangweolense</i> | |
| <i>Combretum collinum</i> | Mlandala |
| <i>Combretum fragrans</i> | Mlozyaminze / Mluziaminzi |
| <i>Combretum molle</i> | Mlama |
| <i>Combretum zeyheri</i> | Msana |
| <i>Commiphora africana</i> | Mponda |
| <i>Crossopteryx febrifuga</i> | Msanza |
| <i>Dalbergia nitidula</i> | Kapondalampassa, Mzoka? |
| <i>Dichrostachys cinerea</i> | Kasunzulu |
| <i>Diospyros cornii</i> | Mnumbulu |
| <i>Diplorhynchus condylocarpon</i> | Msonga |
| <i>Dombeya rotundifolia</i> | Mlalila |
| <i>Ekebergia capensis</i> | Mtuzya |
| <i>Elaeodendron schweinfurthianum</i> | |
| <i>Entada abyssinica</i> | Mfutwamvula / Kamchicha |
| <i>Eriosema sp.</i> | |
| <i>Erythrina abyssinica</i> | Kamchicha |
| <i>Erythrophleum africanum</i> | Mgandongoye |
| <i>Euphorbia matabelensis</i> | Kiponda |

| | |
|--|---------------------------------|
| <i>Flacourtie indica</i> | Msungu |
| <i>Friesodielsia obovata</i> | Msalansi |
| <i>Garcinia huillensis</i> | Myeye |
| <i>Grewia bicolor</i> | Mkoma |
| <i>Hexalobus monopetalus</i> | Mukua, Mukwa? |
| <i>Hymenocardia acida</i> | Kapala |
| <i>Isoberlinia angolensis</i> | Mnembela |
| <i>Julbernardia globiflora</i> | Muva |
| <i>Kigelia africana</i> | Mdungwa |
| <i>Lannea schimperi</i> | Mgumbu |
| <i>Lonchocarpus capassa</i> | Mvalevale |
| <i>Maprounea africana</i> | |
| <i>Maranthes floribunda</i> | Mwasha |
| <i>Markhamia obtusifolia</i> | Mpapa |
| <i>Maytenus senegalensis</i> | Mwesia |
| <i>Memecylon flavovirens</i> | Mseweye |
| <i>Monanthotaxis discolor</i> | Mshenene |
| <i>Monotes africanus</i> | Mkokote |
| <i>Monotes katangensis</i> | Mukokoti |
| <i>Multidentia crassa</i> | Mukukumba |
| <i>Mundelea sericea</i> | |
| <i>Mystroxylon aethiopicum</i> | Kasela |
| <i>Ochna afzelii ssp. afzelii</i> | |
| <i>Ochna holstii</i> | |
| <i>Ochna inermis</i> | |
| <i>Ochna oxyphylla</i> | |
| <i>Olax obtusifolia</i> | Mtundwa |
| <i>Oldfieldia dactylophylla</i> | Mliwamfwengi |
| <i>Ozoroa insignis subsp. reticulata</i> | Mkalakala |
| <i>Parinari curatellifolia</i> | Mbula/Mhula |
| <i>Pavetta stuhlmannii</i> | |
| <i>Pericopsis angolensis</i> | Mbanga |
| <i>Phyllanthus engleri</i> | Mng'ongomtandala |
| <i>Phyllocosmus leimareanus</i> | Msonifya |
| <i>Pleurostylia africana</i> | |
| <i>Protea madiensis</i> | |
| <i>Pseudolachnostylis maprouneifolia</i> | Mtungulu |
| <i>Psorospermum febrifugum</i> | Mvivi |
| <i>Pterocarpus angolensis</i> | Mninga |
| <i>Pterocarpus tinctorius</i> | Mkulungu |
| <i>Rhus longipes</i> | Msilanswagalo |
| <i>Rhus vulgaris</i> | Kankiningi |
| <i>Rytigynia decussata</i> | |
| <i>Rytigynia uhlriegii</i> | Msongwansimba |
| <i>Rothmannia engleriana</i> | Mlozirozi |
| <i>Schrebera trichoclada</i> | Mputeka |
| <i>Sclerocarya birrea</i> | |
| <i>Securidaca longepedunculata</i> | Mteywe |
| <i>Sterculia africana</i> | Msawala |
| <i>Sterculia quinqueloba</i> | Mkungulanga / Msavala / Msawala |
| <i>Stereospermum kunthianum</i> | |
| <i>Strychnos innocua</i> | Mkulwa |
| <i>Strychnos potatorum</i> | Mnyekenyewe, Mgwegue |
| <i>Strychnos pungens</i> | Mkome? |
| <i>Strychnos spinosa</i> | Katonga, Mwaye |
| <i>Syzygium guineense</i> | Kashamongo |
| <i>Tamarindus indica</i> | |
| <i>Terminalia kaiseriana</i> | Kazima |

| | |
|-----------------------------------|------------|
| <i>Terminalia mollis</i> | Mfufu |
| <i>Trichilia emetica</i> | Mkalya |
| <i>Uapaca kirkiana</i> | Mkusu |
| <i>Uapaca nitida</i> | Mkokofinyo |
| <i>Vangueria madagascariensis</i> | Mgelelya |
| <i>Vitex doniana</i> | Mfurū |
| <i>Vitex fischeri?</i> | |
| <i>Vitex madiensis</i> | Mfululegea |
| <i>Vitex mombassae</i> | Mfululegea |
| <i>Vitex payos</i> | Mtalali |
| <i>Xeroderris stuhlmannii</i> | |
| <i>Ximenia americana?</i> | Mtundwa |
| <i>Ximenia caffra</i> | Kaguvguva |
| <i>Ziziphus mucronata</i> | Kagaole |

Appendix D: Vegetation survey 2018 of Mlele Beekeeping Zone

