

新農業展開ゲノムプロジェクトにおける論文一覧

番号	論文名
1	Masaoka T, Okamoto H, Araki K, Nagoya H, Fujiwara A, Kobayashi T. Population distinction of <i>Oryzias latipes</i> and identification of F1 hybrid between populations using nuclear genes and mitochondrial gene region. <i>DNA Testing</i> 3, 13-27 (2011)
2	Ohmori S, Tabuchi H, Yatou O, Yoshida H. Agronomic traits and gene containment ability of rice cleistogamous lines with the superwoman1-cleistogamy mutation. <i>Breeding Science</i> 62,124 (2012)
3	Kobayashi K, Yasuno N, Sato Y, Yoda Y, Yamazaki R, Kimizu M, Yoshida H, Nagamura Y, Kyojuka J. Inflorescence Meristem Identity in Rice Is Specified by Overlapping Functions of Three AP1/FUL-Like MADS Box Genes and PAP2, a SEPALLATA MADS Box Gene. <i>Plant Cell</i> 24, 1848 (2012)
4	Yoshida H. Analysis and Utilization of a Cleistogamous Mutant of Rice. <i>Gamma Field Symposia</i> 49,17 (2012)
5	Yukuhiro K, Iwata K, Kōmoto N, Tomita S, Itoh M, Kiuchi M (2012) Nucleotide sequences of mitochondrial cytochrome C oxidase subunit I (COI) gene show clear differences between the domesticated silkworm <i>Bombyx mori</i> and the wild mulberry silkworm <i>Bombyx mandarina</i> from Japan <i>Journal of Insect Biotechnology and Sericology</i> 81(1):29-35
6	J. Mano, M. Harada, R. Takabatake, S. Furui, K. Kitta, K. Nakamura, H. Akiyama, R. Teshima, H. Noritake, S. Hatano, S. Futo, Y. Minegishi, T. Iizuka, Comprehensive GMO Detection Using Real-Time PCR Array: Single-Laboratory Validation, <i>J. AOAC Intl.</i> , 95(2) 508-516(2012)
7	J. Mano, S. Furui, K. Takashima, T. Koiwa, S. Futo, Y. Minegishi, H. Akiyama, R. Teshima, T. Kurashima, R. Takabatake, K. Kitta, Development and validation of event-specific quantitative PCR method for genetically modified maize MIR604, <i>食品衛生学雑誌</i> , 53(4)166-171(2012)
8	Masaoka T, Okamoto H, Araki K, Nagoya H, Fujiwara A, Kobayashi T. Identification of the hybrid between <i>Oryzias latipes</i> and <i>Oryzias curvinotus</i> using nuclear genes and mitochondrial gene region. <i>Marine Genomics</i> 7, 37-41 (2012)
9	Masaoka T, Okamoto H, Araki K, Nagoya H, Fujiwara A, Kobayashi T. Sterility of triploid F2 hybrid between <i>Oryzias latipes</i> and <i>Oryzias curvinotus</i> . <i>Fish genetics and breeding science</i> 41, 43-49 (2012)
10	小池説夫、大森伸之介、山口知哉、林高見、矢頭治、吉田均.閉花性イネ突然変異体の開花期耐冷性.日本作物学会東北支部会報.55:67-68 (2012年8月21日)
11	R. Takabatake, M. Onishi, T. Koiwa, S. Futo., Y. Minegishi, H. Akiyama, R. Teshima, T. Kurashima, J. Mano, S. Furui, K. Kitta, Development and interlaboratory validation of quantitative PCR method for screening analysis of genetically modified soybeans, <i>Biological & Pharmaceutical Bulletin</i> , 36(1)131-134(2013)
12	Yuhiro K, Sezutsu H, Tamura T, Kosegawa E, Iwata K, Ajimura M, Gu S-H, Wang M, Xia Q, Mita K, Kiuchi M (2013) Little gene flow between domestic silkworm <i>Bombyx mori</i> and its wild relative <i>Bombyx mandarina</i> in Japan, and possible artificial selection on the <i>CAD</i> gene of <i>B. mori</i> <i>Genes & Genetic Systems</i> 87(5):331-340
13	小池説夫、大森伸之介、山口知哉、林高見、矢頭 治、吉田 均. The closed flowering rice is tolerant to high temperature stress at flowering. <i>Plant Production Science</i> .2012年以内に投稿予定
14	A. Matsushita, H. Inoue, S. Goto, A. Nakayama, S. Sugano, N. Hayashi, H. Takatsuji. The nuclear ubiquitin proteasome degradation affects WRKY45 function in the rice defense program. <i>Plant J</i> 73(2), 302-313(2013)

15	Takefumi Hattori, Shinya Murakami, Mai Mukai, Tatsuhiko Yamada, Hirohiko Hirochika, Masakazu Ike, Ken Tokuyasu, Shiro Suzuki, Masahiro Sakamoto, Toshiaki Umezawa. Rapid analysis of transgenic rice straw using near-infrared spectroscopy. <i>Plant Biotechnology</i> 29(4), 359–366(2012)
16	C. J. Jiang, M. Shimono, S. Sugano, M. Kojima, X. Liu, H. Inoue, H. Sakakibara, H. Takatsuji. Cytokinins act synergistically with salicylic acid to activate defense gene expression in rice. <i>Mol Plant Microbe Interact</i> , in press(2012)
17	T. Shimizu, T. Ogamino, A. Hiraguri, E. Nakazono-Nagaoka, T. Uehara-Ichiki, M. Nakajima, K. Akutsu, T. Omura, T. Sasaya. Strong resistance against Rice grassy stunt virus is induced in transgenic rice plants expressing dsRNA of the viral genes for nucleocapsid or movement proteins as targets for RNA interference. <i>Phytopathology</i> , in press(2012)
18	T. Yamamoto, A. Nakamura, H. Iwai, T. Ishii, J. F. Ma, R. Yokoyama, K. Nishitani, S. Satoh, J. Furukawa. Effect of silicon deficiency on secondary cell wall synthesis in rice leaf. <i>J Plant Res</i> 125(6), 771–779(2012)
19	M. Shimono, H. Koga, A. Akagi, N. Hayashi, S. Goto, M. Sawada, T. Kurihara, A. Matsushita, S. Sugano, C. J. Jiang, H. Kaku, H. Inoue, H. Takatsuji. Rice WRKY45 plays important roles in fungal and bacterial disease resistance. <i>Mol Plant Pathol</i> 13(1), 83–94(2012)
20	Katsumi Yazawa, Chang-Jie Jiang, Mikiko Kojima, Hitoshi Sakakibara, Hiroshi Takatsuji. Reduction of abscisic acid levels or inhibition of abscisic acid signaling in rice during the early phase of <i>Magnaporthe grisea</i> infection decreases its susceptibility to the fungus. <i>Physiol Mol Plant Pathol</i> 78, 1–7(2012)
21	K. Takeuchi, A. Gyohda, M. Tominaga, M. Kawakatsu, A. Hatakeyama, N. Ishii, K. Shimaya, T. Nishimura, M. Riemann, P. Nick, M. Hashimoto, T. Komano, A. Endo, T. Okamoto, Y. Jikumaru, Y. Kamiya, T. Terakawa, T. Koshiba. RSOsPR10 expression in response to environmental stresses is regulated antagonistically by jasmonate/ethylene and salicylic acid signaling pathways in rice roots. <i>Plant Cell Physiol</i> 52(9), 1686–1696(2011)
22	T. Konishi, T. Aohara, T. Igasaki, N. Hayashi, Y. Miyazaki, A. Takahashi, H. Hirochika, H. Iwai, S. Satoh, T. Ishii. Down-regulation of UDP-arabinopyranose mutase reduces the proportion of arabinofuranose present in rice cell walls. <i>Phytochemistry</i> 72(16), 1962–1968(2011)
23	Y. Arai-Sanoh, M. Ida, R. Zhao, S. Yoshinaga, T. Takai, T. Ishimaru, H. Maeda, K. Nishitani, Y. Terashima, M. Gau, N. Kato, M. Matsuoka, M. Kondo. Genotypic variations in non-structural carbohydrate and cell-wall components of the stem in rice, sorghum, and sugar vane. <i>Biosci Biotechnol Biochem</i> 75(6), 1104–1112(2011)
24	Y. Sato, Y. Masuta, K. Saito, S. Murayama, K. Ozawa. Enhanced chilling tolerance at the booting stage in rice by transgenic overexpression of the ascorbate peroxidase gene, OsAPXa. <i>Plant Cell Rep</i> 30(3), 399–406(2011)
25	C. J. Jiang, M. Shimono, S. Sugano, M. Kojima, K. Yazawa, R. Yoshida, H. Inoue, N. Hayashi, H. Sakakibara, H. Takatsuji. Abscisic Acid Interacts Antagonistically with Salicylic Acid Signaling Pathway in Rice– <i>Magnaporthe grisea</i> Interaction. <i>Mol Plant Microbe Interact</i> 23(6), 791–798(2010)
26	T. Konishi, M. Ohnishi-Kameyama, K. Funane, Y. Miyazaki, T. Konishi, T. Ishii. An arginyl residue in rice UDP-arabinopyranose mutase is required for catalytic activity and autoglycosylation. <i>Carbohydr Res</i> 345(6), 787–791(2010)
27	Tomoyuki Konishi, Yasumasa Miyazaki, Seiyei Yamakawa, Hiroaki Iwai, Shinobu Satoh, Tadashi Ishii. Purification and biochemical characterization of recombinant rice UDP-arabinopyranose mutase generated in insect cells. <i>Biosci Biotechnol Biochem</i> 74(1), 191–194(2010)
28	H. Kato, G Xie, Y. Sato, R Imai. Isolation of anther-specific gene promoters suitable for transgene expression in rice. <i>Plant Mol. Biol. Rep.</i> 28(3), 381–387(2010)
29	G. Xie, H. Kato, K. Sasaki, R. Imai. A cold-induced thioredoxin h of rice, OsTrx23, negatively regulates kinase activities of OsMPK3 and OsMPK6 in vitro. <i>FEBS Lett</i> 583(17), 2734–2738(2009)
30	C. J. Jiang, M. Shimono, S. Maeda, H. Inoue, M. Mori, M. Hasegawa, S. Sugano, H. Takatsuji. Suppression of the rice fatty-acid desaturase gene OsSSI2 enhances resistance to blast and leaf blight diseases in rice. <i>Mol Plant Microbe Interact</i> 22(7), 820–829(2009)
31	T. Shimizu, M. Yoshii, T. Wei, H. Hirochika, T. Omura. Silencing by RNAi of the gene for Pns12, a viroplasm matrix protein of Rice dwarf virus, results in strong resistance of transgenic rice plants to the virus. <i>Plant Biotechnol J</i> 7(1), 24–32(2009)

32	Y. W. Feng, S. Komatsu, T. Furukawa, T. Koshiba, Y. Kohno. Proteome analysis of proteins responsive to ambient and elevated ozone in rice seedlings. <i>Agric Ecosyst Environ</i> 125(1-4), 255-265(2008)
33	Y. Taniguchi, M. Kawata, I. Ando, T. Shimizu, M. Ohshima. Selecting genetic transformants of indica and indica-derived rice cultivars using bispyribac sodium and a mutated ALS gene. <i>Plant Cell Rep</i> 29(11), 1287-1295(2010)
34	M. Kishi-Kaboshi, K. Okada, L. Kurimoto, S. Murakami, T. Umezawa, N. Shibuya, H. Yamane, A. Miyao, H. Takatsuji, A. Takahashi, H. Hirochika. A rice fungal MAMP-responsive MAPK cascade regulates metabolic flow to antimicrobial metabolite synthesis. <i>Plant J</i> 63(4), 599-612(2010)
35	N. Hayashi, H. Inoue, T. Kato, T. Funao, M. Shirota, T. Shimizu, H. Kanamori, H. Yamane, Y. Hayano-Saito, T. Matsumoto, M. Yano, H. Takatsuji. Durable panicle blast-resistance gene Pb1 encodes an atypical CC-NBS-LRR protein and was generated by acquiring a promoter through local genome duplication. <i>Plant J</i> 64(3), 498-510(2010)
36	S. Sugano, C. J. Jiang, S. Miyazawa, C. Masumoto, K. Yazawa, N. Hayashi, M. Shimono, A. Nakayama, M. Miyao, H. Takatsuji. Role of OsNPR1 in rice defense program as revealed by genome-wide expression analysis. <i>Plant Mol Biol</i> 74(6), 549-562(2010)
37	Michihiro Ito1, Akira Ono, Yoshiyuki Ohtsubo, Masataka Tsuda, Yuji Nagata. Cloning of γ -hexachlorocyclohexane dehydrochlorinase gene with its flanking regions from soil by activity-based screening techniques. 52, 16-19(2012)
38	Yoshihiko NANASATO, Ken-ichi KONAGAYA, Ayako OKUZAKI, Mai TSUDA, Yutaka TABELI. Improvement of Agrobacterium-mediated transformation of cucumber (<i>Cucumis sativus</i> L.) by combination of vacuum infiltration and co-cultivation on filter paper wicks. , <i>Plant Biotechnology Reports (Online First)</i> (2012)
39	R. Takahashi, K. Bashir, Y. Ishimaru, N. K. Nishizawa, H. Nakanishi. The role of heavy-metal ATPases, HMAs, in zinc and cadmium transport in rice. 7(12), (2012)
40	Y. Ishimaru, K. Bashir, H. Nakanishi, N. K. Nishizawa. OsNRAMP5, a major player for constitutive iron and manganese uptake in rice. 7(7), 763-766(2012)
41	T. Mase, H. Yabuki, M. Okai, J. Ohtsuka, F. L. Imai, Y. Nagata, M. Tanokura. Crystallization and preliminary X-ray analysis of the haloalkane dehalogenase DatA from <i>Agrobacterium tumefaciens</i> C58. 68(Pt 6), 652-654(2012)
42	R. Takahashi, Y. Ishimaru, H. Shimo, Y. Ogo, T. Senoura, N. K. Nishizawa, H. Nakanishi. The OsHMA2 transporter is involved in root-to-shoot translocation of Zn and Cd in rice. 35(11), 1948-1957(2012)
43	Y. Ishimaru, R. Takahashi, K. Bashir, H. Shimo, T. Senoura, K. Sugimoto, K. Ono, M. Yano, S. Ishikawa, T. Arao, H. Nakanishi, N. K. Nishizawa. Characterizing the role of rice NRAMP5 in Manganese, Iron and Cadmium Transport. 2, 286(2012)
44	A. T. Nguyen, Y. Sato, T. Iwasaki, K. Miyauchi, M. Tokuda, D. Kasai, E. Masai, M. Fukuda. Characterization of the 1,1-dichloro-2,2-bis(4-chlorophenyl)ethylene (DDE) degradation system in <i>Janibacter</i> sp. TYM3221. 49(6-7), 532-539(2011)
45	Y. Nagata, S. Natsui, R. Endo, Y. Ohtsubo, N. Ichikawa, A. Ankai, A. Oguchi, S. Fukui, N. Fujita, M. Tsuda. Genomic organization and genomic structural rearrangements of <i>Sphingobium japonicum</i> UT26, an archetypal gamma-hexachlorocyclohexane-degrading bacterium. 49(6-7), 499-508(2011)
46	R. Takahashi, Y. Ishimaru, H. Nakanishi, N. K. Nishizawa. Role of the iron transporter OsNRAMP1 in cadmium uptake and accumulation in rice. 6(11), 1813-1816(2011)
47	Y. Ishimaru, K. Bashir, H. Nakanishi, N. K. Nishizawa. The role of rice phenolics efflux transporter in solubilizing apoplasmic iron. 6(10), 1624-1626(2011)
48	H. Shimo, Y. Ishimaru, G. An, T. Yamakawa, H. Nakanishi, N. K. Nishizawa. Low cadmium (LCD), a novel gene related to cadmium tolerance and accumulation in rice. 62(15), 5727-5734(2011)
49	R. Takahashi, Y. Ishimaru, T. Senoura, H. Shimo, S. Ishikawa, T. Arao, H. Nakanishi, N. K. Nishizawa. The OsNRAMP1 iron transporter is involved in Cd accumulation in rice. 62(14), 4843-4850(2011)

50	R. Chaloupkova, Z. Prokop, Y. Sato, Y. Nagata, J. Damborsky. Stereoselectivity and conformational stability of haloalkane dehalogenase DbjA from <i>Bradyrhizobium japonicum</i> USDA110: the effect of pH and temperature. <i>278</i> (15), 2728–2738(2011)
51	Y. Ishimaru, Y. Kakei, H. Shimo, K. Bashir, Y. Sato, N. Uozumi, H. Nakanishi, N. K. Nishizawa. A rice phenolic efflux transporter is essential for solubilizing precipitated apoplasmic iron in the plant stele. <i>286</i> (28), 24649–24655(2011)
52	Y. Nanasato, K. Konagaya, A. Okuzaki, M. Tsuda, Y. Tabei. <i>Agrobacterium</i> -mediated transformation of kabocha squash (<i>Cucurbita moschata</i> Duch) induced by wounding with aluminum borate whiskers. <i>30</i> (8), 1455–1464(2011)
53	M. Tabata, R. Endo, M. Ito, Y. Ohtsubo, A. Kumar, M. Tsuda, Y. Nagata. The <i>lin</i> genes for gamma-hexachlorocyclohexane degradation in <i>Sphingomonas</i> sp. MM-1 proved to be dispersed across multiple plasmids. <i>75</i> (3), 466–472(2011)
54	K. Yamazaki, M. Suzuki, T. Itoh, K. Yamamoto, M. Kanemitsu, C. Matsumura, T. Nakano, T. Sakaki, Y. Fukami, H. Imaishi, H. Inui. Structural basis of species differences between human and experimental animal CYP1A1s in metabolism of 3,3',4,4',5-pentachlorobiphenyl. <i>149</i> (4), 487–494(2011)
55	K. Hasan, A. Fortova, T. Koudelakova, R. Chaloupkova, M. Ishitsuka, Y. Nagata, J. Damborsky, Z. Prokop. Biochemical characteristics of the novel haloalkane dehalogenase <i>DatA</i> , isolated from the plant pathogen <i>Agrobacterium tumefaciens</i> C58. <i>77</i> (5), 1881–1884(2011)
56	T. Prudnikova, R. Chaloupkova, Y. Sato, Y. Nagata, O. Degtjarik, M. Kutý, P. Rezacova, J. Damborsky, I. Smatanova. Development of a crystallization protocol for the <i>DbeA1</i> variant of novel haloalkane dehalogenase from <i>Bradyrhizobium elkanii</i> USDA94. <i>11</i> (2), 516–519(2011)
57	Khurram Bashir, Yasuhiro Ishimaru, Hugo Shimo, Yusuke Kakei, Takeshi Senoura, Ryuichi Takahashi, Yutaka Sato, Yuki Sato, Nobuyuki Uozumi, Hiromi Nakanishi, Naoko K. Nishizawa. Rice phenolics efflux transporter 2 (PEZ2) plays an important role in solubilizing apoplasmic iron. <i>57</i> (6), 803–812(2011)
58	佐藤優花里, 夏目亮, Z. Prokop, J. Brezovsky, R. Chaloupkova, J. Damborsky, 永田裕二, 千田俊哉. ハロアルカン脱ハロゲン酵素DjbAの鏡像異性体選択性機構の解明. <i>53</i> (2), 124–129(2011)
59	Hirotsu Murano, Takashi Otani, Nobuyasu Seike, Mizuki Sakai. Dieldrin uptake and translocation in plants growing in hydroponic medium. <i>29</i> (1), 142–148(2010)
60	M. Okai, K. Kubota, M. Fukuda, Y. Nagata, K. Nagata, M. Tanokura. Crystallization and preliminary X-ray analysis of gamma-hexachlorocyclohexane dehydrochlorinase <i>LinA</i> from <i>Sphingobium japonicum</i> UT26. <i>65</i> (Pt 8), 822–824(2009)
61	T. Prudnikova, T. Mozga, P. Rezacova, R. Chaloupkova, Y. Sato, Y. Nagata, J. Brynda, M. Kutý, J. Damborsky, I. K. Smatanova. Crystallization and preliminary X-ray analysis of a novel haloalkane dehalogenase <i>DbeA</i> from <i>Bradyrhizobium elkanii</i> USDA94. <i>65</i> (Pt 4), 353–356(2009)
62	M. Pavlova, M. Klvana, Z. Prokop, R. Chaloupkova, P. Banas, M. Otyepka, R. C. Wade, M. Tsuda, Y. Nagata, J. Damborsky. Redesigning dehalogenase access tunnels as a strategy for degrading an anthropogenic substrate. <i>5</i> (10), 727–733(2009)
63	Ippei Ogawa, Hiromi Nakanishi, Satoshi Mori, Naoko K. Nishizawa. Time course analysis of gene regulation under cadmium stress in rice. <i>325</i> (1–2), 97–108(2009)
64	G. Fuchu, Y. Ohtsubo, M. Ito, R. Miyazaki, A. Ono, Y. Nagata, M. Tsuda. Insertion sequence-based cassette PCR: cultivation-independent isolation of gamma-hexachlorocyclohexane-degrading genes from soil DNA. <i>79</i> (4), 627–632(2008)
65	Prokop, Z., Y. Sato, J. Brezovsky, T. Mozga, R. Chaloupkova, T. Koudelakova, P. Jerabek, V. Stepankova, R. Natsume, J. G.E. van Leeuwen, D. B. Janssen, J. Florian, Y. Nagata, T. Senda, and J. Damborsky. Enantioselectivity of Haloalkane Dehalogenases and its Modulation by Surface Loop Engineering. <i>Angewandte Chemie</i> . <i>49</i> : 6111–6115 (2010)
66	Okai, M., K. Kubota, M. Fukuda, Y. Nagata, K. Nagata, and M. Tanokura. Crystal structure of g-hexachlorocyclohexane dehydrochlorinase <i>LinA</i> from <i>Sphingobium japonicum</i> UT26. <i>J. Mol. Biol.</i> <i>403</i> : 260–269 (2010)
67	Nagata, Y., Y. Ohtsubo, R. Endo, N. Ichikawa, A. Ankai, A. Oguchi, S. Fukui, N. Fujita, and M. Tsuda. Complete genome sequence of the representative g-hexachlorocyclohexane-degrading bacterium <i>Sphingobium japonicum</i> UT26. <i>J. Bacteriology</i> <i>192</i> : 5852–5853 (2010)

68	Murano H, Otani T, Seike N. Dieldrin-dissolving abilities of the xylem saps of several plant families, particularly Cucurbita pepo L. <i>Environ Toxicol Chem.</i> 2010 Oct;29(10):2269-77. doi: 10.1002/etc.288.
69	Fumio Takaiwa, Chikahiro Hiroi. Rice seed-based allergy vaccines: induction of allergen-specific oral tolerance against cedar pollen and house dust mite allergies. <i>Molecular Vaccines-From Prophylaxis to Therapy</i> , (2013)
70	Y. Shimajiri, K. Ozaki, K. Kainou, K. Akama. Differential subcellular localization, enzymatic properties and expression patterns of gamma-aminobutyric acid transaminases (GABA-Ts) in rice (<i>Oryza sativa</i>). <i>J Plant Physiol</i> 170(2), 196-201(2013)
71	Y. Wakasa, H. Takagi, S. Hirose, L. Yang, M. Saeki, T. Nishimura, O. Kaminuma, T. Hiroi, F. Takaiwa. Oral immunotherapy with transgenic rice seed containing destructured Japanese cedar pollen allergens, Cry j 1 and Cry j 2, against Japanese cedar pollinosis. <i>Plant Biotechnol J</i> 11(1), 66-76(2013)
72	Y. Wakasa, H. Yasuda, F. Takaiwa. Secretory type of recombinant thioredoxin h induces ER stress in endosperm cells of transgenic rice. <i>J Plant Physiol</i> 170(2), 202-210(2013)
73	M. Yoshikawa, K. Ohinata, Y. Yamada. The pharmacological effects of novokinin, a designed peptide agonist of the angiotensin AT2 receptor. <i>Curr Pharm Des</i> , (2012)
74	Y. I. Kwon, K. Abe, K. Osakabe, M. Endo, A. Nishizawa-Yokoi, H. Saika, H. Shimada, S. Toki. Overexpression of OsRecQ14 and/or OsExo1 Enhances DSB-Induced Homologous Recombination in Rice. <i>Plant Cell Physiol</i> 53(12), 2142-2152(2012)
75	C. Cabanos, A. Ekyo, Y. Amari, N. Kato, M. Kuroda, S. Nagaoka, F. Takaiwa, S. Utsumi, N. Maruyama. High-level production of lactostatin, a hypocholesterolemic peptide, in transgenic rice using soybean A1aB1b as carrier. <i>Transgenic Res</i> , (2012)
76	A. Nishizawa-Yokoi, S. Nonaka, H. Saika, Y. I. Kwon, K. Osakabe, S. Toki. Suppression of Ku70/80 or Lig4 leads to decreased stable transformation and enhanced homologous recombination in rice. <i>New Phytol</i> 196(4), 1048-1059(2012)
77	H. Saika, S. Nonaka, K. Osakabe, S. Toki. Sequential monitoring of transgene expression following Agrobacterium-mediated transformation of rice. <i>Plant Cell Physiol</i> 53(11), 1974-1983(2012)
78	S. Hayashi, Y. Wakasa, F. Takaiwa. Functional integration between defence and IRE1-mediated ER stress response in rice. <i>Sci Rep</i> 2, 670(2012)
79	T. Kawakatsu, F. Tamaoki. Reduction of 13 kD prolamins increases recombinant protein yield and recovery rate in rice endosperm. <i>Plant Signal Behav</i> 7(11), 1402-1403(2012)
80	L. Yang, S. Hirose, H. Takahashi, T. Kawakatsu, F. Takaiwa. Recombinant protein yield in rice seed is enhanced by specific suppression of endogenous seed proteins at the same deposit site. <i>Plant Biotechnol J</i> 10(9), 1035-1045(2012)
81	T. Kawakatsu, Y. Wakasa, H. Yasuda, F. Takaiwa. RNA silencing induced by an artificial sequence that prevents proper transcription termination in rice. <i>Plant Physiol</i> 160(2), 601-612(2012)
82	Y. Wakasa, K. Ozawa, F. Takaiwa. Agrobacterium-mediated co-transformation of rice using two selectable marker genes derived from rice genome components. <i>Plant Cell Rep</i> 31(11), 2075-2084(2012)
83	Y. Wakasa, S. Hayashi, F. Takaiwa. Expression of OsBiP4 and OsBiP5 is highly correlated with the endoplasmic reticulum stress response in rice. <i>Planta</i> 236(5), 1519-1527(2012)
84	K. Ozawa, Y. Wakasa, Y. Ogo, K. Matsuo, H. Kawahigashi, F. Takaiwa. Development of an efficient agrobacterium-mediated gene targeting system for rice and analysis of rice knockouts lacking granule-bound starch synthase (Waxy) and beta1,2-xylosyltransferase. <i>Plant Cell Physiol</i> 53(4), 755-761(2012)
85	M. Endo, S. Nakayama, C. Umeda-Hara, N. Ohtsuki, H. Saika, M. Umeda, S. Toki. CDKB2 is involved in mitosis and DNA damage response in rice. <i>Plant J</i> 69(6), 967-977(2012)
86	H. Takahashi, T. Kawakatsu, Y. Wakasa, S. Hayashi, F. Takaiwa. A Rice Transmembrane bZIP Transcription Factor, OsbZIP39, Regulates the Endoplasmic Reticulum Stress Response. <i>Plant Cell Physiol</i> 53(1), 144-153(2012)

87	S. Hayashi, Y. Wakasa, H. Takahashi, T. Kawakatsu, F. Takaiwa. Signal transduction by IRE1-mediated splicing of bZIP50 and other stress sensors in the endoplasmic reticulum stress response of rice. <i>Plant J</i> 69(6), 946–956(2012)
88	Y. Wakasa, K. Hirano, A. Urisu, T. Matsuda, F. Takaiwa. Generation of Transgenic Rice Lines with Reduced Contents of Multiple Potential Allergens Using a Null Mutant in Combination with an RNA Silencing Method. <i>Plant Cell Physiol</i> 52(12), 2190–2199(2011)
89	H. Saika, A. Oikawa, F. Matsuda, H. Onodera, K. Saito, S. Toki. Application of gene targeting to designed mutation breeding of high-tryptophan rice. <i>Plant Physiol</i> 156(3), 1269–1277(2011)
90	Y. Wakasa, C. Tamakoshi, T. Ohno, S. Hirose, T. Goto, S. Nagaoka, F. Takaiwa. The hypocholesterolemic activity of transgenic rice seed accumulating lactostatin, a bioactive peptide derived from bovine milk beta-lactoglobulin. <i>J Agric Food Chem</i> 59(8), 3845–3850(2011)
91	Saika H, Sakamoto W, Maekawa M, Toki S. Highly efficient visual selection of transgenic rice plants using green fluorescent protein or anthocyanin synthetic genes. <i>Plant biotechnology</i> 28(1), 107–110 (2011)
92	H. Takagi, T. Hiroi, S. Hirose, L. Yang, F. Takaiwa. Rice seed ER-derived protein body as an efficient delivery vehicle for oral tolerogenic peptides. <i>Peptides</i> 31(8), 1421–1425(2010)
93	N. Mae, Y. Makino, S. Oshita, Y. Kawagoe, A. Tanaka, T. Akihiro, K. Akama, S. Koike, C. Matsukura, H. Ezura. Stimulation of γ -aminobutyric acid production in vine-ripened tomato (<i>Solanum lycopersicum</i> L.) fruits under an adjusted aerobic atmosphere. <i>Journal of Packaging Science and Technology, Japan</i> 19(5), 375–381(2010)
94	Y. Yuki, D. Tokuhara, T. Nochi, H. Yasuda, M. Mejima, S. Kurokawa, Y. Takahashi, N. Kataoka, U. Nakanishi, Y. Hagiwara, K. Fujihashi, F. Takaiwa, H. Kiyono. Oral MucoRice expressing double-mutant cholera toxin A and B subunits induces toxin-specific neutralising immunity. <i>Vaccine</i> 27(43), 5982–5988(2009)
95	T. Motoyama, N. Maruyama, Y. Amari, K. Kobayashi, H. Washida, T. Higasa, F. Takaiwa, S. Utsumi. α ' Subunit of soybean β -conglycinin forms complex with rice glutelin via a disulphide bond in transgenic rice seeds. <i>J Exp Bot</i> 60(14), 4015–4027(2009)
96	T. Nozoye, F. Takaiwa, N. Tsuji, T. Yamakawa, T. Arakawa, Y. Hayashi, Y. Matsumoto. Production of <i>Ascaris suum</i> As14 protein and its fusion protein with cholera toxin B subunit in rice seeds. <i>J Vet Med Sci</i> 71(7), 995–1000(2009)
97	I. Ikeda, M. Kudo, T. Hamada, K. Nagao, Y. Oshiro, M. Kato, T. Sugawara, T. Yamahira, H. Ito, S. Tamaru, M. Sato, K. Imaizumi, S. Nagaoka, T. Yanagita. Dietary soy protein isolate and its undigested high molecular fraction upregulate hepatic ATP-binding cassette transporter G5 and ATP-binding cassette transporter G8 mRNA and increase biliary secretion of cholesterol in rats. <i>J Nutr Sci Vitaminol (Tokyo)</i> 55(3), 252–256(2009)
98	H. Yasuda, S. Hirose, T. Kawakatsu, Y. Wakasa, F. Takaiwa. Overexpression of BiP has inhibitory effects on the accumulation of seed storage proteins in endosperm cells of rice. <i>Plant Cell Physiol</i> 50(8), 1532–1543(2009)
99	L. Yang, Y. Wakasa, T. Kawakatsu, F. Takaiwa. The 3'-untranslated region of rice glutelin GluB-1 affects accumulation of heterologous protein in transgenic rice. <i>Biotechnol Lett</i> 31(10), 1625–1631(2009)
100	Y. Wakasa, L. Yang, S. Hirose, F. Takaiwa. Expression of unprocessed glutelin precursor alters polymerization without affecting trafficking and accumulation. <i>J Exp Bot</i> 60(12), 3503–3511(2009)
101	K. Suzuki, O. Kaminuma, L. Yang, Y. Motoi, T. Takai, S. Ichikawa, K. Okumura, H. Ogawa, A. Mori, F. Takaiwa, T. Hiroi. Development of transgenic rice expressing mite antigen for a new concept of immunotherapy. <i>Int Arch Allergy Immunol</i> 149 Suppl 1, 21–24(2009)
102	T. Kawakatsu, M. P. Yamamoto, S. M. Touno, H. Yasuda, F. Takaiwa. Compensation and interaction between RISBZ1 and RPBF during grain filling in rice. <i>Plant J</i> 59(6), 908–920(2009)
103	E. Domon, H. Takagi, S. Hirose, K. Sugita, S. Kasahara, H. Ebinuma, F. Takaiwa. 26-Week oral safety study in macaques for transgenic rice containing major human T-cell epitope peptides from Japanese cedar pollen allergens. <i>J Agric Food Chem</i> 57(12), 5633–5638(2009)

104	K. Akama, J. Kanetou, S. Shimosaki, K. Kawakami, S. Tsuchikura, F. Takaiwa. Seed-specific expression of truncated OsGAD2 produces GABA-enriched rice grains that influence a decrease in blood pressure in spontaneously hypertensive rats. <i>Transgenic Res</i> 18(6), 865-876(2009)
105	F. Takaiwa, S. Hirose, H. Takagi, L. Yang, Y. Wakasa. Deposition of a recombinant peptide in ER-derived protein bodies by retention with cysteine-rich prolamins in transgenic rice seed. <i>Planta</i> 229(5), 1147-1158(2009)
106	Y. Wakasa, K. Ozawa, F. Takaiwa. Higher-level accumulation of foreign gene products in transgenic rice seeds by the callus-specific selection system. <i>J Biosci Bioeng</i> 107(1), 78-83(2009)
107	Y. Matsumoto, S. Suzuki, T. Nozoye, T. Yamakawa, Y. Takashima, T. Arakawa, N. Tsuji, F. Takaiwa, Y. Hayashi. Oral immunogenicity and protective efficacy in mice of transgenic rice plants producing a vaccine candidate antigen (As16) of <i>Ascaris suum</i> fused with cholera toxin B subunit. <i>Transgenic Res</i> 18(2), 185-192(2009)
108	Takaiwa F. Allergen-specific immunotherapy with plant-based oral vaccines. <i>Immunotherapy</i> 1(4), 517-519(2009)
109	H. Takagi, T. Hiroi, L. Yang, K. Takamura, R. Ishimitsu, H. Kawauchi, F. Takaiwa. Efficient induction of oral tolerance by fusing cholera toxin B subunit with allergen-specific T-cell epitopes accumulated in rice seed. <i>Vaccine</i> 26(48), 6027-6030(2008)
110	F. Takaiwa, C. Sakuta, S. K. Choi, Y. Tada, T. Motoyama, S. Utsumi. Co-expression of soybean glycinins A1aB1b and A3B4 enhances their accumulation levels in transgenic rice seed. <i>Plant Cell Physiol</i> 49(10), 1589-1599(2008)
111	T. Akihiro, S. Koike, R. Tani, T. Tominaga, S. Watanabe, Y. Iijima, K. Aoki, D. Shibata, H. Ashihara, C. Matsukura, K. Akama, T. Fujimura, H. Ezura. Biochemical mechanism on GABA accumulation during fruit development in tomato. <i>Plant Cell Physiol</i> 49(9), 1378-1389(2008)
112	Y. Johzuka-Hisatomi, R. Terada, S. Iida. Efficient transfer of base changes from a vector to the rice genome by homologous recombination: involvement of heteroduplex formation and mismatch correction. <i>Nucleic Acids Res</i> 36(14), 4727-4735(2008)
113	K. Suzuki, O. Kaminuma, T. Hiroi, F. Kitamura, S. Miyatake, F. Takaiwa, H. Tatsumi, S. Nemoto, N. Kitamura, A. Mori. Downregulation of IL-13 gene transcription by T-bet in human T cells. <i>Int Arch Allergy Immunol</i> 146 Suppl 1, 33-35(2008)
114	Takashi Akihiro, Tatsuhito Fujimura, Hiroshi Ezura, Kazuhito Akama. Oligo microarray analysis of transcription in γ -aminobutyric acid-overaccumulating rice calli. <i>Rice Genet. Newsl.</i> 24, 55-58(2008)
115	赤間一仁, 金藤純子, 道田真帆子, 下崎俊介, 川上浩平. GABA強化米の開発と高血圧自然発症ラット (SHR) を用いた経口投与試験. <i>アミノ酸研究</i> 2(2), 143-146(2008)
116	M. Kuroda, M. Kimizu, C. Mikami. A simple set of plasmids for the production of transgenic plants. <i>Biosci Biotechnol Biochem</i> 74(11), 2348-2351(2010)
117	T. Kawakatsu, S. Wang, Y. Wakasa, F. Takaiwa. Increased lysine content in rice grains by over-accumulation of BiP in the endosperm. <i>Biosci Biotechnol Biochem</i> 74(12), 2529-2531(2010)
118	T. Nochi, Y. Yuki, Y. Katakai, H. Shibata, D. Tokuhara, M. Mejima, S. Kurokawa, Y. Takahashi, U. Nakanishi, F. Ono, H. Mimuro, C. Sasakawa, F. Takaiwa, K. Terao, H. Kiyono. A rice-based oral cholera vaccine induces macaque-specific systemic neutralizing antibodies but does not influence pre-existing intestinal immunity. <i>J Immunol</i> 183(10), 6538-6544(2009)
119	O. Kaminuma, F. Kitamura, N. Kitamura, T. Hiroi, H. Miyoshi, A. Miyawaki, S. Miyatake. Differential contribution of NFATc2 and NFATc1 to TNF- α gene expression in T cells. <i>J Immunol</i> 180(1), 319-326(2008)
120	K. Ohinata, Y. Fujiwata, F. Shingo, I. Masaru, H. Masatsugu, M. Yoshikawa. Orally administered novokinin, an angiotensin AT2 receptor agonist, suppresses food intake via prostaglandin E2-dependent mechanism in mice. <i>Peptides</i> 30(6), 1105-1108(2009)
121	T. Motoyama, Y. Amari, M. R. Tandang-Silvas, C. Cabanos, A. Kimura, M. Yoshikawa, F. Takaiwa, S. Utsumi, N. Maruyama. Development of transgenic rice containing a mutated beta subunit of soybean beta-conglycinin for enhanced phagocytosis-stimulating activity. <i>Peptides</i> 31(7), 1245-1250(2010)

122	Y. Yamada, K. Ohinata, A. W. Lipkowski, M. Yoshikawa. Angiotensin AT(2) receptor agonists act as anti-opioids via EP(3) receptor in mice. <i>Peptides</i> 30(4), 735–739(2009)
123	Y. Wakasa, H. Zhao, S. Hirose, D. Yamauchi, Y. Yamada, L. Yang, K. Ohinata, M. Yoshikawa, F. Takaiwa. Antihypertensive activity of transgenic rice seed containing an 18-repeat novokinin peptide localized in the nucleolus of endosperm cells. <i>Plant Biotechnol J</i> 9(7), 729–735(2011)
124	Y. Oono, Y. Wakasa, S. Hirose, L. Yang, C. Sakuta, F. Takaiwa. Analysis of ER stress in developing rice endosperm accumulating beta-amyloid peptide. <i>Plant Biotechnol J</i> 8(6), 691–718(2010)
125	T. Kawakatsu, F. Takaiwa. Differences in Transcriptional Regulatory Mechanisms Functioning for Free Lysine Content and Seed Storage Protein Accumulation in Rice Grain. <i>Plant Cell Physiol</i> 51(12), 1964–1974(2010)
126	H. Saika, S. Toki. Mature seed-derived callus of the model indica rice variety Kasalath is highly competent in Agrobacterium-mediated transformation. <i>Plant Cell Rep</i> 29(12), 1351–1364(2010)
127	Y. Wakasa, H. Yasuda, Y. Oono, T. Kawakatsu, S. Hirose, H. Takahashi, S. Hayashi, L. Yang, F. Takaiwa. Expression of ER quality control-related genes in response to changes in BiP1 levels in developing rice endosperm. <i>Plant J</i> 65(5), 675–689(2011)
128	T. Kawakatsu, S. Hirose, H. Yasuda, F. Takaiwa. Reducing Rice Seed Storage Protein Accumulation Leads to Changes in Nutrient Quality and Storage Organelle Formation. <i>Plant Physiol</i> 154(4), 1842–1854(2010)
129	D. Tokuhara, Y. Yuki, T. Nochi, T. Kodama, M. Mejima, S. Kurokawa, Y. Takahashi, M. Nanno, U. Nakanishi, F. Takaiwa, T. Honda, H. Kiyono. Secretory IgA-mediated protection against V. cholerae and heat-labile enterotoxin-producing enterotoxigenic Escherichia coli by rice-based vaccine. <i>Proc Natl Acad Sci U S A</i> 107(19), 8794–8799(2010)
130	Y. Fujiwara, Y. Aiki, L. Yang, F. Takaiwa, A. Kosaka, N. M. Tsuji, K. Shiraki, K. Sekikawa. Extraction and purification of human interleukin-10 from transgenic rice seeds. <i>Protein Expr Purif</i> 72(1), 125–130(2010)
131	M. S. Hanafy, S. M. Rahman, Y. Nakamoto, T. Fujiwara, S. Naito, K. Wakasa, M. Ishimoto. Differential response of methionine metabolism in two grain legumes, soybean and azuki bean, expressing a mutated form of Arabidopsis cystathionine gamma-synthase. <i>J Plant Physiol</i> 170(3), 338–345(2013)
132	M. Ishimoto, M. Kuroda, K. Yoza, K. Nishizawa, M. Teraishi, N. Mizutani, K. Ito, S. Moriya. Heterologous expression of corn cystatin in soybean and effect on growth of the stink bug. <i>Biosci Biotechnol Biochem</i> 76(11), 2142–2145(2012)
133	S. Nakamura, F. Abe, H. Kawahigashi, K. Nakazono, A. Tagiri, T. Matsumoto, S. Utsugi, T. Ogawa, H. Handa, H. Ishida, M. Mori, K. Kawaura, Y. Ogihara, H. Miura. A wheat homolog of MOTHER OF FT AND TFL1 acts in the regulation of germination. <i>Plant Cell</i> 23(9), 3215–3229(2011)
134	K. Takagi, K. Nishizawa, A. Hirose, A. Kita, M. Ishimoto. Manipulation of saponin biosynthesis by RNA interference-mediated silencing of beta-amylin synthase gene expression in soybean. <i>Plant Cell Rep</i> 30(10), 1835–1846(2011)
135	Keito Nishizawa, Kyoko Takagi, Masayoshi Teraishi, Akiko Kita, Masao Ishimoto. Application of somatic embryos to rapid and reliable analysis of soybean seed components by RNA interference-mediated gene silencing. <i>Plant Biotechnology</i> 27(5), 409–420(2010)
136	Nakashima, K., Tran, L.-S. P., Nguyen, D.V., Fujita, M., Maruyama, K., Todaka, D., Ito, Y., Hayashi, N., Shinozaki, K., Yamaguchi-Shinozaki, K. (2007) Functional analysis of a NAC-type transcription factor OsNAC6 involved in abiotic and biotic stress-responsive gene expression in rice. <i>Plant J.</i> 51: 617-630.
137	Yamaguchi-Shinozaki, K., Shinozaki, K. (2008) DREB regulons in abiotic-stress-responsive gene expression in plants. In “Molecular Breeding of Forage and Turf”, Yamada T., Spangenberg, G. (eds.) Springer, 15-27.
138	Qin, F., Sakuma, Y., Tran, L.-S. P., Maruyama, K., Kidokoro, S., Fujita, Y., Fujita, M., Umezawa, T., Sawano, Y., Miyazono, K., Tanokura, M., Shinozaki, K., Yamaguchi-Shinozaki, K. (2008) Arabidopsis DREB2A-interacting proteins function as RING E3 ligases and negatively regulate plant drought stress-responsive gene expression. <i>Plant Cell</i> 20: 1693-1707.

139	Yoshida, T., Sakuma, Y., Todaka, D., Maruyama, K., Qin, F., Mizoi, J., Kidokoro, S., Fujita, Y., Shinozaki, K., Yamaguchi-Shinozaki, K. (2008) Functional analysis of an <i>Arabidopsis</i> heat shock transcription factor <i>HsfA3</i> in the transcriptional cascade downstream of the DREB2A stress regulatory system. <i>Biochem. Biophys. Res. Commun.</i> 368: 515-521.
140	Nakashima, K., Ito, Y., Yamaguchi-Shinozaki, K. (2009) Transcriptional regulatory networks in response to abiotic stresses in <i>Arabidopsis</i> and grasses. <i>Plant Physiol.</i> 149: 88-95.
141	Nakashima, K., Yamaguchi-Shinozaki, K. (2009) Promoters and transcription factors in abiotic stress-responsive gene expression. Pareek, A., Sopory, S.K., Bohnert, H.J., Govindjee (eds.) <i>Abiotic stress adaptation in plants</i> . Springer-Verlag Berlin, 199-216.
142	Nakashima, K., Fujita, Y., Kanamori, N., Katagiri, T., Umezawa, T., Kidokoro, S., Maruyama, K., Yoshida, T., Ishiyama, K., Kobayashi, M., Shinozaki, K., Yamaguchi-Shinozaki, K. (2009) Three <i>Arabidopsis</i> SnRK2 protein kinases, SRK2D/SnRK2.2, SRK2E/SnRK2.6/OST1 and SRK2I/SnRK2.3 involved in ABA-signaling are essential for the control of seed development and dormancy. <i>Plant Cell Physiol.</i> 50: 1345-1363.
143	Maruyama, K., Takeda, M., Kidokoro, S., Yamada, K., Sakuma, Y., Urano, K., Fujita, M., Yoshiwara, K., Matsukura, S., Morishita, Y., Sasaki, R., Suzuki, H., Saito, K., Shibata, D., Shinozaki, K., Yamaguchi-Shinozaki, K. (2009) Metabolic pathways involved in cold acclimation identified by integrated analysis of metabolites and transcripts regulated by DREB1A and DREB2A. <i>Plant Physiol.</i> 150: 1972-1980.
144	Fujita, Y., Fujita, M., Yamaguchi-Shinozaki, K., Shinozaki, K. (2009) Transcription factors involved in the crosstalk between abiotic and biotic stress-signaling networks. Yoshioka, K., Shinozaki, K. (eds.) <i>Signal crosstalk in plant stress responses</i> , Wiley-Blackwell, 43-58.
145	Kidokoro, S., Maruyama, K., Nakashima, K., Imura, Y., Narusaka, Y., Shinwari, Z.K., Osakabe, Y., Fujita, Y., Mizoi, J., Shinozaki, K., Yamaguchi-Shinozaki, K. (2009) The phytochrome-interacting factor PIF7 negatively regulates <i>DREB1</i> expression under circadian control in <i>Arabidopsis</i> . <i>Plant Physiol.</i> 151: 2046-2057.
146	Miyazono, K., Miyakawa, T., Sawano, Y., Kubota, K., Kang, H.-J., Asano, A., Miyauchi, Y., Takahashi, M., Zhi, Y., Fujita, Y., Yoshida, T., Kodaira, K., Yamaguchi-Shinozaki, K., Tanokura, M. (2009) Structural basis of abscisic acid signaling. <i>Nature</i> 462: 609-614.
147	Fujita, Y., Nakashima, K., Yoshida, T., Katagiri, T., Kidokoro, S., Kanamori, N., Umezawa, T., Fujita, M., Maruyama, K., Ishiyama, K., Kobayashi, M., Nakasone, S., Yamada, K., Ito, T., Shinozaki, K., Yamaguchi-Shinozaki, K. (2009) Three SnRK2 protein kinases are the main positive regulators of abscisic acid signaling in response to water stress in <i>Arabidopsis</i> . <i>Plant Cell Physiol.</i> 50: 2123-2132.
148	Yoshida, T., Fujita, Y., Sayama, H., Kidokoro, S., Maruyama, K., Mizoi, J., Shinozaki, K., Yamaguchi-Shinozaki, K. (2010) AREB1, AREB2, and ABF3 are master transcription factors that cooperatively regulate ABRE-dependent ABA signaling involved in drought stress tolerance and require ABA for full activation. <i>Plant J.</i> 61: 672-685.
149	Matsukura, S., Mizoi, J., Yoshida, T., Todaka, D., Ito, Y., Maruyama, K., Shinozaki, K., Yamaguchi-Shinozaki, K. (2010) Comprehensive analysis of rice <i>DREB2</i> -type genes that encode transcription factors involved in the expression of abiotic stress-responsive genes. <i>Mol. Genet. Genomics.</i> 283: 185-196.
150	Mizoguchi, M., Umezawa, T., Nakashima, K., Kidokoro, S., Takasaki, H., Fujita, Y., Yamaguchi-Shinozaki, K., Shinozaki, K. (2010) Two closely related subclass II SnRK2 protein kinases cooperatively regulate drought-inducible gene expression. <i>Plant Cell Physiol.</i> 51: 842-847.
151	Takasaki, H., Maruyama, K., Kidokoro, S., Ito, Y., Fujita, Y., Shinozaki, K., Yamaguchi-Shinozaki, K., Nakashima, K. (2010) The abiotic stress-responsive NAC-type transcription factor OsNAC5 regulates stress-inducible genes and stress tolerance in rice. <i>Mol. Genet. Genomics</i> 284: 173-183.

152	Yamamoto, Y.Y., Yoshioka, Y., Hyakumachi, M., Maruyama, K., Yamaguchi-Shinozaki, K., Tokizawa, M., Koyama, H. (2011) Prediction of transcriptional regulatory elements for plant hormone responses based on microarray data. <i>BMC Plant Biol.</i> 11: 39.
153	Fujita, Y., Fujita, M., Shinozaki, K., Yamaguchi-Shinozaki, K. (2011) ABA-mediated transcriptional regulation in response to osmotic stress in plants. <i>J. Plant Res.</i> 124: 509-525.
154	Kodaira, K., Qin, F., Tran, L.-S.P., Maruyama, K., Kidokoro, S., Fujita, Y., Shinozaki, K., Yamaguchi-Shinozaki, K. (2011) Arabidopsis C2H2 zinc-finger proteins AZF1 and AZF2 negatively regulate ABA-repressive and auxin-inducible genes under abiotic stress conditions. <i>Plant Physiol.</i> 157: 742-756.
155	Qin, F., Shinozaki, K., Yamaguchi-Shinozaki, K. (2011) Achievements and challenges in understanding plant abiotic stress responses and tolerance. <i>Plant Cell Physiol.</i> 52: 1569-1582.
156	Yoshida, T., Ohama, N., Nakajima, J., Kidokoro, S., Mizoi, J., Nakashima, K., Maruyama, K., Kim, J.-M., Seki, M., Todaka, D., Osakabe, Y., Sakuma, Y., Schöffl, F., Shinozaki, K., Yamaguchi-Shinozaki, K. (2011) Arabidopsis HsfA1 transcription factors function as the main positive regulators in heat shock-responsive gene expression. <i>Mol. Genet. Genomics</i> 286: 321-332.
157	Kim, J.-S., Mizoi, J., Yoshida, T., Fujita, Y., Nakajima, J., Ohori, T., Todaka, D., Nakashima, K., Hirayama, T., Shinozaki, K., Yamaguchi-Shinozaki, K. (2011) An ABRE promoter sequence is involved in osmotic stress-responsive expression of the <i>DREB2A</i> gene, which encodes a transcription factor regulating drought-inducible genes in Arabidopsis. <i>Plant Cell Physiol.</i> 52: 2136-2146.
158	Qin, F., Kodaira, K., Maruyama, K., Mizoi, J., Tran, L.-S.P., Fujita, Y., Morimoto, K., Shinozaki, K., Yamaguchi-Shinozaki, K. (2011) <i>SPINDLY</i> , a negative regulator of GA signaling, is involved in the plant abiotic stress response. <i>Plant Physiol.</i> 157: 1900-1913.
159	Yamada, K., Kanai, M., Osakabe, Y., Ohiraki, H., Shinozaki, K., Yamaguchi-Shinozaki, K. (2011) The monosaccharide absorption activity of Arabidopsis roots depends on the expression profiles of transporter genes under high salinity conditions. <i>J. Biol. Chem.</i> 286: 43577-43586.
160	Mizoi, J., Shinozaki, K., Yamaguchi-Shinozaki, K. (2012) AP2/ERF family transcription factors in plant abiotic stress responses. <i>BBA - Gene Regul. Mech.</i> 1819: 86-96.
161	Nakashima, K., Takasaki, H., Mizoi, J., Shinozaki, K., Yamaguchi-Shinozaki, K. (2012) NAC transcription factors in plant abiotic stress responses. <i>BBA - Gene Regul. Mech.</i> 1819: 97-103.
162	Maruyama, K., Todaka, D., Mizoi, J., Yoshida, T., Kidokoro, S., Matsukura, S., Takasaki, H., Sakurai, T., Yamamoto, Y.Y., Yoshiwara, K., Kojima, M., Sakakibara, H., Shinozaki, K., Yamaguchi-Shinozaki, K. (2012) Identification of cis-acting promoter elements in cold- and dehydration-induced transcriptional pathways in Arabidopsis, rice and soybean. <i>DNA Res.</i> 9: 37-49.
163	Saint Pierre, C., Crossa, J., Bonnett, D., Yamaguchi-Shinozaki, K., Reynolds, M. (2012) Phenotyping transgenic wheat for drought resistance. <i>J. Exp. Bot.</i> 63: 1799-1808.
164	Todaka, D., Nakashima, K., Shinozaki, K., Yamaguchi-Shinozaki, K. (2012) Toward understanding transcriptional regulatory networks in abiotic stress responses and tolerance in rice. <i>Rice</i> 5:6.
165	Tanaka, H., Osakabe, Y., Katsura, S., Mizuno, S., Maruyama, K., Kusakabe, K., Mizoi, J., Shinozaki, K., Yamaguchi-Shinozaki, K. (2012) Abiotic stress-inducible receptor-like kinases negatively control ABA signaling in Arabidopsis. <i>Plant J.</i> 70(4): 599-613.
166	Miyazono, K., Koura, T., Kubota, K., Yoshida, T., Fujita, Y., Yamaguchi-Shinozaki, K., Tanokura, M. (2012) Purification, crystallization and preliminary X-ray analysis of OsAREB8 from rice, a member of the AREB/ABF family of bZIP transcription factors, in complex with its cognate DNA. <i>Acta Cryst. Sect. F</i> 68: 491-494.

167	Kim, J.-S., Mizoi, J., Kidokoro, S., Maruyama, K., Nakajima, J., Nakashima, K., Mitsuda, N., Takiguchi, Y., Ohme-Takagi, M., Kondou, Y., Yoshizumi, T., Matsui, M., Shinozaki, K., Yamaguchi-Shinozaki, K. (2012) <i>Arabidopsis</i> GROWTH-REGULATING FACTOR 7 functions as a transcriptional repressor of ABA- and osmotic stress-responsive genes including <i>DREB2A</i> . <i>Plant Cell</i> 24: 3393-3405.
168	Todaka, D., Nakashima, K., Maruyama, K., Kidokoro, S., Osakabe, Y., Ito, Y., Matsukura, S., Fujita, Y., Yoshiwara, K., Ohme-Takagi, M., Kojima, M., Sakakibara, H., Shinozaki, K., Yamaguchi-Shinozaki, K. (2012) Rice phytochrome-interacting factor-like protein OsPIL1 functions as a key regulator of internode elongation and induces a morphological response to drought stress. <i>Proc. Natl. Acad. Sci. USA</i> . 109: 15947-15952.
169	Ishizaki, T., Maruyama, K., Obara, M., Fukutani, A., Yamaguchi-Shinozaki, K., Ito, Y., Kumashiro, T. (2012) Expression of <i>Arabidopsis DREB1C</i> improves survival, growth, and yield of upland New Rice for Africa (NERICA) under drought. <i>Mol. Breeding in press</i> . [10.1007/s11032-012-9785-9]
170	Fujita, Y., Yoshida, T., Yamaguchi-Shinozaki, K. (2013) Pivotal role of the AREB/ABF-SnRK2 pathway in ABRE-mediated transcription in response to osmotic stress in plants. <i>Physiologia Plantarum</i> 147: 15-27.
171	Miyakawa, T., Fujita, Y., Yamaguchi-Shinozaki, K., Tanokura, M. (2013) Structure and function of abscisic acid receptors. <i>Trends Plant Sci. in press</i> . [10.1016/j.tplants.2012.11.002]
172	Jan, A., Maruyama, K., Todaka, D., Kidokoro, S., Abo, M., Yoshimura, E., Shinozaki, K., Nakashima, K., Yamaguchi-Shinozaki, K. (2013) OsTZF1, a CCCH-tandem zinc finger protein, confers delayed senescence and stress tolerance in rice by regulating stress-related genes. <i>Plant Physiol. in press</i> .
173	Ishizaki T, Kumashiro T. Genetic transformation of NERICA, interspecific hybrid rice between <i>Oryza glaberrima</i> and <i>O. sativa</i> , mediated by <i>Agrobacterium tumefaciens</i> . <i>Plant Cell Reports</i> 27, 319-327 (2008)
174	Ishizaki T, Kumashiro T. Investigations of copy number of transgene, fertility and expression level of an introduced GUS gene in transgenic NERICA produced by <i>Agrobacterium</i> -mediated methods. <i>In Vitro Cellular and Developmental Biology - Plant</i> 47, 339-347 (2011)
175	Ishizaki T, Maruyama K, Obara M, Fukutani A, Yamaguchi-Shinozaki K, Ito Y, Kumashiro T. Expression of <i>Arabidopsis DREB1C</i> improves survival, growth, and yield of upland New Rice for Africa (NERICA) under drought. <i>Molecular Breeding</i> (DOI 10.1007/s11032-012-9785-9) (2012)
176	Shinozaki, K. and Yamaguchi-Shinozaki, K. (2007) Gene networks involved in drought stress response and tolerance. <i>J Exp Bot</i> , 58, 221-227.
177	Takahashi, F., Yoshida, R., Ichimura, K., Mizoguchi, T., Seo, S., Yonezawa, M., Maruyama, K., Yamaguchi-Shinozaki, K. and Shinozaki, K. (2007) The mitogen-activated protein kinase cascade MKK3-MPK6 is an important part of the jasmonate signal transduction pathway in <i>Arabidopsis</i> . <i>Plant Cell</i> . 19, 805-818.
178	Seki, M., Umezawa, T., Urano, K. and Shinozaki, K. (2007) Regulatory metabolic networks in drought stress responses. <i>Curr. Opin. Plant Biol.</i> , 10, 296-302.
179	Hirayama, T. and Shinozaki, K. (2007) Perception and transduction of abscisic acid signals: keys to the function of the versatile plant hormone ABA. <i>Trends Plant Sci.</i> , 12, 343-351.
180	Fujita, M., Mizukado, S., Fujita, Y., Ichikawa, T., Nakazawa, M., Seki, M., Matsui, M., Yamaguchi-Shinozaki, K. and Shinozaki, K. (2007) Identification of stress-tolerance-related transcription-factor genes via mini-scale Full-length cDNA Over-eXpressor (FOX) gene hunting system. <i>Biochem. Biophys. Res. Commun.</i> , 364, 250-257.
181	Mochida, K., Saisho, D., Yoshida, T., Sakurai, T. and Shinozaki, K. (2008) TriMEDB: a database to integrate transcribed markers and facilitate genetic studies of the tribe Triticeae. <i>BMC Plant Biol.</i> 8, 72.
182	Yasuda, M., Ishikawa, A., Jikumaru, Y., Seki, M., Umezawa, T., Asami, T., Maruyama-Nakashita, A., Kudo, T., Shinozaki, K., Yoshida, S. and Nakashita, H. (2008) Antagonistic interaction between systemic acquired resistance and the abscisic acid-mediated abiotic stress response in <i>Arabidopsis</i> . <i>Plant Cell</i> . 20, 1678-1692.

183	Urano, K., Maruyama, K., Ogata, Y., Morishita, Y., Takeda, M., Sakurai, N., Suzuki, H., Saito, K., Shibata, D., Kobayashi, M., Yamaguchi-Shinozaki, K. and Shinozaki, K. (2009) Characterization of the ABA-regulated global responses to dehydration in <i>Arabidopsis</i> by metabolomics. <i>Plant J.</i> 57, 1065-1078.
184	Takahashi, F., Ichimura, K., Shinozaki, K. and Shirasu, K. (2009) Plant Mitogen-activated protein kinase cascades in signaling crosstalk, Signal Crosstalk in Plant Stress Responses, edited by Keiko Yoshioka and Kazuo Shinozaki, Blackwell Publishing, 23-42.
185	Kawaura, K., Mochida, K., Enju, A., Totoki, Y., Toyoda, A., Sakaki, Y., Kai, C., Kawai, J., Hayashizaki, Y., Seki, M., Shinozaki, K. and Ogihara, Y. (2009) Assessment of adaptive evolution between wheat and rice as deduced from full-length common wheat cDNA sequence data and expression patterns, <i>BMC Genomics</i> , 10, 1-11.
186	Mochida, K., Yoshida, T., Sakurai, T., Ogihara, Y. and Shinozaki, K. (2009) TriFLDB: a database of clustered full-length coding sequences from Triticeae with applications to comparative grass genomics. <i>Plant Physiol.</i> 150, 1135-1146.
187	Shinozaki, K. and Sakakibara, H. (2009) Omics and bioinformatics: an essential toolbox for systems analyses of plant functions beyond 2010, <i>Plant Cell Physiol.</i> , 50, 1177-1180.
188	Umezawa, T., Sugiyama, N., Mizoguchi, M., Hayashi, S., Myouga, F., Yamaguchi-Shinozaki, K., Ishihama, Y., Hirayama, T. and Shinozaki, K. (2009) Type 2C protein phosphatases directly regulate abscisic acid-activated protein kinases in <i>Arabidopsis</i> . <i>Proc. Natl. Acad. Sci. U. S. A.</i> 106, 17588-17593.
189	Mochida, K., Yoshida, T., Sakurai, T., Yamaguchi-Shinozaki, K., Shinozaki, K. and Tran, L. S. (2010) LegumeTFDB: an integrative database of Glycine max, Lotus japonicus and <i>Medicago truncatula</i> transcription factors. <i>Bioinformatics</i> 26, 290-291.
190	Hirayama, T. and Shinozaki, K. (2010) Research on plant abiotic stress responses in the post-genome era: past, present, and future, <i>Plant J</i> , 61, 1041-1062
191	Urano, K., Kurihara, Y., Seki, K. and Shinozaki, K. (2010) 'Omics' analyses of regulatory networks in plant abiotic stress responses, <i>Curr. Opin. Plant Biol.</i> , 13, 132-138.
192	Hirayama, T. and Umezawa, T. (2010) The PP2C-SnRK2 complex: the central regulator of an abscisic acid signaling pathway, <i>Plant Signal Behav.</i> , 5, 160-163.
193	Mochida, K. and Shinozaki, K. (2010) Genomics and bioinformatics resources for crop improvement, <i>Plant Cell Physiol.</i> , 51, 497-523.
194	Mizoguchi, M., Umezawa, T., Nakashima, K., Kidokoro, S., Takasaki, H., Fujita, Y., Yamaguchi-Shinozaki, K. and Shinozaki, K. (2010) Two closely related subclass II SnRK2 protein kinases cooperatively regulate drought-inducible gene expression, <i>Plant Cell Physiol.</i> , 51, 842-847.
195	Shinozaki, K. (2010) The ICAR2010 for development of <i>Arabidopsis</i> research beyond 2010, <i>J Plant Res.</i> , 123, 265-266.
196	Mochida, K., Yoshida, T., Sakurai, T., Yamaguchi-Shinozaki, K., Shinozaki, K. and Tran, L. S. (2010) Genome-wide analysis of two-component systems and prediction of stress-responsive two-component system members in soybean. <i>DNA Res.</i> 17, 303-324.
197	Kuromori, T. and Shinozaki, K. (2010) ABA transport factors found in <i>Arabidopsis</i> ABC transporters. <i>Plant Signal Behav.</i> 5, 1124-1126
198	Betsuyaku, S., Takahashi, F., Kinoshita, A., Miwa, H., Shinozaki, K., Fukuda, H. and Sawa, S. (2010) Mitogen-Activated Protein Kinase Regulated by the CLAVATA Receptors Contributes to Shoot Apical Meristem Homeostasis. <i>Plant Cell Physiol.</i> 52, 14-29.
199	Umezawa, T., Nakashima, K., Miyakawa, T., Kuromori, T., Tanokura, M., Shinozaki, K. and Yamaguchi-Shinozaki, K. (2010) Molecular basis of the core regulatory network in aba responses: sensing, signaling and transport. <i>Plant Cell Physiol.</i> 51, 1821-1839.
200	Takahashi, F., Mizoguchi, T., Yoshida, R., Ichimura, K. and Shinozaki, K. (2011) Calmodulin-dependent activation of MAP kinase for ROS homeostasis in <i>Arabidopsis</i> . <i>Mol. Cell.</i> 41, 649-660.
201	Mochida, K., Uehara-Yamaguchi, Y., Yoshida, T., Sakurai, T. and Shinozaki, K. (2011) Global landscape of a co-expressed gene network in barley and its application to gene discovery in Triticeae crops. <i>Plant Cell Physiol.</i> 52, 785-803.
202	Mochida, K., Yoshida, T., Sakurai, T., Yamaguchi-Shinozaki, K., Shinozaki, K. and Tran, L. S. (2011) <i>In silico</i> analysis of transcription factor repertoires and prediction of stress-responsive transcription factors from six major gramineae plants. <i>DNA Res.</i> 18, 321-332.

203	Kuromori, T., Sugimoto, E. and Shinozaki, K. (2011) Arabidopsis mutants of <i>AtABCG22</i> , an ABC transporter gene, increase water transpiration and drought susceptibility. <i>Plant J.</i> 67, 885-894.
204	Kuromori, T., Ito, T., Sugimoto, E. and Shinozaki, K. (2011) <i>Arabidopsis</i> mutant of <i>AtABCG26</i> , an ABC transporter gene, is defective in pollen maturation. <i>J. Plant Physiol.</i> 168, 2001-2005.
205	Murayama, M., Hayashi, S., Nishimura, N., Ishide, M., Kobayashi, K., Yagi, Y., Asami, T., Nakamura, T., Shinozaki, K. and Hirayama, T. (2012) Isolation of <i>Arabidopsis ahg11</i> , a weak ABA hypersensitive mutant defective in <i>nad4</i> RNA editing. <i>J Exp Bot.</i> , 63, 5301-5310
206	Fujita, M., Fujita, Y., Iuchi, S., Yamada, K., Kobayashi, Y., Urano, K., Kobayashi, M., Yamaguchi-Shinozaki, K. and Shinozaki, K. (2012) Natural variation in a polyamine transporter determines paraquat tolerance in <i>Arabidopsis</i> . <i>Proc. Natl. Acad. Sci. U. S. A.</i> , 109, 6343-6347.
207	Osakabe, Y., Yamaguchi-Shinozaki, K., Shinozaki, K. and Phan Tran, L. S. (2013) Sensing the environment: key roles of membrane-localized kinases in plant perception and response to abiotic stress. <i>J Exp Bot.</i> , 64, 445-458.
208	Mochida, K., Yoshida, T., Sakurai, T., Yamaguchi-Shinozaki, K., Shinozaki, K. and Tran, L. S. (2013) TreeTFDB: An Integrative Database of the Transcription Factors from Six Economically Important Tree Crops for Functional Predictions and Comparative and Functional Genomics. <i>DNA Res.</i> in press.
209	Osakabe, Y., Arinaga, N., Umezawa, T., Katsura, S., Nagamachi, K., Tanaka, H., Ohiraki, H., Yamada, K., Seo, SU., Abo, M., Yoshimura, E., Shinozaki, K. and Yamaguchi-Shinozaki, K. (2013) Osmotic stress responses and plant growth controlled by potassium transporters in <i>Arabidopsis</i> , <i>Plant Cell.</i> in press.
210	Saint Pierre C, Wu H, Yamaguchi-Shinozaki K, Crouch J, Reynolds M. Drought screening and physiological characterization of wheat lines transformed with DREB1A gene driven by RD29A promoter. Joint Annual Meeting of the Geological Society of America, Soil Science Society of America, American Society of Agronomy, Crop Science Society of America, and Gulf Coast Association of Geological Societies. Houston, Texas, 5 –9 October (2008)
211	Saint Pierre C, Yamaguchi Shinozaki K, Reynolds MP. Evaluation of transformed wheat lines under water stress: Results from and open-field trial. In: International Wheat Conference, 8; Abstracts of oral and poster presentations; St. Petersburg (Russia): Vavilov Research Institute of Plant Industry, Abstract only p. 467 (2010)
212	Saint Pierre C, Bonnett D, Yamaguchi-Shinozaki K, Reynolds M. Phenotyping transgenic wheat for drought tolerance. In: GM Crops: From basic research to application; Harpenden (United Kingdom); 28-29 Jun 2011. Association of Applied Biologists, Series: Aspects of Applied Biology (2011)
213	Saint Pierre C, Crossa JL, Bonnett D, Yamaguchi-Shinozaki K, Reynolds MP. Phenotyping transgenic wheat for drought resistance. <i>Journal of Experimental Botany</i> ; doi: 10.1093/jxb/err385 (2012)
214	A.C.M. Gaudin, A. Henry, A. H. Sparks and I. H. Slamet-Loedin. Taking transgenic rice drought screening to the field <i>Journal of Exp Botany</i> 64(1): 109-117 (2012)
215	Shinoyama H, Ichikawa H, Saitoh-Nakashima M, Saito M, Aida R, Ezura H, Yamaguchi H, Mochizuki A, Nakase K, Nishibata Y, Nomura Y, Kamada H. Introduction of male sterility to GM chrysanthemum plants to prevent transgene flow. <i>Acta Horticulturae</i> 937(1), 337–346(2012)
216	M. Hakata, M. Kuroda, A. Ohsumi, T. Hirose, H. Nakamura, M. Muramatsu, H. Ichikawa, H. Yamakawa. Overexpression of a rice TIFY gene increases grain size through enhanced accumulation of carbohydrates in the stem. <i>Biosci Biotechnol Biochem</i> 76(11), 2129–2124(2012)
217	W. Tanaka, T. Toriba, Y. Ohmori, A. Yoshida, A. Kawai, T. Mayama-Tsuchida, H. Ichikawa, N. Mitsuda, M. Ohme-Takagi, H. Y. Hirano. The YABBY gene TONGARI-BOUSHI1 is involved in lateral organ development and maintenance of meristem organization in the rice spikelet. <i>Plant Cell</i> 24(1), 80–95(2012)

218	H. Nakagawa, A. Tanaka, T. Tanabata, M. Ohtake, S. Fujioka, H. Nakamura, H. Ichikawa, M. Mori. Short grain1 decreases organ elongation and brassinosteroid response in rice. <i>Plant Physiol</i> 158(3), 1208–1219(2012)
219	T. Asano, N. Hayashi, M. Kobayashi, N. Aoki, A. Miyao, I. Mitsuhashi, H. Ichikawa, S. Komatsu, H. Hirochika, S. Kikuchi, R. Ohsugi. A rice calcium-dependent protein kinase OsCPK12 oppositely modulates salt-stress tolerance and blast disease resistance. <i>Plant J</i> 69(1), 26–36(2012)
220	H. Tabuchi, Y. Zhang, S. Hattori, M. Omae, S. Shimizu-Sato, T. Oikawa, Q. Qian, M. Nishimura, H. Kitano, H. Xie, X. Fang, H. Yoshida, J. Kyojuka, F. Chen, Y. Sato. LAX PANICLE2 of Rice Encodes a Novel Nuclear Protein and Regulates the Formation of Axillary Meristems. <i>Plant Cell</i> 23(9), 3276–3287(2011)
221	D. Ogawa, K. Abe, A. Miyao, M. Kojima, H. Sakakibara, M. Mizutani, H. Morita, Y. Toda, T. Hobo, Y. Sato, T. Hattori, H. Hirochika, S. Takeda. RSS1 regulates the cell cycle and maintains meristematic activity under stress conditions in rice. <i>Nat Commun</i> 2, 278(2011)
222	R. Morita, M. Kusaba, S. Iida, H. Yamaguchi, T. Nishio, M. Nishimura. Molecular characterization of mutations induced by gamma irradiation in rice. <i>Genes Genet Syst</i> 84(5), 361–370(2010)
223	Asano T, Wakayama M, Aoki N, Komatsu S, Ichikawa H, Hirochika H, Ohsugi R. Overexpression of a calcium-dependent protein kinase gene enhances growth of rice under low-nitrogen conditions. <i>Plant Biotechnology</i> 27(4), 369–373(2010)
224	Hakata M, Nakamura H, Iida-Okada K, Miyao A, Kajikawa M, Imai-Toki N, Pang J, Amano K, Horikawa A, Tsuchida-Mayama T, Song J, Igarashi M, Hiroko K., Kitamoto HK, Ichikawa T, Matsui M, Kikuchi S, Nagamura Y, Hirochika H, Ichikawa H. Production and characterization of a large population of cDNA-overexpressing transgenic rice plants using Gateway-based full-length cDNA expression libraries. <i>Breed. Sci.</i> 60(5), 575–585(2010)
225	Hakata M, Nakamura H, Kajikawa M, Ichikawa H. Application of FTA Card technology for identification of transgenes in transgenic rice. <i>Rice Genet. Newsl.</i> 25, 92–93(2010)
226	H. Nakamura, M. Muramatsu, M. Hakata, O. Ueno, Y. Nagamura, H. Hirochika, M. Takano, H. Ichikawa. Ectopic Overexpression of The Transcription Factor OsGLK1 Induces Chloroplast Development in Non-Green Rice Cells. <i>Plant Cell Physiol</i> 50(11), 1933–1949(2009)
227	Yao S-G, Sonoda Y, Tsutsui T, Nakamura H, Ichikawa H, Ikeda A, Yamaguchi J. Promoter analysis of OsAMT1;2 and OsAMT1;3 implies their distinct roles in nitrogen utilization in rice. <i>Breed. Sci.</i> 58(3), 201–207(2008)
228	C. F. Huang, N. Yamaji, M. Nishimura, S. Tajima, J. F. Ma. A rice mutant sensitive to Al toxicity is defective in the specification of root outer cell layers. <i>Plant Cell Physiol</i> 50(5), 976–985(2009)
229	M. Kusaba, T. Maoka, R. Morita, S. Takaichi. A novel carotenoid derivative, lutein 3-acetate, accumulates in senescent leaves of rice. <i>Plant Cell Physiol</i> 50(8), 1573–1577(2009)
230	Y. Sato, R. Morita, S. Katsuma, M. Nishimura, A. Tanaka, M. Kusaba. Two short-chain dehydrogenase/reductases, NON-YELLOW COLORING 1 and NYC1-LIKE, are required for chlorophyll b and light-harvesting complex II degradation during senescence in rice. <i>Plant J</i> 57(1), 120–131(2009)
231	Y. Ohmori, T. Toriba, H. Nakamura, H. Ichikawa, H. Y. Hirano. Temporal and spatial regulation of DROOPING LEAF gene expression that promotes midrib formation in rice. <i>Plant J</i> 65(1), 77–86(2011)
232	R. Morita, Y. Sato, Y. Masuda, M. Nishimura, M. Kusaba. Defect in non-yellow coloring 3, an alpha/beta hydrolase-fold family protein, causes a stay-green phenotype during leaf senescence in rice. <i>Plant J</i> 59(6), 940–952(2009)
233	T. Asano, M. Hakata, H. Nakamura, N. Aoki, S. Komatsu, H. Ichikawa, H. Hirochika, R. Ohsugi. Functional characterisation of OsCPK21, a calcium-dependent protein kinase that confers salt tolerance in rice. <i>Plant Mol Biol</i> 75(1-2), 179–191(2011)
234	R. Morita, M. Kusaba, S. Iida, T. Nishio, M. Nishimura. Development of PCR markers to detect the glb1 and Lgc1 mutations for the production of low easy-to-digest protein rice varieties. <i>Theor Appl Genet</i> 119(1), 125–130(2009)

235	Y. Sato, N. Namiki, H. Takehisa, K. Kamatsuki, H. Minami, H. Ikawa, H. Ohyanagi, K. Sugimoto, J. I. Itoh, B. A. Antonio, Y. Nagamura. RiceFRIEND: a platform for retrieving coexpressed gene networks in rice. <i>Nucleic Acids Res</i> 41(D1), D1214–D1221(2013)
236	A. J. Nagano, Y. Sato, M. Mihara, B. A. Antonio, R. Motoyama, H. Itoh, Y. Nagamura, T. Izawa. Deciphering and Prediction of Transcriptome Dynamics under Fluctuating Field Conditions. <i>Cell</i> 151(6), 1358–1369(2012)
237	Y. Sato, H. Takehisa, K. Kamatsuki, H. Minami, N. Namiki, H. Ikawa, H. Ohyanagi, K. Sugimoto, B. A. Antonio, Y. Nagamura. RiceXPro Version 3.0: expanding the informatics resource for rice transcriptome. <i>Nucleic Acids Res</i> , (2012)
238	Y. Kawahara, Y. Oono, H. Kanamori, T. Matsumoto, T. Itoh, E. Minami. Simultaneous RNA–Seq Analysis of a Mixed Transcriptome of Rice and Blast Fungus Interaction. <i>PLoS One</i> 7(11), e49423(2012)
239	K. Kobayashi, N. Yasuno, Y. Sato, M. Yoda, R. Yamazaki, M. Kimizu, H. Yoshida, Y. Nagamura, J. Kyojuka. Inflorescence meristem identity in rice is specified by overlapping functions of three AP1/FUL–like MADS box genes and PAP2, a SEPALLATA MADS box gene. <i>Plant Cell</i> 24(5), 1848–1859(2012)
240	C. C. Yang, Y. Kawahara, H. Mizuno, J. Wu, T. Matsumoto, T. Itoh. Independent Domestication of Asian Rice Followed by Gene Flow from japonica to indica. <i>Mol Biol Evol</i> 29(5), 1471–1479(2012)
241	S. Uruguchi, T. Kamiya, T. Sakamoto, K. Kasai, Y. Sato, Y. Nagamura, A. Yoshida, J. Kyojuka, S. Ishikawa, T. Fujiwara. Low–affinity cation transporter (OsLCT1) regulates cadmium transport into rice grains. <i>Proc Natl Acad Sci U S A</i> 108(52), 20959–20964(2011)
242	H. Sakai, H. Mizuno, Y. Kawahara, H. Wakimoto, H. Ikawa, H. Kawahigashi, H. Kanamori, T. Matsumoto, T. Itoh, B. S. Gaut. Retrogenes in rice (<i>Oryza sativa</i> L. ssp. japonica) exhibit correlated expression with their source genes. <i>Genome Biol Evol</i> 3, 1357–1368(2011)
243	H. Takehisa, Y. Sato, M. Igarashi, T. Abiko, B. A. Antonio, K. Kamatsuki, H. Minami, N. Namiki, Y. Inukai, M. Nakazono, Y. Nagamura. Genome–wide transcriptome dissection of the rice root system: implications for developmental and physiological functions. <i>Plant J</i> 69(1), 126–140(2011)
244	Y. Ishimaru, Y. Kakei, H. Shimo, K. Bashir, Y. Sato, N. Uozumi, H. Nakanishi, N. K. Nishizawa. A rice phenolic efflux transporter is essential for solubilizing precipitated apoplasmic iron in the plant stele. <i>J Biol Chem</i> 286(28), 24649–24655(2011)
245	T. Izawa, M. Mihara, Y. Suzuki, M. Gupta, H. Itoh, A. J. Nagano, R. Motoyama, Y. Sawada, M. Yano, M. Y. Hirai, A. Makino, Y. Nagamura. Os–GIGANTEA confers robust diurnal rhythms on the global transcriptome of rice in the field. <i>Plant Cell</i> 23(5), 1741–1755(2011)
246	Y. Sato, B. Antonio, N. Namiki, R. Motoyama, K. Sugimoto, H. Takehisa, H. Minami, K. Kamatsuki, M. Kusaba, H. Hirochika, Y. Nagamura. Field transcriptome revealed critical developmental and physiological transitions involved in the expression of growth potential in japonica rice. <i>BMC Plant Biol</i> 11, 10(2011)
247	Mizuno H, Kawahara Y, Wu J, Katayose Y, Kanamori H, Ikawa H, Itoh T, Sasaki T, Matsumoto T. Asymmetric distribution of gene expression in the centromeric region of rice chromosome 5. <i>Frontiers in Plant Science</i> 2(jun), 1–12(2011)
248	Oono Y, Kawahara Y, Kanamori H, Mizuno H, Yamagata H, Yamamoto M, Hosokawa S, Ikawa H, Akahane I, Zhu Z, Wu J, Itoh T, Matsumoto T. mRNA–Seq reveals a comprehensive transcriptome profile of rice under phosphate stress. <i>Rice</i> 4(2), 50–65(2011)
249	M. Kubota, Y. Saito, T. Masumura, T. Kumagai, R. Watanabe, S. Fujimura, M. Kadowaki. Improvement in the in vivo digestibility of rice protein by alkali extraction is due to structural changes in prolamin/protein body–I particle. <i>Biosci Biotechnol Biochem</i> 74(3), 614–619(2010)
250	T. Sazuka, N. Kamiya, T. Nishimura, K. Ohmae, Y. Sato, K. Imamura, Y. Nagato, T. Koshiba, Y. Nagamura, M. Ashikari, H. Kitano, M. Matsuoka. A rice tryptophan deficient dwarf mutant, tdd1, contains a reduced level of indole acetic acid and develops abnormal flowers and organless embryos. <i>Plant J</i> 60(2), 227–241(2009)
251	Y. Saito, K. Kishida, K. Takata, H. Takahashi, T. Shimada, K. Tanaka, S. Morita, S. Satoh, T. Masumura. A green fluorescent protein fused to rice prolamin forms protein body–like structures in transgenic rice. <i>J Exp Bot</i> 60(2), 615–627(2009)
252	Y. Saito, T. Shigemitsu, K. Tanaka, S. Morita, S. Satoh, T. Masumura. Ultrastructure of mature protein body in the starchy endosperm of dry cereal grain. <i>Biosci Biotechnol Biochem</i> 74(7), 1485–1487(2010)

253	Y. Saito, N. Nakatsuka, T. Shigemitsu, K. Tanaka, S. Morita, S. Satoh, T. Masumura. Thin frozen film method for visualization of storage proteins in mature rice grains. <i>Biosci Biotechnol Biochem</i> 72(10), 2779–2781(2008)
254	H. Mizuno, Y. Kawahara, H. Sakai, H. Kanamori, H. Wakimoto, H. Yamagata, Y. Oono, J. Wu, H. Ikawa, T. Itoh, T. Matsumoto. Massive parallel sequencing of mRNA in identification of unannotated salinity stress-inducible transcripts in rice (<i>Oryza sativa</i> L.). <i>BMC Genomics</i> 11(1), 683(2010)
255	Y. Sato, B. A. Antonio, N. Namiki, H. Takehisa, H. Minami, K. Kamatsuki, K. Sugimoto, Y. Shimizu, H. Hirochika, Y. Nagamura. RiceXPro: a platform for monitoring gene expression in japonica rice grown under natural field conditions. <i>Nucleic Acids Res</i> 39(Database issue), D1141–1148(2011)
256	M. Mihara, T. Itoh, T. Izawa. In silico identification of short nucleotide sequences associated with gene expression of pollen development in rice. <i>Plant Cell Physiol</i> 49(10), 1451–1464(2008)
257	Philippe Leroy, Nicolas Guilhot, Hiroaki Sakai, Aurelien Bernard, Frederic Choulet, Sebastien Theil, Sebastien Reboux, Naoki Amano, Timothee Flutre, Celine Pelegriin, Hajime Ohyanagi, Michael Seidel, Franck Giacomoni, Mathieu Reichstadt, Michael Alaux, Emmanuelle Gicquello, Fabrice Legeai, Lorenzo Cerutti, Hisataka Numa, Tsuyoshi Tanaka, Klaus Mayer, Takeshi Itoh, Hadi Quesneville, Catherine Feuillet. TriAnnot: a versatile and high performance pipeline for the automated annotation of plant genomes. <i>Front. Plant Sci.</i> 3, 5(2012)
258	Yamamoto E, Yonemaru J, Yamamoto T, Yano M. Rice OGRO: The Overview of functionally characterized Genes in Rice online database Rice. <i>Rice</i> 5, 26(2012)
259	S. Y. Yang, M. Gronlund, I. Jakobsen, M. S. Grottemeyer, D. Rentsch, A. Miyao, H. Hirochika, C. S. Kumar, V. Sundaresan, N. Salamin, S. Catausan, N. Mattes, S. Heuer, U. Paszkowski. Nonredundant Regulation of Rice Arbuscular Mycorrhizal Symbiosis by Two Members of the PHOSPHATE TRANSPORTER1 Gene Family. <i>Plant Cell</i> 24(10), 4236–4251(2012)
260	F. Hirose, N. Inagaki, A. Hanada, S. Yamaguchi, Y. Kamiya, A. Miyao, H. Hirochika, M. Takano. Cryptochrome and phytochrome cooperatively but independently reduce active gibberellin content in rice seedlings under light irradiation. <i>Plant Cell Physiol</i> 53(9), 1570–1582(2012)
261	J. Yonemaru, T. Yamamoto, K. Ebana, E. Yamamoto, H. Nagasaki, T. Shibaya, M. Yano. Genome-Wide Haplotype Changes Produced by Artificial Selection during Modern Rice Breeding in Japan. <i>PLoS One</i> 7(3), e32982(2012)
262	C. C. Yang, Y. Kawahara, H. Mizuno, J. Wu, T. Matsumoto, T. Itoh. Independent Domestication of Asian Rice Followed by Gene Flow from japonica to indica. <i>Mol Biol Evol</i> 29(5), 1471–1479(2012)
263	H. Hamada, T. Kurusu, E. Okuma, H. Nokajima, M. Kiyoduka, T. Koyano, Y. Sugiyama, K. Okada, J. Koga, H. Saji, A. Miyao, H. Hirochika, H. Yamane, Y. Murata, K. Kuchitsu. Regulation of a proteinaceous elicitor-induced Ca ²⁺ influx and production of phytoalexins by a putative voltage-gated cation channel, OsTPC1, in cultured rice cells. <i>J Biol Chem</i> 287(13), 9931–9939(2012)
264	A. Miyao, M. Nakagome, T. Ohnuma, H. Yamagata, H. Kanamori, Y. Katayose, A. Takahashi, T. Matsumoto, H. Hirochika. Molecular spectrum of somaclonal variation in regenerated rice revealed by whole-genome sequencing. <i>Plant Cell Physiol</i> 53(1), 256–264(2012)
265	T. Asano, N. Hayashi, M. Kobayashi, N. Aoki, A. Miyao, I. Mitsuhashi, H. Ichikawa, S. Komatsu, H. Hirochika, S. Kikuchi, R. Ohsugi. A rice calcium-dependent protein kinase OsCPK12 oppositely modulates salt-stress tolerance and blast disease resistance. <i>Plant J</i> 69(1), 26–36(2011)
266	G. Chen, T. Komatsuda, J. F. Ma, C. Nawrath, M. Pourkheirandish, A. Tagiri, Y. G. Hu, M. Sameri, X. Li, X. Zhao, Y. Liu, C. Li, X. Ma, A. Wang, S. Nair, N. Wang, A. Miyao, S. Sakuma, N. Yamaji, X. Zheng, E. Nevo. An ATP-binding cassette subfamily G full transporter is essential for the retention of leaf water in both wild barley and rice. <i>Proc Natl Acad Sci U S A</i> 108(30), 12354–12359(2011)
267	D. Ogawa, K. Abe, A. Miyao, M. Kojima, H. Sakakibara, M. Mizutani, H. Morita, Y. Toda, T. Hobo, Y. Sato, T. Hattori, H. Hirochika, S. Takeda. RSS1 regulates the cell cycle and maintains meristematic activity under stress conditions in rice. <i>Nat Commun</i> 2, 278(2011)

268	K. F. Mayer, M. Martis, P. E. Hedley, H. Simkova, H. Liu, J. A. Morris, B. Steuernagel, S. Taudien, S. Roessner, H. Gundlach, M. Kubalaková, P. Suchanková, F. Murat, M. Felder, T. Nussbaumer, A. Graner, J. Salse, T. Endo, H. Sakai, T. Tanaka, T. Itoh, K. Sato, M. Platzer, T. Matsumoto, U. Scholz, J. Dolezel, R. Waugh, N. Stein. Unlocking the barley genome by chromosomal and comparative genomics. <i>Plant Cell</i> 23(4), 1249–1263(2011)
269	T. Matsumoto, T. Tanaka, H. Sakai, N. Amano, H. Kanamori, K. Kurita, A. Kikuta, K. Kamiya, M. Yamamoto, H. Ikawa, N. Fujii, K. Hori, T. Itoh, K. Sato. Comprehensive sequence analysis of 24,783 barley full-length cDNAs derived from 12 clone libraries. <i>Plant Physiol</i> 156(1), 20–28(2011)
270	H. Sakai, H. Ikawa, T. Tanaka, H. Numa, H. Minami, M. Fujisawa, M. Shibata, K. Kurita, A. Kikuta, M. Hamada, H. Kanamori, N. Namiki, J. Wu, T. Itoh, T. Matsumoto, T. Sasaki. Distinct evolutionary patterns of <i>Oryza glaberrima</i> deciphered by genome sequencing and comparative analysis. <i>Plant J</i> 66(5), 796–805(2011)
271	C. C. Yang, H. Sakai, H. Numa, T. Itoh. Gene tree discordance of wild and cultivated Asian rice deciphered by genome-wide sequence comparison. <i>Gene</i> 477(1–2), 53–60(2011)
272	K. Nonomura, M. Eiguchi, M. Nakano, K. Takashima, N. Komeda, S. Fukuchi, S. Miyazaki, A. Miyao, H. Hirochika, N. Kurata. A novel RNA-recognition-motif protein is required for premeiotic G1/S-phase transition in rice (<i>Oryza sativa</i> L.). <i>PLoS Genet</i> 7(1), e1001265(2011)
273	Y. Nagamura, B. A. Antonio, Y. Sato, A. Miyao, N. Namiki, J. Yonemaru, H. Minami, K. Kamatsuki, K. Shimura, Y. Shimizu, H. Hirochika. Rice TOGO Browser: A platform to retrieve integrated information on rice functional and applied genomics. <i>Plant Cell Physiol</i> 52(2), 230–237(2011)
274	R. Babu, C. J. Jiang, X. Xu, K. R. Kottapalli, H. Takatsuji, A. Miyao, H. Hirochika, S. Kawasaki. Isolation, fine mapping and expression profiling of a lesion mimic genotype, <i>spl</i> (NF4050-8) that confers blast resistance in rice. <i>Theor Appl Genet</i> 122(4), 831–854(2011)
275	T. Yamamoto, H. Nagasaki, J. Yonemaru, K. Ebana, M. Nakajima, T. Shibaya, M. Yano. Fine definition of the pedigree haplotypes of closely related rice cultivars by means of genome-wide discovery of single-nucleotide polymorphisms. <i>BMC Genomics</i> 11, 267(2010)
276	H. Sakai, T. Itoh. Massive gene losses in Asian cultivated rice unveiled by comparative genome analysis. <i>BMC Genomics</i> 11, 121(2010)
277	M. Mihara, T. Itoh, T. Izawa. SALAD database: a motif-based database of protein annotations for plant comparative genomics. <i>Nucleic Acids Res</i> 38(Database issue), D835–842(2010)
278	Hiroshi Mizuno, Tsuyoshi Tanaka, Hiroaki Sakai, Hiroyuki Kawahigashi, Takeshi Itoh, Shoshi Kikuchi, Takashi Matsumoto. Characterization of 2159 Unmapped Full-length cDNA Sequences of <i>Oryza sativa</i> L. ssp. <i>japonica</i> 'Nipponbare'. <i>Plant Mol. Biol. Rep.</i> 28(2), 357–362(2010)
279	Yonemaru J, Yamamoto T, Fukuoka S, Uga Y, Hori K, Yano M. Q-TARO: QTL Annotation Rice Online Database. <i>Rice</i> 3(2–3), 194–203(2010)
280	Ebana K, Yonemaru J, Fukuoka S, Iwata H, Kanamori H, Namiki N, Nagasaki H, Yano M. Genetic structure revealed by a whole-genome single-nucleotide polymorphism survey of diverse accessions of cultivated Asian rice (<i>Oryza sativa</i> L.). <i>Breed. Sci.</i> 60(4), 390–397(2010)
281	Nagasaki H, Ebana K, Shibaya T, Yonemaru J, Yano M. Core single-nucleotide polymorphisms—a tool for genetic analysis of the Japanese rice population. <i>Breed. Sci.</i> 60(5), 648–655(2010)
282	Y. Kawahara, R. Sakate, A. Matsuya, K. Murakami, Y. Sato, H. Zhang, T. Gojobori, T. Itoh, T. Imanishi. G-compass: A web-based comparative genome browser between human and other vertebrate genomes. <i>Bioinformatics</i> 25(24), 3321–3322(2009)
283	S. Ohmori, M. Kimizu, M. Sugita, A. Miyao, H. Hirochika, E. Uchida, Y. Nagato, H. Yoshida. MOSAIC FLORAL ORGANS1, an AGL6-Like MADS Box Gene, Regulates Floral Organ Identity and Meristem Fate in Rice. <i>Plant Cell</i> 21(10), 3008–3025(2009)
284	N. Fujita, Y. Toyosawa, Y. Utsumi, T. Higuchi, I. Hanashiro, A. Ikegami, S. Akuzawa, M. Yoshida, A. Mori, K. Inomata, R. Itoh, A. Miyao, H. Hirochika, H. Satoh, Y. Nakamura. Characterization of pullulanase (PUL)-deficient mutants of rice (<i>Oryza sativa</i> L.) and the function of PUL on starch biosynthesis in the developing rice endosperm. <i>J Exp Bot</i> 60(3), 1009–1023(2009)
285	H. Numa, M. Nishimura, T. Tanaka, H. Kanamori, C. C. Yang, T. Matsumoto, Y. Nagamura, T. Itoh. Genome-wide validation of <i>Magnaporthe grisea</i> gene structures based on transcription evidence. <i>FEBS Lett</i> 583(4), 797–800(2009)

286	T. Tanaka, K. O. Koyanagi, T. Itoh. Highly diversified molecular evolution of downstream transcription start sites in rice and Arabidopsis. <i>Plant Physiol</i> 149(3), 1316–1324(2009)
287	M. Yoshii, T. Shimizu, M. Yamazaki, T. Higashi, A. Miyao, H. Hirochika, T. Omura. Disruption of a novel gene for a NAC-domain protein in rice confers resistance to Rice dwarf virus. <i>Plant J</i> 57(4), 615–625(2009)
288	Tsuda K, Ito Y, Yamaki S, Miyao A, Hirochika H, Kurata N. Isolation and mapping of three rice mutants that showed ectopic expression of KNOX genes in leaves. <i>Plant Sci</i> 177(2), 131–135(2009)
289	C. Gutjahr, M. Banba, V. Croset, K. An, A. Miyao, G. An, H. Hirochika, H. Imaizumi-Anraku, U. Paszkowski. Arbuscular Mycorrhiza-Specific Signaling in Rice Transcends the Common Symbiosis Signaling Pathway. <i>Plant Cell</i> 20(11), 2989–3005(2008)
290	M. Banba, C. Gutjahr, A. Miyao, H. Hirochika, U. Paszkowski, H. Kouchi, H. Imaizumi-Anraku. Divergence of evolutionary ways among common sym genes: CASTOR and CCaMK show functional conservation between two symbiosis systems and constitute the root of a common signaling pathway. <i>Plant Cell Physiol</i> 49(11), 1659–1671(2008)
291	S. K. Lee, J. S. Jeon, F. Bornke, L. Voll, J. I. Cho, C. H. Goh, S. W. Jeong, Y. I. Park, S. J. Kim, S. B. Choi, A. Miyao, H. Hirochika, G. An, M. H. Cho, S. H. Bhoo, U. Sonnewald, T. R. Hahn. Loss of cytosolic fructose-1,6-bisphosphatase limits photosynthetic sucrose synthesis and causes severe growth retardations in rice (<i>Oryza sativa</i>). <i>Plant Cell Environ</i> 31(12), 1851–1863(2008)
292	H. Chen, P. P. Samadder, Y. Tanaka, T. Ohira, H. Okuizumi, N. Yamaoka, A. Miyao, H. Hirochika, T. Ohira, S. Tsuchimoto, H. Ohtsubo, M. Nishiguchi. OsRecQ1, a QDE-3 homologue in rice, is required for RNA silencing induced by particle bombardment for inverted repeat DNA, but not for double-stranded RNA. <i>Plant J</i> 56(2), 274–286(2008)
293	Meier S, Gehring C, MacPherson C.R, Kaur M, Maqungo M, Reuben S, Muyanga S, Shih M-D, Wei F-J, Wanchana S, Mauleon R, Radovanovic A, Bruskiwich R, Tanaka T, Mohanty B, Itoh T, Wing R, Gojobori T, Sasaki T, Swarup S, Hsing Y, Bajic V.B. The promoter signatures in rice LEA genes can be used to build a co-expressing LEA gene network. <i>Rice</i> 1(2), 177–187(2008)
294	Kawahara Y, de la Bastide M, Hamilton JP, Kanamori M, McCombie WR, Ouyang S, Schwartz D, Tanaka T, Wu J, Zhou S, Childs KL, Davidson RM, Lin H, Quesada-Ocampo L, Vaillancourt B, Sakai H, Lee SS, Kim J, Numa H, Itoh T, Buell CR, Matsumoto T Improvement of the <i>Oryza sativa</i> Nipponbare reference genome using next generation sequence and optical map data. <i>Rice</i> (in press)
295	Sakai H, Lee SS, Tanaka T, Numa H, Kim J, Kawahara Y, Wakimoto H, Yang CC, Iwamoto M, Abe T, Yamada Y, Muto A, Inokuchi H, Ikemura T, Matsumoto T, Sasaki T, Itoh T Rice Annotation Project Database (RAP-DB): An integrative and interactive database for rice genomics. <i>Plant Cell Physiol</i> . (in press)
296	A. Takahashi, N. Hayashi, A. Miyao, H. Hirochika. Unique features of the rice blast resistance Pish locus revealed by large scale retrotransposon-tagging. <i>BMC Plant Biol</i> 10, 175(2010)
297	N. Amano, T. Tanaka, H. Numa, H. Sakai, T. Itoh. Efficient plant gene identification based on interspecies mapping of full-length cDNAs. <i>DNA Res</i> 17(5), 271–279(2010)
298	T. Hirose, Z. Zhang, A. Miyao, H. Hirochika, R. Ohsugi, T. Terao. Disruption of a gene for rice sucrose transporter, OsSUT1, impairs pollen function but pollen maturation is unaffected. <i>J Exp Bot</i> 61(13), 3639–3646(2010)
299	T. Kurusu, H. Hamada, Y. Sugiyama, T. Yagala, Y. Kadota, T. Furuichi, T. Hayashi, K. Umemura, S. Komatsu, A. Miyao, H. Hirochika, K. Kuchitsu. Negative feedback regulation of microbe-associated molecular pattern-induced cytosolic Ca(2+) transients by protein phosphorylation. <i>J Plant Res</i> 124(3), 415–424(2011)
300	H. Matsui, A. Miyao, A. Takahashi, H. Hirochika. Pdk1 Kinase Regulates Basal Disease Resistance Through the OsOxi1-OsPti1a Phosphorylation Cascade in Rice. <i>Plant Cell Physiol</i> 51(12), 2082–2091(2010)
301	K. Kobayashi, M. Maekawa, A. Miyao, H. Hirochika, J. Kyozuka. PANICLE PHYTOMER2 (PAP2), encoding a SEPALLATA subfamily MADS-box protein, positively controls spikelet meristem identity in rice. <i>Plant Cell Physiol</i> 51(1), 47–57(2010)

302	M. Mihara, T. Itoh, T. Izawa. In silico identification of short nucleotide sequences associated with gene expression of pollen development in rice. <i>Plant Cell Physiol</i> 49(10), 1451–1464(2008)
303	M. Kishi-Kaboshi, K. Okada, L. Kurimoto, S. Murakami, T. Umezawa, N. Shibuya, H. Yamane, A. Miyao, H. Takatsuji, A. Takahashi, H. Hirochika. A rice fungal MAMP-responsive MAPK cascade regulates metabolic flow to antimicrobial metabolite synthesis. <i>Plant J</i> 63(4), 599–612(2010)
304	M. Yoshii, M. Yamazaki, R. Rakwal, M. Kishi-Kaboshi, A. Miyao, H. Hirochika. The NAC transcription factor RIM1 of rice is a new regulator of jasmonate signaling. <i>Plant J</i> 61(5), 804–815(2010)
305	K. Asano, A. Miyao, H. Hirochika, H. Kitano, M. Matsuoka, M. Ashikari. SSD1, which encodes a plant-specific novel protein, controls plant elongation by regulating cell division in rice. <i>Proc Jpn Acad Ser B Phys Biol Sci</i> 86(3), 265–273(2010)
306	K. Sugimoto, Y. Takeuchi, K. Ebana, A. Miyao, H. Hirochika, N. Hara, K. Ishiyama, M. Kobayashi, Y. Ban, T. Hattori, M. Yano. Molecular cloning of Sdr4, a regulator involved in seed dormancy and domestication of rice. <i>Proc Natl Acad Sci U S A</i> 107(13), 5792–5797(2010)
307	Myint K. K. M., H. Yasui, M. Takagi, M. Matsumura. Virulence of long-term laboratory populations of the brown planthopper, <i>Nilaparvata lugens</i> (Stal), and whitebacked planthopper, <i>Sogatella furcifera</i> (Horvath) (Homoptera: Delphacidae), on rice differential varieties. <i>Appl. Entomol. Zool.</i> 44(1), 149–153(2009)
308	S. Fujii, T. Toda, S. Kikuchi, R. Suzuki, K. Yokoyama, H. Tsuchida, K. Yano, K. Toriyama. Transcriptome map of plant mitochondria reveals islands of unexpected transcribed regions. <i>BMC Genomics.</i> 12, 279(2011)
309	Yusaku Uga, Kaworu Ebana, Jun Abe, Shigenori Morita, Kazutoshi Okuno, Masahiro Yano. Variation in root morphology and anatomy among accessions of cultivated rice (<i>Oryza sativa</i> L.) with different genetic backgrounds. <i>Breed. Sci.</i> 59(1), 87–93(2009)
310	Keisuke Nagai, Takeshi Kuroha, Madoka Ayano, Yusuke Kurokawa, Rosalyn B. Angeles-Shim, Jung-Hyun Shim, Hideshi Yasui, Atsushi Yoshimura, Motoyuki Ashikari. Two novel QTLs regulate internode elongation in deepwater rice during the early vegetative stage. <i>Breed. Sci.</i> 62(2), 178–185(2012)
311	Natsuko Iwata, Hiroshi Shinada, Hitoshi Kiuchi, Takashi Sato, Kenji Fujino. Mapping of QTLs controlling seedling establishment using a direct seeding method in rice. <i>Breed. Sci.</i> 60(4), 353–360(2010)
312	Takeuchi Y, Hori K, Suzuki K, Nonoue Y, Takemoto-Kuno Y, Maeda H, Sato H, Hirabayashi H, Ohta H, Ishii T, Kato H, Nemoto H, Imbe T, Ohtsubo K, Yano M, Ando I. Major QTLs for eating quality of an elite Japanese rice cultivar, Koshihikari, on the short arm of chromosome 3. <i>Breed. Sci.</i> 58(4), 437–445(2008)
313	Daisuke Fujita, Kazuyuki Doi, Atsushi Yoshimura, Hideshi Yasui. A major QTL for resistance to green rice leafhopper (<i>Nephotettix cincticeps</i> Uhler) derived from African rice (<i>Oryza glaberrima</i> Steud.). <i>Breed. Sci.</i> 60(4), 336–341(2010)
314	Khin Thanda Win, Takahiko Kubo, Yuta Miyazaki, Kazuyuki Doi, Yoshiyuki Yamagata, Atsushi Yoshimura. Identification of two loci causing F1 pollen sterility in inter- and intraspecific crosses of rice. <i>Breed. Sci.</i> 59(4), 411–418(2009)
315	Akasaka M, Konishi S, Izawa T, Ushiki J. Histological and genetic characteristics associated with the seed-shattering habit of weedy rice (<i>Oryza sativa</i> L.) from Okayama, Japan. <i>Breed. Sci.</i> 61(2), 168–173(2011)
316	I. Ando, H. Sato, N. Aoki, Y. Suzuki, H. Hirabayashi, M. Kuroki, H. Shimizu, T. Ando, Y. Takeuchi. Genetic analysis of the low-amylose characteristics of rice cultivars Oborozuki and Hokkai-PL9. <i>Breed. Sci.</i> 60, 187–194(2010)
317	Asanori Yara, Cong Nguyen Phi, Masaya Matsumura, Atsushi Yoshimura, Hideshi Yasui. Development of near-isogenic lines for BPH25(t) and BPH26(t), which confer resistance to the brown planthopper, <i>Nilaparvata lugens</i> (Stal.) in indica rice 'ADR52'. <i>Breed. Sci.</i> 60(5), 639–647(2010)
318	Daisuke Fujita, Atsushi Yoshimura, Hideshi Yasui. Development of near-isogenic lines and pyramided lines carrying resistance genes to green rice leafhopper (<i>Nephotettix cincticeps</i> Uhler) using the Taichung 65 genetic background in rice (<i>Oryza sativa</i> L.). <i>Breed. Sci.</i> 60(1), 18–27(2010)
319	Abe T, Taguchi-Shiobara F, Kojima Y, Ebitani T, Kuramata M, Yamamoto T, Yano M, Ishikawa S. Detection of a QTL for accumulating Cd in rice that enables efficient Cd phytoextraction from soil. <i>Breed. Sci.</i> 61(1), 43–51(2011)

320	Mayuko Ikeda, Yoshitsugu Hirose, Tomonori Takashi, Yosuke Shibata, Takuya Yamamura, Toshiro Komura, Kazuyuki Doi, Motoyuki Ashikari, Makoto Matsuoka, Hidemi Kitano. Analysis of rice panicle traits and detection of QTLs using an image analyzing method. <i>Breed. Sci.</i> 60(1), 55–64(2010)
321	Hideki Sato, Sachiko Shirasawa, Hiroaki Maeda, Koji Nakagomi, Ryota Kaji, Hisatoshi Ohta, Masayuki Yamaguchi, Takeshi Nishio. Analysis of QTL for lowering cadmium concentration in rice grains from ‘LAC23’. <i>Breed. Sci.</i> 61(2), 196–200(2011)
322	鈴木保宏. 脂質分解酵素欠失イネの開発による米油生産性向上および貯蔵性改良に関する育種の試み. <i>FFIジャーナル</i> . 217(3), 275–283(2012)
323	Akihiro Ohsumi, Toshiyuki Takai, Masashi Ida, Toshio Yamamoto, Yumiko Arai-Sanoh, Masahiro Yano, Tsuyu Ando, Motohiko Kondo. Evaluation of yield performance in rice near-isogenic lines with increased spikelet number. <i>Field Crops Res.</i> 120(1), 68–75(2010)
324	T Takai, M. Yano, T. Yamamoto. Canopy temperature on clear and cloudy days can be used to estimate varietal differences in stomatal conductance in rice. <i>Field Crops Res.</i> 115(2), 165–170(2010)
325	T. Kubo, Y. Yamagata, M. Eguchi, A. Yoshimura. A novel epistatic interaction at two loci causing hybrid male sterility in an inter-subspecific cross of rice (<i>Oryza sativa</i> L.). <i>Genes Genet Syst.</i> 83(6), 443–453(2008)
326	M. Hattori, M. Nakamura, S. Komatsu, K. Tsuchihara, Y. Tamura, T. Hasegawa. Molecular cloning of a novel calcium-binding protein in the secreted saliva of the green rice leafhopper <i>Nephotettix cincticeps</i> . <i>Insect Biochem Mol Biol.</i> 42(1), 1–9(2012)
327	D. Ueno, E. Koyama, N. Yamaji, J. F. Ma. Physiological, genetic, and molecular characterization of a high-Cd-accumulating rice cultivar, Jarjan. <i>J Exp Bot.</i> 62(7), 2265–2272(2011)
328	S. Ishikawa, T. Abe, M. Kuramata, M. Yamaguchi, T. Ando, T. Yamamoto, M. Yano. A major quantitative trait locus for increasing cadmium-specific concentration in rice grain is located on the short arm of chromosome 7. <i>J Exp Bot.</i> 61(3), 923–934(2010)
329	R. D. Taylaran, S. Adachi, T. Ookawa, H. Usuda, T. Hirasawa. Hydraulic conductance as well as nitrogen accumulation plays a role in the higher rate of leaf photosynthesis of the most productive variety of rice in Japan. <i>J Exp Bot.</i> 62(11), 4067–4077(2011)
330	K. K. M. Myint, M. Matsumura, M. Takagi, H. Yasui. Demographic parameters of long-term laboratory strains of the brown planthopper, <i>Nilaparvata lugens</i> Stal, (Homoptera: Delphacidae) on resistance genes, bph20(t) and Bph21(t) in rice. <i>J. Fac. Agr. Kyushu Univ.</i> 54(1), 159–164(2009)
331	Toshiyuki Takai, Akihiro Ohsumi, Yumiko Arai, Norio Iwasawa, Masahiro Yano, Toshio Yamamoto, Satoshi Yoshinaga, Motohiko Kondo. QTL Analysis of Leaf Photosynthesis in Rice. <i>JARQ.</i> , in press(2012)
332	Yusaku Uga, Kazutoshi Okuno, Masahiro Yano. Fine mapping of Sta1, a quantitative trait locus determining stele transversal area, on rice chromosome 9. <i>Mol. Breed.</i> 26(3), 533–538(2010)
333	A. Shomura, T. Izawa, K. Ebana, T. Ebitani, H. Kanegae, S. Konishi, M. Yano. Deletion in a gene associated with grain size increased yields during rice domestication. <i>Nat Genet.</i> 40(8), 1023–1028(2008)
334	Y. Hattori, K. Nagai, S. Furukawa, X. J. Song, R. Kawano, H. Sakakibara, J. Wu, T. Matsumoto, A. Yoshimura, H. Kitano, M. Matsuoka, H. Mori, M. Ashikari. The ethylene response factors SNORKEL1 and SNORKEL2 allow rice to adapt to deep water. <i>Nature.</i> 460(7258), 1026–1030(2009)
335	D. Ueno, I. Kono, K. Yokosho, T. Ando, M. Yano, J. F. Ma. A major quantitative trait locus controlling cadmium translocation in rice (<i>Oryza sativa</i>). <i>New Phytol.</i> 182(3), 644–653(2009)
336	M. J. Milner, E. Craft, N. Yamaji, E. Koyama, J. F. Ma, L. V. Kochian. Characterization of the high affinity Zn transporter from <i>Noccaea caerulea</i> , NcZNT1, and dissection of its promoter for its role in Zn uptake and hyperaccumulation. <i>New Phytol.</i> 195(1), 113–123(2012)
337	Kojima H, Kazama T, Fujii S, Toriyama K. Cytoplasmic male sterility-associated ORF79 is toxic to plant regeneration when expressed with mitochondrial targeting sequence of ATPase gamma subunit. <i>Plant Biotechnology.</i> 27(1), 111–114(2010)
338	A. Sasaki, N. Yamaji, K. Yokosho, J. F. Ma. Nramp5 Is a Major Transporter Responsible for Manganese and Cadmium Uptake in Rice. <i>Plant Cell.</i> 24(5), 2155–2167(2012)

339	D. Ueno, E. Koyama, I. Kono, T. Ando, M. Yano, J. F. Ma. Identification of a novel major quantitative trait locus controlling distribution of Cd between roots and shoots in rice. <i>Plant Cell Physiol.</i> 50(12), 2223–2233(2009)
340	H. Saito, E. Ogiso–Tanaka, Y. Okumoto, Y. Yoshitake, H. Izumi, T. Yokoo, K. Matsubara, K. Hori, M. Yano, H. Inoue, T. Tanisaka. Ef7 encodes an ELF3–like protein and promotes rice flowering by negatively regulating the floral repressor gene Ghd7 under both short– and long–day conditions. <i>Plant Cell Physiol.</i> 53(4), 717–728(2012)
341	S. Fujii, K. Toriyama. DCW11, down–regulated gene 11 in CW–type cytoplasmic male sterile rice, encoding mitochondrial protein phosphatase 2c is related to cytoplasmic male sterility. <i>Plant Cell Physiol.</i> 49(4), 633–640(2008)
342	E. Itabashi, T. Kazama, K. Toriyama. Characterization of cytoplasmic male sterility of rice with Lead Rice cytoplasm in comparison with that with Chinsurah Boro II cytoplasm. <i>Plant Cell Rep.</i> 28(2), 233–239(2009)
343	T. Kazama, T. Nakamura, M. Watanabe, M. Sugita, K. Toriyama. Suppression mechanism of mitochondrial ORF79 accumulation by Rf1 protein in BT–type cytoplasmic male sterile rice. <i>Plant J.</i> 55(4), 619–628(2008)
344	F. Matsuda, K. Yonekura–Sakakibara, R. Niida, T. Kuromori, K. Shinozaki, K. Saito. MS/MS spectral tag–based annotation of non–targeted profile of plant secondary metabolites. <i>Plant J.</i> 57(3), 555–577(2009)
345	K. Fujino, Y. Matsuda. Genome–wide analysis of genes targeted by qLTG3–1 controlling low–temperature germinability in rice. <i>Plant Mol Biol.</i> 72(41276), 137–152(2010)
346	R. D. Taylaran, S. Ozawa, N. Miyamoto, T. Ookawa, T. Motobayashi, T. Hirasawa. Performance of a High–Yielding Modern Rice Cultivar Takanari and Several Old and New Cultivars Grown with and without Chemical Fertilizer in a Submerged Paddy Field. <i>Plant Prod Sci.</i> 12(3), 365–380(2009)
347	S. Adachi, N. Nito, M. Kondo, T. Yamamoto, Y. Arai–Sanoh, T. Ando, T. Ookawa, M. Yano, T. Hirasawa. Identification of Chromosomal Regions Controlling the Leaf Photosynthetic Rate in Rice by Using a Progeny from Japonica and High–yielding Indica Varieties. <i>Plant Prod Sci.</i> 14(2), 118–127(2011)
348	T. Tsukaguchi, Y. Iida. Effects of assimilate supply and high temperature during grain–filling period on the occurrence of various types of chalky kernels in rice plants (<i>Oryza sativa</i> L.). <i>Plant Prod Sci.</i> 11(2), 203–210(2008)
349	T. Hirasawa, S. Ozawa, R. D. Taylaran, T. Ookawa. Varietal Differences in Photosynthetic Rates in Rice Plants, with Special Reference to the Nitrogen Content of Leaves. <i>Plant Production Science.</i> 13(1), 53–57(2009)
350	T. Ookawa, K. Yasuda, M. Seto, K. Sunaga, H. Kato, M. Sakai, T. Motobayashi, T. Hirasawa. Biomass Production and Lodging Resistance in ‘Leaf Star’, a New Long–Culm Rice Forage Cultivar. <i>Plant Production Science.</i> 13(1), 58–66(2009)
351	M. Kano, Y. Inukai, H. Kitano, A. Yamauchi. Root plasticity as the key root trait for adaptation to various intensities of drought stress in rice. <i>Plant Soil.</i> 342(41276), 117–128(2011)
352	K. Fujino, H. Sekiguchi, Y. Matsuda, K. Sugimoto, K. Ono, M. Yano. Molecular identification of a major quantitative trait locus, qLTG3–1, controlling low–temperature germinability in rice. <i>Proc Natl Acad Sci U S A.</i> 105(34), 12623–12628(2008)
353	Y. Yamagata, E. Yamamoto, K. Aya, K. T. Win, K. Doi, Sobrizal, T. Ito, H. Kanamori, J. Wu, T. Matsumoto, M. Matsuoka, M. Ashikari, A. Yoshimura. Mitochondrial gene in the nuclear genome induces reproductive barrier in rice. <i>Proc Natl Acad Sci U S A.</i> 107(4), 1494–1499(2010)
354	K. Asano, M. Yamasaki, S. Takuno, K. Miura, S. Katagiri, T. Ito, K. Doi, J. Wu, K. Ebana, T. Matsumoto, H. Innan, H. Kitano, M. Ashikari, M. Matsuoka. Artificial selection for a green revolution gene during japonica rice domestication. <i>Proc Natl Acad Sci U S A.</i> 108(27), 11034–11039(2011)
355	Takai T, Kondo M, Yano M, Yamamoto T. A quantitative trait locus for chlorophyll content and its association with leaf photosynthesis in rice. <i>Rice.</i> 3(41308), 172–180(2010)

356	Hiroko Yamane, Tomoko Ito, Harumi Ishikubo, Masaki Fujisawa, Harumi Yamagata, Kozue Kamiya, Yukiyo Ito, Masao Hamada, Hiroyuki Kanamori, Hiroshi Ikawa, Yuichi Katayose, Jianzhong Wu, Takuji Sasaki, Takashi Matsumoto. Molecular and Evolutionary Analysis of the Hd6 Photoperiod Sensitivity Gene within Genus <i>Oryza</i> . <i>Rice</i> . 2(1), 56–66(2009)
357	Toda T, Fujii S, Kazama T, Itoh Y, Toriyama K. Prediction of organelle targeting of pentatricopeptide repeat protein and genes tagged with Tos17. <i>Rice Genet. Newsl.</i> 25, 85–86(2010)
358	S. Fukuoka, N. Saka, H. Koga, K. Ono, T. Shimizu, K. Ebana, N. Hayashi, A. Takahashi, H. Hirochika, K. Okuno, M. Yano. Loss of function of a proline-containing protein confers durable disease resistance in rice. <i>Science</i> . 325(5943), 998–1001(2009)
359	K. Ebana, T. Shibaya, J. Wu, K. Matsubara, H. Kanamori, H. Yamane, U. Yamanouchi, T. Mizubayashi, I. Kono, A. Shomura, S. Ito, T. Ando, K. Hori, T. Matsumoto, M. Yano. Uncovering of major genetic factors generating naturally occurring variation in heading date among Asian rice cultivars. <i>Theor Appl Genet.</i> 122(6), 1199–1210(2011)
360	K. Fujino, N. Iwata. Selection for low-temperature germinability on the short arm of chromosome 3 in rice cultivars adapted to Hokkaido, Japan. <i>Theor Appl Genet.</i> 123(7), 1089–1097(2011)
361	S. Fukuoka, R. Mizobuchi, N. Saka, S. Ivan, T. Matsumoto, K. Okuno, M. Yano. A multiple gene complex on rice chromosome 4 is involved in durable resistance to rice blast. <i>Theor Appl Genet.</i> 125(3), 551–559(2012)
362	K. K. Myint, D. Fujita, M. Matsumura, T. Sonoda, A. Yoshimura, H. Yasui. Mapping and pyramiding of two major genes for resistance to the brown planthopper (<i>Nilaparvata lugens</i> [Stal]) in the rice cultivar ADR52. <i>Theor Appl Genet.</i> 124(3), 495–504(2012)
363	T. Ando, T. Yamamoto, T. Shimizu, X. F. Ma, A. Shomura, Y. Takeuchi, S. Y. Lin, M. Yano. Genetic dissection and pyramiding of quantitative traits for panicle architecture by using chromosomal segment substitution lines in rice. <i>Theor Appl Genet.</i> 116(6), 881–890(2008)
364	T. Takai, A. Ohsumi, Y. San-oh, M. R. Laza, M. Kondo, T. Yamamoto, M. Yano. Detection of a quantitative trait locus controlling carbon isotope discrimination and its contribution to stomatal conductance in japonica rice. <i>Theor Appl Genet.</i> 118(7), 1401–1410(2009)
365	田畑美奈子, 飯田幸彦, 奥野眞敏. 根群の一部切除が水稲玄米の外観品質に及ぼす影響. <i>日作紀.</i> 77, 198–203(2008)
366	早田一也, 大川泰一郎, 本林隆, 平沢正. 葉身傾斜角度による水稲の個体群吸光係数の簡易評価法の検討. <i>日本作物学会紀事.</i> 79, 174–183(2010)
367	山口琢也, 表野元保, 蛭谷武志. 低温土中出芽性を高めたコシヒカリ染色体部分置換系統「と系1370」に対する播種前処理の影響. <i>北陸作物学会報.</i> (45), 31–34(2010)
368	S. Adachi, Y. Tsuru, M. Kondo, T. Yamamoto, Y. Arai-Sanoh, T. Ando, T. Ookawa, M. Yano, T. Hirasawa. Characterization of a rice variety with high hydraulic conductance and identification of the chromosome region responsible using chromosome segment substitution lines. <i>Ann Bot</i> 106(5), 803–811(2010)
369	S. Fujii, T. Kazama, M. Yamada, K. Toriyama. Discovery of global genomic re-organization based on comparison of two newly sequenced rice mitochondrial genomes with cytoplasmic male sterility-related genes. <i>BMC Genomics</i> 11, 209(2010)
370	N. Iwata, K. Fujino. Genetic effects of major QTLs controlling low-temperature germinability in different genetic backgrounds in rice (<i>Oryza sativa</i> L.). <i>Genome</i> 53(10), 763–768(2010)
371	S. Adachi, Y. Tsuru, N. Nito, K. Murata, T. Yamamoto, T. Ebitani, T. Ookawa, T. Hirasawa. Identification and characterization of genomic regions on chromosomes 4 and 8 that control the rate of photosynthesis in rice leaves. <i>J Exp Bot</i> 62(6), 1927–1938(2011)
372	Y. Uga, K. Okuno, M. Yano. <i>Dro1</i> , a major QTL involved in deep rooting of rice under upland field conditions. <i>J Exp Bot</i> 62(8), 2485–2494(2011)
373	K. Fujino, J. Wu, H. Sekiguchi, T. Ito, T. Izawa, T. Matsumoto. Multiple introgression events surrounding the Hd1 flowering-time gene in cultivated rice, <i>Oryza sativa</i> L. <i>Mol Genet Genomics</i> 284(2), 137–146(2010)
374	T. Izawa, A. Shomura, S. Konishi, K. Ebana, M. Yano. Reply to “Japonica rice carried to, not from, Southeast Asia”. <i>Nat Genet</i> 40(11), 1265–1266(2008)

375	K. Miura, M. Ikeda, A. Matsubara, X. J. Song, M. Ito, K. Asano, M. Matsuoka, H. Kitano, M. Ashikari. OsSPL14 promotes panicle branching and higher grain productivity in rice. <i>Nat Genet</i> 42(6), 545–549(2010)
376	S. Konishi, K. Ebana, T. Izawa. Inference of the japonica rice domestication process from the distribution of six functional nucleotide polymorphisms of domestication-related genes in various landraces and modern cultivars. <i>Plant Cell Physiol</i> 49(9), 1283–1293(2008)
377	S. Fujii, M. Yamada, M. Fujita, E. Itabashi, K. Hamada, K. Yano, N. Kurata, K. Toriyama. Cytoplasmic–nuclear genomic barriers in rice pollen development revealed by comparison of global gene expression profiles among five independent cytoplasmic male sterile lines. <i>Plant Cell Physiol</i> 51(4), 610–620(2010)
378	S. Fujii, M. Yamada, K. Toriyama. Cytoplasmic male sterility-related protein kinase, OsNek3, is regulated downstream of mitochondrial protein phosphatase 2C, DCW11. <i>Plant Cell Physiol</i> 50(4), 828–837(2009)
379	E. Itabashi, N. Iwata, S. Fujii, T. Kazama, K. Toriyama. The fertility restorer gene, Rf2, for Lead Rice-type cytoplasmic male sterility of rice encodes a mitochondrial glycine-rich protein. <i>Plant J</i> 65(3), 359–367(2011)
380	K. Fujino, H. Sekiguchi. Origins of functional nucleotide polymorphisms in a major quantitative trait locus, qLTG3-1, controlling low-temperature germinability in rice. <i>Plant Mol Biol</i> 75(1–2), 1–10(2011)
381	S. Fujii, K. Toriyama. Suppressed expression of Retrograde-Regulated Male Sterility restores pollen fertility in cytoplasmic male sterile rice plants. <i>Proc Natl Acad Sci U S A</i> 106(23), 9513–9518(2009)
382	D. Ueno, N. Yamaji, I. Kono, C. F. Huang, T. Ando, M. Yano, J. F. Ma. Gene limiting cadmium accumulation in rice. <i>Proc Natl Acad Sci U S A</i> 107(38), 16500–16505(2010)
383	H. Saito, Q. Yuan, Y. Okumoto, K. Doi, A. Yoshimura, H. Inoue, M. Teraishi, T. Tsukiyama, T. Tanisaka. Multiple alleles at Early flowering 1 locus making variation in the basic vegetative growth period in rice (<i>Oryza sativa</i> L.). <i>Theor Appl Genet</i> 119(2), 315–323(2009)
384	K. T. Win, Y. Yamagata, Y. Miyazaki, K. Doi, H. Yasui, A. Yoshimura. Independent evolution of a new allele of F(1) pollen sterility gene S27 encoding mitochondrial ribosomal protein L27 in <i>Oryza nivara</i> . <i>Theor Appl Genet</i> 122(2), 385–394(2010)
385	Q. Yuan, H. Saito, Y. Okumoto, H. Inoue, H. Nishida, T. Tsukiyama, M. Teraishi, T. Tanisaka. Identification of a novel gene ef7 conferring an extremely long basic vegetative growth phase in rice. <i>Theor Appl Genet</i> 119(4), 675–684(2009)
386	H. Mizuno, H. Kawahigashi, Y. Kawahara, H. Kanamori, J. Ogata, H. Minami, T. Itoh, T. Matsumoto. Global transcriptome analysis reveals distinct expression among duplicated genes during sorghum–interaction. <i>BMC Plant Biol.</i> 12, 121(2012)
387	Kawahigashi H, Kasuga S, Okuizumi H, Kanamori H, Ando T, Matsumoto T. Classification of genotypes of the target leaf spot-resistant gene (ds1) in a sorghum collection. <i>Crop Sci.</i> 51(5), 2095–2103(2011)
388	Tomoyuki Takai, Jun-ichi Yonemaru, Hirokazu Kaidai, Shigemitsu Kasuga. Quantitative trait locus analysis for days-to-heading and morphological traits in an RIL population derived from an extremely late flowering F1 hybrid of sorghum. <i>Euphytica.</i> 187(3), 411–420(2012)
389	Hiroyuki Kawahigashi, Shigemitsu Kasuga, Hisahito Okuizumi, Syuntaro Hiradate, Jun-ichi Yonemaru. Evaluation of Brix and sugar content in stem juice from sorghum varieties. <i>Grassland Science.</i> , in press (2012)
390	Tamura K, Kiyoshi T, Yonemaru J. The development of highly transferable intron-spanning markers for temperate forage grasses. <i>Mol. Breed.</i> , 1–8(2011)
391	H. Kawahigashi, S. Kasuga, T. Ando, H. Kanamori, J. Wu, J. Yonemaru, T. Sazuka, T. Matsumoto. Positional cloning of ds1, the target leaf spot resistance gene against <i>Bipolaris sorghicola</i> in sorghum. <i>Theor Appl Genet.</i> 123(1), 131–142(2011)
392	J. Yonemaru, T. Ando, T. Mizubayashi, S. Kasuga, T. Matsumoto, M. Yano. Development of genome-wide simple sequence repeat markers using whole-genome shotgun sequences of sorghum (<i>Sorghum bicolor</i> (L.) Moench). <i>DNA Res.</i> 16(3), 187–193(2009)
393	Tamura K, Yonemaru J. Next-generation sequencing for comparative transcriptomics of perennial ryegrass (<i>Lolium perenne</i> L.) and meadow fescue (<i>Festuca pratensis</i> Huds.). <i>Grassland Science.</i> 56(4), 230–239(2010)

394	Ujiie K, Ishimaru K. Identification of chromosome regions affecting leaf area with rice chromosome segment substitution lines. <i>Plant Production Science</i> 16(1), 31–36(2013)
395	Kazuhiro Ujiie, Takayuki Kashiwagi, Ken Ishimaru. Identification and functional analysis of alleles for productivity in two sets of chromosome segment substitution lines of rice. <i>Euphytica</i> 187(3), 325–337(2012)
396	K. Hori, T. Kataoka, K. Miura, M. Yamaguchi, N. Saka, T. Nakahara, Y. Sunohara, K. Ebana, M. Yano. Variation in heading date conceals quantitative trait loci for other traits of importance in breeding selection of rice. <i>Breed Sci</i> 62(3), 223–234(2012)
397	T. Tanabata, T. Shibaya, K. Hori, K. Ebana, M. Yano. SmartGrain: High-Throughput Phenotyping Software for Measuring Seed Shape through Image Analysis. <i>Plant Physiol</i> 160(4), 1871–1880(2012)
398	T. Kudo, N. Makita, M. Kojima, H. Tokunaga, H. Sakakibara. Cytokinin Activity of cis-Zeatin and Phenotypic Alterations Induced by Overexpression of Putative cis-Zeatin-O-glucosyltransferase in Rice. <i>Plant Physiol</i> 160(1), 319–331(2012)
399	K. Matsubara, E. Ogiso-Tanaka, K. Hori, K. Ebana, T. Ando, M. Yano. Natural variation in Hd17, a homolog of Arabidopsis ELF3 that is involved in rice photoperiodic flowering. <i>Plant Cell Physiol</i> 53(4), 709–716(2012)
400	F. Matsuda, Y. Okazaki, A. Oikawa, M. Kusano, R. Nakabayashi, J. Kikuchi, J. Yonemaru, K. Ebana, M. Yano, K. Saito. Dissection of genotype-phenotype associations in rice grains using metabolome quantitative trait loci analysis. <i>Plant J</i> 70(4), 624–636(2012)
401	S. Marzougui, K. Sugimoto, U. Yamanouchi, M. Shimono, T. Hoshino, K. Hori, M. Kobayashi, K. Ishiyama, M. Yano. Mapping and characterization of seed dormancy QTLs using chromosome segment substitution lines in rice. <i>Theor Appl Genet</i> 124(5), 893–902(2012)
402	K. Matsubara, K. Ebana, T. Mizubayashi, S. Itoh, T. Ando, Y. Nonoue, N. Ono, T. Shibaya, E. Ogiso, K. Hori, S. Fukuoka, M. Yano. Relationship between transmission ratio distortion and genetic divergence in intraspecific rice crosses. <i>Mol Genet Genomics</i> 286(5–6), 307–319(2011)
403	I. Kasajima, K. Ebana, T. Yamamoto, K. Takahara, M. Yano, M. Kawai-Yamada, H. Uchimiya. Molecular distinction in genetic regulation of nonphotochemical quenching in rice. <i>Proc Natl Acad Sci U S A</i> 108(33), 13835–13840(2011)
404	T. Shibaya, Y. Nonoue, N. Ono, U. Yamanouchi, K. Hori, M. Yano. Genetic interactions involved in the inhibition of heading by heading date QTL, Hd2 in rice under long-day conditions. <i>Theor Appl Genet</i> 123(7), 1133–1143(2011)
405	K. Asano, M. Yamasaki, S. Takuno, K. Miura, S. Katagiri, T. Ito, K. Doi, J. Wu, K. Ebana, T. Matsumoto, H. Innan, H. Kitano, M. Ashikari, M. Matsuoka. Artificial selection for a green revolution gene during japonica rice domestication. <i>Proc Natl Acad Sci U S A</i> 108(27), 11034–11039(2011)
406	K. Matsubara, U. Yamanouchi, Y. Nonoue, K. Sugimoto, Z. X. Wang, Y. Minobe, M. Yano. Ehd3, encoding a plant homeodomain finger-containing protein, is a critical promoter of rice flowering. <i>Plant J</i> 66(4), 603–612(2011)
407	K. Ebana, T. Shibaya, J. Wu, K. Matsubara, H. Kanamori, H. Yamane, U. Yamanouchi, T. Mizubayashi, I. Kono, A. Shomura, S. Ito, T. Ando, K. Hori, T. Matsumoto, M. Yano. Uncovering of major genetic factors generating naturally occurring variation in heading date among Asian rice cultivars. <i>Theor Appl Genet</i> 122(6), 1199–1210(2011)
408	Y. Uga, K. Okuno, M. Yano. Dro1, a major QTL involved in deep rooting of rice under upland field conditions. <i>J Exp Bot</i> 62(8), 2485–2494(2011)
409	Taguchi-Shiobara F, Kojima Y, Ebitani T, Yano M, Ebana K. Variation in domesticated rice inflorescence architecture revealed by principal component analysis and quantitative trait locus analysis. <i>Breed. Sci.</i> 61(1), 52–60(2011)
410	Abe T, Taguchi-Shiobara F, Kojima Y, Ebitani T, Kuramata M, Yamamoto T, Yano M, Ishikawa S. Detection of a QTL for accumulating Cd in rice that enables efficient Cd phytoextraction from soil. <i>Breed. Sci.</i> 61(1), 43–51(2011)
411	Ebitani T, Hayashi N, Omoteno M, Ozaki H, Yano M, Morikawa M, Fukuta Y. Characterization of Pi13, a blast resistance gene that maps to chromosome 6 in indica rice (<i>Oryza sativa</i> L. variety, Kasalath). <i>Breed. Sci.</i> 61(3), 251–259(2011)
412	Yozo Okazaki, Yukiko Kamide, Masami Yokota Hirai, Kazuki Saito. Plant lipidomics based on hydrophilic interaction chromatography coupled to ion trap time-of-flight mass spectrometry. <i>Metabolomics</i> , online first(2011)

413	S. Adachi, N. Nito, M. Kondo, T. Yamamoto, Y. Arai-Sanoh, T. Ando, T. Ookawa, M. Yano, T. Hirasawa. Identification of Chromosomal Regions Controlling the Leaf Photosynthetic Rate in Rice by Using a Progeny from Japonica and High-yielding Indica Varieties. <i>Plant Prod Sci</i> 14(2), 118-127(2011)
414	楠岡真理子, 七夕高也, 古谷雅理, 斎藤隆文. 時空間投影画像を用いた植物生長量の抽出. <i>画像電子学会誌</i> 40(5), 808-814(2011)
415	T. Yamamoto, H. Nagasaki, J. Yonemaru, K. Ebana, M. Nakajima, T. Shibaya, M. Yano. Fine definition of the pedigree haplotypes of closely related rice cultivars by means of genome-wide discovery of single-nucleotide polymorphisms. <i>BMC Genomics</i> 11, 267(2010)
416	Ebana K, Yonemaru JI, Fukuoka S, Iwata H, Kanamori H, Namiki N, Nagasaki H, Yano M. Genetic structure revealed by a whole-genome single-nucleotide polymorphism survey of diverse accessions of cultivated Asian rice (<i>Oryza sativa</i> L.). <i>Breed. Sci.</i> 60(4), 390-397(2010)
417	Hirabayashi H, Sato H, Nonoue Y, Takemoto Y, Takeuchi Y, Kato H, Nemoto H, Ogawa T, Yano M, Imbe T, Ando I. Development of introgression lines derived from <i>Oryza rufipogon</i> and <i>O. glumaepatula</i> in the genetic background of japonica cultivated rice (<i>O. sativa</i> L.) and evaluation of resistance to rice blast. <i>Breed. Sci.</i> 60(5), 604-612(2010)
418	Nagasaki H, Ebana K, Shibaya T, Yonemaru J, Yano M. Core single-nucleotide polymorphisms—a tool for genetic analysis of the Japanese rice population. <i>Breed. Sci.</i> 60(5), 648-655(2010)
419	Tanabata T, Taguchi-Shiobara F, Kishimoto N, Chechetka S, Shinomura T, Habu Y. A phenomics approach detected differential epigenetic growth regulation between inbreds and their hybrid in <i>Oryza sativa</i> . <i>Mol Breed</i> 26(4), 729-734(2010)
420	Iwata H, Ebana K, Uga Y, Hayashi T, Jannink J-L. Genome-wide association study of grain shape variation among <i>Oryza sativa</i> L. germplasm based on elliptic Fourier analysis. <i>Mol. Breed.</i> 25(2), 203-215(2010)
421	F. Matsuda, Y. Shinbo, A. Oikawa, M. Y. Hirai, O. Fiehn, S. Kanaya, K. Saito. Assessment of metabolome annotation quality: a method for evaluating the false discovery rate of elemental composition searches. <i>PLoS One</i> 4(10), e7490(2009)
422	H. Shimizu, T. Tanabata, X. Xie, N. Inagaki, M. Takano, T. Shinomura, K. T. Yamamoto. Phytochrome-mediated growth inhibition of seminal roots in rice seedlings. <i>Physiol Plant</i> 137(3), 289-297(2009)
423	T. Takai, A. Ohsumi, Y. San-oh, M. R. Laza, M. Kondo, T. Yamamoto, M. Yano. Detection of a quantitative trait locus controlling carbon isotope discrimination and its contribution to stomatal conductance in japonica rice. <i>Theor Appl Genet</i> 118(7), 1401-1410(2009)
424	M. Iwamoto, K. Higo, M. Takano. Circadian clock- and phytochrome-regulated Dof-like gene, <i>Rdd1</i> , is associated with grain size in rice. <i>Plant Cell Environ</i> 32(5), 592-603(2009)
425	T. Kashiwagi, K. Shindoh, N. Hirotsu, K. Ishimaru. Evidence for separate translocation pathways in determining cadmium accumulation in grain and aerial plant parts in rice. <i>BMC Plant Biol</i> 9, 8(2009)
426	Iwata H, K. Ebana, S. Fukuoka, J-L. Jannink, T. Hayashi. Bayesian multilocus association mapping on ordinal and censored traits and its application to the analysis of genetic variation among <i>Oryza sativa</i> L. germplasm. <i>Theor Appl Genet</i> 118(5), 865-880(2009)
427	F. Matsuda, K. Yonekura-Sakakibara, R. Niida, T. Kuromori, K. Shinozaki, K. Saito. MS/MS spectral tag-based annotation of non-targeted profile of plant secondary metabolites. <i>Plant J</i> 57(3), 555-577(2009)
428	Hori K, Yamamoto T, Ebana K, Takeuchi Y, Yano M. A novel quantitative trait locus, <i>qCL1</i> , involved in semi-dwarfism derived from Japanese rice cultivar <i>Nipponbare</i> . <i>Breed. Sci.</i> 59(3), 285-295(2009)
429	Fumio Matsuda, Henning Redestig, Yuji Sawada, Yoko Shinbo, Masami Yokota Hirai, Shigehiko Kanaya, Kazuki Saito. Visualization of metabolite identifier information. <i>Plant Biotechnology</i> 26(5), 479-483(2009)
430	T. Kanbe, H. Sasaki, N. Aoki, T. Yamagishi, T. Ebitani, M. Yano, R. Ohsugi. Identification of QTLs for Improvement of Plant Type in Rice (<i>Oryza sativa</i> L.) Using Koshihikari/Kasalath Chromosome Segment Substitution Lines and Backcross Progeny F-2 Population. <i>Plant Prod Sci</i> 11(4), 447-456(2008)

431	T. Kawakatsu, M. P. Yamamoto, S. Hirose, M. Yano, F. Takaiwa. Characterization of a new rice glutelin gene GluD-1 expressed in the starchy endosperm. <i>J Exp Bot</i> 59(15), 4233-4245(2008)
432	K. Matsubara, I. Kono, K. Hori, Y. Nonoue, N. Ono, A. Shomura, T. Mizubayashi, S. Yamamoto, U. Yamanouchi, K. Shirasawa, T. Nishio, M. Yano. Novel QTLs for photoperiodic flowering revealed by using reciprocal backcross inbred lines from crosses between japonica rice cultivars. <i>Theor Appl Genet</i> 117(6), 935-945(2008)
433	T. Ando, T. Yamamoto, T. Shimizu, X. F. Ma, A. Shomura, Y. Takeuchi, S. Y. Lin, M. Yano. Genetic dissection and pyramiding of quantitative traits for panicle architecture by using chromosomal segment substitution lines in rice. <i>Theor Appl Genet</i> 116(6), 881-890(2008)
434	Takeuchi Y, Hori K, Suzuki K, Nonoue Y, Takemoto-Kuno Y, Maeda H, Sato H, Hirabayashi H, Ohta H, Ishii T, Kato H, Nemoto H, Imbe T, Ohtsubo K, Yano M, Ando I. Major QTLs for eating quality of an elite Japanese rice cultivar, Koshihikari, on the short arm of chromosome 3. <i>Breed. Sci.</i> 58(4), 437-445(2008)
435	Ebana K, Kojima Y, Fukuoka S, Nagamine T, Kawase M. Development of mini core collection of Japanese rice landrace. <i>Breed. Sci.</i> 58(3), 281-291(2008)
436	蛸谷武志, 山本良孝, 矢野昌裕, 舟根政治. 染色体断片置換系統群を利用したイネの玄米外観品質に關与するQTLの検出. <i>育種学研究</i> 10(3), 91-99(2008)
437	七夕 高也, 清水 久代, 篠村 知子, 高野 誠, 宮村(中村) 浩子, 齋藤 隆文. イネ初期生育期における表現型解析のための画像計測システム. <i>電気学会論文誌C部門</i> 128C(6), 962-969(2008)
438	K. Mochida, T. Furuta, K. Ebana, K. Shinozaki, J. Kikuchi. Correlation exploration of metabolic and genomic diversity in rice. <i>BMC Genomics</i> 10, 568(2009)
439	S. Ishikawa, T. Abe, M. Kuramata, M. Yamaguchi, T. Ando, T. Yamamoto, M. Yano. A major quantitative trait locus for increasing cadmium-specific concentration in rice grain is located on the short arm of chromosome 7. <i>J Exp Bot</i> 61(3), 923-934(2010)
440	M. Kojima, T. Kamada-Nobusada, H. Komatsu, K. Takei, T. Kuroha, M. Mizutani, M. Ashikari, M. Ueguchi-Tanaka, M. Matsuoka, K. Suzuki, H. Sakakibara. Highly sensitive and high-throughput analysis of plant hormones using MS-probe modification and liquid chromatography-tandem mass spectrometry: an application for hormone profiling in <i>Oryza sativa</i> . <i>Plant Cell Physiol</i> 50(7), 1201-1214(2009)
441	K. Hori, K. Sugimoto, Y. Nonoue, N. Ono, K. Matsubara, U. Yamanouchi, A. Abe, Y. Takeuchi, M. Yano. Detection of quantitative trait loci controlling pre-harvest sprouting resistance by using backcrossed populations of japonica rice cultivars. <i>Theor Appl Genet</i> 120(8), 1547-1557(2010)
442	K Tsutsui, K Kaneko, I Hanashiro, K Nishinari, T. Mitsui. Characteristics of opaque and translucent parts of high temperature stressed grains of rice. <i>Journal of Applied Glycoscience</i> , in press(2013)
443	Miyamoto T, Ochiai K, Takeshita S, Matoh T. Identification of quantitative trait loci associated with shoot sodium accumulation in rice plants. <i>Soil Science and Plant Nutrition</i> 59(2), in press(2013)
444	N. I. Kobayashi, K. Tanoi, A. Hirose, T. M. Nakanishi. Characterization of rapid intervascular transport of cadmium in rice stem by radioisotope imaging. <i>J Exp Bot</i> 64(2), 507-517(2013)
445	Mayuko Ikeda, Kotaro Miura, Koichiro Aya, Hidemi Kitano, Makoto Matsuoka. The quantitative trait loci (QTLs) and mutations related to breeding of high-yielding rice. <i>Curr Opin Plant Biol</i> , invited paper(2012)
446	M Tsuyukubo, T Ookura, S Tsukui, T Mitsui, M. Kasai. Elution Behavior Analysis of Starch Degrading Enzymes During Rice Cooking with Specific Antibodies. <i>Food Science and Technology Research</i> 18(5), 659-666(2012)
447	Keitaro Tanoi, Natsuko I. Kobayashi, Takayuki Saito, Naoko Iwata, Atsushi Hirose, Yoshimi Ohmae, Ren Iwata, Hisashi Suzuki, Tomoko M. Nakanishi. Application of 28Mg to the Kinetic Study of Mg Uptake by Rice Plants. <i>Journal of Radioanalytical and Nuclear Chemistry</i> , online first(2012)

448	N. I. Kobayashi, N. Iwata, T. Saito, H. Suzuki, R. Iwata, K. Tanoi, T. M. Nakanishi. Application of 28Mg for Characterization of Mg Uptake in Rice Seedling under Different pH Conditions. <i>Journal of Radioanalytical and Nuclear Chemistry</i> , online first(2012)
449	S Suzuki, J. F Ma, N Yamamoto, T Hattori, M. Sakamoto, T. Umezawa. Silicon deficiency promotes lignin accumulation in rice. <i>Plant Biotechnol</i> 29, 391–394(2012)
450	Kao-Chih She, Mitsuhiro Yaeshima, Takemasa Koumoto, Mariko Ohnuma, Tomoko Hiromasa, Mio Hirai, Takuya Matsunaga, Ryouka Tashiro, Tadamas Sasaki, Hiroaki Kusano, Hiroaki Shimada. High-temperature stress susceptibility of representative japonica rice cultivars derived from Norin-22: inadequate ATP supply during seed development may lead to severe damage. <i>Plant Biotechnology</i> , in press(2012)
451	Mayuko Ikeda, Makoto Matsuoka. Major quantitative trait loci (QTLs) involved in high crop yield in rice. <i>Plant Genetics and Genomics: Crops and Models</i> , in press(2012)
452	T Abiko, M Obara, F Abe, K Kawaguchi, A Oyanagi, T. Yamauchi, M. Nakazono. Screening of candidate genes associated with constitutive aerenchyma formation in adventitious roots of the teosinte <i>Zea nicaraguensis</i> . <i>Plant Root</i> 6, 19–27(2012)
453	S Segami, I Kono, T Ando, M Yano, H Kitano, K. and Iwasaki Miura, Y. Rice SMALL AND ROUND SEED5 gene encodes alpha-tubulin and regulates seed cell elongation. <i>Rice</i> 5(4), 電子ジャーナルのためページがありません(2012)
454	Sriprachote A, Kanyawongha P, Pantuwan G, Ochiai K, Match T. Evaluation of Thai rice cultivars with low-grain cadmium. <i>Soil Science and Plant Nutrition</i> 58, 568–572(2012)
455	Sriprachote A, Kanyawongha P, Ochiai K, Match T. Current situation of cadmium-polluted paddy soil, rice and soybean in the Mae Sot District, Tak Province, Thailand. <i>Soil Science and Plant Nutrition</i> 58, 349–359(2012)
456	M. Hakata, M. Kuroda, A. Ohsumi, T. Hirose, H. Nakamura, M. Muramatsu, H. Ichikawa, H. Yamakawa. Overexpression of a rice TIFY gene increases grain size through enhanced accumulation of carbohydrates in the stem. <i>Biosci Biotechnol Biochem</i> 76(11), 2129–2134(2012)
457	T. Tsutsui, N. Yamaji, C. F. Huang, R. Motoyama, Y. Nagamura, J. F. Ma. Comparative genome-wide transcriptional analysis of Al-responsive genes reveals novel Al tolerance mechanisms in rice. <i>PLoS One</i> 7(10), e48197(2012)
458	N. I. Kobayashi, T. Saito, N. Iwata, Y. Ohmae, R. Iwata, K. Tanoi, T. M. Nakanishi. Leaf senescence in rice due to magnesium deficiency-mediated defect in transpiration rate before sugar accumulation and chlorosis. <i>Physiol Plant</i> , in press(2012)
459	N. Yamaji, Y. Chiba, N. Mitani-Ueno, J. Feng Ma. Functional characterization of a silicon transporter gene implicated in silicon distribution in barley. <i>Plant Physiol</i> 160(3), 1491–1497(2012)
460	M. Hakata, M. Kuroda, T. Miyashita, T. Yamaguchi, M. Kojima, H. Sakakibara, T. Mitsui, H. Yamakawa. Suppression of alpha-amylase genes improves quality of rice grain ripened under high temperature. <i>Plant Biotechnol J</i> 10(9), 1110–1117(2012)
461	L. Luo, W. Li, K. Miura, M. Ashikari, J. Kyojuka. Control of Tiller Growth of Rice by OsSPL14 and Strigolactones, Which Work in Two Independent Pathways. <i>Plant Cell Physiol</i> 53(10), 1793–1801(2012)
462	M. Nagano, H. Uchimiya, M. Kawai-Yamada. Plant sphingolipid fatty acid 2-hydroxylases have unique characters unlike their animal and fungus counterparts. <i>Plant Signal Behav</i> 7(11), in press(2012)
463	Z. C. Chen, N. Yamaji, R. Motoyama, Y. Nagamura, J. F. Ma. Up-regulation of a magnesium transporter gene OsMGT1 is required for conferring aluminum tolerance in rice. <i>Plant Physiol</i> 159(4), 1624–1633(2012)
464	H. Xiong, T. Kobayashi, Y. Kakei, T. Senoura, M. Nakazono, H. Takahashi, H. Nakanishi, H. Shen, P. Duan, X. Guo, N. K. Nishizawa, Y. Zuo. AhNRAMP1 iron transporter is involved in iron acquisition in peanut. <i>J Exp Bot</i> 63(12), 4437–4446(2012)
465	K. Kobayashi, N. Yasuno, Y. Sato, M. Yoda, R. Yamazaki, M. Kimizu, H. Yoshida, Y. Nagamura, J. Kyojuka. Inflorescence meristem identity in rice is specified by overlapping functions of three AP1/FUL-like MADS box genes and PAP2, a SEPALLATA MADS box gene. <i>Plant Cell</i> 24(5), 1848–1859(2012)

466	T. Yamamoto, A. Nakamura, H. Iwai, T. Ishii, J. F. Ma, R. Yokoyama, K. Nishitani, S. Satoh, J. Furukawa. Effect of silicon deficiency on secondary cell wall synthesis in rice leaf. <i>J Plant Res</i> 125(6), 771–779(2012)
467	T. Abiko, L. Kotula, K. Shiono, A. I. Malik, T. D. Colmer, M. Nakazono. Enhanced formation of aerenchyma and induction of a barrier to radial oxygen loss in adventitious roots of <i>Zea mays</i> ssp. <i>mays</i> contribute to its waterlogging tolerance as compared with maize (<i>Zea mays</i> ssp. <i>mays</i>). <i>Plant Cell Environ</i> 35(9), 1618–1630(2012)
468	K. Takanashi, H. Takahashi, N. Sakurai, A. Sugiyama, H. Suzuki, D. Shibata, M. Nakazono, K. Yazaki. Tissue-specific transcriptome analysis in nodules of <i>Lotus japonicus</i> . <i>Mol Plant Microbe Interact</i> 25(7), 869–876(2012)
469	M. Fujii, K. Yokosho, N. Yamaji, D. Saisho, M. Yamane, H. Takahashi, K. Sato, M. Nakazono, J. F. Ma. Acquisition of aluminium tolerance by modification of a single gene in barley. <i>Nat Commun</i> 3, 713(2012)
470	J. Montpetit, J. Vivancos, N. Mitani-Ueno, N. Yamaji, W. Remus-Borel, F. Belzile, J. F. Ma, R. R. Belanger. Cloning, functional characterization and heterologous expression of TaLsi1, a wheat silicon transporter gene. <i>Plant Mol Biol</i> 79(1–2), 35–46(2012)
471	G. Duan, T. Kamiya, S. Ishikawa, T. Arao, T. Fujiwara. Expressing ScACR3 in Rice Enhanced Arsenite Efflux and Reduced Arsenic Accumulation in rice grains. <i>Plant Cell Physiol</i> 53(1), 154–163(2012)
472	S. Marzougui, K. Sugimoto, U. Yamanouchi, M. Shimono, T. Hoshino, K. Hori, M. Kobayashi, K. Ishiyama, M. Yano. Mapping and characterization of seed dormancy QTLs using chromosome segment substitution lines in rice. <i>Theor Appl Genet</i> 124(5), 893–902(2012)
473	C. F. Huang, N. Yamaji, Z. Chen, J. F. Ma. A tonoplast-localized half-size ABC transporter is required for internal detoxification of aluminum in rice. <i>Plant J</i> 69(5), 857–867(2012)
474	C. F. Huang, N. Yamaji, K. Ono, J. F. Ma. A leucine-rich repeat receptor-like kinase gene is involved in the specification of outer cell layers in rice roots. <i>Plant J</i> 69(4), 565–576(2012)
475	H. Takehisa, Y. Sato, M. Igarashi, T. Abiko, B. A. Antonio, K. Kamatsuki, H. Minami, N. Namiki, Y. Inukai, M. Nakazono, Y. Nagamura. Genome-wide transcriptome dissection of the rice root system: implications for developmental and physiological functions. <i>Plant J</i> 69(1), 126–140(2012)
476	S. Uruguchi, T. Kamiya, T. Sakamoto, K. Kasai, Y. Sato, Y. Nagamura, A. Yoshida, J. Kyojuka, S. Ishikawa, T. Fujiwara. Low-affinity cation transporter (OsLCT1) regulates cadmium transport into rice grains. <i>Proc Natl Acad Sci U S A</i> 108(52), 20959–20964(2011)
477	K. Aya, G. Suzuki, K. Suwabe, T. Hobo, H. Takahashi, K. Shiono, K. Yano, N. Tsutsumi, M. Nakazono, Y. Nagamura, M. Matsuoka, M. Watanabe. Comprehensive network analysis of anther-expressed genes in rice by the combination of 33 laser microdissection and 143 spatiotemporal microarrays. <i>PLoS One</i> 6(10), e26162(2011)
478	H. Tabuchi, Y. Zhang, S. Hattori, M. Omae, S. Shimizu-Sato, T. Oikawa, Q. Qian, M. Nishimura, H. Kitano, H. Xie, X. Fang, H. Yoshida, J. Kyojuka, F. Chen, Y. Sato. LAX PANICLE2 of Rice Encodes a Novel Nuclear Protein and Regulates the Formation of Axillary Meristems. <i>Plant Cell</i> 23(9), 3276–3287(2011)
479	T. Sakamoto, Y. T. Inui, S. Uruguchi, T. Yoshizumi, S. Matsunaga, M. Mastui, M. Umeda, K. Fukui, T. Fujiwara. Condensin II Alleviates DNA Damage and Is Essential for Tolerance of Boron Overload Stress in Arabidopsis. <i>Plant Cell</i> 23(9), 3533–3546(2011)
480	M. Tanaka, J. Takano, Y. Chiba, F. Lombardo, Y. Ogasawara, H. Onouchi, S. Naito, T. Fujiwara. Boron-Dependent Degradation of NIP5;1 mRNA for Acclimation to Excess Boron Conditions in Arabidopsis. <i>Plant Cell</i> 23(9), 3547–3559(2011)
481	K. Yokosho, N. Yamaji, J. F. Ma. An Al-inducible MATE gene is involved in external detoxification of Al in rice. <i>Plant J</i> 68(6), 1061–1069(2011)
482	M. Fukuda, M. Satoh-Cruz, L. Wen, A. J. Crofts, A. Sugino, H. Washida, T. W. Okita, M. Ogawa, Y. Kawagoe, M. Maeshima, T. Kumamaru. The small GTPase Rab5a is essential for intracellular transport of proglutelin from the Golgi apparatus to the protein storage vacuole and endosomal membrane organization in developing rice endosperm. <i>Plant Physiol</i> 157(2), 632–644(2011)
483	I. Kasajima, K. Ebana, T. Yamamoto, K. Takahara, M. Yano, M. Kawai-Yamada, H. Uchimiya. Molecular distinction in genetic regulation of nonphotochemical quenching in rice. <i>Proc Natl Acad Sci U S A</i> 108(33), 13835–13840(2011)

484	H. Takahashi, H. Saika, H. Matsumura, Y. Nagamura, N. Tsutsumi, N. K. Nishizawa, M. Nakazono. Cell division and cell elongation in the coleoptile of rice alcohol dehydrogenase 1-deficient mutant are reduced under complete submergence. <i>Ann Bot</i> 108(2), 253–261(2011)
485	K. Oda, M. Otani, S. Uraguchi, T. Akihiro, T. Fujiwara. Rice ABCG43 is Cd inducible and confers Cd tolerance on yeast. <i>Biosci Biotechnol Biochem</i> 75(6), 1211–1213(2011)
486	K. Asano, M. Yamasaki, S. Takuno, K. Miura, S. Katagiri, T. Ito, K. Doi, J. Wu, K. Ebana, T. Matsumoto, H. Innan, H. Kitano, M. Ashikari, M. Matsuoka. Artificial selection for a green revolution gene during japonica rice domestication. <i>Proc Natl Acad Sci U S A</i> 108(27), 11034–11039(2011)
487	T. Ohnishi, M. Yoshino, H. Yamakawa, T. Kinoshita. The biotron breeding system: a rapid and reliable procedure for genetic studies and breeding in rice. <i>Plant Cell Physiol</i> 52(7), 1249–1257(2011)
488	N. Mitani-Ueno, N. Yamaji, J. F. Ma. Silicon efflux transporters isolated from two pumpkin cultivars contrasting in Si uptake. <i>Plant Signal Behav</i> 6(7), 991–994(2011)
489	N. Mitani-Ueno, N. Yamaji, F. J. Zhao, J. F. Ma. The aromatic/arginine selectivity filter of NIP aquaporins plays a critical role in substrate selectivity for silicon, boron, and arsenic. <i>J Exp Bot</i> 62(12), 4391–4398(2011)
490	M. S. Yun, T. Umemoto, Y. Kawagoe. Rice debranching enzyme isoamylase3 facilitates starch metabolism and affects plastid morphogenesis. <i>Plant Cell Physiol</i> 52(6), 1068–1082(2011)
491	K. Ochiai, A. Shimizu, Y. Okumoto, T. Fujiwara, T. Matoh. Suppression of a NAC-like transcription factor gene improves boron-toxicity tolerance in rice. <i>Plant Physiol</i> 156(3), 1457–1463(2011)
492	A. Nagamine, H. Matsusaka, T. Ushijima, Y. Kawagoe, M. Ogawa, T. W. Okita, T. Kumamaru. A role for the cysteine-rich 10 kDa prolamin in protein body I formation in rice. <i>Plant Cell Physiol</i> 52(6), 1003–1016(2011)
493	T. Tsutsui, N. Yamaji, J. Feng Ma. Identification of a cis-acting element of ART1, a C2H2-type zinc-finger transcription factor for aluminum tolerance in rice. <i>Plant Physiol</i> 156(2), 925–931(2011)
494	K. L. Moore, M. Schroder, Z. Wu, B. G. Martin, C. R. Hawes, S. P. McGrath, M. J. Hawkesford, J. Feng Ma, F. J. Zhao, C. R. Grovenor. High-resolution secondary ion mass spectrometry reveals the contrasting subcellular distribution of arsenic and silicon in rice roots. <i>Plant Physiol</i> 156(2), 913–924(2011)
495	K. Miura, M. Ashikari, M. Matsuoka. The role of QTLs in the breeding of high-yielding rice. <i>Trends Plant Sci</i> 16(6), 319–326(2011)
496	Y. Onda, A. Nagamine, M. Sakurai, T. Kumamaru, M. Ogawa, Y. Kawagoe. Distinct roles of protein disulfide isomerase and P5 sulfhydryl oxidoreductases in multiple pathways for oxidation of structurally diverse storage proteins in rice. <i>Plant Cell</i> 23(1), 210–223(2011)
497	M. Kusano, M. Tabuchi, A. Fukushima, K. Funayama, C. Diaz, M. Kobayashi, N. Hayashi, Y. N. Tsuchiya, H. Takahashi, A. Kamata, T. Yamaya, K. Saito. Metabolomics data reveal a crucial role of cytosolic glutamine synthetase 1;1 in coordinating metabolic balance in rice. <i>Plant J</i> 66(3), 456–466(2011)
498	M. Takahashi, M. Teranishi, H. Ishida, J. Kawasaki, A. Takeuchi, T. Yamaya, M. Watanabe, A. Makino, J. Hidema. Cyclobutane pyrimidine dimer (CPD) photolyase repairs ultraviolet-B-induced CPDs in rice chloroplast and mitochondrial DNA. <i>Plant J</i> 66(3), 433–442(2011)
499	R. Ishikawa, T. Ohnishi, Y. Kinoshita, M. Eiguchi, N. Kurata, T. Kinoshita. Rice interspecies hybrids show precocious or delayed developmental transitions in the endosperm without change to the rate of syncytial nuclear division. <i>Plant J</i> 65(5), 798–806(2011)
500	J. Xia, N. Yamaji, J. F. Ma. Further characterization of an aluminum influx transporter in rice. <i>Plant Signal Behav</i> 6(1), 160–163(2011)
501	N. Mitani, N. Yamaji, Y. Ago, K. Iwasaki, J. F. Ma. Isolation and functional characterization of an influx silicon transporter in two pumpkin cultivars contrasting in silicon accumulation. <i>Plant J</i> 66(2), 231–240(2011)
502	Kato T., Segami S., Toriyama M., Kono I., Ando T., Yano M., Kitano H., Miura K., Iwasaki Y. Detection of QTLs for grain length from large grain rice (<i>oryza sativa</i> L.). <i>Breed. Sci.</i> 61(3), 269–274(2011)

503	W Tamura, S Kojima, A Toyokawa, H Watanabe, M Tabuchi-Kobayashi, T. Hayakawa, T. Yamaya. Disruption of a novel NADH-glutamate synthase2 gene caused marked reduction in spikelet number of rice. <i>Front. Plant Sci.</i> 2(57), 1-9(2011)
504	K Kaneko, C Yamada, A Yanagida, T Koshu, U Umezawa, K Itoh, H Hori, T. Mitsui. Differential localizations and functions of rice nucleotide pyrophosphatase/phosphodiesterase isozymes 1 and 3. <i>Plant Biotechnology</i> 28(1), 69-76(2011)
505	田野井 慶太郎, 齊藤 貴之, 岩田 直子, 大前 芳美, 広瀬 農, 小林 奈通子, 岩田 錬, 中西 友子. 28Mgの製造とイネにおけるMg吸収解析への利用. <i>RADIOISOTOPES</i> 60(8), 299-304(2011)
506	Kyoko MIWA, Toru FUJIWARA. Role of overexpressed BOR4, a boron exporter, in tolerance to high level of boron in shoots. <i>Soil Sci. Plant Nutr.</i> 57(4), 558-565(2011)
507	Takehiro Kamiya, Toru Fujiwara. A novel allele of the Arabidopsis phytochelatin synthase 1 gene conferring high sensitivity to arsenic and antimony. <i>Soil Sci. Plant Nutr.</i> 57(2), 272-278(2011)
508	M Obara, T Takeda, T. Hayakawa, T. Yamaya. Mapping quantitative trait loci controlling root length in rice seedlings grown with low or sufficient NH ₄ ⁺ supply using backcross recombinant lines derived from a cross between <i>Oryza sativa</i> L. and <i>Oryza glaberrima</i> Steud. <i>Soil Sci. Plant Nutr.</i> 57(1), 80-92(2011)
509	N. Yamaji, J. F. Ma. Further characterization of a rice Si efflux transporter, Lsi2. <i>Soil Sci. Plant Nutr.</i> 57, 259-264(2011)
510	T. Ookawa, T. Hobo, M. Yano, K. Murata, T. Ando, H. Miura, K. Asano, Y. Ochiai, M. Ikeda, R. Nishitani, T. Ebitani, H. Ozaki, E. R. Angeles, T. Hirasawa, M. Matsuoka. New approach for rice improvement using a pleiotropic QTL gene for lodging resistance and yield. <i>Nat Commun</i> 1, 132(2010)
511	S. Yamaguchi, J. Kyojuka. Branching hormone is busy both underground and overground. <i>Plant Cell Physiol</i> 51(7), 1091-1094(2010)
512	K. Minakuchi, H. Kameoka, N. Yasuno, M. Umehara, L. Luo, K. Kobayashi, A. Hanada, K. Ueno, T. Asami, S. Yamaguchi, J. Kyojuka. FINE CULM1 (FC1) works downstream of strigolactones to inhibit the outgrowth of axillary buds in rice. <i>Plant Cell Physiol</i> 51(7), 1127-1135(2010)
513	W. Tamura, Y. Hidaka, M. Tabuchi, S. Kojima, T. Hayakawa, T. Sato, M. Obara, M. Kojima, H. Sakakibara, T. Yamaya. Reverse genetics approach to characterize a function of NADH-glutamate synthase1 in rice plants. <i>Amino Acids</i> 39(4), 1003-1012(2010)
514	Y. Yamagata, E. Yamamoto, K. Aya, K. T. Win, K. Doi, Sobrizal, T. Ito, H. Kanamori, J. Wu, T. Matsumoto, M. Matsuoka, M. Ashikari, A. Yoshimura. Mitochondrial gene in the nuclear genome induces reproductive barrier in rice. <i>Proc Natl Acad Sci U S A</i> 107(4), 1494-1499(2010)
515	K. Kobayashi, M. Maekawa, A. Miyao, H. Hirochika, J. Kyojuka. PANICLE PHYTOMER2 (PAP2), encoding a SEPALLATA subfamily MADS-box protein, positively controls spikelet meristem identity in rice. <i>Plant Cell Physiol</i> 51(1), 47-57(2010)
516	J. F. Ma, P. Ryan. Understanding how plants cope with acid soils. <i>Funct. Plant Biol.</i> 37(4), iii-vi(2010)
517	Miyagi A, Takahara K, Takahashi H, Kawai-Yamada M, Uchimiya H. Targeted metabolomics in an intrusive weed, <i>Rumex obtusifolius</i> L., grown under different environmental conditions reveals alterations of organ related metabolite pathway. <i>Metabolomics</i> 6(4), 497-510(2010)
518	Miyagi A, Takahashi H, Takahara K, Hirabayashi T, Nishimura Y, Tezuka T, Kawai-Yamada M, Uchimiya H. Principal component and hierarchical clustering analysis of metabolites in destructive weeds; polygonaceous plants. <i>Metabolomics</i> 6(1), 146-155(2010)
519	Uraguchi S, Fujiwara T. Significant contribution of boron stored in seeds to initial growth of rice seedlings. <i>Plant and Soil</i> 340(1-2), 435-442(2010)
520	Kao-Chih She, Hiroaki Kusano, Mitsuhiro Yaeshima, Tadamasasa Sasaki, Hikaru Satoh, Hiroaki Shimada. Reduced rice grain production under high-temperature stress closely correlates with ATP shortage during seed development. <i>Plant Biotechnology</i> 27(1), 67-73(2010)
521	T. Ookawa, K. Yasuda, H. Kato, M. Sakai, M. Seto, K. Sunaga, T. Motobayashi, S. Tojo, T. Hirasawa. Biomass production and lodging resistance in 'Leaf Star', a new long-culm rice forage cultivar. <i>Plant Prod Sci</i> 13(1), 58-66(2010)
522	M. Isa, S. Bai, T. Yokoyama, J. F. Ma, Y. Ishibashi, T. Yuasa, M. Iwaya-Inoue. Silicon enhances growth independent of silica deposition in a low-silica rice mutant, Lsi1. <i>Plant Soil</i> 331(1-2), 361-375(2010)

523	山脇 正人, 廣瀬 農, 菅野 里美, 石橋 弘規, 野田 章彦, 田野井 慶太郎, 中西 友子. 植物の研究のためのリアルタイムRIイメージングシステムの ¹⁰⁹ Cd検出性能評価. <i>RADIOISOTOPES</i> 59(3), 155-162(2010)
524	Y. Abe, Kitano H., Ando T., Kono I., Yano M., Iwasaki Y. Mapping of Small and Round Seed 2 (SRS2) gene in rice. <i>Rice Genet. Newsl.</i> 25, 30-31(2010)
525	N. Yamaji, C. F. Huang, S. Nagao, M. Yano, Y. Sato, Y. Nagamura, J. F. Ma. A zinc finger transcription factor ART1 regulates multiple genes implicated in aluminum tolerance in rice. <i>Plant Cell</i> 21(10), 3339-3349(2009)
526	M. Endo, T. Tsuchiya, K. Hamada, S. Kawamura, K. Yano, M. Ohshima, A. Higashitani, M. Watanabe, M. Kawagishi-Kobayashi. High Temperatures Cause Male Sterility in Rice Plants with Transcriptional Alterations during Pollen Development. <i>Plant Cell Physiol</i> 50(11), 1911-1922(2009)
527	T. Kamiya, T. Fujiwara. Arabidopsis NIP1;1 Transports Antimonite and Determines the Antimonite Sensitivity. <i>Plant Cell Physiol</i> 50(11), 1977-1981(2009)
528	A. Kitajima, S. Asatsuma, H. Okada, Y. Hamada, K. Kaneko, Y. Nanjo, Y. Kawagoe, K. Toyooka, K. Matsuoka, M. Takeuchi, A. Nakano, T. Mitsui. The Rice [alpha]-Amylase Glycoprotein Is Targeted from the Golgi Apparatus through the Secretory Pathway to the Plastids. <i>Plant Cell</i> 21(9), 2844-2858(2009)
529	N. Yamaji, J. F. Ma. A Transporter at the Node Responsible for Intervascular Transfer of Silicon in Rice. <i>Plant Cell</i> 21(9), 2878-2883(2009)
530	M. Kawai-Yamada, Z. Hori, T. Ogawa, Y. Ihara-Ohori, K. Tamura, M. Nagano, T. Ishikawa, H. Uchimiya. Loss of calmodulin binding to Bax Inhibitor-1 affects Pseudomonas-mediated hypersensitive response-associated cell death in Arabidopsis thaliana. <i>J Biol Chem</i> 284(41), 27998-28003(2009)
531	Y. Onda, T. Kumamaru, Y. Kawagoe. ER membrane-localized oxidoreductase Ero1 is required for disulfide bond formation in the rice endosperm. <i>Proc Natl Acad Sci U S A</i> 106(33), 14156-14161(2009)
532	Katsuyuki Oki, Kanako Kitagawa, Yukiko Fujisawa, Hisaharu Kato, Yukimoto Iwasaki. Function of alpha subunit of heterotrimeric G protein in brassinosteroid response of rice plants. <i>Plant Signal Behav</i> 4(2), 126-128(2009)
533	Y. Ito-Inaba, M. Sato, H. Masuko, Y. Hida, K. Toyooka, M. Watanabe, T. Inaba. Developmental changes and organelle biogenesis in the reproductive organs of thermogenic skunk cabbage (<i>Symplocarpus renifolius</i>). <i>J Exp Bot</i> 60(13), 3909-3922(2009)
534	I. Kasajima, K. Takahara, M. Kawai-Yamada, H. Uchimiya. Estimation of the relative sizes of rate constants for chlorophyll de-excitation processes through comparison of inverse fluorescence intensities. <i>Plant Cell Physiol</i> 50(9), 1600-1616(2009)
535	H. Takahashi, K. Takahara, S. Hashida, T. Hirabayashi, T. Fujimori, M. Kawai-Yamada, T. Yamaya, S. Yanagisawa, H. Uchimiya. Pleiotropic modulation of carbon and nitrogen metabolism in Arabidopsis plants overexpressing the NAD kinase2 gene. <i>Plant Physiol</i> 151(1), 100-113(2009)
536	N. Mitani, Y. Chiba, N. Yamaji, J. F. Ma. Identification and characterization of maize and barley Lsi2-like silicon efflux transporters reveals a distinct silicon uptake system from that in rice. <i>Plant Cell</i> 21(7), 2133-2142(2009)
537	R. Y. Li, Y. Ago, W. J. Liu, N. Mitani, J. Feldmann, S. P. McGrath, J. F. Ma, F. J. Zhao. The rice aquaporin Lsi1 mediates uptake of methylated arsenic species. <i>Plant Physiol</i> 150(4), 2071-2080(2009)
538	T. Arite, M. Umehara, S. Ishikawa, A. Hanada, M. Maekawa, S. Yamaguchi, J. Kyojuka. d14, a strigolactone-insensitive mutant of rice, shows an accelerated outgrowth of tillers. <i>Plant Cell Physiol</i> 50(8), 1416-1424(2009)
539	K. Oki, N. Inaba, H. Kitano, S. Takahashi, Y. Fujisawa, H. Kato, Y. Iwasaki. Study of novel d1 alleles, defective mutants of the alpha subunit of heterotrimeric G-protein in rice. <i>Genes Genet Syst</i> 84(1), 35-42(2009)
540	K. Ikeda-Kawakatsu, N. Yasuno, T. Oikawa, S. Iida, Y. Nagato, M. Maekawa, J. Kyojuka. Expression level of ABERRANT PANICLE ORGANIZATION1 determines rice inflorescence form through control of cell proliferation in the meristem. <i>Plant Physiol</i> 150(2), 736-747(2009)
541	K. Toyooka, Y. Goto, S. Asatsuma, M. Koizumi, T. Mitsui, K. Matsuoka. A mobile secretory vesicle cluster involved in mass transport from the Golgi to the plant cell exterior. <i>Plant Cell</i> 21(4), 1212-1229(2009)

542	C. F. Huang, N. Yamaji, M. Nishimura, S. Tajima, J. F. Ma. A rice mutant sensitive to Al toxicity is defective in the specification of root outer cell layers. <i>Plant Cell Physiol</i> 50(5), 976–985(2009)
543	C. F. Huang, N. Yamaji, N. Mitani, M. Yano, Y. Nagamura, J. F. Ma. A bacterial-type ABC transporter is involved in aluminum tolerance in rice. <i>Plant Cell</i> 21(2), 655–667(2009)
544	M. Fujimoto, S. Arimura, S. Mano, M. Kondo, C. Saito, T. Ueda, M. Nakazono, A. Nakano, M. Nishimura, N. Tsutsumi. Arabidopsis dynamin-related proteins DRP3A and DRP3B are functionally redundant in mitochondrial fission, but have distinct roles in peroxisomal fission. <i>Plant J</i> 58(3), 388–400(2009)
545	S. Okamoto, E. Ohnishi, S. Sato, H. Takahashi, M. Nakazono, S. Tabata, M. Kawaguchi. Nod factor/nitrate-induced CLE genes that drive HAR1-mediated systemic regulation of nodulation. <i>Plant Cell Physiol</i> 50(1), 67–77(2009)
546	M Nagano, Y Ihara–Otori, H Imai, N Inada, M Fujimoto, N Tsutsumi, H. Uchimihya, M. Kawai–Yamada. Functional association of cell death suppressor, Arabidopsis Bax Inhibitor–1, with fatty acid 2–hydroxylation through cytochrome b5. <i>Plant J</i> 58(1), 122–134(2009)
547	H. Inoue, T. Kobayashi, T. Nozoye, M. Takahashi, Y. Kakei, K. Suzuki, M. Nakazono, H. Nakanishi, S. Mori, N. K. Nishizawa. Rice OsYSL15 is an iron-regulated iron(III)–deoxymugineic acid transporter expressed in the roots and is essential for iron uptake in early growth of the seedlings. <i>J Biol Chem</i> 284(6), 3470–3479(2009)
548	K. Oki, N. Inaba, K. Kitagawa, S. Fujioka, H. Kitano, Y. Fujisawa, H. Kato, Y. Iwasaki. Function of the alpha subunit of rice heterotrimeric G protein in brassinosteroid signaling. <i>Plant Cell Physiol</i> 50(1), 161–172(2009)
549	T. Kamiya, M. Tanaka, N. Mitani, J. F. Ma, M. Maeshima, T. Fujiwara. NIP1;1, an aquaporin homolog, determines the arsenite sensitivity of Arabidopsis thaliana. <i>J Biol Chem</i> 284(4), 2114–2120(2009)
550	Y. Kato, K. Miwa, J. Takano, M. Wada, T. Fujiwara. Highly boron deficiency-tolerant plants generated by enhanced expression of NIP5;1, a boric acid channel. <i>Plant Cell Physiol</i> 50(1), 58–66(2009)
551	T. Ishimaru, A.K. Horigane, M. Ida, N. Iwasawa, Y.A. San-oh, M. Nakazono, N.K. Nishizawa, T. Masumura, M. Kondo, M. Yoshida. Formation of grain chalkiness and changes in water distribution in developing rice caryopses grown under high-temperature stress. <i>J Cereal Sci</i> 50(2), 166–174(2009)
552	Ito S, Hara T, Kawanami Y, Watanabe T, Thiraporn K, Ohtake N, Sueyoshi K, Mitsui T, Fukuyama T, Takahashi Y, Sato T, Sato A, Ohyama T. Carbon and Nitrogen Transport during Grain Filling in Rice Under High-temperature Conditions. <i>J. Agron. Crop Sci.</i> 195(5), 368–376(2009)
553	M Yamawaki, S Kanno, H Ishibashi, A Noda, A Hirose, K Tanoi, T. M Nakanishi. The development of real-time RI imaging system for plant under light environment. <i>JOURNAL OF RADIOANALYTICAL AND NUCLEAR CHEMISTRY</i> 282(1), 275–279(2009)
554	Hiroaki Shimada, Haruka Takeda, Akina Tsunoda, Tadamasasa Sasaki. Regulatory mechanism on sucrose metabolism in rice endosperm by a specific protein kinase, SPK. <i>The Proceedings of the International Plant Nutrition Colloquium XVI</i> 16, 1117–1123(2009)
555	F. J. Zhao, J. F. Ma, A. A. Meharg, S. P. McGrath. Arsenic uptake and metabolism in plants. <i>New Phytol</i> 181(4), 777–794(2008)
556	M. Tanaka, I. S. Wallace, J. Takano, D. M. Roberts, T. Fujiwara. NIP6;1 is a boric acid channel for preferential transport of boron to growing shoot tissues in Arabidopsis. <i>Plant Cell</i> 20(10), 2860–2875(2008)
557	M. Watanabe. Towards a comprehensive understanding of molecular mechanisms of sexual reproduction in higher plants. <i>Plant Cell Physiol</i> 49(10), 1404–1406(2008)
558	F. J. Munoz, E. Baroja-Fernandez, M. Ovecka, J. Li, T. Mitsui, M. T. Sesma, M. Montero, A. Bahaji, I. Ezquer, J. Pozueta-Romero. Plastidial localization of a potato 'Nudix' hydrolase of ADP-glucose linked to starch biosynthesis. <i>Plant Cell Physiol</i> 49(11), 1734–1746(2008)
559	T Hobo, K Suwabe, K Aya, G Suzuki, K Yano, T Ishimizu, Kikuchi Fujita M, S, K Hamada, M Miyano, T Fujioka, F Kaneko, T Kazama, Y Mizuta, H Takahashi, K Shiono, M Nakazono, N Tsutsumi, Y Nagamura, N Kurata, M Watanabe, M. Matsuoka. Various spatiotemporal expression profiles of anther-expressed genes in rice. <i>Plant Cell Physiol</i> 49(10), 1417–1428(2008)

560	K Suwabe, G Suzuki, H Takahashi, K Shiono, M Endo, Y Yano, M Fujita, M Masuko, H Saito, T Fujioka, F Kaneko, T Kazama, Y Mizuta, Tsutsumi Kawagishi-Kobayashi, N, N Kurata, M Nakazono, M. Watanabe. Separated transcriptomes of male gametophyte and tapetum in rice: validity of a laser microdissection (LM) microarray. <i>Plant Cell Physiol</i> 49(10), 1407–1416(2008)
561	M. Umehara, A. Hanada, S. Yoshida, K. Akiyama, T. Arite, N. Takeda-Kamiya, H. Magome, Y. Kamiya, K. Shirasu, K. Yoneyama, J. Kyojuka, S. Yamaguchi. Inhibition of shoot branching by new terpenoid plant hormones. <i>Nature</i> 455(7210), 195–200(2008)
562	T Fujioka, F. Kaneko, , Kazama, T, K Suwabe, G Suzuki, A Makino, M Endo, M Kawagishi-Kobayashi, M. Watanabe. Identification of small RNAs in late developmental stage of rice anthers. <i>Genes Genet Syst</i> 83(3), 281–284(2008)
563	A. Shimizu, K. Kato, A. Komatsu, K. Motomura, H. Iekashi. Genetic analysis of root elongation induced by phosphorus deficiency in rice (<i>Oryza sativa</i> L.): fine QTL mapping and multivariate analysis of related traits. <i>Theor Appl Genet</i> 117(6), 987–996(2008)
564	J. F. Ma, N. Yamaji, N. Mitani, X. Y. Xu, Y. H. Su, S. P. McGrath, F. J. Zhao. Transporters of arsenite in rice and their role in arsenic accumulation in rice grain. <i>Proc Natl Acad Sci U S A</i> 105(29), 9931–9935(2008)
565	A. Endo, Y. Sawada, H. Takahashi, M. Okamoto, K. Ikegami, H. Koiwai, M. Seo, T. Toyomasu, W. Mitsushashi, K. Shinozaki, M. Nakazono, Y. Kamiya, T. Koshihara, E. Nambara. Drought induction of Arabidopsis 9-cis-epoxycarotenoid dioxygenase occurs in vascular parenchyma cells. <i>Plant Physiol</i> 147(4), 1984–1993(2008)
566	N. Yamaji, N. Mitatni, J. F. Ma. A transporter regulating silicon distribution in rice shoots. <i>Plant Cell</i> 20(5), 1381–1389(2008)
567	T. Kazama, T. Nakamura, M. Watanabe, M. Sugita, K. Toriyama. Suppression mechanism of mitochondrial ORF79 accumulation by Rf1 protein in BT-type cytoplasmic male sterile rice. <i>Plant J</i> 55(4), 619–628(2008)
568	J. Negi, O. Matsuda, T. Nagasawa, Y. Oba, H. Takahashi, M. Kawai-Yamada, H. Uchimiya, M. Hashimoto, K. Iba. CO2 regulator SLAC1 and its homologues are essential for anion homeostasis in plant cells. <i>Nature</i> 452(7186), 483–486(2008)
569	T. Kudo, A. Kawai, T. Yamaya, T. Hayakawa. Cellular distribution of ACT domain repeat protein 9, a nuclear localizing protein, in rice (<i>Oryza sativa</i>). <i>Physiol Plant</i> 133(2), 167–179(2008)
570	N. Mitani, N. Yamaji, J. F. Ma. Characterization of substrate specificity of a rice silicon transporter, Lsi1. <i>Pflugers Arch</i> 456(4), 679–686(2008)
571	Hiromoto Yamakawa, Takeshi Ebitani, Tomio Terao. Comparison between locations of QTLs for grain chalkiness and genes responsive to high temperature during grain filling on the rice chromosome map. <i>Breed. Sci.</i> 58(3), 337–343(2008)
572	T. Sakata, A. Higashitani. Male sterility accompanied with abnormal anther developmental in plants—Genes and environmental stresses with special reference to high temperature injury. <i>Intl. J. Plant Dev. Biol.</i> 2(1), 42–51(2008)
573	H Takahashi, H Matsumura, M. Kawai-Yamada, H. Uchimiya. The cell death factor, cell wall elicitor of rice blast fungus (<i>Magnaporthe grisea</i>) causes metabolic alterations including GABA shunt in rice cultured cells. <i>Plant Signal Behav</i> 3(11), 945–953(2008)
574	K. Tamai, J. F. Ma. Reexamination of silicon effects on rice growth and production under field conditions using a low silicon mutant. <i>Plant Soil</i> 307(1–2), 21–27(2008)
575	K. Shiono, S. Ogawa, S. Yamazaki, H. Isoda, T. Fujimura, M. Nakazono, T. D. Colmer. Contrasting dynamics of radial O2-loss barrier induction and aerenchyma formation in rice roots of two lengths. <i>Ann Bot</i> 107(1), 89–99(2010)
576	K. Tanaka, N. Umeki, T. Mitsui, Z. Fujimoto, S. Maruta. Crystallographic analysis reveals a unique conformation of the ADP-bound novel rice kinesin K16. <i>Biochem Biophys Res Commun</i> 401(2), 251–256(2010)
577	S. Oda, F. Kaneko, K. Yano, T. Fujioka, H. Masuko, J. I. Park, S. Kikuchi, K. Hamada, M. Endo, K. Nagano, Y. Nagamura, M. Kawagishi-Kobayashi, K. Suwabe, G. Suzuki, M. Watanabe. Morphological and gene expression analysis under cool temperature conditions in rice anther development. <i>Genes Genet Syst</i> 85(2), 107–120(2010)
578	T. Ishimizu, H. Kodama, T. Ando, M. Watanabe. Molecular evidence that most RNAs required for germination and pollen tube growth are stored in the mature pollen grain in petunia. <i>Genes Genet Syst</i> 85(4), 259–263(2010)

579	S. N. Hashida, T. Itami, H. Takahashi, K. Takahara, M. Nagano, M. Kawai-Yamada, K. Shoji, F. Goto, T. Yoshihara, H. Uchimiya. Nicotinate/nicotinamide mononucleotide adenylyltransferase-mediated regulation of NAD biosynthesis protects guard cells from reactive oxygen species in ABA-mediated stomatal movement in Arabidopsis. <i>J Exp Bot</i> 61(13), 3813–3825(2010)
580	H. Takahashi, H. Kamakura, Y. Sato, K. Shiono, T. Abiko, N. Tsutsumi, Y. Nagamura, N. K. Nishizawa, M. Nakazono. A method for obtaining high quality RNA from paraffin sections of plant tissues by laser microdissection. <i>J Plant Res</i> 123(6), 807–813(2010)
581	F. J. Zhao, Y. Ago, N. Mitani, R. Y. Li, Y. H. Su, N. Yamaji, S. P. McGrath, J. F. Ma. The role of the rice aquaporin Lsi1 in arsenite efflux from roots. <i>New Phytol</i> 186(2), 392–399(2010)
582	I. Rajhi, T. Yamauchi, H. Takahashi, S. Nishiuchi, K. Shiono, R. Watanabe, A. Mliki, Y. Nagamura, N. Tsutsumi, N. K. Nishizawa, M. Nakazono. Identification of genes expressed in maize root cortical cells during lysigenous aerenchyma formation using laser microdissection and microarray analyses. <i>New Phytol</i> 190(2), 351–368(2011)
583	T. Oikawa, J. Kyojuka. Two-Step Regulation of LAX PANICLE1 Protein Accumulation in Axillary Meristem Formation in Rice. <i>Plant Cell</i> 21(4), 1095–1108(2009)
584	K. C. She, H. Kusano, K. Koizumi, H. Yamakawa, M. Hakata, T. Imamura, M. Fukuda, N. Naito, Y. Tsurumaki, M. Yaeshima, T. Tsuge, K. Matsumoto, M. Kudoh, E. Itoh, S. Kikuchi, N. Kishimoto, J. Yazaki, T. Ando, M. Yano, T. Aoyama, T. Sasaki, H. Satoh, H. Shimada. A novel factor FLOURY ENDOSPERM2 is involved in regulation of rice grain size and starch quality. <i>Plant Cell</i> 22(10), 3280–3294(2010)
585	T. Oshino, S. Miura, S. Kikuchi, K. Hamada, K. Yano, M. Watanabe, A. Higashitani. Auxin depletion in barley plants under high-temperature conditions represses DNA proliferation in organelles and nuclei via transcriptional alterations. <i>Plant Cell Environ</i> 34(2), 284–290(2011)
586	J. I. Park, T. Ishimizu, K. Suwabe, K. Sudo, H. Masuko, H. Hakozaiki, I. S. Nou, G. Suzuki, M. Watanabe. UDP-glucose pyrophosphorylase is rate limiting in vegetative and reproductive phases in Arabidopsis thaliana. <i>Plant Cell Physiol</i> 51(6), 981–996(2010)
587	Z. Hu, H. Yan, J. Yang, S. Yamaguchi, M. Maekawa, I. Takamura, N. Tsutsumi, J. Kyojuka, M. Nakazono. Strigolactones negatively regulate mesocotyl elongation in rice during germination and growth in darkness. <i>Plant Cell Physiol</i> 51(7), 1136–1142(2010)
588	M. S. Yun, Y. Kawagoe. Septum formation in amyloplasts produces compound granules in the rice endosperm and is regulated by plastid division proteins. <i>Plant Cell Physiol</i> 51(9), 1469–1479(2010)
589	K. Kitagawa, S. Kurinami, K. Oki, Y. Abe, T. Ando, I. Kono, M. Yano, H. Kitano, Y. Iwasaki. A novel kinesin 13 protein regulating rice seed length. <i>Plant Cell Physiol</i> 51(8), 1315–1329(2010)
590	S. Ito, N. Kitahata, M. Umehara, A. Hanada, A. Kato, K. Ueno, K. Mashiguchi, J. Kyojuka, K. Yoneyama, S. Yamaguchi, T. Asami. A new lead chemical for strigolactone biosynthesis inhibitors. <i>Plant Cell Physiol</i> 51(7), 1143–1150(2010)
591	T. Ishikawa, K. Takahara, T. Hirabayashi, H. Matsumura, S. Fujisawa, R. Terauchi, H. Uchimiya, M. Kawai-Yamada. Metabolome analysis of response to oxidative stress in rice suspension cells overexpressing cell death suppressor Bax inhibitor-1. <i>Plant Cell Physiol</i> 51(1), 9–20(2010)
592	Y. Izawa, Y. Takayanagi, N. Inaba, Y. Abe, M. Minami, Y. Fujisawa, H. Kato, S. Ohki, H. Kitano, Y. Iwasaki. Function and expression pattern of the alpha subunit of the heterotrimeric G protein in rice. <i>Plant Cell Physiol</i> 51(2), 271–281(2010)
593	H. Yamakawa, M. Hakata. Atlas of rice grain filling-related metabolism under high temperature: joint analysis of metabolome and transcriptome demonstrated inhibition of starch accumulation and induction of amino acid accumulation. <i>Plant Cell Physiol</i> 51(5), 795–809(2010)
594	K. Takahara, I. Kasajima, H. Takahashi, S. N. Hashida, T. Itami, H. Onodera, S. Toki, S. Yanagisawa, M. Kawai-Yamada, H. Uchimiya. Metabolome and photochemical analysis of rice plants overexpressing Arabidopsis NAD kinase gene. <i>Plant Physiol</i> 152(4), 1863–1873(2010)
595	C. F. Huang, N. Yamaji, J. F. Ma. Knockout of a bacterial-type ATP-binding cassette transporter gene, AtSTAR1, results in increased aluminum sensitivity in Arabidopsis. <i>Plant Physiol</i> 153(4), 1669–1677(2010)
596	X. Q. Zhao, N. Mitani, N. Yamaji, R. F. Shen, J. F. Ma. Involvement of silicon influx transporter OsNIP2;1 in selenite uptake in rice. <i>Plant Physiol</i> 153(4), 1871–1877(2010)

597	T. Sakata, N. Yagihashi, A. Higashitani. Tissue-specific auxin signaling in response to temperature fluctuation. <i>Plant Signal Behav</i> 5(11), 1510–1512(2010)
598	K. Nonomura, M. Eiguchi, M. Nakano, K. Takashima, N. Komeda, S. Fukuchi, S. Miyazaki, A. Miyao, H. Hirochika, N. Kurata. A novel RNA-recognition-motif protein is required for premeiotic G1/S-phase transition in rice (<i>Oryza sativa</i> L.). <i>PLoS Genet</i> 7(1), e1001265(2011)
599	J. Xia, N. Yamaji, T. Kasai, J. F. Ma. Plasma membrane-localized transporter for aluminum in rice. <i>Proc Natl Acad Sci U S A</i> 107(43), 18381–18385(2010)
600	K. Sugimoto, Y. Takeuchi, K. Ebana, A. Miyao, H. Hirochika, N. Hara, K. Ishiyama, M. Kobayashi, Y. Ban, T. Hattori, M. Yano. Molecular cloning of <i>Sdr4</i> , a regulator involved in seed dormancy and domestication of rice. <i>Proc Natl Acad Sci U S A</i> 107(13), 5792–5797(2010)
601	T. Sakata, T. Oshino, S. Miura, M. Tomabechei, Y. Tsunaga, N. Higashitani, Y. Miyazawa, H. Takahashi, M. Watanabe, A. Higashitani. Auxins reverse plant male sterility caused by high temperatures. <i>Proc Natl Acad Sci U S A</i> 107(19), 8569–8574(2010)
602	M. Obara, W. Tamura, T. Ebitani, M. Yano, T. Sato, T. Yamaya. Fine-mapping of <i>qRL6.1</i> , a major QTL for root length of rice seedlings grown under a wide range of NH_4^+ concentrations in hydroponic conditions. <i>Theor Appl Genet</i> 121(3), 535–547(2010)
603	K. Hori, K. Sugimoto, Y. Nonoue, N. Ono, K. Matsubara, U. Yamanouchi, A. Abe, Y. Takeuchi, M. Yano. Detection of quantitative trait loci controlling pre-harvest sprouting resistance by using backcrossed populations of japonica rice cultivars. <i>Theor Appl Genet</i> 120(8), 1547–1557(2010)
604	H. Itoh, Y. Nonoue, M. Yano, T. Izawa. A pair of floral regulators sets critical day length for <i>Hd3a</i> florigen expression in rice. <i>Nat Genet</i> 42(7), 635–638(2010)
605	A. J. Nagano, Y. Sato, M. Mihara, B. A. Antonio, R. Motoyama, H. Itoh, Y. Nagamura, T. Izawa. Deciphering and Prediction of Transcriptome Dynamics under Fluctuating Field Conditions. <i>Cell</i> 151(6), 1358–1369(2012)
606	L. Peng, Y. Fukao, F. Myouga, R. Motohashi, K. Shinozaki, T. Shikanai. A chaperonin subunit with unique structures is essential for folding of a specific substrate. <i>PLoS Biol</i> 9(4), e1001040(2011)
607	W. Tanaka, T. Toriba, Y. Ohmori, A. Yoshida, A. Kawai, T. Mayama-Tsuchida, H. Ichikawa, N. Mitsuda, M. Ohme-Takagi, H. Y. Hirano. The YABBY gene <i>TONGARI-BOUSHI1</i> is involved in lateral organ development and maintenance of meristem organization in the rice spikelet. <i>Plant Cell</i> 24(1), 80–95(2012)
608	L. Peng, Y. Fukao, M. Fujiwara, T. Shikanai. Multistep Assembly of Chloroplast NADH Dehydrogenase-Like Subcomplex A Requires Several Nucleus-Encoded Proteins, Including <i>CRR41</i> and <i>CRR42</i> , in <i>Arabidopsis</i> . <i>Plant Cell</i> 24(1), 202–214(2012)
609	N. Tanaka, H. Itoh, N. Sentoku, M. Kojima, H. Sakakibara, T. Izawa, J. Itoh, Y. Nagato. The COP1 ortholog PPS regulates the juvenile-adult and vegetative-reproductive phase changes in rice. <i>Plant Cell</i> 23(6), 2143–2154(2011)
610	T. Izawa, M. Mihara, Y. Suzuki, M. Gupta, H. Itoh, A. J. Nagano, R. Motoyama, Y. Sawada, M. Yano, M. Y. Hirai, A. Makino, Y. Nagamura. <i>Os-GIGANTEA</i> confers robust diurnal rhythms on the global transcriptome of rice in the field. <i>Plant Cell</i> 23(5), 1741–1755(2011)
611	H. Yamamoto, L. Peng, Y. Fukao, T. Shikanai. An Src homology 3 domain-like fold protein forms a ferredoxin binding site for the chloroplast NADH dehydrogenase-like complex in <i>Arabidopsis</i> . <i>Plant Cell</i> 23(4), 1480–1493(2011)
612	L. Peng, Y. Fukao, M. Fujiwara, T. Takami, T. Shikanai. Efficient operation of NAD(P)H dehydrogenase requires supercomplex formation with photosystem I via minor LHCI in <i>Arabidopsis</i> . <i>Plant Cell</i> 21(11), 3623–3640(2009)
613	K. Okuda, A. L. Chateigner-Boutin, T. Nakamura, E. Delannoy, M. Sugita, F. Myouga, R. Motohashi, K. Shinozaki, I. Small, T. Shikanai. Pentatricopeptide repeat proteins with the DYW motif have distinct molecular functions in RNA editing and RNA cleavage in <i>Arabidopsis</i> chloroplasts. <i>Plant Cell</i> 21(1), 146–156(2009)
614	H. Yamasaki, M. Hayashi, M. Fukazawa, Y. Kobayashi, T. Shikanai. <i>SQUAMOSA</i> Promoter Binding Protein-Like7 Is a Central Regulator for Copper Homeostasis in <i>Arabidopsis</i> . <i>Plant Cell</i> 21(1), 347–361(2009)

615	T. Suzuki, A. Yoshida, H. Y. Hirano. Functional diversification of CLAVATA3-related CLE proteins in meristem maintenance in rice. <i>Plant Cell</i> 20(8), 2049–2058(2008)
616	K. Hammani, K. Okuda, S. K. Tanz, A. L. Chateigner-Boutin, T. Shikanai, I. Small. A study of new Arabidopsis chloroplast RNA editing mutants reveals general features of editing factors and their target sites. <i>Plant Cell</i> 21(11), 3686–3699(2009)
617	T. Toriba, T. Suzuki, T. Yamaguchi, Y. Ohmori, H. Tsukaya, H. Y. Hirano. Distinct regulation of adaxial–abaxial polarity in anther patterning in rice. <i>Plant Cell</i> 22(5), 1452–1462(2010)
618	C. Masumoto, S. Miyazawa, H. Ohkawa, T. Fukuda, Y. Taniguchi, S. Murayama, M. Kusano, K. Saito, H. Fukayama, M. Miyao. Phosphoenolpyruvate carboxylase intrinsically located in the chloroplast of rice plays a crucial role in ammonium assimilation. <i>Proc Natl Acad Sci U S A</i> 107(11), 5226–5231(2010)
619	A Yoshida, T. Suzuki, W. Tanaka, H.-Y. Hirano. The homeotic gene LONG STERILE LEMMA (G1) specifies sterile lemma identity in the rice spikelet. <i>Proc Natl Acad Sci U S A</i> 106(47), 20103–20108(2009)
620	M. Takano, N. Inagaki, X. Xie, S. Kiyota, A. Baba-Kasai, T. Tanabata, T. Shinomura. Phytochromes are the sole photoreceptors for perceiving red/far-red light in rice. <i>Proc Natl Acad Sci U S A</i> 106(34), 14705–14710(2009)
621	L. Peng, H. Shimizu, T. Shikanai. The chloroplast NAD(P)H dehydrogenase complex interacts with photosystem I in Arabidopsis. <i>J Biol Chem</i> 283(50), 34873–34879(2008)
622	T. Suzuki, M. Ohneda, T. Toriba, A. Yoshida, H. Y. Hirano. FON2 SPARE1 redundantly regulates floral meristem maintenance with FLORAL ORGAN NUMBER2 in rice. <i>PLoS Genet</i> 5(10), e1000693(2009)
623	K. Okuda, Y. Habata, Y. Kobayashi, T. Shikanai. Amino acid sequence variations in Nicotiana CRR4 orthologs determine the species-specific efficiency of RNA editing in plastids. <i>Nucleic Acids Res</i> 36(19), 6155–6164(2008)
624	M. Ueda, T. Kuniyoshi, H. Yamamoto, K. Sugimoto, K. Ishizaki, T. Kohchi, Y. Nishimura, T. Shikanai. Composition and physiological function of the chloroplast NADH dehydrogenase-like complex in <i>Marchantia polymorpha</i> . <i>Plant J</i> 72(4), 683–693(2012)
625	W. Yamori, C. Masumoto, H. Fukayama, A. Makino. Rubisco activase is a key regulator of non steady-state photosynthesis at any leaf temperature and, to a lesser extent, of steady-state photosynthesis at high temperature. <i>Plant J</i> 71(6), 871–880(2012)
626	H. Shikata, M. Shibata, T. Ushijima, M. Nakashima, S. G. Kong, K. Matsuoka, C. Lin, T. Matsushita. The RS domain of Arabidopsis splicing factor RRC1 is required for phytochrome B signal transduction. <i>Plant J</i> 70(5), 727–738(2012)
627	A. Yoshida, Y. Ohmori, H. Kitano, F. Taguchi-Shiobara, H. Y. Hirano. Aberrant spikelet and panicle1, encoding a TOPLESS-related transcriptional co-repressor, is involved in the regulation of meristem fate in rice. <i>Plant J</i> 70(2), 327–339(2012)
628	K. Ikeda-Kawakatsu, M. Maekawa, T. Izawa, J. I. Itoh, Y. Nagato. ABERRANT PANICLE ORGANIZATION 2/RFL, the rice ortholog of Arabidopsis LEAFY, suppresses the transition from inflorescence meristem to floral meristem through interaction with APO1. <i>Plant J</i> 69(1), 168–180(2012)
629	W. Yamori, N. Sakata, Y. Suzuki, T. Shikanai, A. Makino. Cyclic electron flow around photosystem I via chloroplast NAD(P)H dehydrogenase (NDH) complex performs a significant physiological role during photosynthesis and plant growth at low temperature in rice. <i>Plant J</i> 68(6), 966–976(2011)
630	W. Cai, K. Okuda, L. Peng, T. Shikanai. PROTON GRADIENT REGULATION 3 recognizes multiple targets with limited similarity and mediates translation and RNA stabilization in plastids. <i>Plant J</i> 67(2), 318–327(2011)
631	K. Matsubara, U. Yamanouchi, Y. Nonoue, K. Sugimoto, Z. X. Wang, Y. Minobe, M. Yano. Ehd3, encoding a plant homeodomain finger-containing protein, is a critical promoter of rice flowering. <i>Plant J</i> 66(4), 603–612(2011)
632	A. Takabayashi, N. Ishikawa, T. Obayashi, S. Ishida, J. Obokata, T. Endo, F. Sato. Three novel subunits of Arabidopsis chloroplastic NAD(P)H dehydrogenase identified by bioinformatic and reverse genetic approaches. <i>Plant J</i> 57(2), 207–219(2009)

633	Y. Ohmori, T. Toriba, H. Nakamura, H. Ichikawa, H. Y. Hirano. Temporal and spatial regulation of DROOPING LEAF gene expression that promotes midrib formation in rice. <i>Plant J</i> 65(1), 77–86(2011)
634	Y. Okegawa, Y. Kobayashi, T. Shikanai. Physiological links among alternative electron transport pathways that reduce and oxidize plastoquinone in <i>Arabidopsis</i> . <i>Plant J</i> 63(3), 458–468(2010)
635	K. Okuda, K. Hammani, S. K. Tanz, L. Peng, Y. Fukao, F. Myouga, R. Motohashi, K. Shinozaki, I. Small, T. Shikanai. The pentatricopeptide repeat protein OTP82 is required for RNA editing of plastid <i>ndhB</i> and <i>ndhG</i> transcripts. <i>Plant J</i> 61(2), 339–349(2010)
636	L. Peng, W. Cai, T. Shikanai. Chloroplast stromal proteins, CRR6 and CRR7, are required for assembly of the NAD(P)H dehydrogenase subcomplex A in <i>Arabidopsis</i> . <i>Plant J</i> 63(2), 203–211(2010)
637	Y. Suzuki, A. Makino. Availability of Rubisco small subunit up-regulates the transcript levels of large subunit for stoichiometric assembly of its holoenzyme in rice. <i>Plant Physiol</i> 160(1), 533–540(2012)
638	Y. Oka, S. G. Kong, T. Matsushita. A Non-Covalently Attached Chromophore can Mediate Phytochrome B Signaling in <i>Arabidopsis</i> . <i>Plant Cell Physiol</i> 52(12), 2088–2102(2011)
639	A. Osugi, H. Itoh, K. Ikeda-Kawakatsu, M. Takano, T. Izawa. Molecular dissection of the roles of phytochrome in photoperiodic flowering in rice. <i>Plant Physiol</i> 157(3), 1128–1137(2011)
640	L. Peng, T. Shikanai. Supercomplex formation with photosystem I is required for the stabilization of the chloroplast NADH dehydrogenase-like complex in <i>Arabidopsis</i> . <i>Plant Physiol</i> 155(4), 1629–1639(2011)
641	W. Yamori, S. Takahashi, A. Makino, G. D. Price, M. R. Badger, S. von Caemmerer. The roles of ATP synthase and the cytochrome b6/f complexes in limiting chloroplast electron transport and determining photosynthetic capacity. <i>Plant Physiol</i> 155(2), 956–962(2011)
642	S. Wada, H. Ishida, M. Izumi, K. Yoshimoto, Y. Ohsumi, T. Mae, A. Makino. Autophagy plays a role in chloroplast degradation during senescence in individually darkened leaves. <i>Plant Physiol</i> 149(2), 885–893(2009)
643	K. Matsubara, U. Yamanouchi, Z. X. Wang, Y. Minobe, T. Izawa, M. Yano. Ehd2, a Rice Ortholog of the Maize INDETERMINATE1 Gene, Promotes Flowering by Up-Regulating Ehd1. <i>Plant Physiol</i> 148(3), 1425–1435(2008)
644	E. Ogiso, Y. Takahashi, T. Sasaki, M. Yano, T. Izawa. The role of casein kinase II in flowering time regulation has diversified during evolution. <i>Plant Physiol</i> 152(2), 808–820(2010)
645	A. Makino. Photosynthesis, grain yield, and nitrogen utilization in rice and wheat. <i>Plant Physiol</i> 155(1), 125–129(2011)
646	M. Izumi, S. Wada, A. Makino, H. Ishida. The autophagic degradation of chloroplasts via rubisco-containing bodies is specifically linked to leaf carbon status but not nitrogen status in <i>Arabidopsis</i> . <i>Plant Physiol</i> 154(3), 1196–1209(2010)
647	H. Matsumura, E. Mizohata, H. Ishida, A. Kogami, T. Ueno, A. Makino, T. Inoue, A. Yokota, T. Mae, Y. Kai. Crystal structure of rice Rubisco and implications for activation induced by positive effectors NADPH and 6-phosphogluconate. <i>J Mol Biol</i> 422(1), 75–86(2012)
648	Y. Suzuki, T. Fujimori, K. Kanno, A. Sasaki, Y. Ohashi, A. Makino. Metabolome analysis of photosynthesis and the related primary metabolites in the leaves of transgenic rice plants with increased or decreased Rubisco content. <i>Plant Cell Environ</i> 35(8), 1369–1379(2012)
649	S. Ogawa, Y. Suzuki, R. Yoshizawa, K. Kanno, A. Makino. Effect of individual suppression of RBCS multigene family on Rubisco contents in rice leaves. <i>Plant Cell Environ</i> 35(3), 546–553(2012)
650	R. Saito, H. Yamamoto, A. Makino, T. Sugimoto, C. Miyake. Methylglyoxal functions as Hill oxidant and stimulates the photoreduction of O ₂ at photosystem I: a symptom of plant diabetes. <i>Plant Cell Environ</i> 34(9), 1454–1464(2011)
651	W. Yamori, T. Nagai, A. Makino. The rate-limiting step for CO ₂ assimilation at different temperatures is influenced by the leaf nitrogen content in several C ₃ crop species. <i>Plant Cell Environ</i> 34(5), 764–777(2011)
652	Y. Suzuki, T. Miyamoto, R. Yoshizawa, T. Mae, A. Makino. Rubisco content and photosynthesis of leaves at different positions in transgenic rice with an overexpression of RBCS. <i>Plant Cell Environ</i> 32(4), 417–427(2009)

653	M. Riemann, M. Riemann, M. Takano. Rice JASMONATE RESISTANT 1 is involved in phytochrome and jasmonate signalling. <i>Plant Cell Environ</i> 31(6), 783–792(2008)
654	Y. Suzuki, T. Kihara-Doi, T. Kawazu, C. Miyake, A. Makino. Differences in Rubisco content and its synthesis in leaves at different positions in <i>Eucalyptus globulus</i> seedlings. <i>Plant Cell Environ</i> 33(8), 1314–1323(2010)
655	M. Izumi, H. Tsunoda, Y. Suzuki, A. Makino, H. Ishida. RBCS1A and RBCS3B, two major members within the Arabidopsis RBCS multigene family, function to yield sufficient Rubisco content for leaf photosynthetic capacity. <i>J Exp Bot</i> 63(5), 2159–2170(2012)
656	Y. Taniguchi, H. Ohkawa, C. Masumoto, T. Fukuda, T. Tamai, K. Lee, S. Sudoh, H. Tsuchida, H. Sasaki, H. Fukayama, M. Miyao. Overproduction of C4 photosynthetic enzymes in transgenic rice plants: an approach to introduce the C4-like photosynthetic pathway into rice. <i>J Exp Bot</i> 59(7), 1799–1809(2008)
657	Y. Nishikawa, H. Yamamoto, Y. Okegawa, S. Wada, N. Sato, Y. Taira, K. Sugimoto, A. Makino, T. Shikanai. PGR5-Dependent Cyclic Electron Transport Around PSI Contributes to the Redox Homeostasis in Chloroplasts Rather Than CO2 Fixation and Biomass Production in Rice. <i>Plant Cell Physiol</i> 53(12), 2117–2126(2012)
658	F. Hirose, N. Inagaki, A. Hanada, S. Yamaguchi, Y. Kamiya, A. Miyao, H. Hirochika, M. Takano. Cryptochrome and phytochrome cooperatively but independently reduce active gibberellin content in rice seedlings under light irradiation. <i>Plant Cell Physiol</i> 53(9), 1570–1582(2012)
659	H. Saito, E. Ogiso-Tanaka, Y. Okumoto, Y. Yoshitake, H. Izumi, T. Yokoo, K. Matsubara, K. Hori, M. Yano, H. Inoue, T. Tanisaka. Ef7 encodes an ELF3-like protein and promotes rice flowering by negatively regulating the floral repressor gene Ghd7 under both short- and long-day conditions. <i>Plant Cell Physiol</i> 53(4), 717–728(2012)
660	K. Matsubara, E. Ogiso-Tanaka, K. Hori, K. Ebana, T. Ando, M. Yano. Natural variation in Hd17, a homolog of Arabidopsis ELF3 that is involved in rice photoperiodic flowering. <i>Plant Cell Physiol</i> 53(4), 709–716(2012)
661	S. Ishida, K. Morita, M. Kishine, A. Takabayashi, R. Murakami, S. Takeda, K. Shimamoto, F. Sato, T. Endo. Allocation of absorbed light energy in PSII to thermal dissipations in the presence or absence of PsbS subunits of rice. <i>Plant Cell Physiol</i> 52(10), 1822–1831(2011)
662	N. Endo-Higashi, T. Izawa. Flowering time genes Heading date 1 and Early heading date 1 together control panicle development in rice. <i>Plant Cell Physiol</i> 52(6), 1083–1094(2011)
663	H. Nakamura, M. Muramatsu, M. Hakata, O. Ueno, Y. Nagamura, H. Hirochika, M. Takano, H. Ichikawa. Ectopic Overexpression of The Transcription Factor OsGLK1 Induces Chloroplast Development in Non-Green Rice Cells. <i>Plant Cell Physiol</i> 50(11), 1933–1949(2009)
664	Y. Suzuki, K. Nakabayashi, R. Yoshizawa, T. Mae, A. Makino. Differences in expression of the RBCS multigene family and rubisco protein content in various rice plant tissues at different growth stages. <i>Plant Cell Physiol</i> 50(10), 1851–1855(2009)
665	M. Ishikawa, Y. Ohmori, W. Tanaka, C. Hirabayashi, K. Murai, Y. Ogihara, T. Yamaguchi, H. Y. Hirano. The spatial expression patterns of DROOPING LEAF orthologs suggest a conserved function in grasses. <i>Genes Genet Syst</i> 84(2), 137–146(2009)
666	T. Nagai, A. Makino. Differences between rice and wheat in temperature responses of photosynthesis and plant growth. <i>Plant Cell Physiol</i> 50(4), 744–755(2009)
667	Y. Ohmori, M. Abiko, A. Horibata, H. Y. Hirano. A transposon, Ping, is integrated into intron 4 of the DROOPING LEAF gene of rice, weakly reducing its expression and causing a mild drooping leaf phenotype. <i>Plant Cell Physiol</i> 49(8), 1176–1184(2008)
668	S. Ishida, A. Takabayashi, N. Ishikawa, Y. Hano, T. Endo, F. Sato. A novel nuclear-encoded protein, NDH-dependent cyclic electron flow 5, is essential for the accumulation of chloroplast NAD(P)H dehydrogenase complexes. <i>Plant Cell Physiol</i> 50(2), 383–393(2009)
669	T. Takami, M. Shibata, Y. Kobayashi, T. Shikanai. De novo biosynthesis of fatty acids plays critical roles in the response of the photosynthetic machinery to low temperature in Arabidopsis. <i>Plant Cell Physiol</i> 51(8), 1265–1275(2010)
670	T. Shibaya, Y. Nonoue, N. Ono, U. Yamanouchi, K. Hori, M. Yano. Genetic interactions involved in the inhibition of heading by heading date QTL, Hd2 in rice under long-day conditions. <i>Theor Appl Genet</i> 123(7), 1133–1143(2011)

671	K. Ebana, T. Shibaya, J. Wu, K. Matsubara, H. Kanamori, H. Yamane, U. Yamanouchi, T. Mizubayashi, I. Kono, A. Shomura, S. Ito, T. Ando, K. Hori, T. Matsumoto, M. Yano. Uncovering of major genetic factors generating naturally occurring variation in heading date among Asian rice cultivars. <i>Theor Appl Genet</i> 122(6), 1199–1210(2011)
672	K. Matsubara, I. Kono, K. Hori, Y. Nonoue, N. Ono, A. Shomura, T. Mizubayashi, S. Yamamoto, U. Yamanouchi, K. Shirasawa, T. Nishio, M. Yano. Novel QTLs for photoperiodic flowering revealed by using reciprocal backcross inbred lines from crosses between japonica rice cultivars. <i>Theor Appl Genet</i> 117(6), 935–945(2008)
673	Y. Nonoue, K. Fujino, Y. Hirayama, U. Yamanouchi, S. Y. Lin, M. Yano. Detection of quantitative trait loci controlling extremely early heading in rice. <i>Theor Appl Genet</i> 116(5), 715–722(2008)
674	K. Fujino, U. Yamanouchi, M. Yano. Roles of the Hd5 gene controlling heading date for adaptation to the northern limits of rice cultivation. <i>Theor Appl Genet</i> , (2012)
675	竹内善信, 加藤浩, 根本博, 太田久稔, 佐藤宏之, 平山正賢, 平林秀介, 出田収, 青木法明, 坂井真, 蛭谷武志, 田口文緒, 山本敏央, 矢野昌裕, 井辺時雄, 安東郁男. コシヒカリと同質の遺伝的背景を持つ極早生の水稲品種「コシヒカリ関東HD1号」の育成. <i>作物研究所報告</i> 9, 1–25(2008)
676	Takagi D, Yamamoto H, Sugimoto T, Amako K, Makino A, Miyake C. O ₂ supports 3-phosphoglycerate-dependent O ₂ evolution in chloroplasts from spinach leaves. <i>Soil Sci. Plant Nutr.</i> 58(4), 462–468(2012)
677	Kubo S, Masumura T, Saito Y, Fukayama H, Suzuki Y, Sugimoto T, Makino A, Amako K, Miyake C. Cyclic electron flow around PSI functions in the photoinhibited rice leaves. <i>Soil Sci. Plant Nutr.</i> 57(1), 105–113(2011)
678	Kanno K, Makino A. Increased grain yield and biomass allocation in rice under cool night temperature. <i>Soil Sci. Plant Nutr.</i> 56(3), 412–417(2010)
679	K. Kanno, T. Mae, A. Makino. High night temperature stimulates photosynthesis, biomass production and growth during the vegetative stage of rice plants. <i>Soil Sci. Plant Nutr.</i> 55(1), 124–131(2009)
680	A. Yoshida, H.-Y. Hirano. Mapping of the ABERRANT SPIKELET 1 (ASP1) gene in rice. <i>Rice Genet. Newsl.</i> 25, 12–13(2010)
681	W. Tanaka, A. Ohtsubo, Y. Ohomori, H.-Y. Hirano. Identification of the conserved region in intron 1 of the DROOPING LEAF genes among the species in the grass family. <i>Rice Genet. Newsl.</i> 25, 78–79(2010)
682	H. Shikata, M. Nakashima, K. Matsuoka, T. Matsushita. Deletion of the RS domain of RRC1 impairs phytochrome B signaling in Arabidopsis. <i>Plant Signal Behav</i> 7(8), 933–936(2012)
683	W. Tanaka, T. Toriba, Y. Ohomori, H. Y. Hirano. Formation of two florets within a single spikelet in the rice tongari-boushi1 mutant. <i>Plant Signal Behav</i> 7(7), 793–795(2012)
684	H. Itoh, T. Izawa. A study of phytohormone biosynthetic gene expression using a circadian clock-related mutant in rice. <i>Plant Signal Behav</i> 6(12), 1932–1936(2011)
685	T. Toriba, Y. Ohmori, H. Y. Hirano. Common and distinct mechanisms underlying the establishment of adaxial and abaxial polarity in stamen and leaf development. <i>Plant Signal Behav</i> 6(3), 430–433(2011)
686	M. Riemann, D. Bouyer, A. Hisada, A. Muller, O. Yatou, E. W. Weiler, M. Takano, M. Furuya, P. Nick. Phytochrome A requires jasmonate for photodestruction. <i>Planta</i> 229(5), 1035–1045(2009)
687	H Fukayama, T Fukuda, C Masumoto, Y Taniguchi, H Sakai, W Cheng, T. Hasegawa, M. Miyao. Rice plant response to long term CO ₂ enrichment: Gene expression profiling. <i>Plant Sci</i> 177(3), 203–210(2009)
688	Tomonao Matsushita. A versatile method to prevent transcriptional gene silencing in Arabidopsis thaliana. <i>Plant Biotechnology</i> 5, 515–519(2012)
689	R. Nakano, H. Ishida, M. Kobayashi, A. Makino, T. Mae. Biochemical changes associated with in vivo RbcL fragmentation by reactive oxygen species under chilling-light conditions. <i>Plant Biol (Stuttg)</i> 12(1), 35–45(2010)
690	K. Jumtee, A. Okazawa, K. Harada, E. Fukusaki, M. Takano, A. Kobayashi. Comprehensive metabolite profiling of phyA phyB phyC triple mutants to reveal their associated metabolic phenotype in rice leaves. <i>J Biosci Bioeng</i> 108(2), 151–159(2009)

691	A. Einali, M. Shariati, F. Sato, T. Endo. Cyclic electron transport around photosystem I and its relationship to non-photochemical quenching in the unicellular green alga <i>Dunaliella salina</i> under nitrogen deficiency. <i>J Plant Res</i> 126(1), 179–186(2013)
692	R. Ishikawa, T. Shinomura, M. Takano, K. Shimamoto. Phytochrome dependent quantitative control of Hd3a transcription is the basis of the night break effect in rice flowering. <i>Genes Genet Syst</i> 84(2), 179–184(2009)
693	T. Fujioka, F. Kaneko, T. Kazama, K. Suwabe, G. Suzuki, A. Makino, T. Mae, M. Endo, M. Kawagishi–Kobayashi, M. Watanabe. Identification of small RNAs in late developmental stage of rice anthers. <i>Genes Genet Syst</i> 83(3), 281–284(2008)
694	M. Abiko, Y. Ohmori, H. Y. Hirano. Genome-wide expression profiling and identification of genes under the control of the DROOPING LEAF gene during midrib development in rice. <i>Genes Genet Syst</i> 83(3), 237–244(2008)
695	Seneweera S, Makino A, Hirotsu N, Norton R, Suzuki Y. New insight into photosynthetic acclimation to elevated CO ₂ : The role of leaf nitrogen and ribulose-1,5-bisphosphate carboxylase/oxygenase in rice leaves. <i>Environ Exp Bot.</i> 71(2), 128–136(2011)
696	M. Hakata, M. Kuroda, A. Ohsumi, T. Hirose, H. Nakamura, M. Muramatsu, H. Ichikawa, H. Yamakawa. Overexpression of a Rice TIFY Gene Increases Grain Size through Enhanced Accumulation of Carbohydrates in the Stem. <i>Biosci Biotechnol Biochem</i> 76(11), 2129–2134(2012)
697	Y. Suzuki, T. Mae, A. Makino. RNA extraction from various recalcitrant plant tissues with a cethyltrimethylammonium bromide-containing buffer followed by an acid guanidium thiocyanate-phenol-chloroform treatment. <i>Biosci Biotechnol Biochem</i> 72(7), 1951–1953(2008)
698	K. Imai, Y. Suzuki, T. Mae, A. Makino. Changes in the synthesis of rubisco in rice leaves in relation to senescence and N influx. <i>Ann Bot</i> 101(1), 135–144(2008)
699	Irving LJ, Suzuki Y, Ishida H, Makino A. Protein turnover in grass leaves. <i>Adv. Bot. Res.</i> 54, 139–182(2010)
700	Y. Kobayashi, I. Kobayashi. Microwounding is a pivotal factor for the induction of actin-dependent penetration resistance against fungal attack. <i>Planta</i> , (2013) in press
701	Ryohei Terauchi, Akira Abe, Hiroki Takagi, Kentaro Yoshida, Shunichi Kosugi, Satoshi Natsume, Hiroki Yaegashi, Hiroyuki Kanzaki, Hideo Matsumura, Chikako Mitsuoka, Hiroe Utsushi, Muluneh Tamiru. Whole genome sequencing and future breeding of rice. <i>Journal of Plant Biochemistry and Biotechnology</i> 21(1), S10–S14(2012)
702	MJ. Wamaitha, R. Yamamoto, HL. Wong, T. Kawasaki, Y. Kawano, K. Shimamoto. OsRap2.6 transcription factor contributes to rice innate immunity through its interaction with Receptor for Activated Kinase-C 1 (RACK1). <i>Rice.</i> 5, 35(2012)
703	C. J. Jiang, M. Shimono, S. Sugano, M. Kojima, X. Liu, H. Inoue, H. Sakakibara, H. Takatsuji. Cytokinins act synergistically with salicylic acid to activate defense gene expression in rice. <i>Mol Plant Microbe Interact</i> , (2012) in press
704	T. Sisaphaithong, D. Kondo, H. Matsunaga, Y. Kobae, S. Hata. Expression of plant genes for arbuscular mycorrhiza-inducible phosphate transporters and fungal vesicle formation in sorghum, barley, and wheat roots. <i>Biosci Biotechnol Biochem</i> 76(12), 2364–2367(2012)
705	A. Matsushita, H. Inoue, S. Goto, A. Nakayama, S. Sugano, N. Hayashi, H. Takatsuji. The nuclear ubiquitin proteasome degradation affects WRKY45 function in the rice defense program. <i>Plant J</i> , 73, 302–313 (2012)
706	J. R. Undan, M. Tamiru, A. Abe, K. Yoshida, S. Kosugi, H. Takagi, H. Kanzaki, H. Saitoh, R. Fekih, S. Sharma, J. Undan, M. Yano, R. Terauchi. Mutation in OsLMS, a gene encoding a protein with two double-stranded RNA binding motifs, causes lesion mimic phenotype and early senescence in rice (<i>Oryza sativa</i> L.). <i>Genes Genet Syst</i> 87(3), 169–179(2012)
707	T. Fujikawa, A. Sakaguchi, Y. Nishizawa, Y. Kouzai, E. Minami, S. Yano, H. Koga, T. Meshi, M. Nishimura. Surface alpha-1,3-Glucan Facilitates Fungal Stealth Infection by Interfering with Innate Immunity in Plants. <i>PLoS Pathog</i> 8(8), e1002882(2012)
708	H. Kanzaki, K. Yoshida, H. Saitoh, K. Fujisaki, A. Hirabuchi, L. Alaux, E. Fournier, D. Tharreau, R. Terauchi. Arms race co-evolution of <i>Magnaporthe oryzae</i> AVR-Pik and rice Pik genes driven by their physical interactions. <i>Plant J</i> , in press(2012)
709	T. Okubo, S. Ikeda, A. Yamashita, K. Terasawa, K. Minamisawa. Pyrosequence read length of 16S rRNA gene affects phylogenetic assignment of plant-associated bacteria. <i>Microbes Environ</i> 27(2), 204–208(2012)

710	H. Saitoh, S. Fujisawa, C. Mitsuoka, A. Ito, A. Hirabuchi, K. Ikeda, H. Irieda, K. Yoshino, K. Yoshida, H. Matsumura, Y. Tosa, J. Win, S. Kamoun, Y. Takano, R. Terauchi. Large-scale gene disruption in <i>Magnaporthe oryzae</i> identifies MC69, a secreted protein required for infection by monocot and dicot fungal pathogens. <i>PLoS Pathog</i> 8(5), e1002711(2012)
711	S. Takahashi, S. Kimura, H. Kaya, A. Iizuka, H. L. Wong, K. Shimamoto, K. Kuchitsu. Reactive oxygen species production and activation mechanism of the rice NADPH oxidase OsRbohB. <i>J Biochem</i> 152(1), 37-43(2012)
712	T. Okubo, T. Tsukui, H. Maita, S. Okamoto, K. Oshima, T. Fujisawa, A. Saito, H. Futamata, R. Hattori, Y. Shimomura, S. Haruta, S. Morimoto, Y. Wang, Y. Sakai, M. Hattori, S. Aizawa, K. V. Nagashima, S. Masuda, T. Hattori, A. Yamashita, Z. Bao, M. Hayatsu, H. Kajiya-Kanegae, I. Yoshinaga, K. Sakamoto, K. Toyota, M. Nakao, M. Kohara, M. Anda, R. Niwa, P. Jung-Hwan, R. Sameshima-Saito, S. Tokuda, S. Yamamoto, T. Yokoyama, T. Akutsu, Y. Nakamura, Y. Nakahira-Yanaka, Y. Takada Hoshino, H. Hirakawa, H. Mitsui, K. Terasawa, M. Itakura, S. Sato, W. Ikeda-Ohtsubo, N. Sakakura, E. Kaminuma, K. Minamisawa. Complete genome sequence of <i>Bradyrhizobium</i> sp. S23321: insights into symbiosis evolution in soil oligotrophs. <i>Microbes Environ</i> 27(3), 306-315(2012)
713	S. H. Kim, T. Oikawa, J. Kyozuka, H. L. Wong, K. Umemura, M. Kishi-Kaboshi, A. Takahashi, Y. Kawano, T. Kawasaki, K. Shimamoto. The bHLH <i>Rac Immunity1 (RAI1)</i> Is Activated by <i>OsRac1</i> via <i>OsMAPK3</i> and <i>OsMAPK6</i> in Rice Immunity. <i>Plant Cell Physiol</i> 53(4), 740-754(2012)
714	N. Takeda, T. Maekawa, M. Hayashi. Nuclear-localized and deregulated calcium- and calmodulin-dependent protein kinase activates rhizobial and mycorrhizal responses in <i>Lotus japonicus</i> . <i>Plant Cell</i> 24(2), 810-822(2012)
715	A. Abe, S. Kosugi, K. Yoshida, S. Natsume, H. Takagi, H. Kanzaki, H. Matsumura, C. Mitsuoka, M. Tamiru, H. Innan, L. Cano, S. Kamoun, R. Terauchi. Genome sequencing reveals agronomically important loci in rice using MutMap. <i>Nat Biotechnol</i> 30(2), 174-178(2012)
716	M. Shimono, H. Koga, A. Akagi, N. Hayashi, S. Goto, M. Sawada, T. Kurihara, A. Matsushita, S. Sugano, C. J. Jiang, H. Kaku, H. Inoue, H. Takatsuji. Rice <i>WRKY45</i> plays important roles in fungal and bacterial disease resistance. <i>Mol Plant Pathol</i> 13(1), 83-94(2012)
717	Katsumi Yazawa, Chang-Jie Jiang, Mikiko Kojima, Hitoshi Sakakibara, Hiroshi Takatsuji. Reduction of abscisic acid levels or inhibition of abscisic acid signaling in rice during the early phase of <i>Magnaporthe grisea</i> infection decreases its susceptibility to the fungus. <i>Physiol Mol Plant Pathol</i> 78, 1-7(2012)
718	J. Hirayama, S. Eda, H. Mitsui, K. Minamisawa. Nitrate-dependent N ₂ O emission from intact soybean nodules via denitrification by <i>Bradyrhizobium japonicum</i> bacteroids. <i>Appl Environ Microbiol</i> 77(24), 8787-8790(2011)
719	S. Ikeda, T. Okubo, N. Takeda, M. Banba, K. Sasaki, H. Imaizumi-Anraku, S. Fujihara, Y. Ohwaki, K. Ohshima, Y. Fukuta, S. Eda, H. Mitsui, M. Hattori, T. Sato, T. Shinano, K. Minamisawa. The genotype of the calcium/calmodulin-dependent protein kinase gene (<i>CCaMK</i>) determines bacterial community diversity in rice roots under paddy and upland field conditions. <i>Appl Environ Microbiol</i> 77(13), 4399-4405(2011)
720	M. Anda, S. Ikeda, S. Eda, T. Okubo, S. Sato, S. Tabata, H. Mitsui, K. Minamisawa. Isolation and genetic characterization of <i>Aurantimonas</i> and <i>Methylobacterium</i> strains from stems of hypernodulated soybeans. <i>Microbes Environ</i> 26(2), 172-180(2011)
721	Y. Okuyama, H. Kanzaki, A. Abe, K. Yoshida, M. Tamiru, H. Saitoh, T. Fujibe, H. Matsumura, M. Shenton, D. C. Galam, J. Undan, A. Ito, T. Sone, R. Terauchi. A multifaceted genomics approach allows the isolation of the rice <i>Pia</i> -blast resistance gene consisting of two adjacent NBS-LRR protein genes. <i>Plant J</i> 66(3), 467-479(2011)
722	S. Ikeda, M. Anda, S. Inaba, S. Eda, S. Sato, K. Sasaki, S. Tabata, H. Mitsui, T. Sato, T. Shinano, K. Minamisawa. Autoregulation of nodulation interferes with impacts of nitrogen fertilization levels on the leaf-associated bacterial community in soybeans. <i>Appl Environ Microbiol</i> 77(6), 1973-1980(2011)
723	Letian Chen, Ko Shimamoto. Emerging roles of molecular chaperones in plant innate immunity. <i>J. Gen. Plant Pathol.</i> 77(1), 40917(2011)
724	H. Matsui, A. Miyao, A. Takahashi, H. Hirochika. Pdk1 kinase regulates basal disease resistance through the OsOxi1-OsPti1a phosphorylation cascade in rice. <i>Plant Cell Physiol</i> 51(12), 2082-2091(2010)

725	C. J. Jiang, M. Shimono, S. Sugano, M. Kojima, K. Yazawa, R. Yoshida, H. Inoue, N. Hayashi, H. Sakakibara, H. Takatsuji. Abscisic Acid Interacts Antagonistically with Salicylic Acid Signaling Pathway in Rice–Magnaporthe grisea Interaction. <i>Mol Plant Microbe Interact</i> 23(6), 791–798(2010)
726	S. Masuda, S. Eda, C. Sugawara, H. Mitsui, K. Minamisawa. The cbbL gene is required for thiosulfate-dependent autotrophic growth of <i>Bradyrhizobium japonicum</i> . <i>Microbes Environ.</i> 25, 220–223(2010)
727	T. Isawa, M. Yasuda, H. Awazaki, K. Minamisawa, S. Shinozask., H. Nakshita. <i>Azospirillum</i> sp. strain B510 enhances rice growth and yield. <i>Microbes Environ.</i> 25, 58–56(2010)
728	Kazuhiro S, Seishi I, Shima E, Hisayuki M, Eiko H, Chiharu K, Yuri K, Atsuhiko K, Takuro S, Kiwamu M, Tadashi S. Impacts of plant genotype and nitrogen level on rice growth response to inoculation with <i>Azospirillum</i> sp. strain B510 under paddy field conditions. <i>Soil Science and Plant Nutrition</i> 56(4), 636–644(2010)
729	S. Ikeda, T. Kaneko, T. Okubo, L. E. Rallos, S. Eda, H. Mitsui, S. Sato, Y. Nakamura, S. Tabata, K. Minamisawa. Development of a bacterial cell enrichment method and its application to the community analysis in soybean stems. <i>Microb Ecol</i> 58(4), 703–714(2009)
730	T. Fujikawa, Y. Kuga, S. Yano, A. Yoshimi, T. Tachiki, K. Abe, M. Nishimura. Dynamics of cell wall components of <i>Magnaporthe grisea</i> during infectious structure development. <i>Mol Microbiol</i> 73(4), 553–570(2009)
731	C. J. Jiang, M. Shimono, S. Maeda, H. Inoue, M. Mori, M. Hasegawa, S. Sugano, H. Takatsuji. Suppression of the rice fatty-acid desaturase gene <i>OsSSI2</i> enhances resistance to blast and leaf blight diseases in rice. <i>Mol Plant Microbe Interact</i> 22(7), 820–829(2009)
732	M. Fujiwara, S. Hamada, M. Hiratsuka, Y. Fukao, T. Kawasaki, K. Shimamoto. Proteome analysis of detergent-resistant membranes (DRMs) associated with <i>OsRac1</i> -mediated innate immunity in rice. <i>Plant Cell Physiol</i> 50(7), 1191–1200(2009)
733	K. Yoshida, H. Saitoh, S. Fujisawa, H. Kanzaki, H. Matsumura, K. Yoshida, Y. Tosa, I. Chuma, Y. Takano, J. Win, S. Kamoun, R. Terauchi. Association Genetics Reveals Three Novel Avirulence Genes from the Rice Blast Fungal Pathogen <i>Magnaporthe oryzae</i> . <i>Plant Cell</i> 21(5), 1573–1591(2009)
734	S. A. Hogenhout, R. A. L. Van der Hoorn, R. Terauchi, S. Kamoun. Emerging Concepts in Effector Biology of Plant-Associated Organisms. <i>Mol Plant Microbe Interact</i> 22(2), 115–122(2009)
735	A. Krishnan, E. Guiderdoni, G. An, Y. I. Hsing, C. D. Han, M. C. Lee, S. M. Yu, N. Upadhyaya, S. Ramachandran, Q. Zhang, V. Sundaresan, H. Hirochika, H. Leung, A. Pereira. Mutant resources in rice for functional genomics of the grasses. <i>Plant Physiol</i> 149(1), 165–170(2009)
736	Y. Takahashi, T. Berberich, H. Kanzaki, H. Matsumura, H. Saitoh, T. Kusano, R. Terauchi. Serine Palmitoyltransferase, the First Step Enzyme in Sphingolipid Biosynthesis, Is Involved in Nonhost Resistance. <i>Mol Plant Microbe Interact</i> 22(1), 31–38(2009)
737	M. Yoshii, T. Shimizu, M. Yamazaki, T. Higashi, A. Miyao, H. Hirochika, T. Omura. Disruption of a novel gene for a NAC-domain protein in rice confers resistance to Rice dwarf virus. <i>Plant J</i> 57(4), 615–625(2009)
738	T Ohkubo, S. Ikeda, T. Kaneko, S. Eda, H. Mitsui, S. Sato, S. Tabata, K. Minamisawa. Nodulation-dependent communities of culturable bacteria endophytes from stems of field-grown soybeans. <i>Microbes Environ</i> 24(3), 253–258(2009)
739	C. Gutjahr, M. Banba, V. Croset, K. An, A. Miyao, G. An, H. Hirochika, H. Imaizumi-Anraku, U. Paszkowski. Arbuscular mycorrhiza-specific signaling in rice transcends the common symbiosis signaling pathway. <i>Plant Cell</i> 20(11), 2989–3005(2008)
740	M. Banba, C. Gutjahr, A. Miyao, H. Hirochika, U. Paszkowski, H. Kouchi, H. Imaizumi-Anraku. Divergence of evolutionary ways among common sym genes: <i>CASTOR</i> and <i>CCaMK</i> show functional conservation between two symbiosis systems and constitute the root of a common signaling pathway. <i>Plant Cell Physiol</i> 49(11), 1659–1671(2008)
741	T. Oda, H. Hashimoto, N. Kuwabara, K. Hayashi, C. Kojima, T. Kawasaki, K. Shimamoto, M. Sato, T. Shimizu. Crystallographic characterization of the N-terminal domain of a plant NADPH oxidase. <i>Acta Crystallogr Sect F Struct Biol Cryst Commun</i> 64(Pt 9), 867–869(2008)
742	A. Nakashima, L. Chen, N. P. Thao, M. Fujiwara, H. L. Wong, M. Kuwano, K. Umemura, K. Shirasu, T. Kawasaki, K. Shimamoto. <i>RACK1</i> functions in rice innate immunity by interacting with the <i>Rac1</i> immune complex. <i>Plant Cell</i> 20(8), 2265–2279(2008)

743	H. Kanzaki, H. Saitoh, Y. Takahashi, T. Berberich, A. Ito, S. Kamoun, R. Terauchi. NbLRK1, a lectin-like receptor kinase protein of <i>Nicotiana benthamiana</i> , interacts with <i>Phytophthora infestans</i> INF1 elicitor and mediates INF1-induced cell death. <i>Planta</i> 228(6), 977–987(2008)
744	S. Ikeda, L. E. Rallos, T. Okubo, S. Eda, S. Inaba, H. Mitsui, K. Minamisawa. Microbial community analysis of field-grown soybeans with different nodulation phenotypes. <i>Appl Environ Microbiol</i> 74(18), 5704–5709(2008)
745	S. Masuda, S. Eda, S. Ikeda, H. Mitsui, K. Minamisawa. Thiosulfate-dependent chemolithoautotrophic growth of <i>Bradyrhizobium japonicum</i> . <i>Appl Environ Microbiol</i> 76(8), 2402–2409(2010)
746	M. Yasuda, T. Isawa, S. Shinozaki, K. Minamisawa, H. Nakashita. Effects of colonization of a bacterial endophyte, <i>Azospirillum</i> sp. B510, on disease resistance in rice. <i>Biosci Biotechnol Biochem</i> 73(12), 2595–2599(2009)
747	A. Takahashi, N. Hayashi, A. Miyao, H. Hirochika. Unique features of the rice blast resistance Pish locus revealed by large scale retrotransposon-tagging. <i>BMC Plant Biol</i> 10, 175(2010)
748	L. Chen, S. Hamada, M. Fujiwara, T. Zhu, N. P. Thao, H. L. Wong, P. Krishna, T. Ueda, H. Kaku, N. Shibuya, T. Kawasaki, K. Shimamoto. The Hop/Sti1-Hsp90 chaperone complex facilitates the maturation and transport of a PAMP receptor in rice innate immunity. <i>Cell Host Microbe</i> 7(3), 185–196(2010)
749	Y. Kawano, A. Akamatsu, K. Hayashi, Y. Housen, J. Okuda, A. Yao, A. Nakashima, H. Takahashi, H. Yoshida, H. L. Wong, T. Kawasaki, K. Shimamoto. Activation of a Rac GTPase by the NLR family disease resistance protein Pit plays a critical role in rice innate immunity. <i>Cell Host Microbe</i> 7(5), 362–375(2010)
750	K. Yokota, M. Hayashi. Function and evolution of nodulation genes in legumes. <i>Cell Mol Life Sci</i> 68(8), 1341–1351(2011)
751	T. Kaneko, K. Minamisawa, T. Isawa, H. Nakatsukasa, H. Mitsui, Y. Kawaharada, Y. Nakamura, A. Watanabe, K. Kawashima, A. Ono, Y. Shimizu, C. Takahashi, C. Minami, T. Fujishiro, M. Kohara, M. Katoh, N. Nakazaki, S. Nakayama, M. Yamada, S. Tabata, S. Sato. Complete genomic structure of the cultivated rice endophyte <i>Azospirillum</i> sp. B510. <i>DNA Res</i> 17(1), 37–50(2010)
752	S. Ikeda, T. Okubo, T. Kaneko, S. Inaba, T. Maekawa, S. Eda, S. Sato, S. Tabata, H. Mitsui, K. Minamisawa. Community shifts of soybean stem-associated bacteria responding to different nodulation phenotypes and N levels. <i>Isme J</i> 4(3), 315–326(2010)
753	T. Oda, H. Hashimoto, N. Kuwabara, S. Akashi, K. Hayashi, C. Kojima, H. L. Wong, T. Kawasaki, K. Shimamoto, M. Sato, T. Shimizu. Structure of the N-terminal regulatory domain of a plant NADPH oxidase and its functional implications. <i>J Biol Chem</i> 285(2), 1435–1445(2010)
754	T. Fujiwara, S. Maisonneuve, M. Isshiki, M. Mizutani, L. Chen, H. L. Wong, T. Kawasaki, K. Shimamoto. Sekiguchi lesion gene encodes a cytochrome P450 monooxygenase that catalyzes conversion of tryptamine to serotonin in rice. <i>J Biol Chem</i> 285(15), 11308–11313(2010)
755	R. Terauchi, K. Yoshida. Towards population genomics of effector-effector target interactions. <i>New Phytol</i> 187(4), 929–939(2010)
756	H. Kouchi, H. Imaizumi-Anraku, M. Hayashi, T. Hakoyama, T. Nakagawa, Y. Umehara, N. Sukanuma, M. Kawaguchi. How many peas in a pod? Legume genes responsible for mutualistic symbioses underground. <i>Plant Cell Physiol</i> 51(9), 1381–1397(2010)
757	K. Yokota, T. Soyano, H. Kouchi, M. Hayashi. Function of GRAS proteins in root nodule symbiosis is retained in homologs of a non-legume, rice. <i>Plant Cell Physiol</i> 51(9), 1436–1442(2010)
758	Y. Kobae, S. Hata. Dynamics of periarbuscular membranes visualized with a fluorescent phosphate transporter in arbuscular mycorrhizal roots of rice. <i>Plant Cell Physiol</i> 51(3), 341–353(2010)
759	S. Ikeda, T. Okubo, M. Anda, H. Nakashita, M. Yasuda, S. Sato, T. Kaneko, S. Tabata, S. Eda, A. Momiyama, K. Terasawa, H. Mitsui, K. Minamisawa. Community- and genome-based views of plant-associated bacteria: plant-bacterial interactions in soybean and rice. <i>Plant Cell Physiol</i> 51(9), 1398–1410(2010)
760	L. Chen, K. Shiotani, T. Togashi, D. Miki, M. Aoyama, H. L. Wong, T. Kawasaki, K. Shimamoto. Analysis of the Rac/Rop small GTPase family in rice: expression, subcellular localization and role in disease resistance. <i>Plant Cell Physiol</i> 51(4), 585–595(2010)

761	H. Matsui, M. Yamazaki, M. Kishi-Kaboshi, A. Takahashi, H. Hirochika. AGC kinase OsOxi1 positively regulates basal resistance through suppression of OsPti1a-mediated negative regulation. <i>Plant Cell Physiol</i> 51(10), 1731-1744(2010)
762	M. Kishi-Kaboshi, K. Okada, L. Kurimoto, S. Murakami, T. Umezawa, N. Shibuya, H. Yamane, A. Miyao, H. Takatsuji, A. Takahashi, H. Hirochika. A rice fungal MAMP-responsive MAPK cascade regulates metabolic flow to antimicrobial metabolite synthesis. <i>Plant J</i> 63(4), 599-612(2010)
763	M. Yoshii, M. Yamazaki, R. Rakwal, M. Kishi-Kaboshi, A. Miyao, H. Hirochika. The NAC transcription factor RIM1 of rice is a new regulator of jasmonate signaling. <i>Plant J</i> 61(5), 804-815(2010)
764	T. Hayashi, M. Banba, Y. Shimoda, H. Kouchi, M. Hayashi, H. Imaizumi-Anraku. A dominant function of CCaMK in intracellular accommodation of bacterial and fungal endosymbionts. <i>Plant J</i> 63(1), 141-154(2010)
765	S. Sugano, C. J. Jiang, S. Miyazawa, C. Masumoto, K. Yazawa, N. Hayashi, M. Shimono, A. Nakayama, M. Miyao, H. Takatsuji. Role of OsNPR1 in rice defense program as revealed by genome-wide expression analysis. <i>Plant Mol Biol</i> 74(6), 549-562(2010)
766	Y. Kobae. The use of wideband filters in distinguish green fluorescent protein in roots of arbuscular mycorrhizal symbiosis. <i>Plant Signal Behav</i> 5(9), 1150-1152(2010)
767	M. Kishi-Kaboshi, A. Takahashi, H. Hirochika. MAMP-responsive MAPK cascades regulate phytoalexin biosynthesis. <i>Plant Signal Behav</i> 5(12), 1653-1656(2010)
768	Okubo, T, Fukushima, S., Itakura, M., Oshima, K., Longtonglang, A., Teaumroong, N., Mitsui, H., Hattori, M., Hattori, R., Hattori, T., and Minamisawa, K. Soil oligotrophic bacterium <i>Agromonas oligotrophica</i> (<i>Bradyrhizobium oligotrophicum</i>) is a nitrogen-fixing symbiont of <i>Aeschynomene indica</i> as suggested by genome analysis. <i>Appl. Environ. Microbiol.</i> ,in press(2013)
769	Tsukui, T., S. Eda, T. Kaneko, S. Sato, S. Okazaki, K. Kakizaki-Chiba, M. Itakura, H. Mitsui, A. Yamashita, K. Terasawa, and K. Minamisawa. The type III secretion system of <i>Bradyrhizobium japonicum</i> USDA122 mediates symbiotic incompatibility with Rj2 soybean. <i>Appl. Environ. Microbiol.</i> 79, 1048-1051(2012)
770	Okubo, T., S. Fukushima, and K. Minamisawa. Evolution of <i>Bradyrhizobium</i> - <i>Aeschynomene</i> mutualism: Living testimony of the ancient world or highly evolved state? <i>Plant Cell Physiol.</i> 53(12), 2000-2007(2012)
771	Y. Kawano, L. Chen, K. Shimamoto. The Function of Rac Small GTPase and Associated Proteins in Rice Innate Immunity. <i>Rice.</i> 3, 112-121(2010)
772	S. Sakuma, M. Pourkheirandish, G. Hensel, J. Kumlehn, N. Stein, A. Tagiri, N. Yamaji, J. F. Ma, H. Sassa, T. Koba, T. Komatsuda. Divergence of expression pattern contributed to neofunctionalization of duplicated HD-Zip I transcription factor in barley. <i>New Phytol</i> 197(3), 939-948(2013)
773	C. Li, A. Wang, X. Ma, M. Pourkheirandish, S. Sakuma, N. Wang, S. Ning, E. Nevo, C. Nawrath, T. Komatsuda, G. Chen. An eceriferum locus, cer-zv, is associated with a defect in cutin responsible for water retention in barley (<i>Hordeum vulgare</i>) leaves. <i>Theor Appl Genet</i> , in press(2012)
774	K. F. Mayer, R. Waugh, J. W. Brown, A. Schulman, P. Langridge, M. Platzer, G. B. Fincher, G. J. Muehlbauer, K. Sato, T. J. Close, R. P. Wise, N. Stein. A physical, genetic and functional sequence assembly of the barley genome. <i>Nature</i> 491(7426), 711-716(2012)
775	T. Yuo, Y. Yamashita, H. Kanamori, T. Matsumoto, U. Lundqvist, K. Sato, M. Ichii, S. A. Jobling, S. Taketa. A SHORT INTERNODES (SHI) family transcription factor gene regulates awn elongation and pistil morphology in barley. <i>J Exp Bot</i> 63(14), 5223-5232(2012)
776	S. Taketa, T. Yuo, T. Tonooka, Y. Tsumuraya, Y. Inagaki, N. Haruyama, O. Larroque, S. A. Jobling. Functional characterization of barley betaglucanless mutants demonstrates a unique role for CslF6 in (1,3;1,4)-beta-D-glucan biosynthesis. <i>J Exp Bot</i> 63(1), 381-392(2012)
777	Suzuki T, Sato M, Takeuchi T. Evaluation of the effects of five QTL regions on Fusarium head blight resistance and agronomic traits in spring wheat (<i>Triticum aestivum</i> L.). <i>Breed. Sci.</i> 62(1), 11-17(2012)
778	S. Nakamura, F. Abe, H. Kawahigashi, K. Nakazono, A. Tagiri, T. Matsumoto, S. Utsugi, T. Ogawa, H. Handa, H. Ishida, M. Mori, K. Kawaura, Y. Ogihara, H. Miura. A wheat homolog of MOTHER OF FT AND TFL1 acts in the regulation of germination. <i>Plant Cell</i> 23(9), 3215-3229(2011)

779	G. Chen, T. Komatsuda, J. F. Ma, C. Nawrath, M. Pourkheirandish, A. Tagiri, Y. G. Hu, M. Sameri, X. Li, X. Zhao, Y. Liu, C. Li, X. Ma, A. Wang, S. Nair, N. Wang, A. Miyao, S. Sakuma, N. Yamaji, X. Zheng, E. Nevo. An ATP-binding cassette subfamily G full transporter is essential for the retention of leaf water in both wild barley and rice. <i>Proc Natl Acad Sci U S A</i> 108(30), 12354–12359(2011)
780	K. F. Mayer, M. Martis, P. E. Hedley, H. Simkova, H. Liu, J. A. Morris, B. Steuernagel, S. Taudien, S. Roessner, H. Gundlach, M. Kubalaková, P. Suchanková, F. Murat, M. Felder, T. Nussbaumer, A. Graner, J. Salse, T. Endo, H. Sakai, T. Tanaka, T. Itoh, K. Sato, M. Platzer, T. Matsumoto, U. Scholz, J. Dolezel, R. Waugh, N. Stein. Unlocking the barley genome by chromosomal and comparative genomics. <i>Plant Cell</i> 23(4), 1249–1263(2011)
781	T. Matsumoto, T. Tanaka, H. Sakai, N. Amano, H. Kanamori, K. Kurita, A. Kikuta, K. Kamiya, M. Yamamoto, H. Ikawa, N. Fujii, K. Hori, T. Itoh, K. Sato. Comprehensive sequence analysis of 24,783 barley full-length cDNAs derived from 12 clone libraries. <i>Plant Physiol</i> 156(1), 20–28(2011)
782	K Kakeda, N Ishihara, Y Izumi, K. Sato, S. Taketa. Expression and functional analysis of the barley Nud gene using transgenic rice. <i>Breed. Sci.</i> 61(1), 35–42(2011)
783	Sameri M, Pourkheirandish M, Chen G, Tonooka T, Komatsuda T. Detection of photoperiod responsive and non-responsive flowering time QTL in barley. <i>Breed. Sci.</i> 61(2), 183–188(2011)
784	Insaf Bahri, Motoki Sugisawa, Rie Kikuchi, Taiichi Ogawa, Hiroyuki Kawahigashi, Tomohiro Ban, Hirokazu Handa. Characterization of a wheat transcription factor, TaWRKY45, and its effect on Fusarium head blight resistance in transgenic wheat plants. <i>Breed. Sci.</i> 61(2), 121–129(2011)
785	E Himi, Y Yamashita, N Haruyama, T Yanagisawa, M. Maekawa, S. Taketa. Ant28 gene for proanthocyanidin synthesis encoding the R2R3 MYB domain protein (Hvmyb10) highly affects grain dormancy in barley. <i>Euphytica</i> 188(1), 141–151(2011)
786	Baba T, Tanno K, Furusho M, Komatsuda T. Allelic variation at the EF-G locus among northern Moroccan six-rowed barleys. <i>Plant Genetic Resources: Characterization and Utilization</i> 9(2), 240–242(2011)
787	S. Taketa, T. Yuo, Y. Sakurai, S. Miyake, M Ichii. Molecular mapping of the short awn 2 (lks2) and dense spike 1 (dsp1) genes on barley chromosome 7H. <i>Breed. Sci.</i> 61(1), 80–85(2011)
788	N. Mizuno, M. Yamasaki, Y. Matsuoka, T. Kawahara, S. Takumi. Population structure of wild wheat D-genome progenitor <i>Aegilops tauschii</i> Coss.: implications for intraspecific lineage diversification and evolution of common wheat. <i>Mol Ecol</i> 19(5), 999–1013(2010)
789	Sudha K. Nair, Ning Wang, Yerlan Turuspekov, Mohammad Pourkheirandish, Suphawat Sinsuwongwat, Guoxiong Chen, Mohammad Sameri, Akemi Tagiri, Ichiro Honda, Yoshiaki Watanabe, Hiroyuki Kanamori, Thomas Wicker, Nils Stein, Yoshiaki Nagamura, Takashi Matsumoto, Takao Komatsuda. Cleistogamous flowering in barley arises from the suppression of microRNA-guided HvAP2 mRNA cleavage. <i>Proc Natl Acad Sci U S A</i> 107(1), 490–495(2010)
790	S. Sakuma, M. Pourkheirandish, T. Matsumoto, T. Koba, T. Komatsuda. Duplication of a well-conserved homeodomain-leucine zipper transcription factor gene in barley generates a copy with more specific functions. <i>Funct Integr Genomics</i> 10(1), 123–133(2010)
791	Yuki Fujiwara, Sanae Shimada, Shigeo Takumi, Koji Murai. Differential effects of <i>Aegilops tauschii</i> genotypes on maturing-time in synthetic hexaploid wheats. <i>Breed. Sci.</i> 60(3), 286–292(2010)
792	H. Kouzaki, S. Kidou, H. Miura, K. Kato. Genomic diversity of germinating scutellum specific gene P23k in barley and wheat. <i>Genetica</i> 137(2), 233–242(2009)
793	S Golkari, J Gilbert, T. Ban, Procunier J. QTL-specific microarray gene expression analysis of wheat resistance to Fusarium head blight in Sumai-3 and two susceptible NILs. <i>Genome</i> 52(5), 409–418(2009)
794	M. Sameri, S. Nakamura, S. K. Nair, K. Takeda, T. Komatsuda. A quantitative trait locus for reduced culm internode length in barley segregates as a Mendelian gene. <i>Theor Appl Genet</i> 118(4), 643–652(2009)

795	T. Wicker, S. G. Krattinger, E. S. Lagudah, T. Komatsuda, M. Pourkheirandish, T. Matsumoto, S. Cloutier, L. Reiser, H. Kanamori, K. Sato, D. Perovic, N. Stein, B. Keller. Analysis of Intraspecies Diversity in Wheat and Barley Genomes Identifies Breakpoints of Ancient Haplotypes and Provides Insight into the Structure of Diploid and Hexaploid Triticeae Gene Pools. <i>Plant Physiol</i> 149(1), 258–270(2009)
796	Sven Gottwald, Petra Bauer, Takao Komatsuda, Udda Lundqvist, Nils Stein. TILLING in the two-rowed barley cultivar 'Barke' reveals preferred sites of functional diversity in the gene HvHox1. <i>BMC Research Notes</i> 2, 258(2009)
797	T. Tonooka, E. Aoki, T. Yoshioka, S. Taketa. A novel mutant gene for (1-3, 1-4)- β -D-glucanless grain on barley (<i>Hordeum vulgare</i> L.) chromosome 7H. <i>Breed. Sci.</i> 59(1), 47–54(2009)
798	Takumi S, Nishioka E, Morihiro H, Kawahara T, Matsuoka Y. Natural variation of morphological traits in wild wheat progenitor <i>Aegilops tauschii</i> Coss. <i>Breed. Sci.</i> 59(5), 579–588(2009)
799	Katsuyuki Kakeda, Shin Taketa, Takao Komatsuda. Molecular phylogeny of the genus <i>Hordeum</i> using thioredoxin-like gene sequences. <i>Breed. Sci.</i> 59(5), 595–601(2009)
800	Guoxiong Chen, Takao Komatsuda, Mohammad Pourkheirandish, Mohammad Sameri, Kazuhiro Sato, Tamar Krugman, Tzion Fahima, Abraham B. Korol, Eviatar Nevo. Mapping of the eibi1 gene responsible for the drought hypersensitive cuticle in wild barley (<i>Hordeum spontaneum</i>). <i>Breed. Sci.</i> 59(1), 21–26(2009)
801	Fahimeh Shahinnia, Badraddin Ebrahim Sayed-Tabatabaei, Kazuhiro Sato, Mohammad Pourkheirandish, Takao Komatsuda. Mapping of QTL for intermedium spike on barley chromosome 4H using EST-based markers. <i>Breed. Sci.</i> 59(4), 383–390(2009)
802	Yerlan Turuspekov, Ichiro Honda, Yoshiaki Watanabe, Nils Stein, Takao Komatsuda. An inverted and micro-colinear genomic regions of rice and barley carrying the cly1 gene for cleistogamy. <i>Breed. Sci.</i> 59(5), 657–663(2009)
803	Guoxiong Chen, Mohammad Pourkheirandish, Mohammad Sameri, Ning Wang, Sudha Nair, Yulan Shi, Chao Li, Eviatar Nevo, Takao Komatsuda. Genetic targeting of candidate genes for drought sensitive gene eibi1 of wild barley (<i>Hordeum spontaneum</i>). <i>Breed. Sci.</i> 59(5), 637–644(2009)
804	Kazuhiro Sato, Takashi Matsumoto, Natsuko Ooe, Kazuyoshi Takeda. Genetic analysis of seed dormancy QTL in barley. <i>Breed. Sci.</i> 59(5), 645–650(2009)
805	A. Oikawa, K. Nagai, K. Kato, S. Kidou. Gene silencing of barley P23k involved in secondary wall formation causes abnormal tiller formation and intercalary elongation. <i>Breed. Sci.</i> 59(5), 664–670(2009)
806	Takao Komatsuda, Bjoern Salomon, Roland von Bothmer. Evolutionary process of <i>Hordeum brachyantherum</i> 6x and related tetraploid species revealed by nuclear DNA sequences. <i>Breed. Sci.</i> 59(5), 611–616(2009)
807	Daisuke Saisho, Mohammad Pourkheirandish, Hiroyuki Kanamori, Takashi Matsumoto, Takao Komatsuda. Allelic variation of row type gene Vrs1 in barley and implication of the functional divergence. <i>Breed. Sci.</i> 59(5), 621–628(2009)
808	Ikuo Nakamura, Bhuwan Rai, Hiroko Takahashi, Kenji Kato, Yo-ichiro Sato, Takao Komatsuda. <i>Aegilops</i> section Sitopsis species contains the introgressive PolA1 gene with a closer relationship to that of <i>Hordeum</i> than <i>Triticum-Aegilops</i> species. <i>Breed. Sci.</i> 59(5), 602–610(2009)
809	H Buerstnayr, T. Ban, J. Anderson. QTL mapping and marker-assisted selection for <i>Fusarium</i> head blight resistance in wheat: a review. <i>Plant Breed</i> 128(1), 1–26(2009)
810	S. Takumi, Y. Naka, H. Morihiro, Y. Matsuoka. Expression of morphological and flowering time variation through allopolyploidization: an empirical study with 27 wheat synthetics and their parental <i>Aegilops tauschii</i> accessions. <i>Plant Breed</i> 128(6), 585–590(2009)
811	Katsuyuki Kakeda, Toshiro Ibuki, Junko Suzuki, Hidetaka Tadano, Yuko Kurita, Yosuke Hanai, Yasuo Kowayama. Molecular and genetic characterization of the S locus in <i>Hordeum bulbosum</i> L., a wild self-incompatible species related to cultivated barley. <i>Mol Genet Genomics</i> 280(6), 509–519(2008)
812	A. Torada, M. Koike, S. Ikeguchi, I. Tsutsui. Mapping of a major locus controlling seed dormancy using backcrossed progenies in wheat (<i>Triticum aestivum</i> L.). <i>Genome</i> 51(6), 426–432(2008)

813	S Taketa, S Amano, Y Tsujino, T Sato, D Saisho, K Kakeda, M Nomura, T Suzuki, T Matsumoto, K Sato, H Kanamori, S Kawasaki, K. Takeda. Barley grain with adhering hulls is controlled by an ERF family transcription factor gene regulating a lipid biosynthesis pathway. <i>Proc Natl Acad Sci U S A</i> 105(10), 4062–4067(2008)
814	Hirokazu Handa, Nobukazu Namiki, Donghe Xu, Tomohiro Ban. Dissecting of the FHB resistance QTL on the short arm of wheat chromosome 2D using a comparative genomic approach: from QTL to candidate gene. <i>Mol. Breed.</i> 22(1), 71–84(2008)
815	K. Tanno, R. von Bothmer, K. Yamane, K. Takeda, T. Komatsuda. Analysis of DNA sequence polymorphism at the cMWG699 locus reveals phylogenetic relationships and allopolyploidy within <i>Hordeum murinum</i> subspecies. <i>Hereditas</i> 147(1), 34–42(2010)
816	Y. Kurahashi, A. Terashima, S. Takumi. Variation in Dehydration Tolerance, ABA Sensitivity and Related Gene Expression Patterns in D–Genome Progenitor and Synthetic Hexaploid Wheat Lines. <i>Int J Mol Sci</i> 10(6), 2733–2751(2009)
817	S. Taketa, K. Matsuki, S. Amano, D. Saisho, E. Himi, N. Shitsukawa, T. Yuo, K. Noda, K. Takeda. Duplicate polyphenol oxidase genes on barley chromosome 2H and their functional differentiation in the phenol reaction of spikes and grains. <i>J Exp Bot</i> 61(14), 3983–3993(2010)
818	G. Sun, T. Komatsuda. Origin of the Y genome in <i>Elymus</i> and its relationship to other genomes in Triticeae based on evidence from elongation factor G (EF–G) gene sequences. <i>Mol Phylogenet Evol</i> 56(2), 727–733(2010)
819	T. Kajimura, N. Mizuno, S. Takumi. Utility of leaf senescence–associated gene homologs as developmental markers in common wheat. <i>Plant Physiol Biochem</i> 48(10–11), 851–859(2010)
820	N. Mizuno, N. Hosogi, P. Park, S. Takumi. Hypersensitive response–like reaction is associated with hybrid necrosis in interspecific crosses between tetraploid wheat and <i>Aegilops tauschii</i> coss. <i>PLoS One</i> 5(6), e11326(2010)
821	Y. Katayose, H. Kanamori, M. Shimomura, H. Ohyanagi, H. Ikawa, H. Minami, M. Shibata, T. Ito, K. Kurita, K. Ito, Y. Tsubokura, A. Kaga, J. Wu, T. Matsumoto, K. Harada, T. Sasaki. DaizuBase, an integrated soybean genome database including BAC–based physical maps. <i>Breed Sci</i> 61(5), 661–664(2012)
822	Z. Xia, S. Watanabe, T. Yamada, Y. Tsubokura, H. Nakashima, H. Zhai, T. Anai, S. Sato, T. Yamazaki, S. Lu, H. Wu, S. Tabata, K. Harada. Positional cloning and characterization reveal the molecular basis for soybean maturity locus E1 that regulates photoperiodic flowering. <i>Proc Natl Acad Sci U S A</i> 109(32), E2155–2164(2012)
823	S. Shiro, A. Yamamoto, Y. Umehara, M. Hayashi, N. Yoshida, A. Nishiwaki, T. Yamakawa, Y. Saeki. Effect of rj genotype and cultivation temperature on the community structure of soybean–nodulating bradyrhizobia. <i>Appl Environ Microbiol</i> 78(4), 1243–1250(2012)
824	K. Komatsu, T.–Y. Hwang, M. Takahashi, T. Sayama, H. Funatsuki, N. Oki, M. Ishimoto. Identification of QTL controlling post–flowering period in soybean. <i>Breed. Sci.</i> 61(5), 646–652(2012)
825	A. Kaga, T. Shimizu, S. Watanabe, Y. Tsubokura, Y. Katayose, K. Harada, D. Vaughan, N. Tomooka. Evaluation of soybean germplasm conserved in NIAS genebank and development of mini core collections. <i>Breed. Sci.</i> 61(5), 566–592(2012)
826	T. Yamada, M. Hajika, N. Yamada, K. Hirata, A. Okabe, N. Oki, K. Takahashi, K. Seki, K. Okano, Y. Fujita, A. Kaga, T. Shimizu, T. Sayama, M. Ishimoto. Effects on flowering and seed yield of dominant alleles at maturity loci E2 and E3 in a Japanese cultivar, Enrei. <i>Breed. Sci.</i> 61(5), 653–660(2012)
827	S. Watanabe, Z. Xia, R. Hideshima, Y. Tsubokura, S. Sato, N. Yamanaka, R. Takahashi, T. Anai, S. Tabata, K. Kitamura, K. Harada. A map–based cloning strategy employing a residual heterozygous line reveals that the GIGANTEA gene is involved in soybean maturity and flowering. <i>Genetics</i> 188(2), 395–407(2011)
828	T. Sugimoto, S. Yoshida, A. Kaga, M. Hajika, K. Watanabe, M. Aino, K. Tatsuda, R. Yamamoto, T. Matoh, D.R. Walker, A.R. Biggs, M. Ishimoto. Genetic analysis and identification of DNA markers linked to a novel <i>Phytophthora sojae</i> resistance gene in the Japanese soybean cultivar Waseshiroge. <i>Euphytica</i> 182(1), 133–145(2011)

829	Iwata H, Hayashi T, Terakami S, Takada N, Sawamura Y, Yamamoto T. Potential assessment of genome-wide association study and genomic selection in Japanese pear <i>Pyrus pyrifolia</i> . <i>Breeding Science</i> , in press(2012)
830	H. Fukuoka, K. Miyatake, T. Nunome, S. Negoro, K. Shirasawa, S. Isobe, E. Asamizu, H. Yamaguchi, A. Ohyama. Development of gene-based markers and construction of an integrated linkage map in eggplant by using <i>Solanum</i> orthologous (SOL) gene sets. <i>Theor Appl Genet</i> 125(1), 47–56(2012)
831	Hayashi T, Ohyama A, Iwata H. Bayesian QTL mapping for recombinant inbred lines derived from a four-way cross. <i>Euphytica</i> 183, 277–287(2012)
832	Shirasawa K, Hirakawa H, DNA marker applications to molecular genetics and genomics in tomato, <i>Breed. Sci.</i> , <i>in press</i> , (2013)