

Field Resistance of Six Cultivars of Winter Oilseed Rape against Turnip Yellow Mosaic Virus

JOSEF ŠPAK and DARINA KUBELKOVÁ

Department of Plant Virology, Institute of Plant Molecular Biology, Academy of Sciences
of the Czech Republic, České Budějovice, Czech Republic

Abstract

ŠPAK J., KUBELKOVÁ D. (2002): **Field resistance of six cultivars of winter oilseed rape against Turnip yellow mosaic virus.** Plant Protect. Sci., **38**: 73–75.

The resistance of six cultivars of winter oilseed rape (SL 509, SL 507, Darmor, Solida, Jet Neuf, Silesia) against *Turnip yellow mosaic virus* was studied. The number of over-wintering plants and plants with symptoms of TYMV infection were monitored. All plants were tested by the double diffusion test in agar and by DAS-ELISA to prove infection. ELISA was the most sensitive method, revealing 32–76% of latent virus infections in individual cultivars. The results clearly illustrated that monitoring of plant symptoms is not sufficient to prove TYMV infection in the field and that sensitive, large scale methods like ELISA must be employed to obtain reliable data.

Keywords: *Turnip yellow mosaic virus*; oilseed rape; *Brassica*; ELISA; serology

Abbreviations: **DAS-ELISA** = double antibody sandwich enzyme-linked immunosorbent assay; **PBS** = phosphate-buffered saline; **TYMV** = *Turnip yellow mosaic virus*

Turnip yellow mosaic virus (TYMV) is a seed (DE ASSIS & SHERWOOD 2000) and beetle transmitted virus infecting a wide range of brassicas world-wide (RAYBOULD *et al.* 1999). In our previous studies we estimated losses caused by artificial TYMV infection in several winter oilseed rape and fodder turnip rape cultivars (POLÁK *et al.* 1989) and proved seed transmission of TYMV in these crops (ŠPAK *et al.* 1993). We also searched for natural sources of the virus (PELIKÁNOVÁ *et al.* 1990; ŠPAK *et al.* 1991) and found 50–90% incidence of TYMV in the fodder turnip rape cultivar Perko PVH in the field (ŠPAK & POLÁK 1985). In contrast to Germany (SCHRÖEDER 1994), where TYMV has been observed in low percentage in spring and winter oilseed rape, we have never found naturally TYMV infected winter oilseed rape in the Czech Republic. Also, infected seedlings raised from TYMV-infected seed in our previous experiments remained symptomless (ŠPAK *et al.* 1993).

The aim of this study was to induce an artificial field infection in six oilseed cultivars, to thus test their resistance and the reliability of symptoms, serology and ELISA for the diagnosis of the virus.

MATERIALS AND METHODS

The seed of six winter oilseed rape cultivars (Table 1) were purchased from the Research and Breeding Station Slapy at Tábor. One hundred seeds of each cultivar were sown at the end of August in experimental plots. At the five true leaf stage the number of plants in each plot was reduced to 50. Plants were dusted with carborundum and inoculated with strain TYMV 2 (ŠPAK & POLÁK 1988) using a glass rod. The inoculum was prepared from 1 g of leaves with symptoms of infection ground in a mortar in 2 ml of 0.1 M phosphate buffer pH 7.0. In May of the following year, the number of over-wintering plants and

plants with symptoms of TYMV infection were monitored. All plants were tested by the double diffusion test in agar and by DAS-ELISA to prove the infection.

Double diffusion tests were performed in 0.9% Difco Noble agar in 0.01 M McIlvaine buffer pH 7.0 with 0.02% sodium azide in a Petri dish. Antiserum diluted 1:200 was placed in a central well and antigens (sap from infected plants) in surrounding wells. Antigens were prepared by homogenisation of leaves in 0.01 M McIlvaine buffer pH 7.0 with 0.02% sodium azide, in a 1:1 ratio (w/v). After 24–48 h serological reactions were evaluated and stained with amido black 10B. Antiserum with the titre 1:2048, prepared in our laboratory against the TYMV 2 strain (ŠPAK & POLÁK 1988) was used. DAS-ELISA was performed as described in (ŠPAK *et al.* 1993). Samples with the absorbance at 405 nm higher than $x + 3s$ (x = value of healthy control, s = standard deviation) were scored as positive.

RESULTS AND DISCUSSION

The results summarised in Table 1 show that the overwintering of all cultivars was similar, being the best in cv. Silesia and the worst in cv. Solida. Symptoms of virus infection, not very distinct pale yellowish-green mosaic on leaves, appeared in all cultivars. The number of symptomatic plants was lowest in Silesia. In cv. Solida false symptoms were scored as positive in two plants, as shown later by serology and ELISA. No spur formation was observed in double diffusion serological tests. This indicates that there had been no natural infection of the experimental plants by another strain of the virus occur-

ring in nature (PELIKÁNOVÁ *et al.* 1990; ŠPAK *et al.* 1991) and transmitted e.g. by Phyllotreta beetles (PROESLER 1971).

Although the match between diagnostic methods was best between symptoms and serology (Table 2), ELISA was the most sensitive method, revealing 32–76% of latent virus infections in individual cultivars. Mean absorbance values in positive, negative and healthy control plants were 1.39, 0.18 and 0.08, respectively. We did not observe a significant difference in mean absorbance values (standard deviations) among individual cultivars, which were in cvs SL 509 1.50 (0.15), SL 507 1.15 (0.30), Darmor 1.42 (0.34), Solida 1.29 (0.24), Jet Neuf 1.58 (0.04), Silesia 1.41 (0.29). Although the level of infected plants was the lowest in SL 509, we found no distinct difference in the resistance of the cultivars. The high frequency of latent infections could explain why natural infection of winter oilseed rape has not yet been observed in the Czech Republic.

Recently, seed transmission of brassica viruses (DE ASSIS & SHERWOOD 2000) and their occurrence in natural populations (RAYBOULD *et al.* 1999; MASKELL *et al.* 1999) has attracted interest with respect to risk evaluation of large scale utilisation of genetically modified brassica crops in the field. Similar studies are confronted with a lack of epidemiological data obtained in field studies with cultivated and naturally infected brassicas including weeds. Our results clearly illustrated that monitoring of plant symptoms is not sufficient to prove TYMV infection in the field and that in similar studies sensitive, large scale methods like ELISA must be employed to obtain reliable data.

Table 1. Overwintering and detection of *Turnip yellow mosaic virus* in oilseed rape cultivars

	SL 509		SL 507		Darmor		Solida		Jet Neuf		Silesia	
	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)	N	(%)
Autumn	50	100	50	100	50	100	50	100	50	100	50	100
Spring	38	76	38	76	39	78	34	68	37	74	46	92
Symptoms	13	26	16	32	12	24	7	14	12	24	5	10
Serology positive	16	32	26	52	25	50	5	10	27	54	6	12
ELISA positive	32	64	49	98	48	96	43	86	46	92	43	86

N = number of plants

Table 2. Match of diagnostic methods of *Turnip yellow mosaic virus* in oilseed rape (% of plants)

	SL 509	SL 507	Darmor	Solida	Jet Neuf	Silesia
Symptoms + serology	82	62	48	72	44	84
Symptoms + ELISA	40	32	25	16	26	12
Serology + ELISA	50	56	52	12	58	14

References

- DE ASSIS F.M., SHERWOOD J.L. (2000): Evaluation of seed transmission of turnip yellow mosaic virus and tobacco mosaic virus in *Arabidopsis thaliana*. *Phytopathology*, **90**: 1233–1238.
- MASKELL L.C., RAYBOULD A.F., COOPER J.I. *et al.* (1999): Effects of turnip mosaic virus and turnip yellow mosaic virus on the survival, growth and reproduction of wild cabbage (*Brassica oleracea*). *Ann. Appl. Biol.*, **135**: 401–407.
- PELIKÁNOVÁ J., ŠPAK J., KUBELKOVÁ D. (1990): Comparison of the virulence and serological relationship between the isolate of turnip yellow mosaic virus from *Alliaria officinalis* L. with strains isolated from cultivated brassicas. *Ochr. Rostl.*, **26**: 161–165.
- POLÁK Z., ŠPAK J., PROCHÁZKOVÁ Z., VENCLOVÁ D. (1989): Yield losses of winter rape and turnip rape after infection with some viruses. *Ochr. Rostl.*, **25**: 1–7.
- PROESELER G. (1971): Transmission of the turnip yellow mosaic virus by insects with biting mouthparts. *Arch. Phytopath. Pfl.-Schutz.*, **7**: 391–397.
- RAYBOULD A.F., MASKELL L.C., EDWARDS M.L. *et al.* (1999): The prevalence and spatial distribution of viruses in natural populations of *Brassica oleracea*. *New Phytol.*, **141**: 265–275.
- SCHRÖDER M. (1994): Investigation on the susceptibility of oilseed rape (*Brassica napus* L., ssp. *napus*) to different virus diseases. *Pfl.-Krankh. Pfl.-Schutz.*, **101**: 576–589.
- ŠPAK J., DUSBÁBKOVÁ J., KUBELKOVÁ D., NEČÁSEK J. (1991): Resistance of transformed and non-transformed oilseed rape cv. HM-81 to the infection with cauliflower mosaic, turnip yellow mosaic and turnip mosaic viruses. *Biol. Plant.*, **33**: 234–239.
- ŠPAK J., KOVÁŘOVÁ D., HNILIČKA E. (1993): Seed transmission of turnip yellow mosaic virus in winter turnip and winter oilseed rapes. *Ann. Appl. Biol.*, **123**: 33–35.
- ŠPAK J., POLÁK Z. (1985): The occurrence of turnip mosaic and turnip yellow mosaic viruses on rapes and winter turnip rape. *Ochr. Rostl.*, **21**: 13–20.
- ŠPAK J., POLÁK J. (1988): Identification of six isolates belonging to three serologically distinct strains of turnip yellow mosaic virus on brassicas in Czechoslovakia. *Ochr. Rostl.*, **24**: 251–258.

Received for publication May 24, 2002
Accepted after corrections June 20, 2002

Abstrakt

ŠPAK J., KUBELKOVÁ D. (2002): **Polní rezistence šesti odrůd řepky ozimé vůči Turnip yellow mosaic virus**. *Plant Protect. Sci.*, **38**: 73–75.

V polním pokusu bylo testováno 6 odrůd řepky ozimé (SL 509, SL 507, Darmor, Solida, Jet Neuf, Silesia) na odolnost k viru žluté mozaiky vodnice (TYMV). Výsledky přezimování rostlin, příznaků na listech, sérologie a ELISA testů a vzájemné korelačních vztahy těchto faktorů (%) prokázaly vlastnosti jednotlivých odrůd a hlavně spolehlivost diagnostických metod. Nejlépe přezimovala odrůda Silesia. Příznaky na listech nelze považovat za spolehlivého ukazatele infekce. Jako nejspolehlivější metoda detekce TYMV se osvědčila DAS-ELISA.

Klíčová slova: virus žluté mozaiky vodnice; řepka ozimá; *Brassica*; ELISA; sérologie

Corresponding address:

Doc. Ing. JOSEF ŠPAK, DrSc., Ústav molekulární biologie rostlin Akademie věd České republiky, Branišovská 31, 370 05 České Budějovice, Česká republika
tel.: + 420 38 77 75 54, fax: +420 38 530 03 56, e-mail: spak@umbr.cas.cz