

## LEIOCEPHALIDAE

1989 Leiocephalinae Frost and Etheridge, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 81: 45. Type genus: *Leiocephalus* Gray, 1825.

2001 Leiocephalidae—Frost, Etheridge, Janies, and Titus, Am. Mus. Novit., 3343: 13.

Range: Cuba and nearby islands, the Cayman Islands, the Bahama Islands, Hispaniola, Navassa, and Martinique. Recently extinct species were found on Barbuda and Antigua, Anguilla, Guadeloupe, Hispaniola, Puerto Rico, and Jamaica.

### *Leiocephalus* Gray

1827 *Leiocephalus* Gray, Philos. Mag., London, 2: 207. Type species: *Leiocephalus carinatus* Gray, 1827. (subsequent designation by Fitzinger, Syst. Rept.: 17)

1837 *Holotropis* Duméril and Bibron, Erp. Gen., 4: 259. Type species: none designated.

1837 *Pristinotus* Gravenhorst, Nova Acta Phys. Med. Acad. Caes. Leopold. Carol. Nat. Curios., Halle, 18: 738. Type species (by monotypy): *Pristinotus schreibersii* Gravenhorst, 1837.

1843 *Steironotus* (*Steironotus*)—Fitzinger, Syst. Rept.: 70. Type species (by original designation): *Tropidurus schreibersii* Fitzinger 1826.

1843 *Steironotus* (*Leiocephalus*)—Fitzinger, Syst. Rept.: 70.

1885 *Liocephalus*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 160 (part).

1928 *Hispaniolus* Cochran, Proc. Biol. Soc. Washington, 41: 50. Type species (by original designation): *Hispaniolus pratensis* Cochran, 1928.

1966 *Leiocephalus*—Etheridge, Copeia, 1964: 79.

1992 *Leiocephalus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 1.

Range: Cuba and nearby islands, the Cayman Islands, the Bahama Islands, Hispaniola, and Martinique. Recently extinct species were found on Barbuda and Antigua, Hispaniola, Puerto Rico, and Jamaica.

### *Leiocephalus anonymus* Pregill

1984 *Leiocephalus anonymus* Pregill, Proc. Biol. Soc. Washington, 97: 828, Fig. 1. Type locality: "an unspecified cave(s) near St. Michel de l'Atalye, Département L'Artibonite, Haiti" (Natl. Mus. Nat. Hist. Vert. Paleontol. No. 340182, a right dentary).

1992 *Leiocephalus anonymous*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 37. Incorrect subsequent spelling.

Range: Known only from the type locality in Haiti.

Note: Known only from fossil material but thought to have gone extinct no later than latest Pleistocene.

### *Leiocephalus apertosulcus* Etheridge

1965 *Leiocephalus apertosulcus* Etheridge, Q. J. Florida Acad. Sci., 29: 91.  
Type locality: "Stratum 2, cave in Cerro de San Francisco, Municipio Pedro Santana, Provincia San Rafael, Republica Dominicana" (Holotype: Mus. Comp. Zool. Vert. Paleontol. No. 3404, a right dentary).

1992 *Leiocephalus apertosulcus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 37.

Range: Known only from the type locality in the Dominican Republic.

Note: Known only from fossil material but Stratum 2 of the type locality was estimated to be no older than 4000 ypb.

*Leiocephalus barahonensis* Schmidt

1921 *Leiocephalus barahonensis* Schmidt, Bull. Am. Mus. Nat. Hist., 44: 15; Fig. 12. Type locality: "Barahona, Santo Domingo" (Holotype (original designation): Am. Mus. Nat. Hist. No. 2736).

1967 *Leiocephalus barahonensis*—Schwartz, Tulane Stud. Zool., 14: 34.

1992 *Leiocephalus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 38.

Range: Hispaniola, in southeastern Haiti and southwestern República Dominicana; Isla Alta Velo.

*Leiocephalus barahonensis barahonensis* Schmidt

1932 *Leiocephalus personatus barahonensis*—Cochran, Proc. Biol. Soc. Washington, 45: 177.

1967 *Leiocephalus barahonensis barahonensis*—Schwartz, Tulane Stud. Zool., 14: 35; Fig. 11a.

1992 *Leiocephalus barahonensis barahonensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

Range: República Dominicana from near El Naranjo in the west to Paraíso in the south, both to the north and east of the Sierra de Baoruco.

*Leiocephalus barahonensis altavelensis* Noble and Hassler

1933 *Leiocephalus altavelensis* Noble and Hassler, Am. Mus. Novit., 652: 14.  
Type locality: "Alta Vela Island" (Holotype: Am. Mus. Nat. Hist. No. 51055).

1941 *Leiocephalus personatus altavelensis*—Cochran, Bull. U.S. Natl. Mus., 177: 220; Pl. 7b; Fig. 66a.

1967 *Leiocephalus vinculum altavelensis*—Schwartz, Tulane Stud. Zool., 14: 46; Fig. 12c.

1992 *Leiocephalus barahonensis altavelensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

Range: Isla Alto Velo, República Dominicana.

*Leiocephalus barahonensis aureus* Cochran

1934 *Leiocephalus personatus aureus* Cochran, Occas. Pap. Boston Soc. Nat. Hist., 8: 175. Type locality: "Jacmel, Haiti" (Holotype: U.S. Natl. Mus. No. 75909).

1967 *Leiocephalus barahonensis aureus*—Schwartz, Tulane Stud. Zool., 14: 39; Fig. 11c.

1992 *Leiocephalus barahonensis aureus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

Range: Southeast coast of Haiti from 12 miles southwest of Jacmel east to the vicinity of Pedernales, República Dominicana.

*Leiocephalus barahonensis beatanus* Noble

1923 *Leiocephalus beatanus* Noble, Am. Mus. Novit., 64: 5. Type locality: "Beata Island, Dominican Republic" (Holotype (original designation): Am. Mus. Nat. Hist. No. 24330).

1932 *Leiocephalus personatus beatanus*—Cochran, Proc. Biol. Soc. Washington, 45: 178.

1967 *Leiocephalus barahonensis beatanus*—Schwartz, Tulane Stud. Zool., 14: 41; Fig. 11d.

1992 *Leiocephalus barahonensis beatanus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

Range: Isla Beata, República Dominicana.

*Leiocephalus barahonensis oxygaster* Schwartz

1967 *Leiocephalus barahonensis oxygaster* Schwartz, Tulane Stud. Zool., 14: 36. Type locality: "13.1 mi. (21.1 km) SW Enriquillo, Pedernales Province, República Dominicana" (Holotype: Mus. Comp. Zool. No. 81098).

1992 *Leiocephalus barahonensis oxygaster*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

Range: Península de Barahona from south of Enriquillo in the east to about 22 km southeast of Pedernales in the west, República Dominicana.

*Leiocephalus carinatus* Gray

1827 *Leiocephalus carinatus* Gray, Philos. Mag., London, 2: 208. Type locality: "Cuba" (Holotype: Brit. Mus. Nat. Hist. No. xxii.96a (RR 1946.8.29.75)). Restricted type locality (Schwartz and Ogren, Herpetologica, 12: 102): "Habana, Habana Province, Cuba".

1843 *Steironotus (Leiocephalus) carinatus*—Fitzinger, Syst. Rept.: 70.

1885 *Liocephalus carinatus*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 165.

1914 *Leiocephalus carinatus*—Barbour, Mem. Mus. Comp. Zool., 44: 300.

1992 *Leiocephalus carinatus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

1999 *Leiocephalus carinatus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 115.

Range: Cuba and the Isla de Juventud (= Pinos), the Cayman Islands, and the Bahama Islands.

*Leiocephalus carinatus carinatus* Gray

1837 *Holotropis microlophus* Cocteau in Duméril and Bibron, Erp. Gen., 4: 264. Type locality: "Cuba" (Lectotype (Schwartz, Copeia, 1969: 620): Mus.

- Natl. Hist. Nat. Paris No. 2392). (syn. fide Gray, Ann. Mag. Nat. Hist., Ser. 1, 5: 109).
- 1845 *Leiocephalus macLeayii* Gray, Cat. Spec. Liz. Coll. Brit. Mus.: 218. Type locality: "Cuba" (Syntypes: Brit. Mus. Nat. Hist. No. xxii.97a (RR 1946.8.11.82), xxii.97b (RR 1946.8.10.58)), syn. fide Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 165)
- 1935 *Leiocephalus carinatus carinatus*—Barbour and Shreve, Proc. Boston Soc. Nat. Hist., 40: 360.
- 1959 *Leiocephalus carinatus carinatus*—Schwartz, Reading Public Mus. Art Gallery Sci. Publ. , 10: 16; Fig. 9-11.
- 1992 *Leiocephalus carinatus carinatus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.
- 1999 *Leiocephalus carinatus carinatus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 115.
- Range: The provinces of Pinar del Río, Habana, and Matanzas, western Cuba.

*Leiocephalus carinatus aquarius* Schwartz and Ogren

- 1956 *Leiocephalus carinatus aquarius* Schwartz and Ogren, Herpetologica, 12: 100. Type locality: "Aguadores, near Santiago de Cuba, Oriente Prov., Cuba" (Holotype: Charleston Mus. No. 55.1.62).
- 1992 *Leiocephalus carinatus aquarius*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.
- 1999 *Leiocephalus carinatus aquarius*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 117.
- Range: Eastern Cuba on the ousthern coast of Granma, Santiago de Cuba, and Guantanamo provinces from Cabo Cruz to Punta de Maisí.

*Leiocephalus carinatus armouri* Barbour and Shreve

- 1935 *Leiocephalus carinatus armouri* Barbour and Shreve, Proc. Boston Soc. Nat. Hist., 40: 360. Type locality: "High Rock, Grand Bahama Island, Bahamas" (Holotype: Mus. Comp. Zool. No. 38090).
- 1992 *Leiocephalus carinatus armouri*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.
- Range: Grand Bahama Island (including Wood Cay, Little Sale Cay, Stranger's Cay), Little Abaco Island (including Pensacola Cay, Elbow Cay, Green Turtle Cay) in the northern Bahama Islands; introduced in Florida on Virginia Key and Key Biscayne, Dade County, and Palm Beach, Palm Beach County.

*Leiocephalus carinatus cayensis* Schwartz

- 1959 *Leiocephalus carinatus cayensis* Schwartz, Reading Public Mus. Art Gallery Sci. Publ. , 10: 38; Fig. 15-17. Type locality: "lighthouse on Cayo Cachiboca, Jardin de la Reina, Camagüey Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 77758).
- 1992 *Leiocephalus carinatus cayensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

1999 *Leiocephalus carinatus cayensis*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 117.  
Range: Jardines de la Reina (including Cayo Cachiboca, Cayo Grenada, Cayo Caballones, Cayo Grande, and other unnamed cays) off the south coast of Camagüey Province, Cuba.

*Leiocephalus carinatus coryi* Schmidt

1936 *Leiocephalus carinatus coryi* Schmidt, Field Mus. Nat. Hist. Publ., Zool. Ser., 20: 129; Fig. 11. Type locality: "Bimini Islands, Bahamas" (Holotype: Field Mus. Nat. Hist. No. 260).

1992 *Leiocephalus carinatus coryi*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

Range: The islands of North Bimini, South Bimini, East Bimini, Andros, and Berry (including Great Harbour Cay, Devil's Cay, Frazer's Hog Cay, Cat Cay), and Easter Cay, all in the western Bahamas.

*Leiocephalus carinatus granti* Rabb

1957 *Leiocephalus carinatus granti* Rabb, Herpetologica, 13: 109. Type locality: "Cayman Brac, British West Indies" (Holotype: Univ. Mich. Mus. Zool. No. 114494).

1992 *Leiocephalus carinatus granti*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

Range: Little Cayman Island and Cayman Brac.

*Leiocephalus carinatus hodsdoni* Schmidt

1936 *Leiocephalus carinatus hodsdoni* Schmidt, Field Mus. Nat. Hist. Publ., Zool. Ser., 20: 130; Fig. 12. Type locality: "Salt Pond, Long Island, Bahamas" (Holotype: Field Mus. Nat. Hist. No. 22572).

1992 *Leiocephalus carinatus hodsoni*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

Range: Cat Island, Goat Cay off Little San Salvador, Long Island (including Violet Cay), Guana Cay, Pinders Cay, Cay Verde, and the Ragged Islands (including Flamingo Cay, South Channel Cay, Johnson Cay, Knife Cay, and Little Ragged Island), all in the central Bahamas.

*Leiocephalus carinatus labrossytus* Schwartz

1959 *Leiocephalus carinatus labrossytus* Schwartz, Reading Public Mus. Art Gallery Sci. Publ., 10: 33; Fig. 12-14. Type locality: "five kilometers southeast of Paso Caballo, Las Villas Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 77757).

1992 *Leiocephalus carinatus labrossytus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

1999 *Leiocephalus carinatus carinatus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 117.

Range: Central Cuba, southern region of Matanzas, Cienfuegos, and Sancti Spiritus provinces Cienfuegos in the west, from Península de Zapata to Casilda, including the Sierra de Trinidad.

*Leiocephalus carinatus microcyon* Schwartz

1959 *Leiocephalus carinatus microcyon* Schwartz, Reading Public Mus. Art Gallery Sci. Publ. , 10: 43. Type locality: "Caleta Grande, Isla de Pinos, Habana Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 81271).

1992 *Leiocephalus carinatus microcyon*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

1999 *Leiocephalus carinatus microcyon*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 117.

Range: Isla de Juventud (= Pinos).

*Leiocephalus carinatus mogotensis* Schwartz

1959 *Leiocephalus carinatus mogotensis* Schwartz, Reading Public Mus. Art Gallery Sci. Publ. , 10: 12; Fig. 6-8. Type locality: "Cueva del Cable, San Vicente, Pinar del Río Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 77755).

1992 *Leiocephalus carinatus mogotensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

1999 *Leiocephalus carinatus mogotensis*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 115.

Range: Known only from a restricted area in the Sierra de los Organos in the San Vicente-Viñales-Valle de Ancón region of Pinar del Río Province, western Cuba.

*Leiocephalus carinatus varius* Garman

1887 *Liocephalus varius* Garman, Proc. Am. Philos. Soc., 24: 274. Type locality: "Grand Cayman, British West Indies" (Syntypes: U.S. Natl. Mus. No. 52405; Mus. Comp. Zool. No. 6023 (3)).

1914 *Leiocephalus varius*—Barbour, Mem. Mus. Comp. Zool., 44: 300.

1937 *Leiocephalus carinatus varius*—Barbour, Mem. Mus. Comp. Zool., 44: 135.

1992 *Leiocephalus carinatus varius*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

Range: Grand Cayman Island and Swan Island.

*Leiocephalus carinatus virescens* Stejneger

1901 *Leiocephalus virescens* Stejneger, Proc. U.S. Natl. Mus., 23: 471. Type locality: "Green Cay, Bahamas" (Holotype: U.S. Natl. Mus. No. 26758).

1937 *Leiocephalus carinatus virescens*—Barbour, Bull. Mus. Comp. Zool., 82: 135.

1992 *Leiocephalus carinatus virescnes*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

Range: Green Cay, Eleuthera Island, and the Exuma Cays (including Ship Channel Cay, SW Allan's Cay, Leaf Cay, Warderick Wells Cay, Compass Cay, Triple Cay, Great Exuma Island, and Elizabeth Island), all in the central Bahamas.

*Leiocephalus carinatus zayasi* Schwartz

1959 *Leiocephalus carinatus zayasi* Schwartz, Reading Public Mus. Art Gallery Sci. Publ. , 10: 9; Fig. 3-5. Type locality: "north shore of Ensenada de Corrientes, Pinar del Río Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 77756).

1992 *Leiocephalus carinatus zayasi*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 40.

1999 *Leiocephalus carinatus zayasi*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 117.

Range: Península de Guanahacabibes, from the type locality east to 10 kilometers southeast of Manuel Lazo, Pinar del Río Province, extreme western Cuba.

*Leiocephalus cubensis* (Gray)

1840 *Tropidurus (Leiolaemus) cubensis* Gray, Ann. Mag. Nat. Hist., Ser. 1, 5: 110. Type locality: "Cuba" (Holotype: Brit. Mus. Nat. Hist. No. xxiii.98a). Restricted type locality (Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 104): "vicinity of Guanabacoa, Habana Province, Cuba".

1856 *Holotropis vittatus* Hallowell, Proc. Acad. Nat. Sci. Philadelphia, 8: 151. Type locality: "Cuba" (Holotype: Acad. Nat. Sci. Philad., lost fide Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 104). (syn. fide Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 104)

1862 *Leiocephalus (Holotropis) vittatus*—Reinhardt and Lütken, Vidensk. Medd. Dansk Naturhist. Foren., Ser. 2, 4: 271.

1917 *Leiocephalus cubensis*—Stejneger, Proc. U.S. Natl. Mus., 53: 273.

1959 *Leiocephalus cubensis*—Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 104; Fig. 3-5.

1992 *Leiocephalus cubensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 41.

1999 *Leiocephalus cubensis*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 123.

Range: Cuba and nearby islands, and the Isla de Juventud (= Pinos), Archipiélago de Sabana-Camaguey and Jardines de la Reina.

*Leiocephalus cubensis cubensis* (Gray)

1959 *Leiocephalus cubensis cubensis*—Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 107; Fig. 4.

1992 *Leiocephalus cubensis dubensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 41.

1999 *Leiocephalus cubensis cubensis*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 125.

Range: Cuba, from Artemisia to Sagua de Tánamo.

*Leiocephalus cubensis gigas* Schwartz

1959 *Leiocephalus cubensis gigas* Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 113. Type locality: "Caleta Grande, Isla de Pinos, Habana Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 81056).

1992 *Leiocephalus cubensis gigas*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 41.

1999 *Leiocephalus cubensis cubensis*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 125.

Range: South coast of the Isla de Juventud (= Pinos).

*Leiocephalus cubensis minor* Garrido

1970 *Leiocephalus cubensis minor* Garrido in Varona and Garrido, Poeyana, 75: 18. Type locality: "Cayo Juan García, Cayos de San Felipe, Archipiélago de los Canarreos, Pinar del Río" (Holotype: Inst. Zool. Acad. Cien. Habana No. 2754).

1992 *Leiocephalus cubensis minor*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 41.

Range: Known only from the type locality off the southwest coast of Cuba.

*Leiocephalus cubensis pambasileus* Schwartz

1959 *Leiocephalus cubensis pambasileus* Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 118. Type locality: "Cayo Hicacos, Habana Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 81068).

1992 *Leiocephalus cubensis pambasileus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 41.

1999 *Leiocephalus cubensis pambasileus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 125.

Range: Archipiélago de los Canarreos (Cayo Hicacos and Cayo Campos) off the southwestern coast of Cuba.

*Leiocephalus cubensis paraphrus* Schwartz

1959 *Leiocephalus cubensis paraphrus* Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 111; Fig. 5. Type locality: "southernmost point of large unnamed key 3 kilometers northwest of Cayo Cachiboca lighthouse, Camagüey Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 78005).

1992 *Leiocephalus cubensis paraphrus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 41.

1999 *Leiocephalus cubensis paraphrus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 125.

Range: Jardines de la Reina off the south coast of Camagüey Province, Cuba.

*Leiocephalus cuneus* Etheridge

1964 *Leiocephalus cuneus* Etheridge, Bull. Florida State Mus., Biol. Sci., 9: 48. Type locality: "Cave V, Two Foot Bay on the northeast coast of

Barbuda, British West Indies" (Holotype: Florida State Mus. Vert. Paleontol. No. 8226, a left dentary).

1992 *Leiocephalus cuneus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 41.

Range: Late Pleistocene of Barbuda and Antigua, and possibly Anguilla and Guadeloupe.

*Leiocephalus endomychus* Schwartz

1967 *Leiocephalus vinculum endomychus* Schwartz, Tulane Stud. Zool., 14: 45; Fig. 12b. Type locality: "3.4 mi. (5.5 km) NE Barrage de Péligré, 1100 feet (361 meters), Dépt. de l'Ouest, Haiti" (Holotype: Mus. Comp. Zool. No. 81099).

1992 *Leiocephalus endomychus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 41.

Range: Known only from the type locality.

*Leiocephalus eremitus* Cope

1868 *Liocephalus eremitus* Cope, Proc. Acad. Nat. Sci. Philadelphia, 20: 122. Type locality: "island of Navassa" (Holotype: U.S. Natl. Mus. No. 12016).

1966 *Leiocephalus eremitus*—Thomas, J. Ohio Herpetol. Soc., 5: 80.

1992 *Leiocephalus eremitus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 41.

Range: Navassa Island, extinct fide Thomas, J. Ohio Herpetol. Soc., 5: 80.

*Leiocephalus etheridgei* Pregill

1981 *Leiocephalus etheridgei* Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 71: 35. Type locality: "Backbone 1 Cave, 1.2 km S Barrio de Barahona, Municipio de Morovis, Puerto Rico (18° 20' 57" N, 66° 26' 47" W). Late Pleistocene" (Holotype: Natl. Mus. Nat. Hist. Vert. Paleontol. No. 259190, a right dentary).

1992 *Leiocephalus etheridgei*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 43.

Range: Known only from the type locality from Late Pleistocene fossil material.

*Leiocephalus greenwayi* Barbour and Shreve

1935 *Leiocephalus greenwayi* Barbour and Shreve, Proc. Boston Soc. Nat. Hist., 40: 358. Type locality: "East Plana Cay, Bahamas" (Holotype: Mus. Comp. Zool. No. 36711).

1992 *Leiocephalus greenwayi*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 43.

Range: Known only from the type locality in the southeastern Bahamas.

*Leiocephalus herminieri* (Duméril and Bibron)

1837 *Holotropis Herminieri* Duméril and Bibron, Erp. Gen., 4: 261; Pl. 44. Type locality: "les îles de la Trinité et de la Martinique" (Syntypes: Mus. Hist. Nat. Paris No. 1826, 2389, 6829).

- 1845 *Leiocephalus Herminieri*—Gray, Cat. Spec. Liz. Coll. Brit. Mus.: 217.  
1885 *Liocephalus herminieri*—Boulenger, Cat. Liz. Brit. Mus. Nat. Hist., 2: 166.  
1964 *Leiocephalus herminieri*—Etheridge, Bull. Florida State Mus., Biol. Sci., 9: 56.  
1992 *Leiocephalus herminieri*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 43.

Range: Martinique, extinct.

*Leiocephalus inaguae* Cochran

- 1931 *Leiocephalus inaguae* Cochran, J. Washington Acad. Sci., 21: 39. Type locality: "Man of War Bay, Great Inagua Island" (Holotype: U.S. Natl. Mus. No. 81277).  
1967 *Leiocephalus inaguae*—Schwartz, Ann. Carnegie Mus., 39: 156.  
1992 *Leiocephalus inaguae*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 44.

Range: Great Inagua Island in the southern Bahamas.

*Leiocephalus loxogrammus* Cope

- 1887 *Liocephalus loxogrammus* Cope, Proc. U.S. Natl. Mus., 10: 437. Type locality: "Rum Cay, Bahamas" (Syntypes: U.S. Natl. Mus. No. 14569 (10); Mus. Comp. Zool. No. 10931).  
1914 *Leiocephalus loxogrammus*—Barbour, Mem. Mus. Comp. Zool., 44: 300.  
1992 *Leiocephalus loxogrammus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 45.

Range: Rum Cay and San Salvador in the eastern central Bahamas; possibly Conception Island.

*Leiocephalus loxogrammus loxogrammus* Cope

- 1935 *Leiocephalus loxogrammus loxogrammus*—Barbour and Shreve, Proc. Boston Soc. Nat. Hist., 40: 359.  
1992 *Leiocephalus loxogrammus loxogrammus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 46.

Range: Rum Cay, Bahamas.

*Leiocephalus loxogrammus parnelli* Barbour and Shreve

- 1935 *Leiocephalus loxogrammus parnelli* Barbour and Shreve, Proc. Boston Soc. Nat. Hist., 40: 359. Type locality: "San Salvador or Watling's Island, Bahamas" (Holotype: Mus. Comp. Zool. No. 36748).  
1992 *Leiocephalus loxogrammus parnelli*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 46.

Range: San Salvador, Bahamas.

*Leiocephalus lunatus* Cochran

- 1934 *Leiocephalus personatus lunatus* Cochran, Occas. Pap. Boston Soc. Nat. Hist., 8: 153. Type locality: "Santo Domingo City, Dominican Republic" (Holotype: Field Mus. Nat. Hist. No. 166).

- 1967 *Leiocephalus lunatus*—Schwartz, Tulane Stud. Zool., 14: 24; Fig. 7, 8.  
1992 *Leiocephalus lunatus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 46.

Range: Southern coast and nearby islands of eastern República Dominicana.

*Leiocephalus lunatus lunatus* Cochran

- 1939 *Leiocephalus personatus lunatus*—Mertens, Abh. Senckenb. Naturforsch. Ges., 449: 47; Pl. 8, Fig. 35b, 36d.

- 1967 *Leiocephalus lunatus lunatus*—Schwartz, Tulane Stud. Zool., 14: 24; Fig. 7a, 8a.

- 1992 *Leiocephalus lunatus lunatus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 46.

Range: South coast of República Dominicana between the Río Haina and the Río Ozama, Distrito Nacional.

*Leiocephalus lunatus arenicolor* Mertens

- 1939 *Leiocephalus personatus arenicolor* Mertens, Abh. Senckenb. Naturforsch. Ges., 449: 48; Pl. 8, Fig. 32c, 33, 35c, 36c. Type locality: "Sandstrand bei San Pedro de Macoris" (Holotype: Senckenb. Mus. Frankfurt No. 25715).

- 1967 *Leiocephalus lunatus arenicolor*—Schwartz, Tulane Stud. Zool., 14: 27, Fig. 7b, 8b.

- 1992 *Leiocephalus lunatus arenicolor*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 46.

Range: Southeastern coast of the República Dominicana from San Pedro de Macorís to Boca Chavón, La Altagracia Province.

*Leiocephalus lunatus lewisi* Schwartz

- 1969 *Leiocephalus lunatus lewisi* Schwartz, J. Herpetol., 3: 80. Type locality: "0.9 km E Boca Chica, Distrito Nacional, República Dominicana" (Holotype: Carnegie Mus. No. 45867).

- 1992 *Leiocephalus lunatus lewisi*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 46.

Range: Between the Río Ozama and 7 miles east of Boca Chica, República Dominicana.

*Leiocephalus lunatus louisae* Cochran

- 1934 *Leiocephalus personatus louisae* Cochran, Occas. Pap. Boston Soc. Nat. Hist., 8: 177. Type locality: "Saona Island, Dominican Republic" (Holotype: Mus. Comp. Zool. No. 37551).

- 1967 *Leiocephalus lunatus louisae*—Schwartz, Tulane Stud. Zool., 14: 32; Fig. 7e, 8e.

- 1992 *Leiocephalus lunatus louisae*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 46.

Range: Isla Saona, República Dominicana.

*Leiocephalus lunatus melaenacelis* Schwartz

1967 *Leiocephalus lunatus melaenacelis* Schwartz, Tulane Stud. Zool., 14: 29; Fig. 7c, 8c. Type locality: "western end, Isla Catalina, La Romana Province, República Dominicana" (Holotype: Mus. Comp. Zool. No. 81096).

1992 *Leiocephalus lunatus melaenascelis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 46.

Range: Isla Catalina, off the coast near La Romana, República Dominicana.

*Leiocephalus lunatus thomasi* Schwartz

1967 *Leiocephalus lunatus thomasi* Schwartz, Tulane Stud. Zool., 14: 31; Fig. 7d, 8d. Type locality: "0.5 mi. (0.8 km) NW Boca de Yuma, La Romana Province, República Dominicana" (Holotype: Mus. Comp. Zool. No. 81097).

1992 *Leiocephalus lunatus thomasi*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 46.

Range: Known only from the vicinity of the type locality in southeastern República Dominicana.

*Leiocephalus macropus* Cope

1862 *Liocephalus macropus* Cope, Proc. Acad. Nat. Sci. Philadelphia, 14: 184. Type locality: "Eastern Cuba" (Lectotype (Hardy, J. Washington Acad. Sci., 48: 294): U.S. Natl. Mus. No. 25819). Restricted type locality (Stejneger, Proc. U.S. Natl. Mus., 53: 274): "Monte Verde in Eastern Cuba".

1914 *Leiocephalus macropus*—Barbour, Mem. Mus. Comp. Zool., 44: 301.

1967 *Leiocephalus macropus*—Schwartz and Garrido, Reading Public Mus. Art Gallery, Sci. Publ., 14: 5.

1992 *Leiocephalus macropus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.

1999 *Leiocephalus macropus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 134.

Range: Widespread in isolated populations throughout Cuba.

*Leiocephalus macropus macropus* Cope

1958 *Leiocephalus macropus macropus*—Hardy, J. Washington Acad. Sci., 48: 294; Fig. 1, 5a-f.

1967 *Leiocephalus macropus macropus*—Schwartz and Garrido, Reading Public Mus. Art Gallery, Sci. Publ., 14: 24; Fig. 4c.

1992 *Leiocephalus macropus macropus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.

1999 *Leiocephalus macropus macropus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 137.

Range: Southern coast of the Santiago de Cuba and Guantánamo province, from Bahia de Santiago de Cuba to Cajobabo.

*Leiocephalus macropus aegialus* Schwartz and Garrido

- 1967 *Leiocephalus macropus aegialus* Schwartz and Garrido, Reading Public Mus. Art Gallery, Sci. Publ., 14: 15. Type locality: "Playa Santa Lucía, Camagüey Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 83255).
- 1992 *Leiocephalus macropus aegialus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.
- 1999 *Leiocephalus macropus aegialus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 137.
- Range: Known only from the type locality.

*Leiocephalus macropus asbolomus* Schwartz and Garrido

- 1967 *Leiocephalus macropus asbolomus* Schwartz and Garrido, Reading Public Mus. Art Gallery, Sci. Publ., 14: 30; Fig. 4d. Type locality: "El Guayabo, Baracoa, Oriente Province, Cuba" (Holotype: Inst. Zool. Acad. Cien. Habana No. 568).
- 1992 *Leiocephalus macropus asbolomus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.
- 1999 *Leiocephalus macropus asbolomus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 138.
- Range: Cuba, northern coast of Holguín and Guantánamo provinces, from Guardalavaca to the Río Yumuri, in Baracoa.

*Leiocephalus macropus felinoi* Garrido

- 1979 *Leiocephalus macropus felinoi* Garrido, Poeyana, 188: 2; Fig. 1A, 2A, 3A. Type locality: "Bacunayagua, Provincia de Matanzas, Cuba" (Holotype: Inst. Zool. Acad. Cien. Cuba No. 4751).
- 1992 *Leiocephalus macropus felinoi*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.
- 1999 *Leiocephalus macropus felinoi*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 138.
- Range: Known only from the type locality.

*Leiocephalus macropus hoplites* Zug

- 1959 *Leiocephalus macropus hoplites* Zug, Proc. Biol. Soc. Washington, 72: 140; Fig. 1, 2. Type locality: "12 mi. E. Morón, Loma de Cunagua, Camagüey Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 78020).
- 1967 *Leiocephalus macropus hoplites*—Schwartz and Garrido, Reading Public Mus. Art Gallery, Sci. Publ., 14: 12; Fig. 2b, 3c.
- 1992 *Leiocephalus macropus hoplites*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.
- 1999 *Leiocephalus macropus hoplites*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 137.
- Range: Known only from the type locality in northwestern Camagüey Province, Cuba.

*Leiocephalus macropus hyacinthurus* Zug

- 1959 *Leiocephalus macropus hyacinthurus* Zug, Proc. Biol. Soc. Washington, 72: 145; Fig. 5, 6. Type locality: "Finca La Pastora, 2 kilometers northwest of Trinidad, Las Villas Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 78015).
- 1967 *Leiocephalus macropus hyacinthurus*—Schwartz and Garrido, Reading Public Mus. Art Gallery, Sci. Publ., 14: 10; Fig. 2c, 3b.
- 1992 *Leiocephalus macropus hyacinthurus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.
- 1999 *Leiocephalus macropus hyacinthurus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 137.
- Range: Known only from the type locality in southern Las Villas Province, Cuba.

*Leiocephalus macropus immaculatus* Hardy

- 1958 *Leiocephalus macropus immaculatus* Hardy, J. Washington Acad. Sci., 48: 294; Fig. 1, 5g-l. Type locality: "vicinity of Ocuja, Oriente Province, Cuba" (Holotype: U.S. Natl. Mus. No. 138412).
- 1967 *Leiocephalus macropus immaculatus*—Schwartz and Garrido, Reading Public Mus. Art Gallery, Sci. Publ., 14: 21.
- 1992 *Leiocephalus macropus immaculatus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.
- 1999 *Leiocephalus macropus immaculatus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 137.
- Range: The southern coast of Santiago de Cuba Province from the Río Magdalena east to Bahía de Santiago de Cuba, eastern Cuba.

*Leiocephalus macropus koopmani* Zug

- 1959 *Leiocephalus macropus koopmani* Zug, Proc. Biol. Soc. Washington, 72: 146; Fig. 7, 8. Type locality: "near the base of Cabo Corrientes, Pinar del Río Province, Cuba" (Holotype: Mus. Comp. Zool. No. 55541).
- 1967 *Leiocephalus macropus koopmani*—Schwartz and Garrido, Reading Public Mus. Art Gallery, Sci. Publ., 14: 7; Fig. 2a, 3a.
- 1992 *Leiocephalus macropus koopmani*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.
- 1999 *Leiocephalus macropus hyacinthurus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 137.
- Range: Península de Guanahacabibes, Pinar del Río Province, east to the vicinity of Manuel Lazo, Cuba.

*Leiocephalus macropus lenticulatus* Garrido

- 1973 *Leiocephalus macropus lenticulatus* Garrido, Torreia, N.S., Habana, 30:10. Type locality: "el arenal Los Cocos, a 6 km de Gibara" (Holotype: Inst. Zool. Acad. Cien. Habana No. 2782).
- 1992 *Leiocephalus macropus lenticulatus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.

1999 *Leiocephalus macropus lenticulatus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 138.

Range: Known only from the type locality in eastern Cuba.

*Leiocephalus macropus phylax* Schwartz and Garrido

1967 *Leiocephalus macropus phylax* Schwartz and Garrido, Reading Public Mus. Art Gallery, Sci. Publ., 14: 17; Fig. 4c. Type locality: "Verreón, near Cabo Cruz, Oriente Province, Cuba" (Holotype: Inst. Zool. Acad. Cien. Habana No. 556).

1992 *Leiocephalus macropus phylax*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.

1999 *Leiocephalus macropus phylax*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 138.

Range: Southern coast of Granma Province from the type locality and Cabo Cruz to the Río Puercos and Punta Hicacos, Cuba.

*Leiocephalus macropus torrei* Garrido

1979 *Leiocephalus macropus torrei* Garrido, Poeyana, 188: 7; Fig. 1B, 2B, 3B. Type locality: "las faldas de la loma del Río Paredones, San Miguel de los Baños, Provincia de Matanzas, Cuba" (Holotype: Inst. Zool. Acad. Cien. Cuba No. 3725).

1992 *Leiocephalus macropus torrei*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.

1999 *Leiocephalus macropus torrei*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 138.

Range: Known only from the vicinity of the type locality.

*Leiocephalus melanochlorus* Cope

1862 *Liocephalus melanochlorus* Cope, Proc. Acad. Nat. Sci. Philadelphia, 14: 184. Type locality: "Hayti, near Jeremie" (Syntypes: Mus. Comp. Zool. No. 3598 (10), 84341-50; U.S. Natl. Mus. No. 53402; Cal. Acad. Sci. No. 39392).

1914 *Leiocephalus melanochlorus*—Barbour, Mem. Mus. Comp. Zool., 44: 301.

1966 *Leiocephalus melanochlorus*—Schwartz, J. Ohio Herpetol. Soc., 5: 41.

1992 *Leiocephalus melanochlorus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 47.

Range: The Tiburon Peninsula of southwestern Haiti; Ile-à-Vache.

*Leiocephalus melanochlorus melanochlorus* Cope

1966 *Leiocephalus melanochlorus melanochlorus*—Schwartz, J. Ohio Herpetol. Soc., 5: 41.

1992 *Leiocephalus melanochlorus melanochlorus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 48.

Range: Western part of the Tiburon Peninsula, east to St. Michel du Sud, Dépt. du Sud; also the Ile-à-Vache, Haiti.

*Leiocephalus melanochlorus hypsistus* Schwartz

1966 *Leiocephalus melanochlorus hypsistus* Schwartz, J. Ohio Herpetol. Soc., 5: 44. Type locality: "Furcy, 5600 feet, Dépt. del'Ouest, Haiti" (Holotype: Mus. Comp. Zool. No. 81063).

1992 *Leiocephalus melanochlorus hypsistus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 48.

Range: The Montagne Noire and the southern slope of the Massif de la Selle, and Morne de Cayette near the coast, Dépt. de l'Ouest, Haiti.

*Leiocephalus onaneyi* Garrido

1973 *Leiocephalus onaneyi* Garrido, Poeyana, 116: 4; Fig. 1-3. Type locality: "la cima de la loma de Macambo, entre San Antonio del Sur e Imías, Provincia de Oriente, Cuba" (Holotype: Inst. Zool. Acad. Cien. Habana No. 2869).

1992 *Leiocephalus onaneyi*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 48.

1999 *Leiocephalus onaneyi*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 152.

Range: Known only from the type locality in eastern Cuba.

*Leiocephalus partitus* Pregill

1981 *Leiocephalus partitus* Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 71: 39. Type locality: "Guanica Bat Cave, Reserva Forestal Guanica, 6 km E Barrio de Guanicao, Municipio de Guayanilla, Puerto Rico (17° 57' 40" N, 66° 50' 53" W)" (Holotype: Natl. Mus. Nat. Hist. Vert. Paleontol. No. 259203, a right dentary).

1992 *Leiocephalus partitus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 48.

Range: Known only from the type locality and from Cuevo del Perro, both on Puerto Rico, from Late Pleistocene to sub-Recent fossil material.

*Leiocephalus personatus* Cope

1862 *Liocephalus personatus* Cope, Proc. Acad. Nat. Sci. Philadelphia, 14: 182. Type locality: "Hayti (near Jeremie)" (Syntypes: Mus. Comp. Zool. No. 3615 (2)).

1862 *Liocephalus trigeminatus* Cope, Proc. Acad. Nat. Sci. Philadelphia, 14: 183. Type locality: "Hayti, near Jeremie" (Holotype: Mus. Comp. Zool., now lost). (syn. fide Cochran, Bull. U.S. Natl. Mus., 177: 212)

1941 *Leiocephalus personatus*—Cochran, Bull. U.S. Natl. Mus., 177: 212.

1967 *Leiocephalus personatus*—Schwartz, Tulane Stud. Zool., 14: 4; Fig. 2, 4, 5.

1992 *Leiocephalus personatus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 48.

Range: Haiti and the República Dominicana.

*Leiocephalus personatus personatus* Cope

- 1932 *Leiocephalus personatus personatus*—Cochran, Proc. Biol. Soc. Washington, 45: 177.
- 1967 *Leiocephalus personatus personatus*—Schwartz, Tulane Stud. Zool., 14: 4.
- 1992 *Leiocephalus personatus personatus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: The northern littoral of the Tiburon Peninsula in Haiti, from the type locality in the west to Léogâne, Dépt. de l'Ouest in the east.

*Leiocephalus personatus actites* Schwartz

- 1967 *Leiocephalus personatus actites* Schwartz, Tulane Stud. Zool., 14: 14; Fig. 5d. Type locality: "Sosúa, Puerto Plata Province, República Dominicana" (Holotype: Mus. Comp. Zool. No. 81088).
- 1992 *Leiocephalus personatus actites*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: Along the northern coast of the República Dominicana from Punta Rucia in the west to near Sabaneta de Yásica in the east, Puerto Plata Province.

*Leiocephalus personatus agraulus* Schwartz

- 1967 *Leiocephalus personatus agraulus* Schwartz, Tulane Stud. Zool., 14: 21. Type locality: "1 mi. WSW Constanza, 4000 feet (1311 meters), La Vega Province, República Dominicana" (Holotype: Mus. Comp. Zool. No. 81090).
- 1992 *Leiocephalus personatus agraulus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: Interior uplands of the Cordillera Central in the Valle de Constanza and Valle de Tireo, and the southern slope of the Cordillera Central north of San Juan, República Dominicana.

*Leiocephalus personatus budeni* Schwartz

- 1967 *Leiocephalus personatus budeni* Schwartz, Tulane Stud. Zool., 14: 19; Fig. 5e. Type locality: "12 km NE Jarabacoa, 2000 feet (656 meters), La Vega Province, República Dominicana" (Holotype: Mus. Comp. Zool. No. 81089).
- 1992 *Leiocephalus personatus budeni*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: Known only from the type locality on the north slope of the Cordillera Central, República Dominicana.

*Leiocephalus personatus elattoprosopon* Gali, Schwartz, and Suarez

- 1988 *Leiocephalus personatus elattoprosopon* Gali, Schwartz, and Suarez, Proc. Biol. Soc. Washington, 101: 1. Type locality: "ca. 1 km inland, basal portion of the Morne Dubois 'peninsula,' Dépt. de Sud, Haiti" (Holotype: U.S. Natl. Mus. No. 197370).

1992 *Leiocephalus personatus elattoprosopon*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: Known only from the vicinity of the type locality.

*Leiocephalus personatus mentalis* Cochran

1932 *Leiocephalus personatus mentalis* Cochran, Proc. Biol. Soc. Washington, 45: 178. Type locality: "Jovéro, Dominican Republic" (Holotype: U.S. Natl. Mus. No. 65772).

1967 *Leiocephalus personatus mentalis*—Schwartz, Tulane Stud. Zool., 14: 9; Fig. 4b.

Range: From the type locality east to Juanillo, La Altagracia Province, República Dominicana.

*Leiocephalus personatus poikilometes* Schwartz

1969 *Leiocephalus personatus poikilometes* Schwartz, J. Herpetol., 3: 82. Type locality: "10 km SE El Jorillo, 2050 feet (625 meters), San Juan Province, República Dominicana" (Holotype: U.S. Natl. Mus. No. 165935).

1992 *Leiocephalus personatus poikilometes*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: The northern range of the Sierra de Neiba, and the floor of the Valle de San Juan in the vicinity of Barranca, San Juan Province, República Dominicana.

*Leiocephalus personatus pyrrholaemus* Schwartz

1971 *Leiocephalus personatus pyrrholaemus* Schwartz, Herpetologica, 27: 178. Type locality: "9 km E. Las Galeras, Samaná Province, República Dominicana" (Holotype: Carnegie Mus. No. 52287).

1992 *Leiocephalus personatus pyrrholaemus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: The Peninsula de Samaná, República Dominicana.

*Leiocephalus personatus scalaris* Cochran

1932 *Leiocephalus personatus scalaris* Cochran, Proc. Biol. Soc. Washington, 45: 181. Type locality: "Cap Haitien, Haiti" (Holotype: U.S. Natl. Mus. No. 74054).

1939 *Leiocephalus personatus pulcherrimus* Mertens, Abh. Senckenb. Naturforsch. Ges., 449: 50; Fig. 32b, 35e, 36a. Type locality: "Moncion, 452 m H." (Holotype: Senck. Mus. Frankfurt No. 25757). (syn. fide Schwartz, Tulane Stud. Zool., 14: 16)

1967 *Leiocephalus personatus scalaris*—Schwartz, Tulane Stud. Zool., 14: 16.

1992 *Leiocephalus personatus scalaris*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

1992 *Leiocephalus personatus pulcherrimus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: Northern Hispaniola, from the vicinity of Carosse, near Port Margot, on the northern Haitian coast and inland to Dondon and St. Michel de

l'Atalaye, eastward along the coast to Monte Cristi; inland in the Valle de Cibao to the vicinity of Santiago; inland in Haiti south to Cerca-la-Source and in the República Dominicana to Bánica; Cayos Siete Hermanos (Isla Monte Chico), and Isla Cabras off the coast at Monte Cristi.

*Leiocephalus personatus socoensis* Gali and Schwartz

1982 *Leiocephalus personatus socoensis* Gali and Schwartz, J. Herpetol., 16: 177. Type locality: "25 km E San Pedro de Macorís, Río Cumayasa, La Romana Province, República Dominicana" (Holotype: U.S. Natl. Mus. Nat. Hist. 197371).

1992 *Leiocephalus personatus socoensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: Between the Río Soco and the Río Cumayas, La Romana Province, República Dominicana

*Leiocephalus personatus tarachodes* Schwartz

1967 *Leiocephalus personatus tarachodes* Schwartz, Tulane Stud. Zool., 14: 11; Fig. 5c. Type locality: "6 km SE Nagua, María Trinidad Sánchez Province, República Dominicana" (Holotype: Mus. Comp. Zool. No. 81087).

1992 *Leiocephalus personatus tarachodes*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: From the vicinity of Nagua southeastward to Sabana de la Mar and Hato Mayor, west to the vicinity of Moca and Salcedo, República Dominicana.

*Leiocephalus personatus trujilloensis* Mertens

1939 *Leiocephalus personatus trujilloensis* Mertens, Abh. Senckenb. Naturforsch. Ges., 449: 449: 45; Pl. 8, Fig. 32a, 35a, 36e. Type locality: "Küste am Deutsch-Dominikanischen Tropenforschungs-Institute in Ciudad Trujillo, etwa 4 km westlich der Ozama-Mündung" (Holotype: Senck. Mus. Frankfurt No. 26213).

1967 *Leiocephalus personatus trujilloensis*—Schwartz, Tulane Stud. Zool., 14: 7; Fig. 5b.

1992 *Leiocephalus personatus trujilloensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: South-central República Dominicana from Limonal, Peravia Province, and Sabana Grande de Palenque, San Cristobál Province, east to the Río Ozama, and into the interior to the vicinity of Villa Altagracia, San Cristobál Province, and to near Pedregal, Distrito Nacional.

*Leiocephalus pratensis* (Cochran)

1928 *Hispaniolus pratensis* Cochran, Proc. Biol. Soc. Washington, 41: 50. Type locality: "Atalaye plantation, near St. Michel, Département du Nord, Republic of Haiti" (Holotype: U.S. Natl. Mus. No. 69189). Emended type locality (Schwartz, J. Herpetol., 1: 54): "Atalaye plantation near St. Michel de l'Atalaye, Dépt. de l'Artibonite, Haiti".

- 1941 *Hispaniolus pratensis*—Cochran, Bull. U.S. Natl. Mus., 177: 241; Fig. 66e.  
 1966 *Leiocephalus pratensis*—Etheridge, Copeia, 1966: 79.  
 1968 *Leiocephalus pratensis*—Schwartz, J. Herpetol., 1: 54.  
 1992 *Leiocephalus pratensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: Known from the vicinity of the type locality and Ile à Cabrit in the Golfe de la Gonâve, Haiti.

*Leiocephalus pratensis pratensis* (Cochran)

- 1979 *Leiocephalus pratensis pratensis*—Schwartz, Herpetologica, 35: 252; Fig. 1.  
 1992 *Leiocephalus pratensis pratensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: Known from the vicinity of the type locality.

*Leiocephalus pratensis chimarus* Schwartz

- 1979 *Leiocephalus pratensis chimarus* Schwartz, Herpetologica, 35: 251; Fig. 1.  
 Type locality: "Ile à Cabrit off the Haitian coast, Dépt. de l'Ouest, Haiti" (Holotype: U.S. Natl. Mus. No. 197335).  
 1992 *Leiocephalus pratensis chimarus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 49.

Range: Ile à Cabrit off the Haitian coast in the Golfe de la Gonâve.

*Leiocephalus psammmodromus* Barbour

- 1916 *Liocephalus arenarius* Barbour, Proc. Biol. Soc. Washington, 29: 217.  
 Type locality: "Bastion Cay, Turks Islands, B. W. I." (Holotype: Mus. Comp. Zool. No. 11948) (nec *Steironotus arenarius* Tschudi 1845 = *Ophryoessoides arenarius*)  
 1920 *Leiocephalus psammmodromus* Barbour, Copeia, 85: 73. (substitute name for *Liocephalus arenarius* Barbour, 1916)  
 1967 *Leiocephalus arenarius*—Schwartz, Ann. Carnegie Mus., 39: 159.  
 1992 *Leiocephalus psammmodromus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 50.

Range: Islands of the Caicos Bank and Turks Bank in the southeastern Bahamas.

*Leiocephalus psammmodromus psammmodromus* Barbour

- 1967 *Leiocephalus arenarius arenarius*—Schwartz, Ann. Carnegie Mus., 39: 160; Fig. 2.  
 1975 *Leiocephalus psammmodromus psammmodromus*—Schwartz and Thomas, Spec. Publ. Carnegie Mus. Nat. Hist., 1: 136.  
 1992 *Leiocephalus psammmodromus psammmodromus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 50.

Range: "Bastion Cay", Big Sand Cay, Turks Islands in the eastern Bahamas.

*Leiocephalus psammodromus aphretor* Schwartz

1967 *Leiocephalus arenarius aphretor* Schwartz, Ann. Carnegie Mus., 39: 163; Fig. 3. Type locality: "Long Cay, southeast of Grand Turk Island, Turks Islands" (Holotype: Carnegie Mus. No. 40602).

1975 *Leiocephalus psammodromus aphretor*—Schwartz and Thomas, Spec. Publ. Carnegie Mus. Nat. Hist., 1: 136.

1992 *Leiocephalus psammodromus aphretor*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 50.

Range: Known only from the type locality in the eastern Bahamas.

*Leiocephalus psammodromus apocrinus* Schwartz

1967 *Leiocephalus arenarius apocrinus* Schwartz, Ann. Carnegie Mus., 39: 165; Fig. 4. Type locality: "Big Ambergris Cay, northwest side, Caicos Islands" (Holotype: Carnegie Mus. No. 40601).

1975 *Leiocephalus psammodromus apocrinus*—Schwartz and Thomas, Spec. Publ. Carnegie Mus. Nat. Hist., 1: 136.

1992 *Leiocephalus psammodromus apocrinus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 50.

Range: Big Ambergris Cay and Little Ambergris Cay, Caicos Islands, in the eastern Bahamas.

*Leiocephalus psammodromus cacodoxus* Schwartz

1967 *Leiocephalus arenarius cacodoxus* Schwartz, Ann. Carnegie Mus., 39: 176; Fig. 7. Type locality: "Providenciales Island, Caicos Islands" (Holotype: Mus. Comp. Zool. No. 54185).

1975 *Leiocephalus psammodromus cacodoxus*—Schwartz and Thomas, Spec. Publ. Carnegie Mus. Nat. Hist., 1: 136.

1992 *Leiocephalus psammodromus cacodoxus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 50.

Range: Ft. George Cay, Providenciales Island, and Sugar Loaf Island, all in the Caicos Islands in the southeastern Bahamas.

*Leiocephalus psammodromus hyphantus* Schwartz

1967 *Leiocephalus arenarius hyphantus* Schwartz, Ann. Carnegie Mus., 39: 172; Fig. 6. Type locality: "Pine Cay, Caicos Islands" (Holotype: Univ. Mich. Mus. Zool. No. 126624).

1975 *Leiocephalus psammodromus hyphantus*—Schwartz and Thomas, Spec. Publ. Carnegie Mus. Nat. Hist., 1: 136.

1992 *Leiocephalus psammodromus hyphantus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 50.

Range: Pine Cay, Water Cay, and Stubb Cay, all in the Caicos Islands in the southeastern Bahamas.

*Leiocephalus psammodromus mounax* Schwartz

- 1967 *Leiocephalus arenarius mounax* Schwartz, Ann. Carnegie Mus., 39: 169; Fig. 5. Type locality: "Long Cay, off Cockburn Harbour, South Caicos, Caicos Islands" (Holotype: Carnegie Mus. No. 40603).
- 1975 *Leiocephalus psammodromus mounax*—Schwartz and Thomas, Spec. Publ. Carnegie Mus. Nat. Hist., 1: 136.
- 1992 *Leiocephalus psammodromus mounax*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 50.

Range: Known only from the type locality in the southeastern Bahamas.

*Leiocephalus punctatus* Cochran

- 1931 *Leiocephalus carinatus punctatus* Cochran, J. Washington Acad. Sci., 21: 39. Type locality: "north shore of the bay at Jamaica Wells, Acklin Island" (Holotype: U.S. Natl. Mus. No. 81560).
- 1935 *Leiocephalus carinatus helenae* Barbour and Shreve, Proc. Boston Soc. Nat. Hist., 40: 359. Type locality: "South Cay, Mira Por Vos Islands, Bahamas" (Holotype: Mus. Comp. Zool. No. 38110). (syn. fide Schwartz and Thomas, Spec. Publ. Carnegie Mus. Nat. Hist., 1: 136)
- 1935 *Leiocephalus carinatus picinus* Barbour and Shreve, Proc. Boston Soc. Nat. Hist., 40: 360. Type locality: "Atwood's Cay, Bahamas" (Holotype: Mus. Comp. Zool. No. 38120). (syn. fide Schwartz and Thomas, Spec. Publ. Carnegie Mus. Nat. Hist., 1: 137)
- 1966 *Leiocephalus punctatus*—Etheridge, Copeia, 1966: 79.
- 1975 *Leiocephalus punctatus*—Schwartz and Thomas, Spec. Publ. Carnegie Mus. Nat. Hist., 1: 136.
- 1992 *Leiocephalus punctatus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 50.

Range: Samana Cay, Crooked Island, Goat Cay, Fortune Island, Acklin's Island, Castle Island, Mira Por Vos Island, North Cay, Fish Cay, and Guana Cay, all in the central Bahamas.

*Leiocephalus raviceps* Cope

- 1862 *Liocephalus raviceps* Cope, Proc. Acad. Nat. Sci. Philadelphia, 14: 183. Type locality: "Eastern Cuba" (Syntypes: U.S. Natl. Mus. No. 4162 (5); Mus. Comp. Zool. No. 10928; Acad. Nat. Sci. Philad. No. 8601-3). Restricted type locality (Gundlach, Contr. Erpetol. Cubana: 34): "en la montaña de Guantánamo".
- 1917 *Leiocephalus raviceps*—Stejneger, Proc. U.S. Natl. Mus., 53: 274.
- 1960 *Leiocephalus raviceps*—Schwartz, Proc. Biol. Soc. Washington, 73: 67.
- 1992 *Leiocephalus raviceps*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 50.
- 1999 *Leiocephalus raviceps*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 130.

Range: Cuba.

*Leiocephalus raviceps raviceps* Cope

- 1960 *Leiocephalus raviceps raviceps*—Schwartz, Proc. Biol. Soc. Washington, 73: 74; Fig. 2.
- 1992 *Leiocephalus raviceps raviceps*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 51.
- 1999 *Leiocephalus raviceps raviceps*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 131.
- Range: From the Bahía de Guantánamo east to north of Cajobabo, eastern Cuba.

*Leiocephalus raviceps delavarai* Garrido

- 1973 *Leiocephalus raviceps delavarai* Garrido, Torreia, N.S., Habana, 30: 4.  
Type locality: "el arenal Los Cocos, a seis km de Gibara" (Holotype: Inst. Zool. Acad. Cien. Habana No. 2774).
- 1992 *Leiocephalus raviceps delavarai*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 51.
- 1999 *Leiocephalus raviceps jaumei*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 132.
- Range: Known only from the type locality in eastern Cuba.

*Leiocephalus raviceps jaumei* Schwartz and Garrido

- 1968 *Leiocephalus raviceps jaumei* Schwartz and Garrido, Proc. Biol. Soc. Washington, 81: 24. Type locality: "San Waldo, 4 km N Cortés, on the road between Cortés and Isabel Rubio, Pinar del Río Province, Cuba" (Holotype: Inst. Zool. Acad. Cien. Habana No. 1349).
- 1992 *Leiocephalus raviceps jaumei*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 51.
- 1999 *Leiocephalus raviceps jaumei*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 132.
- Range: Known only from the vicinity of the type locality in western Cuba.

*Leiocephalus raviceps klinikowskii* Schwartz

- 1960 *Leiocephalus raviceps klinikowskii* Schwartz, Proc. Biol. Soc. Washington, 73: 77; Fig. 3. Type locality: "4.5 kilometers southwest of Varadero Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 83326).
- 1992 *Leiocephalus raviceps klinikowskii*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 50.
- 1999 *Leiocephalus raviceps klinikowskii*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 131.
- Range: The Península de Hicacos, northern Matanzas Province, Cuba.

*Leiocephalus raviceps uzzelli* Schwartz

- 1960 *Leiocephalus raviceps uzzelli* Schwartz, Proc. Biol. Soc. Washington, 73: 70; Fig. 1. Type locality: "18.2 kilometers east of Siboney, Oriente Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 79321).
- 1992 *Leiocephalus raviceps uzzelli*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 51.

1999 *Leiocephalus raviceps uzzelli*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 131.  
Range: Along the coast from the Bahía de Guantánamo, Guantánamo Province, west to La Socapa, Santiago de Cuba Province, eastern Cuba.

*Leiocephalus rhutidira* Schwartz

1979 *Leiocephalus rhutidira* Schwartz, Proc. Biol. Soc. Washington, 92: 273; Fig. 1. Type locality: "Lapierre, 10.6 km W Ça Soleil, 122 m, Département de l'Artibonite, Haiti" (Holotype: Carnegie Mus. Nat. Hist. No. 60520).  
1992 *Leiocephalus rhutidira*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 51.

Range: The southern coast of Presqu'île du Nord Ouest, northwestern Haiti.

*Leiocephalus schreibersii* (Gravenhorst)

1826 *Tropidurus schreibersii* Fitzinger, Neue Class. Rept.: 49.(nomen nudum)  
1837 *Pristinotus schreibersii* Gravenhorst, Nova Acta Phys. Med. Acad. Caes. Leopold. Carol. Nat. Curios., Halle, 18: 739; Pl. 54, Fig. 15, 16. Type locality: "St. Domingo" (Holotype: ?Breslau Mus., not located). Restricted type locality (Schwartz, J. Herpetol., 1: 40): "vicinity of Port-au-Prince, Dépt. de l'Oueste, Haiti".  
1843 *Steironotus (Steironotus) Schreibersii*—Fitzinger, Syst. Rept.: 70.  
1868 *Liocephalus schreibersii*—Cope, Proc. Acad. Nat. Sci. Philadelphia, 20: 123.  
1941 *Leiocephalus schreibersii*—Cochran, Bull. U.S. Natl. Mus., 177: 204; Fig. 65a.  
1968 *Leiocephalus schreibersi*—Schwartz, J. Herpetol., 1: 40.  
1992 *Leiocephalus schreibersi*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 51.

Range: Haiti and western República Dominicana.

*Leiocephalus schreibersii schreibersii* (Gravenhorst)

1968 *Leiocephalus schreibersi schreibersi*—Schwartz, J. Herpetol., 1: 41.  
1992 *Leiocephalus schreibersi schreibersi*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 52.

Range: The Cul de Sac—Valle de Neiba plain, from Azua and Barahona (south to La Ciénega on the east coast of the Península de Barahona) in the east to Ca Ira and the type locality in the west, and northwest along the Golfe de la Gonâve to Dessalines, Gonaïves, and Ennery, Dépt. de l'Artibonite; isolated segment on the Haitian Presqu'île du Nord Ouest (Bombardopolis to Port-de-Paix); isolated segment in the Valle de Cibao (Monte Cristi to the vicinity of Guayubín); Cayos Siete Hermanos (Isla Tororú, Muertos, Ratas, and Tercero).

*Leiocephalus schreibersii nesomorus* Schwartz

1968 *Leiocephalus schreibersi nesomorus* Schwartz, J. Herpetol., 1: 47. Type locality: "Palmiste, Ile de la Tortue, Haiti" (Holotype: Mus. Comp. Zool. No. 81120).

1992 *Leiocephalus schreibersi nesomorus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 52.

Range: Ile de la Tortue off the north coast of Haiti.

*Leiocephalus semilineatus* Dunn

1920 *Leiocephalus semilineatus* Dunn, Proc. New England Zool. Club, 7: 33. Type locality: "Thomazeau, Haiti" (Holotype: Mus. Comp. Zool. No. 12748).

1932 *Leiocephalus personatus semilineatus*—Cochran, Proc. Biol. Soc. Washington, 45: 177.

1941 *Leiocephalus personatus semilineatus*—Cochran, Bull. U.S. Natl. Mus., 177: 217; Pl. 2A; Fig. 65c.

1968 *Leiocephalus semilineatus*—Schwartz, J. Herpetol., 1: 50.

1992 *Leiocephalus semilineatus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 52.

Range: Western Hispaniola: the Cul de Sac–Valle de Neiba plain, into the Llanos de Azua in the east; from Port-au-Prince and vicinity in Haiti (ascending the southern slopes of the Montagnes du Trou-d'Eau at Fond Michelle and the northern slopes of the Massif de la Selle at Soliette) in the west, east to 16 km northwest of Baní, Peravia Province, north into the eastern portion of the Valle de San Juan and northeast of Padre las Casas, Azua Province.

*Leiocephalus stictigaster* Schwartz

1959 *Leiocephalus stictigaster* Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 121; Fig. 6. Type locality: "beach on cabo Corrientes, Pinar del Río Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 77864).

1992 *Leiocephalus stictigaster*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 52.

1999 *Leiocephalus macropus asbolomus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 143.

Range: Cuba and the Isla de Juventud (= Pinos).

*Leiocephalus stictigaster stictigaster*—Schwartz

1959 *Leiocephalus stictigaster stictigaster* Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 123; Fig. 7.

1992 *Leiocephalus stictigaster stictigaster*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.

1999 *Leiocephalus stictigaster stictigaster*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 144.

Range: The Península de Guanahacabibes, Pinar del Río Province western Cuba.

*Leiocephalus stictigaster astictus* Schwartz

1959 *Leiocephalus stictigaster astictus* Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 134; Fig. 10. Type locality: "Caleta de Carapachibey, Isla de Pinos, Habana Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 81095).

1992 *Leiocephalus stictigaster astictus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.

1999 *Leiocephalus stictigaster astictus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 145.

Range: Isla de Juventud (= Pinos) south of the Ciénaga de Lanier.

*Leiocephalus stictigaster celeustes* Schwartz and Garrido

1968 *Leiocephalus stictigaster celeustes* Schwartz and Garrido, Nat. Hist. Pap. Natl. Mus. Canada, 37: 14; Pl. 1, Fig. J. Type locality: "Contramaestre, Oriente Province, Cuba" (Holotype: Inst. Zool. Acad. Cien. Habana No. 1182).

1992 *Leiocephalus stictigaster celeustes*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.

1999 *Leiocephalus stictigaster celeustes*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 146.

Range: Along the northern flank of the Sierra Maestra from the vicinity of Bueycitos in the west to the type locality in the east, Cuba.

*Leiocephalus stictigaster exotheotus* Schwartz

1959 *Leiocephalus stictigaster exotheotus* Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 130; Fig. 9. Type locality: "1.5 miles west of Santa Fé, Isla de Pinos, Habana Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 81088).

1992 *Leiocephalus stictigaster exotheotus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.

1999 *Leiocephalus stictigaster exotheotus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 145.

Range: Isla de Juventud (= Pinos) north of the Ciénaga de Lanier.

*Leiocephalus stictigaster gibarensis* Schwartz and Garrido

1968 *Leiocephalus stictigaster gibarensis* Schwartz and Garrido, Nat. Hist. Pap. Natl. Mus. Canada, 37: 18; Pl. 1, Fig. K. Type locality: "Gibara, Oriente Province, Cuba" (Holotype: Inst. Zool. Acad. Cien. No. 1236).

1992 *Leiocephalus stictigaster gibarensis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.

1999 *Leiocephalus stictigaster gibarensis*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 146.

Range: Known only from the type locality in eastern Cuba.

*Leiocephalus stictigaster lipomator* Schwartz and Garrido

- 1968 *Leiocephalus stictigaster lipomator* Schwartz and Garrido, Nat. Hist. Pap. Natl. Mus. Canada, 37: 11; Pl. 1, Fig. F. Type locality: "3 kilometers west of Santa Clara, Las Villas Province, Cuba" (Holotype: Inst. Zool. Acad. Cien. Habana No. 1230).
- 1992 *Leiocephalus stictigaster lipomator*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.
- 1999 *Leiocephalus stictigaster lipomator*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 145.
- Range: Known only from the type locality.

*Leiocephalus stictigaster lucianus* Schwartz

- 1960 *Leiocephalus stictigaster lucianus* Schwartz, Proc. Biol. Soc. Washington, 73: 104. Type locality: "Playa Santa Lucia, Camagüey Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 83583).
- 1992 *Leiocephalus stictigaster lucianus*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.
- 1999 *Leiocephalus stictigaster lucianus*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 145.
- Range: Known only from the vicinity of the type locality.

*Leiocephalus stictigaster naranjoi* Schwartz and Garrido

- 1968 *Leiocephalus stictigaster naranjoi* Schwartz and Garrido, Nat. Hist. Pap. Natl. Mus. Canada, 37: 3; Pl. 1, Fig. E. Type locality: "Los Biasmones, Casilda, Las Villas Province, Cuba" (Holotype: Inst. Zool. Acad. Cien. Habana No. 200).
- 1992 *Leiocephalus stictigaster naranjoi*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.
- 1999 *Leiocephalus stictigaster naranjoi*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 145.
- Range: Southern Sancti Spiritus and Camagüey provinces; vicinity of Juragua and Casilda, Cienfuegos Province, Cuba.

*Leiocephalus stictigaster ophiplacodes* Schwartz

- 1964 *Leiocephalus stictigaster ophiplacodes* Schwartz, Q. J. Florida Acad. Sci., 27: 217; Fig. 3. Type locality: "2.7 mi. SE Bonaó, Camagüey Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 92771).
- 1992 *Leiocephalus stictigaster ophiplacodes*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.
- 1999 *Leiocephalus stictigaster ophiplacodes*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 145.
- Range: The serpentine savannas of Camagüey Province south of the Sierra de Cubitas, Cuba.

*Leiocephalus stictigaster parasphex* Schwartz

- 1964 *Leiocephalus stictigaster parasphex* Schwartz, Q. J. Florida Acad. Sci., 27: 212; Fig. 1. Type locality: "Playa Bonita, east end of Cayo Sabinal, Camagüey Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 92153).
- 1992 *Leiocephalus stictigaster parasphex*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.
- 1999 *Leiocephalus stictigaster parasphex*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 145.
- Range: Known only from the type locality.

*Leiocephalus stictigaster septentrionalis* Garrido

- 1975 *Leiocephalus stictigaster septentrionalis* Garrido, Poeyana, 141: 28; Fig. 6. Type locality: "Cayo Santa María, Archipiélago de Sabana-Camagüey, Provincia de Las Villas, Cuba" (Holotype: Inst. Zool. Acad. Cien. Habana No. 3425).
- 1992 *Leiocephalus stictigaster septentrionalis*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.
- 1999 *Leiocephalus stictigaster septentrionalis*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 146.
- Range: Cayo Santa María, Cayo Francés, and Cayo Guillermo in the Archipiélago de Sabana-Camagüey off the north coast of central Cuba.

*Leiocephalus stictigaster sierrae* Schwartz

- 1959 *Leiocephalus stictigaster sierrae* Schwartz, Bull. Florida State Mus., Biol. Sci., 4: 126; Fig. 8. Type locality: "San Vicente, Pinar del Río Province, Cuba" (Holotype: Am. Mus. Nat. Hist. No. 77813).
- 1992 *Leiocephalus stictigaster sierrae*—Pregill, Misc. Publ. Mus. Nat. Hist. Univ. Kansas, 84: 53.
- 1999 *Leiocephalus stictigaster sierrae*—Rodríguez Schettino, in Rodríguez Schettino (ed.), Iguanid Lizards of Cuba: 145.
- Range: From near Cayuco west to the vicinity of Las Pozas, both in the lowlands and in the massifs of the Sierra de los Organos and the Sierra del Rosario, Pinar del Río Province, Cuba.

*Leiocephalus vinculum* Cochran

- 1928 *Leiocephalus vinculum* Cochran, Proc. Biol. Soc. Washington, 41: 54. Type locality: "Pte. à Roquette, Gonave Island" (Holotype: Mus. Comp. Zool. No. 25435).
- 1941 *Leiocephalus personatus vinculum*—Cochran, Bull. U.S. Natl. Mus., 177: 235; Pl. 6b; Fig. 66b.
- 1967 *Leiocephalus vinculum vinculum*—Schwartz, Tulane Stud. Zool., 14: 43; Fig. 10b.
- Range: Haiti and Isla Alto Velo, República Dominicana.

Literature Cited

- Barbour, T. 1914. A contribution to the zoogeography of the West Indies, with especial reference to amphibians and reptiles. *Memoirs of the Museum of Comparative Zoology* 44: 205–359.
- Barbour, T. 1916. Additional notes on West Indian reptiles and amphibians. *Proceedings of the Biological Society of Washington* 29: 215–220.
- Barbour, T. 1920. A *Leiocephalus* misnamed. *Copeia* 85: 73.
- Barbour, T. 1937. Third list of Antillean reptiles and amphibians. *Bulletin of the Museum of Comparative Zoology* 82: 77–166.
- Barbour, T., and B. Shreve. 1935. Concerning some Bahaman Reptiles, with notes on the Fauna. *Proceedings of the Boston Society of Natural History* 40: 347–366.
- Boulenger, G.A. 1885. Catalogue of the lizards in the British Museum (Natural History). Volume 2. Iguanidae, Xenosauridae, Zonuridae, Anguidae, Anellidae, Helodermatidae, Varanidae, Xantusiidae, Teiidae. London: Printed by Order of the Trustees.
- Cochran, D.M. 1928. The herpetological collections made in Haiti and its adjoining islands by Walter J. Eyerdam. *Proceedings of the Biological Society of Washington* 41: 53–59.
- Cochran, D.M. 1928. A new genus and species of Lizard, *Hispaniolus pratensis*, from the Haitian Republic. *Proceedings of the Biological Society of Washington* 41: 49–51.
- Cochran, D.M. 1931. New Bahaman Reptiles. *Journal of the Washington Academy of Sciences* 21: 39–41.
- Cochran, D.M. 1932. Two new subspecies of lizards of the genus *Leiocephalus* from Hispaniola. *Proceedings of the Biological Society of Washington* 45: 177–182.
- Cochran, D.M. 1934. Herpetological collections made in Hispaniola by the Utowana Expedition, 1934. *Occasional Papers of the Boston Society of Natural History* 8: 163–188.
- Cochran, D.M. 1934. A new lizard, *Leiocephalus personatus lunatus*, from the Dominican Republic. *Occasional Papers of the Boston Society of Natural History* 8: 153–156.
- Cochran, D.M. 1941. The herpetology of Hispaniola. *Bulletin of the United States National Museum* 177: 1–398.
- Cope, E.D. 1862. Contributions to Neotropical saurology. *Proceedings of the Academy of Natural Sciences of Philadelphia* 14: 176–188.
- Cope, E.D. 1868. An examination of the Reptilia and Batrachia obtained by the Orton Expedition to Ecuador and the Upper Amazon, with notes on other species. *Proceedings of the Academy of Natural Sciences of Philadelphia* 20: 96–140.
- Cope, E.D. 1887. List of the Batrachia and Reptilia of the Bahama Islands. *Proceedings of the United States National Museum* 10: 436–439.
- Duméril, A.M.C., and G. Bibron. 1837. *Erpétologie Générale ou Histoire Naturelle Complète des Reptiles*. Volume 4. Histoire de Quarante-six Genres et de Cent Quarante-six Espèces de la Famille des Iguaniens, de l'Ordre des Sauriens. Paris: Librairie Encyclopedique de Roret.

- Dunn, E.R. 1920. A new lizard from Haiti. *Proceedings of the New England Zoölogical Club* 7: 33–34.
- Etheridge, R.E. 1964. Late Pleistocene lizards from Barbuda, British West Indies. *Bulletin of the Florida Museum of Natural History. Biological Sciences* 9: 43–75.
- Etheridge, R.E. 1964. The skeletal morphology and systematic relationships of sceloporine lizards. *Copeia* 1964: 610–631.
- Etheridge, R.E. 1965. Fossil lizards from the Dominican Republic. *Quarterly Journal of the Florida Academy of Sciences* 29: 47–59.
- Etheridge, R.E. 1966. The systematic relationships of West Indian and South American lizards referred to the iguanid lizard genus *Leiocephalus*. *Copeia* 1966: 79–91.
- Fitzinger, L.J.F.J. 1826. *Neue Classification der Reptilien nach ihren Natürlichen Verwandtschaften nebst einer Verwandtschafts-Tafel und einem Verzeichnisse der Reptilien-Sammlung des K. K. Zoologisch Museum's zu Wien*. Wien: J. G. Heubner.
- Fitzinger, L.J.F.J. 1843. *Systema Reptilium. Fasciculus Primus*. Wien: Braumüller et Seidel.
- Frost, D.R., and R.E. Etheridge. 1989. A phylogenetic analysis and taxonomy of iguanian lizards (Reptilia: Squamata). *Miscellaneous Publications. Museum of Natural History, University of Kansas* 81: 1–65.
- Frost, D.R., R.E. Etheridge, D. Janies, and T.A. Titus. 2001. Total evidence, sequence alignment, evolution of polychrotid lizards, and a reclassification of the Iguania (Squamata: Iguania). *American Museum Novitates* 3343: 1–38.
- Gali, F., and A. Schwartz. 1982. A new subspecies of *Leiocephalus personatus* from the Republic Dominicana. *Journal of Herpetology* 16: 177–179.
- Gali, F., A. Schwartz, and A. Suarez. 1988. A new subspecies of *Leiocephalus personatus* (Sauria: Iguanidae) from Haiti. *Proceedings of the Biological Society of Washington* 101: 1–3.
- Garman, S. 1887. On the reptiles and batrachians of Grand Cayman. *Proceedings of the American Philosophical Society* 24: 273–286.
- Garrido, O.H. 1973. Nueva especie de *Leiocephalus* (Lacertilia, Iguanidae) para Cuba. *Poeyana. Habana* 116: 1–19.
- Garrido, O.H. 1973. Nueva subespecie de reptiles para Cuba. *Torreia. Nueva Serie. Habana* 30: 1–28.
- Garrido, O.H. 1975. Nuevos reptiles del Archipiélago Cubano. *Poeyana. Habana* 141: 1–58.
- Garrido, O.H. 1979. Nuevas subespecies de *Leiocephalus macropus* Cope (Lacertilia: Iguanidae) para Cuba. *Poeyana. Habana* 188: 1–16.
- Gravenhorst, J.L.C. 1837. Beiträge zur genauern Kenntniss einiger Eidechsegattungen. *Nova Acta Physico-medica Academiae Caesareae Leopoldino-Carolinae Naturae Curiosorum. Halle* 18: 712–784.
- Gray, J.E. 1827. A description of a new genus and some new species of saurian reptiles: with a revision of the species of chameleons. *The Philosophical*

- Magazine or Annals of Chemistry, Mathematics, Astronomy, Natural history, and General Science 2: 207–214.
- Gray, J.E. 1840. Catalogue of the species of reptiles collected in Cuba by W.S. MacLeay, Esq.;—with some notes on their habits extracted from his MS. Annals and Magazine of Natural History, Series 1, 5: 108–115.
- Gray, J.E. 1845. Catalogue of the Specimens of Lizards in the Collection of the British Museum. London: Trustees of the British Museum.
- Gundlach, J.C. 1880. Contribución a la Erpetología Cubana. Habana, Cuba: G. Montiel & Co.
- Hallowell, E. 1856. Notes on Reptilia in the collection of the Museum of the Academy of Natural Sciences. Proceedings of the Academy of Natural Sciences of Philadelphia 8: 146–153.
- Hardy, J.D., Jr. 1958. A new lizard of the genus *Leiocephalus* from Cuba (Squamata: Iguanidae). Journal of the Washington Academy of Sciences 48: 294–300.
- Mertens, R. 1939. Herpetologische Ergebnisse einer Reise nach der Insel Hispaniola, Westindien. Abhandlungen der Senckenbergischen Naturforschenden Gesellschaft. Frankfurt am Main 449: 1–84.
- Noble, G.K. 1923. Four new lizards from Beata Island, Dominican Republic. American Museum Novitates 64: 1–5.
- Noble, G.K., and W.G. Hassler. 1933. Two new species of frogs, five new species and a new race of lizards from the Dominican Republic. American Museum Novitates 652: 1–17.
- Pregill, G.K. 1981. Late Pleistocene herpetofaunas from Puerto Rico. Miscellaneous Publications. Museum of Natural History. University of Kansas 71: 1–72.
- Pregill, G.K. 1984. An extinct species of *Leiocephalus* from Haiti (Sauria: Iguanidae). Proceedings of the Biological Society of Washington 97: 827–833.
- Pregill, G.K. 1992. Systematics of the West Indian lizard genus *Leiocephalus* (Squamata: Iguania: Tropiduridae). Miscellaneous Publications. Museum of Natural History, University of Kansas 84: 1–69.
- Rabb, G.B. 1957. A new race of the iguanid lizard *Leiocephalus carinatus* from Cayman Brac, B.W.I. Herpetologica 13: 109–110.
- Reinhardt, J.T., and C.F. Lütken. 1863 "1862". Bidrag til det vestindiske Örges og navnlig til de dansk-vestindiske Örs herpetologie. Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening i Kjøbenhavn, Serie 2, 4: 153–291.
- Rodríguez Schettino, L. 1999. Systematic accounts of the species. In L. Rodríguez Schettino (ed.), The Iguanid Lizards of Cuba: 104–380. Gainesville: University Press of Florida.
- Schmidt, K.P. 1921. Notes on the herpetology of Santo Domingo. Bulletin of the American Museum of Natural History 44: 7–20.
- Schmidt, K.P. 1936. Notes on Bahaman reptiles and amphibians. Field Museum of Natural History Publication. Zoological Series 20: 127–133.

- Schwartz, A. 1959. The Cuban lizards of the species *Leiocephalus carinatus* (Gray). Reading Public Museum and Art Gallery. Scientific Publications 10: 1–47.
- Schwartz, A. 1959. Variation in lizards of the *Leiocephalus cubensis* complex in Cuba and the Isla de Pinos. Bulletin of the Florida Museum of Natural History. Biological Sciences 4: 97–143.
- Schwartz, A. 1960. A new subspecies of *Leiocephalus stictiogaster* Schwartz from Central Cuba. Proceedings of the Biological Society of Washington 73: 103–106.
- Schwartz, A. 1960. Variation in the Cuban lizard. *Leiocephalus raviceps* Cope. Proceedings of the Biological Society of Washington 73: 67–82.
- Schwartz, A. 1964. New subspecies of *Leiocephalus* from Cuba. Quarterly Journal of the Florida Academy of Sciences 27: 211–222.
- Schwartz, A. 1966 "1965". The *Leiocephalus* (Lacertilia, Iguanidae) of Hispaniola I. *Leiocephalus melanochlorus* Cope. Journal of the Ohio Herpetological Society 5: 39–48.
- Schwartz, A. 1967. The *Leiocephalus* (Lacertilia, Iguanidae) of Hispaniola. 2. The *Leiocephalus personatus* complex. Tulane Studies in Zoology 14: 1–53.
- Schwartz, A. 1967. The *Leiocephalus* (Lacertilia, Iguanidae) of the southern Bahama Islands. Annals of the Carnegie Museum 39: 153–185.
- Schwartz, A. 1968. The *Leiocephalus* (Lacertilia, Iguanidae) of Hispaniola 3. *Leiocephalus schreibersi*, *L. semilineatus*, and *L. pratensis*. Journal of Herpetology 1: 39–63.
- Schwartz, A. 1969. The status of the name *Holotropis microlophus* Cocteau. Copeia 1969: 620–621.
- Schwartz, A. 1969. Two new subspecies of *Leiocephalus* from Hispaniola. Journal of Herpetology 3: 79–85.
- Schwartz, A. 1971. A new subspecies of *Leiocephalus personatus* (Sauna, Iguanidae). Herpetologica 27: 176–182.
- Schwartz, A. 1979. The herpetofauna of Ile a Cabrit, Haiti, with the description of two new subspecies. Herpetologica 35: 248–255.
- Schwartz, A. 1979. A new species of *Leiocephalus* (Reptilia: Iguanidae) from Hispaniola. Proceedings of the Biological Society of Washington 92: 272–279.
- Schwartz, A., and O.H. Garrido. 1967. A review of the Cuban iguanid lizard *Leiocephalus macropus* Cope. Reading Public Museum and Art Gallery. Scientific Publications 14: 1–41.
- Schwartz, A., and O.H. Garrido. 1968. Four new subspecies of *Leiocephalus stictigaster* from Cuba. Natural History Papers. National Museum of Canada. 37: 1–23.
- Schwartz, A., and O.H. Garrido. 1968. An undescribed subspecies of *Leiocephalus raviceps* Cope (Sauria: Iguanidae) from western Cuba. Proceedings of the Biological Society of Washington 81: 23–30.

- Schwartz, A., and L.H. Ogren. 1956. A collection of reptiles and amphibians from Cuba, with the descriptions of two new forms. *Herpetologica* 12: 91–110.
- Schwartz, A., and R. Thomas. 1975. A check-list of West Indian amphibians and reptiles. *Special Publications of the Carnegie Museum of Natural History* 1: 1–216.
- Stejneger, L. 1901. Diagnosis of a new species of iguanoid lizard from Green Cay, Bahama Islands. *Proceedings of the United States National Museum* 23: 471.
- Stejneger, L. 1917. Cuban amphibians and reptiles collected for the United States National Museum from 1899 to 1902. *Proceedings of the United States National Museum* 53: 259–291.
- Thomas, R. 1966. *Journal of the Ohio Herpetological Society* 5: incl. 80.
- Varona, L.S., and O.H. Garrido. 1970. Vertebrados de los Cayos de San Felipe, Cuba, incluyendo una nueva especie de hutia. *Poeyana. Habana* 75: 1–26.
- Zug, G.R. 1959. Three new subspecies of the lizard *Leiocephalus macropus* Cope from Cuba. *Proceedings of the Biological Society of Washington* 72: 139–150.