

REGULARITY OF SOLUTIONS TO AN ABSTRACT INHOMOGENEOUS LINEAR DIFFERENTIAL EQUATION

G. F. WEBB

ABSTRACT. Let $T(t)$, $t > 0$, be a strongly continuous semigroup of linear operators on a Banach space X with infinitesimal generator A satisfying $T(t)X \subset D(A)$ for all $t > 0$. Let f be a function from $[0, \infty)$ to X of strong bounded variation. It is proved that $u(t) = \text{def} T(t)x + \int_0^t T(t-s)f(s)ds$, $x \in X$, is strongly differentiable and satisfies $du(t)/dt = Au(t) + f(t)$ for all but a countable number of $t > 0$.

1. Introduction. Let $T(t)$, $t \geq 0$, be a strongly continuous semigroup of bounded linear operators on the Banach space X with infinitesimal generator A and let f be an X -valued function on $[0, \infty)$. Our objective is to establish sufficient conditions so that the function

$$(1.1) \quad u(t) \stackrel{\text{def}}{=} T(t)x + \int_0^t T(t-s)f(s) ds, \quad x \in X,$$

is a strong solution of the inhomogeneous linear differential equation

$$(1.2) \quad du(t)/dt = Au(t) + f(t), \quad u(0) = x.$$

It is well known that $u(t)$ satisfies (1.2) for $t \geq 0$ provided that $x \in D(A)$ and f is continuously differentiable (see [4, Theorem 1.19, p. 486] or [5, Theorem 6.5, p. 135]). It is also well known that $u(t)$ satisfies (1.2) for $t > 0$ provided that $x \in X$, $T(t)$, $t \geq 0$, is homomorphic, and f is Hölder continuous (see [4, Theorem 1.27, p. 491] or [5, Theorem 6.7, p. 138]). The theorem which we will prove demonstrates that $u(t)$ satisfies (1.2) under the assumptions that $T(t)X \subset D(A)$ for $t > 0$ and f is of strong bounded variation. The main idea of our proof is to show that under our assumptions the integral in (1.1) lies in $D(A)$ and the image of this integral under A may be represented as a Stieltjes integral.

THEOREM. *Suppose $T(t)X \subset D(A)$ for all $t > 0$ and f is of strong bounded variation on $[0, r]$. For a given $x \in X$ let $u(t)$ be defined on $[0, r]$ by (1.1). Then, $u(t)$ satisfies the following:*

Presented to the Society, September 18, 1975; received by the editors August 14, 1975 and, in revised form, December 16, 1975.

AMS (MOS) subject classifications (1970). Primary 47D05; Secondary 34G05.

Key words and phrases. Strongly continuous semigroup, infinitesimal generator, inhomogeneous equation, strong bounded variation.

© American Mathematical Society 1977

$$(1.3) \quad u(t) \in D(A) \text{ for } t \in (0, r] \text{ and } Au(t) \text{ is continuous on } (0, r];$$

$$(1.4) \quad d^+u(t)/dt = Au(t) + f(t+) \text{ for all } t \in (0, r) \text{ and } d^+u(t)/dt \text{ is continuous from the right on } (0, r);$$

$$(1.5) \quad d^-u(t)/dt = Au(t) + f(t-) \text{ for all } t \in (0, r] \text{ and } d^-u(t)/dt \text{ is continuous from the left on } (0, r);$$

$$(1.6) \quad du(t)/dt = Au(t) + f(t) \text{ for all but a countable number of points in } [0, r] \text{ and } du(t)/dt \text{ is continuous at all but a countable number of points in } [0, r].$$

Before proving our theorem we first state some facts about Banach space-valued functions of strong bounded variation.

2. Vector-valued functions of strong bounded variation. Suppose f is of strong bounded variation from $[0, r]$ to X (according to the definition of [3, p. 59]). The following properties of f may be proved analogously to the case of real-valued functions of bounded variation (for a discussion of real-valued functions of bounded variation the reader is referred to [9, Chapter 2] or [2, Chapter II]):

$$(2.1) \quad f \text{ has a right limit at each } t \in [0, r), \text{ denoted by } f(t+), \text{ and } f(\cdot+) \text{ is right continuous on } [0, r);$$

$$(2.2) \quad f \text{ has a left limit at each } t \in (0, r], \text{ denoted by } f(t-), \text{ and } f(\cdot-) \text{ is left continuous on } (0, r];$$

$$(2.3) \quad f(\cdot-) \text{ is of strong bounded variation on } [0, r] \text{ (where for convenience we define } f(0-) = f(0)), \text{ and if we define } \nu(t) \text{ to be the total variation of } f(\cdot-) \text{ between } 0 \text{ and } t, \text{ then } \nu \text{ is nondecreasing and left continuous on } (0, r];$$

$$(2.4) \quad f \text{ is bounded on } [0, r] \text{ and continuous at all but a countable number of points in } [0, r].$$

3. Proof of the theorem. We first prove the lemmas below, each of which is under the hypothesis of the theorem. In what follows we will suppose that M

is a constant such that $|T(t)| \leq M$ for $0 \leq t \leq r$ (see [4, p. 484]) and ν is defined as in (2.3).

LEMMA 3.1. *If $0 < t \leq r$, then*

$$(3.1) \quad \int_0^t T(t-s)f(s) ds \in D(A);$$

$$A \int_0^t T(t-s)f(s) ds = \int_0^t dT(t-s)f(s-).$$

PROOF. Let $0 < t \leq r$. We observe that $\int_0^t T(t-s)f(s) ds$, the Riemann integral, exists since the integrand is bounded and continuous almost everywhere by virtue of (2.1) and the continuity properties of $T(t)$, $t \geq 0$. The function T from $[0, t]$ to $B(X, X)$ (where $B(X, X)$ denotes the Banach space of bounded linear operators on X) is bounded on $[0, t]$. Further, since $T(t)X \subset D(A)$ for $t > 0$, T is continuous from $(0, t]$ to $B(X, X)$ (see [3, Theorem 10.3.5, p. 310]). By (2.3) the set of discontinuities of $T(t-s)$, considered as a function of s in $[0, t]$ to $B(X, X)$, has ν measure 0. That is, $s \rightarrow T(t-s)$ is discontinuous only at t and, by (2.3), $\lim_{s \rightarrow t-} \nu(s) = \nu(t)$. Thus, the Riemann-Stieltjes integral $\int_0^t dT(t-s)f(s-)$ exists in the sense that for each $\epsilon > 0$ there exists $\delta > 0$ such that if $\{s_i\}_{i=0}^n$ is a chain from 0 to t such that $\sup_{i=1, \dots, n} |s_i - s_{i-1}| < \delta$, and $s_{i-1} \leq s'_i \leq s_i$, then

$$(3.2) \quad \left\| \sum_{i=1}^n (T(t-s_i) - T(t-s_{i-1}))f(s'_i-) - \int_0^t dT(t-s)f(s-) \right\| < \epsilon$$

(see [2, Theorem 13.16, p. 65 and Theorem 11.7, p. 53]).

For each positive integer n let $s_i^n = it/n$, where $i = 0, 1, \dots, n$. Define $g_n: [0, t] \rightarrow X$ by $g_n(s) = T(t-s)f(s_i^n-)$, where $s_{i-1}^n < s \leq s_i^n$, $i = 1, \dots, n$, and $g_n(0) = T(t)f(0)$. By (2.4), $\{g_n\}$ is bounded on $[0, t]$ and $\{g_n\}$ converges to $T(t-s)f(s)$ almost everywhere on $[0, t]$. By the Lebesgue theorem,

$$(3.3) \quad \lim_{n \rightarrow \infty} \int_0^t g_n(s) ds = \lim_{n \rightarrow \infty} \sum_{i=1}^n \int_{s_{i-1}^n}^{s_i^n} T(t-s)f(s_i^n-) ds = \int_0^t T(t-s)f(s) ds$$

(see [3, Theorem 3.7.9, p. 83]). From [4, p. 486], $\int_0^t g_n(s) ds \in D(A)$ and

$$(3.4) \quad A \int_0^t g_n(s) ds = \sum_{i=1}^n (T(t-s_i^n) - T(t-s_{i-1}^n))f(s_i^n-).$$

Then, by (3.2), (3.3), (3.4), and the closedness of A we obtain (3.1).

LEMMA 3.2. *$A \int_0^t T(t-s)f(s) ds$ is continuous from the right in t on $[0, r)$.*

PROOF. Let $0 \leq t < r$. First, we show that

$$(3.5) \quad \lim_{h \rightarrow 0^+} A \int_t^{t+h} T(t+h-s)f(s) ds = 0.$$

We observe that an argument similar to that of Lemma 3.1 shows that

$$\int_t^{t+h} T(t+h-s)f(s)ds \in D(A)$$

and

$$A \int_t^{t+h} T(t+h-s)f(s)ds = \int_t^{t+h} dT(t+h-s)f(s-).$$

Take $h > 0$ and sufficiently small. If $\varepsilon > 0$ there is a chain $\{s_i\}_{i=0}^n$ from t to $t+h$ such that

$$\begin{aligned} & \left\| \int_t^{t+h} (T(h) - T(t+h-s)) df(s-) \right\| \\ (3.6) \quad & < \left\| \sum_{i=1}^n (T(h) - T(t+h-s_{i-1})) (f(s_i-) - f(s_{i-1}-)) \right\| + \varepsilon \\ & < 2M \sum_{i=2}^n \|f(s_i-) - f(s_{i-1}-)\| + \varepsilon \\ & < 2M(\nu(t+h) - \nu(s_1)) + \varepsilon. \end{aligned}$$

Then, (3.6) yields

$$\begin{aligned} & \left\| \int_t^{t+h} (T(h) - T(t+h-s)) df(s-) \right\| \\ (3.7) \quad & < 2M(\nu(t+h) - \lim_{s \rightarrow t^+} \nu(s)). \end{aligned}$$

An integration by parts (see [2, Theorem 11.7, p. 53]) together with (3.7) yields

$$\begin{aligned} & \left\| A \int_t^{t+h} T(t+h-s)f(s)ds \right\| = \left\| \int_t^{t+h} dT(t+h-s)f(s-) \right\| \\ & = \left\| - \int_t^{t+h} T(t+h-s) df(s-) + f((t+h)-) - T(h)f(t-) \right\| \\ (3.8) \quad & = \left\| \int_t^{t+h} (T(h) - T(t+h-s)) df(s-) \right. \\ & \quad \left. + f((t+h)-) - T(h)f((t+h)-) \right\| \\ & < 2M(\nu(t+h) - \lim_{s \rightarrow t^+} \nu(s)) + \|(I - T(h))f((t+h)-)\|. \end{aligned}$$

In order to establish (3.5) we need only show that

$$(3.9) \quad \lim_{h \rightarrow 0^+} \|(I - T(h))f((t+h)-)\| = 0.$$

But (3.9) holds by virtue of the fact that the range of $f(\cdot-)$ on $[0, r]$ lies in a compact set of X and $\lim_{h \rightarrow 0^+} (I - T(h))z = 0$ uniformly for z in a compact set. The right continuity of $A \int_0^t T(t-s)f(s)ds$ in t now follows from (3.5) and the fact that

$$\begin{aligned}
 & A \int_0^{t+h} T(t+h-s)f(s)ds - A \int_0^t T(t-s)f(s)ds \\
 &= (T(h) - I)A \int_0^t T(t-s)f(s)ds + A \int_t^{t+h} T(t+h-s)f(s)ds.
 \end{aligned}$$

LEMMA 3.3. $A \int_0^t T(t-s)f(s)ds$ is continuous from the left in t on $(0, r]$.

PROOF. Let $0 < t \leq r$. Observe that for $c > 0$ and sufficiently small,

$$\begin{aligned}
 (3.10) \quad & \left\| A \int_{t-c}^t T(t-s)f(s)ds \right\| = \left\| \int_{t-c}^t dT(t-s)f(s-) \right\| \\
 &= \left\| - \int_{t-c}^t T(t-s)df(s-) + f(t-) - T(c)f((t-c)-) \right\| \\
 &\leq M(\nu(t) - \nu(t-c)) + M\|f(t-) - f((t-c)-)\| \\
 &\quad + \|(I - T(c))f(t-)\|.
 \end{aligned}$$

If $h > 0$ and $c > 0$ are both sufficiently small, then (3.10) applied twice below yields

$$\begin{aligned}
 (3.11) \quad & \left\| A \int_0^t T(t-s)f(s)ds - A \int_0^{t-h} T(t-h-s)f(s)ds \right\| \\
 &= \left\| AT(c) \left(\int_0^{t-h-c} (T(h) - I)T(t-h-c-s)f(s)ds \right. \right. \\
 &\quad \left. \left. + \int_{t-h-c}^{t-c} T(t-c-s)f(s)ds \right) \right. \\
 &\quad \left. + A \int_{t-c}^t T(t-s)f(s)ds - A \int_{t-h-c}^{t-h} T(t-h-s)f(s)ds \right\| \\
 &\leq |AT(c)| \left(\left\| \int_0^{t-h-c} (T(h) - I)T(t-h-c-s)f(s)ds \right\| \right. \\
 &\quad \left. + Mh \sup_{s \in [t-h-c, t-c]} \|f(s)\| \right) \\
 &\quad + M(\nu(t) - \nu(t-c)) + M\|f(t-) - f((t-c)-)\| \\
 &\quad + \|(I - T(c))f(t-)\| + M(\nu(t-h) - \nu(t-h-c)) \\
 &\quad + M\|f((t-h)-) - f((t-h-c)-)\| \\
 &\quad + \|(I - T(c))f((t-h)-)\|.
 \end{aligned}$$

For a given $\epsilon > 0$ first choose $c > 0$ and then choose $\delta > 0$ such that if $0 < h < \delta$, then (3.11) is $< \epsilon$ (use the fact that ν and $f(\cdot-)$ are left continuous at t and $\lim_{h \rightarrow 0^+} (T(h) - I)z = 0$ uniformly for z in a compact set). The left continuity of $A \int_0^t T(t-s)f(s)ds$ then follows immediately.

To complete the proof of the theorem we see that (1.3) follows from

Lemmas 3.1, 3.2, and 3.3. To prove (1.4) let $0 < t < r$ and observe that for $h > 0$ and sufficiently small we have

$$\begin{aligned} (u(t+h) - u(t))/h &= (T(t+h)x - T(t)x)/h \\ &+ \frac{1}{h} \int_t^{t+h} T(t+h-s)f(s)ds \\ &+ \frac{T(h) - I}{h} \int_0^t T(t-s)f(s)ds. \end{aligned}$$

By (2.1)

$$\lim_{h \rightarrow 0^+} \frac{1}{h} \int_t^{t+h} T(t+h-s)f(s)ds = f(t+)$$

and (1.4) then follows from Lemmas 3.1 and 3.2. To prove (1.5) let $0 < t < r$ and observe that for $h > 0$ and sufficiently small we have

$$\begin{aligned} (u(t-h) - u(t))/(-h) &= (T(t-h)x - T(t)x)/(-h) \\ &+ \frac{1}{h} \int_{t-h}^t T(t-s)f(s)ds \\ &+ \frac{T(h) - I}{h} \int_0^{t-h} T(t-h-s)f(s)ds. \end{aligned}$$

By (2.2)

$$\lim_{h \rightarrow 0^+} \frac{1}{h} \int_{t-h}^t T(t-h-s)f(s)ds = f(t-).$$

Denote

$$z(h) \stackrel{\text{def}}{=} \int_0^{t-h} T(t-h-s)f(s)ds.$$

By Lemma 3.1, $z(h) \in D(A)$ and by Lemma 3.3,

$$\lim_{h \rightarrow 0^+} \frac{T(h) - I}{h} z(h) = \lim_{h \rightarrow 0^+} \frac{1}{h} \int_0^h T(s)Az(h)ds = Az(0),$$

which yields (1.5). Finally, (1.6) follows immediately from (1.4), (1.5), and (2.4).

We conclude with the observation that our theorem may be applied to nonlinear evolution equations of the form $du(t)/dt = Au(t) + B(u(t))$. If $-B$ is an accretive continuous everywhere defined nonlinear operator on X , then there exists a solution $u(t)$ to the Volterra integral equation

$$u(t) = T(t)x + \int_0^t T(t-s)B(u(s))ds, \quad x \in X$$

(see [10, Theorem I]). If we assume that $x \in D(A)$, then it can be shown that $u(t)$ is Lipschitz continuous. If we also assume that B is Lipschitz continuous and $T(t)X \subset D(A)$ for all $t > 0$, then our theorem implies $u(t)$ satisfies $du(t)/dt = Au(t) + B(u(t))$ for all $t \geq 0$. If it is not true that $T(t)X \subset D(A)$, then this conclusion may not hold (see [10, Example 4.1]). A similar

observation is made in [7] for the case that $T(t)$, $t \geq 0$, is a holomorphic semigroup.

REFERENCES

1. P. Benilan, *Opérateurs m -accrétifs hémicontinus dans un espace de Banach quelconque*, C. R. Acad. Sci. Paris Sér. A **278** (1974), 1029–1032. MR **50** #1064.
2. T. H. Hildebrandt, *Introduction to the theory of integration*, Academic Press, New York, 1963. MR **27** #4900.
3. E. Hille and R. S. Phillips, *Functional analysis and semi-groups*, rev. ed., Amer. Math. Soc. Colloq. Publ., vol. 31, Amer. Math. Soc., Providence, R. I., 1957. MR **19**, 664.
4. T. Kato, *Perturbation theory for linear operators*, Springer-Verlag, New York, 1966. MR **34** #3324.
5. S. G. Kreĭn, *Linear differential equations in Banach space*, "Nauka", Moscow, 1967; English transl., Transl. Math. Monographs, vol. 29, Amer. Math. Soc., Providence, R. I., 1972. MR **40** #508.
6. A. Pazy, *Approximation of the identity operator by semigroups of linear operators*, Proc. Amer. Math. Soc. **30** (1971), 147–150. MR **44** #4568.
7. ———, *A class of semi-linear equations of evolution* (to appear).
8. R. S. Phillips, *Perturbation theory for semi-groups of linear operators*, Trans. Amer. Math. Soc. **74** (1953), 199–221. MR **14**, 882.
9. B. Sz.-Nagy, *Introduction to real functions and orthogonal expansions*, Oxford Univ. Press, New York, 1965. MR **31** #5938.
10. G. F. Webb, *Continuous nonlinear perturbations of linear accretive operators in Banach spaces*, J. Functional Analysis **10** (1972), 191–203. MR **50** #14407.

DEPARTMENT OF MATHEMATICS, VANDERBILT UNIVERSITY, NASHVILLE, TENNESSEE 37235