

Neocalyptrella robusta (G.Norman ex Ralfs) Hernández-Becerril & Meave del Castillo

Phycologia 36: 329 (1997)

Rhizosolenia robusta G.Norman ex Ralfs, in Pritchard, *History of the Infusoria, including the Desmidiaceae and Diatomaceae, British and Foreign* 866, pl. 8, fig. 42 (1861)

Type locality: Great Britain

Australian distribution: N & NW Australia, Qld (Moreton Bay), N.S.W. (North Coast, Central Coast, South Coast), Vic., Tas. (Bass. Str.), W.A. (Perth)

References: L.H.Crosby & E.J.F.Wood, Studies on Australian and New Zealand diatoms. I. Planktonic and allied species, *Trans. & Proc. Roy. Soc. New Zealand* 85: 483–530 (1958)

G.M.Hallegraeff & S.W.Jeffrey, Tropical phytoplankton species and pigments of continental shelf waters of north and north-west Australia, *Mar. Ecol. Prog. Ser.* 20: 59–74 (1984)

Extra-Australian distribution: Argentina, S Brazil, Gulf of Mexico, U.S.A. (Virginia), North Atlantic Ocean, British Isles, North Sea, Baltic Sea, Canary Islands, Mediterranean Sea (Spain), Adriatic Sea, Black Sea, Arabian Sea, East China Sea, Yellow Sea, Sea of Japan, South China Sea, New Zealand, W and SE Pacific Ocean, Papua New Guinea, Peru, Southern Ocean