

ACANTHOPONERA

aurea Forel, 1913; see under **HETEROPONERA**.

crassa. *Acanthoponera crassa* Brown, 1958g: 255, fig. 10 (w.) ECUADOR, COSTA RICA.

Type-material: holotype worker, 3 paratype workers.

Type-locality: holotype Ecuador: Pichincha, Santo Domingo de los Colorados, 23.ii.1955 (*E.S. Ross & E.I. Schlinger*); paratypes with same data .

Type-depositories: CASC (holotype); CASC, MCZC (paratypes).

Junior synonym of *minor*: Kempf & Brown, 1968: 90; Kempf, 1972a: 9; Bolton, 1995b: 53.

goeldii. *Acanthoponera goeldii* Forel, 1912c: 34 (w.) BRAZIL (Espírito Santo).

Type-material: holotype worker.

Type-locality: Brazil: Espírito Santo (*Goeldi*) .

Type-depository: MHNG.

Status as species: Borgmeier, 1923: 55; Wheeler, W.M. 1923e: 188; Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 53; Feitosa, 2015c: 98.

Distribution: Brazil.

goyana. *Acanthoponera goeldii* subsp. *goyana* Borgmeier, 1939: 418 (w.) BRAZIL (Goiás).

Type-material: holotype worker.

Type-locality: Brazil: Goyaz (= Goiás), Campinas, ix.1927 (*P. Schwarzmaier*).

Type-depository: MZSP.

Junior synonym of *mucronata*: Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 53.

imbellis Emery, 1895; see under **HETEROPONERA**.

kirki Wheeler, W.M. 1923; see under **HETEROPONERA**.

leae Wheeler, W.M. 1923; see under **HETEROPONERA**.

minor. *Ectatomma (Acanthoponera) mucronatum* var. *minor* Forel, 1899c: 9 (w.) MEXICO (Tabasco).

Type-material: holotype worker.

Type-locality: Mexico: Tabasco, Teapa (*H.H. Smith*) .

Type-depository: MHNG.

[*Acanthoponera minor* Emery, 1896g: 33. *Nomen nudum* (attributed to Forel).]

Combination in *Acanthoponera*: Emery, 1911d: 36.

Subspecies of mucronata: Emery, 1911d: 36; Wheeler, W.M. 1923e: 189.

Status as species: Brown, 1958g: 194; Kempf & Brown, 1968: 89; Kempf, 1972a: 9; Fernández, 1993: 251; Bolton, 1995b: 54; Arias-Penna & Fernández, 2008: 112; Branstetter & Sáenz, 2012: 256; Feitosa & Prada-Achiardi, 2019: 675.

Senior synonym of crassa: Kempf & Brown, 1968: 90; Kempf, 1972a: 9; Bolton, 1995b: 54.

Senior synonym of schwarzi: Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 54.

Senior synonym of spininodis: Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 54.

Distribution: Colombia, Costa Rica, Ecuador, Guatemala, Honduras, Mexico, Trinidad.

mucronata. *Ponera mucronata* Roger, 1860: 299 (q.) BRAZIL (no state data).

Type-material: holotype worker.

Type-locality: Brazil: (no further data) .

Type-depository: MNHU.

Mayr, 1862: 733 (w.); Ketterl & Verhaagh, 2004: 66 (m.).

Combination in *Ectatomma*: Roger, 1863b: 17;

combination in *Ectatomma (Acanthoponera)*: Mayr, 1862: 733; Mayr, 1887: 540;

combination in *Acanthoponera*: Emery, 1911d: 36.

Status as species: Mayr, 1862: 733; Mayr, 1863: 410; Roger, 1863b: 17; Dalla Torre, 1893: 25; Emery, 1894c: 143; Forel, 1895b: 112; Forel, 1899c: 9; Forel, 1905b: 156; Emery, 1911d: 36; Forel, 1912c: 34; Luederwaldt, 1918: 34; Borgmeier, 1923: 55; Wheeler, W.M. 1923e: 189; Borgmeier, 1939: 418; Kusnezov, 1953b: 336; Brown, 1958g: 194; Kempf, 1972a: 9; Fernández, 1993: 251; Bolton, 1995b: 54; Ketterl & Verhaagh, 2004: 66; Wild, 2007b: 29; Arias-Penna & Fernández, 2008: 112; Feitosa, 2015c: 98; Guénard & Economo, 2015: 227; Feitosa & Prada-Achiardi, 2019: 675.

Senior synonym of *goyana*: Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 54.

Senior synonym of *plaumanni*: Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 54.

Senior synonym of *wagneri*: Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 54.

Distribution: Argentina, Bolivia, Brazil, Colombia, Ecuador, Paraguay, Peru, Venezuela.

nigra Clark, 1930; see under **HETEROPONERA**.

occidentalis Clark, 1926; see under **HETEROPONERA**.

peruviana. *Acanthoponera peruviana* Brown, 1958g: 254, figs. 6,7 (w.) PERU.

Type-material: holotype worker.

Type-locality: Peru: Tingo Maria, Monson Valley, 26.x.1954 (*E.S. Ross & E.I. Schlinger*).

Type-depository: CASC.

Status as species: Kempf, 1972a: 9; Bolton, 1995b: 54; Bezděčková, *et al.* 2015: 114; Feitosa, 2015c: 98.

Distribution: Brazil, Peru.

plaumanni. *Acanthoponera plaumanni* Donisthorpe, 1938a: 140 (w.) BRAZIL (Santa Catarina).

Type-material: holotype worker.

Type-locality: Brazil: Nova Teutonia, 27°11'8''S, 52°23'W, ix.1935 (*F. Plaumann*).

Type-depository: BMNH.

Junior synonym of *mucronata*: Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 54.

reichenspergeri Santschi, 1929; see under **GNAMPTOGENYS**.

scabra Wheeler, W.M. 1923; see under **HETEROPONERA**.

schwarzi. *Acanthoponera goeldii* subsp. *schwarzi* Wheeler, W.M. 1923e: 188, fig. 5 (w.) GUATEMALA.

Type-material: holotype worker.

Type-locality: Guatemala: Alta Vera Paz, Trece Aguas (*E.A. Schwarz & H.S. Barber*).

Type-depository: MCZC.

Junior synonym of *minor*: Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 54.

schwebeli Luederwaldt, 1918; see under **HETEROPONERA**.

spininodis. *Acanthoponera (Acanthoponera) spininodis* Weber, 1939a: 97, fig. 2 (w.) TRINIDAD.

Type-material: 2 syntype workers.

Type-locality: Trinidad: El Tucuché summit (3072 ft), 15.xii.1934 (*N.A. Weber*).

Type-depository: MCZC.

Junior synonym of *minor*: Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 54.

wagneri. *Acanthoponera mucronata* var. *wagneri* Santschi, 1921g: 84 (w.) ARGENTINA (Santiago del Estero).

Type-material: 2 syntype workers.

Type-locality: Argentina: Santiago del Estero, Banderas, 55 km. N Icaño, 1910, chaco (*E.R. Wagner*).

Type-depository: MNHN, NHMB.

Subspecies of *mucronata*: Wheeler, W.M. 1923e: 190.

Junior synonym of *mucronata*: Brown, 1958g: 194; Kempf, 1972a: 9; Bolton, 1995b: 54.