

**Biological Assessment  
Imperatrice Property  
City of Ashland, Oregon**

Prepared for:



Prepared by:

Pacific Crest Consulting, LLC

August 2017



## Table of Contents

1.0 INTRODUCTION.....	1
2.0 METHODS.....	2
2.1 Target Species .....	2
2.1.1 Special Status Species .....	2
2.1.2 Noxious Weeds.....	3
2.2 Protocol .....	3
2.3 Study Area.....	3
2.4 Habitat Assessment and Delineation .....	4
2.4.1 Threatened and Endangered Plants.....	4
2.4.2 Sensitive and Strategic Species .....	5
2.4.3 Survey and Manage Species .....	5
2.5 Historical Data Review.....	5
2.6 Survey Schedule .....	5
2.7 Field Survey Methods.....	6
2.7.1 Intuitive Controlled Survey Method.....	6
2.7.2 Complete Survey Method.....	6
2.7.3 Hypogeous Fungi.....	6
2.7.4 Point Counts .....	7
2.7.5 Monumenting Target Species Sites and Recording Site Data .....	7
3.0 RESULTS .....	8
3.1 Current Environment .....	8
3.1.1 Oak Woodlands .....	8
3.1.2 Meadows Between the Oak Woodlands and TID Canal .....	8
3.1.3 Meadows Downslope of the TID Canal .....	9
3.1.4 Other Features of the Study Area.....	10
3.2 Special Status Plants, Lichens, and Fungi.....	11
3.3 Noxious Weeds .....	13
3.4 Birds .....	14
3.5 Other Sites of Interest.....	14
3.6 Inventories.....	15
4.0 REFERENCES.....	16

## List of Tables

Table 1: Special status plants, lichens, and fungi .....	19
Table 2: Special status birds, mammals, reptiles, and amphibians .....	25
Table 3: ODA list of noxious weeds.....	33
Table 4: Avian inventory.....	37
Table 5: Vascular plant inventory .....	40

## List of Figures

Figure 1: Study Area .....	49
Figure 2: ORBIC Rare Plant Form.....	50
Figure 3: California macrophylla site locations .....	51
Figure 4: California macrophylla specimens .....	52
Figure 5: Ranunculus austro-oreganus and Collema quadrifidum site locations .....	53
Figure 6: Ranunculus austro-oreganus and Collema quadrifidum specimens .....	54
Figure 7: Noxious weed locations other than Centaurea solstitialis and Elymus caput-medusae .....	55
Figure 8: Grasshopper sparrow detections.....	56
Figure 9: Species of Phaeocalicium from the Study Area (micrograph).....	57
Figure 10: Petrified log .....	58

## Appendices

Appendix A: Representative Photos of the Property

### Acronyms and Abbreviations

BLM	Bureau of Land Management, US Department of the Interior
CS	Oregon State Conservation Strategy Species
City	City of Ashland, Oregon
GIS	Geographic Information System
GPS	Global Positioning System
GRSPs	Grasshopper Sparrows
ODA	Oregon Department of Agriculture
ODFW	Oregon Department of Fish and Wildlife
ORBIC	Oregon Biodiversity Information Center
Pacific Crest	Pacific Crest Consulting, LLC
POE	Port of Entry, Oregon Department of Transportation
Property	Imperatrice Property, City of Ashland, Oregon
TID	Talent Irrigation District
USFS	Forest Service, US Department of Agriculture
USFWS	US Fish and Wildlife Service

## 1.0 INTRODUCTION

This report presents the methods and results for the biological assessment of the City of Ashland's (City) Imperatrice Property (Property) conducted by Pacific Crest Consulting, LLC (Pacific Crest) during spring and summer of 2017. Efforts requested by the City and undertaken by Pacific Crest included:

- Protocol-level surveys for target species of:
  - Plants (vascular and non-vascular (bryophytes))
  - Lichens
  - Spring fungi (including mushrooms and truffles)
  
- Protocol-level surveys for grasshopper sparrows (*Ammodramus savannarum*; GRSPs)
  
- Informal surveys for target species of:
  - Invertebrates
  - Mammals
  - Reptiles
  - Amphibians
  - Birds other than GRSPs (see further notes in 2.1.1 Special Status Species)
  
- Inventory of all vascular plant species
  
- Inventory of unique biological features, including:
  - Wildlife passage areas and barriers
  - Dense concentrations and large infestations of noxious weeds
  - Potential migratory bird nest sites
  - Bat hibernacula
  - Other habitats or features viewed as unique

## 2.0 METHODS

### 2.1 Target Species

Multiple sources informed the target species lists for each survey included in this report. The categories of target species are described in the following sections. Because many of the plant, lichen, and fungi species addressed in this report do not have common names, all are referred to by scientific names in the text, with common names listed as applicable; a common name is generally given only once for any given species of these taxa groups, at its first occurrence in the text. Primary target species are those for which protocol surveys were conducted; secondary species are those for which informal surveys were conducted.

#### 2.1.1 *Special Status Species*

Special status species of plants, lichens, and fungi were primary targets for the surveys included in this report (Table 1). This included:

- State and federally listed Threatened, Endangered, and Candidate plants
- U.S. Forest Service (USFS) and U.S. Bureau of Land Management (BLM) Sensitive and Strategic plants, lichens, and fungi
- Oregon Biodiversity Information Center (ORBIC) plants, lichens, and fungi.
- Survey and Manage plants, lichens, and fungi

GRSPs were also a primary target. In addition to the species identified in Table 1, Pacific Crest personnel were prepared to identify and document any unexpected, unknown, or out-of-expected-range species that may have been of conservation concern.

Secondary target special status species included mammals, reptiles, amphibians, invertebrates, and birds other than GRSPs.

Although secondary targets, Pacific Crest elected to create a list of special status mammals, birds, reptiles, and amphibians (Table 2) with the potential to occur in, or near to, the Study Area, developed from the following sources:

- United States Fish and Wildlife Service (USFWS) informal list of threatened, endangered, proposed, candidate, species of concern, and migratory birds, generated using the Information, Planning, and Conservation System (IPaC; USFWS 2017).
- Oregon Department of Fish and Wildlife (ODFW) lists of threatened, endangered, candidate, and sensitive animal species in the State of Oregon (ODFW 2017 a and b);
- Oregon Biodiversity Information Center (ORBIC 2016)
- Bald and Golden Eagle Protection Act (16 U.S.C. 668-668c)

Bird and mammal taxa listed by the ODFW, by the USFWS as species of concern, and/or by ORBIC on Lists 1, 2, 3, or 4 are included, but have no legal status or protection on private land in the State of Oregon. On non-federal public lands (e.g., state, county, city lands), animal

species listed by ODFW as threatened or endangered are protected by the Oregon Endangered Species Act (Oregon Revised Statute 497). Federally listed threatened, endangered, candidate, and proposed taxa are protected under the Federal Endangered Species Act, bald and golden eagles are protected under the federal Bald and Golden Eagle Protection Act, and migratory birds are protected under the federal Migratory Bird Treaty Act.

Lists of target species other than the above taxa groups are available at <http://inr.oregonstate.edu/orbic/rare-species/rare-species-oregon-publications>.

### ***2.1.2 Noxious Weeds***

Species of noxious weeds were primary targets. Pacific Crest documented dense concentrations and large infestations of Oregon Department of Agriculture (ODA) target species (ODA 2017a; Table 3) or those that were uncommon or previously unknown in the area, or had a high potential of infestation.

## **2.2 Protocol**

Pacific Crest implemented the following protocols for primary target species during the survey effort:

- *2008 Sporocarp Survey Protocol for Macrofungi, Version 1.0* (Van Norman et al. 2008)
- *Survey Protocols for Protection Buffer Bryophytes* (USFS and BLM 1999a)
- *Survey and Manage Survey Protocols—Vascular Plants* (USFS and BLM 1999b)
- *Survey Protocols for Survey and Manage Category A and C Lichens in the Northwest Forest Plan Area, Version 2.1* (USFS and BLM 2003a)
- *2003 Amendment to the Survey Protocol for Survey & Manage Category A & C Lichens in the Northwest Forest Plan Area, Version 2.1 Amendment* (USFS and BLM 2003b)
- *Survey Protocol Guidance for Conducting Equivalent Effort Surveys under the Northwest Forest Plan Survey and Manage Standards and Guidelines* (USFS and BLM 2006)
- *Survey and Manage Category B Fungi Equivalent-Effort Survey Protocol, Version 1.0* (Van Norman 2010)

Secondary species were searched for informally and concurrently with protocol surveys for other taxa groups. Except for certain bird species, secondary species were not surveyed for under applicable protocols. See 2.7.4 Point Counts for more information on bird survey methods.

## **2.3 Study Area**

The Study Area encompassed the entire Property (Figure 1), consisting of 876 continuous acres across multiple tax lots, immediately north of Interstate Highway 5 and associated north-bound Port of Entry (POE). The Study Area includes portions of sections T38S R1E 27, 28, 32, and 33.



## 2.4 Habitat Assessment and Delineation

The Study Area was assessed and delineated for primary target species of vascular plants, lichens, and fungi. The Study Area exists within the Klamath Mountains level 3 ecoregion, only two miles from the western edge of the West Cascades level 3 ecoregion. It includes portions of the Western Oregon Interior Valleys (Rogue / Illinois) and Oak Savannah Foothills level 4 ecoregions. Therefore, it was considered possible that populations of target species known or suspected from interior valley and oak-associated habitats of both aforementioned level 3 ecoregions could be found in the Study Area. Initial topographical map and orthoquad inspection of the entire Study Area, as required by various protocols, revealed a wide variety of suitable primary target special status species habitats, including:

- Oak woodlands
- Rock outcrops and rock gardens
- Meadows
- Drainages (ephemeral / seasonal)

Field surveys confirmed this diversity of habitat types.

The Study Area was also assessed for secondary target species of animals except invertebrates, the results of which are discussed in Table 2.

### 2.4.1 Threatened and Endangered Plants

Pacific Crest conducted a pre-survey botanical habitat suitability analysis on the Study Area and found that *Fritillaria gentneri* (Gentner's fritillary) was the only species with federal or state listing of Threatened, Endangered, or Candidate with a high likelihood of occurring. *Fritillaria gentneri* has a federal listing of Endangered. The Property is well within the known range of this species and populations have been found nearby. Pacific Crest identified the area containing habitat with highest potential for suitability to be the oak woodlands at the north end of the Property.

Multiple other target special status species with federal or state listing, *Limnanthes floccosa* ssp. *grandiflora* (large-flowered wooly meadowfoam; federally Endangered), *L. floccosa* ssp. *pumila* (dwarf wooly meadowfoam; state Threatened), *Lomatium cookii* (Agate Desert lomatium; federally Endangered), *Meconella oregana* (white fairypoppy), and *Eucephalus vialis* (wayside aster) had limited potential to be found in the survey area. The former three taxa are known from vernal pool habitats nearby to the north in the Rogue Valley: *Limnanthes floccosa* ssp. *grandiflora* and *Lomatium cookii* are known from the Agate Desert while *L. floccosa* ssp. *pumila* is known only from the tops of Upper Table Rock and Lower Table Rock. The only potential habitat for these taxa in the Study Area was initially identified as a small seasonal pond near the eastern property line and a small flat area at the extreme northwest near Butler Creek. The valley bottom near Interstate 5 may have had vernal pool habitat historically, but has been heavily grazed and impacted by livestock and the pre-survey analysis revealed no current habitat. There was low probability to find these taxa in other seasonally moist habitats such as seasonal drainages. *Meconella oregana* had potential to be found in various meadow and oak

woodland communities. *Eucephalus vialis* had potential to be found in the shrub or oak communities within the Study Area. Multiple other species with federal and/or state status were included in Table 1 but had relatively lower potential to be found in the Study Area.

### 2.4.2 Sensitive and Strategic Species

Many of the target Sensitive and Strategic plant and lichen species were found to have a moderate to high likelihood of occurring. Two species, *California macrophylla* (*Erodium macrophyllum*; round-leaved filaree;) and *Ranunculus austro-oreganus* (southern Oregon buttercup) were known to exist in the Study Area (personal communication, Kristi Mergenthaler, Southern Oregon Land Conservancy, 4/25/17). Other species with a moderate to high likelihood of occurring included, but were not limited to, *Calochortus spp* (mariposa lilies), *Camissonia (Tetrapteron) graciliflora* (hill suncup), *Carex spp* (sedges), *Cheilanthes spp* (lipferns), *Cryptantha milobakeri*, *Diplacus spp* (monkeyflowers), *Leptogium burnetiae*, *Limnathes floccosa ssp bellingeriana*, *Pellaea andromedifolia* (cliffbrake), *Plagiobothrys spp* (popcorn flowers), *Orthotrichum euryphyllum*, *Schistidium cinclidodonteum*, and *Solanum parishii* (Parish's nightshade).

### 2.4.3 Survey and Manage Species

The Survey and Manage target species list includes plants, lichens, and fungi. These species were listed with Survey and Manage primarily based on rarity within and dependence on old-growth coniferous or mixed forests. Coniferous and mixed forest habitats do not exist in the Study Area. Therefore, there was very little potential for most Survey and Manage plant, lichen, and fungi species to exist in the Study Area.

## 2.5 Historical Data Review

The BLM Geographic Biotic Observations and USFS Natural Resource Information System databases track observations of noteworthy species on and near BLM and USFS lands. The Oregon Flora Project rare plant and atlas database (OFP 2017) was also consulted. These databases were queried for known site locations of target species within the vicinity of the Study Area (April 26, 2017). The results showed no site locations within the Study Area. Known target species occurrences within a three-mile radius of the Study Area include two of *Fritillaria gentneri*, one of *Martes pennanti* (fisher), one of *Horkelia tridentata* (three-toothed horkelia), and multiple of *Ranunculus austro-oreganus*.

Further results of the data review and literature reviews for plants, lichens, fungi, and animals (except invertebrates) are detailed in Table 1 and Table 2.

## 2.6 Survey Schedule

Two separate survey efforts occurred to coincide with optimum detection of target species of vascular plants and grasshopper sparrows, respectively. Optimal fungi fruiting conditions coincided with optimal vascular plant spring phenology. Lichens and non-vascular plants can

generally be surveyed for any time of year. Therefore, plant, lichen, and fungi surveys occurred concurrently and took place from April 28, 2017 through May 23, 2017, including a first visit throughout the entire study area and revisits to selected parts. Revisits for late-season plant species then occurred occasionally until August 5, 2017.

Surveys for GRSPs were conducted on May 2-4, May 6-9, May 15, and May 18, 2017.

## **2.7 Field Survey Methods**

Survey methods from multiple protocols, listed earlier in this report, were used during the Project surveys. The methods in the protocols are detailed below.

### ***2.7.1 Intuitive Controlled Survey Method***

Multiple protocols recommend the Intuitive Controlled Survey method for plants, lichens, and fungi in all parcels greater than 2.5 acres in size (USFS and BLM 1999b, USFS and BLM 2003a, USFS and BLM 2003b, USFS and BLM 2006, Van Norman 2010, Van Norman et al. 2008). One protocol, *Survey Protocols for Protection Buffer Bryophytes* (USFS and BLM 1999a), requires this method for all parcels, including those less than 2.5 acres in size. The Study Area is greater than 2.5 acres in size; the Intuitive Controlled Survey method was therefore implemented.

This method incorporates lines that traverse the survey area and target the full array of major vegetation types, aspects, topographical features, habitats, and substrate types within a given area. While en-route, the surveyor searches for target species, and when the surveyor arrives at an area of high potential habitat (as defined in the pre-field review or encountered during the field visit), a Complete Survey for the target species is conducted (see below).

### ***2.7.2 Complete Survey Method***

The Complete Survey method for plants, lichens, and fungi was used when special or high potential habitats were encountered. This approach consists of a 100 percent visual examination of the habitat. High potential habitats within the Study Area included large outcroppings, seasonal and perennial drainages, areas with significant native bunchgrass populations, and some areas with *Quercus garryana* (Oregon white oak). A large percentage of the Study Area had moderate potential habitat for target plant species and therefore received a higher intensive survey than that required by the Intuitive Controlled Survey method but not a full 100% examination. The general vascular plant inventory was completed concurrently with surveys for target special status species under the Intuitive and Complete survey protocols.

### ***2.7.3 Hypogeous Fungi***

All implemented fungi survey protocols require searches for hypogeous fungi—truffles. These surveys were conducted by raking microhabitats of higher potential (e.g., small mammal digs and the underside of litter mats in the oak woodlands). Surveyors used four-tine rakes to gently

peel back the litter layer, and soil was inspected for the presence of truffles. If no truffles were found, the area was restored and the surveyor moved to a new area.

### ***2.7.4 Point Counts***

Point count surveys for GRSPs occurred throughout the entire Study Area and were conducted between the hours of 0630 and 1200 during optimal conditions for detecting this species. For the purposes of these surveys, the property was divided into two portions: the area above the Talent Irrigation District (TID) East Canal and the area below the canal. Point count surveys were conducted along north-south running transect lines above the canal and along east-west running transect lines below the canal. Initially, transect lines were spaced 100 meters apart with call stations approximately every 50 meters; however, the call station placement was changed during the course of the survey in the following way: in areas where traffic noise made auditory detection difficult, call stations were maintained at 50-meter intervals; however, in areas where habitat was open and birds were easy to hear, call stations were spaced at 100 meters for efficiency and to avoid duplicate recordings of individuals. Five minutes was spent at each call station recording all birds observed both visually and by sound.

### ***2.7.5 Monumenting Target Species Sites and Recording Site Data***

Target special status plant species sites were generally monumented with orange-glo pin flags placed around population perimeters; exceptions are detailed below in 3.0 Results. Perimeters of most larger populations were recorded on global positioning system (GPS) units for subsequent use in Geographic Information Systems (GIS). All GPS coordinates in this report and associated documents are in Universal Transverse Mercator, Zone 10, North American Datum of 1983 projection. Data for locations of target special status species were recorded on standardized ORBIC report forms (Figure 2), submitted separately from this report. Applicable noxious weed populations were mapped, except for large infestations occupying the entire Study Area, which are noted below in 3.2 Noxious Weeds. GRSP detections were mapped. Special status plant and noxious weed population locations were documented in GIS; the related shapefiles are available upon request. Plant and bird inventory lists were documented in Excel spreadsheets.

## 3.0 RESULTS

The following sections detail the results of the field surveys.

### 3.1 Current Environment

Although the Study Area mostly slopes gently to the south and southwest, it covers a wide variety of aspects, with slopes ranging from approximately flat to steep. The Study Area can be viewed as three distinct habitats:

- Oak woodlands in the far northern part of the Study Area
- Meadows between the oak woodlands and the TID canal to the south
- Meadows downslope of the TID canal

#### 3.1.1 Oak Woodlands

The woodlands generally slope steeply to the north from a broad ridgeline and are dominated by *Quercus garryana*, *Toxicodendron diversilobum* (poison oak), *Symphoricarpos spp* (snowberries), *Prunus subcordata* (Klamath plum), the latter occasionally forming distinct thickets. These woodlands displayed a higher ratio of native versus non-native forb and grass coverage compared to the remainder of the Study Area; *Festuca idahoensis* ssp *roemeri* (Roemer's fescue, Idaho fescue) was found to be common here. Canopy cover varies greatly.

#### 3.1.2 Meadows Between the Oak Woodlands and TID Canal

The meadows between the oak woodlands and the TID canal were dominated primarily by exotic annual grasses and forbs, although dominant native species were also present. Dominant species included *Vicia villosa* (winter vetch), *Vicia sativa* (garden vetch), *Centaurea solstitialis*, *Poa bulbosa* (bulbous bluegrass), *Erodium cicutarium* (reds-stem stork's bill), *Geranium dissectum* (cutleaf geranium), *Geranium molle* (dovefoot geranium), *Avena fatua* (wild oat), *Elymus caput-medusae*, *Trifolium spp* (clovers), *Galium parisiense* (bedstraw), *Lomatium utriculatum* (common lomatium), *Tragopogon dubius* (yellow salsify), *Madia spp* (tarweeds), *Bromus japonicus* (field brome), *B. hordaceus* (soft brome), *B. tectorum* (cheatgrass), and *Vulpia microstachys* (small fescue). Of these, it is difficult to state what species were more dominant than others. These species occurred in varying concentrations across this part of the Study Area. Additionally, different species became more dominant as seasonal phenology progressed and early-bloomers senesced while late-bloomers became more prevalent. There were additional species that were very common, although not as abundant as the above dominants; these included *Calochortus tolmiei* (Tolmie startulip), *Dichelostemma capitatum* (bluedicks), *Calystegia occidentalis* (chaparral false bindweed), *Achyrachaena mollis* (blow-wives), and many others.

This area was historically grazed and likely was previously dominated by native bunchgrass communities. Non-native plant coverage during 2017 was approximately 85% or more on

average, with native species coverage at approximately 40%, on average. It was difficult to determine exact numbers for these percentages and other percentages given in this report with any accuracy, due to constantly changing plant phenology throughout the growing season and associated changes in biomass of any given species. Nonetheless, Pacific Crest personnel were expecting a higher non-native-to-native ratio than observed.

Outside of areas with summer moisture, shrubs comprised a very small amount of the vegetation coverage and consisted mostly of *Prunus subcordata*, *Toxicodendron diversilobum*, and exotic fruit trees, as scattered individuals and small patches.

Islands consisting primarily of native vegetation were found in this area, roughly overlapping with populations of *Ranunculus austro-oreganus* (see 3.2 Special Status Plants, Lichens, and Fungi), and were dominated by varying concentrations of native species including *Festuca idahoensis* ssp *roemeri*, *Horkelia daucifolia* (carrotleaf horkelia), *Eriophyllum lanatum* (Oregon sunshine), *Achnatherum lemmonii* (Lemmon's needlegrass), *Pseudoroegneria spicata* (bluebunch wheatgrass), *Phlox speciosa* (showy phlox), *Lomatium* spp (desertparslies), and *Achillea millefolium* (common yarrow), although non-native species were also common in these areas. These islands are likely not common in most adjacent parcels outside of the Study Area.

Large populations of *Microseris laciniata* ssp *detlingii* (Detling's silverpuffs) were also found, primarily on the flats and gentle slopes in the far northwest part of the Study Area; this species is endemic to southwestern Oregon and adjacent areas in northern California; it was previously a target special status species. *Plectritis congesta* (shortspur seablush) was abundant in the vicinity.

A small seasonal pond was found in the northeast part of T38S R1E S33, approximately 0.25 miles north of the eastern parking area and Property legal access point. Common plants here included *Lolium perenne* (perennial ryegrass), *Hordeum murinum* (mouse barley), and *Eleocharis* spp (spikerushes).

The oak woodlands mentioned above and the slopes between them and the TID canal likely serve as winter range for elk and deer. Elk and deer were observed in the Study Area during the survey efforts; the front cover of this report displays a herd of elk in the Study Area. Additional mammals incidentally observed in the Study Area included one black bear, three coyotes, one grey fox, and many smaller mammals.

### ***3.1.3 Meadows Downslope of the TID Canal***

This area had a much higher amount of moisture than areas upslope of the TID canal. This moisture originated from active irrigation diverted from the canal at multiple points along its length as it runs through the Study Area. Several natural springs and seeps added surface moisture; subterranean seepage from the canal was also a possible contributor. The vast majority of the area was observed to be grazed by livestock. Grazing was heavy throughout most of the area and extensive post-holing by cattle was evident. The exception was a narrow strip set apart by active electric fences located to either side of the drainage that runs south through the center of the Study Area; it is in this strip that the only *California macrophylla*

populations downslope of the canal were found. It was uncertain if this strip was part of the grazing lease, as it was fenced and had only light evidence of grazing, which may have originated from livestock that had escaped the fencing but had been quickly and efficiently recovered.

Vegetation in this area included many of the same species dominant upslope of the canal, but often in very different concentrations, with *Vicia spp*, *Calochortus tolmiei*, *Dichelostemma capitatum*, and others less common, while *Centaurea solstitialis* and others became more abundant. *Brassica rapa* (field mustard), *Shedonorus arundinaceus* (tall fescue), *Alopecurus pratense* (meadow foxtail), and others became dominant downslope of the canal, while existing only in traces upslope of the canal. *Shedonorus arundinaceus* and *Alopecurus pratense* were especially dominant in areas receiving higher volumes of irrigation water, notably at and upslope of the corrals in the southeast part of the Study Area. *Brassica rapa* was especially abundant in the southwest part of the Study Area near the POE. *Juncus effusus* (common rush) was common in some parts. *Rubus armeniacus* and *Rosa canina* (dog rose), with lesser amounts of *Rosa rubiginosa* (*R. eglentaria*; sweetbriar rose) were much more common downslope of the canal than upslope of it. Overall, vegetation in the area downslope of the canal was much denser, taller, and lusher than upslope of the canal; these conditions occasionally impeded foot travel when combined with the often irregular, post-holed, and wet ground surface. This condition receded later in the summer as plants senesced or were grazed down.

As with areas upslope of the TID canal, this area was historically grazed and likely was previously dominated by native bunchgrass communities. Non-native coverage in 2017 was approximately 98%, with native species coverage at approximately 15%, on average.

### ***3.1.4 Other Features of the Study Area***

Most drainages in the Study Area were lined with various concentrations of *Salix spp* (willows), *Rubus armeniacus*, *Carex densa* (dense sedge), *Juncus spp* (rushes), *Dipsacus fullonum* (Fuller's teasel), *Shedonorus arundinaceus*, and other typical riparian species; *Prunus cerasifolia* (cherry plum) was abundant in one drainage. Most of the drainages still had flowing water, at least in the lower stretches, at time of final revisits in early August; Hamby Spring in the southwest area downslope of the TID canal was still flowing strong.

The TID canal traverses the slope through the Study Area. It currently functions as a partial barrier to wildlife travel; certain terrestrial species may find it difficult to cross the flow of relatively deep water when the canal is flowing, although it should be noted that it does not flow for a substantial part of the year and travel may be less impeded then. There are two foot-bridges crossing the canal in the far western and eastern part of the Study Area, respectively, although the western one is composed of metal mesh that would likely inhibit most terrestrial wildlife travel during times of water flow in that canal. A maintenance road follows the canal for its length through the Study Area.

A wooden-pole powerline corridor exists in the northern part of the Study Area and a buried gas pipeline corridor roughly parallels it to the immediate south. Associated maintenance roads

follow these right-of-way corridors. A large pile of treated wood poles, assumingly associated with the powerline corridor construction, was observed at coordinates 524880E/4675620N. A small radio facility exists in the far southeast part of the Study Area near Eagle Mill Road and is accessible by vehicle from it.

A network of trails exists in the Study Area, observed to be used by people on foot, horseback, and OHV. People were seen from distance and personally encountered on the trails throughout the survey efforts, often in relatively large numbers. The trails, for the most part, were found to exist upslope of the TID canal. Most of the OHV use was observed in relation to the grazing leases downslope of the canal. However, OHV use was additionally observed on the trails in the western part of the survey area and their use was evident off-trail in that vicinity as well. The utility right-of-way corridors also had evidence of regular OHV use, much of which was assumingly in relation to infrastructure maintenance. Trails were observed cutting through multiple *California macrophylla* populations (see 3.2 Special Status Plants, Lichens, and Fungi) and trampling was evident at each of those populations. Rerouting of these trails may assist to lessen trampling.

### 3.2 Special Status Plants, Lichens, and Fungi

Fourteen populations (Figure 3) of *California macrophylla*, (Figure 4) totaling approximately 8.0 acres, were found in the Study Area. This species was originally documented in Oregon by Thomas Howell in 1887, with the associated herbarium collection noting “hills near Ashland”. It is possible that his original collection was made at one of the Study Area populations. ORBIC previously listed this species with an “EX” status (assumed to be extirpated in Oregon) until Pacific Crest personnel discovered a new location near the city of Eagle Point, Oregon. Since then, five populations were found in the Study Area by Kristi Mergenthaler and ODA personnel (personal communication, Kristi Mergenthaler, Southern Oregon Land Conservancy, 4/25/17). *California macrophylla* is currently listed by ORBIC (2016) with a “1” status (threatened or endangered throughout its range), the highest list status that ORBIC can assign. This species may soon receive additional listing through the State of Oregon. The Oregon sites represent the northern-most known extent of this species; it is also known from California and Baja California. The California Native Plant Society (2017) lists *California macrophylla* as a 1B.2 (rare, threatened, or endangered in CA and elsewhere).

Nearly all *California macrophylla* plants were found upslope and north of the TID canal. Two small populations were found downslope from the canal. Active grazing by livestock was observed downslope of the canal, where much of the ground had been trampled, whereas there was no current grazing by livestock observed upslope of the canal. Much of the ground downslope of the canal was observed to be irrigated. It is assumed that active grazing, associated trampling, and wet ground make for unfavorable conditions for the growth of *California macrophylla*. The two small populations downslope of the canal were found in an area between electric fences where grazing did not appear to be nearly as heavy as in the areas outside of the fencing, and irrigation was not evident at the time of population discovery.



One small plant rosette, potentially that of *California macrophylla* (Figure 3), was found downslope of the TID canal in early August. Due to immaturity and a lack of flowers and fruit, it was not possible to be certain of an identification. Although the leaves appear to be those of *California macrophylla*, the observed plant was growing well outside of the normal window of phenology for that species, did not have the reddish coloration that the stems and leaves of that species often have, was growing downslope of the canal in less desirable conditions, and all observed *C. macrophylla* plants in verified populations elsewhere in the Study Area were senescent at that time, casting doubt that the rosette in question was *C. macrophylla*. Nonetheless, it was monumented with several strips of yellow/black-striped flagging tied to small rocks in case a revisit would be made in future years.

Five populations (Figure 5) of *Ranunculus austro-oreganus* (Figure 6), totaling approximately 241 acres, were found in the Study Area, all upslope of the TID canal. The oak woodlands to the far north of the Study Area had the greatest concentrations. *Ranunculus austro-oreganus* is currently listed by ORBIC (2016) with a “1” status (threatened or endangered throughout its range), the highest list status that ORBIC can assign; it is also a state Candidate species with ODA. This species is endemic to Jackson County, found primarily in the Rogue Valley and adjacent foothills.

Approximately 633 acres contained vegetative *Ranunculus* plants (Figure 5), including overlap with verified *Ranunculus austro-oreganus* populations. Densities of vegetative plants within the 633 acres varied greatly, often being very widespread and isolated; very few existed downslope of the canal, those plants were also typically observed as depauperate. Due to a lack of flowers (a diagnostic characteristic for discerning *Ranunculus austro-oreganus*), it was not possible to know what species these vegetative plants were. They might flower in future years and a positive identification could then be made. However, it should be noted that no flowering *Ranunculus occidentalis* (western buttercup) was observed in the Study Area, the only other feasible species that the vegetative plants could be.

One site (Figure 5) of *Collema quadrifidum* (Figure 6) was found, present on multiple *Quercus garryana* trunks in approximately one acre of the oak woodlands in the far northern part of the Study Area. This tiny, gelatinous lichen is difficult to discern in the field, blending in with numerous other dark, similarly-sized lichens and blemishes on the tree trunks, and is best identified by its four-celled, polygonal spores as observed under the microscope. Due to the *Collema quadrifidum* being found off the ground on tree trunks, no pin flags were used to monument the site; a labeled set of yellow/black striped flagging was instead used, positioned on a tree trunk near the population center.

No *Fritillaria gentneri* were found. Vegetative *Fritillaria* plants were found, but these plants were impossible to identify to species without flowers. The vegetative plants were found only in the oak woodlands in the far northern part of the Study Area, existed in same vicinity as numerous flowering *Fritillaria affinis* (a non-target species), and possibly may all be that species. No target special status species of spring fungi were found.

### 3.3 Noxious Weeds

*Silybum marianum* (milk thistle; Figure 7) was found in one location along the southern boundary of the Study Area and consisted of approximately 80 specimens covering 10% of a >60 m<sup>2</sup> population area. The population extended from the property fenceline downslope to the POE exit ramp; it is likely that the POE was the vector of introduction. This species has rarely been found in southwestern Oregon. The Medford District BLM (personal communication, Bryan Wender, Medford District BLM Botanist, 8/14/17) has only one record of this invasive species on their lands, found in the Cow Creek Watershed of Douglas County. WeedMapper (ODA 2017b) revealed one site in Jackson County, near Rogue Valley International Airport.

*Spartium junceum* (Spanish broom; Figure 7) was found in two locations along the TID canal. Each location consisted of one plant. Though the populations sizes were very small, this species is reported here due to it being an uncommon invader in southwestern Oregon. WeedMapper (ODA 2017b) shows three sites in Jackson County, all in the far northern part of the county. Pacific Crest personnel know of one site in the City of Ashland, on Siskiyou Boulevard, which had apparently been treated (sprayed) recently.

*Rubus armeniacus* (Himalayan blackberry; Figure 7) was found throughout a substantial portion of the Study Area. The vast majority of the populations were found from the TID canal and downslope to the Study Area boundary. The average percent coverage within the population polygons was 15%. Besides large and dense infestations, multiple smaller infestations were also mapped, primarily in areas upslope of the canal where the species was much less common. A trace amount of *Rubus laciniatus* (cutleaf blackberry) was found mixed in with the *Rubus armeniacus*.

*Centaurea solstitialis* (yellow starthistle) and *Elymus caput-medusae* (medusahead rye) were found throughout the Study Area and are therefore not represented in Figure 7. Both species had an average coverage across the Study Area of approximately 35% each. Concentrations of both species were lighter in the oak woodlands in the far northern part of the Study Area, found most frequently in openings between trees, and heavier downslope of the TID canal.

*Cirsium arvense* (Canada thistle), *Cirsium vulgare* (bull thistle), *Conium maculatum* (poison hemlock), and *Phalaris arundinacea* (reed canarygrass) were all found as widely scattered, very small populations (often only as one isolated plant). These species were found primarily in areas of moisture along the TID canal and irrigated areas downslope of the canal. A small trace of *Hypericum perforatum* (St. Johnswort) was found along the canal. These species are not further documented in this report due to the small population sizes within the Study Area and overall commonness of these species in southwestern Oregon.

One potential population of *Cyperus esculentus* (yellow nutsedge) was found downslope of the canal (Figure 7). It was originally observed early in the season while immature and could not be confidently identified. By the time a return visit was made later in the season, cattle had grazed the plants down beyond recognition. The identification could therefore not be verified.

Although not a target noxious weed species, *Thinopyrum ponticum* (*Elymus elongata*; tall wheatgrass, European quackgrass) was observed infesting the entire gas pipeline right-of-way

in the northern part of the Study Area. The pipeline right-of-way was nearly a complete monoculture of *Thinopyrum ponticum*; it had outcompeted other vegetation and was spreading out from there. It has potential to quickly spread and take out other parts of the Study Area. This species is a pale grey-green color and this infestation is visible on aerial photography as a wide, pale strip cutting across the Study Area. This species is similar to the target noxious species *Elymus repens* (quackgrass, couchgrass), and many of the *Thinopyrum ponticum* specimens in the Study Area exhibited some features characteristic of *Elymus repens*, including very wide leaves and acute glumes, although the majority of features still pointed towards *T. ponticum*. There is potential for *Thinopyrum ponticum* to be considered by ODA for noxious weed listing in the future.

The Study Area has multiple possible vectors of noxious weed introduction including: Interstate 5 and associated POE adjacent to the Study Area, vehicular traffic within the Study Area (OHV's, right-of-way maintenance vehicles), livestock, TID canal, non-vehicular trail traffic (foot, bike, horse). Much of the vegetation between the southern property fenceline and Interstate 5 / POE is mowed annually, possibly slowing the spread of weeds from those two dispersal vectors, although the stretch of exit ramp with the *Silybum marianum* had not been mowed; it may be too steep to maintain.

### 3.4 Birds

Thirty-four GRSPs were detected during the surveys (Figure 8). The majority (thirty-two of thirty-four) were singing males; two GRSPs were flushed from vegetation and the sex of these two birds is unknown. Thirty-two detections were recorded above the TID canal; two were recorded below the canal. GRSPs have Federal Species of Concern, Oregon Department of Fish and Wildlife (ODFW) Conservation Strategy Species, and ORBIC2 status. See Table 4 for ORBIC rank definitions. Other special status bird species detected during the point counts include:

- Acorn woodpecker (*Melanerpes formicivorus*)
- Chipping sparrow (*Spizella passerine*)
- Oak titmouse (*Baeolophus inornatus*)
- Peregrine falcon (*Falco peregrinus*)
- Western meadowlark (*Sturnella neglecta*)
- White-tailed kite (*Elanus leucurus*)
- Yellow-breasted chat (*Icteria virens*)

Table 4 contains further notes on the occurrences of the target special status avian species within the Study Area and includes rank status(es) of each species.

### 3.5 Other Sites of Interest

No other special status target species were found.

Multiple populations of a species of *Phaeocalicium* (Figure 9) were found on twigs of *Quercus garryana* in the oak woodlands in the far northern part of the Study Area. Species of *Phaeocalicium* belong to a group of organisms commonly known as pin lichens. Their spores

are borne atop a small stalk and are distributed by wind and insects travelling the length of the twigs. The collection in question is similar to *Phaeocalicium interruptum*, a species without special status, but differs by multiple morphological and chemical features. It is possibly a new species: one that is new to science, not described, and un-named. A collection has been sent to a pin lichen expert for another opinion; this report will be updated when a determination has been returned. Determinations are also still out for several invertebrate collections.

One large “log” of petrified wood (Figure 10) was found along the boundary of the study area at coordinates 525445E/4675296N, placing it just within the Study Area. The overall length is unknown; it continued underground and its large size and heavy weight prevented movement and further exploration. This feature may serve as an attraction to visitors.

A series of scattered rock outcrops exist on a steep south-facing slope running approximately 0.25 miles west-east through the Study Area in T38S R1E S27. Other, smaller sets of outcrops are occasional throughout much of the Study Area.

No other biological sites of interest, as defined in 1.0 Introduction, were found in the Study Area.

### **3.6 Inventories**

A total of two-hundred-fifty-two vascular plants were recorded during the surveys (Table 5). Note that multiple taxa are not identified past genus. Additionally, several recorded taxa were observed only along Butler Creek; it is uncertain how much of that creek actually exists in the project area due to conflicts in GIS mapping compared to on-the-ground property line evidence. Pacific Crest elected to document all bird species detected during the surveys for GRSPs; fifty-three avian species in total were detected (Table 4).

## 4.0 REFERENCES

- Altman, B. 2015. Oregon Vesper Sparrow Range-wide Inventory and Habitat Assessment Final Report. Prepared by the American Bird Conservancy for the Center for Natural Lands Management. April 2015.
- AmphibiaWeb. 2017. University of California, Berkeley, CA, USA. Accessed 8/22/17. Available: <<http://amphibiaweb.org>>
- Birds of North America. 2017. Online version. Species accounts. Cornell Ornithology Lab and American Ornithologist's Union. Accessed 8/22/17. Available: <http://bna.birds.cornell.edu/bna/>
- Calflora. 2017. Searchable Database of California Flora. Berkeley, California. Accessed 8/12/17. Available: <http://www.calflora.org/>
- Csuti, B., T. A. O'Neil, M.M. Shaughnessy, E.P. Gaines, and J.C. Hak. 2001. Atlas of Oregon wildlife: distribution, habitat, and natural history, Second edition. Oregon State University Press. Corvallis, OR.
- eBird. 2017. eBird: An online database of bird distribution and abundance. eBird, Ithaca, New York. Accessed 8/21/17. Available: <http://www.ebird.org>
- Nafis, G. 2017. California Herps - A Guide to the Amphibians and Reptiles of California. Accessed: 8/21/17. Available: <http://www.californiaherps.com/>
- NatureServe. 2017. NatureServe Explorer Database. Species accounts. Accessed: 8/22/17. Available: <http://www.natureserve.org/explorer/>
- ODA (Oregon Department of Agriculture). 2017a. Noxious Weed Policy and Classification System. Oregon Department of Agriculture Noxious Weed Program. Salem, Oregon. Available: <http://www.oregon.gov/ODA/programs/Weeds/OregonNoxiousWeeds/Pages/Law.aspx>
- ODA (Oregon Department of Agriculture). 2017b. WeedMapper. Searchable Database of Oregon Noxious Weed Occurrences. Oregon Department of Agriculture Noxious Weed Program. Salem, Oregon. Accessed 8/13/17. Available: <http://www.oregon.gov/ODA/programs/Weeds/Pages/WeedMapper.aspx>
- ODFW (Oregon Department of Fish and Wildlife). 2017a. Threatened, Endangered, and Candidate Fish and Wildlife Species in Oregon. Accessed 8/20/17. Available: [http://www.dfw.state.or.us/wildlife/diversity/species/threatened\\_endangered\\_species.asp](http://www.dfw.state.or.us/wildlife/diversity/species/threatened_endangered_species.asp)
- ODFW (Oregon Department of Fish and Wildlife). 2017b. Sensitive Species List (2016). Accessed 8/20/17. Available: [http://www.dfw.state.or.us/wildlife/diversity/species/sensitive\\_species.asp](http://www.dfw.state.or.us/wildlife/diversity/species/sensitive_species.asp)
- ODFW (Oregon Department of Fish and Wildlife). 2017. Wildlife Division Gray Wolves Page. Accessed 8/24/17. Available: <http://www.dfw.state.or.us/Wolves/Packs/Keno.asp>

- ODFW (Oregon Department of Fish and Wildlife). 1984. The Herpetology of Jackson and Josephine Counties, Oregon. Nongame Wildlife Program. Technical Report #84-2-05. By Alan D. St. John.
- OFP (Oregon Flora Project). 2017. Oregon Flora Project Atlas and Rare Plant Guide. Department of Botany and Plant Pathology, Oregon State University, Corvallis, Oregon. Accessed at: <http://www.oregonflora.org>
- ORBIC (Oregon Biodiversity Information Center). 2016. Rare, Threatened, and Endangered Species of Oregon. Portland State University, Portland, Oregon. Available: <http://inr.oregonstate.edu/orbic/rare-species/rare-species-oregon-publications>
- Oregon Wildlife Institute. 2017. Wildlife Conservation in Willamette Valley Grassland & Oak Habitats Species Account. Slender-billed Nuthatch (*Sitta carolineses aculeata*). Accessed 8/24/17. Available: [www.oregonwildlife.org](http://www.oregonwildlife.org).
- Rogue Valley Audubon Society. 2014. Birds of Jackson County, Oregon, Distribution and Abundance. By the Jackson County Checklist Committee. Fourth Edition, 2014. Published by Rogue Valley Audubon Society.
- Stephens, J.L. 2016. Grasshopper Sparrow abundance on the Imperatrice property: Results from 2016 surveys. Klamath Bird Observatory. Rep. No. KBO-2016-0009. September 30, 2016.
- USFS and BLM (US Department of the Interior, Bureau of Land Management). 1999a. Survey and Manage Survey Protocols – Protection Buffer Bryophytes Version 2.0.
- USFS and BLM. 1999b. Survey and Manage Survey Protocol – Vascular Plants.
- USFS and BLM. 2001. Record of Decision and Standards and Guidelines for Amendments to the Survey and Manage, Protection Buffer, and other Mitigation Measures Standards and Guidelines. January 2001.
- USFS and BLM. 2003a. Survey Protocols for Survey and Manage Category A and C Lichens in the Northwest Forest Plan Area. Version 2.1.
- USFS and BLM. 2003b. 2003 Amendment to the Survey Protocol for Survey & Manage Category A & C Lichens in the Northwest Forest Plan Area. Version 2.1 Amendment. September 2003.
- USFS and BLM. 2006. Survey Protocol Guidance for Conducting Equivalent Effort Surveys under the Northwest Forest Plan Survey and Manage Standards and Guidelines.
- USFWS (United States Fish and Wildlife Service). 2017. General Species List Generator. Oregon Fish and Wildlife Office. Accessed 8/20/17. Available: <https://ecos.fws.gov/ipac>
- USFWS (United States Fish and Wildlife Service). 2016. Recovery Outline for the Streaked Horned Lark (*Eremophila alpestris strigata*). Portland, Oregon. 42 pp.
- Van Norman, K., J. Lippert, D. Rivers-Pankratz, R. Holmes, and C. Mayrsohn. 2008. Sporocarp Survey Protocol for Macrofungi, version 1.0. Portland, Oregon. Interagency Special

Status / Sensitive Species Program. US Department of Interior, Bureau of Land Management, Oregon/Washington and US Department of Agriculture, Forest Service, Region 6. 16 pp.

Van Norman, K. 2010. Survey & Manage Category B Fungi Equivalent-Effort Survey Protocol, version 1.0. Portland, OR. US Department of Interior, Bureau of Land Management, Oregon/Washington and US Department of Agriculture, Forest Service, Region 6. 17 pp.

Verts, B.J. and L.N. Carraway. 1998. *Land Mammals of Oregon*. University of California Press. Berkeley and Los Angeles, CA.

## Tables

Table 1: Special status plants, lichens, and fungi

Taxon <sub>1</sub>	Scientific Name	Federal Status <sub>2</sub>	ODA Status <sub>2</sub>	SEN/STR/S&M <sub>2</sub>	Likelihood to Exist in Study Area
VA	<i>Adiantum jordanii</i>			SEN	LOW. Habitat includes moist, shaded hillsides, springs, riparian areas. SW OR populations are found mostly on serpentine. No serpentine present in Study Area, although shaded riparian is found to far north in oak woodland. Most known sites in SW OR are west of Grants Pass.
VA	<i>Agrostis hendersonii</i>			STR	LOW. Found in vernal pools and other moist areas in valley grasslands. Historical collection from Sams Valley, but is presumed extirpated in OR.
FU	<i>Albatrellus ellisii</i>			SM-B	LOW. Typically found in older coniferous forests at higher elevations than the Study Area, although this species is occasionally found in hardwood communities. Known sites exist in coniferous forest in the SW OR Cascades.
VA	<i>Allium bolanderi</i> var. <i>bolanderi</i>			STR	MODERATE. Habitat includes rocky clay soils, although this species typically prefers serpentine. Known sites exist at Howard Prairie and NW of Grants Pass.
VA	<i>Allium peninsulare</i>			SEN	LOW. Habitat includes meadows. Many populations nearby in the Hyatt / Howard Prairie area, although these populations are found in higher elevation snowmelt meadows. Found at lower elevations in CA.
VA	<i>Androsace elongata</i> ssp. <i>acuta</i>			STR	LOW. Habitat includes dry, primarily north-facing meadows. Previously known from one historic site in Jackson County (1887), now assumed to be extirpated.
BR	<i>Anoetangium aestivum</i>			STR	LOW. Lower elevation springs and seeps, often over rock, although this species typically prefers calcareous substrates in SW OR. One known site near Wimer.
VA	<i>Arabis modesta</i>			SEN	MODERATE. Shaded slopes at low to moderate elevations; often associated with rock. Known sites near Shady Cove, Applegate, and NW of Grants Pass.
VA	<i>Astragalus californicus</i>			SEN	LOW. Low to moderate elevation, dry, open, meadows, woodlands, shrub communities; although known sites in SW OR are south of the Siskiyou crest in CSNM (Cascade-Siskiyou National Monument) and found in communities more similar to those of the Great Basin.
VA	<i>Astragalus gambelanus</i>			SEN	MODERATE. Dry, open, grassy areas at low to moderate elevations. Known sites at Sampson Creek and southern part of CSNM.
FU	<i>Balsamia nigrans</i>			STR	MODERATE. Associates with species of oak. Known sites in Jackson and Josephine counties.
BR	<i>Bryum calobryoides</i>			SEN	LOW. Prefers crevices in rock at higher elevations, occasionally found at lower elevations. Known sites along Siskiyou crest and near Hyatt Lake.
LI	<i>Calicium quercinum</i>			STR	LOW. Grows on trunks of oaks at low to moderate elevations. No known sites in SW OR.



VA	<i>California macrophylla</i>			SEN	PRESENT. Known sites documented in Study Area prior to 2017 survey efforts.
VA	<i>Callitriche marginata</i>			SEN	MODERATE. Primarily a species of vernal pool habitats in SW OR. Could exist in seasonal pond in east part of Study Area or vernal pools undetected during desktop analysis.
VA	<i>Calochortus greenii</i>		SC	SEN	LOW. Habitat includes clay soils in meadows, shrub communities, and other areas of exposure as low as 2400' elevation. However, all known sites in SW OR are in the Klamath watershed south of the Siskiyou crest.
VA	<i>Calochortus monophyllus</i>			SEN	MODERATE. One known site upslope of the Study Area, higher in elevation, in forest on the south side of Grizzly Peak. Can grow as low as 1300' elevation.
VA	<i>Calochortus nitidus</i>			STR	LOW. Habitat includes meadows. One known site near Greensprings, although it is much higher in elevation than the Study Area.
BR	<i>Campylopus subulatus</i>			STR	LOW. Found from sea level to moderate elevations. Known sites in Josephine County. Prefers areas without human-induced disturbance or heavy plant competition.
VA	<i>Carex comosa</i>			SEN	LOW. Found in wet areas from sea level to 1200'. Nearest known site is historic, found along the Rogue River.
VA	<i>Carex crawfordii</i>			STR	MODERATE. Found at pond and lake margins that dry up in summer, from sea level to moderate elevations. Rumored site near Grizzly Peak.
FU	<i>Cazia flexiascus</i>			STR	MODERATE. Associates with <i>Quercus garryana</i> and other hardwoods. Known sites in Rogue Valley vicinity.
VA	<i>Cheilanthes covillei</i>			SEN	MODERATE. Rock crevices at a variety of elevations and plant communities. Known sites in Jackson County near Heppsie Mt.
VA	<i>Cheilanthes intertexta</i>			SEN	MODERATE. Rock crevices at a variety of elevations and plant communities. Known sites throughout Jackson County.
VA	<i>Chlorogalum angustifolium</i>			SEN	MODERATE. Clay soils of dry areas with high light exposure at lower elevations. Widely scattered known sites in Jackson and Josephine counties.
FU	<i>Clavariadelphus occidentalis</i>			SM-B	LOW. Typically a species of mixed and coniferous forests, although it is rarely found in hardwood communities. Many known sites in southern Oregon.
FU	<i>Clavariadelphus subfastigiatus</i>			STR	LOW. Typically a species of mixed and coniferous forests, although it is rarely found in hardwood communities. Three known sites in SW OR.
LI	<i>Collema quadrifidum</i>			STR	PRESENT. Prefers <i>Quercus garryana</i> trunks at low to moderate elevations. Many known sites in Jackson County.
VA	<i>Cryptantha milobakeri</i>			SEN	MODERATE. Rocky or gravelly slopes at low to moderate elevations. Known sites in Jackson (Applegate area) and eastern Josephine counties.
VA	<i>Cyperus acuminatus</i>			SEN	LOW. Found at vernal pools, seasonal ponds, ditches, and other wet areas at low elevations. The only previously known sites in SW OR are historic and near Grants Pass.
VA	<i>Delphinium nudicaule</i>			SEN	LOW. Grows in well-drained areas (often talus or gravel) and along river banks and low to moderate elevations. Known sites in Jackson and Josephine counties.

FU	<i>Dendrocollybia racemosa</i>			STR	MODERATE. Found on decayed remains of other mushrooms in a variety of habitats (including hardwood and shrub communities) at low to moderate elevations. Known sites in Jackson (mostly near Shady Cove) and Josephine counties, including one at French Flat found under manzanita.
BR	<i>Didymodon norrisii</i>			STR	MODERATE. Habitat includes a variety of rock substrates in a variety of plant communities from low to moderate elevations. Known sites in Jackson County near Siskiyou Summit and Shady Cove.
VA	<i>Diplacus bolanderi</i>			SEN	MODERATE. Grassy areas and openings in chaparral from low to moderate elevations. Observed in areas of disturbance. Known sites in Applegate Valley.
VA	<i>Diplacus congdonii</i>			SEN	MODERATE. Oak woodlands, grassy areas, and openings in chaparral from low to moderate elevations. Known sites in Applegate Valley.
BR	<i>Entosthodon californicus</i>			STR	MODERATE. Found on clay soils in seasonally wet areas, often associated with disturbance. Known sites at Table Rocks.
BR	<i>Entosthodon fascicularis</i>			SEN	MODERATE. Found on a variety of soils in seasonally wet areas, often associated with disturbance. Known sites near Grants Pass.
BR	<i>Ephemerum crassinervium</i>			SEN	MODERATE. Found on a variety of soils in seasonally wet areas, often associated with disturbance; one Jackson County site was found in water-filled cow tracks. Known sites in Jackson and Josephine counties.
VA	<i>Ericameria arborescens</i>			SEN	LOW. Dry forest, hardwood and shrub communities at low to moderate elevations, often in foothills. Only known sites in OR are in western Curry County; however, it is found throughout CA in a variety of habitats.
VA	<i>Erigeron cervinus</i>			SEN	LOW. Prefers rocky areas, but also grows in open areas. Usually at moderate to higher elevations. Occasionally found in vernal wet areas at lower elevations. Nearest known site is in Josephine County.
VA	<i>Eschscholzia caespitosa</i>			SEN	LOW. Dry, often brushy areas at lower elevations. Nearest known sites are near Glendale and Hellgate.
VA	<i>Eucephalus vialis</i>		ST	SEN	MODERATE. Low to moderate elevation ecotones, but generally involving coniferous and mixed forest.
VA	<i>Fritillaria eastwoodiae</i>			STR	LOW. Dry slopes. Rumored sites at Lower Table Rock and near Gold Hill, otherwise no sites in close proximity.
VA	<i>Fritillaria gentneri</i>	FE	SE	SEN	HIGH. Low to high elevation ecotones, mixed forests, shrub communities. Study Area is well within species range and known sites are in relatively close proximity.
VA	<i>Hackelia bella</i>			SEN	LOW. Moderate to higher elevations. Known from Table Mountain and Grizzly Peak vicinity, but at higher elevations.
VA	<i>Horkelia tridentata</i> ssp. <i>tridentata</i>			SEN	LOW. Dry areas, typically in open forest, on granitic or other igneous soils, at low to high elevations. Known sites are in Ashland Watershed, although these are higher elevation than the Study Area, found exclusively on granite, and favor ridgelines.
VA	<i>Juncus kelloggii</i>			STR	LOW. Vernal pools, springs, meadows at low elevations. ORBIC lists a known site in Josephine County.
LI	<i>Leptogium burnetiae</i>			STR	MODERATE. Found on <i>Quercus garryana</i> trunks at low to moderate elevations. Nearest known verified site is near Shady Cove; another unverified site exists near Buckhorn Springs.

VA	<i>Limnanthes alba ssp. gracilis</i>		SC	SEN	LOW. Wet meadows, streamsides, ditches, cliff bases at typically low elevations. Only one known site in Jackson County, found near City of Rogue River.
VA	<i>Limnanthes floccosa ssp. bellingeriana</i>		SC	SEN	MODERATE. Vernal wet areas with high light exposure, from low to moderate elevations. Many known sites in Cascades of Jackson County.
VA	<i>Limnanthes pumila ssp. grandiflora</i>	FE	SE	SEN	LOW. Vernal pool habitat, but endemic to Agate Desert.
VA	<i>Limnanthes pumila ssp. pumila</i>		ST	SEN	LOW. Vernal pool habitat, but endemic to Table Rocks.
VA	<i>Lomatium cookii</i>	FE	SE	SEN	LOW. Vernal moist habitats, often vernal pools. Known from two concentrations of populations: one in the Agate Desert, the other in the Illinois Valley.
VA	<i>Meconella oregana</i>		SC	SEN	MODERATE. Found in a variety of plant communities, often vernal moist, usually with moderate to high light exposure, at low elevations. Known populations near Medford, Jacksonville, Applegate.
VA	<i>Microseris douglasii ssp. douglasii</i>			STR	LOW. Meadows with heavy clay soils. The only known site in Oregon was near Ashland but has not been seen since the late 1800's and is presumed extirpated.
VA	<i>Nemacladus capillaris</i>			SEN	MODERATE. Dry slopes at a variety of elevations. In SW OR, prefers meadow edges in areas of higher percentages of bare mineral soil. Multiple known sites in Cascades of Jackson County, especially in CSNM.
BR	<i>Orthotrichum bolanderi</i>			STR	MODERATE. Rock features at low to moderate elevations in a variety of plant communities. Known sites near Sampson Creek and Medford.
BR	<i>Orthotrichum euryphyllum</i>			STR	MODERATE. Rocks in seasonal drainages, usually with moderate to high light exposure. Known sites throughout much of the Cascades of southern OR.
BR	<i>Orthotrichum hallii</i>			STR	MODERATE. Rock features at low to moderate elevations in a variety of plant communities. Known sites near Medford.
VA	<i>Pellaea andromedifolia</i>			SEN	MODERATE. Rocky areas at low to moderate elevations. Known sites in Jackson and Josephine counties.
LI	<i>Peltigera pacifica</i>			SM-E	LOW. Typically in coniferous or mixed forests, but can be found in a variety of habitats. Known site in Ashland Watershed; several more in western Jackson County.
LI	<i>Peltula euploca</i>			STR	MODERATE. Rocky areas (basalt, andesite) at lower elevations. Known sites at Upper Table Rock, Applegate Valley, CSNM, Horseshoe Ranch.
BR	<i>Phymatoceros phymatodes</i>			SEN	LOW. Mineral soil substrates that remain wet late into summer. Multiple sites on Medford BLM lands in SW OR.
VA	<i>Pilularia americana</i>			SEN	MODERATE. Vernal wet habitats including vernal pools and pond margins. Known sites at Table Rocks.
VA	<i>Pinus sabiniana</i>			STR	LOW. Foothill woodlands at low to moderate elevations. Common in the Trinity Mountains and elsewhere in CA; very rare in OR as natural sites, but is frequently planted as an ornamental.
VA	<i>Plagiobothrys austiniiae</i>			SEN	MODERATE. Vernal wet areas, such as seeps and ephemeral drainages, typically in meadows, at low to moderate elevations. Known sites at Table Rocks and Cascades of Jackson County.

VA	<i>Plagiobothrys figuratus</i> ssp. <i>corallicarpus</i>		SC	SEN	MODERATE. Vernal wet areas, often rocky, in meadows at low to moderate elevations. Known sites near Greensprings, Medford, Grants Pass.
VA	<i>Plagiobothrys greenei</i>			SEN	MODERATE. Vernal wet areas, such as seeps and ephemeral drainages, typically in meadows, at low to moderate elevations. Known sites in the Cascades of Jackson County.
VA	<i>Plagiobothrys lamprocarpus</i>		SE	STR	LOW. Assumed habitat is vernal wet areas with higher light exposure. Known only from one historic site near Grants Pass (1921) and is assumed extinct.
BR	<i>Porella bolanderi</i>			SEN	MODERATE. Rock outcrops in oak woodlands. Known sites in Ashland Watershed and Cascades of Jackson County.
FU	<i>Psathyrella quercicola</i>			STR	MODERATE. Grows on <i>Quercus garryana</i> at low elevations. Known sites from Jackson and Josephine counties, including the type locality.
BR	<i>Racomitrium depressum</i>			SEN	LOW. Rocks along ephemeral drainages with high light exposure, mostly at moderate to higher elevations in southern OR. Known sites near Howard Prairie.
VA	<i>Rafinesquia californica</i>			SEN	MODERATE. Meadows and post-burn areas in variety of communities from low to high elevations. Large, robust populations were previously found throughout the Squire and Quartz Fire areas in the Applegate.
VA	<i>Ranunculus austro-oreganus</i>		SC	SEN	PRESENT. Known sites documented in Study Area prior to 2017 survey efforts.
VA	<i>Rhamnus ilicifolia</i>			SEN	MODERATE. Chaparral and oak woodlands from low to moderate elevations. Several sites known along the CA border in CSNM and near Applegate Ranger Station.
VA	<i>Rhynchospora alba</i>			SEN	LOW. Wet areas from low to high elevations. Known sites in southern OR are moderate to high elevation, often associated with <i>Sphagnum</i> .
VA	<i>Ribes divaricatum</i> var. <i>pubiflorum</i>			SEN	LOW. Wet areas and forest edges. Multiple known sites west of Grants Pass.
VA	<i>Romanzoffia thompsonii</i>			SEN	LOW. Vernal wet areas, such as seeps and springs, on steep slopes with high light exposure. Known SW OR sites are near Flounce Rock at ~4000' elevation.
FU	<i>Sarcodon fuscoindicus</i>			STR	LOW. Typically found in coniferous forests but occasionally in hardwoods. Widely scattered across western OR, including one site in northern Jackson County.
BR	<i>Schistidium cinclidodonteum</i>			SEN	MODERATE. Rocks in seasonal drainages, usually with moderate to high light exposure. Known sites throughout much of the Cascades of southern OR.
VA	<i>Scirpus pendulus</i>			SEN	MODERATE. Wet areas in a variety of plant communities from low to moderate elevations. Known site near Grizzly Peak.
VA	<i>Sidalcea hickmanii</i> ssp. <i>petraea</i>			SEN	LOW. Dry shrub communities on ridges. One known site: Sams Valley
VA	<i>Solanum parishii</i>			SEN	MODERATE. Found in a variety of dry plant communities at a variety of elevations. Known sites throughout much of Jackson County.
FU	<i>Spathularia flavida</i>			SM-B	LOW. Typically found in coniferous forests and only rarely in hardwoods. Numerous known sites in Jackson County.

VA	<i>Tetrapteron graciliflorum</i>			SEN	MODERATE. Meadows, shrub communities, oak woodlands at low to moderate elevations. Known sites in CSNM, Applegate vicinity, and elsewhere in Jackson County.
BR	<i>Trichostomum tenuirostris</i> var. <i>tenuirostris</i>			STR	LOW. Various moist substrates in various plant communities at a wide variety of elevations. Known site near Wagner Butte is in coniferous forest.
VA	<i>Triteleia ixioides</i> ssp. <i>scabra</i>			STR	LOW. Foothill meadows and woodlands, in clay and granitic soils.
LI	<i>Umbilicaria hirsuta</i>			STR	MODERATE. Rock features in a variety of exposures in a variety of elevations. Known sites in CSNM, near Lake of the Woods, near Wimer.
VA	<i>Wolffia borealis</i>			SEN	LOW. Areas of stagnant water such as ponds, lakes. Known sites at Parsnip Lakes in CSNM, and Sharron Fen, both at ~4500' elevation.
VA	<i>Wolffia columbiana</i>			SEN	LOW. Areas of stagnant water such as ponds, lakes. Known site near Gold Hill.

<sup>1</sup> VA = vascular plant, BR = bryophyte, LI = lichen, FU = Fungus

<sup>2</sup> Federally Listed Species: FE = Endangered, FT = Threatened, SOC = Species of Concern. Oregon Department of Agriculture: SE = Endangered, ST = Threatened, SC = Candidate. STR = Strategic, SEN = Sensitive. S/M Category definitions: Category A = Manage all known sites; pre-disturbance surveys practical, strategic surveys. Category B = Manage all known sites; pre-disturbance surveys not practical and not applicable; strategic surveys; equivalent effort surveys required for most bryophytes, lichens and fungi for habitat-disturbing projects in old growth. Category C = Manage high-priority sites; pre-disturbance surveys practical; strategic surveys. Category D = Manage high-priority sites; pre-disturbance surveys not practical or not necessary; strategic surveys. Category E = Manage all known sites; pre-disturbance survey not applicable; strategic surveys. Category F = known site management and pre-disturbance surveys not applicable; strategic surveys

Table 2: Special status birds, mammals, reptiles, and amphibians

Common Name <i>Scientific Name</i>	Status <sup>1</sup> USFWS STATE ORBIC	Habitat and Ecology	Likelihood to Occur in the Study Area
<b>Amphibians</b>			
Northern red-legged frog <i>Rana aurora</i>	SOC SV, CS 4	Prefers cool and calm or still waters of streams, marshes or ponds, often near or in moist forests. Breed in winter and early spring laying eggs attached to stems of emergent vegetation or submerged branches in permanent water bodies. Highly terrestrial outside of the breeding season. Known historically from Jackson County including records in the Lower Rogue and Applegate watersheds.	<b>Low.</b> Suitable habitat for this species exists along the drainage in the middle of the Study Area and in adjacent wetlands areas; however, it is overall of marginal quality for this species (not within humid woodlands).
western toad <i>Anaxyrus boreas boreas</i>	- SV, CS 4	Historically found throughout Jackson County near aquatic sites (streams, rivers, lakes, ponds, and springs). Occurs in a wide variety of habitats as long as there is suitable aquatic habitat for breeding and is adapted to agricultural environments such as vegetated irrigation canals.	<b>Moderate.</b> Suitable habitat for this species exists along the drainage in the middle of the Study Area and in adjacent wetlands areas; however, this species has disappeared from much of its original range and is now uncommon.
<b>Reptiles</b>			
California kingsnake <i>Lampropeltis californiae</i>	SOC SV 4	Found in a wide variety of habitats. In Oregon, it occurs along the Rogue and Umpqua river valleys, often in dense vegetation along watercourses but also in farmland, chaparral, and deciduous and mixed conifer woodlands.	<b>Moderate.</b> There is suitable habitat in the Study Area and there are historic records of this species in the region.
California mountain kingsnake <i>Lampropeltis zonata</i>	SOC SV, CS 4	Found in a diversity of habitats often pine forests, oak woodlands, and chaparral; commonly in open wooded areas near streams.	<b>Moderate.</b> There is suitable habitat in the Study Area and there are historic records of this species in the region.
western rattlesnake <i>Croatalus oreganus</i> ssp. <i>oreganus</i>	- SC 4	Occurs in a variety of habitats from deserts to chaparral to open forests, usually near rocks, cliffs, or downed logs.	<b>Present.</b> There is suitable habitat for this species in the Study Area and they were observed in the Study Area during surveys.
<b>Birds</b>			
white-tailed kite <i>Elanus leucurus</i>	- - 4	Lower elevation grasslands, agricultural areas, meadow, oak woodlands, riparian woodlands, marshes and wetlands; nest in trees or tall shrubs. Breeding season is approximately February to July.	<b>Present.</b> This species was observed flying over the Study Area and hunting nearby on several occasions during the breeding season (early May). No nest was observed in the Study Area and there is only limited suitable nesting trees/shrubs available within the Study Area; most likely this bird was nesting nearby, possibly in the trees growing on the adjacent property to the east.

Common Name <i>Scientific Name</i>	Status <sup>1</sup> USFWS STATE ORBIC	Habitat and Ecology	Likelihood to Occur in the Study Area
bald eagle <i>Haliaeetus leucocephalus</i>	BCC SV 4	This species is usually found near water and breeds in forested areas adjacent to large bodies of water. Nests in trees, rarely on cliff faces and on the ground in treeless areas.	<b>Present (assumed).</b> There is no suitable nesting habitat for this species in the Study Area, though it is fairly common to the greater region. A bald eagle was recorded in the Study Area on April 29, 2016 and on January 4 and 25, 2014 (eBird 2017).
ferruginous hawk <i>Buteo regalis</i>	SOC SV, CS 4	A rare, but regular winter visitor to Jackson County. Prefers flat, rolling grasslands or shrubsteppe regions including sagebrush shrublands, and edges of western juniper and pinyon-juniper woodlands and other forests. Breeds in northeastern Oregon and found year-round in southeastern Oregon.	<b>Present (assumed).</b> A ferruginous hawk was observed in the Study Area on February 27, 2017 (eBird 2017).
golden eagle <i>Aquila chrysaetos</i>	- - 4	Inhabits a wide variety of open and semi-open habitat types including grasslands, shrublands, woodlands, and coniferous forests. Often nests on cliffs bordering rivers, will also nest in trees, on ground, on river banks, and on human-made structures.	<b>Present (assumed).</b> There is suitable foraging habitat for this species in the Study Area though it would be unlikely to nest there due to a lack of preferred nesting habitat. Two golden eagles were observed in the Study Area on March 4, 2016 and on January 4, 2014 and one was observed in the Study Area on January 25, 2014 (eBird 2017).
short-eared owl <i>Asio flammeus</i>	BCC (year-round) CS 3	Inhabits open terrain, most often marshes, but also grasslands, dunes, agricultural fields, meadows, and pastures. Breeding season is typically from April to August.	<b>Low.</b> Suitable habitat exists in the Study Area; however, this species is a rare to irregular visitor to Jackson County during the non-breeding season (November - April).
burrowing owl <i>Athene cunicularia</i>	SOC SC, CS 4	Habitat includes deserts, open grasslands, shrublands, and other open areas such as vacant lots near human habitation or airports. Nests in abandoned mammal burrows. They have disappeared from the Rogue Valley and are rare in Jackson County, though they once were considered common.	<b>Low.</b> The Study Area contains suitable habitat for this species; however burrowing owls are not currently known to breed in Jackson County and are considered a rare to irregular visitor during the non-breeding season (October - April).
common nighthawk <i>Chordeiles minor</i>	- CS 4	Forage over wide variety of habitats throughout the state. Nest on bare ground in open areas. Breeding season is typically June to August.	<b>Moderate.</b> Species may forage over the Study Area, only reside in the Rogue Valley during the breeding season; unlikely to nest in the Study Area because of limited bare ground.

Common Name <i>Scientific Name</i>	Status <sup>1</sup> USFWS STATE ORBIC	Habitat and Ecology	Likelihood to Occur in the Study Area
rufous hummingbird <i>Selasphorus rufus</i>	BCC (breeding)	Breed in the Rogue Valley, typically in open forest near meadows and riparian thickets in mountainous areas. Breeding typically begins anywhere from April to July, depending on elevation.	<b>Moderate.</b> This species is common in the region however the habitat in the Study Area is only marginally suitable breeding habitat for this species.
Lewis's woodpecker <i>Melanerpes lewis</i>	SOC/BCC (year-round) SC, CS 2	Typically inhabits open forests at lower elevations. Nests in white oak, ponderosa pine, mixed oak-pine, and cottonwood riparian woodlands of eastern Oregon (also in the Klamath River drainage). Common in the Rogue Valley from November through March.	<b>Present (assumed).</b> Limited suitable habitat for this species occurs in the Study Area and it is likely to pass through the Study Area during winter foraging. There is a record of six Lewis's woodpeckers in the Study Area from January 4, 2014 (eBird 2017).
acorn woodpecker <i>Melanerpes formicivorus</i>	SOC CS 4	Occur in oak woodlands, mixed oak-pine woodlands and oak savannah. Primary food is acorns. Very common resident in the Rogue Valley.	<b>Present.</b> This species was detected in the Study Area in the oak woodland area during the breeding season (May) and may nest within the Study Area or nearby.
American peregrine falcon <i>Falco peregrinus</i>	BCC (breeding) SV, CS 4	Inhabits a variety of open habitats. Nests on cliff ledges, or buildings or bridges, usually near water. Breeding season is approximately March through August.	<b>Present.</b> This species was observed flying through the Study Area during the breeding season (early May); there is no suitable nesting habitat within the Study Area, but this species may nest on nearby cliffs.
little willow flycatcher <i>Empidonax trailii brewsteri</i>	- SV, CS 4	Breeds in willows and other riparian vegetation along stream courses, lakes and marshes, also in thickets at edges of forest clearings or fields in proximity to water. Breeding season is typically June to August.	<b>Low.</b> This species is a fairly common migrant but a rare and irregular breeder in Jackson County. The habitat in the Study Area is marginal for this species and it is unlikely to occur.
loggerhead shrike <i>Lanius ludovicianus</i>	BCC (year- round) CS 4	This species occurs in open habitats with shrubs and trees for perching and nesting.	<b>Low.</b> The Study Area contains suitable habitat for this species; however this species is not known to breed in Jackson County and considered a rare and irregular visitor during the non-breeding season.



Common Name <i>Scientific Name</i>	Status <sup>1</sup> USFWS STATE ORBIC	Habitat and Ecology	Likelihood to Occur in the Study Area
streaked horned lark <i>Eremophila alpestris strigata</i>	T SC, CS 1	Nest where there is little or no vegetation including sparsely vegetated agricultural areas, pastures, grasslands, shrublands, playa margins, and alpine areas. Wintering habitats used are very similar to breeding habitats. Breeding season is March to August. The streaked horned lark has been extirpated throughout much of its range, including the Rogue Valley. Although this subspecies was known as a common permanent resident of the Rogue Valley in the early 1900s, the last confirmed breeding record was in 1976. Horned larks are currently only expected as rare and irregular visitors to Jackson County during the non-breeding season (September to February); however, these birds could be any of the four subspecies which frequently form mixed flocks in winter (only <i>E. a. strigata</i> breeds west of the Cascades). A flock of wintering streaked horned larks was confirmed in the Rogue Valley in winter 2015-2016 (USFWS 2016).	<b>Low.</b> There is suitable habitat in the Study Area for this subspecies which used to be a permanent resident of the Rogue Valley, but is currently considered to be extirpated. There is some likelihood that this subspecies could occur in the Study Area in the winter.
purple martin <i>Progne martin</i>	SOC SC, CS 2	Forage in open areas on the wing. Nest in cavities, often using woodpecker nest holes or nest boxes. Breeding season habitat typically open areas (open forest, open water, large meadows, fire scars in forests, or open areas near cities and towns) near to nest cavities (in trees, nest boxes, or crevices in cliffs or buildings).	<b>Moderate.</b> The Study Area has suitable foraging habitat adjacent to limited suitable nesting habitat for this species which is known to breed near the Study Area (breeding birds observed 2015-2017 off Valley View Road approximately two miles to the northwest [eBird 2017]). There is a record from July 26, 2013 of a juvenile hawking insects along an irrigation ditch off Butler Creek Road (mapped location is approximately 0.5 mile west of the northwest corner of the Study Area) (eBird 2017).
oak titmouse <i>Baeolophus inornatus</i>	BCC (year-round) - -	Common resident of the Rogue Valley in oak, mixed oak-pine, and oak-riparian woodlands and in mature chaparral communities. Nest in cavities, usually abandoned woodpecker holes or digs its own nest in soft wood (less common).	<b>Present.</b> This species was detected in the Study Area in the oak woodland area during the breeding season (May) and may nest within the Study Area.

Common Name <i>Scientific Name</i>	Status <sup>1</sup> USFWS STATE ORBIC	Habitat and Ecology	Likelihood to Occur in the Study Area
slender-billed nuthatch <i>Sitta carolinensis aculeata</i>	- SV, CS 3	This subspecies of white-breasted nuthatch generally inhabits the wooded slopes of the major interior valleys west of the Cascades and is a common resident of the Rogue Valley. Inhabits lower elevation deciduous, mixed conifer-deciduous, oak, ponderosa pine, and juniper woodlands. Nests in natural tree cavities or abandoned woodpecker holes.	<b>Present.</b> This species was detected in the Study Area in the oak woodland area during the breeding season (May) and may nest within the Study Area.
golden-crowned kinglet <i>Regulus satrapa</i>	- - 4	Common breeding bird in coniferous forest habitats in Jackson County; often come down in elevation in winter.	<b>High.</b> The Study Area is not suitable breeding habitat for this species, however they are common in the region and could be encountered in the Study Area anytime outside of nesting season.
western bluebird <i>Sialia mexicana</i>	- SV, CS 4	Common nesting bird in Jackson County; occupy a variety of habitats including farms, parks, open woodlands (riparian, oak, and oak-ponderosa pine); require cavities for nesting and typically use abandoned woodpecker holes, natural cavities, or nest boxes.	<b>Present (assumed).</b> There are multiple eBird records of the species in the Study Area. One record is during the breeding season (May 21) and this species would be likely to nest in the Study Area where suitable nesting cavities exist (eBird 2017).
chipping sparrow <i>Spizella passerina</i>	- CS 4	Common breeding bird in Jackson County. Typically found in open woodlands, savannahs, and openings in forests. Most birds have arrived by mid-April and depart by September; rare or irregular in the non-breeding season, though some birds are likely resident.	<b>Present.</b> This species was detected at the eastern edge of the Study Area near the end of North Mountain Avenue in early May; the Study Area may contain some marginally suitable nesting habitat.
Oregon vesper sparrow <i>Pooecetes gramineus affinis</i>	SOC SC, CS 2	This subspecies of vesper sparrow breeds west of the Cascades in Oregon. In Jackson County, it is an uncommon to fairly common summer resident in mountain grasslands; typically arriving in April and departing between July and October. A range-wide inventory and habitat assessment conducted in 2015 found birds to be notably absent from lower elevation grasslands and pasturelands in the Rogue Valley (where they are common in the Umpqua Valley); all detections in the Rogue Basin were above 2,000 feet and were primarily in montane meadows (Altman 2015).	<b>Low.</b> Although the Study Area is suitable habitat for this species, it appears to prefer higher elevation grasslands in the region and has a only low likelihood to occur.
grasshopper sparrow <i>Ammodramus savannarum</i>	-- SV, CS 2	Generally inhabit short to mid-height, open to moderately open grasslands, sometimes with scattered shrubs, and prefer large tracts of habitat to small	<b>Present.</b> This species was thoroughly documented in the Study Area in a study completed by the Klamath Bird Observatory

Common Name <i>Scientific Name</i>	Status <sup>1</sup> USFWS STATE ORBIC	Habitat and Ecology	Likelihood to Occur in the Study Area
		ones. Many pairs often nest close to each other because of limited suitable habitat; territory sizes between one and four acres have been documented. Birds nest on the ground below a dome of grass; frequently have two broods; nesting typically occurs from April to August. A summer resident of limited distribution in Jackson County.	(KBO) in 2016 which found 32 singing males in the Study Area (Stephens 2016). The grasshopper sparrow survey conducted by Pacific Crest Consulting in 2017 also documented 32 singing males in roughly the same locations (Figure 8). An informal survey conducted by KBO in 2014 found 20 grasshopper sparrows mostly in the same area (Stephens 2016).
yellow-breasted chat <i>Icteria virens</i>	SOC CS 4	A summer resident found in brush and thickets in open areas and understory of riparian woodlands along streams. They typically arrive in southern Oregon in May and depart in September. Nest in cups build in dense thickets.	<b>Present.</b> This species was observed in the Study Area along the small drainage in the center of the property as well as the TID in early May and is likely to nest in suitable habitat within the Study Area and nearby. Brush along the TID in the western portion of the Study Area (and likely along the entire length) was mowed this year sometime between mid-May and mid-June (C. Scott pers. observation), removing some of the suitable nesting habitat for chats in this area.
western meadowlark <i>Sturnella neglecta</i>	- CS 4	Very common summer resident in Jackson County and fairly common in winter as well when it may form small flocks. Inhabits open grasslands, pastures, some agricultural fields, meadows, and sometimes open woodlands. Nests are in depressions on the ground under domes of grass; territories sizes reported between several and 10+ acres.	<b>Present.</b> The most commonly encountered species during surveys of the Study Area conducted by Pacific Crest Consulting. Known to breed throughout most of the Study Area where suitable habitat is abundant, particularly above the TID.
tricolored blackbird <i>Agelaius tricolor</i>	BCC (breeding) -- 2	Uncommon to fairly common summer resident (rare but regular year-round resident); though typically resident in most of their range, most birds migrate to Oregon to breed. Prefer freshwater marshes with emergent vegetation or thickets for nesting; often nest in Himalayan blackberry shrubs around wetlands. They breed in colonies, often alongside red-winged blackbirds (may fly as far as four miles from nesting site to forage).	<b>Present.</b> Observed on one occasion (April 29) on the west side of the Study Area below the TID. The latest being from May 1, 2017 where five tricolored blackbirds were observed in the Study Area (eBird 2017). There is suitable nesting habitat in the southern portion of the Study Area for this species and it may breed there or nearby.
<b>Mammals</b>			

Common Name <i>Scientific Name</i>	Status <sup>1</sup> USFWS STATE ORBIC	Habitat and Ecology	Likelihood to Occur in the Study Area
Townsend's big-eared bat <i>Corynorhinus townsendii</i>	SOC SC, CS 2	Known to occur in many habitats but typically inhabits forested regions west of the Cascades. Uses caves, abandoned mines, buildings, and tunnels as roosts.	<b>Low.</b> There is no roosting habitat in the Study Area and this species is not anticipated to forage over the site.
pallid bat <i>Antrozous pallidus</i>	SOC SV, CS 2	This species inhabits a variety of habitats, typically shrublands and woodlands of arid regions but also open woodlands and forests (ponderosa pine, oak), preferably near water. They use narrow crevices in caves, mines, buildings and, less often, rock or debris piles and hollow trees for roosting; night roosts include abandoned buildings, rock overhangs, and bridges.	<b>Moderate.</b> This species may forage in the Study Area; the Study Area does contain some roosting habitat (hollow trees, rock piles), but lacks their more preferred sites (caves, bridges).
hoary bat <i>Lasiurus cinereus</i>	- SV, CS 4	Forest-dweller, day roosts in trees, resides in coniferous and deciduous forests and forages along riparian corridors and brushy areas.	<b>Low.</b> Suitable habitat exists in the limited oak woodland portion of the Study Area and this species may forage along Butler Creek.
California myotis <i>Myotis californicus</i>	- SV, CS 4	This bat typically forages over or near open water; it uses cliff faces, tree crevices, or caves for roosting. Seeks shelter after foraging during active season (does not use fixed roosts), and hibernates during winter in northwest.	<b>Moderate.</b> Suitable foraging and roosting habitat in the Study Area.
long-eared myotis <i>Myotis evotis</i>	SOC - 4	Generally associated with forested habitats or forest edges west of the Cascades; forages in openings in dense forest, between trees in open forest, and over willow-lined streams; roosts in wide variety of refugia including buildings, caves, mines, bridges, hollow trees, loose bark, and rock crevices.	<b>Low.</b> Suitable habitat exists near to the Study Area and this species may forage along Butler Creek. Not expected to roost in the Study Area
little brown myotis <i>Myotis lucifugus</i>	- - 4	Closely associated with water; found in moist forests or riparian woodlands. Commonly roost in structures and maternity colonies often located in structures, caves, or hollow trees; they hibernate in caves.	<b>Low.</b> Limited suitable habitat in the Study Area.
fringed myotis <i>Myotis thysanodes</i>	SOC SV, CS 2	Found in a wide variety of habitats but seems to have a presence for forests or riparian areas; roosts in caves, mines, buildings.	<b>Low.</b> Limited suitable habitat in the Study Area.
long-legged myotis <i>Myotis volans</i>	SOC SV, CS 4	Typically occurs in forests, but also in some desert and riparian habitats. Uses buildings, hollow trees and crevices in rock outcrops for maternity roosts. Uses caves and mines for winter roosts.	<b>Low.</b> Suitable habitat exists in the limited oak woodland portion of the Study Area.

Common Name <i>Scientific Name</i>	Status <sup>1</sup> USFWS STATE ORBIC	Habitat and Ecology	Likelihood to Occur in the Study Area
Yuma myotis <i>Myotis yumanensis</i>	SOC - 4	Highly associated with water; typically forages over open water such as rivers, lakes, ponds. Establish large colonies in buildings, mines, caves and bridges and also take solitary refuge in buildings, tree bark/crevices. In western Oregon, mostly found in Douglas-fir forests, Sitka spruce forest and oak and ponderosa pine woodlands.	<b>Low.</b> Limited suitable habitat in the Study Area.
Brazilian free-tailed bat <i>Tadarida brasiliensis</i>	- - 4	Colonial species that appears to be a permanent resident in Oregon; roosts frequently include caves, hollow trees, and buildings; colonies can be very large. Noted to be common to the Ashland area (Verts and Carraway 1998).	<b>Moderate.</b> May forage in the Study Area; less likely to roost in the Study Area due to lack of large roosting areas preferred by this species (caves, barns), but could roost in the limited oak woodland habitat or in structures nearby.
western gray squirrel <i>Sciurus griseus</i>	- CS 4	Generally inhabits oak woodlands, also mixed forests with hardwoods and conifers, as well as riparian areas and urban parks and orchards adjacent to natural habitats.	<b>Moderate.</b> Suitable habitat exists in the limited oak woodland portion of the Study Area; this species is locally common.
black-tailed jackrabbit <i>Lepus californicus</i>	- - 4	This species is found in open habitats including grasslands, shrubland, pastures, fields, and edges of forests.	<b>Present.</b> This species was observed in a field adjacent to the Study Area and there is suitable habitat throughout the Study Area for this species.
gray wolf <i>Canis lupus</i>	E CS 2	Occur over a wide-variety of habitats, though closely associated with dense coniferous forests west of the Cascades. Wolf territories ranging in size from 25 square miles to more than 1,000 square miles have been reported.	<b>Low.</b> Evidence of gray wolves has been documented as near as approximately 15 miles east of the Study Area (Keno Unit) near Howard Prairie as recently as 2017; gray wolves could hunt in the Study Area but would not be expected to reside there (ODFW 2017c).

<sup>1</sup> Status Code Definitions:**USFWS and STATE:****E:** Endangered**T:** Threatened**C:** Candidate**SOC:** Species of Concern**BCC:** Bird of Conservation Concern**SC:** Sensitive - Critical. SC species are imperiled with extirpation from a specific geographic area of Oregon because of small population sizes, habitat loss or degradation, and/or immediate threats.**SV:** ODFW Sensitive - Vulnerable. SV species are facing one or more threats to their populations and/or habitats.**CS:** ODFW Oregon Conservation Strategy (CS) Species**ORBIC:****List 1:** Taxa that are threatened with extinction or presumed to be extinct throughout their entire range.**List 2:** Taxa that are threatened with extirpation or presumed to be extirpated from the state of Oregon.**List 3:** Taxa for which more information is needed before status can be determined, but which may be threatened or endangered in Oregon or throughout their range.**List 4:** Taxa which are of conservation concern but are not currently threatened or endangered.

Table 3: ODA list of noxious weeds

Common Name	Scientific Name	ODA List
Velvetleaf	<i>Abutilon theophrasti</i>	B
Biddy-biddy	<i>Acaena novae-zelandiae</i>	B
Russian* knapweed	<i>Acroptilon repens</i>	B
Jointed goatgrass	<i>Aegilops cylindrica</i>	B
Ovate goatgrass	<i>Aegilops ovata</i>	A
Barbed goatgrass	<i>Aegilops triuncialis</i>	A, T
Quackgrass	<i>Elymus repens (Agropyron r.)</i>	B
Tree of Heaven	<i>Ailanthus altissima</i>	B
Camelthorn	<i>Alhagi maurorum (A. pseudalhagi)</i>	A
Garlic mustard	<i>Alliaria petiolata</i>	B, T
Yellowtuft	<i>Alyssum murale, A. corsicum</i>	A, T
Ragweed	<i>Ambrosia artemisiifolia</i>	B
Skeletonleaf bursage	<i>Ambrosia tomentosa</i>	A
Indigo bush	<i>Amorpha fruticosa</i>	B
Common bugloss	<i>Anchusa officinalis</i>	B, T
Hoary alyssum	<i>Berteroa incana</i>	A, T
False brome	<i>Brachypodium sylvaticum</i>	B
White bryonia	<i>Bryonia alba</i>	A
Butterfly bush	<i>Buddleja davidii (B. variabilis)</i>	B
Flowering rush	<i>Butomus umbellatus</i>	A, T
Plumeless thistle	<i>Carduus acanthoides</i>	A, T
Smooth distaff thistle	<i>Carduus baeticus</i>	A
Wetted thistle	<i>Carduus crispus</i>	A, T
Musk* thistle	<i>Carduus nutans</i>	B
Italian* thistle	<i>Carduus pycnocephalus</i>	B
Slender-flowered* thistle	<i>Carduus tenuiflorus</i>	B
Smooth distaff thistle	<i>Carthamus lanatus ssp. creticus (C. baeticus)</i>	A
Woolly distaff thistle	<i>Carthamus lanatus</i>	A, T
Purple starthistle	<i>Centaurea calcitrapa</i>	A, T
Diffuse* knapweed	<i>Centaurea diffusa</i>	B
Iberian starthistle	<i>Centaurea iberica</i>	A, T
Meadow* knapweed	<i>Centaurea pratensis</i>	B
Yellow starthistle*	<i>Centaurea solstitialis</i>	B, T
Spotted* knapweed	<i>Centaurea stoebe (C. maculosa)</i>	B, T
Squarrose knapweed	<i>Centaurea virgata</i>	A, T
Rush skeletonweed*	<i>Chondrilla juncea</i>	B, T
Canada* Thistle	<i>Cirsium arvense</i>	B
Bull* thistle	<i>Cirsium vulgare</i>	B
Old man's beard	<i>Clematis vitalba</i>	B
Poison hemlock	<i>Conium maculatum</i>	B

Field bindweed*	<i>Convolvulus arvensis</i>	B
Jubata grass	<i>Cortaderia jubata</i>	B
Common crupina (bearded creeper)	<i>Crupina vulgaris</i>	B
Japanese dodder	<i>Cuscuta japonica</i>	A
Houndstongue	<i>Cynoglossum officinale</i>	B
Yellow nutsedge	<i>Cyperus esculentus</i>	B
Purple nutsedge	<i>Cyperus rotundus</i>	A
Scotch* broom	<i>Cytisus scoparius</i>	B
Portuguese broom	<i>Cytisus striatus</i>	B, T
Spurge laurel	<i>Daphne laureola</i>	B
Cape-ivy	<i>Delairea odorata</i>	A, T
Cutleaf teasel	<i>Dipsacus laciniatus</i>	B
Paterson's curse	<i>Echium plantagineum</i>	A, T
South American waterweed	<i>Egeria densa (Elodea)</i>	B
Giant horsetail	<i>Equisetum telmateia</i>	B
Spanish heath	<i>Erica lusitanica</i>	B
Leafy* spurge	<i>Euphorbia esula</i>	B
Myrtle spurge	<i>Euphorbia myrsinites</i>	B
Oblong spurge	<i>Euphorbia oblongata</i>	A
Japanese (fleece flower) knotweed	<i>Fallopia japonica (Polygonum c.)</i>	B, T
Himalayan knotweed	<i>Fallopia polystachyum (Polygonum p.)</i>	B, T
Giant knotweed	<i>Fallopia sachalinensis (Polygonum s.)</i>	B, T
Goatsrue	<i>Galega officinalis</i>	A
French* broom	<i>Genista monspessulana</i>	B
Herb Robert	<i>Geranium robertianum</i>	B, T
Shiny-leaf geranium	<i>Geranium lucidum</i>	B, T
Halogeton	<i>Halogeton glomeratus</i>	B
Ivy	<i>Hedera helix, H. hibernica</i>	B
Texas blueweed	<i>Helianthus ciliaris</i>	A
Giant hogweed	<i>Heracleum mantegazzianum</i>	A, T
Orange hawkweed	<i>Hieracium (Pilosella) aurantiacum</i>	A, T
Meadow hawkweed	<i>Hieracium (Pilosella) caespitosum</i>	B, T
Yellow hawkweed	<i>Hieracium (Pilosella) floribundum</i>	A, T
Mouse-ear hawkweed	<i>Hieracium (Pilosella) pilosella</i>	A
King-devil hawkweed	<i>Hieracium (Pilosella) piloselloides</i>	A
Meadow hawkweed	<i>Hieracium pratense</i>	A, T
Hydrilla	<i>Hydrilla verticillata</i>	A
Common frogbit	<i>Hydrocharis morsus-ranae</i>	A
St. Johnswort *	<i>Hypericum perforatum</i>	B
Policeman's helmet	<i>Impatiens glandulifera</i>	B
Yellow flag iris	<i>Iris pseudacorus</i>	B
Dyers woad	<i>Isatis tinctoria</i>	B
Kochia	<i>Kochia scoparia</i>	B

Yellow archangel	<i>Lamiastrum galeobdolon</i>	B
Perennial peavine	<i>Lathyrus latifolius</i>	B
Lens-podded whitetop	<i>Lepidium chalepensis (Cardaria)</i>	B
Whitetop (hoary cress)	<i>Lepidium draba (Cardaria)</i>	B
Perennial pepperweed	<i>Lepidium latifolium</i>	B
Hairy whitetop	<i>Lepidium pubescens (Cardaria)</i>	B
West Indian spongeplant	<i>Limnobiium laevigatum</i>	A
Dalmatian* toadflax	<i>Linaria dalmatica (L.genista)</i>	B
Yellow* toadflax	<i>Linaria vulgaris</i>	B
Garden yellow loosestrife	<i>Lysimachia vulgaris</i>	A, T
Purple loosestrife*	<i>Lythrum salicaria</i>	B, T
Spikeweed	<i>Memizonia pungens</i>	B
Parrots feather	<i>Myriophyllum aquaticum</i>	B
Eurasian watermilfoil	<i>Myriophyllum spicatum</i>	B
Matgrass	<i>Nardus stricta</i>	A
Yellow floating heart	<i>Nymphoides peltata</i>	A
Scotch thistle	<i>Onopordum acanthium</i>	B
Taurian thistle	<i>Onopordum tauricum</i>	A, T
Small broomrape	<i>Orobanche minor</i>	B
African rue	<i>Peganum harmala</i>	A
Common reed	<i>Phragmites australis ssp. australis</i>	B
Sulfur cinquefoil	<i>Potentilla recta</i>	B
Kudzu	<i>Pueraria lobata</i>	A, T
Lesser celandine	<i>Ranunculus ficaria</i>	B
Creeping yellow cress	<i>Rorippa sylvestris</i>	B
Himalayan blackberry	<i>Rubus armeniacus (R. procerus, R. discolor)</i>	B
Ravennagrass	<i>Saccharum ravennae</i>	A, T
Mediterranean sage*	<i>Salvia aethiopsis</i>	B
Tansy ragwort*	<i>Senecio jacobaea</i>	B, T
Milk* thistle	<i>Silybum marianum</i>	B
Silverleaf nightshade	<i>Solanum elaeagnifolium</i>	A
Buffalobur	<i>Solanum rostratum</i>	B
Johnsongrass	<i>Sorghum halepense</i>	B
Smooth cordgrass	<i>Spartina alterniflora</i>	A, T
Common cordgrass	<i>Spartina anglica</i>	A, T
Dense-flowered cordgrass	<i>Spartina densiflora</i>	A, T
Saltmeadow cordgrass	<i>Spartina patens</i>	A, T
Spanish broom	<i>Spartium junceum</i>	B
Swainsonpea (Austrian peaweed)	<i>Sphaerophysa salsula</i>	B
Water soldiers	<i>Stratiotes aloides</i>	A
Medusahead rye	<i>Taeniatherum (Elymus) caput-medusae</i>	B
Saltcedar*	<i>Tamarix ramosissima</i>	B
European water chestnut	<i>Trapa natans</i>	A



Puncturevine*	<i>Tribulus terrestris</i>	B
Coltsfoot	<i>Tussilago farfara</i>	A
Gorse*	<i>Ulex europaeus</i>	B
Spiny cocklebur	<i>Xanthium spinosum</i>	B
Syrian bean-caper	<i>Zygophyllum fabago</i>	A

\* Indicates weeds targeted for biocontrol

A-Listed Weed: A weed of known economic importance which occurs in the state in small enough infestations to make eradication or containment possible; or is not known to occur, but its presence in neighboring states make future occurrence in Oregon seem imminent. Recommended action: Infestations are subject to eradication or intensive control when and where found.

B-Listed Weed: A weed of economic importance which is regionally abundant, but which may have limited distribution in some counties. Recommended action: Limited to intensive control at the state, county or regional level as determined on a site specific, case-by-case basis. Where implementation of a fully integrated statewide management plan is not feasible, biological control (when available) shall be the primary control method.

T-Designated Weed: A designated group of weed species that are selected and will be the focus for prevention and control by the Noxious Weed Control Program. Action against these weeds will receive priority. T designated noxious weeds are determined by the Oregon State Weed Board and directs ODA to develop and implement a statewide management plan. T designated noxious weeds are species selected from either the A or B list.

Table 4: Avian inventory

Common Name	Scientific Name	Special Status <sup>1</sup>
		Fed/State/ORBIC
Acorn woodpecker	<i>Melanerpes formicivorus</i>	SOC/CS/List 4
American kestrel	<i>Falco sparverius</i>	--/--/--
American robin	<i>Turdus migratorius</i>	--/--/--
Barn swallow	<i>Hirundo rustica</i>	--/--/--
Bewick's wren	<i>Thryomanes bewickii</i>	--/--/--
Brewer's blackbird	<i>Euphagus cyanocephalus</i>	--/--/--
Brown-headed cowbird	<i>Molothrus ater</i>	--/--/--
Bullock's oriole	<i>Icterus bullockii</i>	--/--/--
California quail	<i>Callipepla californica</i>	--/--/--
California scrub-jay	<i>Aphelocoma californica</i>	--/--/--
California towhee	<i>Melospiza crissalis</i>	--/--/--
Canada goose	<i>Branta canadensis</i>	--/--/--
Cedar waxwing	<i>Bombycilla cedrorum</i>	--/--/--
Chipping sparrow	<i>Spizella passerina</i>	--/--/List 4
Common raven	<i>Corvus corax</i>	--/--/--
Common yellowthroat	<i>Geothlypis trichas</i>	--/--/--
Downy woodpecker	<i>Dryobates pubescens</i>	--/--/--
European starling	<i>Sturnus vulgaris</i>	--/--/--
<b>Grasshopper sparrow</b>	<b><i>Ammodramus savannarum</i></b>	--/SV,CS/List 2
Green-winged teal	<i>Anas crecca</i>	--/--/--
Hairy woodpecker	<i>Leuconotopicus villosus</i>	--/--/--
House finch	<i>Haemorhous mexicanus</i>	--/--/--

House sparrow	<i>Passer domesticus</i>	--/--/--
Killdeer	<i>Charadrius vociferus</i>	--/--/--
Lark sparrow	<i>Chondestes grammacus</i>	--/--/--
Lazuli bunting	<i>Passerina amoena</i>	--/--/--
Lesser goldfinch	<i>Spinus psaltria</i>	--/--/--
Mallard	<i>Anas platyrhynchos</i>	--/--/--
Mourning dove	<i>Zenaida macroura</i>	--/--/--
Northern flicker	<i>Colaptes auratus</i>	--/--/--
Northern harrier	<i>Circus cyaneus</i>	--/--/--
Northern rough-winged swallow	<i>Stelgidopteryx serripennis</i>	--/--/--
Oak titmouse	<i>Baeolophus inornatus</i>	BCC/--/--
Peregrine falcon	<i>Falco peregrinus</i>	BCC/SV/List 4
Red-tailed hawk	<i>Buteo jamaicensis</i>	--/--/--
Red-winged blackbird	<i>Agelaius phoeniceus</i>	--/--/--
Ring-necked pheasant	<i>Phasianus colchicus</i>	--/--/--
Rock pigeon	<i>Columba livida</i>	--/--/--
Savannah sparrow	<i>Passerculus sandwichensis</i>	--/--/--
Song sparrow	<i>Melospiza melodia</i>	--/--/--
Spotted towhee	<i>Pipilo maculatus</i>	--/--/--
Tree swallow	<i>Tachycineta bicolor</i>	--/--/--
Turkey vulture	<i>Cathartes aura</i>	--/--/--
Western kingbird	<i>Tyrannus verticalis</i>	--/--/--
Western meadowlark	<i>Sturnella neglecta</i>	--/--/List 4
Western wood-peewee	<i>Contopus sordidulus</i>	--/--/--
White-breasted nuthatch	<i>Sitta carolinensis</i>	--/--/--

White-tailed kite	<i>Elanus leucurus</i>	--/--/List 4
Wilson's warbler	<i>Cardellina pusilla</i>	--/--/--
Yellow-breasted chat	<i>Icteria virens</i>	SOC/CS/List 4
Yellow-rumped warbler	<i>Setophaga coronata</i>	--/--/--
Chipping sparrow	<i>Spizella passerina</i>	--/--/--
White-crowned sparrow	<i>Zonotrichia leucophrys</i>	--/--/--
Violet-green swallow	<i>Tachycineta thalassina</i>	--/--/--

<sup>1</sup>Status Code Definitions:

**FEDERAL:**

**SOC:** U.S. Fish and Wildlife Service (USFWS) Species of Concern

**BCC:** U.S. Fish and Wildlife Service (USFWS) Bird of Conservation Concern

ORBIC: Oregon Biodiversity Information Center: 1 = taxa that are threatened with extinction or presumed to be extinct throughout their entire range (1-X designating presumed extirpation from Oregon or extinction); 2 = taxa that are threatened with extirpation or presumed to be extirpated from the state of Oregon; these are often peripheral or disjunct species which are of concern (when considering species diversity within Oregon's borders, they can be very significant when protecting the genetic diversity of a taxon)—ORBIC regards extreme rarity as a significant threat and has included species which are very rare in Oregon on this list; 3 = taxa for which more information is needed before status can be determined, but which may be threatened or endangered in Oregon or throughout their range; 4 = taxa which are of conservation concern but are not currently threatened or endangered; this includes taxa which are very rare but are currently secure, as well as List 4 contains taxa which are declining in numbers or habitat but are still too common to be proposed as threatened or endangered. While these taxa may not currently need the same active management attention as threatened or endangered taxa, they do require continued monitoring.

Table 5: Vascular plant inventory

Species	Relative Abundance	Status
<i>Achillea millefolium</i>	frequent	
<i>Achnatherum lemmonii</i>	frequent	
<i>Achyrrachaena mollis</i>	common	
<i>Acmispon wrangelianus</i>	rare	
<i>Agoseris retrorsa</i>	infrequent	
<i>Allium acuminatum</i>	rare	
<i>Allium amplexans</i>	rare	
<i>Allium sp (leaves only)</i>	frequent	
<i>Alopecurus pretense</i>	partially dominant	
<i>Alyssum alyssoides</i>	infrequent	
<i>Amaranthus albus</i>	rare	
<i>Amelanchier alnifolia</i>	infrequent	
<i>Amsinckia menziesii</i>	frequent	
<i>Anthriscus caucalis</i>	infrequent	
<i>Apocynum androsaemifolium</i>	frequent	
<i>Apocynum cannabinum</i>	rare	
<i>Artemesia douglasii</i>	infrequent	
<i>Asclepias fasciculatum</i>	frequent	
<i>Asclepias speciosa</i>	infrequent	
<i>Athysanus pusillus</i>	rare	
<i>Avena fatua</i>	dominant	
<i>Avena sativa</i>	rare	
<i>Balsamorhiza deltoidea</i>	infrequent	
<i>Blepharipappus scaber</i>	rare	
<i>Brassica nigra</i>	infrequent	
<i>Brassica rapa</i>	partially dominant	
<i>Brodiaea coronaria</i>	frequent	

<i>Brodiaea elegans</i>	infrequent	
<i>Bromus carinatus</i>	infrequent	
<i>Bromus diandrus</i>	infrequent	
<i>Bromus hordeaceus</i>	dominant	
<i>Bromus japonicus</i>	dominant	
<i>Bromus tectorum</i>	dominant	
<i>Bromus vulgaris</i>	frequent	
<b><i>California macrophylla (Erodium macrophyllum)</i></b>		SS
<i>Calochortus tolmiei</i>	common	
<i>Calycadenia truncata</i>	rare	
<i>Calystegia occidentalis</i>	common	
<i>Camassia quamash</i>	rare	
<i>Campanula prenanthoides</i>	rare	
<i>Capsella bursa-pastoris</i>	frequent	
<i>Cardamine oligosperma</i>	rare	
<i>Carex densa</i>	frequent	
<i>Carex geyeri</i>	rare	
<i>Carex sp (leaves only)</i>	infrequent	
<i>Carex stipata</i>	rare	
<i>Centaurea solstitialis</i>	dominant	
<i>Cerastium dichotomum</i>	common	
<i>Cerastium glomeratum</i>	frequent	
<i>Cercocarpus betuloides</i>	rare	
<i>Chamaesyce serpyllifolia</i>	rare	
<i>Cichorium intybus</i>	infrequent	
<i>Cirsium arvense</i>	infrequent	NOX
<i>Cirsium cymosum</i>	rare	
<i>Cirsium vulgare</i>	infrequent	NOX
<i>Clarkia gracilis</i>	infrequent	

<i>Clarkia purpurea</i>	infrequent	
<i>Clarkia rhombifolia</i>	infrequent	
<i>Claytonia exigua</i>	rare	
<i>Claytonia parviflora</i>	infrequent	
<i>Claytonia perfoliata</i>	infrequent	
<i>Collinisa parviflora</i>	infrequent	
<i>Collinsia linearis</i>	infrequent	
<i>Collinsia sparsiflora</i>	infrequent	
<i>Collomia grandiflora</i>	rare	
<i>Collomia linearis</i>	infrequent	
<i>Conium maculatum</i>	infrequent	NOX
<i>Conyza canadensis</i>	rare	
<i>Crepis modocensis</i>	frequent	
<i>Crocidium multicaule</i>	infrequent	
<i>Cryptantha intermedia</i>	infrequent	
<i>Cynoglossum grande</i>	rare	
<i>Cynosurus echinatus</i>	frequent	
<i>Cyperus cf esculentus</i>	rare	NOX
<i>Dactylis glomerata</i>	infrequent	
<i>Daucus carota</i>	frequent	
<i>Daucus pusillus</i>	rare	
<i>Delphinium nuttallianum</i>	infrequent	
<i>Dichelostemma capitatum</i>	common	
<i>Dichelostemma congestum</i>	common	
<i>Dipsacus fullonum</i>	frequent	
<i>Dodecatheon hendersonii</i>	rare	
<i>Dowlingia yina</i>	rare	
<i>Draba verna</i>	infrequent	
<i>Echinochloa crus-galli</i>	rare	

<i>Eleocharis acicularis</i>	infrequent	
<i>Eleocharis macrostachya (or palustris?)</i>	infrequent	
<i>Elymus glaucus</i>	infrequent	
<i>Epilobium brachycarpum</i>	common	
<i>Epilobium ciliatum var watsonii</i>	frequent	
<i>Equisetum hyemale</i>	infrequent	
<i>Ericameria nauseosa</i>	rare	
<i>Eriogonum compositum</i>	rare	
<i>Eriophyllum lanatum</i>	frequent	
<i>Erodium cicutarium</i>	dominant	
<i>Erysimum capitatum</i>	rare	
<i>Erythronium hendersonii</i>	frequent	
<i>Eschscholzia californica</i>	rare	
<i>Eurphorbia crenulata</i>	rare	
<i>Festuca roemerii</i>	frequent	
<i>Fraxinus latifolia</i>	rare	
<i>Fritillaria affinis</i>	frequent	
<i>Fritillaria sp (leaves only)</i>	frequent	
<i>Galium aparine</i>	frequent	
<i>Galium bolanderi</i>	rare	
<i>Galium divaricatum</i>	infrequent	
<i>Galium parisiense</i>	dominant	
<i>Geranium dissectum</i>	dominant	
<i>Geranium molle</i>	dominant	
<i>Gilia capitata</i>	rare	
<i>Glyceria sp (leaves only; grazed)</i>	rare	
<i>Gnaphalium palustre</i>	rare	
<i>Hemizonia congesta</i>	rare	
<i>Hieracium albiflorum</i>	rare	



<i>Hieracium scouleri</i>	infrequent	
<i>Holcus lanatus</i>	infrequent	
<i>Hordeum murinum</i>	infrequent	
<i>Horkelia daucifolia</i>	frequent	
<i>Hypericum perforatum</i>	rare	NOX
<i>Juncus effusus</i>	common	
<i>Juncus ensifolius</i>	rare	
<i>Juncus patens</i>	rare	
<i>Koeleria macrantha</i>	infrequent	
<i>Lactuca serriola</i>	frequent	
<i>Lagophylla ramossissima</i>	rare	
<i>Lamium amplexicaule</i>	infrequent	
<i>Lathyrus aphaca</i>	infrequent	
<i>Lathyrus cicera</i>	rare	
<i>Lemna minor</i>	rare	
<i>Lepidium campestre</i>	uncommon	
<i>Leptosiphon bilcolor</i>	rare	
<i>Linum bienne</i>	common	
<i>Lithophragma parviflorum</i>	infrequent	
<i>Lolium perenne</i>	infrequent	
<i>Lomatium californicum</i>	rare	
<i>Lomatium macrocarpum</i>	frequent	
<i>Lomatium nudicaule</i>	infrequent	
<i>Lomatium triternatum</i>	infrequent	
<i>Lomatium utriculatum</i>	dominant	
<i>Lonicera hispidula</i>	infrequent	
<i>Lonicera interrupta</i>	infrequent	
<i>Lotus corniculatus</i>	infrequent	
<i>Lotus micranthus</i>	rare	

<i>Lotus nevadensis</i>	infrequent	
<i>Lupinus albilfrons</i>	infrequent	
<i>Lupinus bicolor</i>	rare	
<i>Lupinus cf microcarpus</i>	rare	
<i>Lupinus latifolius</i>	frequent	
<i>Madia citriodora</i>	frequent	
<i>Madia elegans ssp densiflora</i>	infrequent	
<i>Madia elegans ssp vernalis</i>	frequent	
<i>Madia exigua</i>	infrequent	
<i>Madia gracilis</i>	common	
<i>Madia sativa</i>	rare	
<i>Mahonia aquifolium</i>	rare	
<i>Maianthemum stellatum</i>	rare	
<i>Malus fusca</i>	infrequent	
<i>Malus pumila</i>	rare	
<i>Medicago polymorpha</i>	rare	
<i>Medicago sp (leaves only; perhaps M. sativa)</i>	rare	
<i>Melilotus albus</i>	infrequent	
<i>Micropus californicus</i>	infrequent	
<i>Microseris laciniata ssp detlingii</i>	infrequent	
<i>Mimulus guttatus</i>	infrequent	
<i>Montia linearis</i>	rare	
<i>Myosotis discolor</i>	rare	
<i>Myosotis laxa</i>	rare	
<i>Nemophila parviflora</i>	rare	
<i>Olysnium douglasii</i>	infrequent	
<i>Orobanche uniflora</i>	rare	
<i>Osmorhiza berteroi</i>	rare	
<i>Penstemon sp (leaves only)</i>	rare	


<i>Phacelia hastata</i>	infrequent	
<i>Phacelia ramosissima</i>	rare	
<i>Phalaris arundinacea</i>	infrequent	NOX
<i>Phlox gracilis</i>	infrequent	
<i>Phlox speciosa</i>	frequent	
<i>Phoradendron villosum</i>	frequent	
<i>Piperia sp (leaves only)</i>	rare	
<i>Plagiobothrys tenellus</i>	infrequent	
<i>Plantago lanceolata</i>	frequent	
<i>Plectritis congesta</i>	common	
<i>Plectritis macrocera</i>	infrequent	
<i>Poa bulbosa</i>	dominant	
<i>Poa pratensis</i>	infrequent	
<i>Polygonum douglasii</i>	frequent	
<i>Polypogon monspeliensis</i>	rare	
<i>Populus balsamifera ssp trichocarpa</i>	rare	
<i>Portulaca oleracea</i>	infrequent	
<i>Prunus avium</i>	infrequent	
<i>Prunus cerasifolia</i>	infrequent	
<i>Prunus subcordata</i>	common	
<i>Pseudoroegneria spicata</i>	frequent	
<i>Quercus garryana ssp breweri</i>	rare	
<i>Quercus garryana ssp garryana</i>	common	
<i>Quercus kelloggii</i>	infrequent	
<b><i>Ranunculus austro-oreganus</i></b>	frequent	SS
<i>Ranunculus orthrhynchus</i>	infrequent	
<i>Ranunculus parviflorus</i>	frequent	
<i>Ranunculus sp (leaves only)</i>	frequent	
<i>Ranunculus uncinatus</i>	frequent	

<i>Ribes inerme</i> var. <i>klamathense</i>	rare	
<i>Rosa canina</i>	common	
<i>Rosa eglantina</i>	infrequent	
<i>Rosa gymnocarpa</i>	rare	
<i>Rubus armenicus</i>	common	NOX
<i>Rubus laciniatus</i>	rare	NOX
<i>Rubus ursinus</i>	infrequent	
<i>Rumex acetosella</i>	infrequent	
<i>Rumex crispus</i>	frequent	
<i>Salix cf lucida</i>	rare	
<i>Salix exigua</i>	rare	
<i>Salix lasiandra</i> var. <i>lasiandra</i>	frequent	
<i>Salix scouleriana</i>	rare	
<i>Sambucus cerulea</i>	rare	
<i>Sanguisorba minor</i>	infrequent	
<i>Sanicula crassicaulis</i>	infrequent	
<i>Scandix penctin-veneris</i>	infrequent	
<i>Selaginella wallacei</i>	rare	
<i>Senecio integerrimus</i>	rare	
<i>Shedonorus arundinaceus</i>	partially dominant	
<i>Silybum marianum</i>	rare	NOX
<i>Solanum dulcamara</i>	rare	
<i>Sonchus asper</i>	infrequent	
<i>Spartium junceum</i>	rare	NOX
<i>Stachys ajugoides</i>	infrequent	
<i>Symphoricarpos albus</i>	infrequent	
<i>Symphoricarpos mollis</i>	frequent	
<i>Taeniatherum caput-medusae</i>	dominant	NOX
<i>Taraxicum officinale</i>	infrequent	

<i>Thinopyrum ponticum</i>	common (highly concentrated in one large area)	
<i>Torilis arvensis</i>	frequent	
<i>Toxicodendron diversiloba</i>	common	
<i>Tragopogon dubius</i>	dominant	
<i>Trifolium albopurpureum</i>	frequent	
<i>Trifolium dubium</i>	infrequent	
<i>Trifolium hirtum</i>	common	
<i>Trifolium subterraneum</i>	rare	
<i>Typha latifolia</i>	rare	
<i>Valerianella locusta</i>	frequent	
<i>Verbascum blatteria</i>	rare	
<i>Veronica americana</i>	infrequent	
<i>Vicia americana</i>	frequent	
<i>Vicia sativa</i>	dominant	
<i>Vicia villosa</i>	dominant	
<i>Vulpia bromoides</i>	infrequent	
<i>Vulpia microstachys</i>	dominant	
<i>Wyethia angustifolia</i>	infrequent	
<i>Yabea microcarpa</i>	rare	
<i>Zigadensus venenosus var venenosus</i>	infrequent	

# Figures

Figure 1: Study Area



**Figure 2: ORBIC Rare Plant Form**

**OREGON BIODIVERSITY INFORMATION CENTER  
RARE PLANT FIELD SURVEY FORM**

Please complete all entries in the top section above the heavy line. Please complete as much as possible the more detailed section below the heavy line. You may use the back for comments or additional space. If possible, please attach a map of the location, preferably something of the same quality as a USGS 7.5' map.

Scientific Name: \_\_\_\_\_

Date of Field Work: \_\_\_\_\_ County: \_\_\_\_\_ Collection:  Yes (\_\_\_\_\_),  No  
mo. day year coll #, herbarium

Directions: \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_

Reporter: \_\_\_\_\_ Phone: \_\_\_\_\_

Address: \_\_\_\_\_

**1. LOCATION** - Attach separate map or sketch a map indicating exact site, scale and proximity to prominent features.

- A. Plant found?  Yes  No If no, reason: \_\_\_\_\_
- B. Location: T \_\_\_\_\_ R \_\_\_\_\_ Sec \_\_\_\_\_ 1/4 of \_\_\_\_\_ 1/4 (use back for more TRS)
- C. Source of GPS coordinates (fill one): GPS (make & model \_\_\_\_\_) or map (type & scale \_\_\_\_\_)  
 GPS accuracy distance: \_\_\_\_\_  Feet or  Meters
- Datum:  NAD27  NAD83  Other: \_\_\_\_\_. Easting/Longitude \_\_\_\_\_ Northing/Latitude \_\_\_\_\_
- D. Owner/Manager: \_\_\_\_\_

**2. SPECIES BIOLOGY**

- A. Phenology: \_\_\_\_\_% in flower, \_\_\_\_\_% in fruit, \_\_\_\_\_% in leaf
- B. Population size: Number of plants: \_\_\_\_\_ Area occupied: \_\_\_\_\_
- C. Age Class: \_\_\_\_\_% seedlings, \_\_\_\_\_% immature, \_\_\_\_\_% 1<sup>st</sup> year, \_\_\_\_\_% mature, \_\_\_\_\_% senescent

**3. HABITAT**

- A. Plant communities/Habitat Description/Associated Species: \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_
- B. Aspect: \_\_\_\_\_ (enter compass direction(s) or degrees)
- C. Slope:  Slight (0°-20°)  Moderate (20°-45°)  Extreme (45°+)  Vertical
- D. Topographic position:  Crest  Upper slope  Mid-slope  Lower slope  Bottom
- E. Light:  Open  Filtered  Shaded
- F. Moisture:  Inundated  Saturated  Moist  Dry
- G. Elevation range: \_\_\_\_\_ to \_\_\_\_\_  Feet or  Meters
- H. Substrate/soil: \_\_\_\_\_
- I. Visible threats/potential disturbance: \_\_\_\_\_

**4. DETERMINATION** How was plant identified? (choose one or more; please note the source for each choice)

- Keyed in flora  Compared with specimen  Compared with photo/drawing  Identified by someone else  Other
- Sources: \_\_\_\_\_

**5. PHOTOGRAPHS/SLIDES**

Did you take a photo?  Yes (  Film  Digital )  No If yes, may we obtain duplicates at our cost?  Yes  No

ORBIC-INR / Portland State University / Mail Stop INR / P.O. Box 751 / Portland, Oregon 97201-0751 / 503-725-9950 ph, -9960 fax

Figure 3: *California macrophylla* site locations







Figure 4: *California macrophylla* specimens




Plants, with fruit (lower right inset) and flower (bottom center inset)

Figure 5: *Ranunculus austro-oreganus* and *Collema quadrifidum* site locations



**Figure 6:** *Ranunculus austro-oreganus* and *Collema quadrifidum* specimens



*Ranunculus austro-oreganus* flower (diagnostic petal backs), with *Collema quadrifidum* thalli (upper right inset) and *C. quadrifidum* spore (lower right inset)

Figure 7: Noxious weed locations other than *Centaurea solstitialis* and *Elymus caput-medusae*





Figure 8: Grasshopper sparrow detections




**Legend**

- Imperatrice
- PLS
- Taxlots 2010
- City Limits
- Highways

GRSP detection



**Figure 9: Species of *Phaeocalicium* from the Study Area (micrograph)**



**Figure 10: Petrified log**



## Appendix A: Representative Photos of the Property



Early season view north towards Grizzly Peak (background) from downslope of TID canal



Early season view of the City of Ashland from the Property with Mt Ashland and Ashland Watershed in background





Early season view north towards Grizzly Peak (background) from upslope of TID canal



Early season view of vegetation downslope of TID canal, looking south towards City of Ashland



Oak woodlands at the north end of the Property



Herd of elk, with Bald Mt and Anderson Butte vicinities in far background. (See also photo on front cover page)



Early season view of vegetation upslope of TID canal (yellow flowers are the native *Lomatium utriculatum*)



Looking approximately southeast across the Property, from upslope of the TID canal



View across Property, with seasonal pond (see 3.1 Current Environment) in background; purple flowers in foreground are *Vicia villosa*



View of powerline and gas pipeline corridors on the Property; the pale strip from top to bottom, just left of center, with OHV tracks, is a *Thinopyrum ponticum* monoculture atop the buried gas pipeline (see 3.3 Noxious Weeds)



Pin flags delineating a *California macrophylla* population; background: controlled burning (smoke) from the Ashland Forest Resiliency project within the Ashland Watershed



Field of the native *Plectritis congesta* (shortspur seablush) in far northwest part of property, with Butler Creek in background



Cement “cistern” on broad ridgeline



The native *Calochortus tolmiei* (Tolmie’s startulip)



*Saguisorba minor* (salad burnet)



The native *Calystegia occidentalis* (field bindweed)





The native *Leptosiphon bicolor* (babystars)

*Note: hi-res versions of the above photos are available upon request*