
Common Name: Bullhead Catfish

Scientific Name: Ameiurus spp Size: 62 cm (2 ft)



Field Notes:

**Habitat:** Low oxygen, muddy areas in lakes and ponds

**Fact:** omnivorous bottom feeders: insects, leeches, snails, fish, clams, and many plants. Very tolerant of low oxygen and/or muddy conditions

#### **Identifying Characteristics:**

- non-forked, squared tail
- mottled, brown appearance
- chin barbels

Common Name: Largemouth Bass

**Scientific Name:** *Micropterus salmoides* 

**Size:** 75 cm (2.5 ft)



Field Notes:

**Habitat:** Warm upper waters of lakes, rivers or pools. Seeks logs, rock ledges, vegetation, and man-made structures for shelter.

**Fact:** Top predators-feeds on bluegill, shad, shiners and other minnow species, small catfish, sunfish, large invertebrates, and frogs. Hide among plants, roots or limbs to strike their prey. Will not spawn if the pH is less than 5 and eggs will not hatch if the pH is greater than 9.6

- usually green with dark blotches that form a horizontal stripe along the middle of the body
- underside ranges in color from light green to almost white
- upper jaw reaches far beyond the rear margin of the eye

Common Name: Bluegill

Scientific Name: Lepomis macrochirus

**Size:** 10-41 cm (1 ft)



Field Notes:

**Habitat:** vegetated lakes, ponds, swamps, creeks,

and rivers

Fact:


Lays its eggs in a nest, which is guarded by male fish. A single nest may contain eggs in various stages of development. Young fish feed on plankton, eventually diet shifts to macroinvertebrates. Up to 50% of their diet may consist of midge larvae

## **Identifying Characteristics:**

- dark spot on gill cover
- vertical bars on sides
- small mouth

Common Name: Channel Catfish

Scientific Name: Ictalurus punctatus
Size: up to 1.2 m (4 ft)



Field Notes:

**Habitat:** most abundant in large streams with low or moderate current

**Fact:** Eats organic detritus, aquatic insects, zooplankton and fishes. Spawn in late spring or early summer when water temperatures reach 75°F. One of the most preferred fish to catch in Texas.

- · tail deeply forked
- upper jaw longer than lower
- · barbels around the mouth
- scaleless
- young have body spots

Common Name: Rio Grande Cichlid

Scientific Name: Cichlasoma cyanoguttatum

**Size:** 152 mm (6 in)





**Habitat:** Pools and runs of small to large rivers. Will not survive below 49°F.


**Fact:** eats small fish, insects and crustaceans. Pollution-tolerant omnivorous species, indicator of an unbalanced or stressed ecosystem. May displace native Centrarchidae when aquatic conditions are less than optimal.

### **Identifying Characteristics:**

- color varies from very dark to light olive
- dorsal and anal fins are long and tapered extending behind the fleshy portion of the tail
- males develop a hump on their head
- looks speckled with cream and turquoise spots

Common Name: Mexican Tetra

Scientific Name: Astyanax mexicanus up to 100 mm (4 in)



Field Notes:

**Habitat:** wide range of freshwater habitats. Adults prefer rocky/sandy bottomed pools in creeks, streams and rivers

**Fact:** eats aquatic insects, crustaceans, worms, snails, smaller fish, plants and algae

- silver except for a black lateral band on the caudal fin
- forked tail
- · small dorsal fin
- comparatively large and strong teeth

Common Name: Greenthroat Darter

Scientific Name: Etheostoma lepidum up to 64 mm (2.5 in)

Photo by Chad Thomas, Texas State University - San Marcos

**Habitat:** riffles, non-turbid cool springs and vegetated pools, benthic after hatching

**Fact:** found in spring systems (headwaters and spring runs)

#### **Identifying Characteristics:**


- color ranges
- small and slender body with small rounded head
- females have yellow to white belly
- males have green/blue throat and orange belly

Field Notes:

Common Name: Spotted Gar

Scientific Name: Lepisosteus oculatus

**Size:** 1m (3.6 ft)



Field Notes:

**Habitat:** Mainly clear, quiet waters with aquatic vegetation

**Fact:** move slowly unless trying to catch food, which it grabs in its jaws in a quick sideways lunge. Prey is usually swallowed headfirst. Lifespan up to 18 yrs. Have a specialized swim bladder which allows them to gulp air and live in low oxygen conditions.

- long and cylindrical with long mouths
- back and upper sides olive with numerous dark oval spots on head, body, fins
- large canine teeth in one row on each side of upper jaw
- Young have a brown mid-lateral band

Common Name: Blackstripe Topminnow

Scientific Name: Fundulus notatus
7.4 cm (3 in)

Themas, Bonner, and Whiteside 2007

Fie	ld	Note	es:

**Habitat:** Prefers slow moving creeks

Fact: Eats mainly insects; spawns in plants; eggs

stick to submerged live or dead plants; fairly tolerant of high temperatures and low

oxygen levels

#### **Identifying Characteristics:**

- spots on body and fins
- body with distinct lateral band that extends from snout to tail
- back yellow to light olive-brown, with a few dark spots

Common Name: Mosquito Fish

Scientific Name: Gambusia affinis
Size: up to 5 cm (2 in)



Field Notes:

**Habitat:** Vegetated ponds, lakes, drainage ditches, and backwaters of sluggish creeks. Tends to swim near surface.


Fact: Will rise to water surface and breath thin film of air at very low oxygen levels; have internally fertilized eggs; feed on insect larvae (prefer mosquito larvae), invertebrates, algae and fish fry, including its own progeny; widely introduced for mosquito control, however they've reduced or eliminated native fish population.

- top of head dusky
- bluish black spot below eye
- back/olive color with scattered black spots
- abdomen silvery to whitish

Common Name: Central Stoneroller

**Scientific Name:** Campostoma anomalum

**Size:** up to 23 cm (9 in)



Field Notes:

**Habitat:** Riffles (gravel substrate), benthic, likes cool,

clear water

**Fact:** Eats algae, diatoms and detritus. Males

build nests in gravel riffles by "rolling" larger stones away; eggs adhesive and attached to stones in nest; males defend nests

#### **Identifying Characteristics:**

- snout blunt and rounded
- thick bodies
- olive\brown above, often dark stripe along side
- dark spot on back fin of young
- irregular dark blotches on back and side
- breeding males have orange fins, white lips, & bright red eye

Common Name: Red Shiner or Red-horse

**Minnow** 

**Scientific Name:** Cyprinella lutrensis

**Size:** up to 9 cm (3.5 in)



Field Notes:

**Habitat:** Lakes, rivers, creeks in pools and riffles especially backwaters and creek mouths with sand and silt substrates; ususally not present in clear water

**Fact:** Spawning may occur on riffles, on or near

submerged objects, over vegetation beds, or in association with sunfish nests. Adults typically school in midwater or near the surface. Eats insects and algae. Tolerant of siltation and frequent high turbidity

- olive green above, silver on sides
- spawning males become bluish on the sides and fins get red
- abdomen whitish

Common Name: Gray Redhorse "Sucker"

Scientific Name: Moxostoma congestum up to 70 cm (2 ft)

Fact. Fats along crayfish shalls small clams larval


**Habitat:** Rock, sand, or gravel bottomed pools and

**Fact:** Eats algae, crayfish, snails, small clams, larval insects

## **Identifying Characteristics:**

deep runs

- mouth like a vaccuum
- black grayish or yellowish brown to greenish
- sides silvery
- · abdomen whitish




Field Notes:

Common Name: **Green Sunfish** 

Scientific Name: Lepomis cyanellus

Size: up to 25 cm (10 in)



Field Notes:


**Habitat:** Tolerant of many habitats; common in ponds, lakes, and in areas of creek with little flow

**Fact:** Eats insects, mollusks, and small fishes. Eggs are laid in nests scooped out of gravel or sandy silt by the male who guards the nest.

- dark green, almost blue
- faint vertical bars are apparent on the sides
- some scales have turquoise spots
- dark spot at posterior base of dorsal fin

Common Name: Redbreast Sunfish

Scientific Name: Lepomis auritus
up to 24 cm (9 in)



Field Notes:

Habitat: Ponds, lakes and creeks


**Fact:** Adults feed on insects, snails, crayfish, and small fish. Nests are solitary, usually adjacent to logs or some other structure and are usually constructed over sand.

#### **Identifying Characteristics:**

- yellow, orange, or rusty color belly
- long opercle flap (ear) up to 1 in.
- typically a larger sunfish (up to 1 lb)

Common Name: Longear Sunfish

Scientific Name: Lepomis megalotis up to 20 cm (8 in)



Field Notes:

**Habitat:** Ponds, lakes, clear creeks

**Fact:** Feed at the surface of the water. Eat mainly aquatic insects, mites, microcrustaceans, fish eggs, mollusks, filamentous algae, and small fish.

- brown to olive green back with bright blue spots
- elongated opercle flap is always trimmed in white in adults
- colorful-males are often bright orange or scarlet
- the head and fins usually have turquoise markings