

Phenotypic plasticity triggers rapid morphological convergence

José M. Gómez^{1,2*}, Adela González-Megías^{2,3*}, Eduardo Narbona^{4*}, Luis Navarro^{5*}, Francisco Perfectti^{2,6*}, Cristina Armas^{1*}

¹Estación Experimental de Zonas Áridas (EEZA-CSIC), Almería, Spain.

²Research Unit Modeling Nature, Universidad de Granada, Granada, Spain.

³Dpto. de Zoología, Universidad de Granada, Granada, Spain.

⁴Dpto. de Biología Molecular e Ingeniería Bioquímica, Universidad Pablo de Olavide, Sevilla, Spain.

⁵Dpto. de Biología Vegetal y Ciencias del Suelo, Universidad de Vigo, Vigo, Spain.

⁶Dpto. de Genética, Universidad de Granada, Granada, Spain.

*Corresponding author. Email: jmgreyes@eeza.csic. (J.M.G.); adelagm@ugr.es (A.G.); enarfer@upo.es (E.N.); lnavarro@uvigo.es (L.N.); fperfect@ugr.es (F.P.); cris@eeza.csic.es (C.A.)

Abstract

Phenotypic convergence, the independent evolution of similar traits, is ubiquitous in nature, happening at all levels of biological organizations and in most kinds of living beings. Uncovering its mechanisms remains a fundamental goal in biology. Evolutionary theory considers that convergence emerges through independent genetic changes selected over long periods of time. We show in this study that convergence can also arise through phenotypic plasticity. We illustrate this idea by investigating how plasticity drives *Moricandia arvensis*, a mustard species displaying within-individual polyphenism in flowers, across the morphological space of the entire Brassicaceae family. By compiling the multidimensional floral phenotype, the phylogenetic relationships, and the pollination niche of over 3000 Brassicaceae species, we demonstrated that *Moricandia arvensis* exhibits a plastic-mediated within-individual floral disparity greater than that found not only between species but also between higher taxonomical levels such as genera and tribes. As a consequence of this divergence, *M. arvensis* moves outside the morphospace region occupied by its ancestors and close relatives, crosses into a new region where it encounters a different pollination niche and converges phenotypically with distant Brassicaceae lineages. Our study suggests that, by inducing phenotypes that explore simultaneously different regions of the morphological space, plasticity triggers rapid phenotypic convergence.

Introduction

Phenotypic convergence, the independent evolution of similar traits in different evolutionary lineages, is ubiquitous in nature, happening at all levels of biological organizations and in most kinds of living beings (1-3). Convergent evolution plays a fundamental role in how evolutionary lineages occupy the morphological space (2, 4). The expansion of lineages across the morphological space is a complex process resulting from the ecological opportunities emerging when species enter into different regions of the ecospace and face new ecological niches (5, 6). When this occurs, divergent selection on some phenotypes makes lineages to diversify phenotypically, boosting morphological disparity, triggering a morphological radiation and eventually filling the morphospace (7, 8). Because the ecological space saturate as lineages diversify (9), unoccupied regions become rare in highly diversified lineages (10). Under these circumstances, entering into a new region usually entails sharing it with other species exploiting the same ecological niche (2, 10, 11). In this situation, independent lineages tend to evolve similar phenotypes through convergent evolution (2, 4). In diversified lineages occupying a saturated morphospace, divergent and convergent evolution are ineludibly connected (10, 12), and both processes contribute significantly to shape the geometry of the morphospace occupation (4, 11).

Uncovering the mechanisms triggering convergence remains a fundamental goal in biology. Evolutionary theory shows that convergent phenotypes emerge from several genetic mechanisms, such as independent mutations or gene reuse in different populations or species, polymorphic alleles, parallel gene duplication, introgression or whole-genome duplications, that are selected over long periods of time (13–15). Under these circumstances, the origin of morphological convergence is mostly slow, occurring over evolutionary time and associated with multiple events of speciation and cladogenesis (11). It is increasingly acknowledged, however, that phenotypic plasticity might elicit the emergence of novel phenotypes with new adaptive possibilities, which may be beneficial in some contexts (16, 17). Under these circumstances, plasticity may behave as a facilitator for evolutionary novelty and diversity, shaping the patterns of morphospace occupation (16, 18-21). In this study, we provide compelling evidence showing that phenotypic plasticity also plays a prominent role in the emergence of convergent phenotypes. By inducing the production of several phenotypes, plasticity may cause the species to explore different regions of the morphospace almost simultaneously (18, 19). This opens the opportunity for plastic species to diverge from their lineages and converge with the species already located in other morphospace regions. We illustrate this idea by investigating how plasticity drives *Moricandia arvensis*, a species exhibiting extreme polyphenism in flowers (18), across the

morphological space of the entire Brassicaceae family. *Moricandia arvensis* displays within-individual floral plasticity, with flower morphs varying seasonally on the same individual (18). By studying the multidimensional floral phenotypes, the phylogenetic relationships, and the pollination niches of over 3000 Brassicaceae species, we demonstrate that phenotypic plasticity makes the flowers of this mustard species to diverge from its ancestors and close relatives, to cross into a new region of the ecospace, and to converge morphologically with distant Brassicaceae lineages. This finding has great implications, suggesting that plasticity might not only promote the evolution of novelties and morphological divergence (16, 17, 20, 21) but can also provide an alternative explanation to the pervasiveness of convergence in nature.

Results

Plasticity-mediated floral disparity and divergence

Changes in temperature, radiation and water availability induce the production of different types of flowers by the same *M. arvensis* individuals; large, cross-shaped lilac flowers in spring but small, rounded, white flowers in summer (18). To quantify the magnitude of floral disparity between these two phenotypes of *M. arvensis*, we first assessed floral disparity for the entire mustard family. Brassicaceae is one of the largest angiosperm families, with almost 4000 species grouped in 351 genera and 51 tribes (7, 22–24). We determined the magnitude and extent of floral disparity among 3140 plant species (approx. 80% of the accepted species) belonging to 330 genera (94% of the genera) from the 51 tribes. Because we were interested in floral characters mediating the interaction with pollinators, we recorded for each studied species a total of 31 traits associated with pollination in Brassicaceae (Supplementary Data 1, Methods). We used the resulting phenotypic matrix to generate a family-wide floral morphospace. We first run a principal coordinate analysis (PCoA) to obtain a low-dimensional Euclidean representation of the multidimensional phenotypic similarity existing among the Brassicaceae species (25). Because the raw matrix was composed of quantitative, semi-quantitative and discrete variables, PCoA was based on Gower dissimilarities (25). We optimized this initial Euclidean configuration by running a non-metric multidimensional scaling (NMDS) algorithm with 5000 random starts (25). The resulting morphospace (Figure 1a) was significantly correlated with the initial PCoA configuration ($r = 0.40$, $P < 0.0001$, Mantel test) and was a good representation of the original relationship among the species ($R^2 > 0.95$, $Stress = 0.2$, Figure 1b). The distribution of the species across the morphospace was significantly associated with different pollination traits (Figure S1; Table S1). Species in the central region were mostly medium-sized plants bearing a moderate to high number of small, polysymmetric white flowers with short corolla tubes, exposed nectaries and

visible sepals (Figure 1a, Figure S1). Species in the bottom right corner were small or prostrate, bearing minute flowers, many time apetalous and with just 2 or 4 stamens, whereas species located in the bottom left corner were medium-sized plants with asymmetric flowers arranged in corymbose inflorescences. Plants with yellow flowers were located in the right region of the morphospace. In contrast, large plants with strongly tetradynamous androceum and large, veined, dissymmetrical to asymmetrical, pink to blue flowers with concealed nectaries, long corolla tubes and bullseyes were located in the upper left region (Figure 1a, Figure S1). *Moricandia arvensis*, when blooming in spring (Figure 1c), occupies this later peripheral region of the morphospace, close to other *Moricandia* species (purple dots in Figure 1a). However, during summertime, the individuals of *M. arvensis* are shorter and produce fewer, much smaller flowers with white, unveined and rounded corollas with overlapped petals and green sepals that are mostly arranged along the floral stems (Figure 1d) (18). Due to this radical phenotypic change, the summer phenotype of *M. arvensis* was located in a different, more central position of the floral morphospace (Figure 1a), far away from the region occupied by the *Moricandia* species. As a consequence of this jump, the morphological disparity between the spring and summer phenotypes of *M. arvensis*, calculated as their distance in the morphospace (26), was very high (0.264). In fact, it was much higher than the average pairwise disparities among all studied Brassicaceae species (0.155 ± 0.090 , mean \pm s.e.m., 4,912,545 pairwise disparities) and almost 50% of the largest observed disparity (0.55) (Table S4). This outcome suggests that phenotypic plasticity prompts *M. arvensis* to explore two distant regions of the Brassicaceae floral morphospace simultaneously.

To know how intense is the plasticity-mediated *M. arvensis* disparity, we compared its value with the disparity values observed at different taxonomic levels within Brassicaceae. At the lowest level, discrete changes in pollination traits have been reported between individuals of the same species. In some species, this intraspecific phenotypic change is stable, like the gender polymorphism (27, 28) or the adaptive floral colour polymorphism exhibited as a response to the selective pressures exerted by certain pollinators (29, 30). In other species, discrete phenotypic changes, although affecting pollination traits, seem to be just the consequence of some singular and often unstable mutations affecting floral colour (31), the production of cleistogamous flowers (32) or changes in the expression of homeotic genes that modify the formation of the floral organs (33, 34). We compiled information on the phenotypes of the different morphs in 34 polymorphic species and calculated their values of intraspecific disparities (Figure 1a, Supplementary Data 2). Although several polymorphic species showed considerable values of between-morph disparity, they were significantly smaller than the disparity between spring and summer floral phenotypes of *M. arvensis* (Z-score = 5.06, $P < 0.0001$, Figure 1e, Table S2). We subsequently tested at what

taxonomic level of Brassicaceae the disparity was equivalent to the plasticity-mediated disparity observed in *M. arvensis*. For this, we calculated the floral disparity between pair of species belonging to the genus *Moricandia*, the same genus, the same tribe, and different tribes (Methods). The plasticity-mediated disparity of *M. arvensis* was significantly higher than the disparity existing between the *Moricandia* species (0.057 ± 0.033 , mean ± 1 s.e.m., Z-score = 6.27, $P < 0.0001$) and between the species belonging to the same genus (0.069 ± 0.055 , Z-score = 3.51, $P < 0.0002$). It was marginally different from the disparity existing between species of different genera but the same tribes (0.150 ± 0.085 , Z-score = 1.34, $P = 0.089$) and it was statistically similar to the disparity occurring between species belonging to different tribes (0.167 ± 0.087 , Z-score = 1.11, $P = 0.133$, Figure 1e). These findings suggest that phenotypic plasticity allows *M. arvensis* individuals to jump in the morphospace longer distances than those granted by some macroevolutionary processes.

We explored whether plasticity-mediated disparity may cause evolutionary divergence by calculating the disparity of *M. arvensis* spring and summer phenotypes to their phylogenetic ancestors. We retrieved 80 partial phylogenies from the literature and online repositories (Methods), and assembled them into a supertree comprising 1876 taxa with information on their floral phenotype. We then projected this supertree onto the morphospace to get a family-wide phylomorphospace. We did not find evidence of phylogenetic constraints on morphospace occupation since there was not significant phylogenetic signal for the position occupied by each species (Multivariate *Mantel* test=0.005, $P = 0.34$). The family-wide phylomorphospace was very tangled (Figure 2a), with 492,751 intersections among lineages, suggesting the presence of many events of floral divergence and convergence in the evolution of Brassicaceae pollination traits (11). To calculate the disparity of the *M. arvensis* floral phenotypes to their ancestor, because these analyses are sensitive to the tree topology and the inferred branch lengths (26), we used four independent, time-calibrated phylogenies that included this species (Methods). The results were consistent across phylogenies (Figure 2b,c; Tables S3). The spring phenotype did not significantly diverge neither from the most recent common ancestor (MRCA) of *Moricandia* (Z-score = 0.36, $P = 0.36$) nor from its direct ancestor (Z-score = -1.24, $P = 0.108$). In contrast, the summer phenotype of *M. arvensis* diverged significantly both from *Moricandia* MRCA (Z-score = 2.48, $P = 0.007$) and from its direct ancestor (Z-score = 1.77, $P = 0.038$). Hence, the summer phenotype explores a region of the floral morphospace located out of its phylogenetic clade range (Figure 2b). The ancestral disparity of the summer phenotype was even significantly higher than the ancestral disparity of most other Brassicaceae species (Figure 2c). These findings suggest that phenotypic plasticity causes the appearance of a novel phenotype that diverges radically from its ancestors.

Plastic shifts in pollination niches

Evolutionary divergence is mostly associated with the occupation of new ecological niches (2, 5). Shifts between pollination niches are an important factor driving diversification in angiosperms (35), including Brassicaceae (36, 37). We investigated whether the plasticity-mediated jump of *M. arvensis* across the floral morphospace implicated the exploration of new pollination niches. We compiled a comprehensive database comprising 456,031 visits done by over 800 animal species from 19 taxonomical orders, 276 families and 43 functional groups to 554 Brassicaceae species of 39 tribes (Methods, Supplementary Data 3). Afterwards, we identified the pollination niches of these Brassicaceae plants and determined the niche of each *M. arvensis* floral phenotype by means of bipartite modularity, a complex network tool that identifies the set of plants interacting with similar groups of pollinators (18). This analysis showed that the network was significantly modular ($Modularity = 0.385$, $P < 0.0001$) and identified eight different pollination niches associated with different groups of pollinators (Figure 3a) located in different regions of the morphospace (Figure 3b, $F = 44.4$, $P < 0.001$, $R^2 = 0.39$, Adonis test; Table S4).

Because different insects visited *M. arvensis* in spring and summer (Table S5), this plant species shifted between pollination niches seasonally (Figure 3b). During spring, *M. arvensis* belonged to a niche where most frequent pollinators were long-tongued bees, beeflies, and hawkmoths (pollination niche 5 in Figure 3a) (18). This pollination niche was also shared by the other *Moricandia* species (Figure 3c). In contrast, during summer *M. arvensis* belonged to a niche dominated by short-tongued bees (pollination niche 3 in Figure 3a). This niche shift was substantial. In fact, the overlap between the spring and summer pollinator niches of *M. arvensis* (Czekanowski overlap index = 0.35) was significantly lower than the overlap between congeneric species of Brassicaceae (0.57 ± 0.42 , Z-score = -0.51, $P = 0.003$). This shift even entailed the divergence from the ancestral niche of the *Moricandia* lineage (pollination niche 5 according to a stochastic character mapping inference, Figure 3c). The within-individual floral plasticity allows *M. arvensis* to exploit a pollination niche that differs markedly from that exploited by its closest relatives and that have largely diverged from the ancestral niche.

Plasticity-mediated floral convergence

A common consequence of adaptation to the same niche is convergent evolution (1, 2, 4). We explored the possibility of convergent evolution of *M. arvensis* with other Brassicaceae sharing either the spring niche (pollination niche 5) or the summer niche (pollination niche 3). We first checked for the occurrence of convergence among species belonging to these pollination niches. Because these analyses are extremely sensitive to the inferred branch lengths, we explored

morphological convergence using three time-calibrated large (> 150 spp) phylogenies that included *M. arvensis* (Methods). We tested for the occurrence of floral convergence between the species belonging to each of those two pollination niches using three methods: the angle formed by the phenotypic vectors connecting the position in the floral morphospace of each pair of species with that of their most recent common ancestor (38), the difference in phenotypic distances between convergent species and the maximum distances between all other lineages (39), and the phenotypic similarity of the allegedly convergent species penalized by their phylogenetic distance (*Wheatsheaf* index) (40). The three methods gave similar results, indicating that floral convergence was frequent among the species belonging to any of the two studied niches, irrespective of the method and the time-calibrated tree used (Table S6). These results show that, despite the rampant generalization observed in the pollination system of Brassicaceae, species interacting with similar pollinators converge phenotypically.

Once we determined the occurrence of convergence in these two pollination niches, we assessed whether plasticity caused the evolution of morphological convergence in *M. arvensis*. To do so, we first assessed the convergence region of *Moricandia*, the region that includes the lineages converging morphologically to the *Moricandia* lineage. We found that this region included most species of *Moricandia*, the spring phenotype of *M. arvensis*, and several clades belonging to disparate tribes that interact with pollination niche 5, but excluded the summer phenotype of *M. arvensis* (Figure 4, Table S7). Afterwards, we checked whether any of the two *M. arvensis* floral phenotypes entered the region of the phylomorphospace defined by their pollination niches. We used the C5 index, defined as the number of lineages that cross into the morphospace region of interest from outside³⁹. This index detected between two and six convergent events towards pollination niche 5 depending on the phylogeny used (blue arrows in Figure 4a-c), but none was associated with the spring phenotype of *M. arvensis*. In contrast, the C5 index consistently detected that the summer phenotype of *M. arvensis* has converged with the species belonging to the pollination niche 3 (red arrow in Figure 4d-f). Altogether, these analyses suggest that, whereas the spring phenotype did not show any evidence of convergence, the summer phenotype of *M. arvensis* has converged with other distant Brassicaceae exploiting the same pollination niche.

Conclusions

Convergent selection exerted by efficient pollinators causes the evolution of similar suites of floral traits in different plant species (41–44). Our study shows that plasticity can promote the rapid

convergent evolution of floral traits, providing an additional explanation about how pollination syndromes may evolve. Under this idea, changes in floral traits precede shifts in pollinators, as frequently observed in generalist systems (37, 45). This may explain why many pollination systems are evolutionarily labile, undergoing frequent shifts and evolve multiple times within the same lineages by diverse evolutionary pathways (35, 46).

Morphological convergence is universally acknowledged to be the result of several genetic mechanisms, such as independent mutations in different populations or species, polymorphic genes or introgression (13). We provide in this study compelling evidence suggesting that morphological convergence may also arise as a consequence of phenotypic plasticity. The role of plasticity as a mechanism favouring quick responses of organisms to novel and rapidly changing environments is already beyond doubt (17, 21, 47, 48). Its evolutionary consequences are more debated though (20, 21, 49, 50). The 'plasticity-led evolution' hypothesis states that selection acting on a plastic lineage may either boost its environmental sensitivity and trigger the origin of polyphenisms or alternatively may promote the loss of plasticity and the canalization of the new phenotype through genetic assimilation (21, 49). The related 'flexible stem' hypothesis of adaptive radiation suggests that when a plastic lineage repeatedly colonizes similar niches, the multiple phenotypes fixed by genetic assimilation could converge among them giving rise to a collection of phylogenetically related convergent morphs (16, 50, 51). Our comprehensive study complements these hypotheses by suggesting that plasticity-mediated convergence may even evolve without the existence of basal flexible lineages. Rather, it can occur when plasticity evolving in otherwise non-plastic lineages promotes the colonization of a niche previously occupied by unrelated species. Under these circumstances, contrary to what it is predicted by the previous hypotheses, plasticity-mediated convergence is not circumscribed to phylogenetic-related species arising from a common stem lineage. This overlooked role of phenotypic plasticity may contribute to explain the ubiquity of morphological convergence in nature.

Materials and Methods

Floral traits. We recorded from the literature 31 floral traits in 3140 Brassicaceae plant species belonging to 330 genera and 51 tribes (Supplementary Data 1). All these traits have been proven to be important for the interaction with pollinators (Table S8). These traits were: (1) Plant height; (2) Flower display size; (3) Inflorescence architecture; (4) Presence of apetalous flowers; (5) Number of symmetry axes of the corolla; (6) Orientation of dominant symmetry axis of the corolla; (7) Corolla with overlapped petals; (8) Corolla with multilobed petals; (9) Corolla with visible

sepals; (10) Petal length; (11) Sepal length; (12) Asymmetric petals; (13) Petal limb length; (14) Length of long stamens; (15) Length of short stamens; (16) Stamen dimorphism; (17) Tetrastaminal condition; (18) Visible anthers; (19) Exserted stamens; (20) Number of stamens; (21) Concealed nectaries; (22) Petal carotenoids; (23) Petal anthocyanins; (24) Presence of bullseyes; (25) Presence of veins in the petals; (26) Coloured sepals; (27) Relative attractiveness of petals versus sepals; (28) Petal hue; (29) Petal colour as b CIELAB; (30) Sepal hue; (31) Sepal colour as b CIELAB. A detailed definition and description of these traits and their states is provided in Key Resource Table 1, whereas the original references used to determine the states of each trait per plant species is provided in Supplementary Data 1.

Family-wide floral morphospace. Using the original multidimensional trait-species matrix, we built a floral morphospace. For this, we reduced the high-dimensional matrix of floral traits to a two-dimensional space using an ordination technique (25). Because the set of floral traits included in this study were quantitative, semi-quantitative and qualitative, we used ordination techniques based on dissimilarity values. For this, we first constructed a pairwise square distance matrix of length equal to the number of Brassicaceae species included in the analysis ($n = 3140$). We used the Gower distance, the number of mismatched traits over the number of shared traits. This dissimilarity index is preferable to the raw Euclidean distance when there are discrete and continuous traits co-occurring in the same dataset (52). We reduced the dimensionality of this phenotypic matrix by projecting it in a two-dimensional space. For this, to ensure an accurate description of the distribution of the species in the morphospace, we first run a principal coordinate analysis (PCoA), a technique providing a Euclidean representation of a set of objects whose relationship is measured by any dissimilarity index. We corrected for negative eigenvalues using the Cailliez procedure (25). Afterwards, we used this metric configuration as the initial configuration to run a non-metric multidimensional scaling (NMDS) algorithm (25), a method that will further optimise the sample distribution so as more variation in species composition is represented by fewer ordination axes. Unlike methods that attempt to maximise the variance or correspondence between objects in an ordination, NMDS attempts to represent, as closely as possible, the pairwise dissimilarity between objects in a low-dimensional space. NMDS is a rank-based approach, where the original distance data is substituted with ranks, preserving the ordering relationships among species (25). Objects that are ordinated closer to one another are likely to be more similar than those further apart (53). This method is more robust than distance-based methods when the original matrix includes variables of contrasting nature. However, NMDS is an iterative algorithm that can fail to find the optimal solution. We decreased the potential effect of falling in local optima by running the analysis with 5000 random starts and iterating each run 1×10^6 times (54). The NMDS was run using a

monotone regression minimizing the Kruskal's stress-1 (55, 56), and compared each solution using Procrustes analysis, retaining that with the lowest residual. Because many species did not share trait states, a condition complicating ordination, we used *stepacross* dissimilarities, a function that replaces dissimilarities with shortest paths stepping across intermediate sites while regarding dissimilarities above a threshold as missing data (57). Furthermore, we used weak tie treatment, allowing equal observed dissimilarities to have different fitted values. The scores of the species in the final ordination configuration were obtained using weighted averaging. We checked if the reduction in dimensionality maintained the between-species relationship by checking the stress of the resulting ordination and finding goodness of fit measure for points in nonmetric multidimensional scaling (54). Both PCoA and NMDS ordinations were done using the R package *vegan* (58) and *ecodist* (59). It is important to note that, although the transfer function from observed dissimilarities to ordination distances is non-metric, the resulting NMDS configuration is Euclidean and rotation-invariant (60).

Morphological Disparity. Because we were interested in describing the position of the species in the floral morphospace, we calculated the morphological disparity using indices related to the distance between elements (26, 61). We first determined the absolute position of each of the Brassicaceae species in the morphospace by calculated their Euclidean distance with the overall centroid of the morphospace (61). The disparity between the spring and summer phenotype of *M. arvensis* was also calculated as their Euclidean distance in the floral morphospace. We then calculated the pairwise disparities between all species included in our analysis, between the different morphs of the polymorphic species considered here (Supplementary Data 2), between the species of the genus *Moricandia*, between species of the same genus, between species of different genera but same tribe and between species of different tribes. These disparity values were calculated using the function *dispRity* of the R package *dispRity* using the command *centroid* (62). We checked whether the disparity between spring and summer *M. arvensis* phenotypes was significantly different from the disparities of each of these sets of species using Z-score tests.

Family-wide phylogeny. We retrieved 80 phylogenetic trees from the literature and from the online repositories TreeBase (Table S9). All trees were downloaded in nexus format. The taxonomy of the species included in each tree was checked and updated using the species checklist with accepted names provided by Brassibase (<https://brassibase.cos.uni-heidelberg.de/>) (7, 23, 63). All trees were converted to TreeMan format (64) and concatenated into a single TreeMen file that was then converted into a multiPhylo class. Afterward, we estimated a supertree from this set of trees. Because trees did not share the same taxa, we used the Matrix

representation parsimony method (65). To make this supertree more accurate, it was re-constructed using as backbone phylogeny the tree provided by Walden et al. (7). We removed from the supertree those species without information on floral phenotype, resulting in a tree with 1876 taxa. Because the original trees used to assemble this supertree were very heterogeneous, this supertree was not dated. We finally rooted the supertree using several species belonging to the sister families Capparaceae and Cleomaceae (66). All phylogenetic manipulations were performed using the R libraries *treebase* (67), *ape* (68), *treeman* (64), *phangorn* (69) and *phytools* (70).

We tested whether the position of the Brassicaceae species in the morphospace was associated with the phylogenetic relationship by assessing the phylogenetic signal of the morphospace position. This analysis was performed by means of a multivariate Mantel test, using the pairwise disparity (the Euclidean distance between species in the family-wide morphospace) as a morphological distance and the patristic distances between pairs of tips of the supertree as the phylogenetic distance (71). The correlation method used was Pearson and the statistical significance was found after bootstrapping 999 times the analysis (25). The test was done using the R libraries *vegan* (58) and *ecodist* (59).

Family-wide phylomorphospace. We reconstructed a family-wide phylomorphospace by projecting the phylogenetic relationships provided by the supertree into the floral morphospace. The ancestral character estimation of morphospace coordinate values for each internal tree node was done using maximum likelihood. For this, we used the function *fastAnc* in *phytools*. This function performs fast estimations of the ML ancestral states for continuous traits by re-rooting the tree at all internal nodes and computing the contrasts state at the root each time (70).

We counted the number of intersections between lineages as a measurement of the disorder of the phylomorphospace and evidence of the mode of evolution of the phenotypes (11). For this, we used R codes provided in Ref 11. We compared the observed number of crossings with those expected under several modes of evolution. For this, we counted the number of intersections in 10 simulated sets of species with floral phenotypes following Brownian Motion, Ornstein Uhlenbeck and Early Burst modes of evolution. All simulations were done using as backbone tree the family-wide supertree and considering 1875 species, and by means of the command *mvSIM* in *mvMORPH* (72).

Morphological divergence of the plastic phenotypes. Divergence in floral phenotype was estimated by calculating the disparity of *Moricandia arvensis* and the rest of Brassicaceae species from their ancestors. We first determined the floral phenotype of the Most Recent Common Ancestor (MRCA) using the projection of a recent time-calibrated phylogeny made for

the genus *Moricandia* (73) into the above-described phylomorphospace. We used this phylogeny because it is the only one including all the species of the genus. Once we inferred the coordinates of the MRCA in the morphospace, we calculated the disparity of all the *Moricandia* species and the two plastic phenotypes of *M. arvensis* to it. Afterwards, we calculated the divergence of the two plastic phenotypes from the direct ancestor of *M. arvensis*. This analysis was done for the family-wide supertree and for any of the four time-calibrated phylogenies included in our dataset that had *Moricandia* species (73-76). In addition, we calculated the divergence from the direct ancestors of the rest of Brassicaceae species included in these four phylogenies and in the rest of the time-calibrated trees included in our dataset (Table S9). All floral divergences were calculated using the command *ancestral.dist* of the function *dispRity* in the R package *dispRity* (62).

Pollinator Database. We have compiled a massive database including 21,212 records comprising 455,014 visits done by over 800 animal species from 19 taxonomical orders, 276 families and 43 functional groups to 554 Brassicaceae species belonging to 39 tribes (Supplementary Data 3). Information is coming from literature, personal observation, online repositories and personal communication of several colleagues. The source of information is indicated in the database (Supplementary Data 3, Table S10). In those species studied by us (coded as UNIGEN data origin in the Supplementary Data 3), we conducted flower visitor counts in 1-16 populations per plant species. We visited the populations during the blooming peak, always at the same phenological stage and between 11:00 am and 5:00 pm. In these visits, we recorded the insects visiting the flowers for two hours without differentiating between individual plants. Insects were identified in the field, and some specimens were captured for further identification in the laboratory. We only recorded those insects contacting anthers or stigma and doing legitimate visits at least during part of their foraging at flowers. We did not record those insects only eating petals or thieving nectar without doing any legitimate visit. The information obtained from the literature and online repositories (coded as LITERATURE data origin in the Supplementary Data 3) includes records done during ecological studies, taxonomical studies and naturalistic studies. The reference of every record is included in the dataset. The plant species included in our network do not coexist, implying that this is a clade-oriented network rather than an ecological network (77).

Spatial distribution of pollinator groups. We tested the autocorrelation across the morphospace in the abundance of the functional groups using a multivariate Mantel test. The correlation method used was Pearson, and the statistical significance was found after bootstrapping 999 times the analysis (25). The test was done using the R libraries *vegan* (58).

Pollination niches. In plant species interacting with a diverse assemblage of pollinators, like those included in this study, many pollinator species interact with the flowers in a similar manner, have similar effectiveness and exert similar selective pressures and are thus indistinguishable for the plant (46, 78). These pollinators are thus grouped into functional groups, which are the relevant interaction units in generalised systems (46, 78, 79). We thereby grouped all pollinators visiting the Brassicaceae species using criteria of similarity in body length, proboscis length, morphological match with the flower, foraging behaviour, and feeding habits (46, 78, 79). Table S11 describes the 43 functional groups used in this study. Supplementary Data 4 shows the species with an autogamous pollination system.

We determined the occurrence of different pollination niches in our studied populations and seasons using bipartite modularity, a complex-network metric. Modularity has proven to be a good proxy of interaction niches both in ecological networks, those included coexisting species or population, as well as in clade-oriented network, those including species with information coming from disparate and contrasting sources (77). We constructed a weighted bipartite network, including pollinator data of four populations during the spring and summer flowering. In this network, we pooled the data from the different individuals in a population and did not consider the time difference involved in sampling across different species. We removed all plant species with less than 20 visits. We subsequently determined the modularity level in this weighted bipartite network by using the QuanBiMo algorithm (80). This method uses a Simulated Annealing Monte-Carlo approach to find the best division of populations into modules. A maximum of 10^{10} MCMC steps with a tolerance level = 10^{-10} was used in 100 iterations, retaining the iterations with the highest likelihood value as the optimal modular configuration. We tested whether our network was significantly more modular than random networks by running the same algorithm in 100 random networks, with the same linkage density as the empirical one (81). Modularity significance was tested for each iteration by comparing the empirical versus the random modularity indices using a Z-score test (80). After testing the modularity of our network, we determined the number of modules (82). We subsequently identified the pollinator functional groups defining each module and the plant species ascribed to each module. Modularity analysis was performed using the R package *bipartite* 2.0 (83). We quantified the niche overlap between all pair of Brassicaceae species using the Czekanowski index of resource utilization, an index that measures the area of intersection of the resource utilization histograms of each species pair (84). This index was calculated using the function *niche.overlap* in the R package *spaa* (85).

Estimation of ancestral values of pollination niches. The ancestral states of the pollination niche was inferred for the *Moricandia* lineage by simulate stochastic character mapping of discrete traits with Bayesian posterior probability distribution (86, 87). Three models of character evolution ("ER" - Equal Rates; "SYM" – symmetric; and "ARD" - All Rates Different) were first evaluated using the *fitDiscrete* function of the R package *Geiger* (88). The best model was selected using the Akaike Information Criterion (AIC) and used for stochastic character mapping. The posterior distribution of the transition rate matrix was determined using a Markov chain Monte Carlo (MCMC) simulation, and the stochastic mapping was simulated 100 times. Stochastic character mapping was performed using the *make.simmap* function and a plot of posterior probabilities were mapped using the *describe.simmap* function in R package '*phytools*' (70).

Morphological convergence. To explore morphological convergence, we reconstructed the ancestral states of the species belonging to these two pollination niches and tested for each niche whether the species were morphologically more similar to each other than expected by their phylogenetic relationship (39, 40). We used three different approaches to detect morphological convergence, one based on comparing phenotypic and phylogenetic distances (39) and the other based on comparing the angles formed by two tested clades from their most recent common ancestor with the expected angle according to null evolutionary models (38). Because all these analyses are sensitive to the number of tips in the phylogeny and the inferred branch lengths, we tested for the occurrence of morphological convergence using three independent, time-calibrated phylogenies including more than 45 species (74-76).

Under the first approach, we calculated both distance- and frequency-based measures of convergence (39). Distance-based measures (C1–C4) are calculated between two lineages relative to their distance at the point in evolutionary history where the two lineages were maximally dissimilar. C1 specifically measures the proportion of phenotypic distance closed by evolution, ranging from 0 to 1 (where 1 indicates complete convergence). To calculate C1, ancestral states are reconstructed (via a Brownian motion model of evolution) for two or more putatively convergent lineages, back to their most recent common ancestor. The maximum phenotypic distance between any pair of ancestors (D_{max}) is calculated, and compared with the phenotypic distance between the current putatively convergent taxa (D_{tip}). The greater the difference between D_{max} and D_{tip} , the higher the index. C2 is the raw value of the difference between the maximum and extant distance between the two lineages. C3 is C2 scaled by the total evolution (sum of squared ancestor-to-descendant changes) between the two lineages. C4 is C2 scaled by the total evolution in the whole clade. These four measures quantify incomplete convergence in multidimensional space. Finally, C5, the frequency-based measure, quantifies

and reports the number of convergent events where lineages evolve into a specific region of morphospace (crossing it from outside). C5 sums the number of times through the evolution of a clade that lineages evolve into a given region of phenotypic space. C5 is the number of focal taxa that reside within a limited but convergent region of a phylomorphospace (the phylogenetic connections between taxa represented graphically in a plot of morphological space). The significance of C1–C5 was found by running 1000 simulations for each comparison using Brownian-Motion on a variance–covariance matrix based on data-derived parameters, with convergence measures for each simulation calculated to determine if the observed C value is greater than expected by chance. A priori focal groups forming the basis of convergence tests were the same niche categorizations used in OUwie analyses. These analyses were performed using the R package *convevol* (89).

The second approach to measure convergence was based on comparing the angles formed by two tested clades from their most recent common ancestor with the expected angle according to null evolutionary models (38). Under the “state case”, *search.conv* computes the mean angle over all possible combinations of species pairs using one species per state. Each individual angle is divided by the patristic distance between the species. Significance is assessed by contrasting this value with a family of 1,000 random angles obtained by shuffling the state across the species (38). These analyses were performed using the R package *RRphylo* (90).

The third approach to measure convergence used the Wheatleaf metric (40). This index generates phenotypic (Euclidean) distances from any number of traits across species and penalizes them by phylogenetic distance before investigating similarity (in order to weight close phenotypic similarity higher for distantly related species). It uses an a priori designation of convergent species, which are defined as species belonging to a niche for which the traits are hypothesized to converge. The method then calculates a ratio of the mean (penalized) distances between all species to the mean (penalized) distances between allegedly convergent species. The index detects if convergent species diverge more in phenotypic space from the non-convergent species and show a tighter clustering to each other (40). The significance of this index was found by comparing the empirical values of the index with a distribution of simulated indices obtained running 5000 bootstrap simulations. These analyses were performed using the R package *windex* (91).

Acknowledgments

Authors thank Raquel Sánchez, Angel Caravantes, Isabel Sánchez Almazo, María José Jorquera, and Iván Rodríguez Arós for helping us during several phases of the study. We also thank all contributors to the pollinator database (Table S10) for kindly sending us unpublished

information on Brassicaceae floral visitors. This research is supported by grants from the Spanish Ministry of Science, Innovation and Universities (CGL2015-63827-P, CGL2017-86626-C2-1-P, CGL2017-86626-C2-2-P, UNGR15-CE-3315), Junta de Andalucía (P18-FR-3641, IE19_238 EEZA CSIC), LIFE18 GIE/IT/000755, and Xunta de Galicia (CITACA), including EU FEDER funds. This is a contribution to the Research Unit Modeling Nature, funded by the Consejería de Economía, Conocimiento, Empresas y Universidad, and European Regional Development Fund (ERDF), reference SOMM17/6109/UGR.

Competing interests

The authors declare no competing interests.

References

1. G. R. McGhee, *Convergent Evolution: Limited Forms Most Beautiful*. MIT Press, Cambridge (2011). ISBN:9780262016421
2. J. B. Losos, Convergence, adaptation, and constraint. *Evolution* 65, 1827–1840 (2011). doi:10.1111/j.1558-5646.2011.01289.x
3. S. C. Morris, *Life's solution: Inevitable Humans in a Lonely Universe*. Cambridge University Press, Cambridge (2003). ISBN:9780521603256
4. T. Pearce, Convergence and parallelism in evolution: a Neo-Gouldian account. *Br. J. Philos. Sci.* 63, 429–448 (2011). doi:10.1093/bjps/axr046
5. D. Schluter, *The Ecology of Adaptive Radiation*. Oxford Univ. Press, Oxford, U.K. (2000) ISBN:9780198505228
6. P. Nosil, *Ecological Speciation*. Oxford Univ. Press, Oxford, U.K. (2012) ISBN:9780199587100
7. N. Walden, D. A. German, E. M. Wolf, M. Kiefer, P. Rigault, X. C. Huang, C. Kiefer, R. Schmickl, A. Franzke, B. Neuffer, K. Mummenhoff, Nested whole-genome duplications coincide with diversification and high morphological disparity in Brassicaceae. *Nature Comm.* 11, 3795 (2020). doi:10.1038/s41467-020-17605-7
8. M. Simões, L. Breitzkreuz, M. Alvarado, S. Baca, J. C. Cooper, L. Heins, K. Herzog, B. S. Lieberman, The evolving theory of evolutionary radiations. *Trends Ecol. Evol.* 31, 27–34 (2016). doi:10.1016/j.tree.2015.10.007
9. M. R., Pie, J. S. Weitz, A null model of morphospace occupation. *Am. Nat.* 166, E1–E13 (2005). doi:10.1086/430727
10. K. O. Winemiller, D. B. Fitzgerald, L. M. Bower, E. R. Pianka, Functional traits, convergent evolution, and periodic tables of niches. *Ecol. Lett.* 18, 737–751 (2015). doi:10.1111/ele.12462

- 11.C. T. Stayton, Are our phylomorphospace plots so terribly tangled? An investigation of disorder in data simulated under adaptive and nonadaptive models. *Curr. Zool.* 66, 565–574 (2020). doi:10.1093/cz/zoaa045
- 12.A. L. Pigot, C. Sheard, E. T. Miller, T. P. Bregman, B. G. Freeman, U. Roll, N. Seddon, C. H. Trisos, B. C. Weeks, J. A. Tobias, Macroevolutionary convergence connects morphological form to ecological function in birds. *Nature Ecol. Evol.* 4, 230–239 (2020). doi:10.1038/s41559-019-1070-4
- 13.D. L. Stern, The genetic causes of convergent evolution. *Nat. Rev. Genet.* 14, 751–764 (2013). doi:10.1038/nrg3483
- 14.E. B. Rosenblum, C. E. Parent, E. E. Brandt, The molecular basis of phenotypic convergence. *Annu. Rev. Ecol. Evol. Syst.* 45, 203–226 (2014). doi:10.1146/annurev-ecolsys-120213-091851
- 15.P. A. Christin, D. M. Weinreich, G. Besnard, Causes and evolutionary significance of genetic convergence. *Trends Genet.* 26, 400–405 (2010). doi:10.1016/j.tig.2010.06.005
- 16.M. J. West-Eberhard, *Developmental Plasticity and Evolution*. Oxford Univ. Press, New York, (2003). ISBN: 9780195122343
- 17.S. Sultan, *Organism and Environment: Ecological development, Niche construction, and Adaption*. Oxford Univ. Press, New York, (2015). ISBN:9780199587063
- 18.J. M. Gómez, F. Perfectti, C. Armas, E. Narbona, A. González-Megías, L. Navarro, L. DeSoto, R. Torices, Within-individual phenotypic plasticity in flowers fosters pollination niche shift. *Nature Comm.* 11, 4019 (2020). doi:10.1038/s41467-020-17875-1
- 19.V. Susoy, E. J. Ragsdale, N. Kanzaki, R. J. Sommer, Rapid diversification associated with a macroevolutionary pulse of developmental plasticity, *eLife* 4, e05463 (2015). doi:10.7554/eLife.05463
- 20.N. A. Levis, D. W. Pfennig, Evaluating ‘plasticity-first’ evolution in nature: key criteria and empirical approaches. *Trends Ecol. Evol.* 31, 563–574 (2016). doi:10.1016/j.tree.2016.03.012
- 21.R. J. Sommer, Phenotypic plasticity: from theory and genetics to current and future challenges. *Genetics* 215, 1–13 (2020). doi:10.1534/genetics.120.303163
- 22.M. A. Koch, D. A. German, M. Kiefer, A. Franzke, Database taxonomics as key to modern plant biology. *Trends Plant Sci.* 23, 4–6 (2018). doi:10.1016/j.tplants.2017.10.005
- 23.M. Kiefer, R. Schmickl, D. A. German, T. Mandáková, M. A. Lysak, I. A. Al-Shehbaz, A. Franzke, K. Mummenhoff, A. Stamatakis, M.A. Koch, BrassiBase: introduction to a novel knowledge database on Brassicaceae evolution. *Plant Cell Physiol.* 55, e3–e3 (2014). doi:10.1093/pcp/pct158
- 24.*The Plant List* Version 1.1. Published on the Internet (accessed 1st January 2021). <http://www.theplantlist.org/1.1/browse/A/Brassicaceae/> (2013).

612 25.P. Legendre, L. Legendre, *Numerical Ecology*. Elsevier, Oxford (2012). ISBN:9780080523170

613 26.T. Guillaume, N. Cooper, S. L. Brusatte, K. E. Davis, A. L. Jackson, S. Gerber, A. Goswami, K.

614 Healy, M. J. Hopkins, M. E. Jones, G. T. Lloyd, Disparities in the analysis of morphological

615 disparity. *Biol. Lett.* 16, 20200199 (2020). doi:10.1098/rsbl.2020.0199

616 27.M. Méndez, J. M. Gómez, Phenotypic gender in *Hormathophylla spinosa* (Brassicaceae), a

617 perfect hermaphrodite with tetradynamous flowers, is variable. *Plant Syst. Evol.* 262, 225–23

618 (2006). doi:10.1007/s00606-006-0462-5

619 28. V. L. Soza, V. Le Huynh, V. S. Di Stilio, Pattern and process in the evolution of the sole

620 dioecious member of Brassicaceae. *EvoDevo* 5, 42 (2014). doi:10.1186/2041-9139-5-42

621 29.E. Narbona, H. Wang, P. L. Ortiz, M. Arista, E. Imbert, Flower colour polymorphism in the

622 Mediterranean Basin: occurrence, maintenance and implications for speciation. *Plant Biol.* 20,

623 8–20 (2018). doi:10.1111/plb.12575.

624 30.C. A. Dick, J. Buenrostro, T. Butler, M. L. Carlson, D. J. Kliebenstein, J. B. Whittall, Arctic

625 mustard flower color polymorphism controlled by petal-specific downregulation at the threshold

626 of the anthocyanin biosynthetic pathway. *PLoS One* 6, e18230 (2011).

627 doi:10.1371/journal.pone.0018230

628 31.B. Zhang, C. Liu, Y. Wang, X. Yao, F. Wang, J. Wu, G. J. King, K. Liu, Disruption of a

629 carotenoid cleavage dioxygenase 4 gene converts flower colour from white to yellow in

630 *Brassica* species. *New Phytol.* 206, 1513–1526 (2015). doi:10.1111/nph.13335

631 32.S. Faisal, Y. Guo, S. Zang, B. Cao, G. Qu, S. Hu, Morphological and genetic analysis of a

632 cleistogamous mutant in rapeseed (*Brassica napus* L.). *Genet. Resour. Crop Evol.* 65, 397–

633 403 (2018). doi:10.1007/s10722-017-0598-x

634 33.G. Theissen, Homeosis of the angiosperm flower: studies on three candidate cases of

635 saltational evolution. *Palaeodiversity* 3, 131–139 (2010).

636 34.M. V. Byzova, J. Franken, M. G. Aarts, J. de Almeida-Engler, G. Engler, C. Mariani, M. M. V.

637 L. Campagne, G. C. Angenent, *Arabidopsis* STERILE APETALA, a multifunctional gene

638 regulating inflorescence, flower, and ovule development. *Genes Dev.* 13, 1002–1014 (1999).

639 doi:10.1101/gad.13.8.1002.

640 35.T. Van der Niet, S. D. Johnson, Phylogenetic evidence for pollinator-driven diversification of

641 angiosperms. *Trends Ecol. Evol.* 27, 353–361 (2012). doi:10.1016/j.tree.2012.02.002

642 36.A. Franzke, M. A. Lysak, I. A. Al-Shehbaz, M. A. Koch, K. Mummenhoff, Cabbage family

643 affairs: the evolutionary history of Brassicaceae. *Trends Plant Sci.* 16, 108–116 (2011).

644 doi:10.1016/j.tplants.2010.11.005

645 37.J. M. Gómez, F. Perfectti, J. Lorite, The role of pollinators in floral diversification in a clade of

646 generalist flowers. *Evolution* 69, 863–878 (2015). doi:10.1111/evo.12632

- 38.S. Castiglione, C. Serio, D. Tamagnini, M. Melchionna, A. Mondanaro, M. Di Febbraro, A. Profico, P. Piras, F. Barattolo, P. Raia, A new, fast method to search for morphological convergence with shape data. *PloS One* 14: e0226949 (2019). doi:10.1371/journal.pone.0226949
- 39.C. T. Stayton, The definition, recognition, and interpretation of convergent evolution, and two new measures for quantifying and assessing the significance of convergence. *Evolution* 69, 2140–2153 (2015). doi:10.1111/evo.12729
- 40.K. Arbuckle, C. M. Bennett, M. P. Speed, A simple measure of the strength of convergent evolution. *Methods Ecol. Evol.* 5, 685–693 (2014). doi:10.1111/2041-210X.12195
- 41.K. Faegri, L. Van Der Pijl, *Principles of Pollination Ecology*. Elsevier, Oxford (1980). ISBN:9780080164212
- 42.A. S. Dellinger, Pollination syndromes in the 21st century: where do we stand and where may we go?. *New Phytol.* 228, 1193–1213 (2020). doi:10.1111/nph.16793
- 43.R. D. Phillips, R. Peakall, T. van der Niet, S. D. Johnson, Niche perspectives on plant–pollinator interactions. *Trends Plant Sci.* 25, 779–793 (2020). doi:10.1016/j.tplants.2020.03.009.
- 44.C. A. Wessinger, L. C. Hileman, Parallelism in flower evolution and development. *Annu. Rev. Ecol. Evol. Syst.* 51, 387–408 (2020). doi:10.1146/annurev-ecolsys-011720-124511
- 45.J. D. Thomson, P. Wilson, Explaining evolutionary shifts between bee and hummingbird pollination: convergence, divergence, and directionality. *Int. J. Plant Sci.* 169, 23–38 (2008). doi:10.1086/523361
- 46.J. M. Gómez, F. Perfectti, M. Abdelaziz, J. Lorite, A. J. Muñoz-Pajares, J. Valverde, Evolution of pollination niches in a generalist plant clade. *New Phytol.* 205, 440–453 (2015). doi:10.1111/nph.13016
- 47.E. C. Snell-Rood, M. E. Kobiela, K. L. Sikkink, A. M. Shepherd, Mechanisms of plastic rescue in novel environments. *Annu. Rev. Ecol. Evol. Syst.* 49, 331–354 (2018). doi:10.1146/annurev-ecolsys-110617-062622
- 48.R. J. Fox, J. M. Donelson, C. Schunter, T. Ravasi, J. D. Gaitán-Espitia, Beyond buying time: the role of plasticity in phenotypic adaptation to rapid environmental change. *Phil. Trans. R. Soc. B* 374, 20180174 (2019). doi:10.1098/rstb.2018.0174
- 49.N. A. Levis, D. W. Pfennig, Plasticity-led evolution: evaluating the key prediction of frequency-dependent adaptation. *Proc. Biol. Sci.* 286, 20182754 (2019). doi:10.1098/rspb.2018.2754
- 50.M. R. Warner, L. Qiu, M. J. Holmes, A. S. Mikheyev, T. A. Linksvayer, Convergent eusocial evolution is based on a shared reproductive groundplan plus lineage-specific plastic genes. *Nature Comm.* 10, 1–11 (2019). doi:10.1038/s41467-019-10546-w

51. R. F. Schneider, A. Meyer, How plasticity, genetic assimilation and cryptic genetic variation may contribute to adaptive radiations. *Mol. Ecol.* 26, 330–350 (2017). doi:10.1111/mec.13880

52. T. Guillerme, N. Cooper, Time for a rethink: time sub-sampling methods in disparity-through-time analyses. *Palaeontology* 61, 481–493 (2018). doi:10.1111/pala.12364

53. P. Legendre, D. Borcard, P. R. Peres-Neto, Analyzing beta diversity: Partitioning the spatial variation of community composition data. *Ecol. Monogr.* 75, 435–450 (2005). doi:10.1890/05-0549

54. P. Mair, I. Borg, Rusch T. Goodness-of-fit assessment in multidimensional scaling and unfolding. *Multivar. Behav. Res.* 51, 772–789 (2016). doi:10.1080/00273171.2016.1235966.

55. J. B. Kruskal, Multidimensional scaling by optimizing goodness-of-fit to a nonmetric hypothesis. *Psychometrika* 29, 1–28 (1964).

56. J. B. Kruskal, Nonmetric multidimensional scaling: a numerical method. *Psychometrika* 29, 115–129 (1964).

57. G. De'ath, Extended dissimilarity: a method of robust estimation of ecological distances from high beta diversity data. *Plant Ecol.* 144, 191–199 (1999). doi:10.1023/A:1009763730207

58. J. Oksanen, F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, H. Wagner, 2013 Package 'vegan'. *Community ecology package, version 2*, 1–295 <https://cran.r-project.org>, <https://github.com/vegandevs/vegan> (2019)

59. S. C. Goslee, Urban, D. L. The ecodist package for dissimilarity-based analysis of ecological data. *J. Stat. Softw.* 22, 1–19 (2007). doi:10.18637/jss.v022.i07

60. J. Oksanen, Vegan: an introduction to ordination. URL <http://cran.r-project.org/web/packages/vegan/vignettes/introvegan.pdf> (2020).

61. T. Guillerme, M. N. Puttick, A. E. Marcy, V. Weisbecker, Shifting spaces: Which disparity or dissimilarity measurement best summarize occupancy in multidimensional spaces?. *Ecol. Evol.* 10, 7261–7275 (2020). doi:10.1002/ece3.6452

62. T. Guillerme, dispRity: a modular R package for measuring disparity. *Methods Ecol. Evol.* 9, 1755–1763 (2018). doi:10.1111/2041-210X.13022

63. M. A. Koch, M. Kiefer, D. A. German, I. A. Al-Shehbaz, A. Franzke, K. Mummenhoff, R. Schmickl, BrassiBase: Tools and biological resources to study characters and traits in the Brassicaceae — version 1.1. *Taxon* 61, 1001–1009 (2012). doi:10.1002/tax.615007

64. D. J., Bennett, M. D., Sutton, S. T. Turvey, Treeman: an R package for efficient and intuitive manipulation of phylogenetic trees. *BMC Res. Notes* 10, 30 (2017). doi:10.1186/s13104-016-2340-8

65. M. A. Ragan, Phylogenetic inference based on matrix representation of trees. *Mol. Phylogenetics Evol.* 1, 53–58 (1992). doi:10.1016/1055-7903(92)90035-F

- 66.S. Bayat, M. E. Schranz, E. H. Roalson, J. C. Hall, Lessons from Cleomaceae, the sister of crucifers. *Trends Plant Sci.* 23, 808–821 (2018). doi:10.1016/j.tplants.2018.06.010.
- 67.C. Boettiger, D. Temple Lang, Treebase: an R package for discovery, access and manipulation of online phylogenies. *Methods Ecol. Evol.* 3, 1060–1066 (2012). doi:10.1111/j.2041-210X.2012.00247.x
- 68.E., Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 35, 526–528 (2019). doi:10.1093/bioinformatics/bty633
- 69.K. Schliep, phangorn: phylogenetic analysis in R. *Bioinformatics* 27, 592–593 (2011). doi:10.1093/bioinformatics/btq706
- 70.L. J. Revell, phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol. Evol.* 3, 217–223 (2012). doi:10.1111/j.2041-210X.2011.00169.x
- 71.O. J. Hardy, S. Pavoine, Assessing phylogenetic signal with measurement error: a comparison of Mantel tests, Blomberg et al.'s *K*, and phylogenetic distograms. *Evolution* 66, 2614–2621 (2012). doi:10.1111/j.1558-5646.2012.01623.x
- 72.J. Clavel, G. Escarguel, G. Merceron, mvMORPH: an R package for fitting multivariate evolutionary models to morphometric data. *Methods Ecol. Evol.* 6, 1311–1319 (2015). doi:10.1111/2041-210X.12420
- 73.F. Perfectti, J. M. Gómez, A. González-Megías, M. Abdelaziz, J. Lorite, Molecular phylogeny and evolutionary history of *Moricandia* DC (Brassicaceae). *PeerJ* 5, e3964 (2017). doi:10.7717/peerj.3964
- 74.S. A. Smith, J. W. Brown, Constructing a broadly inclusive seed plant phylogeny. *Am. J. Bot.* 105, 302–314 (2018). doi:10.1002/ajb2.1019
- 75.M. L. Gaynor, J. Ng, R. G. Laport, Phylogenetic structure of plant communities: are polyploids distantly related to co-occurring diploids?. *Frontiers Ecol. Evol.* 6, 52 (2018). doi:10.3389/fevo.2018.00052
- 76.X.-C. Huang, D. A. German, M. A. Koch, Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. *Ann. Bot.* 125, 29–47 (2019). doi:10.1093/aob/mcz123
- 77.J. M. Gómez, M. Verdú, F. Perfectti, Ecological interactions are evolutionarily conserved across the entire tree of life. *Nature* 465, 918–921 (2010). doi:10.1038/nature09113
- 78.C. B. Fenster, W. S. Armbruster, P. Wilson, M. R. Dudash, J. D. Thomson, Pollination syndromes and floral specialization. *Annu. Rev. Ecol. Evol. Syst.* 35, 375–403 (2004). doi:10.1146/annurev.ecolsys.34.011802.132347
- 79.J. M. Gómez, R. Torices, J. Lorite, C. P. Klingenberg, F. Perfectti, The role of pollinators in the evolution of corolla shape variation, disparity and integration in a highly diversified plant family with a conserved floral bauplan. *Ann. Bot.* 117, 889–904 (2016). doi:10.1093/aob/mcv194

754 80.C. F. Dormann, R. Strauss, A method for detecting modules in quantitative bipartite networks.
755 *Methods Ecol. Evol.* 5, 90–98 (2014). doi:10.1111/2041-210X.12139
756 81.R. Guimerà, L. A. N. Amaral, Functional cartography of complex metabolic networks. *Nature*
757 433, 895–900 (2005). doi:10.1038/nature03288
758 82.M. E. J. Newman, Analysis of weighted networks. *Phys. Rev. E* 70, 056131 (2004).
759 doi:10.1103/PhysRevE.70.056131
760 83.C. F. Dormann, B. Gruber, J. Fründ, Introducing the bipartite package: Analysing ecological
761 networks. *R News* 8, 8–11 (2008).
762 84.P. Feinsinger, E. E. Spears, R. W. Poole, A simple measure of niche breadth. *Ecology* 62, 27–
763 32 (1981). doi:10.2307/1936664
764 85.J. Zhang, M. J. Zhang, Package ‘spaa’. *R package version* 1. [https://CRAN.R-](https://CRAN.R-project.org/package=spaa)
765 [project.org/package=spaa](https://CRAN.R-project.org/package=spaa) (2013).
766 86.J. P. Huelsenbeck, R., Nielsen, J. P. Bollback, Stochastic mapping of morphological
767 characters. *Syst. Biol.* 52, 131–158 (2003). doi:10.1080/10635150390192780
768 87.J. P. Bollback, SIMMAP: stochastic character mapping of discrete traits on phylogenies. *BMC*
769 *Bioinformatics* 7, 88 (2006). doi:10.1186/1471-2105-7-88
770 88.L. J. Harmon, J. T. Weir, C. D. Brock, R. E. Glor, W. Challenger, GEIGER: investigating
771 evolutionary radiations. *Bioinformatics* 24, 129–131 (2008). doi:10.1093/bioinformatics/btm538
772 89.C. T. Stayton, Convevol: quantifies and assesses the significance of convergent evolution. R
773 package version 1.0. See <http://cran.r-project.org/web/packages/convevol/index.html> (2018).
774 90.P. Raia, S. Castiglione, C. Serio, A. Mondanaro, M. Melchionna, M. Di Febbraro, RRphylo:
775 Phylogenetic ridge regression methods for comparative studies. *Methods Ecol. Evol.* 9, 974–
776 983 (2019). doi:10.1111/2041-210X.12954
777 91.K., Arbuckle, A. Minter, Windex: Analyzing convergent evolution using the Wheatsheaf index
778 in R. *Evol. Bioinformatics* 11, EBO-S20968 (2015). doi:10.4137/EBO.S20968
779

Figures




Figure 1. Plasticity-mediated floral disparity. (A) Floral morphospace of the Brassicaceae, showed as the projection of 31 traits recorded in 3140 species onto two NMDS axes. The position of the spring and summer phenotypes of *Moricandia arvensis* is linked by a thick lilac dashed line. We have also indicated the movements across this morphospace of several species changing their phenotypes due to floral colour polymorphism (black lines), single mutations in floral colour (blue lines), changes in breeding systems (orange lines), changes in gender expression (green lines), homeotic mutations (brown lines), and plasticity (lilac lines). Numbers matching species are as follow: 1-*Lobularia maritima*; 2-*Raphanus raphanistrum*; 3-*Matthiola incana*; 4-*Mathiola fruticulosa*; 5-*Erysimum cheiri*; 6-*Cakile maritima*; 7-*Matthiola lunata*; 8- *Marcus-kochia littorea*; 9- *Hesperis matronalis*; 10- *Hesperis laciniata*; 11-*Parrya nudicalis*; 12-*Streptanthus glandulosus*; 13- *Eruca vesicaria*; 14- *Capsella bursa-pastoris*; 15-*Hormathophylla spinosa*; 16- *Brassica napus*; 17- *Cardamine hirsuta*; 18- *Lepidium sisymbrioides*; 19-*Lepidium solandri*; 20-*Arabidopsis thaliana*; 21-*Boechera stricta*; 22-*Leavenworthia stylosa*; 23-*Leavenworthia crassa*; 24- *Pachycladon stellatum*; 25- *Pachycladon wallii*; 26-*Cardamine kokairensis*; 27-*Brassica rapa*. (B) Shepard plot showing the goodness of fit of the NMDS ordination. (C) *Moricandia arvensis* in spring. (D) *Moricandia arvensis* in summer. (E) Magnitude of floral disparity between different taxonomic levels of Brassicaceae species. The number above each boxplot shows the number of disparities per level. We have compared this value with the disparity between spring and summer

802 phenotypes of *M. arvensis* (this comparison with boxplots in red is statistically significant at $P <$
803 0.05, in orange is marginally significant at $P < 0.1$, and in grey is non-significant).

804

805




Figure 2. Phylogenetic-mediated floral divergence. (A) Floral phylomorphospace using the supertree that includes 1876 Brassicaceae species. (B) Phylomorphospace considering only the eight *Moricandia* species, using the Perfectti et al.'s phylogeny (phylogeny # 1 in Table S8). (C) Floral disparity to the nearest ancestor, according to the supertree and 18 time-calibrated phylogenies (phylogeny codes in Table S8). We show the disparity between the two *M. arvensis* phenotypes and their direct ancestor (spring: lilac dots; summer: white dots) in those phylogenies

814 that include *Moricandia*. We also show the disparities to their direct ancestors of those
815 Brassicaceae species included in time-calibrated phylogenies of more than 45 species.
816
817




Figure 3. Plasticity-mediated changes in pollination niches. (A) Outcome of the modularity analysis showing the number of pollination niches inferred, the among-niche differences in relative frequency of each pollinator functional group, and the pollinator functional groups defining the niches (n = 511 Brassicaceae species). (B) Morphospatial distribution of the eight pollination niches detected in Brassicaceae. Insect silhouettes were drawn by Divulgare (www.divulgare.net) under a Creative Commons license (<http://creativecommons.org/licenses/by-nc-sa/3.0>). (C) Estimate of the ancestral pollination niche of the *Moricandia* lineage using a stochastic character mapping inference analysis. The numbers underneath each ancestral node indicate the posterior Bayesian probability of belonging to pollination niche 5.





Figure 4. Plasticity-mediated floral convergence. Convergent lineages crossing into the region of the morphospace delimited by the pollination niche of the *M. arvensis* during spring (the shade convex hull) according to (A) Smith & Brown's phylogeny, (B) Gaynor et al.'s phylogeny, and (C) Huang et al.'s phylogeny (phylogenies 2-4, respectively, in Table S8). Convergent lineages crossing into the region of the morphospace delimited by the pollination niche of the *M. arvensis* during summer (the shade convex hull) according to (D) Smith & Brown's phylogeny, (E) Gaynor et al.'s phylogeny, and (F) Huang et al.'s phylogeny. Red arrows indicate the plasticity-mediated convergence, blue arrows the convergence events of the other lineages. The small purple area in all panels is the region of the floral morphospace that includes the lineages that have converged with the entire *Moricandia* clade according to each time-calibrated phylogeny.

SI FIGURES

Figure S1. Association among the 31 pollination traits of 3140 Brassicaceae species. Trait vectors represent the Spearman correlations, with the length and direction indicating the relationship with composite NMDS axes.



SI TABLES

Table S1. Fitting of the floral traits onto the NMDS vectors.

	Floral traits	NMDS1	NMDS2	r²	P value
1	Plant height	-0.20147	0.97949	0.1126	0.001
2	Flower display size	0.80044	0.59942	0.0415	0.001
3	Inflorescence architecture	0.94742	-0.32001	0.0641	0.001
4	Presence of apetalous flowers	0.36770	-0.92994	0.1452	0.001
5	Number of symmetry axes of the corolla	0.66735	-0.74474	0.1121	0.001
6	Orientation of dominant symmetry axis of the corolla	0.98547	0.16987	0.2650	0.001
7	Corolla with overlapped petals	-0.98142	-0.19188	0.0685	0.001
8	Corolla with multilobed petals	-0.45075	-0.89265	0.0110	0.001
9	Corolla with visible sepals	0.70871	-0.7055	0.2729	0.001
10	Petal length	-0.55200	0.83385	0.6287	0.001
11	Sepal length	-0.47963	0.87747	0.5594	0.001
12	Asymmetric petals	-0.49289	-0.87009	0.0604	0.001
13	Petal limb length	-0.67904	0.73410	0.4482	0.001
14	Length of long stamen	-0.48773	0.87300	0.4915	0.001
15	Length of short stamen	-0.42415	0.90559	0.4029	0.001
16	Herkogamy	-0.78612	0.61808	0.1671	0.001
17	Herkogamy category	-0.62172	0.78324	0.3864	0.001
18	Visible anthers	0.76256	-0.64691	0.1526	0.001
19	Exserted stamens	0.29511	-0.95546	0.0033	0.009
20	Number of stamens	-0.28006	0.95998	0.1182	0.001
21	Concealed nectaries	-0.72601	0.68768	0.4101	0.001
22	Petal carotenoids	0.61077	0.7918	0.7264	0.001
23	Petal anthocyanins	-0.98947	0.14476	0.5757	0.001
24	Presence of bullseyes	-0.84653	0.53235	0.2654	0.001
25	Presence of veins in the petals	-0.94487	0.32746	0.3343	0.001
26	Coloured sepal	0.99357	-0.11318	0.0039	0.002
27	Relative attractiveness of petals versus sepals	-0.00049	0.99990	0.1001	0.001
28	Petal hue	0.60720	-0.79455	0.2096	0.001
29	Petal colour as b CIELAB	0.75326	0.65772	0.7232	0.001
30	Sepal Hue	0.61109	-0.79156	0.0791	0.001
31	Sepal colour as b CIELAB	0.87412	0.48571	0.5605	0.001

Table S2. Disparity, calculated as the Euclidean distance in the family-wide floral morphospace, between each of the 38 morphs included in our dataset (see Supplementary Data 2 for details and references) and their respective wild types.

Type of polymorphism	Species	Morph	NMDS
Breeding system	<i>Brassica napus</i>	cleistogamous mutant	0.035856865
Breeding system	<i>Brassica rapa</i>	female sterility mutant	0.133332180
Breeding system	<i>Cardamine kokairensis</i>	cleistogamous mutant	0.111696867
Breeding system	<i>Leavenworthia crassa</i>	Outcrosser morph	0.015771684
Colour mutant	<i>Brassica napus</i>	white mutant	0.173001555
Colour mutant	<i>Moricandia arvensis</i>	white mutant	0.149565069
Flower colour polymorphism	<i>Boechera stricta</i>	pink morph	0.060580070
Flower colour polymorphism	<i>Boechera stricta</i>	purple morph	0.054429426
Flower colour polymorphism	<i>Cakile maritima</i>	white morph	0.016547552
Flower colour polymorphism	<i>Eruca vesicaria</i>	white morph	0.082257145
Flower colour polymorphism	<i>Erysimum cheiri</i>	purple cultivar	0.073997237
Flower colour polymorphism	<i>Erysimum cheiri</i>	white cultivar	0.026460490
Flower colour polymorphism	<i>Hesperis laciniata</i>	white morph	0.069557518
Flower colour polymorphism	<i>Hesperis matronalis</i>	white morph	0.094873544
Flower colour polymorphism	<i>Hormathophylla spinosa</i>	white morph	0.101607789
Flower colour polymorphism	<i>Leavenworthia stylosa</i>	white morph	0.066269050
Flower colour polymorphism	<i>Lobularia maritima</i>	deep purple cultivar	0.090407642
Flower colour polymorphism	<i>Marcus-kochia littorea</i>	light pink morph	0.019542946
Flower colour polymorphism	<i>Matthiola fruticulosa</i>	greenish morph	0.008349920
Flower colour polymorphism	<i>Matthiola incana</i>	white cultivar	0.138357728
Flower colour polymorphism	<i>Matthiola lunata</i>	white morph	0.079964493
Flower colour polymorphism	<i>Parrya nudicaulis</i>	white morph	0.115226060
Flower colour polymorphism	<i>Raphanus raphanistrum</i>	white morph	0.082072459
Flower colour polymorphism	<i>Raphanus raphanistrum</i>	yellow morph	0.032685632
Flower colour polymorphism	<i>Raphanus raphanistrum</i>	pink morph	0.082778860
Flower colour polymorphism	<i>Streptanthus glandulosus</i>	white morph	0.014907493
Gender dimorphism	<i>Hormathophylla spinosa</i>	female morph	0.035783402
Gender dimorphism	<i>Hormathophylla spinosa</i>	male morph	0.025226337
Gender dimorphism	<i>Lepidium sisymbrioides</i>	female morph	0.067913758
Gender dimorphism	<i>Lepidium solandri</i>	female morph	0.065685257
Gender dimorphism	<i>Pachycladon stellatum</i>	female morph	0.014244744
Gender dimorphism	<i>Pachycladon wallii</i>	male morph	0.059320935
Homeotic mutant	<i>Arabidopsis thaliana</i>	AGAMOUS mutant	0.001052532
Homeotic mutant	<i>Arabidopsis thaliana</i>	APETALA1 mutant	0.119617731
Homeotic mutant	<i>Arabidopsis thaliana</i>	APETALA3 mutant	0.102445400
Homeotic mutant	<i>Capsella bursapastoris</i>	Spe mutant	0.051187659
Phenotypic plasticity	<i>Cardamine hirsuta</i>	plastic change in stamens number	0.010624208
Phenotypic plasticity	<i>Cardamine hirsuta</i>	plastic change in petal number	0.060755859

Table S3. Floral disparity of each species of *Moricandia* from the most recent common ancestor (MRCA) of the genus and from the direct ancestor of each species.

Species	Disparity to MRCA	Disparity to direct ancestor
<i>Moricandia foetida</i>	0.039566021	0.13987379
<i>Moricandia moricandioides</i>	0.059142993	0.03730920
<i>Moricandia nitens</i>	0.041727403	0.07347209
<i>Moricandia rytidocarpoides</i>	0.025809374	0.10503623
<i>Moricandia sinaica</i>	0.027589330	0.10550411
<i>Moricandia spinosa</i>	0.019579372	0.20959717
<i>Moricandia suffruticosa</i>	0.063884437	0.20700167
<i>Moricandia arvensis</i> spring phenotype	0.080840848	0.02385532
<i>Moricandia arvensis</i> summer phenotype	0.195061288	0.28741584

Table S4. Significance of the Mantel tests checking for spatial autocorrelation across the morphospace of the pollinator functional groups. Due to the small abundance of some pollinators, the original 43 functional groups have been pooled in 26 main functional groups.

Functional Groups	Mantel R	p value
Ant	0.047	0.055
Autogamy	0.257	0.001
Bug	0.032	0.192
Butterfly	0.089	0.001
Hawkmoth	0.054	0.035
Hoverfly	0.072	0.003
Large bee fly	0.043	0.079
Large beetle	0.046	0.063
Large fly	0.052	0.044
Large wasp	0.053	0.048
Long tongued fly	0.092	0.001
Long tongued large bee	0.242	0.001
Long tongued medium-sized bee	0.051	0.041
Moth	0.039	0.113
Nocturnal moth	0.222	0.001
Other	0.018	0.497
Pollen wasp	0.026	0.313
Small beetle	0.012	0.613
Small diving beetle	0.003	0.899
Small fly	0.112	0.001
Small wasp	0.041	0.112
Short tongued large bee	0.065	0.004
Short tongued medium-sized bee	0.012	0.590
Short tongued small bee	0.073	0.001
Short tongued extra small bee	0.020	0.454
Thrips	-0.014	0.758

Table S5. Differences between the two *Moricandia arvensis* phenotypes in the visitation frequency (both in absolute number of insects and in proportion of visits) of every pollinator functional group. Fifteen censuses of 1 hr and two researchers per phenotype.

Pollinator functional group	spring phenotype	summer phenotype	spring phenotype (proportion)	summer phenotype (proportion)
Hawkmoth	91	0	0.036	0.000
Honeybee	40	0	0.016	0.000
Large beefly	309	72	0.124	0.148
Large beetle	30	40	0.012	0.082
Large butterfly	280	66	0.112	0.135
Large fly	38	0	0.015	0.000
Large hoverfly	8	7	0.003	0.014
Long tongued large bee	1131	5	0.453	0.010
Long tongued medium-sized bee	226	0	0.090	0.000
Small beefly	6	0	0.002	0.000
Small beetle	21	8	0.008	0.016
Small butterfly	11	0	0.004	0.000
Small diving beetle	25	12	0.010	0.025
Small fly	12	0	0.005	0.000
Small hoverfly	49	37	0.020	0.076
Small moth	2	7	0.001	0.014
Short tongued large bee	89	3	0.036	0.006
Short tongued medium-sized bee	36	0	0.014	0.000
Short tongued small bee	78	207	0.031	0.424
Short tongued extra small bee	1	0	0.000	0.000
Thrips	15	24	0.006	0.049

Table S6. Outcome of the analyses to test the occurrence of floral convergence among plants from niches 3 and 5. **Angle** is the mean theta angle between all species belonging to the same niche. **Angle/time** is the angle divided by time distance. The significance of these angles has been found by comparing with a null model consisting in shuffling each niche 1,000 times across the tree tips and calculating a distribution of random angle. **C1** measures the proportion of phenotypic distance closed by evolution, ranging from 0 to 1 (where 1 indicates complete convergence). **C2** is the raw value of the difference between the maximum and extant distance between the lineages. **C3** is C2 scaled by the total evolution (sum of squared ancestor-to-descendant changes) between the two lineages. **C4** is C2 scaled by the total evolution in the whole clade. The significance of C1-C2, was evaluated by running 1000 simulations for each comparison using Brownian-Motion models. **Wheatleaf** is the ratio of the mean (penalized) distances between all species to the mean (penalized) distances between allegedly convergent species. Significance found by running 2000 bootstrapping simulations. In bold, significant values.

Phylogenies	Smith & Brown 2018		Gaynor et al. 2018		Huang et al. 2019	
	Value	p	Value	p	Value	p
Niche 3						
Angle	80.587	0.008	79.431	0.002	64.930	0.055
Angle/time	2.350	0.719	1.645	0.397	4.023	0.815
C1	0.373	0.000	0.472	0.000	0.415	0.000
C2	0.104	0.000	0.142	0.000	0.104	0.000
C3	0.141	0.000	0.166	0.000	0.219	0.000
C4	0.003	0.720	0.002	0.700	0.008	0.600
Wheatleaf	0.830	0.986	0.940	0.715	1.060	0.028
Niche 5						
Angle	70.093	0.002	73.491	0.002	58.313	0.049
Angle/time	1.393	0.021	1.783	0.745	2.474	0.011
C1	0.356	0.000	0.472	0.000	0.240	0.000
C2	0.110	0.000	0.142	0.000	0.075	0.000
C3	0.128	0.000	0.166	0.000	0.118	0.000
C4	0.003	0.727	0.002	0.700	0.006	0.545
Wheatleaf	1.120	0.673	1.170	0.094	0.920	0.978

Table S7. Outcome of the analyses testing for morphological convergence between the *Moricandia* clade and the rest of clades included in each time-calibrated phylogeny. **Clade size** is the number of species within the *Moricandia* clade. θ_{real} is the mean angle over all possible combinations of pairs of species taking one species per clade. θ_{ace} is the mean angle between ancestral states between each pairs of clades. $\text{dist}_{\text{mrca}}$ is the patristic distance (sum of brach length) between the most recent common ancestors of each pair of clade. We indicate the convergent clades and the pollination niches of each species included in the convergent clades. In red *Moricandia* clades including *Moricandia arvensis* spring phenotype. Tribes (E= Erysimeae, A= Anthonieae, C=Cardamineae, M=Malcolmieae, An=Anastatieae).

Moricandia clade	Clade 2	Clade size	θ_{real}	θ_{ace}	$\text{dist}_{\text{mrca}}$	$\theta_{\text{real}}/\text{dist}_{\text{mrca}}$	$\theta_{\text{real}}/\text{dist}_{\text{mrca}}$ p-value	$\theta_{\text{ace}}+\theta_{\text{real}}/\text{dist}_{\text{mrca}}$	$\theta_{\text{ace}}+\theta_{\text{real}}/\text{dist}_{\text{mrca}}$ p-value	Convergent clades	Tribe	Niche
Smith & Brown's phylogeny												
253	347	7	15.200	4.420	124.236	0.122	0.058	0.158	0.012	<i>Erysimum bicolor/ scoparium</i>	E	5,5
254	347	5	19.035	5.389	129.718	0.147	0.058	0.188	0.016	<i>Erysimum bicolor/ scoparium</i>	E	5,5
255	347	4	22.601	6.331	133.342	0.169	0.052	0.217	0.019	<i>Erysimum bicolor/ scoparium</i>	E	5,5
256	347	2	6.180	7.281	133.538	0.046	0.026	0.101	0.005	<i>Erysimum bicolor/ scoparium</i>	E	5,5
256	316	2	12.950	2.109	59.810	0.217	0.061	0.252	0.018	<i>Matthiola</i> clade	A	6,1
256	375	2	8.543	13.786	49.285	0.173	0.066	0.453	0.045	<i>Cardamine penthaphyllo/ pratensis</i>	C	3,7
256	405	2	8.616	1.048	44.577	0.193	0.068	0.217	0.022	<i>Malcolmia maritima— Marcus-kochia ramosissima</i>	M/An	5,7
256	335	2	28.715	33.529	128.518	0.223	0.077	0.484	0.049	<i>Erysimum popovii/ bastetanum/ semperflorens</i>	E	5,5,6
257	347	2	39.021	6.003	133.462	0.292	0.103	0.337	0.015	<i>Erysimum bicolor/ scoparium</i>	E	5,5
258	347	2	5.613	4.410	124.383	0.045	0.012	0.081	0.002	<i>Erysimum bicolor/ scoparium</i>	E	5,5
Gaynor et al.'s phylogeny												
481	334	2	10.943	17.832	78.194	0.140	0.052	0.368	0.037	<i>Erysimum bicolor/ scoparium</i>	E	5,5
479	334	2	9.357	12.654	81.141	0.115	0.052	0.271	0.033	<i>Erysimum bicolor/ scoparium</i>	E	5,5
479	333	2	12.809	14.978	81.151	0.158	0.070	0.342	0.049	<i>Erysimum lagascae/ rondae</i>	E	5,3
Huang et al.'s phylogeny												
143	83	2	4.960	0.985	23.966	0.207	0.023	0.248	0.001	<i>Erucaria</i> clade	B	3,5
143	82	2	7.094	0.554	22.049	0.322	0.041	0.347	0.002	<i>Erucaria</i> clade + <i>Cakile</i> clade	B	3,3,3,5
143	81	2	6.651	40.317	18.008	0.369	0.046	2.608	0.170	<i>Erucaria</i> clade + <i>Cakile</i> clade + <i>Eremophyton chevallieri</i>	B	3,3,3,5,5
143	84	2	9.229	1.876	23.466	0.393	0.066	0.473	0.003	<i>Cakile</i> clade	B	3,3
143	86	2	11.634	12.691	24.281	0.479	0.074	1.002	0.026	<i>Zilla</i> clade	B	5,5
143	85	2	9.383	12.437	24.007	0.391	0.076	0.909	0.029	<i>Zilla</i> clade + <i>Foleyola billotii</i>	B	3,5,5

Table S8. Description of floral traits related to pollinator attraction used to generate the floral morphospace in Brassicaceae. Pollinators respond to the variability of numerous phenotypic traits of plants, and the magnitude of their response shapes the reproductive success of the plants. We estimated for each plant included in our data set the values of several important floral traits.

1) Plant height. Plant height has strong direct and indirect effects on plant fitness in many Brassicaceae. The assessment of plant height for a large number of plant species is not possible without accurate ecological studies. In addition, the information on plant size in general (and plant height in particular) appearing in the floristic catalogues is limited and most time very vague. For this reason, we decided to consider this variable as semi-quantitative, with three levels:

0 = This group includes plants with a prostrate life habit. Plants belonging to this group are those with a cushion shape, displaying flowers located very close to the ground and that thereby can be accessed both by flying and crawling insects (ants, springtails, mites, etc.).

1 = This group includes plants of intermediate size. We included in this group those plants shorter than 50 cm. This threshold is appropriate because it teases apart medium-sized species from those species with a large size. Many pollinators have a specific flight pattern with changes in flight zones occurring around this threshold. Within this group, there are also subshrub species with stunted growth habit.

2 = This group includes plants of large size. We included in this group those plants taller than 50 cm. These are plants particularly big, usually log-lived and sometimes woody species.

(2) Flower display size. The number of flowers produced per individual plant has strong direct and indirect effects on plant fitness in most Brassicaceae species. As occurring with plant height, the assessment of floral display size for a large number of plant species is not possible without accurate ecological studies. In addition, the information on flower number per individual appearing in the floristic catalogues is limited and most time very vague. For this reason, we decided to consider this variable as semi-quantitative, with three levels:

0 = This group includes species with few flowers per individual (pauciflorous), usually less than 50 flowers per individual.

1 = This group includes species with medium number to many flowers per individuals, usually between 50 and 1000 flowers per individual.

2 = This group includes species mass-flowering species, usually with more than 1000 flowers per individual.

(3) Inflorescence architecture. The configuration of flowers along the flowering stems and the inflorescence architecture have been shown to affect the attractiveness and foraging behaviour of pollinators in many angiosperm groups since long time. In Brassicaceae three main types of inflorescences can be distinguished:

0 = Inflorescences where flowers are arranged in solitary. In this species, flowers do not form a dense inflorescence but are solitary usually at the end of the flowering stems.

1 = Inflorescences where flowers are arranged in racemes. A simple inflorescence in which the main axis is indeterminate. This is the most frequent type of inflorescence in Brassicaceae.

2 = Inflorescences where flowers are arranged in corymbs. This is a special case of a panicle where flowers lie in a single plane. Panicles are determinate compound inflorescences in which branching does not occur from the axils of prophylls.

(4) **Presence of apetalous flowers.** Several species from some Brassicaceae genera, especially *Lepidium* and *Rorippa*, and to a lesser extent *Romanschulzia*, *Clypeola*, *Cardamine* and other minor genera, produce flowers without petals. We classified this floral trait as presence (1) or absence (0) of apetalous flowers.

(5) **Number of symmetry axes of the corolla.** Flower symmetry is an important trait in flowering plants. The Brassicaceae flower is defined as a cruciform, actinomorphic or radial flowers with many symmetry axes. However, it is widely acknowledged that some genera such as *Iberis* or *Teesdalia* produce monomorphic or actinomorphic flowers. The number of symmetry axes is even greater in some species. We have distinguished four groups based on number of symmetry axes:

0 = This group includes plants with flowers having no symmetry axis, like many species of *Matthiola*, some *Hesperis*,

1 = This group includes plants bearing flowers with one symmetry axis or actinomorphic flowers. In this group we included *Iberis*, *Teesdalia*, and several species of *Noccaea*, *Thlaspi*, etc.

2 = This group includes plants bearing flowers with two symmetry axes or dissymmetric flowers. This is probably the most abundant group, including most common species of Brassicaceae, like *Erysimum*, *Brassica*, *Diplotaxis*, etc.

4 = This group includes plants bearing flowers with four or more symmetry axes or polysymmetric flowers. This group, including common species of Brassicaceae, like *Lepidium*, some *Erysimum*, many *Heliophila*, many *Sisymbrium*, etc.

(6) **Orientation of dominant symmetry axis of the corolla.** In Brassicaceae, most flowers orientate vertically. Thereby, we classified this floral trait as horizontally- (1) or vertically- (0) orientated flowers.

(7) **Corolla with overlapped petals.** Much like flower symmetry, the presence of overlapped petals and rounded corollas affect fitness in several plant groups, including some Brassicaceae species by mediating the attractiveness of the flowers and the behaviour of pollinators. We classified this floral trait as corolla with overlapped petals (1) or with non-overlapped petals (0).

(8) **Corolla with multilobed petals.** In Brassicaceae petal lobes is not widespread, although it is frequent in some clades such as *Schizopetalon*, *Berteroa*, *Dryopetalon*. We classified this floral trait as corolla with multilobed petals (1) or without them (0).

(9) **Corolla with visible sepals.** Sepals play an important role in the pollination of many plant species. Some plant species, including Brassicaceae, have extended sepals that are visible from the top of the corolla. These visible petals may have important consequences on the behaviour of some pollinators, indirectly influencing the pollination success of the

flower. We scored this floral trait as corolla with visible sepals from the top of the corolla (1) or not (0).

(10) **Petal length.** Different studies have found a significant association between the length of flower petals and the behaviour of pollinators, by increasing corolla size and attractiveness or the floral attraction surface. As a consequence, it has been frequently proven the occurrence of a significant effect of petal length and flower size on the efficiency of pollination. We included in the data set the length of the petal in mm of each plant species. For this, we retrieved from the literature the description of the petal length, and calculated the mean of the values appearing in that description.

(11) **Sepal length.** In Brassicaceae the length of the sepals is positively correlated with the length of the corolla tube and the amount of nectar produced by the flowers. We included in the data set the length of the sepals in mm of each plant species. As in traits 10, we retrieved from the literature the description of the sepal length, and calculated the mean of the values appearing in that description.

(12) **Asymmetric petals.** Brassicaceae is characterized for bearing four symmetric petals. However, some species exhibit corollas with asymmetric petals, a character considered a morphological novelty. Presence of asymmetric petals causes corollas to show zygomorphy. This character, by affecting in an extreme way the number of symmetry axes, have large effects on pollinator preference, pollination efficiency and reproduction success. We scored this floral trait as corolla with asymmetric petals (1) or not (0).

(13) **Petal limb length.** The limb of the petal is the showy part that directly attracts pollinators. We included in the data set the length of the petal limb in mm of each plant species. For this, we retrieved from the literature the description of the petal length, and calculated the mean of the values appearing in that description.

(14) **Length of long stamens.** Brassicaceae has a tetradynamous androceum, with an outer whorl of two short stamens and an inner whorl of four long stamens. The length of the long stamens has been proven to affect pollinator visitation rate and effectiveness, having a strong effect on pollen removal and male fitness. We included in the data set the length of the long stamen in mm of each plant species as appearing in the literature.

(15) **Length of short stamens.** Short stamens may function in outcrossing Brassicaceae to reduce pollen depletion with high rates of pollinator visitation. In self-compatible, short stamens may favour delayed autogamy. In addition, short stamens may also affect pollinator visitation rate and effectiveness, having potential effect on pollen removal and male. We included in the data set the length of the short stamen in mm of each plant species as appearing in the literature.

(16) **Stamen dimorphism.** The difference in length between long and short stamens, hereinafter herkogamy, is related in Brassicaceae with pollinator attraction and evolution of selfing syndrome. We included this trait by estimating the length difference between long and short stamens from the data obtained in the literature.

(17) **Tetradynamous conditions.** In addition, we classified all Brassicaceae included in our dataset as having an androecium with all stamens equally long (0), slightly tetradynamous (1), normal tetradynamous condition (2) and strong tetradynamous condition (3). We used the classification appearing in the floral and formal description of the species.

(18) **Visible anthers.** Most species of Brassicaceae have anthers visible from outside the corolla during anthesis, which ease the magnitude of pollen removal by flower visitors. However, species of some genera (*Matthiola*, *Hesperis*, *Farsetia*, etc.) have stamens well hidden within the corolla tube and imperceptible from outside, a trait that difficult short-tongued insects to collect pollen. We scored this floral trait as corolla with visible anthers (1) or not (0).

(19) **Exserted stamens.** In some Brassicaceae the filaments are very long, causing stamens to be highly exserted. Stamens exsertion influences the behaviour and abundance of certain pollinators, shaping pollinator-mediated selection through male fitness. We scored this floral trait as non-exserted stamens (0) slightly exserted stamens (1) and strongly exserted stamens (2).

(20) **Number of stamens.** The basic number of stamens per Brassicaceae flower is six. However, departure from this number is frequent in some lineages such as *Lepidium* or to a lesser extent *Cardamine* or *Alyssum*, where some species bear 2, 4 or 5 stamens. In addition, some species of the genus *Megacarpa* have flowers with 9 or more stamens. We included for each species in the dataset the number of stamens indicated in the literature.

(21) **Concealed nectaries.** Some Brassicaceae species produce nectar that is concealed in the bottom of long corolla tubes, whereas other species bearing bowl-shaped flowers produce nectar that is freely exposed and easily accessible. This trait may have important consequences for the interaction with pollinators. We scored this floral trait as corolla with concealed nectaries (1) or not (0).

(22) **Petal carotenoids.** Flower colour is a crucial visual cue used by pollinators to locate flowers. In the Brassicaceae, there are numerous studies highlighting the role of flower colour in pollinator attraction and plant reproduction. Petal colour is mainly determined by the presence of pigments; we thereby decided to include the presence or absence of floral pigments in our dataset. Yellow colour is produced in Brassicaceae by the accumulation of carotenoids. We scored this trait as the presence (1) or absence (0) of petal carotenoids.

(23) **Petal anthocyanins.** In the Brassicaceae, species with pink, lilac, blue, purple, orange and red petals are caused by the accumulation of anthocyanins. We scored this floral trait as the presence of petal anthocyanins (1) or absence (0).

(24) **Presence of bullseyes.** Some flowers have circular patterns in the centre of the corolla called bullseyes that is involved in the attraction of pollinators. Bullseyes may be visible to human vision or invisible due to its absorbance in the ultraviolet region of the light spectrum; we considered only the first ones as is the information provide in the consulted Floras. We scored this floral trait as corolla with (1) or without (0) bullseyes.

(25) **Presence of veins in the petals.** In the Brassicaceae, some species may show petals with prominent veins having a different colour from the rest of the petals. The presence of coloured veins in the petals may function as nectar guides, providing visual orientation directing the pollinator to the central landing platform and the entrance to the flower. We scored this floral trait as petals with (1) or without (0) veins.

(26) **Coloured sepals.** As commented in the trait 9, sepals may be involved in pollination attraction in many species. Colouring sepals by accumulating anthocyanins or carotenoids and may help flowers to differentiate from the green background. We scored this floral trait as coloured sepals (1) or green sepals (0).

(27) **Relative attractiveness of petals versus sepals.** In some species of the Brassicaceae, the sepals are bigger and more attractive than the petals. This occurs frequently in some genera such as *Streptanthus*, *Roripa*, *Lepidium* and *Heliophila*. We scored this floral trait as (1) when petals are more attractive than sepals or (0) in the opposite case.

(28) **Petal hue.** Although measuring flower colour with spectrophotometric methods are recommended over methodologies based on human vision, obtaining reflectance data of more than 3000 species widely distributed around the world is virtually unfeasible. We designed a method that allows incorporating colour description in the Floras to generate categorical variables. We used a modification of colour identification with reference standards which are commonly used in comparative studies of flower colour and generates relatively good estimates of flower colour variation. First, we used a subset of 200 species that we have digital photos taken with the same camera and similar light conditions to prevent artificial colour modifications. The colour of petals was assigned to the closest matching Munsell colour chip; the same person performed these measures in order to avoid erroneous assignment due to inter-observer differences in colours perception. A total of 24 colour types were identified covering shades of blue (2.5P7/6, 10PB7/6), lilac-purple (7.5P8/4, 7.5P6/8, 7.5P6/10, 7.5P4/10, 5P6/8, 5P8/4, 5P5/10), pink (7.5RP8/4, 5RP6/10, 2.5RP5/10), yellow (5Y9/6, 5Y9/4, 5Y8.5/12), orange (5Y8/8, 2.5Y8/12, 2.5YR6/14), brown-bronze (10YR6/10, 5YR6/12, 10R5/8), green (2.5G5/5, 10GY6/8) and white (N9). We used spectral characteristics of Munsell colours to transform the categorical colour data to semi-quantitative measures of colour. Hue is one of the best colour descriptors for plant colourimetry; thus, we calculated hue values as the wavelength at peak reflectance. In order to accommodate the Brassicaceae petal colour information provided in the Floras to our 24 Munsell colour types, we generated ten colour categories. The hue of each new colour category was calculated as the mean of the hue values containing each category (i.e., among colour shades). In species with petal colour variation, including petal colour polymorphism, we scored the more common petal colour; if this information is not available, we assigned the colour derived of the presence of floral pigments (anthocyanins, carotenoids or both). The values of the ten hue categories are: 454.31 nm (blue), 503.55 nm (pink), 558.08 nm (lilac-purple), 572.46 nm (yellow), 575.43 nm (pale yellow), 579.38 nm (yellow-orange), 592.74 nm (orange), 589.44 nm (brown-bronze), 546.10 nm (green) and 611.37 nm (white).

(29) **Petal colour as b CIELAB.** We also used a second parameter related to petal colour, the “b*” parameter of the CIE 1976 L*a*b*. In this colour space, b* dimension represent values from -100 (blue colours) to 100 (yellow colours). This metrics is recommendable for the analysis of flower colour, particularly in groups of plant species containing petals with shades of yellow, as occurs in the Brassicaceae. b* values were obtained with the same methodology explained in the previous trait (28). The values of the ten b* categories are: -18.46 (blue), -4.77 (pink), -19.71 (lilac-purple), 45.03 (pale yellow), 80.1 (yellow), 80.45 (yellow-orange), 65.3 (orange), 52.02 (brown-bronze), 29.79 (green) and 0.00 (white).

(30) **Sepal hue.** Sepals of Brassicaceae species are sometimes coloured, differing from the common green. As already mentioned above for traits 9, sepals play an important role in the pollination of many plant species. We used the same method and hue values detailed in the trait 28 to score the sepal colour as hue category.

(31) **Sepal colour as b CIELAB.** For the same reasons mentioned above, we decided to include this trait because of the effect it can have on attracting pollinators. We used the same method and values detailed in the trait 29 to score the sepal colour as “b*” parameter of the CIE 1976 L*a*b*.

Table S9. List of the phylogenies retrieved from the online repositories and from the literature to built up the Brassicaceae supertree. Within brackets appears the number of species included in the analysis of disparity

Code	Species	Dated	Rooted	Focal taxa	Reference
Phylogenies including Moricandia					
1	15 [8]	YES	YES	Moricandia	Perfectti, F., Gómez, J. M., González-Megías, A., Abdelaziz, M., & Lorite, J. (2017). Molecular phylogeny and evolutionary history of Moricandia DC (Brassicaceae). <i>PeerJ</i> , 5, e3964.
2	273 [255]	YES	YES		Smith, S. A., & Brown, J. W. (2018). Constructing a broadly inclusive seed plant phylogeny. <i>American journal of botany</i> , 105(3), 302-314
3	1508 [248]	YES	YES		Gaynor, M. L., Ng, J., & Laport, R. G. (2018). Phylogenetic structure of plant communities: are polyploids distantly related to co-occurring diploids?. <i>Frontiers in Ecology and Evolution</i> , 6, 52.
4	195 [163]	YES	YES	Brassicaceae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
Time-calibrated phylogenies with more than 45 spp					
5	84 [48]	YES	YES	Euclidieae	Chen, H., German, D. A., Al-Shehbaz, I. A., Yue, J., & Sun, H. (2020). Phylogeny of Euclidieae (Brassicaceae) based on plastome and nuclear ribosomal DNA data. <i>Molecular Phylogenetics and Evolution</i> , 153, 106940
6	130 [124]	YES	YES		Durka, W., & Michalski, S. G. (2012). Daphne: a dated phylogeny of a large European flora for phylogenetically informed ecological analyses: <i>Ecological Archives</i> E093-214. <i>Ecology</i> , 93(10), 2297-2297
7	316 [208]	YES	YES		Walden, N., German, D. A., Wolf, E. M., Kiefer, M., Rigault, P., Huang, X. C., Kiefer, C., Schmickl R., Franzke A., Neuffer B., Mummenhoff, K., & Koch, M.A. (2020). Nested whole-genome duplications coincide with diversification and high morphological disparity in Brassicaceae. <i>Nature communications</i> , 11(1), 1-12
8	165 [109]	YES	YES	Alysseae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
9	46 [26]	YES	YES	Anchonieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
10	265 [265]	YES	YES	Arabidae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
11	84 [77]	YES	YES	Boechereae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
12	160 [126]	YES	YES	Cardamineae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
13	57 [23]	YES	YES	Chorisporae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
14	51 [28]	YES	YES	Coluteocarpaceae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
15	110 [89]	YES	YES	Erysimeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
16	75 [55]	YES	YES	Euclidieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
17	56 [53]	YES	YES	Heliophilleae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
18	139 [94]	YES	YES	Lepidieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
19	130 [117]	YES	YES	Thelypodieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
Time-calibrated phylogenies with less than 45 spp					
20	10	YES	YES	Aethionemeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.

21	9	YES	YES	Alyssopsidaeae	events. <i>Annals of Botany</i> , 125(1), pp.29-47. Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
22	30	YES	YES	Anastatiaceae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
23	10	YES	YES	Aphragmeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
24	5	YES	YES	Asteae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
25	19	YES	YES	Biscutelleae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
26	6	YES	YES	Buniadeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
27	8	YES	YES	Calepineae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
28	27	YES	YES	Camelineae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
29	16	YES	YES	Cochleariaeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
30	8	YES	YES	Conringieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
31	29	YES	YES	Cremolobeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
32	13	YES	YES	Crucihimalayaeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
33	41	YES	YES	Descurainieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
34	17	YES	YES	Dontostemoneae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
35	24	YES	YES	Eudemeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
36	25	YES	YES	Eutremeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
37	23	YES	YES	Halimolobeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
38	11	YES	YES	Hesperideae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
39	13	YES	YES	Hillilleae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
40	6	YES	YES	Iberideae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
41	34	YES	YES	Isatideae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
42	5	YES	YES	Kernereae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
43	5	YES	YES	Malcolmieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
44	8	YES	YES	Megacarpaeae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in

					Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
45	26	YES	YES	Microlepidieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
46	4	YES	YES	Notothlaspidieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
47	5	YES	YES	Oreophytoneae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
48	40	YES	YES	Physarieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
49	20	YES	YES	Schizopetaleae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
50	23	YES	YES	Sisymbrieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
51	23	YES	YES	Smelowskieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
52	19	YES	YES	Thlaspidieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
53	5	YES	YES	Turritideae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
54	8	YES	YES	Yinshanieae	Huang, X.C., German, D.A. and Koch, M.A., 2020. Temporal patterns of diversification in Brassicaceae demonstrate decoupling of rate shifts and mesopolyploidization events. <i>Annals of Botany</i> , 125(1), pp.29-47.
Non-time calibrated phylogenies					
55	115	NO	YES		Gómez, J. M., Torices, R., Lorite, J., Klingenberg, C. P., & Perfectti, F. (2016). The role of pollinators in the evolution of corolla shape variation, disparity and integration in a highly diversified plant family with a conserved floral bauplan. <i>Annals of Botany</i> , 117(5), 889-904.
56	44	NO	NO	Erysimum	Gómez, J. M., Perfectti, F., Abdelaziz, M., Lorite, J., Muñoz-Pajares, A. J., & Valverde, J. (2015). Evolution of pollination niches in a generalist plant clade. <i>New Phytologist</i> , 205(1), 440-453.
57	569	NO	YES		Couvreux, T. L., Franzke, A., Al-Shehbaz, I. A., Bakker, F. T., Koch, M. A., & Mummenhoff, K. (2010). Molecular phylogenetics, temporal diversification, and principles of evolution in the mustard family (Brassicaceae). <i>Molecular Biology and Evolution</i> , 27(1), 55-71.
58	115	NO	NO		Salariato, D. L., Manchego, M. A. C., Cano, A., & Al-Shehbaz, I. A. (2019). Phylogenetic placement of the Peruvian-endemic genus <i>Machaerophorus</i> (Brassicaceae) based on molecular data and implication for its systematics. <i>Plant Systematics and Evolution</i> , 305(1), 77-87.
59	53	NO	YES		Guo, X., Liu, J., Hao, G., Zhang, L., Mao, K., Wang, X., ... & Koch, M. A. (2017). Plastome phylogeny and early diversification of Brassicaceae. <i>BMC genomics</i> , 18(1), 176.
60	60	NO	YES	Thysanocarpus	Alexander, P. J., Windham, M. D., Govindarajulu, R., Al-Shehbaz, I. A., & Bailey, C. D. (2010). Molecular phylogenetics and taxonomy of the genus <i>Thysanocarpus</i> (Brassicaceae). <i>Systematic Botany</i> , 35(3), 559-577.
61	56	NO	YES		Huang, C.H., Sun, R., Hu, Y., Zeng, L., Zhang, N., Cai, L., Zhang, Q., Koch, M.A., Al-Shehbaz, I., Edger, P.P. and Pires, J.C., 2016. Resolution of Brassicaceae phylogeny using nuclear genes uncovers nested radiations and supports convergent morphological evolution. <i>Molecular biology and evolution</i> , 33(2), pp.394-412.
62	186	NO	YES		Warwick, S. I., Mummenhoff, K., Sauder, C. A., Koch, M. A., & Al-Shehbaz, I. A. (2010). Closing the gaps: phylogenetic relationships in the Brassicaceae based on DNA sequence data of nuclear ribosomal ITS region. <i>Plant Systematics and Evolution</i> , 285(3-4), 209-232.
63	101	NO	YES		Arias, T., Beilstein, M. A., Tang, M., McKain, M. R., & Pires, J. C. (2014). Diversification times among Brassica (Brassicaceae) crops suggest hybrid formation after 20 million years of divergence. <i>American journal of botany</i> , 101(1), 86-91.
64	27	NO	NO	Microthlaspi	Ali, T., Schmuker, A., Runge, F., Solovyeva, I., Nigrelli, L., Paule, J., Buch, A.K., Xia, X., Ploch, S., Orren, O. and Kummer, V., 2016. Morphology, phylogeny, and taxonomy of Microthlaspi (Brassicaceae: Coluteocarpeae) and related genera. <i>Taxon</i> , 65(1), 79-98.
65	22	NO	YES	Alysseae	Cecchi, L., Gabbriellini, R., Arnetoli, M., Gonnelli, C., Hasko, A., & Selvi, F. (2010). Evolutionary lineages of nickel hyperaccumulation and systematics in European Alysseae (Brassicaceae): evidence from nrDNA sequence data. <i>Annals of Botany</i> , 106(5), 751-767.
66	53	NO	YES		Soza, V. L., & Di Stilio, V. S. (2014). Pattern and process in the evolution of the sole dioecious member of Brassicaceae. <i>EvoDevo</i> , 5(1), 42.
67	38	NO	YES	Descurainia	Goodson, B. E., Rehman, S. K., & Jansen, R. K. (2011). Molecular systematics and

					biogeography of Descurainia (Brassicaceae) based on nuclear ITS and non-coding chloroplast DNA. Systematic Botany, 36(4), 957-980.
68	15	NO	NO	Thlaspi	Koch, M., & Al-Shehbaz, I. A. (2004). Taxonomic and phylogenetic evaluation of the American. Systematic Botany, 29(2), 375-384.
69	101	NO	NO		From TreeBase - d13 [R-package APE, Fri May 31 09:08:01 2019]
70	130	NO	NO		Salariato, D. L., Manchego, M. A. C., Cano, A., & Al-Shehbaz, I. A. (2019). Phylogenetic placement of the Peruvian-endemic genus Machaerophorus (Brassicaceae) based on molecular data and implication for its systematics. Plant Systematics and Evolution, 305(1), 77-87.
71	103	NO	YES		From TreeBase - T3061 [R-package APE, Thu Oct 15 18:34:08 2020]
72	56	NO	YES		From TreeBase - Parrya [R-package APE, Thu Oct 15 19:34:09 2020] - Nikolov, L.A., Shushkov, P., Nevado, B., Gan, X., Al-Shehbaz, I.A., Filatov, D., Bailey, C.D. and Tsiantis, M., 2019. Resolving the backbone of the Brassicaceae phylogeny for investigating trait diversity. New Phytologist, 222(3), pp.1638-1651.
73	223	NO	YES		From TreeBase - varios [R-package APE, Fri Oct 16 07:38:56 2020]
74	97	NO	NO	Vella	Simon-Porcar, V. I., Perez-Collazos, E., & Catalan, P. (2015). Phylogeny and systematics of the western Mediterranean Vella pseudocytisus-V. aspera complex (Brassicaceae). Turkish Journal of Botany, 39(3), 472-486.
75	109	NO	NO	Vella	Crespo, M.B., Lledó, M.D., Fay, M.F. and Chase, M.W., 2000. Subtribe Vellinae (Brassicaceae, Brassicaceae): a combined analysis of ITS nrDNA sequences and morphological data. Annals of Botany, 86(1), pp.53-62.
76	49	NO	YES	Pachycladon	Joly, S., Heenan, P.B. and Lockhart, P.J., 2009. A Pleistocene inter-tribal allopolyploidization event precedes the species radiation of Pachycladon (Brassicaceae) in New Zealand. Molecular phylogenetics and evolution, 51(2), pp.365-372.
77	189	NO	YES		German, D.A., Friesen, N., Neuffer, B., Al-Shehbaz, I.A. and Hurka, H., 2009. Contribution to ITS phylogeny of the Brassicaceae, with special reference to some Asian taxa. Plant Systematics and Evolution, 283(1-2), pp.33-56.
78	195	NO	NO	Brassicaceae	BrassiBase ITS tree- https://brassibase.cos.uni-heidelberg.de/?action=phlv&subaction=Brassicaceae
79	598	NO	YES		Bailey, C.D., Koch, M.A., Mayer, M., Mummenhoff, K., O'Kane Jr, S.L., Warwick, S.I., Windham, M.D. and Al-Shehbaz, I.A., 2006. Toward a global phylogeny of the Brassicaceae. Molecular biology and evolution, 23(11), pp.2142-2160.
80	370	NO	YES		Friesen, N., Čalasan, A.Ž., Neuffer, B., German, D.A., Markov, M. and Hurka, H., 2020. Evolutionary history of the Eurasian steppe plant Schivereckia podolica (Brassicaceae) and its close relatives. Flora, p.151602.

Table S10. List of ecologists kindly sharing unpublished information on Brassicaceae pollinators. The host institutions are those at the time of the contact with our team.

Last Name	First Name	Host institution
Abdelaziz	Mohamed	University of Granada (Spain)
Aizen	Marcelo	Universidad Nacional del Comahue-CONICET (Argentina)
Aguado	Luis Oscar	Castilla y Leon Regional Government (Spain)
Alarcon	Ruben	University Arizona (USA)
Amat	Elena	Real Jardín Botánico de Madrid (Spain)
Arista	Montserrat	University of Seville (Spain)
Banza	Paula	University of Hull (UK)
Barbir	Jelena	ICA-CSIC (Spain)
Bartomeus	Ignasi	EBD-CSIC (Spain)
Bergerot	Benjamin	University of Rennes (France)
Bommarco	Riccardo	Swedish University of Agricultural Sciences (Sweden)
Bosch	Jordi	CREAF-UAB (Spain)
Bruinsma	Maaïke	Leiden University (The Netherlands)
Burkle	Laura	Montana State University (USA)
CaraDonna	Paul	Northwestern University (USA)
Cartar	Ralph	University of Calgary (Canada)
Castro	Silvia	University of Coimbra (Portugal)
Castro-Urgal	Rocio	IMEDEA-CSIC (Spain)
Chacoff	Natacha	Universidad Nacional del Comahue-CONICET (Argentina)
Conner	Jeffrey	Michigan State University (USA)
Cuerda	David	Junta de Andalucía (Spain)
Dennis	Roger L. H.	Staffordshire University (UK)
Ebeling	Anne	University of Jena (Germany)
Escudero	Adrián	Universidad Rey Juan Carlos (Spain)
Evans	Darren	University of Hull (UK)
Fernández	Juande	Greenpeace (Spain)
Ferrero	Victoria	University of León (Spain)
Fründ	Jochen	Georg-August-Universität (Germany)
Fultz	Jessica	Idaho State University (USA)
Garbuzov	Mihail	University Sussex (UK)
García	Begoña	IPE-CSIC (Spain)
García-Camacho	Raúl	Universidad Rey Juan Carlos (Spain)
García	Yedra	CIDE (University of New Brunswick)
García de Lucas	Sandra	Junta de Andalucía (Spain)
Giménez	Luis	Universidad Rey Juan Carlos (Spain)
Iriondo	José María	Universidad Rey Juan Carlos (Spain)
Junker	Robert R.	University of Salzburg (Austria)
Kuppler	Jonas	ULM University (Germany)
Lance	Richard	Northern Arizona University (USA)
Lara	Carlos	Universidad Rey Juan Carlos (Spain)
Lázaro	Amparo	IMEDEA-CSIC (Spain)
Lorite	Juan	University of Granada (Spain)
Louadi	Kamel	University Frères Mentouri Konstantine (Algeria)
Loureiro	João	University of Coimbra (Portugal)
Lucas-Barbosa	Dani	Wageningen University (The Netherlands)
Majetic	Cassey J.	Saint Mary's College Indiana (USA)
Marcos	Maria Ángeles	Universidad de Alicante (Spain)
Medel	Rodrigo	University of Santiago de Chile (Chile)
Meindl	George	Binghamton University (USA)
Melen	Miranda	University of California-Santa Cruz (USA)
Méndez	Marcos	Universidad Rey Juan Carlos (Spain)
Menéndez	Rosa	University of Lancaster (UK)

Milla	Rubén	Universidad Rey Juan Carlos (Spain)
Morales	Carolina	Universidad Nacional del Comahue-CONICET (Argentina)
Morente	Javier	Universidad Rey Juan Carlos (Spain)
Muñoz-Pajares	A. Jesus	University of Coimbra (Portugal)
Norfolk	Olivia	University of Nottingham (UK)
Norton	Nicholas	Washington State University (USA)
O'Malley	Rachel	San Jose State University (USA)
Ojeda	Fernando	University of Cádiz (Spain)
Pelayo	Roxibel	Universidad de las Andes (Venezuela)
Petanidou	Theodora	University of the Aegean (Greece)
Razanajatovo	Mialy	University Konstanz (Germany)
Roberts	S.P.M.	University of Reading (UK)
Santamaría	Silvia	Universidad Rey Juan Carlos (Spain)
Schlinkert	Hella	University Goettingen (Germany)
Schrader	Julian	University Goettingen (Germany)
Schupp	Eugene W.	Utah State University (USA)
Simaika	John P.	Stellenbosch University (South Africa)
Simanonok	Michael P.	MSU- Northern Prairie Wildlife Research Center (USA)
Stang	Martina	University of Leiden (The Netherlands)
Stanley	Dara A.	Trinity College Dublin (Ireland)
Stout	Jane	Trinity College Dublin (Ireland)
Strauss	Sharon	University of California at Davis (USA)
Torices	Rubén	University Lausanne (Switzerland)
Traveset	Anna	IMEDEA-CSIC (Spain)
Tscharntke	Teja	University of Göttingen (Germany)
Tur	Cristina	IMEDEA-CSIC (Spain)
Valido	Alfredo	IPNA-CSIC (Spain)
Valverde	Javier	EBD-CSIC (Spain)
Vargas	Pablo	Real Jardín Botánico de Madrid (Spain)
Warzecha	Daniela	Goethe University (Germany)
Whittall	Justen	Santa Clara University (USA)
Winfree	Rachael	Rutgers University (USA)
Wonneck	Mark	University of Calgary (Canada)
Zink	Lindsay	University of Calgary (Canada)

Table S11. Brief description of the functional groups of the insects visiting the flowers of the studied species.

	Functional Group	Body length	Resource	Behavioural notes	Type of visits	Order	Examples
1	Long-tongued extra-large bees	≥ 15 mm	Nectar + Pollen	Partially introducing the head in the flower	Legitimate	Hymenoptera	Anthophoridae, Apidae
2	Long-tongued large bees	10-15 mm	Nectar + Pollen	Partially introducing the head in the flower	Legitimate	Hymenoptera	Anthophoridae
3	Long-tongued medium-sized bees	< 10 mm	Nectar + Pollen	Partially introducing the head in the flower	Legitimate	Hymenoptera	Anthophoridae
4	Honeybees	6-12 mm	Nectar + Pollen	Introducing the whole head in the flower	Legitimate	Hymenoptera	Apidae (<i>Apis</i> spp.)
5	Short-tongued extra-large bees	≥ 15 mm	Nectar + Pollen	Introducing the head in the flower	Legitimate	Hymenoptera	Apidae
6	Short-tongued large bees	> 10 mm	Pollen + Nectar	Introducing the whole head in the flower	Legitimate	Hymenoptera	Halictidae, Megachilidae, Colletidae Andrenidae
7	Short-tongued medium-sized bees	5 – 10 mm	Pollen + Nectar	Introducing the whole head in the flower	Legitimate	Hymenoptera	Halictidae, Colletidae, Andrenidae , Apidae Xylocopinae, Apidae Nomidinae
8	Short-tongued small bees	2 – 5 mm	Pollen + Nectar	They access the nectar legitimately or from between the sepals	Illegitimate + Legitimate	Hymenoptera	Halictidae, Colletidae, Andrenidae , Apidae Xylocopinae, Apidae Nomidinae
9	Short-tongued extra-small bees	< 2 mm	Nectar + Pollen	They access the nectar legitimately or from between the sepals	Legitimate + Illegitimate	Hymenoptera	Halictidae, Colletidae
10	Large ants	> 2 mm	Nectar	They can introduce the whole body in the flower to reach the nectar	Legitimate + Illegitimate	Hymenoptera	Formicidae
11	Small ants	< 2 mm	Nectar	Mostly sipping nectar from between sepals	Illegitimate + Legitimate	Hymenoptera	Formicidae
12	Large pollen wasps	Variable	Pollen	Partially introducing the head in the flower	Legitimate	Hymenoptera	Massarinae
13	Large nectar-collecting wasps	> 7mm	Nectar	Partially introducing the head in the flower	Legitimate	Hymenoptera	Vespidae
14	Small nectar-collecting wasps	Usually < 3mm	Nectar	Mostly sipping nectar from between sepals	Illegitimate + Legitimate	Hymenoptera	Chalcidoidea, Ichneumonoidea
15	Hovering long-tongued	Variable	Nectar +	Hovering while nectaring and collecting some pollen	Legitimate	Diptera	Bombyliidae (<i>Bombylius</i>)

	flies		Pollen				
16	Non-hovering long tongued flies	Variable	Nectar	Nectaring without hovering; long buccal apparatus	Legitimate	Diptera	Bombyliidae, Tachinidae, Nemestrinidae,
17	Large hoverflies	>5 mm	Pollen	Collect pollen without entering the flower	Legitimate	Diptera	Syrphidae (Eristalini)
18	Small hoverflies	< 5 mm	Pollen + Nectar	Collect pollen without entering the flower and sometimes sip nectar from between the sepals	Legitimate + Illegitimate	Diptera	Syrphidae
19	Large flies	>5 mm	Nectar + Pollen	Collect pollen without entering the flower and nectar	Legitimate + Illegitimate	Diptera	Muscidae, Calliphoridae, Tabanidae, Scatophagidae, Anthomyiidae
20	Small flies	< 5 mm	Nectar + Pollen	Mostly sipping nectar	Illegitimate + Legitimate	Diptera	Muscidae, Anthomyiidae, Micetophyllidae, Drosophilidae, Stratiomyidae
21	Long tongued small flies	< 5 mm	Nectar	Sipping nectar	Illegitimate + Legitimate	Diptera	Bibionidae, Empididae
22	Large beetles	> 7 mm	Mostly Pollen	Consuming not only pollen, also anthers, petals, and other floral parts	Legitimate + Illegitimate	Coleoptera	Cetoniidae, Lagridae, Mylabridae, Alleculinae
23	Small beetles	< 7 mm	Pollen + Nectar	Consuming pollen during legitimate visits and also robbing nectar from the bottom part of the flowers	Legitimate + Illegitimate	Coleoptera	Melyridae (Malachidae, Dasytidae), Cleridae, Oedemeridae, Elateridae, Bruchidae, Buprestidae, Chrysomelidae
24	Small diving beetles	<3 mm	Nectar + Pollen	Entering completely into the flower, crawling down the corolla for nectar	Legitimate	Coleoptera	Nitidulidae, Dermestidae, Phalacridae
25	Large Butterflies	≥ 20 mm	Nectar	Feeding on nectar both from inside the flower and between the sepals	Legitimate	Lepidoptera	Nymphalidae, Papilionidae, Pieridae
26	Small Butterflies	< 20 mm	Nectar	Feeding on nectar both from inside the flower and between the sepals	Legitimate	Lepidoptera	Lycaenidae, Pieridae, Hesperidae
27	Hawkmoths	> 7 mm	Nectar	Hovering to sip nectar	Legitimate	Lepidoptera	Sphingidae
28	Large moths	> 3mm	Nectar	Sipping nectar while landed onto the corolla	Legitimate	Lepidoptera	Crambidae, Noctuidae
29	Small moths	< 3mm	Nectar	Sipping nectar without entering the flower	Illegitimate + Legitimate	Lepidoptera	Adelidae, Plutellidae
30	Nocturnal moths	variable	Nectar	Sipping nectar while landed onto the corolla or by hovering; Visiting the flowers at night	Legitimate	Lepidoptera	Noctuidae

31	Bugs	variable	Nectar	Sipping nectar without entering the flower. Also acting as sapsuckers in vegetative tissues	Legitimate + Illegitimate	Hemiptera	Lygaeidae, Pentatomidae
32	Thrips	< 3 mm	Pollen	Feeding from inside the flowers	Legitimate	Thysanoptera	
33	Grasshoppers	variable	Pollen + Floral parts	Mostly nymphs	Legitimate	Orthoptera	
34	Aphids	< 2 mm	Nectar	Mostly winged individuals	Legitimate	Hemiptera	Aphidoidea
35	Earwig	> 15 mm	Pollen		Legitimate + Illegitimate	Dermaptera	
36	Lacewing	> 15 mm	Pollen + Nectar		Legitimate + Illegitimate	Neuroptera	Chrysopidae
37	Snakeflies	> 8 mm	Pollen + Nectar		Legitimate + Illegitimate	Raphidioptera	
38	Birds	>>> 15 mm	Nectar		Legitimate	Passeriformes	
39	Springtails	< 2 mm	Nectar		Legitimate + Illegitimate		
40	Mites	< 2 mm	Nectar		Legitimate + Illegitimate		
41	Spiders	< 2 mm	Unknown		Illegitimate		
42	Larvae	variable	Unknown		Illegitimate		
43	Others	variable	Unknown		Illegitimate		

