

1 **Chronic disturbance of moist tropical forests favours deciduous over evergreen tree**
2 **communities across a climate gradient in the Western Ghats**

3

4 Nayantara Biswas^{1,*}, Vishal Sadekar¹, Siddharth Biniwale¹, Yukti Taneja^{1,2}, Anand M Osuri¹,
5 Navendu Page³, Kulbhushansingh Suryawanshi^{1,4}, and Rohit Naniwadekar^{1,*}

6

7 ¹ Nature Conservation Foundation, 1311, “Amritha”, 12th Main, Vijayanagar 1st Stage, Mysuru,
8 Karnataka 570017, India

9 ² Interdisciplinary Ecology, School of Natural Resources and Environment & Agronomy
10 Department, University of Florida, Gainesville, FL 32611, USA

11 ³ Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand 248001

12 ⁴ Snow Leopard Trust, 4649, Sunnyside Avenue North, Suite 325, Seattle, WA 98103, USA

13

14 **Corresponding authors:**

15 Nayantara Biswas (nayantara.b.1998@gmail.com) and Rohit Naniwadekar (rohit@ncf-india.org)

16 **Full Postal Address:** Nature Conservation Foundation, 1311, “Amritha”, 12th Main,
17 Vijayanagar 1st Stage, Mysuru, Karnataka 570017, India

18

19 **ABSTRACT**

20 It is well established that climatic factors such as water stress and chronic anthropogenic
21 disturbances such as biomass extraction influence tropical forest tree community structure,
22 richness and composition. However, while the standalone effects of these two drivers on plant
23 communities are well-studied, their interactive effects are not. Moist tropical forests in India’s

24 Western Ghats face a dual threat from increasingly erratic precipitation (and consequent water
25 stress), and an intensifying anthropogenic footprint. Here, we sampled 120 tree plots (0.05 ha
26 each) across forests with varying histories of biomass extraction and a gradient in climate water
27 deficit (CWD, a proxy for water stress) within a 15,000 km² landscape in the northern Western
28 Ghats and examined whether and how disturbance history modulates relationships of tree
29 community structure and composition with climate. As expected, tree species richness increased
30 with decreasing water stress in less- and historically-disturbed forests but remained low in
31 repeatedly-disturbed forests. The increase in evergreen species richness with decreasing water
32 stress was far slower in repeatedly-disturbed forests than other categories, and the relative
33 abundance of evergreens in the repeatedly-disturbed forests (25%) was half that in less-disturbed
34 forests (50%) of comparable water stress in the driest parts. Overall, we show that disturbance
35 can amplify threats from climate change to wet forest-associated evergreen tree species, many of
36 which are threatened, while benefiting more widely distributed dry forest-associated deciduous
37 species. In the northern Western Ghats, where much of the remaining forest cover is disturbed
38 and dominated by deciduous tree species, the persistence of evergreen tree flora hinges on
39 protecting existing evergreen forest patches from future disturbances and restoring locally
40 appropriate evergreen species in secondary forests.

41 INTRODUCTION

42 Plant community assembly is the culmination of abiotic and biotic factors acting over different
43 spatial and temporal scales (Ramos et al., 2023; Zheng et al., 2022). Although climate is the
44 major factor shaping plant diversity and composition (Harrison et al., 2020), in the
45 Anthropocene, human-mediated disturbances act in conjunction with such environmental factors
46 and contribute to altered tree richness, abundance and composition (Bentsi-Enchill et al., 2022),
47 with cascading impacts on ecosystem function (Hooper et al., 2012). However, the effects of
48 anthropogenic disturbances and environmental factors on plant community assembly are often
49 studied in isolation. Few studies have examined their combined effects on plant diversity and
50 structure, and those that have, are primarily focused on seasonally dry tropical forests (Ramos et
51 al., 2023; Rito et al., 2017; Zorger et al., 2019). Despite wet tropical forests (forests receiving >
52 2000 mm of annual rainfall) harbouring more than half of the global terrestrial biodiversity
53 (Malhi et al., 2014) and being increasingly threatened by habitat degradation and biodiversity
54 loss (Saatchi et al., 2021), studies on the interactive effects of anthropogenic and environmental
55 factors on plant communities have been overlooked in these sensitive ecosystems.

56 For plants, annual precipitation is an important factor influencing plant richness and
57 composition across large environmental gradients (Harrison et al., 2020; Krishnadas et al., 2016).
58 Besides annual precipitation, water stress during dry months is critical in determining species
59 distributions (Esquivel-Muelbert et al., 2017), thereby influencing vegetation types (Hirota et al.,
60 2011). On almost half of the Earth's terrestrial surface, climate change is expected to lead to less
61 water availability (McLaughlin et al., 2017), thereby greater water stress. Reduced water
62 availability may limit species distributions (Saiter et al., 2016) and induce deciduousness in the
63 vegetation, particularly in wetter, evergreen forests (Saiter et al., 2016; Seiler et al., 2014),

64 thereby driving tropical biome transitions, like those between wet and dry forests (Dexter et al.,
65 2018). These transition zones represent unstable states where the vegetation is most sensitive to
66 climatic disturbances (Hirota et al., 2011). Like climate, anthropogenic disturbances can mediate
67 shifts in vegetation types. Although deforestation is a major global threat to biodiversity, tropical
68 forests are increasingly imperilled by chronic anthropogenic disturbances like selective logging
69 and fuelwood extraction (Barlow et al., 2016). Despite research indicating that the interactive
70 effects of climatic and anthropogenic factors can have significant consequences on the resilience
71 of plant communities and affect vegetation types (Hirota et al., 2011; Shivaprakash et al., 2018),
72 studies explicitly testing for these effects are sparse. The need for assessing the interactions
73 between climate and anthropogenic factors has been highlighted (Krishnadas et al., 2021). While
74 some studies have investigated the interactive effects of climatic factors like precipitation and
75 chronic anthropogenic disturbances on plant diversity and structure (Rito et al. 2017; Ramos et
76 al. 2023), no studies have examined the effect of these interactive effects on vegetation
77 transitions. Although studies have suggested that fire mediates the transformation of forests to
78 drier vegetation types (savannization) (Sansevero et al., 2020; M. Wang et al., 2023), the
79 implications of chronic anthropogenic disturbances (hereafter, CAD) and environmental factors
80 on vegetation type transitions are less explored.

81 The Western Ghats-Sri Lanka biodiversity hotspot harbours a high diversity of vascular
82 plants, many of which are endemic to the region (Gunawardene et al., 2007). It has lost a
83 substantial extent of its original vegetation cover (<30% remaining) due to human activities
84 (Mittermeier et al., 2011; Myers et al., 2000). The region exhibits high levels of topographic and
85 climatic heterogeneity, creating suitable habitats to support diverse vegetation types along a
86 gradient of deciduous to evergreen forests, each of which harbours unique fauna (Gunawardene

87 et al., 2007). Woody plant diversity in the Western Ghats exhibits a clear latitudinal gradient
88 with decreasing plant richness in northern portions primarily driven by the past geo-climatic
89 history and niche conservatism (Gopal et al., 2023). Additionally, the topography of the
90 mountain ranges leads to a distinct gradient in rainfall from the low elevations to the mountain
91 crest, wherein precipitation increases towards the higher elevations (Venkatesh et al., 2021).
92 Distinct forest types are associated with different rainfall and elevation bands (Joseph et al.,
93 2012). These varying rainfall patterns give rise to water availability gradients, which are known
94 to be important drivers of tree diversity and structure (Terra et al., 2018) and are responsible for
95 a wide diversity of trees and vegetation types in the Western Ghats of India (Joseph et al., 2012;
96 Page & Shanker, 2018).

97 Despite the high biodiversity in this region, the forests in the Western Ghats are
98 threatened by a range of anthropogenic disturbances that have led to the loss of 35.3% of its
99 forest cover from the 1920s to 2013 (Reddy et al., 2016). The forests here are at an elevated risk
100 of degradation as they have the third highest human population density among all global hotspots
101 (Cunningham & Beazley, 2018). This is particularly true for the northern Western Ghats, where
102 the privately-owned forests are periodically clear-felled (every ten years) to harvest fuelwood,
103 resulting in CAD. Apart from the repeatedly disturbed privately-owned forests, the northern
104 Western Ghats also harbour government-owned Reserved Forests and Protected Areas, which
105 were historically disturbed. There are also sacred groves, many of which harbour among the least
106 disturbed forest habitats in the region. These disturbance and water availability gradients in the
107 northern Western Ghats make it a suitable system to study the interplay of climate and CAD on
108 plant vegetation types.

109 Given this background, we investigated the effects of CAD and climate on the diversity
110 and structure of plant communities. To do this, we sampled 120 vegetation plots across three
111 forest categories with varying degrees of protection and along a precipitation gradient. We
112 specifically asked the following questions: (1) How does tree composition differ across forest
113 categories? (2) How do climatic water deficit (CWD) and CAD impact the vegetation structure
114 and overall tree species richness? (3) How do CWD and CAD affect the proportion of evergreen
115 trees and richness of evergreen and deciduous trees? We expected that filtering mediated by
116 human disturbance and climate would result in differences in tree composition and diversity. We
117 expected a higher proportion and richness of evergreen trees to be found in the wetter (low
118 CWD) and cooler regions in the high-elevation forests, as compared to the drier (high CWD),
119 low-elevation forests. Moreover, we predicted that repeated chronic anthropogenic disturbance
120 would further lower evergreen tree richness and proportion, leading to a more deciduous
121 vegetation type.

122

123 **METHODS**

124 **Study Area**

125 We conducted the study in the south-western part of Maharashtra in India (15.72–17.74°N;
126 73.29–74.19°E) between October 2022 and March 2023. The study site forms the northern part
127 of the Western Ghats-Sri Lanka Biodiversity Hotspot, which is classified as one of the “hottest”
128 biodiversity hotspots globally, based on the high degree of endemism and levels of
129 anthropogenic threats (Myers et al. 2000). The climate here is tropical, with average annual
130 rainfall ranging from 2150–7450 mm, and the annual temperature varying between 16–35°C (Jog
131 2009).

132 We sampled sites experiencing varying levels of disturbance across an elevational
133 gradient, ranging from 8 to 1054 m a.s.l. (Fig. S1). The sites were classified as less-disturbed,
134 historically-disturbed and repeatedly-disturbed. For the less-disturbed sites, we sampled the
135 sacred groves. The landscape harbours several sacred groves, locally known as “*Devrais*”, which
136 are protected by the local villages and home to relatively less disturbed patches of forests (Gadgil
137 & Vartak, 1976). They provide refuge to many endemic, evergreen and medicinal plants
138 (Blicharska et al., 2013; Kulkarni et al., 2018). Since some sacred groves experience
139 opportunistic fuelwood and timber collection, these sites were classified as less-disturbed. We
140 classified the government-managed forests (Protected Areas (PAs) and Reserved Forests (RFs))
141 as historically disturbed sites. The PA sites were spread across the Sahyadri Tiger Reserve and
142 Radhanagari Wildlife Sanctuary, and the RF sites were spread across Chiplun and Sawantwadi
143 Forest Divisions. The PAs were designated in the late 1950s (Radhanagari) and mid-1980s
144 (Sahyadri), and these sites have a long human-use history for shifting cultivation and have likely
145 been clear-felled in the past (Chandran, 1997; Ghate et al., 1998), unlike the sacred groves. The
146 existing stunted evergreen forests here result from forest recovery after clear felling (Ghate et al.
147 1998). Since there are no PAs in lower elevations, all the PA sites were located in relatively
148 higher elevations (584–1012 m a.s.l.). The elevation range of RF sites ranged between 26–386 m
149 a.s.l. While no form of resource utilisation by humans is permitted in PAs, some forms of
150 resource use, like cattle grazing and deadwood collection, are still allowed in RFs. We classified
151 the private forests as repeatedly-disturbed forests. The private forests sampled in the region
152 experience the highest level of chronic anthropogenic disturbance among the three forest
153 categories as people clear fell these forests every five to ten years to sell fuelwood (Kulkarni and
154 Mehta, 2013). The prevailing forest type here has been reported to be tropical moist deciduous in

155 the lower elevations, and semi-evergreen to evergreen in the higher elevations, with stunted
156 evergreen trees being present after recovery from clear-felling (Champion & Seth, 1968; Ghate
157 et al. 1998). The dominant tree families in the high-elevation PAs are Melastomataceae and
158 Myrtaceae (Joglekar et al., 2015; Kanade et al., 2008). While there is a considerable amount of
159 information from the higher-elevation protected and non-protected forests (Joglekar et al. 2015;
160 Kulkarni et al. 2018), there is very little information on vegetation type and composition in the
161 low-elevation forests of the northern Western Ghats.

162 **Vegetation Sampling**

163 We conducted the field study in an approximately 15,000 km² landscape in the northern Western
164 Ghats (Fig. S1). To capture the gradients of CAD and climate, we laid 120 50×10m² plots that
165 were evenly distributed across less-disturbed sacred groves (n=40), historically-disturbed
166 government-owned forests (Reserved Forests and Protected Areas) (n=40), and repeatedly-
167 disturbed privately-owned forests (n=40), across low and high elevations. Henceforth, we will
168 refer to these three categories of forests as less-disturbed, historically-disturbed and repeatedly-
169 disturbed forests. In each plot, we recorded the identity, girth and height of all woody stems ≥10
170 cm Girth at Breast Height (GBH). We also recorded the number of cut stems in each plot as an
171 indicator of human disturbance. All trees were classified as evergreen or deciduous using
172 regional floras and expertise within the team (NP). We excluded climbers from further analyses.

173 **Analysis**

174 All the analysis was carried out in R ver. 4.3.1(R Core Team, 2023)

175 **Tree composition across land-use types**

176 To find out the difference in the composition of trees across the six forest categories in low- and
177 high-elevations, we used 3-D non-metric multidimensional scaling (NMDS) with Bray-Curtis

178 dissimilarity metric using the R package ‘vegan’ (Oksanen et al., 2022) and ‘vegan3d’ (Oksanen
179 et al., 2018). We used 3-D NMDS since 2-D NMDS had stress values greater than 0.2. The
180 difference in species composition among the categories was tested using the ‘ANOSIM’ or
181 analysis of similarities function in the R package ‘vegan’, along with a permutation test
182 (permutations = 999).

183 **Influence of CAD and CWD on species richness**

184 We examined the influence of climate and CAD on the species richness per plot. We used the
185 predictor, CWD, to account for the climatic gradient across our study site. We obtained the CWD
186 values for each vegetation plot from a global gridded layer available at [http://chave.ups-](http://chave.ups-tlse.fr/pantropical_allometry.htm#CWD)
187 [tlse.fr/pantropical_allometry.htm#CWD](http://chave.ups-tlse.fr/pantropical_allometry.htm#CWD) (Chave et al., 2014). CWD is a measure of the water
188 stress experienced by plants in the dry months. It is measured as the difference between
189 precipitation and evapotranspiration in dry months when evapotranspiration exceeds
190 precipitation (Krishnadas et al. 2021). The values are always negative, and a less negative CWD
191 indicates a higher water availability for plants. It has been known to play a key role in assessing
192 the sensitivity of vegetation to drought (Vicente-Serrano et al., 2013) and transitions of
193 evergreen forests to dry deciduous forests in Bolivia (Seiler et al., 2014). In the Western Ghats
194 too, studies have shown CWD to influence plant diversity (Krishnadas et al., 2021, Gopal et al.,
195 2023); however, the interactive effects between CWD and CAD are not known. In our area of
196 interest, elevation and CWD are strongly positively correlated (Spearman’s $\rho = 0.79$; $p < 0.001$),
197 indicating that high-elevation sites had lower water deficit than low-elevation sites. We used
198 GLMs to evaluate the combined (interactive) effect of CWD and forest category on the observed
199 species richness per plot with a Poisson error structure (since the response variable was not over-
200 dispersed).

201 **Influence of CAD and CWD on vegetation structure**

202 We estimated the basal area of trees ($\text{m}^2 \text{ha}^{-1}$). We square root transformed the basal area since it
203 was not normally distributed (Shapiro-Wilk's Test; $p < 0.05$). We used the General Linear Model
204 (Gaussian error structure) to test the relationship between basal area, disturbance categories, and
205 CWD.

206 **Influence of CAD and CWD on the proportion of evergreen trees and richness of evergreen 207 and deciduous trees**

208 We examined the influence of CWD and CAD on the proportion of evergreen trees and richness
209 of evergreen and deciduous trees per plot. We used GLMs to evaluate the combined (interactive)
210 effect of CWD and forest category on the proportion of evergreen trees with a binomial error
211 structure and the richness of evergreen and deciduous trees with a Poisson error structure.

212 **RESULTS**

213 We identified 97% of the 7001 (≥ 10 cm GBH) individual trees across our 120 plots. Two
214 individuals were identified till the genus level. Six individuals that could not be identified were
215 excluded from further analyses. We recorded 192 plant species (166 trees) from 52 families
216 (Table S1). The most speciose families were Fabaceae and Moraceae, accounting for 14.6% of
217 all species. We recorded a total of 10,071 stems. The mean (\pm SE) stem density was 1678.5
218 (± 79.8) stems/ha, and the basal area was 32 (± 1.7) m^2/ha . A summary of tree richness is
219 presented in Figure S3.


220 **Woody plant composition across land-use types**

221 The 'ANOSIM' analysis revealed significant differences in tree composition among the six
222 different forest categories ($R_{anosim} = 0.44$, $p = 0.001$, stress = 0.17) (Fig. S2). The NMDS plots
223 suggested that the tree composition in high-elevation forests tended to be different than low-

224 elevation ones (Fig. S2). Even at a particular elevation-level, tree composition differed across
225 disturbance categories.

226 **Influence of CAD and CWD on species richness**

227 The GLM result showed that the interaction between CWD and disturbance category had a
228 significant effect on the species richness of trees (Fig. 1; Table S2). The observed species
229 richness per plot increased with decreasing CWD for less-disturbed sites but was lower and
230 tended to decrease with decreasing CWD for repeatedly-disturbed sites (Fig. 1; Table S2).




231
232 **Figure 1.** Plots showing the relationship between climatic water deficit and observed species
233 richness per plot across the three disturbance categories. The solid line with a band indicates the
234 best-fitted line for the Generalized Linear Model (Poisson error structure) with associated 95%
235 CI. Higher values of CWD (less negative) imply lower water stress experienced by plants.

236

237 **Role of disturbance in influencing forest structure**

238 The interaction between disturbance categories and CWD was significant for the basal area ($R^2 =$
239 0.51; Fig. 2; Table S3). Basal area increased with CWD for historically-disturbed and less-
240 disturbed sites but decreased with CWD for the repeatedly-disturbed site (Fig. 2). This indicates
241 that the trees are larger in wetter regions when chronic anthropogenic disturbance is lower.
242 However, in chronically-disturbed sites trees are smaller in wetter regions. A summary of basal
243 area, tree density, and canopy cover is provided in Table S4.




245 **Figure 2.** Plots showing the relationship between climatic water deficit and square root of basal
246 area per hectare, across the three disturbance categories. The solid line with a band indicates the
247 best-fitted line for the Generalized Linear Model (Gaussian error structure) with associated 95%
248 CI. Higher values of CWD (less negative) imply lower water stress experienced by plants.


249

250 **Influence of CAD and CWD on the proportion of evergreen trees and richness of evergreen**
251 **and deciduous trees**

252 The GLM results showed that the interaction terms between CWD and disturbance category
253 significantly affected the proportion and richness of evergreen trees. The proportion of evergreen
254 trees per plot increased with the increase in CWD values (which implies less water deficit) and
255 this increase was the fastest in less-disturbed forests and slowest in repeatedly-disturbed forests
256 (Pseudo $R^2 = 0.38$, Fig. 3; Table S5). In regions with low CWD values (indicating higher water
257 deficit i.e., drier areas), the predicted proportion of evergreen trees is more than 0.5 for less-
258 disturbed forests but only 0.25 for repeatedly disturbed forests, indicating dominance of
259 deciduous trees in repeatedly-disturbed sites (Fig. 3). In regions with high CWD values
260 (indicating lower water deficit i.e., wetter areas) the predicted proportion of evergreen trees for
261 less-disturbed forests was close to 1 but for repeatedly-disturbed forests it was around 0.75.
262 Similarly, the increase in the number of evergreen tree species per plot with decreasing CWD
263 was very fast for less- and historically-disturbed sites but gradual for the repeatedly-disturbed
264 sites, indicating that disturbance negatively impacted evergreen tree species richness (Pseudo R^2
265 = 0.33, Fig. 4A; Table S6). Consequently, the deciduous species richness per plot rapidly
266 decreased with decreasing CWD for less-disturbed and historically-disturbed sites but this
267 decrease was significantly more gradual in repeatedly disturbed sites (Pseudo $R^2 = 0.36$, Fig. 4B;
268 Table S7).



270 **Figure 3.** Plots showing the relationship between climatic water deficit and proportion of
271 evergreen tree individuals per plot across the three disturbance categories. The solid line with a
272 band indicates the best fitted line for the Generalized Linear Model (binomial error structure)
273 with associated 95% CI. Higher values of CWD (less negative) imply lower water stress
274 experienced by plants.



275

276 **Figure 4.** Plots showing the relationship between climatic water deficit and number of evergreen
277 tree species per plot (A) and number of deciduous tree species per plot (B) across the three
278 disturbance categories. The solid line with a band indicates the best-fitted line for the

279 Generalized Linear Model (Poisson error structure) with associated 95% CI. Higher values of
280 CWD (less negative) imply lower water stress experienced by plants.

281

282 **DISCUSSION**

283 While previous studies have mostly examined the effects of CAD and climate on vegetation in
284 isolation, we examined the interactive effects of the two drivers on diversity, structure and
285 composition of trees in a biodiversity hotspot. Our results demonstrate that CAD and CWD
286 significantly influence woody plant composition, overall tree species richness and structure. We
287 found that CAD disrupts the relationship between CWD and vegetation type, resulting in an
288 increased representation of deciduous trees in chronically disturbed forests, an aspect rarely
289 documented in previous studies. This study provides a scientific basis for planting evergreen
290 trees in ecological restoration efforts in low-elevation, chronically disturbed forests, which
291 predominantly harbour deciduous trees. Given the absence of Protected Areas in low elevations
292 and significant differences in vegetation composition in high- and low-elevation forests, the
293 existing Reserved (historically-disturbed forests) and Community-owned sacred groves (less-
294 disturbed forests) must be protected from further degradation and conversion in partnership with
295 local stakeholders.

296 **CAD-induced shift from evergreen to deciduous forests**

297 Most studies have looked at the impacts of CAD and climate in isolation. Given our poor
298 understanding of the interactive effects between CAD and climate, this study helps fill that
299 knowledge gap. Higher water deficit induces deciduousness in the vegetation and contributes to
300 the transition from evergreen to deciduous vegetation type (Saiter et al. 2015). While CAD is
301 thought to impact plant diversity and structure (Rito et al. 2017), very few studies have

302 documented CAD causing shifts in vegetation types (Bradshaw & Hannon, 1992) as documented
303 in this study. We demonstrate that CAD disrupts this relationship between CWD and
304 evergreenness. We find that for given levels of CWD, CAD results in a significantly lower
305 representation of evergreen trees and species in the community. The greater representation of
306 deciduous trees with CAD could be because of several reasons: source, dispersal and
307 establishment limitation, poor competitive ability of evergreen tree species in more open
308 conditions and greater propensity of deciduous trees to coppice (Chandran 1997). The species
309 composition of neighbouring forests shapes plant composition in a forest patch through seed
310 dispersal (Butaye et al., 2002). In the low-elevation areas of our study site, there are very few
311 remaining patches of intact, evergreen forests. Additionally, our previous study shows that CAD
312 negatively impacts the prevalence of avian frugivores (Biswas et al., 2023), which play an
313 important role in the seed dispersal of fleshy-fruited evergreen plants (Naniwadekar et al., 2019).
314 Thus, the absence of evergreen tree species and reduced habitat use by frugivores in repeatedly-
315 disturbed forests could result in source limitation. Even if the seeds are dispersed in these open
316 degraded patches, evergreen seedlings, which tend to have a higher shade tolerance (Baldochi et
317 al., 2010; Kitajima et al., 2013), may not be able to establish in low canopy cover conditions
318 (Swinfield et al., 2016). Human disturbances such as clear felling of forests initiate secondary
319 succession, which often begins with a resource-rich condition associated with increased light
320 availability due to reduced canopy cover (Dalling, 2008). Often, the pioneer or early successional
321 species in degraded forests are shade-intolerant deciduous trees (Jin et al., 2017). Deciduous tree
322 species usually adopt a resource-acquisitive strategy and have higher growth rates, unlike
323 evergreen tree species that adopt a resource-conservative strategy (L. Wang et al., 2023). Thus,
324 deciduous tree species can be expected to be competitively superior in more open conditions,

325 resulting in the filtering of evergreen tree species recruits in degraded habitats. Furthermore,
326 deciduous trees are known to coppice well (Chandran 1997) and are associated with degraded
327 habitats, as seen in our study. Future studies need to determine the relative influence of different
328 processes in causing the shift from evergreenness to deciduousness. Our findings offer evidence
329 that there is a need for active restoration efforts with focus on evergreen plants in degraded
330 habitats of the northern Western Ghats since natural regeneration may not be able to restore
331 evergreen plants, which were likely present in the past.

332 The distribution of humid tropical forests (HTFs) is best characterised by high rainfall
333 regimes with low water stress environments (Zelazowski et al., 2011). Climate change
334 projections predict that rainfall will decrease in parts of the Western Ghats (Katzenberger et al.,
335 2021; Rajendran et al., 2012). Thus, chronic anthropogenic disturbance and lowered precipitation
336 can trigger shifts in vegetation type to drier vegetation in these humid tropical forests. Moreover,
337 a greater proportion of evergreen tree species are under the threatened category than deciduous
338 tree species (Fig. S4). Thus, the transition from evergreen to deciduous forests also has
339 consequences for threatened plant species in the region.

340

341 **Sacred groves as reservoirs of biodiversity**

342 We found that the community-owned, less-disturbed forests, i.e., sacred groves, had comparable
343 richness and structure as the state-owned, historically-disturbed forests. This contrasts with the
344 central Western Ghats, where state-owned forests performed better on diversity and structure
345 metrics than community-owned sacred groves (Osuri et al., 2014). The drivers of higher plant
346 diversity in sacred groves could be multifold, as they could be remnants of historically
347 contiguous forests or could have been actively restored in the historical past. The existing

348 evergreen tree cover in two sacred groves in central Western Ghats was attributed to cultural
349 practice-driven forest recovery from around 1000 years before the present (Bhagwat et al., 2013).
350 Therefore, there is a need to understand the socio-ecological history of these sacred groves that
351 can throw more light on the observed diversity of woody plants in these groves.

352 Sixty percent of sacred groves in the central Western Ghats originally present in official
353 records were lost and there was a decrease in above-ground biomass and proportion of evergreen
354 species in existing ones (Osuri et al., 2014; Bhagwat et al., 2005). Similarly, there is documented
355 evidence of sacred groves in the northern Western Ghats being cleared and lost for almost 50
356 years (Gadgil and Vartak, 1976). The drivers of loss are many, including logging,
357 encroachments, and habitat conversion. With monoculture plantations replacing private forests
358 and increased development, the pressures on the sacred groves as a source of timber and
359 fuelwood for local communities will increase. Partnerships between conservation practitioners,
360 government departments and local communities are critical to safeguard these groves in a
361 socially just manner.

362

363 **Value of low-elevation forests**

364 We recorded a higher taxonomic diversity of woody plants in our low-elevation sampling sites
365 than the high-elevation ones, especially for the repeatedly-disturbed and less-disturbed forests
366 (Figure S3). Previous studies documenting plant diversity across elevational gradients have also
367 reported similar patterns, with species richness decreasing with elevation (Musciano et al., 2021;
368 Malizia et al., 2020). In the Western Ghats, forests at or below 500 m have the least
369 representation in the protected area network (Bawa et al., 2007). Moreover, only around 1% of
370 land in the northern Western Ghats is legally protected (Blicharska et al., 2013). All the PAs in

371 our study region are in high elevations, leaving the highly diverse, lowland forests unprotected.
372 PAs, which are the high-elevation historically disturbed sites, had the least abundance of cut
373 stems, indicating their efficacy in reducing extractive pressures. The privately or community-
374 owned forests here are prone to habitat degradation (Kulkarni and Mehta, 2013) and conversion,
375 mostly to cash crop plantations (Kale et al., 2016). More than one-third of the geographic area of
376 two tehsils in Sindhudurg District is under cashew plantation (area: 533.5 km²) (Rege et al.,
377 2022). This study did not estimate areas under rubber and mango plantations, which are also
378 prevalent in the region. Most of these plantations were erstwhile private or community-owned
379 forests. Therefore, these remnant patches of RFs and sacred groves in the low elevations are
380 important reservoirs of diverse assemblages of woody plants and other biodiversity. The RFs are
381 relatively less protected than PAs and are more vulnerable to getting denotified and converted to
382 other land uses (Patil 2023). Protection and active restoration, wherever needed, must be
383 prioritised in these forests so that they can continue to sustain high levels of biodiversity.

384 The higher number of cut stems and low basal area of trees in the repeatedly disturbed
385 private forests (Table S4) of high elevations suggests that these forests experience greater CAD
386 than those in the low elevations. As CWD is strongly correlated with elevation, forests with low
387 CWD are high-elevation forests. Generally, an opposite trend is observed globally, where low-
388 elevation forests are more vulnerable to logging, deforestation and habitat conversion, even in
389 the world's biodiversity hotspots (Hamunyela et al., 2020; Tapia-Armijos et al., 2015). This may
390 be attributed to the easy access to large trees by logging companies in the foothills compared to
391 the stunted but steep forests of the highlands (Danielsen et al., 2010). However, in our study
392 area, the high-elevation forests, which receive a high amount of rainfall (> 5000 mm), are
393 relatively easily accessible as they are on the Deccan plateau. These high-elevation, privately-

394 owned forests are also a source of fuelwood for the locals, who may rely on it to provide warmth
395 in their homes during the cooler and wetter seasons. A study across an altitudinal gradient in the
396 western Himalayas revealed a similar occurrence. Fuelwood consumption was found to be 2.6
397 times higher at high elevations (above 2000 m) than its use at low elevations (up to 500 m),
398 owing to people's need to heat spaces and water in harsh weather (Bhatt and Sachan, 2004). The
399 fuelwood is also utilised by the sugar factories in the region, most of which are in the high-
400 elevations. Fuelwood collection can lead to forest degradation (Sassen et al., 2015) and
401 negatively affect regeneration, thereby changing the vegetation type (as documented in this
402 study) and leading to biodiversity decline. Evergreen forests in the Western Ghats harbour a
403 higher diversity of endemic and threatened trees than the deciduous forests (Chandran 1997;
404 Ghate et al. 1998). Thus, the relative needs of industry and local communities for fuelwood must
405 be determined to find suitable alternatives to fuelwood and restore the degraded private forests in
406 these regions.

407 **Conservation implications**

408 Tropical wet forests are vulnerable to climate fluctuations and anthropogenic disturbance. We
409 can expect CAD to shift the wetter forests to drier vegetation types. This effect will be
410 exacerbated by reduced water availability due to climate change. Thus, there is a need to find
411 suitable alternatives to reduce CAD in tropical forests since conversion to deciduous forests will
412 be associated with a significant loss of threatened and endemic plant diversity, which is generally
413 higher in wet tropical forests. Due to the degraded nature of privately owned forests in the
414 northern Western Ghats, primarily driven by periodic clear felling of forests that cater to the
415 fuelwood needs of nearby factories, there is a need to find suitable alternatives to fuelwood in
416 these factories. Given the agroforestry-friendly government policies and the increased advent of

417 technology that allows easy conversion of forests to agroforestry plantations, many existing
418 privately-owned forests will likely be converted to agroforestry plantations. Given compositional
419 and diversity differences in low- and high-elevation forests, ecological restoration efforts of
420 degraded, low-elevation private forests must be prioritised in partnership with local landowners.
421 This is especially important for the lower elevations, which do not harbour any Protected Areas
422 but continue to harbour significant threatened and endemic biodiversity. The existence of
423 restored habitat patches in a predominantly human-modified landscape will enable the threatened
424 biodiversity to persist in the long term. Given that the existing degraded forest patches harbour
425 predominantly deciduous forests due to CAD, ecological restoration efforts should plant
426 evergreen tree species, particularly in the lower-elevation forests.

427

428 **ACKNOWLEDGEMENTS**

429 We are grateful to the Maharashtra Forest Department, particularly Sunil Limaye (CWLW),
430 Nanasahab Ladkat, Uttam Sawant, Vishal Mali, DFOs of Sawantwadi and Chiplun Forest
431 Division, Suhas Patil and Ajit Mali for giving us the necessary permissions (No: Desk-
432 22(8)/WL/Research/CR-53(20-21)/3361/22-22) and support to conduct fieldwork. We thank On
433 the Edge Conservation (UK) for funding this work. We thank Vijay Kumar for helping with
434 fieldwork. We are grateful to Mahesh Sankaran, Jayashree Ratnam, Vijay Karthick, Abhishek
435 Gopal, Suri Venkatachalam, Madhura Niphadkar, Atul Joshi, Aparna Watve, Praveen Desai,
436 Milind Patil, Parag Rangnekar, Hemant Ogale, Kaka Bhise, Monali Mhaskar, Mahesh
437 Mhangore, Gajanan Shetye, Vinayak Sapkal, Dilip Joglekar, Shrikant Ingalhalikar for support

438 and discussions. We thank all the villagers and households who warmly hosted us during our
439 travels.

440 **LITERATURE CITED**

- 441 Baldocchi, D. D., Ma, S., Rambal, S., Misson, L., Ourcival, J.-M., Limousin, J.-M., Pereira, J., &
442 Papale, D. (2010). On the differential advantages of evergreenness and deciduousness in
443 mediterranean oak woodlands: A flux perspective. *Ecological Applications*, 20(6), 1583–
444 1597. <https://doi.org/10.1890/08-2047.1>
- 445 Barlow, J., Lennox, G. D., Ferreira, J., Berenguer, E., Lees, A. C., Nally, R. M., Thomson, J. R.,
446 Ferraz, S. F. de B., Louzada, J., Oliveira, V. H. F., Parry, L., Ribeiro de Castro Solar, R.,
447 Vieira, I. C. G., Aragão, L. E. O. C., Begotti, R. A., Braga, R. F., Cardoso, T. M., de
448 Oliveira, R. C., Souza Jr, C. M., ... Gardner, T. A. (2016). Anthropogenic disturbance in
449 tropical forests can double biodiversity loss from deforestation. *Nature*, 535(7610), 144–
450 147. <https://doi.org/10.1038/nature18326>
- 451 Bawa, K. S., Das, A., Krishnaswamy, J., Karanth, K.U., Kumar, N.S. and Rao, M. (2007).
452 Critical Ecosystem Partnership Fund - Ecosystem Profile. Western Ghats and Sri Lanka
453 Biodiversity Hotspot. Western Ghats region. *ATREE, Bangalore, India*.
- 454 Bentsi-Enchill, F., Dampney, F. G., Pappoe, A. N. M., Ekumah, B., & Akotoye, H. K. (2022).
455 Impact of anthropogenic disturbance on tree species diversity, vegetation structure and
456 carbon storage potential in an upland evergreen forest of Ghana, West Africa. *Trees,*
457 *Forests and People*, 8, 100238. <https://doi.org/10.1016/j.tfp.2022.100238>
- 458 Bhagwat, S. A., Kushalappa, C. G., Williams, P. H., & Brown, N. D. (2005). The role of
459 informal Protected Areas in maintaining biodiversity in the Western Ghats of India.
460 *Ecology and Society*, 10(1), 8. <https://www.jstor.org/stable/26267704>

- 461 Bhagwat, S. A., Nogué, S., & Willis, K. J. (2014). Cultural drivers of reforestation in tropical
462 forest groves of the Western Ghats of India. *Forest Ecology and Management*, 329, 393–
463 400. <https://doi.org/10.1016/j.foreco.2013.11.017>
- 464 Bhatt, B. P., & Sachan, M. S. (2004). Firewood consumption along an altitudinal gradient in
465 mountain villages of India. *Biomass and Bioenergy*, 27(1), 69–75.
466 <https://doi.org/10.1016/j.biombioe.2003.10.004>
- 467 Biswas, N., Biniwale, S., Sadekar, V., Taneja, Y., Suryawanshi, K., Osuri, A. M., &
468 Naniwadekar, R. (2023). Conservation value of low-elevation forests for birds in
469 agroforestry-dominated landscapes in a biodiversity hotspot (p. 2023.08.26.554947).
470 *bioRxiv*. <https://doi.org/10.1101/2023.08.26.554947>
- 471 Blicharska, M., Mikusiński, G., Godbole, A., & Sarnaik, J. (2013). Safeguarding biodiversity and
472 ecosystem services of sacred groves – experiences from northern Western Ghats.
473 *International Journal of Biodiversity Science, Ecosystem Services & Management*, 9(4),
474 339–346. <https://doi.org/10.1080/21513732.2013.835350>
- 475 Bradshaw, R., & Hannon, G. (1992). Climatic change, human influence and disturbance regime
476 in the control of vegetation dynamics within Fiby Forest, Sweden. *The Journal of*
477 *Ecology*, 80(4), 625. <https://doi.org/10.2307/2260854>
- 478 Butaye, J., Jacquemyn, H., Honnay, O., & Hermy, M. (2002). The species pool concept applied
479 to forests in a fragmented landscape: Dispersal limitation versus habitat limitation.
480 *Journal of Vegetation Science*, 13(1), 27–34. [https://doi.org/10.1111/j.1654-](https://doi.org/10.1111/j.1654-1103.2002.tb02020.x)
481 [1103.2002.tb02020.x](https://doi.org/10.1111/j.1654-1103.2002.tb02020.x)
- 482 Champion, S. H. G., & Seth, S. K. (1968). *A Revised Survey of the Forest Types of India*.
483 Manager of Publications.

- 484 Chandran, M. D. (1997). On the ecological history of the Western Ghats. *Current Science*, 73,
485 146–155.
- 486 Chave, J., Réjou-Méchain, M., Búrquez, A., Chidumayo, E., Colgan, M. S., Delitti, W. B. C.,
487 Duque, A., Eid, T., Fearnside, P. M., Goodman, R. C., Henry, M., Martínez-Yrizar, A.,
488 Mugasha, W. A., Muller-Landau, H. C., Mencuccini, M., Nelson, B. W., Ngomanda, A.,
489 Nogueira, E. M., Ortiz-Malavassi, E., ... Vieilledent, G. (2014). Improved allometric
490 models to estimate the aboveground biomass of tropical trees. *Global Change Biology*,
491 20(10), 3177–3190. <https://doi.org/10.1111/gcb.12629>
- 492 Cunningham, C., & Beazley, K. F. (2018). Changes in human population density and protected
493 areas in terrestrial global biodiversity hotspots, 1995–2015. *Land*, 7(4), Article 4.
494 <https://doi.org/10.3390/land7040136>
- 495 Dalling, J. W. (2008). Pioneer Species. In *Encyclopedia of Ecology, Five-Volume Set* (pp. 2779–
496 2782). Elsevier Inc. <https://doi.org/10.1016/B978-008045405-4.00534-6>
- 497 Danielsen, F., Filardi, C. E., Jønsson, K. A., Kohaia, V., Krabbe, N., Kristensen, J. B., Moyle, R.
498 G., Pikacha, P., Poulsen, M. K., Sørensen, M. K., Tatahu, C., Waihuru, J., & Fjeldså, J.
499 (2010). Endemic avifaunal biodiversity and tropical forest loss in Makira, a mountainous
500 Pacific island. *Singapore Journal of Tropical Geography*, 31(1), 100–114.
501 <https://doi.org/10.1111/j.1467-9493.2010.00386.x>
- 502 Dexter, K. G., Pennington, R. T., Oliveira-Filho, A. T., Bueno, M. L., Silva De Miranda, P. L., &
503 Neves, D. M. (2018). Inserting tropical dry forests into the discussion on biome
504 transitions in the tropics. *Frontiers in Ecology and Evolution*, 6, 104.
505 <https://doi.org/10.3389/fevo.2018.00104>
- 506 Esquivel-Muelbert, A., Baker, T. R., Dexter, K. G., Lewis, S. L., ter Steege, H., Lopez-Gonzalez,

- 507 G., Monteagudo Mendoza, A., Brienen, R., Feldpausch, T. R., Pitman, N., Alonso, A.,
508 van der Heijden, G., Peña-Claros, M., Ahuite, M., Alexiades, M., Álvarez Dávila, E.,
509 Murakami, A. A., Arroyo, L., Aulestia, M., ... Phillips, O. L. (2017). Seasonal drought
510 limits tree species across the Neotropics. *Ecography*, *40*(5), 618–629.
511 <https://doi.org/10.1111/ecog.01904>
- 512 Gadgil, M., & Vartak, V. D. (1976). The sacred groves of Western Ghats in India. *Economic*
513 *Botany*, *30*(2), 152–160. <https://doi.org/10.1007/BF02862961>
- 514 Ghate, U., Joshi, N. V., & Gadgil, M. (1998). On the patterns of tree diversity in the
515 Western Ghats of India. *Current Science*, *75*(6), 594–603.
516 <https://www.jstor.org/stable/24100563>
- 517 Gopal, A., Bharti, D. K., Page, N., Dexter, K. G., Krishnamani, R., Kumar, A., & Joshi, J.
518 (2023). Range restricted old and young lineages show the southern Western Ghats to be
519 both a museum and a cradle of diversity for woody plants. *Proceedings of the Royal*
520 *Society B*, *290*(1997), 20222513. <https://doi.org/10.1098/rspb.2022.2513>
- 521 Gunawardene, N. R., Daniels, A. E. D., Gunatilleke, C. V. S., Karunakaran, P. V., Nayak, K. G.,
522 Prasad, S., Puyravaud, P., Ramesh, B. R., Subramanian, K. A., & Vasanthi, G. (2007). A
523 brief overview of the Western Ghats – Sri Lanka biodiversity hotspot. *Current Science*,
524 *93*(11), 1567-1572.
- 525 Hamunyela, E., Brandt, P., Shirima, D., Do, H. T. T., Herold, M., & Roman-Cuesta, R. M.
526 (2020). Space-time detection of deforestation, forest degradation and regeneration in
527 montane forests of Eastern Tanzania. *International Journal of Applied Earth Observation*
528 *and Geoinformation*, *88*, 102063. <https://doi.org/10.1016/j.jag.2020.102063>
- 529 Harrison, S., Spasojevic, M. J., & Li, D. (2020). Climate and plant community diversity in space

- 530 and time. *Proceedings of the National Academy of Sciences*, 117(9), 4464–4470.
- 531 <https://doi.org/10.1073/pnas.1921724117>
- 532 Hirota, M., Holmgren, M., Van Nes, E. H., & Scheffer, M. (2011). Global resilience of tropical
533 forest and savanna to critical transitions. *Science*, 334(6053), 232–235.
- 534 <https://doi.org/10.1126/science.1210657>
- 535 Hooper, D. U., Adair, E. C., Cardinale, B. J., Byrnes, J. E. K., Hungate, B. A., Matulich, K. L.,
536 Gonzalez, A., Duffy, J. E., Gamfeldt, L., & O'Connor, M. I. (2012). A global synthesis
537 reveals biodiversity loss as a major driver of ecosystem change. *Nature*, 486(7401),
538 Article 7401. <https://doi.org/10.1038/nature11118>
- 539 Jog, S. (2009). Sahyadris—Flora and Ethnobotany. Report William Brown Fellowship 2007-08.
540 [https://www.researchgate.net/profile/Suneeti-
541 Jog/publication/238080269_SAHYADRI_-
542 _FLORA_AND_ETHNOBOTANY/links/0a85e534ec60405aed000000/SAHYADRI-
543 FLORA-AND-ETHNOBOTANY.pdf?_sg%5B0%5D=txBdlbGnupoP_2UZYff-
544 hAe3f6ccm1qCAPNmsfbZ6szCfnxZdqsK5ygBPxn-
545 KSR_NelvDoUnxmhcHuTPNIHSmg.2nuCAgS3qHHkHLS9cIsdmig96zHLJM99JtVlr2
546 NXqTFyFfrA8tUxWNLKnxvUhEM1FJ3nHA1YXc0iaGJkqiSEWw&_sg%5B1%5D=16
547 y1C6DmQhW_9UqYKtI5KcVHTHsQax6POOsxYYaudy9SfMBd2Jcqum2A8l4bDDRg
548 kMOHoZZuBXxO1VQmZBercQ3l9QA9DQ3iU2bk9WQvM5T7.2nuCAgS3qHHkHLS9
549 cIsdmig96zHLJM99JtVlr2NXqTFyFfrA8tUxWNLKnxvUhEM1FJ3nHA1YXc0iaGJkqi
550 SEWw&_iepl=](https://www.researchgate.net/profile/Suneeti-Jog/publication/238080269_SAHYADRI-_FLORA_AND_ETHNOBOTANY/links/0a85e534ec60405aed000000/SAHYADRI-_FLORA-AND-ETHNOBOTANY.pdf?_sg%5B0%5D=txBdlbGnupoP_2UZYff-hAe3f6ccm1qCAPNmsfbZ6szCfnxZdqsK5ygBPxn-KSR_NelvDoUnxmhcHuTPNIHSmg.2nuCAgS3qHHkHLS9cIsdmig96zHLJM99JtVlr2NXqTFyFfrA8tUxWNLKnxvUhEM1FJ3nHA1YXc0iaGJkqiSEWw&_sg%5B1%5D=16y1C6DmQhW_9UqYKtI5KcVHTHsQax6POOsxYYaudy9SfMBd2Jcqum2A8l4bDDRgkMOHoZZuBXxO1VQmZBercQ3l9QA9DQ3iU2bk9WQvM5T7.2nuCAgS3qHHkHLS9cIsdmig96zHLJM99JtVlr2NXqTFyFfrA8tUxWNLKnxvUhEM1FJ3nHA1YXc0iaGJkqiSEWw&_iepl=)
- 551 Joglekar, A., Tadwalkar, M., Mhaskar, M., Chavan, B., Ganeshaiyah, K. N., & Patwardhan, A.
552 (2015). Tree species composition in Koyna Wildlife Sanctuary, Northern Western Ghats

553 of India. *Current Science*, 108(9), 1688–1693. <https://www.jstor.org/stable/24905535>

554 Joseph, S., Anitha, K., Srivastava, V. K., Reddy, C. S., Thomas, A. P., & Murthy, M. S. R.
555 (2012). Rainfall and elevation influence the local-scale distribution of tree community in
556 the southern region of Western Ghats Biodiversity Hotspot (India). *International Journal*
557 *of Forestry Research*, 2012, e576502. <https://doi.org/10.1155/2012/576502>

558 Kale, M. P., Chavan, M., Pardeshi, S., Joshi, C., Verma, P. A., Roy, P. S., Srivastav, S. K.,
559 Srivastava, V. K., Jha, A. K., Chaudhari, S., Giri, Y., & Krishna Murthy, Y. V. N. (2016).
560 Land-use and land-cover change in Western Ghats of India. *Environmental Monitoring*
561 *and Assessment*, 188(7), 387. <https://doi.org/10.1007/s10661-016-5369-1>

562 Kanade, R., Tadwalkar, M., Kushalappa, C., & Patwardhan, A. (2008). Vegetation composition
563 and woody species diversity at Chandoli National Park, northern Western Ghats, India.
564 *Current Science*, 95(5), 637–646. <https://www.jstor.org/stable/24102804>

565 Katzenberger, A., Schewe, J., Pongratz, J., & Levermann, A. (2021). Robust increase of Indian
566 monsoon rainfall and its variability under future warming in CMIP6 models. *Earth*
567 *System Dynamics*, 12(2), 367–386. <https://doi.org/10.5194/esd-12-367-2021>

568 Kitajima, K., Cordero, R. A., & Wright, S. J. (2013). Leaf life span spectrum of tropical woody
569 seedlings: Effects of light and ontogeny and consequences for survival. *Annals of Botany*,
570 112(4), 685–699. <https://doi.org/10.1093/aob/mct036>

571 Krishnadas, M., Kumar, A., & Comita, L. S. (2016). Environmental gradients structure tropical
572 tree assemblages at the regional scale. *Journal of Vegetation Science*, 27(6), 1117–1128.
573 <https://doi.org/10.1111/jvs.12438>

574 Krishnadas, M., Sankaran, M., Page, N., Joshi, J., Machado, S., Nataraj, N., Chengappa, S. K.,
575 Kumar, V., Kumar, A., & Krishnamani, R. (2021). Seasonal drought regulates species

576 distributions and assembly of tree communities across a tropical wet forest region. *Global*
577 *Ecology and Biogeography*, 30(9), 1847–1862. <https://doi.org/10.1111/geb.13350>

578 Kulkarni, A., Anuradha, U., Dahanukar, N., & Datar, M. (2018). Floristic uniqueness and effect
579 of degradation on diversity: A case study of sacred groves from northern Western Ghats.
580 *Tropical Ecology*, 59.

581 Kulkarni, J., & Mehta, P. (2013). A study of status, distribution and dynamics of private and
582 community forests in Sahyadri-Konkan corridor of Maharashtra Western Ghats.
583 [Technical Report submitted to CEPF-ATREE]. Wildlife Research and Conservation
584 Society, Pune.

585 Malhi, Y., Gardner, T. A., Goldsmith, G. R., Silman, M. R., & Zelazowski, P. (2014). Tropical
586 forests in the Anthropocene. *Annual Review of Environment and Resources*, 39(1), 125–
587 159. <https://doi.org/10.1146/annurev-environ-030713-155141>

588 Malizia, A., Blundo, C., Carilla, J., Acosta, O. O., Cuesta, F., Duque, A., Aguirre, N., Aguirre,
589 Z., Ataroff, M., Baez, S., Calderón-Loor, M., Cayola, L., Cayuela, L., Ceballos, S.,
590 Cedillo, H., Ríos, W. F., Feeley, K. J., Fuentes, A. F., Álvarez, L. E. G., ... Young, K. R.
591 (2020). Elevation and latitude drives structure and tree species composition in Andean
592 forests: Results from a large-scale plot network. *PLOS ONE*, 15(4), e0231553.
593 <https://doi.org/10.1371/journal.pone.0231553>

594 McLaughlin, B. C., Ackerly, D. D., Klos, P. Z., Natali, J., Dawson, T. E., & Thompson, S. E.
595 (2017). Hydrologic refugia, plants, and climate change. *Global Change Biology*, 23(8),
596 2941–2961. <https://doi.org/10.1111/gcb.13629>

597 Mittermeier, R. A., Turner, W. R., Larsen, F. W., Brooks, T. M., & Gascon, C. (2011). Global
598 biodiversity conservation: The critical role of hotspots. In F. E. Zachos & J. C. Habel

599 (Eds.), *Biodiversity Hotspots* (pp. 3–22). Springer Berlin Heidelberg.
600 https://doi.org/10.1007/978-3-642-20992-5_1

601 Musciano, M. D., Zannini, P., Ferrara, C., Spina, L., Nascimbene, J., Vetaas, O. R., Bhatta, K. P.,
602 d’Agostino, M., Peruzzi, L., Carta, A., & Chiarucci, A. (2021). Investigating elevational
603 gradients of species richness in a Mediterranean plant hotspot using a published flora.
604 *E50007*. <https://doi.org/10.21425/F5FBG50007>

605 Myers, N., Mittermeier, R. A., Mittermeier, C. G., da Fonseca, G. A. B., & Kent, J. (2000).
606 Biodiversity hotspots for conservation priorities. *Nature*, *403*(6772), Article 6772.
607 <https://doi.org/10.1038/35002501>

608 Naniwadekar, R., Chaplod, S., Datta, A., Rathore, A., & Sridhar, H. (2019). Large frugivores
609 matter: Insights from network and seed dispersal effectiveness approaches. *Journal of*
610 *Animal Ecology*, *88*(8), 1250–1262. <https://doi.org/10.1111/1365-2656.13005>

611 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., ... & Wagner,
612 H. (2022). vegan: Community Ecology Package. R package version 2.5-7. 2020. *Preprint*
613 *at*, 3-1.

614 Oksanen, J., Kindt, R., Simpson, G. L., & Oksanen, M. J. (2018). Package ‘vegan3d’. *R package*
615 *version*, 1-0.

616 Osuri, A. M., Madhusudan, M. D., Kumar, V. S., Chengappa, S. K., Kushalappa, C. G., &
617 Sankaran, M. (2014). Spatio-temporal variation in forest cover and biomass across sacred
618 groves in a human-modified landscape of India’s Western Ghats. *Biological*
619 *Conservation*, *178*, 193–199. <https://doi.org/10.1016/j.biocon.2014.08.008>

620 Page, N. V., & Shanker, K. (2018). Environment and dispersal influence changes in species
621 composition at different scales in woody plants of the Western Ghats, India. *Journal of*

- 622 *Vegetation Science*, 29(1), 74–83. <https://doi.org/10.1111/jvs.12586>
- 623 Patil, A. (2023). Temporary stay on Adani Green’s soil testing for Patgaon project. *The Times of*
624 *India*.
- 625 Rajendran, K., Kitoh, A., Srinivasan, J., Mizuta, R., & Krishnan, R. (2012). Monsoon circulation
626 interaction with Western Ghats orography under changing climate. *Theoretical and*
627 *Applied Climatology*, 110(4), 555–571. <https://doi.org/10.1007/s00704-012-0690-2>
- 628 Ramos, M. B., Maciel, M. G. R., Cunha, S. S. D., De Souza, S. M., Pedrosa, K. M., De Souza, J.
629 J. L. L., González, E. J., Meave, J. A., & Lopes, S. D. F. (2023). The role of chronic
630 anthropogenic disturbances in plant community assembly along a water availability
631 gradient in Brazil’s semiarid Caatinga region. *Forest Ecology and Management*, 538,
632 120980. <https://doi.org/10.1016/j.foreco.2023.120980>
- 633 Reddy, C. S., Jha, C. S., & Dadhwal, V. K. (2016). Assessment and monitoring of long-term
634 forest cover changes (1920–2013) in Western Ghats biodiversity hotspot. *Journal of*
635 *Earth System Science*, 125(1), 103–114. <https://doi.org/10.1007/s12040-015-0645-y>
- 636 Rege, A., Warnekar, S. B., & Lee, J. S. H. (2022). Mapping cashew monocultures in the Western
637 Ghats using optical and radar imagery in Google Earth Engine. *Remote Sensing*
638 *Applications: Society and Environment*, 28, 100861.
639 <https://doi.org/10.1016/j.rsase.2022.100861>
- 640 Rito, K. F., Arroyo-Rodríguez, V., Queiroz, R. T., Leal, I. R., & Tabarelli, M. (2017).
641 Precipitation mediates the effect of human disturbance on the Brazilian Caatinga
642 vegetation. *Journal of Ecology*, 105(3), 828–838. [https://doi.org/10.1111/1365-](https://doi.org/10.1111/1365-2745.12712)
643 [2745.12712](https://doi.org/10.1111/1365-2745.12712)
- 644 Roswell, M., Dushoff, J., & Winfree, R. (2021). A conceptual guide to measuring species

- 645 diversity. *Oikos*, 130(3), 321–338. doi:10.1111/oik.07202
- 646 Saatchi, S., Longo, M., Xu, L., Yang, Y., Abe, H., André, M., Aukema, J. E., Carvalhais, N.,
647 Cadillo-Quiroz, H., Cerbu, G. A., Chernela, J. M., Covey, K., Sánchez-Clavijo, L. M.,
648 Cubillos, I. V., Davies, S. J., De Sy, V., De Vleeschouwer, F., Duque, A., Sybille
649 Durieux, A. M., ... Elmore, A. C. (2021). Detecting vulnerability of humid tropical
650 forests to multiple stressors. *One Earth*, 4(7), 988–1003.
651 <https://doi.org/10.1016/j.oneear.2021.06.002>
- 652 Saiter, F. Z., Eisenlohr, P. V., Barbosa, M. R. V., Thomas, W. W., & Oliveira-Filho, A. T.
653 (2016). From evergreen to deciduous tropical forests: How energy–water balance,
654 temperature, and space influence the tree species composition in a high diversity region.
655 *Plant Ecology & Diversity*, 9(1), 45–54. <https://doi.org/10.1080/17550874.2015.1075623>
- 656 Sansevero, J. B. B., Garbin, M. L., Sánchez-Tapia, A., Valladares, F., & Scarano, F. R. (2020).
657 Fire drives abandoned pastures to a savanna-like state in the Brazilian Atlantic Forest.
658 *Perspectives in Ecology and Conservation*, 18(1), 31–36.
659 <https://doi.org/10.1016/j.pecon.2019.12.004>
- 660 Sassen, M., Sheil, D., & Giller, K. E. (2015). Fuelwood collection and its impacts on a protected
661 tropical mountain forest in Uganda. *Forest Ecology and Management*, 354, 56–67.
662 <https://doi.org/10.1016/j.foreco.2015.06.037>
- 663 Seiler, C., Hutjes, R. W. A., Kruijt, B., Quispe, J., Añez, S., Arora, V. K., Melton, J. R., Hickler,
664 T., & Kabat, P. (2014). Modeling forest dynamics along climate gradients in Bolivia.
665 *Journal of Geophysical Research: Biogeosciences*, 119(5), 758–775.
666 <https://doi.org/10.1002/2013JG002509>
- 667 Shivaprakash, K. N., Ramesh, B. R., Umashaanker, R., & Dayanandan, S. (2018). Functional

668 trait and community phylogenetic analyses reveal environmental filtering as the major
669 determinant of assembly of tropical forest tree communities in the Western Ghats
670 biodiversity hotspot in India. *Forest Ecosystems*, 5(1), 25.
671 <https://doi.org/10.1186/s40663-018-0144-0>

672 Swinfield, T., Afriandi, R., Antoni, F., & Harrison, R. D. (2016). Accelerating tropical forest
673 restoration through the selective removal of pioneer species. *Forest Ecology and*
674 *Management*, 381, 209–216. <https://doi.org/10.1016/j.foreco.2016.09.020>

675 Tapia-Armijos, M. F., Homeier, J., Espinosa, C. I., Leuschner, C., & Cruz, M. de la. (2015).
676 Deforestation and Forest Fragmentation in South Ecuador since the 1970s – Losing a
677 Hotspot of Biodiversity. *PLOS ONE*, 10(9), e0133701.
678 <https://doi.org/10.1371/journal.pone.0133701>

679 Terra, M. de C. N. S., Santos, R. M. dos, Prado Júnior, J. A. do, de Mello, J. M., Scolforo, J. R.
680 S., Fontes, M. A. L., Schiavini, I., dos Reis, A. A., Bueno, I. T., Magnago, L. F. S., & ter
681 Steege, H. (2018). Water availability drives gradients of tree diversity, structure and
682 functional traits in the Atlantic–Cerrado–Caatinga transition, Brazil. *Journal of Plant*
683 *Ecology*, 11(6), 803–814. <https://doi.org/10.1093/jpe/rty017>

684 Venkatesh, B., Nayak, P. C., Thomas, T., Jain, S. K., & Tyagi, J. V. (2021). Spatio-temporal
685 analysis of rainfall pattern in the Western Ghats region of India. *Meteorology and*
686 *Atmospheric Physics*, 133(4), 1089–1109. <https://doi.org/10.1007/s00703-021-00796-z>

687 Vicente-Serrano, S. M., Gouveia, C., Camarero, J. J., Beguería, S., Trigo, R., López-Moreno, J.
688 I., Azorín-Molina, C., Pasho, E., Lorenzo-Lacruz, J., Revuelto, J., Morán-Tejeda, E., &
689 Sanchez-Lorenzo, A. (2013). Response of vegetation to drought time-scales across global
690 land biomes. *Proceedings of the National Academy of Sciences*, 110(1), 52–57.

- 691 <https://doi.org/10.1073/pnas.1207068110>
- 692 Wang, L., He, Y., Umer, M., Guo, Y., Tan, Q., Kang, L., Fang, Z., Shen, K., Xia, T., Wu, P.,
693 Liu, Y., Zang, L., Liu, Q., Zhao, Y., Chen, H., & Zhao, Y. (2023). Strategic
694 differentiation of subcommunities composed of evergreen and deciduous woody species
695 associated with leaf functional traits in the subtropical mixed forest. *Ecological*
696 *Indicators*, 150, 110281. <https://doi.org/10.1016/j.ecolind.2023.110281>
- 697 Wang, M., Guo, Q., & Chen, A. (2023). The savannization of tropical forests in mainland
698 Southeast Asia since 2000. *Landscape Ecology*, 38(12), 3961–3971.
699 <https://doi.org/10.1007/s10980-023-01691-1>
- 700 Zelazowski, P., Malhi, Y., Huntingford, C., Sitch, S., & Fisher, J. B. (2011). Changes in the
701 potential distribution of humid tropical forests on a warmer planet. *Philosophical*
702 *Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*,
703 369(1934), 137–160. <https://doi.org/10.1098/rsta.2010.0238>
- 704 Zheng, J., Arif, M., He, X., Ding, D., Zhang, S., Ni, X., & Li, C. (2022). Plant community
705 assembly is jointly shaped by environmental and dispersal filtering along elevation
706 gradients in a semiarid area, China. *Frontiers in Plant Science*, 13.
707 <https://www.frontiersin.org/articles/10.3389/fpls.2022.1041742>
- 708 Zorger, B. B., Tabarelli, M., de Queiroz, R. T., Rosado, B. H. P., & Pinho, B. X. (2019).
709 Functional organization of woody plant assemblages along precipitation and human
710 disturbance gradients in a seasonally dry tropical forest. *Biotropica*, 51(6), 838–850.
711 <https://doi.org/10.1111/btp.12721>

712 **SUPPLEMENTARY MATERIAL**

713 **Table S1.** Checklist of woody plant species detected in 120 (50 × 10 m²) plots. The table also
714 provides information on the number of plots in which the species was detected and the total
715 number of individuals detected.

716

Sr. No	Species	Family	Number of individuals	Number of plots where the species was found
1	<i>Acacia auriculiformis</i>	Fabaceae	3	2
2	<i>Actephila excelsa</i>	Phyllanthaceae	1	1
3	<i>Actinodaphne lanceolata</i>	Lauraceae	21	10
4	<i>Adina cordifolia</i>	Rubiaceae	3	1
5	<i>Aglaia elaeagnoidea</i>	Meliaceae	46	12
6	<i>Aglaia lawii</i>	Meliaceae	51	13
7	<i>Agrostistachys indica</i>	Euphorbiaceae	4	1
8	<i>Albizia lebbek</i>	Fabaceae	20	7

9	<i>Albizia procera</i>	Fabaceae	1	1
10	<i>Alstonia scholaris</i>	Apocynaceae	3	3
11	<i>Anacardium occidentale</i>	Anacardiaceae	8	3
12	<i>Antiaris toxicaria</i>	Moraceae	11	6
13	<i>Aphanamixis polystachya</i>	Meliaceae	1	1
14	<i>Aporosa cardiosperma</i>	Phyllanthaceae	113	19
15	<i>Ardisia solanacea</i>	Primulaceae	163	7
16	<i>Artocarpus gomezianus</i>	Moraceae	1	1
17	<i>Artocarpus heterophyllus</i>	Moraceae	12	6
18	<i>Atalantia racemosa</i>	Rutaceae	43	11
19	<i>Bauhinia racemosa</i>	Fabaceae	5	3
20	<i>Beilschmiedia dalzellii</i>	Lauraceae	42	13
21	<i>Bergera koenigii</i>	Rutaceae	12	8

22	<i>Bischofia javanica</i>	Phyllanthaceae	21	1
23	<i>Blachia denudata</i>	Euphorbiaceae	21	3
24	<i>Bombax ceiba</i>	Malvaceae	19	15
25	<i>Bridelia retusa</i>	Phyllanthaceae	74	32
26	<i>Buchanania lanzan</i>	Anacardiaceae	21	6
27	<i>Butea monosperma</i>	Fabaceae	2	2
28	<i>Callicarpa tomentosa</i>	Lamiaceae	22	12
29	<i>Canarium strictum</i>	Burseraceae	1	1
30	<i>Capparis rotundifolia</i>	Capparaceae	2	1
31	<i>Carallia brachiata</i>	Rhizophoraceae	12	5
32	<i>Careya arborea</i>	Lecythidaceae	114	41
33	<i>Caryota urens</i>	Arecaceae	48	23
34	<i>Casearia graveolens</i>	Salicaceae	52	25

35	<i>Cassia fistula</i>	Fabaceae	2	2
36	<i>Catunaregam spinosa</i>	Rubiaceae	149	30
37	<i>Celtis philippensis</i>	Cannabaceae	7	5
38	<i>Celtis timorensis</i>	Cannabaceae	1	1
39	<i>Chionanthus mala-elengi</i>	Oleaceae	12	4
40	<i>Chukrasia tabularis</i>	Meliaceae	1	1
41	<i>Cinnamomum verum</i>	Lauraceae	14	7
42	<i>Clausena anisata</i>	Rutaceae	1	1
43	<i>Cleidion spiciflorum</i>	Euphorbiaceae	18	2
44	<i>Clerodendrum infortunatum</i>	Lamiaceae	2	2
45	<i>Croton zeylanicus</i>	Euphorbiaceae	1	1
46	<i>Cryptocarya wightiana</i>	Lauraceae	9	3
47	<i>Dalbergia sissoo</i>	Fabaceae	16	5

48	<i>Delonix regia</i>	Fabaceae	1	1
49	<i>Dichapetalum gelonioides</i>	Dichapetalaceae	35	9
50	<i>Dillenia pentagyna</i>	Dilleniaceae	13	9
51	<i>Dimocarpus longan</i>	Sapindaceae	193	23
52	<i>Diospyros candolleana</i>	Ebenaceae	147	27
53	<i>Diospyros montana</i>	Ebenaceae	13	10
54	<i>Diospyros nigrescens</i>	Ebenaceae	123	18
55	<i>Diospyros oocarpa</i>	Ebenaceae	70	10
56	<i>Diospyros sylvatica</i>	Ebenaceae	9	5
57	<i>Donella lanceolata</i>	Sapotaceae	3	1
58	<i>Drypetes venusta</i>	Putranjivaceae	23	6
59	<i>Dysoxylum gotadhora</i>	Meliaceae	46	9
60	<i>Ehretia aspera</i>	Boraginaceae	6	5

61	<i>Elaeocarpus variabilis</i>	Elaeocarpaceae	5	2
62	<i>Elaeodendron paniculatum</i>	Celastraceae	5	4
63	<i>Erinocarpus nimmonii</i>	Malvaceae	1	1
64	<i>Erythrina stricta</i>	Fabaceae	6	4
65	<i>Eugenia kalamii</i>	Myrtaceae	5	2
66	<i>Euonymus indicus</i>	Celastraceae	7	1
67	<i>Falconeria insignis</i>	Euphorbiaceae	12	6
68	<i>Ficus amplissima</i>	Moraceae	1	1
69	<i>Ficus arnottiana</i>	Moraceae	1	1
70	<i>Ficus callosa</i>	Moraceae	1	1
71	<i>Ficus exasperata</i>	Moraceae	5	3
72	<i>Ficus hispida</i>	Moraceae	41	12
73	<i>Ficus microcarpa</i>	Moraceae	1	1

74	<i>Ficus nervosa</i>	Moraceae	7	7
75	<i>Ficus racemosa</i>	Moraceae	32	16
76	<i>Ficus talbotii</i>	Moraceae	1	1
77	<i>Ficus tsjakela</i>	Moraceae	2	2
78	<i>Ficus virens</i>	Moraceae	1	1
79	<i>Firmiana colorata</i>	Malvaceae	1	1
80	<i>Flacourtia montana</i>	Salicaceae	26	13
81	<i>Garcinia gummi-gutta</i>	Clusiaceae	8	4
82	<i>Garcinia indica</i>	Clusiaceae	39	8
83	<i>Garcinia talbotii</i>	Clusiaceae	64	16
84	<i>Glochidion heyneanum</i>	Phyllanthaceae	16	6
85	<i>Glochidion hohenackeri</i>	Phyllanthaceae	9	5
86	<i>Glochidion zeylanicum</i>	Phyllanthaceae	3	3

87	<i>Glycosmis pentaphylla</i>	Rutaceae	15	9
88	<i>Gmelina arborea</i>	Lamiaceae	15	9
89	<i>Gomphandra tetrandra</i>	Stemonuraceae	4	3
90	<i>Grewia serrulata</i>	Malvaceae	14	5
91	<i>Grewia tiliifolia</i>	Malvaceae	66	22
92	<i>Gymnosporia rothiana</i>	Celastraceae	1	1
93	<i>Helicteres isora</i>	Malvaceae	13	8
94	<i>Heterophragma quadriloculare</i>	Bignoniaceae	4	4
95	<i>Holigarna arnottiana</i>	Anacardiaceae	130	20
96	<i>Holigarna grahamii</i>	Anacardiaceae	39	20
97	<i>Homalium ceylanicum</i>	Salicaceae	6	3
98	<i>Hydnocarpus pentandrus</i>	Achariaceae	17	7
99	<i>Ixora brachiata</i>	Rubiaceae	376	45

100	<i>Ixora pavetta</i>	Rubiaceae	7	3
101	<i>Jatropha curcas</i>	Euphorbiaceae	1	1
102	<i>Justicia adhatoda</i>	Acanthaceae	9	1
103	<i>Knema attenuata</i>	Myristicaceae	24	4
104	<i>Kydia calycina</i>	Malvaceae	1	1
105	<i>Lagerstroemia microcarpa</i>	Lythraceae	61	27
106	<i>Lagerstroemia parviflora</i>	Lythraceae	1	1
107	<i>Lannea coromandelica</i>	Anacardiaceae	12	7
108	<i>Lasiosiphon glaucus</i>	Thymelaeaceae	32	12
109	<i>Leea indica</i>	Vitaceae	111	34
110	<i>Lepisanthes tetraphylla</i>	Sapindaceae	23	11
111	<i>Ligustrum robustum</i>	Oleaceae	23	10
112	<i>Litsea glutinosa</i>	Lauraceae	2	1

113	<i>Litsea nigrescens</i>	Lauraceae	3	1
114	<i>Litsea stocksii</i>	Lauraceae	23	7
115	<i>Lophopetalum wightianum</i>	Celastraceae	18	2
116	<i>Macaranga peltata</i>	Euphorbiaceae	125	29
117	<i>Machilus glaucescens</i>	Lauraceae	11	5
118	<i>Mallotus nudiflorus</i>	Euphorbiaceae	1	1
119	<i>Mallotus philippensis</i>	Euphorbiaceae	86	27
120	<i>Mallotus resinousus</i>	Euphorbiaceae	9	2
121	<i>Mammea suriga</i>	Calophyllaceae	49	3
122	<i>Mangifera indica</i>	Anacardiaceae	49	20
123	<i>Mappia nimmoniana</i>	Icacinaceae	25	9
124	<i>Margaritaria indica</i>	Phyllanthaceae	2	2
125	<i>Melastoma malabathricum</i>	Melastomataceae	3	1

126	<i>Melia dubia</i>	Meliaceae	1	1
127	<i>Melicope lunu-ankenda</i>	Rutaceae	1	1
128	<i>Memecylon talbotianum</i>	Melastomataceae	8	4
129	<i>Memecylon umbellatum</i>	Melastomataceae	778	56
130	<i>Memecylon wightii</i>	Melastomataceae	1	1
131	<i>Meyna laxiflora</i>	Rubiaceae	32	15
132	<i>Microcos paniculata</i>	Malvaceae	64	19
133	<i>Miliusa tomentosa</i>	Annonaceae	14	6
134	<i>Mimusops elengi</i>	Sapotaceae	25	14
135	<i>Mitragyna parvifolia</i>	Rubiaceae	3	2
136	<i>Monoon fragrans</i>	Annonaceae	2	2
137	<i>Morinda coreia</i>	Rubiaceae	3	3
138	<i>Murraya paniculata</i>	Rutaceae	3	3

139	<i>Myristica beddomei</i>	Myristicaceae	21	4
140	<i>Myristica malabarica</i>	Myristicaceae	12	2
141	<i>Neolamarckia cadamba</i>	Rubiaceae	3	2
142	<i>Neolitsea zeylanica</i>	Lauraceae	5	2
143	<i>Nothopogia castaneifolia</i>	Anacardiaceae	75	31
144	<i>Ochna obtusata</i>	Ochnaceae	4	2
145	<i>Pavetta sp</i>	Rubiaceae	1	1
146	<i>Phyllanthus emblica</i>	Phyllanthaceae	17	7
147	<i>Pongamia pinnata</i>	Fabaceae	3	1
148	<i>Prunus ceylanica</i>	Rosaceae	1	1
149	<i>Psydrax dicoccos</i>	Rubiaceae	16	11
150	<i>Pterocarpus marsupium</i>	Fabaceae	6	4
151	<i>Pterospermum diversifolium</i>	Malvaceae	5	3

152	<i>Putranjiva roxburghii</i>	Putranjivaceae	3	3
153	<i>Sageraea laurina</i>	Annonaceae	48	12
154	<i>Sapindus trifoliatus</i>	Sapindaceae	1	1
155	<i>Saraca asoca</i>	Fabaceae	86	6
156	<i>Schleichera oleosa</i>	Sapindaceae	39	9
157	<i>Securinega sp</i>	Phyllanthaceae	1	1
158	<i>Semecarpus anacardium</i>	Anacardiaceae	4	1
159	<i>Senegalia catechu</i>	Fabaceae	29	6
160	<i>Solenocarpus indicus</i>	Anacardiaceae	2	2
161	<i>Spondias pinnata</i>	Anacardiaceae	2	2
162	<i>Sterculia foetida</i>	Malvaceae	1	1
163	<i>Sterculia guttata</i>	Malvaceae	9	8
164	<i>Sterculia urens</i>	Malvaceae	1	1

165	<i>Stereospermum colais</i>	Bignoniaceae	38	22
166	<i>Strombosia ceylanica</i>	Olacaceae	12	3
167	<i>Strychnos nux-vomica</i>	Loganiaceae	8	7
168	<i>Symplocos macrophylla</i>	Symplocaceae	89	16
169	<i>Syzygium caryophyllatum</i>	Myrtaceae	18	9
170	<i>Syzygium cumini</i>	Myrtaceae	249	64
171	<i>Syzygium gardneri</i>	Myrtaceae	1	1
172	<i>Syzygium hemisphericum</i>	Myrtaceae	48	12
173	<i>Syzygium rubicundum</i>	Myrtaceae	9	4
174	<i>Tabernaemontana alternifolia</i>	Apocynaceae	116	40
175	<i>Tamilnadia uliginosa</i>	Rubiaceae	3	3
176	<i>Tectona grandis</i>	Lamiaceae	118	13
177	<i>Terminalia bellirica</i>	Combretaceae	60	28

178	<i>Terminalia chebula</i>	Combretaceae	25	11
179	<i>Terminalia elliptica</i>	Combretaceae	298	41
180	<i>Terminalia paniculata</i>	Combretaceae	437	49
181	<i>Tetrameles nudiflora</i>	Tetramelaceae	1	1
182	<i>Tetrapilus dioicus</i>	Oleaceae	125	31
183	<i>Tritaxis glabella</i>	Euphorbiaceae	45	5
184	<i>Vitex altissima</i>	Lamiaceae	7	2
185	<i>Wendlandia thyrsoides</i>	Rubiaceae	2	2
186	<i>Wrightia arborea</i>	Apocynaceae	22	6
187	<i>Wrightia tinctoria</i>	Apocynaceae	23	9
188	<i>Xantolis tomentosa</i>	Sapotaceae	151	37
189	<i>Xylia xylocarpa</i>	Fabaceae	120	16
190	<i>Zanthoxylum rhetsa</i>	Rutaceae	45	15

191	<i>Ziziphus rugosa</i>	Rhamnaceae	1	1
192	<i>Ziziphus xylopyrus</i>	Rhamnaceae	1	1

717

718 **Table S2.** Treatment contrast table showing coefficient estimates and associated 95% CI for the
719 Generalized linear model with Poisson error structure that examined relationship between
720 observed species richness per plot (not including lianas) (≥ 10 cm GBH) and climatic water
721 deficit and land-use categories. Pseudo $R^2 = 0.09$.

Predictor variable	Estimate (95% CI)	p
Intercept: Less-disturbed forests	5.17 (3.41– 6.94)	< 0.001
Historically-disturbed forests	-0.73 (-3.54 – 2.07)	0.61
Repeatedly-disturbed forests	-3.12 (-5.68 – -0.58)	0.016
Climatic water deficit (CWD)	0.002 (0.001 – 0.004)	0.006
CWD \times Historically-disturbed forests	-0.001 (-0.004 – 0.002)	0.62
CWD \times Repeatedly-disturbed forests	-0.003 (-0.006 – -0.0005)	0.02

722

723 **Table S3.** Treatment contrast table showing coefficient estimates and associated 95% CI for the
 724 General linear model with Gaussian error structure that examined the relationship between the
 725 square root of basal area (m^2ha^{-1}) and CWD and land-use categories. $R^2 = 0.51$.

Predictor variable	Estimate (95% CI)	<i>p</i>
Intercept: Less-disturbed forests	13.23 (5.84 – 26.13)	< 0.001
Historically-disturbed forests	2.76 (-1.8×10^{-4} – 0.02)	0.68
Repeatedly-disturbed forests	-29.17 (-44.9 – -19.00)	< 0.001
Climatic water deficit (CWD)	0.01 (-15.9 – 10.36)	0.1
CWD × Historically-disturbed forests	0.003 (-0.048 – -0.019)	0.66
CWD × Repeatedly-disturbed forests	-0.03 (-0.018 – 0.01)	< 0.001

726

727 **Table S4.** Table summarises information (Mean ± SE) on tree density, basal area, canopy cover
 728 and number of cut stems per plot across different land-use categories in low and high elevations.

Structural variables	Repeatedly-disturbed forests (Low)	Repeatedly-disturbed forests (High)	Historically-disturbed forests (Low)	Historically-disturbed forests (High)	Less-disturbed forests (Low)	Less-disturbed forests (High)

Tree density (ha ⁻¹)	1081 (±90.7)	889 (±99.2)	1137 (±81.9)	1565 (±185.8)	1041 (±106.9)	1288 (±112.4)
Basal area (m ² ha ⁻¹)	24.3 (± 2.9)	11.2 (± 2.3)	32.3 (± 3.3)	42.6 (± 2.7)	36.2 (± 2.6)	46.4 (± 5.5)
Canopy cover (%)	39.6 (±2.2)	14.1 (±2.9)	48.6 (±2.4)	45.4 (±1.3)	46.5 (±2.6)	50 (±1.5)
Number of cut stems per plot	15.8 (± 3.3)	24.7 (± 3.3)	6.2 (± 1.3)	2.2 (± 1.3)	5.0 (± 1.1)	10 (± 2.4)

729

730 **Table S5.** Treatment contrast table showing coefficient estimates and associated 95% CI for the
 731 Generalized linear model with binomial error structure that examined relationship between
 732 proportion of evergreen individuals (≥ 10 cm GBH) and climatic water deficit and land-use
 733 categories. Pseudo $R^2 = 0.38$.

Predictor variable	Estimate (95% CI)	<i>p</i>
Intercept: Less-disturbed forests	28.44 (25.28 – 31.74)	< 0.001
Historically-disturbed forests	-7.88 (-11.92 – 3.92)	< 0.001
Repeatedly-disturbed forests	-12.52 (-16.44 – -8.69)	< 0.001
Climatic water deficit (CWD)	0.03 (0.026 – 0.034)	< 0.001

CWD × Historically-disturbed forests	-0.008 (-0.012 – -0.003)	< 0.001
CWD × Repeatedly-disturbed forests	-0.012 (-0.016 – -0.008)	< 0.001

734

735 **Table S6.** Treatment contrast table showing coefficient estimates and associated 95% CI for the
736 Generalized linear model with Poisson error structure that examined the relationship between the
737 number of evergreen tree species (not including lianas) (≥ 10 cm GBH) and climatic water
738 deficit and land-use categories. Pseudo $R^2 = 0.33$.

Predictor variable	Estimate (95% CI)	<i>p</i>
Intercept: Less-disturbed forests	9.96 (7.85 – 12.11)	< 0.001
Historically-disturbed forests	1.16 (-2.36 – 4.69)	0.52
Repeatedly-disturbed forests	-3.99 (-7.41 – -0.56)	0.02
Climatic water deficit (CWD)	0.009 (0.006 – 0.01)	< 0.001
CWD × Historically-disturbed forests	0.0014 (-0.002 – 0.005)	0.48
CWD × Repeatedly-disturbed forests	-0.004 (-0.008 – 0.00005)	0.052

739


740 **Table S7.** Treatment contrast table showing coefficient estimates and associated 95% CI for the
741 Generalized linear model with Poisson error structure that examined the relationship between the

742 number of deciduous tree species (not including lianas) (≥ 10 cm GBH) and climatic water
743 deficit and land-use categories. Pseudo $R^2 = 0.36$.

Predictor variable	Estimate (95% CI)	<i>p</i>
Intercept: Less-disturbed forests	-11.4 (-15.4 – -7.5)	< 0.001
Historically-disturbed forests	1.89 (-3.8 – 7.5)	0.51
Repeatedly-disturbed forests	8.5 (3.8 – 13.3)	< 0.001
Climatic water deficit (CWD)	-0.014 (-0.02 – -0.01)	< 0.001
CWD \times Historically-disturbed forests	0.002 (-0.004 – 0.008)	0.56
CWD \times Repeatedly-disturbed forests	0.009 (0.004 – 0.014)	< 0.001


744

745




746

747 **Figure S1.** Map showing the vegetation plots distributed in our study region in Maharashtra state
748 in India. The three forest categories are spread across low (8–514 m ASL) and high elevations
749 (577–1054 m ASL).



750


751 **Figure S2.** Plot of Non-metric Multidimensional Scaling showing differences in plant
752 composition between low and high elevations and between the different land-use categories.



753

754 **Figure S3.** Hill-Shannon diversity (95% CI) of the three different land-use types, across low and
755 high elevations clearly showing significantly higher diversity of woody plants in less-disturbed
756 low elevation forests. We used the sample-coverage-based method to estimate the species
757 richness of plant communities in the different forest categories (low and high, less-, historically-
758 and repeatedly-disturbed forests) (Roswell et al. 2021). The least sample coverage was in the
759 low-elevation sacred groves (97%). Therefore, we rarefied the diversity measure of all other
760 categories to 97% sample coverage. We bootstrapped the data 50 times to estimate 95%
761 confidence intervals.

762



763

764 **Figure S4.** The stacked bar plot demonstrates that almost 50% of the plant species in this study
765 have not been assessed. It also demonstrates that the proportion of threatened evergreen species

766 is almost twice that of the deciduous species. This study will contribute to generating important
767 information about the not-assessed, Data Deficient and Threatened species for the region.