



An annotated checklist of bryophytes of Europe, Macaronesia and Cyprus

N. G. Hodgetts , L. Söderström , T. L. Blockeel , S. Caspari , M. S. Ignatov , N. A. Konstantinova , N. Lockhart , B. Papp , C. Schröck , M. Sim-Sim , D. Bell , N. E. Bell , H. H. Blom , M. A. Bruggeman-Nannenga , M. Brugués , J. Enroth , K. I. Flatberg , R. Garilleti , L. Hedenäs , D. T. Holyoak , V. Hugonnot , I. Kariyawasam , H. Köckinger , J. Kučera , F. Lara & R. D. Porley

To cite this article: N. G. Hodgetts , L. Söderström , T. L. Blockeel , S. Caspari , M. S. Ignatov , N. A. Konstantinova , N. Lockhart , B. Papp , C. Schröck , M. Sim-Sim , D. Bell , N. E. Bell , H. H. Blom , M. A. Bruggeman-Nannenga , M. Brugués , J. Enroth , K. I. Flatberg , R. Garilleti , L. Hedenäs , D. T. Holyoak , V. Hugonnot , I. Kariyawasam , H. Köckinger , J. Kučera , F. Lara & R. D. Porley (2020) An annotated checklist of bryophytes of Europe, Macaronesia and Cyprus, *Journal of Bryology*, 42:1, 1-116, DOI: [10.1080/03736687.2019.1694329](https://doi.org/10.1080/03736687.2019.1694329)

To link to this article: <https://doi.org/10.1080/03736687.2019.1694329>



© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 28 May 2020.



[Submit your article to this journal](#)



Article views: 2957



[View related articles](#)



[View Crossmark data](#)



Citing articles: 28 [View citing articles](#)

An annotated checklist of bryophytes of Europe, Macaronesia and Cyprus

N. G. Hodgetts^a, L. Söderström^b, T. L. Blockeel^c, S. Caspari^d, M. S. Ignatov^e, N. A. Konstantinova^f, N. Lockhart^g, B. Papp^h, C. Schröckⁱ, M. Sim-Sim^j, D. Bell^k, N. E. Bell^k, H. H. Blom^l, M. A. Bruggeman-Nannenga^m, M. Bruguésⁿ, J. Enroth^o, K. I. Flatberg^p, R. Garilleti^q, L. Hedenäs^r, D. T. Holyoak^s, V. Hugonnot^t, I. Kariyawasam^k, H. Köckinger^u, J. Kučera^v, F. Lara^w and R. D. Porley^x

^aCuillin Views, Isle of Skye, UK; ^bNorwegian University of Science and Technology, Trondheim, Norway; ^cDore, Sheffield, UK; ^dZentrum für Biodokumentation, Schiffweiler, Saarland, Germany; ^eMoscow University and Tsitsin Main Botanica Garden RAS, Moscow, Russia; ^fPolar-Alpine Botanical Garden-Institute RAS, Kirovsk, Russia; ^gNational Parks and Wildlife Service, Dublin, Ireland; ^hHungarian Natural History Museum, Budapest, Hungary; ⁱBiology Centre of Upper Austrian National Museums, Linz, Austria; ^jUniversidade de Lisboa, Faculdade de Ciências, Departamento de Biologia Vegetal, cE3c - Centre for Ecology, Evolution and Environmental Changes, Lisboa, Portugal; ^kRoyal Botanic Garden, Edinburgh, UK; ^lNorwegian Institute of Bioeconomy Research, Bergen, Norway; ^mGriffensteynseplein 23, Zeist, The Netherlands; ⁿFacultat de Biosciències, Universitat Autònoma de Barcelona, Spain; ^oOrganismal and Evolutionary Research Programme, University of Helsinki, Finland; ^pDepartment of Natural History, University Museum, Trondheim, Norway; ^qDepartamento de Botánica y Geología, Facultad de Farmacia, Universidad de Valencia, Spain; ^rSwedish Museum of Natural History, Stockholm, Sweden; ^sRua da Barcoila 274, Cabeçudo, Portugal; ^tConservatoire Botanique National du Massif Central, Chavaniac Lafayette, France; ^uRosegggasse 12, Weisskirchen, Austria; ^vFaculty of Science, University of South Bohemia, České Budějovice, Czech Republic; ^wCentro de Investigación en Biodiversidad y Cambio Global (CIBC-UAM), Madrid, Spain; ^xCerca dos Pomares, Aljezur, Portugal

ABSTRACT

Introduction. Following on from work on the European bryophyte Red List, the taxonomically and nomenclaturally updated spreadsheets used for that project have been expanded into a new checklist for the bryophytes of Europe.

Methods. A steering group of ten European bryologists was convened, and over the course of a year, the spreadsheets were compared with previous European checklists, and all changes noted. Recent literature was searched extensively. A taxonomic system was agreed, and the advice and expertise of many European bryologists sought.

Key results. A new European checklist of bryophytes, comprising hornworts, liverworts and mosses, is presented. Fifteen new combinations are proposed.

Conclusions. This checklist provides a snapshot of the current European bryophyte flora in 2019. It will already be out-of-date on publication, and further research, particularly molecular work, can be expected to result in many more changes over the next few years.

ARTICLE HISTORY

First Published Online
12 May 2020

KEYWORDS

Bryophytes; mosses;
liverworts; hornworts;
Europe; checklist

Introduction

The last published annotated checklist for mosses in Europe was Hill et al. (2006), while for liverworts and hornworts it was Grolle and Long (2000), although a checklist with distributional records was produced by Söderström, Urmi, et al. (2002) and updated by Söderström, Urmi, et al. (2007). In addition, Séneca and Söderström (2009) published a checklist of the Sphagnophyta of Europe and Macaronesia with distribution data. During work for the new European bryophyte Red List, it became apparent that not only was it necessary to publish a new European checklist covering all the bryophytes, but also that this would be relatively little extra work, because spreadsheets of species indicating their distribution by country had been prepared for the Red List, and every effort made to update the nomenclature and taxonomy. A new checklist is also needed because

of the great amount of new molecular work that has been done (and continues to be done) on bryophytes, revealing many new and often unexpected relationships. The nomenclature of liverworts and hornworts in particular has undergone radical changes, as recorded in the recent World Checklist (Söderström, Hagborg, et al. 2016), which, however, did not contain synonyms.

Ironically, the checklist by Hill et al. (2006) was prepared in response to the intention of the European Committee for the Conservation of Bryophytes (ECCB) to produce a new Red List, but the latter had to wait another ten years before being addressed properly, when the ECCB entered into collaboration with the International Union for the Conservation of Nature (IUCN), at which point LIFE funding became available for the Red List project. The Red List has now been

CONTACT N. G. Hodgetts  nick1901@hotmail.co.uk

This article was originally published with a minor error, which has now been corrected in the online version. Please see Correction (<https://doi.org/10.1080/03736687.2020.1812875>).

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

published (Hodgetts et al. 2019), and the names are substantially the same as in this checklist, although there have been some updates and additions to the checklist since the Red List cut-off date of 31 December 2018, and the geographical areas covered by the Red List and the checklist differ slightly (see below).

Methods

Following work on the Red List, funding was secured from the Irish government's National Parks and Wildlife Service to enable author NGH to co-ordinate a new checklist project. A Steering Group was set up consisting of Tom Blockeel, Steffen Caspari, Nick Hodgetts, Misha Ignatov, Nadya Konstantinova, Neil Lockhart, Beata Papp, Christian Schröck, Manuela Sim-Sim and Lars Söderström. Communication has largely been through e-mail, but the Steering Group met once in Budapest, in November 2018, to coincide with the final Red List meeting. After this meeting, NGH led on the moss part of the checklist and author LS led on the liverworts and hornworts.

The existing network of ECCB contacts was crucial to the project, and a large number of bryologists in Europe have contributed, and those making major contributions are included in the authorship.

The first step was to compare the data in the Red List spreadsheets, which had been kept up-to-date taxonomically and nomenclaturally, with that in Hill et al. (2006) and Grolle and Long (2000). Differences between these checklists and current concepts were highlighted and marked for comment. These spreadsheets were then examined closely, species by species, at the Steering Group meeting, with two sub-groups convened, one for mosses, the other for liverworts and hornworts. A team of experts on particular taxonomic groups of mosses was recruited to advise on their areas of expertise. These are listed below and are included in the authorship of this paper:

Neil Bell, Isuru Kariyawasam – Polytrichaceae
 Hans Blom – *Schistidium*
 Ida Bruggeman-Nannenga – Fissidentaceae
 Monserrat Brugués – Funariaceae
 Johannes Enroth – Neckeraceae
 Kjell Ivar Flatberg – Sphagnaceae
 Lars Hedenäs – Amblystegiaceae, Calliergonaceae,
 pleurocarps
 David Holyoak – Bryaceae
 Jan Kučera – Pottiaceae
 Paco Lara, Ricardo Garilleti – Orthotrichaceae
 Ron Porley – *Grimmia* and *Coscinodon*

Specialist advice was either not needed or not available for other groups. It was less crucial to have specialist advice on taxonomic groups of liverworts, as much of the work had been done previously for

the World Checklist (Söderström, Hagborg, et al. 2016). The provisional spreadsheets were also sent to other bryologists for further expert advice (authors DB, VH and HK and those listed in acknowledgements below).

Taxonomic hierarchy

For the liverworts and hornworts, the taxonomic hierarchy follows the Catalogue of Life (Roskov et al. 2019), the online database of the world's known species of animals, plants, fungi and micro-organisms, in which the hierarchy for the hornworts (Anthocerotophyta) and the liverworts (Marchantiophyta) is contributed by the Early Land Plants Today project (Söderström, Hagborg, et al. 2019). The higher classification is substantially the same as the one presented in the bryophyte phylogeny poster by Cole et al. (2019).

It was more complicated to agree on a higher classification of mosses, as current concepts are in a considerable state of flux as more molecular work is done. Pleurocarpous mosses in particular have undergone many changes in recent years, and it is expected that this will continue before relative stability is achieved. Our starting point was Frey and Stech (2009), but it has been necessary to update this treatment with the results of new research. The resultant higher classification is substantially in agreement with the poster produced recently by Cole et al. (2019), which is probably the most accurate representation of the higher classification of mosses as it is currently understood, although there are some differences. For example, Cole et al. (2019) place *Hypopterygium* in the order Hypopterygiales, while we have it in the Hypopterygiaceae within the order Hookeriales.

The final higher classification presented in this checklist is not offered as a definitive view, and it can be expected that there will be further changes with more research.

List of taxa

As Hill et al. (2006) and Grolle and Long (2000) were the starting points for this checklist, taxa that have remained the same since these publications are mostly listed without further comment. All synonyms that have been used since these publications are included. In addition, some synonyms whose synonymy predates these publications, but have continued to be in common usage, have been included. The liverworts in particular have undergone some radical taxonomic and nomenclatural changes since Grolle and Long (2000). Most of these are recorded, but without synonyms, in the World Checklist (Söderström, Hagborg, et al. 2016). The synonymy for the

liverworts is therefore considerably more extensive than that for the mosses.

Hybrid taxa are mentioned only if they have been treated in recent literature.

Author citations

We have tried to follow the International Plant Names Index (www.ipni.org) for author citations, although no doubt we will not always have succeeded. In practice, most author citations, except for very recently treated taxa, are the same as in Hill et al. (2006) for the mosses and Söderström, Hagborg, et al. (2016) for the liverworts and hornworts, but a small number of citations have been corrected.

Bryum s.lat.

It is necessary to make some remarks about *Bryum* s.lat. as the current position is confusing and unsatisfactory. The reviews of world-wide scope by Ochi (1959, 1970, 1972, 1980, 1981, etc.) placed all European Bryaceae in genus *Bryum* with several subgenera, on the basis of extensive morphological studies. Other workers treated his subgenera as genera, *Anomobryum*, *Brachymenium*, *Bryum*, *Plagiobryum*, *Rhodobryum* (e.g. Holyoak in Hill et al. 2006, Guerra, Gallego, et al. 2010). Molecular data (Pedersen and Hedenäs 2002, 2003; Pedersen, Cox and Hedenäs 2003; Pedersen, Holyoak and Newton 2007) showed this generic treatment to be ill-founded, since several of the widely recognised subgenera/genera were shown to be polyphyletic. It also became apparent that morphology in Bryaceae often provides entirely misleading evidence of phylogeny: Examples of this include the morphologically striking '*Plagiobryum*' *zieri* nested within a clade containing e.g. *Ptychostomum pseudotriquetrum*, the superficially similar small tuber-bearing '*Erythrocarpa*' belonging in at least two distinct genera, and some *Bryum dichotomum* being difficult to separate morphologically from *Ptychostomum imbricatum* (syn. *Bryum caespiticium*). Thus, morphological evolution in Bryaceae appears to be 'decoupled' from phylogeny (Holyoak and Pedersen 2007). Generic allocations based on morphology alone are therefore prone to error, some through convergent evolution of distant relatives, others by rapid change among close relatives. These findings have not been refuted by subsequent studies and parts of the sequence data used have been confirmed and supplemented by other workers (Guerra, Jiménez-Martínez, Cano, Brugués 2008; Guerra, Jiménez-Martínez, Cano, Jiménez-Fernández 2011; Wang and Zhao 2009; Bell, Long and Hollingsworth

2013). Thus the generic taxonomy adopted here is based mainly on molecular data.

Nevertheless, Spence (2014) and e.g. Ochyra, Plášek, et al. (2018) have largely ignored the implications of the molecular results and used morphology almost exclusively for their generic classifications. Parts of the classifications by Spence are directly contradicted by molecular data (cf. Shaw 2014: p.10–11); much of the remainder is at risk of being erroneous due to convergent evolution.

It is frustrating that a lack of further molecular data still prevents the allocation of some species to segregate genera, but this problem is declining as more species are sequenced. Indeed, some unpublished DNA barcoding and nuclear genomic sequencing data from one of the authors (DB), along with some last-minute sequencing by another (MSI), and interpretation by a third (DTH), have resulted in a clarification of generic affinity for several species. Further sequencing work should see significant increases in the data available within the next few years. The alternative of reverting to an enormous and still partly polyphyletic genus *Bryum* thus seems to be a poorer option. It would also be out of step with the rapidly improving classifications of some other moss families. Currently recognised European genera of Bryaceae in this checklist are *Anomobryum*, *Brachymenium*, *Bryum*, *Imbribryum*, *Ptychostomum* and *Rhodobryum*. We do not recognise *Gemmabryum*, *Osculatia*, *Plagiobryoides*, *Plagiobryum* or *Rosulabryum*.

Area covered

For the purposes of this checklist, Europe is regarded in the same way as it was in Hill et al. (2006), but with the addition of Cyprus. Thus, the following territories at the margins of Europe are included:

All of Russia west of the Ural watershed and to the north of the Caucasus watershed;
 Western Kazakhstan;
 Iceland;
 Svalbard;
 Jan Mayen;
 Novaya Zemlya;
 Franz Josef Land;
 Malta;
 all the Greek Islands;
 all other Mediterranean islands belonging to European countries (e.g. Pantelleria);
 Turkey west of the Bosphorus (i.e. Turkey-in-Europe);
 Azores;
 Canary Islands;
 Madeira.

Table 1. Bryophytes introduced to Europe

Species	Location	Native range
Liverworts		
<i>Heteroscyphus fissistipus</i>	garden in Ireland	Australia, New Zealand
<i>Lophocolea bispinosa</i>	gardens and naturalised in Britain, Ireland & Spain	Australia
<i>Lophocolea brookwoodiana</i>	cemetery in southern England	Unknown
<i>Lophocolea semiteres</i>	naturalised in western Europe	Southern Hemisphere
<i>Sphaerocarpos stipitatus</i>	Portugal	Southern Hemisphere
<i>Tricholepidozia lindenberghii</i> var. <i>lindenberghii</i>	gardens in Wales	New Zealand, Philippines and Fiji
<i>Tricholepidozia tetradactyla</i>	gardens in Britain	New Zealand
Mosses		
<i>Achrophyllum dentatum</i>	garden in England	Southern Hemisphere
<i>Atrichum crispum</i>	naturalised in Britain & Ireland	Eastern North America
<i>Calomnion complanatum</i>	gardens in Ireland	Australia, New Zealand
<i>Calyptrochaeta apiculata</i>	naturalised in southern England & Ireland	Southern Hemisphere
<i>Campylopus introflexus</i>	naturalised & widespread in much of Europe	Southern Hemisphere
<i>Dicranoloma menziesii</i>	on tree ferns in garden in Ireland	Southern Hemisphere
<i>Hennediella macrophylla</i>	naturalised in Britain	New Zealand
<i>Hennediella stanfordensis</i>	naturalised in Britain, Ireland, France & Greece	California, Mexico & Australia
<i>Hypopterygium tamarisci</i>	gardens in Portugal & Italy	Tropical Africa & America
<i>Leptodontium proliferum</i>	bowling green in northern England	Southern Hemisphere
<i>Leptotheca gaudichaudii</i>	gardens in southern England & Ireland	Southern Hemisphere
<i>Myuroclada maximowiczii</i>	garden in southern Russia	Eastern Asia, Alaska
<i>Orthodontium lineare</i>	naturalised & widespread in western and central Europe	Southern Hemisphere
<i>Sematophyllum adnatum</i>	naturalised in northern Italy, Switzerland & Hungary	America
<i>Thuidiopsis sparsa</i>	former park in Madeira	Southern Hemisphere
<i>Tortula amplexa</i>	naturalised in England	Western North America

The decision to include Cyprus was made because it is politically part of Europe, although geographically closer to Asia. In practice, its inclusion or exclusion is fairly academic, as there are no bryophyte taxa known that have their only 'European' occurrence in Cyprus, except for the dubious species *Funaria anomala*, and possibly *Grimmia ungeri*, which has been considered to be endemic to Cyprus by some authors.

The area covered differs slightly from that covered by the Red List (IUCN 2019), which excludes the Caucasus. Both the checklist and the Red List exclude Cape Verde (Cabo Verde).

Composition of the bryophyte flora

The moss flora of Europe, as described in this checklist, comprises 1392 species (including the 3 hybrids), plus 45 subspecies and 104 varieties. There are 494 species of liverworts, with 35 subspecies and 56 varieties, and 8 species of hornworts, with no subspecies or varieties, giving a total of 1892 species of bryophytes in Europe, plus 80 subspecies and 160 varieties. These figures only include subspecies and varieties where species are represented by more than one subspecies or variety in Europe. There are 187 endemic taxa, with 56 of these being endemic to the Macaronesian islands. The level at which a taxon is recognised is always to some extent arbitrary, even with the benefit of molecular data. In general, we have included subspecies and varieties if they have been validly published and there is no good reason to reject them. However, this self-imposed guidance has been considerably modified according to

expert opinion from many of the authors and those listed in the acknowledgements. It is indicated in the footnotes where disagreement exists.

There are 23 obvious introductions that are now more or less naturalised in at least some part of Europe. These are listed in Table 1. Taxa are considered introductions only if they have clearly been introduced by human agency. Other species that might be regarded as introductions, that may for example be chance ephemeral colonists from wind-blown spores, are regarded as native, as they apparently arrived through natural agency. Whether any particular taxon is native or not to Europe is of course arguable. To some extent the native status of taxa matters less for bryophytes than it does for vascular plants, since their often very efficient dispersal mechanisms mean that many species can potentially occur in suitable habitats worldwide. Several further species were listed by Hill et al. (2006) as non-native (*Anomobryum apiculatum*, *Bryum valparaisense*, *Chenia leptophylla*, *Splachnobryum obtusum*, *Syntrichia bogotensis*, *Tortula bogosica*, *Tortula bolanderi*), but there is significant doubt about their non-native status, and they are not listed in Table 1.

New validations

Three liverwort taxa were published invalidly and therefore require validation.

Riccia sorocarpa subsp. *arctica* R.M.Schust. ex Köckinger & L.Söderstr., **subsp. nov.** [*Riccia sorocarpa* var. *arctica* R.M.Schust., J. Hattori Bot. Lab. 71: 274, 1992, *nom. inval.* ICN Art. 37.7]. Holotype: Greenland, Kangerdlugssuak, Inglefield Bay, NW. Greenland, R.M.Schuster

45831 (F-C0000489F; <http://emuweb.fieldmuseum.org/botany/Display.php?irn=9894&QueryPage=%2Fbotany%2Fdetailed.php>). The description in Schuster (1992a) together with the current data on type specimen validates the name. This subspecies was described from Greenland but also occurs in the Alps (Köckinger 2017).

Riccia sorocarpa subsp. ***erythrophora*** R.M.Schust. ex Konstant. & L.Söderstr., **subsp. nov.** [*Riccia sorocarpa* subsp. *erythrophora* R.M.Schust., J. Hattori Bot. Lab. 71: 274, 1992, *nom. inval.* ICBN Art. 37.7]. Holotype: California: 1.5 km W of Potrero, San Diego Co., 1982, R.M.Schuster 82-146 (F-C1049209F; <http://emuweb.fieldmuseum.org/botany/Display.php?irn=2346544&QueryPage=%2Fbotany%2Fdetailed.php>; type originally filed under *Riccia nigrella*). The description in Schuster (1992a) together with the current data on the type specimen validate the name. It was described from California but is also known from southern Russia (Konstantinova and Doroshina 2011).

Scapania paludicola var. ***rotundiloba*** R.M.Schust. ex Konstant. & L.Söderstr., **var. nov.** Holotype: "West Greenland: east end of Nûk Peninsula, Alángup nunâ, 68°42'N., 52°10'W. Schuster & Damsholt RMS 70-2878" (F-C0171445F; <http://emuweb.fieldmuseum.org/botany/Display.php?irn=2145467&QueryPage=%2Fbotany%2Fdetailed.php>) [*Scapania paludosa* var. *rotundiloba* R.M.Schust., Hepat. Anthocerotae N. Amer. 3: 519, 1974, *nom. inval.* ICBN Art. 37.2; based on more than one gathering]. The description in Schuster (1974: 519), together with the type selected here, validates the name.

New combinations

New combinations are made by individual authors.

Mesoptychia gillmanii var. ***laxa*** (Schiffn. ex Burrell) L.Söderstr. **comb. nov.** Basionym: *Lophozia schultzii* var. *laxa* Schiffn. ex Burrell, J. Bot. 49: 217, 1911 (see Burrell 1911).

Anomobryum apiculatum (Schwägr.) D.Bell & Holyoak **comb. nov.** Basionym: *Bryum apiculatum* Schwägr., Sp. Musc. Frond., Suppl. 1, sect. 2, p. 102, pl. 72 [top], 1816 (see Schwägrichen 1816).

Anomobryum notarisii (Mitt.) D.Bell & Holyoak **comb. nov.** Basionym: *Bryum notarisii* Mitt., J. Linn. Soc., Bot., 7, p. 3, 1864 (see Mitten 1864).

Didymodon glaucus subsp. ***verbanus*** (W.E.Nicholson & Dixon) Jan Kučera **comb. nov.** Basionym: *Eucladium*

verbanum W.E.Nicholson & Dixon, Revue Bryologique 39: 89. 1912 (see Dixon 1912).

Fissidens subgenus ***Pachyfissidens*** (Müll.Hal.) L.Söderstr. & A.Hagborg, **comb. nov.** Basionym: *Fissidens* sect. *Pachyfissidens* Müll.Hal., Syn. Musc. Frond. 1: 45, 1849 [1848] (see Müller 1849). Note: The subgenus has frequently been cited from Kindberg (1897), but in that paper *Pachyfissidens* is an unranked taxon. We have not been able to localise any valid combination at subgeneric rank and thus propose it here.

Imbribryum subapiculatum (Hampe) D.Bell & Holyoak **comb. nov.** Basionym: *Bryum subapiculatum* Hampe, Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn, ser. 3, 4, p. 51, 1872 (see Hampe 1872).

Imbribryum tenuisetum (Limpr.) D.Bell & Holyoak **comb. nov.** Basionym: *Bryum tenuisetum* Limpr., Jahresber. Schles. Ges. Vaterl. Cult., 74(2b), p. 4, 1897 (see Limpricht 1897).

Mnioideae L.Söderstr., N.G.Hodgetts & Ignatov, **comb. et stat. nov.**

Basionym: Trib. Mnieae Müll., Syn. Musc. Frond.: 152, 1858 (Müller 1858) 'Mnioideae'.

≡ Mnioideae A.J.E.Sm., Moss Flora Brit. Ireland, 2nd ed.: 615, 2004 (Smith 2004), *nom. inval.* (ICN Art. 39.1; no Latin description).

≡ Mnioideae Stech & W.Frey, Syll. Pl. Fam. 3: 196 (Frey & Stech 2009), *nom. inval.* (ICN Art. 39.1; no Latin description).

≡ Mnioideae T.J.Kop., Acta Bryolich. Asiatica 5: 39 (Koponen 2014), *nom. inval.* (ICN Art. 38.1(a); no description).

Type genus: *Mnium* Hedw.

Note: The subfamily Mnioideae has been recognized for a long time but as far as we know never been validated. Most authors seem to assume it is an autonym under Mniaceae (e.g. Smith 2004; Frye and Stech 2009). However, autonoms only exist under genera and species (ICN Art. 22.1 and 26.1; Turland et al. 2018) so a formal description of the taxon is needed. No description before 2011 is in Latin, as required then, or refers to Müller (1858) to make it a valid combination. Latin descriptions have been unnecessary since 2011, but we have not found any recent description of any sort that fulfils all the criteria. Frey and Stech (2009) give a good description, but they seem to treat all subfamilies based on the family name as autonoms, contrary to the code (Turland et al. 2018). A reference to Müller (1858), who treats the taxon as a tribus (although with an

incorrect ending according to the current code), validates the name.

Platyhypnum tatrense (Váňa) Hedenäs & Ignatov **comb. nov.** Basionym: *Ochyraea tatrensis* Váňa, J. Bryol. 14: 261. f. 1-3. 1986 (see Váňa 1986).

Ptychostomum austriacum (Köckinger, Holyoak & Suanjak) D.Bell & Holyoak **comb. nov.** Basionym: *Bryum austriacum* Köckinger, Holyoak & Suanjak, J. Bryol. 35(1), p. 57, 2013 (see Köckinger, Holyoak, et al. 2013).

Ptychostomum cellulare (Hook.) D.Bell & Holyoak **comb. nov.** Basionym: *Bryum cellulare* Hook. in Schwägr., Sp. Musc. Frond., Suppl. 3, vol. 1, sect. 1, p. 9, pl. 214a, fig. a, 1827 (see Schwägrichen 1827).

Ptychostomum elegans (Nees.) D.Bell & Holyoak **comb. nov.** Basionym: *Bryum elegans* Nees, in Bridel, Bryol. Univ. 1 (2): p. 849. 1827 (see Bridel 1827).

Ptychostomum minii (Podp. ex Guim.) D.Bell & Holyoak **comb. nov.** Basionym: *Bryum minii* Podp. ex Guim., Rev. Bryol. Lichénol., 8, p. 112–114, 1935.

Ptychostomum schleicheri (DC.) J.R.Spence ex D.Bell & Holyoak **comb. nov.** Basionym: *Bryum schleicheri* DC. in Lam., Fl. Franç., 6, p. 226, 1815 (see Lamarck 1815).

Ptychostomum schleicheri* var. *latifolium (Schwägr.) D.Bell & Holyoak **comb. nov.** Basionym: *Mnium latifolium* Schwägr., Spec. Musc. Suppl. 1(2), p. 138, 1816 ≡ *Bryum schleicheri* var. *latifolium* (Schwägr.) Schimp.

Taxonomic synopsis

This synopsis lists the higher taxa relevant to Europe down to family level.

Anthocerotophyta Stotler & Crand.-Stotl.

Anthocerotopsida de Bary ex Jancz.

Anthocerotidae Rosenv.

Anthocerotales Limpr.

Anthocerotaceae Dumort.

Phymatocerotales R.J.Duff

Phymatocerotaceae R.J.Duff

Notothylatidae R.J.Duff

Notothyladales Hyvönen & Piippo

Notothyladaceae Müll.Frib. ex Prosk.

Phaeocerotideae Häsael

Marchantiophyta Stotler & Crand.-Stotl.

Haplomitriopsida Stotler & Crand.-Stotl.

Haplomitriidae Stotler & Crand.-Stotl.

Calobryales Hamlin

Haplomitriaceae Dědeček

Jungermanniopsida Stotler & Crand.-Stotl.

Jungermanniidae Engl.

Jungermanniales H.Klinggr.

Cephaloziineae Schljakov.

Adelanthaceae Grolle

Anastrophyllaceae L.Söderstr., De Roo & Hedd.

Cephaloziaceae Mig.

Cephaloziellaceae Douin

Lophoziaceae Cavers

Scapaniaceae Mig.

Jungermanniineae R.M.Schust. ex Stotler & Crand.-Stotl.

Acrobolbaceae E.A.Hodgs.

Antheliaceae R.M.Schust.

Arnelliaceae Nakai

Calypogeiaceae Arnell

Endogemmataceae Konstant.

Geocalycaceae H.Klinggr.

Gymnomitriaceae H.Klinggr.

Harpanthaceae Arnell

Hygrobiiellaceae Konstant. & Vilnet

Jungermanniaceae Rchb.

Saccogynaceae Heeg

Solenostomataceae Stotler & Crand.-Stotl.

Southbyaceae Váňa, Crand.-Stotl., Stotler & D.G.Long

Lophocoleineae Schljakov

Blepharostomataceae W.Frey & M.Stech

Herbertaceae Müll.Frib. ex Fulford & Hatcher

Lepidoziaceae Limpr.

Lophocoleaceae Vanden Berghen

Mastigophoraceae R.M.Schust.

Plagiochilaceae Müll.Frib.

Trichocoleaceae Nakai

Myliineae J.J.Engel & Braggins ex Crand.-Stotl.

Myliaceae Schljakov

Porellales Schljakov

Jubulineae Müll.Frib.

Frullaniaceae Lorch

Jubulaceae H.Klinggr.

Lejeuneaceae Cavers

Porellineae R.M.Schust.

Porellaceae Cavers

Radulineae R.M.Schust.

Radulaceae Müll.Frib.

Ptilidiales Schljakov

Ptilidiaceae H.Klinggr.

Metzgeriidae Barthol.-Began

Metzgeriales Chalaud

Aneuraceae H.Klinggr.

Metzgeriaceae H.Klinggr.

Pleuroziales Schljakov

Pleuroziaceae Müll.Frib.

Pelliidae He-Nygrén

Fossombroniales Schljakov

- Calyculariineae He-Nygrén, Juslén, Ahonen,
 Glenny & Piippo
 Calyculariaceae He-Nygrén, Juslén,
 Ahonen, Glenny & Piippo
 Fossombroniineae R.M.Schust. ex Stotler &
 Crand.-Stotl.
 Fossombroniaceae Hazsl.
 Petalophyllaceae Stotler & Crand.-Stotl.
 Pallaviciniales W.Frey & M.Stech
 Pallaviciniineae R.M.Schust.
 Moerckiaceae K.I.Goebel ex Stotler &
 Crand.-Stotl.
 Pallaviciniaceae Mig.
 Pelliales He-Nygrén
 Pelliaceae H.Klinggr.
 Marchantiopsida Cronquist, Takht. & W.Zimm.
 Blasiidae He-Nygrén
 Blasiales Stotler & Crand.-Stotl.
 Blasiaceae H.Klinggr.
 Marchantiidae Engl.
 Lunulariales H.Klinggr.
 Lunulariaceae H.Klinggr.
 Marchantiales Limpr.
 Aytoniaceae Cavers
 Cleveaceae Cavers
 Conocephalaceae Müll.Frib. ex Grolle
 Corsiniaceae Engl.
 Cyathodiaceae Stotler & Crand.-Stotl.
 Dumortieraceae D.G.Long
 Exormothecaceae Müll.Frib. ex Grolle
 Marchantiaceae Lindl.
 Oxymitracae Müll.Frib. ex Grolle
 Ricciaceae Rchb.
 Targioniaceae Dumort.
 Sphaerocarpales Cavers
 Riellaceae Engl.
 Sphaerocarpaceae Heeg
 Bryophyta Schimp.
 Sphagnophytina Doweld
 Sphagnopsida Schimp.
 Sphagnales Limpr.
 Sphagnaceae Dumort.
 Bryophytina Engler
 Andreaeopsida J.H.Schaffn.
 Andreaeidae Engl.
 Andreaeales Limpr.
 Andreaeaceae Dumort.
 Oedipodiopsida Goffinet & W.R.Buck
 Oedipodiales Goffinet & W.R.Buck
 Oedipodiaceae Schimp.
 Tetrphidopsida Goffinet & W.R.Buck
 Tetrphidales M.Fleisch.
 Tetrphidaceae Schimp.
 Polytrichopsida Doweld
 Polytrichales M.Fleisch.
 Polytrichaceae Schwägr.
 Bryopsida Pax
 Buxbaumiidae Doweld
 Buxbaumiales M.Fleisch.
 Buxbaumiaceae Schimp.
 Diphysciidae Ochyra
 Diphysciales M.Fleisch.
 Diphysciaceae M.Fleisch.
 Timmiidae Ochyra
 Timmiales Ochyra
 Timmiaceae Schimp.
 Encalyptidae Ochyra, Żarnowiec & Bedn.-Ochyra
 Encalyptales Dixon
 Encalyptaceae Schimp.
 Funariidae Ochyra
 Funariales M.Fleisch.
 Funariaceae Schwägr.
 Disceliaceae Schimp.
 Gigaspermidae M.Stech & W.Frey
 Gigaspermales Goffinet, Wickett, O.Werner, Ros,
 A.J.Shaw & C.J.Cox
 Gigaspermaceae Lindb.
 Dicranidae Doweld
 Catoscopiales Ignatov & Ignatova
 Catosciaceae Broth.
 Dicranales H.Philib ex M.Fleisch.
 Timmiellaceae Y.Inoue & H.Tsubota
 Distichiaceae Schimp.
 Hymenolomataceae Ignatov & Fedosov
 Flexitrichaceae Ignatov & Fedosov
 Bryoxiphiaceae Besch.
 Archidiaceae Schimp.
 Micromitriaceae Smyth ex Goffinet &
 Budke
 Leucobryaceae Schimp.
 Amphidiaceae M.Stech
 Aongstroemiaceae De Not.
 Dicranellaceae M.Stech
 Fissidentaceae Schimp.
 Dicranaceae Schimp.
 Calymperaceae Lindb.
 Rhabdoweisiaceae Limpr.
 Schistostegaceae Schimp.
 Bruchiaceae Schimp.
 Ditrichaceae Limpr.
 Pottiaceae Schimp.
 Grimmiiales M.Fleisch.
 Saelaniaceae Ignatov & Fedosov
 Seligeriaceae Schimp.
 Ptychomitriaceae Schimp.
 Grimmiaceae Arn.
 Bryidae Engl.
 Hedwigiales Ochyra
 Hedwigiaceae Schimp.
 Bartramiales D.Quandt, N.E.Bell & M.Stech
 Bartramiaceae Schwägr.
 Splachnales Ochyra
 Splachnaceae Grev. & Arn.
 Meesiaceae Schimp.

Bryales Limpr.
 Bryaceae Schwägr.
 Mniaceae Schwägr.
 Orthotrichales Dixon
 Orthotrichaceae Arn.
 Orthodontiales N.E.Bell, A.E.Newton & D.Quandt
 Orthodontiaceae Goffinet
 Aulacomniales N.E.Bell, A.E.Newton & D.Quandt
 Aulacomniaceae Schimp.
 Rhizogoniales Goffinet & W.R.Buck
 Rhizogoniaceae Broth.
 Hookeriales M.Fleisch.
 Hypopterygiaceae Mitt.
 Daltoniaceae Schimp.
 Hookeriaceae Schimp.
 Leucomiaceae Broth.
 Pilotrichaceae Kindb.
 Hypnales W.R.Buck & Vitt
 Fontinalaceae Schimp.
 Plagiotheciaceae M.Fleisch.
 Fabroniaceae Schimp.
 Pterigynandraceae Schimp.
 Habrodontaceae Schimp.
 Climaciaceae Kindb.
 Myriniaceae Schimp.
 Amblystegiaceae G.Roth
 Calliergonaceae Vanderp., Hedenäs,
 C.J.Cox & A.J.Shaw
 Scorpidiaceae Ignatov & Ignatova
 Leskeaceae Schimp.
 Pseudoleskeaceae Schimp.
 Pseudoleskeellaceae Ignatov & Ignatova
 Thuidiaceae Schimp.
 Brachytheciaceae Schimp.
 Hypnaceae Schimp.
 Callicladiaceae Jan Kučera & Ignatov
 Taxiphyllaceae Ignatov
 Pylaisiadelphaceae Goffinet & W.R.Buck
 Jocheniaceae Jan Kučera & Ignatov
 Stereodontaceae Hedenäs, Schlesak &
 D.Quandt
 Pylaisiaceae Schimp.
 Sematophyllaceae Broth.
 Hylocomiaceae M.Fleisch.
 Rhytidiaceae Broth.
 Entodontaceae Kindb.
 Cryphaeaceae Schimp.
 Leucodontaceae Schimp.
 Antitrichiaceae Ignatov & Ignatova
 Neckeraceae Schimp.
 Heterocladellaceae Ignatov & Fedosov
 Lembophyllaceae Broth.
 Echinodiaceae Broth.
 Myuriaceae M.Fleisch.
 Anomodontaceae Kindb.

List of Taxa

Subdivisions ('-phytina'), subclasses ('-ideae') and suborders ('-ineae'), shown in the taxonomic synopsis above, are not listed in the main checklist, but subfamilies ('-oideae') are included. Tribes and subtribes are included in the Lejeuneaceae, as they are fairly useful divisions.

Hornworts

ANTHOCEROTOPHYTA

ANTHOCEROTOPSIDA DE BARY EX JANCZ.

Anthocerotales Limpr.

Anthocerotaceae Dumort.

1 **Anthoceros** L.

- 1 **A. agrestis** Paton [*Anthoceros punctatus* subsp. *agrestis* (Paton) Damsh., *Anthoceros punctatus* var. *douinii* (R.M.Schust.) Damsh.]

- 2 **A. caucasicus** Steph.

- 3 **A. neesii** Prosk.

- 4 **A. punctatus** L.

Phymatocerotales R.J.Duff

Phymatocerotaceae R.J.Duff

2 **Phymatoceros** Stotler

- 1 **P. bulbiculosus** (Brot.) Stotler, W.T.Doyle & Crand.-Stotl. [*Phaeoceros bulbiculosus* (Brot.) Prosk.]

Notothyladales Hyvönen & Piippo

Notothyladaceae Müll.Frib. ex Prosk.

Notothyloideae Grolle

3 **Notothylas** Sull.

subgenus *Notothylas*

- 1 **N. orbicularis** (Schwein.) Sull.

Phaeocerotideae Hässel

4 **Phaeoceros** Prosk.

- 1 **P. carolinianus** (Michx.) Prosk. [*Phaeoceros laevis* subsp. *carolinianus* (Michx.) Prosk.]

- 2 **P. laevis** (L.) Prosk.

Liverworts

MARCHANTIOPHYTA

HAPLOMITRIOPSIDA STOTLER & CRAND.-STOTL.

Calobryales Hamlin

Haplomitriaceae Dědeček

1 **Haplomitrium** Nees

subgenus *Haplomitrium*

section *Haplomitrium*

- 1 **H. hookeri** (Lyell ex Sm.) Nees

a var. *hookeri*¹

JUNGERMANNIOPSIDA STOTLER & CRAND.-STOTL.

Jungermanniales H.Klinggr.

Adelanthaceae Grolle

Adelanthoideae K.Feldberg

2 *Adelanthus* Mitt.section *Lindenbergiani* Grolle1 *A. lindenbergianus* (Lehm.) Mitt.3 *Pseudomarsupidium* Herzog1 *P. decipiens* (Hook.) Grolle [*Adelanthus decipiens* (Hook.) Mitt.]

Jamesonielloideae Inoue

4 *Syzygiella* Spruce [*Jamesoniella* (Spruce) F.Lees, *Crossogyna* (R.M.Schust.) Schljakov]subgenus *Syzygiella*1 *S. autumnalis* (DC.) K.Feldberg, Váňa, Hentschel & Heinrichs [*Jamesoniella autumnalis* (DC.) Steph., *Crossogyna autumnalis* (DC.) Schljakov]2 *S. rubricaulis* (Nees) Steph. [*Jamesoniella rubricaulis* (Nees) Grolle]

Anastrophyllaceae L.Söderstr., De Roo & Hedd.

5 *Anastrepta* (Lindb.) Schiffn.1 *A. orcadensis* (Hook.) Schiffn.6 *Anastrophyllum* (Spruce) Steph.1 *A. alpinum* Steph.²2 *A. assimile* (Mitt.) Steph. [*Anastrophyllum assimile* var. *nardioides* (Lindb.) Damsh.]3 *A. donnianum* (Hook.) Steph.4 *A. joergensenii* Schiffn.5 *A. michauxii* (F.Weber) H.Buch7 *Barbilophozia* Loeskesubgenus *Barbilophozia*1 *B. barbata* (Schmidel ex Schreb.) Loeske [*Lophozia barbata* (Schmidel ex Schreb.) Dumort.]2 *B. hatcheri* (A.Evans) Loeske [*Lophozia hatcheri* (A.Evans) Steph.]3 *B. lycopodioides* (Wallr.) Loeske [*Lophozia lycopodioides* (Wallr.) Cogn.]4 *B. rubescens* (R.M.Schust. & Damsh.) Kartt. & L.Söderstr. [*Lophozia rubescens* R.M.Schust. & Damsh.]subgenus *Sudeticae* (Schljakov) L.Söderstr., De Roo & Hedd. [*Pseudolophozia* Konstant. & Vilnet]5 *B. sudetica* (Nees ex Huebener) L.Söderstr., De Roo & Hedd. [*Lophozia alpestris* auct. non (Schleich. ex F.Weber) A.Evans, *Lophozia sudetica* (Nees ex Huebener) Grolle, *Pseudolophozia sudetica* (Nees ex Huebener) Konstant. & Vilnet, *Lophozia sudetica* var. *anomala* (Schljakov) Schljakov, *Lophozia debiliformis* R.M.Schust. & Damsh., *Pseudolophozia debiliformis* (R.M.Schust. & Damsh.) Konstant. & Vilnet, *Lophozia debiliformis* var. *concolor* R.M.Schust. & Damsh.]³8 *Biantheridion* (Grolle) Konstant. & Vilnet1 *B. undulifolium* (Nees) Konstant. & Vilnet [*Jamesoniella undulifolia* (Nees) Müll.Frib.]9 *Crossocalyx* Meyl1 *C. hellerianus* (Nees ex Lindenb.) Meyl. [*Anastrophyllum hellerianum* (Nees ex Lindenb.) R.M.Schust.]10 *Gymnocolea* (Dumort.) Dumort.1 *G. borealis* (Frisvoll & Moen) R.M.Schust.2 *G. fascinifera* Potemkin⁴3 *G. inflata* (Huds.) Dumort.a subsp. *acutiloba* (Schiffn.) R.M.Schust. ex L.Söderstr. & Váňa [*Gymnocolea acutiloba* (Schiffn.) Müll.Frib.]⁵b subsp. *inflata* [*Gymnocolea inflata* var. *heterostipa* (Carringt. ex Spruce) Müll.-Frib.]11 *Isopaches* H.Buch1 *I. alboviridis* (R.M.Schust.) Schljakov⁶2 *I. bicrenatus* (Schmidel ex Hoffm.) H.Buch [*Lophozia bicrenata* (Schmidel ex Hoffm.) Dumort.]3 *I. decolorans* (Limpr.) H.Buch [*Lophozia decolorans* (Limpr.) Steph.]12 *Neoorthocaulis* L.Söderstr.1 *N. attenuatus* (Mart.) L.Söderstr., De Roo & Hedd. [*Barbilophozia attenuata* (Mart.) Loeske, *Orthocaulis attenuatus* (Mart.) A.Evans, *Lophozia attenuata* (Mart.) Dumort.]¹*Haplomitrium hookeri* var. *minutum* (E.O.Campb.) Barthol.-Began occurs in Japan and New Zealand.²*Anastrophyllum alpinum* was synonymised with *Anastrophyllum joergensenii* by Grolle (1964) but reinstated at species level by Long et al. (2006).³*Lophozia debiliformis* was not recognised in Söderström, Hagborg, et al. (2016), following Söderström, De Roo, et al. (2010), although it was sometimes recognised in later publications. It was nested within *Barbilophozia sudetica* in the molecular studies by de Roo et al. (2007) and Vilnet, Konstantinova, et al. (2010) although neither type material nor material from the type locality was included. The 'taxon' is common in the mountains of Europe (e.g. in Scandinavia, the Alps) and elsewhere and may be a modification of harsh environments.⁴*Gymnocolea fascinifera* was described from Yamal Peninsula and recently recorded for European Russia from Komi Republic by Potemkin (2008).⁵The taxonomic value of *Gymnocolea inflata* subsp. *acutiloba* is unclear. A detailed study is needed to clarify whether this is an extreme form of *Gymnocolea inflata* s.str.⁶*Isopaches alboviridis* was recently recorded from European Russia in Leningrad (Potemkin and Rozantseva 2015) and Murmansk Provinces (Borovichev 2008).

- 2 **N. binsteadii** (Kaal.) L.Söderstr., De Roo & Hedd. [*Barbilophozia binsteadii* (Kaal.) Loeske, *Orthocaulis binsteadii* (Kaal.) H.Buch, *Lophozia binsteadii* (Kaal.) A.Evans]
- 3 **N. floerkei** (F.Weber & D.Mohr) L.Söderstr., De Roo & Hedd. [*Barbilophozia floerkei* (F.Weber & D.Mohr) Loeske, *Orthocaulis floerkei* (F.Weber & D.Mohr) H.Buch, *Lophozia floerkei* (F.Weber & D.Mohr) Schiffn.]
- 4 **N. hyperboreus** (R.M.Schust.) L.Söderstr., De Roo & Hedd. [*Barbilophozia hyperborea* (R.M.Schust.) Potemkin, *Orthocaulis hyperboreus* (R.M.Schust.) Konstant.]
- 13 **Orthocaulis** H.Buch
- 1 **O. atlanticus** (Kaal.) H.Buch [*Barbilophozia atlantica* (Kaal.) Müll.Frib., *Lophozia atlantica* (Kaal.) Müll.Frib.]
- 2 **O. cavifolius** H.Buch & S.W.Arnell [*Sphenolobus cavifolius* (H.Buch & S.W.Arnell) Müll.Frib., *Anastrophyllum cavifolium* (H.Buch & S.W.Arnell) Lammes, *Lophozia cavifolia* (H.Buch & S.W.Arnell) R.M.Schust.]
- 14 **Schizophyllopsis** Váňa & L.Söderstr.
- 1 **S. sphenoloboides** (R.M.Schust.) Váňa & L.Söderstr. [*Anastrophyllum sphenoloboides* R.M.Schust.]⁷
- 15 **Schljakovia** Konstant. & Vilnet
- 1 **S. kunzeana** (Huebener) Konstant. & Vilnet [*Barbilophozia kunzeana* (Huebener) Müll.Frib., *Lophozia kunzeana* (Huebener) A.Evans]
- 16 **Schljakovianthus** Konstant. & Vilnet
- 1 **S. quadrilobus** (Lindb.) Konstant. & Vilnet [*Barbilophozia quadriloba* (Lindb.) Loeske, *Lophozia quadriloba* (Lindb.) A.Evans, *Lophozia quadriloba* var. *glareosa* (Jørg.) Jørg.]
- 17 **Sphenolobopsis** R.M.Schust. & N.Kitag.
- 1 **S. pearsonii** (Spruce) R.M.Schust.
- 18 **Sphenolobus** (Lindb.) Berggr.
- 1 **S. minutus** (Schreb. ex D.Crantz) Berggr. [*Anastrophyllum minutum* (Schreb. ex D.Crantz) R.M.Schust., *Anastrophyllum minutum* var. *weberi* (Mart.) Kartt.]
- 2 **S. saxicola** (Schrad.) Steph. [*Anastrophyllum saxicola* (Schrad.) R.M.Schust.]
- 19 **Tetralophozia** (R.M.Schust.) Schljakov
- 1 **T. filiformis** (Steph.) Urmi
- 2 **T. setiformis** (Ehrh.) Schljakov
- Cephaloziaceae Mig.
Cephalozioidae Müll.Frib.
- 20 **Cephalozia** (Dumort.) Dumort.
- 1 **C. ambigua** C.Massal. [*Cephalozia bicuspidata* var. *paludosa* Jørg. ex Damsh. nom. inval.]
- 2 **C. bicuspidata** (L.) Dumort.
a subsp. **bicuspidata**
b subsp. **lammersiana** (Huebener) R.M.Schust.⁸
- 3 **C. crossii** Spruce⁹
- 4 **C. lacinulata** (J.B.Jack ex Gottsche & Rabenh.) Spruce
- 5 **C. macounii** (Austin) Austin
- 21 **Fuscocephaloziopsis** Fulford [*Pleurocladula* Grolle]
- 1 **F. affinis** (Lindb. ex Steph.) Váňa & L.Söderstr. [*Cephalozia affinis* Lindb. ex Steph., *Pleurocladula affinis* (Lindb. ex Steph.) Konstant., Vilnet & Troitsky]
- 2 **F. albescens** (Hook.) Váňa & L.Söderstr. [*Pleurocladula albescens* (Hook.) Grolle]
a var. **albescens**
b var. **islandica** (Nees) Váňa & L.Söderstr. [*Pleurocladula albescens* var. *islandica* (Nees) L.Söderstr. & Váňa]
- 3 **F. catenulata** (Huebener) Váňa & L.Söderstr. [*Cephalozia catenulata* (Huebener) Lindb., *Pleurocladula catenulata* (Huebener) Konstant., Vilnet & Troitsky]
a subsp. **catenulata**¹⁰
- 4 **F. connivens** (Dicks.) Váňa & L.Söderstr. [*Cephalozia connivens* (Dicks.) Lindb., *Pleurocladula connivens* (Dicks.) Konstant., Vilnet & Troitsky]
a subsp. **connivens** [*Cephalozia connivens* var. *compacta* (Warnst.) W.E.Nicholson ex Macvicar]¹¹

⁷*Schizophyllopsis sphenoloboides* was nested within *Anastrophyllum* in the study by Vilnet, Konstantinova, et al. (2010) which led Stotler and Crandall-Stotler (2017) to retain it in that genus. As it is the type of *Schizophyllopsis*, this means that the whole genus would disappear. However, Vilnet, Konstantinova, et al. (2010) did not include any other member of the genus, or the type of *Anastrophyllum*, in their studies. We thus retain *Schizophyllopsis* pending further studies.

⁸*Cephalozia bicuspidata* subsp. *lammersiana* has a long history of debate on whether it is worth recognizing at any level. The issue requires a molecular study before it can be resolved.

⁹*Cephalozia crossii* was reported from the Canary Islands (Schäfer-Verwimp and Váňa in Ellis, Afonina, et al. 2018).

¹⁰*Fuscocephaloziopsis catenulata* subsp. *nipponica* (S.Hatt.) Váňa & L.Söderstr. occurs in SE Asia.

¹¹*Fuscocephaloziopsis connivens* subsp. *fissa* (Steph.) Váňa & L.Söderstr. occurs in Africa and SE Asia and subsp. *sandwicensis* (Mont.) Váňa & L.Söderstr. in Hawaii, Tahiti and Central America.

- 5 **F. crassifolia** (Lindenb. & Gottsche) Vána & L.Söderstr. [*Cephalozia crassifolia* (Lindenb. & Gottsche) Fulford]
- 6 **F. leucantha** (Spruce) Vána & L.Söderstr. [*Cephalozia leucantha* Spruce, *Pleurocladula leucantha* (Spruce) Konstant., Vilnet & Troitsky, *Cephalozia leucantha* var. *robusta* Schljakov]
- 7 **F. loitlesbergeri** (Schiffn.) Vána & L.Söderstr. [*Cephalozia loitlesbergeri* Schiffn., *Pleurocladula loitlesbergeri* (Schiffn.) Konstant., Vilnet & Troitsky]
- 8 **F. lunulifolia** (Dumort.) Vána & L.Söderstr. [*Cephalozia lunulifolia* (Dumort.) Dumort., *Pleurocladula lunulifolia* (Dumort.) Konstant., Vilnet & Troitsky]
- 9 **F. macrostachya** (Kaal.) Vána & L.Söderstr. [*Cephalozia macrostachya* Kaal., *Pleurocladula macrostachya* (Kaal.) Konstant., Vilnet & Troitsky]
- a subsp. **macrostachya**¹²
- i. var. **macrostachya**
- ii. var. **spiniflora** (Schiffn.) Vána & L.Söderstr.
- 10 **F. pleniceps** (Austin) Vána & L.Söderstr. [*Cephalozia pleniceps* (Austin) Lindb., *Pleurocladula pleniceps* (Austin) Konstant., Vilnet & Troitsky]
- a var. **pleniceps** [*Cephalozia pleniceps* var. *sphagnorum* (C.Massal.) Jørg.]¹³
- 22 **Nowellia** Mitt.
section *Nowellia*
- 1 **N. curvifolia** (Dicks.) Mitt.
- Odontoschismatoideae H.Buch ex Grolle.
- 23 **Odontoschisma** (Dumort.) Dumort. [*Cladopodiella* H.Buch]
section *Cladopodiella* (H.Buch) Gradst., S.C.Aranda & Vanderp.
- 1 **O. francisci** (Hook.) L.Söderstr. & Vána [*Cladopodiella francisci* (Hook.) Jørg.]
- section *Denudata* R.M.Schust.
- 2 **O. denudatum** (Mart.) Dumort.
a subsp. **denudatum**¹⁴
- 3 **O. elongatum** (Lindb.) A.Evans
- 4 **O. macounii** (Austin) Underw.
section *Neesii* Gradst., S.C.Aranda & Vanderp.
- 5 **O. fluitans** (Nees) L.Söderstr. & Vána [*Cladopodiella fluitans* (Nees) H.Buch]
section *Odontoschisma*
- 6 **O. sphagni** (Dicks.) Dumort. [*Odontoschisma prostratum* (Sw.) Trevis.]¹⁵
- Cephaloziellaceae Douin¹⁶
- 24 **Cephaloziella** (Spruce) Schiffn. [*Dichiton* Mont.]¹⁷
subgenus *Cephaloziella*
- 1 **C. arctogena** (R.M.Schust.) Konstant. [*Cephaloziella rubella* subsp. *arctogena* (R.M.Schust.) R.M.Schust. & Damsh.]
- 2 **C. aspericaulis** Jørg.
- 3 **C. baumgartneri** Schiffn.
- 4 **C. divaricata** (Sm.) Schiffn.
a var. **divaricata** [*Cephaloziella divaricata* var. *rupestris* (C.E.O.Jensen) Damsh. *nom. inval.*]
- b var. **scabra** (M.Howe) Haynes [*Cephaloziella divaricata* var. *asperifolia* (Taylor) Damsh.]¹⁸
- 5 **C. elachista** (J.B.Jack ex Gottsche & Rabenh.) Schiffn.
a var. **elachista**¹⁹
- 6 **C. elegans** (Heeg) Schiffn. [*Cephaloziella rubella* var. *elegans* (Heeg) R.M.Schust.]
- 7 **C. grimsulana** (J.B.Jack ex Gottsche & Rabenh.) Lacout.
- 8 **C. hampeana** (Nees) Schiffn. ex Loeske [*Cephaloziella hampeana* var. *subtilis* (Velen) Macvicar]
- 9 **C. massalongi** (Spruce) Müll.Frib. [*Cephaloziella massalongi* var. *compacta* (Jørg.) Müll.Frib.]
- 10 **C. nicholsonii** Douin

¹²*Fuscocephaloziopsis macrostachya* subsp. *australis* (R.M.Schust.) Vána & L.Söderstr. occurs in SE USA.

¹³*Fuscocephaloziopsis pleniceps* var. *caroliniana* (R.M.Schust.) Vána & L.Söderstr. occurs in SE USA.

¹⁴*Odontoschisma denudatum* subsp. *naviculare* (Steph.) Gradst., S.C.Aranda & Vanderp. occurs in E Asia and subsp. *sandvicense* (Ångstr.) Gradst., S.C.Aranda & Vanderp. in Japan and Hawaii.

¹⁵*Odontoschisma prostratum* was recently synonymised by Aranda et al. (2014) based on molecular data.

¹⁶Cephaloziellaceae is here treated in a very broad sense following Vána et al. (2013). Several genera (*Obtusifolium*, *Oleolophozia*, *Protolophozia*) have recently been excluded from other families and been indicated to belong closer to Cephaloziellaceae, but there is no comprehensive study of this phylogenetic region. To avoid monogeneric families of doubtful value the World Checklist (Söderström, Hagborg, et al. 2016) included everything in the only available family name, Cephaloziellaceae, as a sort of 'superfamily', which we are following here.

¹⁷*Cephaloziella* is one of the most difficult liverwort genera. Almost all taxa are difficult to determine, and their taxonomy and distribution is poorly understood. Bell et al. (2013) provided evidence that the molecular data do not match morphological identifications. Type-based studies using molecular data are urgently needed.

¹⁸The taxonomic value of *Cephaloziella divaricata* var. *scabra* is controversial. In Central Europe var. *scabra* intergrades with var. *divaricata* and is possibly only a morph of shady habitats (Köckinger 2017).

¹⁹*Cephaloziella elachista* var. *spinophylla* (C.Gao) C.Gao occurs in China.

- 11 **C. phyllacantha** (C.Massal. & Carestia) Müll.Frib.
- 12 **C. polystratosa** (R.M.Schust. & Damsh.) Konstant. [*Cephaloziella divaricata* var. *polystratosa* (R.M.Schust. & Damsh.) Potemkin]²⁰
- 13 **C. rubella** (Nees) Warnst. [*Cephaloziella rubella* var. *sullivantii* (Austin) Müll.Frib. ex R.M.Schust., *Cephaloziella rubella* var. *bifida* (Lindb.) Douin, *Cephaloziella rubella* var. *pulchella* (C.E.O.Jensen) R.M.Schust.]
- 14 **C. spinigera** (Lindb.) Jørg. [*Cephaloziella spinigera* f. *striatula* (C.E.O.Jensen) Damsh., *Cephaloziella subdentata* Warnst.]
- 15 **C. stellulifera** (Taylor ex Carrington & Pearson) Croz. [*Cephaloziella stellulifera* var. *limprichtii* (Warnst.) Macvicar]
- 16 **C. uncinata** R.M.Schust. a var. **uncinata**²¹
- 17 **C. varians** (Gottsche) Steph. [*Cephaloziella alpina* Douin, *Cephaloziella arctica* Bryhn & Douin, *Cephaloziella varians* var. *arctica* (Bryhn & Douin) Damsh., *Cephaloziella varians* var. *scabra* (S.W.Arnell) Damsh.]
subgenus *Dichiton* (Mont.) Müll.Frib.
- 18 **C. calyculata** (Durieu & Mont.) Müll.Frib.
- 19 **C. integerrima** (Lindb.) Warnst. [*Dichiton integerrimum* (Lindb.) H.Buch, *Cephaloziella integerrima* var. *obtusa* Müll.Frib.]
subgenus *Evansia* (Douin & Schiffn.) Müll.Frib.
- 20 **C. dentata** (Raddi) Steph.
subgenus *Prionolobus* (Spruce) Müll.Frib.
- 21 **C. granatensis** (J.B.Jack ex Steph.) Fulford
- 22 **C. turneri** (Hook.) Müll.Frib.
- 25 **Obtusifolium** S.W.Arnell
1 **O. obtusum** (Lindb.) S.W.Arnell (*Lophozia obtusa* (Lindb.) A.Evans)
- 26 **Oleolophozia** L.Söderstr., De Roo & Hedd.
1 **O. perssonii** (H.Buch & S.W.Arnell) L.Söderstr., De Roo & Hedd. [*Lophozia perssonii* H.Buch & S.W.Arnell, *Lophozia perssonii* (H.Buch & S.W.Arnell) Konstant. & Vilnet]
- 27 **Protolophozia** (R.M.Schust.) Schljakov
1 **P. elongata** (Steph.) Schljakov [*Lophozia elongata* Steph.]
2 **P. herzogiana** (E.A.Hodgs. & Grolle) Váňa & L.Söderstr. [*Lophozia herzogiana* E.A.Hodgs. & Grolle]
- Lophoziaceae Cavers
- 28 **Heterogemma** (Jørg.) Konstant. & Vilnet [*Massularia* Schljakov nom. illeg.]
1 **H. capitata** (Hook.) Konstant. & Vilnet [*Lophozia capitata* (Hook.) Macoun]
2 **H. laxa** (Lindb.) Konstant. & Vilnet [*Lophozia laxa* (Lindb.) Grolle, *Massularia laxa* (Lindb.) Schljakov]
- 29 **Lophozia** (Dumort.) Dumort.²²
1 **L. ascendens** (Warnst.) R.M.Schust. [*Lophozia gracillima* H.Buch]
2 **L. ciliata** Damsh., L.Söderstr. & H.Weibull
3 **L. fuscovirens** Bakalin & Vilnet²³
4 **L. guttulata** (Lindb. & Arnell) A.Evans [*Lophozia porphyroleuca* (Nees) Schiffn. nom. illeg., *Lophozia longiflora* auct. (sensu Grolle and Long 2000; Söderström, Urmi, et al. 2002; Damsholt 2002), *Lophozia longiflora* var. *guttulata* (Lindb. & Arnell) Schljakov]²⁴
5 **L. lantratoviae** Bakalin²⁵
6 **L. longiflora** (Nees) Schiffn. [*Lophozia ventricosa* var. *longiflora* (Nees) Macoun, *Lophozia ventricosa* var. *uliginosa* auct. (sensu Söderström, Urmi, et al. 2002; Damsholt 2002)]²⁶
7 **L. murmanica** Kaal. [*Lophozia groenlandica* auct. (sensu Grolle and Long 2000; Söderström, Urmi, et al. 2002), *Lophozia wenzelii* var. *groenlandica* auct. (sensu Bakalin 2005; Konstantinova, Bakalin, et al. 2009), *Lophozia confertifolia* auct. (sensu Konstantinova, Potemkin, et al. 1992),

²⁰*Cephaloziella polystratosa* was elevated to species rank by Konstantinova (2000b). It is reported from several places in European Russia and Svalbard.

²¹*Cephaloziella uncinata* var. *brevigyra* R.M.Schust. & Damsh. and var. *sphagnicola* R.M.Schust. occur in North America. Schuster (1988; cf. also Damsholt 2013) also described a var. *mamillosa* R.M.Schust. & Damsh. from Greenland, but that name is invalid and we do not validate it here.

²²The genus *Lophozia* includes many taxonomically and nomenclaturally problematic taxa, as outlined in Söderström, Váňa, et al. (2013).

²³*Lophozia fuscovirens* was recently described from the Russian Far East (Bakalin and Vilnet 2019) and is found also on Svalbard (N. Konstantinova in prep.).

²⁴For *Lophozia guttulata*, see footnote on *Lophozia longiflora*.

²⁵*Lophozia lantratoviae* is a recently described species reported from Caucasus by Konstantinova, Akatova, et al. (2009).

²⁶Since Schljakov (1980), *Lophozia longiflora* has been used to include *Lophozia guttulata*, a species mostly restricted to moist dead wood habitats. *Lophozia longiflora* was lectotypified by Bakalin (2016). The lectotype corresponds to the concept of Müller (1954), Saukel (1985), Meinunger and Schröder (2007), Bakalin (2016) and Köckinger (2017), describing a species occurring mostly on peaty soil and in rocky habitats, but not the concept of Grolle and Long (2000), which corresponds to our *Lophozia guttulata*.

- Lophozia heteromorpha* R.M.Schust. & Damsh.]²⁷
- 8 ***L. savicziae*** Schljakov [*Lophozia silvicola* var. *grandiretis* H.Buch & S.W.Arnell, *Lophozia ventricosa* var. *grandiretis* (H.Buch & S.W.Arnell) R.M.Schust. & Damsh., *Lophozia murmanica* auct. (sensu Schljakov 1969)]
 - 9 ***L. schusteriana*** Schljakov [*Lophozia groenlandica* auct. (sensu Schuster 1969)]²⁸
 - 10 ***L. silvicola*** H.Buch [*Lophozia ventricosa* auct. (sensu Müller 1954; Schljakov 1980; Meinunger and Schröder 2007; Köckinger 2017; non Grolle and Long 2000), *Lophozia ventricosa* var. *silvicola* (H.Buch) E.W.Jones]²⁹
 - 11 ***L. silvicoloides*** N.Kitag.³⁰
 - 12 ***L. subapiculata*** R.M.Schust. & Damsh.³¹
 - 13 ***L. ventricosa*** (Dicks.) Dumort. [*Lophozia groenlandica* auct. (sensu Schljakov 1980, 1998), *Lophozia confertifolia* auct. (sensu Schljakov 1975, 1998; Ştefănuţ 2008), *Lophozia murmanica* auct. (sensu Schljakov 1970), *Lophozia ventricosa* var. *confusa* R.M.Schust.]³²
 - 14 ***L. wenzelii*** (Nees) Steph. [*Lophozia groenlandica* auct. (sensu Damsholt 1994, 2002; Ştefănuţ 2008), *Lophozia confertifolia* Schiffn. (sensu Damsholt 2002; Köckinger 2017), *Lophozia ventricosa* var. *uliginosa* Breidl. ex Schiffn., *Lophozia iremelensis* Schljakov]³³
 - a var. ***lapponica*** H.Buch & S.W.Arnell
 - b var. ***litoralis*** (Arnell) Bakalin
 - c var. ***massularioides*** Bakalin
 - d var. ***wenzelii***
- 30 ***Lophozioopsis*** Konstant. & Vilnet
 - 1 ***L. excisa*** (Dicks.) Konstant. & Vilnet [*Lophozia excisa* (Dicks.) Dumort.]
 - a var. ***elegans*** (R.M.Schust.) Konstant. & Vilnet [*Schistochilopsis elegans* (R.M. Schust.) Konstant.]
 - b var. ***excisa*** [*Lophozia excisa* var. *cylindracea* (Dumort.) Müll.Frib.]³⁴
 - 2 ***L. jurensis*** (Meyl. ex Müll.Frib.) Mamontov & Vilnet [*Lophozia jurensis* Meyl. ex Müll. Frib., *Lophozia propagulifera* auct. eur., *Lophozioopsis propagulifera* auct. eur., *Lophozia latifolia* R.M.Schust., *Lophozioopsis latifolia* (R.M.Schust.) Köckinger]³⁵
 - 3 ***L. longidens*** (Lindb.) Konstant. & Vilnet [*Lophozia longidens* (Lindb.) Macoun]
 - a subsp. ***arctica*** (R.M.Schust.) Våňa & L.Söderstr. [*Lophozioopsis rubrigemma* f. *arctica* (R.M.Schust.) Bakalin]
 - b subsp. ***longidens***
 - 4 ***L. pellucida*** (R.M.Schust.) Konstant. & Vilnet [*Lophozia pellucida* R.M.Schust.]

²⁷*Lophozia murmanica* was placed in synonymy with *Lophozia groenlandica* with a question mark (together with *Lophozia heteromorpha*, also with a question mark) by Schljakov (1998). This concept has led to questionable reports of *Lophozia murmanica* from many areas of Europe.

²⁸*Lophozia schusteriana* is a new name given to *Lophozia groenlandica* sensu Schuster (1969) by Schljakov (1975). Schljakov also rejected all reports from Europe but Vilnet, Konstantinova, et al. (2010) reported it later from Svalbard.

²⁹For *Lophozia silvicola*, see footnote under *Lophozia ventricosa*.

³⁰*Lophozia silvicoloides* is originally described from E Asia and recently reported from Svalbard (Vilnet, Milyutina, et al. 2005) and European Russia in Murmansk Province and the Republic of Bashkortostan (Bakalin 2001).

³¹*Lophozia subapiculata* is a critical taxon that was synonymised with *Lophozia ventricosa* by Bakalin (2005) but recognised as a separate species and reported from Svalbard by Konstantinova and Savchenko (2018).

³²The *Lophozia ventricosa/wenzelii* complex has never been studied in detail worldwide using both molecular and morphological methods. Vilnet, Konstantinova, et al. (2010) and de Roo et al. (2007) concentrated on other problems and did not include a sufficient number of specimens of these and related taxa to include the full morphological variation of this group. The taxonomy proposed here is based mostly on morphological studies (which, however, are contradictory!) and is therefore provisional pending future research.

The name *Lophozia ventricosa* has been applied to different concepts since Buch (1929) described *Lophozia silvicola* (see also Buch 1932) and has often included the latter as synonym. The neotype chosen by Grolle and Long (2000) for *Lophozia ventricosa* define it as different from *Lophozia silvicola*. However, there are some doubts if this neotype represents *Lophozia ventricosa* sensu Söderström, Hagborg, et al. (2016) or is a form of *Lophozia wenzelii* (see Meinunger and Schröder 2007; Köckinger 2017). *Lophozia ventricosa* is here understood as a plant with the overall morphology and anatomy of *Lophozia silvicola* but showing homogeneous granular oil bodies. The neotype of Grolle and Long (2000) needs a reinvestigation as it approaches what we understand as *Lophozia wenzelii*.

³³*Lophozia wenzelii* is very close to *Lophozia ventricosa* according to molecular studies by De Roo et al. (2007) and Vilnet, Konstantinova, et al. (2010), but refer to the footnote under *Lophozia ventricosa* concerning the neotype of the latter species. Söderström, Hagborg, et al. (2016) did not accept any subspecific taxa but we list some here accepted in the revision of *Lophozia* for Russia (Bakalin 2005) to draw attention to the need for further studies of this complex.

Lophozia groenlandica has been used in at least three different senses (see Söderström, Våňa, et al. 2013). Although being one of the oldest names in the complex, we refrain from using it in any sense until its affinity is better defined. Schljakov (in Konstantinova, Potemkin, et al. 1992) proposed to reject the name *Lophozia groenlandica* (but never formally did it) and changed it for *Lophozia confertifolia*.

Lophozia confertifolia has also been used in several senses and we here refrain to use it in any specific sense until it is lectotypified (and possibly epi-typtified if needed).

³⁴*Lophozioopsis excisa* var. *infuscata* (R.M.Schust. & Damsh.) Konstant. & Vilnet and var. *succulenta* (R.M.Schust. & Damsh.) Konstant. & Vilnet. occur in arctic North America and Siberia.

³⁵*Lophozioopsis jurensis* is a problematic taxon. It was synonymised with *Lophozia latifolia* by Schljakov (1980) and with *Lophozia propagulifera* by Bakalin (2005). Borovichev and Mamontov (2017; see also Köckinger 2017) rejected the synonymy, as European specimens differed molecularly from specimens named *Lophozioopsis propagulifera* from Kamchatka. However, there is as yet no study to determine whether *Lophozioopsis latifolia* (type from arctic North America), specimens of *Lophozioopsis propagulifera* from the Southern Hemisphere (type from South Georgia) and *Lophozioopsis jurensis* (type from the Alps) are genetically different. We here list only one species, assuming that the Southern Hemisphere taxon is different, and following Schljakov (1980) in regarding *Lophozioopsis latifolia* as a synonym of *Lophozioopsis jurensis*.

- a var. **minor** (R.M.Schust.) L.Söderstr. & Váňa
 b var. **pellucida**
- 5 **L. polaris** (R.M.Schust.) Konstant. & Vilnet
 [*Lophozia polaris* (R.M.Schust.)
 R.M.Schust & Damsh.]
 a var. **polaris**
 b var. **sphagnorum** (R.M.Schust.) Konstant. &
 Vilnet
- 6 **L. rubrigemma** (R.M.Schust.) Konstant. & Vilnet
 [*Lophozia rubrigemma* R.M.Schust.]
- 31 **Trilophozia** (R.M.Schust.) Bakalin
 1 **T. quinqueidentata** (Huds.) Bakalin [*Tritomaria*
quinqueidentata (Huds.) H.Buch, *Trito-*
maria quinqueidentata subsp. *turgida*
 (Lindb.) Damsh., *Trilophozia quinque-*
identata subsp. *turgida* (Lindb.) Kon-
stant., Tritomaria quinqueidentata
 f. *gracilis* (C.E.O.Jensen) R.M.Schust.,
Tritomaria quinqueidentata var.
dentata S.W.Arnell nom. *inval.*, *Trito-*
maria quinqueidentata var. *grandiretis*
 H.Buch & S.W.Arnell]
 a var. **quinqueidentata**³⁶
- 32 **Tritomaria** Schiffn. ex Loeske
 1 **T. exsecta** (Schmidel) Schiffn. ex Loeske
 a subsp. **exsecta**³⁷
 2 **T. exsectiformis** (Breidl.) Schiffn. ex Loeske
 a subsp. **arctica** R.M.Schust.
 b subsp. **exsectiformis**³⁸
 3 **T. scitula** (Taylor) Jørg.
- Scapaniaceae Mig.
 33 **Diplophyllum** (Dumort.) Dumort.
 section *Diplophyllum*
 1 **D. albicans** (L.) Dumort.
 section *Protodiplophyllum* (R.M.Schust.) Váňa &
 L.Söderstr.
 2 **D. obtusatum** (R.M.Schust.) R.M.Schust.³⁹
- 3 **D. obtusifolium** (Hook.) Dumort.
 a subsp. **obtusifolium**⁴⁰
- 4 **D. taxifolium** (Wahlenb.) Dumort.
 a var. **taxifolium** [*Diplophyllum taxifolium*
 var. *macrostictum* H.Buch]⁴¹
- 34 **Douinia** (C.E.O.Jensen) H.Buch
 1 **D. ovata** (Dicks.) H.Buch
- 35 **Pseudotritomaria** Konstant. & Vilnet
 1 **P. heterophylla** (R.M.Schust.) Konstant. &
 Vilnet⁴²
- 36 **Saccobasis** H.Buch
 1 **S. polita** (Nees) H.Buch [*Tritomaria polita*
 (Nees) Jørg.]
 2 **S. polymorpha** (R.M.Schust.) Schljakov [*Tritomaria*
polita subsp. *polymorpha* R.M.Schust.]⁴³
- 37 **Scapania** (Dumort.) Dumort.
 subgenus *Gracilidae* (H.Buch) Váňa, Hentschel,
 Joch.Müll. & Heinrichs
 1 **S. gracilis** Lindb.
 subgenus *Plicicalyx* Müll.Frib.
 section *Planifoliae* (Müll.Frib.) Potemkin
 2 **S. nimbose** Taylor
 3 **S. ornithopodioides**⁴⁴ (With.) Waddell
 subgenus *Scapania*
 section *Aequilobae* (Müll.Frib.) H.Buch
 4 **S. aequiloba** (Schwägr.) Dumort.
 5 **S. aspera** M.Bernet & Bernet
 section *Apiculatae* H.Buch
 6 **S. apiculata** Spruce
 7 **S. carinthiaca** J.B.Jack ex Lindb.⁴⁵
 a var. **carinthiaca**
 b var. **massalongi** Müll.Frib.
 8 **S. umbrosa** (Schrad.) Dumort.
 section *Compactae* (Müll.Frib.) H.Buch
 9 **S. compacta** (Roth) Dumort.
 10 **S. kaurinii** Ryan

³⁶*Trilophozia quinqueidentata* var. *assymetrica* (Horik.) L.Söderstr. & Váňa occurs in E Asia. *Tritomaria quinqueidentata* is a polymorphic species with many infraspecific taxa described, but the value of them are questioned. The only infrataxon known to be included in a molecular study (subsp. *turgida*) was nested among other specimens morphologically belonging to subsp. *quinqueidentata* (De Roo et al. 2007).

³⁷*Tritomaria exsecta* subsp. *novaezealandia* J.J.Engel occurs in New Zealand.

³⁸*Tritomaria exsectiformis* subsp. *camerunensis* S.W.Arnell ex Váňa occurs in Africa.

³⁹*Diplophyllum obtusatum* is close to or possibly conspecific with *Diplophyllum obtusifolium*, differing mainly in being autoicous vs. paroicous. Urmi (2017) regarded them as conspecific but did not formally treat them as such. In addition, Bakalin and Vilnet (2018) treated all reports of both species from Asia as belonging to their new species, *Diplophyllum sibiricum* Vilnet & Bakalin, and regarded *Diplophyllum obtusatum* as an American taxon of unclear status. However, they did not include any American or European specimen of *Diplophyllum obtusatum* to confirm this conclusion.

⁴⁰*Diplophyllum obtusifolium* subsp. *domesticum* (Gottsche) Váňa occurs in the Subantarctic.

⁴¹*Diplophyllum taxifolium* var. *mucronatum* R.M.Schust. occurs in North America.

⁴²*Pseudotritomaria heterophylla* is reported from Franz Josef Land but "the material is poor" (Konstantinova and Potemkin 1996) and the taxon was not included in the list of Grolle and Long (2000). We include it here with some hesitation.

⁴³*Saccobasis polymorpha* was included (as *Tritomaria polymorpha*) in *Tritomaria polita* by Grolle and Long (2000).

⁴⁴Withering (1776) originally spelt the species epithet of *Scapania ornithopodioides* as "*ornithopoides*", a spelling Potemkin (2019) argued should be retained, referring to ICN Art. 60.1 (Turland et al. 2018). However, Grolle (1973) argued that "*ornithopoides*" is bad Latin and in addition that Withering referred to Dillenius' pre-Linnean name "... *ornithopodii minor* ...". Thus Grolle (1973) rejected the original spelling as an orthographic or typographic error, which has been widely accepted and is followed here.

⁴⁵There are two morphotypes of *Scapania carinthiaca* (*Scapania carinthiaca* s.str. and *Scapania massalongi*), which were for a long time treated as separate taxa (Müller 1954), before being synonymised by Potemkin (1999). Most records in Europe belong to the *massalongi*-morphotype, while the *carinthiaca*-morphotype is very rare. Further studies with fertile plants are desirable, especially as two morphotypes can be distinguished in the Alps, following Meuninger and Schröder (2007) and others.

- 11 *S. spitsbergensis* (Lindb.) Müll.Frib.
section *Curtae* (Müll.Frib.) H.Buch
- 12 *S. curta* (Mart.) Dumort.
a var. *curta*
b var. *grandiretis* R.M.Schust.
c var. *isoloba* R.M.Schust.
- 13 *S. helvetica* Gottsche
- 14 *S. irrigua* (Nees) Nees
a subsp. *irrigua* [*Scapania irrigua* var. *rubescens* H.Buch]
b subsp. *rufescens* (Loeske) R.M.Schust.
- 15 *S. lingulata* H.Buch
a var. *lingulata*
b var. *microphylla* (Warnst.) R.M.Schust.
- 16 *S. mucronata* H.Buch
- 17 *S. obcordata* (Berggr.) S.W.Arnell [*Scapania paradoxa* R.M.Schust.]
- 18 *S. parvifolia* Warnst.
- 19 *S. praetervisa* Meyl. [*Scapania mucronata* subsp. *praetervisa* (Meyl.) R.M.Schust.]
- 20 *S. scandica* (Arnell & H.Buch) Macvicar
a var. *argutedentata* H.Buch
b var. *grandiretis* (Schljakov) Schljakov
c var. *scandica*⁴⁶
- 21 *S. uliginosa* (Lindenb.) Dumort.
- 22 *S. zemliae* S.W.Arnell
section *Cuspiduligerae* H.Buch
- 23 *S. cuspiduligera* (Nees) Müll.Frib.
a var. *cuspiduligera*⁴⁷
- section *Hyperboreae* Váňa, Hentschel, Joch.Müll. & Heinrichs
- 24 *S. hyperborea* Jørg.
- 25 *S. paludicola* Loeske & Müll.Frib. [*Scapania paludicola* var. *rufescens* Damsh. nom. inval.]
a var. *paludicola*⁴⁸
b var. *rotundiloba* R.M.Schust. ex Konstant. & L.Söderstr.
- 26 *S. tundrae* (Arnell) H.Buch
section *Kaalaasia* (H.Buch) Jørg.
- 27 *S. calcicola* (Arnell & J.Perss.) Ingham
- 28 *S. gymnostomophila* Kaal.
- 29 *S. ligulifolia* R.M.Schust.
section *Nemorosae* (Müll.Frib.) H.Buch
- 30 *S. crassiretis* Bryhn
- 31 *S. degenii* Schiffn. ex Müll.Frib. [*Scapania brevicaulis* auct. eur. non Taylor, *Scapania degenii* var. *dubia* R.M.Schust., *Scapania brevicaulis* var. *dubia* (R.M.Schust.) Damsh.]⁴⁹
- 32 *S. nemorea* (L.) Grolle
section *Scapania*
- 33 *S. obscura* (Arnell & C.E.O.Jensen) Schiffn.
- 34 *S. paludosa* (Müll.Frib.) Müll.Frib. [*Scapania paludosa* var. *isoloba* Müll.Frib., *Scapania paludosa* var. *rubiginosa* Müll.Frib., *Scapania paludosa* var. *vogesiana* Müll.Frib.]
- 35 *S. subalpina* (Nees ex Lindenb.) Dumort.
a var. *subalpina*⁵⁰
- 36 *S. undulata* (L.) Dumort. [*Scapania undulata* var. *aequatiformis* (De Not.) C.Massal. & Carestia, *Scapania undulata* var. *dentata* (Dumort.) Jørg., *Scapania undulata* var. *oakesii* (Austin) H.Buch]
- section *Scapaniella* (H.Buch) Potemkin
- 37 *S. glaucocephala* (Taylor) Austin
a var. *glaucocephala*⁵¹
- 38 *S. scapanioides* (C.Massal.) Grolle [*Scapania glaucocephala* var. *scapanioides* (C.Massal.) Damsh.]
section *Simmonsiae* (R.M.Schust.) Váňa, Hentschel, Joch.Müll. & Heinrichs
- 39 *S. simmonsii* Bryhn & Kaal.
section *Sphaeriferae* Konstant. & Potemkin
- 40 *S. sphaerifera* H.Buch & Tuom.
section *Verrucosae* Potemkin
- 41 *S. verrucosa* Heeg
- 38 *Schistochilopsis* (N.Kitag.) Konstant.
- 1 *S. grandiretis* (Lindb. ex Kaal.) Konstant. [*Lophozia grandiretis* (Lindb. ex Kaal.) Schiffn., *Lophozia grandiretis* var. *parviretis* R.M.Schust., *Lophozia grandiretis* var. *proteidea* (Arnell) Arnell]
- 2 *S. hyperarctica* Konstant. & L.Söderstr. [*Lophozia hyperarctica* R.M.Schust.]
- 3 *S. incisa* (Schrad.) Konstant. [*Lophozia incisa* (Schrad.) Dumort., *Massularia incisa* (Schrad.) Schljakov, *Schistochilopsis incisa* var. *inermis* (Müll.Frib.) Konstant.]
- 4 *S. opacifolia* (Culm. ex Meyl.) Konstant. [*Lophozia opacifolia* Culm. ex Meyl., *Lophozia incisa* subsp. *opacifolia* (Culm. ex Meyl.) R.M.Schust. & Damsh.]
- Acrobolbaceae E.A.Hodgs.
Acrobolboideae R.M.Schust. ex Briscoe

⁴⁶*Scapania scandica* var. *dimorpha* R.M.Schust. occurs in North America

⁴⁷*Scapania cuspiduligera* var. *diplophyllopsis* R.M.Schust. occurs in North America.

⁴⁸*Scapania paludicola* var. *viridigemma* R.M.Schust. occurs in North America.

⁴⁹*Scapania degenii* was synonymised with *Scapania brevicaulis* by Potemkin (1998) but recognised by Konstantinova, Bakalin, et al. (2009). *Scapania brevicaulis* s.str. is an American taxon that has never been reported from Europe without *Scapania degenii* being included as a synonym.

⁵⁰*Scapania subalpina* var. *haynesiae* Frye & L.Clark and var. *muddiae* C.D.Bird & W.S.Hong occur in North America.

⁵¹*Scapania glaucocephala* var. *saxicola* (R.M.Schust.) Potemkin occurs in North America.

39 **Acrobolbus** Nees [*Tylimanthus* Mitt.]

- 1 **A. azoricus** (Grolle & Perss.) Briscoe [*Tylimanthus azoricus* Grolle & Perss.]
- 2 **A. madeirensis** (Grolle & Perss.) Briscoe [*Tylimanthus madeirensis* Grolle & Perss.]
- 3 **A. wilsonii** Nees
 - a var. **wilsonii**⁵²

Antheliaceae R.M.Schust.

40 **Anthelia** (Dumort.) Dumort.

- 1 **A. julacea** (L.) Dumort.
- 2 **A. juratzkana** (Limpr.) Trevis.

Arnelliaceae Nakai

41 **Arnellia** Lindb.

- 1 **A. fennica** (Gottsche) Lindb.

Calypogeiaceae Arnell

42 **Calypogeia** Raddi

subgenus *Asperifoliae* (Warnst.) R.M.Schust.

- 1 **C. arguta** Nees & Mont.

subgenus *Calypogeia*

- 2 **C. azorica** Bischl.
- 3 **C. azurea** Stotler & Crotz
- 4 **C. fissa** (L.) Raddi
 - a subsp. **fissa**
 - b subsp. **neogaea** R.M.Schust.
 - c var. **paludosa** (Warnst.) Damsh.⁵³
- 5 **C. integristipula** Steph.
- 6 **C. muelleriana** (Schiffn.) Müll.Frib.
 - a subsp. **muelleriana** [*Calypogeia muelleriana* var. **erecta** (Müll.Frib.) Müll.Frib.]⁵⁴
- 7 **C. neesiana** (C.Massal. & Carestia) Müll.Frib.
 - a subsp. **neesiana** [*Calypogeia neesiana* var. **hygrophila** Müll.Frib., *Calypogeia neesiana* var. **repanda** (Müll.Frib.) Meyl.]⁵⁵
- 8 **C. sphagnicola** (Arnell & J.Perss.) Warnst. & Loeske
- 9 **C. suecica** (Arnell & J.Perss.) Müll.Frib.

43 **Mnioloma** Herzog

subgenus *Caracoma* (Bischl.) R.M.Schust.

- 1 **M. fuscum** (Lehm.) R.M.Schust.

Endogemmaceae Konstant.⁵⁶44 **Endogemma** Konstant.

- 1 **E. caespiticia** (Lindenb.) Konstant., Vilnet & A.V.Troitsky [*Jungermannia caespiticia* Lindenb.]

Geocalyceae H.Klinggr.

45 **Geocalyx** Nees

- 1 **G. graveolens** (Schrad.) Nees

Gymnomitriaceae H.Klinggr.

Gymnomitrioideae T.Jensen

46 **Gymnomitrium** Corda [*Apomarsupella* R.M.Schust.]

- 1 **G. adustum** Nees [*Marsupella adusta* (Nees) Spruce]
- 2 **G. alpinum** (Gottsche ex Husn.) Schiffn. [*Marsupella alpina* (Gottsche ex Husn.) Bernet]
- 3 **G. brevissimum** (Dumort.) Warnst. [*Marsupella brevissima* (Dumort.) Grolle]
- 4 **G. commutatum** (Limpr.) Schiffn. [*Marsupella commutata* (Limpr.) Bernet]
- 5 **G. concinnatum** (Lightf.) Corda
- 6 **G. corallioides** Nees
- 7 **G. crenulatum** Gottsche ex Carrington
- 8 **G. obtusum** Lindb.
- 9 **G. revolutum** (Nees) H.Philib. [*Apomarsupella revoluta* (Nees) R.M.Schust.]
 - a subsp. **revolutum**⁵⁷

47 **Marsupella** Dumort.

- 1 **M. andreaeoides** (Lindb.) Müll.Frib.
- 2 **M. apiculata** Schiffn. [*Gymnomitrium apiculatum* (Schiffn.) Müll.Frib.]
- 3 **M. aquatica** (Lindenb.) Schiffn. [*Marsupella emarginata* subsp. **aquatica** (Lindenb.) Meyl., *Marsupella emarginata* var. **aquatica** (Lindenb.) Dumort.]⁵⁸
- 4 **M. arctica** (Berggr.) Bryhn & Kaal.
- 5 **M. boeckii** (Austin) Lindb. ex Kaal.
- 6 **M. condensata** (Ångstr. ex C.Hartm.) Lindb. ex Kaal.
- 7 **M. emarginata** (Ehrh.) Dumort. [*Marsupella emarginata* var. **pearsonii** (Schiffn. ex Macvicar) Jørg.]

⁵²*Acrobolbus wilsonii* var. *andinus* Spruce occurs in South America. Whether it belongs to *Acrobolbus wilsonii* or some other species is unclear, as *Acrobolbus wilsonii* is normally considered to be a European endemic.

⁵³Damsholt (2017) transferred *Calypogeia sphagnicola* f. *paludosa* (Warnst.) R.M.Schust. to *Calypogeia fissa*. Buczkowska et al. (2018) included both 'f. *sphagnicola*' and 'f. *paludosa*' in their study on, primarily, the *Calypogeia* species with blue oil bodies. However, in their molecular phylogenetic tree, '*paludosa*' came out distinctly separated from '*sphagnicola*' and closer to *Calypogeia fissa* (supporting Damsholt's view), although far enough from it that it should perhaps deserve recognition at species level. For now, to avoid premature nomenclatural changes we treat it as a variety without affiliation to any of the subspecies pending further research.

⁵⁴*Calypogeia muelleriana* subsp. *blomquistii* R.M.Schust. occurs in North America.

⁵⁵*Calypogeia neesiana* subsp. *subalpina* (Inoue) Inoue occurs in Japan.

⁵⁶*Endogemma* is feminine not neuter and the family name should therefore be Endogemmaceae, not Endogemmataceae as originally published.

⁵⁷*Gymnomitrium revolutum* subsp. *novoguineanense* (R.M.Schust.) Váňa, Crand.-Stotl. & Stotler occurs in New Guinea.

⁵⁸*Marsupella aquatica* was treated under *Marsupella emarginata* in Grolle and Long (2000) but shown to be sister to it and accepted at species level by Vilnet, Konstantinova, et al. (2010).

- 8 **M. funckii** (F.Weber & D.Mohr) Dumort.
[*Marsupella ramosa* Müll.Frib., *Marsupella funckii* var. *badensis* (Schiffn.) Fam.]
- 9 **M. profunda** Lindb.
- 10 **M. sparsifolia** (Lindb.) Dumort.
a subsp. **sparsifolia**⁵⁹
- 11 **M. sphacelata** (Giesecke ex Lindenb.) Dumort.
- 12 **M. spiniloba** R.M.Schust. & Damsh.
- 13 **M. sprucei** (Limpr.) Bernet [*Marsupella sprucei* var. *neglecta* (Limpr.) Damsh., *Marsupella sprucei* var. *ustulata* (Limpr.) Damsh.]
- 14 **M. stableri** Spruce
- 15 **M. submarginata** Bakalin & Fedosov⁶⁰
- 16 **M. tubulosa** Steph. [*Marsupella emarginata* subsp. *tubulosa* (Steph.) N.Kitag.]⁶¹
- 48 **Prasanthus** Lindb.
1 **P. suecicus** (Gottsche) Lindb.
- Nardioideae Váňa
- 49 **Nardia** Gray
1 **N. breidlerii** (Limpr.) Lindb.
2 **N. compressa** (Hook.) Gray
3 **N. geoscyphus** (De Not.) Lindb.
a var. **bifida** R.M.Schust.⁶²
b var. **geoscyphus**⁶³
c var. **suberecta** (Lindb. ex Kaal.) Váňa
4 **N. insecta** Lindb.
5 **N. japonica** Steph.
6 **N. pacifica** Bakalin⁶⁴
7 **N. scalaris** Gray
a var. **scalaris**⁶⁵
- Harpanthaceae Arnell
- 50 **Harpanthus** Nees
1 **H. flotovianus** (Nees) Nees [*Harpanthus flotovianus* var. *chiloscyphoides* C.E.O.Jensen, *Harpanthus flotovianus* var. *latifolia* Jørg., *Harpanthus flotovianus* var. *retusa* Jørg.]
- 2 **H. scutatus** (F.Weber & D.Mohr) Spruce
- Hygrobilaceae Konstant. & Vilnet
- 51 **Hygrobilla** Spruce
1 **H. laxifolia** (Hook.) Spruce
- Jungermanniaceae Rchb.
- Delavayelloideae Grolle
- 52 **Liochlaena** Nees
1 **L. lanceolata** Nees [*Jungermannia lanceolata* auct. non L., *Jungermannia leiantha* Grolle, *Jungermannia subulata* var. *leiantha* (Grolle) Damsh.]
2 **L. subulata** (A.Evans) Schljakov [*Jungermannia subulata* A.Evans]
- Jungermannioideae Dumort.
- 53 **Eremonotus** Lindb. & Kaal. ex Pearson
1 **E. myriocarpus** (Carrington) Lindb. & Kaal. ex Pearson
- 54 **Jungermannia** L.
1 **J. atrovirens** Dumort. [*Jungermannia lanceolata* L. nom. rejic., *Jungermannia lanceolata* var. *atrovirens* (Dumort.) Damsh.]
2 **J. borealis** Damsh. & Váňa
3 **J. calcicola** Konstant. & Vilnet⁶⁶
4 **J. eucordifolia** Schljakov [*Jungermannia exsertifolia* subsp. *cordifolia* (Dumort.) Váňa]⁶⁷
5 **J. polaris** Lindb. [*Jungermannia pumila* subsp. *polaris* (Lindb.) R.M.Schust.]
6 **J. pumila** With. [*Jungermannia pumila* var. *alpestris* Gottsche & Rabenh.]
- Mesoptychioideae R.M.Schust.
- 55 **Mesoptychia** (Lindb.) A.Evans [*Leiocolea* (Müll.-Frib.) Buch]
1 **M. badensis** (Gottsche ex Rabenh.) L.Söderstr. & Váňa [*Lophozia badensis* (Gottsche ex Rabenh.) Schiffn., *Leiocolea badensis* (Gottsche ex Rabenh.) Jørg., *Lophozia badensis* var. *obtusiloba* (Bernet) Schiffn.]

⁵⁹*Marsupella sparsifolia* subsp. *childii* R.M.Schust. occurs in New Zealand.

⁶⁰*Marsupella submarginata* was described by Bakalin et al. (2019) from Kamchatka but also reported from Switzerland.

⁶¹*Marsupella tubulosa* was treated as a subspecies of *Marsupella emarginata* by Söderström, Hagborg, et al. (2016) but shown to be better treated at species level by Bakalin et al. (2019).

⁶²*Nardia geoscyphus* var. *bifida* was not recognised by Söderström, Hagborg, et al. (2016) but we list it here pending better understanding of the variability of the species.

⁶³*Nardia geoscyphus* var. *dioica* Bakalin occurs in E Asia.

⁶⁴*Nardia pacifica* is a newly described species (Bakalin and Klimova 2016).

⁶⁵*Nardia scalaris* var. *botryoidea* (R.M.Schust.) Váňa occurs in North America and var. *harae* (Amakawa) Váňa in E Asia.

⁶⁶*Jungermannia calcicola* is a newly described species from Russian Caucasus (Konstantinova and Vilnet 2016) and recently also recorded from Albania by Marka et al. (2018).

⁶⁷*Jungermannia eucordifolia* was treated as a subspecies (*Jungermannia exsertifolia* subsp. *cordifolia*) in Söderström, Hagborg, et al. (2016). However, Konstantinova and Vilnet (2016) and later Mamontov, Konstantinova, et al. (2018) showed that it deserves specific status, as was proposed by Schljakov (1981) and followed by Russian bryologists (e.g. Konstantinova, Potemkin, et al. 1992; Konstantinova, Bakalin, et al. 2009).

- 2 **M. bantriensis** (Hook.) L.Söderstr. & Váňa [*Lophozia bantriensis* (Hook.) Steph., *Leiocolea bantriensis* (Hook.) Jørg.]
 a subsp. **bantriensis** [*Lophozia bantriensis* var. *subcompressa* (Limpr.) Damsh. ex L.Söderstr.]
 b subsp. **wallfischii** (Ștefănuț) L.Söderstr. & Váňa [*Leiocolea bantriensis* subsp. *wallfischii* Ștefănuț]⁶⁸
- 3 **M. collaris** (Nees) L.Söderstr. & Váňa [*Leiocolea collaris* (Nees) Schljakov, *Lophozia alpestris* (Schleich ex F.Weber) Evans nom. rej., *Leiocolea alpestris* (Schleich. ex F.Weber) Isov., *Lophozia alpestris* var. *libertae* (Huebener) Damsh.]
- 4 **M. fitzgeraldiae** (Paton & A.R.Perry) L.Söderstr. & Váňa [*Leiocolea fitzgeraldiae* Paton & A.R.Perry]
- 5 **M. gillmanii** (Austin) L.Söderstr. & Váňa [*Lophozia gillmanii* (Austin) R.M.Schust., *Leiocolea gillmanii* (Austin) A.Evans, *Lophozia gillmanii* var. *acutifolia* (Limpr.) R.M.Schust.]
 a var. **gillmanii**
 b var. **laxa** (Schiffn. ex Burrell) L.Söderstr. [*Leiocolea rutheana* var. *laxa* (Schiffn. ex Burrell) Paton, *Lophozia rutheana* var. *laxa* (Schiffn. ex Burrell) Paton ex Damsh. nom. inval.]⁶⁹
- 6 **M. heterocolpos** (Thed. ex Hartm.) L.Söderstr. & Váňa [*Lophozia heterocolpos* (Thed. ex Hartm.) M.Howe, *Leiocolea heterocolpos* (Thed. ex Hartm.) H.Buch]
 a var. **arctica** (S.W.Arnell) L.Söderstr. & Váňa [*Lophozia heterocolpos* var. *arctica* (S.W.Arnell) R.M.Schust. & Damsh.]
 b var. **harpanthoides** (Bryhn & Kaal.) L.Söderstr. & Váňa [*Lophozia heterocolpos* var. *harpanthoides* (Bryhn & Kaal.) R.M.Schust.]
 c var. **heterocolpos**
- 7 **M. rutheana** (Limpr.) L.Söderstr. & Váňa [*Lophozia ruthana* (Limpr.) M.Howe, *Leiocolea rutheana* (Limpr.) Müll.Frib.]
- 8 **M. sahlbergii** (Lindb. & Arnell) A.Evans
- 9 **M. turbinata** (Raddi) L.Söderstr. & Váňa [*Leiocolea turbinata* (Raddi) H.Buch]
- Saccogynaceae Heeg
 56 **Saccogyna** Dumort.
 1 **S. viticulosa** (L.) Dumort.
- Solenostomataceae Stotler & Crand.-Stotl.
 57 **Cryptocolea** R.M.Schust.
 1 **C. imbricata** R.M.Schust.
- 58 **Solenostoma** Mitt. [*Plectocolea* (Mitt.) Mitt.]
 subgenus *Eucalyx* (Lindb.) Váňa, Crand.-Stotl. & Stotler
 1 **S. obovatum** (Nees) C.Massal. [*Jungermannia obovata* Nees, *Plectocolea obovata* (Nees) Mitt.]
 2 **S. subellipticum** (Lindb. ex Heeg) R.M.Schust. [*Jungermannia subelliptica* (Lindb. ex Heeg) Levier, *Plectocolea subelliptica* (Lindb. ex Heeg) A.Evans, *Jungermannia obovata* subsp. *minor* (Carrington) Damsh.]⁷⁰
 subgenus *Metasolenostoma* Váňa, Crand.-Stotl. & Stotler
 3 **S. gracillimum** (Sm.) R.M.Schust. [*Jungermannia gracillima* Sm., *Jungermannia gracillima* var. *crenulata* (Mitt.) Damsh.]
 4 **S. handelii** (Schiffn.) Müll.Frib. [*Jungermannia handelii* (Schiffn.) Amakawa]
 subgenus *Plectocolea* Mitt.
 5 **S. callithrix** (Lindenb. & Gottsche) Steph. [*Jungermannia callithrix* Lindenb. & Gottsche]
 6 **S. hyalinum** (Lyell) Mitt. [*Jungermannia hyalina* Lyell, *Plectocolea hyalina* (Lyell) Mitt.]
 7 **S. paroicum** (Schiffn.) R.M.Schust. [*Jungermannia paroica* (Schiffn.) Grolle]
 subgenus *Solenostoma*
 8 **S. caucasicum** (Váňa) Konstant.⁷¹
 9 **S. confertissimum** (Nees) Schljakov [*Jungermannia confertissima* Nees]
 10 **S. sphaerocarpum** (Hook.) Steph. [*Jungermannia sphaerocarpa* Hook., *Solenostoma pusillum* (C.E.O.Jens.) Steph., *Jungermannia jenseniana* Grolle, *Jungermannia sphaerocarpa* var. *nana* (Nees ex Flot.) Frye & L.Clark nom. illeg., *Solenostoma sphaerocarpum* var. *nanum* (Nees ex Flot.) R.M.Schust.]

⁶⁸*Mesoptychia bantriensis* subsp. *wallfischii* was described as *Leiocolea bantriensis* subsp. *wallfischii* by Ștefănuț (2008) from Romania.

⁶⁹*Mesoptychia gillmanii* var. *laxa* was accepted by Söderström, Hagborg, et al. (2016) as *Mesoptychia rutheana* var. *laxa* although with low confidence. However, the study by Bell et al. (2013) was largely overlooked and the relation to *Mesoptychia gillmanii* rather than to *Mesoptychia rutheana* was not acknowledged.

⁷⁰*Solenostoma subellipticum* was nested in *Solenostoma obovatum* in Shaw et al. (2015) and thus reduced to a synonym, a treatment followed by Söderström, Hagborg, et al. (2016). However, as it seems distinct in at least parts of its distribution area we list it here with some hesitation.

⁷¹*Solenostoma caucasicum* was reported as new to Europe from the northern part of Caucasus by Konstantinova and Savchenko (2013).

Southbyaceae Váňa, Crand.-Stotl., Stotler & D.G.Long

59 **Gongylanthus** Nees

1 **G. ericetorum** (Raddi) Nees

60 **Southbya** Spruce

1 **S. nigrella** (De Not.) Henriq.

2 **S. topfacea** (Spruce) Spruce

Blepharostomataceae W.Frey & M.Stech

61 **Blepharostoma** (Dumort.) Dumort.

1 **B. trichophyllum** (L.) Dumort.

a subsp. **brevirete** (Bryhn & Kaal.) R.M.Schust.
[*Blepharostoma trichophyllum* var.
brevirete Bryhn & Kaal.]

b subsp. **trichophyllum**

Herbertaceae Müll.Frib. ex Fulford & Hatcher

62 **Herbertus** Gray

1 **H. azoricus** (Steph.) P.W.Richards⁷²

2 **H. borealis** Crundw. [*Herbertus delavayi* auct.
eur. non (Steph.) Steph.]

3 **H. hutchinsiae** (Gottsche & Rabenh.) A.Evans
[*Herbertus aduncus* auct. non (Dicks.)
Gray, *Herbertus aduncus* subsp. *hutch-*
insiae (Gottsche & Rabenh.)
R.M.Schust. nom. inval.]⁷³

4 **H. norenius** D.G.Long, D.Bell & H.H.Blom⁷⁴

5 **H. sendtneri** (Nees) Lindb.

6 **H. stramineus** (Dumort.) Trevis.

Lepidoziaceae Limpr.

Bazzanioideae Rodway

63 **Bazzania** Gray

1 **B. azorica** H.Buch & Perss.

2 **B. flaccida** (Dumort.) Grolle

3 **B. pearsonii** Steph.

4 **B. tricrenata** (Wahlenb.) Lindb.

a var. **tricrenata**⁷⁵

5 **B. trilobata** (L.) Gray

a var. **depauperata** (Müll.Frib.) Grolle

b var. **trilobata**

Lembidioideae R.M.Schust.

64 **Kurzia** G.Martens

1 **K. pauciflora** (Dicks.) Grolle

2 **K. sylvatica** (A.Evans) Grolle

3 **K. trichocladus** (Müll.Frib.) Grolle

Lepidozioideae Müll.Frib.

65 **Lepidozia** (Dumort.) Dumort.

1 **L. cupressina** (Sw.) Lindenb.

a subsp. **cupressina** [*Lepidozia cupressina*
subsp. *pinnata* (Hook.) Pócs, *Lepi-*
dozia cupressina var. *dissitifolia*
(Jørg.) Damsh.]⁷⁶

2 **L. pearsonii** Spruce

3 **L. reptans** (L.) Dumort. [*Lepidozia reptans* var.
julacea (Nees) Damsh. nom. inval., *Lepi-*
dozia reptans var. *tenera* (Huebener)
P.Allorge nom. inval.]

4 **L. stuhlmannii** Steph.⁷⁷

66 **Tricholepidozia** (R.M.Schust.) E.D.Cooper

1 **T. lindenberghii** (Gottsche) E.D.Cooper⁷⁸

a var. **lindenberghii**⁷⁹

2 **T. tetradactyla** (Hook.f. & Taylor) E.D.Cooper
[*Telaranea longii* Paton, *Telaranea mur-*
phyae Paton]⁸⁰

Zoopsidoideae R.M.Schust.

67 **Telaranea** Spruce ex Schiffn.

1 **T. azorica** (H.Buch & Perss.) Pócs [*Lepidozia*
azorica H.Buch & Perss.]

2 **T. europaea** J.J.Engel & G.L.Merr. [*Telaranea*
nematodes auct. eur. non (Gottsche
ex Austin) M.Howe]

⁷²Only a single species of *Herbertus*, *Herbertus azoricus*, has to date been published from the Azores. However, Richards (1936) observed that there are two 'forms' present in the archipelago, and it is now clear that two species are present (Hodgetts and Rumsey in prep.), but further analysis is needed to characterize them. Feldberg *et al.* (2004) suggested that *Herbertus azoricus* should be placed in synonymy with *Herbertus sendtneri*. However, further studies show that spurious lumping of taxa in *Herbertus* gives a much too simplified view of this complex genus (F. Rumsey pers. comm. 2018), and *Herbertus sendtneri* is therefore best regarded as an exclusively Northern Hemisphere species, restricted to the Austrian Alps, Himalaya, and the Pacific Northwest of North America. Current indications are that *Herbertus azoricus* is best treated as a synonym or a subspecies of *Herbertus juniperoideus*, that *Herbertus sendtneri* does not occur in Macaronesia, and that this second Azorean species has affinities with South American taxa and plants from the Appalachians previously assigned to *Herbertus tenuis*.

⁷³Reports of *Herbertus aduncus* from Europe belong to *Herbertus hutchinsiae*, treated for a long time as a synonym or subspecies of *Herbertus aduncus*.

⁷⁴*Herbertus norenius* was described from Norway and Scotland by Bell *et al.* (2012).

⁷⁵*Bazzania tricrenata* var. *fulfordiae* W.S.Hong occurs in North America.

⁷⁶*Lepidozia cupressina* subsp. *africana* (Steph.) Pócs occurs in Africa and subsp. *natalensis* (Steph.) Pócs in southern Africa.

⁷⁷*Lepidozia stuhlmannii* was reported new to the Azores by Frahm (2005). *Lepidozia stuhlmannii* subsp. *pulvinata* (Steph.) Pócs and var. *carnosa* (Steph.) Pócs occur in Africa. The subspecific identity of the plants in the Azores is unknown.

⁷⁸*Tricholepidozia lindenberghii* was reported as new to Britain (and the Northern Hemisphere) by Blackstock *et al.* (2019).

⁷⁹*Tricholepidozia lindenberghii* var. *complanata* (J.J.Engel & G.L.Merr.) E.D.Cooper, var. *mellea* (J.J.Engel & G.L.Merr.) E.D.Cooper and var. *papillata* (J.J.Engel & G.L.Merr.) E.D.Cooper, as well as var. *lindenberghii* occur in New Zealand.

⁸⁰*Tricholepidozia* was elevated to a separate genus by Cooper *et al.* (2013) and *Tricholepidozia tetradactyla* was shown to be the same as *Telaranea longii* (Engel and Smith Merrill 2004) and *Telaranea murphyae* (Villarreal *et al.* 2014) from Britain.

Lophocoleaceae Vanden Berghen

68 *Chiloscyphus* Corda⁸¹

- 1 *C. pallescens* (Ehrh.) Dumort.
 - a var. *fragilis* (Roth) Müll.Frib. [*Chiloscyphus fragilis* (Roth) Schiffn.]
 - b var. *pallescens*
- 2 *C. polyanthos* (L.) Corda
 - a var. *polyanthos*
 - b var. *rivularis* (Schrad.) Lindb. & Arnell [*Chiloscyphus rivularis* (Schrad.) Hazl.]

69 *Heteroscyphus* Schiffn.

- 1 *H. denticulatus* (Mitt.) Schiffn.
- 2 *H. fissistipus* (Hook.f. & Taylor) Schiffn.⁸²

70 *Leptoscyphus* Mitt.

- subgenus *Anomylia* (R.M.Schust.) R.M.Schust.
- 1 *L. cuneifolius* (Hook.) Mitt.
- subgenus *Leptoscyphus*
- section *Leptoscyphus*
- 2 *L. porphyrius* (Nees) Grolle
 - a subsp. *azoricus* (H.Buch & Perss.) Vanderp. & Heinrichs [*Leptoscyphus azoricus* (H.Buch & Perss.) Grolle]⁸³

71 *Lophocolea* (Dumort.) Dumort.

- 1 *L. bidentata* (L.) Dumort. [*Lophocolea cuspidata* (Nees) Limpr.]⁸⁴
- 2 *L. bispinosa* (Hook.f. & Taylor) Gottsche, Lindenb. & Nees
- 3 *L. brookwoodiana* Paton & Sheahan⁸⁵
- 4 *L. coadunata* (Sw.) Mont. [*Chiloscyphus coadunatus* (Sw.) J.J.Engel & R.M.Schust., *Lophocolea cuspidata* auct. non (Nees) Limpr.], *Lophocolea bidentata* var. *rivularis* (Raddi) Schiffn., *Chiloscyphus latifolius* (Nees) J.J.Engel & R.M.Schust.]
- 5 *L. fragrans* (Moris & De Not.) Gottsche, Lindenb. & Nees [*Chiloscyphus fragrans* (Moris & De Not.) J.J.Engel & R.M.Schust.]

a subsp. *fragrans*⁸⁶

- 6 *L. heterophylla* (Schrad.) Dumort. [*Chiloscyphus profundus* (Nees) J.J.Engel & R.M.Schust.]
 - a subsp. *heterophylla*⁸⁷
- 7 *L. minor* Nees [*Chiloscyphus minor* (Nees) J.J.Engel & R.M.Schust.]
- 8 *L. semiteres* (Lehm.) Mitt.
 - a var. *semiteres*⁸⁸

Mastigophoraceae R.M.Schust.

- 72 *Mastigophora* Nees
 - 1 *M. woodsii* (Hook.) Nees

Plagiochilaceae Müll.Frib.

- 73 *Pedinophyllum* Lindb. ex Nordst.
 - 1 *P. interruptum* (Nees) Kaal.
- 74 *Plagiochila* (Dumort.) Dumort.

section *Arrectae* Carl

 - 1 *P. bifaria* (Sw.) Lindenb.
 - a var. *bifaria*⁸⁹
 - 2 *P. papillifolia* Steph.⁹⁰
 - 3 *P. punctata* (Taylor) Taylor
 - 4 *P. retrorsa* Gottsche
 - 5 *P. spinulosa* (Dicks.) Dumort.
 - 6 *P. stricta* Lindenb.

section *Fuscoluteae* Carl

 - 7 *P. heterophylla* Lindenb.
 - a var. *heterophylla* [*Plagiochila atlantica* F.Rose]⁹¹

section *Glaucescentes* Carl

 - 8 *P. longispina* Lindenb. & Gottsche

section *Plagiochila*

 - 9 *P. arctica* Bryhn & Kaal. [*Plagiochila asplenioides* subsp. *arctica* (Bryhn & Kaal.) R.M.Schust.]
 - a var. *arctica* [*Plagiochila asplenioides* var. *lobata* (Kaal.) Jørg.]⁹²

⁸¹The *Chiloscyphus polyanthos* complex has been treated in various ways in the past, without any recent solution. Some authors have treated it as one species (*Chiloscyphus polyanthos*) with two subspecies (subsp. *polyanthos* and *pallescens* (e.g. Smith 1990); others as four segregate species (*Chiloscyphus polyanthos*, *Chiloscyphus pallescens*, *Chiloscyphus fragilis* and *Chiloscyphus rivularis* (e.g. Konstantinova, Potemkin, et al. 1992). Grolle and Long (2000) kept two species, but did not deal with any subspecific taxa. Söderström, Urmi, et al. (2002) kept two species but recognised four varieties (*Chiloscyphus polyanthos* var. *polyanthos* and var. *rivularis* and *Chiloscyphus pallescens* var. *pallescens* and var. *fragilis*). Konstantinova, Bakalin, et al. (2009) also used this concept as well as the World Checklist of Hornworts and Liverworts (Söderström, Hagborg, et al. 2016). However, several recent publications do not recognise any varieties of the two species (e.g. Köckinger 2017). The distribution of the taxa recognised here is therefore not recorded consistently over the continent.

⁸²*Heteroscyphus fissistipus* is a neophyte in Europe, introduced from Australia/New Zealand where it occurs in three varieties, var. *fissistipus*, var. *multispinus* (E.A.Hodgs. & Allison) J.J.Engel and var. *repandus* J.J.Engel. Which of these varieties occur in Europe is unknown.

⁸³*Leptoscyphus porphyrius* subsp. *porphyrius* occurs in South America.

⁸⁴The *Lophocolea bidentata* complex has been treated in various ways in the past, and there is also some nomenclatural confusion. Váňa and Engel (2013) clarify the concepts, not only for Europe but world-wide. As we follow the World Check List (Söderström, Hagborg, et al. 2016), we here treat them in the genus *Lophocolea* as *Lophocolea bidentata* (autoicous) and *Lophocolea coadunata* (dioicous). Many earlier reports are confusing and the occurrences in some countries are still questioned.

⁸⁵*Lophocolea brookwoodiana* was described from Britain by Paton and Sheahan (2006).

⁸⁶*Lophocolea fragrans* subsp. *cocosana* G.Dauphin, Gradst. & M.I.Morales occurs on Cocos Island.

⁸⁷*Lophocolea heterophylla* subsp. *cladogyna* R.M.Schust. occurs in North America.

⁸⁸*Lophocolea semiteres* var. *retusa* (J.J.Engel) L.Söderstr. occurs in Australia.

⁸⁹*Plagiochila bifaria* var. *rosea* (R.M.Schust.) Heinrichs occurs in South America.

⁹⁰The primarily neotropical *Plagiochila papillifolia* was reported as new from the Azores by Heinrichs, Rycroft, et al. (2002).

⁹¹*Plagiochila heterophylla* var. *beauverdii* (Steph.) Heinrichs occurs in South America.

⁹²*Plagiochila arctica* var. *intermedia* R.M.Schust. occurs in North America. The synonymy of *Plagiochila asplenioides* var. *lobata* (Damsholt 2002) is problematic, as he states that *Plagiochila arctica* does not occur in Norway and the type of the synonymised variety is from the mountains in central Norway.

- 10 ***P. asplenioides*** (L.) Dumort. [*Plagiochila major* (Nees) S.W.Arnell]
- 11 ***P. britannica*** Paton
- 12 ***P. porelloides*** (Torr. ex Nees) Lindenb. [*Plagiochila asplenioides* subsp. *porelloides* (Torr. ex Nees) Kaal.]⁹³
- a var. ***norvegica*** (H.H.Blom & Holten) Schumacker & Váňa [*Plagiochila norvegica* H.H.Blom & Holten]
- b var. ***porelloides*** [*Plagiochila asplenioides* var. *devexa* Carrington, *Plagiochila asplenioides* var. *minor* Lindenb., *Plagiochila asplenioides* var. *humilis* (Nees) Lindenb.]
- c var. ***subarctica*** (Jørg.) Lammes [*Plagiochila asplenioides* var. *subarctica* Jørg.]
- section *Poeltiae* Inoue
- 13 ***P. carringtonii*** (Balf. ex Carrington) Grolle
- a subsp. ***carringtonii***⁹⁴
- section *Rutilantes* Carl
- 14 ***P. exigua*** (Taylor) Taylor
- 15 ***P. maderensis*** Gottsche ex Steph.⁹⁵
- section *Vagae* Lindenb.
- 16 ***P. virginica*** A.Evans [*Plagiochila dubia* auct. eur. non Lindenb. & Gottsche (= *Plagiochila patula* (Sw.) Lindenb.)]
- a var. ***virginica***⁹⁶

Trichocoleaceae Nakai

- 75 ***Trichocolea*** Dumort.
- 1 ***T. tomentella*** (Ehrh.) Dumort.

Myliaceae Schljakov

- 76 ***Mylia*** Gray
- subgenus *Anomalae* (R.M.Schust. ex Potemkin) L. Söderstr.
- 1 ***M. anomala*** (Hook.) Gray
- subgenus *Mylia*
- 2 ***M. taylorii*** (Hook.) Gray

Porellales Schljakov
Frullaniaceae Lorch

- 77 ***Frullania*** Raddi
- subgenus *Frullania*
- 1 ***F. azorica*** Sim-Sim, Sérgio, Mues & Kraut
- 2 ***F. dilatata*** (L.) Dumort.
- a subsp. ***dilatata***⁹⁷
- 3 ***F. ericoides*** (Nees) Mont.
- a var. ***ericoides***⁹⁸
- 4 ***F. fragilifolia*** (Taylor) Gottsche, Lindenb. & Nees
- 5 ***F. jackii*** Gottsche [*Frullania davurica* subsp. *jackii* (Gottsche) S.Hatt.]
- 6 ***F. oakesiana*** Austin
- a subsp. ***oakesiana***⁹⁹
- 7 ***F. parvistipula*** Steph.
- 8 ***F. riparia*** Hampe [*Frullania cesatiana* De Not.]¹⁰⁰
- 9 ***F. stylifera*** (R.M.Schust.) R.M.Schust.¹⁰¹
- subgenus *Thyopsiella* Spruce
- 10 ***F. acicularis*** Hentschel & von Konrat
- [*Frullania tamarisci* var. *azorica* J.-P.Frahm]¹⁰²
- 11 ***F. calcarifera*** Steph.¹⁰³
- 12 ***F. microphylla*** (Gottsche) Pearson
- [*Frullania microphylla* var. *deciduifolia* Grolle]¹⁰⁴
- 13 ***F. polysticta*** Lindenb.
- 14 ***F. sergiae*** Sim-Sim, Fontinha, Mues & Lion
- 15 ***F. subarctica*** Vilnet, Borovich. & Bakalin¹⁰⁵
- 16 ***F. tamarisci*** (L.) Dumort. [*Frullania tamarisci* var. *atrovirens* Carrington, *Frullania tamarisci* var. *cornubica* Carrington, *Frullania tamarisci* var. *ericetorum* Jørg., *Frullania tamarisci* var. *robusta* Lindb., *Frullania tamarisci* var. *sardoa* (De Not.) De Not.]
- 17 ***F. teneriffae*** (F.Weber) Nees

⁹³*Plagiochila porelloides* is a variable species and many varieties have been described. Their value is questionable but the three included here have frequently been recognised in recent studies.

⁹⁴*Plagiochila carringtonii* subsp. *lobuchensis* Grolle occurs in Himalaya.

⁹⁵*Plagiochila maderensis* was placed in synonymy with *Plagiochila spinulosa* by Grolle (1967) but reinstated by Rycroft et al. (2004).

⁹⁶*Plagiochila virginica* var. *caroliniana* R.M.Schust. and var. *euryphylla* R.M.Schust. occur in North America.

⁹⁷*Frullania dilatata* subsp. *asiatica* S.Hatt. occurs in E Asia.

⁹⁸*Frullania ericoides* var. *laxa* (Gottsche, Lindenb. & Nees) Schiffn. occurs in SE Asia and Mexico, var. *minor* Kamim. in Japan and var. *verrucosa* (Kamim.) Hentschel & von Konrat in Japan.

⁹⁹*Frullania oakesiana* subsp. *takayuensis* (Steph.) R.M.Schust. occurs in E Asia.

¹⁰⁰In spite of being under discussion for over 30 years, the problem of the *Frullania riparia* (American) / *cesatiana* (European) / *musciola* (Asiatic) complex has not yet been solved. As concluded in Bisang et al. (1989), if they are the same, the name *Frullania riparia* has priority. However, if the American populations are different from the European and Asiatic populations (as treated in Bisang et al. 1989), *Frullania cesatiana* would be the correct name (note that their 'combination' *Frullania cesatiana* var. *musciola* (Steph.) Bisang et al. is a *nomen nudum*). Grolle (in Grolle and Long 2000) argued that the American *Frullania riparia* and European *Frullania cesatiana* are conspecific, but separate from the Asiatic *Frullania musciola* Steph. For now we are following this concept pending further research, preferably including molecular data.

¹⁰¹*Frullania stylifera*, until recently considered endemic to eastern North America, has recently been recognised from European and Asian Russia (Konstantinova et al. 2020).

¹⁰²*Frullania acicularis* was described from the Azores by Frahm (2006) as *Frullania tamarisci* var. *azorica* but shown to be separate from *Frullania tamarisci* by Heinrichs, Hentschel, et al. (2010) and Vilnet, Borovichev, et al. (2014) and elevated to species level by Hentschel et al. (2015)

¹⁰³*Frullania calcarifera* was placed in synonymy of *Frullania tamarisci* by Hattori (1972) but shown to be a species separate from it by Heinrichs, Hentschel, et al. (2010) and Vilnet, Borovichev, et al. (2014).

¹⁰⁴*Frullania microphylla* var. *deciduifolia* is not recognized by Sim-Sim (1999) as "all specimens from Portugal or Madeira have deciduous leaves", a treatment followed here as well as in the world checklist (Söderström, Hagborg, et al. 2016).

¹⁰⁵*Frullania subarctica* was described by Vilnet, Borovichev, et al. (2014) from the Russian Far East but shown also to occur in northern Europe.

incertae sedis

18 ***F. bolanderi*** Austin

19 ***F. cleistostoma*** Schiffn. & W.Wollny [*Frullania inflata* auct. eur. non Gottsche]¹⁰⁶

Jubulaceae H.Klinggr.

78 ***Jubula*** Dumort.

1 ***J. hutchinsiae*** (Hook.) Dumort.¹⁰⁷

a subsp. ***caucasica*** Konstant. & Vilnet¹⁰⁸

b subsp. ***hutchinsiae***

Lejeuneaceae Cavers

Lejeuneoideae C.Massal.

trib. Brachiolejeuneae van Slageren & Berendsen

subtrib. Brachiolejeuneinae Gradst.

79 ***Acanthocoleus*** R.M.Schust.

1 ***A. aberrans*** (Lindenb. & Gottsche) Kruijt

a var. ***laevis*** Gradst.¹⁰⁹

trib. Lejeuneae Dumort.

subtrib. Cheilolejeuneinae Gradst.

80 ***Cheilolejeunea*** (Spruce) Steph.

subgenus *Eosmolejeunea* (Spruce) Kachroo

1 ***C. cedercreutzii*** (H.Buch & Perss.) Grolle

subtrib. Cololejeuneinae Gradst.

81 ***Cololejeunea*** (Spruce) Steph. [*Aphanolejeunea* A.Evans]

subgenus *Aphanolejeunea* (A.Evans) Pócs

1 ***C. madeirensis*** Schiffn. [*Aphanolejeunea madeirensis* (Schiffn.) Grolle]

2 ***C. microscopica*** (Taylor) Schiffn. [*Aphanolejeunea microscopica* (Taylor) A.Evans]

a var. ***microscopica***¹¹⁰

3 ***C. sintenisii*** (Steph.) Pócs [*Aphanolejeunea sintenisii* Steph.]

subgenus *Cololejeunea*

4 ***C. calcarea*** (Lib.) Steph.

5 ***C. rosettiana*** (C.Massal.) Schiffn.

6 ***C. schaeferi*** Grolle

subgenus *Diaphanae* R.M.Schust.

7 ***C. azorica*** V.Allorge & Jovet-Ast [*Aphanolejeunea azorica* (V.Allorge & Ast) Bernecker & Pócs]

82 ***Colura*** (Dumort.) Dumort.

subgenus *Colura*

section *Colura*

1 ***C. calyptrifolia*** (Hook.) Dumort.

83 ***Myriocoleopsis*** Schiffn.

1 ***M. minutissima*** (Sm.) R.L.Zhu, Y.Yu & Pócs [*Cololejeunea minutissima* (Sm.) Schiffn.]

a subsp. ***minutissima***¹¹¹

subtrib. Drepanolejeuneinae Gradst.

84 ***Drepanolejeunea*** (Spruce) Steph.

subgenus *Drepanolejeunea*

1 ***D. hamatifolia*** (Hook.) Schiffn.

subtrib. Lejeuneinae Gradst.

85 ***Harpalejeunea*** (Spruce) Schiffn.

subgenus *Harpalejeunea*

1 ***H. molleri*** (Steph.) Grolle

a subsp. ***molleri***¹¹²

86 ***Lejeunea*** Lib.

subgenus *Lejeunea*

1 ***L. cavifolia*** (Ehrh.) Lindb.

2 ***L. eckloniana*** Lindenb. [*Lejeunea holtii* Spruce]¹¹³

3 ***L. flava*** (Sw.) Nees

a subsp. ***moorei*** (Lindb.) R.M.Schust.¹¹⁴

4 ***L. hibernica*** Bischl., H.A.Mill. & Bonner ex Grolle

5 ***L. lamacerina*** (Steph.) Schiffn.

a subsp. ***lamacerina***¹¹⁵

6 ***L. mandonii*** (Steph.) Müll.Frib.

7 ***L. patens*** Lindb.

subgenus *Nanolejeunea* R.M.Schust.

8 ***L. canariensis*** (Steph.) Steph. [*Lejeunea laetevirens* auct.]¹¹⁶

¹⁰⁶*Frullania inflata* Gottsche as treated in Söderström, Hagborg, et al. (2016) is a species complex. *Frullania inflata* s.str. is confined to North America and reports from Europe belongs to the segregate *Frullania cleistostoma* (Mamontov, Potemkin, et al. 2018).

¹⁰⁷*Jubula hutchinsiae* is a complex species with many subspecies. In addition to the two subspecies occurring in Europe, five more are recognised in Söderström, Hagborg, et al. (2016), subsp. *australiae* Pócs & A.Cairns (Australia), subsp. *bogotensis* (Steph.) Verd. (S and C America), subsp. *japonica* (Steph.) Horik. & Ando (E Asia), subsp. *javanica* (Steph.) Verd. (E and SE Asia) and subsp. *pennsylvanica* (Steph.) Verd. (North America).

¹⁰⁸*Jubula hutchinsiae* subsp. *caucasica* was described by Konstantinova and Vilnet (2011). Prior to the description, the subspecies occurring in the Caucasus was regarded as subsp. *javanica* (Steph.) Verd.

¹⁰⁹*Acanthocoleus aberrans* var. *aberrans* seems to be confined to South America.

¹¹⁰*Cololejeunea microscopica* var. *africana* (Pócs) Pócs & Bernecker and var. *exigua* (A.Evans) Pócs occur in Africa and South America.

¹¹¹*Myriocoleopsis minutissima* subsp. *myriocarpa* (Nees & Mont.) R.L.Zhu, Y.Yu & Pócs is widespread in the tropics.

¹¹²*Harpalejeunea molleri* subsp. *integra* (R.M.Schust.) Damsh. occurs in North America.

¹¹³*Lejeunea holtii* was placed in synonymy with *Lejeunea eckloniana* by Dirkse et al. (1993). However, the two taxa differ slightly in morphology and a molecular study is needed to verify the synonymy.

¹¹⁴*Lejeunea flava* subsp. *flava* is widespread in the tropics, subsp. *orientalis* R.M.Schust. occurs in E and SE Asia, subsp. *tabularis* (Spreng.) S.W.Arnell in Africa and var. *pellucida* Lindenb. & Gottsche in South America.

¹¹⁵*Lejeunea lamacerina* subsp. *gemminata* R.M.Schust. occurs in North America.

¹¹⁶*Lejeunea canariensis* from Madeira was nested within *Lejeunea laetevirens* in the study by Heinrichs, Dong, et al. (2013). However, *Lejeunea laetevirens* seems to be a complex species with several other taxa also nested within it. We keep *Lejeunea canariensis* as a separate species for now, recognizing that it seems to be the only element of the *Lejeunea laetevirens* complex occurring in Macaronesia.

- 87 **Microlejeunea** (Spruce) Steph.
1 **M. ulicina** (Taylor) Steph. [*Lejeunea ulicina* (Taylor) Gottsche, Lindenb. & Nees]

Ptychanthoideae Mizut.

- 88 **Marchesinia** Gray
subgenus *Marchesinia*
1 **M. mackaii** (Hook.) Gray

Porellaceae Cavers

- 89 **Porella** L.
1 **P. arboris-vitae** (With.) Grolle
a subsp. **arboris-vitae** [*Porella arboris-vitae* var. *killarniensis* (Pearson) M.F.V. Corley, *Porella arboris-vitae* var. *obscura* (Nees) M.F.V. Corley]¹¹⁷
2 **P. baueri** (Schiffn.) C.E.O. Jensen
3 **P. canariensis** (F. Weber) Underw.
4 **P. cordaeana** (Huebener) Moore [*Porella cordaeana* var. *faeroensis* (C.E.O. Jensen) E.W. Jones, *Porella cordaeana* var. *simplior* (J.E. Zetterst.) Arnell]
5 **P. inaequalis** (Gottsche) Perss.
6 **P. obtusata** (Taylor) Trevis.
7 **P. pinnata** L.
8 **P. platyphylla** (L.) Pfeiff. [*Porella platyphylla* var. *subsquarrosa* (Schiffn.) Arnell]

Radulaceae Müll. Frib.

- 90 **Radula** Dumort.
subgenus *Radula*
1 **R. aquilegia** (Hook. f. & Taylor) Gottsche, Lindenb. & Nees
2 **R. carringtonii** J.B. Jack
3 **R. complanata** (L.) Dumort. [*Radula complanata* var. *alpestris* (Lindb. ex Berggr.) Lindb.]
4 **R. jonesii** Bouman, Dirkse & K. Yamada
5 **R. lindenbergiana** Gottsche ex C. Hartm. [*Radula complanata* subsp. *lindenbergiana* (Gottsche ex C. Hartm.) R.M. Schust., *Radula lindenbergiana* Gottsche ex J.B. Jack *nom. inval.*]

- 6 **R. visianica** C. Massal.¹¹⁸
7 **R. wichurae** Steph.
subgenus *Volutoradula* Devos, M.A.M. Renner, Gradst., A.J. Shaw & Vanderp.
8 **R. holtii** Spruce
9 **R. nudicaulis** Steph.
a var. **delicatula** P. Allorge & V. Allorge
b var. **nudicaulis**
10 **R. voluta** Taylor

Ptilidiales Schljakov

Ptilidiaceae H. Klinggr.

- 91 **Ptilidium** Nees
1 **P. ciliare** (L.) Hampe
2 **P. pulcherrimum** (Weber) Vain. [*Ptilidium pulcherrimum* var. *subpinnatum* (Jørg.) Damsh.]

Metzgeriales Chalaud

Aneuraceae H. Klinggr.

- 92 **Aneura** Dumort. [*Cryptothallus* Malmb.]¹¹⁹
1 **A. latissima** Spruce [*Aneura pseudopinguis* (Herzog) Pócs]¹²⁰
2 **A. maxima** (Schiffn.) Steph.¹²¹
3 **A. mirabilis** (Malmb.) Wickett & Goffinet [*Cryptothallus mirabilis* Malmb.]
4 **A. pinguis** (L.) Dumort. [*Aneura pinguis* var. *angustior* (Hook.) Dumort., *Aneura pinguis* var. *denticulata* (Nees) Godelin., *Aneura pinguis* var. *fuscovirens* (Lindb.) Damsh.]

93 **Riccardia** Gray¹²²

- 1 **R. chamedryfolia** (With.) Grolle [*Riccardia chamedryfolia* var. *major* (Nees) R.M. Schust. *nom. illeg.*, *Riccardia chamedryfolia* var. *submersa* (C.E.O. Jensen ex Müll. Frib.) Damsh. *nom. inval.*]
2 **R. incurvata** Lindb.
3 **R. latifrons** (Lindb.) Lindb.
a subsp. **arctica** R.M. Schust. & Damsh.
b subsp. **latifrons**¹²³
4 **R. multifida** (L.) Gray
a subsp. **multifida**¹²⁴
5 **R. palmata** (Hedw.) Carruth.

¹¹⁷*Porella arboris-vitae* subsp. *nitidula* (C. Massal.) S. Hatt. occurs in E Asia.

¹¹⁸*Radula visianica* was until recently only known from two collections in N. Italy, the type in 1878 and a later collection in the southern Alps in the 1930s, but was recently found in several localities in Austria (Köckinger 2016). It has been shown to be well separated molecularly from other species in subgenus *Radula* (Váňa et al. 2017).

¹¹⁹The genus *Aneura* includes several genetically distinct lineages (cf. Bączkiewicz et al. 2017) and the European '*Aneura maxima*' is genetically different from the population in the type locality on Java (D.G. Long, pers. comm.). It is also unlikely that *Aneura latissima* (syn. *Aneura pseudopinguis*) from South America represents any of the European lineages. However, until the taxonomy is further clarified, the 'traditional' treatment is followed here, as it was in Söderström, Hagborg, et al. (2016).

¹²⁰*Aneura latissima* was reported (as *Aneura pseudopinguis*) new to Europe by Sérgio and Garcia (2009). The type of both names is from South America and it is unlikely that the European plants represent this taxon. It is more probable that they belong to some of the unpublished European lineages (Long et al. in prep.).

¹²¹*Aneura maxima* is a SE Asian taxon that is not conspecific with the taxon in Europe so named. The latter is at present lacking a valid name.

¹²²*Riccardia* is classified into several subgenera and sections in Söderström, Hagborg, et al. (2016). The European species all belong to subgenus *Riccardia*, and mostly also to section *Riccardia*, except for *R. incurvata*, the position of which is unclear.

¹²³*Riccardia latifrons* var. *miyakeana* (Schiffn.) Furuki occurs in Japan.

¹²⁴*Riccardia multifida* subsp. *decreescens* (Steph.) Furuki occurs in E Asia and subsp. *synoica* R.M. Schust. in SE USA.

Metzgeriaceae H.Klinggr.

- 94 **Metzgeria** Raddi [*Apometzgeria* Kuwah.]
- 1 ***M. conjugata*** Lindb. [*Metzgeria conjugata* var. *alipila* Kaal., *Metzgeria conjugata* var. *macvicarii* Kaal.]
 - 2 ***M. consanguinea*** Schiffn. [*Metzgeria temperata* auct. eur. non Kuwah.]¹²⁵
 - 3 ***M. furcata*** (L.) Corda [*Metzgeria furcata* var. *expansa* Douin, *Metzgeria furcata* var. *flexipilis* Kaal., *Metzgeria furcata* var. *ulvula* (Nees) Pavletic]¹²⁶
 - 4 ***M. leptoneura*** Spruce [*Metzgeria hamata* auct. non Lindb.]¹²⁷
 - a var. ***leptoneura***¹²⁸
 - 5 ***M. pubescens*** (Schrank) Raddi [*Apometzgeria pubescens* (Schrank) Kuwah.]
 - 6 ***M. simplex*** Lorb. ex Müll.Frib. [*Metzgeria conjugata* subsp. *simplex* (Lorb. ex Müll.Frib.) R.M.Schust.]
 - 7 ***M. violacea*** (Ach.) Dumort. [*Metzgeria fruticulosa* auct. non (O.F.Müll.) A.Evans]¹²⁹

Pleuroziales Schljakov

Pleuroziaceae Müll.Frib.

- 95 **Pleurozia** Dumort.
subgenus *Constantifolia* B.M.Thiers
- 1 ***P. purpurea*** Lindb.

Fossombroniales Schljakov

Calyculariaceae He-Nygrén, Juslén, Ahonen, Glenny & Piippo

- 96 **Calycularia** Mitt.
- 1 ***C. laxa*** Lindb. & Arnell¹³⁰

Fossombroniaceae Hazsl.

- 97 **Fossombronia** Raddi
- 1 ***F. angulosa*** (Dicks.) Raddi
 - 2 ***F. caespitiformis*** (Raddi) De Not. ex Rabenh.

- a subsp. ***caespitiformis***
- b subsp. ***multispira*** (Schiffn.) J.R.Bray & Cargill [*Fossombronia husnotii* Corb.]
- 3 ***F. crispa*** Nees [*Fossombronia zeyheri* Steph.]¹³¹
- 4 ***F. echinata*** Macvicar
- 5 ***F. fimbriata*** Paton
- 6 ***F. fleischeri*** Osterwald ex Loeske¹³²
- 7 ***F. foveolata*** Lindb.
- 8 ***F. incurva*** Lindb.
- 9 ***F. leucoxantha*** (Lehm.) Lehm. & Lindenb.¹³³
- 10 ***F. maritima*** (Paton) Paton
- 11 ***F. mittenii*** Tind. [*Fossombronia crozalsii* Corb.]
- 12 ***F. pusilla*** (L.) Nees
- 13 ***F. wondraczekii*** (Corda) Dumort. ex Lindb.

Petalophyllaceae Stotler & Crand.-Stotl.

- 98 **Petalophyllum** Nees & Gottsche
- 1 ***P. ralfsii*** (Wilson) Nees & Gottsche

Pallaviciniales W.Frey & M.Stech

Moerckiaceae K.I.Goebel ex Stotler & Crand.-Stotl.

- 99 **Moerckia** Gottsche
- 1 ***M. blyttii*** (Mørch) Brockm.
 - 2 ***M. flotoviana*** (Nees) Schiffn.¹³⁴
 - 3 ***M. hibernica*** (Hook.) Gottsche

Pallaviciniaceae Mig.

Pallavicinioideae Mig. ex Grolle

- 100 **Pallavicinia** Gray
- 1 ***P. lyellii*** (Hook.) Gray

Pelliales He-Nygrén

Pelliaceae H.Klinggr.

- 101 **Apopellia** (Grolle) Nebel & D.Quandt
- 1 ***A. endiviifolia*** (Dicks.) Nebel & D.Quandt
[*Pellia endiviifolia* (Dicks.) Dumort.]

¹²⁵*Metzgeria temperata* was synonymised with the primarily pantropical *Metzgeria consanguinea* by Hill et al. (2008). However, unpublished molecular studies show that the European populations are closer to *Metzgeria violacea* than to the Japanese *Metzgeria temperata* (see Köckinger 2017). Until the position of the European *Metzgeria 'temperata'* is clarified, we use the name *Metzgeria consanguinea*, although with some hesitation.

¹²⁶North American populations of *Metzgeria furcata* have recently been recognised as a separate species, *Metzgeria setigera* R.M.Schust. ex Crand.-Stotl. & L. Söderstr. (Söderström, Vána, et al. 2015). *Metzgeria furcata* var. *pacifica* Brinkm. occurs in E North America, and probably belongs to *Metzgeria setigera*. *Metzgeria furcata* var. *expansa* was recognised in Söderström, Hagborg, et al. (2016), although with low confidence. It is known only from the type specimen in France and we do not recognise it here, as there are several other equally doubtful 'varieties' that could also be included. The whole *Metzgeria furcata* complex needs further study in Europe (and elsewhere).

¹²⁷*Metzgeria hamata* Lindb. is technically a synonym of *Metzgeria procera* Mitt., but identical to *Metzgeria leptoneura* in the sense of Lindberg (1877) and all subsequent authors.

¹²⁸*Metzgeria leptoneura* var. *breviseta* (Schiffn.) O.Yano occurs in Brazil and var. *polychaeta* R.M.Schust. in SE USA.

¹²⁹Grolle and So (2003) demonstrated that the name *Metzgeria fruticulosa* technically belongs to *Riccardia palmata* and that *Metzgeria violacea* is the name that should be used.

¹³⁰*Calycularia laxa* was reported new to European Russia from Nenets (Konstantinova and Lavrinenko 2002) and Murmansk (Sofronova et al. 2013) provinces.

¹³¹*Fossombronia crispa* was first reported from Portugal by Sérgio (1985) but Perold (1997) showed that her concept of the species was wrong. Later, Sérgio (2003) referred the plants to *Fossombronia leucoxantha*. However, Sérgio (1985) also reported *Fossombronia zeyheri*, which Perold (1997) showed was a synonym of *Fossombronia crispa* but Grolle and Long (2000) treated it as a synonym of *Fossombronia foveolata*.

¹³²*Fossombronia fleischeri* has long been considered a synonym of *Fossombronia incurva*, but Stotler et al. (2003) showed that they should be regarded as distinct species.

¹³³*Fossombronia leucoxantha* was reported from Portugal by Sérgio (1985) as *Fossombronia crispa* but Perold (1997) pointed out that her concept of *Fossombronia crispa* was wrong. Sérgio (2003) later referred the plants to *Fossombronia leucoxantha*.

¹³⁴*Moerckia flotoviana* was synonymised with *Moerckia hibernica* by De Sloover (1959) and subsequently usually treated as such in the European literature. However, Crandall-Stotler and Stotler (2007) showed that they are distinct taxa. Thus, many reports of *Moerckia hibernica* belong to *Moerckia flotoviana*.

- 102 **Pellia** Raddi
 1 **P. epiphylla** (L.) Corda
 a subsp. **borealis** (Lorb.) Messe
 b subsp. **epiphylla**
 2 **P. neesiana** (Gottsche) Limpr.

MARCHANTIOPSIDA CRONQUIST, TAKHT. & W.ZIMM.

Blasiales Stotler & Crand.-Stotl.

Blasiaceae H.Klinggr.

- 103 **Blasia** L.
 1 **B. pusilla** L.

Lunulariales H.Klinggr.

Lunulariaceae H.Klinggr.

- 104 **Lunularia** Adans.
 1 **L. cruciata** (L.) Dumort. ex Lindb.
 a subsp. **cruciata**¹³⁵

Marchantiales Limpr.

Aytoniaceae Cavers

- 105 **Asterella** P.Beauv. [*Fimbraria* Nees]
 subgenus *Asterella*
 section *Brachyblepharis* (Nees) D.G.Long
 1 **A. africana** (Mont.) Underw. ex A.Evans
 subgenus *Phragmoblepharis* Grolle
 2 **A. lindenberghiana** (Corda ex Nees) Lindb. ex
 Arnell
 subgenus *Saccatae* (Grolle) D.G.Long
 3 **A. saccata** (Wahlenb.) A.Evans

- 106 **Mannia** Corda
 subgenus *Mannia*
 1 **M. androgyna** (L.) A.Evans
 2 **M. californica** (Gottsche) L.C.Wheeler¹³⁶
 3 **M. controversa** (Meyl.) D.B.Schill¹³⁷
 a subsp. **controversa**¹³⁸
 4 **M. fragrans** (Balb.) Frye & L.Clark
 a subsp. **fragrans** [*Mannia fragrans* var. *bre-vipes* (Kaal.) Damsh.]¹³⁹
 5 **M. sibirica** (Müll.Frib.) Frye & L.Clark
 subgenus *Neesiella*
 6 **M. gracilis** (F.Weber) D.B.Schill & D.G.Long
 [*Asterella gracilis* (F.Weber) Underw.]

- 7 **M. pilosa** (Hornem.) Frye & L.Clark

- 8 **M. triandra** (Scop.) Grolle

- 107 **Plagiochasma** Lehm.
 subgenus *Micropylum*
 1 **P. rupestre** (J.R.Forst. & G.Forst.) Steph.
 a var. **rupestre**¹⁴⁰
 subgenus *Plagiochasma*
 2 **P. appendiculatum** Lehm. & Lindenb.¹⁴¹

- 108 **Reboulia** Raddi
 1 **R. hemisphaerica** (L.) Raddi
 a subsp. **australis** R.M.Schust.
 b subsp. **dioica** R.M.Schust.
 c subsp. **hemisphaerica**¹⁴²

Cleveaceae Cavers

- 109 **Clevea** Lindb. [*Athalamia* auct. eur.]
 1 **C. hyalina** (Sommerf.) Lindb. [*Athalamia hyalina* (Sommerf.) S.Hatt., *Clevea hyalina* var. *suecica* (Lindb.) Lindb., *Athalamia hyalina* var. *suecica* (Lindb.) S.Hatt., *Clevea hyalina* var. *rufescens* (S.W.Arnell) Konstant. *nom. inval.*]
 2 **C. spathysii** (Lindenb.) Müll.Frib. [*Athalamia spathysii* (Lindenb.) S.Hatt.]

- 110 **Peltolepis** Lindb.
 1 **P. quadrata** (Saut.) Müll.Frib.

- 111 **Sauteria** Nees
 Section *Sauteria*
 1 **S. alpina** (Nees) Nees

Conocephalaceae Müll.Frib. ex Grolle

- 112 **Conocephalum** Hill
 1 **C. conicum** (L.) Dumort.
 2 **C. salebrosum** Szwedk., Buczk. & Odrzyk.¹⁴³

Corsiniaceae Engl.

Corsinioideae Schiffn.

- 113 **Corsinia** Raddi
 1 **C. coriandrina** (Spreng.) Lindb.

¹³⁵*Lunularia cruciata* subsp. *thaxteri* (A.Evans & Herzog) R.M.Schust. occurs in South America and New Zealand.

¹³⁶*Mannia californica* was reported new to Europe by Hugonnot and Schill (2006).

¹³⁷*Mannia controversa* (type from Switzerland) was usually regarded as a synonym of *Mannia fragrans* until Schill et al. (2008) showed it to be an independent species.

¹³⁸*Mannia controversa* subsp. *asiatica* D.B.Schill & D.G.Long occurs in Asia.

¹³⁹*Mannia fragrans* subsp. *orientalis* R.M.Schust. occurs in Japan.

¹⁴⁰*Plagiochasma rupestre* var. *volkii* Bischl. occurs in South America and southern Africa.

¹⁴¹*Plagiochasma appendiculatum* was reported new to Europe from the Balears by Cros et al. (2005).

¹⁴²*Reboulia hemisphaerica* subsp. *paradoxa* R.M.Schust. is described from Portugal (Schuster 1992b) but invalidly as it lacks a Latin description. If recognised it needs to be validated, but there are many uncertainties about the segregates of *Reboulia hemisphaerica*. *Reboulia hemisphaerica* subsp. *acrogyna* (R.M.Schust.) R.M.Schust. occurs in North America, subsp. *orientalis* R.M.Schust. in Asia, var. *fissisquama* Herzog in China and var. *turkestanica* C.E.O.Jensen ex Herzog in China.

¹⁴³*Conocephalum salebrosum* was described by Szwedkowski et al. (2005) from Poland but shown to have a wide distribution both in Europe and elsewhere in the Northern Hemisphere. On the other hand, they showed that *Conocephalum conicum* was a near-endemic to Europe.

Cyathodiaceae Stotler & Crand.-Stotl.

- 114 **Cyathodium** Kunze
1 **C. foetidissimum** Schiffn.¹⁴⁴

Dumortieraceae D.G.Long

- 115 **Dumortiera** Nees
1 **D. hirsuta** (Sw.) Nees
a subsp. **hirsuta**¹⁴⁵

Exormothecaceae Müll.Frib. ex Grolle

- 116 **Exormotheca** Mitt.
subgenus *Corbierella* (Douin & Trab.) Schiffn.
1 **E. welwitschii** Steph.
subgenus *Exormotheca*
2 **E. pustulosa** Mitt.

Marchantiaceae Lindl.

Marchantioideae Schiffn.

- 117 **Marchantia** L.
subgenus *Chlamidium* (Corda) Bischl.
section *Paleacea* Bischl.
1 **M. paleacea** Bertol.
a subsp. **paleacea**¹⁴⁶
subgenus *Marchantia*
2 **M. polymorpha** L.¹⁴⁷
a subsp. **montivagans** Bischl. & Boissel.-Dub. [*Marchantia alpestris* (Nees) Burgeff]
b subsp. **polymorpha** [*Marchantia aquatica* (Nees) Burgeff]
c subsp. **ruderalis** Bischl. & Boissel.-Dub. [*Marchantia latifolia* Gray]
subgenus *Preissia* (Corda) D.G.Long, Crand.-Stotl., L.L.Forrest & Villarreal [*Bucegia* Radian, *Preissia* Corda]
3 **M. quadrata** Scop. [*Preissia quadrata* (Scop.) Nees]
a subsp. **hyperborea** (R.M.Schust.) Borovich. [*Preissia quadrata* subsp. *hyperborea* R.M.Schust.]
b subsp. **quadrata**
4 **M. romanica** (Radian) D.G.Long, Crand.-Stotl., L.L.Forrest & J.C.Villarreal [*Bucegia romanica* Radian]

Oxymitraceae Müll.Frib. ex Grolle

- 118 **Oxymitra** Bisch. ex Lindenb.
1 **O. incrassata** (Brot.) Sérgio & Sim-Sim

Ricciaceae Rchb.

- 119 **Riccia** L.
subgenus *Riccia*
section *Riccia*¹⁴⁸
1 **R. atlantica** Sérgio & Perold
2 **R. atromarginata** Levier
a var. **atromarginata**¹⁴⁹
3 **R. beyrichiana** Hampe [*Riccia marginata* Lindb.]
4 **R. bicarinata** Lindb.
5 **R. bifurca** Hoffm. [*Riccia bifurca* var. *subinermis* Heeg]
6 **R. breidlerii** Jur. ex Steph.
7 **R. ciliata** Hoffm. [*Riccia ciliata* var. *epilosa* Warnst., *Riccia ciliata* var. *intumescens* Bisch., *Riccia ciliata* var. *violacea* Kny, *Riccia dalslandica* S.W.Arnell, *Riccia intumescens* (Bisch.) Underw., *Riccia canescens* Steph., *Riccia trichocarpa* M.Howe, *Riccia crinita* auct. eur.]¹⁵⁰
8 **R. ciliifera** Link [*Riccia melitensis* C.Massal.]
9 **R. crozalsii** Levier
10 **R. crustata** Trab.
11 **R. glauca** L. [*Riccia glauca* var. *major* (Roth) Lindenb.]
a var. **ciliaris** Warnst. [*Riccia glauca* var. *subinermis* (Lindb.) Warnst.]
b var. **glauca**
12 **R. gothica** Damsh. & Hallingb.
13 **R. gougetiana** Durieu & Mont.
a var. **armatissima** Levier ex Müll.Frib.
b var. **gougetiana**
14 **R. lamellosa** Raddi
15 **R. ligula** Steph.
16 **R. macrocarpa** Levier
17 **R. michelii** Raddi
18 **R. nigrella** DC.
19 **R. papillosa** Moris
20 **R. sommieri** Levier
21 **R. sorocarpa** Bisch.
a subsp. **arctica** R.M.Schust. ex Köckinger & L. Söderstr. [*Riccia sorocarpa* subsp. *arctica* R.M.Schust. *nom. inval.*, *Riccia lindenberiana* Saut.]

¹⁴⁴*Cyathodium foetidissimum* was recorded new to Europe by Ligrone and Duckett (2005).

¹⁴⁵*Dumortiera hirsuta* subsp. *nepalensis* (Taylor) R.M.Schust. is widespread in the tropics and subsp. *tatunoi* Horik. occurs in Japan.

¹⁴⁶*Marchantia paleacea* subsp. *diptera* (Nees & Mont.) Inoue occurs in E Asia.

¹⁴⁷Recent molecular studies (Cronberg et al., unpubl.) indicate that the subspecies of *Marchantia polymorpha* should be better treated at species level. However, we keep them as subspecies until nomenclatural uncertainties raised by elevating them to species are resolved.

¹⁴⁸As noted by Hugonnot (2010a), the nomenclature and taxonomy of many species in sect. *Riccia* is highly problematic. Difficult taxa include *Riccia bicarinata*, *R. bifurca*, *R. ciliata*, *R. crozalsii*, *R. michelii*, *R. subbifurca*, *R. warnstorffii* and the little-known *R. ligula*. The morphology of these taxa is plastic, and it is unclear to what extent the size of the thallus and the development and structure of cilia are determined by genetic or environmental factors. Jovet-Ast (1986) published an important monograph of the Mediterranean species, but her concepts have been questioned in recent years (e.g. Hugonnot 2010b, 2015). Integrated molecular and morphological studies are urgently needed.

¹⁴⁹*Riccia atromarginata* var. *jovet-astiae* Rauh & Buchloh occurs on the Arabian Peninsula and adjacent parts of Africa.

¹⁵⁰*Riccia trichocarpa* was synonymised with the Australian *Riccia crinita* (the latter having priority) by Jovet-Ast (2000), a synonymisation that has been largely overlooked or rejected by European bryologists. Hugonnot (2010b) argued that Jovet-Ast's (1986) concept of *Riccia trichocarpa* is actually *Riccia ciliata*, and her *Riccia ciliata* belongs to other species. Thus Hugonnot synonymised both taxa under the oldest name, *Riccia ciliata*.

b subsp. **erythrophora** R.M.Schust. ex Konstant. & L.Söderstr. [*Riccia sorocarpa* subsp. *erythrophora* R.M.Schust. *nom. inval.*]

c subsp. **sorocarpa**¹⁵¹ [*Riccia sorocarpa* var. *heegii* Schiffn.]

22 **R. subbifurca** Warnst. ex Croz. [*Riccia oelandica* C.E.O.Jensen]¹⁵²

23 **R. trabutiana** Steph.

24 **R. warnstorffii** Limpr. ex Warnst. [*Riccia warnstorffii* var. *commutata* (J.B.Jack ex Levier) Damsh., *Riccia warnstorffii* var. *subinermis* Warnst., *Riccia warnstorffii* var. *ciliaris* Warnst.]

section *Pilifer* O.H.Volk

25 **R. boumanii** Dirkse, Losada-Lima & M.Stech¹⁵³

subgenus *Ricciella* (A.Braun) Boulay

section *Ricciella* (A.Braun) Bisch.

26 **R. canaliculata** Hoffm.

27 **R. duplex** Lorb. ex Müll.Frib.

a var. **duplex**¹⁵⁴

28 **R. fluitans** L.

29 **R. frostii** Austin

a var. **frostii**¹⁵⁵

30 **R. huebeneriana** Lindenb.

a subsp. **huebeneriana** [*Riccia huebeneriana* var. *pseudo-frostii* Schiffn.]¹⁵⁶

31 **R. perennis** Steph.

32 **R. rhenana** Lorb. ex Müll.Frib.

a var. **rhenana**

b var. **violacea** M.F.Boiko

section *Spongodes* Nees

33 **R. cavernosa** Hoffm. [*Riccia teneriffae* S.W.Arnell, *Riccia cavernosa* var. *angustior* (Nees) Damsh.]

34 **R. crystallina** L.

120 **Ricciocarpos** Corda

1 **R. natans** (L.) Corda

Targioniaceae Dumort.

121 **Targionia** L.

subgenus *Targionia*

1 **T. hypophylla** L.

a subsp. **hypophylla**¹⁵⁷

2 **T. lorbeeriana** Müll.Frib.

Sphaerocarpaceae Cavers

Riellaceae Engl.

122 **Riella** Mont.

subgenus *Riella*

1 **R. battandieri** Trab.

2 **R. bialata** Trab.¹⁵⁸

3 **R. gallica** Balansa ex Trab.

4 **R. helicophylla** (Bory & Mont.) Mont.

a var. **helicophylla**

b var. **macrocarpa** P.Allorge

5 **R. notarisii** (Mont.) Mont.¹⁵⁹

6 **R. reuteri** Mont.

subgenus *Trabutiella* Porsild

7 **R. affinis** M.Howe & Underw.

8 **R. cossoniana** Trab.

9 **R. echinata** (Müll.Frib.) Segarra, Puche & Sabovlj.¹⁶⁰

10 **R. mediterranea** Segarra, Puche, Sabovlj., M.Infante & Heras¹⁶¹

Sphaerocarpaceae Heeg

123 **Sphaerocarpos** Boehm.

subgenus *Austrosphaerocarpos* R.M.Schust.

1 **S. stipitatus** Bisch. ex Lindenb.

subgenus *Sphaerocarpos*

2 **S. europaeus** Lorb. [*Sphaerocarpos texanus* auct. eur. non Austin]¹⁶²

3 **S. michelii** Bellardi

Mosses

BRYOPHYTA

SPHAGNOPSIDA SCHIMP.

Sphagnales Limpr.

Sphagnaceae Dumort.

1 **Sphagnum** L.¹⁶³

¹⁵¹*Riccia sorocarpa* var. *heegii* was accepted with low confidence in the world checklist of liverworts (Söderström, Hagborg, et al. 2016). In Europe it has been reported from Portugal, Spain, France, Corsica, Sardinia, Austria, Serbia, North Macedonia and Romania as well as from the Canary Islands and Madeira.

¹⁵²The name *Riccia subbifurca* has been used for different plants in Europe, and the application of the name is problematic, as is its relationship with other taxa such as *R. oelandica* C.E.O. Jensen. The species was described by Crozals (1903), based primarily on material from a serpentine site in France. It is discussed by Hugonnot (2018), who suggests that it could be a taxon of localised occurrence on serpentine.

¹⁵³*Riccia boumanii* is newly described from the Canary Islands by Dirkse et al. (2016).

¹⁵⁴*Riccia duplex* var. *megaspora* Na-Thalang occurs in Australia.

¹⁵⁵*Riccia frostii* var. *crystallinoides* Schiffn. occurs in W Asia.

¹⁵⁶*Riccia huebeneriana* subsp. *sullivantii* (Austin) R.M.Schust. occurs in North America.

¹⁵⁷*Targionia hypophylla* subsp. *linealis* W.Frey & Kürschner occurs on the Arabian peninsula.

¹⁵⁸*Riella bialata* was reported new to Europe from Spain by Puche and Segarra-Moragues (2013).

¹⁵⁹*Riella notarisii* is a species complex with poorly defined taxonomic boundaries including the European *Riella battandieri*, *Riella gallica* and *Riella reuteri* as well as *Riella cyrenaica* Maire and *Riella sersuensis* Trab. from North Africa.

¹⁶⁰*Riella echinata* is a new species described from North Africa by Segarra-Moragues et al. (2014) but also occurring in Spain, The Balears and Canary Islands.

¹⁶¹*Riella mediterranea* is a new species described from Spain by Segarra-Moragues et al. (2014) but also occurring in the Balears, Malta and Cyprus as well as in Morocco.

¹⁶²European accessions of '*Sphaerocarpos texanus*' were shown to be clearly distinct from the American *Sphaerocarpos texanus* (Bell et al. 2013) and the latter is thus excluded from Europe. *Sphaerocarpos europaeus* is the oldest name based on a European specimen.

¹⁶³The subgeneric and sectional classification of *Sphagnum* follows Laine et al. (2018).

subgenus *Rigida* (Lindb.) A.Eddy

- 1 ***S. compactum*** Lam. & DC.
- 2 ***S. strictum*** Sull.

subgenus *Sphagnum*

- 3 ***S. affine*** Renault & Cardot [*Sphagnum affine* var. *flagellare* (Schlieph. ex Röhl) L.Söderstr. & Hedenäs, *Sphagnum imbricatum* subsp. *affine* (Renault & Cardot) Flatberg]
- 4 ***S. austinii*** Sull. [*Sphagnum imbricatum* subsp. *austinii* (Sull.) Flatberg]
- 5 ***S. centrale*** C.E.O.Jensen [*Sphagnum palustre* var. *centrale* (C.E.O.Jensen) A.Eddy]
- 6 ***S. divinum*** Flatberg & Hassel [*Sphagnum magellanicum* auct. eur. p.p., non Brid.]
- 7 ***S. medium*** Limpr. [*Sphagnum magellanicum* auct. eur. p.p., non Brid.]¹⁶⁴
- 8 ***S. palustre*** L.
- 9 ***S. papillosum*** Lindb.

subgenus *Acutifolia* (Russow) A.J.Shaw

section *Squarrosa* (Russow) Schimp.

- 10 ***S. mirum*** Flatberg & Thinggaard¹⁶⁵
- 11 ***S. squarrosom*** Crome
- 12 ***S. teres*** (Schimp.) Ångstr.
- 13 ***S. tundrae*** Flatberg

section *Polyclada* (C.E.O.Jensen) Horrell

- 14 ***S. wulfianum*** Girg.

section *Insulosa* Isov.

- 15 ***S. aongstroemii*** C.Hartm.

section *Acutifolia* Wilson

- 16 ***S. angermanicum*** Melin
- 17 ***S. arcticum*** Flatberg & Frisvoll
- 18 ***S. beothuk*** R.E.Andrus¹⁶⁶
- 19 ***S. capillifolium*** (Ehrh.) Hedw. [*Sphagnum capillifolium* subsp. *capillifolium*]
- 20 ***S. concinnum*** (Berggr.) Flatberg [*Sphagnum fimbriatum* subsp. *concinnum* (Berggr.) Flatberg & Frisvoll]¹⁶⁷
- 21 ***S. fimbriatum*** Wilson
- 22 ***S. fuscum*** (Schimp.) H.Klinggr.
- 23 ***S. girgensohnii*** Russow
- 24 ***S. molle*** Sull.
- 25 ***S. nitidulum*** Warnst.¹⁶⁸
- 26 ***S. olafii*** Flatberg

- 27 ***S. quinquefarium*** (Braithw.) Warnst.

- 28 ***S. rubellum*** Wilson [*Sphagnum capillifolium* subsp. *rubellum* (Wilson) M.O.Hill]

- 29 ***S. rubiginosum*** Flatberg

- 30 ***S. russowii*** Warnst.

- 31 ***S. skyense*** Flatberg

- 32 ***S. subfulvum*** Sjors

a subsp. ***purpureum*** Flatberg

b subsp. ***subfulvum***

- 33 ***S. subnitens*** Russow & Warnst.

a subsp. ***ferrugineum*** Flatberg [*Sphagnum subnitens* var. *ferrugineum* (Flatberg) M.O.Hill]

b subsp. ***subnitens***

- 34 ***S. tescorum*** Flatberg¹⁶⁹

- 35 ***S. venustum*** Flatberg¹⁷⁰

- 36 ***S. warnstorffii*** Russow

subgenus *Subsecunda* (Lindb.) A.J.Shaw

- 37 ***S. auriculatum*** Schimp. [*Sphagnum denticulatum* Brid.]

- 38 ***S. contortum*** Schultz

- 39 ***S. inundatum*** Russow

- 40 ***S. platyphyllum*** (Lindb. ex Braithw.) Warnst.

- 41 ***S. pylaesii*** Brid.

- 42 ***S. subsecundum*** Nees

subgenus *Cuspidata* Lindb.

- 43 ***S. angustifolium*** (C.E.O.Jensen ex Russow) C.E.O.Jensen

- 44 ***S. annulatum*** H.Lindb. ex Warnst.

- 45 ***S. balticum*** (Russow) C.E.O.Jensen

- 46 ***S. cuspidatum*** Ehrh. ex Hoffm.

a var. ***cuspidatum***

b var. ***viride*** (Flatberg) Lönnell & Hassel [*Sphagnum viride* Flatberg]¹⁷¹

- 47 ***S. fallax*** (H.Klinggr.) H.Klinggr.¹⁷²

a var. ***brevifolium*** (Lindb. ex Braithw.) Lönnell & Hassel [*Sphagnum brevifolium* (Lindb. ex Braithw.) Röhl]

b var. ***fallax***

c var. ***isoviitae*** (Flatberg) Lönnell & Hassel [*Sphagnum fallax* subsp. *isoviitae* (Flatberg) M.O.Hill, *Sphagnum isoviitae* Flatberg]

¹⁶⁴All European records of *Sphagnum magellanicum* are referable to *Sphagnum divinum* or *Sphagnum medium* (Hassel et al. 2018). *Sphagnum magellanicum* s.str. is confined to southern South America.

¹⁶⁵*Sphagnum mirum* is an Arctic species recently found in Nenets Province, European Russia (Laine et al. 2018).

¹⁶⁶*Sphagnum beothuk* is a North American species (Andrus 2006), recently found to occur in oceanic parts of Europe (Kyrkjeeide et al. 2015).

¹⁶⁷Shaw et al. (2012) showed that *Sphagnum concinnum* and *Sphagnum fimbriatum* are distinct species on both morphological and molecular evidence.

¹⁶⁸*Sphagnum nitidulum* is a doubtful species in need of revision, that grows around fumaroles on the island of Terceira in the Azores. It appears distinctive, but the diagnostic characters might merely be the result of these extreme conditions (Séneca and Söderström 2009).

¹⁶⁹*Sphagnum tescorum* was found in subarctic European Russia in 2016 (Laine et al. 2018).

¹⁷⁰*Sphagnum venustum* was described from eastern Canada (Flatberg 2008). It is ampho-Atlantic, with a single location found recently in Norway (Laine et al. 2018).

¹⁷¹*Sphagnum viride* was treated as a synonym of *Sphagnum cuspidatum* by Hill et al. (2006), then as a variety by Lönnell and Hassel (2018). Hanssen et al. (2000) found that the molecular evidence for differentiation is slight. They further suggested that "the different distribution patterns found in north-western Europe speak in favour of a separation at the subspecies rank, but we consider a formal taxonomic decision premature at this point". Laine et al. (2018) state that "the taxonomic status of this species is still somewhat unclear in relation to *Sphagnum cuspidatum*". We follow Lönnell and Hassel (2018) pending further molecular work.

¹⁷²The new combinations for *Sphagnum brevifolium* and *Sphagnum isoviitae* were proposed by Lönnell and Hassel (2018), although further molecular studies are needed to clarify their status conclusively. It has been observed that the three varieties of *Sphagnum fallax* are morphologically rather stable in each of their preferred habitats, and it may be that epigenetic processes are involved in the expression of *Sphagnum fallax* phenotypes.

- 48 *S. flexuosum* Dozy & Molk.
 49 *S. jensenii* H.Lindb.
 50 *S. lenense* H.Lindb. ex L.I.Savicz
 51 *S. lindbergii* Schimp.
 52 *S. majus* (Russow) C.E.O.Jensen
 a subsp. *majus*
 b subsp. *norvegicum* Flatberg
 53 *S. obtusum* Warnst.
 54 *S. pulchrum* (Lindb. ex Braithw.) Warnst.
 55 *S. recurvum* P.Beauv.¹⁷³
 56 *S. riparium* Ångstr.
 57 *S. tenellum* (Brid.) Pers. ex Brid.
 58 *S. troendelagicum* Flatberg

ANDREAEOPSISIDA J.H.SCHAFFEN.

Andreaeales Limpr.

Andreaeaceae Dumort.

- 2 *Andreaea* Hedw.
 section *Chasmocalyx* Lindb. ex Braithw.
 1 *A. nivalis* Hook.
 section *Nerviae* Cardot ex G.Roth
 2 *A. blyttii* Schimp.
 3 *A. crassinervia* Bruch
 4 *A. frigida* Huebener
 5 *A. heinemannii* Hampe & Müll.Hal.
 a subsp. *crassifolia* (Luisier) Sérgio
 b subsp. *heinemannii*
 6 *A. megistospora* B.M.Murray
 7 *A. rothii* F.Weber & D.Mohr
 a subsp. *falcata* (Schimp.) Lindb.
 b subsp. *rothii*
 section *Andreaea*
 8 *A. alpestris* (Thed.) Schimp. [*Andreaea rupestris*
 var. *alpestris* (Thed.) Sharp]
 9 *A. alpina* Hedw. [*Andreaea hartmanii* Thed.,
Andreaea obovata Thed.]¹⁷⁴
 10 *A. flexuosa* R.Br. bis
 a subsp. *luisieri* Sérgio & Sim-Sim¹⁷⁵
 11 *A. hookeri* Schimp. [*Andreaea alpina* auct.
 mult.]¹⁷⁶
 12 *A. mutabilis* Hook.f. & Wilson
 13 *A. rupestris* Hedw.
 a var. *papillosa* (Lindb.) Podp.
 b var. *rupestris*
 14 *A. sinuosa* B.M.Murray

- Oedipodiales Goffinet & W.R.Buck
 Oedipodiaceae Schimp.
 3 *Oedipodium* Schwägr.
 1 *O. griffithianum* (Dicks.) Schwägr.

TETRAPHIDOPSISIDA GOFFINET & W.R.BUCK

Tetraphidales M.Fleisch.

Tetraphidaceae Schimp.

- 4 *Tetraphis* Hedw.
 1 *T. pellucida* Hedw.
 5 *Tetrodontium* Schwägr.
 1 *T. brownianum* (Dicks.) Schwägr.
 2 *T. ovatum* (Funck) Schwägr.
 3 *T. repandum* (Funck) Schwägr.

POLYTRICHOPSISIDA DOWELD

Polytrichales M.Fleisch.

Polytrichaceae Schwägr.

- 6 *Alophosia* Card.
 1 *A. azorica* (Renauld & Cardot) Cardot
 7 *Atrichum* P.Beauv.
 1 *A. androgynum* (Müll.Hal.) A.Jaeger¹⁷⁷
 2 *A. angustatum* (Brid.) Bruch & Schimp.
 3 *A. crispum* (James) Sull.
 4 *A. flavisetum* Mitt.
 5 *A. tenellum* (Röhl.) Bruch & Schimp.
 6 *A. undulatum* (Hedw.) P.Beauv.
 8 *Oligotrichum* DC.
 1 *O. hercynicum* (Hedw.) Lam. & DC.
 9 *Pogonatum* P.Beauv.
 1 *P. aloides* (Hedw.) P.Beauv.
 2 *P. dentatum* (Menzies ex Brid.) Brid.
 3 *P. nanum* (Hedw.) P.Beauv.
 4 *P. neesii* (Müll.Hal.) Dozy
 5 *P. urnigerum* (Hedw.) P.Beauv.
 10 *Polytrichastrum* G.L.Sm.¹⁷⁸
 1 *P. alpinum* (Hedw.) G.L.Sm. [*Polytrichastrum*
norvegicum (Hedw.) Schljakov, *Polytri-*
chum alpinum Hedw., *Polytrichum*
alpinum var. *arcticum* (Sw. ex Brid.)
 Wahlenb.]
 2 *P. altaicum* Ignatov & G.L.Merr.¹⁷⁹

OEDIPODIOPSISIDA GOFFINET & W.R.BUCK

¹⁷³*Sphagnum recurvum* s.str. is essentially an American taxon recently found in the Azores (Dias et al. 2009). Historically the name was used in a wider sense, encompassing *Sphagnum fallax*, *Sphagnum angustifolium* and *Sphagnum flexuosum*.

¹⁷⁴The confused history of *Andreaea alpina* and *Andreaea hookeri* was elucidated by Price and Ellis (2018), who describe the problems associated with the name 'Andreaea alpina' and the type specimens associated with that name. In consequence, the species previously and widely known as 'Andreaea alpina' must now be called *Andreaea hookeri*, while *Andreaea alpina*, for which a lectotype is designated by Price and Ellis (2018), applies to the species previously known as *Andreaea obovata*.

¹⁷⁵*Andreaea flexuosa* subsp. *luisieri* is a new taxon described from Madeira (Sérgio and Sim-Sim 2012); subsp. *flexuosa* occurs in the Southern Hemisphere.

¹⁷⁶See footnote on *Andreaea alpina*, above. *Andreaea hookeri* is therefore the correct name for the plant generally known previously as *Andreaea alpina*.

¹⁷⁷*Atrichum androgynum* was reported from Macaronesia and Portugal by Sérgio et al. (2010).

¹⁷⁸The composition of *Polytrichastrum* in Russia was clarified by Ivanova et al. (2014).

¹⁷⁹*Polytrichastrum altaicum* was described from the Altai Mountains in Asiatic Russia (Ignatov and Smith Merrill (1995), and later also located in European Russia and Finnish Lapland (Ivanova et al. 2014).

- 3 ***P. fragile*** (Bryhn) Schljakov [*Polytrichastrum alpinum* var. *fragile* (Bryhn) D.G.Long, *Polytrichum fragile* Bryhn]¹⁸⁰
- 4 ***P. septentrionale*** (Brid.) E.I.Ivanova, N.E.Bell & Ignatov [*Polytrichastrum alpinum* var. *septentrionale* (Sw. ex Brid.) G.L.Sm., *Polytrichum septentrionale* Brid.]¹⁸¹
- 5 ***P. sexangulare*** (Brid.) G.L.Sm. [*Polytrichum sexangulare* Hedw.]
- 6 ***P. sphaerothecium*** (Besch.) J.-P.Frahm [*Polytrichastrum sexangulare* var. *vulcanicum* (C.E.O.Jensen) G.L.Merr., *Polytrichum sphaerothecium* (Besch.) Müll.Hal.]
- 11 ***Polytrichum*** Hedw.
- 1 ***P. commune*** Hedw. [*Polytrichum commune* var. *commune*, *Polytrichum commune* var. *uliginosum* Wallr., *Polytrichum uliginosum* (Wallr.) Schriebl]¹⁸²
- 2 ***P. densifolium*** Wilson ex Mitt. [*Polytrichastrum formosum* var. *densifolium* (Wilson ex Mitt.) Z.Iwats. & Nog., *Polytrichum formosum* var. *densifolium* (Wilson ex Mitt.) Osada]¹⁸³
- 3 ***P. formosum*** Hedw. [*Polytrichastrum formosum* (Hedw.) G.L.Sm.]¹⁸⁴
- 4 ***P. hyperboreum*** R.Br.
- 5 ***P. jensenii*** I.Hagen
- 6 ***P. juniperinum*** Hedw.
- 7 ***P. longisetum*** Sw. ex Brid. [*Polytrichastrum longisetum* (Sw. ex Brid.) G.L.Sm.]¹⁸⁵
- 8 ***P. pallidisetum*** Funck [*Polytrichastrum pallidisetum* (Funck) G.L.Sm.]¹⁸⁶
- 9 ***P. perigoniale*** Michx. [*Polytrichum commune* var. *humile* Sw., *Polytrichum commune* var. *perigoniale* (Michx.) Hampe]¹⁸⁷
- 10 ***P. piliferum*** Hedw.
- 11 ***P. strictum*** Menzies ex Brid. [*Polytrichum alpestre* Hoppe]
- 12 ***P. swartzii*** Hartm. [*Polytrichum commune* var. *swartzii* (Hartm.) Nyholm]
- 12 ***Psilopilum*** Brid.
- 1 ***P. cavifolium*** (Wilson) I.Hagen
- 2 ***P. laevigatum*** (Wahlenb.) Lindb.
- BRYOPSIDA** PAX
Buxbaumiales M.Fleisch.
Buxbaumiaceae Schimp.
- 13 ***Buxbaumia*** Hedw.
- 1 ***B. aphylla*** Hedw.
- 2 ***B. viridis*** (Moug. ex Lam. & DC.) Brid. ex Moug. & Nestl.
- Diphysciales M.Fleisch.
Diphysciaceae M.Fleisch.
- 14 ***Diphyscium*** D.Mohr
- 1 ***D. foliosum*** (Hedw.) D.Mohr
- Timmiales Ochyra
Timmiaaceae Schimp.
- 15 ***Timmia*** Hedw.
- section *Timmiaurea* Brassard
- 1 ***T. austriaca*** Hedw.
- section *Timmia*
- 2 ***T. bavarica*** Hessel.
- 3 ***T. megapolitana*** Hedw.
- section *Norvegica* Brassard
- 4 ***T. comata*** Lindb. & Arnell
- 5 ***T. norvegica*** J.E.Zetterst.
- 6 ***T. sibirica*** Lindb. & Arnell
- Encalyptales Dixon
Encalyptaceae Schimp.
- 16 ***Bryobrittonia*** Williams
- 1 ***B. longipes*** (Mitt.) D.G.Horton

¹⁸⁰*Polytrichastrum fragile* was treated at species level by Ivanova et al. (2014).

¹⁸¹*Polytrichastrum septentrionale* was treated at species level by Ivanova et al. (2014).

¹⁸²*Polytrichum commune* var. *uliginosum* Wallr. was raised to specific rank by Schriebl (1991) following his observation of stability in the described diagnostic features in culture. Hill et al. (2006) accepted this view, supported partially by Bijlsma et al. (2000), who found two distinct genetic entities within *Polytrichum commune* s.lat. that they recognised as *Polytrichum uliginosum* and *Polytrichum commune*, with *Polytrichum commune* var. *perigoniale* (Michx.) Hampe treated under *Polytrichum commune*. Subsequent molecular and herbarium studies (Bell et al. 2010; Kariyawasam et al. in prep), have confirmed that these two genetic entities exist, although they show that *Polytrichum uliginosum* should be treated under *Polytrichum commune* while *Polytrichum perigoniale* Michx. is the correct name for the other entity. This is because Bijlsma et al. (2000) based their sampling of *Polytrichum commune* on Schriebl's concept of the species, with specimens corresponding to *Polytrichum commune* var. *perigoniale* (as can be seen in their Figure 1C), while the concept of *Polytrichum uliginosum* they used presumably included plants corresponding morphologically to *Polytrichum commune* var. *commune*. The type of *Polytrichum commune* in the Hedwig-Schwägrichen-herbarium comprises nine stems on a single herbarium sheet. Some of these appear to have been added by Schwägrichen for comparison purposes and are assignable to *Polytrichum juniperinum*, *Polytrichum appressum* (= *Polytrichum subpilosum*) and *Polytrichum perigoniale*. Six others match Hedwig's original description of *Polytrichum commune* and descriptions in the literature he cited, as well as the concept of *Polytrichum commune* var. *commune* traditionally used in Europe. Of these, one will be selected as a lectotype for *Polytrichum commune* (Kariyawasam, in prep). See also footnote to *Polytrichum perigoniale*, below.

¹⁸³Originally described in 1859 from Sikkim in India (Mitten 1859), *Polytrichum densifolium* was overlooked in Europe (and elsewhere), where specimens were assigned to *Polytrichum formosum*. Most Russian specimens previously named *Polytrichum formosum* were transferred to *Polytrichum densifolium* by Ivanova et al. (2015), and it seems likely that many other eastern European occurrences of '*Polytrichum formosum*' may also prove to be *Polytrichum densifolium*.

¹⁸⁴*Polytrichastrum formosum* was returned to *Polytrichum* by Bell and Hyvönen (2010), a treatment followed by Ivanova et al. (2014) and others.

¹⁸⁵*Polytrichastrum longisetum* was returned to *Polytrichum* by Bell and Hyvönen (2010), a treatment followed by Ivanova et al. (2014) and others.

¹⁸⁶*Polytrichastrum pallidisetum* was returned to *Polytrichum* by Bell and Hyvönen (2010), a treatment followed by Ivanova et al. (2014) and others.

¹⁸⁷Molecular evidence (e.g. Bell et al. 2010; Kariyawasam et al. in prep.) clearly shows *Polytrichum perigoniale* to be more closely related to a number of non-European species than to *Polytrichum commune* s.str., thus necessitating its recognition as a full species rather than a variety of *Polytrichum commune*. The type (G) has been examined by Kariyawasam.

- 17 **Encalypta** Hedw.
section *Streptothecca* (Kindb.) Broth.
1 ***E. procera*** Bruch
2 ***E. streptocarpa*** Hedw.
section *Pyromitrium* Wallr. ex Hampe
3 ***E. alpina*** Sm.
4 ***E. mutica*** l.Hagen
section *Rhabdothecca* Müll.Hal.
5 ***E. pilifera*** Funck [*Encalypta obovatifolia* Nyholm, *Encalypta intermedia* Jur.]¹⁸⁸
6 ***E. rhaptocarpa*** Schwägr.
7 ***E. spathulata*** Müll.Hal. [*Encalypta rhaptocarpa* var. *spathulata* (Müll.Hal.) Husn.]
8 ***E. trachymitria*** Ripart [*Encalypta rhaptocarpa* var. *leptodon* Lindb., *Encalypta rhaptocarpa* var. *trachymitria* (Ripart) Wijk & Margad.]¹⁸⁹
9 ***E. vulgaris*** Hedw.
section *Megasporae* D.G.Horton
10 ***E. longicolla*** Bruch
section *Encalypta*
11 ***E. affinis*** R.Hedw.
a subsp. ***affinis***
b subsp. ***macounii*** (Austin) D.G.Horton
12 ***E. brevicolla*** (Bruch & Schimp.) Ångstr.
13 ***E. brevipes*** Schljakov
14 ***E. ciliata*** Hedw.
15 ***E. microstoma*** Bals.-Criv. & De Not.

Funariales M.Fleisch.
Funariaceae Schwägr.
Pyramiduloideae O.Werner, Ros & Goffinet¹⁹⁰
18 **Goniomitrium** Hook.f. & Wilson
1 ***G. seroi*** Casas

19 **Pyramidula** Brid.
1 ***P. tetragona*** (Brid.) Brid.

Funarioideae Broth.
20 **Entosthodon** Schwägr.
subgenus *Entosthodon*
1 ***E. abramovae*** Fedosov & Ignatova¹⁹¹
2 ***E. attenuatus*** (Dicks.) Bryhn [*Entosthodon templetonii* (Sm.) Schwägr.]
3 ***E. commutatus*** Durieu & Mont. [*Entosthodon krausei* Besch.]¹⁹²
4 ***E. dagestanicus*** Fedosov & Ignatova¹⁹³
5 ***E. duriaei*** Mont.¹⁹⁴
6 ***E. handelii*** (Schiffn.) Laz. [*Funaria handelii* Schiffn.]¹⁹⁵
7 ***E. hungaricus*** (Boros) Loeske
8 ***E. kroonkurk*** Dirkse & Brugués¹⁹⁶
9 ***E. obtusus*** (Hedw.) Lindb.
10 ***E. stenophyllus*** Fedosov & Ignatova¹⁹⁷
subgenus *Plagiodus* (Mitt.) Fife
11 ***E. convexus*** (Spruce) Brugués
12 ***E. muhlenbergii*** (Turner) Fife
13 ***E. pulchellus*** (H.Philib.) Brugués
14 ***E. schimperii*** Brugués
subgenus *Murcia* Fife
15 ***E. fascicularis*** (Hedw.) Müll.Hal.
16 ***E. mouretii*** (Corb.) Jelenc

21 **Funaria** Schwägr.¹⁹⁸
1 ***F. aequidens*** Lindb. ex Broth.
2 ***F. arctica*** (Berggr.) Kindb.
3 ***F. hygrometrica*** Hedw.
4 ***F. microstoma*** Bruch ex Schimp.

22 **Funariella** Sérgio
1 ***F. curviseta*** (Schwägr.) Sérgio

23 **Physcomitrium** (Brid.) Brid. [*Aphanorrhagma* Sull., *Ephemerella* Mull.Hal., *Physcomitrella* Bruch & Schimp., *Physcomitridium* G.Roth]¹⁹⁹
1 ***P. arenicola*** Laz.
2 ***P. euryostomum*** Sendtn.
a subsp. ***acuminatum*** (Bruch & Schimp.) Giacom.
b subsp. ***euryostomum***
3 ***P. patens*** (Hedw.) Mitt. [*Aphanorrhagma patens* (Hedw.) Lindb., *Physcomitrella patens* (Hedw.) Bruch & Schimp.]
4 ***P. pyriforme*** (Hedw.) Bruch & Schimp.

¹⁸⁸Molecular studies by Fedosov (2012) showed that *Encalypta pilifera* is one of the most isolated taxa in sect. *Rhabdothecca*, thus justifying specific status, and *Encalypta pilifera* is the earliest published name.

¹⁸⁹Fedosov (2012) proposed the recognition of *Encalypta trachymitria* at specific level on the basis of molecular studies on Russian material.

¹⁹⁰The position of *Goniomitrium* and *Pyramidula* in a subfamily within the Funariaceae was confirmed using molecular data by Werner, Ros, et al. (2007).

¹⁹¹*Entosthodon abramovae* was described from the Caucasus by Fedosov, Ignatova, et al. (2010).

¹⁹²*Entosthodon krausei* was synonymised with *Entosthodon commutatus* by Brugués and Sérgio (2010).

¹⁹³*Entosthodon dagestanicus* was described from the Caucasus by Fedosov, Ignatova, et al. (2010).

¹⁹⁴According to Ros, Mazimpaka, et al. (2013), all occurrences of the epithet '*durieui*', and combinations based on it, need to be changed to '*duriaei*', because that is the original spelling.

¹⁹⁵*Entosthodon handelii* was reported from the Caucasus new to Europe by Fedosov, Ignatova, et al. (2010).

¹⁹⁶*Entosthodon kroonkurk* was described from Spain and the Canary Islands by Dirkse and Brugués (2010).

¹⁹⁷*Entosthodon stenophyllus* was described from the Caucasus by Fedosov, Ignatova, et al. (2010).

¹⁹⁸*Funaria anomala* Jur. is a little-known taxon described from Cyprus by Juratzka in Unger and Kotschy (1865). It has also been reported from Egypt and SW Asia. Its status is doubtful. Loeske (1929), who examined the type collection, suggested that it could be a hybrid between female *Entosthodon templetonii* (*Entosthodon attenuatus*) and male *Funaria dentata* (*Funaria muhlenbergii* s.lat.). There have been no subsequent reports from Cyprus, and the taxon is therefore not included in the checklist.

¹⁹⁹Medina et al. (2019) concluded that there are no morphological or molecular grounds for maintaining *Physcomitrella* and *Physcomitridium* as separate genera from *Physcomitrium*.

- 5 ***P. readeri*** Müll.Hal. [*Ephemerella readeri* Müll.Hal., *Physcomitrella readeri* (Müll.Hal.) I.G.Stone & G.A.M.Scott, *Physcomitridium readeri* (Müll.Hal.) G.Roth]²⁰⁰
- 6 ***P. sphaericum*** (C.F.Ludw. ex Schkuhr) Brid.

Disceliaceae Schimp.

- 24 ***Discelium*** Brid.
1 ***D. nudum*** (Dicks.) Brid.

Gigaspermales Goffinet, Wickett, O.Werner, Ros, A.J.Shaw & C.J.Cox

Gigaspermaceae Lindb.

- 25 ***Gigaspermum*** Lindb.
1 ***G. mouretii*** Corb.

26 ***Oedipodiella*** Dixon

- 1 ***O. australis*** (Wager & Dixon) Dixon

Catospiales Ignatov & Ignatova

Catosciaceae Broth.

- 27 ***Catoscopium*** Brid.
1 ***C. nigratum*** (Hedw.) Brid.

Dicranales H.Philib ex M.Fleisch.²⁰¹

Timmiellaceae Y.Inoue & H.Tsubota²⁰²

- 28 ***Timmiella*** (De Not.) Limpr.
1 ***T. anomala*** (Bruch & Schimp.) Limpr.
2 ***T. barbuloides*** (Brid.) Mönk.
3 ***T. flexisetia*** (Bruch) Limpr.

Distichiaceae Schimp.²⁰³

- 29 ***Distichium*** Bruch & Schimp.
1 ***D. capillaceum*** (Hedw.) Bruch & Schimp.
2 ***D. hagenii*** Ryan ex H.Philib.
3 ***D. inclinatum*** (Hedw.) Bruch & Schimp.

Hymenolomataceae Ignatov & Fedosov²⁰⁴

- 30 ***Hymenoloma*** Dusén
1 ***H. compactum*** (Schleich. ex Schwägr.) Ochyra [*Dicranoweisia compacta* (Schleich. ex

Schwägr.) Schimp., *Dicranoweisia crispula* var. *compacta* (Schwägr.) Lindb.]

- 2 ***H. crispulum*** (Hedw.) Ochyra [*Dicranoweisia crispula* (Hedw.) Milde]
- 3 ***H. mulahaceni*** (Höhn.) Ochyra [*Dicranoweisia crispula* var. *intermedia* (J.J.Amann) Podp., *Dicranoweisia intermedia* J.J.Amann, *Hymenoloma intermedium* (J.J.Amann) Ochyra]²⁰⁵

Flexitrichaceae Ignatov & Fedosov²⁰⁶

- 31 ***Flexitrichum*** Ignatov & Fedosov
1 ***F. flexicaule*** (Schwägr.) Ignatov & Fedosov [*Ditrichum flexicaule* (Schwägr.) Hampe]
2 ***F. gracile*** (Mitt.) Ignatov & Fedosov [*Ditrichum crispatissimum* (Müll.Hal.) Paris, *Ditrichum gracile* (Mitt.) Kuntze]

Bryoxiphiaceae Besch.

- 32 ***Bryoxiphium*** Brid.
1 ***B. madeirense*** Á.Löve & D.Löve
2 ***B. norvegicum*** (Brid.) Mitt.

Archidiaceae Schimp.

- 33 ***Archidium*** Brid.
1 ***A. alternifolium*** (Hedw.) Mitt.

Micromitriaceae Smyth ex Goffinet & Budke²⁰⁷

- 34 ***Micromitrium*** Austin
1 ***M. tenerum*** (Bruch & Schimp.) Crosby

Leucobryaceae Schimp.

- 35 ***Atractylocarpus*** Mitt.
1 ***A. alpinus*** (Schimp. ex Milde) Lindb.
2 ***A. subporodictyon*** (Broth.) Bonfim Santos & M.Stech [*Campylopus subporodictyon* (Broth.) B.H.Allen & Ireland, *Dicranodontium subporodictyon* Broth., *Dicranum subporodictyon* (Broth.) C.Gao & T.Gao]²⁰⁸

²⁰⁰*Physcomitrium readeri* was reported, as *Ephemerella readeri*, new to Europe by Hooper et al. (2010). It was moved to *Physcomitridium* by Goffinet and Buck (2011), and is now in *Physcomitrium* (Medina et al. 2019).

²⁰¹The order Dicranales is here pragmatically broadened to include the basal lineages of haplolepidous mosses ('Protohaplolepidae' of Hedderson et al. 2004). This decision necessarily results in merging the order Pottiales with Dicranales, although we retain Grimmiales as one of the crown groups of Dicranidae.

²⁰²Inoue and Tsubota (2014, 2016) showed through molecular studies that *Timmiella* should be accommodated within a new family, Timmiellaceae, along with the tropical genus *Luisierella*. The evidence also suggests that the family does not belong to the Pottiales but rather to the basal haplolepidous mosses ('Protohaplolepidae').

²⁰³Fedosov, Fedorova, Fedosov, et al. (2016) provided molecular evidence that *Distichium* cannot be placed in the Ditrichaceae, They therefore resurrected the family Distichiaceae, which had not been used for more than a century. Further molecular evidence suggests that the Distichiaceae do not belong to the Dicranales but rather to the basal haplolepidous mosses ('Protohaplolepidae').

²⁰⁴Fedosov, Fedorova, Troitsky, et al. (2016) place the genus *Hymenoloma* in a family of its own, the Hymenolomataceae. The relationships of the species in *Hymenoloma* and *Dicranoweisia* were elucidated by Werner, Rams, et al. (2013).

²⁰⁵*Hymenoloma mulahaceni* is a Holarctic species whose taxonomy and distribution is described by Werner, Rams, et al. (2013).

²⁰⁶The family Flexitrichaceae was described as new on the basis of molecular evidence (Fedosov, Fedorova, Fedosov, et al. 2016), which places it apart from Ditrichaceae. As with the Distichiaceae, molecular evidence suggests that the Flexitrichaceae do not belong to the Dicranales but rather to the basal haplolepidous mosses ('Protohaplolepidae').

²⁰⁷The Micromitriaceae is a new family described by Goffinet, Budke, et al. (2011) on molecular grounds. Its affinities with Dicranidae have not yet been clarified but it might be close to taxa basal to Leucobryaceae (cf. Bonfim Santos and Stech 2017; Stech et al. 2012).

²⁰⁸Recent molecular work shows *Campylopus subporodictyon* to be best placed in *Atractylocarpus* (Bonfim Santos and Stech 2017).

- 36 **Campylopus** Brid.
 1 **C. atrovirens** De Not. [*Campylopus atrovirens* var. *falcatus* Braithw.]
 2 **C. brevipilus** Bruch & Schimp.
 3 **C. cygneus** (Hedw.) Brid.
 4 **C. flaccidus** Renauld & Cardot
 5 **C. flexuosus** (Hedw.) Brid.
 6 **C. fragilis** (Brid.) Bruch & Schimp.
 7 **C. gracilis** (Mitt.) A.Jaeger [*Campylopus schwarzii* Schimp.]
 8 **C. incrassatus** Müll.Hal.
 9 **C. introflexus** (Hedw.) Brid.
 10 **C. oerstedianus** (Müll.Hal.) Mitt.
 11 **C. pilifer** Brid.
 12 **C. pyriformis** (Schultz) Brid. [*Campylopus pyriformis* var. *azoricus* (Mitt.) M.F.V.Corley]
 13 **C. schimperi** Milde [*Campylopus subulatus* var. *schimperi* (Milde) Husn.]²⁰⁹
 14 **C. setifolius** Wilson
 15 **C. shawii** Wilson
 16 **C. subulatus** Schimp. ex Milde²¹⁰
- 37 **Dicranodontium** Bruch & Schimp.
 1 **D. asperulum** (Mitt.) Broth.
 2 **D. denudatum** (Brid.) E.Britton [*Dicranodontium denudatum* var. *alpinum* (Schimp.) I.Hagen]
 3 **D. uncinatum** (Harv.) A.Jaeger
- 38 **Leucobryum** Hampe
 1 **L. albidum** (P.Beauv.) Lindb.²¹¹
 2 **L. glaucum** (Hedw.) Ångstr.
 3 **L. juniperoideum** (Brid.) Müll.Hal.
- Amphidiaceae M.Stech
 39 **Amphidium** Schimp.
 1 **A. lapponicum** (Hedw.) Schimp.
 2 **A. mougeotii** (Schimp.) Schimp.
 3 **A. curvipes** (Müll.Hal.) Broth. [*Amphidium tortuosum* auct. eur.]
- Aongstroemiaceae De Not.
 40 **Aongstroemia** Schimp.
 1 **A. longipes** (Sommerf.) Bruch & Schimp.
- 41 **Dichodontium** Schimp.
 1 **D. flavescens** (Dicks.) Lindb.
 2 **D. pellucidum** (Hedw.) Schimp.
- 42 **Diobelonella** Ochyra²¹²
 1 **D. palustris** (Dicks.) Ochyra [*Anisothecium palustre* (Dicks.) I.Hagen, *Dichodontium palustre* (Dicks.) M.Stech, *Dicranella palustris* (Dicks.) Crundw.]
- Dicranellaceae M.Stech
 43 **Dicranella** (Müll.Hal.) Schimp.
 1 **D. campylophylla** (Taylor) A.Jaeger
 2 **D. cerviculata** (Hedw.) Schimp.
 3 **D. crispa** (Hedw.) Schimp.
 4 **D. grevilleana** (Brid.) Schimp.
 5 **D. heteromalla** (Hedw.) Schimp.
 6 **D. howei** Renauld & Cardot
 7 **D. humilis** R.Ruthe
 8 **D. rufescens** (Dicks.) Schimp.
 9 **D. schreberiana** (Hedw.) Dixon
 10 **D. staphylina** H.Whitehouse
 11 **D. subulata** (Hedw.) Schimp.
 12 **D. varia** (Hedw.) Schimp.
- 44 **Microcampylopus** (Müll.Hal.) M.Fleisch.
 1 **M. laevigatus** (Thér.) Giese & J.-P.Frahm
- Fissidentaceae Schimp.
 45 **Fissidens** Hedw.
 subgenus *Pachyfissidens* (Müll.Hal.) L.Söderstr. & A.Hagborg
 section *Pachyfissidens* Müll.Hal.
 1 **F. adianthoides** Hedw.
 2 **F. azoricus** (P.de la Varde) Bizot
 3 **F. dubius** P.Beauv. [*Fissidens cristatus* Wilson ex Mitt.]
 a var. **dubius**
 b var. **mucronatus** (Limpr.) Kartt., Hedenäs & L.Söderstr.²¹³
 4 **F. grandifrons** Brid.
 5 **F. osmundoides** Hedw.
 6 **F. polyphyllus** Wilson ex Bruch & Schimp.
 7 **F. serrulatus** Brid. [*Fissidens luisieri* P.de la Varde]²¹⁴
 8 **F. taxifolius** Hedw. [*Fissidens taxifolius* subsp. *pallidicaulis* (Mitt.) Mönk]²¹⁵

²⁰⁹*Campylopus schimperi* is often treated as a variety of *Campylopus subulatus* (e.g. Meinunger and Schröder 2007) but we follow Hill et al. (2006) in retaining it at species level. Although they are difficult to separate in some parts of their range, *Campylopus schimperi* is usually very distinct in size, habit and habitat.

²¹⁰Because of inconsistencies regarding their respective habitats, there is some doubt that the taxon commonly identified as *Campylopus subulatus* is the same as the type specimen; this therefore requires clarification.

²¹¹Vanderpoorten et al. (2003) suggested that *Leucobryum albidum* is synonymous with *Leucobryum juniperoideum*. However, the two species were included by Hill et al. (2006), and in the absence of further work are retained here. The identification of the three European species is discussed by Simmel and Poschlod (2017).

²¹²Ochyra, Żarnowiec, et al. (2003) proposed a new genus, *Diobelonella*, for *Dichodontium palustre*, which, while it has molecular affinities with *Dichodontium* (Stech 1999), does not sit comfortably there. It is placed alongside *Dichodontium* in the Aongstroemiaceae by Frey and Stech (2009).

²¹³*Fissidens dubius* var. *mucronatus* is morphologically characterised by the mucronate apex of the upper leaves. It also has a different ecology. Recent Dutch research also shows differences in DNA sequences between var. *mucronatus* and the type (H. N. Siebel and M. Stech, pers. comm. 2019).

²¹⁴*Fissidens luisieri* may be distinct but further work is necessary. For the present we follow Werner, Patiño, et al. (2009) in treating it as a synonym of *Fissidens serrulatus*.

²¹⁵*Fissidens taxifolius* subsp. *pallidicaulis* intergrades completely with the type subspecies and is therefore treated as a synonym.

section *Amblyothallia* (Müll.Hal.) Pursell & Brugg.-Nann.

9 ***F. asplenioides*** Hedw.

subgenus *Octodiceras* (Brid.) Broth. [*Octodiceras* Brid.]

10 ***F. fontanus*** (Bach.Pyl.) Steud. [*Octodiceras fontanum* (Bach.Pyl.) Lindb.]

subgenus *Fissidens*

11 ***F. arcticus*** Bryhn

12 ***F. arnoldii*** R.Ruthe

13 ***F. bryoides*** Hedw.

a var. ***bryoides***

b var. ***caespitans*** Schimp. [*Fissidens bryoides* var. *curnovii* (Mitt.) J.J.Amann, *Fissidens curnovii* Mitt.]

14 ***F. coacervatus*** Brugg.-Nann.

15 ***F. crassipes*** Wilson ex Bruch & Schimp.

a subsp. ***crassipes***

b subsp. ***warnstorffii*** (M.Fleisch.) Brugg.-Nann.

16 ***F. crispus*** Mont. [*Fissidens herzogii* R.Ruthe ex Herzog, *Fissidens limbatus* Sull., *Fissidens minutulus* Sull.]

17 ***F. curvatus*** Hornsch. [*Fissidens algarvicus* Solms]

18 ***F. gracilifolius*** Brugg.-Nann. & Nyholm [*Fissidens viridulus* var. *tenuifolius* (Boulay) A.J.E.Sm.]

19 ***F. gymnandrus*** Buse

20 ***F. incurvus*** Starke ex Röhl. [*Fissidens viridulus* var. *incurvus* (Starke ex Röhl.) Waldh.]²¹⁶

21 ***F. jansenii*** Sérgio & Pursell

22 ***F. monguillonii*** Thér.

23 ***F. ovatifolius*** R.Ruthe

24 ***F. pusillus*** (Wilson) Milde [*Fissidens viridulus* var. *pusillus* Wilson]

25 ***F. rivularis*** (Spruce) Schimp.

26 ***F. rufulus*** Bruch & Schimp.

27 ***F. sublimbatus*** Grout

28 ***F. sublineaefolius*** (P.de la Varde) Brugg.-Nann.

29 ***F. viridulus*** (Sw.) Wahlenb. [*Fissidens bambergi* Schimp. ex Milde, *Fissidens exiguus* auct. eur., *Fissidens viridulus* var. *bambergi* (Schimp. ex Milde) Waldh.]²¹⁷

subgenus *Aloma* Kindb.

30 ***F. celticus*** Paton

31 ***F. exilis*** Hedw.

32 ***F. microstictus*** Dixon & Luisier

33 ***F. nobreganus*** Dixon & Luisier

34 ***F. serratus*** Müll.Hal.

Dicranaceae Schimp.

Dicranoideae Lindb.

46 ***Dicranoloma*** (Renauld) Renauld

1 ***D. menziesii*** (Taylor) Broth. ex Renauld²¹⁸

47 ***Dicranum*** Hedw.²¹⁹

section *Dicranum*

1 ***D. bonjeanii*** De Not. [*Dicranum undulatum* Turner, *nom. illeg.*]

2 ***D. crassifolium*** Sérgio, Ochyra & Séneca

3 ***D. leioneuron*** Kindb.

4 ***D. majus*** Sm.

5 ***D. polysetum*** Sw. ex anon. [*Dicranum rugosum* (Hoffm. ex Funck) Brid., *Dicranum undulatum* Ehrh. ex F.Weber & D.Mohr, *nom. illeg.*]

6 ***D. scoparium*** Hedw.

7 ***D. transsylvanicum*** Lüth

section *Spuria* Bruch & Schimp.

8 ***D. acutifolium*** (Lindb. & Arnell) C.E.O.Jensen [*Dicranum bergeri* var. *acutifolium* Lindb. & Arnell, *Dicranum muehlenbeckii* var. *acutifolium* (Lindb. & Arnell) Nyholm]

9 ***D. bardunovii*** Tubanova & Ignatova²²⁰

10 ***D. brevifolium*** (Lindb.) Lindb. [*Dicranum muehlenbeckii* var. *brevifolium* Lindb., *Dicranum muehlenbeckii* var. *cirrhatum* (Schimp.) Lindb.]

11 ***D. dispersum*** Engelmark

12 ***D. drummondii*** Müll.Hal.

13 ***D. septentrionale*** Tubanova & Ignatova²²¹

14 ***D. spurium*** Hedw.

15 ***D. undulatum*** Schrad. ex Brid. [*Dicranum bergeri* Blandow]

section *Fuscescentiformia* (Kindb.) Ochyra

16 ***D. flexicaule*** Brid. [*Dicranum congestum* Brid., *Dicranum fuscescens* var. *congestum* (Brid.) Husn., *Dicranum fuscescens* var. *flexicaule* (Brid.) Wilson]

17 ***D. fuscescens*** Sm.

section *Convolutifolia* (Kindb.) Ochyra

18 ***D. angustum*** Lindb.

19 ***D. muehlenbeckii*** Bruch & Schimp.

²¹⁶Although morphological differences with *Fissidens viridulus* are small, recent molecular studies show that the closest relative of *Fissidens incurvus* is not *Fissidens viridulus* (H. N. Siebel and M. Stech, pers. comm. 2019). It is therefore reinstated as a species.

²¹⁷The status of *Fissidens bambergi* and *Fissidens exiguus*, treated here as synonyms of *Fissidens viridulus*, remains problematic. In a recent DNA study, it appeared that an incompletely limbate species of the *Fissidens bryoides* complex from the Netherlands was different from *Fissidens viridulus* (H. N. Siebel and M. Stech, pers. comm. 2019). However, several incompletely limbate species in this complex have been described. A morphological and molecular revision of these species is needed.

²¹⁸*Dicranoloma menziesii*, a native of Australia and New Zealand, has been recorded on tree ferns in southern Ireland (Holyoak and Lockhart 2009).

²¹⁹Although this sectional classification of *Dicranum* is retained for now, the findings of Lang et al. (2015) suggest that it will have to be revised.

²²⁰*Dicranum bardunovii* was described from Siberia (Tubanova and Ignatova 2011) but later found in the European part of the Urals (E. Ignatova pers. comm. 2018).

²²¹*Dicranum septentrionale* was described from Russia by Tubanova et al. (2010). It has since been found in Scandinavia and Austria.

- 20 ***D. schljakovii*** Ignatova & Tubanova²²²
 21 ***D. spadiceum*** J.E.Zetterst.
 section *Elongata* I.Hagen
 22 ***D. elongatum*** Schleich. ex Schwäger.
 23 ***D. fragilifolium*** Lindb.
 24 ***D. groenlandicum*** Brid.
 25 ***D. laevidens*** R.S.Williams
 section *Crassinervia* G.Roth
 26 ***D. fulvum*** Hook. [*Orthodicranum fulvum*
 (Hook.) G.Roth ex Casares-Gil]
 27 ***D. scottianum*** Turner [*Dicranum canariense*
 Hampe ex Müll.Hal., *Orthodicranum*
scottianum (Turner) G.Roth ex
 Casares-Gil]²²³
 28 ***D. viride*** (Sull. & Lesq.) Lindb.
 section *Montana* Hartm.
 29 ***D. flagellare*** Hedw. [*Orthodicranum flagellare*
 (Hedw.) Loeske]
 30 ***D. montanum*** Hedw. [*Orthodicranum monta-*
num (Hedw.) Loeske]
 31 ***D. tauricum*** Sapjegin [*Dicranum strictum*
 Schleich. ex D.Mohr, *hom. illeg.*, *Orthodi-*
cranum tauricum (Sapjegin) Smirnova]
- 48 ***Paraleucobryum*** (Limpr.) Loeske
 1 ***P. enerve*** (Thed.) Loeske
 2 ***P. longifolium*** (Hedw.) Loeske
 3 ***P. sauteri*** (Bruch & Schimp.) Loeske [*Paraleuco-*
bryum longifolium var. *sauteri* (Bruch &
 Schimp.) C.E.O.Jensen]
- Calymperaceae Kindb.
 49 ***Calymperes*** Michx.
 1 ***C. erosum*** Müll.Hal.
- Rhabdoweisiaceae Limpr. [Oncophoraceae M.Stech]
 50 ***Arctoa*** Schimp.
 1 ***A. anderssonii*** Wich.
 2 ***A. fulvella*** (Dicks.) Bruch & Schimp.
 3 ***A. hyperborea*** (Gunnerus ex Dicks.) Bruch &
 Schimp.
- 51 ***Cnestrum*** I.Hagen
 1 ***C. alpestre*** (Wahlenb. ex Huebener) Nyholm ex
 Mogensen
 2 ***C. glaucescens*** (Lindb. & Arnell) Holmen ex
 Mogensen & Steere
 3 ***C. schisti*** (F.Weber & D.Mohr) I.Hagen
- 52 ***Cynodontium*** Bruch & Schimp.
 1 ***C. asperifolium*** (Lindb. ex Arnell) Paris
 2 ***C. bruntonii*** (Sm.) Bruch & Schimp.
 3 ***C. fallax*** Limpr.
 4 ***C. gracilescens*** (F.Weber & D.Mohr) Schimp.
 5 ***C. jeneri*** (Schimp.) Stirt.
 6 ***C. polycarpon*** (Hedw.) Schimp.
 7 ***C. strumiferum*** (Hedw.) Lindb.
 8 ***C. suecicum*** (Arnell & C.E.O.Jensen) I.Hagen
 9 ***C. tenellum*** (Schimp.) Limpr.
- 53 ***Dicranoweisia*** Milde
 1 ***D. cirrata*** (Hedw.) Lindb.
- 54 ***Glyphomitrium*** Brid.
 1 ***G. daviesii*** (Dicks.) Brid.
- 55 ***Kiaeria*** I.Hagen
 1 ***K. blyttii*** (Bruch & Schimp.) Broth.
 2 ***K. falcata*** (Hedw.) I.Hagen
 3 ***K. glacialis*** (Berggr.) I.Hagen
 4 ***K. riparia*** (H.Lindb.) M.F.V.Corley [*Dicranella*
riparia (H.Lindb.) Mårtensson &
 Nyholm, *Oncophorus riparius* H.Lindb.]
 5 ***K. starkei*** (F.Weber & D.Mohr) I.Hagen
- 56 ***Oncophorus*** (Brid.) Brid.
 1 ***O. demetrii*** (Renauld & Cardot) Hedenäs²²⁴
 2 ***O. dendrophilus*** Hedd. & Blockeel²²⁵
 3 ***O. elongatus*** (I.Hagen) Hedenäs²²⁶
 4 ***O. integerrimus*** Hedenäs [*Oncophorus virens*
 var. *elongatus* Limpr.]²²⁷
 5 ***O. virens*** (Hedw.) Brid.
 6 ***O. wahlenbergii*** Brid. [*Oncophorus compactus*
 (Bruch & Schimp.) Kindb., *Oncophorus*
wahlenbergii var. *compactus* (Bruch &
 Schimp.) Braithw.]²²⁸
- 57 ***Oreas*** Brid.
 1 ***O. martiana*** (Hoppe & Hornsch.) Brid.
- 58 ***Oreoweisia*** (Bruch & Schimp.) De Not.
 1 ***O. torquescens*** (Hornsch. ex Brid.) Wijk &
 Margad.
- 59 ***Rhabdoweisia*** Bruch & Schimp.
 1 ***R. crenulata*** (Mitt.) H.Jameson
 2 ***R. crispata*** (Dicks.) Lindb.
 3 ***R. fugax*** (Hedw.) Bruch & Schimp.

²²²*Dicranum schljakovii* was described from Russia by Ignatova, Tubanova, et al. (2015).

²²³Price et al. (2019) clarified the correct authorship of *Dicranum scottianum*. Recent molecular work suggests that *Dicranum canariense* should be considered a synonym of *Dicranum scottianum* (A. Vanderpoorten pers. comm. 2017). Morphological studies also support this conclusion (Price et al. 2019).

²²⁴*Oncophorus demetrii*, a Scandinavian taxon, was recognised as a species by Hedenäs (2018), and has also been found in Arctic Russia.

²²⁵*Oncophorus dendrophilus* was described by Hedderson and Blockeel (2006) and is so far known only from Crete and Cyprus.

²²⁶The recognition of *Oncophorus elongatus* at species level was confirmed during a revision of Scandinavian *Oncophorus* by Hedenäs (2017a). It has also been found in Russia and France.

²²⁷*Oncophorus integerrimus* was recognised at species level by Hedenäs (2017a).

²²⁸*Oncophorus compactus* was synonymised with *Oncophorus wahlenbergii* by Hedenäs (2018).

Schistostegaceae Schimp.

- 60 **Schistostega** D.Mohr
1 **S. pennata** (Hedw.) F.Weber & D.Mohr

Bruchiaceae Schimp.

- 61 **Bruchia** Schwägr.
1 **B. flexuosa** (Schwägr.) Müll.Hal.
2 **B. vogesiaca** Nestl. ex Schwägr.
- 62 **Trematodon** Michx.
1 **T. ambiguus** (Hedw.) Hornsch.
2 **T. brevicollis** Hornsch.
3 **T. laetevirens** Hakelner & J.-P.Frahm
4 **T. longicollis** Michx.
5 **T. perssoniorum** P.Allorge & Thér. ex V.Allorge

Ditrichaceae Limpr.

- 63 **Ceratodon** Brid.
1 **C. amazonum** Nieto-Lugilde, O.Werner, S.F.McDaniel & Ros²²⁹
2 **C. conicus** (Hampe) Lindb. [*Ceratodon purpureus* var. *conicus* (Hampe) Husn.]²³⁰
3 **C. purpureus** (Hedw.) Brid.
a subsp. **purpureus**
b subsp. **stenocarpus** (Bruch. & Schimp. ex Müll.Hal.) Dixon²³¹
- 64 **Cheilothela** Broth.
1 **C. chloropus** (Brid.) Broth.
- 65 **Cleistocarpidium** Ochyra & Bedn.-Ochyra
1 **C. palustre** (Bruch & Schimp.) Ochyra & Bedn.-Ochyra [*Pleuridium palustre* (Bruch & Schimp.) Bruch & Schimp.]
- 66 **Ditrichum** Timm ex Hampe
1 **D. cornubicum** Paton
2 **D. heteromallum** (Hedw.) E.Britton
3 **D. lineare** (Sw.) Lindb.
4 **D. pallidum** (Hedw.) Hampe
5 **D. plumbicola** Crundw.
6 **D. punctulatum** Mitt.
7 **D. pusillum** (Hedw.) Hampe
8 **D. subulatum** Hampe

- 9 **D. zonatum** (Brid.) Kindb. [*Ditrichum zonatum* var. *scabrifolium* Dixon]

- 67 **Pleuridium** Rabenh.
1 **P. acuminatum** Lindb.
2 **P. subulatum** (Hedw.) Rabenh.
- 68 **Pseudephemerum** (Lindb.) I.Hagen
1 **P. nitidum** (Hedw.) Loeske
- 69 **Rhamphidium** Mitt.
1 **R. purpuratum** Mitt.
- 70 **Trichodon** Schimp.
1 **T. cylindricus** (Hedw.) Schimp. [*Ditrichum cylindricum* (Hedw.) Grout]

Pottiaceae Schimp [Barbuloideae (Herzog) Hilp.]²³²

Merceoideae Broth.

- 71 **Scopelophila** (Mitt.) Lindb.
1 **S. cataractae** (Mitt.) Broth.
2 **S. ligulata** (Spruce) Spruce

Pottioideae Broth.

- 72 **Acaulon** Müll.Hal.
1 **A. casasianum** Brugués & H.A.Crum
2 **A. dertosense** Casas, Sérgio, Cros & Brugués
3 **A. fontiquerianum** Casas & Sérgio
4 **A. mediterraneum** Limpr.
5 **A. muticum** (Hedw.) Müll.Hal.
6 **A. piligerum** (De Not.) Limpr.
7 **A. triquetrum** (Spruce) Müll.Hal.

73 **Aloina** Kindb.

- 1 **A. aloides** (Koch ex Schultz) Kindb.
2 **A. ambigua** (Bruch & Schimp.) Limpr.²³³
3 **A. bifrons** (De Not.) Delgad.
4 **A. brevirostris** (Hook. & Grev.) Kindb.
5 **A. humilis** M.T.Gallego, M.J.Cano & Ros
6 **A. obliquifolia** (Müll.Hal.) Broth.²³⁴
7 **A. rigida** (Hedw.) Limpr.

74 **Barbula** Hedw.²³⁵

- 1 **B. unguiculata** Hedw.

²²⁹*Ceratodon amazonum* was described from Spain by Nieto-Lugilde et al. (2018).

²³⁰*Ceratodon conicus* has been shown to have a hybrid origin (Nieto-Lugilde et al. 2018), with *Ceratodon amazonum* and *Ceratodon purpureus* as parent taxa, and was considered by them a nothospecies, *Ceratodon x conicus*. However, it has evidently been forming self-sustaining populations for some time, whereas *Ceratodon amazonum* is apparently a rare species confined to Spain, so it is here considered a true species of hybrid origin.

²³¹Nieto-Lugilde et al. (2018) presented molecular data suggesting that specimens corresponding to subsp. *stenocarpus* were not molecularly distinct from subsp. *purpureus*, but a formal synonymy was not made, and the two taxa are retained for the present.

²³²Inoue and Tsubota (2016) used molecular techniques to confirm that there are four clades within the Pottiaceae, corresponding to the subfamilies Trichostomoideae, Pottioideae, Merceoideae and Streblotrichoideae. Accordingly, the subfamily Barbuloideae (Herzog) Hilp. is no longer used.

²³³*Aloina ambigua* is doubtfully separate from *Aloina aloides*. The differentiating characters are quantitative and overlapping. Molecular work is needed to elucidate further.

²³⁴Although currently regarded as distinct, there is some evidence to suggest that *Aloina obliquifolia* may be conspecific with *Aloina rigida* (H. Siebel pers. comm. 2017).

²³⁵Other European species of *Barbula* were moved to the reinstated genera *Hydrogonium* and *Streblotrichum* and the newly described genus *Gymnobarbula* by Kučera, Košnar, et al. (2013).

- 75 **Bryoerythrophyllum** P.C.Chen
 1 **B. alpigenum** (Venturi) P.C.Chen
 2 **B. caledonicum** D.G.Long
 3 **B. campylocarpum** (Müll.Hal.) H.A.Crum
 4 **B. duellii** Blockeel²³⁶
 5 **B. ferruginascens** (Stirt.) Giacom.
 6 **B. inaequalifolium** (Taylor) R.H.Zander
 7 **B. recurvirostrum** (Hedw.) P.C.Chen
 8 **B. rubrum** (Jur. ex Geh.) P.C.Chen
- 76 **Chenia** R.H.Zander²³⁷
 1 **C. leptophylla** (Müll.Hal.) R.H.Zander [*Leptophascum leptophyllum* (Müll.Hal.) J.Guerra & M.J.Cano]
 2 **C. ruigtevleia** Hedd. & R.H.Zander²³⁸
- 77 **Cinclidotus** P.Beauv.
 1 **C. aquaticus** (Hedw.) Bruch & Schimp.
 2 **C. danubicus** Schiffn. & Baumgartner
 3 **C. fontinaloides** (Hedw.) P.Beauv.
 4 **C. riparius** (Host ex Brid.) Arn. [*Cinclidotus confertus* Lüth]²³⁹
 5 **C. vivesii** Ederra
- 78 **Crossidium** Jur.
 1 **C. aberrans** Holz. & E.B.Bartram
 2 **C. crassinervium** (De Not.) Jur.
 3 **C. davidai** Catches.
 4 **C. geheebii** (Broth.) Broth.
 5 **C. laevipilum** Thér. & Trab.²⁴⁰
 6 **C. laxefilamentosum** W.Frey & Kürschner²⁴¹
 7 **C. squamiferum** (Viv.) Jur.
 a var. **pottioideum** (De Not.) Mönk.²⁴²
 b var. **squamiferum**
- 79 **Dialytrichia** (Schimp.) Limpr.
 1 **D. mucronata** (Brid.) Broth.
 2 **D. saxicola** (Lamy) M.J.Cano [*Dialytrichia fragilifolia* (Bizot & J.Roux) F.Lara]²⁴³
- 80 **Didymodon** Hedw. [*Aithobryum* R.H.Zander, *Exobryum* R.H.Zander, *Fuscobryum* R.H.Zander, *Geheebia* Schimp., *Trichostomopsis* Cardot, *Vinealobryum* R.H.Zander]²⁴⁴
 1 **D. acutus** (Brid.) K.Saito
 2 **D. asperifolius** (Mitt.) H.A.Crum, Steere & L.E.Anderson [*Exobryum asperifolium* (Mitt.) R.H.Zander]
 3 **D. australasiae** (Hook. & Grev.) R.H.Zander [*Trichostomopsis australasiae* (Hook. & Grev.) H.Rob.]
 4 **D. bistratosus** Hébr. & R.B.Pierrot [*Aithobryum bistratosum* (Hébr. & R.B.Pierrot) R.H.Zander]
 5 **D. brachyphyllus** (Sull.) R.H.Zander [*Didymodon lamyanus* (Schimp.) Thér., *Vinealobryum brachyphyllum* (Sull.) R.H.Zander]²⁴⁵
 6 **D. cordatus** Jur. [*Vinealobryum cordatum* (Jur.) R.H.Zander]
 7 **D. eckeliae** R.H.Zander [*Vinealobryum eckeliae* (R.H.Zander) R.H.Zander]²⁴⁶
 8 **D. fallax** (Hedw.) R.H.Zander [*Geheebia fallax* (Hedw.) R.H.Zander]
 9 **D. ferrugineus** (Schimp. ex Besch.) M.O.Hill [*Geheebia ferruginea* (Schimp. ex Besch.) R.H.Zander]
 10 **D. giganteus** (Funck) Jur. [*Geheebia gigantea* (Funck) Boulay]
 11 **D. glaucus** Ryan
 a subsp. **glaucus**
 b subsp. **verbanus** (W.E.Nicholson & Dixon) Jan Kučera [*Didymodon verbanus* (W.E. Nicholson & Dixon) Loeske]²⁴⁷
 12 **D. icmadophilus** (Schimp. ex Müll.Hal.) K.Saito
 13 **D. insulanus** (De Not.) M.O.Hill [*Vinealobryum insulanum* (De Not.) R.H.Zander]
 14 **D. johansenii** (R.S.Williams) H.A.Crum [*Exobryum johansenii* (R.S.Williams) R.H.Zander]
 15 **D. luridus** Hornsch. [*Vinealobryum luridum* (Hornsch.) R.H.Zander]

²³⁶*Bryoerythrophyllum duellii* was described by Blockeel et al. (2017), and is known from Greece, Crete and Cyprus.

²³⁷The treatment of Hedderson and Zander (2008) is followed for *Chenia*.

²³⁸*Chenia ruigtevleia* is a South African species (Hedderson and Zander 2008) recently identified from Spain (Ellis, Aleffi, et al. 2016). Further work is needed to confirm that the Spanish and South African material are actually conspecific.

²³⁹Recent molecular work has shown *Cinclidotus confertus* to be identical with *C. riparius* (M. Lüth pers. comm. 2017).

²⁴⁰*Crossidium laevipilum* is doubtfully morphologically separable from *Crossidium crassinervium*, but molecular data (Kučera et al. in prep.) confirms its identity.

²⁴¹*Crossidium laxefilamentosum* is doubtfully separable from *Crossidium crassinervium*.

²⁴²*Crossidium squamiferum* var. *pottioideum* is doubtfully separable from var. *squamiferum*.

²⁴³The combination *Dialytrichia saxicola*, which has priority over *Dialytrichia fragilifolia*, was made by Cano (2007).

²⁴⁴The segregation of *Aithobryum*, *Exobryum*, *Fuscobryum*, *Geheebia*, *Trichostomopsis* and *Vinealobryum*, proposed by Zander (2013, 2019), is essentially unnecessary or unsupported by molecular phylogenetic analyses.

²⁴⁵Jiménez (2006) revised the European species of *Didymodon* and synonymised *Didymodon lamyanus* with *Didymodon brachyphyllus*.

²⁴⁶*Didymodon eckeliae* was reported new to Europe by Puche et al. (2006). European plants are morphologically similar to American ones, but preliminary molecular work by J. Kučera shows that they do not belong to the same lineages. The identity of the European plants therefore remains problematic.

²⁴⁷*Didymodon verbanus* is a problematic taxon recorded in Europe from Switzerland, Austria, Italy and Germany. It is probably best recognised as a subspecies of *Didymodon glaucus*, so the new combination is proposed above.

- 16 ***D. maschalogenus*** (Renauld & Cardot) Broth. [*Gehebia maschalogenus* (Ren. & Card.) R.H.Zander].²⁴⁸
- 17 ***D. maximus*** (Syed & Crundw.) M.O.Hill [*Gehebia maxima* (Syed & Crundw.) R.H.Zander]
- 18 ***D. nicholsonii*** Culm. [*Vinealobryum nicholsonii* (Culm.) R.H.Zander]
- 19 ***D. rigidulus*** Hedw.
- 20 ***D. sinuosus*** (Mitt.) Delogne [*Vinealobryum sinuosum* (Mitt.) R.H.Zander]
- 21 ***D. spadiceus*** (Mitt.) Limpr. [*Gehebia spadicea* (Mitt.) R.H.Zander]
- 22 ***D. subandreaeoides*** (Kindb.) R.H.Zander [*Fuscobryum subandreaeoides* (Kindb.) R.H.Zander]
- 23 ***D. tomaculosus*** (Blockeel) M.F.V.Corley
- 24 ***D. tophaceus*** (Brid.) Lisa [*Gehebia tophacea* (Brid.) R.H.Zander]²⁴⁹
- a subsp. ***erosus*** (J.A.Jiménez & J.Guerra) Jan Kučera [*Didymodon erosus* J.A.Jiménez & J.Guerra]
- b subsp. ***sicculus*** (M.J.Cano, Ros, García-Zam. & J.Guerra) Jan Kučera [*Didymodon sicculus* M.J.Cano, Ros, García-Zam. & J.Guerra]
- c subsp. ***tophaceus***
- i. var. ***anatinus*** Hamm.
- ii. var. ***tophaceus***
- 25 ***D. umbrosus*** (Müll.Hal.) R.H.Zander [*Trichostomopsis umbrosa* (Müll.Hal.) H.Rob.]
- 26 ***D. validus*** Limpr.²⁵⁰
- 27 ***D. vinealis*** (Brid.) R.H.Zander [*Vinealobryum vineale* (Brid.) R.H.Zander]
- 81 ***Hennediella*** Paris
- 1 ***H. heimii*** (Hedw.) R.H.Zander [*Desmatodon wilczekii* Meyl., *Tortula rhodonia* R.H.Zander]²⁵¹
- a var. ***arctica*** (Lindb.) R.H.Zander²⁵²
- b var. ***heimii***
- 2 ***H. macrophylla*** (R.Br.bis) Paris
- 3 ***H. stanfordensis*** (Steere) Blockeel
- 82 ***Hilpertia*** R.H.Zander
- 1 ***H. velenovskyi*** (Schiffn.) R.H.Zander
- 83 ***Leptodontium*** (Müll.Hal.) Hampe
- 1 ***L. flexifolium*** (Dicks.) Hampe
- 2 ***L. gemmascens*** (Mitt.) Braithw.
- 3 ***L. proliferum*** Herzog²⁵³
- 4 ***L. styriacum*** (Jur.) Limpr.²⁵⁴
- 84 ***Microbryum*** Schimp. [*Pottia* Ehrh. ex Fűrnr. p.p.]
- 1 ***M. curvicollum*** (Hedw.) R.H.Zander
- 2 ***M. davallianum*** (Sm.) R.H.Zander²⁵⁵
- a var. ***commutatum*** (Limpr.) R.H.Zander [*Pottia commutata* Limpr.]²⁵⁶
- b var. ***conicum*** (Schleich. ex Schwägr.) R.H.Zander [*Pottia conica* (Schleich. ex Schwägr.) Fűrnr., *Pottia starckeana* subsp. *conica* (Schleich. ex Schwägr.) D.F.Chamb.]
- c var. ***davallianum***
- 3 ***M. floerkeanum*** (F.Weber & D.Mohr) Schimp.
- 4 ***M. fosbergii*** (E.B.Bartram) Ros, O.Werner & Rams
- 5 ***M. longipes*** (J.Guerra J.J.Martínez & Ros) R.H.Zander
- 6 ***M. rectum*** (With.) R.H.Zander
- 7 ***M. starckeana*** (Hedw.) R.H.Zander [*Pottia starckeana* (Hedw.)]
- 8 ***M. vlassovii*** (Laz.) R.H.Zander [*Phascum vlassovii* Laz., *Tortula vlassovii* (Laz.) Ros & Herrnst.]²⁵⁷
- 85 ***Pseudocrossidium*** R.S.Williams²⁵⁸
- 1 ***P. hornschurchianum*** (Schultz) R.H.Zander
- 2 ***P. obtusulum*** (Lindb.) H.A.Crum & L.E.Anderson
- 3 ***P. replicatum*** (Taylor) R.H.Zander
- 4 ***P. revolutum*** (Brid.) R.H.Zander

²⁴⁸*Didymodon maschalogenus* was reported new to Europe by Köckinger and van Melick (2007).

²⁴⁹The treatment of *Didymodon tophaceus* and its infraspecific classification, including the reduction of *Didymodon erosus* and *Didymodon sicculus* to subspecies, follows Kučera, Blockeel, et al. (2018).

²⁵⁰Jiménez (2006) clarified the status of *Didymodon validus*, which was not listed by Hill et al. (2006).

²⁵¹*Tortula rhodonia* was synonymised with *Hennediella heimii* by Cano (2008). *Tortula rhodonia* is a replacement name for *Desmatodon wilczekii*, but all possible type specimens seem to have been lost. Therefore a neotypification of *Desmatodon wilczekii* is necessary in order to complete the synonymisation.

²⁵²*Hennediella heimii* var. *arctica* probably has little or no value but is retained for the present.

²⁵³*Leptodontium proliferum* is a Southern Hemisphere species that was reported from a single site in England by Porley and Edwards (2010). No further localities have been discovered, but plants with characters intermediate between this species and *L. gemmascens* have been found at another site in England (Blockeel 2017). The identity of the English material of *Leptodontium proliferum* requires further study.

²⁵⁴*Leptodontium styriacum* has been synonymised with *Leptodontium flexifolium* (Ignatov, Bersanova, et al. 2005), but it appears that *Leptodontium flexifolium* has an Atlantic distribution, whereas *Leptodontium styriacum* has a continental one. Morphological differences between them are very small and further work is necessary to clarify the situation.

²⁵⁵The variability of *Microbryum davallianum* has not been satisfactorily clarified. Ros et al. (1996) argued for the full synonymy of the earlier recognised varieties but no molecular data have been presented so far to support this view. In view of the mostly distinct morphologies and occasional mixed occurrences, as well as their geographical ranges, which do not fully overlap, they are retained at varietal rank pending final taxonomic clarification.

²⁵⁶There are sufficient morphological differences to recognise *Microbryum davallianum* var. *commutatum* at varietal level, and the same applies to *Microbryum davallianum* var. *conicum*.

²⁵⁷*Microbryum vlassovii* was included in *Phascum* by Hill et al. (2006) but placed in *Tortula* by Ros, Mazimpaka, et al. (2013). However, its reddish coloration with KOH suggests that it may be more correctly placed in *Microbryum*.

²⁵⁸Preliminary molecular studies by J. Kučera suggest that what has been recognised as *Pseudocrossidium obtusulum* and also the European records of *Pseudocrossidium replicatum* may prove to be *Pseudocrossidium hornschurchianum*, but until a revision is completed both species are retained.

- 86 ***Pterygoneurum*** Jur.
 1 ***P. kozlovii*** Laz.
 2 ***P. lamellatum*** (Lindb.) Jur.
 3 ***P. ovatum*** (Hedw.) Dixon [*Pterygoneurum compactum* M.J.Cano, J.Guerra & Ros, *Pterygoneurum crossidioides* W.Frey, Herrnst. & Kürschner, *Pterygoneurum squamosum* Segarra & Kürschner]²⁵⁹
 4 ***P. papillosum*** Oesau
 5 ***P. sampaianum*** (Guim.) Guim.
 6 ***P. subsessile*** (Brid.) Jur.
- 87 ***Stegonia*** Venturi
 1 ***S. latifolia*** (Schwägr.) Venturi ex Broth.
 a var. ***latifolia***
 b var. ***pilifera*** (Brid.) Broth.²⁶⁰
- 88 ***Syntrichia*** Brid.
 1 ***S. bogotensis*** (Hampe) R.H.Zander
 2 ***S. calcicola*** J.J.Amann
 3 ***S. caninervis*** Mitt.
 a var. ***abranthesii*** (Luisier) R.H.Zander
 b var. ***astrakhanica*** Ignatov, Ignatova & Suragina
 c var. ***caninervis***
 d var. ***gypsophila*** (J.J.Amann ex G.Roth) Ochyra
 4 ***S. echinata*** (Schiffn.) Herrnst. & Ben-Sasson
 5 ***S. fragilis*** (Taylor) Ochyra
 6 ***S. handelii*** (Schiffn.) S.Agnew & Vondr.
 7 ***S. laevipila*** Brid. [*Syntrichia pagorum* (Milde) J.J.Amann]²⁶¹
 8 ***S. latifolia*** (Bruch ex Hartm.) Huebener
 9 ***S. minor*** (Bizot) M.T.Gallego, J.Guerra, M.J.Cano, Ros & Sánchez-Moya
 10 ***S. montana*** Nees [*Syntrichia intermedia* Brid.]
 a var. ***calva*** (Durieu & Sagot ex Bruch & Schimp.) J.J.Amann
 b var. ***montana***
 11 ***S. norvegica*** F.Weber
 12 ***S. papillosa*** (Wilson) Jur.
 13 ***S. papillosissima*** (Copp.) Loeske
 14 ***S. princeps*** (De Not.) Mitt.
 15 ***S. rigescens*** (Broth. & Geh.) Ochyra
- 16 ***S. ruraliformis*** (Besch.) Mans. [*Syntrichia ruralis* var. *ruraliformis* (Besch.) Delogne]²⁶²
 17 ***S. ruralis*** (Hedw.) F.Weber & D.Mohr [*Syntrichia densa* (Velen.) J.-P.Frahm, *Syntrichia glabra* J.-P.Frahm & M.T.Gallego]²⁶³
 a var. ***epilosa*** (Venturi) J.J.Amann²⁶⁴
 b var. ***ruralis***
 18 ***S. sinensis*** (Müll.Hal.) Ochyra
 19 ***S. submontana*** (Broth.) Ochyra²⁶⁵
 20 ***S. subpapillosissima*** (Bizot & R.B.Pierrot ex W.A.Kramer) M.T.Gallego & J.Guerra²⁶⁶
 21 ***S. virescens*** (De Not.) Ochyra
- 89 ***Tortula*** Hedw. [*Desmatodon* Brid., *Phascum* Hedw., *Pottia* Ehrh. ex Fürnr. p.p., *Protobryum* (Dicks.) J.Guerra & M.J.Cano]
 1 ***T. acaulon*** (With.) R.H.Zander [*Phascum cuspidatum* Hedw.]²⁶⁷
 a var. ***acaulon*** [*Phascum cuspidatum* var. *cuspidatum*]
 b var. ***papillosa*** (Lindb.) R.H.Zander [*Phascum cuspidatum* var. *papillosum* (Lindb.) G.Roth]
 c var. ***pilifera*** (Hedw.) R.H.Zander [*Phascum cuspidatum* var. *piliferum* (Hedw.) Hook. & Taylor]
 d var. ***retortifolia*** (J.Guerra & Ros) R.H.Zander [*Phascum cuspidatum* var. *retortifolium* J.Guerra & Ros]
 e var. ***schreberiana*** (Dicks.) R.H.Zander [*Phascum cuspidatum* var. *schreberianum* (Dicks.) Brid.]
 2 ***T. amplexa*** (Lesq.) Steere [*Syntrichia amplexa* (Lesq.) R.H.Zander]
 3 ***T. ampliretis*** Crundw. & D.G.Long
 4 ***T. atrovirens*** (Sm.) Lindb.
 5 ***T. bogosica*** (Müll.Hal.) R.H.Zander [*Desmatodon bogosicus* Müll.Hal.]
 6 ***T. bolanderi*** (Lesq. & James) M.Howe
 7 ***T. brevissima*** Schiffn.
 8 ***T. canescens*** Mont.
 9 ***T. caucasica*** Broth. [*Pottia intermedia* (Turner) Fürnr., *Tortula modica* R.H.Zander]²⁶⁸

²⁵⁹Synonymies of *Pterygoneurum ovatum* follow Guerra, Brugués, et al. (2006). However, this treatment is not yet supported by molecular data.

²⁶⁰*Stegonia latifolia* var. *pilifera* is reinstated because it appears to be a distinctive and stable taxon, albeit differing from the type in only a single character, the excurrent costa.

²⁶¹*Syntrichia pagorum* (Milde) J.J.Amann, listed as a synonym of *Syntrichia laevipila* by Hill et al. (2006), may prove to be a good species, but more work is needed (Afonina et al. 2014).

²⁶²Molecular and morphological evidence suggest that *Syntrichia ruraliformis* should be regarded as a species (Hedenäs, Heinrichs, et al. 2019).

²⁶³According to Frahm and Sabovljević (2006), *Syntrichia glabra* is a juvenile form of *Syntrichia densa* (Velen.) J.-P.Frahm. *Syntrichia densa* was synonymised with *Syntrichia calcicola* by Gallego et al. (2002), but Kramer (1980) synonymised *Tortula densa* with *Tortula ruralis*. The latter synonymy is followed here.

²⁶⁴*Syntrichia ruralis* var. *epilosa* was resurrected by Gallego et al. (2018). It may represent a special phenotype within *Syntrichia ruraliformis* or *Syntrichia ruralis* but more study is required (Hedenäs et al. 2019).

²⁶⁵*Syntrichia submontana* is essentially a central Asian species recorded from the Caucasus (Afonina et al. 2014).

²⁶⁶*Syntrichia subpapillosissima* may represent a special phenotype within *Syntrichia ruraliformis* or *Syntrichia ruralis* but more study is required (Hedenäs, Heinrichs, et al. 2019).

²⁶⁷We follow the treatment of Ros, Mazimpaka, et al. (2013) and refer the species included in *Phascum* by Hill et al. (2006) to *Tortula*.

²⁶⁸*Tortula modica* was synonymised with *Tortula caucasica* by Ros, Muñoz, et al. (2008).

- 10 ***T. cernua*** (Huebener) Lindb.
 11 ***T. cuneifolia*** (Dicks.) Turner
 12 ***T. freibergii*** Dixon & Loeske
 13 ***T. guepinii*** (Bruch & Schimp.) Broth. [*Desmatodon guepinii* Bruch & Schimp.]
 14 ***T. hoppeana*** (Schultz) Ochyra [*Desmatodon latifolius* (Hedw.) Brid.]
 15 ***T. inermis*** (Brid.) Mont.
 16 ***T. israelis*** Bizot & F.Bilewsky [*Tortula baetica* (Casas & R.Oliva) J.Guerra & Ros]
 17 ***T. laureri*** (Schultz) Lindb. [*Desmatodon laureri* (Schultz) Bruch & Schimp.]
 18 ***T. leucostoma*** (R.Br.) Hook. & Grev. [*Desmatodon leucostoma* (R.Br.) Berggr.]
 19 ***T. lindbergii*** Broth. [*Pottia lanceolata* (Hedw.) Müll.Hal., *Tortula lanceola* R.H.Zander]²⁶⁹
 20 ***T. lingulata*** Lindb.
 21 ***T. marginata*** (Bruch & Schimp.) Spruce [*Desmatodon meridionalis* Luisier]
 22 ***T. mucronifolia*** Schwägr.
 23 ***T. muralis*** Hedw.
 a subsp. ***muralis***²⁷⁰
 i. var. ***aestiva*** Hedw.
 ii. var. ***muralis***
 b subsp. ***obtusifolia*** (Schwägr.) Culm. [*Tortula obtusifolia* (Schwägr.) Mathieu]²⁷¹
 24 ***T. pallida*** (Lindb.) R.H.Zander
 25 ***T. protobryoides*** R.H.Zander [*Protobryum bryoides* (Dicks.) J.Guerra & M.J.Cano]²⁷²
 26 ***T. randii*** (Kenn.) R.H.Zander [*Desmatodon randii* (Kenn.) Laz.]
 27 ***T. revolvens*** (Schimp.) G.Roth
 28 ***T. schimperi*** M.J.Cano, O.Werner & J.Guerra [*Tortula subulata* var. *angustata* (Schimp.) Limpr.]
 29 ***T. solmsii*** (Schimp.) Limpr.
 30 ***T. subulata*** Hedw. [*Tortula subulata* var. *graeffii* Warnst., *Tortula subulata* var. *subinermis* (Bruch & Schimp.) Wilson]
 31 ***T. systylia*** (Schimp.) Lindb. [*Desmatodon systylius* Schimp.]
 32 ***T. truncata*** (Hedw.) Mitt.
 33 ***T. ucrainica*** (Laz.) R.H.Zander [*Desmatodon ucrainicus* Laz.]
 34 ***T. vahliana*** (Schultz) Mont.
 35 ***T. viridifolia*** (Mitt.) Blockeel & A.J.E.Sm. [*Pottia crinita* Bruch & Schimp.]
 36 ***T. wilsonii*** (Hook.) R.H.Zander
 90 ***Triquetrella*** Müll.Hal.
 1 ***T. arapilensis*** Luisier
 Streblotrichoideae Y.Inoue & H.Tsubota²⁷³
 91 ***Streblotrichum*** P.Beauv.²⁷⁴
 1 ***S. convolutum*** (Hedw.) P.Beauv. [*Barbula convoluta* Hedw.]
 a var. ***commutatum*** (Jur.) J.J.Amann [*Barbula convoluta* var. *sardoa* Schimp., *Streblotrichum commutatum* (Jur.) Hilp.]²⁷⁵
 b var. ***convolutum***
 2 ***S. enderesii*** (Garov.) Loeske [*Barbula enderesii* Garov.]
 Trichostomoideae Broth.
 92 ***Anoetangium*** Schwägr.
 1 ***A. aestivum*** (Hedw.) Mitt.
 2 ***A. angustifolium*** Mitt.²⁷⁶
 3 ***A. handelii*** Schiffn.
 93 ***Aschisma*** Lindb.
 1 ***A. carniolicum*** (F.Weber & D.Mohr) Lindb.
 2 ***A. cuynetii*** (Bizot & R.B.Pierrot) J.Guerra & M.J.Cano
 94 ***Chionoloma*** Dixon [*Paraleptodontium* D.G.Long, *Oxystegus* (Limpr.) Hilp.]²⁷⁷
 1 ***C. bombayense*** (Müll.Hal.) P.Sollman [*Pseudosymblepharis bombayensis* (Müll.Hal.) P.Sollman]²⁷⁸
 2 ***C. cylindrotheca*** (Mitt.) M.Alonso, M.J.Cano & J.A.Jiménez [*Chionoloma daldinianum* (De Not.) M.Alonso, M.J.Cano & J.A.Jiménez, *Oxystegus daldinianus* (De Not.) Köckinger, O.Werner & Ros]²⁷⁹
 3 ***C. hibernicum*** (Mitt.) M.Alonso, M.J.Cano & J.A.Jiménez [*Oxystegus hibernicum* (Mitt.) Hilp., *Trichostomum hibernicum* (Mitt.) Dixon]

²⁶⁹*Tortula lanceola* was synonymised with *Tortula lindbergii* by Ros, Muñoz, et al. (2008).

²⁷⁰The two varieties of *Tortula muralis* subsp. *muralis* intergrade but there is a certain level of distinctness, so this treatment follows Košnar and Kolář (2009).

²⁷¹Košnar and Kolář (2009) reduced *Tortula obtusifolia* to a subspecies of *Tortula muralis*.

²⁷²We follow the treatment of Ros, Mazimpaka, et al. (2013) for *Tortula protobryoides*.

²⁷³The subfamily Streblotrichoideae was proposed by Inoue and Tsubota (2016).

²⁷⁴*Streblotrichum* was re-established on molecular evidence by Kučera, Košnar, et al. (2013).

²⁷⁵*Streblotrichum convolutum* var. *commutatum* is retained at varietal rank. Although it is not always morphologically distinct, molecular evidence clearly separates the two taxa (Kučera, Košnar, et al. 2013).

²⁷⁶Unpublished molecular work by J. Kučera shows *Anoetangium angustifolium* to be distinct from *Anoetangium aestivum*.

²⁷⁷While *Oxystegus* is a molecularly well-delimited group, retaining it leaves problems, notably in the delimitation of *Pseudosymblepharis* Broth. Therefore, we follow Alonso et al. (2016) in placing these species in *Chionoloma*.

²⁷⁸*Chionoloma bombayense*, recorded in Europe from the Caucasus, is retained in the checklist for the present, following Ignatova, Kuznetsova, et al. (2012). However, Alonso et al. (2018) circumscribe *Chionoloma bombayense* in a much narrower sense, leaving many other plants in *Chionoloma bombayense* s.lat., including the Caucasus plant, without a name.

²⁷⁹Originally described as *Didymodon cylindricus* var. *daldinianus* De Not., *Chionoloma daldinianum* was treated in various ways historically, including being omitted altogether by Hill et al. (2006); it was resurrected as a species with molecular support by Köckinger, Werner, et al. (2010), then brought into *Chionoloma* by Alonso et al. (2016). Later, Alonso et al. (2019) synonymised *Chionoloma daldinianum* with the more widespread *Chionoloma cylindrotheca*.

- 4 **C. minus** (Köckinger, O.Werner & Ros) M.Alonso, M.J.Cano & J.A.Jiménez [*Oxystegus minor* Köckinger, O.Werner & Ros]²⁸⁰
- 5 **C. recurvifolium** (Taylor) M.Alonso, M.J.Cano & J.A.Jiménez [*Oxystegus recurvifolius* (Taylor) R.H.Zander, *Paraleptodontium recurvifolium* (Taylor) D.G.Long, *Trichostomum recurvifolium* (Taylor) R.H.Zander]
- 6 **C. tenuirostre** (Hook. & Taylor) M.Alonso, M.J.Cano & J.A.Jiménez [*Oxystegus tenuirostris* (Hook. & Taylor) A.J.E.Sm., *Trichostomum tenuirostre* (Hook. & Taylor) Lindb.]
- a var. **holtii** (Braithw.) M.Alonso, M.J.Cano & J.A.Jiménez [*Oxystegus tenuirostris* var. *holtii* (Braithw.) A.J.E.Sm.]
- b var. **tenuirostre**
- 95 **Ephemerum** Hampe²⁸¹
- 1 **E. cohaerens** (Hedw.) Hampe
- 2 **E. crassinervium** (Schwägr.) Hampe²⁸²
- a subsp. **rutheanum** (Schimp.) Holyoak [*Ephemerum hibernicum* Holyoak & V.S.Bryan, *Ephemerum rutheanum* Schimp., *Ephemerum serratum* var. *rutheanum* (Schimp.) Jur.]
- b subsp. **sessile** (Bruch) Holyoak [*Ephemerum sessile* (Bruch) Müll.Hal.]
- 3 **E. recurvifolium** (Dicks.) Boulay
- 4 **E. serratum** (Hedw.) Hampe [*Ephemerum minutissimum* Lindb., *Ephemerum serratum* var. *minutissimum* (Lindb.) Grout]²⁸³
- 5 **E. spinulosum** Bruch & Schimp. ex Schimp.
- 6 **E. stoloniferum** (Hedw.) L.T.Ellis & M.J.Price [*Ephemerum serratum* auct., *Ephemerum stellatum* H.Philib.]²⁸⁴
- 96 **Eucladium** Bruch & Schimp.
- 1 **E. verticillatum** (With.) Bruch & Schimp. [*Eucladium verticillatum* var. *angustifolium* Lindb.]²⁸⁵
- 97 **Gymnobarbula** Jan Kučera²⁸⁶
- 1 **G. bicolor** (Bruch & Schimp.) Jan Kučera [*Barbula bicolor* (Bruch & Schimp.) Lindb.]
- 98 **Gymnostomum** Nees & Hornsch.
- 1 **G. aeruginosum** Sm.
- a var. **aeruginosum**
- b var. **obscurum** J.Guerra
- 2 **G. calcareum** Nees & Hornsch. [*Gymnostomum calcareum* var. *lanceolatum* Sérgio, *Gymnostomum lanceolatum* M.J.Cano, Ros & J.Guerra]²⁸⁷
- a var. **atlanticum** Sérgio²⁸⁸
- b var. **calcareum**
- 3 **G. viridulum** Brid.
- 99 **Gyroweisia** Schimp.
- 1 **G. reflexa** (Brid.) Schimp.
- 2 **G. tenuis** (Hedw.) Schimp.
- 100 **Hydrogonium** (Müll.Hal.) A.Jaeger²⁸⁹
- 1 **H. amplexifolium** (Mitt.) P.C.Chen [*Barbula amplexifolia* (Mitt.) A.Jaeger]²⁹⁰
- 2 **H. bolleanum** (Müll.Hal.) A.Jaeger [*Barbula bolleana* (Müll.Hal.) Broth.]
- 3 **H. consanguineum** (Thwaites & Mitt.) Hilp. [*Barbula consanguinea* (Thwaites & Mitt.) A.Jaeger]²⁹¹
- a var. **kurilense** (Ignatova & Ignatov) Jan Kučera²⁹²
- 4 **H. croceum** (Brid.) Jan Kučera [*Barbula crocea* (Brid.) F.Weber & D.Mohr]

²⁸⁰*Oxystegus minor* was described as a new species by Köckinger, Werner, et al. (2010), then brought into *Chionoloma* by Alonso et al. (2016). However, significant doubts remain about its distinctiveness as a species: in north-western Europe it intergrades morphologically with *Chionoloma hibernicum*.

²⁸¹All recent phylogenies (e.g. Werner, Ros, et al. 2005) agree in placing *Ephemerum* within the Trichostomoideae rather than its own family, however strange this may seem morphologically.

²⁸²The treatment of *Ephemerum crassinervium* and related taxa follows Holyoak (2010).

²⁸³As a consequence of lectotypification, *Ephemerum minutissimum* Lindb. was placed in synonymy with *Ephemerum serratum* (Hedw.) Hampe (Ellis and Price 2015)

²⁸⁴*Ephemerum stellatum* was synonymised with *Ephemerum serratum* by Holyoak (2010). Subsequently, Ellis and Price (2015) lectotypified *E. stoloniferum* and showed that it was the correct name for the large-spored taxon previously treated under the name *E. serratum*.

²⁸⁵*Eucladium verticillatum* var. *angustifolium* is regarded as nothing more than a form of the type.

²⁸⁶*Gymnobarbula* was established as a new genus by Kučera, Košnar, et al. (2013).

²⁸⁷*Gymnostomum lanceolatum* was reduced to a variety of *Gymnostomum calcareum* by Sérgio (2006). We treat it as a synonym, as it appears to be merely a form of that species.

²⁸⁸*Gymnostomum calcareum* var. *atlanticum* was described by Sérgio (2006). On molecular evidence, it is a well-marked variety that may be sufficiently distinct to be elevated to specific status.

²⁸⁹*Hydrogonium* was re-established on molecular evidence by Kučera, Košnar, et al. (2013).

²⁹⁰*Barbula amplexifolia* was reported new to Europe by Köckinger and Kučera (2007) and transferred to *Hydrogonium* by Kučera, Košnar, et al. (2013).

²⁹¹*Hydrogonium consanguineum* was earlier reported from Europe as *Barbula indica*; Köckinger, Kučera, et al. (2012) proved that these reports belong to *Barbula consanguinea*, transferred to *Hydrogonium* by Kučera, Košnar, et al. (2013).

²⁹²The European expression of *Hydrogonium consanguineum* is var. *kurilense*; var. *consanguineum* occurs in Asia; var. *cancellatum* (Müll.Hal.) Jan Kučera is a North American plant (Kučera, Košnar, et al. 2013).

- 101 **Hymenostylium** Brid. [*Ardeuma* R.H.Zander & Hedd.]²⁹³
- 1 ***H. gracillimum*** (Nees & Hornsch.) Köckinger & Jan Kučera [*Gymnostomum boreale* Nyholm & Hedenäs]²⁹⁴
 - 2 ***H. recurvirostrum*** (Hedw.) Dixon [*Gymnostomum recurvirostrum* Hedw.]
 - a var. ***insigne*** (Dixon) E.B.Bartram [*Ardeuma annotinum* (Mitt. ex Dixon) R.H. Zander & Brinda, *Ardeuma insigne* (Dixon) R.H.Zander & Hedd., *Hymenostylium insigne* (Dixon) Podp.]
 - b var. ***recurvirostrum*** [*Ardeuma recurvirostrum* (Hedw.) R.H.Zander & Hedd.]
 - 3 ***H. xerophilum*** Köckinger & Jan Kučera²⁹⁵
- 102 **Hyophila** Brid.
- 1 ***H. involuta*** (Hook.) A.Jaeger
- 103 **Leptobarbula** Schimp.
- 1 ***L. berica*** (De Not.) Schimp.
- 104 **Molendoa** (Müll.Hal.) Hampe
- 1 ***M. hornschurchiana*** (Hook.) Lindb. ex Limpr. [*Anoetangium hornschurchianum* (Hook.) Funck ex Hornsch., *Anoetangium sendtnerianum* Bruch & Schimp., *Molendoa sendtneriana* (Bruch & Schimp.) Limpr., *Molendoa tenuinervis* Limpr.]²⁹⁶
 - 2 ***M. schliephackei*** (Schlieph.) R.H.Zander [*Anoetangium schliephackei* (Schlieph.) Paris]
 - 3 ***M. taeniatifolia*** Herzog [*Anoetangium taeniatifolium* (Herzog) M.O.Hill]
 - 4 ***M. warburgii*** (Crundw. & M.O.Hill) R.H.Zander [*Anoetangium warburgii* (Limpr.) Paris]
- 105 **Pottiopsis** Blockeel & A.J.E.Sm.
- 1 ***P. caespitosa*** (Brid.) Blockeel & A.J.E.Sm. [*Trichostomum pallidisetum* H.Müll., *Trichostomum triumphans* De Not., *Weissia triumphans* (De Not.) M.O.Hill, *Weissia tyrrhena* M.Fleisch.]²⁹⁷
- 106 **Splachnobryum** Müll.Hal.
- 1 ***S. obtusum*** (Brid.) Müll.Hal.
- 107 **Tortella** (Müll.Hal.) Limpr. [*Pleurochaete* Lindb.]
- 1 ***T. alpicola*** Dixon
 - 2 ***T. x cuspidatissima*** (Cardot & Thér.) O.Werner, Köckinger & Ros²⁹⁸
 - 3 ***T. densa*** (Lorentz & Molendo) Crundw. & Nyholm [*Tortella inclinata* var. *densa* (Lorentz & Molendo) Limpr.]
 - 4 ***T. fasciculata*** (Culm.) Culm. [*Tortella bambergeri* auct., non (Schimp.) Broth. p.p., *Tortella tortuosa* subsp. *fasciculata* Culm.]²⁹⁹
 - 5 ***T. flavovirens*** (Bruch) Broth. [*Tortella limosella* (Stirt.) P.W.Richards & E.C.Wallace]³⁰⁰
 - a var. ***flavovirens***
 - b var. ***glareicola*** (T.A.Chr.) Crundw. & Nyholm
 - c var. ***papillosissima*** Sérgio & Casas
 - 6 ***T. fragilis*** (Drumm.) Limpr.³⁰¹
 - 7 ***T. humilis*** (Hedw.) Jenn.
 - 8 ***T. inclinata*** (R.Hedw.) Limpr.
 - 9 ***T. inflexa*** (Bruch) Broth.
 - 10 ***T. limbata*** (Schiffn.) Geh. & Herzog
 - 11 ***T. mediterranea*** Köckinger, Lüth, O.Werner & Ros³⁰²
 - 12 ***T. nitida*** (Lindb.) Broth.
 - 13 ***T. pseudofragilis*** (Thér.) Köckinger & Hedenäs [*Tortella bambergeri* auct., non (Schimp.) Broth. p.p., *Tortella fragilis* var. *moravica* Podp.]³⁰³
 - 14 ***T. rigens*** Alberts.
 - 15 ***T. spitsbergensis*** (Bizot & Thér.) O.Werner, Köckinger & Ros [*Trichostomum arcticum* Kaal.]³⁰⁴
 - 16 ***T. squarrosa*** (Brid.) Limpr. [*Pleurochaete squarrosa* (Brid.) Lindb.]³⁰⁵

²⁹³The transfer of most *Hymenostylium* species, including *Hymenostylium recurvirostrum* and its varieties, to *Ardeuma* (Zander and Hedderson 2016) is not supported according to molecular evidence (Kučera in prep.).

²⁹⁴The position of *Hymenostylium gracillimum* and the synonymy of *Gymnostomum boreale* were clarified by Köckinger and Kučera (2011).

²⁹⁵*Hymenostylium xerophilum* was described from Austria as a new species by Köckinger and Kučera (2011). It has since been found in Germany and Crimea.

²⁹⁶Geissler (1985) proposed the synonymy of *Molendoa sendtneriana* and *Molendoa tenuinervis* with *Molendoa hornschurchiana*. This is followed here, as it is supported by preliminary molecular results by J. Kučera.

²⁹⁷The position of *Pottiopsis caespitosa* and its synonymies were clarified by Ros and Werner (2007). However, *Trichostomum/Weissia triumphans* is a distinctive form, and continues to be recognised by many bryologists.

²⁹⁸Werner, Köckinger, et al. (2014) used molecular techniques to show that *Tortella x cuspidatissima* was of hybrid origin, with *Tortella arctica* (Arnell) Crundw. & Nyholm and *Tortella spitsbergensis* as parents. *Tortella arctica* itself does not occur in Europe. The record of *Tortella arctica* from Jämtland in Sweden (Ellis, Akhoondi Darzikolaei, et al. 2011) proved, according to the molecular data, to be a form of *Tortella tortuosa* (Köckinger and Hedenäs 2017).

²⁹⁹*Tortella fasciculata* was resurrected by Köckinger and Hedenäs (2017) as part of their revision of *Tortella bambergeri*.

³⁰⁰Further studies on *Tortella limosella* are necessary to determine the final position of *Tortella limosella* within *Tortella flavovirens*.

³⁰¹The authorship of *Tortella fragilis* was discussed in Ochyra, Lewis Smith, et al. (2008). The name should correctly be ascribed to Drummond.

³⁰²*Tortella mediterranea* was described from southern Europe by Köckinger, Lüth, et al. (2018).

³⁰³The name *Tortella pseudofragilis* was introduced by Köckinger and Hedenäs (2017) as part of their revision of *Tortella bambergeri*.

³⁰⁴The identity and relationships of *Tortella spitsbergensis* were clarified by Werner, Köckinger, et al. (2014) using molecular techniques.

³⁰⁵We follow Ros, Mazimpaka, et al. (2013) in accepting Werner, Ros, et al. (2005) and Grundmann et al. (2006), who include *Pleurochaete squarrosa* within the genus *Tortella*.

- 17 ***T. tortuosa*** (Hedw.) Limpr. [*Tortella bambergeri* (Schimp.) Broth.]³⁰⁶
- 108 ***Trichostomum*** Bruch.
 1 ***T. brachydontium*** Bruch
 2 ***T. crispulum*** Bruch
- 109 ***Weissia*** Hedw.
 1 ***W. angustifolia*** (Baumgartner) D.A.Callaghan [*Weissia longifolia* var. *angustifolia* (Baumgartner) Crundw. & Nyholm]³⁰⁷
 2 ***W. brachycarpa*** (Nees & Hornsch.) Jur.
 3 ***W. condensata*** (Voit) Lindb.
 a var. ***armata*** (Thér. & Trab.) M.J.Cano, Ros & J.Guerra
 b var. ***condensata***
 4 ***W. controversa*** Hedw.
 a var. ***controversa***
 b var. ***crispata*** (Nees & Hornsch.) Nyholm
 c var. ***densifolia*** (Bruch & Schimp.) Wilson³⁰⁸
 5 ***W. levieri*** (Limpr.) Kindb.
 6 ***W. longifolia*** Mitt.
 7 ***W. x mittenii*** (Bruch & Schimp.) Mitt. emend. A.J.E.Sm.³⁰⁹
 8 ***W. perssonii*** Kindb.
 9 ***W. rostellata*** (Brid.) Lindb.
 10 ***W. rutilans*** (Hedw.) Lindb.
 11 ***W. squarrosa*** (Nees & Hornsch.) Müll.Hal.
 12 ***W. sterilis*** W.E.Nicholson
 13 ***W. wilsonii*** D.A.Callaghan [*Weissia multicapsularis* auct.]³¹⁰
 14 ***W. wimmeriana*** (Sendtn.) Bruch & Schimp.
- Grimmiales M.Fleisch.
 Saelaniaceae Ignatov & Fedosov³¹¹
 110 ***Saelania*** Lindb.
 1 ***S. glaucescens*** (Hedw.) Broth.
- Seligeriaceae Schimp.
 111 ***Blindia*** Bruch & Schimp.
 1 ***B. acuta*** (Hedw.) Bruch & Schimp.
 2 ***B. caespiticia*** (F.Weber & D.Mohr) Müll.Hal.
- 112 ***Blindiadelphus*** (Lindb.) Fedosov & Ignatov [*Seligeria* Bruch & Schimp. subgenus *Blindiadelphus* Lindb.]³¹²
 1 ***B. campylopodus*** (Kindb.) Fedosov & Ignatov [*Seligeria campylopada* Kindb.]
 2 ***B. diversifolius*** (Lindb.) Fedosov & Ignatov [*Seligeria diversifolia* Lindb.]
 3 ***B. polaris*** (Berggr.) Fedosov & Ignatov [*Seligeria polaris* Berggr.]
 4 ***B. recurvatus*** (Hedw.) Fedosov & Ignatov [*Seligeria recurvata* (Hedw.) Bruch & Schimp.]
 5 ***B. subimmersus*** (Lindb.) Fedosov & Ignatov [*Seligeria subimmersa* Lindb.]
- 113 ***Seligeria*** Bruch & Schimp. subgenus *Seligeria*
 1 ***S. acutifolia*** Lindb.
 2 ***S. brevifolia*** (Lindb.) Lindb.
 3 ***S. calcarea*** (Hedw.) Bruch & Schimp.
 4 ***S. calycina*** Mitt. ex Lindb. [*Seligeria paucifolia* auct. non (With.) Carruth.]
 5 ***S. donniana*** (Sm.) Müll.Hal. [*Seligeria galinae* Mogensen & I.Goldberg]³¹³
 6 ***S. pusilla*** (Hedw.) Bruch & Schimp. subgenus *Megalosporia* Vitt [*Trochobryum* Breidl. & Beck]
 7 ***S. austriaca*** T.Schauer
 8 ***S. carniolica*** (Breidl. & Beck) Nyholm [*Trochobryum carniolicum* Breidl. & Beck]
 9 ***S. irrigata*** (H.K.G.Paul) Ochyra & Gos
 10 ***S. oelandica*** C.E.O.Jensen & Medelius
 11 ***S. patula*** (Lindb.) I.Hagen [*Seligeria alpestris* T.Schauer, *Seligeria patula* var. *alpestris* (T.Schauer) Gos & Ochyra, *Seligeria tristichoides* var. *patula* (Lindb.) Broth.]
 12 ***S. trifaria*** (Brid.) Lindb.
 a var. ***longifolia*** (Lindb. ex Broth.) Ochyra & Gos³¹⁴
 b var. ***trifaria***
 13 ***S. tristichoides*** Kindb.
- Ptychomitriaceae Schimp.
 114 ***Brachydontium*** Fűrnr.³¹⁵

³⁰⁶Using morphological and molecular methods, Köckinger and Hedenäs (2017) demonstrated that the type of *Tortella bambergeri* was synonymous with *Tortella tortuosa*. Plants recently treated as *Tortella bambergeri* are referable to *Tortella fasciculata* and *Tortella pseudofragilis*. Up to now, the acceptance of *Tortella tortuosa* var. *fragilifolia* (Jur.) Limpr. has been based on the treatment of Eckel (1998), but this concept is based on plants later referred to *Tortella bambergeri* (Eckel 2010). In general, the name 'var. *fragilifolia*' has been used for morphs of *Tortella tortuosa* with fragile leaves, and the variety is therefore not included in the checklist.

³⁰⁷*Weissia longifolia* var. *angustifolia* was raised to species level by Callaghan et al. (2019) on the basis of morphological and molecular evidence.

³⁰⁸*Weissia controversa* var. *densifolia* is retained as a variety for now, but it is very doubtfully distinct from var. *controversa*, and significantly less so than var. *crispata*.

³⁰⁹Smith (2004) was the first to propose that *Weissia x mittenii* was a hybrid, and this was accepted by Hill et al. (2006). The name was later typified by Callaghan (2019).

³¹⁰*Weissia wilsonii* was newly described by Callaghan et al. (2019) to replace the taxon which was named *Weissia multicapsularis* (Sm.) Mitt. by modern authors. The type of the latter refers to *Tortula acaulon*, into which plants of *Weissia longifolia* were admixed.

³¹¹Molecular studies by Fedosov, Fedorova, Fedosov, et al. (2016) showed that *Saelania* is best placed in its own family within the Grimmiales.

³¹²*Blindiadelphus* was segregated from *Seligeria* on the basis of morphological and molecular evidence (Fedosov, Fedorova, Ignatova, et al. 2017).

³¹³*Seligeria galinae* was synonymised with *Seligeria donniana* by Fedosov, Fedorova, Ignatova, et al. (2017).

³¹⁴*Seligeria trifaria* var. *longifolia* was not mentioned in Hill et al. (2006) but is regarded as a distinct taxon in the central European mountains (Ochyra and Gos 1992).

³¹⁵Molecular evidence places *Brachydontium* in the Ptychomitriaceae (Fedosov et al. 2017a).

- 1 **B. trichodes** (F.Weber) Milde [*Seligeria transylvanica* Plam.]
- 115 **Campylostelium** Bruch & Schimp.
 1 **C. pitardii** (Corb.) E.Maier [*Grimmia pitardii* Corb.]
 2 **C. saxicola** (F.Weber & D.Mohr) Bruch & Schimp.
 3 **C. strictum** Solms
- 116 **Indusiella** Broth. & Müll.Hal.
 1 **I. thianschanica** Broth. & Müll.Hal.
- 117 **Jaffueliobryum** Thér.
 1 **J. latifolium** Lindb. & Arnell ex Thér.
- 118 **Ptychomitrium** Fűrnr.
 1 **P. incurvum** (Schwägr.) Spruce
 2 **P. nigrescens** (Kunze) Wijk & Margad.
 3 **P. polyphyllum** (Dicks. ex Sw.) Bruch & Schimp.
- Grimmiaceae Arn.
 Racomitrioideae Ochyra & Bedn.-Ochyra
- 119 **Racomitrium** Brid.³¹⁶
 subgenus *Racomitrium*
 1 **R. lanuginosum** (Hedw.) Brid.
 subgenus *Niphotrichum* Bedn.-Ochyra [*Niphotrichum* (Bedn.-Ochyra) Bedn.-Ochyra & Ochyra]
 2 **R. canescens** (Hedw.) Brid. [*Niphotrichum canescens* (Hedw.) Bedn.-Ochyra & Ochyra]
 a subsp. **canescens**
 b subsp. **latifolium** (C.E.O.Jensen) Frisvoll [*Niphotrichum canescens* subsp. *latifolium* (Frisvoll) Bedn.-Ochyra & Ochyra]
 3 **R. elongatum** Ehrh. ex Frisvoll [*Niphotrichum elongatum* (Ehrh. ex Frisvoll) Bedn.-Ochyra & Ochyra]
 4 **R. ericoides** (Brid.) Brid. [*Niphotrichum ericoides* (Brid.) Bedn.-Ochyra & Ochyra]
 5 **R. panschii** (Müll.Hal.) Kindb. [*Niphotrichum panschii* (Müll.Hal.) Bedn.-Ochyra & Ochyra]
 subgenus *Cataracta* Vilh. [*Codriophorus* P.Beauv., *Dilutineuron* Bedn.-Ochyra, Sawicki, Ochyra, Szczecińska & Plášek]
 6 **R. aciculare** (Hedw.) Brid. [*Codriophorus acicularis* (Hedw.) P.Beauv.]
 7 **R. aquaticum** (Brid. ex Schrad.) Brid. [*Codriophorus aquaticus* (Brid. ex Schrad.) Bedn.-Ochyra & Ochyra]
 8 **R. fasciculare** (Hedw.) Brid. [*Codriophorus fascicularis* (Hedw.) Bedn.-Ochyra & Ochyra, *Dilutineuron fasciculare* (Hedw.) Bedn.-Ochyra, Sawicki, Ochyra, Szczecińska & Plášek]
 9 **R. hespericum** Sérgio, J.Muñoz & Ochyra [*Codriophorus hespericus* (Sérgio, J.Muñoz & Ochyra) Bedn.-Ochyra & Ochyra] subgenus *Ellipticodryptodon* (Vilh.) Bedn.-Ochyra & Ochyra [*Bucklandiella* Roiv.]
 10 **R. affine** (F.Weber & D.Mohr) Lindb. [*Bucklandiella affinis* (Schleich. ex F.Weber & D.Mohr) Bedn.-Ochyra & Ochyra]
 11 **R. ellipticum** (Turner) Bruch & Schimp. [*Bucklandiella elliptica* (Turner) Bedn.-Ochyra & Ochyra]
 12 **R. heterostichum** (Hedw.) Brid. [*Bucklandiella heterosticha* (Hedw.) Bedn.-Ochyra & Ochyra]
 13 **R. himalayanum** (Mitt.) A.Jaeger [*Bucklandiella himalayana* (Mitt.) Bedn.-Ochyra & Ochyra]
 14 **R. lamprocarpum** (Müll.Hal.) A.Jaeger [*Bucklandiella lamprocarpa* (Müll.Hal.) Bedn.-Ochyra & Ochyra]
 15 **R. lusitanicum** Ochyra & Sérgio [*Bucklandiella lusitanica* (Ochyra & Sérgio) Bedn.-Ochyra & Ochyra]
 16 **R. macounii** Kindb. [*Bucklandiella macounii* (Kindb.) Bedn.-Ochyra & Ochyra]
 a subsp. **alpinum** (E.Lawton) Frisvoll [*Bucklandiella macounii* subsp. *alpina* (E.Lawton) Bedn.-Ochyra & Ochyra]
 b subsp. **macounii**
 17 **R. microcarpon** (Hedw.) Brid. [*Bucklandiella microcarpa* (Hedw.) Bedn.-Ochyra & Ochyra]
 18 **R. nivale** (Köckinger, Bedn.-Ochyra & Ochyra) Köckinger [*Bucklandiella nivalis* Köckinger, Bedn.-Ochyra & Ochyra]³¹⁷
 19 **R. obtusum** (Brid.) Brid. [*Bucklandiella obtusa* (Brid.) Bedn.-Ochyra & Ochyra]
 20 **R. sudeticum** (Funck) Bruch & Schimp. [*Bucklandiella sudetica* (Funck) Bedn.-Ochyra & Ochyra]
- Grimmioideae Broth.
- 120 **Coscinodon** Spreng.
 1 **C. cribrus** (Hedw.) Spruce [*Grimmia cribrus* Hedw.]
 2 **C. horridus** (Muñoz & H.Hespanhol) Hugonnot, R.D.Porley & Ignatov [*Grimmia horrida* Muñoz & H.Hespanhol]³¹⁸

³¹⁶The treatment of *Racomitrium* follows Larraín et al. (2013), who used molecular evidence for a broad treatment of the genus. Sawicki et al. (2015) used mitochondrial DNA to support narrower genera. However, they used only ten species of *Racomitrium* s.lat. (two from each segregate genus), and the results did not contradict the previous treatment. Indeed, significant problems remain (e.g. *Bucklandiella* sensu Ochyra is not monophyletic) and until *Racomitrium* s.lat. is more completely treated, we prefer to retain *Racomitrium* in a broad sense.

³¹⁷Described as *Bucklandiella nivalis* by Köckinger, Bednarek-Ochyra, et al. (2007) but listed as *Racomitrium nivale* by Köckinger, Suanjak, et al. (2008).

³¹⁸*Grimmia horrida* was described by Muñoz et al. (2009), and transferred to *Coscinodon* by Hugonnot et al. (2018).

- 3 **C. humilis** Milde³¹⁹
- 4 **C. monchiquensis** R.D.Porley, Ochyra & Ignatova³²⁰
- 121 **Grimmia** Hedw. [*Dryptodon* Brid., *Guembelia* Hampe, *Hydrogrimmia* (l.Hagen) Loeske, *Orthogrimmia* (Schimp.) Ochyra & Żarnowiec, *Streptocolea* (l.Hagen) Ochyra & Żarnowiec]³²¹
- 1 **G. alpestris** (F.Weber & D.Mohr) Schleich. [*Orthogrimmia alpestris* (F.Weber & D.Mohr) Ochyra & Żarnowiec]
- 2 **G. anodon** Bruch & Schimp.
- 3 **G. anomala** Hampe ex Schimp.
- 4 **G. arenaria** Hampe [*Orthogrimmia arenaria* (Hampe) Ochyra & Żarnowiec]³²²
- 5 **G. atrata** Miel. ex Hornsch. [*Streptocolea atrata* (Miel. ex Hornsch.) Ochyra & Żarnowiec]
- 6 **G. caespiticia** (Brid.) Jur. [*Orthogrimmia caespiticia* (Brid.) Ochyra & Żarnowiec]
- 7 **G. capillata** De Not.³²³
- 8 **G. crinita** Brid.
- 9 **G. crinitoleucophaea** Cardot [*Grimmia poecilostoma* Cardot & Sebillé]³²⁴
- 10 **G. curviseta** Bouman³²⁵
- 11 **G. decipiens** (Schultz) Lindb.
- 12 **G. dissimulata** E.Maier
- 13 **G. donniana** Sm. [*Orthogrimmia donniana* (Sm.) Ochyra & Żarnowiec]
- 14 **G. elatior** Bruch ex Bals.-Criv. & De Not.
- 15 **G. elongata** Kaulf.
- 16 **G. funalis** (Schwägr.) Bruch & Schimp.
- 17 **G. fuscolutea** Hook.
- 18 **G. hartmanii** Schimp.
- 19 **G. incurva** Schwägr.
- 20 **G. laevigata** (Brid.) Brid. [*Guembelia laevigata* (Brid.) Ochyra & Żarnowiec]
- 21 **G. lisae** De Not.
- 22 **G. longirostris** Hook. [*Guembelia longirostris* (Hook.) Ochyra & Żarnowiec]
- 23 **G. meridionalis** (Müll.Hall.) E.Maier [*Grimmia trichophylla* var. *meridionalis* Müll.Hall.]³²⁶
- 24 **G. mollis** Bruch & Schimp. [*Hydrogrimmia mollis* (Bruch & Schimp.) Loeske]
- 25 **G. montana** Bruch & Schimp. [*Orthogrimmia montana* (Bruch & Schimp.) Ochyra & Żarnowiec]
- 26 **G. muehlenbeckii** Schimp.
- 27 **G. nutans** Bruch [*Grimmia meteorae* C.C.Towns.]
- 28 **G. orbicularis** Bruch ex Wilson [*Dryptodon orbicularis* (Bruch ex Wilson) Ochyra & Żarnowiec]
- 29 **G. ovalis** (Hedw.) Lindb. [*Guembelia ovalis* (Hedw.) Müll.Hal.]
- 30 **G. plagiopodia** Hedw.
- 31 **G. pulvinata** (Hedw.) Sm.
- 32 **G. ramondii** (Lam. & DC.) Margad. [*Dryptodon patens* (Hedw.) Brid., *Grimmia curvata* (Brid.) De Sloover]
- 33 **G. reflexidens** Müll.Hal. [*Grimmia sessitana* De Not., *Orthogrimmia sessitana* (De Not.) Ochyra & Żarnowiec]³²⁷
- 34 **G. teretinervis** Limpr.
- 35 **G. tergestina** Tomm. ex Bruch & Schimp. [*Grimmia tergestina* var. *tergestinoides* (Culm.) Podp., *Guembelia tergestina* (Tomm. ex Bruch & Schimp.) Buys.]³²⁸
- 36 **G. torquata** Drumm.
- 37 **G. trichophylla** Grev. [*Grimmia britannica* A.J.E.Sm.]
- 38 **G. triformis** Carestia & De Not. [*Orthogrimmia triformis* (Carestia & De Not.) Ochyra & Żarnowiec]³²⁹
- 39 **G. ungeri** Jur.³³⁰
- 40 **G. unicolor** Hook.

³¹⁹A molecular study by Ignatova, Kuznetsova, et al. (2008) showed that *Coscinodon humilis* is a European endemic known only from the Alps, whereas plants with this name in east Asia belong to *Coscinodon yukonensis* Hastings.

³²⁰*Coscinodon monchiquensis* was described from the Algarve, Portugal, by Ignatov, Porley, et al. (2018).

³²¹Although Ochyra, Żarnowiec, et al. (2003) presented a morphological scheme comprising seven distinct genera, based largely on the traditional subgenera of *Grimmia*, we choose to treat it in the broad sense. There is molecular support for the paraphyletic nature of *Grimmia* (Streiff 2006; Hernández-Maqueda et al. 2008), but there remain some significant incongruities between plastid and nuclear ITS and the morphological schemes. The evolutionary history of *Grimmia* is complex and further sampling and better phylogenetic markers are needed.

³²²*Grimmia arenaria* was synonymised with *Grimmia donniana* by Maier (2010) but treated as a distinct species in Muñoz et al. (2015). Molecular investigations indicate that it should be accepted as a species (D. Callaghan, pers. comm. 2019).

³²³*Grimmia capillata* was synonymised with *G. crinita* by Maier (2010) but this view has not gained acceptance, since molecular studies support it at specific level. It is recognised at species level in Muñoz et al. (2015).

³²⁴*Grimmia poecilostoma* was synonymised with *Grimmia crinitoleucophaea* by Muñoz et al. (2015). However, the synonymy of *G. crinitoleucophaea* with *Grimmia tergestina* (Maier 2010) has not been widely accepted, on morphological grounds.

³²⁵*Grimmia curviseta* was synonymised with *G. orbicularis* by Maier (2010) but this is disputed; Rodríguez-Romero et al. (2016) used DNA sequencing to conclude that it is an allopolyploid derivative with *G. orbicularis* as one of its progenitors.

³²⁶*Grimmia meridionalis* was elevated to the rank of species by Maier (2002), but, unlike *Grimmia dissimulata*, not accepted by Hill et al. (2006). However, its status as a species has subsequently been supported by molecular evidence (Streiff 2006).

³²⁷Taxonomic, nomenclatural and misidentification issues surround *Grimmia reflexidens*. The situation in Europe remains unresolved but we retain *Grimmia reflexidens* for the present.

³²⁸*Grimmia tergestina* var. *tergestinoides* seems distinct in central Europe. More work is needed to determine its taxonomic value.

³²⁹*Grimmia triformis* was synonymised with *Grimmia donniana* by Maier (2010) but differs in several respects, and was accepted as a species by Muñoz et al. (2015).

³³⁰*Grimmia ungeri* was listed as a synonym of *Grimmia montana* by Maier (2010). Muñoz and Pando (2000) listed *Grimmia ungeri* from several countries, including Spain and Italy, but it was not included by Muñoz et al. (2015). Specimens from Mexico were redetermined as *Grimmia montana*

- 122 **Schistidium** Bruch & Schimp.
- 1 **S. abrupticostatum** (Bryhn) Ignatova & H.H.Blom [*Schistidium platyphyllum* var. *abrupticostatum* (Bryhn) H.H.Blom]³³¹
 - 2 **S. agassizii** Sull. & Lesq.
 - 3 **S. andreaeopsis** (Müll.Hal.) Laz.³³²
 - 4 **S. apocarpum** (Hedw.) Bruch & Schimp.
 - 5 **S. atrichum** (Müll.Hal. & Kindb.) W.A.Weber³³³
 - 6 **S. atrofusum** (Schimp.) Limpr.
 - 7 **S. boreale** Poelt
 - 8 **S. brunnescens** Limpr.
 - a subsp. **brunnescens**
 - b subsp. **griseum** (Nees & Hornsch.) H.H.Blom
 - 9 **S. bryhnii** I.Hagen
 - 10 **S. canadense** (Dupr.) Ignatova & H.H.Blom³³⁴
 - 11 **S. confertum** (Funck) Bruch & Schimp.
 - 12 **S. confusum** H.H.Blom
 - 13 **S. convergens** J.Guerra & M.J.Cano³³⁵
 - 14 **S. crassipilum** H.H.Blom
 - 15 **S. crenatum** H.H.Blom
 - 16 **S. dupretii** (Thér.) W.A.Weber
 - 17 **S. echinatum** Ignatova & H.H.Blom³³⁶
 - 18 **S. elegantulum** H.H.Blom
 - a subsp. **elegantulum**
 - b subsp. **wilsonii** H.H.Blom
 - 19 **S. flaccidum** (De Not.) Ochyra
 - 20 **S. flexipile** (Lindb. ex Broth.) G.Roth
 - 21 **S. frigidum** H.H.Blom
 - a var. **frigidum**
 - b var. **havaasii** H.H.Blom
 - 22 **S. frisvollianum** H.H.Blom
 - 23 **S. grande** Poelt
 - 24 **S. grandirete** H.H.Blom
 - 25 **S. helveticum** (Schkuhr) Deguchi [*Schistidium singarense* (Schiffn.) Laz.]
 - 26 **S. holmenianum** Steere & Brassard
 - 27 **S. lancifolium** (Kindb.) H.H.Blom
 - 28 **S. marginale** H.H.Blom, Bedn.-Ochyra & Ochyra³³⁷
 - 29 **S. maritimum** (Sm. ex R.Scott) Bruch & Schimp.
 - a subsp. **maritimum**
 - b subsp. **piliferum** (I.Hagen) B.Bremer
 - 30 **S. obscurum** H.H.Blom, Köckinger & Ignatova³³⁸
 - 31 **S. occidentale** (E.Lawton) S.P.Churchill
 - 32 **S. papillosum** Culm.
 - 33 **S. platyphyllum** (Mitt.) H.Perss.
 - 34 **S. poeltii** H.H.Blom
 - 35 **S. pruinatum** (Wilson ex Schimp.) G.Roth
 - 36 **S. pulchrum** H.H.Blom
 - 37 **S. recurvum** H.H.Blom
 - 38 **S. rivulare** (Brid.) Podp.
 - 39 **S. robustum** (Nees & Hornsch.) H.H.Blom
 - 40 **S. scandicum** H.H.Blom
 - 41 **S. sibiricum** Ignatova & H.H.Blom³³⁹
 - 42 **S. sinensiapocarpum** (Müll.Hal.) Ochyra³⁴⁰
 - 43 **S. sordidum** I.Hagen
 - 44 **S. spinosum** H.H.Blom & Lüth
 - 45 **S. strictum** (Turner) Loeske ex Mårtensson
 - 46 **S. subconfertum** (Broth.) Deguchi³⁴¹
 - 47 **S. subflaccidum** (Kindb.) H.H.Blom³⁴²
 - 48 **S. subjulaceum** H.H.Blom
 - 49 **S. submuticum** H.H.Blom
 - a subsp. **arcticum** H.H.Blom
 - b subsp. **submuticum**
 - 50 **S. succulentum** Ignatova & H.H.Blom³⁴³
 - 51 **S. tenerum** (J.E.Zetterst.) Nyholm
 - 52 **S. tenuinerve** Ignatova & H.H.Blom³⁴⁴
 - 53 **S. trichodon** (Brid.) Poelt
 - a var. **nutans** H.H.Blom
 - b var. **trichodon**
 - 54 **S. umbrosum** (J.E.Zetterst.) H.H.Blom
 - 55 **S. venetum** H.H.Blom
- Hedwigiales Ochyra
Hedwigiaceae Schimp.
- 123 **Braunia** Bruch & Schimp. [*Hedwigidium* Bruch & Schimp.]

(Delgadillo-Moya 2015). On the other hand, it differs from *Grimmia montana* in some respects (J. Muñoz, pers. comm. 2017). Although we retain it in the checklist for now, *Grimmia ungeri* is taxonomically dubious and may ultimately prove to be synonymous with *Grimmia montana*.

³³¹*Schistidium abrupticostatum* is not very close to *Schistidium platyphyllum* according to molecular studies by Ignatova, Blom, et al. (2010).

³³²*Schistidium andreaeopsis* is an Asian Arctic species recorded more recently from the European Arctic (Blom et al. 2006).

³³³*Schistidium atrichum* is principally a North American species with occurrences in France reported by Chavoutier and Hugonnot (2013).

³³⁴*Schistidium canadense* is a North American species with a single European record from Russia (Karelia) (Ignatova, Blom, et al. 2010).

³³⁵*Schistidium convergens* was recently described as a new species from southern Spain and Morocco (Guerra, Jiménez-Martínez, Cano, Alonso, et al. 2019).

³³⁶*Schistidium echinatum* was described from Russia by Ignatova, Blom, et al. (2010), and has also now been found in France and Austria.

³³⁷*Schistidium marginale* was described from Austria by Blom et al. (2016), and has since been found in several montane regions in Europe.

³³⁸*Schistidium obscurum* was described from Austria by Ignatova, Blom, et al. (2010). Elsewhere in Europe it has also been found in Switzerland, the Caucasus and Svalbard.

³³⁹*Schistidium sibiricum* was described from Russia by Ignatova, Blom, et al. (2010).

³⁴⁰*Schistidium sinensiapocarpum* was first reported in Europe from the Caucasus (Blom et al. 2006), and also occurs in the Austrian Alps (Köckinger, Suanjak, et al. 2008).

³⁴¹*Schistidium subconfertum* is primarily an Asian species. However, a herbarium specimen was recently discovered from the Italian Alps (Italy: Stifserjochhöhe, 26 August 1909, Glowacki, GJO), and examined by H. Blom. This species is here published new to Europe.

³⁴²*Schistidium subflaccidum* was reported from the Caucasus in Russia (Blom et al. 2006), and has subsequently been found in mountainous regions of other parts of Europe.

³⁴³*Schistidium succulentum* was described from Russia by Ignatova, Blom, et al. (2010).

³⁴⁴*Schistidium tenuinerve* was described from Asiatic Russia (Ignatova, Blom, et al. 2010), but was recently also recorded from European Russia (E. Ignatova pers. comm. 2018).

- 1 ***B. alopecura*** (Brid.) Limpr.
 2 ***B. imberbis*** (Sm.) N.Dalton & D.G.Long [*Hedwigidium integrifolium* auct., non (P.Beauv.) Dixon, *Hedwigia integrifolia* auct., non P.Beauv.]³⁴⁵
- 124 ***Hedwigia*** P.Beauv.
 1 ***H. ciliata*** (Hedw.) P.Beauv.
 2 ***H. emodica*** Hampe ex Müll.Hal. [*Hedwigia ciliata* var. *leucophaea* Bruch & Schimp.]³⁴⁶
 3 ***H. mollis*** Ignatova, Ignatov & Fedosov³⁴⁷
 4 ***H. nemoralis*** Ignatova, Ignatov & Fedosov³⁴⁸
 5 ***H. stellata*** Hedenäs
 6 ***H. striata*** (Bruch & Schimp.) John Whitehead & Fergusson ex Hobk. & Porritt³⁴⁹
- Bartramiales D.Quandt, N.E.Bell & M.Stech
 Bartramiaceae Schwägr.
 Conostomoideae D.G.Griffin & W.R.Buck
 125 ***Conostomum*** Sw.
 1 ***C. tetragonum*** (Hedw.) Lindb.
- Bartramioideae D.G.Griffin & W.R.Buck
 126 ***Anacolia*** Schimp.
 1 ***A. menziesii*** (Turner) Paris
 2 ***A. webbii*** (Mont.) Schimp.
- 127 ***Bartramia*** Hedw.
 section *Bartramia*
 1 ***B. halleriana*** Hedw.
 2 ***B. pomiformis*** Hedw.
 section *Pyridium* Müll.Hal.
 3 ***B. breviseta*** Lindb.
 4 ***B. ithyphylla*** Brid.
 5 ***B. subulata*** Bruch & Schimp.
 section *Strictidium* Müll.Hal.
 6 ***B. aprica*** Müll.Hal. [*Bartramia rosamrosiae* Damayanti, J.Muñoz, J.-P.Frahm & D.Quandt, *Bartramia stricta* auct. eur., non Brid.]³⁵⁰
 7 ***B. laevisphaera*** (Taylor) Müll.Hal. [*Anacolia laevisphaera* (Taylor) Flowers]³⁵¹
- 128 ***Plagiopus*** Brid.
 1 ***P. oederianus*** (Sw.) H.A.Crum & L.E.Anderson
 a var. ***alpinus*** (Schwägr.) Ochyra³⁵²
 b var. ***oederianus***
- Breutelioideae D.G.Griffin & W.R.Buck
 129 ***Breutelia*** (Bruch & Schimp.) Schimp.
 1 ***B. azorica*** (Mitt.) Cardot
 2 ***B. chrysocoma*** (Hedw.) Lindb.
- 130 ***Philonotis*** Brid.
 section *Bartramidula* (Bruch & Schimp.) Mitt.
 1 ***P. cernua*** (Wilson) D.G.Griffin & W.R.Buck
 section *Philonotula* (Schimp.) A.Jaeger
 2 ***P. rigida*** Brid.
 section *Homomorphae* (Kindb.) Ochyra
 3 ***P. capillaris*** Lindb. [*Philonotis arnellii* Husn.]³⁵³
 4 ***P. falcata*** (Hook.) Mitt.³⁵⁴
 5 ***P. hastata*** (Duby) Wijk & Margad.
 6 ***P. marchica*** (Hedw.) Brid.
 7 ***P. uncinata*** (Schwägr.) Brid.
 section *Philonotis*
 8 ***P. caespitosa*** Jur.
 9 ***P. calcarea*** (Bruch & Schimp.) Schimp.
 10 ***P. fontana*** (Hedw.) Brid.
 11 ***P. seriata*** Mitt.
 12 ***P. tomentella*** Molendo
 13 ***P. yezoana*** Besch. & Cardot³⁵⁵
- Splachnales Ochyra
 Splachnaceae Grev. & Arn.
 Taylorioideae Broth.
 131 ***Tayloria*** Hook.
 1 ***T. acuminata*** Hornsch.
 2 ***T. froelichiana*** (Hedw.) Mitt. ex Broth.
 3 ***T. hornschurchii*** (Grev. & Arn.) Broth.
 4 ***T. lingulata*** (Dicks.) Lindb.
 5 ***T. rudolphiana*** (Garov.) Bruch & Schimp.
 6 ***T. serrata*** (Hedw.) Bruch & Schimp.
 7 ***T. splachnoides*** (Schleich. ex Schwägr.) Hook.
 8 ***T. tenuis*** (Dicks.) Schimp.

³⁴⁵Dalton et al. (2012) found that European specimens identified as *Hedwigidium integrifolium* differed from the type material of *Hedwigia integrifolia* P.Beauv., and were considered to belong to *Braunia imberbis*. De Luna (2016) reported that *Braunia imberbis*, along with some other (non-European) species, group together in a different clade, *Hedwigidium*. Until further research is done on this group, the species is here retained in *Braunia*.

³⁴⁶Hedenäs (1994) suggested that *Hedwigia emodica* was a synonym of *Hedwigia ciliata* var. *leucophaea*. Buchbender et al. (2014) later proved the hybrid origin of some accessions referable to this taxon. Ignatova, Kuznetsova, et al. (2016) recognised the taxon at species level, as *Hedwigia emodica*, but doubts persist that all European material is the same as the plants from Russia (or the Himalayan type of *Hedwigia emodica*).

³⁴⁷Ignatova, Kuznetsova, et al. (2016) described *Hedwigia mollis* as part of their study of the genus in Russia.

³⁴⁸*Hedwigia nemoralis* was described from Russia by Ignatova, Kuznetsova, et al. (2016).

³⁴⁹*Hedwigia striata* was a neglected taxon returned to specific rank by Buchbender et al. (2014).

³⁵⁰Damayanti et al. (2012) described *Bartramia rosamrosiae* for the European plants after finding that *Bartramia stricta* was restricted to South America. This was synonymised with *Bartramia aprica* by Müller (2014). However, recent (as yet unpublished) work by Neumann, Muñoz and Quandt suggests that *Bartramia aprica* and *Bartramia rosamrosiae* are indeed distinct species.

³⁵¹Molecular analysis revealed that *Bartramia laevisphaera* belongs to *Bartramia* section *Strictidium* rather than to *Anacolia* (Damayanti et al. 2012).

³⁵²There is disagreement over the status of *Plagiopus oederianus* var. *alpinus*: it may be nothing more than a high-altitude form of the species, but is retained for the present.

³⁵³*Philonotis capillaris* is the correct name for *Philonotis arnellii* (Koponen and Isoviita 2010).

³⁵⁴*Philonotis falcata* is an Asian species reported from the Caucasus new to Europe by Ignatov, Fedosov, et al. (2010).

³⁵⁵*Philonotis yezoana* is an Asian species reported from Finland new to Europe by Ulvinen and Kypärä (2016).

Splachnoideae Broth.

132 **Aplodon** R.Br.

- 1 **A. wormskioldii** (Hornem.) R.Br.

133 **Splachnum** Hedw.

- 1 **S. ampullaceum** Hedw.
- 2 **S. luteum** Hedw.
- 3 **S. melanocaulon** (Wahlenb.) Schwägr.
- 4 **S. pensylvanicum** (Brid.) Grout ex H.A.Crum
- 5 **S. rubrum** Hedw.
- 6 **S. sphaericum** Hedw.
- 7 **S. vasculosum** Hedw.

134 **Tetraplodon** Bruch & Schimp.

- 1 **T. angustatus** (Hedw.) Bruch & Schimp.
- 2 **T. blyttii** Frisvoll
- 3 **T. mnioides** (Hedw.) Bruch & Schimp.
- 4 **T. pallidus** I.Hagen
- 5 **T. paradoxus** (R.Br.) I.Hagen
- 6 **T. urceolatus** (Hedw.) Bruch & Schimp.

135 **Voitia** Hornsch.

- 1 **V. hyperborea** Grev. & Arn.
- 2 **V. nivalis** Hornsch.

Meesiaceae Schimp.

136 **Amblyodon** P.Beauv.

- 1 **A. dealbatus** (Hedw.) P.Beauv.

137 **Leptobryum** (Bruch & Schimp.) Wilson

- 1 **L. pyriforme** (Hedw.) Wilson

138 **Meesia** Hedw

- 1 **M. hexasticha** (Funck) Bruch
- 2 **M. longiseta** Hedw.
- 3 **M. triquetra** (L. ex Jolycl.) Ångstr.
- 4 **M. uliginosa** Hedw.

139 **Paludella** Brid.

- 1 **P. squarrosa** (Hedw.) Brid.

Bryales Limpr.

Bryaceae Schwägr.³⁵⁶140 **Anomobryum** Schimp.

- 1 **A. apiculatum** (Schwägr.) D.Bell & Holyoak [*Bryum apiculatum* Schwägr., *Gemmabryum apiculatum* (Schwägr.) J.R.Spence & H.P.Ramsay, *Osculatia apiculata* (Schwägr.) Ochyra, Plášek & Bedn.-Ochyra]³⁵⁷
- 2 **A. concinatum** (Spruce) Lindb.³⁵⁸
- 3 **A. julaceum** (Schrad. ex P.Gaertn., E.Mey & Scherb.) Schimp.
- 4 **A. lusitanicum** (I.Hagen) Thér.
- 5 **A. notarisii** (Mitt.) D.Bell. & Holyoak [*Brachymenium notarisii* (Mitt.) A.J.Shaw]³⁵⁹

141 **Brachymenium** Schwägr.

- 1 **B. paradoxum** (Herzog) A.J.Shaw

142 **Bryum** Hedw. [*Gemmabryum* J.R.Spence & H.R.Ramsay, *Osculatia* De Not.]³⁶⁰

- 1 **B. argenteum** Hedw.*
- 2 **B. bavaricum** Warnst. [*Anomobryum bavarium* (Warnst.) Holyoak & Köckinger]³⁶¹
- 3 **B. blindii** Bruch & Schimp.
- 4 **B. canariense** Brid. [*Rosulabryum canariense* (Brid.) Ochyra]³⁶²
- 5 **B. caucasicum** (Schimp. ex Broth.) C.J.Cox & Hedd.*
- 6 **B. demaretianum** Arts [*Gemmabryum demaretianum* (Arts) J.R.Spence, *Osculatia demaretiana* (Arts) Ochyra, Plášek & Bedn.-Ochyra]
- 7 **B. dichotomum** Hedw. [*Bryum barnesii* J.B.Wood ex Schimp., *Gemmabryum barnesii* (J.B.Wood ex Schimp.) J.R.Spence, *Osculatia barnesii* (J.B.Wood ex Schimp.) Ochyra, Plášek & Bedn.-Ochyra, *Osculatia bicolor* (Dicks.) Ochyra, Plášek & Bedn.-Ochyra, *Gemmabryum dichotomum* (Hedw.) J.R.Spence & H.P.Ramsay, *Osculatia dichotoma*

³⁵⁶See note on Bryaceae in introduction.³⁵⁷The only confirmed European record of *Bryum apiculatum* appears to be a specimen from Tenerife (leg. E. Bourgeon s.n., BM), which was reidentified by Ochi (1972). That specimen may nevertheless need critical reappraisal since Ochi did not record *Bryum gemmiparum* which is locally frequent on Tenerife, and there have been no subsequent reports of *Bryum apiculatum* occurring there. Holyoak (2009) and in Blockeel et al. (2014) identified plants from Cornwall as this species and two additional localities have since been found in the same county. However, data from DNA barcoding markers and preliminary analyses of a substantial genomic dataset place Cornish specimens (Holyoak 09-42 and 07-15) in *Bryum* s.str., close to *Bryum dichotomum*, differing markedly from a specimen of *Bryum apiculatum* from Puerto Rico (R.D. Worthington 35254). The Cornish plants thus represent an undescribed taxon allied to *Bryum dichotomum*, an account of which is in preparation.³⁵⁸Holyoak and Köckinger (2010) supported treatment of *Anomobryum concinatum* as a distinct species from *Anomobryum julaceum* based on morphological characters. Data from DNA barcoding markers (D. Bell et al., *in litt.*) has subsequently shown that *Anomobryum concinatum* and *Anomobryum julaceum* form separate clades.³⁵⁹Recent molecular work has shown that *Brachymenium notarisii* is closely related to *Anomobryum lusitanicum*, and is therefore probably best treated in that genus (D.Bell et al. *in litt.*).³⁶⁰Species with adequate molecular data confirming their position in *Bryum* s.str. are annotated with an asterisk (*). The remaining species lack sufficiently informative data, but there are now relatively few remaining 'orphan' species in *Bryum*, several more having been transferred to *Ptychostomum* or *Imbricobryum* on the basis of very recent DNA evidence.³⁶¹Holyoak and Köckinger (2010) placed *Anomobryum bavaricum* in *Anomobryum* on the basis of morphology, but molecular data places it in *Bryum* s.str. (D.Bell et al., *in litt.*).³⁶²Results from DNA barcoding and nuclear genomic datasets (D. Bell et al., *in litt.*) show that *Bryum canariense* does not belong in *Bryum* s.str., but is part of a near-basal clade of Bryaceae allied to *Brachymenium* s.str. and *Leptostomopsis*. However, it is retained in *Bryum* here pending further investigation.

- (Hedw.) Ochyra, Plášek & Bedn.-Ochyra]³⁶³
- 8 ***B. dixonii*** Cardot ex W.E.Nicholson
- 9 ***B. dyffrynense*** Holyoak*
- 10 ***B. gemmiferum*** R.Wilczek & Demaret [*Gemmabryum gemmiferum* (R.Wilczek & Demaret) J.R.Spence, *Osculatia gemmifera* (R.Wilczek & Demaret) Ochyra, Plášek & Bedn.-Ochyra]*
- 11 ***B. gemmilucens*** R.Wilczek & Demaret [*Gemmabryum gemmilucens* (R.Wilczek & Demaret) J.R.Spence, *Osculatia gemmilucens* (R.Wilczek & Demaret) Ochyra, Plášek & Bedn.-Ochyra]*
- 12 ***B. gemmparum*** De Not. [*Imbribryum gemmparum* (De Not.) J.R.Spence]*
- 13 ***B. kikuyense*** (Broth. & Thér.) N.Pedersen [*Brachymenium philonotula* Broth., *Bryum philonotulum* Hampe nom. inval., non *Bryum philonotulum* Müll.Hal.]*³⁶⁴
- 14 ***B. klinggraeffii*** Schimp. [*Gemmabryum klinggraeffii* (Schimp.) J.R.Spence & H.P.Ramsay, *Osculatia klinggraeffii* (Schimp.) Ochyra, Plášek & Bedn.-Ochyra]³⁶⁵
- 15 ***B. marratii*** Hook.f. & Wilson
- 16 ***B. oblongum*** Lindb.³⁶⁶
- 17 ***B. radiculosum*** Brid. [*Gemmabryum radiculosum* (Brid.) J.R.Spence & H.P. Ramsay, *Osculatia radiculosa* (Brid.) Ochyra, Plášek & Bedn.-Ochyra]*
- 18 ***B. reyeri*** Breidl.³⁶⁷
- 19 ***B. riparium*** I.Hagen
- 20 ***B. ruderale*** Crundw. & Nyholm [*Gemmabryum ruderale* (Crundw. & Nyholm) J.R.Spence, *Osculatia ruderalis* (Crundw. & Nyholm) Ochyra, Plášek & Bedn.-Ochyra]*
- 21 ***B. sauteri*** Bruch & Schimp. [*Osculatia sauteri* (Bruch & Schimp.) Ochyra, Plášek & Bedn.-Ochyra]
- 22 ***B. valparaisense*** Thér. [*Gemmabryum valparaisense* (Thér.) J.R.Spence, *Osculatia valparaisense* (Thér.) Ochyra, Plášek & Bedn.-Ochyra]
- 23 ***B. versicolor*** A.Braun ex Bruch & Schimp.³⁶⁸
- 24 ***B. violaceum*** Crundw. & Nyholm [*Gemmabryum violaceum* (Crundw. & Nyholm) J.R.Spence, *Osculatia violacea* (Crundw. & Nyholm) Ochyra, Plášek & Bedn.-Ochyra]*
- 143 ***Imbribryum*** Pedersen³⁶⁹
- 1 ***I. alpinum*** (Huds. ex With.) N.Pedersen [*Bryum alpinum* Huds. ex With.]
- 2 ***I. mildeanum*** (Jur.) J.R.Spence [*Bryum mildeanum* Jur.]
- 3 ***I. miniatum*** (Lesq.) J.R.Spence [*Bryum miniatum* Lesq.]³⁷⁰
- 4 ***I. muehlenbeckii*** (Bruch & Schimp.) N.Pedersen [*Bryum muehlenbeckii* Bruch & Schimp.]³⁷¹
- 5 ***I. subapiculatum*** (Hampe) D.Bell & Holyoak [*Bryum subapiculatum* Hampe, *Gemmabryum subapiculatum* (Hampe) J.R.Spence & H.P.Ramsay, *Osculatia subapiculata* (Hampe) Ochyra, Plášek & Bedn.-Ochyra]³⁷²
- 6 ***I. tenuisetum*** (Limpr.) D.Bell & Holyoak [*Bryum tenuisetum* Limpr., *Gemmabryum tenuisetum* (Limpr.) J.R.Spence & H.P. Ramsay, *Osculatia tenuiseta* (Limpr.) Ochyra, Plášek & Bedn.-Ochyra]³⁷³

³⁶³*Bryum barnesii* is usually regarded as a distinct species in Central Europe, but many plants and populations intermediate between *Bryum barnesii* and *Bryum dichotomum* occur (Holyoak 2003).

³⁶⁴Molecular data show that *Brachymenium philonotula* Broth. should be placed in *Bryum* rather than *Brachymenium*, but since the name *Bryum philonotula* Müll.Hal. (≡ *Pohlia philonotula* (Müll.Hal.) Broth.) had already been used, the valid name for the species becomes *Bryum kikuyense* (Pedersen and Hedenäs 2005).

³⁶⁵According to molecular evidence from Holyoak and Pedersen (2007), *Bryum klinggraeffii* could possibly be placed in *Imbribryum* or its own genus, but is retained in *Bryum* for the present.

³⁶⁶Preliminary molecular data suggest that *Bryum oblongum* may belong to *Ptychostomum* s. lat, but further study is needed to confirm this.

³⁶⁷*Bryum reyeri* is only known from the eastern Italian Alps (type), Austria and Germany (Limpricht 1895; Grims 1999; Meinunger and Schröder 2007; Köckinger, Suanjak, et al. 2008). Provisional data from a nuclear DNA dataset (D. Bell et al., in litt.) imply it is part of a clade basal to the radiation of *Ptychostomum*, as circumscribed here.

³⁶⁸*Bryum versicolor* was included as a synonym of *Bryum dichotomum* by Hill et al. (2006) but this is a widely recognised taxon, so it is reinstated pending further studies.

³⁶⁹*Imbribryum* was recognised at generic level by Holyoak and Pedersen (2007), and new molecular data from a substantial nuclear genomic dataset supports this view (D. Bell et al., in litt.).

³⁷⁰*Imbribryum miniatum* is placed in *Imbribryum* based on molecular data from specimens from both the Faeroes and the U.S.A.

³⁷¹Holyoak (in Hill et al. 2006) synonymised *Bryum muehlenbeckii* with *Bryum alpinum* because it appeared to be only a form resulting from growth under water or beneath snow, and there are numerous intermediate specimens. However, DNA sequence data for *Imbribryum muehlenbeckii* from three exemplars from different localities define a single clade that is well separated from *Imbribryum alpinum*. Hence, they are regarded here as separate species in which occasional apparent morphological intergradation is due to the occurrence of phenotypes ('expressions') that match the appearance of the other taxon. Additional molecular data suggest that it may be better treated in *Ptychostomum*, but further work is needed before this can be confirmed.

³⁷²DNA barcode data show *Bryum subapiculatum* strongly supported in a clade with *Imbribryum alpinum* (D. Bell et al., in litt.), and it is therefore transferred to *Imbribryum*.

³⁷³Along with *Bryum subapiculatum*, DNA barcoding places *Bryum tenuisetum* in a clade with *Imbribryum alpinum* (D. Bell et al., in litt.), so it is here transferred to *Imbribryum*.

- 144 ***Ptychostomum*** Hornsch. [*Plagiobryoides* J.R.Spence, *Plagiobryum* Lindb., *Rosulabryum* J.R.Spence]³⁷⁴
- 1 ***P. arcticum*** (R.Br.) J.R.Spence ex Holyoak & N.Pedersen [*Bryum arcticum* (R.Br.) Bruch & Schimp.]
 - 2 ***P. austriacum*** (Köckinger, Holyoak & Suanjak) D.Bell & Holyoak [*Bryum austriacum* Köckinger, Holyoak & Suanjak]³⁷⁵
 - 3 ***P. bornholmense*** (Wink. & R.Ruthe) Holyoak & N.Pedersen [*Bryum bornholmense* Wink. & R.Ruthe, *Osculatia bornholmensis* (Wink. & R.Ruthe) Ochyra, Plášek & Bedn.-Ochyra, *Rosulabryum bornholmense* (Wink. & Ruthe) J.R.Spence]
 - 4 ***P. calophyllum*** (R.Br.) J.R.Spence [*Bryum calophyllum* R.Br.]³⁷⁶
 - 5 ***P. capillare*** (Hedw.) Holyoak & N.Pedersen [*Rosulabryum capillare* (Hedw.) J.R.Spence, *Bryum capillare* Hedw.]
 - 6 ***P. cellulare*** (Hook.) D.Bell & Holyoak [*Bryum cellulare* Hook., *Plagiobryoides cellularis* (Hook.) J.R.Spence]
 - 7 ***P. cernuum*** (Hedw.) Hornsch. [*Bryum uliginosum* (Brid.) Bruch & Schimp.]
 - 8 ***P. compactum*** Hornsch. [*Bryum algovicum* Sendtn. ex Müll.Hal., *Bryum compactum* (Hornsch.) Kindb.]
 - a var. ***compactum***
 - b var. ***rutheanum*** (Warnst.) Holyoak & N.Pedersen [*Bryum algovicum* var. *rutheanum* (Warnst.) Crundw.]
 - 9 ***P. creberrimum*** (Taylor) J.R.Spence & H.P.Ramsay [*Bryum creberrimum* Taylor]
 - 10 ***P. cryophilum*** (Mårtensson) J.R.Spence [*Bryum cryophilum* Mårtensson]³⁷⁷
 - 11 ***P. cyclophyllum*** (Schwägr.) J.R.Spence [*Bryum cyclophyllum* (Schwägr.) Bruch & Schimp.]
 - 12 ***P. demissum*** (Hook.) Holyoak & N.Pedersen [*Plagiobryum demissum* (Hook.) Lindb.]
 - 13 ***P. donianum*** (Grev.) Holyoak & N.Pedersen [*Bryum donianum* Grev., *Rosulabryum donianum* (Grev.) Ochyra]
 - 14 ***P. elegans*** (Nees) D.Bell & Holyoak [*Bryum elegans* Nees, *Rosulabryum elegans* (Nees) Ochyra]³⁷⁸
 - 15 ***P. funkii*** (Schwägr.) J.R.Spence [*Bryum funkii* Schwägr.]³⁷⁹
 - 16 ***P. imbricatulum*** (Müll.Hal.) Holyoak & N.Pedersen [*Bryum caespiticium* Hedw., *Gemmabryum caespiticium* (Hedw.) J.R.Spence, *Osculatia caespiticia* (Hedw.) Ochyra, Plášek & Bedn.-Ochyra]
 - 17 ***P. inclinatum*** (Sw. ex Brid.) J.R.Spence [*Bryum archangelicum* Bruch & Schimp., *Ptychostomum archangelicum* (Bruch & Schimp.) J.R.Spence]³⁸⁰
 - 18 ***P. intermedium*** (Brid.) J.R.Spence [*Bryum intermedium* (Brid.) Blandow]³⁸¹
 - 19 ***P. knowltonii*** (Barnes) J.R.Spence [*Bryum knowltonii* Barnes]³⁸²
 - 20 ***P. kunzei*** (Hornsch.) J.R. Spence [*Bryum kunzei* Hornsch., *Gemmabryum kunzei* (Hornsch.) J.R.Spence,

³⁷⁴*Ptychostomum* was recognised at generic level by Holyoak and Pedersen (2007). There are no molecular data on the type species of *Rosulabryum*, which is *Rosulabryum albolimbatum* (Hampe) J.R.Spence from Australia. Recent analyses of a nuclear genomic dataset including the rather similar *Rosulabryum billardierei* (Schwägr.) J.R.Spence from the Southern Hemisphere (type from Tasmania), based on a specimen from Ecuador, imply that it forms part of *Brachymenium* s.str. Hence, further study is desirable before placing European and N. American species in *Rosulabryum*.

³⁷⁵*Bryum austriacum* was described from the Alps by Köckinger, Holyoak, et al. (2013). DNA barcode data places it in *Ptychostomum* (D. Bell et al., *in litt.*).

³⁷⁶Results from DNA barcoding (D. Bell et al. *in litt.*) show that *Bryum calophyllum* belongs in *Ptychostomum*.

³⁷⁷DNA barcoding results place *Bryum cryophilum* in *Ptychostomum* (D. Bell et al., *in litt.*).

³⁷⁸Morphological similarities have led authors of numerous floras (e.g. Schimper 1876; Smith 2004; Spence 2014) to place *Bryum elegans* close alongside *Bryum capillare*. Caution induced by lack of molecular data on *elegans* nevertheless prevented Holyoak and Pedersen (2007) transferring the species to *Ptychostomum* alongside *Ptychostomum capillare*, and its treatment in *Bryum* was followed by Ros, Mazimpaka, et al. (2013) and Hodgetts (2015). However, the molecular-phylogenetic study by Guerra, Jiménez-Martínez, Cano, Jiménez-Fernández, (2011) revealed *Bryum elegans* as closer to *Bryum pallescens* (i.e. *Ptychostomum pallescens*) than other Bryaceae they analysed, justifying the new combination that is made here. Newly generated nuclear genomic data also place *elegans* as part of *Ptychostomum* s.lat., but outside the (sub-)clade containing *P. capillare*. Spence (2014) has treated both species within the genus *Rosulabryum*, but Pedersen et al. (2003) and Shaw (2014) noted that species previously attributed by Spence to *Rosulabryum* were placed in two different clades in a phylogenetic tree based on molecular data. For this reason and because of continuing doubts about the affinities of the type species *Rosulabryum albolimbatum*, *Rosulabryum* is treated here as a synonym of the more widely inclusive genus *Ptychostomum*.

³⁷⁹Results from DNA barcoding (D. Bell et al., *in litt.*) show that *Bryum funkii* belongs in *Ptychostomum*. The spelling '*funkii*' was used in Index Muscorum, but it was changed to '*funckii*' because it commemorates Heinrich Christian Funck (1771-1839). Although at first sight this appears to be correction of an orthographical error (under ICBN Art. 60.1) and congruent with the orthography of *Marsupella funckii* and *Funckia* named in honour of the same botanist, Funk or Funck himself used both spellings of his name (biographer E. Hertel, pers. comm. to H. Köckinger), so there is no need to change established usage.

³⁸⁰Ochyra, Lewis Smith, et al. (2008) pointed out that "the epithet *inclinatum* is still available for *Ptychostomum imbricatulum* when it is considered a member of *Ptychostomum*". Since *Pohlia inclinata* Sw. ex Brid. 1803 is the oldest legitimate name available for the species it must therefore be adopted when the species is transferred to *Ptychostomum*, whereas its usage in *Bryum* was blocked by the later homonym *Bryum inclinatum* (Hedw.) Dicks. of 1801 (now *Distichium inclinatum*). Spence (2005) used the epithet *inclinatum* under *Ptychostomum*, but this new combination was invalid since the basionym was incorrectly cited (contrary to ICBN Art. 33.4). Hence, it was not until publication of Spence (2014) that *Ptychostomum inclinatum* (Sw. ex Brid.) J.R.Spence was validated in accordance with ICBN Art. 33.4 and Art. 45.1.

³⁸¹The placing of *Bryum intermedium* in *Ptychostomum* is supported by sequence data from DNA barcoding markers (D. Bell et al., *in litt.*).

³⁸²The placing of *Bryum knowltonii* in *Ptychostomum* is supported by sequence data from DNA barcoding markers (D. Bell et al., *in litt.*).

- Osculatia kunzei* (Hornsch.) Ochyra, Plášek & Bedn.-Ochyra]³⁸³
- 21 ***P. longisetum*** (Blandow ex Schwägr.) J.R.Spence [*Bryum longisetum* Blandow ex Schwägr.]
- 22 ***P. minii*** (Podp. ex Guim.) D.Bell & Holyoak [*Bryum minii* Podp. ex Guim.]
- 23 ***P. moravicum*** (Podp.) Ros & Mazimpaka [*Bryum moravicum* Podp., *Rosulabryum laevifilum* (Syed) Ochyra]
- 24 ***P. pallens*** (Sw. ex anon.) J.R.Spence [*Bryum pallens* Sw. ex anon., *Bryum sibiricum* Lindb. & Arnell]³⁸⁴
- 25 ***P. pallescens*** (Schleich. ex Schwägr.) J.R.Spence [*Bryum pallescens* Schleich. ex Schwägr., *Ptychostomum boreale* (F.Weber & D.Mohr) Ochyra & Bedn.-Ochyra]³⁸⁵
- 26 ***P. pseudotriquetrum*** (Hedw.) J.R.Spence & H.P.Ramsay ex Holyoak & N.Pedersen [*Bryum neodamense* Itzigs., *Bryum pseudotriquetrum* (Hedw.) P.Gaertn. B.Mey. & Scherb. *Ptychostomum neodamense* (Itzigs.) J.R.Spence]³⁸⁶
- a var. ***bimum*** (Schreb.) Holyoak & N.Pedersen [*Bryum pseudotriquetrum* var. *bimum* (Schreb.) Lilj.]
- b var. ***pseudotriquetrum***
- 27 ***P. rubens*** (Mitt.) Holyoak & N.Pedersen [*Bryum rubens* Mitt., *Osculatia rubens* (Mitt.) Ochyra, Plášek & Bedn.-Ochyra, *Rosulabryum rubens* (Mitt.) J.R.Spence]
- 28 ***P. salinum*** (I.Hagen ex Limpr.) J.R.Spence [*Bryum salinum* I.Hagen ex Limpr.]³⁸⁷
- 29 ***P. schleicheri*** (DC.) J.R.Spence ex D.Bell & Holyoak [*Bryum schleicheri* DC.]³⁸⁸
- a var. ***latifolium*** (Schwägr.) D.Bell & Holyoak [*Bryum schleicheri* var. *latifolium* (Schwägr.) Schimp.]
- b var. ***schleicheri***
- 30 ***P. torquescens*** (Bruch & Schimp.) Ros & Mazimpaka [*Bryum torquescens* Bruch & Schimp., *Rosulabryum torquescens* (Bruch & Schimp.) J.R.Spence]
- 31 ***P. turbinatum*** (Hedw.) J.R.Spence [*Bryum turbinatum* (Hedw.) Turner]³⁸⁹
- 32 ***P. warneum*** (Röhl.) J.R.Spence [*Bryum warneum* (Röhl.) Brid.]³⁹⁰
- 33 ***P. weigelii*** (Biehler) J.R.Spence [*Bryum weigelii* Biehler]³⁹¹
- 34 ***P. wrightii*** (Sull. & Lesq.) J.R.Spence [*Bryum wrightii* Sull. & Lesq.]³⁹²
- 35 ***P. zieri*** (Hedw.) Holyoak & N.Pedersen [*Plagiobryum zieri* (Hedw.) Lindb.]
- 145 ***Rhodobryum*** (Schimp.) Limpr.
- 1 ***R. ontariense*** (Kindb.) Kindb.
- 2 ***R. roseum*** (Hedw.) Limpr.
- Mniaceae Schwägr.
- Mielichhoferioideae M.Stech & W.Frey
- 146 ***Mielichhoferia*** Nees & Hornsch.
- 1 ***M. elongata*** (Hoppe & Hornsch. ex Hook.) Hornsch.
- 2 ***M. mielichhoferiana*** (Funck) Loeske

³⁸³DNA barcoding results suggest that *Bryum kunzei* should be transferred to *Ptychostomum* (D. Bell et al., *in litt.*).

³⁸⁴Zolotov (2007a) regarded *Bryum sibiricum* as possibly "a good, but very rare species", with the comment that "it is rather contrastingly different from *B. pallens* which is relatively little variable in gametophytic characters", all this on the basis of study of a single specimen placed as *Bryum sibiricum* from Yakutia because it matched the original description well. Zolotov (2007b) noted it as close *Bryum pallens*, but "differs ... [in the] very narrow leaves". Zolotov made no mention though of the great variability of *Ptychostomum pallens* revealed in the extensive illustrated monograph of the species by Podpěra (1973), which included the new combination *Bryum pallens* subsp. *sibiricum* (Lindb. & Arnell) Podp. (p. 195), redescribed in detail. Other forms of *Bryum pallens* with long narrow leaves and excurrent costa were figured and described in Podpěra's work, including subsp. *prosboreum* Podp. (from arctic Sweden), subsp. *acutiusculum* Podp. (Alaska) and forma *gracile* Podp. (Bohemia), as well as other infraspecific taxa intermediate between these and normal *Ptychostomum pallens*.

³⁸⁵Ochyra & Bednarek-Ochyra (2011) demonstrated that *Hypnum boreale* F.Weber & D.Mohr, 1807, and the subsequent combination *Bryum boreale* (F.Weber & D.Mohr) Funck, 1808, are older names for *Ptychostomum pallescens* than *Bryum pallescens* Schwägr., 1816. Ochyra and Bednarek-Ochyra (2015) published a proposal to maintain established usage of the name *Bryum pallescens* in preference to the unfamiliar and largely forgotten name *Bryum boreale* and this desirable initiative is supported here. The combination *Ptychostomum boreale* (F.Weber & D.Mohr) Ochyra & Bednarek-Ochyra, 2011 was valid, and adopted as correct by Ros, Mazimpaka, et al. (2013) and Hodgetts (2015). However, pending the response to the proposal by Ochyra and Bednarek-Ochyra (2015), the established usage should be maintained for *Ptychostomum pallescens*.

³⁸⁶*Ptychostomum neodamense* has been shown by Holyoak and Hedenäs (2006), from molecular and other data, to represent an inconstant phenotype of *Ptychostomum pseudotriquetrum* occurring locally in highly calcareous habitats prone to inundation, and connected to it by intermediate forms known widely in Europe and also in Asia, Alaska and Greenland. Its retention at species rank by Spence (2014) is a consequence of an over-emphasis on morphological data.

³⁸⁷*Bryum salinum* is very close to *Ptychostomum inclinatum* (Holyoak and Pedersen 2007), and this is supported by sequence data from DNA barcoding markers (D. Bell et al. *in litt.*).

³⁸⁸DNA barcoding results and data from a nuclear genomic dataset show that *Bryum schleicheri* is better placed in *Ptychostomum* (D. Bell et al., *in litt.*). However, relationships between the two varieties remain uncertain, as the two datasets are not directly comparable, and a study is needed to clarify the position of *Ptychostomum schleicheri* var. *latifolium*. As a large, swollen plant with concave leaves, it is the plant that is understood as *Ptychostomum schleicheri* in many parts of Europe; var. *schleicheri* is less well known, sometimes because it is misidentified as *Ptychostomum turbinatum*.

³⁸⁹Spence's placement of *Bryum turbinatum* in *Ptychostomum* is supported by molecular evidence (D. Bell et al., *in litt.*).

³⁹⁰Spence's placement of *Bryum warneum* in *Ptychostomum* is supported by molecular evidence (D. Bell et al., *in litt.*).

³⁹¹*Ptychostomum weigelii* has been considered an invalid combination since Spence (2005) cited Sprengel (1807, July) as the author of the basionym name *Bryum weigelii*, when the name was in fact published validly by Biehler (1807, May) a couple of months earlier. However, since Spence simply cited the wrong publication, and did not omit any clear reference to the basionym, this is a correctable error (cf. ICBN Art. 41.8(a); Turland et al. 2018).

³⁹²*Bryum wrightii* was placed in *Ptychostomum* by Holyoak and Pedersen (2007), following Pedersen et al. (2003).

- 147 **Pohlia** Hedw.
- 1 ***P. andalusica*** (Höhn.) Broth.
 - 2 ***P. andrewsii*** A.J.Shaw
 - 3 ***P. annotina*** (Hedw.) Lindb.
 - 4 ***P. atropurpurea*** (Wahlenb.) H.Lindb.
 - 5 ***P. beringiensis*** A.J.Shaw³⁹³
 - 6 ***P. bolanderi*** (Lesq.) Broth.
 - 7 ***P. bulbifera*** (Warnst.) Warnst.
 - 8 ***P. camptotrachela*** (Renauld & Cardot) Broth.
 - 9 ***P. cruda*** (Hedw.) Lindb.
 - 10 ***P. crudoides*** (Sull. & Lesq.) Broth.
 - 11 ***P. drummondii*** (Müll.Hal.) A.L.Andrews
 - 12 ***P. elongata*** Hedw.
 - a var. ***acuminata*** (Hornsch.) Huebener
 - b var. ***elongata***
 - c var. ***greenii*** (Brid.) A.J.Shaw
 - 13 ***P. erecta*** Lindb.
 - 14 ***P. filum*** (Schimp.) Mårtensson
 - 15 ***P. flexuosa*** Harv.
 - a var. ***flexuosa***
 - b var. ***pseudomuyldermansii*** (Arts, Nordhorn-Richter & A.J.E.Sm.) A.J.E.Sm.
 - 16 ***P. lescuriana*** (Sull.) Ochi
 - 17 ***P. longicolla*** (Hedw.) Lindb.
 - 18 ***P. ludwigii*** (Spreng. ex Schwägr.) Broth.
 - 19 ***P. lutescens*** (Limpr.) H.Lindb.
 - 20 ***P. melanodon*** (Brid.) A.J.Shaw
 - 21 ***P. nutans*** (Hedw.) Lindb.
 - a subsp. ***nutans***
 - b subsp. ***schimperii*** (Müll.Hal.) Nyholm
 - 22 ***P. obtusifolia*** (Vill. ex Brid.) L.F.Koch
 - 23 ***P. prolifera*** (Kindb.) Lindb. ex Broth.
 - 24 ***P. scotica*** Crundw.
 - 25 ***P. sphagnicola*** (Bruch & Schimp.) Broth.
 - 26 ***P. tundrae*** A.J.Shaw
 - 27 ***P. vexans*** (Limpr.) H.Lindb.
 - 28 ***P. wahlenbergii*** (F.Weber & D.Mohr) A.L.Andrews
 - a var. ***calcarea*** (Warnst.) E.F.Warb.
 - b var. ***glacialis*** (Brid.) E.F.Warb.
 - c var. ***wahlenbergii***
- 148 **Schizymenium** Harv.
- 1 ***S. pontevedrense*** (Luisier) Sérgio Casas, Cros & Brugués
- Mnioideae L.Söderstr. & N.G.Hodgetts³⁹⁴
- 149 **Cinclidium** Sw.
- 1 ***C. arcticum*** (Bruch & Schimp.) Schimp.
 - 2 ***C. latifolium*** Lindb.
 - 3 ***C. minutifolium*** Broth.³⁹⁵
- 4 ***C. stygium*** Sw.
 - 5 ***C. subrotundum*** Lindb.
- 150 **Cyrtomnium** Holmen
- 1 ***C. hymenophylloides*** (Huebener) T.J.Kop.
 - 2 ***C. hymenophyllum*** (Bruch & Schimp.) Holmen
- 151 **Epipterygium** Lindb.
- 1 ***E. tozeri*** (Grev.) Lindb.
- 152 **Mnium** Hedw.
- 1 ***M. blyttii*** Bruch & Schimp.
 - 2 ***M. heterophyllum*** (Hook.) Schwägr.
 - 3 ***M. hornum*** Hedw.
 - 4 ***M. lycopodioides*** Schwägr.
 - 5 ***M. marginatum*** (Dicks.) P.Beauv.
 - a var. ***dioicum*** (H.Müll.) Crundw.
 - b var. ***marginatum***
 - 6 ***M. spinosum*** (Voit) Schwägr.
 - 7 ***M. spinulosum*** Bruch & Schimp.
 - 8 ***M. stellare*** Hedw.
 - 9 ***M. thomsonii*** Schimp.
- 153 **Plagiomnium** T.J.Kop. [*Orthomnion* Wils.]³⁹⁶
- section *Plagiomnium*
- 1 ***P. affine*** (Blandow ex Funck) T.J.Kop. [*Orthomnion affine* (Blandow ex Funck) T.J.Kop. & Yu Sun]
 - 2 ***P. curvatulum*** (Lindb.) Schljakov [*Orthomnion curvatulum* (Lindb.) T.J.Kop. & Yu Sun]
 - 3 ***P. cuspidatum*** (Hedw.) T.J.Kop. [*Orthomnion cuspidatum* (Hedw.) T.J.Kop. & Yu Sun]
 - 4 ***P. drummondii*** (Bruch & Schimp.) T.J.Kop. [*Orthomnion drummondii* (Bruch & Schimp.) T.J.Kop. & Yu Sun]
 - 5 ***P. elatum*** (Bruch & Schimp.) T.J.Kop. [*Orthomnion elatum* (Bruch & Schimp.) T.J.Kop. & Yu Sun]
 - 6 ***P. ellipticum*** (Brid.) T.J.Kop. [*Orthomnion ellipticum* (Brid.) T.J.Kop. & Yu Sun]
 - 7 ***P. medium*** (Bruch & Schimp.) T.J.Kop. [*Orthomnion medium* (Bruch & Schimp.) T.J.Kop. & Yu Sun]
- section *Undulata* (Hedw.) T.J.Kop.
- 8 ***P. confertidens*** (Lindb. & Arnell) T.J.Kop. [*Orthomnion confertidens* (Lindb. & Arnell) T.J.Kop. & Yu Sun]

³⁹³*Pohlia beringiensis* was reported from Arctic Russia new to Europe by Ignatov, Afonina and Ignatova (2006).

³⁹⁴The subfamily Mnioideae is in general usage but until now it never seems to have been validly published.

³⁹⁵*Cinclidium minutifolium* was reported from Arctic Russia new to Europe by Koponen and Ignatova (2018).

³⁹⁶The species of *Plagiomnium* were transferred to *Orthomnion* by Koponen and Sun (2017), but there is currently a proposal to conserve the name *Plagiomnium* (Ochyra, Bednarek-Ochyra, et al. 2017). Consequently *Plagiomnium* is retained.

- 9 ***P. undulatum*** (Hedw.) T.J.Kop. [*Orthomnion undulatum* (Hedw.) T.J.Kop. & Yu Sun]
 a var. ***madeirense*** T.J.Kop. & Sérgio [*Orthomnion undulatum* var. *madeirense* (T.J.Kop. & Sérgio) T.J.Kop. & Yu Sun]
 b var. ***undulatum***
 section *Rostrata* (Hedw.) T.J.Kop.
- 10 ***P. rostratum*** (Schrad.) T.J.Kop. [*Orthomnion rostratum* (Schrad.) T.J.Kop. & Yu Sun]
- 154 ***Pseudobryum*** (Kindb.) T.J.Kop.
 1 ***P. cinclidioides*** (Huebener) T.J.Kop.
- 155 ***Rhizomnium*** (Broth.) T.J.Kop.
 1 ***R. andrewsianum*** (Steere) T.J.Kop.
 2 ***R. gracile*** T.J.Kop.
 3 ***R. magnifolium*** (Horik.) T.J.Kop.
 4 ***R. pseudopunctatum*** (Bruch & Schimp.) T.J.Kop.
 5 ***R. punctatum*** (Hedw.) T.J.Kop.
 a var. ***hermanperssonii*** T.J.Kop.³⁹⁷
 b var. ***punctatum***
- 156 ***Trachycystis*** Lindb.
 1 ***T. ussuriensis*** (Maack & Regel) T.J.Kop.
- Orthotrichales Dixon
 Orthotrichaceae Arn.
 Orthotrichoideae Broth.
- 157 ***Codonoblepharon*** Schwägr.
 1 ***C. forsteri*** (Dicks.) Goffinet [*Zygodon forsteri* (Dicks.) Mitt.]³⁹⁸
- 158 ***Lewinskya*** F.Lara, Garilleti & Goffinet³⁹⁹
 1 ***L. acuminata*** (H.Philib.) F.Lara, Garilleti & Goffinet [*Orthotrichum acuminatum* H.Philib.]
 2 ***L. affinis*** (Schrad. ex Brid.) F.Lara, Garilleti & Goffinet [*Orthotrichum affine* Schrad. ex Brid., *Orthotrichum affine* var. *bohemicum* Plášek & Sawicki]
 3 ***L. breviseta*** (F.Lara, Garilleti & Mazimpaka) F.Lara, Garilleti & Goffinet [*Orthotrichum speciosum* var. *brevisetum* F.Lara, Garilleti & Mazimpaka]
- 4 ***L. elegans*** (Schwägr. ex Hook. & Grev.) F.Lara, Garilleti & Goffinet [*Orthotrichum elegans* Schwägr. ex Hook. & Grev.]⁴⁰⁰
- 5 ***L. fastigiata*** (Bruch ex Brid.) Vigalondo, F.Lara & Garilleti [*Orthotrichum fastigiatum* Bruch ex Brid.]⁴⁰¹
- 6 ***L. iberica*** (F.Lara & Mazimpaka) F.Lara, Garilleti & Goffinet [*Orthotrichum ibericum* F.Lara & Mazimpaka]
- 7 ***L. iwatsukii*** (Ignatov) F.Lara, Garilleti & Goffinet [*Orthotrichum iwatsukii* Ignatov, *Orthotrichum laevigatum* var. *japonicum* (Z.Iwats.) Lewinsky, *Orthotrichum macounii* subsp. *japonicum* Z.Iwats.]⁴⁰²
- 8 ***L. laevigata*** (J.E.Zetterst.) F.Lara, Garilleti & Goffinet [*Orthotrichum laevigatum* J.E.Zetterst.]
- 9 ***L. lamyana*** F.Lara, Garilleti, Draper & Mazimpaka⁴⁰³
- 10 ***L. pylaisii*** (Brid.) F.Lara, Garilleti & Goffinet [*Orthotrichum pylaisii* Brid.]
- 11 ***L. rupestris*** (Schleich. ex Schwägr.) F.Lara, Garilleti & Goffinet [*Orthotrichum rupestre* Schleich. ex Schwägr.]
- 12 ***L. shawii*** (Wilson) F.Lara, Garilleti & Goffinet [*Orthotrichum shawii* Wilson]
- 13 ***L. sordida*** (Sull. & Lesq.) F.Lara, Garilleti & Goffinet [*Orthotrichum sordidum* Sull. & Lesq.]
- 14 ***L. speciosa*** (Nees) F.Lara, Garilleti & Goffinet [*Orthotrichum speciosum* Nees]
- 15 ***L. striata*** (Hedw.) F.Lara, Garilleti & Goffinet [*Orthotrichum striatum* Hedw.]
- 16 ***L. tortidontia*** (F.Lara, Garilleti & Mazimpaka) F.Lara, Garilleti & Goffinet [*Orthotrichum tortidontium* F.Lara, Garilleti & Mazimpaka]
- 17 ***L. transcaucasica*** Eckstein, Garilleti & F.Lara⁴⁰⁴
- 18 ***L. vladikavkana*** (Venturi) F.Lara, Garilleti & Goffinet [*Orthotrichum vladikavkana* Venturi]
- 159 ***Nyholmiella*** Holmen & E.Warncke⁴⁰⁵

³⁹⁷*Rhizomnium punctatum* var. *hermanperssonii* was described from Madeira by Koponen (2017).

³⁹⁸Molecular evidence (Goffinet, Shaw, et al. 2004) suggests that *Zygodon forsteri* is best placed in *Codonoblepharon*. This is supported by current thinking (Mazimpaka and Lara 2014).

³⁹⁹*Lewinskya* was established as a genus to accommodate the monoicous and phaneroporos taxa of *Orthotrichum* (Lara, Garilleti, et al. 2016).

⁴⁰⁰Vitt and Darigo (1997) established that *Lewinskya elegans* was distinct from *Lewinskya speciosa* in North America. It is now known to be widespread in European Russia (Fedosov and Doroshina 2018), and probably overlooked elsewhere in Europe.

⁴⁰¹*Lewinskya fastigiata* was reinstated by Vigalondo, Garilleti, et al. (2019), on the basis of integrative taxonomic analyses on the *Lewinskya affinis* complex.

⁴⁰²*Lewinskya iwatsukii* is an Asian taxon described as a subspecies of *Orthotrichum macounii* Austin by Iwatuski (1959), raised to species level by Ignatov, Ivanova, et al. (2001), and later discovered in Europe (Fedosov and Doroshina 2018).

⁴⁰³*Lewinskya lamyana* was described from Spain by Lara, Garilleti, et al. (2018).

⁴⁰⁴*Lewinskya transcaucasica* is a recently described species (Eckstein et al. 2018), which has subsequently been found in the European Caucasus by Fedosov, Doroshina, et al. (2017).

⁴⁰⁵The recognition of *Nyholmiella* at generic level is supported by molecular evidence (Goffinet, Shaw, et al. 2004; Sawicki et al. 2010).

- 1 ***N. gymnostoma*** (Bruch ex Brid.) Holmen & E.Warncke [*Orthotrichum gymnostomum* Bruch ex Brid.]
- 2 ***N. obtusifolia*** (Brid.) Holmen & E.Warncke [*Orthotrichum obtusifolium* Brid.]
- 160 ***Orthotrichum*** Hedw.
- 1 ***O. alpestre*** Bruch & Schimp.
- 2 ***O. anomalum*** Hedw.
- 3 ***O. bistratosum*** (Schiffn.) Guerra [*Orthotrichum cupulatum* var. *bistratosum* Schiffn.]⁴⁰⁶
- 4 ***O. callistomum*** Fisch.-Oost. ex Bruch & Schimp.
- 5 ***O. cambrense*** Bosanquet & F.Lara⁴⁰⁷
- 6 ***O. casasianum*** F.Lara, Garilleti & Mazimpaka
- 7 ***O. columbicum*** Mitt. [*Orthotrichum consimile* auct. eur., non Mitt.]⁴⁰⁸
- 8 ***O. comosum*** F.Lara, R.Medina & Garilleti⁴⁰⁹
- 9 ***O. consobrinum*** Cardot⁴¹⁰
- 10 ***O. crenulatum*** Mitt.
- 11 ***O. cupulatum*** Brid.
- a var. ***cupulatum***
- b var. ***fuscum*** (Venturi) Boulay
- c var. ***riparium*** Huebener
- 12 ***O. dagestanicum*** Fedosov & Ignatova⁴¹¹
- 13 ***O. dentatum*** T.Kiebacher & Lüth⁴¹²
- 14 ***O. diaphanum*** Brid.
- 15 ***O. handiense*** F.Lara, Garilleti & Mazimpaka
- 16 ***O. hispanicum*** F.Lara, Garilleti & Mazimpaka
- 17 ***O. macrocephalum*** F.Lara, Garilleti & Mazimpaka
- 18 ***O. microcarpum*** De Not.
- 19 ***O. moravicum*** Plášek & Sawicki⁴¹³
- 20 ***O. pallens*** Bruch ex Brid.
- 21 ***O. patens*** Bruch ex Brid.
- 22 ***O. pellucidum*** Lindb.
- 23 ***O. philibertii*** Venturi
- 24 ***O. pulchellum*** Brunt.
- 25 ***O. pumilum*** Sw. ex anon.
- 26 ***O. rivulare*** Turner
- 27 ***O. rogeri*** Brid.
- 28 ***O. scanicum*** Grönvall
- 29 ***O. schimperi*** Hammar
- 30 ***O. shevockii*** Lewinsky-Haapasaari & D.H.Norris⁴¹⁴
- 31 ***O. sibiricum*** (Grönvall) Warnst. [*Orthotrichum holmenii* Lewinsky-Haapasaari]⁴¹⁵
- 32 ***O. sprucei*** Mont.
- 33 ***O. stellatum*** Brid.
- 34 ***O. stramineum*** Hornsch. ex Brid.
- 35 ***O. tenellum*** Bruch ex Brid.
- 36 ***O. urnigerum*** Myrin
- 37 ***O. vittii*** F.Lara, Garilleti & Mazimpaka
- 161 ***Plenogemma*** Plášek, Sawicki & Ochyra⁴¹⁶
- 1 ***P. phyllantha*** (Brid.) Sawicki, Plášek & Ochyra [*Ulota phyllantha* Brid.]
- 162 ***Pulviger*** Plášek, Sawicki & Ochyra⁴¹⁷
- 1 ***P. lyellii*** (Hook. & Taylor) Plášek, Sawicki & Ochyra [*Orthotrichum lyellii* Hook. & Taylor]
- 163 ***Ulota*** D.Mohr
- 1 ***U. bruchii*** Hornsch. ex Brid.
- 2 ***U. calvescens*** Wilson
- 3 ***U. coarctata*** (P.Beauv.) Hammar
- 4 ***U. crispa*** (Hedw.) Brid.⁴¹⁸
- 5 ***U. crispula*** Bruch [*Ulota crispa* var. *crispula* (Bruch) Hammar]
- 6 ***U. curvifolia*** (Wahlenb.) Lilj.
- 7 ***U. drummondii*** (Hook. & Grev.) Brid.
- 8 ***U. hutchinsiae*** (Sm.) Hammar
- 9 ***U. intermedia*** Schimp. [*Ulota crispa* var. *intermedia* (Schimp.) Cardot]
- 10 ***U. macrospora*** E.Bauer & Warnst.
- 11 ***U. rehmannii*** Jur.
- 164 ***Zygodon*** Hook. & Taylor
- 1 ***Z. catarinoi*** C.Garcia, F.Lara, Sérgio & Sim-Sim [*Zygodon bistratus* Calabrese & J.Muñoz]⁴¹⁹
- 2 ***Z. conoideus*** (Dicks.) Hook. & Taylor
- a var. ***conoideus***
- b var. ***lingulatus*** S.R.Edwards
- 3 ***Z. dentatus*** (Limpr.) Kartt.
- 4 ***Z. gracilis*** Wilson

⁴⁰⁶*Orthotrichum cupulatum* var. *bistratosum* was raised to species level by Guerra (1985), although it has been recognised as such only recently (Lara and Garilleti 2014).

⁴⁰⁷Bosanquet and Lara (2012) described *Orthotrichum cambrense* from Wales.

⁴⁰⁸*Orthotrichum columbicum* was recognised as a species distinct from *Orthotrichum consimile* Mitt. by Medina et al. (2012). *Orthotrichum consimile* s.str. is confined to the west coast of North America.

⁴⁰⁹*Orthotrichum comosum* was described from southern Europe by Medina et al. (2013).

⁴¹⁰*Orthotrichum consobrinum* is an Asian species detected in Europe (Spain) by Lara, Mazimpaka, et al. (2009).

⁴¹¹*Orthotrichum dagestanicum* was described from the Caucasus by Fedosov and Ignatova (2010).

⁴¹²*Orthotrichum dentatum* was described from the Alps by Kiebacher and Lüth (2017).

⁴¹³*Orthotrichum moravicum* was described from central Europe by Plášek et al. (2009).

⁴¹⁴*Orthotrichum shevockii* is a disjunct species described from California and recently found in the Canary Islands (Vigalondo, Patiño, et al. 2019).

⁴¹⁵*Orthotrichum sibiricum* is a north Asian species with one known occurrence in European Arctic Russia (Fedosov et al. 2017b).

⁴¹⁶Sawicki et al. (2017) provide molecular evidence for the treatment of Plášek et al. (2015), segregating *Ulota phyllantha* in the genus *Plenogemma*.

⁴¹⁷Sawicki et al. (2017) provide molecular evidence for the treatment of Plášek et al. (2015), segregating *Orthotrichum lyellii* in the genus *Pulviger*.

⁴¹⁸The treatment of the *Ulota crispa* complex (*Ulota crispa* s.str., *Ulota crispula*, *Ulota intermedia*) follows Caparrós et al. (2016).

⁴¹⁹*Zygodon catarinoi* was described from Spain, Portugal and Morocco by Garcia et al. (2006), and has subsequently been found elsewhere in the Mediterranean region.

- 5 **Z. rupestris** Schimp. ex Lorentz
- 6 **Z. sibiricus** Ignatov, Ignatova, Z.Iwats. & B.C.Tan
- 7 **Z. stirtonii** Schimp. ex Stirt.
- 8 **Z. viridissimus** (Dicks.) Brid.

Orthodontiales N.E.Bell, A.E.Newton & D.Quandt
Orthodontiaceae Goffinet

- 165 **Leptotheca** Schwägr.
 - 1 **L. gaudichaudii** Schwägr.
- 166 **Orthodontium** Schwägr.
 - 1 **O. gracile** (Wilson) Schwägr. ex Bruch & Schimp.
 - 2 **O. lineare** Schwägr.
 - 3 **O. pellucens** (Hook.) Bruch & Schimp.

Aulacomniales N.E.Bell, A.E.Newton & D.Quandt
Aulacomniaceae Schimp.

- 167 **Aulacomnium** Schwägr.
 - 1 **A. androgynum** (Hedw.) Schwägr.
 - 2 **A. palustre** (Hedw.) Schwägr.
 - 3 **A. turgidum** (Wahlenb.) Schwägr.

Rhizogoniales Goffinet & W.R.Buck
Rhizogoniaceae Broth.

- 168 **Calomnion** Hook.f. & Wilson
 - 1 **C. complanatum** (Hook.f. & Wilson) Lindb.

Hookeriales M.Fleisch. [Hypopterygiales Goffinet]
Hypopterygiaceae Mitt.

- 169 **Hypopterygium** Brid.
 - 1 **H. tamarisci** (Sw.) Brid. ex Müll.Hal.

Daltoniaceae Schimp.

- 170 **Achrophyllum** Vitt & Crosby
 - 1 **A. dentatum** (Hook.f. & Wilson) Vitt & Crosby

- 171 **Calyptrochaeta** Desv.
 - 1 **C. apiculata** (Hook.f. & Wilson) Vitt

- 172 **Daltonia** Hook. & Taylor
 - 1 **D. splachnoides** (Sm.) Hook. & Taylor
 - 2 **D. lindigiana** Hampe [*Daltonia stenophylla* Mitt.]⁴²⁰

- 173 **Distichophyllum** Dozy & Molk.
 - 1 **D. carinatum** Dixon & W.E.Nicholson

Hookeriaceae Schimp.

- 174 **Hookeria** J.E.Sm.
 - 1 **H. lucens** (Hedw.) Sm.

Leucomiaceae Broth.

- 175 **Tetrastichium** (Mitt.) Cardot
 - 1 **T. fontanum** (Mitt.) Cardot
 - 2 **T. virens** (Cardot) S.P.Churchill

Pilotrichaceae Kindb.

Hypnelloideae Broth.

- 176 **Cyclodictyon** Mitt.
 - 1 **C. laetevirens** (Hook. & Taylor) Mitt.

Hypnales W.R.Buck & Vitt

Fontinalaceae Schimp.

- 177 **Dichelyma** Myrin
 - 1 **D. capillaceum** (L. ex Dicks.) Myrin
 - 2 **D. falcatum** (Hedw.) Myrin
- 178 **Fontinalis** Hedw.
 - 1 **F. antipyretica** Hedw.
 - a subsp. **antipyretica**
 - b subsp. **bryhnii** (Limpr.) Podp.
 - c subsp. **gracilis** (Lindb.) Kindb.
 - d subsp. **kindbergii** (Renauld & Cardot) Cardot
 - 2 **F. dalecarlica** Schimp.
 - 3 **F. dichelymoides** Lindb.
 - 4 **F. hypnoides** C.Hartm.
 - a var. **duriaei** (Schimp.) Kindb.
 - b var. **hypnoides**
 - 5 **F. squamosa** Hedw. [*Fontinalis squamosa* var. *curnowii* Cardot, *Fontinalis squamosa* var. *dixonii* (Cardot) A.J.E.Sm.]⁴²¹

Plagiotheciaceae M.Fleisch.

- 179 **Herzogiella** Broth.
 - 1 **H. seligeri** (Brid.) Z.Iwats.
 - 2 **H. striatella** (Brid.) Z.Iwats.
 - 3 **H. turfacea** (Lindb.) Z.Iwats.

- 180 **Isopterygiopsis** Z.Iwats.
 - 1 **I. alpicola** (Lindb. & Arnell) Hedenäs
 - 2 **I. muelleriana** (Schimp.) Z.Iwats.
 - 3 **I. pulchella** (Hedw.) Z.Iwats.

- 181 **Plagiothecium** Bruch & Schimp. [*Buckiella* Ireland]
 - 1 **P. berggrenianum** Frisvoll
 - 2 **P. cavifolium** (Brid.) Z.Iwats.
 - 3 **P. curvifolium** Schlieph. ex Limpr.
 - 4 **P. denticulatum** (Hedw.) Schimp.
 - a var. **denticulatum**
 - b var. **obtusifolium** (Turner) Moore
 - c var. **undulatum** R.Ruthe ex Geh.
 - 5 **P. laetum** Schimp.
 - 6 **P. latebricola** Schimp.

⁴²⁰Majestyk (2011) showed that the correct name for *Daltonia stenophylla* is *Daltonia lindigiana*.

⁴²¹We follow Hill et al. (2008) and Guerra (2014) in treating *Fontinalis squamosa* var. *dixonii* as synonymous with the type, and var. *curnowii* is also considered to have no taxonomic value.

- 7 ***P. neckeroideum*** Schimp. [*Plagiothecium noricum* Molendo ex Limpr.]⁴²²
- 8 ***P. nemorale*** (Mitt.) A.Jaeger
- 9 ***P. piliferum*** (Sw.) Schimp.
- 10 ***P. platyphyllum*** Mönk.
- 11 ***P. rossicum*** Ignatov & Ignatova⁴²³
- 12 ***P. succulentum*** (Wilson) Lindb.
- 13 ***P. svalbardense*** Frisvoll
- 14 ***P. undulatum*** (Hedw.) Schimp. [*Buckiella undulata* (Hedw.) Ireland]
- 182 ***Myurella*** Bruch & Schimp.
- 1 ***M. julacea*** (Schwägr.) Schimp.
- 2 ***M. sibirica*** (Müll.Hal.) Reimers
- 3 ***M. tenerrima*** (Brid.) Lindb.
- 183 ***Ortholimnobia*** Dixon
- 1 ***O. handelii*** (Broth.) C.Schröck & J.T.Wynns [*Plagiothecium handelii* Broth.]⁴²⁴
- 184 ***Orthothecium*** Bruch & Schimp.
- 1 ***O. chryseon*** (Schwägr.) Schimp.
- 2 ***O. intricatum*** (Hartm.) Schimp.
- 3 ***O. lapponicum*** (Schimp.) C.Hartm.
- 4 ***O. rufescens*** (Dicks. ex Brid.) Schimp.
- 5 ***O. strictum*** Lorentz
- 185 ***Platydictya*** Berk.
- 1 ***P. jungermannioides*** (Brid.) H.A.Crum
- 186 ***Pseudotaxiphyllum*** Z.Iwats.
- 1 ***P. elegans*** (Brid.) Z.Iwats.
- 2 ***P. laetevirens*** (Dixon & Luisier ex F.Koppe & Düll) Hedenäs
- Fabroniaceae Schimp.
- 187 ***Fabronia*** Raddi
- 1 ***F. altaica*** Ignatova & Ignatov⁴²⁵
- 2 ***F. ciliaris*** (Brid.) Brid.
- 3 ***F. major*** De Not.⁴²⁶
- 4 ***F. pusilla*** Raddi
- Pterigynandraceae Schimp.
- 188 ***Pterigynandrum*** Hedw.
- 1 ***P. filiforme*** Hedw. [*Pterigynandrum filiforme* var. *majus* (De Not.) De Not.]⁴²⁷
- Habrodontaceae Schimp.
- 189 ***Habrodon*** Schimp.
- 1 ***H. perpusillus*** (De Not.) Lindb.
- Climaciaceae Kindb.
- 190 ***Climacium*** F.Weber & D.Mohr
- 1 ***C. dendroides*** (Hedw.) F.Weber & D.Mohr
- Myriniaceae Schimp.
- 191 ***Myrinia*** Schimp.
- 1 ***M. pulvinata*** (Wahlenb.) Schimp.
- Amblystegiaceae G.Roth
- Cratoneuroideae Vanderp., Hedenäs, C.J.Cox & A.J.Shaw
- 192 ***Cratoneuron*** (Sull.) Spruce
- 1 ***C. curvicaule*** (Jur.) G.Roth
- 2 ***C. filicinum*** (Hedw.) Spruce
- 193 ***Palustriella*** Ochyra
- 1 ***P. commutata*** (Hedw.) Ochyra
- 2 ***P. decipiens*** (De Not.) Ochyra
- 3 ***P. falcata*** (Brid.) Hedenäs [*Palustriella commutata* var. *sulcata* (Lindb.) Ochyra, *Palustriella pluristratosa* M.Stech & J.-P.Frahm]⁴²⁸
- Amblystegioideae Broth.
- 194 ***Amblystegium*** Schimp.
- 1 ***A. serpens*** (Hedw.) Schimp. [*Amblystegium serpens* var. *salinum* Carrington]
- 195 ***Anacamptodon*** Brid.
- 1 ***A. splachnoides*** (Froel. ex Brid.) Brid.
- 196 ***Arvernella*** Hugonnot & Hedenäs⁴²⁹
- 1 ***A. microclada*** Hugonnot & Hedenäs
- 197 ***Campyliadelphus*** (Kindb.) R.S.Chopra
- 1 ***C. chrysophyllus*** (Brid.) R.S.Chopra
- 2 ***C. elodes*** (Lindb.) Kanda
- 198 ***Campylium*** (Kindb.) R.S.Chopra
- 1 ***C. bambergeri*** (Schimp.) Hedenäs, Schlesak & D.Quandt [*Hypnum bambergeri* Schimp.]⁴³⁰
- 2 ***C. laxifolium*** Engelmark & Hedenäs

⁴²²The status of *Plagiothecium noricum* Molendo ex. Limpr., and its relation to *Plagiothecium neckeroideum*, still needs clarification.

⁴²³*Plagiothecium rossicum* was described from Russia by Ignatova, Fedorova, et al. (2019).

⁴²⁴*Plagiothecium handelii* was recorded new to Europe, and its position in *Ortholimnobia* confirmed, by Wynns and Schröck (2018).

⁴²⁵*Fabronia altaica* was described from Russia by Ignatova, Kuznetsova, et al. (2017). In Europe it is restricted to the Caucasus.

⁴²⁶*Fabronia major* was resurrected in the course of a study on the genus *Fabronia* in Russia by Ignatova, Kuznetsova, et al. (2017).

⁴²⁷According to Hill et al. (2008), "Smith (2004) retains *Pterigynandrum filiforme* var. *majus* but expresses severe doubts as to whether it is distinct. It is treated here as a synonym of var. *filiforme*." We follow this treatment.

⁴²⁸The synonymy of *Palustriella pluristratosa* with *Palustriella falcata* follows Hedenäs (2010). The taxonomic value of *Palustriella commutata* var. *sulcata* should be studied in more detail. It is a characteristic plant of high altitude limestone in the Alps.

⁴²⁹*Arvernella* is a new genus, described by Hugonnot and Hedenäs (2015) to accommodate *Arvernella microclada*, a new species found in the Auvergne, France. Its placement in the Amblystegioideae is supported by recent analysis by Kučera, Kuznetsova, et al. (2019).

⁴³⁰Molecular evidence places *Hypnum bambergeri* in *Campylium* (Schlesak et al. 2018).

- 3 **C. longicuspis** (Lindb. & Arnell) Hedenäs
 4 **C. protensum** (Brid.) Kindb.⁴³¹
 5 **C. stellatum** (Hedw.) Lange & C.E.O.Jensen
- 199 **Campylophyllopsis** W.R.Buck [*Campylidium* (Kindb.) Ochyra, *nom. illeg.*]⁴³²
 1 **C. calcarea** (Crundw. & Nyholm) Ochyra [*Campylidium calcareum* (Crundw. & Nyholm) Ochyra, *Campylophyllum calcareum* (Crundw. & Nyholm) Hedenäs]
 2 **C. sommerfeltii** (Myrin) Ochyra [*Campylidium sommerfeltii* (Myrin) Ochyra, *Campylophyllum sommerfeltii* (Myrin) Hedenäs]
- 200 **Campylophyllum** (Schimp.) M.Fleisch.
 1 **C. halleri** (Hedw.) M.Fleisch.
 2 **C. montanum** (Lindb.) B.H.Allen [*Hygrohypnum montanum* (Lindb.) Broth., *Platyhypnum montanum* (Lindb.) Ochyra]⁴³³
- 201 **Conardia** H.Rob.⁴³⁴
 1 **C. compacta** (Drumm. ex Müll.Hal.) H.Rob.
- 202 **Drepanium** (Schimp.) C.E.O.Jensen⁴³⁵
 1 **D. fastigiatum** (Hampe) C.E.O.Jensen [*Hypnum recurvatum* (Lindb. & Arnell) Kindb., *Drepanium recurvatum* (Lindb. & Arnell) G.Roth]
- 203 **Drepanocladus** (Müll.Hal.) G.Roth [*Pseudocalliergon* (Limpr.) Loeske]⁴³⁶
 1 **D. aduncus** (Hedw.) Warnst.
 2 **D. angustifolius** (Hedenäs) Hedenäs & C.Rosborg [*Pseudocalliergon angustifolium* Hedenäs]
 3 **D. arcticus** (R.S.Williams) Hedenäs
 4 **D. brevifolius** (Lindb.) Warnst. [*Pseudocalliergon brevifolium* (Lindb.) Hedenäs]
- 5 **D. capillifolius** (Warnst.) Warnst. [*Drepanocladus longifolius* auct. eur., non (Mitt.) Paris]⁴³⁷
 6 **D. lycopodioides** (Brid.) Warnst. [*Pseudocalliergon lycopodioides* (Brid.) Hedenäs]
 7 **D. polygamus** (Schimp.) Hedenäs⁴³⁸
 8 **D. sendtneri** (Schimp. ex H.Müll.) Warnst.
 9 **D. sordidus** (Müll.Hal.) Hedenäs
 10 **D. trifarius** (F.Weber & D.Mohr) Broth. ex Paris [*Pseudocalliergon trifarium* (F.Weber & D.Mohr) Loeske]
 11 **D. turgescens** (T.Jensen) Broth. [*Pseudocalliergon turgescens* (T.Jensen) Loeske]
- 204 **Hygroamblystegium** Loeske⁴³⁹
 1 **H. fluviatile** (Hedw.) Loeske
 2 **H. humile** (P.Beauv.) Vanderp., Goffinet & Hedenäs [*Hygroamblystegium varium* var. *humile* Vanderp. & Hedenäs]⁴⁴⁰
 3 **H. tenax** (Hedw.) Jenn.
 4 **H. varium** (Hedw.) Mönk. [*Amblystegium varium* (Hedw.) Lindb.]
- 205 **Hygrohypnum** Lindb.
 1 **H. luridum** (Hedw.) Jenn.
 2 **H. styriacum** (Limpr.) Broth.
- 206 **Leptodictyum** (Schimp.) Warnst.
 1 **L. riparium** (Hedw.) Warnst.
- 207 **Microhypnum** Jan Kučera & Ignatov
 1 **M. sauteri** (Schimp.) Jan Kučera & Ignatov [*Anacamptodon sauteri* (Schimp.) Hedenäs, Schlesak & D.Quandt, *Hypnum sauteri* Schimp.]⁴⁴¹
- 208 **Platyhypnum** Loeske [*Ochyraea Váňa*]⁴⁴²
 1 **P. alpestre** (Hedw.) Ochyra [*Hygrohypnum alpestre* (Hedw.) Loeske]
 2 **P. alpinum** (Lindb.) Loeske [*Hygrohypnum alpinum* (Lindb.) Loeske]

⁴³¹*Campylium protensum* and *Campylium stellatum* are distinct in some parts of their range and overlap in others. They are retained as separate species for the present, although there is a strong case for *Campylium protensum* to be reduced to a variety of *Campylium stellatum*, as some authors have done.

⁴³²The separation of *Campylidium* from *Campylophyllum* is supported by molecular studies (Gardiner et al. 2005; Ignatov, Gardiner, et al. 2007). However, *Campylidium* is an illegitimate name and must be changed to *Campylophyllopsis* (Goffinet, Buck, et al. 2009).

⁴³³Ochyra (2013) placed *Hygrohypnum montanum* in *Platyhypnum*. Allen (2014) transferred it to *Campylophyllum*.

⁴³⁴Although here retained in the Amblystegiaceae, *Conardia* does not appear to belong here, and may require a family of its own (see, for example, Vanderpoorten et al. 2002).

⁴³⁵Recent molecular studies place *Hypnum recurvatum* in the genus *Drepanium* (Schlesak et al. 2018).

⁴³⁶This treatment of *Drepanocladus* follows Hedenäs and Rosborg (2008).

⁴³⁷*Drepanocladus longifolius* (Mitt.) Paris is an exclusively Southern Hemisphere species; Saługa et al. (2018) showed that European records of this species should be referred to *Drepanocladus capillifolius*.

⁴³⁸*Campylium decipiens* (Warnst.) Walsemann is a central European plant recognised as a distinct species by Frahm and Walsemann (1973) and Meinunger and Schröder (2007), but otherwise largely disregarded. It may be a form of *Drepanocladus polygamus*, but it requires further study.

⁴³⁹Although *Hygroamblystegium fluviatile*, *Hygroamblystegium humile* and *Hygroamblystegium tenax* have been synonymised with *Hygroamblystegium varium* (Vanderpoorten 2004), we follow Hill et al. (2006) in retaining these species as distinct pending further work.

⁴⁴⁰*Hygroamblystegium humile* was recognised at varietal level by Vanderpoorten and Hedenäs (2009), while *Hygroamblystegium fluviatile* and *Hygroamblystegium tenax* continued to be regarded as synonyms of *Amblystegium varium*.

⁴⁴¹On molecular evidence, Schlesak et al. (2018) placed *Hypnum sauteri* in *Anacamptodon* but Kučera, Kuznetsova, et al. (2019) provided further evidence placing it in the new genus *Microhypnum*.

⁴⁴²We follow the treatment of Ochyra (2013) in transferring several *Hygrohypnum* species to *Platyhypnum*.

- 3 ***P. cochlearifolium*** (Venturi) Ochyra [*Hygrohypnum cochlearifolium* (Venturi) Broth.]
- 4 ***P. duriusculum*** (De Not.) Ochyra [*Hygrohypnum duriusculum* (De Not.) D.W.Jamieson]
- 5 ***P. molle*** (Dicks. ex Hedw.) Loeske [*Hygrohypnum molle* (Dicks. ex Hedw.) Loeske]
- 6 ***P. norvegicum*** (Schimp.) Ochyra [*Hygrohypnum norvegicum* (Schimp.) J.J.Amann]
- 7 ***P. smithii*** (Sw.) Ochyra [*Hygrohypnum smithii* (Sw.) Broth.]
- 8 ***P. tatrense*** (Váňa) Hedenäs & Ignatov [*Ochyraea tatrensis* Váňa]⁴⁴³
- 209 ***Pseudoamblystegium*** Vanderp. & Hedenäs⁴⁴⁴
- 1 ***P. subtile*** (Hedw.) Vanderp. & Hedenäs [*Amblystegium subtile* (Hedw.) Schimp.]
- 210 ***Pseudocampylium*** Vanderp. & Hedenäs⁴⁴⁵
- 1 ***P. radicale*** (P.Beauv.) Vanderp. & Hedenäs [*Amblystegium radicale* (P.Beauv.) Schimp.]
- 211 ***Serpoleskea*** (Limpr.) Loeske
- 1 ***S. confervoides*** (Brid.) Schimp. [*Amblystegium confervoides* (Brid.) Schimp.]⁴⁴⁶
- 212 ***Tomentypnum*** Loeske
- 1 ***T. nitens*** (Hedw.) Loeske
- Calliergonaceae Vanderp., Hedenäs, C.J.Cox & A.J.Shaw
- 213 ***Calliergon*** (Sull.) Kindb.
- 1 ***C. cordifolium*** (Hedw.) Kindb.
- 2 ***C. giganteum*** (Schimp.) Kindb.
- 3 ***C. megalophyllum*** Mikut.
- 4 ***C. richardsonii*** (Mitt.) Kindb.
- 214 ***Loeskytnum*** H.K.G.Paul
- 1 ***L. badium*** (Hartm.) H.K.G.Paul
- 215 ***Sarmentypnum*** Tuom. & T.J.Kop.⁴⁴⁷
- 1 ***S. exannulatum*** (Schimp.) Hedenäs [*Warnstorfia exannulata* (Schimp.) Loeske]
- 2 ***S. procerum*** (Renauld & Arnell) Hedenäs [*Warnstorfia procerum* (Renauld & Arnell) Tuom.]
- 3 ***S. sarmentosum*** (Wahlenb.) Tuom. & T.J.Kop. [*Warnstorfia sarmentosa* (Wahlenb.) Hedenäs]
- 4 ***S. trichophyllum*** (Warnst.) Hedenäs [*Warnstorfia trichophylla* (Warnst.) Tuom. & T.J.Kop.]
- 5 ***S. tundrae*** (Arnell) Hedenäs [*Warnstorfia tundrae* (Arnell) Loeske]
- 216 ***Straminergon*** Hedenäs
- 1 ***S. stramineum*** (Dicks. ex Brid.) Hedenäs
- 217 ***Warnstorfia*** Loeske
- 1 ***W. fluitans*** (Hedw.) Loeske
- 2 ***W. pseudostraminea*** (Müll.Hal.) Tuom. & T.J.Kop.
- Scorpidiaceae Ignatov & Ignatova
- 218 ***Hamatocaulis*** Hedenäs
- 1 ***H. lapponicus*** (Norrl.) Hedenäs
- 2 ***H. vernicosus*** (Mitt.) Hedenäs⁴⁴⁸
- 219 ***Hygrohypnella*** Ignatov & Ignatova⁴⁴⁹
- 1 ***H. ochracea*** (Turner ex Wilson) Ignatov & Ignatova [*Hygrohypnum ochraceum* (Turner ex Wilson) Loeske]
- 2 ***H. polaris*** (Lindb.) Ignatov & Ignatova [*Hygrohypnum polare* (Lindb.) Loeske]
- 220 ***Sanionia*** Loeske
- 1 ***S. nivalis*** Hedenäs [*Sanionia georgicounicinata* auct. eur., non (Müll.Hal.) Ochyra & Hedenäs]⁴⁵⁰
- 2 ***S. orthothecioides*** (Lindb.) Loeske
- 3 ***S. uncinata*** (Hedw.) Loeske
- 221 ***Scorpidium*** (Schimp.) Limpr.
- 1 ***S. cossonii*** (Schimp.) Hedenäs
- 2 ***S. revolvens*** (Sw. ex anon.) Rubers
- 3 ***S. scorpioides*** (Hedw.) Limpr.
- Leskeaceae Schimp.
- 222 ***Claopodium*** (Lesq. & James) Renauld & Cardot
- 1 ***C. rostratum*** (Hedw.) Ignatov [*Anomodon rostratus* (Hedw.) Schimp.]⁴⁵¹
- 2 ***C. whippleanum*** (Sull.) Renauld & Cardot

⁴⁴³*Ochyraea tatrensis* may be nothing more than an extreme form of *Platyhypnum smithii*, but for the present it is retained as a species and transferred to the genus *Platyhypnum* until further work is done.

⁴⁴⁴The genus *Pseudoamblystegium* was established by Vanderpoorten and Hedenäs (2009) to accommodate *Amblystegium subtile*.

⁴⁴⁵The genus *Pseudocampylium* was established by Vanderpoorten and Hedenäs (2009) to accommodate *Amblystegium radicale*.

⁴⁴⁶The placement of *Amblystegium confervoides* in *Serpoleskea* (Vanderpoorten et al. 2002) was confirmed by Vanderpoorten and Hedenäs (2009).

⁴⁴⁷The treatment of *Sarmentypnum* follows Hedenäs (2006).

⁴⁴⁸*Hamatocaulis vernicosus* comprises two well-researched cryptic species in Europe. Careful morphological evaluation has failed to find any morphological difference between the two (Manukjanová et al. 2019).

⁴⁴⁹*Hygrohypnum ochraceum* and *Hygrohypnum polare* are placed in *Hygrohypnella*, following Ignatov and Ignatova (2004).

⁴⁵⁰Molecular studies showed European *Sanionia nivalis* to be distinct from the Southern Hemisphere *Sanionia georgicounicinata* (Müll.Hal.) Ochyra & Hedenäs (Hedenäs 2012).

⁴⁵¹Molecular studies place *Anomodon rostratus* in *Claopodium* (Ignatov, Afonina, Ignatova 2006).

- 223 **Leskea** Hedw.
1 **L. polycarpa** Hedw.
- 224 **Lindbergia** Kindb.⁴⁵²
1 **L. dagestanica** Ignatova & Ignatov
2 **L. grandiretis** (Lindb. ex Broth.) Ignatov & Ignatova [*Lindbergia brachyptera* auct. eur., non (Mitt.) Kindb.]
- 225 **Pseudoleskeopsis** Broth.⁴⁵³
1 **P. artariae** (Thér.) Thér. [*Pseudoleskea artariae* Thér.]
- Pseudoleskeaceae Schimp.
- 226 **Lescurea** Bruch & Schimp. [*Pseudoleskea* Schimp., *Ptychodium* Schimp.]⁴⁵⁴
1 **L. incurvata** (Hedw.) E.Lawton [*Pseudoleskea incurvata* (Hedw.) Loeske]
2 **L. mutabilis** (Brid.) Lindb. ex I.Hagen
3 **L. patens** Lindb. [*Pseudoleskea patens* (Lindb.) Kindb.]
4 **L. plicata** (Schleich. ex F.Weber & D.Mohr) Broth. [*Ptychodium plicatum* (Schleich. ex F.Weber & D.Mohr) Schimp.]
5 **L. radicata** (Mitt.) Mönk. [*Pseudoleskea radicata* (Mitt.) Macoun & Kindb.]
6 **L. saviana** (De Not.) E.Lawton [*Pseudoleskea saviana* (De Not.) Latzel]
7 **L. saxicola** (Schimp.) Molendo
8 **L. secunda** Arnell
- Pseudoleskeellaceae Ignatov & Ignatova
- 227 **Pseudoleskeella** Kindb.
1 **P. catenulata** (Brid. ex Schrad.) Kindb.
2 **P. nervosa** (Brid.) Nyholm
3 **P. papillosa** (Lindb.) Kindb.
4 **P. rupestris** (Berggr.) Hedenäs & L.Söderstr.
5 **P. tectorum** (Funck ex Brid.) Kindb. ex Broth.
- Thuidiaceae Schimp.
- 228 **Abietinella** Müll.Hal.
1 **A. abietina** (Hedw.) M.Fleisch.
a var. **abietina**
b var. **hystricosa** (Mitt.) Sakurai
- 229 **Haplocladium** (Müll.Hal.) Müll.Hal.
1 **H. angustifolium** (Hampe & Müll.Hal.) Broth.
- 2 **H. microphyllum** (Hedw.) Broth.
3 **H. virginianum** (Brid.) Broth.
- 230 **Helodium** (Müll.Hal.) Müll.Hal.⁴⁵⁵
1 **H. blandowii** (F.Weber & D.Mohr) Warnst. [*Elodium blandowii* (F.Weber & D.Mohr) Eckel]
- 231 **Pelekium** Mitt.
1 **P. atlanticum** (Hedenäs) Hedenäs
2 **P. minutulum** (Hedw.) Touw
- 232 **Thuidiopsis** (Broth.) M.Fleisch.
1 **T. sparsa** (Hook.f. & Wilson) Broth.
- 233 **Thuidium** Bruch & Schimp.
1 **T. assimile** (Mitt.) A.Jaeger
2 **T. delicatulum** (Hedw.) Schimp.
3 **T. recognitum** (Hedw.) Lindb.
4 **T. tamariscinum** (Hedw.) Schimp.
- Brachytheciaceae Schimp.
Eurhynchioideae Milde
- 234 **Eurhynchium** Bruch & Schimp.
1 **E. angustirete** (Broth.) T.J.Kop.
2 **E. striatum** (Hedw.) Schimp.
- 235 **Palamocladium** M.Fleisch.
1 **P. euchloron** (Müll.Hal.) Wijk & Margad.
- 236 **Plasteurhynchium** M.Fleisch.
1 **P. meridionale** (Schimp.) M.Fleisch.
2 **P. striatulum** (Spruce) M.Fleisch.
- 237 **Pseudoscleropodium** (Limpr.) M.Fleisch.
1 **P. purum** (Hedw.) M.Fleisch.
- 238 **Rhynchostegium** Bruch & Schimp. [*Platyhypnidium* M.Fleisch.]⁴⁵⁶
1 **R. alopecuroides** (Brid.) A.J.E.Sm. [*Platyhypnidium lusitanicum* (Schimp.) Ochyra & Bedn.-Ochyra, *Platyhypnidium mutatum* Ochyra & Vanderp.]⁴⁵⁷
2 **R. confertum** (Dicks.) Schimp.
3 **R. confusum** K.Cezón, J.Muñoz, Hedenäs & Huttunen⁴⁵⁸

⁴⁵²The treatment of *Lindbergia* follows Ignatova, Ignatov, et al. (2010).

⁴⁵³The type of *Pseudoleskea* is *Pseudoleskea atrovirens* (= *Pseudoleskea incurvata*), now in *Lescurea*, so *Pseudoleskea artariae* cannot be the sole representative of the genus. Therefore we return this taxon to the genus *Pseudoleskeopsis* (placed by Frey and Stech 2009 in Leskeaceae) pending further studies.

⁴⁵⁴Molecular studies suggest that *Pseudoleskea* and *Ptychodium* should be included in *Lescurea* (Gardiner et al. 2005; Ignatov, Gardiner, et al. 2007).

⁴⁵⁵Eckel (2012) proposed that *Elodium* (Sull.) Austin take the place of *Helodium*, on the basis of precedence. However, preliminary research by M. Ignatov indicates that North American and European material is not necessarily congeneric. Globally, there are (traditionally) three species commonly placed in the genus *Helodium*, but they are probably unrelated. As the name *Helodium* is in common usage, it is retained pending further studies.

⁴⁵⁶Molecular studies have shown that the genus *Platyhypnidium* is not sustainable, with all the European species correctly placed in *Rhynchostegium* (Huttunen and Ignatov 2010).

⁴⁵⁷The generic placement of *Rhynchostegium alopecuroides* was confirmed using molecular techniques by Huttunen and Ignatov (2010). Molecular studies have shown that *Platyhypnidium mutatum* is deeply nested within *Rhynchostegium alopecuroides*, although a formal synonymy was not made (Hutsemékers et al. 2012).

⁴⁵⁸*Rhynchostegium confusum* was described from the Iberian Peninsula by Cezón et al. (2010).

- 4 **R. megapolitanum** (Blandow ex F.Weber & D.Mohr) Schimp.
- 5 **R. murale** (Hedw.) Schimp. [*Rhynchostegium arcticum* (I.Hagen) Ignatov & Huttunen]⁴⁵⁹
- 6 **R. riparioides** (Hedw.) Cardot [*Platyhypnidium grolleanum* Ochyra & Bedn.-Ochyra, *Platyhypnidium torrenticola* (Ochyra, C.Schmidt & Bültmann) Ochyra & Bedn.-Ochyra]⁴⁶⁰
- 7 **R. rotundifolium** (Scop. ex Brid.) Schimp.
- 8 **R. strongylense** (Bott.) W.R.Buck & Privitera
- 239 **Scorpiurium** Schimp.
- 1 **S. circinatum** (Bruch) M.Fleisch. & Loeske
- 2 **S. deflexifolium** (Solms) M.Fleisch. & Loeske
- 3 **S. sendtneri** (Schimp.) M.Fleisch.
- Helicodontoideae M.Fleisch.
- 240 **Cirriphyllum** Grout
- 1 **C. crassinervium** (Taylor) Loeske & M.Fleisch.
- 2 **C. piliferum** (Hedw.) Grout
- 241 **Clasmatodon** Hook.f. & Wilson
- 1 **C. parvulus** (Hampe) Sull.⁴⁶¹
- 242 **Hedenasiastrum** Ignatov & Vanderp.⁴⁶²
- 1 **H. percurrens** (Hedenäs) Ignatov & Vanderp. [*Brachythecium percurrens* Hedenäs]
- 243 **Helicodontium** Schwägr.
- 1 **H. capillare** (Hedw.) A.Jaeger
- 244 **Microeurhynchium** Ignatov & Vanderp.⁴⁶³
- 1 **M. pumilum** (Wilson) Ignatov & Vanderp. [*Oxyrrhynchium pumilum* (Wilson) Loeske]
- 245 **Nobregaea** Hedenäs
- 1 **N. latinervis** Hedenäs
- 246 **Oxyrrhynchium** (Schimp.) Warnst.
- 1 **O. hians** (Hedw.) Loeske
- 2 **O. schleicheri** (R.Hedw.) Röhl
- 3 **O. speciosum** (Brid.) Warnst.
- 247 **Pseudorhynchostegiella** Ignatov & Vanderp.⁴⁶⁴
- 1 **P. duriaei** (Mont.) Ignatov & Vanderp. [*Rhynchostegiella durieui* (Mont.) P.Allorge & Perss.]
- 248 **Rhynchostegiella** (Schimp.) Limpr.
- 1 **R. azorica** Hedenäs & Vanderp.⁴⁶⁵
- 2 **R. bourgaeana** (Mitt.) Broth.
- 3 **R. curviseta** (Brid.) Limpr.
- 4 **R. litorea** (De Not.) Limpr. [*Rhynchostegiella tenella* var. *meridionalis* (Boulay) Zodda]⁴⁶⁶
- 5 **R. pseudolitorea** Hedenäs & J.Patiño⁴⁶⁷
- 6 **R. tenella** (Dicks.) Limpr.
- 7 **R. teneriffae** (Mont.) Dirkse & Bouman [*Rhynchostegiella jacquinii* (Garov.) Limpr., *Rhynchostegiella macilenta* (Renauld & Cardot) Cardot, *Rhynchostegiella teesdalei* (Schimp.) Limpr.]⁴⁶⁸
- 8 **R. trichophylla** Dirkse & Bouman
- 9 **R. tubulosa** Hedenäs & J.Patiño⁴⁶⁹
- Brachythecioideae Lotsy
- 249 **Brachytheciastrum** Ignatov & Huttunen
- 1 **B. collinum** (Schleich. ex Müll.Hal.) Ignatov & Huttunen [*Brachytheciastrum fendleri* auct. eur., non (Sull.) Ochyra & Żarnowiec]⁴⁷⁰
- 2 **B. dieckei** (Röhl) Ignatov & Huttunen
- 3 **B. olympicum** (Jur.) Vanderp. et al.
- 4 **B. salicinum** (Schimp.) J.D.Orgaz, M.J.Cano & J.Guerra [*Brachytheciastrum velutinum* var. *salicinum* (Schimp.) Ochyra & Żarnowiec]⁴⁷¹
- 5 **B. trachypodium** (Brid.) Ignatov & Huttunen
- 6 **B. velutinum** (Hedw.) Ignatov & Huttunen [*Brachytheciastrum vanekii* (Šmarda)

⁴⁵⁹Recent molecular work by M. Ignatov suggests that *Rhynchostegium arcticum* should be treated as a synonym of *Rhynchostegium murale*.

⁴⁶⁰*Platyhypnidium grolleanum* was regarded as a rare mutation of *Rhynchostegium riparioides* by Kučera, Váňa, et al. (2012). Molecular studies have shown that *Platyhypnidium torrenticola* is deeply nested within *Rhynchostegium riparioides*, although a formal synonymy was not made (Hutsemékers et al. 2012).

⁴⁶¹*Clasmatodon parvulus* is re-admitted into the European flora following the revision of an old herbarium specimen (collected in Germany in 1851) by Müller (2007). It has not been refound more recently.

⁴⁶²*Hedenasiastrum* was described on molecular grounds (Aigoïn et al. 2009).

⁴⁶³*Microeurhynchium* was described on molecular grounds (Aigoïn et al. 2009).

⁴⁶⁴*Pseudorhynchostegiella* was described on molecular grounds (Aigoïn et al. 2009).

⁴⁶⁵*Rhynchostegiella azorica* is an Azorean endemic described by Vanderpoorten et al. (2015).

⁴⁶⁶The synonymy of *Rhynchostegiella tenella* var. *meridionalis* with *Rhynchostegiella litorea* follows Guerra, Ríos, et al. (2014) and Patiño et al. (2017).

⁴⁶⁷*Rhynchostegiella pseudolitorea* is a Macaronesian endemic described by Patiño et al. (2017).

⁴⁶⁸*Rhynchostegiella macilenta* was synonymised with *R. teneriffae* by Patiño et al. (2017). The difficulties with *Rhynchostegiella jacquinii* and *Rhynchostegiella teesdalei* were discussed by Patiño et al. (2017), who listed them as doubtfully synonymous with *Rhynchostegiella teneriffae*. We continue to treat them as synonyms pending further work.

⁴⁶⁹*Rhynchostegiella tubulosa* is an eastern Mediterranean species described by Patiño et al. (2017), and subsequently also found in Portugal (Ellis, Amélio, et al. 2019).

⁴⁷⁰All European records of *Brachytheciastrum fendleri* are *Brachytheciastrum collinum* (Orgaz et al. 2013). *Brachytheciastrum fendleri* (Sull.) Ochyra & Żarnowiec is a North American species.

⁴⁷¹*Brachytheciastrum salicinum* is reinstated as a species, as it is significantly different from *Brachytheciastrum velutinum* morphologically, molecularly and ecologically (D. Orgaz pers. comm. 2018).

- Ochyra & Żarnowiec, *Brachythecias-
trum velutinum* var. *vagans* (Milde)
Ochyra & Żarnowiec]⁴⁷²
- 250 ***Brachythecium*** Schimp. [*Bryhnia* Kaurin]
1 ***B. albicans*** (Hedw.) Schimp.
2 ***B. buchananii*** (Hook.) A.Jaeger⁴⁷³
3 ***B. campestre*** (Müll.Hal.) Schimp.
4 ***B. capillaceum*** (F.Weber & D.Mohr) Giacom.
[*Brachythecium rotaeanum* De
Not.]⁴⁷⁴
5 ***B. cirrosum*** (Schwägr.) Schimp.
6 ***B. erythrorrhizon*** Schimp. [*Brachythecium*
erythrorrhizon subsp. *asiaticum*
Ignatov, *Brachythecium erythrorrhizon*
subsp. *erythrorrhizon* var. *thedenii*
(Schimp.) Lindb.]⁴⁷⁵
7 ***B. funkii*** Schimp.⁴⁷⁶
8 ***B. geheebii*** Milde
9 ***B. glareosum*** (Bruch ex Spruce) Schimp.
10 ***B. japygum*** (Głow.) Köckinger & Jan
Kučera⁴⁷⁷
11 ***B. laetum*** (Brid.) Schimp.
12 ***B. mildeanum*** (Schimp.) Schimp.
13 ***B. novae-angliae*** (Sull. & Lesq.) A.Jaeger
[*Brachythecium scabridum* (Lindb.)
M.Li & Y.F.Wang, *Bryhnia scabrida*
(Lindb.) Kaurin; *Bryhnia novae-
angliae* (Sull. & Lesq.) Grout]⁴⁷⁸
14 ***B. rivulare*** Schimp.
15 ***B. rutabulum*** (Hedw.) Schimp.
a var. ***atlanticum*** Hedenäs
b var. ***rutabulum***
16 ***B. salebrosum*** (Hoffm. ex F.Weber &
D.Mohr) Schimp.
17 ***B. tauriscorum*** Molendo [*Brachythecium*
coruscum I.Hagen]⁴⁷⁹
18 ***B. tenuicaule*** (Spruce) Kindb. [*Rhynchoste-
giella tenuicaulis* (Spruce) Kartt.]⁴⁸⁰
- 19 ***B. tommasinii*** (Sendtn. ex Boulay) Ignatov &
Huttunen
20 ***B. turgidum*** (Hartm.) Kindb.
21 ***B. udum*** I.Hagen [*Brachythecium mildeanum*
var. *udum* (I.Hagen) Mönk.]⁴⁸¹
- 251 ***Eurhynchiastrum*** Ignatov & Huttunen
1 ***E. diversifolium*** (Schimp.) J.Guerra [*Eurhynch-
iastrum pulchellum* var. *diversifolium*
(Schimp.) Ochyra & Żarnowiec]⁴⁸²
2 ***E. pulchellum*** (Hedw.) Ignatov & Huttunen
[*Eurhynchiastrum pulchellum* var.
praecox (Hedw.) Ochyra & Żarnowiec]⁴⁸³
- 252 ***Homalothecium*** Schimp.
1 ***H. aureum*** (Spruce) H.Rob.
2 ***H. lutescens*** (Hedw.) H.Rob.
a var. ***fallax*** (H.Philib. ex Schimp.) Düll
b var. ***lutescens***
3 ***H. mandonii*** (Mitt.)Geh.⁴⁸⁴
4 ***H. meridionale*** (M.Fleisch. & Warnst.)
Hedenäs⁴⁸⁵
5 ***H. philippeanum*** (Spruce) Schimp.
6 ***H. sericeum*** (Hedw.) Schimp.
- 253 ***Kindbergia*** Ochyra
1 ***K. praelonga*** (Hedw.) Ochyra
- 254 ***Myuroclada*** Besch.
1 ***M. longiramea*** (Müll.Hal.) M.Li, Y.-F.Wang,
Ignatov & Huttunen⁴⁸⁶
2 ***M. maximowiczii*** (G.G.Borshch.) Steere &
W.B.Schofield
- 255 ***Sciuro-hypnum*** (Hampe) Hampe
1 ***S. curtum*** (Lindb.) Ignatov⁴⁸⁷
2 ***S. dovrense*** (Limpr.) Draper & Hedenäs⁴⁸⁸
3 ***S. flotowianum*** (Sendtn.) Ignatov & Huttunen
4 ***S. glaciale*** (Schimp.) Ignatov & Huttunen
5 ***S. latifolium*** (Kindb.) Ignatov & Huttunen

⁴⁷²*Brachytheciastrum vanekii* and *Brachytheciastrum velutinum* var. *vagans* (the latter known only from Poland) are synonymised with *Brachytheciastrum velutinum*, as they do not differ either morphologically or molecularly (J. Kučera and D. Orgaz pers. comm. 2018).

⁴⁷³*Brachythecium buchananii* is an essentially Asiatic species that also occurs on the European side of the Ural mountains (Ignatov and Milyutina 2010).

⁴⁷⁴*Brachythecium capillaceum* is most likely the earlier name for *Brachythecium rotaeanum*, although additional study is needed to confirm this.

⁴⁷⁵In reference to *Brachythecium erythrorrhizon* subsp. *asiaticum*, Ignatov and Milyutina (2010) observed, " ... too broad variation in plant size in Eurasia and North America, making [it] very difficult to segregate this subspecies." Neither *Brachythecium erythrorrhizon* subsp. *asiaticum* nor subsp. *erythrorrhizon* var. *thedenii* are now considered worthy of recognition at any level.

⁴⁷⁶*Brachythecium funkii* was segregated from *Brachythecium cirrosum* on the basis of molecular evidence (Köckinger and Kučera 2016).

⁴⁷⁷*Brachythecium japygum* was segregated from *Brachythecium cirrosum* on the basis of molecular evidence (Köckinger and Kučera 2016).

⁴⁷⁸The Eurasian *Brachythecium scabridum* was synonymised with the North American *Brachythecium novae-angliae* by Huttunen, Kuznetsova, et al. (2015), as no significant differences could be found.

⁴⁷⁹*Brachythecium tauriscorum* is an older name for *B. coruscum* (Hedenäs 2017b).

⁴⁸⁰Molecular work showed *Brachythecium tenuicaule* to be closely related to *Brachythecium tommasinii* (Köckinger and Kučera 2016).

⁴⁸¹*Brachythecium udum* was treated as a variety of *Brachythecium mildeanum* (Podpera 1954; Hill et al. 2006), but Ignatov and Milyutina (2010) found it to be more closely related to *Brachythecium turgidum*. Sporophytes and male gametangia remain unknown, and it needs further study to determine whether it is a good species or an extreme phenotype in the *Brachythecium salebrosum*-*Brachythecium turgidum* complex growing in wet northern environments.

⁴⁸²*Eurhynchiastrum diversifolium* was raised to species level by Guerra (2016).

⁴⁸³The treatment of *Eurhynchiastrum pulchellum* follows Guerra (2016).

⁴⁸⁴*Homalothecium mandonii* was segregated from *H. sericeum* on the basis of molecular and morphological evidence (Hedenäs, Désamoré, et al. 2014).

⁴⁸⁵*Homalothecium meridionale* was segregated from *H. sericeum* on the basis of molecular and morphological evidence (Hedenäs, Désamoré, et al. 2014).

⁴⁸⁶The status of *Myuroclada longiramea* in Europe was clarified by Ignatov, Huttunen, et al. (2015).

⁴⁸⁷*Sciuro-hypnum curtum* was restored from synonymy with *Sciuro-hypnum oedipodium* by Ignatov and Milyutina (2007). It is a widespread species in Europe, whereas *Sciuro-hypnum oedipodium*, which is primarily a western North American species, is very rare, with just a few records from eastern Europe.

⁴⁸⁸*Sciuro-hypnum dovrense* was removed from synonymy with *Brachythecium glaciale* by Draper and Hedenäs (2009).

- 6 ***S. oedipodium*** (Mitt.) Ignatov & Huttunen
 7 ***S. ornellanum*** (Molendo) Ignatov & Huttunen
 8 ***S. plumosum*** (Hedw.) Ignatov & Huttunen
 9 ***S. populeum*** (Hedw.) Ignatov & Huttunen
 10 ***S. reflexum*** (Starke) Ignatov & Huttunen
 11 ***S. starkei*** (Brid.) Ignatov & Huttunen
 12 ***S. tromsoeense*** (Kaurin & Arnell) Draper & Hedenäs⁴⁸⁹
- 256 ***Scleropodium*** Bruch & Schimp.
 1 ***S. cespitans*** (Wilson ex Müll.Hal.) L.F.Koch
 2 ***S. touretii*** (Brid.) L.F.Koch
- Hypnaceae Schimp.
- 257 ***Hypnum*** Hedw.⁴⁹⁰
 1 ***H. andoi*** A.J.E.Sm.
 2 ***H. cupressiforme*** Hedw.
 a var. ***cupressiforme***
 b var. ***filiforme*** Brid.⁴⁹¹
 c var. ***heseleri*** (Ando & Higuchi) M.O.Hill
 d var. ***lacunosum*** Brid.
 e var. ***subjulaceum*** Molendo [*Hypnum subcomplanatum* Hedenäs, Schlesak & D.Quandt, *nom. illeg.*, *Hypnum subjulaceum* (Molendo) Hedenäs, Schlesak & D.Quandt]⁴⁹²
 3 ***H. jutlandicum*** Holmen & E.Warncke
 4 ***H. resupinatum*** Taylor [*Hypnum cupressiforme* var. *resupinatum* (Taylor) Schimp.]⁴⁹³
 5 ***H. uncinulatum*** Jur.
- Callicladiaceae Jan Kučera & Ignatov⁴⁹⁴
- 258 ***Callicladium*** H.A.Crum
 1 ***C. haldanianum*** (Grev.) H.A.Crum
 2 ***C. imponens*** (Hedw.) Hedenäs, Schlesak & D.Quandt [*Hypnum imponens* Hedw.]⁴⁹⁵
- Taxiphyllaceae Ignatov⁴⁹⁶
- 259 ***Taxiphyllum*** M.Fleisch.
 1 ***T. densifolium*** (Lindb. ex Broth.) Reimers
 2 ***T. wissgrillii*** (Garov.) Wijk & Margad.
- Pylaisiadelphaceae Goffinet & W.R.Buck
- 260 ***Brotherella*** M.Fleisch.
 1 ***B. lorentziana*** (Molendo ex Lorentz) Loeske ex M.Fleisch. [*Brotherella henonii* auct. eur. non (Duby) M.Fleisch.]⁴⁹⁷
- 261 ***Heterophyllum*** (Schimp.) Kindb.
 1 ***H. nemorosum*** (W.D.J.Koch ex Brid.) Kindb. [*Heterophyllum affine* (Hook.) M.Fleisch., *nom. illeg.*]⁴⁹⁸
- 262 ***Isopterygium*** Mitt.
 1 ***I. tenerum*** (Sw.) Mitt.
- 263 ***Platygyrium*** Bruch & Schimp.
 1 ***P. repens*** (Brid.) Schimp.
- Jocheniaceae Jan Kučera & Ignatov⁴⁹⁹
- 264 ***Jochenia*** Hedenäs, Schlesak & D.Quandt⁵⁰⁰
 1 ***J. pallescens*** (Hedw.) Hedenäs, Schlesak & D.Quandt [*Hypnum pallescens* (Hedw.) P.Beauv., *Hypnum reptile* Michx., *Hypnum pallescens* var. *reptile* (Michx.) Husn.]
 2 ***J. protuberans*** (Brid.) Jan Kučera & Ignatov [*Hypnum pallescens* var. *protuberans* (Brid.) Austin, *Hypnum protuberans* Brid.]⁵⁰¹
- Stereodontaceae Hedenäs, Schlesak & D.Quandt⁵⁰²
- 265 ***Stereodon*** (Brid.) Mitt. [*Breidleria* Loeske]⁵⁰³

⁴⁸⁹*Sciuro-hypnum tromsoeense* was removed from synonymy with *Brachythecium starkei* by Draper and Hedenäs (2008).

⁴⁹⁰The treatment of *Hypnum* follows Schlesak et al. (2018), apart from *Hypnum cupressiforme* var. *subjulaceum* (see footnote below).

⁴⁹¹*Hypnum cupressiforme* var. *filiforme* may well be nothing more than an environmental modification but is retained as a variety until its position is clarified.

⁴⁹²*Hypnum subjulaceum* was published by Schlesak et al. (2018) as *Hypnum subcomplanatum* rather than *Hypnum subjulaceum* because the latter was thought to be blocked by the name *Hypnum subjulaceum* Besch. However, the latter name was not validly published, so its use at species level is not blocked. This was corrected by Schlesak et al. (2019). Later, Kučera, Kuznetsova, et al. (2019) found that *Hypnum subjulaceum* formed a supported lineage within *Hypnum cupressiforme* s.lat., and so it was returned to varietal status.

⁴⁹³*Hypnum resupinatum* is treated at species level, as it is usually distinct morphologically and has a distinct geographical distribution. However, according to current knowledge, it might equally well be treated as a variety of *Hypnum cupressiforme*, as in Guerra (2018). Ongoing molecular studies should clarify the situation.

⁴⁹⁴The new family Callicladiaceae was established by Kučera, Kuznetsova, et al. (2019) on molecular grounds.

⁴⁹⁵Molecular evidence places *Hypnum imponens* in *Callicladium* (Schlesak et al. 2018).

⁴⁹⁶The family Taxiphyllaceae was established by Ignatov, Afonina, et al. (2012) to accommodate *Taxiphyllum*.

⁴⁹⁷Frahm (2013) synonymised *Brotherella lorentziana* with *Brotherella henonii* (Duby) M.Fleisch., an east Asian species, but a genetic review is still pending. We therefore continue to accept *Brotherella lorentziana* as valid.

⁴⁹⁸The name *Hypnum affine* Hook., basonym of *Heterophyllum affine*, is considered an orthographic variant of the older *Hypnum adfine* Hoffm. ex Schwägr. (Allen 2018). The next available synonym is *Hypnum nemorosum*.

⁴⁹⁹The new family Jocheniaceae was established by Kučera, Kuznetsova, et al. (2019) on molecular evidence.

⁵⁰⁰Schlesak et al. (2018) established the genus *Jochenia* for *Hypnum pallescens*.

⁵⁰¹Using molecular data, Kučera, Kuznetsova, et al. (2019) found that *Jochenia protuberans* was consistently different from *Jochenia pallescens*. Although this taxon has not often been recognised even at varietal level, there is now sufficient evidence to treat it as a species. Ando (1973) correctly distinguished between the taxa, although he treated *Jochenia protuberans* only as a form of *Jochenia pallescens*. His synonymy of *Hypnum reptile* with *Hypnum pallescens* and clarification of the misunderstanding concerning the type of *Leskea pallescens* Hedw. introduced by the authors of *Bryologia Europaea*, is followed here.

⁵⁰²The family Stereodontaceae was established by Schlesak et al. (2018) to legitimate the high statistical support for the clade including *Stereodon*.

⁵⁰³The treatment of *Stereodon* follows Schlesak et al. (2018) and Kučera, Kuznetsova, et al. (2019).

- 1 **S. aemulans** (Bridl.) Broth. [*Hypnum aemulans* Bridl.]⁵⁰⁴
- 2 **S. callichrous** (Brid.) Lindb. [*Hypnum callichroum* Brid.]
- 3 **S. hamulosus** (Schimp.) Lindb. [*Hypnum hamulosum* Schimp.]
- 4 **S. holmenii** (Ando) Ignatov & Ignatova [*Hypnum holmenii* Ando]
- 5 **S. pratensis** (W.D.J.Koch ex Spruce) Warnst. [*Breidleria pratensis* (W.D.J.Koch ex Spruce) Loeske]
- 6 **S. subimponens** (Lesq.) Broth. [*Hypnum subimponens* Lesq.]
- Pylaisiaceae Schimp.
- 266 **Aquilonium** Hedenäs, Schlesak & D.Quandt⁵⁰⁵
- 1 **A. plicatum** (Lindb.) Hedenäs, Schlesak & D.Quandt [*Hypnum plicatum* (Lindb.) A.Jaeger]
- 267 **Buckia** D.Rios, M.T.Gallego & J.Guerra⁵⁰⁶
- 1 **B. vaucheri** (Lesq.) D.Rios, M.T.Gallego & J.Guerra [*Hypnum vaucheri* Lesq.]
- 268 **Calliergonella** Loeske
- 1 **C. cuspidata** (Hedw.) Loeske
- 2 **C. lindbergii** (Mitt.) Hedenäs
- 269 **Homomallium** (Schimp.) Loeske
- 1 **H. incurvatum** (Schrud. ex Brid.) Loeske
- 270 **Pseudohygrohypnum** Kanda⁵⁰⁷
- 1 **P. eugyrium** (Schimp.) Kanda [*Hygrohypnum eugyrium* (Schimp.) Broth.]
- 2 **P. fertile** (Sendtn.) Jan Kučera & Ignatov [*Hypnum fertile* Sendtn., *Stereodon fertilis* (Sendtn.) Lindb.]⁵⁰⁸
- 3 **P. subeugyrium** (Renauld & Cardot) Ignatov & Ignatova [*Hygrohypnum subeugyrium* (Renauld & Cardot) Broth.]⁵⁰⁹
- 271 **Pseudostereodon** (Broth.) M.Fleisch.⁵¹⁰
- 1 **P. procerrimus** (Molendo) M.Fleisch. [*Ctenidium procerrimum* (Molendo) Lindb., *Hypnum procerrimum* Molendo]
- 272 **Ptilium** De Not.⁵¹¹
- 1 **P. crista-castrensis** (Hedw.) De Not.
- 273 **Pylaisia** Schimp.
- 1 **P. polyantha** (Hedw.) Schimp.
- 2 **P. selwynii** Kindb.
- 274 **Roaldia** P.E.A.S.Câmara & Carv.-Silva⁵¹²
- 1 **R. dolomitica** (Milde) Hedenäs, Schlesak & D.Quandt [*Hypnum revolutum* (Mitt.) Lindb. var. *dolomiticum* (Milde) Mönk.]⁵¹³
- 2 **R. revoluta** (Mitt.) P.E.A.S.Câmara & M.Carvalho-Silva [*Hypnum revolutum* (Mitt.) Lindb.]
- 275 **Vesicularia** (Müll.Hal.) Müll.Hal.⁵¹⁴
- 1 **V. reimersiana** Bizot & P.de la Varde
- Sematophyllaceae Broth.
- 276 **Sematophyllum** Mitt.
- 1 **S. adnatum** (Michx.) E.Britton
- 2 **S. demissum** (Wilson) Mitt.
- 3 **S. substrumulosum** (Hampe) E.Britton
- Hylocomiaceae M.Fleisch.
- 277 **Hageniella** Broth.⁵¹⁵
- 1 **H. micans** (Mitt.) B.C.Tan & Y.Jia
- 278 **Hylocomiadelphus** Ochyra & Stebel⁵¹⁶
- 1 **H. triquetrus** (Hedw.) Ochyra & Stebel [*Rhytidadelphus triquetrus* (Hedw.) Warnst.]
- 279 **Hylocomiastrum** Broth.
- 1 **H. pyrenaicum** (Spruce) M.Fleisch.
- 2 **H. umbratum** (Hedw.) M.Fleisch.
- 280 **Hylocomium** Bruch & Schimp.
- 1 **H. splendens** (Hedw.) Schimp.

⁵⁰⁴*Hypnum aemulans* has been treated as a synonym of *Stereodon hamulosus* (*Hypnum hamulosum*) by most European authors, and was therefore omitted by Hill et al. (2006). No useful DNA could be extracted from this species during the recent revision of European *Hypnum* by Schlesak et al. (2018), so its generic placement remains uncertain. However, morphological evidence suggests that it should be located in *Stereodon*.

⁵⁰⁵The genus *Aquilonium* was established by Schlesak et al. (2018) to accommodate, among other species, *Hypnum plicatum*.

⁵⁰⁶The genus *Buckia* was established by Câmara et al. (2018) to accommodate *Hypnum vaucheri*.

⁵⁰⁷*Pseudohygrohypnum*, described by Kanda (1976 (1977)), is accepted, as a broad concept of *Hygrohypnum* can no longer be maintained.

⁵⁰⁸Kučera, Kuznetsova, et al. (2019) placed *Hypnum fertile* in *Pseudohygrohypnum* on molecular evidence.

⁵⁰⁹*Hygrohypnum subeugyrium* was transferred to *Pseudohygrohypnum* by Ignatov and Ignatova (2004).

⁵¹⁰*Pseudostereodon* was re-established by Câmara et al. (2018).

⁵¹¹The placement of *Ptilium* in the Pylaisiaceae is a temporary solution, as its position remains uncertain until further research takes place.

⁵¹²The genus *Roaldia* was established to accommodate *Hypnum revolutum* s.lat. (Câmara et al. 2018).

⁵¹³While *Roaldia dolomitica* is regarded as 'mere morphological variation' by Câmara et al. (2018), it is treated as a full species by Schlesak et al. (2018). The latter view is supported by Kučera, Kuznetsova, et al. (2019).

⁵¹⁴The placement of *Vesicularia* in Pylaisiaceae follows Ignatov, Gardiner, et al. (2007).

⁵¹⁵Although placed in Hylocomiaceae by Frey and Stech (2009), *Hageniella* is probably best placed in Sematophyllaceae.

⁵¹⁶*Hylocomiadelphus* was published by Ochyra and Stebel (2008) to accommodate *Rhytidadelphus triquetrus*. This was not widely accepted, but recent molecular work (Ignatov, Ignatova, et al. 2019) has supported the idea.

- 281 **Loeskeobryum** Broth.
1 **L. brevirostre** (Brid.) M.Fleisch.
- 282 **Pleurozium** Mitt.
1 **P. schreberi** (Willd. ex Brid.) Mitt.
- 283 **Rhytidiadelphus** (Limpr.) Warnst.
1 **R. loreus** (Hedw.) Warnst.
2 **R. squarrosus** (Hedw.) Warnst.
3 **R. subpinnatus** (Lindb.) T.J.Kop.
- Rhytidiaceae Broth.
- 284 **Rhytidium** (Sull.) Kindb.
1 **R. rugosum** (Hedw.) Kindb.
- Entodontaceae Kindb.
- 285 **Entodon** Müll.Hal.
1 **E. challengeri** (Paris) Cardot
2 **E. cladorrhizans** (Hedw.) Müll.Hal.
3 **E. concinnus** (De Not.) Paris
4 **E. schleicheri** (Schimp.) Demet.
- Cryphaeaceae Schimp.
- 286 **Cryphaea** D.Mohr
1 **C. heteromalla** (Hedw.) D.Mohr
- 287 **Dendrocryphaea** Broth.
1 **D. lamyana** (Mont.) P.Rao
- Leucodontaceae Schimp.
- 288 **Leucodon** Schwägr.
1 **L. canariensis** (Brid.) Schwägr.
2 **L. flagellaris** Lindb. ex Broth.⁵¹⁷
3 **L. immersus** Lindb.⁵¹⁸
4 **L. pendulus** Lindb.
5 **L. sciuroides** (Hedw.) Schwägr. [*Leucodon sciuroides* var. *morensis* (Schwägr.) De Not.]⁵¹⁹
6 **L. treleasei** (Cardot) Paris
- 289 **Nogopterium** Crosby & W.R.Buck [*Pterogonium* Sw. nom. illeg.]⁵²⁰
1 **N. gracile** (Hedw.) Crosby & W.R.Buck [*Pterogonium gracile* (Hedw.) Sm.]
- Antitrichiaceae Ignatov & Ignatova
- 290 **Antitrichia** Brid.
- 1 **A. californica** Sull.
2 **A. curtispindula** (Hedw.) Brid.
- Neckeraceae Schimp.
- 291 **Alleniella** S.Olsson, Enroth & D.Quandt⁵²¹
1 **A. besseri** (Lobarz.) S.Olsson, Enroth & D.Quandt [*Neckera besseri* (Lobarz.) Jur.]
2 **A. complanata** (Hedw.) S.Olsson, Enroth & D.Quandt [*Neckera complanata* (Hedw.) Huebener]
- 292 **Exsertotheca** S.Olsson, Enroth & D.Quandt⁵²²
1 **E. baetica** (J.Guerra) Draper, González-Mancebo, O.Werner, J.Patiño & Ros [*Neckera baetica* J.Guerra]⁵²³
2 **E. crispa** (Hedw.) S.Olsson, Enroth & D.Quandt [*Neckera crispa* Hedw.]
3 **E. intermedia** (Brid.) S.Olsson, Enroth & D.Quandt [*Neckera intermedia* Brid.]
- 293 **Homalia** (Brid.) Bruch & Schimp.
1 **H. lusitanica** Schimp.
2 **H. trichomanoides** (Hedw.) Brid.
- 294 **Iwatsukiella** W.R.Buck & H.A.Crum⁵²⁴
1 **I. leucotricha** (Mitt.) W.R.Buck & H.A.Crum
- 295 **Leptodon** D.Mohr
1 **L. corsicus** Enroth, Sotiaux, D.Quandt & Vanderp.⁵²⁵
2 **L. longisetus** Mont. [*Cryptoleptodon longisetus* (Mont.) Enroth]⁵²⁶
3 **L. smithii** (Hedw.) F.Weber & D.Mohr
- 296 **Neckera** Hedw.
1 **N. cephalonica** Jur. & Unger
2 **N. menziesii** Drumm.
3 **N. oligocarpa** Bruch
4 **N. pennata** Hedw.
5 **N. pumila** Hedw.
- 297 **Pseudanomodon** (Limpr.) Ignatov & Fedosov⁵²⁷
1 **P. attenuatus** (Hedw.) Ignatov & Fedosov [*Anomodon attenuatus* (Hedw.) Huebener]

⁵¹⁷Werner, Rodríguez-Atienza, et al. (2015) confirmed the status of *Leucodon flagellaris*.⁵¹⁸Werner, Rodríguez-Atienza, et al. (2015) confirmed the status of *Leucodon immersus*.⁵¹⁹Werner, Rodríguez-Atienza, et al. (2015) found no substantial basis for retaining *Leucodon sciuroides* var. *morensis*.⁵²⁰Although familiar through long use, *Pterogonium* is an illegitimate name. It was replaced by *Nogopterium* (an anagram) by Crosby and Buck (2011).⁵²¹*Alleniella* was segregated from *Neckera* by Olsson et al. (2011).⁵²²*Exsertotheca* was segregated from *Neckera* by Draper, González-Mancebo, et al. (2011).⁵²³*Neckera baetica* was described by Guerra, Jiménez-Martínez, Jiménez (2010) and transferred to *Exsertotheca* by Olsson et al. (2011).⁵²⁴*Iwatsukiella* has been placed in the Heterocladaceae Ignatov & Ignatova, but this family name is illegitimate, having previously been used for a family of algae. Therefore the genus is temporarily placed in Neckeraceae until another name is substituted.⁵²⁵*Leptodon corsicus*, apparently a Corsican endemic, was described by Sotiaux et al. (2009).⁵²⁶*Cryptoleptodon longisetus* was transferred to *Leptodon* by Olsson et al. (2011).⁵²⁷*Pseudanomodon* was established for *Anomodon attenuatus* after molecular studies showed it to be deeply nested in the Neckeraceae (Ignatov, Fedorova, et al. 2019).

- 298 *Thamnobryum* Nieuwl.
 1 *T. alopecurum* (Hedw.) Gangulee
 2 *T. angustifolium* (Holt) Nieuwl.⁵²⁸
 3 *T. cataractarum* N.G.Hodgetts & Blockeel⁵²⁹
 4 *T. fernandesii* Sérgio⁵³⁰
 5 *T. maderense* (Kindb.) Hedenäs
 6 *T. neckeroides* (Hook.) E.Lawton
 7 *T. rudolphianum* Mastracci
 8 *T. subserratum* (Hook. ex Harv.) Nog. & Z.lwats.⁵³¹
- Heterocradiellaceae Ignatov & Fedosov⁵³²
 299 *Heterocradiella* Ignatov & Fedosov
 1 *H. dimorpha* (Brid.) Ignatov & Fedosov [*Heterocradium dimorphum* (Brid.) Schimp.]
- Lembophyllaceae Broth.
 300 *Heterocladium* Bruch & Schimp.⁵³³
 1 *H. flaccidum* (Schimp.) A.J.E.Sm.
 2 *H. heteropterum* (Brid.) Schimp.
 3 *H. wulfsbergii* I.Hagen
- 301 *Isothecium* Brid.
 1 *I. algarvicum* W.E.Nicholson & Dixon
 2 *I. alopecuroides* (Lam. ex Dubois) Isov.
 3 *I. holtii* Kindb.
 4 *I. interludens* Stirt. [*Isothecium myosuroides* var. *brachythecioides* (Dixon) Braithw.]⁵³⁴
- 5 *I. montanum* Draper, Hedenäs, M.Stech, T.Lopes & Sim-Sim⁵³⁵
 6 *I. myosuroides* Brid. [*Isothecium myosuroides* subsp. *brevinerve* Kindb.]⁵³⁶
 7 *I. prolixum* (Mitt.) M.Stech, Sim-Sim, Tangney & D.Quandt [*Echinodium prolixum* (Mitt.) Broth.]⁵³⁷
- Echinodiaceae Broth.
 302 *Echinodium* Jur.
 1 *E. renauldii* (Cardot) Broth.
 2 *E. setigerum* (Mitt.) Jur.
 3 *E. spinosum* (Mitt.) Jur.
- 303 *Pseudomalina* Enroth⁵³⁸
 1 *P. webbiana* (Mont.) Enroth [*Homalia webbiana* (Mont.) Schimp., *Neckera webbiana* (Mont.) Düll]
- Myuriaceae M.Fleisch.⁵³⁹
 304 *Andoa* Ochyra
 1 *A. berthelotiana* (Mont.) Ochyra
- 305 *Ctenidium* (Schimp.) Mitt.
 1 *C. molluscum* (Hedw.) Mitt.⁵⁴⁰
- 306 *Hyocomium* Bruch & Schimp.
 1 *H. armoricum* (Brid.) Wijk & Margad.
- 307 *Myurium* Schimp.
 1 *M. hochstetteri* (Schimp.) Kindb.

⁵²⁸Furness and Gilbert (1980) showed that *Thamnobryum angustifolium* maintains its characters distinct from *Thamnobryum alopecurum* in culture. There are at least three distinct morphological features that distinguish this species from *Thamnobryum alopecurum*. Hodgetts and Blockeel (1992) considered it to be more closely related to *Thamnobryum cataractarum* and the Madeiran *Thamnobryum fernandesii* than to *Thamnobryum alopecurum*. However, more recent molecular work by Olsson et al. (2009) suggests that, while *Thamnobryum angustifolium* is undoubtedly a distinct entity morphologically, colonies of this plant (and of the other narrowly endemic *Thamnobryum* species) may originate from the surrounding subpopulations of *Thamnobryum alopecurum*. In this case, the two subpopulations of *Thamnobryum angustifolium* are independently derived from local *Thamnobryum alopecurum*, and have evolved convergently in response to the rheophilous habitat (Blockeel et al. 2014).

⁵²⁹Olsson et al. (2009) used molecular techniques to suggest that, like *Thamnobryum angustifolium*, *Thamnobryum cataractarum* may be a local derivative of *Thamnobryum alopecurum*, evolving convergently in response to its habitat (Blockeel et al. 2014).

⁵³⁰Recent molecular work by Olsson et al. (2009) suggests that while *Thamnobryum fernandesii* is a distinct entity, colonies may, like *Thamnobryum angustifolium* and *Thamnobryum cataractarum*, originate from surrounding colonies of *Thamnobryum alopecurum*.

⁵³¹Köckinger, Suanjak, et al. (2008), when reporting *Thamnobryum subserratum* from Austria, pointed out that it was first given for Europe by Mastracci (2003), in which Figure 2 (l, m) was drawn from a collection from Latvia (Kurland, leg. Malta, Z.).

⁵³²The new family Heterocradiellaceae was proposed to accommodate *Heterocladium dimorphum*. The taxonomic position of this species is still uncertain, but the genetic differentiation was considered sufficient to place it in a separate genus and family close to Echinodiaceae, Lembophyllaceae and Neckeraceae (Ignatov, Fedorova, et al. 2019).

⁵³³*Heterocladium* has been placed in the Heterocradiaceae Ignatov & Ignatova, but this family name is illegitimate, having previously been used for a family of algae. It was temporarily placed in the Neckeraceae until molecular evidence showed that it is best placed in the Lembophyllaceae (Ignatov, Fedorova, et al. 2019).

⁵³⁴Hodgetts and Vanderpoorten (2018) returned *Isothecium myosuroides* var. *brachythecioides* to species level on the basis of molecular and morphological evidence.

⁵³⁵*Isothecium montanum*, a Madeiran endemic, was described by Draper, Hedenäs, et al. (2015).

⁵³⁶The features characterising *Isothecium myosuroides* subsp. *brevinerve* are very plastic and have no taxonomic value. The authorship of this subspecies is 'Kindb.' rather than 'Lindb.', as it was listed in Hill et al. (2006).

⁵³⁷*Echinodium prolixum* was transferred to *Isothecium* by Stech et al. (2008).

⁵³⁸On the basis of molecular studies, *Pseudomalina* was established as a new genus in the Echinodiaceae to accommodate *Homalia webbiana* (Enroth et al. 2019), a species long suspected not to belong in the Neckeraceae (Olsson et al. 2009).

⁵³⁹*Andoa*, *Ctenidium* and *Myurium* form a well supported clade in Huttunen, Bell, et al. (2012). Furthermore, these three genera plus *Hyocomium* form a clade in Cox et al. (2010).

⁵⁴⁰There continues to be no firm basis for any of the infraspecific taxa of *Ctenidium molluscum*, so the treatment of Hill et al. (2006) is continued.

Anomodontaceae Kindb.

- 308 **Anomodon** Hook. & Taylor [*Anomodontella* Ignatov & Fedosov, *Anomodontopsis* Ignatov & Fedosov, [*Haplohymenium* Schwägr.]⁵⁴¹
- 1 **A. longifolius** (Schleich. ex Brid.) Hartm. [*Anomodontella longifolia* (Schleich. ex Brid.) Ignatov & Fedosov]
 - 2 **A. rugelii** (Müll.Hal.) Keissl. [*Anomodontopsis rugelii* (Müll.Hal.) Ignatov & Fedosov]
 - 3 **A. tristis** (Ces.) Sull. & Lesq. [*Haplohymenium triste* (Ces.) Kindb.]
 - 4 **A. viticulosus** (Hedw.) Hook. & Taylor

Taxa rejected from the covered area

The following taxa have been reported from the area but have been shown not to occur there. Taxa reported from 'Europe' without any further information and which are clearly errors are not included in the list below. There are about 150 liverwort names (mostly subspecific names) with types from Europe that have not been recognised recently, nor synonymized. Some of those names are so old that they may threaten accepted names if their true identity becomes known. Some of the names in *Jungermannia* were enumerated in Söderström, Hagborg, et al. (2016).

Liverworts

Acrobolbus anisodontus (Hook.f. & Taylor) Briscoe [*Tylimanthus anisodontus* (Hook.f. & Taylor) Mitt.] was reported from Madeira by Stephani (1905) but rejected as *Acrobolbus madeirensis* by Grolle and Persson (1966), and from the Azores by Allorge and Allorge (1948) but rejected as *Acrobolbus azoricus*.

Acrobolbos ciliatus (Mitt.) Schiffn. is reported from Britain (coll. D. G. Long 29767, E) by Masuzaki et al. (2010) but this is an error for *Acrobolbus wilsonii* (D. G. Long, pers. comm.).

Acrobolbus laxus (Lehm. & Lindenb.) Briscoe is reported from the Azores and Madeira by Burghardt and Gradstein (2008) but their concept included *Acrobolbus azoricus* and *Acrobolbus madeirensis* to which the reports belong.

Asterella blumeana (Nees) Kachroo is a southeast Asian taxon reported as *Fimbraria blumeana* Nees from Italy by Tassi (1901) but rejected by Aleffi and Schumacker (1995).

Asterella elegans (Spreng.) Trevis. is a South American taxon reported from Corsica as *Fimbraria elegans*

Spreng by Casares-Gil (1919) but this is rejected by Bischler and Jovet-Ast (1973).

Asterella marginata (Nees) S.W.Arnell was reported from Madeira as *Fimbraria marginata* Nees by Gottsche et al. (1846) but this must be rejected. It is a southern African taxon.

Asterella pringlei Underw. is an American taxon reported from Italy as *Fimbraria stahlia* Steph. by Tassi (1901) but this is rejected by Aleffi and Schumacker (1995).

Asterella tenella (L.) P.Beauv. is a North American taxon reported as *Fimbraria tenella* L. from several places in Europe during the 19th century, but all reports are rejected by Long (2006).

Asterella wallichiana (Lehm. & Lindenb.) Grolle is an E Asian taxon reported from Italy as *Fimbraria raddii* Corda ex Nees (type). However, the type is probably from Himalaya and certainly not from Europe (Long 2006).

Bazzania denudata (Lindenb. & Gottsche) Trevis. is a taxon from North America and East Asia reported many times from central and western Europe but rejected from here by Schuster (1969).

Blepharostoma arachnoideum M.Howie is a North American taxon reported from Polar Ural by Zinovjeva (1973) but rejected as *B. trichophyllum* subsp. *brevirete* by Konstantinova and Potemkin (1994).

Cephalozia lucens (A.Evans) Steph. is a taxon from Hawaii reported from Italy by Rhodogher (1896) but rejected by Aleffi (2005).

Cephalozia mammillifera R.M.Schust. & Damsh. is an American taxon reported from 'Europe' by Stotler and Crandall-Stotler (2017). This taxon was treated as a questioned synonym of *Cephalozia varians* var. *scabra* by Damsholt (2002) and if this synonym is correct, the taxon does occur in Europe.

Cephalozia verrucosa Steph. is a taxon from the Southern Hemisphere that has erroneously been reported from Svalbard (Bryhn 1909), probably due to nomenclatural confusion.

Cyathodium spurium (Dicks.) Lindb. ex Braithw. may be conspecific with *Cyathodium cavernarum* (Braithwaite 1878). The type specimen is from Scotland, but it is probably mislabelled as no *Cyathodium* is known from boreal Europe.

Frullania davurica Hampe ex Gottsche, Lindenb. & Nees. has been reported several times from Europe, mostly because it has sometimes included *Frullania jackii* as a subspecies [*F. davurica* subsp. *jackii* (Gottsche) S.Hatt.].

Frullania inflata Gottsche is shown to be a complex species confined in a restricted sense to North America while the European populations must be

⁵⁴¹Ignatov, Fedorova, et al. (2019) established the genera *Anomodontopsis* (for *Anomodon rugelii*) and *Anomodontella* (for *Anomodon longifolius*), and resurrected *Haplohymenium* (for *Anomodon tristis*). Although Ignatov, Fedorova, et al. (2019) show different lineages 'within the Anomodontaceae', the same study gives support for a clade that includes all the species of *Anomodon* s.lat. except *Anomodon attenuatus*, so a relatively conservative approach is adopted here in order to avoid the unnecessary proliferation of small genera.

named *Frullania cleistostoma* (Mamontov, Potemkin, et al. 2018).

Frullania muscicola Steph. is an Asiatic taxon reported from Europe as a putative synonym (variety) of *F. cesatiana* (= *Frullania inflata*; see note under the latter).

Frullania obscurifolia Mitt. is an African taxon reported by Dirkse et al. (1993) but rejected as *Frullania azorica* by González-Mancebo et al. (2008).

Herbertus dicranus (Gottsche, Lindenb. & Nees) Trevis. has (sometimes as *Herbertus sakurii* (Warnst.) S.Hatt.) been reported from Norway and Scotland (Konstantinova 2000a; Hodgetts 2003) due to the inclusion of the now rejected synonymy with *Herbertus borealis*. The unsupported mention of 'Madeira' (Juslén 2006) is an obvious error for Madagascar.

Heteroscyphus integrifolius (Lehm. & Lindenb.) Fulford is a taxon from South America reported from Italy by Rodegher (1896) but rejected by Aleffi (2005).

Jubula hutchinsiae subsp. *javanica* (Steph.) Verd. is an Asiatic taxon. It was supposed to be the subspecies occurring in Caucasus prior to the description of subsp. *caucasica*. It has also erroneously been reported from other parts of Europe and Macaronesia.

Jubula hutchinsiae subsp. *pennsylvanica* (Steph.) Verd. is the North American subspecies that has been reported erroneously from Ireland, France, Macaronesia and Caucasus.

Jungermannia exsertifolia Steph. is an East Asiatic species reported from Europe on the basis that it includes *Jungermannia eucordifolia* as a subspecies, but Mamontov, Konstantinova, et al. (2018) showed that the latter deserves specific rank.

Kurzia makinoana (Steph.) Grolle is a species from North America and East Asia that has been reported from several places in Europe since Grolle (1963) included *Kurzia sylvatica* as synonym. However, this synonymy was rejected by Grolle (1973).

Lejeunea laetevirens Nees & Mont. is a primarily South American taxon but with close affinity to *Lejeunea canariensis*. Unless conspecific with the latter, *Lejeunea laetevirens* is not found in Macaronesia.

Lophocolea humistrata (Hook.f. & Taylor) Gottsche, Lindenb. & Nees is endemic to St Helena but was noted also from Madeira by Stephani (1907).

Lophozia propagulifera (Gottsche) Konstant. & Vilnet is a Southern Hemisphere taxon that was considered conspecific with *Lophozia latifolia* by Bakalin (2005) but the synonymy was rejected by Köckinger (2017) and this is supported by ongoing unpublished molecular research by N. Konstantinova et al.

Metzgeria temperata Kuwah. is a primarily E Asiatic taxon that has been recorded from all around the Northern Hemisphere, including Europe and Macaronesia. However, the European and E Asiatic specimens are not identical (cf. Köckinger 2017) and for the

present the name *Metzgeria consanguinea* is used (see note also under that name).

Microlejeunea diversiloba (Spruce) Müll.Frib. is an American taxon that was described from Mexico under the name *Lejeunea diversifolia* Gottsche and reported from Ireland by Spruce (1876a). However, there exists an older *Lejeunea diversifolia* Mitt. (from the Himalaya), and Spruce (1876b) changed the name to *Lejeunea diversiloba*. It was subsequently reported from the Azores and Ireland by Allorge and Persson (1938). Grolle (1975) concluded that the Irish material was not the same as the Mexican type and re-described it as *Lejeunea hibernica*. All reports outside America should belong here.

Plagiochila javanica (Sw.) Nees & Mont. is a taxon from southeast Asia that Gottsche et al. (1844) reported from the Canary Islands (leg. Webb) but this was rejected by Inoue (1969) who could not locate any specimen supporting the report.

Plagiochila patula (Sw.) Lindenb. is an American species reported from Canary Islands (Dirkse et al. 1993) and Madeira (Nieuwkoop and Arts 1995), both as *Plagiochila dubia* Lindenb. & Gottsche, but this was rejected as *Plagiochila virginica* by Heinrichs, Pröschold et al. (2002).

Plagiochila uniformis Mitt. is a taxon from southeast Asia reported by Stephani (1903) as *Plagiochila ambagiosa* Mitt. from Ireland. Stephani's concept of *P. ambagiosa* was, however, not the same as the taxon described by Mitten, but rather belonging to *Plagiochila spinulosa*.

Porella navicularis (Lehm. & Lindenb.) Pfeiff. is an American taxon reported from several places in central Europe, mostly during the 19th century, but also a few times more recently.

Porella platyphylloidea (Schwein.) Lindb. is frequently reported from Europe. It has sometimes been considered synonymous with *Porella platyphylla* but the study by Heinrichs, Kreier, et al. (2011) shows that the European and American populations of *Porella platyphylla/platyphylloidea* are different and only *Porella platyphylla* occurs in Europe.

Riccia bullosa Link is described with two syntypes, one from South Africa and the other from Portugal. The lectotype is from South Africa and it is not conspecific with the European taxon (Grolle and Long 2000), the latter being *Exormotheca welwitschii*. All reports from Europe are based on this confusion.

Riccia concava Bisch. ex C.Krauss is a taxon from southern Africa reported from the Canary Islands as *Riccia capensis* Steph. by Arnell (1961) and Eggers (1982) but rejected by Perold (1989a).

Riccia limbata Bisch. ex C.Krauss is another taxon from southern Africa reported from the Canary Islands by Arnell (1961) and Eggers (1982) but this is rejected by Perold (1989b) as *Riccia nigrella*.

Riccia oerstediana Lindenb. & Hampe is an American taxon that Schuster (1992b, as *Riccia stenophylla* Spruce) assumed was the taxon reported as fertile *Riccia fluitans* from Europe. However, this has never been verified.

Riella parisii Gottsche is reported twice from Europe, Andalucia in Spain (Müller 1953) and in SE France (Skrzypczak 2001). The Spanish report was rejected by Brugués et al. (2011) and the French report by Hugonnot (2019). It is now known only from Algeria and Tunisia.

Sphaerocarpos texanus Austin is reported as widely distributed in Europe but it is shown by Bell et al. (2013) that the European populations are genetically very distinct from the American populations, although the morphological differences are small. *Sphaerocarpos europaeus* should be used for the European species.

Telaranea nematodes (Gottsche ex Austin) M.Howe is a neotropical-tropical African species which in the past has been reported from several places in south-western Europe and Macaronesia but Engel and Smith Merrill (2004) showed that the European and Macaronesian populations represent a different taxon, *Telaranea europaea*.

Telaranea sejuncta (Ångstr.) S.W.Arnell is an American species that has been treated as a synonym of *Telaranea nematodes* and reported from Europe under this name. Those reports represents *Telaranea europaea*.

Mosses

Amphidium tortuosum (Hornsch.) Cufod. is a widespread species in the tropics. The Macaronesian endemic *Amphidium curvipes* was synonymised with *Amphidium tortuosum* by Frahm et al. (2000), but Sim-Sim et al. (2017) showed that *Amphidium curvipes* is indeed distinct.

Barbula indica (Hook.) Spreng. is a widespread tropical species, which has been transferred to *Hydrogonium*, as *H. orientale* (F.Weber) Kučera, by Kučera, Košnar, et al. (2013). All European records of '*B. indica*' refer to *H. consanguineum* var. *kurilense* (Kučera, Košnar, et al. 2013).

Bartramia stricta Brid. is considered by Damayanti et al. (2012) to be restricted to South America. See footnote on *Bartramia aprica* above.

Brachymenium commutatum (Müll.Hal.) A.Jaeger is excluded following a revision of the specimens from Europe by Ros, Rams, et al. (2007).

Brachytheciastrum fendleri (Sull.) Ochyra & Żarnowiec is a North American species; all European records of *B. fendleri* refer to *B. collinum* (Orgaz et al. 2013).

Brotherella henonii (Duby) M.Fleisch. is an east Asian species; molecular studies are needed to support the proposed synonymy of *Brotherella lorentziana* (Frahm 2013) with this species.

Bryum philonotulum Müll.Hal. is a synonym of *Pohlia philonotula* (Müll.Hal.) Broth., a little-known African species, and not to be confused with *Bryum philonotulum* Hampe, listed here as a synonym of *Bryum kikuyense*.

Cinclidotus pachylomoides Bizot is a mainly Asian species; European records have not been confirmed and it is considered very doubtful that this species occurs in Europe.

Drepanocladus longifolius (Mitt.) Paris is an exclusively Southern Hemisphere species; Satuga et al. (2018) showed that European records of this species should be referred to *Drepanocladus capillifolius*.

Lindbergia brachyptera (Mitt.) Kindb. occurs only in North America (Ignatova, Ignatov, et al. 2010).

Pohlia saphrophila (Müll.Hal.) Broth. has been removed from European list as the European Russian records were based on misidentifications of *Pohlia longicolla* (Czernyadjeva et al. 2017).

Sanionia georgicouninata (Müll.Hal.) Ochyra & Hedenäs is a Southern Hemisphere species distinct from *S. nivalis* (Hedenäs 2012).

Sphagnum lescurii Sull. is a North American species that has been reported from Europe in the past (e.g. by Corley et al. 1981). European records refer to *Sphagnum auriculatum* and *Sphagnum inundatum*.

Sphagnum magellanicum Brid. has been shown to be confined to southern South America (Hassel et al. 2018).

Tortella arctica (Arnell) Crundw. & Nyholm has not been reliably reported from Europe; see note on *T. x cuspidatissima* above.

Taxa only occurring in areas adjacent to the covered area

The following taxa have not yet been found in Europe or northern Macaronesia, but occur just outside the area (in Cape Verde, North Africa, Turkey, Georgia, Azerbaijan, Armenia or western Siberia). Information for liverworts has been obtained from the ELPT database (Söderström, Hagborg, et al. 2019) and for mosses mainly from Ignatov, Afonina, et al. (2006), Kürschner and Erdağ (2005), O'Shea (2006 with updates) and Ros, Mazimpaka, et al. (2013). Some of these taxa are obscure and known only from the original collections; others are mainly American or Pacific species which need to be confirmed in the territories listed.

Liverworts

Acrolejeunea emergens (Mitt.) Steph. (Cape Verde)

Cheilolejeunea xanthocarpa (Lehm. & Lindenb.) Malombe (Cape Verde)

Cyathodium cavernarum Kunze (Cape Verde)

Frullania socotrana Mitt. (Cape Verde)

Frullania spongiosa Steph. (Cape Verde)

Lejeunea capensis Gottsche (Cape Verde)

- Lophozia lacerata* N.Kitag. (Georgia)
Lophozia excisa var. *infusata* (R.M.Schust. & Damsh.) Konstant. & Vilnet (Yamal-Nenets)
Lophozia excisa var. *succulenta* (R.M.Schust. & Damsh.) Konstant. & Vilnet (Yamal-Nenets)
Marchantia debilis K.I.Goebel (Morocco, Egypt)
Marchantia pappeana Lehm. subsp. *pappeana* (Cape Verde)
Plagiochasma eximium (Schiffn.) Steph. (Cape Verde)
Plagiochasma microcephalum (Steph.) Steph. var. *tunesicum* Bischl. (Tunisia)
Pleurozia gigantea (F.Weber) Lindb. (Cape Verde)
Prasanthus jamalicus Potemkin (Yamal-Nenets)
Radula prolifera Arnell (Yamal-Nenets)
Riccia chudoana Steph. (Algeria)
Riccia congoana Steph. (Egypt)
Riccia mamillata Trab. ex Steph. (Algeria)
Riccia polycarpa (Trab.) Jelenc (Algeria)
Riella cyrenaica Maire (Libya)
Riella numidica Trab. (Morocco, Algeria, Libya)
Riella parisii Gottsche (Algeria, Tunisia)
Riella sersuensis Trab. (Algeria)
Scapania brevicaulis Taylor (western Siberia)
Scapania cuspiduligera var. *diplophyllopsis* R.M.Schust. (Yamal-Nenets)
Scapania microdonta (Mitt.) Müll.Frib. (Sverdlovsk)
Solenostoma lignicola (Schiffn.) Váňa, Hentschel & Heinrichs (Turkey)
Solenostoma subtilissimum (Schiffn.) R.M.Schust. (Turkey)
Syzygiella manca (Mont.) Steph. (Cape Verde)
- Mosses**
- Acaulon longifolium* Herrnst. & Heyn (Israel)
Aulacomnium acuminatum (Lindb. & Arnell) Kindb. (Arctic Western Siberia)
Brachymenium acuminatum Harv. (Cape Verde)
Brachymenium commutatum (Müll.Hal.) A.Jaeger (Algeria, Tunisia)
Brachymenium exile (Dozy & Molk.) Bosch. & Sande Lac. (Algeria, Cape Verde, Lebanon)
Brachytheciastrum bellicum (W.R.Buck, J.A.Jiménez, Ros & M.J.Cano) Vanderp., Ignatov, Huttunen & Goffinet (Morocco)
Brachytheciastrum umbilicatum (Jur. & Milde) Orgaz, M.J.Cano & J.Guerra (Turkey)
Bryum anomodon Mont. (Cape Verde)
Bryum atrovirens Brid. (Turkey)
Calohypnum plumiforme (Wilson) Jan Kučera & Ignatov (Georgia)
Cinclidotus bistratosus Kürschner & Lübenau-Nestle (Turkey)
Cinclidotus pachyloma E.S.Salmon (Israel, Lebanon, Turkey)
Cinclidotus vardaranus Erdağ & Kürschner (Turkey)
Crumia latifolia (Kindb.) W.B.Schofield (Armenia)
Didymodon caboverdeanus J.A.Jiménez & M.J.Cano (Cape Verde)
Didymodon tectorum (Müll.Hal.) K.Saito (Egypt)
Entodon pseudoseductrix (Müll.Hal.) A.Jaeger (Cape Verde)
Entodontopsis leucostega (Brid.) W.R.Buck & Ireland (Cape Verde)
Entosthodon angustifolius Jur. & Milde (Jordan, Libya, Turkey)
Entosthodon niloticus Schimp. (Egypt)
Epipterygium rigidum Lindb. ex Broth. (Georgia)
Erpodium grossirete Müll.Hal. (Cape Verde)
Erpodium perrottetii (Mont.) A.Jaeger & Sauerb. (Cape Verde)
Fabronia gueinzii Hampe (Morocco)
Fabronia leikipiae Müll.Hal. (Cape Verde)
Fissidens allorgei P.de la Varde (Cape Verde)
Fissidens androgynus Bruch ex C.Krauss (Cape Verde)
Fissidens dankelmannii Müll.Hal. (Cape Verde)
Fissidens flaccidus Mitt. (Cape Verde)
Fissidens megalotis Schimp. ex Müll.Hal. subsp. *helictocaulos* (Müll.Hal.) Brugg.-Nann. (Cape Verde)
Fissidens sciophyllus Mitt. (Cape Verde)
Fissidens usambaricus Broth. (Cape Verde)
Fontinalis antipyretica var. *heldreichii* (Müll.Hal.) Ruthe (Turkey)
Funaria altissima Dixon (Algeria)
Funaria chevalieri P.de la Varde (Cape Verde)
Groutiella tomentosa (Hornsch.) Wijk & Marg. (Cape Verde)
Gymnostomiella erosula (Müll.Hal. ex Dusén) Arts (Cape Verde)
Gymnostomiella vernicosa (Hook. ex Harv.) M.Fleisch. (Cape Verde)
Herpetineuron toccoae (Sull. & Lesq.) Cardot (Cape Verde)
Hookeria acutifolia Hook. & Grev. (Georgia, Turkey)
Hydrogonium arcuatum (Griff.) Wijk & Margad. (Egypt)
Hydrogonium orientale (F.Weber) Jan Kučera (Cape Verde, Egypt)
Hymenostylium congoanum Dixon & Naveau (Cape Verde)
Hymenostylium hildebrandtii (Müll.Hal.) R.H.Zander (Morocco)
Leptodictyum kurdicum (Schiffn.) Broth. (Turkey)
Leucodon bowringii Mitt. (Turkey)
Leucodon corensis Cardot (Turkey)
Molendoa seravschanica Broth. & Györfy (Arctic Western Siberia)
Orthotrichum urnaceum Müll.Hal. (Armenia, Azerbaijan)
Palamocladium leskeoides (Hook.) E.Britton (Cape Verde)
Perssonia sanguinea Bizot (Cape Verde)
Philonotis laxitexta J.Fröhl. (Lebanon)
Philonotis nanothecioidea Paris & Broth. (Cape Verde)
Physcomitrium immersum Sull. (Egypt)
Physcomitrium niloticum (Delile) Müll.Hal. (Egypt)
Platygyrella densa (Hook.) W.R.Buck (Cape Verde)

Plaubelia sprengelii (Schwägr.) R.H.Zander (Egypt)
Pohlia alba Lindb. & Arnell (Western Siberia)
Pohlia saprophila (Müll.Hal.) Broth. (Kazakhstan)
Pseudoleskeopsis bollei (Broth. & Geh.) P.Rao (Cape Verde)
Pseudoleskeopsis pseudoattenuata (Müll.Hal.) Thér. (Cape Verde)
Rhizomnium striatulum (Mitt.) T.J.Kop. (Turkey)
Schistidium cinclidodonteum (Müll.Hal.) B.Bremer (Morocco)
Sphagnum lescurii Sull. & Lesq. (Turkey, if correct; this is a North American species)
Sphagnum perfoliatum L.I.Savicz (Western Siberia and Arctic Western Siberia)
Splachnobryum aquaticum Müll.Hal. (Jordan)
Splachnobryum limbatum D.H.Norris & R.H.Zander (Egypt)
Syntrichia amphidiacea (Müll.Hal.) R.H.Zander (Cape Verde)
Syntrichia caninervis var. *pseudodesertorum* (Vondr.) M.T.Gallego (Turkey)
Tortella malacophylla (Müll.Hal.) Paris (Turkey)
Tortula acaulon var. *galilaea* (Herrnst. & Heyn) Ros & Herrnst. (Israel)
Tortula grandiretis Broth. (Turkey)
Tortula kneuckeri Broth. & Geh. (Egypt)
Tortula plinthobia (Sull. & Lesq.) Broth. (Egypt)
Weissia breutelii Müll.Hal. (Israel, Turkey)
Weissia leptocarpa Schimp. ex Besch. (Turkey, if accepted as a species – see annotation 160 in Hill et al. 2006)
Weissia ovatifolia Kürschner (Jordan)
Weissia sinaloensis E.B.Bartram (Egypt)

Acknowledgements

Thanks are due to the large number of bryologists whose research, expertise and experience has been drawn on in the preparation of this checklist.

This project would not have been possible, or at least would have been much more difficult, without the initial Red List project, which was financed chiefly through an EC LIFE project. All the bryologists who attended the five workshop meetings or otherwise answered our questions or helped in any way during that project have contributed directly or indirectly to this checklist:

Elvira Baisheva, Ariel Bergamini, Irene Bisang, Patrizia Campisi, Annalena Cogoni, Tomas Hallingbäck, Marko Sabovljević, Norbert Schnyder, Cecilia Sérgio, Olga Afonina, Rosalina Gabriel, César Garcia, Juana González Mancebo, Irina Goldberg, Anders Hagborg, Sanna Huttunen, Elena Ignatova, Marta Infante, Riikka Juutinen, Thomas Kiebacher, Niklas Lönnell, Michael Lüth, Anabela Martins, Oleg Maslovsky, Gordon Rothero, Sorin Ștefănuț, Kimmo Syrjänen, Alain Untereiner, Jiri Váňa[†], Alain Vanderpoorten, Kai Vellak, Michele Aleffi, Jeff Bates, Maude Baudraz, Des Callaghan, Nils Cronberg, Jo Denyer, Jeff Duckett, Heinjo During, Vladimir Fedosov, Anna Ganeva, Piotr Gorski, Urban Gunnarsson, Kristian Hassel, Helena Hesperhol, Mark Hill, Rory Hodd, Kristofer Hylander, Nele Ingerpuu, Sanna Laaka-Lindberg, David Long, Vicente

Mazimpaka, Anna Mežaka, Frank Müller, Jose David Orgaz, Jairo Patiño, Sharon Pilkington, Felisa Puche, Rosa M. Ros, Fred Rumsey, J. G. Segarra-Moragues, Ana Séneca, Adam Stebel, Risto Virtanen, Henrik Weibull, Jo Wilbraham and Jan Żarnowiec.

We are grateful to the following current or former IUCN staff, who led the Red List workshop meetings and did a lot of behind-the-scenes work: David Allen, Marta Cáliz, Eve Englefeld, Catarina Ferreira, Nicholas Fettes, Mariana García Criado and Ana Nieto.

NGH would like to give special thanks to the Irish National Parks and Wildlife Service for providing funding for a year to work on the checklist and to Ed Wymer of the Irish National Parks and Wildlife Service for doing a useful computer check that resulted in the elimination of several errors and inconsistencies. The NTNU, Trondheim, and National Parks and Wildlife Service, Department of Culture, Heritage & the Gaeltacht, Dublin are thanked for funding the Open Access. We are grateful to Phil Stanley for all his help with compiling the index.

ORCID

L. Söderström  <http://orcid.org/0000-0002-9315-4978>
 N.E. Bell  <http://orcid.org/0000-0002-2401-2968>
 R. Garilleti  <http://orcid.org/0000-0002-5977-2908>
 L. Hedenäs  <http://orcid.org/0000-0003-1763-1696>
 J. Kučera  <http://orcid.org/0000-0002-0230-5997>
 F. Lara  <http://orcid.org/0000-0002-1665-5277>

References

- Afonina OM, Ignatova EA, Fedosov VE, Kuznetsova, OI. 2014. Toward a new understanding of *Syntrichia submontana* (Pottiaceae, Bryophyta). *Arctoa*. 23:11–24.
- Aigoín DA, Huttunen S, Ignatov MS, Dirkse GM, Vanderpoorten A. 2009. *Rhynchostegiella* (Brachytheciaceae): molecular re-circumscription of a convenient taxonomic repository. *Journal of Bryology*. 31:213–221.
- Aleffi M. 2005. New check-list of the Hepaticae and Anthocerotae of Italy. *Flora Mediterranea*. 15:485–566.
- Aleffi M, Schumacker R. 1995. Check-list and Red-List of the liverworts (Marchantiophyta) and hornworts (Anthocerotophyta) of Italy. *Flora Mediterranea*. 5:73–161.
- Allen BH. 2014. Maine mosses: Drummondaceae–Polytrichaceae. *Memoirs of the New York Botanical Garden*. 111:1–607.
- Allen BH. 2018. Moss Flora of Central America. Part 4. Fabroniaceae–Polytrichaceae. *Monographs in Systematic Botany from the Missouri Botanical Garden*. 132: i-x, 1–830.
- Allorge V, Allorge P. 1948. Végétation bryologique de l'île de Flores (Açores). *Revue Bryologique & Lichénologique*. 17:126–164.
- Allorge P, Persson H. 1938. Contribution à la flore hépatologique des Îles Açores. *Annales Bryologici*. 11:6–14.
- Alonso M, Jiménez JA, Cano MJ. 2018. New synonyms and typifications in *Chionoloma tenuirostre* (Pottiaceae, Bryophyta). *Phytotaxa*. 373:147–154.
- Alonso M, Jiménez JA, Cano MJ. 2019. Taxonomic revision of *Chionoloma* (Pottiaceae, Bryophyta). *Annals of the Missouri Botanical Garden*. 104:563–632.
- Alonso M, Jiménez JA, Nylander S, Hedenäs L, Cano MJ. 2016. Disentangling generic limits in *Chionoloma*, *Oxystegus*, *Pachyneuropsis* and *Pseudosymblypharis* (Bryophyta: Pottiaceae): an inquiry into their phylogenetic relationships. *Taxon*. 65:3–18.

- Ando H. 1973. Studies on the genus *Hypnum* Hedw. (II). Journal of Science of the Hiroshima University, series B, division 2. 14:165–207.
- Andrus RE. 2006. Six new species of *Sphagnum* (Bryophyta: Sphagnaceae). Sida. 22:959–972.
- Aranda SC, Gradstein SR, Patiño J, Laenen B, Désamoré A, Vanderpoorten A. 2014. Phylogeny, classification and species delimitation in the liverwort genus *Odontoschisma* (Cephaloziaceae). Taxon. 63:1008–1025.
- Arnell SW. 1961. List of hepaticae of the Canary Islands. Svensk Botanisk Tidskrift. 55:379–393.
- Bączkiewicz A, Szczecińska M, Sawicki J, Stebel A, Buczkowska K. 2017. DNA barcoding, ecology and geography of the cryptic species of *Aneura pinguis* and their relationships with *Aneura maxima* and *Aneura mirabilis* (Metzgeriales, Marchantiophyta). PLOS one. 12: e0188837:1–21.
- Bakalin VA. 2001. Notes on *Lophozia* III. Some taxonomic problems in *Lophozia* sect. *Lophozia*. Arctoa. 10:207–218.
- Bakalin VA. 2005. Monograficheskaia obrabotka roda *Lophozia* (Dumort.) Dumort. s. str. [Monograph of the genus *Lophozia* (Dumort.) Dumort. s. str.]. Moscow: Nauka. Russian.
- Bakalin VA. 2016. Notes on *Lophozia* VIII. The Lectotypification of *Lophozia longiflora* (Nees) Schiffn. (Lophoziaceae, Hepaticae). Herzogia. 29:635–643.
- Bakalin VA, Fedosov VE, Fedorova AV, Nguyen VS. 2019. Integrative taxonomic revision of *Marsupella* (Gymnomitriaceae, Hepaticae) reveals neglected diversity in Pacific Asia. Cryptogamie, Bryologie 40:59–85.
- Bakalin VA, Klimova KG. 2016. A note on *Nardia japonica* Steph. (Gymnomitriaceae). Botanica Pacifica. 5:43–50.
- Bakalin VA, Vilnet AA. 2018. A review of the genus *Diplophyllum* (Marchantiophyta) in North and East Asia with the description of a new species (*D. sibiricum*) based on integrative taxonomy. Plant Systematics and Evolution. 304:1269–1287.
- Bakalin VA, Vilnet AA. 2019. *Lophozia fuscovirens* sp. nov. (Lophoziaceae, Marchantiophyta): the second taxon with brown gemmae within *Lophozia* s.s. Nordic Journal of Botany. 37:1–10.
- Bell D, Long DG, Forrest AD, Hollingsworth ML, Blom HH, Hollingsworth PM. 2012. DNA barcoding of European *Herbertus* (Marchantiopsida, Herbertaceae) and the discovery and description of a new species. Molecular Ecology Resources. 12:36–47.
- Bell D, Long DG, Hollingsworth P. 2013. The use of DNA barcoding to address major taxonomic problems for rare British bryophytes. Edinburgh: Royal Botanic Garden.
- Bell NE, Hyvönen J. 2010. A phylogenetic circumscription of *Polytrichastrum* (Polytrichaceae): reassessment of sporophyte morphology supports molecular phylogeny. American Journal of Botany. 97:566–578.
- Biehler JFP. 1807. Plantarum Novarum ex Herbario Sprengelii Centuriam. Halis Saxonum: F.A. Grunerti Patr.
- Bijlsma R, Van der Velde M, Van de Zande L, Boerema AC, Van Zanten BO. 2000. Molecular markers reveal cryptic species within *Polytrichum commune* (common hair-cap moss). Plant Biology. 2:408–414.
- Bisang I, Schumacker R, Sérgio C, Grolle R. 1989. Clé d'identification des espèces du genre *Frullania* Raddi (Hepaticae) en Europe & en Macaronésie. Nuovo Giornale Botanico Italiano. 122:255–266.
- Bischler H, Jovet-Ast S. 1973. Les Hépatiques de Corse. Énumération, notes écologiques & biogéographiques. Revue Bryologique & Lichénologique. 39:43–153.
- Blackstock TH, Blackhall-Miles R, Ruffino L, Watling M. 2019. *Tricholepidozia* (*Telaranea*) *lindenberghii* (Gottsche) E.D.Cooper (Marchantiophyta: Lepidoziaceae), another antipodean liverwort established in Britain. Journal of Bryology. 41:278–280.
- Blockeel TL. 2017. The moss year – 2016. Field Bryology. 117:87–103.
- Blockeel TL, Bosanquet SDS, Hill MO, Preston CD, editors. 2014. Atlas of British and Irish bryophytes. Newbury: Pisces Publications.
- Blockeel TL, Kučera J, Fedosov VE. 2017. *Bryoerythrophyllum duellii* Blockeel (Bryophyta: Pottiaceae), a new moss species from Greece and Cyprus, and its molecular affinities. Journal of Bryology. 39:247–254.
- Blom HH, Bednarek-Ochyra H, Ochyra R. 2016. Studies on *Schistidium* (Grimmiaceae, Bryophyta) in Europe, with particular reference to the Alps: I. A description of *S. marginale* sp. nov. Phytotaxa. 247:210–218.
- Blom HH, Ignatova EA, Afonina OM. 2006. New records of *Schistidium* (Grimmiaceae, Musci) in Russia. Arctoa. 15:187–194.
- Bonfim Santos M, Stech M. 2017. Testing hypotheses on suprageneric relationships and morphological evolution in the Leucobryaceae (Bryophyta). Plant Systematics and Evolution. 303:1383–1397.
- Borovichev EA. 2008. New liverwort records from Murmansk Province. 1. Arctoa 17:191.
- Borovichev EA, Mamontov YS. 2017. Hepaticae Rossicae Exsiccatae. Fasc. XIII. Apatity: Russian Academy of Science, Kola Science Centre, Polar-Alpine Botanical Garden Institute.
- Bosanquet SDS, Lara F. 2012. *Orthotrichum cambrense* sp. nov. (Orthotrichaceae), a distinctive moss from Wales, United Kingdom. Cryptogamie, Bryologie. 33:329–339.
- Braithwaite R. 1878. *Riccia spuria*, Dickson. Journal of Botany, British and Foreign. 16:55.
- Bridel SE. 1827. Bryologia Universa, seu systematica ad novam methodum dispositio, historia et descriptio omnium muscorum frondosorum hucusque cognitorum cum synonymia ex auctoribus probatissimis. Vol. 1, pars 2. Leipzig: J. A. Barth.
- Brugués M, Infante M, Cros RM. 2011. Anotaciones sobre hepáticas de España. Boletín de la Sociedad Española de Biología. 36:3–8.
- Brugués M, Sérgio C. 2010. On the identity of some *Entosthodon* species endemic to Macaronesia or the north of Africa. Cryptogamie, Bryologie. 31:277–280.
- Bryhn N. 1909. Bryophyta pro flora Spitsbergensi nova. Nyt Magazin for Naturvidenskaberne. 47:207–208.
- Buch H. 1929. Eine neue moosystematische Methodik. In: Winge, Ø, editor. Beretning om, det 18. Skandinaviske naturforsker møde i København. 26–31 August 1929. København: Frederiksberg Bogtrykkeri; p. 225–229.
- Buch H. 1932. Vorarbeiten zu einer Lebermoosflora Fennoscandias. I. Ein Versuch zur Aufspaltung der Gattungen *Lophozia* Dum. und *Sphenolobus* Steph. Memoranda Societatis pro Fauna & Flora Fennica. 8:282–297.
- Buchbender V, Hespanhol H, Krug M, Sérgio C, Séneca A, Mau K, Hedenäs L, Quandt D. 2014. Phylogenetic reconstructions of the Hedwigiaceae reveal cryptic speciation and hybridisation in *Hedwigia*. Bryophyte Diversity and Evolution. 36:1–21.
- Buczkowska K, Bakalin VA, Bączkiewicz A, Agüero B, Gonera P, Ślipiko M, Szczecińska M, Sawicki J. 2018. Does *Calypogeia azurea* (Calypogeiaceae, Marchantiophyta) occur outside Europe? Molecular and morphological evidence. PLoS One. 13:1–27.
- Burghardt M, Gradstein SR. 2008. A revision of *Tylimanthus* (Acrobolbaceae, Marchantiophyta) in Tropical America,

- Africa, and Macaronesia. *Fieldiana: Botany* (n.ser.). 47:199–210.
- Burrell WH. 1911. *Lophozia schultzii* (Nees) Schiffn. var. nov. *laxa*. *Journal of Botany*. 49:217–219.
- Callaghan DA. 2019. Typification and diagnosis of *Weissia x mittenii* (Bruch & Schimp.) Mitt. emend. A.J.E.Sm. (*Weissia multcapsularis* x *W. rostellata*) (Pottiaceae, Bryophyta). *Journal of Bryology*. 41:243–248.
- Callaghan DA, Bell NE, Forrest LL. 2019. Taxonomic notes on *Weissia* subgenus *Astomum*, including *Weissia wilsonii* D.A.Callaghan, a new species from Europe. *Journal of Bryology*. 41:135–148.
- Callaghan DA, Masson J, During H. 2020. *Physcomitrium* × *stevensonii* D.A.Callaghan (*Physcomitrium patens* × *P. eurystomum*) (Funariaceae, Bryophyta), a new name for a rarely recorded hybrid moss. *Journal of Bryology*. 42. <https://doi.org/10.1080/03736687.2020.1725325>
- Câmara PEAS, Carvalho-Silva M, Henriques DK, Guerra J, Gallego MT, Poveda DR, Stech M. 2018. Pylaisiaceae Schimp. (Bryophyta) revisited. *Journal of Bryology*. 40:251–264.
- Cano MJ. 2007. Typification of the names of some infraspecific taxa in the *Tortula subulata* complex (Pottiaceae, Bryophyta) and their taxonomic disposition. *Taxon*. 56:949–952.
- Cano MJ. 2008. Taxonomic revision of *Hennediella* Paris (Pottiaceae, Bryophyta). *Bryophytorum Bibliotheca*. 64:1–142.
- Caparrós R, Lara F, Draper I, Mazimpaka V, Garilleti R. 2016. Integrative taxonomy sheds light on an old problem: the *Ulota crispata* complex (Orthotrichaceae, Musci). *Botanical Journal of the Linnean Society*. 180:427–451.
- Casares-Gil A. 1919. *Flora Ibérica Briófitas, Hepáticas*. Madrid: Museo Nacional de Ciencias Naturales.
- Cezón K, Muñoz J, Hedenäs L, Huttunen S. 2010. *Rhynchostegium confusum*, a new species from the Iberian Peninsula and its relation to *R. confertum* based on morphological and molecular data. *Journal of Bryology*. 32:1–8.
- Chavoutier L, Hugonnot V. 2013. *Mousses, hépatiques & anthocérotes du département de la Savoie (France)*. Sevrier: FMBDS.
- Cole TCH, Hilger HH, Goffinet B. 2019. Bryophyte phylogeny poster (BPP). *PeerJ Preprints* 7: e27571v1.
- Cooper ED, Söderström L, Hagborg A, von Konrat MJ. 2013. Notes on Early Land Plants Today. 38. New combinations and synonyms in Lepidoziaceae (Marchantiophyta). *Phytotaxa* 97:52–62.
- Corley MFV, Crundwell AC, Düll R, Hill MO, Smith AJE. 1981. Mosses of Europe and the Azores; an annotated list of species, with synonyms from the recent literature. *Journal of Bryology*. 11:609–689.
- Cox CJ, Goffinet B, Wickett NJ, Boles SB, Shaw AJ. 2010. Moss diversity: a molecular phylogenetic analysis of genera. *Phytotaxa*. 9:175–195.
- Crandall-Stotler B, Stotler RE. 2007. On the identity of *Moerckia hibernica* (Hook.) Gottsche (Moerckiaceae fam. nov., Marchantiophyta). *Nova Hedwigia, Beihefte*. 131:41–59.
- Cros RM, Sáez L, Roselló JA. 2005. *Plagiochasma appendiculatum* Lehm. & Lindenb. (Marchantiales, Ayttoniaceae), a species new to the European bryophyte flora. *Journal of Bryology*. 27:3–6.
- Crosby MR, Buck WR. 2011. *Nogopterium*, a new name for the genus *Pterogonium* (Musci, Leucodontaceae). *Novon*. 21:424–425.
- Crozals A. 1903. *Riccia subbifurca* Warn. in litt. *Revue Bryologique*. 30:62–64.
- Czernyadjeva IV, Kuznetsova OI, Ignatov MS. 2017. On *Pohlia saprophila* (Mielichhoferiaceae, Bryophyta). *Arctoa*. 26:181–186.
- Dalton NJ, Kungu EM, Long DG. 2012. The misapplication of *Hedwigia integrifolia* P.Beauv. and identity of *Gymnostomum imberbe* Sm. (Hedwigiaceae, Bryopsida). *Journal of Bryology*. 34:59–61.
- Damayanti L, Muñoz J, Wicke S, Symmank L, Shaw B, Frahm J-P, Quandt D. 2012. Common but new: *Bartramia rosamrosiae*, a ‘new’ widespread species of apple mosses (Bartramiales, Bryophytina) from the Mediterranean and western North America. *Phytotaxa*. 73:37–59.
- Damsholt K. 1994. On the identity of *Jungermannia groenlandica* Nees. *Journal of the Hattori Botanical Laboratory*. 75:173–178.
- Damsholt K. 2002. *Illustrated flora of Nordic liverworts and hornworts*. Lund: Nordic Bryological Society.
- Damsholt K. 2013. *The liverworts of Greenland*. Lund: Nordic Bryological Society.
- Damsholt K. 2017. The complex liverwort flora of the Faeroe Isles. *Lindbergia*. 40:14–38.
- De Luna E. 2016. Typification, taxonomy and worldwide distribution of *Braunia secunda* (Hook.) Bruch, Schimp. (Hedwigiaceae). *Journal of Bryology*. 38:286–294.
- De Roo RT, Hedderson TA, Söderström L. 2007. Molecular insights into the phylogeny of the leafy liverwort family Lophoziaceae Cavers. *Taxon*. 56:301–314.
- De Sloover JL 1959. Considérations sur la valeur spécifique de *Moerckia flotoviana* (Nees) Schiffn., Dilaenacée nouvelle pour la flore Belge. *Bulletin du Jardin Botanique de l’État, Bruxelles*. 29:157–181.
- Delgadillo-Moya C. 2015. *Grimmia* (Grimmiaceae, Bryophyta) in the Neotropics. México City: Universidad Nacional Autónoma de México.
- Dias E, Mendes C, Shaw J. 2009. *Sphagnum recurvum* P.Beauv. on Terceira, Azores, new to Macaronesia-Europe. *Journal of Bryology*. 31:199–201.
- Dirkse GM, Bouman AC, Losada-Lima A. 1993. Bryophytes of the Canary Islands, an annotated checklist. *Cryptogamie, Bryologie, Lichénologie*. 14:1–47.
- Dirkse GM, Brugués M. 2010. *Entosthodon kroonkurk* (Bryophyta: Funariaceae), a new species from the Iberian Peninsula and Macaronesia. *Journal of Bryology*. 32:133–139.
- Dirkse GM, Losada-Lima A, Stech M. 2016. *Riccia boumanii* Dirkse, Losada & M.Stech sp. nov. (Ricciaceae, Marchantiophyta) in the Canary Islands, the first species of *Riccia* subgenus *Riccia* section *Pilifer* Volk outside South Africa. *Journal of Bryology*. 38:94–102.
- Dixon HN. 1912. *Eucladium verbanum* Nicholson and Dixon, sp. nov. *Revue Bryologique*. 39:89–92.
- Draper I, González-Mancebo JM, Werner O, Patiño J, Ros RM. 2011. Phylogeographic relationships between the mosses *Exsertotheca intermedia* from Macaronesian islands and *Neckera baetica* from southern glacial refugia of the Iberian Peninsula. *Annales Botanici Fennici* 48:133–141.
- Draper I, Hedenäs L. 2008. *Sciuro-hypnum tromsoeense* (Kaurin, Arnell) Draper & Hedenäs, a distinct species from the European mountains. *Journal of Bryology*. 30:271–278.
- Draper I, Hedenäs L. 2009. *Sciuro-hypnum dovrense* (Limpr.) Draper & Hedenäs comb. nov., a distinct Eurasian alpine species. *Cryptogamie, Bryologie*. 30:289–299.
- Draper I, Hedenäs L, Stech M, Patiño J, Werner O, González-Mancebo JM, Sim-Sim M, Lopes T, Ros RM. 2015. How many species of *Isoetecium* (Lembophyllaceae, Bryophyta) are there in Macaronesia? A survey using integrative taxonomy. *Botanical Journal of the Linnean Society*. 177:418–438.

- Eckel P. 1998. A re-evaluation of *Tortella* (Musci, Pottiaceae) in conterminous USA and Canada, with a treatment of the European species *Tortella nitida*. *Bulletin of the Buffalo Society of Natural Sciences*. 36:117–191.
- Eckel P. 2010. *Tortella bambergeri* in North America and an evaluation of its taxonomy. *Bulletin of the Buffalo Society of Natural Sciences*. 39:1–10.
- Eckel P. 2012. The genus *Elodium* (Bryophyta, Helodontiaceae): a history and new combinations. *Phytoneuron*. 70:1–7.
- Eckstein J, Garilleti R, Lara F. 2018. *Lewinskya transcaucasica* (Orthotrichaceae, Bryopsida) sp. nov. A contribution to the bryophyte flora of Georgia. *Journal of Bryology*. 40:31–38.
- Eggers J. 1982. Artenliste der Moose Makaronesiens. *Cryptogamie, Bryologie, Lichénologie*. 3:283–335.
- Ellis LT, Afonina OM, Andriamiarisoa RL, Asthana G, Bharti R, Aymerich P, Bambe B, Boiko M, Brugués M, Ruiz E, et al. 2018. New national and regional records, 56. *Journal of Bryology*. 40:271–296.
- Ellis LT, Akhoondi Darzikolaei S, Shirzadian S, Bakalin VA, Bednarek-Ochyra H, Ochyra R, Claro D, Dulin MV, Eckel PM, Erzberger P, et al. 2011. New national and regional bryophyte records, 29. *Journal of Bryology*. 33:316–323.
- Ellis LT, Aleffi M, Alegro A, Segota V, Asthana AK, Gupta R, Singh VJ, Bakalin VA, Bednarek-Ochyra H, Cykowska-Marzencka B, et al. 2016. New national and regional bryophyte records, 48. *Journal of Bryology*. 38:235–259.
- Ellis LT, Amélio LA, Peralta DF, Bačkor M, Baisheva EZ, Bednarek-Ochyra H, Burghardt M, Czernyadjeva IV, Kholod SS, Potemkin AD, et al. 2019. New national and regional bryophyte records, 59. *Journal of Bryology*. 41:177–194.
- Ellis LT, Price MJ. 2015. Review of the type specimens of species described by J. Hedwig in *Phascum* Hedw. (Pottiaceae). *Journal of Bryology*. 37:23–41.
- Engel JJ, Smith Merrill GL. 2004. Austral hepaticae. 35. A taxonomic and phylogenetic study of *Telaranea* (Lepidoziaceae), with a monograph of the genus in temperate Australasia and commentary on extra-Australasian taxa. *Fieldiana: Botany (n.ser.)*. 44:1–265.
- Enroth J, Olsson S, Huttunen S, Buchbender V, Tangney R, Stech M, Hedenäs L, Quandt D. 2019. Orthostichellaceae fam. nov. and other novelties in pleurocarpous mosses revealed by phylogenetic analyses. *The Bryologist*. 122:219–245.
- Fedosov VE. 2012. *Encalypta* Sect. *Rhabdotheca* in Russia. *Arctoa*. 21:101–112.
- Fedosov VE, Doroshina GYa. 2018. Orthotrichales. In: Ignatov MS, editor. *Moss flora of Russia*. Vol. 4. *Bartramiales–Aulacomniales*. Moscow: KMK Scientific Press Ltd. (Arctoa. 27, supplement 1); p. 67–195.
- Fedosov VE, Doroshina GYa, Tubanova DY, Afonina OM, Ignatova EA. 2017. Four Orthotrichaceae species new for Russian flora. *Arctoa*. 26:154–165.
- Fedosov VE, Fedorova AV, Fedosov AE, Ignatov MS. 2016. Phylogenetic inference and peristome evolution in haplolepidous mosses, focusing on Pseudoditrichaceae and Ditrichaceae s.l. *Botanical Journal of the Linnean Society*. 181:139–155.
- Fedosov VE, Fedorova AV, Ignatova EA. 2017a. On the taxonomic position of the genera *Brachydontium* Fűrnr. and *Campylostelium* Bruch & Schimp. (Bryophyta, Grimmiales). *Journal of Bryology*. 39:161–170.
- Fedosov VE, Fedorova AV, Ignatova EA. 2017b. On the two poorly known *Orthotrichum* species from North Asia. *Arctoa*. 26:144–153.
- Fedosov VE, Fedorova AV, Ignatova EA, Ignatov MS. 2017. A revision of the genus *Seligeria* (Seligeriaceae, Bryophyta) in Russia inferred from molecular data. *Phytotaxa*. 323:27–50.
- Fedosov VE, Fedorova AV, Troitsky AV, Bobrova VK, Ignatov MS. 2016. On the systematic position of *Hymenoloma* (Bryophyta). *Arctoa*. 25:119–130.
- Fedosov VE, Ignatova EA. 2010. *Orthotrichum dagestanicum* sp. nov. (Orthotrichaceae, Musci) – a new species from Dagestan (Eastern Caucasus). *Arctoa*. 19:69–74.
- Fedosov VE, Ignatova EA, Ivanov OV, Doroshina DY. 2010. On the genus *Entosthodon* (Funariaceae, Musci) in the Caucasus. *Arctoa*. 19:75–86.
- Feldberg K, Groth H, Wilson R, Schäfer-Verwimp A, Heinrichs J. 2004. Cryptic speciation in *Herbertus* (Herbertaceae, Jungermanniopsida): range and morphology of *Herbertus sendtneri* inferred from nr ITS sequences. *Plant Systematics and Evolution*. 249:247–261.
- Flatberg KI. 2008. *Sphagnum venustum* (Bryophyta), a noticeable new species in sect. *Acutifolia* from Labrador, Canada. *Lindbergia*. 33:2–12.
- Frahm J-P. 2005. New or interesting records of bryophytes from the Azores. *Tropical Bryology*. 26:45–48.
- Frahm J-P. 2006. *Frullania tamarisci* var. *azorica* (Jubulaceae, Marchantiopsida), a new taxon from the Azores. *Tropical Bryology*. 27:101–105.
- Frahm J-P. 2013. Are there endemic bryophyte species in the Alps? The range of *Brotherella lorentziana* (Lor.) Loeske. *Archive for Bryology*. 183:1–7.
- Frahm J-P, Klöcker T, Schmidt R, Schröter C. 2000. Revision der Gattung *Amphidium* (Musci, Dicranaceae). *Tropical Bryology*. 18:173–184.
- Frahm J-P, Sabovljević M. 2006. Preliminary results of the taxonomic value of *Tortula densa* (Velen.) J-P. Frahm inferred from the Internal Transcribed Spacer (ITS) of the nrDNA. *Cryptogamie, Bryologie*. 27:405–412.
- Frahm J-P, Walsemann E. 1973. Nachträge zur Moosflora von Schleswig-Holstein. *Mitteilungen Arbeitsgemeinschaft Geobotanik in Schleswig-Holstein und Hamburg*. 23:1–203.
- Frey W, Stech M. 2009. Marchantiophyta, Bryophyta, Anthocerotophyta. In: Frey W, editor. *Syllabus of plant families*. A. Engler's *Syllabus der Pflanzenfamilien*, 13th edn, Part 3. Bryophytes and seedless vascular plants. Stuttgart: Gebr. Borntraeger Verlagsbuchhandlung; p. 13–263.
- Furness SB, Gilbert OL. 1980. The status of *Thamnobryum angustifolium* (Holt) Crundw. *Journal of Bryology*. 11:139–144.
- Gallego MT, Cano MJ, Ros RM, Guerra J. 2002. An overview of *Sytrichia ruralis* complex (Pottiaceae: Musci) in the Mediterranean region and neighbouring areas. *Botanical Journal of the Linnean Society*. 138:209–224.
- Gallego MT, Hugonnot V, Cano MJ. 2018. Taxonomic resurrection of an awnless variety of *Sytrichia ruralis* and comparison with other European muticous taxa in this genus. *Journal of Bryology*. 40:244–250.
- Garcia C, Lara F, Sérgio C, Sim-Sim M, Garilleti R, Mazimpaka V. 2006. *Zygodon catarinoi* (Orthotrichaceae, Bryopsida), a new epiphytic species from the Western Mediterranean Basin. *Nova Hedwigia*. 82:247–256.
- Gardiner A, Ignatov MS, Huttunen S, Troitsky A. 2005. On resurrection of the families Pseudoleskeaceae Schimp. and Pylaisiaceae Schimp. (Musci, Hypnales). *Taxon* 54:651–663.
- Geissler P. 1985. Notulae Bryofloristicae Helveticae II. *Candollea*. 40:193–200.
- Goffinet B, Buck WR. 2011. *Physcomitridium readeri* is the correct name for *Ephemerella readeri*. *The Bryologist*. 114:545–546.
- Goffinet B, Buck WR, Shaw AJ. 2009. Morphology, anatomy and classification of the Bryophyta. In: Goffinet B, Shaw

- AJ, editors. *Bryophyte Biology*, 2nd edition. Cambridge: Cambridge University Press; p. 55–138.
- Goffinet B, Budke JM, Newman LC. 2011. Micromitriaceae: a new family of highly reduced mosses. *Taxon*. 60:1245–1254.
- Goffinet B, Shaw AJ, Cox CJ, Wickett NJ, Boles S. 2004. Phylogenetic inferences in the Orthotrichoideae (Orthotrichaceae: Bryophyta) based on variation in four loci from all genomes. *Monographs in Systematic Botany from the Missouri Botanical Garden*. 98:270–289.
- González-Mancebo JM, Romaguera F, Ros RM, Patiño J, Werner O. 2008. Bryophyte flora of the Canary Islands: an updated compilation of the species list with an analysis of distribution patterns in the context of the Macaronesian Region. *Cryptogamie, Bryologie*. 29:315–357.
- Gottsche CM, Lindenberg JBW, Nees CG. 1844. *Synopsis hepaticarum*, fasc. 1. Hamburg: Meissner; p. 1–144.
- Gottsche CM, Lindenberg JBW, Nees CG. 1846. *Synopsis hepaticarum*, fasc. 4. Hamburg: Meissner; p. 465–624.
- Grims F. 1999. Die Laubmoose Österreichs. *Catalogus Florae Austriae*, II. Teil, Bryophyten (Moose), Heft 1, Musci (Laubmoose). Wien: Österreichische Akademie der Wissenschaften.
- Grolle R. 1963. Über *Kurzia v. Martens*. *Revue Bryologique & Lichénologique*. 32:166–180.
- Grolle R. 1964. *Miscellanea hepaticologica* 11–20. *Journal of Japanese Botany*. 39:173–178.
- Grolle R. 1967. *Miscellanea hepaticologica* 71–80. *Transactions of the British Bryological Society*. 5:271–282.
- Grolle R. 1973. *Miscellanea hepaticologica* 131–140. *Herzogia*. 3:75–82.
- Grolle R. 1975. *Miscellanea hepaticologica* 151–160. *Lindbergia*. 3:47–56.
- Grolle R, Long DG. 2000. An annotated check-list of the Hepaticae and Anthocerotae of Europe and Macaronesia. *Journal of Bryology*. 22:103–140.
- Grolle R, Persson H. 1966. Die Gattung *Tylimanthus* auf den Atlantischen Inseln. *Svensk Botanisk Tidskrift*. 60:164–174.
- Grolle R, So ML. 2003. *Riccia fruticulosa* O.F.Müll., 1782 and blue *Metzgeria* (Marchantiophyta) in Europe. *Botanical Journal of the Linnean Society*. 142:229–235.
- Grundmann M, Schneider H, Russell SJ, Vogel JC. 2006. Phylogenetic relationships of the moss genus *Pleurochaete* Lindb. (Bryales: Pottiaceae) based on chloroplast and nuclear genomic markers. *Organisms Diversity and Evolution*. 6:33–45.
- Guerra J. 1985. *Orthotrichum bistratosum* (Schiffn.) Guerra, comb. & stat. nov. *Anales Jardín Botánico de Madrid*. 42:284.
- Guerra J. 2014. Fontinalaceae. In: Guerra J, Cano MJ, Brugués M, editors. *Flora Briofítica Ibérica*. Vol. V. Murcia: Universidad de Murcia, Sociedad Española de Briología; p. 155–163.
- Guerra J. 2016. Notes on the genus *Eurhynchiastrum* (Bryophyta, Brachytheciaceae) in the Iberian Peninsula. *Nova Hedwigia*. 102:359–365.
- Guerra J. 2018. Hypnales. In: Guerra J, Cano MJ, Brugués M, editors. *Flora Briofítica Ibérica*. Vol. VI. Murcia: Universidad de Murcia, Sociedad Española de Briología; p. 365.
- Guerra J, Brugués M, Cano MJ, Cros RM, editors. 2006. *Flora Briofítica Ibérica*. Vol. III, Pottiaceae, Encalyptaceae, Bartramiaceae. Murcia: Universidad de Murcia, Sociedad Española de Briología.
- Guerra J, Gallego MT, Jiménez JA, Cano MJ. 2010. *Bryum* Hedw. In: Guerra, J, Brugués M, Cano MJ, Cros RM, editors. *Flora Briofítica Ibérica*. Vol. IV Murcia: Universidad de Murcia & Vitoria: Sociedad Española de Briología; p. 105–178.
- Guerra J, Jiménez-Martínez JF, Cano MJ, Alonso, M, Gallego MT. 2019. *Schistidium convergens* (Grimmiaceae, Bryophyta), a new species from southern Spain and Morocco. *Nova Hedwigia*. 109:65–80.
- Guerra J, Jiménez-Martínez JF, Cano MJ, Brugués M. 2008. Molecular and morphological evidence on the status of *Bryum mildeanum* Jur. and records revision in the Iberian Peninsula. *Journal of Bryology*. 30:297–302.
- Guerra J, Jiménez-Martínez JF, Cano MJ, Jiménez-Fernández JA. 2011. A contribution to the phylogenetic study of Mielichhoferiaceae-Mniaceae (Bryophyta) based on molecular sequence data. *Nova Hedwigia*. 93:47–56.
- Guerra J, Jiménez-Martínez JF, Jiménez, JA. 2010. *Neckera baetica* sp. nov. (Neckeraceae, Bryophyta) from southern Spain, based on morphological and molecular data. *Nova Hedwigia*. 91:255–263.
- Guerra J, Martínez M, Jiménez JA, Cano MJ, Gallego MT. 2020. A new species of moss emerges from molecular and morphological data: *Schistidium memnonium* sp. nov. (Grimmiaceae, Bryophyta). *Plant Biosystems*. <https://doi.org/10.1080/11263504.2020.1762789>.
- Guerra J, Ríos D, Gallego MT. 2014. A revision of the genus *Rhynchostegiella* (Brachytheciaceae) in the Iberian Peninsula and Balearic Islands. *Phytotaxa*. 183:193–200.
- Hampe E. 1872. *Symbolae ad floram Brasiliae centralis cognoscendam*. Musci frondosi. *Vidensk. Meddel. Dansk Naturhist. Foren. Kjøbenhavn* 4:36–59.
- Hanssen L, Sæstad SM, Flatberg KI. 2000. Population structure and taxonomy of *Sphagnum cuspidatum* and *S. viride*. *The Bryologist*. 103:93–103.
- Hassel K, Kyrkjeeide MO, Yousefi N, Prestø T, Stenøien HK, Shaw JA, Flatberg KI. 2018. *Sphagnum divinum* (sp. nov.) and *S. medium* Limpr. and their relationship to *S. magellanicum* Brid. *Journal of Bryology*. 40:197–222.
- Hattori S. 1972. *Frullania tamarisci* complex and the species concept. *Journal of the Hattori Botanical Laboratory*. 35:202–251.
- Hedderson TA, Blockeel TL. 2006. *Oncophorus dendrophilus*, a new moss species from Cyprus and Crete. *Journal of Bryology*. 28:357–359.
- Hedderson TA, Murray DJ, Cox CJ, Nowell TL. 2004. Phylogenetic relationships of haplolepidous mosses (Dicranidae) inferred from rps4 gene sequences. *Systematic Botany*. 29:29–41.
- Hedderson TA, Zander RH. 2008. *Chenia ruigtevelea* (Pottiaceae), a new moss species from the Western Cape Province of South Africa. *The Bryologist*. 111:496–500.
- Hedenäs L. 1994. The *Hedwigia ciliata* complex in Sweden, with notes on the occurrence of the taxa in Fennoscandia. *Journal of Bryology*. 18:139–157.
- Hedenäs L. 2006. Additional insights into the phylogeny of *Calliergon*, *Loeskygnum*, *Straminergon* and *Warnstorfia* (Bryophyta: Calliergonaceae). *Journal of the Hattori Botanical Laboratory*. 100:125–134.
- Hedenäs L. 2010. ITS distinguishes *Palustriella falcata* from *P. commutata* and *P. decipiens*, but not from *P. pluristratosa* (Bryophyta: Amblystegiaceae). *Nova Hedwigia, Bieheft*. 138:51–60.
- Hedenäs L. 2012. Global phylogeography in *Sanionia uncinata* (Amblystegiaceae: Bryophyta). *Botanical Journal of the Linnean Society*. 168:19–42.
- Hedenäs L. 2017a. Scandinavian *Oncophorus* (Bryopsida, Oncophoraceae): species, cryptic species, and intraspecific variation. *European Journal of Taxonomy*. 315:1–34.
- Hedenäs L. 2017b. *Brachythecium tauriscorum* Molendo, a widely distributed Arctic-alpine species. *Journal of Bryology*. 39:201–203.
- Hedenäs L. 2018. *Oncophorus demetrii*, a fifth Scandinavian species of *Oncophorus* (Musci) possible to recognize by morphology. *Lindbergia*. 41:1–9.

- Hedenäs L. 2020. Disentangling Scandinavian species hidden within *Meesia uliginosa* Hedw. s.l. (Bryophyta, Meesiaceae). *Lindbergia*. 42(1):1–14. DOI: 10.25227/linbg.01125
- Hedenäs L, Désamoré A, Laenen B, Papp B, Quandt D, González-Mancebo JM, Patiño J, Vanderpoorten A, Stech M. 2014. Three species for the price of one within the moss *Homalothecium sericeum* s.l. *Taxon*. 63:249–257.
- Hedenäs L, Heinrichs J, Gallego MT. 2019. The Scandinavian *Syntrichia ruralis* complex (Musci, Pottiaceae): a chaos of diversification. *Plant Systematics and Evolution*. DOI: 10.1007/s00606-019-01596-0
- Hedenäs L, Rosborg C. 2008. *Pseudocalliergon* is nested within *Drepanocladus* (Bryophyta: Amblystegiaceae). *Lindbergia*. 33:67–74.
- Heinrichs J, Dong S, Schäfer-Verwimp A, Pócs T, Feldberg K, Czumaj A, Schmidt AR, Reitner J, Renner MAM, Hentschel J, et al. 2013. Molecular phylogeny of the leafy liverwort *Lejeunea* (Porellales): evidence for a Neotropical origin, uneven distribution of sexual systems and insufficient taxonomy. *PLOS one*. 8(12;E82547):1–14.
- Heinrichs J, Hentschel J, Bombosch A, Fiebig A, Reise J, Edelmann M, Kreier HP, Schäfer-Verwimp A, Caspari S, Schmidt AR, et al. 2010. One species or at least eight? Delimitation and distribution of *Frullania tamarisci* (L.) Dumort. s. l. (Jungermanniopsida, Porellales) inferred from nuclear and chloroplast DNA markers. *Molecular Phylogenetics and Evolution*. 56:1105–1114.
- Heinrichs J, Kreier HP, Feldberg K, Schmidt AR, Zhu RL, Shaw B, Shaw AJ, Wiessman V. 2011. Formalizing morphologically cryptic biological entities: new insights from DNA taxonomy, hybridization, and biogeography in the leafy liverwort *Porella platyphylla* (Jungermanniopsida, Porellales). *American Journal of Botany*. 98:1252–1262.
- Heinrichs J, Pröschold T, Renker C, Groth H, Rycroft DS. 2002. *Plagiochila virginica* A.Evans rather than *P. dubia* Lindenb. & Gottsche occurs in Macaronesia; placement in sect. *Contigua* Carl is supported by ITS sequences of nuclear ribosomal DNA. *Plant Systematics and Evolution*. 230:221–230.
- Heinrichs J, Rycroft DS, Groth H, Cole WJ. 2002. Morphological and phytochemical studies of *Plagiochila papillifolia* Steph., a Neotropical liverwort new to Europe. *Journal of Bryology*. 24:119–126.
- Hentschel J, von Konrat M, Söderström L, Hagborg A, Larraín J, Sukkharak P, Uribe J, Zhang L. 2015. Notes on Early Land Plants Today. 72. Infrageneric classification and new combinations, new names, new synonyms in *Frullania*. *Phytotaxa*. 220:127–142.
- Hernández-Maqueda R, Quandt D, Muñoz J. 2008. Testing reticulation and adaptive convergence in the Grimmiaceae (Bryophyta). *Taxon*. 57:500–510.
- Hill MO, Bell N, Bruggeman-Nannenga MA, Brugués M, Cano MJ, Enroth J, Flatberg KI, Frahm J-P, Gallego MT, Garilleti R, et al. 2006. An annotated checklist of the mosses of Europe and Macaronesia. *Journal of Bryology*. 28:198–267.
- Hill MO, Blackstock TH, Long DG, Rothero GP. 2008. A checklist and census catalogue of British and Irish bryophytes. Middlewich: British Bryological Society.
- Hodgetts NG. 2003. Some synonyms of *Herbertus dicranus* (Taylor ex Gottsche et al.) Trevis. *Journal of Bryology*. 25:138–140.
- Hodgetts NG. 2015. Checklist and country status of European bryophytes – towards a new Red List for Europe. *Irish Wildlife Manuals* 84. Dublin: National Parks & Wildlife Service, Department of Arts, Heritage and the Gaeltacht, Ireland.
- Hodgetts NG, Blockeel TL. 1992. *Thamnobryum cataractarum*, a new species from Yorkshire, with observations on *T. angustifolium* and *T. fernandesii*. *Journal of Bryology*. 17:251–262.
- Hodgetts N, Cáliz M, Englefield E, Fettes N, García Criado M, Patin L, Nieto A, Bergamini A, Bisang I, Baisheva E, et al. 2019. A miniature world in decline: European Red List of mosses, liverworts and hornworts. Brussels: International Union for the Conservation of Nature.
- Hodgetts NG, Vanderpoorten A. 2018. *Isothecium myosuroides* var. *brachythecioides* (Dixon) Braithw. reinstated as a species, *I. interludens* Stirt. *Journal of Bryology*. 40:316–323.
- Holyoak DT. 2003. A taxonomic review of some British coastal species of the *Bryum bicolor* complex, with a description of *Bryum dyffrynense* sp. nov. *Journal of Bryology*. 25:107–113.
- Holyoak DT. 2009. *Bryum apiculatum* (Bryaceae) in East Cornwall: a moss new to Europe. *Journal of Bryology*. 31:117–120.
- Holyoak DT. 2010. Notes on taxonomy of some European species of *Ephemerum* (Bryopsida: Pottiaceae). *Journal of Bryology*. 32:122–132.
- Holyoak DT, Hedenäs L. 2006. Morphological, ecological and molecular studies of the intergrading taxa *Bryum neodamense* and *B. pseudotriquetrum* (Bryopsida: Bryaceae). *Journal of Bryology*. 28:299–311.
- Holyoak DT, Köckinger H. 2010. A taxonomic revision of some European and Asian bulbiferous species of *Anomobryum* (Bryophyta: Bryaceae). *Journal of Bryology*. 32:153–169.
- Holyoak DT, Lockhart N. 2009. Australasian bryophytes introduced to South Kerry with tree ferns. *Field Bryology*. 98:3–7.
- Holyoak DT, Pedersen N. 2007. Conflicting molecular and morphological evidence of evolution within the Bryaceae (Bryopsida) and its implications for generic taxonomy. *Journal of Bryology*. 29:111–124.
- Hooper EJ, Duckett JG, Cuming AC, Kunin WE, Pressel S. 2010. *Ephemerella readeri* Müll.Hal. (*Physcomitrella readeri* (Müll.Hal.) I.G.Stone & G.A.M.Scott, Funariidae, Bryophyta): a genus and species new to Europe. *Journal of Bryology*. 32:256–264.
- Hugonnot V. 2010a. Mousses & hépatiques de Païolive (Ardèche & Gard, France). *Bulletin de la Société Botanique du Centre-Ouest, nouvelle série, numéro special*. 34:1–293.
- Hugonnot V. 2010b. Towards an improved understanding of *Riccia ciliata* Hoffm. (Marchantiopsida: Ricciaceae). *Journal of Bryology*. 32:300–303.
- Hugonnot V. 2015. *Riccia melitensis* Mass. (Marchantiophyta: Ricciaceae), an endemic species of the Maltese archipelago? *Phytotaxa*. 222:238–240.
- Hugonnot V. 2018. Bryophyte flora and vegetation of serpentine sites in the French Massif Central. *Plant Biosystems*. DOI: 10.1080/11263504.2018.1527789.
- Hugonnot V. 2019. New records for the bryophyte flora of Corsica. *Cryptogamie, Bryologie*. 40:153–158.
- Hugonnot V, Hedenäs L. 2015. *Arvernella microclada* Hugonnot & Hedenäs (Amblystegiaceae), a new minute species from France, requiring a separate genus. *Journal of Bryology*. 37:184–191.
- Hugonnot V, Porley RD, Ignatov MS. 2018. A taxonomic conundrum resolved: the transfer of *Grimmia horrida* to *Coscinodon* based on sporophyte discovery in France, with support from molecular data. *The Bryologist*. 121:520–528.
- Hugonnot V, Schill DB. 2006. *Mannia californica* (Gottsche ex Underw.) L.C.Wheeler (Aytoniaceae, Marchantiales) in Ardèche (France), new to Europe. *Cryptogamie, Bryologie*. 27:181–189.

- Hutsemékers V, Vieira CC, Ros RM, Huttunen S, Vanderpoorten A. 2012. Morphology informed by phylogeny reveals unexpected patterns of species differentiation in the aquatic moss *Rhynchostegium riparioides* s.l. *Molecular Phylogenetics and Evolution*. 62:748–755.
- Huttunen S, Bell N, Bobrova VK, Buchbender V, Buck WR, Cox CJ, Goffinet B, Hedenäs L, Ho BC, Ignatov MS, et al. 2012. Disentangling knots of rapid evolution: origin and diversification of the moss order Hypnales. *Journal of Bryology*. 34:187–211.
- Huttunen S, Ignatov MS. 2010. Evolution and taxonomy of aquatic species in the genus *Rhynchostegium* (Brachytheciaceae, Bryophyta). *Taxon*. 59:791–808.
- Huttunen S, Kuznetsova O, Li M, Wang YF, Ignatov MS. 2015. East Asian origin and Northern Hemisphere diversification in the *Brachythecium novae-angliae* complex (Bryophyta). *Taxon*. 64:911–930.
- Ignatov MS, Afonina OM, Ignatova EA. 2006. Check-list of mosses of East Europe and North Asia. *Arctoa*. 15:1–130.
- Ignatov MS, Afonina OM, Kuznetsova OI, Ignatova EA. 2012. The genus *Leptopterigynandrum* (Taxiphyllaceae, Bryophyta) in Russia. *Arctoa*. 21:207–220.
- Ignatov MS, Bersanova AS, Zharzinov ZS, Ignatova EA. 2005. *Leptodontium* (Pottiaceae, Bryophyta), a new genus for Caucasus. *Arctoa*. 14:35–38.
- Ignatov MS, Fedorova AV, Fedosov VE. 2019. On the taxonomy of Anomodontaceae and *Heterocladium* (Bryophyta). *Arctoa*. 28:75–102.
- Ignatov MS, Fedosov VE, Ignatova EA, Doroshina DY, Zolotov VI. 2010. Moss flora of Gunib area in Dagestan, the Eastern Caucasus. *Arctoa*. 19:87–96.
- Ignatov MS, Gardiner AA, Bobrova VK, Milyutina IA, Huttunen S, Troitsky AV. 2007. On the relationships of mosses of the order Hypnales, with special reference to taxa traditionally classified in Leskeaceae. In: Newton AE, Tangney R, editors. *Pleurocarpous mosses: systematics and evolution*. Boca Raton, CRC Press; p. 177–213.
- Ignatov MS, Huttunen S, Li M, Wang Y-F. 2015. How to know *Myuroclada longiramea* (Brachytheciaceae, Bryophyta). *Arctoa*. 24:497–502.
- Ignatov MS, Ignatova EA. 2004. Moss flora of the Middle European Russia. Vol. 2: Fontinalaceae–Amblystegiaceae. *Arctoa*. 11 (Supplement 2):609–960.
- Ignatov MS, Ignatova EA, Kuznetsova OI, Fedosov VE. 2019. On moss genera *Hylocomiadelpus* Ochyra & Stebel and *Rhytidadelphus* (Limpr.) Warnst. *Acta Musei Silesiae, Scientiae Naturales*. 68:123–134.
- Ignatov MS, Ivanova EI, Ignatova EA, Krivoschapkin KK. 2001. On the moss flora of Ust-Maya District (Republic Sakha/Yakutia, East Siberia). *Arctoa*. 10:165–184.
- Ignatov MS, Milyutina IA. 2007. On *Sciuro-hypnum oedipodium* and *S. curtum* (Brachytheciaceae, Bryophyta). *Arctoa*. 16:47–61.
- Ignatov MS, Milyutina IA. 2010. The genus *Brachythecium* (Brachytheciaceae, Musci) in Russia: comments on species and key for identification. *Arctoa*. 19:1–30.
- Ignatov MS, Porley RD, Ochyra R, Kuznetsova OI, Ignatova EA. 2018. *Coscinodon monchiquensis* R.D.Porley, Ochyra & Ignatova (Grimmiaceae), a new species from the Algarve, southern Portugal. *Journal of Bryology*. 40:125–136.
- Ignatov MS, Smith Merrill GL. 1995. Bryophytes of Altai Mountains. VI. The family Polytrichaceae (Musci). *Arctoa*. 5:61–97.
- Ignatova EA, Blom HH, Goryunov DV, Milyutina IA. 2010. On the genus *Schistidium* (Grimmiaceae, Musci) in Russia. *Arctoa*. 19:195–233.
- Ignatova EA, Fedorova AV, Kuznetsova OI, Ignatov MS. 2019. Taxonomy of the *Plagiothecium laetum* complex (Plagiotheciaceae, Bryophyta) in Russia. *Arctoa*. 28:28–45.
- Ignatova EA, Ignatov MS, Milyutina IA. 2010. A revision of the genus *Lindbergia* (Leskeaceae, Bryophyta) in Russia. *Arctoa*. 19:97–116.
- Ignatova EA, Kuznetsova OI, Fedosov VE, Ignatov MS. 2016. On the genus *Hedwigia* (Hedwigiaceae, Bryophyta) in Russia. *Arctoa*. 25:241–277.
- Ignatova EA, Kuznetsova OI, Ignatov MS, Köckinger H. 2012. The genera *Oxystegus* and *Pseudosymblypharis* (Pottiaceae, Bryophyta) in the Caucasus. *Arctoa*. 21:173–180.
- Ignatova EA, Kuznetsova OI, Köckinger H, Hastings R. 2008. A preliminary study of *Coscinodon* (Grimmiaceae, Musci) in Eurasia based on morphology and DNA sequence data. *Arctoa*. 17:1–18.
- Ignatova EA, Kuznetsova OI, Milyutina IA, Fedosov VE, Ignatov MS. 2017. The genus *Fabronia* (Fabroniaceae, Bryophyta) in Russia. *Arctoa*. 26:11–34.
- Ignatova EA, Tubanova DY, Tumurova OD, Goryunov DV, Kuznetsova OI. 2015. When the plant size matters: a new semi-cryptic species of *Dicranum* from Russia. *Arctoa*. 24:471–488.
- Inoue H. 1969. Contributions to the knowledge of Plagiochilaceae of southern Asia. XII. Typification of *Plagiochila javanica* (Swartz) Dum. *Journal of the Hattori Botanical Laboratory*. 32:50–56.
- Inoue Y, Tsubota H. 2014. On the systematic position of the genus *Timmiella* (Dicranidae, Bryopsida) and its allied genera, with the description of a new family Timmiellaceae. *Phytotaxa*. 181:151–162.
- Inoue Y, Tsubota H. 2016. Systematics of the family Pottiaceae (Bryophyta) with special reference to the familial and sub-familial circumscriptions. *Hikobia*. 17:117–129.
- Ivanova EI, Bell NE, Kuznetsova OI. 2015. The genus *Polytrichum* Sect. *Aporotheca* (Polytrichaceae) in Russia. *Arctoa*. 24:67–78.
- Ivanova EI, Bell NE, Kuznetsova OI, Lee H, Park M, Ignatov M. 2014. The genus *Polytrichastrum* (Polytrichaceae) in Russia. *Arctoa*. 23:164–184.
- Iwatsuki Z. 1959. Bryological miscellanies, IX. A note on Japanese *Orthotrichum*. *Journal of the Hattori Botanical Laboratory*. 21:240.
- Jiménez JA. 2006. Taxonomic revision of the genus *Didymodon* Hedw. (Pottiaceae, Bryophyta) in Europe, North Africa and Southeast and Central Asia. *Journal of the Hattori Botanical Laboratory*. 100:211–292.
- Jovet-Ast S. 1986. Les *Riccia* de la région Méditerranéenne. *Cryptogamie, Bryologie Lichénologie*. 7:287–431.
- Jovet-Ast S. 2000. Documents pour la connaissance des *Riccia* Australiens (Hépatiques, Marchantiales) – nouvelles récoltes. *Taxons nouveaux. Commentaires morphologiques & écologiques. Cryptogamie, Bryologie*. 21:289–343.
- Juslén A. 2006. Revision of Asian *Herbertus* (Herbertaceae, Marchantiophyta). *Annales Botanici Fennici*. 43:409–436.
- Kanda H. 1976 (1977). A revision of the family Amblystegiaceae of Japan II. *Journal of Science of the Hiroshima University, Series B, Division 2, Botany*. 16:47–119.
- Kiebacher T, Lüth M. 2017. *Orthotrichum dentatum* T.Kiebacher, Lüth sp. nov. (Orthotrichaceae). *Journal of Bryology*. 39:46–56.
- Kindberg NC. 1893. Notes on Canadian bryology. *Ottawa Naturalist*. 7:17–23.

- Kindberg NC. 1897. European and N. American Bryineae (mosses). Linköping: Linköpings Litografiska Aktiebolag.
- Köckinger, H. 2016. Rediscovery and redescription of the enigmatic *Radula visianica* (Porellales, Marchantiophyta). *Herzogia*. 29:625–634.
- Köckinger H. 2017. Die Horn- und Lebermoose Österreichs (Anthocerotophyta und Marchantiophyta). *Catalogus Florae Austriae*. II(2):1–382.
- Köckinger H, Bednarek-Ochyra H, Ochyra R. 2007. *Bucklandiella nivalis* (Grimmiaceae), a new moss from the Alps of Austria. *The Bryologist*. 110:92–99.
- Köckinger H, Hedenäs L. 2017. A farewell to *Tortella bambergeri* (Pottiaceae) as understood over the last decades. *Journal of Bryology*. 39:213–225.
- Köckinger H, Holyoak DT, Suanjak M. 2013. *Bryum austriacum* Köckinger, Holyoak & Suanjak, a new bulbiferous species from the Alps (Bryaceae, Bryopsida). *Journal of Bryology*. 35:57–61.
- Köckinger H, Kučera J. 2007. *Barbula amplexifolia* (Mitt.) A.Jaeger in Europe. *Journal of Bryology*. 29:33–40.
- Köckinger H, Kučera J. 2011. *Hymenostylium xerophilum* sp. nov., and *H. gracillimum*, comb. nov., two neglected European mosses and their molecular affinities. *Journal of Bryology*. 33:195–209.
- Köckinger H, Kučera J. 2016. *Brachythecium funkii* Schimp. and *B. japygum* (Glow.) Köckinger & Jan Kučera comb. nov., two alpine species hitherto included in *B. cirrosum* (Schwägr.) Schimp. *Journal of Bryology*. 38:267–285.
- Köckinger H, Kučera J, Hofmann H, Müller N, Amann G. 2012. *Barbula consanguinea* discovered in Switzerland and Austria, with a revision of former European records of *B. indica*. *Herzogia*. 25:61–70.
- Köckinger H, Lüth M, Werner O, Ros RM. 2018. *Tortella mediterranea* (Pottiaceae), a new species from southern Europe, its molecular affinities, and taxonomic notes on *T. nitida*. *The Bryologist*. 121:560–570.
- Köckinger H, Suanjak M, Schriebl A, Schröck C. 2008. Die Moose Kärntens. Klagenfurt: Natur Kärnten, Sonderreihe des Naturwissenschaftlichen Vereins für Kärnten.
- Köckinger H, van Melick H. 2007. *Didymodon maschalogenus* (Pottiaceae), a novelty in the European moss flora, reported from the Austrian Alps and southern Norway. *Lindbergia*. 32:62–68.
- Köckinger H, Werner O, Ros RM. 2010. A new taxonomic approach to the genus *Oxystegus* (Pottiaceae, Bryophyta) in Europe based on molecular data. *Nova Hedwigia*, Beiheft. 138:31–49.
- Konstantinova NA. 2000a. [Distribution patterns of the north holarctic hepatics]. *Arctoa*. 9:29–94.
- Konstantinova NA. 2000b. Redkie pechenochniki (Hepaticae) Murmanskogo oblasti i podhody k ih ohrane [Rare liverworts (Hepaticae) of Murmansk province and some approaches to their protection]. *Botanicheskii zhurnal*. Moscow, Leningrad. 85(10):122–135. Russian.
- Konstantinova NA, Akatova TV, Savchenko AN. 2009. Hepatics of Caucasian State Nature Reserve (Western Caucasus, Russia). *Arctoa* 18:121–134.
- Konstantinova NA, Bakalin VA, Andreeva EN, Bezgodov AG, Borovichev EA, Dulin MA, Mamontov YS. 2009. Checklist of liverworts (Marchantiophyta) of Russia. *Arctoa*. 18:1–64.
- Konstantinova NA, Doroshina GYa. 2011. New liverwort records from Republic of Kalmykia. 1. *Arctoa*. 20:256.
- Konstantinova NA, Lavrinenko OV. 2002. [On the flora of hepaticae of the Nenets autonomous area (Archangelsk region, north-eastern European Russia)]. *Botanicheskii zhurnal*. Moscow, Leningrad 87(9):43–50. Russian.
- Konstantinova NA, Mamontov YS, Vilnet AA. 2020. *Frullania stylifera* (R.M.Schust.) R.M.Schust. (Hepaticae), new to Eurasia. *Journal of Bryology*. 42: <https://doi.org/10.1080/03736687.2020.1722511>
- Konstantinova NA, Potemkin AD. 1994. Studies on *Scapania sphaerifera* (Hepaticae). *Annales Botanici Fennici*. 31:121–126.
- Konstantinova NA, Potemkin AD. 1996. Liverworts of the Russian arctic: an annotated check-list and bibliography. *Arctoa*. 6:125–150.
- Konstantinova NA, Potemkin AD, Schljakov RN. 1992. Checklist of the Hepaticae and Anthocerotae of the former USSR. *Arctoa*. 1:87–127.
- Konstantinova NA, Savchenko AN. 2013. Russian Caucasus distribution of fourteen hepatics rare and threatened in Europe. *Polish Botanical Journal*. 58:573–584.
- Konstantinova NA, Savchenko AN. 2018. Contribution to the bryophyte flora of Svalbard: hepatics of Barents Island. *Folia Cryptogamica Estonica* 55:105–115.
- Konstantinova NA, Vilnet AA. 2011. *Jubula hutchinsiae* subsp. *caucasica* subsp. nov. (Jubulaceae, Marchantiophyta) – a new taxon from the western Caucasus. *Arctoa*. 20:227–238.
- Konstantinova NA, Vilnet AA. 2016. A new species of the genus *Jungermannia* (Jungermanniales, Marchantiophyta) from the Caucasus with notes on taxa delimitation and taxonomy of *Jungermannia* s. str. *Phytotaxa*. 255:227–239.
- Koponen T. 2014. Bryophyte flora of Hunan Province, China. 18. Mniaceae subfam. Mnioideae (Musci). *Acta Bryolichenologica Asiatica*. 5:39–72.
- Koponen T. 2017. Lähdelehväsammal (*Rhizomnium dentatum*), jolla oli hampainen lehtilaita, ja sen mahdollinen jälkeläinen, *R. punctatum* var. *hermanperssonii*, var. nov. *Bryobrotherella*. 20:62–66.
- Koponen T, Ignatova EA. 2018. Mniaceae. In: Ignatov MS, editor. Moss Flora of Russia. Vol. 4. Bartramiales-Aulacomniales. Moscow: KMK Scientific Press Ltd. (Arctoa. 27, supplement 1); p. 439–517.
- Koponen T, Isoviita P. 2010. *Philonotis capillaris* Lindb. and *P. arnellii* Husn.; one moss, two names. *Cryptogamie, Bryologie*. 31:75–94.
- Koponen T, Sun Y. 2017. Preliminary study on phylogenetic position and delimitation of the ciliate arthroodontous genera of the moss family Mniaceae. *Journal of Bryology*. 39:23–38.
- Košnar J, Kolář F. 2009. A taxonomic study of selected European taxa of the *Tortula muralis* (Pottiaceae, Musci) complex: variation in morphology and ploidy level. *Preslia*. 81:399–421.
- Kramer W. 1980. *Tortula* Hedw. sect. Rurales De Not. (Pottiaceae, Musci) in der östlichen Holarktis. *Bryophytorum Bibliotheca*. 21:1–165.
- Kučera J, Blockeel TL, Erzberger P, Papp B, Soldán Z, Vellak K, Werner O, Ros RM. 2018. The *Didymodon tophaceus* complex (Pottiaceae, Bryophyta) revisited: new data support the subspecific rank of currently recognized species. *Cryptogamie, Bryologie*. 39:241–257.
- Kučera J, Košnar J, Werner O. 2013. Partial generic revision of *Barbula* (Musci: Pottiaceae): re-establishment of *Hydrogonium* and *Streblotrichum*, and the new genus *Gymnobarbula*. *Taxon*. 62:21–39.
- Kučera J, Kuznetsova OI, Manukjanová A, Ignatov MS. 2019. A phylogenetic revision of the genus *Hypnum*. Towards completion. *Taxon*. 68:1–33.
- Kučera J, Váňa J, Hradílek Z. 2012. Bryophyte flora of the Czech Republic: updated checklist and Red List and a brief analysis. *Preslia*. 84:813–850.
- Kürschner H, Erdağ A. 2005. Bryophytes of Turkey: an annotated reference list of the species with synonyms

- from the recent literature and an annotated list of Turkish bryological literature. *Turkish Journal of Botany*. 29:95–154.
- Kyrkjõeide MO, Hassel, K, Stenøien HK, Prestø T, Boström E, Shaw AJ, Flatberg KI. 2015. The dark morph of *Sphagnum fuscum* (Schimp.) H.Klinggr. in Europe is conspecific with the North American *S. beothuk*. *Journal of Bryology*. 37:251–266.
- Laine J, Flatberg KI, Harju P, Timonen T, Minkkinen K, Laine A, Tuittila ES, Vasander H. 2018. *Sphagnum* mosses – the stars of European mires. Helsinki: University of Helsinki.
- Lamarck, JB. 1815. Flore Française, ou, descriptions succinctes de toutes les plantes qui croissent naturellement en France. Paris: Desray.
- Lang AS, Bocksberger G, Stech M. 2015. Phylogeny and species delimitations in European *Dicranum* (Dicranaceae, Bryophyta) inferred from nuclear and plastid DNA. *Molecular Phylogenetics and Evolution*. 92:217–225.
- Lara F, Garilleti R. 2014. *Orthotrichum* Hedw. In: Guerra J, Cano MJ, Brugués M, editors. Flora Briofítica Ibérica, Vol V. Murcia: Universidad de Murcia, Sociedad Española de Briología; p. 50–135.
- Lara F, Garilleti R, Draper I, Mazimpaka V. 2018. *Lewinskya lamyana* sp. nov. (Orthotrichaceae, Bryopsida), a distinct moss from an exceptional habitat in the southern Iberian Peninsula. *Cryptogamie, Bryologie*. 39:259–270.
- Lara F, Garilleti R, Goffinet B, Draper I, Medina R, Vigalondo B, Mazimpaka V. 2016. *Lewinskya*, a new genus to accommodate the phaneroporous and monoicous taxa of *Orthotrichum* (Bryophyta, Orthotrichaceae). *Cryptogamie, Bryologie*. 37:361–382.
- Lara F, Mazimpaka V, Estébanez B, Garilleti R. 2009. *Orthotrichum consobrinum* Cardot in western Europe and south-western Asia. *Journal of Bryology*. 31:80–85.
- Larraín J, Quandt D, Stech M, Muñoz J. 2013. Lumping or splitting? The case of *Racomitrium* (Bryophytina: Grimmiaceae). *Taxon*. 62:1117–1132.
- Ligrone R, Duckett JG. 2005. *Cyathodium* Kunze (Cyathodiaceae, Marchantiophyta), genere e famiglia nuovi per l'Europa, nella Valle delle Ferriere (Amalfi, Salerno). *Informatore Botanico Italiano*. 37:84–85.
- Limpricht KG. 1895. 1. Die Laubmoose Deutschlands, Oesterreichs und der Schweiz. II. Abteilung: Bryinae. In: Rabenhorst L. Kryptogamen-Flora von Deutschland, Oesterreich und der Schweiz. 4. Leipzig: Eduard Kummer.
- Limpricht G. 1897. Drei neuen Laubmoose. Jahresbericht der Schlesischen Gesellschaft für Vaterländische Cultur. 74 (2b):1–5.
- Lindberg SO. 1877. Monographia *Metzgeriae*. Acta Societatis pro Fauna & Flora Fennica. 1:1–49.
- Loeske L. 1929. Die Laubmoos Europas II Funariaceae. Repertorium Speciarum Novarum Regni Vegetabilis, Sonderbeihft B. 3(2):1–71.
- Long DG. 2006. Revision of the genus *Asterella* P. Beauv. in Eurasia. *Bryophytorum Bibliotheca*. 63:1–299.
- Long DG, Paton JA, Squirrell J, Woodhead M, Hollingsworth PM. 2006. Morphological, ecological and genetic evidence for distinguishing *Anastrophyllum joergensenii* Schiffn. and *A. alpinum* Steph. (Jungermanniopsida: Lophoziaaceae). *Journal of Bryology*. 28:108–117.
- Lönnell N, Hassel K. 2018. New combinations within the genus *Sphagnum* (Sphagnaceae, Bryophyta). *Phytotaxa*. 369:57–58.
- Maier E. 2002. *Grimmia dissimulata* E.Maier sp. nova, and the taxonomic position of *Grimmia trichophylla* var. *meridionalis* Müll.Hal. (Musci, Grimmiaceae). *Candollea*. 56:281–300.
- Maier E. 2010. The genus *Grimmia* Hedw. (Grimmiaceae, Bryophyta) — a morphological anatomical study. *Boissiera*. 63:5–377.
- Majestyk P. 2011. A taxonomic treatment of *Daltonia* (Musci: Daltoniaceae) in the Americas. *Journal of the Botanical Research Institute of Texas*. 5:553–575.
- Mamontov YS, Konstantinova NA, Vilnet AA. 2018. One more species in the genus *Jungermannia* (Marchantiophyta: Jungermanniaceae). *Bryophyte Diversity and Evolution*. 40:79–90.
- Mamontov YS, Potemkin AD, Tubanova DY, Sofronova EV. 2018. [Liverworts of the Dzherginsky Reserve (Republic of Buryatia)]. *Novosti sistematiki nizshikh rastenii*. 52:483–504.
- Manukjanová A, Kouteckí P, Štechová T, Kučera J. 2019. Insights into the distribution patterns, habitat and morphologic differentiation of cryptic species of the moss *Hamatocaulis vernicosus* in the Czech Republic. *Herzogia*. 32:183–199.
- Marka J, Blockeel TL, Long DG, Papp B. 2018. Bryophytes new to Albania from the British Bryological Society field meeting in 2014. *Journal of Bryology*. 40:163–172.
- Mastracci M. 2003. *Thamnobryum neckerooides* (Bryopsida: Neckeraceae): lectotypification, synonymies, diagnostic characters, habitat and distribution. *Journal of Bryology*. 25:115–120.
- Masuzaki H, Shimamura M, Furuki T, Tsubota H, Yamaguchi T, Majid HMA, Deguchi H. 2010. Systematic position of the enigmatic liverwort *Mizutania* (Mizutaniaceae, Marchantiophyta) inferred from molecular phylogenetic analyses. *Taxon*. 59:448–458.
- Mazimpaka V, Lara F. 2014. *Codonoblepharon* Schwägr. In: Guerra J, Cano MJ, Brugués M, editors. Flora Briofítica Ibérica, Vol. V. Murcia: Universidad de Murcia, Sociedad Española de Briología; p. 27–30.
- Medina R, Johnson MG, Liu Y, Wickett NJ, Shaw AJ, Goffinet B. 2019. Phylogenetic delineation of *Physcomitrium* (Bryophyta: Funariaceae) based on targeted sequencing of nuclear exons and their flanking regions rejects the retention of *Physcomitrella*, *Physcomitridium* and *Aphanorrhagma*. *Journal of Systematics and Evolution*. 57:404–417.
- Medina R, Lara F, Goffinet B, Garilleti R, Mazimpaka V. 2012. Integrative taxonomy successfully resolves the pseudo-cryptic complex of the disjunct epiphytic moss *Orthotrichum consimile* s.l. (Orthotrichaceae). *Taxon*. 61:1180–1198.
- Medina R, Lara F, Goffinet B, Garilleti R, Mazimpaka V. 2013. Unnoticed diversity within the disjunct moss *Orthotrichum tenellum* s.l. validated by morphological and molecular approaches. *Taxon*. 62:1133–1152.
- Meinunger L, Schröder W. 2007. Verbreitungsatlas der Moose Deutschlands. Regensburg: O. Dürhammer für die Regensburgische Botanische Gesellschaft.
- Mitten W. 1859. Musci Indiae Orientalis, an enumeration of the mosses of the East Indies. *Journal of the Proceedings of the Linnean Society, Botany*. 1:1–96.
- Mitten W. 1864. Contributions to the Cryptogamic Flora of the Atlantic Islands. *Journal of the Linnean Society, Botany*. 8:1–10.
- Müller C. 1849. Synopsis Muscorum Frondosorum omnium hucusque Cognitorum. Vol. 1. Berolini [Berlin]: A. Foerstner.
- Müller C. 1858. Synopsis Muscorum Frondosorum omnium hucusque Cognitorum. Berlin: A. Foerstner.
- Müller F. 2007. Zum Vorkommen von *Clasmatodon parvulus* (Fabroniaceae, Bryopsida) in Deutschland. *Herzogia*. 20:293–298.

- Müller F. 2014. *Bartramia aprica* - the correct name for the Mediterranean and western North American species historically recognized as "*Bartramia stricta*". *Herzogia*. 27:211–214.
- Müller K. 1953. Hepatikologische Notizen. *Revue Bryologique & Lichénologique*. 22:131–140.
- Müller K. 1954. Die Lebermoose Europas (Dr. L. Rabenhorst's Kryptogamen-Flora von Deutschland, 3. Aufl.) 6. Band, 4. Lieferung. Leipzig: Akademische Verlagsgesellschaft; p. 481–640.
- Muñoz J, Cezón K, Hespanhol H, Quandt D. 2015. *Grimmia*. Flora Briofítica Ibérica. Vol. II. Murcia: Universidad de Murcia, Sociedad Española de Briología.
- Muñoz J, Hespanhol H, Cezón K, Séneca A. 2009. *Grimmia horrida* (Grimmiaceae), a new species from the Iberian Peninsula. *The Bryologist*. 112:325–328.
- Muñoz J, Pando F. 2000. A world synopsis of the genus *Grimmia*. *Monographs in Systematic Botany from the Missouri Botanical Garden*. 83:1–333.
- Nieto-Lugilde M, Werner O, McDaniel SF, Ros RM. 2018. Environmental variation obscures species diversity in southern European populations of the moss genus *Ceratodon*. *Taxon*. 67:673–692.
- Nieuwkoop J, Arts T. 1995. Additions to the bryophyte flora of Madeira. *Lindbergia*. 20:35–39.
- O'Shea BJ. 2006. Checklist of the mosses of sub-Saharan Africa (version 5, 12/06). *Tropical Bryology Research Reports*. 6:1–253. Available at http://www.nhm.ac.uk/hosted_sites/bbstbg/resources_lit_africa.html
- Ochi H. 1959. A revision of the Bryaceae in Japan and adjacent regions. Tottori: The Biological Institute, Faculty of Liberal Arts, Tottori University.
- Ochi H. 1970. A revision of the subfamily Bryoideae in Australia, Tasmania, New Zealand and the adjacent islands. *Journal of the Faculty of Education Tottori University, Natural Science*. 21:7–67.
- Ochi H. 1972. A revision of African Bryoideae, Musci (First Part). *Journal of the Faculty of Education Tottori University, Natural Science*. 23:1–126.
- Ochi H. 1980. A revision of the Neotropical Bryoideae, Musci (First part). *Journal of the Faculty of Education Tottori University, Natural Science*. 29:49–154.
- Ochi H. 1981. A revision of the Neotropical Bryoideae, Musci (Second part). *Journal of the Faculty of Education Tottori University, Natural Science*. 30:21–55.
- Ochyra R. 2013. The new generic name *Torrentaria*, a nomenclatural consequence of the legitimacy of *Platyhypnum* and the illegitimacy of *Platyhypnidium* (Bryophyta). *Nova Hedwigia*. 96:205–212.
- Ochyra R, Bednarek-Ochyra H. 2011. Lectotypification of *Hypnum boreale* (Bryopsida), an early name for a bryalean species and its taxonomic status. *Nova Hedwigia*. 93:525–536.
- Ochyra R, Bednarek-Ochyra H. 2015. (2384) Proposal to conserve the name *Bryum pallescens* against *Hypnum boreale* (*Bryum boreale*) (Bryophyta: Bryophyta). *Taxon*. 64:1053–1054.
- Ochyra R, Bednarek-Ochyra H, Plášek V. 2017. Proposal to conserve the name *Plagiomnium* against *Orthomnion* and *Orthomniopsis* (Bryophyta: Mniaceae). *Taxon*. 66:1218–1220.
- Ochyra R, Gos L. 1992. A new species of *Seligeria* (Musci, Seligeriaceae) from Central Europe. *Fragmenta Floristica & Geobotanica*. 37:371–378.
- Ochyra R, Lewis Smith RI, Bednarek-Ochyra H. 2008. The illustrated moss flora of Antarctica. Cambridge: Cambridge University Press.
- Ochyra R, Plášek V, Bednarek-Ochyra H. 2018. *Osculatia* – an earlier name for a segregate of *Bryum* (Bryaceae, Bryophyta). *Acta Musei Silesiae Scientiae Naturales*. 67:71–79.
- Ochyra R, Stebel A. 2008. *Hylocomiadelphus*, a new genus of pleurocarpous mosses. In: Stebel A, Ochyra R, editors. *Bryophytes of the Polish Carpathians*. Poznań, Sorus. p. 303–308.
- Ochyra R, Żarnowiec J, Bednarek-Ochyra H. 2003. Census catalogue of Polish mosses. *Biodiversity of Poland*. 3:1–372.
- Olsson S, Buchbender V, Enroth J, Huttunen S, Hedenäs L, Quandt D. 2009. Evolution of the Neckeraceae (Bryophyta): resolving the backbone phylogeny. *Systematics and Biodiversity*. 7:419–432.
- Olsson S, Enroth J, Buchbender V, Hedenäs L, Huttunen S, Quandt D. 2011. *Neckera* and *Thamnobryum* (Neckeraceae, Bryopsida) – paraphyletic assemblages. *Taxon*. 60:36–50.
- Orgaz JD, Cano MJ, Guerra J. 2013. Taxonomic revision of *Brachytheciastrum* (Brachytheciaceae, Bryophyta) from the Mediterranean Region. *Systematic Botany*. 38:283–294.
- Patiño J, Hedenäs L, Dirkse GM, Ignatov MS, Papp B, Müller F, González-Mancebo JM, Vanderpoorten A. 2017. Species delimitation in the recalcitrant moss genus *Rhynchostegiella* (Brachytheciaceae). *Taxon*. 66:293–308.
- Paton JA, Sheahan MC. 2006. *Lophocolea brookwoodiana* (Jungermanniales: Geocalyceae), a new species in Britain. *Journal of Bryology*. 28:163–166.
- Pedersen N, Cox CJ, Hedenäs L. 2003. Phylogeny of the moss family Bryaceae inferred from chloroplast DNA sequences and morphology. *Systematic Botany*. 28:471–482.
- Pedersen N, Hedenäs L. 2002. Phylogenetic relationships between *Bryum* and supposedly closely related genera. *Journal of Bryology*. 24:277–289.
- Pedersen N, Hedenäs L. 2003. Phylogenetic investigations of a well supported clade within the acrocarpous moss family Bryaceae: evidence from seven chloroplast DNA sequences and morphology. *Plant Systematics and Evolution*. 240:115–132.
- Pedersen N, Hedenäs L. 2005. Taxonomic and nomenclatural implications of phylogenetic studies of the Bryaceae based on molecular data and morphology. *The Bryologist*. 108:123–128.
- Pedersen N, Holyoak DT, Newton AE. 2007. Systematics and morphological evolution within the moss family Bryaceae: a comparison between parsimony and Bayesian methods for reconstruction of ancestral character states. *Molecular Phylogenetics and Evolution*. 43:891–907.
- Perold SM. 1989a. Studies in the genus *Riccia* (Marchantiales) from South Africa. 14. *R. concava*, section *Pilifer*. *Bothalia*. 19:161–165.
- Perold SM. 1989b. Studies in the genus *Riccia* (Marchantiales) from southern Africa. 11. *R. montana* and *R. alborosa*, a further two new white-scaled species of the group '*Squamatae*'. *Bothalia*. 19:9–16.
- Perold SM. 1997. Studies in the liverwort genus *Fossombronina* (Metzgeriales) from southern Africa. 4. A re-examination of *F. crispa*, *F. leucoantha* and *F. tumida*. *Bothalia*. 27:105–115.
- Plášek V, Sawicki J, Ochyra R, Szczecińska M, Kulik T. 2015. New taxonomical arrangement of the traditionally conceived genera *Orthotrichum* and *Ulota*. *Acta Musei Silesiae Scientiae Naturales*. 64:169–174.
- Plášek V, Sawicki J, Trávníčková V, Pasečná M. 2009. *Orthotrichum moravicum* (Orthotrichaceae), a new moss species from the Czech Republic. *The Bryologist*. 112:329–336.
- Podpěra J. 1954. *Conspectus Muscorum Europaeorum*. Prague: Československé Akademie Ved.

- Podpěra J. 1973. *Bryum* generis monographiae prodromus I. 1. Species Eurasiae septentrionalis. Pars 16. Prague: Academia, Publishing House of the Czechoslovak Academy of Sciences.
- Porley RD, Edwards S. 2010. *Leptodontium proliferum* Herzog (Bryopsida: Pottiaceae), new to Europe. *Journal of Bryology*. 32:46–50.
- Potemkin AD. 1998. On the origin, evolution and classification of the genus *Scapania* (Dum.) Dum. (Hepaticae). *Journal of the Hattori Botanical Laboratory*. 85:33–61.
- Potemkin AD. 1999. An analysis of the practical taxonomy of some critical northern species of *Scapania* (Scapaniaceae, Hepaticae). *The Bryologist*. 102:32–38.
- Potemkin AD. 2008. [New liverwort records from Republic of Komi. 1]. *Arctoa* 17:198.
- Potemkin AD. 2019. The poorly known species *Scapania schljakovii* Potemkin (Scapaniaceae, Marchantiophyta), its gemmae and first record for Bhutan. *Journal of Bryology*. 41:174–176.
- Potemkin AD, Rozantseva EI. 2015. New and little-known for the Leningrad Region liverworts (Marchantiophyta) from Hogland Island (Gulf of Finland, Baltic Sea). *Novosti sistematiki nizshikh rastenii*. 49:369–381.
- Price MJ, Ellis LE. 2018. The disentanglement of *Andreaea alpina* Hedw. (Andreaeaceae): typifications and nomenclatural changes. *Taxon*. 67:989–995.
- Price MJ, Maier E, Ellis LT. 2019. *Dicranum scottianum* Turner (Dicranaceae): synonyms, a type and the correct author. *Journal of Bryology*. 41:149–156.
- Puche F, Casas C, Brugués M. 2006. *Didymodon eckeliae* (Pottiaceae), new to Europe. *The Bryologist*. 109:239–241.
- Puche F, Segarra-Moragues JG. 2013. *Riella bialata* Trab. (Riellaceae, Marchantiophyta): a new addition to the European liverwort flora. *Cryptogamie, Bryologie* 34:341–352.
- Richards PW. 1936. A collection of bryophytes from the Azores. *Annales Bryologici*. 9:131–138.
- Rodegher E. 1896. Elenco delle Epatiche della provincia di Bergamo. *Nuovo Giornale Botanico Italiano* (n.ser.). 3:423–436.
- Rodriguez-Romero A, Hernandez-Ferrer M, Patiño J, Werner W, González-Mancebo, JM. 2016. Characterization of ten polymorphic microsatellite loci for the threatened species *Grimmia curviseta* Bouman (Grimmiaceae, Musci). *Journal of Bryology*. 39:16–32.
- Ros RM, Guerra J, Carrion JS, Cano MJ. 1996. A new point of view on the taxonomy of *Pottia starckeana* agg. (Musci, Pottiaceae). *Plant Systematics and Evolution*. 199:153–165.
- Ros RM, Mazimpaka V, Abou-Salama U, Aleffi M, Blockeel TL, Brugués M, Cano MJ, Cros RM, Dia MG, Dirkse GM, et al. 2007. Hepatics and Anthocerotae of the Mediterranean, an annotated checklist. *Cryptogamie, Bryologie*. 28:351–437.
- Ros RM, Mazimpaka V, Abou-Salama U, Aleffi M, Blockeel TL, Brugués M, Cros RM, Dia MG, Dirkse GM, Draper I, et al. 2013. Mosses of the Mediterranean, an annotated checklist. *Cryptogamie, Bryologie*. 34:99–283.
- Ros RM, Muñoz J, Werner O, Rams S. 2008. New typifications and synonyms in *Tortula* sect. *Pottia* (Ehrh. ex Rchb.) Kindb. (Pottiaceae, Musci). *Taxon*. 57:279–288.
- Ros RM, Rams S, Holyoak DT, Brugués M, Cros RM. 2007. Revision of the records of *Brachymerium commutatum* (Bryophyta, Bryaceae) in Europe. *Nova Hedwigia*. 84:241–247.
- Ros RM, Werner O. 2007. The circumscription of the genus *Pottiopsis* (Pottiaceae, Bryophyta) based on morphology and molecular sequence data. *Nova Hedwigia, Beiheft*. 131:65–79.
- Roskov Y, Ower G, Orrell T, Nicolson D, Bailly N, Kirk PM, Bourgoin T, DeWalt RE, Decock W, Nieukerken E, et al., editors. 2019. Species 2000 and ITIS Catalogue of Life, 2019 Annual Checklist. Digital resource at www.catalogueoflife.org/annual-checklist/2019. Species 2000: Naturalis, Leiden, the Netherlands. ISSN 2405-884X.
- Rycroft DS, Groth H, Heinrichs J. 2004. Reinstatement of *Plagiochila maderensis* (Jungermanniopsida: Plagiochilaceae) based on chemical evidence and nrDNA ITS sequences. *Journal of Bryology*. 26:37–45.
- Saługa M, Ochyra R, Żarnowiec J, Ronikier M. 2018. Do Antarctic populations represent local or widespread phylogenetic and ecological lineages? Complicated fate of bipolar moss concepts with *Drepanocladus longifolius* as a case study. *Organisms Diversity, Evolution*. 18:263–278.
- Saukel J. 1985. Zum Merkmalsbestand einiger mitteleuropäischer Arten der Lebermoosgattung *Lophozia* (Dum.) Dum. (Sektion *Lophozia*). *Stapfia*. 14:149–185.
- Sawicki J, Plášek V, Ochyra R, Szczecińska M, Ślipiko M, Myszczyński K, Kulik T. 2017. Mitogenomic analyses support the recent division of the genus *Orthotrichum* (Orthotrichaceae, Bryophyta). *Scientific Reports*. 7:1–10.
- Sawicki J, Plášek V, Szczecińska M. 2010. Molecular studies resolve *Nyholmia* (Orthotrichaceae) as a separate genus. *Journal of Systematics and Evolution*. 48:183–194.
- Sawicki J, Szczecińska M, Bednarek-Ochyra H, Ochyra R. 2015. Mitochondrial phylogenomics supports splitting the traditionally conceived genus *Racomitrium* (Bryophyta: Grimmiaceae). *Nova Hedwigia*. 100:1–25.
- Schill DB, Long DG, Köckinger H. 2008. Taxonomy of *Mannia controversa* (Marchantiidae, Aytoniaceae) including a new subspecies from East Asia. *Edinburgh Journal of Botany*. 65:35–47.
- Schimper WP. 1876. *Synopsis Muscorum europaeorum*. 2nd edition. 2 vols. Stuttgartiae: E. Schweizerbart.
- Schlesak S, Hedenäs L, Nebel M, Quandt D. 2018. Cleaning a taxonomic dustbin: placing the European *Hypnum* species in a phylogenetic context! *Bryophyte Diversity and Evolution*. 40:37–54.
- Schlesak S, Hedenäs L, Nebel M, Quandt D. 2019. Addendum to *Hypnum subcomplanatum* Hedenäs, Schlesak & Quandt. *Bryophyte Diversity and Evolution*. 41:1.
- Schljakov RN. 1969. [De positione systematica *Lophosiae murmanicae* Kaal.]. *Novosti sistematiki nizshikh rastenii*. 6:241–245.
- Schljakov RN. 1970. [Sectio nova generis *Lophozia* Dum. emend. Loeske]. *Novosti sistematiki nizshikh rastenii*. 7:324–333.
- Schljakov RN. 1975. [Notulae systematicae de hepaticis s. str.]. *Novosti sistematiki nizshikh rastenii*. 12:307–317.
- Schljakov RN. 1980. *Pechenochnye Mkh Severa SSSR*. 3. Leningrad: Nauka.
- Schljakov RN. 1981. *Pechenochnye Mkh Severa SSSR*. 4. Leningrad: Nauka.
- Schljakov RN. 1998. On the *Lophozia groenlandica* (Nees) Macoun (Hepaticae). *Arctoa*. 7:191–196.
- Schriebl A. 1991. Experimentelle Studien über die Laubmoosgattung *Polytrichum*. *Carinthia II*. 181/101:461–506.
- Schuster RM. 1969. *The Hepaticae and Anthocerotae of North America*. II. New York: Columbia University Press.
- Schuster RM. 1974. *The Hepaticae and Anthocerotae of North America*. III. New York: Columbia University Press.
- Schuster RM. 1988. *The hepaticae of South Greenland*. *Beihefte zur Nova Hedwigia*. 92:1–255.

- Schuster RM. 1992a. Studies on Marchantiales. I–III. Journal of the Hattori Botanical Laboratory. 71:267–287.
- Schuster RM. 1992b. The Hepaticae and Anthocerotae of North America. VI. New York: Columbia University Press.
- Schwägrichen CF. 1816. Species Muscorum Frondosorum, Supplementum Primum. Leipzig: Barth.
- Schwägrichen CF. 1827. Species Muscorum Frondosorum, Supplementum Tertium, vol. 1, sect. 1. Leipzig: Barth.
- Segarra-Moragues JG, Puche F, Sabovljević M, Infante M, Heras P. 2014. Taxonomic revision of *Riella* subgenus *Trabutiella* (Riellaceae, Sphaerocarpaceae). Phytotaxa. 159:131–174.
- Séneca A, Söderström L. 2009. Sphagnophyta of Europe and Macaronesia: a checklist with distribution data. Journal of Bryology. 31:243–254.
- Sérgio C. 1985. Notulae Bryoflorae Lusitanicae I. 5 - Notas acerca do género *Fossombronia* Raddi em Portugal. Portugaliae Acta Biologica. Série B. Sistemática, Ecologia, Biogeografia e Paleontologia. 14:188–194.
- Sérgio C. 2003. Notulae Bryoflorae Lusitanicae IX. 1. Acerca sa presença de *Fossombronia leucoxantha* Lehm. e *F. crispa* Ness [sic!] em Portugal. Portugaliae Acta Biologica. Série B. Sistemática, Ecologia, Biogeografia e Paleontologia. 21:279–282.
- Sérgio C. 2006. A review of the *Gymnostomum calcareum* Nees & Hornsch. complex (Bryopsida: Pottiaceae) in southern Europe and the Macaronesian Islands, including *G. calcareum* var. *atlanticum* var. nov. Journal of Bryology. 28:38–45.
- Sérgio C, Garcia, C. 2009. Noteworthy range extensions of two *Aneura* (Jungermanniopsida, Metzgeriales) species new to the Iberian Peninsula: *Aneura maxima* (Schiffn.) Steph. and *A. pseudopinguis* (Herzog) Pócs. Cryptogamie, Bryologie. 30:207–215.
- Sérgio C, Sim-Sim M. 2012. *Andreaea flexuosa* R. Brown bis subsp. *luisieri* Sérgio & Sim-Sim (Andreaeaceae), a new taxon from Madeira Island. Cryptogamie, Bryologie. 33:271–277.
- Sérgio C, Sim-Sim M, Luis L. 2010. *Atrichum androgynum* (Müll. Hal.) A. Jaeger in Portugal, Azores and Madeira, new to the European bryoflora. Cryptogamie, Bryologie. 31:281–288.
- Shaw AJ. 2014. The impact of molecular data on our understanding of the bryophyte flora of North America. Flora of North America Editorial Committee, editors. Flora of North America. Vol. 28. Bryophyta, Part 2. New York: Oxford University Press; p. 3–13.
- Shaw AJ, Flatberg KI, Szövényi P, Ricca M, Johnson MG, Stenøien HK, Shaw B. 2012. Systematics of the *Sphagnum fimbriatum* complex: phylogenetic relationships, morphological variation, and allopolyploidy. Systematic Botany. 37:15–30.
- Shaw B, Crandall-Stotler B, Váňa J, Stotler RE, von Konrat M, Engel JJ, Davis CE, Long DG, Sova P, Shaw AJ. 2015. Phylogenetic relationships and morphological evolution in a major clade of leafy liverworts (phylum Marchantiophyta, order Jungermanniales): suborder Jungermanniineae. Systematic Botany. 40:27–45.
- Simmel J, Poschlod P. 2017. Beiträge zur Bestimmung und Ökologie der mitteleuropäischen *Leucobryum*-Taxa. Herzogia. 30:397–411.
- Sim-Sim M. 1999. The genus *Frullania* Raddi (Hepaticae) in Portugal and Madeira. Cryptogamie, Bryologie 20:83–144.
- Sim-Sim M, Afonina OM, Almeida T, Désamoré A, Laenen B, Garcia CA, González-Mancebo JM, Stech M. 2017. Integrative taxonomy reveals too extensive lumping and a new species in the moss genus *Amphidium* (Bryophyta). Systematics and Biodiversity. 15:451–463.
- Skrzypczak R. 2001. Découverte de *Riella parisii* Gottsche dans le Var (espèce nouvelle pour la France) et d'une deuxième station de *Riella helicophylla* (Bory & Mont.) Mont. dans l'Hérault. Bulletin de la Société Botanique du Centre-Ouest, nouvelle série. 32:277–290.
- Smith AJE. 1990. The liverworts of Britain and Ireland. Cambridge: Cambridge University Press.
- Smith AJE. 2004. The moss flora of Britain and Ireland. 2nd ed. Cambridge: Cambridge University Press.
- Söderström L, De Roo RT, Hedderson TAJ. 2010. Taxonomic novelties resulting from recent reclassification of the Lophoziaceae/Scapaniaceae clade. Phytotaxa. 3:47–53.
- Söderström L, Hagborg A, von Konrat M. 2019. ELPT: Early Land Plants Today (version Jan 2019). In: Roskov Y, Ower G, Orrell T, Nicolson D, Bailly N, Kirk PM, Bourgoin T, DeWalt RE, Decock W, Nieuwerkerken E, et al., editors. Species 2000 and ITIS Catalogue of Life, 2019 Annual Checklist. Digital resource at www.catalogueoflife.org/annual-checklist/2019. Species 2000: Naturalis, Leiden, the Netherlands. ISSN 2405-884X.
- Söderström L, Hagborg A, von Konrat M, Bartholomew-Began S, Bell D, Briscoe L, Brown E, Cargill DC, Costa DP, Crandall-Stotler BJ, et al. 2016. World checklist of hornworts and liverworts. PhytoKeys. 59:1–828.
- Söderström L, Urmi E, Váňa J. 2002. Distribution of Hepaticae and Anthocerotae in Europe and Macaronesia. Lindbergia. 27:3–47.
- Söderström L, Urmi E, Váňa J. 2007. The distribution of hepaticae and anthocerotae in Europe and Macaronesia – Update 1–427. Cryptogamie, Bryologie. 28:299–350.
- Söderström L, Váňa J, Crandall-Stotler B, Hentschel J, Hagborg A, von Konrat MJ. 2015. Notes on Early Land Plants Today. 68. Miscellaneous notes on Marchantiophyta. Phytotaxa. 202:69–72.
- Söderström L, Váňa J, Hagborg A, von Konrat M. 2013. Notes on Early Land Plants Today. 35. Notes on Lophoziaceae (Marchantiophyta). Phytotaxa. 97:27–35.
- Sofronova EV, Abakarova AS, Afonina OV, Akatova TV, Baisheva EZ, Baj X-L, Bezgodov AG, Bochkin VD, Borovichev EA, Chernadjeva IV, et al. 2013. New bryophyte records. 2 — Novye briologičeskie nahdki. 2. Arctoa 22:239–262.
- Sotiaux A, Enroth J, Olsson S, Quandt D, Vanderpoorten A. 2009. When morphology and molecules tell us different stories: a case-in-point with *Leptodon corsicus*, a new and unique endemic moss species from Corsica. Journal of Bryology. 31:186–196.
- Spence JR. 2005. New genera and combinations in Bryaceae (Bryales, Musci) for North America. Phytologia. 87:15–28.
- Spence JR. 2014. Bryaceae. In: Flora of North America Editorial Committee, editors. Flora of North America. Vol. 28. Bryophyta, Part 2. New York: Oxford University Press.
- Sprengel KPJ. 1807. Mantissa Prima Florae Halensis. C. Kumelium: Halae [Halle].
- Spruce R. 1876a. On *Anomoclada*, a new genus of Hepaticae, and on its allied genera, *Odontoschisma* and *Adelanthus* (cont.). Journal of Botany, British and Foreign. 14:193–203.
- Spruce R. 1876b. On *Anomoclada*, a new genus of hepaticae, and on its allied genera, *Odontoschisma* and *Adelanthus* (concluded). Journal of Botany, British and Foreign. 14:230–235.
- Stech M. 1999. *Dichodontium palustre* (Dicks.) Stech comb. nov., a new name for *Dicranella palustris* (Dicks.) Crundw.

- ex Warb. (Dicranaceae, Bryopsida). *Nova Hedwigia*. 69:237–240.
- Stech M, McDaniel SF, Hernández-Maqueda R, Ros RM, Werner O, Muñoz J, Quandt D. 2012. Phylogeny of haplolepidous mosses – challenges and perspectives. *Journal of Bryology*. 34:173–186.
- Stech M, Sim-Sim M, Esquivel MG, Fontinha S, Tangney R, Lobo C, Gabriel R, Quandt D. 2008. Explaining the ‘anomalous’ distribution of *Echinodium* (Bryopsida: Echinodiaceae): independent evolution in Macaronesia and Australasia. *Organisms, Diversity and Evolution*. 8:282–292.
- Ștefănuț S. 2008. The hornwort and liverwort atlas of Romania. Bucharest: Ars Docendi.
- Stephani F. 1903. Species hepaticarum 2. *Bulletin de l’Herbier Boissier (sér. 2)* 3:326–341.
- Stephani F. 1905. Species hepaticarum 3. *Bulletin de l’Herbier Boissier (sér. 2)*. 5:1129–1144.
- Stephani F. 1907. Species hepaticarum 3. *Bulletin de l’Herbier Boissier (sér. 2)*. 7:477–492.
- Stotler RE, Bray JR jr, Cargill DC, Kraysky D, Crandall-Stotler BJ. 2003. Typifications in the genus *Fossombronina* (Marchantiophyta). *The Bryologist*. 106:130–142.
- Stotler RE, Crandall-Stotler BJ. 2017. A synopsis of the liverwort flora of North America north of Mexico. *Annales of the Missouri Botanical Garden*. 102:574–709.
- Streiff A. 2006. Phylogenetic study of *Grimmia* (Grimmiaceae) based on plastid DNA sequences (*trnL-trnF* and *rps4*) and on morphological characters. *The Bryologist*. 109:224–235.
- Sun Y, He X. 2019. On the identity of *Herbertus aduncus* (Dicks.) Gray, a taxonomic update. *Acta Bryolichenologica Asiatica*. 8:145–161.
- Szweykowski J, Buczkowska K, Odrzykoski IJ. 2005. *Conocephalum salebrosum* (Marchantiopsida, Conocephalaceae) – a new holarctic liverwort species. *Plant Systematics and Evolution*. 253:133–158.
- Tassi A. 1901. Epatiche coltivate nell’Orto botanico di Siena. *Bullettino del laboratorio ed orto botanico della R. università di Siena*. 4:17–18.
- Tubanovaya DY, Goryunov DV, Ignatova EA, Ignatov MS. 2010. On the taxonomy of *Dicranum acutifolium* and *D. fuscescens* complexes (Dicranaceae, Bryophyta) in Russia. *Arctoa*. 19:151–164.
- Tubanovaya DY, Ignatova EA. 2011. A new species of *Dicranum* (Dicranaceae, Bryophyta) from Asiatic Russia. *Arctoa*. 20:183–190.
- Turland NJ, Wiersema JH, Barrie FR, Greuter W, Hawksworth DL, Herendeen PS, Knapp S, Kusber W-H, Li D-Z, Marhold K, et al. 2018. International Code of Nomenclature for algae, fungi, and plants (Shenzhen Code). *Regnum Vegetabile*. 159:1–254.
- Ulvinen T, Kypärä T. 2016. Nipponinlähdesammal (*Philonotis yezoana*) Suomen Lapissa, Bryobrotherella. 19:71–76.
- Unger FJAN, Klotschy CGT 1865. *Die Insel Cypren*. Wien: Wilhelm Braun Müller.
- Urmi E. 2017. Betreffend *Diplophyllum obtusatum* (R.M.Schust.) R.M.Schust. in der Schweiz. *Meylania*. 60:5–7.
- Váňa J. 1986. *Ochyraea tatrensis* gen. et spec. nov., a remarkable pleurocarpous moss from Czechoslovakia. *Journal of Bryology*. 14:261–267.
- Váňa J, Bechteler J, Schäfer-Verwimp A, Renner MAM, Heinrichs J. 2017. Integrative taxonomy substantiates the presence of three *Radula* species in Austria: *Radula complanata*, *R. lindenberghiana*, and *R. visianica* (Porellales, Jungermanniopsida). *Cryptogamie, Bryologie* 38:125–135.
- Váňa J, Engel JJ. 2013. The liverworts and hornworts of the Tristan da Cunha group of islands in the south Atlantic Ocean. *Memoirs of the New York Botanical Garden*. 105:1–138.
- Váňa J, Söderström L, Hagborg A, von Konrat M. 2013. Notes on Early Land Plants Today. 40. Notes on Cephaloziellaceae (Marchantiophyta). *Phytotaxa* 112:1–6.
- Vanderpoorten A. 2004. A simple taxonomic treatment for a complicated evolutionary story: the genus *Hygroamblystegium* (Hypnales, Amblystegiaceae). *Monographs in Systematic Botany from the Missouri Botanical Garden*. 98:320–327.
- Vanderpoorten A, Boles S, Shaw AJ. 2003. Patterns of molecular and morphological variation in *Leucobryum albidum*, *L. glaucum*, and *L. juniperoideum* (Bryopsida). *Systematic Botany*. 28:651–656.
- Vanderpoorten A, Hedenäs L. 2009. New combinations in the Amblystegiaceae. *Journal of Bryology*. 31:129–132.
- Vanderpoorten A, Hedenäs L, Cox CJ, Shaw AJ. 2002. Circumscription, classification, and taxonomy of Amblystegiaceae (Bryopsida) inferred from nuclear and chloroplast DNA sequence data and morphology. *Taxon*. 51:115–122.
- Vanderpoorten A, Patiño J, Dirkse G, Blockeel T, Hedenäs L. 2015. Early divergence of an Azorean endemic species in the moss genus *Rhynchostegiella* (Brachytheciaceae). *Phytotaxa*. 210:60–69.
- Vigalondo B, Garilleti R, Vanderpoorten A, Patiño J, Draper I, Calleja JA, Mazimpaka V, Lara F. 2019. Do mosses really exhibit so large distribution ranges? Insights from the integrative taxonomic study of the *Lewinskya affinis* complex (Orthotrichaceae, Bryopsida). *Molecular Phylogenetics and Evolution*. 140:106598.
- Vigalondo B, Patiño J, Draper I, Mazimpaka V, Shevock JR, Losada-Lima A, González-Mancebo JM, Garilleti R, Lara F. 2019. The long journey of *Orthotrichum shevockii* (Orthotrichaceae, Bryopsida): from California to Macaronesia. *PLoS One*. 14(2):e0211017.
- Villarreal JC, Forrest LL, Cooper ED, Long DG. 2014. Phylogenetic affinities and conservation status of *Telaranea murphyae* Paton in Britain. *Journal of Bryology*. 36:191–199.
- Vilnet AA, Borovichev EA, Bakalin VA. 2014. *Frullania subarctica* - a new species of the *Frullania tamarisci* complex (Frullaniaceae, Marchantiophyta). *Phytotaxa*. 173:61–72.
- Vilnet AA, Konstantinova NA, Troitsky AV. 2010. Molecular insight on phylogeny and systematics of the Lophoziaceae, Scapaniaceae, Gymnomitriaceae and Jungermanniaceae. *Arctoa*. 19:31–50.
- Vilnet AA, Milyutina IA, Troitsky AV, Konstantinova NA. 2005. [On phylogeny and systematics of genus *Lophozia* (Dumort.) Dumort. s.str. based on trnL-trnF sequences of chloroplast DNA and ITS 1-2 sequences of nuclear DNA]. In: Vilnet AA, Milyutina IA, Troitsky AV, Konstantinova NA, editors. *Actual problems in Bryology: proceedings of the international meeting devoted to the 90th anniversary of A. L. Abramova* (Saint Petersburg, November 22–25, 2005). St. Petersburg: Komarov Botanical Institute; p. 37–43.
- Vitt DH, Varigo C. 1997. *Orthotrichum elegans*, a taxon worthy of species rank. *Journal of the Hattori Botanical Laboratory*. 82:329–335.
- Wang CY, Zhao JC. 2009. Phylogeny of *Ptychostomum* (Bryaceae, Musci) inferred from sequences of nuclear ribosomal DNA internal transcribed spacer (ITS) and chloroplast *rps4*. *Journal of Systematics and Evolution*. 47:311–320.
- Werner O, Köckinger H, Magdy M, Ros RM. 2014. On the systematic position of *Tortella arctica* and *Trichostomum arcticum* (Bryophyta, Pottiaceae). *Nova Hedwigia*. 98:273–293.
- Werner O, Patiño J, González-Mancebo JM, Gabriel R, Ros RM. 2009. The taxonomic status and the geographical relationships of the Macaronesian endemic moss *Fissidens luisieri*

- (Fissidentaceae) based on DNA sequence data. *The Bryologist*. 112:315–324.
- Werner O, Rams S, Kučera J, Larraín J, Afonina OM, Pisa S, Ros RM. 2013. New data on the moss genus *Hymenoloma*, with special reference to *H. mulahaceni*. *Cryptogamie, Bryologie*. 34:13–30.
- Werner O, Rodriguez-Atienza A, Nieto-Lugilde M, Ros RM. 2015. New data in the systematics of the European-Asian species of *Leucodon* (Leucodontaceae, Bryophyta). *Nova Hedwigia*. 100:333–354.
- Werner O, Ros RM, Goffinet B. 2007. A reconsideration of the systematic position of *Goniomitrium* (Funariaceae) based on chloroplast sequence markers. *The Bryologist*. 110:108–114.
- Werner O, Ros RM, Grundmann M. 2005. Molecular phylogeny of Trichostomoideae (Pottiaceae, Bryophyta) based on nrITS sequence data. *Taxon*. 54:361–368.
- Withering W. 1776. A botanical arrangement of all the vegetables naturally growing in Great Britain. Vol. II. Birmingham: M. Swinney; p. 385–838.
- Wynns JT, Schröck C. 2018. Range extensions for the rare moss *Plagiothecium handelii*, and its transfer to the resurrected genus *Ortholimonium*. *Lindbergia*. 41:1–7.

Index

This index enumerates all names mentioned in the text, accepted taxa as well as synonyms. Taxa in bold are accepted taxa occurring in Europe or northern Macaronesia and taxa in square brackets are taxa only occurring outside the area.

Abietinella 59

A. abietina 59

var. *abietina* 59

var. *hystricosa* 59

Acanthocoleus 22

A. aberrans 22

[var. *aberrans*] 22

var. *laevis* 22

Acaulon 36

A. casasianum 36

A. dertosense 36

A. fontiquerianum 36

[*A. longifolium*] 69

A. mediterraneum 36

A. muticum 36

A. piligerum 36

A. triquetrum 36

Achrophyllum 55

A. dentatum 4, 55

ACROBOLBACEAE 6, 15

Acrobolboideae 15

Acrobolbus 16

[*A. anisodontus*] 66

A. azoricus 16, 66

[*A. ciliatus*] 66

[*A. laxus*] 66

A. madeirensis 16, 66

A. wilsonii 16, 66

[var. *andinus*] 16

var. *wilsonii* 16

[*Acrolejeunea emergens*] 68

ADELANTHACEAE 6, 9

Adelanthoideae 9

Adelanthus 9

A. section *Lindenbergiani* 9

A. *decipiens* = *Pseudomarsupidium decipiens* 9

Zander RH. 2013. A framework for post-phylogenetic systematics. St. Louis: Zetetic Publications. CreateSpace Independent Publishing, Amazon

Zander RH. 2019. Macroevolutionary versus molecular analysis: systematics of the *Didymodon* segregates *Aithobryum*, *Exobryum* and *Fuscobryum* (Pottiaceae). *Hattoria*. 10:1–38.

Zander RH, Hedderson TA. 2016. Re-evaluation of *Hymenostylium xanthocarpum* (Hook.) Brid., and *Ardeuma* R.H.Zander & Hedd., a new name for all other species of *Hymenostylium* (Pottiaceae, Bryophyta). *Journal of Bryology*. 38:295–301.

Zinovieva LA. 1973. K flore pechenochnich mchov poljarnogo i severnogo Urala [On the flora of hepatics of Polar and North Urals]. *Ucenyje Zapiski. Permskij Gosudarstvennyj Universitet imeni A. M. Gor'kogo*. 263:14–37.

Zolotov VI. 2007a. On systematics and distribution of some species of *Bryum* (Bryaceae, Bryophyta) in Russia. *Arctoa*. 15:155–162.

Zolotov VI 2007b. Genus 1. *Bryum* Hedw. – *Brium*, p. 1–67. <http://arctoa.ru/Flora/taxonomy-ru/Bryum1.htm>. [Accessed 2010 May 25].

A. lindenbergianus 9

Aithobryum = *Didymodon* 37

A. bistratosum = *Didymodon bistratosus* 37

Alleniella 64

A. besseri 64

A. complanata 64

Aloina 36

A. aloides 36

A. ambigua 36

A. bifrons 36

A. brevisrostris 36

A. humilis 36

A. obliquifolia 36

A. rigida 36

Alophosia 29

A. azorica 29

Amblyodon 48

A. dealbatus 48

AMBLYSTEGIACEAE 2, 8, 56, 57

Amblystegioideae 56

Amblystegium 56

A. confervoides = *Serpoleskea confervoides* 58

A. radicale = *Pseudocampyllum radicale* 58

A. serpens 56

var. *salinum* = *Amblystegium serpens* 56

A. subtile = *Pseudoamblystegium subtile* 58

A. varium = *Hygroamblystegium varium* 57

AMPHIDIACEAE 7, 33

Amphidium 33

A. curvipes 33, 68

A. lapponicum 33

A. mougeotii 33

[*A. tortuosum*] 68

A. tortuosum auct. eur. = *Amphidium curvipes* 33

Anacamptodon 56, 57

A. sauteri = *Microhypnum sauteri* 57

A. splachnoides 56

Anacolia 47

A. laevisphaera = *Bartramia laevisphaera* 47

A. menziesii 47

A. webbii 47

Anastrepta 9

A. orcadensis 9

ANASTROPHYLLACEAE 6, 9

Anastrophyllum 9, 10**A. alpinum** 9**A. assimile** 9var. *nardioides* = *Anastrophyllum assimile* 9*A. cavifolium* = *Orthocaulis cavifolius* 10**A. donnianum** 9*A. hellerianum* = *Crossocalyx hellerianus* 9**A. joergensenii** 9**A. michauxii** 9*A. minutum* = *Sphenolobus minutus* 10var. *weberi* = *Sphenolobus minutus* 10*A. saxicola* = *Sphenolobus saxicola* 10*A. sphenoloboides* = *Schizophyllopsis sphenoloboides* 10**Andoa** 65**A. berthelotiana** 65**Andreaea** 29A. section *Andreaea* 29A. section *Chasmocalyx* 29A. section *Nerviae* 29**A. alpestris** 29**A. alpina** 29*A. alpina* auct. mult. = *Andreaea hookeri* 29**A. blyttii** 29**A. crassinervia** 29**A. flexuosa** 29[subsp. *flexuosa*] 29subsp. *luisieri* 29**A. frigida** 29*A. hartmanii* = *Andreaea alpina* 29**A. heinemannii** 29subsp. *crassifolia* 29subsp. *heinemannii* 29**A. hookeri** 29**A. megistospora** 29**A. mutabilis** 29**A. nivalis** 29*A. obovata* = *Andreaea alpina* 29**A. rothii** 29subsp. *falcata* 29subsp. *rothii* 29**A. rupestris** 29var. *alpestris* = *Andreaea alpestris* 29var. *papillosa* 29var. *rupestris* 29**A. sinuosa** 29

ANDREAEEACEAE 7, 29

ANDREAEEALES 7, 29

Andreaeidae 7

Andreaeopsida 7, 29

Aneura 23**A. latissima** 23**A. maxima** 23**A. mirabilis** 23**A. pinguis** 23var. *angustior* = *Aneura pinguis* 23var. *denticulata* = *Aneura pinguis* 23var. *fuscovirens* = *Aneura pinguis* 23*A. pseudopinguis* = *A. latissima* 23

ANEURACEAE 6, 23

Anisothecium palustre = *Diobelonella palustris* 33**Anoetangium** 40**A. aestivum** 40**A. angustifolium** 40**A. handelii** 40*A. hornsuschianum* = *Molendoa hornsuschiana* 42*A. sendtnerianum* = *Molendoa hornsuschiana* 42*A. schliephackii* = *Molendoa schliephackei* 42*A. taeniatifolium* = *Molendoa taeniatifolia* 42*A. warburgii* = *Molendoa warburgii* 42**Anomobryum** 3, 48**A. apiculatum** 4, 5, 48*A. bavaricum* = *Bryum bavaricum* 48**A. concinatum** 48**A. julaceum** 48**A. lusitanicum** 48**A. notarisii** 5, 48**Anomodon** 66*A. attenuatus* = *Pseudanomodon attenuatus* 64, 66**A. longifolius** 66*A. rostratus* = *Claopodium rostratum* 58**A. rugelii** 66**A. tristis** 66**A. viticulosus** 66

ANOMODONTACEAE 8, 66

Anomodontella = *Anomodon* 66*A. longifolia* = *Anomodon longifolius* 66*Anomodontopsis* = *Anomodon* 66*A. rugelii* = *Anomodon rugelii* 66**Anthelia** 16**A. julacea** 16**A. juratzkana** 16

ANTHELIACEAE 6, 16

Anthoceros 8**A. agrestis** 8**A. caucasicus** 8**A. neesii** 8**A. punctatus** 8subsp. *agrestis* = *Anthoceros agrestis* 8var. *douinii* = *Anthoceros agrestis* 8

ANTHOCEROTACEAE 6, 8

ANTHOCEROTALES 6, 8

Anthocerotidae 6

ANTHOCEROTOPHYTA 2, 6, 8

ANTHOCEROTOPSIDA 6, 8

Antitrichia 64**A. californica** 64**A. curtispindula** 64

ANTITRICHACEAE 8, 64

Aongstroemia 33**A. longipes** 33

AONGSTROEMIIACEAE 7, 33

Aphanolejeunea = *Cololejeunea* 22*A. azorica* = *Cololejeunea azorica* 22*A. madeirensis* = *Cololejeunea madeirensis* 22*A. microscopica* = *Cololejeunea microscopica* 22*A. sintensisii* = *Cololejeunea sintensisii* 22*Aphanorrhagma* = *Physcomitrium* 31*A. patens* = *Physcomitrium patens* 31**Aplodon** 48**A. wormskioldii** 48*Apomarsupella* = *Gymnomitrium* 16*A. revoluta* = *Gymnomitrium revolutum* 16*Apometzgeria* = *Metzgeria* 24*A. pubescens* = *Metzgeria pubescens* 24**Apopellia** 24**A. endiviifolia** 24**Aquilonium** 63**A. plicatum** 63

ARCHIDIACEAE 7, 32

Archidium 32**A. alternifolium** 32**Arctoa** 35

A. anderssonii 35**A. fulvella** 35**A. hyperborea** 35

Ardeuma = Hymenostylium 42

A. annotinum = Hymenostylium recurvirostrum var. insigne 42

A. insigne = Hymenostylium recurvirostrum var. insigne 42

A. recurvirostrum = Hymenostylium recurvirostrum var. recurvirostrum 42

Arnellia 16**A. fennica** 16

ARNELLIACEAE 6, 16

Arvernella 56**A. microclada** 56**Aschisma** 40**A. carniolicum** 40**A. cuynetii** 40

Asterella = [Fimbraria] 25

A. subgenus Asterella 25

A. subgenus Phragmoblepharis 25

A. subgenus Saccatae 25

A. section Brachyblepharis 25

A. africana 25

[A. blumeana] 66

[A. elegans] 66

A. gracilis = Mannia gracilis 25

A. lindenbergiana 25

[A. marginata] 66

[A. pringlei] 66

A. saccata 25

[A. tenella] 66

[A. wallachiana] 66

Athalamia auct. eur. = Clevea 25

A. hyalina = Clevea hyalina 25

var. suecica = Athalamia hyalina 25

A. spathysii = Clevea spathysii 25

Atractylocarpus 32**A. alpinus** 32**A. subporodictyon** 32**Atrichum** 29

[A. acuminatum] 69

A. androgynum 29**A. angustatum** 29**A. crispum** 4, 29**A. flavisetum** 29**A. tenellum** 29**A. undulatum** 29

AULACOMNIACEAE 8, 55

AULACOMNIALES 8, 55

Aulacomnium 55

[A. acuminatum] 69

A. androgynum 55**A. palustre** 55**A. turgidum** 55

AYTONIACEAE 7, 25

Barbilophozia 9

B. subgenus Barbilophozia 9

B. subgenus Sudeticae 9

B. atlantica = Orthocaulis atlanticus 10

B. attenuata = Neoorthocaulis attenuatus 9

B. barbata 9

B. binsteadii = Neoorthocaulis binsteadii 10

B. floerkei = Neoorthocaulis floerkei 10

B. hatcheri 9

B. hyperborea = Neoorthocaulis hyperboreus 10

B. kunzeana = Schljakovia kunzeana 10

B. lycopodioides 9

B. quadriloba = Schljakovianthus quadrilobus 10

B. rubescens 9**B. sudetica** 9**Barbula** 36

B. amplexifolia = Hydrogonium amplexifolium 41

B. bicolor = Gymnobarbula bicolor 41

B. bolleana = Hydrogonium bolleanum 41

B. consanguinea = Hydrogonium consanguineum 41

B. convoluta = Streblotrichum convolutum 40

var. sardoa = Streblotrichum convolutum var. commutatum 40

B. crocea = Hydrogonium croceum 41

B. enderesii = Streblotrichum enderesii 70

[B. indica] 41, 68

B. unguiculata 36

Barbuloideae = Pottioideae 36

Bartramia 47

B. section Bartramia 47

B. section Pyridium 47

B. section Strictidium 47

B. aprica 47, 68**B. breviseta** 47**B. halleriana** 47**B. ithyphylla** 47**B. laevisphaera** 47**B. pomiformis** 47

B. rosamrosiae = Bartramia aprica 47

[B. stricta] 68

B. stricta auct. eur. = Bartramia aprica 47

B. subulata 47

BARTRAMIACEAE 7, 47

BARTRAMIALES 7, 47

Bartramioidae 47

Bazzania 19**B. azorica** 19

[B. denudata] 66

B. flaccida 19**B. pearsonii** 19**B. tricrenata** 19

[var. fulfordiae] 19

var. tricrenata 19

B. trilobata 19

var. depauperata 19

var. trilobata 19

Bazzanioideae 19

Biantheridion 9**B. undulifolium** 9**Blasia** 25**B. pusilla** 25

BLASIACEAE 7, 25

BLASIALES 7, 25

Blasiidae 7

Blepharostoma 19

[B. arachnoideum] 66

B. trichophyllum 19

subsp. brevirete 19, 66

subsp. trichophyllum 19

var. brevirete = Blepharostoma trichophyllum subsp. brevirete 19

BLEPHAROSTOMATACEAE 6, 19

Blindia 43**B. acuta** 43**B. caespiticia** 43**Blindiadelphus** 43**B. campylopodus** 43**B. diversifolius** 43**B. polaris** 43

- B. recurvatus** 43
B. subimmersus 43
 Brachiolejeuneae (trib.) 22
 Brachiolejeuneinae (subtrib.) 22
Brachydontium 43
B. trichodes 44
Brachymenium 3, 48, 49, 50, 60
 [*B. acuminatum*] 69
 [*B. commutatum*] 68, 69
 [*B. exile*] 69
B. notarisii = *Anomobryum notarisii* 48
B. paradoxum 48
B. philonotula = *Bryum kikuyense* 49
 BRACHYTHECIAACEAE 8, 59
Brachytheciastrum 60
 [*B. bellicum*] 69
B. collinum 60, 68
B. dieckei 60
 [*B. fendleri*] 60, 68
B. fendleri auct. eur. = *Brachytheciastrum collinum* 60
B. olympicum 60
B. salicinum 60
B. trachypodium 60
 [*B. umbilicatum*] 69
B. velutinum 60
 var. *vagans* = *Brachytheciastrum velutinum* 61
 var. *salicinum* = *Brachytheciastrum salicinum* 60
B. vanekii = *Brachytheciastrum velutinum* 61
 Brachythecioideae 60
Brachythecium 61
B. albicans 61
B. buchananii 61
B. campestre 61
B. capillaceum 61
B. cirrosum 61
B. coruscum = *Brachythecium tauriscorum* 61
B. erythrorrhizon 61
 subsp. *asiaticum* = *Brachythecium erythrorrhizon* 61
 var. *thedenii* = *Brachythecium erythrorrhizon* 61
B. funkii 61
B. geheebii 61
B. glareosum 61
B. japygum 61
B. laetum 61
B. mildeanum 61
 var. *udum* = *Brachythecium udum* 61
B. novae-angliae 61
B. percurrens = *Hedenasiastrum percurrens* 60
B. rotaeantum = *Brachythecium capillaceum* 61
B. rivulare 61
B. rutabulum 61
 var. *atlanticum* 61
 var. *rutabulum* 61
B. salebrosum 61
B. scabridum = *Brachythecium novae-angliae* 61
B. tauriscorum 61
B. tenuicaule 61
B. tommasinii 61
B. turgidum 61
B. udum 61
Braunia 46, 47
B. alopecura 47
B. imberbis 47
Breidleria = *Stereodon* 62
B. pratensis = *Stereodon pratensis* 63
Breutelia 47
B. azorica 47
B. chrysocoma 47
 Breutelioideae 47
Brotherella 62
 [*B. henonii*] 62, 68
B. henonii auct. eur. = *Brotherella lorentziana* 62
B. lorentziana 62, 68
Bruchia 36
B. flexuosa 36
B. vogesiaca 36
 BRUCHIACEAE 7, 36
 BRYALES 8, 48
 BRYACEAE 2, 3, 8, 48, 50
Bryhnia = *Brachythecium* 61
B. novae-angliae = *Brachythecium novae-angliae* 61
B. scabrida = *Brachythecium novae-angliae* 61
 Bryidae 7
Bryobrittonia 30
B. longipes 30
Bryoerythrophyllum 37
B. alpigenum 37
B. caledonicum 37
B. campylocarpum 37
B. duellii 37
B. ferruginascens 37
B. inaequalifolium 37
B. recurvirostrum 37
B. rubrum 37
 Bryophyta 7, 27
 Bryophytina 7
 Bryopsida 7, 30
 BRYOXIPHIACEAE 7, 32
Bryoxiphium 32
B. madeirense 32
B. norvegicum 32
Bryum 3, 48
B. algovicum = *Ptychostomum compactum* 50
B. algovicum var. *rutheanum* = *Ptychostomum algovicum* var. *rutheanum* 50
B. alpinum = *Imbribryum alpinum* 49
 [*B. anomodon*] 69
B. apiculatum = *Anomobryum apiculatum* 5, 48
B. archangelicum = *Ptychostomum inclinatum* 50
B. arcticum = *Ptychostomum arcticum* 50
B. argenteum 48
 [*B. atrovirens*] 69
B. austriacum = *Ptychostomum austriacum* 6, 50
B. barnesii = *Bryum dichotomum* 48, 49
B. bavaricum 48
B. blindii 48
B. boreale = *Ptychostomum pallescens* 51
B. bornholmense = *Ptychostomum bornholmense* 50
B. caespiticium = *Ptychostomum imbricatum* 3, 50
B. calophyllum = *Ptychostomum calophyllum* 50
B. canariense 48
B. capillare = *Ptychostomum capillare* 50
B. caucasicum 48
B. cellulare = *Ptychostomum cellulare* 6, 50
B. compactum = *Ptychostomum compactum* 50
B. creberrimum = *Ptychostomum creberrimum* 50
B. cryophilum = *Ptychostomum cryophilum* 50
B. cyclophyllum = *Ptychostomum cyclophyllum* 50
B. demaretianum 48
B. dichotomum 3, 48, 49
B. dixonii 49
B. donianum = *Ptychostomum donianum* 50
B. dyffrynense 49
B. elegans = *Ptychostomum elegans* 6, 50

- B. funkii* = *Ptychostomum funkii* 50
B. gemmiferum 49
B. gemmilucens 49
B. gemmiparum 49
B. inclinatum = *Distichium inclinatum* 50
B. intermedium = *Ptychostomum intermedium* 50
B. kikuyense 49, 68
B. klinggraeffii 49
B. knowltonii = *Ptychostomum knowltonii* 50
B. kunzei = *Ptychostomum kunzei* 50, 51
B. longisetum = *Ptychostomum longisetum* 51
B. marratii 49
B. mildeanum = *Imbribryum mildeanum* 49
B. miniatum = *Imbribryum miniatum* 49
B. minii = *Ptychostomum minii* 6, 51
B. moravicum = *Ptychostomum moravicum* 51
B. muehlenbeckii = *Imbribryum muehlenbeckii* 49
B. neodamense = *Ptychostomum pseudotriquetrum* 51
B. notarisii = *Anomobryum notarisii* 5
B. oblongum 49
B. pallens = *Ptychostomum pallens* 51
 [subsp. *acutiusculum* = *Ptychostomum pallens*] 51
 subsp. *prosboreum* = *Ptychostomum pallens* 51
 subsp. *sibiricum* = *Ptychostomum pallens* 51
 f. *gracile* = *Ptychostomum pallens* 51
B. pallescens = *Ptychostomum pallescens* 51
 [*B. philonotula* Müll.Hal.] 68
B. philonotula Hampe = *Bryum kikuyense* 49
 [*B. philonotulum*] 49, 68
B. philonotulum auct. eur. = *Bryum kikuyense* 49
B. pseudotriquetrum = *Ptychostomum pseudotriquetrum* 51
 var. *bimum* = *Ptychostomum pseudotriquetrum* var.
bimum 51
B. radiculosum 49
B. reyeri 49
B. riparium 49
B. rubens = *Ptychostomum rubens* 51
B. ruderale 49
B. salinum = *Ptychostomum salinum* 51
B. sauteri 49
B. schleicheri = *Ptychostomum schleicheri* 6, 51
 var. *latifolium* = *Ptychostomum schleicheri* var.
latifolium 6, 51
B. sibiricum = *Ptychostomum pallens* 51
B. subapiculatum = *Imbribryum subapiculatum* 5, 49, 50
B. tenuisetum = *Imbribryum tenuisetum* 5, 49, 50
B. torquescens = *Ptychostomum torquescens* 51
B. turbinatum = *Ptychostomum turbinatum* 51
B. uliginosum = *Ptychostomum cernuum* 50
B. valparaisense 4, 49
B. versicolor 49
B. violaceum 49
B. warneum = *Ptychostomum warneum* 51
B. weigeli = *Ptychostomum weigeli* 51
B. wrightii = *Ptychostomum wrightii* 51
Bucegia = *Marchantia* subgenus *Preissia* 26
B. romanica = *Marchantia romanica* 26
Buckia 63
B. vaucheri 63
Buckiella = *Plagiothecium* 55
B. undulata = *Plagiothecium undulatum* 56
Bucklandiella = *Racomitrium* subgenus *Ellipticodryptodon* 44
B. affinis = *Racomitrium affine* 44
B. elliptica = *Racomitrium ellipticum* 44
B. heterosticha = *Racomitrium heterostichum* 44
B. himalayana = *Racomitrium himalayanicum* 44
B. lamprocarpa = *Racomitrium lamprocarpum* 44
B. lusitanica = *Racomitrium lusitanicum* 44
B. macounii = *Racomitrium macounii* 44
 subsp. *alpina* = *Racomitrium macounii* subsp. *alpinum* 44
B. microcarpa = *Racomitrium microcarpon* 44
B. nivalis = *Racomitrium nivale* 44
B. obtusa = *Racomitrium obtusum* 44
B. sudetica = *Racomitrium sudeticum* 44
Buxbaumia 30
B. aphylla 30
B. viridis 30
 BUXBAUMIACEAE 7, 30
 BUXBAUMIALES 7, 30
 Buxbaumiidae 7
 CALLICLADIACEAE 8, 62
Callicladium 62
C. haldanianum 62
C. imponens 62
Calliergon 58
C. cordifolium 58
C. giganteum 58
C. megalophyllum 58
C. richardsonii 58
 CALLIERGONACEAE 2, 8, 58
Calliergonella 63
C. cuspidata 63
C. lindbergii 63
 CALOBRYALES 6, 8
 [*Calohypnum plumiforme*] 69
Calomnion 55
C. complanatum 4, 55
Calycularia 24
C. laxa 24
 Calyculariineae 7
 CALYCVLARIACEAE 7, 24
 CALYMPERACEAE 7, 35
Calymperes 35
C. erosum 35
Calypogeia 16
C. subgenus *Asperifoliae* 16
C. subgenus *Calypogeia* 16
C. arguta 16
C. azorica 16
C. azurea 16
C. fissa 16
 subsp. *fissa* 16
 subsp. *neogaea* 16
 var. *paludosa* 16
C. integristipula 16
C. muelleriana 16
 [subsp. *blomquistii*] 16
 subsp. *muelleriana* 16
 var. *erecta* = *Calypogeia muelleriana* subsp. *muelleriana* 16
C. neesiana 16
 subsp. *neesiana* 16
 [subsp. *subalpina*] 16
 var. *hygrophila* = *Calypogeia neesiana* var. *neesiana* 16
 var. *repanda* = *Calypogeia neesiana* subsp. *neesiana* 16
C. sphagnicola 16
 f. *paludicola* = *Calypogeia fissa* var. *paludosa* 16
C. suecica 16
 CALYPOGEIACEAE 6, 16
Calypstrochaeta 55
C. apiculata 4, 55
Campyliadelphus 56
C. chrysophyllus 56
C. elodes 56
Campylidium = *Campylophyllopsis* 57

C. calcareum = *Campylophyllopsis calcarea* 57
C. sommerfeltii = *Campylophyllopsis sommerfeltii* 57
Campylium 56
C. bambergeri 56
C. decipiens = *Drepanocladus polygamus* 57
C. laxifolium 56
C. longicuspis 57
C. protensum 57
C. stellatum 57
Campylophyllopsis 57
C. calcarea 57
C. sommerfeltii 57
Campylophyllum 57
C. calcareum = *Campylophyllopsis calcarea* 57
C. halleri 57
C. montanum 57
C. sommerfeltii = *Campylophyllopsis sommerfeltii* 57
Campylopus 33
C. atrovirens 33
 var. *falcatus* = *Campylopus atrovirens* 33
C. brevipilus 33
C. cygneus 33
C. flaccidus 33
C. flexuosus 33
C. fragilis 33
C. gracilis 33
C. incrassatus 33
C. introflexus 4, 33
C. oerstedianus 33
C. pilifer 33
C. pyriformis 33
 var. *azoricus* = *Campylopus pyriformis* 33
C. schimperi 33
C. schwarzii = *Campylopus gracilis* 33
C. setifolius 33
C. shawii 33
C. subulatus 33
 var. *schimperi* = *Campylopus schimperi* 33
C. subporodictyon = *Atractylocarpus subporodictyon* 32
Campylostelium 44
C. pitardii 44
C. saxicola 44
C. strictum 44
 CATOSCOPIACEAE 7, 32
 CATOSCOPIALES 7, 32
Catoscopium 32
C. nigratum 32
Cephalozia 10
C. affinis = *Fuscocephaloziopsis affinis* 10
C. ambigua 10
C. bicuspidata 10
 subsp. *bicuspidata* 10
 subsp. *lammersiana* 10
 var. *paludosa* = *Cephalozia ambigua* 10
C. catenulata = *Fuscocephaloziopsis catenulata* 10
C. connivens = *Fuscocephaloziopsis connivens* 10
 var. *compacta* = *Fuscocephaloziopsis connivens* subsp. *connivens* 10
C. crassifolia = *Fuscocephaloziopsis crassifolia* 11
C. crossii 10
C. lacinulata 10
C. leucantha = *Fuscocephaloziopsis leucantha* 11
 var. *robusta* = *Fuscocephaloziopsis leucantha* 11
C. loitlesbergeri = *Fuscocephaloziopsis loitlesbergeri* 11
 [*Cephalozia lucens*] 66
C. lunulifolia = *Fuscocephaloziopsis lunulifolia* 11
C. macounii 10

C. macrostachya = *Fuscocephaloziopsis macrostachya* 11
C. pleniceps = *Fuscocephaloziopsis pleniceps* 11
 var. *sphagnorum* = *Fuscocephaloziopsis pleniceps* var. *pleniceps* 11

CEPHALOZIACEAE 6, 10

Cephaloziella 11

C. subgenus *Cephaloziella* 11
C. subgenus *Dichiton* 12
C. subgenus *Evansia* 12
C. subgenus *Prionolobus* 12
C. alpina = *Cephaloziella varians* 12
C. arctica = *Cephaloziella varians* 12
C. arctogena 11
C. aspericaulis 11
C. baumgartneri 11
C. calyculata 12
C. dentata 12
C. divaricata 11
 var. *asperifolia* = *Cephaloziella divaricata* var. *scabra* 11
 var. *divaricata* 11
 var. *polystratosa* = *Cephaloziella polystratosa* 12
 var. *rupestris* = *Cephaloziella divaricata* var. *divaricata* 11
 var. *scabra* 11
C. elachista 11
 var. *elachista* 11
 [var. *spinophylla*] 11
C. elegans 11
C. granatensis 12
C. grimsulana 11
C. hampeana 11
 var. *subtilis* = *Cephaloziella hampeana* 11
C. integerrima 11
 var. *obtusa* = *Cephaloziella integerrima* 12
 [*C. mammillifera*] 66
C. massalongi 12
 var. *compacta* = *Cephaloziella massalongi* 11
C. nicholsonii 11
C. phyllacantha 12
C. polystratosa 12
C. rubella 12
 subsp. *arctogena* = *Cephaloziella arctogena* 11
 var. *bifida* = *Cephaloziella rubella* 12
 var. *elegans* = *Cephaloziella elegans* 11
 var. *pulchella* = *Cephaloziella rubella* 12
 var. *sullivantii* = *Cephaloziella rubella* 12
C. spinigera 12
 f. *striatula* = *Cephaloziella spinigera* 12
C. stellulifera 12
 var. *limprichtii* = *Cephaloziella stellulifera* 12
C. subdentata = *Cephaloziella spinigera* 12
C. turneri 12
C. uncinata 12
 [var. *brevigyna*] 12
 [var. *mamillosa*] 12
 [var. *sphagnicola*] 12
 var. *uncinata* 12
C. varians 12
 var. *arctica* = *Cephaloziella varians* 12
 var. *scabra* = *Cephaloziella varians* 12, 66
 [*C. verrucosa*] 66
 CEPHALOZIELLACEAE 6, 11
 Cephaloziineae 6
 Cephalozioidaeae 10
Ceratodon 36
C. amazonum 36
C. conicus 36
C. purpureus 36

- subsp. **purpureus** 36
subsp. **stenocarpus** 36
var. *conicus* = *Ceratodon conicus* 36
- Cheilolejeunea** 22
C. subgenus *Euosmolejeunea* 22
- C. cedercreutzii** 22
[C. *xanthocarpa*] 68
Cheilolejeuneinae (subtrib.) 22
- Cheilothela** 36
C. chloropus 36
Chenia 37
C. leptophylla 4, 37
C. ruigtevleia 37
Chiloscyphus 20
C. *coadunatus* = *Lophocolea coadunata* 20
C. *fragilis* = *Chiloscyphus pallescens* var. *fragilis* 20
C. *fragrans* = *Lophocolea fragrans* 20
C. *latifolius* = *Lophocolea bidentata* 20
C. *minor* = *Lophocolea minor* 20
C. pallescens 20
var. *fragilis* 20
var. *pallescens* 20
C. polyanthos 20
var. *polyanthos* 20
var. *rivularis* 20
C. *profundus* = *Lophocolea heterophylla* 20
C. *rivularis* = *Chiloscyphus polyanthos* var. *rivularis* 20
- Chionoloma** 40
C. bombayense 40
C. daldinianum 40
C. hibernicum 40
C. minus 41
C. recurvifolium 41
C. tenuirostre 41
var. *holtii* 41
var. *tenuirostre* 41
- Cinclidium** 52
C. arcticum 52
C. latifolium 52
C. minutifolium 52
C. stygium 52
C. subrotundum 52
Cinclidotus 37
[C. *bistratosus*] 69
C. *confertus* = *Cinclidotus riparius* 37
C. aquaticus 37
C. danubicus 37
C. fontinaloides 37
[C. *pachyloma*] 69
[C. *pachylomoides*] 68
C. riparius 37
[C. *vardaranus*] 69
C. vivesii 37
- Cirriphyllum** 60
C. crassinervium 60
C. piliferum 60
Cladopodiella = *Odontoschisma* 11
C. *francisci* = *Odontoschisma francisci* 11
C. *fluitans* = *Odontoschisma fluitans* 11
- Claopodium** 58
C. rostratum 58
C. whippleanum 58
Clasmatodon 60
C. parvulus 60
Cleistocarpidium 36
C. palustre 36
Clevea 25
- C. hyalina** 25
var. *rufescens* = *Clevea hyalina* 25
var. *suecica* = *Clevea hyalina* 25
- C. spathysii** 25
CLEVEACEAE 7, 25
CLIMACIACEAE 8, 56
Climacium 56
C. dendroides 56
Cnestrum 35
C. alpestre 35
C. glaucescens 35
C. schisti 35
Codonoblepharon 53
C. forsteri 53
Codriophorus = *Racomitrium* subgenus *Cataracta* 44
C. *acicularis* = *Racomitrium aciculare* 44
C. *aquaticus* = *Racomitrium aquaticus* 44
C. *fascicularis* = *Racomitrium fasciculare* 44
C. *hespericus* = *Racomitrium hespericum* 44
- Cololejeunea** 22
C. subgenus *Aphanolejeunea* 22
C. subgenus *Cololejeunea* 22
C. subgenus *Diaphanae* 22
C. azorica 22
C. calcarea 22
C. madeirensis 22
C. microscopica 22
[*var. africana*] 22
[*var. exigua*] 22
var. *microscopica* 22
C. *minutissima* = *Myriocoleopsis minutissima* 22
C. rossettiana 22
C. schaeferi 22
C. sintensisii 22
Cololejeuneinae (subtrib.) 22
- Colura** 22
C. subgenus *Colura* 22
C. section *Colura* 22
C. calyptrifolia 22
Conardia 57
C. compacta 57
CONOCEPHALACEAE 7, 25
Conocephalum 25
C. conicum 25
C. salebrosum 25
Conostomoideae 47
Conostomum 47
C. tetragonum 47
Corsinia 25
C. coriandrina 25
CORSINIACEAE 7, 25
Corsinioideae 25
Coscinodon 2, 44
C. cribrus 44
C. horridus 44
C. humilis 45
C. monchiquensis 45
[C. *yukonensis*] 45
Cratoneuroideae 56
Cratoneuron 56
C. curvicaule 56
C. filicinum 56
Crossidium 37
C. aberrans 37
C. crassinervium 37
C. davidai 37
C. geheebii 37

C. laevipilum 37**C. laxefilamentosum** 37**C. squamiferum** 37var. **pottioideum** 37var. **squamiferum** 37**Crossocalyx** 9**C. hellerianus** 9*Crossogyna* = *Syzygiella* 9*C. autumnalis* = *Syzygiella autumnalis* 9[*Crumia latifolia*] 69**Cryphaea** 64**C. heteromalla** 64

CRYPHAEEACEAE 8, 64

Cryptoleptodon = *Leptodon* 64*C. longisetus* = *Leptodon longisetus* 64**Cryptocolea** 18**C. imbricata** 18*Cryptothallus* = *Aneura* 23*C. mirabilis* = *Aneura mirabilis* 23**Ctenidium** 65**C. molluscum** 65*C. procerrimum* = *Pseudostereodon procerrimum* 63

CYATHODIACEAE 7, 26

Cyathodium 26[*C. cavernarum*] 66, 68**C. foetidissimum** 26[*C. spurium*] 66**Cyclodictyon** 55**C. laetevirens** 55**Cynodontium** 35**C. asperifolium** 35**C. bruntonii** 35**C. fallax** 35**C. gracilescens** 35**C. jenneri** 35**C. polycarpon** 35**C. strumiferum** 35**C. suecicum** 35**C. tenellum** 35**Cyrtomnium** 52**C. hymenophylloides** 52**C. hymenophyllum** 52**Daltonia** 55**D. lindigiana** 55**D. splachnoides** 55*D. stenophylla* = *Daltonia lindigiana* 55

DALTONIACEAE 8, 55

Delavayelloideae 17

Dendrocryphaea 64**D. lamyana** 64*Desmatodon* = *Tortula* 39*D. bogosicus* = *Tortula bogosica* 39*D. guepinii* = *Tortula guepinii* 40*D. latifolius* = *Tortula hoppeana* 40*D. laureri* = *Tortula laureri* 40*D. leucostoma* = *Tortula leucostoma* 40*D. meridionalis* = *Tortula marginata* 40*D. randii* = *Tortula randii* 40*D. systylius* = *Tortula systylia* 40*D. ucrainicus* = *Tortula ucrainica* 40*D. wilczekii* = *Henediella heimii* 38**Dialytrichia** 37*D. fragilifolia* = *Dialytrichia saxicola* 37**D. mucronata** 37**D. saxicola** 37**Dichelyma** 55**D. capillaceum** 55**D. falcatum** 55*Dichiton* = *Cephaloziella* 11*D. integerrimum* = *Cephaloziella integerrima* 12**Dichodontium** 33**D. flavescens** 33*D. palustre* = *Diobelonella palustris* 33**D. pellucidum** 33

DICRANACEAE 7, 34

DICRANALES 7, 32

Dicranella 33**D. campylophylla** 33**D. cerviculata** 33**D. crispa** 33**D. grevilleana** 33**D. heteromalla** 33**D. howei** 33**D. humilis** 33*D. palustris* = *Diobelonella palustris* 33*D. riparia* = *Kiaeria riparia* 35**D. rufescens** 33**D. schreberiana** 33**D. staphylina** 33**D. subulata** 33**D. varia** 33

DICRANELLACEAE 7, 33

Dicranidae 7

Dicranoideae 34

Dicranodontium 33**D. asperulum** 33**D. denudatum** 33var. *alpinum* = *Dicranodontium denudatum* 33*D. subporodictyon* = *Atractylolcarpus subporodictyon* 32**D. uncinatum** 33**Dicranoloma** 34**D. menziesii** 4, 34**Dicranoweisia** 35**D. cirrata** 35*D. compacta* = *Hymenoloma compactum* 32*D. crispula* = *Hymenoloma crispulum* 32var. *compacta* = *Hymenoloma compactum* 32var. *intermedia* = *Hymenoloma mulahaceni* 32*D. intermedia* = *Hymenoloma mulahaceni* 32**Dicranum** 34*D.* section *Convolvutifolia* 34*D.* section *Crassinervia* 35*D.* section *Dicranum* 34*D.* section *Elongata* 35*D.* section *Fuscescentiformia* 34*D.* section *Montana* 35*D.* section *Spuria* 34**D. acutifolium** 34**D. angustum** 34**D. bardunovii** 34*D. bergeri* = *Dicranum undulatum* 34var. *acutifolium* = *Dicranum acutifolium* 34**D. bonjeanii** 34**D. brevifolium** 34*D. canariense* = *Dicranum scottianum* 35*D. congestum* = *Dicranum flexicaule* 34**D. crassifolium** 34**D. dispersum** 34**D. drummondii** 34**D. elongatum** 35**D. flagellare** 35**D. flexicaule** 34**D. fragilifolium** 35**D. fulvum** 35

D. fuscescens 34var. *congestum* = *Dicranum flexicaule* 34var. *flexicaule* = *Dicranum flexicaule* 34**D. groenlandicum 35****D. laevidens 35****D. leioneuron 34****D. majus 34****D. montanum 35****D. muehlenbeckii 34**var. *acutifolium* = *Dicranum acutifolium* 34var. *brevifolium* = *Dicranum brevifolium* 34var. *cirrhatum* = *Dicranum brevifolium* 34**D. polysetum 34***D. rugosum* = *Dicranum polysetum* 34**D. schljakovii 35****D. scoparium 34****D. scottianum 35****D. septentrionale 34****D. spadiceum 35****D. spurium 34***D. strictum* = *Dicranum tauricum* 35*D. subporodictyon* = *Atractylocarpus subporodictyon* 32**D. tauricum 34****D. transsylvanicum 34****D. undulatum 34***D. undulatum* F.Weber & D.Mohr, hom. illeg. = *Dicranum polysetum* 34*D. undulatum* Turner, hom. illeg. = *Dicranum bonjeanii* 34**D. viride 35****Didymodon 5, 37****D. acutus 37****D. asperifolius 37****D. australasiae 37****D. bistratosus 37****D. brachyphyllus 37**[*D. caboverdeanus*] 69**D. cordatus 37***D. cylindricus* var. *daldianus* = *Chionoloma daldinianum* 40**D. eckeliae 37***D. erosus* = *Didymodon tophaceus* subsp. *erosus* 38**D. fallax 37****D. ferrugineus 37****D. giganteus 37****D. glaucus 5, 37**subsp. *glaucus* 37subsp. *verbanus* 5, 37**D. icmadophilus 37****D. insulanus 37****D. johansenii 37***D. lamyanus* = *Didymodon brachyphyllus* 37**D. luridus 37****D. maschalogenus 38****D. maximus 38****D. nicholsonii 38****D. rigidulus 38***D. sicculus* = *Didymodon tophaceus* subsp. *sicculus* 38**D. sinuosus 38****D. spadiceus 38****D. subandreaeoides 38**[*D. tectorum*] 69**D. tomaculosus 38****D. tophaceus 38**subsp. *erosus* 38subsp. *sicculus* 38subsp. *tophaceus* 38var. *anatinus* 38var. *tophaceus* 38**D. umbrosus 38****D. validus 38***D. verbanus* = *Didymodon glaucus* subsp. *verbanus* 37**D. vinealis 38***Dilutineuron* = *Racomitrium* subgenus *Cataracta* 44*D. fasciculare* = *Racomitrium fasciculare* 44**Diobelonella 33****D. palustris 33**

DIPHYSCIACEAE 7, 30

DIPHYSCIALES 7, 30

Diphysciidae 7

Diphyscium 30**D. foliosum 30****Diplophyllum 14***D.* section *Diplophyllum* 14*D.* section *Protodiplophyllum* 14**D. albicans 14****D. obtusatum 14****D. obtusifolium 14**[subsp. *domesticum*] 14subsp. *obtusifolium* 14[*D. sibiricum*] 14**D. taxifolium 14**var. *macrostictum* = *Diplophyllum taxifolium* 14[var. *mucronatum*] 14var. *taxifolium* 14

DISCELIACEAE 7, 32

Discelium 32**D. nudum 32**

DISTICHIACEAE 7, 32

Distichium 32**D. capillaceum 32****D. hagenii 32****D. inclinatum 32, 50****Distichophyllum 55****D. carinatum 55**

DITRICHACEAE 7, 32, 36

Ditrichum 36**D. cornubicum 36***D. crispatisimum* = *Flexitrichum gracile* 32*D. cylindricum* = *Trichodon cylindricus* 36*D. flexicaule* = *Flexitrichum flexicaule* 32*D. gracile* = *Flexitrichum gracile* 32**D. heteromallum 36****D. lineare 36****D. pallidum 36****D. plumbicola 36****D. punctulatum 36****D. pusillum 36****D. subulatum 36****D. zonatum 36**var. *scabrifolium* = *Ditrichum zonatum* 36**Douinia 14****D. ovata 14****Drepanium 57****D. fastigiatum 57***D. recurvatum* = *Drepanium fastigiatum* 57**Drepanocladus 57****D. aduncus 57****D. angustifolius 57****D. arcticus 57****D. brevifolius 57****D. capillifolius 57**[*D. longifolius*] 57, 68*D. longifolius* auct. eur. = *Drepanocladus capillifolius* 57**D. lycopodioides 57****D. polygamus 57**

D. sendtneri* 57**D. sordidus* 57*****D. trifarius* 57*****D. turgescens* 57*****Drepanolejeunea* 22***D.* subgenus *Drepanolejeunea* 22***D. hamatifolia* 22***Drepanolejeuneinae* (subtrib.) 22*Dryptodon* = *Grimmia* 45*D. orbicularis* = *Grimmia orbicularis* 45*D. patens* = *Grimmia ramondii* 45***Dumortiera* 26*****D. hirsuta* 26**subsp. *hirsuta* 26[subsp. *nepalensis*] 26[subsp. *tatunoi*] 26**DUMORTIERACEAE 7, 26****ECHINODIACEAE 8, 65*****Echinodium* 65***E. prolixum* = *Isothecium prolixum* 65***E. renaudii* 65*****E. setigerum* 65*****E. spinosum* 65***Elodium* = *Helodium* 59*E. blandowii* = *Helodium blandowii* 59***Encalypta* 31***E.* section *Encalypta* 31*E.* section *Megasporae* 31*E.* section *Pyromitrium* 31*E.* section *Rhabdotheca* 31*E.* section *Streptotheca* 31***E. affinis* 31**subsp. *affinis* 31subsp. *macounii* 31***E. alpina* 31*****E. brevicolla* 31*****E. brevipes* 31*****E. ciliata* 31***E. intermedia* = *Encalypta pilifera* 31***E. longicolla* 31*****E. microstoma* 31*****E. mutica* 31***E. obovatifolia* = *Encalypta pilifera* 31***E. pilifera* 31*****E. procera* 31*****E. raptocarpa* 31**var. *leptodon* = *Encalypta trachymitria* 31var. *spathulata* = *Encalypta spathulata* 31var. *trachymitria* = *Encalypta trachymitria* 31***E. spathulata* 31*****E. streptocarpa* 31*****E. trachymitria* 31*****E. vulgaris* 31****ENCALYPTACEAE 7, 30****ENCALYPTALES 7, 30****Encalyptidae 7*****Endogemma* 16*****E. caespiticia* 16****ENDOGEEMMATACEAE 6, 16*****Entodon* 64*****E. challengeri* 64*****E. cladorrhizans* 64*****E. concinnus* 64**[*E. pseudoseductrix*] 69***E. schleicheri* 64****ENTODONTACEAE 8, 64**[*Entodontopsis leucostega*] 69***Entosthodon* 31***E.* subgenus *Entosthodon* 31*E.* subgenus *Murcia* 31*E.* subgenus *Plagiodus* 31***E. abramovae* 31**[*E. angustifolius*] 69***E. attenuatus* 31*****E. commutatus* 31*****E. convexus* 31*****E. dagestanicus* 31*****E. duriaei* 31*****E. fascicularis* 31*****E. handelii* 31*****E. hungaricus* 31***E. krausei* = *Entosthodon commutatus* 31***E. kroonkurk* 31*****E. mouretii* 31*****E. mühlenbergii* 31**[*E. niloticus*] 69***E. obtusus* 31*****E. pulchellus* 31*****E. schimperi* 31*****E. stenophyllus* 31***E. templetonii* = *Entosthodon attenuatus* 31*Ephemerella* = *Physcomitrium* 31*E. readeri* = *Physcomitrium readeri* 32***Ephemerum* 41*****E. cohaerens* 41*****E. crassinervium* 41**subsp. *rutheanum* 41subsp. *sessile* 41*E. hibernicum* = *Ephemerum crassinervium* subsp.*rutheanum* 41*E. minutissimum* = *Ephemerum serratum* 41***E. recurvifolium* 41***E. rutheanum* = *Ephemerum crassinervium* subsp.*rutheanum* 41***E. serratum* 41**var. *minutissimum* = *Ephemerum serratum* 41var. *rutheanum* = *Ephemerum crassinervium* subsp. *rutheanum* 41*E. serratum* auct. = *Ephemerum stoloniferum* 41*E. sessile* = *Ephemerum crassinervium* subsp. *sessile* 41***E. spinulosum* 41***E. stellatum* = *Ephemerum stoloniferum* 41***E. stoloniferum* 41*****Eipterygium* 52**[*E. rigidum*] 69***E. tozeri* 52*****Eremonotus* 17*****E. myriocarpus* 17**[*Erpodium grossirete*] 69[*Erpodium perrottetii*] 69***Eucladium* 41***E. verbanum* = *Didymodon glaucus* subsp. *verbanum* 5***E. verticillatum* 41**var. *angustifolium* = *Eucladium verticillatum* 41***Eurhynchiastrum* 61*****E. diversifolium* 61*****E. pulchellum* 61**var. *praecox* = *Eurhynchiastrum pulchellum* 61var. *diversifolium* = *Eurhynchiastrum diversifolium* 61**Eurhynchioideae 59*****Eurhynchium* 59*****E. angustirete* 59*****E. striatum* 59***Exobryum* = *Didymodon* 37

- E. asperifolium* = *Didymodon asperifolius* 37
E. johansenii = *Didymodon johansenii* 37
Exormotheca 26
E. subgenus *Corbierella* 26
E. subgenus *Exormotheca* 26
E. pustulosa 26
E. welwitschii 26, 67
 EXORMOTHEACEAE 7, 26
Exsertotheca 64
E. baetica 64
E. crispa 64
E. intermedia 64
Fabronia 56
F. altaica 56
F. ciliaris 56
 [*F. gueinzii*] 69
 [*F. leikipiae*] 69
F. major 56
F. pusilla 56
 FABRONIACEAE 8, 56
 [*Fimbraria* = *Asterella*] 66
 [*F. blumeana* = *Asterella blumeana*] 66
 [*F. elegans* = *Asterella elegans*] 66
 [*F. marginata* = *Asterella marginata*] 66
 [*F. raddii* = *Asterella wallichiana*] 66
 [*F. stahlia* = *Asterella pringlei*] 66
 [*F. tenella* = *Asterella tenella*] 66
Fissidens 5, 33
F. subgenus *Aloma* 34
F. subgenus *Fissidens* 34
F. subgenus *Octodiceria* 34
F. subgenus *Pachyfissidens* 5, 33
F. section *Amblyothallia* 34
F. section *Pachyfissidens* 5, 33
F. adianthoides 33
F. algarvicus = *Fissidens curvatus* 34
 [*F. allorgei*] 69
 [*F. androgynus*] 69
F. arcticus 34
F. arnoldii 34
F. asplenioides 34
F. azoricus 33
F. bambergi = *Fissidens viridulus* 34
F. bryoides 34
 var. *bryoides* 34
 var. *caespitans* 34
 var. *curnovii* = *Fissidens bryoides* var. *caespitans* 34
F. celticus 34
F. coacervatus 34
F. crassipes 34
 subsp. *crassipes* 34
 subsp. *warnstorffii* 34
F. crispus 34
F. cristatus = *Fissidens dubius* 33
F. curnovii = *Fissidens bryoides* var. *caespitans* 34
F. curvatus 34
 [*F. dankelmannii*] 69
F. dubius 33
 var. *dubius* 33
 var. *mucronatus* 33
F. exiguus auct. eur. = *Fissidens viridulus* 34
F. exilis 34
 [*F. flaccidus*] 69
F. fontanus 34
F. gracilifolius 34
F. grandifrons 33
F. gymnandrus 34
F. herzogii = *Fissidens crispus* 34
F. incurvus 34
F. jansenii 34
F. limbatus = *Fissidens crispus* 34
F. luisieri = *Fissidens serrulatus* 34
 [*F. megalotis* subsp. *helictocaulos*] 69
F. microstictus 34
F. minutulus = *Fissidens crispus* 34
F. monguillonii 34
F. nobreganus 34
F. osmundoides 33
F. ovatifolius 34
F. polyphyllus 33
F. pusillus 34
F. rivularis 34
F. rufulus 34
 [*F. sciophyllus*] 69
F. serratus 34
F. serrulatus 33
F. sublimbatus 34
F. sublineaeifolius 34
F. taxifolius 33
 subsp. *pallidicaulis* = *Fissidens taxifolius* 33
 [*F. usambaricus*] 69
F. viridulus 34
 var. *bambergi* = *Fissidens viridulus* 34
 var. *incurvus* = *Fissidens incurvus* 34
 var. *pusillus* = *Fissidens pusillus* 34
 var. *tenuifolius* = *Fissidens gracilifolius* 34
 FISSIDENTACEAE 2, 7, 33
 FLEXITRICHACEAE 7, 32
Flexitrichum 32
F. flexicaule 32
F. gracile 32
 FONTINALACEAE 8, 55
Fontinalis 55
F. antipyretica 55
 subsp. *antipyretica* 55
 subsp. *bryhnii* 55
 subsp. *gracilis* 55
 subsp. *kindbergii* 55
 [var. *heldreichii*] 69
F. dalecarlica 55
F. dichelymoides 55
F. hypnoides 55
 var. *duriaei* 55
 var. *hypnoides* 55
F. squamosa 55
 var. *curnovii* = *Fontinalis squamosa* 55
 var. *dixonii* = *Fontinalis squamosa* 55
Fossombronia 24
F. angulosa 24
F. caespitiformis 24
 subsp. *caespitiformis* 24
 subsp. *multispira* 24
F. crispa 24
F. crozalsii = *Fossombronia mittenii* 24
F. echinata 24
F. fimbriata 24
F. fleischeri 24
F. foveolata 24
F. husnotii = *Fossombronia caespitiformis* subsp. *multispira* 24
F. incurva 24
F. leucoxantha 24
F. maritima 24
F. mittenii 24

F. pusilla 24**F. wondraczekii** 24*F. zeyheri* 24

FOSSOMBRONIACEAE 7, 24

FOSSOMBRONIALES 6, 24

Fossombroniineae 7

Frullania 21*F.* subgenus *Frullania* 21*F.* subgenus *Thyopsiella* 21**F. acicularis** 21**F. azorica** 21, 67**F. bolanderi** 22**F. calcarifera** 22*F. cesatiana* = *Frullania riparia* 21, 67[var. *musciicola* = *Frullania musciicola*] 21**F. cleistostoma** 22[*F. davurica*] 66subsp. *jackii* = *Frullania jackii* 21, 66**F. dilatata** 21[subsp. *asiatica*] 21subsp. *dilatata* 21**F. ericoides** 21var. *ericoides* 21[var. *laxa*] 21[var. *minor*] 21[var. *verrucosa*] 21**F. fragilifolia** 21*F. inflata* auct. = *F. cleistostoma* 22[*F. inflata*] 22, 66, 67**F. jackii** 21, 66**F. microphylla** 21var. *deciduiifolia* = *Frullania microphylla* 21[*Frullania musciicola*] 21, 67**F. oakesiana** 21subsp. *oakesiana* 21[subsp. *takayuensis*] 21[*F. obscurifolia*] 67**F. parvistipula** 21**F. polysticta** 21**F. riparia** 21**F. sergiae** 21[*F. socotrana*] 68[*F. spongiosa*] 68**F. styliifera** 21**F. subarctica** 21**F. tamarisci** 21var. *atrovirens* = *Frullania tamarisci* 21var. *azorica* = *Frullania acicularis* 21var. *cornubica* = *Frullania tamarisci* 21var. *ericetorum* = *Frullania tamarisci* 21var. *robusta* = *Frullania tamarisci* 21var. *sardoa* = *Frullania tamarisci* 21**F. teneriffae** 21

FRULLANIACEAE 6, 21

Funaria 31[*F. anomala*] 4[*F. altissima*] 69[*F. chevalieri*] 69*F. dentata* = *Entosthodon muhlenbergii* 31*F. handelii* = *Entosthodon handelii* 31**F. aequidens** 31**F. arctica** 31**F. hygrometrica** 31**F. microstoma** 31

FUNARIACEAE 2, 7, 31

FUNARIALES 7, 31

Funariidae 7

Funariella 31**F. curviseta** 31

Funarioideae 31

Fuscobryum = *Didymodon* 37*F. subandreaeoides* = *Didymodon subandreaeoides* 38**Fuscocephaloziopsis** 10**F. affinis** 10**F. albescens** 10var. *albescens* 10var. *islandica* 10**F. catenulata** 10subsp. *catenulata* 10[subsp. *nipponica*] 10**F. connivens** 10subsp. *connivens* 10[subsp. *fissa*] 10[subsp. *sandwicensis*] 10**F. crassifolia** 11**F. leucantha** 11**F. loitlesbergeri** 11**F. lunulifolia** 11**F. macrostachya** 11[subsp. *australis*] 11subsp. *macrostachya* 11var. *macrostachya* 11var. *spiniflora* 11**F. pleniceps** 11[var. *caroliniana*] 11var. *pleniceps* 11*Geheebia* = *Didymodon* 37*G. fallax* = *Didymodon fallax* 37*G. ferruginea* = *Didymodon ferrugineus* 37*G. gigantea* = *Didymodon giganteus* 37*G. maschalogena* = *Didymodon maschalogenus* 38*G. maxima* = *Didymodon maximus* 38*G. spadicea* = *Didymodon spadiceus* 38*G. tophacea* = *Didymodon tophaceus* 38*Gemmabryum* = *Bryum* 3, 48*G. apiculatum* = *Anomobryum apiculatum* 48*G. barnesii* = *Bryum dichotomum* 48*G. caespiticium* = *Ptychostomum imbricatum* 50*G. demaretianum* = *Bryum demaretianum* 48*G. dichotomum* = *Bryum dichotomum* 49*G. gemmiferum* = *Bryum gemmiferum* 49*G. gemmilucens* = *Bryum gemmilucens* 49*G. klinggraeffii* = *Bryum klinggraeffii* 49*G. kunzei* = *Ptychostomum kunzei* 50*G. radiculosum* = *Bryum radiculosum* 49*G. ruderale* = *Bryum ruderale* 49*G. subapiculatum* = *Imbribryum subapiculatum* 49*G. tenuisetum* = *Imbribryum tenuisetum* 49*G. valparaisense* = *Bryum valparaisense* 49*G. violaceum* = *Bryum violaceum* 49

GEOCALYCEAE 6, 16

Geocalyx 16**G. graveolens** 16

GIGASPERMACEAE 7, 32

GIGASPERMALES 7, 32

Gigaspermidae 7

Gigaspermum 32**G. mouretii** 32**Glyphomitrium** 35**G. daviesii** 35**Gongylanthus** 19**G. ericetorum** 19

- Goniomitrium** 31
G. seroi 31
Grimmia 2, 45
G. alpestris 45
G. anodon 45
G. anomala 45
G. arenaria 45
G. atrata 45
G. britannica = *Grimmia trichophylla* 45
G. caespiticia 45
G. capillata 45
G. cribrosa = *Coscinodon cribrus* 44
G. crinita 45
G. crinitoleucophaea 45
G. curvata = *Grimmia ramondii* 45
G. curviseta 45
G. decipiens 45
G. dissimulata 45
G. donniana 45
G. elatior 45
G. elongata 45
G. funalis 45
G. fuscolutea 45
G. hartmanii 45
G. horrida = *Coscinodon horridus* 44
G. incurva 45
G. laevigata 45
G. lisae 45
G. longirostris 45
G. meridionalis 45
G. meteorae = *Grimmia nutans* 45
G. mollis 45
G. montana 45
G. muehlenbeckii 45
G. nutans 45
G. orbicularis 45
G. ovalis 45
G. pitardii = *Campylostelium pitardii* 44
G. plagiopodia 45
G. poecilostoma = *Grimmia crinitoleucophaea* 45
G. pulvinata 45
G. ramondii 45
G. reflexidens 45
G. sessitana = *Grimmia reflexidens* 45
G. teretinervis 45
G. tergestina 45
 var. *tergestinoides* = *Grimmia tergestina* 45
G. torquata 45
G. trichophylla 45
 var. *meridionalis* = *Grimmia meridionalis* 45
G. triformis 45
G. ungeri 45
G. unicolor 45
 GRIMMIACEAE 7, 44
 GRIMMIALES 7, 32, 43
 Grimmioideae 44
 [Groutiella tomentosa] 69
Guembelia = *Grimmia* 45
G. laevigata = *Grimmia laevigata* 45
G. longirostris = *Grimmia longirostris* 45
G. ovalis = *Grimmia ovalis* 45
G. tergestina = *Grimmia tergestina* 45
Gymnobarbula 36, 41
G. bicolor 41
Gymnocolea 9
G. acutiloba = *Gymnocolea inflata* subsp. *acutiloba* 9
G. borealis 9
G. fascinifera 9
G. inflata 9
 subsp. *acutiloba* 9
 subsp. *inflata* 9
 var. *heterostipa* = *Gymnocolea inflata* subsp. *inflata* 9
 GYMNOMITRIACEAE 6, 16
 Gymnomitrioideae 16
Gymnomitrium 16
G. adustum 16
G. alpinum 16
G. apiculatum = *Marsupella apiculata* 16
G. brevissimum 16
G. commutatum 16
G. concinnatum 16
G. corallioides 16
G. crenulatum 16
G. obtusum 16
G. revolutum 16
 [subsp. *novoguineanense*] 16
 subsp. *revolutum* 16
 [Gymnostomiella erosula] 69
 [Gymnostomiella vernicosa] 69
Gymnostomum 41
G. aeruginosum 41
 var. *aeruginosum* 41
 var. *obscurum* 41
G. boreale = *Hymenostylium gracillimum* 42
G. calcareum 41
 var. *lanceolatum* = *Gymnostomum calcareum* 41
 var. *atlanticum* 41
 var. *calcareum* 41
G. lanceolatum = *Gymnostomum calcareum* 41
G. recurvirostrum = *Hymenostylium recurvirostrum* 42
G. viridulum 41
Gyroweisia 41
G. reflexa 41
G. tenuis 41
Habrodon 56
H. perpusillus 56
 HABRODONTACEAE 8, 56
Hageniella 63
H. micans 63
Hamatocaulis 58
H. lapponicus 58
H. vernicosus 58
Haplocladium 59
H. angustifolium 59
H. microphyllum 59
H. virginianum 59
Haplohymenium = *Anomodon* 66
H. triste = *Anomodon tristis* 66
 HAPLOMITRIACEAE 6, 8
 Haplomitriidae 6
 Haplomitriopsida 6, 8
Haplomitrium 8
H. subgenus *Haplomitrium* 8
H. section *Haplomitrium* 8
H. hookeri 8
 var. *hookeri* 9
 [var. *minutum*] 9
Harpalejeunea 22
H. subgenus *Harpalejeunea* 22
H. molleri 22
 [subsp. *integra*] 22
 subsp. *molleri* 22
 HARPANTHACEAE 6, 17
Harpantus 17

H. flotovianus 17var. *chiloscyphoides* = *Harpanthus flotovianus* 17var. *latifolia* = *Harpanthus flotovianus* 17var. *retusa* = *Harpanthus flotovianus* 17**H. scutatus 17****Hedenasiastrum 60****H. percurrans 60****Hedwigia 47****H. ciliata 47**var. *leucophaea* = *Hedwigia emodica* 47**H. emodica 47**[*H. integrifolia*] 47*H. integrifolia* = *Braunia imberbis* 47**H. mollis 47****H. nemoralis 47****H. stellata 47****H. striata 47**

HEDWIGIACEAE 7, 46

HEDWIGIALES 7, 46

Hedwigidium = *Braunia* 46*H. integrifolium* = *Braunia imberbis* 47**Helicodontium 60****H. capillare 60**

Helicodontoideae 60

Helodium 59**H. blandowii 59****Hennediella 38****H. heimii 38**var. *arctica* 38var. *heimii* 38**H. macrophylla 4, 38****H. stanfordensis 4, 38**

HERBERTACEAE 6, 19

Herbertus 19[*Herbertus aduncus*] 19, 116subsp. *hutchinsiae* = *H. hutchinsiae* 19*H. aduncus auct eur* = *Herbertus hutchinsiae* 19**H. azoricus 19****H. borealis 19, 67, 116***H. delavayi auct.* = *Herbertus borealis* 19[*H. delavayi*] 19[*H. dicranus*] 67**H. hutchinsiae 19, 116**[*H. juniperoideus*] 19**H. noreus 19, 116**[*H. sakuraii*] 67**H. sendtneri 19****H. stramineus 19, 116**[*H. tenuis*] 19[*Herpetineuron toccoeae*] 69**Herzogiella 55****H. seligeri 55****H. striatella 55****H. turfacea 55****Heterocладиella 65****H. dimorpha 65**

HETEROCLADIACEAE 8, 65

Heterocladium 65*H. dimorphum* = *Heterocладиella dimorpha* 65**H. flaccidum 65****H. heteropterum 65****H. wulfsbergii 65****Heterogemma 12****H. capitata 12****H. laxa 12****Heterophyllum 62***H. affine* = *Heterophyllum nemorosum* 62**H. nemorosum 62****Heteroscyphus 20****H. denticulatus 20****H. fissistipus 4, 20**[var. *fissistipus*] 20[var. *multispinus*] 20[var. *repandus*] 20[*H. integrifolius*] 67**Hilpertia 38****H. velenovskyi 38****Homalia 64****H. lusitanica 64****H. trichomanoides 64***H. webbiana* = *Pseudomalina webbiana* 65**Homalothecium 61****H. aureum 61****H. lutescens 61**var. *fallax* 61var. *lutescens* 61**H. mandonii 61****H. meridionale 61****H. philippeanum 61****H. sericeum 61****Homomallium 63****H. incurvatum 63****Hookeria 55**[*H. acutifolia*] 69**H. lucens 55**

HOOKERIAEAE 8, 55

HOOKERIALES 2, 8, 55

Hydrogonium 36, 41**H. amplexifolium 41**[*H. arcuatum*] 69**H. bolleanum 41****H. consanguineum 41**[var. *cancellatum*] 41[var. *consanguineum*] 41var. *kurilense* 41, 68**H. croceum 41**[*H. orientale*] 68, 69*Hydrogrimmia* = *Grimmia* 45*H. mollis* = *Grimmia mollis* 45**Hygroamblystegium 57****H. fluviatile 57****H. humile 57****H. tenax 57****H. varium 57**var. *humile* = *Hygroamblystegium humile* 57**Hygrobіella 17****H. laxifolia 17**

HYGROBIELLACEAE 6, 17

Hygrohypnella 58**H. ochracea 58****H. polaris 58****Hygrohypnum 57***H. alpestre* = *Platyhypnum alpestre* 57*H. alpinum* = *Platyhypnum alpinum* 57*H. cochlearifolium* = *Platyhypnum cochlearifolium* 58*H. duriusculum* = *Platyhypnum duriusculum* 58*H. eugyrium* = *Pseudohygrohypnum eugyrium* 63**H. luridum 57***H. molle* = *Platyhypnum molle* 58*H. montanum* = *Campylophyllum montanum* 57*H. norvegicum* = *Platyhypnum norvegicum* 58*H. ochraceum* = *Hygrohypnella ochracea* 58*H. polare* = *Hygrohypnella polaris* 58*H. smithii* = *Platyhypnum smithii* 58

H. styriacum 57*H. subeugyrium* = *Pseudohygrohypnum subeugyrium* 63HYLOCOMIACEAE 8, **63****Hylocomiadelphus** 63**H. triquetrus** 63**Hylocomiastrum** 63**H. pyrenaicum** 63**H. umbratum** 63**Hylocomium** 63**H. splendens** 63**Hymenoloma** 32**H. compactum** 32**H. crispulum** 32*H. intermedium* = *H. mulahaceni* 32**H. mulahaceni** 32HYMENOLOMATACEAE 7, **32****Hymenostylium** 42[*H. congoanum*] 69**H. gracillimum** 42[*H. hildebrandtii*] 69*H. insigne* = *Hymenostylium recurvirostrum* var. *insigne* 42**H. recurvirostrum** 42var. *insigne* 42var. *recurvirostrum* 42**H. xerophilum** 42**Hyocomium** 65**H. armoricum** 65**Hyophila** 42**H. involuta** 42HYPNACEAE 8, **62**HYPNALES 8, **55**Hypnelloideae **55****Hypnum** 62*H. aemulans* = *Stereodon aemulans* 63*H. adfine* = *Heterophyllum nemorosum* 62*H. affine* = *Heterophyllum nemorosum* 62**H. andoi** 62*H. bambergeri* = *Campylium bambergeri* 56*H. boreale* = *Ptychostomum pallescens* 51*H. callichroum* = *Stereodon callichrous* 63**H. cupressiforme** 62var. *cupressiforme* 62var. *filiforme* 62var. *heseleri* 62var. *lacunosum* 62var. *resupinatum* = *Hypnum resupinatum* 62var. *subjulaceum* 62*H. fertile* = *Pseudohygrohypnum fertile* 63*H. hamulosum* = *Stereodon hamulosus* 63*H. holmenii* = *Stereodon holmenii* 63*H. imponens* = *Callicladium imponens* 62**H. jutlandicum** 62*H. nemorosum* = *Heterophyllum nemorosum* 62*H. pallescens* = *Jochenia pallescens* 62var. *protuberans* = *Jochenia protuberans* 62var. *reptile* = *Jochenia pallescens* 62*H. plicatulum* = *Aquilonium plicatulum* 63*H. procerrimum* = *Pseudostereodon procerrimus* 63*H. protuberans* = *Jochenia protuberans* 62*H. recurvatum* = *Drepanium fastigiatum* 57*H. reptile* = *Jochenia pallescens* 62**H. resupinatum** 62*H. revolutum* = *Roaldia revoluta* 63var. *dolomiticum* = *Roaldia dolomitica* 63*H. sauteri* = *Microhypnum sauteri* 57*H. subcomplanatum* = *Hypnum cupressiforme* var. *subjulaceum* 62*H. subimponens* = *Stereodon subimponens* 63*H. subjulaceum* = *Hypnum cupressiforme* var. *subjulaceum* 62**H. uncinulatum** 62*H. vaucheri* = *Buckia vaucheri* 63HYPOPTERYGIACEAE 2, 8, **55**HYPOPTERYGIALES = HOOKERIALES 2, **55****Hypopterygium** 2, **55****H. tamarisci** 4, **55****Imbribryum** 3, **49****I. alpinum** 49*I. gemmiparum* = *Bryum gemmiparum* 49**I. mildeanum** 49**I. miniatum** 49**I. muehlenbeckii** 49**I. subapiculatum** 5, **49****I. tenuisetum** 5, **49****Indusiella** 44**I. thianschanica** 44**Isopaches** 9**I. alboviridis** 9**I. bicrenatus** 9**I. decolorans** 9**Isopterygiopsis** 55**I. alpicola** 55**I. muelleriana** 55**I. pulchella** 55**Isopterygium** 62**I. tenerum** 62**Isothecium** 65**I. algarvicum** 65**I. alopecuroides** 65**I. holtii** 65**I. interludens** 65**I. montanum** 65**I. myosuroides** 65subsp. *brevinerve* = *Isothecium myosuroides* 65var. *brachythecioides* = *Isothecium interludens* 65**I. prolixum** 65**Iwatsukiella** 64**I. leucotricha** 64**Jaffueliobryum** 44**J. latifolium** 44*Jamesoniella* = *Syzygiella* 9*J. autumnalis* = *Syzygiella autumnalis* 9*J. rubricaulis* = *Syzygiella rubricaulis* 9*J. undulifolia* = *Biantheridium undulifolium* 9

Jamesonielloideae 9

Jochenia 62**J. pallescens** 62**J. protuberans** 62JOCHENIACEAE 8, **62****Jubula** 22**J. hutchinsiae** 22[subsp. *australiae*] 22[subsp. *bogotensis*] 22subsp. *caucasica* 22, 67subsp. *hutchinsiae* 22[subsp. *japonica*] 22[subsp. *javanica*] 22, 67[subsp. *pennsylvanica*] 22, 67JUBULACEAE 6, **22**

Jubulineae 6

Jungermannia 17, 66**J. atrovirens** 17**J. borealis** 17*J. caespiticia* = *Endogemma caespiticia* 16**J. calcicola** 17

J. callithrix = *Solenostoma callithrix* 18
J. confertissima = *Solenostoma confertissimum* 18
***J. eucordifolia* 17, 67**
 [*J. exsertifolia*] 67
 subsp. *cordifolia* = *Jungermannia eucordifolia* 17, 67
J. gracillima = *Solenostoma gracillimum* 18
 var. *crenulata* = *Solenostoma gracillimum* 18
J. handelii = *Solenostoma handelii* 18
J. hyalina = *Solenostoma hyalinum* 18
J. jenseniana = *Solenostoma sphaerocarpum* 18
J. lanceolata auct. = *Liochlaena lanceolata* 17
J. lanceolata = *Jungermannia atrovirens* 17
 var. *atrovirens* = *Jungermannia atrovirens* 17
J. leiantha = *Liochlaena lanceolata* 17
J. obovata = *Solenostoma obovatum* 18
 subsp. *minor* = *Solenostoma subellipticum* 18
J. paroica = *Solenostoma paroicum* 18
***J. polaris* 17**
***J. pumila* 17**
 subsp. *polaris* = *Jungermannia polaris* 17
 var. *alpestris* = *Jungermannia pumila* 17
J. sphaerocarpa = *Solenostoma sphaerocarpum* 18
 var. *nana* = *Solenostoma sphaerocarpum* 18
J. subelliptica = *Solenostoma subellipticum* 18
J. subulata = *Liochlaena subulata* 17
 var. *leiantha* = *Liochlaena lanceolata* 17
 JUNGERMANNIACEAE 6, 17
 Jungermanniales 6, 9
 Jungermannidae 6
 Jungermannineae 6
 Jungermannioideae 17
 Jungermanniopsida 6, 9
***Kiaeria* 35**
***K. blyttii* 35**
***K. falcata* 35**
***K. glacialis* 35**
***K. riparia* 35**
***K. starkei* 35**
***Kindbergia* 61**
***K. praelonga* 61**
***Kurzia* 19**
 [*K. makinoana*] 67
***K. pauciflora* 19**
***K. sylvatica* 19, 67**
***K. trichoclados* 19**
Leiocolea = *Mesoptychia* 17
L. alpestris = *Mesoptychia collaris* 18
L. badensis = *Mesoptychia badensis* 17
L. bantriensis = *Mesoptychia bantriensis* 18
 subsp. *wallfischii* = *Mesoptychia bantriensis* subsp.
 wallfischii 18
L. collaris = *Mesoptychia collaris* 18
L. fitzgeraldiae = *Mesoptychia fitzgeraldiae* 18
L. gillmanii = *Mesoptychia gillmanii* 18
L. heterocolpos = *Mesoptychia heterocolpos* 18
L. rutheana = *Mesoptychia rutheana* 18
 var. *laxa* = *Mesoptychia gillmanii* var. *laxa* 18
L. turbinata = *Mesoptychia turbinata* 18
***Lejeunea* 22**
L. subgenus *Lejeunea* 22
L. subgenus *Nanolejeunea* 22
***L. canariensis* 22, 67**
 [*L. capensis*] 68
***L. cavifolia* 22**
 [*L. diversifolia* = *Microlejeunea diversifolia*] 67
 [*L. diversiloba* = *Microlejeunea diversifolia*] 67

***L. eckloniana* 22**
***L. flava* 22**
 [subsp. *flava*] 22
 subsp. ***moorei* 22**
 [subsp. *orientalis*] 22
 [subsp. *tabularis*] 22
 [var. *pellucida*] 22
***L. hibernica* 22**
L. holtii = *Lejeunea eckloniana* 22
 [*L. laetevirens*] 22, 67
L. laetevirens auct. = *Lejeunea canariensis* 22
***L. lamacerina* 22**
 [subsp. *gemminata*] 22
 subsp. ***lamacerina* 22**
***L. mandonii* 22**
***L. patens* 22**
L. ulicina = *Microlejeunea ulicina* 23
 LEJEUNEACEAE 6, 8, 22
 Lejeuneae (trib.) 22
 Lejeuneinae (subtrib.) 22
 Lejeuneoideae 22
 Lembidioideae 19
 LEMBOPHYLLACEAE 8, 65
***Lepidozia* 19**
L. azorica = *Telaranea azorica* 19
***L. cupressina* 19**
 [subsp. *africana*] 19
 subsp. ***cupressina* 19**
 [subsp. *natalensis*] 19
 subsp. *pinnata* = *Lepidozia cupressina* subsp. *cupressina* 19
 var. *dissitifolia* = *Lepidozia cupressina* subsp. *cupressina* 19
***L. pearsonii* 19**
***L. reptans* 19**
 var. *julacea* = *Lepidozia reptans* 19
 var. *tenera* = *Lepidozia reptans* 19
***L. stuhlmannii* 19**
 [subsp. *pulvinata*] 19
 [var. *carnosa*] 19
 LEPIDOZIACEAE 6, 19
 Lepidozioideae 19
***Leptobarbula* 42**
***L. berica* 42**
***Leptobryum* 48**
***L. pyriforme* 48**
***Leptodictyum* 57**
 [*L. kurdicum*] 69
***L. riparium* 57**
***Leptodon* 64**
***L. corsicus* 64**
***L. longisetus* 64**
***L. smithii* 64**
***Leptodontium* 38**
***L. flexifolium* 38**
***L. gemmascens* 38**
***L. proliferum* 4, 38**
***L. styriacum* 38**
Leptophascum = *Chenia* 37
L. leptophyllum = *Chenia leptophylla* 37
 [*Leptostomopsis*] 48
***Leptothea* 55**
***L. gaudichaudii* 4, 55**
***Leptoscyphus* 20**
L. subgenus *Anomylia* 20
L. subgenus *Leptoscyphus* 20
L. section *Leptoscyphus* 20
L. azoricus = *L. porphyrius* subsp. *azoricus* 20

- L. cuneifolius** 20
L. porphyrius 20
 subsp. **azoricus** 20
 [subsp. *porphyrius*] 20
Lescurea 59
L. incurvata 59
L. mutabilis 59
L. patens 59
L. plicata 59
L. radicata 59
L. saviana 59
L. saxicola 59
L. secunda 59
Leskea 59
L. polycarpa 59
 LESKEACEAE 8, 58
 LEUCOBRYACEAE 7, 32
Leucobryum 33
L. albidum 33
L. glaucum 33
L. juniperoideum 33
Leucodon 64
 [*L. bowringii*] 69
L. canariensis 64
 [*L. coreensis*] 69
L. flagellaris 64
L. immersus 64
L. pendulus 64
L. sciuroides 64
 var. *morensis* = *Leucodon sciuroides* 64
L. treleasei 64
 LEUCODONTACEAE 8, 64
 LEUCOMIACEAE 8, 55
Lewinskya 53
L. acuminata 53
L. affinis 53
L. breviseta 53
L. elegans 53
L. fastigiata 53
L. iberica 53
L. iwatsukii 53
L. laevigata 53
L. lamyana 53
L. pylaisii 53
L. rupestris 53
L. shawii 53
L. sordida 53
L. speciosa 53
L. striata 53
L. tortidontia 53
L. transcaucasica 53
L. vladikavkana 53
Lindbergia 59
 [*L. brachyptera*] 68
L. brachyptera auct. eur. = *Lindbergia grandiretis* 59
L. dagestanica 59
L. grandiretis 59
Liochlaena 17
L. lanceolata 17
L. subulata 17
Loeskeobryum 64
L. brevirostre 64
Loeskypnum 58
L. badium 58
Lophocolea 20
L. bidentata 20
 var. *rivularis* = *Lophocolea coadunata* 20
L. bispinosa 4, 20
L. brookwoodiana 4, 20
L. coadunata 20
L. cuspidata = *L. bidentata* 20
L. cuspidata auct. = *L. coadunata* 20
L. fragrans 20
 [subsp. *cocosana*] 20
 subsp. **fragrans** 20
L. heterophylla 20
 [subsp. *cladogyna*] 20
 subsp. **heterophylla** 20
 [*L. humistrata*] 67
L. minor 20
L. semiteres 4, 20
 [var. *retusa*] 20
 var. **semiteres** 20
 LOPHOCOLEACEAE 6, 20
 Lophocoleineae 6
Lophozia 12
L. alpestris auct. = *Barbilophozia sudetica* 9
L. alpestris = *Mesoptychia collaris* 18
 var. *libertae* = *Mesoptychia collaris* 18
L. ascendens 12
L. atlantica = *Orthocaulis atlanticus* 10
L. attenuata = *Neoorthocaulis attenuatus* 9
L. badensis = *Mesoptychia badensis* 17
 var. *obtusiloba* = *Mesoptychia badensis* 17
L. bantriensis = *Mesoptychia bantriensis* 18
 var. *subcompressa* = *Mesoptychia bantriensis* subsp. *bantriensis* 18
L. barbata = *Barbilophozia barbata* 9
L. bicrenata = *Isopaches bicrenatus* 9
L. binsteadii = *Neoorthocaulis binsteadii* 10
L. capitata = *Heterogemma capitata* 12
L. cavifolia = *Orthocaulis cavifolius* 10
L. ciliata 12
L. confertifolia = *Lophozia wenzelii* 13
L. confertifolia [sensu Schljakov 1975, 1998] = *Lophozia ventricosa* 13
L. confertifolia (sensu Damsholt 2002; Köckinger 2017) = *Lophozia wenzelii* 13
L. confertifolia [sensu Konstantinova et al. 1992] = *Lophozia murmanica* 12
L. debiliformis = *Barbilophozia sudetica* 9
 var. *concolor* = *Barbilophozia sudetica* 9
L. decolorans = *Isopaches decolorans* 9
L. elongata = *Protolophozia elongata* 12
L. excisa = *Lophozia excisa* 13
 var. *cylindracea* = *Lophozia excisa* var. *excisa* 13
L. floerkei = *Neoorthocaulis floerkei* 10
L. fuscovirens 12
L. gillmanii = *Mesoptychia gillmanii* 18
 var. *acutifolia* = *Mesoptychia gillmanii* 18
L. gracillima = *Lophozia ascendens* 12
L. grandiretis = *Schistochilopsis grandiretis* 15
 var. *parviretis* = *Schistochilopsis grandiretis* 15
 var. *proteidea* = *Schistochilopsis grandiretis* 15
L. groenlandica [sensu Grolle and Long 2000; Söderström et al. 2002] = *Lophozia murmanica* 12
L. groenlandica [sensu Schuster 1969] = *Lophozia schusteriana* 13
L. groenlandica [sensu Schljakov 1980, 1998] = *Lophozia ventricosa* 13
L. groenlandica [sensu Damsholt 1994] = *Lophozia wenzelii* 13
L. guttulata 12
L. hatcheri = *Barbilophozia hatcheri* 9
L. herzogiana = *Protolophozia herzogiana* 12

- L. heterocolpos* = *Mesoptychia heterocolpos* 18
 var. *arctica* = *Mesoptychia heterocolpos* var. *arctica* 18
 var. *harpanthoides* = *Mesoptychia heterocolpos* var. *harpanthoides* 18
- L. heteromorpha* = *Lophozia murmanica* 13
L. hyperarctica = *Schistochilopsis hyperarctica* 15
L. incisa = *Schistochilopsis incisa* 15
 subsp. *opacifolia* = *Schistochilopsis opacifolia* 15
L. jurensis = *Lophoziaops jurensis* 13
L. iremelensis = *Lophozia wenzelii* 13
L. kunzeana = *Schljakovia kunzeana* 10
 [*L. lacerata*] 69
- L. lantratoviae 12**
L. latifolia = *Lophoziaops jurensis* 13
L. laxa = *Heterogemma laxa* 12
L. longidens = *Lophoziaops longidens* 13
L. longiflora auct. = *Lophozia guttulata* 12
- L. longiflora 12**
 var. *guttulata* = *Lophozia guttulata* 12
L. lycopodioides = *Barbilophozia lycopodioides* 9
- L. murmanica 12**
L. murmanica [sensu Schljakov 1970] = *Lophozia ventricosa* 13
L. murmanica [sensu Schljakov 1969] = *Lophozia savicziae* 13
L. obtusa = *Obtusifolium obtusum* 12
L. opacifolia = *Schistochilopsis opacifolia* 15
L. perssonii = *Oleolophozia perssonii* 12
L. pellucida = *Lophoziaops pellucida* 13
L. polaris = *Lophoziaops polaris* 14
L. porphyroleuca = *Lophozia guttulata* 12
L. propagulifera auct. eur. = *Lophoziaops jurensis* 13
L. quadriloba = *Schljakovianthus quadrilobus* 10
 var. *glareosa* = *Schljakovianthus quadrilobus* 10
L. rubescens = *Barbilophozia rubescens* 9
L. rubrigemma = *Lophoziaops rubrigemma* 14
L. rutheana = *Mesoptychia rutheana* 18
 var. *laxa* = *Mesoptychia gillmannii* var. *laxa* 18
- L. savicziae 13**
L. schultzii = *Mesoptychia rutheana* 5
 var. *laxa* = *Mesoptychia gillmannii* var. *laxa* 5
- L. schusteriana 13**
- L. silvicola 13**
L. silvicola var. *grandiretis* = *Lophozia savicziae* 13
- L. silvicoloides 13**
L. sudetica = *Barbilophozia sudetica* 9
 var. *anomala* = *Barbilophozia sudetica* 9
- L. subapiculata 13**
L. ventricosa auct. = *Lophozia silvicola* 13
- L. ventricosa 13**
 var. *confusa* = *Lophozia ventricosa* 13
 var. *grandiretis* = *Lophozia savicziae* 13
 var. *longiflora* = *Lophozia longiflora* 12
 var. *silvicola* = *Lophozia silvicola* 13
 var. *uliginosa* = *Lophozia wenzelii* 13
 var. *uliginosa* auct. [sensu Söderström et al. 2002] = *Lophozia longiflora* 12
- L. wenzelii 13**
 var. *groenlandica* [sensu Konstantinova et al. 2009] = *Lophozia murmanica* 13
 var. *lapponica* 13
 var. *litoralis* 13
 var. *massularioides* 13
 var. *wenzelii* 13
- LOPHOZIACEAE 6, 12
Lophoziaops 6, 13
L. excisa 13
 var. *elegans* 13
 var. *excisa* 13
 [var. *infuscata*] 13, 69
 [var. *succulenta*] 13, 69
- L. jurensis 13**
L. latifolia = *Lophoziaops jurensis* 13
- L. longidens 13**
 subsp. *arctica* 13
 subsp. *longidens* 13
- L. pellucida 13**
 var. *minor* 14
 var. *pellucida* 14
- L. perssonii* = *Oleolophozia perssonii* 12
- L. polaris 14**
 var. *polaris* 14
 var. *sphagnorum* 14
 [*L. propagulifera*] 13, 67
L. propagulifera auct. eur. = *Lophoziaops jurensis* 13
- L. rubrigemma 14**
 f. *arctica* = *Lophoziaops longidens* var. *arctica* 13
 [*Luisierella*] 32
- Lunularia 25**
L. cruciata 25
 subsp. *cruciata* 25
 [subsp. *thaxteri*] 25
- LUNULARIACEAE 7, 25
 LUNULARIALES 7, 25
- Mannia 25**
 M. subgenus *Mannia* 25
 M. subgenus *Neesiella* 25
- M. androgyna 25**
M. californica 25
M. controversa 25
 [subsp. *asiatica*] 25
 subsp. *controversa* 25
- M. fragrans 25**
 subsp. *fragrans* 25
 [subsp. *orientalis*] 25
 var. *brevipes* = *Mannia fragrans* subsp. *fragrans* 25
- M. gracilis 25**
M. pilosa 25
M. sibirica 25
M. triandra 25
- Marchantia 26**
 M. subgenus *Chlamidium* 26
 M. subgenus *Marchantia* 26
 M. subgenus *Preissia* 26
 M. section *Paleacea* 26
M. alpestris = *Marchantia polymorpha* subsp. *montivagans* 26
M. aquatica = *Marchantia polymorpha* subsp. *polymorpha* 26
 [*M. debilis*] 69
M. latifolia = *Marchantia polymorpha* subsp. *ruderalis* 26
- M. paleacea 26**
 [subsp. *diptera*] 26
 subsp. *paleacea* 26
 [*M. pappeana*] 69
- M. polymorpha 26**
 subsp. *montivagans* 26
 subsp. *polymorpha* 26
 subsp. *ruderalis* 26
- M. quadrata 26**
 subsp. *hyperborea* 26
 subsp. *quadrata* 26
- M. romanica 26**
- MARCHANTIACEAE 7, 26
 MARCHANTIALES 7, 25

- Marchantioideae 7, **26**
 Marchantiidae 7
 Marchantiophyta 2, 6, **8**
 Marchantiopsida 7, **25**
Marchesinia **23**
M. subgenus *Marchesinia* **23**
M. mackaii **23**
Marsupella **16**
M. adusta = *Gymnomitrium adustum* 16
M. alpina = *Gymnomitrium alpinum* 16
M. andreaeoides **16**
M. aquatica **16**
M. arctica **16**
M. boeckii **16**
M. brevissima = *Gymnomitrium brevissimum* 16
M. commutata = *Gymnomitrium commutatum* 16
M. condensata **16**
M. emarginata **16**
 subsp. *aquatica* = *Marsupella aquatica* 16
 subsp. *tubulosa* = *Marsupella tubulosa* 17
 var. *aquatica* = *Marsupella aquatica* 16
 var. *pearsonii* = *Marsupella emarginata* 16
M. funckii **17**
 var. *badensis* = *Marsupella funckii* 17
M. profunda **17**
M. ramosa = *Marsupella funckii* 17
M. sparsifolia **17**
 [subsp. *childii*] 17
 subsp. *sparsifolia* **17**
M. sphacelata **17**
M. spiniloba **17**
M. sprucei **17**
 var. *neglecta* = *Marsupella sprucei* 17
 var. *ustulata* = *Marsupella sprucei* 17
M. stableri **17**
M. subemarginata **17**
M. tubulosa **17**
Massularia = *Heterogemma* 12
M. incisa = *Schistochilopsis incisa* 15
M. laxa = *Heterogemma laxa* 12
Mastigophora **20**
M. woodsii **20**
 MASTIGOPHORACEAE 6, **20**
Meesia **48**
M. hexasticha **48**
M. longiseta **48**
M. minor **116**
M. minutissima **116**
M. triquetra **48**
M. uliginosa **48**
 MEESIACEAE 7, **48**
 Merceyoideae **36**
Mesoptychia **17**
M. badensis **17**
M. bantriensis **17**
 subsp. *bantriensis* **18**
 subsp. *wallfischii* **18**
M. collaris **18**
M. fitzgeraldiae **18**
M. gillmanii **18**
 var. *gillmanii* **18**
 var. *laxa* 5, **18**
M. heterocolpos **18**
 var. *arctica* **18**
 var. *harpanthoides* **18**
 var. *heterocolpos* **18**
M. rutheana **18**
M. sahlbergii **18**
M. turbinata **18**
 Mesoptychioideae **17**
Metzgeria **24**
M. conjugata **24**
 subsp. *simplex* = *Metzgeria simplex* 24
 var. *alipila* = *Metzgeria conjugata* 24
 var. *macvicari* = *Metzgeria conjugata* 24
M. consanguinea **24**, **67**
M. fruticulosa auct. = *Metzgeria violacea* 24
M. fruticulosa = *Riccardia palmata* 24
M. furcata **24**
 var. *expansa* = *Metzgeria furcata* 24
 var. *flexipilis* = *Metzgeria furcata* 24
 [var. *pacifica*] 24
 var. *ulvula* = *Metzgeria furcata* 24
 [M. *hamata* = *Metzgeria procerata*] 24
M. hamata auct. = *Metzgeria leptoneura* 24
M. leptoneura **24**
 [var. *brevisetata*] 24
 var. **leptoneura** **24**
 [var. *polychaeta*] 24
 [M. *procerata*] 24
M. pubescens **24**
 [M. *setigera*] 24
M. simplex **24**
M. temperata auct. = *Metzgeria consanguinea* 24
 [M. *temperata*] 24, **67**
M. violacea **24**
 METZGERIACEAE 6, **24**
 METZGERIALES 6, **23**
 Metzgeriidae 6
Microbryum **38**
M. curvicolllum **38**
M. davallianum **38**
 var. *commutatum* **38**
 var. *conicum* **38**
 var. *davallianum* **38**
M. floerkeanum **38**
M. fosbergii **38**
M. longipes **38**
M. rectum **38**
M. starckeanum **38**
M. vlassovii **38**
Microcampylopus **33**
M. laevigatus **33**
Microeurhynchium **60**
M. pumilum **60**
Microhypnum **57**
M. sauteri **57**
Microlejeunea **23**
 [M. *diversiloba*] 67
M. ulicina **23**
 MICROMITRIACEAE 7, **32**
Micromitrium **32**
M. tenerum **32**
Mielichhoferia **51**
M. elongata **51**
M. mielichhoferiana **51**
 Mielichhoferioideae **51**
 MNIACEAE 5, 8, **51**
 Mnioidae 5, **52**
Mnioloma **16**
M. subgenus *Caracoma* **16**
M. fuscum **16**
Mnium 5, 6, **52**
M. blyttii **52**

M. heterophyllum 52

M. latifolium = *Ptychostomum schleicheri* var. *latifolium* 6

M. hornum 52

M. latifolium = *Ptychostomum schleicheri* var. *latifolium* 6

M. lycopodioides 52**M. marginatum** 52

var. *dioicum* 52

var. *marginatum* 52

M. spinosum 52**M. spinulosum** 52**M. stellare** 52**M. thomsonii** 52**Moerckia** 24**M. blyttii** 24**M. flotoviana** 24**M. hibernica** 24

MOERCKIACEAE 7, 24

Molendoa 42**M. hornsuschiana** 42**M. schliephackei** 42

[*M. seravschanica*] 69

M. sendtneriana = *Molendoa hornsuschiana* 42

M. taeniatifolia 42

M. tenuinervis = *Molendoa hornsuschiana* 42

M. warburgii 42**Mylia** 21

M. subgenus *Anomalae* 21

M. subgenus *Mylia* 21

M. anomala 21**M. taylorii** 21

MYLIACEAE 6, 21

Myliineae 6

Myrinia 56**M. pulvinata** 56

MYRINIACEAE 8, 56

Myriocoleopsis 22**M. minutissima** 22

subsp. *minutissima* 22

[subsp. *myriocarpa*] 22

Myurella 56**M. julacea** 56**M. sibirica** 56**M. tenerrima** 56**Myurium** 65**M. hochstetteri** 65

MYURIACEAE 8, 65

Myuroclada 61**M. longiramea** 61**M. maximowiczii** 4, 61**Nardia** 17**N. breidleri** 17**N. compressa** 17**N. geoscyphus** 17

var. *bifida* 17

[var. *dioica*] 17

var. *geoscyphus* 17

var. *suberecta* 17

N. insecta 17**N. japonica** 17**N. pacifica** 17**N. scalaris** 17

[var. *botryoidea*] 17

[var. *harae*] 17

var. *scalaris* 17

Nardioideae 17

Neckera 64

N. baetica = *Exsertotheca baetica* 64

N. besseri = *Alleniella besseri* 64

N. cephalonica 64

N. complanata = *Alleniella complanata* 64

N. crispa = *Exsertotheca crispa* 64

N. intermedia = *Exsertotheca intermedia* 64

N. menziesii 64**N. oligocarpa** 64**N. pennata** 64**N. pumila** 64

N. webbiana = *Pseudomalina webbiana* 65

NECKERACEAE 2, 8, 64, 65

Neoorthocaulis 9**N. attenuatus** 9**N. binsteadii** 10**N. floerkei** 10**N. hyperboreus** 10

Niphotrichum = *Racomitrium* subgenus *Niphotrichum* 44

N. canescens = *Racomitrium canescens* 44

subsp. *latifolium* = *Racomitrium canescens* subsp. *latifolium* 44

N. elongatum = *Racomitrium elongatum* 44

N. ericoides = *Racomitrium ericoides* 44

N. panschii = *Racomitrium panschii* 44

Nobregaea 60**N. latinervis** 60**Nogopterium** 64**N. gracile** 64

NOTOTHYLADACEAE 6, 8

NOTOTHYLADALES 6, 8

Notothylatidae 6

Notothyas 8

N. subgenus *Notothyas* 8

N. orbicularis 8

Notothyloideae 8

Nowellia 11

N. section *Nowellia* 11

N. curvifolia 11**Nyholmiella** 53**N. gymnostoma** 54**N. obtusifolia** 54

Obtusifolium 11, 12

O. obtusum 12

Ochyraea = *Plathyhypnum* 6, 57

O. tatrensis = *Plathyhypnum tatrense* 6, 58

Octodiceras = *Fissidens* subgenus *Octodiceras* 34

O. fontanum = *Fissidens fontanus* 34

Odontoschisma 11

O. section *Cladopodiella* 11

O. section *Denudata* 11

O. section *Neesii* 11

O. section *Odontoschisma* 11

O. denudatum 11

subsp. *denudatum* 11

[subsp. *naviculare*] 11

[subsp. *sandvicense*] 11

O. elongatum 11**O. fluitans** 11**O. francisci** 11**O. macounii** 11

O. prostratum = *Odontoschisma sphagni* 11

O. sphagni 11

Odontoschismatoideae 11

OEDIPODIACEAE 7, 29

OEDIPODIALES 7, 29

Oedipodiella 32**O. australis** 32

Oedipodiopsida 7, 29

Oedipodium 29

O. griffithianum 29

Oleolophozia 12

O. perssonii 12

Oligotrichum 29

O. hercynicum 29

ONCOPHORACEAE = RHABDOWEISIACEAE 35

Oncophorus 35

O. compactus = *Oncophorus wahlenbergii* 35

O. demetrii 35

O. dendrophilus 35

O. elongatus 35

O. integerrimus 35

O. riparius = *Kiaeria riparia* 35

O. virens 35

var. *elongatus* = *Oncophorus integerrimus* 35

O. wahlenbergii 35

var. *compactus* = *Oncophorus wahlenbergii* 35

Oreas 35

O. martiana 35

Oreoweisia 35

O. torquescens 35

Orthocaulis 10

O. atlanticus 10

O. attenuatus = *Neoorthocaulis attenuatus* 9

O. binsteadii = *Neoorthocaulis binsteadii* 10

O. cavifolius 10

O. floerkei = *Neoorthocaulis floerkei* 10

O. hyperboreus = *Neoorthocaulis hyperboreus* 10

Orthodicranum = *Dicranum* 35

O. flagellare = *Dicranum flagellare* 35

O. fulvum = *Dicranum fulvum* 35

O. montanum = *Dicranum montanum* 35

O. scottianum = *Dicranum scottianum* 35

O. tauricum = *Dicranum tauricum* 35

ORTHODONTIACEAE 8, 55

ORTHODONTIALES 8, 55

Orthodontium 55

O. gracile 55

O. lineare 4, 55

O. pellucens 55

Orthogrimmia = *Grimmia* 45

O. alpestris = *Grimmia alpestris* 45

O. arenaria = *Grimmia arenaria* 45

O. caespiticia = *Grimmia caespiticia* 45

O. donniana = *Grimmia donniana* 45

O. montana = *Grimmia montana* 45

O. sessitana = *Grimmia reflexidens* 45

O. triformis = *Grimmia triformis* 45

Ortholimnobia 56

O. handelii 56

Orthomnion = *Plagiomnium* 52

O. affine = *Plagiomnium affine* 52

O. confertidens = *Plagiomnium confertidens* 52

O. curvatulum = *Plagiomnium curvatulum* 52

O. cuspidatum = *Plagiomnium cuspidatum* 52

O. drummondii = *Plagiomnium drummondii* 52

O. elatum = *Plagiomnium elatum* 52

O. ellipticum = *Plagiomnium ellipticum* 52

O. medium = *Plagiomnium medium* 52

O. undulatum = *Plagiomnium undulatum* 53

var. *madeirense* = *Plagiomnium undulatum* var. *madeirensis* 53

O. rostratum = *Plagiomnium rostratum* 53

Orthothecium 56

O. chryseon 56

O. intricatum 56

O. lapponicum 56

O. rufescens 56

O. strictum 56

ORTHOTRICHACEAE 2, 8, 53

ORTHOTRICHIALES 8, 53

Orthotrichoideae 53

Orthotrichum 54

O. acuminatum = *Lewinskya acuminata* 53

O. affine = *Lewinskya affine* 53

var. *bohemicum* = *Lewinskya affine* 53

O. alpestre 54

O. anomalum 54

O. bistratosum 54

O. callistomum 54

O. cambrense 54

O. casasianum 54

O. columbicum 54

O. comosum 54

[*O. consimile*] 54

O. consimile auct. eur. = *Orthotrichum columbicum* 54

O. consobrinum 54

O. crenulatum 54

O. cupulatum 54

var. *bistratosum* = *Orthotrichum bistratosum* 54

var. *cupulatum* 54

var. *fuscum* 54

var. *riparium* 54

O. dagestanicum 54

O. dentatum 54

O. diaphanum 54

O. elegans = *Lewinskya elegans* 53

O. fastigiatum = *Lewinskya fastigiata* 53

O. gymnostomum = *Nyholmiella gymnostoma* 54

O. handiense 54

O. hispanicum 54

O. holmenii = *Orthotrichum sibiricum* 54

O. ibericum = *Lewinskya iberica* 53

O. iwatsukii = *Lewinskya iwatsukii* 53

O. laevigatum = *Lewinskya laevigata* 53

var. *japonicum* = *Lewinskya iwatsukii* 53

O. lyellii = *Pulvigerella lyellii* 54

O. macounii 53

subsp. *japonicum* = *Lewinskya iwatsukii* 53

O. macrocephalum 54

O. microcarpum 54

O. moravicum 54

O. obtusifolium = *Nyholmiella obtusifolia* 54

O. pallens 54

O. patens 54

O. pellucidum 54

O. philibertii 54

O. pulchellum 54

O. pumilum 54

O. pylaisii = *Lewinskya pylaisii* 53

O. rivulare 54

O. rogeri 54

O. rupestre = *Lewinskya rupestris* 53

O. scanicum 54

O. schimperii 54

O. shawii = *Lewinskya shawii* 53

O. shevockii 54

O. sibiricum 54

O. sordidum = *Lewinskya sordida* 53

O. speciosum = *Lewinskya speciosa* 53

var. *brevisetum* = *Lewinskya breviseta* 53

O. sprucei 54

O. stellatum 54**O. stramineum 54***O. striatum* = *Lewinskya striata* 53**O. tenellum 54***O. tortidontium* = *Lewinskya tortidontia* 53[*O. urnaceum*] 69**O. urnigerum 54****O. vittii 54***O. vladikavkana* = *Lewinskya vladikavkana* 53*Osculatia* = *Bryum* 3, 48*O. apiculata* = *Anomobryum apiculatum* 48*O. barnesii* = *Bryum dichotomum* 48*O. bicolor* = *Bryum dichotomum* 48*O. bornholmensis* = *Ptychostomum bornholmense* 50*O. caespiticia* = *Ptychostomum imbricatum* 50*O. demaretiana* = *Bryum demaretianum* 48*O. dichotoma* = *Bryum dichotomum* 48*O. gemmifera* = *Bryum gemmiferum* 49*O. gemmilucens* = *Bryum gemmilucens* 49*O. klinggraeffii* = *Bryum klinggraeffii* 49*O. kunzei* = *Ptychostomum kunzei* 51*O. radiculosa* = *Bryum radiculosum* 49*O. rubens* = *Ptychostomum rubens* 51*O. ruderalis* = *Bryum ruderale* 49*O. sauteri* = *Bryum sauteri* 49*O. subapiculata* = *Imbriobryum subapiculatum* 49*O. tenuisetia* = *Imbriobryum tenuisetum* 49*O. valparaisense* = *Bryum valparaisense* 49*O. violacea* = *Bryum violaceum* 49**Oxymitra 26****O. incrassata 26**

OXYMITRACEAE 7, 26

Oxyrrhynchium 60**O. hians 60***O. pumilum* = *Microeurhynchium pumilum* 60**O. schleicheri 60****O. speciosum 60***Oxystegus* = *Chionoloma* 40*O. daldinianus* = *Chionoloma daldinianum* 40*O. hibernicus* = *Chionoloma hibernicum* 40*O. minor* = *Chionoloma minus* 41*O. recurvifolius* = *Chionoloma recurvifolium* 41*O. tenuirostris* = *Chionoloma tenuirostre* 41var. *holtii* = *Chionoloma tenuirostre* var. *holtii* 41**Palamocladium 59****P. euchloron 59**[*P. leskeoides*] 69**Pallavicinia 24****P. lyellii 24**

PALLAVICINIACEAE 7, 24

PALLAVICINIALES 7, 24

Pallaviciniineae 7

Pallavicinioideae 24

Paludella 48**P. squarrosa 48****Palustriella 56****P. commutata 56**var. *sulcata* = *Palustriella falcata* 56**P. decipiens 56****P. falcata 56***P. pluristratosa* = *Palustriella falcata* 56*Paraleptodontium recurvifolium* = *Chionoloma**recurvifolium* 41**Paraleucobryum 35****P. enerve 35****P. longifolium 35**var. *sauteri* = *Paraleucobryum sauteri* 35**P. sauteri 35****Pedinophyllum 20****P. interruptum 20****Pelekiium 59****P. atlanticum 59****P. minutulum 59****Pellia 25***P. endiviifolia* = *Apopellia endiviifolia* 24**P. epiphylla 25**subsp. *borealis* 25subsp. *epiphylla* 25**P. neesiana 25**

PELLIACEAE 7, 24

PELLIALES 7, 24

Pelliidae 6

Peltolepis 25**P. quadrata 25**[*Perssonia sanguinea*] 69

PETALOPHYLLACEAE 7, 24

Petalophyllum 24**P. ralfsii 24****Phaeoceros 8***P. bulbiculosus* = *Phymatoceros bulbiculosus* 8**P. carolinianus 8****P. laevis 8**subsp. *carolinianus* = *Phaeoceros carolinianus* 8

Phaeocerotidae 6, 8

Phascum = *Tortula* 39*P. cuspidatum* = *Tortula acaulon* 39var. *cuspidatum* = *Tortula acaulon* var. *acaulon* 39var. *papillosum* = *Tortula acaulon* var. *papillosa* 39var. *piliferum* = *Tortula acaulon* var. *pilifera* 39var. *retortifolium* = *Tortula acaulon* var. *retortifolia* 39var. *schreberianum* = *Tortula acaulon* var.*schreberiana* 39*P. vlassovii* = *Microbryum vlassovii* 38**Philonotis 47***P.* section *Bartramidula* 47*P.* section *Homomorphae* 47*P.* section *Philonotis* 47*P.* section *Philonotula* 47*P. arnellii* = *Philonotis capillaris* 47**P. caespitosa 47****P. calcarea 47****P. capillaris 47****P. cernua 47****P. falcata 47****P. fontana 47****P. hastata 47**[*P. laxitexta*] 69**P. marchica 47**[*P. nanothecioidea*] 69**P. rigida 47****P. seriata 47****P. tomentella 47****P. uncinata 47****P. yezoana 47****Phymatoceros 8****P. bulbiculosus 8**

PHYMATOCEROTACEAE 6, 8

PHYMATOCERATALES 6, 8

Physcomitrella = *Physcomitrium* 31*P. patens* = *Physcomitrium patens* 31*P. readeri* = *Physcomitrium readeri* 32*Physcomitridium* = *Physcomitrium* 31, 32*P. readeri* = *Physcomitrium readeri* 32**Physcomitrium 31**

- P. arenicola** 31
P. eurystomum 31
 subsp. **acuminatum** 31
 subsp. **eurystomum** 31
 [*P. immersum*] 69
 [*P. niloticum*] 69
P. patens 31
P. pyriforme 31
P. readeri 32
P. sphaericum 32
P. × stevensonii 116
 PILOTRICHACEAE 8, 55
Plagiobryoides = *Ptychostomum* 3, 50
P. cellularis = *Ptychostomum cellulare* 50
Plagiobryum = *Ptychostomum* 3, 50
P. demissum = *Ptychostomum demissum* 50
P. zieri = *Ptychostomum zieri* 51
Plagiochasma 25
P. subgenus *Micropylum* 25
P. subgenus *Plagiochasma* 25
P. appendiculatum 25
 [*P. eximium*] 69
 [*P. microcephalum* var. *tunesicum*] 69
P. rupestre 25
 var. **rupestre** 25
 [var. *volkii*] 25
Plagiochila 20
P. section *Arrectae* 20
P. section *Fuscoluteae* 20
P. section *Glauscentes* 20
P. section *Plagiochila* 20
P. section *Poeltiae* 21
P. section *Rutilantes* 21
P. section *Vagae* 21
 [*P. ambagiosa* = *Plagiochila spinulosa*] 67
P. arctica 20
 subsp. *arctica* = *Plagiochila arctica* 20
 var. **arctica** 20
 [var. *intermedia*] 20
P. asplenioides 21
 subsp. *arctica* = *Plagiochila arctica* 20
 subsp. *porelloides* = *Plagiochila porelloides* 21
 var. *devexa* = *Plagiochila porelloides* var. *porelloides* 21
 var. *humilis* = *Plagiochila porelloides* var. *porelloides* 21
 var. *lobata* = *Plagiochila asplenioides* var. *arctica* 20
 var. *minor* = *Plagiochila porelloides* var. *porelloides* 21
 var. *subarctica* = *Plagiochila porelloides* var. *subarctica* 21
P. atlantica = *Plagiochila heterophylla* var. *heterophylla* 20
P. bifaria 20
 var. **bifaria** 20
 [var. *rosea*] 20
P. britannica 21
P. carringtonii 21
 subsp. **carringtonii** 21
 [subsp. *lobuchensis*] 21
P. dubia auct. eur. = *P. virginica* 21
 [*P. dubia* = *Plagiochila patula*] 67
P. exigua 21
P. heterophylla 20
 [var. *beauverdii*] 20
 var. **heterophylla** 20
 [*P. javanica*] 67
P. longispina 20
P. maderensis 21
P. major = *Plagiochila asplenioides* 21
P. norvegica = *Plagiochila porelloides* var. *norvegica* 21
P. papillifolia 20
 [*P. patula*] 21, 67
P. porelloides 21
 var. **norvegica** 21
 var. **porelloides** 21
 var. **subarctica** 21
P. punctata 20
P. retrorsa 20
P. spinulosa 20
P. stricta 20
 [*P. uniformis*] 67
P. virginica 21, 67
 [var. *caroliniana*] 21
 [var. *euryphylla*] 21
 var. **virginica** 21
 PLAGIOCHILACEAE 6, 20
Plagiomnium 52
P. section *Plagiomnium* 52
P. section *Rostrata* 53
P. section *Undulata* 52
P. affine 52
P. confertidens 52
P. curvatulum 52
P. cuspidatum 52
P. drummondii 52
P. elatum 52
P. ellipticum 52
P. medium 52
P. rostratum 53
P. undulatum 53
 var. **madeirense** 53
 var. **undulatum** 53
Plagiopus 47
P. oederianus 47
 var. **alpinus** 47
 var. **oederianus** 47
 PLAGIOTHECIACEAE 8, 55
Plagiothecium 55
P. berggrenianum 55
P. cavifolium 55
P. curvifolium 55
P. denticulatum 55
 var. **denticulatum** 55
 var. **obtusifolium** 55
 var. **undulatum** 55
P. handelii = *Ortholimnobium handelii* 56
P. laetum 55
P. latebricola 55
P. neckeroideum 56
P. nemorale 56
P. noricum = *Plagiothecium neckeroideum* 56
P. piliferum 56
P. platyphyllum 56
P. rossicum 56
P. succulentum 56
P. svalbardense 56
P. undulatum 56
Plasteurhynchium 59
P. meridionale 59
P. striatulum 59
Platydictya 56
P. jungermannioides 56
 [*Platygyrella densa*] 69
Platygyrium 62
P. repens 62
Platyhypnidium = *Rhynchostegium* 59
P. grolleanum = *Rhynchostegium riparioides* 60
P. lusitanicum = *Rhynchostegium alopecuroides* 59

- P. mutatum* = *Rhynchostegium alopecuroides* 59
P. torrenticola = *Rhynchostegium riparioides* 60
Platyhypnum 6, 57
P. alpestre 57
P. alpinum 57
P. cochlearifolium 58
P. duriusculum 58
P. molle 58
P. montanum = *Campylophyllum montanum* 57
P. norvegicum 58
P. smithii 58
P. tatrense 6, 58
[*Plaubelia sprengelii*] 70
Plectocolea = *Solenostoma* 18
P. hyalina = *Solenostoma hyalinum* 18
P. obovata = *Solenostoma obovatum* 18
P. subelliptica = *Solenostoma subellipticum* 18
Plenogemma 54
P. phyllantha 54
Pleuridium 36
P. acuminatum 36
P. palustre = *Cleistocarpidium palustre* 36
P. subulatum 36
Pleurochaete = *Tortella* 42
P. squarrosa = *Tortella squarrosa* 42
Pleurocladula = *Fuscocephaloziopsis* 10
P. affinis = *Fuscocephaloziopsis affinis* 10
P. albescens = *Fuscocephaloziopsis albescens* 10
var. *islandica* = *Fuscocephaloziopsis albescens* var. *islandica* 10
P. catenulata = *Fuscocephaloziopsis catenulata* 10
P. connivens = *Fuscocephaloziopsis connivens* 10
P. leucantha = *Fuscocephaloziopsis leucantha* 11
P. loitlesbergeri = *Fuscocephaloziopsis loitlesbergeri* 11
P. lunulifolia = *Fuscocephaloziopsis lunulifolia* 11
P. macrostachya = *Fuscocephaloziopsis macrostachya* 11
P. pleniceps = *Fuscocephaloziopsis pleniceps* 11
Pleurozia 24
P. subgenus *Constantifolia* 24
[*P. gigantea*] 69
P. purpurea 24
PLEUROZIAEAE 6, 24
PLEUROZIALES 6, 24
Pleurozium 64
P. schreberi 64
Pogonatum 29
P. aloides 29
P. dentatum 29
P. nanum 29
P. neesii 29
P. urnigerum 29
Pohlia 52
[*P. alba*] 70
P. andalusica 52
P. andrewsii 52
P. annotina 52
P. atropurpurea 52
P. beringiensis 52
P. bolanderi 52
P. bulbifera 52
P. camptotrachela 52
P. cruda 52
P. crudoides 52
P. drummondii 52
P. elongata 52
var. *acuminata* 52
var. *elongata* 52
var. *greenii* 52
P. erecta 52
P. filum 52
P. flexuosa 52
var. *flexuosa* 52
var. *pseudomuyldermansii* 52
P. inclinata = *Ptychostomum inclinatum* 50
P. lescuriana 52
P. longicolla 52, 68
P. ludwigii 52
P. lutescens 52
P. melanodon 52
P. nutans 52
subsp. *nutans* 52
subsp. *schimperii* 52
P. obtusifolia 52
P. prolifera 52
[*P. philonotula*] 68
[*P. saprophila*] 68, 70
P. scotica 52
P. sphagnicola 52
P. tundrae 52
P. vexans 52
P. wahlenbergii 52
var. *calcareae* 52
var. *glacialis* 52
var. *wahlenbergii* 52
POLYTRICHACEAE 2, 7, 29
POLYTRICHALES 7, 29
Polytrichastrum 29
P. alpinum 29
var. *fragile* = *Polytrichastrum fragile* 30
var. *septentrionale* = *Polytrichastrum septentrionale* 30
P. altaicum 29
P. formosum = *Polytrichum formosum* 30
var. *densifolium* = *Polytrichum densifolium* 30
P. fragile 30
P. longisetum = *Polytrichum longisetum* 30
P. norvegicum = *Polytrichastrum alpinum* 29
P. pallidisetum = *Polytrichum pallidisetum* 30
P. septentrionale 30
P. sexangulare 30
var. *vulcanicum* = *Polytrichastrum sphaerothecium* 30
P. sphaerothecium 30
Polytrichopsida 7, 29
Polytrichum 30
P. alpestre = *Polytrichum strictum* 30
P. alpinum = *Polytrichastrum alpinum* 29
var. *arcticum* = *Polytrichastrum alpinum* 29
[*P. appressum* = *Polytrichum subpilosum*] 30
P. commune 30
var. *commune* = *Polytrichum commune* 30
var. *humile* = *Polytrichum perigoniale* 30
var. *perigoniale* = *Polytrichum perigoniale* 30
var. *swartzii* = *Polytrichum swartzii* 30
var. *uliginosum* = *Polytrichum commune* 30
P. densifolium 30
P. formosum 30
var. *densifolium* = *Polytrichum densifolium* 30
P. fragile = *Polytrichastrum fragile* 30
P. hyperboreum 30
P. jensenii 30
P. juniperinum 30
P. longisetum 30
P. pallidisetum 30
P. perigoniale 30
P. piliferum 30

- P. septentrionale* = *Polytrichastrum septentrionale* 30
P. sexangulare = *Polytrichastrum sexangulare* 30
P. sphaerothecium = *Polytrichastrum sphaerothecium* 30
***P. strictum* 30**
 [*P. subpilosum*] 20
***P. swartzii* 30**
P. uliginosum = *Polytrichum commune* 30
***Porella* 23**
***P. arboris-vitae* 23**
 subsp. ***arboris-vitae* 23**
 [subsp. *nitidula*] 23
 var. *killarniensis* = *Porella arboris-vitae* subsp.
arboris-vitae 23
 var. *obscura* = *Porella arboris-vitae* subsp. *arboris-vitae* 23
***P. baueri* 23**
***P. canariensis* 23**
***P. cordaeana* 23**
 var. *faeroensis* = *Porella cordaeana* 23
 var. *simplicior* = *Porella cordaeana* 23
***P. inaequalis* 23**
 [*P. navicularis*] 67
***P. obtusata* 23**
***P. pinnata* 23**
***P. platyphylla* 23, 67**
 var. *subsquarrosa* = *Porella platyphylla* 23
 [*P. platyphylloidea*] 67
 PORELLACEAE 6, **23**
 PORELLALES 6, **21**
 Porellineae 6
Pottia = *Tortula* 39
P. commutata = *Microbryum davallianum* var.
commutatum 38
P. conica = *Microbryum davallianum* var. *conicum* 38
P. crinita = *Tortula viridifolia* 40
P. intermedia = *Tortula caucasica* 39
P. lanceolata = *Tortula lindbergii* 40
P. starckeana = *Microbryum starckeanum* 38
 subsp. *conica* = *Microbryum davallianum* var. *conicum* 38
 POTTIACEAE 2, 7, **36**
 Pottioideae **36**
***Pottiopsis* 42**
***P. caespitosa* 42**
***Prasanthus* 17**
 [*P. jamalicus*] 69
***P. suecicus* 17**
Preissia = *Marchantia* subgenus *Preissia* 26
P. quadrata = *Marchantia quadrata* 26
 subsp. *hyperborea* = *Marchantia quadrata* subsp.
hyperborea 26
Protobryum = *Tortula* 39
Protobryum bryoides = *Tortula protobryoides* 40
***Protolophozia* 12**
***P. elongata* 12**
***P. herzogiana* 12**
***Pseudanomodon* 64**
***P. attenuatus* 64**
***Pseudephemerum* 36**
***P. nitidum* 36**
***Pseudoamblystegium* 58**
***P. subtile* 58**
***Pseudobryum* 53**
***P. cinclidioides* 53**
Pseudocalliergon = *Drepanocladus* 57
P. angustifolium = *Drepanocladus angustifolius* 57
P. brevifolium = *Drepanocladus brevifolius* 57
P. lycopodioides = *Drepanocladus lycopodioides* 57
P. trifarium = *Drepanocladus trifarius* 57
P. turgescens = *Drepanocladus turgescens* 57
***Pseudocampyllum* 58**
***P. radicale* 58**
***Pseudocrossidium* 38**
***P. hornschurchianum* 38**
***P. obtusulum* 38**
***P. replicatum* 38**
***P. revolutum* 38**
***Pseudohygrohypnum* 63**
***P. eugyrium* 63**
***P. fertile* 63**
***P. subeugyrium* 63**
Pseudoleskea = *Lescuraea* 59
P. artariae = *Pseudoleskeopsis artariae* 59
P. incurvata = *Lescuraea incurvata* 59
P. patens = *Lescuraea patens* 59
P. radicata = *Lescuraea radicata* 59
P. saviana = *Lescuraea saviana* 59
 PSEUDOLESKEACEAE 8, **59**
***Pseudoleskeella* 59**
***P. catenulata* 59**
***P. nervosa* 59**
***P. papillosa* 59**
***P. rupestris* 59**
***P. tectorum* 59**
 PSEUDOLESKEELLACEAE 8, **59**
***Pseudoleskeopsis* 59**
***P. artariae* 59**
 [*P. bollei*] 70
 [*P. pseudoattenuata*] 70
Pseudolophozia = *Barbilophozia* subgenus *Sudeticae* 9
P. debiliformis = *Barbilophozia sudetica* 9
P. sudetica = *Barbilophozia sudetica* 9
***Pseudomalina* 65**
***P. webbiana* 65**
***Pseudomarsupidium* 9**
***P. decipiens* 9**
***Pseudorhynchostegiella* 60**
***P. duriaei* 60**
***Pseudoscleropodium* 59**
***P. purum* 59**
***Pseudostereodon* 63**
***P. procerrimus* 63**
 [*Pseudosymbplepharis*] 40
P. bombayensis = *Chionoloma bombayense* 40
***Pseudotaxiphyllum* 56**
***P. elegans* 56**
***P. laetevirens* 56**
***Pseudotritomaria* 14**
***P. heterophylla* 14**
***Psilopilum* 30**
***P. cavifolium* 30**
***P. laevigatum* 30**
 PTERIGYNANDRACEAE 8, **56**
***Pterigynandrum* 56**
***P. filiforme* 56**
 var. *majus* = *Pterigynandrum filiforme* 56
Pterogonium = *Nogopterium* 64
P. gracile = *Nogopterium gracile* 64
***Pterygoneurum* 39**
P. compactum = *Pterygoneurum ovatum* 39
P. crossidioides = *Pterygoneurum ovatum* 39
***P. kozlovii* 39**
***P. lamellatum* 39**
***P. ovatum* 39**
***P. papillosum* 39**
***P. sampaianum* 39**

- P. squamosum* = *Pterygoneurum ovatum* 39
***P. subsessile* 39**
PTILIDIACEAE 6, 23
PTILIDIALES 6, 23
***Ptilidium* 23**
***P. ciliare* 23**
***P. pulcherrimum* 23**
var. *subpinnatum* = *Ptilidium pulcherrimum* 23
***Ptilium* 63**
***P. crista-castrensis* 63**
Ptychanthoideae 23
Ptychodium = *Lescurea* 59
P. plicatum = *Lescurea plicata* 59
PTYCHOMITRIACEAE 7, 43
***Ptychomitrium* 44**
***P. incurvum* 44**
***P. nigrescens* 44**
***P. polyphyllum* 44**
***Ptychostomum* 3, 6, 50**
P. archangelicum = *Ptychostomum inclinatum* 50
***P. arcticum* 50**
***P. austriacum* 6, 50**
P. boreale = *Ptychostomum pallescens* 51
***P. bornholmense* 50**
***P. calophyllum* 50**
***P. capillare* 50**
***P. cellulare* 6, 50**
***P. cernuum* 50**
***P. compactum* 50**
var. *compactum* 50
var. *rutheanum* 50
***P. creberrimum* 50**
***P. cryophilum* 50**
***P. cyclophyllum* 50**
***P. demissum* 50**
***P. donianum* 50**
***P. elegans* 6, 50**
***P. funkii* 50**
***P. imbricatum* 3, 50**
P. imbricatum = *Ptychostomum inclinatum* 50
***P. inclinatum* 50**
***P. intermedium* 50**
***P. knowltonii* 50**
***P. kunzei* 50**
***P. longisetum* 51**
***P. minii* 6, 51**
***P. moravicum* 51**
P. neodamense = *Ptychostomum pseudotriquetrum* 51
***P. pallens* 51**
***P. pallescens* 51**
***P. pseudotriquetrum* 3, 51**
var. *bimum* 51
var. *pseudotriquetrum* 51
***P. rubens* 51**
***P. salinum* 51**
***P. schleicheri* 6, 51**
var. *latifolium* 6, 51
var. *schleicheri* 51
***P. torquescens* 51**
***P. turbinatum* 51**
***P. warneum* 51**
***P. weigeli* 51**
***P. wrightii* 51**
***P. zieri* 51**
***Pulviger* 54**
***P. lyellii* 54**
***Pylaisia* 63**
- P. polyantha* 63**
***P. selwynii* 63**
PYLAIACEAE 8, 63
PYLAIADDELPHACEAE 8, 62
***Pyramidula* 31**
***P. tetragona* 31**
Pyramiduloideae 31
Racomitrioideae 44
***Racomitrium* 44**
R. subgenus *Cataracta* 44
R. subgenus *Ellipticodryptodon* 44
R. subgenus *Niphotrichum* 44
R. subgenus *Racomitrium* 44
***R. aciculare* 44**
***R. affine* 44**
***R. aquaticum* 44**
***R. canescens* 44**
subsp. *canescens* 44
subsp. *latifolium* 44
***R. ellipticum* 44**
***R. elongatum* 44**
***R. ericoides* 44**
***R. fasciculare* 44**
***R. hespericum* 44**
***R. heterostichum* 44**
***R. himalayanum* 44**
***R. lamprocarpum* 44**
***R. lanuginosum* 44**
***R. lusitanicum* 44**
***R. macounii* 44**
subsp. *alpinum* 44
subsp. *macounii* 44
***R. microcarpon* 44**
***R. nivale* 44**
***R. obtusum* 44**
***R. panschii* 44**
***R. sudeticum* 44**
***Radula* 23**
R. subgenus *Radula* 23
R. subgenus *Volutoradula* 23
***R. aquilegia* 23**
***R. carringtonii* 23**
***R. complanata* 23**
subsp. *lindenbergiana* = *Radula lindenbergiana* 23
var. *alpestris* = *Radula complanata* 23
***R. holtii* 23**
***R. jonesii* 23**
R. lindenbergiana = *Radula lindenbergiana* 23
***R. lindenbergiana* 23**
***R. nudicaulis* 23**
var. *delicatula* 23
var. *nudicaulis* 23
[*R. prolifera*] 69
***R. voluta* 23**
***R. visianica* 23**
***R. wichurae* 23**
RADULACEAE 6, 23
Radulineae 6
***Reboulia* 25**
***R. hemisphaerica* 25**
[subsp. *acrogyna*] 25
subsp. *australis* 25
subsp. *dioica* 25
subsp. *hemisphaerica* 25
[subsp. *orientalis*] 25
[subsp. *paradoxa*] 25
[var. *fissisquama*] 25

- [var. *turkestanica*] 25
- Rhabdoweisia** 35
- R. crenulata** 35
- R. crispata** 35
- R. fugax** 35
- RHABDOWEISACEAE: 7, 35
- Rhamphidium** 36
- R. purpuratum** 36
- RHIZOGONIACEAE 8, 55
- RHIZOGONIALES 8, 55
- Rhizomnium** 53
- R. andrewsianum** 53
- R. gracile** 53
- R. magnifolium** 53
- R. pseudopunctatum** 53
- R. punctatum** 53
- var. *hermanperssonii* 53
- var. *punctatum* 53
- [*R. striatulum*] 70
- Rhodobryum** 3, 51
- R. ontariense** 51
- R. roseum** 51
- Rhynchostegiella** 60
- R. azorica** 60
- R. bourgaeana** 60
- R. curviseta** 60
- R. durieui* = *Pseudorhynchostegiella durieui* 60
- R. jacquinii* = *Rhynchostegiella teneriffae* 60
- R. litorea** 60
- R. macilenta* = *Rhynchostegiella teneriffae* 60
- R. pseudolitorea** 60
- R. teesdalei* = *Rhynchostegiella teneriffae* 60
- R. tenella** 60
- var. *meridionalis* = *Rhynchostegiella litorea* 60
- R. teneriffae** 60
- R. tenuicaulis* = *Brachythecium tenuicaule* 60
- R. trichophylla** 60
- R. tubulosa** 60
- Rhynchostegium** 59
- R. alopecuroides** 59
- R. arcticum* = *Rhynchostegium murale* 60
- R. confertum** 59
- R. confusum** 59
- R. megapolitanum** 60
- R. murale** 60
- R. riparioides** 60
- R. rotundifolium** 60
- R. strongylense** 60
- RHYTIDIACEAE 8, 64
- Rhytidiadelphus** 64
- R. loreus** 64
- R. squarrosus** 64
- R. subpinnatus** 64
- R. triquetrus* = *Hylocomiadelphus triquetrus* 63
- Rhytidium** 64
- R. rugosum** 64
- Riccardia** 23
- R.* subgenus *Riccardia* 23
- R.* section *Riccardia* 23
- R. chamedryfolia** 23
- var. *major* = *Riccardia chamedryfolia* 23
- var. *submersa* = *Riccardia chamedryfolia* 23
- R. incurvata** 23
- R. latifrons** 23
- subsp. *arctica* 23
- subsp. *latifrons* 23
- [var. *miyakeana*] 23
- R. multifida** 23
- [subsp. *decrescens*] 23
- subsp. *multifida* 23
- [subsp. *synoica*] 23
- R. palmata** 23
- Riccia** 4, 26
- R.* subgenus *Riccia* 26
- R.* subgenus *Ricciella* 27
- R.* section *Pilifer* 27
- R.* section *Riccia* 26
- R.* section *Ricciella* 27
- R.* section *Spongodes* 27
- R. atlantica** 26
- R. atromarginata** 26
- var. *atromarginata* 26
- [var. *jovet-astiae*] 26
- R. beyrichiana** 26
- R. bicarinata** 26
- R. bifurca** 26
- var. *subinermis* = *Riccia bifurca* 26
- R. boumanii** 27
- R. breidleri** 26
- [*R. bullosa*] 67
- [*R. capensis* = *Riccia concava*] 67
- R. cavernosa** 27
- var. *angustior* = *Riccia cavernosa* 27
- R. canaliculata** 27
- R. canescens* = *Riccia ciliata* 26
- [*R. chudoana*] 69
- R. ciliata** 26
- var. *epilosa* = *Riccia ciliata* 26
- var. *intumescens* = *Riccia ciliata* 26
- var. *violacea* = *Riccia ciliata* 26
- R. ciliifera** 26
- [*R. concava*] 67
- [*R. congoana*] 69
- [*R. crinita*] 26
- R. crinita* auct. eur. = *Riccia ciliata* 26
- R. crozalsii** 26
- R. crustata** 27
- R. crystallina** 26
- R. dalslandica* = *Riccia ciliata* 26
- R. duplex** 27
- var. *duplex* 27
- [var. *megaspora*] 27
- R. fluitans** 27, 68
- R. frostii** 27
- [var. *crystallinoides*] 27
- var. *frostii* 27
- R. glauca** 26
- var. *ciliaris* 26
- var. *glauca* 26
- var. *major* = *Riccia glauca* 26
- var. *subinermis* = *Riccia glauca* var. *ciliaris* 26
- R. gothica** 26
- R. gougetiana** 26
- var. *armatissima* 26
- var. *gougetiana* 26
- R. huebeneriana** 27
- subsp. *huebeneriana* 27
- [subsp. *sullivantii*] 27
- var. *pseudo-frostii* = *Riccia huebeneriana* subsp. *huebeneriana* 27
- R. intumescens* = *Riccia ciliata* 26
- R. lamellosa** 26
- R. ligula** 26
- [*R. limbata*] 67

R. lindenbergiana = *Riccia sorocarpa* subsp. *arctica* 26

R. macrocarpa 26

[*R. mamillata*] 69

R. marginata = *Riccia beyrichiana* 26

R. melitensis = *Riccia ciliifera* 26

R. michelii 26

R. nigrella 26, 67

R. oelandica = *Riccia subbifurca* 27

[*R. oerstediana*] 68

R. papillosa 26

R. perennis 27

[*R. polycarpa*] 69

R. rhenana 27

var. *rhenana* 27

var. *violacea* 27

R. sommieri 26

R. sorocarpa 26

subsp. *arctica* 4, 26

subsp. *erythrophora* 5, 27

subsp. *sorocarpa* 27

var. *arctica* = *Riccia sorocarpa* subsp. *arctica* 4

var. *heegii* = *Riccia sorocarpa* 26

R. subbifurca 27

R. teneriffae = *Riccia cavernosa* 27

R. trabutiana 27

R. trichocarpa = *Riccia ciliata* 26

R. warnstorffii 27

var. *ciliaris* = *Riccia warnstorffii* 27

var. *commutata* = *Riccia warnstorffii* 27

var. *subinermis* = *Riccia warnstorffii* 27

RICCIACEAE 7, 26

Riccicarpos 27

R. natans 27

Riella 27

R. subgenus *Riella* 27

R. subgenus *Trabutiella* 27

R. affinis 27

R. battandieri 27

R. bialata 27

R. cossoniana 27

[*R. cyrenaica*] 27, 69

R. echinata 27

R. gallica 27

R. helicophylla 27

var. *helicophylla* 27

var. *macrocarpa* 27

R. mediterranea 27

R. notarisii 27

[*R. numidica*] 69

[*R. parisii*] 68, 69

R. reuteri 27

[*R. sersuensis*] 27, 69

RIELLACEAE 7, 27

Roaldia 63

R. dolomitica 63

R. revoluta 63

Rosulabryum = *Ptychostomum* 3, 50

[*R. albolimbatum*] 50

[*R. billardierei*] 50

R. bornholmense = *Ptychostomum bornholmense* 50

R. canariense = *Bryum canariense* 48

R. capillare = *Ptychostomum capillare* 50

R. donianum = *Ptychostomum donianum* 50

R. elegans = *Ptychostomum elegans* 50

R. laevifilum = *Ptychostomum moravicum* 51

R. rubens = *Ptychostomum rubens* 51

R. torquescens = *Ptychostomum torquescens* 51

Saccobasis 14

S. polita 14

S. polymorpha 14

Saccogyna 18

S. viticulosa 18

SACCOGYNACEAE 6, 18

Saelania 43

S. glaucescens 43

SAELANIACEAE 7, 43

Sanionia 58

S. nivalis 58, 68

[*S. georgicouncinata*] 58, 68

S. georgicouncinata auct. eur. = *Sanionia nivalis* 58

S. orthothecioides 58

S. uncinata 58

Sarmentypnum 58

S. exannulatum 58

S. procerum 58

S. sarmentosum 58

S. trichophyllum 58

S. tundrae 58

Sauteria 25

S. section *Sauteria* 25

S. alpina 25

Scapania 5, 14

S. subgenus *Gracilidae* 14

S. subgenus *Plicicalyx* 14

S. subgenus *Scapania* 14

S. section *Aequilobae* 14

S. section *Apiculatae* 14

S. section *Compactae* 14

S. section *Curtae* 15

S. section *Cuspiduligerae* 15

S. section *Hyperboreae* 15

S. section *Kaalaasia* 15

S. section *Nemorosae* 15

S. section *Planifoliae* 14

S. section *Scapania* 15

S. section *Scapaniella* 15

S. section *Simmonsiae* 15

S. section *Sphaeriferae* 15

S. section *Verrucosae* 15

S. aequiloba 14

S. apiculata 14

S. aspera 14

S. brevicaulis auct. eur. = *Scapania degenii* 15

[*S. brevicaulis*] 15, 69

var. *dubia* = *Scapania degenii* 15

S. calcicola 15

S. carinthiaca 14

var. *carinthiaca* 14

var. *massalongi* 14

S. compacta 14

S. crassiretis 15

S. curta 15

var. *curta* 15

var. *grandiretis* 15

var. *isoloba* 15

S. cuspiduligera 15

var. *cuspiduligera* 15

[var. *diplophyllopsis*] 15

S. degenii 15

var. *dubia* = *Scapania degenii* 15

S. glaucocephala 15

var. *glaucocephala* 14

[var. *saxicola*] 15

var. *scapanoides* = *Scapania scapanioides* 15

- S. gracilis* 14**
***S. gymnostomophila* 15**
***S. helvetica* 15**
***S. hyperborea* 15**
***S. irrigua* 15**
 subsp. *irrigua* 15
 subsp. *rufescens* 15
 var. *rufescens* = *Scapania irrigua* subsp. *irrigua* 15
***S. kaurinii* 14**
***S. ligulifolia* 15**
***S. lingulata* 15**
 var. *lingulata* 15
 var. *microphylla* 15
 [*S. microdonta*] 69
***S. mucronata* 15**
 subsp. *praetervisa* = *Scapania praetervisa* 15
***S. nemorea* 15**
***S. nimbosea* 14**
***S. obcordata* 15**
***S. obscura* 15**
***S. ornithopodioides* 14**
***S. paludicola* 5, 15**
 var. *paludicola* 15
 var. *rotundiloba* 5, 15
 var. *rufescens* = *Scapania paludicola* 15
 [var. *viridigemma*] 15
***S. paludosa* 15**
 var. *isoloba* = *Scapania paludosa* 15
 var. *rubiginosa* = *Scapania paludosa* 15
 var. *vogesiacae* = *Scapania paludosa* 15
S. paradoxa = *Scapania obcordata* 15
***S. parvifolia* 15**
***S. praetervisa* 15**
***S. scandica* 15**
 var. *argutedentata* 15
 [var. *dimorpha*] 15
 var. *grandiretis* 15
 var. *scandica* 15
***S. scapanioides* 15**
***S. simmonsii* 15**
***S. sphaerifera* 15**
***S. spitsbergensis* 15**
***S. subalpina* 15**
 [var. *haynesiae*] 15
 [var. *muddiae*] 15
 var. *subalpina* 15
***S. tundrae* 15**
***S. uliginosa* 15**
***S. umbrosa* 15**
***S. undulata* 15**
 var. *aequatiformis* = *Scapania undulata* 15
 var. *dentata* = *Scapania undulata* 15
 var. *oakesii* = *Scapania undulata* 15
***S. verrucosa* 15**
***S. zemliae* 15**
 SCAPANACEAE 6, 14
Schistidium 46
***S. abrupticostatum* 46**
***S. agassizii* 46**
***S. andreaeopsis* 46**
***S. apocarpum* 46**
***S. atrichum* 46**
***S. atrofusum* 46**
***S. boreale* 46**
***S. brunnescens* 46**
 subsp. *brunnescens* 46
 subsp. *griseum* 46
***S. bryhnii* 46**
***S. canadense* 46**
 [*S. cinclidodonteum*] 70
***S. confertum* 46**
***S. confusum* 46**
***S. convergens* 46**
***S. crassipilum* 46**
***S. crenatum* 46**
***S. dupretii* 46**
***S. echinatum* 46**
***S. elegantulum* 46**
 subsp. *elegantulum* 46
 subsp. *wilsonii* 46
***S. flaccidum* 46**
***S. flexipile* 46**
***S. frigidum* 46**
 var. *frigidum* 46
 var. *havaasii* 46
***S. frisvollianum* 46**
***S. grande* 46**
***S. grandirete* 46**
***S. helveticum* 46**
***S. holmenianum* 46**
***S. lancifolium* 46**
***S. marginale* 46**
***S. maritimum* 46**
 subsp. *maritimum* 46
 subsp. *piliferum* 46
***S. memnonium* 116**
***S. obscurum* 46**
***S. occidentale* 46**
***S. papillosum* 46**
***S. platyphyllum* 46**
 var. *abrupticostatum* = *Schistidium abrupticostatum* 46
***S. poeltii* 46**
***S. pruinatum* 46**
***S. pulchrum* 46**
***S. recurvum* 46**
***S. rivulare* 46**
***S. robustum* 46**
***S. scandicum* 46**
***S. sibiricum* 46**
***S. sinensiapocarpum* 46**
S. singarense = *Schistidium helveticum* 46
***S. sordidum* 46**
***S. spinosum* 46**
***S. strictum* 46**
***S. subconfertum* 46**
***S. subflaccidum* 46**
***S. subjulaceum* 46**
***S. submuticum* 46**
 subsp. *arcticum* 46
 subsp. *submuticum* 46
***S. succulentum* 46**
***S. tenerum* 46**
***S. tenuinerve* 46**
***S. trichodon* 46**
 var. *nutans* 46
 var. *trichodon* 46
***S. umbrosum* 46**
***S. venetum* 46**
Schistochilopsis 15
***S. grandiretis* 15**
S. elegans = *Lophozia excisa* var. *elegans* 13
***S. hyperarctica* 15**
***S. incisa* 15**
 var. *inermis* = *Schistochilopsis incisa* 15

- S. opacifolia** 15
Schistostega 36
S. pennata 36
SCHISTOSTEGACEAE 7, 36
Schizophylloopsis 10
S. sphenoloboides 10
Schizymerium 52
S. pontevedrense 52
Schljakovia 10
S. kunzeana 10
Schljakovianthus 10
S. quadrilobus 10
Sciuro-hypnum 61
S. curtum 61
S. dovrense 61
S. flotowianum 61
S. glaciale 61
S. latifolium 61
S. oedipodium 62
S. ornellanum 62
S. plumosum 62
S. populeum 62
S. reflexum 62
S. starkei 62
S. tromsoeense 62
Scleropodium 62
S. cespitans 62
S. touretii 62
Scopelophila 36
S. cataractae 36
S. ligulata 36
SCORPIDIACEAE 8, 58
Scorpidium 58
S. cossonii 58
S. revolvens 58
S. scorpioides 58
Scorpiurium 60
S. circinatum 60
S. deflexifolium 60
S. sendtneri 60
Seligeria 43
S. subgenus *Blindiadelphus* = *Blindiadelphus* 43
S. subgenus *Megalosporia* 43
S. subgenus *Seligeria* 43
S. austriaca 43
S. acutifolia 43
S. *alpestris* = *Seligeria patula* 43
S. brevifolia 43
S. calcarea 43
S. calycina 43
S. *campylopoda* = *Blindiadelphus campylopodus* 43
S. carniolica 43
S. *diversifolia* = *Blindiadelphus diversifolius* 43
S. donniana 43
S. *galinae* = *Seligeria donniana* 43
S. irrigata 43
S. oelandica 43
S. patula 43
var. *alpestris* = *Seligeria patula* 43
S. *paucifolia* auct. eur. = *Seligeria calycina* 43
S. *polaris* = *Blindiadelphus polaris* 43
S. pusilla 43
S. *recurvata* = *Blindiadelphus recurvatus* 43
S. *subimmersa* = *Blindiadelphus subimmersus* 43
S. *transylvanica* = *Brachydontium trichodes* 43
S. trifaria 43
var. *longifolia* 43
var. *trifaria* 43
S. tristichoides 43
var. *patula* = *Seligeria patula* 43
SELIGERIACEAE 7, 43
SEMATOPHYLLACEAE 8, 63
Sematophyllum 63
S. adnatum 4, 63
S. demissum 63
S. substrumosum 63
Serpoleskea 58
S. confervoides 58
Solenostoma 18
S. subgenus *Eucalyx* 18
S. subgenus *Metasolenostoma* 18
S. subgenus *Plectocolea* 18
S. subgenus *Solenostoma* 18
S. callithrix 18
S. caucasicum 18
S. confertissimum 18
S. gracillimum 18
S. handelii 18
S. hyalinum 18
[*S. lignicola*] 69
S. obovatum 18
S. paroicum 18
S. *pusillum* = *Solenostoma sphaerocarpum* 18
S. sphaerocarpum 18
var. *nanum* = *Solenostoma sphaerocarpum* 18
S. subellipticum 18
[*S. subtilissimum*] 69
SOLENOTOMATAACEAE 6, 18
Southbya 19
S. nigrella 19
S. tophacea 19
SOUTHBYACEAE 6, 19
SPHAEROCARPACEAE 7, 27
SPHAEROCARPALES 7, 27
Sphaerocarpos 27
S. subgenus *Austrosphaerocarpos* 27
S. subgenus *Sphaerocarpos* 27
S. europaeus 27, 67
S. michelii 27
S. stiptatus 4, 27
S. *texanus* auct. = *Sphaerocarpos europaeus* 27, 68
[*S. texanus*] 27, 68
SPHAGNACEAE 2, 7, 27
SPHAGNALES 7, 27
SPHAGNOPHYTINA 7
SPHAGNOPSIDA 7, 27
Sphagnum 27
S. subgenus *Acutifolia* 28
S. subgenus *Cuspidata* 28
S. subgenus *Rigida* 28
S. subgenus *Sphagnum* 28
S. subgenus *Subsecunda* 28
S. section *Acutifolia* 28
S. section *Insulosa* 28
S. section *Polyclada* 28
S. section *Squarrosa* 28
S. affine 28
var. *flagellare* = *Sphagnum affine* 28
S. angermanicum 28
S. angustifolium 28
S. annulatum 28
S. aongstroemii 28
S. arcticum 28
S. auriculatum 28, 68

- S. austinii** 28
S. balticum 28
S. beothuk 28
S. brevifolium = *Sphagnum fallax* var. *brevifolium* 28
S. capillifolium 28
 subsp. *capillifolium* = *Sphagnum capillifolium* 28
 subsp. *rubellum* = *Sphagnum rubellum* 28
S. centrale 28
S. compactum 28
S. concinnum 28
S. contortum 28
S. cuspidatum 28
 var. *cuspidatum* 28
 var. *viride* 28
S. denticulatum = *Sphagnum auriculatum* 28
S. divinum 28
S. fallax 28
 subsp. *isoviitae* = *Sphagnum fallax* var. *isoviitae* 28
 var. *brevifolium* 28
 var. *fallax* 28
 var. *isoviitae* 28
S. fimbriatum 28
 subsp. *concinnum* = *Sphagnum concinnum* 28
S. flexuosum 28
S. fuscum 28
S. girgensohnii 28
 [*S. imbricatum*] 28
 subsp. *affine* = *Sphagnum affine* 28
 subsp. *austinii* = *Sphagnum austinii* 28
S. inundatum 28, 68
S. isoviitae = *Sphagnum fallax* var. *isoviitae* 28
S. jensenii 29
S. lenense 29
 [*S. lescurii*] 68, 70
S. lindbergii 29
 [*S. magellanicum* Brid.] 28, 68
S. magellanicum auct. eur. p.p. = *Sphagnum divinum* 28
S. magellanicum auct. eur. p.p. = *Sphagnum medium* 28
S. majus 29
 subsp. *majus* 29
 subsp. *norvegicum* 29
S. medium 28
S. mirum 28
S. molle 28
S. nitidulum 28
S. obtusum 29
S. olafii 28
S. palustre 28
 var. *centrale* = *Sphagnum centrale* 28
S. papillosum 28
 [*S. perfoliatum*] 70
S. platyphyllum 28
S. pulchrum 29
S. pylaesii 28
S. quinquefarium 28
S. recurvum 29
S. riparium 29
S. rubellum 28
S. rubiginosum 28
S. russowii 28
S. skyense 28
S. squarrosum 28
S. strictum 28
S. subfulvum 28
 subsp. *purpureum* 28
 subsp. *subfulvum* 28
S. subnitens 28
 subsp. *ferrugineum* 28
 subsp. *subnitens* 28
 var. *ferrugineum* = *Sphagnum subnitens* subsp. *ferrugineum* 28
S. subsecundum 28
S. tenellum 29
S. teres 28
S. tescorum 28
S. troendelagicum 29
S. tundrae 28
S. venustum 28
S. viride = *Sphagnum cuspidatum* var. *viride* 28
S. warnstorffii 28
S. wulfianum 28
Sphenolobopsis 10
S. pearsonii 10
Sphenolobus 10
S. cavifolius = *Orthocaulis cavifolius* 10
S. minutus 10
S. saxicola 10
 SPLACHNACEAE 7, 47
 SPLACHNALES 7, 47
Splachnobryum 42
 [*S. aquaticum*] 70
 [*S. limbatum*] 70
S. obtusum 42
 Splachnoideae 48
Splachnum 48
S. ampullaceum 48
S. luteum 48
S. melanocaulon 48
S. pensylvanicum 48
S. rubrum 48
S. sphaericum 48
S. vasculosum 48
Stegonia 39
S. latifolia 39
 var. *latifolia* 39
 var. *pilifera* 39
Stereodon 62
S. aemulans 63
S. callichrous 63
S. fertilis = *Pseudohygrohypnum fertile* 63
S. hamulosus 63
S. holmenii 63
S. pratensis 63
S. subimponens 63
 STERIODONTACEAE 8, 62
Straminergon 58
S. stramineum 58
 Streblotrichoideae 40
Streblotrichum 40
S. convolutum 40
 var. *commutatum* 40
 var. *convolutum* 40
S. commutatum = *Streblotrichum convolutum* var. *commutatum* 40
S. enderesii 40
Streptocolea = *Grimmia* 45
S. atrata = *Grimmia atrata* 45
Syntrichia 39
 [*S. amphidiacea*] 70
S. amplexa = *Tortula amplexa* 39
S. bogotensis 39
S. calcicola 39
S. caninervis 39
 var. *abranchesii* 39
 var. *astrakhanica* 39

- var. **caninervis** 39
var. **gypsophila** 39
[var. *pseudodesertorum*] 70
S. densa = *Syntrichia ruralis* 39
S. echinata 39
S. fragilis 39
S. glabra = *Syntrichia ruralis* 39
S. handellii 39
S. laevipila 39
S. latifolia 39
S. minor 39
S. montana 39
var. **calva** 39
var. **montana** 39
S. norvegica 39
S. pagorum = *Syntrichia laevipila* 39
S. papillosa 39
S. papillosissima 39
S. princeps 39
S. rigescens 39
S. ruraliformis 39
S. ruralis 39
var. **epilosa** 39
var. *ruraliformis* = *S. ruraliformis* 39
var. **ruralis** 39
S. sinensis 39
S. submontana 39
S. subpapillosissima 39
S. virescens 39
Syzygiella 9
S. subgenus *Syzygiella* 9
S. autumnalis 9
[*S. manca*] 69
S. rubricaulis 9
Targionia 27
T. subgenus *Targionia* 27
T. hypophylla 27
subsp. **hypophylla** 27
[subsp. *linealis*] 27
T. lorbeeriana 27
TARGIONIACEAE 7, 27
TAXIPHYLLACEAE 8, 62
Taxiphyllum 62
T. densifolium 62
T. wissgrillii 62
Tayloria 47
T. acuminata 47
T. froelichiana 47
T. hornschurchii 47
T. lingulata 47
T. rudolphiana 47
T. serrata 47
T. splachnoides 47
T. tenuis 47
Taylorioideae 47
Telaranea 19
T. azorica 19
T. europaea 19, 67
T. longii = *Tricholepidozia tetradactyla* 19
T. murphyae = *Tricholepidozia tetradactyla* 19
T. nematodes auct. eur. = *Telaranea europaea* 19
[*T. nematodes*] 67
T. sejuncta auct. eur. = *Telaranea europaea* 19
[*T. sejuncta*] 67
Tetralophozia 10
T. filiformis 10
T. setiformis 10
TETRAPHIDACEAE 7, 29
TETRAPHIDALES 7, 29
Tetraphidopsida 7, 29
Tetraphis 29
T. pellucida 29
Tetraplodon 48
T. angustatus 48
T. blyttii 48
T. mnioides 48
T. pallidus 48
T. paradoxus 48
T. urceolatus 48
Tetrastichium 55
T. fontanum 55
T. virens 55
Tetrodontium 29
T. brownianum 29
T. ovatum 29
T. repandum 29
Thamnobryum 65
T. alopecurum 65
T. angustifolium 65
T. cataractarum 65
T. fernandesii 65
T. maderense 65
T. neckeroides 65
T. rudolphianum 65
T. subserratum 65
THUIDIACEAE 8, 59
Thuidiopsis 59
T. sparsa 4, 59
Thuidium 59
T. assimile 59
T. delicatulum 59
T. recognitum 59
T. tamariscinum 59
Timmia 30
T. section *Norvegica* 30
T. section *Timmia* 30
T. section *Timmiaurea* 30
T. austriaca 30
T. bavarica 30
T. comata 30
T. megapolitana 30
T. norvegica 30
T. sibirica 30
TIMMIACEAE 7, 30
TIMMIALES 7, 30
Timmiella 32
T. anomala 32
T. barbuloides 32
T. flexiseta 32
TIMMIELLACEAE 7, 32
Timmiidae 7
Tomentypnum 58
T. nitens 58
Tortella 42
T. alpicola 42
[*T. arctica*] 42, 68
T. bambergeri = *Tortella tortuosa* 43
T. bambergeri auct. p.p. = *Tortella fasciculata* 43
T. bambergeri auct. p.p. = *Tortella pseudofragilis* 42
T. x cuspidatissima 42, 68
T. densa 42
T. fasciculata 42
T. flavovirens 42
var. **flavovirens** 42

- var. *glareicola* 42
var. *papillosissima* 42
T. fragilis 42
var. *moravica* = *Tortella pseudofragilis* 42
T. humilis 42
T. inclinata 42
var. *densa* = *Tortella densa* 42
T. inflexa 42
T. limbata 42
T. limosella = *Tortella flavovirens* 42
[*T. malacophylla*] 70
T. mediterranea 42
T. nitida 42
T. pseudofragilis 42
T. rigens 42
T. spitsbergensis 42
T. squarrosa 42
T. tortuosa 43
subsp. *fasciculata* = *Tortella fasciculata* 42
var. *fragilifolia* = *Tortella tortuosa* 43
Tortula 39
T. acaulon 39, 43
var. *acaulon* 39
[var. *galilaea*] 70
var. *papillosa* 39
var. *pilifera* 39
var. *retortifolia* 39
var. *schreberiana* 39
T. amplexa 39
T. amplioretis 39
T. atrovirens 39
T. baetica = *Tortula israelis* 40
T. bogosica 4, 39
T. bolanderi 4, 39
T. brevissima 39
T. canescens 39
T. caucasica 39
T. cernua 40
T. cuneifolia 40
T. freibergii 40
[*T. grandiretis*] 70
T. guepinii 40
T. hoppeana 40
T. inermis 40
T. israelis 40
[*T. kneuckeri*] 70
T. lanceola = *Tortula lindbergii* 40
T. laureri 40
T. leucostoma 40
T. lindbergii 40
T. lingulata 40
T. marginata 40
T. modica = *Tortula caucasica* 40
T. mucronifolia 40
T. muralis 40
subsp. *muralis* 40
subsp. *obtusifolia* 40
var. *aestiva* 40
var. *muralis* 40
T. obtusifolia = *Tortula muralis* subsp. *obtusifolia* 40
T. pallida 40
[*T. plintobia*] 70
T. protobryoides 40
T. randii 40
T. revolvens 40
T. rhodonia = *Henediella heimii* 38
T. schimperi 40
T. solmsii 40
T. subulata 40
var. *angustata* = *Tortula schimperi* 40
var. *graefii* = *Tortula subulata* 40
var. *subinermis* = *Tortula subulata* 40
T. systylia 40
T. truncata 40
T. ucrainica 40
T. vahliana 40
T. viridifolia 40
T. vlassovii = *Microbryum vlassovii* 38
T. wilsonii 40
Trachycystis 53
T. ussuriensis 53
Trematodon 36
T. ambiguus 36
T. brevicollis 36
T. laetevirens 36
T. longicollis 36
T. perssoniorum 36
Trichocolea 21
T. tomentella 21
TRICHOCOLEACEAE 6, 21
Trichodon 36
T. cylindricus 36
Tricholepidozia 19
T. lindenbergii 4, 19
[var. *complanata*] 19
var. *lindenbergii* 4, 19
[var. *mellea*] 19
[var. *papillata*] 19
T. tetradactyla 4, 19
Trichostomoideae 40
Trichostomopsis = *Didymodon* 37
T. australasiae = *Didymodon australasiae* 37
T. umbrosa = *Didymodon umbrosus* 38
Trichostomum 43
T. arcticum = *Tortella spitsbergensis* 42
T. brachydontium 43
T. crispulum 43
T. hibernicum = *Chionoloma hibernicum* 40
T. pallidisetum = *Pottiopsis caespitosa* 42
T. recurvifolium = *Chionoloma recurvifolium* 41
T. tenuirostre = *Chionoloma tenuirostre* 41
T. triumphans = *Pottiopsis caespitosa* 42
Trilophozia 14
T. quinquedentata 14
[var. *assymetrica*] 14
var. *quinquedentata* 14
Triquetrella 40
T. arapilensis 40
Tritomaria 14
T. exsecta 14
subsp. *exsecta* 14
[subsp. *novaezealandia*] 14
T. exsectiformis 14
subsp. *arctica* 14
[subsp. *camerunensis*] 14
subsp. *exsectiformis* 14
T. polita = *Saccobasis polita* 14
subsp. *polymorpha* = *Saccobasis polymorpha* 14
T. quinquedentata = *Trilophozia quinquedentata* 14
subsp. *turgida* = *Trilophozia quinquedentata* 14
var. *dentata* = *Trilophozia dentata* 14
var. *grandiretis* = *Trilophozia quinquedentata* 14
f. *gracilis* = *Trilophozia quinquedentata* 14
T. scitula 14

Trochobryum = *Seligeria* 43
T. carniolicum = *Seligeria carniolica* 43
Tylimanthus = *Acrobolbus* 16
[*T. anisodontus* = *Acrobolbus anisodontus*] 66
T. azoricus = *Acrobolbus azoricus* 16
T. madeirensis = *Acrobolbus madeirensis* 16
Ulotia 54
U. bruchii 54
U. calvescens 54
U. coarctata 54
U. crispa 54
var. *crispula* = *Ulotia crispula* 54
var. *intermedia* = *Ulotia intermedia* 54
U. crispula 54
U. curvifolia 54
U. drummondii 54
U. hutchinsiae 54
U. intermedia 54
U. macrospora 54
U. phyllantha = *Phenogemma phyllantha* 54
U. rehmannii 54
Vesicularia 63
V. reimersiana 63
Vinealobryum = *Didymodon* 37
V. brachyphyllum = *Didymodon brachyphyllum* 37
V. cordatum = *Didymodon cordatus* 37
V. eckeliae = *Didymodon eckeliae* 37
V. insulanum = *Didymodon insulanus* 37
V. luridum = *Didymodon luridus* 37
V. nicholsonii = *Didymodon nicholsonii* 38
V. sinuosum = *Didymodon sinuosus* 38
V. vineale = *Didymodon vinealis* 38
Voitia 48
V. hyperborea 48
V. nivalis 48
Warnstorfia 58
W. exannulata = *Sarmentypnum exannulatum* 58
W. fluitans 58
W. procera = *Sarmentypnum procerum* 58
W. pseudostraminea 58
W. sarmentosa = *Sarmentypnum sarmentosum* 58
W. trichophylla = *Sarmentypnum trichophyllum* 58
W. tundrae = *Sarmentypnum tundrae* 58

Addenda

1. *Herbertus* p. 19:

Sun and He (2019) recently proposed synonymising *Herbertus borealis*, *Herbertus norenius* and *Herbertus stramineus* with *Herbertus aduncus* (in which they include *Herbertus hutchinsiae*). While undoubtedly closely related, the small genetic differences between them result in very clear morphological differences, and in the field the species (which sometimes grow in close proximity) are easily distinguished by morphology, ecology and distribution. We therefore reject the synonymies.

Weissia 43

W. angustifolia 43
W. brachycarpa 43
[*W. breutelii*] 70
W. condensata 43
var. *armata* 43
var. *condensata* 43
W. controversa 43
var. *controversa* 43
var. *crispata* 43
var. *densifolia* 43
[*W. leptocarpa*] 70
W. levieri 43
W. longifolia 43
var. *angustifolia* = *Weissia angustifolia* 43
W. x mittenii 43
W. multicapsularis = *Tortula acaulon* 43
W. multicapsularis auct. = *Weissia wilsonii* 43
[*W. ovatifolia*] 70
W. perssonii 43
W. rostellata 43
W. rutilans 43
[*W. sinaloensis*] 70
W. squarrosa 43
W. sterilis 43
W. triumphans = *Pottiopsis caespitosa* 42
W. tyrrhena = *Pottiopsis caespitosa* 42
W. wilsonii 43
W. wimmeriana 43
Zoopsidoideae 19
Zygodon 54
Z. bistratus = *Zygodon catarinoi* 54
Z. catarinoi 54
Z. conoideus 54
var. *conoideus* 54
var. *lingulatus* 54
Z. dentatus 54
Z. forsteri = *Codonoblepharon forsteri* 53
Z. gracilis 54
Z. rupestris 55
Z. sibiricus 55
Z. stirtonii 55
Z. viridissimus 55

2. Additional species:

Physcomitrium (Callaghan et al. 2020) p. 31
23.6i **Physcomitrium** × **stevensonii** (**stevensonii**)
D.A.Callaghan
Schistidium p. 46 (Guerra et al. 2020)
122.29i Schistidium memnonium J.Guerra
Meesia (Hedenäs 2020) p. 48
138.4i **Meesia minor** Brid.
138.4ii **Meesia minutissima** Hedenäs