



AT THE FOREFRONT
UChicago
Medicine

Survive and Thrive 2019

Framework for Understanding Rare Tumors in Ovarian Cancer

Nita Karnik Lee, MD, MPH

Department of Obstetrics and Gynecology

Section of Gynecology Oncology



AT THE FOREFRONT

UChicago**Medicine**

Comprehensive Cancer Center

Objectives:

- Understand basic classification of rare ovarian cancers
- Review incidence and unique features
- Treatment choices in first line treatment
- Surveillance options
- Role of tumor and somatic genetic testing
- Discuss advocacy for rare tumors


NO DISCLOSURES



AT THE FOREFRONT
UChicago
Medicine

Challenges and Opportunities for Rare Ovarian Cancers

- Challenge and frustration of having a rare type of an already rare tumor
- Fewer cases to review and learn
- Harder to accrue and complete clinical trials
- Funding for research
- Diversity of mutations in each cell type
- Opportunities for collaborative research and support networking
- Opportunities for targeted therapies



Not that simple

Ovarian Cancer is not one disease

Epithelial ovarian cancer


What is the Cell of origin?

fallopian tube origin


ovarian surface lining cells

endometriosis


lining of the peritoneal cavity



Source: Drapkin R, Karst AM. The new face of ovarian cancer modeling: Better prospects for detection and treatment. *F1000 Med Reports*. 2011;3:22. Used with permission.




Epithelial Ovarian Cancers (EOC)



- 80% of ovarian cancers are EOC
 - 80% are Serous carcinomas
- Non EOC
 - Germ Cell Tumors
 - Sex Cord Stromal Tumors
 - Younger age
 - Better outcomes



Types of Ovarian Cancer




5-8% of ovarian cancers

~5%

Epithelial Ovarian Cancer
Fallopian Tube Cancer
Primary Peritoneal Cancer

Stage IIC Cancer



© 2005 American Society of Clinical Oncology



AT THE FOREFRONT
UChicago
Medicine

Rare Ovarian Cancers

- Epithelial Ovarian Cancer (high grade serous) ~ 80%
- Malignant Ovarian Germ Cell Tumors 5%
- Sex Cord Stromal Tumors 1.2%
- Rare Epithelial Ovarian Cancers
 - Clear Cell Carcinoma
 - Endometrioid Carcinoma
 - Mucinous carcinomas
 - Low grade serous carcinomas

Cell types of epithelial ovarian cancers

Type	WHC, %
Serous	67.8
Mucinous	3.4
Endometrioid	8.6
Clear Cell	12.6
Transitional	0.6
Mixed	5.7
Undifferentiated	0.6
Other	0

These cell types may have differences in CLINICAL

- Clinical presentation
 - Age; symptoms
- Optimal treatment regimens
- Recurrence risk
- Response to treatment

BIOLOGIC/MOLECULAR

- Different molecular mutations
- Different pathways that are turned on or off that promote cancer development

Framework:

- Treatment Options
 - Primary
 - Maintenance
 - Recurrence
- Follow-up and Surveillance
- Unique Genetics or tumor mutations
- Novel therapies or clinical trials

Use this framework to ask questions about your specific cancer

Malignant Germ Cell Tumors of the Ovary

- Dysgerminoma
- Yolk Sac Tumors
- Embryonal carcinoma
- Choriocarcinoma
- Immature teratoma

Sex Cord Stromal Malignant Ovarian Tumors 5-8%

- Adult granulosa cell tumor
 - Juvenile granulosa cell tumor
 - Sertoli-Leydig cell tumors 0.5% of ovarian cancers
 - Sex cord tumor with annular tubules
-
- Indolent, slower growing
 - Late recurrences

Sex Cord Stromal Tumors

- **Unique features**

 - Younger patients 20-30s

 - Can secrete hormones (estrogen or testosterone)

 - Average size 16 cm ; pelvic mass at presentation

- **Treatment Options**

 - Primary

 - Surgery; Fertility sparing options
 - Chemotherapy

 - Recurrence

 - Repeat surgery
 - Chemotherapy

- **Follow-up and Surveillance**

 - Tumor markers

 - Decision regarding imaging

- **Unique Genetics**

 - Granulosa cell tumors Tumor Mutation FOXL2 mutation

 - Sertoli-Leydig cell tumors DICER 1 genetic mutation

- **Novel therapies or clinical trials**

 - GOG 264**



AT THE FOREFRONT
UChicago
Medicine

DICER 1 – Hereditary Cancer Mutation

- Inherited germline mutation
- Sertoli-Leydig tumors
- Other cancers: pleuropulmonary blastoma, cervical sarcoma

Rare Epithelial Ovarian Cancers

- Endometrioid carcinoma (10 %)
- Clear cell carcinoma (5-10 %)
- Mucinous carcinoma (3-4 %)
- Low grade serous carcinoma (< 5%)
- Carcinosarcoma (2-5%)
- Transitional cell carcinoma (< 2%)
- Small cell carcinoma (< 3%)

Low grade serous carcinoma

- **Unique features**

- Younger patients than average EOC
 - Better overall survival may be seen

- **Treatment Options**

- Primary

- Surgery; Fertility sparing options

- Chemotherapy Stage II – IV

- Concern for chemoresistance

- Consider Hormonal maintenance phase

- Recurrence

- Repeat surgery in select cases

- Chemotherapy ex. Doxil, Avastin,

- **Follow-up and Surveillance**

- Tumor markers

- CA125

- Decision regarding imaging

- **Unique Genetics or tumor characteristics**

- Estrogen or Progesterone receptor positive

- hormonal targets

- Mutations in KRAS/BRAF/MAPK signaling pathway

- Should get genetic testing but BRCA mutations less common ~5%

- **Novel therapies or clinical trials**

- Avastin

- Endocrine therapy

- MEK inhibitors




Low grade serous carcinoma: Endocrine therapy Options

- Aromatase inhibitors
 - inhibits peripheral conversion of steroids to estrogen
- Tamoxifen
 - Selective estrogen receptor modulator (anti-estrogen effects)
- Fulvestrant (faslodex) -Fulvestrant
 - [selective estrogen receptor degrader](#) (SERD)
 - binds to the [estrogen receptor](#) and destabilizing it, causing the cell's normal [protein degradation](#) processes to destroy it.
- Leuprolide – GnRH agonist – chemical menopause

Low grade serous carcinoma: Biologics

- **Avastin** The bevacizumab compound binds to the free VEGF and reduces the concentration of the free VEGF. C, The reduction of available VEGF results in diminished blood supply to the tumor and tumor shrinkage.



Low grade serous carcinoma: Targeted therapies

MEK Inhibitors


- GOG 239 Selumetinib, a MEK1/2 inhibitor

response rate of 15%, with stable disease in 65% and an acceptable toxicity profile

- MILO Study

Phase 3 study of binimetinib or a chemotherapy chosen by a physician (liposomal doxorubicin, paclitaxel or topotecan)

Did not reach PFS improvement compared to advanced ovarian cancer



Source: B. Y. Karlan, R. E. Bristow, A. J. Li: Gynecologic Oncology: Clinical Practice and
www.obgyn.mhmedical.com
Copyright © McGraw-Hill Education. All rights reserved.

Clear cell carcinoma

- **Unique features**

- Perimenopausal age range
- Can be associated with endometriosis
- Risks of hypercalcemia, blood clots

- **Treatment Options**

Primary

Surgery;
Chemotherapy

Concern for chemor esistance

Recurrence

Chemotherapy
rare surgery for secondary

- **Follow-up and Surveillance**

Tumor markers

CA125

Decision regarding imaging

- **Unique Genetics or tumor characteristics**

ARID1 A mutations

PIK3CA mutations (noted in 40% of OCCC)

– Targeting the phosphatidylinositol 3-kinase (PI3K)/Akt/ mammalian target of rapamycin (mTOR) pathway

- **Novel therapies or clinical trials**



AT THE FOREFRONT
UChicago
Medicine

Clear Cell Carcinoma of the Ovary

- A Phase II Study of Tazemetostat (EPZ-6438) (IND # 138671) in Recurrent or Persistent Endometrioid or Clear Cell Carcinoma of the Ovary, and Recurrent or Persistent Endometrioid Endometrial Adenocarcinoma (CIRB) (NRG-GY014)

GOG trials:

- GOG 254-Phase II evaluation of Sunitinib Malate in the treatment of persistent or recurrent clear cell ovarian carcinoma (Closed to patient entry September 2013; late breaking abstract at 2015 SGO))
- GOG 268 A Phase II Evaluation of Temsirolimus (CCI-779) (NCI Supplied Agent: NSC# 683864, IND# 61010) in Combination with Carboplatin and Paclitaxel followed by Temsirolimus (CCI-779) Consolidation as First-line Therapy in the Treatment of Stage III-IV Clear Cell Carcinoma of the Ovary (Closed to patient entry January 2014)
- GOG 283: A Phase II Trial of DCTD-Sponsored Dasatinib (NSC #732517 IND #73969) In Recurrent/Persistent Ovary, Fallopian Tube, Primary Peritoneal, Endometrial, or Endometriosis-Associated Clear Cell Carcinoma Characterized for the Retention or Loss of BAF250a Expression. (suspended)
- GY-001: A randomized phase II study of XL-184 (cabozantinib) in recurrent clear cell carcinoma (Dr Farley)
 - (VEGFR2 inhibitor)



Mucinous ovarian carcinoma

- **Unique features**

- Presents in early stage often
- Larger tumors
- Often can be mets from GI tumors

- **Treatment Options**

- | | | |
|------------|--|-----------------------------|
| Primary | Surgery;
Chemotherapy | Concern for chemoresistance |
| Recurrence | Chemotherapy
rare surgery for secondary | |

- **Follow-up and Surveillance**

- | | |
|----------------------------|-----|
| Tumor markers | CEA |
| Decision regarding imaging | |

- **Unique Genetics or tumor characteristics**

- KRAS mutations >75%

- **Novel therapies or clinical trials**

- | | | |
|------------------------------|---------|-------|
| GI regimens may show promise | Avastin | HIPEC |
|------------------------------|---------|-------|

