

Non-Native Species in Great Britain: establishment, detection and reporting to inform effective decision making

Helen E. Roy, Jim Bacon, Björn Beckmann, Colin A. Harrower, Mark O. Hill, Nick J.B. Isaac, Chris D. Preston, Biren Rathod, Stephanie L. Rorke

Centre for Ecology & Hydrology, Wallingford, Oxfordshire, OX10 8BB

John H. Marchant, Andy Musgrove, David Noble

British Trust for Ornithology

Jack Sewell, Becky Seeley, Natalie Sweet, Leoni Adams, John Bishop

Marine Biological Association

Alison R. Jukes, Kevin J. Walker and David Pearman

Botanical Society of the British Isles

July 2012

Contents

Summary.....	4
GB Non-Native Species Report Card 2011.....	7
1. Introduction.....	10
2. Aims.....	11
3. GB-Non-native Species Information Portal (GB-NNSIP).....	12
3.1 Species register.....	13
Methods of data collation.....	14
Information contained within the species register.....	14
Defining non-native species within the species register.....	17
Pathways of arrival.....	17
Transfer of data from designated experts to the species register.....	19
Limitations of the species register.....	19
3.2 Factsheets.....	20
Selection of species for factsheets and methods for collation of information.....	20
Collation of the factsheet information from the designated experts.....	20
4. Summarising the information within the species register.....	21
4.1 Summary of information within the Species Register.....	22
Totals in major categories.....	22
Species status - categories of non-nativeness.....	23
Established non-native species within broad environmental categories.....	25
Established non-native species within taxonomic groups.....	26
Established non-native species within functional groups.....	29
Date of introduction of established non-native species.....	30
Geographic origin of established non-native species.....	31
Pathways of introduction for established non-native species.....	34
Ecological and human impacts of established non-native species.....	36
Pathways and negative impacts of established non-native species.....	36
5. System for on-line recording of alert species including the outreach component “Recording Invasive Species Counts” (RISC).....	39
6. Key recommendations.....	43
7. Bibliography.....	44
Appendix 1. Glossary.....	77
Appendix 2. Summary of participation by experts, nominated by volunteer zoological scheme and society organisers (and other relevant bodies), to complete the species register entries associated with different informal (species) groups. The environment relates to the main environment of the informal group: terrestrial (T), terrestrial-freshwater (TF) or freshwater (F). The number of non-native species for each informal group is listed alongside the scheme, society or other organisation (and a named individual) contributing information.	80

Appendix 3. List of species selected for factsheets within the GB-NNSIP (294 species)..... 83
Appendix 4. Micro-organisms excluded from the project (250 species) 91
Appendix 5. Species lists, within broad groups, designated as having a negative ecological or human impact. Region of origin and introduction pathway are also provided. 97
Appendix 6. Species designated as having an unknown impact. 105
Appendix 7. Non-native species which are known to have arrived since the onset of the project or have been highlighted by scheme experts as missing from the species register 108

Summary

Here we report on a three-year Defra-funded study to enhance the ability to detect and report non-native species in GB. The specific aims of the project were:

- to document the distribution of non-native species in Great Britain;
- to assess whether there are changes in the rates of these processes;
- to document the impact of non-native species introductions to Great Britain; and
- to make this information available to key data users via web-based media.

The collation of data was coordinated by the Centre for Ecology & Hydrology (CEH) in collaboration with the British Trust for Ornithology, the Marine Biological Association and the Non-Native Species Secretariat. Additional experts, nominated by the Biological Records Centre (within CEH) volunteer schemes and societies, provided further information on many of the species. Of particular note was the involvement of the Botanical Society of the British Isles who provided information for 1873 species.

The project involved populating a database of non-native species (hereafter called “the species register”) with additional information provided in the form of factsheets for 300 of the species. Both these components are delivered through the GB Non-Native Species Information Portal (GB-NNSIP) hosted by the Non-Native Species Secretariat. An on-line system (alongside an e-mail account) for rapid-reporting capability, whereby particularly important new arrivals (so called “alert species”) can be immediately notified to the relevant bodies, was developed as an integral component of the GB-NNSIP. This was linked to an additional on-line recording site called Recording Invasive Species Counts (RISC), primarily designed for public participation.

There are 3758 species included within the species register. Information has been completed for 3204 species which constitutes 85.3% of the total list. The 554 species with incomplete information include 239 insects, 222 fungi (26 other microorganisms), 40 non-insect invertebrates, 24 lower plants, 2 higher plants and 1 alga. Information for the majority of these will be completed by early 2012.

The majority of species (2482 species) within the species register are categorised simply as non-native, that is they are species introduced by human action outside their natural past or present distribution. However, there are a number of other categories of non-nativeness which are relevant to the species register: for the purposes of this report all categories of non-nativeness are included but those designated with certainty as “unknown” (9.7%) are excluded. Therefore, there is a total of 2889 non-native species within the species register with completed information.

Each species occupies a row within the database with information on aspects of the species’ biology such as environment (marine, freshwater, terrestrial etc), functional type (predator, parasite etc), habitats occupied in the invaded range (using EUNIS classification), invasion pathways, establishment status in GB and impacts.

The data compiled within this project are indicative of current knowledge collated by many recognised experts. However, a database such as the species register will be incomplete both because of the number of new species arriving within GB annually and the species which remain cryptic particularly for groups considered to be difficult, such as parasites. Additionally it is difficult to determine the status of some species with respect to whether they are native or non-native, established and the impact that they may have within the invaded range. The involvement of so many experts has undoubtedly minimised the number of omissions and errors.

There are 1875 established non-native species in GB in total. The majority are higher plants (1377 species). Insects are the next most numerous group (278 species) followed by non-insect invertebrates (141 species), vertebrates (50 species), lower plants (25 species) and four other species. It should be noted that the information for 26 established non-native species within the species register is incomplete and so the summaries within this report are based on the 1849 established non-native species for which the information is complete.

Most (1684 species) of the documented established non-native species are found within the terrestrial environment of which higher plants (1350 species) are the largest group within the terrestrial environment. Within the freshwater environment higher plants (23 species) and non-insect invertebrates (23 species) dominate the established species. Non-insect invertebrates (50 species) are also the most numerous of the established non-native species within the marine environment.

There has been a dramatic increase over time in the number of non-native species arriving in Britain and those becoming established. 528 species arrived during 1950-1999 compared to 417 species during 1900-1949 and 250 species during 1850-1899. The majority are not invasive but those that do cause ecological &/or socio-economic impacts generally become permanent burdens if they become established. There have been at least a further 125 new arrivals since 2000 but only 40 of these are known to have established for many of the others information on establishment status is not yet available. Most of the non-native species that are established within Britain originate from Europe. However, in recent decades the rate of new arrivals originating from Europe is slowing and temperate Asia and North America are both becoming major contributors to the non-native fauna and flora of GB.

The majority (1024 species) of non-native species arriving and establishing in terrestrial environments do so as ornamental introductions. The dominant pathways for non-native species in freshwater environments are ornamental (21 species) and aquaculture (23 species). In the marine environment the arrival pathway for many non-native species is unknown but stowaways (40 species) and aquaculture (19 species) are both dominant pathways.

From a preliminary assessment of the 1849 established non-native species in GB, 282 species were found to have either a negative ecological or human impact. 147 species (8.0% of the total number of established non-native species) are considered to have a negative ecological impact and 188 species (10.2% of the total number of established non-native species)


a negative human impact, with 53 species (2.9% of the total number of established non-native species) designated as having both a negative human and ecological impact. There are 108 established non-native plants and 173 established non-native animals (and one other species *Coscinodiscus wailesii*, a marine alga) considered to have a negative ecological or human impact.

A number of key recommendations are provided but particular attention should be given to further enhancement of flow of non-native species distribution data from the many (and diverse) data providers. The on-line systems developed within this project should facilitate data flow but additional development will be required for a comprehensive approach.


GB Non-Native Species Report Card 2011

Confidence assessment: High confidence, medium confidence and low confidence

- More than 2889 non-native species
- 1875 established (self-sustaining populations) non-native species comprising 1402 established non-native plants and 469 established non-native animals and 4 other species
- Total of 7 non-native species are known to have been eradicated from GB
- 282 established non-native species have been designated as having a negative ecological or human impact: 173 established non-native animals, 108 established non-native plants and 1 other species




Establishment status of the species within the GB-NNSIP




Number of established non-native species and the number that are designated as having a negative ecological or human impact against date of first arrival

There has been a dramatic increase in the number of species arriving and those becoming established over the last 400 years and there is no indication of this trend slowing. Indeed over the last 200 years there have been 6.8 additional non-native species arriving and establishing per year in contrast to 0.8 additional non-native species arriving and establishing per year from 1600-1799. The number of established non-native species designated as having a negative ecological or human impact is also increasing with 1.2 of the new species per year causing an impact.



Number of established non-native species originating from different regions against date of first arrival

Most of the non-native species that are established within GB originate from Europe. However, in recent decades the rate of new arrivals originating from Europe is slowing and temperate Asia and North America are becoming both major contributors to the non-native fauna and flora of GB.



Number of established non-native species arriving through different pathways against date of first arrival

Most established non-native species in GB have arrived for ornamental purposes usually as garden plants but also for landscape planting and through the introduction of exotic animals. There has been a dramatic increase in species arriving through this pathway since 1800. There are an increasing number of non-native species for which the pathway of arrival is unknown.

In terrestrial environments 1024 of the established non-native species arrived through ornamental introductions. The dominant pathways for non-native species in freshwater environments are both ornamental (21 species) and aquaculture (23 species). In the marine environment the arrival pathway for many non-native species is unknown but stowaways (40 species) and aquaculture (19 species) are both dominant pathways. Across all environments, biological control, forestry and animal husbandry are very minor pathways with only one, 11 and 13 species arriving and establishing through these methods respectively.

Footnote: The data compiled within this project are indicative of current knowledge collated by many recognised experts. However, a database such as the species register will be incomplete both because of the number of new species arriving within GB annually and the species which remain cryptic particularly for groups considered to be difficult, such as parasites. Additionally it is difficult to determine the status of some species with respect to whether they are native or non-native, established and the impact that they may have within the invaded range. The involvement of so many experts has undoubtedly minimised the number of omissions and errors.

1. Introduction

The Millennium Ecosystem Assessment (Anonymous, 2005) designated invasive non-native (=alien) species, alongside climate change, habitat destruction, pollution and overexploitation, as one of the main drivers of biodiversity loss. Over the last century there has been a dramatic increase in the movement of non-native species around the world, as a consequence of international trade and travel (Hulme *et al.*, 2009). The Convention on Biological Diversity (CBD) suggests a three-stage hierarchical approach to invasive non-native species: prevention, detection/surveillance and rapid response, control and eradication. This approach has been followed in the Invasive Non-Native Species Framework Strategy for Great Britain (Defra, 2008), which outlined nine specific aims including the requirement for improved detection and monitoring capabilities coupled with an effective decision-making framework. Maintaining a list of non-native species within Britain is essential for underpinning decision-making concerning control, mitigation and eradication of invasive non-native species.

An audit, published in 2005, of non-native species in England concluded that there were 2721 non-native species in England (Hill *et al.*, 2005). The total for Britain is estimated to be in excess of 3500 species (Hill *et al.*, 2009). The audit represented a major step towards fulfilling this requirement. Here we report on a three-year Defra-funded study to enhance the ability to detect and report invasive non-native species in GB.

The GB-Non-Native Species Information Portal (GB-NNSIP) has been developed, with funding from Defra, by the Centre for Ecology & Hydrology (CEH) in collaboration with the British Trust for Ornithology, the Marine Biological Association, Non-Native Species Secretariat and the volunteer schemes and societies. The main purpose of the GB-NNSIP is to provide a central point of information about non-native species in Great Britain including: origins, ecology and impacts. It consists of a number of components including an inventory of non-native species called the “species register”, 300 species factsheets, an on-line system for reporting so called “alert species” (species for which the Non-Native Species Programme Board has recommended particular attention in terms of surveillance and reporting) and information to encourage wider participation particularly on-line recording of non-native species through the project Recording Invasive Species Counts (RISC) led by the NBN Trust and CEH. The RISC project has been developed to increase participation in recording non-native species (NNS) and to improve awareness of non-native species.

This report summarises the work, to date, on the ongoing development of the GB-NNSIP, RISC and the on-line Alert system.

2. Aims

The project aims to enhance our ability to detect and report invasive non-native species, and thereby to enable more effective decision-making. The results will be used by the GB Non-native Species Programme Board to determine how its information needs relating to invasive species populations and distributions can be met in the long term.

Specifically, the aims of the project are:

1. to document the arrival and spread of non-native species in Great Britain;
2. to assess whether there are changes in the rates of these processes;
3. to document the impact of non-native species introductions to Great Britain; and
4. to make this information available to key data users via web-based media.

In pursuit of the overall aims, we recognize the following approaches, specified in the invitation to tender:

1. Collate existing data on the introduction, distribution and impact of non-native species.
2. Analyse trends in the rate of non-native species introduction, spread and impact.
3. Summarise and report significant changes in the distribution or abundance of non-native species.
4. Establish a system of alerts for the interception of new introductions.
5. Identify data gaps and instigate additional biological recording.
6. Horizon scan for emerging threats.
7. Compile species profiles for key invasive species.
8. Make species profiles available on web-pages hosted by the Non-native Species Secretariat, alongside information on interceptions, introductions and distribution of non-native species.
9. Assess the scope for participation by the general public in the process of data collection for a number of species.

3. GB-Non-native Species Information Portal (GB-NNSIP)

The GB-NNSIP is an on-line information system involving a network of organisations engaged in monitoring and surveillance of non-native species (Figure 1a and 1b). Key components of the GB-NNSIP are the list of non-native species and associated summary information (database called the “species register”) and nearly 300 factsheets providing detailed information for a selection of species.




Figure 1a. Interactions between the GB-NNSIP and the wider community engaged in monitoring and surveillance of non-native species. Distributional data are collated from various organisations and bodies (statutory bodies, Local Records Centres and national schemes and societies including project collaborators MBA and BTO) through the NBN Gateway. Other information on non-native species is collated in the species register within the Biological Records Centre (Centre for Ecology & Hydrology). Data from the species register and the NBN Gateway are delivered to the GB-NNSIP and from here they can be exchanged with European (such as DAISIE) and Global initiatives such as the Global Invasive Species Database (within ISSG) and Global Invasive Species Information Network (GISIN).



Figure 1b. Structure of the GB-NNSIP illustrating links between databases and web services which provide information to the GB-NNSIP within the Non-Native Species Secretariat (NNS) website.

3.1 Species register

The species register is the basic list of non-native species, together with supporting information such as habitat, country of origin, arrival pathway, ecofunctional type, establishment status in England, Scotland and Wales, first record, human impact and ecological impact. The species register is dynamically linked to the NBN Gateway (<http://www.searchnbn.net/>) which provides maps of the distribution of the non-native species within Britain. It also contains summary information on aspects of the species biology such as environment (marine, freshwater, terrestrial etc), functional type (predator, parasite etc), habitats occupied in the invaded range (using EUNIS classification: <http://eunis.eea.europa.eu/about.jsp>), invasion pathways, establishment status in Britain and impacts.

The non-native species list compiled by Hill *et al.* (2005) formed the basis of the species register with additional species extracted from Delivering Alien Species Inventory in Europe (DAISIE). This list was transferred to an Access database called the “species register”. The list excluded pests of stored crops, human parasites and pests of human habitation unless they were thought likely to be found in the wild). Microorganisms (with the exception of a small number of marine phytoplankton) were also generally not included. The greatest deficiency in the species register was freshwater non-native species but a list of British freshwater non-native species compiled by Philip Lambdon (Centre for Ecology & Hydrology) was added. The list is summarised in Table 1.

Table 1. Numbers of non-native species in environmental groups and species lists derived from the feasibility study. Columns correspond to species lists, or (the last column) to species added but that were not found in any of the lists (taken from Hill *et al.*, 2008).

TAXON_GROUP	All	DAISIE	NBN	BRC	DAISIE not in NBN or BRC	EN audit only	No list
Marine	150	135	97	36	43	2	3
Freshwater	89	4	76	45	1	8	0
Terrestrial	3239	2853	2579	2570	243	33	82
Fungi and microbes	209	0	122	121	0	50	0
Total	3687	2992	2874	2772	287	93	85

Methods of data collation

The project was led and managed initially by Mark Hill (Biological Records Centre (BRC) within Centre for Ecology & Hydrology) and by Helen Roy from July 2010 (also BRC). The main project partners were the British Trust for Ornithology (BTO), Marine Biological Association and the Botanical Society of the British Isles (BSBI); however, the volunteer zoological schemes and societies (hosted through the Biological Records Centre) nominated experts to participate in the project (Appendix 2). BRC supplied an excel template of the species register along with a set of guidelines containing coding information for each column to the numerous experts.

Information contained within the species register

Each species within the species register is assigned a BRC concept code which provides a reference within the BRC database that remains with the species despite changes to taxonomic nomenclature. Where NBN Gateway codes (Taxon Version Keys) were available these were also included within the species register enabling distribution data to be retrieved from the NBN Gateway.

Information was collated for each species under the following categories: informal grouping (such as higher plant, insect etc), phylum, environment (terrestrial, freshwater and marine etc), functional group (predator, herbivore etc), non-native status (non-native, colonist etc), native

range (using Taxonomic Database Working Group (TDWG <http://www.tdwg.org/standards/104/>) categories or country-level data where TDWG provided insufficient resolution), habitats occupied within GB (using Eunis (<http://eunis.eea.europa.eu/about.jsp>) categories), major pathways (stowaway etc), method of introduction (accidental, intentional etc), establishment status in GB, England, Scotland and Wales, ecological and human impact, place and year of first record in GB and key references (Table 2). TDWG and EUNIS are international organisations promoting standards in biodiversity data collection/storage and, therefore, the use of their categorical systems increases the probability of compatibility between GB-NNSIP and other databases such as DAISIE.

Descriptions of the codes used within the species register are summarised in Table 2 with further detail provided for status of non-native species and pathways of introduction in Tables 3 and 4.

Table 2. Main fields within the species register with comments provided to domain experts (additional fields are required to acknowledge the author, date updated and key references).

Field	Notes	Codes
idspecies	DAISIE code	
TVK	the NBN code	
Concept	i.e. the new BRC number	
Scientific name	Binomial aligning with the NBN	
English_name	Dictionary	
Informal group	Preferred name	
Phylum	Informal group	
Env	Environment	Ph = phytoplankton, Alg = algae, LP = land plant, Pr = predator, H = herbivores, O = omnivore, Par = parasite, D = detritivore, F = filter-feeder
Fnct	Functional type	
Factsheet	Factsheet supplied to GB-NNSIP	Supplied by BRC
Notes on name, ecofunction or native range		
Status	Native status	Details within Table 3
Nat	Native range	TDWG categories or country-level data provided (up to 4 regions can be specified in fields Nat_1, Nat_2, Nat_3 and Nat_4)
Hab	Habitat occupied in GB	EUNIS categories provided (up to 4 habitats can be specified in fields Hab_1, Hab_2, Hab_3 and Hab_4)
Path	Invasion pathway	Arrival pathways (pathway to Europe is provided if natural spread to GB from Europe). Details within Table 4 (up to 2 pathways can be specified in fields path1 and path2). GISIN terminology of "method" is used (up to 2 methods can be specified in fields Meth1 and Meth2)
Meth1	Human intention	
GB	Establishment status - GB	
EN	Establishment status - England	
SC	Establishment status - Scotland	
WA	Establishment status - Wales	
Ec_imp	Ecological impact	Based on expert opinion
Hum_imp	Human impact	Based on expert opinion
Place where first found		Place where first found
VC	Vice-county	
current_distribution		Link to NBN Gateway distribution data
Year	Date first reported in the wild	

Defining non-native species within the species register

The CBD definition of a non-native species is clear, however within the species register there are a number of sub-categories employed (Table 3). A few of the species appearing in the species register are considered native in GB but appear as artefacts from previous audits. These are not included within this report but remain in the species register with comments to ensure clarity in the future.

Table 3. Status of non-native species including codes used in the species register, terms and notes based on Hill *et al.* (2005). Species designated as native (N) or Unknown (U) are not included within this report.

Code	Term	Notes
NN	Non-native	Species known to be non-native, including non-native species on the horizon but not yet in GB
NN-N	Non-native/Native	Probably non-native
N-NN	Native/Non-native	Probably native
C	Colonist	Colonist 2000 or later, probably or certainly by natural spread
D	Dependent	With trophic dependence on cultivated or non-native species, arrived possibly naturally before 2000
DNN	Dependent non-native species	With trophic dependence on cultivated or non-native species
DC	Dependent colonist	With trophic dependence on cultivated or non-native species, arrived naturally 2000 or later
HGB	Hybrid GB	GB hybrid origin with non-native parent
NN-S	NN subspecies	Native species that have non-native subspecies within GB
U	Unknown	Origins unknown

Pathways of arrival

The pathway by which a non-native species arrives within a country is particularly relevant to implementing prevention measures. For a minority of species the exact pathway of arrival is unknown but in most cases there is evidence for a specific introduction pathway. The species register uses a detailed set of codes for defining pathways (Table 4) but for the purposes of summarising the data these have been collapsed into broad pathway categories (Table 5).

Table 4. Introduction pathways for non-native species including codes used in the species register, terms and notes based on Hill *et al.* (2005).

Code	Term	Notes
Agr	Agriculture	Not a contaminant e.g. an intended introduction such as a crop or feral goat
AgrS	Agricultural seed contaminant or pest	Such as corncockle
Aq	Aquaculture	Such as crayfish but also aquarium plants but not pond plants which are designated as ornamental.
Bio	Biocontrol	Introduction as a biological control agent or arrival from another region in which it was introduced as a biological control agent
FLG	Fur/lab/greenhouse	Similar to agriculture but with containment e.g. mink but not ornamental plant trade which is Orn
For	Forestry	Not a contaminant e.g. an intended introduction
HF	Hunting/fishing	Such as pheasant, rainbow trout
Hyb	Hybridization	
L	Landscape	Planted into landscape such as pheasant cover or releases of muntjac
Med	Medicinal	Only plants introduced for medicinal reasons
Nat	Natural	Organisms dependent on a non-native species such as rhododendron leafhopper but also natural colonisation from a previously invaded range such as harlequin ladybird
Orn	Ornamental	Garden plant or trade in garden plants, zoo animal, pet
OrnS	Ornamental seed contaminant or component	
P	Produce	Such as harlequin ladybird on flowers, vegetables etc
PS	Seed produce contaminant or component	Such as bird seed contaminant
RM	Raw material	Such as Citrus longhorn beetle on timber
SA	Stowaway – air	Mainly animals
SL	Stowaway – land	Mainly animals
SW	Stowaway – water	Ballast water and hull fouling – mainly animals such as killer shrimp but also algae
U	Unknown	

For the purposes of summarising information in this report the pathways were grouped (Table 5).

Table 5. Grouping of specific pathways into broad pathways

Specific pathway	Broad pathway
Agr – Agriculture	Agriculture
AgrS – Agricultural seed contaminant or pest	Contaminant
Aq – Aquaculture	Aquaculture
Bio – Biocontrol	Biocontrol
FLG – Fur/lab/greenhouse (but not ornamental plant trade which is Orn)	Animal husbandry
For – Forestry	Forestry
HF – Hunting/fishing	Animal husbandry
Hyb – Hybridization	Hybridisation
L – Landscape	Ornamental
Med – Medicinal	Agricultural
Nat – Natural (dependent organisms only)	Natural – Non-Native Dependent
Orn – Ornamental, i.e. garden plant or trade in garden plants, zoo animal, pet	Ornamental
OrnS – Ornamental seed contaminant or component	Contaminant
P – Produce	Contaminant
PS – Seed produce contaminant or component	Contaminant
RM – Raw material	Stowaway
SA – Stowaway – air	Stowaway
SL – Stowaway – land	Stowaway
SM – Stowaway – marine	Stowaway
U – Unknown	Unknown

Transfer of data from designated experts to the species register

Completed templates were collated in excel and uploaded into an Access database at BRC (Figure 1b). A copy of the updated species register entry was then exported and used to update the species register table within the MySQL GBNN database schema on the BRC webserver (WLBRCWEB1). A webservice on the BRC webserver allows for information from this database to be displayed on the Non-Native Species Secretariat website where a flag to display is present. Maps displaying the native range of a species are delivered via an ARCIMS map service on the BRC webserver using information stored within the GBNN database schema.

Limitations of the species register

The data compiled within this project are indicative of current knowledge collated by many recognised experts. However, a database such as the species register will be incomplete both because of the number of new species arriving within GB annually and the species which remain cryptic particularly for groups considered to be difficult, such as parasites. Additionally it is difficult to determine the status of some species with respect to whether they are native or non-native. Many ancient introductions are likely to have been omitted. It is also difficult to

determine whether or not a species is self-sustaining (established). Categorising the impact of species is extremely difficult and, without thorough and systematic assessment of risks, remains subjective for many species. The involvement of so many experts has undoubtedly minimised the number of omissions and errors. Additionally the information for all the “factsheet species” (refer to section 3.2) has been subject to additional external review and conforms to the information provided within the risk assessments and identification sheets available through the Non-Native Species Secretariat website. Finally, it should be noted that some categories for some species were left blank by the designated experts and, therefore, the numbers do not always tally to an expected total.

3.2 Factsheets

The species register contains basic information on a large proportion of the non-native species found in GB. However, detailed information is provided for 294 non-native species (Appendix 3).

Selection of species for factsheets and methods for collation of information

The information was collated by the experts as described above (also see Appendix 2) and the species were selected on the basis of the following criteria:

- DAISIE 100 worst (Europe)
- Horizon Scanning list (Parrott *et al.*, 2009)
- Factsheet prepared for the GB-NNSIP feasibility study (Hill *et al.*, 2008)
- Illustrated in the [ID sheets](#) of the GB Non-native Species Secretariat
- Species subject to legal bans or controls
- Species selected for full risk assessment by the Non-Native Species Secretariat
- Species with strong environmental or other impacts
- Suggestions from GB-NNSIP steering group, project team and others

Collation of the factsheet information from the designated experts

The BRC webserver hosts the database driven website content management system which can be found at http://www.brc.ac.uk/gbnn_admin/. A number of species experts were asked to complete the factsheets and allocated a personal log-in for the factsheet editor allowing them to create and edit the factsheets within a second database schema, drupal_gbnn, on the BRCwebserver. Once complete, BRC staff (Helen Roy, Björn Beckmann and Steph Rorke) checked and edited the factsheets within the factsheet editor ultimately publishing them to the GBNN database schema. At this stage the factsheets were exposed through a webservice to the Non-Native Species Secretariat website (Figure 2). The factsheets can be accessed via this website as pdf files and the generation of the pdf occurs on the Non-Native Species Secretariat webserver. Additionally, a number of images displayed on the website are also stored on this webserver.

It should be noted that Defra will be checking the information provided on control methods and additionally standardising the information on legislation.

Harlequin Ladybird, *Harmonia axyridis*

Overview

Short description of *Harmonia axyridis*, Harlequin Ladybird
Highly variable in appearance. Adults, 5-8 mm in length, occur in three colour forms: yellow-orange to red with 0 to 21 black spots, black with two red spots and black with four red spots. Larvae are covered with branched spines and have orange sides.

Description of *Harmonia axyridis*, Harlequin Ladybird status in GB
The harlequin ladybird is well established in England and Wales and spreading rapidly. Local in Scotland, but likely to invade further.

Impacts summary: *Harmonia axyridis*, Harlequin Ladybird
They prey on some of our native insects and compete for food with others. In autumn and winter, large aggregations enter buildings and can become a nuisance. In other countries, they can be a pest of orchards and vineyards, but the magnitude of this problem in GB is not yet clear.

Habitat summary: *Harmonia axyridis*, Harlequin Ladybird
Harlequin ladybirds are still found mainly in towns but are spreading to hedgerows and woodland. They occur especially on lime and sycamore trees, which support large aphid populations in late summer.

Overview table

Environment:	Terrestrial
Species status:	Non-Native
Native range:	China, Japan, Korea, Democratic People's Republic of Korea, Republic of, Russian Federation
Invasion pathway:	Stowaway - land
Functional type:	Predator
Status in GB:	Established
Status in England:	Established
Status in Scotland:	Established
Status in Wales:	Established
Date of first record:	2004

Author's name:
Helen Roy

Last updated:
March 21st, 2011

Figure 2. Example factsheet on the GB-NNSIP within the GB Non-Native Species Secretariat website.

4. Summarising the information within the species register

The species register represents a resource of detailed information on non-native species. The use of defined codes for each attribute ensures there is considerable potential for detailed analysis. In this report we have summarised the species register in terms of the total numbers

of species but the majority of the summary tables include only the species designated as established in GB.

4.1 Summary of information within the Species Register

Totals in major categories

There are 3758 species included within the species register. Information has been completed for 3204 species which constitutes 85.3% of the total list. The 554 species with incomplete information include 239 insects, 222 fungi (26 other microorganisms), 40 non-insect invertebrates, 24 lower plants, 2 higher plants and 1 alga.

Species and establishment status of the 554 species with incomplete information

There are 250 microorganisms within the species register (Table 6). Micro-organisms (including Oomycete, Mycetozoa, fungi (Ascomycota, Basidiomycota and other fungi), Haplosporea and other micro-organisms) were not included within the project specification and are consequently not well represented within the species register (in terms of species lists and information provided). There are 222 species of fungi (Ascomycota, Basidiomycota and other fungi) listed within the register alongside 28 microorganisms from other taxa (Table 6 and Appendix 4). However, information has only been provided for two species of micro-organism.

The species status (non-nativeness) of many of the remaining 304 species with incomplete information has been designated as “unknown”. Indeed there are only 42 species for which the species status has been ascertained. Of these 31 are considered to be non-native, 4 non-native dependents, 6 are designated as native species and 1 is probably native. The establishment status is documented for the 35 species designated within one of the non-native species categories and 26 are considered to be established.

Table 6. Number of microbial species (total = 250) in the species register within broad taxonomic groups and the number of species with updated information.

	Number of species in species register	Number with updated information
Ascomycota	125	0
Basidiomycota	91	0
Fungi (other)	6	0
Haplosporea	1	1
Mycetozoa	1	0
Oomycete	14	0
Other Microorganisms	12	1

Species status - categories of non-nativeness

The majority of species (2482 species) within the species register are categorised simply as non-native (Table 7). However, there are a number of other categories of non-nativeness which are relevant to the species register. Approximately 7.0% of non-native species in GB are the consequence of hybridisation (Hybrid GB). It has been difficult to assign 4.4% of the species in the species register to a non-native category with complete certainty (non-native/native and native/non-native). However, for the purposes of this report all categories of non-nativeness (outlined in Table 7) are included but those designated with certainty as “unknown” (9.7%) are excluded. Therefore, there is a total of 2889 non-native species within the species register with completed information.

Table 7. Numbers of species (established and non-established but excluding microorganisms) within different environments (T=Terrestrial, F=Freshwater, M=Marine, M(E)=Marine (estuarine), TF=Terrestrial-Freshwater, MF=Marine-Freshwater, TM=Terrestrial-Marine) and subdivided into different categories of non-nativeness (non-native, hybrid within GB, probably non-native, probably native, dependent, new arrival, dependent colonist, dependent non-native and unknown).

	T	F	M	M (E)	TF	MF	TM	Total
Non-native	2194	102	98	1	69	13	5	2482
Hybrid GB	222						1	223
Probably non-native	55		3		8			66
Probably native	43		5		25	1	1	75
Dependent	27							27
New arrival	7		1					8
Dependent colonist	4							4
Dependent non-native	3		1					4
Unknown	263	17	30		1			311
Total	2818	119	138	1	103	14	7	3200

Only a fraction of new arrivals within a country will establish as self-sustaining populations. Of the non-native species, for which we have complete information within the species register, 1795 species are considered to be established in the wild in GB (Table 8). A further 54 non-native species are established indoors. A further 26 species (13 insects, 5 lower plants, 4 mites, 3 platyhelminths, 1 higher plant hybrid) are established non-native species but currently the information within the species register for these is incomplete. Therefore, in total 1875 of the documented species are established. In all subsequent summary tables only established non-native species with complete information (1849 species) will be included unless otherwise stated.

England has more established non-native species than either Scotland or Wales (Table 8). A small proportion (1.5-2.1%) of all non-native species are currently considered extinct (no longer

established) in GB. A total of 7 non-native species have been contained (only present in quarantine or other contained facilities) or exterminated in GB over the last 100 years (Table 9). Musk rat (*Ondatra zibethicus*) and tobacco whitefly (*Bemisia tabaci*) are the only species eradicated from all constituent countries within GB. A number of non-native species listed in the GB-NNSIP are absent from GB. These may enter GB in the future and should be considered 'horizon scanning species'. Similar lists are provided for each individual country.

Table 8. Establishment status of the non-native species in GB, England, Scotland and Wales. Native is included as a category because a species can be non-native in one country within GB but native in another.

	GB	England	Scotland	Wales
Established	1795	1737	1005	1013
Indoors	54	29	14	13
Not established	850	818	605	471
Native	11	10	2	3
Extinct	61	51	49	47
Eradicated	7	6	1	1
Absent	29	63	773	915
Unknown	74	64	89	86

Table 9. Non-native species eradicated.

Scientific name	English name	Notes
<i>Hystrix brachyura</i>	Hodgson's Porcupine	Small colony was briefly established in Devon but successfully eradicated.
<i>Leptinotarsa decemlineata</i>	Colorado Beetle	Probably introduced with potato (or other) shipments, though possibly an immigrant from continental Europe.
<i>Mesocricetus auratus</i>	Golden Hamster	First record from Bath in 1958.
<i>Misgurnus mizolepis</i>	Chinese Weatherfish	A reproducing loach population in a pond in Essex (since eradicated) was found to be <i>M. mizolepis</i> .
<i>Myocastor coypus</i>	Coypu	First brought to England to stock fur farms in 1929. Escapes occurred mainly in the 1930s.
<i>Ondatra zibethicus</i>	Musk Rat	Escaped from fur farms
<i>Bemisia tabaci</i>	Tobacco Whitefly	First reported on <i>Poinsettia</i> cuttings, which have since been main source of interceptions / outbreaks.

The majority (1377 species) of established non-native species are higher plants (Table 10). Insects are the next most numerous group (278 species) followed by non-insect invertebrates (141 species), vertebrates (50 species), and lower plants (25 species).

Table 10. Number of established non-native species within broad taxonomic groups (includes non-native species for which only establishment status is completed within the species register and also the number for which the species registry includes complete information. The latter being a subset of the former).

Broad group	Number of established non-native species	Number of species with complete information in the species register
Higher plants	1377	1376
Lower plants	25	20
Insects	278	265
Non-insect invertebrates	141	134
Vertebrates	50	50
Other	4	4
Total	1875	1849

Established non-native species within broad environmental categories

The majority of established non-native species are terrestrial (Table 11). Indeed there are only 60 freshwater species and 74 marine species. Higher plants are the largest group of established non-native species within the terrestrial environment (Table 11). Within the freshwater environment higher plants and non-insect invertebrates dominate. Non-insect invertebrates also form a high proportion of the established non-native species within the marine environment.

Table 11. Total number of established non-native species in broad taxonomic groups (includes only species designated as non-native) for different environments (T=Terrestrial, F=Freshwater, M=Marine, M(E)=Marine (estuarine), TF=Terrestrial-Freshwater, MF=Marine-Freshwater, T-M=Terrestrial-Marine).

	T	F	M	M (E)	TF	MF	TM	Total
Higher plants	1350	23	0	0	0	0	3	1376
Lower plants	0	0	20	0	0	0	0	20
Insects	255	8	0	0	0	0	2	265
Non-insect invertebrates	52	23	50	1	0	8	0	134
Vertebrates	27	6	0	0	16	1	0	50
Other	0	0	4	0	0	0	0	4
Total	1684	60	74	1	16	9	5	1849

Established non-native species within taxonomic groups

The established non-native species in GB are taxonomically diverse (Table 12). Flowering plants represent the most numerous taxonomic category of established non-native species (1336 species). Birds (15 species) and terrestrial mammals (17 species) are the most species-rich vertebrate taxa. Insects are the most numerous of all the established non-native species of invertebrate with Hemiptera (bugs) and Coleoptera (beetles) dominating. There are a further 239 species of insect to update in the species register and a high proportion of these are beetles. In marine environments algae (20 species) and crustacea (15 species) are the most numerous of all established non-native species. There is also a reasonable number of established non-native crustacean (12 species) within freshwater environments.

Examples of recent new arrivals

There have been 125 new arrivals since 2000 (only 40 of these are known to have established and for many of the others information on establishment status is not yet available). The list of non-native species arriving over the last decade has been dominated by insects and other invertebrates (78 species). A number of these species have received considerable interest from the media and consequently a high profile primarily because of their potentially negative impacts on the environment.

Harmonia axyridis, harlequin ladybird, is native to Asia. The harlequin ladybird was first recorded in England (Essex) in 2004, however assessment of the Rothamsted Aphid Trap archives revealed a specimen from 2003. The spread of this generalist predator has been monitored by people who send their sightings into an on-line survey (www.ladybird-survey.org). This species is known to have spread at 100 km yr⁻¹ and is one of the most abundant species of ladybird through England and Wales.

Dikerogammarus villosus, killer shrimp, was discovered in England (Grafham Water, Cambridgeshire) in 2010. This species, native to the Ponto-Caspian region, has the potential to disrupt freshwater communities and is considered to be an invasive non-native species. There have only been two further records of this species both in Wales (Cardiff Bay and Eglwys Nunydd, Wales). Rapid response to *D. villosus*, including the launch of a campaign (Stop the spread – check, clean, dry) and an on-line system for recording sightings of this species, should provide effective surveillance and rapid response.

In 2008 *Didemnum vexillum*, carpet sea-squirt (native to the north-west Pacific), was recorded in Devon. This species has now been recorded from nine marinas in GB (North Wales, the Clyde, Devon and the Solent). Rapid growth of this species can result in the smothering of habitat and disruption of resident communities.

Pimepales promelas, fathead minnow, is a species which has been reported sporadically across England since 2003 following release from private collections. It is native to North America and Mexico. There are no known impacts of this species in GB but it is a predatory fish which could affect resident communities and additionally it is associated with the spread of enteric red-mouth disease (ERM) to wild and cultured trout and eel in continental Europe.

Two species of crayfish have arrived and established over the last ten years: *Orconectes limosus*, spinycheek crayfish (2001), and *Orconectes virilis*, virile crayfish (2004). Both species are native to North America. The arrival of these two crayfish brings the total number of

established non-native crayfish in GB to seven. *Pacifastacus leniusculus*, signal crayfish, which was first recorded in GB in 1975 is a major threat to the native *Austropotamobius pallipes*, white-clawed crayfish, and is causing declines in diversity and richness of aquatic communities.

Lemna turionifera, red duckweed, was first recorded in 2007 in Stoborough, England. It is spreading rapidly through England and Wales.

Table 12. Numbers of established non-native species in taxonomic groups within different environments (T=Terrestrial, F=Freshwater, M=Marine, M(E)=Marine (estuarine), TF=Terrestrial-Freshwater, TM=Terrestrial-Marine, MF=Marine-Freshwater).

	T	F	M	M (E)	TF	TM	MF	Total
Plants								
Alga			20					20
Clubmoss	1							1
Conifer	29							29
Diatom			4					4
Fern	8	1						9
flowering plant	1311	22				3		1336
Horsetail	1							1
Vertebrates								
Amphibian					8			8
Bird	7				8			15
bony fish (Actinopterygii)		6					1	7
Reptile	3							3
terrestrial mammal	17							17
Insects								
insect - beetle (Coleoptera)	95	1						96
insect - cockroach (Dictyoptera)	1							1
insect - hymenopteran	4							4
insect - moth	44	7						51
insect - stick insect (Phasmida)	3							3
insect - thrips (Thysanoptera)	1							1
insect - true bug (Hemiptera)	103							103
insect - true fly (Diptera)	4					2		6
Non-insect invertebrate								
Annelid		2	11	1				14
Bryozoans			4				1	5
Centipede	3							3
coelenterate (=cnidarian)			2				1	3
Crustacean	2	12	15				3	32
Entoproct			1					1
flatworm (Turbellaria)		1						1
Millipede	2							2
Mollusc	23	8	10				2	43
roundworm (Nematoda)	1						1	2
sea spider (Pycnogonida)			1					1
spider (Araneae)	20							20
springtail (Collembola)	1							1
tunicate (Urochordata)			6					6
Total	1684	60	74	1	16	5	9	1849

Established non-native herbivorous species are prevalent within terrestrial and freshwater systems but appear to be lacking in the marine environment (Table 13). Primary producers (alga and land plants) represent the majority of established non-native species, followed by primary consumers (filter-feeders, herbivores and omnivores) with secondary consumers (parasites and predators) occurring in low proportions in all systems.

Table 13. Number of established species within different functional groups for different environments (T=Terrestrial, F=Freshwater, M=Marine, M(E)=Marine (estuarine), TF=Terrestrial-Freshwater, TM=Terrestrial-Marine, MF=Marine-Freshwater).

	T	F	M	M(E)	TF	TM	MF	Total
No data	9	2	1					12
Detritivore	6	1	3					10
Alga			25					25
Land plant	1350	23				3		1376
Filter-feeder		4	35	1			4	44
Herbivore	217	14			7	2		240
Omnivore	56	5	2		1		2	66
Parasite	1	5	2				1	9
Predator	45	6	6		8		2	67

Established non-native species within functional groups

Marine species are distributed reasonably evenly between littoral rocks, sediment and infralittoral rock habitats (Table 14). This could be an indication of ease of sampling and species could remain largely undiscovered in pelagic zones.

Terrestrial and freshwater non-native species also occupy diverse habitats (Table 14). More than 40% (821 species) of terrestrial established non-native species are found within grassland habitats; a consequence of the high number of plants which occur in ruderal tall-herb communities (categorised by EUNIS as grasslands) (Table 14). Woodlands and forest habitats have 123 established non-native species. Inland surface waters also have a high number of non-native species (140 species) spread across the different environments. Domestic habitats including cultivated gardens and parks also have a high number of non-native species (153 species), as do constructed, industrial habitats (158 species). This reflects the dominance of plants within the species register. Mires and bogs have a very low number of non-native species (7 species).

Table 14. Number of established non-native species occurring in different habitats (EUNIS categories) across different environments (T=Terrestrial, F=Freshwater, M=Marine, M(E)=Marine (estuarine), TF=Terrestrial-Freshwater, TM=Terrestrial-Marine, MF=Marine-Freshwater).

Habitat	T	F	M	M (E)	TF	TM	MF	Total
No data	13							13
Marine			35				3	38
Littoral rock & hard substrate			8					8
Littoral sediment	7		2			2	1	12
Infralittoral rock			19					19
Cirralittoral rock			1					1
Sublittoral sediment			2					2
Pelagic water column							1	1
Coastal habitats	196		3	1		2		202
Inland surface waters	67	53	1		16	1	2	140
Mires, bogs & fens	7							7
Grasslands etc	821							821
Heathland, hedgerow & scrub	81							81
Woodland & Forest	123							123
Inland unveg. or sparsely veg.	48							48
Domestic habitats	153							153
Constructed, industrial etc	158	7						165
Unknown	7							7
Estuarine and brackish habitats	3		3				2	8
Total								1849

Date of introduction of established non-native species

There has been a dramatic increase in the number of species arriving and those becoming established since 1800 and there is no indication of this trend slowing (Figure 3). The number of established non-native species designated as having a negative ecological or human impact is also increasing.




Figure 3. Number of established non-native species and the number that are designated as having a negative ecological or human impact against date of first record. It should be noted that a further 125 species have arrived (only 40 of these are known to have established) between 2000 and 2010 and 14 of these are considered to have a negative impact. There are 170 established non-native species for which there is no date within the species register.

Geographic origin of established non-native species

Most of the non-native species that are established within GB originate from Europe (Figure 4). However, temperate Asia and North America are both major contributors to the non-native fauna and flora of GB.




Figure 4. Number of established non-native species (animals and plants) in GB originating from different regions.

It is interesting to note that non-native species originating from Europe began to arrive and establish in high numbers from the 1700s, whereas non-native species originating from temperate Asia and North America have seen a rapid escalation since the 1800s (Figure 5). The peak in number of species arriving from Europe from 1500 to 1600 is almost certainly a consequence of an increase in human movement from 1500 onwards. It is intuitive that species in close proximity to Britain would arrive earlier than those from greater distances. The patterns in date of first arrival for non-species originating from different regions are entirely consistent with the increase in long-distance transport subsequent to the discovery of the Americas. It is apparent the rate of new species arriving from Europe to Britain is slowing particularly in contrast to the dramatic rate of new arrivals from temperate Asia.




Figure 5. Number of established non-native species originating from different regions and arriving in Britain against date of first record.

In the terrestrial environment most of the non-native species (822 species) within GB have originated from Europe (Table 15). This is dominated by the number of higher plants that have arrived to GB from Europe (666 species) but also the high number of insects arriving from Europe (107 species). In contrast North America is a major contributor of non-native species (29 species), particularly plants (13 species), to GB freshwater environments (Table 15) although a reasonably high number of invertebrates arrive in the freshwater environment from Europe and temperate Asia. Interestingly, within the marine environment the non-native species originate much more evenly from a number of regions with Asia (temperate), North America and the Pacific region dominating (Table 15).

Table 15. Number of established non-native species, within broad taxonomic groups, originating from different regions and environments.

	Africa	Asia- Temperate	Asia- Tropical	Australasia	Europe	North America	Pacific	South America	No data	Total
Terrestrial										
Higher plants	44	197	28	49	666	153		52	161	1350
Vertebrates	1	6	2	1	12	2			4	27
Insects	15	28	9	15	107	32		8	41	255
Other inverts	2		1	2	37	4		2	4	52
Total	62	231	40	67	822	190		62	210	1684
Freshwater										
Higher plants	2	2		1	2	13		3		23
Vertebrates						6				6
Insects		1	5			2				8
Invertebrates		4		1	7	8			3	23
Total	2	7	5	2	9	29		3	3	60
Marine										
Lower plants		9		2	1	1	5		2	20
Invertebrates	4	9		5	5	15	7		5	50
other		1					1		2	4
Total	4	19		7	6	16	13		9	74
Marine (estuarine)										
Invertebrates									1	1
Marine- Freshwater										
Vertebrates					1					1
Invertebrates		3			2	2			1	8
Total		3			3	2			1	9
Terrestrial- Freshwater										
Vertebrates	1	2			9	4				16
Terrestrial- Marine										
Higher plants						2			1	3
Insects						2				2
Total						4			1	5

Pathways of introduction for established non-native species

A very high proportion of the non-native species arriving and establishing in terrestrial environments do so as ornamental introductions (Figure 6, Table 16). The dominant pathways for non-native species in freshwater environments are ornamental and aquaculture (Table 16). In the marine environment the arrival pathway for many non-native species is unknown but stowaways and aquaculture are both dominant pathways (Table 16). Higher plants are the most numerous of all non-native species within terrestrial and freshwater environments and so it is not surprising that the ornamental pathway is the most common because many of the non-native plants arrive through horticulture. However, a high proportion of terrestrial animals have also arrived through this pathway particularly as exotic pets for domestic and zoological collections. There has been a dramatic increase in species arriving through the ornamental pathway since 1800. However, there are an increasing number of non-native species for which the pathway of arrival is unknown (20 species: 1850-1899, 42 species: 1900-1949 and 52 species: 1950-1999).




Figure 6. Number of established non-native species arriving in GB through different pathways against date of first record.

Table 16. Number of non-native species, within broad taxonomic groups, arriving by different pathways (Agr. = agricultural, Anim. husb. = animal husbandry, Aqua. = aquaculture, Bio. = biocontrol, Con. = contamination, For. = forestry, Hyb. = hybridisation, Natural-NND = Natural non-native dependent, Orn. = ornamental, Stow. = stowaway and U = Unknown) into different environments.

	No data	Agr.	Anim. husb.	Aqua.	Bio.	Con.	For.	Hyb.	Natural - NND	Orn.	Stow.	U
Terrestrial												
Higher plants		88	1			145	11	29		955	54	67
Vertebrates			8		1					14	4	
Insects	167	2	1			40			20	51	23	114
Invertebrates			2							4	2	71
Other												
Total	167	90	12		1	185	11	29	20	1024	83	252
Freshwater												
Higher plants				7						14		2
Vertebrates	6		1							6		
Insects				7						1		
Invertebrates	2			9		2					8	4
Total	8		1	23		2				21	8	6
Marine												
Lower plants				7							7	10
Invertebrates				12					1		32	12
Other											1	4
Total				19					1		40	26
M (estuarine)												
Invertebrates											1	
Total											1	
TF												
Vertebrates			1							15		
Total			1							15		
TM												
Higher plants								1		2		
Insects	2											
Total	2							1		2		
MF												
Invertebrates				3							4	
Total				3							4	

Ecological and human impacts of established non-native species

Of the 1849 established non-native species in GB, 282 species (15.3%) are designated as having some negative impact of which 147 species (8.0%) are considered to have a negative ecological impact and 188 species (10.2%) a negative human impact with 53 species (2.9%) designated as having both a negative human and ecological impact. There are 108 (7.7%) established non-native plants and 173 (39.2%) established non-native animals considered to have a negative ecological or human impact (Table 17). The majority of the human impacts are linked to plant health (pests and diseases). Higher plants dominate the list of established non-native species with ecological impacts (Appendix 5) however, a high proportion (48.0%) of the total number of established non-native vertebrates have an ecological impact (24 species). The impact for 106 established non-native species is designated as unknown (Appendix 6) and critical assessment of these species should be a high priority.

Table 17. Number of established non-native species, within broad taxonomic groups, designated as having a **negative ecological or human impact** arriving through each introduction pathway.

	Higher plants	Lower plants	Total plants	Insects	Invertebrates	Vertebrates	Total animals	Other	Total
Agriculture	4		4	1			1		5
Animal husbandry			0			7	7		7
Aquaculture	3	3	6		11		11		17
Biocontrol			0			1	1		1
Contaminant	10		10	38			38		48
Forestry	3		3				0		3
Hybridisation	1		1				0		1
Ornamental	79		79	22	1	13	36		115
Stowaway	3	2	5	12	16	3	31		36
Unknown			0	18	9		27	1	28
No data			0	21			21		21
Total	103	5	108	112	37	24		1	282

Pathways and negative impacts of established non-native species

The ornamental pathway contributes the highest number of non-native species considered to have a negative ecological impact (Table 18). The majority of these are higher plants. A high number of non-native vertebrates arrive through the ornamental pathway and exert a negative ecological impact.

Table 18. Number of established non-native species, arriving by different pathways, within broad taxonomic groupings considered to have a **negative ecological impact**.

	Higher plants	Lower plants	Total plants	Insects	Invertebrates	Vertebrates	Total animals	Total
Agriculture	1		1					1
Animal husbandry						5	5	5
Aquaculture	3	3	6		10		10	16
Biocontrol						1	1	1
Contaminant	1		1					1
Forestry	3		3					3
Hybridisation	1		1					1
Ornamental	76		76	4		12	16	92
Stowaway	2	2	4	3	10	3	16	20
Unknown				1	4		5	5
No data				2			2	2
Total	87	5	92	10	24	21	55	147

The two dominant pathways of arrival for non-native species considered to have a negative human impact are ornamental and contaminant (Table 19). However, there are a large number of established non-native species exerting a negative human impact and for which the introduction pathway is unknown. This is largely a consequence of the paucity of information available for insects and further work is required to elucidate the exact pathways. However, it is apparent that the ornamental pathway is a major contributor of non-native insects and higher plants with a negative impact on humans.

Table 19. Number of established non-native species, arriving by different pathways, within broad taxonomic groupings considered to have a **negative human impact**.

	Higher plants	Lower plants	Total plants	Insects	Invertebrates	Vertebrates	Total animals	Other	Total
Agriculture	3		3	1			1		4
Animal husbandry			0			4	4		4
Aquaculture		2	2		10		10		12
Contaminant	9		9	38			38		47
Ornamental	15		15	22	1	11	34		49
Stowaway	1	1	2	10	12	3	25		27
Unknown			0	17	7		24	1	25
No data			0	20			20		20
Total	28	3	31	108	30	18	156	1	188

Approximately a third of the established non-native species considered to have a negative ecological or human impact within the terrestrial environment originate from Europe (Table 20a). In contrast North America is the major region of origin for established non-native species considered to have a negative ecological or human impact within the freshwater environment (Table 20b). Established non-native species, considered to have a negative ecological or human impact, arriving within the marine environment originate relatively evenly from temperate Asia, North America and the Pacific (Table 20c). It is apparent that the number of non-native species arriving from temperate Asia has been increasing in recent years while the number of new arrivals from Europe has been decreasing. Clearly temperate Asia is becoming a major contributor of established non-native species to all environments.

Table 20. Number of established non-native species, arriving by different pathways and regions of origin, within broad taxonomic groups considered to have a negative ecological or human impact. Subdivided into species from the terrestrial environment (a), freshwater environment (b) and marine environment (c).

	Africa	Asia- Temperate	Asia- Tropical	Australasia	Europe	North America	Pacific	South America	No data	Total
a. Terrestrial										
Higher plants										
Agriculture					4					4
Contaminant					10					10
Forestry					3					3
Ornamental	3	14	4	1	23	12		4	10	71
Stowaway		1		1	1					3
Insects										
No data		8	2		3	7			1	21
Agriculture					1					1
Contaminant	7	3	3	2	5	1		3	14	38
Ornamental	2	5		3	3	2		1	6	22
Stowaway		1		3	3	3			2	12
Unknown	4		1		4	3		2	4	18
Invertebrates										
Ornamental					1					1
Unknown					4					4
Vertebrates										
Animal husbandry		1			2				3	6
Biocontrol									1	1
Ornamental	1	3	1		2	1				8
Stowaway		1	1		1					3
Total	17	37	12	10	70	29	0	10	41	226

b. Freshwater										
	Africa	Asia- Temperate	Asia- Tropical	Australasia	Europe	North America	Pacific	South America	No data	Total
Higher plants										
Aquaculture						1		2		3
Ornamental	1			1		6				8
Invertebrates										
Aquaculture		2			4	1				7
Stowaway		1		1		1				3
Unknown		1			1					2
Vertebrates										
Animal husbandry						1				1
Marine										0
Total	1	4	0	2	5	10	0	2	0	24
c. Marine										
	Africa	Asia- Temperate	Asia- Tropical	Australasia	Europe	North America	Pacific	South America	No data	Total
Lower plants										
Aquaculture		2					1			3
Stowaway		2								2
Invertebrates										
Aquaculture					1	1				2
Stowaway		2		1		3	1		2	9
Unknown	1					1	1			3
Other										
Unknown							1			1
Total	1	6	0	1	1	5	4	0	2	20

5. System for on-line recording of alert species including the outreach component “Recording Invasive Species Counts” (RISC)

An on-line system for reporting occurrences of species that require highlighting so called “alert species” has been developed (<https://www.nonnativespecies/alerts/index.cfm>; Figure 9). The open-access software system Indicia has been used to construct an on-line system, including photo upload and interactive maps, linked to the content management system Drupal for verification of records submitted (Figure 10). The same system is used for species within the RISC project.




Figure 9. Details of mechanism for reporting alert species through an on-line recording system.

There is an e-mail address (alert_nonnative@ceh.ac.uk) that can be used for enquiries on non-native species or for reporting alert species if this is preferable to the on-line mechanism. This e-mail system receives approximately six e-mails a month and these have mainly been in relation to the killer shrimp, *Dikerogammarus villosus*, and the Asian hornet, *Vespa velutina*. In all cases the information has been rapidly disseminated to the Non-Native Species Secretariat. *Dikerogammarus villosus* enquiries are simultaneously relayed to the Environment Agency and David Aldridge (domain expert). *Vespa velutina* enquiries are simultaneously relayed to FERA and the Bees, Wasps and Ants Recording Society.

The on-line Alert system includes the capacity to add a new arrival of any species but there are specific forms for carpet sea-squirt, African clawed frog, Citrus longhorn beetle, emerald ash borer, Indian house crow, oak processionary moth, sacred ibis, Siberian chipmunk, topmouth gudgeon, monk parakeet, prairie dog, eagle owl and killer shrimp.

NON-NATIVE ALERT SYSTEM




Figure 10. Summary of the alert on-line recording system that has been developed using the open-access Indicia software.

The outreach component of the non-native species on-line recording (RISC) was launched in spring 2010 and received considerable attention from the media. Each of the non-native species (Table 23) is linked to a domain expert mainly through the schemes and societies. Some of the species (muntjac and American skunk cabbage) have received reasonable numbers of records. For others the response has been expectedly low (Citrus longhorn beetle and Water primrose). However, for others the response has been disappointing and further promotion is required. Additional non-native species have been added to the RISC project for recording in 2012 and beyond.

Table 21. Non-native species included within RISC with number of records received on-line and the number verified as correct species identification. Verification is only possible if a photograph or specimen is provided.

Species	No. records	No. verified records	Scheme or society involved
Muntjac deer	512	46	Mammal Society
American skunk cabbage	96	47	People's Trust for Endangered Species
Tree of heaven	15	4	BSBI
Floating pennywort	1	0	BSBI
Water fern	4	1	BSBI
Water primrose	0	0	BSBI
Citrus longhorn beetle	5	0	FERA
Western conifer seed bug	5	0	Heteroptera Recording Scheme
Southern green shieldbug	1	0	Heteroptera Recording Scheme
Rhododendron leafhopper	0	0	Auchenorrhyncha Recording Scheme
Zebra mussel	4	1	Conchological Society
Wakame	2	0	MBA
Chinese mitten crab	2	0	MBA
American bullfrog	0	0	Amphibian and Reptile Recording Group

6. Key recommendations

- i. Maintain and develop the database of non-native species (Species Register) to include the non-native species that have arrived since the onset of this project, for example more than 200 species of birds that were recorded in the bird reports data set for NBN Gateway do not yet have an entry in the GB-NNSIP species register.
- ii. Develop factsheets for new species designated as a priority for additional information.
- iii. Increase the flow of distribution data on non-native species particularly developing methods for uploading spreadsheets of non-native species through Indicia.
- iv. Continue to deliver the information from the species register, factsheets and NBN Gateway to the GB-NNSIP through the Non-Native Species Secretariat and enhance search functionality.
- v. Maintain the on-line system of alerts coupled with a mechanism for enquiries through e-mail.
- vi. Develop a system for non-native species information to be disseminated to relevant personnel within other organisations both on the basis of regional locality and environment (freshwater, marine and terrestrial).
- vii. Ensure that mechanisms to enhance data flow will support the development of automated trends and analysis particularly in the context of the indicator of non-native species.

7. Bibliography

- Abbott, P. (2005) *Plant atlas of Mid-West Yorkshire*. Yorkshire Naturalists' Union.
- Abbott, R.J., James, J.K., Irwin, J.A. & Comes, H.P. (2000) Hybrid origin of Oxford Ragwort, *Senecio squalidus* L. *Watsonia*, **23**, 123-138.
- Adams, M.J. (1991) The distribution of barley yellow mosaic virus (BaYMV) and barley mild mosaic virus (BaMMV) in UK winter barley samples, 1987-1990. *Plant Pathology*, **40**, 53-58.
- Agassiz, D. (1978) *Gelechia sabinella* (Zeller) (Lepidoptera: Gelechiidae), a species new to Britain. *Entomologist's Gazette*, **29**, 136-139.
- Agassiz, D.J.L. (1996a) Invasions of Lepidoptera into the British Isles. *The Moths and Butterflies of Great Britain and Ireland, Volume 3* (ed A. M. Emmet), pp. 9-36. Harley Books, Colchester.
- Agassiz, D.J.L. (1996b) Yponomeutidae (excluding Roeslerstammiidae). *The Moths and Butterflies of Great Britain and Ireland, Volume 3* (ed A. M. Emmet), pp. 39-114. Harley Books, Colchester.
- Agassiz, D.J.L. & Tuck, K.R. (1999) The Cypress Tip moth *Argyresthia cupressella* Walsingham, 1890 (Lepidoptera: Yponomeutidae) new to Britain. *Entomologist's Gazette*, **50**, 11-16.
- Ainsworth, G.C. (1949) The *Gladiolus* smut. *Transactions of the British Mycological Society*, **32**, 255-257.
- Alderman, D.J. (1996) Geographical spread of bacterial and fungal disease of crustaceans. *Revue Scientifique et Technique de l'Office International des Epizooties*, **15**, 603-632.
- Alexander, K., Butler, J. & Green, E. (2006) The value of different tree and shrub species to wildlife. *British Wildlife*, **18**, 18-28.
- Alexander, K.N.A. (1989) *Euophryum confine* Broun (Col., Curculionidae) in South Wales. *Entomologist's Monthly Magazine*, **125**, 106.
- Alexander, K.N.A. (1994) The Roman snail *Helix pomatia* L. in Gloucestershire and its conservation. *Gloucestershire Naturalist*, **7**, 9-14.
- Alexander, K.N.A. (2003) *Provisional atlas of the Cantharoidea and Buprestoidea (Coleoptera) of Britain and Ireland*. Biological Records Centre, Huntingdon.
- Allen, A.A. (1976) *Platypus parallelus* F. (= *linearis* Steph.) (Col.: Scolytidae) recaptured in Britain after 150 years. *Entomologist's Record and Journal of Variation*, **88**, 57-58.
- Allen, A.A. (1984) The earliest known British capture of *Cryptopleurum subtile* Sharp (Col.: Hydrophilidae). *The Entomologist's Record and Journal of Variation*, **96**, 35.
- Allen, A.A. (1985) *Platypus parallelus* (F.) (Col., Scolytidae) again captured at light in S.E. London. *Entomologist's Monthly Magazine*, **121**, 141.
- Allen, D.E. & Hatfield, G. (2004) *Medicinal Plants in Folk Tradition: An Ethnobotany of Britain & Ireland*. Timber Press.
- Andrews, C. & Chubb, J.C. (1984) Helminth parasites from Yorkshire fishes. *Bulletin of the European Association of Fish Pathologists*, **4**, 22-23.
- Anon. (2003) Annex 7: Analysis of current biological recording schemes. *Review of non-native species policy, report of the working group*, pp. 117-122. Defra Publications, London.
- Anon. (2003) *Review of non-native species policy, report of the working group*. Defra Publications, London.

- Ansell, A.D. (1963) *Venus mercenaria* L. in Southampton Water. *Ecology*, **44**, 396-397.
- Appiah, A.A., Jennings, P. & Turner, J.A. (2004) *Phytophthora ramorum*: one pathogen and many diseases, an emerging threat to forest ecosystems and ornamental plant life. *Mycologist*, **18**, 145-150.
- Araujo, F.G., Bailey, R.G. & Williams, W.P. (1999) Spatial and temporal variations in fish populations in the upper Thames estuary. *Journal of Fish Biology*, **55**, 836-853.
- Arnold, H.R. (1995) *Atlas of amphibians and reptiles in Britain*. Biological Records Centre, Huntingdon.
- Ashelby, C.W., Worsfold, T.M. & Fransen, C.H.J.M. (2004) First records of the oriental prawn *Palaemon macrodactylus* (Decapoda : Caridea), an alien species in European waters, with a revised key to British Palaemonidae. *Journal of the Marine Biological Association of the United Kingdom*, **84**, 1041-1050.
- Asher, J., Warren, M., Fox, R., Harding, P., Jeffcoate, G. & Jeffcoate, S. (2001) *The millennium atlas of butterflies in Britain and Ireland*. Oxford University Press, Oxford.
- Atkins, P.M., O'Callaghan, D.P. & Kirby, S.G. (1981) *Scolytus laevis* (Chapuis) (Coleoptera, Scolytidae) new to Britain. *Entomologist's Gazette*, **32**, 280.
- Atkins, S.M., Jones, A.M. & Garwood, P.R. (1987) The ecology and reproductive cycle of a population of *Marenzelleria viridis* (Annelida: Polychaeta: Spionidae) in the Tay Estuary. *Proceedings of the Royal Society of Edinburgh*, **92B**, 311-322.
- Aubrook, E. & Fryer, G. (1965) The parasitic copepod *Tracheliastes polycolpus* Nordmann in some Yorkshire rivers: the first British records. *Naturalist*, **893**, 51-56.
- Auge, H. & Brandl, R. (1997) Seedling recruitment in the invasive clonal shrub, *Mahonia aquifolium* Pursh (Nutt.). *Oecologia*, **110**, 205-211.
- Bagnall, R.S. (1912) Report on the field meetings of the Natural History Society for 1911. *Trans.Nat.Hist.Soc.Northumberland, Durham & Newcastle-upon-Tyne*, **4**, 344-365.
- Baker, C.R.B. (1968) Notes on *Epiphyas* (= *Austrotortrix*) *postvittana* (Walker) (Lep., Tortricidae). *Entomologist's Gazette*, **19**, 167-172.
- Baker, R. & Clarke, K. (1997) New amphipod to Norfolk from Wheatfen Broad. *Transactions of the Norfolk and Norwich Naturalists Society*, **31**, 66-67.
- Baker, R.H.A., Black, R., Copp, G.H., Haysom, K.A., Hulme, P.E., Thomas, M.B., Brown, A., Brown, M., Cannon, R.J., Ellis, J., Ellis, M., Ferris, R., Glaves, P., Gozlan, R.E., Holt, J., Howe, L., Knight, J.D., MacLeod, A., Moore, N.P., Mumford, J.D., Murphy, S.T., Parrott, D., Sansford, C.E., Smith, G.C., St-Hilaire, S. & Ward, N.L. (2008) The UK risk assessment scheme for all non-native species. *Biological Invasions – from Ecology to Conservation. (Neobiota, Vol. 7)* (eds W. Rabitsch, F. Essl & F. Klingenstein), pp. 46-57. Berlin.
- Ball, S.G. & Morris, R.K.A. (2000) *Provisional atlas of British hoverflies (Diptera, Syrphidae)*. Biological Records Centre, Huntingdon.
- Bamber, R.N. (1985) The itinerant sea spider *Ammothea hilgendorfi* (Böhm) in British waters. *Proceedings of Hampshire Field Club & Archaeological Society*, **41**, 269-270.
- Banks, B., Foster, J., Langton, T. & Morgan, K. (2000) British bullfrogs? *British Wildlife*, **11**, 327-330.
- Barber, A.D. (2009) *Centipedes*. Field Studies Council, Shrewsbury.
- Barber, A.D. & Keay, A.N. (1988) *Provisional atlas of the centipedes of the British Isles*. Biological Records Centre, Huntingdon.

- Barclay, M.V.L. (2003) *Otiorhynchus* (s. str.) *armadillo* (Rossi, 1792) *Otiorhynchus salicicola* (s. str.) Heyden, 1908 (Curculionidae: Entiminae: Otiorhynchini) – two European vine weevils established in Britain. *The Coleopterist*, **12**, 41-56.
- Barclay, M.V.L. (2004) The green vegetable bug *Nezara viridula* (L., 1758) (Hem.: Pentatomidae) new to Britain. *The Entomologist's Record and Journal of Variation*, **116**, 55-58.
- Barnes, H. & Barnes, M. (1960) Recent spread and present distribution of the barnacle *Elminius modestus* Darwin in north-west Europe. *Proceedings of the Zoological Society of London*, **135**, 137-145.
- Barnes, R.S.K. (1994) *The brackish-water fauna of northwestern Europe*. Cambridge University Press, Cambridge.
- Barnes, R.S.K., Coughlan, J. & Holmes, N.J. (1973) A preliminary survey of the macroscopic bottom fauna of the Solent, with particular reference to *Crepidula fornicata* and *Ostrea edulis*. *Proceedings of the Malacological Society*, **40**, 253-275.
- Barson, G. & Carter, C.I. (1972) A species of Phylloxeridae, *Moritzziella corticalis* (Kalt.) (Homoptera) new to Britain, and a key to the British oak-feeding Phylloxeridae. *The Entomologist*, **105**, 130-134.
- Bartlett, P.W. (1979) Preventing the establishment of Colorado beetle in England and Wales. *Plant Health* (eds D. L. Ebbels & J. E. King), pp. 247-257. Blackwell Scientific Publications, Oxford.
- Bas, K. (2001) The unexpected one jumped the North Sea. *Field Mycology*, **2**, 40-41.
- Bastrop, R., Röhner, M. & Jürss, K. (1995) Are there two species of the polychaete genus *Marenzelleria* in Europe? *Marine Biology*, **121**, 509.
- Beckett, G. & Bull, A. (1999) *Flora of Norfolk*, A. privately published.
- Benson, R.B. (1952) A new *Anoplonyx* destructive to larch in Britain (Hymenoptera, Tenthredinidae). *Bulletin of Entomological Research*, **43**, 543-547.
- Benson, R.B. (1958) Hymenoptera 2. Symphyta. Section (c). *Handbooks for the Identification of British Insects*, **6**, 1-252+vi.
- Benton, T.G. (1991a) The life history of *Euscorpium flavicaudis* (Scorpiones, Chactidae). *Journal of Arachnology*, **19**, 105-110.
- Benton, T.G. (1991b) Reproduction and parental care in the Scorpion, *Euscorpium flavicaudis*. *Behaviour*, **117**, 20-28.
- Benton, T.G. (1992) The Ecology of the scorpion *Euscorpium flavicaudis* in England. *Journal of Zoology*, **226**, 351-368.
- Benvenuti, S. Weed dynamics in the Mediterranean urban ecosystem: ecology, biodiversity and management. *Weed Research*, **44**, 341–354.
- Bertolino, S. & Genovesi, P. (2003) Spread and attempted eradication of the grey squirrel (*Sciurus carolinensis*) in Italy, and consequences for the red squirrel (*Sciurus vulgaris*) in Eurasia. *Biological Conservation*, **109**, 351-358.
- Bevan, D. (1987) *Forest insects. A guide to insects feeding on trees in Britain*. HMSO, Forestry Handbook no.1. London.
- Bew, M. (1993) Hunting for Varroa in the UK. *The Beekeepers Annual* (ed R. Coles), pp. 67-69. Northern Bee Books [cited in Dobson 1998], Mytholmroyd.
- Beyer, K., Gozlan, R.E. & Copp, G.H. ((in prep.)) Divide and conquer: a successful social networking strategy for non-native fishes when invading an established fish assemblage.

- Beyer, K., Gozlan, R.E., Pinder, A.C. & Copp, G.H. ((in prep.)) Early life interactions between native cyprinids and invasive sunbleak *Leucaspis deliniatus* in a three-lake system.
- Billany, D.J. (1978) *Gilpinia hercyniae* – a pest of spruce. HMSO, Forestry Commission Forest Record 117. London.
- Billany, D.J. & Brown, R.M. (1980) The Web-spinning sawfly *Cephalcia lariciphila* (Wachtl) (Hym.: Pamphiliidae) – a new pest of larch in England and Wales. *Forestry*, **53**, 71-80.
- Billings, E. (1976) Fireblight in England. *Proceedings of the Third Workshop on Fire Blight Research, Ithaca, New York, 20-22 September 1976*, pp. 118-121.
- Bishop, M.W.H. (1947) Establishment of an immigrant barnacle in British coastal waters. *Nature*, **159**, 501.
- Bishop, M.W.H. (1950) Distribution of *Balanus amphitrite* Darwin var. *denticulata* (Broch). *Nature*, **165**, 409.
- Blackman, R.L. (1984) Two species of Aphididae (Hem.) new to Britain. *Entomologist's Monthly Magazine*, **120**, 185-186.
- Blackman, R.L. & Eastop, V.F. (1994) *Aphids on the World's Trees*. CABI, Wallingford UK.
- Blackman, R.L. & Eastop, V.F. (2000) *Aphids on World's Crops* John Wiley, Chichester UK.
- Blackman, R.L. & Eastop, V.F. (2006) *Aphids on the Worlds Herbaceous Plants and Shrubs*. John Wiley.
- Blair, M.J., McKay, H., Musgrove, A.J. & Rehfish, M.M. (2000) *Review of the status of introduced non-native waterbird species in the agreement area of the African-Eurasian Waterbird Agreement*. BTO, Thetford.
- Bland, K.P., Corley, M.F.V., Emmet, A.M., Heckford, R.J., Huemer, P., Langmaid, J.R., Palmer, S.M., Parsons, M.S., Pitkin, L.M., Sattler, K. & Simpson, A.N.B. (2002) Gelechiinae. *The Moths and Butterflies of Great Britain and Ireland, Volume 4, Part 2* (eds A. M. Emmet & J. R. Langmaid), pp. 118-203. Harley Books, Colchester.
- Blockeel, T.L. & Long, D.G. (1998) *A check-list and census catalogue of British and Irish bryophytes*. British Bryological Society, Cardiff.
- Blower, G. (1955) Yorkshire Centipedes. *Naturalist* **1955**, 137-146.
- Boalch, G.T. (1987) Changes in the phytoplankton of the western English Channel in recent years. *British Phycological Journal*, **22**, 225-235.
- Bolton, B. & Collingwood, C.A. (1975) Hymenoptera: Formicidae. *Handbooks for the Identification of British Insects, VI Part 3(c)*.
- Booth, R. (2005) <http://www.coleopterist.org.uk/>.
- Booth, R. (2005b) <http://www.coleopterist.org.uk/>.
- Booth, R. (2005c) <http://www.coleopterist.org.uk/>.
- Botham, M.S., Rothery, P., Hill, M.O., Preston, C.D. & Roy, D.B. Do urban areas act as foci for the spread of alien plant species? An assessment of temporal trends in the UK. *Diversity and Distributions*, **15**, 338-345.
- Bowen, H. (2000) *The Flora of Dorset*. Pisces Publications.
- Boxshall, G.A. & Frear, P.A. (1990) *Tracheliastes maculatus* (Kollar, 1836) (Crustacea, Lernaepodidae) a new species for Britain. *Journal of Fish Biology*, **37**, 489-491.
- Bradley, J.D. (2000) *Checklist of Lepidoptera recorded from the British Isles*. D.J.Bradley & M.J.Bradley, Fordingbridge.
- Bradley, J.D., Tremewan, W.G. & Smith, A. (1973) *British Tortricoid Moths*. The Ray Society,

- London.
- Branquart, E. (2007) [Branquart is editor] Guidelines for environmental impact assessment and list classification of non-native organisms in Belgium. http://ias.biodiversity.be/ias/documents/ISEIA_protocol.pdf . Belgian Forum on Invasive Species.
- Brenan, J.P.M. (1961) *Amaranthus* in Britain. *Watsonia*, **4**, 261-280.
- Bretherton, R.F., Goater, B. & Lorimer, R.I. (1979) Noctuidae: Noctuinae and Hadeninae. *The Moths and Butterflies of Great Britain and Ireland, Volume 9* (eds J. Heath & A. M. Emmet), pp. 120-280. Curwen Books, London.
- Brewis, L.A., Bowman, P. & Rose, F. (1996) *The Flora of Hampshire*. Harley Books.
- British Myriapod Group (1988) *Preliminary atlas of the millipedes of the British Isles*. Biological Records Centre, Huntingdon.
- Brock, P.D. (1986) A third New Zealand stick insect (Phasmatodea) established in the British Isles, with notes on the other species, including a correction. *1st International Symposium on Stick Insects* (eds M. Mazzini & V. Scali), pp. 125-132. University of Siena (1985).
- Broennimann, O. & Guisan, A. (2008) Predicting current and future biological invasions: both native and invaded ranges matter. *Biology Letters*, **4**, 585–589.
- Brown, J.A., Moore, W.M. & Quabius, E.S. (2001) Physiological effects of saline waters on zander. *Journal of Fish Biology*, **59**, 1544-1555.
- Brylinski, J.M. (1981) Reports on the presence of *Acartia tonsa* Dana (Copepoda) in the area of Dunkirk and its geographical distribution in Europe. *Journal of Plankton Research*, **3**, 255-261.
- Buck, F.D. (1948) *Pentarthrum huttoni* Woll. (Col., Curculionidae) and some imported Cossoninae. *Entomologist's Monthly Magazine*, **84**, 152-154.
- Buczacki, S.T. & Harris, K.M. (1981) *Collins guide to the pests, diseases and disorders of garden plants*. William Collins and Co., London.
- Buczacki, S.T. & Harris, K.M. (1998) *Pests, diseases and disorders of garden plants*. HarperCollins, London.
- Burrows, E.M. (1991) *Seaweeds of the British Isles. Volume 2 Chlorophyta*. Natural History Museum Publications, London.
- Burton, R.M. (1998) Botanical records for 1997. *The London Naturalist*, **77**, 225-?
- Butler, C. (2002) Breeding parrots in Britain. *British Birds*, **95**, 345-348.
- Cannon, P.F., Hawksworth, D.L. & Sherwood-Pike, M.A. (1985) *The British Ascomycetes: an annotated checklist*. Commonwealth Agricultural Bureaux, Slough.
- Cannon, P.F., Kirk, P.M., Cooper, J.A. & Hawksworth, D.L. (2001) Microscopic fungi. *The changing wildlife of Great Britain* (ed D. L. Hawksworth), pp. 114-125. Taylor & Francis, London & New York.
- Cannon, R.J.C. (1998) The status of the New Zealand flatworm, *Artioposthia triangulata*, in the UK. Central Science Laboratory, unpublished report to MAFF Plant Health Service.
- Cannon, R.J.C., Koerber, D., Ashby, S., Baker, R.H.A., Bartlett, P.W., Brookes, G., Burgess, R., Cheek, S., Evans, H.F., Hammon, R., Head, J., Nettleton, G., Robinson, J., Slawson, D., Taylor, M.C., Tilbury, C.A. & Ward, M. (2004) Gypsy moth, *Lymantria dispar*, outbreak in northeast London, 1995-2003. *International Journal of Pest Management*, **50**, 259-273.

- Cannon, R.J.C., Matthews, L., Cheek, S., Baker, R.H.A., MacLeod, A., Bartlett, P.W. & Savage, D. (2005) Surveying and monitoring western corn rootworm (*Diabrotica virgifera virgifera*) in England and Wales. *2005 BCPC Symposium Proceedings No. 81, Introduction and Spread of Invasive Species*, pp. 155-160.
- Carter, C.I. (1983) Some new aphid arrivals to Britain's forests. *Proceedings and Transactions of the British Entomological and Natural History Society*, **16**, 81-87.
- Carter, C.I. & Eastop, V.F. (1975) *Mindarus obliquus* (Chol.) (Homoptera, Aphidoidea) new to Britain and records of two other aphids recently found feeding on conifers. *Entomologist's Monthly Magazine*, **108**, 202-204.
- Carter, C.I., Fout, D.F. & Bartlett, P.W. (1984) The Lupin Aphid's arrival and consequences. *Antenna*, **8**, 129-132.
- Carter, I., Cross, A.V., Douse, A., Duffy, K., Etheridge, B., Grice, P.V., Newberry, P., Orr-Ewing, D.C., O'Toole, L., Simpson, D. & Snell, N. (2003) Re-introduction and conservation of the red kite (*Milvus milvus*) in Britain: current threats and prospects for future expansion. *Birds of prey in a changing environment* (eds D. B. A. Thompson, S. M. Redpath, A. H. Fielding, M. Marquiss & C. A. Galbraith), pp. 407-416. The Stationery Office, Edinburgh.
- Carter, J. & Leonard, B.P. (2002) A review of the literature on the worldwide distribution, spread of, and efforts to eradicate the coypu (*Myocaster coypus*). *Wildlife Society Bulletin*, **30**, 162-175.
- Centre for Aquatic Plant Management (2003) *Annual report summary 2002*. <http://www.nerc-wallingford.ac.uk/research/capm/pdf%20files/Summary%20Annual%20Report.pdf>.
- Chandler, P.J. (1998) *Checklists of insects of the British Isles (new series). Part 1, Diptera, including a list of Irish Diptera*. Royal Entomological Society, London.
- Chandler, P.J. (1999) *Micropygus vagans* Parent (Diptera: Dolichopodidae), a New Zealand fly in the British Isles. *British Journal of Entomology and Natural History*, **12**.
- Charles, J.G. & Henderson, R.C. (2002) Catalogue of the exotic armoured scale insects (Hemiptera: Coccoidea: Diaspididae) in New Zealand. *Journal of the Royal Society of New Zealand*, **32**, 587-615.
- Chater, A.O. (2010) *Flora of Cardiganshire*. privately published.
- Chater, A.O., Oswald, P.H. & Preston, C.D. (2000) Street floras in Cambridge and Aberystwyth. *Nature in Cambridgeshire*, **42**, 3-26.
- Child, L. & Wade, M. (2000) *The Japanese knotweed manual*. Packard, Chichester.
- China, W.E. (1935) A North American Jassid (Homoptera) in Surrey. *Entomologist's Monthly Magazine*, **71**, 277-279.
- Chytrý, M., Maskell, L.C., Pino, J., Pyšek, P., Vilà, M., Font, X. & Smart, S.M. (2008) Habitat invasions by alien plants: a quantitative comparison among Mediterranean, subcontinental and oceanic regions of Europe. *Journal of Applied Ecology*, **45**, 448-458.
- Clark, M.C. (1982) Further discomycetes in Britain. *Bulletin of the British Mycological Society*, **16**, 60-66.
- Clement, E.J. & Foster, M.C. (1994) *Alien Plants of the British Isles: A Provisional Catalogue of Vascular Plants [excluding Grasses]*. Botanical Society of the British Isles.
- Clemons, L. (2000) *Tephritis matricariae* (Loew, 1844) (Dip.: Tephritidae) new to Britain and breeding in East Kent. *Entomologist's Record and Journal of Variation*, **112**, 225-230.
- Cloudsley Thompson, J.L. & Constantinou, C. (1983) How does the scorpion *Euscorpius*

- flavicaudis (Deg) manage to survive in Britain. *International Journal of Biometeorology*, **27**, 87-92.
- Clover, G.R.G., Ratti, C. & Henry, C.M. (2001) Molecular characterization and detection of European isolates of Soil-borne wheat mosaic virus. *Plant Pathology*, **50**, 761-767.
- Colak, A.H., Cross, J.R. & Rotherham, I.D. (1998) *Rhododendron ponticum* in native and exotic environments, with particular reference to Turkey and the British Isles. *Journal of Practical Ecology and Conservation*, **2**, 34-41.
- Cole, H.A. (1942) The American whelk tingle, *Urosalpinx cinerea* (Say), on British oyster beds. *Journal of the Marine Biological Association of the United Kingdom*, **25**, 477-508.
- Cole, J. (1996) A second British site for *Prosopandrus flavifrons* (Tonnoir & Malloch) (Diptera, Heleomyzidae). *Entomologist's Monthly Magazine*, **132**, 310.
- Cole, S. (2002) 2001 Annual Exhibition, Imperial College, London SW 7- November 2001. Coleoptera. *British Journal of Entomology and Natural History*, **11**, 176.
- Colville, B., Lloyd-Evans, L. & Norris, A. (1974) *Boettgerilla pallens* Simroth, a new British species. *Journal of Conchology, London*, **28**, 203-207.
- Comfort, A. (1950) *Hygromia cinctella* (Draparnaud) in England. *Journal of Conchology, London*, **23**, 99-100.
- Cooter, J. (2005) <http://www.coleopterist.org.uk/>.
- Cope, T. & Gray, A. (2009) *Grasses of the British Isles*. Botanical Society of the British Isles.
- Copp, G.H., Bianco, P.G., Bogutskaya, N., Eros, T., Falka, I., Ferreira, M.T., Fox, M.G., Freyhof, J., Gozlan, R.E., Grabowska, J., Kovác, V., Moreno-Amich, R., Naseka, A.M., Penáz, M., Povž, M., Przybylski, M., Robillard, M., Russell, I.C., Stakenas, S., Šumer, S., Vila-Gispert, A. & Wiesner, C. (2005) To be, or not to be, a non-native freshwater fish? *Journal of Applied Ichthyology*, (in press).
- Copp, G.H., Fox, M.G. & Kovác, V. (2002) Growth, morphology and life history traits of a coolwater European population of pumpkinseed *Lepomis gibbosus*. *Archiv Fur Hydrobiologie*, **155**, 585-614.
- Copp, G.H., Fox, M.G., Przybylski, M., Godinho, F. & Vila-Gispert, A. (2004) Life-time growth patterns of pumpkinseed *Lepomis gibbosus* introduced to Europe relative to native North American populations. *Folia Zoologica*, **53**, 237-254.
- Copp, G.H., Stakenas, S., Davison, P. & Carter, M.G. ((in prep.)) The incidence of non-native fishes in UK rivers, with particular reference to re-occurrence of the white sucker *Catostomus commersoni* in the River Gade (Hertfordshire).
- Copp, G.H., Vaughn, C. & Wheeler, A. (1993) First occurrence of the North American white sucker *Catostomus commersoni* in Great Britain. *Journal of Fish Biology*, **42**, 615-617.
- Copp, G.H., Wesley, K.J., Kovác, V., Ives, M. & Carter, M.G. (2003) Introduction and establishment of the pikeperch *Stizostedion lucioperca* (L.) in Stanborough Lake (Hertfordshire) and its dispersal in the Thames catchment. *The London Naturalist*, **82**, 139-153.
- Copp, G.H., Wesley, K.J. & Vilizzi, L. (2005) Pathways of ornamental and aquarium fish introductions into urban ponds of Epping Forest (London, England): the human vector. *Journal of Applied Ichthyology* (in press).
- Corbet, G.B. & Harris, S. (1991) *The handbook of British mammals*. Blackwell Scientific Publications, Oxford.

- Cotten, J., Bartlett, P.W. & Webb, R.M. (1991) A first record of the root lesion nematode, *Pratylenchus bolivianus* Corbett in England and Wales. *Plant Pathology*, **40**, 311-312.
- Cotten, J. & Hooper, D.J. (1991) Records of nematodes associated with azaleas in England: *Paratrichodorus renifer* Siddiqi, *Tylenchorhynchus claytoni* Steiner. *Plant Pathology*, **40**, 308-310.
- Cotton, A.D. (1908) The appearance of *Colpomenia sinuosa* in Britain. *Bulletin of Miscellaneous Information, Royal Botanic Gardens, Kew*, **1908**, 73-77.
- Countryside Survey Partnership (2008) Countryside Survey: UK Results from 2007. 2. The national picture.
- Cowling, J., Spicer, J.I., Gaston, K.J. & Weeks, J.M. (2004) Current status of an amphipod invader, *Arcitalitrus dorrieni* (Hunt, 1925), in Britain. *Journal of Natural History*, **38**, 1665-1675.
- Cox, M.L. (2004) <http://www.coleopterist.org.uk/>.
- Cox, M.L. ((in prep.)) *Atlas of Chrysomelid and Bruchid beetles*. CEH, Huntingdon.
- Crawley, M. (2005) *The Flora of Berkshire: With Accounts of Charophytes, Ferns, Flowering Plants, Bryophytes, Lichens and Non-lichenized Fungi*. Brambleby Books.
- Crawley, M.J. (1987) What makes a community invasible? *Colonization, succession and stability* (eds A. J. Gray, M. J. Crawley & P. J. Edwards), pp. 429-453. Blackwell Scientific Publications, Oxford.
- Crawley, M.J. (1997) *Aphelonyx cerricola* Giraud (Hym., Cynipidae), an alien gall-former new to Britain. *Entomologist's Monthly Magazine*, **133**, 61.
- Crawley, M.J., Harvey, P.H. & Purvis, A. (1996) Comparative ecology of the native and alien floras of the British Isles. *Philosophical Transactions of the Royal Society of London*, **B 351**, 1251-1259.
- Crisp, D.J. (1958) The spread of *Elminius modestus* Darwin in north-west Europe. *Journal of the Marine Biological Association of the United Kingdom*, **37**, 483-520.
- Cronk, Q.C.B. & Fuller, J.L. (1995) *Plant invaders*. Chapman & Hall, London.
- Crooke, M. & Bevan, D. (1957) Notes on the first occurrence of *Ips cembrae* (Heer) (Col.: Scolytidae). *Forestry*, **30**, 21-28.
- Crowson, R.A. (1988) Observations on Caenoscelini (Col., Cryptophagidae). *Entomologist's Monthly Magazine*, **124**, 72.
- Crundwell, A.C. (1992) The bryophytes of Britain and Ireland in a European context. *Atlas of the bryophytes of Britain and Ireland, Vol. 2* (eds M. O. Hill, C. D. Preston & A. J. E. Smith), pp. 9-16. Harley Books, Colchester.
- DAFF (2002) Database and atlas of freshwater fishes, compiled by C.E. Davies and others. Centre for Ecology and Hydrology. Monkswood, Cambridgeshire. (www.brc.ac.uk/DAFF/daff.htm).
- DAISIE (2008) *Handbook of Alien Species in Europe: Delivering Alien Invasive Species Inventories for Europe*. Springer.
- Davies, B.R., Stuart, V. & Villiers, M.d. (1989) The filtration activity of a serpulid polychaete population (*Ficopomatus enigmaticus* (Fauvel)) and its effects on water quality in a coastal marina. *Estuarine, Coastal and Shelf Science*, **29**, 613-620.
- Davies, C., Shelley, J., Harding, P., McLean, I., Gardiner, R. & Peirson, G. (2004) Freshwater fishes in Britain – the species and their distribution. pp. 176. Harley Books, Colchester.

- Davies, E.H. (1968) Myxosporidian parasites from the fish of the River Lugg. *Journal of Protozoology*, **15**, 471-480.
- Dawson, W., Burslem, D.F.R.P. & Hulme, P.E. (2009) Herbivory is related to taxonomic isolation, but not to invasiveness of tropical alien plants. *Diversity and Distributions*, **15**, 141-147.
- De Ferrari, C.M. & Naiman, R.J. (1994) A multi-scale assessment of the exotic plants on the Olympic Peninsula, Washington. *Journal of Vegetation Science*, **5**, 247-258.
- Deckers, B., Verheyen, K., Vanhellefont, M., Maddens, E., Muys, B. & Hermy, M. (2008) Impact of avian frugivores on dispersal and recruitment of the invasive *Prunus serotina* in an agricultural landscape. *Biological Invasions*, **10**, 717-727.
- Defra (2003) *Review of non-native species policy*. Department for Environment, Food and Rural Affairs, London.
- Dehnen-Schmutz, K. & Williamson, M. (2006) *Rhododendron ponticum* in Britain and Ireland: social, economic and ecological factors in its successful invasion. *Environment and History*, **12**, 325-50.
- Dennis, R.W.G. (1955) The status of *Clathrus* in England. *Kew Bulletin*, **1955**, 101-106.
- Dennis, R.W.G. (1958) New British fungi. *Kew Bulletin*, **1957**, 399-404.
- Dennis, R.W.G. (1975) New or interesting British microfungi, III. *Kew Bulletin*, **30**, 345-365.
- Dennis, R.W.G. (1981) *British Ascomycetes, revised edition, second impression with supplement*. J. Cramer, Vaduz.
- Dennis, R.W.G. (1995) *The fungi of southeast England*. Royal Botanic Gardens, Kew.
- Dennis, R.W.G. & Hubbard, C.E. (1962) A new British smut. *Kew Bulletin*, **15**, 379.
- Dennis, R.W.G. & Wakefield, E.M. (1949) New or interesting British fungi. *Transactions of the British Mycological Society*, **29**, 141-166.
- Dickson, J.H. (1998) Plant introductions in Scotland. *Species history in Scotland* (ed R. A. Lambert), pp. 38-44. Scottish Cultural Press, Edinburgh.
- Dickson, J.H., Macpherson, P. & Watson, K. (2000) *The Changing Flora of Glasgow: Urban and Rural Through the Centuries: Urban and Rural Plants Through the Centuries*. Edinburgh University Press.
- Dickson, R.J. (2002) Blastobasidae. *The Moths and Butterflies of Great Britain and Ireland, Volume 4, Part 2* (eds A. Maitland Emmett & J. R. Langmaid), pp. 196-203. Harley Books, Colchester.
- Dixon, P.S. & Irvine, L.M. (1977) *Seaweeds of the British Isles. Volume 1 Rhodophyta Part 1 Introduction, Nemaliales, Gigartinales*. British Museum (Natural History), London.
- Dobson, J.R. (1998) A 'bee louse' *Braula schmitzi* Örösi-pál (Diptera: Braulidae) new to the British Isles, and the status of *Braula* spp. in England and Wales. *British Journal of Entomology and Natural History*, **11**, 139-148.
- Dolling, W.R. (1972) A new species of *Dicyphus* Fieber (Hem., Miridae) from southern England. *Entomologist's Monthly Magazine*, **107**, 224-225.
- Dony, J.G. (1953) *Flora of Bedfordshire*. Museum & Art Gallery.
- Dring, D.M. (1980) Contributions towards a rational arrangement of the Clathraceae. *Kew Bulletin*, **35**, 1-96.
- Duff, A.G. (2004) <http://www.coleopterist.org.uk/>.
- Duff, A.G. (2004b) <http://www.coleopterist.org.uk/>.
- Duff, A.G. (2004c) <http://www.coleopterist.org.uk/>.

- Duff, A.G. (2004d) <http://www.coleopterist.org.uk/>.
- Durham, M.J. (1947) Occurrence of the land nemertine, *Geonemertes dendyi* Dakin, in Wirral, Cheshire. *North Western Naturalist*, **22**, 182-183.
- Duval, D.M. (1963) The biology of *Petricola pholadiformis* Lamarck (Lamellibranchiata: Petricolidae). *Proceedings of the Malacological Society*, **35**, 89-100.
- Dyrynda, P.E.J., Fairall, V.R., Ambrogi, A.O. & d'Hondt, J.-L. (2000) The distribution, origins and taxonomy of *Tricellaria inopinata* d'Hondt and *Occhipinti* Ambrogi, 1985, an invasive bryozoan new to the Atlantic. *Journal of Natural History*, **34**, 1993-2006.
- Eason, E.H. (1964) *Centipedes of the British Isles*. Warne, London & New York.
- Easy, G. (1978) *Verbascum pyramidatum* Bied. and the hybrid *V. pyramidatum* x *thapsus* along a dismantled railway in Cambs. *BSBI News*, **19**, 21.
- Edwards, C. (1976) A study in erratic distribution: the occurrence of the medusa *Gonionemus* in relation to the distribution of oysters. *Advances in Marine Biology*, **14**, 251-284.
- Eggerling, T.W. (2004) *Paurocotylis pila*. *Field Mycology*, **5**, 41-42.
- Ellis, A.E. (1926) *British snails*. Clarendon Press, Oxford.
- Ellis, G. (1983) *Flowering Plants of Wales*. National Museums and Galleries of Wales.
- Ellis, J.P. (1977a) The genera *Trichothyria* and *Actinopeltis* in Britain. *Transactions of the British Mycological Society*, **68**, 145-155.
- Ellis, J.P. (1977b) The genus *Stomiopeltis* in Britain. *Transactions of the British Mycological Society*, **68**, 157-159.
- Ellis, J.P. (1980) The genus *Morenoina* in Britain. *Transactions of the British Mycological Society*, **74**, 297-307.
- Elphinstone, J.G., Stanford, H. & Stead, D. (1998) Survival and transmission of *Ralstonia solanacearum* in aquatic plants of *Solanum dulcamara* and associated surface water in England. *EPPO Bulletin*, **28**, 93-94.
- Emmet, A.M. (1989) *Phyllonorycter leucographella* (Zeller, 1870) (Lep., Gracillariidae) in Essex, a species new to Britain. *Entomologist's Record and Journal of Variation*, **101**, 189-194.
- Emmet, A.M. (1991) *Phyllonorycter platani* (Staudinger, 1870) (Lepidoptera, Gracillariidae) new to Britain. *Entomologist's Record and Journal of Variation*, **103**, 279-282.
- Emmet, A.M. & Heath, J. (1989) *The moths and butterflies of Great Britain and Ireland Volume 7, Part 1. The butterflies*. Harley Books, Colchester.
- Emmet, A.M., Heckford, R.J., Langmaid, J.R. & Palmer, S.M. (2002) Chelariinae. *The Moths and Butterflies of Great Britain and Ireland, Volume 4, Part 2* (eds A. M. Emmet & J. R. Langmaid), pp. 221-227. Harley Books, Colchester.
- Emmet, A.M., Watkinson, I.A. & Wilson, M.R. (1985) Gracillariidae. *The Moths and Butterflies of Great Britain and Ireland, Volume 2* (eds J. Heath & A. M. Emmet), pp. 244-362. Harley Books, Colchester.
- Engel, J.J. & Merrill, G.L.S. (2004) Austral Hepaticae. 35. A taxonomic and phylogenetic study of *Telaranea* (Lepidoziaceae), with a monograph of the genus in temperate Australasia and commentary on extra-Australasian taxa. *Fieldiana: Botany*, **44**, 1-265.
- Eno, N.C., Clark, R.A. & Sanderson, W.G. (1997) Non-native marine species in British waters: a review and directory. pp. 136. Joint Nature Conservation Committee, JNCC Report. Peterborough.
- Environment Agency (1999) *Category 2 parasite guide*. Environment Agency.

- Equihua, M. & Usher, M.B. (1993) Impact of carpets of the invasive moss *Campylopus introflexus* on *Calluna vulgaris* regeneration. *Journal of Ecology*, **81**, 359-365.
- Essink, K. (1986) Note on the distribution of the American jack knife clam *Ensis directus* (Conrad, 1843) in NW Europe (Bivalvia: Cultellidae). *Basteria*, **50**, 33-34.
- European Environment Agency (2007) *Halting the loss of biodiversity by 2010: proposal for a first set of indicators to monitor progress in Europe*. EEA, Copenhagen.
- Evans, H.F. (1997) The present position of forest entomology in Great Britain. *Forestry*, **70**, 327-336.
- Evans, T.G. (2007) Flora of Monmouthshire. *Chepstow Society, Chepstow, UK*.
- Everett, S. (1999) Putting wild plants where they belong. *Plantlife*, **Spring 1999**, 12-13.
- Eversham, B.C. & Arnold, H.R. (1992) Introductions and their place in British wildlife. *Biological recording of changes in British wildlife* (ed P. T. Harding), pp. 44-59. HMSO, London.
- Farnham, W.F. (1980) Studies on aliens in the marine flora of southern England. *The shore environment, Vol. 2: Ecosystems* (eds J. H. Price, D. E. G. Irvine & W. F. Farnham), pp. 875-914. Academic Press, London and New York.
- Farr-Cox, F., Leonard, S. & Wheeler, A.C. (1996) The status of the recently introduced fish *Leucaspis delineatus* (Cyprinidae) in Great Britain. *Fisheries Management and Ecology*, **3**, 193-199.
- Ferguson, I.K. (1978) *Verbascum speciosum* Schrader x *V. thapsus* L. new to Britain. *Watsonia*, **12**, 160-162.
- Fielding, N.J. & Evans, H.E. (1997) Biological control of *Dendroctonus micans* (Scolytidae) in Great Britain. *Biocontrol News and Information*, **18**, 51N-60N.
- Finch, R.A., Fisk, R.J., Strauss, D.F. & Stevenson, C.R. (2000) *Lophocolea semiteres* new to East Anglia. *Journal of Bryology*, **22**, 146-148.
- Fitter, R.S.R. (1994) Alien shrubs and trees in a Chiltern tetrad. *B.S.B.I. News*, **67**, 54-55.
- Fitter, R.S.R. (1996) Established alien perennials and biennials in a Chiltern tetrad. *B.S.B.I. News*, **73**, 41-42.
- Fletcher, R.L. (1987) *Seaweeds of the British Isles. Volume 3 Fucophyceae (Phaeophyceae)*. British Museum (Natural History), London.
- Ford, T.H., Shaw, M.R. & Robertson, D.M. (2000) Further host records of some west Palaearctic Tachinidae (Diptera). *Entomologist's Record and Journal of Variation*, **112**, 25-36.
- Fortey, R. (2004) *Psilocybe aurantiaca* and a case of mistaken identity. *Field Mycology*, **5**, 77-80.
- Foster, R.W. (1946) The genus *Mya* in the western Atlantic. *Johnsonia*, **2**, 20.
- Fox, R. & Asher, J. (2005) *Butterflies for the New Millennium. Atlas +5. Provisional maps (2000-2004)*. Butterfly Conservation, Lulworth.
- Fox-Wilson, G. (1942) The Lily Beetle *Crioceris* [now *Lilioceris*] *lilii*. *Journal of the Royal Horticultural Society*, **67**, 165-168.
- Francis, S.M., Minter, D.W. & Caine, T.S. (1980) Three new species of *Anthostomella*. *Transactions of the British Mycological Society*, **75**, 201-206.
- Frost, L.C. (1987) The alien Hottentot fig (*Carpobrotis edulis*) in Britain - a threat to the native flora and its conservation control. University of Bristol Lizard Project.
- Fryer, G. & Andrews, C. (1983) The parasitic copepod *Ergasilus briani* Markewitsch in Yorkshire: an addition to the British fauna. *Naturalist*, **108**, 7-10.
- Gaston, K.J., Cush, P., Ferguson, S., Frost, P., Gaston, S., Knight, D., Loram, A., Smith, R.M.,

- Thompson, K. & Warren, P.H. (2007) Improving the contribution of urban gardens for wildlife: some guiding propositions. *British Wildlife*, **18**, 171-177.
- Gates, P. (1992) *The Aliens are Coming - Plant Life and the Greenhouse Effect*. Penguin Books, London.
- Genovesi, P. (2005) Eradications of invasive alien species in Europe: a review. *Biological Invasions*, **7**, 127-133.
- Gibbons, D.W., Reid, J.B. & Chapman, R.A. (1993) *The New Atlas of Breeding Birds in Britain and Ireland: 1988-1991*. T. & A.D. Poyser Ltd, London.
- Gibbs, J.N. (1978) Intercontinental epidemiology of Dutch elm disease. *Annual Review of Phytopathology*, **16**, 287-307.
- Gibbs, J.N., Lipscombe, M.A. & Peace, A.J. (1999) The impact of *Phytophthora* disease on riparian populations of common alder (*Alnus glutinosa*) in southern Britain. *European Journal of Forest Pathology*, **29**, 39-50.
- Gibson, D.I. (1993) *Monobothrium wagneri* Nybelin, 1922: another imported tapeworm established in wild British freshwater fishes. *Journal of Fish Biology*, **43**, 281-285.
- Gilbert, O. (1998) Wildlife reports: lichens. *British Wildlife*, **10**, 130.
- Gilbert, O. (2000) *Lichens*. HarperCollins, London.
- Gledhill, T., Sutcliffe, D.W. & Williams, W.D. (1993) British freshwater Crustacea Malacostraca: a key with ecological notes. *Freshwater Biological Association Scientific Publications*, **52**, 1-173.
- Goater, B. (1986) *British pyralid moths - A guide to their identification*. Harley Books, Colchester.
- Godefroid, S. (2001) Temporal analysis of the Brussels flora as indicator for changing environmental quality. *Landscape and Urban Planning*, **52**, 203-224.
- Goode, D.A. (1989) Urban nature conservation in Britain. *Journal of Applied Ecology*, **26**, 859-873.
- Gosling, L.M. & Baker, S.J. (1989) The eradication of muskrats and coypus from Britain. *Biological Journal of the Linnean Society*, **38**, 39-51.
- Goulding, M.J., Roper, T.J., Smith, G.C. & Baker, S.J. (2003) Presence of free-living wild boar *Sus scrofa* in southern England. *Wildlife Biology*, **9**, 15-20.
- Gozlan, R.E., Flower, C.J. & Pinder, A.C. (2003a) Reproductive success in male sunbleak, a recent invasive fish species in the UK. *Journal of Fish Biology*, **63 (Supplement A)**, 131-151.
- Gozlan, R.E., Pinder, A.C., Durand, S. & Bass, J.A.B. (2003b) Could the small size of *Leucaspius delineatus* be an ecological advantage in invading British watercourses? *Folia Zoologica*, **52**, 99-108.
- Gozlan, R.E., Pinder, A.C. & Shelley, J. (2002) Occurrence of the Asiatic cyprinid *Pseudorasbora parva* in England. *Journal of Fish Biology*, **61**, 298-300.
- Gozlan, R.E., St-Hilaire, S., Feist, S.W., Martin, P. & Kents, M.L. (2005) Disease threat to European fish. *Nature*, **435**, 1046.
- Graddon, W.D. (1977) Some new discomycete species: 4. *Transactions of the British Mycological Society*, **69**, 255-273.
- Graddon, W.D. (1979) Discomycete notes and records 2. *Transactions of the British Mycological Society*, **73**, 180-188.
- Grapow, L.C. & Blasi, C. (1998) A comparison of the urban flora of different phytoclimatic regions in Italy. *Global Ecology and Biogeography*, **7**, 367-378.

- Green, P.R., Green, I.P. & Crouch, G.A. (1997) *The Atlas Flora of Somerset*. Privately published, Wayford & Yeovil.
- Green, P.R., Green, I.P. & Crouch, G.R. (1997) *The Atlas Flora of Somerset*. P.R. Green, I.P. Green, and G.A. Crouch.
- Greenhalgh, G.N. & Morgan-Jones, G. (1964) Some species of *Trochila* and an undescribed discomycete on leaves of *Prunus laurocerasus*. *Transactions of the British Mycological Society*, **47**, 311-320.
- Gregory, S.J. (2000) The land-hopper (*Arcitalitrus dorrieni*) in Britain. *British Myriapod & Isopod Newsletter*, **1**.
- Gregory, S.J. (2000) Non-marine Isopod Recording Scheme. *British Myriapod and Isopod Group Newsletter*, **1**, 3.
- Gregory, S.J. (2009) *Woodlice and Waterlice (Isopoda:Oniscidae & Asellota) in Britain and Ireland*. Field Studies Council, Shrewsbury.
- Hackett, D. (1998) The fern weevil, *Syagrius intrudens* Waterhouse (Curculionidae) in Cornwall. *The Coleopterist*, **7**, 20-21.
- Haes, E.C.M. & Harding, P.T. (1997) *Atlas of grasshoppers, crickets and allied insects in Britain and Ireland*. The Stationary Office, ITE research publication no.11. London.
- Halstead, A.J. (1981) A whitefly pest of sweet bay. *Plant Pathology*, **30**, 123.
- Halstead, A.J. (1992) The 1991 Presidential Address - Part 2. Some horticultural pests new to Britain in recent years. *British Journal of Entomology and Natural History*, **5**, 41-47.
- Halstead, A.J. (1996) Possible breeding by the rosemary beetle, *Chrysolina americana*, in Britain. *British Journal of Entomology and Natural History*, **9**, 107-108.
- Halstead, A.J. (2004) *Berberis* sawfly *Arge berberidis* Schrank (Hymenoptera: Argidae), a pest new to Britain. *British Journal of Entomology and Natural History*, **17**, 131-136.
- Halstead, A.J. & Malumphy, C.P. (2003) Outbreak in Britain of *Stephanitis takeyai* Drake & Mao (Hemiptera, Tingidae), a pest of *Pieris japonica*. *British Journal of Entomology and Natural History*, **16**, 3-6.
- Hancock, E.G. (1983) Further record of *Craspedacusta sowerbyi* Lankester in Lancashire. *Naturalist*, **108**, 119-120.
- Hänfling, B. & Harley, M. (2003) A molecular approach to detect hybridisation between crucian carp (*Carassius carassius*) and non indigenous carp species (*Carassius auratus* and *Cyprinus carpio*) in UK waters, including a consideration of the taxonomic status of gibel carp (*Carassius* spp.). pp. 32. Bristol: Environment Agency R&D Technical Report W2-077/TR.
- Harding, P.T. & Collis, G.M. (2006) The occurrence of *Asellus communis* Say, 1818 (Crustacea, Isopoda) at Bolam Lake, Northumberland. *Bulletin of the British Myriapod and Isopod Group*, **21**, 8-11.
- Hardy, F.G. & Guiry, M.D. (2003) *A check-list and atlas of the seaweeds of Britain and Ireland*. British Phycological Society, London.
- Harlequin Ladybird Survey (2005) The Harlequin Ladybird Survey website <http://www.harlequin-survey.org>.
- Harper, M. (2000) At war with aliens. pp. 11. Plantlife, London.
- Harper, M.W., Langmaid, J.R. & Emmet, A.M. (2002) Oecophoridae. *The Moths and Butterflies of Great Britain and Ireland, Volume 1 (Part1)* (eds A. M. Emmet & J. R. Langmaid), pp.

- 43-177. Harley Books, Colchester.
- Harris, S., Morris, P., Wray, S. & Yalden, D. (1995) *A review of British mammals: population estimates and conservation status of British mammals other than cetaceans*. JNCC, Peterborough.
- Harris, S.A. (2002) Introduction of Oxford ragwort, *Senecio squalidus* L. (Asteraceae), to the United Kingdom. *Watsonia*, **24**, 31-43.
- Harvey, P.R., Nellist, D.R. & Telfer, M.G. (2002) *Provisional atlas of British spiders (Arachnida, Araneae)*. Biological Records Centre, Huntingdon.
- Hasle, G.R. (1983) *Thalassiosira punctigera* (Castr.) comb. nov., a widely distributed marine planktonic diatom. *Nordic Journal of Botany*, **3**, 593-608.
- Hassel, K. & Soderstrom, L. (2005) The expansion of the alien mosses *Orthodontium lineare* and *Campylopus introflexus* in Britain and continental Europe. *Journal of the Hattori Botanical Laboratory*, 183-193.
- Hawkins, R.D. (2001) The Southern Oak Bush-cricket, *Meconema meridionale* Costa (Orthoptera: Tettigoniidae) new to Britain. *British Journal of Entomology and Natural History*, **14**, 207-213.
- Hawkins, R.D. (2002) The hunt for the Southern Oak Bush-cricket *Meconema meridionale* Costa. *Atropos*, **17**, 25-27.
- Hawksworth, D.L. (2001) *The changing wildlife of Great Britain and Ireland*. pp. xv+454. Taylor & Francis, London.
- Heal, N.F. (2003) *Scolytus pygmaeus* (Fabricius, 1787) (Scolytidae) – a new arrival to Britain. *The Coleopterist*, **12**, 57-60.
- Heath, J., Brown, D.J.F. & Boag, B. (1977) *Provisional atlas of the nematodes of the British isles*. Institute of Terrestrial Ecology, Huntingdon.
- Hedenas, L., Herben, T., Rydin, H. & Soderstrom, L. (1989) Ecology of the invading moss species *Orthodontium lineare* in Sweden - substrate preference and interactions with other species. *Journal of Bryology*, **15**, 565-581.
- Henderson, D.M. (2000) *A checklist of the rust fungi of the British Isles*. British Mycological Society, Kew.
- Henderson, D.M. & Bennell, A.P. (1979) British Rust Fungi: additions and corrections. *Notes from the Royal Botanic Garden Edinburgh*, **37**, 475-501.
- Henry, C.M., Bell, G.J. & Hill, S.A. (1992) The effect of methyl bromide fumigation on rhizomania inoculum in the field. *Plant Pathology*, **41**, 483-489.
- Heppell, D. (1961) The naturalization in Europe of the quahog, *Mercenaria mercenaria* (L.). *Journal of Conchology*, **25**, 21-34.
- Herben, T. (1994) Local rate of spreading and patch dynamics of an invasive moss species, *Orthodontium lineare*. *Journal of Bryology*, **18**, 115-125.
- Hill, M.O. (2004) Rare and interesting bryophytes in Britain and Ireland. *Field Bryology*, **83**, 45-47.
- Hill, M.O., Ames, S., Bacon, J., Botham, M.S., Lindsley-Leake, S., Marchant, J.H., Preston, C.D., Rehfisch, M.M., Roy, D.B., Roy, H.E., Tyler-Walters, H. & Wright, L. (2008) *Non-native Species Portal Feasibility Study*. Report to Defra. CEH Monks Wood, Huntingdon.
- Hill, M.O., Baker, R., Broad, G., Chandler, P.J., Copp, G.H., Ellis, J., Jones, D., Hoyland, C., Laing, I., Longshaw, M., Moore, N., Parrott, D., Pearman, D., Preston, C., Smith, R.M. & Waters,

- R. (2005) Audit of non-native species in England. *English Nature Research Reports*, **662**, 1-81.
- Hill, M.O., Cooper, J.M., Hiscock, K., Marchant, J.H., Preston, C.D., Rehfish, M.M., Roy, D.B. & Tyler-Walters, H. (2007) Surveillance of non-native and newly-arrived species: a proposal for a non-native species portal for Britain. *Report to Defra*, pp. 1-41. Centre for Ecology and Hydrology. 41pp, Report to Defra. Huntingdon.
- Hill, M.O. & Edwards, B. (2003) *Mosses and liverworts of Dorset*. Dorset Environmental Records Centre, Dorchester.
- Hill, M.O., Preston, C.D. & Smith, A.J.E. (1991) *Atlas of the bryophytes of Britain and Ireland, Vol. 1. Liverworts (Hepaticae and Anthocerotae)*. Harley Books, Colchester.
- Hill, M.O., Preston, C.D. & Smith, A.J.E. (1992) *Atlas of the bryophytes of Britain and Ireland, Vol. 2. Mosses (except Diplolepideae)*. Harley Books, Colchester.
- Hill, M.O., Preston, C.D. & Smith, A.J.E. (1994) *Atlas of the bryophytes of Britain and Ireland, Vol. 3. Mosses (Diplolepideae)*. Harley Books, Colchester.
- Hill, M.O., Roy, D.B. & Thompson, K. (2002) Hemeroby, urbanity and ruderality: bioindicators of disturbance and human impact. *Journal of Applied Ecology*, **39**, 708-720.
- Hill, M.O., Wright, S.M., Dring, J.C., Firbank, L.G., Manchester, S.J. & Croft, J.M. (1994) The potential for spread of alien species in England following climatic change. *English Nature Research Report*, **90**, 1-76.
- Hillyard, P.D. (2005) *Harvestmen*. Field Studies Council, Shrewsbury.
- Hipkin, C. (2003) Putting our alien flora into perspective. *British Wildlife*, **14**, 413-422.
- Hitchmough, J. & Woudstra, J. (1999) The ecology of exotic herbaceous perennials grown in managed, native grassy vegetation in urban landscapes. *Landscape and Urban Planning*, **45**, 107-121.
- Hodkinson, D.J. & Thompson, K. (1997) Plant dispersal: the role of man. *Journal of Applied Ecology*, **34**, 1484-1496.
- Hodkinson, I.D. (1999) Biocontrol of eucalyptus psyllid *Ctenarytaina eucalypti* by the Australian parasitoid *Psyllaephagus pilosus*: a review of current programmes and their success. *Biocontrol News and Information*, **20**, 129N – 134N <http://pest.cabweb.org/PDF/BNI/bnira51.pdf>.
- Hodkinson, I.D. & Hollis, D. (1980) *Floria variegata* Löw (Homoptera: Psylloidea) in Britain. *Entomologist's Gazette*, **31**, 171-172.
- Hodkinson, I.D. & White, I.M. (1979) *Homoptera, Psylloidea. Handbooks for the Identification of British Insects, volume 2, part 5a*. Royal Entomological Society of London, London.
- Holcík, J., Hensel, K., Nieslanik, J. & Skácel, L. (1988) *The Eurasian huchen, Hucho hucho*. Junk, Dordrecht.
- Holdich, D., Sibley, P. & Peay, S. (2004) The White-clawed Crayfish - a decade on. *British Wildlife*, **15**, 153-164.
- Holdich, D.M. & Reeve, I.D. (1991) The distribution of freshwater crayfish in the British Isles with particular reference to crayfish plague, alien introductions and water quality. *Aquatic Conservation*, **1**, 139-158.
- Holling, M. & Rare Breeding Birds Panel (2007) Non-native breeding birds in the United Kingdom in 2003, 2004 and 2005. *British Birds*, **100**, 638–649.
- Hollingsworth, M.L. & Bailey, J.P. (2000) Hybridisation and clonal diversity in some introduced

- Fallopia* species (Polygonaceae). *Watsonia*, **23**, 111-121.
- Hollis, D. (1978) *Floria variegata* Löw (Homoptera: Psylloidea) on Laburnum in Britain. *Plant Pathology*, **27**, 149.
- Hopkin, S.P. (1991) A key to the woodlice of Britain and Ireland. *Field Studies*, **7**, 599-650.
- Hopkins, D. (1997) A tree aphid (*Tinocallis nevskyi* Remaudière, Quednau & Heie) new to Britain (Hem., Hom., Aphidoidea). *Entomologist's Monthly Magazine*, **133**, 255-256.
- Hopkins, J. (2003) How is the countryside changing? *British Wildlife*, **14**, 305-310.
- Howden, J.C.W. & Jacobs, L. (1973) Report on the rust work at Bath. *The Rose Annual*, **1973**, 113-119.
- Hulme, P.E. (2009) Trade, transport and trouble: managing invasive species pathways in an era of globalization. *Journal of Applied Ecology*, **46**, 10-18.
- Hulme, P.E., Roy, D.B., Cunha, T. & Larsson, T.-B. (2009) A pan-European inventory of alien species: rationale, implementation and implications for managing biological invasions. *Handbook of alien species in Europe* (ed DAISIE), pp. 1-14. Springer, Dordrecht.
- Hussey, N.W. (1961) *Megastigmus flies attacking conifer seed*. Forestry Commission Leaflet No. 8.
- Hutchinson, M. (1974) *Andricus lignicola* (Hartig) (Hymenoptera, Cynipidae) in S.E. England, a new species to Britain. *Entomologist's Record and Journal of Variation*, **86**, 158-159.
- Hyman, P.S. (1987) *Otiorhynchus aurifer* Boheman (Col., Curculionidae) new to the British Isles. *Entomologist's Monthly Magazine*, **123**, 59.
- Hyman, P.S. (1987b) *Bruchela rufipes* (Olivier), (Col., Anthribidae) rediscovered in Great Britain. *Entomologist's Monthly Magazine*, **123**, 90.
- ICES WGITMO (2003) Report of working group on introductions and transfers of marine organisms <http://www.ices.dk/reports/general/2004/ICESCOP2004.pdf>.
- Ing, B. (1985) *Pisolithus* in Ireland. *Bulletin of the British Mycological Society*, **19**, 57-58.
- Ing, B. (1999) *The myxomycetes of Britain and Ireland: an identification handbook*. The Richmond Publishing Co, Slough.
- Ingle, R.W. (1986) The Chinese mitten crab *Eriocheir sinensis* H. Milne Edwards - a contentious immigrant. *The London Naturalist*, **65**, 101-105.
- Inman, A.J., Cook, R.T.A. & Beales, P.A. (2000) A contribution to the identity of rhododendron powdery mildew in Europe. *Journal of Phytopathology*, **148**, 17-27.
- Irvine, L.M. (1983) *Seaweeds of the British Isles. Volume 1 Rhodophyta Part 2A Cryptonemiales (sensu stricto), Palmariales, Rhodomenciales*. British Museum (Natural History), London.
- Irwin, A.G., Cole, J.H. & Ely, W.A. (2001) *Pelomyia occidentalis* Williston (Dip.: Tethinidae) new to Britain and Germany. *The Entomologist's Record and Journal of Variation*, **113**, 153-156.
- Ismay, J. & Smith, D. (1994) *Prosopantrum flavifrons* (Tonnoir and Malloch) (Diptera, Heleomyzidae) new to Britain and the Northern Hemisphere. *Dipterists Digest - second series*, **1**, 1-5.
- IUCN (2000).
- James, C.M. & al., e. (2000) A new hybrid between a European and a Chinese species of *Artemisia* (Asteraceae). *Watsonia*, **23**, 139-147.
- James, T.J. (2009) *Flora of Hertfordshire*. Hertfordshire Natural History Society.
- Jermyn, S.T. (1975) *Flora of Essex*. Essex Wildlife Trust.

- Joerstad, K.E., Farestveit, E., Rudra, H., Agnalt, A.-L. & Olsen, S. (2002) Studies on Red King Crab (*Paralithodes camtschaticus*) introduced to the Barents Sea. *Crabs in cold water regions: biology, management, and economics* (eds A. J. Paul, E. G. Dawe, R. Elnor, G. S. Jamieson, G. H. Kruse, R. S. Otto, B. Sainte-Marie, T. C. Shirley & D. Woodby), pp. 425-438. University of Alaska, 19th Lowell Wakefield Fisheries Symposium. Fairbanks.
- John, D.M. (2002) Order Cladophorales (=Siphonocladales). *The freshwater algal flora of the British Isles* (eds D. M. John, B. A. Whitton & A. J. Brook), pp. 468-470. Cambridge University Press, Cambridge.
- Johnson, C. (1966) *Caenoscelis subdeplanata* Bris. (Col., Cryptophagidae): a beetle new to Britain. *The Entomologist*, **99**, 129-131.
- Johnson, C. & Booth, R.G. (2004) *Luperomorpha xanthodera* (Fairmaire): a new British flea beetle (Chrysomelidae) on garden centre roses. *The Coleopterist*, **13**, 81-86.
- Jolly, R.L. (2000) The predatory mite *Neoseiulus californicus*: its potential as a biocontrol agent for the fruit tree red spider mite *Panonychus ulmi* in the UK. *The BCPC conference - Pests and diseases*, **2000**, 487-489.
- Jones, H.D. (2003) Land flatworms and nemertines in Cornwall. *Kovadha Kernow*, **7**, 3-5 <http://www.ercis.co.uk/download/kovadha7.pdf>.
- Jones, H.D. (2005) Identification: British land flatworms. *British Wildlife*, **16**, 189-194.
- Jones, H.D. & Boag, B. (1996) The distribution of New Zealand and Australian terrestrial flatworms (Platyhelminthes: Turbellaria: Tricladida: Terricola) in the British Isles - The Scottish survey and MEGALAB WORMS. *Journal of Natural History*, **30**, 955-975.
- Jones, P.E. (1980) Provisional atlas of the Arachnida of the British Isles. Part I Pseudoscorpions. Biological Records Centre, Institute of Terrestrial Ecology, Huntingdon.
- Jones, R.A. (1993) The rhododendron lacebug, *Stephanitis rhododendri* Horvath, rediscovered in south-east London. *British Journal of Entomology and Natural History*, **6**, 139-140.
- Jones, R.A. (2001) *Ctenochaeres bicolorus* (L.), an African ichneumonid (Hymenoptera) found in Britain. *British Journal of Entomology and Natural History*, **14**, 96-99.
- Jones, R.A. (2004) *Tephritis praecox* (Loew) (Diptera, Tephritidae) established in Britain. *Dipterists Digest*, **11**, 16.
- Jones, R.E. (1989) On a new species of centipede (Chilopoda, Geophilomorpha) from the Isles of Scilly. *J. Natural History*, **23**, 627-633.
- Karran, A.B. & Rich, T.C.G. (2003) Geographical and temporal distributions of *Alyssum alyssoides* and *Berteroa incana* (Brassicaceae) in the British Isles and the relationship to their modes of introduction. *Watsonia*, **24**, 499-506.
- Keller, R.P., Ermgassen, P.S.E., Aldridge, D.C., Lawrence, W.F. & Wright, S.J. (2009) Vectors and timing of freshwater invasions in Great Britain. *Conservation Biology*, **23**, 1526-1534.
- Kendall, P. (1993) *Bruchela rufipes* (Olivier) (Anthribidae) in the north of England. *The Coleopterist*, **2**, 34.
- Kendle, A.D. & Rose, J.E. (2000) The aliens have landed! What are the justifications for 'native only' policies in landscape plantings? *Landscape and Urban Planning*, **47**, 19-31.
- Kennedy, C.R. (1993) Introductions, spread and colonization of new localities by fish helminth and crustacean parasites in the British Isles: a perspective and appraisal. *Journal of Fish Biology*, **43**, 287-301.
- Kennedy, C.R. & Fitch, D.J. (1990) Colonisation, larval survival, and epidemiology of the

- nematode *Anguillicola crassus*, parasite in the eel *Anguilla anguilla* in Britain. *Journal of Fish Biology*, **36**, 117-131.
- Kent, D.H. (1975) *The Historical Flora of Middlesex : An Account of the Wild Plants Found in the Watsonian Vice-County 21 from 1548 to the Present Time*. The Ray Society.
- Kent, M., Stevens, R.A. & Zhang, L. (1999) Urban plant ecology patterns and processes: a case study of the flora of the city of Plymouth, Devon, U.K. *Journal of Biogeography*, **26**, 1281-1298.
- Kerney, M. (1999) *Atlas of land and freshwater molluscs of Britain and Ireland*. Harley Books, Colchester.
- Kevan, D.K.M. (1953) Notes on the distribution of British Orthopteroids. *Journal of the Society for British Entomology*, **4**, 119-122.
- Kibby, G. (2000) Fungal portraits No. 2: *Amanita inopinata*. *Field Mycology*, **1**, 39-40.
- Kibby, G. (2005) The invasion of *Amanita inopinata* continues. *Field Mycology*, **6**, 31.
- Killick, J., Perry, R. & Woodell, S. (1998) *The Flora of Oxfordshire*. Pisces Publications.
- Kirby, P. (1983) *Neodicyphus rhododendri* (Dolling) (Hem., Miridae) in Derbyshire. *Entomologist's Monthly Magazine*, **119**, 116.
- Kirby, P., Stewart, A.J.A. & Wilson, M.R. (2001) True bugs, leaf- and planthoppers, and their allies. *The changing wildlife of Great Britain and Ireland* (ed D. L. Hawksworth), pp. 262-299. Taylor and Francis, Systematics Association Special Volumes Series 62. London.
- Kirk, P.M. (1981) New or interesting microfungi. III. A preliminary account of microfungi colonising *Laurus nobilis* leaf litter. *Transactions of the British Mycological Society*, **77**, 457-473.
- Kirk, R.S. & Lewis, J.W. (1994) Sanguinicoliasis in cyprinid fish in the UK. *Parasitic diseases of fish* (eds A. W. Pike & J. W. Lewis), pp. 101-117. Samara Publishing Limited, Dyfed.
- Kitchener, G.D. (2003) The relationship between hybridisation in *Epilobium* and Cornish china clay and other mining waste. *Botanical Cornwall*, **12**, 20-32.
- Klaar, M., Copp, G.H. & Horsfield, R. (2004) Autumnal habitat use of non-native pumpkinseed *Lepomis gibbosus* and associations with native fish species in small English streams. *Folia Zoologica*, **53**, 189-202.
- Knight-Jones, P., Knight-Jones, E.W., Thorp, C.H. & Gray, P.W.G. (1975) Immigrant spirorbids (Polychaeta: Sedentaria) on the Japanese *Sargassum* at Portsmouth, England. *Zoologica Scripta*, **4**, 145-149.
- Knops, J.M.H., Griffin, J.R. & Royalty, A.C. (1995) Introduced and native plants of the Hastings Reservation, central coastal California. *Biological Conservation*, **71**, 115-123.
- Kowarik, I. (1990) Some responses of flora and vegetation to urbanization in Central Europe. *Urban Ecology: Plants and plant communities in urban environments* (eds H. Sukopp, S. Hejny [y with acute accent] & I. Kowarik), pp. 45-74. SPB Academic Publishing, Amsterdam.
- Kowarik, I. (1995) On the role of alien species in urban flora and vegetation. *Plant invasions: general aspects and special problems* (eds P. Pysek [inv circumflex on s], K. Prach, M. Rejmánek & M. Wade), pp. 85-103. SPB Academic Publishing, Amsterdam.
- Kucera, J.C.c. (1999) *Didymodon australasiae* var. *umbrosus* in the Czech Republic, with a review of recent records from Central Europe. *Journal of Bryology*, **21**, 71-72.
- Lack, P. (1986) *The atlas of wintering birds in Britain and Ireland*. T. & A.D. Poyser, Calton.

- Laing, F. (1922) *Rhinocola eucalypti* Mask. in England. *Entomologist's Monthly Magazine*, **58**, 141.
- Laing, I. & Gollash, S. (2002) *Coscinodiscus wailesii* - A nuisance diatom in European waters. *Invasive aquatic species of Europe - Distribution, impacts and management* (eds E. Leppäkoski, S. Gollasch & S. Olenin), pp. 53-55. Kluwer Academic Publishers, Dordrecht.
- Lambdon, P.W. (2008) Why is habitat breadth correlated strongly with range size? Trends amongst the alien and native floras of Mediterranean islands. *Journal of Biogeography*, **35**, 1095-1105.
- Lansdown, R.V. (2008) Red duckweed (*Lemna turionifera* Landolt) new to Britain. *Watsonia*, **27**, 127-130.
- Last, F.T. & Watling, R. (1998) First record of *Laccaria fraterna* in Britain. *Mycologist*, **12**, 152-153.
- Lawton, J.H. & Eastop, V.F. (1975) A bracken-feeding *Macrosiphum* new to Britain. *Entomologist's Gazette*, **26**, 135-138.
- Le Maitre, D.C., Van Wilgen, B.W., Chapman, R.A. & McKelly, D.H. (1996) Invasive plants and water resources in the Western Cape Province, South Africa: modelling the consequences of a lack of management. *Journal of Applied Ecology*, **33**, 161-172.
- Le Sueur, F. (1984) *Flora of Jersey*. Société Jersiaise, St Helier.
- Lee, P. (2006) *Atlas of the Millipedes (Diplopoda) of Britain and Ireland*. Pensoft, Sofia-Moscow.
- Leeke, C. (2005) Introduced reptiles and amphibians in the UK. Website <http://www.threadnaught.net/~caleb/ukintro.html>.
- Lever, C. (1977) *The naturalised animals of the British Isles*. Hutchinson, London.
- Lever, C. (1996) *The naturalised fishes of the world*. Academic Press, San Diego.
- Levey, B. (2004) <http://www.coleopterist.org.uk/>.
- Lewis, J.G.E., Jones, R.E. & Keay, A.N. (1988) On a new species and genus of centipede (Chilopoda, Geophilomorpha, Chilenophilidae) from the British Isles. *J.Natural History*, **22**, 1657-1663.
- Lloret, F., Médail, F., Brundu, G., Camarda, I., Moragues, E., Rita, J., Lambdon, P. & Hulme, P.E. (2005) Species attributes and invasion success by alien plants on Mediterranean islands. *Journal of Ecology*, **93**, 512-520.
- Lodge, D.M. (1993) Species invasions and deletions: community effects and responses to climate and habitat change. *Biotic interactions and global change* (eds P. M. Kareiva, J. G. Kingsolver & R. B. Huey), pp. 367-387.
- Long, J.L. (1981) *Introduced birds of the world: the worldwide history, distribution and influence of birds introduced to new environments*. David & Charles, London.
- Lososová, Z., Chytrý, M. & Kühn, I. (2008) Plant attributes determining the regional abundance of weeds on central European arable land. *Journal of Biogeography*, **35**, 177-187.
- Lott, D. (2005) <http://www.coleopterist.org.uk/>.
- Lousley, J.E. (1971) *Flora of the Isles of Scilly*. David & Charles, Newton Abbot.
- Lowe, A.J. & Abbott, R.J. (2003) A new British species, *Senecio eboracensis* (Asteraceae), another hybrid derivative of *S. vulgaris* L. and *S. squalidus* L. *Watsonia*, **24**, 375-388.
- Luff, M.L. (1998) *Provisional atlas of the ground beetles (Coleoptera, Carabidae) of Britain*. Biological Records Centre, Huntingdon.
- Luff, M.L. & Duff, A.G. (2004) <http://www.coleopterist.org.uk/>.

- Lyszkowski, R.M., Owen, J.A. & Taylor, S. (1992) *Corticaria abietorum* Motschulsky (Col.: Lathridiidae) new to Britain. *Entomologist's Record and Journal of Variation*, **104**, 67-69.
- Mabey, R. (1973) *The unofficial countryside*. Collins, London.
- MacNeil, C., Dick, J.T.A., Elwood, R.W. & Montgomery, W.I. (2001) Coexistence among native and introduced freshwater amphipods (Crustacea); habitat utilization patterns in littoral habitats. *Archiv Fur Hydrobiologie*, **151**, 591-607.
- Maggs, C.A. & Guiry, M.D. (1987) An Atlantic population of *Pikea californica* (Dumontiaceae, Rhodophyta). *Journal of Phycology*, **23**, 170-176.
- Maggs, C.A. & Hommersand, M.H. (1993) *Seaweeds of the British Isles. Volume 1 Rhodophyta Part 3A Ceramiales*.
- Mahindapala, R. (1978) Occurrence of Maize Rust, *Puccinia sorghi*, in England. *Transactions of the British Mycological Society*, **70**, 393-399.
- Maitland, P.S. (2004) *Keys to the freshwater fish of Great Britain and Ireland with notes on their distribution and ecology*. Freshwater Biological Association, Scientific Publication No. 62. Ambleside.
- Maitland, P.S. & Price, C.E. (1969) *Urocleidus principalis*, a North American monogenetic trematode new to the British Isles, probably introduced with the largemouth bass (*Micropterus salmoides*). *Journal of Fish Biology*, **1**, 17-18.
- Malumphy, C. (1997) Laurel scale, *Aonidia lauri* (Bouché) (Homoptera, Coccoidea, Diaspididae), a pest of bay laurel, new to Britain. *Entomologist's Gazette*, **48**, 195-198.
- Malumphy, C. (2003) The status of *Bemisia afer* (Priesner & Hosny) in Britain (Homoptera: Aleyrodidae). *Entomologist's Gazette*, **54**, 191-196.
- Malumphy, C. (2005) *Eulecanium excrescens* (Ferris) (Hemiptera: Coccidae), an Asian pest of woody ornamentals and fruit trees, new to Britain. *British Journal of Entomology and Natural History*, **18**, 45-49.
- Malumphy, C. & Halstead, A. (2003) *Cacopsylla fulguralis* (Kuwayama), an Asian plant louse (Hemiptera: Psyllidae), causing damage to *Elaeagnus* in Britain. *British Journal of Entomology and Natural History*, **16**, 89-93.
- Manchester, S.J. & Bullock, J.M. (2000) The impacts of non-native species on UK biodiversity and the effectiveness of control. *Journal of Applied Ecology*, **37**, 845-864.
- Mann, D. (2004) <http://www.coleopterist.org.uk/>.
- Marks, P.L., Wesley, F.R. & Gardescu, S. (2008) The vascular plant diversity of the Finger Lakes region of central New York State: changes in the 1800s and 1900s. *Journal of the Torrey Botanical Society*, **135**, 53-69.
- Marlin (2005) http://www.marlin.ac.uk/marine_aliens.
- Marquiss, M. & Newton, I. (1982) The Goshawk in Britain. *British Birds*, **75**, 243-260.
- Marquiss, M., Petty, S.J., Anderson, D.I.K. & Legge, G. (2003) Contrasting population dynamics of the Northern goshawk (*Accipiter gentilis*) in the Scottish/English Borders and north-east Scotland. *Birds of prey in a changing environment* (eds D. B. A. Thompson, S. M. Redpath, A. H. Fielding, M. Marquiss & C. A. Galbraith), pp. 143-148. The Stationery Office, Edinburgh.
- Marren, P. (2006) The 'global fungal weeds': the toadstools of wood-chip beds. *British Wildlife*, **18**, 98-105.
- Marshall, J.A. (2001) Grasshoppers, crickets and allies. *The changing wildlife of Great Britain*

- and Ireland* (ed D. L. Hawksworth), pp. 328-339. Taylor and Francis, Systematics Association Special Volumes Series 62. London.
- Marshall, J.A. & Haes, E.C.M. (1988) *Grasshoppers and allied insects of Great Britain and Ireland*. Harley Books, Colchester.
- Martin, J. (1993) *Aphis meliloti* (Borner) and *A. oenotherae* Oestlund, two species new to Britain (Homoptera: Aphididae). *Entomologist's Gazette*, **44**, 221-224.
- Martin, J.H. (1981) A new species of *Acyrtosiphon* (Homoptera: Aphididae) from *Primula* in Britain. *Systematic Entomology*, **6**, 97-101.
- Martin, J.H. (1996) *Aphis spiraeicola* Patch, an aphid pest of woody hosts now occurring on *Cotoneaster* in Britain (Homoptera, Aphididae). *Entomologist's Gazette*, **47**, 51- 52.
- Martin, J.H. (2000) Two new British aphid introductions in 1999, in the context of other additions over the preceding thirty years (Sternorrhyncha: Aphidoidea). *Entomologist's Gazette*, **51**, 97-105.
- Martin, J.H. & Malumphy, C.P. (1995) *Trioza vitreoradiata*, a New Zeland jumping plant louse (Homoptera: Psylloidea), causing damage to *Pittosporum* spp. in Britain. *Bulletin of Entomological Research*, **85**, 253-258.
- Martyn, K.P. (1988) *Provisional atlas of the ticks (Ixodidae) of the British Isles*. Biological Records Centre, Huntingdon.
- Maskell, L.C., Bullock, J.M., Smart, S.M., Thompson, K. & Hulme, P.E. (2006) The distribution and habitat associations of non-native plant species in urban riparian habitats. *Journal of Vegetation Science*, **17**, 499-508.
- Maskell, L.C., Firbank, L.G., Thompson, K., Bullock, J.M. & Smart, S.M. (2006) Interactions between non-native plant species and the floristic composition of common habitats. *Journal of Ecology*, **94**, 1052-1060.
- McClintock, D. (1975) *The wild flowers of Guernsey*. Collins, London.
- McCosh, D. & Rich, T. (2011) *Atlas of British and Irish Hawkweeds*. Botanical Society of the British Isles.
- McDonald, J. (1999) *Alien pests of bamboo. As identified from a recent survey of bamboo at holdings across the UK. Plant Pest Notice No.28*. Unpublished, Central Science Laboratory, Sand Hutton.
- McGavin, G.C. (1982) A new genus of Miridae (Hem.: Heteroptera). *Entomologist's Monthly Magazine*, **118**, 79-86.
- McGeoch, M.A., Chown, S.L. & Kalwij, J.M. (2006) A global indicator for biological invasion. *Conservation Biology*, **20**, 1635-1646.
- McMillan, N.F. (1938) Early records of *Crepidula* in English waters. *Proceedings of the Malacological Society*, **23**, 236.
- McNamara, D. (1996) A note on *Bacillus rossius*, the Corsican stick insect. *Bulletin of the Amateur Entomologists' Society*, **55**, 31-32.
- Meikle, R.D. (1984) *Willows and Poplars of Great Britain and Ireland*. BSBI Publications.
- Mendel, H. (2004) <http://www.coleopterist.org.uk/>.
- Menzel, F., Smith, J.E. & Colauto, N.B. (2003) *Bradysia difformis* Frey and *Bradysia ocellaris* (Comstock): two additional neotropical species of black fungus gnats (Diptera: Sciaridae) of economic importance: a redescription and review. *Annals of the Entomological Society of America*, **96**, 448-457.

- Menzies, I. & Spooner, B. (2000) *Henosepilachna argus* (Geoffroy) (Coccinellidae, Epilachninae), a phytophagous ladybird new to the U.K., breeding at Molesey, Surrey. *Coleopterist*, **9**, 1-4.
- Merrifield, K. & Merrifield, R. (2000) A second record of *Chetostoma curvinerve* (Rondani) (Diptera, Tephritidae) from Cornwall. *Dipterists Digest*, **7**, 20.
- Meyer, A.H. & Schmid, B. (1999) Seed dynamics and seedling establishment in the invading perennial *Solidago altissima* under different experimental treatments. *Journal of Ecology*, **87**, 28-41.
- Miller, N.G. & Trigoboff, N. (2001) A European feather moss, *Pseudoscleropodium purum*, naturalized widely in New York State cemeteries. *The Bryologist*, **104**, 98-103.
- Minter, D.W. (1980) Two species of *Lophomerum* on *Rhododendron* leaves. *Transactions of the British Mycological Society*, **74**, 201-204.
- Minter, D.W., Staley, J.M. & Millar, C.S. (1978) Four species of *Lophodermium* on *Pinus sylvestris*. *Transactions of the British Mycological Society*, **71**, 295-301.
- Monro, C.C.A. (1924) A serpulid polychaete from London docks (*Mercierella enigmatica* Fauvel). *Annals and Magazine of Natural History*, **13**, 155-159.
- Moon, H.P. (1970) *Corophium curvispinum* (Amphipoda) recorded again in the British Isles. *Nature*, **226**, 976.
- Moore, J.A. (1986) *Charophytes of Great Britain and Ireland*. Botanical Society of the British Isles, London.
- Moore, W.C. (1959) *British parasitic fungi*. Cambridge University Press, Cambridge.
- Mordue, J.E.M. & Ainsworth, G.C. (1984) *Ustilaginales of the British Isles*. *Mycological Papers no. 154*. Commonwealth Mycological Institute, Kew.
- Morgan, I. (1995) Comments on asilids. *Larger Brachycera Recording Scheme Newsletter*, **13**, 4-5.
- Morgan, M.J. (1979) *Pycnomerus fuliginosus* Er. (Col., Colydiidae) in Devon. *Entomologist's Monthly Magazine*, **114**, 166.
- Morris, R.K.A. (1999) *Chetostoma curvinerve* Rondani (Diptera, Tephritidae) in Surrey. *Dipterists Digest*, **6**, 84.
- Murphy, R.J. (2009) *Fumitories of Britain and Ireland*. Botanical Society of the British Isles.
- Murphy, R.J. & Rumsey, F.J. (2005) *Cystopteris diaphana* (Bory) Blasdel (Woodsiaceae) - an overlooked native new to the British Isles? *Watsonia*, **25**, 255-263.
- Nash, D.R. (2000) *Bruchela rufipes* (Olivier) (Col.: Urodontidae) in west Suffolk. *Entomologist's Record and Journal of Variation*, **112**, 223.
- Nau, B.S. (1980) *Tuponia carayoni* Wagner (Hem., Miridae) new to Britain. *Entomologist's Monthly Magazine*, **116**, 83-84.
- Naylor, E. (1957) Immigrant marine animals in Great Britain. *New Scientist*, **2**, 21-53.
- NBN Species Dictionary (2005) <http://nbn.nhm.ac.uk/nhm/>.
- Newell, G.E. (1949) The occurrence of a species of *Clymenella* Verrill (Polychaeta, fam. Maldanidae) on the north coast of Kent. *Nature*, **163**, 648-649.
- Newton, A. & Randall, R.D. (2004) *Atlas of British and Irish Brambles: A Phytogeographical Analysis of Microspecies of Rubus Sect. Rubus & Sect. Corylifolii*. BSBI Publications.
- Nightingale, B. (2005) The status of Lady Amherst's Pheasant in Britain. *British Birds*, **98**, 20-25.
- Norman, E. (1987) Aliens and adventives. An alien *Silene* in Scotland. *BSBI News*, 26-26.

- Ogilvie, M. & Panel, t.R.B.B. (2003) Non-native birds breeding in the United Kingdom in 2001. *British Birds*, **96**, 620-325.
- Ogilvie, M. & the Rare Breeding Birds Panel (2004) Non-native Birds Breeding in the UK 2002. *British Birds*, **97**, 633-637.
- Olden, J.D. (2006) Biotic homogenization: a new research agenda for conservation biogeography. *Journal of Biogeography*, **33**, 2027-2039.
- Oros, M., Hanzelova, V. & Scholz, T. (2004) The cestode *Atractolytocestus huronensis* (Caryophyllidea) continues to spread in Europe: new data on the helminth parasite of the common carp. *Diseases of Aquatic Organisms*, **62**, 115-119.
- Orton, J.H. & Winckworth, R. (1928) The occurrence of the American oyster pest *Urosalpinx cinerea* (Say) on English oyster beds. *Nature*, **122**, 241.
- Orton, P.D. (1986) *British Fungus Flora Agarics and Boleti. 4 Pluteaceae: Pluteus & Volvariella*. Royal Botanic Garden, Edinburgh.
- Owen, J.A. (1987) *Paratillus carus* Newman (Col. Cleridae) in the open at Windsor. *Entomologist's Record and Journal of Variation*, **99**, 155-156.
- Owen, J.A. (1993) *Magdalis memnonia* (Gyllenhal) (Col.: Curculionidae) in north-west Surrey. *Entomologist's Record and Journal of Variation*, **105**, 283-284.
- Owen, J.A., Allen, A.A., Carter, I.S. & von Hayek, C.M.F. (1985) *Panspoeus guttatus* Sharp (Col., Elateridae) new to Britain. *Entomologist's Monthly Magazine*, **121**, 91-95.
- Palmer, M. (1994) *A UK plant conservation strategy: a strategic framework for the conservation of the native flora of Great Britain and Northern Ireland*. Joint Nature Conservation Committee, Peterborough.
- Parsons, M. (2003) The changing moth and butterfly fauna of Britain during the twentieth century. *The Entomologist's Record and Journal of Variation*, **115**, 49-66.
- Paton, J.A. (1967) *Riccia crystallina* L. and *Riccia cavernosa* Hoffm. in Britain. *Transactions of the British Bryological Society*, **5**, 222-225.
- Paton, J.A. (1992) *Telaranea longii* sp. nov. in Britain, and a comparison with *T. murphyae* Paton. *Journal of Bryology*, **17**, 289-295.
- Paton, J.A. (1999) *The liverwort flora of the British Isles*. Harley Books, Colchester.
- Paton, J.A. & Holyoak, D.T. (2005) *The bryophyte flora of the Isles of Scilly*. Environmental Records Centre for Cornwall and the Isles of Scilly, Truro.
- Payne, R.M. (2002) *The flora of Ely*. Privately published, King's Lynn.
- Pearman, D. (2006) Invasive aliens? Himalayan balsam (alas, now Indian balsam). *BSBI News*, **103**, 31-32.
- Pegler, D.N., Læssøe, T. & Spooner, B.M. (1995) *British puffballs, earthstars and stinkhorns. An account of the British gasteroid fungi*. Royal Botanic Gardens, Kew.
- Pegler, D.N. & Legon, N.W. (1998a) Profiles of fungi. 92 *Agrocybe putaminum*. *Mycologist*, **12**, 60.
- Pegler, D.N. & Legon, N.W. (1998b) Profiles of fungi. 97 *Stropharia aurantiaca*. *Mycologist*, **12**, 180.
- Pegler, D.N. & Legon, N.W. (1998c) Profiles of fungi. 98 *Psilocybe cyanescens*. *Mycologist*, **12**, 180-181.
- Pegler, D.N., Spooner, B.M. & Young, T.W.K. (1993) *British truffles. A revision of British hypogeous fungi*. Royal Botanic Garden, Kew.

- Pelham-Clinton, E.C. (1985) Tineidae. *The Moths and Butterflies of Great Britain and Ireland, Volume 2* (eds J. Heath & A. M. Emmet), pp. 152-207. Harley Books, Colchester.
- Peretti, J.H. (1998) Nativism and nature: rethinking biological invasion. *Environmental Values*, **7**, 183-192.
- Perrins, J., Fitter, A.H. & Williamson, M. (1993) Population biology and rates of invasion of three introduced *Impatiens* species in the British Isles. *Journal of Biogeography*, **20**, 33-44.
- Petersen, K.S., Rasmussen, K.L., Heinemeler, J. & Rud, N. (1992) Clams before Columbus? *Nature*, **359**, 679.
- Philp, E.G. (2010) *New Atlas of the Kent Flora*. Kent Field Club.
- Pinder, A.C. & Gozlan, R.E. (2003) Sunbleak and topmouth gudgeon - two new additions to Britain's freshwater fishes. *British Wildlife Dec*, 77-83.
- Plant, C.W. (1994) *Provisional atlas of the lacewings and allied insects (Neuroptera, Megaloptera, Raphidioptera and Mecoptera) of Britain and Ireland*. Biological Records Centre, Huntingdon.
- Plantlife (2000) *At war with aliens*. Plantlife, London.
- Polaszek, A. & Cotman, H.E. (1983) An aphid new to Britain from *Robinia pseudacacia* L. in London. *Entomologist's Monthly Magazine*, **119**, 251-252.
- Ponder, W.F. (1988) *Potamopyrgus antipodarum*: a molluscan coloniser of Europe and Australia. *Journal of Molluscan Studies*, **54**, 271-285.
- Porley, R.D. & Matcham, H.W. (2003) The status of *Orthotrichum gracile* in Britain and Ireland. *Journal of Bryology*, **25**, 64-66.
- Pratt, M.A. (1979) Potato wart disease and its legislative control in England and Wales. *Plant health* (eds D. L. Ebbels & J. E. King), pp. 199-212. Blackwell Scientific Publications, Oxford.
- Preece, T.F., Weber, R.W.S. & Webster, J. (2000) Origin and spread of the daisy rust epidemic in Britain caused by *Puccinia distincta*. *Mycological Research*, **104**, 576-580.
- Preston, C.D. (1986) An additional criterion for assessing native status. *Watsonia*, **16**, 83.
- Preston, C.D. (2008) The aquatic plants of the River Cam and its riparian commons, Cambridge, 1660-1999. *Nature in Cambridgeshire*, **50**, 18-37.
- Preston, C.D. & Croft, J. (1997) *Aquatic Plants in Britain and Ireland*. Harley Books.
- Preston, C.D. & Croft, J.M. (1998) Britain's changing aquatic flora. *British Wildlife*, **10**, 18-28.
- Preston, C.D., Pearman, D.A. & Dines, T.D. (2002) New atlas of the British and Irish Flora. Oxford University Press, Oxford.
- Preston, C.D., Pearman, D.A. & Hall, A.R. (2004) Archaeophytes in Britain. *Botanical Journal of the Linnean Society*, **145**, 257-294.
- Prichard, A. (2004) *Cameraria ohridella*, Horse Chestnut Leaf Miner, arrives in Suffolk. <http://www.suffolkmothgroup.org.uk/newsletters/33/>. *Suffolk Moth Group Newsletter*, **33**.
- Prichard, A. (2005) *Ectoedemia heringella* (Mariani, 1939) - yet another new leaf miner for the county. <http://www.suffolkmothgroup.org.uk/newsletters/36/>. *Suffolk Moth Group Newsletter*, **36**.
- Pyšek, A., Pyšek, P., Jarošík, V., Hájek, M. & Wild, J. (2003) Diversity of native and alien plant species on rubbish dumps: effects of dump age, environmental factors and toxicity. *Diversity and Distributions*, **9**, 177-189.

- Pyšek, P. (1997) Clonality and plant invasions: can a trait make a difference? *The ecology and evolution of clonal plants* (eds H. de Kroon & J. van Groenendael), pp. 405-427. Backhuys, Leiden.
- Pyšek, P. (1998) Alien and native species in Central European urban floras: a quantitative comparison. *Journal of Biogeography*, **25**, 155-163.
- Pyšek, P. & Richardson, D.M. (2006) The biogeography of naturalization in alien plants. *Journal of Biogeography*, **33**, 2040-2050.
- Pyšek, P., Richardson, D.M., Rejmánek, M., Webster, G.L., Williamson, M. & Kirschner, J. (2004) Alien plants in checklists and floras: towards better communication between taxonomists and ecologists. *Taxon*, **53**, 131-143.
- Pyšek, P., Sádlo, J. & Mandák, B. (2002) Catalogue of alien plants of the Czech Republic. *Preslia*, **74**, 97-186.
- Quinlan, J. & Gauld, I.D. (1981) Symphyta (except Tenthredinidae) Hymenoptera. New Edition. *Handbooks for the Identification of British Insects*, **6**, 1-67.
- Read, R.W.J. (1981) *Furcipes rectirostris* (L.) (Coleoptera: Curculionidae) new to Britain. *Entomologist's Gazette*, **32**, 51-58.
- Read, R.W.J. (1997) *Euophryum confine* Broun (Curculionidae) in west Cumbria. *Coleopterist's Newsletter*, **8**, No. 42.
- Rebele, F. (1994) Urban ecology and special features of urban ecosystems. *Global Ecology and Biogeography Letters*, **4**, 173-187.
- Redfern, M., Shirley, P. & Bloxham, M. (2002) British plant galls: identification of galls on plants and fungi. *Field Studies*, **10**, 207-531.
- Reid, D.A. (1978) Two bambusicolous rust fungi new to Britain. *Transactions of the British Mycological Society*, **70**, 459-463.
- Reid, D.A. (1984) Another British record of *Puccinia longicornis*. *Bulletin of the British Mycological Society*, **18**, 127-129.
- Reid, D.A. (1987) New or interesting records of British hymenomycetes VII. *Notes from the Royal Botanic Garden, Edinburgh*, **44**, 503-540.
- Rejmánek, M. (1989) Invasibility of plant communities. *Biological invasions: a global perspective* (eds J. A. Drake, H. A. Mooney, F. Di Castri, R. H. Groves, F. J. Kruger, M. Rejmánek & M. Williamson). Wiley, Chichester.
- Renault, T. (1996) Appearance and spread of diseases among bivalve molluscs in the northern hemisphere in relation to international trade. *Revue Scientifique et Technique de l'Office International des Epizooties*, **15**, 551-561.
- Rendle, A.B. & West, W. (1899) A new British freshwater alga. *Journal of Botany British and Foreign*, **37**, 289-291.
- Rich, T., Houston, L. & Robertson, A. (2010) *Whitebeams, Rowans and Service Trees of Britain and Ireland: A Monograph of British and Irish Sorbus L.* Botanical Society of the British Isles.
- Rich, T.C.G. (1991) *Crucifers of Great Britain and Ireland*. Botanical Society of the British Isles, London.
- Rich, T.C.G. & Pryor, K.V. (2003) *Galeopsis segetum* Neck. (Lamiaceae), Downy Hemp-nettle: native or introduced in Britain? *Watsonia*, **24**, 401-412.
- Richardson, C.A., Seed, R., Al-Roumaihi, E. & McDonald, L. (1993) Distribution, shell growth and

- predation of the New Zealand oyster *Tiostrea* (= *Ostrea*) *lutaria* Hutton, in the Menai Strait, North Wales. *Journal of Shellfish Research*, **12**, 207-214.
- Ridley, G. (2000) The New Zealand connection - *Amanita inopinata* - the mystery deepens. *Field Mycology*, **1**, 117-118.
- Roach, F.A. (1985) *Cultivated fruits of Britain: their origin and history*. Basil Blackwell, Oxford.
- Roach, F.A. (1985) History and evolution of fruit crops. *HortScience*, **23**, 51-55.
- Robinson, G.S. (1979) Clothes-moths of the *Tinea pellionella* complex: a revision of the world's species (Lepidoptera: Tineidae). *Bulletin of the British Museum, Natural History (Entomology)*, **38**, 175-180.
- Roháček, J. & Marshall, S.A. (2000) A world revision of the seaweed fly genus *Thoracochaeta* Duda (Diptera: Sphaeroceridae: Limosiniinae). Part 2: Palaearctic species. *Studia Dipterologica*, **7**, 313-372.
- Roy, A.S. & Smith, I.M. (1994) Plum pox situation in Europe. *EPPO Bulletin*, **24**, 515-523.
- Roy, H., Rowland, F., Brown, P., Ware, R. & Majerus, M. (2005) Ecology of the Harlequin Ladybird. *British Wildlife*, **16**, 403-407.
- RPS Ecoscope Applied Ecologists (2005) Strategic control of non-native invasive aquatic plants. *English Nature Research Reports*, **686**, 1-54.
- Rumsey, F.J. (2001) *Achrophyllum dentatum* (Hook. f. & Wils.) Vitt & Crosby (Hookeriaceae) naturalized in Britain. *Journal of Bryology*, **23**, 341-344.
- Rushton-Mellor, S.K. (1992) Discovery of the fish louse, *Argulus japonicus* Thiele (Crustacea: Branchiura), in Britain. *Aquaculture and Fisheries Management*, **23**, 269-271.
- Ryves, T.B., Clement, E.J. & Foster, M.C. (1996) *Alien grasses of the British Isles*. Botanical Society of the British Isles, London.
- Salisbury, A. (2003a) Further records of *Chrysolina americana* (Linnaeus) (Chrysomelidae) in Britain. *The Coleopterist*, **12**, 61-62.
- Salisbury, A. (2003b) A further note on the continued spread in Britain of the Lily Beetle *Lilioceris lili* (Scopoli) (Chrysomelidae), with notes on its host plant range. *The Coleopterist*, **12**, 67-75.
- Salisbury, A. & Booth, R.G. (2004) *Rodalia cardinalis* (Mulsant), the Vedalia ladybird (Coleoptera: Coccinellidae) feeding on *Icerya purchasi* Maskell, cottony cushion scale (Hemiptera: Margarodidae), in London gardens. *British Journal of Entomology and Natural History*, **17**, 103-104.
- Salmon, E.S. & Ware, W.M. (1925) The downy mildew of hop and its epidemic occurrence in 1924. *Annals of Applied Biology*, **12**, 121-125.
- Sanford, M. (2007) Another new crop plant in Britain, *Solanum sisymbriifolium* Lam. *BSBI News*, **104**, 47-49.
- Sanford, M.N. & Fisk, R.J. (2010) *A Flora of Suffolk*. D K & M N Sanford.
- Scalera, R. & Zaghi, D. (2004) *Alien species and nature conservation in the EU. The role of the LIFE program*. Office for Official Publications of the European Communities, Luxembourg.
- Scholes, R.J. & Biggs, R. (2005) A biodiversity intactness index. *Nature*, **434**, 45-49.
- Scourfield, D.J. & Harding, J.P. (1966) A key to the British species of freshwater Cladocera, edn 3. *Freshwater Biological Association Scientific Publications*, **5**.
- Sell, P. (2007) Introduced 'look-alikes' and other difficult introduced plants in our

- Cambridgeshire flora. *BSBI News*, **105**, 24-30.
- Sell, P. & Murrell, G. (2006) *Flora of Great Britain and Ireland: Volume 4, Campanulaceae - Asteraceae*. Cambridge University Press.
- Shardlow, M.E.A. & Taylor, R. (2004) Is the Southern Green Shield Bug, *Nezara viridula* (L.) (Hemiptera: Pentatomidae) another species colonising Britain due to climate change? *The British Journal of Entomology and Natural History*, **17**, 143-146.
- Sharrock, J.T.R. (1976) *The Atlas of Breeding Birds in Britain and Ireland*. T. & A.D. Poyser Ltd, Calton, Staffordshire.
- Shaw, M.R. & Kasparyan, D.R. (2003) *Entomologists' Monthly Magazine*, **139**, 17-28.
- Shaw, P.J.A., Butlin, J. & Kibby, G. (2004) Fungi of ornamental woodchips in Surrey. *Mycologist*, **18**, 12-15.
- Shaw, P.J.A. & Kibby, G. (2001) Aliens in the flowerbeds. *Field Mycology*, **2**, 6-11.
- Sheail, J. (2003) Government and the management of an alien pest species: a British perspective. *Landscape Research*, **28**, 101-111.
- Sheath, R.G. & Sherwood, A.R. (2002) Phylum Rhodophyta (Red Algae). *The freshwater algal flora of the British Isles* (eds D. M. John, B. A. Whitton & A. J. Brook), pp. 123-143. Cambridge University Press, Cambridge.
- Shepherd, B. (1990) An invitation to the Isle of Wight. *B.S.B.I. News*, **55**, 9-10.
- Side, A.G. & Whitehouse, H.L.K. (1974) *Tortula amplexa* (Lesq.) Steere in Britain. *Journal of Bryology*, **8**, 15-18.
- Silvertown, J., Franco, M. & Harper, J.L. (1997) *Plant life histories: Ecology, phylogeny and evolution*. Cambridge University Press, Cambridge.
- Simpson, F.W. (1982) *Simpson's flora of Suffolk*. Suffolk Naturalists' Society.
- Simpson, V.R., Gibbons, L.M., Khalil, L.F. & Williams, J.L.R. (2005) Cholecystitis in otters (*Lutra lutra*) and mink (*Mustela vison*) caused by the fluke *Pseudamphistomum truncatum*. *The Veterinary Record*, **157**, 49-52.
- Sinker, C.A. (1985) *Ecological Flora of the Shropshire Region*. Shropshire Wildlife Trust, Shrewsbury.
- Sivanesan, A. (1977) British ascomycetes: *Endoxylina pini* sp. nov., *Scotiosphaeria endoxylinae* gen. et sp. nov. and *Didymosphaeria superapplanata* sp. nov. *Transactions of the British Mycological Society*, **69**, 117-123.
- Smale, M.C. (1990) Ecological role of buddleia (*Buddleja davidii*) in streambeds in Te Urewera National Park. *New Zealand Journal of Ecology*, **14**, 1-6.
- Smart, J. (2000) At war with aliens? *Plantlife Magazine*, **Summer 2000**, 3.
- Smayda, T.J. (1990) Novel and nuisance phytoplankton blooms in the sea: evidence for a global epidemic. *Toxic marine phytoplankton* (eds E. Granéli, B. Sundström, L. Edler & D. M. Anderson), pp. 29-40. Elsevier, Proceedings of the Fourth International Conference on Toxic Marine Phytoplankton, 26-30 June 1989, Lund, Sweden. New York.
- Smith, A.J.E. (2004) *The moss flora of Britain and Ireland*. Cambridge University Press, Cambridge.
- Smith, K.M. (1932) Studies on plant virus diseases XI. Further experiments with a ringspot virus; its identification with spotted wilt of tomato. *Annals of Applied Biology*, **19**, 305-320.
- Smith, P.A., Leah, R.T. & Eaton, J.W. (1998) A review of the current knowledge on the introduction, ecology and management of zander, *Stizostedion lucioperca*, in the UK.

- Stocking and Introduction of Fish* (ed I. G. Cowx), pp. 209-224. Blackwell Science Ltd., Fishing News Books. Oxford.
- Smith, P.R.J., Perrett, J., Garwood, P. & Moore, G. (1999) Two additions to the UK marine fauna: *Desdemona ornata* Banse, 1957 (Polychaeta, Sabellidae) and *Grandidierella japonica* Stephensen, 1938 (Amphipoda, Gammaridea). *Newsletter of the Porcupine Natural History Society*, **2**, 8-11.
- Smith, R.M., Baker, R.H.A., Malumphy, C.P., Hockland, S., Hammon, R.P., Ostojá-Starzewski, J.C. & Collins, D.W. (2005) Non-native invertebrate plant pests established in Great Britain: an assessment of patterns and trends. *BCPC Symposium Proceedings*, **81**, 119-124.
- Snell, C.A. (1991) Disappearance of Britain's tree frog *Hyla arborea* colonies. *British Herpetological Society Bulletin*, **38**, 40.
- Snow, K.R. (1986) Removal of *Aedes aegypti* (Linnaeus, 1762) from the British check list. *British Mosquito Group Newsletter*, **2**, 6.
- Sörensson, M. & Johnson, C. (2004) The first European records of the pantropical genus *Bambara* Vuillet, and a review of the immigrant featherwing beetles in Europe (Coleoptera: Ptiliidae). *Koleopterologische Rundschau*, **74**, 287-302.
- Southwood, T.R.E. & Leston, D. (1959) *Land and water bugs of the British Isles*. Warne, London.
- Spandl, K. (1998) Exploring the round houses of doves. *British Archaeology*, **35**, <http://www.britarch.ac.uk/ba/ba35/ba35feat.html#dovecotes>.
- Speight, M.R. & Nicol, M. (1984) Horse chestnut scale - a new urban menace? *New Scientist*, **1404**, 40-42.
- Spencer, B.E., Edwards, D.B., Kaiser, M. & Richardson, C.A. (1994) Spatfalls of the non-native Pacific oyster, *Crassostrea gigas*, in British waters. *Aquatic Conservation Marine and Freshwater Ecosystems*, **4**, 203-217.
- Spooner, B. (2005) *Aseroe* at Oxshott - a decade on. *Field Mycology*, **6**, 25-26.
- Spooner, B.M. (1981) New records and species of British microfungi. *Transactions of the British Mycological Society*, **76**, 265-301.
- Spooner, B.M. (1994) *Aseroe rubra* at Oxshott. *Mycologist*, **8**, 153.
- Stace, C.A. (1975) *Hybridization and the Flora of the British Isles*. London, New York, San Francisco.: Academic Press.
- Stace, C.A. (1997) *New flora of the British Isles*. Cambridge University Press, Cambridge.
- Stace, C.A. (2005) Plants found in Ireland but not in Britain. *Watsonia*, **25**, 296-298.
- Stead, D.E. (1996) Bacterial diseases of potato - future problems? *Proceedings Crop Protection in Northern Britain*, pp. 303-311.
- Stead, D.E. & Pemberton, A.W. (1987) Recent problems with *Pseudomonas syringae* pv. *pisi* in UK. *EPPO Bulletin*, **17**, 291-294.
- Sterling, D.H. (1989) *Sclerocona acutellus* (Eversmann) (Lepidoptera, Pyralidae) new to Britain. *Entomologist's Gazette*, **40**, 1-3.
- Stewart, A.J.A. (1993) *Placotettix taeniatifrons* (Kbm.) (Hem., Cicadellidae) in Bedfordshire. *Entomologist's Monthly Magazine*, **129**, 80.
- Stieperaere, H., Heylen, O. & Podoor, N. (1997) Differences in species composition of the bryophyte layer of some Belgian and Dutch pinewoods with and without the invading hepatic *Lophocolea semiteres* (Lehm.) Mitt. *Journal of Bryology*, **19**, 425-434.
- Stott, B. (1977) On the question of the introduction of the grass carp (*Ctenopharyngodon idella*

- Val.) into the United Kingdom. *Fisheries Management*, **8**, 63-71.
- Straw, N.A. & Bellet-Travers, M. (2004) Impact and management of the horse chestnut leaf-miner (*Cameraria ohridella*). *Arboricultural Journal*, **28**, 67-83.
- Stroyan, H.L.G. (1971) *Masonaphis* [*Illinoia*] *lambersi* MacGill.: an introduced aphid pest of hybrid rhododendrons. *Plant Pathology*, **20**, 196.
- Stroyan, H.L.G. (1972) Additions and amendments to the checklist of British aphids (Homoptera: Aphidoidea). *Transactions of the Royal Entomological Society of London*, **124**, 37-39.
- Stroyan, H.L.G. (1977) *Homoptera Aphidoidea (Part): Chaitophoridae and Callaphididae*.
- Stroyan, H.L.G. (1979) Additions to the British aphid fauna (Homoptera: Aphidoidea). *Zoological Journal of the Linnean Society*, **65**, 1-54.
- Stroyan, H.L.G. (1984) *Aphids – Pterocommatinae and Aphidinae (Aphidini)*.
- Stubbs, A.E. (2000) [re. *Phytomyza hellebori* Kalténbach]. *Dipterists' Digest*, **7**, 33-35.
- Styles, J. (1959) Notes on the collection of the adults and the web-spinning larvae of *Cephalia alpina* Klug. (Hym.: Pamphiliidae). *Entomologist's Monthly Magazine*, **95**, 152-153.
- Sutcliffe, D.W. (1972) Notes on the chemistry and fauna of water-bodies in Northumberland, with special emphasis on the distribution of *Gammarus pulex* (L.), *G. lacustris* Sars and *Asellus communis* Say (new to Britain). *Transactions of the Natural History Society of Northumberland*, **17**, 222-248.
- Sutton, B.C. & Pirozynski, K.A. (1963) Notes on British microfungi. I. *Transactions of the British Mycological Society*, **46**, 505-522.
- Swale, E.M.F. (1962) Notes on some algae from the Reddish Canal. *British Phycological Bulletin*, **2**, 174-176.
- Swan, G.A. (1993) *Flora of Northumberland*. The Natural History of Northumbria, Newcastle upon Tyne.
- Thébaud, C. & Simberloff, D. (2001) Are plants really larger in their introduced ranges? *American Naturalist*, **157**, 231-236.
- Thomas, S. (2010) Horizon-scanning for invasive non-native plants in Great Britain. *Natural England Commissioned Reports*.
- Thompson, K., Austin, K.C., Smith, R.M., Warren, P.H., Angold, P.G. & Gaston, K.J. (2003) Urban domestic gardens (I): putting small-scale plant diversity in context. *Journal of Vegetation Science*, **14**, 71-78.
- Thompson, K., Hodgson, J.G. & Rich, T.C.G. (1995) Native and alien invasive plants: more of the same? *Ecography*, **18**, 390-402.
- Thompson, R.T. & Styles, J.H. (1958) *Otioryncus niger* [now *coecus*] (F.) (Col., Curculionidae) in Britain. *Entomologist's Monthly Magazine*, **94**, 183.
- Thorp, C.H., Knight-Jones, P. & Knight-Jones, E.W. (1986) New records of tubeworms established in British harbours. *Journal of the Marine Biological Association of the United Kingdom*, **66**, 881-888.
- Thorp, C.H., Pyne, S. & West, S.A. (1987) *Hydroides ezoensis* Okuda, a fouling serpulid new to British coastal waters. *Journal of Natural History*, **21**, 863-877.
- Torchin, M.E., Lafferty, K.D., Dobson, A.P., McKenzie, V.J. & Kuris, A.M. (2003) Introduced species and their missing parasites. *Nature*, **421**, 628-630.
- Torchin, M.E., Lafferty, K.D. & Kuris, A.M. (2002) Parasites and marine invasions. *Parasitology*, **124**, S137-S151.

- Townsend, C.C. (2007) Some further remarks on *Pohlia flexuosa* in Europe, and on other propaguliferous species. *Journal of Bryology*, **29**, 194-195.
- Twinn, P.F.G. & Harding, P.T. (1999) *Provisional atlas of the longhorn beetles (Coleoptera, Cerambycidae) of Britain*. Biological Records Centre, Huntingdon.
- Uhthoff-Kaufmann, R.R. (1990) The occurrence of the sub-family *Aseminae* (Col.: Cerambycidae) in the British Isles. *The Entomologist's Record and Journal of Variation*, **102**, 55-63.
- Uhthoff-Kaufmann, R.R. (1990b) The distribution of the genera *Trinophylum* Bates, *Gracilia* Serv., *Aromia* Serv., and *Hylotrupes* Serv. (Col.: Cerambycidae) in the British Isles. *The Entomologist's Record and Journal of Variation*, **102**, 267-274.
- Uhthoff-Kaufmann, R.R. (1991) The distribution and occurrence of *Acanthocinus* Dej. and *Agapanthia* Serv. (Col.: Lamiidae) in the British Isles. *The Entomologist's Record and Journal of Variation*, **103**, 189-192.
- UK Government (1992) *Biodiversity: the UK action plan*. HMSO, London.
- UK Government (2004) *Plant diversity challenge*. Joint Nature Conservation Committee, Peterborough.
- UK Government (2008) *The invasive non-native species framework strategy for Great Britain*. Department for Environment, Food and Rural Affairs, London.
- Usher, M.B. (2000) The nativeness and non-nativeness of species. *Watsonia*, **23**, 323-326.
- Utting, S.D. & Spencer, B.E. (1992) Introduction of marine bivalve molluscs into the UK for commercial culture - case histories. *ICES Marine Science Symposium*, **194**, 84-91.
- Verkerk, R.H.F. & Bravery, A.F. (2000) The UK termite eradication programme, justification and implementation. *International Research Group on Wood Preservation 31st Annual Meeting Kona, Hawaii 14-19 May 2000*.
- Villeneuve, F., Copp, G.H., Fox, M.G. & Stakenas, S. (2005) Interpopulaton variation in the growth and life history traits of the introduced sunfish, pumpkinseed *Lepomis gibbosus*, in Southern England. *Journal of Applied Ichthyology*, (in press).
- Wadsworth, R.A., Collingham, Y.C., Willis, S.G., Huntley, B. & Hulme, P.E. (2000) Simulating the spread and management of alien riparian weeds: are they out of control? *Journal of Applied Ecology*, **37 (Suppl. 1)**, 28-38.
- Walker, A.J.M. (1972) *Goniadella gracilis*, a polychaete new to British seas. *Marine Biology*, **14**, 85-87.
- Walker, P. (2001) The developing community on the introduced oak, *Quercus cerris*, A catkin gall-forming wasp *Andricus grossulariae* Giraud (Hym., Cynipidae), new to Britain. *Entomologist's Monthly Magazine*, **137**, 145-147.
- Wallentinus, I. (1999) Introduction and transfer of plants. *Status of introductions of non-indigenous marine species to North Atlantic waters 1981-1990* (eds A. L. S. Munro, S. D. Utting & I. Wallentinus), pp. 1-43. Vol. 231, ICES co-operative research report.
- Walls, R. (1996) Aliens are not aggressive. *B.S.B.I. News*, **73**, 36,38 [p. 37 is from another article].
- Waring, P., Townsend, M. & Lewington, R. (2003) *Field Guide to the Moths of Great Britain and Ireland*. British Wildlife Publishing, Hook, Hampshire.
- Watling, R. (1970) *British Fungus Flora Agarics and Boleti. 1 Boletaceae: Gomphidiaceae: Paxillaceae*. Her Majesty's Stationery Office, Edinburgh.

- Watling, R. (2001) Larger fungi. *The changing wildlife of Great Britain* (ed D. L. Hawksworth), pp. 103-113. Taylor & Francis, London & New York.
- Watling, R. & Gregory, N.M. (1987) *British Fungus Flora Agarics and Boleti. 5 Strophariaceae & Coprinaceae p.p.* Royal Botanic Garden, Edinburgh.
- Watling, R. & Turnbull, E. (1998) *British Fungus Flora Agarics and Boleti. 8 Cantharellaceae, Gomphaceae and amyloid-spored and xeruloid members of Tricholomataceae (excl. Mycena).* Royal Botanic Garden, Edinburgh.
- Watson, G.W. & Malumphy, C.P. (2004) *Icerya purchasi* Maskell, cottony cushion scale (Hemiptera: Margarodidae), causing damage to ornamental plants growing outdoors in London. *British Journal of Entomology and Natural History*, **17**, 105-109.
- Weber, E. (1998) The dynamics of plant invasions: a case study of three exotic goldenrod species (*Solidago* L.) in Europe. *Journal of Biogeography*, **25**, 147-154.
- Weber, E.F. (1997) The alien flora of Europe: a taxonomic and biogeographic review. *Journal of Vegetation Science*, **8**, 565-572.
- Welch, D., Carss, D.N., Gornall, J., Manchester, S.J., Marquiss, M., Preston, C.D., Telfer, M.G. & Arnold, H. (2000) An audit of alien species in Scotland. ITE, Banchory.
- Welch, R.C. (1993) Colonisation of introduced oaks by Cynipidae. *Cecidology*, **8**, 58-76.
- Welch, R.C. (2000) *Phytomyza hellebori* Kaltenbach (Dip.: Agromyzidae), a recent addition to the British fauna: further records in east Northamptonshire, Huntingdonshire and Cambridgeshire. *The Entomologist's Record and Journal of Variation*, **112**, 163-166.
- Welcomme, R.L. (1988) International introductions of inland aquatic species. Rome: FAO Fisheries Technical Paper N° 294. FAO, 00100.
- Westbrook, M.A. (1930) Notes on the distribution of certain marine red algae. *Journal of Botany British and Foreign*, **68**, 257-264.
- Westbrook, M.A. (1934) *Antithamnion spirographidis* Schiffner. *Journal of Botany British and Foreign*, **72**, 65-68.
- Wheeler, A.C. (1977) The origin and distribution of the freshwater fishes of the British Isles. *Journal of Biogeography*, **4**, 1-24.
- Wheeler, A.C. (1978) *Ictalurus melas* (Rafinesque, 1820) and *I. nebulosus* (Lesueur, 1819): the North American catfishes in Europe. *Journal of Fish Biology*, **12**, 435-439.
- Wheeler, A.C. (1998) Ponds and fishes in Epping Forest, Essex. *The London Naturalist*, **77**, 107-146.
- Wheeler, A.C. (2000) Status of the crucian carp, *Carassius carassius* (L.), in the UK. *Fisheries Management and Ecology*, **7**, 315-322.
- Wheeler, A.C., Merrett, N.R. & Quigley, D.T.G. (2004) Additional records and notes for Wheeler's (1992) List of the Common and Scientific Names of Fishes of the British Isles. *Journal of Fish Biology*, **65 (Supplement B)**, 1-40.
- Whitehead, P.F. (1991) Some further records of localized British beetles. *Entomologist's Monthly Magazine*, **127**, 158.
- Widgery, J. (2002) New species breeding in Britain. *Orthoptera Recording Scheme for Britain and Ireland Newsletter*, **28**, 1-3.
- Wigginton, M.J. (1999) *British Red Data Books: Vascular Plants No 1*. Joint Nature Conservation Committee.
- Wilcox, M. & Tregale, J. (2008) *Juncus anthelatus* (Wiegand) R.E. Brooks in Britain. *BSBI News*,

- 108**, 50-53.
- Wilkinson, M.J. & Stace, C.A. (1988) The taxonomic relationships and typification of *Festuca brevipila* Tracey and *F. lemanii* Bastard (Poaceae). *Watsonia*, **17**, 289-299.
- Williams, D.J. (1984) Two injurious mealybugs new to Britain (Hem., Homoptera, Coccoidea, Pseudococcidae). *Entomologist's Monthly Magazine*, **120**, 227-228.
- Williams, D.J. (1985) Scale insects (Homoptera, Coccoidea) of Tresco, Isles of Scilly. *Entomologist's Gazette*, **36**, 135-143.
- Williams, W.D. (1972) Occurrence in Britain of *Asellus communis* Say, 1818, a North American freshwater isopod. *Crustaceana Supplement*, **3**, 134-138.
- Williamson, M. (1981) *Island Populations*. Oxford University Press, Oxford.
- Williamson, M. (1996) *Biological Invasions*. Chapman & Hall, London.
- Williamson, M., Preston, C. & Telfer, M. (2003) On the rates of spread of alien plants in Britain. *Plant invasions: ecological threats and management solutions* (eds L. E. Child, J. H. Brock, G. Brundu, K. Prach, P. Pyšek, P. M. Wade & M. Williamson), pp. 63-74. Backhuys, Leiden.
- Williamson, M.H. & Fitter, A. (1996) The characters of successful invaders. *Biological Conservation*, **78**, 163-170.
- Willis, K.J., Cook, E.J., Lozano-Fernandez, M. & Takeuchi, I. (2004) First record of the alien caprellid amphipod, *Caprella mutica*, in the U.K. *Journal of the Marine Biological Association of the United Kingdom*, **84**, 1027-1028.
- Wilmore, G.T.D. (2000) *Alien plants of Yorkshire*. The Yorkshire Naturalists' Union, Weymouth.
- Wilson (1981) Identification of European *Lassus* species (Homoptera, Cicadellidae) with one new species to Britain. *Systematic Entomology*, **6**, 115-118.
- Wilson, I.M., Walshaw, D.F. & Walker, J. (1965) The new groundsel rust in Britain and its relationship to certain Australasian rusts. *Transactions of the British Mycological Society*, **48**, 501-511.
- Wilson, J.B., Steel, J.B., Dodd, M.E., Anderson, B.J., Ullmann, I. & Bannister, P. (2000) A test of community reassembly using the exotic communities of New Zealand roadsides in comparison to British roadsides. *Journal of Ecology*, **88**, 757-764.
- Wilson, M. & Henderson, D.M. (1966) *British Rust Fungi*. Cambridge University Press, Cambridge.
- Winter, T.G. (1990) *Crypturgus subcribrosus* (Eggers)(Col., Scolytidae) a bark beetle new to Britain. *Entomologist's Monthly Magazine*, **126**, 209-211.
- Wolfe, L.M. (2002) Why alien invaders succeed: support for the escape-from-enemy hypothesis. *American Naturalist*, **160**, 705-711.
- Wolff, W.J. (2005) Non-indigenous marine and estuarine species in The Netherlands. *Zoologische Mededelingen*, **79**, 1-116.
- Wood, T.S. & Okamura, B. (2005) A new key to the freshwater bryozoans of Britain, Ireland and continental Europe, with notes on their ecology. *Freshwater Biological Association Scientific Publication*, **63**, 1-105.
- Wurzell, B. (1994) A history of *Conyza* in London. *BSBI News*, **65**, 35-38.
- Wurzell, B. (1995) *x Conyzigeron huelsenii* in east London. *BSBI News*, 32-33.
- Wurzell, B. (2002) The maple bladder gall of the gall mite *Vasates quadripedes* new to Britain. *Cecidology*, **17**, 31-35.

- Wycherley, J. (2003) Water frogs in Britain. *British Wildlife*, **14**, 260-269.
- Yalden, D. (1999) *The history of British Mammals*. University Press, Cambridge.
- Yeomans, W.E., Chubb, J.C. & Sweeting, R.A. (1997) *Khawia sinensis* (Cestoda: Caryophyllidea) - an indicator of legislative failure to protect freshwater habitats in the British Isles? *Journal of Fish Biology*, **51**, 880-885.
- Young, J.O. & Reynoldson, T.B. (1999) Continuing dispersal of freshwater triclads (Platyhelminthes; Turbellaria) in Britain with particular reference to lakes. *Freshwater Biology*, **42**, 247-262.
- Zibrowius, H. & Thorp, C.H. (1989) A review of the alien serpulid and spirorbid polychaetes in the British Isles. *Cahiers de Biologie Marine*, **30**, 271-285.

Appendix 1. Glossary

Alert species (see also sleeper species)

A species which the Non-native Species Programme Board has recommended requires particular attention in terms of surveillance and reporting. Examples of situations in which the Programme Board might recommend an alert species include:

- A high-risk invasive species not yet present in GB but which might enter in the future and cause environmental, economic or social harm.
- A sleeper species considered to be high risk and which might cause environmental, economic or social harm in the future.
- An invasive non-native species for which there is a special concerted effort to provide better-quality distribution information.
- An invasive non-native species against which action is being taken and reporting is required to help facilitate action.

Alert species can be reported using the alert e-mail (alert_nonnative@ceh.ac.uk) or the on-line recording system which results in a cascade of information to relevant stakeholders.

Alien species

See non-native species.

Database (see also species register)

An integrated collection of logically related records and files on non-native species consolidated into a network that provides data for multiple uses. The species register forms the central database of the GB-NNSIP. The NBN Gateway is the central database for species distribution data.

Daisie

Delivering Alien Invasive Species Inventory for Europe - <http://www.europe-aliens.org/>

This website provides a 'one-stop shop' for information on biological invasions in Europe. It is the result of the DAISIE project, funded by the European Commission under the Sixth Framework Programme. CEH maintains the DAISIE database and website.

Content management system (CMS)

A computer system used to edit, track and store electronic documents. The GB-NNSIP uses a CMS called Drupal. Drupal is a Content Management System (CMS) which is an online system for maintaining website content.

Factsheet

In the case of the GB-NNSIP a factsheet refers to the detailed information for each species delivered through the website. The factsheets will be downloadable as a PDF. The information is written for a wide audience but principally stakeholders such as government employees, scientists and volunteer contributors to biological recording schemes. Factsheets within the

GB-NNSIP contain the following information: Invasion History, Ecology and Habitat, Distribution, Impact, Management, Legislation, References and Links.

See for example:

http://www.nonnativespecies.org/01_Fact_File/05_Fact_Sheets.cfm

ID sheet

In the case of the GB-NNSIP the ID sheets are a series of PDFs (focussed purely on providing species identification guidance) that have been produced for the Non-Native Species Secretariat which users can access through the GB-NNSIP. An ID sheet provides information on a species in a structured format which is downloadable as a PDF from the Non-native Species Secretariat website: (http://www.nonnativespecies.org/02_Identification%20Sheets.cfm)

The information is written for a broader audience than the factsheets and includes images and text written for a non-expert audience.

Invasive non-native species (INNS) (= invasive alien species or IAS)

An invasive non-native species is a non-native species that has the ability to spread causing damage to the environment, the economy, our health or the way we live.

Non-native species (= alien species)

A non-native species (= alien species) is a species introduced (i.e. by human action) outside its natural past or present distribution; including any part, gametes, seeds, eggs or propagules of such species that might survive and subsequently reproduce as defined by the Convention on Biological Diversity (CBD). Non-native species covered by the GB-NNSIP include all fauna and flora with the exception of genetically modified organisms (GMOs), fungi, bacteria and viruses. Lower taxonomic ranks such as subspecies, varieties, races or provenances can also be non-native.

Portal

GB-NNSIP. Web-based platform providing a gateway to information, including other websites, factsheets, species nomenclature, and species distribution.

Sleeper species (see also alert species)

Sleeper species are non-native species that have arrived in Great Britain and have currently established only small populations but that have the potential to spread widely and threaten biological diversity.

Species established and persisting in the wild

The term “in the wild” is widely used and generally encompasses both natural and semi-natural habitats in both rural and urban environments. However, this and preceding projects (Hill *et al.*, 2005, Hill *et al.*, 2008) have not attempted to define the term “in the wild” but have only included species that occur outside buildings, captivity or cultivation. In this report we align with the definition provided by Natural England with respect to “the wild” (<http://www.defra.gov.uk/publications/2011/05/26/pb13535wildlife-countryside-act/>):

“The diverse range of natural and semi-natural habitats and their associated wild native flora and fauna in the rural and urban environments in general. This can also be broadly described as the general open environment.”

The term “established” is used for self-sustaining (reproducing) populations. A species is deemed to be “established” if it occurs as a self-sustaining population, persisting for more than four years, not dependent on repeated reintroduction. However, it is useful to include the term “persisting” for species, such as forest trees, that persist for more than four years but are not successfully reproducing.

Species register

Relational database (= central store) containing a list of all non-native species in Great Britain and associated information (excluding distribution information). Principal tables in the database include: Species, Synonyms, Common names, Native range, Habitats, Species in region (invasion information by geographical region), Pathways, Vectors, Donor areas, Date of Introduction etc.

Web services

Web services are web-based software applications that exchange data with other web-based applications. The NBN Gateway maps are a web service to the GB-NNSIP. For more information see:

<http://www.nbn.org.uk/getdoc/4db9ded1-6469-4e2d-a901-87b3686b064f/Web-services.aspx>

Appendix 2. Summary of participation by experts, nominated by volunteer zoological scheme and society organisers (and other relevant bodies), to complete the species register entries associated with different informal (species) groups. The environment relates to the main environment of the informal group: terrestrial (T), terrestrial-freshwater (TF) or freshwater (F). The number of non-native species for each informal group is listed alongside the scheme, society or other organisation (and a named individual) contributing information.

Informal group	Environment	Number of species	Scheme participation
higher plants	All	1873	Botanical Society of the British Isles
Bryophytes	All		British Bryological Society
acarine (Acari)	T	9	Health Protection Agency - Jolyon Medlock
Amphibian	TF	12	Amphibian and Reptile Conservation - John Wilkinson
Annelid	F	6	Earthworm Recording Scheme – Emma Sherlock
Bird	T	119	BTO
bony fish (Actinopterygii)	F	35	CEFAS and ecological consultant Dave Hubble
bryozoan	F	1	To complete
centipede	T	7	British Myriapod and Isopod Group (BMIG) – Tony Barber
coelenterate (=cnidarian)	F	1	Freshwater invertebrate expert - David Aldridge
crustacean	F	15	Freshwater invertebrate expert - David Aldridge
crustacean	T	3	BMIG – Tony Barber
flatworm (Turbellaria)	F	4	Terrestrial Flatworm Recording Scheme - Brian Boag and Hugh Jones
flatworm (Turbellaria)	T	2	Terrestrial Flatworm Recording Scheme - Brian Boag and Hugh Jones
harvestman (Opiliones)	T	2	To complete
insect - beetle (Coleoptera)	F	2	To complete
insect - beetle (Coleoptera)	T	348	Coleoptera Recording Scheme Organisers - Darren Mann, Dave Hubble, Helen Roy, Peter Brown
insect - beetle (Coleoptera)	TF	1	To complete
insect - booklouse (Psocoptera)	T	38	To complete - consulting scheme organiser - Keith Alexander
insect - butterfly	T	3	Lepidoptera expert - David Green
insect - cockroach (Dictyoptera)	T	4	To complete
insect - earwig (Dermaptera)	T	5	Orthoptera Recording Scheme - Björn Beckmann
insect - flea	T	7	To complete

(Siphonaptera)			
insect - hymenopteran (ant)	T	16	Bees Wasp and Ants Recording Society - Mike Fox
insect - hymenopteran (chalcid wasp)	T	16	Parasitic Wasp Recording Scheme Organiser - Gavin Broad
insect - hymenopteran (cynipid wasp)	T	6	Parasitic Wasp Recording Scheme Organiser - Gavin Broad
insect - hymenopteran (ichneumon)	T	1	Parasitic Wasp Recording Scheme Organiser - Gavin Broad
insect - hymenopteran (sawfly)	T	39	Royal Horticultural Society – Andrew Halstead
insect - lacewing (Neuroptera)	T	2	To complete
insect - moth (macro)	T	29	Lepidoptera expert - David Green
insect - moth (micro)	T	108	Lepidoptera expert - David Green
insect - orthopteran	T	4	Orthoptera Recording Scheme - Björn Beckmann
insect - silverfish (Thysanura)	T	1	To complete
insect - stick insect (Phasmida)	T	6	Orthoptera Recording Scheme - Björn Beckmann
insect - termite	T	1	To complete
insect - thrips (Thysanoptera)	T	19	Consulting FERA
insect - true bug (Hemiptera)	T	236	Auchenorrhyncha Scheme Organiser (Alan Stewart), Terrestrial heteroptera scheme organisers (Tristan Bantock with Bernard Nau), Psyllids and Coccids (RHS), Aphids (Rothamsted Research)
insect - true fly (Diptera - fruitfly)	T	6	Dipterists Forum - Ian McLean
insect - true fly (Diptera - fungus gnat)	T	1	Dipterists Forum - Ian McLean
insect - true fly (Diptera - gallfly)	T	5	Dipterists Forum - Ian McLean
insect - true fly (Diptera - gall-midge)	T	16	Royal Horticultural Society – Andrew Halstead
insect - true fly (Diptera - hoverfly)	T	9	Dipterists Forum - Ian McLean
insect - true fly (Diptera - leaf-miner fly)	T	1	Dipterists Forum - Ian McLean
insect - true fly (Diptera - lesser dungfly)	T	1	Dipterists Forum - Ian McLean

insect - true fly (Diptera - mosquito or gnat)	T	2	Mosquito Recording Scheme - Jolyon Medlock
insect - true fly (Diptera - parasite- fly)	T	5	Dipterists Forum - Ian McLean
insect - true fly (Diptera)	T	64	Dipterists Forum - Ian McLean
millipede	T	2	BMIG – Tony Barber
mollusc	F	8	Freshwater invertebrate expert - David Aldridge
mollusc	T	30	Ecological consultant - Dave Hubble
reptile	T	5	Amphibian and Reptile Conservation - John Wilkinson
reptile	TF	1	Amphibian and Reptile Conservation - John Wilkinson
ribbon worm (Nemertinea)	T	1	To complete
roundworm (Nematoda)	F	1	To complete
roundworm (Nematoda)	T	4	To complete
spider (Araneae)	T	44	Ecological consultant - Dave Hubble
springtail (Collembola)	T	7	Collembola Scheme Organiser - Peter Shaw
terrestrial mammal	T	41	BTO

Appendix 3. List of species selected for factsheets within the GB-NNSIP (294 species)

Species	Common name	Informal group	Phylum	Environment
<i>Acaena novae-zelandiae</i>	Pirri-Pirri Bur	flowering plant	Anthophyta	T
<i>Acer pseudoplatanus</i>	Sycamore	flowering plant	Anthophyta	T
	Giant African Land			
<i>Achatina fulica</i>	Snail	mollusc	Mollusca	T
<i>Aedes albopictus</i>	Asian Tiger Mosquito	insect - true fly (Diptera)	Arthropoda	T
<i>Aegopodium podagraria</i>	Ground-elder	flowering plant	Anthophyta	T
<i>Agapornis roseicollis</i>	Rosy-faced lovebird	bird	Chordata	T
<i>Agrilus planipennis</i>	Emerald Ash Borer	insect - beetle (Coleoptera)	Arthropoda	T
<i>Ailanthus altissima</i>	Tree of heaven	flowering plant	Anthophyta	T
<i>Aix galericulata</i>	Mandarin	bird	Chordata	T
<i>Aix sponsa</i>	Carolina Wood Duck	bird	Chordata	T
<i>Alectoris chukar</i>	Chukar	bird	Chordata	T
<i>Alectoris graeca</i>	Rock Partridge	bird	Chordata	T
<i>Alexandrium catenella</i>		alga	Dinophyta	M
<i>Allium paradoxum</i>	Few-flowered Garlic	flowering plant	Anthophyta	T
<i>Allium triquetrum</i>	Three-Cornered Garlic	flowering plant	Anthophyta	T
<i>Alopochen aegyptiacus</i>	Egyptian Goose	bird	Chordata	T
<i>Alytes obstetricans</i>	Midwife Toad	amphibian	Chordata	TF
<i>Ambloplites rupestris</i>	Rock Bass	bony fish (Actinopterygii)	Chordata	F
<i>Ambrosia artemisiifolia</i>	Ragweed	flowering plant	Anthophyta	T
<i>Ameiurus melas</i>	Black Bullhead	bony fish (Actinopterygii)	Chordata	F
<i>Amelanchier lamarckii</i>	Juneberry	flowering plant	Anthophyta	T
<i>Ammothea hilgendorfi</i>	Japanese sea spider			M
	Swimbladder			
<i>Anguillicola crassus</i>	Nematode	roundworm (Nematoda)	Nematoda	MF
<i>Anoplophora chinensis</i>	Citrus longhorn beetle	insect - beetle (Coleoptera)	Arthropoda	T
	Asian Longhorned			
<i>Anoplophora glabripennis</i>	Beetle	insect - beetle (Coleoptera)	Arthropoda	T
<i>Anser caerulescens</i>	Snow Goose	bird	Chordata	T
<i>Anser canagica</i>	Emperor Goose	bird	Chordata	T
<i>Anser indicus</i>	Bar-headed Goose	bird	Chordata	T
<i>Antithamnion nipponicum</i>		alga	Rhodophyta	M
		insect - true bug		
<i>Aphis gossypii</i>	Cotton aphid	(Hemiptera)	Arthropoda	T
	Blue-crowned			
<i>Aratinga acuticaudata</i>	Parakeet	bird	Chordata	T
<i>Arcitalitrus dorrieni</i>		crustacean	Arthropoda	T
<i>Arge berberidis</i>	Berberis Sawfly	insect - hymenopteran	Arthropoda	T
<i>Arion lusitanicus agg.</i>	Lusitanian Slug	mollusc	Mollusca	T
<i>Arthurdendyus triangulata</i>	New Zealand	flatworm (Turbellaria)	Platyhelminthes	T

	Flatworm			
<i>Arundo donax</i>	Giant Reed	flowering plant	Anthophyta	F
<i>Asparagopsis armata</i>	Harpoon weed	alga	Rhodophyta	M
<i>Astacus astacus</i>	Noble Crayfish	crustacean	Arthropoda	F
<i>Astacus leptodactylus</i>	Turkish Crayfish	crustacean	Arthropoda	F
<i>Australoplana sanguinea</i>	Australian Flatworm	flatworm (Turbellaria)	Platyhelminthes	T
	Carolina Mosquito			
<i>Azolla caroliniana</i>	Fern	fern	Pteridophyta	F
<i>Azolla filiculoides</i>	Water Fern	fern	Pteridophyta	F
<i>Balanus improvisus</i>	Barnacle	crustacean	Arthropoda	M
		insect - true bug		
<i>Bemisia tabaci</i>	Tobacco Whitefly	(Hemiptera)	Arthropoda	T
		insect - cockroach		
<i>Blatta orientalis</i>	Oriental Cockroach	(Dictyoptera)	Arthropoda	T
		insect - cockroach		
<i>Blattella germanica</i>	German Cockroach	(Dictyoptera)	Arthropoda	T
<i>Bombina variegata</i>	Yellow-bellied Toad	amphibian	Chordata	TF
<i>Bonnemaisonia hamifera</i>		alga	Rhodophyta	M
<i>Botrylloides cf. diegense</i>				M
<i>Botrylloides violaceus</i>		tunicate (Urochordata)	Chordata	M
<i>Branta canadensis</i>	Canada Goose	bird	Chordata	T
<i>Bubo bubo</i>	Eagle Owl	bird	Chordata	T
<i>Buddleja davidii</i>	Buddleia	flowering plant	Anthophyta	T
<i>Bugula neritina</i>	Brown Bryozoan	bryozoan		M
<i>Bursaphelenchus xylophilus</i>	Pine Wood Nematode	roundworm (Nematoda)	Nematoda	T
<i>Cabomba caroliniana</i>	Carolina Water-shield	flowering plant	Anthophyta	F
<i>Cairina moschata</i>	Muscovy Duck	bird	Chordata	T
	Horse chestnut leaf			
<i>Cameraria ohridella</i>	miner	insect - moth	Arthropoda	T
<i>Campanula rapunculoides</i>	Creeping Bellflower	flowering plant	Anthophyta	T
<i>Campylopus introflexus</i>	Heath Star Moss	moss	Bryophyta	T
<i>Capra hircus</i>	Feral Goat	terrestrial mammal	Chordata	T
	Japanese skeleton			
<i>Caprella mutica</i>	shrimp	crustacean	Arthropoda	M
<i>Carassius auratus</i>	Goldfish	bony fish (Actinopterygii)	Chordata	F
<i>Carpobrotus edulis</i>	Hottentot-Fig	flowering plant	Anthophyta	T
<i>Caulerpa racemosa</i>	Grape Alga	alga	Chlorophyta	M
<i>Caulerpa taxifolia</i>	Killer Alga	alga	Chlorophyta	M
<i>Ceratitis capitata</i>	Medfly	insect - true fly (Diptera)	Arthropoda	T
<i>Cervus nippon</i>	Sika Deer	terrestrial mammal	Chordata	T
<i>Chrysolina americana</i>	Rosemary Beetle	insect - beetle (Coleoptera)	Arthropoda	T
	Lady Amherst's			
<i>Chrysolophus amherstiae</i>	Pheasant	bird	Chordata	T
<i>Chrysolophus pictus</i>	Golden Pheasant	bird	Chordata	T

<i>Cicerbita macrophylla</i>	Blue Sow-Thistle	flowering plant	Anthophyta	T	
<i>Cinara cupressi</i>	Cypress Aphid	insect - true bug (Hemiptera)	Arthropoda	T	
<i>Claytonia sibirica</i>	Pink Purslane	flowering plant	Anthophyta	T	
<i>Codium fragile (fragile)</i>	Green sea fingers	alga	Chlorophyta	M	
<i>Contarinia quinquenotata</i>	Hemerocallis gall	midge	insect - true fly (Diptera)	Arthropoda	T
<i>Corbicula fluminea</i>	Asiatic Clam	mollusc	Mollusca	F	
<i>Cordylophora caspia</i>		coelenterate (=cnidarian)	Cnidaria	M	
<i>Corella eumyota</i>		tunicate (Urochordata)	Chordata	M	
<i>Cornus sericea</i>	Red-Osier Dogwood	flowering plant	Anthophyta	T	
<i>Corophium sextonae</i>		crustacean	Arthropoda	M	
<i>Cortaderia selloana</i>	Pampas Grass	flowering plant	Anthophyta	T	
<i>Corvus splendens</i>	Indian house crow	bird	Chordata	T	
<i>Coscinodiscus wailesii</i>		diatom	Chromista	M	
<i>Cotoneaster horizontalis</i>	Wall Cotoneaster	flowering plant	Anthophyta	T	
<i>Cotoneaster microphyllus</i> <i>s.str.</i>	Small-leaved Cotoneaster	flowering plant	Anthophyta	T	
<i>Cotoneaster simonsii</i>	Himalayan Cotoneaster	flowering plant	Anthophyta	T	
<i>Crangonyx pseudogracilis</i>	Northern River Crangonyctid	crustacean	Arthropoda	F	
<i>Crassostrea gigas</i>	Pacific oyster	mollusc	Mollusca	M	
<i>Crassula helmsii</i>	New Zealand Pigmyweed	flowering plant	Anthophyta	F	
<i>Crepidula fornicata</i>	Slipper limpet	mollusc	Mollusca	M	
<i>Crocasmia pottsii x aurea =</i> <i>C. x crocosmiiflora</i>	Montbretia	flowering plant	Anthophyta	T	
<i>Ctenopharyngodon idellus</i>	Chinese Grass Carp	bony fish (Actinopterygii)	Chordata	F	
<i>Cygnus atratus</i>	Black Swan	bird	Chordata	T	
<i>Cynomys ludovicianus</i>	Black-tailed Prairie Dog	terrestrial mammal	Chordata	T	
<i>Cyprinus carpio</i>	Carp	bony fish (Actinopterygii)	Chordata	F	
<i>Dama dama</i>	Fallow Deer	terrestrial mammal	Chordata	T	
<i>Datura stramonium</i>	Thorn-Apple	flowering plant	Anthophyta	T	
<i>Diabrotica virgifera</i>	Western Corn Rootworm	insect - beetle (Coleoptera)	Arthropoda	T	
<i>Didemnum vexillum</i>	Carpet Sea-squirt	tunicate (Urochordata)	Chordata	M	
<i>Dikerogammarus villosus</i>	Killer Shrimp	crustacean	Arthropoda	F	
<i>Disphyma crassifolium</i>	Purple Dewplant	flowering plant	Anthophyta	T	
<i>Doronicum pardalianches</i>	Leopard's-bane	flowering plant	Anthophyta	T	
<i>Dreissena bugensis</i>	Quagga Mussel	mollusc	Mollusca	F	
<i>Dreissena polymorpha</i>	Zebra Mussel	mollusc	Mollusca	F	
<i>Echinocystis lobata</i>	Wild Cucumber	flowering plant	Anthophyta	T	
<i>Egeria densa</i>	Large-Flowered	flowering plant	Anthophyta	F	

	Waterweed			
<i>Eichhornia crassipes</i>	Water Hyacinth	flowering plant	Anthophyta	F
<i>Eliomys quercinus</i>	Garden Dormouse	terrestrial mammal	Chordata	T
<i>Elminius modestus</i>		crustacean	Arthropoda	M
<i>Elodea canadensis</i>	Canadian Pondweed	flowering plant	Anthophyta	F
<i>Elodea nuttallii</i>	Esthwaite Water-Weed	flowering plant	Anthophyta	F
<i>Emys orbicularis</i>	European Pond Terrapin	reptile	Chordata	T
<i>Ensis americanus</i>	American jack knife clam	mollusc	Mollusca	M
<i>Epilobium brunnescens</i>	New Zealand Willowherb	flowering plant	Anthophyta	T
<i>Eriocheir sinensis</i>	Chinese Mitten Crab	crustacean	Arthropoda	M
<i>Eucalyptus (genus)</i>	Gum Tree	flowering plant	Anthophyta	T
<i>Fallopia baldschuanica</i>	Russian Vine	flowering plant	Anthophyta	T
<i>Fallopia japonica</i>	Janapese Knotweed	flowering plant	Anthophyta	T
<i>Fallopia japonica x sachalinensis = F. x bohemica</i>		flowering plant	Anthophyta	T
<i>Fallopia sachalinensis</i>	Giant Knotweed	flowering plant	Anthophyta	T
<i>Ficopomatus enigmaticus</i>		annelid	Annelida	M
<i>Frankliniella occidentalis</i>		insect - thrips (Thysanoptera)	Arthropoda	T
<i>Gammarus tigrinus</i>	Sideswimmer	crustacean	Arthropoda	MF
<i>Gaultheria shallon</i>	Shallon	flowering plant	Anthophyta	T
<i>Glis glis</i>	Edible Dormouse	terrestrial mammal	Chordata	T
<i>Graphocephala fennahi</i>	Rhododendron leafhopper	insect - true bug (Hemiptera)	Arthropoda	T
<i>Grateloupia filicina subsp. luxurians</i>		alga	Rhodophyta	M
<i>Grateloupia turuturu</i>	Devil's tongue weed	alga	Rhodophyta	M
<i>Gunnera manicata</i>	Brazilian Giant Rhubarb	flowering plant	Anthophyta	T
<i>Gunnera tinctoria</i>	Giant Rhubarb	flowering plant	Anthophyta	T
<i>Gyrodactylus salaris</i>	Salmon fluke	trematode	Platyhelminthes	F
<i>Harmonia axyridis</i>	Harlequin Ladybird	insect - beetle (Coleoptera)	Arthropoda	T
<i>Helianthus tuberosus</i>	Jerusalem Artichoke	flowering plant	Anthophyta	T
<i>Hemigrapsus sanguineus</i>	Japanese shore crab	crustacean	Arthropoda	M
<i>Hemigrapsus takanoi</i>	Brush-clawed shore crab	crustacean	Arthropoda	M
<i>Hemimysis anomala</i>	Bloody-red Mysid	crustacean	Arthropoda	F
<i>Heracleum mantegazzianum</i>	Giant Hogweed	flowering plant	Anthophyta	T
<i>Hyacinthoides hispanica</i>	Garden Bluebell	flowering plant	Anthophyta	T

<i>Hydrocotyle ranunculoides</i>	Floating Pennywort	flowering plant	Anthophyta	F
<i>Hydropotes inermis</i>	Chinese Water Deer	terrestrial mammal	Chordata	T
<i>Hyla arborea</i>	Common Tree Frog	amphibian	Chordata	TF
<i>Hystrix brachyura</i>	Hodgson's Porcupine	terrestrial mammal	Chordata	T
<i>Hystrix cristata</i>	European Porcupine	terrestrial mammal	Chordata	T
<i>Impatiens glandulifera</i>	Himalayan Balsam	flowering plant	Anthophyta	T
<i>Kontikia andersoni</i>		flatworm (Turbellaria)	Platyhelminthes	T
<i>Kontikia ventrolineata</i>		flatworm (Turbellaria)	Platyhelminthes	T
<i>Lacerta bilineata</i>	Western Green Lizard	reptile	Chordata	T
<i>Lagarosiphon major</i>	Curly Waterweed	flowering plant	Anthophyta	F
<i>Lamiastrum galeobdolon</i> <i>subsp. argentatum</i>		flowering plant	Anthophyta	T
<i>Lasius neglectus</i>	Asian Super Ant	insect - hymenopteran	Arthropoda	T
<i>Lemna minuta</i>	Least Duckweed	flowering plant	Anthophyta	F
<i>Lepomis gibbosus</i>	Pond-Perch	bony fish (Actinopterygii)	Chordata	F
<i>Leptinotarsa decemlineata</i>	Colorado Beetle	insect - beetle (Coleoptera)	Arthropoda	T
<i>Leptoglossus occidentalis</i>	Western conifer seed bug	insect - true bug (Hemiptera)	Arthropoda	T
<i>Leucaspis delineatus</i>	Belica	bony fish (Actinopterygii)	Chordata	F
<i>Leuciscus idus</i>	Ide	bony fish (Actinopterygii)	Chordata	F
<i>Lilioceris lillii</i>	Lily Beetle	insect - beetle (Coleoptera)	Arthropoda	T
<i>Linepithema humile</i>	Argentine ant	insect - hymenopteran	Arthropoda	T
<i>Liriomyza huidobrensis</i>	Pea leaf miner	insect - true fly (Diptera)	Arthropoda	T
<i>Lithobates catesbeianus</i>	American Bullfrog	amphibian	Chordata	TF
<i>Lonicera japonica</i>	Japanese Honeysuckle	flowering plant	Anthophyta	T
<i>Lophocolea semiteres</i>	Liverwort			
<i>Ludwigia peploides</i>	Creeping Water- primrose	flowering plant	Anthophyta	F
<i>Lymantria dispar</i>	Gypsy Moth	insect - moth	Arthropoda	T
<i>Lysichiton americanus</i>	American Skunk- cabbage	flowering plant	Anthophyta	T
<i>Macropus rufogriseus</i>	Red-necked Wallaby	terrestrial mammal	Chordata	T
<i>Mahonia aquifolium</i>	Oregon-grape	flowering plant	Anthophyta	T
<i>Marenzelleria neglecta</i>	Red-gilled Mud Worm	annelid	Annelida	M
<i>Marenzelleria viridis</i>	Red-gilled Mud Worm	annelid	Annelida	M
<i>Marsupenaeus japonicus</i>	Caribbean mud crab	crustacean	Arthropoda	M
<i>Melopsittacus undulatus</i>	Budgerigar	bird	Chordata	T
<i>Mercenaria mercenaria</i>	American hard-shelled clam	mollusc	Mollusca	M
<i>Meriones unguiculatus</i>	Mongolian Gerbil	terrestrial mammal	Chordata	T
<i>Mesotriton alpestris</i>	Alpine Newt	amphibian	Chordata	TF
<i>Micropterus salmoides</i>	Largemouth Bass	bony fish (Actinopterygii)	Chordata	F
<i>Mimulus guttatus</i>	Monkeyflower	flowering plant	Anthophyta	T
<i>Mnemiopsis leidyi</i>	comb jelly			M

<i>Muntiacus reevesi</i>	Muntjac	terrestrial mammal	Chordata	T
<i>Mus domesticus</i>	House Mouse	terrestrial mammal	Chordata	T
<i>Mustela furo</i>	Feral Ferret	terrestrial mammal	Chordata	T
<i>Mustela vison</i>	American Mink	terrestrial mammal	Chordata	T
<i>Myiopsitta monachus</i>	Monk Parakeet	bird	Chordata	T
<i>Myocastor coypus</i>	Coypu	terrestrial mammal	Chordata	T
<i>Myriophyllum aquaticum</i>	Parrot's Feather	flowering plant	Anthophyta	F
<i>Nasua nasua</i>	Coati	terrestrial mammal	Chordata	T
<i>Neogobius melanostomus</i>	Round Goby	bony fish (Actinopterygii)	Chordata	F
<i>Netta rufina</i>	Red-crested Pochard	bird	Chordata	T
<i>Nezara viridula</i>	Southern green stink bug	insect - true bug (Hemiptera)	Arthropoda	T
<i>Numida meleagris</i>	Helmeted Guinea fowl	bird	Chordata	T
<i>Nyctereutes procyonoides</i>	Raccoon Dog	terrestrial mammal	Chordata	T
<i>Nycticorax nycticorax</i>	Black-crowned Night Heron	bird	Chordata	T
<i>Odontella sinensis</i>		diatom	Chromista	M
<i>Ondatra zibethicus</i>	Musk Rat	terrestrial mammal	Chordata	T
<i>Orconectes limosus</i>	Spinycheek Crayfish	crustacean	Arthropoda	F
<i>Orthodontium lineare</i>	Cape Thread-moss	moss	Bryophyta	T
<i>Oryctolagus cuniculus</i>	Rabbit	terrestrial mammal	Chordata	T
<i>Oxalis pes-caprae</i>	Bermuda-buttercup	flowering plant	Anthophyta	T
<i>Oxyura jamaicensis</i>	Ruddy Duck	bird	Chordata	T
<i>Pachygrapsus marmoratus</i>	Marbled shore crab	crustacean	Arthropoda	M
<i>Pacifastacus leniusculus</i>	Signal Crayfish	crustacean	Arthropoda	F
<i>Paralithodes camtschaticus</i>	Red King Crab	crustacean	Arthropoda	M
<i>Parthenocissus inserta</i>	False Virginia-Creeper	flowering plant	Anthophyta	T
<i>Parthenocissus quinquefolia</i>	Virginia-creeper	flowering plant	Anthophyta	T
<i>Paspalum distichum</i>	Water Finger-grass	flowering plant	Anthophyta	T
<i>Pelophylax esculentus</i>	Edible Frog	amphibian	Chordata	TF
<i>Pelophylax ridibundus</i>	Marsh frog	amphibian	Chordata	TF
<i>Periplaneta americana</i>	American Cockroach	insect - cockroach (Dictyoptera)	Arthropoda	T
<i>Periplaneta australasiae</i>	Australian Cockroach	insect - cockroach (Dictyoptera)	Arthropoda	T
<i>Persicaria campanulata</i>	Lesser Knotweed	flowering plant	Anthophyta	T
<i>Persicaria wallichii</i>	Himalayan Knotweed	flowering plant	Anthophyta	T
<i>Petasites albus</i>	White Butterbur	flowering plant	Anthophyta	T
<i>Petasites fragrans</i>	Winter Heliotrope	flowering plant	Anthophyta	T
<i>Petasites japonicus</i>	Giant Butterbur	flowering plant	Anthophyta	T
<i>Phagocata woodworthi</i>		flatworm (Turbellaria)	Platyhelminthes	F
<i>Phasianus colchicus</i>	Common Pheasant	bird	Chordata	T
<i>Phasianus versicolor</i>	Green Pheasant	bird	Chordata	T

<i>Picea sitchensis</i>	Sitka Spruce	conifer	Pinophyta	T
<i>Pikea californica</i>	Captain Pike's weed	alga	Rhodophyta	M
<i>Pimepales promelas</i>	Fathead Minnow	bony fish (Actinopterygii)	Chordata	F
<i>Pinus contorta</i>	Lodgepole Pine	conifer	Pinophyta	T
<i>Pinus nigra</i>	Black Pine	conifer	Pinophyta	T
<i>Pinus pinaster</i>	Atlantic Maritime Pine	conifer	Pinophyta	T
<i>Pistia stratiotes</i>	Water Lettuce	flowering plant	Anthophyta	F
<i>Planaria torva</i>		flatworm (Turbellaria)	Platyhelminthes	F
<i>Podarcis muralis</i>	Wall Lizard	reptile	Chordata	T
<i>Pontederia cordata</i>	Pickerelweed	flowering plant	Anthophyta	F
<i>Potamopyrgus antipodarum</i>	Jenkins' Spire Snail	mollusc	Mollusca	F
<i>Procambarus clarkii</i>	Red Swamp Crayfish	crustacean	Arthropoda	F
<i>Procambarus marmoratus</i>	Marbled Crayfish	crustacean	Arthropoda	F
<i>Procyon lotor</i>	Raccoon	terrestrial mammal	Chordata	T
<i>Prokelisia marginata</i>	Spartina Planthopper	insect - true bug (Hemiptera)	Arthropoda	TM
<i>Prunus laurocerasus</i>	Cherry Laurel	flowering plant	Anthophyta	T
<i>Prunus serotina</i>	Rum Cherry	flowering plant	Anthophyta	T
<i>Pseudorasbora parva</i>	Topmouth Gudgeon	bony fish (Actinopterygii)	Chordata	F
<i>Pseudotsuga menziesii</i>	Douglas Fir	conifer	Pinophyta	T
<i>Psittacula eupatria</i>	Alexandrine Parakeet	bird	Chordata	T
<i>Psittacula krameri</i>	Ring-Necked Parakeet	bird	Chordata	T
<i>Pudu puda</i>	Pudu Deer	terrestrial mammal	Chordata	T
<i>Pulvinaria regalis</i>	Horse Chestnut Scale	insect - true bug (Hemiptera)	Arthropoda	T
<i>Quercus cerris</i>	Turkey Oak	flowering plant	Anthophyta	T
<i>Quercus ilex</i>	Evergreen Oak	flowering plant	Anthophyta	T
<i>Rapana venosa</i>	Mangrove oyster	mollusc	Mollusca	M
<i>Rattus norvegicus</i>	Brown Rat	terrestrial mammal	Chordata	T
<i>Rattus rattus</i>	Ship Rat	terrestrial mammal	Chordata	T
<i>Reticulitermes lucifugus</i>		insect - termite	Arthropoda	T
<i>Rhithropanopeus harrisi</i>	Dwarf crab	crustacean	Arthropoda	M
<i>Rhodeus amarus</i>	Bitterling	bony fish (Actinopterygii)	Chordata	F
<i>Rhododendron luteum</i>	Yellow Azalea	flowering plant	Anthophyta	T
<i>Rhododendron ponticum</i>	Rhododendron	flowering plant	Anthophyta	T
<i>Ribes sanguineum</i>	Flowering Currant	flowering plant	Anthophyta	T
<i>Robinia pseudoacacia</i>	False Acacia	flowering plant	Anthophyta	T
<i>Rosa rugosa</i>	Japanese Rose	flowering plant	Anthophyta	T
<i>Rubus spectabilis</i>	Salmonberry	flowering plant	Anthophyta	T
<i>Sagittaria latifolia</i>	Duck-potato	flowering plant	Anthophyta	F
<i>Salvelinus fontinalis</i>	American Brook Trout	bony fish (Actinopterygii)	Chordata	F
<i>Salvinia molesta</i>	Giant Salvinia	fern	Pteridophyta	F

<i>Sander lucioperca</i>	Pikeperch	bony fish (Actinopterygii)	Chordata	F
<i>Sargassum muticum</i>	Wireweed	alga	Chromista	M
<i>Sciurus carolinensis</i>	Grey Squirrel	terrestrial mammal	Chordata	T
<i>Sedum album</i>	White Stonecrop	flowering plant	Anthophyta	T
<i>Selenochlamys ysbryda</i>	Ghost Slug	mollusc	Mollusca	T
	Narrow-Leaved			
<i>Senecio inaequidens</i>	Ragwort	flowering plant	Anthophyta	T
<i>Senecio squalidus</i>	Oxford Ragwort	flowering plant	Anthophyta	T
<i>Silurus glanis</i>	Catfish	bony fish (Actinopterygii)	Chordata	F
<i>Smyrniium olusatrum</i>	Alexanders	flowering plant	Anthophyta	T
<i>Smyrniium perfoliatum</i>	Perfoliate Alexanders	flowering plant	Anthophyta	T
<i>Solidago canadensis</i>	Canadian Goldenrod	flowering plant	Anthophyta	T
<i>Solidobalanus fallax</i>		crustacean	Arthropoda	M
<i>Spartina anglica</i>	Common Cord-grass	flowering plant	Anthophyta	TM
	Spiraea Bridewort			
<i>Spiraea salicifolia agg.</i>	group	flowering plant	Anthophyta	T
	African Cotton			
<i>Spodoptera littoralis</i>	Leafworm	insect - moth	Arthropoda	T
<i>Styela clava</i>	Leathery sea squirt	tunicate (Urochordata)	Chordata	M
<i>Symphoricarpos albus</i>	Snowberry	flowering plant	Anthophyta	T
<i>Symphytum x uplandicum</i>	Russian Comfrey	flowering plant	Anthophyta	T
<i>Syringa vulgaris</i>	Lilac	flowering plant	Anthophyta	T
<i>Syrmaticus reevesii</i>	Reeves's Pheasant	bird	Chordata	T
<i>Tadorna ferruginea</i>	Ruddy Shelduck	bird	Chordata	T
<i>Tamarix gallica</i>	Tamarisk	flowering plant	Anthophyta	T
<i>Tamias sibiricus</i>	Siberian Chipmunk	terrestrial mammal	Chordata	T
<i>Tapes philippinarum</i>	Manila clam	mollusc	Mollusca	M
<i>Teredo navalis</i>		mollusc	Mollusca	M
<i>Thaumetopoea processionea</i>	Oak Processionary			
	Moth	insect - moth	Arthropoda	T
<i>Threskiornis aethiopicus</i>	Sacred Ibis	bird	Chordata	T
<i>Trachemys scripta</i>	Red-eared Terrapin	reptile	Chordata	T
<i>Tricellaria inopinata</i>		bryozoan	Bryozoa	M
<i>Triturus carnifex</i>	Italian Crested Newt	amphibian	Chordata	TF
<i>Undaria pinnatifida</i>	Japanese kelp	alga	Chromista	M
<i>Urosalpinx cinerea</i>	American tingle	mollusc	Mollusca	M
<i>Varroa destructor</i>	Varroa Mite	acarine (Acari)	Arthropoda	T
<i>Vespa velutina</i>	Asian Hornet	insect - hymenopteran	Arthropoda	T
<i>Watersipora subtorquata</i>		bryozoan	Bryozoa	M
<i>Xenopus laevis</i>	African Clawed Toad	amphibian	Chordata	TF
<i>Zamenis longissimus</i>	Aesculapian Snake	reptile	Chordata	T

Appendix 4. Micro-organisms excluded from the project (250 species)

Species	English name	Phylum	Establishment status	Status
<i>Aglaospora profusa</i>		Ascomycota	E	U
<i>Anthostomella trachycarpi</i>		Ascomycota	E	U
<i>Apiognomonina errabunda</i>		Ascomycota		U
<i>Arachnopeziza aranea</i>		Ascomycota	E	U
<i>Arthrocladiella mougeotii</i>		Ascomycota	E	U
<i>Astrosphaeriella trochus</i>		Ascomycota	E	U
<i>Aulographina eucalypti</i>		Ascomycota	E	U
<i>Blumeriella jaapii</i>		Ascomycota		U
<i>Botryosphaeria rhodora</i>		Ascomycota	E	U
<i>Botryotinia draytonii</i>		Ascomycota	N	U
<i>Botryotinia sphaerosperma</i>		Ascomycota	E	U
<i>Botryotinia squamosa</i>		Ascomycota	N	U
<i>Calonectria kyotensis</i>		Ascomycota		U
<i>Ciboria americana</i>		Ascomycota	E	U
<i>Cryptodiaporthe aesculi</i>		Ascomycota	E	U
<i>Cryptodiaporthe castanea</i>		Ascomycota	E	U
<i>Cryptodiaporthe robergeana</i>		Ascomycota	N	U
<i>Cryptosporella platanigera</i>		Ascomycota	E	U
<i>Cryptostroma corticale</i>		Ascomycota		U
<i>Cucurbitaria laburni</i>		Ascomycota	E	U
<i>Cucurbitaria piceae</i>		Ascomycota	E	U
<i>Cylindrocladium buxicola</i>		Ascomycota		U
<i>Diaporthe aucubae</i>		Ascomycota	E	U
<i>Diaporthe nobilis</i>		Ascomycota	E	U
<i>Diaporthe oncostoma</i>		Ascomycota	E	U
<i>Diaporthe skimmiae</i>		Ascomycota	E	U
<i>Didymascella thujina</i>	Thuja Needle Scorch	Ascomycota	E	U
<i>Didymella exitialis</i>		Ascomycota	E	U
	Tomato Stem and Fruit			
<i>Didymella lycopersici</i>	Rot	Ascomycota	E	U
<i>Discohainesia oenotherae</i>		Ascomycota	E	U
<i>Dothidotthia celtidis</i>		Ascomycota	E	U
<i>Drepanopeziza populi-albae</i>		Ascomycota	E	U
<i>Drepanopeziza punctiformis</i>		Ascomycota		U
<i>Elsinoe ampelina</i>	Grape Anthracnose	Ascomycota	E	U
<i>Entoleuca mammata</i>		Ascomycota		U
<i>Epibelonium gaeumannii</i>		Ascomycota	E	U
<i>Erysiphe alphitoides</i>		Ascomycota		U
<i>Erysiphe arcuata</i>		Ascomycota		U
<i>Erysiphe euonymi-japonici</i>		Ascomycota	E	U
<i>Erysiphe flexuosa</i>		Ascomycota		U
<i>Erysiphe necator</i>	Vine Powdery Mildew	Ascomycota	E	U
<i>Erysiphe palczewskii</i>		Ascomycota		U
<i>Erysiphe platani</i>		Ascomycota	E	U
<i>Erysiphe rayssiae</i>		Ascomycota	E	U
<i>Erysiphe russellii</i>		Ascomycota		U
<i>Erysiphe syringae</i>		Ascomycota	E	U
<i>Erysiphe vanbruntiana</i> var.		Ascomycota		U

<i>sambuci-racemosae</i>				
<i>Eupropoella arundinariae</i>		Ascomycota	E	NN
<i>Eupropoella britannica</i>		Ascomycota	E	U
<i>Glomerella cingulata</i>		Ascomycota		U
<i>Gnomonia leptostyla</i>		Ascomycota	E	U
<i>Gnomonia tetraspora</i>		Ascomycota		U
<i>Golovinomyces orontii</i>		Ascomycota	E	U
<i>Guignardia aesculi</i>		Ascomycota	E	U
<i>Hapalocystis berkeleyi</i>		Ascomycota	E	U
<i>Hyalopeziza spinicola</i>		Ascomycota	E	U
<i>Hyaloscypha mirabilis</i>		Ascomycota	E	U
<i>Hysterostegiella lauri</i>		Ascomycota	E	U
<i>Kabatiella caulivora</i>		Ascomycota		U
<i>Kabatina thujae</i>		Ascomycota		U
<i>Khuskia oryzae</i>		Ascomycota		U
<i>Lachnellula resinaria</i>		Ascomycota	E	U
<i>Lachnellula willkommii</i>		Ascomycota	E	U
<i>Lachnum castaneicola</i>		Ascomycota	E	U
<i>Lanzia coracina</i>		Ascomycota	E	U
<i>Lanzia echinophila</i>		Ascomycota	E	U
<i>Lembosina aulographoides</i>		Ascomycota	E	U
<i>Leptosphaeria lunariae</i>		Ascomycota		U
<i>Leucostoma curreyi</i>		Ascomycota	E	U
<i>Leucostoma kunzei</i>		Ascomycota	E	U
<i>Lichenopeltella fimbriata</i>		Ascomycota	E	U
<i>Lophodermium piceae</i>		Ascomycota	E	U
<i>Lophodermium pini-excelsae</i>		Ascomycota	E	U
<i>Lophodermium vagulum</i>		Ascomycota	E	U
<i>Lophomerum ponticum</i>		Ascomycota	E	U
<i>Macrophomina phaseolina</i>		Ascomycota		U
<i>Melanomma rhododendri</i>		Ascomycota	E	U
<i>Microthyrium lauri</i>		Ascomycota	E	U
<i>Moellerodiscus advenulus</i>		Ascomycota	E	U
<i>Monilinia mespili</i>		Ascomycota	E	U
<i>Morenoina chamaecyparidis</i>		Ascomycota	E	U
<i>Mycosphaerella chrysanthemi</i>	Chrysanthemum Ray Blight	Ascomycota	E	NN
<i>Mycosphaerella cydoniae</i>		Ascomycota		U
<i>Mycosphaerella pini</i>		Ascomycota		U
<i>Mycosphaerella pinodes</i>		Ascomycota	E	U
<i>Mycosphaerella podagrariae</i>		Ascomycota	E	U
<i>Mycosphaerella rhododendri</i>		Ascomycota	E	U
<i>Natrassia mangiferae</i>		Ascomycota		U
<i>Nectriella consolationis</i>		Ascomycota	E	U
<i>Neobulgaria undata</i>		Ascomycota		U
<i>Niptera subbiatorina</i>		Ascomycota	E	U
<i>Oidium hortensiae</i>		Ascomycota		U
<i>Oidium (Pseudoideum) sp.</i>	Powdery mildew of rhododendron	Ascomycota	E	U
<i>Ophiostoma novo-ulmi</i>	Dutch elm disease	Ascomycota	E	U

	Dutch elm disease (older less virulent strain)			
<i>Ophiostoma ulmi</i>		Ascomycota	E	U
<i>Orbilia retrusa</i>		Ascomycota	E	U
<i>Paurocotylis pila</i>		Ascomycota	E	U
<i>Pestalotiopsis guepinii</i>		Ascomycota		U
<i>Pezicula houghtonii</i>		Ascomycota	E	U
<i>Phaeocryptopus gaeumannii</i>		Ascomycota	E	U
<i>Phloeospora robiniae</i>		Ascomycota		U
<i>Phomopsis juniperivora</i>		Ascomycota		U
<i>Podosphaera mors-uvae</i>	American Mildew	Ascomycota	E	U
<i>Podosphaera xanthii</i>		Ascomycota		U
<i>Protoventuria arxii</i>		Ascomycota	E	U
<i>Pseudomassaria thistletonia</i>		Ascomycota	E	U
<i>Pseudonectria pachysandricola</i>		Ascomycota		U
<i>Pseudophacidium piceae</i>		Ascomycota	E	U
<i>Pseudovalsa modonia</i>		Ascomycota	E	U
<i>Psilachnum auranticolor</i>		Ascomycota	E	U
<i>Pyrenophora chaetomioides</i>		Ascomycota		U
<i>Ramularia collo-cygni</i>		Ascomycota		U
<i>Rhabdocline pseudotsugae</i>		Ascomycota	E	U
<i>Sarcotrochila alpina</i>		Ascomycota	E	U
<i>Seiridium cardinale</i>		Ascomycota		U
<i>Septoria betulae</i>		Ascomycota		U
<i>Stigmatea aegopodii</i>		Ascomycota	E	D
<i>Stomiopeltis cupressicola</i>		Ascomycota	E	U
<i>Stromatinia gladioli</i>		Ascomycota		U
<i>Taphrina deformans</i>	Peach Leaf Curl	Ascomycota	E	U
<i>Trochila laurocerasi</i>		Ascomycota	E	U
<i>Tympanis laricina</i>		Ascomycota	E	U
<i>Valsa laurocerasi</i>		Ascomycota	E	U
<i>Venturia saliciperda</i>		Ascomycota		U
<i>Xenomeris nicholsonii</i>		Ascomycota	E	U
<i>Agrocybe putaminum</i>		Basidiomycota	E	NN
<i>Amanita inopinata</i>		Basidiomycota	E	U
<i>Aseroe rubra</i>	Starfish Fungus	Basidiomycota	E	U
<i>Calocera pallidospathulata</i>	Pale Stagshorn	Basidiomycota		NN
<i>Chroogomphus helveticus</i>		Basidiomycota		U
<i>Chrysomyxa abietis</i>	Spruce Needle Rust	Basidiomycota	E	U
<i>Chrysomyxa ledi</i> var. <i>rhododendri</i>	Rhododendron Rust	Basidiomycota	E	U
<i>Clathrus archeri</i>	Devil's Fingers	Basidiomycota	E	U
<i>Clathrus ruber</i>	Red Cage	Basidiomycota	E	U
<i>Collybia biformis</i>		Basidiomycota		NN
<i>Cronartium ribicola</i>	White Pine Blister Rust	Basidiomycota	E	U
<i>Cumminsella mirabilissima</i>		Basidiomycota	E	NN
<i>Endophyllum sempervivi</i>	Houseleek Rust	Basidiomycota	E	U
<i>Entyloma calendulae</i>	Calendula Smut	Basidiomycota	E	U
<i>Entyloma compositarum</i>	Gaillardia Smut	Basidiomycota	N	U
<i>Entyloma dahliae</i>	Dahlia Smut	Basidiomycota	E	U

<i>Entyloma fuscum</i>		Basidiomycota		U
<i>Flaviporus brownei</i>		Basidiomycota	E	U
<i>Gastrosporium simplex</i>	Steppe Truffle	Basidiomycota	N	U
<i>Geastrum campestre</i>	Field Earthstar	Basidiomycota	N	U
<i>Geastrum floriforme</i>	Daisy Earthstar	Basidiomycota	E	U
<i>Guepinia helvelloides</i>	Salmon Salad	Basidiomycota	E	U
<i>Gymnosporangium confusum</i>		Basidiomycota	E	U
<i>Gymnosporangium juniperi-virginianae</i>		Basidiomycota	E	NN
<i>Gymnosporangium sabinae</i>		Basidiomycota	E	U
<i>Hydnangium carneum</i>		Basidiomycota	E	U
<i>Ileodictyon cibarium</i>	Basket Fungus	Basidiomycota	E	U
<i>Laccaria fraterna</i>		Basidiomycota	E	U
<i>Leucocoprinus birnbaumii</i>	Plantpot Dapperling	Basidiomycota	E	U
<i>Leucocoprinus tenellus</i>		Basidiomycota		U
<i>Leucocoprinus zeylanicus</i>		Basidiomycota		U
<i>Lysurus cruciatus</i>	Lizard's Claw	Basidiomycota	N	U
<i>Melampsora amygdalinae</i>		Basidiomycota	E	U
<i>Melampsora lini</i> var. <i>liniperda</i>		Basidiomycota	N	U
<i>Melampsora ribesii-viminalis</i>		Basidiomycota	E	U
<i>Melampsora salicis-albae</i>		Basidiomycota	E	U
<i>Melampsidium hiratsukanum</i>		Basidiomycota		U
<i>Melanotaenium hypogaeum</i>		Basidiomycota		U
<i>Melanotaenium jaapii</i>		Basidiomycota	E	U
<i>Microstroma juglandis</i>		Basidiomycota	E	U
<i>Mikronegeria fagi</i>		Basidiomycota		U
<i>Mutinus ravenelii</i>	Red Stinkhorn	Basidiomycota	E	NN
<i>Omphalotus olearius</i>		Basidiomycota	E	U
<i>Phragmidium fusiforme</i>		Basidiomycota		U
<i>Pisolithus arrhizus</i>	Dyeball	Basidiomycota	E	U
<i>Psilocybe cyanescens</i>	Blueleg Brownie	Basidiomycota	E	NN
<i>Puccinia aegopodii</i>		Basidiomycota	E	U
<i>Puccinia antirrhini</i>	Antirrhinum Rust	Basidiomycota	E	U
<i>Puccinia cyani</i>	Centaurea Rust	Basidiomycota		U
<i>Puccinia distincta</i>	Rust of daisy	Basidiomycota	E	U
<i>Puccinia gentianae</i>		Basidiomycota		U
<i>Puccinia gladioli</i>		Basidiomycota		U
<i>Puccinia horiana</i>	Chrysanthemum White Rust	Basidiomycota	N	U
<i>Puccinia kusanoi</i>		Basidiomycota	E	U
<i>Puccinia lagenophorae</i>	Rust of groundsel	Basidiomycota	E	U
<i>Puccinia ljulinica</i>		Basidiomycota	E	U
<i>Puccinia longicornis</i>		Basidiomycota	E	U
<i>Puccinia malvacearum</i>		Basidiomycota	E	NN
<i>Puccinia mariana</i>		Basidiomycota		U
<i>Puccinia oxalidis</i>		Basidiomycota	E	U
<i>Puccinia pazschkei</i> var. <i>pazschkei</i>	Saxifrage Rust	Basidiomycota		U

<i>Puccinia pelargonii-zonalis</i>	Pelargonium Rust	Basidiomycota	E	U
<i>Puccinia porri</i>		Basidiomycota	E	U
<i>Puccinia prostii</i>		Basidiomycota	N	U
<i>Puccinia smyrnii</i>		Basidiomycota	E	U
<i>Puccinia sorghi</i>	Maize Rust	Basidiomycota	E	NN
<i>Puccinia vincae</i>	Periwinkle Rust	Basidiomycota	E	U
<i>Queletia mirabilis</i>	Quelet's Stalk Puffball	Basidiomycota	N	U
<i>Sporisorium destruens</i>	Millet Smut	Basidiomycota		NN
<i>Sporisorium sorghi</i>	Sorghum Smut	Basidiomycota		NN
<i>Strobilurus esculentus</i>	Sprucecone Cap	Basidiomycota	E	U
<i>Stropharia aurantiaca</i>	Redlead Roundhead	Basidiomycota	E	U
<i>Stropharia percevalii</i>		Basidiomycota	E	U
<i>Suillus amabilis</i>		Basidiomycota		D
<i>Suillus grevillei</i>	Larch Bolete	Basidiomycota	E	D
<i>Suillus placidus</i>		Basidiomycota		U
<i>Suillus viscidus</i>	Sticky Bolete	Basidiomycota	E	D
<i>Tilletia caries</i>	Bunt of wheat	Basidiomycota	N	U
<i>Urocystis eranthidis</i>		Basidiomycota	E	U
<i>Urocystis gladiolicola</i>	Gladiolus Smut	Basidiomycota	E	U
<i>Urocystis occulta</i>	Stripe Smut of Rye	Basidiomycota	E	U
<i>Urocystis syncocca</i>		Basidiomycota		U
<i>Uromyces aecidiiformis</i>		Basidiomycota	N	U
<i>Uromyces appendiculatus</i>		Basidiomycota	E	U
<i>Uromyces colchici</i>		Basidiomycota	N	U
<i>Uromyces dianthi</i>	Carnation Rust	Basidiomycota	E	D
<i>Uromyces erythronii</i>	Erythronium Rust	Basidiomycota	N	U
<i>Ustilago cynodontis</i>		Basidiomycota	E	U
<i>Ustilago maydis</i>	Maize Smut	Basidiomycota	E	U
	Loose Smut of Wheat and Barley	Basidiomycota	E	N-NN
<i>Zaghouania phillyreae</i>		Basidiomycota	X	U
<i>Batrachochytrium dendrobatidis</i>		Fungi (other)		U
<i>Circinotrichum britannicum</i>		Fungi (other)	E	U
<i>Colletotrichum acutatum</i>		Fungi (other)	E	U
<i>Sphaeropsis sapinea</i>		Fungi (other)		U
<i>Synchytrium endobioticum</i>	Wart disease of potato	Fungi (other)	E	NN
<i>Bonamia ostreae</i>	None known	Haplosporea	E	NN
<i>Aerococcus viridans</i>		Microorganisms	E	U
<i>Barley mild mosaic virus</i>	Barley mild mosaic	Microorganisms	E	U
<i>Barley yellow mosaic virus</i>	Barley yellow mosaic	Microorganisms	E	U
<i>Beet necrotic yellow vein virus</i>		Microorganisms	E	U
<i>Dickeya dianthicola</i>	Rhizomania	Microorganisms	E	U
	Slow wilt of potato	Microorganisms	E	U
<i>Erwinia amylovora</i>	Fireblight of pome fruit	Microorganisms	E	NN
<i>Plum pox virus</i>	Plum pox	Microorganisms	E	U
<i>Pseudomonas syringae pv. pisi</i>	Pea blight	Microorganisms	E	U
<i>Ralstonia solanacearum race 3 biovar 2</i>	Brown rot of potato	Microorganisms	E	NN
<i>Rhabdovirus carpio</i>	Spring Viremia of Carp	Microorganisms		

<i>Soil-borne cereal mosaic virus</i>	Soil-borne cereal mosaic	Microorganisms	E	U
<i>Tomato spotted wilt virus</i>	Tomato spotted wilt	Microorganisms	E	U
<i>Cornuvia serpula</i>		Mycetozoa	N	U
<i>Aphanomyces astaci</i>	Crayfish plague	Oomycete	E	NN
<i>Phytophthora alni</i>	Dieback of alder	Oomycete	E	U
<i>Phytophthora cambivora</i>		Oomycete		U
<i>Phytophthora cinnamomi</i>		Oomycete		U
<i>Phytophthora citricola</i>		Oomycete		U
<i>Phytophthora citrophthora</i>		Oomycete		U
<i>Phytophthora drechsleri</i>		Oomycete		U
<i>Phytophthora ilicis</i>		Oomycete		U
<i>Phytophthora infestans</i>	Potato Blight	Oomycete	E	NN
<i>Phytophthora kernoviae</i>	Trunk canker of beech	Oomycete	E	U
<i>Phytophthora quercina</i>		Oomycete		U
<i>Phytophthora ramorum</i>	Sudden oak death	Oomycete	E	U
<i>Phytophthora syringae</i>		Oomycete		U
<i>Pseudoperonospora humuli</i>	Downy mildew of hop	Oomycete	E	U
<i>Ostracoblabe implexa</i>		Zygomycota	U	U

Appendix 5. Species lists, within broad groups, designated as having a negative ecological or human impact. Region of origin and introduction pathway are also provided.

	Species	Ecological	Human	Region	Pathway
Higher plants					
Pirri-Pirri Bur	<i>Acaena novae-zelandiae</i>	N		Australasia	Stowaway
Bear's-Breech	<i>Acanthus mollis</i>	N		Europe	Ornamental
Sycamore	<i>Acer pseudoplatanus</i>	N		Europe	Ornamental
Ground-elder	<i>Aegopodium podagraria</i>		N	Europe	Agriculture
Tree-Of-Heaven	<i>Ailanthus altissima</i>	N		Asia-Temperate	Ornamental
Garden Lady's-mantle	<i>Alchemilla mollis</i>	N		Europe	Ornamental
Few-flowered Garlic	<i>Allium paradoxum</i>	N		Asia-Temperate	Ornamental
Rosy Garlic	<i>Allium roseum</i>	N		Europe	Ornamental
Three-Cornered Garlic	<i>Allium triquetrum</i>	N		Europe	Ornamental
Black Twitch	<i>Alopecurus myosuroides</i>		N	Europe Northern	Contaminant
Juneberry	<i>Amelanchier lamarckii</i>	N		America	Ornamental
Great Brome	<i>Anisantha diandra</i>	N		Europe	Contaminant
Barren Brome	<i>Anisantha sterilis</i>		N	Europe	Stowaway
Stinking Chamomile	<i>Anthemis cotula</i>		N	Europe	Contaminant
Loose Silky-bent	<i>Apera spica-venti</i>		N	Europe Northern	Contaminant
Michaelmas-daisies	<i>Aster (alien N American taxa)</i>	N		America	Ornamental
Late Michaelmas-Daisy	<i>Aster laevis x novi-belgii (A. x versicolor)</i>	N			Ornamental
Narrow-Leaved				Northern	
Michaelmas-Daisy	<i>Aster lanceolatus</i>	N		America	Ornamental
Common Michaelmas-daisy	<i>Aster lanceolatus x novi-belgii (A. x salignus)</i>	N			Ornamental
Hairy Michaelmas-Daisy	<i>Aster novae-angliae</i>	N		Northern America	Ornamental
Confused Michaelmas-Daisy	<i>Aster novi-belgii</i>	N		Northern America	Ornamental
Wild Oat	<i>Avena fatua</i>		N	Europe	Contaminant
Animated Oat	<i>Avena sterilis</i>		N	Europe Northern	Contaminant
Water Fern	<i>Azolla filiculoides</i>	N		America	Ornamental
Buddleia	<i>Buddleja davidii</i>	N	N	Asia-Temperate	Ornamental
American Bellbine	<i>Calystegia silvatica</i>		N	Europe	Ornamental
Hottentot-Fig	<i>Carpobrotus edulis</i>	N	N	Africa	Ornamental
Corn Marigold	<i>Chrysanthemum segetum</i>		N	Europe	Contaminant
Bladder-senna	<i>Colutea arborescens</i>	N		Europe Northern	Ornamental
Red-Osier Dogwood	<i>Cornus sericea</i>	N		America	Ornamental
Wall Cotoneaster	<i>Cotoneaster horizontalis</i>	N		Asia-Temperate	Ornamental
Entire-leaved Cotoneaster	<i>Cotoneaster integrifolius</i>	N		Asia-Temperate	Ornamental
Himalayan Cotoneaster	<i>Cotoneaster simonsii</i>	N		Asia-Tropical	Ornamental
New Zealand Pigmyweed	<i>Crassula helmsii</i>	N	N	Australasia	Ornamental

Montbretia	<i>Crocsmia aurea x pottsii (C. x crocosmiiflora)</i>	N			Ornamental
Purple Dewplant	<i>Disphyma crassifolium</i>	N	N	Africa	Ornamental
Large-Flowered Waterweed	<i>Egeria densa</i>	N		Southern America	Aquaculture
Canadian Pondweed	<i>Elodea canadensis</i>	N		Northern America	Ornamental
Esthwaite Water-Weed	<i>Elodea nuttallii</i>	N		America	Ornamental
Russian Vine	<i>Fallopia baldschuanica</i>	N		Asia-Temperate	Ornamental
Japanese Knotweed	<i>Fallopia japonica</i>	N	N	Asia-Temperate	Ornamental
Conolly's Knotweed	<i>Fallopia japonica x sachalinensis (F. x bohémica)</i>	N	N		Ornamental
Giant Knotweed	<i>Fallopia sachalinensis</i>	N		Asia-Temperate Northern	Ornamental
Shallon	<i>Gaultheria shallon</i>	N		America Southern	Ornamental
Brazilian Giant Rhubarb	<i>Gunnera manicata</i>	N		America Southern	Ornamental
Giant Rhubarb	<i>Gunnera tinctoria</i>	N		America	Ornamental
Giant Hogweed	<i>Heracleum mantegazzianum</i>	N	N	Asia-Temperate	Ornamental
Garden Bluebell	<i>Hyacinthoides hispanica</i>	N		Europe	Ornamental
Bluebell	<i>Hyacinthoides hispanica x non-scripta</i>	N			Ornamental
Floating Pennywort	<i>Hydrocotyle ranunculoides</i>	N		Northern America	Ornamental
Himalayan Balsam	<i>Impatiens glandulifera</i>	N	N	Asia-Tropical	Ornamental
Small Balsam	<i>Impatiens parviflora</i>	N		Asia-Temperate	Stowaway
Curly Waterweed	<i>Lagarosiphon major</i>	N		Africa	Ornamental
	<i>Lamiastrum galeobdolon subsp. argentatum</i>	N	N		Ornamental
Least Duckweed	<i>Lemna minuta</i>	N		Northern America	Ornamental
Uruguay Water-primrose	<i>Ludwigia grandiflora</i>	N		Northern America	Aquaculture
Wireplant	<i>Muehlenbeckia complexa</i>	N		Australasia Southern	Ornamental
Parrot's Feather	<i>Myriophyllum aquaticum</i>	N		America	Aquaculture
Pheasant's-eye Daffodil	<i>Narcissus poeticus</i>	N		Europe	Ornamental
	<i>Narcissus poeticus x pseudonarcissus subsp. pseudonarcissus (N. x incomparabilis)</i>	N		Europe	Ornamental
Nonesuch Daffodil	<i>Narcissus poeticus x tazetta (N. x medioluteus)</i>	N		Europe	Ornamental
Primrose-Peerless	<i>Narcissus pseudonarcissus subsp. major</i>	N		Europe Southern	Ornamental
Spanish Daffodil		N		Europe Southern	Ornamental
Large-Flowered Pink-Sorrel	<i>Oxalis debilis</i>	N	N	America Southern	Ornamental
Garden Pink-sorrel	<i>Oxalis latifolia</i>	N	N	America	Ornamental
Bermuda-buttercup	<i>Oxalis pes-caprae</i>	N	N	Africa	Ornamental

Virginia-creeper	<i>Parthenocissus quinquefolia</i>	N		Northern America	Ornamental
Green Alkanet	<i>Pentaglottis sempervirens</i>	N		Europe	Ornamental
Lesser Knotweed	<i>Persicaria campanulata</i>	N		Asia-Tropical	Ornamental
Himalayan Knotweed	<i>Persicaria wallichii</i>	N		Asia-Tropical	Ornamental
White Butterbur	<i>Petasites albus</i>	N		Europe	Ornamental
Winter Heliotrope	<i>Petasites fragrans</i>	N		Europe	Ornamental
Black Pine	<i>Pinus nigra</i>	N		Europe	Forestry
Corsican Pine	<i>Pinus nigra subsp. laricio</i>	N		Europe	Forestry
Austrian Pine	<i>Pinus nigra subsp. nigra</i>	N		Europe	Forestry
Atlantic Maritime Pine	<i>Pinus pinaster</i>	N		Europe	Ornamental
Cherry Laurel	<i>Prunus laurocerasus</i>	N		Europe	Ornamental
Portugal Laurel	<i>Prunus lusitanica</i>	N		Europe	Ornamental
				Northern America	
Rum Cherry	<i>Prunus serotina</i>	N		America	Ornamental
Arrow Bamboo	<i>Pseudosasa japonica</i>	N		Asia-Temperate	Ornamental
Turkey Oak	<i>Quercus cerris</i>	N		Europe	Ornamental
Evergreen Oak	<i>Quercus ilex</i>	N		Europe	Ornamental
Rhododendron	<i>Rhododendron ponticum</i>	N	N		Ornamental
				Northern America	
False Acacia	<i>Robinia pseudoacacia</i>	N		America	Ornamental
Dutch Rose	<i>Rosa 'Hollandica'</i>	N			Ornamental
Japanese Rose	<i>Rosa rugosa</i>	N		Asia-Temperate	Ornamental
				Northern America	
Duck-potato	<i>Sagittaria latifolia</i>	N		America	Ornamental
Broad-Leaved Bamboo	<i>Sasa palmata</i>	N		Asia-Temperate	Ornamental
Hairy Bamboo	<i>Sasaella ramosa</i>	N		Asia-Temperate	Ornamental
White Mustard	<i>Sinapis alba</i>		N	Europe	Agriculture
	<i>Sinapis alba subsp. alba</i>		N	Europe	Agriculture
Charlock	<i>Sinapis arvensis</i>		N	Europe	Contaminant
Alexanders	<i>Smyrniolum olusatrum</i>	N		Europe	Agriculture
	<i>Spartina maritima x alterniflora = S. x townsendii</i>	N			Hybridisation
Corn Spurrey	<i>Spergula arvensis</i>		N	Europe	Contaminant
Spiraea	<i>Spiraea</i>	N		Europe	Ornamental
	<i>Spiraea alba x douglasii (S. x billardii)</i>	N			Ornamental
				Northern America	
Steeplebush	<i>Spiraea douglasii</i>	N		America	Ornamental
	<i>Spiraea douglasii x salicifolia (S. x pseudosalicifolia)</i>	N			Ornamental
Confused Bridewort	<i>Spiraea salicifolia</i>	N		Europe	Ornamental
				Northern America	
Snowberry	<i>Symphoricarpos albus</i>	N		America	Ornamental
Scentsless Mayweed	<i>Tripleurospermum inodorum</i>		N	Europe	Contaminant
Slender Speedwell	<i>Veronica filiformis</i>	N	N	Asia-Temperate	Ornamental
Lesser Periwinkle	<i>Vinca minor</i>	N		Europe	Ornamental

Lower plants

Green sea-fingers (tomentosoides)	<i>Codium fragile fragile</i>	N	N	Asia-Temperate Pacific	Aquaculture
	<i>Polysiphonia harveyi</i>	N			Aquaculture

Wire weed	<i>Sargassum muticum</i>	N	N	Asia-Temperate	Aquaculture
Japanese kelp	<i>Undaria pinnatifida</i>	N		Asia-Temperate	Stowaway
Insects					
American Seed Beetle	<i>Acanthoscelides obtectus</i>		N	Southern America	Contaminant
Douglas Fir Woolly Aphid	<i>Adelges cooleyi</i>		N	Northern America	
Lesser Mealworm Beetle	<i>Alphitobius diaperinus</i>		N		Unknown
Black Fungus Beetle	<i>Alphitobius laevigatus</i>		N		Unknown
Australian Carpet Beetle	<i>Anthrenocerus australis</i>		N	Australasia	Contaminant
	<i>Anthrenus (Anthrenus)</i>				
Furniture Carpet Beetle	<i>flavipes</i>		N	Asia-Temperate	Contaminant
	<i>Anthrenus (Anthrenus)</i>				
Common Carpet Beetle	<i>scrophulariae</i>		N	Europe	Contaminant
	<i>Anthrenus (Florilinus) olgae</i>		N	Europe	Unknown
	<i>Anthrenus oceanicus</i>		N	Asia-Tropical	Unknown
Black legume aphid	<i>Aphis craccivora</i>		N	Asia-Temperate	
Cotton aphid / Melon aphid	<i>Aphis gossypii</i>		N	Africa	Ornamental
Locust Bean Moth	<i>Apomyelois ceratoniae</i>		N		Contaminant
Berberis Sawfly	<i>Arge berberidis</i>	N	N	Europe	Ornamental
	<i>Attagenus brunneus</i>		N	Africa	Unknown
	<i>Attagenus cyphonoides</i>		N		Contaminant
Wardrobe Beetle	<i>Attagenus fasciatus</i>		N		Contaminant
Two-spotted Carpet Beetle	<i>Attagenus pellio</i>		N		Unknown
Brown Carpet Beetle	<i>Attagenus smirnovi</i>		N	Africa	Unknown
	<i>Attagenus trifasciatus</i>		N	Europe	Unknown
Black Carpet Beetle	<i>Attagenus unicolor</i>		N	Africa	Contaminant
	<i>Aulacorthum(Neomyzus)</i>				
Mottled Arum aphid	<i>circumflexus</i>		N	Asia-Temperate	
Common Cockroach	<i>Blatta orientalis</i>		N	Africa	Contaminant
Lentil Seed Beetle	<i>Bruchus ervi</i>		N	Asia-Temperate	Contaminant
Pea Beetle	<i>Bruchus pisorum</i>		N	Europe	Contaminant
Bean Seed Beetle	<i>Bruchus rufimanus</i>		N	Europe	Unknown
Dried fruit moth	<i>Cadra calidella</i>		N		Contaminant
Dried currant moth	<i>Cadra cautella</i>		N		Contaminant
Raisin moth	<i>Cadra figulilella</i>		N		Contaminant
Adzuki Beanseed Beetle	<i>Callosobruchus chinensis</i>		N	Asia-Temperate	Contaminant
	<i>Cerataphis orchidearum</i>		N	Asia-Tropical	
	<i>Chaetosiphon fragaefolii</i>		N	Northern America	
Soft scale	<i>Coccus hesperidum</i>	N	N		Ornamental
an orchid scale insect	<i>Coccus pseudoheperidum</i>		N	Southern America	Ornamental
				Northern	
Spruce Groundling	<i>Coleotechnites piceaella</i>		N	America	Unknown
a palm scale insect	<i>Colobopyga kewensis</i>		N	Australasia	Ornamental
Rice Moth	<i>Corcyra cephalonica</i>		N		Contaminant
Asparagus Beetle	<i>Crioceris asparagi</i>		N	Europe	Unknown
Honeydew moth	<i>Cryptoblabes gnidiella</i>		N	Europe	Contaminant
Great Spruce Bark Beetle	<i>Dendroctonus micans</i>		N	Europe	Stowaway

Black Larder Beetle	<i>Dermestes (Dermestes) ater</i>			N	Europe	Contaminant
	<i>Dermestes (Dermestes)</i>				Southern	
Peruvian Larder Beetle	<i>peruvianus</i>			N	America	Contaminant
	<i>Dermestes (Dermestinus)</i>				Northern	
	<i>carnivorus</i>			N	America	Contaminant
	<i>Dermestes (Dermestinus)</i>					
	<i>frischii</i>			N		Contaminant
	<i>Dermestes (Dermestinus)</i>					
Hide Beetle	<i>maculatus</i>			N	Africa	Contaminant
rhododendron whitefly	<i>Dialeurodes chittendeni</i>			N		Ornamental
Dark Marbled Tabby	<i>Duponchelia fovealis</i>			N	Africa	Contaminant
	<i>Dysaphis tulipae</i>			N	Europe	
Green Spruce Aphid	<i>Elatobium abietinum</i>			N	Europe	
Cacao Moth	<i>Ephestia elutella</i>			N		Contaminant
Mediterranean Flour Moth	<i>Ephestia kuehniella</i>			N		Contaminant
Light Brown Apple Moth	<i>Epiphyas postvittana</i>			N	Australasia	Contaminant
					Northern	
Woolly Aphid	<i>Eriosoma lanigerum</i>			N	America	
New Zealand weevil	<i>Euophryum confine</i>			N	Australasia	Stowaway
	<i>Euophryum rufum</i>			N	Australasia	Stowaway
Peach Knot-horn	<i>Euzophera bigella</i>			N		Contaminant
a fern whitefly	<i>Filicaleyrodes williamsi</i>			N		Ornamental
Laburnum sucker	<i>Floria variegata</i>			N	Europe	Ornamental
					Northern	
	<i>Frankliniella occidentalis</i>			N	America	Ornamental
					Southern	
Broad-horned Flour Beetle	<i>Gnatocerus cornutus</i>			N	America	Unknown
					Northern	
Rhododendron leafhopper	<i>Graphocephala fennahi</i>	N			America	Unknown
Harlequin Ladybird	<i>Harmonia axyridis</i>	N	N		Asia-Temperate	Stowaway
	<i>Hemiberlesia palmae</i>			N		Ornamental
	<i>Homotoma ficus</i>			N	Europe	Agriculture
					Northern	
The Dump Fly	<i>Hydrotaea aenescens</i>	N			America	
Large Larch Bark Beetle	<i>Ips cembrae</i>			N	Europe	Stowaway
Invasive garden ant	<i>Lasius neglectus</i>	N	N		Asia-Temperate	Ornamental
Long-Headed Flour Beetle	<i>Latheticus oryzae</i>			N	Asia-Tropical	Contaminant
					Northern	
Western Conifer Seed Bug	<i>Leptoglossus occidentalis</i>			N	America	Stowaway
	<i>Macrosiphoniella sanborni</i>			N	Asia-Temperate	
					Northern	
Lupin Aphid	<i>Macrosiphum albifrons</i>			N	America	
					Northern	
	<i>Macrosiphum euphorbiae</i>	N	N		America	
Pharo's ant	<i>Monomorium pharaonis</i>			N	Africa	Unknown
	<i>Monomorium salomonis</i>			N	Africa	Unknown
Large peach aphid	<i>Myzus varians</i>			N	Asia-Temperate	
Plain shortwing beetle	<i>Nathrius brevipennis</i>	N			Europe	Stowaway
Southern Green Shield Bug	<i>Nezara viridula</i>			N	Africa	Contaminant
	<i>Oceanaspidiotus spinosus</i>			N		
Small-Eyed Flour Beetle	<i>Palorus ratzeburgii</i>			N	Africa	Contaminant

Stored Nut Moth	<i>Paralipsa gularis</i>		N	Asia-Tropical	Contaminant
Azalea whitefly	<i>Pealius azaleae</i>		N	Asia-Temperate	Ornamental
				Northern	
	<i>Pentarthrum huttoni</i>		N	America	Stowaway
Indian Meal Moth	<i>Plodia interpunctella</i>		N		Contaminant
				Northern	
Spartina Planthopper	<i>Prokelisia marginata</i>	N		America	Stowaway
Citrus Mealybug	<i>Pseudococcus calceolariae</i>		N	Australasia	Ornamental
Australian Spider Beetle	<i>Ptinus tectus</i>		N	Australasia	Stowaway
Hydrangea Scale	<i>Pulvinaria hydrangeae</i>		N	Asia-Temperate	Ornamental
Green shield scale	<i>Pulvinaria psidii</i>		N		Ornamental
Horse Chestnut Scale	<i>Pulvinaria regalis</i>	N	N	Asia-Temperate	Ornamental
	<i>Pulvinariella</i>				
Hottentot fig scale	<i>mesembryanthemi</i>		N	Africa	Ornamental
				Northern	
Museum Nuisance	<i>Reesa vespulae</i>		N	America	Unknown
	<i>Rhopalosiphoninus latysiphon</i>		N	Europe	
				Northern	
	<i>Rhopalosiphum insertum</i>		N	America	
	<i>Rhopalosiphum maidis</i>		N	Asia-Tropical	
Olive hemispherical scale	<i>Saissetia oleae</i>		N		Ornamental
Greater Rice Weevil	<i>Sitophilus zeamais</i>		N		Stowaway
Biscuit Beetle	<i>Stegobium paniceum</i>		N		Stowaway
				Northern	
Rhododendron Lacebug	<i>Stephanitis rhododendri</i>		N	America	Ornamental
Andromeda Lacebug	<i>Stephanitis takeyai</i>		N	Asia-Temperate	Ornamental
	<i>Takecallis arundicolens</i>		N	Asia-Temperate	
	<i>Takecallis taiwanus</i>		N	Asia-Temperate	
Dark Mealworm Beetle	<i>Tenebrio obscurus</i>		N		Contaminant
				Southern	
Large Pale Clothes Moth	<i>Tinea pallescentella</i>		N	America	Unknown
	<i>Tinocallis ulmiparvifoliae</i>		N	Asia-Temperate	
	<i>Tinocallis zelkowae</i>		N	Asia-Temperate	
Rust-red Flour Beetle	<i>Tribolium castaneum</i>		N	Asia-Tropical	Contaminant
Confused Flour Beetle	<i>Tribolium confusum</i>		N	Africa	Contaminant
				Southern	
Dark Flour Beetle	<i>Tribolium destructor</i>		N	America	Contaminant
Bay Sucker	<i>Trioza alacris</i>		N	Europe	Ornamental
Pittosporum Psyllid	<i>Trioza vitreoradiata</i>		N	Australasia	Ornamental
Coloured Cabinet Beetle	<i>Trogoderma glabrum</i>		N	Europe	Contaminant
Large Cabinet Beetle	<i>Trogoderma inclusum</i>		N		Unknown
Warehouse beetle	<i>Trogoderma variabile</i>		N		Contaminant

Invertebrates

Eel swim-bladder nematode	<i>Anguillicola crassus</i>	N	N	Asia-Temperate	Aquaculture
Turkish Crayfish	<i>Astacus leptodactylus</i>		N	Asia-Temperate	Aquaculture
Asiatic Clam	<i>Corbicula fluminea</i>	N		Asia-Temperate	Unknown
Freshwater hydroid	<i>Cordylophora caspia</i>	N	N	Europe	Stowaway
Northern River				Northern	
Crangonyctid	<i>Crangonyx pseudogracilis</i>	N		America	Stowaway
Brazilian oyster	<i>Crassostrea brasiliana</i>		N	Pacific	Unknown

Slipper limpet	<i>Crepidula fornicata</i>	N	N	Northern America	Aquaculture
Chestnut Slug	<i>Deroceras (Deroceras) panormitanum</i>		N	Europe	Unknown
Carpet Sea-squirt	<i>Didemnum vexillum</i>	N	N		Stowaway
Killer Shrimp	<i>Dikerogammarus villosus</i>	N		Europe	Unknown
Zebra Mussel	<i>Dreissena polymorpha</i>	N	N	Asia-Temperate	Stowaway
Say mud crab	<i>Dyspanopeus sayi</i>	N		Northern America	Stowaway
	<i>Ergasilus briani</i>	N	N	Europe	Aquaculture
	<i>Ergasilus gibbus</i>	N	N	Europe	Aquaculture
	<i>Ergasilus sieboldi</i>	N	N	Europe	Aquaculture
Chinese Mitten Crab	<i>Eriocheir sinensis</i>	N	N	Asia-Temperate	Stowaway
Australian tube worm	<i>Ficopomatus enigmaticus</i>		N		Stowaway
Sideswimmer	<i>Gammarus tigrinus</i>	N		Northern America	Stowaway
Orange-striped anemone	<i>Haliplanella lineata</i>		N	Pacific	Stowaway
Common Garden Snail	<i>Helix aspersa</i>		N	Europe	Unknown
A tube worm	<i>Hydroides dianthus</i>		N	Northern America	Stowaway
A polychaete worm	<i>Hydroides elegans</i>	N	N	Australasia	Stowaway
A tube worm	<i>Hydroides ezoensis</i>		N	Asia-Temperate	Stowaway
A parasitic copepod	<i>Mytilicola intestinalis</i>	N	N	Europe	Aquaculture
False dark mussel	<i>Mytilopsis leucophaeta</i>	N	N	Northern America	Stowaway
	<i>Neoergasilus japonicus</i>	N	N	Europe	Aquaculture
Signal Crayfish	<i>Pacifastacus leniusculus</i>	N	N	Northern America	Aquaculture
	<i>Paraergasilus longidigitus</i>	N		Asia-Temperate	Aquaculture
Jenkins' Spire Snail, New Zealand Mudsnail	<i>Potamopyrgus antipodarum</i>	N		Australasia	Stowaway
Root Lesion Nematode	<i>Pratylenchus bolivianus</i>		N	Europe	Ornamental
Dwarf crab	<i>Rhithropanopeus harrisi</i>	N	N	Northern America	Unknown
A barnacle	<i>Solidobalanus fallax</i>	N	N	Africa	Unknown
Leathery sea squirt	<i>Styela clava</i>		N	Asia-Temperate	Stowaway
Budapest Slug	<i>Tandonia budapestensis</i>		N	Europe	Unknown
Keeled Slug	<i>Tandonia sowerbyi</i>		N	Europe	Unknown
American Oyster Drill	<i>Urosalpinx cinerea</i>	N	N	Northern America	Aquaculture
	<i>Watersipora subtorquata</i>		N		Stowaway
Vertebrates					
French Partridge	<i>Alectoris rufa</i>	N		Europe	Animal husbandry
Egyptian Goose	<i>Alopochen aegyptiacus</i>	N	N	Africa	Ornamental
Canada Goose	<i>Branta canadensis</i>	N	N	Northern America	Ornamental
Barnacle Goose	<i>Branta leucopsis</i>	N	N	Europe	Ornamental
Feral Goat	<i>Capra hircus</i>		N		Animal husbandry
Sika Deer	<i>Cervus nippon</i>	N	N	Asia-Temperate	Ornamental

Fallow Deer	<i>Dama dama</i>	N	N	Europe	Ornamental
Feral Cat	<i>Felis catus</i>	N			Biocontrol
Edible Dormouse	<i>Glis glis</i>		N	Europe	Ornamental
Chinese Water Deer	<i>Hydropotes inermis</i>	N	N	Asia-Temperate Northern	Ornamental
American Bullfrog	<i>Lithobates catesbeianus</i>	N	N	America	Ornamental
Reeves's Muntjac	<i>Muntiacus reevesi</i>	N	N	Asia-Temperate	Ornamental
House Mouse	<i>Mus domesticus</i>	N	N	Europe	Stowaway Animal
Feral Ferret	<i>Mustela furo</i>	N	N		husbandry Animal
American Mink	<i>Mustela vison</i>	N	N		husbandry
Rainbow Trout	<i>Oncorhynchus mykiss</i>	N		Northern America	Animal husbandry
Rabbit	<i>Oryctolagus cuniculus</i>		N	Europe Northern	Animal husbandry
Ruddy Duck	<i>Oxyura jamaicensis</i>	N		America	Ornamental Animal
Common Pheasant	<i>Phasianus colchicus</i>	N		Asia-Temperate	husbandry
Ring-Necked Parakeet	<i>Psittacula krameri</i>	N		Asia-Tropical	Ornamental
Brown Rat	<i>Rattus norvegicus</i>	N	N	Asia-Temperate	Stowaway
Ship Rat	<i>Rattus rattus</i>	N	N	Asia-Tropical Northern	Stowaway
Grey Squirrel	<i>Sciurus carolinensis</i>	N	N	America	Ornamental
African Clawed Toad	<i>Xenopus laevis</i>	N	N	Africa	Ornamental
Other					
	<i>Coscinodiscus wailesii</i>		N	Pacific	Unknown

Appendix 6. Species designated as having an unknown impact.

	Species
	<i>Acartia (Acartiura) margalefi</i>
	<i>Acartia tonsa</i>
	<i>Agardhiella subulata</i>
Pile worm	<i>Alitta succinea</i>
A sea spider	<i>Ammonothea hilgendorfi</i>
	<i>Anthrenus flavidus</i>
	<i>Aphis oenotherae</i>
	<i>Asellus communis</i>
Harpoon weed	<i>Asparagopsis armata</i>
Little Owl	<i>Athene noctua</i>
Big-eyed Sand-smelt	<i>Atherina boyeri</i>
	<i>Atheta harwoodi</i>
Worm Slug	<i>Boettgerilla pallens</i>
	<i>Bonnemaisonia hamifera</i>
tubificid worm	<i>Branchiura sowerbyi</i>
Wrinkled Snail	<i>Candidula intersecta</i>
Japanese skeleton shrimp	<i>Caprella mutica</i>
Common White Sucker	<i>Catostomus commersoni</i>
Vineyard Snail	<i>Ceruella virgata</i>
	<i>Chromaphis juglandicola</i>
	<i>Corticeus linearis</i>
A brown seaweed	<i>Corynophlaea umbellata</i>
Lesser White-toothed Shrew	<i>Crocidura suaveolens</i>
	<i>Cryphalus asperatus</i>
A red seaweed	<i>Cryptonemia hibernica</i>
	<i>Crypturgus subcribrosus</i>
None known	<i>Desdemona ornata</i>
Killer Shrimp	<i>Dikerogammarus villosus</i>
	<i>Drepanosiphum acerinum</i>
	<i>Dysaphis tulipae</i>
Say mud crab	<i>Dyspanopeus sayi</i>
	<i>Eulepidosaphes pyriformis</i>
Cypress Pug	<i>Eupithecia phoeniceata</i>
Wautier's Limpet	<i>Ferrissia wautieri</i>
	<i>Frankliniella occidentalis</i>
Sideswimmer	<i>Gammarus tigrinus</i>
	<i>Goniadella gracilis</i>

	<i>Gonionemus vertens</i>
Devil's Tongue Weed	<i>Grateloupia turuturu</i>
Siphoned Japan Weed	<i>Heterosiphonia japonica</i>
Marbled Cellar Spider	<i>Holocnemus pluchei</i>
	<i>Idiopterus nephrolepidis</i>
	<i>Illinoia liriodendri</i>
	<i>Illinoia rhododendri</i>
Western Green Lizard	<i>Lacerta bilineata</i>
Greenhouse Slug	<i>Lehmannia valentiana</i>
Pond-Perch	<i>Lepomis gibbosus</i>
tubificid worm	<i>Limnodrilus cervix</i>
	<i>Lithocharis nigriceps</i>
A solitary entoproct	<i>Loxosomella kefersteinii</i>
	<i>Macrorhyncholus littoralis</i>
	<i>Melaphis rhois</i>
Trumpet Ramshorn	<i>Menetus dilatatus</i>
Slender-tube amphipod	<i>Monocorophium acherusicum</i>
Oblong Orb Mussel	<i>Musculium transversum</i>
	<i>Myzaphis turanica</i>
	<i>Myzocallis boeneri</i>
	<i>Myzocallis schreiberi</i>
An amphipod	<i>Orchestia cavimana</i>
	<i>Otiorhynchus (Otiorhynchus) aurifer</i>
	<i>Otiorhynchus (Otiorhynchus) salicicola</i>
Privet weevil	<i>Otiorhynchus (Tournieria) crataegi</i>
	<i>Otiorhynchus armadillo</i>
Oriental prawn	<i>Palaemon macrodactylus</i>
None known	<i>Paralaeospira malardi</i>
Marsh frog	<i>Pelophylax ridibundus</i>
	<i>Pemphigus populitransversus</i>
	<i>Periphyllus acericola</i>
	<i>Periphyllus aceris</i>
	<i>Periphyllus californiensis</i>
	<i>Periphyllus lyropictus</i>
	<i>Philonthus rectangulus</i>
A parasitic copepod	<i>Phyllodictya petiti</i>
European physa	<i>Physella acuta</i>
Tadpole physa	<i>Physella gyrina</i>
A tube worm	<i>Pileolaria berkeleyana</i>
An annelid worm	<i>Pileolaria militaris</i>
	<i>Pineus similis</i>

	<i>Pineus strobi</i>
	<i>Planaria torva</i>
Wall Lizard	<i>Podarcis muralis</i>
A brown seaweed	<i>Pseudolithoderma roscoffense</i>
	<i>Rhodobium porosum</i>
	<i>Rhopalosiphum rufulum</i>
Veitch's Bamboo	<i>Sasa veitchii</i>
	<i>Scolytus laevis</i>
Pygmy Elm Bark Beetle	<i>Scolytus pygmaeus</i>
Lesser Caucasian-Stonecrop	<i>Sedum stoloniferum</i>
	<i>Sitobion ptericolens</i>
	<i>Sitticus pubescens</i>
	<i>Takecallis arundicolens</i>
	<i>Takecallis taiwanus</i>
A centric diatom	<i>Thalassiosira punctigera</i>
A centric diatom	<i>Thalassiosira tealata</i>
White Snail	<i>Theba pisana</i>
A Kelp Fly	<i>Thoracochaeta johnsoni</i>
A Kelp Fly	<i>Thoracochaeta seticosta</i>
	<i>Tinocallis ulmiparvifoliae</i>
	<i>Tinocallis zelkowae</i>
	<i>Tupiocoris rhododendri</i>
	<i>Tuponia brevirostris</i>
	<i>Tuponia mixticolor</i>
	<i>Uroleucon erigeronensis</i>
	<i>Utamphorophora humboldti</i>
	<i>Watersipora subtorquata</i>
Aesculapian Snake	<i>Zamenis longissimus</i>

Appendix 7. Non-native species which are known to have arrived since the onset of the project or have been highlighted by scheme experts as missing from the species register

Arocatus longiceps
Closterotomus trivialis
Dicyphus escalerae
Nysius huttoni
Macrotylus horvathi
Brachynotocoris punctipennis
Dicyphus pallicornis
Reuteria marqueti
Rhyparochromus vulgaris
Conostethus venustus
Eremocoris fenestratus
Cyphostethus tristriatus
Mytilus galloprovincialis
Mytilus trossulus
Anastrepha obliqua
Anastrepha species indet.
Bactrocera cucurbitae
Ceratitis cosyra
Cheilosia caerulescens
Chymomyza amoena
Dacus ciliatus
Dasineura oleae
Dasineura oxycoccana
Didactylomyia longimana
Lestodiplosis pini
Macrolabis aquilegiae
Medetera grisescens
Megaselia dimorphica
Phytomyza astrantiae
Phytomyza gymnostoma
Scaptomyza adusta
Sciophila fractinervis
Sphegina sibirica
Taomyia marshalli
Tephritis divisa
Elaphe guttata
Triturus marmoratus

Bufo (Pseudepidalea) viridis
Cynops spp. and Paramesotriton spp.
Bombina orientalis
Lampropeltis sspp.
Thamnophis spp.
Synophropsis lauri
Liguropia juniperi
Arocatus longiceps
Closterotomus trivialis
Dicyphus escalerae
Nysius huttoni
Zyginella pulchra
Idiocerus heydenii
Edwardsiana nigriloba
Eurhadina loewii
Batracomorphus allionii
Zygina nivea
Dryodurgades antoniae
Macrotylus horvathi
Brachynotocoris punctipennis
Dicyphus pallicornis
Reuteria marqueti
Rhyparochromus vulgaris
Conostethus venustus
Eremocoris fenestratus
Cyphostethus tristriatus
Synophropsis lauri
Liguropia juniperi
Zyginella pulchra
Acericerus heydenii
Edwardsiana nigriloba
Eurhadina loewii
Batracomorphus allionii
Zygina nivea
Dryodurgades antoniae
Fieberiella florii
Fieberiella septentrionalis
Viridicerus ustulatus
Acericerus ribauti
Anoterostemma ivanhofi

