Pathology & Laboratory Medicine Children's Hospital Los Angeles 4650 Sunset Boulevard Los Angeles, CA 90027 Laboratory Service Center: 323-361-2423



877-KIDZLAB

Alexander R. Judkins, MD Pathology-in-Chief

Richard Ko, MDMedical Director,
Clinical Coagulation Lab

PLATELET FUNCTION / PLATELET AGGREGATION WITH ATP RELEASE TEST REQUEST FORM

Institution/Ordering Physician: Billing Address:	Report Address: (if different from billing address)
Phone:Fax: Account Number: PATIENT INFORMATION	Phone: Fax: CA License #: NPI#: PHYSICIAN INFORMATION
Name:(Last name) (First Name)	Requesting Physician: Physician Signature: (not required for institutions)
Institution Patient ID/MR#: Date of Birth: Gender: Male Female	Diagnosis Code(s):
FAX THE REQUISITION AND PATIENT BLEEDING HISTORY FORM TO 323-361-6157 FOR INTERNAL USE ONLY	
Collection Date: Time:Collector ID:	:FIN: MR#:
TEST(S) REQUESTED	
□ *PLATELET FUNCTION SCREEN (DADE BEHRING PFA 100) (Available M-F, 7AM to 6PM, no appointment needed) Test includes Collagen/Epinephrine and/or Collagen/ADP. If Collagen/Epinephrine is abnormal, reflex to Collagen/ ADP.	*PLATELET AGGREGATION AND ATP RELEASE STUDIES (Available Tuesdays only, please schedule an appointment) Full Panel (ADP, COLLAGEN, EPINEPHRINE, RISTOCETIN, ARACHNIDONIC ACID) Ristocetin (Induced Platelet Aggregation Panel)

INSTRUCTIONS

- 1. A Platelet Aggregation blood test is ordered by your doctor.
- 2. This test is performed at Children's Hospital Los Angeles by appointment only.
 - Call 323-361-2423 from 6:00 AM 7:00 PM to schedule an appointment.
 - The test is performed on Tuesdays only.
- 3. Arrive at Children's Hospital Los Angeles Laboratory 30 minutes prior to your appointment time.
 - We are located at 4650 Sunset Boulevard, Los Angeles, CA 90027.
 - Check-in at the front desk and bring this form and the patient bleeding history to the Outpatient Laboratory.
 - The Outpatient Laboratory is located on the 1st Floor of the Gateway Building.
- 4. Do not ingest Aspirin or NSAIDs (Non-Steroidal Anti-Inflammatory Drugs) for 10 days prior to this test unless your physician specifically instructs you to do so.
 - If you have been taking Aspirin or NSAIDs, please make sure the information is included on this form.
- 5. Have a light, low-fat breakfast on day of testing.
- 6. Bring ID and insurance information on day of testing.