



# Cotton Pest Management Guide 2021-22



The latest in cotton RD&E  
brought to you by  
**CRDC and CottonInfo**




*Best Practice*





**More resilience,  
more productivity,  
powered by biology**



**EndoFuse™** from Sumitomo Chemical is a plant and soil enhancement product that contains arbuscular mycorrhizae fungi (AMF). Mycorrhizae are beneficial fungi that naturally exist in soils colonising the root systems of plants. EndoFuse includes 4 high performing endo-mycorrhizae species that have been proven to increase crop resilience, productivity and overall plant and soil health.

**KEY AREAS ENDOFUSE HAS BEEN SHOWN TO IMPACT:**

- Crop resilience under plant stress conditions
- Reduce long fallow disorder
- Crop yield
- Root and shoot biomass
- N, P, K and trace mineral uptake
- Water uptake during moisture stress
- Soil health
- Re-populate mycorrhizae following cultivation

 **SUMITOMO CHEMICAL**  
AgroSolutions Division

[www.sumitomo-chem.com.au](http://www.sumitomo-chem.com.au)

™ EndoFuse is the trademark of Sumitomo Chemical Australia.

# PLAN FOR GROWTH

## Longterm, growing success requires dedicated planning

With over 40 years of dedicated cotton experience, CGS's expertise and local agronomic knowledge provides you the insights needed to grow consistent and high yielding crops.

CGS is committed to service the Australian cotton growing community from our strategically located branch network servicing all cotton valleys, nationwide.

Seed, technology, chemistry,

crop nutrition, agronomy, digital tools and local advice, CGS is here to help you plan for growth.

Find out more: please contact your local CGS cotton specialist today.

Growing Success

#cgsaustralia



#growingsuccess



**CGS**

[cgs.com.au](http://cgs.com.au)



## Bollgard® 3 Roundup Ready Flex® cotton

Combining leading insect and weed control technology to enable growers to manage their crop with more flexibility.



To find out more visit [bollgard3.com.au](http://bollgard3.com.au)








Bollgard® and Roundup Ready Flex® are registered trademarks of the Bayer Group. Insect control technology incorporated into these seeds is commercialised under a license from Syngenta Crop Protection AG. Bayer CropScience Pty Ltd, ABN 87 000 226 022 Level 1, 8 Redfern Road, Hawthorn East, VIC 3123. Phone: 1800 636 001 [crop.bayer.com](http://crop.bayer.com) Copyright © 2021 the Bayer Group. All rights reserved.

# Contents...




|                 |   |
|-----------------|---|
| Foreword        | 4 |
| Index of tables | 4 |

## INSECTS

|   |    |  |
|---|----|--|
| Integrated Pest Management (IPM)                              | 5  | Sponsored by –  |
| Protecting beneficials  | 9  | Sponsored by –  |
| Management of key insect and mite pests                       | 16 | Sponsored by –  |
| Insecticide Resistance Management Strategy (IRMS) for 2021-22 | 54 | Sponsored by –  |
| Bollgard® 3 Resistance Management Plan (RMP)                  | 61 | Sponsored by –  |
| Managing unsprayed pigeon pea refuges                         | 71 |  |


## WEEDS

|                                  |    |  |
|----------------------------------|----|--|
| Integrated Weed Management (IWM) | 73 |  |
| Key weeds                        | 80 |  |
| Herbicide resistance management  | 81 | Sponsored by –  |
| Herbicide tolerant technology    | 89 |  |


## VOLUNTEER & RATOON COTTON

|   |    |  |
|---|----|--|
| Management of volunteer and ratoon cotton | 92 | Sponsored by –  |
|---|----|--|


## DISEASES

|                                     |     |  |
|-------------------------------------|-----|--|
| Integrated Disease Management (IDM) | 96  |  |
| Key diseases of cotton              | 99  | Sponsored by –  |
| Regional disease update 2020-21     | 109 |  |


## BIOSECURITY

|  |     |  |
|--|-----|--|
| Biosecurity – we all have a responsibility | 114 |  |
| High priority exotic pests and diseases    | 118 | Sponsored by –  |

## PGRs & HARVEST AIDS

|   |     |  |
|---|-----|--|
| Cotton growth regulators and harvest aids | 122 | Sponsored by –  |
|---|-----|--|

## SPRAY APPLICATION

|                                     |     |  |
|-------------------------------------|-----|--|
| Best practice for spray application | 123 |  |
| Pesticides and regulation           | 130 |  |
| Using pesticides safely             | 132 | Sponsored by –  |
| Pesticides and the environment      | 136 |  |
| Protecting pollinators              | 138 |  |

|                               |     |
|-------------------------------|-----|
| Glossary and acronyms         | 140 |
| Coordinators and contributors | 142 |
| Index                         | 143 |

Cover photos courtesy Tonia Grundy and Linda Smith.



# Foreword

**Tonia Grundy** (DAF), **Susan Maas** (CRDC), and **Warwick Waters** (CottonInfo)

**W**elcome to the 2021-22 *Cotton Pest Management Guide*. This Guide provides you with a comprehensive summary of the key cotton crop protection issues, and is brought to you by the organisations responsible for cotton industry research, development and extension (RD&E): the Cotton Research and Development Corporation (CRDC) and CottonInfo.

CRDC invests in RD&E projects for the Australian cotton industry. A partnership between the Australian cotton industry and the Australian Government, CRDC exists to enhance the industry's performance. In 2021-22, CRDC will invest \$18.9 million into RD&E projects on behalf of growers and the Government, in collaboration with over 100 research partners.

CottonInfo is an initiative of the joint venture partners: CRDC, Cotton Australia and Cotton Seed Distributors. It is designed to connect you – our cotton growers and consultants – with research and provide you with information, when and where you need it. The CottonInfo team takes the research and development invested in by CRDC and turns it into practical information and knowledge, applicable to you and your farm.

CottonInfo integrates closely with the industry's best management practices program, *myBMP*, supported by Cotton Australia and CRDC, which sets the industry's best practice performance criteria and provides a framework by which growers can participate in, and be accredited in, best practice.

This Guide, along with its sister publication, the *Australian Cotton Production Manual*, are two of the key ways that CRDC and CottonInfo provide the latest in cotton industry RD&E out to you each year. The manual contains additional information on spray application and integrated management and is available to download from the CottonInfo and CRDC websites ([www.cottoninfo.com.au/publications](http://www.cottoninfo.com.au/publications) or [www.crdc.com.au/publications](http://www.crdc.com.au/publications)).

This year the editing team has conducted a major review of all chapters. Notable additions to this edition include the new silverleaf whitefly (SLW) decision support tool, information on reoccurring wilt, and many additional photographs of key pest and beneficial species.

Remember, the CottonInfo team of regional extension officers, technical leads and *myBMP* experts are standing by to assist you with all your cotton information needs (you can find our contact details on the inside back cover).

You can also find information from the CottonInfo team online at our website ([www.cottoninfo.com.au](http://www.cottoninfo.com.au)), while best practice information for your farm is available at the *myBMP* website ([www.mybmp.com.au](http://www.mybmp.com.au)). And you can find information about all of CRDC's investments online at [www.crdc.com.au](http://www.crdc.com.au).

On behalf of CRDC and CottonInfo, thank you to the team of authors, reviewers and contributors from across the cotton research community and the wider industry for their invaluable assistance with this publication. We hope you find this year's *Cotton Pest Management Guide* a valuable and informative reference.



## INDEX OF TABLES

| No. | Title   | Page |
|-----|---|------|
| 1   | Seasonal activity plan for IPM  | 8    |
| 2   | Friends in the field  | 9    |
| 3   | Impact of insecticides at planting or as seed treatments on beneficials | 12   |
| 4   | Impact of insecticides and miticides on beneficials                     | 14   |
| 5   | Control of aphids   | 22   |
| 6   | Control of green vegetable bug  | 26   |
| 7   | Control of <i>Helicoverpa</i>   | 30   |
| 8   | Control of solenopsis mealybug  | 31   |
| 9   | Mirid thresholds  | 35   |
| 10  | Control of mirids   | 36   |
| 11  | Estimates of yield loss due to two-spotted spider mites                 | 39   |
| 12  | Control of two-spotted spider mites                                     | 40   |
| 13  | Control of wireworm   | 44   |
| 14  | Control of thrips   | 46   |
| 15  | Control of silverleaf whitefly  | 51   |
| 16  | Herbicides for use in pigeon pea  | 72   |
| 17  | Control of weeds in dry channels  | 73   |
| 18  | Herbicide plant-backs – rotations to cotton                             | 75   |
| 19  | Herbicide plant-backs – seedbed preparation                             | 77   |
| 20  | Knockdown efficiency  | 77   |
| 21  | Herbicide plant-backs – cotton to rotations                             | 78   |
| 22  | Glyphosate resistant weeds  | 81   |
| 23  | Paraquat resistant weeds  | 82   |
| 24  | 2,4-D resistant weeds   | 83   |
| 25  | Resistance risk of cotton herbicides                                    | 88   |
| 26  | Herbicides registered for volunteer cotton                              | 94   |
| 27  | Control of large volunteer or ratoon plants                             | 94   |
| 28  | Fungicides in cotton  | 108  |
| 29  | Regional disease update – QLD   | 111  |
| 30  | Regional disease update – NSW   | 113  |
| 31  | Commonly used pesticides  | 135  |

### DISCLAIMER

This document has been prepared by the authors for CRDC in good faith on the basis of available information. While the information contained in the document has been formulated with all due care, the users of the document must obtain their own advice and conduct their own investigations and assessments of any proposals they are considering, in the light of their own individual circumstances.

The document is made available on the understanding that the CRDC, the authors and the publisher, their respective servants and agents accept no representation, statement or information whether expressed or implied in the document, and disclaim all liability for any loss, damage, cost or expense incurred or arising by reason of any person using or relying on the information claimed in the document or by reason of any error, omission, defect or mis-statement (whether such error, omission or mis-statement is caused by or arises from negligence, lack of care or otherwise).

While the information is considered true and correct as at June 30, 2021, changes in circumstances after the time of publication may impact on the accuracy of the information. The information may change without notice and the CRDC, the authors and the publisher and their respective servants and agents are not in any way liable for the accuracy of any information contained in this document.

Recognising that some of the information is provided by third parties, the CRDC, the authors and the publisher take no responsibility for the accuracy, currency, reliability and correctness of any information included in the document provided by third parties.

The product trade names in this publication are supplied as examples on the understanding that no preference between equivalent products is intended and that the inclusion of a product does not imply endorsement by CRDC over any other equivalent product from another manufacturer. Any information regarding pesticides or their use in this publication does not constitute a recommendation for that particular use by the authors, the author's organisations or CRDC. All pesticide applications must accord with the currently registered label or permit for that particular pesticide, crop, pest and region.

All work in this publication is copyright – however, CRDC encourages wide dissemination of its research. In line with the fair dealing provisions of the *Copyright Act 1968*, information in this publication can be used for personal, educational or any other non-commercial purposes, providing CRDC is clearly acknowledged. Information used for any other purpose must only be done so with the prior written permission of CRDC. Requests and enquiries concerning reproduction and rights should be addressed to the CRDC Communications Manager.

Bollgard® 3, Roundup Ready Flex®, Roundup Ready PLUS® and PLANTSHIELD® are registered trademarks of the Bayer Group.

ISSN 1442-8462 (Print)

Published by the Cotton Research and Development Corporation

Production by Greenmount Press, August 2021 (Ph: 07 4659 3555)

# Integrated Pest Management (IPM)

Integrated Pest Management could really be called Intelligent Pest Management as it aims to use our knowledge and understanding of the pest, the crop, and the environment to minimise the likelihood of pest outbreak and reduce our reliance on insecticides. Some pest populations (such as mites, mealybug and whitefly) have limited chemical options for control and populations can quickly increase if IPM is not implemented.

IPM is not a recipe, but instead requires the crop manager to implement a mix of pre-emptive and responsive pest management actions to reduce the risk of crop loss and improve the health of both the environment and ourselves.

The outcome of an effective IPM system is long term stable management of pests and beneficials, reducing the risk of resistance, so that economic losses of crop yield and quality and threats to human health and the environment can be minimised.

## IPM – the basics

While IPM is not a recipe, there are some time-honoured practices that form a solid basis for implementing IPM on your farm. These are:

- **Farm hygiene.** Minimising cotton volunteers, ratoons and other weeds in fallow fields, field edges, roadways and drainage lines on your farm is a simple step that can limit the survival and spread of overwintering pests such as mealybugs or resistant aphids that might also carry bunchy top disease. **PUT SIMPLY, CLEAN FARMS TEND TO HAVE LESS PEST PROBLEMS THAN DIRTY ONES.**
- **Preventing problems.** Sowing date, crop sequences and field selection are just some of the tactics that you can use to disadvantage pests on your farm. Avoiding back-to-back cotton in fields that had mealybugs to limit season carryover, or not planting as late so that your crop is not exposed to displaced populations of SLW are just two examples of how to strategically avoid giving pests the upper hand.
- **Grow a healthy crop.** Cotton plants have a significant ability to recover from damage (especially early season damage) with no reduction in yield or delay in maturity. Plant monitoring in conjunction with regular insect monitoring allows an assessment of pest effects that might be difficult to detect in regular sampling. Plant monitoring can

## Dryland cotton...

- **Dryland cotton has the same pests and beneficials as irrigated crops. Effective IPM and resistance management are critical regardless of where or how cotton is grown.**
- **A key IPM challenge for dryland crops in minimum tillage systems is to ensure 100% crop destruction. Ratoon cotton or volunteers that emerge in a subsequent grain crop can provide a green bridge for pests and diseases. Crop mulching or slashing alone at the end of the season is not sufficient to prevent pest carryover unless it is followed with tactics that prevent regrowth.**

## Best practice...

- **KNOWLEDGE is the key that unlocks effective IPM.**
- **Seek to PREVENT pest outbreaks through good farm hygiene, complementary crop rotation and conserving beneficial populations.**
- **Regularly MONITOR pest species, beneficials, crop stage, crop growth and the weather.**
- **Bring it all together with effective management ACTION that is mindful of pest thresholds, resistance risks and potential impacts on natural enemies, bees and the environment.**

assist in decision making where pest levels are just below threshold or where there are combinations of pests present. Acceptable damage levels will vary depending on crop physiology stage, yield expectations and climatic conditions.

- **Sample your crop effectively.** Detecting and being able to quantify both pests and beneficials is a fundamental requirement for good decision making. Depending on the crop stage and pest or beneficial type, sampling techniques may vary. While some thresholds only require monitoring of one lifecycle stage, it can be useful to be aware of all life stages to identify if a population has built up within the crop or has migrated in recently.
- **Take note of where your crop is at.** The development stage of your crop has a big bearing on (i) whether it is susceptible to yield loss and (ii) if it can recover from pest damage without intervention.
  - Minimal squares on a normal crop at 10-12 nodes might indicate poor retention due to pest or weather damage, but if the plant has been developmentally delayed, the retention levels may be appropriate.
  - Cotton can compensate for early season damage, particularly vegetative plant parts, if growing conditions remain favourable.
  - It is important to be able to distinguish between vegetative and fruiting branches when measuring retention during early crop development. View a CottonInfo video that explains the difference between branch types in young crops: [youtu.be/7PXIAuinmeY](https://youtu.be/7PXIAuinmeY)
  - Many pests become less important as bolls open, but SLW or aphids (that can contaminate open cotton with honeydew) become more important.
- **Know your enemy** – this guide outlines the key pests that you would expect to deal with on a regular basis when growing cotton. A key to managing these pests is to understand what they are, what damage they cause, preventative steps you can take to minimise their numbers or impact, how best to sample for them, when control is likely to be required (action thresholds), and what types of control techniques (chemical, biological or physical etc) are likely to be suitable and effective. Knowledge of the pest's interaction with the environment can be important in determining whether a spray is warranted. For example, two-spotted mite populations can be suppressed by cool conditions, but will increase rapidly when it is hot and dry.
- **Know your friends.** Cotton crops host a diverse array of beneficials that can suppress pests, particularly mites, SLW, aphids and mealybugs. Noting which beneficials are present when sampling can help inform your control decisions. Conserving beneficials can have a significant impact on whether you develop a mite or mealybug problem later in the season. Table 4 (see pages 14-15) allows you to contrast and compare the potential impact of different products so that you can make a balanced control decision.

- **Do you even need to spray?** Many pest insects have scientifically-based action thresholds that can help you decide whether a given level of pests or crop damage might require control action. Resist the urge to use insecticides prophylactically with herbicide application operations. Economic thresholds are defined as the pest density or damage level at which control must be implemented to prevent economic loss, and should be considered in the context of other factors that may influence the need to spray. For pest abundance just over threshold but with low damage and high beneficial populations, consider delaying control for several days. This is a low risk strategy to allow time for beneficials to reduce pest levels to below threshold, thereby avoiding a potentially disruptive spray and reducing insecticide costs and selection for resistance. Conversely, for high pest damage with low beneficial populations, immediate control with an insecticide may be the best option. Thresholds for cotton aphid, two-spotted mite and SLW are based on cumulative population changes and require comparison of multiple samplings to determine if action thresholds have been reached.
- **Choose an insecticide wisely.** When insecticide control is warranted make sure the product you have selected appropriately balances effectiveness against the target pest with any potential harm to natural enemies. Being aware of the product's mode of action (MoA) group, and actively avoiding the use of broad-spectrum disruptive insecticides (e.g. organophosphates, pyrethroids and some neonicotinoids) wherever possible, especially early to mid-season, will go a long way to reducing mite, SLW and mealybug numbers later in the season. Also be aware of restrictions on the use of certain products as part of the Industry's Insecticide Resistance Management Strategies (IRMS). If several pesticide options are available, examine their selectivity profiles (see Table 4). For example, clothianidin will reduce populations of ladybird beetles (aphid predators) and *Eretmocerus* wasps (whitefly parasitoids) but conserve predatory bugs and thrips (mite predators). In contrast, the low rate of fipronil will reduce predatory bug populations, and conserve ladybird beetles, but its impact on the key wasp parasitoids of whitefly is unknown. Increases in populations of non-target pests such as aphid, mite and whitefly may follow insecticide applications if the beneficial populations keeping them in check are disrupted. Lower registered rates of a product may provide sufficient efficacy against the target pest, while minimising impact on beneficials.
- **Preserve the usefulness of insecticides and Bollgard.** The ability of insecticides or *Bacillus thuringiensis* (Bt) protein-expressing Bollgard crops to provide control depends on pest populations being susceptible to those insecticides or insecticidal compounds. Follow the label directions and consider both your application technique and environmental conditions to optimise chemical efficacy. Adhering to industry resistance management programs is important to ensure that the industry continues to have access to pest control tactics that give good control.
- **Look after bees.** Bees are particularly susceptible to many of the insecticides used on cotton farms, such as abamectin, fipronil, indoxacarb, pyrethroids and profenofos. The productivity of hives can be damaged by direct or indirect contact with the applied product by foraging bees, or when insecticide drifts over hives or neighbouring vegetation. Always look for and follow label directions regarding impact on bees and refer to the *Protecting pollinators* section for more information on how to manage the risk.

- **Working with your neighbours.** Pests and beneficials do not recognise farm boundaries. Working with your neighbours to better coordinate planting, farm hygiene and crop spraying can provide benefits for the management of pests such as SLW on an area-wide basis. In some regions growers meet regularly at area wide management (AWM) meetings to discuss pest issues and work towards shared solutions. Talk with your local CottonInfo Regional Extension Officer (REO) about any AWM groups in your valley.

## How do I implement IPM?

The great thing about IPM is that it does not have to be complicated – it's an evolving process that builds on taking practical steps to reduce the survival of pests. For example, a simple first step could be improved on-farm hygiene. This might be followed by a decision to avoid the use of a particular type of disruptive insecticide.

For first time growers, the easiest way to start your IPM journey is to employ the services of an experienced consultant to help guide you through the process of pest management decision making. A consultant will be able to provide information on what is going on in your crop and advise you on various management options that might be applicable to your situation.

For the more experienced manager familiar with pest management principles, consider challenging your boundaries by asking yourself some simple 'WHY' questions such as "Why were silverleaf whitefly (SLW) higher or lower than last season?" or "Why am I using this particular insecticide?" The answers might provide insight into the most appropriate next steps for your management program. These types of questions are also good starting points in discussions with your advisor.

The important thing about IPM is to appreciate that biological systems are continually evolving along with our knowledge about pest species and control options. Keeping up to date with the latest information is essential for effective IPM. The latest research findings are made available to growers and advisors via industry meetings and a range of CottonInfo information products, including this guide.

As your IPM program evolves, new practices will bring prospective benefits as well as potential trade-offs (e.g. the use of a more selective insecticide may cost more to apply than a broad-spectrum option but it reduces the likelihood of secondary pest outbreak which may require further spraying). The key is to learn as you go, build on the successes that you have, and when things change, be prepared to adapt what you are doing to suit new challenges.

### Useful resources:

***Pest and Beneficial Insects in Australian Cotton Landscapes.* Available from [www.cottoninfo.com.au](http://www.cottoninfo.com.au)**

### CottonInfo videos:

**What is IPM?** [youtu.be/BdQRLx4hN5o](https://youtu.be/BdQRLx4hN5o)

**IPM in action** [youtu.be/k5-0XrzBEeg](https://youtu.be/k5-0XrzBEeg)

**Conserving beneficials** [youtu.be/q3GYrt6QoN8](https://youtu.be/q3GYrt6QoN8)

III



# What's to Admire?



## Applaud<sup>®</sup>

**INSECTICIDE**

**Give Silverleaf Whitefly the send-off it deserves.**

Silverleaf whitefly is a major pest in cotton, it has the ability to contaminate and downgrade lint quality. Help protect the reputation of Australian cotton and rotate to **Applaud<sup>®</sup> Insecticide** to manage whitefly resistance.

- Unique Mode of Action for SLWF
- No known resistance in cotton crops
- Selective to key beneficial insects
- Won't flare secondary pests
- Use early as part of a resistance management strategy

**TABLE 1: Seasonal activity plan for IPM**

|  | <b>Overwinter &amp; planning</b>   | <b>Planting – first flower</b>   | <b>Flower – first open boll</b>   | <b>Open cotton – harvest</b>   |
|--|--|--|---|--|
| <b>Develop an IPM strategy</b>   | Review last season's IPM approach.<br>Communicate IPM goals and pesticide application management plan (PAMP) for the coming season.  | Good record keeping supports PAMP, regulatory requirements and allows end of season assessment of IPM strategy.  |   |  |
| <b>Know your enemy</b>   | Get the latest guides and IPM related information.   | Participate in IPM training, field days, or workshops: Contact your local CottonInfo Regional Extension Officer (REO) (see inside back cover for contact details) to join mailing list or go to <a href="http://www.cottoninfo.com.au/subscribe">www.cottoninfo.com.au/subscribe</a> |   |  |
| <b>Take a year round approach</b>  | Manage winter crops carefully to avoid disrupting beneficial populations.<br>Plan ahead to ensure insecticides are available.  | Consider the summer cropping plan and pest risk.   | Begin planning for rotation crops.  | Reduce pest risk for next season by considering rotation crop type and location.   |
| <b>Think beyond the crop</b>   | Participate in area wide management (AWM) all year round.<br>Apply IPM to all crops.<br>Consider rotation crops (type, location, and potential to host pests and disease).<br>Establish and maintain communication with bee keepers in the region.<br>Avoid spray drift.<br>Consider native vegetation when managing pests. Maximise its value by improving its health, linking patches of vegetation, controlling weeds and keeping it diverse for a range of species (including birds and bats). |  |   |  |
| <b>Have good on-farm hygiene</b>   | Zero tolerance to volunteer cotton in entire landscape all year.<br>Ensure a host free period for pests and diseases.<br>Keep farm weed free all year.<br>Where practical remove weeds from native vegetation areas.   | Consider pre-irrigation, to allow control of cotton volunteers and other weeds with non-glyphosate control prior to planting.<br>Consider in-crop cultivation where necessary.   | Continue to manage volunteer cotton in entire landscape (e.g. fence lines, channels, perennial vegetation and pastures).<br>Consider chipping.            | Conduct effective crop removal to prevent ratoons.   |
| <b>Practice Come Clean. Go Clean. all year round</b>   |  |  |   |  |
| <b>Consider options to avoid or reduce pests</b>   | When planning cotton, consider paddock history, proximity to sensitive areas, pest hosts and beneficial habitats.<br>Manage areas of vegetation to encourage beneficials.<br>Assess risk of soil pests before planting.  | Use a suitable variety for your region.<br>Provide optimum planting conditions to promote healthy seedlings.<br>Consider summer trap crop.<br>Consider insecticide choice or releasing natural enemies to build beneficial numbers.  | Monitor crop development to maintain a healthy crop.<br>Maintain high beneficial numbers.   | Slash and pupae bust last generation summer trap crop (follow guidelines).<br>Follow pupae busting guidelines for Bt cotton.<br>Practice Come Clean. Go Clean. to prevent spread of pests on, off and around farm. |
| <b>Sample crops effectively and regularly</b>  | Remain up-to-date with key pests, beneficials, crop sampling and plant damage monitoring.  | Sample for pests, beneficials, parasitism, fruit load and plant damage at least twice weekly throughout the season.<br>Consider both current numbers and population trends.<br>Use pest and damage thresholds and the beneficial to pest ratio.                                      |   |  |
| <b>Grow a healthy crop</b>   | Consider the best rotation crop for your situation.<br>Soil test to determine fertiliser requirements for cotton crop.<br>Consider potential disease risks.  | Provide optimum planting conditions to promote healthy seedlings that can outgrow damage.<br>Monitor leaf and tip damage and development of first squaring node.   | Monitor crop development, fruit retention, nodes above white flower and vegetative growth.<br>Manage nutrition and irrigation to maintain a healthy crop. | Monitor crop development, nodes above cracked boll and percentage of open bolls for defoliation decisions.<br>Manage nutrition and irrigation to avoid or reduce regrowth that may harbour pests.                  |
| <b>Use established thresholds</b>  | Use thresholds and careful spray selection for all crops.  | Use pest and damage thresholds relevant to region, time of season and sampling method.<br>Consider the beneficial to pest ratio taking parasitism into account.  |   |  |
| <b>Choose insecticides wisely</b>  | Use thresholds and careful spray selection for all crops.  | Consider insecticide selectivity and impact on beneficials and bees.<br>Avoid early season use of broad-spectrum sprays.<br>Consider edge or patch spraying for aphids and mites.<br>Avoid prophylactic sprays.  | When choosing insecticides think about impact on beneficials, including bees.   | Defoliation may be a late season alternative to an insecticide.  |
| <b>Apply good resistance management principles</b>   | Complete pupae busting (follow guidelines).<br>Zero tolerance of volunteer and ratoon cotton in the entire landscape.  | Adhere to refuge requirements.<br>Consider choice of at-planting insecticides/seed dressings and implications for later sprays.  | Ensure that Bt cotton refuges are attractive/effective.   | Follow pupae busting guidelines for Bt cotton.   |
| <b>Use pest thresholds and follow your Insecticide Resistance Management Strategy (IRMS) for every spray</b> |  |  |   |  |

# Protecting beneficials

**N**ot all arthropods found in cotton are pests. Some are just passing through, and some are actively providing benefits, either to the crop directly (e.g. predators or parasitoids of crop pests; often referred to as ‘natural enemies’) or to the local ecosystem (e.g. pollinators).

Building and/or conserving beneficial populations is at the heart of integrated pest management (IPM). Pest management decisions need to be well informed and supported by effective sampling and valid control thresholds. Knowledge of the beneficials present, their activity on pests, and the potential impacts of other pest control measures is essential to ensuring the best management decisions are made for your crop.

## Choose insecticides wisely to conserve beneficials

The contribution of beneficials should be balanced against the need to protect the crop from significant loss. Where insecticide control is warranted (based on industry recommended monitoring and thresholds), product selection should carefully consider:

- Efficacy on the pest.
- Impact on the beneficial population(s), including bees (see Tables 3 and 4).
- Allowable usage in the cotton industry’s Insecticide Resistance Management Strategy (IRMS).

Always aim to maximise effectiveness against the pest while minimising impact on beneficials. Take into consideration other factors such as rate, application technique and other products used in tank mixes that could either improve or hinder efficacy, along with the potential for unintended consequences such as flaring of non-target pests or increased resistance (including cross-resistance) in the target or secondary pests.

Some insecticides are very selective and have little impact on beneficial insects (often referred to as ‘soft’), while others are highly disruptive to beneficial populations (‘broad-spectrum or ‘hard’). Knowing the selectivity of an insecticide helps to assess the risk that following its use, populations of other pests may ‘flare’ (increase rapidly). The relative selectivity of insecticides available for use in cotton can be found in Tables 3 and 4 (which have been largely based on industry-funded research).

In addition to general impact, Table 4 lists the product’s selectivity to specific types of beneficials. For example, if mealybugs are present, look for insecticides that have less effect on parasitic wasps and key mealybug predators such as lacewings and ladybird beetles. When products are registered for a range of rates, using lower rates may provide sufficient efficacy against some target pests, while minimising impact on beneficials. Note that these ratings are based on results after a single application, and multiple applications of a product with a low rank can still have a cumulative disruptive impact.

When selecting an insecticide, refer to the IRMS (pages 56-57) to reduce the risk of resistance, and adhere to label directions for the specific product.

**CottonInfo video on conserving beneficials in cotton [youtu.be/q3GYrt6QoN8](https://youtu.be/q3GYrt6QoN8)** III

**TABLE 2: Friends in the field**

|                               | <i>Helicoverpa</i> | Aphids | Mealybug | Spider mites | SLW | Green mirids | Jassids | Thrips | Notes   |
|-------------------------------|--------------------|--------|----------|--------------|-----|--------------|---------|--------|---|
| Ladybird beetles              |                    | X      | X        | X            | X   |              | X       | X      | Also feed on scale insects.   |
| Red and blue beetle           | X                  | X      |          |              | X   |              |         |        | Also predators of slow moving insects.  |
| Damsel bug                    | X                  | X      |          | X            |     | X            |         |        | Will feed on <i>Helicoverpa</i> eggs and up to medium sized caterpillars.                                       |
| Big-eyed bug                  | X                  | X      |          | X            | X   | X            |         |        | Important predator of eggs and small soft-bodied insects.   |
| Brown smudge bug              | X                  |        | X        | X            | X   |              |         |        | Important predator of eggs and small soft-bodied insects.   |
| Apple dimpling bug            |                    |        |          | X            | X   |              |         |        | Can also cause damage, but threshold is 5 times greater than green mirids. Monitor fruit retention.             |
| Glossy shield bug             | X                  |        |          |              |     |              |         |        | Also predators of other caterpillars.   |
| Spined predatory bug          | X                  |        |          |              |     | X            |         |        | Also predators of other caterpillars.   |
| Minute pirate bug             |                    |        |          | X            | X   |              |         | X      | Effective predators of eggs.  |
| Assassin bug                  | X                  |        |          |              |     | X            |         |        | Also predators of other caterpillars.   |
| Lacewings                     | X                  | X      | X        | X            | X   |              |         | X      | In green lacewings, the larval stage is the predator. In brown lacewings, both adults and larvae are predatory. |
| Spiders                       | X                  |        |          | X            | X   | X            | X       |        | Generalist predators.   |
| Parasitoids                   | X                  | X      | X        |              | X   |              |         |        | Species of parasitoid are specific in pests targeted. Monitor for parasitised pests.                            |
| Hoverfly and silverfly larvae |                    | X      |          |              |     |              |         |        | The adult will lay eggs on the crop when there are near aphid colonies.   |
| Thrips                        |                    |        |          | X            | X   |              |         |        | Can be a pest early season.   |

# Examples of beneficial species



**Red and blue beetle.**  
5 mm. (D. McClenaghan)



**Striped ladybird.**  
4 mm. (L. Wilson)



**3 banded ladybird.**  
8 mm. (P. Grundy)



**Variable ladybird.**  
5 mm. (P. Grundy)



**Apple dimpling bug (also known as yellow mirid).**  
3 mm. (J. Wessels)



**Damsel bug.**  
8 mm. (D. McClenaghan)




**Big-eyed bug.**  
3 mm. (M. Dillon)



**Brown smudge bug.**  
4 mm. (C. Mares)



**Glossy shield bug.**  
12 mm. (J. Wessels)



**Spined predatory bug.**  
12 mm. (L. Wilson)



**Adult green lacewing (larval stage is predatory).**  
12 mm. (T. Grundy)



**Green lacewing larva.**  
6 mm. (M. Miles)

# FASCINATE™ DRY

HERBICIDE

**NEW** easy-to-use herbicide for broad-spectrum weed control prior to planting



## Why go DRY?



- 1** Use only 8 kg versus a 20 litre\* drum
- 2** No drums to wash, store & recycle
- 3** Built in surfactant + low foam so its easy to use
- 4** Works on even the hardest to kill grasses
- 5** Give glyphosate a break!

**...it's EASY**



[www.upl-ltd.com/au](http://www.upl-ltd.com/au)

\* Compared to 200g ai/L glufosinate formulations

™ Fascinate is a registered trademark of UPL Australia Pty Ltd



Examples of beneficial species (continued)



**Christmas spider.**  
15 mm. (P. Grundy)



**Minute pirate bug nymph feeding on adult thrips.**  
1-2 mm. (L. Wilson)




**Tangle web spider.**  
5 mm. (P. Grundy)



**Minute pirate bug adult.**  
3 mm. (T. Grundy)



**Two-toned parasite.**  
20 mm. (P. Reid)



**Assassin bug.**  
25 mm. (L. Wilson)



**Adult hoverfly (the larval stage is predatory).**  
9 mm. (P. Grundy)



**Thrips feeding on mite eggs.** 1 mm. (L. Wilson)

**TABLE 3: Impact of insecticides at planting or as seed treatments on natural enemies in cotton**

| Active   | MoA group    | Rate           | Target pest(s) |       |        |        |        | Persistence <sup>1</sup> | Impact <sup>2</sup> |                                |
|--|--------------|----------------|----------------|-------|--------|--------|--------|--------------------------|---------------------|--------------------------------|
|  |              |                | Wireworm       | Mites | Mirids | Aphids | Thrips |                          | Thrips              | Other beneficials <sup>3</sup> |
| <b>At planting (rate = g ai/ha)</b>              |              |                |                |       |        |        |        |                          |                     |                                |
| Phorate  | 1B           | 600            | Y              | Y     | Y      | Y      | Y      | Medium-long              | Very high           | Very low                       |
| Carbosulfan                                      | 1A           | 750-1000       | Y              |       | Y      |        | Y      | Medium-long              | Very high           | Very low                       |
| Chlorpyrifos                                     | 1B           | 250-750        | Y              |       |        |        |        | Medium                   | —                   | Very low                       |
| <b>Seed treatments (rate = g ai/100 kg seed)</b> |              |                |                |       |        |        |        |                          |                     |                                |
| Imidacloprid + Thiodicarb + Fipronil             | 4A + 1A + 2B | 350 + 250 + 50 | Y              |       |        |        | Y      | —                        | —                   | —                              |
| Thiodicarb                                       | 1A           | 500            |                |       |        |        | Y      | Short                    | High                | Very low                       |
| Thiodicarb + Fipronil                            | 1A + 2B      | 259 + 12       | Y              |       |        |        | Y      | Short-medium             | High                | Very low                       |
| Imidacloprid                                     | 4A           | 525            | Y              |       |        | Y      | Y      | Medium                   | Very high           | Very low                       |
|  |              | 700            | Y              |       |        | Y      | Y      | Medium                   | Very high           | Very low                       |
| Thiamethoxam                                     | 4A           | 280            | Y              |       |        | Y      | Y      | Medium                   | Very high           | Very low                       |

1. Persistence: short, 2-3 weeks; medium, 3-4 weeks; long, 4-6 weeks.

2. Impact rating (% reduction in beneficials following application): very low, <10%; low, 10-20%; moderate, 20-40%; high, 40-60%; very high, >60%; —, no data.

3. Other beneficials include predatory beetles, predatory bugs, spiders, wasps and ants.

**Pick Your**  
**MainMan<sup>®</sup>**  
**First!**



Pick MainMan first in your insect control program to help keep beneficial insect populations working for you this season

✓ Proven Performer ✓ IPM Compatible ✓ Unique Mode of Action

***Superior control of mirids and aphids***



[www.upl-ltd.com/au](http://www.upl-ltd.com/au)

© Registered Trademark of Ishihara Sangyo Kaisha, Ltd. Always read and follow the product label prior to use.




TABLE 4: Impact of insecticides and miticides on predators, parasitoids and bees in cotton

| Insecticides<br>(in increasing rank order<br>of impact on beneficials) | Rate (g a/ha) | Target pest(s) |       |        |        |        |                     | Overall ranking <sup>10</sup> | Beneficials              |                    |                   |                                  |                        |                    |                |               |                      |                |                 |         |               |                           | Pest resurgence <sup>12</sup> |      |                      | Toxicity to bees <sup>14</sup> |                 |                 |             |
|--|---------------|----------------|-------|--------|--------|--------|---------------------|-------------------------------|--------------------------|--------------------|-------------------|----------------------------------|------------------------|--------------------|----------------|---------------|----------------------|----------------|-----------------|---------|---------------|---------------------------|-------------------------------|------|----------------------|--------------------------------|-----------------|-----------------|-------------|
|  |               | Helicoverpa    | Mites | Mirids | Aphids | Thrips | Silverleaf whitefly |                               | Persistence <sup>8</sup> | Total <sup>1</sup> | Red & blue beetle | Minute 2-spotted ladybird beetle | Other ladybird beetles | Total <sup>2</sup> | Damselfly bugs | Big-eyed bugs | Other predatory bugs | Apple dimpling | Lacewing adults | Spiders | Total (wasps) | Eretmocerus <sup>18</sup> | Trichogramma                  | Ants | Thrips <sup>21</sup> |                                | Mites           | Aphids          | Helicoverpa |
| Bt <sup>11</sup>   |               | ✓              |       |        |        |        |                     | Very low                      | VL                       | VL                 | VL                | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | VL                        | VL                            | VL   | VL                   |                                |                 |                 | VL          |
| Nucleopolyhydrovirus NPV   |               | ✓              |       |        |        |        |                     | Very low                      | VL                       | VL                 | VL                | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | VL                        | VL                            | VL   | VL                   |                                |                 |                 | VL          |
| Pririmcarb   | 250           |                |       |        | ✓      |        | Short               | Very low                      | VL                       | VL                 | VL                | L                                | L                      | M                  | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | M                         | VL                            | L    |                      |                                |                 | VL              |             |
| BioPest oil  | 2-4%          |                |       | ✓      | ✓      |        | Short               | Very low                      | VL                       | VL                 | VL                | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | L                         | VL                            | M    | VL                   |                                |                 | VL              |             |
| PSO (Canopy) <sup>16</sup>   | 2%            | ✓              |       | ✓      | ✓      |        | Short               | Very low                      | VL                       | L                  | VL                | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | L                         | VL                            | H    | VL                   |                                |                 | VL              |             |
| Pyriproxyfen   | 50            |                |       |        |        |        | Long                | Very low                      | M                        | M                  | M                 | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | VL                        | VL                            | VL   | VL                   |                                |                 | L               |             |
| Sero-X   | 800           | ✓              | ✓     | ✓      |        |        | Short               | Low                           | VL                       | VL                 | L                 | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | VL                        | VL                            | VL   | VL                   |                                |                 | VL              |             |
| Etoazole   | 38.5          |                | ✓     |        |        |        | Short               | Low                           | VL                       | VL                 | L                 | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | VL                        | VL                            | VL   | VL                   |                                |                 | VL              |             |
| Buprofezin   | 440           |                |       |        | ✓      |        | Long                | Low                           | L                        | H                  | VL                | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | VL                        | VL                            | VL   | VL                   |                                |                 | VL              |             |
| Indoxacarb (low+/-salt, oil)   | 60            |                |       |        | ✓      |        | Medium              | Low                           | L                        | L                  | H                 | M                                | VL                     | L                  | L              | M             | VL                   | VL             | H               | M       | VL            | VL                        | H                             | VL   |                      |                                | VL              |                 |             |
| Chlorantraniliprole  | 52.5          | ✓              |       |        |        |        | Long                | Low                           | L                        | M                  | M                 | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VH              | VL      | VL            | L                         | L                             | VL   |                      |                                | VL              |                 |             |
| Dicofol <sup>3</sup>   | 960           |                | ✓     |        |        |        | Long                | Low                           | L                        | L                  | L                 | L                                | L                      | L                  | L              | L             | L                    | L              | L               | L       | L             | L                         | L                             | L    | L                    |                                |                 | VL              |             |
| Afdopropen   | 10            |                |       | ✓      |        |        | Medium              | Low                           | L                        | L                  | M                 | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | VL                        | VL                            | VL   | VL                   |                                |                 | VL              |             |
| Spinetoram   | 48            | ✓              |       |        | ✓      |        | Long                | Low                           | M                        | M                  | H                 | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | M                         | H                             | VH   | VH                   |                                |                 | VL              |             |
| Diafenthiuron  | 350           |                |       | ✓      |        |        | Medium              | Low                           | M                        | H                  | VL                | M                                | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | L                         | H                             | VL   | H                    |                                |                 | VH              |             |
| Pymetrozine  | 150           |                |       | ✓      |        |        | Short               | Low                           | M                        | M                  | M                 | L                                | L                      | L                  | L              | L             | L                    | L              | L               | L       | L             | L                         | L                             | M    | VL                   |                                |                 | M               |             |
| Indoxacarb   | 127.5         | ✓              | ✓     | ✓      |        |        | Medium              | Low                           | H <sup>13</sup>          | L                  | VH                | VL                               | VL                     | VL                 | VL             | L             | M                    | L              | VL              | VL      | VL            | VL                        | VL                            | VL   | VL                   |                                |                 | VL              |             |
| Cyantraniliprole   | 60            | ✓              |       |        | ✓      |        | Long                | Moderate                      | M                        | M                  | VL                | L                                | M                      | M                  | M              | M             | M                    | M              | VH              | VL      | VL            | VL                        | VL                            | VL   | H                    |                                |                 | VH              |             |
| Sulfoxaflor (low)  | 48            |                |       | ✓      | ✓      |        | Medium              | Moderate                      | L                        | L                  | L                 | M                                | VL                     | VL                 | VL             | VL            | VL                   | VL             | VL              | VL      | VL            | M                         | H                             | VL   | H                    |                                |                 | VH              |             |
| Spirotetramat  | 96            |                |       | ✓      | ✓      |        | Medium              | Moderate                      | M                        | L                  | H                 | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | VH              | VL      | VL            | M                         | H                             | M    | M                    |                                |                 | H <sup>15</sup> |             |
| Fonicamid  | 70            |                |       | ✓      | ✓      |        | Medium              | Moderate                      | VL                       | VL                 | VL                | H                                | H                      | H                  | H              | H             | H                    | H              | H               | H       | H             | H                         | H                             | H    |                      |                                | H <sup>15</sup> |                 |             |
| Sulfoxaflor (mid)  | 72            |                |       | ✓      | ✓      |        | Medium              | Moderate                      | M                        | L                  | M                 | H                                | L                      | VL                 | L              | VL            | VL                   | VL             | H               | VL      | VL            | M                         | H                             | H    |                      |                                | VH              |                 |             |
| Dinotefuran (low)  | 18            |                |       | ✓      | ✓      |        | Medium              | Moderate                      | M                        | M                  | M                 | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | H               | VL      | VL            | M                         | M                             | M    |                      |                                | VH              |                 |             |
| Abamectin  | 5.4           | ✓              | ✓     |        |        |        | Medium              | Moderate                      | L                        | M                  | H                 | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | H               | VL      | VL            | M                         | H                             | H    |                      |                                | VL              |                 |             |
| Enamectin benzoate   | 8.4           | ✓              |       |        |        |        | Medium              | Moderate                      | L                        | VL                 | M                 | VL                               | VL                     | VL                 | VL             | VL            | VL                   | VL             | H               | VL      | VL            | M                         | H                             | L    |                      |                                | VL              |                 |             |
| Propargite   | 1500          |                | ✓     |        |        |        | Medium              | Moderate                      | M                        | H                  | H                 | M                                | M                      | H                  | H              | H             | H                    | H              | H               | VL      | VL            | M                         | M                             | M    |                      |                                | VH              |                 |             |
| Acetamiprid  | 22.5          |                | ✓     |        |        |        | Medium              | Moderate                      | M                        | M                  | VH                | H                                | H                      | M                  | M              | M             | M                    | M              | H               | VL      | VL            | M                         | M                             | M    |                      |                                | VH              |                 |             |
| Clothianidin (low)   | 25            |                | ✓     | ✓      |        |        | Medium              | Moderate                      | M                        | VL                 | L                 | M                                | VL                     | VL                 | VL             | VL            | VL                   | VL             | H               | VL      | VL            | M                         | M                             | M    |                      |                                | VH              |                 |             |
| Voliam Flex <sup>23</sup>  |               | ✓              |       | ✓      |        |        | Medium              | Moderate                      | H                        | M                  | VH                | H                                | H                      | L                  | L              | M             | M                    | M              | H               | VL      | VL            | M                         | M                             | M    |                      |                                | VH              |                 |             |
| Amitraz  | 400           | ✓              |       | ✓      | ✓      |        | Medium              | Moderate                      | H                        | M                  | VH                | H                                | H                      | L                  | L              | M             | M                    | M              | H               | VL      | VL            | M                         | M                             | M    |                      |                                | VH              |                 |             |
| Skope (low) <sup>22</sup>  |               |                | ✓     | ✓      |        |        | Medium              | Moderate                      | H                        | H                  | VH                | H                                | H                      | L                  | M              | M             | M                    | M              | H               | VL      | VL            | M                         | M                             | M    |                      |                                | VH              |                 |             |



|                            |      |                 |              |           |    |    |    |    |    |    |    |    |    |    |    |    |    |    |     |     |                 |
|----------------------------|------|-----------------|--------------|-----------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|-----|-----|-----------------|
| Fipronil (low)             | 12.5 | ✓               | Medium       | Moderate  | L  | L  | H  | L  | L  | L  | VH | L  | M  | M  | —  | M  | VH | VH | +ve | +ve | M <sup>15</sup> |
| Chlorfenapyr (low)         | 200  | ✓               | Medium       | Moderate  | M  | L  | VH | VL | M  | H  | VH | L  | L  | M  | —  | VH | H  | M  | +ve |     | VH              |
| Sulfoxaflor (full)         | 96   | ✓               | Medium       | Moderate  | H  | L  | M  | H  | L  | VL | VH | L  | L  | M  | —  | H  | VH | H  | +ve |     | L               |
| Fipronil (high)            | 25   | ✓               | Medium       | Moderate  | L  | VL | H  | L  | M  | H  | VH | L  | L  | M  | M  | M  | VH | VH | +ve |     | VH              |
| Imidacloprid               | 49   | ✓               | Medium       | Moderate  | H  | L  | VH | H  | H  | M  | VH | L  | L  | L  | L  | VH | M  | H  | +ve |     | H               |
| Clothianidin (high)        | 50   | ✓               | Medium       | Moderate  | H  | VL | —  | VH | M  | M  | H  | H  | M  | M  | M  | VH | H  | VH | +ve |     | H               |
| Dimoteuran (high)          | 75   | ✓               | Medium       | Moderate  | H  | H  | H  | H  | M  | H  | VH | M  | L  | M  | —  | VH | VH | H  | +ve |     | VH              |
| Methomyl                   | 169  | ✓               | Very short   | High      | H  | L  | VH | VH | M  | L  | VH | L  | L  | M  | M  | VH | H  | H  | +ve |     | VH              |
| Thiodicarb                 | 750  | ✓               | Long         | High      | VH | M  | VH | VH | M  | M  | VH | VL | M  | M  | —  | M  | H  | M  | +ve |     | M               |
| Dimethoate                 | 200  | ✓ <sup>17</sup> | Short        | High      | M  | M  | H  | H  | M  | H  | H  | VH | M  | H  | H  | H  | VH | M  | +ve |     | VH              |
| Chlorfenapyr (high)        | 400  | ✓               | Medium       | High      | H  | M  | VH | L  | H  | H  | VH | L  | M  | M  | —  | VH | VH | M  | +ve |     | VH              |
| Skope (high) <sup>22</sup> |      | ✓               | Medium       | High      | VH | H  | VH | VH | H  | —  | —  | —  | VH | H  | L  | VL | VH | VH |     | +ve | H <sup>15</sup> |
| OPs <sup>5</sup>           |      | ✓               | Short-medium | High      | H  | M  | H  | H  | M  | H  | H  | H  | M  | H  | H  | VH | H  | VH | +ve |     | M <sup>15</sup> |
| Carbaryl <sup>3</sup>      |      | ✓               | Short        | High      | H  | —  | —  | —  | H  | —  | —  | —  | —  | —  | —  | —  | —  | —  | —   | —   | H               |
| Pyrethroids <sup>4</sup>   |      | ✓ <sup>7</sup>  | Long         | Very high | VH | —  | —  | —  | VH | —  | —  | —  | VH | VH | VH | VH | VH | VH | +ve |     | H               |

1. Total predatory beetles – ladybeetles, red and blue beetles, other predatory beetles.  
 2. Total predatory bugs – big-eyed bugs, minute pirate bugs, brown smudge bugs, glossy shield bug, damselfield bug, assassin bug, apple dimpling bug.  
 3. Information: Citrus pests and their natural enemies, edited by Dan Smith; University of California Statewide IPM project. Cotton, Selectivity and persistence of key cotton insecticides and miticides.  
 4. Pyrethroids: alpha-cypermethrin, cypermethrin, beta-cyfluthrin, cyfluthrin, bifenthrin, fenvalerate, estenvalerate, deltamethrin, lambda-cyhalothrin.  
 5. Organophosphates: omethoate, monocrotophos, profenofos, chlorpyrifos-methyl, methidathion, parathion-methyl, thiometon.  
 6. *Helicoverpa punctigera* only.  
 7. Bifenthrin is registered for mite and silverleaf whitefly control; alpha-cypermethrin, beta-cyfluthrin, bifenthrin, deltamethrin and lambda-cyhalothrin are registered for control of mirids.  
 8. Persistence of pest control; very short about 24 hours; short, less than 3 days; medium, 3-7 days, long, greater than 10 days.  
 9. Suppression of mites and aphids only.  
 10. Impact rating (% reduction in beneficials following application, based on scores for the major beneficial groups): VL (very low), less than 10%; L (low), 10-20%; M (moderate), 20-40%; H (high), 40-60%; VH (very high), >60%. A '—' indicates no data available for specific local species.  
 11. *Bacillus thuringiensis*  
 12. Pest resurgence is +ve if repeated applications of a particular product are likely to increase the risk of pest outbreaks or resurgence. Similarly sequential applications of products with a high pest resurgence rating will increase the risk of outbreaks or resurgence of the particular pest species.  
 13. Very high impact on minute two-spotted ladybeetle and other ladybeetles for wet spray, moderate impact for dried spray.  
 14. Data Sources: APVMA Public Release Summaries and British Crop Protection Council. 2003. The Pesticide Manual: A World Compendium (Thirteenth Edition). Where LD50 data is not available impacts are based on comments and descriptions. Where LD50 data is available impacts are based on the following scale: very low = LD50 (48h) >100 ug/bee, low = LD50 (48h) <100 ug/bee, moderate = LD50 (48h) <10 ug/bee, high = LD50 (48h) <1 ug/bee, very high = LD50 (48h) <0.1 ug/bee. Refer to the *Protecting pollinators* section in this guide.  
 15. Wet residue of these products is toxic to bees, however, applying the products in the early evening when bees are not foraging will allow spray to dry, reducing risk to bees the following day.  
 16. May reduce survival of ladybeetle larvae – rating of moderate for this group.  
 17. Will not control organophosphate resistant pests (e.g. mites, some cotton aphid (*Aphis gossypii*) populations).  
 18. Rankings for *Eretmocerus* based on data from Jamie Hopkinson in semi-laboratory replicated experiments (Qld DAF) and on ranking for *E. mundus* (P. De Barro, CSIRO, unpublished) and for *E. eremicus* (Koppert B.V., The Netherlands).  
 19. Suppression only.  
 20. Transform is registered for control of greenhouse whitefly at the 96 g ai/ha rate.  
 21. Effects on thrips are for populations found on leaves. This is relevant to seedling crops, where thrips damage leaves, and to mid-late season when thrips adults and larvae help control mites by feeding on them as well as on leaf tissue. Note that flowers are a protected sites, so live adult thrips may be found in flowers even after crops have been treated with products that would control them on leaves.  
 22. Skope is a mixture of acetamiprid and emamectin benzoate. At the low rate (175 ml/ha) this is 38.2 g ai/ha acetamiprid and 5.7 g ai/ha emamectin benzoate. At the high rate (350 ml/ha) this is 76.3 g ai/ha acetamiprid and 11.4 g ai/ha emamectin benzoate. Skope may flare mite populations where abamectin resistant two-spotted mites are present.  
 23. Voliam Flexi is a mixture of thiamethoxam and chlorantraniliprole. At the 250 g/ha rate this is 40 g ai/ha thiamethoxam and 40 g ai/ha chlorantraniliprole.  
**DISCLAIMER:** Information provided is based on the current best information available from research data. Users of these products should check the label for further details of rate, pest spectrum, safe handling and application. Further information on the products can be obtained from the manufacturer.  
**AUTHORS:** Simone Heimoana – CSIRO; Lewis Wilson – formerly CSIRO; Jamie Hopkinson – DAF Qld; Robert Mensah – formerly NSW DPI; Moazzem Khan – formerly DAF Qld; Martin Dillon – formerly CSIRO; Mark Wade – formerly CSIRO; Brad Scholz – formerly DAF Qld; Dave Murray – formerly DAF Qld; Viliami Heimoana – formerly NSW DPI; Richard Lloyd – formerly DAF Qld; Richard Sequeira – DAF Qld; Paul De Barro – CSIRO and Jonathan Holloway – formerly NSW DPI.

# Management of key insect and mite pests

This chapter is presented as a guide to assist growers and advisors in planning their integrated pest management (IPM) programs and provides specific management information for the key insect and mite pests of Australian cotton. Cotton production now covers quite a diverse range of environmental conditions, planting dates and agronomic strategies. Be aware of any regional differences in pest activity or damage potential that may impact your insect management decisions.

## Crop damage

Visible damage indicates that a pest could be influencing crop development and possibly yield potential. In some instances, damage symptoms will be observed without the pest, meaning either the pest is there but cannot be observed or that the pest that caused the damage has since left the crop. In other cases, a pest (or incidental non-pest insect) may be observed with little or no associated crop damage. Use knowledge of the pests/beneficials present in combination with crop damage to make appropriate pest management decisions. Be aware that some pest damage symptoms (such as square shedding/abscission) can easily be confused with damage from physiological/environmental or related causes.

**CottonInfo video: What is causing my squares to shed?** [youtu.be/XMQr2mC25zc](https://youtu.be/XMQr2mC25zc)

## Sampling and thresholds

Sampling is the process of collecting the day-to-day information on pest and beneficial abundance and crop damage used to make pest management decisions, and thresholds provide a rational basis for making those decisions and keeping them consistent. Correctly identifying pest species and knowing the key beneficial predators and parasitoids for each pest is important for developing confidence in IPM approaches to pest management. Thresholds aim to ensure a positive economic return from a pest management action (the benefit gained by the action is more than it costs), and can be based on insect numbers, crop damage, or predicted population levels.

**CottonInfo video on the types of insects that might be found when beat sheeting** [youtu.be/3r4uqD5hp-w](https://youtu.be/3r4uqD5hp-w)



**Knowledge is the key to effective pest management. The shedding of these squares was caused by environmental factors, not pest damage. (Paul Grundy, Qld DAF)**

## Product selection

Selecting an insecticide (or miticide) can be a complex decision based on trade-offs between preventing pest damage and conserving beneficials, or reducing levels of one pest but risking the outbreak of another.

All pests have survival strategies that allow them to live and breed in cotton farming systems. Understanding how pests can survive, including knowing their resistance status and risks, overwintering habit and alternative hosts can help with good decision making for the longer term.

Highly hazardous pesticides (HHP) refers to pesticides that are acknowledged to present particularly high levels of acute or chronic hazards to health or environment, according to internationally accepted classification systems or relevant international agreements or conventions, such as those listed in the Stockholm convention, Montreal protocol and Rotterdam convention. The Better Cotton Initiative (BCI) has banned or is looking to phase out the use of certain pesticides. Phorate and abamectin have been categorised by BCI as HHPs and BCI growers are encouraged to find alternatives where possible. Where an alternative is not available or appropriate (e.g. for resistance management), growers should use a closed transfer system as well as recommended sampling and thresholds as part of a broader IPM strategy.

**Registration of a pesticide is not a recommendation** for the use of a specific pesticide in a particular situation. The pesticide chosen should be the best one for the crop, pest and current circumstances. Carefully study the container label before using any pesticide, and note any specific instructions or restrictions relating to the rate, timing, application and safety. Confirm that the registration or permit is current prior to use.

Ensure your insecticide program fits in with the **Insecticide Resistance Management Strategy** (see pages 54-60 for more information). Insecticides can be a costly part of cotton production. Follow industry thresholds (when available) to prevent unnecessary spraying.

## Important – avoid spray drift

For legal requirements and best practice information on reducing spray drift, refer to the *Spray application* chapter. Insecticides can be highly toxic to humans and the environment. Carefully follow all label directions. ■■■

## KEY INSECT PESTS

|                                       |         |
|---------------------------------------|---------|
| Aphids                                | Page 18 |
| Armyworm ( <i>Spodoptera</i> species) | Page 24 |
| Green vegetable and other stink bugs  | Page 25 |
| <i>Helicoverpa</i>                    | Page 27 |
| Mealybug                              | Page 31 |
| Mirids                                | Page 33 |
| Mites                                 | Page 37 |
| Pale cotton stainers                  | Page 41 |
| Soil and establishment pests          | Page 42 |
| Thrips                                | Page 45 |
| Whitefly                              | Page 47 |
| Other pests                           | Page 52 |

# Hard on mirids Soft on beneficials



## Transform<sup>®</sup> WG Isoclast<sup>®</sup> active

**INSECTICIDE**

Cotton pests can suck the profitability right out of your crop. That's why growers have been turning to Transform<sup>®</sup> WG Isoclast<sup>®</sup> active insecticide as part of a strategic Integrated Pest Management program.

Transform controls a range of cotton pests while proving soft on beneficials. Because of its unique Mode of Action, Transform can control insects that are resistant to other insecticides.

## Aphids

**Cotton aphid** – *Aphis gossypii*

**Cowpea aphid** – *Aphis craccivora*

**Green peach aphid** – *Myzus persicae*

Cotton aphid is the most common aphid pest in cotton. Green peach aphid and cowpea aphid are occasional pests of young cotton but populations of both species decline as temperatures increase (generally early December). While the use of Bollgard 3 and IPM practices have reduced the incidence of aphids in most fields, occasional outbreaks may appear later in the season.

### Damage symptoms

Nymphs and wingless adults of cotton aphid feed on the undersides of leaves, in the terminals, on young stems and on developing fruit. They may cause stunting and downward cupping of the leaves, and in severe cases portions of a damaged leaf's upper surface will turn red. Feeding on terminals and fruit can also cause stunting. Populations of aphids that develop early and increase quickly can inhibit photosynthesis and reduce yield. The sugary 'honeydew' excreted by aphids can promote sooty mould. Cotton aphids have also been shown to transmit the disease cotton bunchy top (CBT; see page 102). Once bolls begin to open, the honeydew can contaminate the lint, which is why sampling at this stage of crop development is still critical. Green peach aphid causes more severe damage to plant growth than cotton aphid.



**Aphids indirectly damage crops by transmitting cotton diseases.**  
(Lewis Wilson)

### Sampling

Aphid populations are sampled visually. Focus on non-winged adults and nymphs. Winged adults may be transitory, while the presence of non-winged adults together with their nymphs indicates that a population has settled in the crop.

### Species identification

As the different aphid species differ in their potential to damage cotton or spread CBT, identification is important. Verify which aphid species is present (particularly early in the season) before implementing any management strategies. Note that the species composition may change during the season. Aphid species can be distinguished by close examination with a hand lens. Green peach aphids are pale green and more oval than cotton aphids, and have W-shaped tubercles (bumps on the head between the antennae), and long siphunculi (tubes between the back legs). Cotton aphid and cowpea aphid don't have tubercles (the head is smooth between the antennae) and the siphunculi are shorter. Adults of cowpea aphid are shiny black and nymphs are always

dusky matt grey, while adults and nymphs of cotton aphid are matt and vary widely from yellow, green, brown to dull black. The siphunculi of adult cotton aphids are always black. If you are unable to make a determination, or suspect more than one species could be present, contact Simone Heimoana, CSIRO, Narrabri, to arrange sending a sample. Contact details are provided at the end of this section.

**CottonInfo video on aphid identification** [youtu.be/Vi07GhSoQtg](https://youtu.be/Vi07GhSoQtg)

### Methods

**Check the population at least weekly:** Begin aphid sampling at seedling emergence and continue until defoliation. Sample for non-winged adults and nymphs on the underside of mainstem leaves 3-4 nodes below the plant terminal. If a high proportion of plants have only the winged form, recheck within a few days to see if they have settled and young are being produced.

**Seedling to first open boll:** Use a 0-5 scoring system based on the number of aphids/leaf (see page 20). If hotspots of cotton aphid are found early season, monitor cotton in these areas for symptoms of CBT.

**First open boll to harvest:** Use a presence/absence scoring system. Check one leaf/plant. Choose a recently expanded leaf, from the 3rd, 4th or 5th node below the terminal. If necessary, mites can be sampled using the same leaf. Sample at least 20 leaves per location. Score a plant as infested if the leaf has 4 or more wingless aphids in any 2 cm<sup>2</sup> area. As wingless females surround themselves with their offspring, the indication of a growing aphid population is the clumping of differently sized aphids in a small area. Aphids are most abundant on the edges of fields so ensure some perimeter sampling occurs. Also assess plants for the presence of honeydew.

### Thresholds

#### Cotton aphid

From the seedling stage until first open boll, thresholds are based on the potential for feeding damage of the aphid population to reduce yield. These thresholds are dynamic, allowing the grower/consultant to consider the value of the crop and the cost of control as part of the decision. After first open boll the thresholds aim to protect the quality of the lint by avoiding contamination of open bolls with honeydew. Since penalties for honeydew contamination are severe, thresholds aim to limit honeydew contamination to trace amounts.

**CottonInfo video on sampling aphids** [youtu.be/L9N64u1yi8E](https://youtu.be/L9N64u1yi8E)

There is also a risk that yield loss can occur through crop infection with CBT, which these thresholds do not take into account. Recent research has shown that the risk of CBT spreading through crops and affecting yield are low unless significant populations of ratoon cotton or alternative weed hosts are nearby or within the field.

In situations where there are many hosts of CBT near the field and a large influx of aphids occurs from these hosts into the cotton crop, the development and spread of aphids should be monitored intensively (at least twice weekly), and any hotspots checked for plants with CBT symptoms. Mark aphid hotspot areas and return to them to check aphid survival. If it is low, then no action may be needed; if populations are healthy, increasing and spreading, control may be required to prevent transmission of CBT within the crop. If control is needed, choose a selective option to conserve beneficials. Removing cotton ratoons/volunteers and weeds in and around fields well before cotton planting will reduce winter survival of aphids and carryover of CBT.

| SEEDLING TO FIRST OPEN BOLL   | FIRST OPEN BOLL TO HARVEST                                      |
|---|---|
| Calculate the Cumulative Season Aphid Score (see page 20), and use to determine an estimate of yield loss | 50% plants infested or 10% if trace amounts of honeydew present |

## Green peach aphid

This species can severely stunt young cotton plants. As it is more damaging than cotton aphid, the threshold for control is lower: 25% of plants infested. Populations can occasionally occur on seedling cotton. However as populations usually decline naturally when temperatures increase, it is unusual for control to be necessary.

## Cowpea aphid

This species can occur on seedling cotton crops, sometimes in quite high numbers. However, populations usually decline quickly as temperatures increase. Control would only be needed if cowpea populations persisted (e.g. cooler temperatures) and plants were showing signs of damage and stunting.

## Key beneficials

**Predators** – ladybird beetle adults and larvae, red and blue beetles, damsel bugs, big-eyed bugs, lacewing larvae, hoverfly and silverfly larvae.

**Parasitoids** – (these cause mummification) *Aphidius colemani*, *Lysiphlebus testaceipes*.



**Spot the parasitoid (mummy) amongst the aphids.**  
(Paul Grundy, Qld DAF)

## Selecting an insecticide

The insecticide products registered for the control of aphids in cotton in Australia are presented on page 22 (check label for target species). If aphid control is required early season, use a selective option to help conserve beneficial populations, in accordance with the IRMS (pages 56-57). These beneficials can assist in controlling any survivors from the insecticide.

## Resistance profile

Aphids reproduce asexually. All the progeny of a resistant individual will be resistant. Once resistance is selected for in a population, it can quickly dominate and give rise to new, entirely resistant populations.

## Resistance profile – cotton aphid

Neonicotinoid resistance was once widespread and is now essentially under control but there remains cross-resistance between acetamiprid, thiamethoxam, clothianidin and imidacloprid.

Resistance is being inadvertently selected in two ways. The first has been through the widespread use of neonicotinoid seed treatments and the second is through the use of foliar-applied products targeting mirids. Even when aphids are present at very low levels, resistance is being selected.

It therefore remains critical to follow the recommendations of the IRMS and rotate insecticide chemistries, taking into account the insecticide group of any seed treatment at-planting insecticide (see Table 3 on page 12).

There is cross-resistance in cotton aphid between pirimicarb and dimethoate, and in the early 2000s this resistance rendered these compounds ineffective. In recent years, resistance to these compounds has declined in southern growing regions but has re-emerged in northern growing regions (i.e. northern WA and north Qld). Re-selection of resistance is a risk, and the IRMS stipulates that omethoate/dimethoate should not be used in rotation with pirimicarb, or vice versa. Neonicotinoid resistance may put pressure on pirimicarb and dimethoate/omethoate and attention should be paid to the effective management of these valuable products.

When choosing an aphicide, consider previous insecticide choices for mirids as well as for aphids and rotate chemical groups. If a phorate side dressing was used then pirimicarb or dimethoate/omethoate should not be used as the first foliar spray since there is cross-resistance between them. Dimethoate/omethoate use will re-select for catastrophic pirimicarb resistance in aphids so do not use pirimicarb and dimethoate/omethoate in the same field.

## Overwintering habit

Aphids don't have an overwintering form, but cool temperatures slow their growth rate dramatically. In cotton growing areas, aphids persist through winter on whatever suitable host plants are available, including cotton volunteers and ratoons, and ornamental garden species.

## Alternative hosts

Cotton aphid has a broad host range, including many common weeds. Winter weed hosts include: marshmallow, capeweed and thistles. Ratoon or volunteer cotton is a host and can act as a green bridge for CBT. Some legume crops such as faba beans are also potential winter hosts. Spring and summer weed hosts include: thornapples, nightshades, paddymelon, bladder ketmia and Bathurst burr. Sunflower crops and volunteers also accommodate cotton aphid.

Cowpea aphid is more abundant in winter and has a broad host range. Populations in winter can be found on burr medic, marshmallow, dwarf amaranth, caustic weed, volunteer cotton and in summer on these hosts as well as hogweed, cathead, volunteer cotton, cobbler's peg, datura, tarvine, small crumbleweed, paddy melon and sowthistle.

Winter weeds that support green peach aphids include turnip weed and marshmallow. Spring germinations of peach vine and thornapples also host green peach aphid. Canola is an attractive host crop through late winter and early spring.

### Further information:

**CSIRO Agriculture and Food, Narrabri**  
**Simone Heimoana: 02 6799 1592 or 0427 992 466.**


**NSW DPI, Tamworth**  
**Lisa Bird: 02 6763 1128.**

**SAMPLING PROTOCOLS FOR COTTON APHID – FOR USE UNTIL FIRST OPEN BOLL**

**STEP 1: Collect and score leaves.**

Collect 20 leaves from each of several locations, taking only one mainstem leaf per plant from 3, 4 or 5 nodes below the terminal (these leaves can also be used for mite scoring). Sample regularly, as yield loss estimates will be most accurate when using the time aphids first arrive in the crop.

Estimate the number of aphids on the leaf and allocate each leaf a score of 0 to 5 (ignore pale brown bloated aphids as these are parasitised). Sum the scores and divide by the number of leaves to calculate the **Average Aphid Score**.



**STEP 2: Calculate the cumulative season aphid score.**

Convert the Average Aphid Score calculated in Step 1 to a **Sample Aphid Score** using the look-up table below to account for the length of time the aphids have been present in the crop. Find the value where the Average Aphid Scores from 'this check' and the 'last check' intersect, and multiply by the number of days that have lapsed between checks to calculate the Sample Aphid Score. If aphids are found in the first assessment of the season, assume the 'Score last check' was '0' and that it occurred 5 days ago.

As the season progresses, add each check's Sample Aphid Score to the previous value to give the **Cumulative Season Aphid Score**. Note that aphids can disappear due to predation by beneficials, changes in the weather or insecticide application. When aphids are sprayed, or if the Average Aphid Scores return to '0' in 2 consecutive checks during the season, reset the Cumulative Season Aphid Score to '0'.

|                          |     | Average score this check |     |     |     |     |     |     |     |     |     |     |
|--------------------------|-----|--------------------------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
|                          |     | 0                        | 0.5 | 1.0 | 1.5 | 2.0 | 2.5 | 3.0 | 3.5 | 4.0 | 4.5 | 5.0 |
| Average score last check | 0   | 0.0                      | 0.3 | 0.5 | 0.8 | 1.0 | 1.3 | 1.5 | 1.8 | 2.0 | 2.3 | 2.5 |
|                          | 0.5 | 0.3                      | 0.5 | 0.8 | 1.0 | 1.3 | 1.5 | 1.8 | 2.0 | 2.3 | 2.5 | 2.8 |
|                          | 1.0 | 0.5                      | 0.8 | 1.0 | 1.3 | 1.5 | 1.8 | 2.0 | 2.3 | 2.5 | 2.8 | 3.0 |
|                          | 1.5 | 0.8                      | 1.0 | 1.3 | 1.5 | 1.8 | 2.0 | 2.3 | 2.5 | 2.8 | 3.0 | 3.3 |
|                          | 2.0 | 1.0                      | 1.3 | 1.5 | 1.8 | 2.0 | 2.3 | 2.5 | 2.8 | 3.0 | 3.3 | 3.5 |
|                          | 2.5 | 1.3                      | 1.5 | 1.8 | 2.0 | 2.3 | 2.5 | 2.8 | 3.0 | 3.3 | 3.5 | 3.8 |
|                          | 3.0 | 1.5                      | 1.8 | 2.0 | 2.3 | 2.5 | 2.8 | 3.0 | 3.3 | 3.5 | 3.8 | 4.0 |
|                          | 3.5 | 1.8                      | 2.0 | 2.3 | 2.5 | 2.8 | 3.0 | 3.3 | 3.5 | 3.8 | 4.0 | 4.3 |
|                          | 4.0 | 2.0                      | 2.3 | 2.5 | 2.8 | 3.0 | 3.3 | 3.5 | 3.8 | 4.0 | 4.3 | 4.5 |
|                          | 4.5 | 2.3                      | 2.5 | 2.8 | 3.0 | 3.3 | 3.5 | 3.8 | 4.0 | 4.3 | 4.5 | 4.8 |
|                          | 5.0 | 2.5                      | 2.8 | 3.0 | 3.3 | 3.5 | 3.8 | 4.0 | 4.3 | 4.5 | 4.8 | 5.0 |

**STEP 3: Estimate the yield loss.**

Use the Cumulative Season Aphid Score to estimate the **percentage yield loss** that aphids have already caused (excluding risks of yield loss from CBT) in the table below. The 'Days remaining' in the season is determined using the first time aphids are found in the crop, based on 165 days from planting to 60% open bolls.

For example, if aphids are first found 9 weeks (63 days) after planting, the days remaining would be ~100. When aphids are sprayed, or if they disappear from the crop then reappear later in the season, reassess the days remaining based on the number of days left in the season from the time of their reappearance.

| Cumulative Season Aphid Score | Days remaining (until 60% open bolls from the time when aphids are first observed)* |    |    |    |    |    |    |    |    |    |  |
|-------------------------------|---|----|----|----|----|----|----|----|----|----|--|
|                               | 100   | 90 | 80 | 70 | 60 | 50 | 40 | 30 | 20 | 10 |  |
| 0                             | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  |  |
| 5                             | 0   | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  | 0  |  |
| 10                            | 2   | 2  | 1  | 1  | 1  | 0  | 0  | 0  | 0  | 0  |  |
| 15                            | 5   | 4  | 3  | 3  | 2  | 1  | 1  | 0  | 0  | 0  |  |
| 20                            | 7   | 6  | 5  | 4  | 3  | 2  | 1  | 1  | 0  | 0  |  |
| 25                            | 9   | 8  | 7  | 6  | 5  | 3  | 2  | 1  | 0  | 0  |  |
| 30                            | 11  | 10 | 8  | 7  | 6  | 5  | 3  | 2  | 1  | 0  |  |
| 40                            | 15  | 13 | 12 | 10 | 8  | 7  | 5  | 3  | 1  | 0  |  |
| 50                            | 19  | 17 | 15 | 13 | 11 | 9  | 7  | 5  | 2  | 0  |  |
| 60                            | 23  | 21 | 18 | 16 | 13 | 11 | 8  | 6  | 3  | 1  |  |
| 80                            | 31  | 28 | 25 | 22 | 18 | 15 | 12 | 8  | 5  | 1  |  |
| 100                           | 38  | 34 | 31 | 27 | 23 | 19 | 15 | 11 | 7  | 2  |  |
| 120                           | 45  | 41 | 37 | 32 | 28 | 23 | 18 | 13 | 9  | 3  |  |

\*Crop sensitivity to yield loss declines as the crop gets older. These estimates take into account factors affecting aphid population development, such as beneficials, weather and variety. Yield reductions >4% are highlighted, however use individual crop value and cost of control to determine how much yield loss can be tolerated before intervention is required.

# Support Food & Fibre Research & Development

**SERO-**<sup>®</sup>  
**PESTICIDE**

  
Allowed Input  
Cert. No. 20021

**Active constituent:** 400 g/L *Clitoria ternatea* Extract  
For the control or suppression of green mirids, silver leaf white fly (biotype b) and heliothis in cotton, and suppression of diamondback moth in brassicas as specified by the Directions for Use table. Also for use in cotton for the reduction in formation of the microsclerotia of *verticillium dahliae* assisting in the management of Verticillium wilt.

Now registered  
as a tool for  
**Verticillium Wilt  
Management**

Contact us for  
more details



Safe to pollinators  
and predators

Unique production  
and manufactured  
in rural Australia



For every hectare  
of Australian Cotton  
treated with Sero-X  
funds are returned  
to the CRDC for  
future innovations.



New class of active compounds  
with innovative pest control and  
virtually no chance of resistance

- Direct Mortality
- Inhibits the development of  
Microsclerotia of *Verticillium dahliae*
- Repellency and anti feedant
- Mating disruption

  
**INNOVATE AG**

**INNOVATE AG & GROWTH AGRICULTURE**  
77a Rose Street, Wee Waa NSW 2388  
[www.innovate-ag.com.au](http://www.innovate-ag.com.au)

**TABLE 5: Control of aphids (refer to label for target species)**

| Active ingredient                 | MoA group | Cotton aphid resistance  | Overall impact on beneficials* | Comments#  |
|-----------------------------------|-----------|--|--------------------------------|--|
| Paraffinic oil                    | No group  | Unknown  | <b>Very low</b>                | Apply by ground rig using a minimum of 80 L/ha of water. If populations exceed 20 aphids per terminal shoot, in a mixture with another aphicide.   |
| Pirimicarb                        | 1A        | Very low in southern growing regions. High in northern WA and north Qld. Cross-resistance to dimethoate/phorate                  | <b>Very low</b>                | Thorough spray coverage essential for best results. Maximum 2 applications per season. Do not use as consecutive applications.   |
| Afidopyropen                      | 9D        | Unknown  | <b>Low</b>                     | Will disrupt insect behaviour and feeding. Provides a slow knockdown. Maximum 4 applications per season.   |
| Diafenthiuron                     | 12A       | Very low   | <b>Low</b>                     | Apply before damage occurs. Lower rate for ground rig application only. Maximum 2 applications per season.   |
| Pymetrozine                       | 9B        | Unknown  | <b>Low</b>                     | Apply prior to row closure on an actively growing crop with a developing population before honeydew damage has occurred. Maximum 2 applications per season.  |
| Cyantraniliprole                  | 28        | Unknown  | <b>Moderate</b>                | Suppression only. Maximum 2 applications per season.   |
| Chlorantraniliprole/ Thiamethoxam | 28/4A     | Cross-resistance between all the neonicotinoids  | <b>Moderate</b>                | Apply in early stages of population development. Maximum 2 applications per season. If a neonicotinoid seed treatment has been used, do not use as first foliar spray.   |
| Spirotetramat                     | 23        | Unknown  | <b>Moderate</b>                | Use the higher rate when periods of high pest pressure or rapid crop growth are evident, when longer residual control is desired or when crops are well advanced. Do not re-apply within 14 days of a previous spray. Maximum 2 applications per season.   |
| Sulfoxaflor                       | 4C        | No resistance  | <b>Moderate</b>                | Use higher rate under heavy aphid infestations and/or when water volume is reduced such as aerial application. Maximum 4 applications per season.  |
| Fonicamid                         | 9C        | Unknown  | <b>Moderate</b>                | Apply to an aphid population in the early stages of development before honeydew is evident or aphid damage occurs. Thorough spray coverage is essential. Maximum 2 applications per season.  |
| Emamectin benzoate/ acetamiprid   | 6/4A      | Cross-resistance between all the neonicotinoids  | <b>Moderate</b>                | Use the high rate under sustained heavy aphid pressure. Maximum 2 applications per season (note maximum of 3 applications in total of Group 6 insecticides). Use organosilicone adjuvant as per label.   |
| Phorate                           | 1B        | Occasional – low. Phorate is cross resistant to pirimicarb and dimethoate  | <b>Moderate</b>                | For short residual control at time of planting. Irrigate as soon as possible after treatment.<br>For extended period of control. Only use the highest rate on heavy soils when conditions favour good emergence, use higher rates (NSW and WA only). Maximum 1 application per season.                     |
| Acetamiprid                       | 4A        | Cross-resistance between all the neonicotinoids  | <b>Moderate</b>                | Ensure good coverage. Apply with 0.2% Incide penetrant. Use high rate under sustained heavy pressure. Do not use as first foliar if neonicotinoid seed treatment used. Maximum 2 applications per season. If repeat applications are required, alternate with products from a different insecticide group. |
| Amitraz                           | 19        | Unknown  | <b>Moderate</b>                | Suppression when used for controlling <i>Helicoverpa</i> ( <i>Helicoverpa</i> rates). Maximum 4 applications per season.   |
| Clothianidin                      | 4A        | Occasional – low. Cross-resistance between all the neonicotinoids  | <b>Moderate</b>                | Apply when aphid numbers are low and beginning to build. Do not use as first foliar if neonicotinoid seed treatment used. Maximum 2 applications per season.   |
| Imidacloprid                      | 4A        | Cross-resistance between all the neonicotinoids  | <b>Moderate</b>                | Add Pulse penetrant at 0.2% v/v (2 mL/L water) or equivalent organosilicon surfactant. Do not use as first foliar if neonicotinoid seed treatment used. Apply early in the establishment of an aphid infestation. Maximum 2 applications per season.   |
| Dimethoate                        | 1B        | Very low in southern growing regions. High in northern WA and north Qld. Dimethoate is cross resistant to pirimicarb and phorate | <b>High</b>                    | Do not use where resistant strains are present. Do not harvest for 14 days after application. Do not graze or cut for stockfeed for 14 days after application. Maximum 2 applications per season.  |
| Chlorpyrifos                      | 1B        | Occasional – low   | <b>High</b>                    | Use higher rates on heavy infestations. Maximum 3 applications per season.   |

#For all control options always refer to the label for instructions.

\*For more details about impact on beneficial insects (including bees), refer to Table 4 in this guide.





up to 1,500 cotton farms

family farms



90% =

(Source: Cotton Australia, 2020)



Producing a bale of irrigated cotton now requires:

1992

2021

LESS WATER

49%

LESS LAND

34%

LESS INCECTICIDES

97%

(Source: Cotton Australia, 2020)

The Australian cotton industry employs more than:




X 1000

(Source: Cotton Australia, 2020)

Water efficiency in agricultural irrigation:

50% —



— 92%

FLOOD IRRIGATION

CENTRE PIVOT IRRIGATION

(Source: USDA, 2001)

FEWER RESOURCES.  
MORE FIBRE.  
THIS IS VALLEY.

WWW.ANZ.VALLEYIRRIGATION.COM

## Armyworm

Cluster caterpillar – *Spodoptera litura*

Lesser armyworm – *S. exigua*

Fall armyworm – *S. frugiperda*

Armyworm species found in cotton have a wide host range. Moths are a mottled grey-brown and lay clusters of eggs that they cover with scales, giving a ‘furry’ appearance. Individual larvae (particularly up to medium size) are often mistaken for *Helicoverpa*.

**Cluster caterpillar** has been a pest of cotton in Northern Australia since the 1950s during pre-commercial crop testing at Kununurra. During commercial cropping in the 1960s and 1970s, cotton growers were challenged with a complex of caterpillar species including *S. litura*, with the combined spraying of these pests leading to resistance in *Helicoverpa armigera*.

Larger larvae have three lengthwise yellow/orange stripes, two bands of distinctive dark half-moon shapes along their back and a row of dark spots along their sides. Primarily leaf feeders, they occasionally attack fruiting structures, particularly open flowers, however damage in Bollgard 3 crops often appears to be superficial, with larvae rarely penetrating the bolls like *Helicoverpa* do in conventional cotton.

*Spodoptera* species have a naturally higher tolerance to Bt endotoxins than *Helicoverpa*, but although there is low and ongoing larval survival, the advent of Bollgard 3 (which is currently only registered in Australia for control of *Helicoverpa*) appears to have reduced the frequency of *S. litura*. However, when crops are subjected to environmental stress (e.g. several weeks of extreme heat with cloudy weather, or plant self-shading in the canopy), significant populations of *S. litura* are able to survive due to poor Bt expression in these plants. Maximising plant health and subsequent Bt endotoxin expression in tropical environments is a key management practice for achieving control of this pest.

Research is planned to understand the feeding behaviour of *S. litura* larvae in Bollgard 3 crops and assess potential damage.

**Lesser armyworm** (also known as beet armyworm) can be present in low numbers in young cotton, occasionally defoliating seedlings, but generally prefers weed hosts. Larvae are green to brown, with a white stripe along each side of the back. Feeding by groups of larvae is often associated with webbing and frass.

**Fall armyworm (FAW)** is a migratory tropical species that was first detected in Australia in early 2020, and has since spread to many crop production districts. Although FAW has the capacity to feed on cotton, it appears to prefer grass species with whorls (such as maize and sorghum), and to date no impact on cotton crops has been reported in Australia. Large larvae have plump, smooth skin, a white inverted Y on the head, and four dark raised dots in a square on the second last segment.

For current registrations and permits applying to armyworm species, search the relevant tab on the APVMA portal ([portal.apvma.gov.au](http://portal.apvma.gov.au))



**Lesser armyworm.** (Melina Miles, Qld DAF)



**Fall armyworm.** (Melina Miles, Qld DAF)



**Cluster caterpillar.** (Paul Grundy, Qld DAF)



**Cluster caterpillar eggs.** (Paul Grundy, Qld DAF)

## Green vegetable bug (and other stink bugs)

*Nezara viridula*

Several species of stink bug can be found in cotton, but the most common and most damaging is the green vegetable bug (GVB; *Nezara viridula*). There are also several species of predatory shield bugs that can be present, so correct identification is essential to making the most appropriate management decisions.

### Damage symptoms

Both adult and nymph green vegetable bugs (GVB) cause dull to black shiny spots on the outer boll, warty growth inside the carpels and brown staining of lint in developing bolls. Severe cases may result in tight lock and yield loss. Damage symptoms cannot be distinguished from those caused by mirids. Bolls aged up to 7 days old are usually shed. Bolls 8 to 24 days can suffer significant damage resulting in incomplete development of one or more sectors (locks), stained lint and reduced yield. Older bolls are not susceptible to damage.

### Sampling

Sample for adults and nymphs and monitor fruit retention at least weekly from the start of squaring, more often if numbers are close to threshold. The crop is most susceptible to damage from flowering through until one open boll/m. Smaller instars are less damaging than older instars so it is important to note the size of nymphs in order to use the thresholds correctly.

GVBs are most visible early to mid morning. Visual sampling and beat sheets are equally effective checking methods from the start of squaring until flowering. However, from flowering onwards, beat sheeting is twice as efficient at detecting GVBs, although it may still underestimate the actual field population, as the first and second instars tend to hide in the bracts and may be difficult to dislodge. Bug distribution is often uneven - increasing the number of samples will give more accurate results.

### Thresholds

| Sampling method                     | Adult GVB thresholds from flowering to harvest |
|-------------------------------------|--|
| Visual                              | 0.5 adults/m                                   |
| Beat sheet                          | 1.0 adult/m                                    |
| Damage to small bolls (14 days old) | 20%  |

### Converting GVB nymphs and other stink bugs to GVB adult equivalents

Other stink bugs that cause damage similar to that of GVB occasionally occur in cotton. However, their damage potential is less than that of GVB so their counts need to be adjusted to GVB adult equivalents:

| GVB nymphs and other stink bugs               | GVB adult equivalent |
|---|----------------------|
| Green stink bug                               | 0.5                  |
| Red banded shield bug                         | 0.33                 |
| Pale cotton stainer bug                       | 0.33                 |
| Brown stink bug                               | 0.25                 |
| Harlequin bug                                 | 0.25                 |
| 1st or 2nd instar GVB nymphs (cluster of 10+) | 1.0                  |
| 3rd instar GVB nymphs                         | 0.5                  |
| 4th or 5th instar GVB nymphs                  | 1.0                  |

### Key beneficials



GVB with 4 *Trichopoda* parasite eggs. (Hugh Brier, Qld DAF)

**Parasitoids** – *Trissolcus* is a wasp that inserts its eggs inside GVB eggs. After hatching, *Trissolcus* larvae remain inside the GVB egg and continue to feed and mature. *Trichopoda* is a fly which parasitises later instar nymphs and adults of GVB, laying its eggs on the outside of the bugs. The hatched larvae bore into the host, dramatically reducing the bug's feeding and egg production and ultimately killing them.



Shield bugs instars can change dramatically in colour and pattern as they develop. From left: 1st, 4th and 5th instar GVB nymphs.

## Selecting an insecticide

Insecticide products registered for the control of GVBs in Australian cotton are presented in Table 6. Avoid mid-season use of dimethoate for GVB control if possible as this compound also selects for organophosphate/carbamate resistance in aphids, particularly if using an OP/carbamate in early season furrow treatments for thrips control.

## Resistance profile

No GVB resistance to insecticides has been detected in Australia.

## Overwintering habit

A high proportion of GVB adults enter a dormant phase (bronze colour) during late autumn and overwinter in a variety of sheltered locations such as under bark, in sheds, and under the leaves of unharvested maize crops. A small proportion will remain green and active, feeding on whatever hosts are available.

## Alternative hosts

In Queensland there are two GVB generations during the warmer part of the year. The preferred weed hosts of the first (spring) generation include turnip weed, wild radish, variegated thistle and early mungbeans. The second generation breeds in late summer and early autumn, with pulse crops (particularly soybeans and mungbeans) and blackberry nightshade (*Solanum nigrum*) as key hosts. GVB populations are usually much lower in mid summer, mainly due to a lack of suitable hosts. In NSW the two generations occur a little later compared to Queensland.

### Further information:

Qld DAF, Paul Grundy: 0427 929 172

## ALERT: EXOTIC BUG THREAT



**Brown marmorated stink bug.**  
(Kristie Graham, USDA ARS, Bugwood.org)

## Brown marmorated stink bug (BMSB)

An exotic pest that could cause damage to crops including cotton, BSMB has been associated with cargo imports from shipping containers and machinery to household goods (including shoes and barbie dolls).

### Monitor and report

If you spot any unusual stink bugs, Catch (if possible) and call the Exotic Plant Pest Hotline: 1800 084 881.

CottonInfo insect ID guide on brown marmorated stink bug [www.cottoninfo.com.au/publications/insect-id-guide-endemics-exotics-brown-marmorated-stink-bug](http://www.cottoninfo.com.au/publications/insect-id-guide-endemics-exotics-brown-marmorated-stink-bug)

**TABLE 6: Control of green vegetable bug**

| Active ingredient              | MoA group | Overall impact on beneficials* | Comments#   |
|--------------------------------|-----------|--------------------------------|---|
| Fipronil                       | 2B        | Moderate                       | Apply when pests appear. Use higher rate when higher infestations are present. Avoid repeated use of this insecticide group.  |
| Clothianidin                   | 4A        | Moderate                       | Use higher rate when heavy infestations are expected and longer control is required. Treated insects may still be on plant 2 or 3 days after application but will have stopped feeding. Maximum 2 applications per season.                                      |
| Emamectin benzoate/Acetamiprid | 6/4A      | Moderate                       | Use higher rate on heavier populations and for a faster knockdown. Use non-ionic surfactant or organosilicone adjuvant as per label. Maximum 2 applications per season.   |
| Dinotefuran                    | 4A        | Moderate                       | Apply when thresholds are reached. Ensure thorough spray coverage of the crops. Performance can be reduced in stressed crops, or when senescing late season or when pests are not actively feeding in the upper crop canopy. Maximum 2 applications per season. |
| Dimethoate                     | 1B        | High                           | Apply when pests appear. Use higher rates for heavier infestations. Do not harvest for 14 days after application. Do not graze or cut for stockfeed for 14 days after application. Maximum 2 applications per season.   |

#For all control options ALWAYS refer to the label for instructions.

\*For more details about impact on beneficial insects (including bees), refer to Table 4 in this guide.

## Helicoverpa

*Helicoverpa armigera* (cotton bollworm)

*H. punctigera* (native budworm)

*Helicoverpa* larvae are major pests of conventional cotton, capable of dramatically reducing yield. Widespread industry adoption of Bt cotton has dramatically reduced the need to control these larvae with insecticides, however even Bollgard 3 crops may have *Helicoverpa* present, and the risk of resistance requires careful management.

### Damage symptoms

Larvae attack all stages of plant growth. In conventional cotton (non-Bt varieties), larval feeding can result in seedlings being tipped out, chewing damage to squares and small bolls causing them to shed, and chewed holes in maturing bolls, preventing normal development and encouraging boll rot. In any year an average of 15% of Bt cotton area may carry *Helicoverpa* larvae at or above the recommended threshold levels at some point during peak to late flower. In Bt cotton, chewing damage is mostly confined to fruit and may lead to yield loss.

### Sampling

Visually sample the whole plant for *Helicoverpa* from seedling emergence in both conventional and Bt cotton crops. Check at least 2 times/week until the crop has 30-40% open bolls.

Sample at least 30 plants or 3 separate metres of row for every 50 ha of crop. Fields may not be uniform (e.g. lush areas may occur in head ditches that are more attractive to insects), and variability within the field may determine the minimum number of sampling points required. Larger samples will give more accurate estimates.

Note the individual egg and larval growth stages of the pest. The growth stages are defined as:

|                   |    |                    |
|-------------------|----|--------------------|
| White egg         | WE | pearly white       |
| Brown egg         | BE | off-white to brown |
| Very small larvae | VS | 0-3 mm             |
| Small larvae      | S  | 3-7 mm             |
| Medium larvae     | M  | 7-20 mm            |
| Large larvae      | L  | >20 mm             |

Eggs are laid on plant terminals, leaves, stems and the bracts of fruit. Larvae may be found on terminals, the upper or lower surface of leaves, inside squares, flowers and bolls and along stems. Note the species where possible.



***H. armigera* larvae (left) have pale hairs compared to darker hairs on *H. punctigera* larvae (right).** (Hugh Brier, Qld DAF)  
**More detailed information on identifying *Helicoverpa* is available at [thebeatsheet.com.au/key-pests/helicoverpa-identification-and-behaviour](http://thebeatsheet.com.au/key-pests/helicoverpa-identification-and-behaviour)**

Once squaring begins, note fruit retention or fruiting factors, to gauge what level of damage is being caused to the crop.

Also sample key beneficials to allow the beneficial to pest ratio to be applied (see page 28). Collect brown eggs to check for parasitism by *Trichogramma* spp. (the eggs will turn black).



**Normal eggs (left) and parasitised eggs (right).** (Photo: B. Scholtz)

### Thresholds

*Helicoverpa* thresholds are based on the assumption of potentially mixed populations of *H. armigera* and *H. punctigera*.

#### Conventional cotton

Larval thresholds are preferable as not all eggs hatch. Early in the season eggs are particularly prone to desiccation and being washed or blown from the small plants. Natural mortality can be as high as 80% early season, 75% mid season and 60% late season.

Parasitism and predation can dramatically reduce egg survival (*Trichogramma* parasitoids have the potential to reduce egg survival by over 90%), and larval thresholds are also affected by beneficial insects. It is important to assess beneficial insect numbers when making pest control decisions. Also consider fruit retention when determining whether *Helicoverpa* have caused, or are at risk of causing economic damage.

| SEEDLING TO FLOWERING  | FLOWERING TO CUT-OUT   |
|--|--|
| 2 larvae/m<br>or<br>1 larvae >8 mm/m                         | 2 larvae/m<br>or<br>1 larvae >8 mm/m<br>or<br>5 brown eggs/m |
| CUT-OUT TO 15% OPEN BOLLS                                    | 15% TO 40% OPEN BOLLS  |
| 3 larvae/m<br>or<br>1 larvae >8 mm/m<br>or<br>5 brown eggs/m | 5 larvae/m<br>or<br>2 larvae >8 mm/m<br>or<br>5 brown eggs/m |

#### Bt cotton

Objectively assess larval size and exclude any very small larvae (<3 mm). There are no egg thresholds in Bt cotton, as research has shown no effect on yield from high egg lays (over 100/m).

| SEEDLING TO 40% OPEN BOLLS   |
|--|
| 2 larvae >3 mm/m in 2 consecutive checks<br>or<br>1 larvae >8 mm/m |

Where larvae between 3 and 8 mm are observed on Bt cotton, consecutive checks are essential for decision making. *Helicoverpa* must feed in order to ingest the Bt toxin. If the number of 3-8 mm larvae are above threshold on a given check, chances are that a large portion of these will ingest a sufficient dose of the toxin and die before the next check.

## Using the beneficial to pest ratio

The beneficial to pest ratio can be applied in conventional and Bt cotton. The ratio is calculated as:

$$\frac{\text{Total beneficials}^*}{\text{Helicoverpa (eggs - (\% parasitised) + VS + S larvae)}}$$

At least 30 plants or 3 to 4 separate metres of row by visual sampling or 20 metres of row by suction sampling is needed in order to use the ratio. The total number of beneficials **must only** include the key beneficial insects (marked with an asterisk) and **at least** 3 of the key beneficial species need to be present.

When the beneficial to pest ratio is 0.5 or higher, the *Helicoverpa* population should remain below the threshold of 2 larvae/m.

The beneficial to pest ratio incorporates parasitoids, particularly *Trichogramma*, in the calculation. The level of egg parasitism should be deducted from the number of *Helicoverpa* eggs before the beneficial to pest ratio is calculated. Levels of egg parasitism can vary greatly from farm to farm, region to region and from season to season. Generally levels decline as the season progresses. To assess parasitism, keep collected brown eggs at 25°C until they hatch (healthy) or turn black (parasitised).

### Key beneficials

**Predators of eggs** – red and blue beetle\*, damsel bug\*, green lacewing larvae\*, brown lacewing\*, ants, yellow night stalker spiders, ladybird beetles and apple dimpling bugs.

**Predators of larvae** – glossy, brown\* and predatory shield bugs, big-eyed bug\*, damsel bug\*, assassin bug\*, red and blue beetle\*, brown lacewing\*, common brown earwig, lynx, tangleweb and jumping spiders.

**Predators of pupae** – common brown earwig.

**Predators of moths** – orb-weaver spiders and bats.

**Parasitoids of eggs** – *Trichogramma* spp., *Telenomus* spp.

**Parasitoids of larvae** – *Microplitis demolitor*, orange caterpillar parasite, two-toned caterpillar parasite.

**Predators of late instar larvae** – wolf spiders and carabids.

**Parasitoids of pupae** – banded caterpillar parasite.

\*Species for use in the beneficial to pest ratio formula.

### Selecting an insecticide

Insecticides registered for the control of *Helicoverpa* in cotton are presented in Table 7 on page 30. The use of more selective insecticide options will help to conserve beneficial insects (see Table 4).

Be aware of resistance status in *H. armigera* and follow the insecticide resistance management strategy (IRMS) (pages 56-57).

### Resistance profile

#### *H. armigera* – conventional insecticides

With the introduction of Bt cultivars, *H. armigera* is no longer front of mind as a major pest of concern. However maintaining effective control is critical for conventional cotton and other crops such as pulses. Importantly, large plantings of Bt cotton does not change the overall frequencies of insecticide resistance genes in the *Helicoverpa* population and is unlikely to influence the rate at which *H. armigera* will develop resistance to conventional insecticides if significant selection pressure is imposed.

Results from an industry-wide resistance surveillance program, implemented by NSW DPI and supported by CRDC and GRDC, show levels of resistance to indoxacarb in *H. armigera* are 5.7%. Importantly, populations

of *H. armigera* in Central and Northern Qld had levels of indoxacarb resistance 2.4-fold higher than Southern Qld and NSW during 2018-19, although resistance in most regions declined during the 2019-20 and 2020-21 seasons. Research suggests that indoxacarb resistant *H. armigera* are less likely to survive diapause which may partly explain why resistance has increased in regions where overwintering does not occur to any large extent.

Chlorantraniliprole resistance is stable (>0.5%) and there has been no recent detection of emamectin benzoate resistance. The industry has ceased monitoring broad-spectrum chemistry for resistance. However, field failures are likely for synthetic pyrethroids and carbamates if used to target *H. armigera*.

The use of conventional insecticides for control of *H. armigera* in conventional and Bt cotton crops should be based on relevant thresholds and the IRMS. As well as the cotton IRMS, it is important to adhere to the resistance management strategy (RMS) for the grains industry. [www.ipmguidelinesforgrains.com.au/ipm-information/resistance-management-strategies/#heli](http://www.ipmguidelinesforgrains.com.au/ipm-information/resistance-management-strategies/#heli)

Adoption of resistance management tactics in grains will be critical for maintaining efficacy of indoxacarb, particularly in Central and Northern Qld where resistance risk is highest.

Pupae busting should be a priority post-harvest operation on all cotton farms. The IRMS recommends pupae busting as soon as possible after harvest. For Bt cotton crops, follow the pupae busting guidelines in the Bollgard Resistance Management Plan (RMP).

#### *H. armigera* – Bt cotton

Several genes are currently present in field populations of *H. armigera* that confer high-level resistance to the discrete proteins in Bollgard cotton, however none confer cross-resistance to the other proteins. Individuals must carry two copies of these resistance genes to survive the protein or Bollgard 3, and the vast majority only carry one copy. However, the continued efficacy of Bt cotton relies on how the industry manages its refuges and implements the other elements of the RMP. For further details, including information about recent changes in the frequency of Bt resistance genes in *H. armigera*, refer to the Bollgard 3 RMP for cotton on pages 61-70.

#### *H. punctigera* survival strategies

##### Conventional cotton

Resistance to insecticides has only rarely been detected in *H. punctigera*. In conventional cotton, the tendency for *H. punctigera* to occur in mixed populations with *H. armigera* often limits insecticide control options to those that are also efficacious on *H. armigera*.

##### Bt cotton

Several genes are currently present in field populations of *H. punctigera* that confer resistance to the discrete proteins in Bollgard cotton, however none confer cross-resistance to the other proteins. Individuals must carry two copies of these resistance genes to survive the protein or Bollgard 3, and the vast majority only carry one copy. However, the continued efficacy of Bt cotton relies on how the industry manages its refuges and implements the other elements of the RMP. For further details refer to the Bollgard 3 RMP for cotton.

## Overwintering habit

In colder regions, *H. armigera* over-winters in cotton fields as diapausing pupae. These pupae are the major carriers of resistance from one season to the next. Diapause initiation is caused by falling temperatures and shortening day lengths. The proportion of pupae entering diapause increases from 0% in late February to over 90% in late April – early May, depending on the region. Diapause termination is based on rising soil temperatures beginning in mid to late September in most regions. Emergence from diapause usually occurs over a 6 to 8 week period.

Though *H. punctigera* has always had the capacity to over-winter as diapausing pupae, extensive research conducted in the early 1990s found this species was rarely found late season and over-wintered pupae were rarely found in the ground in the temperate cotton growing areas.

Spring migrations from western inland have largely been thought to be the main source of this species in the eastern cropping regions, and *H. punctigera* have historically never shown any trends towards insecticide resistance in the past. However, in recent times this species has been persisting throughout the whole cotton-growing season, with significant overwintering now taking place.

Migration flights also appear to have lessened, reducing the natural genetic dilution factor. Research continues to investigate whether *H. punctigera* is more likely to develop resistance in the future.

## Alternative hosts

Spring host crops of *H. armigera* include: faba beans, chickpeas, safflower, linseed and canola. Pastures and weed flushes also sustain emerging spring populations. Summer host crops include: soybeans, mungbeans, pigeon pea, sunflower, sorghum and maize. *H. armigera* will attack flowering crops of sorghum and maize preferentially over most other crop hosts.

*H. punctigera* moths are able to utilise a vast selection of host plants (mostly broad-leaved) and do not appear to be as closely associated with crop hosts as *H. armigera*. Though favourable weather and non-crop hosts such as daisies appear to be critical for the early successful survival of *H. punctigera*, spring crops such as chickpea, canola, faba beans and linseed can be heavily infested by this species. Summer crop hosts can include pigeon pea, sunflower and various legume crops (mungbean, soybean etc).

### Further information:

**CSIRO Entomology, Narrabri**  
**Sharon Downes: 02 6799 1576 or 0427 480 967**

**Qld DAF, Paul Grundy: 0427 929 172**

**NSW DPI, Tamworth**  
**Lisa Bird: 02 6763 1128.**



**Helicoverpa larva damaging a maturing boll. (Paul Grundy, Qld DAF)**

**TABLE 7: Control of *Helicoverpa***

| MoA                                       | MoA group  | Resistance ( <i>H. armigera</i> )                                   | Overall impact on beneficials* | Comments#   |
|---|------------|---|--------------------------------|---|
| <i>Bacillus thuringiensis</i>             | No group   | None detected   | Very low                       | Restrictions apply – refer to Bt cotton resistance management plan.   |
| <i>Helicoverpa</i> NPV                    | No group   | None detected   | Very low                       | Use alone or with compatible larvicide. Target application to coincide with egg hatching.   |
| Paraffinic oil                            | No group   | None detected   | Very low                       | Use a minimum of 80 L/ha of water. Apply only by ground rig before crop closure.  |
| Magnet                                    | Attractant | None detected   | Very low                       | Use with insecticides as per label instructions.  |
| Indoxacarb                                | Group 22A  | Widespread – moderate   | Low                            | Maximum 3 applications per season.  |
| <i>Clitoria ternatea</i> extract (Sero-x) | No group   | Unknown   | Low                            | Ensure good coverage. Treatment effects may not be seen for 3 or more days. Applications should be timed to coincide with egg hatch and when small larvae are present. Maximum 5 applications per season.   |
| Amorphous silica                          | No group   | Unknown   | Low                            | Apply during egg lay to egg hatch. Best results are obtained from two sequential applications 6-7 days apart.   |
| Chlorantraniliprole                       | 28         | Occasional – low  | Low                            | Target brown eggs or hatching to 2nd instar larvae before they become entrenched in squares, flowers and bolls. Use high rate where the potential is for >2 larvae/m and to achieve longer residual control. Maximum 3 applications per season.   |
| Spinetoram                                | 5          | Occasional – low  | Low                            | Use the higher rate for heavy infestations. Larvae >8 mm or feeding within bolls & squares may not be controlled. Maximum 2 applications per season.  |
| Cyantraniliprole                          | 28         | Occasional – low  | Moderate                       | Target eggs and hatchlings to 2nd instar larvae before they become entrenched in hidden feeding sites. Maximum 2 applications per season.   |
| Chlorantraniliprole/<br>thiamethoxam      | 28/4A      | Occasional – low<br>Cross-resistance between all the neonicotinoids | Moderate                       | Apply at brown egg to 2nd instar larvae before becoming entrenched in squares, flowers and bolls. Use high rate where the potential is for >3.5 larvae/m and to achieve longer residual control. Maximum 2 applications per season.   |
| Abamectin                                 | 6          | Occasional – low  | Moderate                       | Use the higher rate alone or the lower rate with a suitable mixing partner. Some labels indicate control of <i>H. punctigera</i> only. Maximum 2 applications per season.   |
| Emamectin benzoate                        | 6          | Occasional – low  | Moderate                       | Apply at or just prior to hatching. Use non-ionic surfactant as per label. Maximum 4 applications per season.   |
| Emamectin benzoate/<br>acetamiprid        | 6/4A       | Occasional – low<br>Cross-resistance between all the neonicotinoids | Moderate                       | Apply at or just prior to hatching. Use non-ionic surfactant as per label. Maximum 2 applications per season.   |
| Amitraz                                   | 19         | Unknown   | Moderate                       | Apply as an ovicide with larvicide when eggs or very small larvae are detected. Repeat treatments at 5 and 7 day intervals, when necessary. May suppress mites. Maximum 4 applications per season.  |
| Methomyl                                  | 1A         | Widespread – moderate   | High                           | Higher rate of larvicidal rate may cause reddening of foliage. Do not use more often than every 14 days, if excessive use an alternative. Do not apply during periods of plant stress. Maximum 2 applications per season.   |
| Thiodicarb                                | 1A         | Widespread – moderate   | High                           | This product has ovicidal and larvicidal activity. See label for details. Lower rate is on light to moderate infestations and the higher rate on heavier infestations. Maximum 2 applications per season.   |
| Alpha-cypermethrin                        | 3A         | Widespread – high   | Very high                      | Use low rate for eggs or newly hatched larvae. Use higher rates for higher egg pressure or larger larvae. Maximum 1 application per season.   |
| Bifenthrin                                | 3A         | Widespread – moderate<br>Cross-resistance                           | Very high                      | Time spray to coincide with egg hatch. DO NOT apply to larvae >5 mm. Use higher rate when pest pressure is high, conditions favour pest development and when increased residual protection is required. <i>H. armigera</i> resistance to bifenthrin has increased. Field failures are likely. Maximum 1 application per season. |
| Cypermethrin                              | 3A         | Widespread – high   | Very high                      | See label for specific concentrations and higher rate situations. Maximum 1 application per season.   |
| Deltamethrin                              | 3A         | Widespread – high   | Very high                      | Use low rate as ovicide and high rates for small to medium larvae. Maximum 1 application per season.  |
| Esfenvalerate                             | 3A         | Widespread – high   | Very high                      | Use low rate when larvae are small and pressure is low. Maximum 1 application per season.   |
| Gamma-cyhalothrin                         | Group 3A   | Widespread – high   | Very high                      | Use low rate as ovicide and high rate when egg lay is heavy and/or <i>H. punctigera</i> >10 mm and/or <i>H. armigera</i> <5 mm. Maximum 1 application per season.   |
| Lambda-cyhalothrin                        | Group 3A   | Widespread – high   | Very high                      | Use low rate as ovicide and/or for newly hatched larvae. Maximum 1 application per season.  |

#For all control options always refer to the label for instructions.

\*For more details about impact on beneficial insects (including bees), refer to Table 4 in this guide.



## Mealybug

### *Phenacoccus solenopsis*

The solenopsis mealybug occurs in nearly all regions where cotton is grown in Australia.

### Damage symptoms

Mealybug can affect plant growth at all stages of crop development. When infested during early development, plants exhibit distorted terminal growth, crinkled and bunched leaves, and in severe cases, plant death. On older plants, mealybugs can cause shedding of leaves, squares and small bolls, boll deformity and premature crop senescence. Heavy infestations (>500 mealybug in the top 8 nodes at cut-out) will result in around 80% reduction in harvestable bolls. Honeydew excreted onto the leaves is high in melezitose sugar, which is very sticky and can promote the development of black sooty mould and render seed cotton un-ginnable.

Infestation with mealybug up to the early boll setting stage has the potential to be highly damaging. Population densities of around 25, 110 and 150 mealybugs per plant in the seedling, squaring and early boll stages, respectively, appear to be sufficient to cause significant economic damage.

### Sources and sampling

At low densities, mealybug can be present anywhere on the plant. However, they are likely to aggregate on the underside of leaves and inside bracts of squares or bolls within the top 10 leaf nodes. Hence, checking in the upper half of the crop canopy is an effective sampling strategy.

Ratoon cotton, broadleaf weed species and volunteer cotton can be sources of mealybug within the crop. Ratoon cotton stubs are a major source of infection in following cotton crops. Uncontrolled volunteer cotton growing between seasons can be a significant infestation source. Mealybug populations from volunteer cotton and weeds are easily dispersed onto nearby cotton. Checking volunteer and adjacent cotton will help to detect early infestations in the field. Mealybugs often appear to be more prevalent in crops that are stressed from lack of moisture, waterlogging and/or nitrogen deficiency, so it is important to include stressed areas (e.g. tail drains) when checking. Investigate patches of stunted or dead plants.

#### CottonInfo videos:

Solenopsis mealybug in cotton [youtu.be/vQTJ2XkAsug](https://youtu.be/vQTJ2XkAsug)

Mealybug hotspots [youtu.be/r4aeGPcVHk0](https://youtu.be/r4aeGPcVHk0)

What to do if you find a hotspot [youtu.be/pR7U\\_1ng71E](https://youtu.be/pR7U_1ng71E)

### Management strategy

Research done to date has shown that solenopsis mealybug is best controlled through good crop destruction, farm hygiene and the conservation of naturally-occurring beneficial insects. Minimising impacts of pesticides on beneficial insects is critical for the cost effective control of this pest.



**Solenopsis mealybug adults and crawlers.** (Tonia Grundy, Qld DAF)



**Mealybug predators: Cryptolaemus ladybird beetle larva (left) and lacewing larvae (right), can look very similar to mealybugs.** (Zara Hall and Paul Grundy, Qld DAF)

### What to do if you find them...

- **Correct identification.** While it is likely that observed mealybug are solenopsis mealybug, other similar looking species occasionally occur in cotton, but they are rarely damaging.
- **Mark the plants they are on,** as they can be difficult to relocate when in low numbers. Re-checking marked plants will enable you to judge the efficacy of any natural enemies that are present.
- **Drones can be useful in identifying hotspots.**
- **Look for the presence of natural enemies.** Evidence of predation and parasitism and whether the solenopsis colony is growing or dwindling in size will provide a picture of the effectiveness of natural enemies in your field.
- **Come Clean. Go Clean. Be mindful of spreading mealybug with passage of people and machinery.**

**TABLE 8: Control of solenopsis mealybug**

| Active ingredient | MoA group | Overall impact on beneficials* | Comments#  |
|-------------------|-----------|--------------------------------|--|
| Buprofezin        | 16        | Low                            | Aim application at early nymph stages. Thorough spray coverage is essential. Maximum of 2 applications per crop.   |
| Spirotetramat     | 23        | Moderate                       | Do not re-apply within 14 days of a previous Movento spray. Do not apply more than 2 applications per crop. The re-application of spirotetramat presents a risk for SLW resistance management. |
| Sulfoxaflor       | 4C        | Moderate                       | Thorough spray coverage is essential. Addition of an adjuvant may improve control.   |

#For all control options ALWAYS refer to the label for instructions.

\*For more details about impact on beneficial insects (including bees), refer to Table 4 in this guide.

Upon first detection, mark infested plants/spots. Monitor regularly for mealybug and key beneficials (lacewings, ladybird beetles and parasitoids) to provide information for ongoing management.

Adhere strictly to good farm hygiene principles and ensure everyone entering the farm practices Come Clean. Go Clean. Brush down clothing between fields and ensure farm equipment is cleaned down after entering fields with mealybugs.

Beneficial insects are highly effective in keeping mealybug populations in check. Chemical insecticides should be used for mealybug control only in very specific circumstances. Note that no insecticide option provides 100% control, and mealybug management over the life of the crop is dependent on beneficials. Good coverage is critical for effective control of mealybug with insecticides. Depending on plant size, higher water volumes will increase efficacy. Sequential application 10-14 days apart may be required to achieve a satisfactory level of mealybug control with some insecticides. Refer to the product label for directions. For management advice in specific situations or circumstances, contact the CottonInfo team.

### Management options until early flowering

Mealybug infestation carries a significant risk of plant death during early development as the cotton plant is highly susceptible and seasonal build-up of beneficial insects has not yet occurred. If there is a growing risk of mealybug infestation/damage, and beneficials are not adequately building up, chemical intervention may be required.

Recommended insecticidal options for specifically targeting mealybug include sulfoxaflor, spirotetramat and buprofezin. Product selection should consider the presence of other pests:

- If mirids are at or nearing threshold in a crop with mealybug, use sulfoxaflor for mirid control as it is unlikely to flare mealybug.
- If whitefly are present and at risk of being flared: Spirotetramat (+ 3-5% oil v/v) is effective on mealybug and has efficacy on SLW. Buprofezin has suppressive effects on mealybug and can suppress SLW.

### Management options from late flowering to defoliation

Mealybug hotspots in mid-late season cotton may be indicative of failure to detect and manage infestations early, and/or disruption of beneficials. In the absence of other pests (e.g. mirids, SLW), insecticidal control of mealybug is generally not recommended. Allow beneficials to build and control mealybug.

Aim to control mirids and/or SLW without flaring mealybug by following the product choices recommended above. AVOID USE OF BROAD-SPECTRUM PRODUCTS.

### End-of-season recommendations for a mealybug infested crop

- Prior to crop harvest, destroy all non-crop vegetation that may harbour mealybug (head, tail drains, sides) to minimise mealybug survival and carry-over to next crop.
- At harvest, pick infested fields last. If possible lift picker heads above mealybug hotspots or infested sections.
- After harvest, destroy crop residue thoroughly and root cut to a depth of at least 10 cm to minimise the survival of plant material that could harbour mealybug.
- Consider pupae busting if you are planning on growing cotton again in the same or neighbouring field within 1-2 years.

### Key beneficials

**Predators** – Lacewings, three banded ladybird beetles, white collared ladybird beetles, transverse ladybird beetles, red and blue beetles, cryptolaemus, smudge bugs and earwigs. Lacewings and three banded ladybird beetles are the most common and effective predators.

**Parasitic wasps** – *Aenaisus bambawalei*, a parasitoid of solenopsis mealybug that attacks adult females is effective in suppressing populations.



**Three banded ladybirds are very effective mealybug predators.** (Paul Grundy, Qld DAF)

### Key features that influence survival and pest status

Mealybug have a high reproductive rate and all hatched stages can cause damage. One female can produce hundreds of offspring; eggs hatch within an hour and take about two weeks to develop into adults. Mealybug shelter in protected positions on the cotton plant; in squares, bracts and under surfaces of leaves and their waxy coating is water repellent, making insecticide contact difficult. They can be spread in the field by wind, surface water runoff, rain splash, birds, people and farm equipment. Mealybugs disperse most readily as first instar 'crawlers', but all nymph and adult stages can move between plants by crawling.

Adults and large nymphs can survive for long periods without a host. The crawler stage can live for up to 6 days, and the third instar stage for up to 50 days without food or water.

Overwintering mealybugs can be found in the root zone of weed hosts; loose soil and ant nests on the ground can also provide them with shelter during winter. Once the weather begins to warm, breeding and dispersal begins.

### Alternative hosts

The solenopsis mealybug has an extremely wide host range and has been recorded on pigweed, sowthistle, bladder ketmia, native rosella, vines (cow, bell and potato), crownbeard, stagger weed, marshmallow, verbena, raspweed, and volunteer cotton.

**Further information:** Qld DAF, Richard Sequeira 0407 059 066, Paul Grundy 0427 929 172.



**The pupal stage (left) and adult wasp stage (right) of *Aenaisus bambawalei* which is a parasitoid of the solenopsis mealybug.** (Zara Hall, Qld DAF)

## Mirids

**Green mirid** – *Creontiades dilutus*

**Brown mirid** – *Creontiades pacificus*

**Apple dimpling bug (yellow mirid)** –  
*Campylomma liebknechti*

Green and brown mirids look similar, but brown mirids are slightly larger and many have darker pigmentation. Both species cause similar damage at the boll stage, however green mirids do more damage during squaring. Green mirids are usually more abundant in cotton and they move into crops earlier than brown mirids. The apple dimpling bug (ADB) is about 1/3 the size of other mirids, and can damage small squares (4 ADBs do equivalent damage to 1 adult green mirid). ADB is also a predator of mites, silverleaf whitefly eggs and *Helicoverpa* eggs.


### Damage symptoms

Mirid adults and nymphs cause blackening and death of terminals of young plants, and mid-season damage to squares and small bolls including blackening of pinhead squares and square shedding. Older (4th and 5th) instars can cause as much damage as adults.

Square loss depends upon where the mirids are feeding and size of the squares. Feeding on small or medium sized squares is very likely to directly damage the developing ovules and anthers, resulting in shedding. Feeding on large squares may not result in shedding but still damage the developing ovules, resulting in poor fertilisation and seed development in several locks. The resulting misshapen bolls are commonly referred to as being 'parrot beaked', a deformity that can be also be caused by high temperatures. The relationship between mirid numbers and square loss



**Cut bolls open to determine if black spots correspond to internal lock damage.** (Paul Grundy, Qld DAF)



**Green mirid adult (top) and nymph (bottom).** (C. Mares & M. Khan)

is not always the same due to the different responses from various sized squares, as well as changes in mirid feeding, likely due to a range of factors including temperature and the presence of natural enemies. For this reason both retention and mirid numbers should be considered when making a control decision.

Bolls that are damaged during the first 7-10 days of development may be shed. Bolls damaged between 10-20 days old will be retained but may not develop normally resulting in one or more stunted, brown locks. Apart from reduced weight, damaged bolls may not open sufficiently to enable efficient spindle picking. Black, shiny spots on the outside of bolls can indicate feeding sites for a number of species (mirids, green vegetable bugs and pale cotton stainers), however environmental conditions can also give rise to similar looking marks. Cut bolls open with a sharp knife and inspect the underlying tissue to confirm whether the boll marks are the result of feeding damage. Warty growths can often be found beneath the spots or there might be light brown discolouration of developing lint.

Once bolls exceed 20 days of development, susceptibility of the developing seed and lint to feeding damage becomes less as fibre elongation ceases and seed is located deeper within the boll relative to the boll wall.

**CottonInfo video on mirid identification and damage** [youtu.be/FdCIVONjh\\_Y](https://youtu.be/FdCIVONjh_Y)

### Sampling

Mirids are a very mobile pest and are easily disturbed so care must be taken during sampling, otherwise numbers may be underestimated (discussed later in this section). It is important to distinguish between nymphs and adults. The presence of nymphs can indicate that a mirid population has become entrenched.

Once squaring commences, regularly assess fruit retention. Monitor for all types of plant damage potentially caused by feeding such as tip damage (early-season) and boll damage (mid-season).

## Frequency

Sampling at least twice a week is important in order to identify sudden changes in abundance that may indicate rapid influxes of mirid adults. The greatest risk from mirids is through the period of peak fruit production, from first flower until first open boll.

## Methods and sample size

Sample throughout the field to gain a reliable estimate of overall density. Visual inspection of whole plants, using a beat sheet or sweep netting all give comparable estimates of mirid abundance when plants are small. Once the crop reaches 9-10 nodes the efficiency of visual whole plant sampling declines because the plants are too big to sample quickly and effectively without disturbing the mirids. From 10 nodes onwards, a beat sheet is the recommended option.

When beat sheeting, place the sheet against the base of a row of plants and drape across the furrow and up over the adjacent row. Vigorously shake a section of row with a 1 m stick onto the area of beat sheet lying across the furrow. Quickly count the number of mirids dislodged onto the beat sheet, looking carefully for first and second instar nymphs, which are light green in colour and can be difficult to see against the yellow sheet. Mirids are faster moving and have longer antennae than most other small green insects.

**CottonInfo video on using a beatsheet** [youtu.be/vl-Y8qj\\_xXs](https://youtu.be/vl-Y8qj_xXs)

Accurate estimation of mirid numbers is closely linked to sample size – more is better. A minimum of 12-15 beat sheet samples are required per management unit (approximately 50 ha) to obtain reasonable estimates of the mirid density. Crop sampling protocols should aim to get as close to the recommended sample number as possible.

When using a standard (380 mm diameter) sweep net, collect 20 sweeps along a single row of cotton per sample. Preliminary research has shown that at least 6 sweep samples are required per management unit (approximately 50 ha) to achieve a good estimation of mirid numbers.

It is essential to also monitor fruit retention and signs of fruit damage to assess if mirid presence is affecting the crop. Note that other stresses (e.g. day or night high temperatures or cloudiness) can also cause square and young boll shedding. If mirid damage is suspected, a sample of bolls from multiple locations should be cut open and checked for internal damage.

**CottonInfo video on mirid sampling and thresholds** [youtu.be/iK63neuw7rU](https://youtu.be/iK63neuw7rU)

## Thresholds

Economic thresholds are based on when the amount of damage by a given population of pests will exceed the costs (product and application) associated with controlling that pest. However this is only an estimate of the true cost of deciding to spray, which includes other costs that may arise later as a result of the control operation (e.g. the flaring of secondary pests such as silverleaf whitefly). Cheap broad-spectrum products may appear to justify low pest control trigger points, but it may be more cost-effective overall to delay control until mirid numbers are higher and use a more expensive but selective product option. Alternatively, the use of a more selective option at a particular pest threshold could be considered on the basis of potential future savings on consequent controls or secondary pest control.

A recent review has reconsidered the historical basis for mirid thresholds in the context of modern high-yielding Bollgard 3 production systems, allowing existing mirid threshold data to be interpreted in a way that better reflects current economics and production practices.

These revised thresholds take into account mirid damage and give the user greater flexibility in determining the relative economics of pest control. They are based on thorough sampling twice weekly, the cost of control (product and application costs/ha), crop stage and commodity value.

For example, previous work suggests that at squaring, the crop will incur a loss of ~0.026 bales/ha/mirid, while at flowering it will incur a loss of ~0.021-0.042 bales/ha/mirid. When the estimated cost of mirid control is \$45/ha, and cotton prices are at \$550/bale, using a threshold of 3 mirids/m until flowering aims to prevent an economic yield loss (see Table 9).

## Deciding when to spray

Under field conditions, the link between mirid density and damage is not straightforward. Sometimes few mirids seem to be causing a lot of damage, while at other times, large populations of mirids cause little damage.

When making a mirid management decision, it is also important to consider:

- **The crop's ability to compensate for damage**, which varies with the season length and the plant's development stage. Well-managed crops with optimised agronomic inputs have a very high capacity for compensation to overcome fruit loss during squaring, but this reduces as the crop progresses through early flowering. Ultimately the capacity for compensation depends on the extent of damage and crop stage, and how these two factors relate to available season length and availability of additional resources such as soil water.
- **Regional differences**. Shorter season areas have a smaller window for crop compensation, particularly after the commencement of flowering. Severe early season damage (tipping and/or fruit loss) may also cause a maturity delay. In regions with shorter seasons, additional caution may be required when managing mirids e.g. using lower thresholds or higher levels of retention.
- **Other factors** that may influence the likelihood of mirid damage include the presence of beneficials (predation and disruption of mirid feeding behaviour), the presence of other pests (mirids may preferentially feed on insect eggs), temperature (possible effects on mirid behaviour), and the behaviour of mirid predators (work on this is on-going). There are also a range of other factors that might be relevant for mirid management such as nymphal stages, the influence of time of day on sampling accuracy, previous spray history, recent weather events or trends, and the likelihood that secondary pests (e.g. whitefly or mealybug) might be flared by ongoing control practices.

**THE FIRST STEP** is to confirm that the damage in the crop is caused by mirids (or other sucking pests) and not environmental factors. Low temperatures can stop cotton plants developing squares, and lack of water, high temperatures, or cloudiness may cause the plant to shed squares. High temperatures on fruit may cause infertility, resulting in beaked bolls. Bolls get a number of black dots on them during the season that are unrelated to mirid activity. If you find damage, but not mirids, it is unlikely that mirids caused the damage.

Table 9 provides a checklist to assist with making the decision to spray based on economic thresholds. The 'Cost of control' should incorporate not only the cost of the insecticide and application, but also likely longer term costs, such as triggering other pests and the need for subsequent sprays. Each farm will vary in this respect so consider your unique system.

## Key beneficials

Damsel bugs and larger lynx spiders, yellow night stalkers and jumping spiders attack mirid adults. These predators also attack smaller instars, as do big-eyed bugs and smaller instars of these spiders. Predatory shield bugs may also attack mirids. None of these beneficials are considered to be specialist mirid predators, however their presence can provide an overall reduction of mirid numbers and reduce the survival of developing nymphs.

## Selecting an insecticide

Insecticide products registered for the control of mirids in cotton in Australia are presented in Table 10. The use of more selective insecticide options will help to conserve beneficial insects (see Table 4 on pages 14-15). A lower rate of indoxacarb mixed with salt provides similar efficacy against mirid and stink bug to the full rate alone but with reduced negative effects on beneficials.

A factor to consider when using reduced rates is that residual efficacy might be reduced and if an influx of adults has had time to deposit eggs within the crop, hatching nymphs may escape control and require a follow up treatment 5-10 days later, potentially negating any benefits. Be cautious when considering reduced rates where populations have built up over time to reach threshold levels, as a mixed population of adults and eggs is more likely to exist compared with treating sudden influxes.

**CottonInfo video on 'soft' options for mirids [youtu.be/zmz70fX4v5U](https://youtu.be/zmz70fX4v5U)**

## Resistance profile

Mirids are not known to have developed resistance to insecticides in Australian cotton, however they are difficult to bioassay and so there is currently limited resistance assessment occurring. It is possible that resistance could develop and the principles underlying the IRMS should be followed in making mirid management decisions.

Many of the products registered for mirid control in cotton are also registered for the control of other pests. It is critical that mirid management decisions also consider sub-threshold populations of other pests that are present in the field, as application against mirids will also select for resistance in these other pests. For example a number of neonicotinoid (Group 4A) insecticides (acetamiprid, clothianidin, dinotefuran) could cause resistance to aphids and silverleaf whitefly. Any decision to use an additional active ingredient (either in a co-formulation or a mix) should be threshold-based. Not only can additional active ingredients be unnecessarily disruptive, but they can lead to resistance. For example use of abamectin (Group 6) when treating mirids has caused high-level resistance in mites.

## Overwintering habit

Mirids survive on weeds and native plant hosts surrounding cotton fields. They also breed on native hosts in inland (central) Australia in winter and can migrate to cotton growing areas in spring in a similar way to *Helicoverpa punctigera*. Understanding whether there are many local hosts or if there has been inland rain (and therefore an abundance of inland hosts), can help with IPM planning for heavy pressure seasons.

## Alternative hosts

Other crop hosts include soybeans, mungbeans, pigeon pea, safflower and sunflowers. It is assumed that mirids migrate between these crops. Weed hosts include turnip weed, Noogoora burr, yellow vine, variegated thistle and volunteer sunflowers.

### Further information:

**Qld DAF, Emerald – Richard Sequeira: 0407 059 066.**

**CSIRO, Narrabri – Simone Heimoana: 0427 992 466.**

**NSW DPI, Tamworth – Lisa Bird: 02 6763 1128.**

**TABLE 9: Mirid thresholds**

| Economic factors                     |                                    | Thresholds (adults or nymphs/m)  |                            |
|--------------------------------------|------------------------------------|--|----------------------------|
| Product value                        | Cost of control (1:1 cost benefit) | Planting to 1 flower/m   | Flowering to 1 open boll/m |
| \$450                                | \$15                               | 1.5  | 1                          |
| \$550                                |                                    | 1  | 0.5                        |
| \$650                                |                                    | 1  | 0.5                        |
| Consider reducing the thresholds if: |                                    | <ul style="list-style-type: none"> <li>• No plant compensation time (e.g. short, cold season)</li> <li>• &lt;60% retention up to 1st flower</li> <li>• &gt;50% light tip damage (black embryo leaves within terminal)</li> <li>• &gt;20% heavy tip damage (terminal and 2-3 nodes dead)</li> <li>• &lt;60-70% retention 1st flower to 4 NAWF</li> <li>• &gt;20% boll damage</li> </ul> |                            |
| Economic factors                     |                                    | Thresholds (adults or nymphs/m)  |                            |
| Product value                        | Cost of control (1:1 cost benefit) | Planting to 1 flower/m   | Flowering to 1 open boll/m |
| \$450                                | \$45                               | 4  | 2.5                        |
| \$550                                |                                    | 3  | 2                          |
| \$650                                |                                    | 2.5  | 1.5                        |
| Consider raising the thresholds if:  |                                    | <ul style="list-style-type: none"> <li>• Time for plants to compensate (e.g. long, warm season)</li> <li>• Many beneficials</li> <li>• High levels of retention (~80%)</li> <li>• High threat of whitefly damage (raises the cost of control)</li> </ul>   |                            |
| Economic factors                     |                                    | Thresholds (adults or nymphs/m)  |                            |
| Product value                        | Cost of control (1:1 cost benefit) | Planting to 1 flower/m   | Flowering to 1 open boll/m |
| \$450                                | \$60                               | 5  | 3                          |
| \$550                                |                                    | 4  | 2.5                        |
| \$650                                |                                    | 3.5  | 2                          |

The thresholds are based on beat sheet sampling, however they are also applicable to sweep net sampling. Visual sampling is not recommended when mirid densities are low. In high mirid density situations, visual thresholds will be approximately half of the corresponding thresholds for beat sheet sampling.

**TABLE 10: Control of mirids**

| Active ingredient                    | MoA group | Overall impact on beneficials* | Comments#   |
|--------------------------------------|-----------|--------------------------------|---|
| Paraffinic oil                       | No group  | <b>Very low</b>                | Apply low rate for suppression of fewer than 0.5 mirids/m. Apply high rate if population reaches threshold of 0.5 mirids/m or apply 2 successive low rate sprays not more than 7 days apart.  |
| <i>Clitoria ternatea</i> extract     | No group  | <b>Low</b>                     | Apply as indicated by field checks and pest presence. Ensure good coverage. Maximum 5 applications per season. Treatment effects may not be seen for 3 or more days. A repeat application may be required at 14-20 days if conditions favour pest development.    |
| Indoxacarb                           | 22A       | <b>Low</b>                     | Under high populations suppression only may be observed. Maximum 3 applications per season.   |
| Indoxacarb + salt                    | 22A       | <b>Low</b>                     | For controlling green mirids ONLY. Use the higher rate on infestations exceeding economic spray threshold levels and/or large canopy crops. Maximum 3 applications per season.  |
| Chlorantraniliprole/<br>Thiamethoxam | 28/4A     | <b>Moderate</b>                | If pest pressure remains high additional control measures may be required from 7 days after application. Do not use as first foliar if neonicotinoid seed treatment used. Maximum 2 applications per season.  |
| Sulfoxaflor                          | 4C        | <b>Moderate</b>                | Use lower rate when infestation is predominately nymphs. Maximum 4 applications per season.   |
| Fipronil                             | 2B        | <b>Moderate</b>                | Apply spray to achieve thorough coverage. Use higher rate under sustained heavy pressure, 3-4 days to reach full effectiveness. Compatible with early season IPM with lower rate having less impact on beneficials. Avoid repeated use of this insecticide group. |
| Fonicamid                            | 29        | <b>Moderate</b>                | Thorough spray coverage is essential. Maximum 2 applications per season.  |
| Emamectin benzoate                   | 6         | <b>Moderate</b>                | For suppression only. Apply to developing populations that are predominantly nymphs. Use non-ionic surfactant at label rate. Maximum effect may take 5 to 7 days. Maximum 2 applications per season.  |
| Emamectin benzoate/<br>acetamiprid   | 6/4A      | <b>Moderate</b>                | Use higher rate on heavier populations and for longer residual control. Use non-ionic surfactant as per label. Maximum 2 applications per season.   |
| Acetamiprid                          | 4A        | <b>Moderate</b>                | Apply with 0.2% Incide penetrant. Target nymphs and/or adults when they reach economic thresholds. On above threshold or increasing populations, suppression only when they reach economic thresholds. Maximum 2 applications per season.                         |
| Clothianidin                         | 4A        | <b>Moderate</b>                | Apply when numbers reach threshold levels requiring treatment. Maximum 2 applications per season.   |
| Imidacloprid                         | 4A        | <b>Moderate</b>                | Do not use as first foliar if neonicotinoid seed treatment used. Maximum 2 applications per season.   |
| Dinotefuran                          | 4A        | <b>Moderate</b>                | When mirids and SLW are present and SLW is at or above threshold levels, use SLW. Performance can be reduced in stressed crops, when senescing late season, or when pests are not actively feeding in the upper crop canopy. Maximum 2 applications per season.   |
| Phorate                              | 1B        | <b>High</b>                    | Qld and WA only. Suppression only. Apply into seed furrow at planting or incorporate into the soil as side dressing deep enough to avoid disturbance by future cultivations. Maximum 1 application per season.  |
| Dimethoate (high rate)               | 1B        | <b>High</b>                    | Do not use where resistant strains are present. Do not harvest for 14 days after application. Do not graze or cut for stockfood for 14 days after application. Maximum 2 applications per season.   |
| Gamma-cyhalothrin                    | 3A        | <b>Very high</b>               | Apply at recommended threshold levels as indicated by field check. Maximum 1 application per season.  |
| Lambda-cyhalothrin                   | 3A        | <b>Very high</b>               | Apply at recommended threshold levels as indicated by field checks. Maximum 1 application per season.   |
| Alpha-cypermethrin                   | 3A        | <b>Very high</b>               | Apply at recommended threshold levels as indicated by field checks. Use the higher rate when pest pressure is high and increased residual protection is required. Maximum 1 application per season.   |
| Bifenthrin                           | 3A        | <b>Very high</b>               | Apply at recommended threshold levels as indicated by field checks. Use the higher rate for increased pest pressure and longer residual control. Maximum 1 application per season.  |
| Deltamethrin                         | 3A        | <b>Very high</b>               | Suppression only. Maximum 1 application per season.   |

#Target pest is predominantly green mirid (*Creontiades dilutus*) For specific species registrations and all other control options ALWAYS refer to the label for instructions.  
 \*For more details about impact on beneficial insects (including bees), refer to Table 4 in this guide.

## Mites

**Two-spotted spider mite** – *Tetranychus urticae*

**Bean spider mite** – *T. ludeni*

**Strawberry spider mite** – *T. lambi*

**Broad mite** – *Polyphagotarsonemus latus*

Two-spotted spider mite is the main pest species, other spider mite species seldom cause economic damage, even in high numbers. In recent years strawberry spider mite has become more common. The difference in damage potential means correct identification of the species present is crucial for good management decisions.

Broad mites are sporadic pests of cotton. They are so small they are difficult to see, even with a hand lens.

Other mite species, including blue oat mite, redlegged earth mite and brown wheat mite occasionally occur on seedling cotton in southern areas.

### Damage symptoms

Mites feed on the underside of leaves but the damage symptoms differ.

**Two-spotted mite** damage appears as brownish areas on the lower leaf surface, usually starting at the junction of the petiole and leaf blade or in leaf folds. Damaged areas show reddening on the upper surface. At high mite densities, leaves become completely red and fall off. Diapausing two spotted mites turn bright orange.

**Bean spider mite** damage results in white, intensely stippled areas on the lower leaf surface, but there is generally no reddening of the upper surface. Severe damage may result in some leaf shedding.

**Strawberry spider mite** can be very abundant (>90% of plants infested) but rarely, if ever, affects yield. Damage is a light, sparse stippling or white dots on the underside of the leaf. There is generally no reddening of the upper leaf surface.

**Broad mite** infestations can resemble herbicide damage. Leaves may distort, develop shiny undersides and become hard or brittle.

### Sampling

Sampling protocols for spider mites in cotton are presented in full on page 40. There is currently no sampling protocol for broad mites, but they tend to infest terminals rather than older leaves, so visually check for broad mite symptoms in the upper canopy.



The top and underside of leaves showing two-spotted spider mite damage (left) and strawberry spider mite damage (right). (Carlos Trapero Ramirez, CSIRO)



Two-spotted mite with egg (mite is 0.5 mm long). (Lewis Wilson)



Bean spider mite. 0.5 mm. (M. Hill)



Strawberry spider mite. 0.3 mm. (C. Mares)



Broad mites. 0.2 mm. (T. Grundy)



Leaves damaged by broad mites become brittle with shiny 'wet' undersides. (Tonia Grundy, Qld DAF)

Look for the presence of any mite stages on the underside of leaves. Use a hand lens, especially if there is doubt about the species present. In young plants, sample the oldest leaf. As plants grow, sample leaves from node 3, 4 or 5 below the terminal. Also look for fine webbing that protects the developing mite population.

Determine the mite species present. Two-spotted spider mite is a pale yellowish-green with one dark spot on either side of its body. Strawberry spider mite is also pale yellowish-green but has three dark spots on each side, which may merge with age, however the adults are only about half the size of adult two-spotted spider mites. Bean spider mite is the same size as two-spotted mite but is dark red.

With nymphs of all ages present, identification can be challenging. It is therefore important to also consider any other signs of damage. For example, leaf reddening is a good indication of two-spotted spider mite.

### Frequency

Sample at least weekly, beginning at seedling emergence. Sample more frequently if mite populations begin to increase, if conditions are hot and dry, or if sprays that impact natural enemy abundance are used.

### Methods

Presence/absence sampling allows many plants to be sampled quickly, thus increasing the likelihood of finding mites if they are present. It is helpful to plot the development of mite populations on a graph to allow changes in mite populations to be seen at a glance.

## Thresholds for two-spotted mites

The following thresholds and yield loss charts have been developed for two-spotted mite only. These thresholds should NOT be applied to bean spider mite or strawberry spider mite. Bean spider mite causes half the damage of two spotted mite, and strawberry spider mite does not cause significant damage.

A general threshold of 30% of plants infested is advocated through the bulk of the season (squaring to first open boll). Yield loss due to mites depends on when mite populations begin to increase and how quickly they increase. Use Table 11 to determine whether the rate of increase in two-spotted mite populations warrants control (adjust for bean spider mite if required).

**Seedling emergence to squaring:** Mites are normally suppressed by predators (particularly thrips) during this period. Mite populations only need to be controlled if they begin to increase, which indicates that natural controls are not keeping them in check.

**Squaring to first open boll:** Implement control if mite populations increase at greater than 1% of plants infested per day in two consecutive checks, or if more than 30% of plants are infested.

**First open bolls to 20% open bolls:** Control is only warranted if mites are well established (greater than 60% plants infested) and are increasing rapidly (faster than 3% of plants infested per day).

**Crop exceeds 20% open bolls:** Control is no longer warranted.

## Two-spotted spider mite yield reduction charts

A simple relationship allows prediction of yield loss from two-spotted spider mites, based on knowledge of the rate of population increase and the time remaining until defoliation. These 'look-up' charts have been provided in Table 11 for areas with different season lengths.

The rate of increase of the mite population is calculated by dividing the change in the percentage of plants infested between consecutive checks by the number of days between the checks. For example, if a field had 10% of plants infested a week ago and 24% infested now, this gives a rate of increase of 2% of plants infested per day.

### To use the charts

1. Select the chart appropriate for your region.
2. Go to the section that is closest to the current infestation level of the field i.e. 10, 30 or 60%.
3. Select the column with the rate of increase closest to that of the mite population in the field.
4. Look down this column to the value that corresponds with the current age of the crop.

This value is the predicted percentage yield loss that the mite population is likely to cause if left uncontrolled. It must be stressed that these charts only provide a guide for potential yield losses caused by mites.

You will need to take into account the vigour of the crop, other pests (you may be about to spray with a pyrethroid which may flare mites) and the conditions (e.g. mite populations develop faster in hot dry conditions). The effect of beneficials is also built-in as high predation on mites will result in lower rates of mite population growth and less risk of yield loss.

## Key beneficials

**Predators** – thrips, minute two-spotted ladybird beetle, mite-eating ladybird beetle, damsel bug, big-eyed bug, brown lacewing adults, brown smudge bug, apple dimpling bug, minute pirate bug, predatory mites, tangleweb spiders.

## Selecting a miticide

The miticide products registered for the control of spider mites in cotton in Australia are presented in Table 12 on page 40. There are currently no registrations for control of broad mites. Amitraz, used for the control of *Helicoverpa* early in the season, will tend to slow, or suppress, the development of mite populations that may also be in the field. Conversely, mite infestations may increase after the application of some broad-spectrum insecticides used for *Helicoverpa* or mirid control, such as synthetic pyrethroids, fipronil, and organophosphates (OP). This occurs because those sprays kill key beneficial species allowing resistant mite populations to flourish (see Table 4 for the impact of pesticides on key beneficials).

Two-spotted mite causes economic damage and has a long history of developing resistance to miticides. While current resistance levels are low for all products excluding OPs, abamectin and pyrethroids, resistance can be selected very quickly. Avoid consecutive sprays of the same miticide. If mite numbers rebuild after a miticide application, rotate to a product from a different chemical group. Once cotton is ~8-10 nodes, thrips cease to be a seedling pest and become important predators of mites. Where thrips are preserved, they can provide sustained suppression of mite populations below damaging levels.

Abamectin resistance has occasionally been detected at high levels in two-spotted mite in horticulture, and frequencies in cotton continue to rise. Bifenthrin and chlorfenapyr resistance in mites occurred largely due to the use of these compounds against other pests. This is true also for abamectin when used in combination with mirid sprays to prevent mite flare.

## Overwintering habit

While the lifecycle slows in cool temperatures, mites are adapted to exploit a wide range of plant hosts and to produce large numbers of offspring, especially as conditions warm up in spring. Control of winter hosts on farms will reduce carry-over of mites between seasons.

## Alternative hosts

Spider mites have a wide host range and can survive through the year on a wide range of broad-leaved weeds. In summer, two-spotted spider mite can also attack sorghum, maize, pulses, sunflowers and many horticultural crops. Preferred winter weed hosts are turnip weed, marshmallow, deadnettle, medics, wireweed and sowthistle, although they can be found on almost any broad-leaved weed species. Alternative winter and spring host crops include safflower, faba beans and field peas. Broad mites also have a wide host range, including many horticultural crop and ornamental species.

### Further information:

**CSIRO Narrabri, Simone Heimoana: 02 6799 1592 or 0427 992 466.**

**NSW DPI Narrabri, Lisa Bird: 02 6763 1128.**

### CottonInfo videos:

**Spider mites in cotton** [youtu.be/oUcatfA0QQM](https://youtu.be/oUcatfA0QQM)

**Sampling spider mites** [youtu.be/2hB84S7p6vY](https://youtu.be/2hB84S7p6vY)

**Broad mites in cotton** [youtu.be/PexkYjLQn0U](https://youtu.be/PexkYjLQn0U)



**TABLE 11: Yield reduction caused by two-spotted spider mites**

The charts below can be used to estimate the percentage of yield reduction, for different cotton growing regions.

| Days from planting   | Current % plants infested with mites |     |      |      |      |       |       |                                   |     |      |      |      |       |       |                                   |      |      |      |      |       |       |
|--|--------------------------------------|-----|------|------|------|-------|-------|-----------------------------------|-----|------|------|------|-------|-------|-----------------------------------|------|------|------|------|-------|-------|
|  | 10                                   |     |      |      |      |       |       | 30                                |     |      |      |      |       |       | 60                                |      |      |      |      |       |       |
|  | Observed rate of increase (%/day)    |     |      |      |      |       |       | Observed rate of increase (%/day) |     |      |      |      |       |       | Observed rate of increase (%/day) |      |      |      |      |       |       |
|  | 0.5                                  | 1   | 1.5  | 2    | 3    | 5     | 7     | 0.5                               | 1   | 1.5  | 2    | 3    | 5     | 7     | 0.5                               | 1    | 1.5  | 2    | 3    | 5     | 7     |
| <b>Warmer regions; planting to 60% bolls open in 134-154 days.</b>                               |                                      |     |      |      |      |       |       |                                   |     |      |      |      |       |       |                                   |      |      |      |      |       |       |
| <b>Biloela, Bourke, Emerald, Macintyre, Mungindi, St. George, Theodore and Walgett</b>           |                                      |     |      |      |      |       |       |                                   |     |      |      |      |       |       |                                   |      |      |      |      |       |       |
| 10   | 1.1                                  | 4.0 | 8.6  | 14.9 | 32.8 | 89.3  | 100.0 | 1.8                               | 5.2 | 17.2 | 10.3 | 36.1 | 94.7  | 100.0 | 3.1                               | 7.3  | 13.2 | 20.8 | 41.2 | 100.0 | 100.0 |
| 20   | 1.0                                  | 3.5 | 7.4  | 12.9 | 28.2 | 76.7  | 100.0 | 1.6                               | 4.6 | 9.0  | 14.9 | 31.2 | 81.6  | 100.0 | 2.6                               | 5.8  | 10.3 | 16.0 | 31.2 | 76.7  | 100.0 |
| 30   | 0.9                                  | 3.0 | 6.3  | 10.9 | 23.9 | 65.0  | 100.0 | 1.5                               | 4.0 | 7.8  | 12.9 | 26.7 | 69.6  | 100.0 | 2.6                               | 5.8  | 10.3 | 16.0 | 31.2 | 76.7  | 100.0 |
| 40   | 0.7                                  | 2.5 | 5.3  | 9.2  | 20.0 | 54.3  | 100.0 | 1.3                               | 3.5 | 6.7  | 10.9 | 22.6 | 58.4  | 100.0 | 2.4                               | 5.2  | 9.0  | 13.9 | 26.7 | 65.0  | 100.0 |
| 50   | 0.6                                  | 2.1 | 4.4  | 7.6  | 16.5 | 44.5  | 86.2  | 1.1                               | 3.0 | 5.6  | 9.2  | 18.8 | 48.3  | 91.5  | 2.2                               | 4.6  | 7.8  | 11.9 | 22.6 | 54.3  | 99.6  |
| 60   | 0.5                                  | 1.7 | 3.6  | 6.1  | 13.3 | 35.7  | 69.1  | 1.0                               | 2.5 | 4.7  | 7.6  | 15.4 | 39.1  | 73.8  | 2.0                               | 4.0  | 6.7  | 10.0 | 18.8 | 44.5  | 81.1  |
| 70   | 0.4                                  | 1.4 | 2.8  | 4.8  | 10.4 | 27.9  | 53.9  | 0.9                               | 2.1 | 3.8  | 6.1  | 12.3 | 30.9  | 58.0  | 1.8                               | 3.5  | 5.6  | 8.4  | 15.4 | 35.7  | 64.5  |
| 80   | 0.3                                  | 1.1 | 2.2  | 3.7  | 7.9  | 21.0  | 40.5  | 0.7                               | 1.7 | 3.1  | 4.8  | 9.5  | 23.7  | 44.1  | 1.6                               | 3.0  | 4.7  | 6.8  | 12.3 | 27.9  | 49.9  |
| 90   | 0.3                                  | 0.8 | 1.6  | 2.7  | 5.7  | 15.1  | 29.1  | 0.6                               | 1.4 | 2.4  | 3.7  | 7.1  | 17.4  | 32.2  | 1.5                               | 2.5  | 3.8  | 5.5  | 9.5  | 21.0  | 37.1  |
| 100  | 0.2                                  | 0.6 | 1.1  | 1.9  | 3.9  | 10.2  | 19.5  | 0.5                               | 1.1 | 2.8  | 2.7  | 5.1  | 12.1  | 22.1  | 1.3                               | 2.1  | 3.1  | 4.2  | 7.1  | 15.1  | 26.2  |
| 110  | 0.1                                  | 0.4 | 0.7  | 1.2  | 2.4  | 6.3   | 11.9  | 0.4                               | 0.8 | 1.3  | 1.9  | 3.4  | 7.7   | 13.9  | 1.1                               | 1.7  | 2.4  | 3.2  | 5.1  | 10.2  | 17.2  |
| 120  | 0.1                                  | 0.2 | 0.4  | 0.6  | 1.3  | 3.3   | 6.1   | 0.3                               | 0.6 | 0.8  | 1.2  | 2.0  | 4.3   | 7.6   | 1.0                               | 1.4  | 1.8  | 2.3  | 3.4  | 6.3   | 10.0  |
| 130  | 0.1                                  | 0.1 | 0.2  | 0.3  | 0.5  | 1.2   | 2.3   | 0.3                               | 0.4 | 0.5  | 0.6  | 1.0  | 1.9   | 3.2   | 0.9                               | 1.1  | 1.3  | 1.5  | 2.0  | 3.3   | 4.8   |
| 140  | 0.0                                  | 0.0 | 0.0  | 0.1  | 0.1  | 0.2   | 0.3   | 0.2                               | 0.2 | 0.3  | 0.3  | 0.3  | 0.5   | 0.6   | 0.7                               | 0.8  | 0.8  | 0.9  | 1.0  | 1.2   | 1.5   |
| <b>Average regions; planting to 60% bolls open in 161-170 days.</b>                              |                                      |     |      |      |      |       |       |                                   |     |      |      |      |       |       |                                   |      |      |      |      |       |       |
| <b>Dalby, Gwydir, Lockyer, Lower Namoi</b>   |                                      |     |      |      |      |       |       |                                   |     |      |      |      |       |       |                                   |      |      |      |      |       |       |
| 10   | 1.5                                  | 5.3 | 11.5 | 20.0 | 44.1 | 100.0 | 100.0 | 2.3                               | 6.7 | 13.5 | 22.6 | 47.9 | 100.0 | 100.0 | 3.7                               | 9.0  | 16.7 | 26.7 | 53.9 | 100.0 | 100.0 |
| 20   | 1.3                                  | 4.7 | 10.1 | 17.6 | 38.8 | 100.0 | 100.0 | 2.0                               | 6.0 | 12.0 | 20.0 | 42.3 | 100.0 | 100.0 | 3.4                               | 8.2  | 15.0 | 23.9 | 47.9 | 100.0 | 100.0 |
| 30   | 1.2                                  | 4.1 | 8.8  | 15.4 | 33.8 | 92.0  | 100.0 | 1.9                               | 5.3 | 10.6 | 17.6 | 37.1 | 97.4  | 100.0 | 3.2                               | 7.4  | 13.5 | 21.3 | 42.3 | 100.0 | 100.0 |
| 40   | 1.0                                  | 3.6 | 7.7  | 13.3 | 29.1 | 79.1  | 100.0 | 1.7                               | 4.7 | 9.3  | 15.4 | 32.2 | 84.2  | 100.0 | 2.9                               | 6.7  | 12.0 | 18.8 | 37.1 | 92.0  | 100.0 |
| 50   | 0.9                                  | 3.1 | 6.5  | 11.3 | 24.8 | 67.3  | 100.0 | 1.5                               | 4.1 | 8.0  | 13.3 | 27.6 | 71.9  | 100.0 | 2.7                               | 6.0  | 10.6 | 16.5 | 32.2 | 79.1  | 100.0 |
| 60   | 0.8                                  | 2.6 | 5.5  | 9.5  | 20.8 | 56.3  | 100.0 | 1.3                               | 3.6 | 6.9  | 11.3 | 23.4 | 60.6  | 100.0 | 2.5                               | 5.3  | 9.3  | 14.3 | 27.6 | 67.3  | 100.0 |
| 70   | 0.6                                  | 2.2 | 4.6  | 7.9  | 17.2 | 46.4  | 89.9  | 1.2                               | 3.1 | 5.8  | 9.5  | 19.5 | 50.3  | 95.2  | 2.3                               | 4.7  | 8.0  | 12.3 | 23.4 | 56.3  | 100.0 |
| 80   | 0.5                                  | 1.8 | 3.7  | 6.4  | 13.9 | 37.4  | 72.4  | 1.0                               | 2.6 | 4.9  | 7.9  | 16.0 | 40.9  | 77.2  | 2.0                               | 4.1  | 6.9  | 10.4 | 19.5 | 46.4  | 84.7  |
| 90   | 0.4                                  | 1.4 | 3.0  | 5.1  | 10.9 | 29.4  | 56.8  | 0.9                               | 2.2 | 4.0  | 6.4  | 12.9 | 32.5  | 61.0  | 1.9                               | 3.6  | 5.8  | 8.7  | 16.0 | 37.4  | 67.7  |
| 100  | 0.4                                  | 1.1 | 2.3  | 3.9  | 8.4  | 22.3  | 43.0  | 0.8                               | 1.8 | 3.2  | 5.1  | 10.0 | 25.0  | 46.8  | 1.7                               | 3.1  | 4.9  | 7.1  | 12.9 | 29.4  | 52.6  |
| 110  | 0.3                                  | 0.8 | 1.7  | 2.9  | 6.1  | 16.2  | 21.2  | 0.6                               | 1.4 | 2.5  | 3.9  | 7.6  | 18.6  | 34.4  | 1.5                               | 2.6  | 4.0  | 5.7  | 10.0 | 22.3  | 39.5  |
| 120  | 0.2                                  | 0.6 | 1.2  | 2.0  | 4.2  | 11.1  | 21.3  | 0.5                               | 1.1 | 1.9  | 2.9  | 5.5  | 13.1  | 23.9  | 1.3                               | 2.2  | 3.2  | 4.5  | 7.6  | 16.2  | 28.2  |
| 130  | 0.2                                  | 0.4 | 0.8  | 1.3  | 2.7  | 7.0   | 13.3  | 0.4                               | 0.8 | 1.4  | 2.0  | 3.7  | 8.5   | 15.4  | 1.2                               | 1.8  | 2.5  | 3.4  | 5.5  | 11.1  | 18.8  |
| 140  | 0.1                                  | 0.3 | 0.5  | 0.7  | 1.5  | 3.8   | 7.1   | 0.4                               | 0.6 | 0.9  | 1.3  | 2.3  | 4.9   | 8.7   | 1.0                               | 1.4  | 1.9  | 2.4  | 3.7  | 7.0   | 11.3  |
| 150  | 0.1                                  | 0.1 | 0.2  | 0.3  | 0.6  | 1.6   | 2.9   | 0.3                               | 0.4 | 0.6  | 0.7  | 1.2  | 2.3   | 3.9   | 0.9                               | 1.1  | 1.4  | 1.6  | 2.3  | 3.8   | 5.7   |
| 160  | 0.0                                  | 0.0 | 0.1  | 0.1  | 0.2  | 0.3   | 0.5   | 0.2                               | 0.3 | 0.3  | 0.3  | 0.4  | 0.7   | 1.0   | 0.8                               | 0.8  | 0.9  | 1.0  | 1.2  | 1.6   | 2.0   |
| <b>Cooler regions; planting to 60% boll open in &gt;170 days.</b>                                |                                      |     |      |      |      |       |       |                                   |     |      |      |      |       |       |                                   |      |      |      |      |       |       |
| <b>Upper Namoi, Boggabri, Breeza, Cecil Plains, Pittsworth, Trangie, Macquarie, Southern NSW</b> |                                      |     |      |      |      |       |       |                                   |     |      |      |      |       |       |                                   |      |      |      |      |       |       |
| 10   | 1.7                                  | 6.3 | 13.6 | 23.7 | 52.2 | 100.0 | 100.0 | 2.6                               | 7.7 | 15.7 | 26.5 | 56.3 | 100.0 | 100.0 | 4.1                               | 10.2 | 19.2 | 30.9 | 62.8 | 100.0 | 100.0 |
| 20   | 1.6                                  | 5.6 | 12.1 | 21.0 | 46.4 | 100.0 | 100.0 | 2.3                               | 7.0 | 14.1 | 23.7 | 50.3 | 100.0 | 100.0 | 3.8                               | 9.4  | 17.4 | 27.9 | 56.3 | 100.0 | 100.0 |
| 30   | 1.4                                  | 4.9 | 10.7 | 18.6 | 40.9 | 100.0 | 100.0 | 2.1                               | 6.3 | 12.6 | 21.0 | 44.5 | 100.0 | 100.0 | 3.5                               | 8.5  | 15.7 | 25.0 | 50.3 | 100.0 | 100.0 |
| 40   | 1.2                                  | 4.3 | 9.4  | 16.2 | 35.7 | 97.4  | 100.0 | 1.9                               | 5.6 | 11.1 | 18.6 | 39.1 | 100.0 | 100.0 | 3.3                               | 7.7  | 14.1 | 22.3 | 44.5 | 100.0 | 100.0 |
| 50   | 1.1                                  | 3.8 | 8.1  | 14.1 | 30.9 | 84.2  | 100.0 | 1.7                               | 4.9 | 9.8  | 16.2 | 34.1 | 89.3  | 100.0 | 3.0                               | 7.0  | 12.6 | 19.8 | 39.1 | 97.4  | 100.0 |
| 60   | 0.9                                  | 3.3 | 7.0  | 12.1 | 26.5 | 71.9  | 100.0 | 1.6                               | 4.3 | 8.5  | 14.1 | 29.4 | 76.7  | 100.0 | 2.8                               | 6.3  | 11.1 | 17.4 | 34.1 | 84.2  | 100.0 |
| 70   | 0.8                                  | 2.8 | 5.9  | 10.2 | 22.3 | 60.6  | 100.0 | 1.4                               | 3.8 | 7.3  | 12.1 | 25.0 | 65.0  | 100.0 | 2.6                               | 5.6  | 9.8  | 15.1 | 29.4 | 71.9  | 100.0 |
| 80   | 0.7                                  | 2.3 | 4.9  | 8.5  | 18.6 | 50.3  | 97.4  | 1.2                               | 3.3 | 6.3  | 10.2 | 21.0 | 54.3  | 100.0 | 2.3                               | 4.9  | 8.5  | 13.1 | 25.0 | 60.6  | 100.0 |
| 90   | 0.6                                  | 1.9 | 4.1  | 7.0  | 15.1 | 40.9  | 79.1  | 1.1                               | 2.8 | 5.3  | 8.5  | 17.4 | 44.5  | 84.2  | 2.1                               | 4.3  | 7.3  | 11.1 | 21.0 | 50.3  | 92.0  |
| 100  | 0.5                                  | 1.6 | 3.3  | 5.6  | 12.1 | 32.5  | 62.8  | 0.9                               | 2.3 | 4.3  | 7.0  | 14.1 | 35.7  | 67.3  | 1.9                               | 3.8  | 6.3  | 9.4  | 17.4 | 40.9  | 74.3  |
| 110  | 0.4                                  | 1.2 | 2.6  | 4.3  | 9.4  | 25.0  | 48.3  | 0.8                               | 1.9 | 3.5  | 5.6  | 11.1 | 27.9  | 52.2  | 1.7                               | 3.3  | 5.3  | 7.7  | 14.1 | 32.5  | 58.4  |
| 120  | 0.3                                  | 0.9 | 1.9  | 3.3  | 7.0  | 18.6  | 35.7  | 0.7                               | 1.6 | 2.8  | 4.3  | 8.5  | 21.0  | 39.1  | 1.5                               | 2.8  | 4.3  | 6.3  | 11.1 | 25.0  | 44.5  |
| 130  | 0.2                                  | 0.7 | 1.4  | 2.3  | 4.9  | 13.1  | 25.0  | 0.6                               | 1.2 | 2.1  | 3.3  | 6.3  | 15.1  | 27.9  | 1.4                               | 2.3  | 3.5  | 4.9  | 8.5  | 18.6  | 32.5  |
| 140  | 0.2                                  | 0.5 | 0.9  | 1.6  | 3.3  | 8.5   | 16.2  | 0.5                               | 0.9 | 1.6  | 2.3  | 4.3  | 10.2  | 18.6  | 1.2                               | 1.9  | 2.8  | 3.8  | 6.3  | 13.1  | 22.3  |
| 150  | 0.1                                  | 0.3 | 0.6  | 0.9  | 1.9  | 4.9   | 9.4   | 0.4                               | 0.7 | 1.1  | 1.6  | 2.8  | 6.3   | 11.1  | 1.1                               | 1.6  | 2.1  | 2.8  | 4.3  | 8.5   | 14.1  |
| 160  | 0.1                                  | 0.2 | 0.3  | 0.5  | 0.9  | 2.3   | 4.3   | 0.3                               | 0.5 | 0.7  | 0.9  | 1.6  | 3.3   | 5.6   | 0.9                               | 1.2  | 1.6  | 1.9  | 2.8  | 4.9   | 7.7   |
| 170  | 0.0                                  | 0.1 | 0.1  | 0.2  | 0.3  | 0.7   | 1.2   | 0.2                               | 0.3 | 0.4  | 0.5  | 0.7  | 1.2   | 1.9   | 0.8                               | 0.9  | 1.1  | 1.2  | 1.6  | 2.3   | 3.3   |

**SAMPLING PROTOCOLS FOR SPIDER MITES IN COTTON**

**Population monitoring**

1. Walk into the field about 40 m. Early in the season also sample near the field edges to see if significant influxes of mites have occurred.
2. Take a leaf from the third, fourth or fifth main-stem node below the terminal of the first plant. If the plant has less than three leaves, sample the oldest. Early in the season until the plant has about five true leaves, it is simplest to pull out whole plants.
3. Walk five steps and take a leaf from the next plant, on your opposite side and so on until you have 50 leaves.
4. Once all the leaves have been collected, score each leaf by turning it over and looking at the underside, firstly near the stalk, then scanning the rest of the leaf. If mites of any stage (including eggs) are present, score the leaf as infested. A hand lens will be needed as mite eggs cannot be seen with the naked eye.
5. Repeat at several widely separated places in the field to allow for differences in mite abundance. Depending on the field size, 4-6 sites are needed to obtain a good estimate of mite abundance.
6. Calculate the percentage of plants infested in the field. Continue presence/absence sampling until 20% open bolls.

**Additional recommendations for mites in seedling cotton**

On seedling cotton (up to 6-8 true leaves) sample regularly to determine the level of infestation using the presence/absence technique above.

When more than 5% of plants are infested also count the numbers of mites on plants, and score the damage level (i.e. estimate the percentage of total leaf area per plant that is damaged by mites). Continue to monitor mite numbers, damage levels and infestation levels at least weekly, or more frequently if >30% of plants are infested.

If the level of infestation, damage level or mite number per plant declines then control is unnecessary, but monitoring should continue.

If mite numbers per plant do not decline after about 6 weeks, if damage levels exceed an average of 20% of plant leaf area, or if infestation levels increase, then predators are not abundant enough to control mites and a miticide should be applied.

**Miticide resistance monitoring**

1. If mites are being collected after a miticide application, ensure sufficient time has lapsed for the miticide to be fully activated. Depending on the product, this may take 7 to 10 days. Avoid making collections and sending samples on Thursdays or Fridays.
2. Collect 50 infested leaves from different areas across the field. Keep samples from different fields separate. If mite numbers per leaf are very low, consider collecting up to 100 leaves.
3. Phone Lisa Bird (see below) to let her know you are sending the sample.
4. Clearly label samples with the following information:

Farm Name .....

Field .....

Region (e.g. Gwydir) .....

Collector's Name.....

Phone No.....

Fax No.....

Email address .....

Date of collection ...../...../.....

Comments (e.g. details of the problem if a control failure has occurred).

**Samples for testing**

5. Pack the leaves loosely in a paper bag, fold and staple the top. Pack this in a 6-pack esky. Attach the sample details and send by overnight courier to:

**Lisa Bird**  
Department of Primary Industries  
Tamworth Agricultural Institute  
4 Marsden Park Rd  
Calala NSW 2340  
Ph: 02 67631128 or 0438 623 906

**TABLE 12: Control of two-spotted spider mites**

| Active ingredient  | MoA group | Mite resistance                | Overall impact on beneficials* | Comments#  |
|--------------------|-----------|--------------------------------|--------------------------------|--|
| Etoazole           | 10B       | Occasional – low               | Low                            | Good coverage is essential. Refer to label for no-spray zones and record keeping. Best on low to increasing populations. Maximum 1 application per season.   |
| Dicofol            | 2B        | No data                        | Low                            | NSW registration only. Apply by ground rig at first appearance of mites before row closure. Maximum 2 applications per season.   |
| Diafenthiuron      | 12A       | No resistance                  | Low                            | Treatment at higher infestation levels may lead to unsatisfactory results. Maximum 2 applications per season.  |
| Abamectin          | 6         | Widespread – med/high          | Moderate                       | Best results will be obtained when applied to low mite populations. Maximum 2 applications per season.   |
| Emamectin benzoate | 6         | Abamectin                      | Moderate                       | When applied for <i>Helicoverpa</i> control will reduce the rate of mite population development. Suppression only. Maximum 2 applications per season.  |
| Propargite         | 12C       | Occasional – low               | Moderate                       | Apply spray before mite infestations reach damaging levels as maximum efficacy is not reached until 2 weeks after spraying. Maximum of 2 non-consecutive applications per season.  |
| Amitraz            | 19        | No data                        | Moderate                       | Suppression when used for controlling <i>Helicoverpa</i> . Maximum 4 applications per season.  |
| Dimethoate         | 1B        | No data                        | High                           | Will not control organophosphate-resistant mites. Do not harvest for 14 days after application. Do not graze or cut for stockfood for 14 days after application. Maximum 2 applications per season.  |
| Phorate            | 1B        | No data                        | High                           | Use lower rate for short residual control at time of planting. For extended period of control use higher rates. Only use the highest rate on heavy soils when conditions favour good emergence. Maximum 1 application per season. Note that rates differ by state. |
| Bifenthrin         | 3A        | Widespread – med/high          | Very high                      | Applications against <i>Helicoverpa</i> will give good control of low mite populations. Maximum 1 application per season.  |
| Deltamethrin       | 3A        | Cross-resistance to Bifenthrin | Very high                      | Suppression only. Maximum 1 application per season.  |

#For all control options ALWAYS refer to the label for instructions.

\*For more details about impact on beneficial insects (including bees), refer to Table 4 in this guide.

## Pale cotton stainers

### *Dysdercus sidae*

Pale cotton stainers are occasional pests of cotton in Australia. Economic damage is rare because of their:

- Susceptibility to insecticides used for other pests, especially in conventional cotton;
- Need for free water to be present; and,
- Inability to survive high temperatures (>40°C).

However in mild seasons Bt crops may present a favourable environment for cotton stainers and management may be required.

Usually cotton is infested by adults flying into fields around the time of first open boll from up to 15 km away. Sometimes, perhaps due to seasonal conditions, populations can be found earlier, during boll maturation. Adults will mate soon after arrival and the expanding population of developing nymphs is likely to cause economic damage.

### Damage symptoms

Pale cotton stainers can penetrate the boll wall of young and mature bolls to feed on cotton seeds and will also feed on seeds in open bolls. Seed weight, oil content and viability all decline as a result. Loss of seed viability should be a consideration in pure seed crops.

Damage to bolls up to 20 days old causes small black marks on the outer surface and may cause warty growths on the inner boll wall. Damage to older bolls usually shows no external symptoms and only small dark marks will be seen on the inside of the boll wall. Most damage is to seeds, reducing their size and vigour, and sometimes lint production is affected. Tight lock can result around damaged seeds, preventing the lint from fluffing out as the boll opens, and damaged locks are often yellow or stained.



Damage to cotton seed. (Simone Heimoana, CSIRO)

### Sampling

Where adults and nymphs are observed feeding, monitor the percentage of damaged bolls. Sample at least weekly once bolls are present, and more often if pale cotton stainer numbers approach threshold.

Distribution through the field and through the canopy can be quite patchy, as adult females lay eggs in clusters in the soil or sometimes in open bolls. Sample with a beat sheet at multiple sites spread throughout the field. Search visually for younger instars in the lower canopy.

Cut open bolls of varying ages to monitor for signs of damage. Examine seeds for browned, dried damage areas. A week after damage the lint may begin to yellow and locks will be stuck to the boll wall.

The mild, wet conditions that favour the survival of pale cotton stainers in cotton will also favour the occurrence of secondary infections by fungi and bacteria in cracked bolls. These infections can also cause tight lock and lint staining, so in older crops, do not automatically assume these symptoms are caused by stainers.



Juvenile (left) and adult pale cotton stainers. (Lewis Wilson)

### Thresholds

**Boll development:** When stainers are observed in the crop and damage to developing bolls is detected, an action threshold of 3 pale cotton stainers/m is recommended (based on beat sheet sampling – see GVB adult equivalents on page 25). Adults and older nymphs (3rd to 5th instars) cause similar levels of damage.

**After first open boll:** When stainers are observed feeding in open bolls, consider the potential for quality downgrades of the lint as well as the loss of seed weight and seed viability. Where staining is observed, use a threshold of 30% affected bolls to prevent a colour downgrade.

### Key beneficials

A range of natural enemies such as parasitic flies and predatory bugs (e.g. assassin bugs) have been recorded overseas. The role of natural enemies in Australia has not been studied.

### Selecting an insecticide

There are few products registered for pale cotton stainer control apart from the synthetic pyrethroids lambda-cyhalothrin and gamma-cyhalothrin, however their status as an occasional pest is influenced by their susceptibility to insecticides used for the control of *Helicoverpa* and other pests such as mirids. Any decision to use broad-spectrum insecticides should take into account their impact on beneficial insects and the subsequent risk of flaring whitefly and other secondary pests.

Worldwide, there are few records of resistance to insecticides developing in the field, however cotton stainers will react to selection pressure under laboratory conditions.

### Overwintering habit

As there is no resting stage in the cotton stainer's lifecycle, cultural controls, particularly management of alternative hosts between cotton seasons, assist greatly in limiting population development.

### Alternative hosts

Fuzzy cotton seed used for stockfeed is an important alternative source of food for cotton stainers. Avoid storing fuzzy seed in exposed places accessible to cotton stainers over long periods. Controlling volunteer and ratoon cotton is also important for limiting cotton stainer's access to alternative food sources.

#### Further information:

CSIRO, Narrabri

Simone Heimoana: 02 6799 1592 or 0427 992 466

## Soil and establishment pests

**True wireworms** – *Agrypnus* spp.

**False wireworms** – *Gonocephalum* spp.

*Pterohelaeus* spp.

**Black field earwig** – *Nala lividipes*

**Symphyla** – *Hanseniella* spp.

Soil pests can reduce plant establishment, row density and vigour. Symptoms can be confused with other establishment problems, and may be worse if seedling development is slow due to climate or other factors such as allelopathy, disease or soil constraints. Control measures must usually be implemented at planting, so pre-plant sampling for these pests is required, particularly in fields with a history of soil pest activity.

### Sampling

Sampling for earwigs and wireworms is best conducted using a baiting technique. Soil surveying can also be used for detecting pests such as symphylans and scarab larvae.

**Grain or potato baiting** can be conducted following planting rain or irrigation:

1. Soak insecticide-free crop seed in water for at least two hours to initiate germination or cut medium sized potatoes in half. The type of seed used makes no noticeable difference in attracting soil-dwelling insects.
2. Bury a dessert spoon full of soaked seed or a potato half (with the cut side facing down) under 1 cm of soil at each corner of a 5x5 m square at five widely-spaced sites per 100 ha.
3. Mark the position of the baits to easily find them again (high populations of soil insects can completely destroy them).
4. One day after seedling emergence or 5-7 days after placing the potato baits, dig up the remaining bait and count the insects. Feeding on the potato bait will be immediately obvious.

**CottonInfo video on sampling for soil insects** [youtu.be/i\\_ODxTAeASw](https://youtu.be/i_ODxTAeASw)

**Soil surveying** involves directly digging to sample for pests. Insert a shovel to full depth at the plant line on the hill and carefully lever the soil out for closer inspection. Start with the deepest soil. Holding a soil clod in one hand, use your other hand to carefully break the soil apart.

### Factors influencing soil pest numbers

Tillage and farm management practices can influence the composition and abundance of pest species. For example weedy fallows encourage the abundance of soil pests whereas clean fallows generally cause a decline in pest insect numbers due to a lack of food.

The influence of field stubble is contentious as high stubble loads within fields will promote soil pests, however stubble can also provide a diversionary food source as well as increase the diversity of other soil fauna such as predatory beetles (carabidae), centipedes and earthworms. The incorporation of grains stubble prior to planting cotton may increase the damage potential of black field earwig populations as it can cause them to switch feeding activity from stubble to seedlings. Wireworms are found under a range of cultivation and stubble retention regimes.

### True and false wireworms

Wireworms are thin, cream-coloured beetle larvae about 20 mm long.

Larvae attack germinating seeds, the hypocotyl and roots of young cotton plants resulting in seedling death, young plant 'felling' and patchy plant stands. The adult beetles can also damage seedlings by chewing at or just above ground level.



**True wireworm larva.** (Tonia Grundy, Qld DAF)

Conduct bait sampling prior to planting to determine wireworm abundance. There are no specific thresholds in cotton, but densities of one or more larvae per baiting site are considered damaging in summer grains.

Wireworm larvae are unlikely to be controlled with standard seed treatments, so where populations are high, an in-furrow insecticide treatment at planting should be considered. Infestations detected after crop emergence cannot be controlled with baiting or surface spraying, so early detection is essential for effective management.

### Black field earwigs



**Black field earwigs can be difficult to see in dark soils.**

Black field earwigs are about 15 mm long. They are an occasional pest of seedling cotton, predominantly feeding on germinating seed and seedling roots, resulting in poor establishment. Nymphs look similar to adults.

Conduct bait sampling prior to planting to determine earwig presence. No thresholds have been defined for cotton, but thresholds used for maize and sorghum suggest that control maybe warranted when more than 50 earwigs are found across 20 baits or 2-3 earwigs per bait sample.

If earwig numbers are high the application of insecticide treated grain baits at sowing may offer protection. In-furrow insecticide treatments have generally been ineffective if dense populations are present. The efficacy of seed dressings for black field earwig control is unknown.

### Symphylans

Symphyla are small (up to 7 mm), white, 'millipede-like' arthropods with 12 pairs of legs. Relatively common in most soils, they may be confused with other soil organisms such as diplurans or collembola (springtails).

Symphyla generally feed on decomposing organic matter but can feed on rootlets and root hairs and have been associated with crop establishment issues in some fields, often as part of a broader complex of soil pests. Damage is more likely in dry soil conditions. They can move up and down in the soil profile and distribution within a field is generally patchy. If plants show symptoms of damage, conduct a basic soil survey, keeping a close eye on newly broken-apart clods for movement. Symphyla are fast moving and will rapidly shift to avoid sunlight.



# A PROFESSIONAL NETWORK OF AGRONOMISTS AND CROP CONSULTANTS

SERVICING THOSE NEW TO THE INDUSTRY THROUGH TO EXPERIENCED ADVISORS

Crop Consultants Australia (CCA) supports those that provide agronomic advice to Australia's cotton, grain, pulse and oilseed producers. For over 30 years the association has provided:

-  Professional development and upskilling
-  Research, product updates and industry news
-  Networking opportunities



**Symphylan (left) versus a dipluran (right).** (Paul Grundy Qld DAF)

Recent crop surveys and pot trials suggest that symphyla are unlikely to be the lone cause of crop establishment issues. In the absence of other soil pests and diseases or under more optimal field conditions their impact may be minimal.

There are no recommended chemical control options. The in-furrow application of insecticide at planting will not provide protection for establishing seedlings as symphyla are active to depths of up to one metre and will easily avoid exposure. Standard seed dressings offer limited protection when high densities of soil pests are present.

Plant fields where symphyla have been abundant last, so that the warmer conditions aid more rapid establishment. Roots that grow deeper into the profile more quickly are less likely to become stranded in dry soil through root pruning. If plants show signs of moisture stress where symphyla are present, a quick flush with irrigation may assist. Irrigation can also decrease symphyla activity in the upper profile for about 7-14 days, which may assist crop recovery.

If establishment is low enough to warrant replanting, consider alternate fibrous-rooted crops such as maize or sorghum that are less susceptible.

**Other soil pests**



**Black field cricket.** (J. Wessels)

**Black field cricket** (*Teleogryllus commodus*) adults and late instar nymphs may occasionally damage cotton seedlings if occurring in large numbers, but damage leading to replanting is uncommon.



**Cutworm larvae.** (P. Room)

**Cutworms** (*Agrotis* spp.) chew stems at ground level and can be a pest of emerging cotton but is rare. This pest is typically found along field margins that adjoin pastures or where cotton has been sown into recently sprayed out weedy fallows.



**Whitegrub attacking peanut.** (J. Wessels)

**Whitegrubs** are the larvae of scarab beetles and can feed on the roots of crops, causing a loss of vigour and promoting lodging. Damage in cotton is rare and likely only if sown into fields that were previously a weedy fallow or a summer sorghum crop.

**TABLE 13: Control of wireworm (refer to label for target species)**

| Active ingredient           | MoA group | Overall impact on beneficials* | Comments#  |
|-----------------------------|-----------|--------------------------------|--|
| Phorate                     | 1B        | High                           | Apply into the seed furrow at sowing. Maximum 1 application per season.  |
| Chlorpyrifos                | 1B        | High                           | Use the higher rate with extreme population numbers. Maximum 3 applications per season. Use minimum spray volume of 20 L per sown ha.  |
| Bifenthrin                  | 3A        | Very high                      | Apply as spray into the furrow at planting. Use a spray nozzle which will deliver a coarse spray in a total volume of 60-100 L/ha. Maximum 1 application per season.   |
| Bifenthrin/<br>chlorpyrifos | 3A/1B     | Very high                      | Apply as directed spray into the furrow at planting. Use a spray nozzle which will deliver a coarse spray in a total volume of 60-100 L/ha. Use the higher rate with extreme population numbers. Maximum 1 application per season. |

#For all control options ALWAYS refer to the label for instructions.

\*For more details about impact on beneficial insects (including bees), refer to Table 4 in this guide.

## Thrips

**Tobacco thrips** – *Thrips tabaci*

**Tomato thrips** – *Frankliniella schultzei*

**Western flower thrips** – *F. occidentalis*

Thrips are small thin insects that are primarily pests of cotton seedlings, but can also act as key predators of mite eggs later in the season.

### Damage symptoms

Thrips larvae and adults cause early season damage to terminals, leaves and stems. The most obvious damage is smaller, crinkled leaves and a visible 'silvering' on the undersides of leaves. At very high densities, thrips can also kill the growing terminal, which delays the plant's growth until it can establish a new terminal.

Occasionally thrips can also build to high numbers in flowers and on leaves mid-late season, particularly in crops with minimal pesticide use. High numbers on leaves can lead to stunting and damage especially along leaf veins, but late season thrips damage rarely justifies control.

**CottonInfo video on thrips identification and damage:** [youtu.be/e8B0wpimUvA](https://youtu.be/e8B0wpimUvA)

### Sampling

Sample seedlings and count the number of thrips/plant. Adults thrips can migrate into the crop from surrounding vegetation, however presence of larvae in crops that have had an insecticide seed treatment or in-furrow insecticide treatment indicates that the population is actively breeding and the early insecticide offered poor control. No larvae and little plant damage indicates that the insecticide is effectively controlling the thrips.

Score the severity of damage to the seedlings by estimating the percentage reduction in leaf area.

### Frequency

Sample weekly from seedling emergence until thrips abundance declines and plants begin to recover (usually by about 4-8 nodes, but sometimes up to 10 nodes). In the mid to late season monitor for the presence of thrips in flowers and on the undersides of leaves in the upper canopy. Also look for thrips when sampling mites, as the presence of thrips adults and larvae in mite colonies is a good indicator of potential natural control of the mites.



**Thrips damage to lower nodes with terminal showing new growth without damage. Plant is likely to recover, however, continue to monitor.** (Lewis Wilson)



**Thrips nymph on underside of leaf and adult in cotton flower.** (Tonia Grundy, Qld DAF)

### Methods

Use a hand lens to count the number of adult and larval thrips on 20-30 separate seedlings for every 50 ha of crop. Also assess leaf damage (if the damaged young leaves are less than 1 square centimetre then the reduction in seedling leaf area is usually greater than 80%).

If thrips are present in high numbers (>30/plant), check the plant terminal. Complete blackening of the embryonic leaves means the terminal has died.

**CottonInfo video on thrips sampling and management:** [youtu.be/FKkwWID4iQA](https://youtu.be/FKkwWID4iQA)

### Thresholds

As thrips occur in cotton in most years the most effective management option is to use a seed treatment or an at-planting insecticide applied with the seed. This protects plants during the establishment phase and has the advantage of minimal negative effects on beneficial species compared to an insecticide applied to the crop after emergence (see Tables 3 and 4).

Severe damage to leaves can result in delayed maturity or yield loss. In northern and central regions with warmer climates, the risk is lower because plants can outgrow and compensate for damage, so yield loss is only likely to occur about one year in 10.

In cooler, shorter season areas (Downs, Upper Namoi, Macquarie and Liverpool Plains), the risk of delayed maturity and/or yield loss is higher because there is less time to compensate, hence yield loss may occur one year in every two. Southern NSW regions have a similar risk, however other issues relating to a cool start and associated maturity delays can further compound thrips' impact, so the ability of the crop to compensate from damage will need to be considered.

In all instances a seed treatment or a planting insecticide applied with the seed should provide sufficient control for plants to establish. Neonicotinoid (imidacloprid) resistance has been detected in tobacco thrips from cotton, and may effect neonicotinoid seed dressing efficacy. Control may be required if significant leaf damage continues past 6-8 nodes.

Thrips populations will normally naturally decline in early December. In situations where populations of thrips remain high and plant growth is delayed by cool, wet weather, seed treatments or at-planting insecticides may run out and supplementary control may be necessary according to the thresholds below.

Western flower thrips is not controlled by the current seed treatments, but this species is not normally abundant in early season cotton.

### Thresholds

| SEEDLING TO 6 TRUE LEAVES  |
|--|
| 80% reduction in leaf area<br>+<br>10 thrips/plant (adults and larvae) |

Thrips found in cotton in the mid and late season are usually *Frankliniella* spp. Adult thrips feed on pollen, but are unlikely to affect pollination or fruit set. Eggs are laid on leaves and the hatching larvae may cause damage to the undersides, resulting in distorted, smaller leaves. As these larvae are also predatory, high levels of damage would be required to affect yield, and control should not be considered unless >30% of leaf area is damaged in the top 6 nodes in pre-cut-out crops or more than 50% of leaf area damaged after the crop has cut-out.

CottonInfo video on whether thrips are friends or foes: [youtu.be/UsWxHB94NN4](https://youtu.be/UsWxHB94NN4)

### Key beneficials

**Predators** – minute pirate bug, green lacewing larvae, brown lacewing, ladybird beetles.

### Selecting an insecticide

Insecticide products registered for the control of thrips in cotton in Australia are presented in Table 14. Insecticide seed treatments will provide protection from thrips early in the crop. A neonicotinoid seed treatment should not be used if resistance to neonicotinoids in thrips is suspected. Consider the benefit of thrips as predators of spider mites when deciding whether to control thrips in cotton with an insecticide. **It should be noted that every western flower thrips tested in Australia since resistance screening commenced has been pyrethroid resistant, so insecticides in that mode of action group (3A) must be avoided.**

In Australia, pyrethroid and OP resistance has been detected in some tobacco thrips associated with bulb onion but resistance in tomato thrips has not been detected.

### Overwintering habit

Thrips prefer milder temperatures. Populations decline at temperatures greater than 30°C. Thrips are active and common through winter on a range of hosts.

### Alternative hosts

Thrips continue to feed and reproduce on a range of weed hosts during winter and spring. Adult thrips may migrate from these weeds into flowering wheat crops but generally don't reproduce there. In spring, as weeds and wheat crops dry out, large numbers of adults are forced to seek new hosts, and transfer to cotton. Cotton crops planted adjacent to cereal crops are particularly at risk of infestation by thrips.

**Further information:**

**CSIRO Agriculture, Narrabri –  
Simone Heimoana: 02 6799 1592 or 0427 992 466  
Lisa Bird: 02 6763 1128.**

**TABLE 14: Control of thrips**

| Active ingredient   | MoA group | Overall impact on beneficials* | Comments#  |
|---|-----------|--------------------------------|--|
| <b>Tobacco thrips (<i>Thrips tabaci</i>) and tomato thrips (<i>Frankliniella schultzei</i>)</b> |           |                                |  |
| Fipronil  | 2B        | Moderate                       | Fipronil will take 3-4 days to reach full effectiveness. Use higher rates under high pressure. Avoid repeated use of this insecticide group.   |
| Phorate   | 1B        | High                           | Use low rates for short residual control at time of planting. Use high rates for extended period of control. Only use the highest rate on heavy soils when conditions favour good emergence. Maximum 1 application per season. |
| Dimethoate  | 1B        | High                           | Do not harvest for 14 days after application. Maximum 2 applications per season.   |
| <b>Western flower thrips (<i>Frankliniella occidentalis</i>)</b>                                |           |                                |  |
| Spinetoram  | 5         | Low                            | Maximum 2 applications per season. Refer to mandatory no-spray zone on label.  |

#For all control options ALWAYS refer to the label for instructions.

\*For more details about impact on beneficial insects (including bees), refer to Table 4 in this guide.



## Whitefly

**Silverleaf whitefly – *Bemisia argentifolii***

**Greenhouse whitefly – *Trialeurodes vaporariorum***

Silverleaf whitefly (SLW) is a major pest that infests mid to late-stage cotton. If populations build up, their honeydew becomes a serious contaminant of cotton lint. SLW also have a track record of developing insecticide resistance, increasing the difficulty of their management. Greenhouse whitefly is a rare minor pest of cotton.

### Damage symptoms

Both adults and nymphs excrete honeydew, which dries to a matte, non-sticky consistency on lint, making contamination difficult to detect visually. The sugar composition, in particular the high concentration of trehalulose, makes SLW-contaminated lint 'sticky' and more difficult to process at gins and textile mills compared to aphid honeydew-contaminated cotton, usually leading to downgrading or rejection. Whitefly honeydew can also promote the growth of sooty mould.

**CottonInfo video on implications for lint quality and colour** [youtu.be/uVSSlqUg5q0](https://youtu.be/uVSSlqUg5q0)

### Sampling

#### Sample for species and population

While SLW can be visually differentiated from greenhouse whitefly by the wing shape in adults and the absence of hairs on the nymphs (see photographs on page 50), they cannot be easily distinguished from endemic *Bemisia* species that are occasionally present in cotton crops. These endemic species are less common and tend to have slower population growth so are rarely important in pest management decisions.

Effective sampling is the key to successful management. Keep an eye out for SLW while sampling for other pests pre-flowering to assist in making decisions about product selections; conservation of beneficial species may help delay build-up of SLW or other pests. Begin sampling between 1000 and 1300 day degrees (DD) and continue for several weeks to record changes in nymph abundance in relation to cotton development.

SLW sampling aims to monitor changes in the nymph population density over time. SLW densities naturally fluctuate and during hot and dry conditions populations can rapidly increase, so sample twice weekly.

### Sampling protocol







A management unit can be a whole field or part of a field. Start at least 20 metres from the field edge and collect one leaf per plant from the lower canopy (11-14 nodes down from the main stem terminal shoot), gathering at least 30 main stem leaves per management unit from plants 3-5 metres apart.

**CottonInfo video on sampling SLW nymphs** [youtu.be/gWkTKMmJYLw](https://youtu.be/gWkTKMmJYLw)

Count the total number of large nymphs on each leaf, and note the number that are viable ('red eye') versus non-viable (dead or parasitised). Nymph density on leaves is easier to assess prior to significant wilting and with magnification (e.g. hand lens). A phone app that can be used to image and record nymph densities is under development. Refer to the CottonInfo website for more information [www.cottoninfo.com.au/insect-and-mite-management](http://www.cottoninfo.com.au/insect-and-mite-management)




## Silverleaf whitefly: healthy vs parasitised late stage (4<sup>th</sup> instar) nymphs

Healthy SLW










Fourth instar nymph with healthy mycetomes      Red eye nymph development (the white wing buds appear just before emergence)

Parasitised SLW











Adult *Encarsia* (left) and *Eretmocerus* sp.      Empty nymph case with chewed exit hole





Empty pupal case (ripped open) and adult SLW



Signs of parasitism:

Displaced mycetomes      Parasitoid larva      Red faecal waste      Examples of parasitoid pupae (colours may vary)



With support from:

Last revised July 2021

### The new SLW decision support tool

An industry-developed decision support tool (DST), figure below, is available from [www.cottoninfo.com.au/silverleaf-whiteflydecision-support-tool](http://www.cottoninfo.com.au/silverleaf-whiteflydecision-support-tool) to guide control decisions. The DST generates two population density profiles showing viable (red symbol) and total (black symbol) nymphs, plotted against crop development (which is predicted using daily temperature data). The magnitude of difference between the red and black dots is an indicator of mortality factors (such as parasitism) that may continue to suppress the SLW nymph population growth.

To use the DST:

1. Count the total number of large nymphs (3rd & 4th instar with red eyes) including nonviable for each leaf and determine the number of viable (healthy) nymphs. Ignore empty pupal cases.
2. Input the total and viable nymph counts for each leaf into the DST spreadsheet.
3. Enter the accumulated day degrees (base 12) for each check.

### Late season considerations

SLW management involving late season mass-immigration of adults into crops with open bolls should be based on (a) expected time to defoliated leaf drop, (b) lint contamination level, and (c) prior chemical use from a resistance management perspective (i.e. avoid multiple applications from the same mode of action). During defoliation, adult SLW will leave the crop, while falling leaves will take the nymphs with them to the ground. These displaced adult SLW relocate to nearby host crops and the nymph-based DST method will be of limited value for crops in this situation.

When selecting products for late season SLW management at or prior to the first pass of defoliant, consider potential efficacy and any control delay, residual impact and withholding periods (WHPs). If the risk of honeydew contamination is high, consider commencing defoliation earlier.


Also monitor for the presence of honeydew on lower leaves, particularly if there is open cotton, as this indicates some remedial action should be taken to prevent contamination of bolls. If defoliation is not due for 10-14 days and honeydew is collecting on the leaves, managing SLW numbers with a knockdown product may be prudent. If defoliation is 15-21 days away, consider using a knockdown spray in the first 7 days, followed by continued monitoring. Commencing defoliation earlier (at 50% open bolls) may be warranted if significant population resurgence is evident in the week post-treatment. If defoliation is more than 3 weeks away, the use of a systemic product with residual activity would be a prudent choice as it would have efficacy on both nymphs and adults within the canopy.

### Key beneficials

The tiny wasps, *Encarsia* and *Eretmocerus* are important parasitoids of SLW nymphs. Predators of nymphs include big-eyed bugs, pirate bugs, lacewing larvae, apple dimpling bugs, brown smudge bugs and ladybird beetles.

Natural enemies can play a vital role in the successful management of whitefly. Avoid early season use of broad-spectrum insecticides, particularly synthetic pyrethroids and organophosphates.

Find out how to identify parasitised SLW nymphs in this CottonInfo video [youtu.be/S00cedrGIQI](https://youtu.be/S00cedrGIQI)





Large 'red eye' nymphs (3rd or 4th instar) are used in the SLW decision support tool. (Tonia Grundy, Qld DAF)

## Selecting an appropriate insecticide

The insecticides registered for control of SLW are presented in Table 15. One of the challenges in SLW management is the timing of insecticide application as it's often a compromise between delaying treatment to reduce the risk of re-infestation (to avoid the cost of additional sprays), versus the need to control populations prior to open cotton (reducing the risk of sticky cotton).

Considerations when applying insecticides to control SLW include:

- **Impact on natural enemies** that can assist in controlling SLW that survive the insecticide.
- **Mode of action** of the insecticide (e.g. insect growth regulators may require up to several weeks to take full effect, so need to be timed to have completed control prior to open cotton).
- **Coverage and insecticide activity.** Post cut-out, insecticides with limited plant penetration and/or poor application may not provide adequate control.

### NOTES

**Sampling protocol** The minimum sample size per management unit is 30 leaves collected between nodes 11-14 from the terminal.

**Day degrees** Daily day degrees (DD) are calculated using the formula;  $DD = [(Max\ ^\circ C - 12) + (Min\ ^\circ C - 12)] \div 2$ .  
More accurate day degree calculations can be made using the online calculator at [www.csd.net.au/ddc](http://www.csd.net.au/ddc)

### PATHWAY

### CHEMISTRY

#### Green

The green pathway is a conservative indicator of low population density. Due to inherent variability associated with sampling, one or two data points slightly above the upper boundary of the green pathway does not constitute a 'breakout' and cause for concern that the population is increasing rapidly. These populations still require regular checking as a sharp increase in viable nymph density above the green pathway for two or more consecutive checks (spanning 7 days) is deemed a breakout population.

#### 1250-1450 DD

Insecticide use is not warranted for fields with low SLW densities.

#### 1600+ DD

If SLW are in the upper portion of the green pathway and you are targeting other pests, consider choosing products with modes of action effective against SLW activity:

- Diafenthiuron (vapour activity)
- Dinotefuran (high rate) (knockdown)
- Acetamiprid/emamectin (knockdown)
- Spirotetramat (use adjuvant as per label).

#### Amber

A population tracking within or above the bounds of the amber pathway for two (spanning 7 days) or more checks is likely to require control action. The exact timing of control should take into account crop stage and the insecticide mode of action. If the crop is at 1250-1450 DD, a slower acting Insect Growth Regulator (IGR) product would be an appropriate choice. These products have a low impact on beneficial insects that can continue to exert biological control after the efficacy of the insecticide residues have degraded. If the crop is nearing or has open bolls (1600+ DD), a faster acting insecticide would be a more appropriate choice. A follow up treatment for SLW may be necessary during the late stages of boll opening if natural enemies are disrupted and SLW populations recur. The difference between viable and total SLW nymphs will provide an estimate of natural mortality of which biological control is a component.

#### 1250-1450 DD

Partial selectivity

- Pyriproxyfen (IGR) (Refer to recommended regional 30 day window)
- Spirotetramat (use adjuvant as per label)
- Buprofezin

#### 1600+ DD

- Diafenthiuron (vapour activity)
- Dinotefuran (high rate) (knockdown)
- Acetamiprid/emamectin (knockdown)
- Spirotetramat (use adjuvant as per label).

#### Red

Red pathway populations represent the highest risk of lint contamination, but they are the simplest to manage because:

- High SLW numbers occurring well before boll opening allow time for control action without the immediate risk of lint contamination.
- Insecticides such as IGRs, that have a slower mode of action and reduce population recruitment, can be used to best effect prior to canopy closure thereby improving spray penetration.
- The efficacy provided by early IGR usage can then be reinforced by natural enemies that are largely undisrupted by these products, suppressing populations for much longer.

#### 1250-1450 DD

Partial selectivity

- Pyriproxyfen (IGR) (Refer to recommended regional 30 day window)
- Cyantraniliprole (if used twice in succession within 10-15 days)
- Spirotetramat (use adjuvant as per label)
- Buprofezin

#### 1600+ DD

- Partial selectivity to broad-spectrum
- Diafenthiuron (knockdown)
- Acetamiprid/emamectin (knockdown)
- Spirotetramat (control nymphs).

Products listed in order of selectivity based on Table 4 (pages 14-15).

### Resistance profile – SLW

When silverleaf whitefly was first identified in Australia in 1994 it already possessed resistance to many older insecticide groups, including pyrethroids (SP) and organophosphates (OPs). Selection of resistance in SLW populations can happen very quickly. Currently there is widespread, low level resistance to imidacloprid and bifenthrin. Rare instances of resistance to acetamiprid has been detected in previous seasons.

Resistance to pyriproxyfen is a significant concern for industry with increase in number of regions where resistance was detected with some reports of field failures. In an effort to maintain product efficacy Transgenic & Insect Management Strategy (TIMS) has recommended a continuation of the regional 30 day pyriproxyfen window and restrictions on pyriproxyfen use in open cotton. Rare and low level resistance to spirotetramat is of concern due to the dominant target site resistance mechanism. This means resistance could develop rapidly and reversal of resistance is unlikely.

Compliance with the IRMS (see pages 56-57) will ensure the products available for SLW control will remain efficacious into the future.

**ENSURE ONLY SINGLE APPLICATIONS OF PYRIPROXYFEN AND SPIROTETRAMAT OCCUR WITHIN A SEASON.**

### Overwintering habit

Whitefly does not have an overwintering diapause stage. It relies on alternative host plants to survive. Generation times are temperature dependent, slowing down during winter months. In central Queensland, the winter generation time is 40-45 days while in the Macintyre, Gwydir and Namoi valleys, generation time increases to 65-70 days. At Griffith in southern NSW, development during winter takes just over 100 days.

### Alternative hosts

The availability of a continuous source of hosts is the major contributing factor to a severe whitefly problem. Even a small area of a favoured host can maintain a significant whitefly population. Preferred weed hosts include: sowthistle, melons, bladder ketmia, native rosella, rhynchosia, vines (cow, bell and potato), rattlepod, native jute, burr gerkin and other cucurbit weeds, Josephine burr, young volunteer sunflowers, *Euphorbia* weeds, poinsettia and volunteer cotton.

In cotton growing areas the important alternative crop hosts are soybeans, sunflowers and all cucurbit crops. Spring plantings of these crops may provide a haven for SLW populations to build up in and then move into cotton. Autumn plantings of these crops may be affected by large



Adults and nymphs of SLW (left) and greenhouse whitefly (right). (Richard Lloyd, Qld DAF)

populations moving out of cotton. Do not plant cotton near preferred SLW host crops such as melons. Destroy crop residue from all susceptible crops immediately after harvest.

Minimising winter hosts, particularly sowthistle and volunteer cotton, is important in reducing the base population at the start of the cotton season. Smaller base populations will take longer to reach outbreak levels and reduce the likelihood that a particular field will need to be treated.

Further information – SLW Factsheet available: [www.cottoninfo.com.au](http://www.cottoninfo.com.au)

Qld DAF, Toowoomba

Jamie Hopkinson: 07 4529 4152 or 0475 825 340 (resistance enquiries).

Paul Grundy: 0427 929 172.

Qld DAF, Emerald – Richard Sequeria: 07 4991 0810 or 0407 059 066.



**Natural enemies play a vital role in the successful management of whitefly. Minimise impact on these beneficials by using thresholds for all pests and consider the impact on beneficials when selecting an insecticide.**

**TABLE 15: Control of silverleaf whitefly**

| Active ingredient                | MoA group | SLW resistance                                       | Overall impact on beneficials* | Comments#  |
|----------------------------------|-----------|--|--------------------------------|--|
| Paraffinic oil                   | No group  | Unknown  | Very low                       | Most effective when targeting low, early season populations. Apply in a minimum of 100 L/ha for ground applications. Use in combination with another registered insecticide when applying from the air. Multiple applications are more effective.  |
| Pyriproxyfen                     | 7C        | Widespread – low                                     | Very low                       | Ensure thorough coverage. Maximum 1 application per season.  |
| <i>Clitoria ternatea</i> extract | No group  | Not detected   | Low                            | Apply as indicated by field checks and pest presence. Ensure good coverage. Maximum 5 applications per season. Treatment effects may not be seen for 3 or more days. A repeat application may be required at 14-20 days if conditions favour pest development.   |
| Buprofezin                       | 17A       | Not detected   | Low                            | Application should be aimed at the early nymph stages of whitefly. Spray in sufficient volume to provide adequate penetration and coverage. Maximum 1 application per season. Thorough spray coverage is essential.  |
| Diafenthiuron                    | 12A       | Not detected   | Low                            | Provides suppression of SLW. Apply when population densities are low to moderate. Suppression may not be satisfactory once population densities exceed moderate infestation, or when high numbers of adults are invading from nearby fields. Maximum 2 applications per season.  |
| Afidopyropen                     | 9D        | Not detected   | Low                            | Registered to provide suppression of both adult and nymph stages of whitefly, however it is recommended to target the nymph stage.   |
| Cyantraniliprole                 | 28        | Not detected   | Moderate                       | Target early developing populations. Maximum 2 applications per season.  |
| Spirotetramat                    | 5         | Very rare  | Moderate                       | Use the higher rate when periods of high pest pressure or rapid crop growth are evident, and when crops are well advanced. Do not re-apply within 14 days. Spirotetramat controls nymphs and has a sterility effect on female SLW. Maximum 1 application per season. Ensure thorough coverage and use of adjuvant as specified on label.     |
| Emamectin benzoate/acetamiprid   | 6/4A      | Acetamiprid – very rare – low                        | Moderate                       | Use prior to heavy populations becoming established in the crop. Activity primarily on nymphs and therefore evidence of activity will be slower than typical contact insecticides. Use adjuvant as per label. Maximum 2 applications per season and use an insecticide from another mode of action between applications.                     |
| Dinotefuran                      | 4A        | Not detected   | Moderate                       | When mirids and SLW are present and SLW is at or above threshold levels, use SLW rate. Performance can be reduced in stressed crops, when senescing late season, or when pests are not actively feeding in the upper crop canopy. Maximum 2 applications per season and use an insecticide from another mode of action between applications. |
| Bifenthrin                       | 3A        | Widespread – low<br>Cross-resistance with other SPs. | Very high                      | The adult stage should be targeted. Do not spray crops with a high population of the juvenile stages. Thorough coverage of the crop canopy is essential. Maximum 1 application per season.   |


#For all control options ALWAYS refer to the label for instructions.

\*For more details about impact on beneficial insects (including bees), refer to Table 4 in this guide.

## Other insect pests

### Rutherglen bug (RGB) – *Nysius vinitor*

RGB's wide host range includes fruits, vegetables, oilseeds and grains, and many common weeds. Adult RGBs are often found in cotton, sometimes in large numbers, but generally do not feed and can't reproduce in the crop. Starving nymphs migrating from maturing sunflower or canola crops may damage border areas of adjacent cotton.



Adult Rutherglen bugs are 3-4 mm long; nymphs are reddish-brown. (Qld DAF)

Cage experiments on 5 day old bolls demonstrated that boll blemishes were superficial and caused no impact to the developing seeds or lint. Damage such as blackened stamens or darkened seeds is unlikely to be caused by RGB, and the crop should be monitored for other sucking pests.

### Leafhoppers (jassids)

Several species of leafhoppers are found in cotton. The wedge-shaped adults are quick to move when disturbed. Leafhoppers occasionally damage seedlings and cause stippling on mature leaves. Leafhoppers are unlikely to reduce yield unless in very high numbers (damage to >50% of the leaf surface) just before cut-out. Unnecessary sprays for leafhoppers may adversely affect beneficial species and flare other pests.



Leafhoppers are about 3 mm long and vary in colour from green to yellow-green to brown. (C.Mares)

### Redshouldered leaf beetle – *Monolepta australis*

An uncommon visitor to the majority of cotton-growing regions, redshouldered leaf beetle adults chew leaves, terminals, squares and the surface of cotton bolls. In the tropics (particularly after good rains), swarms of beetles can cause significant damage to patches of plants.



Adult redshouldered leaf beetles are about 6 mm long. (Joe Wessels)

## Bollworms


While this term is often used for *Helicoverpa*, several other species are also referred to as bollworms.

**Rough bollworm** (*Earias* spp.) larvae tunnel into squares, bolls or the main stem, destroying the growing point. While larvae can cause serious damage to unsprayed cotton, they are normally incidentally controlled by *Helicoverpa* sprays in conventional cotton and Bt toxins in Bollgard crops.



Rough bollworm larvae grow to about 18mm long and their favoured host is the weed, bladder ketmia. (Lewis Wilson)

**Pink spotted bollworm** (*Pectinophora scutigera*) occurs in coastal and central Queensland where it attacks cottonwood and bottle trees as well as cotton. **Pink bollworm** (*P. gossypiella*) occurs in cotton and other Malvaceae hosts in the Northern Territory and northern Western Australia. The larvae of both species tunnel into squares, flowers or bolls, with pupation occurring in bolls, stems or surface trash. Pink bollworm larvae can diapause while sheltering in bolls, lint, seed or the soil. While pink spotted bollworm does not have an overwintering diapause phase, larvae can survive by feeding on dry cotton seed in trash or cotton modules. |||



Pink spotted bollworm moths and larva (moth – right 12mm). (P. Room)



Damaged lint due to pink spotted bollworm feeding and the entry of boll-rotting fungi. (D. Ironside)

# SCHOLAR DOLLARS PROGRAM

Together with local growers,  
we are giving back  
to our local schools!

## What is Scholar Dollars?

Here at Growth Ag we've always prided ourselves on giving back to the communities in which we live & work. Our Scholar Dollar program is all about providing much needed funding support to regional schools by you, the grower supporting us, a regional manufacturer of sustainable quality products.

All growers need to do is complete a simple claim form. Alternatively, each participating school can co-ordinate the completion of the form and submit to Growth Ag directly.



## CASH BACK on Eligible Products

For every litre of the products listed below, cash will be given back to the nominated school along the following levels:

**B&B**  
Super-Fine  
Liquid Blood and Bone

Up to 499L - **5c/L**  
500L to 1999L - **10c/L**  
Over 2000L - **15c/L**

**SERO-**  
PESTICIDE

Up to 499L - **5c/L**  
500L to 1999L - **10c/L**  
Over 2000L - **15c/L**

**Zinc Xtra**  
Amino Acid Chelates

Up to 499L - **2c/L**  
500L to 999L - **5c/L**  
Over 1000L - **7c/L**

**STEADFAST**  
A Better Wetter

Up to 400L - **5c/L**  
Over 401L - **15c/L**

Maximum amount payable is \$5,000.00 per claim, per quarter.



**SCAN QR CODE**

For Claim Form & Further Details

**Head Office**  
02 6795 3787

accounts@growthag.com.au

**QLD - Fergus Tweedy**  
0447 502 520

ftweedy@growthag.com.au

**NSW - Martin Murray**  
0428 211 114

mmurray@growthag.com.au

growthag.com.au



# Insecticide Resistance Management Strategy (IRMS) for 2021-22

In every population of every pest species there is a small proportion of individuals with naturally occurring genetic resistance. These genes remain rare until they are selected for by a toxin, either from an applied pesticide or from within Bt cotton. Exposure controls the susceptible insects, leaving behind the resistant individuals, which can then build up as a larger proportion of the overall population.

Over-reliance on an insecticide can lead to an increase in the proportion of resistant individuals to the point where the insecticide fails to provide satisfactory control. This simple scenario is obviously more complex in a field situation as products applied not only select for resistance in the target pest, but also in other pests present.

The IRMS aims to assist users to:

- Lower the risk of inadvertent selection of resistance in pests that are not the primary target of the insecticide application.
- Delay the evolution of pest resistance to key chemical groups, by minimising the survival of individuals with resistance.
- Manage entrenched resistance problems, such as the now widespread resistance in silverleaf whitefly (SLW) to pyriproxyfen.

The cotton industry IRMS seeks to manage the risk of resistance in all major pests of cotton including aphids, mites, SLW and *Helicoverpa* spp. in both conventional and Bt cotton (note that in this document, the term 'insecticide' refers generally to pesticides used for insect or mite control). Additional resistance management requirements are also in place for managing the risk of *Helicoverpa* developing resistance to Bt cotton (pages 61-70).

**The IRMS includes all active ingredients commercially available for use in cotton at the time of publication and should be consulted for EVERY insecticide/miticide decision in both Bt and conventional cotton.**

## Key changes for the 2021-22 cotton season

- **Fipronil bee risk.** While bees are susceptible to many insecticides used on cotton, fipronil risk to managed hives has been highlighted due to the extended residue risk. The industry recommends fipronil should not be used on flowering crops. Refer to label for statement about bees.
- **Incidental resistance from fall armyworm spraying.** Use of insecticides for fall armyworm in other crops may inadvertently select for resistance in cotton pests. Apply resistance management principles to FAW management.

## Principles underlying the IRMS

- **Monitor pest and beneficial populations.** Consider species as well as abundance.
  - **Monitor fruit retention and determine if losses are likely to be due to pest damage or environmental factors.**
  - **Use recommended thresholds for all pests (do not use 'insurance' sprays).**
  - **Aim to use the most selective insecticide options first, delaying the use of broad-spectrum insecticides for as long as possible.**
  - **Comply with all directions for use on product labels.**
  - **Avoid repeated applications of similar products, even when targeting different pests. Rotate between mode of action (MoA) groups.**
  - **Do not respray an apparent failure with the same product or another product from the same insecticide group.**
  - **Do not exceed the maximum recommended use limits.**
  - **Control weeds and cotton volunteers in fields and around the farm all year to minimise alternate hosts.**
  - **Pupae bust as soon as possible after harvest.**
- **Continuation of pyriproxyfen window.** Resistance to pyriproxyfen in SLW is a significant concern for industry with an increase in the number of regions where resistance was detected and field failures reported. In an effort to maintain product efficacy, TIMS has recommended a continuation of the regional 30 day pyriproxyfen window and restrictions on pyriproxyfen use in open cotton. After consultation, appropriate window dates will be published on CottonInfo and Cotton Australia websites. Limit pyriproxyfen use to no more than ONE application per season.
  - **SLW products.** With the move from the SLW threshold matrix to the DST, SLW products are listed against the pathways as examples only. The SLW management fact sheet (available from the CottonInfo website) should be consulted when making decisions regarding SLW products. Rare and low level resistance to spirotetramat has again been detected. The dominant target site resistance mechanism means resistance can develop rapidly and reversal of resistance is unlikely. With pyriproxyfen resistance remaining a concern, and increased reliance on spirotetramat, TIMS has recommended usage be limited to one application per season. The 'double knock' use for mealybug – 2 applications 14 days apart is an allowable exception.
  - **Northern IRMS.** To avoid confusion, the previous 'Northern Regions' IRMS has been renamed Central Qld IRMS. An IRMS strategy has not been yet been developed for cotton regions in Northern Australia (NQ, NT, WA).

## IRMS in grains

Resistance management strategies have also been developed for key grains pests and should be used in conjunction with the Cotton IRMS. They are available at [ipmguidelinesforgrains.com.au/ipm-information/resistance-management-strategies](http://ipmguidelinesforgrains.com.au/ipm-information/resistance-management-strategies)

For other resistance management strategies and a list of insecticide MoA groups visit CropLife Australia at [www.croplife.org.au](http://www.croplife.org.au)





# A command performance every time.

In farming, incremental improvements in efficiency make all the difference. That's why Ag Leader's DirectCommand gives growers the edge. Designed to deliver unparalleled accuracy and efficiency, it features the latest technology developed to give growers complete control while minimising wastage.

#### FEATURES INCLUDE:

**AUTOSWATH** - increased control with minimal wastage

**PRESSURE FALLBACK** - ensures correct application rate at all times

**ACCURATE FLOW AND RATE CONTROL** - ensure the right rate in the right place

**ISOBUS AND SERIAL CONNECTIVITY** - highly compatible and flexible

**SMARTREPORTS** - generate as applied maps right from the display.



To find out more, contact Doug Amos, sales manager, on 0428 409 885.

[agleader.com](http://agleader.com)

**Ag Leader**<sup>®</sup>

# Insecticide Resistance Management Strategy 2021/22

## Best Practice Product Windows and Use Restrictions to Manage Insecticide Resistance in Insect Pests of Australian Cotton

**CENTRAL QLD REGIONS:** Belyando, Callide, Central Highlands, Dawson

INCREASING

SELECTIVITY

DECREASING

CONSIDER IMPACT ON BENEFICIALS & BEES (TABLE 4 CPMG)

| Stage 1  | Stage 2                | Stage 3   | Stage 4                                    |
|--|------------------------|---|--|
|  | 1-Nov                  | 1-Dec   | 1-Jan                                      |
| <b>Helicoverpa viruses (Virus)</b>   |                        |   |  |
| Pirimicarb Group 1A  |                        |   | Note 1                                     |
| Paraffinic Oil (Canopy, Biopest)   |                        |   |  |
|  | Pyriproxyfen Group 7C  |   | Use an alternative from open cotton        |
| Sero-X   |                        |   |  |
| Etoxazole  |                        |   |  |
| Buprofezin (Applaud) Group 16  |                        |   | Note 3, 9                                  |
|  | GROUP 28: Max 4/season | Chlorantraniliprole (Altacor) Group 28                                |  |
| Afidopyropen (Versys) Group 9D   |                        |   |  |
| Spinetoram (Success Neo) Group 5   |                        |   |  |
| start date = canopy closure  |                        | Diafenthiuron Group 12A   |  |
| Pymetrozine Group 9B   |                        |   |  |
| Indoxacarb Group 22A   |                        |   | Dec-31                                     |
|  | GROUP 28: Max 4/season | Cyantraniliprole (Exirel) Group 28                                    |  |
| Spirotetramat (Movento) Group 23   |                        |   |  |
| Fonicamid (MainMan) Group 29   |                        |   |  |
| Abamectin Group 6  |                        |   | GROUP 6: Max 3 / season                    |
| Emamectin Group 6  |                        |   |  |
| start date = squaring  |                        | Propargite Group 12C  |  |
| Amitraz Group 19   |                        |   |  |
| Sulfoxaflor (Transform) Group 4C   |                        |   |  |
| Fipronil Group 2B  |                        | Do not apply to flowering crops                                       |  |
| Neonicotinoids (acetamiprid, clothianidin, dinotefuran, imidacloprid, thiamethoxam) Group 4A |                        |   |  |
|  |                        | Chlorantraniliprole + Thiamethoxam (Voliam Flexi) Group 4A + Group 28 |  |
| Acetamiprid + Emamectin (Skope) Group 4A + Group 6   |                        |   |  |
| Phorate  | Note 1                 | Feb-01  | Carbamates (methomyl, thiodicarb) Group 1A |
|  |                        |   | Dimethoate Group 1B                        |
|  |                        |   | OPs (chlorpyrifos, methidathion) Group 1B  |
|  |                        |   | Synthetic pyrethroids Group 3A             |

**Note 1** If a phorate side dressing is used at planting then do not use a pirimicarb or dimethoate first foliar spray as there is cross resistance between them all. Dimethoate use will select catastrophic pirimicarb resistance in aphids so do not use pirimicarb and dimethoate in the same field.

**Note 2** Failures of neonicotinoids against aphids have been confirmed. DO NOT follow a neonicotinoid seed treatment with a foliar neonicotinoid when aphids are present. If there is an alternative do not follow a neonicotinoid with sulfoxaflor.

**Note 3** Refer to CottonInfo SLW fact sheet for additional guidance [www.cottoninfo.com.au/publications/insects-managing-silverleaf-whitely-australian-cotton](http://www.cottoninfo.com.au/publications/insects-managing-silverleaf-whitely-australian-cotton)

**Note 4** Imidacloprid (neonicotinoid) resistance in cotton seedling thrips is likely. If resistance is suspected, phorate is an appropriate at-planting alternative. Consider non-neonicotinoid alternatives for first foliar spray.

**Note 5** Additional applications can be made if targeting *Helicoverpa* moths using Magnet.

**Note 6** High SP resistance is present in *Helicoverpa armigera* populations. Expect field failures. Low resistance is present in SLW.

**Note 7** Addition of abamectin to mirid sprays has caused high level resistance in mites. Base miticide decisions on thresholds only.

**Note 8** Rare and low level resistance to spirotetramat has been detected in some regions. The dominant target site resistance mechanism means resistance can develop rapidly and reversal of resistance is unlikely.

**Note 9** Where buprofezin usage is targeting mealybugs, maximum of 2 sprays per field.

**Note 10** Use of insecticides for fall armyworm in other crops may inadvertently select for resistance in cotton pests. Apply resistance management principles to FAW management.

**Note 11** Refer to label statement about bees.

**ALWAYS FOLLOW LABEL DIRECTIONS**  
**CONSIDER IMPACT ON BENEFICIALS & BEES; (TABLE 4, COTTON PEST MANAGEMENT GUIDE)**  
**IMPLEMENT AN IPM STRATEGY INCLUDING GOOD FARM HYGIENE AND CONTROL OF OVERWINTER HOSTS.**  
**PUPAE BUST CONVENTIONAL COTTON CROPS AFTER HARVEST**

|   |                                  |
|---|----------------------------------|
| <span style="background-color: #c8e6c9; border: 1px solid black; display: inline-block; width: 15px; height: 10px;"></span> | Avoid repeated use of same group |
| <span style="background-color: #ffe0b2; border: 1px solid black; display: inline-block; width: 15px; height: 10px;"></span> | No more than 1 application       |
| <span style="background-color: #bbdefb; border: 1px solid black; display: inline-block; width: 15px; height: 10px;"></span> | No more than 2 applications      |
| <span style="background-color: #c5cae9; border: 1px solid black; display: inline-block; width: 15px; height: 10px;"></span> | No more than 3 applications      |
| <span style="background-color: #e91e63; border: 1px solid black; display: inline-block; width: 15px; height: 10px;"></span> | No more than 4 applications      |

Note 1  
 Note 3  
 Note 3, 9  
 Note 10  
 Note 3  
 Note 3  
 Note 10  
 Note 3  
 Note 3, 8  
 Note 7  
 Note 2  
 Note 11  
 Note 2, 3, 4, 11  
 } Consider risk to each group  
 Note 5  
 Note 1  
 Note 3, 6

# Insecticide Resistance Management Strategy 2021/22

## Best Practice Product Windows and Use Restrictions to Manage Insecticide Resistance in Insect Pests of Australian Cotton

**CENTRAL & SOUTHERN REGIONS:** Balonne, Bourke, Darling Downs, Gwydir, Lachlan, Upper & Lower Namoi, MacIntyre, Macquarie, Murrumbidgee, Murray

**INCREASING**
**SELECTIVITY**
**DECREASING**
**CONSIDER IMPACT ON BENEFICIALS & BEES (TABLE 4 CPMG)**

| Stage 1  | Stage 2                                | Stage 3                                    | Stage 4    |
|--|--|--|------------|
|  | 15-Dec                                 | 15-Jan                                     | 15-Feb     |
| <b>Helicoverpa viruses (Virus)</b>   |  |  |            |
| Pirimicarb Group 1A  |  |  | Note 1     |
| Paraffinic Oil (Canopy, Biopest)   |  |  |            |
| Pyriproxyfen Group 7C <b>Regional 30 day window</b>  |  |  | Note 3, 8  |
| Use an alternative from open cotton  |  |  |            |
| Sero-X   |  |  |            |
| Etoxazole  |  |  |            |
| Buprofezin (Applaud) Group 16  |  |  | Note 3, 10 |
| GROUP 28: Max 4/season   | Chlorantraniliprole (Altacor) Group 28 |  |            |
| Note 11  |  |  |            |
| Afidopyropen (Versys) Group 9D   |  |  |            |
| Spinetoram (Success Neo) Group 5   |  |  |            |
| start date = canopy closure  |  | Diafenthiuron Group 12A                    |            |
| Note 3   |  |  |            |
| Pymetrozine Group 9B   |  |  |            |
| Indoxacarb Group 22A   |  |  | Jan-31     |
| GROUP 28: Max 4/season   | Cyantraniliprole (Exirel) Group 28     |  |            |
| Note 11  |  |  |            |
| Spirotetramat (Movento) Group 23   |  |  |            |
| Note 3, 9  |  |  |            |
| Flonicamid (MainMan) Group 29  |  |  |            |
| Abamectin Group 6  |  | GROUP 6: Max 3 / season                    |            |
| Note 7   |  |  |            |
| Emamectin Group 6  |  |  |            |
| start date = squaring  |  | Propargite Group 12C                       |            |
| Amitraz Group 19   |  |  |            |
| Sulfoxaflor (Transform) Group 4C   |  |  |            |
| Note 2   |  |  |            |
| Fipronil Group 2B  |  | Do not apply to flowering crops            |            |
| Note 12  |  |  |            |
| Neonicotinoids (acetamiprid, clothianidin, dinotefuran, imidacloprid, thiamethoxam) Group 4A |  |  |            |
| Note 2, 3, 4, 12   |  |  |            |
| Chlorantraniliprole +Thiamethoxam (Voliam Flexi) Group 4A + Group 28                         |  |  |            |
| } Consider risk to each group  |  |  |            |
| Acetamiprid + Emamectin (Skope) Group 4A + Group 6   |  |  |            |
| Phorate  | Note 1                                 | Carbamates (methomyl, thiodicarb) Group 1A |            |
|  |  | Dimethoate Group 1B                        |            |
|  |  | OPs (chlorpyrifos, methidathion) Group 1B  |            |
|  |  | Synthetic pyrethroids Group 3A             |            |
|  |  | Note 5                                     |            |
|  |  | Note 1                                     |            |
|  |  | Feb-01                                     |            |
|  |  | Note 3, 6                                  |            |

- Avoid repeated use of same group
- No more than 1 application
- No more than 2 applications
- No more than 3 applications
- No more than 4 applications

**Note 1** If a phorate side dressing is used at planting then do not use a pirimicarb or dimethoate first foliar spray as there is cross resistance between them all. Dimethoate use will select catastrophic pirimicarb resistance in aphids so do not use pirimicarb and dimethoate in the same field.

**Note 2** Failures of neonicotinoids against aphids have been confirmed. DO NOT follow a neonicotinoid seed treatment with a foliar neonicotinoid when aphids are present. If there is an alternative do not follow a neonicotinoid with sulfoxaflor.

**Note 3** Refer to CottonInfo SLW fact sheet for additional guidance [www.cottoninfo.com.au/publications/insects-managing-silverleaf-whitefly-australian-cotton](http://www.cottoninfo.com.au/publications/insects-managing-silverleaf-whitefly-australian-cotton).

**Note 4** Imidacloprid (neonicotinoid) resistance in cotton seedling thrips is likely. If resistance is suspected, phorate is an appropriate at-planting alternative. Consider non-neonicotinoid alternatives for first foliar spray.

**Note 5** Additional applications can be made if targeting *Helicoverpa* moths using Magnet.

**Note 6** High SP resistance is present in *Helicoverpa armigera* populations. Expect field failures. Low resistance is present in SLW.

**Note 7** Addition of abamectin to mirid sprays has caused high level resistance in mites. Base miticide decisions on thresholds only.

**Note 8** Resistance to pyriproxyfen is now widespread. To avoid complete loss of product efficacy, adhere to the 30 day regional window. Limit pyriproxyfen use to no more than 1 application per season.

**Note 9** Rare and low level resistance to spirotetramat has been detected in some regions. The dominant target site resistance mechanism means resistance can develop rapidly and reversal of resistance is unlikely.

**Note 10** Where buprofezin usage is targeting mealybugs, maximum of 2 sprays per field

**Note 11** Use of insecticides for fall armyworm in other crops may inadvertently select for resistance in cotton pests. Apply resistance management principles to FAW management.

**Note 12** Refer to label statement about bees

**ALWAYS FOLLOW LABEL DIRECTIONS**

**CONSIDER IMPACT ON BENEFICIALS & BEES; (TABLE 4, COTTON PEST MANAGEMENT GUIDE)**

**IMPLEMENT AN IPM STRATEGY INCLUDING GOOD FARM HYGIENE AND CONTROL OF OVERWINTER HOSTS.**

**PUPAE BUST CONVENTIONAL COTTON CROPS AFTER HARVEST**

## Region

The IRMS aims to reduce the chance that highly mobile pests would be repeatedly exposed to the same insecticide group by limiting the timing of insecticide availability. The two strategies accommodate the different growing seasons: from southern Queensland through to southern NSW, and in Central Queensland where the different stage dates account for early planting and quicker crop development.

A specific IRMS strategy has not yet been developed for cotton regions in northern Australia. Apply the IRMS principles in these regions.

## Stage

The dates shown on the strategy charts indicate the start of each stage. Specific start &/or end dates are listed for individual insecticides and miticides that start or end outside window boundaries. All windows start at 12 am.

## Selectivity

The products in the IRMS are listed in order of decreasing selectivity. For all pest species, aim to use the most selective option, delaying or avoiding the use of broad-spectrum insecticides.

## Use restrictions

Colours represent the maximum number of applications per crop per season for any given product or product group. Additional restrictions can be found to the right of the table, with links to specific footnotes.

# More about resistance

## The role of IPM, refuges and pupae-busting

IPM principles promote the conservation of beneficials to help prevent over-reliance on chemical control that leads to insecticide resistance and renders insecticidal control options ineffective. The benefits of preserving beneficials are particularly important for mites and SLW where there is increasing concern about resistance to key products. Early season pest control decisions can lead to flaring of these pests later in the season. Aim to preserve beneficials through the use of thresholds for all pests and consider the impact on beneficials when selecting insecticides. Refer to the *IPM* section on page 5 for more information.

Growing refuge crops is a pre-emptive resistance management strategy implemented to retard the evolution of field-scale resistance to Bt cotton (see the Bollgard RMP for more information). While the dilution effect of refuges is far less effective when insecticide resistance is already widespread and prevalent in the field population (such as with synthetic pyrethroids), there may be some benefit from the unsprayed refuge options for new chemistries.

Pupae busting is an effective, non-chemical method of preventing resistance carryover in diapausing *H. armigera* pupae. In conventional cotton, it is recommended that pupae busting be conducted as soon as possible after harvest. For Bollgard crops, refer to the pupae-busting requirements for your region in the current Resistance Management Plan (RMP).

## Cross-resistance versus multiple resistance

Cross-resistance occurs when selection for resistance against one pesticide also confers resistance to another pesticide, either from the same mode of action or a different group. For example, the mechanism for pirimicarb resistance (Group 1A) in aphids also gives resistance to dimethoate (Group 1B). Cross-resistance is important as it means that a pest could be resistant to a chemical to which it has never been exposed.

Multiple resistance means that an insect is resistant to more than one MoA group. For instance, *H. armigera* can have metabolic resistance to synthetic pyrethroids (Group 3A) and nerve insensitivity to organophosphates (Group 1B).

The development of both cross-resistance and multiple resistance can be minimised by following the IRMS. For example, in the strategy for aphids, there is a break between the use of pirimicarb and dimethoate during which other chemistries should be used to minimise the number of pirimicarb-resistant aphids being exposed to dimethoate.

## What about insecticide mixtures?

Tank mixtures are high-risk and a controversial strategy for managing resistance. They can undermine the IRMS by repeatedly selecting for resistance to their common components and across multiple chemical groups. When mixtures are used frequently, it also becomes difficult to determine whether each component is contributing equally to efficacy.

Some mix partners provide more than additive kill (synergism), but this is not always the case. As a general rule, mixtures are unnecessary in situations where individual products provide adequate control.

For mixtures to be effective, their components should:

- Be equally persistent;
- Have different modes of action (and not be subject to the same routes of metabolic detoxification);
- Not be subject to resistance in the majority of the target pest population; and,
- Be tank-mix compatible.

**It is illegal to use rates above those recommended on the label of an insecticide either alone or in mixtures.** Efficacy will not always improve at rates above the highest label rate or if two insecticides of the same chemical group are applied as a mixture.

## What to do if you suspect a spray failure

Carefully assess the situation (the presence of live pests does not necessarily indicate insecticide failure). What is the insecticide's mode of action? Has it been given enough time to work? Was it applied correctly and in the right conditions?

Stomach poisons such as thiodicarb, foliar Bt, NPV and indoxacarb may not give maximum control until 5-7 days after application. Similarly, propargite, abamectin and diafenthiuron are slow acting and may take 7-10 days or longer to achieve maximum control. Insect growth regulators (IGRs) such as pyriproxifen can take up to three weeks. In some cases pests may have ceased feeding following a treatment even though infestation levels remain high, meaning little if any economic damage to the crop.



# Set up for success.

SURESPEED™

SUREFORCE™

SUREDRIIVE™

SURESTOP™

INCOMMAND DISPLAY™

To find out more, contact Doug Amos,  
sales manager, on 0428 409 885.

[agleader.com](http://agleader.com)

Little efficiencies yield the biggest results. That's why Ag Leader's SeedCommand technology makes all the difference. Each product in the range is designed to give you more control, allowing you to plant faster with greater accuracy and consistency – maximising your seed and your time.

Thanks to the InCommand Display, next-level visibility and control over your entire operation is at your fingertips. Plus, the Ag Leader range is compatible with most planter makes and models, giving you the flexibility to adapt and enhance your setup on your terms.

When you plant with Ag Leader, you're perfectly placed for the season ahead.

**Ag Leader®**

A wide range of variables influence insecticide efficacy, including species complex, population density and age, crop canopy structure, application timing, the application method, carrier and solution pH (and their effects on coverage and the insecticide dose delivered to the target), environmental conditions, assessment timing and insecticide resistance expressed in the pest population. The interaction of all these factors determines the outcome of a pesticide application. While it is not possible to optimise all of these variables all of the time, the more compromises that are made, the greater likelihood of unsatisfactory efficacy.

It is also important to maintain realistic expectations of the efficacy that can be achieved. Satisfactory control of medium and large *Helicoverpa* larvae is unlikely, regardless of the insecticide treatment used. If a field failure is suspected to be due to insecticide resistance, contact the relevant researcher for advice on how to collect and send in samples:

- *Helicoverpa*, mites and aphids: Lisa Bird 02 6763 1128.
- Whitefly: Jamie Hopkinson 07 4529 4152.

Sending samples for testing can help confirm or rule out resistance as the cause of the spray failure and is an important part of assessing the presence of resistance across the industry.

**After any spray failure, do not follow with an application that includes a product from the same insecticide MoA group.**

## Note: the IRMS includes multiple pests

The IRMS aims to minimise selection across consecutive generations of the pest, therefore pest life cycles determine the length of the 'windows'. As the life cycles of *Helicoverpa* and various sucking pests are very different, the strategy for one will not manage resistance for the other.

### *Helicoverpa*

While ideally the window length would be 42 days (average time from egg to moth), most chemicals are restricted to windows of 1-2 generations to account for the practicalities of pest control. To counteract this compromise there are additional restrictions on the maximum number of applications for each chemical group.

### Mites and aphids

The resistance strategy for short life cycle pests depends on rotation between different MoA groups to avoid selection over successive generations. Non-consecutive use of chemistries is particularly important for aphids as they reproduce asexually (all offspring from a resistant aphid will be resistant). There are also restrictions on the maximum number of uses for individual products and chemical groups to further encourage rotation of chemistries.

### Mirids

It is possible that resistance could develop and the industry has begun resistance monitoring in mirids as a precaution. Many of the products registered for mirid control in cotton include registration for the control of other pests, so it is critical that mirid control decisions also consider sub-threshold populations of other pests that are present. Using dimethoate for mirid control can inadvertently select for both dimethoate and pirimicarb resistance in aphids. Use of neonicotinoid products for mirid control (particularly if also used as a seed treatment) can inadvertently select for resistance in aphids. When selecting an insecticide, consider the options remaining for subsequent aphid control, if required.

## Silverleaf whitefly (SLW)

The IRMS includes all commercially available products registered for use against SLW in cotton. Refer to CottonInfo ([www.cottoninfo.com.au](http://www.cottoninfo.com.au)) or Cotton Australia ([cottonaustralia.com.au](http://cottonaustralia.com.au)) websites for details of the current Cotton Grower Association (CGA)-nominated voluntary pyriproxyfen application window.

## In-season troubleshooting

Ratification of the IRMS prior to the start of each season is the responsibility of Cotton Australia's Transgenic and Insect Management Strategies (TIMS) Committee. A Troubleshooting sub-committee is empowered to act on TIMS' behalf during the cotton season to respond to emergency requests to vary the IRMS for a district (or part of a district) for the remaining season. A request can be initiated by any grower or consultant, but will not be considered unless it is presented with clear evidence of discussion and majority support at a local level. For further information contact Cotton Australia (02 9669 5222).



# Bollgard 3 Resistance Management Plan (RMP)

**R**esistance is the greatest threat to the continued availability and efficacy of Bt cotton in Australia. The RMP was established as a pre-emptive measure to mitigate the risks of resistance developing to any of the insecticidal proteins contained in Bt varieties, and aims to prevent field level changes in resistance.

Crops with multiple toxins should be relatively robust, especially if the 'stacked' toxins kill insects in different ways. But the resilience of a stack depends on how well each toxin controls larvae and the levels of resistance to each toxin at the time that the technology is introduced. Bollgard 3 cotton contains Cry1Ac, Cry2Ab and Vip3A, however with over 90% of the industry using this technology, it is not impossible that *Helicoverpa* could adapt. It is therefore imperative that the RMP is implemented effectively by all growers to ensure the longevity of the product.

## Current status of Bt resistance in Australia

While field resistance to Cry1Ac is still relatively low, the baseline frequency of Cry2Ab and Vip3A resistance genes for F1 screening in *H. armigera* and *H. punctigera* populations is currently about 1%. Additionally, the expression of Cry1Ac and Vip3A varies throughout the season and is not always high enough for each insecticidal protein to individually control susceptible insects. None of the resistance genes identified confer cross-resistance to the other insecticidal proteins.

The data for current frequencies is sourced from Bayer's 2020-21 monitoring program. Unless otherwise stated, other information around trends in resistance frequencies draws on data collected over time from CSIRO and Bayer. The frequencies of these recessive resistance genes are currently not at a level that impacts performance of the technology (there is currently no resistance to Bollgard 3 that affects its performance against *H. armigera* or *H. punctigera*).

**Cry1Ac:** A gene present in field populations of *H. armigera* has the potential to confer high-level resistance to Cry1Ac. This largely recessive gene occurs at a low frequency – probably less than 5 in 10,000 (<0.05%).



*H. armigera*. (Melina Miles, Qld DAF)

It also has a high fitness cost (i.e. resistant individuals develop slowly and are more likely to die) but this disadvantage is unlikely to greatly impact on the development of resistance.

At least 2 different forms of resistance to Cry1Ac have been isolated from field populations of *H. punctigera*. Frequency of both is 1 in 1000 (0.1%).

**Cry2Ab:** A recessive gene that confers high level resistance to Cry2Ab is present in field populations of *H. armigera*. The mechanism is likely to be an alteration of a binding site in the gut of the insect. The current estimate of resistance frequency is approximately 1 in 100 (1%) or less.

A recessive gene that confers high level resistance to Cry2Ab is present in field populations of *H. punctigera*, and demonstrates the same broad characteristics as the *H. armigera* Cry2Ab-resistance, and also occurs at a frequency of approximately 1%.

**Vip3A:** At least two different types of Vip3A resistance have been identified in both *H. armigera* and *H. punctigera* populations. Data by CSIRO and Bayer suggests that the frequency of all Vip3A resistances is around 1 in 100 (1%).

Both CSIRO and Bayer data sets show that there is no significant difference in the frequencies of Cry2Ab and Vip3A resistance alleles over the longer term. Irrespective of changes through time, the frequencies of Cry2Ab and Vip3A in *H. armigera* and *H. punctigera* are higher than expected and of concern. Until the widespread adoption of Bt cotton there was presumably little exposure of *Helicoverpa* to this toxin, and therefore minimal selection for resistance. Although the Cry2Ab and Vip3A toxins are present in some Australian soils, they are not common compared to the Cry1Ac toxin, yet resistance to the Cry1Ac toxin in *Helicoverpa* is rare.

*H. punctigera* has the capacity to develop resistance to insecticide sprays but it has been presumed that any resistance selection in cotton regions was kept in check by dilution from susceptible immigrants from central Australia each spring. There may be some recent changes to the ecology of *H. punctigera* that could impact on their ability to develop resistance including a greater tendency to overwinter in cotton regions and less immigration of inland individuals than in the past due to low rainfall inland.

## How is resistance monitored?

CSIRO, with funding support from CRDC, implemented a monitoring program from 2003-18, that adapted to new varieties and circumstances each season. Bayer also invests in an annual program monitoring field populations of *Helicoverpa* spp. for resistance of all three proteins contained in Bollgard 3 cotton (Cry1Ac, Cry2Ab and Vip3A), providing an early warning of the onset of resistance. The programs have been used by industry to make decisions about the need to modify the RMP each season to ensure its ongoing effectiveness.

Originally, all resistance monitoring used F2 screens (which involve testing the grandchildren of pairs of moths raised from eggs collected from field populations). This method detects all previously isolated and potentially new types of resistances but takes about 10 weeks and is very labour-intensive, so protocols were developed to halve the testing time. F1 screens involve testing the offspring of single-pair matings between a moth from resistant strains maintained in the laboratory and a moth raised from eggs collected from field populations. This method works as the various isolates of Cry2Ab, Cry1Ac and Vip3A detected to date are of the same kind. Both methods are still utilised to measure known forms of resistance and to detect any new forms present in *Helicoverpa* populations.

Although some individuals that carry two copies of the resistance genes (homozygotes) have been detected for all toxins in both species, the vast majority of detected resistant individuals carry only one copy (heterozygotes) and therefore are controlled by Bollgard 3. Any homozygote (resistant to one of the proteins in Bollgard 3 cotton) is also controlled by Bollgard 3.

A molecular tool for Bt resistance testing using historical samples is currently under development, supported by CSIRO, CRDC and Bayer.

## Elements of the Bollgard 3 RMP

The RMP imposes limitations and requirements for management on farms that grow Bollgard 3 cotton. Its five elements are:

- Mandatory growing of refuges;
- Control of volunteer and ratoon plants;
- Planting window or planting restrictions;
- Restrictions on the use of foliar Bt; and,
- Mandatory cultivation of crop residues.

In theory the interaction of all of these elements should effectively slow the evolution of resistance. While there have been no reported field failures of Bt cotton due to resistance in Australia, the higher than expected baseline frequency of Cry2Ab and Vip3A is a major concern, and it is imperative that all Bollgard 3 users are responsible stewards of the technology. In particular, it is critical that close attention is paid to managing associated refuges, and that if required, effective pupae busting occurs in a timely fashion.

Bayer and the TIMS Bt Technical Panel will continue to work together to annually assess new information on resistance frequencies in *Helicoverpa* species and provide background information and recommendations for the Cotton Australia convened TIMS Committee.

For full details of how to practically implement the RMP, please refer to the RMP document, your Technology Service Provider (TSP) or your Bayer Territory Business Manager.

## RMP tactics

### 1. Using refuges

While the use of planting windows and use of two or three Bt genes are aimed at reducing selection pressure for Bt resistance, refuge crops are used to balance or counter the selection that will still occur. Refuge crops assist genetic dilution by generating significant numbers of susceptible moths (SS) that have not been exposed to selection pressure from the Bollgard 3 insecticidal proteins. Moths produced in refuge crops disperse to form part of the local mating population where they reduce the chance that resistant moths (RR) emerging from Bollgard 3 crops will meet and mate. The offspring from matings between one resistant and one susceptible moth carry one gene from each parent (RS) and are referred to as heterozygotes. In the cases of Bt resistance identified so far, heterozygotes and homozygotes are still controlled by Bt cotton. **It is crucial that the timing of the production of moths from refuges matches that of Bollgard 3 crops.**

Refuge crop areas must be in close proximity to the Bollgard 3 crop(s) to maximise the likelihood that moths emerging from the Bollgard 3 crop will mate with susceptible moths from the nearby refuge crop. During the summer cropping season a significant part of the *Helicoverpa* population may move only a few kilometres, depending on the mix of crops and their attractiveness at the time of moth emergence. Therefore the best location for a refuge crop is within 2 km of a Bollgard 3 crop, with a road or drain



*H. punctigera*. (Melina Miles, Qld DAF)

between the two to minimise the movement of large larvae from the refuge to the crop.

Managing resistance is a population level activity, and every refuge makes an important contribution to both the overall RMP for the valley and, because *Helicoverpa* disperse widely, for the whole industry. It is imperative that all refuges produce their quota of susceptible (SS) moths. Bayer audits the quality of refuges on every farm that grows Bollgard 3 to ensure that they are well maintained and effective.

The relative sizes of refuge crops required are based on *Helicoverpa* moth emergence for different crop types, and use a refuge of 10% unsprayed cotton as an initial reference point for Bt crops. On average, pigeon pea produces twice as many moths as the same area of unsprayed cotton, hence only a 5% refuge of pigeon pea is required.

Refuge requirements for Bollgard 3 have been reduced to 5% unsprayed cotton and 2.5% pigeon pea, representing the industry's confidence in the robustness of a 3 gene product in managing resistance risk combined with an industry commitment to improve the quality of refuges. Note that Bollgard 3 refuges must be a minimum of 0.5 ha and at least 24 m wide to account for possible insecticide drift onto the refuge.

The productivity of refuges will vary considerably across regions and seasons. Improving the production potential of each individual refuge is an integral component of the RMP. Growers must ensure that on farm refuge management is a priority. Looking after refuges, including nutrition, weed control, timely irrigation and all factors that make the refuge 'attractive' to female moths laying eggs, is the key to ensuring that they are effective. Guidelines on refuge management are provided in the RMP and the following unsprayed pigeon pea refuge agronomy guide (page 71).

Note that the term 'unsprayed' encompasses all management activities which are likely to reduce the survival of *Helicoverpa* in the refuge. For example, insecticides with activity against *Helicoverpa*; inundative beneficial releases; or, food sprays to attract natural enemies; cannot be used.

### 2. Controlling volunteers and ratoons

It is important to prevent both the establishment of conventional cotton in Bollgard 3 fields and the presence of Bollgard 3 cotton volunteer plants in a conventional crop or unsprayed refuge. Larger larvae are less susceptible to Bollgard technology. Heterozygous (RS) larvae that emerge from eggs laid on conventional cotton may grow and during their development move onto Bollgard 3 cotton plants. In this way RS larvae become exposed to Bt insecticidal protein at later growth stages when they can survive to produce offspring. This will lead to an increase in the frequency of resistant individuals (both RS and RR) in the population. If the field is designated as a refuge crop, the presence of the Bollgard 3 cotton volunteers will diminish the value of the refuge.





ADAMA



**RX-380™**

**Cotton Growth Regulator**

# The super concentrate.

**ADAMA Reward®**  
(38 g/L mepiquat chloride)

**ADAMA RX-380**  
(380 g/L mepiquat chloride)

Application rate (mL/ha)

|      |     |
|------|-----|
| 250  | 25  |
| 500  | 50  |
| 1000 | 100 |
| 1500 | 150 |
| 2000 | 200 |

New ADAMA RX-380 is a highly concentrated formulation of the cotton growth regulator, mepiquat chloride.

- 10 times the concentration of ADAMA REWARD®
- 380 g/L mepiquat chloride in a soluble concentrate (SL) formulation
- Significantly reduces packaging wastage and handling
- Compatible with a wide range of ADAMA products
- Look for the new dark red formulation colour and 10 L pack size!

™Trademark or ®Registered trademarks of an ADAMA Agricultural Solutions Ltd Company.  
Please note: This information is not intended to replace the product label.  
Always read the complete product label appearing on the container before opening or using products. ADA20438A



Scan here for more information



**GROWTH  
REGULATOR**

**ADAMA.COM**



### 3. Planting windows

The planting window limits the number of generations of *Helicoverpa* exposed to Bollgard 3 cotton in any one season, restricting the selection pressure to develop resistance.

In developing the Bollgard 3 RMP, late crops were identified as having the highest potential to increase resistance risk due to increased length of exposure. So the emphasis shifted toward using mitigation tactics that reduce the risk of late crops and Bollgard 3 volunteers, with less emphasis on reducing season length at the start of the growing season. However, planting windows still remain an important mitigation tactic for Bollgard 3, particularly in warmer climates where cotton and *Helicoverpa* can survive and reproduce all year round.

### 4. No Bt sprays

Preventing the use of foliar Bt on all refuges (sprayed and unsprayed), reduces the exposure of *Helicoverpa* to Bt outside the plant and maximises the likelihood of producing moths that are susceptible (SS) rather than resistant (RR) to Bt. This is an important part of the RMP because susceptible refuge moths are presumed to mate with any resistant moths in the population to produce heterozygotes (RS) that are killed by Bt cotton.

### 5. Pupae destruction

Pupae busting is a highly effective mitigation tactic for reducing resistance risk in areas where *Helicoverpa* have a diapause stage, provided it is performed well and at the right time. Although minimal larvae are expected to survive in Bollgard 3 cotton, those that do are most likely carrying resistance genes and are precisely the ones that must be killed so that the next generation of moths (emerging the following spring) are not carrying resistance alleles.


Introducing a defoliation date for Bollgard 3 that determines whether a field requires pupae busting ensures only those fields most likely to contain

highest risk pupae in diapause are being cultivated. Based on the *Helicoverpa* 'Diapause Induction and Emergence model' that was developed from field research, 31 March is when the likelihood of diapause occurring is 50% (NSW and southern Qld). Growers who defoliate earlier are still encouraged to pupae bust. A review of research into pupae busting also indicated that the majority of pupae were found within the hill, closer to the plant line, compared to the furrow, so the pupae busting guidelines have been changed to reflect this. Overall, the changes are intended to make pupae busting more targeted and effective while also improving the practicalities of the operation for growers. Refer to the RMP for details on pupae busting requirements for Bollgard 3.

Do not pupae bust unsprayed refuges. Once Bollgard 3 crops begin flowering and are highly attractive to *Helicoverpa* moths, the corresponding refuge should not be cultivated (e.g. for weed control, row formation etc). Destruction of refuges must not occur until after the destruction of corresponding Bollgard 3 crops.

Sprayed refuges and late conventional cotton crops should be pupae busted as soon as possible after picking to manage resistance to other insecticide products.

In warmer areas (central Queensland and northern Australia), *Helicoverpa* pupae produced late in the cotton season do not remain in the soil, but emerge within 15 days of pupating, so trap crops are required as an alternative. Trap crops of pigeon pea are planted after the cotton and are timed to be at their most attractive after the cotton has cut-out. Moths emerging from Bollgard 3 cotton fields at the end of the season will be attracted to the trap crops and are likely to lay their eggs there. Once the cotton has been harvested, the trap crop should be destroyed, removing the food source from the larvae (which will then die) and the soil then cultivated to destroy any pupae. It is critical to time the destruction so that it corresponds with the period of most effective kill of the range of life stages of *Helicoverpa*. See the RMP for more details.





ADAMA

**Skope<sup>®</sup>**

**Excellent performance  
on key pests including  
Silverleaf Whitefly  
and Green Mirid with  
flexible use timings.**

- Choosing new Skope insecticide gives you excellent performance against key cotton pests, including Green Mirid, Silverleaf Whitefly, Heliothis spp, Cotton Aphid and Green Vegetable Bug
- With two modes of action, IPM fit and no mite flaring, Skope makes no compromises when managing multi-pest scenarios in Bollgard# cotton.



Scan here for  
more information



**INSECTICIDE**

**ADAMA.COM**



©Registered trademarks of an ADAMA Agricultural Solutions Company.

## Guidelines for *Helicoverpa* management in Bollgard 3 cotton

Since 2005–06 there have been occasional reports of larvae surviving for several weeks at threshold levels in Bt fields. All affected fields were at mid-flowering to late-flowering and included *H. armigera* and *H. punctigera*.

Work conducted by CSIRO and Bayer demonstrated that this survival was not due to Bt resistance or because of the absence of Bt genes in the cotton. Larvae exhibit strong behavioural responses to the Bt insecticidal protein in Bt cotton plants, and attempts to avoid the Bt toxins can result in movement of larvae within and between plants, resulting in an apparent feeding preference for flowers. These behaviours, coupled with the sometimes temporal and spatial variability of Bt toxin expression, can result in some larvae becoming established.

For resistance management reasons, larvae that reach thresholds in Bollgard 3 fields should be controlled by spraying. However, work conducted by Bayer suggests that a yield penalty associated with larvae survival in Bollgard 3 fields is unlikely. Bt cotton plants could tolerate up to 100% square loss at early flowering, up to 100% square removal alone or in combination with 30% boll damage at peak flowering, and 30% boll damage at late flowering, without impacting yield or quality. Therefore Bt cotton seems to compensate well for damage caused by larvae and the current threshold can be used in most situations without causing significant yield reduction.

It is critical that the distribution and proportions of fields that are affected by surviving larvae, and the number of fields that are sprayed to control *Helicoverpa* is recorded. Part of the end of season general survey of Crop Consultants Australia (CCA) members includes questions about control of *Helicoverpa* in Bollgard 3 fields.

If you experience above threshold levels of *Helicoverpa* in your Bt fields please immediately contact:

- **Sharon Downes: 0427 480 967 or**
- **Kristen Knight: 0429 666 086**

## Insecticide selection for Bollgard 3 crops

When controlling *Helicoverpa* within Bollgard 3 crops, insecticide selection should comply with the cotton industry's Insecticide Resistance Management Strategy (pages 56–57). The beneficial/pest ratio (described on page 28) should also be given careful consideration before applying insecticides. If an insecticide is required, choose the most effective product that is the least disruptive to the beneficial complex (refer to Table 4). While foliar Bt can be used on Bollgard 3 crops, it is a requirement of the Bollgard 3 cotton Resistance Management Plan that foliar Bt not be used on any refuge crops.

### *Helicoverpa* thresholds

Do not include eggs or any larvae <3 mm long in spray threshold counts. For economic management of *Helicoverpa*, larval populations should be controlled with an insecticide if a threshold of:

- 2 larvae/m >3 mm long are found over 2 consecutive checks; or,
- 1 larvae/m >8 mm long is found in any check.

Application of these thresholds requires careful and accurate assessment. Checks should be made over the whole plant including the terminals, squares and especially flowers and small bolls. Be sure to objectively assess larval size. A complete description of the sampling protocols for *Helicoverpa* can be found on page 27.

**For the Bollgard 3 Schedule, refer to your Technology User Agreement (TUA). Information on the Resistance Management Plan for Bollgard 3 cotton is available at [www.bollgard3.com.au](http://www.bollgard3.com.au)**



**Rinse them out  
Round them up  
Run them in**

**drummuster** provides Australian agricultural and veterinary chemical users with a recycling pathway for eligible empty agvet chemical containers.

**drumMUSTER** is a national product stewardship program that is supported by agvet chemical manufacturers, industry stakeholders including member and farming associations as well as state and local governments.

Recycle your empty eligible agvet containers. Visit the **drumMUSTER** website to find the nearest collection site to you. Remember, every container counts.



[drummuster.org.au](http://drummuster.org.au)



**drumMUSTER**  
rinse them out, round them up and run them in.

[info@drummuster.org.au](mailto:info@drummuster.org.au)  
1800 008 707

**BOLLGARD 3 RESISTANCE MANAGEMENT PLAN (NOTE Northern Australia RMP is available at [www.bollgard3.com.au](http://www.bollgard3.com.au))**

**Developed by Monsanto Australia Pty Ltd**

The Resistance Management Plan is based on three basic principles: (1) minimising the exposure of *Helicoverpa* spp. to the *Bacillus thuringiensis* (Bt) proteins Cry1Ac, Cry2Ab and Vip3A, (2) providing a population of susceptible individuals that can mate with any resistant individuals, hence diluting any potential resistance, and (3) removing resistant individuals at the end of the cotton season. These principles are supported through the implementation of five elements that are the key components of the Resistance Management Plan. These elements are:

1. Planting restrictions;
2. Refuge crops;
3. Control of volunteers and ratoon cotton;
4. Pupae destruction/trap crops; and,
5. Spray limitations.

Growers of Bollgard 3 cotton are required to practice preventative resistance management as set out below. Compliance with the Resistance Management Plan is required under the terms of the Bollgard 3 Technology User Agreement and per the Conditions of Registration for Bollgard 3 under the Agricultural and Veterinary Chemicals Act 1994.

**1. Planting Restrictions**

**Victoria, New South Wales and Southern Queensland**

All Bollgard 3 crops and refuges must be planted into moisture or watered-up between August 1 and December 31 each year, unless otherwise specified in this Resistance Management Plan.

**Central Queensland**

All Bollgard 3 crops and refuges must be planted into moisture or watered-up between August 1 and before October 31 each year, unless otherwise specified in this Resistance Management Plan. Bollgard 3 can only be planted from August 1 to October 31 each year. Seed cannot be planted wet or dry prior to August 1. Any Bollgard 3 crops planted into moisture or watered-up after October 31 and up to December 31 must plant additional refuge as specified in Table 3 and 4. Bollgard 3 cannot be planted dry prior to December 31 if not watered up.

**2. Refuges**

Growers planting Bollgard 3 cotton will be required to grow a refuge crop that is capable of producing large numbers of *Helicoverpa* spp. moths which have not been exposed to selection with the Bt proteins Cry1Ac, Cry2Ab and Vip3A. These unselected moths are expected to dominate matings with any survivors from Bollgard 3 crops and thus help to maintain resistant alleles to the Bt proteins Cry1Ac, Cry2Ab and Vip3A at low frequencies.

All refuge options are based on the requirement of a 5% unsprayed cotton refuge or its equivalent, as determined by the relative production of *Helicoverpa* spp. from each of the refuge types as described in Tables 1 and 2 for irrigated and dryland production scenarios, respectively.

For each area of irrigated Bollgard 3 cotton planted, a grower is required to plant one or more of the following:

**TABLE 1: Irrigated Bollgard 3 cotton refuge options**

| Crop       | Conditions                               | % of Bollgard 3 |
|------------|--|-----------------|
| Cotton     | Irrigated, sprayed conventional cotton   | 100             |
|            | Irrigated, unsprayed conventional cotton | 5               |
| Pigeon pea | Fully irrigated, unsprayed               | 2.5             |

**TABLE 2: Dryland Bollgard 3 cotton refuge options**

| Crop       | Conditions  | % of Bollgard 3 |
|------------|---|-----------------|
| Cotton     | Dryland or irrigated, sprayed conventional cotton   | 100             |
|            | Dryland or irrigated, unsprayed conventional cotton   | 5               |
| Pigeon pea | Dryland or fully irrigated, unsprayed. Dryland pigeon peas can only be planted with an approved plan from Monsanto Australia. | 2.5             |

**TABLE 3: Irrigated Bollgard 3 cotton refuge options for Central Queensland planted after October 31**

| Crop       | Conditions                               | % of Bollgard 3 |
|------------|--|-----------------|
| Cotton     | Irrigated, sprayed conventional cotton   | 100             |
|            | Irrigated, unsprayed conventional cotton | 10              |
| Pigeon pea | Fully irrigated, unsprayed               | 5               |

**TABLE 4: Dryland Bollgard 3 cotton refuge options for Central Queensland planted after October 31**

| Crop       | Conditions  | % of Bollgard 3 |
|------------|---|-----------------|
| Cotton     | Dryland or irrigated, sprayed conventional cotton   | 100             |
|            | Dryland or irrigated, unsprayed conventional cotton   | 10              |
| Pigeon pea | Dryland or fully irrigated, unsprayed. Dryland pigeon peas can only be planted with an approved plan from Monsanto Australia. | 5               |

**Note:** Unsprayed means not sprayed with any insecticide that targets any life stage of *Helicoverpa* spp. Bt products must not be applied to any refuge (including sprayed cotton). If the viability of an unsprayed refuge is at risk due to early or late season pressure by *Helicoverpa* spp., or any other caterpillar species, contact Monsanto Australia immediately. With prior approval from Monsanto Australia, a non-Bt helioicid can be applied. For the purposes of this Resistance Management Plan, conventional cotton includes any cotton varieties that do not have Bt proteins in the plant that control *Helicoverpa* spp. larvae.

## BOLLGARD 3 RESISTANCE MANAGEMENT PLAN

### General conditions for all refuges

- (a) Refuge crops are to be planted and managed so that they are attractive to *Helicoverpa* spp. during the growing period of the Bollgard 3 cotton varieties.
- Irrigated:** It is preferable that all refuge is planted within the 2 week period prior to planting Bollgard 3. If this is not possible, refuge planting must be completed within 3 weeks of the first day of sowing of Bollgard 3. At this time, sufficient refuge must have been planted to cover all of the Bollgard 3 cotton proposed to be planted for the season (including Bollgard 3 already planted and any that remains unplanted). If additional Bollgard 3 is planted after this date, which is not already covered by refuge, additional refuge must be planted as soon as possible and no more than 2 weeks after sowing of the additional Bollgard 3.
- Dryland:** A dryland refuge must be planted within the 2 week period prior to the first day of planting Bollgard 3 cotton.
- (b) Pigeon pea refuges should not be planted until the soil temperature reaches 17°C, which is a requirement for germination, and should also be planted into moisture to ensure successful germination. If soil temperatures are not suitable to allow germination of pigeon peas in line with condition (a), an alternative refuge must be planted in its place within the prescribed period (under (a) above).
- (c) All refuges should preferably be planted into a fallow or rotation field that has not been planted to Bt cotton in the previous season to avoid volunteer and ratoon cotton. See Refuge Management Guide for all unsprayed refuges.
- (d) Once Bollgard 3 cotton begins to flower, the corresponding refuge must not be cultivated.
- (e) All refuges are to be planted within the farm unit growing Bollgard 3 cotton no more than 2 km from the associated Bollgard 3 cotton field. For any cases where it may not be possible to plant the refuge within 2 km from the associated Bollgard 3, approval must be sought from Monsanto Australia.
- (f) To minimise the possibility of refuge attractiveness being affected by herbicide drift, non-herbicide tolerant refuges should be separated from herbicide tolerant Bollgard 3 cotton crops by a sufficient distance to minimise such drift, but no more than 2 km from the Bollgard 3 cotton.
- (g) To account for possible insecticide drift, the options for the width of refuge crops vary according to spray regime. If any sprayed conventional cotton is grown on the same farm unit, Bollgard 3 refuge crops must be at least 48 metres wide and each refuge area must be a minimum of 2 hectares. If sprayed conventional cotton is not grown on the same farm unit, Bollgard 3 refuge crops must be at least 24 metres wide and each refuge area must be a minimum of 0.5 hectares. Different unsprayed refuge options may be planted in the same field as a single unit; however a sprayed conventional cotton refuge must not be planted in a field that is also planted to an unsprayed refuge type unless a sufficient buffer is in place to prevent insecticide drift.
- (h) In all regions, destruction of refuges must only be carried out after Bollgard 3 has been harvested (refer to section 4 Pupae Destruction).
- (i) Refuges for dryland Bollgard 3 cotton crops must be planted in the same row configuration as the Bollgard 3 crop unless the refuge is irrigated. If an irrigated option is utilised for a dryland Bollgard 3 crop, then that refuge may be planted in a solid configuration. Dryland cotton is measured as green hectares.

### 3. Control of volunteer and ratoon cotton

Volunteer and ratoon cotton may impose additional selection pressure on *Helicoverpa* spp. to develop resistance to the Bt proteins Cry1Ac, Cry2Ab and Vip3A produced by Bollgard 3 cotton.

As soon as practical after harvest, Bollgard 3 cotton crops must be destroyed by cultivation, root cutting or herbicide so that they do not continue to act as hosts for *Helicoverpa* spp.

Growers must ensure that volunteer and ratoon plants are removed prior to flowering from all fields, including fallow areas, Bollgard 3 crops, conventional cotton crops and all refuges. **The presence of Bollgard 3 volunteers/ratoon cotton in any refuge will diminish the value of the refuge and must be removed as soon as possible.**

**Note:** The refuge should preferably be planted into fallow or rotation fields that have not been planted to cotton in the previous season.

### 4. Pupae destruction/trap crops

#### Victoria, New South Wales and Southern Queensland

To further mitigate the risk of resistance, each grower of Bollgard 3 must undertake *Helicoverpa* spp. pupae destruction in fields with a higher probability of carrying over wintering pupae according to the following key guidelines:

- If first defoliation of a Bollgard 3 field occurs on or before March 31, the Bollgard 3 field must be slashed or mulched and controlled to prevent regrowth within 4 weeks of harvesting.
- If first defoliation of a Bollgard 3 field occurs after March 31, the Bollgard 3 field must be slashed or mulched and controlled to prevent regrowth within 4 weeks of harvesting and pupae busting must be completed by July 31 for all valleys except regions including the Lachlan, Murrumbidgee, Menindee and Murray Valleys and Victoria where pupae busting must be complete by August 31.
- Ensure disturbance of the soil surface to a depth of 10 cm to a distance of 30 cm both sides of the plant line.

#### Option for an alternative pupae destruction management for Lachlan, Murrumbidgee, Menindee, Murray Valleys and Victoria

If opting to apply a registered attract and kill for *Helicoverpa* spp. instead of pupae destruction for the Lachlan, Murrumbidgee, Menindee and Murray Valleys and Victoria:

- Grower must advise at planting audit if opting in for attract and kill strategy but can opt out up to and including February 1.
- Grower must apply 3 weekly applications commencing no earlier than February 10 with the final application being no later than March 1.
- Grower must make applications as per the label of the registered attract and kill for *Helicoverpa*.
- For all other valleys contact your local Bayer Cotton Territory Business Manager.

**BOLLGARD 3 RESISTANCE MANAGEMENT PLAN**

**Central Queensland**

**Crop destruction**

All Bollgard 3 crops must be slashed or mulched and controlled to prevent regrowth within 4 weeks of harvesting.

**End of season management of refuges/trap crops**

End of season pupae busting practices are not effective in the Central Queensland region as *Helicoverpa* spp. are less likely to diapause. A late summer trap crop (pigeon pea) must be planted for all Bollgard 3 cotton grown in Central Queensland. The planting configuration of the trap crop should be the same as that of the Bollgard 3 crop. Irrigated Bollgard 3 must have an irrigated trap crop. Table 5 shows the requirements for the late summer pigeon pea trap crop. **Dryland Bollgard 3 growers who do not have any irrigated cotton on their farm should contact Monsanto Australia for alternative options.**

Refuge and late summer trap crops have different purposes. Where a pigeon pea refuge is utilised, the full pigeon pea refuge area must be managed to become the late summer trap crop. If unsprayed cotton is used as the refuge, an additional area of 1% pigeon pea must be planted as the late summer trap crop. Requirements for late summer trap crops are detailed in Table 5 (below).

**TABLE 5: Late summer pigeon pea trap crop requirements in Central Queensland**

| Criterion                              | Trap crop*   |
|--|--|
| Minimum area & dimension (Requirement) | A minimum trap crop of 1% of planted Bollgard 3 cotton crop is required.<br>If sprayed conventional cotton is grown on that farm unit: the trap crop must be at least 48 m x 48 m.<br>If no sprayed conventional cotton is grown on that farm unit: the trap crop must be at least 24 m x 24 m.  |
| Planting time                          | The trap crop should preferably be planted 4 weeks after the associated Bollgard 3. Note: if growers choose to plant their trap crop to coincide with the planting of pigeon pea refuges, they must manage the trap crop in such a way that it remains attractive to <i>Helicoverpa</i> spp. 2-4 weeks after final defoliation.  |
| Planting rate**                        | 35 kg/ha (recommended establishment greater than 4 plants per metre)   |
| Insect control                         | The trap crop can be sprayed with virus after flowering, while avoiding insecticide spray drift, <b>EXCEPT</b> where a pigeon pea refuge is converted to a trap crop. <b>In this case</b> the full 2.5% pigeon pea (or 5% pigeon pea if cotton is planted in the later window after October 31) refuge area managed to become the late summer trap crop can only be sprayed with virus after the first defoliation of Bollgard 3 cotton. |
| Irrigation                             | The refuge/trap crop must be planted into an area where it can receive the additional irrigation required to keep the trap crop attractive to <i>Helicoverpa</i> spp. until after the cotton is defoliated.  |
| Weed control                           | The trap crop should be kept free of weeds and particularly volunteer Bollgard 3 cotton. When using the full pigeon pea refuge area as the trap crop, weed control must not be carried out by cultivation once flowering of the associated Bollgard 3 cotton crop has commenced.   |
| Crop destruction                       | The trap crop must be destroyed 2-4 weeks (but not before 2 weeks) after final defoliation of the Bollgard 3 cotton crop, (slash and pupae bust – full soil disturbance to a depth of 10 cm across the entire trap crop area). All Bollgard 3 and associated trap crops must be destroyed by July 31.  |

\*A pigeon pea trap crop is to be planted so that it is attractive (flowering) to *Helicoverpa* spp. after the cotton crop has cut out, and as any survivors from the Bollgard 3 crop emerge. Planting pigeon pea too early (e.g. before November) or too late (e.g. mid December) is not adequate for cotton crops planted during September through to October.  
\*\*The planting rate is a recommendation based on a minimum of 85% seed germination.

**Failed crops – all regions**

Bollgard 3 crops that will not be grown through to harvest for various reasons and are declared to, and verified by, Monsanto as failed must be destroyed within two weeks after verification, in such a way that prevents regrowth. Crops that are abandoned before February 28 should be slashed and mulched within 4 weeks.

**5. Spray limitations**

Insecticide preparations containing Bt may be used on Bollgard 3 cotton throughout the season BUT NOT on any refuge crops.

An unsprayed refuge should not be planted in the same field as any crop sprayed with a rate of insecticide that is registered for *Helicoverpa* spp., with the exception of Bollgard 3. Sprayed crops and unsprayed refuges that are planted in adjacent fields must be separated by sufficient distance to minimise the likelihood of insecticide drift onto the unsprayed refuge.

If the viability of an unsprayed refuge is at risk due to early or late season pressure by *Helicoverpa* spp., or any other caterpillar species, contact Monsanto Australia immediately. With prior approval from Monsanto Australia, a non-Bt heliocide can be applied.

**Note:** If any grower encounters problems in complying with the Resistance Management Plan, please contact Monsanto Australia.

Background information on the various components of this plan are provided earlier in the RMP section of this guide.

**Please check with Bayer for the most up to date version as the RMP was under review at the time of this guide’s publication.**



# Managing unsprayed pigeon pea refuges

**E**stablishing and growing an attractive refuge is a mandatory component of the Resistance Management Plan (RMP) for Bollgard 3. The purpose of a refuge is to generate significant numbers of *Helicoverpa* moths that have not been exposed to selection pressure from any insecticidal Bt proteins. Attractive, fully irrigated, unsprayed flowering pigeon pea will, on average, produce twice as many moths as the same area of unsprayed non-Bt cotton. As well as producing high numbers of moths, it is also crucial that moth production from refuges has synchrony with Bollgard 3 cotton crops. A well-watered refuge with adequate nitrogen is most likely to sustain larvae through to pupation and consequently produce the most moths. This is the key to delaying Bt resistance.

**The following information is intended to assist growers to establish and maintain effective pigeon pea refuges. Refer to the RMP for guidance on mandatory refuge requirements.**

## Field selection

Pigeon pea can be grown on many soil types but can be susceptible to waterlogging, so select fields that have good post-irrigation/rainfall drainage. Avoid fields that were sown to cotton during the previous season to reduce the likelihood of volunteer and ratoon cotton occurring in refuges. The presence of Bollgard 3 cotton in refuge areas diminishes the resistance mitigation potential of the refuge. Similarly, selecting fields with a low weed seed bank enables easier management of weeds that could compete with the pigeon pea and reduce refuge effectiveness.

Ideally, refuges should be sown adjacent to the Bollgard 3 crop. Ensure sufficient separation to buffer against the drift of herbicides or insecticides applied to the cotton or other crops onto the refuge area.

As with many other legumes, pigeon pea can have allelopathic effects on subsequent crops, so take this into account when making field selections.

## Crop establishment

### Timing

Similar to mungbean and soybean, a minimum soil temperature of 17°C and rising is optimal for pigeon pea establishment. In most cotton production regions these conditions occur during October–November. Under the RMP, pigeon pea should be sown within the two week period prior to planting Bollgard 3, or if not possible, completed within 3 weeks of the first day of sowing for irrigated crops.

### Sowing and inoculation

Nitrogen fixation by pigeon pea is optimal in soils with very low residual soil N. The use of peat-based Group J inoculation formulations on seed just prior to planting will help to ensure effective rootzone colonisation by active strains of rhizobium bacteria. Effective nodulation of the root system can reduce crop susceptibility to waterlogging. To ensure inoculant efficacy, follow all label requirements and directions regarding storage, handling and application.

Match pigeon pea row spacing to that of the corresponding Bt cotton crop.



**Sunrise™ (left) is a strongly indeterminate variety compared with the varieties traditionally used for refuges. (Paul Grundy)**

The continual recycling and saving of seed from undamaged refuges caused an evolutionary shift towards pigeon peas that flower much later (or at times not at all). Sunrise™ is a new variety of pigeon pea that exhibits excellent vigour under furrow irrigation across a range of soil types, and should commence flowering in early January or typically within 75 days of sowing. Sunrise™ is strongly indeterminate and has the ability to flower repeatedly, particularly after sustaining insect attack.

Sunrise™ seed production is being undertaken annually by Associated Grains to ensure that the planting seed that is available to industry has excellent germination characteristics and remains true to type to preserve the heritage of this variety for years to come. Sowing rates for Sunrise™ typically fall within the range of 25–40 kg/ha (depending on germination statistics and field conditions at the time of planting). Growers concerned about crop residues should consider using planting rates at the higher end of this range to promote plants with thinner stalks, which makes later crop destruction much easier.

Comparisons between Sunrise™ pigeon peas and the original pea cultivar type Quest (under commercial conditions at a range of sites over several seasons) have demonstrated that Sunrise™ flowers much longer than the determinate Quest, and on average generates 2–3 times more pupae per hectare of refuge.

## Seed bed preparation and planting

Ensure that the seedbed has good till to maximise seedling emergence and establishment. Seed should not be sown deeper than 5 cm. Levelling of any seed trenches created during planting is important, particularly when residual herbicides have been used and/or the field is to be watered up. The use of press wheels with light pressure has been shown to improve emergence.

## Pre-irrigation

Pre-irrigation and planting into moisture is generally recommended over watering up. Some growers choose to water up the refuge with the rest of the field, then replant into this moisture if a replant is required.

## Crop nutrition

Pigeon pea requires inoculation with Group J inoculant. Nodulation will be limited in high nitrogen soils. A well-grown crop of pigeon pea can add up to 38 kg/ha of nitrogen. Pigeon pea is much more sensitive to phosphorus deficiency than cotton. In soils with long cropping histories where soil P may be depleted, pigeon pea is likely to respond to the addition of phosphorus and zinc.

## Weed management

Pigeon pea are poor competitors with weeds during establishment, particularly when planted under cool conditions.

As well as herbicides, inter-row cultivation can be a useful tactic. However, because cultivation can inadvertently kill (the Bt-susceptible) *Helicoverpa* pupae present in the soil, it is a requirement that once Bt cotton begins to flower, the corresponding refuge should not be cultivated.

The presence of volunteers/ratoon Bollgard 3 cotton in any refuge will diminish the value of the refuge and must be removed as soon as possible.

## Irrigation

Pigeon pea can be very sensitive to waterlogging on heavier soil types. Select a site with good drainage, avoid irrigation prior to heavy rainfall predictions and only water every second row to reduce the risk of waterlogging on heavy soils. Use the same best management tactics on pigeon pea as those being used for cotton crops, e.g. getting water on and off the field in a timely and effective manner.

Being a drought-tolerant plant, pigeon pea generally has a lower water requirement than cotton. However, ensure crops do not become moisture stressed as this reduces attractiveness and truncates the flowering period. Sunrise™ is an indeterminate variety, and in some circumstances irrigating too frequently can prolong the vegetative growth stage delaying the onset of flowering. A good rule of thumb is to plant Sunrise™ on a full profile of moisture and then apply the first in-crop irrigation as the plants begin to develop flower buds. On lighter loamy soil types with lower moisture holding capacity (<140 mm PAWC) or in hotter climates, two irrigations prior to flowering may be required. Sunrise™ should be exhibiting signs of budding by the time it reaches 50–70 cm in height. After the initiation of buds, Sunrise™ can be irrigated on a similar schedule to the adjacent cotton to prolong flowering and ensure rapid regrowth after insect attack.



Larvae in pigeon pea refuge. (Johnele Rogan)

## Destruction and harvest of pigeon pea refuge crops

Harvest or destruction of a pigeon pea refuge should only be carried out after the corresponding Bollgard 3 cotton crop has been fully picked. In NSW and Southern Queensland, soil disturbance should only occur after Bollgard 3 cotton fields have been pupae busted (to ensure maximum emergence of pupae from refuges).

In Central Queensland soil disturbance of refuge crops can only occur 2 weeks after the final defoliation of the Bollgard 3 cotton. Growers in Central Queensland using pigeon pea for trap crop purposes should refer to the late summer pigeon pea trap crop requirements of the RMP for full details.

**Do not feed crop product or crop residue to livestock.** III

**TABLE 16: Herbicides available for use in pigeon pea**

| Active ingredient | MoA group | Comment   |
|-------------------|-----------|---|
| Prometryn*        | C         | Apply up to the maximum rate pre planting and incorporate, or as a post-emergent directed spray towards the base of established plants.   |
| Trifluralin       | D         | Apply up to the maximum rate pre planting and incorporate. NSW and ACT only.  |
| Butoxydim*        | A         | Apply the specified rate as a post-emergence spray over the top of the pigeon pea crops. Refer to label as rates are different depending on weed being controlled.  |
| Fluazifop-p*      | A         |   |
| Haloxyfop*        | A         |   |
| Sethoxydim*       | A         | Apply specified rate as a post-emergence spray over the top of the pigeon pea crops.  |
| Clethodim*        | A         | Always apply with D-C-trate at 2 L/100 L or Hasten or Kwickin at 1 L/100 L. Uptake at 500 mL/100 L spray volume. The lower doses will provide effective control if applied under ideal conditions to weed that are smaller, actively growing and free from temperature or water stress.   |
| Quizalofop*       | A         | Refer to permit for growth stages of species and critical comments.   |
| Diquat            | L         | Harvest aid   |
| Diquat/paraquat   | L         | Apply pre-sowing, in minimum 50-100 L water. Apply specified rates for certain weeds at particular growth stages, refer to label.   |
| Pendimethalin     | D         | Incorporate into the soil within 24 hours of application. Use higher rate on heavy textured soils or those high in organic matter. May be applied by aerial or ground spraying. In Macquarie Valley area, only apply by air when ground is too wet for ground application.  |
| Metribuzin        | C         | Furrow irrigated: apply after furrowing out, within 2 weeks before sowing and incorporate. For post-emergence: apply to actively growing seedling stage weeds provided crop plants have at least 2 trifoliolate leaves. Do not spray if rain is likely to fall within several hours. Overhead irrigated: apply pre-emergence then irrigate. |

\* Use of these products is under permit PER13758 (valid until 30 September 2022).

Permit comments: Only apply to pigeon pea crops that are to be destroyed at the end of the season or to be harvested for seed for refuge replanting. No crop product or crop residue is to be fed to livestock. Unless otherwise stated in the permit, use must be in accordance with the product label.

# Integrated Weed Management (IWM)

It is important to strategically plan how different tactics will be utilised to give the best overall results for the existing weed spectrum. A short term approach to weed management may reduce costs for the immediate crop or fallow, but is unlikely to be cost effective over a five or ten year cropping plan. Over this duration, problems with species shift and the development of herbicide resistant weed populations are likely to occur where weed control has not been part of an integrated plan.

## Develop a strategy

Having good records on crop rotations, herbicides and other tactics used as well as weed species present will help in developing a plan that identifies where there are particular risks in the system and also where there might be opportunities to incorporate additional tactics. The herbicide resistance management strategy (HRMS pages 81-87) can help to inform the effectiveness of combinations of tactics on reducing the weed seed bank as well as the risk of herbicide resistance.

## Know your enemy

Consider what weed species are present. Ensure that weeds are correctly identified, and consider which tactics, or combination of tactics, are going to be most effective for your weed spectrum. Key resources that can assist with these decisions include:

**Weeds of Australian Cotton App.**

[www.cottoninfo.com.au/weeds-australian-cotton-app](http://www.cottoninfo.com.au/weeds-australian-cotton-app)

**Weed ID guide** [www.cottoninfo.com.au/publications/weedpak-weed-id-guide](http://www.cottoninfo.com.au/publications/weedpak-weed-id-guide)

**WEEDpak** [www.cottoninfo.com.au/publications/weedpak](http://www.cottoninfo.com.au/publications/weedpak)

Identify any particular problem areas. Managing these patches more intensively may help prevent a problem weed or resistance from spreading.

## Time your tactics

Often the timeliness of a weed control operation has the largest single impact on its effectiveness. For example, herbicides are far more effective on rapidly growing small weeds, and may be quite ineffective in controlling large or stressed weeds.

## Think about the whole farming system

It is important to consider weed management in the context of other in-crop agronomic issues, other crops and across the whole farm.

## Consider crop competition

An evenly established, vigorously growing cotton crop can compete strongly with weeds, especially later in the season. Delaying planting on weedy fields until last gives more opportunity to control weeds that emerge prior to planting, and better conditions for cotton emergence and early vigorous growth. In irrigated crops, weed-free periods of 8-9 weeks from planting provide enough time for cotton to out-compete most later emerging weeds and significantly reduce weed seed production. Refer to critical periods for weed control (page 68, 2021 ACPM) for more information.

**The MoA classification system for herbicides will be changing from mid-2021. See the *Herbicide resistance management* section for more details.**

## Plan weed management to fit with other operations

Look for opportunities in the cropping system to coordinate weed control tillage with other operations, such as pupae busting, fertiliser incorporation, seed bed preparation and irrigation furrow maintenance.

## Consider the wider impacts of weeds

Weed management is also an important consideration for pest and disease management. Many cotton pests rely on weed hosts and cotton volunteers prior to migrating into cotton fields. Some weeds and cotton volunteers/ratoons can act as a reservoir for plant viruses such as cotton bunchy top disease which can cause significant loss of yield. Some weeds can also allow disease inoculum to build up in the soil, increasing the risk for subsequent crops.

## Mix it up with rotation crops

Rotation crops provide an opportunity to introduce a range of different tactics into the system, particularly herbicide groups that are not available in cotton. Mixing rotations may also vary the time of year non-selective measures can be used and the time of year that crop competition suppresses weed growth. Rotation between summer and winter cropping provides opportunities to use cultivation and knockdown herbicides in-fallow at all times of the year. Where cotton is grown in rotation with other crops, such as winter cereals or maize, retaining the stubble cover from these rotation crops for as long as possible reduces weed establishment and encourages more rapid breakdown of weed seed on the soil surface. In terms of the HRMS, treat rotation crops like a fallow and aim to use at least 2 non-glyphosate tactics within the crop.

## Don't forget non-crop areas

Weeds in non-crop areas on the farm such as channels, tail drains, fence lines and roadsides can develop and introduce herbicide resistance into the farming system. Manage these areas as a fallow, using a range of tactics including residual herbicides, cultivation and chipping of weeds. Do NOT rely on glyphosate only to manage weeds in non-crop areas.

**TABLE 17: Control of weeds in dry channels**

| Active ingredient               | Mode of Action group |
|---------------------------------|----------------------|
| Amitrole + ammonium thiocyanate | Q                    |
| Glyphosate                      | M                    |
| Imazapyr + glyphosate           | B + M                |
| Pendimethalin                   | D                    |
| Flumioxazin                     | G                    |

## Dryland cotton

Moisture is the key ingredient for dryland cotton and weeds are very efficient at robbing the soil of vital moisture and nutrients. Controlling weeds in winter crops pre-cotton and maintaining a clean winter fallow are crucial in providing sowing opportunities for dryland cotton. Running the weed seed bank down in the previous winter crop or fallow is vital to keep in-crop weed management costs at a minimum. Including residual herbicides in fallows and ensuring that pre-emergent herbicides are used at planting will provide ongoing in-crop weed control and reduce the reliance on glyphosate.

Rotating to other modes of action (MoA) for knockdown weed control is an important management strategy to prolong the life of glyphosate. The development of new Group G herbicides also provides additional control options pre-sowing. Cultivation is often seen as detrimental in dryland systems due to moisture losses from cultivation passes.

Late in-crop strategic cultivation to remove weed escapes is an ideal time to introduce non-chemical weed control, where the impacts of moisture loss are reduced. Alternatively, consider the inclusion of cover crops to reduce weed numbers and improve soil health while also trapping and storing valuable soil moisture. As with irrigated cotton, it is important to ensure that any surviving weeds are prevented from setting seed.

### Come Clean. Go Clean.

To minimise the entry of new weeds into fields, clean down boots, vehicles, and equipment between fields and between properties. Pickers and headers require special attention. Eradicate any new weeds that appear while they are still in small patches. Monitor patches frequently for new emergences. More biosecurity information is available at [www.cottoninfo.com.au](http://www.cottoninfo.com.au)

Irrigation water can be a source of weed infestation with weed seeds being carried in the water. Control weeds that establish on irrigation storages, supply channels and head ditches.

### Control survivors before they set seed

To be effective in preventing resistance, weeds that survive a herbicide application must be controlled by another tactic before they are able to set seed. Monitor the field after each spray application to assess efficacy. Weeds may need to be closely examined, as some are capable of setting seed while very small and many weeds respond to varying day-length, so a winter weed emerging in late winter or spring may rapidly enter the reproductive phase of growth in response to lengthening daylight hours.

Preventing survivors from setting seed is critical to the longer term success of an IWM strategy.

In terms of survivor control, research indicates that high efficacy with an alternative tactic is good, but high frequency control is better than reliance on efficacy. Cultivation after glyphosate application is predicted to achieve 80% survivor control, whereas cultivation plus chipping is predicted to achieve 99.9% survivor control. Other tactics for survivor control could

be equally effective, such as shielded or spot-spraying with an effective knockdown herbicide.

### Manual chipping

Ideally suited to dealing with low densities of weeds, especially those that occur within the crop row, manual chipping is normally used to supplement inter-row cultivation or spraying. As a tool to prevent survivors setting seed, chipping has been shown to be a very cost effective option.


### Spot spraying

Spot sprayers may be used as a cheaper alternative to manual chipping for controlling low densities of weeds in-crop. Ideally, spray weeds with a relatively high label rate of a herbicide from a different herbicide group to the herbicides most recently used to ensure that all weeds are controlled. This intensive tactic can be particularly useful for new weed infestations where weed numbers are low, or where weeds are outside the field and difficult to get to, such as roadside culverts.


New weed detection technologies provide an opportunity to use spot spraying (optical sprayers; e.g. WEEDit®, WeedSeeker®) across large areas of fallow. This can provide opportunity to reduce herbicide costs, while still ensuring robust label rates are applied to problem weeds. In addition, the development of green-on-green technology has the potential to add flexibility into the spraying operation by targeting weeds in-crop. Refer to the herbicide label for plant-back limitations relevant to the rate applied. Follow manufacturer recommendations for speed and nozzle type, as well as allowable products to ensure each application is effective. Permits for herbicide application in fallow are available on the APVMA website [portal.apvma.gov.au/permits](http://portal.apvma.gov.au/permits)

### Fallow management


Weed management in the fallow is an important component of a weed management plan. Summer fallows following a Roundup Ready Flex® cotton crop where only glyphosate herbicide is used for weed control, poses the greatest risk to glyphosate resistance developing. The continued use



# THE SUMMER BIG 6



|   |   |
|---|---|
| <h4 style="text-align: center; color: #0056b3;">1. USE DIVERSE ROTATIONS</h4> <ul style="list-style-type: none"> <li>Use rotation crops, fallow and pasture phases to drive the weed seedbank down over consecutive years</li> </ul>  | <h4 style="text-align: center; color: #0056b3;">2. DOUBLE KNOCK TO STOP SURVIVORS</h4> <ul style="list-style-type: none"> <li>Use two weed control tactics with different modes of action (eg. glyphosate followed by paraquat) on a single flush of weeds to stop any survivors from the first application setting seed.</li> <li>Also use non-herbicidal tactics - cultivation can be the second knock</li> <li>When executed well (correct rates, timing and application) the double knock tactic can provide 100% control of target weeds.</li> </ul>                       |
| <h4 style="text-align: center; color: #0056b3;">3. MIX AND ROTATE HERBICIDES AND TILLAGE (2+2+0)</h4> <ul style="list-style-type: none"> <li>Rotate between herbicide groups</li> <li>Use different modes of action within same herbicide tank mix</li> <li>Strategic use of tillage can complement a herbicide programme and help get on top of weeds in a reset situation.</li> <li>In cotton systems aim to target both grasses and broadleaf weeds using 2 non-glyphosate tactics for broadleaf &amp; grasses in crop and 2 non-glyphosate tactics during the summer fallow and always remove any survivors. (2+2+0)</li> <li>For cotton post picking, ensure complete survivor control using all options, including at pupae busing for extra weed seed burial.</li> </ul> | <h4 style="text-align: center; color: #0056b3;">4. STOP WEED SEED SET IN CROP AND IN FALLOW</h4> <ul style="list-style-type: none"> <li>Aim for 100% control of weeds and diligently monitor for survivors in all post weed control inspections.</li> <li>Crop top or pre harvest spraying in canola, pulses, feed barley and wheat in weedy paddocks.</li> <li>Consider hay production, brown manure or long fallow in high pressure situations.</li> <li>Spray top/spray fallow pasture prior to cropping phases to ensure a clean start to any seeding operation.</li> </ul> |
| <h4 style="text-align: center; color: #0056b3;">5. CROP COMPETITION IN ROTATIONAL GRAIN CROPS</h4> <ul style="list-style-type: none"> <li>Adopt at least 1 competitive strategy (2 is better), including reduced row spacing, higher seeding rates, precision seed placement, east west sowing and competitive varieties.</li> </ul>  | <h4 style="text-align: center; color: #0056b3;">6. HARVEST WEED SEED CONTROL IN ROTATIONAL GRAIN CROPS</h4> <ul style="list-style-type: none"> <li>Capture weed seed survivors at harvest using chaff lining, chaff tramlining, chaff carts, or integrated weed seed destruction</li> </ul>   |



### WEEDSMART WISDOM

- Never cut the rate;
- Spray well – choose correct nozzles, adjuvants and water rates for efficacy and drift control;
- Plant clean seed - don't plant weed seeds.
- Maintain farm hygiene - clean borders, fence lines, channels and contour banks;
- Test for resistance - know your levels;
- Come Clean. Go Clean. - don't let weeds hitch a ride with visitors;
- Be weed aware!

**TABLE 18: Herbicide plant-backs from rotation crops to cotton**

| Active ingredient                             | MoA | Plant-back to cotton          | Notes  |
|---|-----|-------------------------------|--|
| Aminopyralid + fluroxypyr                     | I   | 9 months                      | Plant-back interval on black cracking clay soils. When rainfall is less than 100 mm for a period of 4 months or greater the plant-back period may be significantly longer.   |
| Aminopyralid + picloram + 2,4-D               | I   | 12 months (northern NSW, Qld) | Plant-back periods for rotational crops and fallow following application of FallowBoss Tordon Herbicide up to 20 months (Southern NSW) 700 mL/ha. Plant-back periods are based on normal rainfall pattern. During drought conditions (or when rainfall is less than 100 mm for a period of 4 months or greater) the plant-back period may be significantly longer. Under such circumstances a bioassay is required, before planting the next crop. |
| Atrazine                                      | C   | 6 months                      | Following treatments of up to 1.4 kg/ha.   |
|   |     | 18 months                     | Following treatments of 1.4 kg/ha to 3.3 kg/ha.  |
| Atrazine + S-metolachlor                      | C+K | 6 months                      | When rates up to 3.2 L/ha are used.  |
|   |     | 18 months                     | When rates up to 3.2 L/ha are used. On alkaline soils, a bioassay or analytical test should be undertaken.   |
| Chlorsulfuron                                 | B   | 18 months                     | Where soil pH is 6.6-7.5 and 700 mm of rain has fallen.<br>For soil pH >7.5 only grow cotton after growing a test strip.   |
|   |     | 3 months                      | When rates up to 30 g/ha are used.   |
| Clopyralid                                    | I   | 6 months                      | When rates of 30-120 g/ha are used   |
|   |     | 24 months                     | When rates above 120 g/ha are used. For all rates at least 100 mm rain required during plant-back period.  |
|   |     | Refer to label                | Do not replant treated areas within 2 years of application of diuron except when otherwise stated on the label.  |
| Diuron  | C   | Refer to label                | Do not replant treated areas to any crop within 1 year after last spray except cotton (along with corn or grain sorghum) which may be planted in the spring following year.  |
| Flumioxazin                                   | G   | 0 months                      | For zero plant-back for knockdown spike rates of 45 g/ha or below.   |
|   |     | 2 months                      | For residual rates above 210 g/ha.   |
| Flumetsulam                                   | B   | 2 years                       | For NSW and Qld a minimum of 50 mm and preferably 100 mm rain or more must have fallen over the warm months of the year.   |
| Imazamox                                      | B   | 10 months                     | Must have 800 mm of rainfall or irrigation   |
| Imazamox + imazapyr                           | B   | Unknown                       | Registered for use in Clearfield crops (all other – 34 months).  |
| Imazethapyr                                   | B   | 22 months                     | Dryland cotton.  |
|   |     | 18 months                     | Irrigated only. (Providing rainfall and irrigation exceeds 2000 mm)  |
| Isoxaflutole                                  | H   | 7 months                      | 350 mm rainfall (do not include flood/furrow irrigation) between application and planting the subsequent crop.   |
| Mefenpyr-diethyl + iodosulfuron-methyl sodium | B   | 12 months                     | Rainfall of less than 500 mm may result in extended re-cropping intervals for summer crops sown in the following season.   |
| Metsulfuron methyl                            | B   | Unknown                       | Registered for use in cereal crops: wheat, barley, triticale. Legume crops: chickpeas (desiccant).   |
| Metribuzin                                    | C   | 6 months                      | Rates <1.5 L/ha. This could be longer if there have been long dry periods between crops.   |
|   |     | 12 months                     | Rates >1.5 L/ha.   |
| Metsulfuron-methyl + mefenpyr-diethyl         | B   | 12 months                     | Rainfall of less than 500 mm following application may result in extended re-cropping intervals for summer crops sown in the following year.   |
| Pyroxasulfone                                 | K   | 5 months                      | + 150 mm rainfall. Less total rainfall between application and planting of the following crop than 150 mm may require extended plant-back period.  |
| Picloram + 2,4-D                              | I   | 12 months                     | (Nth NSW & Qld). Do not use on land to be cultivated for growing susceptible crops within 12 months of application. Based on normal rainfall.  |
| Simazine                                      | C   | 9 months                      | When up to 2.5 kg/ha are used. When rates exceed 2.5 kg/ha, plantings may not be possible for very long periods of time afterwards.  |
| Sulfosulfuron                                 | B   | Unknown                       | Registered for use in cereal crops: wheat, triticale.  |
| Tribenuron methyl                             | B   | Unknown                       | Registered for use in fallows.   |
| Triasulfuron                                  | B   | 15 months                     | Soil pH <7.5, 700 mm rainfall between application and sowing the plant-back crop.  |
|   |     | 18 months                     | Soil pH 7.6-8.5.   |
| Triclopyr + picloram + aminopyralid           | I   | 4 months                      | 0.2 L/ha. During drought conditions (<100 mm rainfall in a 4 month period) the plant-back is significantly longer.   |
|   |     | 6 months                      | 0.4 L/ha.  |


This is a guide only – always read and follow product label directions.

Where fields have been treated with herbicides with no plant-back recommendations to cotton, determine cotton tolerance by growing through to maturity on a smaller scale before sowing larger areas.

# Integrated weed management tactics

## NON GLYPHOSATE WEED TACTICS FOR THE COTTON FARMING SYSTEM

An integrated weed management system relies on a large number of interrelated, complementary components including both chemical and non-chemical tactics as well as cultural practices such as rotation, crop competition, farm hygiene, and crop scouting.



## 2 non-glyphosate tactics in fallow + 2 non-glyphosate tactics in crop & NO SURVIVORS

| NON-GLYPHOSATE TACTICS  |  | NO SURVIVORS  |  |
|---|--|---|--|
| <p><b>Fallow</b></p> <p>Strategic cultivation<br/>Double knock<br/>Optical sprayers<br/>Patch management<br/>Cover crops</p> <p>Non Glyphosate options***<br/>Group C (bromoxynil, terbutylazine)<br/>Group G (flumioxazin, pyraflufen, saflufenacil)<br/>Group H (isoxaflutole)<br/>Group L (paraquat, paraquat/diquat)<br/>Group L/Q (amitrole/paraquat)<br/>Group N (glufosinate)<br/>Group I (2,4-D, dicamba, fluroxypyr)</p> <p><b>Rotation Crops*</b><br/>Crop competition<br/>Rotate MoA<br/>Plant-back restrictions cover crops</p> | <p><b>Pre/at plant</b></p> <p>Cultivation</p> <p><b>Knockdown MoA</b><br/>Group C (bromoxynil)<br/>Group D (chloral dimethyl)<br/>Group G (pyraflufen, flumioxazin, saflufenacil, oxyfluorfen, carfentrazone)<br/>Group I (dicamba, fluroxypyr)<br/>Group L (paraquat, paraquat/diquat)<br/>Group L/Q (amitrole/paraquat)<br/>Group N (glufosinate)</p> <p><b>Residual MoA*</b><br/>Group C (flumeturon, prometryn, terbutylazine, diuron)<br/>Group D (pendimethalin, trifluralin)<br/>Group K (S-metolachlor, metolachlor)</p> | <p><b>Post-emergent*</b></p> <p>In-crop cultivation<br/>Manual chipping<br/>Rogueing</p> <p><b>Post-emergent OTT MoA</b><br/>Group A* (sethoxydim, clethodim, butoxydim, propaquizafop, haloxyfop)<br/>Group K (S-metolachlor**) <br/>Group Z (MSMA)</p> <p><b>Lay-by, directed or shielded spray</b><br/>Group C (flumeturon, prometryn, terbutylazine, diuron)<br/>Group D (pendimethalin)<br/>Group G (flumioxazin**) <br/>Group K (S-metolachlor)<br/>Group J (2,2-DPA)</p> | <p><b>Survivor control</b></p> <p>Aim for 100% control<br/>Cultivation, chipping or spot spraying</p>  |
| <p><b>In fallow survivor control</b><br/>Cultivation, chipping or spot spraying</p> <p>Refer above for NON glyphosate options<br/>Optical sprayer</p>   | <p><b>Pre-plant survivor control</b><br/>Cultivation, chipping or spot spraying<br/>flumioxazin<br/>carfentrazone-ethyl<br/>bromoxynil</p>   | <p><b>Survivor control</b></p> <p>Aim for 100% control</p> <p><b>Post-emergent</b><br/>Inter-row cultivation, chipping or spot spraying</p>   | <p><b>Comments</b></p> <p>Survivors MUST be controlled by another tactic prior to seed set.<br/>Read and follow label directions.<br/>Rotate herbicide Mode of Action (MoA).<br/>Come Clean. Go Clean. to stop weed seeds.<br/>Scout fields to look for weed survivors.<br/>Keep accurate field records.<br/>Ensure effective volunteer/ratoon management.<br/>DO NOT rely on glyphosate to control weeds in non-field areas<br/>Weed management requires a farming systems approach, winter, summer and non-field areas.<br/>*Group A herbicides already exhibit widespread resistance in several species. Controlling survivors is essential.<br/>**Mix full rates of different MoA, and rotate to alternative MoA in following years.<br/>*Refer to label for plant-back restrictions to following crop.<br/>** Limited formulations are registered for this use. Please check label.<br/>*** APVMA permits: 13460, 12941, 89322, 88120 for fallow use.</p> |
| <p><b>Post harvest</b></p> <p>Root cutting for crop destruction<br/>Cultivation, chipping or spot spraying<br/>carfentrazone-ethyl<br/>bromoxynil<br/>fluroxypyr</p>  |  |   |  |



of glyphosate for controlling summer weeds means that these weeds are only exposed to one mode of herbicide action. The Herbicide Resistance Management Strategy recommends at least two non-glyphosate tactics in summer fallows in addition to two non-glyphosate tactics in the cotton crop. Residual herbicides and double knock tactics provide good alternatives to a glyphosate-only fallow herbicide (see Table 18). To control larger weeds that may be tolerant to herbicides, a strategic cultivation or manual chipping is recommended. Field activities such as fertiliser placement and bed cultivators should be set up to have adequate soil disturbance to eradicate weeds during these mechanical tasks, and this will lessen the pressure to control weeds with further actions.

## In-crop tactics

### Pre-plant/at planting

Prior to planting there is an excellent opportunity to incorporate a non-glyphosate herbicide or combination of herbicides, or to integrate cultivation with a pre-planting operation such as seed bed preparation. In irrigation systems, consider utilising pre-irrigating to cause a flush of weeds to emerge and be controlled using a non-glyphosate tactic before the cotton emerges.

Herbicides from Groups C, G, I, L, M, N and Q can be used to target weeds that have emerged in the field (see the Integrated weed management diagram). This can be made more effective when used as a double knock.

Residual herbicides remain active in the soil for months and can act on successive weed germinations. This can be particularly effective in managing the earliest flushes of in-crop weed, when the crop is too small to compete. Broadleaf and grass weeds can be targeted with residual herbicides from Groups C, D or K.

Most residual herbicides need to be incorporated into the soil for optimum activity. Adequate incorporation of some residual herbicides is achieved through rainfall or irrigation, but others require cultivation. Soil surfaces that are cloddy or covered in stubble may need some pre-treatment such as light cultivation or burning to prevent 'shading' during application. Ash from burnt stubble may inactivate the herbicide, and therefore must be dissipated with a light cultivation or rainfall prior to herbicide application.

Crop safety is an important consideration when using residuals. How the herbicide moves in the soil following incorporation will depend on soil type, bed formation, solubility of the herbicide, the ability of the herbicide to bind to the soil and organic matter content, and the volume and timing of rainfall/irrigation, in addition to the method of applying irrigation. Growers can influence crop safety by the choice of herbicide, when it is applied, application rate, planting depth, planting date (to promote rapid crop establishment) and moisture management. Always follow label direction and if you are inexperienced in the use of residuals in cotton you should discuss your circumstances with your consultant, chemical supplier or the manufacturer.

The persistence of residual herbicides needs to be considered in order to avoid impacts on rotation crops. Persistence is determined by a range of factors including application rate, soil texture, organic matter levels, soil pH, rainfall/irrigation, temperature and the herbicide's characteristics. It

can be quite complex. For example, it is not usually the volume of rain, but the length of time the soil is moist that is the critical factor. Microbes that degrade many herbicides live near the soil surface and require moist soil to flourish. A couple of storms, where the soil surface dries out quickly won't contribute as much to residue breakdown as a period of rain that moistens the soil surface for days. Refer to product labels for more information. If you suspect that a residual may still be active in the lead up to planting, look for the presence of susceptible weeds in the treated paddock or pot up soil from treated and untreated areas, sow the susceptible crop and compare emergence. Where there is a concern, consider planting an alternative crop that is tolerant of the herbicide, or if cotton is to be used, plant the paddock last and pre-irrigate if it is to be irrigated. Always ensure that best practice is followed in terms of capture and management of runoff water.

### Post-emergence

Once cotton has emerged there are still many opportunities to incorporate different tactics. Check labels for application restrictions based on node development.

When targeting the over the top (OTT) application of glyphosate (Roundup Ready Flex<sup>®</sup>), aim to treat actively growing weeds, and do not allow weeds to become too large. Avoid using the same herbicide to control successive generations of weeds, and ensure survivors are not able to set seed. Do not apply more than the allowable number of OTT applications. Refer to the *Herbicide tolerant technology* section (page 89) for more information.

Grass selective herbicides (Group A) can be applied over the top of cotton, however repeated use is likely to lead to the development of Group A resistance. It is important that in managing glyphosate resistance, that resistance to other herbicides doesn't develop. Use Group A herbicides sparingly and ensure any survivors are controlled before they set seed, using another tactic, such as manual chipping.

**TABLE 20: Efficacy of knockdowns in winter fallows measured 6 weeks after treatment**

| Herbicide  | Weed control (%) |             |
|--|------------------|-------------|
|  | 1-month-old      | 3-month-old |
| Glyphosate + 2,4-D   | 84 (62-100)      | 76 (63-96)  |
| Glyphosate + Tordon 75-D <sup>®</sup>                              | 93 (86-99)       | 84 (62-98)  |
| Glyphosate + 2,4-D (fb) Spray.Seed <sup>®</sup>                    | 96 (93-100)      | 93 (87-97)  |
| Glyphosate + Tordon 75-D <sup>®</sup> (fb) Spray.Seed <sup>®</sup> | 99 (97-100)      | 97 (92-100) |
| Glyphosate + 2,4-D (fb) Alliance <sup>®</sup>                      | 96 (92-99)       | 90 (78-100) |
| 2,4-D (fb) Spray.Seed <sup>®</sup>                                 | 97 (97-98)       | 83 (68-97)  |
| 2,4-D#   | 88 (81-95)       | 53 (48-57)  |
| Amitrole <sup>®</sup> #  | 90 (84-95)       | 96 (95-97)  |
| Spray.Seed <sup>®</sup> #  | 84 (78-89)       | 22 (13-30)  |

Brackets indicate the range of efficacy across the experiments.

fb = followed by a 7-day interval

# = applied in only two of the four field experiments

Source: Steve Walker (QAAFI, University of Queensland),

Michael Widderick, Andrew McLean and Jeff Werth (Qld DAF).

**TABLE 19: Plant-backs to cotton for herbicides used in seedbed preparation**

| Active ingredient              | 2,4-D amine 700 g/L<br>(2,4-D amine 300 g/L) |                      |                      | dicamba 700 g/kg<br>(dicamba 500 g/L) |              |              | fluroxypyr 200 g/L<br>(fluroxypyr 333 g/L) |                |              | triclopyr<br>600 g/L |
|--------------------------------|--|----------------------|----------------------|---------------------------------------|--------------|--------------|--|----------------|--------------|----------------------|
| Rate L or g/ha                 | 0.5<br>(1.1)                                 | 0.5-1.0<br>(1.1-2.3) | 1.0-1.5<br>(2.3-3.4) | 140<br>(200)                          | 200<br>(280) | 400<br>(560) | 0.375<br>(0.225)                           | 0.75<br>(0.45) | 1.5<br>(0.9) | 0.16                 |
| Plant-back <sup>1</sup> (days) | 10   | 14                   | 21                   | 7                                     | 7            | 14           | 14   | 14             | 28           | 14                   |

<sup>1</sup> If applied to dry soil, at least 15 mm rain is required before plant-back period begins.

Some metolachlor registrations now include over the top use in-crop from 4 node up to 18 node crop growth and can be used with glyphosate to provide additional residual control of grass weeds. If leaf spotting is a concern, use a directed or shielded spray. Other lay-by/shielded spray options include herbicides in Groups C, D, G, K and J. Check label to confirm usage is allowed for each product in your situation and for crop safety directions.

In-crop cultivation, and (if required) chipping, provide important non-herbicide options for controlling herbicide survivors. Cultivating when the soil is drying out is the most successful strategy for killing weeds and will reduce the damage to soil caused by tractor compaction and soil smearing from tillage implements. Take care during set-up to minimise the crop damage. Inter-row cultivation can also increase issues with some soil-borne pathogens.

**Post-harvest**

Some weeds are likely to be present in the crop later in the season – even in the cleanest crop. These weeds will produce few seeds in a competitive cotton crop, but can be very problematic in skip-row configurations and can take advantage of the open canopy created by defoliation and picking. Removing crop residues and weeds as soon after picking as practical greatly reduces the opportunity for these weeds to set seed. See also *Management of volunteers and ratoons* (page 92).

**Notes on herbicide use**

Registration of a herbicide is not a recommendation for the use of a specific herbicide in a particular situation. The information in this publication is presented as a guide only to assist growers in planning their herbicide programs. Satisfy yourself that the product (or products) used is the best choice for the crop and weed. Carefully study the container label before using any herbicide, and note any specific instructions relating to the rate, timing, application and safety.

**IMPORTANT – avoid spray drift**

Take every precaution to minimise the risk of causing or suffering spray drift damage:

- Plan your crop layout to avoid sensitive areas.
- Ensure spray contractors have details of any nearby sensitive areas.
- Consult with neighbours to minimise risks from spraying near property boundaries (refer to SataCrop).
- Carefully follow all label directions.
- Pay particular attention to weather conditions before application.
- Use buffer zones to reduce the impact of spray drift or over-spray.
- Keep detailed records.
- Adhere to relevant 2,4-D application regulations.

See the *Spray application* section (page 123) for more details.

**For more information:**

Refer to IWM tactics diagram on page 76.

SataCrop website <https://satacrop.com.au>

The *Australian Cotton Production Manual* also includes information on weed control tactics.

**CottonInfo videos:**

Late season weeds [youtu.be/4gqLpEu5uTE](https://youtu.be/4gqLpEu5uTE)

Sources of weed seed [youtu.be/TLSpFIZZaqU](https://youtu.be/TLSpFIZZaqU)



**TABLE 21: Cotton herbicide plant-backs to rotation crops**

| Active ingredient        | Plant-backs from cotton to rotation crops (months) |                |        |      |                |           |       |              |          |        |           |           |         |       |         |          |            |             |        |           |         |
|--------------------------|--|----------------|--------|------|----------------|-----------|-------|--------------|----------|--------|-----------|-----------|---------|-------|---------|----------|------------|-------------|--------|-----------|---------|
|                          | Cereal grain-crops                                 |                |        |      |                |           |       | Legume crops |          |        |           |           |         |       |         |          |            | Other crops |        |           |         |
|                          | Barley   | Maize          | Millet | Oats | Sorghum        | Triticale | Wheat | Azuki bean   | Chickpea | Cowpea | Faba bean | Field pea | Lablaba | Lupin | Lucerne | Mungbean | Pigeon pea | Soybean     | Canola | Safflower | Linseed |
| Chlorthal dimethyl       | 8  | 8              | 8      | 8    | 8              | 8         | 8     | 8            | 8        | 8      | 8         | 8         | 8       | FH    | FH      | 8        | FH         | 8           | 8      | 8         | 8       |
| Diuron                   | 24   | 24             | 24     | 24   | 24             | 24        | 24    | 24           | 24       | 24     | 24        | 24        | 24      | 12    | 24      | 24       | 24         | 24          | 24     | 24        | 24      |
| Fluometuron              | 6  | 6              | 6      | 6    | 6              | 6         | 6     | 6            | 6        | 6      | 6         | 6         | 6       | 6     | 6       | 6        | 6          | 6           | 6      | 6         | 6       |
| Fluometuron + prometryn  | 6  | 6              | 6      | 6    | 6              | 6         | 6     | 6            | 6        | 6      | 6         | 6         | 6       | 6     | 6       | 6        | 6          | 6           | 6      | 6         | 6       |
| Halosulfuron-methyl      | 24   | 2              | 24     | 24   | 2              | 24        | 3     | 24           | 24       | 24     | 24        | 24        | 24      | 24    | 24      | 24       | 24         | 24          | 24     | 24        | 24      |
| Metolachlor              | 6  | 0              | 6      | 6    | 0 <sup>1</sup> | 6         | 6     | 6            | 6        | 6      | 6         | 6         | 6       | 6     | 6       | 6        | 0          | 6           | 6      | 6         | 0       |
| Norflurazon <sup>2</sup> | 24   | 21             | —      | 24   | 21             | 24        | 24    | —            | 3        | —      | 24        | —         | —       | —     | 21      | —        | 3          | —           | 18     | 18        | 27      |
| Pendimethalin            | 6  | 0 <sup>3</sup> | 12     | 12   | 12             | —         | —     | —            | —        | —      | —         | —         | —       | 6     | —       | —        | —          | 6           | —      | —         | —       |
| Prometryn                | 6  | 6              | 6      | 6    | 6              | 6         | 6     | 6            | 6        | 6      | 6         | 6         | 6       | 6     | 6       | 6        | 6          | 6           | 6      | 6         | 6       |
| S-metolachlor            | 6  | 0              | 6      | 6    | 0 <sup>1</sup> | 6         | 6     | 6            | 6        | 6      | 6         | 6         | 6       | 6     | 6       | 6        | 0          | 6           | 6      | 6         | 0       |
| Trifloxysulfuron sodium  | 6  | 22             | 22     | 6    | 22             | 22        | 6     | 22           | 18       | 22     | 7         | 22        | 22      | 22    | 9       | 15       | 15         | 22          | 22     | 22        | 22      |
| Trifluralin              | 12   | 12             | 12     | 12   | 12             | 12        | 12    | FH           | FH       | FH     | FH        | FH        | FH      | FH    | FH      | FH       | FH         | FH          | FH     | FH        | FH      |

1 Concep II treated seed only.

2 For rates up to 3.5 kg/ha. Where higher rates, up to 4.2 kg/ha are used, increase plant-back period by 6 months.

3 Maize can be resown immediately after use in a failed crop provided the seed is sown below the treated band of soil.

FH = following cotton harvest

— = no information

Further information available in Weed control in Summer and Winter Crop Publications from NSW DPI.





# MIX WITH THE BEST

**HASTEN** MADE FROM AUSTRALIAN  
CANOLA OIL AND OVER 48 CROSS LABELS.

**A HIGHLY TRUSTED AND WIDELY COMPATIBLE  
SPRAY ADJUVANT MADE FOR AUSSIE CONDITIONS.**

WWW.VICCHEM.COM | PH: 03 9301 7000

**VICCHEM**  
The Right Chemistry

# Key weeds

**W**atch out for these common weeds of cotton with a high risk of herbicide resistance. Make your herbicide decisions early and target young growth stages. Controlling larger plants with glyphosate may be too late, as seeds may have already set.



Graham Charles, NSW DPI

**Liverseed grass** (*Urochloa panacoides*) is a prolific seeding annual summer grass favoured by reduced tillage systems. Fresh seed has a strong dormancy and therefore most seed germinates in the season following seed production. Emergence usually occurs in one large flush in spring. Seed viability on the surface is short but persistence increases with seed burial.



Graham Charles, NSW DPI

**Awnless barnyard grass** (*Echinochloa colona*) is an annual summer grass germinating multiple times during the season and is capable of producing >40,000 seeds per plant. Dormancy breaks down after four to six months and seedlings emerge between October and March. In a zero-till system, stopping seed set will quickly reduce seed numbers in the soil.



Graham Charles, NSW DPI

**Flaxleaf fleabane** (*Conyza bonariensis*) is an annual or short-lived perennial plant germinating from the soil surface. It has become a difficult weed to control with the move to reduced or zero tillage. Individual plants can produce >100,000 seeds. Plants can grow to 1 metre tall and commonly have a branching habit. It can emerge in late autumn or early winter, growing very slowly above ground but at the same time developing an extensive root system allowing it to grow quickly when spring temperatures increase.



Graham Charles, NSW DPI

**Feathertop Rhodes grass** (*Chloris virgata*) is a small-seeded surface-germinating annual that can produce >40,000 seeds per plant. It germinates after small amounts of rain and can produce viable seed within 6 weeks of emergence.



Graham Charles, NSW DPI

**Sowthistle** (*Sonchus oleraceus*) is a surface-germinating, prolific seed producer, up to 25,000 per plant and with no innate dormancy (able to germinate straight away), although 30-50% of seed buried below 2 cm will persist and remain viable. It flourishes in zero-till situations and seed is readily dispersed by wind. Able to germinate and set seed year-round, it requires multiple control efforts.

**Annual ryegrass** (*Lolium rigidum*) is predominantly a winter annual that has developed widespread herbicide resistance to a number of modes of action. Reports are now emerging of germinations in irrigated cotton fields over spring/summer, exposing annual ryegrass to more herbicide pressure. Stopping seed set for two years will deplete the seed bank to very low levels.



Eric Koetz, NSW DPI

**Red pigweed** (*Portulaca oleracea*) is a prostrate annual herb germinating in spring and summer that often forms dense mats and flowers from September through to March. Red pigweed is becoming harder to control with glyphosate and is appearing as a more common weed in surveys of field edges. It spreads via very small seeds that normally don't move far from the parent.



Eric Koetz, NSW DPI

# Herbicide resistance management

The classification system for agricultural chemicals will be changing from mid-2021. The main reason is there are currently 25 MoA groups listed and with new chemistry to enter the market in the near future we will run out of letters. The Australian letter system will be replaced with the International numbering MoA classification. The alignment will commence in July 2021 and is anticipated to be complete by 2024. A mobile App is available to cross reference the current Australian letter system with the new numbering format. [hracglobal.com/tools/classification-lookup](http://hracglobal.com/tools/classification-lookup)

**W**eed populations are naturally genetically diverse. Due to this diversity it is likely that a small number of individuals may exist that are able to survive exposure to a particular herbicide mode of action (MoA). When a herbicide from this MoA is used upon the population, individuals that have this gene present may survive and set seed, whereas the majority of plants without the gene (susceptible plants) are killed. While it might only be one or two individuals surviving at first, continued use of the same MoA herbicide will result in an ever-increasing proportion of the population being able to survive those herbicide applications. In Australia, herbicides are currently grouped according to their MoA which is represented by a letter code on the label (see box above) and are ranked according to their resistance risk. Research has shown that weeds can develop resistance to any single control tactic used alone, not only herbicidal ones. For example, regular mowing of annual bluegrass, *Poa annua*, in golf courses selected strains for lower grass seed heads, which is essentially a resistance to mechanical control.

There are currently 521 unique cases (species x mode of action) of herbicide resistant weeds globally across 263 species (152 dicots and 111 monocots). Weeds have evolved resistance to 23 of the 26 known herbicide sites of action and to 164 different herbicides. Costs of weed control in cotton in the US have increased from \$150 to \$400 per acre due to the evolution of resistance to glyphosate (Group M). While historically the Australian cotton industry has had a strong integrated weed management system, the extensive use of herbicide tolerant (HT) cotton varieties since 2006 has meant that glyphosate now accounts for more than 70% of all herbicide used within cotton – herbicide resistance is a reality in the Australian cotton industry. 18 weed species have now been confirmed as glyphosate resistant (see Table 22), 11 of which occur widely in cotton farming systems. A NSW DPI survey has confirmed the presence of glyphosate resistant sowthistle (*S. oleraceus*) within at least three cotton growing regions.

#### For more information:

C. Preston, The Australian Sustainability Working Group. March 2020. [facebook.com/AGSWG](https://www.facebook.com/AGSWG)

I. Heap, The International Survey of Herbicide Resistant Weeds. March 2021. [weeds-science.org](http://weeds-science.org)

## Why the need for an industry-wide strategy?

Experience with conventional insecticide resistance has encouraged a proactive culture to resistance issues in the Australian cotton industry. The increased use of glyphosate and escalating incidence of resistance has brought about the need for an industry-wide Herbicide Resistance Management Strategy (HRMS). This strategy draws together available information, enabling growers to understand and manage the risks of herbicide resistance in Australian cotton farming systems.

## How was the HRMS strategy developed?

The strategy was developed by the TIMS herbicide technical panel to help the Australian cotton industry manage the risk of herbicide resistance, especially the risks associated with glyphosate. The strategy indicates how different combinations of weed control tactics affect the timeframe to resistance developing as well as their impact on the weed seed bank.

The modeling used as the foundation of the HRMS is based on barnyard grass control in glyphosate tolerant cotton where three over the top (OTT) glyphosate applications are made in any one season. The time to resistance developing and effect on the weed seed bank was predicted using combinations of weed control tactics used in-crop and in the summer fallow phase for both irrigated and dryland cropping systems. The models indicate that in irrigated cotton, crop competition provides higher weed control than in dryland systems. The model demonstrates that the weed control tactics used in the summer fallow phase have the greatest impact on the time to glyphosate resistance developing.

## HRMS review

The HRMS is reviewed annually by the Herbicide Resistance Tech Committee in light of weed field surveys, CCA and grower surveys and in consultation with registrants of glyphosate registered for over the top (OTT) use in Roundup Ready Flex.

**TABLE 22: Glyphosate resistant weeds in Australia**

| Weed species   | Year first documented | Confirmed populations |
|--|-----------------------|-----------------------|
| Annual ryegrass ( <i>Lolium rigidum</i> )            | 1996                  | 854                   |
| Awnless barnyard grass ( <i>Echinochloa colona</i> ) | 2007                  | 103                   |
| Liverseed grass ( <i>Urochloa panicoides</i> )       | 2008                  | 4                     |
| Flaxleaf fleabane ( <i>Conyza bonariensis</i> )      | 2010                  | 64                    |
| Windmill grass ( <i>Chloris truncata</i> )           | 2010                  | 11                    |
| Wild radish ( <i>Raphanus raphanistrum</i> )         | 2010                  | 2                     |
| Great brome ( <i>Bromus diandrus</i> )               | 2011                  | 5                     |
| Sowthistle ( <i>Sonchus oleraceus</i> )              | 2014                  | 23                    |
| Red brome ( <i>Bromus rubens</i> )                   | 2014                  | 1                     |
| Sweet summer grass ( <i>Moorochloa eruciformis</i> ) | 2014                  | 1                     |
| Prickly lettuce ( <i>Lactuca serriola</i> )          | 2014                  | 1                     |
| Feathertop Rhodes grass ( <i>Chloris virgata</i> )   | 2015                  | 5                     |
| Tridax daisy ( <i>Tridax procumbens</i> )            | 2016                  | 1                     |
| Tall fleabane ( <i>Conyza sumatrensis</i> )          | 2017                  | 3                     |
| Winter grass ( <i>Poa annua</i> )                    | 2017                  | 2                     |
| Willow-leaved lettuce ( <i>Lactuca saligna</i> )     | 2017                  | 2                     |
| Northern barley grass ( <i>Hordeum glaucum</i> )     | 2018                  | 2                     |
| Capeweed ( <i>Arctotheca calendula</i> )             | 2021                  | 1                     |

Preston, C. The Australian Glyphosate Sustainability Working Group. March 2020

### Why does the HRMS only focus on glyphosate?

The strong reliance on glyphosate in the current farming system and the increasing number of cases of glyphosate resistance has meant that for the first HRMS, glyphosate was the key focus. Tables 23 and 24 show resistance has also been identified in Group L (paraquat) and I (2,4-D) herbicides to weeds common in cotton farming systems. The HRMS has been updated with notes to address risks from other MoAs, especially the emerging concerns around Group A resistant grass species such as feathertop Rhodes grass, annual ryegrass and windmill grass.

It is important to note that the principles behind the strategy, particularly the use of a diverse range of tactics and the control of survivors are applicable to other groups as well as Group M (glyphosate). There is concern that glyphosate resistance may result in reliance on other herbicidal groups, leading to multiple resistance, for example, reliance on the repeated use of Group A grass selective herbicides can quickly lead to development of Group A resistance.

A NSW DPI survey has confirmed Group A resistance in windmill grass (*Chloris truncata*), feathertop Rhodes grass (*Chloris virgata*) and barnyard grass (*Echinochloa crus-galli*) in three cotton growing regions. The intent is to continue to expand the HRMS to incorporate other modes of action and multi-trait HR cotton varieties.



A patch of glyphosate resistant awnless barnyard grass, likely to have started near a road. Consider whole of farm use of herbicides. (T.Cook, NSW DPI)

### How to use the HRMS

Given that modelling shows glyphosate resistance takes around 19 years to develop when used alone in an irrigated cotton cropping system and 13 years in dryland, it is important to identify the likelihood of resistance development in your own operation. The HRMS enables you to determine which other weed control tactics can be incorporated into your management system by providing guidance as to how much extra time they will give you until resistance develops, and demonstrating the effect they will have on the weed seed bank, which is critical to effectively managing resistance.

### How do non-cropping areas relate to the HRMS?

Areas adjacent to cotton fields such as irrigation channels, head ditches, tail drains, roadways, fence lines and areas next to stock routes can be a significant entry source for resistant weed seeds. Where possible, use a range of tactics to manage weeds in these non-crop areas, and do NOT rely on glyphosate to manage weeds in these non-crop areas. Prevent survivors of herbicide application from setting seed.

### Why does the strategy include weed seed bank as well as herbicide resistance risk?

The key to good weed management is having low weed seed bank numbers. Not only does this reduce impact on the crop, but it also reduces the herbicide resistance risk. The more weed seeds present, the more likely that an individual containing herbicide resistance genes will be present and hence likely to become a problem.



Glyphosate resistant barnyard grass was confirmed in 2007. This infestation 'blow-out' was due to an extremely wet summer that prevented access to the paddock and hence no effective treatment at an early growth stage. (T.Cook, NSW DPI)

**TABLE 23: Species that have developed resistance to paraquat (group L) in Australia**

| Species                         | Common name           | Year confirmed | State           | Crop                | Resistance to other MoAs/herbicides |
|---------------------------------|-----------------------|----------------|-----------------|---------------------|-------------------------------------|
| <i>Hordeum glaucum</i>          | Northern barley grass | 1983           | Victoria        | Lucerne             | Diquat (L)                          |
| <i>Arctotheca calendula</i>     | Capeweed              | 1984           | Victoria        | Lucerne             | Diquat (L)                          |
| <i>Hordeum leporinum</i>        | Barley grass          | 1988           | Victoria        | Lucerne             | Diquat (L)                          |
| <i>Vulpia bromoides</i>         | Silver grass          | 1990           | Victoria        | Lucerne             | Diquat (L)                          |
| <i>Mitracarpus hirtus</i>       | Small square weed     | 2007           | Queensland      | Mangoes             | Diquat (L)                          |
| <i>Lolium rigidum</i>           | Annual ryegrass       | 2010           | South Australia | Pasture seed        | A/M – 2 populations                 |
| <i>Gamochoaeta pensylvanica</i> | Cudweed               | 2015           | Queensland      | Tomatoes, sugarcane |                                     |
| <i>Solanum nigrum</i>           | Blackberry nightshade | 2015           | Queensland      | Tomatoes, sugarcane |                                     |
| <i>Eleusine indica</i>          | Crowsfoot grass       | 2015           | Queensland      | Tomatoes, sugarcane |                                     |
| <i>Conyza bonariensis</i>       | Flaxleaf fleabane     | 2016           | NSW             | Grape vines         |                                     |

## Seed bank control/resistance risk

A high weed burden contributes to herbicide resistance risk, as the more weeds that are present, the more likely that a resistant individual will be present and multiply. Strategies are best aimed at driving down the seed bank and preventing seed bank replenishment.

### Seed bank control key:

- Very high control = <10 seeds/m<sup>2</sup>
- High control = 10-100 seeds/m<sup>2</sup>
- Medium control = 100-500 seeds/m<sup>2</sup>
- Low control = 500-1500 seeds/m<sup>2</sup>
- Very low control = >1500 seeds/m<sup>2</sup>

## Do I have to adhere to the HRMS?

The HRMS is not intended to be prescriptive, and is aimed to be an industry mechanism for communicating the herbicide resistance risks from different tactics. It has been designed to present the risk related to a range of tactic combinations, to allow growers and consultants to make their own informed decisions.

## Assessing your own risk

For a more detailed assessment of the resistance risks for individual paddocks, use Qld DAF's Online Glyphosate Resistance Toolkit, available via [www.cottoninfo.com.au](http://www.cottoninfo.com.au). This tool allows you to check what your current level of risk is for developing glyphosate-resistant weed populations on your farm. The tool allows you to enter information on your current practices (including crop rotation, crop density, and weed control tactics) and to identify which weed species you usually have to control. It will then calculate a glyphosate resistance risk score for the paddock, and a level of risk for each weed identified.

The Barnyard Grass Understanding and Management (BYGUM) tool, available via [www.cottoninfo.com.au](http://www.cottoninfo.com.au), enables the resistance risk from summer weed control to be considered in the context of economics and seed bank management. This weed management scenario testing tool combines biological, agronomic and economic factors to examine the economics of current summer grass management strategies, and compare with new tactics.

## What does herbicide resistance look like?

Resistance begins with the survival of one plant and the seed that it produces. Early in the development of a resistant population, resistant plants are likely to occur only in isolated patches. These are often surrounded by dead 'susceptible' plants of the same species, or other species usually



**Surviving glyphosate resistant awnless barnyard grass plants amongst dead susceptible plants and dead plants of other species.**

**TABLE 24: Weed species with populations resistant to 2,4-D (Group I)**

| Species                      | Common Name          | Year | State             | Situation     | Herbicide                  | Also resistant to MoAs |
|------------------------------|----------------------|------|-------------------|---------------|----------------------------|------------------------|
| <i>Raphanus raphanistrum</i> | Wild radish          | 1999 | Western Australia | Winter cereal | 2,4-D                      |                        |
|                              |                      | 2006 | South Australia   | Winter cereal | 2,4-D, MCPA                | B, F                   |
|                              |                      | 2009 | Victoria          | Winter cereal | 2,4-D                      | B                      |
|                              |                      | 2010 | Western Australia | Winter cereal | 2,4-D                      | B, F, M                |
|                              |                      | 2011 | Victoria          | Winter cereal | 2,4-D                      |                        |
|                              |                      | 2011 | New South Wales   | Winter cereal | 2,4-D                      |                        |
|                              |                      | 2013 | New South Wales   | Winter cereal | 2,4-D                      |                        |
|                              |                      | 2020 | Western Australia | Winter cereal | 2,4-D                      | B, H, K                |
| <i>Sisymbrium orientale</i>  | Indian hedge mustard | 2005 | South Australia   | Winter cereal | 2,4-D, MCPA                | B                      |
|                              |                      | 2015 | South Australia   | Winter cereal | 2,4-D                      | F                      |
|                              |                      | 2016 | Victoria          | Winter cereal | 2,4-D                      |                        |
|                              |                      | 2016 | Victoria          | Winter cereal | 2,4-D                      | B, F                   |
|                              |                      | 2015 | Victoria          | Winter cereal | 2,4-D                      |                        |
| <i>Sonchus oleraceus</i>     | Sowthistle           | 2015 | South Australia   | Winter cereal | 2,4-D, dicamba, clopyralid |                        |
| <i>Arctotheca calendula</i>  | Capeweed             | 2015 | South Australia   | Winter cereal | 2,4-D                      |                        |

C.Preston. The Australian Sustainability Working Group. March 2020.

controlled by the herbicide applied. This is the critical time to identify the problem. For other resistance mechanisms, the symptoms may appear as a 'sick' plant that subsequently recovers and may look similar to 'underdosing' or poor application. If a higher application rate is required to kill these individuals in subsequent years this indicates non-target site resistance is present. Many of the symptoms of herbicide resistance can also be explained by other causes of spray failure. Regularly maintain and calibrate spray equipment, spray under appropriate conditions and keep good records to ensure maximum spray efficacy. Resistant weed seeds can also be transported in to a management unit through irrigation channels, vehicle tyres, or blow in on the wind (in the case of species such as fleabane), and consequently can be relatively widespread before they are noticed.

### Why should I get weeds tested?

Testing a plant population for the presence of herbicide resistant individuals is an important tool for growers and advisors. The results from these tests can confirm if there is still efficacy within some of the MoA groups. Generally, seed is collected from the suspect plants and is sent for testing (see below). Bayer also offers free resistance testing – see page 90 for more details.

An alternative sampling method is to collect actual plants from the field for the 'Quick Test'. This process is limited to grass weeds only and is best targeted at seedlings or small plants as large numbers need to be collected and posted. Upon arrival they are potted up and once re-established, herbicide treatments are applied. In mid-summer conditions, plants are less likely to survive the trip than if collected in cooler times of the year. It is recommended to take seed samples from the surviving plants in summer and mark these sites to enable seedling collections in the following autumn or spring if they are needed. The timeline for obtaining results from sending seed samples can be several months. Results are usually available by the end of April when samples are received before January. When plants are sent for Quick Tests, results are usually available within 4-8 weeks.

#### Collecting seed samples:

- Collect 2000-3000 seeds from plants you suspect are resistant. Barnyard grass = 1 cup full.
- If testing >3 modes of action, collect additional seed.
- Avoid collecting large amounts of seed from just a few large plants.
- Follow a 'W' shaped pattern stopping every ~20 m if survivors are widespread. If survivors are localised, collect from within this area.
- Shake seed heads into a bucket to ensure only ripe seed is collected.
- Store samples in a paper bag at room temperature, away from sunlight, moisture and heat.
- Post as soon as possible.

#### Collecting plant samples for the Quick Test:

- For each mode of action to be tested: collect 50 plants/field from areas where you suspect resistance.
- Gently pull out plants and wash roots.
- Wrap in paper towel. Do not moisten.
- Place in waterproof plastic bag.
- Collect weeds early in the week, and Express Post as soon as possible. Do not store or post over the weekend. If plants cannot be posted on the same day, store overnight in the fridge.

### Sending samples to resistance testing services

Contact the testing service via their website and complete the sample registration request so they know to expect the sample. Follow the instructions above and send samples together with sample registration, contact details, field and weed management history and testing required to one of the testing services below

**Dr Peter Boutsalis** (seed or Quick Test)  
**Plant Science Consulting**  
22 Linley Avenue,  
Prospect SA 5082  
Phone: 0400 664 460  
Email: [info@plantscienceconsulting.com](mailto:info@plantscienceconsulting.com)  
Website: [www.csu.edu.au/weedresearchgroup/herbicide-resistance](http://www.csu.edu.au/weedresearchgroup/herbicide-resistance)

**Dr John Broster** (seed test only)  
**Charles Sturt University**  
Herbicide Resistance Testing Service,  
PO Box 588  
Wagga Wagga NSW 2678  
Phone: 02 6933 4001  
Email: [jbroster@csu.edu.au](mailto:jbroster@csu.edu.au)



An annual ryegrass survivor. (Eric Koetz, NSW DPI)

### How do I manage glyphosate resistant weeds?

The strategy to **manage** glyphosate resistant weeds is similar to the strategy to **prevent** glyphosate resistance – integrate a range of different tactics throughout the weed lifecycle to rapidly deplete the soil weed seed bank, and prevent further seed set/recruitment. This means that the HRMS is just as relevant to managing resistant weeds as it is preventing them. If detected early, managing known patches of herbicide resistant weeds by applying an intensive program of different tactics and ensuring weeds do not set seed, may be effective in preventing the problem from spreading.

Refer to **Integrated weed management section**.

For more information on herbicide resistance visit [www.weedsmart.org.au](http://www.weedsmart.org.au)  
**BYGUM** (barnyard grass understanding & management) [www.cottoninfo.com.au](http://www.cottoninfo.com.au)  
Contact: [david@innokasintellectual.com.au](mailto:david@innokasintellectual.com.au)

**CottonInfo videos:**  
Minimising glyphosate resistance [youtu.be/cke-mamGe2o](https://youtu.be/cke-mamGe2o)  
A demonstration of weed resistance [youtu.be/y7Ji1aiSLk](https://youtu.be/y7Ji1aiSLk)

# COTTON GROWING CALLS FOR THE RIGHT CHEMISTRY

## HASTEN<sup>†</sup>-COTTON SPRAY ADJUVANT

Patented technology including cottonseed oil & endorsed by leading defoliant brands as preferred adjuvant.

## IN-BOUND<sup>†</sup> SPRAY ADJUVANT

Highly refined Mineral Oil with over 20% Non-Ionic Surfactants. Provides excellent performance enhancement of certain agricultural products, including cotton defoliation.

## TRUMP<sup>†</sup> SPRAY OIL

A blend of highly refined paraffinic oil (USR>98%) & non-ionic surfactants for cotton defoliation and managing pests in summer crops.

**VICCHEM**  
The Right Chemistry

Dugald Macfarlane (*Nth NSW / Sth Qld Territory Manager*)

0421 901 424

Jeshua Smith (*Sth & Central NSW Territory Manager*)

0428 710 400

Jim Wark (*Business Development Manager*)

0429 149 039

Owen Connelly (*National Sales Manager*)

0427 129 572

# Herbicide Resistance Management Strategy

## Explanatory Notes:

2021-22

**The HRMS is designed as a tool to manage the risk of herbicide resistance in irrigated and dryland farming systems incorporating herbicide tolerant (HT) cotton, to delay glyphosate resistance.**

The strategy has been developed in response to the escalating problem of glyphosate herbicide resistance. This version of the HRMS focuses on a glyphosate tolerant cotton system; however the future availability of multi-trait herbicide tolerant varieties has not been considered in the design of the strategy, and may require a more sophisticated strategy to follow into the future.

### The formula to manage/delay glyphosate resistance

The most effective way to delay resistance is to use:

**2** non-glyphosate tactics targeting both grasses and broadleaf weeds during the cotton crop

+

**2** non-glyphosate tactics in summer fallow targeting both grasses and broadleaf weeds

and

**NO survivors**, control survivors of glyphosate applications and do not allow them to set seed.

### Increased time to resistance:

Research indicates that typically glyphosate failure may appear in grass weeds after 13 years (dryland) and 19 years (irrigated) in a glyphosate only system. Resistance in broadleaf weeds is slower to emerge and usually takes around 18 years in both irrigated and dryland systems when cotton is grown in rotation with a summer fallow. Glyphosate resistance is delayed by 4-6 years if residual + double knock regularly implemented in summer fallow.

**Cropping system** – The HRMS models two systems,

- Continuous back to back irrigated glyphosate-tolerant cotton with no summer fallow; and,
- Dryland glyphosate tolerant cotton grown every second year, alternating with long summer fallows.

With many farms now reporting glyphosate resistance, it is important to note that the strategies identified to avoid resistance are similar to those required to manage it. However, recent research has found that to eradicate populations, additional tactics such as patch management are required.

In the dryland scenario, rotation cropping should be considered similar to a fallow, with 2 non-glyphosate tactics recommended. Rotation crops provide an opportunity to incorporate other tactics, rotate herbicide groups, vary the time of year crop competition suppresses weeds and produce stubble loads that reduce subsequent weed germinations.

### In-crop tactics

- The control of survivors and use of two non-glyphosate tactics is critical to the HRMS.
- Aim for 100% control of glyphosate survivors after glyphosate application. Cultivation after glyphosate application is predicted to achieve 80% survivor control, whereas cultivation plus chipping is predicted to achieve 99.9% survivor control. Other tactics for survivor control could be equally effective, such as shielded or spot-spraying with an effective knockdown herbicide.
- A key principle of herbicide usage in an IWM system is to rotate herbicide groups.
- Residual herbicides need back up, such as tillage, chipping and non-glyphosate knockdowns. When using residuals, consider plant-back periods and crop safety.

### Summer fallow tactics

- Summer fallows (and rotations) may include any two non-glyphosate tactics such as residual or knockdown herbicides or tillage that are effective on the weed species present.

### Other management recommendations:

- Control weeds in adjacent areas (channels, tail drains, fencelines and roadsides) to minimise the seed bank and eliminate unknown weed seed sources. Do NOT rely on glyphosate to manage weeds in non-crop areas.
- Be aware of weed seed contamination sources (e.g. waterways, vehicle/machinery, and farm inputs). Establish and maintain COME CLEAN. GO CLEAN to prevent introduction and transport of resistant seeds.
- Monitor and follow up to ensure weeds that survive glyphosate applications are controlled using a non-glyphosate tactic before they are able to set seed. Get suspect weed survivors tested for resistance.
- Patch control – control weeds in isolated patches
- Use IWM best practice when employing tactics, including:
  - Regular scouting and correct weed identification;
  - Good record keeping;
  - Timely implementation of tactics;
  - Rotating herbicide mode of action groups;
  - Always following label recommendations; and,
  - Considering other aspects of crop agronomy.

### Assessing your own risk

Refer to page 84 of this publication for information on how to get weeds tested for resistance. For more information and tools on herbicide resistance and weed management in cotton refer to:

[www.cottoninfo.com.au](http://www.cottoninfo.com.au) or [www.weedsmart.org.au](http://www.weedsmart.org.au)



**Irrigated back to back cotton**

| Risk  | In-crop tactics<br>3 x OTT glyphosate applications<br>PLUS       | Seed bank<br>control | Comments   |
|---|--|----------------------|--|
| High - Resistance risk - Low →<br>↓ Decreasing - Survivor control - Increasing →<br>↓ | Very high survivor control after <b>each</b> OTT glyphosate      | Very high            | Control all survivors of OTT glyphosate applications. Don't use glyphosate alone to control the last in-crop flush                       |
|   | 2 x strategic in crop cultivations                               | Very high            | Time the second cultivation to control last weed flush and escapes prior to row closure  |
|   | Pre-plant residual plus residual layby                           | Very high            | Consider plant-back period restrictions  |
|   | Very high survivor control after <b>first</b> OTT glyphosate     | Very high            | Control survivors from first flush which has highest weed germination  |
|   | Grass selective in-crop herbicide + cultivation                  | High                 | Resistance to Group A herbicides may already be present in some populations. Controlling survivors is essential; follow with cultivation |
|   | Moderate survivor control after <b>first</b> OTT glyphosate only | Low                  | Survivors allowed to set seed will increase the speed of selection for resistance. Test survivors for glyphosate resistance              |
|   | Glyphosate only  | Very low             | Survivors allowed to set seed will increase the speed of selection for resistance. Test survivors for glyphosate resistance              |

**Dryland cotton every second summer**

| Risk  | Summer fallow tactics    | In-crop tactics<br>3 x OTT glyphosate applications PLUS     | Seed bank control in cotton phase | Comments   |
|---|--------------------------|---|-----------------------------------|--|
| High - Resistance risk - Low →<br>↓ Decreasing - Survivor control - Increasing →<br>↓ | 2 non-glyphosate tactics | Very high survivor control after <b>each</b> OTT glyphosate | Very high                         | The most effective scenario for delaying glyphosate resistance   |
|   | Glyphosate only fallow   | Very high survivor control after <b>each</b> OTT glyphosate | Very high                         | Very high frequency & efficacy of survivor control is required if in-crop only tactics are used  |
|   | 2 non-glyphosate tactics | Moderate survivor control after <b>each</b> OTT glyphosate  | High                              | Lower intensity in-crop tactics can give excellent results if backed up with robust control in summer fallows. Specific, frequent, well-timed control of glyphosate survivors provides long term resistance delay/management |
|   | Glyphosate only fallow   | 2 strategic cultivations                                    | Low                               | Time last cultivation to control late flushes and escapes  |
|   | Glyphosate only fallow   | Pre-plant residual + layby                                  | Very low                          | These tactics give limited increased time to resistance and poor seed bank control   |
|   | Glyphosate only fallow   | Moderate survivor control after <b>each</b> OTT             | Very low                          |  |
|   | 2 non-glyphosate tactics | Glyphosate only   | Very low                          |  |
|   | Glyphosate only fallow   | Glyphosate only   | Very low                          |  |

Glyphosate (Group M) resistance has been confirmed and is widespread in the following cotton weeds:

- Windmill grass
- Awnless barnyard grass
- Fleabane
- Sowthistle
- Feathertop Rhodes grass
- Liverseed grass
- Annual ryegrass is a significant issue in Southern valleys and is emerging as a problem in Northern NSW. There are reports of cross resistance to glyphosate and Group A herbicides.
- Group A resistance is widespread throughout broadacre farming systems and is increasing in cotton farming systems, especially in hard to control weeds such as feathertop Rhodes grass, annual ryegrass and windmill grass.
- Emerging herbicide resistance to Group L (paraquat) has been reported in other farming systems, especially in grasses. Resistance has not been reported in cotton farming systems, however the increase in double knock strategies makes it essential that **all survivors** of a double knock involving paraquat need to be controlled. Two populations of flaxleaf fleabane collected during surveys have tested as resistant to a glyphosate + paraquat double knock.
- Increasing use of Group I herbicides in summer fallows is a concern with a population of sowthistle reported as resistant to 2,4-D in winter cereals.
- Hit weeds where it hurts, use WeedSmart BIG 6 (page 74).

**TABLE 25: Resistance risk for herbicides used in cotton**

| Herbicide active ingredient         | Pre plant | At plant | Post plant | Mode of Action | Years to resistance | Resistance status |
|-------------------------------------|-----------|----------|------------|----------------|---------------------|-------------------|
| MSMA                                | N         | N        | Y          | Z              | N/A                 | Rare              |
| Amitrole + paraquat                 | Y         | N        | N          | Q+L            | >15                 | Rare              |
| Amitrole + ammonium thiocyanate     | Y         | N        | N          | Q              | N/A                 | Rare              |
| Paraquat                            | Y         | Y        | Y          | L              | >15                 | Occasional        |
| Paraquat + diquat                   | Y         | Y        | N          | L              | >15                 | Occasional        |
| Glufosinate-ammonium                | Y         | N        | N          | N              | 10-15               | Rare              |
| Glyphosate(a)                       | Y         | Y        | Y          | M              | >12                 | Widespread        |
| s-Metolachlor or Metolachlor (b)    | Y         | Y        | Y          | K              | >15                 | Rare              |
| 2,4-D (c)                           | Y         | N        | N          | I              | 10-15               | Occasional        |
| Dicamba (c)                         | Y         | N        | N          | I              | 10-15               | Rare              |
| Fluroxypyr                          | Y         | N        | N          | I              | 10-15               | Rare              |
| Fluroxypyr+ aminopyralid            | Y         | N        | N          | I              | 10-15               | Rare              |
| Triclopyr                           | Y         | N        | N          | I              | 10+                 | Rare              |
| Triclopyr + picloram                | Y         | N        | N          | I              | 10+                 | Rare              |
| Triclopyr + picloram + aminopyralid | Y         | N        | N          | I              | 10+                 | Rare              |
| Carfentrazone-ethyl                 | Y         | N        | N          | G              | 10                  | Rare              |
| Flumioxazin (d)                     | Y         | N        | Y*         | G              | 10                  | Rare              |
| Oxyfluorfen                         | Y         | N        | N          | G              | 10                  | Rare              |
| Saflufenacil                        | Y         | N        | N          | G              | N/A                 | Rare              |
| Norflurazon                         | Y         | N        | N          | F              | 15+                 | Occasional        |
| Pendimethalin (e)                   | Y         | Y        | Y          | D              | 10-15               | Occasional        |
| Trifluralin                         | Y         | Y        | N          | D              | 10-15               | Occasional        |
| Chlorthal dimethyl                  | Y         | Y        | Y          | D              | 10-15               | Occasional        |
| Bromoxynil                          | Y         | N        | N          | C              | 10-15               | Occasional        |
| Diuron                              | Y         | Y        | Y          | C              | 10-15               | Rare              |
| Fluometuron                         | Y         | Y        | Y          | C              | 10-15               | Rare              |
| Fluometuron + prometryn             | Y         | Y        | Y          | C              | 10-15               | Rare              |
| Prometryn                           | Y         | Y        | Y          | C              | 10-15               | Rare              |
| Isoxaflutole (c)                    | Y         | N        | N          | H              | 10                  | Rare              |
| Halosulfuron-methyl                 | N         | N        | Y          | B              | 4                   | Widespread        |
| Trifloxysulfuron sodium             | N         | N        | Y          | B              | 4                   | Widespread        |
| Butroxydim                          | N         | N        | Y          | A              | 6-8                 | Widespread        |
| Clethodim                           | N         | N        | Y          | A              | 6-8                 | Widespread        |
| Fluazifop-p                         | N         | N        | Y          | A              | 6-8                 | Widespread        |
| Haloxifop                           | N         | N        | Y          | A              | 6-8                 | Widespread        |
| Propaquizafop                       | N         | N        | Y          | A              | 6-8                 | Widespread        |

**Lowest resistance risk**
**Moderate resistance risk**
**Highest resistance risk**

**Group A herbicides already exhibit widespread resistance in several species. Controlling survivors is essential.**

**Always read the label for detailed use patterns and application rates.**

- a) Roundup Ready Flex® varieties only.
- b) Bouncer® and Dual Gold formulations.
- c) See label for rainfall required before plant-back period begins.
- d) Valor® formulation only.
- e) Rifle® formulations.

**Refer to Tables 18, 19 and 21 for plant-back periods.**

# Herbicide tolerant technology

## Roundup Ready Flex<sup>®</sup> technology

### Bayer Crop Science

#### How does Roundup Ready Flex cotton work?

The primary effect of glyphosate on plants is the inhibition of the production of EPSPS. EPSPS is an enzyme responsible for the production of amino acids essential for protein construction and plant growth.

Bayer CropScience identified a soil bacterium that produces a modified form of the EPSPS enzyme, the CP4 strain. The CP4 strain of EPSPS is not inhibited by glyphosate. Roundup Ready Flex cotton plants produce the modified form of EPSPS, so are able to continue producing amino acids and proteins after glyphosate has been applied. Roundup Ready Flex cotton contains two copies of the CP4 EPSPS gene and a promoter sequence resulting in expression in both the vegetative and reproductive parts of the plant. Roundup Ready Flex cotton is therefore able to tolerate applications of glyphosate in its vegetative (pre-squaring) and reproductive (squaring, flowering, boll development and maturation) stages. Only glyphosate products registered for usage in Roundup Ready Flex can be used over the top (OTT) of Roundup Ready Flex cotton. Usage must be in accordance with the label. Registered glyphosate products may be applied OTT of Roundup Ready Flex cotton up to four times between emergence and 60% boll open stage. One application is permitted OTT between 60% bolls open and harvest. However, the total amount of herbicide applied to any one crop must not exceed 6 kg/ha of Roundup Ready Herbicide with PLANTSHIELD<sup>®</sup> in a total of 4 applications as illustrated in the Application Guidelines figure (page 90). Crops that are intended for seed production must not have an application of glyphosate past the 60% bolls open stage.

The full-plant glyphosate tolerance of Roundup Ready Flex means that applications of glyphosate can be made irrespective of the rate of crop growth or the number of days between applications with no effect on fruit retention, fibre quality parameters or yield.

#### Weed management in Roundup Ready Flex

Roundup Ready Flex cotton offers growers an increased margin of crop safety, a more flexible window for OTT applications of glyphosate, and the potential to improve the efficacy of weed control. However Roundup Ready Flex cotton should be viewed as a component of an Integrated Weed Management (IWM) system, not as a solution to all weed management scenarios. Weed species with natural tolerance to glyphosate will be selected for with repeated glyphosate applications, resulting in species shift. The most effective, economic and sustainable weed management system for growers will, therefore, be achieved using an integrated (IWM) approach. Refer to the *Integrated weed management* section for more detailed information.

#### Know your field history

A combination of the relative effectiveness of previous herbicide programs and other agronomic practices employed on a farm is likely to influence the weed species present in any field. The correct identification and a basic understanding of the biology and ecology of the weeds

present in a field are essential elements in the design of a successful weed management program. It is critical that the appropriate herbicide and herbicide rate are chosen for the target weed species. By knowing field history, growers can determine which weed control tools they should use and when they should be employed to achieve the best results.

#### Pre-plant knockdown

Starting with a 'clean' field provides seedling cotton with the best possible conditions to emerge and to develop, unhindered by the competitive effects of weeds. Pre-plant weed control can be achieved using tillage and/or the appropriate registered herbicides. The use of glyphosate tank-mixes or herbicides with other modes of action is encouraged prior to planting to strengthen the IWM program. It is important that any cotton volunteers are controlled at this stage.

#### The role of residual herbicides

Residual herbicides should be used where appropriate in the Roundup Ready Flex system. The nature of pre-emergence residual herbicides often requires that they be applied in anticipation of a weed problem. Consideration for the use of residual herbicides in a weed control program for any given field should be determined based on the knowledge of the field's history.

#### The first OTT (over-the-top) application

Cotton is a very poor competitor and is sensitive to early season weed competition. The longer OTT window with Roundup Ready Flex may tempt growers to delay the first OTT application of a registered glyphosate product in the hope that multiple weed germinations can be controlled with a single spray. While competitive effects will vary according to weed species and weed density, it is commonly recognised that good weed control in the first 6-8 weeks following crop emergence maximises cotton yield potential. Delaying the initial OTT application may result in growers having to target weeds later in the season that are beyond the growth stage for optimum control. After the first spray is completed a thorough check of each field is recommended to note the presence of survivors and evaluate the efficacy of the spray.

#### Subsequent OTT applications

After the first OTT application, the use of subsequent OTT applications (up to a maximum of four), should be made according to the presence of new weed germinations. In any field, a mix of weed species will commonly exist. Correct identification of weeds is very important as this will have a direct impact on the rate selection and application timing(s) chosen. Select the timing and application rate of the registered glyphosate product based upon the most difficult to control weed species in each field. Refer to label for more information. Scouting of fields after each spray to evaluate the effectiveness of any glyphosate application, and to identify if any weeds have survived the spray is essential for monitoring and managing resistance.

#### Inter-row cultivation

Inter-row cultivation is a relatively cheap and non-selective method of weed control. In irrigated cotton, it also assists in maintaining furrows to facilitate efficient irrigation. In a Roundup Ready Flex crop, inter-row cultivation contributes to the diversity of weed control methods being employed and, as such, is a valuable component of an IWM strategy.

#### Lay-by residual application

Growers and their advisors are encouraged to scout fields prior to row closure and to combine these observations with their historical knowledge of individual fields to ascertain the need for a lay-by herbicide application. A lay-by application should be used on fields where there is an expectation of a significant emergence of weeds later in the season.

**Pre-harvest application**

One application of a registered glyphosate product may be made OTT between 60% boll open and harvest. In most circumstances, good weed control earlier in the crop should render the pre-harvest application redundant. However, if late season weeds are present, a pre-harvest application can be used to reduce seed set and improve harvest efficiency. Pre-harvest applications of glyphosate will not provide regrowth control in Roundup Ready Flex cotton.

**Post-spray surveillance**

Scouting of fields after each spray to evaluate the effectiveness of any glyphosate application and to identify if any weeds have survived the spray is essential for monitoring and managing resistance. Samples should be collected and sent for resistance testing. Surviving weeds should be controlled with an alternative tactic. To assist growers and consultants to confirm the status of suspected resistant weeds on farm, Bayer is offering a free herbicide resistance testing program for Australian cotton growers. The aim of the program is to provide access to free herbicide resistance testing for cotton growers and their consultants to assist decision making and IWM strategy development.

Testing is available to all farmers who have grown Roundup Ready Flex cotton in any of the previous 3 seasons. Growers can test five major weeds (annual ryegrass, awnless barnyard grass, liverseed grass, red pigweed, milk (sow) thistle) present in cotton systems for resistance to any of the key herbicides included in the Roundup Ready PLUS® program.


The Roundup Ready PLUS® program is designed to reward cotton growers who use herbicides sustainably and help slow or prevent the development of glyphosate resistance in key weed species. The program encourages growers to use a range of weed control practices through product recommendations, education and stewardship campaigns and financial rebates. Visit [www.roundupreadyplus.com.au](http://www.roundupreadyplus.com.au) or email [australia.cotton@bayer.com](mailto:australia.cotton@bayer.com) for further information.

Testing will be anonymous. Bayer will not be provided with grower identification information.

**Managing Roundup Ready Flex volunteers**

A major consideration in the development of an IWM plan for Roundup Ready Flex is the management of herbicide tolerant cotton volunteers. Plans need to be made to use cultural control options and herbicides with alternate modes of action in fallows and subsequent crops to control volunteers. Refer to pages 92-94 for more information.

**FIGURE A: Roundup Ready Flex cotton allows you to spray a registered glyphosate product over the top (OTT) of your cotton up to four times between emergence and 60% boll open stage with one application permitted OTT between 60% bolls open and harvest.**



## Application guidelines

### Timing options

The glyphosate label for OTT usage in Roundup Ready Flex permits:

- Applications in fallow, prior to sowing the Roundup Ready Flex crop, with the maximum rate applied dependent on the targeted weed/s. Application may be made by ground rig sprayer or by aircraft.
- Up to four applications of Roundup Ready herbicide with PLANTSHIELD or Roundup Ready® PL herbicide with PLANTSHIELD technology between crop emergence and 60% boll open stage, with a maximum of 1.5 kg/ha or 1.9 L/ha respectively, being applied in any single spray event.
- An option for a pre-harvest application, alone or in tank mix with a registered defoliant, once the crop is 60% open. The maximum herbicide rate for pre-harvest use is 1.5 kg/ha of Roundup Ready herbicide with PLANTSHIELD, or 1.9 L/ha of Roundup Ready PL herbicide with PLANTSHIELD. Application may be made by ground rig sprayer or by aircraft.
- Not more than four applications and 6.0 kg/ha of Roundup Ready herbicide with PLANTSHIELD or 7.6 L/ha of Roundup Ready PL herbicide with PLANTSHIELD may be applied through all growth stages of Roundup Ready Flex cotton in any one growing season.

### Over-the-top applications

Before an OTT application, it is absolutely essential to thoroughly decontaminate the sprayer of any products which might damage the crop, particularly sulfonylurea and phenoxy herbicides. For ground rig sprayers, a spray volume of 50-80 litres per sprayed hectare is recommended for optimum performance. Nozzles and pressure settings must be selected to deliver a minimum of a COARSE spray quality (American Society of Agricultural Engineers, ASAE S572) at the target. For aerial application, nozzles and pressure settings must also be selected to deliver a minimum of a COARSE spray quality at the target. A minimum total application volume of 40 L per hectare needs to be used. Do not apply by aircraft at temperatures above 30°C or if relative humidity falls below 35%.

**Other sources of information:** Please refer to [bollgard3.com.au](http://bollgard3.com.au) and [crop.bayer.com.au](http://crop.bayer.com.au) for further information on stewardship and resistance management.

[www.weedsmart.org.au](http://www.weedsmart.org.au)




**LDC**  
 Louis Dreyfus Company



**Head Office:** LDC Enterprises Australia Pty Ltd  
 Level 3, 1 Breakfast Creek Rd, Newstead Qld 4006  
 P O Box 2142, Fortitude Valley BC Qld 4006  
 T: 07 3253 5999

**Toowoomba Office:** 2A Phillip St  
 East Toowoomba Qld 4350  
 T: 07 4602 0355

### OUR COMMITMENT

Getting The Right Product, To The Right Location, At The Right Time.

#### Cotton Ginning and Marketing

|                  |              |
|------------------|--------------|
| Cathy Cook       | 0407 253 784 |
| Michael Thompson | 0407 712 880 |
| Andrew Cook      | 0428 441 676 |
| Juanita Park     | 0428 753 282 |
| Trent Raymond    | 0447 194 125 |
| Rodney StClair   | 0438 768 609 |
| Alex Dalton      | 0409 546 712 |
| Hollie Gall      | 0477 005 704 |

#### Cotton Seed

|                  |              |
|------------------|--------------|
| Dan Sullivan     | 0418 750 973 |
| Michael Thompson | 0407 712 880 |

YOUR  
 TRUSTED  
 PARTNER  
 SINCE  
**1851**

# Management of volunteer and ratoon cotton

**C**ontrolling volunteer and ratoon cotton is an essential part of good integrated pest and disease management and general farm hygiene. Maintaining a farm that is free of volunteer cotton (cotton that has established unintentionally) and ratoon cotton (cotton that has regrown from leftover root stock from a previous season) breaks the 'green bridge' that enables pests and pathogens to overwinter between seasons.

## Control of volunteers

Cultivation and herbicides are the two most common methods of volunteer cotton control. Both methods require the cotton plants to have germinated and emerged. Planning to control volunteers is a key part of an integrated weed management strategy and should consider issues such as rotational crops and other weeds present in the field. Reducing the amount of viable seed left in fields (through clean pick and stubble management) and around farm (through clean up after module removal and spillages) will reduce the amount of volunteers that germinate. It is also important to remember that volunteers and ratoons that are left to set seed will also contribute to additional volunteers.

## Cultural

Broadacre cultivation controls established seedling volunteers as well as larger volunteer cotton plants in a fallow, however effective control only occurs when applied across the entire field (both furrow and hill). Cultivation also manages other weeds, making it an excellent non-chemical option to include within an integrated weed management program. A disadvantage of this control method, particularly in dryland farming systems, is soil moisture loss. Cultivation is also relatively slow and can lead to soil damage if conducted at an inappropriate time.

### Best practice...

- Control volunteer cotton as part of an integrated weed management strategy.
- Control volunteer and ratoon cotton plants in crop and non-cropping areas.
- Target plants when small, using the appropriate herbicide option applied in a sufficient spray volume to achieve good coverage. Read all labels before use to confirm timing and rates. ALWAYS FOLLOW LABEL DIRECTIONS.
- Undertake crop destruction operations as soon as practical after picking to minimise the number of residual stalks that can regrow into ratoon cotton.
- Ensure implements are set up to cultivate both hill and furrow, to avoid leaving uncultivated strips.
- Manual removal (i.e. chipping) may be necessary where isolated plants remain in non-field areas.

Seedling volunteers can be controlled reasonably well with less invasive physical removal such as Kelly chains, which break the young stems and can be used close to planting. In-crop cultivation with sweeps is also effective on small volunteer cotton plants.

Rotation of crops enables the use of alternate and residual herbicides. Ensure good control is achieved as cotton plants hidden within subsequent crops can continue to host pests and diseases.

Where isolated plants remain in fallow or non-field areas such as along roadsides and fences, physical removal by chipping can be effective.

## Chemical

Pre-watering is a method used to encourage volunteer cotton germination prior to planting the next cotton crop, providing a window for appropriate herbicide control.

Most herbicides work best on seedling volunteers. Tables 26 and 27 provide a list of herbicide actives registered for the control of volunteer cotton plants. Not all products containing these actives will have volunteer cotton on the label, so refer to the label for specific use information and ensure all label directions are followed.

When using contact herbicides to control volunteers, excellent spray coverage is essential. Coverage may be compromised by the presence of nearby stubble, lint or other weeds.

Do not apply glyphosate to volunteer cotton seedlings in fields that have previously been planted to Roundup Ready Flex<sup>®</sup> cotton.

For more detailed information on chemical options for controlling volunteer cotton, refer to section F4 in WEEDpak.



**Volunteer cotton plants enable pests to survive between seasons.** (Lewis Wilson)

## Ratoon cotton

Ratoon cotton is normally a product of minimum tillage where either conventional cotton is double-cropped back to a winter cereal, or consecutive cotton crops are grown. Mealybug hotspots are often associated with ratoon cotton plants.

In theory, ratoon cotton should not occur as harvested cotton is required to be controlled with adequate cultivation and soil disturbance as soon as practical after picking (usually mulching and/or root cutting followed by cultivation to destroy the root system).

In conducting this cultivation an additional aim is to destroy overwintering *Helicoverpa* pupae. This pupae control is a strategy in managing insecticide resistance for the cotton industry and is mandatory for

**Exirel<sup>®</sup>**  
INSECTICIDE

*To perfect this*



**Now a more  
affordable  
solution.**

*you need to protect this.*

**Exirel<sup>®</sup> insecticide provides outstanding and extended crop protection.**

When applied early in pest infestation cycle, Exirel<sup>®</sup> helps to keep pest populations below damaging levels. It has cross spectrum activity to control Cotton bollworm, Native budworm, Silverleaf whitefly and Cotton aphid. Pests stop feeding rapidly for quick protection. Exirel<sup>®</sup> also has selectivity for key beneficial insects making it ideal for IPM programs. **Visit [www.fmccrop.com.au](http://www.fmccrop.com.au) for more information.**

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. Copyright © 2021. All rights reserved. The FMC Logo and Exirel<sup>®</sup> are Trademarks or Registered trademarks of FMC Corporation or its affiliates.



An Agricultural  
Sciences Company



FMC Australasia Pty Ltd  
Phone: 1800 066 355  
[www.fmccrop.com.au](http://www.fmccrop.com.au)

Bollgard 3 crops where the first defoliation occurs after March 31. Thorough crop destruction can be particularly challenging in a zero-till situation, where the only soil disturbance is pupae busting. This operation should be conducted carefully to minimise the number of residual stalks that can regrow the following spring.

Ratoon cotton plants (regrowth/stub cotton) that have survived crop destruction can be difficult to control, having developed a large root system and small leaf surface area. Three herbicide options are available for the control of large volunteer cotton or ratoon cotton amongst stubble or in fallow. These options are registered for both optical booms and broadcast application, enabling all growers to conserve soil moisture. Refer to Table 27 for further information.

### ALWAYS FOLLOW LABEL DIRECTIONS

For more information (see [www.cottoninfo.com.au/publications](http://www.cottoninfo.com.au/publications)):

\* *Australian Cotton Production Manual: Chapter 13, Integrated Disease Management and Chapter 23, Managing cotton stubble/residues.*

- WEEDpak
- Managing ratoons and volunteers

CottonInfo videos ([www.youtube.com/cottoninfoaustr](http://www.youtube.com/cottoninfoaustr)):

- Keep your farm free from pests, weeds and diseases: Come Clean. Go Clean. [youtu.be/gR8hf8-hYQA](https://youtu.be/gR8hf8-hYQA)
- Rogue cotton plants in the farming community [youtu.be/CJP14\\_swggE](https://youtu.be/CJP14_swggE)
- Checking your farm for volunteer plants [youtu.be/9C2Uf5mUxk](https://youtu.be/9C2Uf5mUxk)



**Inadequate end of season crop destruction can lead to ratoon cotton. (Murray Sharman, Qld DAF)**

**TABLE 26: Herbicides for control of volunteer cotton**

| Active ingredient    | MoA group | Comments (always refer to product labels)   |
|----------------------|-----------|---|
| Amitrole + paraquat  | Q + L     | Can be applied after an initial spray of a glyphosate herbicide (double knockdown). Refer to label for spot spray rates.  |
| Bromoxynil           | C         | Apply in minimum of 80 L/ha water for Roundup Ready cotton. Refer to label for rain-fastness and restrictions on spray quality & condition.   |
| Carfentrazone-ethyl  | G         | Apply minimum spray volume of 80 L/ha to ensure effective coverage. To broaden weed spectrum may be tank mixed with the recommended rate of a knockdown herbicide. Refer to label for adjuvant recommendation.      |
| Paraquat + diquat    | L         | Apply in 50-100 L water/ha. Avoid spraying under hot dry conditions. For best results, spray in the evenings or in humid conditions.  |
| Flumetsulam          | B         | May be banded (>40%) over the row or broadcast. Minimum spray volume 150 L/ha for optimum results.  |
| Flumioxazin          | G         | Do not apply post-sowing pre-emergent. Do not sow crops for at least one hour after application. Can be tank mixed with glyphosate to control other weeds that may be present. Refer to label for adjuvant details. |
| Glufosinate-ammonium | N         | Good coverage is essential. Do not apply more than three applications per season. Best results are achieved when applied under warm humid conditions.   |
| Metribuzin           | C         | Registered for control of volunteer cotton in pigeon pea. Refer to label for critical comments.   |
| Fluroxypyr           | I         | Summer fallow.  |
| Saflufenacil         | G         | Do not apply post-sowing pre-emergent. Always apply with adjuvant or high quality methylated seed oil. See label for mandatory no-spray zone and spraying rates.  |
| Pyraflufen-ethyl     | G         | Prior to sowing summer crop or starting a summer fallow. Apply by ground rig only. Good spray coverage is essential. Do not sow crops for at least 1 hour after application.  |

For any changes to labels, refer to [portal.apvma.gov.au/pubcris](http://portal.apvma.gov.au/pubcris)

**TABLE 27: Herbicides for control of large (15 to 30 node) volunteer cotton and ratoon cotton in fallow**

| Active ingredient | MoA group | Rates                                  | Comments  |
|-------------------|-----------|--|---|
| Fluroxypyr        | I         | 1 L/ha followed by 1 L/ha OR           | For control of large cotton plants or ratoon cotton a sequential application of Comet followed by Comet is required for maximum control. Ensure sufficient leaf regrowth has occurred on the ratoon cotton to maximise chemical uptake.   |
|                   |           | 1 L/ha followed by Shirquat 2 L/ha OR  | For control of large cotton plants or ratoon cotton a sequential application of Comet followed by Shirquat is required for maximum control. The sequential application interval should be 7-14 days. Ensure sufficient leaf regrowth has occurred on the ratoon cotton to maximise chemical uptake. |
|                   |           | 1 L/ha + 1 L/ha Amicide Advance 700/ha | For a single pass operation apply Comet + Amicide Advance 700. Ensure sufficient leaf regrowth has occurred on the ratoon cotton to maximise chemical uptake.   |

Refer to the Comet 400 registration label for further details on control rates for optical spot spray technologies. Note that control rates are based on L/ha for broadcast application and L/100L (spot spraying rate) for optical sprayers. For any changes to labels, refer to [portal.apvma.gov.au/pubcris](http://portal.apvma.gov.au/pubcris)





# Parachute®

INSECTICIDE

## An Essential Partner for IPM and Weed Control

**Parachute® takes the science of paraffinic oil formulation to new heights.**

With an unrivaled combination of paraffinic loading, molecular weight efficacy and hyper-spreading of hydro-carbons, Parachute® works in partnership with beneficial insects to suppress chewing and sucking pests of Cotton and other broadacre crops, whilst also getting the most out of defoliants at the end of the season.

**Visit [www.fmccrop.com.au](http://www.fmccrop.com.au) for more information.**

ALWAYS READ AND FOLLOW LABEL DIRECTIONS. Copyright © 2021. All rights reserved. The FMC Logo and Parachute® are Trademarks or Registered trademarks of FMC Corporation or its affiliates.



An Agricultural  
Sciences Company



FMC Australasia Pty Ltd  
Phone: 1800 066 355  
[www.fmccrop.com.au](http://www.fmccrop.com.au)

# Integrated Disease Management (IDM)

## Developing an IDM strategy

A plant disease occurs when there is an interaction between a plant host, a pathogen, and the environment. Effective IDM involves a range of control strategies which must be integrated with management of the whole farming system.

Most disease control strategies should be implemented regardless of whether or not a disease problem is evident, as the absence of visible symptoms does not necessarily indicate an absence of disease.

Appropriate disease management does not always mean complete control of the disease through eradication of the pathogen, but rather aims at improving the resilience of the farming system. Many strategies may help to reduce the disease burden for subsequent crops and minimise the risk of diseases spreading within and between farms and regions.

## IDM at planting

### Optimal seed bed conditions

- Check field history for disease problems and avoid back-to-back cotton.
- Prepare firm, high beds to optimise establishment and seedling vigour.
- Position fertiliser and herbicides in the bed to avoid root damage.
- Ensure fields have good drainage to avoid waterlogging.

### Sowing date/temperature

Sowing in cool and/or wet conditions slows plant growth and favours disease development. Aim to plant when soil temperatures are 16°C at 8am and rising for the next 7 days. The *Australian Cotton Production Manual* (ACPM) includes information on crop establishment.

Also see the CSD Faststart Cotton Soil Temperature Network for your region <https://www.csd.net.au/soil-temperature>

### Variety selection

A number of varieties have some resistance to Verticillium wilt or Fusarium wilt (indicated by higher V rank and F rank respectively). Consider the disease status of each field when selecting varieties. Also consider the variety's seedling vigour, particularly when pre-irrigating or planting early. Refer to CSD's variety notes for more information ([www.csd.net.au/disease-ranks](http://www.csd.net.au/disease-ranks)).

When black root rot is present, use the more indeterminate varieties that have the capacity to catch up later in the season.

Different seed treatment options are also available and must be chosen when seed is ordered. See the CSD website for more information.

### Replanting

Make replant decisions early, based on stand losses, not on seedling size. Refer to the ACPM Crop establishment chapter for more information.

## IDM in crop

### Fungicides

All cotton seed sold in Australia for planting is treated with a standard fungicide for broad-spectrum disease control. See Table 28 (page 108) for a list of fungicides registered in cotton or available under permit.

## Best practice...

- Monitor in-crop and over time for disease symptoms. Identify areas with a history of disease.
- Select disease resistant varieties where possible.
- Practice IDM at planting, in-crop and post-harvest.
- Implement a reporting system for unusual symptoms.
- Be aware of and control insect vectors according to industry thresholds.
- Implement good farm hygiene and biosecurity practices.

## Irrigation scheduling

Early season, the temperature of irrigation water is considerably lower than soil temperatures, so pre-irrigating fields before planting can avoid cold shock and provide better conditions for seedling emergence, provided the hills or beds sub-up adequately. Irrigations late in the season to extend plant maturity can result in a higher incidence of Verticillium wilt.

Irrigate as required if disease has weakened the root systems and there are signs of water stress, but avoid waterlogging, particularly when temperatures are cool. Manage tail water to minimise the risk of spreading pathogens; if possible water the least infected fields first. General traffic, sprays, and picking should also follow the same order.

## Agronomic management

High planting rates can compensate for seedling mortality, but a dense canopy favours development of bacterial blight, Alternaria leaf spot, Sclerotinia and boll rots. Optimise nutrition and irrigations and consider the use of growth regulators when required.

If black root rot is present, either manage for earliness to get the crop in on time (short season areas) or manage for delayed harvest to allow catch up (longer season areas).

## Balanced crop nutrition

A healthy crop is more able to express its natural resistance to disease. Excessive nitrogen will favour Fusarium and Verticillium wilt, and will also greatly increase the risk of boll rot, particularly in fully irrigated situations. Potassium is important for natural plant defences and a deficiency can lead to the expression of more severe disease symptoms. Refer to NUTRIpak or the ACPM for more information on cotton nutrition.

## Conduct your own in-field disease survey

Monitoring and recording each disease's incidence, prevalence, and severity allows you to make a comparison over time and adjust your management strategies accordingly. The following section includes examples of the key diseases in cotton and their common symptoms.

If disease is suspected, sample healthy plants as well for submission to cotton pathologists. Occasionally, symptoms of nutrient deficiency can be misinterpreted for disease, so if the disease diagnostics turn out to be negative, consider a nutritional analysis.

**Early season:** Start monitoring when seedlings reach the 2 true leaf stage. Look for plants with signs of poor vigour or unusual symptoms. Examine roots by digging up the seedlings – DO NOT pull seedlings as this may damage the root system, or remove characteristic symptoms or infected parts of the roots. Examine multiple plants at multiple locations as diseases are often unevenly distributed.

Compare number of plants established per metre with number of seeds planted. Refer to the ACPM for replanting considerations.

**During the season** conduct disease surveys as frequently as possible. Look for plants that show unusual symptoms, have poor vigour, or have died.

Cut both diseased and healthy-looking plants in several places along the stem and look for discolouration. Plants infected with *Verticillium* or *Fusarium* wilt may display minimal or no external symptoms, particularly in varieties with high disease rankings.

Perform a **late season** check after the final irrigation but before defoliation to give an indication of the disease burden that field has been under and assess disease risk in the following years.

**If a suspect plant is located, please contact your local CottonInfo REO (see inside back cover) for more information or to determine the appropriate pathologist and process for submitting a sample.**

## IDM post harvest

### Control alternative hosts and volunteers

Having a host free period prevents build-up of pathogen inoculum and carryover of disease from one season to the next. The pathogens that cause *Verticillium* wilt, *Fusarium* wilt, tobacco streak virus (TSV), *Alternaria* leaf spot, some rots, and cotton bunched top can also infect common weeds found in cotton growing areas. Refer to WEEDpak F5 for more details.

It is particularly important to have a host-free period as some diseases, such as cotton bunched top, can only survive on living plants. Controlling alternative hosts, especially cotton volunteers and ratoons will help reduce the risk of quality downgrades and yield loss.

Utilise the off-season to rotate herbicide chemistries and explore alternate mechanical means of weed control.

**CottonInfo video on checking your farm for volunteer plants [youtu.be/9C2Utt5mUxk](https://youtu.be/9C2Utt5mUxk)**

### Manage crop residues

The pathogens that cause *Verticillium* wilt, *Fusarium* wilt, black root rot, boll rots, seedling disease and *Alternaria* leaf spot can all survive in cotton and some rotation crop residues. Carefully manage crop residues to minimise carryover of pathogens into subsequent crops.

If *Fusarium* wilt is known to be present in a field, residues should be slashed and retained on the surface for at least one month (60-120 days if possible) prior to mulching, in order to disinfect the stalks through UV light exposure.


In all other circumstances (including the presence of *Verticillium* wilt and other diseases), incorporate crop residues as soon as possible after harvest to promote faster decomposition.

### Use crop rotations to assist in managing disease

Successive crops of cotton (or other host species) can contribute to a rapid increase in disease incidence, particularly if susceptible varieties are used. Employ a sound crop rotation strategy using crops that are not hosts for the pathogens present.

The pathogen that causes *Verticillium* wilt has a large host range, and most legume crops host the black root rot and *Fusarium* wilt pathogens. Some alternative crops such as vetch, canola and mustards can provide a biofumigation effect against black root rot under specific management regimes.

Australian trial work with cotton has shown that crop rotation influences the abundance and diversity of soil microbial populations. After two years



**Aerial photo of Fusarium wilt damage.**

of rotation, *Verticillium* wilt disease levels were lowest in cotton following bare fallow compared to corn, sorghum or continuous cotton. However, soil analysis indicated the fallow also had the lowest diversity of soil microbial communities which is not good for long term soil health.

Cotton is highly dependent on arbuscular mycorrhiza fungi (AMF) which form beneficial associations with plant roots and can act as agents in nutrient exchange. Bare fallow for more than 3 to 4 seasons or removal of top-soil (especially more than 40 cm), such as in the development of a bankless channel system, may result in a severe lack of AMF. A cereal or green-manure crop may restore sufficient mycorrhizal fungi.

Pages 78-79 of the ACPM list the potential disease implications of rotation crops with cotton and can assist in developing a rotation strategy.

## IDM all year round

### Control insect vectors

Diseases caused by a virus or phytoplasma are often preventable by controlling the vector that carries the pathogen. Cotton bunched top (CBT) can be transmitted by aphids feeding on infected plants before migrating to healthy plants. Transmission of TSV to plants relies on the virus from infected pollen entering plant cells through the feeding injury caused by thrips. Consider integrated pest management principles and resistance risks when managing vectors.

Viruses can only survive in living plants. Control of cotton ratoons and volunteers throughout winter will reduce pathogen levels and also lower vector insect populations, drastically reducing disease risk.

### On-farm biosecurity and hygiene

Minimise the risk of moving pathogens on or off your farm, from field to field or farm to farm by considering vehicle and machinery movements within the farm. Have a strategy for ensuring clean movement of vehicles and machinery onto and around the farm. Minimise spillage and loss when transporting modules, hulls, cotton seed or gin trash.

Ensure all staff, contractors and visitors are aware of the requirements and your commitment to 'Come Clean. Go Clean.' before entering the farm.

**Useful resources:**

[www.cottoninfo.com.au](http://www.cottoninfo.com.au) and [www.mybmp.com.au](http://www.mybmp.com.au)

**CottonInfo video:**

**Keep your farm free from pests, weeds and diseases: Come Clean. Go Clean.**  
[youtu.be/qR8hf8-hYQA](https://youtu.be/qR8hf8-hYQA)





An industry partnership connecting growers with research and best practice.

**Keeping Australian cotton growers out-standing in their field.**



*Best Practice*



For more, visit [www.cottoninfo.net.au](http://www.cottoninfo.net.au) and [www.mybmp.com.au](http://www.mybmp.com.au).

# Key diseases of cotton

## Alternaria leaf spot (ALS)

Pathogen: *Alternaria alternata* and other minor spp.

### Symptoms

Cotton plants are prone to infection at all growing stages, but are most susceptible as seedlings and late in the season as the crop begins cut-out. Symptoms start as pinhead spots that develop into necrotic lesions surrounded by a purple halo. They can be seen on cotyledons, young to mature leaves, squares and bolls. Under favourable conditions, lesions continue to enlarge and coalesce to form irregular shapes. Severe infection may result in leaf desiccation.

### Favoured by

Spores can only germinate when there is several hours of free moisture on the leaf surface. Epidemic development is therefore favoured by either repeated heavy dews or extended periods of wet weather.

- Symptom development is suppressed by periods of very hot weather.
- Symptom development is favoured by physiological or nutritional stress e.g. heavy fruit load or premature senescence.
- Back-to-back fields are at greater risk of developing the disease.

### Host range

*A. alternata* has a wide host range and is globally distributed. Around 100 plants species have been reported as hosts, however, some strains show host specificity.

### IDM tactics

- Avoid planting into fields containing unincorporated infected residues from a previous crop.
- Provide balanced crop nutrition (especially potassium).
- Manage crop to avoid extremely rank growth.
- Apply registered or permit foliar fungicides according to label.
- Incorporate crop residues as soon as possible after harvest.
- Control alternative weed hosts, volunteer and ratoon cotton plants before, during and post season.



**Alternaria leaf spot of seedlings. Cotton trash is a common source of Alternaria inoculum.**



**Typical lesions associated with Alternaria leaf spot of cotton. (Linda Scheikowski, Qld DAF)**

## Black root rot (BRR)

Pathogen: *Berkeleyomyces rouxiae*  
(formerly *Thielaviopsis basicola*)

### Symptoms

Affected crops may be slow growing or stunted, especially early season. The disease causes destruction of the root cortex (outer layer), seen as blackening of the roots. Some roots may die but the fungus does not kill seedlings by itself. Severe black root rot can increase the susceptibility of plants to *Pythium* or *Rhizoctonia* spp. Plants that are badly affected early in the season may not continue to show symptoms later in the season as the dead cells of the root cortex are sloughed off when growth resumes in warmer weather.

### Favoured by

- Cool wet conditions – soil temperatures below 20°C are most favourable, but infection will still progress at temperatures up to 25°C.
- Medium to heavy clay soils.
- Back-to-back cotton or following susceptible crops including most legumes.

### Host range

All cotton varieties are susceptible, as are most legumes, including faba bean, soybean, cowpea, field pea, chickpea, mungbean, lablab and lucerne. *Datura* weeds (thornapple, castor oil) are also hosts. Non-hosts include cereals, sunflower, brassicas such as canola and broccoli, and onions.



**Black root rot on seedlings.** (Duy Le, NSW DPI)

### IDM tactics

- Choose more indeterminate varieties that have the capacity to 'catch up' later in the season.
- Use Bion® Plant Activator seed treatment to assist the plant's natural resistance mechanisms.
- Plant into well prepared, high, firm beds.
- Pre-irrigate/plant into moisture in preference to 'watering up' to avoid a cold shock.
- Time sowing to avoid cool soil temperatures (<18°C) if possible. Sow early only if conditions are warm enough (minimum soil temperature of 16°C at 8am and rising over consecutive days). Temperature measurements should be taken in all fields.
- Make replanting decisions on the basis of stand losses, not the size of the seedlings. Watch for early onset of water stress and irrigate accordingly, but avoid waterlogging.
- Anticipate delayed growth and later maturity and manage the crop accordingly (black root rot 'steals' time from your crop).
- Minimise tailwater.
- Practice good farm hygiene – Come Clean. Go Clean.
- Crop rotations have many benefits and increase microbial diversity in the soil, but will not reduce black root rot inoculum levels (apart from a rice rotation, where prolonged flooding can reduce inoculum levels).
- Rotate with biofumigation crops such as vetch or mustard between consecutive cotton crops or after a wheat fallow. Success depends upon the growth of the biofumigation crop and good incorporation (at least four weeks before planting cotton). Check that the biofumigant is not a host of other cotton diseases.
- Avoid legumes.
- Control weeds, cotton volunteers, and cotton ratoons.
- Flooding of fields for 30 days when maximum temperatures are above 30°C significantly reduces the population of *B. rouxiae*. Even better results will be seen if fields are flooded for 60 days.

## Boll rot, tight lock and seed rot

Boll rots are caused by a number of pathogens, including fungi and bacteria. Tight lock refers to a type of boll rot, where the lock remains hard and fails to fluff out. The term seed rot is used to describe a boll rot which begins in the seed.

### Phytophthora boll rot

Infected bolls quickly turn brown and become blackened (sometimes with areas of white mould on the surface before opening prematurely). The locks, which remain compact and do not fluff out, can be easily dislodged and fall to the ground. Symptoms are most prevalent on the lower bolls. *Phytophthora* boll rot usually occurs when soil is splashed up onto low bolls that are beginning to crack open or when low bolls are subject to inundation by tail water backing up into rows.

### Sclerotinia boll rot

Characteristically has black fungal structures (2-10 mm diameter) within and/or on the surface of the rotted bolls. A white fluffy fungal growth may be present and the branch adjacent to the boll may also be affected. The sclerotia germinate to produce apothecia (small cream coloured 'golf tees') which release clouds of microscopic spores that can infect the plant via dead or dying tissue (e.g. flower petals). The fungus then grows into healthy plant tissue such as the developing boll and down the fruiting branch towards the main stem.

CottonInfo video on *Sclerotinia* in cotton [youtu.be/yt\\_542Kj3\\_0](https://youtu.be/yt_542Kj3_0)



**Sclerotinia boll rot is more likely to be a problem in wet seasons as spores can be moved around the plant through rain-splash.** (Linda Smith/Tony Cooke)

### Fusarium boll rot

Not to be confused with Fusarium wilt, Fusarium boll rot causes similar boll rots to Phytophthora, with mould often appearing as a covering of orange or pink spore masses.



**Fusarium affected boll (left) showing orange spore masses.** (Duy Le, NSW DPI)

### Diplodia boll rot

Diplodia boll rot starts as dark brown lesions that rapidly expand to cover the whole boll as the rot progresses. In the later stages of development, bolls become covered with a black smut-like fungal growth which can easily be rubbed off the boll surface.

Several other fungi can cause secondary boll rots in cotton, taking advantage of injury or wounds in the boll wall, often caused by insect pests.



**Bolls affected by Diplodia are covered in 'sooty' black spores.** (Linda Smith, Qld DAF)

### Anthracnose boll rot

Characterised by large spreading lesions on bolls, often with a pink spore mass in the centre. The pathogen is able to infect all parts of the cotton plant and at any stage of growth. Seedling stems may be girdled at or near the base of the stem. Anthracnose boll rot is uncommon in Australia, but is occasionally seen in Queensland cotton crops.

### Seed rots

Seed rot refers to boll rot that begins in the seed. Pathogens gain entry to the unopened boll when sucking insects (such as green vegetable bug, mirids or pale cotton stainers) feed on the developing seeds through the boll wall, as indicated by small black spots (1-2 mm diameter) on the surface of the boll. Seeds within the maturing green bolls are swollen and discoloured yellow or brown. When the affected bolls open, the locks with infected seed fail to fluff out and remain compact and discoloured. Seed rots do not necessarily affect the whole boll, rather they may be limited to one or two locks.

### Favoured by

- Wet and humid conditions, especially from a thick rank canopy and high moisture from rains and dews.
- Rainfall on exposed soil that splashes soil up onto the plant. Low mature bolls and lodged plants are at higher risk of infection.
- High numbers of sucking pests soon after flowering (seed rots).
- Boll rots and tight locks can also develop when bolls that are opening are exposed to wet weather.

### Host range

A broad range of fungal and bacterial species are associated with boll rots and their host ranges vary. For example, Phytophthora hosts include safflower, pineapple, tomato and citrus as well as a large number of ornamental plants derived from Australian native flora. Sclerotinia hosts include sunflower, safflower, soybeans and most pasture legumes.

### IDM tactics

- Do not allow water to back-up into the field and inundate low bolls on plants near the tail drain.
- Avoid very low plant populations leading to exposed soil that can be splashed up onto low bolls at the end of the season.
- Avoid rank growth and a dense crop canopy if possible, especially in wet seasons.
- Assess incidence prior to or after defoliation by counting all of the bolls on ten plants from each of ten randomly selected sites across the field. Note that plants near the tail drain may bias the result.
- Thoroughly incorporate crop residues as soon as possible.
- Practice good farm hygiene and Come Clean. Go Clean.

## Cotton bunchy top (CBT)

CBT is a viral disease spread by the cotton aphid, and can only survive in living plants.

### Symptoms

Leaves usually have pale green angular patterns around the margins and darker green centres, and can be leathery and brittle compared to the leaves on healthy plants.

After the plant is infected, subsequent growth is characterised by small leaves, short internodes and small bolls.

When plants are affected at a very early stage (e.g. as seedlings) the whole plant has a compact, severely stunted appearance. Roots appear hairy and dark brown compared to the light yellow-brown colour of healthy roots, and small knots form on the secondary root branches.

Symptoms are difficult to distinguish in perennial volunteer cotton and late crops (post cut-out) where there has been insufficient new growth. It usually takes 3-8 weeks from infection until symptoms become obvious.

Patches of infected plants may occur around ratoon plants that were affected by CBT and survived from the previous season. These infected ratoon plants often harbour aphids which can then move to adjacent plants, spreading the disease.



Pale green angular patterns around the margins and darker green centres is typical of cotton bunchy top. (Linda Smith, Qld DAF)



CBT infection leads to a distinctive leaf mottling. (Stephen Allen)

### Favoured by

- High aphid populations, in close proximity to ratoon cotton. Ratoons act as both a preferred host for the aphids and a reservoir for the disease, creating a source of infection in the new season.
- Climatic conditions suitable for aphid reproduction, feeding and spread.
- Wet or mild winters that enable more volunteer and ratoon cotton and aphids to survive between season and cropping cycles.

Cotton aphid has a broad host range, including many weeds. The presence of weed hosts allow cotton aphid populations to persist over winter, increasing the likelihood of aphids moving into cotton early in the season.

### Host range

The most critical alternative host plant is ratoon or volunteer cotton that survives between seasons, retaining leaves through winter and supporting infected aphid populations from one season to the next. The importance of other host plants is not well understood but in some situations marshmallow weed (*Malva parviflora*) may be an important overwintering host for virus and aphids.

Thirteen natural field hosts of CBT have been identified including: cotton, *Abutilon theophrasti* (velvetleaf), *Anoda cristata* (spurred anoda), *Chamaesyce hirta* (asthma plant), *Gossypium sturtianum* (Sturt's desert rose), *Hibiscus sabdariffa* (rosella), *Hibiscus trionum* (bladder ketmia), *Lamium amplexicaule* (deadnettle), *Malva parviflora* (marshmallow weed), *Malvastrum coromandelianum* (prickly malvastrum), *Medicago polymorpha* (burr medic), *Sida rhombifolia* (Paddy's lucerne), and *Trianthema portulacastrum* (black pigweed). *Gossypium australe* and *Cicer arietinum* (chickpea) were also found to be hosts in field or pot experiments.

These are currently the only known hosts of CBT. However the virus may have a wider host range than originally thought and include further non-Malvaceae species.

### IDM tactics

**Step 1. Avoid the problem** – elimination of hosts, particularly over winter, is the most effective means of minimising the risk of CBT. Break the green bridge and step 2 will not be required.

- CBT virus can only survive in living plants. A break in the presence of hosts between cotton seasons will reduce the risk of CBT surviving on-farm through winter. Cotton volunteers, regrowth and ratoons are important hosts of CBT. Good crop destruction and control of ratoons and volunteers is critical for CBT control. This also removes an important overwinter host for cotton aphid.



- Good on-farm management of broadleaf weeds is important as they can also host aphids and some may be hosts for CBT.
- Control volunteers/ratoons before cotton emerges to reduce the risk of winged aphids moving into nearby crops.

**Step 2. Manage the risk** – aphid control should not be the primary means of preventing infection.

- Don't over-react to aphids. Excessive use of aphicides will select for resistance and restrict control options.
- Sample young cotton regularly for aphids and assess aphid spread within the field.
- Maintain the beneficial complex (many beneficials present, high aphid mortality and little population spread) and keep monitoring. If aphids are healthy then consider selective control so that beneficials can provide ongoing mortality.
- If experiencing a high influx of aphids, consider a quick selective control to reduce the risk of CBT infection.

**CottonInfo videos:**

Identifying cotton bunchy top disease [youtu.be/xrp1420cilk](https://youtu.be/xrp1420cilk)

Cotton bunchy top management [youtu.be/3DQ561-wqqA](https://youtu.be/3DQ561-wqqA)

## Fusarium wilt

Pathogen: *Fusarium oxysporum* f. sp. *vasinfectum* (Fov)

### Symptoms

External symptoms include stunted growth and dull and wilted leaves followed by leaf yellowing or browning and eventual death from the top of the plant. Some affected plants may reshoot from the base of the stem. These symptoms most commonly become apparent in the seedling phase when plants are beginning to develop true leaves, or after flowering during boll fill. Symptoms can appear as individual plants or as a small patch, often near the tail drain or low-lying areas of the field.

Internal symptoms can be checked by cutting the stem. Infected plants will reveal continuous brown discoloration of the stem tissues from the main root up into the stem. The discoloration is similar to Verticillium wilt but usually appears as continuous browning rather than flecking.

### Favoured by

- Use of susceptible varieties.
- Stresses in the crop – e.g. waterlogging, root damage through cultivation, cool and wet growing conditions.
- Poor farm hygiene on and between farms and districts.



**Fusarium wilt: Wilting and dying plants are often observed at the tail drain.** (Linda Smith, Qld DAF)



**Dark brown discoloration observed in cross sections of cotton stems infected with Fov.** (Janelle Montgomery, CottonInfo)

## Host range

The Fov pathogen is specific to cotton but can also live in the residues of most non-host crops. Bladder ketmia, sesbania pea, dwarf amaranth, bellvine and wild melon are alternative weed hosts that show no external symptoms. These weeds may act as an on-farm reservoir for the disease and need to be constantly managed.

## IDM tactics

- If your farm is free from this disease, keep it that way! Ensure all staff and contractors practice good farm hygiene and Come Clean. Go Clean.
- Select varieties with a high F rank and use BION® Plant Activator.
- If possible, delay planting until soil temperatures are 16°C and rising.
- Manage the crop to avoid stresses such as waterlogging, over-fertilisation and root damage.
- Avoid mechanical inter-row cultivations if possible, as this can cause root damage that provides an entry point for the pathogen.
- Regularly inspect fields for early detection and containment of isolated outbreaks. Send any suspect samples to Dr Linda Smith (Qld DAF).
- Isolate affected areas from irrigation flows and traffic.
- Minimise tail water from affected fields.
- After harvest, root pull and retain crop residues on the surface for at least a month (60-120 days if possible) prior to incorporation. Raking and burning the whole field is likely to spread any pathogen present.
- Fusarium can survive on non-host crop residues. Avoid green manure crops as this returns organic matter to the field which Fusarium can survive on as a saprophyte.
- Rotate with non-hosts for up to 3 years. Hosts such as legumes can potentially increase disease. A summer sorghum/maize-fallow-cotton rotation can increase cotton plant survival, reduce disease incidence and increase yield in the third year compared to continuous cotton.

### CottonInfo videos:

Managing fusarium wilt [youtu.be/Cygy6XIRDew](https://youtu.be/Cygy6XIRDew)

Post-harvest management of Fusarium wilt [youtu.be/NGeQ4FTH7IM](https://youtu.be/NGeQ4FTH7IM)

## Reniform nematode

Pathogen: *Rotylenchulus reniformis*

A plant parasitic nematode that feeds on the plant root using retractable, hollow, spear-like mouthparts. Distributed worldwide within tropical and subtropical regions, reniform nematodes are one of the most damaging nematode pests. They are capable of attacking a wide range of crop plants, as well as many weed species.

## Symptoms

Feeding causes damage resulting in stunting and generally poor plant growth. Reniform nematodes typically reduce crop productivity rather than causing complete plant death. Populations can be quite uniform in their distribution across a field, making detection of early plant symptoms difficult.

## Favoured by

Warm climates (tropical and subtropical region), although it can be found in warm temperate regions as well.

Damage potential differs widely according to soil type. Sandy soils tend to promote the greatest level of damage, while nematode survival and reproductive success is favoured by soils with higher (20-40%) silt or clay.

## Host range

The very wide host range includes chickpeas, mungbeans, pigeon pea, sunflower and vetch. Certain crops are considered to be non-hosts,

including corn, canola, faba beans, safflower, sorghum, soybean, wheat, barley, triticale and oats.

## IDM tactics

- Come Clean. Go Clean. Good farm hygiene is the key to minimising the spread.
- Rotating with non-host crops such as wheat or sorghum to reduce base populations. Long fallows can also help to break the life cycle; however all weeds, cotton volunteers and ratoons must be controlled.
- Plant into good conditions, including optimum soil temperature, no water stress and well-formed beds.
- Monitor crops for patches of stunted plants and submit suspicious root samples for testing to Dr Linda Smith (Qld DAF).
- Manage cotton stubble. Cut cotton stalks and till soil through the stubble zone as soon as possible after harvest to destroy breeding sites. Ensure root cutting is successful and there is no re-growth.

## Assessment

Monitor for patches of unexplained unthrifty or stunted plants and send a sample of soil if concerned. Nematodes cannot be seen with the naked eye, but affected roots may have small nodules/knots.

- Mark patches with a GPS or on a map so you can re-monitor next season.
- Scrape off the dry top soil and sample 10-15 cm deep using a small trowel or soil corer.
- If there is more than one patch in a field, collect multiple samples from these areas in a bucket, and mix through.
- Place approximately 400 g of the soil in a clearly labelled plastic bag.
- The extraction process relies on live nematodes so please keep cool in an esky without an ice brick, DO NOT STORE SAMPLES IN THE FRIDGE.
- Include information about the sample sheet (see page 107 for a form and checklist on sending plant samples for diagnosis).

CottonInfo video on reniform nematodes [youtu.be/QgBn4vfkOzl](https://youtu.be/QgBn4vfkOzl)

## Reoccurring wilt

Pathogen: Yet to be formally confirmed

A new disease of cotton first observed in the 2017/18 season in Central Queensland has since been detected in the Namoi, Gwydir, St George and Border Rivers regions. It was initially thought to be an exotic disease based on symptoms, but all major biosecurity threats have been ruled out.

## Symptoms

The disease typically presents as wilting and sudden death of plants, either as single plants or in a patch similar to a lightning strike, but some



**Initial symptoms of wilting with leaves remaining attached.**  
(Linda Smith, Qld DAF)

plants within the patch may not be affected. Leaves turn bronze-brown, die and remain attached to the plant. A section of bark on affected plants will become blackened.

Internal symptoms can be observed by cutting the stem. A reddish-grey vascular discolouration can be seen and may not appear evenly through the cross-section.

### Favoured by

The etiology is still being established. At this stage it is unclear what factors are contributing to the disease.



**Single dead plant amongst healthy plants.** (Linda Smith, Qld DAF)



**In cross-section the infected tissue may be V-shaped with reddish-grey discolouration.** (Linda Smith, Qld DAF)



**Blackening of the stem.** (Linda Smith, Qld DAF)

### Host range

It is too early to make assumptions on the host range of the pathogen that causes reoccurring wilt.

### IDM tactics

There are no specific management tools yet determined for this disease. However, monitor fields for symptoms as described.

Contact your local REO or one of the pathologists if you suspect the disease on your farm.

Come Clean, Go Clean. Implement and maintain good farm biosecurity practices between fields and farms to help prevent the spread of all pathogens.

Please send any suspicious plants to one of the cotton pathologists:

- Linda Smith (Qld DAF) – 0457 547 617.
- Duy Le (NSW DPI) – 0439 941 542 (do not send suspected Fusarium samples to Duy at ACRI).
- Beth Shakeshaft (NSW DPI and CottonInfo) – 0409 477 303.

## Seedling diseases

Seedling diseases are caused by numerous pathogens either acting alone or in combination that commonly cause 'damping off' (death of seedlings) and reduced plant stands. The main pathogens infecting cotton seedlings are *Rhizoctonia solani*, *Pythium ultimum* and *Fusarium* spp. (not the *Fusarium* wilt pathogen).

### Symptoms

- Pre-emergent seed rots.
- Post-emergent damping off (wilting, collapse and death of seedlings).
- Slow early season growth, small cotyledons and reddened hypocotyls.
- Lesions on roots.

Affected plants may be scattered across the field or concentrated in poorly drained areas. In some situations seedling disease may be particularly evident in rows where other factors such as fertiliser placement, herbicide application, planting depth etc have had an effect.

### Favoured by

Anything that slows down germination and seedling growth, including cool and/or wet weather, poorly formed beds, compaction, waterlogging, incorrect planting depth, poor placement of fertiliser (under the plant line), excessive rates of herbicide at planting, movement of herbicide into the root zone (i.e. by rain) and infection by other pathogens.

### Host range

Seedling disease pathogens have a wide host range and can survive on the residues of many crops and weeds. International research has determined that winter legume crops do not appear to increase the risk of cotton seedling diseases sufficiently to deter their use in reducing soil erosion and providing nitrogen to the following crop.

The pathogens can survive on plant residues, so incorporation early enough to allow breakdown is important.

### IDM tactics

- Use a variety with good seedling vigour.
- Use effective seed treatment fungicides.
- Plant into well prepared, high, firm beds.
- Carefully position fertiliser in the bed (not under the plant line).
- Plant into moisture rather than planting dry and watering-up.
- Delay planting until temperature and moisture conditions are optimum.
- Be careful with the use of herbicides at planting.
- Incorporate rotation crop residues as soon as possible after harvest.


## Verticillium wilt

### *Verticillium dahliae*

Verticillium wilt is caused by the soil-borne fungal pathogen *Verticillium dahliae*. In Australian cotton there are currently three strains of *V. dahliae*, however virulence varies greatly within some strains. All incidences of Verticillium wilt should be taken seriously regardless of the strain as yield losses can still be severe without defoliation occurring.

### Symptoms

Symptoms of Verticillium wilt and Fusarium wilt are similar. Verticillium wilt has dark brown to black streaks through the centre of the stem when cut diagonally. When cut lengthways, stems show brown flecking of the inner tissues. As Verticillium wilt and Fusarium wilt can be difficult to tell apart, plant/s suspected of being infected with *F. oxysporum* f. sp. *vasinfectum* or *V. dahliae* need to be diagnosed by a pathologist. In some instances there



Typical leaf symptoms of Verticillium wilt. (Stephen Allen)



Dark grey peppery discolouration observed in cross sections of cotton stems infected with *V. dahliae*. (Duy Le, NSW DPI)



Characteristic symptoms of Verticillium wilt include wilted plants, leaf mottle and necrosis. As disease progresses, defoliation and the death of plants may occur. (Linda Smith, Qld DAF)

are fields with both Verticillium and Fusarium wilt present, so send multiple stems in the sample.

*V. dahliae* can also cause a characteristic yellow mottle between the veins and around the leaf margins. Lower leaves are usually affected first. Dead tissue develops at the leaf edges and may replace the mottled areas.

### Favoured by

- Cooler temperatures (cotton varieties that are resistant at 25-27°C are susceptible at 20-22°C).
- Extended wet weather.
- Waterlogging (particularly in late-maturing crops).
- Potassium deficiency.

Extended crop growth late in the season, including through the use of excessive nitrogen, increases the risk of this disease.

## Host range

*V. dahliae* has a large host range and causes vascular wilt on more than 250 plant species including volunteer cotton, ratoon cotton, soybeans, canola and safflower. Weeds that host the disease include Noogoora and Bathurst burr, saffron thistle, thornapple, caustic weed, bladder ketmia, burr medic, black bindweed, pigweed, devils claw, turnip weed, mintweed and blackberry nightshade. There is some host specificity between strains. International literature and pot trials with Australian strains also suggest that wheat, barley, oats, vetch, sunflower, maize, mungbean, chickpea and faba beans may be hosts.

## IDM tactics

- Select varieties with a high V rank.
- Manage for earliness, including optimising nutrition and water inputs.
- Avoid over-watering, waterlogging and late season irrigations that extend maturity.
- Minimise tailwater to reduce risk of spread.
- Avoid inter-row cultivation with knives if possible. This causes root damage and provides an entry point for the pathogen.
- Ensure that crop destruction occurs soon after picking to reduce the build-up of inoculum.
- Rotate with non-host crops (i.e. sorghum).
- Avoid frequent fallows to maintain microbial diversity and populations of beneficial bacteria and fungi.
- Avoid/control alternative hosts, including volunteer and ratoon cotton.
- Send any suspicious plant samples to your state pathologist for correct identification of pathogen.

- If your farm is free of *V. dahliae*, try to keep your farm that way. Ensure all farm staff and contractors practice good farm hygiene and Come Clean. Go Clean.

**CottonInfo video on Verticillium wilt** [youtu.be/hqiatA52nNw](http://youtu.be/hqiatA52nNw)

## Other diseases

Other diseases that can affect cotton include bacterial blight, bacterial stunt, post-harvest rot, sudden wilt, tobacco streak virus (TSV) and tropical rust. These diseases are much less common than the others mentioned in this chapter, largely thanks to cotton breeding programs and biosecurity measures.

Not all unusual plant symptoms are a sign of disease. Some herbicides, insects, soil constraints and environmental factors can also contribute to abnormal plant growth.

More detailed information on cotton diseases and other symptoms can be found at the CottonInfo website ([www.cottoninfo.com.au/disease-management](http://www.cottoninfo.com.au/disease-management)) with links to the Cotton Symptoms Guide and various disease factsheets, along with information on how to send a sample in for disease diagnostics.

**CottonInfo video on collecting stem and leaf samples for disease testing** [youtu.be/4mCk2883100](http://youtu.be/4mCk2883100)

### SENDING A SAMPLE FOR DIAGNOSIS BY A PATHOLOGIST – ATTACH A COMPLETED FORM TO EACH SAMPLE

|  |                                    |
|--|------------------------------------|
| Collected/Submitted by: (e.g. Cotton Regional Extension Officer) | Address/Email/Fax/Telephone:       |
| Property name and field number:                                  | Date collected:                    |
| Grower/Agronomist  | Grower's address or area/locality: |

Mark (X) as appropriate

| SYMPTOMS  | DISTRIBUTION  | INCIDENCE/SEVERITY                                   | CROP GROWTH STAGE                         |
|---|---|--|---|
| <input type="checkbox"/> Poor emergence or seedling depth       | <input type="checkbox"/> One field only             | <input type="checkbox"/> All plants                  | <input type="checkbox"/> Irrigated        |
| <input type="checkbox"/> Leaves: spots or dead areas            | <input type="checkbox"/> In several fields          | <input type="checkbox"/> Scattered single plants     | <input type="checkbox"/> Dryland          |
| <input type="checkbox"/> Leaves: discoloured                    | <input type="checkbox"/> In all fields              | <input type="checkbox"/> Scattered patches of plants | <input type="checkbox"/> Seedling stage   |
| <input type="checkbox"/> Leaves: mottled                        | <input type="checkbox"/> One variety only           | <input type="checkbox"/> In a large patch (>5 m)     | <input type="checkbox"/> Setting squares  |
| <input type="checkbox"/> Leaves or shoots: distorted or curled  | <input type="checkbox"/> Several varieties affected | <input type="checkbox"/> In a small patch (1-5 m)    | <input type="checkbox"/> Early flowering  |
| <input type="checkbox"/> Plants stunted                         | <input type="checkbox"/> Some rows more affected    | <input type="checkbox"/> In a small patch (<1 m)     | <input type="checkbox"/> Peak flowering   |
| <input type="checkbox"/> Plants wilting                         | <input type="checkbox"/> On lighter soil types      | <input type="checkbox"/> Plants dead                 | <input type="checkbox"/> First bolls open |
| <input type="checkbox"/> Premature plant death                  | <input type="checkbox"/> On heavier soil types      | <input type="checkbox"/> Plants defoliating          | <input type="checkbox"/> Defoliated       |
| <input type="checkbox"/> Bolls: spots or dead areas             | <input type="checkbox"/> In poorly drained area(s)  | <input type="checkbox"/> One to a few plants only    | <input type="checkbox"/> Ready to pick    |
| <input type="checkbox"/> Roots: discoloured, bent, pruned, etc. | <input type="checkbox"/> Other: (please specify)    |  |   |

**OTHER INFORMATION**

- Cultivar .....
- Paddock History .....
- Nearby crops .....
- Rainfall in last 10 days .....
- Average temperature range over the last 10 years .....
- Date of last irrigation .....
- Date of last cultivation .....

**Please contact your local CottonInfo REO to determine the appropriate pathologist and address for submitting sample**

**IF FUSARIUM WILT IS SUSPECTED, DO NOT SEND SAMPLES TO ACR!**

When sending samples:

- Send multiple samples (e.g. more than 1 leaf, stem or plant).
- If possible include a healthy plant as well as the diseased plant material.
- It is better to despatch samples early in the week rather than just before the weekend.
- Never wrap samples in plastic. Dry or slightly dampened newspaper is better.

- When collecting seedlings – dig them up rather than pull them out. Include some soil.
- Several sections of stem (10-15 cm long) are usually adequate for wilt diseases.
- Keep the sample cool and send as soon as possible.

**TABLE 28: Fungicides registered for use in cotton**

|                   | Active ingredient   | Trade name                             | MoA group    | Targets  | Comments   |
|-------------------|---|--|--------------|--|--|
| Seed treatments*  | Azoxystrobin 75 g/L + Metalaxyl-M 75 g/L + Sedaxane 35 g/L + Fludioxonil 12.5 g/L | Vibrance (V)<br>Vibrance Complete (V2) | 4, 7, 11, 12 | Damping-off caused by <i>Pythium</i> spp. and <i>Rhizoctonia solani</i>                              | Seed treatment available through CSD when ordering seed.   |
|                   | Acibenzolar-S-methyl 500 g/L  | Bion                                   | No group     | Fusarium wilt, black root rot  | Seed treatment available through CSD when ordering seed. Suppression only. Activates the plant's natural resistance mechanisms.  |
|                   | Metalaxyl-M 350 g/L   | Various                                | 4            | Damping-off caused by <i>Pythium</i> and <i>Phytophthora</i> .<br>Fusarium wilt disinfection on seed | Seed dressing. QLD & NSW only. Does not protect against damping off by <i>Rhizoctonia</i> , <i>Fusarium</i> or other fungi.<br>97% R-enantiomer. Much more effective in controlling diseases. Lower rates can be used.   |
| At planting       | Quintozene 750 g/kg   | Terraclor                              | 14           | Damping-off caused by <i>Rhizoctonia</i>   | QLD, NSW & WA only. Soil fungicide. Spray over seed and surrounding soil at planting.  |
|                   | Tolclofos-methyl 500 g/L  | Tolex<br>Rhizolex In Furrow            | 14           | <i>Rhizoctonia solani</i> , <i>Rhizoctonia</i> spp.  | QLD & NSW only. Mix with seed immediately before planting.<br>Applied as an in-furrow spray or water injection at planting.  |
| Foliar fungicides | <i>Clitoria ternatea</i> extract 400 g/L  | Sero-X                                 | No group     | Verticillium wilt microsclerotia   | Will reduce the amount of microsclerotia (inoculum) returning to the soil. QLD, NSW & WA only. Three applications are required:<br>1. When majority of plants are between first square and first flower.<br>2. When majority of plants are between mid to peak flowering.<br>3. With the first defoliation.<br>Foliar application only. Will not reduce the severity of the disease in the season it is applied. |
|                   | Prothioconazole 150 g/L + Bixafen 75 g/L  | Aviator Xpro                           | 3, 7         | Sclerotinia stem rot   | PERMIT 87169. Expires 30 September 2022. NSW, QLD & WA only. WHP: 5 weeks. Boomspray application only. Maximum 2 applications per season.  |
|                   | Tebuconazole 430 g/kg   | Various                                | 3            | Alternaria leaf spot   | PERMIT 82660. Expires 31 March 2022. Southern cotton growing valleys: Bourke, Gwydir, Lachlan, Macquarie, Murrumbidgee, Murray and Namoi Valleys only. WHP: 6 weeks. Maximum of 2 applications per season, 14 days apart.  |
|                   | Mancozeb 750 g/kg   | Various                                | M3           | Alternaria leaf spot ( <b>pima cotton varieties only</b> )   | Foliar spray. Do not apply more than 4 sprays per season. <b>Please note PER87168 for use on <i>Gossypium hirsutum</i> varieties expired on 31 March 2021 and is currently under revision for renewal.</b>   |
| Seed export       | Carboxin 200 g/L + Thiram 200 g/L   | Vitavax 200FF                          | 7, M3        | Seedborne disease control  | Cotton seed destined for export to countries that specify such treatment only – not registered for seed used in Australia.   |

\*Penflufen 154 g/L + trifloxystrobin 154 g/L (EverGol Xtend), also registered as a seed treatment, has been superseded by Vibrance.

**Industry pathology team leaders:**

Qld DAF pathologist, Linda Smith 0457 547 617

NSW DPI pathologists, Duy Le 0439 941 542, Karen Kirkby 0428 944 500

NSW DPI pathologist and CottonInfo Tech Lead for Diseases, Beth Shakeshaft 0409 477 303

[www.cottoninfo.com.au/publications/cotton-symptoms-guide](http://www.cottoninfo.com.au/publications/cotton-symptoms-guide)

# Regional disease update 2020-21

**C**ommercial cotton crops across New South Wales and Queensland were inspected early and late in the 2020-21 season. This represents the 19th and 38th consecutive seasons of cotton disease surveys in Queensland and New South Wales and respectively.

The incidence and severity of diseases present were assessed and field history, ground preparation, cotton variety, planting date and seed rate recorded for each field. The endemic diseases monitored include seedling diseases (caused by *Rhizoctonia solani*, *Pythium* spp. and *Fusarium* spp.), black root rot (BRR), Fusarium wilt, Verticillium wilt, Alternaria leaf spot, tobacco streak virus (TSV), cotton bunchy top (CBT) and boll rots, plus anything unusual.

## Cotton industry biosecurity plan – Crop surveillance for priority pests

Particular attention was given to surveying fields for the presence/absence of exotic diseases including cotton leaf curl disease, blue disease, Phymatotrichopsis (Texas) root rot, the hypervirulent strains of the bacterial blight pathogen and exotic strains of the Verticillium wilt and Fusarium wilt pathogens. None of these diseases and/or pathogens were observed.

**Acknowledgement:** The regional surveys were undertaken by Linda Smith, Linda Scheikowski & Dinesh Kafle (Qld DAF), Duy Le, Aphrika Gregson & Beth Shakeshaft (NSW DPI), and CottonInfo Regional Extension Officers. Thank you also to regional consultants who assisted with collection of disease incidence data. This work is being undertaken as part of the CRDC funded project DAQ2002 Characteristics of disease suppressive cotton farming systems and soils understood.

## Queensland

A total of 61 fields across 46 farms were surveyed early season from October to December in 2020. Rain and timing impacted the ability to complete surveys of all fields late season (where possible, alternative fields were surveyed). A total of 45 fields across 35 farms were inspected from February to May. The key disease issues in the 2020/21 season for Queensland cotton varied for each region:

- **Central Queensland:** Poor root development and high reniform nematode pressure impacted seedling development. Reoccurring wilt incidence is increasing and of concern.
- **Darling Downs:** Fusarium wilt and the increase in the incidence of Verticillium wilt are key issues. Severe leaf spot associated with the presence of pathogens other than Alternaria is of concern and requires monitoring to determine impact.
- **St George:** The incidence of common diseases was low. The detection of reoccurring wilt and leaf spot associated with fungi other than Alternaria requires monitoring.
- **Border Rivers:** Fusarium and Verticillium wilts are of major concern, with high incidence in several fields surveyed. Reoccurring wilt is also increasing.

The amount of cotton trash present at planting was estimated using a standardised photograph guide to provide information on the potential carryover of pathogen inoculum. Pathogens that cause Verticillium wilt, Fusarium wilt, black root rot, boll rots, seedling disease and Alternaria leaf spot can all survive in association with undecomposed trash from previous cotton crops. It is important that cotton residues are incorporated, in time

to allow for microbial breakdown, however this season was challenging as another dry winter meant that cotton trash was evident in some fields.

## Central Queensland

With insufficient rain and the Fairbairn Dam hitting a historic low of 7.83% capacity, plantings in the Emerald region in 2020 were limited. Therefore, disease surveys concentrated efforts in the Theodore region, surveying a total of 14 fields across 7 farms. Fusarium wilt and Pythium were each detected in one field. Black root rot incidence ranged up to 100% for infected seedlings (Figure 1). Alternaria leaf spot was detected in 57% of fields surveyed. Leaf spots were observed on the cotyledons only (Figure 2).




**Figure 1: Severe BRR in tap root of seedlings.**



**Figure 2: Planting into infected cotton residues increases risk of Alternaria leaf spot in emerging seedlings.**

Reniform nematode was observed in roots of seedlings in 12/14 fields surveyed (86%) (Figure 3). Roots of mungbean volunteers were also infected with reniform, confirming this crop as a host (Figure 3). 'Witches broom' roots were observed in 6/14 (43%) of fields surveyed (Figure 4). The level of reniform infection would not only have impacted plant development, but the root damage could also have provided entry to seedling disease pathogens.




**Figure 3: Reniform nematode infected cotton (left) and mungbean (right).**



**Figure 4: 'Witches broom' roots.**

Verticillium wilt was not detected in late season surveys, and the field in which Fusarium wilt was detected early season was not surveyed late season. Boll rot, seed rot and tight lock were observed in every field surveyed. The average percentage of leaf surface with Alternaria leaf spot was 0.4%. Reoccurring wilt was detected in four of the eight fields surveyed (Figure 5). In three fields the incidence was low, however the fourth field had a significant patch of disease that was 10 x 20 m in size. Mealybugs were observed on cotton in 34% of fields surveyed.



**Figure 5: Reoccurring wilt. Symptoms include (A, B) sudden wilting and death of plant with leaves remaining attached, and (C) discoloration of the vascular tissue – reddish/grey.**

## Darling Downs


A total of 20 fields across 16 farms were surveyed. Fusarium wilt was detected in eight fields. Pythium was detected in three fields at a low incidence. Alternaria leaf spot was detected on the cotyledons of seedlings in one field at a low infection rate. There were moisture issues resulting in seedling death following germination. In some fields where a lack of moisture was an issue, soil crusting at the hypocotyl resulted in stunted seedlings with a pinched hypocotyl (Figure 6).



**Figure 6: Pinched hypocotyl in soils with surface crusting.**

Both Fusarium and Verticillium wilts were detected in fields surveyed late season. Fusarium wilt is a key issue for this region. To better understand the impact of the pathogen on the cotton plant, the severity of vascular discoloration visible in a cross section of the main plant stem was rated using a vascular discoloration index. Before harvest, 28.7% of infected plants had greater than 40% vascular discoloration of the stem cross section and 11.7% were dead (Figure 7). Hence Fusarium wilt is still having a significant impact. Although Verticillium wilt is not typically prevalent in this region, there are some fields for which Verticillium is the

most significant disease issue. One field known to have Fusarium wilt, had Verticillium wilt detected for the first time. Boll rot, seed rot and tight lock were observed in most of fields surveyed, at a low level. Reoccurring wilt was not detected.



**Figure 7: Percentage of Fusarium wilt infected plants rated 1 to 5 for vascular discoloration during late season surveys on the Darling Downs.**

This rating system is based on the severity of vascular discoloration visible in a cross section of the main plant stem cut as close as practicable to ground level where:

- 1 = discoloration restricted to an area less than 5% of the stem cross section;
- 2 = discoloration of between 5% and 20% of the stem cross section;
- 3 = discoloration of between 20% and 40% of the stem cross section;
- 4 = greater than 40% vascular discoloration of the stem cross section; and,
- 5 = dead plant.

Several farms on the Darling Downs this season suffered significant leaf spot (Figure 8). One field had leaves with 20% of the leaf area infected and a second field had 7.5%.



**Figure 8: A high incidence of leaf spot on the Darling Downs (4 Feb 2021). Disease was widespread and yield impact was of concern.**

## St George/Dirranbandi

A total of 14 fields across 10 farms were surveyed. Fusarium wilt was detected in three fields. BRR was not detected at the time of the surveys due to plant age. Tipping out (caused by insects) was observed across all farms surveyed.

In late season surveys, Fusarium wilt was detected in four fields. No Verticillium wilt was detected. The average incidence of Alternaria leaf spot was 5%, with most leaf spots observed in the top half of the canopy. The incidence of boll rot, seed rot and tight lock was low this season. The highest incidence of tight lock (12.7%) was detected in a late planted crop. Reoccurring wilt was detected in one field, and on a second farm in a field not included in the surveys. The incidence was low, with the disease observed in a small number of single plants amongst healthy plants.

## Border Rivers

A total of 13 fields across 8 farms were surveyed. Fusarium wilt was detected in three fields, at a low incidence. Rhizoctonia incidence was detected in 10 fields. BRR was not detected at the time of the surveys due to plant age. A trace of Alternaria leaf spot was detected on the cotyledons of seedlings in one field.

Twelve fields across the 8 farms were surveyed late season. Fusarium wilt was detected in five fields, and Verticillium wilt in three (two fields



**TABLE 29a: Overall average disease incidence (% of plants infected) detected during early season surveys in Queensland, with incidence range for fields where disease was present. Carry over of cotton trash is also provided.**

|                               | Seedlings/m        | Fusarium          | Pythium    | Rhizoctonia        | Black root rot*                  | Alternaria**                 | Cotton trash (g/m <sup>2</sup> ) |
|-------------------------------|--------------------|-------------------|------------|--------------------|----------------------------------|------------------------------|----------------------------------|
| Central Queensland (Theodore) | 9.9<br>(6.1-13.4)  | 0.3<br>(4.7)      | 0.1<br>(2) | 1.7<br>(1.3-5)     | 15<br>(0.7-100)<br>Severity = 8% | 0.4 cotyledons               | 65.8<br>(8-140)                  |
| Darling Downs                 | 10.7<br>(4.7-12.9) | 1.6<br>(0.7-12.3) | 0.1        | 3.4<br>(0.33-14.3) | 24<br>(2-67)<br>Severity = 16%   | 0.01<br>(0.22)<br>cotyledons | 39.7<br>(0.8-128)                |
| St George/<br>Dirranbandi     | 11.7<br>(9-14.9)   | 0.2<br>(0.5-2.3)  | 0          | 0.5<br>(0.33-2.5)  | Growing out of<br>disease        | 0                            | 8<br>(0.25-27.6)                 |
| Border Rivers                 | 9.8<br>(7.2-14.8)  | 0.3<br>(0.33-3)   | 0          | 0.5<br>(0.33-1.3)  | Growing out of<br>disease        | trace<br>(1 field)           | 38.3<br>(1.3-189.1)              |

\*BRR also includes a severity ranking (from no blackening of the tap root to 100% of tap root blackened).  
\*\*Alternaria incidence based on % of leaf surface with leaf spotting.

**TABLE 29b: Average disease incidence detected during late season disease surveys in Queensland, with incidence range for fields where disease was present.**

|                               | Fusarium         | Verticillium    | Boll rot          | Seed rot         | Tight lock         | Alternaria and other leaf spots | Reoccurring wilt                         |
|-------------------------------|------------------|-----------------|-------------------|------------------|--------------------|---------------------------------|--|
| Central Queensland (Theodore) | 0                | 0               | 1.7<br>(0.35-6.7) | 2.3<br>(0.8-7.6) | 1.4<br>(0.5-2.5)   | 0.4                             | 0.4<br>(0.33-2.3 + high incidence patch) |
| Darling Downs                 | 3.4<br>(0.5-14)  | 1.4<br>(0.5-19) | 2.8<br>(0.4-7.5)  | 1.4<br>(0.4-3.4) | 3.9<br>(0.35-14.4) | 2.4                             | 0  |
| St George/<br>Dirranbandi     | 0.8<br>(0.5-3.5) | 0               | 2.4<br>(0.4-3.3)  | 0.5<br>(1-1.45)  | 3.6<br>(1.2-12.7)  | 5                               | 0.05<br>(0.5)                            |
| Border Rivers                 | 3.5<br>(0.5-20)  | 5<br>(3-48)     | 2.8<br>(0.6-6.6)  | 1.7<br>(0.4-7.9) | 5.7<br>(0.4-13.5)  | 0.4                             | 0.5<br>(0.5-4)                           |

contained both). Both Fusarium and Verticillium wilts are key diseases of concern for this region. Boll rot, seed rot and tight lock were observed at a low incidence in most of fields surveyed. Reoccurring wilt was detected in three fields.

## New South Wales

A total of 56 fields from 39 NSW farms were surveyed during the months of November and December 2020. After many years without cotton, growers in Bourke were able to plant around 8,000 ha of cotton this 2020/21 season. Common seedling diseases such as black root rot, Rhizoctonia rot and Alternaria leaf spot were detected across all cotton growing regions in NSW. Black root rot and Rhizoctonia rot were still major seedling diseases across all regions. Though detected with relatively high incidence, most of the Rhizoctonia infection was superficial and the cotton seedlings grew out of it. Alternaria leaf spot remained prevalent and the disease damage caused minor concerns in most of the surveyed fields. Additionally, unforeseeable weather events such as hailstorms and heavy rainfall caused some significant seedling damage and several fields in the Lachlan and Namoi were replanted.

The late season disease survey in NSW was conducted during mid-March to mid-April 2021. A total of 46 fields across 34 farms were revisited. A wet weather event and subsequent localised flooding early in March 2021 was a challenge for our disease surveys in Gwydir, Namoi and Macquarie valleys, and created conducive conditions for late season diseases such as Fusarium and Verticillium wilts, and boll rot development. Alternaria leaf spot was prevalent across NSW but the severity was relatively mild. Reoccurring wilt was detected again in three different properties in the Namoi and Gwydir valleys.

## Gwydir

A total of 13 fields were surveyed early season. Black root rot and Rhizoctonia rot incidence varied widely; of the infected fields, black root rot severity was from 2 to 35% and Rhizoctonia from 2 to 27.5%. Alternaria leaf spot was detected in most of the surveyed fields, usually as only minor pin head lesions, and the severity was well below 1%. Fusarium wilt on seedlings was also detected outside of surveyed transects in several fields in the Gwydir this season (Figure 9).



**Figure 9: Cotton seedling exhibited necrosis on cotyledons and profound vascular discolouration.**

Again this season, Gwydir was a hotspot for Fusarium wilt. The disease was detected late season in nine out of 11 surveyed fields, with incidence up to 96% in one field, significantly reducing yield. The mean incidence across the valley was at 20.83%. Verticillium was detected in seven fields with the incidence below 3% (mean 0.46%). Six fields contained both wilt diseases. Reoccurring wilt disease was detected in three fields from one new property with a disease incidence of less than 2%. Of the affected

fields, dead plants occurred in single or small batches that resembled lightning strike damage at first (Figure 10). This typical symptom could assist in field inspections for reoccurring wilt occurrence. Alternaria leaf spot was minor, the mean incidence across the valley was 0.81%. By late March 2021, parts of the valley were flooded, and several fields included in our survey were also affected (Figure 11), coinciding with a particularly high incidence of boll rot and tight lock in these fields. In a couple of fields, tight lock incidence up to 23% was recorded, but as a whole the incidence was sitting below 10%. Seed rot was detected in several fields with a mean incidence of 0.68%.



**Figure 10: Reoccurring wilt observed in multiple dead plants resembling lightning strike damage was spotted in a Gwydir field.**



**Figure 11: Flood affected bolls in Gwydir fields.**

### Bourke

Four selected fields from the two biggest growers in Bourke were included in the disease survey this season. Only minor pin head Alternaria lesions were observed in infested fields and the severity was well below 1%.

Unfortunately, due to unexpected weather conditions and a shortage of human resources the late season survey for Bourke could not be undertaken.

### Namoi

A total of 11 fields were surveyed early season. Alternaria leaf spot was detected in most of the surveyed fields, with the severity mostly less than 3%.

Verticillium wilt remained a major disease in the valley, detected in eight of the 10 fields surveyed late season. While the disease incidence was especially high (above 50%) in a couple of fields which were likely to experience significant yield reduction, overall the mean incidence across the valley remained below 20%. No Fusarium wilt was detected during the late season disease survey. Reoccurring wilt often occurred in small batches (Figure 12) and was detected in three fields from two properties with an incidence in the affected fields of less than 3% (although up to six

dead plants in a row were recorded). Of the two properties, one was newly detected and the other a targeted surveillance field due to its previously known history of reoccurring wilt. Boll rot incidence remained below 10%. Tight lock incidence was low and seed rot was only detected in one field.



**Figure 12: Reoccurring wilt found in a small hotspot in a Namoi field.**

### Macquarie

A total of 11 fields were surveyed early season. Alternaria leaf spot was detected in all surveyed fields, but only minor pin head lesions were observed in most fields and the severity was well below 1%. Seedling death was also detected in some fields, but most cases were associated with insect feeding damage.

No Fusarium wilt was detected in the valley during the late season disease survey. Verticillium was detected in three fields across three properties with an incidence lower than 10%, and Alternaria leaf spot was prevalent. Boll rot incidence was up to 15% in one field, but the mean incidence across the valley remained below 5%. Tight lock was also common. Seed rot was only detected in one field.

### Murrumbidgee

A total of 10 fields were surveyed early season. Alternaria leaf spot was detected in most of the surveyed fields; leaf spot incidence ranged from less than 1% to 43% and the severity was mostly less than 1%.

Late season disease pressure in Murrumbidgee was mild. Boll rot was detected in seven out of 11 surveyed fields. Tight lock was less common and seed rot only detected once. Both wilt diseases were recorded during the survey: Fusarium wilt was detected in two fields located on two separate properties (where the disease occurred previously), while Verticillium wilt was detected in a new block. Additionally, a number of send-in samples were also positive to both wilt pathogens, indicating a widespread presence of the two wilt diseases in the valley.

### Lachlan

A total of seven fields were surveyed early season. Alternaria leaf spot was detected in all surveyed fields, but only minor pin head lesions were observed in most fields and the severity was less than 1%.

Disease pressure was relatively mild during the late season survey. Alternaria leaf spot was evident in the top half of the canopy in all seven surveyed fields. Boll rot and tight lock were detected in the lower canopy in many surveyed fields but the incidence remained low at around 1%. Seed rot was detected in a couple of fields at low incidence. No Verticillium was detected during the survey. Fusarium wilt was detected in one field with an incidence of 1%.



**TABLE 30a: A summary of seedling diseases recorded during the early season survey in NSW.**

| Season 20/21 | Plant stands/m      | Black root rot     |                  | Rhizoctonia rot  |                  | Alternaria       |                   |
|--------------|---------------------|--------------------|------------------|------------------|------------------|------------------|-------------------|
|              |                     | Incidence (%)      | Severity (%)     | Incidence (%)    | Severity (%)     | Incidence (%)    | Severity (%)      |
| Gwydir       | 9.4<br>(7.9-12.1)   | 44.4<br>(9-80)     | 14.8<br>(2-35)   | 61.1<br>(5-87)   | 16.4<br>(2-27.5) | 50.1<br>(15-98)  | 0.7<br>(0.1-3.2)  |
| Bourke       | 7.2<br>(6-8.4)      | 36.8<br>(0-60)     | 4.6<br>(2-7)     | 22.4<br>(0-55)   | 5.7<br>(3-10)    | 28.2<br>(0-34)   | 0.8<br>(0-0.8)    |
| Namoi        | 9.3<br>(6.2-10.9)   | 33.5<br>(0-96)     | 12.1<br>(0.5-32) | 31.7<br>(0-71)   | 8.0<br>(2-17.5)  | 35.0<br>(0.92)   | 2.1<br>(0-8.4)    |
| Macquarie    | 11.9<br>(8.5-14.5)  | 33.5<br>(8-71)     | 9.5<br>(2-30)    | 61.3<br>(25-90)  | 14.3<br>(1-5)    | 10.2<br>(1-16)   | 0.2<br>(0.01-0.5) |
| Murrumbidgee | 12.4<br>(10.5-14.2) | 31.6<br>(0-98)     | 14.6<br>(0.5-79) | 35.5<br>(2.5-78) | 2.2<br>(1-5)     | 13.7<br>(0.3-43) | 0.6<br>(0.03-2.1) |
| Lachlan      | 12.5<br>(8.3-14.5)  | 46.8<br>(0.6-96.5) | 18.4<br>(1-40)   | 43.9<br>(10-62)  | 5.4<br>(2.4-12)  | 8.1<br>(0.8-26)  | 0.3<br>(0.01-0.8) |

**TABLE 30b: A summary of disease incidence recorded during the late season disease survey in NSW.**

| Valleys      | Verticillium wilt incidence (%) <sup>1</sup> | Fusarium wilt incidence (%) <sup>1</sup> | Reoccurring wilt <sup>2</sup> | Alternaria left spot severity (%) | Boll rot incidence (%) | Tight lock incidence (%) | Seed rot incidence (%) |
|--------------|--|--|-------------------------------|-----------------------------------|------------------------|--------------------------|------------------------|
| Gwydir       | 0.46<br>(0-3)                                | 20.83<br>(0-96)                          | Detected                      | 0.81<br>(0-3.6)                   | 6.17<br>(0-12.8)       | 7.04<br>(0.4-23)         | 0.68<br>(0-2.8)        |
| Namoi        | 19.55<br>(0-67)                              | 0.00                                     | Detected                      | 3.60<br>(1-7)                     | 2.31<br>(0-7.4)        | 0.56<br>(0-2.5)          | 0.03<br>(0-0.6)        |
| Macquarie    | 2.00<br>(0-12)                               | 0.00                                     | Not detected                  | 3.64<br>(3-7)                     | 4.80<br>(0-15)         | 2.64<br>(0-10)           | 0.04<br>(0-0.5)        |
| Lachlan      | 0.00   | 0.13<br>(0-2)                            | Not detected                  | 1.80<br>(0.1-5.9)                 | 0.93<br>(0-3.5)        | 0.87<br>(0-6.1)          | 0.13<br>(0-1.4)        |
| Murrumbidgee | 0.42<br>(0-7)                                | 0.42<br>(0-6)                            | Not detected                  | 2.44<br>(1-10)                    | 0.47<br>(0-2.9)        | 0.15<br>(0-2.2)          | 0.03<br>(0-0.8)        |

1. Both Verticillium and Fusarium wilt incidences presented here were collated directly from the field survey data, which might be subjected to adjustment upon the confirmation of laboratory diagnoses.

2. Reoccurring wilt was only indicated as 'Detected' and 'Not detected' in each valley. Where detected the disease occurred in small hotspots; the disease incidence recorded in formal survey transects was less than 3%.

# Biosecurity – we all have a responsibility

**B**iosecurity is the protection of your property and the entire industry from the entry, spread and establishment of exotic and endemic pests (including insects, weeds and diseases). Exotic pests can affect everyone – farmers, agronomists and the community, so it is important that everyone plays a part in preparing for and minimising their biosecurity risks. While Australia's national quarantine system helps to prevent the introduction of harmful exotic pests, the threat they pose is still very real. Our ongoing reliance on imported goods, as well as existing natural routes (e.g. winds), makes future incursions of exotic pests and diseases almost inevitable.

## Biosecurity – a legal responsibility

All Australian states and territories have their own respective biosecurity legislation to minimise the risk of the entry and spread of pests, pathogens and weed seeds. Legislation in Queensland and NSW is based on the principle of a 'shared responsibility' where everyone has a responsibility to contribute to biosecurity. These regulations are designed to provide more capability, flexibility and innovation in the management of biosecurity risks. Everyone in the community has a responsibility to ensure they minimise the risk of spreading pathogens, pests and weeds ('biosecurity matter') which could impact the environment, the economy and the broader community.

It is your responsibility to ensure you meet the requirements of your state or territory. If you have any questions or concerns, contact your State or Territory Agriculture Department.

**New South Wales Department of Primary Industries: 1800 680 244**

**Queensland Department of Agriculture and Fisheries: 13 25 23**

**Northern Territory Department of Primary Industries and Resources: (08) 8999 2118**

**Western Australia Department of Primary Industries and Regional Development: (08) 9334 1800**

## Anyone working on or visiting farms has a biosecurity responsibility

- **Come Clean. Go Clean.** Vehicles, farm equipment and people can carry weed seeds and diseases. Clean down between farms, including vehicles and footwear. Suggest using an on-farm vehicle where possible.
- **Spotted anything unusual?** Ensure any unusual plant symptoms or suspected exotic pests are reported to the Exotic Plant Pest Hotline (1800 084 881). Vigilance is vital for an early detection of an exotic plant pest threat.

## Growers have a biosecurity responsibility

- **Check your crop frequently** looking out for unusual crop symptoms and if you find anything suspicious, report it immediately. Make sure that you and your employees are familiar with the most important cotton pests. Don't move the infected material.
- **Call the Exotic Plant Pest Hotline 1800 084 881**, a dedicated reporting line that will be answered by an officer from your state or territory's agriculture department. Early reporting improves the chance of effective control and eradication.
- **Communicate your biosecurity requirements using clear signage** to ensure only essential vehicles and equipment gain access to production areas.

- **Provide wash down facilities** for visitors to use with appropriate agricultural detergents and/or disinfectants to clean equipment and tools prior to entry and exit.
- **Develop a farm biosecurity management plan** to assess how pests, weeds and diseases could enter the farm and what practices are in place to manage or reduce these risks.
- **Declare visits to farms overseas on re-entry to Australia.** All clothes and footwear worn on-farm should be thoroughly washed before returning, or left behind.
- **Ensure all seed is pest free.** This includes cotton and other refuge and commercial crops. Keep records of all farm inputs just in case.
- **Maintain zero tolerance of cotton volunteer plants and other weeds** at all times throughout the year to prevent pests harbouring there.

## Come Clean. Go Clean.

There are two different types of products used in wash down procedures: agricultural detergents and agricultural disinfectants.

**Agricultural detergents** (e.g. Bio-Cleanse™ or most industrial and domestic brands of soaps and detergents) provide optimal soil removal.

**Agricultural disinfectants** (e.g. Path-X™, Sporekill®, Steri-maX®, or products containing quaternary ammonium compounds) are effective against a range of pathogens found on agricultural farms.

To work properly, disinfectant products need to be applied to a surface that has all soil and trash removed. Effective decontamination (wash down) of items involves two steps:

- (1) Cleaning with detergent, followed by
- (2) Disinfection.

Product users must also satisfy themselves that the product they choose is appropriate for their situation, and ensure that all label requirements are read and followed.

## Biosecurity Plan for the Cotton Industry

Plant Health Australia (PHA) is the national coordinator of the government-industry partnership for plant biosecurity in Australia. Cotton Australia is a member and CRDC is an associate member of PHA. The PHA-facilitated Biosecurity Plan for the Cotton Industry is a framework to coordinate biosecurity activities and investment for Australia's cotton industry. It provides a mechanism for industry, governments and stakeholders to better prepare for and respond to incursions of pests that could have significant impacts on the cotton industry.

**For further information on cotton biosecurity stewardship contact Cotton Australia on 02 9669 5222 or go to [cottonaustralia.com.au/biosecurity](http://cottonaustralia.com.au/biosecurity)**

**For more information on the NSW's General Biosecurity Duty: [www.dpi.nsw.gov.au/biosecurity/your-role-in-biosecurity](http://www.dpi.nsw.gov.au/biosecurity/your-role-in-biosecurity)**

**For more information on Qld's General Biosecurity Obligation: [www.daf.qld.gov.au/business-priorities/biosecurity/policy-legislation-regulation/biosecurity-act-2014/general-biosecurity-obligation](http://www.daf.qld.gov.au/business-priorities/biosecurity/policy-legislation-regulation/biosecurity-act-2014/general-biosecurity-obligation)**

**To find out more about on-farm biosecurity measures to help prevent endemic and exotic pests entering and establishing on farm, go to Farm Biosecurity Program: [www.farmbiosecurity.com.au](http://www.farmbiosecurity.com.au)**

# Come Clean. Go Clean.

Dirty vehicles, machinery and equipment carry pests, weeds and diseases

## A GUIDE TO EFFECTIVE WASH DOWN OF VEHICLES AND MACHINERY

### 1 WASH DOWN

- Use compressed air or high pressure water to remove caked on trash and mud
- Get into crevices where mud or trash might be trapped
- Clean out the inside of the car, particularly foot pedals and mats regularly in contact with dirty footwear

#### WHERE

- ✓ On a clean wash down pad with a hard surface
- ✓ Located away from production areas
- ✓ Where wash off contaminants can be trapped



### 2 CLEAN

- Use a sponge or spray to cover all surfaces with an agricultural detergent
- Leave the detergent to work for 10 minutes\* before rinsing, making sure to remove any remaining soil or plant material

\*unless otherwise directed by product label



#### REMEMBER

To wash all equipment, floor mats, tools and footwear kept in the vehicle as well

### 3 DECON

- After removing physical dirt, consider using an agricultural decontaminant to kill any remaining pests or pathogens
- Refer to the APVMA for registered decontaminants and follow label instructions
- An additional rinse step may be necessary following disinfection

#### NOTE

Make sure vehicles and equipment are clean and free of mud and trash before applying a decontaminant



### 4 RINSE

- Rinse off vehicle, machine and/or other washed equipment
- Use high pressure water to remove mud and debris from the wash down area so it is clean for the next person



#### CHECK

Equipment that has not been cleaned on farm should be thoroughly inspected to ensure cleanliness

Images courtesy of Sharna Holman, QDAF, unless otherwise stated

Together we can stop the spread of pests, weeds and diseases.

BE A GOOD MATE STOP IT AT THE GATE

## Farm biosecurity management plans

A farm biosecurity management plan is a practical way of showing how you are preventing pests, weeds and diseases from:

- Being introduced to your property;
- Spreading around your property; or,
- Spreading from your property to others.

Developing a farm biosecurity management plan can help growers or farm managers to:

- Identify and assess the biosecurity risks to their farm and business;
- Outline biosecurity practices currently implemented; and,
- Identify biosecurity practices that can be implemented in the short and long-term to reduce biosecurity risks.

The suggested practices in a farm biosecurity management plan cover a range of farming scenarios and can be adapted to suit your particular circumstances. It is unlikely that all biosecurity risks or practices will apply to your farm, and in some cases alternative practices may be better suited to your farm and business. The important thing is that you are aware of your biosecurity risks and are taking reasonable steps to minimise those risks. If you build your plan around daily, monthly and yearly farming routines, biosecurity practices will become a habit rather than an additional task.


Completing a farm biosecurity management plan provides a point of discussion and clear instructions of your biosecurity expectations for farm visitors such as agronomists, researchers, utility workers and hunters. Farm biosecurity management plans are available on the CottonInfo website ([www.cottoninfo.com.au](http://www.cottoninfo.com.au)). If you require assistance with developing a farm biosecurity plan, please contact the CottonInfo Technical Lead for Biosecurity.

III

## Biosecurity Top Tips:

There can be several ways to mitigate a biosecurity risk, with the practices you implement reflecting what suits you and your farming business. Check out the Biosecurity Top Tip videos on the CottonInfo YouTube channel to see short videos of growers, consultants and researchers sharing how they are implementing biosecurity in their day-to-day routines.

Visit [www.youtube.com/CottonInfoAust](http://www.youtube.com/CottonInfoAust) for more information.



March 2021

## Biosecurity Management Plan for broadacre industries

|  |  |
|--|--|
| Property Name:   | Property Address:  |
| Owner or Manager:                                      | Contact number:  |
| Review date:<br><small>Review every 12 months.</small> |  |
| Biosecurity Management Plan completed by:              | <b>Emergency Plant Pest Hotline</b><br><b>1800 084 881</b> |
| <small>(Name, signature and date)</small>              |  |



### Contents

|  |    |
|--|----|
| Why develop a farm biosecurity management plan?    | 2  |
| Biosecurity legislation                            | 2  |
| Developing your farm biosecurity management plan   | 3  |
| People, vehicles and equipment                     | 4  |
| Cropping and production practices                  | 8  |
| Training, planning and recording                   | 11 |
| Action Plan for implementing biosecurity practices | 14 |

*This is a Biosecurity Management Plan in accordance with Section 94G(4) of the Queensland Biosecurity Regulation 2016 and clause 44B of the New South Wales Biosecurity Regulation 2017. The purpose of this Biosecurity Management Plan is to state the measures to prevent, control or stop the spread of biosecurity matter into, at, or from the management areas defined in the plan; as described in relevant Australian state and territory biosecurity legislation. Various obligations apply to areas where a Biosecurity Management Plan is in place. For details of these of these obligations, refer to the relevant legislation in your jurisdiction. People entering areas where a Biosecurity Management Plan applies need to be aware of their obligations under the plan and must comply with the measures outlined.*





*Ground Up Protection for the Australian Cotton Industry*

 eChem Australia Pty Ltd is;

-  100% Australian owned company
-  Australian Products for Australian conditions
-  Longstanding relationships with the highest quality manufacturers
-  Fully QA tested
-  Committed to Australian cotton growers

## Our Products

### PGR'S & Defoliants

- e-Tron Defoliation Oil
- Ethephon 900
- Mepiquat 38
- Mepiquat 380
- Parasol Insecticide & Defoliant Oil
- Thidiazuron 500 SC
- Thi-Ultra SC

### Herbicides

- Dicamba 500
- Glyphosate 450
- Glyphosate 540 K
- Halox 520
- Paraquat 250
- Pendimethalin 330
- S-Metol 960
- Staroxy 200 EC
- Staroxy 400 EC
- Triclopyr 600

### Insecticides

- Abamectin
- AceTam 225
- Alpha-Cyp 100 Duo.
- Amitraz 200 EC/ULV
- Bifenthrin 100 EC
- Chlorpyrifos 500
- Difen 500
- Indox 150
- Pyrip 100

### Seed Treatments

- Genero 600 SC
- Methoxam 600



**Contact us now**

[www.echem.com.au](http://www.echem.com.au)

 1300 781 649

# High priority exotic pests and diseases

**A** risk assessment carried out during the development of the Biosecurity Plan for the Cotton Industry identified exotic pests that could significantly impact cotton production if they arrived and established in Australia. It is important that people working on-farm are aware of these pests as well as what to do if they see any unusual plant symptoms or pests to improve the chance of an effective control and eradication response.

## Exotic pests

**These pests have not yet been detected in Australia. Report suspected incursions to the exotic plant pest hotline on 1800 084 881.**

### Cotton boll weevil

*Anthonomus grandis*

A specific pest of cotton, the cotton boll weevil damages developing bolls leading to large yield losses. In the USA, the cotton boll weevil eradication program has been largely successful, but at a cost of hundreds of millions of dollars.



**Boll weevil.** (Alton N. Sparks, Jr., University of Georgia, [Bugwood.org](http://Bugwood.org))

### Brown marmorated stink bug (BMSB)

*Halyomorpha halys*

Stink bugs have a shield-shaped body and emit a pungent odour when disturbed. A number of native stink bugs look similar to BMSB, but lack the distinctive white bands on the antennae (and legs of nymphs). BMSB can damage more advanced bolls than other stink bug species.

**CottonInfo insect ID guide on brown marmorated stink bug**  
[www.cottoninfo.com.au/publications/insect-id-guide-endemics-exotics-brown-marmorated-stink-bug](http://www.cottoninfo.com.au/publications/insect-id-guide-endemics-exotics-brown-marmorated-stink-bug)



**Brown marmorated stink bug.** (Steven Valley, Oregon Department of Agriculture, [Bugwood.org](http://Bugwood.org))

### Cotton stainer

*Dysdercus* spp.

Related to the already endemic pale cotton stainers (see page 41), these colourful red bugs are often found in groups. While feeding on cotton bolls they can transfer microorganisms that increase lint staining.



**Adults and nymphs of the cotton stainer, *Dysdercus suturellus*.** (Lyle J. Buss, University of Florida. [entnemdept.ufl.edu/creatures/field/bugs/cotton\\_stainer.htm](http://entnemdept.ufl.edu/creatures/field/bugs/cotton_stainer.htm))

### False codling moth

*Thaumatotibia leucotreta*

False codling moth is a pest of economic importance to many crops in its native habitat including avocado, citrus, corn, macadamia, peach and plum. Adult false codling moths are small, brownish-gray moths up to 20 mm, with a triangular mark on the outer part of the wing with a crescent shaped mark above it.



**False codling moth larvae (left) and female moth (right).** (Marja van der Straten, NVWA Plant Protection Service [bugwood.org](http://bugwood.org) (left) and [idtools.org](http://idtools.org) (right))



## Lygus bugs

*Lygus lineolaris* and *L. hesperus*

Tarnished plant bug and western plant bug occur predominantly in North America and have a wide host range. In cotton, feeding causes seed abortion, stem or leaf wilting and poor seed germination. It is likely that control of these plant bugs would be very disruptive to the current Australian IPM system.



**Tarnished plant bug.** (Scott Bauer, USDA Agricultural Research Service, Bugwood.org)

## Cotton leaf curl disease (CLCuD)

CLCuD, sometimes referred to as Gemini virus, can cause yield losses of up to 35% in cotton. It is spread by whiteflies. There are at least seven different begomoviruses and several different DNA satellite molecules associated with CLCuD. A cotton plant needs to be infected with at least one begomovirus and one satellite to develop CLCuD.

Symptoms of CLCuD initially appear as a swelling and darkening of leaf veins, followed by a deep downward cupping of the youngest leaves then either an upward or downward curling of the leaf margins. Leaf-like structures (enations) on the veins are common and vary in size from only a few millimetres in diameter to almost the size of a normal leaf. These larger structures are often cup-shaped.



**Leaf curl disease.** (Cherie Gambley, Qld DAF)

## Texas root rot

*Phymatotrichopsis omnivore*

Texas root rot is an extremely damaging fungal disease with a wide host range. It causes sudden death, usually during the warmer months. In cotton, infection can result in 100% crop loss. If this disease became established in Australia, control would be extremely difficult as management using rotations and fungicides is usually only partially effective.

Symptoms include yellowing or bronzing of leaves, which then wilt and die but usually remain attached to the plant. At this stage, the roots are dead and their surface is covered with a network of tan fungal strands.



**Texas root rot.** (Chris Anderson, NSW DPI)

## Blue disease

*Cotton leafroll dwarf virus*

Blue disease is a virus specific to cotton that can reduce yield potential by up to 20%. It is spread by a vector, the cotton aphid. It has been associated with plants infected with cotton leafroll dwarf virus (CLRVD) and has similarities with cotton bunchy top, anthocyanosis and cotton leaf roll. It is not known if the same pathogen causes all these diseases or if there are multiple pathogens causing similar symptoms. CLRVD has not been detected in Australian cotton affected by cotton bunchy top disease.

Affected leaves tend to be smaller, thicker, more leathery and brittle and are intense green to bluish with yellow veins. Reddening of stem petioles and leaf veins can occur in some infections. Leaf edges tend to roll downwards and under and plants become stunted (due to a shortened branch internodes) and produce many branches, giving a bunchy zig-zag stem habit. Symptoms and stunting are more obvious in plants infected at an early age. Infected plants also produce smaller bolls and boll shedding may occur. Single infected plants can be difficult to spot if overgrown by nearby healthy plants.



**Blue disease.** (Murray Sharman, Qld DAF)

# Are your usual sprays or control measures not working?

Some of the industry's high priority exotics are variants of endemic pests and diseases. These exotic strains could become a problem if they reached Australia, either through different insecticide resistance profiles or the potential to overcome host plant resistance bred into Australian cotton varieties.

It is important to contact the relevant researcher if you've experienced an insecticide spray failure without a reasonable explanation, or are seeing new or unusual disease symptoms to organise the appropriate sampling and testing.

## Bt-resistant cotton bollworm

*Helicoverpa armigera* (carrying Bt resistance alleles)

The introduction of Bt cotton has dramatically reduced the need to control *Helicoverpa*. Exotic strains of *H. armigera* that carry resistance alleles (e.g. dominant resistance to Cry1Ac) would have a significant impact on Australia's cotton industry if they were to establish here. With the movement of *H. armigera* into South America, there is also some concern that *Helicoverpa zea* (American cotton bollworm) may hybridise with *H. armigera*.

## Bt-resistant fall armyworm

*Spodoptera frugiperda* (carrying Bt resistance alleles)

Fall armyworm was detected in Australia in early 2020 and has been declared endemic. Similar to *H. armigera*, there are also exotic Bt resistant strains of *S. frugiperda* that could have an effect on Australia's cotton industry if they were to become established in Australia.

## Cotton aphid

*Aphis gossypii* (exotic strains)

Cotton aphids are potential vectors for exotic pathogens such as blue disease. As well as the risk of disease, there is a risk that new aphid strains entering the country will have different insecticide resistance profiles, making control more difficult.

## Bacterial blight

*Xanthomonas axonopodis* or *X. campestris* pv. *mavacearum* (exotic strains)

Although strains of bacterial blight are already present in Australia, they are no longer a problem due to varietal resistance. Exotic strains (races) occur, however, that are 'hypervirulent' and, if established in Australia, would cause large yield losses.

Initial symptoms include angular water-soaked lesions on the undersides of leaves that dry and darken with age, followed by leaf shedding. Black lesions spread along stem and bolls are often infected at the base or tip (where the dried lesions prevent the boll from opening). This seed-borne pathogen is capable of symptomless transfer and could therefore remain undetected through quarantine.



**Bacterial blight.**

## Verticillium and Fusarium wilts

*Verticillium dahliae* (exotic strains)

*Fusarium oxysporum* f. sp. *vasinfectum* (exotic strains)

While Fusarium and Verticillium wilts already affect Australian cotton, it is important to get new or unusual wilt symptoms tested as this could indicate incursion of exotic strains.

## Whitefly

Exotic *Bemisia* species

There is a risk of further incursions of exotic *Bemisia* species, with different genetic profiles to current pest populations (e.g. additional forms of insecticide resistance) entering the country. Invasive whitefly could potentially introduce viruses (such as GLCuD) that they vector.



**IF YOU SEE ANYTHING UNUSUAL,  
CALL THE EXOTIC PLANT PEST HOTLINE**



**1800 084 881**




## Quality **CROP DEFOLIATION** products for the Australian Cotton Industry



eChem specialising in high quality agricultural chemicals specifically developed for Australian conditions. eChem draws upon a wealth of expertise and industry knowledge to bring you the very best for your farming operation.

### **Our PGR'S & Defoliants Products**

- 
- e-Tron Defoliation Oil
  - Ethephon 900
  - Mepiquat 38
  - Mepiquat 380
  - Parasol Insecticide & Defoliant Oil
  - Thidiazuron 500 SC
  - Thi-Ultra SC

*Ground Up Protection for the Australian Cotton Industry*

**☎ 1300 781 649**

**[www.echem.com.au](http://www.echem.com.au)**

# Cotton growth regulators and harvest aids

## Plant growth regulators

Cotton is very responsive to management and environmental changes. Excessive vegetative growth may reduce the retention of fruit and delay crop maturity, and can also inhibit the efficacy of insecticide by poor penetration into the canopy. Mepiquat chloride can help to manage crop growth by reducing the production of gibberellic acid. Gibberellins are natural growth regulators in plants, and play an important role in cell expansion.

There are many factors that should be considered when making the decision to apply mepiquat chloride, the most critical being whether there are other sources of stress already controlling growth (such as temperature or soil moisture), and the rate and timing of the application.

For more information refer to the latest *Australian Cotton Production Manual* (ACPM). Note: Some defoliant products containing ethephon are labelled as a 'Growth Regulator'. Ethephon is used for preparing the crop for harvest and may cause significant fruit loss if used at inappropriate times.

## Types of harvest aids

The categories of harvest-aid chemicals include herbicidal and hormonal defoliants, boll conditioners/openers and desiccants. Each has a different mode of action.

### Defoliants

Defoliants (e.g. thidiazuron, diuron, dimethipin) induce leaf abscission (a break in the cellular structure joining the leaf to the stem allowing the leaf to fall off). Leaf removal is critical for reducing the amount of leaf trash in machine harvesters. Boll opening is also accelerated by defoliation as removal of leaves exposes bolls to more direct sunlight, promoting increased temperatures for maturation, and drying and cracking of the boll walls. Hormonal defoliants reduce auxin and/or enhance ethylene production, while herbicide defoliants injure or stress the plant into increasing ethylene production (similar to waterlogging or drought effects). The safe timing of defoliation is when the youngest boll expected to reach harvest is physiologically mature. This usually occurs when 60-65% of bolls are open.

In addition to timing of harvest aids, it is important to consider product, rate and application issues. If herbicide rates are too high, the plant material may die before releasing enough ethylene to cause defoliation, resulting instead in leaf desiccation (leaf death).

A number of factors improve the performance of defoliation products: defoliate before the onset of frost; aim to have soil moisture at refill at defoliation; and, avoid using defoliants when there is a risk of rainfall shortly afterwards. For more information on defoliation refer to the ACPM.

### Boll conditioning/openers

Boll conditioning/openers (e.g. ethephon, cyclanillide, amino methanamide dihydrogen tetraoxosulfate) specifically enhance ethylene production by providing a chemical precursor, leading to quicker separation of boll walls (carpels). Only consider using boll conditioning/opener products if the bolls that will be forced open are mature; if applied prior to boll maturation these products may cause bolls to shed and reduce yield.

### Desiccants and herbicides

Desiccants and herbicides (e.g. sodium chlorate, magnesium chlorate, glyphosate, diquat, paraquat, carfentrazone-ethyl) cause disruption of leaf membrane integrity, leading to rapid loss of moisture. Usually, desiccants should be avoided as they dry all plant parts (including stems) which can increase the trash content of harvested lint, however their use may be necessary if conditions do not enable the effective use of defoliants (e.g. very cold weather). Desiccants are also a reliable method to reduce leaf regrowth. High rates of some defoliants can act as desiccants.

### Wetting agents and spray adjuvants

Wetting agents and spray adjuvants (e.g. paraffinic oil, non-ionic surfactants or petroleum oil) are used to assist with defoliation as cool temperatures, low humidity and water stress prior to defoliant application can result in increased wax and thickness of the leaf cuticle, reducing the efficiency of chemical uptake.

Many growers use combinations of defoliants with different modes of action and multiple applications to enhance defoliation. Multiple applications are becoming more common due to the more indeterminate growth habit of current varieties.

Registration of a chemical is not a recommendation for the use of a specific chemical in a particular situation. Satisfy yourself that the chemical you choose is the best one for the crop and situation.

Carefully study the container label before use and note any specific instructions relating to the rate, timing, application and safety. Take every precaution to minimise the risk of causing or suffering spray drift damage

**More information is available in a CottonInfo factsheet on defoliation**  
[www.cottoninfo.com.au/blog/defoliation-what-do-consultants-say](http://www.cottoninfo.com.au/blog/defoliation-what-do-consultants-say)

**APVMA's PubCRIS database can be searched for current registrations and permits.**  
[portal.apvma.gov.au/pubcris](http://portal.apvma.gov.au/pubcris)

**CottonInfo videos:**

**What is pix?** [youtu.be/cIFpyyWW6Q](https://youtu.be/cIFpyyWW6Q)

**Using harvest aids in cotton** [youtu.be/8st4yggcCNY](https://youtu.be/8st4yggcCNY)



# Best practice for spray application

**W**hen using pesticides, best practice means not only doing the best job you possibly can, but also being able to demonstrate what you have done and how it has impacted others.

Movement of spray beyond the target area is undesirable as it represents wastage of product and exposure of sensitive non-target areas to potentially damaging materials. Achieving the best outcome from spray application requires the careful consideration of many factors. Application technique needs to be matched to the target, tank mix and weather conditions.

## Plan ahead

The development of a comprehensive Pesticide Application Management Plan (PAMP) prior to the season is an important part of the Best Management Practice (*myBMP*) program for cotton. Include:

- Farm layout;
- Record keeping;
- Weather monitoring;
- Identification of sensitive areas and potential hazards; and,
- Communications procedures and complaint handling.

Having a PAMP in place helps to ensure that everyone involved in pesticide application has a clear understanding of their responsibilities.

## Legal requirements

Be aware of federal and state regulations for chemical application.

### Top tips for agronomists...

1. Run through a pre-season spray plan with your growers (see module 5 of the GRDC spray application manual) and have a conversation if you suspect that spray practices can be improved.
2. Ensure the grower has realistic expectations of timeframes for pest control.
3. Check that your grower's spray rig set-up is fit-for-purpose to deliver your spray recommendations.
4. Highlight the importance of identifying spray drift factors and not spraying under inversion conditions (including the necessity of halting the spray operation if conditions become unfavourable).
5. Ensure the grower is aware of factors that may impact on spray coverage and performance (nozzle selection, pressures, speed, adjuvants, water quality, tank mix compatibilities etc)
6. Recommend a coarser nozzle to reduce spray drift risk and higher water rates to improve efficacy of pest control.
7. Highlight the importance of reading and understanding the labels of any products used.
8. Follow up on recommendations by assessing efficacy and efficiency post-spray.

Staff responsible for handling and applying pesticides must be qualified according to relevant state and federal requirements.

There may also be workplace health and safety requirements related to storage and use of hazardous chemicals, such as completing a hazard analysis, maintaining a chemical inventory and easily accessible current copies of relevant Safety Data Sheets.

**Always read and follow the label when handling and applying chemicals.** Label conditions may specify spray quality, spray conditions including mandatory wind speed range, and no-spray zones/buffers.

Users are not absolved from compliance with the directions on the label or the conditions of the permit by reason of any statement made or not made in this publication.

## Mandatory label instructions

Many product labels now include a range of mandatory requirements. Some examples include:

### Spray quality

A typical requirement is the use of a minimum spray quality according to the American society of Agricultural and Biological Engineers (ASABE) or British Crop Production Council (BCPC) classification systems. Select nozzles from charts that refer to either of these standards and ensure that equipment is set up and used appropriately to achieve the required spray quality.

### Wind speeds

Labels often include that the wind speed must be above 3 km/h and less than either 15 km/h or 20 km/h (depending on the product) as measured at the site of application.

### No-spray zones

Products must not be applied if a sensitive area (such as a residence, school, public area, water body, pasture, terrestrial vegetation or another crop) is within a nominated downwind no-spray zone. The distance required for the no-spray zone may differ for the various types of sensitive areas.

Always check the label for no-spray zone requirements for the product and the specific tank mix you wish to use. No-spray zones for aerial applications can be larger than those required for ground application.

### Record keeping

Record keeping requirements are included on the label or permit conditions for many products. These records must be maintained in addition to any state-based requirement for record keeping.

### Good communication

Prior to spray application and product selection, check the proximity of susceptible crops and sensitive areas. BeeConnected can help identify nearby location of hives and facilitate communication between spray applicators and beekeepers ([www.beeconnected.org.au](http://www.beeconnected.org.au)). It is good practice to notify neighbours and staff of your spray intentions, regardless of label requirements.

Cotton is extremely sensitive to phenoxy herbicide drift.

**SataCrop is a free online mapping tool that maps all crops, not just cotton, allowing growers and applicators of all products to be better informed of the location of potentially sensitive crops before spraying.** <https://satacrop.com.au>

## Monitoring and recording conditions

Labels contain a legal requirement to measure weather parameters (including temperature, humidity, and wind speed and direction) at the site and time of application. This can be done with handheld equipment (e.g. Kestrel or equivalent) or portable weather stations. On-board weather stations that provide live weather information while the sprayer is operating (such as the Watchdog systems) are also available.

There are also subscription websites that provide forecasts of conditions for spraying up to 10 days in advance. These sites evaluate a range of factors to produce tables indicating suitable times for spraying. For example, see [spraywisecisions.com.au](http://spraywisecisions.com.au) or [syngenta.com.au](http://syngenta.com.au)  
**myBMP resources:** [Fact sheet on weather monitoring equipment.](#)

## Temperature and humidity

Higher ambient air temperatures and lower relative humidity conditions increase evaporation rates, which will decrease droplet size and increase the risk of drift. Water-based sprays should not be applied under these conditions. Spraying is best conducted when the Delta T (the difference between the wet bulb and dry bulb) is in the range 2-10°C (or up to 12°C where targets are not stressed and a coarse spray quality or larger is used).  
**myBMP resources:** [Fact sheet on Tips to Reduce Spray Drift for a Delta T Chart.](#)

When using coarse sprays at high water volumes, evaporation may be less significant, which may allow some applications to continue into marginal Delta T conditions (where soil moisture exists, and the targets are not in a stressed condition). Never start a spraying operation when the Delta T is below 2 or above 10-12.

**Further information:** [www.grdc.com.au/GRDC-FS-SprayPracticalTips](http://www.grdc.com.au/GRDC-FS-SprayPracticalTips)


## Surface temperature inversions

DO NOT apply if there are surface temperature inversion conditions present at the application site during the time of application. These conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

**For more information refer to the GRDC factsheets on Surface Temperature Inversions ([grdc.com.au/GRDC-FS-SprayInversions](http://grdc.com.au/GRDC-FS-SprayInversions)) or refer to the 24 hour spray diagram on page 128.**

### Three common states of atmospheric stability and their relationship to spray application.

(SOURCE: GRDC Weather Essentials for Pesticide Application booklet, 2017; author: Graeme Tepper)



## Vegetative plantings as spray drift barriers

Effective vegetative barriers (a mixture of tree and shrub species with foliage all the way to the ground) can reduce drift by up to 90%. The planting arrangement and density should allow for air to partly flow through – a barrier that is too dense will act like a wall, directing wind containing the spray drift up and over the top and increasing how far drift may travel. Do not locate vegetative barriers where airflow will be obstructed by adjacent objects such as turkey's nests, water storages or large banks.

Trees and shrubs with long thin or needle-like leaves, or hairy leaf surfaces are the most effective at trapping airborne droplets. Choose hardy and drought tolerant species with thick cuticles to help them survive small doses of pesticide.

Most guidelines suggest that the optimum barrier width is 20 m with a 10 m maintenance strip on either side. A barrier with a porosity of around 50% (you can see 50% light and 50% dark when you look through the vegetation) should be at least 1.5 times the release height of the spray. As vegetation density increases, the height of the barrier needs to be increased (e.g. at 40% the height should be 2 times the release height for the spray).

Do not utilise remnant native vegetation or environmental corridors as a vegetative barrier. This habitat vegetation should be identified as 'sensitive' along with riparian areas and waterways and protected from spray drift.

**CottonInfo NRM/Pesticide Input Efficiency fact sheet – Using vegetative barriers to minimise spray drift on cotton farms [www.cottoninfo.com.au/publications/nrmpesticide-input-efficiency-using-vegetative-barriers-minimise-spray-driftcotton](http://www.cottoninfo.com.au/publications/nrmpesticide-input-efficiency-using-vegetative-barriers-minimise-spray-driftcotton)**

## Factors that influence spray drift

Spray application needs to maximise efficacy against the target pest, while minimising any off-target movement or effect. Application technique needs to be matched to the target and weather conditions. Achieving the best outcome from spray application requires the careful consideration of many factors:

- Appropriate spray release height;
- Avoiding excessive travel speed for ground rigs;
- Pressure at the nozzle;
- Suitable water volumes and quality;
- Nozzle selection; and,
- Maintenance and hygiene.

See the *Australian Cotton Production Manual* for more details.

## Decontaminating equipment

Thoroughly decontaminate application equipment after use, particularly when herbicides have been used. The *Australian Cotton Production Manual* includes a decontamination/cleaning product guide. Follow decontamination instructions on pesticide and cleaning product labels.

**CottonInfo video on decontaminating a spray rig [youtu.be/zDQwoIEcB0](https://youtu.be/zDQwoIEcB0)**

PRECISION PAYS



JOHN DEERE

# SEE. SPRAY. SAVE.

**See & Spray™ Select.**  
Gain more efficient herbicide usage.

Available on the new John Deere 400 and 600 Series Sprayers, See & Spray Select's integrated camera technology rapidly detects green in fallow ground automatically triggering a targeted spot application on weeds. That's a significant saving on your herbicide inputs. Make more informed agronomic decisions and gain the leading edge with this first factory installed, complete solution.



Visit our virtual showroom at  
[JohnDeere.com.au/SeeandSpray](https://www.johndeere.com.au/SeeandSpray)

## Ground application

Spray contractors can be used if you do not have the equipment or expertise to apply pesticides yourself. Ensure your spray contractor has suitable training and accreditations and has the appropriate state license. You can search to see if a ground or aerial contractor is licensed in NSW at [apps.epa.nsw.gov.au/prrdlapp/plpr.aspx](http://apps.epa.nsw.gov.au/prrdlapp/plpr.aspx) and in Qld at [www.business.qld.gov.au/industries/farms-fishing-forestry/agriculture/land-management/chemical-controls/licence-search](http://www.business.qld.gov.au/industries/farms-fishing-forestry/agriculture/land-management/chemical-controls/licence-search).

Commonly asked questions are:

- What is the actual application rate?
- How much chemical to put in the tank?
- How many paddock hectares can be treated with each tank load?
- What rate to put in the spray controller? and,
- What nozzles to use to achieve required application rate?

## Calculating banded sprays

Banded sprays present an opportunity for strategic and economical use of products; applying the equivalent rate of active per hectare to the target, but using less pesticide overall.

Label rates are always given as litres of active/sprayed ha. Advisors also give recommendations as L/sprayed ha, however confusion over banded spraying can lead to big differences between the consultant's recommendation, the applicator's instincts and what the application equipment can physically achieve.

How to apply the correct amount of product:

1. The strategic application rate (L of active/ha) is based on the average width (in metres) sprayed by individual nozzles.
2. How much chemical to put in the tank is based on the application rate.
3. What to put into the controller is based on the area of paddock able to be sprayed per tank.

## Calculations

### Application rate

You need to know:

Band width in metres: e.g. **0.7 m band** ÷ 1 m row spacing = band width (m) ÷ row spacing (m).

Sprayed width per nozzle (m): = **band width (m) ÷ number nozzles per band** (e.g. 3 nozzles per 70% band of a 1 m row = 0.7 m ÷ 3 = 0.23 m)

Calculate:

$$\text{L/sprayed ha} = \text{L/min/nozzle} \times 600 \div \text{speed (km/h)} \div \text{sprayed width per nozzle (m)}$$

L/sprayed ha applies to each band (row), whether you spray 1 band (row), or many rows, whether it is a solid plant, single skip or double skip.

### How much chemical per tank

You need to know:

Number of sprayed ha per tank = **Tank size (L) ÷ L/sprayed ha**.

Calculate:

$$\text{Amount of chemical to add per tank} = \text{Sprayed ha per tank} \times \text{chemical rate/ha}$$

## Paddock hectares per tank

Calculate:

$$\text{Paddock ha per tank (solid plant)} = \text{Sprayed ha per tank} \div \text{band width (m)}$$

For skip row configuration, you need to calculate an adjusted band width:

Adjusted band width (m) = **band width (m) x width of boom ÷ row width (m) ÷ number of planted rows under the boom**.

Calculate:

$$\text{Paddock ha per tank (skip)} = \text{Sprayed ha per tank} \div \text{the band width (m) x width of boom} \div \text{row width (m) ÷ number of planted rows under the boom}$$

## Rate to put in the Controller

Calculate:

$$\text{Rate to put in the Controller} = \text{Tank Size (L)} \div \text{Paddock ha per tank}^*$$

\*This works if you don't want to change the section widths in the controller.

## Nozzle selection

To select the correct spray nozzle, you need to calculate the required flow rate of each nozzle (L/min). This is simple if all nozzles are the same size, as the flow rate will be the same for each nozzle.

You need to know:

The average sprayed width per nozzle:

e.g. For 5 nozzles per 1 m row at 100% band: **1 m ÷ 5 = 0.2 m**

For 4 nozzles per 1 m row and a 70% band: **0.7 m ÷ 4 = 0.17 m**

Calculate:

$$\text{Required flow rate per nozzle} \\ \text{L/min/nozzle} = \text{L/sprayed ha} \div 600 \times \text{speed (km/h)} \times \text{average sprayed width per nozzle (m)}$$

Always remember to check the spray quality produced to ensure it is consistent with what is required by the product label.

Useful resources:

- The *myBMP* Pesticide application module, [www.mybmp.com.au](http://www.mybmp.com.au)
- Nufarm Australia Ltd: 03 9282 1000, [www.nufarm.com.au](http://www.nufarm.com.au)
- GRDC fact sheets on:
  - Spray Mixing Requirements
  - Spray Water Quality
  - Pre-season check and Controller Settings
- Information on weather:
  - Weather essentials for pesticide application, Graeme Tepper, GRDC.
  - GRDC Fact Sheet on Weather Monitoring Equipment
- Information on weather forecasting tools:
  - [www.spraywisedecisions.com.au](http://www.spraywisedecisions.com.au)
  - Agricast
- Information on pesticide application:
  - Spraywise Broadacre Application Handbook, Dr Jorg Kitt, Nufarm Australia
- Information on nozzle selection tools:
  - Teejet Nozzle Selection App
  - Hardi Nozzle App

See also [www.grdc.com.au/resources-and-publications/grownotes/technical-manuals/spray-application-manual](http://www.grdc.com.au/resources-and-publications/grownotes/technical-manuals/spray-application-manual)



## Aerial application

Aircraft are extensively used for pesticide application in agriculture. Their advantage lies in being able to cover large areas quickly and in conditions that may not suit ground application (e.g. dense crop cover or wet ground). Aerial operators are professional spray applicators that understand the parameters required for getting the best out of a spray job. Discuss your requirements with the aerial operator, including the target, timeliness, coverage required, and sensitive crops nearby to assist them in ensuring optimal set up for the aircraft. The best results are achieved when growers and agronomists work closely with their aerial operator in planning, executing and following up on their spray applications.

Some key points that should be discussed are:

- **Target** – e.g. the pest's location within crop canopy or field, or if there are time constraints related to the pest's development.
- **Water rates** – efficiency is increased by using lower water rates, but it is important to understand the potential impacts on efficacy. It is critical that the correct coverage is achieved (as specified on the product label). Minimum water rates should be adhered to and in some cases will need to be increased (e.g. to match weather parameters or to provide sufficient volume to carry all of the products in the tank).
- **Droplet Size** – aircraft are good at producing small droplets that are generally good for efficacy but may increase the risk of off-target movement. Some product labels specify a certain number of droplets per square cm on the target. Aerial operators are well positioned to select the correct droplet size to achieve this.

- **Swath width** – A spray plane's swath width is wider than the wingspan, due to the vortex effect created by the wings as they move through the air. Each aircraft has recommended swath widths that they can use in various conditions. Many aerial operators do regular pattern testing as well to optimise the swath width that they use for each aircraft. It is good to review this data with your aerial operator to ensure that all parties are satisfied with the spray parameters being used.

It is highly recommended that growers select an aerial applicator with Aerial Application Association of Australia (AAAA) membership, Spraysafe or AIMS accreditations to ensure that they are getting the most professional service.

The AAAA has produced some comprehensive documents about aerial spraying to assist with working more effectively with your aerial operator. Visit the AAAA website [www.aaaa.org.au](http://www.aaaa.org.au).

**Further information:**

**2021 Australian Cotton Production Manual**

**GRDC Spray application manual** [www.grdc.com.au/resources-and-publications/grownotes/technical-manuals/spray-application-manual](http://www.grdc.com.au/resources-and-publications/grownotes/technical-manuals/spray-application-manual)

**Mary O'Brien Rural Spray Drift resources** [www.maryobrienrural.com.au/resources/](http://www.maryobrienrural.com.au/resources/)

**SPRAYpak – Cotton Grower's Spray Application Handbook, 2nd edition, available from** [www.cottoninfo.com.au/publications/spraypak](http://www.cottoninfo.com.au/publications/spraypak)

**Spraywise – Broadacre Application Guide – Available through Croplands Distributors.**

**Nozzle selection for boom, band and shielded spraying. The Back Pocket Guide published by GRDC** [www.grdc.com.au/resources-and-publications/all-publications/publications/2019/grdc-nozzle-selection-guide](http://www.grdc.com.au/resources-and-publications/all-publications/publications/2019/grdc-nozzle-selection-guide)

**For more information about using vegetative barriers to minimise spray drift on cotton farms, see factsheet:** [www.cottoninfo.com.au/publications/nrmpesticideinput-efficiency-using-vegetative-barriers-minimise-spray-drift-cotton](http://www.cottoninfo.com.au/publications/nrmpesticideinput-efficiency-using-vegetative-barriers-minimise-spray-drift-cotton)

**Additional resources can be found at** [www.myBMP.com.au](http://www.myBMP.com.au)


|||



For more information on best practice for aerial and ground spray application go to [www.myBMP.com.au](http://www.myBMP.com.au). (Jack Hawkins, Cotton Australia)

# 24 Hour risk profile for summer spraying

Always follow label instructions



For more information visit [nufarm.com.au](http://nufarm.com.au)

© 2018 Nufarm Australia Ltd. All trade marks (®, ™) are owned by Nufarm Australia Ltd or used under license

## The new 400 and 600 Series Sprayers turn technology into application quality.

Reduce overlaps and gain chemical cost savings with the ExactApply™ nozzle-control system that helps you spray with greater accuracy. New boom pressure recirculation and product reclaim save precious product, time and expense when rinsing and priming the boom while switching solutions. Plus an all-new cab with enhanced operator comfort means you'll cover more ground per day in maximum comfort.



**JOHN DEERE**



Visit our virtual showroom at  
[JohnDeere.com.au/Sprayers](https://JohnDeere.com.au/Sprayers)

**GAIN ACCURACY**  
**GAIN SAVINGS**  
**GAIN INSIGHT**

**PRECISION PAYS**

# Pesticides and regulation

## APVMA product registration

The federal, state and territory governments work together to regulate agricultural and veterinary (agvet) chemicals in Australia. The Australian Pesticides and Veterinary Medicines Authority (APVMA) evaluates the safety and performance of agvet chemicals intended for sale in Australia to protect the health and safety of people, animals, crops, trade and the environment.

Before agvet chemical products can be legally sold, supplied, marketed or bought in Australia, they must be registered by the APVMA. The assessment uses broad risk analysis, including how human and environmental exposure can be minimised through instructions for use and safety directions on the label of registered products. The APVMA approves product labels that explain how the product is to be used, stored, disposed of and managed in the event of poisoning.

Agvet chemical product labels are legally binding. State and territory governments are responsible for monitoring post-sale compliance with label instructions and limitations.

### Using the APVMA website

The APVMA Public Chemicals Registration Information System (PubCRIS) database can be searched to find agvet chemical products, active constituents and labels approved and registered for use in Australia at: [portal.apvma.gov.au/pubcris](http://portal.apvma.gov.au/pubcris)

Products approved for minor use or under temporary emergency use permits can be searched at [portal.apvma.gov.au/permits](http://portal.apvma.gov.au/permits)

### Applying for variations or new registrations

The APVMA site includes guidance for applicants on the data and supporting information required to meet APVMA criteria when varying a registered product label or submitting a new product.

Access tailored guidance for applicants at: [apvma.gov.au/node/27441](http://apvma.gov.au/node/27441)

### Chemical reviews

The APVMA may undertake a formal review (or reconsideration) to scientifically reassess the risks and determine whether changes are needed to ensure the chemical can continue to be used safely and effectively. This may include modifying the way chemicals are used or if risks cannot be managed, removing chemicals from the market.

A list of products currently under review is available at: [apvma.gov.au/chemicals-and-products/chemical-review/listing](http://apvma.gov.au/chemicals-and-products/chemical-review/listing)

## Legal responsibilities

All agricultural chemical applications MUST accord with the approved label instructions or permit conditions for that particular registered agricultural chemical, crop, pest and region. State regulations detail additional requirements associated with the use of agricultural chemicals including record keeping, and training and licensing requirements for applicators.

### New South Wales

The *Pesticides Act 1999* is the primary legislative instrument controlling the use of pesticides in NSW and is administered by the Environment Protection Authority (EPA). The underlying principle of the Pesticides Act is that pesticides must only be used for the purpose described on the product label and all the instructions on the label must be followed. Consequently, all label directions must be read by or explained to the user prior to each use of the pesticide.

All pesticide users should take reasonable care to protect their own health and the health of others when using a pesticide. They should also make every reasonable attempt to prevent damage occurring from the use of a pesticide, such as off-target drift onto sensitive areas or harm to endangered and protected species.

The Pesticides Regulation 2017 contains licensing and training provisions and requires all commercial pesticide users, i.e. all farmers and spray contractors, to keep records of their pesticide application.

No set form is required for records but they must include:

- Full product name;
- Description of the crop or situation;
- Rate of application and quantity applied;
- Description of the equipment used;
- Address of the property, identification of the area treated and order of paddocks treated;
- Date and time of the application (including start and finish);
- Name, address, and contact details of the applicator and of the employer or owner if an employee or contractor is the applicator;
- Estimated wind speed and direction (including any significant changes during application); and,
- Other weather conditions specified on label as being relevant (e.g. temperature, rainfall, relative humidity).

Some products have additional record keeping requirements determined on the label (so are national requirements) e.g. the new 2,4-D label requires that nozzle records are also kept.

An example form that captures all the information required by the Pesticides Regulation 2017 is available on page 134 or can be downloaded from the NSW DPI website ([www.dpi.nsw.gov.au/agriculture/chemicals/farm-chemical-management/records](http://www.dpi.nsw.gov.au/agriculture/chemicals/farm-chemical-management/records)).

The record must be made as soon as practicable after the use of the pesticide concerned and, in any event, no later than 48 hours after the pesticide is used. A copy of the record must be kept for at least three years.

**For further information about NSW Pesticides Regulation 2017 please refer to [legislation.nsw.gov.au/#/view/regulation/2017-448/part4/sec36](http://legislation.nsw.gov.au/#/view/regulation/2017-448/part4/sec36)**

All fee-for-service pesticide users (aerial and ground application) must also be licensed. You can search to see if a ground or aerial contractor is licensed in NSW at [apps.epa.nsw.gov.au/prrdlapp/plpr.aspx](http://apps.epa.nsw.gov.au/prrdlapp/plpr.aspx)

The Pesticides Regulation 2017 also requires all commercial pesticide users to be trained in pesticide application and hold a prescribed qualification. Only domestic use, such as home gardens, is excluded, provided the pesticide is a specific domestic/home garden product and used in quantities allowed through the domestic like use exemption.

The minimum level of competency in pesticide use required under the Pesticides Regulation 2017 is Australian Qualifications Framework Level 2 (AQF2). This is for people applying pesticides under supervision (AHCCHM201) where the supervisor must be AQF3 certified.

If you are working as an unsupervised operator or independent business person, you need training in Australian Qualifications Framework Level 3 (AQF3) (AHCCHM 303 and 304).

**Further detail is available at [www.epa.nsw.gov.au/your-environment/pesticides/compulsory-training-pesticides](http://www.epa.nsw.gov.au/your-environment/pesticides/compulsory-training-pesticides)**

Growers are recommended to undertake the SMARTtrain course, Chemical Application, or the standard ChemCert course, both of which cover the higher AQF3 competencies.

An important change in the 2017 regulation is cotton growers who are accredited under a quality assurance (QA) program (*myBMP* accreditation), that includes chemical recording and management records are no longer obliged to retrain every 5 years.

**Further information available at:**  
**[www.epa.nsw.gov.au/your-environment/pesticides/compulsory-training-pesticides](http://www.epa.nsw.gov.au/your-environment/pesticides/compulsory-training-pesticides)**

The EPA can be contacted on [www.epa.nsw.gov.au](http://www.epa.nsw.gov.au) phone 131 555 or [info@epa.nsw.gov.au](mailto:info@epa.nsw.gov.au) or your regional EPA office.

**For additional information on legal responsibilities in applying pesticides in NSW, refer to Primefact 1464, [www.dpi.nsw.gov.au/data/assets/pdf\\_file/0009/186390/legal-responsibilities-in-applying-pesticides-F.pdf](http://www.dpi.nsw.gov.au/data/assets/pdf_file/0009/186390/legal-responsibilities-in-applying-pesticides-F.pdf)**

## Queensland

In Queensland, the *Chemical Usage (Agricultural and Veterinary) Control Act 1988* (Chem Usage Act) sets legal requirements on how agricultural chemicals must be used. All chemical users are required to:

- Use agricultural chemical products which are currently registered with the Australian Pesticides and Veterinary Medicines Authority (APVMA);
- Use the products for the crop or situations specified on the approved label instructions or under the conditions of a permit granted by the APVMA; and,
- Apply agricultural chemical products according to all other label instructions, including any use instructions or restraints that may be listed, such as those which specify droplet size, wind speed and direction, mandatory downwind no-spray zones and other off-target spray drift reduction, risk management practices.

On 19 June 2020, the Chemical Usage (Agricultural and Veterinary) Control Regulation 2017 was amended to incorporate nationally agreed record-keeping and training requirements, as part of national reforms to the agvet chemical regulatory system. All chemical users are therefore also required to:

- Make and keep spray records for a minimum of two years, for all commercial use of any agricultural chemical. For details of the relevant particulars which must be recorded, please refer to [www.business.qld.gov.au/industries/farms-fishing-forestry/agriculture/land-management/chemical-controls/using-chemicals/keeping-records](http://www.business.qld.gov.au/industries/farms-fishing-forestry/agriculture/land-management/chemical-controls/using-chemicals/keeping-records) (Workplace health and safety also requires spray records to be maintained).
- Ensure they have completed appropriate training for any use of certain prescribed hazardous chemicals. For the majority of cotton growers, the relevant chemicals will be those which contain Schedule 7 poisons,

such as paraquat, MSMA and methomyl. The training requirements for use of agricultural chemicals containing Schedule 7 poisons take effect from 19 June 2021. The most appropriate vocational education and training competency units for cotton growers to meet this training obligation will generally be AHCCHM304 (Transport and store chemicals) and AHCCHM307 (Prepare and apply chemicals to control pest, weeds and diseases). Any training believed to be equivalent will be considered. For example, AHCCHM307 has superseded and is regarded as equivalent to AHCCHM303 (Prepare and apply chemicals).

There are significant penalties that apply to anyone found to have breached the Chem Usage Act for failing to follow label instructions.

Under the *Agricultural Chemicals Distribution Control Act 1966* (ACDC Act) aerial distribution contractors and ground distribution contractors must be licensed. Pilots and ground spray operators working for these contractors must also be licensed. The ACDC Act makes provision for the licensing which applies to unmanned aerial vehicles (UAVs) or “drones” for the application of pesticides in Queensland. An aerial distribution contractor licence and pilot chemical rating licence are required for agricultural chemical spraying with a UAV. Exemptions which may apply to Commonwealth Civil Aviation Safety Authority (CASA) authorisations, for certain types of UAVs used on privately-owned land, do not apply for State chemical applicator licensing purposes.

In most instances, cotton growers applying agricultural chemicals with ground equipment on their own land do not need to hold a licence. However, growers are strongly encouraged to complete chemical application training to improve their skills and knowledge in application technology, handling, storing and transporting chemicals.

**For additional advice on legal responsibilities in applying pesticides in Qld, contact Biosecurity Queensland on 13 25 23.**

## Other states

Please refer to your relevant state department for information on legal responsibilities for use of pesticides:

**NT** – [nt.gov.au/industry/agriculture/farm-management/using-chemicals-responsibly](http://nt.gov.au/industry/agriculture/farm-management/using-chemicals-responsibly)

**VIC** – [agriculture.vic.gov.au/farm-management/chemicals](http://agriculture.vic.gov.au/farm-management/chemicals)

**WA** – [www.agric.wa.gov.au/pests-weeds-diseases/control-methods/chemicals](http://www.agric.wa.gov.au/pests-weeds-diseases/control-methods/chemicals)

## Keeping records

Keeping records of pesticide applications is a legal requirement (the amount of detail and length of time records need to be maintained will depend on the label and your state regulations). An example of a pesticide application record sheet developed by NSW DPI can be found on page 134 or at [www.dpi.nsw.gov.au/agriculture/chemicals/farm-chemical-management/records](http://www.dpi.nsw.gov.au/agriculture/chemicals/farm-chemical-management/records)

Several commercial websites, including Spraywise ([www.sprayshop.com.au/product/spraywise-logbook-spraywiselog](http://www.sprayshop.com.au/product/spraywise-logbook-spraywiselog)) have record forms available, or there are various paddock recording programs that can be used to keep records on computers or mobile devices.

# Using pesticides safely

## Post-spray periods

Pesticide and other agrochemical labels will usually specify a minimum amount of time after application before:

- The treated field can be safely re-entered by unprotected persons (re-entry period); and,
- The crop can be harvested, or grazed/cut for domestic stock fodder (withholding period).

Always check the label of each product in the tank mix for these periods before use. Note they are subject to change and may be different according to concentration or formulation, even if the products have the same active ingredient.

Non-compliance with current label restrictions may result in prosecution under relevant state or territory legislation.

## Re-entry period

Re-entry periods and any specific personal protective equipment (PPE) to be worn provide a guide to the risk associated with dermal (skin) contact with spray deposits. Consider re-entry periods when developing workplace health and safety procedures and protocols. More information on WH&S is available in the *Australian Cotton Production Manual*.

All workers must be advised of the appropriate time to wait before re-entering a sprayed field. It is particularly important to observe the re-entry period when contact between foliage and skin is likely. Also exercise caution when entering previously sprayed crops if foliage is wet from dew or light rain, irrespective of the time between application and re-entry.

While many herbicides do not specify re-entry periods, some herbicide actives are quite toxic, so care should still be taken to read label safety requirements and avoid contact with wet spray deposits.

For older products that don't specify a re-entry period, avoid re-entering the field for at least 24 hours after treatment, particularly for products with a signal heading of 'Caution', 'Poison' or 'Dangerous Poison'. Newer products will usually state a re-entry period; either 'Do not allow entry into treated areas until the spray has dried' or a specific number of hours – often 0, 12, or 24 (a notable exception is profentofos, which has a re-entry period of 34 days); if no re-entry period is listed, wait until after the spray has dried before entering the field (see Table 31 on page 135).

If the field needs to be entered prior to the stated re-entry period or before the spray is dry, limit the duration of entry and wear the PPE specified on the label. Even after the re-entry period has been observed, some protective clothing may be advisable. If not specified, appropriate PPE should be determined using a risk assessment.

## Withholding period

A withholding period (WHP) is the minimum period from when a pesticide is applied to when the treated area is allowed to be grazed, cut for fodder or harvested. This period is to prevent the presence of inappropriate levels of chemical residues in the harvested crop or from developing in stock that graze the stubble or are fed by-products of the treated crop.

The APVMA sets maximum residue limits (MRLs) for agricultural and veterinary chemicals in agricultural produce, particularly produce entering the Australian food chain (such as cotton seed). MRLs are unlikely to be exceeded as long as the products are used in accordance with approved label instructions (see Table 31). Residue limits for exported products are set by the importing country.

Label WHPs assume label compliance and effective application techniques, including even coverage with well-maintained equipment. Excessive residues may still occur due to a range of factors, including mixing issues, inappropriate application volumes, worn nozzles, or an overlap in sprayed areas.

Crop residues **MUST NOT** be fed to stock if any product that specifically prohibits grazing or cutting for stock food has been applied to that crop. Crops treated with a product that has a no grazing WHP specified should also not be grazed either before or after harvest. Contact the relevant chemical manufacturer for advice on managing chemical residues in stock.

**ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS.**

## Handling & storage

A critical part of responsible use of pesticides is their safe storage, transport and handling, as well as appropriate disposal of product that is no longer wanted or able to be used. Storing, handling and applying pesticides correctly greatly reduces any potential negative impacts to you, your staff, your business, your neighbours and the environment.

Many registered pesticides are classified as hazardous chemicals, or pose some degree of health risk to those who use them or are exposed to them. Workplace health and safety regulations exist to protect workers from both short and long term health effects of exposure to hazardous chemicals. *myBMP* provides guidance and resources to meet legal requirements for handling, storage and application of agrochemicals. It also includes templates to help document farm-specific procedures to minimise harm and respond in the event of an injury, fire, or spill. These procedures must be communicated to all staff.

Ensure anyone handling or applying pesticides has the appropriate training/accreditations, and maintain a collection of easily accessible copies of Safety Data Sheets (SDS) for all products used.

## Safe disposal

### Disposing of empty containers

Empty chemical containers present a risk to people and the environment. A critical part of responsible pesticide use is their safe transport, storage, and handling. Correct handling of pesticides greatly reduces potential negative impacts to you, your staff, your business, your neighbours and the environment.

Clean empty containers according to the steps below and store securely until they can be recycled. Containers marked 'returnable' should be returned to the supplier. Many 'non-returnable' metal and plastic containers used for farm chemicals can be recycled via Agsafe's national *drumMUSTER* program (look for the logo on the container).

Containers presented at a *drumMUSTER* receive site will be inspected to ensure that there is no visible product residue (including powder, flake, coloured/dark or clear fluid) on the inside or outside of the container. Some stains (including dirt, dust or mould) are acceptable.

- Triple rinse or pressure rinse your containers immediately after use (residues are more difficult to remove when dry). Pour the rinse water back into the spray tank.
- Thoroughly clean the container thread and outside surfaces with a hose into the spray tank. Rinse all caps separately in a bucket of clean water, and pour the rinsate from the caps into the spray tank.
- Inspect the container, particularly the thread and screw neck to ensure all chemical residues have been removed.
- Use a steel rod or crowbar to puncture metal containers. Pass it through the neck/pouring opening and out the base of the container.
- Allow the containers to drain completely and air dry.
- Store capless cleaned containers in a sheltered place to keep them clean and dry until they can be delivered to a *drumMUSTER* collection centre.

Intermediate Bulk Containers (IBCs) are not recycled by the *drumMUSTER* program, however a quick and easy Agsafe guide is available on how to send IBCs back for recycling or reuse.

[www.drummuster.org.au/container-recycling/the-abcs-for-your-ibcs](http://www.drummuster.org.au/container-recycling/the-abcs-for-your-ibcs)

## Disposing of unwanted chemicals

Unwanted rural chemicals may result from discontinued use due to changes in cropping or animal practices, development of newer/more effective/safer chemicals, or changes in a chemical's registration and/or banning from use. They may also be due to unknown product, property sale; inherited product or deceased estate. Any unwanted or unknown chemicals held on farm are potential hazards to people, the environment and the community, and must be disposed of responsibly and lawfully.

ChemClear is an industry stewardship program that collects currently registered agricultural and veterinary chemicals at the end of their life cycle, or when they become surplus. The program aims to meet disposal requirements of agvet chemical users and prevent potential hazardous chemicals from being dumped in landfills, creeks or otherwise inappropriately disposed of.

1. Take an inventory of any unwanted rural chemicals, including all identifiable container features (e.g. label, manufacturer, expiry date, container size and the remaining quantity left in the container). An inventory form can be obtained from the ChemClear website.
2. Register the inventory for the next collection in your area. Book on freecall 1800 008 182; fax 03 9371 8501 or at [www.chemclear.org.au](http://www.chemclear.org.au)
3. Continue to store your registered chemicals safely and securely. ChemClear will contact you direct to advise the location for retrieval.
4. Deliver chemicals ensuring that transportation is safe (never place chemicals in the boot of a car or back of a station wagon). Refer to the ChemClear website for more information.

Group 1 chemicals (currently registered, within 2 years of expiry/ deregistration, or containers displaying the *drumMUSTER* logo) are collected free of charge under the program. Group 2 chemicals (no longer registered, do not display the *drumMUSTER* logo, unknown, unlabelled, out of date, or mixed product) attract a fee for disposal, although the ChemClear

program covers the transport costs. This fee is payable directly to the contractor after a quote has been accepted by the holder of the chemicals. While not guaranteed, individual state EPA funding may be available from time to time to subsidise disposal of Group 2 chemicals.

### Further information:

AgSafe Limited 02 6206 6888 [www.agsafe.org.au](http://www.agsafe.org.au)

*drumMUSTER* program 1800 008 707 [www.drummuster.org.au](http://www.drummuster.org.au) or contact your local representative ([www.drummuster.org.au/our-people/your-local-regional-consultant](http://www.drummuster.org.au/our-people/your-local-regional-consultant))

ChemClear program 1800 008 182 [www.chemclear.org.au](http://www.chemclear.org.au)



A step by step guide to effective and economic waste management

Proper chemical container rinsing can save you money. It's simple.

Uncleaned containers with left over residue could contain up to **\$10 worth** of product.

*drumMUSTER* suggests the following steps to prevent you pouring money down the drain:

1.  Rinse out your *drumMUSTER* containers immediately after using them.
2.  Pour the rinse water back into the spray tank and ensure the container is clean throughout.
3.  Remove lids and puncture metal containers.
4.  Allow drums to dry.
5.  Check drums again for any left over residue and rinse if residue remains.
6.  Return them to your nearest *drumMUSTER* site. This can be found at [www.drummuster.org.au](http://www.drummuster.org.au)

# Pesticide Application Record Sheet



Industry &  
Investment

## Location, Applicator, Date of Application

|   |              |        |   |       |        |   |              |   |  |   |  |  |  |  |
|---|--------------|--------|---|-------|--------|---|--------------|---|--|---|--|--|--|--|
| Property/Holding: (residential address)   |              |        |   | Date: |        |   |              |   |  |   |  |  |  |  |
| Applicator's Full Name:   |              |        | Owner (if not applicator):                                      |       |        |   |              |   |  |   |  |  |  |  |
| Address:  |              |        | Address:  |       |        |   |              |   |  |   |  |  |  |  |
|   |              | Phone: |   |       | Phone: |   |              |   |  |   |  |  |  |  |
| Mobile:   | Fax:         | Email: | Mobile:   | Fax:  | Email: |   |              |   |  |   |  |  |  |  |
| Sensitive Areas (including distances, buffers):   |              |        | Comments (including risk control measures for sensitive areas): |       |        |   |              |   |  |   |  |  |  |  |
| <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td></td> <td>N</td> <td></td> </tr> <tr> <td>W</td> <td>Treated Area</td> <td>E</td> </tr> <tr> <td></td> <td>S</td> <td></td> </tr> </table> |              |        |   | N     |        | W | Treated Area | E |  | S |  |  |  |  |
|   | N            |        |   |       |        |   |              |   |  |   |  |  |  |  |
| W   | Treated Area | E      |   |       |        |   |              |   |  |   |  |  |  |  |
|   | S            |        |   |       |        |   |              |   |  |   |  |  |  |  |

## Host/Pest

|                       |   |                            |
|-----------------------|---|----------------------------|
| Paddock Number/Name:  | Paddock Area:   | Order of Paddocks Sprayed: |
| Crop/Situation:       | Type of Animals:  |                            |
| Crop/Pasture Variety: | Age/Growth Stage:   |                            |
| Growth Stage:         | Mob/Paddock/Shed:   |                            |
| Pest/Disease/Weed:    | Animals — Number Treated:   |                            |
|                       | Pest Density/Incidence: Heavy <input type="checkbox"/> Medium <input type="checkbox"/> Light <input type="checkbox"/> |                            |

## Application Data

|                          |                                    |                    |                         |           |
|--------------------------|------------------------------------|--------------------|-------------------------|-----------|
| Full Label Product Name: |                                    | Rate/Dose:         | Water Rate L/ha:        |           |
| Permit No.:              | Expiry Date:                       | Additives/Wetters: |                         |           |
| Total L or kg:           | WHP:                               | ESI*:              | Date Suitable for Sale: |           |
| Equipment Type:          |                                    | Nozzle Type:       | Nozzle Angle:           | Pressure: |
| Date Last Calibrated:    | Water Quality (pH or description): |                    |                         |           |

## Weather

|  |                |                       |            |           |                            |
|--|----------------|-----------------------|------------|-----------|----------------------------|
| Showers <input type="checkbox"/> Overcast <input type="checkbox"/> Light Cloud <input type="checkbox"/> Clear Sky <input type="checkbox"/> |                |                       |            |           |                            |
| Rainfall (24 hours before and after)   |                |                       |            |           |                            |
| Before:  | mm             | During:               | mm         | After:    | mm                         |
| Time (show time in this column)  | Temperature °C | Relative Humidity (%) | Wind Speed | Direction | Variability (e.g. gusting) |
| Start  |                |                       |            |           |                            |
|  |                |                       |            |           |                            |
| Finish   |                |                       |            |           |                            |
| Comments:  |                |                       |            |           |                            |

\* When using herbicides in mixtures with fungicides and insecticides, an ESI may apply to the non-herbicide component of the mixture.



**TABLE 31: Commonly used pesticides**

| Active                               | MoA group | Re-entry | Withholding (days)  |        |
|--------------------------------------|-----------|----------|---------------------|--------|
|                                      |           |          | Harvest             | Stock* |
| Abamectin                            | 6A        | dry      | 20                  | 20     |
| Acetamiprid                          | 4A        | dry      | 10                  | NG     |
| Afidopyropen                         | ??        | dry      | 7                   | —      |
| Alpha-cypermethrin                   | 3A        | 24 hrs   | 14                  | —      |
| Amitraz                              | 19        | dry      | 21                  | —      |
| Amorphous silica                     | —         | dry      | 0                   | 0      |
| Bacillus thuringiensis               | 11A       | N/A      | 0                   | 0      |
| Bifenthrin                           | 3A        | dry      | 14                  | NG     |
| Beta-cypermethrin                    | 3A        | —        | 14                  | —      |
| Chlorantraniliprole                  | 28        | N/A      | 28                  | NG     |
| Chlorantraniliprole/<br>Thiamethoxam | 28/4A     | dry      | 28                  | NG     |
| Chlorfenapyr                         | 13        | 12 hrs   | 28                  | NG     |
| Chlorpyrifos                         | 1B        | dry      | 28                  | 28     |
| Chlorpyrifos/<br>bifenthrin          | 1B/3A     | dry      | 28                  | NG     |
| Chlorpyrifos-methyl                  | 1B        | 24 hrs   | 28                  | NG     |
| <i>Clitoria ternatea</i><br>extract  | —         | dry      | 0                   | —      |
| Clothianidin                         | 4A        | —        | 5                   | NG     |
| Cyantraniliprole                     | 28        | dry      | 14                  | NG     |
| Cypermethrin                         | 3A        | 12 hrs   | 14                  | —      |
| Deltamethrin                         | 3A        | dry      | 7                   | NG     |
| Diafenthiuron                        | 12A       | 24 hrs   | 14                  | NG     |
| Dimethoate                           | 1B        | dry      | 14                  | NG     |
| Dinotefuran                          | 4A        | dry      | 14                  | NG     |
| Emamectin benzoate                   | 6A        | 12 hrs   | 28                  | NG     |
| Esfenvalerate                        | 3A        | dry      | 7                   | —      |
| Etoxazole                            | 10B       | dry      | 21                  | NG     |
| Fipronil                             | 2B        | dry      | 28                  | NG     |
| Gamma-cyhalothrin                    | 3A        | dry      | 21                  | —      |
| <i>Helicoverpa</i> NPV               | —         | dry      | 0                   | 0      |
| Imidacloprid                         | 4A        | 12 hrs   | 91                  | NG     |
| Indoxacarb                           | 22        | dry      | 28                  | NG     |
| Lambda-cyhalothrin                   | 3A        | dry      | 21                  | —      |
| Magnet®                              | —         | 24 hrs   | See insecticide mix |        |
| Methidathion                         | 1B        | 2 days   | 3                   | —      |
| Methomyl                             | 1A        | 24 hrs   | 0                   | NG     |
| Paraffinic oil                       | —         | —        | 1                   | —      |
| Pirimicarb                           | 1A        | dry      | 21                  | 21     |
| Phorate                              | 1B        | 48 hrs   | 70                  | 70     |
| Profenofos                           | 1B        | 34 days  | 28                  | —      |
| Propargite                           | 12C       | dry      | 28                  | NG     |
| Pymetrozine                          | 9B        | dry      | 28                  | NG     |
| Pyriproxyfen                         | 7C        | 24 hrs   | 28                  | NG     |
| Quintozene                           | 14        | dry      | N/A                 | NG     |
| Spinetoram                           | 5         | dry      | 28                  | NG     |
| Spirotetramat                        | 23        | dry      | 21                  | NG     |
| Sulfoxaflor                          | 4C        | dry      | 14                  | NG     |
| Thiamethoxam                         | 4A        | dry      | 28                  | NG     |
| Thiodicarb                           | 1A        | 24 hrs   | 21                  | 21     |

| Active  | MoA group            | Re-entry | Withholding (days) |        |
|---|----------------------|----------|--------------------|--------|
|   |                      |          | Harvest            | Stock* |
| <i>Clitoria ternatea</i><br>extract             | —                    | dry      | 0                  | —      |
| Mancozeb  | M3                   | dry      | 14                 | NG     |
| Metalaxyl-M                                     | 4                    | —        | N/A                | NG     |
| Prothiconazole/<br>Bixafen (PERMIT)             | 3/7                  | dry      | N/A                | 35     |
| Tebuconazole<br>(PERMIT)                        | 3                    | dry      | 42                 | NG     |
| Tolclofos-methyl                                | 14                   | dry      | N/A                | N/A    |
| Carfentrazone-ethyl                             | dessicant (G)        | dry      | 1                  | NG     |
| Diquat  | dessicant (L)        | dry      | N/A                | 1 day  |
| Ethephon  | conditioner          | 4 hrs    | 14                 | NG     |
| Ethephon + AMADS                                | conditioner          | —        | 1                  | NG     |
| Mepiquat  | PGR                  | dry      | 28                 | NG     |
| Paraffinic oil & other<br>non-ionic surfactants | adjuvant             | —        | 1                  | —      |
| Paraquat + diquat                               | dessicant (L)        | dry      | 7                  | 1-7    |
| Petroleum oil + guar<br>gum derivative**        | adjuvant             | —        | —                  | —      |
| Pyraflufen-ethyl                                | defoliant (G)        | dry      | 7                  | 7      |
| Sodium chlorate                                 | dessicant            | dry      | 0                  | NG     |
| Thidiazuron                                     | defoliant            | dry      | N/A                | NG     |
| Thidiazuron + diuron                            | defoliant<br>(- + C) | —        | N/A                | NG     |

dry = until spray has dried.  
 NG = no grazing.  
 — = not specified or unknown.  
 N/A = not required when used as directed.  
 \*Refer to label for export market requirements.  
 \*\* polygalactomannan hydroxy propyl ether.  
 Refer to the Weeds chapter for herbicide registrations, and page 108 for a more comprehensive list of registered fungicides.  
 Note: This information is provided as a guide only. The post-spray periods given may not be the same for all products with that active ingredient.  
**ALWAYS CHECK THE LABEL.**

# Pesticides and the environment

**T**he cotton industry's guidelines for minimising risk to the environment are a component of the *myBMP* program.

Most insecticides are toxic to aquatic organisms, bees, birds and other fauna. Fungicides and herbicides are relatively safe to bees in terms of their active ingredients, but their carriers and surfactants may be toxic. The risk that a particular product poses to the environment (native terrestrial and aquatic plants and animals) are reflected in statements on the label under headings like 'Protecting wildlife, fish, crustaceans and the environment'.

Pesticide residues may accumulate in animals treated with any pesticides or fed any crop product, including crop residues, which have been sprayed with pesticides. Animals that test positive for chemical residues (i.e. with readings which exceed maximum residue limits) at slaughter will be rejected.

Pesticide residues may also contaminate other plant products for human use and consumption. Observe harvest withholding periods on the pesticide label and do not assume that in the absence of a withholding period or after the expiry of a withholding period that plant products will be free of residues.

## Protecting the aquatic environment

Aquatic fauna and flora are often highly sensitive to agrochemicals, as are many other species that utilise aquatic areas as part of their habitat. All spray operations should consider the risk of contaminating nearby standing water.

- Prevent drift onto surface waters during application;
- Locate mixing/loading and decontamination facilities well away and use bunding and sumps to prevent movement of either concentrate or rinsate into surface waters;
- Install valves that prevent back-flow in equipment used to fill spray tanks from surface waters and in suction lines for chemigation systems that draw directly from surface waters;
- Avoid aerial application of spray on fields during irrigation;
- Build sufficient on-farm storage capacity (including provision for storm runoff) to contain pesticide-contaminated tail water from irrigation;
- Spray in an upstream direction (when spraying near water is necessary), to reduce the maximum concentration at any one point in the watercourse;
- Use only registered products to control aquatic weeds, e.g. Roundup Bioactive® rather than Roundup®; and,
- Avoid disposal of used containers in surface waters and on flood plains and river catchments.

## Protecting birds

Broad-spectrum insecticides can be particularly toxic to birds, especially in granular formulations, and insecticidal seed dressings can pose similar risks. Spillages can be very hazardous as birds can easily ingest a toxic dose from a small area.

- Ensure complete incorporation beneath the soil, particularly at row ends where spillage may occur; and,
- Immediately clean up spillage, however small.

Bait materials for control of rodents (not registered in cotton, but relevant to rotation crops) or soil insect pests can also be hazardous to birds, either through direct consumption of the bait or from feeding on bait-affected animals or pests.

- Ensure even bait distribution, with no locally high concentrations;
- Don't bait over bare ground or in more open situations, such as near crop perimeters, where birds may see the baits;
- Don't bait near bird habitat such as remnant native vegetation;
- Use bait stations to prevent bird access, particularly near bird habitat;
- Only bait where pest pressure is high;
- Bait late in the evening when birds have finished feeding; and,
- Promptly collect and bury rodent carcasses in open situations.

Foliar-applied insecticide sprays can also be hazardous to birds, either because of direct contact with the sprayed chemical, or by feeding on sprayed insect pests or crops. Even where birds are not killed, they may be more vulnerable to predation. Contaminated seed and insects collected from sprayed fields by parent birds can also be lethal to young chicks still in the nest.

- Minimise drift into remnant vegetation, wildlife corridors, nesting sites, or other bird habitats; and,
- Actively discourage birds from feeding in crops that are to be sprayed.

Pesticides can also indirectly impact on bird populations through the loss of plants and animals on which they feed and through loss of habitat.

- Protect sensitive areas such as remnant vegetation, riparian areas and waterways from spray drift;
- Where possible use target-specific as opposed to broad-spectrum pesticides (that are more likely to impact on non-target organisms and plants in the environment) and adopt an Integrated Pest Management (IPM) approach to controlling pests;
- Spray late in the day when birds have finished feeding; and,
- Use only low toxicity chemicals when large concentrations of birds are nesting nearby.



Whistling ducks. (Anne Palfreyman)



**Protect aquatic environments from pesticides; many aquatic birds such as egrets contribute towards natural pest control.**  
(Stacey Vogel, CottonInfo)

### Protecting other fauna

Sodium fluoroacetate (1080) baiting has been an essential tool for protecting and recovering threatened species, especially mammals threatened by foxes. There are non-target risks associated with almost every poison, including 1080. However based on years of scientific research the APVMA concluded that although individual poisoning of non-target animals can occur this does not adversely affect the overall population of non-target wildlife. For more information on the impacts of 1080 baits on wildlife visit the PestSmart website [pestsmart.org.au/domestic-animal-safety/the-facts-of-1080-baiting](http://pestsmart.org.au/domestic-animal-safety/the-facts-of-1080-baiting)

- Tie or bury baits where applicable to limit removal by non-target species;
- Do not lay baits at times or locations where native wildlife are likely to be harmed;
- Place baits and/or design bait stations so that wildlife access is minimised; and,
- Regularly check and replace baits to reduce excess baiting and to be more effective.

### Protecting remnant native vegetation

Remnant native vegetation, i.e. any native patches of trees, shrubs or grasses that still remain in the landscape, can be damaged by herbicide and defoliant poisoning either via leaching through the soil or absorption through the leaves.

- Prevent drift by implementing best practices for spray application;
- Install natural or artificial barriers to intercept spray drift;
- Adhere to product label no-spray zone instructions;
- Select target-specific herbicides;
- Select non-soil active herbicides when remnant vegetation is nearby; and,
- Use an Integrated Weed Management (IWM) approach when managing environmental weeds.

The best way to manage any long term adverse environmental risk is to follow the protection statements on labels, minimise spray drift, and to dispose of chemical containers and waste in accordance with label directions and codes of practice.



**Top arrow – vegetative barrier.**  
**Bottom arrow – remnant vegetation.**

### Cotton Landcare Innovations project

**Under the National Landcare Program's Smart Farming Partnerships initiative, CRDC secured a \$1.3 million grant to bring Cotton Landcare Tech-Innovations 2021 onto Australian cotton farms to enhance natural resources and biodiversity.**

**The project consists of four key activities: biodiversity, innovation, technology, and collaboration, and builds on international best practice to implement and develop cutting-edge technologies, such as drone mapping and aerial seeding, acoustic monitoring and big data, to help Australian cotton better report on and improve on-farm biodiversity.**

**For more information, visit [www.crdc.com.au/cotton-landcare-tech-innovations](http://www.crdc.com.au/cotton-landcare-tech-innovations)**

# Protecting pollinators

**A**lthough cotton is commonly regarded as being largely self-fertile and self-pollinating, bees provide an important service to agriculture generally. Native stingless bees are resistant to diseases and parasites common in honeybees (including varroa mite) and therefore provide an ‘insurance policy’ for pollination. Many solitary wild insect pollinators are also present on and around farms but are less easily seen because they do not have the obvious hives of managed honeybees and social native bees. Remnant vegetation and other bushland on the farm should be thought of as pollinator habitat and protected accordingly.

As with many other crops, cotton can be a high-risk environment for bees as they are particularly susceptible to many insecticides.

Contamination can occur through:

- Direct contact with spray or spray drift;
- Subsequent contact while foraging; and,
- Carrying of contaminated material back to the hive.

Insecticide contamination can result in individual bee death, diminished productivity or even destruction of the whole hive. Insecticides that are particularly toxic to bees (including abamectin, fipronil, indoxacarb and pyrethroids) are identified with a special statement on the label such as:

**Dangerous to bees. DO NOT spray any plants in flower while bees are foraging.**

The cotton IRMS highlights insecticides with a label warning about bee safety. The relative toxicity of cotton insecticides to honeybees is listed in Table 4 on pages 14-15.

The residual toxicity (the amount of time the product remains toxic to bees after the time of application), should also be considered. Many insecticides are safe to bees once the spray deposits have dried on the crop. The label provides information about any residual activity on bees and what action must be taken to avoid impacting bees (note label recommendations may vary between different formulations containing the same active ingredient). Refer to the APVMA PubCRIS database to search for information about individual insecticides ([portal.apvma.gov.au/pubcris](http://portal.apvma.gov.au/pubcris)).

**ALWAYS READ AND FOLLOW LABEL INSTRUCTIONS.**

## Bee ecology

Honeybee field activity is temperature and food-related. The bees become more active above 12-13°C with maximum flight activity reached at 18°C. With water gatherers, the main flight activity occurs when temperatures are above the mid-thirties. Honeybees usually forage within 2-4 km of their hive, although they will fly long distances (7-10 km) in search of good food.

Native bee species often have much shorter flight ranges; as little as 500 m for some of the smaller species.

Bees collect nectar from extra-floral nectaries (e.g. under leaves) as well as from cotton flowers so they may be foraging in cotton crops before, during and after flowering. They may also visit weeds in flower in and around fields. Damage can also occur to bees if pesticides drift over hives, water surfaces or neighbouring vegetation being foraged by bees.

Coolibah trees (*Eucalyptus microtheca*), black box (*E. largiflorens*) and river red gums (*E. camaldulensis*) are a primary source of nectar and pollen for honeybees. These trees grow on the black soil plains along many of the river courses in cotton growing areas, with flowering occurring in response

to good spring rains. In northern NSW trees begin to flower mid to late December, finishing about the end of January. Flowering times may vary by a few weeks in the southern and central areas of Queensland. When heavy flowering is expected, beekeepers may move large numbers of hives into these regions.



**Cotton can be a high-risk environment for bees.** (Susan Maas, CRDC)

## Protect bees when using fipronil

While bees are susceptible to many insecticides used on cotton, fipronil risk to managed hives has been highlighted due to the extended residue risk. The industry IRMS recommends *fipronil should not be applied to flowering crops*. Refer to label for statement about bees.

**Example of fipronil label statement:**

**“Dangerous to bees. DO NOT apply where bees from managed hives are known to be foraging, and crops, weeds or cover crops are in flower at the time of spraying, or are expected to flower within 28 days (7 days for pastures and sorghum).**

**Before spraying, notify beekeepers to move hives to a safe location with an untreated source of nectar, if there is any potential for managed bees to be affected by the spray or spray drift. If an area has been sprayed inadvertently, in which the crop, weeds or cover crop were in flower or subsequently came into flower, notify beekeepers in order to keep managed bees out of the area for at least 28 days (7 days for pastures and sorghum) from the time of spraying. Where the owner of managed hives in the vicinity of a crop to be sprayed is not known, contact your State Department of Primary Industries/Agriculture, citing the hive registration number, for assistance in contacting the owner.”**

## Reducing the risk

- Follow the registered label directions.
- Register on BeeConnected (see details below).
- If beehives are in the vicinity of crops, notify apiarists at least 48 hours before spraying, to allow time for removal of the hives.
- Inform contract pesticide applicators of apiary locations.
- Pay particular attention to wind speed and direction, air temperature and time of day before applying pesticides.
- Use buffer zones to reduce the impact of spray drift or overspray on non-target crops and native vegetation.
- Avoid contamination of surface waters where bees may drink.
- Use an IPM framework and only apply pesticides when necessary.
- Favour pesticides that have lower toxicity to bees and other beneficial insects (see Table 4).
- Where possible, use EC or granular formulations that dissolve completely in preference to wettable powders or microencapsulated formulations as bees can transport the particles back to the hive.

With good communication and good will, it is possible for apiarists and cotton growers to work together to minimise risks to bees, as both honey (including pollination services) and cotton are important agricultural industries.



## BeeConnected app

Communication between growers and bee keepers is critical in reducing the risk of unintended exposure of bees to products that may have the potential to negatively impact bee health. BeeConnected is a nation-wide, user-driven smart-phone app and website that enables collaboration between beekeepers, farmers and spray contractors to facilitate best-practice pollinator protection. Growers log the location of their properties and beekeepers can log the present or future locations of their beehives through a Google Maps-based platform with GPS capability. When a beehive is logged near a farmer's property, both users are sent automated notifications and can chat further about their activities via a secure internal messaging service. CropLife Australia provide BeeConnected to the community free of charge as part of their Pollinator Protection Initiative. The effectiveness of the BeeConnected system is greatest when as many growers and beekeepers as possible use the service. It is important to keep an eye out for bees or apiaries near your property that might not be logged in the system.

For more information and to participate in this great service go to: [www.beeconnected.org.au](http://www.beeconnected.org.au)

## Other information sources

### Insecticide-treated seed

CropLife Australia's Seed Treatment Stewardship Strategy outlines measures to reduce risks from dust generated during handling and planting of treated seed, and provides guidance on minimising off-target movement of crop protection products to protect important pollinators. More information is available at [www.croplife.org.au/resources/programs/seed-treatment-stewardship-strategy](http://www.croplife.org.au/resources/programs/seed-treatment-stewardship-strategy)



## BeeAware

BeeAware is a hub of information for beekeepers and growers about honeybee biosecurity and pollination of agricultural and horticultural crops. The site contains information about the pollination of crops and how beekeepers and growers can work together to provide and receive best practice pollination services.

BeeAware was funded by the Australian honeybee industry, pollinator-reliant plant industries, Plant Health Australia, governments and R&D agencies ([beeaware.org.au](http://beeaware.org.au)).

## Native bees

More detailed information on native beekeeping is available in the Australian Native Bee Book. [nativebeebook.com.au](http://nativebeebook.com.au)

## Australian cotton industry sustainability reporting: Bees

The environmental toxic load (ETL) is an indicator to assess the average amount of toxic pressure on human health and the environment from pesticides applied on one hectare. ETL for bees has been chosen as the target indicator for pesticides applied in the cotton industry, as bees are a critical provider of pollination services and a key indicator for terrestrial ecosystem health. The lower the ETL score, the lower the toxic pressure.

The Australian cotton industry's ETL for bees decreased by 18.2% (from 11 to 9) in the four years to 2018, continuing a long-term, although slowing, industry trend. It is important that the industry continues to practice IPM, and where possible avoid use of products that are highly toxic to bees such as fipronil, neonicotinoids (e.g. clothianidin), abamectin, chlorpyrifos and bifenthrin. The industry's voluntary IRMS now includes a specific recommendation on fipronil use (see page 54).

CottonInfo video on bees in cotton: [youtu.be/LBRUaqZaVoQ](https://youtu.be/LBRUaqZaVoQ)



**For further information about protecting bees or to notify the owner of beehives not logged in to BeeConnected, take a note of the identification/registration number and contact your state agricultural department or apiarist/beekeeping association:**

### New South Wales

NSW DPI Beekeeper registrations 02 6552 3000

NSW Apiarists' Association

[www.nswaa.com.au](http://www.nswaa.com.au) E: [info@nswaa.com.au](mailto:info@nswaa.com.au)

Apiaries on public land in NSW are displayed on the BPASS map ([bpass.dpi.nsw.gov.au/s/view-sites-public](http://bpass.dpi.nsw.gov.au/s/view-sites-public))

### Queensland

Biosecurity Queensland 13 25 23

Queensland Beekeepers' Association

[www.qbabees.org.au](http://www.qbabees.org.au) E: [qbainc@bigpond.com](mailto:qbainc@bigpond.com)

### Western Australia

DPIRD Apiary and biosecurity officers

E: [PBHoney@dpird.wa.gov.au](mailto:PBHoney@dpird.wa.gov.au)

Bee Industry Council of Western Australia

[www.bicwa.com.au](http://www.bicwa.com.au) E: [info@bicwa.com.au](mailto:info@bicwa.com.au)

### Northern Territory

Plant Biosecurity Branch (08) 8999 2036

### Victoria

Beekeeper registration helpline 1800 356 761

Victorian Apiarists' Association

[www.vicbeekeepers.com.au](http://www.vicbeekeepers.com.au) E: [vaa@vicbeekeepers.com.au](mailto:vaa@vicbeekeepers.com.au)

# Glossary & acronyms

- ACPM.** *Australian Cotton Production Manual* (available from CottonInfo).
- Action threshold.** See threshold.
- Adjuvant.** A substance (e.g. wetting agent, sticker, thickener, buffering agent) added to a spray mixture to enhance performance or overcome an inhibiting factor.
- AMF** (formerly VAM). Arbuscular mycorrhiza fungi symbiotically colonise plant roots, transferring extra nutrients, especially phosphorus, from the soil in exchange for photosynthates from the plant.
- Aphid colony.** Four or more aphids within 2 cm<sup>2</sup> on a leaf or terminal.
- APVMA.** Agricultural Pesticides and Veterinary Medicines Authority.
- Area wide management (AWM).** Growers working together in a region to manage pest populations.
- At-planting insecticide.** Insecticides applied as a granule or spray in the seed furrow with the seed during planting.
- BCI.** Better Cotton Initiative.
- Beat sheet.** Large sheet of yellow or white canvas, placed in the furrow and extended over the adjacent row. Plants are beaten against the sheet with a stick, dislodging insects which are quickly counted.
- Beneficials.** Biological agents that are beneficial to the crop, including natural enemies and pollinators.
- Beneficial to pest ratio.** Ratio used to incorporate predatory insect activity into pest management decisions.
- Biological insecticides.** Insecticides based on or derived from living organisms (e.g. NPV or Bt).
- Bollgard cotton.** Genetically modified cotton containing Bt insecticidal proteins.
- Box mapping.** See plant mapping.
- Broad-spectrum.** Insecticides that kill a wide range of insects, including both pest and beneficial species. Their use, particularly early in the season, often leads to pest resurgence or flaring.
- Bt.** Toxin derived from *Bacillus thuringiensis* that is toxic to many caterpillar species.
- Buffer zone.** An area where pesticide application does not occur between the application site and an identified sensitive area which is downwind from the application site. For boom and aerial spraying, it is measured from the edge of the sprayer swath closest to the downwind sensitive area.
- Calendar sprays.** Application on a timed schedule, regardless of pest density or the actual need for pest control.
- Come Clean. Go Clean.** Industry slogan promoting thorough cleaning to prevent transfer of pests between fields and farms.
- Conditioner.** Product used to hasten boll opening.
- Consecutive checks.** Successive insect checks taken from the same field or management unit.
- Conventional cotton.** Cotton varieties that are not genetically modified (in this guide the term is usually used to indicate absence of genes that produce Bt-derived insecticidal proteins).
- Cotton bunchy top (CBT).** Cotton virus vectored by aphids.
- Crop compensation.** Plant's capacity to 'catch-up' after insect damage without affecting yield or maturity.
- Cut-out.** Assimilate (photosynthesis products) demand exceeds supply and no more harvestable fruit is set.
- Damage threshold.** Level of damage leading to economic loss of yield or delay in maturity. Usually used in conjunction with pest thresholds to account for both pest numbers and plant growth. See also threshold.
- Damping off.** Wilting, collapse and death of seedlings, usually due to infection by pathogens.
- Day degree (DD).** Unit combining temperature and time used to monitor and compare crop development.
- Defoliation.** Leaf removal in preparation for harvest by artificially enhancing the natural process of senescence and abscission.
- Dessicant.** Harvest aid chemical that damages the leaf membrane, causing moisture loss and desiccation.
- Diapause.** Period of physiologically controlled dormancy in insects (e.g. at the pupal stage in the soil for *Helicoverpa armigera*).
- Double knock.** Sequential application of two pesticides with different modes of action in a short timeframe.
- DST.** Decision support tool used with consecutive silverleaf whitefly sampling to consider management options based on population pathways.
- D-vac.** Small portable suction sampler or blower/vacuum machine used to suck insects from the cotton plants into a fine mesh bag.
- Economic threshold.** See threshold.
- Efficacy.** Effectiveness of a product or natural enemy against pests.
- Egg parasitoids.** Parasitoid species that specifically target insect eggs, killing the developing host.
- Eretmocerus.** Parasitoid wasp of SLW that occurs naturally in most cotton regions or can be purchased for commercial release.
- Food sprays.** Natural food products applied to attract beneficials, particularly predators, into crops to help control pests. Yeast-based sprays attract; sugar-based sprays retain beneficials already present.
- F rank.** Ranking of each cotton variety in accordance with its resistance to Fusarium wilt.
- Frass.** Insect faeces is an indicator of insect activity, and sometimes helpful for identification.
- Fruit retention.** Percentage of fruit (squares or bolls) retained compared with the overall number produced.
- GVB.** Green vegetable bug.
- Herbicide Resistance Management Strategy (HRMS).** Voluntary industry strategy designed to delay herbicide resistance.
- Honeydew.** Sticky sugar-rich waste excreted by sap-sucking insects that can interfere with photosynthesis, promote sooty mould, affect lint quality and cause problems with fibre processing.
- In-furrow insecticide.** See at-planting insecticide.
- Insect growth regulator (IGR).** Compound that influences insect growth and development.
- Insecticide Resistance Management Strategy (IRMS).** Voluntary industry strategy that imposes restrictions on pesticide use to help prevent the development of insecticide resistance.
- Instar.** Juvenile insect development stages between moults.
- Integrated disease management (IDM).** The integration of appropriate disease control strategies with management of the whole farming system.

- Integrated pest management (IPM).** Using knowledge of pest biology, behaviour and ecology to implement a range of tactics in an integrated way to suppress or reduce pest populations.
- Integrated weed management (IWM).** Strategy to plan and implement a range of different tactics to achieve sustainable weed management.
- Lay-by herbicide.** Residual herbicide used to control weeds during crop growth.
- Larva.** Immature 'grub' stage of insects that undergo complete metamorphosis (e.g. moths, beetles, flies, lacewings). See also pupa.
- Lesion.** A localised, defined area of diseased tissue.
- Lock.** Segment of cotton boll.
- Management unit.** Farm area managed in the same way for the purposes of sampling and insect management.
- Maximum residue limit (MRL).** Maximum amount of pesticide residue legally permitted on food products, based on levels considered toxicologically acceptable.
- Mepiquat chloride.** Plant growth regulator used to prevent excessive growth or node elongation.
- Mode of action (MoA).** Pesticide classification scheme based on target site that aims to prevent the development of cross-resistance to related pesticide compounds.
- Mycorrhizae.** See AMF.
- Natural enemies.** Predators, parasitoids and pathogens of pests.
- Natural mortality.** Expected death rate of insects in the field due to environmental factors (including natural enemies).
- No-spray zone.** Mandatory unsprayed distance between sprayed and downwind sensitive areas. See also buffer zone.
- NPV (nucleopolyhedrovirus).** An entomopathogenic virus. Commercial formulations targeting *Helicoverpa* are available in Australia.
- Nursery.** Crop or natural habitat that attracts and sustains pests or beneficials through multiple generations.
- Nymph.** Immature insect stage that looks like the adult but without wings (e.g. bugs, thrips, mites). Nymphs go through a series of moults and do not pass a pupal stage.
- Organophosphates (OPs).** Broad-spectrum insecticides classified into MoA Group 1B.
- OTT.** 'Over the top' application of herbicides.
- PAMP.** Pesticide application management plan.
- Parasitoid.** Insect that lays its eggs on or in another arthropod; the larva develops within the host, killing it on emergence.
- Pathogen.** Microorganism that causes disease (e.g. Fusarium wilt is the disease caused by a soil-inhabiting fungus *Fusarium oxysporum* f.sp. *vasinfectum* (Fov)).
- Peak flowering.** Period where the plant has the highest numbers of flowers opening per day.
- Pest flaring.** Increase in a pest population (usually in species with very fast life cycles) following a pesticide application targeted at another species.
- Pest resurgence.** Increase in a targeted pest population following a pesticide application. Usually due to a reduction in beneficial populations.
- Pest threshold.** Pest population at which a control measure is needed to prevent eventual economic loss to the crop. See also threshold and damage threshold.
- Pix.** See mepiquat chloride.
- Plant growth regulator (PGR).** Chemical that can be applied to reduce growth rate and prevent rank growth.
- Plant mapping.** Recording the fruiting dynamics of a cotton plant to help interpret the effects of pest damage or environmental factors.
- Planting window.** Restriction on planting time used to restrict the number of pest generations exposed to particular control measures.
- Post-emergent knockdown herbicide.** Herbicide used to rapidly control weeds after they emerge.
- PPE.** Personal protective equipment.
- Pre-plant knockdown herbicide.** Herbicide used to rapidly control weeds prior to planting.
- Presence/absence.** Noting presence/absence of a numerous pest rather than counting absolute numbers.
- Prophylactic sprays.** Insecticides applied in anticipation of a potential pest problem. These 'insurance' sprays are usually not recommended due to increased costs, selection for insecticide resistance, and the risk of secondary pest outbreaks.
- PSO (petroleum spray oil).** Petroleum-derived oil used to control insect pests. Can also be used to deter egg lay of some pests.
- PubCRIS (Public Chemicals Registration Information System).** APVMA's searchable database of agrochemicals registered for use in Australia including products with off-label permits.
- Pupae.** Stage where larva undergo transformation into an adult (e.g. from caterpillar to moth).
- Pupae busting.** Tillage to reduce the survival of overwintering pupae.
- Rank growth.** Excessive vegetative growth; usually managed using mepiquat chloride.
- Ratoon cotton.** Regrowth from the previous season's rootstock.
- Re-entry period.** Mandatory period between chemical application and re-entry to the treated area.
- Refuge.** Area left untreated with an insecticide or technology that generates unexposed susceptible individuals to act as a dilution factor when mating with resistant adults.
- Resistance.** When individual pests develop a mechanism for dealing with a particular pesticide or mode of action group, potentially resulting in poor control (less than expected) or control failures (much less than expected).
- Resistance management plan (RMP).** Proactive plan to mitigate the risks of resistance developing to any of the proteins contained in Bt cotton. Compliance is required under the Bt cotton Technology User Agreement.
- Retention.** See fruit retention.
- Rotation crops.** Other crop types grown before or after the cotton crop.
- Scouting.** Checking crops (for insects, damage, weeds, growth etc). Often refers to the gathering of data only; interpretation and management recommendations are provided by crop agronomists/consultants.
- Secondary pests.** Pests that do not usually become a problem unless their natural enemies (predators or parasitoids) are reduced in number by insecticides. See also pest flaring.
- Seed treatment.** Insecticide/fungicide coating on cotton seeds that offers protection during germination and establishment.

**Selection pressure.** Pesticide sprays remove susceptible individuals. More selection events from a particular chemical group leads to greater 'pressure' or chance of selecting a resistant population.

**Shedding.** Abortion/loss of reproductive structures (squares, flowers or bolls) caused by environmental factors or pest damage.

**Siphunculi.** Pair of wax-secreting tubular organs on the abdomen of aphids (also called cornicles).

**SLW.** Silverleaf whitefly.

**'Soft' approach.** Managing insect and mite pests using options that have minimal impact on beneficial populations.

**Square.** Cotton flower bud.

**Stacked genes.** Two or more genes that express similar activity (e.g. Bollgard 3), or different activities in the same variety (e.g. Bollgard + Roundup Ready®).

**Sucking pests.** Bugs with piercing mouthparts (e.g. mirids, aphids, whitefly).

**Sweep net.** Large cloth net with long handle used to sample insects in the upper canopy.

**Synthetic insecticides.** Non-biological insecticides (artificial versions of natural insecticides or synthetic molecules with insecticidal/miticidal activity).

**Systemic pesticide.** Pesticide that is taken up through plant tissue (as opposed to remaining on the surface) and transported through the plant's vascular system.

**Terminal.** Uppermost growing point, particularly on the main stem.

**Threshold.** The point at which an action is taken. Action thresholds are based on a level/number of pests/damage. Economic thresholds consider the commodity value and treatment cost. See also pest threshold and damage threshold.

**TIMS.** Transgenic & Insect Management Strategy (Committee).

**Tippling.** Loss of the terminal, causing the plant to develop multiple stems.

**Trap crop.** Small area of more attractive host plants that concentrates a pest population to allow easier management.

**Tubercles.** Small knob-like or rounded protuberances between the antennae of some insects.

**Vegetative barrier.** Deliberately planted vegetation strips designed to protect adjacent sensitive areas (remnant vegetation, waterways, other crops) by capturing airborne spray droplets.

**Visual sampling.** Counting insects in the field without the use of additional equipment such as a beat sheet.

**Volunteer cotton.** Plants that have germinated and established unintentionally in-field or elsewhere (roadsides, fencelines etc).

**V rank.** Ranking of each cotton variety in accordance with its resistance to Verticillium wilt.

**Withholding period.** Minimum period allowed between application of an agrochemical and harvest or grazing by stock.

## Coordinating authors for this edition

Insects: Paul Grundy, Qld DAF & CottonInfo

Weeds: Eric Koetz, NSW DPI & CottonInfo

Volunteers & ratoon cotton: Sharna Holman, Qld DAF & CottonInfo

Diseases: Beth Shakeshaft, NSW DPI & CottonInfo

Biosecurity: Sharna Holman, Qld DAF & CottonInfo

PGRs & harvest aids: Tonia Grundy, Qld DAF

Sprays & application: Susan Maas, CRDC

## Contributors to current and/or previous editions

Stephen Allen

Carly Ambler

Fiona Anderson

Katie Aspin

Jon Baird

Michael Bange

Bartley Bauer

Donald Benn

Graham Betts

Lisa Bird

Sally Ceeney

Graham Charles

Greg Constable

Nicola Cottee

Sharon Downes

Vesna Gagic

Bill Gordon

Stuart Gordon

Tim Green

Aphrika Gregson

Paul Grundy

Annabelle Guest

Simone Heimoana

Grant Herron

Andrew Hewitt

Sharna Holman

Jamie Hopkinson

Peter Hughes

Phil Hurst

Dinesh Kafle

Jenene Kidston

Karen Kirkby

Kristen Knight

Eric Koetz

Kate Langfield

Duy Le

Tracey Leven

Rob Long

Doug McCollum

Susan Maas

Robert Mensah

Melina Miles

Geoff Naylor

Mary O'Brien

Ruth Redfern

Linda Scheikowski

Richard Sequeira

Murray Sharman

Brad Siebert

Linda Smith

Tanya Smith

Frank Taylor

Ian Taylor

David Thornby

Phil Tucker

Rene van der Sluijs

Stacey Vogel

Jeff Werth

Mary Whitehouse

Sandra Williams

Lewis Wilson

Bayer Crop Science

CottonInfo Regional Extension Officers



# Index

## A

- Alternaria leaf spot 96-97, **99**, 108, 109-113
- Annual ryegrass **80**, 81, 82, 84, 87, 90
- Anthraco-nose boll rot 101
- Aphids 6, 8, 9, 12, 14-15, **18-22**, 102-103
  - As vectors 18, 97, 102, 119
  - Cotton aphid 6, 15, 18-19, 22, 102, 119, 120
  - Cowpea aphid 18-19
  - Green peach aphid 18-19
  - Honeydew 6, 18, 22, 47
  - Resistance 5, 19, 22, 26, 35, 54-60
- Apple dimpling bug 9, 10, 14-15, 28, 33, 38, 48
- APVMA (Australian Pesticide and Veterinary Medicines Authority) **130**, 137
- Area wide management 6, 8
- Assassin bugs 9, 12, 15, 28, 41
- Awnless barnyard grass **80**, 81, 82, 83, 87, 90 – see also barnyard grass

## B

- Bacterial blight 96, 107, 109, **120**
- Barnyard grass 81, 82, 83, 84
- Bees 5-6, 8, 9, 14-15, 54, 56-57, 136, **138-139**
- Beneficial (insects) 5-6, 8, **9-15**, 16, 54-58, 66, 103, 139 – see also content within each key pest
  - Inundative release 62
- Beneficial (microbes) 97, 107
- Beneficial to pest ratio 8, 27, **28**, 66
- Big-eyed bug 9, 10, 14-15, 19, 28, 35, 38, 48
- Biofumigation 97, 100
- Biosecurity 74, 96-97, 104, 105, 107, 109, **114-120**, 131, 139
- Birds 8, 32, **136-137**
- Black root rot 96-97, **100**, 108, 109-113
- Blue disease (exotic) 109, **119**, 120
- Bollgard 6, 18, 24, 27-28, 34, 52, 58, **61-70**, 71-72, 91, 94
- Boll rots 27, 52, 96-97, **100-102**, 109-113
- Boll weevil (exotic) 118
- Bollworms – see cotton bollworm, pink bollworm, pink spotted bollworm, rough bollworm and *Helicoverpa*
- Brown marmorated stink bug (exotic) 26, 118
- Bt resistance 28, 61-70, 71, 120
- Buffer zone 69, 71, 78, 123, 134, 139

## C

- Capeweed 19, 81, 82, 83
- ChemClear 133
- Come Clean. Go Clean. 8, 31, 32, 74, 76, 86, 94, 97, 100, 102, 104, 105, 107, **114-115**
- Cotton bollworm (see also *Helicoverpa*)
- Cotton bunchy top (CBT) 5, 18, 73, 97, **102-103**, 109, 119
- Cotton leaf curl disease (exotic) 109, 119
- Cotton stainer (exotic) 118
- Cricket (black field) 44
- Cutworm 44

## D

- Damsel bug 9, 10, 14-15, 19, 28, 35, 38
- Decontamination 114-115, 124, 135-136
- Defoliation 8, 18, 32, 38, 48, 64, 69, 70, 72, 78, 94, 97, 102, 106, 108, **122**
- Dessicants 122, 135
- Diapause 28, 29, 50, 52, 64, 70
- Diplodia boll rot 101
- Drift – see spray drift
- drumMUSTER* 133

## E

- Earwig 28, 32, 42
- Encarsia* 47, 48
- Eretmocerus* 6, 14-15, 47, 48
- Exotic pests 26, 104, 109, 114, **118-120**

## F

- Fall armyworm **24**, 54, 56-57, 120
- False codling moth (exotic) 118
- Feathertop Rhodes grass **80**, 81, 82, 87
- Fleabane 81, 84, 87
- Flaxleaf flebane **80**, 81, 82, 87
- Food sprays 62
- Fruiting factor 27
- Fruit retention 8, 9, 25, 27, 33-35, 54, 89, 122
- Fungicides 96, 99, 106, 108, 119, 136
- Fusarium 106, 108, 109
  - Fusarium boll rot 101
  - Fusarium wilt 96-97, **103-104**, 105, 106, 107, 108, 109-113, 120

## G

- Glossy shield bug 9, 10, 15, 28
- Green vegetable bug **25-26**, 33, 41, 101

## H

- Harlequin bug 25
- Harvest aids 16, 72, **122**
- Hazardous chemicals 123, 131, 132-133, 136
- Helicoverpa* 9, 14-15, 22, 24, **27-30**, 33, 35, 38, 40, 41, 52, 54-60, 61-70, 71-72, 92, 120, 135
- Herbicide Resistance Management Strategy (HRMS) 73, 77, 81-84, **86-87** – see also herbicides (resistance)
- Herbicides 6, 69, 73-78, 80, 81-88, 89-91, 92-94, 97, 106, 122, 124, 132, 137
  - Plant-back periods 74-78, 86-87, 88
  - Resistance – 73-77, 80, 81-84, 88, 89-91, 92, 96-97, 103, 120 – see also HRMS
- Honeydew 6, 18-19, 22, 31, 47-48
- Host plant resistance 100, 108, 120

## I

- Insect growth regulator (IGR) 49, 58
- Insecticides
  - Post-spray periods 135
  - Resistance 5, 6, 8, 9, 16, 54-60, 61-70, 71 – see also IRMS, RMP and individual sections for key insects
  - Toxicity 14-15, 136, 138-139 – see also hazardous chemicals and safety
- Insecticide Resistance Management Strategy (IRMS) 6, 8, 9, 16, 19, 28, 35, 50, **54-60**, 66, 138-139
- Integrated Disease Management (IDM) **96-97** – see also IDM tactics for key diseases
- Integrated Pest Management (IPM) 5-8, 9, 16, 18, 35, 56-57, 58, 119, 136, 139
- Integrated Weed Management (IWM) **73-78**, 81, 86, 89-90, 92, 137
- Inversion (weather) 123, 124, 128

## J

- Jassids – see leafhoppers

## L

- Lacewings 9-10, 14-15, 19, 28, 31, 32, 38, 46, 48
- Ladybird beetles 6, 9, 10, 14-15, 19, 28, 31-32, 38, 46, 48
- Leafhoppers 52
- Liverseed grass **80**, 81, 87, 90
- Lygus bugs (exotic) 119

## M

- Marshmallow weed 19, 32, 38, 102
- Mealybug (Solenopsis) 5-6, 9, **31-32**, 34, 54, 56-57, 92, 110
- Minute two-spotted ladybird beetle 14-15, 38
- Minute pirate bug 9, 12, 15, 38, 46
- Mirids 9, 12, 14-15, 19, 25, 32, **33-36**, 38, 41, 51, 56-57, 60, 101
- Mites 5-6, 8, 9, 12, 14-15, 16, 18, 30, 33, 35, **37-40**, 45, 54, 56-57, 58, 60, 138
- myBMP* 4, 98, 123, 124, 126, 127, 131, 132, 136

## N

Nematode – see Reniform nematode  
No-spray zone 40, 46, 94, 123, 131, 137

## O

Overwintering 5, 8, 16, 28, 32, 52, 56-57, 61, 92, 102 –  
see also diapause and overwintering content within each key pest

## P

Pale cotton stainers 25, 33, **41**, 101, 118  
Parasitoids 6, 9, 14-15, 16, 19, 25, 27, 28, 32, 47, 48  
Parrot beaked boll 33  
Pesticide application management plan (PAMP) 8, 123  
Petroleum spray oil (PSO) 14, 122  
Pigeon pea 29, 35, 62-64, 68-70, **71-72**, 78, 94, 104  
Pigweed 32, 102, 107 – see also red pigweed  
Pink bollworm 52  
Pink spotted bollworm 52  
Plant growth regulators (PGR) 96, 122, 135  
Planting window 62-64, 70  
Pollinators 138, 139  
Predators 6, 9-12, 14-15, 16, 19, 25, 28, 31, 32, 33, 34, 35, 38, 40, 41, 42,  
45-46, 48  
Pupae busting 8, 28, 32, 54, 56-57, 58, 62-64, 69-70, 72, 73, 94  
Pythium 100, 106, 108, 109-111

## R

Ratoon cotton 5, 8, 18-19, 31, 41, 62, 68-69, 71-72, 73, 76, 78, **92-94**, 97, 99,  
100, 102-103, 104, 107  
Red and blue beetle 9, 10, 14-15, 19, 28, 32  
Red pigweed 80, 90  
Redshouldered leaf beetle 52  
Re-entry period 132, 135  
Refuges 8, 28, 58, 62-66, 68-70, 71-72, 114  
Reniform nematode **104**, 109  
Reoccurring wilt **104-105**, 109-113  
Resistance – see Herbicides (Resistance), HRMS, IRMS and RMP  
Resistance management plan (RMP) 28, 58, 61-70, 71-72  
Retention (stubble) 42 – see also fruit retention  
*Rhizoctonia* 100, 106, 108, 109-113  
Rough bollworm 52  
Roundup Ready Flex 74, 77, 81, 88, **89-91**, 92  
Rutherglen bug 52

## S

Safety  
Crop 77-78, 86, 89  
Human and Animal 16, 122, 123, 130-133  
Safflower 29, 35, 38, 78, 101, 104, 107  
Sampling (disease) 96, 97, 104, 105, 106, **107**, 112, 120  
Sampling (insects) 5-6, 8, 9, 16, 60, 66, 103, 120 –  
see also content within each key pest  
Sampling (weeds) **84**, 90  
Sclerotinia 96, 100-101, 108  
Seedling disease 97, 99, 100, 101, 102, 103, **106**, 109-113  
Seed rots **100-102**, 106, 110-113  
Seed treatments 12, 19, 42, 45-46, 56-57, 60, 96, 100, 106, 108, 139  
Shield bugs 9, 10, 15, 25, 28, 35  
Silverleaf whitefly (SLW) – see whitefly  
Soil pests 8, 42-44  
Solenopsis mealybug – see mealybug  
Sooty mould 18, 31, 47  
Sowthistle 19, 32, 38, 50, 51, **80**, 81, 83, 87, 90  
Spray drift 8, 16, 62, 69-70, 71, 74, 78, 122, 123-124, 127, 130-131, 136-137,  
138-139  
Spray window 49, 50, 54, 56-60, 89, 92  
Stink bugs 25-26, 35, 118  
Symphyla 42-44

## T

Texas root rot (exotic) 109, **119**  
Thistles 19, 26, 35, 107 – see also sowthistle  
Thresholds 5-6, 8, 9, 16, 54, 56-60, 66, 73, 96 –  
see also content within each key pest  
Thrips 6, 9, 12, 14-15, 26, 38, **45-46**, 56-57, 97  
Tight lock 25, 41, 100, 101, 110-113  
TIMS Committee 50, 54, 60, 62, 81  
Tobacco streak virus 97, 107, 109  
Trap crops 8, 64, 68-70, 72  
*Trichogramma* 14-15, 27-28

## V

Vectors 96-97, 119, 120  
Verticillium wilt 96-97, 103, **106-107**, 108, 109-113, 120  
Volunteer cotton 5, 8, 18-19, 31-32, 41, 50-51, 54, 62-64, 68-70, 71-72, 73,  
76, 78, 89, 90, **92-94**, 97, 99, 100, 102-103, 104, 107, 114

## W

Whitefly  
Exotic *Bemisia* 120  
Greenhouse 15, 47, 50  
Silverleaf (SLW) 4, 5-6, 9, 14-15, 33, 34, 35, **47-51**, 54, 56-57, 60  
Whitegrubs 44  
Wild radish 26, 81, 83  
Windmill grass 81, 82, 87  
Wireworm 12, **42-44**  
Withholding periods 48, **132**, 136

## ADVERTISERS

|                           |               |
|---------------------------|---------------|
| Ag Leader .....           | 55, 59        |
| Adama .....               | 63, 65        |
| Agsafe .....              | 67            |
| Bayer .....               | 2             |
| CCA .....                 | 43            |
| Corteva Agriscience ..... | 7, 17, Insert |
| CGS .....                 | 1             |
| CottonInfo Team .....     | IBC           |
| eChem .....               | 117, 121      |
| FMC .....                 | 93, 95        |
| GrowthAg .....            | 53            |
| Innovate Ag .....         | 21            |
| John Deere .....          | 125, 129      |
| Louis Dreyfus .....       | 91            |
| Sumitomo .....            | IFC, OBC      |
| UPL .....                 | 11, 13        |
| Valley/Valmont .....      | 23            |
| Vicchem .....             | 79, 85        |



# Meet our team

Led by CottonInfo Program Manager Warwick Waters (0437 937 074, [warwick.waters@crdc.com.au](mailto:warwick.waters@crdc.com.au)), and supported by Communications Lead Brad Pfeffer ([brad.pfeffer@crdc.com.au](mailto:brad.pfeffer@crdc.com.au)) the CottonInfo team of Regional Extension Officers, Technical Leads & myBMP experts are all here to help!

## Regional Extension Officers

Regional Extension Officers provide cotton research outcomes and information directly to growers, agronomists, consultants and agribusinesses in each region. Contact your local Regional Extension Officer for the latest research, trials and events in your area.

| Elsie Hudson   | Amanda Thomas   | Andrew McKay   |
|--|---|--|
| <i>Namoi</i><br>P: 0458 142 777<br>E: <a href="mailto:elsie.hudson@cottoninfo.net.au">elsie.hudson@cottoninfo.net.au</a>           | <i>Macquarie and Bourke</i><br>P: 0417 226 411<br>E: <a href="mailto:amanda.thomas@cottoninfo.net.au">amanda.thomas@cottoninfo.net.au</a> | <i>Border Rivers, St George, Dirranbandi</i><br>P: 0407 992 495<br>E: <a href="mailto:andrew.mckay@cottoninfo.net.au">andrew.mckay@cottoninfo.net.au</a> |
| Annabel Twine  | Kieran O'Keeffe   | Janelle Montgomery   |
| <i>Darling Downs</i><br>P: 0447 176 007<br>E: <a href="mailto:annabel.twine@cottoninfo.net.au">annabel.twine@cottoninfo.net.au</a> | <i>Southern NSW</i><br>P: 0427 207 406<br>E: <a href="mailto:kieran.okeeffe@cottoninfo.net.au">kieran.okeeffe@cottoninfo.net.au</a>       | <i>Gwydir and Mungindi</i><br>P: 0428 640 990<br>E: <a href="mailto:janelle.montgomery@cottoninfo.net.au">janelle.montgomery@cottoninfo.net.au</a>       |

## Technical Leads

Technical leads are experts in their fields and provide in-depth analysis, information and research to the industry, for the benefit of all growers. Contact the technical leads to learn more about water use efficiency, nutrition, soil health and much, much more.

| Eric Koetz  | Jon Welsh  | Paul Grundy   |
|---|--|---|
| <i>Weed Management</i><br>P: 0413 256 132<br>E: <a href="mailto:eric.koetz@dpi.nsw.gov.au">eric.koetz@dpi.nsw.gov.au</a>                                | <i>Energy and Climate</i><br>P: 0458 215 335<br>E: <a href="mailto:jon@agecon.com.au">jon@agecon.com.au</a>                | <i>Integrated Pest Management</i><br>P: 0427 929 172<br>E: <a href="mailto:paul.grundy@daf.qld.gov.au">paul.grundy@daf.qld.gov.au</a> |
| Stacey Vogel  | Sharna Holman  | René van der Sluijs   |
| <i>Natural Resources and Catchments</i><br>P: 0428 266 712<br>E: <a href="mailto:staceyvogel.consulting@gmail.com">staceyvogel.consulting@gmail.com</a> | <i>Biosecurity</i><br>P: 0477 394 116<br>E: <a href="mailto:sharna.holman@daf.qld.gov.au">sharna.holman@daf.qld.gov.au</a> | <i>Fibre Quality</i><br>P: 0408 885 211<br>E: <a href="mailto:sluijs@optusnet.com.au">sluijs@optusnet.com.au</a>                      |

| Beth Shakeshaft   | Ben Crawley   | Oliver Knox  | Jon Baird  |
|---|---|--|--|
| <i>Disease Management</i><br>P: 0409 477 303<br>E: <a href="mailto:beth.shakeshaft@dpi.nsw.gov.au">beth.shakeshaft@dpi.nsw.gov.au</a> | <i>Irrigation</i><br>P: 0439 247 605<br>E: <a href="mailto:ben.crawley@dpi.nsw.gov.au">ben.crawley@dpi.nsw.gov.au</a> | <i>Soil Health</i><br>P: 0490 045 326<br>E: <a href="mailto:oknox@une.edu.au">oknox@une.edu.au</a> | <i>Nutrition</i><br>P: 0429 136 581<br>E: <a href="mailto:jon.baird@dpi.nsw.gov.au">jon.baird@dpi.nsw.gov.au</a> |

## myBMP team

The myBMP team run the industry's best management practice program, myBMP. Contact the myBMP team to learn more about - or to participate in - myBMP.

| Nicole Scott   | Polly Quinn   |
|--|---|
| <i>myBMP Customer Service Officer</i><br>P: 1800cotton (1800 268 866)<br>E: <a href="mailto:nicoles@cotton.org.au">nicoles@cotton.org.au</a> | <i>myBMP Lead Auditor</i><br>P: 0418 385 656<br>E: <a href="mailto:pollygibbons@gmail.com">pollygibbons@gmail.com</a> |



For the 2021/22 season, cotton growers may be eligible for a rebate on Valor sprayed in the Roundup Ready FLEX® cotton system under the Roundup Ready PLUS® program.

Visit [www.roundupreadyplus.com.au](http://www.roundupreadyplus.com.au) for full terms and conditions.

## RESIDUAL AND BURNDOWN CONTROL WITH FLEXIBILITY

- 6-8 weeks residual control for late fallow and residual carryover control into the emerged crop.
- Enhanced knockdown control when used at spike rates with non-selective herbicides.
- Excellent control of Roundup Ready cotton volunteers.
- Registered for Lay-by application in emerged cotton.
- Short re-cropping intervals to a wide range of summer crops, including cotton.
- Strength against difficult to control weeds including feathertop Rhodes grass and fleabane.



[www.sumitomo-chem.com.au](http://www.sumitomo-chem.com.au)



Scan here to see more information about Valor 500WG Herbicide

© Valor is the registered trademark of Sumitomo Chemical Company, Japan. Roundup Ready, Roundup Ready PLUS and Roundup Ready FLEX are registered trademarks of the Bayer Group.