

## Venerid Bibliography

References for the "Generic Names" database as well as other selected works on the family Veneridae

Accorsi Benini, C. (1974). I fossili di Case Soghe - M. Lungo (Colli Berici, Vicenza); II, Lamellibranchi. *Memorie Geopaleontologiche dell'Università de Ferrara* 3(1): 61-80.

Adachi, K. (1979). Seasonal changes of the protein level in the adductor muscle of the clam, *Tapes philippinarum* (Adams and Reeve) with reference to the reproductive seasons. *Comparative Biochemistry and Physiology* 64A(1): 85-89

Adams, A. (1864). On some new genera and species of Mollusca from the seas of China and Japan. *Annals and Magazine of Natural History* 13(3): 307-310.

Adams, C. B. (1845). Specierum novarum conchyliorum, in Jamaica repertorum, synopsis. Pars I. *Proceedings of the Boston Society of Natural History* 12: 1-10.

Ahn, I.-Y., G. Lopez, R. E. Malouf (1993). Effects of the gem clam *Gemma gemma* on early post-settlement emigration, growth and survival of the hard clam *Mercenaria mercenaria*. *Marine Ecology -- Progress Series* 99(1/2): 61-70 (2 Sept.)

Ahn, I.-Y., R. E. Malouf, G. Lopez (1993). Enhanced larval settlement of the hard clam *Mercenaria mercenaria* by the gem clam *Gemma gemma*. *Marine Ecology -- Progress Series* 99(1/2): 51-59

Alemany, J. A. (1986). Estudio comparado de la microestructura de la concha y el enrollamiento espiral en *V. decussata* (L. 1758) y *V. rhomboides* (Pennant, 1777) (Bivalvia: Veneridae). *Bollettino Malacologico* 22(5-8): 139-152.

Alemany, J. A. (1987). Estudio comparado de la microestructura de la concha y el enrollamiento espiral en *Dosinia exoleta* (L. 1758) y *Dosinia lupinus* (Poli, 1791), (Bivalvia: Veneridae). *Bollettino Malacologico* 23(5-8): 223-236.

Alemany, J. A. (1987). Microestructura de la concha de *Venerupis aurea* Gmelin, 1791; Revision microestructural de la subfamilia Tapetinae, Bivalvia; Veneridae. *Boletin de la Real Sociedad Espanola de Historia Natural*, Sección Geológica 83(1-4): 15-24.

Ali Zade, A. A. and A. Aleskerov Dzh (1974). Novyye dannyye o sostave sarmatsko fauny Azerbaydzhan. *Doklady - Akademiya Nauk Azerbaydzhanskoy SSR = Me'ruzeler - Azerbaydzhan SSR Elmler Akademiyasy* 30(12): 31-34.

- Amano, K. (1998). First discovery of fossil *Humilaria* (Bivalvia: Veneridae) from Japan. *Venus, The Japanese Journal of Malacology* 57(4): 271-280.
- Anderson, G. J., M. B. Miller, K. K. Chew. (1982). A guide to Manila clam aquaculture in Puget Sound, Washington. Washington, University of Washington, *Sea Grant Program, Publications WSG-82-4*: 45 pp.
- Anderson, R. C. (1986). Resistance to predation of a clam *Humilaria kennerleyi* (Reeve, 1863) [abst.]. *Annual Report of the Western Society of Malacologists* 18: 7
- Anderson, R. C. (1994). *Humilaria kennerleyi* survives drill attack. *The Festivus* 25(8): 95
- Andres, I. (1983). Los Veneridae, Bivalvia, en el Plioceno de Bonares. *Studia Geologica, Universidad de Salamanca* 19: 101-131.
- Anonymous (1922). Cold weather kills Pismo clams. *California Fish and Game* 8(8): 124-125
- Ansell, A. D. (1961). Reproduction, growth and mortality of *Venus striatula* (da Costa) in Kames Bay, Millport. *Journal of the Marine Biological Association of the United Kingdom* 41(1): 191-215
- Ansell, A. D. (1961). The functional morphology of the British species of Veneracea (Eulamellibranchia). *Journal of the Marine Biological Association of the United Kingdom* 41(2): 489-515
- Ansell, A. D. (1961). The development of the primary gonad in *Venus striatula* (da Costa). *Proceedings of the Malacological Society of London* 34(5): 243-247
- Ansell, A. D. (1962). The functional morphology of the larva, and the post-larval development of *Venus striatula* (da Costa). *Journal of the Marine Biological Association of the United Kingdom* 42(2): 419-443
- Ansell, A. D. (1962). Observations on burrowing in the Veneridae (Eulamellibranchia). *Biological Bulletin* 123(3): 521-530
- Ansell, A. D. (1964). Some parameters of growth of mature *Venus mercenaria* L. Conseil Permanent International pour l'Exploration de la Mer, *Journal du Conseil* 29(2): 214-220
- Ansell, A. D. (1967). Egg production of *Mercenaria mercenaria*. *Limnology and Oceanography* 12(1): 172-176

Ansell, A. D. (1968). The rate of growth of the hard clam *Mercenaria mercenaria* (L.) throughout the geographical range. Conseil Permanent International pour l'Exploration de la Mer, *Journal du Conseil* 31(3): 364-409

Ansell, A. D. and K. F. Lander (1967). Studies on the hard-shell clam, *Venus mercenaria*, in British waters. III. Further observations on the seasonal biochemical cycle and on spawning. *Journal of Applied Ecology* 4(?): 425-435

Ansell, A. D. and F. A. Loosmore (1963). Preliminary observations on the relationship between growth, spawning and condition in experimental colonies of *Venus mercenaria* L. Conseil Permanent International pour l'Exploration de la Mer, *Journal du Conseil* 28(2): 285-293

Ansell, A. D., F. A. Loosmore, K. F. Lander (1964). Studies on the hard-shell clam, *Venus mercenaria*, in British waters. I. Growth and reproduction in natural and experimental colonies. *Journal of Applied Ecology* 1: 63-82

Ansell, A. D., F. A. Loosmore, K. F. Lander (1964). Studies on the hard-shell clam, *Venus mercenaria*, in British waters. II. Seasonal cycle in condition and biochemical composition. *Journal of Applied Ecology* 1: 83-95

Ansell, A. D. and E. R. Trueman (1967). Burrowing in *Mercenaria mercenaria* (L.). *Journal of Experimental Biology* 46(1): 105-115

Aplin, J. A. (1947). Pismo clams of San Quintin, Lower California. *California Fish and Game* 33(1): 31-33

Aplin, J. A. (1947). Pismo clam increase. *California Fish and Game* 33(3): 129-131

Arakawa, K. Y. (1986). Bibliography on Japanese necked clam *Ruditapes philippinarum* (Adams & Reeve, 1850) with annotations. *Hiroshima Shell Club*. 45 pp.

Arneri, E., G. Giannetti, B. Antolini (1998). Age determination and growth of *Venus verrucosa* L. (Bivalvia: Veneridae) in the southern Adriatic and the Aegean Sea. *Fisheries Research* 38(2): 193-198.

Asson-Batres, M. A. (1986). The feeding behavior of the juvenile Dungeness crab, *Cancer magister* Dana, on the bivalve, *Transennella tantilla* (Gould), and a determination of its daily consumption rate. *California Fish and Game* 72(3): 144-152

Asson-Batres, M. A. (1988). Reproduction and growth of the brooding bivalve *Transennella tantilla*. *The Veliger* 30(3): 257-266

Baxter, J. L. (1961). Results of the 1955 to 1959 Pismo clam census. *California Fish and Game* 47(2): 153-162

Baxter, J. L. (1962). The Pismo clam in 1960. *California Fish and Game* 48(1): 35-37

Beentjes, M. P. and B. G. Williams (1986). Endogenous circatidal rhythmicity in the New Zealand cockle *Chione stutchburyi* (Bivalvia, Veneridae). *Marine Behaviour And Physiology* 12(3): 171-180.

Belding, D. L. (1912). A report upon the quahaug and oyster fisheries of Massachusetts. Massachusetts, *Report for Commissioners on Fisheries and Game* 1911: 1-134

Bello, G. (1986). Porcentajes teoricos de retencion de *Venus verrucosa* L. (Bivalvia: Veneridae) con cedazos de varillas. *Investigacion Pesquera* 50(2): 167-177.

Bemis, B. E. (1992). *Stable isotopic records of venerid bivalve shells: environmental information from the eastern Pacific Ocean and southern Caribbean Sea*. M.S. Thesis, University of Wisconsin, Madison: xi, 95 leaves.

Bemis, B. E. and D. H. Geary (1996). The usefulness of bivalve stable isotope profiles as environmental indicators: data from the eastern Pacific Ocean and the southern Caribbean Sea. *Palaios* 11(4): 328-339.

Berkeley, C. (1959). Some observations of *Cristispira* in the crystalline style of *Saxidomus giganteus* Deshayes and in that of some other Lamellibranchiata. *Canadian Journal of Zoology* 37(1): 53-58

Bernard, F. R. (1969). The parasitic copepod *Mytilicola orientalis* in British Columbia bivalves. Canada, *Journal of the Fisheries Research Board of Canada* 26(1): 190-191

Bernard, F. R. (1982). *Nutricola* n. gen. for *Transennella tantilla* (Gould) from the northeastern Pacific (Bivalvia: Veneridae). *Venus, The Japanese Journal of Malacology* 41(2): 146-149

Berry, S. S. (1907). Molluscan fauna of Monterey Bay, California. *The Nautilus* 21(2): 17-22.

Berta, A. (1976). An investigation of individual growth and possible relationships in a population of *Protothaca staminea* (Mollusca: Pelecypoda). California, University of California at Berkeley, Museum of Paleontology, *PaleoBios* 21: 26 pp.

Blainville (1828). *Venus*. *Dictionnaire des Sciences Naturelles* 57: 261-294.

Borsa, P., Y. Jousselin, B. Delay (1992). Relationships between allozymic heterozygosity, body size, and survival to natural anoxic stress in the palourde *Ruditapes decussatus* L. (Bivalvia: Veneridae). *Journal of Experimental Marine Biology and Ecology* 155(2): 169-181.

Borsa, P. and C. Thiriot Quievreux (1990). Karyological and allozymic characterization of *Ruditapes philippinarum*, *R. aureus* and *R. decussatus* (Bivalvia, Veneridae). *Aquaculture* 90(3-4): 209-227.

Borzone, C. A. (1989). El ciclo gonadal de *Venus antiqua* King & Broderip 1835 (Veneridae: Bivalvia) en el golfo San Jose. *Physis* (113A): 61-72.

Botton, M. L. (1984). Spacial distribution of three species of bivalves on an intertidal flat: the interaction of life-history strategy with predation and disturbance. *The Veliger* 26(4): 282-287

Bourne, N. F. (1971). The effects of temperature, salinity and food on the development of the larvae of butter clams (*Saxidomus giganteus*, Deshayes) [abst.]. *Proceedings of the National Shellfisheries Association* 61: 1

Bourne, N. F. (1982). Distribution, reproduction, and growth of Manila clam, *Tapes philippinarum* (Adams and Reeves [*sic!*]), in British Columbia. *Journal of Shellfish Research* 2(1): 47-54

Bourne, N. F. (1983). Clam predation by scoter ducks in the Strait of Georgia, British Columbia [abst.]. *Journal of Shellfish Research* 3(1): 84

Bourne, N. F. (1984). Pinnotherid crabs in butter clams: a commensal or a parasite [abst.]. *Journal of Shellfish Research* 4(1): 82

Bower, S. M. (1992). Winter mortalities and histopathology in Japanese littlenecks [*Tapes philippinarum* (A. Adams and Reeve, 1850)] in British Columbia due to freezing temperatures. *Journal of Shellfish Research* 11(2): 255-263

Bower, S. M., J. Blackhorn, Gary R. Meyer (1992). Parasite and symbiont fauna of Japanese littlenecks, *Tapes philippinarum* (Adams and Reeve, 1850), in British Columbia. *Journal of Shellfish Research* 11(1): 13-19

Bradley, W. H. and P. Cooke (1959). Living and ancient populations of the clam *Gemma gemma* in a Maine coast tidal flat. United States, Department of the Interior, Fish and Wildlife Service, *Fishery Bulletin* 58(137): ii + 305-334

Breese, W. P. and F. D. Phibbs (1970). Some observations on the spawning and early development of the butter clam, *Saxidomus giganteus* (Deshayes). *Proceedings of the National Shellfisheries Association* 60: 95-98

Broderip, W. J. and G. B. Sowerby, I. (1835). [... new species of shells contained in Mr. Cuming's collection]. *Proceedings of the Zoological Society of London*, for 1835[3](27): 41-47

Bryan, W. A. (1919). A Hawaii form of *Tapes philippinarum*. *The Nautilus* 32(4): 124-125

Calabrese, A. and H. C. Davis (1966). The pH tolerance of embryos and larvae of *Mercenaria mercenaria* and *Crassostrea virginica*. *Biological Bulletin* 131(3): 427-436

Canapa, A., I. Marota, F. Rollo, E. Olmo (1996). Phylogenetic analysis of Veneridae (Bivalvia): comparison of molecular and palaeontological data. *Journal of Molecular Evolution* 43(5): 517-522.

Canapa, A., I. Marota, F. Rollo, E. Olmo (1999). The Small-Subunit rRNA Gene Sequences of Venerids and the Phylogeny of Bivalvia. *Journal of Molecular Evolution* 48: 463-468.

Cantera, K. J. R. (1991). Shallow-water venerid clams (Bivalvia: Veneridae) from the Pacific coast of Colombia. *The Veliger* 34(1): 78-84.

Carballo, H. M. A. and R. Fisher (1978). La Formacion San Miguel (Miocene, Costa Rica). *Informe Semestral Instituto Geográfico Nacional*: 45-144

Carlisle, J. G. Jr. (1973). Results of the 1971 Pismo clam census. *California Fish and Game* 59(2): 138-139

Carriker, M. R. (1956). Biology and propagation of young hard clams, *Mercenaria mercenaria*. *Journal of the Elisha Mitchell Scientific Society* 72(1): 57-60

Carriker, M. R. (1961). Interrelation of functional morphology, behavior, and autecology in early stages of the bivalve *Mercenaria mercenaria*. *Journal of the Elisha Mitchell Scientific Society* 77(2): 168-241

Carter J. G., E. Barrera, M. J. S. Tevesz (1998). Thermal potentiation and mineralogical evolution in the Bivalvia (Mollusca). *Journal of Paleontology* 72(6): 991-1010.

Casey, R. F. G. S. (1952). Some genera and subgenera, mainly new, of mesozoic heterodont Lamellibranchs. *Proceedings of the Malacological Society* 29: 121-176.

Castagna, M. and J. N. Kraeuter (1981). Manual for growing the hard clam *Mercenaria*. College of William and Mary, Virginia Institute of Marine Science, *Sea Grant Program, Special Report in Applied Marine Science and Ocean Engineering* 249: 110 pp.

Cassab Rita de Cassia, T. (1984). Sobre a ocorrência de *Periglypta multicostata* na Formação Pirabas, Oligoceno-Mioceno do Brasil. *Anais da Academia Brasileira de Ciencias* 56(3): 283-286.

Castro Ortiz, J. L. and F. Garcia Dominguez (1993). Crecimiento de *Chione californiensis* (Broderip, 1835) (Veneridae) en la Ensenada de la Paz, Baja California Sur, Mexico. *Investigaciones Marinas Cicimar* 8(2): 45-52.

Castro Ortiz, J. L. and A. Tripp Quezada (1994). Crecimiento de la almeja chocolata roja *Megapitaria aurantiaca* (Bivalvia: Veneridae). *Investigaciones Marinas Cicimar* 9(2): 79-84.

Chanley, P. E. (1961). Inheritance of shell markings and growth in the hard clam, *Venus mercenaria*. *Proceedings of the National Shellfisheries Association* 50: 163-169

Cheney, D. P. (1971). A summary of invertebrate leucocyte morphology with emphasis on blood elements of the Manila clam, *Tapes semidecussata*. *Biological Bulletin* 140(3): 353-368

Chestnut, A. F. (1952). Growth rates and movements of hard clams, *Venus mercenaria*. *University of Miami, Marine Laboratory, Gulf and Caribbean Fisheries Institute, Proceedings* 4: 49-59

Chestnut, A. F., W. E. Fahy, H. J. Porter (1957). Growth of young *Venus mercenaria*, *Venus campechiensis*, and their hybrids. *Proceedings National Shellfisheries Association* 47: 50-56

Chew, K. K. (1989). Manila clam biology and fishery development in western North America. pp. 243-261, in: J. J. Manzi and Michael Castagna, *Clam culture in North America*. Amsterdam, etc. (Elsevier). x + 461 pp.

Chiba, K. and Y. Ohshima (1957). Effect of suspending particles on the pumping and feeding of marine bivalves, especially of Japanese neck-clam. *Bulletin of the Japanese Society of Scientific Fisheries* 23(7-8): 348-353

Choe, S. (1965). On the morphological variations and special features of the elongated and stunted forms in the short necked clam, *Tapes japonica*. *Korean Journal of Zoology* 8(1): 1-7

- Chryploff, G. (1957). Über die Makrofauna der Unterkreide aus der Tiefbohrung Werle 8. *Zeitschrift für Angewandte Geologie* 3(11-12): 500-505.
- Chung, E. Y., D. K. Ryou, J. H. Lee (1994). Gonadal development, age and growth of the shortnecked clam, *Ruditapes philippinarum* (Pelecypoda: Veneridae), on the coast of Kimje, Korea. *Korean Journal of Malacology* 10(1): 38-54.
- Clark, F. N. (1932). The present status of the Pismo clam. *California Fish and Game* 18(2): 170-180
- Clifton, H. E. and S. Boggs, Jr. (1970). Concave-up pelecypod (*Psephidia*) shells in shallow marine sand, Elk River beds, southwestern Oregon. *Journal of Sedimentary Petrology* 40(3): 888-897
- Coan, E. V. (1996). Designation of a neotype for the Pismo clam, *Tivela stultorum* (Mawe, 1823), and lectotypes for synonymous nominal taxa (Bivalvia: Veneridae). *The Veliger* 39(3): 267-269.
- Coates, H. (1925). Catalogue of the Conchological Collection in the Perthshire Natural History Museum. *Transactions and Proceedings of the Perthshire Society of Natural Science* 8(2): 17-95.
- Coe, W. R. (1947). Nutrition, growth and sexuality of the Pismo clam (*Tivela stultorum*). *Journal of Experimental Zoology* 104(1): 1-24
- Coe, W. R. (1948). Nutrition, environmental conditions, and growth of marine bivalve mollusks. *Journal of Marine Research* 7(3): 586-601
- Coe, W. R. (1953). Resurgent populations of littoral marine invertebrates and their dependence on ocean currents and tidal currents. *Ecology* 34(1): 225-229
- Coe, W. R. (1956). Fluctuations in populations of littoral marine invertebrates. *Journal of Marine Research* 15(3): 212-232
- Coe, W. R. and J. E. Fitch (1950). Population studies, local growth rates and reproduction of the Pismo clam (*Tivela stultorum*). *Journal of Marine Research* 9(3): 188-210
- Commito, J. A., C. A. Currier, L. R. Kane, K. A. Reinsel, I. M. Ulm (1995). Dispersal dynamics of the bivalve *Gemma gemma* in a patchy environment. *Ecological Monographs* 65(1): 1-20

- Conrad, T. A. (1837). Descriptions of new Marine Shells, from Upper California. *Journal of the Academy of Natural Sciences of Philadelphia* 7(1): 227-268.
- Conrad, T. A. (1856). Notes on shells, with descriptions of three recent and one fossil species. *Proceedings of the Academy of Natural Sciences of Philadelphia* 7: 31-32.
- Conrad, T. A. (1862). Descriptions of New Generea, Subgenera and Species of Tertiary and Recent Shells. *Proceedings of the Academy of Natural Sciences of Philadelphia* 14: 284-291.
- Conrad, T. A. (1862). Catalogue of the Miocene Shells of the Atlantic Slope. *Proceedings of the Academy of Natural Sciences of Philadelphia* 14: 559-586.
- Conrad, T. A. (1865). Catalogue of the eocene and oligocene Testacea of the United States. *American Journal of Conchology* 1(1): 1-35.
- Conrad, T. A. (1865). Descriptions of new eocene shells of the United States. *American Journal of Conchology* 1(1): 142-149, pl. 10, 11, 26.
- Conrad, T. A. (1865). Catalogue of the older eocene shells of Oregon. *American Journal of Conchology* 1(1): 150-154.
- Conrad, T. A. (1868). Notes on recent and fossil shells, with descriptions of new genera. *American Journal of Conchology* 4(3): 246-249, pl. 1, 18, 19, 20.
- Conrad, T. A. (1869). Descriptions of new fossil Mollusca, principally cretaceous. *American Journal of Conchology* 5(2): 96-108, pl. 1, 2, 9, 12, 13.
- Conrad, T. A. (1870). Notes on recent and fossil shells, with descriptions of new species. *American Journal of Conchology* 6(1): 71-78, pl. 1, 3, 10, 11.
- Corni, M. G., M. Farneti, et al. (1985). Histomorphological aspects of the gonads of *Chamelea gallina* (Linne) (Bivalvia; Veneridae) in autumn. *Journal of Shellfish Research* 5(2): 73-80.
- Corni, M. G. and M. Trentini (1986). A chromosomal study of *Chamelea gallina* (L.) (Bivalvia, Veneridae). *Bollettino di Zoologia* 53(1): 23-24.
- Corni, M. G. and M. Trentini (1990). The chromosomes of *Venerupis aurea* and *Ruditapes philippinarum* of the northern Adriatic Sea (Bivalvia, Heterodontida, Veneridae). *Venus, The Japanese Journal of Malacology* 49(3): 258-261.

Cosel, R. von (1995). Fifty-one new species of marine bivalves from tropical West Africa. *Iberus* 13(1): 1-115.

Cossmann, M. (1885). Catalogue des Coquilles Fossiles de l'éocène des environs de Paris. *Annales de la Société Royale Malacologique de Belgique Tome XXI* (Quatrième Série, Tome I, Année 1886): 17-186.

Coutts, P. J. (1974). Growth characteristics of the bivalve *Chione stutchburyi*. *New Zealand Journal of Marine and Freshwater Research* 8(2): 333-339

Cox, L. R. (1930). *Comus*, a new Genus of the Veneridae from South and East Africa. *Proceedings of the Malacological Society* 19: 20-22.

Cox, L. R., N. D. Newell, D.W. Boyd, C.C. Branson, C. Raymond, A. Chavan, A.H. Coogan, C. Dechaseaux, C.A. Fleming, F. Haas, L.G. Hertlein, E.G. Kauffman, A.M. Keen, A. LaRocque, A.L. McAlester, R.C. Moore, C.P. Nuttall, B.F. Perkins, H.S. Puri, L.A. Smith, T. Soot-Ryen, H.B. Stenzel, E.R. Trueman, R.D. Turner, J. Weir (1969). Bivalvia. In: Moore, Raymond C. (ed.), *Treatise on Invertebrate Palaeontology* (2), Boulder, Colorado, The Geological Society of America, Inc. and The University of Kansas. pp. N491-N952

Crane, J. M. Jr., L. G. Allen, C. Eisemann (1975). Growth rate, distribution, and population density of the northern quahog *Mercenaria mercenaria* in Long Beach, California. *California Fish and Game* 61(2): 68-91(April) [Errata, 61(4): 261, Oct.]

Crenshaw, M. A. and N. Watabe (1969). The muscle attachment to the shell of *Mercenaria mercenaria* [abst.]. *American Zoologist* 9(4): 1139

Crosse, H. A. A. (1870). On the species of Veneridae found in Japan. *Journal de Conchyliologie* 18: 437-438.

Cumber, R. (1983). Colour patterns of *Tawera spissa* (Deshayes, 1835) (Veneridae) in the far North. *Poirieria* 13(2): 8-9.

D'Asaro, C. N. (1968). The morphology of larval and postlarval *Chione cancellata* Linné (Eulamellibranchia: Veneridae) reared in the laboratory. *Bulletin of Marine Science* 17(4): 949-972

Dall, W. H. (1871). Revision of the classification of the Mollusca of Massachusetts. *Proceedings of the Boston Society of Natural History* 13: 240-257.

Dall, W. H. (1883). On a collection of shells sent from Florida by Mr. Henry Hemphill. *Proceedings of the United States National Museum* 6: 318-342.

Dall, W. H. (1891). On a new subgenus of Meretrix, with descriptions of two new species from Brazil. *The Nautilus* 5(3): 26-29.

Dall, W. H. (1902). Illustrations and descriptions of the new, unfigured, or imperfectly known shells, chiefly American, in the U. S. National Museum. *Proceedings of the United States National Museum* 24: 499-566.

Dall, W. H. (1902). New species of pacific coast shells. *The Nautilus* 16(4): 43-44.

Dall, W. H. (1902). On the Genus Gemma, Deshayes. *Journal of Conchology* 10: 238-243.

Dall, W. H. (1902). Synopsis of the family Veneridae and of the North American recent species. *Proceedings of the United States National Museum* 26: 335-412.

Dall, W. H. (1905). Note on Lucina (Miltha) childreni Gray and on a new species from the Gulf of California. *The Nautilus* 18(10): 110-113.

Dall, W. H. (1905). Notes on some preoccupied names of molluscs. *The Nautilus* 18(10): 113.

Dall, W. H. (1909). Notes, Paradione, n.n., vice Chionella. *Proceedings of the Malacological Society* 8: 197.

Dall, W. H. (1911). Some remarks on the nomenclature of the Veneridae. *Proceedings of the Malacological Society* 9(6): 349-351

Dall, W. H. (1914). Note on Clementia obliqua Jukes-Browne. *The Nautilus* 27(9): 103-104

Dall, W. H. (1916). Two new bivalve shells from Uruguay. *The Nautilus* 29(10): 112-113.

Davis, H. C. (1960). Effects of turbidity-producing materials in sea water on eggs and larvae of the clam (Venus (Mercenaria) mercenaria). *Biological Bulletin* 118(1): 48-54

De Boer, T. W. (1987). Venus fasciata (da Costa, 1778) (Bivalvia, Veneridae) in the Netherlands. *Basteria* 51(1-3): 57-58.

De Cassia, T. C. R. (1983). Sobre a ocorrência de Periglypta multicostata (Sowerby) na Formação Pirabas, Oligo-Mioceno do Brasil (Mollusca-Bivalvia). *Anais da Academia Brasileira de Ciências* 55(1): 140.

De Cassia, T. C. R. (1984). Sobre a ocorrência de Periglypta multicostata na Formação Pirabas, Oligoceno-Mioceno do Brasil. *Anais da Academia Brasileira de Ciências* 56(3): 283-286.

De la Rosa-Vélez, J. and A. M. de la Torre (1995). Genetic differentiation between the clams *Chione fructifraga* [sic] and the endemic *Chione cortezi* from the Gulf of California. *Unitas Malacologica, 12th International Malacological Congress, Abstracts*. 105

De Oliveira Castro Gueron, C. and A. C. dos Santos Coelho (1989). Consideracoes taxonomicas e morfologia de *Dosinia (Dosinia) concentrica* (Born, 1778) (Mollusca, Bivalvia, Veneridae). *Boletim Museu Nacional Rio de Janeiro Zoologia* No. 334: 1-19.

Defossez, J. M. and J. Daguzan (1996). In situ orientation of the Manila clam *Ruditapes philippinarum* (Bivalvia, Veneridae): preliminary results. *Vie et Milieu* 46(1): 21-24.

Del Pilar de Castro Fernandes, M. (1981). Contribuicao para o estudio de bivalves de Mocambique. *Garcia de Orta Serie de Zoologia* 10(1-2): 55-102.

Del Rio, C. J. (1994). Middle Miocene bivalves of the Puerto Madryn Formation, Valdes Peninsula, Chubut Province, Argentina (Lucinidae-Pholadidae). *Palaeontographica. Abteilung A: Palaeozoologie-Stratigraphie* 231(4-6): 93-132.

Del Rio, C.-J. (1997). Cenozoic biogeographic history of the eurythermal genus *Retrotapes*, new genus (subfamily Tapetinae) from southern South America and Antarctica. *The Nautilus* 110(3): 77-93.

Denis, A. (1979). L'utilisation du Microscope electronique a balayage dans l'examen des tissus mineralises chez les lamellibranches. Proceedings of the Sixth European malacological congress. *Malacologia* 18: 19-21.

Deshayes, G. P. (1853). *Catalogue of the Conchifera or bivalve shells in the collection of the British Museum*. Part I. Veneridae, Cyprinidae and Glauconomidae. London (British Museum). 216 pp. [concerning: Kabat (1989)].

Deshayes, G. P. (1854). Descriptions of new species of shells in the collection of Mr. Cuming. 1853[21](248): 1-11, pls. 18, 19

Deshayes, G. P. (1854). Observations sur les animaux de quelques genres de mollusques acéphales (*Chamostrea*, *Glaucone*, *Circe* and *Capsa*). *Proceedings of the Zoological Society of London*, for 1853[21](258): 167-172, pl. 21

Diaz, J. M. (1988). Occurrence of *Ventricolaria listeroides* Fischer-Piette and Testud, 1967 (Bivalvia: Veneridae) on the Caribbean coast of Colombia. *Caribbean Journal of Science* 24(1-2): 83-85.

- Dillon, R. T. (1992). Minimal hybridization between populations of the hard clams, *Mercenaria mercenaria* and *Mercenaria campechiensis*, co-occurring in South Carolina. *Bulletin of Marine Science* 50(3): 411-416
- Dillon, R. T. and J. J. Manzi (1989). Genetics and shell morphology in a hybrid zone between the hard clams *Mercenaria mercenaria* and *M. campechiensis*. *Marine Biology* 100(2): 217-222
- Dillon, R. T. and J. J. Manzi (1989). Genetics and shell morphology of hard clams (genus *Mercenaria*) from Laguna Madre, Texas. *The Nautilus* 103(2): 73-77
- Dillon, R. T. and J. J. Manzi (1992). Population genetics of the hard clam, *Mercenaria mercenaria*, the northern limit of its range. Canada, Department of Fisheries and Oceans, *Canadian Journal of Fisheries and Aquatic Sciences* 49(12): 2574-2578
- Dimmick, C. W. (1969). *Hinge and early shell development of representative venerid Bivalvia*. Doctoral Thesis, Mar. 1969, New Orleans, LA, Tulane University.: 89; Format: v, 84 numb. 1 incl. photos, tables, diagr.
- Dockery, D. T., III (1980). Color patterns of some Eocene molluscs. *Mississippi Geology* 1(1): 3-7.
- Doering, P. H. (1976). A burrowing response of *Mercenaria mercenaria* (Linnaeus, 1758) elicited by *Asterias forbesi* (Desor, 1848). *The Veliger* 19(2): 167-175
- Doering, P. H. (1981). Reduction of attractiveness to the sea star *Asterias forbesi* (Desor) by the clam *Mercenaria mercenaria* (Linnaeus). *Journal of Experimental Marine Biology and Ecology* 60(1): 47-61
- Doering, P. H. (1981). Observations on the behavior of *Asterias forbesi* feeding on *Mercenaria mercenaria*. *Ophelia* 20(2): 169-177
- Dohrn, H. (1880). Beiträge zur Kenntnis der Seeconchylien von Westafrika. *Jahrbücher der deutschen Malakozoologischen Gesellschaft* 7: 161-183.
- Drivas, J. and M. Jay (1990). Descriptions of new mollusks from Reunion Island. 2. Cerithidae, Lamellaridae, Buccinidae, Mitridae and Veneridae. *Venus, The Japanese Journal of Malacology* 49(4): 271-279.
- DuShane, H. (1966). Erroneous range extension for *Tivela stultorum* (Mawe, 1823). *The Veliger* 9(1): 86-87

- Eagar, R. M. (1978). Shape and function of the shell: a comparison of some living and fossil bivalve molluscs. *Biological Reviews of the Cambridge Philosophical Society* 53(2): 169-210
- Eldridge, P. J. and A. G. Eversole (1982). Compensatory growth and mortality of the hard clam, *Mercenaria mercenaria* (Linnaeus, 1758). *The Veliger* 24(3): 276-278
- Elston, R. A. (1986). Occurrence of branchial rickettsiales-like infections in two bivalve molluscs, *Tapes japonica* and *Patinopecten yessoensis*, with comments on their significance. *Journal of Fish Diseases* 9(2): 69-71
- Emerson, W. K. and M. K. Jacobson (1976). *The American Museum of Natural History guide to shells*. Land, freshwater, and marine, from Nova Scotia to Florida. Alfred A. Knopf, Inc. New York. pp. 1-482
- Ente di Sviluppo Agricolo del Veneto (1990). *Tapes philippinarum. Biologia e sperimentazione*. Venice (Ente di Sviluppo del Veneto). 299 pp.
- Fang, Y. (1982). Tidal zonation and cardiac physiology in four species of bivalves from Hong Kong. Pp. 849-857, in: Brian S. Morton and C. K. Tseng, *Proceedings of the First International Marine Biology Workshop: the marine fauna of Hong Kong and southern China*, Hong Kong, 1980. Vol. 2. Ecology, Morphology, Behaviour and Physiology. Hong Kong (Hong Kong University). viii + 557-933 pp.
- Farinati, E. A. (1978). Microfauna de moluscos querandinenses (Holoceno), Ingeniero White, Provincia de Buenos Aires. Revista de la Asociacion Geologica Argentina 33(3): 211-231.
- Feder, H. M., J. C. Hendee, P. Holmes, G. J. Mueller, A. J. Paul (1979). Examination of a reproductive cycle of *Protothaca staminea* using histology, wet weight-dry weight ratios, and condition indices. *The Veliger* 22(2): 182-187
- Feder, H. M. and A. J. Paul (1973). Abundance estimations and growth-rate comparisons for *Protothaca staminea* from three beaches in Prince William Sound, Alaska, with additional comments on size-weight relationships, harvesting and marketing. Alaska, University of Alaska, *Technical Report of the Institute of Marine Science* R73-3: 34 pp.
- Finlay, H. J. (1927). A further commentary on New Zealand Molluscan Systematics. *Transactions and Proceedings of the New Zealand Institute* 57: 320-485.
- Finlay, H. J. (1928). The recent Mollusca of the Chatham Islands. *Transactions and Proceedings of the New Zealand Institute* 59: 232-286.

Finlay, H. J. (1930). Additions to the recent molluscan fauna of New Zealand, No.3. *Transactions and Proceedings of the New Zealand Institute* 61: 222-247.

Finlay, H. J. (1930). Invalid molluscan names. *Transactions and Proceedings of the New Zealand Institute* 61(1): 37-48.

Fischer, P.-H., II and É. Fischer-Piette (1938). Sur quelques espèces de Tivela (Veneridae) et sur l'extension géographique de ce genre. *Muséum National d'Histoire Naturelle (Paris), Bulletin* (2)10(1): 86-92

Fischer-Piette, É. (1969). Mollusques récoltés par M. Hoffstetter sur les côtes de l'Équateur et des Îles Galapagos. Veneridae. *Muséum National d'Histoire Naturelle (Paris), Bulletin* (2)40(5): 998-1018, 3 pls.

Fischer-Piette, É. (1975). Révision des Venerinae, s. s. (Mollusques Lamellibranches). *Muséum National d'Histoire Naturelle (Paris), Mémoires* (n.s.) (A)93: 64 pp., 8 pls.

Fischer-Piette, É. and P.-H. Fischer, II. (1942). Révision des espèces vivants de Tivela et Eutivela du Muséum National d'Histoire Naturelle. *Journal de Conchyliologie* 85[(4)39](1): 5-49

Fischer-Piette, É. and B. Métivier (1971). Révision des Tapetinae (Mollusques bivalves). *Muséum National d'Histoire Naturelle (Paris), Mémoires* (n.s.) (A)71: 106 pp., 16 pls.

Fischer-Piette, É. and D. Vukadinovic (1972). Révision des Veneridae appartenant aux Comus, Saxidomus, Cyclininae, Gemminae et Clementiinae. *Journal de Conchyliologie* 109[(4)62](4): 130-167

Fischer-Piette, É. and D. Vukadinovic (1977). Suite des révisions des Veneridae (Moll. Lamellibr.) Chioninae, Samaranginae et complément aux Vénus. *Muséum National d'Histoire Naturelle (Paris), Mémoires* (n.s.) (A)106: 186 pp., 22 pls.

Fitch, J. E. (1950). The Pismo clam. *California Fish and Game* 36(3): 285-312

Fitch, J. E. (1959). The plight of the Pismo clam. *Outdoor California* 18(1): 3-4

Fitch, J. E. (1961). The Pismo clam. California, *Department of Fish and Game, Marine Resources Leaflet* 1: 23 pp.

Fitch, J. E. (1965). A relatively unexploited population of Pismo clams, Tivela stultorum (Mawe, 1823) (Veneridae). *Proceedings of the Malacological Society of London* 36(5): 309-312, pl. 13

Foster, N. R. (1991). *Intertidal bivalves: a guide to the common marine bivalves of Alaska*. University of Alaska Press, Fairbanks. pp. 1-152

Fraser, C. McLean. (1929). The spawning and free swimming larval periods of Saxidomus and Paphia. *Proceedings and Transactions of the Royal Society of Canada* (3)23[for 1929](5)(2): 195-198

Fraser, C. McLean and G. M. Smith (1928). Notes on the ecology of the little neck clam, Paphia staminea Conrad. *Proceedings and Transactions of the Royal Society of Canada* (3)22[for 1928](5)(2): 249-269, 5 pls.

Fraser, C. McLean and G. M. Smith (1928). Notes on the ecology of the butter clam, Saxidomus giganteus Deshayes. *Proceedings and Transactions of the Royal Society of Canada* (3)22[for 1928](5)(2): 271-286, 2 pls.

Frizzell, D. L. (1930). A new Pleistocene fossil from Port Blakely, Washington. *The Nautilus* 43(4): 120-121

Frizzell, D. L. (1930). The status of Paphia tenerrima alta Waterfall. *The Nautilus* 44(2): 48-50

Frizzell, D. L. (1931). A molluscan species new to the Recent West Coast fauna. *Transactions of the San Diego Society of Natural History* 6(21): 319-324, pl. 22

Frizzell, D. L. (1936). Preliminary reclassification of veneracean pelecypods. *Musée Royal d'Histoire Naturelle de Belgique, Bulletin* 12(34): 84 pp.

Fujita, T. (1898). On the colour pattern of the shell of Tapes philippinarum. *Zoological Magazine* (Tokyo) [Dobutsugaku Zasshi] 10(113): 85-90

Fulton, H. C. (1922). A list of the species and genera of recent Mollusca first described in 'Le Naturaliste'. *Proceedings of the Malacological Society* 15: 19-31.

Garcia Dominguez, F., J. Arvizu Martinez, S. Rodriguez Astudillo, R. Ramirez Sevilla (1994). Distribucion y abundancia de Chione californiensis (Pelecypoda: Veneridae) en la Ensenada de La Paz, Baja California Sur, Mexico. *Revista de Biología Tropical* 42(3): 531-536.

Garcia Dominguez, F., B. P. Ceballos Vazquez, M. Villalejo Fuerte, M. Arellano Martinez (1998). Reproductive cycle of the giant reef clam Periglypta multicostata (Sowerby, 1835) (Pelecypoda: Veneridae) at Isla Espiritu Santo, Baja California Sur, Mexico. *Journal of Shellfish Research* 17(4): 1009-1013.

Garcia Dominguez, F., S. A. Garcia Gasca, J. L. Castro Ortiz (1994). Spawning cycle of the red clam *Megapitaria aurantiaca* (Sowerby, 1831) (Veneridae) at Isla Espiritu Santo, Baja California Sur, Mexico. *Journal of Shellfish Research* 13(2): 417-423.

Garcia Dominguez, F. A., E. F. Felix Pico, O. Cruz del Carmen Juarez, N. Romero Ibarra (1994). Alimentacion de *Chione gnidia* (Broderip & Sowerby, 1829), *C. undatella* (Sowerby 1835) y *C. californiensis* (Broderip, 1835) (Veneridae) en la Ensenada de la Paz, B.C.S., Mexico. *Investigaciones Marinas Cicimar* 9(2): 113-117.

Garcia Gasca, S. A. and F. A. Garcia Dominguez (1995). [Comparative morphology of the marginal and paleal zones of the mantle of the clams *Megapitaria aurantiaca* (Sowerby, 1831) and *M. squalida* (Sowerby, 1835) (Bivalvia: Veneridae).]. *Anales de la Escuela Nacional de Ciencias Biologicas Mexico* 40(1-4): 173-181.

Gerard, A. (1978). Etude des garnitures chromosomiques de deux Veneridae: *Ruditapes decussatus* (L.) et *Ruditapes philippinarum* (Adams et Reeve). *Haliotis* 9(1): 69-71.

Giese, A. C. (1958). Myoglobin in the adductor of *Tivela* [abst.]. *Anatomical Record* 132(3): 442

Giese, A. C., M. A. Hart, A. M. Smith, M. A. Cheung (1967). Seasonal changes in body component indices and chemical composition in the Pismo clam *Tivela stultorum*. *Comparative Biochemistry and Physiology* 22(2): 549-561

Gillilan, W. (1964). Pismo clam survey. *Outdoor California* 25(2): 10-11

Girscher, W. (1938). Die am Strand der Nordsee-Insel Juist angespülten fossilen und rezenten Cardiidae, Veneridae und Mactridae. *Archiv für Molluskenkunde der Senckenbergischen Naturforschenden Gesellschaft* 70(1): 60-72.

Gladenkov, Y. B. and V. N. Sinelnikova (1990). Miocene mollusks and climatic in Kamchatka. *Trudy Geologicheskogo Instituta Akademii Nauk SSSR* 453: 1-173.

Glemarec, M. and D. Bouron (1978). Evolution de la maturite sexuelle chez six especes de bivalves des Glenan. *Haliotis* 9(1): 45-48.

Glibert, M. and L. van d. Poel (1966). *Les Bivalvia fossiles du cenozoique étranger des collections de l'Institut Royal des Sciences Naturelles de Belgique*; IV, Heteroconchia, 2 (super eme) partie; Corbiculidae a Petricolidae (fin). Brussels, Belgium, Institut Royal des Sciences Naturelles de Belgique. pp. 1-108

Glock, J. W. and K. K. Chew (1979). Growth, recovery, and movement of Manila clams, *Venerupis japonica* (Deshayes) at Squaxin Island, Washington. *National Shellfisheries Association, Proceedings* 69: 15-20

Glude, J. B. (1967). The effect of scoter duck predation on a clam population in Dabob Bay, Washington. *Proceedings of the National Shellfisheries Association* 55: 73-86

Gonzalez de Canales, M. L. (1990). Enzimas hidrolíticas en el aparato digestivo de las almejas *Ruditapes decussatus* (Linnaeus, 1758) y *Ruditapes philippinarum* (Adams & Reeve, 1850) (Pelecypoda: Veneridae). *Scientia Marina* 54(1): 89-93.

Goncharova, I. A. (1986). [*Biogeographical characteristics of Neogene Veneridae (Bivalvia) from the Mediterranean Basin.*] In: Kafanov, A.I. (Ed.) [Palaeogene Neogene Bivalvia of the Far East and eastern Paratethys. Collection of scientific works.] Akademiya Nauk SSSR, Vladivostok. pp. 101-106

Goncharova, I. A. (1986). [*The systematics and history of Tapetinae (Bivalvia, Veneridae) from the Neogene seas of western Eurasia.*] In: Kafanov, A.I. (Ed.) [Palaeogene Neogene Bivalvia of the Far East and eastern Paratethys. Collection of scientific works.] Akademiya Nauk SSSR, Vladivostok. pp. 75-100

Goodsell, J. G., S. C. Fuller, A. G. Eversole, M. Castagna, R. A. Lutz (1992). Larval and early postlarval shell morphology of several venerid clams. *Journal of the Marine Biological Association of the United Kingdom* 72(1): 231-255.

Goodwin, C. L. (1973). Distribution and abundance of subtidal hard-shell clams in Puget Sound, Washington. Washington, *Department of Fisheries, Technical Report* 14: v + 81 pp.

Gordon, J. and M. R. Carriker (1978). Growth lines in a bivalve mollusk: subdaily patterns and dissolution of the shell. *Science* 202(4367): 519-521

Grabyan, R. J. (1973). Variations in concentrations of magnesium and strontium in Recent shells of *Tivela stultorum*. *Bulletin of the Southern California Academy of Sciences* 72(1): 42-48

Gray, J. E. (1838). Catalogue of the species of the genus *Cytherea*, of Lamarck, with the description of some new genera and species. *The Analyst, a Quarterly Journal of Science, Literature, Natural History, and the Fine Arts* 8(24): 302-309

Gray, S. (1980). Comparative morphology and systematics of two species of *Transennella* (Bivalvia: Veneridae). *Western Society of Malacologists Annual Report* 13: 11.

- Gray, S. (1982). Morphology and taxonomy of two species of the genus *Transennella* (Bivalvia: Veneridae) from western North America and a description of *T. confusa* sp. nov. *Malacological Review* 15: 107-117.
- Green, R. H. and K. D. Hobson (1970). Spatial and temporal structure in a temperate intertidal community, with special emphasis on *Gemma gemma* (Pelecypoda: Mollusca). *Ecology* 51(6): 999-1011
- Gregorio, M. A. d. (1884). Studi su talune Conchiglie Mediterranee, viventi e fossili. Con una rivista del Gen. Vulsella. *Società Malacologica Italiana, Bulletino* 10-11, 1884-85: 27-203.
- Gustafson, A. H. (1955). Growth studies in the quahog, *Venus mercenaria*. *Proceedings of the National Shellfisheries Association* 45: 140-150
- Haas, F. (1955). *Bivalvia Teil 2* (hier: Veneridae). Dr. H. G. Bronns Klassen und Ordnungen des Tierreichs. Leipzig, Akademische Verlagsgesellschaft Geest and Portig K.-G. 3 (Mollusca); 3. Abteilung: Bivalvia: 306-308.
- Hagn, H. and E. Martini (1990). A lower Eocene shell bed and some quaternary mollusks from ODP Leg 112 off Peru. *Proceedings of the Ocean Drilling Program*, Peru continental margin; covering Leg 112 of the cruises of the Drilling Vessel JOIDES Resolution, Callao, Peru to Valparaiso, Chile, sites 679-688, 20 October 1986-25 December 1986. E. Suess, R. von Huene, K. C. Emeis et al. College Station, TX, United States, Texas A & M University Ocean Drilling Program. 112: 329-333.
- Hall, C. A. Jr. (1975). Latitudinal variation in shell growth patterns of bivalve molluscs: implications and problems. pp. 163-175, in: G. D. Rosenberg and S. K. Runcorn, *Growth rhythms and the history of the Earth's rotation*. London, etc. (Wiley). xvi + 559 pp.
- Hall, C. A., Jr., W. A. Dollase, C. E. Corbató (1974). Shell growth in *Tivela stultorum* (Mawe, 1823) and *Callista chione* (Linnaeus, 1758) (Bivalvia): annual periodicity, latitudinal differences, and diminution with age. *Palaeogeography, Palaeoclimatology, Palaeoecology* 15(1): 33-61
- Hamwi, A. and H. H. Haskin (1969). Oxygen consumption and pumping rates in the hard clam *Mercenaria mercenaria*: a direct method. *Science* 163(3869): 823-824
- Hansen, B. (1953). Brood protection and six ratio of *Transennella tantilla* (Gould), a Pacific bivalve. *Dansk Naturhistorisk Forening (Copenhagen), Videnskabelige Meddelelser* 115: 313-324

Harbo, R. M. and N. F. Bourne (1992). A newly developing commercial fishery for the subtidal clam, Compsomyax subdiaphana, in British Columbia [abst.]. *Journal of Shellfish Research* 11(2): 553

Harrington, R. J. (1986). *Growth patterns within the genus Protothaca (Bivalvia: Veneridae) from the Gulf of Alaska to Panama: Paleotemperatures, Paleobiogeography and Paleolatitudes*, PhD Thesis, University of California Santa Barbara: pp. 1-249.

Harrington, R. J. (1987). Skeletal growth histories of Protothaca staminea (Conrad) and Protothaca grata (Say) throughout their geographic ranges, northeastern Pacific. *The Veliger* 30(2): 148-158

Harte, M. E. (1992). An eastern Pacific Mercenaria and notes on other chionine genera (Bivalvia: Veneridae). *The Veliger* 35(2): 137-140

Harte, M. E. (1992). A new approach to the study of bivalve evolution. *American Malacological Bulletin* 9(2): 199-206

Harte, M. E. (1998). Translating trees into taxonomy within Veneridae (Bivalvia): a critique of two recent papers. *Malacologia* 39(1-2): 215-219.

Harte, M. E. (1992). The Veneridae of Polynesia. *Xenophora* No. 59: 4-10.

Harte, M. E. (1993). A review of Pitar (Hyphantosoma) Dall, 1902 (Veneridae: Pitarinae) and a description of Pitar (H.) festoui sp. nov. *The Veliger* 36(4): 343-350.

Harte, M. E. (1998). Is Cyclininae a monophyletic subfamily of Veneridae (Bivalvia)? *Malacologia* 40(1-2): 297-304.

Harte, M. E. (1998). Translating trees into taxonomy within Veneridae (Bivalvia): a critique of two recent papers. *Malacologia* 39(1-2): 215-219.

Harte, M. E. and K. L. Lamprell (1999). Lioconcha (Sulcilioconcha) caledonensis sp. nov., a species of Veneridae (Bivalvia) from New Caledonia. *The Veliger* 42(1): 97-100.

Hawbecker, A. C. (1939). Feeding of gulls on Pismo clams. *Condor* 41(3): 120

Hayward, P. J. (1990). Mollusca 2: Bivalvia and Cephalopoda. In: Hayward, P.J. and Ryland, J.S. (Eds). *The marine fauna of the British Isles and north west Europe* 2, Clarendon Press, Oxford, pp. 628-996.

- Hayward, P. J., G. D. Wigham, N. Yonow (1995). Molluscs. In: Hayward, P.J. and Ryland, J.S. (Eds). *Handbook of the marine fauna of north west Europe*. Oxford University Press, Oxford, New York etc. pp. 484-628
- Healy, J. and K. Lamprell (1992). New species of Veneridae, Cardiidae, Crassatellidae, Tellinidae and Mactridae from Australia (Veneroida, Bivalvia, Mollusca). *Journal of the Malacological Society of Australia* 13: 75-97.
- Hedley, C. (1906). The Mollusca of Mast Head Reef, Capricorn Group, Queensland. *Proceedings of the Linnean Society of New South Wales* 31: 453-479.
- Henry, M., M. Auffret, E. Boucaud Camou (1990). Aspects ultrastructuraux et fonctionnels des hemocytes de quatre familles de bivalves (Ostreidae, Veneridae, Mytilidae, Pectinidae). *Haliotis* 20: 195-196.
- Herrington, W. C. (1925). The Pismo clam crop. *California Fish and Game* 11(3): 103-109
- Herrington, W. C. (1926). Depletion of the Pismo clam in California. *California Fish and Game* 12(3): 117-124
- Herrington, W. C. (1930). The Pismo clam. Further studies of its life history and depletion. *California, Department of Natural Resources, Division of Fish and Game, Fish Bulletin* 18: 67 pp.
- Hertlein, L. G. and A. M. Strong (1948). Eastern Pacific Expeditions of the New York Zoological Society. XXXIX. Mollusks from the West Coast of Mexico and Central America. Part VI. *Zoologica* 33(4): 163-198, pl. 1-2.
- Hikida, Y. (1996). Shell structure and its differentiation in the Veneridae (Bivalvia). *Journal of The Geological Society of Japan* 102(10): 847-865.
- Hillman, R. E. (1961). Formation of the periostracum in *Mercenaria mercenaria*. *Science* 134(3492): 1754-1755
- Hillman, R. E. (1964). The functional morphology of the fourth fold of the mantle of the northern quahog, *Mercenaria mercenaria* (L.). *Journal of the Elisha Mitchell Scientific Society* 80(1): 8-12
- Hillman, R. E. and C. N. Shuster, Jr. (1962). Observations on the mantle of the northern quahog, *Mercenaria mercenaria* L. *Proceedings of the National Shellfisheries Association* 51: 15-22

- Hily, A. (1983). Remarques sur l'ultrastructure de la glande digestive chez des juvéniles de Ruditapes philippinarum. *Journal of Molluscan Studies, Supplement 12A*: 51-59
- Hodson, F., H. K. Hodson, G. D. Harris (1927). Some Venezuelan and Caribbean Mollusks. *Bulletin of American Paleontology* 13(49): 45-153.
- Holland, D. A. and K. K. Chew (1974). Reproductive cycle of the Manila clam (Venerupis japonica) from Hood Canal, Washington. *Proceedings of the National Shellfisheries Association* 64: 53-58
- Houghton, J. P. (1973). *The intertidal ecology of Kiket Island, Washington, with emphasis on age and growth of Protothaca staminea and Saxidomus giganteus (Lamellibranchia; Veneridae)*. PhD Thesis, University of Washington. Seattle, WA, pp. 1-179.
- Houghton, J. P. (1977). Age and growth of Protothaca staminea (Conrad) and Saxidomus giganteus (Deshayes) at Kiket Island, Washington [abst.]. *Proceedings of the National Shellfisheries Association* 69: 119
- Howard, A. D. (1953). Some viviparous pelecypod mollusks. *Wasmann Journal of Biology* 11(2): 233-240
- Ieyama, H. (1980). Studies on the chromosomes in three species of the Veneridae (Bivalvia, Heterodontia). *Venus, The Japanese Journal of Malacology* 39(1): 49-55.
- Ieyama, H. (1985). Karyotype in Ruditapes philippinarum (Bivalvia: Veneridae). *Special Publication of the Mukashima Marine Biological Station*. 83-85
- Ihering, H. v. (1923). Transgression des Meeres während der Ablagerung der Pampas. *Archiv für Molluskenkunde* 55: 183-193, pl. 7.
- Ikebe, N. (1941). On Katelysia (Nipponomarcia) nakamurai, subgen. et sp. nov., from the Togarian Miocene of Japan. *Venus, The Japanese Journal of Malacology* 11(2-3): 49-53.
- Ikeda, S. (1893-1894). On the colour pattern of the shell of Tapes philippinarum. *Zoological Magazine* (Tokyo) [Dobutsugaku Zasshi] 5(56): 222-226; (58): 292-295; 6(63): 13-18; 6(64): 39-47
- Ikematsu, W. (1957). Ecological studies on the clam, Tapes japonica (Reeve) -- II. On the setting season and the growth in early young stage. *Bulletin of the Japanese Society of Scientific Fisheries* 22(12): 736-741

- Ionesi, B. and M. Ciobanu (1978). Contributii la studiul faunei sarmatiene din Platforma Moldoveneasca (Dealul lui Stan - Bozieni); Partea a II-a. *Anuarul Muzeului de Stiinte Naturale Piatra Neamt, Seria Geologie Geografie* 4: 159-183.
- Iredale, T. (1915). Some more misused molluscan generic names. *Proceedings of the Malacological Society* 11: 291-306.
- Iredale, T. (1917). More molluscan name-changes, generic and specific. *Proceedings of the Malacological Society* 12: 322-330.
- Iredale, T. (1924). Results from Roy Bell's molluscan collections. *Proceedings of the Linnean Society of New South Wales* 49: 179-278.
- Iredale, T. (1927 - 1929). Strange molluscs in Sydney Harbour. *The Australian Zoologist* 5: 337-354.
- Iredale, T. (1930). More notes on the marine mollusca of New South Wales. *Records of the Australian Museum* 17(9): 384-407, pl. lxii-lxv.
- Iredale, T. (1936). Australian Molluscan Notes No. 2. *Records of the Australian Museum* 19(5): 267-340, pl. xx-xxiv.
- Jackson, J. B. C. (1968). Bivalves: spatial and size-frequency distributions of two intertidal species. *Science* 161(3840): 479-480
- Jagnow, B. and F. Gosselck (1987). Bestimmungsschlüssel für die Gehäuseschnecken und Muscheln der Ostsee. *Mitteilungen aus dem Zoologischen Museum in Berlin* 63(2): 191-268.
- Janssen, A. W., G. A. Peeters, L. van der Slik (1984). [The fossil shells of the Netherlands stranded in rivermouths, second series, 8 (conclusion).]. *Basteria* 48(4-5): 91-219.
- Jay, J. C. (1852). *Supplement to the fourth edition of the catalogue of the shells in the collection of John C. Jay*, M.D. New York (Bailliére). pp. 465-479.
- Jiang, J. and F. Xu (1992). A new species of Venefidae [Veneridae] (Bivalvia) from Hong Kong waters. *Journal of Oceanography in Taiwan Strait* 11(4): 291-293.
- Johnson, M. S. and R. Black (1990). *Genetic divergence in venerid clams in Shark Bay, Western Australia*. In: Berry, P.F., Bradshaw, S.D. and Wilson, B.R. (Eds). Research in Shark Bay. Report of the France Australe Bicentenary Expedition Committee. Western Australian Museum. pp. 1-325

Jones, C. C. (1979). Anatomy of *Chione cancellata* and some other chionines (Bivalvia, Veneridae). *Malacologia* 19(1): 157-199.

Jones, C. C. (1984). *Ventricolaria judithae* n. sp. (Bivalvia: Veneridae) from the Oligocene of North Carolina with comments on ecology. *Proceedings of the Academy of Natural Sciences of Philadelphia* 136: 152-164.

Jones, C. and K. K. Chew (1975). Planting hatchery spawned Manila clams (*Venerupis japonica*) in Puget Sound beaches [abst.]. *Proceedings of the National Shellfisheries Association* 65: 8-9

Jousseaume, M. I. D. (1891). Reflexions sur la Faune Malacologique de la Mer Rouge. *Annales des sciences naturelles* (Zoologie et Paleontologie) 12: 343-350.

Jukes-Browne, A. J. (1908). On the Genera of Veneridae represented in the cretaceous and older tertiary deposits. *Proceedings of the Malacological Society of London* 8: 148-177.

Jukes-Browne, A. J. (1912). The Genus *Dosinia* and its subdivisions. *Proceedings of the Malacological Society of London* 10: 95-106.

Jukes-Browne, A. J. (1913). On *Tivela* and *Gratelouphia*. *Proceedings of the Malacological Society of London* 10(4): 266-273

Jukes-Browne, A. J. (1913). On a new species of *Clementia*. *Annals and Magazine of Natural History* (8)12(67): 58-62, pl. 1

Jukes-Browne, A. J. (1913). On *Callista*, *Amiantis*, and *Pitaria*. *Proceedings of the Malacological Society of London* 10: 335-347.

Jukes-Browne, A. J. (1913). On the shells known as *Gemma*, *Parastarte*, and *Psephidia*. *The Annals and Magazine of Natural History, including Zoology, Botany, and Geology*. 12(8): 473-480.

Jukes-Browne, A. J. (1914). A synopsis of the family Veneridae. Part I. *Proceedings of the Malacological Society of London* 11: 58-74.

Jukes-Browne, A. J. (1914). A synopsis of the family Veneridae. Part II. *Proceedings of the Malacological Society of London* 11: 75-94.

Jukes-Browne, A. J. (1914). Note on *Clementia subdiaphana*, Carp. *Annals and Magazine of Natural History* (8)13(75): 338-339

Kabat, A. R. (1985). The allometry of brooding in *Transennella tantilla* (Gould) (Mollusca: Bivalvia). *Journal of Experimental Marine Biology and Ecology* 91(3): 271-279

Kabat, A. R. (1986). Effects of trematode parasitism on reproductive output of the bivalve *Transennella tantilla*. *Canadian Journal of Zoology* 64(1): 267-270

Kadolsky, D. (1988). Stratigraphy and mollusc faunas of "Landschneckenkalk" and "Cerithienschichten" in the Mainz Basin (Late Oligocene to Early Miocene?) - stratigraphical, paleogeographical and paleoecological results. *Geologisches Jahrbuch Reihe A* 110: 69-133.

Kafanov, A. I. (2000). A review of Cenozoic North Pacific *Liocyma* (Bivalvia, Veneridae). *Bulletin of the Russian Far-East Malacological Society* 3: 5-10 [pl. published in Kafanov et al. (1999)]

Kafanov, A. I. and K. A. Lutaenko (1997). New data on the bivalve mollusc fauna of the North Pacific Ocean. 3. On some species of the subfamily Dosiniinae (Veneridae). *Ruthenica* 7(2): 149-153.

Kafanov, A. I. and K. A. Lutaenko (1998). New data on the bivalve mollusc fauna of the North Pacific. 6. *Callista trigonoovata* Scarlato in Volova et Scarlato, 1980 is a new synonym of *Callista (Ezocallista) brevisiphonata* (Carpenter, 1864) (Veneridae). *Ruthenica* 8(1): 75-76.

Kalashnikova, S. A. and N. A. Aizdaicher (1993). On the larval development of *Spisula sachalinensis* (Cardida, Mactridae), *Ruditapes philippinarum* (Cardida, Veneridae), and *Crenomytilus grayanus* (Mytilida, Mytilidae) in culture. *Zoologicheskii Zhurnal* 72(7): 13-19.

Kamada, Y. (1980). Tertiary molluscan fossils from the coalfields of Kyushu 2. (Okinoshiman molluscan fossils). *Atlas of Japanese Fossils* 56(332).

Katkansky, S. C. and R. W. Warner (1969). Infestation of the rough-sided little neck clam, *Protothaca laciniata*, in Morro Bay, California, with larval cestodes (*Echeneibothrium* sp.). *Journal of Invertebrate Pathology* 13(1): 125-128

Katkansky, S. C., R. W. Warner, R. L. Poole (1969). On the occurrence of larval cestodes in the Washington clam, *Saxidomus nuttalli*, and the gaper clam, *Tresus nuttalli*, from Drakes Estero, California. *California Fish and Game* 55(4): 317-322

- Kauffman, E. G., W. P. Elder, J. I. Kirkland, B. B. Sageman, N. F. Sohl (1987). *Evolutionary patterns and biostratigraphy of Cretaceous bivalves and gastropods, Western Interior Cretaceous basin, North America*. Geological Association of Canada, Mineralogical Association of Canada; 1987 joint annual meeting; program with abstracts--Association Geologique du Canada, Association Mineralogique du Canada, reunion annuelle conjointe; programme et resumes. Anonymous. Waterloo, ON, Canada, Geological Association of Canada. 12: 60.
- Kautsky, F. (1929). Ein neues Veneridengenus "Gomphomarcia" aus dem europäischen Miozän nebst Bemerkungen über die systematische Stellung von Tapes gregarius Partsch und Tapes senescens Dod. *Annalen des Naturhistorischen Museums in Wien* 43: 379-387.
- Kautsky, F. (1929). Biologische Studien über den Schlossapparat von Tapes. *Palaeobiologica* 2(4-5): 202-212, pls. 15-18
- Kautsky, F. (1936). Die Veneriden und Petricoliden des niederösterreichischen Miozäns; eine biologisch-stratigraphische Untersuchung. *Bohrtechniker-Zeitung* 54(7-11): 157-160, 173-177, 202-205, 225-228, 250-252.
- Kautsky, F. (1937). Die Veneriden und Petricoliden des niederösterreichischen Miozäns; eine biologisch-stratigraphische Untersuchung. *Bohrtechniker-Zeitung* 1: 11-14.
- Kawamoto, T. (1934). The colour change of the shell of Venerupis philippinarum [in Japanese]. *Venus* 4(5): 310-311
- Keck, R., D. Maurer, R. E. Malouf (1974). Factors influencing the setting behavior of larval hard clams, Mercenaria mercenaria. *Proceedings of the National Shellfisheries Association* 64: 59-67
- Keen, A. M. (1951). Outline of a proposed classification of the pelecypod family Veneridae. *Conchological Club of Southern California, Minutes* 113: 2-11 [1-10]
- Keen, A. M. (1954). Nomenclatural notes on the pelecypod family Veneridae. *Conchological Club of Southern California, Minutes* 139: 50-55
- Keen, A. M. (1954). Ventricolaria and Similivenus insolida, new names in the pelecypod family Veneridae. *Journal of Paleontology* 28(2): 217-218.
- Keen, A. M. (1958). New mollusks from tropical West America. *Bulletin of American Paleontology* 38(172): 235-255.

Keen, A. M. (1969). Superfamily Veneracea. pp. 670-690, in: Leslie Reginald Cox et al., Part N [Bivalvia], Mollusca 6, vols. 1 and 2: xxxvii + 952 pp. In: Raymond C. Moore, ed., *Treatise on invertebrate paleontology*. Lawrence, Kansas (Geological Society of America and University of Kansas).

Kellogg, J. L. (1903). Feeding habits and growth of Venus mercenaria. *Bulletin of the New York State Museum* 71(10[296]): 28 pp.

Kennish, M. J. (1980). Shell microgrowth analysis: Mercenaria mercenaria as a type example for research in population dynamics. pp. 255-294, in: Donald C. Rhoads and Richard Arthur Lutz, eds., *Skeletal growth of aquatic organisms*. New York and London (Plenum). xiii + 750 pp.

Kilburn, R. and E. Rippey (1982). *Sea shells of southern Africa*. Macmillan South Africa, Johannesburg. pp. 1-249

Kinoshita, T. and S. Shibuya (1939). [Scientific name and spawning season of Japanese necked clam in Hokkaido] [in Japanese]. *Hokkaido Fisheries Experimental Station, Report* 410: 3-7

Kishinouye, K. (1908). On the colour pattern of the shell of Tapes philippinarum. *Conchological Magazine* (Kyoto) 2(12): 410-411

Kline, T. C. (1983). The effect of population density on the growth of the butter clam Saxidomus gigantus [sic] [abst.]. *Journal of Shellfish Research* 3(1): 112

Kō, Y. (1957). Some histological notes on the gonads of Tapes japonica Deshayes. *Bulletin of the Japanese Society of Scientific Fisheries* 23(7-8): 394-399, 1 pl.

Kobelt, W. M., O. von (1903). Catalog der Familie Buliminidae. 'Literatur'. *Nachrichtenblatt der deutschen Malakozoologischen Gesellschaft* 3-4 36-63.

Kozloff, E. N. (1987). *Marine invertebrates of the Pacific northwest*. University of Washington Press, Seattle and London. pp. 1-511

Kubo, H. (1993). New record of two Japanese venerids from Okinawa-Jima Island. *Chiribotan* 23(4): 97-99.

Küster, H. C. and L. G. C. L. Pfeiffer (1841-1872). Die Familie der Venusmuscheln. Veneracea. Lucinen. Galateen. Corbis. Systematisches Conchylien-Cabinet von Martini und Chemnitz 11(1): 302 pp., 42 pls. [pp. 1-32, pls. 1-3, 5-10, 1841, Küster; pls. 11-15, 1842, Küster; 1-32 (reissue), 33-56; pls. 4, 16-24, 1868; 57-160, pls. 25-34, 1869; 161-224, pls. 35-38, 1870; 225-302, pls. 39-42, 1872, Pfeiffer]

- Kuroda (1945). [New species of Japanese shells.] *Venus* 14: 29-42.
- Kuroda, T. H., Tadashige (1951). Donacidae and Myidae in Japan. *Illustrated Catalogue of Japanese Shells* 12: 71-78.
- Kuroda, T. H., Tadashige (1951). Nomenclatorial notes. *Illustrated Catalogue of Japanese Shells* 13: 86.
- Kvitek, R. G. (1992). Paralytic shellfish poisoning toxins as a chemical defense in the butter clam (*Saxidomus giganteus*) [abst.]. *Annual Report of the Western Society of Malacologists* 24: 11-12.
- Lajoie K. R., J. F. Wehmiller, G. L. Kennedy (1980). Inter- and intrageneric trends in apparent racemization kinetics of amino acids in Quaternary mollusks. In: P. E. Hare, T. C. Hoering and K. King, Jr. (Eds.) *Biogeochemistry of amino acids*. New York, NY, United States, John Wiley and Sons: pp. 305-340.
- Lam, V. W. W. (1980). Shell form and diagnostic differences in the structure of the siphons and ciliary currents of the ctenidia in coastal species of the Tapetinae (Bivalvia: Veneracea) in Hong Kong. Pp. 11-31, 9 pls., in: Brian S. Morton, ed., *The malacofauna of Hong Kong and southern China*. Hong Kong (Hong Kong University). vi + 345 pp.
- Lambiotte, M. (1978). Espèces européennes récentes de la super-famille des Veneroidea C.S. Rafinesque, 1815 (Mollusca: Lamellibranchiata). *Informations de la Société Belge de Malacologie* 6(2): 45-60.
- Lamprell, K. L. and J. M. Healy (1997). New species of Pteriidae, Pinnidae, Veneridae, Euciroidae and Corbulidae from Australia (Mollusca: Bivalvia: Veneroida). *Memoirs of the Queensland Museum* 42(1): 271-281.
- Lamprell, K. L. and J. M. Healy (1997). *Pitar* Römer from Australian and adjacent waters, with descriptions of four new species (Mollusca: Bivalvia: Veneridae). *Memoirs of the Queensland Museum* 42(1): 283-306.
- Lamprell, K. L. and J. M. Healy (1999). *Granicorium* from Australian waters, with descriptions of a new species (Mollusca: Bivalvia: Veneridae). *Molluscan Research* 20(1): 73-78.
- Lamprell, K. L. and J. Stanisic (1996). *Callista*, *Lioconcha* and *Pitar* in New Caledonia and adjacent waters (Mollusca, Veneridae). *Molluscan Research* 17(30): 27-48.

Lamy, E. (1923). Révision des Venerupis vivants du Muséum National d'Histoire Naturelle de Paris. *Journal de Conchyliologie* 67[(4)21](4): 275-308, pl. 3 [part]

Lamy, E. (1936). Note sur le Cytherea callosa Conrad (Moll. Lamellibr.). *Muséum National d'Histoire Naturelle (Paris)*, *Bulletin* (2)8(2): 158-159

Le Pennec, M. (1980). The larval and post-larval hinge of some families of bivalve molluscs. *Journal of the Marine Biological Association of the United Kingdom* 60(3): 601-618

Le Pennec, M. (1983). Ontogenèse et morphogenèse de l'isthme palléal chez Ruditapes philippinarum (Adams et Reeve) (Bivalvia, Veneridae). *Cahiers de Biologie Marine* 24(3): 327-335, 4 pls.

Le Pennec, M. and A. Lucas (1971). Comparative growth and morphology of some venerid larvae (Bivalvia, Veneridae). *Malacological Review* 3(2): 175-183

Le Pennec, M. and H. Petit (1983). Etude de l'isthme palleal de Ruditapes philippinarum, Adams & Reeve (Bivalvia, Veneridae) au microscope electronique à balayage. *Haliotis* 13: 83-89.

Leach, W. E. (1847). The classification of the British Mollusca. *The Annals and Magazine of Natural History, including Zoology, Botany, and Geology* 20: 267-273.

Lee, S. Y. (1996). Distribution pattern and interaction of two infaunal bivalves, Tapes philippinarum (Adams and Reeve) and Anomalocardia squamosa (Linnaeus) (Bivalvia: Veneridae). *Journal of Experimental Marine Biology and Ecology* 201(1-2): 253-273.

Lindberg, D. R. (1990). Transennella Dall versus Nutricola Bernard (Bivalvia: Veneridae): an argument evolutionary systematics. *Journal of Molluscan Studies* 56(1): 129-132.

Liu, H. P., K. Y. Liao, W. I Wu (1992). Comparison on the reproductive ecology of Meretrix lusoria from Tam-shui and Lu-kang area (Bivalvia [Bivalvia]: Veneridae). *Bulletin of Malacology Republic of China* 17: 69-77.

Loesch, J. G. and D. S. Haven (1973). Estimated growth functions and size-age relationships of the hard clam, Mercenaria mercenaria, in the York River, Virginia. *The Veliger* 16(1): 76-81

Loosanoff, V. L. (1936). Sexual phases in the quahog. *Science* 83(2151): 287-288

Loosanoff, V. L. (1937). Development of the primary gonad and sexual phases in Venus mercenaria Linnaeus. *Biological Bulletin* 72(3): 389-405

Loosanoff, V. L. (1937). Seasonal gonadal changes of adult clams, *Venus mercenaria* (L.). *Biological Bulletin* 72(3): 406-416

Loosanoff, V. L. (1937). Spawning of *Venus mercenaria* (L.). *Ecology* 18(4): 506-515

Loosanoff, V. L. (1959). The size and shape of metamorphosing larvae of *Venus Mercenaria* grown at different temperatures. *Biological Bulletin* 117(2): 303-318

Loosanoff, V. L. and H. C. Davis (1950). Conditioning *V. mercenaria* for spawning in winter and breeding its larvae in the laboratory. *Biological Bulletin* 98(1): 60-65

Loosanoff, V. L. and H. C. Davis (1963). Rearing of bivalve mollusks. *Advances in Marine Biology* 1: 1-136

Losi, M. (1974). Tendenze evolutive nei Veneridi (Lamellibranchi) terziari della Venezia. *Atti e Memorie dell'Accademia Patavina di Scienze Lettere ed Arti* 86(2): 41-56.

Lozada, E. and H. Bustos (1984). Madurez sexual y fecundidad de *Venus antiqua antiqua* King & Broderip 1835 en la bahía de Ancud (Mollusca: Bivalvia: Veneridae). *Revista de Biología Marina* 20(2): 91-112.

Luckens, P. A. (1990). Distribution, size-frequency, and growth-ring analyses of *Tawera mawsoni* (Bivalvia: Veneridae) at Macquarie Island. *New Zealand Journal of Marine and Freshwater Research* 24(1): 59-73.

Luckens, P. A. (1991). Distribution, growth rate, and death from octopod and gastropod predation of *Tawera bollonsi* (Bivalvia: Veneridae) at the Auckland Islands. *New Zealand Journal Of Marine And Freshwater Research* 25(3): 255-268.

Lussi, M. (1996). A new species of *Tivela* Link, 1807 from South Africa (Mollusca, Bivalvia, Veneridae). *Conchiglia* 28(279): 40-42.

MacKenzie, C. L., Jr. (1977). Predation on hard clam (*Mercenaria mercenaria*) populations. *Transactions of the American Fisheries Society* 106(6): 530-537

Maes, P. and C. Paillard (1992). Effet du Vibrio P1, pathogène de *Ruditapes philippinarum*, sur d'autres espèces de bivalves. *Ifremer* 14: 141-148.

Malakhov, V. V., L. A. Medvedeva, R. V. Gareeva (1989). Embryonal and early larval development of clam *Callista brevisiphonata* (Cardiida, Veneridae). *Zoologicheskii Zhurnal* 68(6): 130-134.

- Mann, R. (1979). The effect of temperature on growth, physiology, and gametogenesis in the Manila clam *Tapes philippinarum* (Adams & Reeve, 1850). *Journal of Experimental Marine Biology and Ecology* 38(2): 121-133
- Mano, K. (1969). (Structure of hinge teeth of the bivalve Veneridae). Chishitsugaku Zasshi = *Journal of the Geological Society of Japan* 75(2): 109.
- Marsden, I. D. and R. M. Pilkington (1995). Spatial and temporal variations in the condition of *Austrovenus stutchburyi* Finlay, 1927 (Bivalvia: Veneridae) from the Avon-Heathcote estuary, Christchurch. *New Zealand Natural Sciences* 22(0): 57-67.
- Martinez, E. R. (1984). Notas acerca del guacuco *Tivela mactroides* Born (Mollusca-Bivalvia de la familia Veneridae). *Natura* 76: 41-42.
- Marwick, J. (1927). The Veneridae of New Zealand. *Transactions and Proceedings of the New Zealand Institute* 57: 567-636.
- Marwick, J. (1928). The tertiary mollusca of the Chatham Islands including a generic revision of the New Zealand Pectinidae. *Transactions and Proceedings of the New Zealand Institute* 58: 432-506.
- Marwick, J. (1938). *Notocallista* and its allies. *Transactions and Proceedings of the Royal Society of New Zealand* 68: 60-81.
- Matsukuma, A. (1996). Transposed hinges: A polymorphism of bivalve shells. *Journal of Molluscan Studies* 62(4): 415-431.
- Matsukuma, A., T. Ozawa, W. Yoosukh (1988). *Paphia (Protapes) irrediviva* Makiyama, an extinct tropical embayment element of Japanese Cenozoic mollusks, and allied species from the Indo-West Pacific. In: J. A. Grant Mackie, K. Masuda, K. Mori and K. Ogasawara (Eds.). *Professor Tamio Kotaka commemorative volume on Molluscan paleontology*. Sendai, Japan, Saito Ho-On Kai Museum. 2: 405-423.
- Matsukuma, A. and W. Yoosukh (1988). Living *Bassina* and *Placamen* (Bivalvia; Veneridae) from the Australasia and the Indo-West Pacific (studies on the Kawamura Collection (Mollusca) stored in the National Science Museum, Tokyo-V). In: J. A. Grant Mackie, K. Masuda, K. Mori and K. Ogasawara (Eds.). *Professor Tamio Kotaka commemorative volume on Molluscan paleontology*. Sendai, Japan, Saito Ho-On Kai Museum. 2: 567-587.
- Matsumoto, E. (1971). Oligocene Molluscs from the Setogawa Group in Central Japan. *Bulletin of the National Science Museum, Tokyo = Kokuritsu Kagaku Hakubutsukan Kenkyu Hokoku* 14(4): 661-669.

Maurer, D. (1967). Filtering experiments on marine pelecypods from Tomales Bay, California. *The Veliger* 9(3): 305-309

Maurer, D. (1967). Burial experiments on marine pelecypods from Tomales Bay, California. *The Veliger* 9(4): 376-381

Maurer, D. (1967). Mode of feeding and diet, and synthesis of studies on marine pelecypods from Tomales Bay, California. *The Veliger* 10(1): 72-76

Maurer, D. (1969). Pelecypod-sediment association in Tomales Bay, California. *The Veliger* 11(3): 243-249

McHugh, J. L. and M. W. Sumner (1988). Annotated bibliography II of the hard clam *Mercenaria mercenaria*. United States, Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Technical Report NMFS-68: 59 pp.

McHugh, J. L., M. W. Sumner, P. J. Flagg, D. W. Tipton, W. J. Behrens (1982). Annotated bibliography of the hard clam (*Mercenaria mercenaria*). *United States, Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Technical Report* SSRF-756: 845 pp.

Menzel, R. W. (1970). The species and distribution of quahog clams *Mercenaria* [abst.]. *Proceedings of the National Shellfisheries Association* 60: 8

Menzel, R. W. (1989). The biology, fishery and culture of quahog clams, *Mercenaria*. pp. 201-242, in: J. J. Manzi and Michael Castagna, *Clam culture in North America*. Amsterdam, etc. (Elsevier). x + 461 pp.

Menzel, R. W. and M. Y. Menzel (1965). Chromosomes of two species of quahog clams and their hybrids. *Biological Bulletin* 129(1): 181-188

Milera, J. F. (1982). Localizacion de *Pitar (Hysteroconcha) dione* (Linneo 1758) (Mollusca: Eulamelibranchia: Veneridae) en Cuba. *Miscelanea Zoologica* 14: 1-2.

Miller, M. B., C. H. Hanson, K. K. Chew (1975). Shell growth of butter clams and littleneck clams near Madrona Beach on Camano Island [abst.]. *Proceedings of the National Shellfisheries Association* 65: 10

Moisescu, V., M. Chivu, V. Dragu, E. Margarit (1979). Studiul faunei de moluste egeriene din Bazinul Petrosani. *Memoires - Institut de Geologie et de Geophysique* 29: 65-120.

Monterosato, M. d. (1870). Sulla scoperta del genere Dacrydium nel Mediterraneo. *Bullettino Malacologico Italiano*, Pisa 3(2): 43-46.

Monti, D., L. Frenkiel, M. Moueza (1991). Demography and growth of Anomalocardia brasiliiana (Gmelin) (Bivalvia: Veneridae) in a mangrove, in Guadeloupe (French West Indies). *Journal of Molluscan Studies* 57(2): 249-257.

Monti, D., S. Salvidio, N. Pasteur, M. Amanieu (1986). Modification alternative rapide du polymorphisme des malatedehydrogenases dans un population de palourdes de l'etang de Thau (Ruditapes decussatus, Bivalvia: Veneridae). *Comptes Rendus de l'Academie des Sciences Serie III Sciences de la Vie* 302(14): 513-518.

Moore, H. B. and N. N. López (1969). The ecology of Chione cancellata. *Bulletin of Marine Science* 19(1): 131-148

Moore, H. B. and N. N. López (1975). An additional study of the ecology of Chione cancellata L. *Bulletin of Marine Science* 25(1): 126-130

Mora de la Torre, A. and F. Correa Sandoval (1996). Estructura genética poblacional de Chione cortezi (Bivalvia), in el alto Golfo de California [abst.]. *Sociedad Mexicana de Genética y Sociedad Mexicana de Toxicología Genética, Congreso Conjunto* 4:32

Mörch, O. A. L. (1853). *Acephala. Annulata. Cirripedia. Echinodermata. Catalogus Conchyliorum*. D. A. D. A. G. C. d. Yoldi. Fasciculus Secundus: 1-76.

Mörch, O. A. L. (1861). Beiträge zur Molluskenfauna Central-Amerika's. *Malakozoologische Blätter* 7: 170-213.

Moroz, S. A. and L. I. Jermokhina (1991). Vikopni molyuski paleotsenu Povolzhya. *Paleontologicheskiy Sbornik* (L'vov) 28: 8-17.

Morton, B. (1983). Aspects of the biology and functional morphology of Irus irus (Bivalvia: Veneridae: Tapetinae) with a comparison of Bassina calophylla (Chioninae). In: Morton, B. and Dudgeon, D. (Eds). *The malacofauna of Hong Kong and southern China*. 2: 6-24.

Morton, B. (1985). Aspects of the biology and functional morphology of Irus irus (Bivalvia: Veneridae: Tapetinae) with a comparison of Bassina calophylla (Chioninae). pp. 321-336, in: Brian S. Morton and D. Dudgeon, eds., *The malacofauna of Hong Kong and southern China*. II, Vol. 1. Hong Kong (Hong Kong University). viii + 361 pp.

Morton, B. (1990). The life cycle and sexual strategy of Gafrarium pectinatum (Bivalvia: Veneridae) in a Hong Kong mangrove. *Malacological Review* 23(1-2): 53-62.

- Morton, B. (2000). The Anatomy of *Callocardia hungerfordi* (Bivalvia: Veneridae) and the origin of its shell camouflage. *Journal of Molluscan Studies* 66: 21-30.
- Moueza, M., O. Gros, L. Frenkiel (1999). Embryonic, larval and postlarval development of the tropical clam, *Anomalocardia brasiliiana* (Bivalvia, Veneridae). *Journal of Molluscan Studies* 65(1): 73-88.
- Muralidharan, G. and P. Vivek-Raja (1997). Trace element concentration in the meat of the edible clam *Marcia recens* Chemnitz (Pelecypoda: Veneridae). *Indian Journal of Marine Sciences* 26(4): 383-385.
- Murphy, R. C. (1985). Factors affecting the distribution of the introduced bivalve, *Mercenaria mercenaria*, in a California lagoon -- the importance of bioturbation. *Journal of Marine Research* 43(3): 673-692
- Nakazima, M. (1958). On the differentiation of the crenated-folds in the midgut-gland of Eulamellibranchia (II). *Venus* 20(1): 101-109, pls. 4-6
- Nakazima, M. (1959). On the differentiation of the crenated-folds in the midgut-gland of Eulamellibranchia III. Various developmental stages of *Meretrix lusoria*. *Venus* 20(4): 356-360
- Narchi, W. (1971). The presence of byssus in adults of *Transennella tantilla* (Gould) (Bivalvia: Veneridae). *Wasmann Journal of Biology* 28(2): 233-236
- Narchi, W. (1972). Structure and adaptation in *Transennella tantilla* (Gould) and *Gemma gemma* (Totten) (Bivalvia: Veneridae). *Bulletin of Marine Science* 21(4): 866-885
- Narchi, W. (1972). Comparative study of the functional morphology of *Anomalocardia brasiliiana* (Gmelin, 1791) and *Tivela mactroides* (Born, 1778) (Bivalvia, Veneridae). *Bulletin of Marine Science* 22(3): 643-670
- Narchi, W. (1980). On the biology of *Veremolpa scabra* (Hanley 1845) (Bivalvia: Veneridae) from the South China Sea. In: Morton, B. (Ed.) *The malacofauna of Hong Kong and southern China*. Hong Kong University Press for the Department of Zoology, University of Hong Kong, Hong Kong.
- Narchi, W. (1980). Sobre *Veremolpa scabra* (Hanley, 1845) (Bivalvia, Veneridae). *Revista Nordestina de Biologia (especial)*: 233.
- Narchi, W. and M. A. Gabrieli (1981). Sobre anatomia funcional de *Chione subrostrata* (Lamarck, 1818). *Revista Nordestina de Biologia* 3(Espec.): 25-46

- Nayak, K. K. and R. M. Badve (1985). Bivalvia from Nimar Sandstone, Bagh Beds, Jhabua District, Madhya Pradesh. *Biovigyanam* 11(1): 77-86.
- Neave, F. (1942). The butter-clam (*Saxidomus giganteus* Deshayes): studies in productivity. British Columbia, *Provincial Fisheries Department, Report for 1941*: J79-81
- Neave, F. (1944). The spread of the Japanese little-neck clam in British Columbia waters. Canada, *Fisheries Research Board of Canada, Pacific Coast Stations, Progress Reports* 61: 3
- Neave, F. (1944). The legal size-limit in relation to the size at which butter clams mature. Canada, *Fisheries Research Board of Canada, Pacific Coast Stations, Progress Reports* 61: 4-5
- Neave, F. (1946). Conditions of the butter-clam fishery in British Columbia. *British Columbia, Department of Fisheries, Report for 1945*: N67-74
- Nevesskaya, L. A., I. A. Goncharova, N. P. Paramonova, S. V. Popov, E. V. Babak, K. G. Bagdasaryan, A. A. Voronina (1993). [A key to the Miocene bivalve molluscs of South-Western Eurasia.]. *Trudy Paleontologicheskogo Instituta Akademii Nauk SSSR* 247: 1-412.
- Nichols, F. H. and J. K. Thompson (1985). Persistence of an introduced mudflat community in south San Francisco Bay, California. *Marine Ecology -- Progress Series* 24(1/2): 83-97
- Nickerson, R. B. (1977). A study of the littleneck clam (*Protothaca staminea* Conrad) and the butter clam (*Saxidomus giganteus* Deshayes) in a habitat permitting coexistence, Prince William Sound, Alaska. *Proceedings of the National Shellfisheries Association* 67: 85-102
- Nicol, D. and S. Jones Douglas (1984). *Chione (Chione) craspedonia* Dall in the Crystal River Formation (Eocene) in peninsular Florida. *Tulane Studies in Geology and Paleontology* 18(1-2): 73-75.
- Nicolay, K. (1979). Synoptic tables of Mediterranean and European shells. Superfam. Veneroidea, fam. Veneridae, gen. *Callista*, *Irus*, *Pitar*, *Dosinia*. *Conchiglia* 11(126-127): 20-21.
- Nicolay, K. (1979). Synoptic tables of Mediterranean and European shells. Superfamily Veneroidea. Family Veneridae, genera *Venerupis*, *Paphia*. *Conchiglia* 11(128-129).
- Nicolay, K. (1980). Synoptic tables of Mediterranean and European shells. *Conchiglia* 12(130-131): 20-21.

- Nistratova, S. N. (1981). Mechanisms of adaptation of the cardiac muscle in molluscs to nervous influences. *Pacific Science Congress Proceedings* 14(3): 161-169.
- Nomura, S. (1937). On some Recent venerid mollusks from northeast Honshû, Japan. *Research Bulletin of the Saito Ho-on Kai Museum* 13: 7-10, pl. 3
- Nomura, S. (1940). *Clementia (Compsomyax) iizukai* (Yokoyama). *Research Bulletin of the Saito Ho-on Kai Museum* 19: 13-16, pl. 1
- Norman, A. M. (1860). The Mollusca of the Firth of Clyde. *The Zoolologist* 16.
- Nosho, T. Y. and K. K. Chew (1972). The setting and growth of the Manila clam, *Venerupis japonica* (Deshayes), in Hood Canal, Washington. *Proceedings of the National Shellfisheries Association* 62: 50-58
- Nybakken, J. and M. D. Stephenson (1975). Effects of engineering activities on the ecology of Pismo clams. United States, *Army Corps of Engineers, Coastal Engineering Research Center, Miscellaneous Paper* 8-75: 65 pp
- Ohba, S. (1959). Ecological studies in the natural population of a clam, *Tapes japonica*, with special reference to seasonal variations in size and structure of the population and to individual growth. *Biological Journal of the Okayama University* 5(1-2): 13-42
- Ohno, T. (1996). Intra-periostracal calcified needles of the bivalve family Veneridae. *Bulletin De L'Institut Oceanographique Numero Special* 14(4): 305-314.
- Ohta, N., S. Ode, K. Tomura, M. Omori (1980). Relative abundance of lanthanide in recent and fossil shells. In: Omori, M. and Watabe, N. *The mechanisms of biomineralization in animals and plants*. Proceedings 3rd International Biomineralization Symposium. 1-310
- Oliver, W. R. B. (1923). Notes on New Zealand Pelecypods. *Proceedings of the Malacological Society* 15: 179-188.
- Olsson, A. (1961). *Mollusks of the Tropical Eastern Pacific*. Ithaca, N.Y., Paleontological Research Institution.
- Olsson, A. A. (1932). Contributions to the Tertiary Paleontology of Northern Peru: Part 5. The Peruvian Miocene. *Bulletin of American Paleontology* 19(68): 1-241.
- Olsson, A. A. (1942). Tertiary and Quarternary Fossils from the Burica Peninsula of Panama and Costa Rica. *Bulletin of American Paleontology* 27(106): 1-97.

Olsson, A. A. (1944). Contributions to the Paleontology of Northern Peru. Part VII. The Cretaceous of the Paita Region. *Bulletin of American Paleontology* 28(111): 1-127.

Ortlieb, L., T. DeVries, A. Diaz (1990). Ocurrencia de Chione broggi (Pilsbry & Olsson, 1943) (Pelecypoda) en depositos litorales cuaternarios del sur del Peru; implicaciones paleoceanograficas. *Boletin de la Sociedad Geologica del Peru* 81: 127-134.

Osorio, C., D. Frassinetti, E. Bustos (1983). Taxonomia y morfometria de Venus antiqua antiqua King y Broderip, 1835 (Mollusca, Bivalvia, Veneridae). *Tethys* 11(1): 49-56.

Osorio, R. C., P. V. Bustos, R. E. Bustos (1987). Contenido gastrico de la almeja (Venus antiqua antiqua King y Broderip, 1835) en Ancud, Chile (Mollusca: Veneridae). *Investigacion Pesquera* 34: 139-147.

Otto, S. V. (1973). Hermaphroditism in two species of pelecypod mollusks. *Proceedings of the National Shellfisheries Association* 63: 96-98

Owen, S. (1992). Pismo clams rebound after decade of decline. *Outdoor California* 53(4): 1-4

Oyama, K. (1986). On Periglypta magnifica (Sowerby). *Chiribotan* 17(1): 7-8.

Padilla, M. (1984). Desintegracion neurohormonal 'in vitro' de la vesicula germinal del oocito de Venus antiqua antiqua (Mollusca: Pelecypoda: Veneridae). *Revista De Biologia Marina* 20(1): 39-60.

Palacios, J. A., J. A. Rodriguez, R. A. Cruz, J. Bolanos (1986). Estudio sobre la biologia de Protothaca asperrima (Pelecypoda: Veneridae) 1. Distribucion y cuantificacion. *Brenesia*. pp. 1-11

Palacios, J. A., J. Bolanos, J. A. Rodriguez, R. A. Cruz (1986). Estudio sobre la biologia de Protothaca asperrima (Pelecypoda: Veneridae) 2. Estructura poblacional. *Brenesia*. pp. 13-22

Palacios, J., R. A. Cruz, J. Bolanos, J. A. Rodriguez (1986). Estudio sobre la biologia de Protothaca asperrima (Pelecypoda: Veneridae) 3. Ciclo reproductivo. *Brenesia*. pp. 23-32

Palmer, K. V. W. (1927). The Veneridae of eastern America; Cenozoic and recent. *Palaeontographica Americana* 1(5): 209-428.

Pamatmat, M. M. (1969). Seasonal respiration of Transennella tantilla Gould. *American Zoologist* 9(2): 419-426

- Parker, P. (1949). Fossil and Recent species of the pelecypod genera *Chione* and *Securella* from the Pacific coast. *Journal of Paleontology* 23(6): 577-593, pls. 89-95
- Parker-Schnitzler, M. (2000). Population genetics of *Protothaca staminea* and *Macoma balthica* in Puget Sound, WA. [abst.]. *Journal of Shellfish Research* 19(2)
- Passamonti, M., B. Mantovani, V. Scali (1997). Allozymic characterization and genetic relationships among four species of Tapetinae (Bivalvia, Veneridae). *Italian Journal of Zoology* 64(2): 117-124.
- Passamonti, M., B. Mantovani, V. Scali (1998). Characterization of a highly repeated DNA family in Tapetinae species (Mollusca Bivalvia: Veneridae). *Zoological Science* 15(4): 599-605.
- Passamonti, M., B. Mantovani, V. Scali (1999). Allozymic analysis of some Mediterranean Veneridae (Mollusca: Bivalvia): Preliminary notes on taxonomy and systematics of the family. *Journal of the Marine Biological Association of the United Kingdom* 79(5): 899-906.
- Paul, A. J. and H. M. Feder (1973). Growth, recruitment, and distribution of the littleneck clam, *Protothaca staminea*, in Galena Bay, Prince William Sound, Alaska. United States, Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, *Fishery Bulletin* 71(3): 665-677
- Paul, A. J., J. McDonald Paul, H. M. Feder (1976). Recruitment and growth in the bivalve *Protothaca staminea*, at Olson Bay, Prince William Sound, ten years after the 1964 earthquake. *The Veliger* 18(4): 385-392
- Paul, A. J., J. McDonald Paul, H. M. Feder (1976). Age, growth, and recruitment of the butter clam, *Saxidomus giganteus*, on Porpoise Island, southeast Alaska. *Proceedings of the National Shellfisheries Association* 66: 26-28
- Paul, A. J., J. McDonald Paul, H. M. Feder (1976). Growth of the littleneck clam, *Protothaca staminea*, on Porpoise Island, southeast Alaska. *The Veliger* 19(2): 163-166
- Perkins, G. H. (1871). The molluscan fauna of New Haven. Part II. Acephala and Bryozoa. *Proceedings of the Boston Society of Natural History* 13: 139-164.
- Peterson, C. H. (1977). Competitive organization of the soft-bottom macrobenthic communities of southern California. *Marine Biology* 43(4): 343-359

Peterson, C. H. (1982). The importance of predation and intra- and interspecific competition in the population biology of two infaunal suspension-feeding bivalves, *Protothaca staminea* and *Chione undatella*. *Ecological Monographs* 52(4): 437-475

Peterson, C. H. (1983). Interactions between two infaunal bivalves, *Chione undatella* (Sowerby) and *Protothaca staminea* (Conrad), and two potential enemies, *Crepidula onyx* Sowerby and *Cancer anthonyi* (Rathbun). *Journal of Experimental Marine Biology and Ecology* 68(2): 145-158

Peterson, C. H. and W. G. Ambrose, Jr. (1958). Potential habitat dependence in deposition rate of presumptive annual lines in shells of the bivalve *Protothaca staminea*. *Lethaia* 18(3): 257-260

Peterson, C. H. and M. L. Quammen (1982). Siphon nipping: its importance to small fishes and its impact on growth of the bivalve *Protothaca staminea* (Conrad). *Journal of Experimental Marine Biology and Ecology* 63(3): 249-268

Pezant, A. (1908). Mollusques fossiles de Monneville (Oise). *La Feuille des Jeunes Naturalistes* 4(38): 224-229.

Pfeiffer, L. G. C. L. (1868). Studien über die Familie der Veneraceen oder Venusmuscheln. *Malakozoologische Blätter* 15(4): 141-155

Pfeiffer, L. G. C. L. (1869). Ueber die Gattung *Clementia* Gray. *Malakozoologische Blätter* 16(5): 190-194

Philippi, R. A. (1843-1849). [*Monograph of Venus*. Abbildungen und Beschreibungen neuer oder wenig gekannter Conchylien 1(2): 39-42, pl. 1 (March 1843); 1(5): 127-129, pl. 2 (April 1844); 1(7): 175-178, pl. 3 (Nov. 1844); 2(2): 61-63, pl. 4 (Nov. 1845); 2(4): 107-110, pl. 5 (Aug. 1846); 2(5): 151-152, pl. 6 (Oct. 1846); 3(3): 75-78, pl. 7 (Feb. 1848); 3(4): 21-25, pl. 8 (Jan. 1849); 3(5): 59-61, pl. 9 (March 1849); 3(6): 81-84, pl. 10 (Oct. 1849) [also numbered pp. 1-40]

Philippi, R. A. (1844-1848). [*Monograph of Cytherea*. Abbildungen und Beschreibungen neuer oder wenig gekannter Conchylien 1(6): 149-150, pl. 1 (July 1844); 1(7): 169-173, pl. 2 (Oct. 1844); 1(8): 197-200, pl. 3 (Jan. 1845); 2(4): 95-97, pl. 4 (Aug. 1846); 2(6): 179-182, pl. 5 (Feb. 1847); 2(8): 299-230, pl. 6 (April 1847); 2(8): 231, pl. 7 (April 1847); 3(1): 23-25, pl. 8 (Sept. 1847); 3(3): 71-74, pl. 9 (Feb. 1848) [also numbered pp. 1-42]

Philippi, R. A. (1849-1851). Centuria quarta testaceorum novorum. *Zeitschrift für Malakozoologie* 6(2): 27-32 (May 1849); 8(2): 29-32 (June 1851); (3): 39-48 (July); (4): 49-64 (15 July); (5): 65-74 (July)

- Picariello, O. and M. Paolucci (1983). Comparative electrophoretic analysis of esterase molecular forms in the digestive gland of some venerids (Mollusca: Bivalvia). *Rendiconto dell'Accademia delle Scienze Fisiche e Matematiche Serie IV* 56(2): 381-387.
- Pilsbry, H. A. (1922). *Gemma gemma purpurea*. *The Nautilus* 36(1): 32
- Pizarro, J. F. and R. A. Cruz (1987). Ciclo reproductivo de la almeja *Protothaca grata* (Pelecypoda: Veneridae). *Brenesia* 27: 23-34.
- Pochon-Masson, J. and I.-D. Gharagozlov (1970). Particularité morphologique de l'acrosome dans le spermatozoïde de *Tapes decussatus* L. (mollusque lamellibranche). *Annales des Sciences Naturelles (Zoologie)* (12)12(2): 171-180, 2 pls.
- Polakovic, G. (2000). Small clams court. *Los Angeles Times, Ventura County News* July 17, 2000: B1-2
- Pollock, L.-W. (1998). *A practical guide to the marine animals of northeastern North America*. Rutgers University Press, New Brunswick, New Jersey and London. pp. 1-367
- Ponurovsky, S. K. and Y. M. Yakovlev (1992). The reproductive biology of the Japanese littleneck, *Tapes philippinarum* (A. Adams and Reeve, 1850) (Bivalvia: Veneridae). *Journal of Shellfish Research* 11(2): 265-277.
- Popov, S. V. (1986). Composite prismatic structure in bivalve shell. *Acta Palaeontologica Polonica* 31(1-2): 3-26.
- Popov, S. V. (1986). [Microstructure of bivalve shells from the family Veneridae.]. In: Kafanov, A.I. (Ed.) [*Palaeogene Neogene Bivalvia of the Far East and eastern Paratethys. Collection of scientific works.*] Akademiya Nauk SSSR, Vladivostok. pp. 1-132
- Popov, S. V. (1992). [The shell microstructure of several groups of bivalve molluscs.]. *Trudy Paleontologicheskogo Instituta Akademii Nauk SSSR* 245: 1-46.
- Porter, H. J. (1967). Seasonal gonadal changes of adult clams, *Mercenaria mercenaria* (L.), in North Carolina. *Proceedings of the National Shellfisheries Association* 55: 35-52
- Poutiers, J. M. (1978). Introduction a l'étude faunistique des bivalves du littoral français: les espèces marines du golfe d'Aigues Mortes. Deuxième partie. *Notes et Contributions centre d'études et de recherches de Paleontologie Biostratigraphique* No. 15(2): 290-563.

Poutiers, J. M. (1978). *Introduction à l'étude faunistique des Bivalves du littoral français; les espèces marines du golfe d'Aigues Mortes*. Orsay, France, Université de Paris Sud CERPAB. pp. 1- 272.

Pratt, D. M. (1953). Abundance and growth of *Venus mercenaria* and *Calloocardia morrhuanus* in relation to the character of bottom sediments. *Journal of Marine Research* 12(1): 60-74

Pratt, D. M. and D. A. Campbell (1956). Environmental factors affecting growth in *Venus mercenaria*. *Limnology and Oceanography* 1(1): 2-17

Quayle, D. B. (1938). *Paphia bifurcata*, a new molluscan species from Ladysmith Harbour, B. C. Canada, *Journal of the Fisheries Research Board of Canada* 4(1): 53-54

Quayle, D. B. (1939). Note re *Paphia bifurcata*, a new molluscan species from Ladysmith Harbour, B. C. *The Nautilus* 52(4): 139-140

Quayle, D. B. (1941). The Japanese "little neck" clam accidentally introduced into British Columbia waters. Canada, *Progress Reports of the Fisheries Research Board of Canada, Pacific Coast Stations* 48: 17-18

Quayle, D. B. (1943). Sex, gonad development and seasonal gonad changes in *Paphia staminea* Conrad. Canada, *Journal of the Fisheries Research Board of Canada* 6(2): 140-151

Quayle, D. B. (1949). Movements in *Venerupis (Paphia) pullastra* (Montagu). *Proceedings of the Malacological Society of London* 28(1): 31-37

Quayle, D. B. (1952). The relation of digging frequency to productivity of the butter clam, *Saxidomus giganteus* Deshayes. Canada, *Journal of the Fisheries Research Board of Canada* 8(5): 369-373

Quayle, D. B. (1952). Structure and biology of the larva and spat of *Venerupis pullastra* (Montagu). *Transactions of the Royal Society of Edinburgh* 62(1): 255-297, 1 pl.

Rajan, A., N. Krishnaveni, et al. (1986). Seasonal variation in the concentration of copper, zinc, manganese and iron in the clam *Katelysia opima* (Gmelin) (Mollusca: Bivalvia, Veneridae) collected from Vellar Estuary, S. India. *Salem Institute of Experimental Biology Memoir No 2*: 39-48.

Ramon, M. and C. A. Richardson (1992). Age determination and shell growth of *Chamelea gallina* (Bivalvia: Veneridae) in the western Mediterranean. *Marine Ecology Progress Series* 89(1): 15-23.

Récluz, C. A. (1846). De la famille des lithophages de Lamarck et des genres qui la composent. *Société Cuvierienne, Revue Zoologique* 9(11): 405-425

Reeve, L. A. (1850). Description of a new species of *Cytherea*. *Annals and Magazine of Natural History* (2)6(34): 291

Reeve, L. A. (1863-1864). Monograph of the genus *Dione*. In: L. A. Reeve, ed., *Conchologia iconica*; or, illustrations of the shells of molluscous animals 14: 12 pls. (pls. 2-8, Aug. 1863; 9-12, Oct.; 1, Feb. 1864)

Reeve, L. A. (1863-1864). Monograph of the genus *Venus*. In: L. A. Reeve, ed., *Conchologia iconica*; or, illustrations of the shells of molluscous animals 14: 26 pls. (pls. 2-11, April 1863; 12-23, June; 1, Feb. 1864; 24-26, March)

Reeve, L. A. (1864). Monograph of the genus *Tapes*. In: L. A. Reeve, ed., *Conchologia iconica*; or, illustrations of the shells of molluscous animals 14: 13 pls. (pls. 2-6, Jan.; 7-11, Feb.; 1, 12, 13, March)

Rehder, H. A. (1939). New marine mollusks from the west Atlantic. *The Nautilus* 53(1): 16-21, pl. 6

Rehder, H. A. (1943). New marine mollusks from the Antillean Region. *Proceedings of the United States National Museum* 93(3161): 187-203.

Rice, T. R. and R. J. Smith (1958). Filtering rates of the hard clam (*Venus mercenaria*) determined with radioactive phytoplankton. United States, *Department of the Interior, Fish and Wildlife Service, Fishery Bulletin* 58(129): 82 pp.

Roberts, D. (1981). The effects of substrate and shell form on burrowing in the genus *Katelysia* (Bivalvia: Veneridae). *Journal of the Malacological Society of Australia* 5(1-2): 33-40.

Roberts, D. (1984). The genus *Katelysia* (Bivalvia: Veneridae) in southern Australia. *Journal of the Malacological Society of Australia* 6(3-4): 191-204.

Robinson, A. M. and W. P. Breese (1982). The spawning season of four species of clams in Oregon. *Journal of Shellfish Research* 2(1): 55-57

Robinson, A. M. and W. P. Breese (1984). Gonadal development and hatchery rearing techniques for the Manila clam *Tapes philippinarum* (Adams & Reeve). *Journal of Shellfish Research* 4(2): 161-163

Robinson, W. E. (1983). Quantification of ingestion by *Mercenaria mercenaria* (L.) veligers feeding on mixed suspensions of inert material and algae using microspectrofluorimetry. *Journal of Molluscan Studies, Supplement 12A*: 167-171

Robinson, W. E. and R. W. Langton. (1980). Digestion in a subtidal population of *Mercenaria mercenaria* (Bivalvia). *Marine Biology* 8(3): 173-179

Roesijadi, G. (1980). Influence of copper on the clam *Protothaca staminea*: effects on gills and occurrence of copper-binding proteins. *Biological Bulletin* 158(2): 233-247

Rodriguez-Juiz, A. M., M. Torrado, et al. (1996). Genome-size variation in bivalve molluscs determined by flow cytometry. *Marine Biology* 126(3): 489-497.

Römer, E. (1857). *Kritische Untersuchung der Arten des Molluskengeschlechts Venus bei Linné und Gmelin mit Berücksichtigung der später beschriebenen Arten*. Cassel (Luckhardt). xiii + 135 + 1 pp.

Römer, E. (1861). Kritische Übersicht sämmtlicher bis jetzt bekannten Arten aus der Cythereengruppe *Tivela* Lam. *Malakozoologische Blätter* 8(1): 17-28

Römer, E. (1861). Über *Saxidomus*. *Malakozoologische Blätter* 8(2): 63-70

Römer, E. (1861). Beschreibung neuer *Venus*-Arten. *Malakozoologische Blätter* 7: 148-165.

Römer, E. (1862). Kritische Übersicht der Arten aus der Cythereengruppe *Caryatis* Römer (*Pitar olim*). *Malakozoologische Blätter* 9: 58-86.

Römer, E. (1864). Kritische Übersicht sämmtlicher Arten der zur Gattung *Venus* gehörenden Untergattung *Tapes*. *Malakozoologische Blätter* 11: 19-44.

Römer, E. (1864). Kritische Übersicht sämmtlicher *Tapes*-Arten. *Malakozoologische Blätter* 11: 58-98.

Römer, E. (1864-1869). *Monographie der Molluskengattung Venus*, Linné. I. Band. Subgenus *Cytherea* Lamarck. Novitates Conchologicae, Supplement 3. Cassel (Fischer). 219 pp., 59 pls. (Pp. 1-12, pls. 1-3, June 1864; 13-24, pls. 4-6, Jan. 1865; 25-32, pls. 7-9, June 1865; 33-42, pls. 10-12, April 1866; 43-58, pls. 13-18, Sept. 1866; 59-78, pls. 19-21, Feb. 1867; 79-102, pls. 22-27, Sept. 1867; 103-126, pls. 28-33, Dec. 1867; 127-146, pls. 34-39, Sept. 1868; 147-172, pls. 40-46, Nov. 1868; 173-190, pls. 47-50, Feb. 1869; 191-198, June 1869; 199-206, pls. 51-59, July 1869; 207-217, Oct. 1869)

Römer, E. (1865). Kritische Übersicht sämmtlicher Arten der zur Gattung Venus gehörenden Untergattung Mercenaria und Gemma. *Malakozoologische Blätter* 12(3): 134-141

Römer, E. (1867). Kritische Übersicht aller zum Subgenus Chione gehörenden Arten von Venus. Teil 1. *Malakozoologische Blätter* 14: 28-62.

Römer, E. (1867). Kritische Übersicht aller zum Subgenus Chione gehörenden Arten von Venus. Teil 2. *Malakozoologische Blätter* 14: 92-126.

Römer, E. (1869-1872). *Monographie der Molluskengattung Venus*, Linné. II. Band. Subgenus Sunetta Link. Subgenus Tapes Megerle von Mühlfeld. Novitates Conchologicae, Supplement 3. Cassel (Fischer). 126 pp., 40 pls. (pl. 1, Oct. 1869; pp. 1-16, pls. 2-16, Feb. 1870; 17-32, April 1870; 33-48, June 1870; 49-64, pls. 17-21, Oct. 1870; 65-80, pls. 22-26, Jan. 1871; 81-96, pls. 27-31, July 1871; 97-112, pls. 32-35, Jan. 1872; 113-128, pls. 36-40, June 1872 [all completed; ends in mid-sentence])

Römer, E. (1870). Beschreibung neuer Arten von Dosinia und Tapes. *Malakozoologische Blätter* 17(1): 1-11

Roopnarine, P. D. (1998). Translating trees into taxonomy within Veneridae (Bivalvia): a reply to Harte. *Malacologia* 39(1-2): 221-224.

Roopnarine, P. D. (1995). A re-evaluation of evolutionary stasis between the bivalve species Chione erosa and Chione cancellata (Bivalvia: Veneridae). *Journal of Paleontology* 69(2): 280-287.

Roopnarine, P. D. (1996). Systematics, biogeography and extinction of chionine bivalves (Bivalvia: Veneridae) in tropical America: Early Oligocene-Recent. *Malacologia* 38(1-2): 103-142.

Roopnarine, P. D. (1997). Endemism and extinction of a new genus of chionine (Veneridae: Chioninae) bivalve from the Late Neogene of Venezuela. *Journal of Paleontology* 71(6): 1039-1046.

Roopnarine, P. D. and G. J. Vermeij (2000). One species becomes two: the case of Chione cancellata, the resurrected C. elevata, and a phylogenetic analysis of Chione. *Journal of Molluscan Studies* 66(4): 517-534

Roth, B. (1975). What is Pitaria ida Tegland? *The Veliger* 17(4): 389-392, 1 pl.

Russell, M. P. and J. P. Huelsenbeck (1989). Seasonal variation in brood structure of Transennella confusa (Bivalvia: Veneridae). *The Veliger* 32(3): 288-295.

- Rybakov, A. V. (1983). Partenity i lichinki trematod dvustvorchatogo molliuska *Ruditapes philippinarum* v Zalivie Petra Velikogo Iaponskogo Moria. [Reproduction and larvae of a trematode in the bivalve mollusk *Ruditapes philippinarum* in the Bay of Peter the Great and the Japanese Sea]. *Biologija Moria* 9(1): 12-20 [transl.: Soviet Journal of Marine Biology 9(1): 9-16]
- Salas, C. and S. Gofas (1998). Description of four new species of *Neolepton* Monterosato, 1875 (Mollusca: Bivalvia: Neoleptonidae), with comments on the genus and on its affinity with the Veneracea. *Ophelia* 48(1): 35-70.
- Salchak, A. and J. Haas (1971). Occurrence of the northern quahog, *Mercenaria mercenaria*, in Colorado Lagoon, Long Beach, California. *California Fish and Game* 57(2): 126-128
- Santulli, A., N. P. Wilkins, et al. (1992). Two tissue-specific loci for octopine dehydrogenase in *Talpes decussatus* (Bivalvia, Veneridae). *Comparative Biochemistry And Physiology B Comparative Biochemistry* 102(2): 409-411.
- Sato, S. (1994). Analysis of the relationship between growth and sexual maturation in *Phacosoma japonicum* (Bivalvia: Veneridae). *Marine Biology* 118(4): 663-672.
- Sato, S. i. (1996). Genetic variability and population structure of *Phacosoma japonicum* (Bivalvia: Veneridae). *Venus, The Japanese Journal of Malacology* 55(1): 51-63.
- Sato, S. i. (1999). Genetic and environmental control of growth and reproduction of *Phacosoma japonicum* (Bivalvia: Veneridae). *Veliger* 42(1): 54-61.
- Saul, L. R. (1973). *Parallel evolution and continental drift*. Cordilleran Section, 69th Annual Meeting. Boulder, CO, United States, Geological Society of America (GSA). 5: 99-100.
- Saul, L. R. (1991). What do we know about Cretaceous Pacific slope venerid bivalves? *Western Society of Malacologists Annual Report* 24(8): 3-7.
- Saul, L. R. (1993). Pacific slope Cretaceous bivalves; eight venerid species. *Journal of Paleontology* 67(6): 965-979.
- Saul, L. R. and W. P. Popenoe (1992). Pacific Slope Cretaceous Bivalves of the Genus *Calva*. *Contributions in Science* 433: 1-68.
- Savage, N. B. (1976). Burrowing activity in *Mercenaria mercenaria* (L.) and *Spisula solidissima* (Dillwyn) as a function of temperature and dissolved oxygen. *Marine Behaviour and Physiology* 3(4): 221-234

Savitskii, V. O. (1979). Kainzoiskie *Liocyma hokkaidoensis* Habe Dal'nego Vostoka. [Cenozoic *Liocyma hokkaidoensis* Habe of the Far East]. pp. 75-80, in: A. I. Ponomareva, ed., *Iskopaemye molliuski Dal'nego Vostoka i ikh stratigraficheskoe znachenie*. [Fossil mollusks of the Far East and their stratigraphic significance]. Akademii Nauk SSSR, Dalnevostochnyi Nauchnyi Tsentr, Dalnevostochnyi Geologischeskii Institut. 114 pp., 18 pls.

Schmidt, R. R. and J. E. Warme (1969). Population characteristics of *Protothaca staminea* (Conrad) from Mugu Lagoon, California. *The Veliger* 12(2): 193-199

Schultz, O. (1978). Veneridae. In: A. Papp, I. Cicha, J. Senes and F. Steininger (Eds.). *Chronostratigraphie und Neostratotypen; Miozaen der Zentralen Paratethys; M4--Badenien (Moravien, Wielicien, Kosovien)*. Bratislava, Czechoslovakia, VEDA Verl. Slow. Akad. Wiss.: 352-358.

Sellmer, G. P. (1967). Functional morphology and ecological life history of the gem clam, *Gemma gemma* (Eulamellibranchia: Veneridae). *Malacologia* 5(2): 137-223

Semper, O. (1862). Du genre *Woodia*, Deshayes. *Journal de Conchyliologie* 10: 141-146.

Shaw, R. F. (1956). The polymorphism of the Japanese littleneck clam. *The Nautilus* 70(2): 53-59

Shaw, W. N. (1986). Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest). Common littleneck clam. United States, *Department of the Interior, Fish and Wildlife Service, Biological Report* 82(11.46): vi + 11 pp.

Shaw, W. N. and T. J. Hassler (1989). Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest). Pismo clam. United States, *Department of the Interior, Fish and Wildlife Service, Biological Report* 82(11.95): 12 pp.

Shimamoto, M. (1986). Shell microstructure of the Veneridae (Bivalvia) and its phylogenetic implications. *Science Reports of the Tohoku University. Series 2: Geology = Tohoku Daigaku Rika Hokoku. Dai 2: Shu Chishitsugaku* 56(1): 1-39.

Shimamoto, M. (1988). Shell microstructure in some genera of the Veneridae (Bivalvia); coexistence of different shell microstructure types within a single shell layer. In: J. A. Grant Mackie, K. Masuda, K. Mori and K. Ogasawara (Eds.). *Professor Tamio Kotaka commemorative volume on Molluscan paleontology*. Sendai, Japan, Saito Ho-On Kai Museum. 2: 239-255.

Shimamoto, M. (1991). Coexistence of different shell microstructure types in a single cell layer of the Veneridae (Bivalvia, Mollusca). In: Suga, S. and Nakahara, H. (Eds.). *Mechanisms and phylogeny of mineralization in biological systems*. Springer Verlag, Tokyo, Berlin etc. pp. 421-425

Shimamoto, M. (1996). Phylogenetic implication of shell microstructures and amino acid compositions in the Veneridae (Bivalvia, Mollusca). *Bulletin de l'Institut Oceanographique Numero Special 14(4)*: 263-270.

Shulenberger, E. (1970). Responses of *Gemma gemma* to a catastrophic burial. *The Veliger* 13(2): 163-170

Shuto, T. (1979). Neogene molluscan fossils from South Kyushu. (Molluscan fossils from the Miyazaki group. 3). *Atlas of Japanese Fossils* 57(339).

Singer, B. S. and H. K. Mienis (1995). *Ventricolaria toreuma* (Gould) reported for the first time from the Red Sea (Bivalvia, Veneridae). *Levantina* 82: 9-11.

Skarlato, O. A. (1981). [Bivalve molluscs of the temperate latitudes of the western part of the Pacific Ocean.]. *Opredeliteli po Faune SSSR No. 126*: 1-461.

Skarlato, O. A. and Y. I. Starobogatov (1972). [Bivalvia.]. *Vodyanitzkii*: 1-340.

Skoglund, C. C. (1988). Deep water shells from off Isla Smith, Bahía de Los Angeles, Baja California, Mexico. *The Festivus* 20(11): 110-116

Smith, G. M. (1928). Food material as a factor in growth rate of some Pacific clams. *Proceedings and Transactions of the Royal Society of Canada*, (3)22[for 1928](2)(5): 287-291

Smith, G. M. (1933). Further observations on the ecology, rate of growth and food supply of some Pacific clams. *Proceedings and Transactions of the Royal Society of Canada* (3)27[for 1933](5): 229-245

Smith, L. S. and J. C. Davis (1965). Haemodynamics in *Tresus nuttallii* and certain other bivalves. *Journal of Experimental Biology* 43(1): 171-180

Solis, U. I., A. P. Sanchez, V. S. Navarrete (1976). Identification y descripcion de larvas de moluscos bivalvos en el plancton del Estero de Castro. *Boletin de la Sociedad de Biología de Concepcion* 50: 185-195.

Sowerby, G. B., I. (1835). [...] new species of shells contained in the collection of Mr. Cuming]. *Proceedings of the Zoological Society of London* for 1835[3](26): 21-23

- Sowerby, G. B., I. (1851). Monograph of the genus *Cythereaea*. In: G. B. Sowerby II, ed., *Thesaurus conchyliorum*; or, monographs of genera of shells 2(12): 611-648, pls. 127-136
- Sowerby, G. B., I. (1852). Monograph of the genus *Tapes*. In: G. B. Sowerby II, ed., *Thesaurus conchyliorum*; or, monographs of genera of shells 2(14): 678-702, pls. 145-151
- Sowerby, G. B., I. (1853). Monograph of the genus *Venus*; supplement; index [to Veneridae]. In: G. B. Sowerby II, ed., *Thesaurus conchyliorum*; or, monographs of genera of shells 2(14): 703-762, pls. 152-163
- Sowerby, G. B., I. (1854). Monograph of the genus *Venerupis*. In: G. B. Sowerby II, ed., *Thesaurus conchyliorum*; or, monographs of genera of shells 2(15): 763-770, pls. 164-165
- Sowerby, G. B., I. (1874). Monograph of the genus *Venerupis*. In: G. B. Sowerby II, ed., *Conchologia iconica*; or, illustrations of the shells of molluscous animals 19: 4 pls.
- Sparks, A. K. and K. K. Chew (1966). Gross infestation of the little neck clam, *Venerupis staminea*, with a larval cestode (*Echeneibothrium* sp.). *Journal of Invertebrate Pathology* 8(3): 413-416
- Stanley, S. M. (1975). Why clams have the shape they have: an experimental analysis of burrowing. *Paleobiology* 1(1): 48-58
- Stanley, S. M. (1981). Infaunal survival: alternative functions of shell ornamentation in the Bivalvia (Mollusca). *Paleobiology* 7(3): 384-393
- Stead, R.-A., E. Clasing, J. M. Navarro, G. Asencio (1997). Reproductive cycle and cohort formation of *Venus antiqua* (Bivalvia: Veneridae) in the intertidal zone of southern Chile. *Revista Chilena de Historia Natural* 70(2): 181-190.
- Stearns, R. E. C. (1891). Edible shell notes, for the nautilus. *Nautilus* 5(3): 25-29.
- Stearns, R. E. C. (1898). Notes on *Cytherea (Tivela) crassatelloides* Conrad, with descriptions of many varieties. *Proceedings of the United States National Museum* 21(1149): 371-378, pls. 23-25
- Stearns, R. E. C. (1899). *Donax stultorum*, Mawe = Conrad's species, *Cytherea crassatelloides*. *The Nautilus* 13(7): 73-75

- Stephenson, M. D. (1977). Sea otter predation on Pismo clams in Monterey Bay. *California Fish and Game* 63(2): 117-120
- Stephenson, R. L. and P. E. Chanley (1979). Larval development of the cockle *Chione stutchburyi* (Bivalvia: Veneridae) reared in the laboratory. *New Zealand Journal of Zoology* 6(4): 553-560.
- Stock, J. H. (1995). Copepoda Poecilostomatoida associated with Bivalvia from New Guinea. *Hydrobiologia* 312(1): 37-45.
- Stohler, R. (1960). Fluctuations in mollusk populations after a red tide in the Estero de Punta Banda, Lower California, Mexico. *The Veliger* 3(1): 23-28, pl. 4
- Strand, E. (1926). Miscellanea nomenclatorica zoologica et palaeontologica. *Archiv für Naturgeschichte* Abt. A (8. Heft): 30
- Stresemann, E., H.-J. Hannemann, B. Klausnitzer, K. Senglaub (1992). *Fauna von Deutschland* [Fauna of Germany. Volume 1. Invertebrates (without Insecta). 8th edition]. Volk und Wissen Verlag GmbH, Berlin: 1-637.
- Strougo, A. (1975). Les Corbulidae de l'Eocene superieur d'Egypte. *Bulletin du Museum National d'Histoire Naturelle* 280: 29-40.
- Subba Rao, N. V., K. V. Surya Rao, S. Maitra (1991). Marine molluscs. In: Chhotani, O.B. (Ed.). *Fauna of Orissa*. Part 3: 1-262.
- Szoor, G. (1972). Analyses of molluscan shells by the derivatographic fingerprint method. *Geologicky Sbornik* 23(1): 15-38.
- Takagi, T. (1986). On some *Dosinia* from the Pliocene and Pleistocene formations in Hokkaido, Japan. *Journal of the Faculty of Science, Hokkaido University, Series 4: Geology and Mineralogy* 21(4): 599-617.
- Takagi, T. (1990). Taxonomic and paleozoogeographic significance of Cenozoic hunch-back like venerids. Part 1. *Gigantocallista*, gen. nov. (Mollusca: Bivalvia), a new endemic venerid genus from the Pliocene Tatsunokuchi Formation, Sendai City, Miyagi Prefecture, northeastern Japan. *Transactions and Proceedings of the Palaeontological Society of Japan New Series No. 159*: 541-553.
- Taki, I. (1941). On the variation in the colour pattern of a bivalve, *Venerupis philippinarum*, with special reference to its bilateral asymmetry. *Venus* 11(2-3): 70-87.

- Tanabe, K. and T. Oba (1988). Latitudinal variation in shell growth patterns of *Phacosoma japonicum* (Bivalvia: Veneridae) from the Japanese coast. *Marine Ecology Progress Series* 47(1): 75-82.
- Tanaka, Y. (1954). Spawning season of important bivalves in Ariake Bay -- III. *Venerupis semidecussata* (Reeve). *Japanese Society of Scientific Fisheries, Bulletin* 19(12): 1165-1167
- Tashiro, M. and T. Kozai (1989). Bivalve fossils from the type Monobegawa Group (part 4). *Research Reports of Kochi University Natural Science* 38: 113-156.
- Tashiro, M. and M. Otsuka (1982). Bivalve fossils from the uppermost formation of the Upper Himenoura Subgroup in Amakusa-Shimojima Island, Kyushu (Part 2). *Memoirs of the Faculty of Science Kochi University Series E Geology* 3: 7-22.
- Tata, B. A. and A. A. S. Prieto (1991). Production secundaria en una poblacion del tropical *Tivela mactroides* (Veneridae) en el Oriente de Venezuela. *Caribbean Journal of Science* 27(1-2): 28-34.
- Tegland, N. M. (1928). *Pitar ida*, a new Recent species from Sitka, Alaska. *The Nautilus* 42(1): 4-6
- Tegland, N. M. (1929). Correlation and affinities of certain species of *Pitaria*. California, *Bulletin of the University of California, Publications, Department of Geology* 18(10): 275-290, pls. 21-23
- Teranes, J.-L., D.-H. Geary, B.-E. Bemis (1996). *The oxygen isotopic record of seasonality in Neogene bivalves from the Central American isthmus*. Jackson, Jeremy B.C. pp.1-425
- Tevesz, M. J. S. (1972). Implications of absolute age and season of death data compiled for Recent *Gemma gemma*. *Lethaia* 5(1): 31-38
- Thompson, I. (1975). Biological clocks and shell growth in bivalves. pp. 149-162, in: G. D. Rosenberg and S. K. Runcorn, eds., *Growth rhythms and the history of the Earth's rotation*. London, etc. (Wiley). xvi + 559 pp.
- Thompson, J. K. (1982). Population structure of *Gemma gemma* (Bivalvia: Veneridae) in south San Francisco Bay, with a comparison to some northeastern United States estuarine populations. *Veliger* 24(3): 281-290.
- Tiffany, W. J., III (1982). Excretory concretions in the sunray venus clam, *Macrocallista nimbosa* (Bivalvia: Veneridae). *Veliger* 25(1): 77-78.

- Toba, D. R., D. S. Thompson, K. K. Chew, G. J. Anderson, M. B. Miller (1992). Guide to Manila clam culture in Washington. Washington, *University of Washington, Sea Grant Program, Report 92-01*: viii + 80 pp.
- Tomlin, J. R. L. B. (1923). Some Synonyms in the Veneridae. *Proceedings of the Malacological Society* 15: 310-313.
- Tomlinson, P. K. (1977). Mortality, growth, and yield per recruit for Pismo clams. *California Fish and Game* 54(2): 100-107
- Tryon, G. W., Jr. (1865). Description of a new species of *Mercenaria*. *American Journal of Conchology* 1(4): 297, pl. 26
- Verduin, J. (1969). Hard clam pumping rates: energy requirement. *Science* 166(3910): 1309-1310
- Vermeij, G. J. (1980). Drilling predation of bivalves in Guam; some paleoecological implications. *Malacologia* 19(2): 329-333.
- Villarreal-Chavez, G., F. Garcia-Dominguez, F. Correa, N. Castro-Castro (1999). Note on the geographic distribution of *Chione cortezi* (Carpenter, 1864) (Mollusca: Pelecypoda: Veneridae). *Ciencias Marinas* 25(1): 153-159.
- Vokes, H. E. (1939). Molluscan faunas of the Domengine and Arroyo Hondo Formations of the California Eocene. *Annals of the New York Academy of Sciences* 38: 1-246; pl.1-22.
- Voloshina, M. I. (1973). [Late Tortonian bivalve molluscs of Moldavia and aspects of their existence]. Shtiintza, Kishinev. pp. 1-191
- Wang, H. (1989). Veneridae. In: Cai, R. and Huang, W. (Eds). *Fauna of Zhejiang: mollusks*. Zhejiang Science and Technology Publishing House, Hangzhou, Zhejiang Province. pp. 243-256
- Warner, R. W. and S. C. Katkansky (1969). Infestation of the clam *Protothaca staminea* by two species of tetrphyllidian cestodes (*Echeneibothrium* spp.). *Journal of Invertebrate Pathology* 13(1): 129-133
- Warner, R. W. and S. C. Katkansky (1969). A larval cestode from the Pismo clam, *Tivela stultorum*. *California Fish and Game* 55(3): 248-251

- Webb, C. M. (1987). Post-larval development of the bivalves *Nucula turgida*, *Venus striatula*, *Spisula subtruncata* and *S. elliptica* (Mollusca: Bivalvia), (with reference to the late larva). *Proceedings of the Marine Biological Association of the United Kingdom* 67(2): 441-459
- Weinberg, J. R. (1984). Interactions between functional groups in soft substrata: do species differences matter? *Journal of Experimental Marine Biology and Ecology* 80(1): 11-28
- Weinberg, J. R. (1985). Factors regulating population dynamics of the marine bivalve *Gemma gemma*: intraspecific competition and salinity. *Marine Biology* 86(2): 173-182
- Weinberg, J. R. and R. B. Whitlach (1983). Enhanced growth of a filter-feeding bivalve by a deposit-feeding polychaete by means of nutrient regeneration. *Journal of Marine Research* 41(3): 557-569
- Wells, H. W. (1957). Abundance of the hard clam *Mercenaria mercenaria* in relation to environmental factors. *Ecology* 38(1): 123-128
- Weymouth, F. W. (1921). The abundance of young Pismo clams. *California Fish and Game* 7(2): 103-106
- Weymouth, F. W. (1923). Certain features of the physiology of growth as illustrated by the lamellibranch *Tivela*. *American Journal of Physiology* 63(3): 412
- Weymouth, F. W. (1923). The life-history and growth of the Pismo clam (*Tivela stultorum* Mawe). California, *Fish and Game Commission, Fish Bulletin* 7: 120 pp.
- Whitehead, T. (1983). The genus *Periglypta* Jukes-Brown 1914 in Australia. *Australian Shell News* No. 41: 4-5.
- Whitehead, T. (1997). A puzzling *Periglypta*. *Australasian Shell News* 94(7).
- Willett, G. (1944). Two new west American pelecypods. *Bulletin of the Southern California Academy of Sciences* 43(1): 19-22, pls. 7, 8
- Williams, J. G. (1980). The influence of adults on the settlement of spat of the clam, *Tapes japonica*. *Journal of Marine Research* 38(4): 729-741
- Williams, J. G. (1980). Growth and survival in newly settled spat of the Manila clam, *Tapes japonica*. United States, *Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Fishery Bulletin* 77(4): 891-900

Williamson, M. B. (1893). On Clementia subdiaphana Cpr. in San Pedro Bay. *The Nautilus* 6(10): 116

Woodring, W. P. (1926). American Tertiary mollusks of the genus Clementia. United States, *Department of the Interior, Geological Survey, Professional Paper* 147C: 23-49, pls. 14-17

Woodring, W. P. (1982). Geology and paleontology of Canal Zone and adjoining parts of Panama; description of Tertiary mollusks; pelecypods, Propeamussiidae to Cuspidariidae; additions to families covered in P 306-E; additions to gastropods; cephalopods. Reston, VA, United States, *U. S. Geological Survey*. pp. 541-759

Wu, W. L. and H. P. Liu (1992). Developmental rhythm on gamete and gonad of Meretrix lusoria from Taiwan (Bivalvia: Veneridae). *Bulletin of Malacology Republic of China* 17: 79-86.

Wu, W. L., H. P. Liu, K. Y. Liao (1992). The morphology and anatomy of Meretrix lusoria from Taiwan (Bivalvia: Veneridae). *Bulletin of Malacology Republic of China* 17: 91-98.

Yang, X. (1986). *Evolutionary stasis in Neogene bivalve lineages; a morphometric evaluation*. Doctoral Thesis, The Johns Hopkins University. Baltimore, MD, Pages: 185.

Yap, W. G. (1978). Population biology of the Japanese little-neck clam, Tapes philippinarum, in Kaneohe Bay, Oahu, Hawaiian Islands. *Pacific Science* 31(3): 223-244

Yaseen, A. E., M. A. El Helow, et al. (1995). A cytogenetical study on Red Sea marine bivalve valves [bivalves] Dosinia radiata (Veneridae: Mollusca). *Assiut Veterinary Medical Journal* 32(64): 71-79.

Yasuda, J., I. Hamai, H. Hotta (1954). A note on the spawning season in Venerupis philippinarum. *Bulletin of the Japanese Society of Scientific Fisheries* 20(4): 277-279

Yasuda, J. and S. Takamori (1952). Study on seeds of asari (Venerupis philippinarum). *Bulletin of the Naikai Regional Fisheries Laboratory (Fisheries Agency)* 2(12): 1-11

Yoo, S. K., Y. J. Chung, H. Y. Ryu (1978). Biological studies on the propagation of important bivalves. 6. Morphological characteristics of the short-necked clam, Tapes japonica. Pusan, *Bulletin of the National Fisheries University, (Natural Science)* 18(1-2): 89-94

Yoshida, H. (1935). On the full-grown veligers and early young-stage of Venerupis (Amygdala) philippinarum (Adams & Reeve). *Venus* 5(5): 264-273, pl. 12

Yoshida, H. (1953). Studies on larvae and young shells of industrial bivalves in Japan. *Journal of the Shimonoseki College of Fisheries* 3(1): 106 pp.

Zakhvatkina, K. A. (1972). [Larvae of Bivalvia]. *Vodyanitzkii*: 1-340.

Zhuang, Q. (1964). Studies on Chinese species of Veneridae (Class Lamellibranchia). *Studia Marina Sinica* 5: 43-106, 10 pls.

Zhuang, Q., H. Lin, X. Liang (1981). On the species of the genus *Ruditapes* (Mollusca, Lamellibranchia, Veneridae) off China coast. *Studia Marina Sinica* 18: 207-218.

Zinsmeister, W. J. (1984). Late Eocene bivalves (Mollusca) from the La Meseta Formation, collected during the 1974-1975 joint Argentina-American Expedition to Seymour Island, Antarctic Peninsula. *Journal of Paleontology* 58(6): 1497-1527.