



CW[®]

Continuous Wormer

(pyrantel tartrate)
Equine Anthelmintic

- Medicated Pellets for Daily Feeding
- Provides Continuous Worm Control

For the prevention of Strongylus vulgaris larval infections in horses

For control of the following parasites in horses:

LARGE STRONGYLES (adults) *S. vulgaris*, *S. edentatus*, *Triodontophorus* spp.

SMALL STRONGYLES (adults and fourth-stage larvae) *Cyathostomum* spp., *Cylicocycylus* spp.,
Cylicostephanus spp., *Cylicodontophorus* spp., *Poteriostomum* spp.

PINWORMS (adults and fourth-stage larvae) *Oxyuris equi*

ASCARIDS (adults and fourth-stage larvae) *Parascaris equorum*

ACTIVE DRUG INGREDIENT

Pyrantel Tartrate 2.11% (9.6 grams per pound)

GUARANTEED ANALYSIS

Crude Protein (min)	17%	Crude Fat (min)	2%
Crude Fiber (max)	27%	Moisture (max)	12%

INGREDIENTS

Dehydrated Alfalfa Meal, Wheat Middlings, Brewers Dried Yeast, Brewers Dried Grains, Natural and Artificial Flavors, and preserved with Propionic Acid.

ANADA 200-168

DIRECTIONS FOR USE: SEE BACK PANEL

NET WT. 50 lb (22.7 kg)

EQUI AID

CW[®] Continuous Wormer

(pyrantel tartrate)

Equine Anthelmintic

CW[®] Continuous Wormer (pyrantel tartrate) is to be administered on a continuous basis as a top dress on the horse's daily ration. Feed at the rate of 1.2 mg pyrantel tartrate per pound of body weight per day. To achieve this dose administer 0.5 ounces of CW[®] Continuous Wormer per 250 lb of body weight (a CW[®] Continuous Wormer measuring scoop is enclosed). CW[®] Continuous Wormer should be administered for the entire period that the animal is at risk to internal parasites. Unprotected animals that have grazed may already have an established *S. vulgaris* larval infection. Before administering CW[®] Continuous Wormer, these animals should be treated with a therapeutic dose of a larvicidal product.

Foals may be fed CW[®] Continuous Wormer at such time when consistent intake of grain mix is occurring. This is generally between two and three months of age. CW[®] Continuous Wormer may be used in mares at any stage of pregnancy or lactation. Stallion fertility is not affected by the use of CW[®] Continuous Wormer.

TOP-DRESS DIRECTIONS	
<i>(1 oz Measuring Scoop Enclosed – 1 Level Scoop per 500 lb Body Weight)</i>	
POUNDS BODY WEIGHT	Ounces per day of CW [®] Continuous Wormer
250	0.5 oz
500	1.0 oz
750	1.5 oz
1,000	2.0 oz
1,250	2.5 oz

CAUTION:

Consult your veterinarian before using in severely debilitated animals and for assistance in the diagnosis, treatment and control of parasitism. Do not mix in feeds containing Bentonite.

WARNING:

Not for use in horses intended for food.
Not for Human Use.

Questions?

Call with questions or comments: 1-800-735-3399
Store in dry, cool place.

RM#: 99.0000407.00 09-1432


©2006, 2009 Equi Aid Products

Central Garden & Pet and its logo are trademarks of Central Garden & Pet Company.
Farnam, the Farnam logo, CW and the Equi Aid logo are trademarks of Farnam Companies, Inc.



Distributed by:
Farnam Companies, Inc. • Phoenix, AZ 85013
equiaid.com

3002626



CENTRAL
Garden & Pet