

# 'Reverse' identification key for mosquito species



More and more people are getting involved in the **surveillance of invasive mosquito species in the EU/EEA**, not just professionals with formal training in entomology. There are many taxonomic keys available for identifying mosquitoes of medical and veterinary importance, but they are almost all designed for professionally trained entomologists.

The current identification key aims to provide non-specialists with a simple mosquito recognition tool for distinguishing between invasive mosquito species and native ones. On the overview displays ([p. 4, 5 and 6](#)) you can select the species that best resembles your specimen. On the species-specific pages you will find additional information on those species that can easily be confused with that selected, so you can check these additional pages as well.

This key provides the non-specialist with **reference material to help recognise an invasive mosquito species** and gives details on the morphology (in the species-specific pages) to help with verification and the compiling of a final list of candidates. The key displays six invasive/exotic mosquito species that are present in the EU/EEA or have been intercepted in the past. It also contains 16 native species. The native species have been selected (1) based on their morphological similarity with the invasive species; (2) on the likelihood of encountering them; and (3) to include representatives of all mosquito genera present in Europe.

If you want to further develop your identification skills, you can consult 'Mosquitoes - Identification, Ecology and Control. Third Edition' [1] or identification tools available online, such as MosKeyTool: <https://www.medilabsecure.com/moskeytool.html>.

Different scientific names are available for the species included in this key. Table 1 provides an overview of the scientific names used in the key [2] and names based on the revisions made by Reinert and colleagues [3-7] that can be found in scientific literature. We hope that this will aid communication between health professionals and scientists.

| Species name used in the key               | Synonyms currently in use               | Common name                         |
|--|---|-------------------------------------|
| <i>Aedes aegypti</i>                       | <i>Stegomyia aegypti</i>                | Yellow fever mosquito               |
| <i>Aedes albopictus</i>                    | <i>Stegomyia albopicta</i>              | Asian tiger mosquito                |
| <i>Aedes atropalpus</i>                    | <i>Georgecraigius atropalpus</i>        | American rock pool mosquito         |
| <i>Aedes japonicus japonicus</i>           | <i>Hulecoeteomyia japonica japonica</i> | Asian bush or rock pool mosquito    |
| <i>Aedes koreicus</i>                      | <i>Hulecoeteomyia koreica</i>           |                                     |
| <i>Aedes triseriatus</i>                   | <i>Ochlerotatus triseriatus</i>         | American Eastern tree hole mosquito |
| <hr style="border-top: 1px dashed #000;"/> |   |                                     |
| <i>Aedes cantans</i>                       | <i>Ochlerotatus cantans</i>             |                                     |
| <i>Aedes caspius</i>                       | <i>Ochlerotatus caspius</i>             | Salt marsh mosquito                 |
| <i>Aedes communis</i>                      | <i>Ochlerotatus communis</i>            | Snowpool mosquito                   |
| <i>Aedes cretinus</i>                      | <i>Stegomyia cretina</i>                |                                     |
| <i>Aedes geniculatus</i>                   | <i>Dahlia geniculata</i>                |                                     |
| <i>Aedes pulchritarsis</i>                 | <i>Ochlerotatus pulchritarsis</i>       |                                     |
| <i>Aedes vexans</i>                        | <i>Aedimorphus vexans</i>               |                                     |
| <i>Aedes vittatus</i>                      | <i>Fredwardsius vittatus</i>            |                                     |
| <i>Aedes zammitii</i>                      | <i>Acartomyia zammitii</i>              |                                     |
| <i>Anopheles plumbeus</i>                  |   |                                     |
| <i>Coquillettidia richiardii</i>           |   |                                     |
| <i>Culex pipiens</i>                       |   | (Northern) house mosquito           |
| <i>Culiseta annulata</i>                   |   |                                     |
| <i>Culiseta longiareolata</i>              |   |                                     |
| <i>Orthopodomyia pulcripalpis</i>          |   |                                     |
| <i>Uranotaenia unguiculata</i>             |   |                                     |

**Note:** *Aedes* above dashed line are invasive species; *Aedes* below dashed line are native *Aedes* species. The solid lines separate the genera.




# What is/is not a mosquito?

## Scientific classification of mosquitoes

| Kingdom  | Phylum     | Class   | Order   | Suborder   | Family    | Genus             | Species                |
|----------|------------|---------|---------|------------|-----------|-------------------|------------------------|
| Animalia | Arthropoda | Insecta | Diptera | Nematocera | Culicidae | e.g. <i>Aedes</i> | e.g. <i>albopictus</i> |



## Body structures




### **Culicidae** or True (biting) mosquitoes (*Culex pipiens* female)

Females with elongated 'needle-like' mouth parts [1]; wings with scales; long legs; size of mosquito 3–6mm;

## Female vs male mosquitoes



## Diptera that look similar to Culicidae



### **Mycetophilidae** or Fungus gnats

Short mouth parts [1]; markedly humped thorax [2]; 2–14mm



### **Limoniidae** or Limonid crane flies

Short mouth parts [1]; very long legs [2]; slender body [3]; 2–11mm



### **Ceratopogonidae** or Biting midges

Short proboscis, not needle-like [1]; hooped thorax [2]; wing without scales [3]; 1–3mm



### **Chironomidae** or Chironomids or Non-biting midges

Short mouth parts [1]; no scales on wing [2]; shape of thorax, 'hooped' [3]; ~10mm



### **Anisopodidae** or Wood gnats


Short mouth parts [1]; ~4–12 mm



### **Tipulidae** or Crane flies

No needle-like proboscis; looks like an oversized mosquito; slender body [1]; long legs [2]; wing span 1–6.5 cm

# Life cycle of a floodwater *Aedes* mosquito




# Female mosquitoes

Representatives of the seven mosquito genera present in Europe

*Aedes albopictus* ●  
p.8



*Anopheles plumbeus* ●  
p.22



*Coquillettidia richiardii* ●  
p.23



*Culex pipiens* ●  
p.24



*Culiseta annulata* ●  
p.25



*Culiseta longiareolata* ●  
p.26



*Orthopodomyia pulcripalpis* ●  
p.27



*Uranotaenia unguiculata* ●  
p.28



0 2.5 5 mm

● Invasive

● Native

# Female *Aedes* mosquitoes

*Aedes aegypti* •  
p.7



*Aedes albopictus* •  
p.8



*Aedes atropalpus* •  
p.9



*Aedes japonicus* •  
p.10



*Aedes koreicus* •  
p.11



*Aedes triseriatus* •  
p.12



*Aedes cantans* •  
p.14



*Aedes caspius* •  
p.15



*Aedes communis* •  
p.16



*Aedes cretinus* •  
p.13



*Aedes geniculatus* •  
p.17



*Aedes pulcritarsis* •  
p.18



*Aedes vexans* •  
p.19



*Aedes vittatus* •  
p.20



*Aedes zammiti* •  
p.21



- Invasive
- Exotic
- Native

# Male invasive *Aedes* mosquitoes

*Aedes aegypti* •  
p.7



*Aedes albopictus* •  
p.8



*Aedes japonicus* •  
p.10



*Aedes koreicus* •  
p.11




*Aedes triseriatus* •  
p.12



*Aedes atropalpus* •  
p.9



0 2.5 5 mm



- Invasive
- Exotic
- Native



Yellow fever mosquito

# *Aedes aegypti*

*Stegomyia aegypti*




© Anders Lindström/SVA



© Anders Lindström/SVA

## Morphological characteristics

1. The genus *Aedes* has a pointy abdomen
2. Scutum (dorsal part of the thorax) has silver scales in the shape of a lyre on a black background.
3. Small size.
4. Contrasting black-and-white colouration.
5. Silvery-white markings on legs and abdomen.



© Disa Eklof/SVA

## Easily-confused species

*Aedes albopictus*; *Aedes cretinus*

## Status in Europe

- Exotic, invasive
- Origin: Tropical Africa

## Distribution



<https://bit.ly/3MAZ3un>

## Likely point of entry

International trade and travel, airports.

## Ecology (habitat, breeding sites)

- Originally *Aedes aegypti* was found in forested areas, using tree holes as habitats. The species is now commonly found in tropical and sub-tropical areas, in close proximity to humans. It thrives well in urban and peri-urban environments.

- In Europe, female *Aedes aegypti* will lay their eggs in artificial water containers, much like *Aedes albopictus*. Suitable habitats include earthenware pots and water tanks, uncovered cisterns, empty cans, flower pots or dishes, broken bottles or discarded tyres.
- On the island of Madeira, *Aedes aegypti* is active throughout the year, with a peak in abundance from August to October.
- The eggs are resistant to desiccation. Unlike *Aedes albopictus*, *Aedes aegypti* cannot produce diapausing eggs resistant to frost.

## Biting habits

- Human, occasionally other mammals.
- The females feed predominantly during the day in shaded places and only occasionally during the night.
- Females can feed multiple times between egg laying.

(Asian) tiger mosquito

# *Aedes albopictus*

*Stegomyia albopicta*




© Anders Lindström/SVA



© Anders Lindström/SVA

## Morphological characteristics

1. Scutum (dorsal part of the thorax) with a median silver-scale line on a black background.
2. Medium size.
3. White stripes beside and posterior to the median white line on the scutum do not reach the middle of the scutum.
4. Mid leg: white basal bands only on tarsomere 1 and 2.



© Disa Eklof/SVA

## Easily-confused species

*Aedes cretinus*; *Aedes aegypti*

## Status in Europe

- Exotic, invasive
- Origin: Asia

## Distribution



<https://bit.ly/3MAZ3un>

## Likely point of entry

Eggs of *Aedes albopictus* are often imported in used tyres or hydroponic plant containers. Other ways of introduction include vehicles in which adult *Aedes albopictus* individuals can 'hitchhike' across borders.

## Ecology (habitat, breeding sites)

- In Europe *Aedes albopictus* prefers urban and suburban habitats.

- In a temperate climate *Aedes albopictus* has been shown to be most active during the period May-September.
- Adult *Aedes albopictus* females can produce eggs able to survive periods of frost during the winter (diapausing eggs).
- Diapausing eggs of European *Aedes albopictus* have been shown to be able to survive a cold spell of -10°C, whereas eggs of tropical *Aedes albopictus* can only survive -2°C.
- Larvae develop in natural or artificial water containers. Artificial aquatic habitats include tyres, barrels, rainwater gully catch basins and flower pot dishes.

## Biting habits

- Adult females bite aggressively, usually during the day outdoors, but also during the night indoors.
- *Aedes albopictus* feeds on humans, domestic and wild animals, reptiles, birds and amphibians, depending on host availability.



American rock pool mosquito

# *Aedes atropalpus*

*Georgeraigijs atropalpus*

♀  
Exotic




© Anders Lindström/SVA



© Anders Lindström/SVA

## Morphological characteristics

1. Presence of two lateral lines of pale scales on a black background on the scutum (dorsal part of the thorax).
2. Legs have inter-articular pale rings.
3. Wings have a patch or short line of pale scales at the base of the first principal longitudinal vein of the wing (costa).
4. Dorsal plates of abdomen with regular pale basal bands



© Disa Eklof/SVA

## Easily-confused species

*Aedes caspius*; *Aedes pulcritarsis*

## Status in Europe

- Exotic
- Origin: North and Central America

## Distribution



<https://bit.ly/3MAZ3un>

## Likely point of entry

International used tyre trade.

## Ecology (habitat, breeding sites)

- *Aedes atropalpus* appear early in the season.
- Females can lay their first eggs without taking a bloodmeal (autogenous egg production).
- *Aedes atropalpus* larvae are most often associated with soft water rock pool habitats along mountain streams in North America.

- The species is also known to breed in a variety of artificial containers, especially discarded tyres and other man-made water collectors such as concrete septic tanks.
- Eggs are desiccation-resistant and can therefore survive outside of water until conditions are suitable for them to hatch.

## Biting habits

- *Aedes atropalpus* will readily bite humans and the species has a preference for mammalian hosts.
- Females bite at night and during the day and are known to be a pest in the vicinity of aquatic habitats.

Asian bush or rock pool mosquito

# *Aedes japonicus japonicus*


*Ochlerotatus japonicus japonicus*, *Hulecoeteomyia japonica japonica*

  
Invasive



## Morphological characteristics

1. Scutum (dorsal part of the thorax) with several lines of yellowish scales on a black background.
2. Relatively large.
3. White scale patches on black legs.
4. Tarsomeres 4 and 5 of hind leg almost entirely dark (different from *Aedes koreicus*).
5. Dorsal plates of the abdomen with lateral and median pale patches at the base of each segment that do not form complete bands.



© Disa Eklof/SVA

## Easily-confused species

*Aedes koreicus*

## Status in Europe

- Exotic, invasive
- Origin: Asia

## Distribution



<https://bit.ly/3MAZ3un>

## Likely point of entry

Can be imported by international tyre trade.

## Ecology (habitat, breeding sites)

- Adults are often found in forested areas.
- *Aedes japonicus* prefers to lay eggs in shady rock holes but exploits also a large range of both habitats including tree holes, tyres, bird baths, and all breeding sites rich in organic matter.
- *Aedes japonicus* can produce frost- and desiccation-resistant diapausing eggs that can remain dormant over winter and hatch once environmental conditions become favourable.
- *Aedes japonicus* can also overwinter as larva.

## Biting habits

- *Aedes japonicus* females feed mainly on mammals.
- Female *Aedes japonicus* feed during the day. This species is an aggressive biter and will readily bite humans outside —mainly in forested areas, but occasionally also inside houses.

# Aedes koreicus

*Ochlerotatus koreicus*, *Hulecoeteomia koreica*




Invasive



## Morphological characteristics

1. Strongly resembles *Aedes japonicus* in that it also has clear longitudinal lines on the scutum (dorsal part of the thorax).
2. Relatively large.
3. The presence of a complete 4th basal band on hind-tarsomere distinguishes the species from *Aedes japonicus*.
4. Can have an incomplete pale band at the base of hind tarsomere 5.



## Easily-confused species

*Aedes japonicus*

## Status in Europe

- Exotic, invasive
- Origin: Asia

## Distribution



<https://bit.ly/3MAZ3un>

## Likely point of entry

International trade routes, although for several established populations in Europe the introduction pathway is not clear.

## Ecology (habitat, breeding sites)

- *Aedes koreicus* overwinters as eggs which hatch during spring. Adults are most active between May and October.
- The species survives in the same manner as *Aedes japonicus* - frost- and desiccation-resistant eggs that are dormant during winter and hatch once environmental conditions become favourable.
- Larvae of *Aedes koreicus* can be found in both natural and artificial water containers such as garden ponds, water drums and other vessels and unused metal construction equipment. Natural sites include tree holes and stone cavities containing rain water and decaying tree leaves.

## Biting habits

- *Aedes koreicus* bites humans both during the day and at night.
- Hosts are humans and mammals.

# American Eastern tree hole mosquito

## *Aedes triseriatus*

*Ochlerotatus triseriatus*

  
Exotic




♀



♂

### Morphological characteristics

1. Presence of two pale-scaled stripes on the sides of the scutum (dorsal part of the thorax).
2. Absence of pale bands on the legs.
3. Dorsal plates of abdomen with baso-lateral pale patches.



© Disa Eklof/SVA

### Easily-confused species

*Aedes geniculatus*

### Status in Europe

- Exotic
- Origin: North America

### Distribution

Not introduced into Europe in the last five years.

### Likely point of entry

International tyre trade.

### Ecology (habitat, breeding sites)

- In its native range of North America, *Aedes triseriatus* is widely distributed. *Aedes triseriatus* breeds in tree-holes, tyres and other artificial containers. Adults are commonly encountered in forested areas.

- Hatching is thought to be dependent upon flooding and may be staggered, resulting in only a proportion of an egg batch hatching in response to a particular flooding event at a certain time. This allows *Aedes triseriatus* to survive in a variety of environments across its range.
- Eggs can survive prolonged periods without water. The species overwinters by diapausing eggs.

### Biting habits

- *Aedes triseriatus* feeds on a multitude of hosts (birds, mammals, reptiles) and readily bites humans.
- Females mostly bite during the day in shaded areas.

# Aedes cretinus

*Ochlerotatus cretinus*, *Stegomyia cretina*




Native



## Morphological characteristics

1. Scutum (dorsal part of the thorax) has a central narrow white stripe which forks at the end.
2. The scutum is bordered by a fine line of white scales, with a minute break at the scutal angle.
3. Resembles *Aedes albopictus* but differs through the longer lateral white lines on the scutum.
4. Hind tarsi with basal white rings; fifth tarsomere entirely white.
5. Small size.



© Dtsa Eklöf/SVA

## Easily-confused species

*Aedes albopictus*; *Aedes aegypti*

## Status in Europe

Native

## Distribution

*Aedes cretinus* is found in Cyprus, Greece and Turkey.

## Ecology (habitat, breeding sites)

Larvae can be found in tree holes and used tyres but also in small ground pools with thick vegetation which is not typical for container breeding mosquitoes.

## Biting habits

Females are aggressive human biters during the day, both in shaded and open places.


# *Aedes cantans*

*Ochlerotatus cantans*



## Morphological characteristics

1. Dark blackish-brown scaling with scattered white or yellow scales on body and wings.
2. Scutum covered with dark-brown or bronze-brown scales and the lateral parts with greyish-white or creamy scales.
3. Tarsomere 1 (tarsomere is the individual sub-segment of a tarsus) of all the legs has more or less mixed scales.
4. Tarsomeres 2–5 have moderately broad white basal rings, except for Tarsomere 5 of the forelegs which is entirely dark-scaled.



© Disa Eklöf/SVA

## Easily-confused species

- *Aedes cantans* is part of the *annulipes* group: This group includes *Aedes annulipes*, *Aedes behningi*, *Aedes cantans*, *Aedes cyprius*, *Aedes euedes*, *Aedes excrucians*, *Aedes flavescens*, *Aedes mercurator*, *Aedes riparius* and *Aedes surcoufi*.
- This species is generally not confused with any of the current invasive mosquito species in Europe.

## Status in Europe

Native

## Distribution

The species is widespread in Europe.

## Ecology (habitat, breeding sites)

- *Aedes cantans* produces only one generation (or in some cases two) per year. After this the species hibernates as diapausing eggs.
- The larval habitat of *Aedes cantans* is meadow or forest pools without much vegetation but with a layer of organic material at the bottom.

## Biting habits

Females feed on mammals and occasionally on birds.



Salt marsh mosquito


# *Aedes caspius*

*Ochlerotatus caspius*



## Morphological characteristics

1. Scutum (dorsal part of thorax) is covered with golden scales and has two dorsocentral white stripes.
2. Legs have inter-articular pale rings.
3. Dorsal plates of the abdomen have golden bands at the base and end of each plate and these are widest in the middle.
4. Small size.



© Disa Eklof/SVA

## Easily-confused species

- *Aedes atropalpus*
- *Aedes caspius* is part of the *caspius* group. Members of this group are difficult to distinguish based on morphology. Species included in this group are *Aedes berlandi*, *Aedes caspius*, *Aedes dorsalis*, *Aedes mariaae*, *Aedes phoeniciae*, *Aedes pulcritarsis* and *Aedes zammitii*.

## Status in Europe

Native

## Distribution



<https://bit.ly/3MAZ3un>

## Ecology (habitat, breeding sites)

- Adult *Aedes caspius* can be found in many habitats since they disperse over long distances from their larval hatching sites.
- Larvae develop mainly in coastal marshes (brackish water) with temporary flooding. They can also be found in rice fields or meadows flooded with fresh water. They can withstand substantial salt concentrations of up to 150g/L.
- The species overwinters as diapausing eggs.

## Biting habits

- Females feed mostly outside, but can be found indoors if there are large numbers. They feed both during the day and at night and are most active around dusk.
- Hosts are both humans and animals.



Snowpool mosquito

# *Aedes communis*

*Ochlerotatus communis*



Native




© Anders Lindström/SVA



© Anders Lindström/SVA

## Morphological characteristics

1. Medium size.
2. Dark-scaled tarsi.
3. Scutum (dorsal part of thorax) has yellow or golden scales.
4. Dorsal plates of abdomen with large pale basal bands.



© Disa Eklof/SVA

## Easily-confused species

- This species is part of the *communis* group. The members of this group are morphologically difficult to distinguish. The most common species in this group are *Aedes cataphylla*, *Aedes detritus*, *Aedes punctor* and *Aedes sticticus*.
- This species is generally not confused with any of the current invasive mosquito species in Europe.

## Status in Europe

Native

## Distribution

In Europe the species is found from the northern European region to the Mediterranean.

## Ecology (habitat, breeding sites)

- *Aedes communis* only completes one generation per year and is mainly found in swampy forests. The species prefers breeding in acid waterbodies, filled with water from melting snow or spring rainfall.
- Larvae can be found in small water bodies without vegetation, but with a dense layer of dead leaves. They can be found in strongly acidic waters with a pH as low as three.
- *Aedes communis* larvae can hatch in temperatures of little more than 0°C.

## Biting habits

- Hosts are warm-blooded forest inhabitants.
- Females are most active during the twilight period.



# *Aedes geniculatus*

*Dahlia geniculata*




© Anders Lindström/SVA



© Anders Lindström/SVA

## Morphological characteristics

1. Large mosquito
2. Conspicuous white knee spots.
3. Scutum has two central black stripes, sometimes fused into one, otherwise completely separated by a pale medio-dorsal stripe.
4. Tibiae and tarsi are entirely black-scaled.
5. Dorsal plates of abdomen with baso-lateral pale triangular patches



© Disa Eklof/SVA

## Easily-confused species

- *Aedes triseriatus*
- This species can be confused with the closely related species *Aedes echinus* and *Aedes gilcolladoi* (occurring in southern Europe only; not represented in this key).

## Status in Europe

Native

## Distribution

Europe

## Ecology (habitat, breeding sites)

- Adult *Aedes geniculatus* are mainly found in deciduous or mixed forests, rarely in conifer forests. Although they can be a nuisance to humans they rarely enter urban areas.
- The species mainly lays eggs in tree holes and open tree stumps, but can also colonise artificial containers, such as tyres. Breeding sites are usually rich in organic matter and tannins.

- Eggs are resistant to both frost and desiccation. *Aedes geniculatus* hibernates as eggs in northern climates and as larvae in southern climates.

## Biting habits

- Feeds on various mammals, including humans and cattle, but also on birds and reptiles.
- Females bite during daytime and during twilight hours. In south-eastern Europe, the species can be a nuisance for humans in forested areas.
- The species is also found in peri-urban sites.

# *Aedes pulcritarsis*

*Ochlerotatus pulchritarsis*




Native



## Morphological characteristics

1. Medium sized.
2. Dark palps with white tips.
3. Silvery-white inter-articular bands on tarsi.
4. Tarsomere 5 entirely white.
5. Scutum (dorsal part of the thorax) with large anterior yellow-brown patch and white posterior dots.



© Disa Eklöf/SVA

## Easily-confused species

- *Aedes atropalpus*
- This species can be confused with the closely related species *Aedes berlandi* (not represented in this key).

## Status in Europe

Native

## Distribution

Principally distributed in the Mediterranean region but found as far north as the Czech Republic. The species is also present in Central and South East Asia.

## Ecology (habitat, breeding sites)

- This species hibernates during the egg stage and has one to two generations per year.
- The larvae are mostly found in tree holes of deciduous and olive trees. Larval development can last up to two months. A suitable breeding site has a water temperature which never exceeds 21°C.
- In the absence of woodland areas, adult mosquitoes are often found near stables or in villages.

## Biting habits

Anthropophilic mosquitoes that bite outdoors and mainly during the daytime.



# Aedes vexans

*Aedimorphus vexans*




© Anders Lindström/SVA



© Anders Lindström/SVA

## Morphological characteristics

1. Scutum has an indefinite pattern with yellowy-cream-coloured scales.
2. Tarsi have narrow pale basal rings.
3. Dorsal plates of the abdomen have pale bi-lobed basal bands.
4. Small-to-medium sized.



© Disa Eklof/SVA

## Easily-confused species

- This species is generally not confused with any of the current invasive mosquito species in Europe.

## Status in Europe

Native

## Distribution



<https://bit.ly/3MAZ3un>

## Ecology (habitat, breeding sites)

- Adults can migrate long distances from breeding sites, up to 15 km, entering human settlements in large numbers.
- This species breeds mostly on flood plains, exhibiting fast larval development. It is often found in flooded areas and around the margins of lakes with fluctuating water levels.

## Biting habits

Feeds aggressively on humans and cattle during the daytime.


# *Aedes vittatus*

*Fredwardsius vittatus*



## Morphological characteristics

1. Palps with white apex and a few white scales in the middle.
2. Scutum (dorsal part of thorax) blackish brown with six silvery white spots arranged in three pairs.
3. White knee spots on all legs.
4. Hind leg with white rings on tarsomeres 1–4; Tarsomere 5 entirely white.



© Dina Eklof/SVA

## Easily-confused species

This species is generally not confused with any of the current invasive mosquito species in Europe.

## Status in Europe

Native

## Distribution

This species is present in the Mediterranean sub region and the Afrotropical and Oriental regions. Its distribution is probably limited by the low temperature in more northern areas.

## Ecology (habitat, breeding sites)

- *Aedes vittatus* larvae are found in rock pools containing clear water, with mud and organic material on the bottom. Other breeding sites are gardening utensils, boats, wells or tree holes.
- *Aedes vittatus* eggs can withstand desiccation and are laid above the water level.

## Biting habits

This species bites both indoors and outdoors, *Aedes vittatus* females can sometimes attack people in large numbers. Biting activity occurs mainly during the twilight period.



# *Aedes zammitii*

*Acartomyia zammitii*




© Anders Lindström/SVA



© Anders Lindström/SVA

## Morphological characteristics

1. Scutum (dorsal part of the thorax) covered with rust-brown to gold scales.
2. Scutellum with three groups of white sickle-shaped scales and golden-brown or dark setae.
3. Ventral surface of femora is white scaled.
4. Anterior surface of tibia is black scaled with speckled white scales.



© Dina Eklof/SVA

## Easily-confused species

- *Aedes caspius*
- This species is hardly distinguishable from the closely related species *Aedes mariaae* and *Aedes phoeniciae* (not represented in this key).
- This species is generally not confused with any of the current invasive mosquito species in Europe.

## Status in Europe

Native

## Distribution

Coastal areas of the Mediterranean region.

## Ecology (habitat, breeding sites)

- The eggs enter diapause at temperatures below 16°C and during short-day photoperiods.
- Larvae are found in rock pools along the Mediterranean seashore with concentrations of up to 20% of salt.
- *Aedes zammitii* produces multiple generations per year and can often be a nuisance in rocky coastal areas.

## Biting habits

Readily bites humans.

# Anopheles plumbeus

Native




© Anders Lindström/SVA



© Anders Lindström/SVA

## Morphological characteristics

1. Palps of the female mosquitoes from the genus *Anopheles* are the same length as the proboscis.
2. Global dark appearance.
3. Black proboscis and palps.
4. Vertex (dorsal part of the head) with a tuft of white scales and yellowish setae.
5. Pale / ash-grey median part of the scutum (dorsal part of the thorax).



## Easily-confused species

- *Anopheles claviger* (not shown in this key).
- This species is generally not confused with any of the current invasive mosquito species in Europe.

## Status in Europe

Native

## Distribution

Widely distributed throughout Europe. It can also be found in the northern Caucasus, the Middle East south to Iran, Iraq and North Africa.

## Ecology (habitat, breeding sites)

- Breeding sites are tree holes with decomposing organic matter. Other breeding sites can be artificial containers such as water catch basins and tyres, or unused stable slurry pits.
- Eggs are laid slightly above the water surface on the side of the of breedings sites and only hatch when the breeding site is flooded.
- *Anopheles plumbeus* overwinters as egg or larvae. Larvae can survive periods during which water surfaces are frozen.

## Biting habits

Females are most active during twilight and prefer to feed on mammalian blood, however they also feed on birds and reptiles. Some populations have a strong preference for human hosts.

# Coquillettidia richiardii




Native



## Morphological characteristics

1. The scales on the wings of this genus are broad and clearly visible.
2. Wing veins covered with broad, intermixed yellowish and brown scales.
3. Apex of the proboscis slightly broader and darker than the preceding part.
4. Pale ring in the middle of tarsomere 1 of all legs (sometimes hardly visible).
5. Broad pale basal rings usually present on all tarsomeres.



© Dina Eklöv/SVA

## Easily-confused species

This species is generally not confused with any of the current invasive mosquito species in Europe. There is only one other species of this genus present in Europe: *Coquillettidia buxtoni*.

## Status in Europe

Native

## Distribution

Common species in the western Palearctic region.

## Ecology (habitat, breeding sites)

- *Coquillettidia richiardii* has one to three generations per season. The larvae obtain oxygen from the air channels of aquatic plants by attaching themselves to the plants' roots.
- Breeding sites are various freshwater or slightly saline permanent water bodies such as marshes, lakes, riverbeds and estuaries.

## Biting habits

- Mostly feed on mammals but also birds and amphibians.
- Females are mostly active around the breeding sites but can fly a few kilometers to villages, where they can be a severe nuisance.
- Bite after sunset, at night, and just after sunrise.



House mosquito, Northern house mosquito

# Culex pipiens




Native



## Morphological characteristics

1. The genus *Culex* has a rounded abdomen end whereas the genus *Aedes* has a pointy abdomen.
2. Brownish-yellow, with no pattern on scutum.
3. Abdominal dorsal plates bearing yellowish basal bands.



© Disa Eklof/SVA

## Easily-confused species

- *Culex pipiens* is part of a group of species which are very hard to distinguish based on morphological characteristics. For more information see
- *Culex pipiens* is hard to confuse with any of the current invasive mosquito species in Europe.

- Females lay their eggs on water surfaces in batches as egg rafts of around 200 eggs. These eggs are non-dormant and the larvae hatch rapidly as soon as the embryonic development is completed (around 24 hours).
- Larvae can be found from mid-spring until the first frosts. In summer and autumn *Culex pipiens* can be found in abundance.

## Status in Europe

Native

For more information see fact sheet:



<https://bit.ly/3LmZQ1L>

## Distribution



<https://bit.ly/3MAZ3un>

## Ecology (habitat, breeding sites)

- *Culex pipiens* can inhabit nearly every type of water source. They can breed in clear water but also in water highly polluted with organic matter, and can even tolerate a small amount of salinity (e.g. coastal marshes or rock pools).
- Females overwinter in frost-free shelters, such as cellars, caves, bunkers, or ground burrows.

## Biting habits

- Females of the *Culex pipiens* form *pipiens* mainly bite birds (ornithophilic), feed outdoors (exophagic) and rest outdoors (exophilic).
- The *Culex pipiens* form *molestus* is characterised by females that mainly bite humans and other mammals indoors (endophagic) or occasionally outdoors. They frequently rest indoors (endophilic). They can lay a batch of eggs without a bloodmeal (autogenous).
- They are most active after dusk and before dawn.



# Culiseta annulata

Native




© Anders Lindström/SVA



© Anders Lindström/SVA

## Morphological characteristics

1. The genus *Culiseta* has a rounded abdomen whereas the genus *Aedes* has a pointy abdomen.
2. Generally large mosquitoes
3. Dark brown with whitish markings.
4. Tarsomere I (the individual sub-segment of a tarsus) with a noticeable white ring in the middle and white rings at the bases of Tarsomeres 2–4.
5. Tarsomeres 5 of all legs entirely dark-scaled.
6. Wings largely covered with scales, some of which clustered, forming distinct dark spots.
7. Dorsal plates of the abdomen with whitish basal bands.



© Disa Eklof/SVA

## Easily-confused species

*Culiseta longiareolata*; *Culiseta subochrea*;  
*Culiseta alaskaensis*.

## Status in Europe

Native

## Distribution

Occurs throughout Europe, but is more common in the North than in the South, where it is largely replaced by *Culiseta longiareolata*.

## Ecology (habitat, breeding sites)

- This species hibernates at the adult stage in the cellars or attics of dwellings or in domestic animal sheds, where it can be a nuisance even during the winter.

- Eggs are laid on water surfaces in stagnant pools, ponds, ditches, water troughs and other artificial containers, such as rainwater collection barrels. Manure basins can also act as a larval habitat. The larvae can even survive in water with a high level of salinity.
- Eggs are laid in rafts, like *Culex pipiens*.
- Adults can be encountered from early spring, with the population peak occurring in September.

## Biting habits

- Females bite humans, indoors and outdoors.
- Also bites birds occasionally.
- They are active during the night.

# Culiseta longiareolata




Native



## Morphological characteristics

1. The genus *Culiseta* has a rounded abdomen, whereas the genus *Aedes* has a pointy abdomen.
2. Very large size.
3. Lines on scutum.
4. White spots on legs.
5. Wing veins covered with dark scales, some of which clustered, forming dark spots which are less distinct than in *Culiseta annulata*.



© Disa Eklöf/SVA

## Easily-confused species

*Culiseta annulata*; *Culiseta subochrea*, (not shown in this key); *Culiseta alaskaensis* (not shown in this key)

## Status in Europe

Native

## Distribution

In Europe the species is widely distributed in the Mediterranean region.

## Ecology (habitat, breeding sites)

- Breeding occurs in rock holes, wooden barrels, concrete tanks, wells and other artificial containers.
- Larvae are rarely found in natural water bodies such as ditches or drain canals. Larvae are able to survive in conditions of high salinity and polluted waters.
- Hibernation takes place during the larval stage.

## Biting habits

- *Culiseta longiareolata* do not enter human dwellings and rarely bite humans.
- *Culiseta longiareolata* prefer feeding on birds.


# Orthopodomyia pulcripalpis

Native



## Morphological characteristics

1. Typical for this genus: length of tarsomere 1 of the foreleg longer than tarsomeres 2 to 5 together and tarsomere 4 of forelegs is shorter.
2. Palps nearly half as long as the proboscis.
3. White ring on apical half of the proboscis.
4. Scutum with three pairs of narrow white stripes.
5. Legs covered in black scales with a metallic shine. Has white spots on the joint between femur and tibia.
6. Dark brown scales on the wing veins.



© Disa Eklöf/SVA

## Easily-confused species

- This species is generally not confused with any of the current invasive mosquito species in Europe.
- This is the only species of this genus present in Europe.

## Status in Europe

Native

## Distribution

Palaearctic, mostly present around the Mediterranean area. The only species of this genus present in Europe.

## Ecology (habitat, breeding sites)

- The species is closely associated with forest.
- Larvae breed in tree holes and holes in tree roots. The species prefers large tree holes with water permanently present (e.g. *Platanus* spp.).
- The species overwinter in the larval stage and can survive when water surfaces freeze.

## Biting habits


Females are mainly ornithophilic and rarely bite humans; they are most active during the day in shaded places.

# Uranotaenia unguiculata



## Morphological characteristics

1. Genus characterised by the anal vein which is sharply bent at the end.
2. Proboscis swollen at the apex.
3. Dark head with silver line along the eye margins.
4. Silvery white line running across the lateral margin of the scutum (see photograph).
5. White scales on the tip of femur and tibia.
6. Pale longitudinal stripe on the anterior surface of the femora.
7. Terga covered with dark-brown iridescent scales.
8. Very small size.



© Disa Eklof/SVA

## Easily-confused species

This species is generally not confused with any of the current invasive mosquito species in Europe.

## Status in Europe

Native

## Distribution

Mediterranean region and as far north as Luxembourg.

There are only three species of *Uranotaenia* present in the western Palearctic region:

*Uranotaenia balfouri* is present in Morocco and *Uranotaenia mashonaensis* in Israel. *Uranotaenia unguiculata* is present all around the Mediterranean.

## Ecology (habitat, breeding sites)

- Preferred breeding sites of this species are pools, ditches, canals or shallow shores of lakes with stagnant or slow running water that are rich in aquatic vegetation. Larvae prefer fresh water.
- The adults are most abundant in late summer.
- The species overwinters as adults in shelters, caves, piles of cut reeds, or dense vegetation.

## Biting habits

Feeds on amphibians and rarely bites humans or mammals.

# References

1. Becker N, Petric D, Zgomba M, Boase C, Madon MB, Dahl C, et al. Mosquitoes. Identification, Ecology and Control: Springer Nature Switzerland AG 2020; p. 570.
2. Wilkerson RC, Linton YM, Fonseca DM, Schultz TR, Price DC, Strickman DA. Making Mosquito Taxonomy Useful: A Stable Classification of Tribe Aedini that Balances Utility with Current Knowledge of Evolutionary Relationships. *PLoS One*. 2015;10(7):e0133602.
3. Reinert JF. New classification for the composite genus *Aedes* (Diptera : Culicidae : Aedini), elevation of subgenus *Ochlerotatus* to generic rank, reclassification of the other subgenera, and notes on certain subgenera and species. *J Am Mosquito Contr*. 2000;16(3):175-88.
4. Reinert JF, Harbach RE, Kitching IJ. Phylogeny and classification of Aedini (Diptera: Culicidae), based on morphological characters of all life stages. *Zoological Journal of the Linnean Society*. 2004;142(3):289-368.
5. Reinert JF, Harbach RE, Kitching IJ. Phylogeny and classification of *Finlaya* and allied taxa (Diptera: Culicidae: Aedini) based on morphological data from all life stages. *Zoological Journal of the Linnean Society*. 2006;148(1):1-101.
6. Reinert JF, Harbach RE, Kitching IJ. Phylogeny and classification of *Finlaya* and allied taxa (Diptera: Culicidae: Aedini) based on morphological data from all life stages. *Zoological Journal of the Linnean Society*. 2008;148:1-101.
7. Reinert JF, Harbach RE, Kitching IJ. Phylogeny and classification of tribe Aedini (Diptera: Culicidae). *Zoological Journal of the Linnean Society*. 2009;157(4):700-94.

Further reading. Wilkerson RC, Linton YM, Stricker D. Mosquitoes of the world: Johns Hopkins University Press; 2021.

This report was commissioned by the European Centre for Disease Prevention and Control (ECDC) Specific contract No 2 ECD.10468 and No 3 ECD.11885 implementing inter-agency framework contract for services No ECDC/2019/020 (VectorNet). The report was coordinated by Olivier Briët, and produced with contributions from Anders Lindström, Disa Eklöf, Nick Van Hul, Marieta Braks, Arjan Stroo, Dušan Petrić, Francis Schaffner and Wim Van Bortel.

The key is based on illustrations made by Disa Eklöf and photos by Anders Lindström from the National Veterinary Institute/SVA. Anders Lindström owns the copyright of all photographs and Disa Eklöf owns the copyright of all illustrations. These photographs and illustrations are public and may be reproduced, adapted and/or distributed, totally or in part, irrespective of the means and/or the formats used, provided that the copyright owners are always acknowledged as the original source of the material. Such acknowledgement must be included in each copy of the material.

