

# DANILON® Équidos

## QUALITATIVE AND QUANTITATIVE COMPOSITION

Suxibuzone (microcapsulated) 150 mg

Pharmaceutical form  
Granules



## CLINICAL PARTICULARS

Target species  
Horses

### Indications of use, specifying the target species

Horses: Inflammations. Tendonitis. Contusions. Arthritis. Bursitis, Osteoarthritis, Laminitis, Sprains and soft tissue inflammation.

- Product indicated generally for treatment of pain and inflammation associated with musculo-skeletal conditions in horses.

### Contraindications

Special warnings for each target species

### Adverse reactions (frequency and seriousness )

Use during pregnancy, lactation or lay

### Posology and route of administration

Loading Dose: 2 sachets/horse, twice daily (2 in the morning and 2 in the evening), for 2 days.

Maintenance Dose: 1 sachet/horse, twice daily (1 in the morning and 1 in the evening), for 3 - 6 days.

Young Horses and Ponies:

Administer half the recommended doses for adult horses.

### Withdrawal period(s)

Do not administer to horses which meat is used for human consumption

### Pharmaceutical particulars

Shelf life

<Shelf-life of the veterinary medicinal product as packaged for sale> 4 years

<Shelf-life after first opening the immediate packaging >

<Shelf-life after dilution or reconstitution according to directions >

### Special precautions for storage

Pack containing 18 sachets of 10 g.

Pack containing 60 sachets of 10 g.

<Not all pack sizes may be marketed.>

### The marketing authorization number (s)

EXPORT-1