



**An evaluation of the vegetation and plant diversity
along the proposed route between the Eskom
Merensky and Foskor substations**

February 2012



An evaluation of the vegetation and plant diversity along the proposed route between the Eskom Merensky and Foskor substations

Prepared for:

Nsovo Environmental Consulting

by

G.J. Bredenkamp DSc., Pr.Sci.Nat.

EcoAgent CC

PO Box 23355

Monument Park

0181

Tel 012 4602525

Fax 012 460 2525

Cell 082 5767046

March 2012



Contents

DECLARATION OF INDEPENDENCE	5
EXECUTIVE SUMMARY	6
THE PROPOSED DEVELOPMENT	8
2. ASSIGNMENT	8
3. RATIONALE	10
4. STUDY AREA.....	11
4.1. General.....	11
6. METHODS	15
6.1 Vegetation and habitat survey.....	15
6.2 Conservation Priority / Sensitivity.....	16
7. RESULTS:	18
7.1 Vegetation Classification.....	18
7.2 Description of the vegetation types	21
Sekhukhune Plains Bushveld (Vulnerable)	21
Species of Conservation Concern	24
Conclusion	25
2. Sekhukhune Mountain Bushveld.....	26
Species of Conservation Concern	28
Conclusion	29
3. Ohrigstad Mountain Bushveld	29
a. Plains Bushveld.....	30
b. Mountain Slope Bushveld.....	32
Species of Conservation Concern	34
Conclusion	35
4. Lydenburg Thornveld.....	36
Species of Conservation Concern	38
Conclusion	39
5. Pong Dolomite Mountain Bushveld	40
Species of Conservation Concern	42
Conclusion	42

6. Northern Escarpment Quartzite Sourveld.....	43
7. Northern Mistbelt Forest	43
8. Tzaneen Sour Bushveld.....	44
Species of Conservation Concern	45
Conclusion	45
9. Granite Bushveld	47
Species of Conservation Concern	49
Conclusion	49
10. Lowveld Rugged Mopaneveld.....	51
Species of Conservation Concern	52
Conclusion	53
11. Phalaborwa-Timbavati Mopaneveld	53
Species of Conservation Concern	55
Conclusion	55
12 River and Spruit Systems	56
Conclusion	58
8 DISCUSSION	59
11. REFERENCES.....	61
ABRIDGED CURRICULUM VITAE: GEORGE JOHANNES BREDEKAMP	64



DECLARATION OF INDEPENDENCE

I, George Johannes Bredenkamp, Id 4602105019086, declare that I:

- Am the owner of Eco-Agent CC, CK 95/37116/23
- am committed to biodiversity conservation but concomitantly recognize the need for economic development. Whereas we appreciate the opportunity to also learn through the processes of constructive criticism and debate, we reserve the right to form and hold our own opinions and therefore will not willingly submit to the interests of other parties or change our statements to appease them;
- abide by the Code of Ethics of the S.A. Council for Natural Scientific Profession;
- Act as independent specialist consultants respectively in the fields of ecology, vegetation science and botany,
- am assigned as specialist consultant by Nsovo Environmental Consulting for the proposed project "An evaluation of the vegetation and plant diversity along the proposed route between the Eskom Merensky and Foskor substations" described in this report;
- do not have or will not have any financial interest in the undertaking of the activity other than remuneration for work performed;
- have or will not have any vested interest in the proposed activity proceeding;
- have no and will not engage in conflicting interests in the undertaking of the activity;
- undertake to disclose to the client and the competent authority any material information that have or may have the potential to influence the decision of the competent authority required in terms of the Environmental Impact Assessment Regulations 2006;
- will provide the client and competent authority with access to all information at our disposal, regarding this project, whether favourable or not.



C J

EXECUTIVE SUMMARY

The vegetation of all four alternatives was investigated. From the desktop study, confirmed by the field survey, option 3, which runs from Burgersfort to Ohrigstad along the R555, was eliminated. This is because the route along the R555 runs for most of the way in a narrow valley, with the Mabitsana River and the tarred R555 in this valley. The line will have to run for most of the way on the sensitive mountain foot slopes and cross the river and road several times. Furthermore, many irrigated agricultural enterprises occur in the Ohrigstad area, stretching all the way to Marapeng. This mosaic of narrow river valley, river, mountain slopes and agriculture where-ever the valley is a bit broader, causes the route to be unsuitable. From an ecological perspective both the riverine vegetation and the vegetation of the mountain slopes have a high ecological sensitivity. Therefore this entire valley forms an ecologically sensitive ecosystem. This is also a much longer route.

Furthermore, from the desktop study, confirmed by the field survey, option 4, was eliminated. The line of this option runs through nine vegetation types, and over very high and steep mountains of Sekhukhune Mountain Bushveld and Ohrigstad Mountain Bushveld with two endangered ecosystems (Sekhukhune Mountainlands and Sekhukhune Norite Bushveld, SANBI & DEAT 2009), Pong Dolomite Mountain Bushveld with endangered Malmani Karstland (SANBI & DEAT 2009), the vulnerable Northern Escarpment Quartzite Sourveld (Mucina & Rutherford (2006), Northern Mistbelt Forest area and the vulnerable Tzaneen Sour Bushveld (SANBI & DEAT 2009). Especially the Great Escarpment area consists of very rugged and high mountains, resulting in a very difficult route with several threatened ecosystems.

Alternative Routes 1 and 2 were further investigated in more detail by field surveys. The vegetation along these routes is described in detail, including the protected and red data species. Medicinal plants and aliens and weeds are indicated.

The most difficult part of the route is from the Merensky substation through Orichstad Mountain Bushveld which is an extremely mountainous area with sensitive vegetation. This part of the line transects quite sensitive vegetation and it is suggested that a walkthrough in this area is essential.

The most serious limitation on the Lowveld plains where the line transects the Granite Lowveld vegetation type, is the abundance of the protected tree *Sclerocarya birrea*. It is certain that several of these trees will be in the way of the transect.

Locally are also the river and spruit crossings. No river or spruit is very wide, so the lines can easily cross these rivers or spruits systems. Care should be taken to place pylons adequately away from river or spruit banks, avoiding any damage to the banks or water courses. Erosion should be avoided at all times.

Another factor in this area is that large properties are game farms and lodges. These areas are effectively conserved by the owners, and it is realised that the public participation is an important issue. After finalisation of the exact transect, a walkthrough will have to confirm any issues regarding vegetation.

From an ecological perspective, Alternative 1 is the preferred route.

THE PROPOSED DEVELOPMENT

Eskom proposed the development of a new power line from the Merensky substation near Steelpoort to the Foskor substation near Phalaborwa. This power line is more than 120 km long. Alternative routes were proposed.

2. ASSIGNMENT

EcoAgent Ecological Consultants CC was appointed by Nsovo Environmental Consulting to assess the vegetation and flora for the entire route of proposed routes. This assignment is in accordance with the 2010 EIA Regulations (No. R. 543-546, Department of Environmental Affairs and Tourism, 18 June 2010) emanating from Chapter 5 of the National Environmental Management Act, 1998 (Act No. 107 of 1998).

The assignment is interpreted as follows: Compile a study of the vegetation and flora of the route, with emphasis on Red Data or Protected plant species that occur or may occur along the route. In order to compile this, the following had to be done:

2.1. Initial preparations:

- Obtain relevant maps and information on the natural environment of the concerned area.
- This includes information on Red Data plant species that may occur in the area.
- Obtain the Vegetation Types (Mucina & Rutherford, 2006) along the routes.

2.2. Vegetation and habitat survey:

- Use the Vegetation Types as basis for ecosystem delimitation.
- List the plant species (trees, shrubs, grasses and herbaceous species) present in the ecosystem recognised.
- Identify potential red data plant species, alien plant species, and medicinal plants.

2.3. Plant community delimitation and description

- Describe the habitat and vegetation.
- Determine the sensitivity of the site for biodiversity, veld condition and presence of rare or protected species.

2.4. General

- Identify and describe particular ecologically sensitive areas.
- Identify problem areas in need of special treatment or management, e.g. bush encroachment, erosion, water pollution, degraded areas, reclamation areas.
- This includes information on Red Data plant species that may occur in the area.

3. RATIONALE

It is widely recognised that it is of the utmost importance to conserve natural resources in order to maintain ecological processes and life support systems for plants, animals and humans. To ensure that sustainable development takes place, it is therefore important that the environment is considered before relevant authorities approve any development. This led to legislation protecting the natural environment. The Environmental Conservation Act (Act 73 of 1989), the National Environmental Management Act, 1998 (NEMA) (Act 107 of 1998) and the National Environmental Management Biodiversity Act, 2004. (Act 10 Of 2004) ensure the protection of ecological processes, natural systems and natural beauty as well as the preservation of biotic diversity in the natural environment. They also ensure the protection of the environment against disturbance, deterioration, defacement or destruction as a result of man-made structures, installations, processes or products or human activities. A draft list of Threatened Ecosystems was published (Government Gazette 2009) as part of the National Environmental Management Biodiversity Act, 2004 (Act 10 Of 2004). These Threatened Ecosystems are described by SANBI & DEAT (2009).

All components of the ecosystems (physical environment, vegetation, animals) of a site are interrelated and interdependent. A holistic approach is therefore imperative to include effectively the development, utilisation and where necessary conservation of the given natural resources in an integrated development plan, which will address all the needs of the modern human population (Bredenkamp & Brown 2001).

It is therefore necessary to make an inventory of the plant communities and biodiversity on the alternative routes, in order to evaluate the biodiversity and possible rare species. This inventory should then serve as a scientific and ecological basis for the planning exercises.

4. STUDY AREA

4.1. General

The different options suggested for the Merensky - Foskor Eskom power line are indicated in Figure 1. The northern parts of all options will traverse through conservation areas (Figure 2). All options start at the Merensky substation near Steelpoort. As the area is very large, many spruit systems are found along the routes (Figure 3). A few larger rivers or spruits are mentioned, but the numerous smaller spruit systems will have to be identified during a walk-down.

Option 1 (the preferred option) will run along the Steelpoort – Burgersfort road (R555) but before reaching Burgersfort it will turn north-eastwards and cross the rugged mountainous area towards the Strydom tunnel on the R36. It will then cross the mountains east of the Strydom tunnel and run towards Mica and from there towards the Foskor substation south of Phalaborwa.

Option 2 will follow the same route as option 1 up to the Mica area, but will then turn more east to run south and east of option 1, in the direction of Hoedspruit, and then turn northwards to the Foskor substation.

Option 3 will initially also follow the same route as option 1 for a short distance, but at Burgersfort it will turn eastwards and follow the R555 to Ohrigstad and further on to the Strydom tunnel. From here on the route is similar to that of option 1.

Option 4 will run northwards from the Merensky substation, over undulating and mountainous area towards Penge and then over the rugged mountains over the great escarpment to cross the R36 between Trichardsdal and Diputhi and then run through the Kapama / Madrid Nature Reserve area toward Phalaborwa.




Figure 1: A locality map showing the different options for the Merensky Foskor Eskom power line




Figure 2: A locality map showing the different options for the Merensky Foskop Eskom power line with conservation and sensitive areas




Figure 3: The hydrology of the area showing major river systems and numerous smaller spruits along all the route options

6. METHODS

6.1 Vegetation and habitat survey

A desktop study was made on the vegetation and flora fauna and general ecology and ecological sensitivity of the area along all options suggested for this power line. A site visit followed the desktop study.

The routes were driven on 24 and 25 October 2011 by Prof G.J. Bredenkamp, a delegation of Nsovo Environmental Consulting and also a delegation of Eskom. Staff members of Eskom gave guidance in the field on the location of the various options. Parts of the route, especially the south-western mountainous parts, were again visited during February 2012.

The vegetation of the route was stratified into relatively homogeneous units based on Vegetation Type Units (Mucina & Rutherford 2006). Regular stops were made in each vegetation unit identified, to record vegetation and plant species present and also on the conservation status, sensitivity and condition of the vegetation. Special features were identified as major river crossings, wetlands, rocky ridges or any other features considered to be of importance for the biodiversity assessment.

The general vegetation of the unit was described using both the desktop study and the field observations. For the particular vegetation type a description of the dominant and characteristic species was made at several sites within each Vegetation Type unit. These descriptions were based on total floristic composition, following established vegetation survey techniques (Mueller-Dombois & Ellenberg 1974; Westhoff & Van der Maarel 1978). Data recorded included a list of the plant species present, including trees, shrubs, grasses and forbs. Comprehensive species lists were therefore derived for each plant community / ecosystem present on the site. These vegetation survey methods have been used as the basis of a national vegetation survey of South Africa (Mucina *et al.* 2000) and are considered to be an efficient method of describing vegetation and capturing species information. Notes were additionally made of any other features that might have an ecological influence.



The identified systems are not only described in terms of their plant species composition, but also evaluated in terms of the potential habitat for red data plant species.

Red data plant species for the area were obtained from the SANBI data bases, with updated threatened status, (Raimondo *et al* 2009). These lists were then evaluated in terms of habitat available on the site, and also in terms of the present development and presence of man in the area.

Protected trees are according to the list provided by the National Forests Act 1998 (Act 84 of 1998). Other protected plants are according to the National Environmental Management: Biodiversity Act, 2004 (Act No. 10 of 2004), or provincial regulations.

Alien invasive species, according to the Conservation of Agricultural Resources Act (Act No.43 of 1983) as listed in Henderson (2001), are indicated.

Medicinal plants are indicated according to Van Wyk, Van Oudthoorn & Gericke (1997), these are mentioned in the species lists.

6.2 Conservation Priority / Sensitivity

The following **conservation priority / ecological sensitivity** categories were used for each site:

High: Ecologically sensitive and valuable land with high species richness and/or sensitive ecosystems or red data species that should be conserved and no developed allowed.

Medium-high: Land where sections are disturbed but which is in general ecologically sensitive to development/disturbances.

Medium: Land on which low impact development with limited impact on the vegetation / ecosystem could be considered for development. It is recommended that certain portions of the natural vegetation be maintained as open space.

Medium-low: Land of which small sections could be considered to conserve but where the area in general has little conservation value.

Low: Land that has little conservation value and that could be considered for developed with little to no impact on the vegetation.

Species status

Plant species recorded in each plant community with an indication of the status of the species by using the following symbols:

A = Alien woody species

D = Dominant

d = subdominant

G = Garden or Garden Escape

M = Medicinal plant species

P = Protected trees species

p = provincially protected species

RD = Red data listed plant

W = weed

The field observations were supplemented by literature studies from the area (Bredenkamp 1982, Gertenbach 1983a, 1983b, Mathews 1991, Siebert 2001, Siebert *et al.* 2002a, 2002b, 2002c, 2002d 2002e and 2003).



7. RESULTS:

7.1 Vegetation Classification

According to the new vegetation map of South Africa (Mucina & Rutherford 2006) the routes transect the following vegetation types, Threatened Ecosystem Status is according to SANBI & DEAT (2009):

Vegetation Type	Route	Conservation status (Mucina & Rutherford 2006)	Threatened Ecosystems (SANBI & DEAT 2009)	Threatened Status
1. Sekhukhune Plains Bushveld	1, 3, 4	Vulnerable		
2. Sekhukhune Mountain Bushveld	4	Least Threatened	Sekhukhune Mountainlands Sekhukhune Norite Bushveld	Endangered Endangered
3. Ohrigstad Mountain Bushveld	1, 3, 4	Least Threatened	Sekhukhune Mountainlands	Endangered
4. Lydenburg Thornveld	1	Vulnerable		
5. Pong Dolomite Mountain Bushveld	1, 3, 4	Least Threatened	Malmani Karstlands	Endangered
6. Northern Escarpment Quartzite Sourveld	4	Vulnerable		
7. Northern Mistbelt Forest	4	Least Threatened		
8. Tzaneen Sour Bushveld	4 (1, 2)	Least Threatened	Tzaneen Sour Bushveld	Vulnerable
9. Granite Bushveld	1, 2, 4	Least Threatened		
10. Lowveld Rugged Mopaneveld	1, 2	Least Threatened		
11. Phalaborwa-Timbavati Mopaneveld	1, 4	Least Threatened		
12. River Crossings	all			






Figure 4: The vegetation types (Mucina & Rutherford 2006) along the optional routes

From the desktop study, confirmed by the field survey, option 3, which runs from Burgersfort to Ohrigstad along the R555, was eliminated. This is because the route along the R555 runs for most of the way in a narrow valley, with the Mabitsana River and the tarred R555 in this valley. The line will have to run for most of the way on the sensitive mountain foot slopes and cross the river and road several times. Furthermore, many irrigated agricultural enterprises occur in the Ohrigstad area, stretching all the way to Marapeng. This mosaic of narrow river valley, river, mountain slopes and agriculture where-ever the valley is a bit broader, causes the route to be unsuitable. From an ecological perspective both the riverine vegetation and the vegetation of the mountain slopes have a high ecological sensitivity. Therefore this entire valley forms an ecologically sensitive ecosystem. This is also a much longer route.

Furthermore, from the desktop study, confirmed by the field survey, option 4, was eliminated. The line of this option runs through nine vegetation types, and over very high and steep mountains of Sekhukhune Mountain Bushveld and Ohrigstad Mountain Bushveld with two endangered ecosystems (Sekhukhune Mountainlands and Sekhukhune Norite Bushveld, SANBI & DEAT 2009), Pong Dolomite Mountain Bushveld with endangered Malmani Karstland (SANBI & DEAT 2009), the vulnerable Northern Escarpment Quartzite Sourveld (Mucina & Rutherford (2006), Northern Mistbelt Forest area and the vulnerable Tzaneen Sour Bushveld (SANBI & DEAT 2009). Especially the Great Escarpment area consists of very rugged and high mountains, resulting in a very difficult route with several threatened ecosystems.

7.2 Description of the vegetation types

Sekhukhune Plains Bushveld (Vulnerable)

The Merensky substation is located in the Sekhukhune Plains Bushveld. Although the vegetation of these plains falls within the Sekhukhune Centre of plant endemism (Siebert 2001, Siebert *et al.* 2002a-e) this area is highly transformed by many villages and their agricultural fields. Within the study area this bushveld is restricted to the valley floors of the rivers that dissect the mountains. These areas are heavily grazed and often not in prime condition. This resulted in Mucina & Rutherford (2006) labelling the conservation status of this vegetation as Vulnerable.

Large parts of these plains are dominated by *Dichrostachys cinerea*, *Acacia tortilis*, *Acacia mellifera* and *Acacia nilotica*. Other plant species found here include the trees *Boscia foetida*, *Euclea linearis*, *Searsia batophylla* (along spruities and dongas) with the forbs *Felicia clavipilosa*, *Hermannia odorata*, *Gisekia africana*, *Melhania rehmannii* and the grasses *Aristida congesta*, *Enneapogon cenchroides*, *Urochloa mosambicensis*. Alien plant species are often found close to villages or along roads and tracks.

The following species were listed for this plant community:

TREES AND SHRUBS

<i>Acacia gerrardii</i>		<i>Grewia bicolor</i>	
<i>Acacia mellifera</i>	d	<i>Grewia flava</i>	
<i>Acacia nilotica</i>	d	<i>Lantana camara</i>	A
<i>Acacia tortilis</i>	D	<i>Melia azedarach</i>	A
<i>Agave americana</i>	A	<i>Rhigozum obovatum</i>	
<i>Aloe castanea</i>	p	<i>Sarcostemma viminale</i>	
<i>Aloe cryptopoda</i>	p	<i>Schotia latifolia</i>	RD
<i>Aloe globuligemma</i>	p	<i>Sclerocarya birrea</i>	P
<i>Balanites maughamii</i>	P	<i>Searsia batophylla</i>	RD
<i>Boscia foetida</i>		<i>Searsia engleri</i>	
<i>Dichrostachys cinerea</i>	D	<i>Tinnea rhodesiana</i>	
<i>Ehretia rigida</i>		<i>Triaspis glaucophylla</i>	
<i>Euphorbia tirucalli</i>		<i>Ziziphus mucronata</i>	M

GRASSES

<i>Aristida adscensionis</i>		<i>Eragrostis lehmanniana</i>	
<i>Aristida congesta s. barbicollis</i>		<i>Eragrostis superba</i>	
<i>Cenchrus ciliaris</i>		<i>Fingerhuthia africana</i>	
<i>Chloris virgata</i>		<i>Melinis repens s. repens</i>	
<i>Dactyloctenium aegyptium</i>		<i>Panicum maximum</i>	
<i>Enneapogon cenchroides</i>		<i>Sporobolus ioclados</i>	
<i>Enneapogon scoparius</i>		<i>Stipagrostis hirtigluma</i>	
<i>Eragrostis heteromera</i>		<i>Themeda triandra</i>	

Tragus berteronianus

Urochloa mosambicensis

FORBS

Abutilon angulatum

Kohautia cynanchica

Acalypha indica

Lantana rugosa

Achyranthes aspera v. sicula

Kleinia longiflora

Asparagus suaveolens

Leonotis ocymifolia

Bidens bipinnata W

Melhania acuminata

Blepharis integrifolia

Melhania rehmannii

Clerodendrum ternatum

Monechma divaricatum

Corchorus asplenifolius

Ocimum americanum

Datura stramonium WM

Pavonia burchellii

Felicia clavipilosa

Phyllanthus maderaspatensis

Flaveria bidentis W

Pollichia campestris

Galenia sarcophylla

Schkuhria pinnata W

Geigeria burkei

Seddera fruticosa

Gossypium herbaceum

Sesamum triphyllum W

Hermannia modesta

Sesbania bispinosa

Hermbstaedtia odorata

Sida alba

Hibiscus caesius

Solanum panduriforme W

Hibiscus micranthus

Tephrosia purpurea

Jamesbrittenia atropurpurea

Tribulus terrestris W

Jatropha latifolia

Vernonia poskeana

Justicia flava

Waltheria indica W

Justicia protracta s. rhodesiana

Zinnia peruviana W

Sekhukhune Plains Bushveld			
Status	Dense to Disturbed plains bushveld		
Soil	Clay-loam	Rockiness %	5-25
Conservation priority:	Medium	Sensitivity:	Medium
Agricultural potential:	Medium-Low	Need for rehabilitation	Medium



Dominant spp.	<i>Acacia tortilis</i> , <i>Acacia mellifera</i> , <i>Dichrostachys cinerea</i> , <i>Euclea linearis</i>
----------------------	----------------------------------------------------------------------------------------------------------

Species of Conservation Concern

A list of Species of Conservation Concern for the Grid 2627BB was obtained from the database on the SANBI website. Threatened species are those that are facing high risk of extinction, indicated by the categories Critically Endangered (CE), Endangered (EN) and Vulnerable (VU). Species of Conservation Concern include the Threatened Species, but additionally have the categories Near Threatened (NT), Data Deficient (DD), Critically Rare (CR), Rare (R) and Declining (D). This is in accordance with the new Red List for South African Plants (Raimondo *et al.* 2009).

The following species of conservation concern were previously recorded from the Grid 2430CA (SANBI, POSA website):

Species	Status
<i>Dicliptera fruticosa</i> K.Balkwill	NT
<i>Elaeodendron transvaalense</i> (Burt Davy) R.H.Archer	NT
<i>Lydenburgia cassinoides</i> N.Robson	NT
<i>Adenia fruticosa</i> Burt Davy subsp. <i>fruticosa</i>	NT
<i>Searsia sekhukuniensis</i> (Moffett) Moffett	Rare
<i>Combretum petrophilum</i> Retief	Rare
<i>Euphorbia sekukuniensis</i> R.A.Dyer	Rare
<i>Searsia batophylla</i> (Codd) Moffett	VU
<i>Zantedeschia jucunda</i> Letty	VU
<i>Gladiolus sekukuniensis</i> P.J.D.Winter	VU
<i>Acacia sekhukuniensis</i> P.J.H.Hurter	CR
<i>Delosperma rileyi</i> L.Bolus	DDD
<i>Asparagus intricatus</i> (Oberm.) Fellingham & N.L.Mey.	DDT
<i>Acalypha caperonioides</i> Baill. var. <i>caperonioides</i>	DDT
<i>Myrothamnus flabellifolius</i> Welw.	DDT
<i>Ilex mitis</i> (L.) Radlk. var. <i>mitis</i>	Declining
<i>Drimia altissima</i> (L.f.) Ker Gawl.	Declining
<i>Hypoxis hemerocallidea</i> Fisch., C.A.Mey. & Avé-Lall.	Declining
<i>Eulophia speciosa</i> (R.Br. ex Lindl.) Bolus	Declining

Searsia batophylla, *Hypoxis hemerocallidea* and *Eulophia speciosa* were observed within the transect area. For most of the other species the plains habitat is not suitable, they are present on the mountain areas of Sekhukhuneland.

Balanites maughamii and *Sclerocarya birrea* are nationally protected trees observed along the route while the *Aloe* species are all provincially protected.

Conclusion

The vegetation within on the plains are quite disturbed, there are often villages, roads, tracks and current or old agricultural fields present. As the pylons of the power line will have a relatively small footprint, the impact on the vegetation will be small. However, due to the presence of red data and possibly protected plant species, a walkthrough is recommended for this area, to ensure that sensitive areas are excluded for construction of pylons.



Figure 5: Sekhukhune Plains Bushveld in the foreground and Sekhukhune Mountain Bushveld in the Background

2. Sekhukhune Mountain Bushveld

This Open Mountain Bushveld occurs patchy throughout the area, the soils contain high concentrations of heavy metals and the area is often prone to mining operations. In this study very small part of this Bushveld is affected, the largest piece is just north of Steelpoort along route option 4, which was already eliminated. However, being a very sensitive ecosystem, due to several endemic and threatened species, the description is included.

Trees and Shrubs

<i>Acacia ataxacantha</i> ,		<i>Elephantorrhiza praetermissa</i>	
<i>Acacia gerrardii</i>		<i>Elaeodendron transvaalense</i>	RD
<i>Acacia mellifera s. detinens</i>		<i>Euclea linearis</i>	
<i>Acacia nigrescens</i> ,		<i>Euclea undulata</i>	M
<i>Acacia nilotica</i>		<i>Grewia flava</i>	
<i>Acacia senegal v. leiorrhachis</i>		<i>Grewia vernicosa</i>	
<i>Acacia senegal v. rostrata</i>		<i>Hippobromus pauciflorus</i>	
<i>Acacia tortilis s. heteracantha</i>		<i>Kirkia wilmsii</i>	
<i>Aloe arborescens</i>	p	<i>Maerua cafra</i>	
<i>Aloe castanea</i>	p	<i>Maytenus undata</i>	
<i>Aloe cryptopoda</i>	p	<i>Lydenburgia cassinoides</i>	RDP
<i>Boscia foetida</i>		<i>Ormocarpum trichocarpum</i>	
<i>Brachylaena ilicifolia</i>		<i>Ozoroa sphaerocarpa</i>	
<i>Carissa bispinosa</i>		<i>Rhoicissus sekhukhuniensis</i>	
<i>Celtis africana</i>		<i>Sclerocarya birrea</i>	P
<i>Combretum apiculatum</i>		<i>Searsia keetii</i>	
<i>Combretum hereroense</i>		<i>Searsia sekhukhuniensis</i>	RD
<i>Combretum molle</i>		<i>Searsia wilmsii</i>	
<i>Combretum petrophilum</i>	RD	<i>Terminalia prunioides</i>	
<i>Commiphora mollis</i>		<i>Tinnea rhodesiana</i>	
<i>Croton gratissimus</i>		<i>Vitex obovata subsp wilmsii</i>	
<i>Cussonia transvaalensis</i>			
<i>Dichrostachys cinerea</i>			

Grasses

<i>Aristida canescens</i>		<i>Eragrostis lehmanniana</i>
<i>Aristida canescens</i> ,		<i>Eragrostis superba</i>
<i>Aristida transvaalensis</i>		<i>Fingerhuthia africana</i>
<i>Bothriochloa insculpta</i>		<i>Heteropogon contortus</i>
<i>Brachiaria eruciformis</i>		<i>Loudetia simplex</i>
<i>Digitaria eriantha</i>		<i>Melinis repens</i>
<i>Diheteropogon amplexans</i>		<i>Panicum deustum</i>
<i>Elionurus muticus</i>		<i>Sporobolus ioclados</i>
<i>Enneapogon scoparius</i>		<i>Themeda triandra</i>

Forbs

<i>Abutilon angulatum</i>		<i>Kohautia cynanchica</i>	
<i>Adenia fruticosa</i>	RD	<i>Kyphocarpa angustifolia</i>	
<i>Asparagus cooperi</i>		<i>Melhania rehmannii</i>	
<i>Asparagus suaveolens</i>		<i>Merwillia plumbea</i>	RD
<i>Barleria kaloxytonea</i>		<i>Monechma divaricatum</i>	
<i>Barleria saxatilis</i>		<i>Myrothamnus flabellifolius</i>	RD
<i>Berkheya insignis</i>		<i>Ocimum americanum</i>	
<i>Blepharis aspera</i>		<i>Phyllanthus glaucophyllus</i>	
<i>Blepharis integrifolia</i>		<i>Polygala hottentotta</i>	
<i>Clerodendrum ternatum</i>		<i>Ptychlobium plicatum</i>	
<i>Commelina africana</i>		<i>Rhynchosia minima</i>	
<i>Corchorus asplenifolius</i>		<i>Sansevieria hyacinthoides</i>	
<i>Crabbea angustifolia</i>		<i>Seddera capensis</i>	
<i>Cyphostemma coddii</i>		<i>Senna italica</i>	
<i>Ectadiopsis oblongifolia</i>		<i>Stylochiton natalensis</i>	
<i>Euphorbia enormis</i>		<i>Stylochiton sp</i>	
<i>Euphorbia schinzii</i>		<i>Syncolostemon concinnus</i>	
<i>Evolvulus alsinoides</i>		<i>Tephrosia purpurea</i>	
<i>Geigeria burkei</i>		<i>Tetradenia brevispicata</i> ,	
<i>Gerbera jamesonii</i>		<i>Tragia dioica</i>	
<i>Hibiscus aethiopicus</i>		<i>Waltheria indica</i>	W
<i>Hypoestes forskalii</i>		<i>Xerophyta retinervis</i>	

Sekhukhune Mountain Bushveld			
Status	Dense to Disturbed mountain bushveld		
Soil	Clay-loam	Rockiness %	5-25
Conservation priority:	High	Sensitivity:	High
Agricultural potential:	Low	Need for rehabilitation	Low
Dominant spp.	<i>Combretum apiculatum</i> , <i>Grewia vernicosa</i> , <i>Dichrostachys cinerea</i> , <i>Euclea linearis</i> and <i>Euclea undulata</i>		

Species of Conservation Concern

A Threatened species and Species of Conservation Concern list for the Grid 2627BB was obtained from the POSA database on the SANBI website. Threatened species are those that are facing high risk of extinction, indicated by the categories Critically Endangered (CE), Endangered (EN) and Vulnerable (VU). Species of Conservation Concern include the Threatened Species, but additionally have the categories Near Threatened (NT), Data Deficient (DD), Critically Rare (CR), Rare (R) and Declining (D). This is in accordance with the new Red List for South African Plants (Raimondo *et al.* 2009).

The following species of conservation concern were previously recorded from the Grid 2430CA (SANBI, POSA website):

Species	Status
<i>Dicliptera fruticosa</i> K.Balkwill	NT
<i>Elaeodendron transvaalense</i> (Burt Davy) R.H.Archer	NT
<i>Lydenburgia cassinoides</i> N.Robson	NT
<i>Adenia fruticosa</i> Burt Davy subsp. <i>fruticosa</i>	NT
<i>Searsia sekhukhuniensis</i> (Moffett) Moffett	Rare
<i>Combretum petrophilum</i> Retief	Rare
<i>Euphorbia sekukuniensis</i> R.A.Dyer	Rare
<i>Searsia batophylla</i> (Codd) Moffett	VU
<i>Zantedeschia jucunda</i> Letty	VU
<i>Gladiolus sekukuniensis</i> P.J.D.Winter	VU



<i>Acacia sekhukhuniensis</i> P.J.H.Hurter	CR
<i>Delosperma rileyi</i> L.Bolus	DDD
<i>Asparagus intricatus</i> (Oberm.) Fellingham & N.L.Mey.	DDT
<i>Acalypha caperonioides</i> Baill. var. <i>caperonioides</i>	DDT
<i>Myrothamnus flabellifolius</i> Welw.	DDT
<i>Ilex mitis</i> (L.) Radlk. var. <i>mitis</i>	Declining
<i>Drimia altissima</i> (L.f.) Ker Gawl.	Declining
<i>Hypoxis hemerocallidea</i> Fisch., C.A.Mey. & Avé-Lall.	Declining
<i>Eulophia speciosa</i> (R.Br. ex Lindl.) Bolus	Declining

Elaeodendron transvaalense, *Combretum petrophilum*, *Lydenburgia cassinoides*, *Adenia fruticosa*, *Myrothamnus flabellifolius* and *Searsia sekhukhuniensis* were observed close to the transect area. *Merwillia plumbea*, not listed above, was also seen. The habitat is also suitable for most of the other threatened species listed.

The protected trees *Lydenburgia cassinoides* and *Sclerocarya birrea* are present.

Conclusion

The vegetation on the mountains contains several red data and protected plant species, where-ever possible the mountain areas of Sekhukhuneland should be avoided. It seems that although this type of Bushveld is prominent between Steelpoort and Burgersfort, the proposed line will seldom transect these mountains, except for Route 4, which is already eliminated. However, should any line cross this vegetation type, a walkthrough is recommended for this area, to ensure that sensitive areas are excluded for construction of pylons.

3. Ohrigstad Mountain Bushveld

Ohrigstad Mountain Bushveld is present in the Burgersfort – Ohrigstad and Penge areas. (present for options 1, 3 and 4, not for option 2). This Mountain Bushveld also has, as the Sekhukhune Mountain Bushveld, high species diversity and several plant species of conservation concern. However, this area consists of mountains and valleys, both being important for this survey. Option 3 runs for most of the way in a prominent valley, containing plains bushveld, but often has to run over the footslopes of the mountains.

Option 3, which runs from Burgersfort to Ohrigstad along the R555, was eliminated. This is because the route along the R555 runs for most of the way in a narrow valley, with the Mabitsana River and the tarred R555 in this valley. The line will have to run for most of the way on the sensitive mountain foot slopes and cross the river and road several times. Furthermore, many irrigated agricultural enterprises occur in the Ohrigstad area, stretching all the way to Marapeng. This mosaic of narrow river valley, river, mountain slopes and agriculture where-ever the valley is a bit broader, causes the route to be unsuitable. From an ecological perspective both the riverine vegetation and the vegetation of the mountain slopes have a high ecological sensitivity. Therefore this entire valley forms an ecologically sensitive ecosystem. This is also a much longer route.

The two main plant communities found in this area are: a) The Plains Bushveld in the Valleys and b) the Mountain slope Bushveld. These are described separately.

a. Plains Bushveld

The Dense Plains Bushveld is restricted to the Ohrigstad valley, the route of Option 3. This is degraded to pristine bushveld with a dense woody cover. The vegetation is dominated by *Acacia tortilis*, *Dichrostachys cinerea*, *Combretum apiculatum*, *Euclea linearis* and *Euclea undulata*, while *Eragrostis rigidior*, *Enneapogon scoparius* and *Themeda triandra* are prominent in the grass layer.

The following plant species were recorded from this plant community:

Trees and Shrubs

<i>Acacia tortilis</i>	d	<i>Euphorbia tirucalli</i>	M
<i>Aloe castanea</i>	p	<i>Grewia monticola</i>	
<i>Aloe marlothii</i>	p	<i>Gymnosporia senegalensis</i>	M
<i>Berchemia zeyheri</i>	M	<i>Hexalobus monopetalus</i>	
<i>Combretum apiculatum</i>	M	<i>Karomia speciosa</i>	
<i>Dichrostachys cinerea</i>	d	<i>Mundulea sericea</i>	M
<i>Ehretia rigida</i>	M	<i>Ormocarpum trichocarpum</i>	
<i>Euclea linearis</i>	d	<i>Peltophorum africanum</i>	M
<i>Euclea undulata</i>	dM	<i>Ptaeroxylon obliquum</i>	M
<i>Euphorbia cooperi</i>		<i>Rhoicissus tridentata</i>	M

<i>Sclerocarya birrea</i>	PM	<i>Ximenia americana</i>	M
<i>Searsia leptodictya</i>		<i>Ziziphus mucronata</i>	M
<i>Tarchonanthus camphoratus</i>	M		

Grasses

<i>Aristida congesta</i>		<i>Eragrostis superba</i>	
<i>Aristida congesta</i> subsp <i>barbicollis</i>		<i>Heteropogon contortus</i>	d
<i>Bothriochloa insculpta</i>		<i>Melinis repens</i>	
<i>Brachiaria nigropedata</i>		<i>Panicum maximum</i>	
<i>Cymbopogon excavatus</i>		<i>Pogonarthria squarrosa</i>	
<i>Digitaria eriantha</i>		<i>Setaria sphacelata</i>	
<i>Enneapogon scoparius</i>	d	<i>Themeda triandra</i>	d
<i>Eragrostis rigidior</i>	d	<i>Urochloa mosambicensis</i>	

Forbs

<i>Abutilon austroafricanum</i>		<i>Hypoestes aristata</i>	M
<i>Aloe cryptopoda</i>	p	<i>Ipomoea magnusiana</i>	
<i>Aloe fosteri</i>	p	<i>Kyphocarpa angustifolia</i>	
<i>Barleria cf guenzii</i>		<i>Pentarrhinum insipidum</i>	M
<i>Chascanum hederaceum</i>		<i>Senecio tamoides</i>	M
<i>Commelina africana</i>	M	<i>Solanum incanum</i>	M
<i>Cucumis zeyheri</i>	M	<i>Solanum nigrum</i>	WM
<i>Datura stramonium</i>	WM	<i>Solanum panduriforme</i>	M
<i>Evolvulus alsinoides</i>	M	<i>Tephrosia sp</i>	
<i>Hibiscus micrantha</i>			
<i>Hibiscus trionum</i>			

Ohrigstad Mountain Bushveld: Plains Bushveld			
Status	Dense to Disturbed plains bushveld		
Soil	sandy-loam	Rockiness %	1
Conservation priority:	Medium-High	Sensitivity:	Medium-High
Agricultural potential:	Medium to High (irrigation)	Need for rehabilitation	Low
Dominant spp.	<i>Acacia tortilis</i> , <i>Combretum apiculatum</i> , <i>Dichrostachys cinerea</i> , <i>Euclea linearis</i> and <i>Euclea undulata</i>		



Figure 6: The Dense Plains Bushveld of the area

b. Mountain Slope Bushveld

The mountain slopes, facing in all directions, as the valley curves through the mountains are covered with dense Mountain Slope Bushveld. This is often pristine mountain bushveld with a dense woody cover, though the herbaceous layer is poorly



developed due to the dense woody layer. Many woody species occur in this plant community, with *Combretum apiculatum* and *Tarchonanthus camphoratus* prominent. *Aristida congesta* subsp *barbicollis*, *Eragrostis rigidior*, *Eragrostis lehmanniana* and *Enneapogon scoparius* are the most conspicuous grass species in the scanty herbaceous layer.

The following plant species were recorded from this plant community:

Trees and Shrubs

<i>Acacia exuvialis</i>		<i>Gymnosporia senegalensis</i>	M
<i>Acacia tortilis</i>	d	<i>Hexalobus monopetalus</i>	
<i>Berchemia zeyheri</i>	M	<i>Karomia speciosa</i>	
<i>Combretum apiculatum</i>	DM	<i>Mundulea sericea</i>	M
<i>Crotalaria monteiroi</i>		<i>Opuntia ficus-indica</i>	A
<i>Dichrostachys cinerea</i>		<i>Ormocarpum trichocarpum</i>	
<i>Ehretia rigida</i>	M	<i>Pappea capensis</i>	
<i>Elaeodendron transvaalense</i>	PM	<i>Peltophorum africanum</i>	M
<i>Euclea linearis</i>	d	<i>Phyllanthus reticulatus</i>	
<i>Euclea natalensis</i>	M	<i>Ptaeroxylon obliquum</i>	M
<i>Euclea undulata</i>	dM	<i>Rhoicissus tridentata</i>	M
<i>Euphorbia ingens</i>		<i>Sclerocarya birrea</i>	P
<i>Euphorbia tirucalli</i>	M	<i>Searsia leptodictya</i>	
<i>Flueggea virosa</i>	M	<i>Tarchonanthus camphoratus</i>	dM
<i>Grewia bicolor</i>		<i>Ximenia americana</i>	M
<i>Grewia monticola</i>		<i>Ziziphus mucronata</i>	M

Grasses

<i>Aristida congesta</i> subsp <i>barbicollis</i>	d	<i>Eragrostis rigidior</i>	d
<i>Digitaria eriantha</i>		<i>Heteropogon contortus</i>	
<i>Enneapogon scoparius</i>	d	<i>Panicum maximum</i>	
<i>Eragrostis lehmanniana</i>	d	<i>Sporobolus fimbriatus</i>	

Forbs

<i>Abutilon austroafricanum</i>		<i>Asparagus</i> sp	
<i>Aloe cryptopoda</i>	p	<i>Barleria cf guenzii</i>	
<i>Aloe fosteri</i>	p	<i>Commelina africana</i>	M

<i>Evolvulus alsinoides</i>	M	<i>Pentarrhinum insipidum</i>	M
<i>Hibiscus micrantha</i>		<i>Solanum incanum</i>	M
<i>Kalanchoe paniculata</i>	M	<i>Stylosanthes fruticosa</i>	
<i>Kyphocarpa angustifolia</i>		<i>Waltheria indica</i>	W
<i>Melhania prostrata</i>			

Ohrigstad Mountain Bushveld: Mountain Slope Bushveld			
Status	Pristine mountain bushveld		
Soil	Rocky shallow sandy	Rockiness %	5-20
Conservation priority:	High	Sensitivity:	High
Agricultural potential:	Low	Need for rehabilitation	Low
Dominant spp.	<i>Combretum apiculatum</i> , <i>Euclea linearis</i> , <i>Euclea undulata</i> , <i>Tarchonanthus camphoratus</i>		

Species of Conservation Concern

A Threatened species and Species of Conservation Concern list for the Grid 3325DB was obtained from the POSA database on the SANBI website. Threatened species are those that are facing high risk of extinction, indicated by the categories Critically Endangered, Endangered and Vulnerable. Species of Conservation Concern include the Threatened Species, but additionally have the categories Near Threatened, Data Deficient, Critically Rare, Rare and Declining. This is in accordance with the new Red List for South African Plants (Raimondo *et al.* 2009).

The following species of conservation concern were previously recorded from the Grid 2430DA (SANBI, POSA website April 2011):

Species	Status
<i>Adenia fruticosa</i> Burt Davy subsp. <i>fruticosa</i>	NT
<i>Aloe fouriei</i> D.S.Hardy & Glen	DDT
<i>Ansellia africana</i> Lindl.	Declining
<i>Ceropegia distincta</i> N.E.Br. subsp. <i>verruculosa</i> R.A.Dyer	DDD

<i>Combretum petrophilum</i> Retief	Rare
<i>Dicliptera fruticosa</i> K.Balkwill	NT
<i>Dracaena transvaalensis</i> Baker	Rare
<i>Eulophia speciosa</i> (R.Br. ex Lindl.) Bolus	Declining
<i>Euphorbia sekukuniensis</i> R.A.Dyer	Rare
<i>Gladiolus macneilii</i> Oberm.	CR
<i>Gladiolus pavonia</i> Goldblatt & J.C.Manning	CR
<i>Indigofera leendertziae</i> N.E.Br.	DDT
<i>Jamesbrittenia macrantha</i> (Codd) Hilliard	NT
<i>Khadia alticola</i> Chess. & H.E.K.Hartmann	Rare
<i>Lydenburgia cassinoides</i> N.Robson	NT
<i>Ocimum tubiforme</i> (R.D.Good) A.J.Paton	CR
<i>Orbea gerstneri</i> (Letty) Bruyns subsp. <i>gerstneri</i>	Rare
<i>Pentatrachia alata</i> S.Moore	DDD
<i>Rhoicissus laetans</i> Retief	Rare
<i>Searsia batophylla</i> (Codd) Moffett	VU
<i>Thesium davidsonae</i> Brenan	VU

Lydenburgia cassinoides was found during the field visit in this area. There is suitable habitat for several of the threatened plant species along the transect of this Option of the power line, especially for *Adenia fruticosa* subsp. *fruticosa*, *Ocimum tubiforme*, *Orbea gerstneri* subsp. *gerstneri* and *Combretum petrophilum*, but none of these species were found on the site during the field visit.

Conclusion

This plant community is high in species richness with several red data species. A few individuals of the protected *Elaeodendron transvaalensis*, *Sclerocarya birrea*, *Aloe fosteri* and *Aloe cryptopoda* are found in this vegetation. The conservation value and sensitivity are regarded as being high, due to biodiversity and also due to the ecological function of the mountains and the valley as a dispersal corridor for plants and animals. There is also abundant farming activities in the valley, including irrigated crops. There will be several turning points for the line, as the road winds through the valley. It is suggested that this option not be used due to the several constraints.



Figure 7: Cleared Open Plains Bushveld in the foreground, Open Plains Bushveld in the middle and Mountain Slope Bushveld in the background

4. Lydenburg Thornveld

The Option 1 route crosses a small section of the Lydenburg Thornveld on the undulating mountain plateau area above the Strydom tunnel. This area is a wooded grassland, quite cold with frost during winter. Species such as the frost hardy *Acacia karroo*, *Acacia caffra*, *Cussonia paniculata* *Diospyros lycioides* and *Euclea crispa* are prominent.

The following plant species were recorded from this plant community:

Trees and Shrubs

<i>Acacia caffra</i>	d	<i>Euclea crispa</i>	M
<i>Acacia karroo</i>	dM	<i>Dombeya rotundifolia</i>	
<i>Acacia robusta</i>		<i>Rubus transvaalensis</i>	
<i>Cussonia paniculata</i>		<i>Ehretia rigida</i>	M
<i>Diospyros lycioides</i>		<i>Gymnosporia buxifolia</i>	M

<i>Mundulea sericea</i>	M	<i>Searsia pyroides</i>	
<i>Rhoicissus tridentata</i>	M	<i>Ziziphus mucronata</i>	M
<i>Searsia leptodictya</i>			

Grasses

<i>Aristida congesta</i>		<i>Melinis repens</i>	
<i>Aristida congesta</i> subsp <i>barbicollis</i>		<i>Microchloa caffra</i>	
<i>Aristida diffusa</i>		<i>Monocymbium ceresiiforme</i>	
<i>Bewsia biflora</i>		<i>Panicum maximum</i>	
<i>Brachiaria serrata</i>		<i>Panicum natalense</i>	
<i>Cymbopogon excavatus</i>		<i>Pogonarthria squarrosa</i>	
<i>Digitaria eriantha</i>		<i>Schizachyrium sanguineum</i>	
<i>Diheteropogon amplexans</i>		<i>Setaria sphacelata</i>	
<i>Eragrostis superba</i>		<i>Themeda triandra</i>	d
<i>Heteropogon contortus</i>	d	<i>Tristachya leucothrix</i>	

Forbs

<i>Anthospermum rigidum</i>		<i>Lippia javanica</i>	
<i>Commelina africana</i>	M	<i>Pentarrhinum insipidum</i>	M
<i>Dicoma anomala</i>		<i>Schistostephium crataegifolium</i>	
<i>Elephantorrhiza elephantina</i>	M	<i>Senecio coronatus</i>	
<i>Euphorbia clavarioides</i>		<i>Senecio microglossus</i>	
<i>Evolvulus alsinoides</i>	M	<i>Senecio tamoides</i>	M
<i>Helichrysum cephaloideum</i>		<i>Solanum incanum</i>	M
<i>Helichrysum rugulosum</i>		<i>Solanum panduriforme</i>	M
<i>Hibiscus trionum</i>		<i>Tephrosia</i> sp	
<i>Kohautia amatymbica</i>		<i>Vernonia oligocephala</i>	

Lydenburg Thornveld			
Status	Open wooded grassland		
Soil	sandy-loam	Rockiness %	1
Conservation priority:	Medium	Sensitivity:	Medium
Agricultural potential:	Medium	Need for rehabilitation	Low
Dominant spp.	<i>Acacia karroo</i> , <i>Acacia caffra</i> , <i>Euclea crispa</i>		

Species of Conservation Concern

A Threatened species and Species of Conservation Concern list for the Grid 2430CB was obtained from the POSA database on the SANBI website. Threatened species are those that are facing high risk of extinction, indicated by the categories Critically Endangered, Endangered and Vulnerable. Species of Conservation Concern include the Threatened Species, but additionally have the categories Near Threatened, Data Deficient, Critically Rare, Rare and Declining. This is in accordance with the new Red List for South African Plants (Raimondo *et al.* 2009).

The following species of conservation concern were previously recorded from the Grid 2430DA (SANBI, POSA website April 2011):

Species	Status
<i>Adenia fruticosa</i> Burt Davy subsp. <i>fruticosa</i>	NT
<i>Aloe fouriei</i> D.S.Hardy & Glen	DDT
<i>Ansellia africana</i> Lindl.	Declining
<i>Ceropegia distincta</i> N.E.Br. subsp. <i>verruculosa</i> R.A.Dyer	DDD
<i>Combretum petrophilum</i> Retief	Rare
<i>Dicliptera fruticosa</i> K.Balkwill	NT
<i>Dracaena transvaalensis</i> Baker	Rare
<i>Eulophia speciosa</i> (R.Br. ex Lindl.) Bolus	Declining
<i>Euphorbia sekukuniensis</i> R.A.Dyer	Rare
<i>Gladiolus macneillii</i> Oberm.	CR
<i>Gladiolus pavonia</i> Goldblatt & J.C.Manning	CR

<i>Indigofera leendertziae</i> N.E.Br.	DDT
<i>Jamesbrittenia macrantha</i> (Codd) Hilliard	NT
<i>Khadia alticola</i> Chess. & H.E.K.Hartmann	Rare
<i>Lydenburgia cassinoides</i> N.Robson	NT
<i>Ocimum tubiforme</i> (R.D.Good) A.J.Paton	CR
<i>Orbea gerstneri</i> (Letty) Bruyns subsp. <i>gerstneri</i>	Rare
<i>Pentatrichia alata</i> S.Moore	DDD
<i>Rhoicissus laetans</i> Retief	Rare
<i>Searsia batophylla</i> (Codd) Moffett	VU
<i>Thesium davidsonae</i> Brenan	VU

None of these species occur in this plant community as the habitat is not suitable for any of them.

Conclusion

This plant community is fairly high in species richness with no red data species or protected species found in the survey. It is also a very small area that will be crossed by the line. The construction of the line can be supported.



5. Pong Dolomite Mountain Bushveld

This is a narrow band of bushveld on dolomite, in the vicinity of the Strydom tunnel. It is situated along the western drier part of the Escarpment. The vegetation is woodland with a dense shrub layer. It occurs on the low and high mountain slopes in the area. The geology is dolomite and the shallow, rocky soils are of the Mispah soil Form. A very small, almost negligible part of this vegetation is crossed by Options 1, 3 and 4.

Pong Dolomite Mountain Bushveld			
Status	Open to dense bushveld		
Soil	sandy-loam	Rockiness %	1
Conservation priority:	Medium-High	Sensitivity:	Medium-High
Agricultural potential:	Low	Need for rehabilitation	Low
Dominant spp.	<i>Acacia nigrescens, Acacia nilotica, Combretum apiculatum</i>		

The vegetation is dominated by the woody layer with several woody species present.

Trees and Shrubs

<i>Acacia ataxacantha,</i>		<i>Combretum molle</i>	
<i>Acacia gerrardii</i>		<i>Croton gratissimus</i>	
<i>Acacia nigrescens</i>	d	<i>Cussonia spicata</i>	
<i>Acacia nilotica</i>	d	<i>Dichrostachys cinerea</i>	
<i>Acacia tortilis</i>		<i>Dombeya rotundifolia</i>	
<i>Aloe cryptopoda</i>	p	<i>Euclea crispa</i>	M
<i>Boscia albitrunca</i>	M	<i>Euclea undulata</i>	M
<i>Brachylaena ilicifolia</i>		<i>Euphorbia tirucalli</i>	M
<i>Carissa bispinosa</i>		<i>Grewia bicolor</i>	
<i>Celtis africana</i>		<i>Grewia flava</i>	
<i>Combretum apiculatum</i>	d	<i>Gymnosporia senegalensis</i>	M
<i>Combretum hereroense</i>		<i>Hippobromus pauciflorus</i>	

Kirkia wilmsii
Ozoroa albicans
Pouzolzia mixta
Rhoicissus tridentata

Searsia leptodictya
Senna petersiana
Tecoma capensis
Vitex obovata subsp wilmsii

Grasses

Aristida canescens
Aristida congesta
Aristida transvaalensis
Bewisia biflora
Bothriochloa insculpta
Brachiaria serrata
Digitaria eriantha
Diheteropogon amplexans
Elionurus muticus
Enneapogon scoparius

Eragrostis lehmanniana
Eragrostis superba
Heteropogon contortus
Loudetia simplex
Melinis nerviglume
Melinis repens s. repens
Panicum deustum
Panicum maximum
Themeda triandra

Forbs

Abutilon angulatum
Asparagus intricatus
Asparagus suaveolens
Barleria saxatilis
Blepharis integrifolia
Cheilanthes dolomitica
Clerodendrum ternatum
Commelina africana M
Corchorus asplenifolius
Euphorbia schinzii
Evolvulus alsinoides
Geigeria burkei
Hibiscus aethiopicus
Hypoestes forskalii

Kohautia cynanchica
Kyphocarpa angustifolia
Ocimum americanum
Phyllanthus glaucophyllus
Polygala hottentotta
Rhynchosia nitens
Sansevieria hyacinthoides
Stylochiton natalensis
Tephrosia purpurea
Tetradenia brevispicata,
Tragia dioica
Waltheria indica W
Xerophyta retinervis

Species of Conservation Concern

Although Matthews (1991), Van Wyk & Smith (2001) and Mucina & Rutherford (2006) mention that several endemic plant species occur on this dolomite area, the area crossed by the lines is so small that none of these species were recorded.

Conclusion

As far as vegetation is concerned, the development of the power lines can be supported in this area.



Figure 8: Pong Dolomite Mountain Bushveld

6. Northern Escarpment Quartzite Sourveld

A very small part of this veld type is close to the proposed route of Route 1, though it is so small that this vegetation is not discussed further.

7. Northern Mistbelt Forest

The proposed lines cannot go through indigenous forest. The proposed Route 4 seems to cross a patch of forest, but this will not be allowed by the authorities. This is one of the reasons why Route 4 was eliminated as an option. This vegetation is not discussed further.



8. Tzaneen Sour Bushveld

This vegetation stretches in a narrow band on the plains and all along the lower footslopes and hills of the escarpment. The vegetation is tall open bushveld with a tall grass layer. The geological substrate is gneiss and granite and the soils are shallow sandy and rocky lithosols. This is very dense, often tall bushveld, merging into forest-like vegetation. Only the route of Option 4 transects this vegetation, albeit a very narrow band. This vegetation contains many large trees and also has a very high species richness. Large *Ficus* trees (various species) together with several other large trees are prominent. Several threatened species occur in this vegetation.

The following species are present:

Trees and shrubs

<i>Acacia davyi</i>		<i>Ficus sansibarica</i>	
<i>Acacia polyacantha</i>		<i>Heteropyxis natalensis</i>	
<i>Acacia sieberiana</i>		<i>Parinari curatellifolia</i>	
<i>Albizia versicolor</i>		<i>Peltophorum africanum</i>	
<i>Antidesma venosum</i>		<i>Piliostigma thonningii</i>	
<i>Bauhinia galpinii</i>		<i>Pterocarpus angolensis</i>	P
<i>Catha edulis</i>	M	<i>Pterocarpus rotundifolius</i>	
<i>Faurea rochetiana</i>		<i>Sclerocarya birrea</i>	P
<i>Faurea saligna</i>		<i>Searsia pentheri</i>	
<i>Ficus burkei</i>		<i>Terminalia sericea</i>	
<i>Ficus petersii</i>		<i>Trichilia emetica</i>	

Grasses

<i>Alloteropsis semialata</i>		<i>Heteropogon contortus</i>	
<i>Andropogon schirensis</i>		<i>Hyparrhenia cymbaria</i>	
<i>Aristida congesta</i>		<i>Hyperthelia dissoluta</i>	
<i>Bothriochloa bladhii</i>		<i>Setaria nigrirostris</i>	
<i>Cymbopogon caesius</i>		<i>Setaria sphacelata</i>	
<i>Cynodon dactylon</i>		<i>Themeda triandra</i>	
<i>Diheteropogon amplexans</i>			

Forbs

Some forb species occur scattered in the grassy layer, and these are not abundant:

Merensky Foskor February 2012



Agathisanthemum bojeri

Barleria elegans

Dicliptera clinopodia

Polygala producta

Species of Conservation Concern

Species	Status
<i>Aloe hardyi</i> H.F.Glen	Rare
<i>Combretum petrophilum</i> Retief	Rare
<i>Dracaena transvaalensis</i> Baker	Rare
<i>Encephalartos brevifoliolatus</i> Vorster	EW
<i>Encephalartos cupidus</i> R.A.Dyer	CR
<i>Encephalartos paucidentatus</i> Stapf & Burtt Davy	VU
<i>Gladiolus macneilii</i> Oberm.	CR
<i>Helichrysum junodii</i> Moeser	Rare
<i>Thesium davidsonae</i> Brenan	VU

Several species of conservation concern are present in this dense vegetation. Although none of these were recorded during the field survey, possibly due to the very narrow band of this vegetation, but also due to the inaccessibility, it is certain that some of them will occur on the route. This is one of the reasons why Route 4 is eliminated as an option.

Conclusion

Due to the dense vegetation, species richness and possible presence of threatened species, it is suggested to avoid this route.





Figure 9: Dense Tzaneen Sour Bushveld



9. Granite Bushveld

The granite bushveld on the granite Lowveld plains cover the largest part of all the transects in the study site. This large granite plain is covered with the “typical” Lowveld bushveld, well know, for example, from large parts of the Kruger National Park.

Prominent species are *Combretum apiculatum* on the sandy or gravelly upland sites, while *Acacia nigrescens* and other *Acacia* species are prominent on the more clayey bottomland sites. Many species could be recorded, as this area is often close to roads and accessible.

The following plant species were recorded in this plant community:

Trees and shrubs

<i>Acacia exuvialis</i>		<i>Dodonaea angustifolia</i>	M
<i>Acacia gerrardii</i>		<i>Dombeya rotundifolia</i>	M
<i>Acacia nigrescens</i>	d	<i>Ehretia amoena</i>	
<i>Acacia nilotica</i>		<i>Ehretia rigida</i>	m
<i>Acacia sieberiana</i>	M	<i>Erythrina lysistemon</i>	
<i>Acacia tortilis</i>	d	<i>Euclea divinorum</i>	M
<i>Agave sisalana</i>	A	<i>Euclea natalensis</i>	
<i>Albizia harveyi</i>	d	<i>Ficus stuhlmanni</i>	
<i>Bolusanthus speciosa</i>		<i>Flueggea virosa</i>	
<i>Capparis tomentosa</i>		<i>Gardenia volkensii</i>	
<i>Cassia abbreviata</i>		<i>Grewia bicolor</i>	
<i>Combretum apiculatum</i>	D	<i>Grewia flava</i>	
<i>Combretum collinum</i>		<i>Grewia flavescens</i>	
<i>Combretum hereroense</i>		<i>Grewia monticola</i>	
<i>Combretum imberbe</i>	P	<i>Gymnosporia senegalensis</i>	M
<i>Combretum zeyheri</i>	M	<i>Lannea discolor</i>	
<i>Commiphora africana</i>		<i>Lannea schweinfurthii</i>	
<i>Cordia ovalis</i>		<i>Lantana camara</i>	A
<i>Daibergia melanoxydon</i>		<i>Ozoroa engleri</i>	
<i>Dichrostachys cinerea</i>	dM	<i>Pappea capensis</i>	

<i>Pavetta schumanniana</i>		<i>Searsia guenzii</i>	
<i>Pavetta schumanniana</i>	M	<i>Searsia leptodictya</i>	
<i>Peltophorum africanum</i>	M	<i>Senna didymobotrya</i>	A
<i>Philenoptera violacea</i>	PM	<i>Senna petersiana</i>	DM
<i>Pterocarpus angolensis</i>	PM	<i>Spirostachys africana</i>	p
<i>Pterocarpus rotundifolius</i>		<i>Terminalia prunoides</i>	
<i>Schotia brachypetala</i>		<i>Terminalia sericea</i>	DM
<i>Schotia capitata</i>		<i>Xeromphis obovata</i>	
<i>Sclerocarya birrea</i>	PM	<i>Ziziphus mucronata</i>	M

Grasses

<i>Aristida adscensionis</i>		<i>Hyperthelia dissoluta</i>	
<i>Aristida congesta</i>		<i>Melinis repens</i>	
<i>Aristida diffusa</i>		<i>Panicum maximum</i>	
<i>Bothriochloa insculpta</i>		<i>Perotis patens</i>	
<i>Brachiaria nigropedata</i>		<i>Pogonarthria squarrosa</i>	
<i>Cenchrus ciliaris</i>		<i>Sporobolus africanus</i>	
<i>Digitaria eriantha</i>		<i>Sporobolus ioclados</i>	
<i>Enneapogon cenchroides</i>		<i>Tricholaena monachne</i>	
<i>Enneapogon scoparius</i>		<i>Trichoneura grandiglumis</i>	
<i>Eragrostis rigidior</i>		<i>Urochloa mosambicensis</i>	d
<i>Heteropogon contortus</i>			

Forbs

<i>Achyranthes aspera</i>		<i>Geigeria burkei</i>	
<i>Acrotome inflata</i>		<i>Gomphrena celosioides</i>	W
<i>Agathisanthemum bojeri</i>		<i>Helichrysum</i> sp	
<i>Aspilia mossambicensis</i>		<i>Heliotropium steudneri</i>	
<i>Becium filamentosum</i>		<i>Heliotropium strigosum</i>	
<i>Bidens pilosa</i>	W	<i>Hermannia</i> sp	
<i>Bulbostylis hispidula</i>		<i>Hermannia tomentosa</i>	
<i>Chamaecrista absus</i>		<i>Hermbstaedtia odorata</i>	
<i>Commelina benghalensis</i>		<i>Hibiscus cannabinus</i>	
<i>Commelina erecta</i>		<i>Hibiscus</i> sp	
<i>Cyperus esculentus</i>	W	<i>Indigofera filipes</i>	
<i>Dicerocaryum zanguebarium</i>	M	<i>Kohautia virgata</i>	
<i>Evolvulus alsinoides</i>	M	<i>Kyphocarpa angustifolia</i>	

<i>Leucas glabrata</i>		<i>Solanum panduriforme</i>	W
<i>Ocimum americanum</i>		<i>Tagetes minuta</i>	W
<i>Phyllanthus maderaspatensis</i>		<i>Tephrosia grandiflora</i>	
<i>Rhynchosia totta</i>		<i>Waltheria indica</i>	W
<i>Richardia braziliensis</i>	W	<i>Zornia milneana</i>	
<i>Schkruhria pinnata</i>	WM		

Species of Conservation Concern

Asphodelaceae	<i>Aloe hardyi</i> H.F.Glen	Rare
Asphodelaceae	<i>Aloe thompsoniae</i> Groenew.	Rare
Apocynaceae	<i>Brachystelma parvulum</i> R.A.Dyer	VU
Combretaceae	<i>Combretum petrophilum</i> Retief	Rare
Celastraceae	<i>Elaeodendron transvaalense</i> (Burt Davy) R.H.Archer	NT
Zamiaceae	<i>Encephalartos dyerianus</i> Lavranos & D.L.Goode	CR
Zamiaceae	<i>Encephalartos lebomboensis</i> I.Verd.	EN
Proteaceae	<i>Protea laetans</i> L.E.Davidson	VU
Rosaceae	<i>Prunus africana</i> (Hook.f.) Kalkman	VU
Anacardiaceae	<i>Searsia batophylla</i> (Codd) Moffett	VU

None of these species were recorded during the field survey. The flat granite plains are not suitable habitat for these species, which are all niche specialists, or occur on the mountain areas to the west of the granite plains.

Conclusion

Several game farms and cattle farms are found in this area.

Special care will be needed in the crossing of the spruit systems within this area.

The construction of the line can be supported.



Figure 10: Typical Granite Lowveld vegetation



10. Lowveld Rugged Mopaneveld

Only Routes 1 and 2 will transect this vegetation type in the north-eastern extreme of the study area. The Foskop substation is located on the northern boundary of this vegetation type. This is the rugged hilly area of the Olifants River valley, south of Phalaborwa. The landscape is irregular plains and rocky hills, with moderate to steep slopes. *Colophospermum mopane* is often restricted to valleys, while the hills are dominated by *Acacia nigrescens* with *Combretum apiculatum* also present.

The following plant species were recorded in this plant community:

Trees and shrubs

<i>Acacia exuvialis</i>		<i>Gardenia volkensii</i>	
<i>Acacia nigrescens</i>		<i>Grewia bicolor</i>	
<i>Acacia nilotica</i>		<i>Grewia flavescens</i>	
<i>Berchemia discolor</i>		<i>Grewia hexamita</i>	
<i>Boscia albitrunca</i>	P	<i>Grewia monticola</i>	
<i>Colophospermum mopane</i>	D	<i>Grewia villosa</i>	
<i>Combretum apiculatum</i>		<i>Gymnosporia senegalensis</i>	M
<i>Combretum hereroense</i>		<i>Hexalobus monopetalus</i>	
<i>Combretum imberbe</i>	P	<i>Kirkia wilmsii</i>	
<i>Combretum zeyheri</i>	M	<i>Lannea discolor</i>	
<i>Commiphora africana</i>		<i>Manilkara mochisia</i>	
<i>Commiphora mollis</i>		<i>Ozoroa engleri</i>	
<i>Dalbergia melanoxylon</i>		<i>Pappea capensis</i>	
<i>Dichrostachys cinerea</i>	M	<i>Pavetta schumanniana</i>	M
<i>Dodonaea angustifolia</i>	M	<i>Peltophorum africanum</i>	M
<i>Dombeya rotundifolia</i>	M	<i>Philenoptera violacea</i>	PM
<i>Ehretia amoena</i>		<i>Pterocarpus rotundifolius</i>	
<i>Ehretia rigida</i>	M	<i>Rhigozum zambesiaceum</i>	
<i>Erythrina lysistemon</i>		<i>Sclerocarya birrea</i>	PM
<i>Euclea natalensis</i>		<i>Terminalia prunoides</i>	
<i>Ficus abutilifolia</i>		<i>Terminalia sericea</i>	DM
<i>Flueggea virosa</i>			

Grasses

<i>Aristida adscensionis</i>	<i>Aristida congesta</i>
------------------------------	--------------------------



Bothriochloa radicans
Cenchrus ciliaris
Digitaria eriantha
Enneapogon cenchroides
Enneapogon scoparius
Eragrostis rigidior
Fingerhuthia africana
Heteropogon contortus

Melinis repens
Panicum maximum
Pogonarthria squarrosa
Sporobolus panicoides
Tricholaena monachne
Trichoneura grandiglumis
Urochloa mosambicensis d

Forbs

Achyranthes aspera
Agathisanthemum bojeri
Aspilia mossambicensis
Chamaecrista mimosoides
Commelina benghalensis
Commelina erecta
Crabbea velutina
Evolvulus alsinoides
Geigeria burkei
Gomphrena celosioides
Gossypium africanum
Heliotropium steudneri
Heliotropium strigosum
Hemizygia elliotii

Hermannia tomentosa
Hermestaedtia odorata
Hibiscus sidiformis
Kohautia virgata
Kyphocarpa angustifolia
Leucas glabrata
Melhania forbesii
Melhania rehmannii
Ocimum americanum
Phyllanthus asperulatus
Solanum panduriforme
Waltheria indica
Xerophyta retinervis
Zornia milneana

Species of Conservation Concern

Species	Status
<i>Ansellia africana</i> Lindl.	Declining

This epiphytic orchid is often found growing on trees within this plant community.

Conclusion

Game farming is a special feature in this area.

Special care will be needed in the crossing of spruit systems.

The construction of the line can be supported.



Figure 11: Lowveld Rugged Mopaneveld in the background

11. Phalaborwa-Timbavati Mopaneveld

Only Routes 1 (small part) and 4 will transect this vegetation type in the north-eastern extreme of the study area. The Foskor substation is located on the southern boundary of this vegetation type. This is the flat plains west of Phalaborwa. The landscape is an undulating plain with *Colophospermum mopane* and *Acacia nigrescens* in the lower lying areas, while *Combretum apiculatum* and *Terminalia sericea* becomes more prominent on upland sites.

The following plant species were recorded in this plant community:

Trees and shrubs

Acacia exuvialis

Acacia nigrescens

<i>Acacia tortilis</i>		<i>Gardenia volkensii</i>	
<i>Albizia harveyi</i>		<i>Grewia bicolor</i>	
<i>Boscia albitrunca</i>	P	<i>Grewia flavescens</i>	
<i>Cassia abbreviata</i>		<i>Grewia hexamita</i>	
<i>Cissus cornifolia</i>		<i>Grewia monticola</i>	
<i>Colophospermum mopane</i>	D	<i>Grewia villosa</i>	
<i>Combretum apiculatum</i>	d	<i>Gymnosporia senegalensis</i>	M
<i>Combretum hereroense</i>		<i>Lannea discolor</i>	
<i>Combretum imberbe</i>	P	<i>Maerua parvifolia</i>	
<i>Combretum zeyheri</i>	M	<i>Ozoroa engleri</i>	
<i>Commiphora africana</i>		<i>Pappea capensis</i>	
<i>Commiphora mollis</i>		<i>Pavetta schumanniana</i>	M
<i>Dalbergia melanoxylon</i>		<i>Peltoporum africanum</i>	M
<i>Dichrostachys cinerea</i>	M	<i>Philenoptera violacea</i>	PM
<i>Ehretia amoena</i>		<i>Sclerocarya birrea</i>	PM
<i>Ehretia rigida</i>		<i>Strychnos madagascariensis</i>	
<i>Euclea divinorum</i>		<i>Terminalia prunoides</i>	
<i>Flueggea virosa</i>		<i>Terminalia sericea</i>	dM

Grasses

<i>Aristida adscensionis</i>		<i>Melinis repens</i>	
<i>Aristida congesta</i>		<i>Panicum maximum</i>	
<i>Bothriochloa radicans</i>		<i>Perotis patens</i>	
<i>Brachiaria nigropedata</i>		<i>Pogonarthria squarrosa</i>	
<i>Cenchrus ciliaris</i>		<i>Schmidtia pappophoroides</i>	
<i>Digitaria eriantha</i>		<i>Themeda triandra</i>	
<i>Andropogon gayanus</i>		<i>Tricholaena monachne</i>	
<i>Fingerhuthia africana</i>		<i>Trichoneura grandiglumis</i>	
<i>Enneapogon scoparius</i>		<i>Urochloa mosambicensis</i>	d
<i>Eragrostis rigidior</i>			
<i>Heteropogon contortus</i>			

Forbs

<i>Achyranthes aspera</i>		<i>Chamaecrista mimosoides</i>	
<i>Acrotome inflata</i>		<i>Clerodendrum ternatum</i>	
<i>Agathisanthemum bojeri</i>		<i>Commelina benghalensis</i>	
<i>Aspilia mossambicensis</i>		<i>Commelina erecta</i>	

Evolvulus alsinoides
Gomphrena celosioides
Heliotropium steudneri
Heliotropium strigosum
Hemizygia elliotii
Hermannia glanduligera
Hermannia tomentosa
Hermbstaedtia odorata
Ipomoea magnusiana

Kohautia virgata
Kyphocarpa angustifolia
Leucas glabrata
Melhania forbesii
Ocimum americanum
Solanum panduriforme
Tephrosia polystachya
Waltheria indica
Zornia milneana

Species of Conservation Concern

Species	Status
<i>Aloe thompsoniae</i> Groenew.	Rare
<i>Encephalartos dyerianus</i> Lavranos & D.L.Goode	CR
<i>Encephalartos lebomboensis</i> I.Verd.	EN

None of these species were found during the field survey.

Conclusion

Game farming is a special feature in this area.

Special care will be needed in the crossing of spruit systems.

The construction of the line can be supported.



Figure 12: Game farming in the Lowveld areas.

12 River and Spruit Systems

Several River or spruit crossings occur along the transect routes. At the smaller spruits no riparian zone is present, the vegetation is continuous with the adjacent bushveld vegetation. However, the larger rivers have a distinct riparian zone, with large trees for example *Faidherbia albida*, *Acacia galpinii*, *Acacia robusta* and *Philenoptera violacea*.

The following species were recorded at a larger river:

Woody species

<i>Acacia galpinii</i>		<i>Diospyros mespiliformis</i>	
<i>Acacia robusta</i>		<i>Faidherbia albida</i>	D
<i>Acacia tortilis</i>		<i>Ficus sycomorus</i>	
<i>Combretum apiculatum</i>		<i>Grewia bicolor</i>	
<i>Combretum imberbe</i>	P	<i>Grewia monticola</i>	
<i>Combretum microphyllum</i>		<i>Peltophorum africanum</i>	
<i>Dichrostachys cinerea</i>		<i>Philenoptera violacea</i>	P

Phoenix reclinata
Sclerocarya birrea MP

Terminalia sericea
Ziziphus mucronata M

Grasses

Aristida congesta
Cynodon dactylon
Dactyloctenium aegyptium
Digitaria eriantha
Eragrostis heteromera
Eragrostis rigidior d

Eragrostis superba
Heteropogon contortus D
Melinis repens
Pogonarthria squarrosa d
Tricholaena monachne
Trichoneura grandiglumis

Forbs

Agathisanthemum bojeri
Ageratum conyzoides W
Cyperus sexangularis
Lippia rehmannii
Schkruhria pinnata W

Melanthera scandens
Solanum incanum W
Solanum panduriforme W
Xanthium strumarium W

River and Spruit crossings			
Status	River and spruit linear wetlands		
Soil	Sandy or clayey	Rockiness	0%
Conservation priority:	High	Sensitivity:	High
Agricultural potential:	Low	Need for rehabilitation	Low
Dominant spp.	<i>Combretum imberbe</i> , <i>Faidherbia albida</i> , <i>Acacia galpinii</i> , <i>Diospyros mespiliformis</i> , <i>Ficus sycomorus</i> , <i>Acacia robusta</i> , <i>Philenoptera violacea</i> , <i>Phoenix reclinata</i>		

Conclusion

No river or spruit is very wide, so the lines can easily cross these river or spruit systems. Care should however be taken to place pylons adequately away from river or spruit banks, avoiding any damage to the banks or water courses. Erosion should be avoided at all times.

8 DISCUSSION

The vegetation of all four alternatives was investigated. From the desktop study, confirmed by the field survey, option 3, which runs from Burgersfort to Ohrigstad along the R555, was eliminated. This is because the route along the R555 runs for most of the way in a narrow valley, with the Mabitsana River and the tarred R555 in this valley. The line will have to run for most of the way on the sensitive mountain foot slopes and cross the river and road several times. Furthermore, many irrigated agricultural enterprises occur in the Ohrigstad area, stretching all the way to Marapeng. This mosaic of narrow river valley, river, mountain slopes and agriculture where-ever the valley is a bit broader, causes the route to be unsuitable. From an ecological perspective both the riverine vegetation and the vegetation of the mountain slopes have a high ecological sensitivity. Therefore this entire valley forms an ecologically sensitive ecosystem. This is also a much longer route.

Furthermore, from the desktop study, confirmed by the field survey, option 4, was eliminated. The line of this option runs through nine vegetation types, and over very high and steep mountains of Sekhukhune Mountain Bushveld and Ohrigstad Mountain Bushveld with two endangered ecosystems (Sekhukhune Mountainlands and Sekhukhune Norite Bushveld, SANBI & DEAT 2009), Pong Dolomite Mountain Bushveld with endangered Malmani Karstland (SANBI & DEAT 2009), the vulnerable Northern Escarpment Quartzite Sourveld (Mucina & Rutherford (2006), Northern Mistbelt Forest area and the vulnerable Tzaneen Sour Bushveld (SANBI & DEAT 2009). Especially the Great Escarpment area consists of very rugged and high mountains, resulting in a very difficult route with several threatened ecosystems.

Alternative Routes 1 and 2 were further investigated in more detail by field surveys. The vegetation along these routes is described in detail, including the protected and red data species. Medicinal plants and aliens and weeds are indicated.

The most difficult part of the route is from the Merensky substation through Orichstad Mountain Bushveld which is an extremely mountainous area with sensitive vegetation. This part of the line transects quite sensitive vegetation and it is suggested that a walkthrough in this area is essential.

The most serious limitation on the Lowveld plains where the line transects the Granite Lowveld vegetation type, is the abundance of the protected tree *Sclerocarya birrea*. It is certain that several of these trees will be in the way of the transect.

Locally are also the river and spruit crossings. No river or spruit is very wide, so the lines can easily cross these rivers or spruits systems. Care should be taken to place pylons adequately away from river or spruit banks, avoiding any damage to the banks or water courses. Erosion should be avoided at all times.

Another factor in this area is that large properties are game farms and lodges. These areas are effectively conserved by the owners, and it is realised that the public participation is an important issue. After finalisation of the exact transect, a walkthrough will have to confirm any issues regarding vegetation.

From an ecological perspective, Alternative 1 is the preferred route.

11. REFERENCES

Acocks, J.P.H. 1988. Veld types of South Africa, 3rd ed. *Memoirs of the Botanical Survey of South Africa*. 57: 1–146.

Bredenkamp, G.J. 1982. 'n Plantekologiese studie van die Manyeleti Wildtuin. DSc thesis, University of Pretoria, Pretoria.

Bredenkamp, G.J. & Brown, L.R. 2001. Vegetation – A reliable ecological basis for environmental planning. *Urban Greenfile* Nov-Dec 2001: 38-39.

Department of Environmental Affairs and Tourism. 2007. National Environmental Management: Biodiversity Act, 2004 (Act 10 of 2004): Publication of Lists of Critically Endangered, Endangered, Vulnerable and Protected Species. Government Notices.

Gertenbach, W.P.D. 1983a. Ekologiese studie van die suidelikste Mopaneveld in the Kruger Nasionale Wildtuin. DSc thesis, University of Pretoria, Pretoria.

Gertenbach, W.P.D. 1983b. Landscapes in the Kruger National park, *Koedoe* 26: 9-121

Henderson, L. 2001. Alien weeds and invasive plants. ATC, Pretoria.

Low, A.B. & Rebelo, A.G. (eds) 1996 *Vegetation of South Africa, Lesotho and Swaziland*. Dept Environmental Affairs & Tourism, Pretoria.

Matthews, W.S. 1991. Phytosociology of North-eastern Mountain Sourveld. MSc thesis, University of Pretoria, Pretoria.

Mucina, L. & Rutherford, M.C. (Eds.) 2006. The vegetation of South Africa, Lesotho and Swaziland. *Strelitzia* 19. South African National Biodiversity Institute, Pretoria.

Mucina, L., Bredenkamp, G.J., Hoare, D.B. & McDonald, D.J. 2000. A National vegetation database for South Africa. *South Africa Journal of Science* 96:497-498.

Mueller-Dombois, D. & Ellenberg, H. 1974. Aims and methods of vegetation ecology. Wiley, New York.

Raimondo, D., Von Staden, L., Foden, W., Victor, J.E., Helme, N.A., Turner, R.C. Kamundi, D.A. & Manyama, P.A. (Eds.). 2009. Red list of South African plants 2009. *Strelitzia* 25:1-668.

Siebert, S.J. 1998. Ultramafic substrates and floristic patterns in Sekhukhuneland, South Africa. MSc thesis, University of Pretoria, Pretoria

Siebert, S.J. 2001 Vegetation on the ultramafic soils of the Sekhukhuneland Centre of Endemism. PhD thesis, University of Pretoria, Pretoria

Siebert, S.J., Van Wyk, A.E. & Bredenkamp, G.J. 2001. Endemism in the flora of ultramafic areas of Sekhukhuneland, South Africa *South African Journal of Science* 97:529-532.

Siebert, S.J., Victor, J., Van Wyk, A.E. & Bredenkamp, G.J. 2002a. An assessment of threatened plants and conservation in Sekhukhuneland. *PlantLife* 26:7-18.

Siebert, S.J. Van Wyk, A.E. & Bredenkamp, G.J. 2002b. The physical environment and major vegetation types of Sekhukhuneland, South Africa. *South African Journal of Botany* 68: 127-142.

Siebert S.J., Van Wyk A.E., Bredenkamp G.J. & Du Plessis F. 2002c. Grasslands and wetlands of the Sekhukhuneland Centre of Plant Endemism. *Bothalia* 32(2): 211–231.

Siebert S.J., Van Wyk A.E., & Bredenkamp G.J. 2002d. Vegetation ecology of Sekhukhuneland, South Africa: *Combretum hereroense* – *Grewia vernicosa* Open Mountain Bushveld. *South African Journal of Botany* 68(4): 497-517.

Siebert S.J., Van Wyk A.E., & Bredenkamp G.J. 2002e. Vegetation ecology of Sekhukhuneland, South Africa: *Kirkia wilmsii* – *Terminalia prunioides* Closed Mountain Bushveld. *South African Journal of Botany* 68(4): 474-496.



Siebert, S.J., Van Wyk, A.E., Bredenkamp, G.J. & Siebert, F. 2003. Vegetation of the rock habitats of the Sekhukhuneland Centre of Plant Endemism, South Africa. *Bothalia* 33(2): 207-228.



The Conservation of Agricultural Resources Act, 1983 (Act No. 43 of 1983)

The Environmental Conservation Act, 1989 (Act No. 73 of 1989)

The National Environment Management Act, 1998 (Act No. 107 of 1998)

The National Environmental Management Biodiversity Act, 2004. (Act 10 Of 2004).
Government Gazette RSA Vol. 467, 26436, Cape Town, June 2004.

The National Environmental Management: Biodiversity Act, 2004. (Act 10 Of 2004),
Draft list of threatened ecosystems. Government Gazette RSA Vol. 1477, November
2009.

The National Forests Act, 1998 (Act 84 of 1998)

Van Wyk, A.E. & Smith, G.F. 2001. Regions of floristic endemism in southern Africa.
Umdaus Press, Hatfield, Pretoria.

Van Wyk, B.E., Van Oudtshoorn, B. & Gericke, N. 1997. Medicinal plants of South
Africa. Briza, Pretoria.

Westhoff, V. & Van der Maarel, E. 1978. The Braun-Blanquet approach. In: Whittaker,
R.H. (ed.) Classification of plant communities. W. Junk, The Hague.

ABRIDGED CURRICULUM VITAE: GEORGE JOHANNES BREDEKAMP

Born: 10 February 1946 in Johannesburg, South Africa.

Citizenship: South African

Marital status: Married, 1 son, 2 daughters

Present work address

Department of Botany, University of Pretoria, Pretoria, 0002, South Africa

Tel:(27)(12)420-3121 Fax: (27)(12)362 5099

E-Mail: gbredenk@postino.up.ac.za

or

EcoAgent CC

PO Box 25533, Monument Park, 0105, South Africa

Tel: (27)(12) 346 3180

Fax: (27)(12) 460 2525

Cell 082 5767046

E-Mail: ecoagent@mweb.co.za

Qualifications:

1963 Matriculation Certificate, Kemptonpark High School

1967 B.Sc. University of Pretoria, Botany and Zoology as majors,

1968 B.Sc. Hons. (cum laude) University of Pretoria, Botany.

1969 T.H.E.D. (cum laude) Pretoria Teachers Training College.

1975 M.Sc. University of Pretoria, Plant Ecology .

1982 D.Sc. (Ph.D.) University of Pretoria, Plant Ecology.

Theses: (M.Sc. and D.Sc.) on plant community ecology and wildlife management in nature reserves in South African grassland and savanna.

Professional titles:

- MSAIE South African Institute of Ecologists and Environmental Scientists
 - 1989-1990 Council member
- MGSSA Grassland Society of Southern Africa
 - 1986 Elected as Sub-editor for the Journal
 - 1986-1989 Serve on the Editorial Board of the Journal
 - - 1990 Organising Committee: International Conference: Meeting Rangeland challenges in Southern Africa
 - 1993 Elected as professional member
- PrSciNat. South African Council for Natural Scientific Professions **Registration Number 400086/83**
 - 1993-1997 **Chairman** of the Professional Advisory Committee: Botanical Sciences
 - 1993-1997: **Council** Member
 - 1992-1994: Publicity Committee
 - 1994-1997: Professional Registration Committee

Professional career:

- Teacher in Biology 1970-1973 in Transvaal Schools
- Lecturer and senior lecturer in Botany 1974-1983 at University of the North
- Associate professor in Plant Ecology 1984-1988 at Potchefstroom University for CHE

• Professor in Plant Ecology 1988-2008 at University of Pretoria.

• 2009 – current Professor Extra-ordinary in the Dept of Plant Science, University of Pretoria

- Founder and owner of the Professional Ecological Consultancy firms Ecotrust Environmental Services CC and Eco-Agent CC, 1988-present.

Academic career:

- Students:
 - Completed post graduate students: M.Sc. 53; Ph.D. 14.
 - Presently enrolled post-graduate students: M.Sc. 4; Ph.D. 2.
- Author of:
 - 175 scientific papers in refereed journals
 - >150 papers at national and international congresses
 - >250 scientific (unpublished) reports on environment and natural resources
 - 17 popular scientific papers.
 - 39 contributions in books
- Editorial Committee of
 - South African Journal of Botany,
 - Journal Grassland Society of Southern Africa,
 - Bulletin of the South African Institute of Ecologists.
 - Journal of Applied Vegetation Science.(Sweden)
 - Phytocoenologia (Germany)
 -
- FRD evaluation category: C2 (=leader in South Africa in the field of Vegetation Science/Plant Ecology)

Membership:

- International Association of Vegetation Science.
- British Ecological Society
- International Society for Ecology (Intecol)
- Association for the Taxonomic study of the Flora of Tropical Africa (AETFAT).
- South African Association of Botanists (SAAB)
 - 1988-1993 Elected to the **Council** of SAAB.
 - 1989-1990 Elected as **Chairman** of the Northern Transvaal Branch
 - 1990 Elected to the Executive Council as **Vice-President**
 - 1990- Sub-editor Editorial Board of the Journal
 - 1991-1992 Elected as **President** (2-year period)
 - 1993 **Vice-President** and Outgoing President
- Wildlife Management Society of Southern Africa
- Suid-Afrikaanse Akademie vir Wetenskap en Kuns
(=South African Academy for Science and Art).
- Wildlife Society of Southern Africa
 - 1975 - 1988: Member
 - 1975 - 1983: Committee member, Pietersburg Centre
 - 1981 - 1982: **Chairman**, Pietersburg Centre
- Dendrological Society of Southern Africa
 - 1984 - present: Member
 - 1984 - 1988: Committee member, Western Transvaal Branch
 - 1986 - 1988: **Chairman**, Western Transvaal Branch
 - 1987 - 1989: Member, Central Committee (National level)
 - 1990 - 2000: Examination Committee
- Succulent Society of South Africa
 - 1987 - 2000
- Botanical Society of South Africa
 - 2000 - present: Member
 - 2001- 2008: Chairman, Pretoria Branch



2002 – 2006: Chairman, Northern Region Conservation Committee
2002- 2007: Member of Council

Special committees:

- Member of 10 special committees re ecology, botany, rangeland science in South Africa.
- Member of the International Code for Syntaxonomical Nomenclature 1993-present.

Merit awards and research grants:

1968 Post graduate merit bursary, CSIR, Pretoria.
1977-1979 Research Grant, Committee re Research Development, Dept. of Co-operation and Development, Pretoria.
1984-1989 Research Grant, Foundation for Research Development, CSIR, Pretoria.
1986-1987 Research Grant, Dept. of Agriculture and Water Supply, Potchefstroom.
1990-1997 Research Grant, Dept. of Environmental Affairs & Tourism, Pretoria.
1991-present Research Grant, National Research Foundation , Pretoria.
1991-1993 Research Grant, Water Research Commission.
1999-2003 Research Grant, Water Research Commission.
2006 South African Association of Botanists Silver Medal for outstanding contributions to South African Botany

Abroad:

1986 Travel Grant, Potchefstroom University for Christian Higher Education, Potchefstroom
Visits to Israel, Italy, Germany, United Kingdom, Portugal.
1987 Travel Grant, Potchefstroom University for Christian Higher Education, Potchefstroom.
Visits to Germany, Switzerland, Austria, The Netherlands, United Kingdom.
1990 Travel Grant, FRD.
Visit to Japan, Taiwan, Hong-Kong.
1991 Travel Grant, FRD.
Visits to Italy, Germany. Switzerland, Austria, France, The Netherlands, United Kingdom.
1993 Travel Grant, University of Pretoria.
Visits to the USA, Costa Rica, Czech Republic, Austria.
1994 Travel Grant FRD.
Visits to Switzerland, The Netherlands, Germany, Czech Republic.
1995 Travel Grant FRD, University of Pretoria
Visits to the USA
1996 Travel Grant, University of Pretoria
Visit to the UK.
1997 Travel Grant University of Pretoria, Visit Czech Republic, Bulgaria
1998 Travel Grant, University of Pretoria, Visit Czech Republic, Italy, Sweden
1999 Travel Grant, University of Pretoria, Visit Hungary, Spain, USA
2000 Travel Grant, University of Pretoria, Visit Poland, Italy, Greece.
2001 Travel Grant, NRF, Visit Brazil
2006 German Grant Invited lecture in Rinteln, Germany

Consultant

Founder and owner of Ecotrust Environmental Services CC and Eco-Agent CC
Since 1988 >250 reports as consultant on environmental matters, including:

- Game Farm and Nature Reserve planning,
- Environmental Impact Assessments,



- Environmental Management Programme Reports,
- Vegetation Surveys,
- Wildlife Management,
- Veld Condition and Grazing Capacity Assessments,
Red data analysis (plants and animals).



