

OFFICIAL ORGAN
OF THE



EUROPEAN GROUP OF LYMPHOLOGY
LATIN-MEDITERRANEAN CHAPTER OF ISL
SOCIETÀ ITALIANA DI LINFANGIOLOGIA
CZECH SOCIETY OF LYMPHOLOGY
ROMANIAN SOCIETY OF LYMPHOLOGY
GREEK SOCIETY OF LYMPHOLOGY

THE EUROPEAN JOURNAL OF lymphology

and related problems

VOLUME 30 • No. 78 • 2019

INDEXED IN EXCERPTA MEDICA

SUMMARY

Clinical Sciences

Original articles

- Podoconiosis in Ethiopia. A pilot study to improve the management of Lymphedema
Isabel Forner-Cordero, MD. PhD., Marta Lopez-Agustin, MD., Marta Gutierrez-Delgado, MD. p. 1
- Dermal-edema: the micro-edema
Marcello Izzo, Enrico Oliva p. 5
- Suggestions for Education - Training in Lymphology - Do we need it?
Evangelos P. Dimakakos MD, PhD, EDA/VM, MLD/CDT p. 10
- Measuring leg volume in Lymphoedema using optoelectronic volumetry and the sequential tape method
Nyree Dunn, Jane H. Davies, Michelle Fishbourne, Gina Dolan, and E. Mark Williams p. 14
- The patient with primary and secondary lower extremity lymphedema; A 1-year follow-up study
Pt. Cemile Uydur, Pt. Msc. Gulbala Nakip p. 19
- Values and limits of surgery in primary and secondary lymphedema
Michellini S., Fiorentino A., Fantegrossi M.R., Failla A., Moneta G., Cardone M., Gobbetti G. p. 22
- The effect of lower limb Lymphedema on postural stabilization
Alis Kostanoglu, Meltem Ramoglu, Hikmet Ucgun p. 27

THE EUROPEAN JOURNAL OF LYMPHOLOGY AND RELATED PROBLEMS (EJLRP)

The EJLRP - official organ of the European Group of Lymphology (ESL), Czech Society of Lymphology, Romanian Society of Lymphology, Greek Society of Lymphology, the Latin-Mediterranean Chapter of Lymphology (LMCL), the Società Italiana di Linfangiologia (SIL) covers all fields of Lymphology and aims to present a multidisciplinary approach to diseases of the lymphatic system, with information on the analysis, control and treatments of such diseases.

Topics

The topics include:

- anatomy and anatomopathology
- physiology and physiopathology
- pharmacology
- diagnostic methods (conventional radiology, nuclear medicine, ultrasonography, computed tomography, biopsy, nuclear magnetic resonance)
- therapy (surgery, medicine, radiotherapy, physical)
- oncology (primary lymphatic system diseases, lymphonodal metastatic process)
- immunology
- post-therapeutic complications
- upper and lower limb edemas

Manuscripts publications

Submitted manuscripts will be published in the form of Editorial, Review article, Original article, Teaching article, Special article, Work in progress, Case Report, Short Communications, Letter to the Editor (in English), Abstract (in English)

They will be subdivided in Clinical and Basic Sciences.

Send manuscripts to:
the Executive Editor

Dr. S. MICHELINI

Department of Vascular Rehabilitation
S. Giovanni Battista Hospital
Via L.E. Morselli, 13 - 00148 Rome, Italy
Tel. +39 06 655961 - Fax +39 06 65596235
e-mail: sandro.michelin@fastwebnet.it

The Editor-in-Chief

Prof. Dr. F. BOCCARDO

Department of Surgery, Lymphatic Surgery and Microsurgery
S. Martino Hospital, University of Genoa
Largo R. Benzi, 8 - 16132 Genoa, Italy
Fax 0039010532778 - e-mail: Francesco.boccardo@unige.it

Associate-Editors also can receive and promote articles and start the review process.

Publications languages

Official language of the Journal is English.

Publication rate

The EJLRP is published on a quarterly basis.

Subscription rates - All members of European Group of Lymphology or of National societies (with which the ESL has a cooperation agreement and whose fee includes a subscription to the EJLRP) receive the Journal free of charge.

Subscription rate for non-members is:

- for all issues, 30 € within European Countries, 50 € elsewhere,
- for single issue, 15 € within European Countries, 18 € elsewhere.

Annual subscription rate of ESL: 80 €

Please make cheque (in euro) to order of the ESL and to be sent to the Treasurer of the ESL: Mr J.P. BELGRADO, Treasurer of the ESL, Service de Kinésithérapie, Avenue Paul Héger, 28, OF 168, 1050, Brussels, Belgium.

E-mail: belgrado@ulb.ac.be or transfer the corresponding amount on the following Bank Account of the ESL n. 210-0557380-70 N° IBAN BE60 2100 5573 8070 BIC GEBABEBB (Générale de Banque), with mention of your name and of the year(s) subscription.

Change of address - Please notify the Secretary and the Treasurer of the ESL of any change of address and telephone number at least 30 days prior to the issue date by sending both the old and new address.

Data base & Treasurer of the ESL - J.P. BELGRADO: Université Libre de Bruxelles, CP 168, Av. F.D. Roosevelt, 50, 1050 Bruxelles, Belgium. Tel. +32 2 650.24.34 - Fax: +32 2 280.13.33 - Mobile +32 475 63.34.34

Business communications - Business communications concerning advertising, subscriptions, change of address, and permission requests should be sent to the Secretary, O. LEDUC, Service de Kinésithérapie, Avenue Paul Héger, 28 CP 168, 1050 Brussels, Belgium. Tel. (32) (2) 650.24.70 - Fax: (32) (2) 650.24.73.

Advertisements are subject to editorial approval and restricted to products or services pertinent to lymphology.

Advertising rates can be obtained from the Secretary and Treasurer.

Miscellaneous - The use of general descriptive names, trade names, trademarks, etc., in the publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this Journal is believed to be true and accurate at the date of its going to press, neither the authors, the Editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

The Editors do not accept any responsibility for opinions that may be made by the authors.

Areas of distribution - Austria, Belgium, Czech Republic, Denmark, Egypt, France, Germany, Greece, Holland, Hungary, Israel, Italy, Japan, Norway, Poland, Portugal, Rumania, Russia, Spain, Sweden, UK, USA.

E.S.L. EXECUTIVE COMMITTEE

Distinguished Honorary President
LEDUC ALBERT (Belgium)

President

BOCCARDO FRANCESCO (Italy)

Vice-Presidents

BAUMEISTER RUEDIGER (Germany)

FÖLDI ETHEL (Germany)

Secretary

LEDUC OLIVIER (Belgium)

Treasurer

BELGRADO JEAN PAUL (Belgium)

Scientific Committee President

PISSAS ALEXANDRE (France)

Scientific Committee Vice-President

DIMAKAKOS EVANGELOS (Greece)

FORNER CORDERO ISABEL (Spain)

Members

BOURGEOIS PIERRE (Belgium)

BRORSON HÅKAN (Sweden)

BRUN JEAN PATRICE (France)

CESTARI MARINA (Italy)

ELIŠKA OLDŘICH (Czech Rep.)

HAMADÉ AMER (France)

JOHANSSON KARIN (Sweden)

MICHELINI SANDRO (Italy)

MONETA GIOVANNI (Italy)

RIQUET MARC (France)

THIBAUT GILBERT (France)

WALD MARTIN (Czech Rep.)

EUROPEAN JOURNAL OF LYMPHOLOGY AND RELATED PROBLEMS

Editor-in-Chief: P. BOURGEOIS (Belgium)

Executive-Editor-in-Chief: S. MICHELINI (Italy)

Assistant Executive Editor: O. ELIŠKA (Czech R.), M. CESTARI (Italy)

Assistant-Editors: J.P. BELGRADO (Belgium), I. FORNER CORDERO (Spain), A. HAMADÉ (France), E. IKER (USA), O. LEDUC (Belgium)

National delegates and Scientific Committee: G. AZZALI (Italy) - A. BEHAR (France) -

K. BENDA (Czech. Rep.) - E. DIMAKAKOS (Greece) - E. ELIŠKA (Czech Rep.) - E. FÖLDI (Germany) -

M. FÖLDI (Germany) - I. FORNER-CORDERO (Spain) - P.S. MORTIMER (Great-Britain) -

W. OLSZEWSKI (Poland) - A. PECKING (France) - A. PISSAS (France) - A. SOUSA PEREIRA (Portugal) -

G. THIBAUT (France) - M. WALD (Czech. Rep.)

International Board of Trustees: MFC ANDRADE (Brazil) - M. WITTE (USA) - C. PAPENDIECK (Argentina) - M. OHKUMA (Japan)

Secretary: O. LEDUC (Belgium)

Treasurer: J.P. BELGRADO (Belgium)

ESL Awards: Caplan price (anatomie, clinical) one year and the other year Papamiltiades price (physiology or patho physiology).

Graphic, Design & Printing: Benjamin GD&P srl - Via Salvatore Rebecchini, 5 - 00148 Rome - Italy

Instructions to authors

General

Submission of an original article implies: that the work described has not been published before (except in the form of an abstract or as part of a published lecture, review, or thesis); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institute where the work has been carried out (including ethical committees and national licencing authorities); that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher; and that the manuscript will not be published elsewhere in any language without the consent of the copyright holders.

Manuscripts should be submitted in triplicate (original and two copies); they should be double-spaced, with wide margins on one side of the paper only, and should be carefully prepared in the style of this journal and checked before submission. Typing errors should be corrected legibly.

All manuscripts are subject to copy editing and, if necessary, will be returned to the authors for open questions to be answered or for missing information to be supplied before being sent to the printers. When extensive corrections are necessary, authors are responsible for having manuscripts retyped.

Pages should be consecutively numbered, starting with the title page. The desired position of figures and tables should be marked in the margin.

Changes in the proofs should be kept to a minimum: a charge will be made for changes introduced after the manuscript has been set in type.

Organization of the manuscript

The speed of publication depends greatly upon following these guidelines precisely.

1. The manuscript should be divided clearly into an Introduction, Materials and Methods, Results, Discussion and Conclusion and References. The text should be concise and consistent as to spelling, abbreviations, etc...

2. The **title page** should include the title of the work, first and last name(s) of author(s), name of institution, any footnotes referring to the title (marked with an asterisk), and the address of the author to whom the proofs are to be sent.

To facilitate communication between the authors, editors and publisher, the author should furnish a **telex or fax number** on the title page of the manuscript.

3. The **abstract** should be a summary of the hypothesis or aims of the work, the basic material and methods and the conclusion of the study.

4. Immediately following the abstract, up to 7 relevant **key words** should be supplied for subject indexing.

5. **Footnotes**, other than those referring to the title heading, should be numbered consecutively.

6. The accuracy of the **References** is the responsibility of the authors.

The list of references should only include works that are cited in the text and that have been published or accepted for publication. Personal communications should be mentioned in the text only. The list should be in alphabetical order according to the first author's name. Works by two authors should be listed alphabetically according to the second author's name, then chronologically; those by three or more authors should be in chronological order. References should be styled as follows.

Biancos J.A., Eimaleh D.R., Leppo J.I.A. (1986) Effect of glucose and insulin infusion on the myocardial extraction of a radioiodinated methyl-substituted fatty acid. *Eur. J. Nucl. Med.* 12: 120-124.
Gullberg G.T., Malko J.A., Eisner R.L. (1983) Bounday determination methods for attenuation correction in single photon emission computed tomography. In: Esser PD (ed). *Emission computed tomography: current trends*. Society of Nuclear Medicine, New-York, pp. 33-53.
Meltzer YL (1971) Hormonal and attractant pesticide technology. Noyes data, Park Ridge, New Jersey.

Citations in the text should be given in parentheses (Child 1941; Godwin and Cohen 1969; MacWilliams et al., 1970), except when the author is mentioned, as in "and the study of Hiliman and Tasca (1977)".

7. **Tables** should be submitted on separate sheets. Numerical data given in graphs and tables must not be duplicated.

8. All **figures**, whether photographs, graphs or diagrams, should be numbered consecutively throughout and submitted on separate sheets. Plate layouts or single figures may either match the width of the column (9 cm) or be 11.8 cm in width with the legend at the side.

The maximum height for a figure or plate is 23 cm, including the legend printed at its foot. Photographs can be grouped into plates. They must be mounted on regular bond paper, not on cardboard.

All photographs and electron micrographs should be supplied as high-contrast glossy prints trimmed at right angles. Inscriptions on illustrations should allow for reduction if this is necessary; figures and letters should have a final height of 2 mm after reproduction.

Color illustrations will be accepted: however, the authors will be expected to make a contribution (approximately BF 7.500 per page) to the additional costs involved.

9. Typewritten **mathematical equations** should be clear, so that there is no opportunity for misinterpretation by the printer.

All letters contained in formulae as well as single letters in the text are automatically set in italics and therefore require no underlining. Hence, abbreviations that appear in formulae and are to be set in roman type (the type normally used for the text) should be specially marked by underlining in yellow, if possible.

It will be helpful to the printer if *Greek characters are underlined in red and script in green*. Lowercase letters should then be underlined once and capital letters twice; this applies also to Latin letters in formulae (in pencil). Boldface type (heavy type) should be marked by wavy underlining.

Subscripts and superscripts should be indicated by an inverted caret below the line, or a caret above the line, respectively: 12_{12} ; a subscript to a subscript is styled: 12_{12} .

Obscure primes and dots must be clarified for the printer. The following must be differentiated clearly: number 1 and letter l; zero 0 and letters O, o, e, c, n, u, v, primes and apostrophes. Fractional exponents should be used in, stead of root signs and the solidus (/) for fractions whenever they are horizon. tal; an exp notation must be numbered sequentially in arabic numerals in parentheses on the right-hand side of the page.

10. Fifty (50) *offprints* of each paper with additional copies are available in lots of 100, (provided the order is teceived with the corrected proofs) may be supplied charged to the authors.

11. Enclose the picture of the first author of each article.

PODOCONIOSIS IN ETHIOPIA. A PILOT STUDY TO IMPROVE THE MANAGEMENT OF LYMPHEDEMA

ISABEL FORNER-CORDERO, MD. PHD., MARTA LOPEZ-AGUSTIN, MD.,
MARTA GUTIERREZ-DELGADO, MD.

Lymphedema Unit, Hospital Universitari I Politècnic La Fe, University of Valencia, Valencia, SPAIN.
Fontilles & Fundación ATA, SPAIN

Correspondence: Isabel Forner-Cordero, MD, PhD.
Physical Medicine and Rehabilitation Service
Hospital Universitari I Politècnic La Fe
Avda. Fernando Abril Martorell nº 106
46026 - Valencia
SPAIN

Telephone number: +34-961244000 extension: 411157.
E-mail: ifornercordero@gmail

ABSTRACT

Podoconiosis is a non-infectious cause of lymphedema, frequent in Africa. The aim of this study is to implement the management of lymphedema secondary to podoconiosis in a rural hospital in Southern Ethiopia. A call for donation of compression garments was made in Spain, to give any used and retired garment among lymphedema patients. The working plan was to train the staff in the lymphatic system, the causes of lymphedema, prevention strategies, skin care, intensive treatment and maintenance at long-term, Decongestive lymphatic treatment of the patients affected from podoconiosis and the adaptation of the donated garments for maintenance phase. Fifteen patients were treated with 10 sessions of Decongestive Lymphatic Therapy (DLT) consisting of Manual Lymphatic Drainage and multilayer bandages. The Percentage of Volume Reduction was 11.3% (95%CI: 8.8-13.9). Fifty percent of the patients showed an increase of the volume of lymphedema as they did not comply with the compression treatment. This pilot study has had good results in helping lymphedema management in Southern Ethiopia. The donation of used garments by chronic patients can be a feasible and effective step to improving the maintenance phase of patients with lymphedema in places with limited resources.

Keywords: lymphedema, podoconiosis, Decongestive Lymphatic Therapy, compression garments.

BACKGROUND

Lymphedema is a misdiagnosed disease even in occidental countries that leads to chronic and disabling complications when the treatment is delayed^{1,2}.

Podoconiosis is a non-infectious cause of lymphedema, frequent in some African countries (Figure 1). In Southern Ethiopia with a



Figure 1 - Podoconiosis.

prevalence of 5.46%, podoconiosis is an important chronic public health problem affecting both men and women³.

It is caused by the exposure of bare feet to irritant alkalic clay soils while working⁴. Colloid-sized silicate particles appear to enter through the skin, are taken up into macrophages in the lower limb lymphatics and cause endolymphangitis and obliteration of the lymphatic lumen⁴. It provokes not only medical problems of elephantiasis and repetitive dermatolymphangitis, but also a severe disability and a social stigma.

According to Davey, people with podoconiosis are excluded from school, denied participation in meetings, churches and mosques, and barred from marriage⁵.

There were considerable variations among community members' understanding of the causes of podoconiosis including genetic susceptibility, snakebite, direct contact of feet with clay soil, contagious, curses from God and poor nutrition. Other less commonly mentioned reasons were injuries (e.g., cut with axe), exposure to condensation, washing feet in hot water and the evil eye⁶.

Podoconiosis is unique in being an entirely preventable non-communicable disease⁴.

As the International Society of Lymphology states, decongestive lymphatic therapy is the base for conservative treatment of lymphedema⁷, and most of the guidelines of different countries follow this model of treatment^{8,9,10,11}.

OBJECTIVES

This pilot study aims to improve /to implement the management of lymphedema secondary to podoconiosis in a rural hospital in Southern Ethiopia.

METHODS

A call for donation of compression garments was made in Spain, to give any used and retired garment, even if deteriorated, in different associations of lymphedema patients (i.e. ACVEL) and in the Forum of Lymphedema in the internet (<http://www.med-foren.de/>). The collected garments, principally stockings were sent to the Rural Hospital in Gambo, Southern Ethiopia.

A physiatrist Dr. M Lopez-Agustin worked there for 6 months, between November 2010 to April 2011, sponsored by Fontilles and Fundacion ATA, to start a Rehabilitation centre in the hospital. Concerning podoconiosis, the working plan was:

- To train the staff in the lymphatic system, the causes of lymphedema, prevention strategies, skin care, intensive treatment and maintenance at long-term;
- Decongestive lymphatic treatment of the patients affected from podoconiosis;
- Adaptation of the donated garments for maintenance phase.

The staff received some theory lessons and instructions were put on the wall to be referred to as needed. The doctor was in charge of the diagnosis, of planning the treatment schedule and of the assessment of the results. At first, she performed the Manual Lymphatic Drainage and the bandages herself (Figure 2) but, then the physiotherapist was employed and she trained him in the bandage technique and in the adaptation of the stockings (Figure 3). Patients received 10 sessions of Decongestive Lymphatic Therapy (DLT) consisting of Manual Lymphatic Drainage and multilayer bandages¹². Once volume reduction was achieved, compression was necessary to maintain the reduction of the volume, and another member of the staff helped adapt the stockings using a sewing machine to start maintenance treatment (Figure 4 and 5). He was skillful in this job.



Figure 2 - Dr Lopez performing the bandages.



Figure 3 - The physiotherapist is assessing a patient.



Figure 4 and 5 - The adaptation of the donated compression garments to the patients.



Figure 5.

The assessment of the results was made by taking the volume of the limb (with tape measurements from the foot until the knee using Kunhke formula¹³) and recording the degree of satisfaction with the treatment.

RESULTS

Fifteen patients (14 women and 1 man) (30 limbs) with bilateral lower limb lymphedema secondary to podoconiosis attended the hospital and were treated with DLT.

The mean age was 20 years (range: 11-55), the mean of volume at baseline was 2319 ml (95%CI: 2154-2485) and the volume after treatment was 2083 ml (95%CI: 1889-2277).

The reduction of the lymphedema volume was significant in all the patients, with 10 sessions of treatment.

The Percentage of Volume Reduction, calculated as the reduction of absolute volume related to the volume at baseline, was 11.3% (95%CI: 8.8-13.9).

The maintenance phase was made possible to perform by means of the donated garments that were adapted by the staff.

After one month the Change of Volume was 0.8% from the post-treatment volume (95%CI: -9.1-10.7). Fifty percent of the patients showed an increase of the volume of lymphedema as they did not comply with the compression treatment.

DISCUSSION

The main limitation of this study is the small size of the sample. But it has to be taken into account that medical attendance in Ethiopia is reserved for few people. Therefore, we could only treat patients that were obliged by their husbands to come to the hospital due to their uselessness to work or by their families due to the difficulties in marrying them, or they live in the same village and did not need to travel to the hospital.

The published studies to the date show no difference between men and women affected by podoconiosis³, but the real situation at Rural Hospital of Gambo is very different. As Dr Lopez treated 14 women and a man with Podoconiosis, she realized that it was the disease of poverty because it affects people with very low resources, with no possibility to buy shoes and that go to work barefoot. It is also a women disease because it affects women that are triple exploited: by their race, their social class and their gender. On the other hand, she observed that their adherence to treatment depended on the possibility to be taken to the hospital by their husband to the prescribed sessions.

Despite the fact that Spanish patients only receive one garment per year, 42 garments were donated. It is nice to see, their disposition to help and their solidarity with poorer countries.

All the patients of our sample showed an improvement of their lymphedema volume, but the effectiveness was smaller than in our country. The main reasons for this smaller effectiveness could be the severity and chronicity of lymphedema in this sample, the presence of cutaneous complications such as hypodermatitis and papilomatosis and the several episodes of lymphangitis attacks they have suffered before treatment. Nevertheless, all the patients were very satisfied with the results.

This improvement in the management of lymphedema has generated:

- The appointment of a physiotherapist trained in lymphedema in the Rural Hospital of Gambo.
- An increase in the number of the patients coming to the hospital looking for attendance.
- A start to reduction of the social isolation of patients.
- A start in information and self-care

The most important limitation to the management of lymphedema in Gambo is the lack of patient's adherence to treatment. Possible reasons may include: the belief that there is no effective medical treatment, long distances to the hospital setting, and the little awareness of disease that these patients have. Moreover, the social status of women does not allow them to spend money and time on their health, considering that the economic cost of the treatment is high for them. The patients have to pay 60 birrs for the treatment (equivalent to 1.78 €) when their salary is 300 birrs per month (equivalent to 8.82 €). The maintenance treatment is also very difficult as patients continue walking barefoot and their garments get easily deteriorated. As other projects have reported in filarial elephantiasis, and despite the limitations, our results suggest that basic lymphedema management is feasible and effective in podoconiosis areas where resources are limited¹⁴.

As Ryan *et al.* reported after their experience in Ethiopia, also clean water and research in washing materials and soap are essential to improve the management of podoconiosis^{15,16}. A randomised controlled trial is being conducted in Ethiopia by Negussie *et al.* to demonstrate that community-based treatment of podoconiosis can reduce the frequency of acute dermatolymphangioadenitis episodes and improve other clinical, social and economic outcomes.

CONCLUSIONS

This pilot study has had good results in helping lymphedema management in Southern Ethiopia. The donation of used garments by chronic patients can be a feasible and effective step to improve the maintenance phase of patients with lymphedema in places with limited resources.

Future projects are ongoing. We are planning to send more compression stockings and bandages, shoes that are essential for the prevention of the disease, and educating material for skin care, prevention, bandages. Education and training in prevention strategies focusing in the importance on wearing shoes at work have to be planned. Given the importance of the problem, other help has been asked to go on with this project.

Table 1 - Patients in Hospital of Gambo.

Patients	(n=15)
Age (mean; 95%CI)	20 (11-55)
Gender: Women	14
Men	1
Etiology podoconiosis	15
LL Volume baseline (ml) (mean; 95%CI)	2319 (2154-2485)
LL Volume after treatment (ml) (mean; 95%CI)	2083 (1889-2277)

REFERENCES

- Moffatt C.J., Franks P.J., Doherty D.C. *et al.*: *Lymphoedema: an underestimated health problem*. QJM, 2003; 96(10): 731-8.
- Forner-Cordero I., Navarro-Monsoliu R., Muñoz-Langa J. *et al.*: *Early or late diagnosis of lymphedema in our Lymphedema Unit*. Eur J Lymphol Relat Probl., 2006; 16(49): 19-23.
- Destas K., Ashine M., Davey G.: *Prevalence of podoconiosis (endemic non-filarial elephantiasis) in Wolaitta, Southern Ethiopia*. Trop Doct., 2003; 33(4): 217-20.
- Davey G., Tekola F., Newport M.J.: *Podoconiosis: non-infectious geochemical elephantiasis*. Trans R Soc Trop Med Hyg., 2007; 101(12): 1175-80.
- Davey G.: *Podoconiosis, non-filarial elephantiasis, and lymphology*. Lymphology, 2010; 43(4): 168-77.
- Tekola F., Bull S., Farsides B. *et al.*: *Impact of social stigma on the process of obtaining informed consent for genetic research on podoconiosis: a qualitative study*. BMC Med Ethics, 2009; 10:13.
- International Society of Lymphology. *The diagnosis and treatment of peripheral lymphedema*. 2009 Consensus Document of the International Society of Lymphology. Lymphology, 2009; 42(2): 51-60.
- Eliska O., Benda K., Houdova H. *et al.*: *The founding and development of the Czech Lymphology Society*. Brief guidelines of the Czech Lymphology Society. Eur J Lymphol Relat Probl. 2006; 16(47): 1-6.
- Lee B., Andrade M., Bergan J. *et al.*: *Diagnosis and treatment of primary lymphedema*. Consensus Document of the International Union of Phlebology (IUP)-2009. Int Angiol., 2010; 29(5): 454-70.
- Campisi C., Michelini S., Boccardo F.: *Società Italiana di Linfangiologia. Guidelines of the Società Italiana di Linfangiologia: excerpted sections*. Lymphology, 2004; 37(4): 182-4.
- The Swedish Council on Technology Assessment in Health Care. *Manual lymph drainage combined with compression therapy for arm lymphedema following breast cancer treatment*. In: SBU alert [database online]. Stockholm: SBU; 2005. SBU alert report no 2005-04. Available: http://www.sbu.se/upload/Publikationer/Content0/3/Manual_Lymph_Drainage_Compression_Arm_Lymphedema_Breast_Cancer_200504.pdf
- Földi E., Földi M.: *Lymphostatic diseases*. In: Földi M. Textbook of lymphology for physicians and lymphedema therapists. Urban & Fisher. Munchen, 2003. p. 231-319.
- Kurz I.: *Textbook of Dr. Vodder's manual lymph drainage*. In: Harris RS, editors. Therapy. 2nd ed. Heidelberg (Germany): Karl S. Haug Publishers; 1989.
- Addiss D.G., Louis-Charles J., Roberts J. *et al.*: *Feasibility and effectiveness of basic lymphedema management in Leogane, Haiti, an area endemic for bancroftian filariasis*. PLoS Negl Trop Dis. 2010; 4(4): e668.
- Ryan T., Fuller C., Matts P.: *Management of mossy foot in Ethiopia*. J Lymphoedema, 2011; 6: 58-61.
- Ryan T.J.: *Dermatological care for all: Ethiopia November 24-27, 2009*. Int J Dermatol., 2010; 49(2): 236-7.
- Negussie H., Kassahun M.M., Fegan G., Njuguna P., Enquselassie F., McKay A. *et al.*: *Podoconiosis treatment in northern Ethiopia (GoLBet): study protocol for a randomised controlled trial*. Trials, 2015; 16: 307.

DERMAL-EDEMA: THE MICRO-EDEMA

MARCELLO IZZO*, ENRICO OLIVA**

*Research Center “Mathematics for Technology, Medicine and Biosciences”, University of Ferrara, Palazzo Manfredini - Via Muratori 9, 44121 Ferrara (corresponding author, email: zzimcl@unife.it)

*Compression Therapy study Group (CTG)

**Accademia di Storia dell’Arte Sanitaria – ASAS – Roma

**Gruppo di studio sul Linfedema - Società Scientifica Alleanza Flebolinefologica A.F.

ABSTRACT

The presence of the lower limbs of “fovea” indicates the objective presence of edema, however, the most recent literature has reported the presence of a subclinical edema called Dermal-Edema or Papillaredema. The relief of this form of edema was the prerogative of ultrasound examinations with high frequency probes (<20MHz), while today it is possible to identify it with the common ultrasound devices equipped with harmonics already with frequencies of 10-14MHz. This sort of microedema affects above all the papillary part of the dermis more than the reticular part because the particular richness of Glycosaminoglycan (hyaluronic acid, etc) of this dermal zone represents a sort of “dermal sponge” for the storage of water.

The increase in this area that can be identified on the ultrasound as “subepidermal-low-echogenic-band (SLEB)” immediately below the epidermal line may represent an early sign of future tissue dystrophy and is therefore useful during the execution of a venous echocolor Doppler of the lower limbs, dwell on the evaluation, above all comparative between the two arts, of the SLEB.

INTRODUCTION

Edema, also spelled “oedema” is an abnormal accumulation of fluid in the interstitium, in various tissues localized. The edema arises in just one area, for example a leg, or it can attack/infect different parts of the body, in that case, first the edema needs to be clinically evident, second litres of liquid have to accumulate; for this reason, most of the time, the weight grows before the edema comes out on lower limbs (fovea, etc.)

The formation of fluid in the interstitium, from the Staling’s equation:

Net ultrafiltration (F) = $K [(PI_{cap} - PI_{int}) - \sigma (PO_{cap} - PO_{int})]$

where **K** = coefficient of capillary filtration; **PI_{cap/int}** = hydrostatic and interstitial pressure; **σ** = theoretical limitation coefficient to the passage of proteins through endothelial pores; **PO_{cap/int}** = blood and interstitial oncotic pressure.

The pressure ultrafiltration effective, produces an ultrafiltration in extravascular direction, with a loss water containing small cells. On the other hand, **the effective reabsorption pressure** draws the interstitial liquids in the bloodstream. The amount of ultrafiltered water is not the same quantity of the absorbed water: the ultrafiltered water is 10% higher of the absorbed water.

Gross ultrafiltration means the quality of ultrafiltered water and **net ultrafiltration** means the difference between gross ultrafiltration and the quantity of resorbed liquid.

That net ultrafiltration matches to the physiological lymphatic load.

In the end we have a state of non-equilibrium condition, and a state of continuous water movement from the plasma to the interstitial compartment. This slight surplus, so the net filtration, however, is balanced by the amount of liquid who/which returns to the circulation through the lymphatics. So we can talk about the Edema when we have causes of increased oncotic pressure outside the blood vessels (for example, the inflammation), or reduces the oncotic pressure in the blood (for example, the cirrhosis). (Tab. 1)

Table 1 - Edema causes.

PHYSIOLOGY	POSSIBLE CAUSE	EFFECTS
↑ Capillary permeability (c)	Cellulite, arthritis, cyclic hormonal edema	Inflammatory edema, “idiopathic edema”
↑ Venous (capillary) pressure (Pc)	Heart failure, venous insufficiency, abstinence syndrome	Cardiac edema, venous edema
↑ Tissue oncotic oncotic (πt)	Insufficiency of lymphatic drainage (albumine increase)	Lymphedema
↓ Capillary oncotic pressure (πc)	Hypoalbuminemia, nephrotic syndrome, liver failure	Hypoproteinemic edema

An increase in hydrostatic pressure within the blood vessels (phlebotasis, orthostatic venous hypertension) or a reduction in the interstitial hydrostatic pressure eventually has the same proedemigenous effect. The lymphatic edema (lymphedema) is a hyperproteic edema and depends on the alteration of the lymphatic vessels. It is characterized by a reduced transport capacity, by a normal lymphatic load and by a high protein content edema. If the permeability of the capillary walls increases, will be released in the latter case we speak of inflammatory exudate or edema. Non-inflammatory edema, caused by alteration of the hemodynamic forces, is called “transudate”. The transudate has a

lower specific weight to 1012, it is poor in protein and does not contain inflammatory cells. sed very fluid from the capillary, such as when it happens in inflammations. Instead, the exudate is characterized by the other specific weight, greater than 1020, is rich in proteins and cells of inflammation. Venous edema, which is characterized by a normal transport capacity, a normal lymphatic load and a low protein concentration, an example is an edema due to increased hydrostatic pressure. Physiologically, the pressure at the arteriolo-capillary end is about 32-35 mmHg, while at the venular one it is about 12-15mmHg with a gradient of about 25mmHg. When forming a venous stasis for chronic venous insufficiency venular the hydrostatic pressure reaches values of approximately 35-50mmHg with the triggering of interstitial edema, which subsequently may 'cloinicamente become evident. In this way, chronic venous stenosis Microangiopathy is determined, a prelude to trophic sequences such as Lipodermatosclerosis (LDS) and varicose ulcers. When the lymphatic transport capacity (Földi safety valve) exceeds that of drainage of the lymphatic system, there is a stagnation of liquid in the interstitium with the genesis of edema (hyperprotein edema typical of lymphedema).

In the last decade the Starling model enunciated in 1986 has been replaced by the Michel-Weibbaum model ⁽¹⁻¹¹⁾ due to the importance of glycocalyx (venous glycocalyx consists essentially of 50-90% of heparan-sulfate and for a lower aliquot of chondroitin-sulfate and hyaluronic acid or Hyluronan) that would function as a sort of molecular sieve. In the new model of Michel-Weinbaum the true osmotic barrier does not correspond to the entire vessel wall with its endothelium, as defined by Starling, but rather to glycocalyx (molecular sieve) while the inter-endothelial pathways represent the preferential pathways of passage for albumin, etc. These recent scientific evidences point out that the damage of glycocalyx (glycocalyx-shedding), is a harbinger of strong increase in permeability with creation of interstitial edema, eritrodiapedesis, passage of albumine, etc. The extracellular and extravascular interstitial fluid represents 15% of the total body fluids (6-7 litres), and essentially consists of water and salts (especially NaCl) with a low protein content (0.3- 0.5 g/dl), ie in practice it is plasma without proteins. Albumins unlike globulins have different osmotic power ⁽¹²⁾:

1 gr of albumine = oncotic pressure of 6 mmHg

1 gr of globulins = oncotic pressure of 1.5 mmHg

a lower volume and molecular weight (compared to globulins) and therefore capable of attracting a greater quantity of water. In addition, albumins are proteins with negative electrical charges (anionic protein gap) able to attract cation ions (Fe ++, Fe +++) and to induce tissue acidosis ^(13,14). One of the most used formulas for the calculation of "the protein anionic gap" is as follows:

ANION GAP CALCULATION FOR ALBUMIN:


$$AG = (Na+) - (Cl- + HCO_3) \text{ (v.n. = 10-10mmmoL/L)}$$

Correzione per i valori dell'albumina = valore AG + [2,5 x (4- albumina)]

AG : Anion Gap

Human albumin is the most abundant serum protein (60% of the total protein) with an average concentration of about 3-5g / 100ml equal to it has a half-life of about 20 days and weighs about 69kDa. At physiological pH of the blood, albumin, besides

binding fatty acids and various other molecules, binds to water, playing a fundamental role in the determinism of blood protein oncotic pressure. The water is an electric dipole, and is the main constituent of the body and corresponds to 60/65% of body weight in men and 55-60% in women. The total amount of water tends to decrease with age. The total body water is distributed in two compartments, the intracellular and extracellular divided into intravascular and extravascular or interstitial. The interstitial liquid is essentially made up of water and salts (especially NaCl) with few proteins (0.3-0.5 g / dl), so is a plasma without proteins. The interstitial fluid is extracellular and extravascular and accounts for 15% of total body fluids (6-7 liters). As already mentioned before, weight gain, even of several kgs, precedes the appearance of the fovea, the demonstrative sign of edema. Some classifications in use differentiate according to the depth of the fovea and the time of disappearance 4 different stages of edema:



Dermal-Edema or Microedema.

In recent decades, literature has reported a new entity of non-clinically visible edema, therefore not improntabile with the fovea, called "DERMAL_OEDEMA" (DE) or PAPILLARY DERMAL-EDEMA (PPDE), but can be highlighted with 25MHz echography. The author described this from edema confined to the Derma as "subepidermal-low-echogenic-band (SLEB)", a thin hypoechoic band (water) immediately below the epidermis, later also referred to as "Sub- Epidermal- Non-Echogenic Band (SENEB)". The author in his studied had carried out ultrasound findings on the skin of the forearm in a large number of patients with the age between 0 years (newborn) and 90 years, remarking that this type of dermal edema was an early and important sign of skin and fibrosis aging. This form of invisible edema (or invisible swelling) has subsequently become an important marker in some rheumatological diseases, such as dermatomyositis, multiple sclerosis, etc., in which the dermal edema is a prelude to the subsequent dermal fibrosis. Furthermore, other authors (Gniadecka, 1996) reported the importance of Dermal-Edema (DE) in both lipodermatosclerosis from chronic venous insufficiency, in lymphedema ⁽¹⁶⁻¹⁹⁾ and in venous ulcers ⁽²⁰⁾. The dermis (thickness varying from 0.3 to 4mm depending on the different anatomical sites) consists of two different layers, the one near the epidermis called "papillary or superficial dermis" that contains cells (fibroblasts, mastocytes, histiocytes, etc.) and fibers (collagen fibers and elastin) also the amorphous substance

consisting of electrolytes, glycoproteins, proteoglycans, etc., (known as GAGs or mucopolysaccharides) strongly hydrophilic and therefore able to absorb water (hyaluronic acid an unperforated GAG) equipped with the highest hydrophilicity, 1 gram of hyaluronic acid is able to retain a volume of 3 liters of water). The concentration of hyaluronic acid varies with the years, in fact when we have 20 years old, the hyaluronic acid that we have in our body captures about 100% of the water, while at 30 years old it drops to 65% of the concentration of water held back by the fabric skin, at 50 years old at 45% and infinite at 65 years old the value drops to 25%. The papillary dermis is more cellular and less fibrillary and is constituted prevalently by loose connective fabric skin with collagen fibers, elastic fibers, rich in blood-lymphatic vessels and nerve endings (amyelinated nerve fibers, tactile corpuscles of Meissner, etc.). The Reticular Derma, is or deep or fibrillary, and is constituted mainly by robust collagen and elastic fibers oriented in a spatial way (so in the same direction) instead. This spatial organization ensures resistance to deformation of the skin. One need only think that the surgical incisions along the lines of orientation of these fibers, called Langer lines, have a better cicatrization. The organization and composition of the papillary edema explains why it is also called “dermal sponge”, as it has the ability to retain a lot of water is maximum in this segment of the dermis. Therefore the connective fabric skin (Derma Papillare) is essential for the storage of water in the body:

- a man with a body weight of 70 kg this storage is 3-6 lt. and the extracellular matrix holds 2/3 of it;
- in the healthy subjects in the ortho-clinostasis passage, the SKIN-THICKNESS or DERMAL THICKNESS (the thickness of the whole dermis measured between the epidermis and the hypodema) varies from 6-20% and after the physical activity the DERMAL-OEDEMA is reduced.

So, when water accumulates inside the skin, especially in PAPILLARY DERMAL-EDEMA (PPDE), will also increase the DERMAL THICKNESS or SKIN-THICKNESS, and if this increase in thickness exceeds 50-100% (mm) of its thickness, edema can be clinically visible, this is how invisible edema turns into visible edema (fovea). The presence of “DERMAL-EDEMA” determines the disruption of the collagen bundles (spatial orientation of the collagen fibers) with reduction of the density to the ultrasounds (low echodensity) and such reduction of the dermal density to the ultrasounds (as consequence of the “DERMAL-EDEMA”) has been confirmed both with MRI and spectroscopy⁽²¹⁻²³⁾. This storage of water in the dermis (in particular in the papillary dermis) with the presence of the “DERMAL-EDEMA” is also confirmed by other authors⁽²⁴⁾ which point out the marginal in terms of the hypodermis in water storage. Postural edema, which tends to increase with age⁽²⁵⁾, may be visible (fovea) or remain invisible (dermal-edema) and this may depend on the reduction of the absorbing capacity of the papillary dermis for the progressive reduction of hyaluronic acid. Another aspect is the fact that the edema to be able to reabsorb with the position of the lower limbs antideclive requires a time not less than 3 hours⁽²⁴⁾. So in the end we become more susceptible to developing postural edema with the years and, moreover the edema is formed faster than its disappearance with the antideclive position.

An example of what is described is shown in fig. 1.




Fig. 1 - In (A) and (B) we can see how is changing rapidly (5min.) the echogenicity and thickness of the SLEB detected on the back of the hand after 5 minutes of adducted and abducted upper limb (14MHz with THI: Thissue Harmonic Image). Today with the modern ultrasound equipment equipped with the harmonic frequencies (THI: Tissue Harmonic Image) with frequencies 10-14 MHz, it is possible to see the SLEB that corresponds to the Dermal-Edema (DE).

HYPODERMIC EDEMA: ECOGRAPHIC DIFFERENCES

The edema of the subcutis, even if it is not able to give exact information, can manifest itself with different ultrasound images as reported in Fig. 2.




Fig. 2 - The different possible ultrasound images of the subcutaneous edema.

Some authors⁽²⁷⁾ in the lymphedema of the lower limbs describe, with an 11 MHz ultrasound probe, different degrees of echogenicity (SEG: Subcutaneous Echogenicity Grade) of the hypodermic edema (Fig. 3).




Fig. 3 - Different degree of echogenicity (grade 0, 1, 2) of the subcutis in secondary limb secondary lymphedema (from Suehiro K, 2013).

With ultrasound equipment with a higher frequency (20MHz) some authors⁽¹⁷⁾ show that intradermal echogenicity shows characteristic patterns in different types of edema (lymphedema, lipodermatosclerosis secondary to chronic venous insufficiency, heart failure edema) which indicates that the localization of fluid (water) can be disclosed in different ways:

- SURFACE PAPILLARY DERMA in Lipodermatosclerosis (LDS);
- DERMO-HYPODERMIC in Lymphedema;
- DEEP RETICULAR DERMA in heart failure.

These differences can help to understand the etiology but above all to prevent the possible temporal skin sequelae that edema can induce. The presence of Dermal-Edem (SLEB: subepidermal-low-echogenic-band) constant in a patient with chronic venous insufficiency is a marker of dystrophic risk and of dermal elastosis and therefore it is necessary to get used during the execution of the ecodoppler test, which is normally performed in such phlebopathic patients, the presence of Dermal - EDEMA (SLEB: subepidermal-low-echogenic-band) so the subepidermal hypoechoic band and compare it with the other healthy limb or with different zones of the same limb (Fig. 4).




Fig. 4 - Comparison between peri-lipodermatosclerosis (LDS) and lipodermatosclerotic tissue, on the right note the presence of dermal- edema above the image of the varices, while in the left image the disappearance of SLEB with replacement of hyperechoic reticular dermal tissue (fibrosis) with increased dermis (DERMAL THICKNESS or SKIN-THICKNESS). D2 = papillary dermis = 2.4mm; D3: reticular dermis = 9.3 mm.

Recently, some authors⁽²⁸⁾ report the importance of early echographic skin changes in the different CEAP-C states of chronic venous disease (MVC), highlighting the importance of the evaluation of SLEB (Dermal- Edema) sometimes present in the CEAP-C2 stage. as an early marker with respect to subsequent ultrasound alterations typical of the more advanced stages (Fig. 5).

CEAP -	PARAMETRI ECOGRAFICI
CEAP-C2	Dermal edema -flogosi cute e/o ipoderma
CEAP-C3	Dermal edema- lacune anecogene- ispessimento (thickening) omogeneo dell'ipoderma.
CEAP-C4a	non alterazioni tipiche per la melanos Se eczema -ispessimento flogistico cute- sottocute, giunzione dermo-ipodermica poco visibile o assente.
LDS (CEAP-C4b)	-epidermide poco visibile -ispessimento cute-sottocute (iperecogeno) -giunzione dermo-ipodermica assente - sclerosi e assottigliamento cute o -iperproliferazione sottocute(iperecogeno) -dermal -edema (aumento skin thickness) -rarefazione sottocute (RISCHIO UL.)

Fig. 5.

It is also reported that patients in the CEAP-C2 class who have dermal-edema (SLEB) would be more symptomatic than those who do not have this form of edema and also that a small difference in diameter (+ 1.0mm) of a varicose vein may be responsible for the appearance of dermal-edema (Fig. 6).



Fig. 6 - In orthostatism the evaluation of the SLEB and the subsequent SKIN or DERMAL-THICKNESS of two varices positioned at the same height with respect to the plantar support surface and the same cutaneous depth in the two limbs in the same patient, where the minimum increase of diameter (+ 1.0mm) determines the appearance of dermal thickening.

In conclusion, with the arrival of modern ultrasound equipment equipped with harmonics it is possible to highlight early extravenous alterations of the surrounding peri-varicose tissues, premonents of possible dystrophic disturbances already with 12 12 MHz frequencies. Moreover, as previously mentioned, the presence of greater damage to glycocalyx (Michel-Weibaum model) may lead to greater permeability with edema stasis. So today, For all these reasons, the ecocolor Doppler study of phlebopathic patients must include in addition to hemodynamics, also the evaluation of perivenous tissues with comparison with

other skin areas and / or between the two limbs. This ultrasound evaluation allows to go beyond the clinical evaluation until now (CEAP-C) representing a sort of “ultrasonic biopsy” and the microedema (DERMAL-EDEMA) is certainly an important datum to consider useful also for therapeutic monitoring⁽³⁰⁾.

BIBLIOGRAPHY

- Weinbaum S., Tarbell J.M., Damiano E.R.: *The Structure and Function of the Endothelial Glycocalyx Layer*. Annu. Rev. Biomed. Eng., 2007; 9: 121-67.
- Woodcock T.E., Woodcock T.M.: *Revised Starling equation and the glycocalyx model of transvascular fluid exchange: an improved paradigm for prescribing intravenous fluid therapy*. Br J Anaesth., 2012; 108(3): 384-94.
- Levick J.R., Michel C.C.: *Microvascular fluid exchange and the revised Starling principle*. Cardiovasc Res., 2010; 15; 87(2): 198-210.
- Olson K.R., Kinney D.W., Dombkowski R.A., Duff D.W.: *Transvascular and intravascular fluid transport in the rainbow trout: revisiting Starling's forces, the secondary circulation and interstitial compliance*. J Exp Biol., 2003; 206(Pt 3): 457-67.
- Facchini L., Bellin A., Toro E.F.: *A mathematical model for filtration and macromolecule transport across capillary walls*. Microvasc Res., 2014; 94: 52-63.
- Jacob M., Chappell D.: *Reappraising Starling: the physiology of the microcirculation*. Curr Opin Crit Care, 2013; 19(4): 282-9.
- Pillinger N.L., Kam P.: *Endothelial glycocalyx: basic science and clinical implications*. Anaesth Intensive Care, 2017; 45(3): 295-307.
- Damiano E.R.: *The effect of the endothelial-cell glycocalyx on the motion of red blood cells through capillaries*. Microvasc Res, 1998; 55(1): 77-91.
- Tuma M., Canestrini S., Alwahab Z., Marshall J.: *Trauma and Endothelial Glycocalyx: The Microcirculation Helmet?* Shock, 2016; 46(4): 352-7.
- Chappell D., Jacob M.: *Role of the glycocalyx in fluid management: Small things matter*. Best Pract Res Clin Anaesthesiol, 2014; 28(3): 227-34.
- Lennon F.E., Singleton P.A.: *Hyaluronan regulation of vascular integrity*. Am J Cardiovasc Dis, 2011; 1(3): 200-13.
- Izzo M., Napolitano L., Coscia V. et al.: *The role of diamagnetic pump (CTU mega 18) in the physical treatment of limbs lymphoedema. A clinical study*. The European Journal of Lymphology, 2010, Vol. XXI, Nr. 61; 24-29.
- Feldman M., Soni N., Dickson B.: *Influence of hypoalbuminemia or hyperalbuminemia on the serum anion gap*. J Lab Clin Med, 2005; 146(6): 317-20.
- Carvounis C., Feinfeld D.: *A simple estimate of the effect of the serum albumin level on the anion gap*. Am J Nephrol 2000; 20: 369-72.
- de Rigal J., Escoffier C., Querleux B. et al.: *Assessment of aging of the human skin by in vivo ultrasonic imaging*. J Invest Dermatol, 1989; 93(5): 621-5.
- Gniadecka M.: *Dermal oedema in lipodermatosclerosis: distribution, effects of posture and compressive therapy evaluated by high-frequency ultrasonography*. Acta Derm Venereol, 1995; 75(2): 120-4.
- Gniadecka M.: *Localization of dermal edema in lipodermatosclerosis, lymphedema, and cardiac insufficiency. High-frequency ultrasound examination of intradermal echogenicity*. J Am Acad Dermatol, 1996; 35(1): 37-41.
- Gniadecka M., Karlsmark T., Bertram A.: *Removal of dermal edema with class I and II compression stockings in patients with lipodermatosclerosis*. J Am Acad Dermatol, 1998; 39(6): 966-70.
- Gniadecka M., Serup J., Søndergaard J.: *Age-related diurnal changes of dermal oedema: evaluation by high-frequency ultrasound*. Br J Dermatol, 1994; 131(6): 849-55.
- Hu D., Phan T.T., Cherry G.W., Ryan T.J.: *Dermal oedema assessed by high frequency ultrasound in venous leg ulcers*. Br J Dermatol, 1998; 138(5): 815-20.
- Knaus R., Fleischer G., Grunder W. et al.: *Pulsed field gradient NMR and nuclear magnetic relaxation studies of water mobility in hydrated collagen*. Magn Reson Med, 1996; 36: 241-8.
- Richard S., Querleux B., Bittoun J. et al.: *Characterization of skin in vivo by high resolution magnetic resonance imaging: water behaviour and age-related effects*. J Invest Dermatol, 1993; 100: 705-9.
- Gniadecka M., Quistorff B.: *Assessment of dermal water by high frequency ultrasound: comparative studies with nuclear magnetic resonance*. Br J Dermatol, 1996; 135: 218-24.
- Eisenbeiss C., Welzel J., Eichler W., Klotz K.W.: *Influence of body water distribution on skin thickness: measurements using high-frequency ultrasound*. British Journal of Dermatology, 2001; 144: 947-51.
- Gniadecka M.I., Gniadecki R., Serup J., Søndergaard J.: *Impairment of the postural venoarteriolar reflex in aged individuals*. Acta Derm Venereol, 1994; 74(3): 194-6.
- Becker M. et al.: *Sonography of subcutaneous tissue cannot determine causes of lower limb edema*. Vasa, 2015; 44(2): 122-8.
- Suehiro K., Morikage N., Murakami M. et al.: *Significance of ultrasound examination of skin and subcutaneous tissue in secondary lower extremity lymphedema*. Ann Vasc Dis, 2013; 6(2): 180-8.
- Caggiati A.: *Ultrasonography of Skin Changes in Legs with Chronic Venous Disease*. Eur J Vasc Endovasc Surg, 2016; 1-9.
- Labropoulos N.: *Commentary on “Ultrasonography of Skin Changes in Legs with Chronic Venous Disease”*. Eur J Vasc Endovasc Surg, 2016; 52(4): 543.

SUGGESTIONS FOR EDUCATION - TRAINING IN LYMPHOLOGY - DO WE NEED IT?

EVANGELOS P. DIMAKAKOS MD, PHD, EDA/VM, MLD/CDT

Vascular Internist

Vascular Unit of 3rd Internal Medicine Department, Sotiria General Hospital Athens School of Medicine, Athens, Greece

Incoming President of Greek Society of Lymphology

Member of Scientific Committee of European Society of Lymphology

Member of E.C. of ESL and of ISL

Corresponding Author: edimakakos@yahoo.gr

ABSTRACT

The European Society of Lymphology (ESL) is a society for lymphology in Europe and it is the official European scientific organ on Lymphology. The educational level of the Lymphology among the European countries varies from very low –poor until very high. The ESL must stimulates and gives incentives to the persons, societies, countries in order to develop the lymphology in Europe. Lymphology has begun to develop in Europe and generally all over the world especially after the establishment and the big efforts of the International Society of Lymphology, European Society of Lymphology and others very significant relevant societies. But in our days have begun to appear bad phenomena concerning malpractice both for the diagnosis and treatment of the lymphatic diseases and for the management of the patients. So it is more needed to have educational program accepted from all the Europe in order to establish limits and to define the terms of certified lymphologists and/or the therapists of lymphology and to exclude not certified persons. In this article are described some suggestions for the educational program of the scientific lymphology: what, how and why to give education and training and in Lymphology.

Key words: Lymphology, Education, Training, Program

INTRODUCTION

The European Society of Lymphology (ESL) is a society for lymphology in Europe and it is the official European scientific organ on Lymphology. There are many countries in Europe without any educational level Lymphology or educational Lymphology with a very low level. The ESL must stimulates and gives incentives to the persons, societies, countries in order to develop the lymphology in Europe. Lymphology and general the lymphatic diseases with the most known lymphedema employs many health professionals and especially surgeons, plastic surgeons, pathologists, oncologists, internists, vascular internists, vascular surgeons, radiologists, radiotherapists, dermatologists, medical persons that are specialized in Infectious Diseases,

gynecologists, anesthetists and more other medical specializations and nurses, physiotherapists, psychologists, persons for diet and for gymnastic⁽¹⁾. Lymphoedema is a serious disease that since its appearance not only creates intense psychological problems to the patients but it changes their lifestyle and relationships with other people and it affects very significantly in the quality of life of the patient. Unfortunately, the above complications- mainly occur after surgical or conservative treatment of cancer- has been abandoned by physicians and health professionals all over the world. Patients feel helpless and disappointment when they find out that lymphedema could have been prevented and now they have the disease and moreover they have no someone specialist to advise and guide them.

The lymph is drained to the lymph vessels and lymph nodes to the bloodstream via the thoracic duct. When the lymphatic system is malfunctioning then the fluid accumulates in the interstitial space and increases the osmotic pressure resulting in edema and swelling of the limb. Lymphedema is the abnormal accumulation of protein-rich fluid in the interstitial space which creates chronic inflammation and reactive fibrosis tissue array⁽²⁾.

Thus edema increase slow but steady increase the weight of the limb, decrease the mobility of the limb, it will appearance pain, weakness, stiffness and finally leads to the loss of functionality of the limb.

Once the patient is to display or has been diagnosed as suffering from lymphedema should then come in contact directly with the specialist physician or medical professionals to provide the appropriate treatment. Lymphedema is a medical condition with many other diseases and complications! We need a close cooperation of a specialist of Lymphology (which can be anyone as long as he specialized in Lymphology) with other medical specialists as well as nurses, physiotherapists, trainers, psychologists, nutritionists and other specifications for the proper and efficient treatment of lymphedema⁽³⁾. It is prohibited any treatment unless it is under the close supervision of qualified physician of lymphology or of qualified specialist of lymphology!⁽³⁾ The treatment aims to decrease the edema (fluid, fibrosis),

restoring the mobility of the limb, the prevention of infection, improve the aesthetics of the limb and generally improve the quality of life of the patient.

According to the ISL consensus the treatment of the lymphatic disorders is separated in conservative and in surgical treatment⁽³⁾.

In addition, according to internationally accepted guidelines the conservative treatment of choice is the Complete Decongestive Therapy (CDT). The treatment is divided into two stages, the intensive phase that lasts at least two to four weeks with at least one session a day and the maintenance phase to maintain the results achieved in the first stage. The first stage consists of:

- a) the care of the skin and nails,
- b) manual drainage of lymphatic fluid,
- c) elastic bandage,
- d) exercises,
- e) education of the patient.

The last element of CDT is very important and should always be taught to the patient.

The second stage, conservative, takes place in the patient's home where the same applies with the first stage only the position of strapping the patient has an elastic glove or sock scheduled.

Because lymphedema is a chronic disease, the patient should be monitored regularly by a qualified physician of lymphology or of qualified specialist of lymphology and its team and many times, if it is necessary, repeat the first intensive stage.

It is forbidden to carry out the above treatment if physicians, therapists and other trainers are not specially trained⁽³⁾. Effective treatments are those that lead to a reduction of more than 50% of the initial swelling at the end of the intensive phase. It should be noted that treatment failure when followed only one or two of these components of CDT as reported in the literature^(4, 5, 6).

Lymphology has begun to develop in Europe and generally all over the world especially after the establishment and the big efforts of the International Society of Lymphology, European Society of Lymphology and others very significant relevant societies. But in our days have begun to appear ugly phenomena concerning lymphology and malpractice of the treatment of lymphedema. Phenomena involving malpractice both for the diagnosis and treatment of the lymphatic diseases and for the management of the patients. So everyone has to follow some necessary Lymphology instructions which are involved in the diagnosis and the treatment of Lymphatic Disorders. Moreover the World Health Organization (WHO) wants everyone will be educated with special guidelines⁽⁷⁾. Below are suggestions according WHO according National Lymphedema Network, according ISL and ESL and according the educational programs-seminars existing in Europe⁽⁸⁾:

1. Everyone who is involved in the diagnosis and the treatment of Lymphatic Disorders must be certified and expertise in them. Because currently there is no systematic training in Europe proposed, for the time being, to be certified those that they have already certification from seminar or faculty as Foeldi, Leduc, Vodder and many others famous and unknown 'schools'. These training centers must meet some of the criteria listed below and must have the approval of ESL.

2. When someone takes a certification from any above specialized center (approval by ESL) can provide it to the specially committee of ESL in order to be a member of ESL and to be recognized by all over the Europe and especially in his local place.
3. He will take a Diploma called European fellowship on Lymphology. If the committee of ESL believes that the candidate or the center that has graduated has any deficiency can suggest more examination or anything else more.
4. In order to help to spread out the lymphology the ESL needs more instructors and specially in the countries with low level of Lymphology. For the begging ESL can define 2-3 instructors in each country (national members of ESL and/or national or local committee can decides or suggests about the number and names of the instructor). After that the every new instructor has to meet some of the criteria listed below.
5. The treatment should be done by a certified team of lymphology (must be at least a collaboration of a one physician and a physiotherapist).
6. Each treatment on any lymphatic disease which is made by other certified therapists (except physicians) should always be done under the close supervision and recommendations of a certified physician.
7. Every bad, false and immoral practice concerning the treatment of the lymphatic disorders without the certification and the appropriate experience and it will lead to the impairment of lymphatic disease of the patients and their quality of life, should be punished by the special committee of ESL and his local lymphology society.
8. The special committee of ESL for the recognition of the diploma is required to have: at least three members where all members are already certified and include at least one physician. The committee will examine all relevant certification and training of Lymphology in Europe. All require approval by the Board of Scientific Committee of ESL and from the general assembly of ESL.
9. In the event that questions arise regarding treatment or education or anything else related to the Lymphology can be solved if officially asked the ESL.
10. The certifications must be given during the European Congress of ESL and/or in other event (if executive committee of ESL decides) with a celebration.

As the knowledge and technical approach to diagnosis and treatment for lymphatic diseases develops we need continuing education from certificate training centers with high level of standards for those persons dealing with lymphatic diseases and generally with the Lymphology.

To have an adequate knowledge base in pathophysiology of lymphatic function and its disorders, as well as adequate training in the techniques and principles of manual massage, compression bandaging, together with other components of complete decongestive therapy, the minimum-criteria for training in the treatment of lymphedema are as follows:

- A) Students in the treatment of patients with lymphedema have successfully completed at least 135 hours of complete

decongestant therapy into a single program. The hours should not be confused with others hours that one can do: additional course or courses at advanced levels. It is important to have a single education program and that must have at least the following topics:

- Anatomy, physiology and pathophysiology of the lymphatic system
- The classification and staging of lymphedema
- Diagnosis and differential diagnosis of lymphedema
- Manual lymphatic drainage (mld)
- Bandaging and compression therapy
- Practice of bandaging of upper and lower extremities
- Indications and contraindications of manual lymphatic drainage and CDT
- Exercises for patients with lymphedema
- Patient evaluation and weekly measuring (circumference and volume)
- Basic and advanced techniques and sequences of mld
- Treatment of primary and secondary lymphedema of the extremities
- Other methods of treatment except CDT
- Treatment of head and neck cancer and genital lymphedema
- Skin care and nail for patients with lymphedema
- The importance of healthy skin and nail care for lymphedema
- Measurement techniques and how to wear gloves and stockings
- Teaching and training the patient to continue treatment at home.

B) It is necessary that one third (1/3) of hours of training, at least 45 hours, be theoretical training. The rest (2/3) of hours of training, at least 90 hours should be practical training with patients. During the course there will be oral and written examinations to assess the adequacy of the skills of the learner, while at the end of the study will be the final exam on the theoretical and practical part of education.

C) The book study of the students could be either the text book of lymphology of Foeldi or other book (it depends on each country) or the book of the VAS which is preparing. VAS is the official organ of UEMS for the specialization of European Angiology. ESL has strong collaboration with VAS and the VAS with the UEMS. All the books must be approval by ESL.

D) The trainee after the completion of the examination should continue to follow courses, seminars, congresses related to lymphology and with credits of CME.

E) The trainee shall be good after the end of his training to become member of ESL.

F) To be Instructor must a) be trained according to international standards and at least according to above basic program, b) be certified as a member (if there is in each country) in local or national society, c) to be member of ESL, d) be certified in lymphatic diseases by ESL with the diploma of European Fellowship on Lymphology, e) have proven at least 5 years experience in dealing with lymphatic disorders and proven credits of continuing education (CME), f) have proven teaching

experience in Lymphology least two years at the local level (each country), g) to make teaching courses in recognized educational centers abroad in European educational congress as ESL as workshops, conferences related to lymphology or teaching educational courses under the auspices of Foeldi or Leduc or Vodder or anything else - always all approval by ESL. The instructor will take a certification called European Fellowship on Lymphology-Instructor.

If someone has already the certification of instructor of Lymphology from recognized schools he must be recognized by ESL also, with the above criteria, in order to take the above certification.

Concerning the surgical educational side of Lymphology: There are different techniques regarding to the surgical management of lymphedema and of others lymphatic disorders and different official centers. Surgical techniques could be microsurgery, transplantation of lymph vessels and or lymph nodes, liposuction or others techniques.

A) The trainee must attend, study and follow the works of the surgical center.

B) The duration of the training depends on different specialization (ex microsurgery or transplantation) of surgical center. The centers must send its program of training to ESL and the center must has the official approval of training of ESL.

The material that the trainee has to study depends on each center but it must be approval by ESL.

C) After having complete the training the student, the head of the center must give a certification of attendance.

D) The student will have the possibility to take a more Diploma from ESL called European Fellowship on Microsurgery of Lymphology or European Fellowship on Liposuction of Lymphology ecc.

E) To be Instructor must a) be trained according to international standards and at least according to basic surgical program of the official center, b) be certified as a member (if there is in each country) in local or national society, c) to be member of ESL, d) be certified in lymphatic diseases by ESL with the diploma of European Fellowship on 'Surgical' Lymphology, e) have proven at least 5 years experience in dealing with lymphatic disorders and proven credits of continuing education (CME), f) have proven teaching experience in Lymphology least two years at the local level (each country), g) to make teaching courses in recognized educational centers abroad in European educational congress as ESL as workshops, conferences related to lymphology or teaching educational courses under the auspices of Campisi, Brosnor, Baumeister, Bacer or anything else - always all approval by ESL. The instructor will take a certification called European Fellowship on 'Surgical' Lymphology-Instructor.

F) If someone has already the certification of instructor of 'Surgical' Lymphology from recognized schools he must be recognized by ESL also, with the above criteria, in order to take the above certification.

G) The special committee for diploma of surgical training of ESL

is required to have: at least three members where all members are already certified and include at least two physician of surgical specialization. The committee will examine all relevant certification and training of Lymphology in Europe. All the latter is required approval by the Board of Scientific Committee of ESL and from the general assembly of ESL.

REFERENCES

1. Ethel Foeldi, Evangelos P. Dimakakos: *Interdisciplinary Lymphology: The Best Place for Each Discipline in a Team*. Int J Low Extrem Wounds, 2012 Mar; 11(1): 28-30.
2. Foeldi M., Foeldi E.: *Physiology and pathophysiology of the lymphatic system*. In: Foeldi M., Foeldi E., Kubik S. (eds) "Textbook of Lymphology". 1st ed. Munich, Germany: Urban & Fischer; 2003: 195-230.
3. International Society of Lymphology: *The Diagnosis and Treatment of Peripheral Lymphedema: 2013 Consensus Document of the International Society of Lymphology*. Lymphology, 2013; 46(1): 1-11.
4. Zimmermann A., Wozniowski M., Szklarska A., Lipowicz A., Szuba A.: *Efficacy of Manual Lymphatic Drainage in Preventing Secondary Lymphedema after Breast Cancer Surgery*. Lymphology, 2012; 45(3): 103-112.
5. Johansson K., Albertsson M., Ingvar C., Ekdahl C.: *Effects of compression bandaging with or without manual lymph drainage treatment in patients with postoperative arm lymphedema*. Lymphology, 1999; 32(3): 103-110.
6. Johansson K., Tibe K., Weibull A., Newton R.U.: *Low intensity resistance exercise for breast cancer patients with arm lymphedema with or without compression sleeve*. Lymphology, 2005; 38(4): 167-180.
7. *Transforming and scaling up health professionals' education and training: Recommendations by WHO* <http://www.who.int/hrh/education/en/>
8. National Lymphedema Network Position Papers: *The Diagnosis and Treatment of Lymphedema*. www.lymphnet.org

MEASURING LEG VOLUME IN LYMPHOEDEMA USING OPTOELECTRONIC VOLUMETRY AND THE SEQUENTIAL TAPE METHOD

NYREE DUNN¹, JANE H DAVIES², MICHELLE FISHBOURNE³, GINA DOLAN¹, AND E MARK WILLIAMS¹

¹Faculty of Life Sciences, University of South Wales, ²Centre for Trials Research, Cardiff University,

³Dewi Sant Hospital, Cwm Taf University Health Board, Pontypridd, UK

Correspondence: Nyree Dunn,
Faculty of Life Sciences,
University of South Wales
Pontypridd, CF37 1DL, UK
nyree.dunn@southwales.ac.uk

ABSTRACT

Objective: To assess the level of agreement between two techniques for measuring leg volume, infrared optical volumetry (OEV) (Perometer 400T, Pero-System GmbH, Wupertal, Germany) and a sequential tape measure (STM) method in lymphoedema patients.

Design: Data were collected during a 6 month randomised control feasibility trial which assessed the effect of intermittent pneumatic compression (IPC) plus standard lymphoedema versus standard lymphoedema treatment alone, on lower limb volume.

Leg volume data were categorised into three groups, A) controls, non-oedematous legs from those patients with unilateral disease, B) legs with lymphoedema from patients with unilateral and bilateral lymphoedema but not receiving IPC, and C) oedematous legs as B, but receiving IPC.

Setting: Lymphoedema outpatient clinic, Dewi Sant Hospital, Pontypridd, UK.

Participants: Twenty adults (age range: 25-78 years, male/female: 30:70) with stage II or III lower limb lymphoedema were recruited. Ten participants had bilateral lymphoedema, and 10 unilateral lymphoedema. Non-oedematous legs were assigned to be control limbs (Group A).

Interventions: Observational data.

Main Outcome Measures: Lower limb volume.

Results: The range of leg volume measured by both methods were 2862 to 7021 mls. The level of agreement between the OEV and STM methods was 7.2% with lower limb volume underestimated by 324 mls using the STM method. The correlation between the two measurement techniques was strong with a fitted linear regression correlation coefficient, r , 0.99.

Conclusion: The study shows that the two methods are strongly correlated however between the STM and OEV methods there is an offset of 324 mls'. Both methods are easy to use, both for the patients and health staff, but the automated OEV method is quicker and provides an instant measure, while the STM is slower, it is more economically suited for the clinic.

Keywords: Perometer, tape measure, lymphoedema, leg, volume.

INTRODUCTION

Lymphoedema is a chronic condition that causes swelling in the body tissues due to an excess accumulation of protein rich fluid called lymph. While it is more commonly seen in the arms and legs, it can affect any part of the body with a key characteristic being an increase in limb volume (Lymphoedema Framework, 2006).

An accurate measurement of limb volume is essential for both clinical and research settings. Current treatment for lymphoedema is based on decongestive lymphatic therapy (DLT); there are four components to such therapy, compression, skin care, exercise and manual lymph drainage (Lanski, 2013). More recently, pneumatic compression devices have been considered as an additional treatment modality for treating lymphoedema (Zaleska et al., 2014). Thus, limb volume is an important outcome measure to ascertain the effectiveness of such decongestive treatments, such as compression therapy (Williams and Whitaker, 2015).

The gold standard technique considered for measuring limb volume is water displacement (Kaulesar et al., 1993). However, this method is time consuming and is not always seen as a practical technique to measure limb volume in a clinical setting. Circumferential measurements using a tape measure are more commonly used in clinical settings and have been shown to correlate with the water displacement method (Tewari et al., 2008). The preferred technique for circumferential measurements involves measuring the circumference of the leg just above the malleolus and then at 4cm intervals ascending the leg up to a point that is clinically significant for each patient (Lymphoedema Framework, 2006). Measurements can then put into preprogrammed calculators or computer software to calculate limb volume. The most common formula to calculate limb volume is the volume of a truncated cone; this is where the limb is seen as a series of frustum shaped segments ($\text{segment volume} = L/12\pi(C1^2 + C1 C2 + C2^2)$). $C1$ and $C2$ are the circumferences at either end of the segment length (L). Total limb volume is the sum of the segment volumes (Williams and Whitaker, 2015). However, the reliability of this method is reliant on the technique, with a tendency to overestimate or underestimate limb volume (Deltombe et al., 2007).

Alternative measurements of limb volume can be obtained from an optoelectric volumeter, otherwise known as a perometer. The perometer was first developed in the 1980s and validated using geometric objects (Stanton et al., 1997). It has been used in many clinical studies but recently Tan et al., (2013) questioned its level of agreement with the widely used tape measure method, albeit they tested a healthy population.

An aim of this feasibility study was to assess the agreement between the two methods of volume measurement in patients with unilateral and bilateral lymphoedema in a clinical setting.

METHODS AND MATERIALS

Participants

Twenty adult participants (Age range: 25 -78 years, mean \pm SD, 50 ± 15 ; male to female ratio = 30:70, weight 94 ± 36 Kg, BMI 33 ± 11) with stage II or III lower limb lymphoedema were recruited onto the study. Half of the participants (n=10) had unilateral and half bilateral lymphoedema (n=10). Population data were taken at baseline, along with measured leg volumes which were repeated at 3 month and 6 month time-points. Ethical approval for this study was obtained from the local NHS Research Ethics Committee (LREC No 17/WA/0076). All participants provided written consent. Three participants withdrew from the study at 3 months and one at 6 months.

Protocol

Participants' gender, age, height and weight were recorded at baseline (Table 1).

Next participants underwent both right and left leg volumes (including the nonoedematous legs in participants with unilateral lymphoedema) measurement using the two different methods, optoelectronic volumetry, OEV (Perometer 400T, Pero-System GmbH, Wupertal, Germany) (Figure 1) (Stanton et al., 1997) and the sequential tape measure method, STM. With OEV, the exposed leg to be scanned is positioned on a platform with the person standing. Each leg was scanned from the ankle to the upper thigh in a single vertical sweep of the optoelectronic measuring frame (Stanton et al., 1997). A plastic cylinder with a geometric volume of 2833ml was scanned during each clinic to provide a standard.

The sequential tape measure method used a metric non-stretch tape measure (Medi Germany) to measure the leg circumference at 4cm intervals from the top of the malleolus up the leg to 40cm from the first mark. To assure consistent measurement a registered nurse with experience in circumferential leg measurements using a tape measure performed the measurements. The volume of each segment was calculated from the leg circumference measures (LymCalc V 4.0, UK), and combined to provide the overall leg volume (Williams and Whitaker, 2015). The recorded OEV scans, allowed the selection of the same leg length as used by STM method, ensuring that the same leg sections were compared.

Table 1 - Participant demographics at baseline.

	Control (n=10)	Intervention (n=10)	All (n=20)	P value
Mean age (years) (SD)	41.3 \pm 13.2	58.3 \pm 11.5	49.8 \pm 14.9	0.16*
Gender (M:F) (%)	50:50	10:90	30:70	0.14
Weight (KG) (SD)	92.9 \pm 29.2	95.5 \pm 43.8	94.2 \pm 36.2	0.88*
Height (CM) (SD)	169.6 \pm 8.5	162.9 \pm 10.4	166 \pm 9.8	0.16*
BMI (SD)	30.2 \pm 8.4	36.2 \pm 12.2	33.2 \pm 10.6	0.21*

Comparison of means \pm SD, differences between groups were made using a T-test or a Mann Whitney Rank sum test. A Chi-squared test was used to calculate ratio.



Figure 1. Perometer 400T, Pero-System GmbH, Wupertal, Germany.

Statistical Analysis

Comparison of means and correlation between groups was performed using computer software (IBM SPSS, version 22, New York, USA and Sigmaplot, Version 14, Sysstat, UK). Agreement plots were used to determine the agreement between the two methods (Bland and Altman, 1986), linear regression was used to test for correlation and ANOVA's to test for significance between groups. Statistical significance was set at $p < 0.05$.

RESULTS

In the twenty bipedal participants measured on three occasions, a total of 104 paired measures were collected, 16 measurements were not collected due to participant withdrawal from the study (Table 2). Measured leg volumes were allocated to one of three groups, normal legs (non-oedematous legs from patients with unilateral oedema, group A, $n = 24$), oedematous legs acting as controls not receiving the intervention, (from uni and bi lateral cases, untreated group B, $n=37$), and oedematous legs receiving the intervention (from unilateral and bilateral cases, treated group C, $n=43$). The model cylinder with a calculated volume of 2833 ml was scanned on ten occasions with the OEV, recording a mean volume of 2946 ± 10 ml (range: 2935 – 2960), with the OEV typically overestimating the volume by 113 ± 10 ml. A comparison of the agreement between the OEV and STM measures of leg volume in the three groups showed a difference of 437, 412 and 458 mls in groups A,B, and C respectively (Figure 2, Table 2). In all cases the OEV provided the largest measure; the mean difference of the combined data (Groups A, B and C) being 437 ml (Figure 2 and Table 2). This offset was consistent across all leg volumes, large and small, and irrespective of group. Despite this offset, the correlation between the two methods was strong, a fitted linear regression analysis produced a correlation coefficient, r , of 0.99 (Figure 3).

Table 2 - Comparison between methods.

Groups	Agreement between two measurements (LymCalc, Pero) mean offset (mls) \pm 1.96 SD	Median volumes Percentiles 25-75% shown (LymCalc, Pero)	Coefficient of variation (%) (LymCalc, Pero)
A) Lymphoedema free legs (n = 24)	-437 \pm 201	3363 (2862-4175) 3853 (3223-4624)	28, 26
B) Oedematous Legs Untreated (n= 37)	-412 \pm 259	3764 (3052-6426) 4054 (3417-7021)*	38, 36
C) Oedematous Legs Treated (n=43)	-458 \pm 239	3711 (3222-5604) 4217 (3772-6114)	38, 35
All data (n=104)	-437 \pm 239	3643 (3031-5372) 4084 (3454-5811)	39, 36

*Significantly different from Group 1, lymcalc measurement, see text (Figure 4).




Figure 2. Agreement plots between STM (LymCalc) and OEV (Perometer) measurements in A; unaffected legs with no lymphoedema, B; legs with lymphoedema but untreated with Lymphassist, and C; legs with lymphoedema and treated with the Lymphassist. Panel D shows all data. Note, for all plots baseline, 3 and 6 month data are shown together. See Table 1 for details.

The range of leg volumes for the three groups are shown in Figure 4. A Kruskal-Wallis one way ANOVA on ranks was performed (data not normally distributed). There was a difference between the medians of group A and B, between the OEV and STM measure (p 0.046). Otherwise there were no notable differences between techniques or three groups, despite Group B (oedematous legs, not treated), exhibiting the larger volumes and Group A (non-oedematous legs), the smallest, this observation is reflected in the coefficient of variation (Table 2).




Figure 3. Correlation between the two methods, all data shown, fitted with a linear regression line. The line of identity is shown.




Figure 4 - Box plots showing the distribution of data, the median and mean, for 3 groups, A, lymphoedema free legs, B, lymphoedema legs untreated, C legs with lymphoedema and treated. The subscript LC and PE denote Lymcalc (STM) and perometer (OEV) measures respectively.

DISCUSSION

The agreement comparisons (Figure 2) show that the OEV method overestimated lower limb volume by around 437mls compared to the STM, a difference of 9.5 %. The OEV also overestimated the cylindrical model volume by 113 ml, if this OEV based error is subtracted from the limb OEV measures; the agreement between the methods is closer, with a difference of 324 mls, or 7.2%. The correlation between the two measurement techniques is strong and linear, so for the two techniques to agree in this study then a constant, of 324 mls should be extracted from the OEV data, assuming the STM to be the reference standard.

Mayrovitz et al., (2000) found the tape measure method and perometer were closely correlated ($r=0.98$) in patients with both upper and lower lymphoedema. A total of 184 measurements were included in their study, however a horizontally orientated perometer was used. Overall, the volumes obtained by the tape measure and perometer methods differed by less than 5% for legs and less than 7% for arms. Stanton et al., (1997), found a 6.8% difference in oedematous arms. These results are similar in magnitude to the results in our study, hence showing that the perometer consistently overestimates limb volume compared to the tape measure method. This may be partly explained by the different formula used to calculate the leg volume. With STM the 4 cm limb segments are treated as truncated cones. Both Mayrovitz et al., (2000) and this feasibility study used the truncated cone formula ($\text{segment volume} = L/12\pi(C_1^2 + C_1 C_2 + C_2^2)$, where L is the segment length and C_1 and C_2 are the circumference at the end of each segment). The perometer uses a cylinder or disc formula, $V = \pi r^2 L$ (where r = segment radius). Tiereny et al., (1996) compared the horizontally orientated perometer with the tape

measure method but using the disc model formula and showed that the tape measure overestimated limb volume. The truncated cone formula calculates a relatively smaller volume in comparison to the cylinder formula. This indicates that a correction factor of 7% is required to compare the two techniques as the actual difference will depend on the total limb length scanned or measured.

Tan et al., (2013) found that the tape measure method overestimated lower limb volume in healthy volunteers compared to a vertically oriented perometer, with a mean volume difference of 157mls. The coefficients of variation were 0.14 and 0.13 for the tape measure and perometer method respectively. This has also been supported by Labs et al., (2000) suggesting the two methods need a different correction factor, however these studies used the cylinder formula to calculate the volume for the circumferential measurements.

An accurate measurement of limb volume is essential for lymphoedema patients in both clinical and research settings. Limb volume is an important marker of both disease progression and effectiveness of prescribed treatments. Although the OEV method is a quick and efficient way of obtaining leg volume and allows a digital image of the scanned limb, financial considerations may prohibit its use as a typical perometer costs in the region of £16,731. The tape measure circumferential method in the hands of a skilled person is also simple, and requires little equipment, but only provides an answer after a series of calculations. This study shows that a perometer is not necessarily needed as the tape measure produces reliable results and enables whole legs to be measured when patients have a limited range of motion.

The limitations of this study were that leg volume measurements were performed only once at each assessment period. Participants were lymphoedema patients so it cannot be assumed that the same results would be found in healthy participants (Burton et al., 2012).

CONCLUSION

Results have demonstrated that OEV and STM provide comparable measures of lower limb volume in people with lymphoedema, with a 7% difference between the methods; this corroborates the existing base relating to this subject. The advantage of the OEV method is that it is fast, non-operator dependent and automatically provides a limb volume. The disadvantage is that it is not a portable system and cost issues may prohibit its use in many settings. The tape circumferential method is easy to use by a trained person but can be time consuming and requires manual calculation of leg volume. Further studies on leg volume using these techniques should focus in creating predicted leg volumes, allowing swollen limbs to be compared with non-oedematous legs.

Acknowledgements

The study team would like to thank Cwm Taf University Health Board for allowing access to their patients and the staff of the lymphoedema clinic at Dewi Sant Hospital, Pontypridd for their assistance with data collection. Sincere thanks are also extended to the patients that took part in the study.

Ethical Approval: Wales 6 (LREC No 17/WA/0076)

Funding: ND is supported by KESS which includes a contribution

by Huntleigh Ltd (Cardiff, UK). Knowledge Economy Skills Scholarships (KESS) is a pan- Wales higher level skills initiative led by Bangor University on behalf of the HE sector in Wales. It is part funded by the Welsh Government's European Social Fund (ESF) convergence programme for West Wales and the Valleys.

Conflict of Interest: None declared

REFERENCES

1. Bland J.M., Altman D.: *Statistical methods for assessing agreement between two methods of clinical measurement*. The Lancet, 1986; 8, 327(8476): 307-10.
2. Burton R.F., Nevill A.M., Stewart A.D., Daniell N., Olds T.: *A negative relationship between leg length and leg cross-sectional areas in adults*. American Journal of Human Biology, 2012; 24(4): 562-564.
3. Deltombe T., Jamart J., Recloux S., Legrand C., Vandenbroeck N., Theys S., Hanson P.: *Reliability and limits of agreement of circumferential, water displacement and optoelectric volumetry in the measurement of upper limb lymphedema*. Lymphology, 2007; 40: 26-34.
4. Kaulesar Sukul. D.M., den Hoed P.T., Johannes E.J., van Dolder R., Benda E.: *Direct and indirect methods for the quantification of leg volume: comparison between water displacement volumetry, the disk model method and the frustum sign model method, using the correlation coefficient and the limits of agreement*. J Biomed Eng, 1993; 15: 477-80.
5. Labs K.H., Tschoepf M., Gamba G., Aschwanden M., Jaeger K.A.: *The reliability of leg circumference assessment: a comparison of spring tape measurements and optoelectronic volumetry*. Vascular Medicine, 2000; 5(2): 69-74.
6. Lasinski B.B.: *Complete decongestive therapy for treatment of lymphedema*. Seminars in oncology nursing, 2013; 29(1): 20-7.
7. Lymphoedema Framework: *Best Practice for the management of lymphoedema*. London, 2006, MEP Ltd. 2.
8. Mayrovitz H., Sims N., Macdonald J.: *Limb volume measurement in patients with edema*. Adv Skin Wound Care, 2000; 113(272): 6.
9. Stanton A.W.B., Northfield J.W., Holroyd B., Mortimer P.S., Levick J.R.: *Validation of an optoelectronic limb volumeter (Perometer®)*. Lymphology, 1997; 30(2): 77-97.
10. Tan C.W., Coutts F., Bulley C.: *Measurement of lower limb volume: agreement between the vertically oriented perometer and a tape measure method*. Physiotherapy, 2013; 99(3): 247-251.
11. Tewari N., Gill P.G., Bochner M.A., Kollias J.: *Comparison of volume displacement versus circumferential arm measurements for lymphoedema: implications for the SNAC trial*. ANZ journal of surgery, 2008; 78(10): 889-893.
12. Tierney S., Aslam M., Rennie K., Grace P.: *Infrared optoelectronic volumetry, the ideal way to measure limb volume*. European Journal of Vascular and Endovascular Surgery, 1996; 12(4): 412-417.
13. Williams A.F., Whitaker J.: *Measuring change in limb volume to evaluate lymphoedema treatment outcome*. EWMA Journal, 2015; 15(1).
14. Zaleska M., Olszewski W.L., Durlik M.: *The effectiveness of intermittent pneumatic compression in long-term therapy of lymphedema of lower limbs*. Lymphatic research and biology, 2014; 12(2): 103-109.

THE PATIENT WITH PRIMARY AND SECONDARY LOWER EXTREMITY LYMPHEDEMA; A 1-YEAR FOLLOW-UP STUDY

PT. CEMILE UYDUR¹, PT. MSC.GULBALA NAKIP²

¹Avrupa Safak Hospital, Physical Therapy and Rehabilitation Clinic, Istanbul

²Hacettepe University, Faculty of Health Sciences, Department of Physiotherapy and Rehabilitation

Corresponding author: Cemile UYDUR , Avrupa Safak Hospital,
Physical Therapy and Rehabilitation Clinic, Gaziosmanpasa, Istanbul
Fax number: +90 212 614 86 77

ABSTRACT

Purpose: The aim of this study was to investigate the 1-year results of the complex decongestive therapy (CDT) in patient with stage 3 primary and secondary lymphedema in the lower extremities.

Methods: A 24-year-old female patient (body mass index-BMI 83 kg/m²). Her height of 1.68 cm and weight of 255 kg. CDT program was administered for 8 weeks, 7 days a week. As a home program, compression garment for day time and compression bandage at night was recommended.

Conclusion: Regularly repeated CDT in patients with severe lymphedema is important for softening of fibrotic tissue, improvement of skin folds, and reduction of extremity volume.

Key words: Lymphedema, Liposuction, CDT, Complex Decongestive Therapy, Edema

INTRODUCTION

Lymphedema, described as abnormal accumulation of protein rich fluid, due to lymphatic dysfunction. It can be present in extremities, the head and neck, trunk, abdomen, external genitalia, and inner organs. The most common reason for lymphedema in the U.S. are surgical interventions in combination with dissection of lymph nodes due to breast cancer (12.5% in the US). Approximately 40% of patients who underwent treatment for breast cancer develop some degree of upper extremity lymphedema. LE can be categorized as congenital LE or secondary LE. Congenital LE is caused by abnormal lymphatic system formation due to a genetic mutation. Meanwhile, secondary LE is caused by damage to the lymphatic system due to infection, damage, cancer, lymph node incision, or radiotherapy. Lymphedema, in most cases, will gradually progress through stages – see Table.

Stages	Phase I (decongestion)	Phase II (self-maintenance)
Latency	Patient Instruction	
Stage I	MLD daily short-stretch bandages skin care decongestive exercises patient instruction	MLD if necessary compression garments skin care decongestive exercises
Stage II	MLD daily short-stretch bandages skin care decongestive exercises patient instruction	MLD compression garments bandages at night skin care decongestive exercises repeat Phase I
Stage III	MLD daily short-stretch bandages skin care decongestive exercises patient instruction	compression garments (in combination w/ bandages) bandages at night skin care decongestive exercises repeat Phase I if indicated: reduction surgery

Therefore, we report the case of patient with primary and secondary lower extremity lymphedema.

CASE REPORT

We report a 24 year old female was consulted to our department of physical and rehabilitation medicine for the bilateral lower extremity swelling and difficulty in walking. The patient 's parents report a sudden onset of swelling in her left lower extremity when

she was 3 years old. She was diagnosed with lymphedema at that time. She has never been any treatment before. She had morbid obesity (body mass index-BMI 83 kg/m²). Her height of 1.68 cm and weight of 255 kg. When she was 14 year old, developed lymphedema in the right lower extremity resulting from conduction excision. Her family's past medical history was unremarkable. In her physical examination she had skin folds under the knee and fibrotic tissues in the lower dorsum and under the knee. Her diffuse edema was non pitting. Stemmer was positive. The skin was dry and intact. However but there was no evidence of any infection. There was no pain but she had difficulty walking alone. The patient had liposuction and debulking operation 6 years ago. We started to Complex Decongestive Therapy (CDT) program which include manuel lymphatic drainage, skin care, multilayered bandaging (with short tension) and exercisses techniques, were applied daily for a duration of eight weeks (Figure 1). The circumferences of the right and left lower extremity were measured before and after treatment. During this time she was taught self manual lymph drainage, skin care, and breathing and leg exercises.



Figure 1. Multilayered bandaging (with short tension).

The patient responded very well to her MLD/CDP. After the treatment, both lower extremity edema was reduced (Figure 2, 3). The compression garment was planned as special measurement panty, class 3 (Figure 4). BMI was also reduced 57.1 kg/m². As a home program, compression garment for day time and compression bandage at night was recommended. After 6 months, BMI of the patient who applied to our clinic for compression garment control was measured as 29.4 kg /m². CDT was performed again for 4 weeks, 6 days a week.



Figure 2. a) Before CDT (anterior), b) After CDT (anterior).



Figure 3. a) Before CDT, b) After CDT (Right leg).



Figure 4. a) Before CDT, b) After CDT, c) special measurement panty compression garments.

RESULTS

After Complex Decongestive Therapy (CDT), both lower extremity volumes decreased significantly (table 1). Fibrotic tissues and skin folds of extremities were improved positively (Figure 5). Body Mass Index (BMI) of 83.2 kg/m² to 29.4 kg/m² and her weight of 255 kg to 72 kg decreased. By the time she was discharged, the patient had new, well fitted compression garments. While the patient was discharged, she stated that she now had hope and that her daily life activities (ADL) were more independent.



Figure 5. a) Before CDT, b) After CDT, c) compression garments.

Table 1 - Before and 6 months after right and left leg circumference measurement (cm).

Left lower extremity (cm)	Before treatment	After treatment
Malleol (0)	36	34
+10	53	37
+20	60,7	42
+30	63,4	47
+40	88	54
Left lower extremity (cm)	Before treatment	After treatment
Malleol (0)	35	31
+10	55	35
+20	67	41
+30	74	50
+40	94	56

DISCUSSION

Complex decongestive therapy (CDT) which has individual components of skin care, manual lymph drainage, compression bandages, exercises and compression garments, is the golden standard treatment for reducing both primary and secondary

lymphedema. In this case we aimed to indicate the long term effects of manual lymphatic drainage and multilayer bandaging in the both lower extremities. Regularly repeated CDT in patients with severe lymphedema is important for softening of fibrotic tissue, improvement of skin folds, and reduction of extremity volume.

REFERENCES

1. Consensus Document of the International Society of Lymphology, *The diagnosis and treatment of peripheral Lymphedema*, Volume 21, No. 59, 2010.
2. Maclellan R.A., Zurakowski D., Grant F.D., Greene A.K.: *Massive Localized Lymphedema: A Case-Control Study*. J Am Coll Surg., 2017 Feb; 224(2): 212-216.
3. Kurt H., Arnold C.A., Payne J.E., Miller M.J., Skoracki R.J., Iwenofu O.H.: *Massive localized lymphedema: a clinicopathologic study of 46 patients with an enrichment for multiplicity*. Mod Pathol., 2016 Jan; 29(1): 75-82.
4. Greene A.K., Voss S.D., Maclellan R.A.: *Liposuction for Swelling in Patients with Lymphedema*. N Engl J Med., 2017 Nov 2; 377(18): 1788-1789.
5. Boyages J., Kastanias K., Koelmeyer L.A., Winch C.J., Lam T.C., Sherman K.A., Munnoch D.A., Brorson H., Ngo Q.D., Heydon-White A., Magnussen J.S., Mackie H.: *Liposuction for Advanced Lymphedema: A Multidisciplinary Approach for Complete Reduction of Arm and Leg Swelling*. Ann Surg Oncol. 2015 Dec; 22 Suppl 3: S1263-70.
6. Brorson H., Svensson H.: *Liposuction combined with controlled compression therapy reduces arm lymphedema more effectively than controlled compression therapy alone*. Plast Reconstr Surg. 1998 Sep; 102(4): 1058-67.

VALUES AND LIMITS OF SURGERY IN PRIMARY AND SECONDARY LYMPHEDEMA

MICHELINI S.*; FIORENTINO A.*; FANTEGROSSI M.R.*; FAILLA A.*; MONETA G.*; CARDONE M.*; GOBBETTI G.**

* San Giovanni Battista Hospital - ACISMOM - Rome - (Italy).

** Villa Regina Clinic' - Arco di Trento - Italy

SUMMARY

Surgical treatment in Lymphedema is an approach proposed in synergy with the physical treatment that still requires further investigation regarding the actual indications. 95 patients with lymphedema were studied (23 males and 72 females), aged between 19 and 74 years, 31 primary and 64 secondary, 35 to lower limbs, 60 to upper limbs with average period between clinical onset of the disease and surgery for 3.8 years (0.3 to 11 years). The observation was performed on a minimum follow-up of 1 year from the intervention.

The observed results suggest that the indications for intervention must be further investigated in order to avoid (taking into account same technique, operator and clinical status) the high incidence of non-success compared to the persistent clinical improvement (though maintained also with the continuation of the physical treatment and of the permanent elastic garment); the observation is in line with what is recommended by the ISL Consensus Document and other guidelines and partly justifies the decision of some states not to recognize the surgical treatment itself as effective.

Key words: Primary and secondary lymphoedema. Surgical treatment.

INTRODUCTION

The surgical treatment in the primary and secondary lymphedema is becoming more frequent in its various techniques. From the demolitive surgery, always less employed, to derivative or reconstructive microchirurgia, to supermicrosurgery to lymphatic transplantation to liposuction, there are many therapeutic proposals that the medical and scientific world present. Professor BB Lee, introducing this field in the Consensus Document of International Union of Phlebology, remember us: "A new approach to restore normal lymphatic flow with direct surgical correction of disabled lymph transport has been a dream among lymphatic/vascular surgeons since Tosatti, Donini, Olszewski et al. made a landmark report on direct anastomosis between lymphatic and venous system using microscopic surgical technique a half century ago. This breakthrough gave new hope to the management of chronic lymphedema."

This new approach became popular throughout the past century because of its potential to restore normal lymphatic flow when performed properly, providing a chance of a "cure" in theory. However, over the past several decades, an often cavalier approach with limited knowledge of lymphatic anatomy and physiology, adopted by many surgeons, resulted in disastrous outcomes and added more confusion with erroneous prejudice. Many surgeons condemned the procedure due to the poor outcomes, which were believed to be a result of the surgery itself rather than an ill-planned treatment strategy and lack of proper knowledge regarding lymphatic anatomy and physiology. I (B.B.L.) was no exception, being among the first group of surgeons who began performing lympho-venous anastomosis (n = 15, patients operated) in early 1980s. Interim follow-up results were embarrassingly poor despite technical success in performing flawless surgery without a clear explanation. It took almost a decade to figure out why and what went wrong to make ourselves better prepared before resuming lymphatic reconstructive surgery" ⁽¹⁾. The consensus document of the International Society of Lymphology underlines the importance of the methods and recommends that these are being performed in specialized centers and asks even more evidence of results in long time ⁽²⁾.

MATERIALS AND METHODS

The study includes ninety eight subjects who underwent surgery after surgical treatment 31 affected from primary lymphoedema and 64 from secondary one). 35 of lower limbs, sixty of upper limbs. Follow up was performed at least one year after surgical treatment.

They had been considered the following aspects:

- Modifications of Volume of treated limb,
- Modifications of tissue consistency,
- Incidence of phlogistic complications,
- Patients expectations.

All patients undergone combined physical treatment after surgery and had worn the prescribed garment. In this regard, must underline the tendency to better respect the garment after surgery respect to after only physical treatment.

RESULTS

The data emerged regarding the various types of intervention undergone demonstrated:

- The most frequently controlled type of intervention is represented by Supermicrosurgery (47/98, 47,9%). Less often: Microsurgery (38 cases of which 34 ‘derivative microsurgery’ and 4 ‘reconstructive microsurgery’, 38,7%), Liposuction (11 cases 11,2%) and Lymphatic transplantation (5 cases 5,1%) (Tab.1) ^(3,4,5,6).
- In relation to the reduction of volume, Secondary shapes demonstrated best results, compared to the primary (Fig. 1).
- Similar results we have observed about the tissue consistency (Fig. 2).
- Best results, absolute, for both the forms, primary and secondary, have been observed regarding the decrease of the episodes of lymphangitis (Fig. 3).
- Results of the decongestive physical treatment after surgical therapy seem to be encouraging, especially in secondary shapes. analogy to other therapies with the passing of time the same decreases (Fig. 4).
- The patient's judgements on clinical results after surgical treatment, on average, did not reflect adequate satisfaction with expectations. This is highly listened by patients with primary lymphedema when you ask them for hundred percention of expectations as soon as they have obtained by surgical intervention (Fig. 5).

	Reconstructive Microsurgery	Derivative Microsurgery	Lymph node transplantation	Supermicrosurgery	Liposuction
Primary OK		2		3	1
Primary indifferent	2	8	1	7	1
Primary KO		3	1	2	1
Secondary OK		8	1	11	5
Secondary indifferent	1	6	1	10	1
Secondary KO	1	7	1	10	1

OK: persistent volume and consistency reduction - lymphangitis regression
Indifferent No changes in volume and consistency or incidence of lymphangitis
KO Increase in volume and / or consistency - lymphangitic complications




Fig. 1 - Medium Volume decrease in primary and secondary lymphoedema.




Fig. 2 - Medium tissue consistency decrease.




Fig. 3 - Decrease of lymphangitis episodes.




Fig. 4 - Results of physical treatment after surgery.




Fig. 5 - Expressed expectations of patient by the surgical treatment.

DISCUSSION AND CONCLUSIONS

In this preliminary study (that's still going on) the first observation is that by surgical treatment, on the contrary with other kind of surgery, there is not the surely clinical result, at least according with the medium expectations of patients. Best results are obtained in Lymphangitis decrease, as in a paradigmatic case (Fig. 6).

53 years patient, with secondary lymphedema of the upper limb starting from over 9 years ago, with considerable increase of tissue consistency and up to nine lymphangitis per year, despite antibiotic therapies. At follow up, after three years from the surgical treatment (supramicrosurgery), the patient demonstrated excellent results below all points of view, but in particular in the total disappearance of lymphangitis^(7,8). But, some other considerations must do. In some cases the indication to surgery is a mistake, incorrect. How in a case of inguinal lymphadenitis (Fig. 7) in which the presurgical lymphoscintigraphy and clinic before the surgical treatment of lymphatic venous derivative microsurgery was 'normal'; after the intervention the lymphoscintigraphy showed important dermal back flow and disappearance of left inguinal lymphoneds, corresponding to an important increase of the edema, especially of left thigh. In other case of Klippel Trenaunay Syndrome two lymphoscintigraphies, carried out in an incorrect way, had made the diagnosis of hypogenesis of the inguinal left lymphonods^(3,9,10,11). A third exam, carried out with the 'Bourgeois method' with further injection of the tracer at the third lower of thigh, showed the presence of inguinal left lymphonods and, according with ultrasound exam, the prevalent role of venous insufficiency in the etiopathogenesis of vascular malformation. A possible surgical treatment would have made worse the clinical picture (Fig. 8). In a last case of Lipedema of the lower limbs a lymphatic derivative microsurgery intervention has been performed in the right lower limb and it transformed the lipedema in lipolymphoedema of the right lower limb, with the pre-operative lymphoscintigraphy absolutely normal. In this case the indication to the surgical treatment is absolutely a mistake (Fig. 9). The observed results suggest that the indications for intervention must be further investigated in order to



Fig. 6 - Large arm three years follow up after supramicrosurgery.




Fig. 7a,b,c,d - Lymphoscintigraphy and clinical aspect before and after derivative microsurgical treatment.




Fig. 8 - No Lymphatic surgery indication.




Fig. 9 - Bad surgical indication (Lipedema of lower limbs): preoperative lymphoscintigraphy showed normal transport capacity of lymphatic system of both lower limbs. After microsurgery evidence of Lipolymphoedema in right lower limb.

avoid (with the same technique, the same operator and the same clinical picture) the high incidence of non-success compared to the persistent clinical improvement (though maintained also with the continuation of the physical treatment and of the permanent elastic garment)^(12,13,14); The observation is in line with what is recommended by the consensus document of International Society of Lymphology and other guidelines and partly justifies the decision of some countries of not recognize the surgical treatment itself as an average effective (ie: Belgium who removed the intervention from the National Sanitary System of derivative and / or reconstructive surgery from November 2017). On the basis of these observations can conclude that in the lymphedema the clinical variables are many and not always the indications to a types of therapy or to the other are discounted regarding the clinical results. Probably, it will be useful to study together (surgeons, clinicians and radiologists) which are the factors which, according to the clinical picture, surgical treatment, operator and general conditions of the patient and adhesion to the entire proposed protocol, positively or negatively influences the final outcome of the treatment with the aim to highly select the candidates to safe positive result.

REFERENCES

1. Lee B. B., Andrade M., Antignani P. L., Boccardo F., Bunke N., Campisi C., Damstra R., Flour M., Forner-Cordero J., Gloviczki P., Laredo J., Partsch H., Piller N., Michelini S., Mortimer P., Rabe E., Rockson S., Scuderi A., Szolnoky G., Villavicencio J. L.: *Diagnosis and Treatment of Primary Lymphedema*. Consensus Document of the International Union of Phlebology (IUP)-2013. International Angiology, 2013 December; 32(6): 541-74
2. *The diagnosis and treatment of peripheral lymphedema: 2016 consensus document of the international society of lymphology*. Lymphology 49 (2016): 170-184.
3. Roman M.M., Barbieux R., Nogaret J.M., Bourgeois P.: *Use of lymphoscintigraphy to differentiate primary versus secondary lower extremity lymphedema after surgical lymphadenectomy: A retrospective analysis*. World Journal of Surgical Oncology 16 (1): 75. 2018.

4. Boccardo F., Bellini C., Campisi C., Eretta C., Pertile D., Benatti E., Campisi M., Talamo G., Macciò A., Campisi C.: *Role of Lymphoscintigraphy in the indications to microsurgical treatment of peripheral lymphedema*. The European Journal of Lymphology, 2006; Vol. 16, N. 49: 24-28.
5. Campisi C., Jiménez Cossio J.A., Pissas A., Leduc A., Michelini S., Boccardo F., Zilli A.: *Prevention of Secondary Lymphedema: Prospects for the Future*. Lymphology, 1998; 31 (Suppl): 513-515.
6. Michelini S., Failla A., Paroni Sterbini G.L., Micci A., Santoro A., Valle G.: *Limb phlebolymphedema: diagnostic non invasive approach and therapeutical implications*. The European of Lymphology, 1995; Vol. 5, N. 20: 103-8.
7. Wald M., Jarolim L., Adámek J., Hyklova L., Vachová H.: *The surgical management of chronic penoscrotal lymphedema*. The European Journal of Lymphology, 2009; Vol. 20, N. 56: 19-22.
8. Foldi M., Foldi E., Kubik S.: *Textbook of lymphology*. Elsevier, Munich, 2005.
9. Cormier J.N., Rourke L., Crosby M., Chang D., Armer J.: *The surgical treatment of lymphoedema: a systematic review of the contemporary Literature (2004-2010)*. Annals of Surgical Oncology, 2012; Vol. 12, 2: 642-651.
10. Brorson H., Svensson H.: *Liposuction combined with controlled compression therapy reduces arm lymphedema more effectively than controlled compression therapy alone*. Plast Reconstr Surg, 1998; 102: 1058-1067.
11. Granzow J.W., Soderberg J.M., Dauphine C.: *A novel two-stage surgical approach to treat chronic lymphedema*. Breast J, 2014; 20: 420-422.
12. Granzow J.W., Soderberg J.M., Kaji A.H., Dauphine C.: *Review of current surgical treatments for lymphedema*. Ann Surg Oncol, 2014; 21: 1195-1201.
13. Granzow J.W., Soderberg J.M., Kaji A.H., Dauphine C.: *An effective system of surgical treatment of lymphedema*. Ann Surg Oncol, 2014; 21: 1189-1194.
14. Brorson H., Svensson H., Norrgren K., Thorsson O.: *Liposuction reduces arm lymphedema without significantly altering the already impaired lymph transport*. Lymphology, 1998; 31: 156-172.

THE EFFECT OF LOWER LIMB LYMPHEDEMA ON POSTURAL STABILIZATION

ALIS KOSTANOGLU¹, MELTEM RAMOGLU¹, HIKMET UCGUN¹

Affiliations:

¹Division of Physiotherapy and Rehabilitation, Faculty of Health Sciences,
Bezmiâlem Vakif University, Turkey

Correspondence: A Kostanoglu

Division of Physiotherapy and Rehabilitation,
Faculty of Health Sciences,
Bezmiâlem Vakif University,
Turkey

Gsm: 90 543 493 18 48

Fax: 90 212 453 18 70

E-mail: aliskostanoglu@yahoo.com

ABSTRACT

Objectives: Lymphedema is a chronic and serious condition that causes asymmetric weight dispersion between extremities. Patients commonly report symptoms such as decreased mobility of the limb, pain, tissue fibrosis and associated skin changes due to the swelling. These symptoms may involve postural disorders that affect the whole body. The aim of this study was to investigate whether there is a change in overall postural stability and limits of stability in patients with unilateral lower limb lymphedema.

Methods: Fifteen (10 female, 5 male) patients (43.2±13.08 years) diagnosed with lower limb lymphedema were included in the study. The mean volume difference between the two extremities was 548±492 ml. Lymphedema severity was evaluated and classified as mild, moderate and severe for 2, 7 and 6 patients respectively. Fifteen healthy people with similar ages (43.33±8.29) were recruited for the control group. All the participants were assessed by using the Biodex Balance System. Postural stability, which is the ability of the patient to maintain a static postural balance, and limit of stability, which is the ability of directional control of the body by displacing the center of gravity, were assessed. All evaluations were carried out on a firm and stable platform with open eyes. The data were analyzed with the SPSS 21.0 program. Mann-Whitney U Test was used for intergroup comparisons.

Results: There was no significant difference between the patients and the control group in overall, anterior-posterior and mediolateral postural stability indexes ($p < 0.05$). However, the lymphedema group had worse overall, forward and backward limits of stability scores in comparison to the control group (42.33±12.5 vs 60.29±14.1; 50.73±18.6 vs 73.93±16.08; 49.60±22.8 vs 67.79±21.2 respectively) ($p < 0.05$).

Conclusion: These results lead us to think that asymmetric fluid distribution in the lower body parts deteriorate directional control

in patients with unilateral lymphedema in comparison to the healthy group. Further studies are needed to confirm the changes of postural stability and limits of stability in these patients.

Abbreviations: BW = body weight, CoG = center of gravity, LLL = lower limb lymphedema, LoS = limits of stability, PoS = postural stability

Keywords: Unilateral Lower Limb Lymphedema, Biodex Balance System, Postural Stability,

1. INTRODUCTION

Lower limb lymphedema (LLL) is a chronic condition with no cure. The impact is not only limited to the related region, but it can also negatively affect other systems in a multidirectional way. Balance is a skill of total body performance, which depends on somatosensory, vestibular and visual systems, on the motor output and a central processing system¹. Any deficit of these systems could be responsible for poor balance control. Maintaining upright stance in humans is a complex task as it provides postural stability. Postural stability may deteriorate due to any change in the center of gravity (CoG), and the asymmetric distribution of body weight (BW) is one of the causes². Asymmetric distribution of BW such as amputation or unilateral volume change, plays a specific role in disruption of the postural stability in the lower body. Postural sway may be defined as all deviation in the location of the CoG.³. It is assumed that asymmetric BW distribution increases postural sway and therefore, it is important to emphasize that postural asymmetry is characterized by postural disturbances and balance problems. In other words, such balance problems are expressed as an individuals' stability limitation problem and may be evaluated with limits of stability (LoS) tests, one of the related evaluation methods⁴. Review of studies in the literature provided inadequate studies investigating the effects of unilateral volume increase on postural stability in lower limb lymphedema. A similar study,

which described the effects of unilateral upper extremity lymphedema on postural stability, was found³. Most studies in the literature focused on lymphedema severity, volume reduction and subjective sensations such as pain, heaviness or tension. Problems such as balance and postural stability disorders should be a part of general assessment. Studies have extensively assessed upper limb lymphedema^{4,5}. In this study, we aimed to assess the effects of unilateral LLL on postural stabilization.

2. MATERIALS AND METHODS

2.1. Sample

Fifteen patients aged 18-63 years old who were clinically diagnosed with unilateral LLL and fifteen healthy people were included in the study. A volume difference of the lower limb of more than 200 ml compared to the contralateral limb was defined as lymphedema.

The lymphedema severity levels of the patients were evaluated and classified as Stage 1 (mild), Stage 2 (moderate) and Stage 3 (severe). The clinical stages of lymphedema of our cases were classified based on the International Society of Lymphology⁶. All participants were informed of the study, and written consent was obtained. Based on the Declaration of Helsinki, the Ethics Committee approved this study, which was performed in accordance with the ethical principles for human research. The study was designed as a prospective and single-center trial. All tests were applied by the same physiotherapist. Static postural stability and limits of stability were evaluated by the Biodex Balance System.

The inclusion criteria were:

- Diagnosis of unilateral lower limb lymphedema
- Minimum of 200 ml volume difference between lower limbs
- Conserved lymphedema with a duration of at least 6 months

The exclusion criteria were:

- Diagnosis of bilateral lower limb lymphedema
- Having vestibular or visual problems
- Having a previous diagnosis of neurological disorders or severe arthritis
- A history of spinal surgery

2.2. Biodex balance system

The Biodex Balance System (Biodex Medical Systems, Shirley, New York) is a testing machine which has a multiaxial standing platform to evaluate situations relevant to balance, proprioception and neuromuscular control⁷. To achieve a static situation, a proper stable platform surface is selected. The patients' ability of maintaining static postural balance on this stable platform was assessed.

(1) anterior/posterior index, (2) medial/lateral index and (3) overall index are electronically generated. The overall index is assumed to be the best indicator of the general ability of the patient's balance. The total duration of evaluation of the test takes about 10 minutes on average.

2.2.1. The postural stability test

Deviations from the center create the test score of the patient, thus a lower score is more creditable than a higher score. The patients stood for 20 seconds on both legs in the most stable position while the test was being performed. After the patient had accommodated to the platform, 3 trial repetitions were performed. The patients' performance was noted as a stability index. It was made in 3 directions (overall, anterior-posterior and mediolateral) and 1 condition with open eyes. The stability index represents the change in grade from the initial level of platform displacement. A high score means the individual had difficulty to maintain their balance.

2.2.2. The limits of stability test

This test assesses the movement and control of patients' CoG within their base of support. In the test, patients must switch their weight to move the cursor from the center to a blinking target. Return to the starting position should be in a short time and with as little deviation as possible. For all targets, the same process is valid. Targets blink randomly.

Scores are based on the patient's time to complete the test and the effort to reach the goal. Low scores indicate that the patient needs more effort and time to complete the test.

2.3. Statistical analyses

The data were analyzed using the "Statistical Package for the Social Sciences (SPSS) Version 20.0" (SPSS Inc., Chicago). Sample size calculation was performed by using G*Power version 3.1.9.2 for Windows (Universitat Kiel, Germany). The survey and test results of the individuals were analyzed by calculating frequencies, minimum and maximum values, and 95% confidence intervals. Data distribution was assessed with Shapiro-Wilk test. Independent-Samples t-Test was used for intergroup comparisons. The results were considered significant with P values <0.05.

3. RESULTS

Fifteen patients (10 female, 5 male) diagnosed with lower limb lymphedema were included in the study. The mean duration of lymphedema was 74 months [7-360] ([min-max]). The mean volume difference between the two extremities was 548±492 ml. Lymphedema severity was evaluated and classified as mild, moderate and severe for 2, 7 and 6 patients respectively. Fifteen healthy people (12 female, 3 male) with similar ages were recruited for the control group. The subjects' characteristics are shown in Table 1. There were no significant differences in the demographic data between the groups ($p > 0.05$).

The results of postural stability tests and limits of stability tests in a static and eyes open condition for the lymphedema group are presented in Table 2. There was no significant difference between the lymphedema group and the control group in overall, anterior-posterior and mediolateral postural stability indices ($p > 0.05$). However, the lymphedema group had worse overall, forward, backward, forward/left, forward/right, backward/left and backward/right limits of stability scores in comparison to the

Table 1. Demographic characteristics of the lymphedema and control group.

Demographic characteristics	Lymphedema Group (n = 15)	Control Group (n = 15)	p
	X ± SD	X ± SD	
Age (years)	43.27 ± 13.08	42.00 ± 9.3	0.51
Body weight (kg)	86.22 ± 13.31	79.35 ± 10.4	0.13
Body height (cm)	167.22 ± 9.31	169.17 ± 5.67	0.24
BMI (kg/m ²)	30.04 ± 4.57	25.13 ± 4.82	0.07

Table 2. Results of postural stability and limits of stability tests for lymphedema group.

Tests Indexes	Lymphedema Group (n = 15), X ± SD	Control Group (n = 15), X ± SD	p
Overall PoS index	0.96 ± 1.27	0.46 ± 0.25	0.14
Anterior-posterior PoS index	0.78 ± 1.28	0.40 ± 0.19	0.27
Mediolateral PoS index	0.26 ± 0.29	0.21 ± 0.16	0.54
Overall LoS index	42.33 ± 12.57	61.06 ± 14.00	0.001
Forward LoS index	50.73 ± 18.65	73.53 ± 15.57	0.001
Backward LoS index	49.60 ± 22.82	69.46 ± 21.50	0.001
Left LoS index	57.53 ± 17.92	65.66 ± 11.19	0.14
Right LoS index	59.80 ± 18.25	62.86 ± 20.72	0.67
Forward/Left LoS index	48.00 ± 12.77	69.60 ± 13.46	0.000
Forward/Right LoS index	49.86 ± 15.24	71.86 ± 16.07	0.001
Backward/Left LoS index	42.73 ± 16.95	66.06 ± 17.05	0.001
Backward/Right LoS index	50.66 ± 16.44	69.93 ± 12.22	0.001
Abbreviation: PoS – Postural Stability; LoS – Limits of Stability; X – Mean; SD - Standard Deviation (p< 0.05)			

control group (p<0.05). In addition to all these significant differences, the changes in the LoS scores for the right and left sides were not statistically significant (p>0.05).

4. DISCUSSION

The main objective of this study was to evaluate postural stabilization changes in unilateral LLL patients in comparison to healthy individuals. Our results indicated that, while unilateral LLL affected the overall, forward and backward limits of stability indices negatively, it did not affect CoG and the overall, anterior-posterior and mediolateral postural stability indices. Preservation of postural stability is related to control over the center of gravity and resulting momentum⁸. Momentum may be explained as a sway that is actively trying to stay in the best position. Stability while standing upright may be affected by weight asymmetry, unilateral volume change and weight bearing⁹.

Although lymphedema is a chronic and widespread problem that negatively impacts patients' functional status^{10,11}, there was not enough evidence about the changes in postural stability, postural sway and directional control in patients with LLL. In this study, the results demonstrated that there was no significant displacement of CoG when the lymphedema group and healthy individuals were compared.

Similarly to the findings in this study for asymmetrical weight dispersion in the literature, Bakar et al. found no significant differences overall for anterior-posterior and mediolateral postural stability indices with eyes opened in a breast cancer-related lymphedema group in comparison to healthy individuals, but we found different results in our study showing a significant difference between the eyes closed test and a foam surface¹². Visual and specific sensory inputs are important to compensate balance impairment. Standing with eyes closed or standing on different surfaces may make it difficult to maintain an upright

stance posture. Nadollek et al. reported that increased CoG sway with and without visual input affected patients with lower leg amputation¹³. Arifin et al. showed significant postural instability of amputees in an eyes-closed condition, followed by standing on a foam surface¹⁴. In contrast to these studies of unilateral weight bearing, similar changes were observed in CoG sway for patients with chronic subdural hematoma and unilateral lower limb amputees for the open and closed-eyes tasks^{15,16}. We think that there are a few factors that may be effective on maintaining postural stability in unilateral LLL. One of these factors is that our patients were very young. The other one is that there were no abnormalities in the patients' visual, vestibular and somatosensory systems. Additionally, we think patients with chronic lymphedema may have developed adaptive responses to maintaining postural stability.

Limits of stability tests assess dynamic balance control during voluntary movement in daily activities such as climbing stairs, getting up or walking. Dynamic balance failures that increase postural sway occur as a result of the absence of motor and proprioceptive senses, asymmetrical body weight distribution and posture. It is known that individuals with increased postural sway have less ability to maintain balance control¹⁷. On the other hand, some studies showed that high BMI may impair dynamic postural control and be a risk factor of balance loss^{18,19}. According to the results of the evaluation, the lymphedema patients had a higher body mass index than the healthy individuals. All these findings confirm the decline of dynamic balance control in unilateral LLL patients because of weight gain and asymmetric mass distribution. In the literature, we could not find any study which has been done on static and dynamic balance for unilateral lower limb lymphedema patients. Nevertheless, similar results were found in previous studies that focused on postural stability and postural sway in unilateral upper extremity lymphedema in breast cancer survivors. Angin et al. found that, in the case of positional changes, the dynamic balance was distorted due to increased volume towards the lymphedema and a heavier upper extremity³. Similarly, Basar et al.¹² assessed the postural stability between lymphedema and healthy groups, and significant differences were observed in all displacement parameters. Despite the difference in the upper and lower extremities, these results indicate that unilateral lymphedema may have effects in the way that it may impair the dynamic balance.

As in the results of our study, in other studies where postural stabilization and dynamic balance in individuals with asymmetric body weight distributions were discussed, dynamic balance parameters deteriorated^{2,20}. Factors that were thought to be the cause of these disturbances were deterioration in vestibular, visual, tactile and proprioceptive senses and reduced control of adaptive strategies to maintain an upright stance.

Based on our study, we think limits of stability impairments in unilateral LLL patients could be associated with weight bearing problems due to asymmetric body weight distribution and increased postural sway. So, treatment programs should be planned accordingly, not only by focusing on volume reduction approaches but also considering all other possible balance problems.

One of the limitations of the study was that the Biodex Balance System assessments were not evaluated for closed-eyes and foam

surface conditions. The other one was that postural disorders that may develop due to musculoskeletal problems were not evaluated. By evaluating these parameters in future studies and using a larger sample size, it will be possible to increase the level of knowledge in this subject and obtain more objective results.

5. CONCLUSION

According to the results of this study, asymmetric fluid distribution may be related to balance distortions. Static and dynamic balances, which are related to postural stabilization, may be impaired in patients with unilateral lower limb lymphedema. The authors declare no conflicts of interest.

REFERENCES


1. Bouche K., Stevens V., Cambier D., Caemaert J., Danneels L.: *Comparison of postural control in unilateral stance between healthy controls and lumbar discectomy patients with and without pain*. European Spine Journal, 2006; 15: 423-432.
2. Rougier P.R., Genthon N.: *Dynamical assessment of weight-bearing asymmetry during upright quiet stance in humans*. Gait Posture, 2009; 29: 437-443.
3. Angin S., Karadibak D., Yavuzşen T., Demirbüken İ.: *Unilateral upper extremity lymphedema deteriorates the postural stability in breast cancer survivors*. Contemp Oncol, 2014; 18: 279.
4. Hormes J.M., Bryan C., Lytle L.A., Gross C.R., Ahmed R.L., Troxel A.B., et al.: *Impact of lymphedema and arm symptoms on quality of life in breast cancer survivors*. Lymphology, 2010; 43: 1-13.
5. Ahmed R.L., Prizment A., Lazovich D., Schmitz K.H., Folsom A.R.: *Lymphedema and quality of life in breast cancer survivors: the Iowa Women's Health Study*. J Clin Oncol, 2008; 26: 5689-5696.
6. International Society of Lymphology. *The diagnosis and treatment of peripheral lymphedema: 2013 Consensus Document of the International Society of Lymphology*. Lymphology, 2013; 46: 1-11.
7. Road R.: *Biodex Balance System Unpacking Instructions*. Shirley, NY: Biodex Medical System; 1999.
8. Riley P.O., Benda B.J., Gill-Body K.M., Krebs D.E.: *Phase plane analysis of stability in quiet standing*. J Rehabil Res Dev, 1995; 32: 227.
9. Zhang X.A., Ye M., Wang C.T.: *Effect of unilateral load carriage on postures and gait symmetry in ground reaction force during walking*. Comput Methods Biomech Biomed Engin, 2010; 13: 339-344.
10. Symvoulakis E.K., Anyfantakis D.I., Lionis C.: *Primary lower limb lymphedema: a focus on its functional, social and emotional impact*. Int J Med Sci, 2010; 7: 353.
11. Franks P.J., Moffatt C.J., Doherty D.C., Williams A.F., Jeffs E., Mortimer P. S.: *Assessment of health-related quality of life in patients with lymphedema of the lower limb*. Wound Repair Regen, 2006; 14: 110-118.

12. Basar S., Bakar Y., Keser I., Kaba H., Güzel N.A., Özdemir Ö. Ç., et al.: *Does Lymphedema Affect the Postural Stability in Women After Breast Cancer?*. Top Geriatr Rehabil, 2012; 28: 287-294.
13. Duclos C., Roll R., Kavounoudias A., Mongeau J.P., Roll J.P., Forget R.: *Postural changes after sustained neck muscle contraction in persons with a lower leg amputation*. J Electromyogr Kinesiol, 2009; 19: 214–222.
14. Arifin N., Osman A., Azuan N., Ali S., Gholizadeh H., Abas W., et al.: *Postural stability characteristics of transtibial amputees wearing different prosthetic foot types when standing on various support surfaces*. The Scientific World Journal, 2014.
15. Sundström N., Djerf L., Olivecrona Z., Koskinen L.O.D.: *Postural stability in patients with chronic subdural hematoma*. Acta Neurochir, 2016; 158: 1479-1485.
16. Vrieling A.H., Van Keeken H.G., Schoppen T., Otten E., Hof A.L., Halbertsma J.P.K., et al.: *Balance control on a moving platform in unilateral lower limb amputees*. Gait Posture, 2008; 28: 222-228.
17. Liaw M.Y., Chen C.L., Pei Y.C., Leong C.P., Lau Y.C.: *Comparison of the static and dynamic balance performance in young, middle-aged, and elderly healthy people*. Chang Gung Med J, 2009; 32: 297-304.
18. Singh D., Park W., Levy M.S., Jung E.S.: *The effects of obesity and standing time on postural sway during prolonged quiet standing*. Ergonomics, 2009; 52: 977-986.
19. Ku P.X., Abu Osman N.A., Yusof A., Abas W.W.: *Biomechanical evaluation of the relationship between postural control and body mass index*. J Biomech, 2012; 45: 1638-1642.
20. Mansfield A., Danells C.J., Zettel J.L., Black S.E., McIlroy W.E.: *Determinants and consequences for standing balance of spontaneous weight-bearing on the paretic side among individuals with chronic stroke*. Gait Posture, 2013; 38: 428-432.

A COMPLETE PROGRAM FOR THE TREATMENT OF LYMPHOEDEMA

CIZETA PRESS®

Short extensibility bandages



VARISAN® FLAT

Flat knitting elastic garments,
standard and made to measure



LINFORoll®

Medical device for the
lymphatic drainage



CIZETA MEDICALI S.p.A. - Via IV Novembre, 46
20012 Cuggiono (MI) Italy

Tel. +39 02.97.21.81.1 - Fax +39 02.97.24.07.34

www.cizetamedicali.com - info@cizetamedicali.com


**CIZETA
MEDICALI**
**SCIENCE & MEDICAL
QUALITY**