



Colurodontis paxmani Hutchins, 1977

En - Paxman's leatherjacket.

Maximum total length about 15 cm. Inhabits shallow seagrass beds. Found in northern Western Australia to Cape York.




(after Hutchins, 1977)




Eubalichthys mosaicus (Ramsay and Ogilby, 1886)

En - Mosaic leatherjacket.

Maximum total length about 60 cm. Inhabits inshore and offshore reefs to depths of 170 m. Marketed occasionally in southeastern Australia. Found in Australia's southern half.




(after Allen, 1985)




Monacanthus chinensis (Osbeck, 1765)

En - Fanbellied leatherjacket.

Maximum total length about 38 cm. Inhabits shallow seagrass beds and coastal reefs. Marketed in Australia, Indonesia, and the Philippines. Occurs in northern Australia, New Guinea, Indonesia, Malaysia, Thailand, Philippines, China (including Taiwan Province), and southern Japan.




(after Schroeder, 1980)




Nelusetta ayraudi (Quoy and Gaimard, 1824)**En** - Chinaman leatherjacket.


Maximum total length about 71 cm. Inhabits shallow embayments as a juvenile, adult moving into offshore waters at depths to 120 m. Marketed in southeastern Australia. Found only in southern Australia.




(after Kailola et al., 1990)

***Oxymonacanthus longirostris*** (Bloch and Scheider, 1801)**En** - Harlequin filefish.


Maximum total length about 9 cm. Inhabits shallow coral reefs. A popular aquarium fish. Widespread in the Indo-West Pacific, from East Africa to Samoa, and northern Australia to southern Japan.




(after Schroeder, 1980)

***Paraluteres prionurus*** (Bleeker, 1851)**En** - Blacksaddle filefish.

Maximum total length about 10 cm. Inhabits shallow coral reefs. Widespread throughout the Indo-West Pacific, from East Africa to the Marshall Islands, and from northern Australia to southern Japan.




(after Debelius, 1993)





***Paramonacanthus choirocephalus* (Bleeker, 1852)**

En - Pigface leatherjacket.

Maximum total length about 14 cm. Inhabits trawling grounds at depths between 10 and 58 m. Marketed in Thailand. Occurs in northwestern Australia, New Guinea, Indonesia, Malaysia, Thailand, and the Philippines.




(after Gloerfelt-Tarp and Kailola, 1984)




***Paramonacanthus japonicus* (Tilesius, 1810)**

En - Japanese leatherjacket.

Maximum total length about 13.5 cm. Inhabits trawling grounds at depths between 8 and 80 m. Marketed in Thailand. Found in parts of northern Australia, New Guinea, Indonesia, Malaysia, Thailand, Philippines, and southern Japan.





(after Masuda et al., 1984)



***Paramonacanthus lowei* Hutchins, 1997**



En - Lowe's leatherjacket.

Maximum standard length about 18.7 cm. Inhabits embayments and inshore trawling grounds at depths to 62 m. Occurs along the east coast of Australia.





Paramonacanthus otisensis Whitley, 1931**En** - Dusky leatherjacket.

Maximum total length about 14 cm. Inhabits inshore trawling grounds at depths between 10 and 23 m. Occurs in eastern Australia only.

***Paramonacanthus pusillus*** (Rüppell, 1828)**En** - Sinhalese leatherjacket.



Maximum standard length about 18.8 cm. Inhabits trawlings grounds at depths between 28 and 79 m. Occurs in the Indo-West Pacific, from East Africa to northwestern Australia, and north to Japan. In the past, this species has been often referred to as *Laputa cingalensis*, which is a junior synonym.



(after Sainsbury et al., 1985)

Pervagor janthinosoma (Bleeker, 1854)**En** - Gillblotch leatherjacket.

Maximum total length about 14 cm. Inhabits shallow coral and rocky reefs. A popular aquarium fish. Widespread in the Indo-West Pacific, from East Africa to Samoa, and from northern Australia to southern Japan.



(after Randall, Allen, and Steene, 1990)