

Forgotten Books

— www.forgottenbooks.com —

Copyright © 2016 FB &c Ltd.

All rights reserved. No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the publisher, except in the case of brief quotations embodied in critical reviews and certain other noncommercial uses permitted by copyright law.

GULIELMI HUDSONI,

Regiæ Societatis Socii et Pharmacopæi Londinensis,

⁴⁷²⁶
F L O R A
A N G L I C A ;

EXHIBENS

P L A N T A S

P E R

Regnum BRITANNIÆ sponte crescentes,

DISTRIBUTAS

Secundum Systema Sexuale :

C U M

DIFFERENTIIS SPECIERUM,
SYNONYMIS AUCTORUM,
NOMINIBUS INCOLARUM,
SOLO LOCORUM,
TEMPORE FLORENDI,
OFFICINALIBUS PHARMACOPEORUM.

*Robert. Group, Pharmacopæi
to Harward Catalogue*
L O N D O N :

PRINTED FOR R. FAULDER; WM. OTRIDGE AND SON; R.
LEA; E. JEFFERY; OGILVIE AND SON; J. CUTHELL;
LACKINGTON, ALLEN, AND CO. VERNOR AND HOOD;
J. WALKER; AND J. NUNN.

MDCCXCVIII.

~~1879~~

1879. April 9.
Paris request.

NOBILISSIMO VIRO
HUGONI DUCI
NORTHUMBRIÆ,

ORDINIS PERICELIDIS EQUITI, &c.

ARTIUM TUM UTILIUM

TUM ELEGANTIORUM

JUDICI ET PATRONO,

HOC OPUS HUMILLIME DEDICAT

GULIEL. HUDSON.

P R Æ F A T I O.

PAUCIS lectorem præfari necessarium duxi, tum ut eum admonerem quid in hac Florâ expectare debeat, tum ut notas quasdam compendii gratiâ usurpatis explicarem.

Fundamentum hujus operis quoad *Catalogum* Plantarum Raii synopsis agnoscitur, et agnosci debet ; quantulum enim intactum reliquit vir ille diligentissimus, accuratissimus, et addere licet, sagacissimus, contentibus tum popularibus suis, tum exteris ; qui floram ejus inter optimas et perfectissimas uno ore numerant ? si nullum igitur post ejus tempora augmentum accepisset ipsa scientia botanica, nihil pene pro plantis Anglicanis dignoscendis ulterius desideraretur, quam nova synopsis editio, cujus exemplaria hodie apud bibliopolas omnino non inveniuntur. At exortum est his diebus novum sidus, quod orbi botanico lucem affudit ne in somnis quidem antea visam ;

visam ; magnus scilicet ille Linnæus, qui partes plantarum minutissimas, vel prius neglectas, vel non omnino observatas in conspectum producendo, plene illustrando, et rite nominando fundamentum jecit novæ methodo simplici, compendiosæ, certæ ; quam et constantiâ plane admirandâ toti regno vegetabili adhuc cognito feliciter applicuit.

Hunc igitur virum egregium quoad *methodum* per omnia, quatenus licuit, secutus sum.

Essentiales characteres ex Systemate Naturæ, edit. XII: adduxi ; quando autem novum genus occurrebat, characterem naturalem ad ejus exemplar, prout in *Genera Plantarum* exhibetur, adjunxi.

Quoad Synonyma, ut vocantur, sive Characteres Specificos, primo loco eos adhibui quos idem author in *Species Plantarum* dedit ; quotiescunque certum fuit eum plantam descripsisse, aliter novum characterem, ejus vestigiis insistens construxi ; postea synonyma adjeci ex Bauhini pinace, Raio, Gerardo, Parkinsono et varijs authoribus hodiernis deprompta.

Nomina plerumque Anglica ex Raio, Gerardo, &c. apposui, quædam mutavi exemplo et hortatu Domini Stillingfleet motus, ut generica obtinerem, quod optandum esset per omnia, ut confusio, quæ hodie regnat, finis aliquando imponatur ;

natur ; quando autem nulla occurrebant meo jure, ut opinor, imposui.

Quoad loca natalia plantarum satis duxi indicasse solum, quo maxime vulgares sponte proveniunt ; locum verò rariorum semper notavi ex Raii Synopsi ultima editione et ex observationibus, sive propriis, sive amicorum.

Plantarum naturam quoad durationem indicavi signis sequentibus ; viz. ☉ annum, ♂ biennem, ♃ perennem innuit.

Hoc signum ♁ fruticem vel arborem designat.

Numeri I, II, III, &c. tempus florendi indicant, viz. Januarium, Februarium, Martium, &c.

Plantas cultas non autem indigenas omnino omisi, ut Avenam sativam, Triticum sativum, Solanum tuberosum, &c.

Plantæ quæ sine culturâ non raro inveniuntur extra hortorum pomæria, sed quæ ab origine propagationem eorum rejectamentis debeat asterisco * distinguuntur.

Hanc occasionem lubens amplector gratias agendi amicis, qui mihi plantas pro eorum benevolentia suppeditarunt inter quos nominandi sunt Domini Alchorne, Bolton, Collinson, Lyons, Miller, Pultney, Watson, M.D. Wilmer, M.D. præcipue vero D. Stillingfleet, qui et hoc et aliis amicitiae privatim et publice functus est.

Cæterum quamvis nihil sciens prætermisisti quod conferre potuit elucidando rem subjectam, non la-
bori

bori pepercit, nullam occasionem neglexi libros consulendi, Herbaria sicca evolvendi et Plantas vivas examinandi et desecandi, ut opus omnibus numeris absolutum prodiret; conscius sum tamen me nullo modo scopum attigisse, præsertim quoad classem cryptogamiam, quæ ad hæc usque tempora quantum scio, obscura manet, et mihi plurimum molestiæ creavit. In uno istius ordine, viz. Fungorum me multum deficere agnosco; nec sine aliis Dilleniis, aliis Micheliis extricare se quisquam poterit, cum hunc ordinem experientia edoctus affirmare possim, ut Linnæi verbis utar, etiamnum Chaos esse nescientibus Botanicis in his, quid sit Species, quid varietas.

In hac secunda editione plurima emendavi et plantas a me noviter detectas vel ab amicis communicatas in propriis locis inserui.

T E R M I N I

B O T A N I C I.

RADIX organum nutriens plantam.

- | | |
|-----------------|--|
| Durati- one. | 1. <i>Annua</i> intra annum emoritura. |
| | 2. <i>Biennis</i> altero anno florens et dein peritura. |
| | 3. <i>Perennis</i> per plures annos regerminans. |
| Figura. | 4. <i>Fibrosa</i> tota constans filamentis. |
| | 5. <i>Ramosa</i> in fibras subdivisa. |
| | 6. <i>Fusiformis</i> simpliciuscula attenuata. |
| | 7. <i>Præmorsa</i> apice quasi truncata. |
| | 8. <i>Repens</i> longe excurrentis, hinc inde germinans. |
| | 9. <i>Articulata</i> geniculis intercepta. |
| | 10. <i>Dentata</i> moniliformis ex articulis concatenatis. |
| | 11. <i>Globosa</i> subrotunda (156) radiculis lateralibus. |
| | 12. <i>Tuberosa</i> e partibus carnosis filo basi connexis. |
| | 13. <i>Fascicularis</i> e partibus carnosis basi sessili (150) connexis. |
| | 14. <i>Palmata</i> carnosa lobata (182). |
| | 15. <i>Bulbosa</i> radix bulbo (648) instructa. |
| | 16. <i>Granulata</i> particulis carnosis adspersa. |

TRUNCUS organum multiplicans plantam.

- | | |
|-----------------|--|
| Specie. | 17. <i>Caulis</i> , truncus elevans et fructificationem et folia. |
| | 18. <i>Culmus</i> graminibus proprius. |
| | 19. <i>Scapus</i> elevans fructificationem, nec folia. |
| | 20. <i>Stipes</i> truncus in folia transiens. |
| Durati- one. | 21. <i>Herbaceus</i> etiamnum annus (non lignosus) |
| | 22. <i>Suffruticosus</i> basi permanens, ramis quotannis marcescens. |

23. *Fruticosus* perennis cum caudicibus pluribus.
 24. *Arboreus* perennis caudicibus simplici.
 25. *Solidus* interne factus.
 26. *Inanis* interne medulla spongiosus.
 27. *Fistulosus* interne tubulosus.
 Directi- 28. *Erectus* fere ad perpendiculum se attollens.
 one. 29. *Strictus* omnino perpendicularis absque flexura.
 30. *Rigidus* flexionis impatiens.
 31. *Laxus* libere in arcum flexibilis.
 32. *Obliquus* a perpendiculari horizontalive linea discedens.
 33. *Adscendens* arcuatim sursum versus.
 34. *Declinatus* arcuatim descendens.
 35. *Incurvatus* introrsum nutans.
 36. *Nutans* apice extrorsum reflexus.
 37. *Diffusus* ramulis patentibus.
 38. *Procumbens* debilis terra innitens.
 39. *Stoloniferus* turiones ad radicem edens.
 40. *Sarmentosus* filiformis geniculis radicans.
 41. *Repens* terræ incumbens radicalesque agens.
 42. *Radicans* alis se affigens radiculis lateralibus.
 43. *Geniculatus* internodiis interceptus.
 44. *Flexuosus* a gemma ad gemmam horsum versus actus.
 45. *Scandens* alta petens, aliis sustinendus.
 46. *Volubilis* spiraliter adscendens per alia corpora.
 Dextrorsum a dextra ad sinistram.
 Sinistrorsum a sinistra ad dextram.
 Figura. 47. *Teretis* angulis destitutus.
 48. *Semiteretis* hinc planus; inde teretiusculus.
 49. *Compressus* duobus lateribus oppositis planis.
 50. *Anceps* angulis duobus oppositis acutiusculus.
 51. *Angulatus* pluribus quam duobus angulis cavis longitudinaliter excavatus.
 Acutangulus figura angulorum.
 52. *Trigonus*, &c. angulis tribus prominentibus longitudinalibus.
 53. *Triqueter* lateribus tribus exacte planis.
 Vestitu. 54. *Nudus* oppositus 5, 199. (55, 56, 57, 58, 59).
 55. *Aphyllus* foliis destitutus.
 56. *Foliatus* foliis instructus.
 57. *Vaginatus* foliorum vaginis cinctus.
 58. *Squamosus*

Superfi-
cie,

58. *Squamosus* squama adpersus.
 59. *Imbricatus* tectus, ut nudus non appareat.
 60. *Suberosus* cortice exteriori molliore, sed elastico indutus.
 61. *Rimosus* cortice exteriori fissuras sponte agens.
 62. *Tunicatus* membranis vestitus.
 63. *Lævis* superficie æquali.
 64. *Striatus* lineis tenuissimis excavatis inscriptus.
 65. *Sulcatus* lineis profundis exaratus.
 66. *Glaber* (213).
 67. *Scaber* (239).
 68. *Muricatus* punctis subulatis adpersus.
 69. *Tomentosus* (234).
 70. *Villosus* (223).
 71. *Hispidus* (240).
 72. *Aculeatus* (241).
 73. *Spinosus* spinis (382) armatus.
 74. *Urens* stimulus (389) adpersus.
 75. *Stipulatus* stipulis (289) notatus.
 76. *Membranaceus* complanatus more folii.
 77. *Bulbiferus* bulbos (648) gerens.
- Compo-
sitione.
78. *Enodis* continuus absque articulis.
 79. *Simplicissimus* ramis vix ullis.
 80. *Simplex* continua serie versus apicem extensus.
 81. *Integer* simplicissimus ramis vix angustatis.
 82. *Articulatus* internodiis geniculatus.
 83. *Prolifer* ex apicis centro tantum emittens ramos.
 84. *Dichotomus* bifariam semper divisus.
 85. *Brachiatus* ramis decussatim oppositis.
 86. *Subramosus* ramis paucissimis lateralibus.
 87. *Ramosus* ramis pluribus lateralibus.
 88. *Ramosissimus* ramis multis absque ordine cumulatus.
 89. *Virgatus* ramusculis debilibus inæqualibus.
 90. *Paniculatus* ramis varie subdivisis.
 91. *Fastigiatus* ramis æqualis altitudinis.
 92. *Patens* (131).
 93. *Divaricatus* (103).

RAMI partes caulis.

94. *Alterni* (113).
95. *Distichi* (114).
96. *Sparsi* (116).

97. *Conferti* (117).
 98. *Oppositi* (124).
 99. *Verticillati* plures truncum ad genicula circumdantes.
 100. *Erecti* (127).
 101. *Coarctati* versus summitatem fere incumbentes.
 102. *Divergentes* a trunco ad angulum rectum discedentes.
 103. *Divaricatis* ad angulum obtusum a trunco discedentes.
 104. *Deflexi* in arcum deorsum inclinati.
 105. *Reflexi* perpendiculariter dependentes.
 106. *Retroflexi* horsum vorsum divaricati.
 107. *Fulcrati fulcris* (287) instructum.

FOLIA organa motus plantæ.

- Determina- 108. *Radicale* radici insidens.
 tione: 109. *Caulinum* cauli insertum.
 loco. 110. *Rameum* ramis insidens.
 111. *Axillare* sub rami basi insertum.
 112. *Florale* flori proximum.
 Numero. *Unicum, duo, tria, pauca, plurima.*
 Situ 113. *Alterna* per gradus circum ramum exorta.
 114. *Disticha* duo latera rami respicientia, licet undique
 inserta.
 115. *Bifaria* ad latera rami opposita tantum enata.
 116. *Sparsa* absque certo ordine constituta.
 117. *Conferta* plurima totum fere occultantia ramum vel
 caulem.
 118. *Imbricata* qua dimidiam partem invicem tecta.
 119. *Fasciculata* plura ex eodem puncto ad genicula ra-
 morum.
 120. *Bina, trina, quina, &c.* de numero ad genicula
 ramorum.
 121. *Confluentia* ad basin inter se cohærentia.
 122. *Approximata* ad se invicem proxime accedentia.
 123. *Remota* spatio a se invicem distantia.
 124. *Opposita* per paria decussatim collocata.
 125. *Decussata* ita disposita opposite, ut rami, apice
 inspecto, quatuor ordines referant.
 126. *Stellata* plura quam duo folia caulem ambiente.
 Directi- 127. *Erectum* fere ad perpendicularum constitutum.
 one. 128. *Strictum* omnino perpendicularare absque flexura.

129. *Rigidum*

TERMINI BOTANICI.

129. *Rigidum* flexionis impatiens.
130. *Appressum* disco cauli approximatum.
131. *Pateens* ad angulum acutum cauli insidens.
132. *Horizontale* ad angulum rectum a caule descendens.
133. *Assurgens* arcuatim erectum, primum declinatum deinde apice erectum.
134. *Inflexum* sursum arcuatum versus apicem.
135. *Reclinatum* deorsum flexum, ut arcus sit basi inferior, apice ascendente.
136. *Recurvatum* deorsum, ut arcus superiora spectat.
137. *Revolutum* in spiram recurvatum.
138. *Dependens* terram recta spectans.
139. *Obliquum* basi cœlum, apice horizontem spectans.
140. *Adversum* paginam superiorem meridiei (non cœlo) obvertens.
141. *Verticale* obversum, ut regio basis angustior evadat regione apicis.
142. *Resupinatum* pagina superiore inferiore et contra inferiore superiore facta.
143. *Submersum* intra aquæ superficiem absconditum.
144. *Natans* superficiem aquæ incumbens.
145. *Radicans* radices agens.
- Infer- 146. *Petiolum* petiolo (288) ad basin inserto.
- one. 147. *Petitum* petiolo disco folii inserto.
148. *Sessile* immediate cauli (absque petiolo) insidens.
149. *Adnatum* pagina superiore basi rami adnexum.
150. *Coadunatum* plura inter se connata.
151. *Decurrens* basi folii deorsum per caulem extensa.
152. *Amplexicaule* basi caulem ambiens.
153. *Perfoliatum* basi transversim cingente (nec antice dehiscente) caulem.
154. *Connata* oppositorum paribus basi utrinque coadunatis.
155. *Vaginans* basi formante tubum caulem vestientem.
- Stru- 156. *Subrotundum* figura orbicularæ (157) proxima.
- tu- 157. *Orbiculatum* peripheria circinnata (diameter longi-
- ra.Figura 158. *Ovatum* cujus diameter longitudinalis superat trans-
159. *Ovale* ex orbiculato oblongo utraque extremitate rotundata æquali.

160. *Oblongum*, diametro longitudinali aliquoties superante transversalem.
161. *Parabolicum*, versus apicem sensim angustius rotundatum.
162. *Cuneiforme* sensim versus basin angustatum.
163. *Spathulatum* subrotundum (156) basi angustiore lineari (167).
164. *Rotundatum* angulis orbium.
165. *Lanceolatum* oblongum ab utraque extremitate attenuatum.
166. *Ellipticum* lanceolatum, latitudine ovati folii.
167. *Lineare* æquali ubique latitudine.
168. *Acerosum* lineare (167) persistens (271).
- Angulis. 169. *Integrum* indivisum, omnique sinu destitutum.
170. *Triangulare*, &c. a numero angulorum.
171. *Deltoideum* rhombeum (172) ex quatuor angulis, e quibus laterales minus a basi distant, quam reliqui.
172. *Rhombeum* forma rhombi.
173. *Trapeziforme* forma trapezii.
- Sinubus. 174. *Cordatum* subovatum basi sinu exsculpta absque angulis posticis.
175. *Reniforme* subrotundum basi sinu exsculpta absque angulis posticis.
176. *Lunatum* subrotundum basi sinu divisa angulis posticis acutis.
177. *Sagittatum* triangulare, angulis posticis acutis sinu divisis.
178. *Hastatum* sagittatum (177) angulis posticis sinu divisis ad latera prominentibus.
179. *Runcinatum* pinnatifidum (187) ita, ut lobi antice convexi, postice sint transversi. e. g. *Leontodon*.
180. *Panduriforme* oblongum lateribus inferne coarctatum.
181. *Fissum* divisum sinubus linearibus, marginibus rectis.
182. *Lobatum* ad dimidium divisum in partes distantes.
183. *Bi-Quinquesidum* pro numero fissurarum (181).
184. *Partitum* ad basin fere divisum.
185. *Palmatum* divisum ultra dimidium in lobos subæquales.
186. *Lyratum*

186. *Lyratum* transversim divisum in lacinias, quarum inferiores minores remotiores.
187. *Pinnatifidum* transversim divisum in lacinias horizontales oblongas.
188. *Sinuatum* a lateribus sinibus distatis.
189. *Laciniatum* varie et indeterminate in partes sectum.
190. *Squarrosum* divisum in lacinias elevatas, nec plano parallelas.
- Margine.** 191. *Integerrimum* ipso margine lineari, nec minimum secto.
192. *Crenatum* margine incisuris sine respectu extremitatis secto.
193. *Serratum* margine omnibus incisuris respicientibus extremitatem.
194. *Ciliatum* margine setis parallelis longitudinaliter digestis.
195. *Dentatum* margine acuminibus patentibus remotis.
196. *Spinosum* margine acuminibus subulatis rigidis pungentibus.
197. *Cartilagineum* marginē subosseo.
198. *Repandum* margine flexuoso, tamen plano.
199. *Lacerum* margine varie divisum, segmentis difformibus.
200. *Erosum* sinuatum (188) sinibus minimis obtusis laciniisque inæqualibus.
201. *Dædaleum* una flexuosum lacerumque (199).
202. *Obtusum* intra segmentum circuli terminatum.
203. *Emarginatum* terminatum crena.
204. *Retusum* terminatum sinu obtuso.
205. *Præmorsum* terminatum obtuse incisuris inæqualibus.
206. *Truncatum* terminatum linea transversali.
207. *Acutum* terminatum angulo acuto.
208. *Acuminatum* terminatum apice subulato (266).
209. *Cuspidatum* terminatum apice setaceo.
210. *Mucronatum* terminatum mucrone extante.
211. *Cirrhosum* cirrho (290) terminatum.
- Superficie.** *Pagina* superior vulgo cælum, inferior terram spectat.
212. *Nudum* setis et pilis destitutum.
213. *Glabrum* superficie lubrica.
214. *Nitidum* glabricie lucente.

TERMINI BOTANICI.

215. *Lucidum* quasi illuminatum.
216. *Coloratum* alio corpore quam viridi.
217. *Nervosum* vasis simplicissimis a basi ad apicem.
218. *Trinerve* tribus nervis in basi folii supra basin concurrentibus.
219. *Triplinerve* nervosum tribus nervis pone basin folii concurrentibus.
220. *Enerve* nervoso (217. 219) oppositum.
221. *Lineatum* nervis depressis.
222. *Striatum* lineis parallelis leviter excavatum.
223. *Sulcatum* lineis profundis excavatum.
224. *Venosum* vasis multifariam divisis.
225. *Rugosum* rugis refertum.
226. *Bullatum* ex rugoso (225) venis contractis, ab altera parte concavis.
227. *Lacunosum* disco depresso inter venas interjectas.
228. *Avene* venoso (224) oppositum.
229. *Punctatum* adpersum punctis excavatis.
230. *Papillatum* tectum punctis carnosis.
231. *Papulosum* tectum punctis vesicularibus.
232. *Viscidum* oblitum tenaci humore.
233. *Villosum* tectum pilis mollibus.
234. *Tomentosum* villis intertextis nec discernendis obductum.
235. *Sericeum* tectum pilis adpressis mollissimis.
236. *Lanatum* indutum quasi tela aranæ (pilis sponte curvatis).
237. *Barbatum* pilis parallelis obsitum.
238. *Pilosum* pilis distinctis elongatis tectum.
239. *Scabrum* punctis eminentibus rigidiusculis exasperatum.
240. *Hispidum* setis rigidis adpersum.
241. *Aculeatum* aculeis (376) armatum.
242. *Strigosum* aculeis lanceolatis (165) rigidis (30).
243. *Planum* superficie æqualis.
244. *Canaliculatum* supra sulco profundo longitudinaliter excavatum.
245. *Concavum* margine disco arctiore, ut deprimatur discus.
246. *Convexum* margine disco arctiore, ut elevatur discus.
247. *Cucullatum* lateribus ad basin conniventibus apice vero dilatatis.

248. *Plicatum*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

275. *Conjugatum*, pinnatum (279) tantum binis foliolis lateralibus.
276. *Digitatum* petiolo simplici apice adnectente foliola.
277. *Binatum* digitatum (276) foliolis duobus terminatum, &c.
278. *Pedatum* petiolo bifido, latere tantum interiore adnectente foliola plura.
279. *Pinnatum* petiolo simplici lateribus adnectente foliola plura.
280. *Bijugum* sic *trijuga*, *quadrijuga*, *quinquejuga*, *sejuga*, &c. pinnatum, sed foliolis tantum quatuor, &c.
cum impari pinnatum, foliolo unico (impari) terminatum.
abrupte pinnatum, nec cirrho, nec foliolis, terminatum.
cirrhosum cirrho (290) terminatum.
Foliolis oppositis (124) *alternis* (113).
ruptis foliolis alternis minoribus.
decurseis foliolis petiolo decurrentibus.
- Decom- 281. *Bigeminum* petiolo dichotomo (84) apice adnectente foliola plura.
 posita.
282. *Biternatum* duplicato-ternatum (277).
283. *Bipinnatum* duplicato pinnatum (279).
284. *Tergeminum* triplicato geminatum. Petiolus bifidus utroque apice foliola 2-ferens, et insuper foliola 2 ad divaricationem petioli communis.
285. *Triternatum* triplicato ternatum.
286. *Tripinnatum* triplicato pinnatum.
287. **FULCRA** adminicula plantæ pro commodiore sustentatione.
288. *Petiulus* fulcrum sustinens folium.
289. *Stipula* squama basi petiolorum enascentium adstans.
290. *Cirrhus* vinculum filiforme spirale, quo planta alii corpori alligatur.
291. *Pæbes* hirsuties omnis in planta.
292. *Arma* mucrones arcentes animalia, ne lædant plantam.
293. *Bractea* folium florale (112) facie a ceteris foliis recedens.
294. *Pedunculus* fulcrum sustinens fructificationem.

PETIOLUS

- Figura. 295. *Linearis* (167).
 296. *Alatus* lateribus dilatatus.
 297. *Clavatus* versus apicem incrassatus.
 298. *Membranaceus*, complanatus.
 299. *Teres* (254).
 300. *Semiteres* (48).
 301. *Triquetus* (53).
- Magnitudine. 302. *Brevissimus*, qua magnam partem non attingens longitudinem folii.
 303. *Brevis* nondum longitudinem folii attingens.
 304. *Mediocris* longitudine folii.
 305. *Longus* superans longitudinem folii.
 306. *Longissimus*, aliquoties superans longitudinem folii.
- Insertione. 307. *Insertus* perpendiculariter ramo infidens.
 308. *Adnatus* (149).
 309. *Decurrens* (151).
 310. *Amplexicaulis* (152).
 311. *Appendiculatus*, ramentis foliaceis ad basin.
- Directione. 312. *Erectus* (127).
 313. *Patens* (131).
 314. *Assurgens* (133).
 315. *Recurvatus* (136).
- Superficie. 316. *Glaber* (213).
 317. *Aculeatus* (241).
 318. *Nudus* (212).
 319. *Articulatus* (82).
 320. *Spinescens* indurefcens et pungens.
- STIPULÆ. 321. *Geminæ* duæ et duæ per paria.
 322. *Solitariæ* simplices sparsæ.
 323. *Laterales* lateribus insertæ.
 324. *Extrafoliaceæ* infra folium collocatæ.
 325. *Intrafoliaceæ* supra folium collocatæ.
 326. *Oppositifoliæ* in latere folii opposito-collocatæ.
 327. *Caducæ* (270).
 328. *Deciduae* (269).
 329. *Persistentes* post defoliationem restantes.
 330. *Spinescentes*

TERMINI BOTANICI.

330. *Spinescentes* (320).
 331. *Sessiles* (148).
 332. *Adnatæ* (149).
 333. *Decurrentes* (151).
 334. *Vaginantes* (155).
 335. *Subulatæ* (266).
 336. *Lanceolatæ* (165).
 337. *Sagittatæ* (177).
 338. *Lunatæ* (176).
 339. *Ereclæ* (127).
 340. *Patentes* (131).
 341. *Integerrimæ* (191).
 342. *Serratæ* (193).
 343. *Ciliatæ* (194).
 344. *Dentatæ* (195).
 345. *Fissæ* (181).
CIRRHUS. 346. *Axillaris* (111).
 347. *Foliaris* folio insidens.
 348. *Petiolearis* petiolo (288) insidens.
 349. *Peduncularis* pedunculo (294) insidens.
 350. *Simplex* indivisus.
 351. *Trifidus* in tres partes divisus.
 352. *Multifidus* multoties divisus.
 353. *Convolutus* in annulos contortus.
 354. *Revolutus* spira dimidio itinere retorta.
PUBES. 355. *Pili* ductus excretorei plantæ setacei.
 356. *Lana* pili curvi densi.
 357. *Barba* pili paralleli.
 358. *Tomentum*, villi intertexti vix conspicui.
 359. *Strigæ* pili rigidiusculi planiusculi.
 360. *Setæ*, pili rigidiusculi teretiusculi.
 361. *Simplices*, continuo extensæ longitudinaliter.
 362. *Hamosæ*, animalibus facile adhærentes.
 363. *Ramosæ*, subdivisæ quasi in ramulos.
 364. *Plumosæ*, villosæ compositæ.
 365. *Stellatæ*, decussatim depositæ.
 366. *Hami*, mucrones acuminati curvati.
 367. *Glochides*, mucrones apice retrorsum multidentati.
 368. *Glandulæ*.

368. *Glandula papilla* humorem excernens.

369. *Utriculus*, vasculum liquore secreto repletum.

370. *Foliaceæ* foliis insertæ.

371. *Petiolares* (348).

372. *Pedunculares* (349).

373. *Stipulares* stipulis (289) insertæ.

374. *Viscositas* qualitas humoris tenacis.

375. *Glutinositas* qualitas humoris lubrici.

ARMA. 376. *Aculei* mucrones pungentes plantæ cortici tantum affixi.

377. *Recti*, flexuis destituti.

378. *Incurvi*, introrsum flexi.

379. *Recurvi* extrorsum reflexi.

380. *Furca* aculeis divisis in plures.

381. *Bifidæ* et *Trifidæ* a numero divisionis.

382. *Spina*, mucro e ligno plantæ protrusus.

383. *Terminalis* ad apicem plantæ posita.

384. *Axillaris* (111).

385. *Calycina* Calyci (491) insidens.

386. *Foliaris* (347).

387. *Simplex* (361).

388. *Divisa* apice partita.

389. *Stimuli* mucrones puncturas exflamatorias efficientes, unde pruriginosæ evadunt partes.

BRACTEÆ folia sequentis anni.

390. *Coloratæ* (216).

391. *Caducæ* (270).

392. *Decidua* (269).

393. *Persistentes* (271).

394. *Coma* Bracteæ caulem terminantes magnitudine insignes.

PEDUNCULUS. 395. *Partialis*, aliquot flores pedunculi communis proferens.

396. *Communis*, communis pluribus floribus.

397. *Pedicellus*,

TERMINI BOTANICI.

397. *Pedicellus*, proprius floribus in
Pedunculo communi.
398. SCAPUS *pedunculis* radicatis caulis faciem in-
duens.
- Loco. 399. *Radicalis* (108).
400. *Caulinus* (109).
401. *Rameus* (110).
402. *Petiolaris* (348).
403. *Cirrhiferus* (290).
404. *Terminalis* (383).
405. *Axillaris* (111).
406. *Oppositifolius* (326).
407. *Lateriflorus* (323).
408. *Interfoliaceus* (325).
409. *Extrastriaceus* (324).
- Situ. 410. *Alternus* (113).
411. *Sparsus* (116).
412. *Oppositus* (124).
413. *Verticillatus* (99).
- Numero. 414. *Solitarius* (322).
415. *Geminatus* (321).
416. *Umbellula sessilis*, plures pedunculi ex
eodem centro ambitu æquali pro-
deuntes.
- Directione. 417. *Adpressus* (130).
418. *Erectus* (127).
419. *Patens* (131).
420. *Cernuus* apice terram spectat.
421. *Resupinatus* (142).
422. *Declinatus* (34).
423. *Nutans* (36).
424. *Flaccidus*, debilis, ut a proprio floris
pondere dependeat.
425. *Ascendens* (33).
426. *Pendulus*, laxus, ut cum folio deorsum
tendat.
427. *Strictus* (29).
428. *Flexuosus* a flore ad florem flexus.
429. *Retrofractus* vi quasi ad dependentiam
redactus.
- Multiflorus* pro numero florum, quos profert
Pedunculus.

Structura.

- Structura. 430. *Teres* (47).
 431. *Triqueter* (53).
 432. *Tetragonus* (52).
 433. *Filiformis* æquabili ubique crassitie.
 434. *Attenuatus* versus apicem crassitiem perdens sensim.
 435. *Clavus* (297).
 436. *Incrassatus*, apice sensim majorem molem acquirens.
 437. *Nudus* (212).
 438. *Squamosus* (58).
 439. *Foliatus* (56).
 440. *Bracteatus* Bracteis (293) instructus.
 441. *Geniculatus* (43).
 442. *Articulatus* (82).
 443. **INFLORESCENTIA** est modus, quo flores pedunculo plantæ adnectuntur.
 444. **VERTICILLUS** floribus pluribus caulem annulatim ambientibus.
 445. *Sessilis* absque pedicellis manifestis.
 446. *Pedunculatus* pedunculis flores elevantibus.
 447. *Nudus* (448, 449) oppositus.
 448. *Involucratus* involuero (518) instructus.
 449. *Bracteatus* (440).
 450. *Confertus* pedunculis approximatis.
 451. *Distans*, pedunculis remotis.
 452. **CAPITULUM** floribus pluribus in globum congestis.
 453. *Subrotundum*, fere globosum (452).
 454. *Globosum*, ab omni parte rotundum.
 455. *Dimidiatum* ab altera parte rotundatum, ab altera planum.
 456. *Foliosum*, foliis intermixtis floribus.
 457. *Nudum* foliis et setis destitutum.
 458. *Fasciculus* colligens flores erectos, parallelolos, fastigiato approximatos.
 459. **SPICA**, flores sessiles alterni in pedunculo communi simplici.
 460. *Simplex*, continua indivisa.
 461. *Composita*, pluribus spicillis pedunculo insidentibus.
 462. *Glomerata*

TERMINI BOTANICI.

462. *Glomerata* spicillis varie congesta.
 463. *Ovata* (158).
 464. *Ventricosa*, a lateribus gibba (253).
 465. *Cylindrica* teretiuscula. *Secunda* vero dimidiata.
 466. *Interrupta* spicis minoribus alternis distantibus.
 467. *Imbricata* (118).
 468. *Articulata* (82).
 469. *Ramosa*, varie divisa.
 470. *Linearis* (167).
 471. *Ciliata* (194).
 472. *Poliacea* foliis interstinctis.
 473. *Comosa* terminata foliolis.
 474. **CORYMBUS** fit ex *spica* (459) dum singuli flores petiolis propriis instruuntur, situ elevato proportionali.
 475. **THYRSUS**, *Panicula* (487) coarctata in formam ovatam.
 476. **RACEMUS** pedunculo ramis lateralibus instructo.
 477. *Simplex* indivisus.
 478. *Compositus*, divisus in plures.
 479. *Unilateris*, flores ad alterum latus omnes inferti.
 480. *Secundus* flores omnes ad alterum latus flexi.
 481. *Pedatus* (278).
 482. *Conjugatus* (275).
 483. *Erectus* (127).
 484. *Laxus* (31).
 485. *Nudus* (456).
 486. *Foliatus* (56).
 487. **PANICULA** flores sparsi, in pedunculis diverse divisi.

Trunci Structura.

488. **FRUCTIFICATIO** vegetabilium pars temporaria Germinationi dicata.
 489. *Simplex* paucis floribus constans.
 490. *Composita*, pluribus flosculis confluentibus.

491. **CALYX**

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

518. *Involucrum*, calyx a flore remotus.
519. *Universale* umbellæ universali
subjectum.
520. *Partiale* umbellæ partiali sub-
jectum.
521. *Proprium*, cuilibet flori subjec-
tum.
522. *Gluma* Calyx Graminis; valvis amplexanti-
bus.
523. *Uniflora* unicum florem amplectens.
524. *Multiflora* plures flores includens.
525. *Univalvis* unica constans squama.
526. *Bivalvis* duabus constans squamis.
527. *Multivalvis* pluribus quam duabus
squamis.
528. *Colorata* (216).
529. *Glabra* (213).
530. *Hispida* (240).
531. *Arista*, mucro subulatus Glumæ insidens.
532. *Mutica* mucrone destituta.
533. *Terminalis* apice glumæ affixa.
534. *Dorsalis* lateri exteriori glumæ im-
posita.
535. *Recta* perpendiculariter exiens.
536. *Tortilis* flexa funis instar.
537. *Geniculata* (43).
538. *Recurvata* (136).
539. *Amentum ex Receptaculo* (629) communi palea-
ceo gemmaceo.
540. *Spatha* Calyx longitudinaliter ruptus.
541. *Univalvis* uno latere dehiscens.
542. *Dimidiata* latere tantum inferiore
Fructificationem obvestiens.
543. *Calyptra* Calyx MUSCI cucullatus antheræ super-
impositus.
544. *Recta* undique æquales.
545. *Obliqua* ad aliquod latus flexa.
546. *Volva* Calyx FUNGI membranaceus.
547. *Approximata* ad capitulum accedens.
548. *Remotissima* a capitulo spatio recedens.
549. COROLLA

549. **COROLLA** Liber plantæ in flore præfens.
550. *Petalum* pars corollæ in plures divisæ.
551. *Tubus* Corollæ monopetalæ pars inferior.
552. *Unguis* Corollæ polypetalæ pars inferior receptaculo affixa.
553. *Limbus* Corollæ pars superior dilatata.
554. *Lamina* Corollæ monopetalæ pars superior patula. Monopetala.; vel polypetala a numero petalorum.
555. *Regularis* æqualis figura, magnitudine et proportione partium.
556. *Irregularis* Limbi partibus, figura, magnitudine et proportione partium diversa.
557. *Inæqualis* partibus non magnitudine sed proportione respondentibus.
558. *Globosa* globum referens.
559. *Campanulata* ventricosa (464) tubo destituito.
560. *Infundibuliformis*, conica tubo imposita.
561. *Rotata* plana nulli tubo imposita.
562. *Ringens* irregularis in' duo labia hians.
563. *Galea Ringentis* Labium superius.
Labium pro ringentis labio inferiore sæpius sumitur.
564. *Faux* hiatus inter lacinias corollæ, ubi tubus terminatur.
565. *Personata* Ringens (562) sed inter labia palato clausa.
566. *Cruciata* petalis quatuor æqualibus patens.
567. *Concava* (249).
568. *Patens* (131).
569. *Papilionacea* irregularis, petalo inferiore cymbiformi (*carina*), superiore adscendente (*vexillum*) lateribus solitariis (*alæ*).
570. *Composita* pluribus constans flosculis intra commune perianthium, supra receptaculum commune.
571. *Ligulata* Corollulis flosculorum omnibus planis versus exterius latus.

TERMINI BOTANICI.

572. *Tubulosa* corollulis flosculorum omnibus tubulosis subæqualibus.
573. *Radiata* corollulis disci tubulosus ambitus vero ligulatis difformibus Color notetur.
574. *Nectarium* pars mellifera flori propria.
575. *Proprium* a petalis & ceteris partibus distinctum.
576. *Petalinum* Petalis insertione.
577. STAMEN Viscus pro Pollinis præparatione.
578. *Filamentum* pars elevans adnectensque Antheram (organum genitale masculinum).
579. *Æqualia* eandem servantia longitudinem.
580. *Inæqualia* aliquot majoribus vel minoribus.
581. *Connata* in unum corpus coalita.
582. *Anthera* pars floris gravida polline, quod matura dimittit.
583. *Distinctæ* nec inter se cohærentes.
584. *Connatæ* in unum plures concretæ.
585. *Pollen*, pulvis floris, humore rumpendus, atomosque elasticos ejaculans, (genitura plantarum).
586. PISTILLUM, viscum fructui adhærens pro pollinis receptione (organum genitale femineum).
587. *Germen*, rudimentum fructus immaturi in flore
588. *Superum* corollæ inclusum.
589. *Inferum*, infera corollam positum.
590. *Stylus*, pars pistilli elevans stigma a germine.
591. *Stigma*, summitas pistilli madida humore.
592. PERICARPIUM, viscus plantæ gravidum seminibus, quæ matura dimittet.
593. *Capsula* pericarpium cavum determinate dehiscens.
594. *Vacuola*, paries, quo tegitur fructus externe.
595. *Loculamentum*, concameratio cava pro seminum loco.
596. *Dissepimentum*

596. *Dissepimentum*, paries, quo fructus interne distinguitur in concamerationes plures.
597. *Bicapsularis* a numero capsularum.
598. *Bilocularis* a numero loculamentorum.
599. *Tricocca* capsula tribus nodis protuberans, intusque in tria loculamenta divisa.
600. *Didyma*, Capsula duobus nodis extus gibba.
601. *Siliqua*, Pericarpium bivalve affigens semina secundum utramque futuram.
602. *Compressa* lateribus oppositis proxime ad se invicem accedentibus.
603. *Torulosa* prominentibus hinc inde gibba.
604. *Articulata* geniculis coarctatis intercepta.
605. *Paralielum dissepimentum* latitudine et diametro transversali ad valvulas accedens.
606. *Transversum dissepimentum* angustius ubi valvulae coarctatae evadunt concavae.
607. *Legumen*, pericarpium bivalve affigens semina secundum alteram tantum.
608. *Isthmis interceptum*, in varia loculamenta interne transversim distinctum.
609. *Folliculus*, pericarpium univalve altero latere longitudinaliter dehiscens, nec futurae semina affigens.
610. *Drupa* pericarpium factum e valve continens nucem.
604. *Succulenta*, humorem continens.
605. *Sicca* praecedenti opposita.
611. *Pomum*, pericarpium factum e valve continens capsulam.
612. *Bacca*, pericarpium factum e valve, semina ceteroquin nuda continens.
613. *Nidulantia*,

TERMINI BOTANICI.

613. *Nidulantia*, semina per pulpam sparsa.
614. *Strobilus*, ex Amento factum, squamis induratis.
615. **SEMEN**, rudimentum novæ plantæ.
616. *Hilum*, cicatrix externa seminis ab ejusdem affixione in fructu.
617. *Corculum*, primordium novæ plantæ intra semen.
618. *Corona Calyculus* supra adhærens, quo volitat.
619. *Pappus Corona* peniacea pilosave volitans.
620. *Stipitatus* filo elevante et adnectente pappum et semen.
621. *Capillaris*, pilis indivisis.
622. *Plumosus* pilis pennatis constans.
623. *Cauda* semen terminatum filo.
624. *Hamus*, quo adhæret animalibus.
625. *Calyculus tegmentum* seminis proprium.
626. *Nux* semine tecto epidermide osseo.
627. *Arillus*, tunica propria exterior seminis sponte secedens.
628. **RECEPTACULUM**, basis qua partes Fructificationis connectuntur.
629. *Commune* plures flores eorumque fructus continens.
630. *Punctatum*, punctis excavatis adpersum.
631. *Pilosum* (238).
632. *Paleaceum* lamellis innatis distinguens flosculos.
633. *Planum* (243).
634. *Convexum* (246).
635. *Conicum*, teres versus apicem attenuatum.
636. *Subulatum* (266).
637. *Compositus Flos*, receptaculo dilatato integro, Flosculis sessilibus.
638. *Aggregatus Flos*, receptaculo dilatato, Flosculis subpetiolatis.
639. *Umbellia*, receptaculum ex eodem centro elongatum in pedunculos filiformes proportionatos.
640. *Simplex*

640. *Simplex*, pedunculis omnibus ex uno eodemque receptaculo ortis.
641. *Composita* omnibus pedunculis apice umbellulas gerentibus.
642. *Universalis*, ex pluribus simplicibus facta composita.
643. *Partialis* umbellula pars universalis.
644. *Prolifera*, umbella plus quam decomposita.
645. *Cyma*, receptaculum ex centro eodem universalis, partialibus vero vagis, elongatum, in pedunculos fastigiatos.
646. *Rachis*, receptaculum filiforme fasciculos longitudinaliter adnectens in spicam.
647. *Spadix* receptaculum PALMÆ enatum intra spatham, in ramulos fructificantes divisum.
648. **BULBUS**, hybernaculum plantæ e rudimento foliorum præteritorum.
649. *Solidus* carnosus intus indivisus.
650. *Tunicatus*, bulbus tunica extra tunicas.
651. *Squamatus*, squamis imbricatis.
652. *Caulinus* cauli insidens.
- petiolaris*, e petiolorum rudimentis.
- stipularis* e stipulis confecta.
- corticalis* e corticis ramentis.
- foliaris*, folia continens, non flores.
- floralis*, flores non folia continens.
- communis*, et folia et flores continens.
653. **GEMMA**, hybernaculum plantæ ex rudimentis foliorum futurorum.
654. **VERNATO** foliorum dispositio intra gemmam.
655. *Conduplicata*, foliorum lateribus parallele approximatis.
656. *Convoluta*, cuculi in modum spiralis.
657. *Involuta*, lateribus versus paginam superiorem utrinque spiraliter contortis.
658. *Equitantia*, marginibus conniventia situ opposito, ut alterum includat alterum.
659. *Obvoluta*, pagina superiore lateribus approximatis

TERMINI BOTANICI.

proximatis, ita ut alterum latus distinguat alterum folium.

660. *Plicatum* in plicas varias coarctatum.

661. *Circinnale* in spiram transversalem coarctatum, ut apex centrum obtineat.

Mensura.

662. *Linearis*, longitudinaliter lunulæ, ab unguis radice versus unguem (non in Pollice).

663. *Unguicularis*, longitudine unguis.

664. *Pollicaris*, longitudine extimi articuli Pollicis.

665. *Palmaris*, longitudine manus transversæ.

666. *Spithameus*, spatium inter apicem Pollicis et indicis extensum.

667. *Dodrantalis*, spatium inter apicem Pollicis et digiti minimi extensum.

668. *Pedalis* a flexuosa cubitis ad basin Pollicis.

669. *Orgyalis*, longitudine hominis.

NOMINA.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

- Blackf. spec. bot.* Joannis Blackstone Specimen Botanicum. Lond. 1746. 8vo.
- Blackf. cat. pl.* Ejusdem Catalogus Plantarum circa Harefield. Lond. 1737. 8vo.
- Boch. Lips.* Georgii Rudolphi Boehmeri Flora Lipsiæ Indigena. Lips. 1750. 8vo.
- Boeh. ind.* Hermannii Boerhaave Index alter Plantarum. Lugdb. 1720. 4to.
- Bocc. mus.* Pauli Boctone Musæum Plantarum rariorum et Fificæ, &c. 2. Venet. 1697. 4to.
- Bocc. pl. rar.* Ejusdem Plantarum rariorum Siciliæ Icones et Descriptiones. Oxon. 1674. 4to.
- Breyn. cent.* Jacobi Breynii Centuria Plantarum Gedani. 1678. fol.
- Breyn. pr.* Ejusdem Prodrromus Fasciculus Plantarum. Ibid. 1689. 4to.
- Burm. Ger.* Nicolai Laurentii Burmanni Specimen Botanicum de Geraniis. Lugdb. 1759. 4to.
- Buxb. em.* Joannis Christiani Buxbaum Enumeratio Plantarum in Agro Hallensi. Halæ Magd. 1728. 8vo.
- Buxb. cent.* Ejusdem Plantarum minus cognitarum Centuriæ V. Petrop. 1728—1740. 4to.
- Budd. hort. sic.* Adamæ Budleii Hortus Siccus Plantarum Angliæ.
- Camer. epit.* Joachim Camerarii Epitome utilissima de Plantis. Franc. 1586.
- Cæsalp.* Andreas Cæsalpinus de Plantis Florent. 1583. 4to.
- Clus. hist.* Caroli Clusi Atrebatensis rariorum Plantarum Historia. Antw. 1601. fol.
- Col. eph.* Fabii Columnæ Ephraesis I. et II. minus cognitarum rariorumque Stirpium. Romæ. 1626. 4to.
- Col. phytob.* Ejusdem Phytobasanos. Neapoli. 1592. 4to.
- Comm. cat.* Joannis Commelini Catalogus Plantarum Hollandiæ. Amstel. 1685. 12mo.
- Cord. hist.* Valerii Cordii Historiæ Stirpium Liber IV. Argent. 1564. fol.
- Crantz. aust.* Henrici Joh. Nepomuceni Crantz Stirpium Austriacarum. 2. Vien. 1762. 4to. et 8vo.
- Daleb. hist.* Jacobi Dalechampii Historia generalis Plantarum. Lugdb. 1584. fol.

- Daleb. Paris.* Dionys. Dalibordi Floræ Parisiensis prodromus. Paris. 1750. 8vo.
- Deer. nott.* Caroli Deering Catalogus Plantarum circa Nottinghamiam. Nottingh, 1738. 8vo.
- Dill. giss.* Joannis Jacobi Dillenii Catalogus Plantarum circa Gissam. Gissæ. 1719. 8vo.
- Dill. eth.* Ejusdem Hortus Elthamensis. 2. Lond. 1732. fol.
- Dill. musc.* Ejusdem Historia Muscorum. Oxon. 1741. 4to.
- Dod. pempt.* Remberti Dodonæi Stirpium Historiæ Pemptades sex s. Libri xxx. Antw. 1616. fol.
- Fabric. en. pl.* Philippi Conradi Fabricii Enumeratio Methodica Plantarum Horti Medici Helmstadiensis. Helmst. 1763. 8vo.
- Fl. dan.* Flora Danica. Havnæ. 1761, &c.
- Fuchs. hist.* Leonardi Fuchsii Historia Plantarum. Basl. 1542.
- Ger. em.* Joannis Gerardi Historia Plantarum, a Thoma Johnson Emaculata. Lond. 1633. et 1636. fol.
- Ger. gallopr.* Ludovici Gerardi Flora Gallo-provincialis. Paris. 1761. 8vo.
- Gesn. ic.* Conradi Gesneri Opera Botanica. Neuremb. 1752. fol.
- Gled. fung.* Joannis Gottlieb Gleditchii Methodus fungorum. Berol. 1753. 8vo.
- Gmel. sib.* Joannis Georgii Gmelini Flora Siberica. 4. Petrop. 1750—1769. 4to.
- Gmel. fac.* Samuelis Gottlieb Gmelini Historia Fucorum. Petrop. 1768. 4to.
- Gmel. enum.* Joannis Frederici Gmelini enumeratio Stirpium Agro Tubingensi Indigenarum. Tubing. 1776. 8vo.
- Gort. Belg.* Davidi De Gorteri Flora Belgica. Traj. ad Rhenum. 1767. 8vo.
- Gouan. monsp.* Antonii Gouan Flora Mospeliaca. Lugd. 1765. 8vo.
- Gouan. Hort. monsp.* Ejusdem Hortus Regius Mospeliensis. Lugd. 1766. 8vo.
- Gron. virg.* Joannis Fred. Gronovii Flora Virginica. 2. Leid. 1739. 8vo.
- Guett.*

- Guett. stamp.* D. Guettard Observations sur les Plantes.
2. Paris. 1747. 8vo.
- Gunn. Norveg.* J. Ernesti Gunneri Flora Norvegica.
Nidros, 1766. fol.
- Hall. hist.* Alberti v. Haller Historia Stirpium Indigenarum Helvetiæ. 3. Bern. 1768. fol.
- Hall. nomencl.* Ejusdem Nomenclator. Bern. 1769. 8vo.
- Hall. opusc.* Ejusdem Opuscula Botanica. Gotting.
1749. 8vo.
- Hall. orch.* Ejusdem Orchidum Classis Constituta.
- Herm. parad.* Pauli Hermanni Paradisus Batavus,
Lugd. 1698.
- Hist. Ox.* Roberti Morisoni et Jacobi Bobarti Historia Plantarum Oxoniensis. 3. Oxon. 1680. fol.
- Hunt. Ev. sylv.* Sylva, or a Discourse of Forest-Trees, by John Evelyn, Esq. with Notes by A. Hunter, M.D. F.R.S. York. 1776. 4to.
- Jacq. inum.* Nicolai Jacquin Enumeratio Stirpium in agro Vindobonensis sponte nascentium. Vind. 1762. 8vo.
- Jacq. obs.* Ejusdem Observationes Botanicae. Vind. 1764, &c. fol.
- Jacq. austr.* Ejusdem Floræ Austriacæ sive Plantarum selectarum in Austriæ Archiducatu sponte crescentium Icones. 4. Vindob. 1773. fol.
- Jacq. hort. vindob.* Ejusdem Hortus Botanicus Vindobonensis. 2. Vindob. 1770. fol.
- Johns. merc. bot. et it.* Thomæ Johnsoni Mercurius et Itinera Botanica. Lond. 1632, 1634. 8vo.
- Kram elench.* Guil. Henrich. Kramer Elenchus Vegetabilium et Animalium Austriæ inferioris. Vien. 1756. 8vo.
- Krasch. Ingr.* Stephani Krascheninnikof Flora Ingrica. Peterfb. 1761. 8vo.
- Leyf. hal.* Frederici Wilhelm. Leyser Flora Halensis. Hal. 1761. 4to.
- Lin. gen. pl.* Caroli Linnæi Genera Plantarum. Holm. 1764.
- Lin. sp. pl.* Ejusdem Species Plantarum. 2. Holm. 1762.
- Lin. syst. nat.* Ejusdem Systema Naturæ. 3. Holm. 1766, &c.
- Lin. mantiss.* Ejusdem Mantissa Plantarum, ad finem Tomi

- Tomii 2di Systematis Naturæ et Mantissa altera Plantarum. Holm. 1771. 8vo.
- Lin. Fl. lapp.* Ejusdem Flora Lapponica. Amstel. 1737. 8vo.
- Lin. Fl. suec.* Ejusdem Flora Suecica. Stockh. 1755. 8vo.
- Lin. Hort. cliff.* Ejusdem Hortus Cliffortianus. Amstel. 1737. fol.
- Lin. virid. cliff.* Ejusdem Viridiarum Cliffortianum. Amstel. 1737. 8vo.
- Lin. hort. upf.* Ejusdem Hortus Upsaliensis. Holmiæ. 1748. 8vo.
- Lin. mat. med.* Ejusdem Materia Medica. Holmiæ. 1749. 8vo.
- Lin. it. oeland.* Ejusdem Iter Oelandicum. Holmiæ. 1745. 8vo.
- Lin. it. W-goth.* Ejusdem Iter Westrogothicum. Holmiæ. 1745. 8vo.
- Lin. it. scan.* Ejusdem Iter Scanicum. Holmiæ. 1751. 8vo.
- Lin. Amœn. acad.* Ejusdem Amœnitates Academicæ. I. Lips. 1749. II. Holm. 1751. III. Amstel. 1756. IV—VII. Lugdb. 1760, 1764, 1769. 8vo.
- Lob. ic.* Matthiæ Lobelii Icones Plantarum. Antw. 1580. 4to.
- Lob. ob. et adv.* Ejusdem Adversaria et Observationes. Lond. 1605.
- Loef. it.* Petri Loefling Iter Hispanicum. Stockh. 1758. 8vo.
- Loef. pruss.* Joannis Loeselii Flora Prussica. Regimont. 1703. 4to.
- Lyons fasc.* Israelis Lyons Fasciculus Plantarum circa Cantabrigiam. Lond. 1763. 8vo.
- Magn. bot. monsp.* Petri Magnol Botanicum Monspeliense. Monspel. 1686. 8vo.
- Magn. hort. reg. monsp.* Ejusdem Hortus Regius Monspeliensis. Monspel. 1697. 8vo.
- Mart. cent.* Joannis Martyn Centuria Plantarum. Lond. 1728. fol.
- Mart. tourn.* Ejusdem Tournefort's History of Plants. 2. Lond. 1732.
- Mart. plant. Cant.* Thomæ Martyn Plantæ Cantabrigiæ. Lond. 1763. 8vo.

NOMINA AUCTORUM EXPLICATA.

- Meez. frisi* Davidi Meeze Flora Frisica. Frank. 1760. 8vo.
- Mentz. pug.* Christiani Mentzelii Index Nominum Plantarum multilinguis et Pugillus rariorum Plantarum. Berol. 1682. fol.
- Merr. pin.* Christophori Merret Pinax rerum Naturalium Britannicarum. Lond. 1667. 8vo.
- Mich. gen.* Petri Antonii Michellii Nova Plantarum Genera. Florent. 1729. 4to.
- Mont. prod.* Josephi Monti Catalogi Stirpium Agri Bononiensis Prodomus. 1719. 4to.
- Mull. frid.* Ot. Frederici Muller Flora Friderichsdalina. Argent. 1767. 8vo.
- Murr. gctt.* Joannis Andreæ Murray Prodomus Designationis Stirpium Gottingensium. Gottingæ. 1770.
- Nec. delic. gallob.* Natalis Josephi de Necker Deliciæ Gallobelgice sylvestres. 2. Argent. 1748. 8vo.
- Nec. musci* Ejusdem Methodus Muscorum. Mannh. 1771. 8vo.
- Pall. elench.* P. S. Pallas Elenchus Zoophytorum. Hagæ. 1766. 8vo.
- Park. th.* Joannis Parkinsoni Theatrum Botanicum. Lond. 1640. fol.
- Park. par.* Ejusdem Paradisus terrestris. Lond. 1629. fol.
- Penn. Tour.* A Tour in Scotland, and Voyage to the Hebrides, by Thomas Pennant. 2. Chester. 1774. 8c. 4to.
- Pet. herb. brit.* Jacobi Petiveri Herbarium Britannicum. Lond. 1702. fol.
- Pet. conc. gr.* Ejusdem Concordia Graminum. Lond.
- Pet. gaz.* Ejusdem Gazophylaceum. 2. Lond. 1711. fol.
- Pet. mus.* Ejusdem Musæum. Lond. 1703. 8vo.
- Pluk. alm. et phyt.* Leonardi Pluknetii Almagestum Botanicum et Phytographia. Lond. 4to. diverso tempore edita.
- Phyt. br.* Phytologia Britannica. Lond. 1650.
- Pont. Julii.* Pontideræ Compendium Tabularum Botanicarum. Patavii. 1720. 4to.
- Raii. syn.* Joannis Raii Synopsis Methodica Stirpium Britannicarum, ed. I. 1690. II. 1696. III. Lond. 1724. 8vo.

- R. cat. pl. angl.* Ejusdem Catalogus Plantarum Angliæ. Lond. 1688. 12mo.
- R. bist.* Ejusdem Historia Plantarum. 3. Lond. 1686. et 1704. fol.
- R. meth.* Ejusdem Methodus Plantarum emendata et aucta. Lond. 1703. 8vo.
- Renealm. spec.* Pauli Renealmi Specimen Historiæ Plantarum. Paris. 1611. 4to.
- Riv. irr. mon. ic.* Augusti Quirini Rivini Ordines Plantarum Flore Irregulari Monopetalo, Tetrapetalo et Pentapetalo. Lipsiæ 1691, et 1699. fol.
- Royen. Leyd.* Adriani Van Royen Floræ Leydenfis Prodromus. Lugdb. 1740. 8vo.
- Rupp. Jen.* Henrici Bernardi Ruppil Flora Jenensis. Jenæ. 1745. ed. Haller. 8vo.
- Schæff. fung.* Jacobi Christiani Schæfferi Fungorum qui in Bavaria et Palatinatum circa Ratisbonam nascuntur Icones. 4. Ratisb. 1762, &c. 4to.
- Sauv. monsp.* Fran. Sauvages Flora Monspeliensis. Hagæ. 1751. 8vo.
- Sch. agr.* Joannis Jacobi Scheuchzeri Agrostographia Tigurii. 1719. et 1775. 4to.
- Sch. it. alp.* Ejusdem Itinera Alpina. 2. Lond. 1708. 4to.
- Schmid. icon.* Cassimir Christophori Schmidel Icones Plantarum et Analyses æri incisæ. Norimb. 1748. fol.
- Schmid. jung.* Ejusdem de Jungermannia. 1761. 4to.
- Schreb. gram.* Joannis Christiani Schreber Descriptiones et Icones Graminum. Lips. 1769. fol.
- Schreb. spir.* D. Joannis Christiani Dan. Schreberi Spicilegium Floræ Lipsicæ. Lips. 1771. 8vo.
- Scop. carn.* Joannis Antonii Scopoli Flora Carniolica. 2. Vindob. 1772. 8vo.
- Seg. pl. Veron.* J. Franc. Segulier Plantæ Veronenses. 3. Veronæ. 1745. 8vo.
- Sib. scot.* Roberti Sibbaldi Scotia Illustrata. Edinb. 1684. fol.
- Spig. isag.* Andreae Spigellii Isagoge in Rem Herbariam. Lugdb. 1633.
- Sterb. fung.* Francisci Van Sterbeck Theatrum Fungorum. Antw. 1675. 4to.

- Still. misc. Tract.* Benjamin Stillingfleet's Miscellaneous Tracts. Lond. 1762. 8vo.
- Tab. ic.* Jacobi Theodori Tabernæmontani Icones Plantarum. Franc. 1690. 4to.
- Thal. herc.* Joannis Thalii Hercynia 'Sylvia. Franc. 1588. 4to.
- Tourn. inst.* Josephi Pitton Tournefort Institutiones Rei Herbariæ. 3. Paris. 1700. 4to.
- Trag. hist.* Hieronymi Tragi Stirpium Historia. Argentor. 1552. 4to.
- Turn.* Gulielmi Turneri Historia Plantarum. Lond. 1551. fol.
- Vaill. paris.* Sebastiani Vaillant Botanicon Parisiense. Lugdb. et Amstel. 1727. fol.
- Volk. Norib.* Joannis Georgii Volckameri Flora Noribergensis. Noribergæ. 1700. 4to.
- Warn. pl. Woodf.* Richardi Warner Plantæ Woodfordienses. Lond. 1771. 8vo.
- Weis. crypt.* D. Friderici Gulielmi Weis Plantæ Cryptogamiæ Flora Gottingensis. Gotting. 1770. 8vo.
- Wulff. boruss.* Christoph. Wulff Flora Borussica Reigim. 176 . 8vo.
- Zan. hist.* Giacom. Zanoni Historia Botanica. Bologn. 1675, 1742. fol.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

MONANDRIA MONOGYNIA.

HIPPURIS. *Lin. Gen. pl. 11.*

Calyx nullus. Petala nulla. Stigma simplex. Semen unicum.

vulgaris. 1. HIPPURIS. Lin. Fl. lapp. 1. suec. 2. dan. 87. Sp. pl. 6.

Limnopeuce. Cord. hist. 150. R. syn. 136.

Equisetum palustre, brevioribus foliis, polyspermum.

Bauh. pin. 15.

Equisetum palustre alterum, brevioribus foliis. Park. 1200.

Cauda equina foemina. Ger. em. 1114.

Anglis, Mare's-tail.

Habitat in fontibus, paludibus, solo profundo limoso. In stagnis prope Hornsey. N. V. V.

Classis II.

D I A N D R I A

M O N O G Y N I A.

* Flores inferi, monopetali, regulares.

LIGUSTRUM. *Lin. Gen. pl.* 18.

Corolla quadrifida. *Bacca* tetrasperma.

1. LIGUSTRUM. *Ger. em.* 1394. *Park.* 1446. *vulgare.*

R. syn. 465. *Fl. suec.* 5. *Spec. pl.* 10. *germanicum.* *B. pin.* 475.

Anglis, Privet, or Prim.

Habitat in sylvis et sepibus. *h.* V.

** Flores inferi, monopetali, irregulares. *Fruetus* capsulares.

V E R O N I C A. 25.

Cor. quadrifida rotata, lacinia infima angustiore.

Capsula bilocularis.

* *Spicatae.*

1. VERONICA spica terminali, foliis oppositis crenatis *spicatae*, obtusis; caule ascendente simplicissimo. *Sp. pl.* 14. *Fl. suec.* 10. *Fl. dan.* 52.

Veronica foliis ellipticis serratis, conjugatis, floribus spicatis. *Hall. belv.* 542.

Veronica spicata minor. *B. pin.* 247. *R. syn.* 279.

Veronica recta minima. *Ger. em.* 627.

Veronica recta angustifolia. *Park.* 550.

2 *Veronica* spica terminali, foliis oppositis obtuse serratis hybrida-scabris, caule erecto. *Sp. pl.* 14. *Fl. suec.* 11.

Veronica spicata cambrobritannica, bugulae subhirsuto folio. *R. syn.* 278. *t.* 11.

DIANDRIA MONOGYNIA.

Anglis, spiked Speedwell. *β* welsh Speedwell.

Habitat in pratis et pascuis montosis. In several closes adjoining to Newmarket heath, beyond Bottesham.

R. syn. in pascuis circa Horseheath Hall, in comitatu Cantabrigienti. D. Miller, prope Snowdon in Wallia. *β* Ad latera montis cujusdam, Craig Wreidhin dicti in comitatu Montis Gomerici Walliæ, D. Lhwyd invenit. 4. VI. VIII.

officinalis. 2. VERONICA spicis lateralibus pedunculatis, foliis oppositis, caule procumbente. *Sp. pl.* 14. *Fl. dan.* 248.

Veronica mas supina et vulgatissima. *B. pin.* 246. *R. syn.* 281.

Veronica vera et major. *Ger. em.* 626.

Veronica mas vulgaris supina. *Park.* 550.

Anglis, male Speedwell.

Habitat in pascuis sterilibus et ericetis. 4. V. VI.

Pharm. Veronicæ Herba.

** Corymbo-racemosa.

fruticulo- 3. VERONICA corymbo terminali, foliis lanceolatis
sa. obtusiusculis, caulibus fruticulosus. *Sp. pl.* 15. *Fl. dan.* 34?

Veronica caulibus fruticosis declinatis, foliis glabris ellipticis, dentatis, floribus in summitate confertis, petiolatis. *Hall. hist.* 545. t. 16?

Veronica tertia fruticans. *Clus. hist.* 347? *Park.* 551?

Veronica frutescens serpyllifolia. *Ger. em.* 628?

Veronica alpina frutescens. *B. pin.* 247? *Hist. Ox.* II. 318. j. 3. t. 22. f. 5?

Veronica saxatilis, *B. hist.* III. 284? *R. hist.* 848? *Anglis*, shrubby Speedwell.

Habitat in montibus, in monte Ben Nevis dicto in Scotia. &c. 4. V. VI.

serpyllisfo- 4. VERONICA racemo terminali subspicato, foliis
lia. ovatis glabris crenatis. *Sp. pl.* 15. *Fl. dan.* 492.

Veronica caule recto, foliis ovatis glabris crenatis, petiolis ex alis unifloris, brevissimis. *Hall. hist.* 546.

Veronica

DIANDRIA MONOGYNIA.

5

Veronica pratensis serpyllifolia. *B. pin.* 247.
Veronica pratensis minor. *Park* 551. *R. syn.* 279.
Veronica minor. *Ger. em.* 627.
Anglis, smooth Speedwell, or Paul's Betony.
Habitat in pratis et pascuis. 4. V. VI.

5. *VERONICA* racemis lateralibus, foliis ovatis planis, *becabung-*
 caule repente. *Sp. pl.* 16. *Mat. med.* 12. *Fl. ga.*
dan. 511.

Veronica aquatica rotundifolia, *Becabunga dicta mi-*
nor. *R. syn.* 280.

Anagallis aquatica minor, folio subrotundo. *B. pin.*
 252.

Anagallis aquatica vulgaris, f. *Becabunga.* *Park.*
 1236.

Anagallis sive Becabunga. *Ger. em.* 620.

Anglis, common Speedwell, or Brooklime.

Habitat in fossis et rivulis. 4. VI.

Pharm. *Becabungæ herba.*

6. *VERONICA* racemis lateralibus, foliis lanceolatis *anagallis.*
 serratis, caule erecto. *Sp. pl.* 16. *Fl. suec.* 13. Δ.

Veronica foliis lanceolatis, serratis, glabris, ex alis
racemosa. *Hall. hist.* 533.

Veronica aquatica longifolia media. *R. syn.* 280.

Anagallis aquatica minor folio oblongo. *B. pin.* 252.

Anagallis aquatica folio oblongo crenato. *Park* 1237.

Anagallis aquatica major. *Ger. em.* 620.

Anglis, water Speedwell.

Habitat in fossis et rivulis. 4. VII.

7. *VERONICA* racemis lateralibus, pedicellis pendulis, *scutellata.*
 foliis linearibus integerrimis. *Sp. pl.* 16. *Fl. dan.*
 209.

Veronica foliis lineari-lanceolatis; racemis ex alis
paucifloris. *Hall. hist.* 532.

Veronica aquatica angustifolia minor. *R. syn.* 280.

Veronica aquatica angustifolia scutellata. *B. pin.* 252.

Anglis, marsh Speedwell.

Habitat in palustribus. 4. VI.

*chamæ-
drys.*

8. VERONICA racemis lateralibus, foliis ovatis rugosis dentatis sessilibus, caule debili. *Sp. pl.* 17.

Fl. dan. 448.

Veronica foliis cordatis, subrotundis, hirsutis, nervosis, ex alis racemosa. *Hall. hist.* 536.

Chamædrys sylvestris. *Ger. em.* 657. *R. syn.* 282.

Chamædrys spuria sylvestris. *Park.* 107.

Chamædrys spuria minor rotundifolia. *B. pin.* 249.

Anglis, wild Speedwell or Germander.

Habitat in pratis et pascuis. 4. V.

montana.

9. VERONICA racemis lateralibus paucifloris, calycibus hirsutis, foliis ovatis rugosis crenatis petiolatis, caule debili. *Sp. pl.* 17. *Jac. austr.* II. 6. t. 109.

Veronica caule procumbente, foliis hirsutis, cordatis, racemis paucifloris. *Hall. hist.* 539.

Veronica chamædryoides, foliis pediculis oblongis insidentibus. *R. syn.* 281.

Chamædrys spuria affinis rotundifolia scutellata. *Bauh. pin.* 242.

Anglis, mountain Speedwell. *Pet. H. Br.* t. 51. f. 4.

Habitat in sylvis et sepibus opacis. In Charlton Wood, the lower part plentifully. *Mr. J. Sherard.* *R. syn.*

4. V. VI.

*** *Pedunculis unifloris.*

arvensis.

10. VERONICA floribus solitariis, foliis cordatis incisiss sessilibus. *Fl. angl. o. dan.* 515.

Veronica floribus solitariis, foliis cordatis incisiss, pedunculo longioribus. *Sp. pl.* 18. *Fl. suec.* 19.

Veronica caule erecto, foliis ovatis, subhirsutis, dentatis, petiolis brevissimis. *Hall. hist.* 548.

Veronica flosculis singularibus, cauliculis adhærentibus. *R. syn.* 279.

Alsine veronicæ foliis, flosculis cauliculis adhærentibus. *Bauh. pin.* 250.

Alsine foliis veronicæ. *Ger. em.* 613.

Alsine foliis subrotundis veronicæ. *Park.* 762.

Anglis,

DIANDRIA MONOGYNIA,

7

Anglis, corn Speedwell, or Chickweed.

Habitat in arvis et muris antiquis. ☉. V.

11. VERONICA floribus solitariis, foliis cordatis in-agrestis. citis, petiolatis, *Fl. angl.* 6. *dan.* 449.

Veronica floribus solitariis, foliis cordatis incis, pedunculo brevioribus. *Sp. pl.* 18. *Fl. suec.* 20.

Veronica floribus singularibus, in oblongis pediculis chamædryfolia. *R. syn.* 279.

Veronica caule procumbente, foliis petiolatis, ovatis, crenatis. *Hall. hist.* 549.

Alfne foliis Triflaginis. *Ger. em.* 616. *Park.* 764.

Alfne chamædryfolia flosculis pediculis oblongis insidentibus. *Baub. pin.* 250.

Anglis, Germander-Speedwell, or Chickweed.

Habitat in arvis, ruderatis, cultis frequens. ☉. V.

12. VERONICA floribus solitariis, foliis cordatis hederifol. planis quinquelobis. *Sp. pl.* 19. *Fl. suec.* 21. *dan.* liq. 428.

Veronica flosculis singularibus, Hederulæ folio, Morsus gallinæ minor dicta. *R. syn.* 280.

Alfne Hederulæ folio. *B. pin.* 250. minor. *Park.* 760.

Alfne hederacea. *Ger. em.* 616.

Anglis, Ivy-leaved Speedwell, or small Henbit.

Habitat in agris, ruderatis, cultis frequens. ☉. IV. V.

13. VERONICA floribus solitariis, foliis digitato-par-triphyll. titis pedunculo brevioribus. *Sp. pl.* 19. *Fl. dan.* 627.

Veronica foliis ovatis, tripartitis et quinquepartitis, floribus longe petiolatis. *Hall. hist.* 551.

Veronica flosculis singularibus, foliis laciniatis, erecta. *R. syn.* 280.

Alfne triphylls cærulea. *Baub. pin.* 250.

Alfne triphylls s. laciniata. *Park.* 760.

Alfne recta. *Ger. em.* 612.

Anglis, trifold Speedwell.

Habitat in arvis arenosis; at Rowton in Norfolk, betwixt the town and the highway; and at Methwold in Suffolk, between the two windmills and the

DIANDRIA MONOGYNIA.

warren-lodge, &c. T. Willifel. R. syn. in arvis prope Rossington in comitatu Eboracensi. D. To-field. ☉. IV. V.

- verna.* 14. VERONICA floribus solitariis, foliis digitato-partitis pedunculo longioribus. *Sp. pl.* 19.
Veronica foliis ovatis, tripartitis et quinquepartitis, floribus brevissime petiolatis. *Hall. hist.* 552.
Anglis, vernal Speedwell.
Habitat in arvis arenosis, circa Bury in comitatu Suffolk. D. Cullum. ☉. IV.

PINGUICULA. 30.

- Cor. ringens, calcarata. Cal. bilabiatus, 5-fidus. Caps. unioocularis.*

- vulgaris.* 1. PINGUICULA nectario cylindraco longitudine petali. *Fl. lapp.* 11. *succ.* 25. *dan.* 93. *Sp. pl.* 25.
Pinguicula calcare floris longitudine. *Hall. hist.* 292.
Pinguicula Gesneri. *B. hist.* 111. 546. *R. syn.* *281.
Pinguicula s. Sanicula Eboracensis. *Park.* 532. *Ger. em.* 788.
Sanicula montana, flore calcari donato. *B. pin.* 243.
Anglis, common Butterwort, or Yorkshire Sanicle.
Habitat in uiginosis in comitatibus Eboracensi, Westmorlandico, Cumberlandico, &c. 4. V.

- villosa.* 2. PINGUICULA scapo villoso. *Fl. lapp.* 13. t. 12. f. 2. *Sp. pl.* 25.
Pinguicula flore minore carneo. *R. syn.* *281.
Anglis, hairy Butterwort.
Habitat in palustribus in comitatibus Hantoniensi, Dorsetsiensi, Devoniensi, et Cornubiensi frequens. *Prope Air et in insula Lamash in Scotia. D. Hope. 4. VI. VII.*
Obs. Folia tenella, marginibus conniventibus: venis purpurascensibus.

UTRICULARIA. 31.

Cal. diphyllus, æqualis. Cor. ringens, calcarata. Caps. unilocularis.

- vulgaris.* 1. UTRICULARIA nectario conico, scapo paucifloro. *Sp. pl.* 26. *Fl. succ.* 28. *dan.* 138.

Lentibularia.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

SALVIA. 39,

Cor. inæqualis. *Filamenta* transverse pedicello affixa.

- pratensis.* 1. SALVIA foliis cordato-oblongis crenatis: summis amplexicaulibus, verticillis subnudis, corollis galea glutinosis. *Sp. pl.* 35. *Fl. suec.* 32.
Sciarea pratensis foliis ferratis, flore cæruleo. *Tourn.* 179. *R. syn.* 237.
Horminum pratense, foliis ferratis. *Bauh. pin.* 238.
Horminum sylvestre vulgare. *Park.* 55.
Horminum sylvestre Fuchsii. *Ger. em.* 769.
Anglis, meadow Sage, or Clary.
Habitat in septo Cobhamiano in Essexia; à D. Watso observatus. 4. VI. VII.

- verbena-* 2. SALVIA foliis ferratis læviusculis, corollis calyce
ca. angustioribus, *Sp. pl.* 35. *Dalib. paris.* 9.
Horminum sylvestre. *Ger. em.* 771.
Anglis, wild Sage, or Clary.
Habitat in pascuis et pratis frequens. 4. V. VI,
Pharm. Hormini semina.

*** Flores superi.

CIRCÆA. 24.

Cal. diphyllus, superus. *Cor.* dipetala. *Sem.* 1. biloculare.

- lutetiana.* 1. CIRCÆA caule erecto, racemis pluribus, foliis ovatis. *Syst. nat.* 56. *Sp. pl.* 12. *Fl. dan.* 210.
Circæa lutetiana. *Lob. ic.* 266. *Ger. em.* 351. *R. syn.* 289. major. *Park* 351.
Solanifolia Circæa dicta major. *B. pin.* 168.
Anglis, common Enchanters-Nightshade.
Habitat in locis umbrosis humidis et ad sepes. 4. VI. VII.

- alpina.* 2. CIRCÆA caule prostrato, racemo, subunico, foliis cordatis. *Syst. nat.* 56. *Sp. pl.* 12. *Fl. suec.* 7. *Fl. dan.* 256.
Circæa foliis cordatis, acute dentatis. *Hell. hist.* 814.
 Circæa

DIANDRIA MONOGYNIA.

11

Circæa lutetiana minor. *Park.* 351.

Circæa minima. *Column. Ecphr.* 2. p. 79. t. 80.

Solanifolia Circæa alpina. *B. pin.* 168.

Anglis, mountain Enchanters-Night shade.

Habitat ad radices montium in comitatibus Westmorelandico, Eboracensi, &c. circa Dallam Tower in agro Westmorlandico. In Hamilton-wood in Scotland.
D. Hope. 4. VI. VII.

DIGYNIA.

ANTHOXANTHUM. 42.

Cal. Gluma bivalvis. uniflora. *Cor.* Gluma bivalvis, acuminata. *Sem.* unicum.

1. ANTHOXANTHUM spicâ oblonga, flosculis sub-odoratum, pedunculatis arista longioribus. *Sp. pl.* 40. *Still. misc.* t. 1. *Schreb. gram.* 49. t. 5.

Gramen vernum spica brevi laxa. *R. syn.* 398.

Gramen pratense spica flavescente. *B. pin.* 3. *Tb.* 44. *Sch. agr.* 88.

Gramen Anthoxanthon spicatum. *B. hist.* II. 466.

Gramen alopecurum vernum, spica flavescente. *Hist.*

Ox. III. p. 193. f. 8. t. 7. f. 25.

Anglis, Vernal, or Spring Grass.

Habitat in pratis et pascuis. 4. V.

Classis

Classis III.

T R I A N D R I A
M O N O G Y N I A.

* *Flores superi.*

V A L E R I A N A. 44.

Cal. nullus. *Cor.* monopetala, basi hinc gibba, supera.
Sem. unicum.

*? *rubra.* 1. VALERIANA floribus monandris caudatis, foliis
lanccolatis integerrimis. *Sp. pl.* 44.

Valeriana foliis glaberrimis, floribus calcaratis. *Hall.*
hist. 213.

Valeriana rubra. *Dod. purg.* 91. *B. pin.* 165. *Ger.*
em. 678. *Park.* 122.

Anglis, red Valerian.

Habitat in muris antiquis et ruderatis, in Devoniam et
Cornubiam frequens. 4. V.—VIII.

officinalis. 2. VALERIANA floribus triandris, foliis omnibus
pinnatis. *Sp. pl.* 45. *Fl. succ.* 34. *dan.* 57.

Valeriana sylvestris major. *Ger. em.* 1275. *Park.*
122. *B. pin.* 164. *R. syn.* 200.

β Valeriana sylvestris major montana. *B. pin.* 164.

Anglis, wild Valerian.

Habitat in sepibus et nemoribus paludosis frequens. 4. VI.

Pharm. Valerianæ Radix.

dioica. 3. VALERIANA floribus triandris dioicis, foliis radi-
calibus ovatis, caulinis pinnatis. *Fl. angl.* 12.

Valeriana floribus triandris dioicis, foliis pinnatis in-
tegerrimis. *Sp. pl.* 44. *Fl. succ.* 35.

Valeriana foliis radicalibus petiolatis ovatis; caulinis
pinnatis, sexu distincta. *Hall. hist.* 208.

Valeriana palustris minor. *B. pin.* 164.

Valeriana

- Valeriana sylvestris minor. *Park.* 122. *R. syn.* 200.
 Valeriana minor. *Ger. em.* 1075. *Riv. mon.* 3.
 β Valeriana sylvestris f. palustris minor altera. *R. syn.*
 200.
Anglis, marsh Valerian.
 Habitat in pascuis humidis. 4. VI.

4. VALERIANA floribus triandris, caule dichotomo, *locusta*.
 foliis linearibus. *Sp. pl.* 47. *Fl. suec.* 36.
 Valerianella foliis oblongis, rariter incisus, corona fe-
 minis simplici, acuminata. *Hall. hist.* 214.
 Valerianella arvensis præcox humilis semine com-
 pressa. *Mor. Umb. R. syn.* 201.
 Valeriana campestris inodora major. *B. pin.* 165.
 Lactuca agnina. *Ger. em.* 310. *Park.* 812.
 β Valerianella vulgaris species major serotina. *Mor.*
præhud. 319.
 γ Lactuca agnina f. valerianella foliis serratis. *R. syn.*
 201.
Anglis, corn Valerian or Sallet, Lambs-Lettuce.
 Habitat in arvis et agris. ☉. IV.—VIII.

C R O C U S. 55.

- Cor.* sexpartita, æqualis. *Stigmata* convoluta.
 1. CROCUS spatha univalvi radicale, corollæ tubo *officinalis*.
 longissimo. *Sp. pl.* 50.
 Crocus autumnalis foliis angustioribus margine revo-
 lutis. *Syst. natur.* 75.
 Crocus. *Ger. em.* 151. Sativus. *B. pin.* 65.
 β Crocus vernalis foliis latioribus margine patulo. *Syst. sylvestris.*
nat. 75.
 Crocus tuba brevissima trifida. *Hall. hist.* 1257.
 Crocus vernus. *Ger. em.* 153. latifolius purpureus
 variegatus. *B. pin.* 65.
Anglis, saffron. β Crocus.
 Habitat in agro Cantabrigiensi et Essexiensi; Saffron-
 walden oppidum editur. *R. syn.* in pratis et pascuis
 passim. 4. VIII.
Pharmac. Croci stigma.

TRIANDRIA MONOGYNIA.

IRIS. 59.

Cor. 6-partita. Petalis alternis reflexis. Stigmata petaliformia.

*pseud-
acorus.*

1. IRIS corollis imberbibus; petalis interioribus stigmatate minoribus, foliis ensiformibus. *Sp. pl.* 56. *Fl. dan.* 494.

Iris palustris lutea. Ger. em. 50. *R. syn.* 374.

Acorus adulterinus. B. pin. 74.

Acorus palustris, sive Pseudo-iris et Iris lutea palustris. Park. 1219.

β *Iris palustris pallida. R. syn.* 375.

Anglis, yellow Iris or Flower-de-luce.

Habitat in udis, fluviis, paludosis frequens. 4. VII.

Pharm. Acori palustris Radix.

*fœtidissi-
ma.*

2. IRIS corollis imberbibus: petalis interioribus patentissimis, caule unangulato, foliis ensiformibus. *Sp. pl.* 57.

Gladiolus fœtidus. B. pin. 30. *R. syn.* 375.

Xyris. Ger. em. 60: s. *spatula fœtida, Park.* 256.

Anglis, stinking Iris or Gladwyn.

Habitat in sylvis et ad sepes, prope Hornsey, in sylva prope Charlton, in Cantio. 4. VI. VII.

Pharm. Iris fœtidæ Radix.

•• *Flores Graminci valvulis Glumæ calycinæ.*

SCHOENUS. 65.

Glumæ paliacæ, univalves, congestæ. Cor. 0. Sem. unicum subrotundum inter glumas.

* *Culmo tereti.*

mariscus. 1. **SCHOENUS** culmo tereti, foliis margine dorsoque aculeatis. *Fl. suec.* 38. *Sp. pl.* 62.

Cyperus longus inodorus sylvestris. Ger. em. 29. *Park.* 1263. *R. syn.* 426.

Cyperus longus inodorus germanicus. B. pin. 14.

Cyperus longus inodorus major, foliis et carina serratis. Hist. Ox. III. 237. t. 11. f. 24.

Pseudo Cyperus palustris, foliis et carina serratis. Sch. agr. 375.

Anglis,

Anglis, prickly Bog-rush or bastard Cyperus.

Habitat in paludibus; on Hinton-Moor, near Cambridge; near Tamworth, Warwickshire; near Pensans, Cornwall. R. syn. 4. VII. VIII.

2. SCHOENUS culmo tereti nudo capitulo ovato, nigricans, involucri diphylli valvula altera subulata longa.

Sp. pl. 64. *Fl. suec.* 39.

Cyperus spicis conglomeratis, calyce multifolio. *Hall. hist.* 1347.

Juncus lævis minor, panicula glomerata nigricante.

R. cat. 174. *syn.* 430. *Hist. Ox.* III. 243. t. 10. f. 28.

Junco affinis capitulo glomerato nigricante. *Sch. agr.* 349.

Anglis, black Bog-rush.

Habitat in paludibus caespitosis et turfosis. 4. VI.—VIII.

3. SCHOENUS culmo tereti nudo, spica duplici, inferrugivolucri valvula majore spicam æquante. *Sp. pl.* 64. *neus.*

Anglis, ferrugineus Bog-rush.

Habitat in paludibus turfosis, in Westmorlandia, in the island of Skye. D. Hope. 4. VII.

** Culmo triquetro.

4. SCHOENUS culmo subtriquetro folioso, spica distica, spiculis multifloris, involucri monophyllo. *compres-*

Schoenus culmo subtriquetro nudo, spica disticha, involucri monophyllo. *Sp. pl.* 65. *jus.*

Gramen cyperoides spica simplici compressa disticha.

Pluk. pb. t. 34. f. 9. *Sch. Agrest.* 490. *Hist. Ox.* III. 245. *R. syn.* 425.

Anglis, compressed Bog-rush.

Habitat in paludibus caespitosis in comitatibus Eboracensi, et Westmorlandico frequens; about Chislehurst in Kent. R. syn. 4. VI. VII.

5. SCHOENUS culmo subtriquetro folioso, spica sub-*ovata*, spiculis unifloris, involucri monophyllo. *ovata.*

Anglis, brown Bog-rush.

Habitat

Habitat in paludibus, in insula Skye et prope Glasgōw in Scotia. 4. VII.

Desc. Radix perennis, ramosa, horizontalis, repens. Culmo plures, semipedales, erecti, subtriquetri, basi foliati, simplices. Folia radicalia linearia, acuta, nuda, parum canaliculata, longitudine fere culmi; caulina duo, petiolata, linearia, acuta, culmo dimidio breviora. Petiolus vaginans, cylindricus, striatus, glaber. Spica terminalis, subovata, spiculæ quinque aut sex, sessiles, alternæ, imbricatæ, rusæ. Involucrum monophyllum, acutum, subpersistens, longitudine spicæ. Cal. Gluma uniflora, bivalvis: valvulis erectis, oblongis, concavis, persistentibus. Cor. Petala sex, lanceolata, acuta, membranacea calyce dimidio breviora. Stam. Filamenta tria, capillaria corolla breviora. Antheræ oblongæ, erectæ, flavæ. Pist. Germen ovatum, triquetrum. Stylus brevissimus. Stigmata tria, capillaria. Sem. unicum, triquetrum, acuminatum. affinis præcedenti.

albus.

6. SCHOENUS culmo subtriquetro folioso, floribus fasciculatis, foliis setaceis. *Sp. pl.* 65. *Fl. dan.* 320.

Scirpus foliis setaceis, panicula plana, locustis teretibus. *Hall. hist.* 1341.

Cyperus minor palustris hirsutus, paniculis albis paleaceis. *Hist. Ox. III.* 239. t. 9. f. 39. *R. syn.* 427.

Gramen cyperoides palustre, leucanthemum. *Park.* 1295. *Scb. agr.* 503.

Gramen junceum leucanthemum. *Ger. em.* 30.

fuscus.

β Schoenus culmo tereti folioso, spiculis fasciculatis, foliis filiformibus canaliculatis. *Sp. pl.* 1664.

Cyperella palustris, capitulis umbellatis, primum albis deinde fulvis. *Mich. gen.* 53.

Cyperus minor angustifolius palustris, capitulis fuscis, paliaceis. *Hist. Ox. III.* 239. t. 11. f. 40. *R. syn.* 427.

Anglis, white-headed Bog-rush.

Habitat in palustribus on Bagshot Heath, Mr. Doody; between Wickham and Croydon, Mr. Petiver.

R. syn. in comitatibus Cornubiensi, Hantoniensi, Eboracensi

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Habitat in paludibus, in insula Skye et prope Glasgōw in Scotia. 4. VII.

Desc. Radix perennis, ramosa, horizontalis, repens. Culmo plures, semipedales, erecti, subtriquetri, basi foliati, simplices. Folia radicalia linearia, acuta, nuda, parum canaliculata, longitudine fere culmi; caulina duo, petiolata, linearia, acuta, culmo dimidio breviora. Petiolus vaginans, cylindricus, striatus, glaber. Spica terminalis, subovata, spiculæ quinque aut sex, sessiles, alternæ, imbricatæ, rufæ. Involucrum monophyllum, acutum, subpersistens, longitudine spicæ. Cal. Gluma uniflora, bivalvis: valvulis erectis, oblongis, concavis, persistentibus. Cor. Petala sex, lanceolata, acuta, membranacea calyce dimidio breviora. Stam. Filamenta tria, capillaria corolla breviora. Antheræ oblongæ, erectæ, flavæ. Pist. Germen ovatum, triquetrum. Stylus brevissimus. Stigmata tria, capillaria. Sem. unicum, triquetrum, acuminatum. affinis præcedenti.

albus.

6. SCHOENUS culmo subtriquetro folioso, floribus fasciculatis, foliis setaceis. *Sp. pl.* 65. *Fl. dan.* 320.

Scirpus foliis setaceis, panicula plana, locustis teretibus. *Hall. hist.* 1341.

Cyperus minor palustris hirsutus, paniculis albis paleaceis. *Hist. Ox. III.* 239. t. 9. f. 39. *R. syn.* 427.

Gramen cyperoides palustre, leucanthemum. *Park.* 1295. *Scb. agr.* 503.

Gramen junceum leucanthemum. *Ger. em.* 30.

fuscus.

β Schoenus culmo tereti folioso, spiculis fasciculatis, foliis filiformibus canaliculatis. *Sp. pl.* 1664.

Cyperella palustris, capitulis umbellatis, primum albis deinde fulvis. *Mich. gen.* 53.

Cyperus minor angustifolius palustris, capitulis fuscis, paliaceis. *Hist. Ox. III.* 239. t. 11. f. 40. *R. syn.* 427.

Anglis, white-headed Bog-rush.

Habitat in palustribus on Bagshot Heath, Mr. Doody; between Wickham and Croydon, Mr. Petiver. *R. syn.* in comitatibus Cornubiensi, Hantoniensi, Eboracensi

Eboracensi, Westmorlandico, &c. copiose. 4. VII.
IX.

Obs. α. β. *Folia subsetacea parum canaliculata; Flores primum albi postea fusci.*

CYPERUS. 66.

Glumæ paliaciæ, distiche, imbricatæ. Cor. o. Sem. unicum nudum.

1. CYPERUS culmo triquetro folioso, umbella foliosa *longus*.
supradecomposita; pedunculis nudis, spicis alternis.

Sp. pl. 67. Mat. Med. 29.

Cyperus longus. Ger. em. 30. odoratus. Park. 145. R. Syn. 425.

Cyperus odoratus radice longa. f. Cyperus Off. B. pin. 14. Sch. agr. 378.

Anglis, sweet Cyperus or English Galingale.

Habitat in paludibus. Found by Mr. Newton in the Isle of Purbeck, Dorsetshire, near a chapel on the side that looks towards Portland Island. R. Syn. on St. Faith's Newton Bogs, near Norwich. Mr. Humphrys. 4. VII.

SCIRPUS 67.

Glumæ paliaceæ, undique imbricatæ. Cor. o. Sem. unicum imberbe.

* *Spica unica.*

1. SCIRPUS culmo tereti nudo, spica subovali termi-*palustris*.
nali. *Sp. pl. 70. Fl. Carn. 56. dan. 273.*

Scirpus equiseti capitulo, major. Tourn. 528. R. Syn. 429. Sch. agr. 360.

Juncus equiseti capitulis, major. Baub. pin. 12.

Juncus aquaticus, capitulis equiseti. Park. 1196.

Juncus minor, capitulis equiseti. Ger. em. 35.

Anglis, marsh Club-rush.

Habitat in fossis, paludibus, fluviiis, frequens. 4. VII.

2. SCIRPUS culmo striato nudo, spica bivalvi termi-*cestojus*.
nali longitudine calycis, radicibus squamula in-
terstinctis.

terstinctis. *Fl. Suec.* 43. *Sp. pl.* 71. *Fl. Dan.* t. 167.

Scirpus caule tereti aphylo, spica pauciflora acuminata, calycinæ glumæ longitudine. *Hall. hist.* 1334.

Scirpus montanus capitulo breviori. *Tourn.* 528. *Sch. agr.* 363. t. 7. f. 18. *R. Syn.* 429.

Juncus parvus palustris, cum parvis capitulis equiseti. *Pluk. ph.* t. 40. f. 6.

Anglis, dwarf Club-rush.

Habitat in paludibus cespitosis et ericetis udis. 4. VII.

acicularis. 3. SCIRPUS culmo quadrangulo nudo, spica ovata terminali.

Scirpus culmo tereti nudo setiformi, spica ovata bivali terminali, seminibus nudis. *Sp. pl.* 71. *Fl. dan.* 287.

Scirpus minimus capitulis equiseti. *Cat. Gifs.* 165. *R. Syn.* 429.

Juncellus omnium minimus, capitulis equiseti. *Hist. Ox. III.* 145. t. 10. f. 37. *Pluk. ph.* t. 40. f. 7.

Anglis, least Club-rush.

Habitat in palustribus; in ericeto Hounsejano versus Hampton, invenit D. Doody. *R. Syn.* 4. VIII.

fluitans. 4. SCIRPUS culmis teretibus nudis alternis, caule folioso flaccido. *Sp. pl.* 71.

Scirpus foliis linearibus planis alternatim fasciculatis, spica terminali. *Guett. stamp.* 141.

Scirpus equiseti capitulo minor. *Sch. agr.* 365. t. 7. f. 20. *R. Syn.* 431.

Juncellus, capitulis equiseti, minor fluitans. *B. pin.* 12.

Gramen junceum clavatum minimum. *R. Hist.* 1310. *Hist. Ox. III.* 230. t. 10. f. 31. *Pluk. ph.* t. 35. f. 1.

Anglis, floating Club-rush.

Habitat in fossis et paludibus, in stagnis prope Wandsworth, ericeto Putnejano et Hounsejano. 4. VII. VIII.

** *Culmo tereti polystachio.*

5. SCIRPUS culmo tereti nudo, spicis ovatis pluribus *lacustris*.
 - pedunculatis terminalibus. *Fl. suec.* 46. *Sp. pl.* 72.
 Scirpus caule tereti, panicula laterali ramosa, locustis
 ovatis. *Hall. hist.* 1337.
 Scirpus palustris altissimus. *Tourn.* 528. *R. Syn.*
 428. *Sch. agr.* 154.
 Juncus maximus f. Scirpus major. *B. pin.* 12. *th.* 178.
 Juncus lævis maximus. *Park.* 1191.
 Juncus aquaticus maximus. *Ger. em.* 35.
 β Juncus f. Scirpus medius. *B. pin.* 12. *R. Syn.* 428.
 • γ Juncus aquaticus major, caule carinato. *Dood. R.*
syn.
Anglis, tall Club-rush or Bull-rush.
 Habitat in stagnis fluviis et lacubus. 4. VII. VIII.
 Obs. In α. Involucrum submembranaceum panicula bre-
 vius, Spicæ oblongiusculæ; in β. Involucrum paniculæ
 subæquale, Spicæ ovatæ; in γ. unum latus culmi sub-
 planiusculum.
6. SCIRPUS culmo tereti nudo, spicis subglobosis glo- *Holoscæ-*
 meratis pedunculatis, involucro diphylo inæquali *nus.*
 mucronato. *Sp. pl.* 72. *Fl. dan.* 454.
 Scirpus maritimus, capitulis rotundioribus glomeratis.
Tourn. 528. *R. Syn.* 429. *Sch. agr.* 371.
 Juncus acutus maritimus, capitulis rotundis. *B.*
pin. 11.
 Juncus acutus maritimus alter. *Park.* 1196.
Anglis, round-headed Club-rush.
 Habitat in litoribus maritimis. Found by Mr. Stephens
 in Brounton Boroughs, in Devonshire; in Somers-
 setshire and Hampshire. *Pet. conc. Gr. R. Syn.*
 429. VII. VIII.
7. SCIRPUS culmo tereti nudo, capitulo laterali con- *romanus.*
 globato, bractea reflexa. *Sp. pl.* 72. *Fl. Gallopr.*
 116?

Scirpus capitulo singulari radice tomentosa. *Barr. rar.* 1249. t. 255. f. 3.

Scirpoides acutum maritimum, capitulo solitario. *Sch. agr.* 373. *Mich. gen.* 52.

Anglis, single-headed Club-rush.

Habitat in palustribus juxta Throgmorton in agro Worcesterensi. *Rev. D. Sheffield.* 4. VII.

setaceus. 8. SCIRPUS culmo nudo setaceo, spicis lateralibus subgeminis sessilibus. *Sp. pl.* 73. *Fl. suec.* 45.

Scirpus omnium minimus, capitulo brevior. *Sch. agr.* 358.

Scirpus foliaceus humilis. *C. Gifs.* 158. *R. Syn.* 430.

Juncellus omnium minimus. *Chamæschænus. Lob. adv.* 44. *Hist. Ox. III.* 232. t. 10. f. 23.

Gramen junceum minimum capitulo squamoso. *B. pin.* 6. pr. 13.

Gramen junceum maritimum exile Plimmostii. *Park.* 1270. ic. 1271. descr.

Anglis, least Club-rush or Bull-rush.

Habitat in arenosis humidioribus. ☉. VII. VIII.

Obs. variat numero spicularum, ab uno ad quatuor.

*** Culmo triquetro, Panicula nuda.

mucronatus. 9. SCIRPUS culmo triquetro nudo acuminato spicis conglomeratis lateralibus. *Sp. pl.* 73.

Scirpus caule triquetro, panicula laterali ramosa, locustis ovatis. *Hall. hist.* 1338.

Juncus acutus maritimus, caule triangulo. *B. pin.*

11. *Tb.* 175. *Park.* 1194. *Hist. Ox. III.* 232. t. 10. f. 20.

Juncus acutus maritimus, caule triquetro, rigido, mucrone pungente. *Pluk. ph. t.* 40. f. 1. *R. Syn.* 429.

triqueter. β Scirpus culmo triquetro nudo, spicis subsessilibus pedunculatisque mucronem æquantibus. *Syst.* 83.

Juncus acutus maritimus, caule triquetro maximo molli, procerior nostras. *Pluk. ph. t.* 40. f. 2.

Anglis, pointed Club-rush, or Bull-rush.

Habitat

Habitat in litoribus maritimis; ad ripam Thameſis fluvii tam infra quam ſupra Londinum. 4. VIII.

**** Culmo triquetro, Panicula foliacea.

10. SCIRPUS culmo triquetro, panicula conglobata *maritima* foliacea, ſpicularum ſquamis trifidis: intermedia *mus. ſubulata*. *Sp. pl.* 51.

Gramen cyperoides panicula ſparſa, majus. *B. pin.* 6.

Gramen cyperoides paluſtre, panicula ſparſa, majus. *Park.* 1266. *R. Syn.* 425.

β *Cyperus rotundus inodorus anglicus*. *B. pin.* 14.

Anglis, round rooted Club-ruſh, or Baſtard Cyperus.

Habitat ad ripas fluviorum et in litoribus maritimis. 4. VIII.

Obſ. variat paniculâ ſimplici et ramoſa.

11. SCIRPUS culmo triquetro folioſo, umbella foliacea *ſylvaticus*. pedunculis nudis ſuperadecompoſitis, ſpicis confertis. *Sp. pl.* 51. *Fl. ſuec.* 48. *dan.* 307.

Scirpus planifolius, paniculâ ramoſiſſima, locuſtis ovatis. *Halt. hiſt.* 1340.

Gramen cyperoides miliaceum. *B. pin.* 6. *Th.* 90.

· Pſeudo-cyperus miliaceus. *Park.* 1171.

Cyperus gramineus. *B. hiſt.* II. 504. *R. ſyn.* 426. *miliaceus Ger. em.* 30.

Anglis, wood Club-ruſh, or Millet Cyperus Graſs.

Habitat in aquoſis et in ſylvis humidis; prope Hampſtead; by the Thame ſide, near Tamworth, Warwickſhire; near Haverford-Weſt, Pembrokeſhire; near Bocking, Eſſex. *Mr. Dale.* *R. ſyn.* 4. VII.

ERIOPHORUM. 68.

Glumæ paleaceæ, undique imbricatæ. *Cor.* O. *Sem.*

1. lana longiſſima cinctum.

1. ERIOPHORUM culmis teretibus, foliis planis, *polyſtachion* ſpicis pedunculatis. *Sp. pl.* 76. *Dalib. pariſ.* 18.

Eriophorum foliis planis, ſpicis pendulis. *Hall. hiſt.* 1331.

TRIANDRIA MONOGYNIA.

Gramen pratense tomentosum, panicula sparsa. *B. pin.* 4.

Gramen junceum lanatum, vel juncus bombycineus vulgaris. *Park.* 1271.

Gramen tomentarium. *Ger. em.* 29.

Linagrostis. *Tab. hist.* 599. *R. syn.* 435. Panicula ampliori. *Vaill. parif. t.* 16. *f.* 1. *Sch. agr.* 306.

Anglis. common Cotton-Grass.

Habitat in pratis udis, paludibus, cespitosis. 4. VI. VII.

vagina-
tum.

2. ERIOPHORUM culmis vaginatis teretibus, spica scariosa. *Sp. pl.* 76. *Dalib. parif.* 18.

Juncus capitulo lanuginoso. f. Schoenolaguros. *B. pin.* 12. *Sch. agr.* 302. *pr. t.* 7.

Juncus alpinus cum cauda leporina. *B. hist. II.* 514.

Gramen juncoides lanatum alterum danicum. *Park.* 1271.

Anglis, mountain Cotton-Grass, or Hairs-tail-rush.

Habitat in paludibus turfosis alpinis, in comitatibus Westmorlandico, Cumberlandico, &c. frequens; on the great bog beyond Joan Cole's, towards Croydon. Mr. Sherard. *R. syn.* 4. II.—IV.

N A R D U S. 69.

Calyx nullus. *Cor.* bivalvis.

stricta.

1. NARDUS spica setacea recta secunda. *Sp. pl.* 77. *Hall. hist.* 1410. *Schreb. gram.* 65. *t.* 7.

Gramen sparteum juncifolium. *B. pin.* 5. *R. syn.* 393. *Sch. agr.* 90.

Spartum parvum Batavicum et Anglicum. *Park.* 1199.

Spartum nostras parvum Lobelio. *Ger. em.* 43.

Anglis, Mat-grass.

Habitat in ericetis humidiusculis, frequens. 4. VI.—VIII.

DIGYNIA.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

TRIANDRIA DIGYNIA.

Anglis, reed Canary-grass.

Habitat ad ripas fluviorum et in paludibus. 4. VII.

P A N I C U M. 76.

Cal. trivalvis ; valvula tertia minima.

viride.

1. PANICUM spica tereti, involucellis bifloris fasciculato-pilosis, seminibus nervosis. *Sp. pl.* 83.

Panicum spica unica, flosculis sessilibus solitariis, setis numerosis. *Hall. hist.* 1542.

Gramen paniceum spica simplici lævi. *R. syn.* 394.

Gramen panici effigie spica simplici. *Ger. em.* 17.

Gramen paniceum spica simplici. *B. pin.* 8. *Park.* 1154. *Sch. agr.* 46.

Anglis, green Panick-grass.

Habitat in arvis arenosis ; prope Battersea copiose, et circa Martha's Chapel duobus millearibus, à Guildford distans. ☉. VII.

verticillatum.

2. PANICUM spica verticillata, racemulis quaternis, involucellis unifloris bisetis, setis scabris, culmis diffusis. *Sp. pl.* 82.

Panicum spica unica paniculata, setis paucioribus. *Hall. hist.* 1543.

Gramen paniceum spica aspera. *B. pin.* 8. *Th.* 139. *Sch. agr.* 97. *R. syn.* 394.

Anglis, rough Panick-grass.

Habitat in arvis, inter Putney et Roughampton. *R. syn.* ☉. VI. VII.

erugalli.

3. PANICUM spicis alternis conjugatisque, spiculis subdivisis, glumis aristatis hispidis, rachi quinquangulari. *Sp. pl.* 83.

Gramen paniceum spica divisa. *B. pin.* 8. *R. syn.* 394.

Panicum vulgare. *Ger. em.* 85.

Panicum sylvestre Herbariorum. *Park.* 1154.

Anglis, loose Panick-grass.

Habitat in arvis, prope Battersea ; et circa Martha's Chapel, prope Guildford. ☉. VII.

4. PANICUM

4. PANICUM spicis digitatis basi interiore nodosis, *sanguina-*
flosculis geminis muticis, vaginis foliorum punc- *le.*
tatis. *Sp. pl.* 84. *Schreb. gram.* 119. t. 16.

Digitaria foliis subhirsutis, caule debili, spicis verticil-
latis, scapo ancipiti. *Hall. hist.* 1526.

Gramen dactylon latiore folio. *B. pin.* 8. *Sch. agr.*
101. *R. syn.* 399.

Ischæmon sylvestre latiore folio. *Park.* 1178.

Ischæmon vulgare. *Ger. em.* 27.

Anglis, Cock's-foot Panick-grafs.

Habitat in arvis; about Elden, Suffolk; at Witching-
ham near Norwich, *R. S. prope* Battersea; et prope
Martha's Chapel, juxta Guildford. ☉. VII. VIII.

5. PANICUM spicis digitatis patentibus; basi inte- *dactylon.*
riore villosis, floribus solitariis, sarmentis repenti-
bus. *Sp. pl.* 85.

Digitaria stolonibus teretibus, foliis patulis glabris,
spicis digitatis. *Hall. hist.* 1527.

Gramen dactylon, folio arundinaceo, majus aculea-
tum. *B. pin.* 7.

Gramen dactyloides radice repente. *Ger. em.* 28.

Gramen canarium ischæmi paniculis. *Park.* 1179,

Gramen repens cum panicula graminis Mannæ. *B.*
hist. II. 459. *R. syn.* 399.

Anglis, creeping Panick-grafs.

Habitat in arenosis, inter Penzance et Marketjeu in
Cornubia. D. Newton. *R. syn.* 4. VII.

PHLEUM. 77.

Cal. bivalvis, sessilis, linearis, truncatus, apice bicuspi-
dato. *Cor.* inclusa.

1. PHLEUM spica cylindrica longissima, glumarum *pratense.*
carina ciliata.

α. Phleum spica cylindrica longissima ciliata, culmo
erecto. *Sp. pl.* 87. *Schreb. gr.* 102. t. 14.

Gramen typhoides maximum, spica longissima. *B.*
pin. 4. *Tb.* 49. *Hist. Ox.* III. 188. t. 4. f. 1.
Sch. agr. 65.

Gramen typhinum majus seu primum. *Ger. em.* 11.
R. syn. 398.

Gramen

Gramen typhinum medium s. vulgatissimum. *Park.*
1170.

β Gramen typhinum minus. *Ger. em.* 11. *Park.*
1170.

nodosum. γ Phleum spica cylindrica, culmo adscendente, foliis
obliquis, radice bulbosa. *Syst. nat.* 88. *Sp. pl.*
88.

Gramen nodosum spica parva. *B. pin.* 2. *pr.* 3.
Anglis, α Meadow Cat's-tail grass γ Bulbosa Cat's-
tail-grass.

Habitat in pratis et pascuis, et γ in pascuis sterilibus, et
ad vias. 4. VII. VIII.

Obs. α. et β. Radix sabbulbosa, Culmus erectus; glumæ
carinatae, rectae, carinis ciliatis. γ Radix bulbosa,
Culmus adscendens. Glumæ carinatae, carinis glu-
marum ciliatis; at in horto culta, Radix fibrosa et
Culmus erectus evadit.

panicula- 2. PHLEUM panicula cylindrica spiciformi, glumis
tum. nudis, culmo subramoso. *Fl. Angl.* 23.

Phleum culmo recto, spicis cylindricis, glabris, caly-
cibus brevissime mucronatis. *Hall. hist.* 1536.

Phalaris panicula cylindrica spiciformi. *Sp. pl.* 80.

Gramen typhoidides asperum primum. *B. pin.* 4.
Th. 51.

Anglis, bulbosa Cat's-tail-grass.

Habitat in pratis, infra King's Weston prope Bris-
tolium; on Newmarket-heath. *D. C. Miller.*
4. VII.

Desc. Radix fibrosa. Culmus pedalis, subobliquus,
teres, glaber, geniculatus, ramosus. Folia plurima,
alterna, erecta, petiolata, ensiformia, acuta, nervosa,
scabra. Petiolus cylindricus, longissimus, vaginans,
scaber. Stipula solitaria, intrafoliacea, vaginans,
obtusula, membranacea. Panicula cylindrica, longissima.
spiciformis, ad alterutrum flexa in lobos divisibilis.
Gluma sessilis, imbricata, bivalvis uniflora, acumi-
nata, superne vinctricosa, nuda. Corolla bivalvis
mutica, glumis brevior.

ALOPECURUS. 78.

Calix bivalvis. *Corolla* univalvis.

1. ALOPECURUS culmo erecto, panicula cylindrica *pratensis*.
spiciformi, glumis villosis.

Alopecurus culmo spicato erecto, glumis villosis. *Sp. pl.* 80. corollis muticis. *Syst. nat.* 89. *Schreb. gr.* 133. t. 19. f. 1.

Gramen alopecuroides majus. *Ger. em.* 18.

Gramen phalaroides majus, f. Italicum. *B. pin.* 4.

Park. 1164. *Hist. Ox.* III. 191. t. 4. f. 8.

Anglis, Meadow Fox-tail-grass.

Habitat in pratis et pascuis. 4. V.

2. ALOPECURUS culmo erecto, spica lineari-lanceolata, glumis subnudis. *agrestis*.

Alopecurus culmo spicato erecto, glumis nudis. *Sp. pl.* 89. *Schreb. gram.* 140. t. 19. f.

Gramen myosuroides majus, spica longiore, aristis rectis. *R. syn.* 307.

Gramen typhoides spica angustiore. *B. pin.* 4.

Gramen alopecuroides, spica longa, majus et minus. *Park.* 1169.

Gramen alopecurinum minus. *Ger. em.* 11.

β Gramen myosuroides minus, spica brevior, aristis recurvis. *R. syn.* 397.

Anglis, field Fox-tail-grass.

Habitat in arvis et ad vias. 4. VII. VIII.

3. ALOPECURUS culmo ascendente, spica cylindrica, glumis apice divergentibus pilosis. *geniculatus*.

Alopecurus culmo spicato infraeto. *Sp. pl.* 89. corollis muticis. *Syst. nat.* 89.

Gramen aquaticum geniculatum spicatum. *B. pin.* 3. *Tb.* 41. *Sch. agr.* 72. *R. syn.* 396.

Gramen fluviatile spicatum. *Ger. em.* 14.

Gramen aquaticum spicatum. *Park.* 1273.

β Alopecurus spica cylindrica, culmo erecto, radice bulbosus. bulboso. *F. Angl.* 24. *Sp. pl.* 1665.

Gramen myosuroides nodosum. *R. syn.* 397. t. 22. f. 2. *Anglis*,

Anglis, Flote β bulbose Fox-tail-grass.

Habitat in aquosis et stagnis. β In the first field next the road before you go into Northfleet. Dr. Wilmer. *R. syn.* V.—VIII.

Obs. *A. bulbosus* in aquosis degenerat in geniculatum, cuius varietas est.

aristatus. 4. **ALOPECURUS** culmo suberecto, panicula coarctata spiciformi, glumis aristatis, aristis longissimis.

monspe-liensis. Alopecurus panicula subspicata, glumis scabris, corollis aristatis. *Sp. pl.* 89.

Alopecurus maxima anglica. *Park.* 1166. *R. syn.* 396.

Alopecurus altera maxima anglica paludosa. *Ger. em.* 88. *Hist. Ox.* III. 191. t. 4. f. 3.

Phleum crinitum, calycibus linearibus basi subventricosis hispidis aristatis, aristis capillateis longissimis. *Schreb. gram.* 151. t. 20. f. 3.

Gramen alopecuroides anglobritannicum maximum. *B. pin.* 4.

paniceus. β Alopecurus panicula subspicata, glumis villosis, corollis aristatis. *Sp. pl.* 90.

Anglis, bearded Fox-tail grass.

Habitat in pratis humidiusculis. β aridis, in riguis herbidis Comitatus Southamptoniæ proxime salinas et antiquas ædes Drayton vocatas, cis mare duobus milliaribus anglicis a Portsmouth ex adverso Vectis insulæ plurima; at Purfleet, over against the mill towards Raynam on the other side of the great ditch. Mr. Newton. *R. syn.* circa Clay in Norfolk. D. Humphrey. ☉. VI.—VIII.

Obs. Gluma bivalvis, valvulis linearibus, apice divergentibus aristatis; arista terminali longissima; Corolla minima aristata; arista dorsali brevi.

ventricosus. 5. **ALOPECURUS** culmo obliquo, panicula lanceolata spiciformi, glumis basi ventricoli, acuminatis.

lendigerum. Miliun panicula subspicata, floribus aristatis. *Sp. pl.* 91. *Schreb. gram.* II. 14. t. 23. f. 3.

Agrostis calycibus inferne ventricosis, valvula exteriore acuminata longitudine aristæ. *Gouan Hort. Mensp.* 39. t. 1.

Panicum

Panicum serotinum arvense, spica pyramidata. Tourn.
515. *R. syn.* 394.

Gramen serotinum spica laxa pyramidali. R. hist. 1288.

*Gramen alopecuro accedens, ex culmi geniculis spicas
cum petiolis longiusculis promens. Pluk. ph. t. 33.*
f. 6.

Anglis, panick Fox-tail-grass.

Habitat in pratis et pascuis. In Insula Shepey copiose.

○. VII.—VIII.

*Obs. Gluma bivalvis, inæqualis, basi ventricosa, acu-
minata; Corolla brevissima, ovata, aristata; arista
infra apicem longitudine glumæ.*

S T I P A. 90.

*Cal. bivalvis, uniflorus. Cor. valvula exteriori arista
terminali. basi articulata.*

1. *STIPA aristis lanatis. Sp. pl.* 125. *Hall. hist.* 1541. *pennata.*

Gramen spartheum pennatum. B. pin. 5. *R. Syn.*
393. *Sch.* 153.

Spartum austriacum pennatum. Clus. hist. II. 221.
Ger. em. 42.

Anglis, Feather-grass.

*Habitat in montosis. Found by Dr. Richardson in
company with T. Lawson, on the limestone rocks
hanging over a little valley called Longleadale, about
six miles north of Kendal, Westmoreland. 4.*
VII. VIII.

M I L I U M. 79.

*Calix bivalvis, uniflorus. Cor. brevissima. Stigmata
penicilliformia.*

1. *MILIUM floribus paniculatis dispersis muticis. Sp. effusum.*
pl. 90. *Fl. suec.* 61. *Dalib. par.* 23.

Milium paniculis raris, longissime petiolis. Hall. hist.
1525.

Gramen miliaceum. Lob. ic. 3. *Ger. em.* 6. *B.*
hist. II. 462. *R. Syn.* 402.

Gramen sylvaticum, panicula miliacea sparsa. B. pin.
8. *Sch. agr.* 133.

Gramen

TRIANDRIA DIGYNIA.

Gramen miliaceum vulgare. *Park.* 1153.

Anglis, Millet-grafs.

Habitat in *fykvis humidiusculis frequens.* ☉. VI. VII.

A G R O S T I S. 80.

Cal. bivalvis uniflorus : valvulis acutis æqualibus.

Cor. bivalvis inæqualis, gluma paulo minor. *Stigmata* plumosa.

* *Aristatæ.*

spica-ven- 1. AGROSTIS petala exteriori arista recta stricta longissima, panicula patula. *Sp. pl.* 90.

Avena monantha, panicula ascendente, multiflora, calyce levi, flori arista longissima. *Hall. hist.* 1480.

Gramen segetum panicula arundinacea. *B. pin.* 3.

Gramen arundinaceum. *Ger. em.* 5.

Gramen agrorum venti spica. *Park.* 1151.

Gramen miliaceum majus, glumis aristatis, spadiceis et pallidis. *Pet. conc. gr. n.* 122. 123. *R. Syn.* 405.

Anglis, silky Bent-grafs.

Habitat in *arvis solo arenoso. passim.* ☉. VI.—VIII.

canina. 2. AGROSTIS panicula patente, corolla nuda, valvula altera dupla longiore aristata, arista dorsali rectiuscula.

α Agrostis calycibus elongatis, petalarum arista dorsali recurva, culmis prostratis subramosis. *Sp. pl.* 92. foliis planis. *Syst. nat.* 90.

Avena monantha, panicula diffusa, multiflora, ascendente, gluma calycina exasperata. *Hall. hist.* 1479.

Gramen supinum caninum paniculatum, folio varians. *B. pin.* 1. *Tb.* 12.

Gramen paniculatum supinum, ad infima culmorum genicula, foliorum capillarum fasciculis donatum. *Sch. agr.* 141. t. 3. f. 9. C.

β Agrostis foliis radicalibus filiformibus ; caulinis planis, culmo suberecto.

Avena monantha, paniculata, gluma calycina exasperata arista flori æquali. *Hall. hist.* 1477.

Gramen

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Poa stolonifera monantha, calycibus leviter exasperatis.
Hall. hist. 1473.

Gramen montanum miliaceum minus, radice repente,
R. Syn. 402.

Gramen caninum supinum minus. *B. pin.* 1. *Sch. agr.* 128.

Gramen miliaceum folio angustissimo. *Pet. conc. gr.* 116.

palustris. ζ *Agrostis panicula coarctata mutica*, calycibus æqualibus hispidiusculis coloratis, culmo repente. *Fl. Angl.* 27.

Agrostis (alba) panicula laxa, calycibus muticis æqualibus. *Roy. lugdb.* 50. *Sp. pl.* 93.

Gramen miliaceum majus, panicula spadicea. *Pet. conc. gr.* 118. *R. Syn.* 464.

Gramen miliaceum majus, panicula viridi. *Pet. conc. gr.* 118. *R. Syn.* 464.

sylvatica. η *Agrostis panicula coarctata mutica*, calycibus æqualibus corolla ante florescentiam calyce brevior, postea duplo longiore. *Fl. Angl.* 28.

Agrostis panicula coarctata mutica, calycibus æqualibus: virgineis corolla brevioribus, fœcundatis duplo longioribus. *Sp. pl.* 1665.

Gramen miliaceum sylvestre, glumis oblongis. *Pet. conc. gr.* 121. *R. Syn.* 404.

Anglis, fine, γ little, δ creeping, ζ marsh η wood Bent-grass.

Habitat in pratis et pascuis ζ in fossis et uliginosis η in sylvis humidiusculis. in Bishops Wood Hampstead. Mr. Buddle. *R. syn. in sylvula Hornsey-Wood dicta.* 4. VI.—VIII.

minima. 4. AGROSTIS panicula mutica filiformi. *Sp. pl.* 93.
Nardus panicula spicata, floribus solitariis alternis.
Guett. Stamp. 155. *Dalib. paris.* 153.

Gramen minimum paniculis elegantissimis. *B. pin.* 2.
Sch. agr. 40. *Hist. Ox.* III. 200. t. 2. f. 10.

Gramen minimum. *B. hist.* II. 465. *Dalech. hist.* 425. anglo-britannicum. *Lob. ill.* 29. *R. Syn.* 11. dub.
Anglis, small Bent-grass.

Habitat in Wallia. D. Stillingfleet invenit. ©. VII.
A I R A.

A I R A. 81.

Cal. bivalvis, biflorus. Flosculi absque interjecto rudimento.

* *Mutica.*

1. AIRA panicula spicata, calycibus pedunculo longi-*cristata.*
 oribus, petalis subaristatis inæqualibus. *Sp. pl.* 94.
 Poa panicula spicata, calycibus subpilosis subquadri-
 floris pedunculo longioribus, petalis aristatis. *Syst.*
Nat. 94.
 Festuca locustis bifloris, mucronatis, confertis, imbricatis paniculatis, in spicam congestis. *Hall. bist.* 1444.
 Gramen spica cristata, subhirsutum. *B. pin.* 2. *Hist. Ox. III.* 194. t. 4. f. 7. *Scb. agr.* 166.
 Gramen pumilum, spica purpuro-argentea molli. *R. syn.* 396.
Anglis, cristed Hair-grass.
 Habitat in pratis montosis sterilibus. 4. VII. VIII.
2. AIRA panicula coarctata, floribus pedunculatis muti-*cærulea.*
 cis convoluto-subulatis, foliis planis. *Sp. pl.* 95.
 Poa spiculis subulatis, panicula rara contracta. *Fl. lapp.* 29. *Roy. lugdb.* 61.
 Gramen pratense serotinum, panicula longa purpurascente. *R. Syn.* 404. *Hist. Ox. III.* 201. t. 5. f. 22.
 Gramen pratense spica Lavendulæ. *Merr. pin.*
Anglis, purple Hair-grass.
 Habitat in pratis et pascuis uliginosis, sterilissimis frequens. 4. VII. VIII.
3. AIRA panicula patente, floribus muticis lævibus ob-*aquatica.*
 tusis calyce longioribus, foliis planis. *Sp. pl.* 95.
 Aira culmo inferiore repente, flosculis muticis calyce longioribus : altero pedunculato. *Roy. lugdb.* 60.
 Gramen miliaceum aquaticum. *R. syn.* 402. *Scb. agr.* 176. *Vaill. paris.* 89. t. 17. f. 7.
 Gramen canium supinum paniculatum dulce. *B. pin.* 1.

Gramen exile tenuifolium Canario simile. f. Gramen dulce. *Park.* 1174.

distans. β *Poa* panicula ramis subdivisis, floribus quinquefloris, flosculis distantibus obtusis. *Mant. I.* 32. *Syst. Nat.* 94. *Anglis*, water Hair-grass.

Habitat in aquosis et ad ripas fluviorum. β in arenosis prope Exmouth; circa Northfleet; in agro Cantiano; in comitatibus Eboracensi et Lancasterensi passim. 4. VI. VII.

Desc. β. Radix fibrosa. Culmi plures, obloqui et erectiusculi, teretes, glabri. Folia radicalia, pauca ensiformia, erecta, obtusa, nuda, caulina petiolata, obtusa, subinde involuta, glauca. Petiolus vaginans, cylindricus, striatus, glaber. Stipula membranacea, brevissima, obtusa. Panicula diffusa patens, ramosa, ramis subquaternis. Spiculae lineares. Flosculi a tribus ad sex vel septem, remoti obtusi quasi truncati, subinde purpurei, margine albo. Glumae bivalvis, inequalis, valde obtusa.

caespitosa. 4. *AIRA* foliis planis, panicula patente, petalis basi villosis aristatisque: arista recta brevi. *Sp. pl.* 96. *Fl. dan.* 24.

Avena diantha, panicula amplissima, patula, bicolore, *Hall. hist.* 1487.

Gramen segetum altissimum panicula sparsa. *B. pin.* 3. *Hist. Ox. III.* 199. t. 5. f. 1. ?

Gramen segetum panicula speciosa. *Park.* 1158.

Gramen segetale. *Ger. em.* 5.

Gramen miliaccum segetale majus. *Pet. conc. gr.* 114. *R. syn.* 403.

Anglis, turfy Hair-grass.

Habitat in pratis et sylvis humidis. 4. VII.

flexuosa. 5. *AIRA* foliis setaceis, culmis subnudis, panicula patente, pedunculis flexuosis, flosculis basi pilosis aristatis: arista tortili longiore.

Aira foliis setaceis, culmis subnudis, panicula divaricata, pedunculis flexuosis. *Sp. pl.* 96. *Fl. dan.* 157.

Gramen alpinum nemorosum paniculatum, foliis angustissimis, locustis splendentibus aristatis. *Seb. agr.* 218. pr. t. 6. f. 1.

Gramen

Gramen nemorosum paniculis albis, capillaceo folio.

B. pin. 7. pr. 14. th. 97. Hist. Ox. III. 300. f. 8. t. 7. f. 9. R. syn. 407.

Gramen paniculatum locustis parvis purpureo-argenteis, majus et perenne. *R. syn. 407.*

Avena diantha, paniculis flexuosis, floribus basi villosis, locustis ovatis. *Hall. hist. 1486.*

Anglis, heath Hair-grass.

Habitat in ericetis, sylvis et pascuis sterilibus. 4. VII. VIII.

Obs. variat floribus fusco-purpureis et virescentibus.

6. AIRA foliis setaceis, panicula erecta angusta subsecunda, flosculis basi pilosis aristatis; arista tortili longiore. *montana.*

Aira foliis setaceis, panicula angusta, flosculis basi pilosis aristatis: aristas tortili longiore. *Sp. pl. 96.*

Gramen avenaceum capillaceum minoribus glumis. *B. pin. 10.*

Gramen avenaceum paniculatum alpinum, foliis capillaceis brevibus, locustis purpureo-argenteis et aristatis. *Sch. agr. 216. pr. 1. 4. f. 4.*

β Aira foliis setaceis, panicula secunda coarctata, aristatis flosculis duplo-longioribus recurvis. *Fl. angl. 30.*

Anglis, mountain Hair-grass.

Habitat in ericetis et pascuis sterilibus et arenosis; in ericeto Strattoniensi in comitatu Norfolk, D. Stillingfleet; in ericetis arenosis in comitatibus Eboracensi et Lancasterensi passim. 4. VII. VIII.

Desc. Radix perennis, fibrosa. Culmi plures, semipedales et sesquipedales erectiusculi, teretes, laeves, bi s. triarticulati. Folia radicalia plurima, fasciculata, setacea, nuda, glauca, rigidiuscula; caulina duo aut tria patentia, petiolata, involuta, ut setacea apparent, rigida. Petiolus cylindricus, longissimus, vaginans, striatus, nudus. Stipula intrasoliacea, persistens, membranacea, erecta, acuta. Panicula angusta, erecta secunda, coarctata ramosa; Ramis flexuosis, scabris, Spiculæ bifloræ; Flosculo altero sessili altero pedunculato. Receptaculum pilosum. Cal. Gluma bivalvis oblonga, concava, acutiuscula, nuda, purpurascens; valvulis æqualibus, corolla longioribus, Corolla bivalvis; valvulis oblongis, æqualibus, concavis, acutis, glabris, valvula exteri-

TRIANDRIA DIGYNIA.

ore basi aristata : arista corolla duplo longior, medio flexa.
 Stam. Filamenta tria capillaria, brevia. Antheræ
 hastatæ purpuræ Stig. villosa.

canescens. 7. AIRA foliis setaceis, panicula coarctata erecta flos-
 culis aristatis gluma duplo brevioribus,

Aira foliis setaceis: summo paniculam inferne obvol-
 vente. *Fl. suec. 74. Sp. pl. 97.*

Avena diantha, foliis setaceis, panicula stricta, aristis
 clavatis. *Hall. hist. 1483.*

Gramen foliis junceis, radice jubata. *B. pin. 5. Th.*
72. Sch. agr. 243.

Gramen miliaceum maritimum molle. *Pet. Conc. gr.*
126. R. syn. 405.

Anglis, grey Hair-grass.

Habitat in arenosis maritimis. u. VII. VIII.

*Desc. Rad. perennis, fibrosa; Culmi plures, semipedales,
 obloqui, glabri. Folia radicalia fasciculata, setacea, sca-
 bra, rigida, glauca; caulina erecta, petiolata, invo-
 luta, acuta, scabra. Petiolus cylindricus vaginans
 scaber fere longitudine internodii: supremus spathecus
 paniculam inferne obvolvens. Panicula oblonga, erecta,
 coarctata, ramosa; Spiculae bifloræ, flosculo altero sessili,
 altero pedunculato, gluma duplo brevior. Gluma bival-
 vis: valvulis æqualibus, oblongis, acutis, carinatis, carina
 scabra. Corolla bivalvis: valvulis inæqualibus ob-
 longis concavis, acutiusculis, glabris; valvula exterior
 media aristata; Arista longitudine glumæ, in medio cincla
 denticulis, infra crassior fusca, supra subclavata albida.*

præcox. 8. AIRA foliis setaceis: vaginis angulatis, floribus pa-
 niculato-spicatis, flosculis basi aristatis. *Sp. pl. 97.*

Gramen parvum præcox, panicula (potius spica) laxa
 canescente. *R. Syn. 407. Pluk. ph. t. 33. f. 9.*

Anglis, early Hair-grass.

Habitat in arenosis et sterilibus frequens. O. V. VI.

caryophyl- lea. 9. AIRA foliis setaceis, panicula trichotoma, floribus
 aristatis divergentibus.

Aira foliis setaceis, panicula divaricata, floribus arista-
 tis distantibus. *Sp. pl. 97.*

Aira

Aira foliis subulatis, aristis e basi flosculorum tortilibus, flosculo altero pedunculato. *Roy Lugdb.* 60.

Gramen paniculatum, locustis purpuro-argenteis, annuum. *R. Syn.* 407.

Gramen paniculatum purpuro-argenteum, locustis parvis, annuum. *Hist. Ox. III.* 200. t. 5. f. 11.

Caryophyllus arvensis glaber minimus. *B. pin.* 210. pr. 105.

Anglis, silver Hair-grass.

Habitat in pascuis arenosis frequens. ☉. VII.

MELICA. 82.

Cal. bivalvis, biflorus, Rudimentum floris inter flosculos.

1. MELICA petalis imberbibus, panicula secunda nutante, gluma uniflora.

Gramen avenaceum locustis rarioribus. *B. pin.* 10.

Gramen avenaceum spica mutica rariore gluma. *Hist. Ox. III.* t. 7. f. 49.

Gramen avenaceum nemorense, glumis rarioribus ex fusco zerampelinis. *R. Syn.* 403.

Anglis, wood Melic-grass.

Habitat in sylvis et sepibus frequens. ♄. VI. VII.

2. MELICA petalis imberbibus, panicula coarctata, secunda, erecta, subsimplici, floribus pendulis.

Melica petalis imberbibus, panicula nutante simplici. *Sp. pl.* 98. *Schreb. gram.* 62. t. 6. f. 1.

Melica floribus sub culmo pendulis. *Fl. lapp.* 36.

Gramen avenaceum, locustis rubris, montanum. *B. pin.* 10. pr. 20.

Gramen locustis rubris. *B. hist.* 11. 434. *Park.* 1151.

Gramen avenaceum gluma mutica montanum, locustis rubris. *Hist. Ox. III.* t. 7. f. 48.

Anglis, mountain Melic-grass.

Habitat in nemoribus montosis in comitatibus Eboracensi, Westmorlandico et Cumberlandico frequens. ♄. VI. VII.

TRIANDRIA DIGYNIA.

BRIZA. 84.

Cal. bivalvis, multiflorus. Spicula disticha: valvulis cordatis, obtusis: interiore minore.

media.

1. BRIZA spiculis ovatis calyce flosculis brevioribus. *Fl. suec.* 86. *dan.* 258. *Sp. pl.* 103.
Gramen tremulum. *B. hist.* II. 469. majus. *B. pin.* 2.
R. Syn. 412. *Sch. agr.* 204.
Gramen tremulum s. phalaris media anglica prima an secunda. *Park.* 1165.
Phalaris pratensis. *Ger. em.* 86.
Anglis, middle Quaking-grass, Cow-quakes and Ladies-hair.
Habitat in pratis et pascuis. 4. VI.

minor.

2. BRIZA spiculis triangulis, calyce flosculis longioribus. *Hort. cliff.* 23. *Roy. lugdb.* 63. *Sp. pl.* 102.
Gramen tremulum minus, panicula parva. *B. pin.* 2.
Hist. Ox. III. 208. t. 6. f. 47.
Gramen tremulum minus, panicula ampla, locustis parvis triangulis. *R. Syn.* 412.
Anglis, small Quaking-grass.
Habitat in pascuis. *In Jersey.* Dr. Sherard. *R. Syn.* prope Bath. D. Alchorne. 4. VII.

POA. 83.

Cal. bivalvis, multiflorus. Spicula ovata: valvulis ovato-acuminatis.

- aquatica.* 1. POA panicula erecta ramosa, spiculis ovato-linearibus, culmo erecto.
Poa panicula diffusa, spiculis sexfloris linearibus. *Fl. suec.* 81. *Sp. pl.* 98.
Poa altissima, foliis latissimis, panicula amplissima, locustis distichis multifloris. *Hall. hist.* 1454.
Gramen paniculatum aquaticum latisolium. *B. pin.* 3.
Hist. Ox. III. 201. t. 6. f. 25.
Gramen aquaticum majus. *Ger. em.* 6. *R. Syn.* 411.
Anglis, water Meadow-grass.
Habitat in paludibus et ad ripas fluviorum. 4. VII. VIII.
Desc. Culmus quatuorpedalis et sexpedalis, erectus, anceps, glaber. Folia ensiformia erecta, acuta, carinata,
nata,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Poa panicula diffusa; spiculis subtrifloris basi pubescentibus, culmo erecto tereti. *Sp. pl.* 99.

Poa panicula diffusa, locustis trifloris, glabris. *Hall. hist.* 1465.

Gramen pratense paniculatum medium. *B. pin.* 2. *R. Syn.* 409. *Sch. agr.* 180.

Gramen pratense minus. *B. hist. II.* 542. *Ger. em.* 2. *Park.* 1156.

palustris. β *Poa panicula diffusa*, spiculis subtrifloris pubescentibus, foliis subtus scabris. *Sp. pl.* 98.

Gramen paniculatum aquaticum angustifolium, panicula speciosa. *Sch. agr.* 184.

Gramen palustre panicula speciosa. *B. pin.* 3. *pr.* 7. *Hist. Ox. III.* 201. t. 6. f. 27.

setacea. γ *Poa panicula diffusa*, spiculis trifloris pubescentibus, foliis radicalibus-setaceis. *Fl. Angl.* 34.

Anglis, common Meadow-grass, β . marsh Meadow-grass, γ . hair-leaved Meadow-grass.

Habitat in pratis et pascuis, β . in aquosis inter *Arundines* et *Poam aquaticam* frequens. α . VI. VII.

Desc. Radix fibrosa; Culmi plures a semi ad quadripedales erecti, subinde ascendentes, ancipites, glabri. Folia petiolata ensiformia acuminata, scabra, in aridis involuta ut setacea apparent. Petiolus vaginans carinatus striatus, scaber; Stipula intrasoliacea membranacea, acuminata. Panicula patens ramosa: ramis triangulis scabris; Spiculæ ovatæ bi s. triflores rarissime quatuor, glomeratæ, subpedunculatæ, pedunculis brevissimis, flosculi acuti, basi et parte inferiore carinæ lanati.

angustifolia. 4. *POA panicula diffusa* erectiuscula, spiculis ovatis pedunculatis, pedunculis divaricatis, flosculis basi pilosis, foliis acutis, culmo erecto.

Poa panicula diffusa, spiculis quadrifloris pubescentibus, culmo erecto tereti. *Sp. pl.* 99. *Fl. suec.* 81.

Poa foliis duris se convolventibus, locustis quadrifloris villosis. *Hall. hist.* 1460.

Gramen pratense paniculatum majus angustiore folio. *B. pin.* 2. *pr.* 5. *Hist. Ox. III.* 201. t. 5. f. 19.

Sch. 178.

β *Poa panicula attenuata*, spiculis subbifloris mucrona- nemora-
tis scabris, culmo incurvo. *Sp. pl.* 102. lis.

Poa culmo debili panicula nutante, locustis bifloris,
folliculis subvillosis. *Hall. hist.* 1469.

Gramen paniculatum angustifolium alpinum, locustis
rarioribus et angustioribus non aristatis. *Sch. agr.*
164. *pr. t.* 2.

Anglis, narrow-leaved, β. wood Meadow-grass.

Habitat in sylvis et sepibus frequens. 4. VI. VII.

Desc. Radix repens, Culmi plures erecti, ancipites, ge-
niculati, glabri, nitidi. Folia petiolata, ensiformia, a-
cuta, scabriuscula; Petiolus vaginans carinatus, gla-
ber, nitidus. Stipula intrafoliacea brevissima, obtusa.

• Panicula diffusa, erectiuscula, ramosa, ramis angu-
latis divaricatis, scabris. Spiculæ ovatæ, acutæ, pe-
dunculatæ; Flosculi ab uno ad sex acuti basi pilosi.

5. POA panicula secunda patente, spiculis ovatis, flos- bulbosa.
culis acutis glabriusculis, culmo erecto basi bulboso.

Poa panicula secunda patentiuscula, spiculis quadri-
floris. *Sp. pl.* 102.

Poa foliis bulbosis, panicula diffusa, locustis quadri-
floris folliculis subvillosis. *Hall. hist.* 1461.

Gramen arvense panicula crispa. *B. pin.* 3. *pr.* 6.
Barr. ic. 730. longiore nostras. *Pluk. alm.* 176.

Gramen vernalis, radice ascalonitis. *Vaill. paris.* 1. 17.
f. 8.

Anglis, bulbose Meadow-grass.

Habitat in pratis et pascuis. 4. V. VI.

Desc. Radix fibrosa, Culmi plures erecti, teretes, basi
bulbosi. Folia ensiformia, acutiuscula, nuda, glauca.
Panicula ramosa, ramis angulatis, scabriusculis, spi-
culis ovatis acutis, flosculi tres, quatuor subinde quin-
que acuti, basi subpilosi.

6. POA panicula coarctata secunda erecta, culmo ad- compressa.
scendente compresso.

Poa panicula coarctata secunda, culmo obliquo com-
presso. *Fl. suec.* 84. *Sp. pl.*

Poa culmo compresso declinato, locustis distichis, sex-
floris. *Hall. hist.* 1455.

Gramen

Gramen pratense medium, culmo compresso. *Bud. H. S. R. syn.* 409.

Gramen murorum, radice repente. *B. pin.* 2. *pr.* 2.

Gramen paniculatum radice repente, culmo compresso. *Vaill. paris.* 91. t. 18. f. 5.

Anglis, creeping Meadow-grass.

Habitat in locis siccis, tectis, muris. 4. VI. VII.

annua.

7. POA panicula secunda patente, ramis binatis, culmo obliquo ancipiti.

Poa panicula diffusa, angulis rectis, spiculis obtusis, culmo obliquo compresso. *Sp. pl.*

Poa culmo infracto, panicula triangulari, locustis trifloris glabris. *Hall. hist.* 1466.

Gramen pratense minus s. vulgatissimum. *R. syn.* 408.

Gramen pratense paniculatum minus. *B. pin.* 3.

Gramen pratense minimum album et rubrum. *Ger. em.* 3. *Park.* 1156.

Anglis, annual Meadow-grass.

Habitat in pratis et pascuis ubique. ☉. II.—X.

maritima.

8. POA panicula coarctata secunda ramosa, ramis binatis, spiculis oblongis; flosculis obtusis, foliis involutis acutis, culmo obliquo tereti.

Gramen paniculatum maritimum vulgatissimum. *R. syn.* 409.

β Gramen caninum maritimum paniculatum. *R. syn.* 410.

Anglis, sea Meadow-grass.

Habitat in littoribus maritimis frequens. 4. VI. VII.

Desc. Radix perennis repens. Culmus obliquus semipedalis & pedalis, teres, glaber. Folia petiolata ensiformia, patentia, involuta, acuminata, nuda. Panicula secunda oblonga, coarctata, erecta, ramosa, Ramis binatis, infimis subinde patentiusculis. Spiculæ oblongæ. Flosculi a tribus ad septem vel octo, obtusiusculi glabri.

rigida.

9. POA panicula lanceolata secunda ramosa, ramis solitariis, spiculis linearibus, culmo suberecto.

Poa panicula lanceolata, subramosa secunda: ramulis alternis secundis. *Syst. Nat.* 93. *Sp. pl.* 101.

Gramen

Gramen panicula multiplici. *B. pin.* 3. *Park.* 1157.

Gramen loliaceum murorum duriusculum, spica erecta rigida. *Hist. Ox.* III. 182. t. 2. f. 9.

Gramen exile duriusculum in muris et aridis proveniens. *R. syn.* 410.

Anglis, hard Meadow-grass.

Habitat in muris, tectis et arenosis frequens. ☉. VII.

Obs. Numerus flosculorum in spiculis a tribus ad novem variat.

10. POA spicata; spiculis secundis erectis sessilibus. *loliacea.*

Triticum maritimum, calycibus multifloris mucronatis, spica ramosa. *Syst. Nat.* 102.

Gramen pumilum loliaceo simile. *R. syn.* 395.

Gramen exile duriusculum maritimum, foliolis circumvolutis, veluti junceis brevibus. *Pluk. phyt.* 32. t. 7.

Gramen maritimum, panicula loliacea. *B. pin.* 8. pr. 19. *Hist. Ox.* III. t. 2. f. 6.

Anglis, spiked Meadow-grass.

Habitat in arenosis maritimis frequens. ☉. VI. VII.

Obs. Spica subsecunda basi subinde ramosa, ramis brevissimis erectis, Spiculæ ovatae; Flosculi a tribus ad decem, glabri.

DACTYLIS, 86.

Cal. bivalvis, compressus: altera valvula majore carinata.

1. DACTYLIS spicis sparsis secundis scabris numerosis. *Læst. hisp.* 115. *Sp. pl.* 104. *cynosuroides.*

Gramen sparteum serotinum, capite bifido vel gemino. *Merr. pin.*

Sparteum Essexianum, spica gemina clausa. *Pet. conc. gr.* 35. *R. Hist.* III. *App.* 248. *Syn.* 393.

Anglis, smooth Cocks-foot-grass.

Habitat in palustribus maritimis passim. 4. VII. VIII.

2. DACTYLIS panicula secunda glomerata. *Sp. pl. glomeratus.* 105. *Schreb. gram.* 72. t. 8. f. 2.

Poa flosculis confertis uno versu dispositis. *Hort. Cliff.* 28. *Roy. lugdb.* 62.

Bromus

Bromus locustis tetranthis fasciculatis imbricatis. *Hall. hist.* 1512.

Gramen spicatum folio aspero. *B. pin.* 3. *pr.* 59. *th.* 45.

Gramen pratense spica multiplici rubra. *Park.* 1161.

Gramen asperum. *J. Baub. hist.* II. 496. *R. syn.* 400.

Gramen panicula torosa pratense asperum. *Hist. Ox.* III. 202. *t.* 6. *f.* 38.

Anglis, rough Cocks-foot-grass.

Habitat in pratis et pascuis. 4. VI.—VIII.

FESTUCA. 88.

Cal. bivalvis. *Spicula* oblonga teretiuscula, glumis acuminatis.

* *Panicula secunda.*

ovina. 1. FESTUCA panicula secunda coarctata subaristata, glumis acuminatis, culmo tetragono nudiusculo, foliis setaceis. *Sp. pl.* 108.

Festuca foliis perangustis, panicula stricta, locustis quinquefloris, hirsutis. *Hall. hist.* 1442.

Gramen capillaceum, locustellis pennatis non aristatis. *Pluk. ph. t.* 34. *f.* 2. *R. syn.* 410.

Gramen loliaceum, foliis brevibus junceis, minus. *Hist. Ox.* III. 182. *t.* 3. *f.* 13.

Gramen, foliolis junceis brevibus, majus, radice nigra. *B. pin.* 5. *Sch. agr.* 276.

vivipara. β Festuca spiculis viviparis. *Fl. juec.* 94.

Gramen sparteum-montanum, spica foliacea graminea, majus et minus. *R. syn.* 410. *t.* 22. *f.* 1.

Anglis, sheeps Fescue-grass. *Still. misc. t.* 8.

Habitat in pratis et pascuis succioribus, præsertim glareosis. β in montibus Cambro-britannicis, Eboracensibus, Westmorlandicis. 4. VI. VII.

Obs. Numerus flosculorum in singula spicula a tribus ad sex. Flosculi frequentissime acuminati sive acuminato-aristati, rarissime aristati.

duriuscula. 2. FESTUCA panicula secunda oblonga ramosa, ramis subbinatis, spiculis oblongis aristatis, foliis radicalibus setaceis.

Festuca

- Festuca panicula secunda*, oblonga, spiculis sexfloris oblongis lævibus, foliis setaceis. *Sp. pl.* 108.
- Festuca* foliis hirsutis latiusculis, locustis paniculatis quinquefloris, aristatis, glabris. *Hall. hist.* 1435.
- Gramen pratense panicula duriore laxa unam partem spectante. *R. syn.* 413. t. 19. f. 1. *Sch. agr.* 285.
- β *Festuca* panicula spiciformi pubescente, foliis filiformibus. *Sp. pl.* 109? dumetorum.
- Anglis*, hard Fescue-grass.
- Habitat in pratis et pascuis siccis β in sylvis et ad sepes. 4. VI. VII.
- Obs. Variant Foliis radicalibus setaceis et filiformibus, scabriusculis, caulinis planis et canaliculatis, nudis; Spiculis a quatuor ad decem flores; glabris et pubescentibus.
3. **FESTUCA** panicula secunda scabra, spiculis sexfloris rubra. aristatis, flosculo ultimo mutico, culmo semitereti. *Sp. pl.* 109.
- Festuca* foliis perangustis, locustis teretibus, quinquefloris, aristatis oris membranaceis. *Hall. hist.* 1440.
- Gramen alpinum pratense, panicula duriore laxa spadicæ, locustis majoribus. *Sch. agr.* 287. t. 6. f. 9.
- Anglis*, purple Fescue-grass.
- Habitat in ericetis et pascuis montosis in comitatu Westmorlandico. 4. VII.
4. **FESTUCA** panicula secunda oblonga erecta ramosa, *cambrica*, spiculis aristatis lævibus, foliis planiusculis, nudis. *Anglis*, welsh Fescue-grass.
- Habitat in summis montibus circa Llanberis copiose. 4. VII. VIII.
- Desc. Radix perennis fibrosa. Culmus subpedalis obliquus, teres, foliatus, glaber, bis. triarticulatus. Folia radicalia erectiuscula, ensiformia, acutiuscula, canaliculata nuda; caulina duo aut tria, petiolata, acutiuscula plana, nervosa, nuda. Petiolus vaginans, cylindricus striatus, hirsutus. Stipula membranacea, vaginans, obtusa. Panicula secunda, oblonga, subcoarctata basi ramosa. Spiculæ pedunculatæ, subsexfloræ, ovatæ subulato-aristatæ, læves; aristis brevibus.

bus. Pedunculi longitudine fere Spiculæ. affinis Festucæ duriusculæ.

bromoi-
des.

5. **FESTUCA** panicula secunda, spiculis erectis lævibus, calycis altera valvula integra, altera acuminata. *Sp. pl.* 110.

Festuca foliis setaceis, panicula erecta, locustis glabris, longius aristatis. *Hall. hist.* 1443.

Gramen paniculatum bromoides minus, paniculis aristatis unam partem spectantibus. *R. syn.* 415. *hist.* 1287. *Pluk. phyt. t.* 33. *f.* 10. *Sch.* 297.

Anglis, barren Fescue-grass.

Habitat in muris et pascuis arenosis. ☉. VI. VII.

myurus.

6. **FESTUCA** panicula secunda lineari scabra, ramis subsolitariis adpressis, glumæ altera valvula brevissima mutica, floribus aristatis; aristis longis.

Festuca panicula spicata nutante, calycibus minutissimis muticis, floribus scabris; aristis longis. *Sp. pl.* 109. *Roy. lugdb.* 68.

Gramen murorum spica longissima. *Ger. em.* 31. *Hist. Ox. III.* 215. *t.* 7. *f.* 43. *R. syn.* 415.

Gramen spica nutante longissima. *Park.* 1162.

Anglis, wall Fescue-grass.

Habitat in muris et locis agrorum sterilioribus et siccioribus. ☉. VI.

fluitans.

7. **FESTUCA** panicula secunda erecta subramosa, ramis solitariis binatisque, spiculis teretibus linearibus muticis; foliis planis; culmo ancipiti. *Fl. dan.* 237.

Festuca panicula ramosa erecta, spiculis subsessilibus teretibus muticis. *Sp. pl.* 111. *Schreb. gram.* 37. *t.* 3.

Poa locustis teretibus multifloris, glumis floralibus exterioribus truncatis, interioribus bifidis. *Hall. hist.* 1453.

Gramen fluviale. *Ger. em.* 14. *Park.* 1275.

Gramen aquaticum fluitans, multiplici spica. *B. pin.* 2.

Gramen aquaticum cum longissima panicula. *B. hist. II.* 490. *R. syn.* 412.

Festuca

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Gramen avenaceum parvum procumbens, paniculis non aristatis. *R. syn.* 408. *Pluk. ph. t.* 34. *f.* 1.

Gramen triticeum palustre, humilius, spica mutica brevior. *Hist. Ox. III.* 177. *t.* 1. *f.* 6.

Anglis, decumbent Fescue-grass.

Habitat in pascuis sterilibus subhumidis. 4. VIII.

pinnata. 10. FESTUCA spicata, subdisticha; spiculis alternis, teretibus, aristatis.

Bromus culmo indiviso, spicis alternis subsessilibus teretibus flosculis arista longioribus. *Fl. angl.* 41.

Bromus culmo indiviso, spicis alternis, subsessilibus teretibus. *Fl. suec.* 100. *Sp. pl.* 115.

Triticum spicis distichis, locustis teretibus hirsutis, aristatis, duodecimfloris. *Hall. hist.* 1431.

Gramen spica brizæ majus, *B. pin.* 9. *pr.* 19. *th.* 133. *R. syn.* 392.

sylvatica. β Festuca spicata, spiculis alternis subdistichis sessilibus aristatis. *Fl. angl.* 38. *Danic.* 164.

Triticum locustis remotis, hirsutis, teretibus exiguis, culmis floriferis gracilibus. *Hall. hist.* 1432.

Festuca dumetorum. *B. pr.* 69. *th.* 148.

Gramen avenaceum dumetorum spicatum. *R. syn.* 394. *Anglis*, spiked Fescue-grass.

Habitat in pratis, solis cretaceo et calcareo: β in sylvis et sepibus. 4. VII.

Obs. α. Planta glabra, luteo-viridis, Folia inferiora subinde parum pilosa, arista flosculis brevior, β. Planta pilosa arista flosculis longior. Flores in α. et β. α. sex ad duodecim.

B R O M U S. 89.

polymor-
phus. Cal. bivalvis. Spicula oblonga, teres, disticha; arista infra apicem.

mollis. 1. BROMUS panicula erectiuscula subsimplici, spiculis ovatis, floribus obtusis, aristis rectis.

α Bromus panicula erectiuscula, spicis ovatis aristis rectis, foliis mollissime villosis. *Sp. pl.* 112. *Schreb. gram.* 60. *t.* 6. *f.* 2.

Bromus hirsutus, locustis septifloris, ovata-conicis. *Hall. hist.* 1504.

Festuca

Festuca avenacea hirsuta, paniculis minus sparsis. *R. syn.* 414.

Gramen avenaceum pratense, panicula squamata et villosa. *Hist. Ox. III.* 213. t. 7. f. 18.

β *Festuca avenacea* spicis strigosioribus, e glumis glabris compactis. *R. syn.* 414.

γ *Bromus* panicula patente, spiculis ovatis, aristis rectis. *Jecalinus. Sp. pl.* 112.

Bromus foliis glabris, locustis duodecimfloris. *Hall. hist.* 1502.

Festuca avenacea spicis habitioribus, glumis glabris. *R. syn.* 414.

Gramen avenaceum segetale majus, gluma turgidiore. *Hist. Ox. 3. III.* 212. t. 7. f. 16.

δ *Gramen avenaceum pratense*, gluma tenuiore glabra. *Hist. Ox. III.* 213. t. 7. f. 19.

Anglis, field Brome-grass.

Habitat in pratis et pascuis, γ. δ. in arvis. ☉. V. VI.

Obs. α. Tota plana mollissime hirsuta. β. Foliis hirsutis, panicula glabriuscula. γ. Tota planta glabra. δ. Foliis inferioribus parum hirsutis, panicula glabra.

2. BROMUS panicula simplici subnutante, spicis ova-squarrosus, floribus obtusis; aristis divaricatis.

Bromus panicula nutante, spicis ovatis: aristis divaricatis. *Sp. pl.* 112.

Festuca graminea, glumis vacuis. *Sch. agr.* 251. t. 5. f. 11.

Gramen phalaroides acerosum, nutante spica. *Bar. ic.* 24. f. 1.

Anglis, corn Brome-grass.

Habitat in arvis prope Glastenbury, in comitatu Somersetensi et Marshfield in Suffexia. ☉. VII.

3. BROMUS panicula subsimplici erecta, spiculis ovato-erectus, oblongis, floribus acuminato-aristatis glabris.

Bromus racemo simplicissimo, pedunculis unifloris, floribus sublexfloris lævibus aristatis. *Sp. pl.* 114?

Festuca avenacea sterilis spicis erectis. *R. syn.* 413. *Pluk. ph.* t. 299. f. 2.

Festuca avenacea sterilis, pediculis brevibus et spicis erectis. *Hist. Ox. III.* 212. t. 7. f. 13.

arvensis. β Bromus panicula nutante, spiculis ovato-oblongis.
Sp. pl. 113. *Fl. succ.* 97?

Festuca elatior paniculis minus sparsis, locustis oblongis strigosis, aristis purpureis splendentibus. *R. syn.* 414.

Anglis, upright Brome-grass.

Habitat in pratis et pascuis cretaceis frequens β . in arvis. γ . VI. VII.

Obi. Folia plana, et in locis siccioribus involuta, pilosa; Panicula simplex β basi subramosa glabra, Spiculis ovato-oblongis levibus, flosculi u quinque ad octo aristati; aristis rectis brevibus.

muralis. 4. BROMUS panicula patulo-erecta simplici, spiculis linearibus, floribus acuminatis scabris, aristis longis.

madri- Bromus panicula patulo-erecta, spiculis linearibus: ten- intermediis geminis, pedicellis incrassatis. *Amen. ac.* IV. 265. *Sp. pl.* 114.

Bromus sterilis, erecta panicula major. *Barr. ic.* 76. f. 1.

Gramen bromoides pumilum, locustis erectis majoribus aristatis. *Sch. agr.* 260.

Anglis, wall Brome-grass.

Habitat in muris antiquis; circa Londinum et Oxonium. \odot . V.

Desc. Radix annua, fibrosa; Culmi plures pedales erecti, teretes, villosi. Folia petiolata, ensiformia, villosa; Petiolus vaginans, villosus. Stipula membranacea, obtusa. Panicula subsimplex, parte florente patula, deflorata erecta. Pedunculi subunispiculi, scabri. Spiculæ lineares; Gluma multiflora, bivalvis, inæqualis carinata, carina scabra. Corolla bivalvis, inæqualis; Valvula exterior, aristata, villososcabra, acuta carinata, arista longa, scabra. Valvula interior oblonga, ciliata, obtusa, plana; variat colore, viridi et rubente.

Conf. *Bromus rubens?*

sterilis. 5. BROMUS panicula nutante ramosa, ramis quaternis, spiculis oblongis, glumis subulato-aristatis.

Bromus panicula patula, spiculis oblongis distichis, glumis subulato-aristatis. *Sp. pl.* 113.

Festuca

Festuca avenacea sterilis elatior seu Bromos Dioscoridis. B. pin. 9. Th. 146. Hist. Ox. III. 212. t. 7. f. 11. R. syn. 412.

Bromos herba sive avena sterilis. Park. 1147.

Bromos sterilis. Ger. em. 76.

Anglis, barren Brome-grass.

Habitat in sylvis, sepibus et ad margines agrorum frequens. ☉. VI. VII.

6. **BROMUS** panicula ramosa nutante, ramis binatis, *giganteus*.
spiculis subquadrifloris, arista brevioribus.

Bromus panicula nutante, spiculis quadrifloris aristis brevibus. Fl. suec. 99. It. scan. 394. Sp. pl. 114. Sch. gr. 88. t. 11.

Bromus glaber, locustis quadrifloris nutantibus, aristis longissimis. Hall. hist. 1510.

Gramen sylvaticum glabrum panicula recurva. Vaill. paris. 93. t. 18. f. 3.

Gramen avenaceum glabrum, panicula e spicis raris strigosis composita, aristis tenuissimis. R. syn. 415.

Gramen bromoides aquaticum latifolium, panicula sparsa tenuissime aristata. Sch. agr. 264. t. 5. f. 17.

Anglis, tall Brome-grass.

Habitat in sylvis et sepibus subhumidis. ☿. VII. VIII.

Obs. habet multa communia cum sequente; an species distincta?

BROMUS panicula ramosa nutante scabra, ramis bi-nemoralis.
natis, spiculis linearibus teretibus arista longioribus.

Bromus foliis hirsutis, per oras asperrimis, locustis glabris, teretibus, novemfloris. Hall. hist. 1503.

Festuca graminea nemoralis latifolia mollis. B. pin. 9. Th. 144.

Gramen avenaceum dumetorum, panicula sparsa. R. syn. 415.

Gramen avenaceum dumetorum, juba longiore, spica divisa. Hist. Ox. III. 213. t. 7. f. 27.

Anglis, wood Brome-grass.

Habitat in sylvis et sepibus frequens. ☉. VII. VIII.

A V E N A. 91.

Cal. bivalvis, multiflorus; arista dorsali contorta.

- nuda.* 1. AVENA paniculata, calycibus trifloris, receptaculo calycem excedente, petalis dorso aristatis. *Sp. pl.* 118. Avena nuda. *B. pin.* 23. *hist.* II. 433. *Ger. em.* 75. *Park.* 1134. *R. syn.* 389. *Anglis*, naked Oats or Pilcorn. *Habitat in arvis sed rarius. In extrema Cornubia prope Belerium, promontorium copiose feritur. R. syn. in Staffordshire, Dr. Plot. ☉. VII.*
- satua.* 2. AVENA paniculata, calycibus trifloris, flosculis basi pilosis: aristis totis lævibus. *Sp. pl.* 118. *Fl. suec.* 101. Avena sylvestris pilosa, aristis recurvis. *Hist. Ox.* III. 209. t. 7. f. 5. Avena triantha, locustis patulis, folliculis villosis. *Hall. hist.* 1495. Festuca utriculis lanugine flavescens. *B. pin.* 10. Ægilops quibusdam aristis recurvis, f. Avena pilosa. *B. hist.* II. 433. *R. syn.* 489. Ægilops bromoides. *Ger. em.* 77. *Belgarum. Park.* 1148. *Anglis*, bearded Oat-grass. *Habitat in arvis. ☉. VII. VIII.*
- pratensis.* 3. AVENA panicula simplici, glumis quinquefloris, foliis involutis nudis. Avena subspicata, calycibus quinquefloris. *Sp. pl.* 119. Avena periolis unifloris, brevibus, locustis teretibus quinquefloris, basi pubescentibus. *Hall. hist.* 1499. Gramen avenaceum montanum, spica simplici, aristis recurvis. *R. syn.* 405. t. 21. f. 1. *Anglis*, meadow Oat-grass. *Habitat in ericetis, pratis montosis præsertim solo cretaceo. 4. VII.*
- pubescens.* 4. AVENA panicula ramosa, glumis subtrifloris, receptaculis barbatis, foliis planis pubescentibus. Avena calycibus subtrifloris basi pilosis, foliis planis pubescentibus. *Fl. angl.* 42. *Sp. pl.* 1165. Avena triantha, locustis teretibus erectis, petalis ferricis; gluma exteriori lacera. *Hall. hist.* 1498. Gramen avenaceum hirsutum, panicula purpuro-argentea splendente. *R. syn.* 406. t. 21. f. 2. *Anglis*,

Anglis, rough Oat-grass.

Habitat in pratis et pascuis cretaceis, prope Northfleet et supra Banstead Downs copiose. 4. VI.

5. AVENA panicula laxa, calycibus trifloris brevibus, *flavescens*. flosculis omnibus aristatis. *Sp. pl.* 118. *Schreb. gram.* 76. t. 9.

Avena triantha, locustis teretibus, calycina gluma altera minima, petiolo villosa. *Hall. hist.* 1497.

Gramen avenaceum pratense elatius, panicula flavescente, locustis parvis. *R. syn.* 407. *Sch. agr.* 223.

Gramen avenaceum, spica sparsa flavescente, locustis parvis. *Hist. Ox. III.* 215. t. 7. f. 24.

Anglis, yellow Oat-grass.

Habitat in pratis et pascuis. 4. VII.

Obs. variat gluma biflora.

6. AVENA paniculata, calycibus bifloris, flosculo hermaphrodito submutico, masculino aristato. *Sp. pl.* 116. *Schreb. gram.* 25. t. 1.

Avena diantha, folliculis basi villosis, majoris arista geniculata. *Hall. hist.* 1492.

Gramen caninum nodosum. *Ger. em.* 23. f. bulbosum vulgare. *Park.* 1175.

Gramen nodosum avenacea panicula. *B. pin.* 2. pr. 3. *R. syn.* 406.

Gramen avenaceum elatius, juba longa splendente. *R. meth.* 179.

Anglis, tall Oat-grass.

Habitat in pratis et pascuis. 4. VI—VIII.

ARUNDO. 93.

Cal. bivalvis. Flosculi congesti, lana cincti.

1. ARUNDO calycibus quinquefloris, panicula laxa. *phragmites*. *Roy. lugdb.* 66. *Gren. virg.* 137. *Sp. pl.* 120.

Arundo foliis secantibus, locustis trifloris, papposis, muticis. *Hall. hist.* 1515.

Arundo vulgaris f. *phragmites* Dioscoridis. *B. pin.* 17.

Arundo vallatoria. *Ger. em.* 36. *R. syn.* 401.

Arundo vulgaris sive *vallatoria*. *Park.* 1208.

Anglis, common Reed-grafs.

Habitat in fluviis, lacubus, fossis. 4. VII.

Calamagrostis.

2. ARUNDO calycibus unifloris lævibus, corollis lanuginosis, culmo ramoso. *Sp. pl.* 120. *Fl. dan.* 280.

Arundo locustis unifloris, muticis, panicula alterne contracta. *Hall. hist.* 1519.

Gramen arundinaceum, panicula molli spadicea, majus. *B. pin.* 7. *Sch. agr. pr. t.* 5. *R. syn.* 401.

Gramen tomentosum arundinaceum. *Ger. em.* 9.

Calamagrostis f. gramen tomentosum. *Park.* 1182.

Anglis, wood Reed-grafs.

Habitat in sylvis et sepibus subhumidis. Inter Newington et Hornsey. 4. VI. VII.

Obs. nunquam vidi culmo ramoso.

epigejos.

3. ARUNDO calycibus unifloris, panicula erecta, foliis subtus glabris. *Sp. pl.* 120. *Fl. suec.* 106.

Arundo locustis unifloris, sericeis, muticis, panicula stricta. *Hall. hist.* 1520.

Gramen arundinaceum paniculatum montanum, panicula spadiceo-viridi, semine papposo. *Sch. agr.* 124.

Calamagrostis minor glumis rufis et viridibus. *Pet. conc. gr. n.* 69. *R. syn.* 401.

Anglis, small Reed-grafs.

Habitat in sylvis et sepibus humidis, circa Oundle in comitatu Leicestrensi. *R. syn.* circa Ranaugh in Norfolk. 4. VI. VII.

arenaria.

4. ARUNDO calycibus unifloris, foliis involutis mucronato-pungentibus. *Sp. pl.* 121.

Gramen spartheum spicatum, foliis mucronatis longioribus vel spica fecalina. *B. pin.* 5. *th.* 67. *R. syn.* 393.

Spartum marinum nostras. *Park.* 1198.

Spartum anglicanum. *Ger. em.* 42.

Anglis, sea Reed-grafs.

Habitat in littoribus maritimis arenosis. 4. VI. VII.

LOLIUM.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

tis, scabris. Gluma univalvis, linearis, concava, aristata. Corolla bivalvis, inæqualis: Valvula exterior major linearis, carinata, aristata, scabra, Valvula interior brevior, linearis, plana, mutica. Arista flosculo duplo longior.

E L Y M U S. 96.

Cal. lateralis, bivalvis, aggregatus, multiflorus.

arenari- 1. ELYMUS spica erecta arcta, calycibus tomentosis
us. flosculo longioribus. *Spec. pl. 122.*

Elymus foliis mucronato-pungentibus. It. scan. 336.

Gramen sparteum spicatum latifolium. B. pin. 5. th. 67?

Spartum herba IV. Batavicum. Clus. hist. 221?

Anglis, sea Lyme-grass.

Habitat in littoribus maritimis, in arena mobili, in Insula Bute; et prope Exmouth, in Devoniam. 4. VH. VIII.

H O R D E U M. 98.

Cal. lateralis, bivalvis, uniflorus, ternus.

murinum. 1. HORDEUM flosculis lateralibus masculis aristatis, involucris intermediis ciliatis. *Sp. pl. 126.*

Hordeum spica crassa, longe aristata, calycinis glumis aristatis. Hall. hist. 1536.

Hordeum spureum vulgare. Park. 1147.

Gramen secalinum et secale sylvestre. Ger. em. 73. R. syn. 391.

Gramen hordeaceum minus et vulgare. B. pin. 9.

Anglis, wall Barley-grass, or Way-bennet.

Habitat ad vias et muros. 0. IV—VIII.

pratense. 2. HORDEUM flosculis lateralibus masculis muticis, involucris setaceis scabris. *Sp. pl. 126.*

Hordeum spica disticha, folliculo brevi, glumis calycinis aristatis. Hall. hist. 1538.

Gramen secalinum. Ger. em. 73. R. syn. 392. majus et minus. Park. 1144.

Gramen secalinum minus pratense. Hist. Ox. III. 179. t. 2. f. 6.

Anglis,

Anglis, meadow Barley-grass.

Habitat in pratis et pascuis humidiusculis. 4. VI.

3. HORDEUM flosculis lateralibus masculis aristatis, *marinum*.
involucro interno florum lateralium semiovato.

Gramen secalinum palustre et maritimum. *R. syn.* 392.

Gramen secalinum maritimum minus. *Hist. Ox. III.*

179. t. 6. f. 5.

Anglis, sea Barley-grass.

Habitat in pratis et pascuis. ☉. V. VI.

4. HORDEUM flosculis omnibus hermaphroditis aris-*sylvatici-*
tatis, involucris basi connatis arista brevioribus. *cum.*

Elymus spica erecta: spiculis bifloris involucro æqua-*europæ-*
libus. *Mantiss. I.* 35. *us.*

Gramen secalinum majus sylvaticum. *R. syn.* 392.

Hist. Ox. III. 180.

Gramen secalinum maximum. *Park.* 1144. *Merr.*

Gramen hordeaceum montanum s. majus. *B. pin.* 9.

Gramen hordeaceum montanum, spica strigosiore,
brevibus aristata. *Sch. agr.* 16. *pr. t. I. f. bona.*

Anglis, wood Barley-grass.

Habitat in sylvis, solo calcareo. In Stoken-Church

Woods, Oxfordshire, circa Marlow et Henley.

4. VI.

Obs. Pedicellus in omnibus speciebus elongatus in sulco longitudinali florum reconditus, ut alteram foliolam involucris apparct, et subinde florem abortivum sustinet.

TRITICUM. 99.

Cal. bivalvis, solitarius, subtriflorus. Flos obtusiusculus, acutus.

1. TRITICUM calycibus subulatis multifloris, floscu-*repens.*
lis mucronatis.

Triticum calycibus quadrifloris subulatis acuminatis
foliis planis. *Sp. pl.* 128. *Schreb. gram. II.* 24.
t. 26.

Gramen caninum. *Ger. em.* 23. vulgaris. *Park.* 1173.

Hist. Ox. III. 178. *t. I. f. 8.*

Gramen caninum arvense. s. Gramen Dioscoridis.

B. pin. I. th. 7.

Gramen

TRIANDRIA DIGYNIA.

Gramen spica triticea repens vulgare, caninum dictum. *R. syn.* 390.

Triticum radicibus repentibus vaginosis, foliis hirsutis, locustis quinquefloris. *Hall. hist.* 1426.

β Triticum radicibus repentibus, vaginosis, foliis hirsutis, locustis octofloris. *Hall. hist.* 1427.

γ Gramen caninum maritimum, spica triticea nostras. *R. syn.* 390.

Anglis, common Wheat-grass; Dog's-grass, Quick-grass, or Couch-grass.

Habitat in agris. γ in littoribus maritimis. 4. VI—IX.

Pharmac. Graminis canini Radix.

Obs. variat calycibus a tribus ad octo flores flosculis acuminatis, in γ mucronato-aristatis et subinde aristatis. Foliis in γ glaucis, rigidis, pungentibus.

caninum. 1. TRITICUM calycibus subulatis subquadri-floris, flosculis aristatis.

Triticum radice fibrosa, foliis hirsutis, locustis quinquefloris aristatis. *Hall. hist.* 1429.

caninus. Elymus spica nutante arcta, spiculis rectis involucre destitutis: infimis geminis. *Sp. pl.* 124. *Fl. suec.* 122.

Gramen caninum aristatum radice non repente, sylvaticum. *R. syn.* 390.

Gramen caninum non repens elatius, spica aristata. *Hist. Ox.* III. 177. t. 1. f. 2.

Anglis, bearded Wheat-grass.

Habitat in sylvis et sepibus, inter Greenwich et Woolwich. 4. VI. VII.

juncum. 3. TRITICUM calycibus truncatis quinquefloris, foliis involutis. *Sp. pl.* 128. *Cent. pl.* 13.

Gramen maritimum, spica loliacea foliis pungentibus nostras. *Pluk. ph.* t. 33. f. 4. *R. syn.* 391.

Gramen phœnicoides foliis convolutis, juncis ac pungentibus alterum. *B. hist.* II. 478.

β Gramen loliaceum maritimum supinum, spica foliacea. *R. syn.* 391.

γ Gramen caninum maritimum, spica foliacea. *R. s.* 391. *Anglis*, sea Wheat-grass.

Habitat in littoribus maritimis passim. 4. VI. VII.

CYNO-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

T R I G Y N I A.

M O N T I A. 101.

Cal. diphyllus. *Cor.* monopetala, irregularis, *Caps.* unilocularis, trivalvis.

fontana. 1. MONTIA. *Sp. pl.* 129. *Fl. lapp.* 30. *Suec.* 115.
Hall. hist. 301. *aquatica minor.* *Mich. gen.* 18. t. 13.
f. 2.

Cameraria arvensis minor. *Dill. giff.* 46.

Alfina parva palustris tricoccus, portulacæ aquaticæ similis. *R. syn.* 352.

Portulaca arvensis. *B. pin.* 282.

Anglis, Water-chickweed.

Habitat ad fontes et scaturigines. ☉. IV. V.

P O L Y C A R P O N. 105.

Cal. pentaphyllus. *Petala* quinque, minima, ovata;
Caps. unilocularis, trivalvis.

tetraphyl- 1. POLYCARPON. *Sp. pl.* 131. *Gouan. Fl. monf.* 211.
lum. *Mollugo foliis quaternis obovatis, paniculis dichotomis.* *Hort. cliff.* 28.

Anthyllis marina alsinifolia. *B. pin.* 282.

Anthyllis alsinifolia polygonoides major. *Barr. car.* 103. t. 534.

Anglis, four-leaved Polycarpon.

Habitat in pratis aridis et locis arenosis, circa Lymston prope Exeter, in Devonia; et in Insula Portlandica.

☉. V—VIII.

Obs. variat floribus a petalis.

*Classis IV.*TETRANDRIA
MONOGYNIA.

* *Flores monopetali, monospermi, superi. Aggregati.*

DIPSACUS. 114.

Cal. communis polyphyllus; proprius superus. Receptaculum paleaceum.

1. DIPSACUS foliis sessilibus serratis. *Sp. pl.* 140. *fullonum.*
 Dipsacus foliis sessilibus serratis, aristis fructus erectis. *sylvestris.*
Fl. angl. 49.
 Dipsacus capitulis ovatis, foliis arcuatis circumvallatis,
 aristis squamarum rectis. *Hall. hist.* 198.
 Dipsacus sylvestris. *Ger. em.* 1167. *Park.* 984. *R. syn.*
 192. *aut* Virga pastoris major. *B. pin.* 385.
- β Dipsacus foliis sessilibus serratis, aristis fructus recurvis. *fativus.*
Fl. angl. 49.
 Dipsacus fativus. *Ger. em.* 1167. *Park.* 983. *B. pin.*
 385. *R. syn.* 192.
 Carduus fullonum. *Lob. ic.* 17.
Anglis, wild β manured Teasel.
 Habitat *ad sepes.* ♂. VII.
Pharmac. Dipsaci fativi *Radix.*
2. DIPSACUS foliis petiolatis appendiculatis. *Sp. pl. pilosus.*
 141.
 Dipsacus foliis biauribus, capitulis hemisphæricis.
Hall. hist. 199.
 Dipsacus sylvestris capitulo minore, vel Virga pastoris
 minor. *B. pin.* 385.
 Dipsacus minor s. virga pastoris. *Ger. em.* 1168. *R.*
syn. 192.
 Virga Pastoris. *Park.* 984. *Cam. epit.* 423.
Anglis,

Anglis, Small Teasel.

Habitat in aquosis et ad sepes. 4. VIII.

S C A B I O S A. 115.

Cal. communis polyphyllus; proprius duplex superus.

Receptaculum paleaceum s. nudum.

succisa.

1. SCABIOSA corollulis quadrifidis æqualibus, caule simplici, ramis approximatis, foliis lanceolato-ovatis integerrimis. *Mat. Med. 41. Sp. pl. 142. Fl. dan. 279.*

Succisa caule trifloro, floribus convexis, foliis radicalibus ovatis, caulinis lanceolatis. Hall. hist. 201.

Succisa glabra et hirsuta. B. pin. 269.

Scabiosa radice succisa, flore globofo. R. syn. 191.

Morsus Diaboli. Ger. em. 726. vulgaris flore purpureo. Park. 491.

Anglis, Devil's-bit Scabious.

Habitat in pratis, pascuis et sylvis. 4. VI—VIII.

arvensis.

2. SCABIOSA corollulis quadrifidis radiantibus, caule hispido, foliis pinnatifidis: lobis distantibus. *Sp. pl. 143. Fl. suec. 117. dan. 447.*

Scabiosa foliis petiolatis, ovato-lanceolatis dentatis, superioribus semipinnatis. Hall. hist. 206.

Scabiosa major vulgaris. Ger. em. 719.

Scabiosa vulgaris pratensis. Paris. 484.

Scabiosa major communior folio laciniato. B. hist. III. 2. R. syn. 191.

Anglis, field Scabious.

Habitat in pratis et arvis. 4. VIII.

Pharmac. Scabiosæ Herba.

columbaria.

3. SCABIOSA corollulis quinquefidis radiantibus, foliis radicalibus ovatis, crenatis, caulinis pinnatis setaceis. *Sp. pl. 143. Fl. suec. 118. dan. 314.*

Scabiosa minor. Cam. epit. 711. s. Columbaria. Ger. em. 719.

Scabiosa minor campestris. Park. 484. R. syn. 191.

Scabiosa capitulo globofo major et minor. B. pin. 270.

Succisa foenis imis ovatis, superioribus pinnatis, pinnis semipinnatis acutis, ciliis flosculorum longitudine.

Hall. hist. 202.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

scapo tereti, spica cylindrica. *Sp. pl.* 163. *Hall.* 472.
Fl. dan. 581.

Plantago major incana. *Park.* 493. *R. syn.* 314.

Plantago incana. *Ger. em.* 419.

Plantago latifolia incana. *B. pin.* 189.

Anglis, hoary Plantain,

Habitat in pascuis et pratis, solo calcareo. μ . V—VIII.

lanceolata. 3. PLANTAGO foliis lanceolatis, spica subovata nuda,
scapo angulato. *Sp. pl.* 164. *Hall. hist.* 656. *Fl. dan.*
437.

Plantago quinquenervia. *Ger. em.* 422.—*R. syn.* 314.
major. *Park.* 495.

Plantago angustifolia major. *B. pin.* 189.

β Plantago angustifolia minor. *Tab. ic.* 732. *Ger.* 420.
Anglis, Ribwort-plantain.

Habitat in pratis et pascuis. μ . VI.

maritima. 4. PLANTAGO foliis linearibus subintegerrimis basi
lanatis, spica cylindrica, scapo tereti.

Plantago foliis semicylindraceutis integerrimis basi la-
natis, scapo tereti. *Sp. pl.* 165. *Fl. dan.* 243.

Coronopus. *Ger. em.* 425. *maritima* major. *B. pin.*
190.

læffingia. β Plantago foliis linearibus subdentatis, scapo tereti,
spica cylindrica, bracteis carinatis membranaceis.
Fl. angl. 52. *Sp. pl.* 166. *Læff. hisp.* 124.

Plantago marina. *Ger. em.* 423. *R. syn.* 315. *vulgaris.*
Park. 498.

γ Plantago alpina angustifolia. *B. hist.* III. 506. *R. syn.*
315.

Anglis, sea, β narrow, γ mountain Plantain.

Habitat in littoribus maritimis. γ in rupibus Tryghfyl-
chau prope Llanberis *R. syn.* in montibus prope Set-
tle in comitatu Eboracensi. μ . VII. VIII.

Obs. Folia in α linearia crassa subinde subtus convexa.
in γ Folia subinde rariter dentata in δ rigidiuscula acu-
ta et in aridis involuta. Spica variat in aridis oblonga
et in plantis minimis subrotunda.

Corono- PLANTAGO foliis linearibus pinnato-dentatis, scapo
pus. tereti. *Fl. angl.* 52. *Dan.* 272.

Plantago

Plantago foliis linearibus dentatis, scapo tereti. Sp. pl.
166. *Fl. suec.* 134.

Plantago foliis subhirsutis, sempinnatis, pinnis raris lanceolatis. Hall. hist. 658.

Plantago foliis laciniatis, Coronopus dicta. R. syn.
315.

Coronopus vulgaris f. *Cornu cervinum. Park.* 502.

Coronopus sylvestris hirsutior. B. pin. 190.

Cornu Cervinum. Ger. em. 316.

β *Plantago coronopus dicta, foliis hirsutioribus magis laciniatis, capitulis brevibus laxis. Blackf. spec.* 76.

γ *Plantago graminea folio hirsuto minor, capitulo rotundo brevi. R. syn.* 316.

Anglis, Buckshorn-plantain, Star of the Earth.

Habitat in glareosis, γ in pratis aridis maritimis. ☉.
VI—VIII.

Pharmac. Coronopi Herba.

Obs. Folia in γ. linearia integerrima, hirsuta, spica subrotunda.

*** *Flores monopetali, monocarpi superi.*

SANGUISORBA. 146.

Cal. diphyllus. Germen inter calycem corollamque.

1. *SANGUISORBA spicis ovatis. Sp. pl.* 169. *Fl. officinalis: dan.* 197.

Sanguisorba major flore spadiceo. B. hist. III. 120.
R. syn. 203.

Pimpinella tetrastemon, spica brevi. Hall. hist. 705.

Pimpinella sylvestris. Ger. em. 1045.

Pimpinella major vulgaris. Park. 582.

Pimpinella sanguisorba, major. B. pin. 160.

Anglis, meadow sanguisorba, or Burnet-saxifrage.

Habitat in pratis et pascuis subhumidis. 4. VII. VIII.

**** *Flores monopetali, dicocci, superi. Stellatæ.*

RUBIA. 127.

Cor. monopetala, campanulata. Baccæ duæ, monopermæ.

1. *RUBIA foliis perennantibus ellipticis, supra lævibus. peregrina. Syst. nat.* 119.

Rubia foliis quaternis. Sp. pl. 158. *Fl. angl.* 54.

Rubia foliis ellipticis, asperis, quinis senisque. *Hall. hist.* 708.

Rubia sylvestris. *Park.* 274.

Rubia sylvestris aspera, quæ sylvestris Dioscoridis. *R. syn.* 223.

Anglis, wild Madder.

Habitat in rupe S. Vincentii prope Bristolium et in Devonía frequens. *R. syn.* prope Exmouth copiose. 4. VI. VII.

Obs. variat Foliis a quatuor ad octo lanceolatis. Corolla pentaphylla; Stamina quinque fida; nimis affinis Rubiæ tinctorum. an vere distincta sit species?

SHERARDIA. 120.

Cor. monopetala infundibuliformis. Semina duo, tridentata.

arvensis. 1. SHERARDIA foliis omnibus verticillatis, floribus terminalibus. *Sp. pl.* 149. *Fl. dan.* 439.

Sherardia foliis senis, lanceolatis, floribus sessilibus, umbellatis. *Hall. hist.* 734.

Rubeola arvensis repens cærulea. *B. pin.* 334. *R. syn.* 225.

Rubia minor pratensis cærulea. *Park.* 276.

Anglis, little Sherardia or Field-madder.

Habitat in arvis. ☉. V—VII.

ASPERULA. 121.

Cor. monopetala, infundibuliformis, sem. duo globosa.

odorata. 1. ASPERULA foliis octonis lanceolatis, florum fasciculis pedunculatis. *Sp. pl.* 150. *Dalib. paris.* 46. *F. dan.* 562.

Asperula caule erecto, foliis octonis; petiolis ramosis erectis, seminibus hirsutis. *Hall. hist.* 728.

Asperula. *Ger. em. II.* 24. *R. syn.* 224. *odorata.*

Park. 563. flore albo. *Dod. pempt.* 355.

Asperula s. Rubeola montana odorata. *B. pin.* 334.

Anglis, sweet Woodroof.

Habitat in sylvis et umbrosis. 4. V.

Pharmac. Asperulæ odoratæ Herbae.

cynanchi- 2. ASPERULA foliis quarternis linearibus: superioribus
ca. oppositis;

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- spurium.* 3. **GALIUM** foliis senis lanceolatis carinatis scabris retrorsum aculeatis, geniculis simplicibus, fructibus glabris. *Sp. pl.* 154
 Aparine semine læviore. *R. syn.* 225. *hist.* 484.
 Aparine lævis. *Park.* 567.
Anglis, corn Ladies Bedstraw.
 Habitat in arvis in Insula Thanet. prope Leatherhead et alibi in comitatu Surrensi. et prope Stamford. ☉. VI. VII.
- uüginosum.* 4. **GALIUM** foliis senis lanceolatis retrorsum serrato-aculeatis mucronatis rigidis, corollis fructu majoribus. *Fl. suec.* 127. *Sp. pl.* 153.
 Aparine palustris minor parisiensis, flore albo. *Vaill. par.* 14. *R. syn.* 225.
 Rubia quædam minor. *B. hist.* III. 2. 716.
Anglis, marsh Ladies Bedstraw.
 Habitat in pratis, pascuis et ericetis humidiusculis. ☿. VII.
 Obs. Variat foliis septenis et octonis.
- erectum.* 5. **GALIUM** foliis suboctonis lanceolatis aculeato-serrulatis mucronatis, paniculis trichotomis, caule scabro flaccido.
 Galium austriacum. *Jac. Fl. austr.* 1. 51. t. 80.
Anglis, upright Ladies Bedstraw.
 Habitat in pratis et pascuis subhumidis. ☿. VI. VII.
 Desc. Radix perennis ramosa; Caules plures flaccidi, erectiusculi, geniculati: geniculis tumidis; quadranguli angulis scabriusculis; parum hirsuti, ramosi, ramis oppositis floriferis. Folia verticillata, octona, subinde sena, sessilia lanceolata et lineari-lanceolata, aculeato-serrulata, mucronata, nuda. Panicula terminalis trichotoma. Flores albi quadrifidi. Semina glabra, parva.
- Mollugo.* 6. **GALIUM** foliis octonis ovato-linearibus subserratis patentissimis mucronatis, caule flaccido, ramis patentibus. *Sp. pl.* 155. *Fl. dan.* 455.
 Galium foliis octonis ellipticis, caule anguloso. *Hall. hist.* 711. *Scop. earn.* 104.
 Mollugo vulgator. *Park.* 565. *R. syn.* 223.
 Rubia sylvestris. *Ger. em.* 1118. lævis. *B. pin.* 333.
Anglis,

Anglis, great Ladies Bedstraw, or bastard Madder.
Habitat in *Jepibus et ericetis frequens.* 4. VI.—VII.

7. GALIUM foliis octonis hispidis linearibus acuminatis subimbricatis, pedunculis dichotomis. *Sp. pl.* 154.
Rubeola saxatilis. *B. pin.* 334. *pr.* 145.
Aparine minima s. Rubia saxatilis minima. *Mag. mons.* 291.

Anglis, least Ladies Bedstraw.

Habitat in montibus calcareis. prope Kendal, in comitatu Westmorlandico. 4. VIII.

Desc. Radix ramosa perennis. Caules plures, digitales, ramosi, quadranguli, angulis aculeato-scabris; Folia verticillata octona et septena lanceolata et linearia, scabra, inferioribus imbricatis. Panicula subdichotoma. Flores albi. Semina parva glabra. affinis Galio uliginoso.

8. GALIUM foliis subsenis, lanceolatis, aculeato-ciliatis acuminatis, reflexis, caulibus diffusis retrorsum aculeato-scabris, fructibus glabris.

Aparine minima. *R. syn.* 225. t. 9. f. 1.

Anglis, small Ladies Bedstraw, or Goose-grass.

Habitat in muris et pratis arenosis. inter Dartford et Northfleet, et supra murum in oppido Farningham in agro Cantiano. 6. VI. VII.

Desc. Radix annua ramosa. Caules pedales et sesquipedales diffusi, tetragoni, angulis retrorsum aculeato-scabris; geniculati ramosi, ramis alternis, floriferis oppositis. Folia verticillata sena et septena reflexa lanceolata et lineari-lanceolata, margine aculeato-ciliata, acuminata, nuda. Panicula terminalis, pedunculi trichotomi, Flores viridi-lutei, tetrapetali, parvi. Semina duo, corolla majora, glabra.

9. GALIUM foliis octonis linearibus sulcatis, ramis verum, floriferis brevibus. *Sp. pl.* 155. *Mat. med.* 46.

Galium foliis linearibus perangustis, octonis, racemis multifloris spicatis. *Hall. hist.* 710.

Galium luteum. *Ger. em.* 1126. *Park.* 565. *B. pin.* 335. *R. syn.* 224. *Mill. Ic.* 125.

Anglis, yellow Ladies Bedstraw, or Cheese-rening.

TETRANDRIA MONOGYNIA.

Habitat in pratis et ad margines viarum agrorumque frequens. 4. VII. VIII.

Pharmac. Galii Herba.

** Fructu hispido.

- boreale.* 10. GALIUM foliis quaternis lanceolatis trinerviis glabris, caule suberecto, seminibus hispidis. *Sp. pl.* 156. *Hall. hist.* 722. *Scop. carn.* 102.
 Rubia pratensis lævis, acuto folio. *B. pin.* 333.
 Mollugo montana erecta quadrifolia. *R. syn.* 224.
Anglis, cross-leaved Ladies Bedstraw.
 Habitat in pratis montosis, in Westmorlandia et Cambria, prope Kirby-Lonsdale, supra Pontem, in rupibus prope Llanberys. 4. VII. VIII.

- aparine.* 11. GALIUM foliis octonis lanceolatis carinatis scabris retrorsum aculeatis, geniculis villosis, fructu hispido. *Sp. pl.* 157. *Fl. dan.* 495.
 Galium caule ferrato, foliis senis linearibus, ferratis, petiolis unifloris. *Hall. hist.* 723.
 Aparine. *Ger. em.* 1122. *Dod. pempt.* 353. *R. syn.* 225. vulgaris. *Park.* 567. *B. pin.* 334.
Anglis, common Ladies Bedstraw, Goose-grass or Cleavers.
 Habitat in cultis et ruderatis frequens. ☉. V—VIII,
 Pharmac. Aparines Herba.

Flores tetrapetali superi.

C O R N U S. 149.

Involutum tetraphyllum sæpius. *Petala* supera quatuor. *Drupa* nucleo biloculari.

- sanguinea.* 1. CORNUS arborea, cymis nudis. *It. W-goth.* 32. *Sp. pl.* 171. *Hall. hist.* 816. *Fl. dan.* 481.
 Cornus femina. *Ger. em.* 1467. *Park.* 1521. *B. pin.* 447. *R. syn.* 460.
Anglis, female Cornel, Dog-berry, Gatter or Prickwood Tree.
 Habitat in sylvis et sepibus. ♀. VI.

2. COR-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

TETRANDRIA DIGYNIA.

landia, in vallicula umbrosa Longledale dicta. D. Richardson. *R. syn.* 4. VII.

APHANES. 166.

Cal. quadrifidus. *Cor.* O. *Pist.* unicum. *Sem.* duo, nuda.

arvensis. 1. APHANES. *Fl. sæc.* 143. *Sp. pl.* 179. *Gron. vir.* 17.

Alchemilla hirsuta, foliis trilobatis, lobis bipartitis et tripartitis. *Hall. hist.* 1569.

Percepier anglorum. *Lob.* 727. *Ger. em.* 1594. *R. syn.* 159.

Polygonum felinoides. *Park.* 449.

Chærophyllo nonnihil similis. *B. pin.* 152.

Anglis, Pursley-pierl.

Habitat in arvis et pascuis glareosis. ☉. V—VIII.

DIGYNIA.

BUFONIA. 168.

Cal. tetraphyllus. *Cor.* tetrapetala. *Caps.* unilocularis disperma.

tenuifolia. 1. BUFONIA. *Sp. pl.* 179. *Sauv. monsp.* 141. *Gouan. Fl. monsp.* 145. *Læst. it.* 44.

Alfene polygonides tenuifolia, flosculis ad longitudinem caulis velut in spicam dispositis. *Pluk. ph.* 1, 75. f. 3. *R. syn.* 346. ●

Herniaria angustissimo gramineo folio erecta. *Magn. hort.* 97. t. 97.

Anglis, small Buffonia.

Habitat in maritimis circa Bostoniam agri Lincolnensis, *R. syn.* ☉. V. VI,

TETRAGYNIA.

SAGINA. 176.

Cal. tetraphyllus. *Petala* quatuor. *Caps.* quadrilobularis, quadrivalvis.

1. SAGINA ramis procumbentibus, petalis brevissimis. *procumbens.*
Sp. pl. 185. *Ard. sp.* II 23. t. 8. f. 2.
 Alfine tetrastemon, foliis connatis, lanceolatis. *Hall. hist.* 861.
 Alfinella muscoso flore repens. *Cat. giff.* 81. *R. syn.* 345.
 Saxifraga anglicana alfine folio. *Ger. em.* 568.
 Caryophyllus minimus muscosus nostras. *Park.* 1340.
 β Sagina caule erectiusculo pubescente, floribus alternis apetalis, apetalis. *Mantiss.* II. 559.
 Sagine caulibus erectis, radice annua, floribus apetalis. *Ard. spec.* II. 22. t. 8. f. 1.
 Saxifraga anglica alinesfolia annua. *Plot. Ox.* t. 9. f. 7. *R. syn.* 345.
 Alfine saxifraga graminifolia, flosculis tetrapetalis herbidis et muscosis. *Pluk. ph.* t. 74. f. 2.
Anglis, procumbent Pearl-work.
 Habitat in muris et pratis arenosis ubique. ☉. V—VIII.
 Obs. variat floribus apetalis.

2. SAGINA caule erecto unifloro triflorove. *Sp. pl.* 185. *erecta.*
 Alfinella foliis caryophylleis. *Cat. giff.* 47. *R. syn.* 344. t. 15. f. 4.
 Alfine verna glabra. *Vaill. paris.* 6. t. 3. f. 2.
Anglis, upright Pearl-wort or least Stickwort.
 Habitat in pascuis et ericetis glareosis. ☉. V.

TILLÆA. 177.

Cal. tripartitus, *Petala* tria, *Caps.* tres, polyspermæ.

1. TILLÆA procumbens. *Sp. pl.* 186. *muscosa*
 Tillæa muscosa annua perfoliata. *Mich. gen.* 22. t. 20.
 Crassula foliis sessilibus connatis, floribus aggregatis in foliorum alis. *Guett. stamp.* I. 97.

Polygonum

Polygonum muscosum minimum. *Bocc. sic. 59? t. 29.*
Anglis, procumbent Tillæa.

Habitat in pratis et ericetis arenosis; in ericetis Drayton
 et Mushold in comitatu Norfolk. D. Rose. circa
 Bury in comitatu Suffolkiensi. D. Cullum. 4. V.
 VI.

POTAMOGETON. 174.

Cal. O. Petala quatuor. Stylus nullus. Sem. quatuor.

natans.

1. POTAMOGETON foliis oblongo-ovatis petiolatis
 natantibus. *Sp. pl. 182. Fl. suec. 145.*

Potamogeton foliis firmis, petiolatis; ovatis natanti-
 bus. *Hall. hist. 843.*

Potamogeton rotundifolium. *B. pin. 193. R. syn. 148.*

Potamogeton latifolium. *Ger. em. 821.*

Fontinalis major latifolia vulgaris. *Park. 1254.*

Anglis, broad-leaved Pondweed.

Habitat in fluviis et stagnis frequens. 4. VII.

*perfolia-
 tum.*

2. POTAMOGETON foliis cordatis amplexicaulibus.
Sp. pl. 182. Fl. suec. 145. dan. 196.

Potamogeton perfoliatum. *R. hist. 181. syn. 149.*

Potamogeton foliis tenuibus, cordatis, amplexicauli-
 bus. *Hall. hist. 845.*

Potamogeton foliis latis splendentibus. *B. pin. 193.*

Potamogeton tertium. *Dod. 582. Ger. em. 122.*

β Potamogeton rotundifolium alterum. *Losel. fl. pr. 205.*
t. 65. Hall. hist. 844.

Anglis, perfoliate Pondweed.

Habitat in fluviis frequens. 4. VI—VIII.

lucens.

3. POTAMOGETON foliis lanceolatis planis in pe-
 tiolos desinentibus. *Sp. pl. 183. Fl. dan. 195.*

Potamogeton foliis tenuibus, longissime lanceolatis.
Hall. hist. 847.

Potamogeton longis acutis foliis. *Ger. em. 822.*

Potamogeton foliis angustis splendentibus. *B. pin. 193.*

Potamogeton aquis immersum folio pellucido, lato,
 oblongo acuto. *R. syn. 148.*

Anglis, shining Pondweed.

Habitat in fluviis et lacubus. 4. VI. VII.

4. POTAMOGETON

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

TETRANDRIA TETRAGYNIA.

Potamogeton caulibus compressis, foliis gramineis lanceolatis, spicis paucifloris. *Hall. hist.* 851.

Potamogiton caule compresso, folio graminis canini. *R. syn.* 149. *Hist.* 189.

Anglis, flat Pondweed.

Habitat in fluviis leniter fluentibus et fossis : prope Cambridge. *R. syn.* circa Stains, Middlesex copiose. 4. VI. VII.

gramineum. 8. POTAMOGETON foliis lineari-lanceolatis alternis sessilibus stipula latioribus. *Sp. pl.* 184. *Fl. dan.* 222.

Potamogeton foliis gramineis, petiolis erectis, vaginis laxis, numerosissimis. *Hall. hist.* 850.

Potamogiton gramineum latiusculum, foliis et ramificationibus densissime stipatis. *Buddl. Hort. Sic. Vol.* 27. *R. syn.* 149. t. 4. f. 3.

Potamogiton gramineum latifolium. *Fl. pr.* 206. t. 66. *R. syn.* 149.

Potamogiton gramineum latum. *Pet. H. br. t.* 5. f. *Anglis*, grassy Pondweed.

Habitat in fluviis et fossis. prope Deptford. 4. VII.

pectinatum. 9. POTAMOGETON foliis setaceis parallelis approximatis distichis. *Sp. pl.* 183. *Fl. suec.* 150.

Potamogiton millefolium, seu foliis gramineis ramosum. *R. syn.* 150.

Potamogiton gramineum ramosum. *B. pin.* 193. pr. 101.

Potamogiton fluviatile longissimo gramineo folio nostras. *Pluk. Mant.* 155. *Amalth.* 177. *R. syn.* 150.

Millefolium tenuifolium. *Ger. em.* 828.

Anglis, fine Pondweed.

Habitat in fluviis. 4. VI. VII.

setaceum. 10. POTAMOGETON foliis lanceolatis oppositis acuminatis. *Roy. lugdb.* 213. *Sp. pl.* 184.

Potamogiton ramosum angustifolium. *B. pin.* 193. *Anglis*, setaceous Pondweed.

Habitat in fossis turfosis in comitatu Lancastriensi. 4. VII. VIII.

marinum. 11. POTAMOGETON foliis linearibus alternis distinctis

tinctis inferne vaginantibus. *Sp. pl.* 184. *Fl. dan.* 186.

Potamogeton caule tereti, foliis linearibus planis, spica multiflora, senescente longissima et interrupta. *Hall. hist.* 853.

Potamogeton ingens gramineo folio longiori. *Vaill. paris. t.* 32. *f.* 3.

Potamogiton maritimum ramosissimum, grandiusculis capitulis, capillaceo folio nostras. *Pluk. ph. t.* 216. *f.* 5. *R. syn.* 150.

Anglis, sea Pondweed. *Pet. H. br. t.* 5. *f.* 13.

Habitat in fossis maritimis. In insula Shepey abunde. 4. VII. VIII.

12. POTAMOGETON foliis linearibus oppositis alternisque distinctis: basi patentibus, caule tereti. *Sp. pl.* 184. *Gouan. Fl. monsp.* 148.

Potamogeton caule tereti, foliis perangustis, spicis paucifloris minimis. *Hall. hist.* 852.

Potamogiton pusillum, gramineo folio, caule tereti. *R. syn.* 150. *hist.* 190.

Potamogiton minimum capillaceo folio. *B. pin.* 193.

Potamogiton pusillum, gramineo folio breviori. *Vaill. paris. t.* 32. *f.* 4.

Anglis, small Pondweed. *Pet. h. br. t.* 5. *f.* 11.

Habitat in fossis et stagnis. 4. VI—VIII.

R U P P I A. 175.

Cal. o. *Cor.* o. *Sem.* quatuor pedicellata.

1. RUPPIA. *It. W-goth.* 186. *Sp. pl.* 184. *Fl. suec.* 154. *maritima.* Buccaferra maritima foliis acutissimis. *Mich. gen.* 72. *t.* 35.

Potamogiton maritimum gramineis longioribus foliis fructu fere umbellato. *Pluk. Alm.* 305. *R. syn.* 135.

Potamogiton maritimum pusillum alterum, seminibus singulis longis pediculis insidentibus. *Pluk. ph. t.* 248. *f.* 4.

Anglis, sea Ruppia.

Habitat in fossis maritimis, in insula Shepey copiose. 4. VII. VIII.

Classis

*Classis V.*PENTANDRIA
MONOGYNIA.

* *Flores monopetali, inferi, tetraspermi. Asperifoliæ.*

MYOSOTIS. 180.

*Cor. hypocrateriformis, quinquefida, emarginata ;
fauce clausa fornicibus.*

- scorpioides.*
arvensis. 1. MYOSOTIS seminibus nudis, foliorum apicibus cal-
lulis. *Sp. pl.* 188. *Vir. cliff.* 149. *Fl. dan.* 583.
Myosotis foliis hirsutis. *Hort. cliff.* 46.
Myosotis scorpioides hirsuta. *Ger. em.* 337. *Park.*
691. *R. syn.* 229.
Scorpiurus annuus, radice exigua. *Hall. hist.* 590.
Echium scorpioides arvense. *B. pin.* 254.
B Myosotis scorpioides hirta minor. *Mer. pin.* *R. syn.*
229.
Echium scorpioides minus flosculis luteis. *B. pin.* 254.
γ Myosotis scorpioides latifolia hirsuta. *Mer. pin.* *R. syn.*
229. *t. 9. f. 2.*
palustris. δ Myosotis foliis glabris. *Hort. cliff.* 46.
Myosotis scorpioides palustre. *Ger.* 337. *R. syn.* 229.
Scorpiurus radice longa fibrata perenni. *Hall. hist.*
591.
Echium scorpioides palustre. *B. pin.* 254.
Anglis, mouse-ear Scorpion-grass.
Habitat in pratis aridis, δ in fossis, et ad scaturigines et
fontes. 4. IV—VIII.

LITHOSPERMUM. 181.

Cor. infundibuliformis fauce perforata, nuda. Cal.
quinquepartitus.

1. LITHOS-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

PENTANDRIA MONOGYNIA.

ANCHUSA. 182.

Cor. infundibuliformis fauce clausa fornicibus. *Sem.*
basi insculpta.

*semper-
virens.*

1. ANCHUSA pedunculis diphyllis capitatis. *Sp. pl.*
192.

Buglossum latifolium sempervirens. *B. pin.* 256. *R.*
syn. 227.

Borrago sempervirens. *Ger. em.* 797.

Anglis, evergreen Alkanet.

Habitat in ruderatis. Near Hornsplace, near Roches-
ter. *R. syn.* Juxta Vauxhall prope Londinum, circa
Norwich. *D. Rose.* 4. V. VI.

CYNOGLOSSUM. 183.

Cor. infundibuliformis, fauce clausa fornicibus. *Sem.*
depressa, inferiore tantum latere stylo affixa.

officinale.

1. CYNOGLOSSUM staminibus corolla brevioribus,
foliis lato-lanceolatis tomentosis sessilibus. *Sp. pl.*
192.

Cynoglossum foliis ellipticis, lanceolatis sericetis, caule
foliofo. *Hall. hist.* 587.

Cynoglossum majus vulgare. *Ger. em.* 804. *Park.* 511.
B. pin. 257. *R. syn.* 226.

β Cynoglossum foliis planis, sessilibus elliptico-lanceola-
tis, asperis. *Hall. hist.* 588.

Cynoglossum folio virente. *B. hist.* III. 600. *R. syn.*
226.

Anglis, common Hound's-tongue.

Habitat in ruderatis et ad vias β in Essexia frequens. In
a heage facing the road on Stamford-hill near Tot-
tenham. *R. syn.* 4. VI.

Pharmac. Cynoglossi Radix.

PULMONARIA. 184.

Cor. infundibuliformis, fauce pervia. *Cal.* prismatico-
pentagonus.

1. PULMO-

1. PULMONARIA foliis radicalibus ovato-cordatis *officinalis*.
scabris. *Sp. pl.* 164. *Hall. hist.* 597. *Fl. dan.* 482.

Pulmonaria foliis Echii. *Ger. em.* 808. *R. syn.* 226.

Pulmonaria latifolia. *Park. par.* 248.

Symphytum maculosum f. *Pulmonaria latifolia.* *B. pin.*
259.

Anglis, common Buglos-cowslip.

Habitat in nemorosis. In a Wood by Holbury-house, in
the new forest, Hampshire. *R. syn.* 4. V.

2. PULMONARIA calycibus abbreviatis, foliis ovatis, *maritima*.
caule ramoso procumbente. *Sp. pl.* 195. *Fl. dan.*
25.

Cerithe maritima procumbens. *Dill. elth.* 75. t. 65.

Echium marinum. *Phys. br.* *R. syn.* 228.

Cynoglossum procumbens glaucophyllum maritimum
nostras, floribus purpuro-cæruleis, seminibus lævi-
bus. *Pluk. ph. t.* 172. f. 3.

Buglossum dulce ex Insulis Lancastræ. *Park.* 765.

Anglis, sea Buglos-cowslip.

Habitat in littoribus maritimis arenosis. Near White-
haven and against Bigger in the Isle of Walney, in
Lancashire, Mr. Lawson; near Threfarthen An-
glesea, and by the River Euyfni, between Dinardinell
and Clynog in Carnarvonshire, Mr. Llhwyd. *R.*
Syn. In Scotia frequens. D. Hope. 4. V. VI.

S Y M P H Y T U M. 185.

Corollæ limbus tubulato-ventricosus: fauce clausa ra-
diis sublatus.

1. SYMPHYTUM foliis ovato-lanceolatis decurrenti-*officinale*.
bus. *Sp. pl.* 125. *Mat. med.* 61. *Hall. hist.* 600.

Symphytum majus vulgare. *Park.* 523.

Symphytum, consolida major. *B. pin.* 259. *Ger.* 806.

Symphytum flore purpureo. *Merr. pin.*

Anglis, common Comfrey.

Habitat ad ripas fluviorum et in aquosis. 4. VII.

Pharmac, Consolidæ majoris, Radix.

BORAGO. 188.

Cor. rotata : fauce radiis clausa.

- officinalis.* * 1. BORAGO foliis omnibus alternis, calycibus patentibus. *Sp. pl.* 197.
 Borago hortensis. *Ger. em.* 797. *R. syn.* 228.
 Buglossum latifolium, Borago. *B. pin.* 256.
Anglis, common Borage.
 Habitat in ruderatis et muris frequens. 4. VI—VIII.
Pharmac. Boraginis Folia, Flores.

ASPERUGO. 189.

Cal. fructus compressus, lamellis plano-parallelis, sinuatis.

- procumbens.* 1. ASPERUGO calycibus fructus compressis. *Sp.* 198.
Hall. hist. 606. *Fl. dan.* 552.
 Buglossum sylvestre, caulibus procumbentibus. *B. pin.* 257. *Hist. Ox.* III. 439. t. 26. f. 13.
 Borago minor sylvestris. *Park.* 765.
 Aparine major Plinii. *Ger. em.* 1122.
Anglis, great Goose-grass, or German Madwort.
 Habitat in ruderatis pinguibus. By Boxley in Suffex, and in Holy Island. *R. syn.* 228. prope Purfleet in Essex. D. Alchorne. ☉. IV. V.

LYCOPSIS. 190.

Corollæ tubo incurvato.

- arvensis.* 1. LYCOPSIS foliis lanceolatis hispidis, calycibus florescentibus erectis. *Sp. pl.* 199. *Fl. dan.* 435.
 Lycopsis foliis asperrimis, undulatis, serratis, linguiformibus. *Hall. hist.* 605.
 Buglossum sylvestre minus. *B. pin.* 256. *Park.* 765.
 Buglossa sylvestris minor. *Ger. em.* 799. *R. syn.* 227.
Anglis, small Bugloss.
 Habitat in arvis et ad vias frequens. ☉. VI—IX.

ECHIUM. 191.

Corolla irregularis, fauce nuda.

1. ECHIUM

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

PENTANDRIA MONOGYNIA.

Primula foliis dentatis rugosis. scapo nullo. Sp. pl.
205.

Primula foliis hirsutis, rugosis, dentatis, scapis unifloris. Hall. hist. 608.

Primula veris minor. Ger. em. 781.

Primula veris vulgaris. Park. 539. R. syn. 284.

Verbasculum sylvarum majus singulari flore. B. pin.
241.

β *Primula foliis rugosis dentatis, scapis multifloris, floribus exterioribus nutantibus. Hall. hist. 609. Fl. dan. 433.*

Primula pratensis inodora lutea. Ger. em. 780. R. syn.
284.

Verbasculum pratense aut sylvaticum inodorum. B.
pin. 241.

Anglis, common Primrose β Oxslips.

Habitat in sylvis, sepibus et ericetis. 4. IV. V.

veris.

2. PRIMULA foliis dentatis rugosis, scapo multifloro, limbo corollarum concavo. *Sp. pl. 204. Fl. dan.*
434.

Primula foliis rugosis, dentatis hirsutis, scapis multifloris, floribus omnibus nutantibus. Hall. hist. 610.

Primula veris major. Ger. em. 781. R. syn. 284.

Verbasculum pratense odoratum. B. pin. 241.

Paralysis vulgaris pratensis, flore flavo simplici odorato. Park. th. 535. Par. 244.

Anglis, common Cowslips or Pagils.

Habitat in pratis et pascuis. 4. IV. V.

Pharmac. Paralysis Flores.

farinosa.

3. PRIMULA foliis crenatis glabris, florum limbo plano. *Sp. pl. 205. Fl. succ. 172. Fl. dan. 125.*

Primula veris flore rubro. Ger. em. 783.

Aretia foliis glabris, ellipticis, rugosis, subtus farinosis, floribus umbellatis. Hall. hist. 625.

Verbasculum umbellatum alpinum minus. B. pin.
247. *R. syn. 285.*

Paralysis minor flore rubro. Park. par. 246.

Anglis, Bird's-eye.

Habitat in paludibus montosis in comitatibus Westmordandico Cumberlandico et Eboracensi. 4. V. VI.

MENYANTHES

MENYANTHES. 202.

Corolla hirsuta. *Stigma* bifidum. *Caps.* unilocularis.

1. MENYANTHES foliis ternatis. *Sp. pl.* 208. *Hall. trifoliata. hist.* 633. *Fl. dan.* 551.

Menyanthes palustre triphyllum latifolium et angustifolium. Tourn. 117. *R. syn.* 285.

Trifolium palustre. B. pin. 327. *Cord. hist.* 96.

Trifolium paludosum. Ger. em. 1194. *Park.* 1212.

Anglis, marsh Trefoil or Buckbean.

Habitat in palustribus et aquosis frequens. 4. VI. VII.

Pharmac. Trifolii paludosi *Herba.*

2. MENYANTHES foliis cordatis integerrimis, corollis ciliatis. *Sp. pl.* 207. *Fl. dan.* 339. *Nymphoides.*

Nymphaea lutea minor flore fimbriato. B. pin. 194. *hist. II.* 772. *R. syn.* 368.

Anglis, fringed Buckbean.

Habitat in fluviis et fossis majoribus, circa Kingston et Hampton. 4. VII.

HOTTONIA. 203.

Corolla hypocrateriformis. *Stamina* tubo corollæ imposita. *Caps.* unilocularis.

1. HOTTONIA pedunculis verticillato-multifloris. *Sp. palustris. pl.* 208.

Hottonia florum verticillis spicatis. Hall. hist. 632.

Hottonia. Boerb. ind. alt. 207. *R. syn.* 285. *Fl. dan.* 484.

Millefolium aquaticum s. Viola aquatica caule nudo. B. pin. 141.

Millefolium aquaticum floridum s. Viola aquatica. Park. 1256.

Viola palustris. Ger. em. 826.

Anglis, Water-violet.

Habitat in fossis et paludibus frequens. 4. VII. VIII.

LYSIMACHIA. 205.

Cor. rotata. *Caps.* globosa, mucronata, decemvalvis.

* *Pedunculis multifloris.*

- vulgaris.* 1. **LYSIMACHIA** paniculata, racemis terminalibus.
Sp. pl. 209. *Scop. Carn.* 138.
 Lysimachia foliis ovato-lanceolatis, spicis paniculatis.
Hall. hist. 630.
 Lysimachia lutea. *Ger. em.* 474. *Hort. æst. Ord. X.*
t. 6. f. 3. *R. syn.* 282. major. *B. pin.* 246. vul-
 garis. *Park.* 545.
 β Lysimachia lutea foliis brevioribus et obtusis, floribus
 in summitate congestis. *R. syn.* 282.
Anglis, yellow Loosestrife or Willow-herb.
 Habitat in paludibus nemorosis et ad ripas fluviorum. 4.
 VI. VII.
Pharm. Lysimachiae Herba.

- thyrsiflo-
ra.* 2. **LYSIMACHIA** racemis lateralibus pedunculatis.
Sp. pl. 209. *Fl. dan.* 517.
 Lysimachia ramis densissimis, alaribus, folio breviori-
 bus. *Hall. hist.* 631.
 Lysimachia lutea flore globoso. *Ger. em.* 475. *Park.*
 544. *R. syn.* 283.
 Lysimachia bifolia, flore globoso luteo. *B. pin.* 245.
Anglis, tufted Loosestrife.
 Habitat in paludibus et ad ripas fluviorum sed rarius.
 Near York. *Dr. White.* Anglelea. *Mr. Llhwyd.* By
 Kings-Langley in Hertfordshire. *Ph. br. R. syn.*
 4. VI.

** *Pedunculis unifloris.*

- nemorum.* 3. **LYSIMACHIA** foliis ovatis, acutis, floribus solita-
 riis, caule procumbente. *Sp. pl.* 211. *Fl. dan.* 174.
 Lysimachia caule decumbente, foliis ovato-lanceolatis
 petiolis alaribus unifloris. *Hall. hist.* 628.
 Anagallis lutea. *Ger. em.* 618. *R. syn.* 282. nemo-
 rum. *B. pin.* 252.
 Anagallis flore luteo. *Park.* 558.
Anglis, wood Loosestrife or Pimpernel.
 Habitat in sylvis et umbrosis humidis. 4. V—VII.

4. **LYSIMACHIA**

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

A Z A L E A. 211.

Cor. campanulata. *Stamina* receptaculo inferta. *Caps.* quinquelocularis.

procumbens. 1. AZALEA ramis diffuso-procumbentibus. *Fl. lapp.* 90. t. 6. f. 2. *Sp. pl.* 215. *Fl. carn.* 140. *Dan.* 9. *Penn. voy.* 43. t. 5.

Azalea caule procumbente foliis ovatis, margine introverso. *Hall. hist.* 666.

Chamæcistus serpyllifolia flore carneo. *Park.* 657. *B. pin.* 466.

Chamæcistus serpyllifolius. *Ger. em.* 1284.

Anglis, procumbent Azalea.

Habitat in montibus. Glen Ciren, D. Hope, Ben. Croachan Argyleshire; Ben-more, Bredalbane; near Arnisdale, *Pennant's Voy.* 5. VI. VII.

C O N V O L V U L U S. 215.

Cor. Campanulata, plicata. *Stigmata* duo. *Caps.* bilocularis: loculis dispermis.

arvensis. 1. CONVOLVULUS foliis sagittatis utrinque acutis, pedunculis unifloris. *Sp. pl.* 218. *Fl. dan.* 459.

Convolvulus foliis sagittatis, latefcentibus, petiolis unifloris stipulis remotis, subulatis. *Hall. hist.* 664.

Convolvulus minor vulgaris. *Park.* 171. *R. syn.* 275.

Convolvulus minor arvensis. *B. pin.* 294.

Smilax lævis minor. *Ger. em.* 861. *Dod. purg.* 213.

β Convolvulus arvensis minimus. *D. Dent.* *R. syn.* 276.

γ Convolvulus flore minimo ad unguem fere secto. *R. syn.* 276.

Anglis, small Bindweed.

Habitat in arvis et ad vias ubique. 4. VI. VII.

sepium. 2. CONVOLVULUS foliis sagittatis postice truncatis, pedunculis tetragonis unifloris. *Sp. pl.* 218. *Fl. dan.* 458.

Convolvulus foliis sagittatis, hamis emarginatis angulosis, petiolis unifloris, stipulis cordatis maximis. *Hall. hist.* 663.

Convolvulus

Convolvulus major albus. *Park.* 163. *B. pin.* 294.
R. syn. 275.

Smilax lævis major. *Ger. em.* 861. *Dod. purg.* 210.
Anglis, great Bindweed.

Habitat in sepibus subhumidis. 4. VII. VIII.

3. CONVOLVULUS foliis reniformibus, pedunculis *Soldanella*
unifloris. *Sp. pl.* 159. *Fl. Carn.* 142.

Convolvulus maritimus Soldanella dicta. *R. syn.* 276.

Soldanella marina. *Ger. em.* 838.

Soldanella vulgaris. *Park.* 168.

Soldanella maritima minor. *B. pin.* 295.

Anglis, sea Bindweed, or Scottish Scurvy-grass.

Habitat in littoribus maritimis. 4. VII. VIII.

POLEMONIUM. 216.

Cor. quinquepartita. *Stam.* squamis basin corollæ
claudentibus inserta. *Stigma* trifidum. *Caps.* trilobularis.

1. POLEMONIUM foliis pinnatis, floribus erectis, ca- *cærule-*
lycibus corollæ tubo longioribus. *Sp. pl.* 230. *Fl. um.*
Dan. 255.

Polemonium foliis pinnatis, floribus erectis, elliptico-
lanceolatis. *Hall. hist.* 665.

Polemonium vulgare cæruleum. *Sch. It.* 460. *Tourn.*
146. *R. syn.* 288.

Valeriana cærulea. *B. pin.* 164.

Valeriana græca. *Ger. em.* 1076. *Park.* 122.

Anglis, blue Polemonium, or greek-Valerian.

Habitat in dumetis infra Malham-cove. 'Dr. Richard-
son. *R. syn.* 4. VI.

VERBASCUM. 244.

Cor. rotata, subinæqualis. *Caps.* bilocularis, bivalvis.

1. VERBASCUM foliis decurrentibus utrinque tomen- *Thapsus.*
tosis, caule simplici. *Sp. pl.* 252. *Hall. hist.* 681.
Fl. dan. 631.

Verbasculum. *Cam. ep.* 878. *mas latifolium luteum.*
B. pin. 239. *R. syn.* 287.

Verbascum

Verbascum album vulgare, sive Thapsus barbatus communis. *Park.* 60.

Thapsus barbatus. *Ger. em.* 773.

Anglis, great Mullein, or High-taper.

Habitat in locis cretaceis et glareosis. §. VII. VIII.

Pharmac. Verbasci. *Folia.*

Thapsoi-
des.

2. VERBASCUM foliis decurrentibus, caule ramosa.

Sp. pl. 1670.

Verbascum angustifolium ramosum, flore aureo, folio crassiore. *B. hist.* III. 856.

Verbascum album mas, flore luteo. *Dalech. hist.* 1301.

Verbascum angustius. *Dod. pempt.* 143.

Verbascum foliis viridibus crassioribus. *Hist. Ox.* II. 436.

Anglis, bastard Mullein.

Habitat in pratis et pascuis cretaceis et arenosis, in comitatu Cantiano passim. §. VII. VIII.

Lychnitis. 3. VERBASCUM foliis cuneiformi-oblongis. *Sp. pl.*

253. *Fl. dan.* 586.

Verbascum spica ramosa, foliis ovato-lanceolatis inferioribus petiolatis. *Hall. hist.* 583.

Verbascum Lychnitis Matthioli. *Ger. em.* 775.

Verbascum Lychnitis flore albo parvo. *B. pin.* 240.

Verbascum mas foliis longioribus. *Park.* 60.

Verbascum flore albo parvo. *B. hist.* III. 813. *R. syn.* 287.

β Verbascum pulverulentum flore luteo parvo. *B. hist.*

III. 872. *R. syn.* 287.

Anglis, white Mullein.

Habitat in pratis et pascuis cretaceis et arenosis in agro Cantiano frequens; β circa mœnia Nordovici urbis; et ad Buriam oppidum. *R. syn.* §. VII.

nigrum. 4. VERBASCUM foliis oblongo-cordatis petiolatis. *Sp.*

pl. 253.

Verbascum foliis imis petiolatis, cordato lanceolatis, superioribus sessilibus, ovatis, acutis. *Hall. hist.* 584.

Verbascum nigrum. *Ger. em.* 775. vulgare. *Park.* 61.

Verbascum nigrum flore ex luteo purpurascente. *B. pin.* 240.

Verbascum

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

PENTANDRIA MONOGYNIA,

Habitat in sylvis et sepibus. 4. V. VI.

HYOSCYAMUS. 246.

Cor. infundibuliformis, obtusa. Stamina inclinata.
Caps. operculata, bilocularis.

niger.

1. HYOSCYAMUS foliis amplexicaulibus sinuatis, floribus sessilibus. *Sp. pl.* 257. *Hall. hist.* 580.
Hyoscyamus vulgaris. *B. hist.* III. 627. *R. syn.* 274.
Hyoscyamus niger. *Ger. em.* 353. s. vulgaris. *B. pin.* 169. *Park.* 362.
Anglis, common Henbane.
Habitat ad pagos, vias, ruderata. ☉. VI.
Pharmac. Hyoscyami Radix, Herba, Semina.

DATURA. 245.

Cor. infundibuliformis plicata. Cal. tubulosus, angulatus, deciduus. Caps. quadrivalvis.

Stramonium.

- *1. DATURA pericarpis spinosis erectis ovatis, foliis ovatis glabris. *Sp. pl.* 255. *Fl. dan.* 460.
Stramonium foliis angulosis, fructu erecto, muricato, calyce pentagono. *Hall. hist.* 586.
Stramonium spinosum. *Ger. em.* 349.
Solanum foetidum, pomo spinoso oblongo, flore albo. *B. pin.* 164.
Solanum pomo spinoso oblongo, flore calathoide, Stramonium vulgo dictum. *R. syn.* 266.
Anglis, common Datura, or Thorny Apple.
Habitat in sinitis, et ruderatis passim. ☉. VII.
Pharmac. Stramonii Herba.

SOLANUM. 250.

Cor. rotata. Antheræ subcœalitæ, apice poro gemina debiscentes. Bacca bilocularis.

nigrum.

1. SOLANUM caule inermi herbaceo, foliis ovatis dentato-angulatis, umbellis nutantibus. *Sp. pl.* 266. *Hall. hist.* 576. *Fl. dan.* 460.
Solanum hortense. *Ger. em.* 339. s. vulgare baccis nigris. *B. hist.* III. 608.
Solanum vulgare. *Park.* 346. *R. syn.* 264.

Solanum

Solanum officinarum. *B. pin.* 166.

β *Solanum hortense* f. *vulgare* baccis luteis. *B. hist.* III. 608.

Anglis, common or garden Nightshade.

Habitat in *runderatis et fimetis.* ☉. VI. VII.

Pharmac. Solani Herba, *Fruetus.*

2. SOLANUM caule inermi frutescente flexuoso, foliis *dulcamara*-superioribus hastatis, racemis cymosis. *Sp. pl.* 264. *ra.*
Fl. dan. 607.

Solanum caule flexuoso, frutescente, foliis superioribus tripartitis et cordato-lanceolatis. *Hall. hist.* 576.

Solanum lignosum f. *Dulcamara.* *Park.* 350. *R. syn.* 65. *Blakw.* 34.

Solanum scandens f. *Dulcamara.* *B. pin.* 167.

Amara dulcis. *Ger. em.* 350.

Solanum lignosum f. *Dulcamara marina.* *R. syn.* 265.

Anglis, woody Nightshade, or Bitter-sweet.

Habitat in *sepibus humidis*, β in *littoribus maritimis passim.* 4. VI. VII.

Pharmac. *Dulcamaræ Radix, Folia, Cortex.*

Obs. *variat flore albo.* β *foliis hirsutis.*

A T R O P A. 248.

Cor. campanulata. *Stamina* distantia. *Bacca* globosa, bilocularis.

1. ATROPA caule herbaceo, foliis ovatis integris. *Belladonna.*
Sp. pl. 260. *na.*

Belladonna. *Clus. hist.* II. 86. *R. syn.* 265. caule herbaceo, brachiato, foliis ovato-lanceolatis integerrimis. *Hall. hist.* 579.

Solanum lethale. *Ger. em.* 340. *Park.* 346.

Solanum melanocerasus. *B. pin.* 166.

Anglis, common Dwale or deadly Nightshade.

Habitat in *sepibus et ruderatis.* In Chuckstone, near Rochester, about Fulborn, Cambridgeshire. Sutton Cofield, Warwickshire. *R. syn.* Among the box going down to Dorking, Surry, Dr. Wilmer., By the Park wall between Temsford-mills and Wellwyn,

PENTANDRIA MONOGYNIA.

wyn, Herts. Dr. Watson. *Bl. spec. bot.* 93. 4.
VI.

*** *Flores monopetali superi.*

S A M O L U S. 221.

Cor. hypocrateriformis. *Stamina* munita squamulis
corollæ. *Caps.* unilocularis.

- valerandi.* 1. SAMOLUS. *Sp. pl.* 243. *Fl. dan.* 148. *Hall.* 707.
• Samolus valerandi. *B. hist.* III. 791. *R. syn.* 283.
Anagallis aquatica rotundo folio non crenato. *B. pin.*
252.
• Anagallis aquatica rotundifolia. *Ger. em.* 620.
Anagallis aquatica 3. Lobeli. folio subrotundo non
crenato. *Park.* 1237.
Anglis, water Samolus or Pimpernel.
Habitat in palustribus, in pratis humidis; inter Green-
wich et Woolwich. 4. VI.

L O N I C E R A. 232.

Cor. monopetala, irregularis. *Bacca* polysperma, bi-
locularis.

- periclyme-* 1. LONICERA capitulis ovatis imbricatis terminalibus,
num. foliis omnibus distinctis. *Sp. pl.* 247. *Hall. hist.* 673.
Caprifolium Germanicum. *Dod. pempt.* *R. syn.* 458.
Periclymenum. *Ger. em.* 891. *Fuchs. hist.* 646. *f.* Ca-
prifolium vulgare. *Park.* 1460.
Periclymenum non perfoliatum germanicum. *B. pin.*
302.
β Periclymenum foliis quernis. *Merr. pin.*
Anglis, common Honeysuckle or Woodbine.
Habitat in sepibus. 2. V. VII.
Pharmac. Caprifolii. *Folia, Flores.*

C A M P A N U L A. 217.

Cor. campanulata fundo clauso valvis staminiferis.
Stigma trifidum. *Caps.* infera, poris lateralibus
dehiscens.

• *Foliis lævioribus angusticrilus.*

I. CAM-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

PENTANDRIA MONOGYNIA.

Litchfield et Meriden copiose ; prope Worcester. 3.
VII.

** *Foliis scabris latioribus.*

latifolia. 4. CAMPANULA foliis ovato-lanceolatis, caule simplicissimo tereti, floribus solitariis pedunculatis, fructibus cernuis. *Sp. pl.* 234. *Vir. cliff.* 17. *Fl. dan.* 85.

Campanula foliis ovato-lanceolatis serratis, petiolis unifloris, calycibus glabris. *Hall. hist.* 691.

Campanula maxima foliis latissimis. *B. pin.* 94. *R. syn.* 276.

Trachelium majus Belgarum. *Lob. ic.* 278. *Park.* 643. sive gigantium. *Ger. em.* 448.

Anglis, giant Bell-flower, or Throatwort.

Habitat in dumetis et sepibus. In the mountainous part of Derbyshire, Staffordshire, Yorkshire, and elsewhere in the north of England plentifully. *R. syn.* prope Holywell, Flintshire. 4. VII.

trachelium. 5. CAMPANULA caule angulato, foliis petiolatis, calycibus ciliatis, pedunculis trifidis. *Sp. pl.* 235.

Campanula foliis cordato-lanceolatis, acute serratis, hirsutis, petiolis ex alis paucifloris, calycibus hispida. *Hall. hist.* 690.

Campanula vulgatiore foliis Urticæ, vel major et asperior. *B. pin.* 94. *R. syn.* 276.

Trachelium majus. *Ger. em.* 448. flore purpureo. *Park.* 354.

Anglis, great Bell-flower, or Canterbury-Bells.

Habitat in sylvis et sepibus. 4. VII. VIII.

glomerata 6. CAMPANULA caule angulato simplici, floribus sessilibus, capitulo terminali. *Sp. pl.* 235.

Campanula caule simplici, aspero, foliis amplexicaulibus, floribus capitatis. *Hall. hist.* 685.

Campanula pratensis flore conglomerato. *B. pin.* 94. *R. syn.* 277.

Trachelium minus. *Ger. em.* 449. *Park.* 644.

Anglis, lesser Bell-flower, or Canterbury-Bells.

Habitat

Habitat in pascuis montosis, cretaceis frequens. 4. VII.

*** Caule subdiviso.

7. CAMPANULA caule basi subramoso stricto, foliis *hybrida*. oblongis crenatis, calycibus aggregatis corolla longioribus, capsulis prismaticis. *Sp. pl.* 239.

Campanula caule erecto, fructibus prismaticis, floribus rotatis, sessilibus, congestis. *Hall. hist.* 704.

Campanula arvensis erecta, vel speculum Veneris minus. *Park.* 1331. *Ger. em.* 439. *R. syn.* 278.

Campanula arvensis minima erecta. *Hist. Ox. II.* 457. t. 2. f. 22.

Anglis, corn Bell-flower.

Habitat in arvis cretaceis passim. ☉. VI. VII.

8. CAMPANULA foliis cordatis quinquelobis petiolaribus *hederacea*. glabris, caule laxo. *Sp. pl.* 240. *Fl. dan.* 330.

Campanula Cymbalariae foliis. *Ger. em.* 452. *Park.* 652. *R. syn.* 277. vel folio hederaceo. *B. pin.* 93.

Anglis, ivy-leaved Bell-flower.

Habitat in locis umbrosis humidis, in Cornubia et Devonia frequens. In Bagley Wood, near Oxford. *R. syn.* In Wallia frequens, D. Stillingfleet. prope Halifax, D. Bolton. prope Bayly, Marshfield et alibi, in Suffexia. 4. VI.—VIII.

PHYTEUMA. 219.

Cor. rotata laciniis linearibus, quinquepartita. *Stigma* trifidum. *Caps.* trilocularis.

1. PHYTEUMA capitulo subrotundo, foliis serratis : orbicularibus radicalibus cordatis. *Sp. pl.* 242. *ris.*

Rapuntium foliis imis cordatis, caulinis sessilibus acutis, bracteis ovato-lanceolatis. *Hall. hist.* 681.

Rapunculus corniculatus montanus. *Ger. em.* 455. *R. syn.* 278.

Rapunculus alopecuroides orbiculatus. *Park.* 648.

Rapunculus folio oblongo, spica orbiculari. *B. pin.* 92. *Hist. Ox. II.* 463. t. 5. f. 47.

Anglis, horn'd Rampions.

Habitat in pascuis cretaceis. On the downs of Suffex and

PENTANDRIA MONOGYNIA.

and Hampshire. *R. syn. prope Leatherhead in comitatu Surriensi. 4. VI. VII.*
Obs. variat spica oblonga.

* *Flores pentapetali inferi.*

RHAMNUS. 264.

Cal. tubulosus: squamis stamina munientibus. Cor. nulla. Bacca.

- catharticus.* 1. RHAMNUS spinis terminalibus, floribus quadrifidis, dioicis, foliis ovatis, caule erecto. *Sp. pl. 279.*
 Rhamnus spinosis, foliis ovato-lanceolatis ferratis. *Hall. hist. 824.*
 Rhamnus catharticus. *B. pin. 478. R. syn. 466.*
 Rhamnus solutivus. *Ger. em. 1337. f. spina infectoria vulgaris. Park. 243.*
Anglis, purging Rhamnus, or Buckthorn.
Habitat in sylvis et sepibus. 4. IV. V.
Pharmac. Spinæ cervinæ Bacca.

- frangula.* 2. RHAMNUS inermis, floribus monogynis hermaphroditis, foliis integerrimis. *Sp. pl. 280, Fl. dan. 278.*
 Rhamnus inermis, foliis ovato-lanceolatis, integris, floribus quinquefidis, androgynis. *Hall. hist. 821.*
 Frangula f. Alnus nigra baccifera. *Park. 240. R. syn. 465.*
 Alnus nigra f. Frangula. *Ger. em. 1469.*
 Alnus nigra baccifera. *B. pin. 428. Hist. I. 560.*
Anglis, common Rhamnus, or berry-bearing Alder.
Habitat in sylvis et sepibus. 4. IV. V.
Pharmac. Frangulæ Cortex,

EUONYMUS. 270.

Cor. pentapetala. Caps. pentagona, quinquelocularis, quinquevalvis, colorata. Sem. calyptrata.

- europæus.* 1. EUONYMUS floribus plerisque quadrifidis. *Sp. pl. 286. Fl. suec. 204.*
 Euonymus foliis ovato-lanceolatis ferratis, ramis tetragonis. *Hall. hist. 829.*
 Euonymus vulgaris. *Park. 241. R. syn. 468.*
 Euonymus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Ribes nigrum vulgo dictum, folio olente. *B. hist. II.*
98. *R. syn.* 456.

Ribes fructu nigro. *Park.* 1562. *Ger. em.* 1593.

Grossularia non spinosa fructu nigro. *B. pin.* 455.

Anglis, black Currants, or Squinancy berries.

Habitat in sylvis humidis et ad ripas fluviorum. At
Abington, Cambridgeshire; Blunham, Bedford-
shire; in Warwickshire, Westmorland and Cum-
berland; and by the Hoppet Bridge near Braintree,
Essex. *R. syn.* h. V.

H E D E R A. 280.

Petala quinque, oblonga. *Bacca* quinquesperma, ca-
lyce cincta.

helix.

1. HEDERA foliis ovatis lobatisque. *Sp. pl.* 292.

Hedera foliis sterilis trilobatis, fructiferis ovato-lan-
ceolatis. *Hall. hist.* 826.

Hedera helix. *Math.* 627. *Ger. em.* 858.

Hederá communis major et minor. *B. hist. II.* 111.
R. syn. 459.

Hedera arborea, et major sterilis. *B. pin.* 305.

Hedera arborea sive scandens et corymbosa communis.
Park. 678.

Anglis, common Ivy.

Habitat in sylvis et sepibus. h. X.

Pharmac. Hederæ Folia, Bacca, Gummi.

* *Flores incompleti superi.*

I L L E C E B R U M. 287.

Cal. pentaphyllus, cartilagineus. *Cor.* ○. *Stigma*
simplex. *Caps.* quinquevalvis, monosperma.

verticil- 1. ILLECEBRUM floribus verticillatis nudis, caulibus
latum. procumbentibus. *Sp. pl.* 298. *Fl. dan.* 335.

Corrigiola. *Cat. giff.* 169. *R. syn.* 160.

Polygonum serpyllifolium verticillatum. *R. f. ed. I.*
160.

Polygala repens. *Park.* 1333. *nivea.* *B. pin.* 215.
Paronychia

Paronychia serpyllifolia palustris. Vaill. 157. t. 15.
f. 1.

Anglis, marsh Illecebrum.

Habitat in palustribus, udis et humidioribus pascuis circa Pensans oppidum et alibi versus extremum Cornubiæ angulum. R. syn. 4. VII.

GLAUX. 288.

Cal. monophyllus. Cor. o. Caps. unilocularis, quinquevalvis, pentasperma.

1. GLAUX. *Sp. pl. 301. Fl. suec. 210. dan. 269.*

Glaux maritima. B. pin. 215. R. syn. 285. minor. maritima. Park. 1283.

Glaux exigua maritima. Ger. em. 562.

Anglis, sea Glaux.

Habitat in palustribus maritimis frequens. 4. VI. VII.

** *Flores incompleti inferi.*

THESIUM. 289.

Cal. monophyllus, cui stamina inserta. Sem. unicum.

1. THESIUM *panicula foliacea, foliis lineari-lanceolatis. Sp. pl. 301. Fl. suec. 207. dan. 328. Linophyllum.*

Thesium caule erecto, paniculato, foliis lanceolatis. Hall. hist. 1573.

Linaria adulterina. Ger. em. 555. R. syn. 202.

Linaria montana flosculis albicantibus. B. pin. 213.

Pseudo-linaria montana alba. Park. 459.

Anglis, common Thesium or bastard Toad-flax.

Habitat in pascuis montosis præsertim solo cretaceo. Gogmagog Hills, Newmarket Heath. R. syn. prope Basingstoke in Hampshire, et prope Croydon et alibi in agro Surriensi. 4. VI. VII.

DIGYNIA.

* *Flore monopetali inferi.*

SWERTIA. 318.

Cor. rotata. Nectariferi pori ad basin laciniarum corollæ. Caps. unilocularis, bivalvis.

- perennis.* 1. SWERTIA corollis quinquefidis, foliis radicalibus ovalibus. *Sp. pl.* 329. *Fac. austr.* III. 25. t. 243.
 Gentiana foliis radicalibus petiolatis, ovatis, floribus paniculatis, rotatis, spicatis. *Hall. hist.* 636.
 Gentiana palustris latifolia. *B. pin.* 188.
 Gentiana XII. punctato flore. *Clus. hist.* 1. 316.
 Gentiana Pennei minor. *Ger. em.* 433.
Anglis, marsh Swertia.
 Habitat in Wallia. D. Richardson invenit. 4. VII. VIII.

GENTIANA. 319.

Cor. monopetala. *Caps.* bivalvis, unilocularis. *Receptaculis* duobus longitudinalibus.

* *Corollis quinquefidis subcampaniformibus.*

- Pneumonanthe.* 1. GENTIANA corollis quinquefidis campanulatis oppositis pedunculatis, foliis linearibus. *Sp. pl.* 330.
Fl. suec. 328. *Dan.* 269.
 Gentiana alis floriferis, foliis linearibus. *Hall. hist.* 641.
 Gentiana palustris angustifolia. *B. pin.* 118. *R. syn.* 274.
 Gentiana autumnalis Pneumonanthe dicta. *Park.* 406.
Pneumonanthe. *Cord. hist.* I. 162. *Ger. em.* 438.
Anglis, marsh Gentian or Calathian Violet.
 Habitat in pratis humidioribus, in pratis Strattoniensibus in Norfolk, D. Stillingfleet; prope Tunbridge, Dr. Wilmer. prope Doncaster, D. Tosfield; prope Milthorp in comitatu Westmorlandico et in Lancastria passim. 4. VIII.

** *Corollis quinquefidis infundibuliformibus.*

- Centauri-* 2. GENTIANA corollis quinquefidis infundibuliformibus caule dichotomo. *Sp. pl.* 332. *Fl. dan.* 617.
um. Centaurium minus. *Cam. epit.* 428. *B. pin.* 278. *R. syn.* 286. vulgare. *Park.* 272.
 Centaurium parvum. *Ger. em.* 547.
Anglis, Centory Gentian,

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

C U S C U T A.

Cal. quinquefidus. *Cor.* monopetala quinquefida. *Caps.* bilocularis. *Sem.* duo.

europæa. 1. CUSCUTA floribus sessilibus. *Sp. pl.* 180. *Hall. hist.* 654. *Fl. dan.* 427.

Cuscuta major. *B. pin.* 219. *R. syn.* 281. f. *Castu-*
tha. *Ger. em.* 577. *Park.* 10.

Anglis, europæan Dodder or Hellweed.

Habitat in arvis et ericetis passim. ☉. VI—VIII.

Pharmac. *Cuscutæ* f. *Epithymi Herba.*

* *Flores incompleti.*

C H E N O P O D I U M. -306.

Cal. pentaphyllus, pentagonus. *Cor.* 0. *Sem.* unicum
lenticulare, superum.

* *Foliis angulosis.*

Bonus
Henricus. 1. CHENOPODIUM foliis triangulari-sagittatis inte-
gerrimis, spicis compositis aphyllis. *Sp. pl.* 318.
Fl. dan. 579.

Chenopodium foliis triangularibus, undulatis, inte-
gerrimis, subtus farinosis. *Hall. hist.* 1578.

Blitum perenne Bonus Henricus dictum. *Hist. Ox.*
II. 599. t. 30. f. 1. *R. syn.* 156.

Bonus Henricus. *B. hist.* II. 965. *Ger. em.* 329.

Lapathum unctuosum. *Park.* 1225.

Lapathum sylvestre latifolium f. *unctuosum folio tri-*
angulo. *B. pin.* 115.

Anglis, triangular Goosefoot, or English Mercury.

Habitat in ruderatis, et ad vias et muros passim. ☉.
VII. VIII.

urbicum. 2. CHENOPODIUM foliis triangularibus subdentatis,
racemis confertis strictissimis caule approximatis
longissimis. *Sp. pl.* 318. *Fl. succ.* 215.

Chenopodium latifolium minus ramosum, petiolis lon-
gissimis ex foliorum alis confertim nascentibus.

Buxb. ball. 69. t. 1.

Chenopodium

Chenopodium erectum, foliis triangularibus dentatis, spicis e foliorum alis plurimis longis erectis tenuibus. *R. syn.* 155.

Anglis, upright Goosefoot.

Habitat in fimetis et ruderatis. ☉. VII. VIII.

3. *CHENOPODIUM* foliis cordato-triangularibus obtusiusculis dentatis, racemis erectis compositis subfoliosis caule brevioribus. *Sp. pl.* 318.

Chenopodium erectum, foliis subtriangularibus antice sinuato-dentatis, racemis erectis foliosis lateralibus. *Hort. cliff.* 85. *Roy. lugdb.* 219.

Blitum Pes anserinus dictum. *R. syn.* 154.

Atriplex sylvestris latifolia. *B. pin.* 119.

Pes anserinus. *Dalech. hist.* 542. t. 542. *Fuchs. hist.* 653. *Ger. em.* 328. *B. hist.* II. 975. *Park.* 749.

- β Blito Pes anserinus dicto similis. *R. syn.* 154.

Anglis, red Goosefoot or Sowbane.

Habitat in fimetis, ruderatis, cultis. ☉. VIII.

Obs. an eadem cum priore?

4. *CHENOPODIUM* foliis ovatis acutis dentatis, racemis ramosis nudis. *Fl. suec.* 216. *Sp. pl.* 318.

Blitum Pes anserinus acutiore folio. *R. syn.* 154.

Atriplex sylvestris latifolia acutiore folio. *B. pin.* 119.

Atriplex sylvestris latifolia altera. *Ger. em.* 328.

Atriplex dicta Pes anserinus alter s. ramosior. *B. hist.* 976. *cum fig. bona*.

- β Blitum Morifono atriplex procumbens folio sinuato lucido, crasso dictum. *Hort. Blas.*

Anglis, wall Goosefoot.

Habitat in fimetis et ruderatis. ☉. VII. VIII.

5. *CHENOPODIUM* foliis cordatis angulato-acuminatis, racemis ramosis nudis. *Fl. suec.* 220. *Sp. pl.* 319.

Chenopodium Stramonii folio. *R. syn.* 154. *Vaill. paris.* 36. t. 7. f. 2.

Chenopodium foliis glabris septangulis, floribus paniculatis. *Hall. hist.* 1581.

Anglis, bastard Goosefoot.

Habitat

Habitat in ruderatis cultis, arvis passim; prope Northfleet copiose: ☉. VIII.

album.

6. CHENOPODIUM foliis inferioribus rhomboideis dentato sinuatis, superioribus oblongis, racemis ramosis subnudis.

Chenopodium foliis rhomboideo-triangulis erectis postice integris: summis oblongis, racemis erectis. *Sp. pl.* 319.

Blitum atriplex sylvestre dictum. *R. syn.* 154.

Atriplex folio sinuato candicante. *B. pin.* 119.

Atriplex sylvestris. *Fuchs. hist. t.* 119. vulgatiores sinuata major. *Park.* 748.

Atriplex vulgaris. *Ger. em.* 326.

- β Chenopodium foliis rhomboideis dentato-sinuatis, racemis ramosis nudis. *Sp. pl.* 319.

Chenopodium sylvestre, opuli folio. *Vaill.* 36. t. 7. f. 1.

Blitum folio subrotundo. *Pet. b. br.* 8. 4. *R. syn.* 155.

- γ Chenopodium foliis integris racemosum. *R. syn.* 155.

δ Chenopodium crasso et obtuso oleæ folio. *R. syn.* 156. *Anglis,* common Goosefoot.

Habitat in fimetis antiquis, ruderatis. ☉. VIII.

- serotinum.* 7. CHENOPODIUM foliis deltoideis sinuato-dentatis rugosis glabris uniformibus, racemis terminalibus. *Cent. pl. II.* *Sp. pl.* 319.

Chenopodium foliis glabris trilobatis dentatis. *Hall. hist.* 1582.

Blitum ficus folio. *Pet. b. br. t.* 8. f. 3. *R. syn.* 155.

- β Chenopodium erectum Chrysanthemi segetum folio. *R. syn.* 155.

Anglis, fig-leaved Goosefoot.

Habitat in fimetis et ruderatis. ☉. VIII. IX.

- glaucum.* 8. CHENOPODIUM foliis ovato-oblongis repandis, racemis nudis simplicibus glomeratis. *Sp. pl.* 320.

Chenopodium angustifolium laciniatum minus. *Tourn.* 506. *R. syn.* 155.

Atriplex angustifolia laciniata. *B. hist. II.* 473.

Anglis, oak-leaved Goosefoot.

Habitat in ruderatis. ☉. VIII.

•• Foliis

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

fruticosa. 2. **SALSOLA** erecta fruticosa, foliis filiformibus obtusiusculis. *Sp. pl.* 324.

Chenopodium foliis linearibus teretibus, carnosis, caule fruticoso. *Hort. cliff.* 806.

Blitum fruticosum maritimum; *Vermicularis frutex dictum.* *R. syn.* 156.

Sedum minus fruticosum. *B. pin.* 284.

Vermicularis frutex minor. *Ger. em.* 523.

Vermicularis fruticosa altera. *Park.* 731.

Anglis, shrubby Saltwort.

Habitat in littoribus maritimis *Norfolchiæ, Devonix, Cornubiæ, &c.* h. VII. VIII.

B E T A. 307.

Cal. pentaphyllus. *Cor.* o. *Sem.* reniforme, intra substantiam baseos calycis.

maritima. 1. **BETA** caulibus decumbentibus, floribus geminis. *Sp. pl.* 322.

Beta sylvestris maritima. *B. pin.* 157. *Park.* 750. *R. syn.* 156.

Anglis, sea Beet.

Habitat in littoribus maritimis frequens. h. VIII.

H E R N I A R I A. 305.

Cal. quinquepartitus. *Cor.* o. *Stamina* quinque sterilia. *Caps.* monosperma.

glabra. 1. **HERNIARIA** glabra herbacea. *Sp. pl.* 317. *Fl. dan.* 529.

Herniaria. *Ger. em.* 569. *R. syn.* 160. *glabra.* *B. hist. III.* 378.

Polygonum minus, s. *Millegrana major.* *B. pin.* 281.

Millegrana major, s. *Herniaria vulgaris.* *Park.* 446.

β *Herniaria lenticulata.* *Sp. pl.* 317.

Polygonum maritimum, longius radicum nostras, serpyllifolia circinato crasso nitente. *R. syn.* 161.

Polygonum minus lenticulatum. *B. pin.* 282. *pr.* 131.

Polygonum folio circinato. *Pet. h. br. t.* 10. f. 6.

Anglis, smooth Rupture-wort.

Habitat in glareosis. In promontorio *Cornubiensi*, the Lizard-point dicto inveni. h. VII.

2. HERNIARIA

2. **HERNIARIA** hirsuta. *B. hist. III. 379. R. syn. 161. hirsuta.*
 herbacea. *Sp. pl. 312.*
Anglis, hairy Rupture-wort.
Habitat in pratis glareosis; prope Colney-Hatch, juxta
Barnet. ☉. VII. VIII.

U L M U S. 313.

Cal. quinquefidus. Cor. 0. Bacca exsucco compresso-
membranacea.

1. **ULMUS** foliis duplicato terratis: basi inæqualibus. *campestris.*
Sp. pl. 225. Hunt. Evel. Sylv. 118. Pl. dax. 632. tris.
Ulmus foliis scabris, ovato-lanceolatis, dentibus serratis. *Hall. hist. 1586.*
Ulmus. D. pemp. 837. Cam. epit. 70. vulgatissima folio lato scabro. *Ger. em. 1480. R. syn. 468.*
Ulmus campestris et Theophrasti. B. pin. 426.
Ulmus vulgaris. Park. 1404.
 β *Ulmus minor folio angusto scabro. Ger. em. 1480. R. syn. 469.*
 γ *Ulmus folio latissimo scabro. Ger. em. 1481. R. syn. 469.*
 δ *Ulmus folio glabro. Ger. em. 1481. R. syn. 469.*
Anglis, common β Cornish, γ Wych-hazel, δ smooth Elm.
Habitat in sepibus. ♀. IV.
Pharmac. Ulmi Folia, Cortex, medius.

* *Flores pentapetali superi dispermi. Umbellatæ.*

• *A Involucro universali partialique.*

ERYNGIUM. 321.

Flores capitati. Receptaculo paleaceo.

1. **ERYNGIUM** foliis radicalibus subrotundis plicatis *maritimo-*
spinosis, capitulis pedunculatis. Sp. pl. 337. It. æl. mum. 151.
Eryngium maritimum. B. pin. 386.
Eryngium marinum. Cam. epit. 448. Clus. hist. II. 169. *Ger. em. 1162. Park. 986. R. syn. 222.*
Anglis, sea Eryngo, or Holly.
Habitat in littoribus maritimis. ♀. VII. VIII.
Pharmac.

Pharmac. Eringii Radix.

campestre. 2. ERYNGIUM foliis amplexicaulibus pinnato-lacinatis. *Sp. pl.* 337. *Fl. dan.* 554. *Jac. austr.* II. t. 155.

Eryngium foliis semipinnatis, pinnis pinnatis, pinnulis lobatis, spinosis. *Hall. hist.* 735.

Eryngium vulgare. *B. hist.* III. 85. *pin.* 386. *R. syn.* 222.

Eryngium mediterraneum. *Ger. em.* 1162. *γ. campestre.* *Park.* 986.

Anglis, common Eryngo.

Habitat in pascuis maritimis; near the ferry from Plymouth into Cornwall; by the old Roman Way called Watling-Street, by Brookhall near Daventry, Northamptonshire, Mr. Thornton; observed by Mr. Lawson, on the shore called Friar-goose, near Newcastle upon Tyne. *R. syn.*; below Melling in Yorkshire plentifully. *Blackf. Spec. bot.* 4. VII. VIII.

HYDROCOTYLE. 322.

Umbella simplex: Involucro tetraphyllo. Petala integra. Sem. semiorbiculato-compressa.

vulgaris. 1. HYDROCOTYLE foliis pelatis, umbellis quinquefloris. *Sp. pl.* 338. *Fl. dan.* 90.

Hydrocotyle foliis rotundis, emarginatis, petiolis centralibus, umbellis fastigiatis. *Hall. hist.* 812.

Hydrocotyle vulgaris. *R. syn.* 222. *Tourn.* 328.

Cotyledon palustris. *Ger. em.* 530. *Park.* 1214.

Ranunculus aquaticus, Cotyledonis folio. *B. pin.* 180.

Anglis, marsh Pennywort, or White-rot.

Habitat in palustribus et inundatis. 4. V. VI.

SANICULA. 323.

Umbellæ confertæ subcapitatæ. Fructus scaber. Flores disci abortientes.

europæa. 1. SANICULA foliis radicalibus simplicibus, flosculis omnibus sessilibus. *Sp. pl.* 339. *Fl. dan.* 283.

Sanicula

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

ECHINOPHORA. 326.

Flores laterales masculi, centrali hermaphrodito. *Sem.*
unicum, involucello immersum.

- spinosa.* 1. ECHINOPHORA foliolis subulato-spinosis integerimis. *Sp. pl.* 344.
Echinophora maritima spinosa. *Tourn.* 656. *R. syn.* 220.
Crithmum spinosum. *Ger. em.* 533.
Crithmum maritimum spinosum. *B. pin.* 288. seu
Pastinaca marina. *Park.* 1286.
Anglis, prickly Echinophora, or Sampire.
Habitat in littoribus maritimis. Observed by Mr. Lawson at Roolebeck in Low-Furness, Lancashire; on the sea-beach between Feverham and Sea Salter. *Blackf. Spec. bot.* 4. VII.

TORDYLIUM. 327.

Cor. radiatæ. *Fructus* orbiculatus, planus, margine
crenatus.

- officinale.* *1. TORDYLIUM involucris partialibus longitudine
florum, foliolis ovatis laciniatis. *Sp. pl.* 345.
Tordylium narbonense minus. *Tourn.* 320. *R. syn.* 206.
Seseli creticum minimum. *B. pin.* 161. *Park.* 906.
Ger. em. 1050.
Anglis, small Tordylium.
Habitat in arvis. Plentifully on the banks of the road to Heddington, about half a mile from Oxford. Mr. J. Sherard. ☉. VI.

CAUCALIS. 328.

Cor. radiatæ. *Fructus* subovatus, striatus, setis rigidis hispidus.

- daucoides.* 1. CAUCALIS umbellis trifidis aphyllis, umbellulis trispermis triphyllis. *Syst. nat.* 205. *Fac. austr.* II. 37. t. 157.
Caucalis involucro universali nullo, umbella trifida involucellis triphyllis. *Fl. Gall. prov.* 236.

Caucalis

Caucalis tenuifolia, flosculis subrubentibus. *R. syn.*
219. *hist.* 467.

Caucalis tenuifolia purpurea. *Park.* 920.

Echinophora tertia leptophyllon purpurea. *Col. echp.*
I. 96. t. 97.

Anglis, small *Caucalis*.

Habitat in arvis et sterilibus. Circa Kingston in agro
Cantabrigiensi. *R. syn. prope Reading.* ☉. V. VI.

2. CAUCALIS umbella universali trifida, partialibus *latifolia*.
penta-permis, foliis pinnatis serratis. *Hort. cliff.*
91. *Syst. nat.* 205. *Jac. Hort. vind.* II. 59. t.
128.

Caucalis involucris et involucellis pentaphyllis, foliis
pinnatis. *Ger. gall. prov.* 227.

Caucalis foliis asperis pinnatis, pinnis serratis, involu-
cris ovato-lanceolatis. *Hall. hist.* 738.

Caucalis arvensis echinata latifolia. *B. pin.* 151. *R.*
syn. 219.

Caucalis apii foliis, flore rubro. *Ger. em.* 1021.

Echinophora arvensis latifolia purpurea. *Park.* 920.

Anglis, broad-leaved *Caucalis*.

Habitat in arvis in agro Cantabrigiensi et alibi. *R. syn.*
circa Crooks Easton in comitatu Hamptoniensi. ☉.
VII.

3. CAUCALIS involucro universali monophyllo, um-*arvensis*.
bella conferta, seminibus rotundato-ovatis, foliolis
ovato-lanceolatis pinnatifidis, ramis divaricatis.

Caucalis foliis duplicato-pinnatis, pinnulis longe
confluentibus. *Hall. hist.* 742.

Caucalis segetum minor, anthrisco hispido similis.
R. syn. 220. *hist.* 468.

Caucalis arvensis humilior, ramosior. *Hist. Ox.* III.
338.

Scandix umbellulis confertissimis hemisphæricis, se-*infesta*.
mine exteriori hispido. *Syst. nat.* 732. *Jac. Fl.*
austr. II. 28. T. 46.

Anglis, corn *Caucalis*.

Habitat in arvis, cretaceis frequens. ☉. VII. VIII.

*anthrif-
cus.*

4. CAUCALIS involucris multifidis, umbella conferta, seminibus oblongis, foliolis ovatis pinnatifidis, ramis erectis.

Tordylium umbella conferta, foliolis ovato-lanceolatis pinnatifidis. *Sp. pl.* 346. *Jac. austr.* III. 24. t. 261.

Caucalis foliis duplicato-pinnatis, nervo multoties latioribus. *Hall. hist.* 741.

Caucalis minor flosculis rubentibus. *Ger. em.* 1022, *R. syn.* 219.

Caucalis minor flore rubente, *Park.* 921.

Caucalis semine aspero, flosculis rubentibus. *B. pin.* 153.

Anglis, hedge Caucalis.

Habitat in sepibus. ♂, VIII.

nodosa.

5. CAUCALIS umbellis lateralibus simplicibus sessilibus, Tordylium umbellis simplicibus sessilibus, seminibus exterioribus hispidis. *Sp. pl.* 346.

Caucalis nodosa echinato semine. *B. pin.* 153. *pr.* 80, *Ger. em.* 1022. *Park.* 921. *R. syn.* 220.

Anglis, knotted Caucalis.

Habitat ad margines agrorum frequens, ☉, V, VII.

carota.

6. CAUCALIS involucro universali polyphylo pinnatifido, umbellis defloratis conniventibus, foliis supradecompositis, petiolis subtus nervosis.

Daucus carota, seminibus hispidis, petiolis subtus nervosis. *Sp. pl.* 348.

Daucus vulgaris. *Clus. hist.* II. 198. *R. syn.* 218.

Daucus involucris cavis, communibus pinnatis, peculiaribus lineari-lanceolatis. *Hall. hist.* 746.

Pastinaca tenuifolia sylvestris. *B. pin.* 155. *Ger. em.* 1028. *Park.* 902.

Anglis, wild Caucalis, or Carrot or Bird's-nest,

Habitat in pratis et pascuis. ♂, VI.—VIII.

Pharmac. Dauci sylvestris Semina,

CONIUM. 333.

Involucella dimidiata, subtriphyllo. *Fructus* subglobosus, quinquestriatus, utrinque crenatus.

I. CONIUM

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Cambridgeshire. *inter* St. Albans *et* Stoney-Stratford. 4. VII. VIII.

meum.

2. ATHAMANTA foliis capillaribus, seminibus glabris striatis. *Sp. pl.* 353. *Mat. med.* 126. *Jauv. monsp.* 182.

Meum. *Dod. pempt.* 305. *Ger. em.* 1052. *R. syn.* 207. *Hall. hist.* 761. *vulgatius.* *Park.* 888.

Anglis, common Spignel, Meu or Bawd-money.

Habitat in pratis montosis in comitatibus Westmorlandico, Cumberlandico, Lancastriensi frequens, near Dolgehle in Merionethshire. R. syn. 4. V.

Pharmac. Athamantæ Radix.

PEUCEDANUM. 336.

Fruetus ovatus, utrinque striatus, ala cinctus. *Involucra* brevissima.

- officinale.* 1. PEUCEDANUM foliis quinquies tripartitis filiformi-linearibus. *Sp. pl.* 353.

Peucedanum. *Ger. em.* 1054. *R. syn.* 206. *germanicum.* *B. pin.* 149.

Peucedanum vulgare. *Park.* 880.

Anglis, common Sulphur-wort, or Hogs-fennel.

Habitat in palustribus maritimis. Prope Shoreham *in* Suffexia, Walton *non procul ab* Harvico *in* Essexia; *about a quarter of a mile below* Feversham, *by the river side,* Mr. J. Sherard. *R. syn. circa* Yarmouth *et* Clay *in* Norfolk. *D. Humphrey.* 4. VI. VII.

Pharmac. Peucedani Radix.

2. PEUCEDANUM foliis pinnatifidis: laciniis oppositis, involucro universalis diphyllo. *Sp. pl.* 354. *Leis. ball.* 232. *Jac. austr. I.* 12. t. 15.

Peucedanum foliis triplicato-pinnatis, pinnulis nervo distinctis, lanceolatis, impari trilobatis. Hall. hist. 797.

Seseli pratense. *B. pin.* 162. *Park.* 905. *R. syn.* 216. *Crantz. fasc. III.* t. 6. f. 1. *Riv.* 57.

Saxifraga anglica facie seseli pratensis. *Ger. em.* 1047.

Anglis, meadow Sulphur-wort or Saxifrage.

Habitat in pratis et pascuis humidiusculis. 4. VII.

CRITH-

CRITHMUM. 337.

Fructus ovalis, compressus. *Flosculi* æquales.

1. CRITHMUM foliolis lanceolatis carnosis. *Sp. pl. maritimum.*
354. *Sauv. monsp.* 358. *Gouan. Fl. monsp.*

Crithmum marinum. *Ger. em.* 533. *Dod. pemp.* 705.

R. syn. 217. vulgare. *Park.* 1286.

Crithmum L *Foeniculum marinum minus.* *B. pin.*
288.

Anglis, sea Sampire.

Habitat in littoribus maritimis. 4. VIII.

Pharmac. *Crithmi Herba.*

HERACLEUM. 342.

Fructus ellipticus, emarginatus, compressus, striatus,
marginatus. *Cor.* difformis, inflexo-emarginata.
Involucrum caducum.

1. HERACLEUM foliolis pinnatifidis. *Hort. cliff. sphondylium.*
103. *Sp. pl.* 358. *Fl. suec.* 243. lævibus, floribus
radiatis. *Syst. nat.* 210.

Sphondylium foliis hirsutis pinnatis, pinnis quinque-
lobis. *Hall. hist.* 809.

Sphondylium. *Ger. em.* 1009. *R. syn.* 205.

Sphondylium vulgare. *Park.* 953. hirsutum. *B. pin.*
157.

- β *Heracleum* foliis cruciatio-pinnatis: foliolis lineari-angusti-
bus, corollis flosculosis. *Syst. nat.* 210. folium.

Sphondylium majus aliud laciniatis foliis. *Park.* 953.

Anglis, common Cow-parsnep.

Habitat in sepibus, pratis et pascuis. 8. VII.

Obs. β variat corollis frequentissime radiatis, rarissime
flosculosis, et in hortis β in α degenerat.

LIGUSTICUM. 343.

Fructus oblongus, quinquesulcatus utrinque. *Corollæ*
æquales. *Petalis* involutis, integris.

1. LIGUSTICUM foliis biternatis. *Sp. pl.* 359. *Fl. scoticum.*
dan. 207.

Ligusticum scoticum apii folio. *R. syn.* 214.

Ligusticum humilius scoticum. *Pluk. ph. 96. f. 2.*
 Seseli maritimum scoticum. *Herm. par. 227. t. 227.*
Anglis, scottish Lovage.

Habitat in clivis et littoribus maritimis, sed rarius. In
 tumulo quodam saxoso et arenoso ad mare sex ab Edin-
 burgo miliaribus versus Queens-ferry, et alibi in
 Scotia. *Æ. VII.*

cornubi- .2. **LIGUSTICUM** foliis decompositis ineisis: radicali-
ense. bus ternatis lanceolatis integerrimis. *Sp. pl. 359.*
 Smernium tenuifolium nostras. *R. hist. III. 254. syn.*
209. t. 8.
 Saxifraga cornubiæ. *Pet. h. br. t. 26. f. 9.*
Anglis, cornish Lovage.
 Habitat in Cornubia. *Æ. VII.*

ANGELICA. 344.

Fructus subrotundus, angulatus, solidus, styli re-
 flexis. *Corollæ* æquales: petalis incurvis.

sylvestris. 1. **ANGELICA** foliis æqualibus ovato-lanceolatis ser-
 ratis. *Sp. pl. 361. Fl. suec. 246.*
 Angelica foliis duplicato-pinnatis, ovatis, serratis.
Hall. hist. 806.
 Angelica sylvestris. *Ger. em. 999. Park. 249. R.*
syn. 208. major. B. pin. 145.
Anglis, wild Angelica.
 Habitat in sylvis et sepibus paludosis. *Æ. VII. VIII.*

SIUM. 345.

Fructus subovatus, striatus. *Invalucrum* polyphyllum.
Petala cordata.

latifolium. 1. **SIUM** foliis pinnatis, foliolis oblongo-lanceolatis
 serratis, umbellis terminalibus.
 Sium foliis pinnatis, umbellis terminalibus. *Sp. pl.*
361. Fl. suec. 235. Dan. 246. Jac. austr. I. 42.
t. 66.
 Sium foliis pinnatis, argute dentatis, umbellis erectis.
Hall. hist. 777.
 Sium latifolium. *B. pin. 154. foliis variis. R. syn.*
211.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Petroselinum macedonicum Fuchsi. *Ger. em.* 1016.
Anglis, field Honewort, or Stone-Parley.
Habitat in sylvis et sepibus subhumidis. ☉. VIII. IX.
Pharmac. Amomi semina.

segetum. 2. **SISON** foliis pinnatis, umbellis cernuis. *Sp. pl.* 362.
Hall. hist. 779. *Jac. hort. vind.* II. 63. t. 134.
Sium arvense sive *segetum.* *R. syn.* 211. *Tourn.* 308.
Sium terrestre, umbellis rarioribus. *Hist. Ox.* III.
 283. t. 5. f. 6.
Selinum sili foliis. *Ger. em.* 1018.
Selinum segetale. *Park.* 932.
Anglis, corn Honewort.
Habitat in arvis et ad sepes solo cretacco et lutofo. ☉.
 VII.

inundatum. 3. **SISON** repens umbellis bifidis. *Sp. pl.* 363. *Fl.*
Dan. 69.
Sium foliis submersis capillaribus, emerfis pinnatis.
Sauv. monsp. 249. *R. lugdb.* 105.
Sium foliis radicalibus, capillaribus multifidis, caulini-
nis pinnatis, ultima pinna trilobata. *Hall. hist.* 780.
Sium minimum, foliis imis ferulaceis. *Hist. Ox.* III.
 223. t. 9. f. 5.
Sium pusillum foliis variis. *Pb. br. R. syn.* 212.
Anglis, water Honewort.
Habitat in fossis, stagnis et inundatis passim. In ericetis
Putnejano et Hounsejano; on Wandsworth common,
inter Newington et Hornsey. ☉. V. VI.

verticillatum. 4. **SISON** foliolis verticillatis capillaribus. *Sp. pl.* 263.
Carum foliolis setaceis verticillatis, radice napiformi.
Sauv. monsp. 168.
Daucus pratensis, Dalech. hist. 718. *millefolii palustris*
folio. *B. pin.* 150.
Anglis, verticillate Honewort.
Habitat in pratis et pascuis. Plentifully in the west of
Scotland, near Greenock. Mr. W. Houston, prope
Air in Scotia; in pascuis humidiusculis in Carmar-
thenshire; inter Treacastle et St. Davids in Pem-
brokeshire copiose, D. Banks. 4. VIII.

OENANTHE. 349.

Flosculi difformes : in disco sessiles, steriles. *Fructus*
calyce coronatus.

1. OENANTHE stolonifera, foliis caulinis pinnatis *fi- fistulosa*.
liformibus fistulosis. *Sp. pl.* 365.
Oenanthe foliis radicalibus rotunde lobatis planis,
caulinis fistulosis. *Hall. hist.* 755.
Oenanthe aquatica. *B. pin.* 162. *R. syn.* 210.
Oenanthe. *Riv. pent.* 66. *palustris.* *Park.* 895.
Filipendula aquatica. *Ger. em.* 1060.
- β Oenanthe aquatica triflora, caulibus fistulosis. *Hist.*
Ox. III. 289. t. 7. f. 8.
Anglis, common Water-dropwort.
Habitat in fossis et palustribus frequens. 4. VII.
2. OENANTHE foliis radicalibus cuneatis fissis ; cau- *pimpinell-*
linis integris linearibus longissimis canaliculatis. *loides.*
Sp. pl. 367.
Oenanthe foliis radicalibus duplicato-pinnatis lineari-
bus, recurvis. *Hall. hist.* 756.
Oenanthe apii folio. *B. pin.* 162.
Oenanthe aquatica pimpinellæ saxifragæ divisura nos-
tras. *Pluk. ph. t.* 49. f. 4.
Anglis, parsley Water-dropwort.
Habitat in fossis et palustribus, in parochia Quaplod
agri Lincolnensis non procul ab oppido Spalding ; D.
Pluk. *R. syn.* in paludibus maritimis frequens. 4.
VII. VIII.
3. OENANTHE foliis omnibus multifidis obtusis sub- *crocata.*
æqualibus. *Hort. cliff.* 99. *Sp. pl.* 365.
Oenanthe cicutæ facie Lobelii. *Park.* 894. *R. syn.*
210.
Oenanthe chærophylli foliis. *B. pin.* 162.
Oenanthe cicutæ facie, succo viroso crocante. *Lob.*
adv. 326. *Act. angl.* 1747. N. 480. 235. t. 3.
Filipendula cicutæ facie. *Ger. em.* 1059.
Anglis, hemlock Water-dropwort.

Habitat

PENTANDRIA DIGYNIA.

Habitat in aquosis et ad ripas fluviorum passim. ad ripas Thamesis frequens. 4. VI. VII.

B U N I U M. 332.

Cer. uniformis; *Umbella* conferta. *Fruetus* ovatus.

- Bulbocastanum*.
vanum. 1. BUNIMUM. *Sp. pl.* 349. *Fl. dan.* 220. *Hall. hist.* 783.
Bulbocastanum. *B. hist.* III. 30. *R. syn.* 209.
Bulbocastanum majus apii folio. *B. pin.* 1065.
Bulbocastanum majus et minus. *Ger. em.* 1065.
Nucula terrestris. *Lob. hist.* 429. *major*. *Park.* 893.
Anglis, Earth, Kipper, Pig, or Hawk-nut.
Habitat in pratis, pascuis et sylvis. 4. V. VI.

B. Involucris, partialibus: universali nulla.

C I C U T A. 351.

Fruetus subovatus, sulcatus.

- virosa*. 1. CICUTA umbellis oppositifoliis, petiolis marginatis obtusis. *Sp. pl.* 366. *Fl. dan.* 208.
Cicuta aquatica. *Gesn. hort.* 253. *coll.* 68.
Sium foliis duplicato-pinnatis, pinnulis trifidis et simplicibus, acute serratis. *Hall. hist.* 781.
Sium alterum. *Dod.* 589. *olusatrici facie*. *Lob. ic.* 208.
Ger. em. 256. *R. syn.* 212.
Sium majus alterum angustifolium. *Park.* 1241.
Sium erucæ folio. *B. pin.* 154.
Anglis, water Cicuta.
Habitat in aquosis et fluviis. In some pools of water at Thistleworth. *R. syn.* in fluvio Coln dicto prope Colnbrook, Middlesex. 4. VII. VIII.

P H E L L A N D R I U M. 350.

Flosculi disci minores, *Fruetus* ovatus, lævis, coronatus perianthio et pistillo.

- aquatium*. 1. PHELLANDRIUM foliorum ramificationibus divaricatis. *Sp. pl.* 366.
Phellandrium foliis refractis. *Hall. hist.* 757.
Phellandrium vel Cicutaria aquatica quorundam. *B. hist.* III. 181. *R. syn.* 215.

Cicutaria

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Anglis, corn Cicely or Shepherd's-needle, or Venus's-comb.

Habitat in arvis. ☉. VI. VII.

- odorata.* *2. SCANDIX seminibus sulcatis angulatis. *Sp. pl.* 368. Myrrhis major, Cicutaria odorata. *B. pin.* 160. Myrrhis. *Dodd. pempt.* 701. major vulgaris s. cerefolium majus. *Park.* 935. Cerefolium magnum s. myrrhis. *Ger. em.* 1039. *Anglis*, sweet Cicely. Habitat in sepibus et pomariis in comitatibus Westmorlandico, Cumberlandico et Lancasteriensi frequens. ☉. VI.

- Anthriscus.* 3. SCANDIX seminibus ovatis hispidis, corollis uniformibus, caule lævi. *Sp. pl.* 368. *Fac. austr. II.* 35. t. 154. *Caucalis vaginis lanuginosis, foliis triplicato-pinnatis, seminibus rostratis, Hall. hist.* 743. Myrrhis sylvestris seminibus asperis. *B. pin.* 160. *R. syn.* 220. Myrrhis sylvestris neapolitana atque anglicana. *Park.* 935. Cerefolium sive Myrrhis nova æquicolorum Columnæ. *Ger. em.* 1038. *Anglis*, rough Cicely, or Chervil. Habitat in aggeribus ruderatis; circa Londinum ubique. ☉. V. VI.

CHÆROPHYLLUM. 355.

Involucrum reflexum, concavum. *Petala* inflexo-cordata. *Fruetus* oblongus, lævis.

- sylvestre.* 1. CHÆROPHYLLUM caule lævi striato, geniculis tumidiusculis. *Sp. pl.* 369. *Fac. austr. II.* 31. t. 149. Cerefolium foliis acute dentatis, triplicato pinnatis, glabris, nervis hirsutis. *Hall. hist.* 748. Cicutaria vulgaris. *B. hist. III.* 181. *R. syn.* 207. Cicutaria alba Lugdunensis. *Ger. em.* 1038. Myrrhis sylvestris. *Park.* 935. seminibus lævibus. *B. pin.* 160.

Anglis,

Anglis, wild Chervil, Cicely, or Cow-weed.
Habitat ad sepes. ☉. V. VI.

2. CHÆROPHYLLUM caule scabro: geniculis tu-temulen-
 midis. *Sp. pl.* 370. umbellis cernuis. *ed. I. tum.*
 258.

Chærophyllum sylvestre. *B. pin.* 152.

Cerofolium sylvestre. *Ger. em.* 1038. *Park.* 915.

R. syn. 207.

Myrrhis foliis hirsutis, laciniis obtusis, caule genicu-
 lato. *Hall. hist.* 750.

Anglis, rough Chervil.

Habitat ad sepes frequens. ☉. VII. VIII.

C. Involucro nullo; nec universali, nec partialibus.

PASTINACA. 359.

Fructus ellipticus, compresso planus. *Petala* invo-
 luta, integra.

1. PASTINACA foliis simpliciter pinnatis. *Sp. pl. sylvestris.*
 376. *Fl. suec.* 259. *Gart. Gelr.* 60.

Pastinaca foliis pinnatis. *Hall. hist.* 808.

Pastinaca sylvestris latifolia. *B. pin.* 155. *R. syn.*
 206.

Pastinaca latifolia sylvestris. *Ger. em.* 1025. *Park.*
 944.

Pastinaca germanica sylvestris, quibusdam Elaphobos-
 cum. *B. hist. III.* 149.

- β Pastinaca latifolia sativa. *Ger. em.* 1025. *Park.*
 944.

Anglis, wild Parsnep. β garden Parsnep.

Habitat in marginibus agrorum, solo calcareo. ☉.
 VII.

SMYRNIUM.

PENTANDRIA DIGYNIA.

SMYRNIUM. 360.

Fruetus oblongus, striatus. *Petala* acuminata, carinata.

Olusatrum.

1. SMYRNIUM foliis caulinis ternatis petiolatis serratis. *Sp. pl.* 376.

Smyrniium. *Matth.* 773. *R. syn.* 208.

Hippofelinum. *Ger. em.* 1019. f. Smyrniium vulgare. *Park.* 930.

Hippofelinum Theophrasti f. Smyrniium Dioscoridis. *B. pin.* 154.

Anglis, common Alexanders.

Habitat in fossis et in rupibus maritimis; about Scarborough Castle, near Deptford, Battersea, Vauxhall, about Nottingham plentifully. *R. syn.* 8. V. VI.

Pharmac. Smyrnii Radix, Herba, semina.

ANETHUM. 361.

Fruetus subovatus, compressus, striatus. *Petala* involuta, integra.

Fœniculum.

1. ANETHUM fructibus ovatis. *Sp. pl.* 377.

Fœniculum. *Cam. epit.* 534. vulgare. *Ger. em.* 1032. *Park.* 884. *R. syn.* 217. Germanicum. *B. pin.* 147.

β Fœniculum dulce. *B. pin.* 147.

Anglis, common Fennel, or Finckle.

Habitat in clivis cretaceis frequens. 8. VII. VIII.

Pharmac. Fœniculi semina.

CARUM. 362.

Fruetus ovato-oblongus, striatus. *Involucrum* monophyllum. *Petala* carinata, inflexo-emarginata.

Carvi.

*1. CARUM. *Dod. pempt.* 299. *Cam. epit.* 516. *Sp. pl.* 378. *Hall. hist.* 789. vulgare. *Park.* 910.

Carum seu Careum. *Ger. em.* 1934. *R. syn.* 213.

Cuminum pratense Carui officinarum. *B. pin.* 158.

Anglis, Caraways.

Habitat in pratis et pascuis passim. 8. V. VI.

Pharmac. Carui semina.

PIMPINELLA,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

dioica.

3. PIMPINELLA foliis subbipinnatis : foliolis linearibus simplicibus ternatisque, floribus dioicis.

Pimpinella pumilla umbellis numerosissimis compositis simplicibusque. *Jacq. enum.* 52. et 227. *austr.* I. 19. t. 28.

Seseli petiolis ramiferis membranaceis oblongis integris, foliolis ternatis linearibus subcarnosis petioli longitudine. *Sp. pl.* 373. *Fl. gall. prov.* 253.

Fœniculum minimum patulum. *Clus. hist.* II. 200.

Peucedanum pumilum. *Ger. em.* 1054.

Peucedanum minus. *Park.* 880. *R. syn.* 217.

Daucus montanus, multifido folio, felini semine. *B. pin.* 150.

Anglis, least Pimpinell, or Burnet-saxifrage.

Habitat in pratis montosis. On St. Vincent's Rock, near Bristol. *R. syn. supra* Uphill in agro Somersetensi. 4. V. VI.

Desc. Radix perennis, fusiformis, perpendicularis, basi subramosus, capite villoso. Caulis semicubitalis et cubitalis, erectus, angulatus, striatus, foliatus, glaber, ramosissimus ; ramis patulis. Folia inferiora opposita, petiolato, pinnata et bipinnata ; foliolis, simplicibus, ternatisque, linearibus, canaliculatis, nudis, glaucis. Petiolus membranaceus vaginans oblongus, striatus, integer. Umbellæ pedunculatæ axillares terminalesque, universales radice circiter sex longis, partiales confertæ, radiis circiter quinque, brevibus. Flores lutescentes vel albidii dioici. Flo. Mas. Cal. obsoletus. Corolla petala quinque, oblonga, apice inflexa, æqualia. Stam. Filamenta quinque, subulata. Antheræ subrotundæ. Pist. &c. nulla. Flo. Fem. Cal. Cor. ut in masculo. Stam. Filamenta quinque subulata. Antheræ steriles. Pist. germen inferum ; styli duo distantes. Stigmata globosa. Pericarpium nullum. Fructus ovatus, striatus, bipartibilis. Sem. duo, ovata, hinc convexa, striata, inde plana.

A P I U M. 364.

Fructus ovatus, striatus. *Involucrum* monophyllum. *Petala* æqualia.

• I. APIUM

1. APIUM foliis caulinis cuneiformibus, umbellis ses- *graveo-*
silibus. *Sp. pl.* 379. *lens.*
 Apium foliis pinnatis pinnis trilobatis. *Hall. hist.* 784.
 Apium palustre et off. *B. pin.* 154. *R. syn.* 214.
 Apium vulgare s. palustre. *Park.* 926.
 Eleoselinum seu Paludapium. *Ger. em.* 1014.
Anglis, smallage Parsley.
 Habitat in palustribus et ad fossas. 8. VIII.

Æ G O P O D I U M. 365.

Fruetus ovato-oblongus, striatus.

1. ÆGOPODIUM foliis caulinis summis ternatis. *Sp. Podagra-*
pl. 379. *ria.*
 Podagraria. *Riv. pent. t.* 46. *Hall. hist.* 759. *vulgare.*
Park. 943.
 Angelica sylvestris minor s. erratica. *B. pin.* 155. *R.*
syn. 208.
 Herba Gerardi. *Dod. pemp.* 320. *Ger. em.* 1001.
Anglis, Gout-weed, Herb-gerard, or Ash-weed.
 Habitat ad sepes et in oleraceis, hortis, frequens. 4. V.
 VI.

T R I G Y N I A.

* *Flores superi.*

V I B U R N U M. 367.

Cal. quinquepartitus. Cor. quinquefida. Bacca mo-
nosperma.

1. VIBURNUM foliis cordatis serratis venosis subtus *Lantana.*
tomentosis. *Sp. pl.* 381. *Hall. hist.* 669.
 Viburnum. *Cam. epit.* 122. *B. pin.* 426. *Park.* 1448.
R. syn. 460.
 Lantana sive Viburnum. *Ger. em.* 1490.
Anglis, common Viburnum, or mealy or way-faring
Tree.
 Habitat in sylvis et sepibus solo calcareo. 1/2. V.

K

2. VIBURNUM

- Opulus.* 2. **VIBURNUM** foliis lobatis petiolis glandulosis. *Sp. pl.* 384.
Opulus. *Ruellii* 281. *R. syn.* 460. *Hall. hist.* 668.
Sambucus aquatica, flore simplici. *B. pin.* 456.
Sambucus palustris. *Cam. ep.* 977. *f. aquatica.* *Park.* 208.
Sambucus aquatilis *f. palustris.* *Ger. em.* 1424.
Anglis, water Viburnum, or Elder.
Habitat in sylvis et sepibus humidis. h. V. VI.

S A M B U C U S. 369.

Cal. quinquepartitus. *Cor.* quinquefida. *Bacca* triperma.

- Ebulus.* 1. **SAMBUCUS** cymis tripartitis, stipulis foliaceis, caule herbaceo. *Sp. pl.* 385.
Sambucus herbacea floribus umbellatis. *Hall. hist.* 671.
Sambucus humilis *f. Ebulus.* *B. pin.* 456. *R. syn.* 461.
Ebulus. *Fuchs. hist.* 65. *Cam. epit.* 979. *f. sambucus humilis.* *Ger. em.* 1426. *Park.* 209.
Anglis, dwarf Elder, or Danewort.
Habitat in sepibus et ad vius. u. VII.
Pharmac. *Ebuli Radix, Folia, Flores, Bacca.*

- nigra.* 2. **SAMBUCUS** cymis quinquepartitis, caule arboreo.
Sp. pl. 385. *Fl. dan.* 545.
Sambucus arborea floribus umbellatis. *Hall. hist.* 670.
Sambucus. *Cam. epit.* 975. *Ger. em.* 1422. *vulgaris.* *Park.* 207. *R. syn.* 461.
Sambucus fructu in umbella nigri. *B. pin.* 456.
B *Sambucus fructu in umbella viridi.* *B. pin.* 456.
γ *Sambucus laciniato folio.* *B. pin.* 456.
Anglis, common Elder, *γ* Parsley Elder.
Habitat in sylvis et sepibus humidis. h. V.
Pharmac. *Sambuci Cortex, Folia, Flores, Bacca.*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

PENTANDRIA PENTAGYNIA.

Parnassia vulgaris et palustris. *Tourn.* 246. *R. syn.* 555.

Gramen parnassi minus. *Ger. em.* 840.

Gramen parnassi vulgare. *Park.* 429.

Gramen parnassi albo simplici flore. *B. pin.* 309.

Anglis, grass of Parnassus.

Habitat in pratis humidis in Wallia, et comitatibus Lancastriensi, Westmorlandico, Cumberlandico, etc. frequens. 4. VIII.

PENTAGYNIA.

* *Flores inferi.*

STATICE. 384.

Cal. monophyllus, integer, plicatus, scariosus. Petala quinque. Sem. unicum.

Armeria. 1. STATICE scapo simplici capitato, foliis linearibus. *Sp. pl.* 394. *Hall. hist.* 835.

Statice montana minor. *Tourn.* 341. *R. syn.* 203.

Limonium majus flore globofo. *Hist. Ox. III.* 601. t. 1. f. 29.

Caryophyllus montanus minor. *B. pin.* 211.

Caryophyllus marinus minimus. *Lob. Ger. em.* 602.

Gramen marinum minus. *Park.* 1279.

Anglis, common Thrift, or sea Gilly-flower.

Habitat in pratis et rupibus maritimis et alpinis, in monte Snowdon et alibi in Wallia et in comitatu Eboracensi prope Settle et Ingleborough. 4. VII. VIII.

Limonium. 2. STATICE scapo paniculato tereti, foliis lævibus enerviis. *Sp. pl.* 394. *Fl. dan.* 315.

Limonium. *Ger. em.* 411. *R. syn.* 201. majus vulgatius. *Park.* 1234.

Limonium maritimum majus. *B. pin.* 192.

β *Limonium anglicum minus, caulibus ramosioribus, floribus in spicis rarius sitis.* *R. hist. III.* 247. *syn.* 202.

γ *Limonium minus.* *R. syn.* 202. *Anglis, sea Thrift, or Lavender.*

Habitat

Habitat in littoribus maritimis. 4. VII. IX.

Pharmac. Limonii Radix.

3. STATICE scapo paniculato prostrato, ramis sterilibus *reticulata* retroflexis nudis, foliis cuneiformibus. *Sp. pl.* 394. *ta.*

Limonium minus, virgulis retiformiter se invicem implicatis. *Pluk. ph. t.* 42. *f.* 4.

Limonium reticulatum. *R. hist.* 396. *Bocc. sic.* 82. *t.* 44.

Anglis, matted Thrift or Sea-lavender.

Habitat in littoribus maritimis in comitatu Norfolk. 4. VII. VIII.

L I N U M. 385.

Cal. pentaphyllus. *Petala* quinque. *Caps.* quinquevalvis, decemlocularis. *Sem.* solitaria.

* *Foliis alternis.*

1. LINUM calycibus capsulisque mucronatis, petalis *usitatissimo* crenatis, foliis lanceolatis alternis, caule subsolitario. *Sp. pl.* 397.

Linum arvense. *B. pin.* 214. *R. hist.* 1073.

Linum sylvestre vulgatius. *Park.* 1334.

- β Linum sativum. *Ger. em.* 556. *R. syn.* 362.

Anglis, common Flax.

Habitat in arvis et pascuis arenosis in comitatibus Dorsetsiensi et Devonienſi passim. ☉. VII.

Pharmac. Lini semina.

2. LINUM calycibus capsulisque obtusiusculis, foliis *perenne* alternis lanceolatis integerrimis. *Sp. pl.* 397. *Mill. fig. t.* 166. *f.* 2.

Linum sylvestre cæruleum perenne erectius, flore et capitulo majore. *R. syn.* 362.

Linum perenne majus cæruleum capitulo majore. *Hist. Ox. II.* 573.

- β Linum sylvestre cæruleum perenne procumbens, flore et capitulo minore. *R. syn.* 362.

Anglis, perennial Flax.

Habitat in pascuis et pratis solo calcareo. In agri Cantabrigienſis collibus Gogmagog, dictis copiose; at Crosby Ravensworth, and between Shap and Threapland,

PENTANDRIA PENTAGYNIA.

land; Westmoreland, Mr. Lawson. *R. syn. prope Stamford copiose* 24. VI. VII.

Obs. variat caule erecto et decumbente ex eadem Radice; Staminibus pistillis longioribus in planta erecta, et Staminibus pistillis brevioribus in planta decumbente.

angustifolium. 3. LINUM calycibus capsulisque acuminatis, foliis lineari-lanceolatis acuminatis, caule basi ramoso subdeclinato.

Linum sylvestre angustifolium floribus dilute purpurascens vel carneis. *B. pin. 214. R. syn. 362.*

Linum sylvestre VI angustifolium. *Clus. hist. 1. 318.*

Linum foliis linearibus, calycibus longe aristatis. *Hall. hist. 838.*

Anglis, narrow-leaved Flax.

Habitat in pratis et pascuis aridis. In Cornubia et Devoniam frequens; prope Minster in insula Shepey, prope Deal, et alibi in Cantio. 24. VI. VII.

*** *Foliis oppositis.*

catharticum. 4. LINUM foliis oppositis ovato-lanceolatis, caule dichotomo, corollis acutis. *Sp. pl.*

Linum foliis conjugatis, ovatis, calycibus aristatis patulis, lanceolatis. *Hull. hist. 839.*

Linum sylvestre catharticum. *Ger. em. 560. R. syn. 362.*

Linum sylvestre, flosculis exiguis. *B. pin. 214.*

Chamælinum Clusii flore albo, s. Linum sylvestre catharticum. *Park. 1336.*

Anglis, purging Flax.

Habitat in pratis et pascuis siccioribus. ☉. V.—VIII. *Pharmac. Lini Cathartici Herba.*

radiola. 5. LINUM foliis oppositis, caule dichotomo, floribus tetrandris tetragynis. *Sp. pl. 402. Fl. dan. 178.*

Linum tetrapetalon, caule ramosissimo, dichotomo. *Hall. hist. 840.*

Chamælinum vulgare. *Vaill. paris. 33. t. 4. f. 6.*

Radiola vulgaris serpyllifolia. *R. syn. 345. t. 15. f. 3.*

Mille-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

PENTANDRIA POLYGYNIA:

Habitat in paludosis in comitatibus Westmorlandico, Cumberlandico, Eboracensi, Lancastriensi, Devoniensi, Hantonensi, et Norfolkensi passim. 4. VII. VIII.

Obs. habet multa communia cum precedente, sed major et capsulis quadrivalvis.

SIBBALDIA. 389.

Cal. pentaphyllus decemfidus. Petala quinque calyce inserta. Styli e latere germinis. Sem. quinque.

procumbens.

1. SIBBALDIA foliis tridentatis. Fl. lapp. III. suec. 216. Dan. 32. Sp. pl. 406. Pen. voy, II. 43. t. 5.

Fragaria foliis ternatis, retusis tridentatis, floris calyci equali pentastemon. Hall. hist. 1116.

Fragariæ sylvestri affinis planta, flore luteo. Sibb. pr. hist. nat. scot. p. 11. 25. t. 6. f. 1.

Pentaphylloides pumila foliis ternis ad extremitates trifidis. R. syn. 256.

Pentaphylloides fruticosum minimum procumbens, flore luteo, foliis sericeis fragariæ ternis. Pluk. ph. t. 212. f. 3.

Anglis, procumbent Sibbaldia.

Habitat in montosis; in sylvis regionis Jernensis Scotiæ sponte provenit. R. syn. in monte Ben Lomond dicto in Scotia. 4. VII. VIII.

POLYGYNIA.

MYOSURUS. 390.

Cal. pentaphyllus, basi adnatus. Neclaria quinque subulata petaliformia. Sem. numerosa.

minimus.

1. MYOSURUS foliis integris. Sp. pl. 407. Fl. dan. 406.

Myosurus. B. hist. III. 512. R. syn. 251.

Holosteum affinis, Cauda muris. B. pin. 190.

Holosteum Loniceri, Cauda muris vocatum, Park. 500. Cauda muris. Ger. em. 426.

Anglis, Mouse-tail.

Habitat in arvis, pratis et pascuis solo glareoso; inter segetes circa Wimbleton, inter Kentish-Town et Hampstead; prope Marylebon capiose. ①. IV. V.

Obs. variat foliis hirsutis.

Classis

Classis VI.

H E X A N D R I A
M O N O G Y N I A.

- *Flores Caliculati calyce corollaque instructi.*

B E R B E R I S. 442.

Cal. hexaphyllus. *Petala* sex: ad unguis glandulis duobus. *Stylus* 0. *Bacca* disperma.

1. BERBERIS pedunculis racemosis. *Sp. pl.* 541. *vulgaris*
 Berberis racemifera, foliis ciliatis. *Hall. hist.* 828.
 Berberis. *Park.* 1559. dumetorum. *B. pin.* 454. *R.*
syn. 465.
 Spina acida sive Oxyacantha. *Ger. em.* 1325.
Anglis, common Barberry, or Pipperidge-bush.
 Habitat in sylvis et sepibus. *h.* V. VI.
Pharmac. Barberidis Cortex, Bacca.

F R A N K E N I A. 445.

Cal. quinquefidus, infundibuliformis. *Petala* quinque. *Stigmata* tria. *Caps.* unilocularis, trivalvis.

1. FRANKENIA foliis linearibus confertis basi ciliatis. *lævis.*
Sp. pl. 473.
 Franca maritima supina saxatilis glauca ericoides, sempervirens, flore purpureo. *Mich. gen.* 23. t. 22. f. 1.
 Lychnis supina maritima ericæ facie. *R. syn.* 338.
 Erica supina maritima. *Ph. br. anglica.* *Park.* 1484.
 Polygonum maritimum minus, foliolis serpylli. *B. pin.* 281.
 Polygonum pusillo vermiculato serpyllifolio. *Ger. em.* 567.
Anglis, smooth Frankenia or Sea-Heath.
 Habitat in palustribus maritimis solo lutofo. In Lovingsland, just over the water at Yarmouth. *R. syn.* . In insulis Shepey et Thanet copiose, et in comitatibus Essexiensi,

Essexiensi, Suffexiensi et Cantiano frequens. 4.
VII. VIII.

- pulveru-
lenta.* 2. **FRANKENIA** foliis obovatis retusis subtus pulveru-
lentis. *Sp. pl.* 474.
Franca maritima quadrifolia supina, chamæsyces folio
et facie. *Mich. gen.* 23.
Alsine maritima supina foliis chamæsyces. *Tourn.*
665. *R. syn.* 352.
Anthyllis valentina. *Clus. hist.* II. 186.
Anthyllis marina, chamæsyces similis. *B. pin.* 282.
Anglis, dusty Frankenia or Sea-heath.
Habitat in littoribus maritimis. Found on the coast of
Sullex, and sent by Mr. Brewer. *R. syn.* In littore
Suffexiæ inter Bognor et Brightelmston inveni. ④.
VII.

* *Flores spathacei.*

A L L I U M. 409.

Cor. sexpartita, patens. *Spatha* multiflora. *Umbella*
congesta. *Caps.* supera.

* *Foliis caulibus planis.*

- ampelo-
prasum.* 1. **ALLIUM** caule planifolio umbellifero, umbella
globosa, staminibus tricuspидatis, petalis carina
scabris. *Sp. pl.* 423.
Allium radice solumifera, foliis gramineis, spica sphæ-
rica, staminibus alterne trifidis. *Hall. hist.* 1218.
Allium holmense sphærico capite. *R. syn.* 370.
Allium sphærico capite, folio latiore s. Scorodopra-
sum. *B. pin.* 74. *Rudb. ethn.* II. 151.
Scorodoprasum primum Clusii. *Ger. em.* 180.
Anglis, round-headed Garlick.
Habitat in parva quadam insula Holms dicta in Sabri-
næ æstuario copiose, D. Newton. *R. syn.* 4. VII.

- arena-
rium.* 2. **ALLIUM** caule planifolio, vaginis teretibus, spatha
mutica, staminibus tricuspидatis. *Sp. pl.* 426. *It.*
scan. 227. *Fl. dan.* 290.

Allium

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

oleraceum. 5. **ALLIUM** caule teretifolio bulbifero, foliis scabris semiteretibus subtus sulcatis, staminibus simplicibus. *Sp. pl.* 429.

Allium bulbo simplici, foliis fistulosis, spatha bicorni, umbella bulbifera. *Hall. hist.* 1223.

Allium umbella bulbifera, vagina bicorni, foliis teretibus. *Hall. opusc. All.* 26. t. 1. f. 4.

Allium montanum bicorne, flore exalbido. *B. pin.* 75.

Allium sylvestre bicorne flore exherbaceo albicante, cum triplici in singulis petalis stria atro-purpurea.

R. syn. 370.

Anglis, herbaceous Garlick.

Habitat in pratis et pascuis, et inter segetes. In a cornfield in Black-notley in Essex, belonging to the hall called Westfield, adjoining to Leeze-lane plentifully.

R. syn. In arvis in Westmorlandia passim; prope Bristolium copiose. 4. VII.

*** *Foliis radicalibus, scapo nudo.*

ursinum. 6. **ALLIUM** scapo nudo triquetro, foliis lanceolatis petiolatis, umbella fastigiata. *Sp. pl.* 431.

Allium foliis radicalibus petiolatis, ovato-lanceolatis, floribus umbellatis. *Hall. hist.* 1228.

Allium sylvestre latifolium. *B. pin.* 74. *R. syn.* 370.

Allium ursinum. *Ger. em.* 179. *Park. par.* 514.

Anglis, wild Garlick or Ramson.

Habitat in sylvis et sepibus frequens. 4. V.

G A L A N T H U S. 401.

Petala tria concava. *Nectarium* ex petalis tribus, parvis, emarginatis. *Stigma* simplex.

nivalis. *1. **GALANTHUS.** *Sp. pl.* 413.

Leucoium bulbosum trifolium minus. *B. pin.* 56.

Leucoium bulbosum præcox minus. *Ger. em.* 147.

Anglis, Snow-drop.

Habitat in pratis, et ad sepes; in pomariis frequens, in comitatibus Westmorlandico, Cumberlandico, Lancasteriensi, Gloucesterensi. 4. III, IV.

NARCISsus. 403.

*Petala sex. Nectario infundibuliformi, monophyllo.
Stam. intra nectarium.*

*1. NARCISsus spatha uniflora, nectario rotato brevissimo scariofo crenulato. *Hort. ups. 74. Sp. pl. 414.*

Narcissus uniflorus, foliis ensiformibus, scypho brevissimo. *Hall. hist. 1250.*

Narcissus medio luteus. *Ger. em. 124. vulgaris. Park. par. 74. R. syn. 371.*

Narcissus pallidus circulo luteo. *B. pin. 51.*

Anglis, pale Narcissus or Daffodil.

Habitat in pratis et ad sepes, sed rarius. 4. V.

2. NARCISsus spatha uniflora, nectario campanulato erecto crispo æquante petala ovata. *Sp. pl. 414. pseudo-narcissus.*

Narcissus foliis ensiformibus, scapo unifloro, scypho undulato, ferrato, petales æquali. *Hall. hist. 1252.*

Narcissus sylvestris pallidus calyce luteo. *B. pin. 52. R. syn. 371.*

Pseudo-Narcissus anglicus. *Ger. em. 133. vulgaris. Park. par. 100.*

Anglis, common Narcissus or Daffodil.

Habitat in sylvis et ad sepes. - In sylvis prope Erith in agro Cantiano. 4. IV.

*** *Flores nudi.*

HYACINTHUS. 427.

Cor. campanulata: pori tres melliferi an apicem germinis.

1. HYACINTHUS corollis cylindraneo-campanulatis sexpartitis apice revolutis, bracteis binatis. *nonscriptus.*

Hyacinthus corollis campanulatis sexpartitis apice revolutis. *Sp. pl. 453.*

Hyacinthus anglicus. *Ger. em. 111. R. syn. 373. belgicus. s. hispanicus. Park. par. 122.*

Hyacinthus oblongo flore cæruleus major. *B. pin. 43. Anglis,*

Anglis, common Hyacinth or Harebells.
Habitat in sylvis et ad sepes. 4. V.

S C I L L A. 419.

Cor. hexapetala; patens, decidua. *Filamenta* filiformia.

verna.

1. SCILLA corymbo hemisphærico paucifloro bracteato, foliis subulato-linearibus canaliculatis.

Ornithogalum hispanicum minus. *Clus. hist.* 188.

Ger. em. 166. *Park. par.* 139. *R. hist.* 1153.

Ornithogalum umbellatum flosculis ex albo subcæruleis. *B. pin.* 70.

Anglis, vernal Squil.

Habitat in pratis et pascuis; in clivis maritimis in insula Mona, et in Cambria; circa St. Ives copiose et alibi in Cornubia; hæc planta ex Scotia sub nomine Scillæ bifoliæ accipi, quæ nec Scotiæ nec Angliæ indigena est. 4. IV. V.

Desc. Radix bulbosa, tunicata fusca, magnitudine nucis avellanæ. Scapus a duabus ad sex pollicares, erectus, teres, aphyllus, lævis, glaber. Folia plurima subulato-linearia, acuta, canaliculata, longitudine fere Scapi. Flores corymbosi a tribus ad octo; et in hortis a sex ad sedecim pallide cærulei: Corymbus hemisphæricus. Bracteæ persistentes, membranaceæ, concavæ, acutæ longitudine fere pedunculi.

- autumna-* 2. SCILLA foliis filiformibus linearibus, floribus corymbosis, pedunculis nudis adscendentibus longitudine floris. *Sp. pl.* 243.

Hyacinthus autumnalis minor. *Ger. em.* 110. *Park. par.* 132. *R. syn.* 373.

Hyacinthus stellaris autumnalis minor. *B. pin.* 47.

Anglis, autumnal Squil or Star-hyacinth.

Habitat in pascuis siccis. In rupe S. Vincentii prope Bristolum inque promontorio, Lizard Point, dicto copiose; eundem in ericeto nigro in comitatu Cantii invenit D. Plukenet, near Ditton, on Moulsey-Hurst, over against Hampton-Court. *R. syn.* 4. VIII, IX.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

HEXANDRIA MONOGYNIA.

Ornithogalum stipulis maximis, petiolis lateralibus longissimis. *Hall. hist.* 1215.

Ornithogalum umbellatum medium angustifolium. *B. pin.* 70. *R. syn.* 372.

Ornithogalum vulgare. *Ger. em.* 165. *Park. par.* 136. *Anglis*, common Star of Bethlehem.

Habitat in sylvis, pratis et pascuis, in Norfolk. D. Stillingfleet. 4. IV. V.

FRITILLARIA. 411.

Cor. hexapetala, campanulata, supra unguem cavitate nectarifera. *Stum.* longitudine corollæ.

meleagris. *1. FRITILLARIA foliis omnibus alternis, caule unifloro. *Sp. pl.* 346. *Syst. nat.* 141.

Fritillaria caule paucifloro, foliis caulinis gramineis alternis. *Hall. hist.* 1235.

Fritillaria præcox purpurea variegata. *B. pin.* 64.

Fritillaria vulgaris. *Park. par.* 40.

Fritillaria variegata. *Ger. em.* 149.

Anglis, common Fritillary, checquered Daffodil or Snake's-head.

Habitat in pratis et pascuis: In Mawde Fields near Rislip Common, Middlesex. *Blackf. Spec. bot.* 23. *prope* Bury, D. Cullum; in pratis inter Mortlake et Kew, et *prope* Enfield. 4. IV. V.

ANTHERICUM. 422.

Cor. hexapetala, patens. *Caps.* ovata.

serotinum. 1. ANTHERICUM foliis planiusculis, scapo unifloro. *Sp. pl.* 444.

Bulbocodium foliis subulato-linearibus. *Roy. lugdb.* 41.

Bulbocodium alpinum juncifolium, flore unico: intus albo extus squalide rubente. *R. syn.* 274. t. 17.

f. 1.

Pseudo-Narcissus gramineo folio. *B. pin.* 51. *pr.* 27.

Phalangium foliis crassulis, scapo unifloro. *Hall. hist.* 1209.

Anglis, bulbous Anthericum.

Habitat

Habitat in excelsis rupibus montis Snowdon, et prope Llanberrys. v. g. Trigvylchau y. Clogwyn du yn-hem y Glydir, Clogwyn yr Ardhu, Crib y Distil, &c. L. Lwyd. ad latis occidentale montis Trigvylchau. D. Richardson. R. syn. in rupibus Crib y Distil inveni. 2. IV. V.

NARTHECIUM. *Fl. angl.* 127.

CAL. nullus.

COR. Petala sex, æqualia, lanceolata, acuta, erectiuscula persistentia.

STAM. Filamenta sex, subulata, lanata, Antheræ parvæ incumbentes.

PIST. Germen triquetrum. Stylus nullus. Stigma obtusum.

PER. Capsula avata, acuta, triquetra, trilocularis, trivalvis.

SEM. numerosa paleacea.

I. NARTHECIUM. *Moehr. E. N. C.* 1742. 389. *offisragum.*
1. 5. f. 1.

Anthericum foliis ensiformibus, filamentis lanatis. *Sp. pl.* 446. *Fl. suec.* 287. *sibir.* I. 73. t. 18. f. 1. *dan.* t. 42.

Phalangium anglicum palustre Iridis folio. *R. syn.* 375.

Asphodelus Lancastriæ verus. *Ger. em.* 95.

Pseudo-Asphodelus palustris anglicus. *B. pin.* 29.

Anglis, yellow Narthesium or Lancashire Asphodel.

Habitat in paludibus cæspitosis et turfosis. 2. VII. VIII.

ASPARAGUS. 424.

Cor. sexpartita, erecta: petalis tribus interioribus apice reflexis. *Bacca* trilocularis, disperma.

I. ASPARAGUS caule herbaceo suberecto, foliis setaceis, stipulis duabus interioribus, una exteriori. *officinalis.*
Sp. pl. 448.

Asparagus. *Cam. epit.* 259. *Ger. em.* 1110. *Park.* 454. *R. syn.* 267. sativus. *B. pin.* 489.

Asparagus sylvestris, tenuissimo folio. *B. pin.* 490.

Asparagus maritimus, crassiore folio. *B. pin.* 400.

L

Anglis,

Anglis, common Sparagus.

Habitat in pratis et rupibus maritimis. Ad promontorium dictum Lizard Point invenimus; in the marshes near Bristol, below Looks Folly; near Gravesend; R. syn. in insula Portlandica. 4. VII. VIII.

Obs. variat caule erecto et prostrato, floribus masculis, feminis et hermaphrodites; Baccis monospermis et dispersis.

CONVALLARIA. 425.

Cor. sexfida. Bacca maculosa, trilocularis.

- majalis.* 1. CONVALLARIA scapo nudo. *Sp. pl.* 451.
 Polygonitum scapo diphylo, floribus spicatis, nutantibus, campaniformibus. *Hall. hist.* 1241.
 Lilium convallium. *Cam. epit.* 618. *Ger. em.* 410.
R. syn. 264. album. *B. pin.* 304.
 Lilium convallium flore albo. *Park. par.* 349.
 β Lilium convallium angustifolium. *R. syn.* 264.
Anglis, May-Lily-convally or May-lily.
Habitat in sylvis et ericetis frequens. prope Hampstead.
 4. V.
Pharmac, Lili convallii Flores.
- polygonatum.* 2. CONVALLARIA foliis alternis amplexicaulibus, caule ancipiti, pedunculis axillaribus subunifloris.
Sp. pl. 451. *Phil. bot.* 218. *Fl. dan.* 377.
 Polygonatum caule simplici, anguloso, cernuo, foliis ovato-lanceolatis, rigidis, alis unifloris. *Hall. hist.* 1242,
 Polygonatum latifolium II. Clusii. *Ger. em.* 904.
 Polygonatum latifolium flore majore odoro. *B. pin.* 303.
 Polygonatum floribus ex singularibus pediculis. *B. hist.* III. 529. *R. syn.* 263.
 Polygonatum majus flore majore. *Park.* 696.
 β Polygonatum Hellebori albi folio, caule purpurascente.
R. syn. 263.
Anglis, sweet Lily-convally or Solomon's Seal.
Habitat in fissuris rupium prope Settle, Skipton, &c. in comitatu Eboracensi. 4. V. VI.

3. CON-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Habitat in fluviis fundo cœnosis. In the river Yare near Norwich, about Hedley in Surry, and in Cheshire. R. syn. in ericeto Hounsejano; prope Harefield. 4. V. VI.

Pharmac. Calami aromatici Radix.

J U N C U S. 437.

Cal. hexaphyllus. Cor. o. Caps. unilocularis.

* Culmis nudis.

- acutus.* 1. JUNCUS culmo subnudo tereti mucronato, panicula terminali, involucrio diphylo spinoso. *Sp. pl.* 463.
 Juncus acutus capitulis Sorghi. *B. pin.* 10. *R. syn.* 431.
 Juncus maritimus capitulis Sorghi. *Park.* 1192.
 β Juncus acutus maritimus anglicus. *R. hist.* 1303.
Hist. Ox. III. 232. t. 10. f. 15.
Anglis, sea Rush.
 Habitat in paludosis maritimis. 4. VII. VIII.
- conglomeratus.* 2. JUNCUS culmo nudo stricto, capitulo laterali. *Sp. pl.* 464. *Fl. suec.* 298.
 Juncus caule nudo, foliis teretibus mollibus, panicula conglomerata. *Hall. hist.* 1312.
 Juncus lævis vulgaris panicula compactiore. *R. syn.* 432.
 Juncus lævis panicula non sparsa. *B. pin.* 12.
 Juncus lævis glomerato flore. *Lob. ic.* 84. *Park.* 1192.
Anglis, common Rush.
 Habitat in pratis et ericetis humidiusculis. 4. VII. VIII.
- effusus.* 3. JUNCUS culmo nudo stricto: panicula laterali. *Sp. pl.* 464. *Fl.* 299.
 Juncus caule nudo, foliis teretibus striatis, panicula laterali sparsa. *Hall. hist.* 1311.
 Juncus lævis vulgaris, panicula sparsa nostras. *R. syn.* 432.
 Juncus lævis. *Ger. em.* 39. panicula sparsa major.
B. pin. 12. *Park.* 1191.

β Juncus

- β *Juncus acutus*. *Ger. em.* 39. *R. syn.* 432. vulgaris.
Park. 1193.
Anglis, soft β hard Rush.
Habitat in pratis et pascuis paludosis. 4. VI.—VIII.
4. **JUNCUS** culmo nudo filiformi nutante, panicula *filiformis*.
lateralis. *Sp. pl.* 465. *Gron. virg.* 152.
Juncus caule nudo, foliis teretibus mollibus, panicula
lateralis pauciflora. *Hall. hist.* 1313.
Juncus lævis, panicula sparsa, minor. *B. pin.* 12.
tb. 183. *Sch. agr.* 347. *t.* 7. *f.* 1.
Juncus parvus, calamo supra paniculam longius pro-
ductio. *R. syn.* 432. *Pluk. ph. t.* 40. *f.* 8.
Anglis, least Rush.
Habitat in montibus turfosis. Found by Mr. Newton
not far from Ambleside, in Westmorland. 4. VIII.
5. **JUNCUS** culmo nudo, foliis floribusque tribus ter-*trifidus*.
minalibus. *Fl. lapp.* 119. *Sp. pl.* 465. *Fl. dan. t.*
107. *Schm. Gesn. t.* 4. *f.* 16.
Juncus acumine reflexo trifidus. *B. pin.* 12. *pr.* 22.
tb. 186. *Hall. hist.* 1315.
Anglis, trifid Rush.
Habitat in montibus paludosis, ex Scotia recepi. 4. VI.
6. **JUNCUS** culmo nudo, foliis setaceis, capitulis glo-*squarros-*
meratis aphyllis. *Sp. pl.* 465. *Fl. suec.* 302. *dan.* 430. *sus.*
Juncus rigidus, panicula spicata, floribus cartilagi-
neis. *Hall. hist.* 1317.
Juncus montanus palustris. *R. syn.* 432. *Fl. lapp.* 121.
Gramen junceum maritimum. *Lob. ic.* 18. *Ger. em.*
21. majus. *Park.* 1270.
Gramen junceum, foliis et spica junci. *B. pin.* 5.
Anglis, moss Rush or Goose-corn.
Habitat in ericetis frequens. 4. VI. VII.

•• *Culmis foliosis.*

7. **JUNCUS** foliis nodoso-articulatis, petalis obtusis. *articula-*
Sp. pl. 465. *Fl. suec.* 305. *tus.*
Juncus foliis articulosis, panicula inæquali, capitulis
sessilibus, capsulis subulatis triquetris. *Roy. lugdb.* 43.
L 3
Juncus

HEXANDRIA MONOGYNIA.

Juncus foliis articulatis, compressis, panicula semel ramosa. *Hall. hist.* 1322.

Juncus foliis articulosis, floribus umbellatis. *Tourn.* 247. *R. syn.* 433.

Gramen aquaticum. *Tab. ic.* 214. *Ger. em.* 22.

Gramen junceum folio articulato aquaticum. *B. pin.* 5.

Gramen junceum aquaticum. *Park.* 1270.

β Juncus foliis articulatis, teretibus, panicula repetito ramosa. *Hall. hist.* 1323.

Juncus nemorosus folio articuloso. *R. syn.* 433.

γ Juncus caule brachiato, foliis setaceis, floribus fasciculatis, ad ramas sessilibus. *Hall. hist.* 1320.

Gramen junceum minimum, capsulis triangularibus.

R. syn. 434. *Hist. Ox. III.* 227. t. 9. f. 3.

Graminis juncei varietas minor. *Ger. em.* 4.

Anglis, jointed Rush.

Habitat in pratis et pascuis humidis β in sylvis γ in ericetis arenosis. 4. VII. VIII.

bulbosus. 8. JUNCUS foliis linearibus canaliculatis, capsulis obtusis. *Sp. pl.* 466. *Mees. Fl. fris.* 160. *dan.* 431.

Juncus compressus, foliis linearibus canaliculato-concavis, capsulis ovatis, culmo compresso. *Jac. enum.* 235.

Juncus foliis carinatis, panicula sparsa, fructo globoso. *Hall. hist.* 1318.

Juncus parvus cum pericarpis rotundis. *B. hist.* II. 522. *R. syn.* 433.

Anglis, bulbose Rush.

Habitat in pratis et ericetis humidis. 4. VIII.

bufonius. 9. JUNCUS culmo dichotomo, foliis angulatis, floribus solitariis sessilibus. *Sp. pl.* 466. *Fl. suec.* 305.

Juncus caule brachiato, foliis setaceis, floribus solitariis ad ramum sessilibus. *Hall. hist.* 1319.

Juncus palustris humilior erectior. *Tourn.* 246. *R. syn.* 434.

Gramen junceum. *Ger. em.* 4. parvum s. *Holosteum Matthioli* et *Gramen bufonium Flandrorum.* *Park.* 1190.

Gramen nemorosum, calyculis paleaceis. *B. pin.* 7.

β Gramen

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

HEXANDRIA TRIGYNIA.

Gramen nemorosum hirsutum latifolium maximum.

R. syn. 416. *Sch. agr.* 317.

Gramen Luxulæ maximum. *B. hist.* II. 493.

Anglis, wood Rush.

Habitat in sylvis et sepibus; prope Hampstead. 4. V.

campes-
tris.

13. JUNCUS foliis planis pilosis, spicis sessilibus pedunculatisque. *Sp. pl.* 468. *Fl. suec.* 309.

Juncus foliis gramineis hirsutis, spicis petiolatis nutantibus, petalis aristatis. *Hall. nom. et hist.* 1330.

Gramen hirsutum capitulis psyllii. *B. pin.* 7. pr. 15.

Gramen exile hirsutum. *Ger. em.* 17.

Gramen nemorosum hirsutum minus angustifolium. *Park.* 1185.

β Gramen hirsutum elatius, panicula juncea compacta.

R. syn. 416.

Anglis, field Rush.

Habitat in pratis, pascuis et ericetis, β in paludibus turfosis frequens. 4. IV. V.

T R I G Y N I A.

* Flores inferi.

T R I G L O C H I N. 453.

Cal. triphyllus. *Petala* tria calyciformia. *Stylus* 0.
Caps. basi dehiscens.

palustre. 1. TRIGLOCHIN capsulis trilocularibus sublinearibus

Sp. pl. 482. *Fl. suec.* 321. *dan.* 490.

Juncajo palustris et vulgaris. *Tourn.* 266. *R. syn.* 435.

Gramen junceum spicatum seu Triglochm. *B. pin.* 6.

Gramen marinum spicatum. *Ger. em.* 20. alterum.

Park. 1279.

Anglis, marsh Triglochm or Arrow-headed Grass.

Habitat in pratis et pascuis udis. 4. VII. VIII.

mariti-
mum.

2. TRIGLOCHIN capsulis sexlocularibus ovatis. *Sp.*

pl. 483. *Fl. suec.* 322. *dan.* 306.

Gramen marinum spicatum. *Lob. ic.* 16. *Park.* 1278.

R. syn. 435.

Gramen

Gramen junceum et spicatum alterum. *B. pin.* 6.
Anglis, sea Triglochin, or spiked Grass.
 Habitat in pratis maritimis. *Ꝟ.* V—VIII.

RUMEX. 451.

Cal. triphyllus. *Petala* tria, conniventia. *Sem.* unicum triquetrum.

• *Hermaphroditi* : valvulis granulo notatis.

1. RUMEX floribus hermaphroditis : valvulis integer-sanguin-
 rimis : unica granifera, foliis cordato-lanceolatis. *us.*

Sp. pl. 476. *Roy. lugdb.* 229.

Lapathum folio acuto rubente. *B. pin.* 115. *R. syn.*
 142.

Lapathum sativum sanguineum. *Ger. em.* 390.

Anglis, blood Dock.

Habitat in sylvis, sed rarius, prope Hampstead, et prope
 Maidston in Cantio. *Ꝟ.* VII.

Pharmac. *Lapathi sanguinei Folio, Semina.*

2. RUMEX floribus hermaphroditis : valvulis integris *crispus.*
 graniferis, foliis lanceolatis undulatis acutis. *Sp. pl.*
 476.

Lapathum foliis crispis, imis ovatis, supremis lanceo-
latis, calycibus verrucosis. *Hall. hist.* 1589.

Lapathum folio acuto crispo. *B. pin.* 115. *R. syn.*
 141.

Lapathum acutum minus. *Park.* 1226.

Lapathi acuti varietas folio crispo. *Ger. em.* 387.

Anglis, curled Dock.

Habitat in pratis, pascuis et ad vias. *Ꝟ.* VI. VII.

3. RUMEX floribus hermaphroditis : valvulis dentatis : *pulcher.*
 subunica granifera, foliis radicalibus panduriformi-
 bus. *Sp. pl.* 477.

Lapathum foliis radicalibus utrinque emarginatis, ca-
lycibus reticulatis, ciliatis, verrucosis. *Hall. hist.*
 1593.

Lapathum pulchrum bononiense sinuatum. *B. hist.*
II. 988. *R. syn.* 142.

Anglis,

Anglis, fiddle Dock.

Habitat in ruderatis, pratis siccis et ad vias. 4. VII.
VIII.

paludosus. 4. RUMEX floribus hermaphroditis : valvulis integris graniferis, foliis ovato-lanceolatis acutiusculis. Lapathum petiolis latescentibus, foliis longe lanceolatis floribus, verticillatis verrucosis. *Hall. hist. 1590. ?*

Lapathum aquaticum minus. *B. hist. II. 987 ?*

Lapathum acutum angustifolium non crispum. *R. syn. 141.*

Anglis, marsh Dock.

Habitat in palustribus et ad ripas fluviorum. 4. VIII.

Desc. Radix perennis ramosa. Caulis erectus quadripedalis striatus, ramosus ; ramis alternis erectis. Folia radicalia, ovata, obtusiuscula, subcrenulata. Caulina alterna, petiolata, ovato-lanceolata, subcrenulata, acuti-
uscula, plana, nuda, petiolis brevibus. Stipula vaginans, membranacea. Flores verticillati, pedunculati, verticillis fere distinctis. Valvula exterior florum granum magnum rubrum habet, reliquæ granum minimum habent.

Hydrolapathum. 5. RUMEX floribus hermaphroditis : valvulis integris graniferis, foliis lanceolatis acutis

Lapathum foliis longe lanceolatis, paniculis densissimis. *Hall. hist. 1588.*

Lapathum maximum aquaticum s. Hydrolapathum. *B. hist. II. 986. R. syn. 140.*

Lapathum aquaticum folio cubitali. *B. pin. 116.*

Lapathum longifolium nigrum palustre, s. Britannica antiquorum vera, vel Hydrolapathum nigrum. *Munt. brit. t. 1.*

Hydrolapathum magnum. *Ger. em. 389.*

Hydrolapathum majus. *Park. 1225.*

Anglis, water Dock.

Habitat in palustribus, et ad ripas fluviorum. 4. VII.
VIII.

Pharmac. Hydrolapathi Radix.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

••• *Hermaphroditi* : *valvulis granulo destitutis.*

digynus.

9. RUMEX floribus hermaphroditis digynis. *Sp. pl.* 480. *Fl. suec.* 317. *Dan.* 14.

Lapathum foliis antice emarginatis. *Hall. hist.* 1595.

Acetosa cambrobritannica montana. *Park.* 745.

Acetosa rotundifolia repens eboracensis, folio in medio deliquium patiente. *Hist. Ox. II.* 583. f. 5. t. 36. f. penult.

Acetosa repens Westmorlandica, cochleariæ foliis apicibus nonnihil sinuatis. *Pluk. ph. t.* 252. f. 2.

Anglis, mountain Dock, or Sorrel.

Habitat in montibus Wallicis, Eboracensibus, Westmorlandicis & Cumberlandicis, frequens. 4. VI. VII.

••• *Floribus dioicis.*

Acetosa. 10. RUMEX floribus dioicis, foliis oblongis sagittatis. *Sp. pl.* 481.

Lapathum sexubus separatis, foliis sagittatis, hamis retrorsum porrectis. *Hall. hist.* 1597.

Lapathum acetosum vulgare. *R. syn.* 143.

• *Acetosa pratensis.* *B. pin.* 114.

Acetosa vulgaris. *Park.* 742.

Oxalis seu *Acetosa.* *Ger. em.* 396.

β *Acetosa montana maxima.* *B. pin.* 114. *R. syn.* 143.

Anglis, common Dock, or Sorrel.

Habitat in pratis et pascuis. β in sylvis montosis humidiusculis passim. 4. VI.

Pharmac. *Acetosæ pratensis.* *Radix, Folia, Semina.*

Acetosella. 11. RUMEX floribus dioicis, foliis lanceolato-hastatis. *Sp. pl.* 481. *Fl. lapp.* 131. *suec.* 319.

Lapathum sexubus separatis, foliis sagittatis, hamis acutis, recurvis. *Hall. hist.* 1596.

Lapathum acetosum repens lanceolatum. *R. syn.* 143.

Acetosa arvensis lanceolata. *B. pin.* 114.

Acetosa minor lanceolata. *Park.* 744.

Oxalis tenuifolia. *Ger. em.* 397.

Anglis, sheep's Dock or Sorrel.

Habitat in pratis et pascuis arenosis. 4. V—VII.

Pharmac. *Acetosæ lanceolatæ Folia.*

TOFIELDIA.

T O F I E L D I A.

CAL. nullus.

COR. Petala sex, æqualia, oblonga, obtusa, concava, persistencia.

STAM. Filamenta sex subulata, glabra, longitudine petalorum *Antheræ* parve, subrotundæ, incumbentes.

PIST. Germen trigonum. *Styli* tres subulati, patentes. *Stigmata* obtusa.

PER. Capsula subrotunda, subtrigona, trilocularis, sexvalvis.

SEM. numerosa, oblonga, subtrigona, parva.

I. T O F I E L D I A.

Anthericum, *calyculatum*, foliis ensiformibus, perianthis trilobis, filamentis glabris. *Sp. pl.* 447. *Fl. dan.* 36. *palustris.*

Anthericum filamentis glabris. *Hall. hist.* 1205.

Scheuchzeria Pseudo-asphodelus. *Scop. Carn.* 263.

Phalangium scoticum palustre minimum iridis folio.

Tourn. 369. *R. syn.* 375.

Phalangium alpinum palustre iridis folio. *Segu. ver.*

II. 61. t. 14.

Pseudo-asphodelus alpinum. *B. pin.* 29.

Anglis, marsh Tofieldia, or Asphodel.

Habitat in palustribus montosis. Juxta rivulum non procul Bervico in Scotia. *R. syn.* on Glenmore, Ross-shire and Ben Grihum, Sutherland. *D. Hope.* 4.

VII—IX.

C O L C H I C U M. 457.

Spatha. *Cor.* sexpartita; tubo radicato. *Caps.* tres connexæ, inflatæ.

I. COLCHICUM foliis planis lanceolatis erectis. *Sp. autumnale.*
pl. 485. *Bœhm. lips.* 29.

Colchicum flore folia longe præcedente, petalis ovatis.

Hall. hist. 1255.

Colchicum commune. *B. pin.* 67. *R. syn.* 373.

Colchicum. *Dod. purg.* 371. anglicum purpureum et album. *Ger. em.* 157. *Park. par.* 153.

Anglis, autumnal Colchicum, or meadow-Saffron.

Habitat in pratis tam planis quam montosis, solo pingui in Occidentalibus

Occidentalibus ut et Septentrionalibus Angliæ. R. syn. 4. VIII. IX.

P O L Y G Y N I A.

A L I S M A. 460.

Cal. triphyllus. Petala tria. Sem. plura.

Plantago. Δ. 1. ALISMA foliis ovatis acutis, fructibus obtuse trigonis. Sp. pl. 468. Fl. dan. 561.

Damasonium foliis ellipticis, lanceolatis, capitulo rotunde triquetro. Hall. hist. 1184.

Plantago aquatica. Cam. epit. 264. R. syn. 257. major. Ger. em. 417. Park. 1245.

Plantago aquatica latifolia. B. pin. 190.

β Plantago aquatica longifolia. Pet. h. br. t. 43. l. 7.

γ Plantago aquatica, λεπτομακρόφυλλος. Fl. prus. I. 99. ic. 66. R. syn. 258.

Anglis, great Water-plantain.

Habitat in aquosis et ad ripas fluviorum. 4. VII.

ranuncu- 2. ALISMA foliis lineari-lanceolatis, fructibus globoso-
loides. squarrosis. Sp. pl. 487. Fl. dan. 122.

Plantago aquatica minor. Park. 1245. R. syn. 257.

Plantago aquatica humilis. Ger. em. 417.

Plantago aquatica angustifolia. B. pin. 190.

β Plantago aquatica minor, ranunculi flammei majoris folio. Pluk. Amalth. 175. R. syn. 258.

Anglis, lesser Water-plantain.

Habitat in aquosis et in paludibus turfosis. 4. VI—IX.

natans. 3. ALSIMA foliis ovatis obtusis, pedunculis solitariis. Sp. pl. 487. Fl. succ. 324.

Alisma repens, foliis gramineis et subrotundis. Mart. Tourn. 17.

Damasonium repens, potamogetonis rotundifolii folio. Vaill. Act. paris. 1719. 29. t. 4. f. 8.

Anglis, creeping Water-plantain.

Habitat in lacubus et jossis majoribus. In the Great Lake, below the old Castle at Llanberys. Mr. Brewer. Mart. Tourn. 4. VII. VIII.

Obi. Folia submersa linearia ; natantia ovata.

4. ALISMA

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Classis VII.

H E P T A N D R I A

M O N O G Y N I A.

T R I E N T A L I S. 461.

Cal. heptaphyllus. *Cor.* septem-partita, æqualis, plana. *Bacca.* exsucca.

- europæa.* 1. TRIENTALIS foliis lanceolatis integerrimis. *Sp. pl.* 488. *Fl. suec.* 326. *dan.* 84.
 Alsinanthemos. *Thal. sylv.* 15. *Moerb. hort.* 8. III.
R. syn. 286.
 Pyrola alfine flore europæa. *B. pin.* 191. *Park.* 509.
 Pyrola alfine flore major. *B. pr.* 100.
 Herba trientalis. *B. hist.* III. 506.
 Alfine alpina. *Swenk. files* 17. *descr.*
 β Pyrola alfine flore Brasiliana. *B. pin.* 191. *pr.* 100.
R. syn. 286.
Anglis, common Trientalis, or Chickweed-Wintergreen.
 Habitat in sylvis et ericetis turfosis in comitatibus Eboracensi, Northumberlandico, Cumberlandico et Westmorlandico; prope Bingley in comitatu Eboracensi copiose. 4. VI. VII.

Classis

*Classis VIII.*O C T A N D R I A
M O N O G Y N I A,* *Flores completi: superi.*

EPILOBIUM. 471.

Cal. quadrifidus. *Petala* quatuor. *Caps.* oblonga. *Sem.* papposa.* *Staminibus declinatis.*1. EPILOBIUM foliis sparsis lineari-lanceolatis, flori- *angustifolium.*
bus inæqualibus. *Sp. pl.* 493. *Fl. dan.* 289.*Epilobium* flore difformi, foliis lanceolatis transversim
nervosis. *Hall. hist.* 1000.*Lysimachia Chamænerion* dicta *angustifolia.* *B. pin.*
245.*Lysimachia speciosa*, quibusdam *Onagra* dicta, filiquo-
sa. *B. hist.* II. 906: *R. syn.* 310.*Chamænerion* flore *delphinii.* *Park.* 548.*Anglis*, narrow-leaved *Epilobium*, or *Rosebay Wil-*
*low-herb.**Habitat in sylvis et sepibus. In the meadows near Shef-*
field, in Yorkshire, and in several other places of the
north, plentifully. R. syn. On Maize-hill, beyond
*Greenwich. 4. VII. VIII.*** *Staminibus erectis regularibus, petalis bifidis.*2. EPILOBIUM foliis lanceolatis serratis subdecurren- *hirsutum.*
tibus; inferioribus oppositis, caule subsimplici. *Fl.*
angl. 140.*Chamænerion* foliis oppositis lanceolatis serratis sessili-
bus cauleque villosis. *Schreb. Spic. Fl. Lips.* 146.

M

Lysimachia

Lyfimachia filiquosa hirsuta, parvo flore. B. pin. 245.

R. syn. 311. Hist. Ox. II. 270. t. 10. f. 4.

Lyfimachia sylvatica. Ger. em. 476.

Lyfimachia filiquosa sylvestris hirsuta. Park. 549.

Anglis, hairy Epilobium, or Willow-herb.

Habitat in aquosis et ad fossas. 4. VII.

Obs. Caulis villosus, simplex, rarissime ramosus nisi injuriam recepit; Folia villosa, Flores parvi purpurei.

ramosum. 3. **EPILOBIUM** foliis lanceolatis ferratis decurrentibus inferioribus oppositis, caule ramoso. *Fl. angl. 141.*

Epilobium foliis oppositis lanceolatis ferratis decurrenti-amplexicaulibus. Sp. pl. 494. Fl. dan. 326.

Chamænerion foliis oppositis alternisque lanceolatis glabris ferratis basi decurrentibus. Schreb. Spic. Fl. lips. 147.

Lyfimachia filiquosa hirsuta magno flore. B. pin. 245.

R. syn. 311.

Lyfimachia filiquosa. Ger. em. 476.

Anglis, branched Epilobium, or Codlings and Cream.

Habitat in sepibus humidis et ad ripas fluviorum. 4. VII.

Obs. Caules ramosi foliaque pubescentes; Flores variant colore carneo et purpureo.

montanum. 4. **EPILOBIUM** foliis oppositis ovatis dentatis. *Sp. pl. 494.*

Epilobium foliis ovato-lanceolatis, glabris, dentatis. Hall. hist. 996.

Lyfimachia campestris. Ger. em. 478.

Lyfimachia filiquosa glabra major. B. pin. 245.

Lyfimachia filiquosa major. Park. 548.

Anglis, mountain Epilobium, or Willow-herb.

Habitat in pratis humidis glareosis. 4. VII.

tetragonum. 5. **EPILOBIUM** foliis lanceolatis denticulatis: imis oppositis, caule tetragono. *Sp. pl. 494.*

Epilobium foliis lanceolatis, glabris dentatis. Hall. hist. 998.

Lyfimachia filiquosa glabra media f. minor. Ger. em. 479. R. syn. 311.

Lyfimachia filiquosa glabra minor. B. pin. 245.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Vitis idæa angulosa. B. hist. I. 520. R. syn. 457.
Anglis, common Bil, Whortle, or Blea-berry.
Habitat in ericetis et sylvis frequens. h. IV.

uliginosum.

2. **VACCINIUM** pedunculis unifloris, foliis integerrimis ovalibus obtusis lævibus. *Syst. Nat. 267. Sp. pl. 498. Fl. dan. 231.*

Vaccinium pedunculis unifloris, foliis integerrimis ovatis venosis. Hall. hist. 1021.

Vaccinia nigra fructu majore. Park. 1455.

Vitis idæa foliis subrotundis major. Ger. em. 1416.

Vitis idæa foliis subrotundis exalbidis. B. pin. 470.

Vitis idæa magna quibusdam, s. Myrtillis grandis. B. hist. I. 518. R. syn. 457.

Anglis, marsh Bil-berry.

Habitat in ericetis et sylvis humidiusculis; in montosis Cumberlandiæ medio itinere inter Hexham et Pereth oppida; sed copiosius prope vicum Gambleby, ad sextum ab oppido Pereth lapidem, via quæ Novum Castellum ducit: utrobique in pascuis palustribus. R. syn. In Whinfield Forest, Westmorland: in Septentrionali parte Scotiæ frequens. h. IV.

*** Foliis semper virentibus.*

Vitis idæa.

3. **VACCINIUM** racemis terminalibus nutantibus, foliis obovatis revolutis integerrimis subtus punctatis. *Sp. pl. 500. Fl. dan. 40.*

Vaccinium foliis perennantibus ovatis, subtus punctatis, oris contractis. Hall. hist. 1022.

Vaccinia rubra. Dod. 770. Ger. em. 1415. buxeis foliis. Park. 1458.

Vitis idæa foliis subrotundis non crenatis, baccis rubris. B. pin. 470.

Vitis idæa sempervirens fructu rubro. B. hist. I. 522. R. syn. 457.

Anglis, red Bil-berry, or Whorts.

Habitat in ericetis paludosis et montosis in boreali parte Angliæ et in Scotia frequens. h. IV. V.

Oxycoccis. 4. **VACCINIUM** foliis integerrimis revolutis ovatis, caulibus repentibus filiformibus nudis. *Sp. pl. 500. Fl. suec. 335. dan. 80.*

Vaccinium

Vaccinium caule prostrato, foliis ovato-lanceolatis acutis, flore quadripartito. Hall. hist. 1023.

Vaccinium palustre. Park. 1229.

Vaccinia palustris. Dod. 770. Lob. ic. 109. Ger. em. 1419.

Vitis idæa palustris. B. pin. 471.

Oxycoccus s. Vaccinia palustris. B. hist. I. 525. R. syn. 267.

Anglis, creeping Bil-berry, or Cran-berries.

Habitat in palustribus turfosis in septentrionali parte Angliæ copiose. prope Glastenbury. h. V. VI.

E R I C A. 484.

Cal. tetraphyllus. Cor. quadrifida. Filamenta receptaculo inserta. Antheræ bifidæ. Caps. quadrilocularis.

* *Antheris aristatis.*

1. ERICA antheris aristatis inclusis, stylo exserto, corollis inæqualibus campanulatis mediocribus, foliis oppositis sagittatis. *Lin. diss. de Erica I.*

Erica antheris bicornibus inclusis, corollis inæqualibus campanulatis mediocribus, foliis oppositis sagittatis. Sp. pl. 501.

Erica foliis imis adpressis simplicibus, floralibus calcatis. Hall. hist. 1012.

Erica vulgaris. Park. 1480. R. syn. 470. s. pumila. Ger. em. 1380.

Erica vulgaris glabra. B. pin. 485.

β *Erica vulgaris hirsuta. Ger. em. 1382. R. syn. 471. Anglis, common Heath.*

Habitat in ericetis, et sylvis. h. VI—VIII.

2. ERICA antheris cristatis styloque inclusis, corollis cinereis ovatis, foliis ternis, floribus glomeratis. *Mant. II. 372. Fl. dan. 28.*

Erica antheris bicornibus inclusis, corollis ovatis racemosis, foliis ternis glabris linearibus. Sp. pl. 501.

Erica tenuifolia. Ger. em. 1380. R. syn. 471.

Erica humilis, cortice cinereo, arbuti flore. B. pin. 486.

Erica virgata f. VI. Clusii. *Park.* 1483.

Anglis, fine-leaved Heath.

Habitat in ericetis et nemorosis siccioribus. ½. VI—VIII.

Tetralix. 3. *ERICA* antheris aristatis, stylo incluso, corolla ovata, foliis quaternis, floribus capitatis. *Mont.* II. 372.

Erica antheris bicornibus inclusis, corollis subglobosis confertis folio longioribus, foliis quaternis ciliatis patentibus. *Sp. pl.* 502. *dan.* 81.

Erica major flore purpureo. *Ger. em.* 1381.

Erica ex rubro nigricans scoparia. *B. pin.* 486.

Erica brabantica foliis coridis hirsuto quaterno. *B. hist.* I. 358. *R. syn.* 471.

Anglis, cross-leaved Heath.

Habitat in ericetis humidis. ½. VI—VIII.

** *Antheris muticis obtusis emarginatis.*

multiflora. 4. *ERICA* antheris muticis styloque exsertis, corolla cylindrica, foliis quinis, floribus sparsis. *Mant.* II. 372.

Erica antheris simplicibus bifidis exsertis, corollis cylindricis, foliis quinis patentibus. *Sp. pl.* 503.

Erica foliis corios multiflora. *B. hist.* I. 356. *R. syn.* 471.

Erica maxima purpurascens longioribus foliis. *B. pin.* 485.

Erica juniperifolia narbonensis dense fruticans. *Lob. hist.* 620. *Garid. aix.* 160. t. 32. *Sauv. monsp.* 46.

Anglis, many-flowered Heath.

Habitat in ericetis. On Goon hilly Downs, going from Helston to the Lizard Point in Cornwall. *R. syn.* ½. VI—VIII.

daboecii. 5. *ERICA* antheris muticis styloque inclusis, corolla ovata, foliis alternis lanceolatis revolutis, floribus racemosis secundis.

Erica racemo terminali, foliis alternis subtus tomentosis. *Sp. pl.* 509.

Andromeda racemis secundis, floribus quadrifidis ovatis, foliis alternis lanceolatis revolutis. *Syst. Nat.* 300.

Erica

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

DIGYNIA.

CHLORA. *Lin. Mant.* 10.*Cal.* octophyllus. *Cor.* monopetala, octofida. *Caps.* unilocularis, bivalvis, polysperma.*perfoliata.* 1. CHLORA foliis perfoliatis. *Syst. Nat.* 267.Chlora. *Reuealm. Spec.* 80. t. 76.Blackstonia perfoliata. *Fl. angl.* 146.Gentiana corollis octofidis, foliis perfoliatis. *Sp. pl.* 335.Centaureum luteum perfoliatum. *B. pin.* 278. *R. syn.* 287.Centaureum parvum luteum Lobelli. *Ger. em.* 547.Centaureum minus luteum et perfoliatum non ramosum, *Park.* 271.*Anglis,* perfoliate Yellow-wort.

Habitat in pratis et pascuis montosis, solo calcareo. ☉. VII.

TRIGYNIA.

POLYGONUM. 495.

Cal. o. *Cor.* quinquepartita, calycina. *Sem.* unicum angulatum.

** Bistortæ spica unica.

Bistorta. 1. POLYGONUM caule simplicissimo monostachio, foliis ovatis in petiolum decurrentibus. *Sp. pl.* 516.*Fl. dan.* 421.Polygonum radice lignosa intorta, spica ovata, foliosis petiolis alatis. *Hall. hist.* 1559.Bistorta major. *Ger. em.* 399. *R. syn.* 147. vulgaris. *Park.* 391.Bistorta radice minus intorta. *B. pin.* 192.*Anglis,* Bistort Knot-grass, or Snakeweed.

Habitat in pratis humidiusculis in comitatibus Cumberlandico, Westmorlandico, Eboracensi etc. frequens; prope Battersea. 4. V. VI.

Pharmac. Bistortæ Radix.

2. POLYGONUM.

2. POLYGONUM caule simplicissimo monostachio, vivipar-
foliis lanceolatis. *Sp. pl.* 516. *Fl. suec.* 340. *Dan. rum.*
13.

Polygonum radice lignosa intorta, foliis nervosis, radi-
calibus ovata-lanceolatis, caulinis linearibus. *Hall.*
hist. 1558.

Bistorta minor. *Ger. em.* 369. *R. syn.* 147. nostras.
Park. 392.

Bistorta alpina minor. *B. pin.* 192.

- β Bistorta minima alpina, foliis imis subrotundis et mi-
nutissime serratis. *R. syn.* 147.

Anglis, viviparus Knot-grass, or Bistort.

Habitat in pratis montosis; in several places in West-
moreland, as at Crosby Ravensworth, and near Set-
tle in Yorkshire. β in pascuis clivosis excelsæ rupis y
Crib-gôch dictæ supra lacum ffynnon frêch prope
Llanberris. *R. syn.* 4. VI.

** *Persicariæ* pistillo bifido, aut stamina minus 8.

3. POLYGONUM floribus pentandris semidigynis, *amphibi-*
spica ovata. *Syst. Nat.* 275. staminibus corolla lon- *um.*
gioribus. *Sp. pl.* 517. *Fl. suec.* 341. *Dan.* 282.

- Polygonum foliis ovato-lanceolatis ciliatis, spicis o-
vatis. *Hall. hist.* 1565.

Persicaria salicis folio perennis, *Potamogiton* angusti-
folium dicta. *R. syn.* 145.

Potamogiton angustifolium. *Ger. em.* 821.

Potamogiton salicis folio. *B. pin.* 131.

Fontalis major longifolia. *Park.* 1254.

Anglis, amphibius Knot-grass, or *Persicaria*.

Habitat in fossis, paludibus et lacubus. 4. VI—VIII.

Obs. varietas aquatica staminibus flore brevioribus, ter-
restribus longioribus. *Jacq. enum.*

4. POLYGONUM floribus hexandris digynis, spicis *Persica-*
ovato-oblongis, foliis lanceolatis, stipulis ciliatis. *ria.*
Sp. pl. 518.

Polygonum foliis ovato-lanceolatis, subhirsutis, spicis
ovatis, vaginis ciliatis. *Hall. hist.* 1557.

Persicaria maculosa. *Ger. em.* 821. *R. syn.* 145.

Persicaria. *Dodon. pempt.* 108. nitida maculosa et non
maculosa. *B. pin.* 191.

Persicaria

- Perficaria vulgaris mitis seu maculosa.* *Park.* 856.
- β *Perficaria angustifolia.* *B. pin.* 101. *pr.* 43. ex singulis geniculis florens. *M. P.* 90. *R. syn.* 145.
- γ *Perficaria folio subtus incano.* *Tourn.* 510. *R. syn.* 145.
- δ *Perficaria maculosa procumbens foliis subtus incanis.* *R. syn.* 146.
- *Perficaria foliis salicis albæ vulgaris.* *Budell. hort. sic.* *W.* 20. *R. syn.* 145. *argentea.* *Pet. ber. br. t.* 3. *f.* 8.
- pensylva- ζ *Polygonum floribus octandris digynis, pedunculis bif-*
nicum. *pidis, foliis lanceolatis, stipulis muticis.* *Sp. pl.* 518.
- Perficaria mitis major, foliis pallidioribus.* *R. syn.* 145.
- *Perficaria latifolia geniculata, caulibus maculatis.* *R. syn.* 145.
- Anglis, spotted Knot-grass, Perficaria or Arsmart.*
Habitat in pratis, pascuis, arvis, fimetis et ruderatis humidiusculis ubique. γ. δ. i. in arenosis et glareosis aridis. ©. VII.—IX.
- Obs. ζ. Caulis supra geniculos tumidus; Stipulæ solitariae intrasolitariae, persistentes, vaginante, adnatae, truncatae, ciliatae, sed frequentissime evenit; ut in medio abruptæ, et tum mutice apparent; numerus staminum variat a sex ad octo.*
- hydropi- 5. POLYGONUM floribus hexandris semidigynis, fo-
per. *liis lanceolatis, stipulis submuticis.* *Sp. pl.* 517. *Fl. suec.* 343.
- Polygonum foliis ovato-lanceolatis, vaginis calvis truncatis.* *Hall. hist.* 1554.
- Perficaria urens seu Hydropiper.* *B. pin.* 101.
- Perficaria vulgaris acris s. Hydropiper.* *B. hist.* III. 780. *R. syn.* 144.
- Perficaria vulgaris acris seu minor.* *Park.* 856.
- Hydropiper.* *Ger. em.* 445.
- β *Polygonum floribus hexandris digynis, foliis lanceo-*
latis, stipulis ciliatis, caule divaricato patulo. *Fl. angl.* 148.
- Polygonum foliis ovato lanceolatis glabris, spicis strigosis, vaginis ciliatis.* *Hall. hist.* 1555.
- Perficaria pusilla repens.* *Ger. em.* 446. *R. syn.* 145.
Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

*fagopy-
rum.*

- 8. POLYGONUM foliis cordato-sagittatis, caule erectiusculo inermi; seminum angulis æqualibus. *Sp. pl.* 522.

Polygonum caule erecto, foliis sagittatis, valvulis integerrimis. *Hall. hist.* 1563.

Fagopyrum. *Dod. pempt.* 511. *cer.* 80. *R. syn.* 144.

Fagopyrum. *Ger. em.* 89. *Park.* 1141.

Erysimum Theophrasti folio hederaceo. *B. pin.* 27.

Anglis, Buck-wheat Knot-grass.

Habitat in arvis cultis passim. ☉. VII. VIII.

TETRAGYNIA.

A D O X A. 501.

Cal. bifidus, inferus. *Cor.* quadrifida seu quinquefida, supera. *Caps.* quadrilocularis s. quinquelocularis, calyce coalita.

*moschatel-
lina.* 1. ADOXA. *Sp. pl.* 527. *Fl. dan.* 94. *Hall. hist.* 1005.

Moschatellina foliis fumarizæ bulbosæ. *B. hist.* III. 206. *R. syn.* 267.

Ranunculus nemorosus Moschatella dictus. *Park.* 226.

Ranunculus nemorum Moschatellina dictus. *B. pin.* 178.

Radix cava minima viridi flore. *Ger. em.* 1091.

Anglis, tuberous Moschatel.

Habitat in sylvis et umbrosis. ♃. IV. V.

P A R I S. 500.

Cal. tetraphyllus. *Petala* quatuor angustiora. *Bacca* quadrilocularis.

*quadrifo-
lia.* 1. PARIS. *Sp. pl.* 526. *Fl. dan.* 139. *Hall. hist.* 1006.

Herba paris. *Matth.* 1193. *Ger. em.* 405. *Park.* 390.

B. hist. III. 316. *Till. ab.* 33. *ic.* 32. *R. syn.* 264.

Solanum quadrifolium bacciferum. *B. pin.* 167.

Anglis, Herb-Paris, True-love or One-berry.

Habitat in sylvis et umbrosis. In sylvula non longe ab

Hampstedio. C. Miller. ♃. V. VI.

Pharmac. Herba Paris *Fructus*, *Radix*.

ELATINE.

ELATINE. 502.

Cal. tetraphyllus. *Petala* quadrilocularis quadrivalvis,
depressa.

1. ELATINE foliis verticillatis. *Sp. pl.* 527. *Fi. succ. alsinastrum.*
342. *Boem. lepf.* 127. *Hall. hist.* 857.

Elatine foliis emerfis linearibus, immerfis capillaceis.

Sauv. monf. 1743. p. 50.

Alsinastrum gallii folio. *Vail. paris.* 6. t. 1. 6.

Alsinastrum gratiolæ folio. *Tourn.* 244. *R. syn.* 346.

Equisetum palustre linariæ scopariæ folio. *B. pin.* 15.

Anglis, Waterwort.

Habitat in fossis et paludibus. Found by Mr. J. She-
rard on the boggy ground, on the common just by the
road from Eltham to Chiffelhurst. R. syn. 24.
VIII.

Classis IX.

E N N E A N D R I A
H E X A G Y N I A.

BUTOMUS. 507.

Cal. 0. Petala sex. Caps. sex, polysperma.

- umbella-*
tus. 1. BUTOMUS. *Cæsalp.* 556. *R. syn.* 70. *Sp. pl.*
532. *Fl. succ.* 350. *dan.* 604.
Juncus floridus. *Cam. epit.* 781. *B. hist.* II. 524.
Park. 1197. *major.* *B. pin.* 12.
Gladiolus palustris *Cordi.* *Ger. em.* 29.
Anglis, Butomus or flowering Rush.
Habitat in fossis limosis et ad ripas fluviorum et stagno-
rum. 4. VI.

Classis

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Pyrola. *Cam. epit.* 723. *Ger. em.* 408. *R. syn.* 363.
rotundifolia major. *B. pin.* 191.

Pyrola nostras vulgaris. *Park.* 508.

Anglis, common Winter-green.

Habitat in sylvis et nemoribus. In septentrionalibus
Angliæ locis reperitur; in meridionalibus rarius oc-
currit; near Halifax plentifully, and in Stoken-
church woods, Oxfordshire. *R. syn.* 4. VI. VII.

minor.

2. PYROLA floribus racemosis dispersis, staminibus
pistillisque rectis. *Sp. pl.* 567. *Fl. dan.* 55.

Pyrola foliis subrotundis, tubis rectis. *Hall. hist.* 1009.

Pyrola minor. *Riv. pent.* 149. *R. syn.* 363.

Pyrola folio minore et duriore. *B. pin.* 191.

Anglis, lesser Winter-green.

Habitat in sylvis; in sylvula prope Clapham in comitatu
Eboracensi. 4. VI. VII.

secunda.

3. PYROLA racemo unilaterali. *Sp. pl.* 567. *Fl. suec.*
362. *dan.* 402. *Hall. hist.* 1008.

Pyrola folio mucronato serrato. *B. pin.* 181. *R. syn.*
363.

Pyrola secunda tenerior. *Clus. hist.* 117. *Ger. em.*
408.

Pyrola tenerior. *Park.* 509.

Anglis, serrated Winter-green.

Habitat in sylvis et nemorosis. In Haslewood, near
Sir Walter Vavasor's park, Yorkshire. *R. syn.*
4. VI.

** Flores monopetali æquales.

ANDROMEDA. 549.

Cal. quinquepartitus. *Cor.* ovata: ore quinquefido.

Caps. quinquelocularis.

po. ifolia.

1. ANDROMEDA pedunculis aggregatis, corollis ova-
tis, foliis alternis lanceolatis revolutis. *Sp. pl.* 564.

Fl. suec. 353. *dan.* 54.

Andromeda foliis alternis, lineari-lanceolatis. *Fl. lapp.*
163. t. 1. f. 3. *Hall. hist.* 1017.

Iedum

Ledum palustre nostras, arbuti flore. *R. syn.* 472.

Rosmarinum sylvestre minus nostras. *Park.* 76.

Erica humilis, rosmarini foliis, unedonis flore, capsula cistoide. *Pluk. ph. t.* 175. *f.* 1.

Anglis, marsh Andromeda or wild Rosemary.

Habitat in paludibus turfosis in comitatibus Eboracensi, Lancastriensi, Westmorlandico, Cumberlandico frequens. *4. VI.*

ARBUTUS. 552.

Cal. quinquepartitus. *Cor.* ovata : ore basi pellucida.

Bacca quinquelocularis.

1. ARBUTUS caule arboreo, foliis glabris ferratis, *unedo.*

baccis polyspermis. *Sp. pl.* 566. *Sauv. monsp.* 56.

Arbutus. *Cam. epit.* 168. *Ger. em.* 1496. *Park.* 1489.

R. syn. 464. folio ferrato. *B. pin.* 460.

Anglis, common Arbutus or Strawberry-tree.

Habitat in sylvis montosis, in occidentali Hiberniæ parte.

R. syn. 4. IX.

2. ARBUTUS caulibus procumbentibus, foliis rugosis *alpina.*

ferratis. *Sp. pl.* 566. *Fl. dan.* 73. *Hall. hist.* 1019.

Vaccinia rubra, foliis myrtinis crispis. *R. syn.* 457.

Vitis idæa foliis oblongis albicantibus. *B. pin.* 470.

Vitis idæa. *Clus. hist. I.* 61. *pann.* 77. 78.

Anglis, mountain Arbutus.

Habitat in montibus succis; in that end of Mull next to

Eycolumb-Kill, Mr. Llhwyd. *R. syn.* in monte

Ben Grihum dicta, et alibi in comitatu Sutherland.

D. Hope. 4. V.

3. ARBUTUS caulibus procumbentibus, foliis integerrimis. *uva ursi.*

Fl. lapp. 162. *t. 6. f.* 3. *dan.* 33. *Sp. pl.* 566.

Arbutus caulibus procumbentibus, foliis duris integerrimis. *Hall. hist.* 1018.

Uva ursi. *Clus. hist. I.* 63. *Park.* 1458.

Radix idæa putata et uva ursi. *B. hist. I.* 524.

Anglis, trailing Arbutus or Bearberry.

Habitat in sylvis et ericetis siccioribus. In the Duke of

Portland's wood, near Hexham, in Northumber-

land plentifully. *Wall. N. hist. Northumb.* 148. at

- Drummelzier in Twedale. Sir J. Nasmyth; *in ericetis ficcioribus in boreali parte Scotiae copiose* D. Hope. h. VI.

D I G Y N I A.

SCLERANTHUS. 562.

Cal. monophyllus. *Cor.* o. *Sem.* duo calyce inclusa.

- annuus.* 1. SCLERANTHUS calycibus fructus patulis. *Sp. pl.* 580. *Fl. suec.* 377. *dan.* 504.

Scleranthus floribus patulis. *Hall. hist.* 1551.

Knawel. *Trag.* 393. *R. syn.* 159.

Polygonum germanicum, vel Knawel germanorum. *Park.* 446.

Polygonum selinoides sive Knawel. *Ger. em.* 566.

Polygonum gramineo folio majus erectum. *B. pin.* 281.

Anglis, annual Knawel.

Habitat in arvis et locis arenosis. ☉. VII. VIII.

- perennis.* 2. SCLERANTHUS calycibus fructus clausis. *Sp. pl.* 580. *Fl. suec.* 378. *dan.* 563.

Scleranthus floris segmentis erectis. *Hall. hist.* 1550.

Knawel incanum flore majore perenne. *R. hist.* 213. *syn.* 160. t. 5. f. 1.

Alchimilla gramineo folio majori flore. *Tourn.* 509.

Vaill. paris. 4. t. 1. f. 5.

Anglis, perennial Knawel.

Habitat in arvis arenosis; about Elden in Suffolk, plentifully. ♀. VII. VIII.

CHRYSOSPLENIUM. 558.

Cal. quadrifidus s. quinquefidus, coloratus. *Cor.* o.

Caps. birostris, unilocularis polysperma.

- oppositifolium.* 1. CHRYSOSPLENIUM foliis oppositis. *Sp. pl.* 569. *Fl. suec.* 365. *dan.* 365. *Hall. hist.* 1549.

Chrysoisplenium s. Saxifraga aurea. *Tabern. hist.* 1224.

Saxifraga aurea. *Ger. em.* 841. *Park.* 425. *R. syn.* 158.

Saxifraga

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

nivalis. 2. SAXIFRAGA foliis obovatis crenatis subsessilibus, caule nudo, floribus congestis. *Sp. pl.* 573. *Fl. dan.* 28.

Saxifraga caule nudo simplici, foliis elliptico-subrotundis crenatis, floribus capitatis. *Fl. lapp.* 176. t. 2. f. 5. 6.

Saxifraga foliis oblongo-rotundis dentatis, floribus compactis. *R. syn.* 354. t. 16. f. 1.

Anglis, mountain Saxifrage.

Habitat in summis alpibus Arvonizæ et rupibus Clogwyn-y-Garnedd, Crib-y-ddescil, Glyder, etc. juxta Llanberris. D. Lhwyd. *R. syn.* in monte Ben-Buy dicto, D. Oakes; et in insulis Skye et Mull in Scotia. D. Robertson. 4. VII. VIII.

** Foliis indivisis, caule folioso.

oppositifolia. 3. SAXIFRAGA foliis caulinis ovatis oppositis imbricatis: summis ciliatis. *Sp. pl.* 575. *Fl. succ.* 369. *dan.* 34.

Saxifraga caule repente, foliis quadrifariam imbricatis, cartilagineis ciliatis. *Hall. hist.* 980.

Saxifraga alpina ericoides, flore cæruleo. *Tourn.* 253. *R. syn.* 353. *Sch. It.* VI. 428.

Sedum alpinum ericoides cæruleum. *B. pin.* 284. *pr.* 132. *Hist. Ox.* III. 480. f. 12. t. 10. f. 36.

Anglis, purple Saxifrage.

Habitat in rupibus et petris. In rupibus montis Ingleborough non ad summitatem, sed ad latus occidentale. D. Richardson; *R. syn.* In rupibus circa Llanberris copiose. 4. III. IV.

autumnalis. 4. SAXIFRAGA foliis caulinis linearibus alternis ciliatis, radicalibus aggregatis. *Sp. pl.* 575. *Fl. carn.* 293. t. 14. *dan.* 72. *Jacq. enum.* 71. et. 240.

Saxifraga foliis ellipticis ciliatis, caule simplici folioso, paucifloro. *Hall. hist.* 971.

Saxifraga alpina angustifolio, flore luteo guttato. *R. syn.* 353.

Sedum alpinum I. *Clus. pan.* 484. *Ger. em.* 516.

Sedum minus VI. f. alpinum I. *Clus. hist.* II. 60.

Sedum

Sedum alpinum flore pallido. *B. pin.* 284. *Hist. Ox.* III. 477. f. 12. t. 7. f. 3. *R. hist.* 1042.

Sedum alpinum floribus luteis maculosis. *B. pin.* 284?

Sedum parvum montanum luteum. *B. hist.* III. 693.

Sedum minus alpinum luteum nostras. *R. cat. angl.* 263. *hist.* 1042. *Hist. Ox.* III. 477.

Sedula montana pulchra. *Gesn. fasc.* 1. f. 4.

Anglis, autumnal Saxifrage.

Habitat in montosis; on Ingleborough-hill plentifully, and in Westmorland. *R. syn.* On Beeston Castle, Cheshire. *Blackf. Spec. bot.* 4. VII. VIII.

5. **SAXIFRAGA** foliis alternis lanceolatis integerrimis *hirculus*. acutis nudis, caule adscendente folioso.

Saxifraga foliis caulinis lanceolatis alternis nudis inermibus, caule erecto. *Sp. pl.* 576. *Fl. suec.* 370. *dan.* 120. *Gmel. fib.* IV. t. 65. f. 3.

Saxifraga foliis ellipticis, caule unifloro. *Hall. hist.* 972. t. 11.

Saxifraga angustifolia autumnalis, flore luteo guttato. *Breyn. Cent.* 104. *R. hist.* 1048. foliis florum magis acuminatis. *Breyn. Cent.* 106.

Hirculus friscus Dortmanni. *Clus. cur. post.* 5.

Hirculus qui chamæcisti genus. *Clus. cur. post.* 5. 6.

Chamæcistus friscus, foliis nardi celticæ. *B. pin.* 466.

Sedum palustre luteum bicornè nardi celticæ foliis. *Hist. Ox.* III. 477. f. 12. t. 8. f. 5.

Sedum angustifolium autumnale, flore luteo guttato. *Hist. Ox.* III. 477. f. 12. t. 8. f. 6.

Geum angustifolium autumnale, flore luteo guttato. *Tourn.* 252. *R. syn.* 355.

Anglis, marsh Saxitrage.

Habitat in palustribus turfosis; on Knutsford-moor, Cheshire. 4. VII. VIII.

*** *Foliis lobatis, caulibus erectis.*

6. **SAXIFRAGA** foliis radicatis aggregatis linearibus *cæspitosa*. integris trifidisque, caule erecto subnudo subifloro. *Sp. pl.* 578. *Fl. dan.* 71.

Saxifraga foliis integris et trifidis, caule subnudo, paucifloro.

paucifloro. *Hall. hist. 988. opusc. bot. 291. t. 1. f. 1.*

Saxifraga foliis partim integris partim trifidis. *Segu. ver. 1. 451. t. 9. f. 4. Sch. It. II. 49. VII. 518.*

Sedum tridactylites alpinum minus. B. pin. 284.

Anglis, matted Saxifrage.

Habitat in montibus supra Ambleside in comitatu Westmorlandico. 4. VII. VIII.

tridactyli- 7. **SAXIFRAGA** foliis caulinis cuneiformibus trifidis
tes. alternis, caule erecto ramoso. *Sp. pl. 578.*

Saxifraga foliis petiolatis, trilobatis, caule erecto, ramoso et folioso. *Hall. hist. 986.*

Saxifraga verna annua humilior. Journ. 252. R. syn. 354.

Sedum tridactylites tectorum. B. pin. 285.

Paronychia rutaceo folio. Ger. em. 624.

Paronychia foliis incis. Park. 556.

Anglis, rue-leaved Saxifrage or Whitlow-grass.

Habitat in muris, tectis, ruderatis. 6. IV.

Pharmac. Paronychiæ Herba.

granulata. 8. **SAXIFRAGA** foliis caulinis reniformibus lobatis,
ta. caule ramoso, radice granulata. *Sp. pl. 576. Fl. succ. 372. dan. 514. Mat. med. 215.*

Saxifraga foliis radicalibus reniformibus, obtuse dentatis, caulinis palmatis. *Hall. hist. 976.*

Saxifraga rotundifolia alba. B. pin. 309. R. syn. 354.

Saxifraga alba. Dod. pempt. 316. Ger. em. 841. vulgaris. Park. 424.

Anglis, white Saxifrage.

Habitat in pratis et pascuis siccioribus; prope Wandsworth copiose. 4. IV. V.

Pharmac. Saxifragæ albæ Radix, Folia, Flores, Semina.

*** *Foliis lobatis, caulibus procumbentibus.*

hypnoides. 9. **SAXIFRAGA** foliis caulinis linearibus trifidisve, stolonibus procumbentibus, caule erecto nudiusculo.
Sp. pl. 579. Fl. dan. 348.

Saxifraga foliis petiolatis trifidis caule subfolioso viscido. *Hall. hist. 989.*

Saxifraga

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Caryophyllus barbatus sylvestris. *B. pin.* 209.

Anglis, Deptford pink.

Habitat in pratis et pascuis glareosis, in sylva Charleton-wood dicta et alibi in Cantio; prope Croydon copiose.

☉. VII.

prolifer. 2. **DIANTHUS** floribus aggregatis capitatis, squamis calycinis ovatis obtusis muticis tubum superantibus. *Sp. pl.* 587. *Fl. dan.* 221.

Tunica capitulo compacto, ovali, calyce universali quadrifido. *Hall. hist.* 901.

Caryophyllus sylvestris prolifer. *B. pin.* 209. *Park.* 1338. *R. syn.* 337.

Armeria prolifera. *Lob. ic.* 449. *Ger. em.* 599.

Anglis, proliferus Pink.

Habitat in pratis et pascuis arenosis. Found by the Rev. Mr. Manningham in Selsey island, Suffex. *R. syn.* prope Norwich. *D. Humphrey.* ☉. VII.

*** Flores solitarii plures in eodem caule.

caryophyllus. *3. **DIANTHUS** floribus solitariis, squamis calycinis subovatis brevissimis, corollis crenatis. *Sp. pl.* 587.

Tunica ramis unifloris et bifloris, petalis levibus, stipulis calycinis brevissimis. *Hall. hist.* 896.

Caryophyllus hortensis simplex, flore major. *B. pin.* 208.

Caryophyllus simplex major. *Ger. em.* 590.

β *Caryophyllus simplex flore minore pallido rubente.* *B. pin.* 208.

Caryophyllus sylvestris simplex. *Ger. em.* 591.

Anglis, Clove Pink, July-flower or Carnation.

Habitat in muris antiquis; on Rochester, Deal, Sandown, &c. castles, plentifully. 4. VII.

Pharmac. Caryophylli Flores.

deltoides. 4. **DIANTHUS** floribus solitariis: squamis calycinis lanceolatis binis, corollis crenatis. *Sp. pl.* 588. *Fl. dan.* 577. *succ.* 382.

Caryophyllus minor repens nostras. *R. syn.* 335. *Dill. elth.* 402.

Caryo-

- Caryophyllus simplex supinus latifolius. *B. pin.* 209.
 β Tunica ramosior, flore candido cum corolla purpureo.
Dill. elth. 400. t. 298. f. 348.
Anglis, maiden Pink.
 Habitat in pratis et ericetis arenosis passim, supra Dup-
 pers-hill, prope Croydon. ♂. VI.—VIII.

5. DIANTHUS floribus subsolitariis, squamis calycinis *glaucus*.
 lanceolatis quaternis brevibus, corollis crenatis. *Sp.*
pl. 588.
 Tunica rupestris, folio cæsius molli, flore carneo.
Dill. elth. 401. tab. 298. f. 385.
 Armeriæ species flore in summo caule singulari. *R.*
syn. 336.
 Caryophyllus sylvestris, alines holostæ arvensis gla-
 bræ foliis, flore unico, calyce barbato. *Pluk. pb.*
 t. 81. f. 3.
Anglis, mountain Pink.
 Habitat in montosis, on Chedder-rocks, in Somerset-
 shire, Mr. Brewer. VI. VII.

*** *Caule unifloro herbaceo.*

- *6. DIANTHUS caulibus subunifloris, squamis caly- *arenarius*
 cinis ovatis obtusis, petalis multifidis, foliis lineari-
 bus. *Sp. pl.* 589. *Fl. suec.* 384.
 Caryophyllus sylvestris humilis, flore unico. *B. pin.*
 209?
 Caryophyllus sylvestris I. *Clus. hist.* I. 282.
 Caryophyllus plumarius albus. *Ger. em.* 591?
 Armerius flos tertius. *Dod. pempt.* 176.
Anglis, common Pink.
 Habitat in muris antiquis. ♀. VII. VIII.

TRIGYNIA.

CUCUBALUS. 566.

Cal. inflatus. *Petala* quinque unguiculata absque co-
 rona ad faucem. *Caps.* trilocularis.

1. CUCUBALUS calycibus campanulatis, petalis dis- *bacciferus*
 tantibus,

tantibus, pericarpiis coloratis, ramis divaricatis.

Sp. pl. 591.

Cucubalus. *Mill. Ic. t.* 112. *Plinii. Lugd.* 1429.

Tourn. 339. *R. syn.* 267.

Visago baccifera, petalis ferratis. *Hall. hist.* 912.

Alfne scandens baccifera. *B. pin.* 250.

Anglis, berry-bearing Champion.

Habitat in sepibus in insula Mona. *D. Richardson.*

R. syn. VI. VII.

Behen.

2. CUCUBALUS calycibus subglobosis glabris reticulato-venosis, capsulis trilocularibus, corollis subnudis. *Sp. pl.* 591.

Viscago caule brachiato, nodoso, calycibus inflatis venosis. *Hall. hist.* 913.

Lychnis sylvestris, quæ Behen album vulgo. *B. pin.* 205. *R. syn.* 337.

Behen album officinarum. *Ger. em.* 678.

Papaver spumeum sive Behen album vulgo. *Park.* 263.

Anglis, bladder Champion or Spatling-Poppy.

Habitat in arvis, pratis et pascuis siccioribus. 4. VIII.

Obs. variat foliis scabris.

viscosus.

3. CUCUBALUS floribus lateralibus undique decumbentibus, caule indiviso, foliis basi reflexis. *Sp. pl.* 592. *Fl. succ.* 386.

Lychnis major noctiflora Dubrensis perennis. *R. syn.* 340. *hist.* 995.

Lychnis montana noctiflora hirsuta latifolia alba, floribus absque appendicibus. *Till. pis.* 105.

Anglis, Dover Champion.

Habitat in montosis. Found on Dover-cliffs, by Mr. Newton. *R. syn.* 4. VII.

atites.

4. CUCUBALUS floribus dioicis, petalis linearibus indivisis. *Sp. pl.* 594. *Fl. dan.* 518.

Viscago floribus verticillatis, spicatis, sexu distinctis petalis linearibus. *Hall. hist.* 920.

Lychnis viscosa flore muscosa. *B. pin.* 206.

Sesamoides salamanticum magnum. *Ger. em.* 493.

Muscipula salamantica major. *Park.* 636.

Anglis, Spanish Champion or Catchfly.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Habitat in arvis arenosis, passim, prope Coombe in agro Surriensi. ☉. VI. VII.

Obs. Petala obtusa non crenata, subinde emarginata; Calycis fructus maturi horizontaliter reflexi, non penduli, angulati, visceri, pilosi.

quinque-
vulnera. 2. SILENE petalis integerrimis subrotundis, fructibus erectis alternis. *Sp. pl.* 595.

Lychnis sylvestris lanuginosa minor. *B. pin.* 206.

Lychnis hirsuta flore eleganter variegato. *R. hist.* 997. *Anglis*, variegated Catchfly.

Habitat in arvis arenosis juxta Wrotham in Cantio. ☉. VII.

** Floribus lateralibus confertis.

nutans. 3. SILENE petalis bifidis, floribus lateralibus secundis cernuis, panicula nutante. *Sp. pl.* 596. *Fl. dan.* 242.

Viscago foliis lanceolatis hirsutis, floribus paniculatis nutantibus. *Hall. hist.* 915.

Lychnis sylvestris alba IX. clusii. *Ger. em.* 470.

Lychnis montana viscosa alba latifolia. *B. pin.* 205.

Lychnis sylvestris alba f. ocymoides minus album. *Park.* 631.

Anglis, Nottingham Catchfly.

Habitat in pratis montosis. On the walls of Nottingham-Castle. *R. syn.* 4. VI. VII.

ameena. 4. SILENE petalis bifidis: coronula subcoadunata, floribus secundis, pedunculis oppositis trifloris, ramis alternis. *Sp. pl.* 596.

Lychnis marina anglica. *Ger. em.* 469. *R. syn.* 337.

Lychnis marina repens alba. *Park.* 638.

β *Lychnis maritima anglica amplioribus foliis.* *D. Brewer. Rand. hort. Chels.*

Anglis, sea Catchfly.

Habitat in littoribus maritimis arenosis frequens. 4. VI—VIII.

*** Floribus

*** *Floribus ex dichotomia caulis.*

5. SILENE calycibus fructus globosis acuminatis *conoides*.
 striis triginta, foliis glabris, petalis integris. *Sp. pl.*
 598.

Lychnis sylvestris angustifolia caliculis turgidis striatis. *B. pin.* 205. *R. syn.* 341.

Lychnis sylvestris altera incana caliculis striatis. *Lob. ic.* 338.

Anglis, corn Catchfly.

Habitat in arvis arenosis ; a little to the north of Sandown-castle plentifully. *R. syn.* ☉. VI. VII.

6. SILENE calycibus decemangularibus : dentibus tubum æquantibus, caule dichotomo, petalis bifidis. *noctiflora*.
Sp. pl. 599. *Syst. Nat.* 311.

Viscago foliis ovato-lanceolatis hirsutis, caule brachiato, calycibus venosis. *Hall. hist.* 911.

Lychnis noctiflora. *B. pin.* 205. *Park.* 632. *R. syn.* 340.
Anglis, night-flowering Catchfly.

Habitat in arvis. Found by Mr. Dent among corn, on the left hand of the road leading from Newmarket to Canvass-hall in Wood-ditton. *R. syn. circa* Norwich. *D. Rose.* ☉. VII.

7. SILENE floribus fasciculatis fastigiatis, foliis superioribus cordatis glabris, petalis integris. *Armeria*.
Syst. Nat. 311. *Sp. pl.* 601. *Fl. dan.* 559.

Viscago foliis glabris ovato-lanceolatis, floribus umbellatis. *Hall. hist.* 916.

Lychnis vilcosa, purpurea latifolia lævis. *B. pin.* 205.
R. syn. 341.

Lychnis sylvestris I. *Clus. hist.* 288.

Anglis, common Catchfly.

Habitat in arvis et muris antiquis ; on the banks of the river half a mile below Chester. *D. Richardson.*
R. syn. ☉. VII. VIII.

STELLARIA. 568.

Cal. pentaphyllus, patens. *Petala* quinque, bipartita.
Caps. unilocularis polysperma.

I. STELLARIA

Holostea. 1. **STELLARIA** foliis lanceolatis ferrulatis, petalis bifidis. *Sp. pl.* 603. *Fl. suec.* 392.

Alfinae foliis gramineis, ciliatis. *Hall. hist.* 884.

Caryophyllus holosteus arvensis glaber, flore majore.

B. pin. 210. *R. syn.* 346.

Gramen leucanthemum. *Ger. em.* 47. *Park.* 1325.

Anglis, greater Stitchwort.

Habitat in sylvis et sepibus frequens. 4. IV. V.

graminea. 2. **STELLARIA** foliis linearibus integerrimis, floribus paniculatis. *Fl. suec.* 393. *Sp. pl.* 604.

Alfinae foliis lanceolatis lineatis, petiolis ramosis. *Hall. hist.* 882.

Caryophyllus holosteus arvensis, glaber, flore minore.

B. pin. 210. *R. syn.* 346.

Gramen leucanthemum alterum. *Ger. em.* 47.

Gramen leucanthemum minus. *Park.* 1325.

β Caryophyllus holosteus arvensis medius. *R. syn.* 347.
Fl. dan. 414.

γ Alfinae foliis lanceolatis, petalis bipartitis, petiolis unifloris. *Hall. hist.* 881.

Alfinae longifolia uliginosis proveniens locis. *B. hist.*

III. 565. *R. syn.* 347. *Fl. dan.* 415.

Anglis, lesser Stitchwort.

Habitat in pratis, pascuis & sepibus frequens; β in palustribus. γ. in pratis humidis ad fontes et ad scaturigines frequens. 4. VIII.

nemorum. 3. **STELLARIA** foliis cordatis petiolatis, panicula pedunculis ramosis. *Sp. pl.* 603. *Fl. dan.* 271.

Myosotis foliis petiolatis, cordatis, tubis ternis. *Hall. hist.* 886.

Alfinae montana, folio smilacis instar, flore lacineato.

Hist. Ox. II. 550. f. 5. t. 23. f. 2. *R. syn.* 347.

Alfinae altissima nemorum. *B. pin.* 250.

Alfinae hederacea montana maxima. *Col. Ecpbr. I.* p. 289. t. 290. *Park.* 761.

Anglis, wood Stitchwort.

Habitat in sylvis et sepibus humidis, et ad ripas fluviorum in comitatibus Eboracensi, Lancastriensi, Westmorlandico, Cumberlandico, passim. 4. VII. VIII.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Alfina alpina glabra tenuissimis foliis, floribus albis.
Herm. par. 12. t. 12.

Alfina caryophylloides tenuifolia, flore albo punctato.
Pluk. ph. t. 7. f. 3.

β *Arenaria* foliis linearibus erectis, subtus striatis, floribus fastigiatis, inæqualiter pedunculatis. *Ger. gallopr.* 405. t. 15. f. 1.

Anglis, mountain Sandwort.

Habitat in montosis Eboracensibus circa Settle, ut et Derbiansibus, copiose; circa Kendal in comitatu Westmorlandico, prope Holywell, et circa Llanberris in Wallia copiose. 4. V—VIII.

laricifolia.

5. *ARENARIA* foliis setaceis, caule superne nudiusculo, calycibus subhirsutis. *Sp. pl.* 607. *Jac. austr.* III. 39. t. 272.

Alfina foliis linearibus, angustissimis, calycibus tubulosis villosis. *Hall. hist.* 869.

Alfina alpina junceo folio. *B. pin.* 251. *pr.* 118. *R. syll.* 60. *hist.* 1031.

Auricula muris pulchro flore, folio tenuissimo. *B. hist.* III. 360.

Lychnoides juniperi folio perennis. *Vaill. paris.* 121.

Anglis, larch-leaved Sandwort.

Habitat in montibus Westmorlandicis. 4. VII. VIII.

tenuifolia.

6. *ARENARIA* foliis subulatis, caule paniculato, capsulis erectis, petalis calyce brevioribus lanceolatis. *Sp. pl.* 607. *Syst. Nat.* 313. *Fl. dan.* 389.

Alfina foliis linearibus, glabris, petalis calyce brevioribus. *Hall. hist.* 866.

Alfina tenuifolia. *B. hist.* III. 364. *R. syn.* 350. *Vaill. paris.* 7. t. 3. f. 1. *Sigu. ver.* I. 418. t. 6. f. 2.

Anglis, fine-leaved Sandwort.

Habitat in pratis et pascuis arenosis; in the corn-fields on the borders of Triplow-heath, Gogmagog-hills, &c. in Cambridgeshire. *R. syn.* In Cornbury Quarry, near Charlbury, Oxfordshire, Blacks. *supra murum in oppido Battersea, prope Londinum.* 4. VI. VII.

rubra.

7. *ARENARIA* foliis filiformibus, stipulis membranaceis vaginantibus. *Sp. pl.* 606.

Alfina foliis linearibus, stipulis ovato-lanceolatis, argenteis. *Hall. hist.* 872.

- Spergula purpurea*. *B. hist.* III. 722. *R. syn.* 351.
Saginæ spergula minima. *Park.* 561.
 Alfine, *Spergulæ facie minor*. f. *Spergula minor sub-
 cæruleo flore*. *B. pin.* 251, *pr.* 119.
 β *Arenaria foliis linearibus carnosis, stipulis membrana-
 ceis, caulibus pubescentibus*. *Syst. Nat.* 312: *Sp.*
pl. 606.
Spergula maritima flore parvo cæruleo, semine vario.
Dill. giss. 30. *R. syn.* 351.
 γ *Arenaria foliis linearibus longitudine internodiorum marina*.
Hort. cliff. 173. *Gron. virg.* 161. *Sp. pl. ed. I.* 423.
 Alfine *spergulæ facie media*. *B. pin.* 251. *R. syn.* 351.
Spergula marina: *Ger. em.* 1125. nostras. *B. hist.* III.
 772.
Anglis, purple Sandwort, or Spurry γ Sea Spurry.
 Habitat in pratis et arvis arenosis β in pascuis marinis,
 γ in littoribus marinis. ☉: VI—VIII.

CHERLERIA. 570:

- Cal.* pentaphyllus. *Nectararia* quinque bifida, petaloi-
 dea. *Antheræ* alternæ steriles. *Caps.* trilocularis,
 trivalvis.
- i. **CHERLERIA.** *Sp. pl.* 608. *Hall. hist.* 859. t. 21. *Sedoides*.
opusc. 300. t. 1. f. 3. *Segu. ver.* III. 180. t. 4. f. 3.
Pen. Voy. t. 33. *Jac. austr.* III. 46. t. 284.
Sedum montanum perpusillum; luteolis floribus. *Park.*
 738.
Sedum montanum perpusillum luteum: *Hist. Ox.* III.
 471. f. 12. t. 6. f. 14.
Lychnis alpina, muscosis foliis densius stipatis, flori-
 bus parvis, calyce duriore. *Pluk. ph.* t. 42. f. 8.
Anglis, dwarf Cherleria.
 Habitat in montibus in boreali parte Scotiæ. D. Hope.
 Found on Baikeval in the island of Rum. *Penn. Voy.*
 4. VII. VIII.



P E N T A G Y N I A.

C O T Y L E D O N. 578.

Cat. quinquefidus. *Cor.* monopetala. Squamæ nectari-
feræ quinque ad basin germinis. *Caps.* quinque.

umbilicus 1. COTYLEDON foliis peltatis crenatis, caule sub-
simplici, floribus pendulis, bracteis integris.

Cotyledon *Dod. stirp.* 39. vera radice ruberosa. *B.*
hist. III. 683. *R. syn.* 271.

Cotyledon major. *B. pin.* 285.

Cotyledon umbilicus veneris. *Clas. hist.* II. 63.

Umbilicus repens. *Cam. epit.* 858.

Umbilicus veneris. *Ger. em.* 528. vulgaris. *Park.* 740.

Sedum luteum murale spicatum, folio umbilicato ro-
tundo. *Hist. Ox.* III. 470. f. 12. t. 10. f. 4.

Anglis, common navel-wort, or Wall Penny-wort.

Habitat in muris antiquis et saxosis humidiusculis passim;
in occidentali parte Angliæ et in Arvoniam frequens, circa
Troutbeck in Westmorlandia. *N.* VI—VIII.

Obs. variat Foliis imis reniformibus.

luteo. 2. COTYLEDON foliis peltatis crenatis caule sub-
simplici, floribus erectis bracteis dentatis.

Cotyledon foliis cucullato-peltatis, serrato-dentatis
alternis, caule ramoso, floribus erectis. *Sp. pl.* 615.

Cotyledon flore luteo radice repente. *Dodart. Mem.*
265. t. 73.

Cotyledon radice tuberosa longa repente. *R. hist.*
1878.

Cotyledon luteum umbilicatum spicatum, radice re-
pente majus. *Hist. Ox.* III. 471.

Anglis, yellow Navel-wort.

Habitat in muris et saxosis humidiusculis, in occidentali
parte comitatus Eboracensis, D. Tofield. in horto
D. Clement, qui recepit ex agro Somersensi, vidi.
N. VI.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Sedum minus e rupe s. Vincentii. *R. fyn.* 271.

Sedum minus vincentii. *Pet. H. br. t.* 41. *f.* 8.

Anglis, rock Stonecrop.

Habitat in rupibus St. Vincentii prope Bristolium; in rupibus Hysfæ vallecule Nantphrancon imminentibus. *R. fyn.* VIII.

album.

4. SEDUM foliis oblongis obtusis teretiusculis sessilibus patentibus, cyma ramosa. *Sp. pl.* 619. *Fl. dan.* 66.

Sedum caule glabro, foliis teretibus. umbellis ramosis, floribus petiolatis. *Hall. hist.* 959.

Sedum minus officinarum. *Ger. em.* 512.

Vermicularis flore albo. *Park.* 733.

Anglis, white Stonecrop.

Habitat in muris, tectis, passim. On a wall just beyond the chapel Kentish-town. *Blacks.* 4. VI. VII.

Pharmac. Vermicularis Herba.

acris.

5. SEDUM foliis subovatis adnato-sessilibus gibbis erectiusculis alternis, cyma trifida. *Sp. pl.* 619.

Sedum foliis conicis confertis, caulibus ramosis, summis trifidis. *Hall. hist.* 966.

Sedum parvum acre flore luteo, *B. hist.* III. 694. *R. fyn.* 270.

Sempervivum minus vermiculatum acre. *B. pin.* 283.

Vermicularis seu Illecebra minor acris. *Ger. em.* 517.

Illecebra minor s. Sedum III. Dioscoridis, *Park.* 735.

β Sedum foliis subovatis adnato-sessilibus gibbis erectiusculis sexfariam imbricatis. *Sp. pl.* 620.

Sedum foliis teretibus, ternatis, caulibus simplicibus trifidis. *Hall. hist.* 965.

Sempervivum minus vermiculatum insipidum. *B. pin.* 284.

Sempervivum minimum. *Cam. epit.* 856.

Anglis, wall Stonecrop, or Pepper.

Habitat in muris tectis, rupibus et pascuis siccioribus prope Northfleet et Sheerness; in insula Shepey capiose. VI. VII.

Pharmac. Sedi minoris Herba.

-*anglicum.* 6. SEDUM foliis subovatis adnato-sessilibus gibbis alternis, cyma ramosa bifida.

Sedum

Sedum minimum non acre flore albo. R. syn. 270. t. 12. f. 2.

Sedum minimum, flore mixto ex albo et rubro. Merr. pin.

Anglis, English Stonecrop.

Habitat in rupibus, saxosis et arenosis vitrescentibus in comitatibus Westmorlandico, Cumberlandico, Lancastriensi, Devonensi, et Cornubiensi, copiose in Wallia frequens; prope Deal in Cantia. 4. VII. VIII.

Desc. Radix ramosa perennis. Caulis procumbens, teres, foliatus, glaber, ramosus, rubescens; ramis alternis. Folia alterna, subtrifariam imbricata, adnato-sessilia, subovata, acutiuscula superne planiuscula, subtus gibba, pulposa, nuda. Cyma ramosa, bifida, subrecurva. Flores sessiles secundi; Corolla alba seu purpurascens-alba. Capsulæ purpurascens.

7. *SEDUM* foliis oppositis ovatis obtusis carnosis, caule *dasyphyllum* infirmo, floribus sparsis. *Sp. pl. 618. Jac. Hort. lum. vind. II. 71. t. 153.*

Sedum foliis conicis obtusis glaucis, reticulatis, caule ramoso viscido. Hall. hist. 961.

Sedum minus circinato folio. B. pin. 283. Hist. Ox. III. 473. f. 12. t. 17. f. 35. R. syn. 271.

Sedum minus VII. Clus. hist. I. 1. 60.

Aizoon dasyphyllum. Dalech. hist. 1133.

Anglis, round-leaved Stonecrop.

Habitat in muris et tectis, plentifully on the houses and walls at Market-eit, near Market-street. Mr. Th. Knowlton. R. syn. in muris prope Hammersmith et Kew prope Londinum, et in muris apud Bugden in agro Huntingtoniensi. ☉. VII.

8. *SEDUM* caule erecto, foliis planiusculis, pedunculis *villosum*, que subpilosis. *Sp. pl. 620. Fl. dan. 24.*

Sedum hirsutum et viscidum, foliis linearibus obtusis planiusculis. Hall. hist. 962.

Sedum purpureum pratense. B. hist. III. 692. R. syn. 270.

Sedum palustre subhirsutum purpureum. B. pin. 285.

Sedum minus palustre. Ger. em. 515.

Sedum arvense s. palustre flore rubente. Park. 734.

DECANDRIA PENTAGYNIA.

Anglis, hairy Stonecrop.

Habitat in pratis et pascuis montosis, humidis, on the moist rocks about Ingleborough-hill, on Hartside-hill, near Gambleby, in the way to Osten in Cumberland; in et juxta montem Hincklethwaite prope Settle. Dr. Richardson, R. syn. In Scotia frequens. ☉. VI. VII.

O X A L I S. 582.

Cal. pentaphyllus. *Petala* unguibus connexa. *Caps.* angulis dehiscens, pentagona.

Acetosella. i. OXALIS scapo unifloro, foliis ternatis, radice squamoso articulata. *Sp. pl.* 620. *Hall. hist.* 928.

Oxys alba. *Ger. em.* 1201. *R. syn.* 281.

Trifolium acetosum vulgare. *B. pin.* 330. *Park.* 746.

β Trifolium acetosum vulgare. flore purpureo. *Merr.*

Anglis, common Wood-forrel.

Habitat in sylvis, sepibus, ericetis umbrosis. ☿. IV.

Pharmac. Lujulæ Folia.

A G R O S T E M A. 583.

Cal. monophyllus, coriaceus. *Petala* quinque unguiculata: Limbo obtuso, indiviso. *Caps.* unitocularis.

Githago. i. AGROSTEMA hirsuta calycibus corollam æquantibus, petalis integris nudis. *Sp. pl.* 624. *Fl. dan.* 576.

Lychnis calycibus longissime caudatis. *Hall. hist.* 926.

Lychnis segetum major. *B. pin.* 204. *R. syn.* 338.

Lychnoides segetum f. Nigellastrum. *Park.* 632.

Pseudo-melanthium. *Dalech. hist.* 438. *Ger. em.* 1087.

B. hist. III. 341.

Anglis, corn Cockle.

Habitat in arvis frequens. ☉. VI. VII.

L Y C H N I S. 584.

Cal. monophyllus, oblongus, lævis. *Petala* quinque unguiculata: Limbo subbisido. *Caps.* quinquelocularis.

LYCHNIS

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

CERASTIUM. 585.

Cal. pentaphyllus. *Petala* bifida. *Caps.* unilocularis, apice dehiscens.

• *Capsulis oblongis.*

*pulga-
tum.*

1. CERASTIUM foliis ovatis, panicula dichotoma, petalis calycē æqualibus, caulibus diffusis.

Cerastium foliis ovatis, petalis calyce æqualibus caulibus diffusis. *Sp. pl.* 627.

Myosotis foliis ovato-lanceolatis, petalis calycis longitudine. *Hall. hist.* 893.

Myosotis arvensis hirsuta, parvo flore albo. *Vaill. paris.* 142. t. 30. f. 1.

Alfne hirsuta, magno flore. *B. pin.* 251.

Alfne hirsuta myosotis. *Lob. adv.* 195. *R. syn.* 349.

Anglis, common Cerastium, or Mouse-ear-chickweed.

Habitat in pratis, pascuis et muris. ☉. VI.

viscosum.

2. CERASTIUM erectum villos-viscosum, panicula dichotoma, floribus pedunculo longioribus.

Cerastium erectum villos-viscosum, *Sp. pl.* 627.

Myosotis hirsuta et viscosa. *Hall. hist.* 895.

Myosotis hirsuta altera viscosa. *Vaill. paris.* 142. t. 30. f. 3.

Alfne viscosa. *Park.* 764.

Alfne hirsuta altera viscosa. *B. pin.* 251.

Alfne hirsuta myosotis latifolia precosior. *R. syn.* 348.

Anglis, viscos Cerastium, or Mouse-ear-chickweed.

Habitat in pratis, pascuis et muris. ☉. IV. V.

*semide-
candrium.*

3. CERASTIUM floribus pentandris, petalis emarginatis. *Sp. pl.* 627. *Hort. cliff.* 173. *Fl. suec.* 416.

Cerastium hirsutum minus, parvo flore. *Dill. giss.* 80. *R. syn.* 348. t. 15. f. 1.

Myosotis caule simplici, foliis ovatis hirsutis, tubis ternis. *Hall. hist.* 894.

Myosotis arvensis hirsuta minor. *Vaill. paris.* 142. t. 30. f. 2.

Alfne hirsuta minor. *B. pin.* 251.

Anglis,

Anglis, least Cerastium, or Mouse-ear-chickweed.
Habitat in muris et pascuis frequens. ☉. IV.

4. CERASTIUM foliis lineari-lanceolatis obtusis glabris, corollis calyce majoribus. *Sp. pl.* 628. *Fl. suec.* 417. *dan.* 626.

Myosotis foliis linearibus lanceolatis, petalis calyce duplo longioribus. *Hall. hist.* 889.

Myosotis arvensis hirsuta, flore majore. *Vaill. paris.* 141. t. 50. f. 4.

Caryophyllus arvensis hirsutus, flore majore. *B. pin.* 210. *R. syn.* 348.

Caryophyllus holosteus. *Ger. em.* 595. arvensis hirsutus. *Park.* 1339.

Anglis, Corn Cerastium, or Mouse-ear-chickweed.

Habitat in arvis, pratis et pascuis arenosis supra Dupper's-hill prope Croydon. ☿. V.

5. CERASTIUM foliis ovato-lanceolatis, caule diviso, alpinum, capsulis oblongis. *Sp. pl.* 628. *Fl. dan.* 6.

Myosotis foliis ovato-lanceolatis subhirsutis, petalis calyce duplo longioribus. *Hall. hist.* 888.

Alfene myosotis facie, Lychnis alpina, flore amplo nigro repens. *R. syn.* 349. t. 15. f. 2.

Anglis, alpine Cerastium, or Mouse-ear-chickweed.

Habitat in montibus, et juxta scaturigines, ad latera montis Snowdon. Nullibi frequentius quam juxta scaturigines aquarum montis excelsi Wyddfa boream versus. Dr. Richardson. *R. syn.* in monte Clogwyn dŷ yn yr Arddu dicto prope Llanberris. ☿. VI. VII.

•• Capsulis subrotundis.

6. CERASTIUM foliis ovatis, floribus umbellatis, petalis subintegris, fructibus pendulis. *umbellatum.*

Holosteum floribus umbellatis. *Læst. it.* 120. *Sp. pl.* 130.

Holosteum caryophyllæum arvense. *Tabern. ic.* 233.

Caryophyllus arvensis umbellatus, folio glabro. *B. pin.* 210.

Caryophyllus holosteus arvensis. *Ger. em.* 595.

Alfene

Alfinae floribus umbellatis, petalis ferratis. *Hall. hist.* 878.

Anglis, umbellated Cerastium.

Habitat in muris et arvis arenosis, in muris antiquis et aggeribus circa Norwich D. Rose, et circa Bury in Suffolgia. D. Cullum. \odot . IV. V.

Obs. variat floribus triandris, tetrandris et pentandris trigynis, et floribus decandris pentagynis.

latifolium.

7. CERASTIUM foliis ovatis subtomentosis, ramis subunifloris, capsulis globosis. *Sp. pl.* 629.

Myosotis foliis tomentosis ovatis. *Hall. hist.* 887.

Caryophyllus holosteus alpinus latifolius. *B. pin.* 210. *pr.* 104.

Alfinae myosotis lanuginosa alpina grandiflora, seu Auricula muris villosa, flore amplo membranaceo. *R. syn.* 349.

Herba facie auriculæ muris. *Mart. Spitz. T. G. f. d.* *Anglis*, mountain Cerastium.

Habitat in rupe prope Clogwyn y Garnedd. totius Cambriæ altissima juxta Llanberris in Arvoniam copiose *R. syn.* 4. V. VI.

aquaticum.

8. CERASTIUM foliis cordatis sessilibus floribus solitariis, fructibus pendulis. *Sp. pl.* 628.

Alfinae foliis ovato-cordatis, imis petiolatis, tubis quinque. *Hall. hist.* 885.

Alfinae aquatica major. *B. pin.* 251.

Alfina major repens perennis. *B. hist.* III. 362. *R. syn.* 347.

Alfina major. *Cam. epit.* 851. *Tabern.* 707. *Ger. em.* 611. et maxima. *Park.* 759.

Anglis, water Cerastium.

Habitat in aquis et ad ripas fluviorum. 4. VII.

SPERGULA. 586.

Cal. pentaphyllus. *Petala* quinque integra. *Caps.* ovata, unilocularis, quinquevalvis.

arvensis. 1. SPERGULA foliis verticillatis, fructibus pendulis. *Spergula* foliis verticillatis, floribus decandris. *Sp. pl.* 630.

Spergula.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

DECANDRIA PENTAGYNIA.

Habitat in pratis et ericetis arenosis passim ; prope Cobham et Esler in comitatu Surriensi ; in Devoniam et Cornubiam frequens. 4. VI. VII.

Desc. Radix perennis, ramosa, repens. Caules plures ramosi procumbentes. Folia opposita, connata, subulata, ciliata, acuta, nuda in axillis foliorum, foliis minoribus fasciculatis. Pedunculi laterales, ad apicem rami, solitarii, longissimi uniflori, teretes. Flores cernui, albi, Stamina quinque raris decem.

*Classis **

Classis XI.

D O D E C A N D R I A

Stamina inter XI. et XIX. inclusive.

M O N O G Y N I A.

A S A R U M. 589.

Cal. quadrifidus, germi ni insidens. *Cor.* o. *Capsf.*
coriacea, coronata.

I. ASARUM foliis reniformibus obtusis binis. *Sp. pl. europaeum.*
633. *Fl. suec.* 421. *dan.* 633.

Asarum foliis reniformibus subhirsutis. *Hall. hist.*
1547.

Asarum. *B. pin.* 197. *hist. III.* 548. *R. syn.* 158.

Asarum vulgare. *Ger. em.* 836. *Park.* 266.

Anglis, common Asarabacca.

Habitat in nemorosis. In a hedge on the left hand, as soon as you are within the gate, the first field from Charnell's green, going to Sir Thomas Seabright's beach wood, Th. Knowlton; in several woods in Lancashire. R. syn. 4. V.

Pharmac. Asari Radix, Folia.

L Y T H R U M. 604.

Cal. duodecim-fidus. *Petala* sex, calyci inserta.
Capsf. bilocularis, polysperma.

I. LYTHRUM foliis oppositis cordato-lanceolatis, floribus spicatis dodecandris. *Sp. pl.* 640.

Salicaria foliis lanceolatis, subhirsutis, floribus spicatis. *Hall. hist.* 854.

Salicaria vulgaris purpurea, foliis oblongis. *Tourn.* 253. *R. syn.* 367.

Lyfimachia

Lyfimachia purpurea. *Ger. em.* 476. *Park.* 546.

Lyfimachia spicata purpurea. *B. pin.* 246.

β *Lyfimachia purpurea trifolia, caule hexagono.* *Spig. isag. R. syn.* 367.

Anglis, purple Loosestrife.

Habitat in paludibus et ad ripas fluviorum. 4. VII. VIII.

hyssopifolia.

2. **LYTHRUM** foliis alternis linearibus, floribus hexandris. *Sp. pl.* 642. *Fac. austr. II.* 20 t. 133.

Salicaria foliis linearibus, floribus per alas sessilibus. *Hall. hist.* 583.

Salicaria hyssopifolia latiore et angustiore. *Tourn.* 253. *R. syn.* 367. *Hall. jen.* 147. t. 6. f. 3.

Hyssopifolia. *B. pin.* 218. *aquatica.* *B. hist.* III. 792.

Gratiola angustifolia. *Ger. em.* 581. s. minor. *Park.* 220.

Anglis, hedge Loosestrife or hyssop.

Habitat in aquosis et ubi aquæ per hyemem stagnarunt, sed rarius, in ericeto Hounsejano. ☉. VIII.

D I G Y N I A.

AGRIMONIA. 607.

Cal. quinquedentatus, altero obvallatus. *Petala* quinque. *Sem.* duo infundo calycis.

eupatoria. 1. **AGRIMONIA** foliis caulinis pinnatis: impari petiolato, fructibus hispidis. *Sp. pl.* 643.

Agrimonia foliis pinnatis, pinnulis alterne minimis. *Hall. hist.* 991.

Agrimonia. *Ger. em.* 712. *R. syn.* 202. *vulgaris.* *Park.* 594.

Eupatorium. *Fuchs. hist.* 244. *Cam. ep.* 759. *veterum* s. *Agrimonia.* *B. pin.* 321.

Anglis, common Agrimony.

Habitat in marginibus agrorum et nemorosis. 4. VI VII.

Pharmac. Agrimonix Herba.

TRIGYNIA.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Peplis maritima folio obtuso. *B. pin.* 293.

Anglis, purple Spurge.

Habitat in littoribus maritimis arenosis; inter Pensans et Marketjeu in Cornubia; prope Exmouth in Devonia copiose. ☉. VII. VIII.

** *Umbella trifida.*

peplus.

2. EUPHORBIA umbella trifida: dichotoma, involu-
cellis ovatis, foliis integerrimis obovatis petiolatis.

Sp. pl. 653. *Amœn. Acad.* III. 115.

Tithymalus foliis rotundis, stipulis floralibus cordatis, petalis argute corniculatis. *Hall. hist.* 1049.

Tithymalus parvus annuus, foliis subrotundis non crenatis, *Peplus* dictus. *R. syn.* 313.

Peplus s. *Esula* rotunda. *B. pin.* 292. *Ger. em.* 503.

Anglis, petty Spurge.

Habitat in cultis oleraceis. ☉. VII. VIII.

exigua.

3. EUPHORBIA umbella trifida: dichotoma, involu-
cellis lanceolatis, foliis linearibus. *Sp. pl.* 654.

Amœn. Acad. III. 118. *Fl. dan.* 592.

Tithymalus foliis linearibus, stipulis lanceolatis aristatis. *Hall. hist.* 1048.

Tithymalus s. *Esula* exigua. *B. pin.* 295.

Tithymalus leptophyllus. *Park.* 193. *R. syn.* 313.

Esula exigua *Tragi.* *Ger. em.* 502.

Anglis, dwarf Spurge.

Habitat in arvis. ☉. VII.

*** *Umbella quinquefida.*

*portlan-
dica.*

4. EUPHORBIA umbella quinquefida: dichotoma, involu-
cellis subcordatis concavis, foliis lineari-lanceolatis acutis glabris patentibus. *Sp. pl.* 656.

Tithymalus maritimus minor, *Portlandicus.* *R. syn.* 313. t. 24. f. 6.

Anglis, Portland Spurge.

Habitat in arenosis maris littoribus. Found by the Rev. Mr. Stonestreet in the narrow neck of land which joins Portland to Dorsetshire. *R. syn.* prope Exmouth,

mouth, in Devon; in littoribus Cornubiæ passim; prope Carnarvon in Wallia, &c. ☉. VII. VIII.

5. EUPHORBIA umbella subquinquefida; bifida, in-paralias. involucellis cordato-reitiformibus, foliis sursum imbricatis. *Sp. pl.* 657. *Amæn. Acad.* III. 129.

Tithymalus maritimus: *B. pin.* 291.

Tithymalus paralius. *Ger. em.* 498. *R. syn.* 312.

Tithymalus paralius s. maritimus. *Park.* 184.

Anglis, sea Spurge.

Habitat in littoribus maritimis. ☿. VII. VIII.

6. EUPHORBIA umbella quinquefida: trifida: dichohelioscopia, involucellis obovatis, foliis cuneiformibus ferratis. *Sp. pl.* 658. *Amæn. Acad.* III. 124.

Tithymalus foliis petiolatis, subrotundis ferratis, stipulis rotundis ferratis. *Hall. hist.* 1050.

Tithymalus helioscopius. *Cam. epit.* 963. *Ger. em.* 498. *B. pin.* 291. *R. syn.* 313.

Anglis, sun Spurge or Water-wort.

Habitat in cultis oleraceis. ☉. VII. VIII.

7. EUPHORBIA umbella quinquefida: subtrifida: biverrucosafido, involucellis ovatis, foliis lanceolatis ferrulatis villosis, capsulis verrucosis. *Sp. pl.* 658. *Amæn. Acad.* III. 120.

Tithymalus foliis ellipticis ferratis, stipulis umbellaribus quinis, floralibus obtuse quadrangulis, fructibus undique exasperatis. *Hall. hist.* 1052.

Tithymalus myrsinites fructu verrucæ simili: *B. pin.* 291.

Tithymalus verrucosus. *B. hist.* III. 673. *Park.* 181. *R. syn.* 312.

Anglis, warty Spurge:

Habitat in arvis. Found by Mr. Dale in Essex; and by Dr. Robinson near York. *R. syn.* ♂. VIII.

8. EUPHORBIA umbella quinquefida: trifida: dichoplatyphyltoma, involucellis carina pilosis, foliis ferratis lanceolatis, capsulis verrucosis. *Sp. pl.* 660.

Tithymalus foliis lanceolatis ferratis, stipulis floralibus

DODECANDRIA TRIGYNIA.

bus cordatis, capsulis asperis lineis levibus divisis.
Hall. hist. 1053.

Tithymalus platyphyllos. *Fuchs. hist.* 813. *B. hist.*
III. 670. *R. syn.* 312.

Tithymalus arvensis latifolius germanicus. *B. pin.*
 291.

β *Tithymalus segetum longifolius.* *Cat. Cant. app. R.*
syn. 312.

Anglis, corn Spurge.

Habitat in arvis; at Black Notley in Essex. *R. syn.*
In the corn fields between Harefield-common and Bat-
tle's-Well. Blackf. Sp. bot. prope Northfleet. ☉
 VII. VIII.

Obs. in β Folia fere integerrima, basi tantum minutissi-
ma serrulata.

**** *Umbella multifida.*

byberna. 9. *EUPHORBIA umbella sexfida: dichotoma involu-*
cellis ovalibus, foliis integerrimis, ramis nullis,
capsulis verrucosis. *Sp. pl.* 662.

Tithymalus latifolius hispanicus. *Tab. hist.* 987. *B.*
pin. 291.

Tithymalus bybernicus. *Merr. pin.* *R. syn.* 312.
vasculis muricatis erectis. *Dill. eltb.* 387. t. 290.
 f. 374.

Anglis, Irish Spurge.

Habitat in arvis; amongst the corn near Twickenham-
park, against Richmond; and near Otterspool. *Mr.*
Dood. R. syn. inter Ecyerham et Sittingborn in
Cantio. 4. VIII.

amygda- 10. *EUPHORBIA umbella multifida: dichotoma, in-*
volucellis perfoliatis orbiculatis, foliis obtusis. *Sp.*
pl. 662.

Tithymalus characias amygdaloides. *Ger. em.* 500.
B. pin. 290. *R. syn.* 312.

Tithymalus sylvaticus toto anno folia retinens. *B.*
hist. III. 671.

Tithymalus characias vulgaris. *Park.* 186.

Anglis, wood Spurge.

Habitat in sylvis et sepibus frequens. 4. IV. V.

II. EU-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Classis XII.

I C O S A N D R I A
M O N O G Y N I A.

P R U N U S. 620.

Cal. quinquefidus, inferus. *Petala* quinque. *Drupa*
nux futuris prominulis.

communis. 1. PRUNUS pedunculis subgeminis, foliis lanceolato-
ovatis convolutis, ramis submuticis.

insititia. Prunus pedunculis geminis, foliis ovatis subvillosis
convolutis, ramis spinascentibus. *Sp. pl.* 680.
Hall. hist. 1081.

Prunus sylvestris major. *B. hist. I.* 196. *R. syn.* 462.

Prunus sylvestris præcocia. *B. pin.* 444.

Prunus sylvestris fructu majore nigro. *R. syn. ed.*
II. 301.

β Prunus sylvestris fructu rubro, acerbo et ingrato. *R.*
syn. 463.

γ Prunus sylvestris fructu majore albo. *R. syn.* 462.

domestica. δ Prunus pedunculis subsolitariis, foliis lanceolato-ova-
tis convolutis, ramis muticis. *Sp. pl.* 680.

Prunus domestica. *Ger. em.* 1497.

Prunus. *B. pin.* 443.

Pruna majora dulcia et parva atro-cæruleo. *B. pin.*
443.

Anglis, common Prunus or Plumb; α. β. γ. Bullace-
tree δ. Plumb-tree.

Habitat in sepibus. h. IV.

spinosa. 2. PRUNUS pedunculis solitariis, foliis lanceolatis gla-
bris, ramis spinosis. *Sp. pl.* 681. *Fl. suec.* 432.

Prunus sylvestris. *Ger. em.* 1497. *Park.* 1033. *B.*
pin. 444. *hist. I.* 193. *R. syn.* 462.

Anglis, thorny Prunus, Black-thorn or Sloe-tree.

Habitat

Habitat in sepibus. h. IV.

Pharmac. Pruni sylvestris, Cortex, Flores, Fructus.

3. PRUNUS umbellis subpedunculatis, foliis ovato-cerasus lanceolatis conduplicatis.

Prunus umbellis subpedunculatis, foliis ovato lanceolatis conduplicatis glabris. *Sp. pl.* 679.

Cerasus foliis glabris serratis, ovato-lanceolatis, mucrone producto. *Hall. hist.* 1083.

Cerasus sylvestris fructu rubro. *B. hist.* 1220. *R. syn.* 463.

Cerasus lativa rotunda, rubra et acida. *B. pin.* 449.

Cerasus acida. *Tab. ic.* 985.

- β Cerasus sylvestris fructu minimo cordiformi. *Ph. brit. R. syn.* 463.

- γ Prunus umbellis sessilibus, foliis ovato-lanceolatis conduplicatis subtus pubescentibus. *Sp. pl.* 680.

Cerasus foliis ovato-lanceolatis, serratis, inferne subhirsutis, mucrone producto. *Hall. hist.* 1082.

Cerasus sylvestris ac major, fructu subdulci nigro colore insufficiente. *B. pin.* 450.

Cerasus sylvestris fructu nigro. *B. hist. I.* 220. *R. syn.* 463.

Cerasus nigra. *Tab. ic.* 986.

- δ Cerasus sylvestris septentrionalis fructu parvo serotino. *R. syn.* 463.

Anglis, red γ black Prunus or Cherry. *Hunt. Ev. sylv.* 188.

Habitat in sylvis et sepibus. h. V.

Pharmac. Cerasorum nigrorum Fructus.

4. PRUNUS floribus racemosis, foliis deciduis basi subtus biglandulosis. *Sp. pl.* 677.

Padus foliis ovato-lanceolatis, serratis. *Hall. hist.* 1086.

Cerasus avium nigra et racemosa. *Ger. em.* 1504. *R. syn.* 463.

Cerasus avium-racemosa. *Park.* 1517.

Cerasus racemosa sylvestris fructu non eduli. *B. pin.* 451.

Anglis, birds Prunus or Cherry.

ICOSANDRIA DIGYNIA.

Habitat in sylvis et sepibus ; in comitatibus Westmorlandico, Cumberlandico, Eboracensi, Lancastriensi, &c. frequens. In the lane going from Templemills to Epping Forest, Dr. Wilmer. h. V.

D I G Y N I A.

C R A T Æ G U S. 622.

Cal. quinquefidus. Petala quinque. Bacca infera disperma.

aria.

1. CRATÆGUS foliis ovatis incisiss ferratis subtus tomentosis. *Sp. pl.* 681. *Fl. suec.* 433. *dan.* 302.

Mespilus foliis ovatis ferratis, subtus tomentosis. *Hall. hist.* 1089.

Mespilus alni folio subtus incano, Aria Theophrasti dicta. *R. syn.* 453.

Aria Theophrasti. *Ger. em.* 1327.

Sorbus sylvestris Aria Theophrasti dicta. *Park.* 1421.

Alni effigie, lanato folio major. *B. pin.* 452.

Anglis, white Haw or Beam-tree. *Hunt. Ev. sylv.* 181.

Habitat in sylvis et sepibus præsertim montosis et solo calcareo. h. V.

torminalis.

2. CRATÆGUS foliis cordatis septangulis : lobis infimis divaricatis. *Sp. pl.* 681.

Mespilus foliis ferratis, septilobis ; lobis primis divergentibus. *Hall. hist.* 1088.

Mespilus apii folio sylvestris non spinosa, seu Sorbus torminalis. *B. pin.* 454. *R. syn.* 453.

Sorbus torminalis. *Cam. epit.* 162. *Ger. em.* 1471. seu vulgaris. *Park.* 1420.

Anglis, wild Haw-tree or Service. *Hunt. El. sylv.* 182.

Habitat in sylvis et sepibus. h. IV. V.

Oxyacantha.

3. CRATÆGUS foliis obtusis subtrifidis ferratis. *Sp. pl.* 683. *Fl. suec.* 434. *dan.* 634. floribus monogynis.

Mespilus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Habitat in montibus nemorosis. In the mountainous part of Cornwall. Walter Moyle, Esq. In the Morelands in Staffordshire. D. Plot. R. syn. h. V. Pharmac. Sorbi Fructus.

- hybrida.* 3. SORBUS foliis angulatis semipinnatisque, subtus tomentosis.
 Sorbus foliis semipinnatis subtus tomentosis. *Sp. pl.* 684. *Lin. dec. II. t. 6.*
Anglis, bastard Service-tree.
 Habitat in montosis. On the walls of Castel Dinas brân Denbighshire. Mr. Waring. h. V.

PENTAGYNIA.

P Y R U S. 626.

Cal. quinquefidus. *Petala* quinque. *Pomum* inferum quinqueloculare, polyspermum.

- communis.* 1. PYRUS foliis serratis, pedunculis corymbosis. *Sp. pl.* 686. *Fl. suec.* 436.
 Pyrus foliis serratis, pomis basi productis. *Hort. cliff.* 190.
 Pyrus foliis ovato-lanceolatis serratis glabris. *Hall. hist.* 1096.
 Pyrus sylvestris. *Ger. em.* 1457. *Park.* 1500. *B. pin.* 439.
 Pyraister s. Pyrus sylvestris. *B. hist. I.* 57. *R. syn.* 452.
 β Pyrus lativa. *R. syn.* 452.
Anglis, common Pear-tree.
 Habitat in sylvis et sepibus. h. IV. V.

- malus.* 2. PYRUS foliis serratis umbellis sessilibus. *Sp. pl.* 686.
 Pyrus foliis serratis, pomis basi concavis. *Hort. cliff.* 186.
 Pyrus foliis ovatis, acuminatis subtus hirsutis, petiolis florigeris brevissimis. *Hall. hist.* 1097.
 Malus sylvestris. *Ger. em.* 1461. *Park.* 1502. *R. syn.* 452.

Malus

Malus sylvestria. *B. pin.* 451.

β *Malus sativa.* *R. syn.* 451.

Anglis, Crab-tree or Wilding, β *Apple-tree.*

Habitat in sylvis et sepibus. *h. V.*

Pharmac. *Mali sylvestris Fructus.*

MESPILUS. 625.

Cal. quinquefidus. *Petala* quinque. *Bacca* infera,
pentasperma.

1. MESPILUS inermis, foliis lanceolatis subtus tomentosis, floribus solitariis sessilibus. *Sp. pl.* 684. *germanica.*

Mespilus foliis elliptico-lanceolatis serratis, calycibus longissimis, persistentibus. *Hall. hist.* 1094.

Mespilus. *Dod.* 801. *Ger. em.* 1453. *R. syn.* 453. *vulgaris.* *Park.* 1422.

Mespilus germanica folio laurino non serrato, s. *Mespilus sylvestris.* *B. pin.* 453.

Anglis, Medlar-tree.

Habitat in sepibus. *In all the hedges about Minchiville.*
Mr. Du Bois. *R. syn.* *h. V.*

SPIRÆA. 630.

Cal. quinquefidus. *Petala* quinque. *Caps.* polysperma.

1. SPIRÆA foliis pinnatis: foliolis uniformibus serratis, caule herbaceo, floribus cymosis. *Sp. pl.* 702. *Filipendula.*
Fl. suec. 439. *dan.* 635.

Filipendulo foliis pinnatis minimis intermissis, extrema trilobata. *Hall. hist.* 1136.

Filipendula. *Ger. em.* 1058. *Park.* 434. *R. syn.* 259. *vulgaris.* *B. pin.* 163.

Anglis, common Spiræa or Drop-wort.

Habitat in pratis et pascuis montosis, sola cretaceo. *Supra Dupper's-Hill, prope Croydon.* *D. Caute; juxta Clapham prope Londinum.* *h. VII.*

Pharmac. *Filipendulæ Radix, Folia.*

2. SPIRÆA foliis pinnatis: impari majorę lobato, floribus cymosis. *Sp. pl.* 702. *Fl. dan.* 547.

Filipendula foliis pinnatis, acute serratis, minimis intermissis,

ICOSANDRIA POLYGYNIA.

termistis, extrema trilobata maxima. *Hall. hist.*
1135.

Ulmaria. *Clus. hist. II.* 198. *B. hist. III.* 488. *R.*
syn. 259. vulgaris. *Park.* 592.

Regina prati. *Ger. em.* 1043.

Barba capri floribus compactis. *B. pin.* 164.

Anglis, meadow Spiræa or -sweet.

Habitat in pratis humidis et ad ripas fluviorum. *H.*
VI. VII.

Pharmac. Ulmariae Herba.

POLYGYNIA.

R O S A . 631.

Cal. urceolatus, quinquefidus, carnosus, collo coarctatus. *Petala* quinque. *Sem.* plurima, hispida, calycis interiori lateri affixa.

* *Germinibus globosis.*

*eglante-
ria.*

1. ROSA germinibus subglobosis petiolisque aculeatis, aculeis recurvis, foliolis ovatis subtus rubiginosis.

Rosa rubiginosa, germinibus globosis petiolisque aculeatis, aculeis recurvis, foliis subtus rubiginosis.

Lin. Mantiss. II. 564. *Fac. austr. I.* 31. t. 50.

Rosa spinis aduncis, foliis subtus rubiginosis. *Hall. hist.* 1103.

Rosa sylvestris odora. *Ger. em.* 1269. *R. syn.* 454. seu Eglenteria flore simplici. *Park. par.* 418.

Rosa sylvestris foliis odoratis. *B. pin.* 483.

Anglis, sweet Rose or Briar.

Habitat in sepibus et ericetis. *H.* VI. VII.

*spinosissi-
ma.*

2. ROSA germinibus globosis glabris, pedunculis hispidis, caule petiolisque aculeatissimis. *Syst. nat.* 348.

Sp. pl. 705. *Fl. dan.* 398.

Rosa spinis rectis confertis, foliis novenis glabris, pomis et petiolis subspinosis. *Hall. hist.* 1106.

Rosa pumila, spinosissima, foliis pimpinellæ glabris, flore albo. *B. hist. II.* 40. *R. syn.* 455.

Rosa

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Rosa sylvestris inodora f. *canina*. *Park.* 1017. *R. syn.* 454.

Rosa sylvestris vulgaris flore odorato incarnata. *B. pin.* 483.

Anglis, dog Rose or Hep-tree.

Habitat in sylvis et sepibus. h. VI.

Pharmac. Rosæ sylvestris, *Flores Spongiolæ (insectorum niduli)* f. *Bedeguar, Cynosbati.*

R U B U S. 632.

Cal. quinquefidus. *Petala* quinque. *Bacca* composita acinis monospermis.

* *Frutescentes.*

idæus.

1. RUBUS foliis quinato-pinnatis ternatisque, caule aculeato, petiolis canaliculatis. *Sp. pl.* 706.

Rubus caule spinoso, subrecto, foliis quinatis et ternatis, subtus tomentosus, fructibus hirsutis. *Hall. hist.* 1108.

Rubus idæus spinosus. *B. pin.* 470. fructu rubro, *B. hist. II.* 59. *R. syn.* 467.

Rubus idæus. *Ger. em.* 1272. *Park.* 557.

Anglis, Raspberry-bramble or Framboise.

Habitat in sylvis et sepibus, montibus saxosis et udis Walliæ, et septentrionalium Angliæ, &c. h. V. VI.

Pharmac. Rubi idæi *Fructus.*

cæsius.

2. RUBUS foliis ternatis subnudis: lateralibus bilobis, caule aculeato tereti. *Sp. pl.* 706.

Rubus caule aculeato prostrato, foliis ternatis, subtus hirsutis. *Hall. hist.* 1110.

Rubus repens fructu cæsius. *B. pin.* 479. *Ger. em.* 1271.

Rubus minor fructu cæruleo. *B. hist. II.* 59. *R. syn.* 467.

Rubus minor, chamærubus f. Humirubus. *Park.* 1013.

Anglis, small Bramble, or Dewberry-bush.

Habitat in sylvis et sepibus passim. h. VI. VII.

fruticosus. 3. RUBUS foliis quinato-digitatis ternatisque, caule petiolisque aculeatis. *Sp. pl.* 707. *Fl. succ.* 444.

Rubus.

Rubus caule spinoso serpente, foliis quinatis et ternatis, subtus subtomentosis, bacca levi. *Hall. hist.* 1109.

Rubus. *Cam. epit.* 751. *Lob. ic. II.* 211. vulgaris major. *Park.* 1013. fructu nigro. *B. hist. II.* 57. *R. syn.* 467.

Rubus vulgaris f. Rubus fructu nigro. *B. pin.* 479.

β Rubus vulgaris major, fructu albo. *R. syn.* 467.
Anglis, common Bramble.

Habitat in sylvis et sepibus. *h.* V—IX.

** *Herbacei.*

4. RUBUS foliis ternatis nudis, flagellis reptantibus saxatilis. herbaceis. *Sp. pl.* 708. *Fl. dan.* 138.

Rubus caule herbaceo, foliis ternatis, glabris, acinis baccarum distinctis, paucissimis. *Hall. hist.* 1111.

Rubus saxatilis. *Ger. em.* 1273.

Rubus alpinus saxatilis. *Park.* 1014.

Chamærubus saxatilis. *B. pin.* 479. *R. syn.* 261.

Anglis, stone Bramble.

Habitat inter lapides ad latera montium in comitatibus Eboracensi, Cumberlandico et Westmorlandico. *u.* VI.

5. RUBUS foliis simplicibus lobatis, caule inermi unichamæmorus. *Sp. pl.* 708. *Fl. suec.* 449. *dan.* 1. *morus.*

Rubus caule bifolio unifloro, foliis simplicibus. *Fl. lapp.* 208. *t.* 5. *f.* 1.

Chamæmorus. *Clus. hist. I.* 118. *Ger. em.* 1273. *R. syn.* 260. *anglica.* *Park.* 1014.

Chamærubus foliis ribes *anglica.* *B. pin.* 480.

Anglis, mountain Bramble or Cloud-berry.

Habitat in paludibus turfosis supra montes Eboracenses, Westmorlandicos, Cumberlandicos, Lancastreses, Arvonicos. *u.* V. VI.

FRAGARIA. 633.

Cal. decemfidus. *Petala* quinque. *Receptaculum* seminum ovatum, baccatum, deciduum.

1. FRAGARIA flagellis reptantibus. *Sp. pl.* 708. *vesca.*
Fragaria

ICOSANDRIA POLYGYNIA.

Fragaria foliis ternatis, flagellis reptans. Hall. hist. 1112.

Fragaria. Ger. em. 997. R. syn. 254. vulgaris. B. pin. 758. Park. 758. Fl. lapp. 209.

β *Fragaria fructu hispido. Ger. em. 998.*

γ *Fragaria fructu parvi pruni magnitudine. B. pin. 327.*

δ *Fragaria fructu albo. B. pin. 326.*

Anglis, common Strawberry.

Habitat in sylvis et ad sepes frequens. 4. V. VI.

sterilis. 2. FRAGARIA caule decumbente, ramis floriferis laxis. *Syst. nat. 350.*

Fragaria caule decumbente repente. Sp. pl. 709.

Fragaria caule prostrato, foliis ternatis retusis sericeis. Hall. hist. 1113.

Fragaria sterilis. B. pin. 327. R. syn. 254.

Fragaria minime vesca. Park. 758. f. steriles. Lob. ic. 696. Ger. em. 998.

Anglis, barren Strawberry.

Habitat in priscuis et ericetis sterilioribus. 4. IV. V.

POTENTILLA. 634.

Cal. decemfidus. Petala quinque. Sem. subrotunda, nuda, receptaculo parvo exsucco affixa.

fruticosa 1. POTENTILLA foliis pinnatis, caule fruticoso. *Sp. pl. 709. Fl. succ. 451. Kalm. it. II. 263.*

Pentaphylloides rectum fruticosum. Hist. Ox. II. 193. f. 2. t. 23. f. 5. Hortul. angl. 54. t. 54.

Pentaphylloides fruticosa. Cat. pl. angl. 228. ic. R. syn. 256.

Anglis, shrubby Cinquefoil.

Habitat ad ripam meridionalem Tese fluvii infra vicum Thorp dictum variis in locis, nec non infra cœnobium Athelstani, vulgo Eggleston-abbey, in agro Eboracensi. Mr. Lawson observed thousands of these plants by Mickle-force, in Teasdale. R. syn. 2. VI.

anferina. 2. POTENTILLA foliis pinnatis serratis, caule repente. *Sp. pl. 710. pedunculis unifloris. Syst. nat. 350. Fl. dan. 544.*

Fragaria

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Fragaria caule subrecto, foliis quinis, septenis, serratis, obtusis. *Hall. hist.* 1120.

Pentaphyllum parvum hirsutum. *B. hist.* II. 598.
R. syn. 255.

Pentaphyllum incanum minus repens. *Ger. em.* 989.

Quinquefolium minus repens lanuginosum. *Park.* 399. luteum. *B. pin.* 325.

Quinquefolium IV. flavo flore 2 species. *Clus. hist.* II. 106.

Anglis, hairy Cinquefoil.

Habitat in pascuis, circa Kippax in agro Eboracensi. *Ꝟ.* VI. VII.

verna. 6. POTENTILLA foliis radicalibus quinatis acute serratis retusis, caulinis ternatis, caule declinato. *Sp. pl.* 712. *Fl. suec.* 455.

Fragaria caule decumbente, foliis quinatis serratis subhirsutis. *Hall. hist.* 1119.

Pentaphyllum s. Quinquefolium minus. *Tabern. ic.* 123.

Quinquefolium minus repens luteum. *B. pin.* 325.

Anglis, Spring Cinquefoil.

Habitat in pratis et pascuis sterilioribus prope Preston in comitatu Lancastrensi, D. Tofield; in the King's Park, Edinburgh, Dr. Hope. prope Giggleswick in comitatu Eboracensi. *Ꝟ.* V. VI.

alba. 7. POTENTILLA foliis quinatis apice conniventiserratis, caulibus filiformibus procumbentibus, receptaculis hirsutis. *Sp. pl.* 713. *Fac. austr.* II. 10. t. 115.

Fragaria foliis ovatis apice serratis, oris sericeis, pappo placentæ longissimo. *Hall. hist.* 1123.

Quinquefolium album majus. *B. pin.* 325.

Quinquefolium sylvaticum majus flore albo. *Ger. em.* 987.

Anglis, white Cinquefoil.

Habitat in Wallia, D. Haviland. *Ꝟ.* VII. VIII.

reptans. 8. POTENTILLA foliis digitatis, caule repente, pedunculis unifloris. *Sp. pl.* 714. *Fl. dan.* 589.

Fragaria

- Fragaria foliis quinatis, ferratis, petiolis uniflori-, caule repente. *Hall. hist.* 1118.
- Pentaphyllum vulgatissimum. *Park.* 398. *R. syn.* 255.
- Quinquefolium majus. *Ger. em.* 987. repens. *B. pin.* 325.
- β Tormentilla reptans, caule repente, foliis petiolatis. *Fl. angl.* 198. *Sp. pl.* 176.
- Tormentilla reptans. *Pet. h. br. t.* 41. *f.* 10. *R. syn.* 357.
- Pentaphyllum reptans alatum foliis profundius serratis. *Plot. hist. Ox. c.* 6. *p.* 7. *t.* 9. *f.* 5.
- Pentaphyllum minus viride, flore aureo tetrapetalo, radículas in terram e geniculis demittens. *Hist. Ox.* II. 190.
- γ Pentaphyllum aureum minus, sylvaticum, nostras, foliis tripartito divisis ex cauliculis geniculis radicescens. *Pluk. alm.* 285.
- Anglis, common Cinquefoil, or Five-leaved-grass.*
Habitat in pratis et pascuis et ad vias frequens. β in sylvis et pascuis arenosis et sterilioribus passim. 4. VI.—VIII.
- Pharmac. Pentaphylli Radix.*
Obl. in β variat floribus tetrapetalis et pentapetalis in eadem planta.

TORMENTILLA. 635.

Cal. octofidus. Petala quatuor. Sem. subrotunda, nuda receptaculo parvo exsucco affixa.

1. TORMENTILLA caule erectiusculo, foliis sessili. *erecta.*
bus. Sp. pl. 716.
- Tormentilla. *Cam. epit.* 685. *Ger. em.* 992. *R. syn.* 257. vulgaris. *Park.* 394.
- Tormentilla sylvestris. *B. pin.* 326.
- Fragaria tetrapetala, foliis caulinis sessilibus, quinatis. *Hall. hist.* 1117.
- Anglis, common Tormentil, or Septfoil.*
Habitat in pascuis, et nemorosis sterilioribus. 4. VI.—VIII.
- Pharmac. Tormentillæ Radix, Folia,*

G E U M. 636.

Cal. decemfidus. *Pctala* quinque. *Sem.* arista geniculata.

- urbanum.* 1. GEUM floribus erectis, fructu globoso villosa : aristis uncinatis nudis, foliis lyratis. *Sp. pl.* 716.
 Geum foliis pinnatis, pinna ultima trilobata, floribus patulis, tubis aduncis. *Hall. hist.* 1130.
 Caryophyllata. *Ger. em.* 994. *R. syn.* 253. vulgaris.
B. pin. 321. *R. syn.* 136.
 2. Caryophyllata vulgaris majore flore. *B. pin.* 321.
Anglis, common Aven, or Herb-bennet.
 Habitat in fylvis et ad sepes frequens. 4. VI—VIII.
Pharmac. Caryophyllatæ Radix.

- rivale.* 2. GEUM floribus nutantibus, fructu oblongo : aristis plumosis tortis. *Sp. pl.* 717.
 Geum foliis pinnatis, pinnis acutis, extrema maxima trilobata : floribus conniventibus. *Hall. hist.* 1129.
 Caryophyllata montana purpurea. *Ger. em.* 995. *R. syn.* 253.
 Caryophyllata aquatica nutante flore. *B. pin.* 321.
 Caryophyllata montana s. palustris purpurea. *Park.* 136.
Anglis, water Avens.
 Habitat in pascuis montosis subhumidis, in septentrionalibus Angliæ circa Settle, Ingleton, &c. Alio in Wales, about Snowdon, &c. *R. syn.* 4. VI. VII.

D R Y A S. 637.

Cal. octofidus. *Cor.* octo. *Sem.* caudata, pilosa.

- octopetala.* 1. DRYAS octopetala, foliis simplicibus. *Sp. pl.* 718.
Fl. suec. 262. *dan.* 31.
 Dryas floribus octopetalis, foliis ovatis subtus albissimis. *Hall. hist.* 1133.
 Caryophyllata alpina chamædryos folio. *Hist. Ox.* II. 432. *R. syn.* 253.
 Chamædryos alpina, cisti flore. *B. pin.* 248.
 Chamædryos spuria montana, cisti flore. *Park.* 106.
 Teucrium

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Classis XIII.

P O L Y A N D R I A
 M O N O G Y N I A.

* *Flores tetrapetali.*

A C T Æ A. 664:

Cal. tetraphyllus. *Cor.* tetrapetala. *Bacca.* unilocularis. *Sem.* femiorbiculata.

- spicata.* I. ACTÆA racemo ovato, fructibus baccatis. *Sp. pl.* 722. *Fl. suec.* 464. *dan.* 498.
 Actæa foliis circumferratis, duplicato-pinnatis, pinnula extrema trilobata. *Hall. hist.* 1076.
 Christophoriana. *Clus. hist.* II. 86. *Ger. em.* 979. *Till. ic.* 148. *R. syn.* 262. *Weinm. t.* 380. vulgaris. *Park.* 379.
 Aconitum racemosum. *B. pin.* 182.
Anglis, Herb-christopher, or Bane-berries.
 Habitat in sylvis et nemorosis. Prope Malham-cove. *R. syn. in sylvula prope Clapham in agro Eboracensi.* 4. V. VI.

CHELIDONIUM. 647.

Cal. diphyllus. *Cor.* tetrapetala. *Siliqua* unilocularis, linearis.

- majus.* I. CHELIDONIUM pedunculis umbellatis. *Sp. pl.* 723. *Fl. dan.* 542.
 Chelidonium lobis foliorum subrotundis. *Hall. hist.* 1069.

Chelidonium

Chelidonium majus. *Fuchs. hist.* 865. *Ger. em.* 1069.
vulgare. *B. pin.* 144. *Park.* 616.

Papaver corniculatum luteum, *Chelidonia dictum.* *R. syn.* 309.

β *Chelidonium majus foliis quernis.* *B. pin.* 144.

Anglis, common Celendine.

Habitat in ruderatis. 4. V. VI.

Pbarmac. *Chelidonii majoris Radix, Herba.*

2. CHELIDONIUM pedunculis unifloris, foliis am-plexicaulibus sinuatis, caule glabro. *Sp. pl.* 724. *Glaucium.*

Glaucium foliis radicalibus semipinnatis, caulinis amplexicaulibus. *Hall. hist.* 1060.

Papaver corniculatum. *Cam. epit.* 805. luteum. *B. pin.* 171. *hist.* III. 398. *Park.* 261. *R. syn.* 309.

Papaver cornutum flore luteo. *Ger. em.* 367.

Anglis, sea Celendine, or yellow Horned-poppy.

Habitat in littoribus maritimis arenosis. ☉. VIII.

3. CHELIDONIUM pedunculis unifloris, foliis sessilibus pinnatifidis, caule hispido. *Sp. pl.* 724. *corniculatum.*

Papaver corniculatum phœniceum hirsutum. *B. pin.* 171.

Papaver cornutum flore rubro. *Ger. em.* 367.

Anglis, red Celendine, or Horned-poppy.

Habitat in arvis arenosis. *D. Stillingfleet ex Norfolkia cum sequente misit.* ☉. VII. VIII.

4. CHELIDONIUM pedunculis unifloris, foliis pinnatifidis linearibus, caule lævi, siliquis trivalvibus. *hybridum.*

Papaver corniculatum violaceum. *B. pin.* 172. *hist.* III. 393. *R. syn.* 309.

Papaver cornutum flore violaceo. *Ger. em.* 367.

Papaver corniculatum flore violaceo. *Park.* 262.

Anglis, violet Celendine, or Horned-poppy.

Habitat in arvis arenosis; in agro Cantabrigiensi inter segetes inter Swafham et Burwel vicos. *R. syn.* ☉. VII. VIII.

P A P A V E R. 648.

Cal. diphyllus. *Cor.* tetrapetala. *Caps.* unilocularis,
sub stigmatate persistente poris dehiscens.

* *Capsulis hispidis.*

bybridum. 1. PAPAVER capsulis subglobosis torosis hispidis, caule folioso multifloro. *Sp. pl.* 725.

Papaver foliis hispidis, pinnatis, pinnis lobatis, capsulis ellipticis, hispidis. *Hall. hist.* 1061.

Papaver laciniato folio, capitulo hispido rotundiore. *R. syn.* 308.

Argemone capitulo torulo. *Ger. em.* 372.

Argemone capitulo brevior. *B. pin.* 172.

Argemone capitulo rotundiore. *Park.* 369.

Anglis, bastard Poppy.

Habitat in arvis passim. ☉. VI. VII.

Argemone. 2. PAPAVER capsulis clavatis hispidis, caule folioso multifloro. *Sp. pl.* 725. *Fl. suec.* 466.

Papaver laciniato folio capitulo hispido longiore. *R. syn.* 308.

Argemone capitulo longiore. *Lob. ic.* 276. *Ger. em.* 273. *B. pin.* 172. *Park.* 370.

Anglis, rough Poppy.

Habitat in arvis passim. ☉. VI. VII.

** *Capsulis glabris.*

Rhoeas. 3. PAPAVER capsulis glabris globosis, caule piloso multifloro, foliis pinnatifidis incis. *Sp. pl.* 726.

Papaver foliis semipinnatis, hispidis, fructu ovato glabro. *Hall. hist.* 1064.

Papaver laciniato folio, capitulo brevior glabro, annum Rhoeas dictum. *R. syn.* 308.

Papaver rhoeas. *Ger. em.* 371.

Papaver erraticum rhoeas sive sylvestre. *Park.* 367.

Papaver erraticum majus. *B. pin.* 171.

Anglis, corn, or red Poppy.

Habitat inter segetes. ☉. VI. VII.

Pharmac. Papaveris erratici Flores.

dubium. 4. PAPAVER capsulis oblongis glabris, caule multifloro

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

- Tilia foemina. *Ger. em.* 1483. major. *Park.* 1407.
 Tilia foeminá folio majore. *B. pin.* 426.
 β Tilia foemina folio minore. *B. pin.* 426.
 γ Tilia foemina foliis molliter hirsutis, viminibus rubris,
 fructu tetragono. *R. syn.* 473.
 δ Tilia ulmifolia, semine hexagono. *Merr. pin.*
Anglis, the Lime or Linden-tree. Hunt. Ev. sylv. 201.
 • Habitat in sylvis, sepibus et areis. *h.* VII.

C I S T U S. 673.

Cal. pentaphyllus : foliolis duobus minoribus. *Cor.*
 pentapetala. *Caps.* triquetra, trivalvis.

* *Exstipulati herbacei.*

- guttatus.* 1. CISTUS herbaceus exstipulatus, foliis oppositis lan-
 ceolatis, trinerviis, racemis ebracteatis. *Sp. pl.* 741.
 Cistus flore pallido, punicante macula insignito. *B.*
pin. 465. *R. syn.* 343.
 Cistus annuus flore maculato. *Ger. em.* 1281.
 Cistus annuus flore guttato. *Park.* 661.
 Helianthemum flore maculoso. *Col. ecphr.* II. 78. t.
 77.
Anglis, annual Cistus.
 Habitat in pratis arenosis, in monte Llech ddû prope Ho-
 lyhead in insula Mona. *D. Brewer.* ☉. VI.

** *Exstipulati suffruticosi.*

- hirsutus.* 2. CISTUS suffruticosus procumbens exstipulatus, foliis
 oppositis oblongis subrevolutis pilosis, subtus inca-
 nis, floribus racemosis. *Fl. angl.* 206. *Lin. mantif.*
 II. 245.
 Cistus caulibus duris, procumbentibus, foliis ellipticis
 hirsutis, inferne tomentosus, petiolis confertis uni-
 floris, calycibus tomentosus. *Hall. hist.* 1035.
 Helianthemum alpinum folio Pilosellæ minoris Fuch-
 sii. *B. hist.* II. 18. *R. syn.* 342. *Segu. ver.* III,
 196.
Anglis, hairy Cistus.
 Habitat in pratis et rupibus montosis; uno circiter milli-
 aria Kendalia, loco saxoso Betham-bank dicto occiden-
 tem

tem versus copiose, D. Richardson. At Buckbarrow-bank-scar, between Bridgsteer and Conzick; also on the rocks about Cartmel-Wells in Lancashire. Mr. Fitz-Roberts. R. syn. 4. V.

*** *Stipulati herbacei.*

3. CISTUS herbaceus patulus villosus stipulatus, flori-*salicifoli-*bus racemosis erectis: pedicellis horizontalibus. *us.*

Sp. pl. 742.

Cistus folio salicis. *B. pin. 465.*

Cistus annuus. *Ger. em. 1280. folio salicis. Park. 661.*

Helianthemum annum humile, foliis ovatis, flore fugaci. *Seg. ver. III. 397. t. 6. f. 3.*

Anglis, willow-leaved Cistus.

Habitat in pratis et pascuis arenosis. juxta Brean-downs in agro Somersettensi. *©. VI. VII.*

Obs. Petala lutea calyce minora fugacissima; variat floribus apetalis.

**** *Stipulati suffruticosi.*

4. CISTUS suffruticosus procumbens, stipulis lanceola-*Helian-*tis, foliis oblongis revolutis subpilosis. *Sp. pl. 744. themum;* *Fl. dan. 101.*

Cistus foliis conjugatis ellipticis, hirsutis, integerrimis, petiolis unifloris subhirsutis. *Hall. hist. 1033.*

Helianthemum vulgare. *Park. 656. R. syn. 341.*

Helianthemum anglicum luteum. *Ger. em. 1282.*

Chamæcistus vulgaris flore luteo. *B. pin. 465.*

- β Helianthemum foliis majoribus, flore albo. *B. hist. II. 16.*

γ Helianthemum ampliore folio, flore roseo. *D. Sher-rard.*

- δ Cistus suffruticosus procumbens stipulatus, foliis ova-*furreja-*to-oblongis subpilosis, petalis lanceolatis. *Sp. pl. nus. 743.*

Helianthemum vulgare petalis florum perangustis.

R. syn. 341. Dill. elth. 177. t. 145. f. 174.

Anglis, dwarf Cistus, or little Sun-flower.

Habitat in pratis et pascuis montosis, præsertim solo calca-*reo. 4. VI—VIII.*

- polifolius.* 6. CISTUS suffruticosus procumbens stipulatus, foliis oblongo-ovatis revolutis incanis, calycibus lævibus, petalis ferratis. *Fl. angl.* 205. *Sp. pl.* 745.
 Cistus foliis crassis tomentosiss ellipticis, linea exaratis, *Hall. hist.* 1036.
 Cistus humilis alpinus durior, polii nostratis folio candidante. *Pluk. ph.* 23. f. 6.
 Chamæcistus montanus polii folio. *R. syn.* 342.
 Helianthemum montanum, polii folio incano, flore candido. *Dill. elth.* 175. t. 145. f. 172.
Anglis, mountain Cistus.
 Habitat in pratis montosis. On Brean-downs, in Somersetshire, near the Severn-sea. D. Plukenet. *R. syn.* 4. VI.

*** Flores polypetali.

NYMPHÆA. 653.

Cal. tetra f. pentaphyllus. *Cor.* polypetala. *Bacca* multilocularis, truncata.

- lutea.* 1. NYMPHÆA foliis cordatis integerrimis, calyce pentaphyllo majore. *Sp. pl.* 729. *Fl. dan.* 603.
 Nymphæa calyce pentaphyllo, petalis majori, fructu conico. *Hall. hist.* 1066.
 Nymphæa lutea. *Ger. em.* 819. *R. syn.* 368.
 Nymphæa major lutea. *B. pin.* 193. *Park.* 1252.
Anglis, yellow Water-lilly.
 Habitat in fluviis, stagnis, fossis. 4. VII. VIII.
- alba.* 2. NYMPHÆA foliis cordatis integerrimis, calyce quadrifido. *Sp. pl.* 729. *Fl. dan.* 602.
 Nymphæa calyce quadrifido, petalis minori, fructu spherico. *Hall. hist.* 1067.
 Nymphæa alba. *Cam. em.* 634. *Ger. em.* 819. *R. syn.* 368. major. *B. pin.* 193. vulgaris. *Park.* 1251.
Anglis, white Water-lilly.
 Habitat in fluviis et stagnis. 4. VII. VIII.

TRIGYNIA.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

POLYANDRIA POLYGYNIA.

Anglis, mountain Columbine.

Habitat in sylvis montosis in comitatu Westmorlandico.
4. VI.

Obs. nimis affinis præcedenti, an distincta.

H E X A G Y N I A.

S T R A T I O T E S. 687.

Spatha diphylla. Perianthium trifidum. Petala tria.
Bacca sexlocularis, infera.

Aloides.

1. STRATIOTES foliis ensiformi-triangularis, ciliato-
 aculeatis. *Sp. pl. 754. Fl. dan. 337.*

Stratiotes foliis aloes, semine longo. Gundesb. ap.
Johren. R. syn. 290.

Stratiotes five militaris aizoides. Park. 1249.

Militaris aizoides. Ger. em. 825.

Aloe palustris. B. pin. 286.

Anglis, common Water-aloe, or fresh-water Soldier.

Habitat in fluentibus pigrioribus et palustribus fossis, in
Insula Eliensi copiose. R. syn. 4. VI. VII.

P O L Y G Y N I A.

A N E M O N E. 694.

Cal. o. Petala sex-novem. Sem. plura.

nemorosa. 1. ANEMONE feminibus acutis, foliolis incis, caule
 unifloro. *Sp. pl. 762. Fl. dan. 514.*

Anemone feminibus nudis, caule unifloro, foliis radi-
calibus nullis, caulinis quinquepartitis, lobis tripar-
titis, incis. Hall. hist. 1154.

Anemone nemorosa, flore majore. B. pin. 176.

Anemone nemorum album. Ger. em. 383. R. syn.
259.

Ranunculus nemorosus albus simplex. Park. 325.

Anglis, wood Anemone.

Habitat in sylvis et ad sepes. 4. IV.

2. ANEMONE

2. ANEMONE feminibus acutis, foliolis incisis, petalis *apennina*. lanceolatis numerosis. *Sp. pl.* 762.
 Anemone geranii robertiani folio, cærulea. *B. pin.* 174.
 Ranunculus nemorosus flore purpureo cæruleo. *Park.* 325. *R. syn.* 259.
 Ranunculus nemorosus, flore cæruleo, foliis majoribus apennini montis. *Mentz. pug. t.* 8.
Anglis, mountain Anemone.
 Habitat in sylvis et nemorosis. In Wimbleton woods, Mr. Rand; near Harrow-on-the-hill. Mr. Dubois; in a wood near Luton-hoe, in Bedfordshire, Mr. T. Knowlton. *R. syn.* 4. IV.
3. ANEMONE feminibus acutis, foliolis incisis, petalis *ranuncu-* subrotundis, caule sub-bifloro. *Sp. pl.* 762. *Fl. loides.* *dan.* 140.
 Anemone biflora, feminibus nudis, foliis radicalibus nullis, caulinis quinquepartitis, lobis tripartitis, incisis. *Hall. hist.* 1153.
 Ranunculus nemorosus luteus. *B. pin.* 178.
 Anemone nemorosum luteum. *Ger. em.* 383.
Anglis, yellow Anemone.
 Habitat in nemorosis et ad sepes. Prope Kings-Langley in comitatu Hertfordiensi; juxta Wrotham in Cantia. 4. V.
4. ANEMONE pedunculo involucrato, petalis rectis, *Pulsatil-* foliis bipinnatis. *Sp. pl.* 758. *Fl. suec.* 481. *dan. la.* 152.
 Anemone tubis caudatis, involucris unifoliis multifidis, foliis linearibus, hirsutis, duplicato-pinnatis, pinnis tripartitis et quadripartitis. *Hall. hist.* 1146.
 Pulsatilla folio crassiore et majore flore. *B. pin.* 177. *R. syn.* 260.
 Pulsatilla vulgaris. *Ger. em.* 385. *Park.* 341.
Anglis, Pasque-flower Anemone.
 Habitat in pratis montosis; on Gogmagog-hills; on Berhack-heath, near Stamford; in pascuis montosis et siccis juxta Leadstone-hall, supra lacum loco the Close dicto abunde prope Pontemfractum. D. Richardson.

POLYANDRIA POLYGYNIA.

chardson. *R. syn. in* Cornbury park, near Charlbury, Oxfordshire. Blacks. spec. bot. 24. IV.

CLEMATIS. 696.

Cal. 0. Petala quatuor. Sem. caudata.

- Vitalba.* 1. **CLEMATIS** foliis pinnatis: foliolis cordatis, scandentibus. *Sp. pl.* 766.
Clematis sylvestris latifolia. B. pin. 300. f. *Viorna. Park.* 380.
Clematis latifolia seu Atragene quibusdam. B. hist. II. 125. *R. syn.* 258.
Viorna. Ger. em. 886.
Anglis, wild Climber, or Travellers-joy.
Habitat in sepibus et nemorosis, solo calcareo. 2. VII.
 VIII.

THALICTRUM. 697.

Cal. 0. Petala quatuor s. quinque. Sem. ecaudata.

- alpinum.* 1. **THALICTRUM** caule simplicissimo subnudo, racemo simplici terminali. *Sp. pl.* 767. *Fl. dan.* 11.
Thalictrum montanum minimum præcox, foliis splendentibus. Hist. Ox. III. 325. f. 9. t. 20. f. 14.
Thalictrum minimum montanum atro-rubens, foliis splendentibus. R. syn. 204. *Boerb. ind. alt.* 44. ic.
Anglis, mountain Meadow-rue.
Habitat in udis scopulis et ad rivulos in alpibus in Arvoniam planta est vulgatissima. D. Lhwyd. R. syn. in boreali parte Scotiæ frequens. 2. VI.
- minus.* 2. **THALICTRUM** foliis sexpartitis, floribus cernuis. *Sp. pl.* 769. *Fl. dan.* 244.
Thalictrum foliis patulis, imis obtuse trilobis, supremis acutis, paniculis diffusis, floribus nutantibus. Hall. hist. 1139.
Thalictrum minus. Ger. em. 1251. *Park.* 264. *B. pin.* 337. *R. syn.* 203.
 3 *Thalictrum montanum minus, foliis latioribus. R. syn.* 204.
Anglis, lesser Meadow-rue.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Ranunculus caule declinato, foliis elliptico-lanceolatis subserratis. *Hall. hist.* 1182.

Ranunculus flammeus minor. *Ger. em.* 961. *R. syn.* 250.

Ranunculus palustris flammeus minor sive angustifolius. *Park.* 1214.

Ranunculus longifolius palustris minor. *B. pin.* 180.

β Ranunculus flammeus serratus. *Ger. em.* 961. *Park.* 1214.

γ Ranunculus flammeus, latiori plantaginis folio, marginibus pilosis. *Pluk. alm.* 312.

Anglis, procumbent Crowfoot or Spearwort.

Habitat in paludosis ubique. 4. VI.—IX.

lingua.

2. RANUNCULUS foliis lanceolatis, caule erecto. *Sp. pl.* 773. *Fl. suec.* 493.

Ranunculus erectus, ramosus, foliis ensiformibus subserratis. *Hall. hist.* 1181.

Ranunculus flammeus major. *Ger. em.* 961. *R. syn.* 250.

Ranunculus palustris flammeus major. *Park.* 1215.

Ranunculus longifolius palustris major. *B. pin.* 180.

Anglis, upright Crowfoot or Spearwort.

Habitat in pascuis udis et ad ripas lacuum. *In the bogs on Iwer-heath and near Uxbridge.* *Blackf. Spec. bot. Inter Rotherhithe et Deptford.* 4. VI. VII.

•• *Foliis dissectis et divisis.*

repens.

3. RANUNCULUS calycibus patulis, pedunculis sulcatis, sarmentis repentibus, foliis compositis. *Sp. pl.* 779.

Ranunculus caule repente radicato, foliis semitrilobatis, lobis petiolatis. *Hall. hist.* 1173.

Ranunculus pratensis. *Ger. em.* 951. *repens.* *Park.* 329. *R. syn.* 247. *hirsutus.* *B. pin.* 179.

β Ranunculus repens major. *Pet. h. br. t.* 38. *f.* 8.

Anglis, creeping Crowfoot.

Habitat in pratis, pascuis, et ruderatis. 4. VI.—VIII.

RANUN-

4. RANUNCULUS calycibus retroflexis, pedunculis *bulbosus*.
 fulcatis, caule erecto, foliis compositis. *Sp. pl.*
 778. *Fl. suec.* 504. *dan.* 552.
 Ranunculus radice subrotunda, foliis hirsutis semi-
 septilobis petiolatis. *Hall. hist.* 1174.
 Ranunculus bulbosus. *Ger. em.* 953. *Park.* 329. *R.*
syn. 247.
 Ranunculus pratensis radice verticilli modo rotunda.
B. pin. 179.
 Ranunculus rectus foliis pallidioribus hirsutus. *B. hist.*
III. 417. *R. syn.* 247.
Anglis, bulbous Crowfoot or Butter-cups.
 Habitat in pratis et pascuis. 4. V.
Pharmac. Ranunculi bulbosi *Herba*.
5. RANUNCULUS calycibus patulis, pedunculis te-*acris*
 retibus, foliis tripartito-multifidis: summis lineari-
 bus. *Sp. pl.* 779.
 Ranunculus foliis hirsutis, semitrilobatis, lobis late-
 ralibus bipartitis, foliis caulinis semitrilobis. *Hall.*
hist. 1169.
 Ranunculus pratensis erectus acris. *B. pin.* 178.
R. syn. 248. vulgaris. *Park.* 329.
 Ranunculus surrectis cauliculis. *Ger. em.* 951.
Anglis, upright Crowfoot.
 Habitat in pratis et pascuis. 4. VI. VII.
 Obs. variat in summis alpibus, in Arvoniam et West-
 morlandiam, caule uni aut bifloro, magno, et calycem
 valde hirsuto.
6. RANUNCULUS foliis inferioribus palmatis; sum-*sceleratus*
 mis digitatis, fructibus oblongis. *Sp. pl.* 776. *Fl.*
dan. 371.
 Ranunculus foliis levibus, semitrilobatis, rotunde ser-
 ratis, fructu ovato. *Hall. hist.* 1175.
 Ranunculus palustris rotundifolius. *Ger. em.* 962:
R. syn. 249:
 Ranunculus palustris apii folio lævis. *B. pin.* 180.
 Ranunculus palustris sardonius lævis. *Park.* 1215.
Anglis, marsh Crowfoot.

R

Habitat

Habitat in aquosis ubique. ☉. V. VI.
Pharmac. Ranunculi palustris *Herba.*

aurico-
mus.

7. RANUNCULUS foliis radicalibus reniformibus crenatis incisus; caulinis digitatis linearibus, caule multifloro. *Sp. pl.* 775.

Ranunculus foliis radicalibus integris et semitrilobatis, rotundè crenatis, caulinis multipartitis, linearibus, integerrimis. *Hall. hist.* 1177.

Ranunculus auricomus. *Ger. em.* 954.

Ranunculus nemorosus dulcis, II. Tragi. *Park.* 326.
R. syn. 248.

Ranunculus nemorosus vel sylvaticus folio rotundo.
B. pin. 178.

Anglis, wood Crowfoot or Goldilocks.

Habitat in sylvis, nemorosis, et ad scpes. ☉. IV. V.

parviflo-
rus.

8. RANUNCULUS seminibus muricatis, foliis simplicibus laciniatis acutis hirsutis, caule prostrato. *Sp. pl.* 780.

Ranunculus hirsutus annuus, flore minimo. *R. syn.* 248. t. 12. f. 1. *Pluk. ph. t.* 55. f. 1.

Ranunculus arvensis annuus hirsutus, floriflore omnium minimo. *Hist. Ox. II.* 440. f. 4. t. 28. f. 21.

Anglis, small-flowered Crowfoot.

Habitat in arvis et pratis solo glareoso; prope Greenstreet Green, juxta Dartford in comitatu Cantiano copiose. ☉. V. VI.

arvensis.

9. RANUNCULUS feminibus aculeatis, foliis superioribus decompositis linearibus. *Sp. pl.* 780. *Fl. dan.* 216.

Ranunculus feminibus aculeatis, foliis tripartitis, lobis longe petiolatis, bipartitis et tripartitis, acute incisus. *Hall. hist.* 1176.

Ranunculus arvorum. *Ger. em.* 951. *Park.* 328. *R. syn.* 249.

Ranunculus arvensis echinatus. *B. pin.* 179.

Anglis, corn Crowfoot.

Habitat in arvis. ☉. VI.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

F I C A R I A.

CAL. Perianthium triphyllum: foliolis ovatis, concavis, caducis.

COR. Petala circiter octo, lanceolata, nitida. Nectarium fovea in singulo petalo supra unguem squamula emarginata clausum.

STAM. Filamenta plurima brevissima. Antheræ rectæ, oblongæ, didymæ.

PIST. Germina numerosa, in capitulum collecta, styli nulli. Stigmata reflexa minima.

PER. nullum. Receptaculum pedunculis minutissimis semina adnectit.

SEM. plurima, oblonga, apice reflexo.

verna.

- I. FICARIA. *Dill. cat. giff. 39. Hall. hist. 1160. Ranunculus foliis cordatis angulatis petiolatis. Sp. pl. 774 caule unifloro. Syst. nat. 378. Fl. dan. 499. Chelidonium minus. Ger. em. 816. Park. 617. R. syn. 246.*

Chelidonia rotundifolia minor. B. pin. 309. Anglis, common Pilewort or lesser Celendine. Habitat in pratis et pascuis. 4. IV. Pharmac. Chelidonii minoris Radix, Herba.

T R O L L I U S. 700.

Cal. o. Petala circiter quatuordecim. Caps. plurimæ, ovatæ, monospermæ.

- europæus. I. TROLLIUS corollis conniventibus, nectariis longitudine staminum. Sp. pl. 782. Hall. hist. 1189. Fl. dan. 133.*

Ranunculus globosus. Ger. em. 955. Park. par. 218. R. syn. 272.

Ranunculus montanus, aconiti folio, flore globoso. B. pin. 182.

Anglis, Globe-flower or Locker-gowlons.

Habitat ad latera montium, inque pratis montosis in septentrionalibus Angliæ et Cambriæ copiosissime. 4. V. VI.

CALTHA.

CALTHA. 703.

Cal. o. *Petala* quinque. *Nectaria.* o. *Caps.* plures, polyspermæ.

1. CALTHA, *Sp. pl.* 784. *Fl. suec.* 511. *Hall. hist. palustris.* 1188.

Caltha palustris. *B. hist.* III. 419. flore simplici, *pin.* 276.

Caltha palustris vulgaris simplex. *Park.* 1213.

Caltha palustris major. *Ger. em.* 817.

Populago. *Tab. ic.* 753. *R. syn.* 272.

Anglis, Marsh-marigold.

Habitat in pratis et pascuis humidis et ad ripas fluviorum. 4. IV. V.

HELLEBORUS. 702.

Cal. o. *Petala* quinque s. plura. *Nectaria* bilabiata tubulata. *Caps.* polyspermæ, erectiusculæ.

1. HELLEBORUS caule multifloro folioso, foliis pe-fœtidus. datis. *Sp. pl.* 784.

Helleborus ramosus, multiflorus, foliis multipartitis ferratis, stipulis ovato lanceolatis coloratis. *Hall. hist.* 1193.

Helleborus niger fœtidus. *B. pin.* 185.

Helleboraster maximus. *Ger. em.* 976. *R. syn.* 271. f. *Confiligo.* *Park.* 212,

Anglis, stinking Hellebore, Bearsfoot or Setterwort.

Habitat in pratis, nemorosis et ad sepes. Near Cherryhinton, Cambridgeshire; upon the downs in Suffex, towards Chichester. *D. Dillenius.* *R. syn. in sylvis inter Gounsbery et Blackwell in agro Somersetiensi copiose.* 4. IV.

2. HELLEBORUS caule multifloro folioso, foliis digi-viridis. tatis. *Sp. pl.* 784. *Jac. austr.* II. 4. t. 106.

Helleborus foliis multipartitis, ferratis, caule paucifloro. *Hall. hist.* 1192.

Helleborus niger hortensis flore viridi. *B. pin.* 185. *R. syn.* 271.

POLYANDRIA POLYGYNIA.

Helleboraster minor, flore viridente. *Park.* 212.

Helleborastrum. *Ger. em.* 976.

Anglis, green Hellebore.

Habitat in sylvis. In Bigwyn closes, near Cambridge; about Arundel-castle, Suffex; in Sir Nath. D'Aeth's woods at Knowlton, East-Kent; in Stocken-church woods Oxfordshire. R. syn. circa Ditton et Whitwell, in comitatu Cantabrigiensi. 4. III. IV.

Classis

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

DIDYNAMIA GYMNOSPERMIA.

Bugula foliis imis linearibus, caulinis tripartitis. *Hall. hist.* 284.

Chamæpitys vulgaris. *Park.* 283. *R. syn.* 244.

Chamæpitys mas. *Ger. em.* 525.

Chamæpitys lutea vulgaris f. folio trifido. *B. pin.* 249.
Anglis, ground-pine Germander.

Habitat in agris restibilibus. On the lays about the borders of Triplow-heath in Cambridgeshire; about Rochester, Dartford, &c. *R. syn.* ☉. VI. VII.
Pharmac. Chamæpityos Herba.

scorodonia.

4. TEUCRIUM foliis cordatis serratis petiolatis, racemis lateralibus secundis, caule erecto. *Sp. pl.* 789.
Fl. dan. 485.

Chamædryas foliis cordatis productis, spicis longissimis nudis heteromallis. *Hall. hist.* 287.

Scorodonia f. Salvia agrestis. *Ger. em.* 662. *R. syn.* 245.

Scorodonia f. Scordium alterum quibusdam et Salvia agrestis. *Park.* III.

Scordium alterum f. Salvia agrestis. *B. pin.* 247.

Anglis, wood Germander or Sage.

Habitat in sylvis, dumetis et ericetis frequens. 4. VII.
Pharmac. Scorodiniæ Herba.

A J U G A. 705.

Corollæ labium superius minimum. Stamina labio superiore longiora.

reptans. 1. AJUGA glabra, stolonibus reptantibus. *Syst. n. Sp. pl.* 785.

Bugula flagellifera, foliis ovatis dentatis. *Hall. nomencl.* 282. *hist.* 282.

Bugula. *Dod. p.* 135. *Ger. em.* 631. *R. syn.* 245.
Riv. t. 75. vulgaris. *Park.* 525.

Consolida media pratensis cærulea. *B. pin.* 260.
Anglis, common Bugle.

Habitat in pratis et pascuis humidis. 4. V.

pyramidalis.

2. AJUGA tetragono-pyramidalis. *Sp. pl.* 785. villosa, foliis radicalibus maximis. *Syst. nat.* 387. *Fl. dan.* 185.

Bugula

Bugula foliis angulosis hirsutis calycibus villosis.
Hall. hist. 283.

Bugula cærulea alpina. *Park.* 525. *Pluk. ph. t.* 18.
f. 3. *R. syn.* 245.

Consolida media cærulea alpina. *B. pin.* 260.
Anglis, mountain Bugle.

Habitat in montibus; on Carnedh-lhewllyn in Carnar-
vonshire; Dr. Johnson. *R. syn. supra montem Ben-*
Nevis dictum in Scotia. D. Hope. 4. VI,

N E P E T A. 710.

Corollæ labium inferius lacinula intermedia crenata :
faux margine reflexo. *Stamina* approximata.

†. NEPETA floribus spicatis, verticillis subpedicellatis, *cataria*,
foliis petiolatis cordatis dentato-serratis. *Sp. pl.*
796. *Fl. suec.* 514. *dan.* 580.

Nepeta major vulgaris. *Park.* 38. *R. syn.* 237.

Cataria foliis cordatis petiolatis, verticillis spicatis,
Hall. hist. 246.

Mentha cataria vulgaris et major. *B. pin.* 228.

Mantha felina s. cataria. *Tabern.* 348. *Ger. em.* 682.
Anglis, common Nep or Cat-mint.

Habitat in pascuis et sepibus solo calcareo. 4. VII.

V E R B E N A. 32.

Cor. infundibuliformis subæqualis, curva. *Calycis*
unico dente truncato. *Sem.* quatuor.

†. VERBENA spicis filiformibus paniculatis, foliis *officinalis*,
multifido-laciniatis, caule solitario. *Sp. pl.* 29.
Fl. suec. 30. *dan.* 628.

Verbena foliis tripartitis rugosis, spicis nudis graci-
lissimis. *Hall. hist.* 219.

Verbena vulgaris. *B. hist.* III. 443. *R. syn.* 236.

Verbena mas s. recta et vulgaris. *Park.* 674.

Verbena communis. *Ger. em.* 664. cæruleo flore.
B. pin. 269.

Anglis, common Vervain.

Habitat in arvis et ad vias publicas. ♂. VI. VII.

Pharmac. Verbenæ Herba.

MENTHA.

M E N T H A. 713.

Corolla subæqualis, quadrifida: lacinia latiore emarginata. *Stamina* erecta distantia.

* *Spicatae.*

- viridis.* 1. MENTHA spicis oblongis, foliis lanceolatis nudis ferratis sessilibus, staminibus corolla longioribus, *Sp. pl.* 804.
 Mentha foliis elliptico-lanceolatis, spicis cylindricis glaberrimis. *Hall. hist.* 229.
 Mentha. *Cam. epit.* 477. angustifolia spicata. *B. pin.* 227.
 Mentha romana. *Ger. em.* 680. angustifolia s. cardiaca. *Park.* 31.
Anglis, spear Mint.
 Habitat in aquosis et ad ripas fluviorum; prope Exmouth in Devoniam et ad ripas Thamesis. 4. VII. VIII.
Pharmac. Menthæ vulgaris Herba.

- villosa.* 2. MENTHA spica oblonga, foliis sessilibus ovato-lanceolatis ferratis acutis villosis, staminibus corollam brevioribus.
 Mentha foliis elliptico-lanceolatis, subtus tomentosis spicis cylindricis, staminibus floris tubo æqualibus. *Hall. hist.* 228?
 Mentha angustifolia spicata, glabra, folio rugosiore odore graviore. *R. syn.* 233.
 Mentha spicata nostras, cardiacæ sativæ forma et odore æmula, folio rugosiore. *Pluk. mant.* 129.
 β Mentha spicata angustifolia glabra, spica latiore. *R. syn.* 233.
 γ Mentha spicata glabra latiore folio. *R. syn.* 234.
 δ Mentha sylvestris candicans, odore sativi. *Doddy, Rand. Hort. Chelf.* 129.
Anglis, hairy Mint.
 Habitat in aquosis et ad ripas fluviorum; circa Bocking in Essex, Maidstone in agro Cantiano. 4. VIII.
 Obs. variat staminibus corolla brevioribus et æqualibus foliis lanceolatis ovato-lanceolatis et ovatis, villosis et in δ tomentosis, affinis præcedenti, an distincta sit species?

- sylvestris.* 3. MENTHA spicis oblongis, foliis oblongis tomentosis ferratis sessilibus, staminibus corolla longioribus. *Sp. pl.* 804.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

DIDYNAMIA GYMNOSPERMIA.

Habitat in aquosis et ad rivulos; inter Newington et Hornsey, D. Alchorne. 4. VIII.

Pharmac. Menthæ piperitæ Herba.

Obs. variat floribus verticillato-spicatis et foliis fere nudis in hortis.

aquatica. 6. MENTHA spicis capitatis, foliis ovatis serratis petiolatis, staminibus corolla longioribus. *Sp. pl.* 805.

Mentha foliis ovato-lanceolatis serratis, verticillis paucissimis terminante maximo. *Hall. hist.* 225.

Mentha aquatica s. Sisymbrium. *Ger. em.* 684. *B. hist. III.* 223. *R. syn.* 233.

Mentha rotundifolia palustris s. aquatica major. *B. pin.* 227. *Hist. Ox. III.* 270. t. 7. f. 6.

Mentha aquatica rubra. *Park.* 1243.

hirsuta. β Mentha floribus capitatis, foliis ovatis serratis subsessilibus pubescentibus staminibus corolla longioribus. *Syst. nat.* 393. *Mantiff. I.* 88.

Mentha floribus capitatis globosis, foliis ovatis serratis, staminibus corolla brevioribus. *Fl. angl.* 223.

Mentha aquatica sive Sisymbrium hirsutius. *B. hist. III.* 2. 224.

Sisymbrium hirsutum. *R. syn.* 233.

γ Mentha sisymbrium dicta hirsuta glomerulis ac foliis minoribus ac rotundioribus. *R. syn.* 233. t. 10. f. 1.

Anglis, water Mint.

Habitat in aquosis et ad ripas fluviorum. 4. VIII.

Pharmac. Menthæ aquaticæ s. Sisymbrii Herba.

Obs. variat foliis pilosis et hirsutis; staminibus corolla brevioribus, æquantibus et longioribus.

*** Verticillatæ.

rubra. 7. MENTHA floribus verticillatis, caulibus diffusis, foliis subsessilibus ovato-lanceolatis serratis acutis subnudis.

Mentha floribus verticillatis, foliis glabris, ovato-lanceolatis, verticillis subhirsutis. *Hall. hist.* 224.

gentilis. Mentha floribus verticillatis, foliis ovatis acutis serratis, staminibus corolla brevioribus. *Sp. pl.* 805.

Mentha

Mentha hortensis verticillata ocymi odore. *B. pin.*
227.

Mentha sativa rubra. *Ger. em.* 680.

Mentha fusca sive vulgaris. *Park.* 31. *R. syn.* 232.

β *Mentha aquatica exigua.* *Trag. lib.* 1. c. 6. ic. *R.*
syn. 232.

Mentha hortensis IV. *Fuchs. hist.* 291.

Anglis, red Mint.

Habitat in aquosis et ad rivulos. In the hop-ground at
Bocking, Essex. Mr. Dale. *R. syn.* 4. VII.
VIII.

Obf. *Caulibus diffusis, rubris glabris; variat foliis nu-*
dis et parum pilosis; staminibus corolla brevioribus;
æquantibus et parum longioribus.

8. MENTHA floribus verticillatis, caulibus diffusis, fo- *arvensis*
liis petiolatis ovatis ferratis villosis.

Mentha floribus verticillatis, foliis ovatis acutis fer-
ratis, staminibus corolla æquantibus. *Sp. pl.* 806.
Fl. dan. 512.

Mentha floribus verticillatis, foliis hirsutis ovato-lan-
ceolatis, verticillis tomentosis. *Hall. hist.* 222.

Mentha arvensis verticillata. *B. pin.* 229. hirsuta.
Hist. Ox. III. 329. f. 11. t. 7. f. 5.

Mentha s. *Calamintha aquatica.* *Ger. em.* 684. *R.*
syn. 232.

Calamintha arvensis verticillata s. *aquatica* Belgarum.
Lob. ic. 515. *Park.* 36.

β *Mentha arvensis verticillata folio rotundiore odore*
aromatico. *R. syn.* 232.

Anglis, corn Mint.

Habitat in aquosis et arvis humidis frequens. 4. VII. →
IX.

Obf. *variat staminibus corolla brevioribus, æquantibus*
et longioribus.

9. MENTHA floribus verticillatis, caulibus erectis, fo- *sativa*
liis petiolatis ovatis ferratis acutis villosis.

Mentha floribus verticillatis, foliis ovatis acutiufcu-
lis ferratis, staminibus corolla longioribus. *Sp. pl.*
805.

Mentha crispa. *Park.* 31. verticillata. *B. pin.* 227.
Mentha

DIDYNAMIA GYMNOSPERMIA.

Mentha verticillata. Riv. mon. irr. t. 48. R. syn. 232.

Anglis, marsh Mint.

Habitat in palustribus et ad rivulos. 4. VII. VIII.

pulegium. 10. MENTHA floribus verticillatis, foliis ovatis obtusis subcrenatis, caulibus subteretibus repentibus, staminibus corolla longioribus. *Sp. pl.* 807.

Mentha caule prostrato, foliis subrotundis, obiter dentatis, staminibus exsertis. *Hall. hist.* 221.

Pulegium. *Fuchs. hist.* 199. *B. hist.* III. 256. *Riv. mon. irr. t.* 22. *R. syn.* 235. *latifolium.* *B. pin.* 222.

Pulegium regium. *Ger. em.* 671.

Pulegium vulgare. *Park.* 29.

exigua.

Mentha floribus verticillatis, foliis lanceolato-ovatis glabris acutis integerrimis. *Sp. pl.* 806.

Anglis, pennyroyal Mint.

Habitat in ericetis et pascuis humidis. 4. VIII. IX.

GLECOMA. 714.

Antherarum singulum par in formam crucis connivans. *Cal.* quinquefidus.

hederacea. 1. GLECOMA foliis reniformibus crenatis. *Sp. pl.* 807:

Chamæclema caule procumbente radicato, foliis reniformibus rotunde crenatis. *Hall. hist.* 245.

Hedera terrestris. *Ger. em.* 856. *vulgaris.* *Park.* 676. *B. pin.* 306.

Calamintha humilior, folio rotundiore. *Tourn.* 167. *R. syn.* 243.

β *Hedera terrestris montana.* *B. pin.* 306. *Park.* 677.

Chamæclema majus hirsutius. *Vaill. parif.* 33. t. 6. f. 4.

Anglis, common Ground-ivy.

Habitat in umbrosis et ad sepes. 4. IV. V.

Pharmac. Hederae terrestris Herba.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Alfne hederulæ folio major. *Park.* 762.
Anglis, perfoliate Archangel, or great Henbil.
 Habitat in arvis et cultis. ☉. V. VI.

GALEOPSIS. 717.

Corollæ labium superius subcrenatum, fornicatum;
 labium inferius supra bidentatum.

ladanum. 1. GALEOPSIS internodiis caulinis æqualibus, verticillis remotis, foliis lineari-lanceolatis subnudis.

Galeopsis internodiis caulinis æqualibus, verticillis omnibus remotis. *Sp. pl.* 810.

Galeopsis foliis linearibus serratis, verticillis diffitis. *Hall. hist.* 266.

Sideritis arvensis angustifolia rubra. *B. pin.* 233.

Sideritis arvensis rubra. *Park.* 587: *R. syn.* 242.

Ladanum segetum. *Riv. mon. ir.* 24.

Anglis, red Dead-Nettle; or Nettle-hemp.

Habitat in arvis solo calcareo frequens. ☉. VI—VIII.

Desc. Caulis erectus tetragonus subhirsutus, brachiatus: Folia opposita, petiolata, lineari-lanceolata et lanceolata, serrata et integerrima, acuminata nuda, et parum hirsuta, serraturis utrinque tribus s. quatuor. Flores rubri villosiusculi. Galea dentata; labia crenata; lacinia intermedia rubra albaque picta. Calycis dentes acuminati s. spinosi.

villosa. 2. GALEOPSIS internodiis caulinis æqualibus, verticillis remotis, foliis lanceolatis serratis villosis.

Galeopsis foliis rhomboidis, serratis, sericeis, verticillis diffitis. *Hall. hist.* 267.

Sideritis arvensis latifolia hirsuta lutea. *R. syn.* 242.

Sideritis lutea. *Pet. h. br. t.* 33. f. 10.

Ladanum segetum folio latiore. *Riv. mon. irr.* 24.

Anglis, hairy Dead-Nettle.

Habitat in arvis arenosis in comitatibus Eboracensi et Lancastrensi etc. frequens; prope Newark in agro Nottinghamensi; circa Bangor in Wallia. ☉. VII. VIII.

Desc. Caulis erectus, tetragonus, ramosus, villosus, ramis erectis. Folia opposita, petiolata radicalia ovata caulina lanceolata, serrata, acuminata, villosa aut sericea, venosa,

venosa, venis rectis. Calycis dentes spinosi. Corollae flavæ, villosæ : galea dentata, labia crenata. Habitus præcedentis, sed differt latitudine, serraturis, venis vilisque foliorum, et colore florum.

3. GALEOPSIS internodiis caulinis superne incrassatis, *tetrahit.* verticillis summis contiguis. *Sp. pl.* 810.

Galeopsis foliis ovato lanceolatis serratis, caule hirsuto. *Hall. hist.* 268.

Lamium cannabinum folio vulgare. *R. syn.* 240.

Cannabis spuria. *Ger. em.* 709. *Park.* 399. *Riv.* 3

Urtica aculeata, foliis serratis. *B. pin.* 232.

β Lamium cannabinum, floribus albis, verticillis purpurascens. *R. syn.* 241.

γ Cannabis spuria flore albo magno eleganti. *Merr.*

δ Galeopsis corolla flava, labio inferiore maculato. *Fl. lapp.* 193.

Galeopsis foliis ovato-lanceolatis, serratis, caule hirsuto, flore calycis quadruplo longiore. *Hall. hist.* 269.

Lamium cannabinum folio, flore amplo luteo : labio purpureo. *R. syn.* 241.

Anglis, common Dead-Nettle, or Nettle-hemp.

Habitat inter Jegetes et ad margines agrorum, δ in a corn-field, at the bottom of Ribton Lane, near Woodford.

Warn. cat. Woodf. 77. in arvis arenosis in comitatibus Eboracensi, Lancastriensi, &c. ☉. VII. VIII.

GALEOBDOLON.

CAL. *Involucrum* verticillis subiectum : foliolis linearibus, acutis calyce brevioribus.

Perianthum monophyllum, tubulatum, campanulatum; quinquedentatum, acuminatum : *dente* superiore erecto distanti ; duobus inferioribus reflexo-patulis.

COR. *monopetala, ringens. Tubus* cylindricus, brevis. *Labium* superius ovale, fornicatum, subintegrum, ciliatum, villosum. *Lab. inferius* brevius, trifidum, inæquale : *laciniis* lateralibus ovatis, acuminatis, latere exteriori reflexo ; *lacinia* media longiori recta, acuminata.

S

STAM.

DIDYNAMIA GYMNOSPERMIA.

STAM. *Filamenta* quatuor subulata, sub labio superiori recondita, quorum duo longiora. *Antheræ* oblongæ binæ, didymæ, supra convexæ, subtus concavæ.

PIST. *Germen* quadripartitum. *Stylus* filiformis longitudine et situ staminum. *Stigma* bifidum, acutum.

PER. nullum. *Calyx* immutatus, semina infundo calycis.

SEM. quatuor, brevia, triquetra, truncata.

luteum.

1. GALEOBDOLON.

Galeopsis verticillis sexfloris, involucro tetraphyllo.

Sp. pl. 810.

Leonurus foliis ovatis serratis acutis. *Hort. cliff.*

313.

Leonurus foliis cordatis. *Scop. carn.* 469. 2d. II. 409.

Cardiaca foliis petiolatis, cordatis, verticillis foliosis.

Hall. hist. 275.

Lamium luteum. *Ger. em.* 671. *Park.* 606. *R. syn.*

240.

Lamium folio oblongo luteum. *B. pin.* 231.

Anglis, yellow Galeobdolon or Dead-nettle.

Habitat in nemorosis et ad sepes. 4. V.

Obs. Folia inferiora, cordata; superiora ovata, acuta; Flores verticillati sessiles: verticilli a sex ad duodecim flores involucrati. Corolla lutea, lacinia intermedia labii inferioris fulva lineata: lineis tribus.

BETONICA. 718.

Cal. aristatus. *Corollæ* labium superius adscendens planiusculum: tubus cylindricus.

officinalis. 1. BETONICA spica interrupta, corollarum lacinia labii intermedia emarginata. *Sp. pl.* 810.

Betonica foliis petiolatis, imis cordatis, superioribus ovatis crenatis, spica brevi, foliis insidentibus.

Hall. hist. 264.

Betonica. *Ger.* 714. *R. syn.* 234. purpurea. *B. pin.* 235.

Betonica vulgaris flore purpureo. *Park.* 614.

Anglis, wood Betony.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Richardson, four miles south of Grantham, near the London road, opposite Easton. 4. VII.

- arvensis.* 4. STACHYS verticillis sexfloris, foliis obtusis nudiusculis corollis longitudine calycis, caule debili. *Syst. nat.* 395. *Sp. pl.* 814. *Fl. dan.* 587.
 Glechoma foliis cordatis oblongis crenatis, caule hispido erecto. *Sp. pl. ed. 1.* 578.
 Trixago foliis ovatis, cordatis, rotunde dentatis hispidis. *Hall. hist.* 231.
 Sideritis humilis lato obtuso folio. *Ger. em.* 699. *R. syn.* 242.
 Sideritis hederulæ folio. *Park.* 587.
 Sideritis alines trixaginis folio. *B. pin.* 233.
Anglis, corn Stachys.
 Habitat in arvis. ☉. VII. VIII.

BALLOTA. 720.

Cal. hypocrateriformis, quinquedentatus, decemstriatus. *Cor.* labium superius crenatum concavum.

- nigra.* 1. BALLOTA foliis cordatis indivisis serratis, calycibus acuminatis. *Sp. pl.* 814. *Fl. suec.* 529.
 Ballote foliis cordatis, serratis, verticillis nudis. *Hall. hist.* 259.
 Ballote. *Matth.* 825. *Fuchs.* 154. *R. syn.* 244.
 Marrubium nigrum. *Ger. em.* 701. foetidum, Ballote dictum. *Park.* 1230.
 Marrubium nigrum foetidum, Ballote Dioscoridis. *B. pin.* 230.
alba. 2. Ballota foliis cordatis indivisis serratis, calycibus subtruncatis. *Sp. pl.* 814. *Fl. suec.* 530.
 Ballota flore albo. *Tourn.* 185.
Anglis, stinking Ballote, or black Horehound.
 Habitat in ruderatis et sepibus. 4. VII.
Pharmac. Marrubii nigri Herba.

MARRUBIUM. 721.

Cal. hypocrateriformis, rigidus, decemstriatus. *Corolla* labium superius bidentum, lineare rectum.

1. MARRUBIUM

1. **MARRUBIUM** dentibus calycinis setaceis uncinatis *vulgare*.
 tis. *Sp. pl.* 816. *Hall. hist.* 258.

Marrubium album. *Ger. em.* 693. *R. syn.* 239. *vul-*
gare. *B. pin.* 230. *Park.* 44.

Anglis, white Horehound.

Habitat in ruderatis et ad vias; in ericeto Hampstediano.
 4. VII.

Pharmac. Marrubii albi *f. Prassii Herba,*

LEONURUS. 722.

Antherae punctis nitidis adspersae.

1. **LEONURUS** foliis caulinis lanceolatis trilobis, *Sp. Cardiaca.*
pl. 817. *Fl. suec.* 532. *Scop. carn.* 409.

Cardiaca foliis tripartitis lanceolatis. *Hall. hist.* 274.
Cardiaca. *Dorst. bot.* 651, *B. hist. III.* 320. *Park.*
 41. *R. syn.* 239.

Marrubium Cardiaca dictum. *B. pin.* 230.

Anglis, common Lion's tail, or Mother-wort.

Habitat in ruderatis et ad sepes. in insula Selsey prope
Chichester; inter Tickhill et Worktop in comitatu
Eboracensi. 8. VII.

Pharmac. Cardiacæ *Herba.*

•• *Calycis bilabiati,*

CLINOPODIUM. 725.

Involucrum multifetum verticillo subjectum.

1. **CLINOPODIUM** capitulis subrotundis hispidis, *vulgare.*
 bracteis setaceis. *Sp. pl.* 821.

Clinopodium foliis ovatis, acuminatis obiter dentatis
stipulis setaceis. *Hall. hist.* 239.

Clinopodium origano simile. *B. pin.* 224. *R. syn.*
 239.

Acinos. *Ger. em.* 675. *f. Clinopodium majus.* *Park.*
 22.

Anglis, wild Basil.

Habitat in pratis et ad sepes præsertim solo calcareo. 4.
 VII.

ORIGANUM. 726.

Strobilus tetragonus, spicatus calyces colligens.

- vulgare.* 1. **ORIGANUM** spicis subrotundis paniculatis conglomeratis, bracteis calyce longioribus ovatis. *Sp. pl.* 824. *Fl. suec.* 534. *Dan.* 638.
Origanum foliis ovatis, umbellis coloratis, staminibus exsertis. *Hall. hist.* 233.
Origanum anglicum. *Ger. em.* 666.
Origanum sylvestre. *B. pin.* 223.
Origanum vulgare spontaneum. *B. hist.* III. 236. *R. syn.* 236.
Majorana sylvestris. *Park.* 12.
Anglis, common Marjoram.
Habitat in dumetis et ad sepes solo calcareo. 4. VII.
Pharmac. *Origani vulgaris Herba.*

- onites.* 2. **ORIGANUM** spicis oblongis aggregatis hirsutis, foliis cordatis tomentosis. *Sp. pl.* 824.
Origanum onites. *B. pin.* 223. *R. syn.* 236.
Majorana major anglica. *Ger. em.* 664.
Majorana latifolia s. major anglica. *Park.* 12.
Anglis, pot Marjoram.
Habitat in pratis aridis; on the left hand of the road from Braintree to Raine, beyond the bridge, Mr. Dale.
R. syn. 4. VIII.

THYMUS. 727.

Calycis bilabiati, faux villis clausa.

- Serpyllum.* 1. **THYMUS** floribus capitatis, caulibus decumbentibus foliis planis obtusis basi ciliatis. *Sp. pl.* 825.
Thymus foliis ovatis ad basin ciliatis. *Hall. hist.* 235.
Serpyllum vulgare. *Ger. em.* 570. *B. hist.* III. 269.
R. syn. 230. *B. pin.* 220. *Park.* 8.
β Serpyllum vulgare flore albo. *Tourn.* 197.
γ Serpyllum vulgare flore amplo. *R. syn.* 230.
δ Serpyllum vulgare majus. *B. pin.* 220.
• Serpyllum citratum. *Ger. em.* 575.
ζ Serpyllum angustifolium glabrum. *B. pin.* 220.
• Serpyllum vulgare hirsutum. *R. syn.* 231.

Serpyllum

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

DIDYNAMIA GYMNOSPERMIA.

Calamintha odore pulegii. *Ger. em.* 687. *R. syn.* 243.

Calamintha pulegii odore s. Nepeta. *B. pin.* 228.

Calamintha altera odore pulegii, foliis maculosis.
Park. 36.

Anglis, field Calamint.

Habitat in marginibus viarum agrorumque. 4. VIII.

Pharmac. Calaminthæ officinalis Herba.

MELITTIS. 731.

Calyx tubo corollæ amplior. *Corollæ* labium superius planum; labium inferius crenatum. *Antheræ* cruciatæ.

*melisso-
phyllum.*

1. MELITTIS. *Sp. pl.* 832. *Hall. hist.* 224. *Jac. austr.* I. 18. t. 26.

Melissa Fuchsii. *Lob. ic.* 515. *Ger. em.* 690. *R. syn.* 242.

Melissophyllon Fuchsii. *Park.* 41.

Lamium montanum, melissæ folio. *B. pin.* 231.

Anglis, common Bastard-Balm.

Habitat in sylvis et sepibus. In several woods in the west of England, about Totness in Devonshire, and Haverford-west, Pembrokeshire. *R. syn. in novo salutu in comitatu Hantonensi.*

PRUNELLA. 732.

Filamenta bifurca: altero apice antherifera. *Stigma* bifidum.

vulgaris. 1. PRUNELLA foliis omnibus ovato-oblongis petiolatis. *Sp. pl.* 837. *Fl. succ.* 540.

Prunella foliis ovatis oblongis, calycibus superne truncatis. *Hall. hist.* 277.

Prunella. *Ger. em.* 631. *R. syn.* 238. *vulgaris.* *Park.* 1680.

Prunella major folio non dissecto. *B. pin.* 260.

Anglis, common Self-heal.

Habitat in pratis et pascuis. 4. VIII.

Pharmac. Prunellæ Herba.

SCUTELLARIA.

SCUTELLARIA. 734.

Calyx ore integro: post florentiam clauso, operculato.

1. SCUTELLARIA foliis cordato-lanceolatis crenatis *galerica*-rugosis, floribus axillaribus. *Sp. pl.* 835. *Fl. dan. lata.* 637.

Cassida foliis oblonge-cordatis crenatis, verticillis nudis bifloris. *Hall. bist.* 280.

Cassida palustris vulgatiores flore cæruleo. *Tourn.* 182. *R. syn.* 244.

Lysimachia cærulea galericulata, seu *Gratiola cærulea*. *B. pin.* 246.

Lysimachia galericulata. *Ger. em.* 477.

Gratiola cærulea. s. *latifolia* major. *Park.* 221.

Anglis, common Skull-cap, or Hooded-willow-herb.

Habitat in aquosis et ad ripas fluviorum. 4. VIII.

2. SCUTELLARIA foliis cordato-ovatis subintegerrimis, floribus axillaribus. *Fl. angl.* 232. *Sp. pl.* 835.

Cassida palustris minima, flore purpurascente. *Tourn.* 182. *R. syn.* 244.

Gratiola latifolia. *Ger. em.* 585. s. nostras minor. *Park.* 221.

Anglis, lesser Skull-cap, or Hooded-willow-herb.

Habitat in palustribus. 4. VII. VIII.

ANGIOSPERMIA.

* *Calyces bifida.*

OROBANCHE. 779.

Cal. bifidus. *Cor.* ringens. *Caps.* unilocularis, bivalvis polysperma. *Glandula* sub basi germinis.

1. OROBANCHE caule simplici pubescente, corollis *major*. quadrifidis.

Orobanche caule simplicissimo pubescente, staminibus subexsertis. *Sp. pl.* 882.

Orobanche

Orobánche caule simplici, stipula unica, calyce quadrifido. *Hill. hist.* 295.

Orobanche major *Garyophyllum olens*. *B. pin.* 87. *R. syn.* *288.

Rapum genistæ. *Ger. em.* 1311. *Park.* 229.

Anglis, common Broom-rape.

Habitat in pratis siccis ad radices genistæ et inter segetes. 4, V. VI.

ramosa. 2. OROBANCHE caule subramoso, corollis quinquefidis.

Orobanche caule ramoso, corollis quinquefidis. *Sp. pl.* 882.

Orobanche caule ramoso, flore quinquepartito. *Hall. hist.* 296.

Orobanche. *Cam. epit.* 311. *ramosa*. *B. pin.* 88. *Park.* 1363. *R. syn.* *288.

β Orobanche flore minore. *B. hist. II.* 781. *R. syn.* *288.

Anglis, branched Broom-rape.

Habitat in arvis et pascuis siccis. Near Beccles, in Suffolk. J. Sherrard. *R. syn.* In insula Shepey et prope Feverham et Rochester in Cantio; circa Glastonbury; et in comitatibus Devonensi et Hantoniensi. 4? VI. VII.

Desc. Caulis erectus, angulatus, squamosus, flavescens, basi ramosus, sed frequens simplex; ramis, erectis fere longituaine caulis; squamis ovatis, adpressis: Flores spicati, sessiles, fusco-purpurascens, labio superiore subrotundo bifido, inferiore trifido. Bractææ solitariae, ovatae, membranaceae, purpurascens; Stamina corollâ breviora,

** Calyces quadrifidi.

LATHRÆA. 743.

Cal. quadrifidus. Glandula depressa ad basin futuræ germinis. *Caps.* unilocularis.

squamaria. 1. LATHRÆA caule simplicissimo, corollis pendulis: labio inferiore trifido. *Sp. pl.* 884.

Anblatum. *Cord. bist.* 89. *f. Aphyllon*. *B. bist. II.* 783. *R. syn.* 288.

Orobanche

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

DIDYNAMIA ANGIOSPERMIA.

enti copiose. Near Arncastle, the seat of Lord F. Campbell, near Loch-Gyle. Dt. Hope. ☉. VII. VIII.

odonites.

3. BARTSIA foliis superioribus alternis lanceolatis serratis, floribus racemosis secundis.

Euphrasia foliis linearibus: omnibus serratis. *Sp. pl.* 841. *Fl. dan.* 625.

Euphrasia pratensis rubra. *B. pin.* 234. *R. syn.* *284. major. *Park.* 1329.

Cratæogonon Euphrosine. *Ger. em.* 92.

Odontites bracteis serratis hirsutis. *Hall, hist.* 304. *Anglis*, red Bartsia.

Habitat in pratis et pascuis. ☉. VII—IX.

RHINANTHUS. 740.

Calyx quadrifidus, ventricosus. *Caps.* bilocularis, obtusa compressa.

Crista galli.

1. RHINANTHUS corollarum labio superiore compresso brevior. *Sp. pl.* 840. *Fl. lapp.* 248. *fuec.* 542.

Alectorolophus calycibus glabris. *Hall. hist.* 313.

Pedicularis s. Crista galli lutea. *Park.* 713. *R. syn.* 284.

Pedicularis pratensis lutea s. Crista galli. *B. pin.* 163.

Crista galli. *Ger. em.* 1071.

β Pedicularis major angustifolia ramosissima, flore minore luteo, labello purpureo. *B. syn.* 284.

Anglis, common Yellow-Rattle, or Cock's-comb.

Habitat in pratis et pascuis. ☉. VI. VII.

EUPHRASIA. 741.

CAL. *Perianthium* monophyllum, tubulatum, campanulatum, angulatum, rectum, quadrifidum: laciniis acuminatis, æqualibus.

COR. *monopetala*, ringens. *Tubus* longitudine calycis. *Labium* superius, concavum, emarginatum.

Lab.

Lab. inferius patens, semitrifidum: laciniis inæqualibus, bifidis; media majori.

STAM. *Filamenta* quatuor, filiformia, sub labio superiori recondita. *Antheræ* bilobæ basi bifidæ; lobis basi aristato-acuminatis.

PIST. *Germen* oblongum. *Stylus* filiformis situ et figura staminum. *Stigma* capitatum, obtusum, integrum.

PER. *Capsula* oblonga, ima quadrangula, supra compressa, emarginata, bilocularis, bivalvis, despicimentero valvis contrario, marginibus dehiscens.

SEM. numerosa, oblonga.

1. EUPHRASIA foliis ovatis lineatis argute dentatis. *officinalis*

Sp. pl. 841. *Hall. hist.* 303.

Euphrasia. Tabern. 862. *Cam. epit.* 767. *Ger. em.* 663.

R. syn. 284. vulgaris. *Park.* 1329.

Euphrasia officinarum. B. pin. 233.

Anglis, common Eye-bright.

Habitat in pratis et pascuis. ☉. VII—IX.

Pharmac. Euphrasizæ Herba.

MELAMPYRUM. 742.

Cal. quadrifidus. *Corollæ* labium superius compressum, margine replicato. *Caps.* bilocularis obliqua, hinc dehiscens. *Semina* duo gibba.

1. MELAMPYRUM spicis quadrangularibus: bracteis cordatis compactis denticulatis imbricatis. *Sp. cristatum*
pl. 842.

Melampyrum foliis integerrimis, floribus spicatis, bracteis duplicatis cristatis imbricatis. *Hall. hist.* 311.

Melampyrum cristatum, flore albo et purpureo. B. hist. III. 440. *R. syn.* 286.

Melampyrum luteum angustifolium. B. pin. 234.

Melampyrum angustifolium cristatum, spica quadrata, flosculis ex luteo pallefcientibus. Pluk. ph. t. 99. *f.* 2.

Anglis, crested Cow-wheat.

Habitat in sylvis Cantabrigiensibus et Bedfordiensibus frequens. ☉. VII.

2. MELAMPYRUM

arvense. 2. MELAMPYRUM spicis conicis laxis: bracteis dentato-setaccis. *Sp. pl.* 842.

Melampyrum foliis integerrimis, bracteis semipinnatis. *Hall. hist.* 310.

Melampyrum purpurascente coma. *B. pin.* 234. *R. syn.* 285.

Anglis, purple Cow-wheat.

Habitat in arvis. In the corn, on the right hand, just before you come to Lycham in Norfolk. *R. syn.* on a bill near Horsley-bath, near Beeston Castle, Cheshire. *Blacks. Spec. bot. inter Norwich et Cosley.* D. Rose. ☉. VII.

pratense. 3. MELAMPYRUM floribus secundis lateralibus, conjugationibus remotis, corollis clausis. *Sp. pl.* 843.

Melampyrum foliis imis integerrimis, mediis dentatis, floralibus hastatis. *Hall. hist.* 308.

Melampyrum luteum latifolium. *B. pin.* 234.

Melampyrum latifolium flore albo: labio inferiore duabus maculis luteis distincto. *R. syn.* 286.

Anglis, meadow Cow-wheat.

Habitat in pratis, pascuis et dumetis. In the wood by Dr. Richardson's house at North Bierley. *R. syn.* ☉. VII. VIII.

sylvaticum. 4. MELAMPYRUM floribus secundis lateralibus, conjugationibus remotis, corollis hiantibus. *Sp. pl.* 843. *Fl. dan.* 145.

Melampyrum foliis integerrimis, floribus hiantibus. *Hall. hist.* 307.

Cratæogonon album. *Ger. em.* 91.

Cratæogonon vulgare. *Park.* 1326.

Anglis, yellow Cow-wheat.

Habitat in sylvis et nemorosis. ☉. VI—VIII.

*** Calycibus quinquefidis.

PEDICULARIS. 746.

Cal. quinquefidus. *Caps.* bilocularis, mucronata, obliqua. *Sem.* tunicata.

palustris. 1. PEDICULARIS caule ramoso, calycibus callosopunctatis, corollis labio obliquis. *Sp. pl.* 845.

Pedicularis

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Elatine. *Cam. epit.* 754. altera. *Ger. em.* 623.

Anglis, sharp pointed Toadflax or Fluellin.

Habitat in arvis. ☉. VII.—IX.

Pharmac. Elatines *Herba.*

Obs. *variat floribus regularibus quinquefidis; calcaribus tribus quatuor aut quinque, ut in Peloria.*

spurius

3. ANTIRRHINUM foliis ovatis alternis, caulibus procumbentibus. *Sp. pl.* 851.

Antirrhinum caule procumbente, foliis villosis ovatis, imis conjugatis, superioribus alternis. *Hall. hist.*

341.

Linaria Elatine dicta folio subrotundo. *R. syn.* 282.

Elatine folio subrotundo. *B. pin.* 252. *Park.* 553.

Veronica foemina. *Cam. ep.* 462. *Dod. pempt.* 42.

f. Elatine. *Ger. em.* 625.

Anglis, round-leaved Toadflax or Fluellin.

Habitat in arvis. ☉. VII.—IX.

Obs. *variat floribus regularibus, &c. ut in præcedenti.*

•• *Foliis oppositis; Florum calcari brevi obtusiusculo.*

minus.

4. ANTIRRHINUM foliis plerisque alternis, lanceolatis, obtusis caule ramosissimo diffuso. *Sp. pl.* 852. *Fl. dan.* 502.

Antirrhinum viscidum, foliis inferioribus conjugatis ellipticis obtusis hirsutis, calcare dimidii floris longitudine. *Hall. hist.* 335.

Antirrhinum arvense minus. *B. pin.* 212.

Antirrhinum minimum repens. *Ger. em.* 549.

Antirrhinum sylvestre minimum. *Park.* 1334.

Linaria antirrhinum dicta. *R. syn.* 283.

Anglis, least Toadflax.

Habitat in arvis solo arenoso. . VI.—IX.

*** *Foliis oppositis; Florum calcari exquisito.*

repens.

5. ANTIRRHINUM foliis lanceolato-linearibus confertis; inferioribus verticillatis, caule paniculato, pedunculis spicatis nudis.

*monspe-
sulanum.*

Antirrhinum foliis linearibus confertis caule nitido paniculato, pedunculis spicatis nudis. *Sp. pl.* 854.

Linaria

- Linaria capillaceo folio, odora. *B. pin.* 213. *pr.* 106.
Dill. elth. 199.
- Linaria odorata monspessulana. *B. hist.* III. 456.
R. syn. *282.
- β Antirrhinum foliis linearibus confertis: inferioribus repens, quaternis, calycibus capsulæ æqualibus. *Sp. pl.* 854.
- Linaria angustifolia flore cinereo striato. *Dill. elth.* 190. *t.* 163. *f.* 197.
- Linaria cærulea foliis brevioribus et angustioribus.
R. syn. *282.
Anglis, creeping Toadflax.
Habitat in pratis pascuis et ad sepes; in Cornubia non longe ab oppido Penryn occidentem versus; et circa Henley. R. syn. 4. VII.—IX.
6. ANTIRRHINUM foliis sublinearibus: inferioribus *arvense.* quaternis, calycibus piloso-viscidis, floribus spicatis, caule erecto. *Sp. pl.* 855.
- Linaria arvensis cærulea. *B. pin.* 213. *pr.* 107. *Dill. elth.* 200.
- β Linaria pumilla, foliolis carnosis, flosculis minimis flavis. *B. pin.* 213. *Dill. elth.* 200.
- Linaria tetraphylla lutea. *Col. scphr.* I. 299. *t.* 300.
Anglis, corn Toadflax.
Habitat in arvis solo arenoso. ☉. VI. VII.
7. ANTIRRHINUM foliis lanceolato-linearibus confertis, caule erecto, spicis terminalibus sessilibus, floribus imbricatis. *Sp. pl.* 858. *Fl. suec.* 557.
- Antirrhinum foliis linearibus ascendentibus congestis, caule erecto spicato. *Hall. hist.* 336.
- Linaria lutea vulgaris. *Ger. em.* 550. *R. syn.* 281.
- Linaria vulgaris nostras. *Park.* 458.
- Linaria vulgaris lutea flore majore. *B. pin.* 212.
- β Peloria. *Amœn. Acad.* I. 55. *t.* 3.
Anglis, common Toadflax.
Habitat in pratis et pascuis sterilioribus. 4. VII.
Pharmac. Linariæ Herba.

*** Corollis ecaudatis.

majus.

8. ANTIRRHINUM corollis ecaudatis, floribus spicatis, capsulis rotundatis. *Sp. pl.* 859.

Antirrhinum foliis ellipticis obtusis, floribus spicatis, calycibus obtusis, calcare brevissimo. *Hall. hist.* 333.

Antirrhinum purpureum. *Ger. em.* 549.

Antirrhinum majus alterum folio longiore. *B. pin.* 211.

Anglis, greater Toadflax or Snapdragon.

Habitat in muris antiquis; in clivis cretaceis prope Dover, et inter Northfleet et Gravesend. *♁. VI. VII.*

arenitum.

9. ANTIRRHINUM corollis ecaudatis; floribus subspicatis, calycibus digitatis corolla longioribus. *Sp. pl.* 860.

Antirrhinum foliis ellipticis obtusis, floribus spatosis, calycibus subulatis longissimis, calcare brevissimo. *Hall: hist.* 334.

Antirrhinum angustifolium sylvestre. *B. hist.* III. 464. *R. syn.* 383.

Antirrhinum arvense majus. *B. pin.* 212.

Antirrhinum minus. *Ger. em.* 549.

Antirrhinum sylvestre medium. *Park.* 1334.

Anglis, lesser Toadflax or Snapdragon.

Habitat in arvis, solo arenoso. *♁. VII. VIII.*

SCROPHULARIA. 756.

Cal. quinquefidus. *Cor.* subglobosa, resupinata. *Caps.* bilocularis.

nodosa.

1. SCROPHULARIA foliis cordatis trinervatis, caule obtusangulo. *Sp. pl.* 863.

Scrophularia caule quadrangulari, paniculato, foliis cordato-oblongis. *Hall. hist.* 326.

Scrophularia major. *Ger. em.* 716. *R. syn.* 283.

Scrophularia nodosa foetida. *B. pin.* 235.

Scrophularia vulgaris. *Park.* 610.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

DIDYNAMIA ANGIOSPERMIA.

Digitalis foliis calycinis ovatis, galea simplici. Hall. hist. 330.

Digitalis purpurea. Fuchf. hist. 893. Ger. em. 790. R. syn. 283. folio aspero. B. pin. 243.

Digitalis purpurea vulgaris. Park. 653.

β *Digitalis alba folio aspero. B. pin. 243. Anglis, purple Foxglove.*

Habitat ad vias et sepes. ♂. VII.

Pharmac. Digitalis Folia, Flores.

SIBTHORPIA. 775.

Cal. quinquepartitus. Cor. quinquepartita æqualis. Stam. paribus remotis. Caps. compressa, orbicularis, bilocularis : dissipimento transverso.

europæa. 1. SIBTHORPIA foliis reniformibus subpeltatis crenatis. Sp. pl. 880. Læf. it. 150.

Chryso-splenium cornubiense. Pet. h. br. t. 6. f. 11.

Alfina spuria pusilla repens foliis saxifragæ aureæ. R. cat. Angl. Pluk, ph. t. 7. f. 6. R. syn. 352.

Anglis, cornish Siphthorpia or Money-wort.

Habitat in palustribus umbrosis, et ad scaturigines et rivulos in Cornubia et Devoniam frequens. Circa Longsledale in Westmorlandia. Will. syn. 4. VII.—IX.

LIMOSELLA. 776.

Cal. quinquefidus. Cor. quinquefida, æqualis. Stam. per paria approximata. Caps. unilocularis, bivalvis, polyspermia.

aquatica. 1. LIMOSELLA. Sp. pl. 881. Fl. dan. 89. Hall. hist. 300.

Plantaginella palustris. B. pin. 190. R. syn. 278.

Spergula perpusilla, lanceolatis foliis. Læf. pr. 261. ic. 11.

Alfina palustris repens, foliis lanceolatis, floribus perexiguis. Pluk. ph. t. 74. f. 4.

Anglis, common Mudwort.

Habitat in innundatis et ubi aquæ pluviales stagnant. ☉. VII.—IX.

Classis

*Classis XV.*TETRADYNAMIA
SILICULOSA.* *Silicula integra, nec apice emarginata.*

MYAGRUM. 796.

Silicula stylo conico terminata, loculo submonospermo.

- 1. MYAGRUM siliculis obovatis pedunculatis polyspermis. *Sp. pl.* 894.

Myagrū. *Ger. em.* 273. *R. syn.* 302.Myagrū sylvestre. *B. pin.* 109. f. Pseudo-myagrū. *Park.* 868.Alyssum foliis sagittatis, siliquis cordiformibus. *Hall. hist.* 489.*Anglis, common Myagrū or Gold of Pleasure.**Habitat in agris inter Linum sæpe copiose, cum quo ab exteris terris introducitur; prope Bridport et Lime in agro Dorsetsiensi. ☉. V. VI.*

SUBULARIA. 799.

Silicula integra, ovata: valvis, ovatis, concavis, dissipimento contrariis. Stylus silicula brevior.

1. SUBULARIA. *Sp. pl.* 896. *Fl. suec.* 566. *dan.* 35. *aquatica.*

Subularia erecta, junci foliis acutis mollibus. *R. syn.* 307.Graminifolia aquatica, thlaspeos capitulis rotundis, septo medio siliculam dirimente. *Pluk. ph. t.* 188. f. 5.Gramen juncum hybernum, thlaspeos capitulis. *Hist. Ox. III. f.* 8. t. 10. f. ult.*Anglis, Awl-wort.*

TETRADYNAMIA SILICULOSA.

Habitat *sub aqua, in lacu* Hibernico Lough-Neagh,
R. *syn. in lacubus alpinis circa* Llanberris *copiose* ;
prope Loch Carran *in* Scotia. D, Hope. ☉. VII.

V E L L A. 797.

Silicula dissepimento valvulis duplo majore, extus
ovato.

annua.

1. VELLA foliis pinnatifidis, filiculis pendulis. *Sp. pl.*
895.

Nasturtium sylvestre erucæ affine. *B. pin.* 105. *R.*
syn. 304.

Nasturtium sylvestre valentinum. *Clus. hist.* II. 130.
B. hist. II. 920. *Park.* 830.

Eruca nasturtio cognita tenuifolia. *Ger. em.* 247.
Anglis, annual Cresse-rocket.

Habitat *on* Salisbury-plain, *not far from* Stonehenge,
Mr. Lawson. *R. syn.* ☉. VI.

D R A B A. 800.

Silicula integra, ovali-oblonga : valvis planiusculis,
dissepimento parallelis. *Stylus* nullus.

verna.

1. DRABA scapis nudis, foliis lanceolatis subincisis.
Sp. pl. 896.

Draba cauliculis nudis, foliis subhirsutis subdentatis.
Hall. hist. 496.

Paronychia vulgaris. *Ger. em.* 624. *R. syn.* 292.

Paronychia vulgaris, alfinis folio. *Park.* 556.

Bursa pastoris minor loculo oblongo. *B. pin.* 108.

Anglis, common Whitlow-grass.

Habitat *in muris et siccioribus.* h. IV.

muralis.

2. DRABA caule ramoso, foliis cordatis dentatis am-
plexicaulibus. *Sp. pl.* 897. *Hall. hist.* 499.

Draba minima muralis discoides. *Column. ec.* 274.

Bursa pastoris major loculo oblongo. *B. pin.* 108.
pr. 50. *R. syn.* 292.

Thlaspi veronicæ folio. *Park.* 843.

Anglis, wall Whitlow-grass.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

TETRADYNAMIA SILICULOSA.

Nasturtium apetalum, foliis pinnatis, radicalium pinnis semipinnatis, caulinis simplicibus. *Hall. hist.* 508.

Nasturtium sylvestre, osyridis folio. *B. pin.* 105.

Park. 829. *R. syn.* 303. capsulis minimis. *Hist.*

Ox. II. 301. *f.* 3. *c.* 19. *f.* 7.

Thlaspi minus. *Ger. em.* 269.

Anglis, narrow-leaved Pepperwort or Dittander.

Habitat in maritimis, ruderatis. At Maldon, Essex;

and Yarmouth, Lynne and Clay, in Norfolk. And

Truro, Cornwall. *R. syn.* ♂. VI. VII.

petræum. 3. **LEPIDIUM** foliis pinnatis integerrimis, petalis emarginatis calyce minoribus. *Sp. pl.* 899. *Jac. austr.* II. 19. t. 131.

Nasturtium foliis pinnatis, radicalibus ovatis, lanceolatis, caulinis linearibus. *Hall. hist.* 515.

Nasturtium pumilum vernalium. *B. pin.* 105.

Nasturtium petræum annuum. *Pluk. ph.* 206. *f.* 4.

Nasturtiolum montanum annuum tenuissime divisum.

R. syn. 304.

Cardamine pusilla saxatilis montana discoides. *Col. ec.* I. 271. t. 274.

Anglis, mountain Pepperwort.

Habitat in rupibus, saxosis et muris; in rupe St. Vincent's rock dicto et muris circa Bristol passim; ad

Uphill, in agro Somersetensi. IV. V.

anglicum. 4. **LEPIDIUM** floribus diandris tetrapetalis, foliis pinnatifidis, caule procumbente piloso.

Anglis, procumbent Pepperwort.

Habitat in ruderatis, circa Exeter. D. Newberry; circa Truro et Penryn in Cornubia. ♂. VI. VII.

Desc. Radix annua ramosa. Caules plures, herbacei, pedales, procumbentes, trigoni, foliati, pilosi, ramosi; Ramis alternis, longis. Folia alterna sessilia, pinnatifida, nuda; Laciniis suboppositis, acutis, serratis, præcipue margine superiore; dentibus duobus aut tribus. Flores racemosi, minimi, albi, sparsi, pedunculati; pedunculis horizontalibus, longis. Racemi laterales, simplices, oppositifolii, folio longiores, pilosi. Cal. Perianthium tetraphyllum, foliolis ovatis, concavis, deciduis, margine tenui. Cor. tetrapetala. Petala æqualia, ovata, calyce paulo breviora

breviora. Stam. Filamenta duo, opposita, subulata, calyce longiora. Antheræ subrotundæ, magnæ simplices, flavæ. Pist. Germen obcordatum. Stylus brevissimus; Stigma obtusum. Pericarpium silicula semibifida, lobis subrotundis, compressis, rugosis, bivalvibus. Valvulis navicularibus, carinatis; Dissipimento lanceolato. Sem. solitaria, reniformia utrinque acuta, fusca.

Obs. variat floribus apetalis et subinde tetrandris. Hæc planta Lepidium bonariense Lin. Sp. pl. 901. videtur, sed cum Thlaspi bonariensi, &c. Dill. elth. 281, t. 286. f. 370. non confundenda, cui folia linearia, serrata.

THLASPI. 802.

Silicula emarginata, obcordata, polysperma: valvulis navicularibus, marginato-carinatis.

1. **THLASPI** siliculis subrotundis, foliis sagittatis *den-campestre*.
tatis incanis. *Sp. pl. 903.*

Thlaspi vulgatus. B. hist. II. 921. R. syn. 305.

Thlaspi arvense, vaccariæ folio, majus. B. pin. 106.

Thlaspi vulgatissimum. Ger. em. 262.

Thlaspi mithridaticum f. vulgatissimum, vaccariæ folio. Park. 835.

Anglis, mithridate Bastard-cresse or Mustard.

Habitat in arvis, apricis, solo argilloso et arenoso. ♂.
VI. VII.

Pharmac. Thlaspeos vulgaris semina.

2. **THLASPI** siliculis subrotundis pilosis, foliis caulinis *hirtum*.
sagittatis villosis. *Sp. pl. 901.*

Thlaspi capsulis hirsutis. B. hist. II. 922.

Thlaspi villosum, capsulis hirsutis. B. pin. 106.

Thlaspi vaccariæ incano folio perenne. R. syn. 305.

Nasturtium foliis imis petiolatis ovatis, caulinis sagittatis dentatis. Hall. hist. 509.

Anglis, hairy Bastard-cresse.

Habitat in montosis Cambrobritannia et alibi observavi.
R. syn. ♂. VII.

Obs. habitus præcedenti, sed siliculis villosis.

3. **THLASPI** siliculis orbiculatis, foliis oblongis *denta-arvense*.
tis glabris. *Sp. pl. 901.*

Thlaspi

TETRADYNAMIA SILICULOSA.

Thlaspi arvense, siliquis latis. *B. pin.* 105.

Thlaspi Dioicoridis. *Ger. em.* 262. *R. syn.* 305.

Thlaspi drabæ folio. *Park.* 836.

Nasturtium siliquis orbiculatis, planis, foliis oblongis dentatis glabris. *Hall. hist.* 511.

Anglis, treacle Bastard-creffe or Mustard.

Habitat in agris, solo lutofo passim. ☉, VI. VII.

Pharmac. *Thlaspeos semina*.

alpestre.

4. **THLASPI** siliculis obcordatis, foliis subdentatis; caulinis amplexicaulibus, petalis longitudine calycis, caule simplici. *Sp. pl.* 903.

Lepidium caule erecto, foliis radicalibus ovatis petiolatis, caulinis ovatis amplexicaulibus. *Hall. hist.* 519.

Thlaspi vaccariæ folio, bursæ pastoris siliquis. *B. pin.* 105. *pr.* 47.

Thlaspi minus Clusii. *Ger. em.* 268.

Thlaspi pumilum H. *Clus. hist.* II. 31.

Anglis, perfoliate Bastard-creffe.

Habitat in pratis et pascuis montosis et saxosis solo calcareo.

Among the stone-pits between Whitney and Burford, Oxfordshire; on the moist limestone pastures in Yorkshire, Westmorland and Cumberland. *R. syn.* ☉. V. VI.

Obs. *Caule erecto simplici in montosis; ramoso in hortis et solo fertilique. Folia ima petiolata ovata: an Conf. Thlaspi perfoliatum.*

montanum.

5. **THLASPI** siliculis obcordatis, foliis glabris, radicalibus carnosis obovatis integerrimis, caulinis amplexicaulibus corollis calyce majoribus. *Sp. pl.* 903. *Jac. austr.* III. 22. t. 237.

Lepidium caule erecto foliis radicalibus petiolatis ovatis, caulinis amplexicaulibus, sagittatis. *Hall. hist.* 518.

Thlaspi montanum, glasti folio minus. *Park.* 842.

Thlaspi alpinum bellidis cæruleæ folio. *B. pin.* 106.

Thlaspi foliis globulariæ. *B. hist.* II. 926. *R. syn.* 305.

Thlaspeos albi supini varietas. *Gr. em.* 268.

Anglis, mountain Bastard-creffe.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- anglica.* 2. **COCHLEARIA** foliis omnibus ovato-lanceolatis sinuatis. *Fl. angl.* 248. *Sp. pl.* 903. *Fl. dan.* 329. Cochlearia folio sinuato. *B. pin.* 110. *R. syn.* 303. Cochlearia vulgaris. *Park.* 285. Cochlearia britannica. *Ger. em.* 401. *Anglis*, sea Scurvy grass. *Habitat in littoribus maritimis solo lutofo.* ♂. V. *Pharmac.* Cochleariæ marinæ s. britannicæ *Herba.* *Obs.* An varietas præcedentis?
- danica.* 3. **COCHLEARIA** foliis omnibus deltoïdibus. *Fl. dan.* 100. Cochlearia foliis hastatis angulatis. *Sp. pl.* 905. Cochlearia marina folio anguloso parvo. *R. syn.* 303. Thlaspi hederaceum. *Lob. ic.* 615. *Anglis*, danish Scurvy-grass. *Habitat in littoribus marinis.* In insula Walney, Lancastriæ; near Llanbadrick Church in Anglesea. Mr. Lhwyd. *R. syn.* ♂. V. VI.
- ceronopus.* 4. **COCHLEARIA** foliis pinnatifidis. *Sp. pl.* 904. *Fl. dan.* 202. *suec.* 579. Coronopus. *Ruel.* 373. *Ger. em.* 427. *Hal. hist.* 502. recta vel repens Ruellii. *Park.* 502. Nasturtium supinum capsulis verrucosis. *R. syn.* 304. Ambrosia campestris repens. *B. pin.* 138. *Anglis*, swine's Scurvy-grass or Cresse. *Habitat in ruderatis et ad vias.* ☉. VI.—VIII.
- armoracia.* 5. **COCHLEARIA** foliis radicalibus lanceolatis crenatis, caulinis incis. *Sp. pl.* 904. *Hall. hist.* 504. Raphanus rusticanus. *Ger. em.* 241. *Park.* 860. *B. pin.* 96. *R. syn.* 301. *Anglis*, horle Scurvy-grass or Radish. *Habitat in ruderatis et ad fossas; circa Londinum frequens; about Alnwick and elsewhere in Northumberland; upon the banks of Skipton-beck, and elsewhere in Bolland Craven.* *R. syn.* 4. V. *Pharmac.* Raphani rusticani *Radix.*

I B E R I S. 804.

Cor. irregularis: petalis duobus exterioribus majoribus. *Silicula* polysperma emarginata.

*1. **IBERIS** herbacea foliis lanceolatis acutis subdentatis, *amara*.
floribus racemosis. *Sp. pl.* 906.

Iberis foliis dilatatis dentatis, floribus umbellatis.
Hall hist. 520.

Thlaspi umbellatum arvense iberidis folio. *B. pin.*
106.

Thlaspi amarum. *Tab. ic.* 462. *Ger. em.* 263.

Anglis, bitter Candy-tuft.

Habitat in arvis circa Henley et alibi comitatu Oxoniensi. ☉. VI. VII.

2. **IBERIS** herbacea foliis sinuatis, caule nudo simplici. *nudicaulis.*
Sp. pl. 907. *It. æland.* 139. *Fl. dan.* 322. *lis.*

Iberis foliis pinnatis, pinnis acutis. *Hall. hist.* 521.

Bursa pastoris minor. *Ger. em.* 251. *Park.* 806. foliis
incisis. *B. pin.* 108.

Anglis, naked Candy-tuft or Rock-creffe.

Habitat in glareosis et ericetis sterilissimis; near Hampton-court, Richmond and Barnes commons. *R. syn.*
in ericeto Putnejano. *D. Ph. Miller.* ☉. V. et
VII.

Obs. variat caule erecto et procumbente nudo et folioso,
foliis duobus tribusve oblongis integris.

S I L I Q U O S A.

* *Calyx clausus: foliolis longitudinaliter conniventibus.*

D E N T A R I A. 811.

Siliqua elastice defiliens valvulis revolutis. *Stigma*
emarginatum. *Cal.* longitudinaliter connivens.

1. **DENTARIA** foliis inferioribus pinnatis, summis *bulbifera*.
simplicibus. *Sp. pl.* 912. *Fl. dan.* 367. *Hall. hist.*
470.

Dentaria

Dentaria bulbifera. *Ger. em.* 984.

Dentaria heptaphylla baccifera. *B. pin.* 322.

Coralloides. *Cord. hist.* 151.

Anglis, bulbiferus Coral-wort.

Habitat in sylvis et umbrosis. In old Park wood, near Harefield, Blackf. fasc. 23. in silvis inter Beconfield in Wickham copiose. 4. IV. V.

ERYSIMUM. 814.

Siliqua columnaris exacte tetraedra. *Cal.* clausus.

officinale. 1. ERYSIMUM siliquis spicæ adpressis foliis runcinatis. *Sp. pl.* 922. *Fl. dan.* 260.

Erysinum foliis pinnatis, pinnis rectangulis, acutis extrema triangulari maxima, siliquis adpressis. *Hall. hist.* 878.

Erysinum vulgare. *B. pin.* 298.

Erysimum Dioscoridis Lobelio. *Ger. em.* 254.

Iris I. *Tabern.* 448. s. Erysimum. *Dod. pempt.* 714.

Lob. ic. 206. vulgare. *Park.* 833.

Eruca hirsuta, siliqua caule appressa, Erysimum dicta. *R. syn.* 298.

Anglis, common Hedge-mustard,

Habitat in ruderatis et ad vias. ☉. V. VI.

Pharmac. Erysimi Herba.

barbarea. 2. ERYSIMUM foliis lyratis: extimo subrotundo. *Sp. pl.* 922.

Erysimum foliis pinnatis, pinna extrema maximo subrotunda. *Hall. hist.* 479.

Eruca lutea latifolia s. Barbarea. *B. pin.* 98. *R. syn.* 297.

Barbarea. *Dod. pempt.* 712. *Ger. em.* 243. *B. hist.* II. 868. flore simplici. *Park.* 819.

β Barbarea foliis minoribus et frequentius sinuatis. *R. syn.* 297. *hist.* 809.

Anglis, winter Hedge-mustard or Cresse.

Habitat in palustribus et ad fossas. 4. V.

alliarica. 3. ERYSIMUM foliis cordatis. *Sp. pl.* 922. *Hall. hist.* 480.

Alliaria.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

TETRADYNAMIA SILIQUOSA.

Leucojum foliis glabris lanceolatis, integerrimis. *Hall. hist.* 443.

Leucojum luteum, vulgo Cheiri flore simplici. *B. hist.* II. 872. *R. syn.* 291.

Leucojum luteum vulgare. *B. pin.* 202.

Viola lutea. *Ger. em.* 456.

Keira s. Leucojum vulgare luteum. *Park.* 625.

Anglis, wall Stock or Flower.

Habitat in muris antiquis. *U. V.*

Pharmac. Cheiri Flores.

sinuatus. 3. CHEIRANTHUS foliis tomentosis obtusis subsinuatis: rameis integris, siliquis muricatis. *Sp. pl.* 926.

Leucojum maritimum sinuato folio. *B. pin.* 201.

Leucojum marinum majus. *Park.* 622. *R. syn.* 291.

Anglis, sea Stock.

Habitat in littoribus maritimis arenosis, Walliæ et Cornubiæ. *U. V. VI.*

HESPERIS. 817.

Petala oblique flexa. *Glandula* intra stamina breviora. *Siliqua* stricta. *Stigma* basi bifurca apice connivente. *Cal.* clausus.

inodora. 1. HESPERIS caule simplici erecto, foliis subhastatis dentatis, petalis obtusis. *Sp. pl.* 927.

Hesperis caule hirsuto, foliis subsperis ovato-lanceolatis serratis petiolatis. *Hall. hist.* 191.

Hesperis sylvestris inodora. *B. pin.* 202. *R. syn.* 293. *Rupp. jen.* 78. t. 1.

Hesperis pannonica inodora. *B. hist.* II. 878. *Park.* 628.

Anglis, wild Rocket, or Dame's violet.

Habitat in pascuis et ad sepes. On the banks of the rivers about Dalehead and Grassmire, Westmorland.

R. syn. *U. V. VI.*

Obs. Foliis ovatis, dentato-serratis, acutis, subsessilibus, hastatis non vidi. nimis affinis *H. matronali*.

RAPHANUS. 822.

Cal. clausus. *Siliqua* torosa, subarticulata, teres. *Glandulæ*

dulæ melliferæ duo inter stamina breviora et pistillum, totidem inter stamina longiora et calycem.

1. RAPHANUS siliquis teretibus articulatis lævibus unilocularibus. *Sp. pl.* 935. *Hall. hist.* 468. *Raphanistrum*

Raphanus sylvestris. *Ger. em.* 240.

Raphanistrum siliqua articulata glabra, majore et minore. *Hist. Ox. II.* 265. *R. syn.* 296.

Rapistrum album articulatum. *Park.* 863.

Rapistrum flore albo siliqua articulata. *B. pin.* 95.

- β Rapistrum flore luteo siliqua articulata. *R. syn.* 296.

- γ Raphanus maritimus flore luteo, siliquis articulatis. *R. syn.* 296.

Anglis, corn Radish, or Charlock.

Habitat inter segetes. ☉. VI. VII.

BRASSICA. 820.

Cal. erectus, connivens. *Sem.* globosa. *Glandula* inter stamina breviora et pistillum, interque longiore et calycem.

1. BRASSICA radice caulescente tereti carnosa. *Sp. pl. oleracea.* 932.

Brassica maritima arborea s. procerior ramosa. *Hist. sylvestris Ox. II.* 208. *R. syn.* 293.

Brassica radice alba. *Tourn.* 219.

Anglis, sea Cabbage. *Pet. h. br. t.* 45. *f.* 6.

Habitat in clivis maritimis. On Dover Cliffs, &c. *R. syn.* prope castellum Harleck in comitatu Mervinia, et Penzanse et alibi in Cornubia. ♀. V. VI.

2. BRASSICA radice caulescente orbiculari depressa carnosa. *Sp. pl.* 931. *Rapa*

Rapa sylvestris. *B. hist.* 800.

Rapa sativa rotunda. *B. pin.* 89. *R. syn.* 294.

Rapum majus. *Ger. em.* 232.

- β Rapa sativa oblonga s. foemina. *B. pin.* 90.

Anglis, Turneps.

Habitat in arvis et ad margines agrorum. ♂. IV.

Pharmac. Rapi Radix, Semina.

- Napus.* 3. BRASSICA radice caulescente fusiformi. *Sp. pl.* 931. *Napus sylvestris.* *B. pin.* 95. *R. syn.* 295. *Bunias seu Napus sylvestris nostras.* *Park.* 865. *Bunias sylvestris Lobelii.* *Ger. em.* 235. *Anglis,* wild Cabbage, Rape, Nape, or Navew. *Habitat in aggeribus fossarum et inter segetes.* ♀. V.
- campestris.* 4. BRASSICA radice cauleque tenui, foliis caulinis uniformibus cordatis sessilibus. *Sp. pl.* 931. *Jac. austr.* III. 45. t. 283. *Brassica campestris perfoliata flore luteo.* *Loes pruf.* 29. *Eruca foliis ovatis integerrimis amplexicaulibus.* *Hall. hist.* 457. *Perfoliata siliquosa.* *Ger. em.* 536. *B. hist.* 335. *Perfoliata siliquosa vulgaris* t. *Brassica campestris.* *Park.* 580. *Anglis,* field Cabbage. *Habitat in arvis et clivis maritimis prope Harvicum oppidum in Essexia, et in clivis juxta Bardsey prope Orford in Suffolcia a D. Dale.* *R. syn.* ☉. VI. *Obs. Folia radicalia, oblonga, sinuata, obtusa, scabra; Flores lutei, aut albi.*
- orientalis.* 5. BRASSICA foliis cordatis amplexicaulibus glabris radicalibus scabris integerrimis, siliquis tetragonis. *Sp. pl.* 931. *Jac. austr.* III. 45. t. 282. *Brassica campestris perfoliata flore albo.* *B. pin.* 112. *Brassica orientalis' perfoliata, flore albo, siliqua quadrangula.* *Tourn. cor.* 16. *Anglis,* perfoliate Cabbage. *Habitat in arvis et clivis maritimis; inter segetes prope Godstone et Marshfield in Sufflexia.* ☉. VI. *Obs. nimis affinis præcedenti; Flores albi.*
- muralis.* 6. BRASSICA foliis lanceolatis sinuato-ferratis læviusculis, caule erecto glabro. *Sisymbrium subacaule, foliis lanceolatis sinuato-ferratis læviusculis, scapis-subscabris ascendentibus.* *Sp. pl.* 918. *Eruca sylvestris.* *Ger. em.* 246. *R. syn.* 296. *vulgarior.* *Park.* 818.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

TETRADYNAMIA SILIQUOSA:

Turritis muralis minor. *Pet. h. br. t. 47. f. 12. R. syn. 294.*

Erysimo similis hirsuta, non laciniata alba. *B. pin. 101. pr. 101. ic.*

Erysimo similis hirsuta planta. *Park. 834.*

Anglis, rough Tower-mustard.

Habitat in rupibus, saxosis et muris antiquis. In Switham-bottom prope Croydon. 4. VI.

ARABIS. 818.

Glandulæ nectariferæ quatuor, singulæ intra calycis foliola, squamæ instar reflexæ.

italiana. 1. ARABIS foliis petiolatis lanceolatis integerrimis. *Sp. pl. 929.*

Arabis foliis radicalis petiolatis, ovatis, dentatis, caule subnudo ramoso. *Hall. hist. 452.*

Turritis vulgaris ramosa. *Tourn. 224. R. syn. 294.*

Bursæ pastoris similis siliquosa major et minor. *B. pin. 108.*

Paronychia major et altera minor. *Park. 556.*

β Turritis minor foliosa. *Pet. h. br. t. 48. f. 2.*

Anglis, common Wall-Cresse.

Habitat in muris, tectis et pascuis arenosis et aridis. ☉. V.

stricta.

2. ARABIS foliis dentatis hispidis: radicalibus spatulatis, caulinis semiamplexicaulibus oblongis, siliquis ancipitibus erectis.

Arabis multicaulis, foliis radicalibus scabris dentatis, dentibus ciliatis. *Hall. nom. et hist. 453.*

Hesperis alpina minor, flore albo, siliquis longis. *R. syll. stirp. exter. 296.*

Anglis, rough Wall-Cresse.

Habitat in rupibus et saxosis montosis. In rupe St. Vincentii prope Bristol. ☉. V.

Desc. Radix annua, ramosa. Caules plures, semipedales, herbacei, erecti, teretes, basi villosi, supra glabri, simplices. Folia radicalia plurima, spatulata, dentata et inciso-dentata, hispida aut piloso-scabra: pilis longis, albis. Caulina tria aut quatuor, semiamplexicaulia, oblonga, dentata, hispida. Flores racemosi, pedunculati, magni, albi; pedunculis brevibus. Petala obovata integerrima

*gerrima calyce duplo longiora. Siliquæ erectæ, striatæ
sesquipollicares, ancipites, basi quadrangulæ, striatæ.
Semina subrotunda, fusca.*

3. ARABIS foliis amplexicaulibus, siliquis decurvis pla- *Turrita,*
nis linearibus, calycibus subrugosis. *Sp. pl. 930.*
Fl. dan. 62.

*Brassica sylvestris foliis integris et hispidis. B. pin.
Hist. Ox. II. 211. f. 3. t. 2. f. 23.*

*Brassica sylvestris, albido flore, nutante siliqua. Bocc.
mus. II. 81. t. s. 72. Barr. pl. rar. 353.*

Turrita major plateau. Clus. hist. II. 126.

Anglis, tower Wall-Cresse.

*Habitat in muris antiquis et saxosis. On Trinity and St.
John's Colledge Walls, Cambridge. Dr. Martyn.*

©. VI.

•• *Calyx hians : foliolis superne distantibus.*

CARDAMINE. 812.

*Siliqua elastice diffiliens valvulis revolutis. Stigma in-
tegrum. Cal. subhians. Glandulæ nectariferæ duæ,
una utrinque inter stamina breviora et calycem.*

* *Foliis simplicibus.*

1. CARDAMINE foliis simplicibus ovatis integerrimis : *bellidifo-*
petiolis longis. *Sp. pl. 913. Fl. dan. 20.* *lia.*

*Cardamine pumila bellidis folio, alpina. Ger. em.
260.*

Plantula cardamines alterius æmula Clusii. Park. 828.

Nasturtium alpinum bellidis folio minus. B. pin. 105.

Anglis, daisie Ladies-smock.

*Habitat in rupibus et saxosis alpinis. In variis locis jux-
ta Rippon ad monasterium ejus et in Denbighshire.*

Merr. pin. on St. Vincent's-rock, near Bristol. R.

syn. circa Denbeigh. R. it. 4. IV. V.

2. CARDAMINE foliis simplicibus oblongis dentatis. *petræa.*
Sp. pl. 913. Fl. dan. 386. suec. 591.

*Cardamine petræa cambrica, nasturtii facie. Dill.
elth. 70. t. 61. f. 71.*

TETRADYNAMIA SILIQUOSA.

Nasturtium petræum. *Merc. bot. Pluk. ph. t. 101. f. 3. R. syn. 300.*

Anglis, mountain Ladies-smock.

Habitat in excelsis rupibus Arvonæ: in rupibus udis supra lacum Llyn-dû copiose. D. Lhwyd. 4. V. VI.

** *Foliis pinnatis.*

pratense. 3. **CARDAMINE** foliis pinnatis: foliolis radicalibus subrotundis, caulinis lanceolatis. *Sp. pl. 915. integerrimis.*

Cardamine foliis pinnatis, radicalibus subrotundis caulinis linearibus. *Hall. hist. 473.*

Cardamine. *Ger. em. 259. R. syn. 299.*

Nasturtium pratense majus s. Cardamine latifolia, *Park. 825.*

Nasturtium pratense magno flore. *B. pin. 104.*

Flos cuculi. *Dod. pempt. 592.*

Anglis, common Ladies smock, or Cuckow-flower.

Habitat in pratis et pascuis humidiusculis. 4. IV. V.

Pharmac. Cardamines Herba, Flores.

amara. 4. **CARDAMINE** foliis pinnatis, axillis stoloniferis. *Sp. pl. 915.*

Cardamine foliis pinnatis, subrotundis angulosis. *Hall. hist. 474.*

Cardamine flore major elatior. *Tourn. 224. R. syn. 299.*

Nasturtium aquaticum majus et amarum. *B. pin. 104.*

Anglis, bitter Ladies-smock, or Cresse.

Habitat in pratis et pascuis humidis, circa Lewisham D.

Alchorne; by the river side at Harefield, and about

Uxbridge plentifully. Blacks, in salicetis prope Battersea. 4. IV. V.

impatiens. 5. **CARDAMINE** foliis pinnatis, petiolis basi dilatatis dentatis, foliolis lanceolatis dentatis.

Cardamine foliis pinnatis incisiss stipulatis, floribus apetalis. *Sp. pl. 914.*

Cardamine foliis pinnatis, imis obtuse lobatis, caulibus acute dentatis, floribus apetalis. *Hall. hist. 471.*

Cardamine

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

- Nasturtium aquaticum.* 1. **SISYMBRIUM** siliquis declinatis, foliis pinnatis, foliolis subcordatis. *Sp. pl.* 916.
Sisymbrium foliis pinnatis, pinnis subrotundis, spicis sessilibus. *Hall. hist.* 482.
Sisymbrium cardamine s. *Nasturtium aquaticum.* *B. hist.* II. 884. *R. syn.* 300.
Nasturtium aquaticum supinum. *B. pin.* 104.
Nasturtium aquaticum vulgare. *Park.* 1239.
Nasturtium aquaticum s. cratevæ sium. *Ger. em.* 297.
 β *Nasturtium aquaticum* pinnulis paucioribus. *R. syn.* 300.
 γ *Nasturtium aquaticum* foliis minoribus præcotius. *R. syn.* 301.
Anglis, common Water-cresse.
Habitat in aquosis et rivulis. 6. VI. VII.
Pharmac. Nasturtii aquatici Herba.

- Sylvestre.* 2. **SISYMBRIUM** siliquis declinatis, foliis pinnatis, foliolis lanceolatis serratis. *Sp. pl.* 916.
Sisymbrium foliis pinnatis, pinnis dentatis dissitis. *Hall. hist.* 485.
Eruca aquatica. *Ger. em.* 248. *Park.* 1242. *R. syn.* 297.
Eruca sylvestris. *Fuchs.* 263. minor, luteo parvoque flore. *B. pin.* 98.
Anglis, wild Water-cress, or Rocket.
Habitat in palustribus et aquosis et ad ripas fluviorum.
In Tothil-fields. Blacks. ad ripam Thamesis frequens. 4. VI. VII.

- amphibi-um.* 3. **SISYMBRIUM** siliquis declinatis oblongo-ovatis, foliis pinnatifidis serratis. *Sp. pl.* 917.
palustre. α *Sisymbrium* foliis pinnatifidis serratis. *Fl. suec.* 551.
Radicula foliis pinnatis, pinnis dentatis, petalis calyce brevioribus. *Hall. hist.* 487.
Raphanus aquaticus foliis in profundas lacinias divisis. *B. pin.* 97. *pr.* 38. *R. syn.* 301.
aquaticum. β *Sisymbrium* foliis simplicibus dentatis serratis. *Fl. suec.* 550.
Radicula foliis integris aut pinnatifidis, petalis calyce longioribus. *Hall. hist.* 486.
Raphanus aquaticus alter. *B. pr.* 38. *Park.* 1229.
 Rapistrum

- Rapistrum aquaticum. *Tab. ic.* 408. *Ger. em.* 240.
 ♣ Sisymbrium aquaticum foliis variis. *Vaill. paris.* 135. terrestre.
Anglis, amphibious Water-cresse, or Radish.
 Habitat in aquosis et ad ripas fluviorum. 4. VI—
 VIII.

4. SISYMBRIUM foliis runcinatis dentatis nudis cau- *Irio.*
 le lævi, siliquis erectis. *Sp. pl.* 922.
 Erysimum latifolium majus glabrum. *B. pin.* 101.
 Erysimum latifolium neapolitanum. *Park.* 834. *R.*
syn. 298.
Irio lævis Apulas, erucæ folio. *Col. ecphr.* I. 264.
Anglis, broad-leaved Water-cresse, or Hedge Mustard.
 Habitat in ruderatis et muris antiquis; circa Londinum
 frequens. ☉. V—VIII.

5. SISYMBRIUM petalis calyce minoribus, foliis de- *Sophia.*
 composito-pinnatis. *Sp. pl.* 920. *Fl. dan.* 528.
 Sisymbrium pinnis et pinnulis foliorum pinnatis, laci-
 niis linearibus subhirsutis. *Hall. hist.* 484.
 Erysimum Sophia dictum. *R. syn.* 298.
 Sophia chirurgorum. *Ger. em.* 1068. *Park.* 830.
 Nasturtium sylvestre tenuissimo divisum. *B. pin.*
 105.
Anglis, Flix-weed.
 Habitat in maceris, ruderatis, frequens. ☉. VII.
Pharmac. Sophiæ Semina.

SINAPIS. 821.

- Cal.* patens. *Cor.* unguis recti. *Glandula* inter fla-
 mina breviora et pistillum, interque longiora et ca-
 lycem.
1. SINAPIS siliquis glabris apice tetragonis racemo ad- *nigra.*
 pressis. *Syst. Nat.* 448. *Sp. pl.* 933. *Hall. hist.*
 465.
 Sinapi rapi folio. *B. pin.* 99.
 Sinapi sativum secundum. *Ger. em.* 243. *R. syn.*
 295.
 Sinapi siliqua latiuscula glabra, semine rufo s. vulgare.
B. hist. II. 885,
Anglis,

Anglis, common Mustard.

Habitat in arvis, aggeribus fossarum et juxta vias. ☉.

VI

Pharmac. Sinapis Semina.

alba.

2. SINAPIS filiquis hispidis: rostro obliquo longissimo.

Sp. pl. 933. Hall. hist. 466.

Sinapi apii folio. B. pin. 99.

Sinapi album. Ger. em. 244. filiqua hirsuto, semine albo vel rufo. B. hist. II. 856. R. syn. 295.

Anglis, white Mustard.

Habitat in agris et ad vias. ☉. VIII.

Pharmac. Sinapis albi Semina.

arvensis.

3. SINAPIS filiquis multangulis toroso-turgidis rostro longioribus. *Sp. pl. 933. Hall. hist. 467.*

Rapistrum arvorum. Ger. em. 233. Park. 862. R. syn. 295.

Rapistrum flore luteo. B. pin. 95. hist. II. 844.

Anglis, wild Mustard, or Charlock.

Habitat in arvis et marginibus agrorum. ☉. V.

BUNIAS. 823.

Siliqua decidua, tetraëdra, angulis inæqualibus acuminatis, muricata.

cakile.

1. BUNIAS filiquis ovatis lævibus ancipitibus. *Sp. pl. 936.*

Crambe foliis succulentis linearibus pinnato-dentatis.

Sauv. monf. 280.

Cakile quibusdam, aliis Eruca marina et Raphanus marinus. B. hist. II. 867. R. syn. 307.

Eruca marina. Ger. em. 284. anglica. Park. 821.

Eruca maritima italica, filiqua hastæ cuspidi simili. B. pin. 99.

Anglis, sea Bunias, or Rocket.

Habitat in littoribus maritimis passim. ☉. VI.

ISATIS. 824.

Siliqua lanceolata, unilocularis, monosperma, decidua, bivalvis: valvulis navicularibus.

*1. ISATIS

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Classis XVI.

MONADELPHIA
DECANDRIA.

GERANIUM. 832.

Monogynia. *Stigmata* quinque. *Caps.* rostrata, pentacocca.

* *Staminibus* quinque *fertilibus.*

sicutari-
um.

1. GERANIUM pedunculus multifloris, floribus pentandris, foliis pinnatis incisus obtusis. *Sp. pl.* 951. *Burm. ger.* 33.

Geranium petiolis multifloris, caule procumbente, foliis duplicato-pinnatis pinnulis acute incisus. *Hall. hist.* 944.

Geranium cicutæ folio inodorum. *Ger. em.* 945. *R. syn.* 357.

Geranium cicutæ folio, minus et supinum. *B. pin.* 319.

Geranium moschatum inodorum, *Park.* 708.

β Geranium inodorum album. *Ger. em.* 946.

γ Geranium pimpinellæ folio. *Cat. gifs.* 173.

Anglis, hemlock Cranebill.

Habitat in arenosis ruderatis et muris. ☉. IV—VII.

moscha-
tum.

2. GERANIUM pedunculis multifloris, floribus pentandris, foliis pinnatis incisus, cotyledonibus pinnatifidis. *Sp. pl.* 951. *Ger. em.* 29. *Jac. Hort. vind.* I. 22. t. 55.

Geranium petiolis multifloris, caule diffuso, foliis ovatis pinnatifidis. *Hall. hist.* 945.

Geranium

- Geranium moschatum. *Ger. em.* 941. *Park.* 706.
Riv. pent. irr. 110. *R. syn.* 358.
- Geranium cicutæ folio, moschatum. *B. pin.* 319.
Anglis, musk Cranesbill, or Muscovy.
Habitat in pratis siccioribus in comitatibus Westmorlan-
dico et Eboracensi passim. ☉. V. VI.
Pharmac. Geranii Moschati *Herba.*
3. GERANIUM pedunculis subtrifloris, floribus pentan- *mariti-*
 dris, foliis cordatis incisissimis crenatis scabris, caulibus *num.*
 procumbentibus. *Sp. pl.* 951.
 Geranium pusillum supinum maritimum althææ aut
 betonicæ folio nostras. *R. syn.* 357. *Pluk. ph. t.* 31.
f. 4.
 Geranium minimum anglicum procumbens, foliis be-
 tonicæ. *Hist. Ox. II.* 512. *f.* 5. *t.* 35. *f.* 10.
Anglis, sea Cranesbill.
Habitat in littoribus maritimis arenosis frequens. ☉.
 VI. VII.
- *** *Staminibus decem fertibus.*
4. GERANIUM pedunculis bifloris, foliis caulinis tri- *nodosum.*
 lobis integris serratis : summis subsessilibus. *Sp. pl.*
 953. *Burm. ger.* 7.
 Geranium V. nodosum plateau. *Clus. hist. II.* 101.
Ger. em. 947. *R. syn.* 361.
 Geranium nodosum. *B. pin.* 318. *Park.* 704.
Anglis, knotty Cranesbill.
Habitat in montosis Cumberlandiæ D. Archergeren. *R.*
syn. 4. VII. VIII.
5. GERANIUM pedunculis bifloris foliisque alternis, *phæum.*
 calycibus subaristatis, caule erecto, petalis undula-
 tis. *Sp. pl.* 953. *Burm. Ger. II.*
 Geranium foliis rugosis hirsutis semiseptilobis, lobis
 dentatis, petalis serratis acuminatis. *Hall. hist.* 934.
 Geranium montanum fuscum. *B. pin.* 318. *R. syn.*
 361.
 Geranium maculatum sive fuscum. *Ger. em.* 945.
 Geranium pullo flore. *Park.* 704.
Anglis, spotted Cranesbill.

Habitat

Habitat in pascuis montosis. Found at Tovel, in the valley by the fulling-mill, by Mr. Drayton, Apothecary of Maidstone. R. syn. circa Clapham et Ingleton in agro Eboracensi. 4. V. VI.

*florati-
cum.*

6. GERANIUM pedunculis bifloris, foliis subpeltatis quinquelobis inciso-ferratis, caule erecto, petalis emarginatis. *Sp. pl. 954. Burm. ger. 12. Fl. dan. 124.*

Geranium caule erecto, foliis rugosis septilobis, lobis acute ferratis, petalis emarginatis. *Hall. hist. 932.*

Geranium batrachioides montanum nostras. *Ger. em. 942. R. syn. 360.*

Geranium batrachioides minus. *Park. 704.*

Geranium batrachioides folio aconiti. *B. pin. 317. Anglis, wood Cranesbill.*

Habitat in dumetis et pascuis montosis in agris Westmorlandico, Cumberlandico, et Eboracensi frequens. 4. VL VII.

pratense.

7. GERANIUM pedunculis bifloris, foliis subpeltatis multipartitis rugosis acutis, petalis integris. *Sp. pl. 954. Burm. ger. 13.*

Geranium caule erecto, foliis rugosis hirsutis multilobis, lobis trifidis, lobulis semipinnatis, floribus umbellatis. *Hall. hist. 931.*

Geranium batrachioides. *Ger. em. 922. R. syn. 360. flore cæruleo. Park. 704.*

Geranium batrachioides, Gratia Dei Germanorum. *B. pin. 318.*

Anglis, meadow Cranesbill.

Habitat in pratis et pascuis subhumidis prope Battersea. R. syn. 4. VI. VII.

*pyrenai-
cum.*

8. GERANIUM pedunculis bifloris, petalis bilobis, foliis inferioribus quinquepartito-multifidis, superioribus trilobis, calycibus apice subglandulosis, caule erecto.

Geranium pedunculis bifloris, petalis bilobis, calycibus apice glandulosis, antheris exterioribus sterilibus. *Lin. Mantiss. 1. 97.*

Geranium pedunculis bifloris, foliis multifidis; laciniis

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Pes columbinum. *Dod. pempt.* 61.

Anglis, dovesfoot Cranebill.

Habitat in muris, tectis et pascuis solo arenoso circa Bath et Bristol copiose; inter Battersea et Wandsworth; et inter Mortlack et Kew. ☉. VII.

Pharmac. Geranii columbini *Herba.*

lucidum. 11. GERANIUM pedunculis bifloris, calycibus pyramidatis angulatis elevato-rugosis, foliis quinquelobis rotundatis. *Sp. pl.* 955. *Fl. dan.* 218. *Burm. ger.* 19.

Geranium foliis subrotundis semiquinquelobis, lobis obtusis, calycibus transversim rugosis. *Hall. hist.* 942.

Geranium saxatile. *Ger. em.* 938. *Park.* 707. *R. syn.* 361.

Geranium lucidum saxatile. *B. pin.* 318.

Anglis, shining Cranebill.

Habitat in muris, tectis, umbrosis arenosis. Inter Mortlack et Kew. ☉. VI—VIII.

columbinum. 12. GERANIUM pedunculis bifloris folio longioribus, foliis quinquepartito multifidis, laciniis acutis, arillis glabris, calycibus aristatis. *Sp. pl.* 956. *Burm. ger.* 22.

Geranium caule procumbente, foliis quinquepartitis lobis trilobis, lobulis trifidis. *Hall. hist.* 938.

Geranium columbinum, dissectis foliis, pediculis florum longissimis. *R. syn.* 359. *Vaill. par.* 79. t. 15. f. 4.

Geranium columbinum annum minus, folio tenuius laciniato, flore pediculo longissimo insistente. *Hort. Ox. II.* 512. f. 5. t. 15. f. 3.

Anglis, long-stalked Cranebill.

Habitat in arvis, pascuis et ad sepes. ☉. VI—VIII.

dissectum. 13. GERANIUM pedunculis bifloris, foliis quinquepartito-trifidis, petalis emarginatis longitudine calycis, arillis villosis. *Sp. pl.* 956. *Burm. ger.* 15.

Geranium caule erecto, foliis quinquelobis, lobis trifidis acutis. *Hall. hist.* 937.

Geranium

Geranium columbinum majus dissectis foliis. *Ger. em.* 938. *R. syn.* 359.

Geranium malacoides s. columbinum alterum. *Park.* 706.

Geranium columbinum majus foliis imis longe usque ad pediculum divisis. *Hist. Ox.* II. 511. *Vaill. par.* 79. t. 15. f. 3.

β Geranium columbinum maximum foliis dissectis. *Plot. Ox.* 145. t. 9. f. 3. *R. syn.* 340.

Anglis, jagged Cranesbill.

Habitat in pascuis, arvis et ad sepes. ☉. V—VIII.

14. GERANIUM pedunculis bifloris, calycibus pilosis *robertianum.* decemangularibus. *Sp. pl.* 955. *Burm. ger.* 18.

Geranium foliis duplicato-pinnatis, pinnis ultimis confluentibus, calycibus striatis, hirsutis. *Hall. hist.* 943.

Geranium robertianum. *Ger. em.* 939. *Dod. pempt.* 69. *B. pin.* 319. *R. syn.* 358. vulgare. *Park.* 710.

β Geranium lucidum saxatile, foliis geranii robertiani. *R. syn.* 358.

Anglis, stinking Cranesbill, or Herb-Robert.

Habitat in ruderatis, maris, saxosis. ☉. IV—VIII.

Pharmac. Geranii robertiani *Herba.*

15. GERANIUM pedunculis unifloris, foliis quinque *sanguineum.* partitis trifidis orbiculatis. *Sp. pl.* 958. *Fl. suec.* 616. *Burm. ger.* 3.

Geranium pedunculis unifloris, foliis septilobis, petalis bifidis. *Hall. hist.* 930.

Geranium sanguineum. *Ger. em.* 955. maximo flore. *B. pin.* 318.

Geranium hæmatodes. *Park.* 705. *R. syn.* 360.

β Geranium hæmatodes, foliis majoribus pallidioribus et altius incis. *R. syn.* 360.

γ Geranium hæmatodes Lancastriense flore eleganter striato. *R. syn.* 360. *Dill. elth.* 163. t. 136. f. 163.

Anglis, bloody Cranesbill.

X

Habitat

MONADELPHIA POLYANDRIA.

Habitat in rupibus calcareis, saxosis et pascuis siccioribus.
 ⁊ in Insula Walney. 4. VII. VIII.

POLYANDRIA.

ALTHÆA. 839.

Cal. duplex: exterior novemfidus. *Arilli* plurimi monospermi.

officinalis. 1. ALTHÆA foliis simplicibus tomentosis. *Sp. pl.* 966. *Fl. dan.* 530. *Gort. gelr.* 410.

Althæa tomentosa, herbacea, caule erecto, foliis cordato-lanceolatis. obsolete trifido. *Hall. bist.* 1074.

Althæa vulgaris. *Park.* 303. *R. syn.* 252.

Althæa ibiscus. *Ger. em.* 933.

Althæa dioscoridis et plinii. *B. pin.* 315.

β Althæa vulgari similis folio retuso brevi. *R. syn.* 252.
Anglis, marsh Althæa, or Mallow.

Habitat in palustribus maritimis. 4. VIII.

Pharmac. Althæa Radix, Herba, Semina.

LAVATERA. 842.

Cal. duplex: exterior trifidus. *Arilli* plurimi monospermi.

arborea. 1. LAVATERA caule arborea, foliis septemangularibus tomentosis plicatis, pedunculis confertis unifloris axillaribus. *Sp. pl.* 972. *Hort. ups.* 202.

Malva arborea minima nostras. *Merr. R. syn.* 252.

Malva arborea veneta dicta, parvo flore. *B. pin.* 315.
Anglis, tree Lavatera, or sea-tree Mallow.

Habitat in littoribus maritimis. At Hurst-castle, over against the Isle of Wight; in Portland island; on the rocks of Caldey island and Basse island, near Edinburgh. *R. syn.* In Cornubia et Devoniam passim. 8. VII. VIII.

MALVA. 841.

Cal. duplex: exterior triphyllus. *Arilli* plurimi, monospermi.

1. MALVA

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

MONADELPHIA POLYANDRIA.

Alcea vulgaris major. *B. pin.* 316.

Alcea vulgaris f. *Malva verbenacea.* *Park.* 301.

Malva verbenacea. *Ger. em.* 930.

Anglis, vervain Mallow.

Habitat in sepibus et ad margines agrorum in comitatibus
Warwicensi, Leicestriensi et Nottinghamensi. 4.
VII. VIII.

Pharmac. Alceæ Herba.

mōschata. 5. *MALVA* foliis radicalibus reniformibus incis, cau-
linis quinquepartitis pinnato multifidis. *Sp. pl.* 971.
Hall. hist. 1072.

Malva montana f. *Alcea rotundifolia laciniata.* *Col.*
ecphr. I. 148. t. 147. f. 147.

Alcea folio rotundo laciniato. *B. pin.* 316.

Alcea tenuifolia crispa. *B. hist.* II. ap. 1067. *R. syn.*
353.

Anglis, musk Mallow.

Habitat in pratis, pascuis et ad sepes. 8. VII. VIII.

Classis XVII.

DI A D E L P H I A

H E X A N D R I A.

F U M A R I A. 849.

Cal. diphyllus. *Cor.* ringens. *Filamenta* duo, membranacea, singula.

1. FUMARIA pericarpis monospermis racemosis, caule *officinalis*.
diffuso. *Sp. pl.* 984. *Fl. suec.* 630.

Fumaria foliis multifidis, lobis subrotunde lanceolatis, fructibus monospermis. *Hull. hist.* 346.

Fumaria. *Matth.* 1158. *Fuchs.* 338. *Cam. ep.* 890. vulgaris. *B. hist.* III. 101. *Park.* 287. *R. syn.* 204.

Fumaria purpurea. *Ger. em.* 1088.

Fumaria officinarum et Dioscoridis. *B. pin.* 143.

β Fumaria pericarpis monospermis racemosis, foliis capreolata scandentibus subcirrhosis. *Sp. pl.* 984. ta.

Fumaria major scandens, flore pallidior. *R. syn.* 204.

Anglis, common Fumitory.

Habitat in arvis et sepibus frequens. ☉. V.—VIII.

Pharmac. Fumarizæ Herba, Semina.

Obt. nullam differentiam inter α et β observavi; nisi in colore florum; Flos albus cum apice purpureo in β est; nunquam folia cirrhosa inveni; sed caules scandentes et petioli contorti sive volubiles sunt.

γ FUMARIA siliquis linearibus, petiolis cirrhiferis. *claviculata*

Fumaria siliquis linearibus, foliis cirrhiferis. *Sp. pl.* ta. 985. *Fl. dan.* 340.

Fumaria alba latifolia. *Park.* 288. *R. syn.* 335. claviculata. *Ger. em.* 1088.

DIADELPHIA DECANDRIA.

Fumaria clavicularis donata. *B. pin.* 143. *Hist. Ox. II,*
'266. *f.* 3. *t.* 12. *f.* 3.

Anglis, climbing Fumitory.

Habitat in sylvis, sepibus subhumidis et arenosis. ☉. VI,
VII.

OCTANDRIA.

POLYGALA. 851.

Cal. pentaphyllus: foliolis duobus alæformibus, coloratis. *Legumen* obcordatum, biloculare.

vulgaris. 1. POLYGALA floribus cristatis racemosis, caulibus herbaceis simplicibus procumbentibus, foliis linearilanceolatis. *Sp. pl.* 986. *Fl. dan.* 516.

Polygala caule erecto, foliis linearibus acutis. *Hall. hist.* 344.

Polygala. *Ger. em.* 563. *R. syn.* 287. minor. *Park.* 1332.

Polygala vulgaris. *B. pin.* 215

β *Polygala major.* *B. pin.* 215.

γ *Polygala violacea.* *Boerb. ind. alt.* 236.

δ *Polygala carnea.* *Boerb. ind. alt.* 236.

ε *Polygala alba.* *Tourn.* 174.

ζ *Polygala myrtifolia palustris humilis et ramosior.* *R. syn.* 287.

Anglis, common Milkwork.

Habitat in pascuis et ericetis. ♀. VI. VII.

DECANDRIA.

SPARTIUM. 858.

Stigma longitudinale, supra villosum. *Filamenta* germini adhærentia. *Cal.* deorsum productus.

scoparium. 1. SCOPARIUM foliis ternatis solitariisque, ramis infermibus angulatis. *Sp. pl.* 996. *Fl. dan.* 313.

Spartium foliis inferioribus ternatis hirsutis, superioribus simplicibus. *Hall. hist.* 354.

Genista angulosa et scoparia. *B. pin.* 395.

Genista.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Genista minor aspalathoides, f. *Genista spinosa anglica*. *B. pin.* 395. *R. syn.* 475.

Genista aculeata. *Ger. em.* 1320.

Genistella aculeata. *Park.* 1005. *Lob. ic.* 2. t. 93. *Anglis*, needle Greenweed, Furze, or Petty-whin.

Habitat in ericetis et pascuis subhumidis. $\frac{1}{2}$. V. VI.

U L E X. 881.

Cal. diphyllus. *Legumen* vix calyce longius.

europæus. 1. ULEX foliis villosis acutis, spinis sparsis. *Sp. pl.* 1043. *Fl. dan.* 608.

Genista spinosa vulgaris. *Ger. em.* 1319. *R. syn.* 475. f. *Scoparius Theophrasti*. *Park.* 1003.

Genista spinosa major, longioribus aculeis. *B. pin.* 394.

β *Genista spinosa minor*. *Park.* 1003. *Anglis*, common Furze, Whin or Gorse.

Habitat in ericetis et pascuis. $\frac{1}{2}$. IV.—X.

O N O N I S. 863.

Cal. quinquepartitus: laciniis linearibus. *Vexillum* striatum. *Legumen* turgidum sessile. *Filamenta* connata absque fissura.

spinosa. 1. ONONIS floribus racemosis subsolitariis, foliis ternatis: superioribus solitariis nudis, ramis spinosis glabris.

Anonis floribus racemosis solitariis, foliis ternatis, ramis spinosis. *Syst. nat. ed. XI.* 1156. *Sp. pl.* 1006.

Anonis caule decumbente, ramis senescentibus spinosis. *Hall. hist.* 356.

Anonis spinosa flore purpureo. *B. pin.* 389. *Park.* 994. *R. syn.* 332.

Anonis. *Fuchs. hist.* 60. f. *Resta bovis*. *Ger. em.* 1323.

Anglis, common Restharrow or Cammock.

Habitat in pascuis sterilioribus. $\frac{1}{2}$. VI—VIII.

Pharmac. *Anonidis Radix*.

inermis. 2. ONONIS floribus racemosis subgeminis, foliis ternatis: superioribus solitariis, ramisque villosis inermibus.

Ononis

Ononis floribus racemosis geminis, foliis ternatis : arvensis. superioribus solitariis, ramis inermibus subvillosis.

Sp. pl. 1006.

Anonis non spinosa purpurea. *Ger. em.* 1323. *Park.* 993. *R. syn.* 332.

Anonis spinis carens purpurea. *B. pin.* 389.

β Ononis caulibus diffusis, ramis erectis : foliis superioribus solitariis, stipulis ovatis. *Sp. pl.* 1006.

Anonis maritima procumbens, foliis hirtutis pubescentibus. *R. syn.* 332. *Dill. elth.* 29. t. 25. f. 28.

Anglis, hairy Restharrow.

Habitat in arvis, pratis et pascuis et littoribus maritimis arenosis frequens. h. VI.—VIII.

Obs. α et β in horto cultæ per quatuor annos, nullam specificam differentiam præbuerunt. Plantæ villosæ subviscidæ. Floribus solitariis, geminatis et subinde ternatis.

ANTHYLLIS. 864.

Cal. ventricosus. *Legumen* subrotundum, tectum.

1. ANTHYLLIS herbacea, foliis pinnatis inæqualibus, vulnera- capitulo duplicato. *Sp. pl.* 1012. *ria.*

Anthyllis leguminosa. *Ger. em.* 1240. vulgaris. *Park.* 1093.

Vulneraria foliis radicalibus ovatis, caulinis pinnatis, umbellis ovatis. *Hall. hist.* 398.

Vulneraria rustica. *B. hist.* II. 362. *R. syn.* 323.

Loto affinis Vulneraria pratensis. *B. pin.* 332.

β Vulneraria supina flore coccineo. *R. syn.* 323. *Dill. elth.* 431. t. 320. f. 413.

Anglis, common Ladies-finger or Kidney-vetch.

Habitat in pratis et pascuis cretaceis et calcareis β. in In- sula Anglesea. 4. VII.

PISUM. 870.

Stylus triangulus, supra carinatus pubescens. *Calycis* lacinia superiores duæ breviores.

1. PISUM petiolis supra planiusculis, caule angulato, mariti- stipulis sagittatis, pedunculis multifloris. *Sp. pl. mum* 1027. *Fl. dan.* 338. *It. W-goth.* 190.

Pisum

DIADELPHIA DECANDRIA.

Pisum marinum. *Ger. em.* 1250. *R. syn.* 319.
Pisum spontaneum maritimum anglicum. *Park.* 1059.
Pisum spontaneum perenne repens humile. *Hist. Ox.*
II. 47. f. 2. t. 1. f. 5.
Anglis, sea Pea.
Habitat in littoribus maritimis. On the long balck of
stones running from Aldburgh towards Orford in Suff-
olk; near Hastings in Suffex; among the stones on
the west side of Dengeness, near Lyd in Kent. R.
syn. 4. VIII.

O R O B U S. 871.

Stylus linearis. Cal. basi obtusus: laciniis superiori-
bus profundioribus, brevioribus.

tuberosus. 1. OROBUS foliis pinnatis lanceolatis, stipulis semisa-
gittatis, caule simplici. Sp. pl. 1028.
Orob. caule simplici, foliis senis ellipticis, radice tu-
berosa. Hall. hist. 417.
Orob. sylvaticus foliis oblongis glabris. Tourn. 393.
R. syn. 324.
Astragalus sylvestris foliis oblongis glabris. B. pin.
351.
Astragalus sylvaticus. Ger. em. 1237.
Lathyrus sylvestris lignosior. Park. 1061.
Anglis, common Orobus or Heath-pea.
Habitat in pratis, pascuis, sylvis et ad sepes. 4. V.

sylvaticus. 2. OROBUS caulibus decumbentibus hirsutis ramosis.
Cent. pl. 67. Sp. pl. 1029.
Orob. sylvaticus nostras. R. syn. 324. hist. 1892.
Aët. paris. 1706. 87. t. 90.
Anglis, wood Orobus or bitter Vetch.
Habitat in sylvis et sepibus. At Gambleby in Cumber-
land, about six miles from Pereth in the way to New-
castle, observed by Tho. Willisel below Brecknock
hills in the way to Caerdyff, and in Merionethshire,
near Bala. R. syn. in Scotia frequens. D. Hope.
4. V. VI.

LATHY-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Essex. R. syn. *In the border of a field between the road and Munden church; likewise among the bushes within the gate leading to Laydon-hall, both in Dengy hundred, Essex, Mr. S. Dale. Blackf. Spec. bot. ©. VII.*

Obs. *variat pedunculis uni et trifloris.*

*** *Pedunculis multifloris.*

latifolius. 4. LATHYRUS pedunculis multifloris, cirrhis diphyllis: foliolis lanceolatis, internodis membranaceis, *Sp. pl.* 1033. *Fl. suec.* 645.

Lathyrus caulibus alatis, foliis ovato-lanceolatis rigidis, scapis multifloris. *Hall. hist.* 433.

Lathyrus major latifolius. *Ger. em.* 1229. *R. syn.* 319.

Lathyrus latifolius. *B. pin.* 344.

Lathyrus major perennis. *Park.* 1061.

Anglis, broad-leaved Lathyrus.

Habitat *in sylvis et sepibus in Madingley and other woods near Cambridge; Mr. Lawson observed it in the rocks near Red Neele by Whitehaven, plentifully. R. syn.* 4. VII. VIII.

sylvestris. 5. LATHYRUS pedunculis multifloris, cirrhis diphyllis: foliolis ensiformibus. *Sp. pl.* 1033. *Fl. dan.* 325.

Lathyrus caule alato; foliis geminis ensiformibus. *Hall. hist.* 434.

Lathyrus sylvestris. *Dod. pempt.* 523. *Clus. hist.* II. 129. *Park.* 1061. major. *B. pin.* 344.

Lathari majoris species flore rubente et albido minore dumetorum s. Germanicus. *B. hist.* II. 129. *R. syn.* 319.

Anglis, narrow-leaved or wild Lathyrus.

Habitat *in sylvis et sepibus; Mr. Dale observed it in the hedges between Castle-Campes and Bartlow, in Cambridgeshire; inter Rochester et Northfleet in Cantio; inter Bath et Bristol copiose; circa Conway in Wallia. Æ. VII. VIII.*

6. LATHYRUS pedunculis multifloris, cirrhis poly-*palustris*.
 phyllis; stipulis lanceolatis. *Sp. pl.* 1034. *Fl. dan.*
 399.
 Lathyrus foliis senis, floribus racemosis. *Hall. hist.*
 431.
 Lathyrus peregrinus, foliis viciæ, flore subcæruleo
 pallidive purpurascente. *B. pin.* 344.
 Lathyrus viciæ formis f. *Vicia lathyroides nostras.* *R.*
syn. 320.
Vicia Lathyroides f. *Lathyrus viciæ formis.* *Pluk.*
ph. t. 71. *f.* 2.
Anglis, marsh Lathyrus.
 Habitat in pratis et pascuis humidis; in a wood near Ab-
 ington, Berkshire. Blackf. upon the bogs in Char-
 ley forest, near Bardon-hill, Leicestershire. *D.*
 Pultney; near Ranaugh in Norfolk. Mr. Hum-
 phry; in comitatibus Lancastriensi et Eboracensi
 passim. 4. VII. VIII.
7. LATHYRUS pedunculis multifloris, cirrhis diphyllis *pratensis*.
 simplicissimis: foliolis lanceolatis. *Sp. pl.* 1033.
Fl. dan. 527.
 Lathyrus scapis multifloris, foliis lanceolatis, capreo-
 lis simplicibus. *Hall. hist.* 436.
 Lathyrus sylvestris flore luteo. *Ger. em.* 1231. *Park.*
 1062.
 Lathyrus sylvestris luteus. *B. pin.* 344.
 Lathyrus luteus sylvestris dumetorum. *B. hist.* II.
 304. *R. syn.* 320.
Anglis, meadow Lathyrus.
 Habitat in pratis, pascuis, sylvis. 4. VII. VIII.

V I C I A. 873.

Stigma latere inferiore transverse barbatum.

* *Pedunculis elongatis.*

1. VICIA pedunculis multifloris, floribus imbricatis, *cracca*.
 foliolis lanceolatis pubescentibus, stipulis integris.
Sp. pl. 1035. *Fl. succ.* 652.

Vicia

DIADELPHIA DECANDRIA:

Vicia foliis lanceolatis sericeis, racemis multifloris reflexis, stipulis integerrimis. *Hall. hist.* 424.

Vicia multiflora. *B. pin.* 345. f. *spicata.* *Park.* 1072.

Cracca. *Riv.* 49. *Rupp. jen.* 212. *R. syn.* 322.
Anglis, tufted Vetch.

Habitat in nemorosis et sepibus. 4. VII. VIII.

sylvatica. 2. *VICIA* pedunculis multifloris, foliolis ovalibus, stipulis denticulatis. *Sp. pl.* 1035. *Fl. dan.* 277.

Vicia foliis ovatis, stipulis argute dentatis, siliquis racemosis pendulis. *Hall. hist.* 426.

Vicia sylvatica multiflora. *Ph. br. R. syn.* 322.

Vicia multiflora maxima perennis, tetro odore, floribus albescentibus lineis cæruleis striatis. *Pluk. ph. t.* 41. f. 1.

Anglis, wood Vetch.

Habitat in sylvis et sepibus. *In a wood near Caerwent, Monmouthshire; in the woods about Greta bridge, Yorkshire, Mr. Johnson; in Smokhall wood, Bath, and about the Devises, Wiltshire, Mr. Goodyer; supra pontem Kirby Londale, in Westmorlandia. Dr. Richardson. R. syn.* 4. VII. VIII.

** *Floribus axillaribus sessilibus.*

fativa.

3. *VICIA* leguminibus sessilibus erectis, foliis retusis, stipulis notatis. *Sp. pl.* 1037. *Fl. dan.* 522.

Vicia. *Ger. em.* 1227. *R. syn.* 320. *vulgaris fativa.* *Park.* 1072.

Vicia fativa vulgaris, semine nigro. *B. pin.* 344.

β *Vicia fativa vulgaris*, semine albo. *B. pin.* 344.

Anglis, common Vetch or Tare.

Habitat in agris et inter segetes. ☉. V. VI.

Pharmac. Viciæ semine.

Obs. variat foliis superioribus lanceolatis; stipulæ maculatae, macula atro-rubescente.

latbyroides.

4. *VICIA* leguminibus sessilibus subsolitariis erectis glabris, foliolis inferioribus retusis, superioribus linearilanceolatis. *Fl. dan.* 58.

. *Vicia*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Vicia grandiflora. Scop. carn. 65. t. 40.

Anglis, bastard Vetch.

Habitat in pratis et pascuis; circa Weymouth in agro Dorsetiensi. ☉. VII. VIII.

sepium. 7. VICIA leguminibus pedicellatis subquaternis erectis, foliolis ovatis integerrimis: exterioribus decrescen-
tibus. *Sp. pl.* 1038. *Fl. suec.* 651.

*Vicia foliis ovatis, oblongis, scapis brevibus quadri-
floris.* Hall. hist. 429.

Vicia maxima dumetorum. Ger. em. 1227. Park. 1072.

Vicia sepium folio rotundiore acuto. B. pin. 345.

Vicia sepium perennis. B. hist. II. 713. R. syn. 320.

β *Vicia folio subrotundo, brevi obtuse mucronato, pe-
dunculo brevi infidente, flore viciæ sepium seu du-
metorum vulgaris.* R. syn. 320.

Anglis, bush Vetch.

Habitat in sylvis, sepibus, pascuis et pratis. ♀. V.

bithynica. 8. VICIA leguminibus pedunculatis solitariis erectis, foliolis quaternis ovali-lanceolatis, stipulis denta-
tis. *Sp. pl.* 1038. *Jac. hort. vind.* II. 69. t. 147.

*Lathyrus pedunculis unifloris, cirrhis tetraphyllis sti-
pulis dentatis.* Roy. lugdb. 363.

*Clymenum bithynicum, siliqua singulari, flore mi-
nore.* Boerh. ind. alt. 43.

Anglis, rough Vetch.

Habitat in arvis et pascuis glareosis, et ad sepes prope Doncaster in agro Eboracensi, D. Tosfield. in insu-
la Purbeck. ♂. VII.

ERVUM. 874.

Cal. quinquepartitus, longitudine corollæ.

tetrasper- 1. ERVUM pedunculis subbifloris, seminibus globosis
mum. quaternis. *Sp. pl.* 1039. *Fl. dan.* 95.

Vicia foliis linearibus, siliquis gemellis glabris. Hall. hist. 423.

Vicia segetum singularibus siliquis glabris. B. pin. 345.

Vicia minor segetum, cum siliquis paucis glabris. Hist. Ox. II. 64. f. 2. t. 4. f. 16.

Cracca minor siliquis singularibus, flosculis cæruleo-centibus. *Hoff. cat. h. alt. R. syn. 322.*

Anglis, smooth Tare.

Habitat in arvis, sepibus et ad margines agrorum. ☉. VI.

2. ERVUM pedunculis multifloris, seminibus globosis *hirsutum* binis. *Sp. pl. 132.*

Vicia foliis linearibus, siliquis racemosis dispermis, hirsutis. *Hall. hist. 422.*

Vicia segetum cum siliquis plurimis hirsutis. *B. pin. 345.*

Vicia sylvestris s. Cracca minima. *Ger. em. 1228.*

Aracus s. Cracca minor. *Park. 1070.*

Cracca minor. *Riv. tetr. 55. R. syn. 322.*

Anglis, hairy Tare.

Habitat in arvis et pratis arenosis. ☉. VI.

ORNITHOPUS 884.

Legumen articulatum, teres, arcuatum.

1. ORNITHOPUS foliis pinnatis, leguminibus incur-^{perpusil-}vis. *Syst. nat. 491. Sp. pl. 1049.* ^{ius.}

Ornithopodium caule prostrato, foliis pinnatis, siliquis umbellatis, articulatis. *Hall. hist. 393.*

Ornithopodium minus. *Ger. em. 1241. B. pin. 350.*

Ornithopodium radice nodosa. *Park. 1093. R. syn. 326.*

Ornithopodium majus. *B. pin. 350. Park. 193.*

Anglis, common Bird's-foot.

Habitat in pascuis et arenosis. ☉. V.—VIII.

HIPPOCREPIS. 885.

Legumen compressum, altera sutura pluries emargina- ^{comosa.} *tum, curvum.*

1. HIPPOCREPIS leguminibus pedunculatis confertis arcuatis: margine exteriori repandis. *Sp. pl. 1050.*

Ferrum equinum germanicum siliquis in summitate.

B. pin. 346. R. syn. 325.

Y

Ferrum

DIADELPHIA DECANDRIA.

Ferrum equinum comosum. *Park.* 1091. f. capitatum. *Col. ephr.* l. 302. t. 301.

Anglis, tufted Horfeshoe-vetch.

Habitat in pratis et pascuis solo calcareo; on the chalky grounds about Gogmagog hills, and elsewhere in Cambridgeshire; in collibus cretaceis in comitatibus Cantiano, Surriensi, Bedfordiensi, &c. passim. *8.* VII.

HEDYSARUM. 887.

Cor. carina transverse obtusa. *Legumen* articulis monospermis.

onobrychis.

1. HEDYSARUM foliis pinnatis, leguminibus monospermis aculeatis, corollarum alis calycem æquantibus, caule elongato. *Sp. pl.* 1059.

Onobrychis caule erecto ramoso, floribus spicatis. *Hall. hist.* 396.

Onobrychis. *Dod. cereal.* 166. *hist.* 558. f. caput galinaceum. *Ger. em.* 1243. *R. syn.* 327.

Onobrychis vulgaris. *Park.* 1082.

Onobrychis foliis viciæ, fructu echinato major. *B. pin.* 350.

Polygalon Gesneri. *B. hist. II.* 335.

Anglis, common Saint-Foin or Cock's-head.

Habitat in pratis et pascuis cretaceis. *4.* VII.

ASTRAGALUS. 892.

Legumen biloculare, gibbum.

glycyphyllos. 1. ASTRAGALUS caulescens prostratus, leguminibus subtriquetris arcuatis, foliis ovalibus pedunculo longioribus. *Sp. pl.* 1067.

Astragalus caule prostrato foliis ovatis, scapis brevibus racemosis, siliquis subtriquetris incurvis. *Hall. hist.* 413.

Astragalus luteus perennis procumbens, vulgaris, f. sylvestris. *Hist. Ox. II.* 107. *R. syn.* 326.

Hedysarum glycyrrhizatum. *Ger. em.* 1233.

Glycyrrhiza sylvestris floribus luteo pallefcentibus. *B. pin.* 352.

Glaux vulgaris f. Glycyrrhiza sylvestris. *Park.* 1098. *Anglis,*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Melilotus filiquis racemosis, pendulis, rugosis, dispersis acutis. *Hall. hist.* 362.

Melilotus officinarum germaniæ. *B. pin.* 331.

Melilotus vulgaris. *Park.* 719. *R. syn.* 331.

Melilotus germaniæ. *Ger. em.* 1205.

Anglis, Melilot Trefoil.

Habitat in arvis et ad sepes. ♂. VI. VII.

Pharmac. Meliloti Herba.

ornithopodioides. 2. TRIFOLIUM leguminibus nudis octospermis subternis calyce duplo longioribus caule prostrato. *Sp. pl.* 1078. *Fl. dan.* 368.

Trifolium filiquis ornithopodii. *R. syn. ed. II.* 195.

Trifolium filiquosum loto affine filiquis ornithopodii. *Pluk. ph. t.* 68. *f.* 1.

Fœnumgræcum humile repens ornithopodii filiquis brevibus erectis. *R. syn.* 331. *t.* 14. *f.* 1.

Anglis, birdsfoot Trefoil.

Habitat in pratis et pascuis arenosis. Found by Tho. Willisel among the corn, half a mile on this side Tadcaster towards Sherborn; also near Oxford; on the sandy banks by the sea side, at Tottlesbury, in Essex, plentifully. Mr. Newton; Mr. Doody in Tothilfields, Westminster, *R. syn.* on Mushold-heath near Norwich, Mr. Pitchford, in ericeto nigro prope Londinum; circa Marazion et Penzance in Cornubia. ©. VI. VII.

** Lotoidea leguminibus tectis polyspermis.

repens. 3. TRIFOLIUM capitulis umbellaribus, leguminibus tetraspermis, caule repente. *Sp. pl.* 108.

Trifolium caule repente, spicis depressis, filiquis tetraspermis. *Hall. hist.* 367.

Trifolium pratense album. *B. pin.* 327. *Park:* 1110. *R. syn.* 327.

Trifolium minus pratense, flore albo s. 2. *Ger. em.* 1185.

hybridum. β Trifolium capitulis umbellaribus, leguminibus tetraspermis caule ascendente. *Sp. pl.* 1099.

Trifolium.

Trifolium caulibus suberectis, spicis depresso, filiquis dispermis. *Hall. hist.* 368.

Trifolium pratense corymbiferum. *Mich. gen.* 28. t. 25. f. 2. 6.

Anglis, white Trefoil, or Dutch Clover.

Habitat in pratis et pascuis ubique, β in pratis humidis; prope Peckham et Battersea observavi. γ . V—IX.

*** Lagopoda calycibus villosis monospermis.

4. TRIFOLIUM spicis villosis globosis, corollis mono-^{ochroleu-}petalis, calycum infimo dente longissimo recto, ^{cum.} caule subdeclinato, folisque villosis infimis obcordatis.

Trifolium spicis villosis, caule erecto pubescente, foliis infimis obcordatis. *Syst. Nat.* III. 233. *Jac. Austr.* I. 26. t. 40.

Trifolium caule erecto, foliis hirsutis, supremis conjugatis, spicis oblongis. *Hall. hist.* 378.

Trifolium pratense hirsutum majus, flore albo-sulphureo seu $\omega\chi\rho\lambda\acute{\epsilon}\kappa\omega$. *R. syn.* 328.

Trifolium lagopoides annum hirsutum pallide luteum seu ochroleucum. *Hist. Ox.* II. 141. f. 2. t. 12. f. 12. *Anglis*, yellow Trefoil.

Habitat in pratis et pascuis cretaceis in comitatibus Essexiensi, Cantabrigiensi, Bedfordiensi, Hertfordiensi copiose; prope Stamford; et supra Duppens Hill, Croydon. δ . VI. VII.

5. TRIFOLIUM spicis globosis villosis, cinctis stipulis ^{pratense,} oppositis membraceis, corollis monopetalis. *Sp. pl.* 1082. *Fl. suec.* 666.

Trifolium caule obliquo, foliis hirsutis supremis conjugatis, vaginis aristatis. *Hall. hist.* 377.

Trifolium pratense. *Dod. hist.* 365. *Cam. ep.* 582. *Ger. em.* 1185. purpureum. *B. pin.* 327. *R. syn.* 328. vulgare. *Park.* 1110.

β Trifolium pratense purpureum minus, foliis cordatis. *R. syn.* t. 13. f. 1.

γ Trifolium purpureum majus sativum, pratense simile. *R. syn.* 328.

Anglis, purple Trefoil, Honeyfuckle, or Clover.

Habitat in pratis et pascuis. ♂. V—VIII.

Pharmac. Trifolii pratensis Herba.

alpestre. 6. TRIFOLIUM spicis subglobosis villosis terminalibus sessilibus, caule erecto, foliis lanceolatis serrulatis. *Sp. pl.* 1082.

Trifolium spicis villosis globosis terminalibus, corollis monopetalis, foliis lanceolatis subintegerrimis, caule erecto. *Fl. angl.* 284.

Trifolium foliis ovatis nervosis, supremis conjugatis, vaginis lanceolatis. *Hall. hist.* 376.

Trifolium montanum purpureum majus. *B. pin.* 328.

Trifolium purpureum majus, foliis longioribus et angustioribus, floribus saturatoribus. *R. syn.* 328.

Anglis, perennial Trefoil, Clover, or Marle-grass.

Habitat in pratis et pascuis montosis. ♀. VII.

stellatum. 7. TRIFOLIUM spicis pilosis globosis, calycibus patentibus æqualibus, caule obliquo foliisque villosis infimis obcordatis.

Trifolium spicis pilosis, calycibus patentibus, caule diffuso, foliolis obcordatis. *Sp. pl.* 1083.

Trifolium stellatum. *B. pin.* 329. *pr.* 143. glabrum.

Ger. em. 1208. *Pluk. pb. t.* 113. *f.* 4. *R. syn.* 329.

Anglis, star Trefoil.

Habitat in pratis et pascuis maritimis. Observed by Dr. Johnson in Dartsford salt-marsh; I found it by the water-side at Leigh and at Little-Holland, Essex; at Tilbury-Fort, Mr. Petiver; plentifully about Sheerness. Mr. Doody. *R. syn. inter Greenhithe et Northfleet et alibi in Cantio; prope Bristolium copiose; ♀. VI. VII.*

arvense. 8. TRIFOLIUM spicis villosis ovalibus, dentibus calycinis setaceis villosis æqualibus. *Sp. pl.* 1083.

Trifolium caule ramoso, foliis lanceolatis, serratis, spicis villosis ovatis. *Hall. hist.* 373.

Trifolium arvense humile spicatum s. Lagopus. *B. pin.* 328. *R. syn.* 330.

Lagopus. *Fuchf.* 494. *Dod. cer.* 212. vulgaris. *Park.* 1107.

Lagopodium sive Pes leporis. *Ger. em.* 1193.

β Lagopus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

subterra- 12. TRIFOLIUM capitulis villosis quinquefloris, invo-
neum. lucro centrali reflexo rigido fructum obvolvente.

Sp. pl. 1087.

Trifolium pumilum supinum flosculis longis albis.

Ph. br. R. syn. 327. t. 13 f. 2.

Trifolium album tricoccum subterraneum reticulatum.

Hist. Ox. II. 138. f. 2. t. 14. f. 5.

Anglis, dwarf Trefoil.

Habitat in pascuis et ericetis sterilioribus circa Londinum frequens. ☉. V.

**** Vesicaria calycibus inflatis ventricosis.

fragife- 13. TRIFOLIUM capitulis subrotundis, calycibus in-
rum. flatis bidentatis reflexis, caulibus repentibus. *Sp. pl.*
1086.

Trifolium caule repente, spicis glabris, calycibus sericeis ampullescentibus. *Hall. hist.* 370.

Trifolium fragiferum. *Ger. em.* 1208. *R. syn.* 329. friscum. *B. pin.* 329. *Clus. curr. app.* 38. t. 39.

Park. 1109.

Trifolium fragiferum purpureum, folio oblongo. *Hist.*

Ox. II. 144. f. 2. t. 13. f. 2. *Vaill. paris.* 195. t. 22. f. 2.

Anglis, strawberry Trefoil.

Habitat in pratis et pascuis humidis. ♃. VIII.

**** Lupulina vexillis corollis reflexis.

agrarium. 14. TRIFOLIUM spicis ovalibus imbricatis; vexillis deflexis persistentibus, calycibus nudis, caule erecto.

Sp. pl. 1087. *Fl. dan.* 558.

Trifolium spicis ovatis densissimis, strepentibus, caulibus diffusis. *Hall. hist.* 363.

Trifolium luteum minimum. *Ger. em.* 1186. *Lob. ic.* 29. *Park. III.*

Trifolium pratense luteum, capitulo lupuli, vel agrarium. *B. pin.* 328. *R. spn.* 330. *Vaill. par.* 196.

t. 22. f. 3.

Anglis, hop Trefoil.

Habitat in pratis et pascuis glareosis. ☉. VI. VII.

procum- 15. TRIFOLIUM spicis ovalibus imbricatis, vexillis
bens. deflexis

deflexis persistentibus, caulibus procumbentibus.

Sp. pl. 1088. *It. goth.* 257.

Trifolium spicis strepentibus paucifloris, caulibus erectis. *Hall. hist.* 364.

Trifolium lupinum alterum minus. *R. syn.* 330.

t. 14. *f.* 3. *Hall. ofsc.* 193.

Anglis, procumbent Trefoil.

Habitat in pascuis et pratis. ☉. V—VIII.

16. TRIFOLIUM spicis subimbricatis : vexillis deflexis *filiforme*. persistentibus, calycibus pedicellatis, caulibus procumbentibus. *Sp. pl.* 1088.

Trifolium lupulinum minimum. *R. syn.* 331. *t.* 14.

f. 4.

Anglis, small Trefoil.

Habitat in pratis et pascuis arenosis. ☉. VI.

L O T U S. 879.

Legumen cylindricum strictum. *Alæ* sursum longitudinaliter conniventes. *Cal.* tubulosus.

1. LOTUS capitulis depressis, caulibus decumbentibus, *corniculatus*. leguminibus cylindricis patentibus. *Sp. pl.* 1092. *tus.*

Lotus siliquis umbellatis rectissimis, cylindricis. *Hall. nom.* 385. *hist.* 385.

Lotus corniculata glabra minor. *B. hist.* II. 356. *R. syn.* 334.

Lotus s. Melilotus pentaphyllos minor glabra. *B. pin.* 332.

Trifolium filiquosum minus. *Ger. em.* 1190.

β Lotus pentaphyllos minor angustioribus foliis fruticosior. *R. syn.* 334.

Lotus pentaphyllos frutescens tenuissimis glabris foliis. *B. pin.* 332.

γ Lotus pentaphyllos, flore majore luteo splendente. *B. pin.* 332.

δ Lotus corniculata major, minus hirsuta. *R. syn.* 334.

• Lotus corniculata minor, foliis subtus incanis. *R. syn.* 334.

Anglis, common Bird's-foot-trefoil.

Habitat in pratis, pascuis et ericetis. γ. in pratis nemorosis et sepibus humidis. ☉. VI—VIII.

MEDICAGO.

M E D I C A G O. 899.

Ligumen compressum, cochleatum, *Carina* corollæ a vexillo deflectens.

fativa.

*1. MEDICAGO pedunculis racemosis, leguminibus contortis, caule erecto, glabro. *Sp. pl.* 1096.

Medica caule recto, foliis oblongis serratis, racemis erectis, siliquis planis, iterato contortis. *Hall. hist.* 382.

Medica legitima. *Clus. hist. II.* 242.

Medica fativa. *Hist. Ox. II.* 150. *f.* 2. *t.* 16. *f.* 2.

Fœnum burgundiacum. *Lob. ic.* 2. *p.* 38.

Anglis, manured Medick, or Lucern.

Habitat in pratis et pascuis et ad sepes. 4. VI. VII.

falcata.

2. MEDICAGO pedunculis racemosis, leguminibus lunatis, caule prostrato. *Sp. pl.* 1096. *Fl. dan.* 233.

Medica foliis oblongis serratis, siliquis semilunaribus racemosis. *Hall. hist.* 381.

Medica sylvestris. *B. hist. II.* 383. *R. syn.* 333.

Medica frutescens flavo flore. *Clus. hist. II.* 243. *Park.*

1114.

Trifolium luteum siliqua cornuta. *Ger. em.* 1191.

Trifolium luteum sylvestre, siliqua cornuta et Medica frutescens. *B. pin.* 330.

Anglis, yellow Medick.

Habitat in aggeribus agrorum et pascuis arenosis in Norfolkia; circa Norwich copiose D. Role; betwixt Watford and Bushyhill, the footway; Mr. Doody. *R. syn.*

lupulina.

3. MEDICAGO spicis ovalibus, leguminibus reniformibus monospermis, caulibus procumbentibus. *Sp. pl.* 1097.

Medica caule diffuso, capitulis hemisphæricis, siliquis reniformibus. *Hall. hist.* 380.

Medica polycarpus fructu minore compresso scabro, *R. syn.* 333.

Medica cochleata minor polycarpus annua, capsula nigra minus hispida. *Hist. Ox. II.* 154. *f.* 2. *t.* 15. *f.* 14.

Trifolium

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Classis XVIII.

POLYADELPHIA

POLYANDRIA.

HYPERICUM. 902.

Cal. quinquepartitus. *Petala* quinque. *Filamenta* multa in quinque phalanges basi connata. *Caps.* subrotunda.

Androsæmum.

1. HYPERICUM floribus trigynis, fructibus baccatis, caule fruticoso ancipiti. *Sp. pl.* 1102.

Hypericum maximum Androsæmum vulgare dictum.

R. syn. 343.

Androsæmum maximum frutescens. *B. pin.* 280.

Androsæmum. *Dod. pempt.* 78. vulgare. *Park.* 575.

Clymenum italorum. *Ger. em.* 548.

Anglis, Tutstan Saint-John's-wort.

Habitat in sylvis et sepibus humidis. 4. VII.

humifusum.

2. HYPERICUM floribus trigynis axillaribus solitariis, caulibus ancipitibus prostratis filiformibus, foliis glabris. *Sp. pl.* 1105. *Fl. dan.* 141.

Hypericum caule prostrato, foliis ovatis, calycibus serratis punctatis. *Hall. hist.* 1039.

Hypericum supinum glabrum. *Ger. em.* 541. *B. pin.* 279.

Hypericum minus supinum. *Park.* 572. *R. syn.* 343.

Anglis, trailing Saint-John's-wort.

Habitat in pratis pascuis et ericetis. 4. VII.

pulchrum.

3. HYPERICUM floribus trigynis, calycibus ferrato-glandulosis,

glandulosus, caule tereti, foliis perfoliatis cordatis, glabris. *Sp. pl.* 1106. *Fl. dan.* 25.

Hypericum foliis amplexicaulibus, cordatis, calycibus ovatis ferratis glanduligeris. *Hall. hist.* 1041.

Hypericum minus rectum. *B. pin.* 279.

Hypericum pulchrum Tragi. *B. hist.* III. 383. *R. syn.* 342.

Hypericum quintum s. pulchrum Tragi. *Ger. em.* 540.

Anglis, upright Saint-John's-wort.

Habitat in sylvis et ericetis. 4. VII.

4. **HYPERICUM** floribus trigynis, caule ancipiti, fo-
liis obtusis pellucido-punctatis. *Sp. pl.* 1104. *perforatum.*

Hypericum caule tereti alato ramosissimo, foliis ovatis perforatis. *Hall. hist.* 1037.

Hypericum. *Dod. pempt.* 76. *Ger. em.* 540. *R. syn.* 342. vulgare. *B. pin.* 279. *Park.* 572.

Anglis, common Saint-John's-wort.

Habitat in dumetis et ad sepes. 4. VII.

Pharmac. *Hyperici Herba, Flores, Semina.*

5. **HYPERICUM** floribus trigynis, calycibus ferrato-
glandulosus, caule tereti erecto, foliis ovatis subpu-
bescentibus. *pl.* 1105. *hirsutum.*

Hypericum foliis ovatis hirsutis, calycibus lanceolatis ferratis glanduliferis. *Hall. hist.* 1043.

Hypericum Androsæmum dictum. *B. hist.* III. 382. *R. syn.* 343.

Androsæmum hirsutum. *B. pin.* 280.

Androsæmum alterum hirsutum. *Column. ecphr.* I. 74.

Androsæmum ascyron dictum, caule rotundo hirsuto.

Hist. Ox. II. 271. f. 5. t. 6. f. 11.

Anglis, hairy Saint-John's-wort.

Habitat in dumetis et sepibus. VI. VII.

6. **HYPERICUM** floribus trigynis, calycibus ferraturis
glandulosus, caule tereti erecto glabro, foliis ovatis. *num.*
Sp. pl. 1105. *Fl. dan.* 173.

Hypericum foliis ovatis, per oram punctatis, calycibus lanceolatis, ferratis globuligeris. *Hall. hist.* 1042.

Hypericum elegantissimum non ramosum folio lato. *B. hist.* III. 383. *R. syn.* 343.

Androsæmum

Androsæmum campoclarense. *Col. ecphr. I. 73.*

Ascyron. *Fuchs. hist. 74. f. Hypericum bifolium glabrum non perforatum. B. pin. 280.*

Anglis, mountain Saint-John's-wort.

Habitat in sylvis et dumetis montosis passim. In Charlton Wood. Blackf. sp. bot. 4. VII.

quadran- 7. HYPERICUM floribus trigynis, caule quadrato her-
gulum. baceo. *Sp. pl 1104. Fl. dan. 640.*

Hypericum caule quadrangulati, foliis ovatis perforatis, punctatis. *Hall. hist. 1038.*

Hypericum vulgare minus, caule quadrangulo, foliis non perforatis. *B. pin. 279.*

Hypericum Ascyron dictum, caule quadrangulo. *B. hist. III. 382. R. syn. 344.*

Ascyron. *Dod. pempt. 78. Ger. em. 542. vulgare. Park. 575.*

Anglis, quadrangular Saint-John's-wort, or Saint-Peter's-wort.

Habitat in sepibus et nemorosis humidis. 4. VII.

elodes. 8. HYPERICUM floribus trigynis, caule tereti repente
foliisque villosis subrotundis. *Sp. pl. 1106.*

Ascyron supinum villosum palustre. *B. pin. 280. Park. 574. R. syn. 344.*

Ascyron supinum *ελωδες.* *Clus. hist. app. alt. Ger. em. 542.*

Anglis, marsh Saint-John's-wort, or Saint-Peter's-wort.

Habitat in palustribus solo putrido et spongioso, passim. 4. VII.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Long-Reach, below Woolwich, and in some meadows near Edmonton. Blackf. spec. bot. ♂. V.

SONCHUS. 908.

Recept. nudum. *Cal.* imbricatus, ventricosus. *Pappus* plumosus.

oleraceus. 1. SONCHUS pedunculis tomentosus, calycibus glabris. *Sp. pl.* 1116.

lævis. α Sonchus foliis amplexicaulibus dentatis integris aut semipinnatis, calycibus levibus. *Hall. hist.* 21.

Sonchus lævis. *Ger. em.* 292. *R. syn.* 162. vulgaris. *Park.* 805.

Sonchus lævis laciniatus latifolius. *B. pin.* 124.

β Sonchus lævis minor paucioribus laciniis. *B. pin.* 124.

γ Sonchus lævis laceratus. *Pet. H. br. t.* 14. f. 10.

asper. δ Sonchus foliis amplexicaulibus, rigidiusculis, integris aut semipinnatis, dentatis, calycibus levibus. *Hall. hist.* 22.

Sonchus asper laciniatus. *B. pin.* 124. *Park.* 804.

ε Sonchus asper non laciniatus. *B. pin.* 123.

ζ Sonchus subrotundo folio nostras, levissimis spinulis circa foliorum oras exasperatus. *Pluk. alm.* 354. t. 61. f. 5. *R. syn.* 163.

Sonchus ἀφυλλόκαυλος angusto et oblongo folio nostras, per foliorum ambitum creberrimis spinulis asperatus. *Pluk. alm.* 354. t. 62. f. 4.

Sonchus asper dentatus. *Pet. H. br. t.* 14. f. 4.

Anglis, common Sow-thistle α—δ smooth δ, &c. rough Sow-thistle.

Habitat in ruderalis, cultis, arvis. ☉. VI.—VIII.

Pharmac. Sonchi Herba.

alpinus. 2. SONCHUS pedunculis squamosis, floribus racemosis, foliis runcinatis. *Sp. pl.* 1117.

Sonchus foliis amplexicaulibus petiolatis triangularibus, petiolo pinnato, floribus spicatis. *Hall. hist.* 20.

Sonchus lævis laciniatus cæruleus, f. Sonchus alpinus cæruleus. *B. pin.* 124.

Sonchus cæruleus. *Cam. epit.* 284.

Anglis,

Anglis, alpine Sow-thistle.

Habitat in pascuis montosis. On the borders of corn-fields about Willington and Howden-Pans. Wall. Northumb. 186. ☉.

3. SONCHUS pedunculis calycibusque hispidis subum-*arvensis*. bellatis, foliis runcinatis basi cordatis. *Sp. pl.* 1116.

Fl. suec. 687. *dan.* 606.

Sonchus foliis amplexicaulibus semipinnatis ferratis, calycibus hispidis. *Hall. hist.* 23.

Sonchus repens, multis hieracium majus. *B. hist.* II. 170. *R. syn.* 163.

Sonchus arborescens. *Ger. em.* 294.

Hieracium majus folio sonchi. *B. pin.* 126.

Anglis, corn Sow-thistle.

Habitat in arvis et ad sepas. 4. VIII.

4. SONCHUS pedunculis calycibusque hispidis subum-*palustris*. bellatis, foliis runcinatis basi sagittatis. *Sp. pl.* 1116.

Sonchus tricubitalis, folio cuspidato. *Merr. pin. R. syn.* 163.

Sonchus arborescens alter. *Ger. em.* 294.

Sonchus asper arborescens. *B. pin.* 124.

Anglis, marsh Sow-thistle.

Habitat in aquosis et ad ripas fluviorum; ad ripas Thamesis fluvii non procul Greenvico et circa Blackwall. *R. syn.* 4. VIII.

LACTUCA. 909.

Receptaculum nudum. *Cal.* imbricatus, cylindricus, margine membranaceo. *Pappus* simplex, stipitatus.

1. LACTUCA foliis horizontalibus carina aculeatis *virosa*. dentatis. *Sp. pl.* 119. *Gouan. monf.* 408.

Lactuca foliis ovatis patulis, nervo spinoso. *Hall. hist.* 15.

Lactuca sylvestris major odore opii. *Ger. em.* 309. *R. syn.* 161.

Lactuca sylvestris odore viroso. *B. pin.* 123.

Lactuca virosa. *Park.* 813.

Anglis, stinking Lettuce.

Habitat ad sepas et margines agrorum. 8. VII. VIII.

- scariola.* 2. **LACTUCA** foliis verticalibus, carina aculeatis. *Sp. pl.* 1119.
Lactuca foliis semipinnatis nervo spinoso. Hall. hist. 14.
Lactuca sylvestris costa spinosa. B. pin. 123. R. syn. 161.
Lactuca sylvestris foliis dissectis. Ger. em. 309.
Lactuca sylvestris laciniata. Park. 814.
 β *Lactuca sylvestris folio non laciniato. R. syn. 162.*
Anglis, prickly Lettuce.
Habitat in ruderatis et ad margines agrorum et ad sepes.
 ♂. VIII.

- saligna.* 3. **LACTUCA** foliis hastato-linearibus sessilibus carina aculeatis. *Sp. pl.* 1119. *Fac. austr. III. 28. t. 250.*
Lactuca foliis imis semipinnatis, superioribus amplexicaulibus hamatis, nervis spinosis. Hall. hist. 13.
Lactuca sylvestris altera angusto saligno folio, costa albicante. Hall. jen. 207. t. 4.
Lactuca sylvestris minima. R. cat. cant. ; syn. 160.
Chondrilla viscosa humilis. B. pin. 130. p. 68. Ger. em. 287. Park. 783.
Anglis, least Lettuce.
Habitat in aggeribus fossarum et pascuis solo cretaceo passim. ☉. VIII.

PRENANTHES. 911.

Recept. nudum. Cal. calyculatus. Pappus simplex, subsessilis. Flosculi simplici serie.

- muralis.* 1. **PRENANTHES** flosculis quinis, foliis runcinatis. *Sp. pl.* 1121. *Fl. dan. 509.*
Prenanthes foliis serratis pinnatis, pinna suprema triangulari trilobata. Hall. hist. 18.
Lactuca sylvestris murorum flore luteo. B. hist. II. 1004. R. syn. 162.
Sonchus lævis muralis. Ger. em. 293.
Sonchus lævis laciniatus muralis, parvis floribus. B. pin. 124.
Sonchus lævis alter parvis floribus. B. pin. 124.
Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Habitat in pratis et pascuis aridis; circa Cantabrigiam et alibi sed rarius. R. syn. On Banstead Downs, by the side of the road to Darking in Surry; in Northfleet chalk-pits, Kent. Blackf. Spec. bot. prope Greenhithe. δ . VI. VII.

H E D Y P N O I S.

CAL. Communis calyculatus, oblongus. Squamis exterioribus linearibus, subæqualibus, laxis, brevibus.

Squamis interioribus linearibus, æqualibus, laxis, conniventibus.

COR. Composita, imbricata, uniformis, Corollulis hermaphroditis, numerosis, æqualibus.

Propria monopetala, ligulata, linearis, truncata, quinquentata.

STAM. Filamenta quinque, capillaria, brevissima. Anthera cylindræa, tubulosa.

PIST. Germen ovale. Stylus filiformis longitudine staminum. Stigmata duo reflexa.

PER. nullum. Calyx connivens, erectus.

SEM. solitaria, oblonga, reticulato-sulcata. Pappus subplumosus, sessilis.

REC. nudum, punctatum.

hispidum. α . HEDYPNOIS scapo nudo unifloro, foliis lanceolatis dentatis hispidis.

Leontodon calyce toto erecto hispido, foliis dentatis integerrimis hispidis: pilis furcatis. *Sp. pl.* 1124.

Picris caule nudo unifloro, foliis asperis dentatis. *Hall. hist.* 25.

Dens leonis hirsutus λεπτόκαυλος, Hieracium dictus. *R. syn.* 171.

Hieracium dentis leonis folio hirsutum. *Ger. em.* 303.

Hieracium asperum flore magno dentis leonis. *B. pin.* 127.

Hieracium asperum foliis et floribus dentis leonis bulbosi. *Park.* 788.

hirtum. β Leontodon calyce toto erecto, foliis dentatis hirtis: pilis simplicissimis. *Sp. pl.* 1123.

Hieracium pumilum saxatile asperum præmorsa radice.

B. pin. 128. *pr.* 66. *ic.* *R. syn.* 167.

Anglis,

Anglis, rough Hedypnois.

Habitat in pratis et pascuis. 4. V. VI.

2. HEDYPNOIS caule nudo ramoso, foliis lanceolatis *autum-*
dentatis nudis. *nale.*

Leontodon caule ramoso, pedunculis squamosis, foliis
lanceolatis dentatis integerrimis glabris. *Sp. pl.*
1123. *Fl. dan.* 501.

Picris caule nudo brachiato, foliis glabris semipinna-
tis. *Hall. hist.* 28.

Hieracium minus. *Fuchs. hist.* 320. præmorfa radice.
Park. 794. *R. syn.* 164.

Hieracium minus s. leporinum. *Ger. em.* 296.

Hieracium chondrillæ folio glabro, radice succisa, ma-
jus. *B. pin.* 127.

β Hieracium præmorsum laciniatum. *Pet. h. br. t.* 12.
f. 2.

γ Hieracium folio acuto minus. *Pet. h. br. t.* 12. f. 3.

δ Hieracium folio obtuso minus. *Pet. h. br. t.* 12. f. 4.

• Hieracium foliis lanceolatis dentatis glabris, scapo sub- *Taraxa-*
nudo unifloro, calyce hirsuto. *Sp. pl.* 1125. *ci.*

Anglis, autumnal Hedypnois.

Habitat in pratis et pascuis, • in pascuis montosis in
Wallia et Scotia. 4. VII. VIII.

Obs. variat foliis dentatis, sinuato-dentatis et pinnatifidis,
dentibus integerrimis. Calyce glabro et hirsuto et in
hirsutissimo, pilis nigris. Caule simplici et ramoso,
nudo et squamoso. In : caule plerisque unifloro.

3. HEDYPNOIS caule folioso ramoso, foliis runcinatis *teflorum.*
nudis, radicalibus lanceolatis, caulinis sagittatis
acutis sessilibus.

Crepis foliis lanceolato-runcinatis sessilibus lævibus :
inferioribus dentatis. *Sp. pl.* 1135.

Crepis foliis ad terram pinnatis, superne amplexicau-
libus pinnatis hastatis. *Hall. hist.* 31.

Hieracium luteum glabrum s. minus hirsutum. *B. hist.*
211. *R. syn.* 165.

Hieracium aphacoides. *Tabern.* 182. *Ger. em.* 297.

Cichoreum pratense luteum lævius. *B. pin.* 126.
Park. 778.

β Hieracium aphacoides acutum. *Pet. h. br. t.* 12. f. 7.

- γ Hieracium lactucæ folio. *Pet. h. br. t. 12. f. 5.*
Anglis, yellow Hedypnois or Devil's-bit.
 Habitat in pratis, pascuis, muris et tectis. ☉. VI.—IX.
 Obs. Pappus simplex.

- Hieraci-* 4. HEDYPNOIS caule folioso corymboso, foliis sessilibus oblongis integris hispidis.
oides. Picris perianthiis laxis, foliis integris. *Sp. pl. 1115.*
 Picris aspera foliis oblongis dentatis, superioribus amplexicaulibus. *Hall. hist. 24.*
 Hieracium asperum, majore flore, in agrorum limitibus. *B. hist. II. 1029. R. syn. 167.*
 Hieracium asperum. *Ger. em. 298.*
 Cichoreum pratense luteum hirsutie asperum. *B. pin. 126.*
 Cichoreum pratense luteum asperum. *Park. 777.*
Anglis, succory Hedypnois.
 Habitat in agrorum marginibus solo calcareo. ☉. VII.

- biennis.* 5. HEDYPNOIS caule folioso corymboso, foliis runcinatis hispidis basi superne dentatis, calycibus muricatis.
 Crepis foliis runcinato-pinnatifidis scabris basi superne dentatis, calycibus muricatis. *Sp. pl. 1136.*
 Crepis foliis subscabris, semipinnatis, pinna extrema maxima. *Hall. hist. 30.*
 Hieracium maximum Chondrillæ folio, asperum. *B. pin. 127. pr. 64. R. syn. 166.*
 Hieracium, erucæ folio hirsutum. *B. hist. II. 1025.*
 β Hieracium cichorei folio minus. *Pet. h. br. t. 12. f. 9.*
Anglis, tall Hedypnois.
 Habitat in pratis et pascuis solo calcareo; circa Northfleet et alibi in Cantio frequens. ♂. VII. VIII.

P I C R I S. 907.

Recept. nudum. Cai. duplicatus : exterior pentaphyllus maximus ; interior imbricatus. Pappus plumosus stipitatus.

- echioides.* 1. PICRIS perianthiis exterioribus pentaphyllis interiore aristato majoribus. *Sp. pl. 1114.*

Hieracium

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

** Scapo nudo multifloro.

dubium.

3. **HIERACIUM** foliis integris ovato-oblongis, stolonibus repentibus, scapo nudo multifloro. *Sp. pl.* 1125. Hieracium caule nudo paucifloro, stolonibus reptantibus, foliis longe pilosis lanceolatis. *Hall. hist.* 53. Pilosella major. 1. *Tabern.* 196. repens minus hirsuta. *B. pin.* 262.

Anglis, creeping Hawkweed.

Habitat in montosis humidiusculis; in monte Fairfield dicta prope Rydall in comitatu Westmorlandica. 4. VII. VIII.

Obs. habet multa communia cum Pilosella; sed major est, et folia subtus pilosa non tomentosa. Scapus a tribus ad sex flores; semipedalis aut dodrantalis glaber.

auricula.

4. **HIERACIUM** foliis lanceolatis integerrimis, scapo nudo multifloro, stolonibus reptantibus. *Sp. pl.* 1126. Hieracium caule nudo, stolonibus reptantibus, foliis longe pilosis, lanceolatis, floribus umbellatis. *Hall. hist.* 52.

Pilosella major erecta altera. *B. pin.* 262.

Pilosella major umbellifera μακροκαυλος montana et pratensis. *Col. ecphr.* 248. 249.

Anglis, narrow-leaved Hawkweed.

Habitat in montosis; supra Dalehead non longe a Grassmere in Westmorlandia, sed sparsim. 4. VII.

Desc. Radix præmorsa, radiculis numerosis, simplicibus. Stolones repentis foliosæ, Scapus erectus, nudus, subpilosus, pilis raris, vix semipedalis. Folia lanceolata subintegerrima, acuta, nuda. Flores paniculati a tribus ad sex, lutei. Calycis foliola linearia, pilosa.

*** Caule folioso.

murorum.

5. **HIERACIUM** caule ramoso, foliis radicalibus ovatis dentatis, caulino minori. *Sp. pl.* 1128. Hieracium caule subnudo paucifloro, foliis ovato-lanceolatis, circa petiolum dentatis. *Hall. hist.* 46. Hieracium murorum folio pilosissimo. *B. pin.* 129. *R. syn.* 168.

Hieracium murorum Bauhini, quod est Pulmonaria Gallorum Lobelii. *Park.* 801.

Pulmonaria gallica s. aurea latifolia. *Ger. em.* 304.

β Hieracium

- β Hieracium Pulmonaria dictum angustifolium. *R. syn.* 168.
 Pulmonaria gallica f. angustifolia. *Ger. em.* 304.
- γ Hieracium macrocaulon hirsutum folio rotundiore. *R. syn.* 169.
 Hieracium fruticosum folio subrotundo. *B. pin.* 126.
- δ Hieracium λεπτόκαυλον hirsutum folio longiore. *R. syn.* 169.
- Pilosellæ majoris seu Pulmonariæ luteæ species angustifolia. *B. hist.* II. 1034.
Anglis, wall Hawkweed, French or Golden Lungwort.
 Habitat in sylvis, muris et aggeribus agrorum. 4. VII.
6. HIERACIUM caule paniculato, foliis amplexicaulibus dentatis glabris, calycibus hispidis. *Sp. pl. lum.* 1129.
 Hieracium foliis glabris dentatis, imis petiolatis, superioribus amplexicaulibus, calycibus hirsutis. *Hall. hist.* 45.
 Hieracium montanum latifolium glabrum minus. *B. pin.* 129. *Hist. Ox.* III. 69. f. 7. t. 5. f. 47.
 Hieracium montanum cichorei folio. *R. syn.* 165.
Anglis, marsh Hawkweed.
 Habitat in pratis humidis et juxta rivulos Cravoniæ montosos abundat. *R. syn.* 6. VII.
7. HIERACIUM caule erecto multifloro, foliis ovato-sabaulanceolatis dentatis semiamplexicaulibus. *Sp. pl. dum.* 1131. *Hort. ups.* 238.
 Hieracium foliis hirsutis dentatis, inferioribus ellipticis, superioribus ovato-lanceolatis. *Hall. hist.* 35.
 Hieracium fruticosum latifolium hirsutum. *B. pin.* 129. *Park.* 802. *R. syn.* 167.
- β Hieracium fruticosum latifolium glabrum. *Park.* 168.
- γ Hieracium f. Pilosellæ majoris species humilis, foliis longioribus rarius dentatis plurimis simul, flore singulari nostras. *R. syn.* 170. *Pluk. ph. t.* 37. f. 3.
Anglis, shrubby Hawkweed.
 Habitat in sylvis et sepibus, β in sylvis humidis opacis. 4. VII. VIII.

- umbellatum.* 8. **HIERACIUM** foliis linearibus subdentatis sparsis, floribus subumbellatis. *Sp. pl.* 1131. *Hist.* 34.
 Hieracium fruticosum angustifolium majus. *B. pin.* 129. *Park.* 801. *R. syn.* 298.
 Hieracium intibaceum. *Ger. em.* 298.
 β Pulmonaria angustifolia glabra. *Pet. h. br. t.* 13. f. 11.
 γ Pulmonaria graminea. *Pet. h. br. t.* 13. f. 12.
Anglis, umbellated Hawkweed.
Habitat in sepibus, nemorosis siccis. 4. VII. VIII.

H Y O S E R I S. 916.

Recept, nudum. Cal. subæqualis. Pappus pilosus vel obsoletus.

- minima.* 1 **HYOSERIS** caule diviso nudo, pedunculis incrassatis.
Sp. pl. 1138. *Fl. dan.* 201.
 Hyoseris mascula. *Ger. em.* 288.
 Hieracium minimum. *Clus. hist.* 142. Hyoseris Tabernæmontani et Gerardi. *Park.* 791. *R. syn.* 173.
 Hieracium minus folio subrotundo. *B. pin.* 127.
 Lapsana foliis ovatis, dentatis, caulibus nudis sub flore tumentibus. *Hall. hist.* 4.
Anglis, least Hyoseris or Swine's Succory.
Habitat in pratis et arvis solo glareoso. Circa Hampton-court copiose. D. Doody. In the common corn-field, Walthamstow, Essex, Dr. Watson. In Teddington field, near Hampton. ☉. V. VI.

H Y P O C H Æ R I S. 918.

Recept. peleaceum. Cal. subimbricatus. Pappus plumosus.

- maculata.* 1. **HYPOCHÆRIS** caule subnudo : ramo solitario, foliis ovato-oblongis integris dentatis. *Sp. pl.* 1143. *Fl. dan.* 149.
 Hypochæris caule unifloro, foliis dentatis, hirsutis. *Hall. hist.* 2.
 Hieracium 1. latifolium. *Clus. hist.* II. 139. *R. syn.* 167.
 Hieracium alpinum latifolium hirsutie incanum, flore magno. *B. pin.* 128. *Seb. it.* 11.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

SYNGENESIA : POLYGAMIA ÆQUALIS.

Soncho affinis *Lampfana domestica*. *B. pin.* 124.

Anglis, common Nipplewort.

Habitat in sepibus, nemorosis et ruderatis. ☉. VI. VII.

C I C H O R I U M. 921.

Recept. subpaleaceum. *Cal.* calyculatus. *Pappus* sub-
quinqüedentatus, obsolete pilosus.

Intybus. 1. CICHORIUM floribus geminis sessilibus, foliis run-
cinatis. *Sp. pl.* 1142.

Cichorium foliis pinnatis pinnis triangularibus denta-
tis, floribus sessilibus. *Hall. hist.* 1.

Cichorium sylvestre. *Ger. em.* 284. *Park.* 776. *R.*
syn. 172. f. officinarum. *B. pin.* 125.

Anglis, wild Succory.

Habitat ad margines agrorum. ♂. VII. VIII.

Pharmac. Cichorei Radix, Herba.

** Capitati.

A R C T I U M. 923.

Cal. globosus : squamis apice hamatis inflexis.

Lappa. 1. ARCTIUM foliis cordatis inermibus petiolatis. *Sp.*
pl. 1143. *Fl. dan.* 642.

Lappa. *Hall. hist.* 161. major seu Arctium Dioscoridis.
B. pin. 198. *R. syn.* 197.

Bardana major. *Ger. em.* 809.

Bardana vulgaris major. *Park.* 1222.

β *Lappa* major capitulo glabro maximo. *R. syn.* 196.

γ *Lappa* major montana, capitulis tomentosus. *B. pin.*
198. *Mill. ic.* 159.

δ *Lappa* major capitulis parvis glabris. *R. syn.* 197.

ε *Lappa* major montana, capitulis minoribus, rotundio-
ribus et magis tomentosus. *Tourn.* 450.

ζ *Lappa* major ex omni parte minor, capitulis parvis ele-
ganter reticulatis. *Pluk. alm.* 205.

Anglis, common Burdock or Clot-bur.

Habitat in viis, ruderatis, cultis. ♂. VII. VIII.

Pharmac. Bardanae Radix.

SERRATULA. 924.

Cal. subcylindricus, imbricatus, muticus.

1. SERRATULA foliis lyrato-pinnatifidis : pinna terminali maxima, flosculis conformibus. *Sp. pl.* 1144.
Fl. dan. 281.
 Serratula. *B. pin.* 225. *R. syn.* 196. purpurea. *Ger. em.* 713.
 Serratula vulgaris flore purpureo. *Park.* 474.
 Carduus inermis foliis glabris ferratis, imis ovatis, superioribus semipinnatis. *Hall. hist.* 163.
 β Serratula flore candido. *Boerb. ind. alt.* 144.
Anglis, common Saw-wort.
 Habitat in sylvis et pascuis. 4. VII.
2. SERRATULA calycibus subhirsutis ovatis, foliis alpina. indivisis. *Sp. pl.* 1145. *Fl. dan.* 37.
 Cirsium foliis ovato-lanceolatis dentatis subtus tomentosis, calycibus villosis. *Hall. hist.* 179. t. VI.
 Cirsium humile montanum cynoglossi folio polyanthemum. *R. syn.* 193. *Dill. elth.* 82. t. 70. f. 81.
 Carduo-cirsium minus britannicum, floribus plurimis summo caule congestis. *Pluk. ph. t.* 154. f. 3.
 Carduus mollis lapathi folio. *B. pin.* 377.
 β Cirsium montanum polyanthemum salicis folio angustodenticulato. *R. syn.* 193.
Anglis, mountain Saw-wort.
 Habitat in rupibus montosis. On the highest rocks of Snowdon, and Brerecliff, near Burnly, Lancashire.
R. syn. 4. VII.
3. SERRATULA foliis dentatis spinosis. *Sp. pl.* 1149. *arvensis.*
Fl. dan. 644.
 Cirsium caule anguloso ramoso, foliis semipinnatis, pinnis angulosis, spinosis, calycibus longis inermibus. *Hall. hist.* 171.
 Carduus vulgatissimus viarum. *Ger. em.* 1173. *R. syn.* 194.
 Carduus vinearum repens, fonchi folio. *B. pin.* 387.
 Carduus ceanothos s. viarum et vinearum repens.
Park. 959.
Anglis,

Anglis, corn Saw-wort, or Way-thistle.
Habitat in arvis et ad vias. 24. VII.

C A R D U U S. 925.

Cal. ovatus, imbricatus, squamis, spinosis. *Receptaculum pilosum.*

• *Foliis decurrentibus.*

lanceolatus.

1. CARDUUS foliis decurrentibus pinnatifidis hispidis, laciniis divaricatis, calycibus ovatis spinosis villosis, caule piloso. *Sp. pl.* 1149.
 Cirsium foliis semipinnatis, pinnis bilobatis spinosis, capitulis congestis tomentosus conicis. *Hall. hist.* 169. *Nom.* 169.
 Carduus lanceolatus. *Tabern.* 699. *Ger. em.* 1174. *R. syn.* 195. latifolius. *B. pin.* 385. s. majus. *Park.* 982.
 β Carduus lanceatus major. *R. syn.* 195.
 γ Carduus lanceatus flore et capite minoribus. *Merr. pin.* *R. syn.* 195.
Anglis, spear Thistle.
Habitat in ruderatis et marginibus viarum. ♂. VII.

nutans.

2. CARDUUS foliis semidecurrentibus spinosis, floribus cernuis, squamis calycinis superne patentibus. *Sp. pl.* 1150.
 Carduus foliis semipinnatis, pinnis angulosis spinosis, floribus nutantibus, calycinis squamis pungentibus patulis. *Hall. hist.* 167.
 Carduus nutans. *B. hist.* III. 56. *R. syn.* 193.
 Carduus muscatus. *Ger. em.* 1174.
 Carduus spinosissimus latifolius sphærocephalus vulgaris. *B. pin.* 385.
Anglis, musk Thistle. *Pet H. br.* t. 21. f. 1.
Habitat in pascuis solo calcareo. ☉. VII.
Obs. Flores pedunculati; pedunculi tomentosi, nudi.

hispidus.

3. CARDUUS foliis decurrentibus sinuatis: margine spinosis, calycibus globosis pedunculatis solitariis erectis.

Carduus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

erectis, axillaribus. Folia alterna utrinque decurrentia, oblonga, sinuato-dentata, laciniis oppositis dentatis; margine spinosa, subtus tomentosa, supra villosa. Flores terminales, aggregati, sessiles, pallide purpurei. Calyx ovatus; subtomentosus; squamis ovatis acuminatis, erectiusculis, persistentibus.

- palustris.* 5. **CARDUUS** foliis decurrentibus dentatis: margine spinosis, floribus racemosis erectis, pedunculis iner-
mibus. *Sp. pl.* 1151.
Cirsium caule recto subnudo alato, foliis semipinnatis spinosis, calycibus aristatis. *Hall. nom. et. hist.* 170.
Carduus palustris. *B. pin.* 377. *pr.* 156. *Park.* 983.
R. syn. 194.
β Carduus palustris mitior, bardanæ capitulo in summo caule singulari. *Pluk. R. syn.* 194.
Anglis, marsh Thistle.
Habitat in pratis paludosis et nemorosis humidis. 4.
VII.

** *Foliis amplexicaulibus.*

- helenioides.* 6. **CARDUUS** foliis radicalibus lanceolatis, caulinis sessilibus cordatis acutis dentatis spinoso-ciliatis subtus niveis, caule subsimplici.
Carduus foliis amplexicaulibus lanceolatis dentatis, spinulis inæqualibus ciliatis, caule iner-
mi. *Sp. pl.* 1155. *Fl. dan.* 109. *Jac. austr. I. t.* 43.
Cirsium foliis ciliatis subtus tomentosis, radicalibus petiolatis ovato-lanceolatis, caulinis lanceolatis et semipinnatis. *Hall. hist.* 180. t. 7.
Cirsium britannicum clusii repens. *B. hist.* III. 46.
R. syn. 193.
Cirsii anglici alia icon Pennei. *Ger. em.* 1183.
Cirsium aliud anglicum. *Park.* 961.
Cirsium singulari capitulo squamato, vel incanum alterum. *B. pin.* 377. *Mill. ic.* 94.
Heterophyllus. β Carduus foliis amplexicaulibus lanceolatis ciliatis integris laciniatisque, caule subunifloro, calyce iner-
mi. *Sp. pl.* 1154. *Gmel. sib. II.* 51. *Jac. austr. I.* 27. 43.
Cirsium maximum asphodeli radice. *B. pin.* 377.
Anglis, melancholy Thistle.

Habitat

Anglis, melancholy Thistle.

Habitat in pascuis montosis agri Eboracensis, Westmorlandici, Cumberlandici et Cambrobritannici passim. VI. VII.

Obs. Folia in α lanceolata, integra et dentata; in β , folia radicalia dentata et pinnatifida; caulina inferiora pinnatifida, media dentata, suprema, integra, omnia spinoso-ciliata, subtus tomentosa, nivea; Caules subramosi, ab unico ad sex flores; α et β ex eoaem radice observavi.

7. **CARDUUS** foliis sessilibus semiamplexicaulibus lanceolatis subdentatis spinulis inæqualibus ciliatis; caule subunifloro. *Jac. austr. I. t. 42.*

Cirsium pannonicum I. pratense. *Clus. hist. 148.*

Cirsium angustifolium non laciniatum. *B. pin. 377.*

Cirsium montanum anglicum. *Ger. em. 1183.*

Cirsium anglicum. *Lob. ic. 583. Park. 961. radice hellebõri nigri modo fibroso, folio longo. B. hist. III.*

45.

Anglis, meadow Thistle.

Habitat in pratis et pascuis humidis. 4. VI.

Desc. Radix perennis, fibrosa, repens. Caulis solitarius, herbaceus, pedalis et bipedalis, erectus, foliatus, tomentosus, simplex; subinde ramo unico. Folia radicalia pauca, petiolata, lanceolata, subdentata, margine inæqualiter aculeato-ciliata, acuminata, supra hispidiuscula subtus tomentosa; Caulina a tribus ad sex vel septem, alterna, semiamplexicaulia, lanceolata, subintegerrima, margine inæqualiter aculeato ciliata, supra hispidiuscula, subtus tomentosa; non nivea; Flos terminalis purpureus; Calyx ovatus tomentosus, squamis imbricatis, ovatis, acuminatis, purpurascens.

8. **CARDUUS** foliis amplexicaulibus hastato-pinnatifidis spinosis, calycibus aphyllis. spinis canaliculatis duplicato-spinosis. *Sp. pl. 1152.*

Silybum nervis foliorum albis. *Hall. hist 181.*

Carduus mariæ. *Trag. hist. 850. Ger. em. 1150. R. syn. 195. vulgaris. Park. 979.*

Carduus albis maculis notatus vulgaris. *B. pin. 281.*

β *Carduus mariæ hirsutus non maculatus. R. syn. 194.*

Anglis, milk Thistle.

A a

Habitat

Habitat in aggeribus fossarum et ruderatis. ☉. VIII.
Pharmac. Cardui mariae Herba, Semina.

*eriopho-
rus.*

9. **CARDUUS** foliis sessilibus bifariam pinnatifidis: laciniis alternis erectis, calycibus globosis villosis.

Sp. pl. 1153.

Cirsium foliis pinnatis, pinnis bilobatis lanceolatis spinosis, calycibus sphaericis spinosis tomentosus.

Hall. hist. 168.

Carduus eriocephalus. Dod. purg. 445. Ger. em. 1152.

Carduus tomentosus corona fratrum dictus. Lob. ic. 2.

p. 9. Park. 978. R. syn. 195.

Carduus capite tomentolo. B. pin. 382.

Anglis, globe Thistle.

Habitat in pratis et pascuis montosis, solo calcareo. ♂. VII.

acaulis.

10. **CARDUUS** acaulis, calyce glabro. *Sp. pl. 1156.*

Cirsium floribus acaulis. *Hall. hist. 178.*

Carduus acaulis, minore purpureo flore. B. pin. 380.

Ger. em. 1158. R. syn. 195.

Carduus acaulis septentrionalium. Park. 969.

Anglis, dwarf Thistle.

Habitat in pascuis montosis praecipue solo calcareo. In
 Eficeto nigro prope Londinum. ♀. VII.

ONOPORDUM. 927.

Recept. favosum. Col. squamæ mucronatæ.

*Acanthi-
um.*

1. **ONOPORDUM** calycibus squarrosis: squamis patentibus, foliis ovato-oblongis sinuatis. *Sp. pl. 1158.*

Onopordon caule alato, foliis ovatis dentatis, dentibus angulosis aristatis. *Hall. hist. 159.*

Carduus tomentosus, Acanthium dictus, vulgaris. R. syn. 196.

Acanthium. Dod. pem. 221. album. Ger. em. 1149.

Acanthium vulgare. Park. 1149.

Spina alba. Fuchf. hist. 57. tomentosa latifolia sylvestris. B. pin. 382. Loef. pr. 261. t. 82.

Anglis, woolly Onopordum or Cotton-thistle.

Habitat in ruderatis. ♂. VII.

CARLINA.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- cernua.* 2. **BIDENS** foliis lanceolatis amplexicaulibus, floribus cernuis, seminibus erectis. *Sp. pl.* 1165.
 Bidens foliis sessilibus serratis, floribus nutantibus circumvallatis. *Hall. hist.* 120.
 Verbesina pulchriore flore luteo. *B. hist. III.* 1074.
R. syn. 187.
 Cannabina aquatica folio non diviso. *B. pin.* 321.
 Eupatorium cannabinum fœminæ varietas altera. *Ger. em.* 711.
 Eupatorium aquaticum folio integro. *Park.* 596.
- Bidens.* β **Coreopsis** foliis lanceolatis serratis oppositis amplexicaulibus. *Sp. pl.* 1281.
 Chrysanthemum aquaticum bidens, s. Hepatorium aquatile folio non diviso, flore aureis undique petalis radiato. *Triumph.* 65. *Hist. Ox. III.* 17. *f. 6. n. 23. f. 2.*
 Conyza palustris foliis serratis. *Loef. pr.* 54. *t. 11. Anglis,* nodding Bidens.
 Habitat in fossis et palustribus; β circa Tapperley et alibi in comitatu Cestriensi. ☉. VII. VIII.

EUPATORIUM. 935.

Recept. nudum. Pappus plumosus. Cal. imbricatus, oblongus. Stylus semibifidus, longus.

- cannabinum.* 1. **EUPATORIUM** foliis digitatis. *Sp. pl.* 1173.
 Eupatorium foliis ovato-lanceolatis tripteris. *Hall. hist.* 136.
 Eupatorium cannabinum. *B. pin.* 320. *Park.* 595.
R. syn. 179. *mas. Ger. em.* 711.
 β Eupatorium cannabinum folio integro, seu non digitato. *R. syn.*
Anglis, common Eupatorium.
 Habitat ad ripas fluviorum et rivulorum. ☿. VII. VIII.

SANTOLINA. 942.

Recept. paleaceum: Pappus nullus. Cal. imbricatus, hemisphæricus.

- maritima.* 1. **SANTOLINA** pedunculis unifloris subcorymbosis, foliis oblongis tomentosis.

Athanasia

Athanasia pedunculis unifloris subcorymbosis, foliis lanceolatis indivisis crenatis obtusis tomentosis. Sp. pl. 1182.

Filago tomentosa, corymbo subramoso, foliis oblongis obtusis crenatis. Sp. pl. ed. I. 927. Mill. ic. 135.

Gnaphalium maritimum. B. pin. 263. R. syn. 180.

Gnaphalium marinum. Ger. em. 640. f. Cotonaria. Park. 687.

Anglis, sea Santolina.

Habitat in littoribus maritimis. Near Abermeney ferry in Anglesea; and between Pensans and St. Michael's mount in Cornwall. R. syn. in insula Shepey; prope Pool in comitatu Dorsetiensi. 4. VII.

POLYGAMIA SUPERFLUA.

* *Discoidei.*

TANACETUM. 944.

Recept. nudum. Pappus submarginatus. Cal. imbricatus hemisphæricus. Cor. radii obsoletæ, trifidæ.

1. TANACETUM foliis bipinnatis incisissimis ferratis. *Sp. vulgare. pl. 1183.*

Tanacetum foliis pinnatis, semipinnatis acute dentatis. Hall. hist. 132.

Tanacetum. Ger. em. 650. R. syn. 188. vulgare. Park. 81. luteum. B. pin. 132.

β *Tanacetum foliis crispis. B. pin. 132.*

Anglis, common Tansy.

Habitat in pratis et pascuis montosis. 4. VII. VIII.

Pharmac. Tanaceti Herba, Flores, Semina.

ARTEMISIA. 945.

Recept. subvillosum vel nudiusculum. Pappus nullus. Cal. imbricatus, squamis rotundatis, conniventibus. Cor. radii nullæ.

* *Procumbentes ante florescentiam.*

1. ARTEMISIA foliis multifidis linearibus, caulibus *campestris.* procumbentibus virgatis. *Sp. pl. 1185.*

Artemisia caulibus procumbentibus, foliis primis fericeis adultis viridibus pinnatis, pinnis multifidis lanceolatis. *Hall. hist.* 131.

Abrotanum campestre. *Tabern.* 16. *B. pin.* 136.

Ger. em. 1106. *Park.* 94. *R. syn.* 190.

Anglis, field Wormwood or Southernwood.

Habitat in pratis, prope Elden in Suffolk. *R. syn.* 4. VII.

maritima. 2. *ARTEMISIA* foliis multipartitis tomentosis, racemis cernuis, flosculis femineis ternis. *Sp. pl.* 1186.

Absinthium marinum album. *Ger. em.* 1099. *R. syn.* 188.

Absinthium seriphinum belgicum. *B. pin.* 139.

β *Absinthium* maritimum seriphio belgico simile, latiore folio, odoris grati. *Pluk. R. syn.* 188.

γ *Absinthium* maritimum gallicum. *B. pin.* 139.

δ *Absinthium* seriphium tenuifolium marinum narbonense. *B. hist.* III. 177. *R. syn.* 189.

• *Absinthium* maritimum nostras. *R. syn.* 189.

Anglis, sea Wormwood.

Habitat in littoribus maritimis. 4. VIII.

Pharmac. *Absinthii marini Herba.*

** *Erectæ herbaceæ, foliis compositis.*

Absinthium. 3. *ARTEMISIA* foliis compositis multifidis, floribus subglobosis pendulis, receptaculo villosa. *Sp. pl.* 1188.

Absinthium foliis incanis pinnatis, pinnis primis bilobatis, pinnulis incisiss lanceolatis, spicis heteromallis, floribus pendulis. *Hall. hist.* 124.

Absinthium vulgare. *Park.* 98. *R. syn.* 188.

Absinthium latifolium, s. Ponticum. *Ger. em.* 196.

Absinthium ponticum s. romanum officinarum s. dioscoridis. *B. pin.* 138. *Tourn.* 457.

Anglis, common Wormwood.

Habitat in ruderatis et ad vias. 4. VIII.

Pharmac. *Absinthii vulgaris Herba.*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Habitat in muris et aridis; in insula Jersey copiose.
R. syn. ☉. VII. VIII.

** *Herbacea Argyrocoma.*

*margari-
taceum.*

2. GNAPHALIUM herbaceum, foliis lineari-lanceo-
latis acuminatis alternis, caule superne ramoso,
corymbis fastigiatis. *Sp. pl.* 1198.

Filago foliis lanceolatis viridibus subtus tomentosis,
floribus umbellatis. *Hall. hist.* 146.

Gnaphalium americanum. *Clus. hist.* 327. *Ger. em.*
641. *R. syn.* 182. latifolium. *B. pin.* 263.

Argyrocome s. Gnaphalium americanum. *Park.* 690.
Anglis, American Cudweed.

Habitat in pratis et pascuis; in prato prope Bocking,
in vicum agri Essexiensis, D. Dale; on the banks of
Rymny river, for the space of at least twelve miles,
Mr. Lhwyd. *R. syn.* 4. VIII.

dioicum.

3. GNAPHALIUM sarmentis procumbentibus, caule
simplicissimo, corymbo simplici, floribus dioicis.
Sp. pl. 1199. *Syst. nat.* 545.

Filago flagellis reptans sexu distincta, flosculis omni-
bus androgynis. *Hall. hist.* 157.

Gnaphalium montanum album. *Ger. em.* 640. *R.*
syn. 181.

Gnaphalium montanum s. Pes cati. *Park.* 690.

Gnaphalium montanum flore rotundiore. *B. pin.* 263.

Gnaphalium montanum longiore folio et flore. *B. pin.*
263. *R. syn.* 182.

Anglis, mountain Cudweed or Cat's-foot.

Habitat in pratis montosis aridis. On Newmarket heath,
not far from Botolpham beacon; in boreali parte
Angliæ frequens, et in Wallia et Cornubia. 4.
V. VI.

*** *Filaginoidea.*

*silvati-
cum*

4. GNAPHALIUM caule herbaceo simplicissimo erecto,
floribus sparsis. *Syst. nat.* 545. *Sp. pl.* 1200.

Filago foliis linearibus, alis spiciferis. *Hall. hist.* 148.

Gnaphalium anglicum. *Ger. em.* 639. *R. syn.* 180.
vulgare majus. *Park.* 685.

Gnapha-

Gnaphalium majus, angusto oblongo folio, alterum.

B. pin. 263.

Anglis, upright Cudweed.

Habitat in pascuis et sylvis arenosis. In ericeto Hampstediano, et prope Charleton in Cantio. δ . VIII.

5. GNAPHALIUM caule herbaceo simplicissimo pro-supinum. cumbente, floribus sparsis. *Syst. nat.* III. 234.

Gnaphalium fuscum. *Scop. carn.* 1048. t. 57?

Filago caule simplicissimo, spica pauciflora. *Hall. hist.*

149.

Gnaphalium supinum, lavendulæ folio. *Bocc. mus.*

107. t. 85.

Anglis, dwarf Cudweed.

Habitat in pascuis et pratis alpinis, aridis; at the foot of a hill, about five miles from Fort George, on the west side of the road leading to Aberdeen. D. Oakes; et in boreali parte Scotiæ passim. D. Hope. 4. VII. VIII.

6. GNAPHALIUM caule herbaceo ramoso diffuso, floribus confertis terminalibus. *Syst. nat.* 545. *Sp. pl. sum.* 1200.

Filago caule ramosissimo diffuso, umbella sessili terminali. *Hall. hist.* 151.

Gnaphalium longifolium humile ramosum, capitulis nigris. *R. syn.* 181.

Filago minor. *Park.* 686.

Anglis, marsh Cudweed.

Habitat in aquosis præcipue ubi aquæ hyemali tempore stagnant. \odot . VIII.

7. GNAPHALIUM caule herbaceo dichotomo erecto, gallicum. floribus subulatis axillaribus, foliis filiformibus.

Filago caule erecto dichotomo, floribus subulatis axillaribus, foliis filiformibus. *Sp. pl.* 1312.

Filago florum acervulis insidentibus, foliis filiformibus. *Hall. hist.* 156.

Gnaphalium parvum ramosissimum foliis angustissimis polyspermon. *R. syn.* 181.

Gnaphalium minimum alterum nostras, stoechadis citrinæ foliis tenuissimis. *Pluk. ph.* t. 298. f. 2.

Anglis,

Anglis, corn Cudweed.

Habitat in arvis glareosis; among corn in sandy ground about Castle-Heveningham in Essex. D. Dale. R. syn. ☉. VII. VIII.

*monta-
num.*

8. GNAPHALIUM caule herbaceo subdichotomo erecto, floribus conicis axillaribus terminalibusque.

Filago caule erecto diviso, floribus conicis lateralibus terminalibusque. *Sp. pl.* 1311.

Filago caule dichotomo, acervulis spicatis. *Hall. hist.* 155.

Filago caule dichotomo, floribus glomeratis spicatis. *Hall. nom.* 155.

Filago minor. *Ger. em.* 641.

Gnaphalium minimum. *Lob. ic.* 481. *B. hist.* III. 159. *R. syn.* 181.

Gnaphalium minus repens. *B. pin.* 363.

Anglis, least Cudweed.

Habitat in pratis et pascuis arenosis. ☉. VII. VIII.

*germani-
cum.*

9. GNAPHALIUM caule herbaceo erecto ramoso, panicula dichotoma, capitulis rotundatis axillaribus terminalibusque.

Filago panicula dichotoma, floribus rotundatis axillaribus hirsutis, foliis acutis. *Sp. pl.* 1311.

Filago caule dichotomo, corymbo sphaerico, in angulo sessili. *Hall. hist.* 153.

Gnaphalium. *Fuchf. hist.* 222. minus s. Herba impia. *Park.* 685. *R. syn.* 180.

Gnaphalium vulgare majus. *B. pin.* 263.

Filago s. herba impia. *Ger. em.* 642.

Anglis, common Cudweed.

Habitat in pratis et pascuis sterilioribus et ad vias publicas. ☉. VII. VIII.

Obs. variat caule erecto simplicissimo, floribus axillaribus, sessilibus.

C O N Y Z A. 950.

Recept. nudum. *Pappus* simplex. *Cal.* imbricatus subrotundus. *Cor.* radii trifidæ.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

TUSSILAGO. 952.

Recept. nudum. Pappus simplex. Cal. squamæ æquales, discum æquantes, submembranaceæ.

Farfara. 1. TUSSILAGO scapo imbricato unifloro, foliis subcordatis angulatis denticulatis. *Sp. pl.* 1214. *Fl. dan.* 595.

Tussilago. *Cam. ep.* 490. 491. *Ger. em.* 811. *Park.* 1220. *R. syn.* 173. vulgaris. *B. pin.* 197.

Petasites scapo unifloro, flore radiato. *Hall. hist.* 143. *Anglis*, common Coltsfoot.

Habitat in pascuis et locis humidiusculis. 4. III. IV. *Pharmac.* Tussilaginis s. Farfaræ Herba, Flores.

Petasites. 2. TUSSILAGO thyrsu ovato, flosculis omnibus hermaphroditis. *Sp. pl.* 1214.

Petasites floribus dense spicatis, flosculis androgynis. *Hall. hist.* 138.

Petasites. *Dod. pempt.* 597. *Ger. em.* 814. *R. syn.* 179. vulgaris. *Park.* 419.

Petasites major et vulgaris. *B. pin.* 107. *Anglis*, greater Coltsfoot or Butter-bur.

Habitat in pratis et pascuis humidis; et ad ripas fluviorum. 4. III. IV.

Pharmac. Petasitidis Herba.

hybrida. 3. TUSSILAGO thyrsu oblongo, flosculis femineis nudis plurimis. *Sp. pl.* 1214.

Petasites floribus spicatis pendulis, petiolis multifloris, calycibus æqualibus, flosculis paucissimis androgynis. *Hall. hist.* 140.

Petasites major, floribus pediculis longis insidentibus. *R. syn.* 179. *Dilh. elth.* 309. t. 230. f. 297.

Anglis, stalked Coltsfoot or Butter-bur.

Habitat in pratis humidis; prope Loughborough et alibi in comitatu Leicestriensi passim. 4. IV.

SENECIO. 953.

Recept. nudum. Pappus simplex. Cal. cylindricus, calyculatus; squamis apice sphacelatis.

* *Floribus*

* *Floribus flosculosis.*

1. SENECEO corollis nudis, foliis pinnato-sinuatis am-*vulgaris.*
plexicaulibus, floribus sparsis. *Sp. pl.* 1217. *Fl.*
dan. 513. *Hall. hist.* 58.

Senecio vulgaris. *Park.* 671. *R. syn.* 178.

Senecio minor vulgaris. *B. pin.* 131.

Erigeron. *Ger. em.* 278.

Anglis, common Groundsel or Simpson.

Habitat in cultis et ruderatis. ☉. IV—X.

Pharmac. Senecionis Herba.

** *Floribus radiatis : radio revolutis.*

2. SENECEO corollis revolutis, foliis pinnatifidis denti-*viscosus.*
culatis subviscidis, caule erecto corymboso.

Senecio corollis revolutis, foliis pinnatifidis viscidis,
squamis calycinis laxis longitudine perianthii. *Sp.*
pl. 1217.

Senecio foliis viscidis pinnatis, pinnis semipinnatis an-
gulosis, calycibus hirsutis. *Hall. hist.* 60.

Senecio incanus pinguis. *B. pin.* 131.

Senecio hirsutus viscidus major odoratus. *B. hist.* III.
190. *R. syn.* 178.

Senecio foetidus. *Park.* 671.

Senecio hirsutus viscidus graveolens. *Dil. elth.* 347.
t. 258. f. 336.

Erigerum tomentosum. *Ger. em.* 278.

- β Senecio corollis revolutis, foliis pinnatifidis denticula-*sylvati-*
tis, caule corymboso erecto. *Sp. pl.* 1217. *cus.*

Senecio foliis pinnatis, pinnis alternis majoribus, se-
mipinnatis angulosis, calycibus levibus, radiis bre-
vissimis revolutis. — *Hall. hist.* 59.

Senecio minor, latiore folio five montanus. *B. pin.*
131. *R. syn.* 178. *Dill. elth.* 258. t. 258. f. 337.

Jacobæa, senecionis folio incano, perennis. *R. hist.*
Rupp. jen. 177. t. 4.

Anglis, stinking β mountain Groundsel.

Habitat in ericetis, nemorosis et sepibus. ☉. VII.

*** *Floribus radiatis : radio patente, foliis pinnatifidis.*

3. SENECEO corollis radiantibus, foliis pinnato-lyratis *Jacobæa.*
laciniis lacinulatis, caule erecto. *Sp. pl.* 1219.

Jacobæa

SYNGENESIA: POLYGAMIA SUPERFLUA.

Jacobæa. *Ger. em.* 280. vulgaris. *B. hist. II.* 1057.
R. syn. 177. major. *Park.* 668.

Jacobæa vulgaris laciniata. *B. pin.* 131.

Anglis, Ragwort Groundsel.

Habitat in pratis et pascuis et ad vias. 4. VII.

aquaticus. 4. SENECEO corollis radiantibus, foliis dentatis, radicalibus ovatis, caulinis pinnatifidis: lacinia extrema majora. *Fl. angl.* 317.

Senecio foliis lineatis pinnatis, pinnis dentatis, extrema maxima. *Hall. hist.* 62.

Jacobæa latifolia. *Ger. em.* 280. *B. hist. II.* 1056.
palustris. *R. syn.* 178.

Jacobæa barbæræ instar laciniata. *Fl. prus.* 129. t. 35.

Jacobæa latifolia pannonica prima. *Park.* 668.

Jacobæa alpina laciniata, flore buphthalmi, *B. pin.*
131.

Anglis, water Groundsel or Ragwort.

Habitat in pratis et pascuis humidis. 4. VII. VIII.

erucifolius. 5. SENECEO corollis radiantibus, foliis pinnatifidis dentatis subhirtis, caule erecto. *Sp. pl.* 1218.

Jacobæa senecionis folio incano perennis. *R. syn.* 177.

Jacobæa altissima, foliis erucæ artemisiæve similibus et æmulis. *Rupp. jen.* 164.

Jacobæa incana altera. *B. pin.* 131.

Anglis, hoary Groundsel.

Habitat in sylvis et ad sepes. 4. VII. VIII.

*** Floribus radiatis: radio patente, foliis indivisis.

paludosus. 6. SENECEO corollis radiantibus, foliis ensiformibus acute serratis, subtus subvillosis, caule stricto. *Sp. pl.* 1220. *Fl. dan.* 385.

Senecio foliis linearibus serratis, floribus umbellatis.
Hall. hist. 66.

Jacobæa foliis longis integris et mucronatis. *Hist. Ox. III.* 110. f. 7. t. 7. f. 22.

Virgæ aureæ s. Solidagini angustifoliæ affinis, Lingua avis Dalechampii. *B. hist. II.* 1063. *R. syn.* 176.

Conyza palustris. *Park.* 1231. serratifolia. *B. pin.*
266.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Anglis, Welch Golden-rod. Pet. b. br t. 16. f. 11.
Habitat in montibus circa Llanberris copiose. In montibus in comitatibus Eboracensi et Westmorlandico.
 4. VIII.

Obs. variat pedunculo unifloro in summis montibus; an Solidago minuta Linnæi.

A S T E R. 954.

Recept. nudum. Pappus simplex. Cor. radii plures decem. Cal. imbricati, squamæ inferiores patulæ.

Tripolium.

1. ASTER foliis lanceolatis integerrimis carnosis glabris, ramis inæquatis, floribus corymbosis. *Sp. pl. 1226. Fl. dan. 615.*

Aster maritimus cæruleus Tripolium dictus. R. syn. 175.

Tripolium vulgare majus et minus. Ger. em. 413.

Tripolium majus cæruleum et minus. B. pin. 267.

Tripolium majus s. vulgare et minus. Park. 673.

Anglis, sea Starwort.

Habitat in littoribus maritimis, solo lutofo. 4. VIII.

I N U L A. 956.

Recept. nudum. Pappus simplex. Antheræ basi in setas duas desinentes.

Helenium.

1. INULA foliis amplexicaulibus ovatis rugosis, subtus tomentosis, calycum squamis ovatis. *Sp. pl. 1236.*

Aster foliis ovato-lanceolatis serratis subtus tomentosis, calycinis ovato-lanceolatis maximis. Hall. hist. 72.

Helenium. Cam. ep. 35. Ger. em. 793. R. syn. 176.

s. Enula campana. Park. 674. B. hist. III. 108.

Anglis, common Elecampane.

Habitat in pratis et pascuis humidis; in Essexia frequens; circa St. Ives in Cornubia et Bugden in comitatu Huntingtoniensi; inter Denbigh et St. Asaph. 4. VII. VIII.

Pharmac. Enulæ campanæ Radix.

dysenterica.

2. INULA foliis amplexicaulibus cordato-oblongis subtomentosis, caule villoso paniculato, squamis calycinis setaceis. *Sp. pl. 1237. Fl. dan. 410.*

Aster

Aster foliis amplexicaulibus undulatis subtus tomentosis. *Hall. hist.* 79.

Conyza media. *Ger. em.* 482. *R. syn.* 174. asteris flore luteo, vel tertia *Dioscoridis.* *B. pin.* 265.

Herba Dysenterica. *Hoff. altd.*

Anglis, middle Elecampane or Fleabane.

Habitat in pratis et pascuis humidis. 4. VIII.

3. *INULA* foliis amplexicaulibus oblongis undulatis, caule erecto, floribus globosis radio brevissimo. *Pulicaria.*

Inula foliis amplexicaulibus undulatis, caule prostrato, floribus subglobosis radio brevissimo. *Sp. pl.* 1238. *Fl. dan.* 613.

Aster foliis amplexicaulibus undulatis hirsutis, radiis brevissimis. *Hall. hist.* 80.

Conyza minor. *Matth. p. II.* 223. *R. syn.* 174.

Conyza minima. *Dod. pempt.* 52. *Ger. em.* 482.

Conyza minor, flore globoso. *B. pin.* 266.

Anglis, lesser Elecampane or Fleabane.

Habitat ad vias et in locis ubi per hyemem aquæ stagnant. . VIII. IX.

4. *INULA* foliis linearibus carnosis tricuspидatis. *Sp. crithmoides.*

Aster maritimus flavus, *Crithmum chrysanthemum,* dictus. *R. syn.* 174.

Crithmum chrysanthemum. *Ger. em.* 533. *Park,* 1287.

Crithmum maritimum, flore asteris attici. *B. pin.* 288.

Anglis, golden Elecampane or Sampire.

Habitat in palustribus maritimis, solo lutofo. 4. VIII.

C I N E R A R I A. 957.

Recept. nudum. *Pappus simplex.* *Cal. simplex,* polyphyllus, æqualis.

1. *CINERARIA* floribus corymbosis, foliis lato-lanceo-palustris latis dentato-sinuatis, caule villoso. *Sp. pl.* 1243. *Fl. dan.* 573.

Othonna foliis lanceolatis dentato-sinuatis. *Sp. pl. ed. 1.* 924.

• *Conyza foliis laciniatis.* *Ger. em.* 483. *R. syn.* 174.

Conyza aquatica laciniata. *B. pin.* 266.

Conyza helenitis foliis laciniatis. *Park.* 126.

Anglis, marsh Cineraria or Fleabane.

Habitat in palustribus in comitatu Lincolnensi copiose.

In the fen-ditches about Marsh and Chatterers in the isle of Ely; at Caister near Yarmouth; in the ditches about Pillin-moss, Lancashire; at Aberavon in Merionethshire in Wales. *R. syn.* 4. VI. VII.

alpina.

2. **CINERARIA** umbella involuocrata, pedunculo communi nudiusculo, foliis oblongis villosis. *Sp. pl.* 1243. *Jac. austr. II.* 48. t. 170. et 189.

*integri-
folia.*

Othonna foliis lanceolatis subdenticulatis villosis, floribus umbellatis. *Sp. pl. ed. I.* 925.

Senecio foliis imis petiolatis cordatis obiter dentatis, caulibus ligulatis sessilibus. *Hall. hist.* 68.

Jacobæa montana lanuginosa angustifolia non laciniata. *B. pin.* 131. *Hist. Ox. III.* 111. t. 7. f. 12.

Jacobæa pannonica folio non laciniata. *R. syn.* 178.

Jacobæa angustifolia. *Ger. em.* 280. *pannonica non laciniata.* *Park.* 668.

Anglis, mountain Cineraria or Ragwort.

Habitat in pratis montosis. On Gogmagog hills, Newmarket heath, &c. R. syn. prope Basingstoke et Andover. 4. VI.

B E L L I S. 962.

Recept. nudum, conicum. *Pappus* nullus. *Cal.* hemisphæricus: squamis æqualibus. *Sem.* obovata.

perennis. 1. **BELLIS** scapo nudo. *Sp. pl.* 1248. *Fl. dan.* 503. *Hall. hist.* 93.

Bellis sylvestris minor. *B. pin.* 262. *R. syn.* 184.

Bellis minor sylvestris. *Ger. em.* 635, simplex. *Park.* 530.

Anglis, common Daisie.

Habitat in pratis et pascuis. 4. III—IX.

Pharmac. *Bellidis minoris Herba, Flores, Radix.*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Anglis, common Feverfew.

Habitat in ruderatis et ad sepas. ♂. f. 4. VI. VII.

Pharmac. Matricariæ Herba.

*Chamo-
milla.*

2. MATRICARIA receptaculis conicis, seminibus nudis, squamis calycinis margine æqualibus, caule erecto.

Matricaria receptaculis conicis, radiis patentibus, seminibus nudis, squamis calycinis margine æqualibus. Sp. pl. 1256.

Matricaria foliis capillaribus duplicato-pinnatis, pinnis lanceolatis bifidis trifidisque. Hall. nom. et hist. 101.

Chamæmelum. Ger. em. 754. R. syn. 184. vulgare. Dod. pempt. 257. Park. 85. f. Leucanthemum Dioscoridis. B. pin. 135.

*suaveo-
lens.*

- β *Matricaria receptaculis conicis, radiis deflexis, seminibus nudis, squamis calycinis margine æqualibus. Sp. pl. 1256.*

Matricaria leucanthemos annua suaveolens, chamæmeli folio, ovariis albicantibus. Vaill. aët. paris. 1720. 556.

Chamæmelum tenuè incisum, foliis floris deorsum protensis : umbone extuberante suavius olens. Cup. cath. suppl. 17.

Anglis, corn Feverfew.

Habitat in arvis et ad vias. ☉. V. VI.

Pharmac. Chamæmeli vulgaris Herba, Flores.

inodora.

3. MATRICARIA receptaculis hemisphæricis, seminibus coronato-marginatis, squamis calycinis margine obsolete, caule diffuso.

Chrysanthemum foliis pinnatis multifidis, caule ramoso diffuso. Sp. pl. 1253.

Matricaria inodora. Fl. suec. 765.

Chamæmelum inodorum annum humilius, foliis obscure virentibus. Hist. Ox. III. 36. R. syn. 186.

Chamæmelum inodorum f. Cotula non foetida. B. hist. III. 120.

- β *Chamæmelum majus, folio tenuissimo, caule rubente. Magn. hort. monsp. 53. R. syn. 18.*

Matri-

- γ *Matricaria* receptaculis hemisphæricis, foliis bipinna-maritima-
tis subcarnosis, supra convexis, subtus carinatis.

Sp. pl. 1256.

Chamæmelum maritimum perenne humilius, foliis
brevibus crassis obscure virentibus. *R. syn.* 186. t. 7.

f. 1.

Anglis, corn Feverfew.

Habitat in arvis et ad vias. ☉. ♂. VII. VIII.

ANTHEMIS. 970.

Recept. peleaceum. *Pappus* nullus. *Cal.* hemisphæ-
ricus, subæqualis. *Flosculi* radii plures quam quin-
que.

* *Radio* discolore f. albo.

1. ANTHEMIS foliis pinnato-compositis linearibus *nobilis*.
acutis subvillosis. *Sp. pl.* 1260. seminibus nudis.

Chamæmelum foliis subhirsutis, nervo duro, pinnis
pinnatis, pinnulis lanceolatis incis. *Hall. hist.* 102.

Chamæmelum nobile seu *Leucanthemum* odoratius.
B. pin. 135.

Chamæmelum odoratissimum repens flore simplici. *B.*
hist. III. 118. *R. syn.* 185.

Chamæmelum romanum. *Ger. em.* 755.

Anglis, common Camomile.

Habitat in pratis et pascuis apricis. ♀. VIII. IX.

Pharmac. Chamæmeli Flores.

2. ANTHEMIS receptaculis conicis, paleis setaceis, *arvensis*.
seminibus coronato-marginatis. *Sp. pl.* 1261. fo-
liis villosiusculis.

Chamæmelum inodorum. *B. pin.* 135. *R. syn.* 184.

Bupthalam album. *Pet. h. br. t.* 19. f. 8.

Cotula. *Tab. ic.* 21.

Anglis, corn Camomile.

Habitat in arvis et agris restibilibus. ♂. VI. VII.

3. ANTHEMIS receptaculis conicis, paleis setaceis, se- *Cotula*.
minibus nudis. *Sp. pl.* 1261. foliis nudis.

Chamæmelum foliis glabris, duplicato-pinnatis, nervo
foliaceo,

foliaceo, pinnulis lanceolatis, seminibus exasperatis.

Hall. hist. 104.

Chamæmelum foetidum. *B. pin.* 135. *R. syn.* 185.

Cotula foetida. *Ger. em.* 756. *B. hist.* III. 120.

Anglis, stinking Camomile or May-weed.

Habitat in arvis et ad vias. ☉. VI. VII.

maritima. 4. **ANTHEMIS** foliis pinnatis denticulatis carnosis nudis punctatis, caule prostrato, calycibus subtomentosis. *Sp. pl.* 1259.

Anthemis maritima perennis, foliis crassis et punctatis. *Mich. gen.* 33.

Chamæmelum maritimum. *B. hist.* III. 122. latifolium ramosissimum, flore albo. *Pluk. alm.* 97. *R. syn.* 186.

Matricaria maritima. *B. pin.* 134.

Anglis, sea Camomile.

Habitat in pratis maritimis. 4. VII. VIII.

** *Radio concolore f. luteo.*

tinctoria. 5. **ANTHEMIS** foliis bipinnatis ferratis subtus tomentosis, caule corymboso. *Sp. pl.* 1263.

Chamæmelum foliis pinnatis, hirsutis, pinnulis simplicibus, lanceolatis. *Hall. hist.* 105.

Bupthalmum tanacetii minoris folio. *B. pin.* 134.

Bupthalmum vulgare. *Ger. em.* 747. *R. syn.* 182.

Bupthalmum matthioli f. vulgare millefolii foliis. *Park.* 1370.

Anglis, yellow Camomile or Ox-eye.

Habitat in pratis apricis, sed rarius. I found this on a bank near the river Tees, not far from Sogburn in the bishoprick of Durham. *R. syn.* 4. VII. VIII.

A C H I L L E I A. - 971.

Recept. paleaceum. *Pappus* nullus. *Cal.* ovatus, imbricatus. *Flosculi* radii circiter quinque.

Millefolium. 1. **ACHILLEA** foliis bipinnatis nudis: laciniis linearibus dentatis, caulibus sulcatis. *Sp. pl.* 1267.

Achillea foliis pinnatis, pinnis longe æqualibus pinnatis, pinnulis trifidis et quinquefidis. *Hall. hist.* 107.

Millefo-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Jacea foliis radicalibus semipinnatis, caulinis ovato-lanceolatis. *Hall. hist.* 184.

Jacea nigra. *Ger. em.* 721. *R. syn.* 198. vulgaris. *Park.* 468.

Jacea nigra pratensis latifolia. *B. pin.* 272.

β Jacea nigra minor, tomentosa laciniata. *R. syn.* 199.

Jacea nigra laciniata. *B. pin.* 271.

Rhaponticum foliis tomentosis radicalibus semipinnatis, caulinis lanceolatis. *Hall. hist.* 195.

γ Centaurea calycibus ciliatis, squamula ovata, ciliis capillaribus erectis, foliis lanceolatis dentatis, floribus radiatis.

Cyanus foliis radicalibus semipinnatis, caulinis lanceolatis, squamis calycinis oblongis obtusis. *Hall. hist.* 185.

Anglis, black Centaury or Knapweed.

Habitat in pratis et pascuis. 4. VII. VIII.

Scabiosa. 3. CENTAUREA calycibus ciliatis: foliis pinnatifidis, pinnis lanceolatis. *Sp. pl.* 1291.

Cyanus foliis pinnatis, pinnis dentatis obtusis, squamis calycinis triangulis. *Hall. hist.* 186.

Jacea major. *Ger. em.* 727. *R. syn.* 198.

Jacea nigra vulgaris laciniata. *Park.* 470.

Scabiosa major squamatis capitulis. *B. pin.* 269.

Anglis, greater Centaury or Knapweed.

Habitat in arvis et ad margines agrorum. 4. VII.

** *Calcitrapæ: calycinis spinis compositis.*

Calcitrapa. 4. CENTAUREA calycibus subduplicato-spinosis sessilibus, foliis pinnatifidis linearibus dentatis, caule piloso. *Sp. pl.* 1297.

Calcitrapa foliis pinnatis, pinnis dentatis, floribus in alis sessilibus. *Hall. hist.* 194.

Carduus stellatus. *Dod. purg.* 448. *Ger. em.* 1166.

R. syn. 196. s. calcitrapa vulgaris. *Park.* 989.

Carduus stellatus foliis papaveris erratici. *B. pin.* 387.

Anglis, star Centaury or Thistle.

Habitat in pratis sterilioribus et ad vias. ○. VIII.

Pharmac. Calcitrapæ Herba, Semina.

5. CENTAU.

5. **CENTAUREA** calycibus duplicato-spinosis solita-solstitialis. riis foliis rameis decurrentibus inermibus lanceolatis, radicalibus lyrato-pinnatifidis. *Sp. pl.* 1297.
Calcitrapa caule alato, foliis tomentosis, radicalibus semipinnatis caulinis lanceolatis dentatis, *Hall. hist.* 193.
Carduus stellatus luteus, foliis cyani. *B. pin.* 387. *R. syn.* 196.
Carduus solstitialis. *Ger. em.* 1166. *Park.* 989.
Anglis, St. Barnaby's Centaury or Thistle.
Habitat in arvis et ad sepes. *In the hedges not far from Cirencester in Gloucestershire.* *D. Robarb.* *R. syn.* *Juxta Norfleet in Cantio.* ☉. VII. VIII.

M O N O G A M I A,

J A S I O N E.

Cal. communis decemphyllus. *Cor.* pentapetala, regularis. *Capsula* infera, bilocularis.

1. **JASIONE.** *Sp. pl.* 1316. *Fac. enum.* 289. *Fl. dan. montana.* 319.
Rapunculus foliis linearibus subsperis, spica planiuscula petalis liberis. *Hall. hist.* 678.
Rapunculus scabiosæ capitulo cæruleo. *B. pin.* 92. *Park.* 646. *R. syn.* 723.
Scabiosa. *D. u. pem.* 122. minima hirsuta. *Ger. em.* 723.
Anglis, mountain Jasionne or Sheep's Scabious.
Habitat in pratis et ericetis. ☉. VI. VII.

L O B E L I A. 1006.

Cal. quinquefidus. *Cor.* monopetala, irregularis. *Caps.* infera bi-f. trilocularis.

1. **LOBELIA** foliis linearibus bilocularibus integerrimis, *Dort-* caule subnudo. *Sp. pl.* 1318. *Fl. dan.* 39. *manna.*
Dortmanna palustris, floribus sparsis pendulis. *Rudb. æt. ups. an.* 1720. p. 97. t. 2.

Gladiolus

Gladiolus lacustris Dortmanni. *Clus. cur. post.* 40.
R. syn. 287.

Gladiolus lacustris Clusii s. Leucojum palustre, flore subcæruleo. *Park.* 1250.

Leucojum palustre, flore subcæruleo. *B. pin.* 41.
Anglis, water Lobelia or Gladiole.

Habitat in lacubus. In omnibus Cambriæ montibus.
Dr. Richardson. *R. syn. in lacubus Westmorlandiæ,*
Cumberlandiæ et Scotiæ. 4. VII. VIII.

urens.

2. LOBELIA caule erectiusculo, foliis inferioribus subrotundis crenatis, superioribus lanceolatis serratis, floribus racemosis. *Sp. pl.* 1321.

Rapunculus galeatus blesensis s. soloniensis, flore violaceo minimo. *Hist. Ox.* II. 407. t. 5. f. 56.

Rapuntium urens soloniense. *Hort. R. bl.* 300. *Bacc. sic.* 20. t. II. *R. hist.* 746. *Monn. obs.* 131.

Draba flore cæruleo galeato. *B. pin.* 110. *pr.* 53.
Anglis, stinging Lobelia.

Habitat in pratis montosis. *Supra Shute common inter Axminster et Honiton.* D. Newberry. ☉. VII. VIII.

V I O L A. 1007.

Cal. pentaphyllus. *Cor.* pentapetala irregularis, postice cornuta. *Caps.* supera, trivalvis, unilocularis.

* *Acaules.*

odorata.

1. VIOLA acaulis, foliis cordatis, stolonibus reptantibus. *Sp. pl.* 1324. *Fl. dan.* 309.

Viola acaulis stolonifera, foliis cordatis. *Hall. hist.* 558.

Viola martia purpurea. *B. hist.* II. 542. *R. syn.* 364. flore simplici odoro. *B. pin.* 199.

Viola nigra s. purpurea. *Dod. pempt.* 156. *Ger. em.* 850.

Viola simplex martia. *Park.* 755.

B Viola martia alba. *B. pin.* 199.

Anglis, sweet Violet.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

*** *Stipulis pinnatifidis. Stigmate urceolato.*

tricolor.

5. VIOLA caule triquetro diffuso, foliis oblongis dentatis, stipulis dentato-pinnatifidis, lacinia suprema oblonga subpetiolata.

Viola caule triquetro diffuso, foliis oblongis incis, stipulis pinnatifidis. *Sp. pl.* 1326.

Viola caule anguloso diffuso, foliis ovatis dentatis, flore calyce paulo majore. *Hall. hist.* 569.

Viola bicolor arvensis. *B. pin.* 200. *R. syn.* 366.

Viola tricolor sylvestris. *Ger. em.* 853? *Park.* 755.

- β Viola caule anguloso diffuso, foliis ovatis dentatis, flore calyce duplo longiore. *Hall. hist.* 568.

Viola tricolor. *Ger. em.* 854. *R. syn.* 365. major et vulgaris. *Park.* 756.

- Viola tricolor hortensis repens. *B. pin.* 199.

Jacea tricolor s. Irinitatis flos. *B. hist.* III. 546.

Anglis, pansy Violet Heart's-ease or three Faces under a hood.

Habitat in arvis. ☉. V.—IX.

grandiflora.

6. VIOLA caule triquetro simplici, foliis subcordatis dentatis, stipulis pinnatifidis.

Viola caule triquetro simplici, foliis oblongiusculis, stipulis pinnatifidis. *Syst. nat.* 585. *Mant. I.* 120.

Viola montana lutea grandiflora. *B. pin.* 200. nostras. *R. syn.* 356.

Viola lutea. *Ger. em.* 850. desc. non icon.

Anglis, yellow Violet or Pansies.

Habitat in montosis pascuis. In septentrionalibus et Wallicis, solo putrido et palustri multoties observavimus.

4. V.—IX.

I M P A T I E N S. 1008.

Cal. diphyllus. *Cor.* pentapetala, irregularis, nectario cucullato. *Caps.* supera quinquevalvis.

noli me tangere.

1. IMPATIENS pedunculis multifloris solitariis, foliis ovatis, geniculis caulinis tumentibus. *Sp. pl.* 1329. *Fl. dan.* 582.

Impatiens

Impatiens caule geniculato, pedunculis ramosis, foliis petiolatis ovatis crenatis. Hall. hist. 557.

Perficaria filiquosa. Ger. em. 446.

Balsamine lutea s. noli me tangere. B. pin. 306. R. syn. 316.

Balsamita altera, Noli me tangere. Col. ecphr. I. 149. t. 50.

Mercurialis sylvestris, Noli me tangere dicta sive Perficaria filiquosa. Park. 296.

Anglis, yellow Balsam, Quick in the Hand or Touch-me-not.

Habitat in umbrosis humidis et ad rivulos; on the banks of Winandermere, near Ambleside. R. syn. prope Rydall-hall; et alibi in Westmorlandia. ☉. VII. VIII.

Classis XX.

GYNANDRIA

DIANDRIA.

ORCHIS. 1009.

Nectarium corniforme pone florem.* *Bulbis indivisis.**bifolia.*

1. ORCHIS bulbis indivisis, nectarii labio lanceolato integerrimo : cornu longissimo, petalis patentibus.

Sp. pl. 1331. *Fl. dan.* 235.*Orchis* radicibus oblongis, labello lineari. *Hall. hist.* 1285. t. 35. *Orch.* 82.*Orchis* alba bifolia minor, calcari oblongo. *B. pin.* 83. *R. syn.* 380. *Vaill. par.* 151. t. 30. f. 7.*Orchis* psychodes. *Ger. em.* 211.*Orchis* sphegodes f. *Testiculus vulpinus* 1. *Park.* 1351.

- β *Orchis* hermaphroditica bifolia. *B. hist.* II. 772.

Anglis, butterfly *Orchis*.*Habitat in pratis, pascuis et dumetis.* 4. V. VI.*pyrami-
dalis.*

2. ORCHIS bulbis indivisis, nectarii labio bicorni trifido æquali integerrimo, cornu longo, petalis sublanceolatis. *Syst. nat.* 589. *Sp. pl.* 1332. *Jac. austr.* III. 37. t. 266.

Orchis radicibus subrotundis, spica densa, calcare longissimo, labello bicorni æqualiter tripartito. *Hall. hist.* 1286. t. 35. *Orch.* 83.*Orchis* purpurea spica congesta pyramidali. *R. syn.* 377. t. 18. *Seg. ver.* t. 15. f. XI.*Cynosorchis*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Anglis, female Orchis.

Habitat in pratis et pascuis subhumidis. 4. V. VI.

ustulata. 6. ORCHIS bulbis indivisis, nectarii labio quadrifido punctis scabro : cornu obtuso, petalis distinctis.

Sp. pl. 1333. Fl. dan. 103.

Orchis radicibus subrotundis, labello quadrifido, calcare brevissimo. *Hall. hist. 1273. t. 28. Orch. 77.*

Orchis pannonica IV. *Clus. hist. 268. R. syn. 377.*

Cynorchis militaris pratensis humilior. *B. pin. 81.*

Cynorchis militaris pannonica. *Park. 1345.*

Cynorchis minor pannonica. *Ger. em. 207.*

Anglis, dwarf Orchis.

Habitat in pratis et pascuis siccioribus. 4. V. VI.

militaris. 7. ORCHIS bulbis indivisis, nectarii labio quinquefido punctis scabro ; cornu obtuso, petalis confluentibus.

Sp. pl. 1333.

Orchis bulbis indivisis, nectarii labio quinquefido lineari punctis scabro : petalis confluentibus. *Fl. angl. 335.*

Orchis radicibus subrotundis, spica conica, labello quadrifido, brachiolis et pectore perangustis. *Hall. hist. 1277. t. 28. Orch. 79.*

Orchis galea et alis fere cinereis. *B. pin. II. 755. R. syn. 378.*

Cynorchis latifolia hiante cuculo major. *B. pin. 80.*

Cynorchis latifolia minor. *Park. 1344.*

Cynorchis major altera. *Ger. em. 205.*

purpurea. β Orchis bulbis indivisis, nectarii labio barbato-quinquelobo lateralibus obtusis crenulatis, medio brevissimo acuto, petalis confluentibus. *Fl. angl. 334.*

Orchis radicibus subrotundis, spica longa, labello quadrifido, brachiolis angustis crúsculis latis ferratis. *Hall. hist. 1276. t. 31. Orch. 78.*

Orchis magna latis foliis, galea fusca vel nigricante: *B. hist. II. 759. R. syn. 378. t. 19. f. 2.*

Anglis, man Orchis β purple man Orchis.

Habitat in pratis et pascuis solo cretaceo. On Cawsham hills by the Thames side near Reading, Berkshire.

Near

*Near the old chalk-pit by the paper-mill at Harefield.
Blackf. β in sylvis et dumetis prope Northfleet in Can-
tio. 4. V.*

**** Bulbis palmatis.**

8. ORCHIS bulbis subpalma:is rectis, nectarii cornu *latifolia.*
conico: labio trilobo lateribus reflexo, bracteis flore
longioribus. *Sp. pl.* 1334. *Fl. dan.* 246.

Orchis radicibus palmatis, caule fistuloso, bracteis
maximis, labello trifido ferrato: medio segmento
obtusio. *Hall. hist.* 1279. t. 32. *Orch.* 70.

Orchis palmata pratensis latifolia, longis calcaribus.
B. pin. 85. *R. syn.* 380. *Vaill. par.* 152. t. 31.
f. 1—5.

Orchis palmata mas s. Palma Christi mas. *Park.* 1356.
Palma Christi mas. *Ger. em.* 220.

Anglis, broad-leaved Orchis.

Habitat in pratis et pascuis. 4. V. VI.

9. ORCHIS bulbis palmatis patentibus, nectarii cornu *maculata.*
germinibus brevioribus: labio plano, petalis dorsalibus
patulis. *Sp. pl.* 1335.

Orchis radicibus palmatis, caule solido, labello trifido
ferrato medio segmento acuminato. *Hall. hist.*
1278. t. 32. *Orch.* 69.

Orchis palmata pratensis maculata. *B. pin.* 85. *R.*
syn. 381.

Orchis palmata foemina s. Palma Christi foemina; fo-
liis maculatis. *Park.* 1357.

Palma Christi foemina. *Ger. em.* 220.

Anglis, spotted Orchis.

Habitat in pratis, pascuis et nemorosis humidis. 4. VI.

10. ORCHIS bulbis palmatis, nectarii cornu setaceo ger- *conopsea.*
minibus longiore: labio trifido, petalis duobus pa-
tentissimis. *Sp. pl.* 1335. *Fl. dan.* 224.

Orchis radicibus palmatis, calcare longissimo, labello
trifido obtuso unicolore. *Hall. hist.* 1287. t. 29.
Orch. 84.

Orchis palmato minor calcaribus oblongis. *B. pin.* 85.
Vaill. par. t. 30. f. 8—10. *R. syn.*

Orchis palmata minor flore rubro. *Park.* 1358.

Serapias minor nitente flore. *Ger. em.* 222.
Anglis, sweet Orchis.
 Habitat in pratis et pascuis. 4. VI.

** *Bulbis fasciculatis.*

abortiva. 11. ORCHIS bulbis fasciculatis filiformibus, nectarii labio ovato integerrimo. *Sp. pl.* 1336. *Jac. austr.* II. 55. t. 193.
 Epipactis aphylla, nectario longo, labello ovato-lanceolato. *Hall. hist.* 1288. t. 36.
 Orchis abortiva violacea. *B. pin.* 86.
 Limodorum austriacum. *Clus. pan.* 242. *R. syn.* 383.
 Nidus avis flore et caule violaceo purpureo colore. *Ger. em.* 228.
 Nidus avis purpureus. *Park.* 1362.
Anglis, purple Orchis or Bird's-nest.
 Habitat in pratis siccioribus; in Southamptoniæ comitatu prope Alton oppidum invenit. D. Goodyer. *R. syn.* ? 4. V. VI.

S A T Y R I U M. 110.

Nectarium scrotiforme s. inflato-didymum pone florem.

hircinum. 1. SATYRIUM bulbis indivisis, foliis lanceolatis, nectarii labio trifido: intermedia lineari obliqua præmorsa. *Sp. pl.* 1337.
 Orchis radicibus subrotundis, labello longissimo tripartito plicato. *Hall. hist.* 1268. t. 25. *Orch.* 62.
 Orchis barbata odore hirci, brevior latioreque folio. *B. pin.* 82. *Hist. Ox.* III. 491. f. 12. t. 12. f. 9.
 Orchis barbata foetida. *B. hist.* II. 756. *R. syn.* 376.
 Tragorchis maxima et T. vulgaris. *Park.* 1348.
 Tragorchis maximus et T. mas. *Ger. em.* 210.
Anglis, Lizard Satyrion.
 Habitat in pratis et pascuis cretaceis; circa Dartford et Crayford in Cantio sed rarius. *R. syn.* VI. VII.

viride. 2. SATYRIUM bulbis palmatis, foliis oblongis obtusis, nectario labio lineari trifido: intermedia obsoleta. *Sp. pl.* 1337. *Fl. dan.* 77.

Orchis

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

* *Bulbis ramosis.**Nidus
avis.*

1. OPHRYS bulbis fibroso-fasciculatis, caule vaginato aphylo, nectarii labio bifido. *Sp. pl.* 1339. *Fl. dan.* 181.

Epipactis aphylla, flore inermi, labello bicorni. *Hall. hist.* 1290. t. 37. *Orch.* 47.

Nidus avis. *Lob. ic.* 195. *Lugd.* 673. *R. syn.* 382.

Orchis abortiva fusca. *B. pin.* 86.

Orchis abortiva rufa. f. *Nidus avis.* *Park.* 1362.

Satyrion abortivum f. *Nidus avis.* *Ger. em.* 228.

Anglis, bird's-nest Ophrys.

Habitat in sylvis et umbrosis. In sylvis circa Charlton, in Cantio. 4. V. VI.

Corallorhiza.

2. OPHRYS bulbis ramosis flexuosis, caule vaginato, nectarii labio trifido. *Sp. pl.* 1339. *Fl. dan.* 451.

Corallorhiza. *Hall. hist.* 1301. t. 44. *Orch.* 12.

Orobanche radice f. coralloide. *B. pin.* 88.

Orobanche spuria f. Corallorhiza. *Rup. gen.* 284. t. 2.

Dentaria minor. *Ger. em.* 1585. *Park.* 1387.

Anglis, corall Ophrys.

Habitat in sylvis pinetis in boreali parte Scotiæ. 4. VIII. IX.

ovata.

3. OPHRYS bulbo fibroso, caule bifolio, foliis ovatis, nectarii labio bifido. *Sp. pl.* 1340. *Fl. dan.* 137.

Epipactis foliis binis ovatis, labello bifido. *Hall. hist.* 1291. t. 37. *Orch.* 45.

Ophrys bifolia. *Ger. em.* 402. *B. pin.* 87.

Bifolium sylvestre vulgare. *Park.* 504. *R. syn.* 385.

Anglis, common Ophrys or Twayblade.

Habitat in sylvis et nemorosis humidis. 4. V. VI.

cordata.

4. OPHRYS bulbo fibroso, caule bifolio, foliis cordatis, nectarii labio bifido.

Ophrys bulbo fibroso, caule bifolio, foliis cordatis. *Sp. pl.* 1340. *Fl. suec.* 809.

Epipactis foliis binis cordatis, labello bifido, postice bidentato. *Hall. hist.* 1292. t. 22. f. 4. *Orch.* 43.

Ophrys minima. *B. pin.* 87. pr. 31.

Bifolium

Bifolium minimum. *B. hist. III. 534. R. syn. 385.*

Anglis, mountain Ophrys or Twayblade.

Habitat in ericetis montosis humidis, præsertim solo turfoso in comitatibus Eboracensi, Lancastriensi et Westmorlandico passim. 4. VI. VII.

5. OPHRYS bulbo fibroso, caule angulato, foliis vagi-paludosa. natis apice scabris, nectarii labio integro.

Ophrys bulbo subrotundo, scapo subnudo pentagono, foliis apice scabris, nectarii labio integro. *Sp. pl.*

1341.

Orchis minima bulbosa. *R. syn. 375.*

Orchis bifolia minor palustris. *Pluk. ph. t. 247. f. 2.*

Bifolium palustre. *Park. 505. R. syn. 385.*

Anglis, marsh Ophrys or Twayblade.

Habitat in paludibus. Found by Mr. Dent and Mr. Dale, on the boggy grounds near Gamlingay in Cambridgeshire; by Parkinson on the low wet grounds between Hatfield and St. Albans; in Rumney marsh; by Mr. Dubois at Hursthill, Tunbridge-wells. R. syn. prope Norwich. D. Pitchford. 4. VI. VII.

•• *Bulbis rotundis.*

6. OPHRYS bulbis aggregatis oblongis, caule subfolio-spiralis. so, floribus secundis, nectarii labio indiviso crenato.

Sp. pl. 1340. Fl. dan. 387.

Epipactis bulbis cylindricis, spica spirali, labello crenulato. *Hall. hist. 1294. t. 38. Orch. 36.*

Orchis spiralis alba odorata. *B. hist. II. 769. R. syn. 378.*

Triorchis. *Ger. em. 218. alba odorata minor. B. pin. 87. Park. 1354.*

Anglis, Triple Ophrys or Ladies Traces.

Habitat in pratis et pascuis siccis. 4. VII. VIII.

7. OPHRYS bulbo ovato, scapo angulato nudo, foliis lanceolatis, petalis linearibus, nectarii labio subrotundo integro.

Ophrys bulbo subrotundo, scapo nudo, foliis lanceolatis, nectarii labio integro, petalis dorsalibus linearibus. *Sp. pl. 1341.*

Epipactis foliis binis ovatis, bulbis piriformibus foliosa vagina cinctis. *Hall. Orch.* 40.

Pseudo-orchis bifolia palustris. *R. syn.* 382. *hist.* 1226.

Chamæorchis lilifolia. *B. pin.* 84.

Chamæorchis lilifolius minor fabuletorum Zeelandiæ, &c. *Lob. adv. pars. alt.* 506.

Helleborine bifolia, radice ovali, cucullo sparso. *Hall. act. gott.* 1751. 215. t. xiv.

Læselii. β Ophrys bulbo subrotundo scapo nudo trigono, nectarii labio ovato. *Sp. pl.* 1341.

Ophrys diphyllus bulbosa. *Læf. Fl. pruss.* 180. t. 58. *Anglis*, dwarf Ophrys.

Habitat in palustribus. On Hinton and Teversham moors near Cambridge. *R. syn. prope Norwich.* D. Pitchford. 4. VI. VII.

Desc. Radix bulbosus, turicatus, bulbo ovato infra radicato. Scapus semipedalis, erectus, tri et quinquangulatus, nudus, glaber. Folia radicalia duo lanceolata, integerrima, nudus, fere longitudine scapi. Flores racemosi a quatuor ad sex vel septem, luteo-viridi. Petala quinque, tria exteriora linearia, reflexa, viridia; duo interiora lateralia, filiformia subpurpurascens. Nectarii labium magnum, subrotundum integerrimum, subpurpurascens; Capsula erecta, oblonga, angulata, Semina numerosa, parva.

Monorchis. 8. OPHRYS bulbo globoso, scapo nudo, nectarii labio trifido. *Sp. pl.* 1342. *Fl. dan.* 102.

Orchis bulbo unico subrotundo, labello cruciformi. *Hall. hist.* 1262. t. 22. *Orch.* 53.

Orchis odorata moschata, s. monorchis. *B. pin.* 84. *R. syn.* 378.

Orchis pusilla odorata. *Park.* 1354. *Anglis*, musk Ophrys.

Habitat in pascuis sterilibus solo calcareo. 4. VII.

anthrophora. 9. OPHRYS bulbis subrotundis, scapo folioso, nectarii labio lineari tripartito: medio elongato bifido. *Sp. pl.* 1343. *Fl. dan.* 103.

Orchis radicibus subrotundis, spica longa, flore inermi, labello perangusto quadrifido. *Hall. hist.* 1264. *Orch.* 56.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Orchis sphegodes seu fucum referens. *Park.* 1350.
Anglis, bee Ophrys.

Habitat in pratis et pascuis cretaceis. 4. VII. VIII.

Obs. Corolla petala quinque; tria exteriora patentia oblonga purpurascens, trinervia, nervis viridibus; duo interiora lateralia, linearia, villosa, viridia; Nectarii labium magnum, subrotundum quinquelobum, coloratum, sericeum, atro-purpureum, lobis subtus inflexis, macula ad basin bilobata luteo variegata, glabra, nitida, utrinque maculae dente obtuso exerto.

aranifera.

12. OPHRYS bulbo subrotundo, scapo folioso, nectarii labio subrotundo integro emarginato convexo.

Orchis five Testiculus sphægodes hirsuto flore. *B. hist.* II. 767. *R. syn.* 380.

Orchis fucum referens colore rubiginoso. *B. pin.* 83. *Vaill. par. t.* 31. *f.* 15. 16.

Orchis araneam referens. *B. pin.* 84. *Vaill. par.* 146. *t.* 30. *f.* 10.

Orchis sphegodes altera. *Park.* 1351.

Testiculus vulpinus major sphægodes. *Ger. em.* 212.
Anglis, spider Ophrys.

Habitat in pratis et pascuis cretaceis. 4. IV. V.

Obs. Corolla petala quinque; tria exteriora oblonga, patentia, viridia; duo interiora, lateralia, linearia, herbacea, breviora. Nectarii labium magnum, subrotundum, integrum, emarginatum, convexum, coloratum, sericeum, ad basin macula bilobata, nitida, glabra, et utrinque maculae dente obtuso exerto, subtus concavo, supra atro purpureo, margine viride, infra herbacea.

CYPRIPEDIUM. 1015.

Nectarium ventricosum, inflatum, cavum.

Calceolus. 1. CYPRIPEDIUM radicibus fibrosis, foliis ovato-lanceolatis caulinis. *Sp. pl.* 1346. *Mill. ic. t.* 242. *Hall. hist.* 1300. *Orch.* 14.

Calceolus mariae. *Ger. em.* 443. *R. syn.* 385.

Helleborine flore rotundo s. *Calceolus.* *B. pin.* 187.

Elleborine major seu *Calceolus mariae.* *Park.* 217.

Anglis,

Anglis, Ladies-slipper.

Habitat in sylvis et dumetis, sed rarius, in sylvis circa Clapham et Ingleton in comitatu Eboracensi, sed sparsim. 4. VII.

S E R A P I A S. 1012.

Nectarium ovatum, gibbum, labio ovato. /

1. SERAPIAS bulbis fibrosis, nectarii labio obtuso petalis æquali, fructibus pendulis. *Fl. angl.* 341.
 Serapias bulbis fibrosis, foliis ovatis amplexicaulibus, floribus pendulis. *Syst. nat.* 593.
 Epipactis foliis amplexicaulibus ovato-lanceolatis, labello lanceolato. *Hall. hist.* 1297. t. 40. *Orch.* 28.
 Helleborine. *Dod. pempt.* 387. *Ger. em.* 442. latifolia montana. *B. pin.* 186. *R. syn.* 383.
 Elleborine flore viridente. *Park.* 218.
 β Helleborine altera atro-rubente flore. *B. pin.* 186. *R. syn.* 383.
- γ Serapias bulbis fibrosis, foliis ensiformibus sessilibus, palustris. floribus pendulis. *Syst. nat.* 593. *Fl. dan.* 267.
 Epipactis foliis ensiformibus caulinis, floribus pendulis, labello obtuso per oras plicato. *Hall. hist.* 1296. *Orch.* 31.
 Helleborine palustris nostras. *R. syn.* 384.
 Helleborine angustifolia palustris. *B. pin.* 87.
Anglis, common γ marsh Helleborine.
 Habitat in sylvis, nemorosis, et sepibus γ in palustribus. 4. VII. VIII.
 Obs. γ in horto aut solo arido illata, secundo anno in ævadit.
3. SERAPIAS bulbis fibrosis, nectarii labio obtuso petalis brevioribus, fructibus erectis. *Fl. angl.* 341.
 Serapias bulbis fibrosis, foliis ensiformibus, floribus grandierectis, nectarii labio obtuso petalis brevioribus. *Syst. flora. nat.* 594.
 Epipactis caule paucifloro, lineis in obtuso labello levibus. *Hall. hist.* 1298. t. 41. *Orch.* 25.
 Helleborine flore albo. *Ger. em.* 442. *R. syn.* 383. vel Damasonium montanum latifolium. *B. pin.* 187.
 Elleborine minor flore albo. *Park.* 218.
 β Helle-

GYNANDRIA HEXANDRIA.

- β Helleborine latifolia, flore albo clauso. *R. syn.* 384.
 γ Helleborine foliis prælongis angustis acutis. *R. syn.* 385.
 δ Helleborine montana angustifolia purpurascens. *B.*
pin. 187.

Anglis, white Helleborine.

Habitat in sylvis et dumetis; in the woods near Stokenchurch, Oxfordshire. *R. syn.* in sylvis circa Marlow et in sylvis in comitatibus Bedfordiensi, Buckinghamensi, &c. frequens; γ δ in dumetis ad latus mortium circa Clapham et Ingleton, &c. in comitatu Eboracensi. 4. VIII.

H E X A N D R I A.

A R I S T O L O C H I A. 1022.

Hexagynia Cal. o. *Cor.* monopetala, lingulata, integra. *Caps.* sexlocularis infera,

clematidis. 1. A R I S T O L O C H I A foliis cordatis, caule erecto, floribus axillaribus confertis. *Sp. pl.* 1364. *Hall. hist.* 1029.

Aristolochia clematidis. *Ger. em.* 847. *Park.* 292. *recta.* *B. pin.* 307.

Anglis, climbing Birth-wort.

Habitat in sylvis et sepibus; in a wood two miles from Thorndon, Essex; and also in Cambridgeshire. *Blackf. spec. prope Maidston et alibi in Cantio.* 4, VII.

Pharmac. *Aristolochiæ clematidis Radix.*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- *Alga angustifolia vitrariorum.* *R. syn.* 53.
Anglis, Grass-wrack.
Habitat in littoribus maritimis. 24. VI—VIII.
Obs. variat seminibus pedicellatus et sessilibus in eodem
spadice sed plerisque pedicellatus.

Classis XXI.

M' O N O E C I A :

M O N A N D R I A.

ZANNICHELLIA. 1034.

Masculi. *Cal.* o. *Cor.* o. *Stam.* Filamentum unicum longum, erectum. *Anthera* ovata, erecta.

Feminei. *Cal.* monophyllus. *Cor.* o. *Germina* circiter quatuor. *Semina* totidem.

I. ZANNICHELLIA. *Sp. pl.* 1375. *Hall. hist.* 1640. *palustris.*

Zannichellia palustris major, foliis gramineis acutis, &c. *Mich. gen.* 71. t. 34. f. 1. 2.

Aponogiton aquaticum graminifolium, staminibus singularibus. *Pont. anth.* 117. *R. syn.* 135.

Potamogetoni similis graminifolia ramosa et ad genicula polyceratos. *Pluk. ph. t.* 132. f. 7.

Patamogeton capillaceum, capitulis ad alas trifidis. *B. pin.* 193. *pr.* 101.

Anglis, common Zannichellia.

Habitat in fossis et stagnis frequens. ☉. VII.

Obs. variat seminis dorso integro et dentato.

CHARA. *Mantiss. I.* 23. *syst. nat.* 615.

Masc. *Cal.* o. *Cor.* o. *Anthera* germini subjecta.

Fem. *Cal.* tetraphyllus. *Cor.* o. *Stigma* quinquefidum. *Sem.* unicum.

I. CHARA caulibus lævibus, frondibus interne dentatis. *vulgaris.*

Sp. pl. 1624. *Fl. dan.* 150.

Chara caule levi fragile. *Hall. hist.* 1681.

Chara vulgaris foetida. *Act. paris.* 1719. p. 17. t. 3.

f. 1. *R. syn.* 132.

Equisetum

Equisetum foetidum sub aqua repens. *Ger. em.* 1115.

B. pin. 16. *pr.* 25. *hist.* III. 731. *Park.* 1201.

β Chara minor caulibus et foliis tenuissimis. *Act. paris.* 1719. p. 18.

Anglis, common Chara.

Habitat in fossis et stagnis. ☉. VII. VIII.

tomentosa. 2. CHARA aculeis caulinis ovatis. *Sp. pl.* 1624.

Chara major subcinerea fragilis. *Act. paris.* 1719. p. 18.

R. syn. 132.

Chara caule tomentoso, fragili. *Hall hist.* 1683.

Equisetum fragile majus subcinereum aquis immersum.

Hist. Ox. III. 621. t. 4. f. 9.

Equisetum s. Hippuris coralloides. *Ger. em.* 1115.

Equisetum s. Hippuris lacustris, foliis mansu arenosis.

Pluk. ph. t. 21. f. 4.

Anglis, brittle Chara.

Habitat in fossis et stagnis; in fossis turfosis in comitatibus Lancastriensi et Westmorlandico frequens. ☉. VI—X.

bispida. 3. CHARA aculeis caulinis capillaribus confertis. *Sp. pl.* 1624. *Pl. den.* 154.

Chara caule fragili, spinoso. *Hall. hist.* 1682.

Chara major caulibus spinosis. *Act. paris.* 1719. p. 18.

t. 3. f. 3. *R. syn.* 132.

Equisetum s. Hippuris muscosus sub aqua repens.

Pluk. ph. t. 193. f. 6.

Anglis, prickly Chara.

Habitat in fossis et stagnis; in comitatibus Eboracensi, Lancastriensi et Westmorlandico. VI—X.

flexilis. 4. CHARA caule trichotomo diaphano inermi.

Chara caulium articulis inermibus diaphanis superne latioribus. *Sp. pl.* 1624.

Chara translucens major flexilis. *Act. paris.* 1719. p. 18. f. 8.

Chara translucens minor flexilis. *Act. paris.* 1719. p. 18. f. 9. *R. syn.* 133.

Anglis, smooth Chara.

Habitat in fossis et stagnis. In the ponds about Ipswich, Suffolk.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

MONOECIA TRIANDRIA.

Lenticula aquatica. *B. pin.* 362. *R. syn.* 129.
 Hederula aquatica. *Ger. em.* 830.
 Ranunculus hederaceus aquaticus. *Park.* 1260.
Anglis, Ivy-leaved Duck's-meat.
 Habitat in fossis et stagnis. ☉. VI—IX.

T R I A N D R I A.

T Y P H A. 1040.

Mas. *Amentum* cylindricum. *Cal.* obsoletus, tri-
 phyllus. *Cor.* 0.

Fem. *Amentum* cylindricum, infra masculos. *Cal.*
 capillo villoso. *Cor.* 0. *Sem.* unicum, insidens
 pappo capillari.

latifolia. 1. TYPHA foliis subensiformibus, spica mascula femi-
 neaque approximatis. *Sp. pl.* 1377.
 Typha clava unica. *Hall. hist.* 1205.
 Typha. *Cam emp.* 677. *Ger. em.* 46. palustris major.
B. pin. 20. *hist. II.* 539. *R. syn.* 436.
 Typha palustris maxima. *Park.* 1204.
Anglis, greater Cat's-tail or Reed-mace.
 Habitat ad ripas fluviorum, piscinarum et paludium fre-
 quens. ♀. VII.

angustifolia. 2. TYPHA foliis semicylindricis, spica mascula feminea-
 que remotis. *Sp. pl.* 1377.
 Typha clava mascula a femina remota. *Hall. hist.*
 1306.
 Typha palustris media. *B. hist. II.* 540. *R. syn.* 436.
 Typha palustris clava gracili. *B. pin.* 20.
 Typha minor. *Park.* 1204.
 • β Typha minor. *Lob. ic.* 81. *B. hist. II.* 540.
 Typha palustris minor. *B. pin.* 20. *R. syn.* 436.
Anglis, lesser Cat's-tail or Reed-mace.
 Habitat in fossis et stagnis. ♀. VII.

S P A R G A N I U M. 1041.

Mas. *Amentum* subrotundum. *Cal.* triphyllus. *Co.* 0.

Fem. *Amentum* subrotundum. *Cal.* triphyllus. *Cor.* 0.
Stigma bitidum. *Drupa* exsucca, monosperma.

1. SPAR-

1. SPARGANIUM foliis ensiformibus triquetris, caule *ramosum*.
ramoso.

Sparganium foliis erectis triquetris. *Sp. pl.* 1378. erectum.

Sparganium caule foliisque erectis. *Hall. hist.* 1303.

Sparganium. *Cam. ep.* 732. ramosum. *Ger. em.* 45.

Park. 1205. *B. pin.* 15. *R. syn.* 437.

Anglis, greater Bur-reed.

Habitat in fossis, paludibus et ad ripas fluviorum. 24.

VII.

2. SPARGANIUM foliis ensiformibus planis, caule *simplex*.
simplici.

Sparganium non ramosum. *B. pin.* 15. *Park.* 1205.

R. syn. 437. five latifolium. *Ger. em.* 45.

3. Sparganium foliis decumbentibus planis. *Sp. pl.* 1378. natans.

Fl. dan. 260.

Sparganium foliis planis mollibus decumbentibus.

Hall. hist. 1304.

Sparganium minimum. *R. syn.* 437. *hist.* 1910..

Anglis, lesser Bur-reed.

Habitat in palustribus et ad ripas stagnorum et lacuum &
in stagnis, lacubus et fluviis leniter fluentibus; circa
Norwich. D. Rose. Circa Llanberris, et in comi-
tatibus Eboracensi, Westmorlandico passim. 24, VII.

C A R E X. 1046.

Maf. *Amentum* imbricatum. *Cal.* monophyllus. *Cor.* 0.

Fem. *Amentum* imbricatum. *Cal.* monophyllus, *Cor.* 0.

Nectarium inflatum, tridentatum. *Stigmata* tria.

Sem. triquetrum, intra nectarium.

• *Spica unica simplici.*

1. CAREX spica simplici dioica. *Sp. pl.* 1379. *Fl. dan. dioica.*
369.

Gramen cyperoides minimum, spica simplici casta.

Hist. Ox. III. 244. *f.* 8. *t.* 12. *f.* 22.

Gramen cyperoides minus, ranunculi capitulo lon-
giore. *R. syn.* 425.

Cyperoides parvum, caulibus et foliis tenuissimis, tri-

D d

angularibus,

angularibus, spica longiori, capsulis oblongis, in angustum collum vix bifidum attenuatis. *Mich. gen. 56. t. 32. f. 1.*

Anglis, small Carex.

Habitat in paludibus turfosis. 4. VI. VII.

capitata. 2. CAREX spica simplici androgyna: superne mascula, capsulis imbricato-patulis. *Sp. pl. 1379. Fl. dan. 372?*

Carex spica unica subrotunda, seminibus inflatis. Hall. hist. 1351.

Cyperoides parvum, caulibus et foliis tenuissimis triangularibus, spica subrotunda, capsulis oblongis in angustum collum vix bifidum attenuatis. Mich. gen. 56. t. 32. f. 2.

Gramen cyperoides minimum, ranunculi capitulo rotundo. Hist. Ox. III. 245. f. 8. t. 12. f. 36. R. syn. 425.

Anglis, round-headed Carex.

Habitat in paludibus turfosis; in palustribus prope Hinton et Feverham in agro Cantabrigienſi; in Sherley common, prope Croydon. 4. VI.

patula. 3. CAREX spica simplici androgyna oblonga: superne mascula, capsulis oblongis acuminatis patulis.

Anglis, patulous Carex.

Habitat in ericetis montosis; prope Glasgow in Scotia. 2. VI.

Desc. Radix perennis ramosa. Culmus quadripollicaris, erectus, triangularis, foliatus, striatus. Folia radicalia pauca, ensiformia, acuta; nuda; caulina duo culmo breviora. Spica terminalis erecta, oblonga, androgyna, superne mascula, bi s. triflora fusca; feminea tri s. quinqueflora virescentia; Capsula oblonga, patula.

pulicaris. 4. CAREX spica simplici androgyna: superne mascula, capsulis divaricatis retroflexis. *Sp. pl. 1380. Fl. dan. 166.*

Carex spica unica terete, seminibus rostratis, Hall. hist. 350.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- arenaria.* 7. CAREX spica composita, spiculis subrotundis androgynis: inferioribus remotioribus foliolo longiori instructis, culmo triquetro. *Sp. pl.* 1381.
 Carex spicis stipula brevioribus, imis distinctis, superioribus congestis. *Hall. hist.* 1363?
 Carex maritima humilis, radice repente, caule trilatero, spica spadicea nonnihil foliosa, capitulis crassioribus capsulis marginatis. *Mich. gen.* 67. t. 33. f. 4.
 Gramen cyperoides ex monte Ballon simile humilius, in maritimis et arenosis nascens. *R. syn.* 423. *Pluk. ph.* t. 34. f. 7.
Anglis, sea Carex.
 Habitat in arenosis maritimis. 4. VI.
- leporina.* 8. CAREX spica composita, spiculis ovatis sessilibus approximatis androgynis nudis. *Sp. pl.* 1381. *Fl. dan.* 371?
 Carex spicis nudis subrotundis distinctis, in summo caule congestis. *Hall. hist.* 1361.
 Carex angustifolia, caule triquetro, spicis pluribus elegantibus parum inter se distantibus. *Seg. per.* I. 124. t. 1. f. 2.
 Gramen cyperoides spica e pluribus spicis brevibus mollibus composita. *R. syn.* 422. *Sch. agr.* 456. t. 10. f. 15.
 Gramen cyperoides palustre majus, spica divisa. *B. pin.* 6. *Hist. Ox.* III. 244. f. 8. t. 12. f. 29.
Anglis, naked Carex.
 Habitat in palustribus et aquosis. 4. VI.
 Obs. Spica e spiculis a tribus ad sex s. septem composita; foliolo s. bractea, ad spiculam inferiorem spica æquante, caduco.
- vulpina.* 9. CAREX spica supradecomposita inferne laxiore, spiculis ovatis androgynis glomeratis, superne masculis. *Sp. pl.* 1382. *Fl. dan.* 304.
 Carex spicis subrotundis echinatis, supremis confluentibus, capsulis rostratis bifidis. *Hall. hist.* 1364.
 Carex palustris major, radice fibrosa, caule exquisite, triangulari,

triangulari, spica brevi, habitiori compacta. *Mich. gen.* 69.

Gramen cyperoides palustre majus. *Park.* 1266. spica compacta. *B. pin.* 6. *R. syn.* 423. *Hist. Ox.* III. 244. *f.* 8. *t.* 12. *f.* 24.

Gramen palustre cyperoides. *Ger. em.* 18.

Anglis, great Carex.

Habitat in palustribus et ad ripas fluviorum. *4. V. VII.*

10. CAREX spiculis subrotundis androgynis contiguis, *spicata.* capsulis ovatis acutis. *Fl. angl.* 349.

Carex palustris media, radice fibrosa, caule exquisite triangulari, spica brevi compactiori. *Mich. gen.* 69. *t.* 33. *f.* 14.

Gramen cyperoides parvum. *Ger. em.* 21.

Gramen cyperoides spicis minoribus minusque compactis. *B. pin.* 6. *pr.* 13. *hist.* II. 497.

Gramen cyperoides spicatum minus. *R. syn.* 424.

Anglis, spiked Carex.

Habitat in aquosis et ad margines fossarum. *4. VI.*

Obs. variat spica nuda et bracteata.

11. CAREX spica composita, spiculis rotundis remotiusculis androgynis, foliolo longiori instructis, capsulis acutis, culmo teretiusculo. *Fl. angl.* 348. *Fl. dan.* 371. *divisa.*

Gramen cyperoides palustre spica divulsa. *B. pin.* 6.

Gramen cyperoides ex monte Ballon spica divulsa. *B. hist.* II. 498. *R. syn.* 423.

Gramen cyperoides palustre minus. *Park.* 1287?

Anglis, marsh Carex.

Habitat in palustribus maritimis; in the meadows near the Hithe at Colchester in Essex; by Hithe in Kent. *R. syn.* *4. VI. VII.*

12. CAREX spiculis subrotundis remotis sessilibus obtusiusculis androgynis, capsulis ovatis obtusiusculis. *Sp. pl.* 1385. *Fl. dan.* 285. *canescens.*

Carex spicis ovatis alternis, supremis se contingentibus. *Hall. hist.* 1560.

MONOECIA TRIANDRIA.

Carex nemorosa, fibrosa radice, caule exquisite triangulari, spica longa divulsa seu interrupta, capitulis solitariis præterquam ultimo. *Mich. gen.* 69. t. 23. f. 10.

Gramen cyperoides spicatum minus, spica longa divulsa seu interrupta. *R. syn.* 424.

Gramen cyperoides spicis curtis divulsis. *Læs. pr.* 117. t. 32.

Anglis, grey Carex.

Habitat in nemorosis humidis. 4. V. VI.

muricata. 13. CAREX spiculis subovatis sessilibus remotis androgynis, capsulis acutis divergentibus spinosis. *Sp. pl.* 1382. *Fl. dan.* 284.

Carex spicis ternis echinatis, glumis lanceolatis, capsulæ mucrone simplici. *Hall. hist.* 1366.

Carex nemorosa, fibrosa radice, angustifolia minima, caule exquisite triangulari, spica brevi interrupta. *Mich. gen.* 69. t. 33. f. 12.

Gramen cyperoides spicatum minimum, spica divulsa aculeata. *R. syn.* 424. *Sch. agr.* 485. t. 11. f. 3.

Gramen cyperoides echinatum minimum. *Park.* 1272.

Gramen nemorosum spicis parvis asperis. *B. pin.* 7. *tb.* 100. *Hist. Ox. III.* 344. f. 8. t. 12. f. 26.

Anglis, prickly Carex.

Habitat in aggeribus fossarum et palustribus. 4. VII.

brizoides. 14. CAREX spica composita disticha nuda, spiculis androgynis oblongis cōtiguis, culmo nudo. *Sp. pl.* 1381.

Carex spicis distichis teretibus repandis. *Hall. hist.* 1358.

Carex fibrata radice, angustifolia, caule exquisite triangulari, capitulis pulchellis albicantibus, paululum inter se distantibus. *Mich. gen.* 70. t. 33. f. 18.

Gramen cyperoides palustre elegans, spica composita asperiore. *R. syn.* 423. *Hist. Ox. III.* 245.

Anglis, rough Carex.

Habitat in palustribus et aquosis, sed rarius. 4. VI.

15. CAREX

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

MONOECIA TRIANDRIA.

perna parte pilosis, caule tenuissimo rotundo-tri-
quetro, spicis parvis atris, squamis obtusioribus,
medio nervo tanquam in aristam prolongato, capsu-
lis oblongis crassioribus subhirsutis trilateris turbi-
natis. *Mich. gen.* 64. *N.* 68. et 69. t. 32. f. 3.
Gramen spicatum angustifolium montanum. *B. pin.*
4. th 47.

Anglis, mountain Carex.

Habitat in pratis et ericetis alpinis; prope Sedberg in
comitatu Eboracensi. 4. VI.

Desc. Radix repens; Culmus spithameus, erectus, tri-
queter, basi foliatus, superne nudus. Folia radicalia
plurima, caulina duo vel tria, linearia, margine scabra,
nuda, glauca. Spica mascula terminalis, oblonga, atro-
fusca. Spicæ femineæ duæ vel tres approximate, ses-
siles, ovatæ; squamæ atro-purpuræ, obtusiusculæ, nu-
dæ. Capsulæ imbricatæ ovatæ, triquetrae, obtusius-
culæ, pallidæ, herbacæ, villosæ. Bractea ad infimam
spicam femineam longitudine spicæ. Habitus Caricis
piluliferæ.

pilulifera. 18. CAREX spicis terminalibus confertis subrotundis :
masculâ oblonga. *Sp. pl.* 1385.

Gramen cyperoides tenuifolium spicis ad summum
caulem sessilibus globulorum æmulis. *R. syn.* 422.
Pluk. ph. t. 91. f. 8.

β Gramen cyperoides spicis brevibus congestis, folio
molli. *R. syn.* 421.

Anglis, round-headed Carex.

Habitat in pascuis et ericetis humidis passim. 4. VI.

saxatilis. 19. CAREX spicis tribus ovatis sessilibus alternis, mas-
cula oblonga. *Sp. pl.* 1385: *Fl. dan.* 159.

Carex spicis maribus ovatis, femininis gracilibus erec-
tis, capsulis triquetro-conicis. *Hall. hist.* 1406.

Gramen cyperoides venum minimum. *R. syn.* 421.

Gramen cyperoides spicatum. *Ger. em.* 22.

Gramen caryophyllatæ foliis, spica divulsa. *B. pin.* 3.
th. 46.

Gramen spicatum foliis caryophylleis. *Park.* 1160.

Anglis, vernal Carex.

Habitu

Habitat in pratis et ericetis montosis. 4. IV. V.

*** Spicis sexu distinctis: femineis pedunculatis.

20. CAREX spica androgynis terminalibus peduncula- *atrata*
tis: florentibus erectis, fructiferis pendulis. *Sp. pl.*
1386. *Fl. dan.* 158.

Carex spicis ovatis petiolatis congestis, senescentibus
pendulis. *Hall. hist.* 1369.

Gramen alpinum pulchrum, foliis caryophyllæis, spicis
atris et tumentibus. *Sch. agr.* 481. t. 11. f. 1. 2.

Anglis, black Carex.

Habitat in montibus Cambrobritannicis; circa Llan-
berris; copiose ex Scotia recepi. 4. VI. VII.

21. CAREX spicis linearibus erectis: mascula brevior *digitata*
inferioreque, bracteis aphyllis, capsulis distantibus.
Sp. pl. 1384.

Carex spicis femininis petiolatis raris masculam supe-
rantibus. *Hall. hist.* 1376.

Cyperoides montanum nemorosum, caule triquetro-
compresso, spicis ferrugineis tenuioribus inter se
distantibus, capsulis rarius dispositis oblongis tur-
binatis trilateris. *Mich. gen.* 65. t. 32. f. 9.

Gramen caryophyllatum montanum, spica varia. *B.*
pin. 4. *pr.* 23. *th.* 48. *Sch. agr.* 448.

Gramen caryophyllatum polycarpon fructu triangulo.
Læf. pruss. 112. t. 17.

Anglis, digitated Carex.

Habitat in sylvis et nemorosis; prope Bath. D. Sole. 4.
V. VI.

22. CAREX spicis ovatis pendulis: mascula longiore *limosa*
erectiore, radice repente. *Sp. pl.* 1386.

Carex spicis femininis pendulis, capsulis ovatis com-
pressis. *Hall. hist.* 1392.

Cyperoides spica pendula brevior, squamis e spadiceo
vel fusco rutilante viridibus. *Sch. agr.* 443. t. 10.
f. 13.

Anglis, brown Carex.

Habitat in paludibus turfosis in comitatibus Eboracensi,
Lancastriensi, Westmorlandico, &c. *passim.* 4. VI.

23. CAREX

panicea. 23. CAREX spicis pedunculatis erectis remotis, feminis linearibus, capsulis obtusiusculis inflatis. *Sp. pl.* 1387. *Fl. dan.* 267.

Carex spica mare una pluribusve, femininis petiolatis, erectis, capsulis raris maximis ovato-triquetris bidentatis. *Hall. hist.* 1405.

Gramen cyperoides foliis caryophylleis, spicis e rarioribus et tumidioribus granis compositis. *R. syn.* 418.

Pluk. ph. t. 91. *f.* 7.

Cyperoides foliis caryophylleis (caule rotundo-triquetro) spicis e rarioribus et tumidioribus vesicis compositis. *Mich. gen.* 61, *t.* 32. *f.* 11.

Anglis, pink Carex.

Habitat in pratis et pascuis humidis. 4. VI.

pallidescens. 24. CAREX spicis pendulis : mascula erecta, femineis ovatis imbricatis, capsulis confertis obtulis. *Sp. pl.* 1386.

Carex spicis femininis pendulis, capsulis ovato-conicis. *Hall. hist.* 1393.

Gramen cyperoides polystachyon flavicans, spicis brevibus prope summitatem caulis. *R. syn.* 419. *Hist. Ox. III.* 243. *f.* 8. *t.* 12. *f.* 16. *Pluk. ph. t.* 37. *f.* 5.

Mich. gen. 61, *t.* 32. *f.* 13.

Anglis, pale Carex.

Habitat in pratis et pascuis humidis. 4. VI.

Pseudo-cyperus. 25. CAREX spicis confertis pedunculatis pendulis, mascula erecta, femineis cylindricis, capsulis aristatis furcatis.

Carex spicis pendulis, pedunculis geminatis. *Sp. pl.* 1387.

Carex spicis femininis pendulis asperis, capsulis longe rostratis retroversis. *Hall. hist.* 1397.

Gramen cyperoides spica pendula brevior. *B. pin.* 6. *th.* 86. *Hist. Ox. III.* 242. *f.* 8. *t.* 12. *f.* 5. *R. syn.* 419.

Pseudo cyperus. *Dod. pempt.* 339. *Ger. em.* 29.

Cyperus s. Pseudo-cyperus spica pendula brevior. *Park.* 1266.

Anglis;

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Obs. *Squamæ amenti ovatæ, acuminatæ.*

distans.

29. CAREX spicis remotissimis subsessilibus, bractea vaginante, capsulis angulatis mucronatis. *Sp. pl.* 1387.

Carex spicis brevissime petiolatis diffisis. *Hall. hist.* 1482.

Gramen cyperoides palustre, spicis tribus subrotundis vix aculeatis, spatio distantibus. *Hist. Ox. III.* 243. f. 8. t. 12. f. 18.

Gramen cyperoides spicis parvis longissime distantibus. *R. syn.* 420.

Anglis, loose Carex.

Habitat in palustribus. 4. VI.

inflata.

30. CAREX spicis remotis subpedunculatis erectiusculis, mascula erecta lanceolata, capsulis ovatis acuminatis calycis duplo longioribus.

Cyperoides sylvarum angustifolium, caule trilatere, spicis parvis, capsulis rarius dispositis obesis, in angustum collum vix ab una parte, ab altera vero magis bifidum coarctatis. *Mich. gen.* 56. t. 33. f. 5?

Anglis, bottle Carex.

Habitat in palustribus, in Wallia. D. Davies. In comitatibus Eboracensi et Westmorlandico passim. 4. VI. VII.

Obs. *Affinis Carici distantis. Spica mascula frequentissime solitaria, rarissime duæ.*

caespitosa. 31. CAREX spicis erectis cylindricis ternis subsessilibus: mascula terminali, culmo triquetro. *Sp. pl.* 1388.

Carex spicis masculis duabus, femininis sessilibus, capsulis ovatis brevissime mucronatis. *Hall. hist.* 1400.

Gramen cyperoides foliis caryophylleis, spicis erectis sessilibus, e feminibus confertis compositis. *R. syn.* 418.

Gramen caryophylleum angustissimis foliis, spicis sessilibus brevioribus erectis non compactis. *Hist. Ox. III.* 243.

Anglis, turfy Carex.

Habitat in pratis, pascuis et nemorosis humidis. 4. V. VI.

Obs.

Obs. *variat frequentissime spicis masculis duabus.*

32. CAREX spicis confertis pedunculatis cylindricis sub-*recurva*, pendulis, mascula terminali, capsulis imbricatis obtusiusculis.

Carex spicis maribus una pluribusve, femininis pendulis, capsulis ovato-triquetris confertis. *Hall hist.* 1408.

Gramen cyperoides foliis caryophylleis, spicis oblongis e pediculis longioribus pendulis. *R. syn.* 418.

Gramen cyperoides nemorosum, spica subnigra recurva. *Hist. Ox. III.* 243. *f.* 8. *t.* 12. *f.* 14.

Anglis, heath Carex.

Habitat in pratis, pascuis et ericetis. γ . V. VI.

Obs. Culmus erectus, triqueter, superne nudus; Calyx obtusiusculus, atro-fuscus. Capsula obtusa pallide herbacea, *variat spicis masculis duabus.*

**** Spicis sexu distinctis: masculis pluribus.

33. CAREX spicis masculis pluribus, femineis subpedunculatis erectis, capsulis ovato-lanceolatis aristato-acuminatis furcatis.

Carex spicis masculis pluribus, femineis subsessilibus, capsulis obtusiusculis. *Sp. pl.* 1388.

Carex spicis masculis ternis, femininis numerosis erectis, brevissime petiolatis capsulis bicornibus. *Hall. hist.* 1404.

Gramen cyperoides. *Ger. em.* 12. latifolium spica rufa s. caule triangulo. *B. pin.* 6.

Gramen cyperoides cum paniculis nigris. *B. hist.* II. 494. *R. syn.* 417.

Gramen cyperoides majus latifolium. *Park.* 1265.

β Gramen cyperoides majus angustifolium. *Park.* 1265. *Ger. em.* 29.

γ Gramen cyperoides minus angustifolium. *Park.* 1266. *Anglis*, acute Carex.

Habitat in aquosis et ad rivulos. γ . V.

34. CAREX spicis masculis pluribus, femineis subpedunculatis, capsulis globosis inflatis acuminatis.

Carex spicis masculis pluribus, femineis pedunculatis, capsulis inflatis acuminatis. *Sp. pl.* 1388. *Fl. dan.* 647.

Carex

MONOECIA TETRANDRIA.

Carex spicis maribus una pluribusve, femininis erectis teretibus, capsulis ampullatis rostratis. *Hall. hist.* 1401.

Gramen cyperoides majus præcox, spicis turgidis teretibus flavéscentibus. *Hist. Ox. III. 242. f. 8. t. 12. f. 6. R. syn. 420.*

β Gramen cyperoides polystachyon majus, spicis teretibus erectis. *R. syn. 419.*

Anglis, bladder Carex.

Habitat in palustribus et aquosis. 4. V.

Obs. variat spicis sessilibus et pedunculatis.

hirta.

35. CAREX spicis remotis, masculis pluribus, femineis subpedunculatis erectis, capsulis hirtis. *Sp. pl. 1388. Fl. dan. 319.*

Carex caule foliis capsulisque hirsutis. Hall. hist. 1403.

Gramen cyperoides polystachyon lanuginosum. *R. syn. 418. Hist. Ox. III. 243. f. 8. t. 12. f. 10.*

Gramen cyperoides norvegicum parium lanosum. *Park. 1172.*

Anglis, hairy Carex. 4.

Habitat in pratis et pascuis humidis. 4, VI.

T E T R A N D R I A.

N A S M Y T H I A.

CAL. *Perianthium commune* hemisphæricum : foliolis orbiculatis, subæqualibus, appressis ; persistens, continens *Florem* hemisphæricum, aggregatum, ex flosculis numerosissimis.

Masculi Flosculi in medio disci.

CAL. *Perianthium proprium* diphyllum ; foliolis cuneiformibus, concavis.

COR. Monopetala, infundibuliformis, bipartita, obtusa.

STAM. *Filamenta* quatuor, corollulæ inserta, breviora. *Antheræ* subrotundæ.

Feminei Flosculi in ambitu copiosissimi.

CAL. *Perianthium proprium* diphyllum : foliolis cuneiformibus, concavis.

COR.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

BETULA. 1052.

Masc. *Cal.* monophyllus, trifidus, triflorus. *Cor.*
quadripartita.

Fem. *Cal.* monophyllus, subtrifidus, biflorus. *Sem.*
utrinque membrana alata.

alba.

1. BETULA foliis ovatis acuminatis serratis. *Sp. pl.*
1393.

Betula foliis cordato-lanceolatis serratis, *Hall. hist.*
1628.

Betula. *B. pin.* 427, *hist.* 1. 148. *Ger. em.* 1478. *R.*
syn. 443.

Anglis, common Birch-tree. *Hunt. ev. sylv.* 225.

Habitat in sylvis et sepibus humidis. *h.* VII.

nana.

2. BETULA foliis orbiculatis crenatis. *Fl. lapp.* 266.
t. 6. *f.* 4. *Sp. pl.* 1394. *Hall. hist.* 1629. *Fl. dan.*
91.

Betula pumila, foliis subrotundis. *Amm. ruth.* 180.
cor. petrop. IX. 319. *t.* 14.

Anglis, dwarf Birch-tree.

Habitat in montosis et ericetis humidis. Found by Sir J.
Nasmyth in Twedale. *h.*

alnus.

3. BETULA pedunculis ramosis. *Sp. pl.* 1394.
Alnus foliis glabris rotundis, rotunde crenatis, spon-
giolis ad nervum angulis. *Hall. hist.* 1630.

Alnus. *Cam. ep.* 60. *Ger. em.* 1477. *R. syn.* 442.
vulgaris. *Park.* 1408.

Alnus rotundifolia glutinosa vulgaris. *B. pin.* 428.

β Alnus vulgaris, sub conis ligulis membranaceis rubris.
R. syn. 442.

Anglis, common Alder-tree. *Hunt. ev. sylv.* 240.

Habitat in aquosis. *h.* VII.

BUXUS. 1053.

Masc. *Cal.* triphyllus. Petala duo, Germinis rudi-
mentum.

Fem.

Fem. *Cal.* tetraphyllus. *Petala* tria. *Styli* tres. *Caps.* trilobris, trilocularis. *Sem.* duo.

1. BUXUS. *Cam. ep.* 601. *Ger. em.* 1410. *R. syn. semperdi-*
445. *Hall. hist.* 1610. *Sp. pl.* 1394: arborescens, rens.
| *B. pin.* 471.

Buxus arbor vulgaris. *Park.* 1428.

β *Buxus angustifolia.* *R. syn.* 445.

Anglis, Box-tree.

Habitat in sylvis et sepibus. h. IV:

Pharmac. *Buxi Folia, Lignum.*

U R T I C A. 1054:

Masc. *Cal.* tetraphyllus. *Cor.* nulla. *Neclarium* cen-
trale cyathiforme.

Fem. *Cal.* bivalvis. *Cor.* o. *Sem.* unicum, nitidum.

1. URTICA foliis oppositis ovatis serratis, amentis fruc- *pilulife-*
tiferis globosis. *Sp. pl.* 1395. *ra.*

Urtica urens pilulas ferens. *B. pin.* 232.

Urtica romana. *Ger. em.* 706. *Park.* 440.

Urtica pilulifera folio profundius urticæ majoris in
modum serrato, semine magno lini. *R. syn.* 140.

Anglis, roman Nettle.

Habitat in ruderatis; plentifully at Yarmouth in Nor-
folk; Aldborough in Suffolk. *R. syn.* ☉. VIII.

Pharmac. *Urticæ romanæ Semina.*

2. URTICA foliis oppositis ovalibus. *Sp. pl.* 1396. *urens;*

Urtica sexubus sede disjunctis, foliis ovato-lanceolatis
serratis, spicis oblongis. *Hall. hist.* 1615.

Urtica urens minor. *B. pin.* 232.

Urtica minor. *Ger. em.* 707. *R. syn.* 140.

Anglis, lesser Nettle.

Habitat in ruderatis; cultis. ☉. VII—IX.

3. URTICA foliis oppositis cordatis, racemis geminis. *dioica;*
Sp. pl. 1396.

Urtica stirpibus sexu distinctis, foliis ovato-lanceolatis
productis serratis. *Hall. hist.* 1614.

- Urtica. *Cæsalp.* 156. urens. *Ger. em.* 706. maxima
B. pin. 232.
 Urtica racemifera major perennis. *R. syn.* 139.
Anglis, common Nettle.
 Habitat in ruderatis et ad sepes. 4. VII.
Pharmac. Urticæ Semina, Folia, Cymæ.

PENTANDRIA.

XANTHIUM. 1056.

Masc. *Cal.* communis imbricatus. *Cor.* monopetalæ,
 quinquefidæ, infundibuliformes. *Receptaculum* pa-
 leaceum.

Fem. *Cal.* involucreum diphyllum, biflorum. *Cor.* o.
Drupa sicca muricata, bifida. *Nucleus* bilocularis.

struma-
rium.

1. XANTHIUM caule inermi, foliis cordatis trinerva-
 tis. *Sp. pl.* 1400.

Xanthium foliis semitrilobatis, aculeis uncinatis.

Xanthium. *Cam. ep.* 926. f. Lappa minor. *B. hist.*
III. 572. *Park.* 1222.

Lappa minor Xanthium Dioscoridis. *B. pin.* 198.

Bardana minor. *Ger. em.* 809.

Anglis, lesser Xanthium or Burdock.

Habitat in siccis et pinguibus. ☉. VI—IX.

Pharmac. Strumatix Herba.

AMARANTHUS. 1060.

Masc. *Cal.* triphyllus. *Cor.* o. *Stam.* tria f. quinque.

Fem. *Cal.* triphyllus. *Cor.* o. *Styli* tres. *Caps.* uni-
 locularis circumscissa. *Sem.* unicum.

- Blitum.* 1. AMARANTHUS glomerulis lateralibus, floribus
 trifidis, foliis ovatis retusis, caule diffuso. *Sp. pl.*
 1405.

Blitum rubrum minus. *Cam. ep.* 235. *Ger. em.* 321.
B. pin. 118. *R. syn.* 187.

β Blitum album minus. *Cam. ep.* 236.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Millefolium aquaticum pennatum spicatum. *B. pin.*
141. *pr.* 73. *Park.* 1257.

- β Millefolium aquaticum pennatum minus, foliolis singularibus latiusculis, flosculis subjectis donatum.
Hist. Ox. III. 622. *f.* 15. *t.* 4. *f.* 7.
Anglis, spiked Water-millfoil.
Habitat in fossis, lacubus et aquis quietus. 4. VII.

verticillatum.

2. MYRIOPHYLLUM floribus omnibus verticillatis.
Sp. pl. 1410. hermaphroditis. *Syst. nat.* 628.
Pentapteris floribus alaribus. *Hall. hist.* 992.
Myriophyllum aquaticum minus. *Clus. hist. II.* 252.
• *Ger. em.* 827. *Vaill. act. paris.* 1719. 31.
Millefolium aquaticum minus. *Park.* 1257.
Millefolium aquaticum flosculis ad foliorum nodos.
B. pin. 141. *Loef. pr.* 165.
Pentapterophyllum aquaticum flosculis ad foliorum nodos.
Cat. giff. 112. *R. syn.* 316.
Anglis, verticillated Water-millfoil.
Habitat in fossis et aquis stagnantibus. 4. VI. VII.

SAGITTARIA. 1067.

Masc. *Cal.* triphyllus. *Cor.* tripetala. *Filamenta* fere viginti quatuor.

Fem. *Cal.* triphyllus. *Cor.* tripetala. *Pist.* multa.
Sem. multa, nuda.

Sagittifolia.

1. SAGITTARIA foliis sagittatis acutis. *Sp. pl.* 1410.
Fl. dan. 172.
Sagitta foliis acuminatis. *Hall. hist.* 1185.
Sagitta. *Cam. ep.* 814. *B. hist. III.* 789. *R. syn.* 258. minor latifolia. *Park.* 1247.
Sagitta major. *Ger. em.* 416.
Sagitta aquatica minor latifolia. *B. pin.* 194.
β Sagitta aquatica omnium minima. *D. Pluk.* *R. syn.* 258.
Anglis, common Arrow-head.
Habitat in fossis et ad ripas fluviorum. 4. VII.

P O T E R I U M. 1069.

Masc. *Cal.* tetraphyllus. *Cor.* quadripartita. *Stam.*
a triginta ad quadraginta.

Fem. *Cal.* tetraphyllus. *Cor.* quadripartita. *Pist.* duo.
Bacca e tubo corollæ indurato.

1. POTERIUM inerme, caulibus subangulosis. *Sp. pl.* Sanguisorba.

1411. *Hort. ups.* 288. *cliff.* 446.

Pimpinella polystemon. *Hall. hist.* 706.

Sanguisorba minor. *B. hist. III.* 113. *R. syn.* 203.

Pimpinella sanguisorba minor hirsuta. *B. pin.* 160.

Pimpinella vulgaris minor. *Park.* 582.

Anglis, common Burnet.

Habitat in pratis et pascuis montosis, solo calcareo. 4.
VII.

Pharmac. *Pimpinellæ Herba.*

Q U E R C U S. 1070.

Masc. *Cal.* quinquefidus fere. *Cor.* o. *Stam.* a quinque ad decem.

Fem. *Cal.* monophyllus, integerrimus, scaber. *Cor.* o.
Stylus unicus. *Sem.* unicum ovatum.

1. QUERCUS foliis deciduis oblongis superne latioribus: sinibus acutioribus: angulis obtusis. *Robur.*
Sp. pl.

1414.

Quercus foliis glabris semipinnatis, pinnis obtusis, superioribus majoribus. *Hall. hist.* 1626.

Quercus vulgaris. *Ger. em.* 1339.

Quercus latifolia. *Park.* 1385. *R. syn.* 440.

Quercus cum longo pediculo. *B. pin.* 420.

Quercus latifolia mas quæ brevi pediculo est. *B. pin.*
419.

Anglis, common Oak. *Hunt. ev. sylv.* 69.

Habitat in sylvis et sepibus. h. IV.

Pharmac. *Quercus Cortex, Folia, Glandes, Cupulæ.*

MONOECIA POLYANDRIA.

F A G U S. 1072.

Masc. *Cal.* quinquefidus, campanulatus. *Cor.* 0.
Stam. duodecim.

Fem. *Cal.* quadridentatus. *Cor.* 0. *Styli* tres. *Caps.*
(*Calyx* antea) muricata, quadrivalvis. *Sem.* duo.

sylvatica. 1. FAGUS foliis ovatis obsolete ferratis. *Sp. pl.* 1416.
Fagus foliis ovato-lanceolatis, oris undulatis. *Hall.*
hist. 1622.

Fagus. *Cam. ep.* 112. *Ger. em.* 1444. *B. pin.* 419.
Park. 1403. *R. syn.* 439.

Anglis, common Beach-tree. *Hunt. ev. sylv.* 136.

Habitat in sylvis et sepibus solo calcareo. *h.* IV. V.

Castanea. *2. FAGUS foliis lanceolatis acuminato-ferratis subtus
nudis. *Sp. pl.* 1416.

Castanea foliis ovato-lanceolatis, ferratis dentibus
aduncis. *Hall. hist.* 1623.

Castanea. *Cam. ep.* 118. *B. hist. I.* 121. *Ger. em.*
1442. *R. syn.* 440. vulgaris. *Park.* 1400.

Castanea sylvestris. *B. pin.* 419.

Anglis, Chesnut-tree. *Hunt. ev. sylv.* 159.

Habitat in sylvis et sepibus; in Cantio frequens. *h.* V.
Pharmac. Castaneæ Fructus.

C A R P I N U S. 1073.

Masc. *Cal.* monophyllus: squama ciliata. *Cor.* 0.
Stam. viginti.

Fem. *Cal.* monophyllus: squama ciliata. *Cor.* 0.
Germina duo. *Styli* singulis duo. *Nux* ovata.

Betulus. 1. CARPINUS squamis strobilorum planis. *Sp. pl.* 1416.
Carpinus. *Cam. ep.* 71. *Hall. hist.* 1627.

Ostrya ulmo similis, fructu in umbilicis foliaceis. *B.*
pin. 427. *R. syn.* 451.

Ostrys sive Ostrya. *Park.* 1406.

Betulus sive Carpinus. *Ger. em.* 1479,

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

MONOECIA MONODELPHIA,

Abies. *Ger. em.* 1363. *Park.* 1539. conis sursum
spectantibus, s. mas. *R. pin.* 505.

Anglis, yew-leaved Fir.

Habitat in alpidibus Scoticis. h. V. VI.

Pharmac. Abietis Summitates, Resina.

Abies.

*3. PINUS foliis solitariis subulatis mucronatis lævibus
bifariam versis. *Sp. pl.* 1421. *Fl. dan.* 193.

Pinus foliis solitariis, tetragonis mucronatis. *Hall. hist.*
1656.

Abies mas Theophrasti. *R. syn.* 441.

Picea. *Park.* 1538. major. *Ger. em.* 1354.

Picea major, sive *Abies* rubra. *B. pin.* 493.

Anglis, common Fir or Pitch-tree.

Habitat in alpidibus Scoticis. h. V. VI.

Pharmac. Abietis Resina.

Classis XXII.

D I O E C I A

D I A N D R I A.

S A L I X. 1098.

Masc. *Amenti squamæ. Cor. o. Glandula baseos
nectarifera.*

Fem. *Amenti squamæ. Cor. o. Stylus bifidus. Caps.
unilocularis, bivalvis. Sem. papposa.*

• • *Foliis glabris serratis.*

1. SALIX foliis serratis glabris, floribus hermaphroditis *herma-
diandris. Fl. suec. 878. Sp. pl. 1442? phroditica*

Salix latifolia folio splendente. R. syn. 450.

Anglis, shining Willow.

*Habitat in sepibus humidis. About Ashton in Cumber-
land. R. syn. h. III. IV.*

2. SALIX foliis serratis glabris, floribus triandris. *Sp. triandria,
pl. 1442.*

*Salix foliis glabris, elliptico-lanceolatis, serratis, sti-
pulis dentatis, julis gracilibus triandris. Hall, hist.
1637?*

*Salix foliis elliptico-lanceolatis utrinque glabris ap-
pendiculatis. Gmel. sib. I. 155. t. 34. f. 3.*

*Salix folio Amygdalino utrinque aurito corticem ab-
jiciens. R. syn. 448.*

Anglis, smooth Willow.

Habitat in sylvis sepibus et ad ripas fluviorum. h. IV.

*Desc. Truncus arboreus, mediocris. Ramuli virgati,
glabri. Folia alterna, petiolata, lanceolata, serrata,
acuta, nuda, subtus glauca, polycaria et tripolycaria.*

Petiolus

Petiolus *brevis, canaliculatus, nudus, nunc stipulatus nunc exstipulatus*. Stipulæ *duæ laterales, sessiles, subrotundæ dentatæ*. Glandulæ *duæ ad apicem petioli*. Mas. Amentum *pedunculatum, lineare, unciale, villosiusculum*. Filamenta *tria longissima*. Antheræ *subrotundæ flavæ*. Fem. amentum *pedunculatum, lineare, unciale, glabriusculum: pedunculis foliaceis*. *Variat floribus diandris et triandris in eodem Amento.*

pentan-
dia.

3. SALIX foliis ferratis glabris, flosculis pentandris.

Sp. pl. 1442. Fl. lapp. 370. t. 8. f. 3.

Salix foliis glabris ovato-lanceolatis, petiolis glandulosis, floribus hexandris. *Hall. hist. 1639.*

Salix folio laureo, seu lato glabro odorato. *Pb. br.*

R. syn. 449.

Anglis, Sweet Willow.

Habitat *in sylvis et sepibus, in septentrionali parte Angliæ passim; in sylva Coomb-wood dicta in comitatu Surriensi. h. IV.*

vitellina.

4. SALIX foliis ferratis ovatis acutis glabris: ferraturis cartilagineis, petiolis calloso-punctatis. *Sp. pl. 1442.*

Salix sativa lutea, folio crenato. *B. pin. 473.*

Anglis, yellow Willow.

Habitat *in salicetis passim. h. V.*

amygdali-
na.

5. SALIX foliis ferratis glabris lanceolatis petiolatis, stipulis trapeziformibus. *Sp. pl. 1443.*

Salix stipulis ferratis, foliis lanceolatis ferratis glabris. *Hill. hist. 1636.*

Salix folio auriculato splendente flexilis. *R. cat. cantab. syn. 448.*

Salix folio amygdalino utrinque virente aurito. *B. pin. 473.*

Anglis, almond-leaved Willow.

Habitat *in salicetis et ad ripas fluviorum. h. IV. V.*

fragilis.

6. SALIX foliis ferratis glabris ovato-lanceolatis, petiolis dentato-glandulosis. *Sp. pl. 1443.*

Salix fragilis. *B. pin. 474. pr. 159. Park. 1431.*

Salix

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

DIOECIA DIANDRIA.

Salix foliis orbiculatis ferratis glabris, julis paucifloris.
Hall. hist. 1649.

Salix alpina alni rotundo folio. *Bocc. mus.* II. 19.
t. 1. *R. syn.* 448.

Salix saxatilis minima. *B. pin.* 474. *pr.* 159.
Anglis, herbaceous Willow.

Habitat in ascensu montis Snowdon copiose. D. G. She-
rard. *R. syn. et in montibus Westmorlandicis, Ebo-*
racensibus et Scoticis. 4. V.

*** *Foliis glabris integerrimis.*

reticula- 11. SALIX foliis integerrimis glabris ovatis obtusis.
ta. *Sp. pl.* 1446. *Fl. suec.* 888. *dan.* 212.

Salix foliis integris ovatis subtus reticulatis. *Fl. lapp.*
359. t. 8. f. 4. et t. 7. f. 1. 2.

Salix foliis rotundis integerrimis, subtus tomentosis,
juli gracilibus longe petiolatis. *Hall. hist.* 1650.

Salix pumila folio rotundo. *B. hist.* I. 12. *Sch. it.*
alp. 43. 340. *R. syn.* 449.

Anglis, round-leaved Willow.

Habitat in montibus Cambrobritannicis, Eboracensi-
bus, Westmorlandicis passim. 5. V.

rubra. 12. SALIX foliis integerrimis glabris lineari-lanceolatis
acutis. *Fl. angl.* 364.

Salix minime fragilis foliis longissimis utrinque viridi-
bus, non ferratis, *R. syn.* 449.

Anglis, red Willow.

Habitat in salicetis sed rarius. In the Osier-holt between
Maidenhead and Windsor; and near Salisbury. Mr,
Sherard and Mr. Rand. *R. syn.* 5. IV. V.

*** *Foliis integerrimis villosis.*

repens. 13. SALIX foliis integerrimis lanceolatis: subtus villo-
sis nitidis.

Salix foliis integerrimis lanceolatis utrinque subpilosis,
caule repente. *Sp. pl.* 1447.

Salix procumbens, foliis lanceolatis subtus sericeis.
Hall. hist. 1644.

Salix pumila angustifolia inferne lanuginosa. *B. hist.*
I. 212. *R. syn.* 447.

Salix

Salix humilis angustifolia repens. *Park.* 1434.

Salix pumila brevi angustoque folio incano. *B. pin.* 474.

β *Salix alpina pumila rotundifolia repens, inferne subcinerea.* *B. pin.* 474. *R. syn.* 448.

Salix humilis. *Ger. em.* 1391.

γ *Salix foliis crassis elliptico-lanceolatis integerrimis, utrinque sericeis.* *Hall. hist.* 1643.

Salix pumila foliis utrinque candicantibus et lanuginosis. *R. syn.* 447. *t.* 19. *f.* 3.

δ *Salix foliis integerrimis ovatis: subtus villosis nitidis, fusca.* *Sp. pl.* 1447. *Fl. lap.* 364. *t.* 8. *f.* 13.

Salix pumila angustifolia prona parte cinerea. *B. hist.* 1. 213. *R. syn.* 447.

Salix pumila linifolia incana. *B. pin.* 474.

Salix pumila angustifolia recta. *Park.* 1434.

Chamæitea f. *Salix pumila.* *Ger. em.* 1391.

ε *Salix foliis integerrimis lanceolato-linearibus strictis rosmarinifolia sessilibus subtus tomentosis.* *Sp. pl.* 1448. nifolia.

Salix oblongo incano aucto folio. *B. pr.* 159.

Salix humilis repens angustifolia. *Lob. ic. p.* 2. 137.

Salix pumila rhamni secundi Clusii folio. *R. syn.* 447.

Anglis, creeping Willow.

Habitat in ericetis et paludibus turfosis frequens; γ by Sandown castle, near Deal. *R. syn.* 2. V.

*** *Foliis subserratis villosis.*

14. SALIX foliis ellipticis subserratis rugosis, supra *caprea* villosis subtus tomentosis.

Salix foliis ovatis rugosis: subtus tomentosis, undatis superne denticulatis. *Sp. pl.* 1448.

Salix foliis ovatis rugosis, subtus reticulatis tomentosis, julis ovatis, stipulis amplexicaulibus serratis. *Hall. hist.* 1653.

Salix folio ex rotunditate acuminato. *B. pin.* 474. *R. syn.* 449.

Salix caprea latifolia. *Ger. em.* 1390.

β *Salix caprea acuto longoque folio.* *R. syn.* 450.

γ *Salix foliis integerrimis, ovato lanceolatis subtus sericeis, julis ovatis.* *Hall. hist.* 1651.

Salix latifolia rotunda. *B. pin.* 474. *Park.* 1432. *R. syn.* 449.

Salix

- Salix caprea rotundifolia. *Ger. em.* 1390.
- aurita. ♂ Salix foliis integerrimis utrinque villosis obovatis appendiculatis. *Sp. pl.* 1446. *Fl. lap.* 369. t. 8. f. y.
Salix foliis rugosis reticulatis integerrimis, inferne tomentosis, julis ovatis. *Hall. hist.* 1652.
Salix caprea pumila folio subrotundo, subtus incano. *R. syn.* 450.
- Salix folio rotundo minore. *R. syn.* 450.
Anglis, common Willow or Sallow.
Habitat in sylvis et sepibus humidis; ♂, • in ericetis montosis siccioribus. h. IV. V.
- viminalis. 15 SALIX foliis subintegerrimis lanceolato-linearibus longissimis acutis subtus sericeis, ramis virgatis. *Sp. pl.* 1448.
Salix foliis prælongis obscure dentatis, subtus tomentosis et albicantibus. *Hall. hist.* 1641.
Salix folio longissimo. *R. cat. ant.*
Salix angustis et longissimis foliis crispis subtus albicantibus. *B. hist.* I. 212.
Anglis, osier Willow.
Habitat in sylvis, sepibus et salicetis. h. IV. V.
- alba. 16. SALIX foliis lanceolatis acuminatis ferratis utrinque pubescentibus: ferraturis infimis glandulosis, *Sp. pl.* 1449.
Salix arborea foliis ellipticis lanceolatis, subtus sericeis, dentibus crassescantibus. *Hall. hist.* 1635.
Salix. *Ger. em.* 1389. *R. syn.* 447. arborea angustifolia alba vulgaris. 1430.
Salix vulgaris alba arborecens. *B. pin.* 473.
Anglis, white Willow.
Habitat in sylvis et sepibus. h. IV.
Pharmac. Salicis Folia, Cortex.

T R I A N D R I A.

E M P E T R U M. 1100.

Mas. Cal. tripartitus, Cor, tripetala. Stam. longa.
Fem.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

DIOËCIA PENTANDRIA:

Rhamnoides fructifera, foliis falcis, baccis leviter flavescens. *Tourn. Cor.* 55. *R. syn.* 445.

Rhamnus 2. *Clus. hist.* 110. *Ger. em.* 1334.

Rhamnus falcis folio angusto, fructu flavescens. *B. pin.* 477.

Rhamnus primus Dioscoridis Löbelio, sive littoralis. *Park.* 1006.

Anglis, common Swallow-thorn or Sea Buck-thorn.

Habitat in littoribus maritimis arenosis. On the sea-banks on Lindsey-coast, Lincolnshire. About Sandwich, Deal and Folkston in Kent; between Whitby and Lyth, Yorkshire. *R. syn. in insula Sheepey.* h. IV. V.

M Y R I C A. 1107.

Mas. *Amenti squama lunata.* *Cor.* 0.

Fem. *Amenti squama lunata.* *Cor.* 0. *Styli duo.* *Bacca monosperma.*

Gale.

1. MYRICA foliis lanceolatis subserratis, caule suffruticoso. *Sp. pl.* 1453. *Fl. dan.* 327.

Gale frutex odoratus septentrionalium, Elæagnus Cordi. *B. hist.* I. 223. *R. syn.* 443.

Rhus myrtifolia belgica, *B. pin.* 414.

Rhus sylvestris s. Myrtus Brabantica vel Anglica. *Park.* 1451.

Myrtus brabantica s. Elæagnus Cordi. *Ger. em.* 1414. *Anglis*, sweet Gale, Willow or Dutch Myrtle.

Habitat in paludibus, in Cornubia, Devoniam, frequens. h. V.

P E N T A N D R I A.

H U M U L U S. 1116.

Mas. *Cal.* pentaphyllus. *Cor.* 0.

Fem. *Cal.* monophyllus, oblique patens, integer. *Cor.* 0. *Styli duo.* *Sem.* unicum intra calycem foliatum.

1. HUMULUS.

1. HUMULUS. *Sp. pl.* 1457. *Fl. suec.* 908. *Lupulus.*
Lupulus. *Hall. hist.* 1618. mas et femina. *B. hist.*
II. 151. *pin.* 298. *R. syn.* 137.
Lupulus salictarius. *Fuchs. hist.* 124. *Ger. em.* 885.
Lupulus sylvestris. *Park.* 176.
Anglis, Hops.
Habitat in sepibus. 4. VI.
Pharmac. *Lupuli Strobili.*

H E X A N D R I A.

T A M U S. 1119.

- Maf. Cal.* sexpartitus. *Cor.* 0.
Fem. Cal. sexpartitus. *Cor.* 0. *Stylus* trifidus. *Bacca*
trilocularis, infera. *Sem.* duo.
 1. TAMUS foliis cordatis indivisis. *Sp. pl.* 1458. *Hall. communis.*
hist. 1620.
Tamus racemosa, flore minore luteo pallefcente.
Tourn. 103. *R. syn.* 262.
Bryonia nigra. *Ger. em.* 871.
Bryonia sylvestris nigra. *Park.* 178.
Bryonia lævis f. nigra racemosa. *B. pin.* 297.
Anglis, common Tamus or black Bryony.
Habitat in dumetis et ad sepes. 4. VI.
Pharmac. *Bryoniæ nigræ Folia, Radix.*

O C T A N D R I A.

P O P U L U S. 1123.

- Maf. Amenti Cal.* lamina lacera. *Cor.* turbinata, ob-
liqua, integra.
Fem. Amenti Cal. Cor. maris. *Stigma* quadrifidum.
Caps. bilocularis. *Sem.* multa, papposa.
 1. POPULUS foliis subrotundis dentato-angulatis sub-*alba.*
tus tomentosus. *Sp. pl.* 1463. *Hall. hist.* 1634.
 F f Populus

MONOECIA MONODELPHIA,

Abies. *Ger. em.* 1363. *Park.* 1539. conis sursum
spectantibus, s. mas. *R. pin.* 505.

Anglis, yew-leaved Fir.

Habitat in alpidibus Scoticis. h. V. VI.

Pharmac. Abietis Summitates, Resina.

Abies.

*3. PINUS foliis solitariis subulatis mucronatis lævibus
bifariam versis. *Sp. pl.* 1421. *Fl. dan.* 193.

Pinus foliis solitariis, tetragonis mucronatis. *Hall. hist.*
1656.

Abies mas Theophrasti. *R. syn.* 441.

Picea. *Park.* 1538 major. *Ger. em.* 1354.

Picea major, sive *Abies* rubra. *B. pin.* 493.

Anglis, common Fir or Pitch-tree.

Habitat in alpidibus Scoticis. h. V. VI.

Pharmac. Abietis Resina.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Petiolus brevis, canaliculatus, nudus, nunc stipulatus nunc exstipulatus. Stipulæ duæ laterales, sessiles, subrotundæ dentatæ. Glandulæ duæ ad apicem petioli. Mas. Amentum pedunculatum, lineare, unciale, villosiusculum. Filamenta tria longissima. Antheræ subrotundæ flavæ. Fem. amentum pedunculatum, lineare, unciale, glabriusculum: pedunculis foliaceis. Variat floribus diandris et triandris in eodem Amento.

pentan-
dria.

3. SALIX foliis ferratis glabris, flosculis pentandris. *Sp. pl.* 1442. *Fl. lapp.* 370. t. 8. f. 3.

Salix foliis glabris ovato-lanceolatis, petiolis glandulosis, floribus hexandris. *Hall. hist.* 1639.

Salix folio laureo, seu lato glabro odorato. *Ph. b.* *R. syn.* 449.

Anglis, Sweet Willow.

Habitat in sylvis et sepibus, in septentrionali parte Angliæ passim; in sylva Coomb-wood dicta in comitatu Surriensi. *h.* IV.

vitellina.

4. SALIX foliis ferratis ovatis acutis glabris: ferraturis cartilagineis, petiolis calloso-punctatis. *Sp. pl.* 1442.

Salix sativa lutea, folio crenato. *B. pin.* 473.

Anglis, yellow Willow.

Habitat in salicetis passim. *h.* V.

amygdali-
na.

5. SALIX foliis ferratis glabris lanceolatis petiolatis, stipulis trapeziformibus. *Sp. pl.* 1443.

Salix stipulis ferratis, foliis lanceolatis ferratis glabris. *Hall. hist.* 1636.

Salix folio auriculato splendente flexilis. *R. cat. cantab. syn.* 448.

Salix folio amygdalino utrinque virente aurito. *B. pin.* 473.

Anglis, almond-leaved Willow.

Habitat in salicetis et ad ripas fluviorum. *h.* IV. V.

fragilis.

6. SALIX foliis ferratis glabris ovato-lanceolatis, petiolis dentato-glandulosis. *Sp. pl.* 1443.

Salix fragilis. *B. pin.* 474. *pr.* 159. *Park.* 1431.

Salix

Salix folio longo latoque splendente fragilis. R. syn.
448.

Anglis, crack Willow.

Habitat in sylvis, sepibus et ad ripas fluviorum. h.
IV. V.

7. **SALIX** foliis ferratis glabris lanceolatis : inferioribus *purpurea*.
oppositis. *Sp. pl.* 1444. *Fl. succ.* 884.

Salix vulgaris nigricans folio non ferrato. B. pin. 473.

Salix folio longo subluteo non auriculato viminibus
rubris. R. syn. 450.

Anglis, purple Willow.

Habitat in salicetis. h. IV, V.

8. **SALIX** foliis ferratis glabris lanceolato-linearibus : *Helix*.
superioribus oppositis obliquis. *Sp. pl.* 1444.

Salix monandra, foliis glabris lineari-lanceolatis fer-
ratis, superne conjugatis, julis tomentosis. Hall.
hist. 1640.

Salix humilior, foliis angustis subcæruleis ex adverso
binis. R. cat. cant. 144. *Syn.* 448.

Salix tenuior folio minore utrinque glabro, fragilis.
B. hist. I. 213. *f.* 2.

Anglis, rose Willow.

Habitat in sepibus et aquosis. h. IV. V.

9. **SALIX** foliis ferratis glabris ovatis venosis. *Fl. lapp. myrsinites.*
353. *t.* 8. *f.* *F. et t.* 7. *f.* 6. *Sp. pl.* 1445.

Salix foliis ovato-lanceolatis venosis ferratis, julis to-
mentosis. Hall. hist. 1645.

Salix alpina pumilla myrsinites. Pont. comp. 146.

Salix alpina pumilla repens, foliis oblongis exiguis :
superne splendentibus, inferne cinereis creberrimis
et tenuissimis crenis. Till. hort. piss. 151.

Anglis, mountain Willow.

Habitat in montibus Scoticis, in Breadalbane. D. Mil-
lar. h. V. VI.

10. **SALIX** foliis ferratis glabris orbiculatis. *Sp. pl. herbacea.*
1445. *Fl. lapp.* 355. *t.* 8. *f.* *H. et t.* 7. *f.* 3. 4. *Dan.*
212.

Salix

Salix foliis orbiculatis ferratis glabris, julis paucifloris.

Hall. hist. 1649.

Salix alpina alni rotundo folio. *Bocc. mus.* II. 19.

t. 1. *R. syn.* 448.

Salix saxatilis minima. *B. pin.* 474. *pr.* 159.

Anglis, herbaceous Willow.

Habitat in ascensu montis Snowdon copiose. D. G. Sherard. *R. syn. et in montibus Westmorlandicis, Eboracensibus et Scoticis.* 4. V.

*** *Foliis glabris integerrimis.*

reticulata. 11. SALIX foliis integerrimis glabris ovatis obtusis.
Sp. pl. 1446. *Fl. suec.* 888. *dan.* 212.

Salix foliis integris ovatis subtus reticulatis. *Fl. lapp.* 359. t. 8. f. 4. et t. 7. f. 1. 2.

Salix foliis rotundis integerrimis, subtus tomentosis, juli gracilibus longe petiolatis. *Hall. hist.* 1650.

Salix pumila folio rotundo. *B. hist.* I. 12. *Sch. it. alp.* 43. 340. *R. syn.* 449.

Anglis, round-leaved Willow.

Habitat in montibus Cambrobritannicis, Eboracensibus, Westmorlandicis passim. 6. V.

rubra. 12. SALIX foliis integerrimis glabris lineari-lanceolatis acutis. *Fl. angl.* 364.

Salix minime fragilis foliis longissimis utrinque viridibus, non ferratis, *R. syn.* 449.

Anglis, red Willow.

Habitat in salicetis sed rarius. In the Osier-holt between Maidenhead and Windsor; and near Salisbury. Mr. Sherard and Mr. Rand. *R. syn.* 2. IV. V.

*** *Foliis integerrimis villosis.*

repens. 13. SALIX foliis integerrimis lanceolatis: subtus villosis nitidis.

Salix foliis integerrimis lanceolatis utrinque subpilosis, caule repente. *Sp. pl.* 1447.

Salix procumbens, foliis lanceolatis subtus sericeis. *Hall. hist.* 1644.

Salix pumila angustifolia inferne lanuginosa. *B. hist.* I. 212. *R. syn.* 447.

Salix

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

- aurita. δ Salix caprea rotundifolia. *Ger. em.* 1390.
 Salix foliis integerrimis utrinque villosis obovatis appendiculatis. *Sp. pl.* 1446. *Fl. lap.* 369. t. 8. f. y.
 Salix foliis rugosis reticulatis integerrimis, inferne tomentosis, julis ovatis. *Hall. hist.* 1652.
 Salix caprea pumila folio subrotundo, subtus incano. *R. syn.* 450.
- \bullet Salix folio rotundo minore. *R. syn.* 450.
Anglis, common Willow or Sallow.
 Habitat in sylvis et sepibus humidis; δ , \bullet in ericetis montosis succioribus. *h.* IV. V.
- viminalis. 15 SALIX foliis subintegerrimis lanceolato-linearibus longissimis acutis subtus sericeis, ramis virgatis. *Sp. pl.* 1448.
 Salix foliis prælongis obscure dentatis, subtus tomentosis et albicantibus. *Hall. hist.* 1641.
 Salix folio longissimo. *R. cat. cant.*
 Salix angustis et longissimis foliis crispis subtus albicantibus. *B. hist.* I. 212.
Anglis, osier Willow.
 Habitat in sylvis, sepibus et salicetis. *h.* IV. V.
- alba. 16. SALIX foliis lanceolatis acuminatis ferratis utrinque pubescentibus: ferraturis infimis glandulosis, *Sp. pl.* 1449.
 Salix arborea foliis ellipticis lanceolatis, subtus sericeis, dentibus crassescens. *Hall. hist.* 1635.
 Salix. *Ger. em.* 1389. *R. syn.* 447. arborea angustifolia alba vulgaris. 1430.
 Salix vulgaris alba arborecens. *B. pin.* 473.
Anglis, white Willow.
 Habitat in sylvis et sepibus. *h.* IV.
Pharmac. Salicis Folia, Cortex.

T R I A N D R I A.

E M P E T R U M. 1100.

Mas. Cal. tripartitus, Cor. tripetala. Stam. longa.
 Fem.

DIOECIA TETRANDRIA.

431

Fem. *Cal.* tripartitus. *Cor.* tripetala. *Styli* novem.
Sem. 9-sperma.

1. EMPETRUM procumbens. *Sp. pl.* 1450. *nigrum.*
Empetrum procumbens, foliis ovato-lanceolatis obtu-
sis. *Hall. hist.* 1605.
Empetrum montanum, fructu nigro. *Tourn.* 579.
R. syn. 444.
Erica baccifera. *Cam. ep.* 77. procumbens. *Ger. em.*
1383. nigra. *B. pin.* 486.
Erica baccifera nigra. *Park.* 1485.
Anglis, black Crow, Crake or Heath-berry.
Habitat in ericetis montosis in boreali parte Angliæ fre-
quens. *h.* IV. V.

T E T R A N D R I A .

V I S C U M . 1105.

Mas. *Cal.* quadripartitus. *Cor.* 0. *Filamenta* 0. *An-*
theræ calyci adnatæ.

Fem. *Cal.* tetraphyllus, superus. *Stylus* 0. *Cor.* 0.
Bacca monosperma.

1. VISCUM foliis lanceolatis obtusis, caule dichotomo, *album.*
spicis axillaribus. *Sp. pl.* 1451. *Hall. hist.* 1609.
Viscum. *Cam. ep.* 555. *Ger. em.* 1350. *R. syn.* 464.
vulgare. *Park.* 1392. baccis albis. *B. pin.* 423.
Anglis, white Mistletoe.
Habitat in arboribus; frutex parasiticus. *h.* V.
Pharmac. Visci Folia.

H I P P O P H A E . 1106.

Mas. *Cal.* bipartitus. *Cor.* 0.

Fem. *Cal.* bifidus. *Cor.* 0. *Stylus* unicus. *Bacca*
monosperma.

1. HIPPOPHAE foliis lanceolatis. *Sp. pl.* 1452. *Fl. Rhamnoi-*
dan. 265. *des.*
Hippophae foliis linearibus, subtus rubiginosis. *Hall.*
hist. 1603.
Rhamnoides

DIOECIA PENTANDRIA:

Rhamnoides fructifera, foliis salicis, baccis leviter flavescens. *Tourn. Cor.* 55. *R. syn.* 445.

Rhamnus 2. *Clus. hist.* 110. *Ger. em.* 1334.

Rhamnus salicis folio angusto, fructu flavescens. *B. pin.* 477.

Rhamnus primus Dioscoridis Lobelio, sive littoralis. *Park.* 1006.

Anglis, common Swallow-thorn or Sea Buck-thorn.

Habitat in littoribus maritimis arenosis. On the sea-banks on Lindsey-coast, Lincolnshire. About Sandwich, Deal and Folkston in Kent; between Whitby and Lyth, Yorkshire. *R. syn. in insula Sheepey.* h. IV. V.

M Y R I C A. 1107.

Mas. *Amenti* squama lunata. *Cor.* 0.

Fem. *Amenti* squama lunata. *Cor.* 0. *Styli* duo. *Bacca* monosperma.

Gale.

1. MYRICA foliis lanceolatis subserratis, caule suffruticoso. *Sp. pl.* 1453. *Fl. dan.* 327.

Gale frutex odoratus septentrionalium, *Elæagnus Cordi.* *B. hist.* I. 223. *R. syn.* 443.

Rhus myrtifolia belgica, *B. pin.* 414.

Rhus sylvestris s. *Myrtus Brabantica* vel *Anglica.* *Park.* 1451.

Myrtus brabantica s. *Elæagnus Cordi.* *Ger. em.* 1414.

Anglis, sweet Gale, Willow or Dutch Myrtle.

Habitat in paludibus, in Cornubia, Devoniam, frequens. h. V.

P E N T A N D R I A.

H U M U L U S. 1116.

Mas. *Cal.* pentaphyllus. *Cor.* 0.

Fem. *Cal.* monophyllus, oblique patens, integer. *Cor.* 0. *Styli* duo. *Sem.* unicum intra calycem foliatum.

1. HUMULUS.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Populus alba. *Cam. ep.* 61. *Ger. em.* 1486. *Park.* 1410. *R. syn.* 446. majoribus foliis. *B. pin.* 429. *Anglis*, white Poplar. *Hunt. Ev. sylv.* 208. *Habitat in sepibus et sylvis.* h. III.

tremula. 2. POPULUS foliis subrotundis, dentato-angulatis utrinque glabris. *Sp. pl.* 1464.

Populus foliis glabris orbiculatis acuminatis laxe serratis. *Hall. hist.* 1633.

Populus lybica. *Plin. lib. XVI. c. 23.* *Cam. ep.* 67. *Ger. em.* 1487. *Park.* 1411. *R. syn.* 446.

Populus tremula. *B. pin.* 429.

Anglis, trembling Poplar or Asp.

Habitat in sylvis humidis. h. III.

nigra. 3. POPULUS foliis deltoidibus acuminatis serratis. *Sp. pl.* 1464.

Populus foliis glabris cordato-rhomboides serratis. *Hall. hist.* 1632.

Populus nigra. *Plin. Lib. XVI. c. 23.* *Cam. ep.* 66. *Ger. em.* 1486. *B. pin.* 429. *Park.* 1410. *R. syn.* 446.

Anglis, black Poplar.

Habitat in umbrosis humidis et ad fluvios. h. III.

Pharmac. Populi nigri Folia, Germinæ.

R H O D I O L A. 1124.

Maf. *Cal.* quadripartitus. *Cor.* quadripetala.

Fem. *Cal.* quadripartitus. *Cor.* o. Nectaria quatuor.

Pist. quatuor. *Caps.* quatuor, polysperma.

Rosea. 1. RHODIOLA. *Sp. pl.* 1464. *Fl. lapp.* 378. *dan.* 181.

Sedum sexu distinctum, foliis serratis, floribus densissime umbellatis. *Hall. hist.* 953.

Anacampteros radice rosam spirante major. *Tourn.* 264. *R. syn.* 260.

Rhodia radix. *B. pin.* 286. *hist.* III. 683. *Ger. em.* 532. *Park.* 729.

Anglis, Rosewort.

Habitat

Habitat in montibus Westmorlandicis, Cumberlandicis, Eboracensibus et Cambrobritannicis frequens. 4. VI. VII.

ENNEANDRIA.

MERCURIALIS. 1125.

Mas. Cal. tripartitus. Cor. o. Stam. a novem ad duodecim. Antheræ globosæ, didymæ.

Fem. Cal. tripartitus. Cor. o. Styli duo. Caps. dicocca, bilocularis, monosperma.

1. MERCURIALIS caule simplicissimo, foliis scabris. *perennis.*

Sp. pl. 1465. *Fl. dan.* 400.

Mercurialis caule perenni simplici, foliis ovato-lanceolatis hirsutis. *Hall. hist.* 1601.

Mercurialis perennis repens, Cynocrambe dicta. *R. syn.* 138.

Mercurialis montana spicata et mercurialis montana testiculata. *B. pin.* 122.

Mercurialis sylvestris Cynocrambe dicta vulgaris mas et femina. *Park.* 295.

Cynocrambe mas et femina. *Cam.* 998. *Ger. em.* 333. *Anglis*, dog's Mercury.

Habitat in sylvis et ad sepes. 4. IV. V.

2. MERCURIALIS caule brachiato, foliis glabris, floribus spicatis. *Sp. pl.* 1465. *annua.*

Mercurialis caule annuo brachiato, foliis conjugatis ovato-lanceolatis glabris. *Hall. hist.* 1600.

Mercurialis annua glabra vulgaris. *R. syn.* 139.

Mercurialis spicata s. femina. *B. pin.* 121.

Mercurialis testiculata s. mas. *B. pin.* 121.

Mercurialis mas et femina. *Ger. em.* 332.

Mercurialis vulgaris mas et femina. *Park.* 295.

Anglis, annual or french Mercury.

Habitat in ruderatis, cultis, circa Londinum frequens.

©. VIII. IX.

DIOECIA MONADELPHIA.

HYDROCHARIS. 1126.

Maf. *Spatha* diphylla. *Cal.* trifidus. *Cor.* tripetala.
Filamenta tria interiora stylifera.

Fem. *Cal.* trifidus. *Cor.* tripetala. *Styli* sex. *Caps.*
sex locularis, polysperma, infera.

Morsus
ranæ.

1. HYDROCHARIS. *Sp. pl.* 1466. *Hall. hist.* 1068.

Nymphæa alba minima. *B. pin.* 193. f. *Morsus* ranæ.

Park. 1252.

Morsus ranæ. *Ger. em.* 818.

Stratiotes foliis asari, semine rotundo. *R. syn.* 290.

Anglis, Frog bit.

Habitat in fossis et aquis leniter fluentibus. 4. VI.

MONADELPHIA.

JUNIPERUS. 1134.

Maf. *Amenti Calyx* squamæ. *Cor.* 0. *Stamina* tria.

Fem. *Cal.* tripartitus. *Petala* tria. *Styli* tres. *Bacca*
trisperma, tribus tuberculis calycis inæqualibus.

communis. 1. JUNIPERUS foliis ternis patentibus mucronatis bac-
ca longioribus. *Sp. pl.* 1470.

Juniperus foliis convexo-concavis aristatis, baccis ala-
ribus. *Hall. hist.* 1662.

Juniperus. *Cam. ep.* 55. *Ger. em.* 1372. vulgaris.

Park. 1028. baccis parvis purpureis. *B. hist.* I.

293. *R. spn.* 440.

Juniperus vulgaris fruticosa. *B. pin.* 488.

β *Juniperus* alpina. *B. hist.* I. 301.

Anglis, common Juniper.

Habitat in ericetis et montosis. 5. V.

Pharmac. *Juniperi* *Baccæ*, *Lignum*, *Resina*.

TAXUS. 1135.

Maf. *Cal.* triphyllus gemmæ. *Cor.* 0. *Stamina* multa.

Antheræ peltatæ octofidæ.

Fem.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

DIOECIA SYNGENESIA.

Bryonia alba. *Dod. purg.* 234. *Ger. em.* 869. *R.*
syn. 261. *vulgaris.* *Park.* 178.

Bryonia aspera s. *alba*, *baccis rubris.* *B. pin.* 297,
Anglis, white Bryony.

Habitat in sepibus et dumetis. 4. VI.

Pharmac. *Bryoniæ albæ Radix.*

Classis XXIII.

P O L Y G A M I A

M O N O E C I A.

CALLITRICHE. 13.

Masculus flos.

CAL. nullus.

COR. *Petala* duo, opposita, oblonga, concava, conniventia.STAM. *Filamentum* unicum, longissimum rectum.*Anthera* subrotunda, simplex.

PIST. PER. SEM. nulla.

Femineus flos in eadem planta.

CAL. COR. ut in mare.

PIST. *Germen* superum compressum. *Styli* duo, filiformes longissimi.PER. *Capsula* obcordata, compressa, bilocularis quadrivalvis.

SEM. solitaria, oblonga.

Hermaphroditus flos in eadem planta.

CAL. COR. STAM. ut in mare.

PIST. PER. SEM. ut in femineo.

1. CALLITRICHE foliis emarginatis.

Callitriche foliis superioribus ovalibus : floribus androgynis. *Sp. pl.* 6. *Fl. dan.* 129. *aquatica.*
*verna.**Stellaria* foliis imis linearibus, superioribus subrotundis. *Hall. hist.* 554.*Stellaria.* *Dill. giff.* 119. t. 6. *R. syn.* 289. *aquatica.* *Lob. ic.* 792. *Park.* 1282. *B. pin.* 141. *Ger. em.* 830.2. Callitriche Plinii. *Col. ecph.* 315.

Stellaria foliis petiolatis subrotundis. *Hall. hist.* 553.

Stellaria minor et repens. *R. syn.* 289.

autum-
nalis.

γ *Callitriche* foliis omnibus linearibus apice bifidis, floribus hermaphroditis. *Sp. pl.* 6.

Stellaria foliis omnibus linearibus. *Hall. hist.* 555.

Stellaria aquatica longifolia. *R. syn.* 290.

Lenticula aquatica angustifolia, folio in apice dissecto.

Læs. pr. 140. t. 38.

• *Anglis*, common Starwort.

Habitat in fossis et aquis quietus, β in aquis fluentibus.

⊙. III—X.

H O L C U S. 1146.

Hermaphroditus. *Calyx* Gluma biflora. *Cor.* Gluma aristata. *Stam.* tria. *Styli* duo, *Sem.* unicum.

Masc. *Cal.* Gluma bivalvis. *Cor.* o. *Stam.* tria.

lanatus.

i. **HOLCUS** glumis bifloris villosis: hermaphrodito mutico, masculo arista recurva. *Sp. pl.* 1485. *Schr. gr.* 145. t. 20. f. 1.

Avena diantha, floribus ovatis, perfecto mutico, imperfecto aristato. *Hall. hist.* 1484.

Gramen pratense paniculatum molle. *B. pin.* 2. pr. 3. th. 37. *Park.* 1555. *Sch. agr.* 234.

Gramen miliaceum paniculatum molle. *Pet. conc. gr.* 124. *R. syn.* 404.

Anglis, meadow Soft-grass.

Habitat in pratis et pascuis. 4. VI. VII.

mollis.

2. **HOLCUS** glumis bifloris nudiusculis: flosculo hermaphrodito mutico, masculo arista geniculata. *Sp. pl.* 1485. *Schr. gram.* 149. t. 20. f. 2.

Gramen caninum longius radicum. *B. pin.* 1.

Gramen paniculatum molle, radice graminis canini repente. *Hist. Qx.* III. 202.

Gramen miliaceum aristatum molle. *Pet. conc. gr.* 125. *R. syn.* 401.

Gramen caninum paniculatum molle. *R. hist.* 1285. *Sch. agr.* 235. t. 4. f. 25.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

PARIETARIA. 1152.

Hermaphrod. *Cal.* quadrifidus. *Cor.* o. *Stam.* quatuor. *Stylus* unicus. *Sem.* unicum, superum, elongatum.

Fem. *Cal.* quadrifidus. *Cor.* o. *Stam.* o. *Stylus* unicus. *Sem.* unicum, superum, elongatum.

officinalis. 1. PARIETARIA foliis lanceolato-ovatis, pedunculis dichotomis, calycibus diphyllis. *Syst. nat.* 672. *Sp. pl.* 1492.

Parietaria foliis hirsutis, elliptico-lanceolatis. *Hall. hist.* 1612.

Parietaria. *Ger. em.* 331. *B. hist. II.* 976. *R. syn.* 158. vulgaris. *Park.* 437.

Parietaria officinarum et Dioscoridis. *B. pin.* 121, *Anglis*, common Pellitory of the Wall.

Habitat in ruderatis et muris. 4. V—IX, *Pharmac.* Parietarizæ Herba.

ATRIPLEX. 1153.

Hermaphrod. *Cal.* pentaphyllus. *Cor.* o. *Stam.* quinque. *Stylus* bipartitus. *Sem.* unicum, depressum.

Fem. *Cal.* diphyllus. *Cor.* o. *Stylus* bipartitus. *Sem.* unicum compressum.

portulacoides. 1. ATRIPLEX caule fruticoso, foliis obovatis. *Sp. pl.* 1493.

Atriplex maritima fruticosa, Halimus et Portulaca marina dicta, angustifolia. *R. syn.* 153.

Halimus seu portulaca marina. *B. pin.* 123.

Halimus vulgaris s. portulaca marina. *Ger. em.* 523.

Portulaca marina nostras. *Park.* 724.

Anglis, shrubby Orache or sea Purslane.

Habitat in littoribus maritimis, solo lutofo. 2. VII. VIII.

laciniata. 2. ATRIPLEX caule herbaceo, foliis deltoidibus sinuato-dentatis undulatis subtus argenteis.

Atriplex

Atriplex caule herbaceo, foliis deltooidibus dentatis subtus argenteis. *Sp. pl.* 1494.

Atriplex maritima. *B. hist.* II. 979. *R. syn.* 152. laciniata. *B. pin.* 120.

Atriplex marina. *Ger. em.* 324. repens. *Park.* 758.

β Atriplex caule herbaceo, foliis deltooidibus sinuato-tartarica, dentatis undulatis alternis. *Sp. pl.* 1493.

Atriplex maritima nostras procerior, foliis angulosis incanis, admodum sinuatis. *Pluk. alm.* 60.

Anglis, sea Orache.

Habitat in littoribus maritimis. ☉. VII. VIII.

Obs. α, Caulis, bipedalis, erectiusculus, ramosus; ramis patulis; β. Caulis tri et quinquepedalis, erectus, ramosus; ramis erectis. Folia in α et β, deltoidea, farinosa, argentea; Calyx fructus, magnus, deltoideus. compressus, basi utriusque bidentatus. α, Sata in horto in β mutatur.

3. ATRIPLEX caule herbaceo, calycis valvulis femineis hastata, magnis deltooidibus sinuatis. *Sp. pl.* 1494.

Atriplex foliis triangularibus, basi producta, valvulis triangularibus subasperis. *Hall. hist.* 1617.

Atriplex sylvestris, foliis hastato s. deltoide. *R. syn.* 151.

Atriplex sylvestris annua, folio deltoide triangulari sinuato et mucronato, hastæ cuspidi simili. *Hist. Ox.* II. 607. f. 5. t. 32. f. 14.

β Atriplex maritima perennis, folio deltoide seu triangulari minus incano. *Hist. Ox.* II. 607.

γ Atriplex maritima ad foliorum basin auriculata procumbens et ne vix sinuato. *Pluk. alm.* 61.

Anglis, wild Orache.

Habitat in ruderatis, fimetis et littoribus maritimis. ☉. VIII. IX.

Obs. variat caule erecto et decumbente, foliis deltooidibus sinuato-dentatis, dentatis et integris.

4. ATRIPLEX caule herbaceo patulo, foliis subdeltoideo-lanceolatis, calycibus feminum disco-dentatis.

Sp. pl. 1494.

Atriplex caule herbaceo, foliis superioribus lanceolatis integerrimis. *Hort. cliff.* 469. *Roy. lugdb.* 218.

Atriplex

Atriplex foliis imis hamatis, reliquis lingulatis, valvulis feminalibus dentatis. *Hall. hist.* 1616.

Atriplex sylvestris angustifolia. *Ger. em.* 326. *Park.* 748. *R. syn.* 151.

Atriplex angusto oblongo folio. *B. pin.* 119.

β Atriplex caule herbaceo, foliis hastatis dentatis. *Fl. angl.* 376.

Atriplex angustifolia laciniata. *R. syn.* 152.

Anglis, spreading Orache.

Habitat in ruderatis et ad sepes. ☉. VIII.

serrata. 5. ATRIPLEX caule herbaceo erectiusculo, foliis linearibus serratis. *Fl. angl.* 377.

Atriplex angustifolia maritima dentata. *R. syn.* 152.

Atriplex maritima angustifolia. *B. pin.* 152.

Anglis, serrated Orache. *Pet. h. br. t.* 7. f. 4.

Habitat in ruderatis et littoribus maritimis. ☉. VIII.

littoralis. 6. ATRIPLEX caule herbaceo erecto, foliis omnibus linearibus integerrimis. *Sp. pl.* 1494.

Atriplex angustissimo et longissimo folio. *Herm. lugd.* 79. *Tourn.* 505. *R. syn.* 153.

β Atriplex maritima scopariæ folio. *R. syn.* 153.

γ Atriplex maritima angustifolia, obtusiore folio. *R. syn.* 153.

Anglis, grass Orache. *Pet. h. br. t.* 7. f. 6.

Habitat in ruderatis et littoribus maritimis. ☉. VIII.

pedunculata. 7. ATRIPLEX caule herbaceo divaricato, foliis lanceolatis obtusis integris, calycibus femineis pedunculatis. *Sp. pl.* 1675.

Atriplex marina semine lato. *Ph. br.* *R. syn.* 153.

Atriplex maritima Halimus dicta, humilis erecta, semine folliculis membranaceis bivalvibus in latitudinem expansis et utrinque recurvis, longo pediculo insidentibus, clauso. *Pluk. ph. t.* 36. f. 1.

β Atriplex maritima nostras Ocymi minoris folio. *D. Pluk.* *R. syn.* 253.

Anglis, pedunculated Orache.

Habitat in littoribus maritimis; prope Schirbeck vicum milliare a Bostoniam distantem, copiose provenit. *D. Pluknet.*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Fem. *Pist.* unicum. *Sem.* unicum, lanceolatum.

- excelsior.* 1. FRAXINUS foliis ferratis, floribus apetalis. *Sp. pl.* 1509. *Fl. suec.* 926.
 Fraxinus apetalâ, foliis pinnatis ferratis. *Hall. hist.* 528.
 Fraxinus. *Dod. pempt.* 771. *Cam. ep.* 64. *Ger. em.* 1472. *R. syn.* 469. *excelsior.* *B. pin.* 416.
Anglis, common Ash-tree. *Hunt. Ev. sylv.* 150.
 Habitat in sylvis et sepibus. *h.* III. IV.
 Obs. variat foliis pinnatis et simplicibus.

I L E X.

Hermaphrod. *Cal.* quinquedentatus. *Cor.* quinquepartita, rotata. *Stam.* quinque, rotata. *Stylus* nullus. *Stigmata* quatuor. *Bacca* tetrasperma.

Masc. *Cal.* quadridentatus. *Cor.* quadripartita. *Stam.* quatuor.

- aquifolium.* 1. ILEX foliis ovatis acutis spinosis. *Sp. pl.* 181.
 Ilex aculeata baccifera. *B. pin.* 425.
 Aquifolium. *Cam. ep.* 84. *Hall. hist.* 667.
 Agrifolium. *Ger. em.* 1338. s. Aquifolium. *Part.* 1486. *R. syn.* 466.
Anglis, common Holly. *Hunt. Ev. sylv.* 383.
 Habitat in sylvis, sepibus et ericetis. *h.* V.

Classis XXIV.

CRYPTOGAMIA

FILICES.

EQUISETUM. 1169.

Spica fructificationibus peltatis, basi dehiscentibus multivalvi.

1. EQUISETUM caule spicato, frondibus compositis. *sylvaticum.*
Sp. pl. 1516.

Equisetum foliis repetito ramosis, vaginis levissimis.
Hall. hist. 1680.

Equisetum sylvaticum. *Tab. ic. 253. Ger. em. 1114.*
R. syn. 130. tenuissimis setis. B. pin. 16.

Equisetum omnium minus tenuifolius. *Park. 1201.*

β Equisetum sylvaticum, procumbens, setis uno versu dispositis, *R. syn. 131.*

γ Equisetum palustre tenuissimis et longissimis setis.
B. pin. 16. pr. 24.

Anglis, wood Horse-tail.

Habitat in sylvis et umbrosis humidis. 4. IV. V.

2. EQUISETUM scapo fructificante nudo: sterili *arvense.*
frondoso. *Sp. pl. 1516.*

Equisetum caule florifero nudo, sterili verticillato, radiorum duodecim. *Hall. hist. 1676.*

Equisetum arvense longioribus setis. *B. pin. 16. Park., 1202. R. syn. 130.*

Equisetum segetale. *Ger. em. 1114.*

β Equisetum pratense longissimis setis. *B. pin. 16.*

γ Equisetum nudum minus variegatum basiliense. *B. pin. 16. pr. 25. R. syn. 130.*

Anglis,

Anglis, corn Horse-tail.

Habitat in arvis humidis. 4. III. IV.

- palustre.* 3. EQUSETUM caule angulato, frondibus simplicibus. *Sp. pl.* 1516. *Fl. suec.* 928.
 Equisetum caule sulcato, ramis multifloris, foliis indivisis. *Hall. hist.* 1677.
 Equisetum palustre brevioribus setis. *B. pin.* 16.
 Equisetum palustre. *Lob. ic.* 795. *Ger. em.* 1114.
R. syn. 131. minus. *Park.* 1200.
 β Equisetum palustre minus polytacion. *B. pin.* 16.
pr. 24. *R. syn.* 131. t. 5. f. 3.
Anglis, marsh Horse-tail.
 Habitat in palustribus et aquosis. 4. VI. VII.

- fluviale.* 4. EQUSETUM caule striato, frondibus subsimplicibus. *Sp. pl.* 1517.
 Equisetum caule florifero nudo, sterili verticillato, radiorum quadraginta. *Hall. hist.* 1675.
 Equisetum. *Cam. epit.* 770. majus. *Ger. em.* 1113.
R. syn. 130. palustre. *Park.* 1290.
 Equisetum palustre longioribus setis. *B. pin.* 15.
Anglis, river Horse-tail.
 Habitat in palustribus et aquosis et ad ripas fluviorum et lacuum.

- limosum.* 5. EQUSETUM caule subnudo lævi. *Sp. pl.* 1517.
 Equisetum nudum lævius nostras. *R. syn.* 131. t. 5. f. 6.
Anglis, smooth Horse-tail.
 Habitat in aquosis. 4. VI. VII.

- hyemale.* 6. EQUSETUM caule nudo scabro basi subramoso. *Sp. pl.* 1517.
 Equisetum caule subnudo asperrimo, vaginis caulinis indivisis, ramis foliatis. *Hall. hist.* 1679.
 Equisetum nudum. *Ger. em.* 1113. *R. syn.* 131.
 Equisetum junceum ramosum. *Park.* 1207.
 Equisetum foliis nudum ramosum. *B. pin.* 16.
 β Equisetum foliis nudum non ramosum s. junceum. *B. pin.* 16.

Equisetum

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

CRYPTOGAMIA FILICES.

Filix florida s. Osmunda regalis. *Ger. em.* 1131.
Anglis, common Osmund-royal, or flowering fern.
Habitat in aquosis et palustribus putridis. 4. VIII.

*** *Frondebis aliis foliaceis. aliis fructificantibus.*

Spicant. 3. OSMUNDA frondibus lanceolatis pinnatifidis : laciniis confluentibus integerrimis parallelis. *Sp. pl.* 1522. *Fl. suec.* 936. *dan.* 217.

Struthiopteris. *Hall. hist.* 1687.

Lonchitis aspera. *Ger. em.* 1140. *R. syn.* 118. minor.
Cam. ep. 665. *B. pin.* 359. *Park.* 1042.

Anglis, spleenwort Osmund-royal.

Habitat in nemorosis et ericetis humidis. 4. VIII.

crispa. 4. OSMUNDA frondibus supradecompositis : pinnis alternis subrotundis incis. *Sp. pl.* 1522. *Fl. dan.* 496.

Filix foliis triplicato-pinnatis, sterilibus obtusis dentatis, floralibus ovatis integerrimis. *Hall. hist.* 1689.

Adiantum album crispum alpinum. *Schwenkf. Cat.* 10. *R. syn.* 126.

Adiantum album floridum s. *Filicula petræa crispa per-elegans.* *R. syn. ed. II.* 51. *Pluk. ph. t.* 9. *f.* 3.

Anglis, curl'd Osmund-royal, or stone-fern.

Habitat in petris et rupibus. *In comitatibus Eboracensi, Westmorlandico et Wallia frequens.* 4. VIII.

ACROSTICHUM. 1173.

Fructificationes discum totum frondis tegentes.

* *Fronde simplici divisa.*

septentri-onaie. 1. ACROSTICHUM frondibus nudis linearibus laciniatis. *Sp. pl.* 1524. *Fl. dan.* 60.

Acrostichum caule bis bifurcato. *Hall. hist.* 1714.

Filix saxatilis Tragi. *Lob. ic.* 47. *B. hist.* III. 755.
Park. 1045. *R. syn.* 120.

Filix saxatilis corniculata. *B. pin.* 358.

Anglis, forked or horned Acrostichum.

Habitat in fissuris rupium. *Ad cacumen montis Carnedh-Lhewelyn prope Lhan-Lhechyd in agro Arvonensi invenit,*

invenit D. Lhwyd. in muris antiquis Lhan Dethylæ undè circiter milliari a Lhan-Rhoost aquilonem versus D. Richardson. R. syn. in monte Ingleborough, D. Tofield, in montibus supra Ambleside in agro Westmorlandico. 4. VII.

** *Fronde subbipinnata.*

2. ACROSTICHUM? frondibus subbipinnatis; pinnis *ilvense.* opposito-coadunatis obtusis subtus hirsutis: basi integerrimis. *Sp. pl.* 1528.

Filix alpina pedicularis rubræ foliis subtus villosis. D. Lhwyd. R. syn. 118.

Filix pumila lonchitidis maranthæ species cambrobri-tannica. *Pluk. ph. t.* 89. f. 8.

Filicula alpina tenerior, alis latiusculis brevioribus integris profunde dentatis. *Hist. Ox. III.* 576. t. 3. f. 23.

Anglis, hairy Acrostichum.

Habitat in fissuris rupium. It grows on a moist black rock, almost at the top of Clogwyn y Garnedd, facing North-west, directly above the lower lake. D. Richardson, R. syn. in eodem loco inveni. 4. VII—IX,

P T E R I S. 1174.

Fruclificationes in lineis marginalibus.

1. PTERIS frondibus supradecompositis: foliolis pin-*aquilina.* natis: pinnis lanceolatis: infimis pinnatifidis, superioribus minoribus. *Sp. pl.* 1533.

Filix foliis triplicato-pinnatis, pinnis nervosis integerrimis, ultimis lanceolatis. *Hall. bist.* 1688.

Filix femina. *Fuchf. hist.* 596. *Cam. ep.* 992. *Ger. em.* 1128. R. syn. 124. vulgaris. *Park.* 1037.

Filix ramosa major, pinnulis obtusis non dentatis. *B. pin.* 357.

2 Filicula saxatilis ramosa maritima nostras. R. syn. 125. *Pluk. ph. t.* 182. f. 1.

Anglis, female Brakes, or Fern.

Habitat in sylvis et ericetis. 4. VIII.

ASPLENIUM. 1178.

Frustrificationes in lineolis disci frondis sparsis.

* *Fronde simplici.*

- Scolopendrium.* 1. ASPLENIUM frondibus simplicibus cordato-lingulatis integerrimis, stipitibus hirsutis. *Sp. pl.* 1537.
 Asplenium petiolis hirsutis, folio longe lineari-lanceolato integerrimo postice cordato. *Hall. hist.* 1695.
 Phyllitis. *Cam. ep.* 579. *Ger. em.* 1138. *R. syn.* 116.
 f. *Lingua cervina vulgaris.* *Park.* 1046.
Lingua cervina officinarum. *B. pin.* 353.
Anglis, Hart's tongue Spleen-wort.
Habitat in umbrosis saxosis. 4. VIII.
Pharmac. Phyllitidis Folia.

** *Fronde pinnatifida.*

- Ceterach.* 2. ASPLENIUM frondibus pinnatifidis: lobis alternis confluentibus. *Sp. pl.* 1538.
 Asplenium foliis pinnatis, pinnis in basi dilatatis obtusis, subtus muscosis. *Hall. hist.* 1694.
 Asplenium. *Cam. ep.* 640. f. *Ceterach.* *B. hist.* III. 749. *Ger. em.* 1139. *Park.* 1046. *R. syn.* 118.
Ceterach officinarum. *B. pin.* 354.
Anglis, common Spleenwort.
Habitat in muris antiquis et fissuris rupium humidis.
Circa-Bristolium copiose. 4. V—X.
Pharmac. Ceterach f. Asplenii Herba.

*** *Fronde pinnata.*

- Trichomanes.* 3. ASPLENIUM frondibus pinnatis: pinnis subrotundis crenatis. *Sp. pl.* 1540. *Hall. hist.* 1693.
 Trichomanes. *Fuchs. hist.* 796. *Cam. ep.* 925. *Park.* 1051. *R. syn.* 119. f. *Polytrichum officinarum.*
B. pin. 356.
Trichomanes mas. *Ger. em.* 1146.
 β *Trichomanes ramosum.* *B. hist.* III. 755. *R. syn.* 119. *majus et minus.* *B. pin.* 356.
Anglis, Trichomanes Spleenwort.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

POLYGAMIA MONOECIA.

Atriplex foliis imis hamatis, reliquis lingulatis, valvulis feminalibus dentatis. *Hall. hist.* 1616.

Atriplex sylvestris angustifolia. *Ger. em.* 326. *Park.* 748. *R. syn.* 151.

Atriplex angusto oblongo folio. *B. pin.* 119.

β Atriplex caule herbaceo, foliis hastatis dentatis. *Fl. angl.* 376.

Atriplex angustifolia laciniata. *R. syn.* 152.

Anglis, spreading Orache.

Habitat in ruderatis et ad sepes. ☉. VIII.

serrata. 5. ATRIPLEX caule herbaceo erectiusculo, foliis linearibus serratis. *Fl. angl.* 377.

Atriplex angustifolia maritima dentata. *R. syn.* 152.

Atriplex maritima angustifolia. *B. pin.* 152.

Anglis, serrated Orache. *Pet. h. br. t.* 7. f. 4.

Habitat in ruderatis et littoribus maritimis. ☉. VIII.

littoralis. 6. ATRIPLEX caule herbaceo erecto, foliis omnibus linearibus integerrimis. *Sp. pl.* 1494.

Atriplex angustissimo et longissimo folio. *Herm. lugd.* 79. *Tourn.* 505. *R. syn.* 153.

β Atriplex maritima scopariæ folio. *R. syn.* 153.

γ Atriplex maritima angustifolia, obtusiore folio. *R. syn.* 153.

Anglis, grass Orache. *Pet. h. br. t.* 7. f. 6.

Habitat in ruderatis et littoribus maritimis. ☉. VIII.

pedunculata. 7. ATRIPLEX caule herbaceo divaricato, foliis lanceolatis obtusis integris, calycibus femineis pedunculatis. *Sp. pl.* 1675.

Atriplex marina femine lato. *Ph. br.* *R. syn.* 153.

Atriplex maritima Halimus dicta, humilis erecta, femine folliculis membranaceis bivalvibus in latitudinem expansis et utrinque recurvis, longo pediculo insidentibus, clauso. *Pluk. ph. t.* 36. f. 1.

β Atriplex maritima nostras Ocymi minoris folio. *D. Pluk.* *R. syn.* 253.

Anglis, pedunculated Orache.

Habitat in littoribus maritimis; prope Schirbeck vicum milliare a Bostoniam distantem, copiose provenit. *D.*

Pluknet.

Pluknet. *Found in the Isle of Thanet, near the ferry-house, in great plenty.* Mr. Sherard. R. syn. *prope Yarmouth.* ☉. VII—IX.

A C E R. 1155.

Hermaphrod. *Cal.* quinquefidus. *Cor.* pentapetala. *Stam.* octo. *Pist.* unicum. *Caps.* duæ s. tres, monospermæ, ala terminatæ.

Masc. *Cal.* quinquefidus. *Cor.* pentapetala. *Stam.* octo.

1. ACER foliis quinquelobis inæqualiter ferratis, floribus racemosis *Sp. pl.* 1495. *Pseudo-platanus.*

Acer foliis quinquelobis, acute ferratis, racemis pendulis. *Hall. hist.* 1029.*****

Acer montanum candidum. *B. pin.* 430.

Acer majus. *Dod. pem.* 840. *Ger. em.* 1484. *R. syn.* 470. *latifolium, Sycomorus falso dictum.* *Park.* 1425.

Anglis, greater Mapple or Sycamore. *Hunt. Evel. syl.* 200.

Habitat in sepibus et ad domos. ♀. V. VI.

2. ACER foliis lobatis obtusis emarginatis. *Sp. pl.* 1497. *campestre.*

Acer foliis semitrilobis, obtusis, lobis lateralibus emarginatis. *Hall. hist.* 1029*****.

Acer minus. *Dod. pem.* 840. *Ger. em.* 1484. *R. syn.* 470. *et vulgare.* *Park.* 1426.

Acer campestre et minus. *B. pin.* 431.

β *Acer campestre et minus, fructu rubente.* *Vail. paris. R. syn.* 470.

Anglis, common Mapple. *Hunt. Evel. syl.* 190.

Habitat in sepibus et dumetis. ♀. VI.

D I O E C I A.

F R A X I N U S. 1160.

Hermaphrod. *Cal.* quadripartitus. *Cor.* o. *Stam.* duo. *Pist.* unicum. *Sem.* unicum, lanceolatum.

Fem.

Atriplex foliis imis hamatis, reliquis lingulatis, valvulis feminalibus dentatis. *Hull. hist.* 1616.

Atriplex sylvestris angustifolia. *Ger. em.* 326. *Park.* 748. *R. syn.* 151.

Atriplex angusto oblongo folio. *B. pin.* 119.

β Atriplex caule herbaceo, foliis hastatis dentatis. *Fl. angl.* 376.

Atriplex angustifolia laciniata. *R. syn.* 152.

Anglis, spreading Orache.

Habitat in ruderatis et ad sepes. ☉. VIII.

serrata. 5. ATRIPLEX caule herbaceo erectiusculo, foliis linearibus serratis. *Fl. angl.* 377.

Atriplex angustifolia maritima dentata. *R. syn.* 152.

Atriplex maritima angustifolia. *B. pin.* 152.

Anglis, serrated Orache. *Pet. h. br. t. 7. f. 4.*

Habitat in ruderatis et littoribus maritimis. ☉. VIII.

littoralis. 6. ATRIPLEX caule herbaceo erecto, foliis omnibus linearibus integerrimis. *Sp. pl.* 1494.

Atriplex angustissimo et longissimo folio. *Herm. lugd.* 79. *Tourn.* 505. *R. syn.* 153.

β Atriplex maritima scopariæ folio. *R. syn.* 153.

γ Atriplex maritima angustifolia, obtusiore folio. *R. syn.* 153.

Anglis, grass Orache. *Pet. h. br. t. 7. f. 6.*

Habitat in ruderatis et littoribus maritimis. ☉. VIII.

pedunculata. 7. ATRIPLEX caule herbaceo divaricato, foliis lanceolatis obtusis integris, calycibus femineis pedunculatis. *Sp. pl.* 1675.

Atriplex marina femine lato. *Ph. br. R. syn.* 153.

Atriplex maritima Halimus dicta, humilis erecta, femine folliculis membranaceis bivalvibus in latitudinem expansis et utrinque recurvis, longo pediculo insidentibus, clauso. *Pluk. ph. t. 36. f. 1.*

β Atriplex maritima nostras Ocymi minoris folio. *D. Pluk. R. syn.* 253.

Anglis, pedunculated Orache.

Habitat in littoribus maritimis; prope Schirbeck vicum milliare a Bostoniam distantem, copiose provenit. *D. Pluknet.*

Pluknet.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Fem. *Pist.* unicum. *Sem.* unicum, lanceolatum.

- excelsior.* 1. FRAXINUS foliis ferratis, floribus apetalis. *Sp. pl.* 1509. *Fl. suec.* 926.
 Fraxinus apetalà, foliis pinnatis ferratis. *Hall. hist.* 528.
 Fraxinus. *Dod. pempt.* 771. *Cam. ep.* 64. *Ger. em.* 1472. *R. syn.* 469. *excelsior.* *B. pin.* 416.
Anglis, common Ash-tree. *Hunt. Ev. sylv.* 150.
 Habitat in sylvis et sepibus. *h.* III. IV.
 Obs. variat foliis pinnatis et simplicibus.

I L E X.

Hermaphrod. *Cal.* quinquedentatus. *Cor.* quinquepartita, rotata. *Stam.* quinque, rotata. *Stylus* nullus. *Stigmata* quatuor. *Bacca* tetrasperma.

Masc. *Cal.* quadridentatus. *Cor.* quadripartita. *Stam.* quatuor.

- aquifolium.* 1. ILEX foliis ovatis acutis spinosis. *Sp. pl.* 181.
 Ilex aculeata baccifera. *B. pin.* 425.
 Aquifolium. *Cam. ep.* 84. *Hall. hist.* 667.
 Agrifolium. *Ger. em.* 1338. s. Aquifolium. *Part.* 1486. *R. syn.* 466.
Anglis, common Holly. *Hunt. Ev. sylv.* 383.
 Habitat in sylvis, sepibus et ericetis. *h.* V.

Classis XXIV.

CRYPTOGAMIA

FILICES.

EQUISETUM. 1169.

Spica fructificationibus peltatis, basi dehiscentibus multivalvi.

1. EQUISETUM caule spicato, frondibus compositis. *sylvaticum.*
Sp. pl. 1516.

Equisetum foliis repetito ramosis, vaginis levissimis.
Hall. hist. 1680.

Equisetum sylvaticum. *Tab. ic. 253. Ger. em. 1114.*
R. syn. 130. tenuissimis setis. B. pin. 16.

Equisetum omnium minus tenuifolius. *Park. 1201.*

β Equisetum sylvaticum, procumbens, setis uno versu dispositis, *R. syn. 131.*

γ Equisetum palustre tenuissimis et longissimis setis.
B. pin. 16. pr. 24.

Anglis, wood Horse-tail.

Habitat in sylvis et umbrosis humidis. 4. IV. V.

2. EQUISETUM scapo fructificante nudo: sterili *arvense.*
frondoso. *Sp. pl. 1516.*

Equisetum caule florifero nudo, sterili verticillato, radiorum duodecim. *Hall. hist. 1676.*

Equisetum arvense longioribus setis. *B. pin. 16. Park. 1202. R. syn. 130.*

Equisetum segetale. *Ger. em. 1114.*

β Equisetum pratense longissimis setis. *B. pin. 16.*

γ Equisetum nudum minus variegatum basiliense. *B. pin. 16. pr. 25. R. syn. 130.*

Anglis,

Anglis, corn Horse-tail.

Habitat in arvis humidis. 4. III. IV.

palustre. 3. EQUSETUM caule angulato, frondibus simplicibus. *Sp. pl.* 1516. *Fl. suec.* 928.

Equisetum caule sulcato, ramis multifloris, foliis indivisis. *Hall. hist.* 1677.

Equisetum palustre brevioribus setis. *B. pin.* 16.

Equisetum palustre. *Lob. ic.* 795. *Ger. em.* 1114.

R. syn. 131. minus. *Park.* 1200.

β Equisetum palustre minus polystachion. *B. pin.* 16.

pr. 24. *R. syn.* 131. t. 5. f. 3.

Anglis, marsh Horse-tail.

Habitat in palustribus et aquosis. 4. VI. VII.

fluviale. 4. EQUSETUM caule striato, frondibus subsimplicibus. *Sp. pl.* 1517.

Equisetum caule florifero nudo, sterili verticillato, radiorum quadraginta. *Hall. hist.* 1675.

Equisetum. *Cam. epit.* 770. majus. *Ger. em.* 1113.

R. syn. 130. palustre. *Park.* 1290.

Equisetum palustre longioribus setis. *B. pin.* 15.

Anglis, river Horse-tail.

Habitat in palustribus et aquosis et ad ripas fluviorum et lacuum.

limosum. 5. EQUSETUM caule subnudo lævi. *Sp. pl.* 1517.

Equisetum nudum lævius nostras. *R. syn.* 131. t. 5. f. 6.

Anglis, smooth Horse-tail.

Habitat in aquosis. 4. VI. VII.

hyemale. 6. EQUSETUM caule nudo scabro basi subramoso.

Sp. pl. 1517.

Equisetum caule subnudo asperrimo, vaginis caulinis indivisis, ramis foliatis. *Hall. hist.* 1679.

Equisetum nudum. *Ger. em.* 1113. *R. syn.* 131.

Equisetum junceum ramosum. *Park.* 1207.

Equisetum foliis nudum ramosum. *B. pin.* 16.

β Equisetum foliis nudum non ramosum s. junceum.

B. pin. 16.

Equisetum

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

CRYPTOGAMIA FILICES.

Filix florida s. Osmunda regalis. *Ger. em.* 1131.
Anglis, common Osmund-royal, or flowering fern.
Habitat in aquosis et palustribus putridis. 4. VIII.

*** *Fronibus aliis foliaceis. aliis fructificantibus.*

Spicant. 3. OSMUNDA frondibus lanceolatis pinnatifidis : laciniis confluentibus integerrimis parallelis. *Sp. pl.* 1522. *Fl. suec.* 936. *dan.* 217.

Struthiopteris. *Hall. hist.* 1687.

Lonchitis aspera. *Ger. em.* 1140. *R. syn.* 118. minor.
Cam. ep. 665. *B. pin.* 359. *Park.* 1042.

Anglis, spleenwort Osmund-royal.

Habitat in nemorosis et ericetis humidis. 4. VIII.

crispa. 4. OSMUNDA frondibus supradecompositis : pinnis alternis subrotundis incis. *Sp. pl.* 1522. *Fl. dan.* 496.

Filix foliis triplicato-pinnatis, sterilibus obtusis dentatis, floralibus ovatis integerrimis. *Hall. hist.* 1689.

Adiantum album crispum alpinum. *Schwenkf. Cat.* 10. *R. syn.* 126.

Adiantum album floridum s. *Filicula petræa crispa per-elegans.* *R. syn. ed. II.* 51. *Pluk. ph. t.* 9. *f.* 3.

Anglis, curl'd Osmund-royal, or stone-fern.

Habitat in petris et rupibus. *In comitatibus Eboracensi, Westmorlandico et Wallia frequens.* 4. VIII.

ACROSTICHUM. 1173.

Fruetificationes discum totum frondis tegentes.

* *Fronde simplici divisa.*

septentri-onaie. 1. ACROSTICHUM frondibus nudis linearibus laciniatis. *Sp. pl.* 1524. *Fl. dan.* 60.

Acrostichum caule bis bifurcato. *Hall. hist.* 1714.

Filix saxatilis Tragi. *Lob. ic.* 47. *B. hist.* III. 755.
Park. 1045. *R. syn.* 120.

Filix saxatilis corniculata. *B. pin.* 358.

Anglis, forked or horned Acrostichum.

Habitat in fissuris rupium. *Ad cacumen montis Carnedh-Lhewelyn prope Lhan-Lhechyd in agro Arvonensi invenit,*

invenit D. Lhwyd. *in muris antiquis* Lhan Dethylæ und circiter milliari a Lhan-Rhoost aquilonem versus D. Richardson. R. syn. *in monte* Ingleborough, D. Tosfield, *in montibus supra* Ambleside *in agro* Westmorlandico. 4. VII.

•• *Fronde subbipinnata.*

2. ACROSTICHUM? frondibus subbipinnatis; pinnis *ilvense.* opposito-coadunatis obtusis subtus hirsutis: basi integerrimis. *Sp. pl.* 1528.

Filix alpina pedicularis rubræ foliis subtus villosis. D. Lhwyd. R. syn. 118.

Filix pumila lonchitidis maranthæ species cambrobri-tannica. *Pluk. ph. t.* 89. f. 8.

Filicula alpina tenerior, alis latiusculis brevioribus integris profunde dentatis. *Hist. Ox. III.* 576. t. 3. f. 23.

Anglis, hairy Acrostichum.

Habitat in fissuris rupium. *It grows on a moist black rock, almost at the top of* Clogwyn y Garnedd, *facing North-west, directly above the lower lake.* D. Richardson, R. syn. *in eodem loco inveni.* 4. VII—IX.

P T E R I S. 1174.

Fruetificationes in lineis marginalibus.

1. PTERIS frondibus supradecompositis: foliolis pin-aquilina. natis: pinnis lanceolatis: infimis pinnatifidis, superioribus minoribus. *Sp. pl.* 1533.

Filix foliis triplicato-pinnatis, pinnis nervosis integerrimis, ultimis lanceolatis. *Hall. hist.* 1688.

Filix femina. *Fuchs. hist.* 596. *Cam. ep.* 992. *Ger. em.* 1128. R. syn. 124. vulgaris. *Park.* 1037.

Filix ramosa major, pinnulis obtusis non dentatis. *B. pin.* 357.

2 Filicula saxatilis ramosa maritima nostras. R. syn. 125. *Pluk. ph. t.* 182. f. 1.

Anglis, female Brakes, or Fern.

Habitat in sylvis et ericetis. 4. VIII.

ASPLENIUM. 1178.

Frustrificationes in lineolis disci frondis sparsis.

* *Fronde simplici.*

- Scolopendrium.* 1. ASPLENIUM frondibus simplicibus cordato-lingulatis integerrimis, stipitibus hirsutis. *Sp. pl.* 1537.
 Asplenium petiolis hirsutis, folio longe lineari-lanceolato integerrimo postice cordato. *Hall. hist.* 1695.
 Phyllitis. *Cam. ep.* 579. *Ger. em.* 1138. *R. syn.* 116.
 f. Lingua cervina vulgaris. *Park.* 1046.
 Lingua cervina officinarum. *B. pin.* 353.
Anglis, Hart's tongue Spleen-wort.
Habitat in umbrosis saxosis. 4. VIII.
Pharmac. Phyllitidis Folia.

** *Fronde pinnatifida.*

- Ceterach.* 2. ASPLENIUM frondibus pinnatifidis: lobis alternis confluentibus. *Sp. pl.* 1538.
 Asplenium foliis pinnatis, pinnis in basi dilatatis obtusis, subtus muscosis. *Hall. hist.* 1694.
 Asplenium. *Cam. ep.* 640. f. *Ceterach.* *B. hist.* III. 749. *Ger. em.* 1139. *Park.* 1046. *R. syn.* 118.
 Ceterach officinarum. *B. pin.* 354.
Anglis, common Spleenwort.
Habitat in muris antiquis et fissuris rupium humidis.
Circa-Bristolium copiose. 4. V—X.
Pharmac. Ceterach f. Asplenii Herba.

*** *Fronde pinnata.*

- Trichomanes.* 3. ASPLENIUM frondibus pinnatis: pinnis subrotundis crenatis. *Sp. pl.* 1540. *Hall. hist.* 1693.
 Trichomanes. *Fuchs. hist.* 796. *Cam. ep.* 925. *Park.* 1051. *R. syn.* 119. f. Polytrichum officinarum. *B. pin.* 356.
 Trichomanes mas. *Ger. em.* 1146.
 β Trichomanes ramosum. *B. hist.* III. 755. *R. syn.* 119. majus et minus. *B. pin.* 356.
Anglis, Trichomanes Spleenwort.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

liis, superioribus semitribus rhomboideis circumferratis. *Hall. hist.* 1691.

Ruta muraria. *B. pin.* 356. *Ger. em.* 1154. *R. syn.* 122. f. *Salvia vitæ.* *Park.* 1050.

Adiantum album. *Tab ic.* 796.

β Adiantum album tenuifolium, rutæ murariæ accedens.

B. hist. III. 753. *R. syn.* 123. *Pluk. ph.* 1. 3. f. 3.

Anglis, white Spleenwort, or Maidenhair.

Habitat in muris antiquis et rimis petrarum humidis.

4. VI—X.

Pharmac. Adianti albi *Herba.*

Adiantum nigrum.

7. ASPLENIUM frondibus subtriplicato-pinnatis deltoideis, foliolis alternis, pinnis ovatis inciso-ferratis.

Asplenium frondibus tripinnatis : foliolis alternis : pinnis lanceolatis inciso-ferratis. *Sp. pl.* 386. *Fl. dan.* 250.

Asplenium foliis triangularibus, pinnis pinnatis, pinnulis semipinnatis, lobulis ovatis ferratis. *Hall. hist.* 1692.

Adiantum nigrum officinarum. *B. hist.* III. 743. *R. syn.* 126.

Adiantum foliis longioribus pulverulentis, pediculo nigro. *B. pin.* 355.

Adiantum nigrum vulgare. *Park.* 1406.

Onopteris mas. *Ger. em.* 1137.

β Filix minor longifolia, tarsis raris, pinnulis longis, tenuissimis et oblongis laciniis fimbriatis. *R. syn.* 127.

γ Filix pumila petræa nostras adianti nigri foliorum æmula, saxorum interuenis prorumpens. *Pluk. alm.* 91. *R. syn.* 127.

Anglis, black Spleenwort, or Maidenhair.

Habitat in umbrosis et muris antiquis. 4. IV—X.

Pharmac. Adianti nigri *Herba.*

lanceolatum.

8. ASPLENIUM frondibus duplicato-pinnatis lanceolatis : foliolis alternis ; pinnis obovatis inciso-crenatis.

Asplenium frondibus duplicato-pinnatis : pinnis obovatis

vatis crenatis : foliolis inferioribus minoribus. *Sp. pl.* 1541.

Filix elegans, adianto nigro accedens segmentis rotundioribus. *R. syn.* 127?

Anglis, Spear Spleenwort.

Habitat in muris antiquis et rupibus circa St. Ives et alibi in Cornubia copiose. 4. V—IX.

POLYPODIUM. 1179.

Frustrificationes in punctis subrotundis sparsis per discum frondis.

* *Fronde pinnatifida lobis coadunatis.*

1. POLYPODIUM frondibus pinnatifidis : pinnis oblongis subserratis obtusis, radice squamata. *Sp. pl.* 1544.

Polypodium foliis pinnatis lanceolatis, radice squamata. *Hall. hist.* 1696.

Polypodium vulgare, *B. pin.* 359. *Park.* 1039.

Polypodium. *Matth.* 1292. *Ger. em.* 1132. *R. syn.* 117.

- β *Polypodium murale* pinnulis serratis. *Pct. bot. hort.* II. *R. syn.* 117.

- γ *Polypodium frondibus pinnatis, pinnis lanceolatis lacero-pinnatifidis serratis.* *Sp. pl.* 1546. *cambricum.*

Polypodium cambrobritannicum, pinnulis ad margines laciniatis. *R. syn.* 117.

Polypodium cambrobritannicum, lobis foliorum profunde dentatis. *Hist. Ox.* III. 565 f. 14. t. 2. f. 8.

Filix amplissima, lobis foliorum laciniatis cambrica. *Pluk. ph.* t. 30. f. 1.

Anglis, common γ jagged Polypody.

Habitat in muris antiquis umbrosis et ad radices arborum.

γ on a rock in a wood, near Dennys Powis Castle, not far from Caerdiff in Glamorganshire. *R. syn.* 4. VI—X.

Pharmac. Polypodii. Radix, Herba.

** *Fronde pinnata.*

2. POLYPODIUM frondibus pinnatis : pinnis lunulatis, *Lonchitis.*

CRYPTOGAMIA FILICES.

tis, ciliato-ferratis declinatis, stipitibus strigosis.
Sp. pl. 1548. *Fl. dan.* 497.

Polypodium foliis pinnatis, pinnis ciliatis dentatis, appendiculatis. *Hall. hist.* 1711.

Lonchitis aspera. *B. pin.* 359. major. *Cam. ep.* 664.
Ger. em. 1140. *R. syn.* 118. *Matthioli.* *Park.* 1042.

Anglis, rough Polypody, or Spleenwort.

Habitat in fissuris rupium; in montibus circa Llanberis, copiose. 4. V—IX.

*** *Fronde subbipinnata.*

fontanum. 3. **POLYPODIUM** frondibus subbipinnatis lanceolatis, foliolis subrotundis argute incis, stipite lævi.
Sp. pl. 1550.

Polypodium duriusculum, pinnis laxè pinnatis, pinnulis bidentatis et tridentatis argutis. *Hall. hist.* 1706.

Adiantum album. *Lob. ic.* 810.

Filicula fontana. *Tabern.* 792. minor. *B. pin.* 358.

Filix saxatilis non ramosa minima. *R. cat. eur.* *Pluk. ph. t.* 89. f. 2.

Filix saxatilis omnia minima elegantissima. *Hort. reg. paris.* *Pluk.* 89. f. 3.

Anglis, rock Polypody.

Habitat in muris antiquis et rupibus. supra Hammertham Church, D. Bradney; in locis saxosis, prope Wybourn in Westmorlandia. 4. VI—IX.

phogopteris. 4. **POLYPODIUM** frondibus subbipinnatis, foliolis infimis reflexis: paribus pinnula quadrangulati coadunatis. *Sp. pl.* 1550. *Fl. dan.* 497.

Polypodium foliis pinnatis reflexis, pinnis ovatis hirsutis, primis cum nervo confluentibus. *Hall. hist.* 1698.

Filix minor britannica, pediculo pallidiore, alis inferioribus, deorsum spectantibus. *Hist. Ox. III.* 575.

f. 14. t. 4. f. 17. *R. syn.* 122.

Anglis, wood Polypody.

Habitat in rimis petrarum humidiusculis et umbrosis in comitatibus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

rhæticum. 8. **POLYPODIUM** frondibus bipinnatis : foliolis pinnisque remotis lanceolatis : ferraturis acuminatis. *Sp. pl.* 1552.

Polypodium triplicato-pinnatum, pinnulis tertiis semipinnatis lobulis bifidis. Hall. hist. 1710.

Filix montana ramosa minor argute denticulata. R. syn. 124.

Filix alpina myrrhidis facie cambrobritannica. Pluk. ph. t. 89. f. 4.

Filix fontana major s. Adiantum album, filicis folio. B. pin. 358?

Filix rhætica tenuissime denticulata. B. hist. III. 740?

Anglis, stone Polypody.

Habitat in montibus saxosis ; in Westmorlandia passim ; ad summitatem montis Glyder. R. syn. 2. VI—IX.

Obs. nimis affinis præcedenti, sed multo minor, an distincta sit species ?

Filix mas. 9. **POLYPODIUM** frondibus bipinnatis ; pinnis obtusis crenulatis, stipite paleaceo. *Sp. pl.* 1551.

Polypodium pinnis pinnatis obtusis dentatis. Hall. hist. 1701.

Filix mas vulgaris. Park. 1036. *R. syn.* 120.

Filix mas non ramosa dentata. B. pin. 258.

Filix mas non ramosa, pinnulis latis densis minutim dentatis. Ger. em. 1130.

Anglis, male Polypody, or Fern.

Habitat in sylvis, ericetis, et locis saxosis. 4. VI—IX.

Pharmac. Filicis maris Radix.

Filix femina. 10. **POLYPODIUM** frondibus bipinnatis : pinnulis lanceolatis pinnatifidis acutis. *Sp. pl.* 1552.

Polypodium pinnis pinnatis, pinnulis lanceolatis semipinnatis, lobis acute bisectis. Hall. hist. 1740.

Filix mas non ramosa, pinnulis angustis raris profunde dentatis. Ger. em. 1130. *Hist. Ox.* III. 579.

f. 14. t. 3. f. 8. R. syn. 121. *Pluk. ph.* 180. f. 4.

Anglis, female Polypody, or Fern.

11. POLYPODIUM frondibus bipinnatis; pinnis lu-*aculea-*
nulatis ciliato-dentatis, stipite strigoso. *Sp. pl. tum.*
1552.

Filix mas aculeata major. *B. pin. 558. pr. 151.*

Filix mas non ramosa, pinnulis latis auriculatis spino-
sis. *Ger. em. 1130. Pluk. ph. 179. f. 6. R. syn.*
121.

β Filix aculeata major, pinnulis auriculatis crebrioribus,
foliis integris angustioribus. *R. syn. 121.*

γ Filix mas aculeata nostras, alis expansis, muscosa la-
nugine aspersa. *Pluk. ph. 180. f. 1. R. syn. 121.*

Anglis, prickly Polypody.

Habitat in sylvis et umbrosis. 4. VI—IX.

12. POLYPODIUM frondibus bipinnatis: pinnulis lobatum.
ovatis ciliato-ferratis suprema basi auriculatis, sti-
pite strigoso.

Polypodium pinnis pinnatis ciliatis ferratis appendi-
culatis. *Hall. hist. 1712?*

Filix lonchitidis affinis. *Pluk. ph. 180. f. 3. R. syn.*
121.

Anglis, lobed Polypody.

Habitat in umbrosis et ad sepes. 4. VI—IX.

Obs. affinis præcedenti, an distincta sit species?

13. POLYPODIUM frondibus bipinnatis: foliolis re-*fragile.*
motis: pinnis subrotundis incis. *Sp. pl. 1553.*

Filix saxatilis caule tenui fragile. *R. syn. 125. Pluk.*
ph. 180. f. 5.

Filix pumila saxatilis altera clusii. *Park. 1043.*

Filix saxatilis non ramosa, nigris maculis punctata.
B. pin. 358.

β Adiantum nigrum pinnulis cicutarizæ divisura. *D.*
Bobart. R. syn. 126.

Anglis, brittle Polypody.

Habitat in aridis lapidosis in boreali parte Britanniz
et in Wallia; circa Bristol. 4. VI—IX.

**** *Fronde supradecomposita.*

- Dryopteris.* 14. **POLYPODIUM** frondibus supradecompositis : foliolis ternis bipinnatis. *Sp. pl.* 1555.
 Polypodium pinnis pinnatis conjugatis, pinnulis ovatis obtusis crenatis. *Hall. hist.* 1699.
 Filix ramosa minor. *B. hist. III.* 741. *R. syn.* 125.
 Filix saxatilis ramosa nigris maculis punctata. *B. pin.* 358.
 Filix pumila saxatilis prima Clusii. *Park.* 1043.
 Dryopteris Tragi. *Ger. em.* 1135.
Anglis, branched Polypody.
 Habitat in aridis saxosis in comitatibus Eboracensi, Lancastriensi, Westmorlandico, &c. frequens. 4. VI—IX.

A D I A N T U M. 1180.

Fruetificationes in maculis terminalibus, sub replicato margine frondis.

- Capillus veneris.* 1. **ADIANTUM** frondibus decompositis : foliolis alternis : pinnis cuneiformibus lobatis pedicellatis. *Sp. pl.* 1559.
 Adiantum petiolis ramosis, foliis flabelliformibus lobatis. *Hall. hist.* 1713.
 Adiantum. *Cam. ep.* 924. verum s. Capillus veneris verus. *Park.* 1049.
 Adiantum foliis coriandri. *B. pin* 355.
 Capillus veneris verus. *Ger. em.* 1143. *R. syn.* 123.
Anglis, true Maidenhair.
 Habitat in rupibus et muris humidiusculis ; Found by Mr. Lhywd at Barry island, and Porth Kirig in Glamorganshire. *R. syn.* 4. V—IX.
- trapeziforme.* 2. **ADIANTUM** frondibus supradecompositis : foliolis alternis : pinnis rhombeis incisive utrinque fructificationibus. *Sp. pl.* 1559.
 Adiantum majus coriandri folio, adianto vero affine, pediculo pallide rubente. *Sibb. pr.* 7. *R. syn.* 124.
 Adianto

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

PILULARIA. 1183.

Flores *Masculi* ad latus frondis.

Fructificatio *Feminea* ad radicem, globosa, quadrilobularis.

globulifera.

1. PILULARIA. *Sp. pl.* 1563. *Fl. dan.* 223.

Pilularia palustris juncifolia. *Vaill. paris.* 159. t. 15. f. 6. *Dill. musc.* 538. t. 79. f. 1.

Graminifolia palustris repens, vasculis granorum piperis æmulans. *R. syn.* 130.

Anglis, Pepper-grass.

Habitat in inundatis præsertim solo arenoso; prope fontes medicatos Strethamenses, D. Wilmer. in ericeto Hounsejano. *R. syn.* 4. V—IX.

ISOETES. 1184.

Flores *Masculi* *Anthera* intra basin frondis.

Flores *Feminei* *Capsula* intra basin frondis.

lacustris. 1. ISOETES. *Sp. pl.* 1563. *it. scan.* 417. *ic. Fl. dan.* 191.

Calamaria folio longiore et graciliore. *Dill. musc.* 541. t. 80. f. 2.

Subularia vulgaris erecta, folio rigidissimo. *R. syn.* 306.

2 *Subularia repens* folio minus rigido. *R. syn.* 306.

Subularia fragilis, folio longiore et tenuiore. *R. syn.* 307.

Anglis, common Quillwort.

Habitat in fundo lacuum, circa Llanberris copiose; in Loch Tay, in Scotia. D. Arch. Menzies. 4. V—IX.

MUSCI.

LYCOPODIUM. 1185.

Anthera bivalvis, sessilis. *Calyptra* nulla.

clavatum. 1. LYCOPODIUM foliis sparsis filamentosis, spicis teretibus pedunculatis geminis. *Sp. pl.* 1564. *Fl. dan.* 126. *Weis Crypt.* 269.

Lycopodium

Lycopodium ramis prolixis, spicis petiolatis gemellis, foliis linearibus pilo aristatis. *Hall. hist.* 1722.

Lycopodium. *Tab. ic.* 814. *R. syn.* 107. vulgare pilosum anfragosum et repens. *Dill. musc.* 441. t. 58.

f. 1.

Muscus terrestris clavatus. *B. pin.* 360.

Muscus clavatus s. *Lycopodium*. *Ger. em.* 1562. *Park.* 1307.

Anglis, common Club-moss.

Habitat in montibus et ericetis, in ericeto Hounsejano et prope Esber. 4. VII. VIII.

Pharmac. Musci clavati *Herba Lycopodii Semina*.

2. LYCOPODIUM foliis sparsis integerrimis, spicis inundantibus terminalibus foliosis. *Sp. pl.* 1565. *Fl. dan.* 336. *tum.*

Lycopodium spicis sessilibus, foliis linearibus confertissimis incurvis. *Hall. hist.* 1721.

Lycopodium palustre repens, clava singulari. *Vaill. paris.* 123. t. 16. f. 11. *Dill. musc.* 452. t. 62. f. 7.

Muscus terrestris repens, clavis singularibus foliolis erectis. *R. syn.* 108.

Anglis, marsh Club moss.

Habitat in ericetis humidis passim. in ericeto Hounsejano et prope Esber. 4. VI—IX.

3. LYCOPODIUM foliis sparsis ciliatis lanceolatis, selaginoidibus spicis solitariis terminalibus foliosis. *Sp. pl.* 1565. *des.* *Fl. dan* 70.

Lycopodium spicis sessilibus, foliis ovato-lanceolatis serratis confertis. *Hall. hist.* 1717. t. 45. f. 1.

Selaginoides foliis spinosis. *R. syn.* 1066. *Dill. musc.* 460. t. 88. f. 1.

Anglis, prickly Club-moss.

Habitat in pascuis et ericetis montosis; in boreali parte Britanniae et in Wallia frequens. 4. VI—IX.

4. LYCOPODIUM foliis sparsis octofariis, caule Selago. dichotomo erecto fastigiato, floribus sparsis. *Sp. pl.* 1565. *Weis. cryp.* 273.

Lycopodium spica sessili, foliis linearibus lanceolatis confertis. *Hall. hist.* 1716.

Selago

Selago vulgaris abietis rubræ facie. *Dill. musc.* 435. t. 56. f. 1.

Selago foliis et facie abietis. *Fl. jen.* 330. *R. syn.* 106.

Chamæpeuce. *Turn. hist. par.* I. 129.

Muscus erectus ramosus saturate viridis. *B. pin.* 360. *Anglis,* fir Club-moss.

Habitat in ericetis montosis, in comitatibus Eboracensi, Lancastriensi, Westmorlandico, Cambrobritannico. etc. frequens. 4. IV—X.

*annoti-
zum.*

5. **LYCOPODIUM** foliis sparsis quinquefariis subserratis, furculis annotino-articulatis, spicis terminalibus glabris erectis. *Sp. pl.* 1566.

Lycopodium ramis prolixis, spicis sessilibus foliis linearibus subserratis. *Hall. hist.* 1720.

Lycopodium elatius juniperinum, clavis singularibus, sine pediculis. *Dill. musc.* 455. t. 63. f. 9. *R. syn.* 107,

Muscus clavatus, juniperinis foliis reflexis, clavis singularibus sine pediculis. *Hist. Ox.* III. 624. f. 15. t. 5. f. 3.

Anglis, Welch Club-moss.

Habitat in alpibus Arvonizæ sed rarius; copiosam observavimus in monte Rhiwr Glyder, qua super lacum Llyn y cown, prope ecclesiam S. Peris, nec rarior in depressis ejusdem montis Glyder, qua rupem y Triwan spectat. *D. Llhwyd.* *R. syn.* 4. VI—IX.

alpinum. 6. **LYCOPODIUM** foliis quadrifariam imbricatis acutis, furculis erectis dichotomis, spicis sessilibus teretibus. *Sp. pl.* 1567. *Fl. dan.* 79. *Weis. Cryp.* 275.

Lycopodium spicis sessilibus, foliis lanceolatis adpressis, utrinque ternatis. *Hall. hist.* 1719.

Lycopodium sabinæ facie. *Dill. musc.* 435. t. 58. f. 2.

Muscus clavatus cupressiformis. *Park.* 1309.

Muscus clavatus foliis cupressi. *Ger. em.* 1562. *B. pin.* 360.

Anglis, mountain Club-moss.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

P H A S C U M. 1189.

Anthera operculata : ore ciliata. *Calyptra* caduca, minuta.

subula-
tum.

1. PHASCUM acaule, anthera sessili, foliis subulatis setaceis patulis. *Sp. pl.* 1570. *Weis. crypt.* 268. *Fl. dan.* 249.

Sphagnum acaulon trichoides. *Dill. giff.* 229. *Musc.* 251. t. 32. f. 10. *R. syn.* 105. *Hall. hist.* 1727.

Anglis, subulated Phascum.

Habitat in ericetis solo arenoso. III—V.

monta-
num.

2. PHASCUM acaule, anthera sessili, foliis imis patulis oblongis, supremis erectis lanceolatis acuminatis.

Phascum sessile foliis radicalibus obtusis, centralibus ciliatis. *Hall. hist.* 1725. t. 45. f. 3.

Sphagnum acaulon maximum foliis in centro ciliariis. *Hall. it. alp.* 83. *Dill. musc.* 253. t. 32. f. 13.

Anglis, mountain Phascum.

Habitat in rupibus alpinis (terrâ substrata); circa Llanberris in Arvoniam, *Dill. in monte Hart-fell dicta prope Moffat. D. Sheffield.* ☉. V—VIII.

acaulon.

3. PHASCUM acaule, anthera sessili, foliis ovatis acutis conniventibus. *Sp. pl.* 1570. *Fl. dan.* 249. *Weis. crypt.* 266.

Sphagnum acaulon, foliis ovato-lanceolatis, pilo aristatis. *Hall. hist.* 1726.

Sphagnum acaulon bulbiforme, majus. *Dill. musc.* 251. t. 32. f. 11.

Sphagnum acaulon foliis in bulbi formam congestis, majus. *Dill. giff.* 230. *R. syn.* 105.

- β *Sphagnum* acaulon bulbiforme, minus. *Dill. musc.* 252. t. 32. f. 12.

Sphagnum acaulon, foliis in bulbi formam congestis, minus. *Dill. giff.* 230. *R. syn.* 105.

Muscus trichoides acaulos, minor, latifolius. *Vaill. par.* 128. t. 27. f. 2.

Anglis, oval Phascum.

Habitat.

Habitat in ericetis et aggeribus fossarum solo arenoso.
 ☉. I—V.

4. PHASCUM caulescens repens. *Sp. pl.* 1571. *repent.*
 Sphagnum vermiculare repens capsulis intra foliorum
 squamosorum alas minimis. *Dill. musc.* 550. t. 85.
f. 16.
Anglis, creeping Phascum.
 Habitat in arborum truncis, in comitatu Eboracensi.
 D. Richardson. 4. V.

F O N T I N A L I S. 1190.

Anthera operculata, calyptrata, sessilis, perichætio
 inclusa.

1. FONTINALIS foliis complicato-carinatis trifariis *antipyre-*
acutis, antheris lateralibus. *Sp. pl.* 1571. *Weis. tica.*
crypt. 260.

Fontinalis triangularis major complicata e foliorum
 alis capsulifera. *Dill. musc.* 254. t. 33. *f.* 1.

Fontinalis major foliis triangularibus complicatis ca-
 pitulis in foliorum alis sessilibus. *R. syn.* 79.

Hypnum foliis complicatis ovato-lanceolatis, capsulis
 in alis petiolatis ovatis. *Hall. hist.* 1794.

Muscus squamosus foliis acutissimis, in aquis nascens.
Tourn. 554. *Vaill. par.* 140. t. 33. *f.* 5.

Anglis, greater Fontinalis, or Water-moss.

Habitat in fluviis leniter fluentibus et stagnis. 4.
 VI—X.

2. FONTINALIS foliis imbricatis subulato-lanceola- *squamosa.*
tis, antheris lateralibus. *Sp. pl.* 1571.

Fontinalis squamosa tenuis sericea atro-virens. *Dill.*
musc. 250. t. 33. *f.* 3.

Hypnum foliis ovato-lanceolatis, pilo terminatis, cap-
 sulis per alas sessilibus. *Hall. hist.* 1796.

Anglis, squamous Fontinalis.

Habitat in rivulis alpinis in Wallia & septentrionali
 parte Britanniae frequens. 4. VI—X.

Obs. variat foliis subulato-lanceolatis et lanceolatis acutis ;
 nimis affinis præcedenti.

minor.

3. FONTINALIS foliis ovatis concavis trifariis acutis, passim geminis, antheris terminalibus. *Sp. pl.* 1571.

Fontinalis triangularis minor carinata, e cymis capsuliferis. *Dill. musc.* 257. t. 33. f. 2.

Fontinalis minor foliis triangularibus minus complicatis capitulis in summis ramulis sessilibus. *R. syn.* 79.

Hypnum foliis ovato-lanceolatis carinatis, apicibus capsuliferis. *Hall. hist.* 1795.

Anglis, lesser Fontinalis, or Water-moss.

Habitat ad ripam Thamesis circa Londinum, et in Iside prope Oxonium frequens. 4. VIII—X.

pennata.

4. FONTINALIS foliis bifariis patentibus, capsulis lateralibus. *Sp. pl.* 171.

Sphagnum pennatum undulatum; vagina squamosa. *Dill. musc.* 250. t. 32. f. 9.

Hypnum caulibus distichis ramosis, foliis ovatis transversim undosis. *Hall. hist.* 1797.

Anglis, feathered Fontinalis.

Habitat in arborum truncis, in sylvis inter Troutbeck et Ambleside in Westmorlandia. 4. VIII—X.

S P L A C H N U M. 1191.

Anthera incidens apophysi maximæ coloratæ. *Calyptra* caduca.

Stella feminea in distincto individuo.

ampullaceum.

1. SPLACHNUM umbraculo ampullaceo obconico. *Sp. pl.* 1572. *Fl. dan.* 192.

Bryum foliis ovato-lanceolatis, capsula pyriformi terminata cylindro. *Hall. hist.* 1830.

Bryum ampullaceum foliis thymi pellucidis, collo strictiore. *Dill. musc.* 343. t. 44. f. 3.

Bryum erectis gigantinis capitulis, foliis serpylli pellucidis acutis. *R. syn.* 93.

Muscus aureus capillaris minor, capitulis geminatis erectis mutuo incubitu adnatis. *Hist. Ox.* III. 629. f. 15. t. 16. f. 10.

Anglis, purple Splachnum.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Polytrichum montanum et minus capsula quadrangulari. *Dill. giss.* 221. *R. syn.* 90.

Polytrichum aureum medium. *B. pin.* 356.

γ Polytrichum quadrangulare minus, juniperi foliis pilosis. *Dill. musc.* 426. *t.* 54. *f.* 3.

Mnium calyptra villosa, foliis serratis pilo aristatis capsula quadrangula insidente disco. *Hall. hist.* 1836.

Anglis, common Polytrichum, or greater Golden Maiden-hair.

Habitat in sylvis, ericetis, uliginosis et sterilissimis. *u.* V. VI.

Pharmac. Adianti aurei Herba.

nanum. 2. POLYTRICHUM caule simplici, anthera oblonga. Polytrichum parvum, aloes folio serrato capitulis oblongis. *Dill. musc.* 429. *t.* 55. *f.* 7.

Polytrichum capsula cylindrica erecta, surculis simplicibus brevissimis, foliis serrulatis. *Weis. crypt.* 173.

polytri-
choides. Mnium calyptra villosa. *Sp. pl.* 1576. acaulon, foliis serratis capsulis cylindricis erectis. *Hall. hist.* 1837.

β Polytrichum nanum, capsulis subrotundis gateritis, aloes folio non serrato. *Dill. musc.* 428. *t.* 55. *f.* 6.

Polytrichum minus, capsulis subrotundis, calyptra quasi lacera coronatis. *Dill. giss.* 221. *R. syn.* 91.

Mucos coronatus rigidus minor et humilior capitulis villosis brevibus. *Hist. Ox. III.* 630. *f.* 15. *t.* 7. *f.* 7.

Anglis, dwarf Polytrichum.

Habitat in ericetis. VIII—XI.

alpinum. 3. POLYTRICHUM caule ramoso, pedunculis subterminalibus.

Polytrichum caule ramosissimo pedunculis terminalibus. *Sp. pl.* 1573. *Fl. dan.* 296.

Polytrichum alpinum ramosum, capsulis e summitate ellipticis. *Dill. musc.* 427. *t.* 55. *f.* 4.

Bryum foliis serratis, capsulis ovatis, basi turbinatis. *Hall. hist.* 1800. *t.* 46. *f.* 6.

Polytrichum

β Polytrichum caule ramosissimo, pedunculis axillaribus. urnigerum. *Sp. pl.* 1573.

Polytrichum ramosum fetis ex alis urnigeris. *Dill. musc.* 427. t. 55. f. 5.

Muscus erectius juniperi folio ramosus. *Vaill. par.* 131. t. 28. f. 13.

Bryum caule ramoso, ex alis florigerum, foliis ferratis, capsulis cylindricis. *Hall. hist.* 1801.

Anglis, alpine Polytrichum.

Habitat in ericetis montosis in boreali parte Angliæ, et in montibus Wallicis et Scoticis frequens. 4. VI—VIII.

Obs. Pedunculi in α et β non ex apice surculorum sed paulo infra oriuntur.

4. POLYTRICHUM caule ramoso, antheris subsessilibus subterminalibus, calyptra sursum pilosa. *striatum.*

Polytrichum bryi ruralis facie, capsulis sessilibus majus. *Dill. musc.* 430. t. 55. f. 8.

Polytrichum capsulis subrotundis, pediculis brevissimis insidentibus, calyptra striata, arboreum ramosum majus. *Dill. giss.* 222. *R. syn.* 91.

Bryum antheris sessilibus sparsis, calyptris striatis sursumve pilosis. *Sp. pl.* 1579. *Fl. dan.* 537. f. 3.

Bryum fetis brevissimis alaribus, calyptris villosis conicis. *Hall. hist.* 1799.

Muscus apocarpos arboreus ramosus. *R. hist.* III. 40. *Vaill. par.* 129. t. 25. f. 5, 6.

β Polytrichum bryi ruralis facie, capsulis sessilibus, minus. *Dill. musc.* 431. t. 55. f. 9.

Polytrichum capsulis subrotundis, pediculis brevissimis insidentibus, calyptra striata, arboreum et terrestre minus ramosum et breve. *Dill. giss.* 222. *R. syn.* 91.

Muscus capillaceus minimus acaulos, calyptra striata. *Tourn.* 552. *Vaill. par.* 130. t. 27. f. 10.

γ Polytrichum capsulis sessilibus, foliis brevibus rectis carinatis. *Dill. musc.* 432.

δ Polytrichum capillaceum crispum, calyptris acutis pilosissimis. *Dill. musc.* 433. t. 55. f. 11.

Polytrichum capsulis oblongo-rotundis, calyptris pilosissimis. *R. syn.* 91.

Bryum setis brevissimis alaribus, calyptris cylindricis pilosissimis. *Hall. hist.* 1798.

Muscus capillaceus minimus, calyptra villosa. *Vaill. par.* 130. t. 27. f. 9.

Anglis, striated *Polytrichum*.

Habitat in arboribus et rupibus. II—V.

M N I U M. 1593.

Anthera operculata, *Calyptra* lævis.

Fem. capitulum nudum, pulverulentum, remotum.

pellucidum.

1. MNIUM caule simplici foliis ovatis. *Sp. pl.* 1574. *Fl. dan.* 300. *Weis. crypt.* 161.

Mnium caule simplicissimo, foliis ovato-lanceolatis, seta fungifera foliosa. *Hall. hist.* 1853. t. 45. f. 8.

Mnium serpylli foliis tenuibus et pellucidis. *Dill. musc.* 232. t. 31. f. 2.

Mnium minus non ramosum, angustioribus et pellucidis foliis. *R. syn.* 78.

β *Mnium* minimum non ramosum angustissimis et pellucidis foliis. *R. syn.* 78.

Anglis, pellucid *Mnium*.

Habitat in sylvis et umbris udis. ☉. III. IV.

androgynum.

2. MNIUM caule ramoso, androgynum. *Sp. pl.* 1574. *Fl. dan.* 299. *Weis. crypt.* 163.

Mnium caule simplici, cauliculis rosiferis. *Hall. hist.* 1854.

Mnium perangustis et brevibus foliis. *Dill. gifs.* 214. *musc.* 230. t. 31. f. 1. *R. syn.* 78.

Muscus coronatus humilis minimus, foliolis, pediculis et capitulis minimis erectis. *Hist. Ox.* III. 631.

f. 15. t. 7. f. 2.

Anglis, upright *Mnium*.

Habitat in sylvis, ericetis et muris. ☉. III. IV.

palustre.

3. MNIUM caule dichotomo, foliis subulatis. *Sp. pl.* 1574. *Weis. crypt.* 165.

Mnium caulibus rubiginosis bipartitis, foliis lanceolatis

tis

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Jungermannia foliis amplexicaulibus subrotundis.
Hall. hist. 1861. *Weis. crypt.* 123.

Anglis, purple Mnium.

Habitat in rivulis alpinis et palustribus; in fossulis in ericetis turfosis frequentissime cum capitulis femininis observavi. 4. III—VII.

B R Y U M. 1594.

Anthera operculata; *Calyptra* lævis. *Filamentum* e tuberculo terminali ortum.

Stella feminea in distincto individuo.

* *Antheris sessilibus.*

apocar-
cum.

1. BRYUM antheris sessilibus terminalibus, calyptra minima. *Sp. pl.* 1115. *Fl. dan.* 480.

Sphagnum nodosum hirsutum incanum. *Dill. musc.* 246. t. 32. f. 5.

Sphagnum cauliferum et ramosum saxatile hirsutum, capitulis virentibus. *R. syn.* 105.

Hypnum caulibus ramosis, foliis lanceolatis hirsutis, operculo aristato. *Hall. hist.* 1793.

♣ *Sphagnum subhirsutum, obscure virens, capsulis rubellis.* *Dill. musc.* 245. t. 32. f. 4.

Sphagnum cauliferum et ramosum saxatile, hirsutum virens, capitulis obscuris rubris. *Dill. giss.* 229. *R. syn.* 104.

Muscus apocarpos hirsutus, saxi adnascens, capitulis obscure rubris. *Vaill. par.* 129. t. 27. f. 15.

Anglis, sessile Bryum.

Habitat in rupibus et arboribus. XII—IV.

** *Antheris pedunculatis erectis.*

pomifor-
me.

2. BRYUM antheris erectis sphaericis. *Sp. pl.* 1580.

Bryum capillaceum capsulis sphaericis. *Dill. musc.* 339. t. 44. f. 1.

Bryum foliis subulatis, setis verticalibus, capsulis sphaericis. *Hall. hist.* 1803.

Bryum trichoides virens, erectis majusculis capitulis maliformibus. *R. syn.* 97.

Muscus trichoides minimus sericeus capillaceus, capitulis

capitulis sphaericis. *Hist. Ox. III. 628. f. 15. t. 6. f. 6.*
Anglis, apple Bryum.
 Habitat in ericetis; on Hampstead Heath. *R. syn.*
 ©. III. IV.

2. BRYUM antheris erectis obovatis; calyptra subula-pyriforme.
 ta, turculis acaulibus, foliis ovatis muticis. *Sp. pl.*
 1589. *Fl. dan. 537. Weis. crypt. 183.*
Bryum caule brevissimo, foliis ovato-lanceolatis, cap-
sulis ovatis, operculo brevi. Hall. hist. 1832.
Bryum serpyllifolium pellucidum, capsulis pyriform-
mibus. Dill. musc. 345. t. 44. f. 6.
Bryum parvum erectis pyriformibus majusculis capi-
tulis, foliolis serpylli pellucidis. R. syn. 93.
Muscus coronatus humilis stellaris, foliis latiusculis,
capitulis pyriformibus erectis turgidiusculis. Hist.
Ox. III. 631. f. 15. t. 7. f. 16.
Anglis, pear Bryum.
 Habitat in ericetis et ad aggeres viarum. ©. III. IV.

3. BRYUM antheris obliquis subrotundis; caule sim-fontanum.
 plici geniculis inflexo.
Bryum palustre, scapis teretibus stellatis, capsulis mag-
nis subrotundis. Dill. musc. 340. t. 44. f. 2.
Mnium caule simplici geniculis inflexo. Sp. pl.
 1574. *Fl. dan. 298.*
Mnium caulibus teretibus erectis ex vertice multifidis.
Hall. hist. 185 f.
Muscus palustris adianto aureo affinis, scapis tenui-
bus foliolis brevibus. R. syn. 98.
Anglis, marsh Bryum.
 Habitat in palustribus turfosis et ad fontes. 4. V—
 VIII.

4. BRYUM anthera erecta oblonga minori, calyptris *extincto-*
 laxis aequalibus. *Sp. pl. 1581. Weis. crypt. 185. rium.*
Bryum caule brevissimo, calyptra cylindrica, longe
aristata. Hall. hist. 1829.
Bryum calyptra extinctorii forma minus. Dill. musc.
 349. *t. 45. f. 8.*
Bryum erectis capitulis, calyptra conica foliis serpylli
pellucidis angustioribus. R. syn. 92.

Muscus

Muscus capillaceus minimus calyptra longa conoide nitida. *Vaill. par. 137. t. 26. f. 1.*

♁ Bryum calyptra extensorii figura, majus et ramosum. *Dill. musc. 350. t. 45. f. 9.*

Anglis, extinguisher Bryum.

Habitat in ericetis montosis et rupibus circa Llanberris frequens; in monte Ingleborough et alibi in comitatu Eboracensi. ☉. X—VIII.

subula-
tum.

5. BRYUM antheris erectis subulatis, furculis acaulibus. *Sp. pl. 1581. Neck. dilic. Gallob. 456. Weis. crypt. 187.*

Bryum caule brevissimo, foliis lanceolatis, capsulis longissimis, operculo prælongo. *Hall. hist. 1827.*

Bryum capsulis longis subulatis. *Dill. musc. 350. t. 45. f. 10.*

Bryum erectis longis et acutis falcatis capitulis, calyptra subfusca, foliis serpylli pellucidis. *R. syn. 92.*

Muscus capillaris, corniculis longissimis incurvis. *R. hist. III. 34. Vaill. par. 133. t. 25. f. 8.*

Anglis, subulated Bryum.

Habitat in aggeribus terrenis humidiusculis et sylvis. III,

rurale.

6. BRYUM antheris erectiusculis, foliis piliferis recurvis. *Sp. pl. 1581.*

Bryum rurale unguiculatum hirsutum elatius et ramosius. *Dill. musc. 352. t. 45. f. 12.*

Bryum majus erectis falcatis capitulis, foliis latiusculis extantibus, in pilum canescentem desinentibus. *R. syn. 94.*

Hypnum caulibus erectis, foliis lanceolatis pilo fluente terminatis, capsulis erectis longe rostratis. *Hall. hist. 1789.*

Muscus erectus hirsutis capitulis longis acutis unguiculis avicularum similibus, vulgatissimus. *Hist. Ox. III. 626. f. 15. t. 6. f. 1.*

Muscus capillaris tectorum densis cæspitibus nascens capitulis oblongis, foliis in pilum oblongum desinentibus. *Vaill. paris. 133. t. 25. f. 3.*

Anglis, hairy Bryum.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

cronato. *Sp. pl.* 1584. *Weis. crypt.* 191. *Fl. dan.* 537.

Bryum foliis ovato-lanceolatis, capsulis erectis ovatis aristatis. *Hall. hist.* 1833.

Bryum exiguum, creberrimis capsulis rufis. *Dill. musc.* 347. t. 45. f. 7.

Bryum exiguum, erectis parvis subrotundis creberrimis capitulis rufis, foliolis serpylli angustis pellucidis. *R. syn.* 94.

Muscus coronatus minimus, foliis et capitulis oblongis in pediculis brevissimis. *Hist. Ox. III.* 631. f. 15. t. 7. f. 18.

β Bryum parvum erectis subrotundis majusculis capitulis subfuscis, foliis serpylli pellucidis. *R. syn.* 93.

Anglis, brown Bryum.

Habitat in pratis, pascuis et ericetis. ☉. III.

viridulum.

10. BRYUM antheris erectis ovatis foliis lanceolatis acuminatis imbricato-patulis. *Sp. pl.* 1584.

Bryum foliis capillaribus, capsulis ovatis aristatis. *Hall. hist.* 1809.

Bryum capillaceum breve, pallide et læte virens, capsulis ovatis. *Dill. musc.* 380. t. 48. f. 43.

Bryum trichoides exile, erectis capitulis in pediculis brevissimis. *Cat. giss.* 224. *R. syn.* 97.

Anglis, green Bryum.

Habitat in pratis, aggeribus fossarum et ad vias. ☉. XII.—III.

paludosum.

11. BRYUM acaulon, foliis setaceis, antheris obtusissimis patulis. *Sp. pl.* 1584. *Weis. crypt.* 194. *Neck. dillic. Gallob.* 454.

Bryum foliis capillaribus, cauliculo sessili, capsulis ovatis aristatis. *Hall. hist.* 1810.

Bryum trychoides acaulon palustre minimum, setis et capsulis brevissimis. *Dill. musc.* 387. t. 49. f. 53.

Anglis, dwarf Bryum.

Habitat in palustribus arenosis et turfosis in comitatu Eboracensi. D. Richardson. ☉. III—V.

12. BRYUM

12. BRYUM antheris erectis oblongis, foliis linearibus incurvum. carinatis integerrimis incurvis.

Anglis, incurved Bryum.

Habitat in pascuis et rupibus circa Llanberris. D. Davies. 4. VI—X.

Desc. Caulis erectus simplex, uncialis; Folia alterna erectiuscula, incurva, linearia, integerrima, acuta, carinata, nuda; Pedunculus terminalis, solitarius, erectus, uncialis; Calyptra conica, acuta, pallida. Anthera erecta oblonga s. cylindracea; ore denticulato-ciliata; Operculum conicum acutiusculum.

13. BRYUM antheris erectiusculis, pedunculis subso- undulati-
litariis, foliis lanceolatis carinatis undulatis patentibus ferratis. *Sp. pl.* 1582. *Weis. crypt.* 197. *Fl. dan.* 477.

Bryum foliis lanceolatis ferratis, capsulis cylindricis inclinatis aristatis. *Hall. hist.* 1823.

Bryum phyllitidis folio rugoso acuto, capsulis incurvis. *Dill. musc.* 360. t. 46. f. 18.

Bryum capitulis oblongis rubentibus, foliis oblongis angustis pellucidis rugosis. *R. syn.* 95.

Anglis, undulated Bryum.

Habitat in aggeribus umbrosis et sylvis. 4. XII—VI.

14. BRYUM antheris erectiusculis, operculo arcuato, glaucum. foliis erectis imbricatis, furculis ramosis. *Sp. pl.* 1582.

Bryum albidum et glaucum fragile majus, foliis erectis, setis brevibus. *Dill. musc.* 362. t. 46. f. 20.

Bryum trichoides, erectis capitulis, albidum fragile. *Dill. giss.* 225. *R. syn.* 97.

Hypnum caulibus erectis teretibus, foliis imbricatis fragilibus, capsulis inclinatis aristatis. *Hall. hist.* 1785. *Weis. crypt.* 208.

Anglis, white Bryum.

Habitat in ericetis et pascuis montosis. 4. IX—XI.

15. BRYUM antheris erectis, operculo aciformi, foliis acicularibus erectis subsecundis. *Sp. pl.* 1583.

Bryum

Bryum montanum hemiheterophyllum, operculis acutis. *Dill. musc.* 366. t. 46. f. 25.

Bryum hypnoides erectum montanum, erectis capitulis acutis. *R. syn.* 94.

β *Bryum hypnoides aquaticum*, calyptris nigris acutis. *Dill. musc.* 367. t. 46. f. 26.

Bryum hypnoides repens aquaticum, erectis capitulis acutis. *R. syn.* 94.

Hypnum caulibus teretibus, foliis imbricatis lanceolatis, capsulis erectis ovatis aristatis. *Hall. hist.* 1781.

Anglis, mountain *Bryum*.

Habitat in rupibus humidis et ad ripas rivulorum, in boreali parte Angliæ frequens, et in Wallia. 4. V—X.

Obs. in α, caules et folia erecti; in β caules decumbentes et folia patentiuscula.

hypnoides. 16. **BRYUM** antheris erectis, caule erectiusculo, ramis lateralibus brevibus fertilibus. *Sp. pl.* 1585. *Weis. crypt.* 211.

Bryum hypnoides hirsutie canescens vulgare. *Dill. musc.* 368. t. 47. f. 27.

Bryum trichoides erectis capitulis, lanuginosum. *R. syn.* 97.

β *Bryum hypnoides*, hirsutie virescens, fasciculare alpinum. *Dill. musc.* 370. t. 47. f. 28.

γ *Bryum hypnoides alpinum*, setis et capsulis exiguis. *Dill. musc.* 370. t. 47. f. 29.

Hypnum caulibus teretibus, foliis pilo aristatis, setis brevissimis. *Hall. hist.* 1782.

δ *Bryum hypnoides alpinum* operculis obtusis. *Dill. musc.* 371. t. 47. f. 30.

ε *Bryum hypnoides*, ericæ facie, capsulis barbatis, alpinum. *Dill. musc.* 371. t. 47. f. 31.

Bryum hypnoides, capitulis plurimis erectis non lanuginosum. *R. syn.* 458.

ζ *Bryum hypnoides polycephalon*, lanuginosum montanum. *Dill. musc.* 372. t. 47. f. 32.

Bryum hypnoides, capitulis plurimis erectis, lanuginosum. *Dill. musc.* 372. t. 47. f. 32.

Hypnum

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Bryum trichoides exile pallidum, erectis capitulis e furculis annotinis egredientibus. *R. syn.* 96.

Mnium antheris erectis, operculis filiformibus longitudine antherarum. *Sp. pl.* 1575.

Hypnum cauliculis teretibus, foliis lanceolatis repandis, capsulis erectis cylindricis aristatis. *Hall. hist.* 1788. *Weis. crypt.* 206.

β Bryum stellare lacustre, foliis rubris capillaceis. *Dill. musc.* 382. t. 48. f. 45.

Anglis, setaceous Bryum.

Habitat in aggeribus fossarum, sepibus et muris. ☉. III.

unguiculatum.

20. BRYUM antheris erectis oblongis, operculo acuminato obliquo, foliis lineari-lanceolatis carinatis.

Bryum capsulis urceolatis operculo longo obliquo, foliis lanceolatis tortilibus, cauliculis erectis. *Schreb. spic.* 78.

Bryum unguiculatum et barbatum, furculis in summitate crassioribus. *Dill. musc.* 383. t. 48. f. 47.

Bryum angustis viridibus foliis capitulis erectis, brevibus pediculis insidentibus, calyptra falcata vel avium unguiculos referente. *R. syn.* 96.

Muscus capillaris, capitulis erectis lageniformibus. *Buxb. cent.* 11. t. 2. f. 9.

Anglis, birds-claw Bryum.

Habitat in muris et arenosis. ☉. III. IV.

purpureum.

21. BRYUM antheris erectiusculis ovatis, caule bifido, axillis pedunculiferis, foliis lanceolatis acutis carinatis.

Bryum foliis lanceolatis subulatis, capsulis erectis operculis conicis. *Hall. hist.* 1816.

Bryum tenue stellatum, setis purpureis. *Dill. musc.* 386. t. 49. f. 51.

Bryum perangustis foliis et cauliculis, foliis crebrioribus et circa summitates magis congestis, capitulis erectis e furculis annotinis egredientibus. *R. syn.* 99.

Mnium caule dichotomo, axillis pedunculiferis, anthera erecta, foliolis carinatis. *Sp. pl.* 1075.

α Bryum tenue barbatum, foliis angustioribus et rarioribus. *Dill. musc.* 384. t. 48. f. 49.

Bryum

Bryum foliis lanceolatis imbricatis capsulis cylindricis aristatis. *Hall. hist.* 1817.

Bryum perangustis foliis et cauliculis, foliis rarioribus, cinctis, capitulis erectis, e furculis annotinis egredientibus. *Cat. giss.* 225. *R. syn.* 99.

7 Bryum foliis lanceolatis, laxe imbricatis, capsulis cylindricis obliquis, operculis conicis. *Hall. hist.* 1818.

Bryum polytrichoides palustre, setis longioribus rubris sericeis. *Dill. musc.* 387. t. 49. f. 52.

Bryum perangustis crebrioribus foliis, capitulis erectis longiusculis, pediculis e furculis annotinis nascentibus. *R. syn.* 99.

Anglis, purple Bryum.

Habitat in muris, ericetis et pascuis glareosis. ☉. III. IV.

22. BRYUM antheris erectiusculis, pedunculis aggregato-scopariatis, foliis secundis recurvatis, caule declinato. *um.* *Sp. pl.* 1582.

Bryum reclinatum, foliis falcatis, scoparium effigie. *Dill. musc.* 357. t. 46. f. 16.

Bryum erectis capitulis, angustifolium, caule reclinato. *Dill. giss.* 222. *R. syn.* 95.

Hypnum foliis falcatis heteromallis, vaginis multifloris. *Hall. hist.* 1777. *Weis. crypt.* 213.

Muscus coronatis medius, foliis tenuissimis pallidis longioribus, capitulis erectis acutis. *Hist. Ox.* III. 630. f. 15. t. 7. f. 11.

Anglis, broom Bryum.

Habitat in ericetis et sylvis. ☿. VI. VII.

23. BRYUM antheris erectis ovalibus, pedunculis lateralibus foliis setaceis secundis.

Bryum foliis mollibus subulatis, setis brevissimis alaribus, capsulis ovatis. *Hall. hist.* 1202. t. 46. f. 8.

Anglis, lateral Bryum.

Habitat in sylvis et ad ripas rivulorum in Scotia; in monte Hart-fell prope Moffat. D. Sheffield. ☿. VII. IX.

Desc. Caules *cæspitosi*, *unciales* et *triunciales*, *erectiasculi* *purpurascens*, *basi tomentosi*; *tomento fulvo*, *subsimplici*. *Folia numerosa*, *secunda*, *setacea*, *viridissima*, *mollia*, *longa*. *Pedunculi laterales*, *axiliares*, *erecti*, *breves*, *filiformes*, *purpurei*. *Calyptra conica*, *acuta*, *pallida*, *caduca*. *Anthera erecta*, *subrotunda*, *ciliata*, *fulva*. *Operculum conicum*, *brevissimum*, *rubescens*.

heteromallum.

24. BRYUM antheris erectis ovatis, pedunculis terminalibus, foliis setaceis secundis.

Bryum antheris erectis, foliis setaceis secundis. *Sp. pl.* 1583. *Fl. dan.* 479.

Bryum heteromallum. *Dill. musc.* 371. t. 47. f. 37.

Bryum trichoides reclinatis cauliculis, capitulis erectis acutis. *R. syn.* 96.

Muscus trichoides, foliis capillaceis, capitulis minoribus. *Hist. Ox.* III. 628. f. 15. t. 6. f. 5.

β Bryum pilosum interrupte falcatum. *Dill. musc.* 376. t. 47. f. 38.

Anglis, heath Bryum.

Habitat in ericetis. ☉. III—V.

flexuosum.

25. BRYUM antheris erectis oblongis, pedunculis terminalibus flexuosis, foliis setaceis erectis.

Bryum antheris erectis foliis setaceis, pedunculis flexuosis. *Sp. pl.* 1583.

Bryum foliis capillaceis, capsulis cylindricis, operculis aristatis. *Hall. hist.* 1804.

Bryum pilosum molle, setis intortis. *Dill. musc.* 373. t. 47. f. 33.

Bryum trichoides capitulis erectis, pediculis intortis tenuibus virentibus. *Dill. giss.* 225. *R. syn.* 97.

Anglis, waved Bryum.

Habitat in rupibus circa Llanberris. *Dill.* 4. IV—VI.

acutum.

26. BRYUM antheris erectis subrotundis, foliis setaceis erectis, furculis ramosis acutis.

Bryum pilosum sphagni subulatū facie. *Dill. musc.* 374. s. 47. t. 34.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Bryum trichoides exile erectis capitulis in pediculis longioribus rubris. *Dill. giss.* 224. *R. syn.* 97.

Mnium foliis arefactione revolutis. *Sp. pl.* 1576.

Hypnum foliis capillaceis congestis, crispulentibus, capsulis ovato-cylindricis, brevissime ciliatis. *Hall. hist.* 1786. *Weis. crypt.* 207.

Muscus coronatus minimus, capillaceis foliis, capitulis oblongis. *Hist. Ox. III.* 631. *f.* 15. *t.* 7. *f.* 19.

β Bryum cirratum, setis et capsulis brevioribus et pluribus. *Dill. musc.* 378. *t.* 48. *f.* 41.

Bryum trichoides erectis sublongis capitulis, extremitatibus per siccitatem stellatis. *R. syn.* 98.

γ Bryum perangustis et brevissimis foliis, extremitatibus stellatis. *Dill. giss.* 226. *R. syn.* 98.

Anglis, starry Bryum.

Habitat in sylvis et ericetis montosis. 4. III—VII.

tortuosum.

30. BRYUM antheris erectis, foliis setaceis imberbibus arefactione retortis. *Syst. nat.* 702. *Sp. pl.* 1583.

Bryum cirratum, setis et capsulis longioribus. *Dill. musc.* 377. *t.* 48. *f.* 40.

Bryum trichoides longifolium, crassiusculis cauliculis, capitulis erectis aduncis acutis. *R. syn.* 68.

Hypnum foliis subulatis cincinnatis, capsulis cylindricis, longissime ciliatis. *Hall. hist.* 1787. *t.* 45. *f.* 2.

Muscus alpinus cirrhosus s. crinum tortorum instar crispatus. *Sch. it. alp.* 65. *t.* 19. *f.* 5.

Anglis, frizzled Bryum.

Habitat in ericetis et rupibus montosis. 4. VIII—X.

*** *Antheris nutantibus.*

simplex.

31. BRYUM antheris oblongis nutantibus, foliis subulatis, surculo simplicissimo medio pedunculifero. *Sp. pl.* 1587. *Schreb. spicil.* 81.

Bryum cauliculis cernuis, foliis capillaceis heteromallis, capsulis curvulis cylindricis, operculis conicis. *Hall. hist.* 1808.

Bryum trichoides, capsulis rubris cernuis. *Dill. musc.* 390. *t.* 50. *f.* 59.

Anglis,

Anglis, simple Bryum.

Habitat in locis gramineis arenosis. ☉. III. IV.

32. BRYUM antheris subrotundis, pedunculis reflexis, *pulvinatum.*
foliis piliferis. *Sp. pl.* 1586. *Weis. crypt.* 201.

Bryum foliis lanceolatis pilo terminatis, capsulis pendulis ovatis aristatis. *Hall. hist.* 1822.

Bryum orbicularis pulvinatum hirsutie canescens, capsulis immersis. *Dill. musc.* 395. t. 50. f. 65.

Bryum trichoides hirsutie canescens, capitulis subrotundis reflexis, in perbrevis pediculis. *Dill. giss.* 226. *R. syn.* 100.

Muscus capillaceus lanuginosus minimus. *Vaill. paris.* 133. t. 29. f. 2.

Anglis, grey Bryum.

Habitat in muris et tectis. 4. III. IV.

33. BRYUM antheris nutantibus turbinatis, foliis seta-*aureum.*
ceis.

Bryum foliis capillaceis capsulis pyriformibus pendulis. *Hall. hist.* 1813. t. 45. f. 7. operculo brevissimo rotundata. *Schreb. spicil.* 81.

Bryum trichoides aureum, capitulis pyriformibus nutantibus. *Deer. cat. Nottingh. App. Dill. Musc.* 391. t. 50. f. 60.

Mnium antheris pendulis turbinatis, stipite filiformi, pyriformibus femineis setiferis. *Sp. pl.* 1576. *forme.*

Anglis, golden Bryum.

Habitat in rupibus *Parci* Nottinghamensis. D. Deering. 4. IV—VI.

34. BRYUM antheris pendulis oblongis, foliis lan-*sericium.*
ceolatis acutis carinatis.

Bryum foliis lanceolatis subulatis, capsulis urceolatis cernuis, operculo conico. *Schreb. spicil.* 81.

Bryum trichoides læte virens, capitulis cernuis oblongis. *R. syn.* 100. *Dill. musc.* 391. t. 50. f. 61.

Anglis, silky Bryum.

Habitat in ericetis montosis, prope Woolwich; *Dill. circa Llanberris.* D. Davies. in monte Ingleborough et alibi in comitatu Eboracensi. 4. III—VII.

Bryum trichoides exile erectis capitulis in pediculis longioribus rubris. *Dill. giss.* 224. *R. syn.* 97.

Mnium foliis arefactione revolutis. *Sp. pl.* 1576.

Hypnum foliis capillaceis congestis, crispescentibus, capsulis ovato-cylindricis, brevissime ciliatis. *Hall. hist.* 1786. *Weis. crypt.* 207.

Muscus coronatus minimus, capillaceis foliis, capitulis oblongis. *Hist. Ox. III.* 631. *f.* 15. *t.* 7. *f.* 19.

β Bryum cirratum, setis et capsulis brevioribus et pluribus. *Dill. musc.* 378. *t.* 48. *f.* 41.

Bryum trichoides erectis sublongis capitulis, extremitatibus per siccitatem stellatis. *R. syn.* 98.

γ Bryum perangustis et brevissimis foliis, extremitatibus stellatis. *Dill. giss.* 226. *R. syn.* 98.

Anglis, starry Bryum.

Habitat in sylvis et ericetis montosis. 4. III—VII.

tortuosum.

30. BRYUM antheris erectis, foliis setaceis imberbibus arefactione retortis. *Syst. nat.* 702. *Sp. pl.* 1583.

Bryum cirratum, setis et capsulis longioribus. *Dill. musc.* 377. *t.* 48. *f.* 40.

Bryum trichoides longifolium, crassiusculis cauliculis, capitulis erectis aduncis acutis. *R. syn.* 98.

Hypnum foliis subulatis cincinnatis, capsulis cylindricis, longissime ciliatis. *Hall. hist.* 1787. *t.* 45. *f.* 2.

Muscus alpinus cirrhosus s. crinum tortorum instar crispatus. *Sch. it. alp.* 65. *t.* 19. *f.* 5.

Anglis, frizzled Bryum.

Habitat in ericetis et rupibus montosis. 4. VIII—X.

*** Antheris nutantibus.

simplex.

31. BRYUM antheris oblongis nutantibus, foliis subulatis, surculo simplicissimo medio pedunculifero. *Sp. pl.* 1587. *Schreb. spicil.* 81.

Bryum cauliculis cernuis, foliis capillaceis heteromallis, capsulis curvulis cylindricis, operculis conicis. *Hall. hist.* 1808.

Bryum trichoides, capsulis rubris cernuis. *Dill. musc.* 390. *t.* 50. *f.* 59.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

cæspitici-um. 35. BRYUM antheris pendulis, foliis lanceolatis, acuminato-setaceis, pedunculis longissimis. *Sp. pl.* 1586.

Bryum pendulum ovatum cæspiticium et pilosum, seta bicolori. *Dill. musc.* 396. t. 50. f. 66.

Bryum trichoides capitulis reflexis, pediculis ima medietate rubris, summa luteo-virentibus. *R. syn.* 100.

Hypnum foliis ovato-lanceolatis aristatis patulis, capsulis ovatis obtusis pendulis. *Hall. hist.* 1790. *Weis. crypt.* 203.

Muscus trichoides, capitulo parvo reflexo, pediculo ima medietate rubro, summa luteo-viridi. *Hist. Ox. III.* 629. f. 15. t. 6. f. 15.

Muscus capillaceus minimus, capitulo nutante, pediculo purpureo. *Vaill. paris.* 134. t. 29. f. 7.

Anglis, matted Bryum.

Habitat in muris et glareosis. 4. III. IV.

capillare. 36. BRYUM antheris pendulis, foliis ovatis fetiferis carinatis, pedunculis longissimis. *Sp. pl.* 1586.

Bryum foliis latiusculis congestis, capsulis longis nutantibus. *Dill. musc.* 398. t. 50. f. 67.

Bryum capitulis reflexis, foliis latiusculis congestis. *Dill. giss.* 227. *R. syn.* 100.

Hypnum foliis ovato-lanceolatis aristatis, capsulis ovatis pendulis, operculis conicis. *Hall. hist.* 1791. *Weis. crypt.* 205.

Muscus capillaris, foliis latiusculis congestis, capitulis longis reflexis. *Hist. Ox. III.* 629. f. 15. t. 6. f. 19.

Muscus capillaceus major, capitulis crassioribus cylindraceis nutantibus. *Vaill. paris.* 134. t. 24. f. 6.

Anglis, capillary Bryum.

Habitat in sylvis, ericetis, solo glareoso. 4. IV. V.

Obs. nimis affinis præcedenti.

bygrometricum. 37. BRYUM antheris nutantibus turbinatis, calyptra tetragona, foliis ovatis conniventibus.

Bryum bulbiforme aureum, calyptra quadrangulâri, capsulis pyriformibus nutantibus. *Dill. musc.* 407. t. 52. f. 75.

Bryum

Bryum aureum, capitulis reflexis pyriformibus, calyptra quadrangulati, foliis in bulbi formam congestis.

R. syn. 101.

Mnium acaule anthera nuda, calyptrae reflexa tetragona. *Sp. pl.* 1575. *Weis. crypt.* 147.

Mnium foliis ovato-lanceolatis conniventibus, capsulis ovatis reflexis. *Hall. hist.* 1852.

Muscus capillaris s. Adiantum aureum minus. *Ger. em.* 1559.

Polytrichon aureum minus. *Park.* 1052.

Anglis, yellow Bryum.

Habitat in sylvis, ericetis, hortis et ad seminas. ☉. III—V.

38. BRYUM antheris pendulis, furculis cylindricis imbricatis laevibus. *Sp. pl.* 1586. *Weis. crypt.* 199. *argenteum.*

Bryum caulibus teretibus, capsulis ovatis acuminatis pendulis. *Hall. hist.* 1821.

Bryum pendulum julaceum argenteum et sericeum. *Dill. musc.* 392. t. 50. f. 62.

Bryum capitulis subrotundis reflexis, cauliculis teretibus argenteis. *Dill. giss.* 226. *R. syn.* 100.

Muscus minimus e viridi argenteus, capitulis oblongis. *Hist. Ox. III.* 629. f. 15. t. 6. f. 17.

β Bryum pendulum furculis teretibus viridibus. *Dill. musc.* 314. t. 50. f. 69.

Anglis, silver Bryum.

Habitat in muris, tectis, apricis et rupibus. ♃. X—II.

39. BRYUM antheris pendulis subglobosis, foliis acutis alternis. *Sp. pl.* 1587. *carneum.*

Bryum foliis lanceolatis rarioribus, capsulis ovatis pendulis. *Hall. hist.* 1834. t. 45. f. 6.

Bryum lanceolatum pellucidum capsulis rotundis pendulis carneis. *Dill. musc.* 400. t. 50. f. 69.

Bryum nitidum, foliis serpylli pellucidis angustioribus, reflexis capitulis, carnei coloris. *R. syn.* 102.

β Bryum nitidum perangustis serpyllinis pellucidis foliis, reflexis capitulis subrotundis virentibus, longioribus pediculis insidentibus. *R. syn.* 102.

Angles, shining Bryum.

Habitat in pascuis, et locis udis. ♃. III. IV.

40. BRYUM

alpinum. 40. BRYUM antheris pendulis oblongis, foliis imbricatis ovatis acutis carinatis, furculis ramosis, axillis pedunculiferis. *Fl. Angl.* 415. *Lin. Mant.* 309.

Bryum hypnoides pendulum sericeum, coma insigni atro-rubente. *Dill. musc.* 394. t. 50. f. 64.

Anglis, alpine Bryum.

Habitat in rupibus tenui terra tectis, prope Castellum quoddam dirutum, non longe a Llanberris. *Dill. et in boreali parte Britanniae.* 4. IV—VI.

annotinum.

41. BRYUM antheris nutantibus oblongis, pedunculis subradicalibus, foliis ovatis acuminatis pellucidis.

Bryum annotinum lanceolatum pellucidum, capsulis oblongis pendulis. *Dill. musc.* 399. t. 50. f. 68.

Mnium foliis ovatis acuminatis pellucidis, pedunculis radicalibus, anthera nutante. *Sp. pl.* 1576.

Weis. crypt. 151.

Anglis, wood Bryum.

Habitat in sylvis et umbrosis humidis. 4. III. IV.

triquestrum.

42. BRYUM antheris pendulis oblongis, foliis trifariis patulis subulato-lanceolatis carinatis, furculis ramosis.

Bryum annotinum palustre, capsulis ventricosus pendulis. *Dill. musc.* 404. t. 51. f. 72.

Bryum nitidum foliis serpylli angustioribus, majus. *Dill. giff.* 228. *R. syn.* 102.

Mnium foliis trifariis patulis subulato-lanceolatis carinatis. *Sp. pl.* 1578.

Mnium caulibus prælongis rubiginosis, foliis ovato-lanceolatis, capsulis ovato pendulis. *Hall. hist.* 1849.

Muscus capillaris major et elatior, capitulis longis et obtusis deorsum reflexis et veluti pendulis præaltis pediculis rubris. *Hist. Ox.* III. 629. f. 15. t. 6. f. 20.

β Bryum lanceolatum bynum, setis et capsulis longis pendulis. *Dill. musc.* 405. t. 51. f. 73.

Bryum nitidum foliis serpylli pellucidis angustis, capitulis tumidis nutantibus, præaltis pediculis e furculis annotinis egredientibus. *R. syn.* 102.

Mnium

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- serpyllifolium.* 45. BRYUM antheris pendulis oblongis, pedunculis subaggregatis, foliis patentibus pellucidis. Mnium pedunculis aggregatis, foliis patentibus pellucidis. *Sp. pl.* 1577.
- punctatum.* Mnium pedunculis aggregatis, foliis obovatis integerrimis obtusis punctatis. *Sp. pl.* 1577. *Weis. crypt.* 153.
Mnium foliis ovatis integerrimis, capsulis pendulis ovatis aristatis. *Hall. hist.* 1845.
Bryum pendulum serpyllifolio rotundiore pellucido, capsulis ovatis. *Dill. musc.* 416. t. 53. f. 81.
Bryum nitidum serpylli rotundis et latioribus foliis pellucidis. *R. syn.* 103.
Muscus polytrichoides palustris major, serpylli latiore folio pellucido. *Hist. Ox. III.* 627. f. 15. t. 6. f. 39.
- β Bryum pendulum serpylli folio longiore pellucido, capsulis oblongis cuspidatis. *Dill. musc.* 416. t. 53. f. 80.
- cuspidatum.* γ Mnium pedunculis aggregatis, foliis alternis acutis serratis. *Sp. pl.* 1577. *Weis. crypt.* 155.
Mnium foliis ovato-lanceolatis confertis serratis, capsulis nutantibus obtusis. *Hall. hist.* 1841.
Bryum pendulum, foliis variis pellucidis, capsulis ovatis. *Dill. musc.* 413. t. 53. f. 79.
- proliferum.* δ Mnium pedunculis aggregatis, foliis rosaceo-congestis lanceolatis acutis. *Sp. pl.* 1578.
Bryum stellare roseum majus, capsulis ovatis pendulis. *Dill. musc.* 411. t. 52. f. 77.
Bryum roseum majus, foliis oblongis. *R. syn.* 92.
Muscus coronatus humilis, foliolis latioribus stellatim nascentibus donatus. *Hist. Ox. III.* 630. f. 15. t. 7. f. 9.
- Bryum stellare roseum pendulum minus, capsulis et setis longioribus. *Dill. musc.* 412. t. 52. f. 78.
Bryum roseum minus, foliis subrotundis. *R. syn.* 92.
- 1 Muscus parvus stellaris. *B. pin.* 61. *Ger. em.* 1563. *Park.* 1308.
- undulatum.* ζ Mnium pedunculis aggregatis, foliis oblongis undulatis. *Sp. pl.* 1578.

Bryum dendroides polycephalon phyllitidis folio undulato pellucido, capsulis ovatis pendulis. Dill. musc. 410. t. 52. f. 76.

Bryum nitidum foliis oblongis undulatis, capitulis cernuis arbusculam referens. R. syn. 103.

Muscus ad polytrichoidon accedens ramosus, foliis longis lucidis et veluti crispis. Hist. Ox. III. 630. f. 15. t. 6. f. 1.

Anglis, thyme-leaved Bryum.

Habitat in sylvis et ericetis humidiusculis. 4 IV.

HYPNUM. 1195.

Masc. Anthera operculata. Calyptra lævis. Filamentum ortum e perichætio.

Fem. Pistila solitaria, ex alis foliorum, nudæ plumulæ.

* *Frondebis pinnatis.*

1. HYPNUM fronde simplicissima pinnata lanceo-bryides. lata apice pedunculifera. *Sp. pl. 1588. Weis. crypt. 217. Fl. dan. 473. f. 1.*

Hypnum foliis planis pinnatis, parium septem, setis ex summitate. Hall. hist. 1776.

Hypnum taxiforme exiguum, versus summitatem capsuliferum. Dill. musc. 262. t. 34. f. 1.

Hypnum repens filicifolium non ramosum, pediculis brevibus versus foliorum summitatem egredientibus. R. syn. 88.

Muscus polytrichoides, exiguis capitulis in extremis cauliculis s. foliis, subrotundis erectis. Hist. Ox. III. f. 15. t. 6. f. 11. Vaill. paris. 136. t. 24. f. 13.

Anglis, dwarf Hypnum.

Habitat in umbrosis et ad sepes. ☉. II—V.

2. HYPNUM fronde simplicissima pinnata lanceolata taxifoliosi-basi pedunculifera. *Sp. pl. 1587. Weis. crypt. 216. um. Fl. dan. 473. f. 2.*

Hypnum foliis planis pinnatis, parium duodecim, setis radicalibus. Hall. hist. 1775.

Hypnum taxiforme minus, basi capsulifera. Dill. musc. 263. t. 34. f. 2.

Hypnum

- Hypnum repens filicifolium non ramosum, pediculis brevioribus ad radicem egredientibus. *R. syn.* 88.
Muscus pennatus, capitulis adiante. *Vaill. paris.* 136. t. 2. f. 11.
Anglis, yew-leaved Hypnum.
Habitat in sylvis et umbrosis. 4. II-V.
- adiantoides.* 3. HYPNUM fronde pinnata ramosa erecta medio pedunculifera. *Sp. pl.* 1588.
Hypnum foliis planis, pinnarum paribus quadraginta. *Hall. hist.* 1774.
Hypnum taxiforme palustre ramosum, majus et erectum. *Dill. musc.* 264. t. 34. f. 3.
Hypnum erectum filicifolium ramosum, pinnulis acutis. *R. syn.* 87.
Muscus taxiformis ramosus. *Vaill. paris.* 136. t. 28. f. 5.
Anglis, upright Hypnum.
Habitat in paludosis et ericetis humidis. 4. III. IV.
- sylvaticum.* 4. HYPNUM fronde pinnata ramosa decumbente, foliolis acutis medio pedunculifera. *Fl. Angl.* 419. *Lin. Mant.* II. 310. *Schreb. spicil.* 87.
Hypnum denticulatum pinnatum, pinnis simplicibus rectioribus. *Hall. hist.* 1773.
Hypnum denticulatum pinnatum, pinnulis simplicibus rectioribus. *Dill. musc.* 267. t. 34. f. 6.
Hypnum repens filicifolium ramosum pedunculis et capitulis longioribus e foliorum alis egredientibus, foliolis utrinque simplicibus. *R. syn.* 88.
Anglis wood Hypnum.
Habitat in sylvis ad arborum radices. 4. IV. V.
- denticulatum.* 5. HYPNUM fronde pinnati simplici, pinnis duplicatis basi pedunculifera. *Sp. pl.* 1588. *Schreb. spicil.* 86.
Hypnum foliis laxis pinnatis ovato-lanceolatis aristatis retroversis. *Hall. hist.* 1772.
Hypnum denticulatum pinnatum, pinnulis duplicatis recurvis. *Dill. musc.* 266. t. 34. f. 5.
Hypnum repens filicifolium non ramosum pediculis et capitulis

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Hypnum caule plano, foliis distichis ovatis rugosis nitidis, operculo conico. *Hall. hist.* 1770.

Hypnum repens crispum, cauliculis compressis, lycopodii in morem per terram sparsis. *R. syn.* 88.

Anglis, waved Hypnum.

Habitat in sylvis, umbrosis et rupibus humidis. 4. III. IV.

crispum.

9. HYPNUM furculis ramosis, frondibus subpinnatis : foliolis undulatis planis. *Sp. pl.* 1589. *Weis. crypt.* 221.

Hypnum caule plano, foliis distichis ovatis nitidis, transversim undulatis operculis aristatis. *Hall. hist.* 1769. t. 46. f. 5.

Hypnum pennatum undulatum crispum, setis et capitulis brevibus. *Dill. musc.* 273. t. 36. f. 12.

Hypnum repens crispum, ramulis compressis, filicinorum more dispositis. *R. syn.* 80.

Muscus terrestris major, ramulis compressis, foliis superficie crispis. *Hist. Ox.* III. 625. f. 15. t. 5. f. 10.

Anglis, curled Hypnum.

Habitat in sylvis, ericetis, rupibus, et nemorosis cretaceis frequens. Prope Gravesend copiose. *R. syn.* 4. III. IV.

triangulum.

10. HYPNUM ramis vagis recurvis, foliis ovatis recurvatis patulis. *Sp. pl.* 1589. *Weis. crypt.* 222.

Hypnum foliis hispidis ovato-lanceolatis aristatis. *Hall. hist.* 1729.

Hypnum vulgare triangulum maximum et pallidum. *Dill. musc.* 293. t. 38. f. 28.

Hypnum repens triangularibus majoribus et pallidioribus foliis. *Dill. giss.* 219. *R. syn.* 80.

Muscus squamosus major s. vulgaris. *Vaill. paris.* 137. t. 28. f. 9.

Anglis, triangular Hypnum.

Habitat in sylvis et pascuis sterilibus. 4. IX—I.

rutabulum.

11. HYPNUM ramis vagis subrepentibus, foliolis ovatis mucronatis imbricatis. *Sp. pl.* 1589. *Weis. crypt.* 224.

Hypnum

Hypnum foliis ovato-lanceolatis patulis striatis breviter aristatis, operculis conicis. *Hall. hist.* 173.

Hypnum dentatum vulgatissimum, operculis obtusis. *Dill. musc.* 295. t. 38. f. 29.

Hypnum repens triangularibus angustioribus foliis. *R. syn.* 80.

Muscus myosuroides, rutabuli fructu. *Vaill. paris.* 27. f. 8.

Muscus erectus major, foliis angustioribus acutis. *Vaill. paris.* t. 23. f. 2.

8 Hypnum palustre erectum breve, foliis brevibus angustis tenuibus confertis. *R. syn.* 83.

7 Hypnum striatum, ramis vagis recurvis, foliis ovatis lineatis acutis undique patentibus, operculis cuspidatis. *Schreb. spicil.* 91.

Hypnum vulgare dentatum, operculis cuspidatis. *Dill. musc.* 297. t. 38. f. 30.

Hypnum repens triangularibus minoribus foliis, pediculis et capitulis brevioribus et tumidioribus, minus. *R. syn.* 80.

8 Hypnum ruscifolium, furculo repente, ramis imbricatis fetiferis, foliis carinatis, capitulo subrotundo incurvo acuminato. *Neck. meth. musc.* 181.

Hypnum foliis ovato-lanceolatis acutissimis, operculis aristatis. *Hall. hist.* 1731. *Weis. crypt.* 225.

Hypnum foliis rusciformibus, capsulis subrotundis. *Dill. musc.* 298. t. 38. f. 31.

Anglis, toothed Hypnum.

Habitat in sylvis, ericetis et sepibus. 4. IX—XII.

12. HYPNUM furculis vagis ramosissimis, ramis filiformibus teretibus, antheris obliquis. *Fl. angl.* 421. *Mant. II.* 308.

Hypnum furculo ramosissimo decumbente, ramis teretibus, capitulis erectis acutis. *Neck. meth. musc.* 179.

Hypnum furculis ramosissimis: ramis rectis teretibus multifloris, capsulis erectis. *Schreb. spicil.* 97.

Hypnum foliis perangustis, filo terminatis, capsulis inclinatis breviter aristatis. *Hall. hist.* 1755.

Hypnum sericeum ramosius et tenuius, capsulis acuminatis. *Dill. musc.* 327. t. 42. f. 62.

Hypnum repens trichoide terrestre, priori viridius et minus, capitulis cernuis minus tumidis. *R. syn.* 84.

Anglis, filiforme *Hypnum*.

Habitat in arborum truncis. 4. II.

*** *Surculis pinnatis.*

filicinum. 13. **HYPNUM** furculis pinnatis : ramulis distantibus foliis imbricatis incurvis acutis secundis. *Sp. pl.* 1590. *Nick. dilic. gallob.* 470.

Hypnum caulibus pinnatis, pinnis distantibus alterne longioribus, ex apice confertis, foliis cincinnatis.

Hall. hist. 1766.

Hypnum repens filicinum crispum. *R. syn.* 85. *Dill. musc.* 282. t. 136. f. 19.

Muscus filicinus palustris. *Vaill. paris.* 138. t. 29. f. 9.

β *Hypnum repens filicinum*, trichoides palustre. *R. syn.* 85. *Dill. musc.* 286. t. 35. f. 21.

γ *Hypnum repens trichoides palustre vernum fuscum* capitulis oblongis incurvis. *R. syn.* 85.

Anglis, bog *Hypnum*.

Habitat in palustribus et ad fontes. 4. III.

compressum. 14. **HYPNUM** furculis pinnatis compressis, foliis mucronatis incurvis, antheris erectiusculis, *Fl. angl.* 423. *Lin. mant.* II. 310.

Hypnum ramis pinnatis, pinnis ramosis, foliis cincinnatis. *Hall. hist.* 1767.

Hypnum filicinum sericeum molle et pallidum mucronibus aduncis. *Dill. musc.* 286. t. 36. f. 22.

Anglis, compressed *Hypnum*.

Habitat in sylvis et arborum truncos; in sylva prope Tottenham non longe a Londino. *Dill.* III. IV.

crista castrensis. 15. **HYPNUM** furculis pinnatis : ramulis approximatis apicibus recurvis. *Sp. pl.* 1595. *Weis. crypt.* 233.

Hypnum foliis pinnatis, pinnis simplicibus æqualibus foliis cincinnatis. *Hall. hist.* 1768.

Hypnum filicinum, cristam castrensem representans. *Dill. musc.* 284. t. 36. f. 20.

Muscus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- Hypnum repens filicinum, triangularibus parvis foliis, prælongum. *R. syn.* 80. *Dill. musc.* 278. t. 35. f. 15.
 Muscus filicinus minor. *Vaill. paris.* 140. t. 23. f. 9.
Anglis, trailing Hypnum.
 Habitat in arborum truncis, lignis putrescentibus et terris
 udis. 4. XI—II.

abieti-
num.

19. HYPNUM furculis pinnatis teretiusculis remotis
 inæqualibus. *Sp. pl.* 1591. *Weis. crypt.* 236.
 Hypnum ramis duris teretibus, foliis appressis ovato-
 lanceolatis pilo aristatis. *Hall. hist.* 1762.
 Hypnum lutescens, alis subulatis tenacibus. *Dill.*
musc. 280. t. 35. f. 17.
 Hypnum repens filicinum trichoides montanum, ra-
 mulis teretibus lutescentibus non divisis. *R. syn.* 86.
 Muscus terrestris furculis filamentosis tenacibus abie-
 tinis, semel tantum divisis. *Hist. Ox. III.* 626. f.
 15. t. 5. f. 22.
 Muscus pennatus minor, caulibus ramosis, in summi-
 tate velut spicatus. *Vaill. paris.* t. 29. f. 12.
Anglis, fir Hypnum.
 Habitat in nemorosis aridis et collibus cretaceis passim;
 circa Northfleet copiose. 4. III.

*** Foliis reflexis.

cupressi-
forme.

20. HYPNUM furculo subpinnato, foliis secundis re-
 curvis apice subulatis. *Sp. pl.* 1592. *Weis. crypt.*
 1737. *Fl. dan.* 535.
 Hypnum caule compresso, foliis imbricatis falcatis
 pilo aristatis. *Hall. hist.* 1737.
 Hypnum crispum cupressiforme, foliis aduncis. *Dill.*
musc. 287. t. 32. f. 23.
 Hypnum repens crispum cupressiforme. *R. syn.* 89.
 Muscus terrestris aureus minor, flagellis clavisque bre-
 vibus. *Hist. Ox. III.* 624. f. 15. t. 5. f. 8.
 Muscus squamosus ramosus, minor et crispus. *Vaill.*
paris. 139. t. 27. f. 13.
 B Hypnum repens trichoides terrestre, foliolis uno versu
 dispositis. *R. syn.* 85.
Anglis, cypress Hypnum.

Habitat

Habitat in sylvis et ad arborum radices. 4. II. III.

21. HYPNUM surculo erectiusculo subramoso, foliis *aduncum*. secundis recurvis subulatis, ramis recurvatis. *Sp. pl.* 1592. *Weis. crypt.* 240.

Hypnum caule erecto, foliis falcatis pilo aristatis, operculis conicis. *Hall. hist.* 1736.

- Hypnum palustre erectum summitatibus aduncis. *R. syn.* 82. *Dill. musc.* 292. t. 37. f. 26.

Anglis, hooked Hypnum.

Habitat in palustribus. 4. IV—VIII.

22. HYPNUM ramis vagis procumbentibus recurvis, *scorpioides*. foliis secundis acuminatis. *Sp. pl.* 1593.

Hypnum scorpioides palustre magnum, Lycopodii instar sparsim. *Dill. musc.* 290. t. 37. f. 25.

Anglis, scorpion Hypnum.

Habitat in palustribus turfosis; prope Bishops-castle in agro Salopiensi et circa Pentir, prope Bangor. *Dill. prope Norwich.* D. Pitchford, in septentrionali parte Angliæ passim. 4. XI—IV.

23. HYPNUM surculis repentibus, ramis vagis tereti-viticulibus, foliis patulis acuminatis. *Sp. pl.* 1592. *Weis. sum. crypt.* 241.

Hypnum foliis lanceolatis crispis, capsulis erectis, operculo conico. *Hall. hist.* 1739.

Hypnum subhirsutum, viticulis gracilibus erectis, capsulis teretibus. *Dill. musc.* 307. t. 39. f. 43.

Hypnum repens trichoides arboreum majus, capsulis et surculis erectis, minus ramosis. *R. syn.* 85.

Muscus montanus, flagellis longis tenuibus cupressinis, ut plurimum indivisis. *Hist. Ox. III.* 624. f. 15. t. 5. f. 7.

Muscus squamosus, viticulis longioribus glabris. *Vaill. paris.* 137. t. 23. f. 1.

Anglis, cylindric Hypnum.

Habitat in arborum stipitibus et ericetis. in collibus cretaceis inter Northfleet et Gravesend in Cantio. *Dill.* III. IV.

- palustre.* 24. HYPNUM furculis repentibus, ramis confertis erectis, foliis ovatis secundis, antheris erectiusculis. *Fl. angl.* 429. *Sp. pl.* 1593.
 Hypnum foliis ovato-lanceolatis, falcatis, capsulis ovatis. *Hall. hist.* 1738.
 Hypnum heterophyllum aquaticum polycephalum repens. *Dill. musc.* 293. t. 37. f. 27.
 Hypnum repens filicifolium ramosum, ramulis surrectis et minus complanatis. *R. syn.* 87.
Anglis, marsh Hypnum.
 Habitat ad arborum radices et in saxosis udis. ad ripam Thamesis prope Batterseam. 4. I—IV.
- squarrosum.* 25. HYPNUM ramis vagis, foliis ovatis apice subulatis undique reflexis. *Sp. pl.* 1593. *Weis. crypt.* 242. *Fl. dan.* 535.
 Hypnum foliis patulis subulatis, operculis conicis. *Hall. hist.* 1732.
 Hypnum repens triangularibus reflexis foliis, majus. *R. syn.* 82. *Dill. musc.* 303. t. 39. f. 38.
 Muscus terrestris erectior, foliis reflexis. *Hist. Ox.* III. 624. f. 15. t. 5. f. 2.
 β Hypnum foliis cincinnatis, patulis reflexis. *Hall. hist.* 1733.
 Hypnum repens triangularibus reflexis foliis, minus. *R. syn.* 82. *Dill. musc.* 304. t. 39. f. 39.
Anglis, common Hypnum.
 Habitat in pratis et pascuis humidis. 4. III—VI.
- loreum.* 26. HYPNUM furculis repentibus, ramis vagis erectis, foliis secundis, antheris subrotundis. *Sp. pl.* 1593. *Weis. crypt.* 244.
 Hypnum loreum montanum, capsulis subrotundis. *Dill. musc.* 305. t. 39. f. 40.
 Hypnum repens, furculis magis erectis, foliis reflexis longioribus cinctis, operculo capituli magno. *R. syn.* 82.
 Muscus terrestris vulgaris. *Ger. em.* 1370. alter. *Park.* 306.
 Muscus repens major, foliis et flagellis longis et tenuibus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

***** *Surculus teretiusculus.*

curtipen- 29. HYPNUM furculis vagis teretibus, foliis ovatis
dulum. acutis patulis, antheris pendulis. *Sp. pl.* 1594.
Weis. crypt. 248.

Hypnum ramis teretibus, foliis ovato-lanceolatis pilo
aristatis, capsulis ovatis pendulis, operculis arista-
tis. *Hall. hist.* 1740.

Hypnum dentatum curtipendulum, viticulis rigidis.
Dill. musc. 333. t. 43. f. 69.

Hypnum arboreum repens, capsulis reflexis brevibus
pediculis insidentibus. *R. syn.* 89.

Anglis, pendulous Hypnum.

Habitat ad radices arborum et in locis saxosis. 4. II—IV.

purum. 30. HYPNUM furculis pinnato-sparfis subulatis, foliis
ovatis obtusis capsulis obliquis. *Sp. pl.* 1594. *Weis.*
crypt. 249.

Hypnum ramis teretibus, foliis ovato-lanceolatis,
setis prælongis, capsulis inclinatis aristis. *Hall.*
hist. 1743.

Hypnum cupressiforme vulgare, foliis obtusis. *Dill.*
musc. 309. t. 40. f. 45.

Hypnum terrestre erectum, ramulis teretibus, foliis
inter rotunda et acuta medio modo se habentibus.
Dill. giff. 220. *R. syn.* 81.

Muscus squamosus, cupressiformis. *Vaill. paris.* 138.
t. 28. f. 3.

• Hypnum ramis teretibus, foliis ovato-lanceolatis setis
prælongis, capsula crassa, operculo conico. *Hall.*
hist. 1742.

Hypnum cupressiforme tenuius et compressius. *Dill.*
musc. 312. t. 40. f. 47.

Hypnum longum erectum, foliis angustis caulibus ad-
pressis. *R. syn.* 83.

Anglis, meadow Hypnum.

Habitat in pascuis, pratis et sylvis. 4. II.

illece- 31. HYPNUM furculis ramisque vagis teretibus erec-
brum. tiusculis obtusis. *Sp. pl.* 1594. *Schreb. spicil.* 95.
Hypnum foliis ovato-lanceolatis concavis, ramis te-
retibus. *Hall. hist.* 1744.

Hypnum

Hypnum cupressiforme rotundius vel illecebræ æmulum. *Dill. musc.* 311. t. 40. f. 46.

Hypnum terrestre erectum, ramulis teretibus, foliis subrotundis albo-virentibus cinctis. *R. syn.* 81.

Muscus terrestris, furculis kali geniculati aut illecebræ æmulis foliolis subrotundis squamatis incumbentibus. *R. syn. ed.* 11. 37. *Hist. Ox. IH.* 626. *Vaill. paris.* 137. t. 25. f. 7.

Anglis, glass-wort Hypnum.

Habitat in ericetis, nemorosis, et pascuis humidiusculis. 4. III. IV.

32. HYPNUM furculis teretibus ramosis, foliis acutis *riparium* patulis distantibus. *Sp. pl.* 1595. *Weis. crypt.* 250.

Hypnum ramis teretibus et pinnatis, foliis ovato-lanceolatis pilo terminatis, setis caulinis brevibus. *Hall. hist.* 1746.

Hypnum aquaticum, flagellis et teretibus et pennatis. *Dill. musc.* 308. t. 40. f. 44.

Hypnum ramolium fluitans pennatum. *R. syn.* 81. *Anglis,* water Hypnum.

Habitat in muris et saxosis humidis ad ripas fluviorum et in fluviis. 4. IX—IV.

33. HYPNUM furculis vagis: apice foliis convolutis *cuspidatum* acuminatis. *Sp. pl.* 1595. *Weis. crypt.* 251.

Hypnum foliis ovato-lanceolatis, ramis teretibus, supremis conicis acuminatis. *Hall. hist.* 1741.

Hypnum palustre extremitatibus cuspidatis et pungentibus. *Dill. musc.* 300. t. 39. f. 34.

Hypnum repens palustre, foliis triangularibus per caules expansis, extremitatibus convolutis et acuminatis. *Dill. giss.* 219. *R. syn.* 82.

Anglis, pointed Hypnum.

Habitat in palustribus pascuis et sylvis udis. 4. III—V.

***** *Surculis confertis.*

34. HYPNUM furculo repente, ramis confertis erectis, *sericeum* foliis subulatis, antheris erectis. *Sp. pl.* 1595. *Weis. crypt.* 254.

Hypnum

Hypnum ramis teretibus, foliis pilo præpilatis, capsulis cylindricis erectis aristatis. *Hall. hist.* 1750.

Hypnum vulgare sericeum recurvum, capsulis erectis cuspidatis. *Dill. musc.* 323. t. 42. f. 59.

Hypnum repens trichoides terrestre luteo-virens vulgare majus, capitulis erectis. *R. syn.* 84.

Muscus terrestris luteo-viridis sericeus repens. *R. syn. ed. II.* 38. *Hist. Ox. III.* 626. f. 15. t. 5. f. 25.

Muscus arboreus splendens sericeus. *Vaill. paris.* 123. t. 27. f. 3.

plumosum. *B* Hypnum furculis pinnatis repentibus, ramis confertis, foliis imbricatis subulatis, antheris erectis. *Fl. angl.* 423. *Sp. pl.* 237. *Weis. crypt.* 237.

Hypnum repens filicinum plumosum. *R. syn.* 86. *Dill. musc.* 280. t. 35. f. 16.

lutescens. *γ* Hypnum furculis vagis procumbentibus, foliis subulatis antheris obliquis. *Fl. angl.* 421.

Hypnum caulibus teretibus, foliis pilo terminatis, capsulis obliquis aristatis. *Hall. hist.* 1753.

Hypnum sericeum, furculis longioribus et rectioribus, capsulis incurvis. *Dill. musc.* 325. t. 42. f. 60.

Anglis, silky Hypnum.

Habitat in muris, truncis arborum et pascuis, *γ* in collibus cretaceis inter Northfleet et Gravesend. 4. I—IV.

velutinum. 35. HYPNUM furculo repente, ramis confertis erectis, foliis subulatis, antheris subnutantibus. *Sp. pl.* 1595. *Weis. crypt.* 255. *Fl. dan.* 475.

Hypnum ramis teretibus, foliis pilo terminatis, capsulis ovatis inclinatis mucronatis, *Hall. hist.* 754.

Hypnum velutinum, capsulis ovatis cernuis. *Dill. musc.* 326. t. 42. f. 61.

Hypnum repens trichoides terrestre viridius minus, capitulis tumidioribus cernuis. *R. syn.* 84.

Muscus squamosus ramosus tenuior, capitulis incurvis. *Vaill. paris.* 138. t. 26. f. 9.

B Hypnum repens trichoides terrestre minimum et breve, capitulis tumidioribus cernuis. *R. syn.* 84.

Anglis, velvet Hypnum.

Habitat ad radices arborum et sepes, et in umbrosis. 4. II. III. .

36. HYPNUM

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Muscus trichoides humilis ramosus, capitulis oblongis tumidiusculis donatus. *Hist. Ox. III. 628. f. 15. t. 6. f. 3.*

Muscus squamosus minor, myosuroides, capitulis incurvis. *Vaill. paris. 137. t. 27. f. 6.*

♁ Hypnum myosuroides crassius, capsulis erectis. *Dill. musc. 316. t. 41. f. 50.*

Hypnum repens triangularibus angustis foliis, ramulis subrotundis. *R. syn. 83.*

γ Hypnum myosuroides brevius et crassius, capsulis cernuis. *Dill. musc. 318. t. 41. f. 52.*

♂ Hypnum myosuroides sericeum tenuius, capsulis erectis. *Dill. musc. 318. t. 41. f. 53.*

Anglis, mouse-tail Hypnum.

Habitat in sylvis, saxosis et muris. 2. I—IV.

*ornitho-
poides.*

39. HYPNUM furculis repentibus, ramis fasciculatis teretibus erectiusculis, antheris erectis ovatis. *Fl. angl. 430. Lin. mant. 310.*

Hypnum gracile ornithopoides. *Dill. musc. 320. t. 41. f. 55.*

Anglis, birds-foot Hypnum.

Habitat in truncis arborum et rupibus passim. in Fagis in saltu Enfieldiensi copiose; ad saxa per campos elatos Grey-weather dictos. S. Brewer. 2. II—IV.

*clavella-
tum.*

40. HYPNUM repens, ramulis erectis confertissimis, antheris incurvis, operculis inflexis. *Sp. pl. 1596?*

Hypnum furculo repente, ramis contiguis teretibus erectis duplo seta brevioribus, capitulo incurvo acuminato. *Neck. meth. muse. 176.*

Hypnum clavellatum parvum repens, setis brevibus. *Dill. musc. 551. t. 81. f. 17.*

Anglis, club Hypnum.

Habitat in ramis mortuis et truncis arborum passim. 2. XII—II.

ALGÆ.

A L G Æ.

JUNGERMANNIA. 1169.

Masc. Pedunculatus nudus. *Anthera* quadrivalvis.
Fem. Sessilis nudus, seminibus subrotundis.

* *Frondebis pinnatis secundis.*

1. JUNGERMANNIA frondibus simpliciter pinnatis, *asplenio-*
 foliolis ovatis subciliatis. *Sp. pl.* 1597. *Weis. crypt. des.*
 110.

Jungermannia foliis pinnatis subrotundis serratis, ex
 apice floriferis. *Hall. hist.* 1868.

Lichenastrum asplenii facie, pinnis laxioribus. *Dill.*
musc. 482. t. 69. f. 5.

- β Lichenastrum asplenii facie, pinnis confertioribus.
Dill. musc. 483. t. 69. f. 6.

Lichenastrum trichomanis facie, capitulis e foliorum
 summitate nascentibus, majus. *R. syn.* 112.

Musculus trichomanoïdes, foliis rotundioribus pelluci-
 dis, squamatim conjuncte sibi incumbentibus. *Hist.*
Ox. III. 627. f. 15. t. 6. f. 42.

Hepaticoides polytrichi facie. *Vaill. paris.* 99. t. 19.
 f. 7.

Anglis, spleenwort Jungermannia.

Habitat in sylvis et umbrosis udis. 4. II—IV.

2. JUNGERMANNIA frondibus simpliciter pinnatis, *viticulosa.*
 foliolis subulatis. *Sp. pl.* 1597.

Jungermannia foliis pinnatis subrotundis, ex caule
 florifera. *Hall. hist.* 1869.

Jungermannia terrestris, viticulis longis, foliis perexi-
 guis, densissimis, ex rotunditate acuminatis. *Mich.*
gen. 8. t. 5. f. 4.

Lichenastrum trichomanis facie, e basi et medio flo-
 rens. *Dill. musc.* 484. t. 69. f. 7.

Lichenastrum capitulis nudis, trichomanis facie, fo-
 liolis densius congestis majus. *R. syn.* 111.

Anglis, straggling Jungermannia.

Habitat

Habitat in sylvis et umbrosis humidis; in comitatu Eboracensi. D. Richardson. circa Slingsfoldiam in Suffexia. 4. III. IV.

multiflora. 3. JUNGERMANNIA frondibus simpliciter pinnatis basi floriferis, foliolis fetaceis.

Jungermannia fronde repente ramosa, foliolis alternis geminis fetaceis æqualibus. *Lin. mann. II. 310.*

Lichenastrum multiflorum exile, foliis angustissimis. *Dill. musc. 481. t. 69. f. 4.*

Anglis, many-flowered Jungermannia.

Habitat in sylvula collis Shooter's-hill dicti, prope Londinum. *Dill. III.*

polyan-
thos. 4. JUNGERMANNIA frondibus simpliciter pinnatis, foliolis integerrimis imbricatis convexis. *Sp. pl. 1597. Weis. crypt. 113.*

Jungermannia foliis pinnatis ovatis ex caule florifera. *Hall. bist. 1870.*

Jungermannia palustris repens, foliis densissimis ex rotunditate acuminatis. *Mich. gen. 8. t. 5. f. 5.*

Lichenastrum trichomanis facie, polyanthemum breve et repens. *Dill. musc. 486. t. 70. f. 9.*

Anglis, imbricated Jungermannia.

Habitat in sylvis et umbrosis humidis et ad ripas rivulorum. 4. III. IV.

lanceola-
ta. 5. JUNGERMANNIA frondibus simpliciter pinnatis lanceolatis apice floriferis, foliolis integerrimis. *Sp. pl. 1597. Weis. crypt. 114.*

Jungermannia foliis pinnatis ovatis confertis ex apice floriferis. *Hall. bist. 1871.*

Jungermannia palustris minima repens, foliis subrotundis densissimis læte viridibus. *Mich. gen. 8. t. 5. f. 6.*

Lichenastrum trichomanis facie minus, ab extremitate florens. *Dill. musc. 486. t. 70. f. 10.*

Lichenastrum capitulis nudis, trichomanis facie, foliolis densius congestis, minus. *R. syn. 112.*

Anglis, lanceolate Jungermannia.

Habitat in umbrosis humidis. II. III.

6. JUNGER-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Florifera, foliolis quinquedentatis. *Fl. angl.* 443.
Sp. pl. 1598.

Lichenastrum multifidum majus, ab extremitate florens. *Dill. musc.* 494. t. 71. f. 23.

Lichenastrum trichomanis facie, foliolis multifidis, capitulis e summis ramulis nascentibus. *R. syn.* 113.

Muscoides terrestre repens, ex obscuro virescens, foliis superioribus inferioribusque ad apicem dentatis.

Mich. gen. 10. t. 6. f. 2.

Anglis, toothed Jungermannia.

Habitat in sylvis et umbrosis udis, in comitatu Eboracensi.

D. Richardson. 4. IV.

•• *Frondebis pinnatis : pinnulis auriculatis superne tectis.*

undulata. 10. JUNGERMANNIA frondibus suprabipinnatis apice floriferis, foliolis subrotundis integerrimis.

Sp. pl. 1598. *Weis. crypt.* 120.

Lichenastrum pinnulis auriculatis majoribus et non crenatis. *Dill. musc.* 490. t. 71. f. 17.

Hepatica saxatilis undulata seminifera. *Vaill. paris.* t. 19. f. 6.

Anglis, waved Jungermannia.

Habitat in umbrosis. 4. III. IV.

nemorosa. 11. JUNGERMANNIA frondibus supra bipinnatis apice floriferis, foliolis ciliatis. *Sp. pl.* 1598. *Weis. crypt.* 121.

Jungermannia foliis bipartitis, auriculatis imbricatis supremis punctiferis. *Hall. hist.* 1859.

Jungermannia nemorosa, foliis acutioribus auritis, tenuissime denticulatis, flore insidente pediculo brevioris. *Mich. gen.* 7. t. 5. f. 8.

Lichenastrum auriculatum, pinnis minoribus crenatis. *Dill. musc.* 490. t. 71. f. 18.

Anglis, wood Jungermannia.

Habitat in sylvis et umbrosis udis; in comitatu Westmorlandico. 4. III. IV.

resupina- 12. JUNGERMANNIA frondibus supra bipinnatis
sa. inferis

- inferne floriferis, foliolis crenulatis imbricatis rotundis. *Sp. pl.* 1599.
 Lichenastrum auriculatum, pinnulis rotundis, crispum. *Dill. musc.* 491. t. 71. f. 19.
 Lichenastrum trichomanis facie, capitulis foliorum summitate enascentibus medium. *R. syn.* 112.
Anglis, curled Jungermannia.
 Habitat in fissuris rupium et ericetis turfosis. 4. IV.
13. JUNGERMANNIA frondibus supra bipinnatis *albicans.* apice floriferis, foliolis linearibus recurvatis. *Sp. pl.* 1599. *Weis. crypt.* 122.
 Lichenastrum auriculatum, pinnulis angustis planis recurvis. *Dill. musc.* 492. t. 71. f. 20.
 Lichenastrum trichomanis facie, capitulis e foliorum summitate enascentibus minus. *R. syn.* 112.
 Hepaticoides albescens, foliis pinnatis. *Vuill. paris.* 57. t. 19. f. 5.
Anglis, white Jungermannia.
 Habitat in sylvis et umbrosis udis. 4. III. IV.
14. JUNGERMANNIA frondibus subtus bipinnatis, *trilobata.* foliolis quadratis obsolete trilobatis. *Sp. pl.* 1599. *Weis. crypt.* 118.
 Jungermannia caulibus convexis, foliis descendentes imbricatis tridentatis. *Hall. hist.* 1866.
 Jungermannia foliis latiusculis obtusis undulatis et veluti angulosis. *Mich. gen.* 7. t. 5. f. 10.
 Lichenastrum pinnulis obtuse trifidis nervo geniculato. *Dill. musc.* 493. t. 71. f. 22.
Anglis, trid Jungermannia.
 Habitat in montibus Cambrobritannis, in monte Crib y ddiscil. prope Llanberris. 4. III. IV.
15. JUNGERMANNIA frondibus subtus bipinnatis *reptans.* apice reptantibus, foliolis quadridentatis. *Sp. pl.* 1599. *Weis. crypt.* 119.
 Jungermannia minima ramosa, polyanthos, foliis imbricatis trifidis. *Hall. hist.* 1879.
 Lichenastrum multifidum exiguum ad basin florens

per siccitatem imbricatam. *Dill. musc.* 494. t. 71.
f. 24.

Lichenastrum trichomanis facie foliolis multifidis,
capitulis ex imis cauliculis nascentibus. *R. syn.* 113.

Anglis, creeping *Jungermannia*.

Habitat in umbrosis humidis. 4. IV.

*** *Fronibus imbricatis.*

complanata. 16. *JUNGERMANNIA* furculis repentibus, foliolis
inferne auriculatis duplicato-imbricatis, ramis

æqualibus. *Sp. pl.* 1599. *Weis. crypt.* 124.

Jungermannia foliis rotundis, alterne imbricatis, caule
plano multifloro, fetis brevissimis. *Hall. hist.*
1860.

Jungermannia foliis cincinnatis auritis, imbricatis
dispositis, ex viridi flavescentibus. *Mich. gen.* 7. t.
5. f. 21.

Lichenastrum imbricatum majus, squamis compres-
sis et planis. *Dill. musc.* 496. t. 72. f. 26.

Lichenastrum imbricatum majus. *R. syn.* 111.

Anglis, flat *Jungermannia*.

Habitat in arborum truncis frequens. 4. IV.

dilatata. 17. *JUNGERMANNIA* furculis repentibus, foliolis
inferne auriculatis duplicato-imbricatis, ramis apice
lterioribus. *Sp. pl.* 1600. *Weis. crypt.* 126.

Jungermannia foliis imbricatis alternis orbiculatis,
fetis brevissimis. *Hall. hist.* 1874.

Lichenastrum imbricatum minus, squamis convexo-
concavis. *Dill. musc.* 497. t. 72. f. 27.

Lichenastrum imbricatum minus. *R. syn.* 111.

Hepaticoides furculis et foliolis thuyæ instar compres-
sis, minor. *Vaill. paris.* 99. t. 19. f. 10.

Anglis, scaly *Jungermannia*.

Habitat ad truncos arborum. 4. III. IV.

tamarisci-folia. 18. *JUNGERMANNIA* foliis imbricatis serie duplici,
superioribus subrotundis convexis obtusis quadruplo-
majoribus. *Sp. pl.* 165. *Weis. crypt.* 129.

Junger-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- varia.* 21. JUNGERMANNIA furculis erectiusculis bifariam imbricatis, foliolis bipartitis. *Sp. pl.* 1601.
 Jungermannia repens, foliis cordatis carinatis. *Mich. gen.* 8. t. 5. f. 9.
 Lichenastrum foliis variis. *R. syn.* 113. *Dill. musc.* 505. t. 73. f. 36.
Anglis, various-leaved Jungermannia.
 Habitat in sylvis et umbrosis humidis. III. IV.

*** Surculis undique imbricatis foliolis sparsis.

- julacea.* 22. JUNGERMANNIA furculis teretibus, foliolis undique imbricatis, floribus pedunculatis. *Sp. pl.* 1601.
 Jungermannia caule filiformi, foliis adpressis inconspicuis. *Hall. hist.* 1882.
 Lichenastrum alpinum brji julacei argentei facie. *Dill. musc.* 506. t. 73. f. 38.
Anglis, silver Jungermannia.
 Habitat in rupibus udis, in montibus Snowdon, Glyder et Cader-Ideris in Wallia. 4. IX. X.

- rupestris.* 23. JUNGERMANNIA furculis teretibus, foliolis subulatis secundis. *Sp. pl.* 1601.
 Lichenastrum alpinum nigricans, foliis capillaceis reflexis. *Dill. musc.* 507. t. 73. f. 40.
Anglis, rock Jungermannia.
 Habitat in rupibus humentibus; in montibus Snowdon et Glyder in Arvoniam, et in boreali parte Angliæ passim. 4. III—X.

- trichophylla.* 24. JUNGERMANNIA furculis teretibus, foliolis capillaceis æqualibus. *Sp. pl.* 1601. *Weis. crypt.* 130.
 Jungermannia minima foliis capillariter multifidis. *Hall. hist.* 1880.
 Lichenastrum trichoides minimum in extremitate florens. *Dill. musc.* 505. t. 73. f. 37.
Anglis, hairy Jungermannia.
 Habitat in ericetis turfosis prope North Bierly in comitatu Eboracensi, D. Richardson. 4. IV.

25. JUNGER-

25. JUNGERMANNIA furculis teretibus, foliolis *alpina*.
 ovatis patentibus, calycibus imbricatis. *Sp. pl.*
 1602.

Lichenastrum alpinum atro-rubens teres, calycibus
 squamosis. *Dill. musc.* 506. t. 73. f. 39.

Anglis, alpine Jungermannia.

Habitat in rupibus udis in montibus Arvonix et septen-
 trionali parte Angliæ frequens. 4. IV—X.

**** Acaules frondibus simplicibus.

26. JUNGERMANNIA acaulis, foliolo frondi inna-*epiphylla*.
 to. *Sp. pl.* 1602. *Weis. crypt.* 106.

Jungermannia fronde foliacea, obtuse lobata ex medio
 florifera. *Hall. hist.* 1883. *Fl. dan.* 359.

Marsilea major atro-virens, floribus albicantibus e fo-
 liorum medio egredientibus. *Mich. gen.* 5. t. 4.
 f. 1.

Lichenastrum capitulis rotundis e foliorum medio e-
 nascentibus. *R. syn.* 110. *Dill. musc.* 508. t. 74.
 f. 41.

Lichen petræus, cauliculo calceato. *B. pin.* 362.

Lichen alter minor, folio calceato. *Park.* 1314.

Anglis, broad-leaved Jungermannia.

Habitat in umbris udis et ad rivulos. 4. III. IV.

27. JUNGERMANNIA acaulis, fronde oblonga sinu-*pinguis*.
 ata pingui. *Sp. pl.* 1602. *Weis. crypt.* 107.

Jungermannia fronde foliacea lacera ex latere flori-
 fera. *Hall. hist.* 1884.

Marsilea media pinguis, pallide virens, floribus majo-
 ribus nigricantibus ad foliorum latera egredientibus.
Mich. gen. 5. t. 4. f. 2.

Lichenastrum capitulis oblongis, juxta foliorum di-
 visuras enascentibus. *R. syn.* 110. *Dill. musc.* 509. t.
 74. f. 2.

Anglis, slippery Jungermannia.

Habitat in paludibus. IV.

28. JUNGERMANNIA acaulis fronde bipinnatifida. *multifida*.
Sp. pl. 1602. *Weis. crypt.* 109.

Lichenastrum ambrosiæ divisura. *R. syn. III. Dill. musc. 511. t. 74. f. 43.*

Anglis, multifid Jungermannia.

Habitat in sylvis et umbrosis humidis; circa Hampstead et Charleton prope Londinum copiose. 4. IV.

furcata. 29. JUNGERMANNIA acaulis, fronde lineari ramosa: extremitatibus furcatis obtusiusculis. *Sp. pl. 1602. Weis. crypt. 108.*

Jungermannia foliis linearibus dichotomis, inferne floriferis. Hall. hist. 1885.

Marsilea minima angustifolia, floribus nigricantibus ex inferiori foliorum parte, e subhirsuta et turbinata vagina erumpentibus. Mich. gen. 5. t. 4. f. 4.

Lichenastrum tenuifolium furcatum, thecis globosis. Dill. musc. 512. t. 74. f. 45.

Ulva saxatilis furcata, latiusculis et tenerioribus segmentis. R. syn. 63.

Anglis, linear Jungermannia.

Habitat in truncis arborum et saxosis udis et umbrosis. 4. IV.

pufilla. 30. JUNGERMANNIA acaulis, fronde supinnatifida, lobis imbricatis, perichætio plicato. *Sp. pl. 1136.*

Jungermannia foliis palmatis digitatis, ex basi florifero. Hall. hist. 1887.

Lichenastrum exiguum, capitulis nigris lucidis e cotylis parvis nascentibus. Dill. musc. 513. t. 74. f. 46.

Anglis, dwarf Jungermannia.

Habitat in sylvis et ericetis arenosis humidis et umbrosis; prope Woolwich. Dill. In sylva Wray-wood dicta prope Castle-Howard in comitatu Eboracensi. D. Teasdale. 4. III—VI.

B L A S I A. 1199.

Masc. Cal. cylindricus granis repletus.

Fem. Cal. nudus. Fructu subrotundo, foliis immerso, polyspermo.

BLASIA.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Lichen fontanus major stellatus æquæ ac umbellatus et cyathophorus. *Dill. musc.* 524. t. 76. f. 6.

Lichen petræus latifolius s. Hepatica fontana. *B. pin.* 362. *R. syn.* 115.

β Marchantia capitulo stellato, radiis teretibus. *Mich. gen.* 2. t. 1. f. 2.

Lichen domesticus minor stellatus æque ac umbellatus et cyathophorus. *Dill. musc.* 527. t. 77. f. 7.

Lichen petræus stellatus. *B. pin.* 362. *R. syn.* 115.

γ Marchantia calyce communi octopartito : laciniis e plano-convexis. *Fl. lapp.* 423. *Roy. lugdb.* 507.

Marchantia capitulo non dissecto. *Mich. gen.* 1. t. 1. f. 5.

Marchantia fronde æquabili, cyathis ferratis, bacillis umbellatis et stellatis octifidis. *Hall. hist.* 1892.

Anglis, common Marchantia.

Habitat *juxta aquas, ad fontes et in locis umbrosis udis ;*
β *in areis, ambulacris et muris umbrosis et humidis.*

4. VI—VIII.

cruciata. 2. MARCHANTIA calyce communi quadripartito : laciniis tubulosis. *Sp. pl.* 1604.

Marchantia scyphis lunatis, capitulis quadrifidis. *Hall. hist.* 1888.

Lunaria vulgaris. *Mich. gen.* 4. t. 4.

Lichen seminifer lunulatus, florifer pileatus tandem cruciatus. *Dill. musc.* 521. t. 75. f. 5.

Lichen pileatus parvus, capitulo crucis instar se expandente. *R. syn.* 115.

Anglis, cross Marchantia.

Habitat *in areis et ambulacris hortorum et, umbrosis.* 4.
VI—X.

*hemi-
spherica.*

3. MARCHANTIA calyce communi quinquefido hemispherico, perichætio nullo. *Sp. pl.* 1604.

Marchantia fronde crenata, capitulis conicis hirsutis. *Hall. hist.* 1890.

Hepatica media capitulo hemisphærico. *Mich. gen.* 3. t. 2. f. 2.

Lichen

Lichen pileatus parvus, foliis crenatis. *R. syn.* 114.
Dill. musc. 519. t. 75. f. 2.
Anglis, hemisphæric Marchantia.
 Habitat ad ripas fluviorum et fossarum humidas, et in
 rupibus udis. 4. IV. V.

4. MARCHANTIA calyce communi quinqueloculari *conica*.
 subovato. *Sp. pl.* 1604. *Fl. dan.* 274.

Marchantia verrucis asperis, capitulis conicis. *Hall.*
hist 1889.

Hepatica vulgaris major f. officinarum Italiæ. *Mich.*
gen. 3. t. 2. f. 1.

Lichen vulgaris major pileatus et verrucosus. *Dill.*
musc. 516. t. 75. f. 1.

Lichen petræus cauliculo pileolum sustinente. *B. pin.*
 362.

Lichen petræus pileatus. *Park.* 1314. *R. syn.* 114.
Anglis, conic Marchantia.

Habitat ad ripas rivulorum et in umbrosis udis. 4. IV.

R I C C I A. 1200.

Cal. 0. nisi cavitatem vesicularem intra substantiam.

Cor. 0.

Stam. Filamenta nulla. *Anthera* cylindrica sessilis,
 germiini imposita.

Pistil. Germen turbinatum. *Stylus* filiformis, anthe-
 ram perforans.

Peric. Sphæricum, coronatum anthera emarcida.

Sem. hemisphærica pedicellata.

1. RICCIA frondibus glabris bipartitis acutis. *Sp. pl. minima.*
 1605.

Riccia minima nitida, segmentis angustioribus acutis.
Mich. gen. 107. t. 57. f. 6.

Lichen omnium minimus, foliolis fissis super terram
 expansis. *R. syn.* 115. *Dill. musc.* 534. t. 78. f. 11.
Anglis, least Riccia.

Habitat in inundatis; in ericeto nigro prope Londinum.
R. syn. ☉. XI. XII.

2. RICCIA

- glauca.* 2. RICCIA frondibus glabris canaliculatis, bilobis obtusis. *Sp. pl.* 1605. *Schreb. spicil.* 111.
 Riccia fronde sulcodivisa, lobis bifurcatis, lobulis lanceolatis obtusis. *Hull. hist.* 1897.
 Riccia minima pinguis foliis latiusculis ample sulcatis e glauco virescentibus. *Mich. gen.* 107. t. 54. f. 4.
 Hepatica palustris bifurca, lobis brevioribus carinatis. *Vaill. paris.* 98. t. 19. f. 1.
 Lichen minimus, foliis venosis bifariam vel trifariam se dividendo progredientibus. *R. syn. ed.* 11. 41.
Dill. musc. 533. t. 78. f. 10.
 Lichen s. Hepatica foliis crassis rutæ murariæ aut chamedryos foliis laciniatis quodammodo similibus. *R. hist.* 111. 48. *R. syn.* 116.
Anglis, glaucous Riccia.
 Habitat in ericetis arenosis, humidis. ☉. X—IV.

- natans.* 3. RICCIA frondibus obcordatis ciliatis. *Syst. nat.* 708.
 Lichen parvus vernus cordiformis, ima parte fimbriatus, lentis palustris modo aquæ innatans. *Buddl. hort. sicc. vol. II.* 15. *R. syn.* 116.
Anglis, fringed Riccia.
 Habitat in aquis stagnantibus prope Henley in comitatu Suffolciensi. *R. syn.* ☉. VIII—X.
 Obs. valde affinis Targioniæ hypophyllæ.

- fluitans.* 4. RICCIA frondibus dichotomis lineari-filiformibus. *Sp. pl.* 1686. *Fl. dan.* 275.
 Jungermannia fronde plana pinnata dichotoma, ramulis brevissime bicornibus. *Hull. hist.* 1886.
 Lichenastrum aquaticum fluitans tenuifolium furcatum. *Dill. musc.* 514. t. 74. f. 47.
 Ulva palustris furcata, angustioribus et firmioribus segmentis. *R. syn.* 63.
Anglis, flote Riccia.
 Habitat in fossis et stagnis passim. ♀. I—XII.

ANTHOCEROS. 1201.

Masc. *Cal.* sessilis, cylindricus, integer. *Anthera* subulata, longissima, bivalvis.

Fem.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- rugosus.* 3. LICHEN leprofus albicans, lineolis simplicibus punctisque nigris confertis. *Sp. pl.* 1607. *Weis. crypt.* 41.
Lichen punctatum et rugosum nigrum. *Dill. musc.* 125. t. 18. f. 2.
Anglis, wrinkled Lichen.
Habitat in corticibus arborum. 4. I—XII.
- niger.* 4. LICHEN leprofus niger, tuberculis concoloribus. Lichen crusta continua aterrima, globulis subrotundis. *Hall. hist.* 2084.
Anglis, black Lichen.
Habitat in rupibus et saxis grandioribus passim, circa St. Ives in Cornubia copiose. 4. I—XII.
Desc. Crusta granulata, dura, sicca, atra, latissime expansa. Tubercula convexa, atra, magnitudine seminis sinapis.
- sanguinari-
rius.* 5. LICHEN leprofus cinereo-virescens, tuberculis atris. *Sp. pl.* 1607.
Lichen crusta farinacea virescente, scutellis atris. *Hall. hist.* 2065.
Lichenoides leprosum, crusta cinereo-virescente tuberculis nigerrimis. *R. syn.* 71. *Dill. musc.* 126. t. 18. f. 3.
Anglis, sanguineous Lichen.
Habitat in rupibus, muris, arborum corticibus et ericetis turfosis. 4. I—XII.
- fusco-
ater.* 6. LICHEN leprofus fuscus, tuberculis atris. *Sp. pl.* 1607.
Lichen crusta farinacea inseparabili cinerea, scutis nigerrimis confertis. *Hall. hist.* 2062.
Anglis, fuscous Lichen.
Habitat in rupibus et saxis. 4. V—XII.
- calcareus.* 7. LICHEN leprofus candidus, tuberculis atris. *Sp. pl.* 1607.
Lichenoides tartareum tinctorium candidum, tuberculis atris. *Dill. musc.* 128. t. 18. f. 8.
Anglis,

- Anglis, calcareous Lichen.*
Habitat in rupibus calcareis, in septentrionali parte Angliæ et in Wallia. ʒ. I—XII.
8. LICHEN leprofus viridis margine tuberculisque atris. *atro-virens.*
Syst. nat. 709. Sp. pl. 1607.
Anglis, greenish Lichen.
Habitat in rupibus, in comitatu Eboracensi passim. ʒ. I—XII.
9. LICHEN leprofus, tuberculis nigris albo marginatis, *cinereus.*
Mant. I. 132. Syst. nat. 709.
Anglis, grey Lichen.
Habitat in rupibus et saxis grandioribus. ʒ. I—XII.
10. LICHEN leprofus niger, tuberculis atro alboque *atro-albus,*
mixtis. Syst. nat. 709. Sp. pl. 1607.
Anglis, mixed Lichen.
Habitat in rupibus. ʒ. I—XII.
11. LICHEN leprofus, verrucis subtessellatis lævigatis *pertusus.*
pertusis poro uno alterove cylindrico. Mant. I. 131.
Lichenoides verrucosam et rugosum cinereum glabrum. Dill. musc. 128. t. 18. f. 9.
Anglis, pierced Lichen.
Habitat in corticibus arborum, rupibus et muris. ʒ. I—XII.
12. LICHEN leprofus albidus, tuberculis pallidis albo *rupicola.*
marginatis. Mant. I. 132. Syst. nat. 709.
Anglis, rock Lichen.
Habitat in rupibus calcareis. ʒ. I—XII.
13. LICHEN leprofus albus, tuberculo centrali con- *betulinus.*
colori.
Anglis, birch Lichen.
Habitat in truncis et ramis Betulæ. ☉. IX—IV.
Desc. Crusta fere semiuncialis diametro, orbicularis, alba, tenua, fibrosa. Tuberculum solitarium, centrale, hemisphæricum, album, magnitudine fere seminis cannabis.
14. LICHEN

- lacteus.* 14. LICHEN leprofus albus, tuberculis concoloribus hemispæricis. *Mant. I.* 132. *Syst. nat.* 709.
Anglis, white Lichen.
Habitat in rupibus et saxis. 4. I—XII.
- Fagineus.* 15. LICHEN leprofus albus, tuberculis albis farinaceis
Sp. pl. 1608.
Lichen verrucosus crustaceus albus, orbiculis villosis concoloribus. *Hall. hist.* 2086.
Lichen crustaceus, arboribus adnascens, farinaceus albus, superficie in acetabulis pulverulentis velut efflorescente. *Mich. gen.* 99. t. 53. f. 1. 2.
Lichenoides candidum et farinaceum, scutellis fere planis. *Dill. musc.* 131. t. 18. f. 11. A. B.
Anglis, beach Lichen.
Habitat in corticibus Fagi, carpini, &c. 4. I—XII.
- Carpinus.* 16. LICHEN leprofus cinereus, tuberculis albidis rugosis. *Sp. pl.* 1608.
Lichenoides candidum et farinaceum, scutellis fere planis. *Dill. musc.* 131. t. 18. f. 11. C. D?
Anglis, horn-beam Lichen.
Habitat in corticibus Carpini, Fagi et Fraxini. 4. I—XII.
- corallinus.* 17. LICHEN leprofus ramosus teres fasciculatus fastigiatus confertissimus albus. *Mant. I.* 131. *Syst. nat.* 709.
Anglis, coral Lichen.
Habitat in rupibus, saxosis, et ericetis montosis in Wallia et boreali parte Angliæ passim. 4. I—XII.
Obs. an varietas lichenis tartarei?
- ferrugineus.* 18. LICHEN leprofus albidus, tuberculis subrotundis ferrugineus. *Syst. nat. III.* 234. *Fl. angl.* 440.
Lichenoides leprofum, tuberculis fuscis et farinaceis.
Dill. musc. 126. t. 18. f. 4. t. 55. f. 8.
Anglis, rusty Lichen.
Habitat in corticibus arborum. 4. I—XII.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

β Coralloides fungiforme ex ungula equina, livide rubescens. *Dill. musc.* 78. t. 14. f. 5.

Fungi parvi globosi, ex ungue equino putrescente enati. *R. syn.* 13. t.

Anglis, pedunculated Lichen.

Habitat in corticibus arborum. 4. I—XII.

** *Leprosi scutellati.*

candelarius. 23. LICHEN crustaceus flavus, scutellis luteis. *Sp. pl.* 1608. *Weis. crypt.* 46.

Lichen fronde laciniata rotunde lobata flava concolore. *Hall. hist.* 2023.

Lichenoides crustosum orbiculis et scutellis flavis. *Dill. musc.* 236. t. 18. f. 18.

Anglis, golden Lichen.

Habitat in rupibus, muris, et arborum truncis. 4. I—XII.

flavescens. 24. LICHEN crustaceus flavo-virescens, scutellis rufoluteis.

Anglis, yellow Lichen.

Habitat in corticibus arborum, muris et rupibus. 4. I—XII.

Obs. *Habitus præcedentis sed crusta orbicularis, rugosa, virescens; scutellæ fusco-flavescentes, seu testaceo-luteæ convexæ.*

flavo-rubescens. 25. LICHEN crustaceus cinereo-virescens, scutellis flavo rubescentibus.

Anglis, orange Lichen.

Habitat in truncis arborum, rupibus et saxis. 4. I—XII.

glidus. 26. LICHEN crustaceus albicans, peltis tuberculosis rugosis testaceis. *Mant. I.* 133. *Syst. nat.* 709.

Lichenoides tartareum lividum, scutellis lividis. *Dill. musc.* 133. t. 18. f. 14?

Anglis, broad Lichen.

Habitat in rupibus et saxis. prope Malham et Ingleton in comitatu Eboracensi. 4. I—XII.

Obs. *Crusta albida, rugosa, crassa, scutellæ convexæ, magnæ rufæ, margine reflexæ ut tubercula apparent; habitus sequentis.*

27. LICHEN

27. LICHEN crustaceus albidus, scutellis luteis, mar-tartareus-gine albo. *Sp. pl.* 1608.

Lichenoides crustaceum et leprosum acetabulis majoribus luteis, limbis argenteis. *R. syn.* 71. *Dill. musc.* 132. t. 18. f. 13.

Anglis, tartareous Lichen.

Habitat in rupibus et saxis grandioribus in boreali parte Angliæ, et in Devoniam et Cambria passim. 4. I—XII.

28. LICHEN crustaceus albicans, scutellis pallidis. *pallefcens.* *Sp. pl.* 1608. *Weis. crypt.* 47.

Lichen fronde laciniata appressa, cervina concolore. *Hall. hist.* 2024.

Lichenoides crustosum orbiculare incanum. *Dill. musc.* t. 18. f. 17.

Lichenoides cinereum mere crustaceum eleganter expansum. *R. syn.* 71.

Lichenoides crustaceum et leprosum, scutellis cinereo-virescentibus. *R. syn.* 71.

Anglis, pale Lichen.

Habitat in rupibus, muris et arborum truncis. 4. I—XII.

29. LICHEN crustaceus cinereus, scutellis concavis fa-rinaceus albidis. *albesfcens.*

Anglis, powdered Lichen.

Habitat in muris et corticibus arborum. 4. I—XII.

Obs. *Habitus* Lichenis carpinei.

30. LICHEN crustaceus albicans, scutellis subfuscis, *subfuscus.* margine cinereo subcrenato. *Sp. pl.* 1609. *Weis. crypt.* 48.

Lichen crusta tenace verrucosa cinerea, scutellis rufis marginatis. *Hall. hist.* 2049.

Lichenoides crustaceum et leprosum, scutellis sub-fuscis. *R. syn.* 71. *Dill. musc.* 134. t. 18. f. 16.

Anglis, brownish Lichen.

Habitat in corticibus arborum et muris. I—XII.

Parellus. 31. LICHEN crustaceus albus, peltis concavis obtusis pallidis. *Mant. I.* 132. *Syst. nat.* 710.
 Lichen verrucosus tenax albus, scutellis excavatis marginatis concoloribus. *Hall. hist.* 2048.
 Lichenoides leprosum tinctorium scutellis lapidum cancri figura. *Dill. musc.* 130. t. 18. f. 10.
Anglis, dyers Lichen.
 Habitat in muris et truncis arborum. 4. I—XII.

ater. 32. LICHEN crustaceus albicans rugosus, scutellis atris margine albo subcrenato. *Fl. angl.* 445.
 Lichen crusta glauca tenace verrucosa. *Hall. hist.* 2956.
 Lichenoides crustaceum et leprosum, scutellis nigricantibus majoribus et minoribus. *Dill. gifs.* 205. *musc.* 133, t. 18. f. 15. *R. syn.* 71.
Anglis, rugged Lichen.
 Habitat in muris, rupibus, et corticibus arborum. 4. I—XII.
 Obs. Variat scutella margine integra et crenata.

*** *Imbricati.*

centrifugus. 33. LICHEN imbricatus, foliolis obsolete multifidis lævibus albidis centrifugis, scutellis rufo-fuscis. *Sp. pl.* 1609. *Weis. crypt.* 54.
 Lichen fronde profunde laciniata, semipinnata, rotunde lobata, crasse villosa, scutellis peramplis. *Hall. hist.* 2018.
 Lichenoides imbricatum viridans, scutellis badiis. *Dill. musc.* 180. t. 24. f. 75.
Anglis, spreading Lichen.
 Habitat in rupibus, muris, saxis grandioribus et truncis arborum passim. 4. I—XII.

crassus. 34. LICHEN imbricatus, foliolis lobatis rotundatis crenatis virescentibus, scutellis planis fulvis.
 Lichen cartilagineus, scutellis congestis cervinis, fronde brevi rotunde lobata. *Hall. hist.* 2009.
 Lichenoides cartilagineum, scutellis fulvis planis. *Dill. hist.* 179. t. 24. f. 74.
Anglis, thick Lichen.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Anglis, stone Lichen.

Habitat in *saxis, rupibus et truncis, arborum.* 4. I—XII.

omphalodes.

37. LICHEN imbricatus, foliolis multifidis glabris obtusis incanis, punctis vagis eminentibus. *Sp. pl.* 1609.

Lichen pulmonarius tinctorius, foliis eleganter tenuiterque divisis, inferne nigricantibus et cirrhosis, superne purpureis et lacunatis, receptaculis concoloribus. *Mich. gen.* 90. t. 49. f. 2.

Lichen nigricans omphalodes. *Vaill. par.* t. 20. f. 10.

Lichen petræus purpureus derbiensis. *Park.* 1315.

Lichenoides saxatile tinctorium, foliis pilosis, purpureum. *R. syn.* 74. *Dill. musc.* 185. t. 24. f. 80.

Anglis, purple Lichen.

Habitat in *rupibus et saxis.* 4. I—XII.

fahlunensis.

38. LICHEN imbricatus, foliolis linearibus dichotomis planiusculis acutis nigris, scutellis atris. *Sp. pl.* 1610.

Lichen fronde laciniata undique nigerrima, scutellis concoloribus. *Hall. hist. et Nomencl.* 2020.

Lichenoides tinctorium atrum, foliis minimis crispis. *Dill. musc.* 188. t. 24. f. 81.

Anglis, cork Lichen.

Habitat in *rupibus et saxis grandioribus.* Prope Langdale in comitatu Lancastrensi et ex Scotia recepti. 4. I—XII.

- olivaceus.* 39. LICHEN imbricatus, foliolis lobatis nitidis lividis. *Sp. pl.* 1610. *Weis. crypt.* 58.

Lichen fronde semipinnata rotunda lobata olivacea, scutellis lævibus. *Hall. hist.* 2013.

Lichenoides olivaceum, scutellis lævibus. *Dill. musc.* 182. t. 24. f. 775.

Lichenoides crusta foliosa scutellata, pullum. *Dill. gifs.* 206. *R. syn.* 72.

- β Lichen fronde semipinnata rotunde lobata olivacea pulla, scutellis crenatis. *Hall. hist.* 2014.

Lichen crusta modo arboribus adnascens olivaceus, *Vaill. par.* t. 20. f. 8.

γ Lichenoides

- 37 Lichenoides arboreum et saxatile, crusta foliosa tenui, fusco virente, in segmenta latiora plerumque divisa.
R. syn. 74.
Anglis, olive Lichen.
 Habitat in rupibus saxis et arborum truncis. 4. I—XII.
40. LICHEN imbricatus, foliis crispis fulvis, peltis concoloribus fulvis. *Sp. pl.* 1610. *Weis. crypt.* 59. *parietinus.*
 Lichen fronde rugosa, rotunde lobata, flava concolore.
Hall. hist. 2021.
 Lichenoides vulgare sinuosum, foliis et scutellis luteis.
Dill. musc. 180. t. 24. f. 76.
 Lichenoides crusta foliosa, scutellata, flavescens. *Dill. giss.* 206. *R. syn.* 72.
Anglis, wall Lichen.
 Habitat in arborum truncis, muris et saxis. 4. I—XII.
41. LICHEN imbricatus, laciniis obtusis subinflatis. *physodes.*
Sp. pl. 1610. *Weis. crypt.* 57.
 Lichen fronde semipinnata cava, cornubus pulverulentis. *Hall. hist.* 2018.
 Lichen pulmonarius arboribus adnascens: desuper cinereus, subtus anthracinus, segmentis teretibus tubulosis, ad extremitates clausis, receptaculis florum subrubentibus. *Mich. gen.* 92. t. 50.
 Lichen arboreus albescens, segmentis cornigeris, subtus aterrimus. *Pet. gag. t.* 14. f. 6.
 Lichenoides ceratophyllon obtusius et minus ramosum. *R. syn.* 76. *Dill. musc.* 154. t. 20. f. 49.
Anglis, inflated Lichen.
 Habitat in truncis arborum, saxis et ericetis. 4. I—XII.
42. LICHEN imbricatus, foliolis multifidis linearibus fuscis subfuscis, scutellis fusco-atris.
 Lichen pulmonarius saxatilis farinaceus ex fusco-nigricans, foliolis angustissimis et creberrimis, receptaculis florum ferme nigris. *Mich. gen.* 95. t. 51. f. 6.
 Lichenoides angustifolium fuscum, scutellis pullis.
Dill. musc. 175. t. 24. f. 69.
M m 3
Anglis,

Anglis, brown Lichen.

Habitat in rupibus et saxis grandioribus, prope Eangor et alibi in Wallia et in Westmorlandia. 4. I—XII.

Stellaris. 43. LICHEN imbricatus, foliolis oblongis laciniatis angustis cinereis, scutellis pullis. *Sp. pl.* 1611. *Weis. crypt.* 60.

Lichen fronde cinerea, profundius divisa laciniata obtuse lobata, scutellis glaucis nigrescentibus. *Hall. hist.* 2017.

Lichen fronde cinerascente, lineari-multifida; scutellis fuscis. *Nick. meth. musc.* 89.

Lichen pulmonarius vulgarissimus, superne albo cinereus, inferne nigricans, segmentis angustis et eleganter divisis, receptaculis florum nigricantibus. *Mich. gen.* 91. t. 48. f. 2.

Lichenoides arboreum, crusta foliosa albo-cinerea tenuiter et eleganter dissecta, scutellis nigris. *R. syn.* 74.

Lichenoides cinereum, segmentis argutis stellatis, scutellis nigris. *Dill. musc.* 176. t. 24. f. 70.

β Lichen imbricatus, foliolis lobatis crenatis obtusis viridibus, polline conspersis, scutellis nigris margine crenatis. *Schreb. spicil.* 128.

Lichen fronde semipinnata rotunde lobata cinerea, scutellis nigris. *Hall. hist.* 2016.

Lichenoides arboreum, crusta foliosa virescente, tenuiter et eleganter dissecta, scutellis nigris. *Dill. giss.* 207. *R. syn.* 72.

Lichenoides glaucum orbiculare, segmentis latiusculis, scutellis nigris. *Dill. musc.* 177. t. 24. f. 71.

γ Lichenoides arboreum, crusta foliosa informi scutellis subnigris, limbo cinereo crispo cinctis. *R. syn.* 75. *Anglis*, starry Lichen.

Habitat in arborum truncis, muris et saxis. 4. I—XII, *Obs. color in humidis viridescens in siccis cinereus est.*

marginalis. 44. LICHEN imbricatus, foliolis multifidis dentato-crenatis atro-viridibus, scutellis marginalibus planiusculis atro-fuscis.

Lichen gelatinosus ramosus laciniatus, scutellis rubris. *Hall. hist.* 2039.

Lichen

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Anglis, sinuated Lichen.

Habitat in rupibus, saxosis et umbrosis, prope Wigmore, in Herefordia Littl. Brown; circa Ashburton in Devoniam. ☉. IX—VII.

fascicularis. 48. LICHEN imbricatus, foliolis palmatis erectiusculis, scutellis terminalibus turbinatis fronde majoribus.

Lichen foliaceus gelatinosus, tuberculis turbinatis fasciculatis fronde majoribus. *Mant.* 133.

Lichenoides gelatinosum palmatum, tuberculis conglomeratis. *Dill. musc.* 141. t. 19. f. 27.

Anglis, clustered Lichen.

Habitat in arborum truncis et saxis. 4. I—XII.

granulatus. 49. LICHEN imbricatus, foliolis subrotundis crenatis scabriusculis atro-viridibus, scutellis concavis fulvis.

Lichenoides gelatinosum atro-virens, auriculatum et granolum. *Dill. musc.* 140. t. 19. f. 24.

Anglis, granulated Lichen.

Habitat in rupibus, muris, et arenosis humidis passim; in ambulacris horti Oxoniensis. *Dill.* 4. I—XII.

fluviatilis. 50. LICHEN imbricatus, foliolis lobatis subcrenatis convexis atro-viridibus, scutellis globosis cavis concoloribus.

Lichenoides gelatinosum, foliis angustioribus tuniformibus. *Dill. musc.* 142. t. 19. f. 28.

Lichenoides gelatinosum opuntioides. *R. syn.* 72.

Anglis, river Lichen.

Habitat in saxis sub aqua in rivulis alpinis. Rich. Richardson, in rivulo e spelunco Malham-cove dicta, in comitatu Eboracensi, erumpente primus observavit; in torrentibus alpinis Montis Snowdon. *Dill.* 4. I—XII.

palmatus. 51. LICHEN imbricatus, foliolis subpalmatis subcrenatis glauco-purpurascensibus, scutellis fulvis.

Lichen terrestris minimus fuscus. *R. syn. ed. II.* 331. *hist. III.* 48. *Vaill. paris.* 144. t. 21. f. 15.

Lichenoides

- Lichenoides gelatinosum tenerius laciniatum, ex fusco-purpurascens. *R. syn.* 72.
- Lichenoides pellucidum fuscum corniculatum. *Dill. musc.* 143. t. 19. f. 30.
Anglis, palmated Lichen.
Habitat in ericetis et truncis arborum inter alios muscos.
☉. IX—VI.

●●●● Foliacei.

52. LICHEN foliaceus membranaceus erectiusculis *tremeloides.*
margine crispo lacinulato, scutellis concavis fulvis.
- Lichen foliaceus gelatinosus foliorum segmentis capilliceis teretibus ramosis. *Scop. carn. ed. II.* 1412.
Weis. crypt. 52.
- Lichen atro-cæruleus laciniatus et crispus. *Hall. hist.* 2032.
- Lichen terrestris membranaceus mollior fuscus, receptaculis florum sordide rubris per exiguis. *Mich. gen.* 76. t. 38.
- Lichenoides pellucidum endiviæ foliis tenuibus crispis. *Dill. musc.* 143. t. 19. f. 31.
- Lichenoides sativæ tenue rufescens. *R. syn.* 77.
- Tremella *Lichenoides*, frondibus erectis planis margine crispo lacinulato. *Sp. pl.* 1625.
- Musco-fungus terrestris minor fuscus, foliis e latitudine crenatis musco innascens. *Hist. Ox. III.* 632.
f. 15. t. 7. *f.* 4.
Anglis, fringed Lichen.
Habitat in umbrosis, saxis et truncis arborum inter alios muscos passim. ☉. I—XII.
- Obs. Scutella magnitudine seminis rapi, orbiculata, concava, marginata, margine glauco, subintegro; fune plano fulvo.

53. LICHEN foliaceus membranaceus subrotundus lobatus rugosus atro-virens, scutellis confertis rufis. *scens.*
Fl. angl. 450.
- Lichen gelatinosus membranaceus exasperatus convolutus atroviridis, scutellis rubris. *Hall. hist.* 2037.

Lichenoides

Lichenoides gelatinosum membranaceum tenue nigricans. *Dill. musc.* 138. t. 19. f. 20.

Lichenoides saxatile membranaceum gelatinosum tenue nigricans. *R. syn.* 72.

Anglis, blackish Lichen.

Habitat in arborum truncis, rupibus, saxis et muris. ☉.
X—VI.

Burgesii. 54. LICHEN foliaceus subimbricatus crispus, scutellis elevatis muricato-crispis fundo depresso plano. *Syst. nat. ed.* 13. 827.

Anglis, Burges's Lichen.

Habitat in arborum truncis, in Dumfriesshire, Dr. Burges. 4. I—XII.

ciliaris. 55. LICHEN foliaceus erectiusculus; laciniis linearibus ciliatis, scutellis pedunculatis crenatis. *Sp. pl.* 1611. *Weis. crypt.* 62.

Lichen corniculatus planus, ramulis capillaribus scutellis petiolatis. *Hall. hist.* 1980.

Lichen cinereus arboreus, marginibus pilosis, major. *Vaill. paris.* 115. t. 20. f. 4.

Lichenoides arboreum foliosum cinereum, scutellis nigris, foliorum extremitatibus hispidis et pilosis. *Dill. giss.* 207. *R. syn.* 73.

Lichenoides hispidum majus et rigidius, scutellis nigris. *Dill. musc.* 150. t. 20. f. 45.

Musco-fungus arboreus cinereus scutellatus, marginibus pilosis. *Hist. Ox.* III. 634. f. 15. t. 7. f. 6.

⚡ Lichen foliaceus laciniatus albus: laciniis ascendentibus obtusis pilosis, scutellis sessilibus. *Scop. carn.* 1406. *Schreb. spicil.* 127.

Lichen fronde sempinnata canaliculata cinerea. *Hall. hist.* 2012.

Lichen cinereus arboreus, marginibus pilosis, minor. *Vaill. paris.* 115. t. 20. f. 5.

Lichenoides hispidum minus et tenerius, scutellis nigris. *Dill. musc.* 152. t. 20. f. 46.

Lichenoides saxatile et arboreum tenerius, foliis pilosis, scutellis in averfa foliorum superficie apparentibus. *R. syn.* 69.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Lichen alpinus cornua cervi referens, subtus anthracinus desuper cinereus, receptaculis florum amplioribus, intus fuscis. *Mich. gen.* 76. t. 38. f. 1.

Lichenoides cornutum amarum, superne cinereum, inferne nigrum. *Dill. musc.* 157. t. 21. f. 52.

Muscus amarus absinthii folio. *B. hist.* III. 764.
Anglis, branny Lichen.

Habitat in truncis arborum, in Arvoniam, et ad palos vivarii Croft-castle in Herefordia. *Littl. Brown.* in Westmorlandia et Cumberlandia observavi. 4. I—XII.

farina-
ceus.

58. LICHEN foliaceus erectus compressus ramosus farinaceus, lateribus verrucosis. *Sp. pl.* 1613. *Weis. crypt.* 64.

Lichen lacunofus complanatus, ramis acutissimis, orbiculis farinosis, marginalibus frequentissimis. *Hall. hist.* 1981.

Lichen cinereus angustior, scutis in marginibus segmentorum. *Vaill. paris.* 115. t. 20. f. 13, 14, 15.

Lichenoides segmentis angustioribus ad margines verrucosis et pulverulentis. *Dill. musc.* 172. t. 23. f. 63.

Lichenoides arboreum ramosum cinereo-candicans, segmentis angustioribus. *R. syn.* 76.

β Lichenoides arboreum ramosum, cinereo-virescens, segmentis angustioribus, circa margines crispis. *R. syn.* 76.

γ Lichenoides non tubulosum πλατυδαύφυλλοι, tuberculis pulverulentis donatum. *R. syn.* 66.

Anglis, mealy Lichen.

Habitat in truncis et ramis arborum. 4. I—XII.

calicaris. 59. LICHEN foliaceus erectus linearis ramosus lacunofus convexus mucronatus. *Sp. pl.* 1613. *Weis. crypt.* 67.

Lichen utrinque lacunatus farinosus et scutellifer. *Hall. hist.* 1983.

Lichen cinereus latifolius ramosus. *Vaill. paris.* 115. t. 20. f. 6.

Lichenoides coralliforme rostratum et canaliculatum. *Dill. musc.* 170. t. 23. f. 62.

Lichenoides

Lichenoides arboreum ramosum, angustioribus cinereo-virescentibus ramulis. *R. syn.* 76.

Musco-fungus arboreus, capitulis rostratis. *Hist. Ox. III.* 634. *f.* 15. *t.* 7. *f.* 5.

Anglis, channeled Lichen.

Habitat in rupibus et corticibus arborum. In Cambria.

∇. Lhwyd ; circa Slingfoldiam in Suffexia, et copiose ad rupes montis Steiperstone in Salopiensi comitatu. *Dill.* 4. I—XII.

60. LICHEN foliaceus erectiusculus lacunofus : subtus *prunastri*. tomentosus albus. *Sp. pl.* 1614. *Weis. crypt.* 73.

Lichen complanatus, utrinque lacunofus undique farinosus. *Hall. hist.* 1984.

Lichen pulmonarius mollior dichotomus, superne cinereus, inferne albus, receptaculis florum interna parte ferrugineis. *Mich. gen.* 75 *t.* 36. *f.* 3.

Lichen cinereus vulgatissimus cornua damæ referens. *Vaill. paris.* 115. *t.* 20. *f.* 11, 12.

Lichenoides cornutum bronchiale molle subtus incanum. *Dill. musc.* 160. *t.* 21. *f.* 55.

Lichenoides arboreum ramosum majus et mollius, colore candicante. *R. syn.* 75.

Lichen cornua damæ referens, angustifolius. *Vaill. paris.* *t.* 20. *f.* 7.

Lichenoides corniculatum candidum molle, segmentis angustis. *Dill. musc.* 159. *t.* 21. *f.* 54.

Anglis, thorn Lichen.

Habitat in truncis et ramis arborum. 4. I—XII.

61. LICHEN foliaceus erectus oblongus lanceolatus *fraxi*-lacunofus glaber, scutellis subpedunculatis. *Sp. pl. new.* 1614. *Weis. crypt.* 72.

Lichen utrinque lacunofus fronde latissima, scutellis creberrimis. *Hall. hist.* 1985.

Lichen pulmonarius cinereus mollior, in amplas lacinias divisus. *Tourn.* 549. *t.* 325. *f.* A. B.

Lichen pulmonarius cinereus durior, in amplas lacinias divisus. *Tourn.* 549. *Mich. gen.* 74. *t.* 36. *f.* 1.

Lichenoides longifolium rugosum rigidum. *Dill. musc.* 165. *t.* 22. *f.* 59.

Lichenoides

Lichenoides arboreum ramosum scutellarum majus et
rigidius, colore virescente. - *R. syn.* 75.

Musco-fungus quernus latifolius cinereus. *Hist. Ox.*
III. 634. *f.* 15. *t.* 7. *f.* 3, 4, 5, 9, 14, 19.

Anglis, alh Lichen.

Habitat in arborum truncis, præsertim Fraxinis et Quer-
ci. 4. I—XII.

pulmona- 62. LICHEN foliaceus laciniatus obtusus glaber, supra
rius. lacunofus, subtus tomentosus. *Sp. pl.* 1612. *Weis.*
crypt. 64.

Lichen lacunatus inferne gibbosus, reticulo farinoso,
scutellis lateralibus. *Hall. hist.* 1986.

Lichen s. pulmonaria arborea minor et angustifolia.
Mich. gen. 86. *t.* 45.

Lichen arborum. *Ger. em.* 1566.

Lichenoides pulmoneum reticulatum vulgare, mar-
ginibus peltiferis. *Dill. musc.* 212. *t.* 29. *f.* 113.

Lichenoides peltatum arboreum maximum. *R. syn.* 70.

Muscus pulmonarius. *B. pin.* 361. Lichen arbo-
reum. *Park.* 1311.

Anglis, Lungwort, or tree Lichen.

Habitat in truncis arborum. 4. I—XII.

juniperi- 63. LICHEN foliaceus laciniatus crispus fulvus, peltis
mus. lividis. *Fl. suec.* 1093. *Sp. pl.* 1614.

Lichen fulvus sinibus dædalcis lacinatus. *Fl. lapp.*
451. *Roy. lugdb.* 50.

Anglis, juniper Lichen.

Habitat in truncis et ramis arborum. 4. I—XII.

Obs. convenit in multitis, cum L. parietino sed differt
colore luteo-viridi; foliis laxis, erectiusculis, sublacu-
nosis; scutellis disco bruneis s. olivaceis.

ampulla- 64. LICHEN foliaceus planiusculus lobatus crenatus,
ceus. peltis globosis inflatis. *Sp. pl.* 1613.

Lichen foliis laciniatis, marginibus convolutis in ve-
siculam abeuntibus. *Roy. lugdb.* 509.

Lichenoides tinctorum glabrum vesiculosum. *Dill.*
musc. 188. *t.* 24. *f.* 82.

Lichenoides

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

- Lichen pallide viridis rugosus margine undulatus.
Sp. pl. 1614. *Guet. stamp.* 1. t. 31. *Weis. crypt.* 74.
 Lichen fronde rotunde lobata rugosa, inferne atra et
 aspersa. *Hall. hist.* 2005.
 Lichen pulmonarius saxi et arboribus adnascens ma-
 jor, inferne nigricans, superne e sulphureo cinereus
 receptaculis florum amplioribus, interne sordide et
 obsolete viridis. *Mich. gen.* 89. t. 48. f. 1.
 Lichenoides caperatum rufaceo expansum e sulphureo
 virens. *Dill. musc.* 193. t. 25. f. 97.
 Lichenoides crusta foliosa, ex cinereo et luteo vire-
 scente, inferne nigra et lævi. *R. syn.* 73.
 Musco-fungus lichenoides, crustæ modo adnascens
 major cinereus. *Hist. Ox.* III. 633. y. 15. t. 7,
 f. 1.
Anglis, rose Lichen.
 Habitat in saxi et arboribus passim. 4. I—XII.

- laciniatus.* 68. LICHEN foliaceus repens sinuato-lobatus glaber,
 lobis sinuatis, scutellis fulvis.
 Lichen fronde rotunde lobata, inferne scabra, arbus-
 culis obscuris. *Hall. hist.* 2004.
 Lichen foliaceus repens lobatus, supra cinereus, infra
 roseus tomentosus, scutellis rubris. *Scop. carn.*
 1393.
 Lichen pulmonarius arboribus adnascens maximus
 inferne obscurus, desuper e glauco cinereus; recep-
 taculis florum rubris; seminibus nigricantibus
Mich. gen. 88. t. 46.
 Lichenoides subglaucum cumatile, foliis tenacibus,
 eleganter laciniatis. *Dill. musc.* 197. t. 26. f. 99.
Anglis, jagged Lichen.
 Habitat in truncis et ad radices arborum; secus viam inter
 Carno et Main-Lloyne in Mervinia. *Dill.* 4.
 I—XII.

- herbaceus.* 69. LICHEN foliaceus repens obtuse lobatus crenatus
 glaber herbaceus, scutellis rufis.
 Lichen fronde rotunde lobata virente, inferne scabra.
Hall. hist. 2002.

Lichenoides

Lichenoides læte-virens, scutellis fulvis. *Dill. musc.* 195. t. 25. f. 98.

Lichenoides arboreum cinereo-virens, tenue et læve ubique, scutellis minoribus. *R. syn.* 73.

Musco-fungus lichenoides arborum Hibernicus scutellatus. *Hist. Ox. III.* 633. f. 15. t. 7. f. 3.

Anglis, green Lichen.

Habitat in *juvis, truncis et ad radices arborum, et ad sepes umbrosas prope Bishops-castle in agro Salopiensi; juxta viam inter Carno et Main-L'oyne in Mervinia. Dill. Prope Wakehurst in Suffexia. D. Rand. Prope Ivy-bridge in Devonia, et in comitatibus Eboracensi et Westmorlandico observari. 4. I—XII.*

70. LICHEN foliaceus repens subrotundus sublobatus *verrucosus* crenatus rugosus verrucosus, verrucis farinaceis, *scutellis minutissimis fulvis.*

Lichen scrobiculatus. *Scop. carn.* 1319.

Lichen fronde obtuse lobata, inferne scabra, maculis rotundis varia. *Hall. hist.* 1989.

Lichen pulmonarius arboribus adnascens, inferne obscurus, desuper lacunatus, ex glauco et cinereo flavescens, receptaculis florum rubris. *Mich gen.* 90. t. 49.

Lichenoides pulmonicum villosum, superficie scrobiculata et peltata. *Dill musc.* 216. t. 29. f. 114.

Lichenoides arboreum foliosum cinereum et sinuatum, inferne scabrum. *R. syn.* 77.

Anglis, warty Lichen.

Habitat in *truncis arborum et saxis grandioribus passim. 4. I—XII.*

Obs. *Scutellæ minutissimæ, hemisphærica, cavæ, lutescentes, fundo fulvæ sunt.*

**** *Coriacei peltati.*

71. LICHEN coriaceus repens lobatus, peltis marginalibus posticis. *Syst. nat.* 711. *Sp. pl.* 1615. *resupinatus.*

Lichen pulmonarius major ex obscuro purpureus: inferne ex albo rufescens, receptaculis florum rubris

N n

amphoribus,

amplioribus, ad iatera oblongis. *Mich. gen.* 86.
t. 44. f. 42.

Lichenoides fuscum, peltis posticis ferrugineis. *Dill. musc.* 206. t. 28. f. 105.

Lichenoides saxatile fuscum, peltis in averfa foliorum superficie locatis. *R. syn.* 77.

Anglis, reversed Lichen.

Habitat in truncis arborum saxis et rupibus. 4. I—XII.

venosus. 72. LICHEN coriaceus repens ovatus planus, subtus venosus villosus, peltis marginalibus horizontalibus. *Syst. nat.* 711. *Sp. pl.* 1615. *Weis. crypt.* 82.

Lichen flabelliformis, subtus reticulatis, scutellis marginalibus planis. *Hall. hist.* 1993.

Lichen pulmonarius minimus inferne albus et niger reticulatus, superne e cinereo virescens, receptaculis florum nigricantibus ad latera oblongis. *Mich. gen.* 85. t. 44. f. 3. et 5.

Lichenoides parvum virescens, peltis nigricantibus. *Dill. musc.* 208. t. 28. f. 109.

Anglis, vein'd Lichen.

Habitat in saxis et rupibus humidiusculis, circa Dunkeld et alibi in Scotia. D. Oakes. 4. I—XII.

caninus. 73. LICHEN coriaceus repens lobatus obtusus planus, subtus venosus villosus, pelta marginali ascendente. *Syst. nat.* 712. *Sp. pl.* 1616. *Weis. crypt.* 78.

Lichen fronde subrotunde lobata, inferne reticulata, peltis convexo-concavis. *Hall. hist.* 1988.

Lichen pulmonarius saxatilis digitatus. *Vaill. paris.* t. 21. f. 16.

Lichen terrestris cinereus. *R. Cat. angl.* 185.

Lichenoides digitatum cinereum, lactucæ foliis sinuosis. *Dill. musc.* 200. t. 27. f. 102.

Lichenoides peltatum terrestre cinereum majus, foliis divis. *R. syn.* 76.

Lichen fronde lobata obtusa avenia margine sinuata subtus villosa, peltis subrotundis. *Neck. meth. musc.* 79.

Lichen fronde obtuse lobata, superne verrucosa, inferne villosa. *Hall. hist.* 1990.

Lichenoides

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- horizontalis.* 76. LICHEN coriaceus repens planus subtus avenius, peltis marginalibus horizontalibus. *Fl. angl.* 543. *Lin. mantiss.* I. 132.
Lichen fronde rotunde lobata, peltis planissimis. *Hall. hist.* 1991.
Lichen pulmonarius maximus e cinereo rufescens inferne subrufus, receptaculis florum rubris ad latera oblongis. *Mich. gen.* 85. t. 44. f. 1.
Lichen pulmonarius minor, inferne obscurus, desuper cinereus, receptaculis florum nigricantibus circinnatis. *Mich. gen.* 85. t. 44. f. 6.
Lichenoides subfuscum, peltis horizontalibus. *Dill. musc.* 205. t. 28. f. 104.
Anglis, flat Lichen.
Habitat in rupibus et saxis humidis et ad radices arborum passim. 4. I—XII.

- saccatus.* 77. LICHEN coriaceus repens subrotundus, peltis depressis subtus saccatis. *Sp. pl.* 1616. *Fl. dan.* 532. f. 3.
Lichen fronde coriacea rotunde lobata, scutellis immerfis. *Hall. hist.* 1995.
Lichen pulmonarius alpinus terristris glauco-virescens, receptaculis florum fuscis. *Mich. gen.* 95. t. 52.
Lichenoides lichenis facie, peltis acetabulis immerfis. *Dill. musc.* 221. t. 30. f. 121.
Anglis, pitted Lichen.
Habitat in rupibus et saxis, in rupibus Chedder-rocks dictis in Somersetia; circa Llandberris et alibi in Cambria, in boreali parte Britanniae frequens. 4. I—XII.

- croceus.* 78. LICHEN coriaceus repens subrotundus planus subtus venosus villosus croceus, peltis sparsis adnatis. *Sp. pl.* 1616. *Fl. dan.* 263.
Lichen foliis subrotundis planis lævissime incis, calycibus orbiculatis disco folii adnatis. *Fl. lapp.* 443. t. 11. f. 3.
Lichen fronde rotunde lobata plana subtus venosa minutata. *Hall. hist.* 1994.

Lichenoides

Lichenoides subtus croceum, peltis appressis. *Dill. musc.* 211. t. 30. f. 120.

Anglis, saffron Lichen.

Habitat in rupibus et saxosis et nemorosis montosis, prope Money-musk, Aberdeenshire. 4. I—XII.

***** Umbilicati, squalentes quasi fuligine.

79. LICHEN umbilicatus gibbus punctatus, subtus *miniatus*. fulvus. *Sp. pl.* 1617. *Fl. dan.* 532. f. 1.

Lichen fronde imbricata rotunde lobata cinerea punctata, inferne ochrea subaspera. *Hall. hist.* 1999. t. 47. f. 2.

Lichen pulmonarius saxatilis e cinereo fuscus, minimus. *Mich. gen.* 101. *ord.* 36. t. 54. f. 1.

Lichenoides coriaceum nebulosum cinereum punctatum subtus fulvum. *Dill. musc.* 223. t. 30. f. 127.

β Lichen pulmonarius saxatilis sordide virens, foliis minimis. *Mich. gen.* 101. *ord.* 36. t. 54. f. 2.

Lichenoides imbricatum lividum. *Dill. musc.* 224. t. 30. f. 128. *Fl. dan.* 532. f. 2.

Anglis, cloudy Lichen.

Habitat in rupibus et saxis grandioribus passim. 4. I—XII.

80. LICHEN umbilicatus subtus lacunofus, fursure *pustulatus*. nigro aspersus. *Sp. pl.* 1617. *Fl. dan.* 597. f. 2.

Lichen fronde cinerea inferne lacunosa, arbusculis superne nigris. *Hall. hist.* 1996.

Lichen pulmonarius saxatilis, inferne reticulatus et lacunofus superne cinereus ac verrucosus, receptaculis florum et seminibus nigricantibus et veluti deustus. *Mich. gen.* 89. t. 47.

Lichen crustæ modo saxis adnascens verrucosus cinereus et veluti deustus. *Vaill. paris.* 115. t. 20. f. 9.

Lichenoides pustulatum cinereum et veluti ambustum. *Dill. musc.* 226. t. 30. f. 131.

Anglis, blistered Lichen.

Habitat in rupibus, saxis et muris, in rupibus meridiem spectantibus, subter arcem Keven-lees-castle dictam, in comitatu Radnorienſi. Littl. Brown; prope Halifax, D. Bolton; in montibus Scotiſis paſſim. 4. I—XII.

- deustus.* 81. LICHEN umbilicatus undique lævis. *Sp. pl.* 1618. Lichen fronde rotunde lobata inferne nigerrima glebosa. *Hall. hiſt.* 1998. Lichen pulmonarius saxatilis e cinereo fuscus minimus. *Vailt. paris.* 116. t. 21. f. 14. Lichenoides coriaceum cinereum, peltis atris compressis. *Dill. muſc.* 218. t. 29. f. 117. Lichenoides saxatile, foliis minus diviſis cinereo-fuscis. *R. ſyn.* 73. *Anglis,* sooty Lichen. Habitat in rupibus et saxis; in rupe Sancti Vincentii prope Bristolium. D. Dare. circa Llanberris. D. Davies. 4. I—XII.

- polyrrhizos.* 82. LICHEN umbilicatus polyphyllus utrinque lævis polyrrhizos. *Sp. pl.* 1618. *Fl. dan.* 471. f. 3? Lichenoides rugosum durum pullum, peltis atris verrucosis. *Dill. muſc.* 220. t. 30. f. 118. *Anglis,* finged Lichen. Habitat in rupibus et saxis, in rupe Sancti Vincentii prope Bristolium; circa Llanberris. D. Davies, et in montibus Scotiſis. 4. I—XII.

- velleus.* 83. Lichen umbilicatus subtus hirsutissimus. *Sp. pl.* 1617. Lichen fronde coriacea inferne pustulata villosa, scutellis atris. *Hall. hiſt.* 1997. Lichenoides pullum superne glabrum, inferne nigrum et cirrosum. *Dill. muſc.* 226. t. 30. f. 130. β Lichenoides coriaceum, latissimo folio umbilicato et verrucoso. *Dill. muſc.* 545. t. 82. f. 5. *Anglis,* fleecy Lichen. Habitat in rupibus et saxis circa Llanberris, Dill. in Clark's Park et Paradise prope Money-musk, Aberdeenshire. 4. I—XII.

84. LICHEN

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

CRYPTOGAMIA ALGÆ.

- Lichenoides tubulosum pyxidatum proliferum. *Dill. giss.* 204. *musc.* 80. t. 14. f. 6. D—H. *R. syn.* 60.
- γ Lichen scyphiger infundibulorum fungis fuscis. *Hall. hist.* 1919.
- Lichen pyxidatus major, acetabulo fimbriato et tuberculoso. *Vaill. par.* 115. t. 21. f. 11.
- Lichenoides tubulosum pyxidatum, tuberculis fusconigris. *Dill. giss.* 204. *musc.* 81. t. 14. f. 6. c. proliferum. *Dill. giss.* 204. *musc.* 81. t. 14. f. 6. K—M. *R. syn.* 69.
- fimbriatus. δ Lichen scyphifer simplex denticulatus, stipite cylindrico. *Sp. pl.* 1619.
- Lichen scyphiger infundibulis ferratis. *Hall. hist.* 1915. coronatis. *Hall. hist.* 1916.
- Lichen pyxidatus minor. *Vaill. paris.* 115. t. 21. f. 6. *Mich. gen.* 82. t. 41. f. 4, 5.
- Lichenoides tubulosum pyxidatum, marginibus serratis. *Dill. giss.* 204. *R. syn.* 69.
- Coralloides scyphiforme gracile, marginibus ferratis. *Dill. musc.* 84. t. 14. f. 8.
- Lichen infundibulis ferratis proliferis. *Hall. hist.* 1924.
- Lichen infundibulis proliferis ferratis foliosis, et verrucosis. *Hall. hist.* 1925.
- Lichenoides pyxidatum proliferum, marginibus ferratis. *R. syn.* 69.
- filiformis. ζ Lichen scyphifer simplex integerrimis filiformis, tuberculis fuscis. *Fl. angl.* 456.
- Lichen corniculo simplici apice depresso scutellato. *Hall. hist.* 1908.
- Lichenoides tubulosum pyxidatum exiguum, fusco-virens. *Dill. giss.* 206. *Ry. syn.* 70.
- Coralloides scyphis gracilibus tubiformibus, pedicularis folio. *Dill. musc.* 85. t. 14. f. 10.
- exiguus. η Lichen stipite brevissimo scyphifero.
- Coralloides scyphis humilibus intus fuscis. *Dill. musc.* 86. t. 14. f. 11.
- foliaceus. θ Lichen foliis suberectis laciniatis crispis scyphiferis, scyphis brevissimis conicis. *Fl. angl.* 457.
- Lichen infundibulis explanatis fungiferis. *Hall. hist.* 1934.

- Lichen fronde semipinnata rotunde lobata, ex medio scyphigera. *Hall. hist.* 1935.
- Lichen pyxidatus, endiviæ crispæ folio prolifer, acetabulorum oris crispis. *Vaill. paris.* 115. t. 21. f. 3.
- Lichen pyxidatus, endiviæ crispæ folio prolifer, superne flavescens, inferne albidus, acetabulorum oris crispis. *Mich. gen.* 82. t. 42. f. 1, 2.
- Lichenoides cartilaginofum, tubulis et pyxiduis exiguis. *R. syn.* 70.
- Coralloides scyphiforme, foliis alcicorniformibus cartilaginosis. *Dill. musc.* 87. t. 14. f. 12.
- Musco-fungus terrestris minor crispus, foliis superne e flavo virescentibus, subtus albicantibus. *Hist. Ox. III.* 632. f. 15. t. 7. f. 3.
- Lichen scyphifer simplex integerrimis, stipite cylindrico, tuberculis coccineis. *Sp. pl.* 58. **cocciferus.**
- Lichen scyphiger infundibulorum fungis coccineis. *Hall. hist.* 1918.
- Lichen pyxidatus, acetabulorum oris coccineis et tumentibus. *Vaill. paris.* 115. t. 21. f. 4, 10, 11. *Mich. gen.* 82. t. 41. f. 3, 4.
- Lichenoides tubulosum pyxidatum tuberculis amæne coccineis. *R. syn.* 69.
- Coralloides scyphyforme, tuberculis coccineis. *Dill. musc.* 82. t. 14. f. 7.
- Musco-fungus pyxidatus, calyce altero alteri innato, apicibus nonnunquam coccineis. *Hist. Ox. III.* 632. f. 15. t. 7. f. 4.
- Lichen scyphifer simplex folio brevior, tuberculis coccineis. *Sp. pl.* 1691. **cornucopioides.**
- Lichen infundibulis proliferis, foliosis et fungosis. *Hall. hist.* 1929.
- Lichenoides pyxidatum, marginibus eleganter foliatis. *R. syn.* 69.
- Coralloides scyphiforme, marginibus foliatis et radiatis. *Dill. musc.* 85. t. 14. f. 9.
- Lichen scyphifer simplicifolius subventricosus, calycibus integris. *Sp. pl.* 1620. **cornutus.**
- Lichen corniculis simplicissimis acutis farinosis. *Hall. hist.* 1902.

- Lichen cornubus acutis farinosis subramosis. *Hall. hist.* 1903.
- Coralloides vix ramosum, scyphis obscuris. *Dill. musc.* 90. t. 15. f. 14.
- Lichenoides tubulosum cinereum, non ramosum. *R. syn.* 68
- deformis. μ Lichen scyphifer simpliciusculus subventricosus, calycibus dentatis. *Sp. pl.* 1620. *Fl. lapp.* t. 11. f. 5.
- Lichen pyxidatus difformis, acetabulis cornigeris. *Hall. hist.* 1938.
- Lichen pyxidatus teres, acetabulis minoribus repandis. *Mich. gen.* 80. t. 41. f. 1.
- Lichenoides tubulosum minus ramosum cauliculis crassioribus. *R. syn.* 68.
- Lichenoides tubulosum magis ramosum, maxime difforme. *R. syn.* 68.
- Coralloides crassius subincanum, calycibus dentatis. *D. ll. hist.* 95. t. 15. f. 18.
- ventrico-
sus. Lichen scyphifer ramosissimus ventricosus, scyphis dentatis, tuberculis fuscis. *Fl. angl.* 457.
- Lichen infundibulis proliferis palmatis serratis et punctatis. *Hall. hist.* 1937.
- Lichenoides tubulosum cinereum, valde crustaceum, ramulis brevioribus et crebrioribus ex acetabulis enascentibus. *Dill. giss.* 203. *R. syn.* 68.
- Coralloides cornucopioides incanum, scyphis cristatis. *Dill. musc.* 94. t. 15. f. 17.
- Muscus cornucopioides cristatus alpinus saxatilis. *Bocc. musc.* 11. 21. t. 8.
- digitatus. ξ Lichen scyphifer ramosissimus, ramis cylindricis, calycibus integris nodosis. *Sp. pl.* 1620. *Weis. crypt.* 91.
- Lichen infundibulis palmatis, fungulis coccineis. *Hall. hist.* 1939.
- Lichenoides coralliforme, apicibus coccineis. *R. syn.* 68.
- Coralloides ramulosum, tuberculis coccineis. *Dill. musc.* 96. t. 15. f. 19.
- Coralloides parum ramosum, tuberculis fuscis. *Dill. musc.* 97. t. 13. f. 20.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Lichen coralloides tubulosus albidus major et mollior, caulibus crassioribus minus ramosis, receptaculis florum perexiguis rufescentibus. *Mich. gen.* 79. t. 40. f. 2.

Lichenoides tubulosum, cauliculis mollioribus et crassioribus, majus *R. syn.* 67.

Coralloides perforatum majus, molle et crassum. *Dill. musc.* 98. t. 16. 5. 21.

Musco-fungus ceranoides mollior et elatior albidus tubulosus. *Hist. Ox. III.* 633. f. 15. t. 7. f. 7.

spinosus. γ Lichen fruticulosus tubulosus ramosissimus spinosus, ramulis tuberculatis subdigitatis. *Fl. angl.* 459.

Lichen fruticosus vage ramosus foliosus, extremis ramis palmatis fungiferis. *Hall. hist.* 1952.

Lichen coralloides tubulosus major, ex cinereo fusco-virens caulibus crassioribus et durioribus, minus ramosis, extremitatibus palmatis, receptaculis florum exiguis rufescentibus. *Mich. gen.* 79. t. 40. f. 3, 5.

Lichenoides tubulosum ramosum rigidius, majus et crassius cinereo-fuscum. *R. syn.* 67.

Coralloides sparsum, caulibus tortuosis et spinosis. *Dill. musc.* 101. t. 16. f. 25.

Anglis, perforated Lichen.

Habitat in ericetis et saxosis, montosis. 4. I—XII.

Obf. medius inter uncialem et subulatum est.

subulatus. 88. LICHEN fruticulosus subdichotomus, ramis simplicibus subulatis. *Sp. pl.* 1620. *Weis. crypt.* 97.

Lichen caule fruticoso erecto, ramis adscendentibus dichotomis. *Hall. hist.* 1955.

Lichenoides tubulosum cinereum minus crustaceum minusque ramosum. *R. syn.* 67.

Coralloides corniculis longioribus et rarioribus. *Dill. musc.* 102. f. 26.

Muscus corniculatus. *Tab. ic.* 800. *Ger. em.* 1560. *Park.* 1308.

furcatus. β Lichen fruticulosus tubulosus ramosus, ramis erectis furcatis. *Fl. angl.* 458.

Lichen caule fruticoso subfolioso ramis adscendentibus repetito bifurcatis punctiferis. *Hall. hist.* 1954.

Lichen

Lichen coralloides tubulosus medius e cinereo viridans ramulis erectis nonnihil foliosis, extremitatibus apertis et palmatis, receptaculis florum rufescentibus exiguis. *Mich. gen.* 80. t. 40. f. 4. et *D.*

Lichenoides tubulosum cinereum, ramosius et crustaceum. *R. syn.* 67.

Coralloides corniculis brevioribus et crebrioribus. *Dill. musc.* 104. t. 16. f. 27.

Musco-fungus montanus corniculatus. *Hist. Ox.* III. 632. f. 15. t. 7. f. 1. *Vaill. paris.* t. 26. f. 7.

Anglis, horned, β forked Lichen.

Habitat in ericetis et pascuis montosis. 4. I—XII.

89. LICHEN fruticulosus perforatus ramosissimus, ramulosis nutantibus. *Sp. pl.* 1620. *Weis. crypt.* 95. *Fl. dan.* 180.

Lichen fruticosus farinosus, furculis innumeris nutantibus. *Hall. hist.* 1957, alopetris.

Lichen coralloides tubulosus major ramosissimus receptaculis florum rufescentibus perexiguis. *Mich. gen.* 79. t. 40. f. 1.

Lichenoides tubulosum ramosissimum, fruticuli specie, candicans. *R. syn.* 66.

Coralloides montanum fruticuli specie, ubique candicans. *Dill. musc.* 107. t. 16. f. 29.

Musco-fungus coralloides montanus ramosissimus cinereus vulgaris. *Hist. Ox.* III. 633. f. 15. t. 7. f. 9.

8 Lichen caule ramoso, alis perforatis, ramis capitatis. *Roy. lugdb.* 511. sylvaticus.

Lichen caule fruticoso folioso, ramis subnutantibus, minute multifidis fungiferis. *Hall. hist.* 1956.

Lichenoides tubulosum ramosissimum, fruticuli specie, candicans, corniculis rufescentibus. *R. syn.* 67.

Coralloides fruticuli specie candicans, corniculis rufescentibus. *Dill. musc.* 110. t. 16. f. 30.

Muscus terrestris coralloides erectus, cornibus rufescentibus. *B. pin.* 361. *Park.* 1310.

Anglis, Rhendeer Lichen.

Habitat in ericetis, pascuis montosis et sylvis. 4. I—XII.

90. LICHEN

- pschalis.* 90. LICHEN fruticulosus solidus, tectus foliolis crustaceis. *Sp. pl.* 1621. *Weis. crypt.* 98. *Fl. dan.* 151. Lichen caule fruticoso, fungifero, verrucis calcariis tecto. *Hall. bist.* 1963.
Lichen alpinus glaucus ramosus botryoides. *Sch. it. alp.* 11. 137. t. 19. f. 4. *Mich. gen.* 78. t. 35. f. 5—8.
Lichenoides non tubulosum cinereum ramosum, totum crustaceum. *R. syn.* 66.
Coralloides crispum et botryforme alpinum. *Dill. musc.* 114. t. 17. f. 33.
Musco-fungus coralloides fruticosior et lignosior. *Hist. Ox.* III. 635. f. 15. t. 7. f. 12.
Anglis, crisp Lichen.
Habitat in rupibus et saxosis montosis. 4. I—XII.

- fragilis.* 91. LICHEN fruticulosus solidus lævis, ramulis teretibus obtusis, tuberculis terminalibus globosis crevis.
Lichen fruticulosus solidus ramulis teretibus obtusis. *Sp. pl.* 1621. *Fl. lapp.* t. 11. f. 1.
Lichen caule fruticoso ramo erecto densissime cespitoso. *Hall. bist.* 1962.
Lichenoides non tubulosum ramosissimum, fruticuli specie, cinereo-fuscum. *R. syn.* 65.
Coralloides alpinum corallinae minoris facie. *Dill. musc.* 116. t. 17. f. 34.
globiferus. β Lichen fruticulosus solidus ramosissimus ramulis teretibus acutis furcatis, tuberculis terminalibus globosis. *Fl. angl.* 460. *Lin. Mantiss.* 133.
Lichen fruticosus coralloides non tubulosus cinereus ramosissimus, receptaculis florum sphaericis concoloribus. *Mich. gen.* 103. t. 39. f. 6.
Lichenoides non tubulosum, ramulis scutellis nigris terminatis. *R. syn.* 66.
Coralloides cupressiforme, capitulis globosis. *Dill. musc.* 117. t. 17. f. 35.
Anglis, brittle, β globe Lichen.
Habitat in rupibus, saxis et ericetis montosis. 4. I—XII.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

- floridus.* 95. LICHEN filamentosus ramosissimus erectus, tuberculis farinaceis sparsis, scutellis radiatis.
Lichen filamentosus ramosus erectus, scutellis radiatis. *Sp. pl.* 1624. *Weis. crypt.* 102.
Lichen barbatus, ramis teretibus, furculis parallelis simplicibus, scutellis radiatis. *Hall. hist.* 1972.
Usnea vulgatissima tenuior et brevior cum orbiculis. *Dill. musc.* 69. t. 13. f. 13.
Muscus ramosus floridus. *Tab. ic.* 808. *Ger. em.* 1560.
Muscus arboreus cum orbiculis. *B. pin.* 361. *R. syn.* 65.
Musco-fungus arborum capillaceus, scutis amplis, per ambitum pilis radiatis. *Hist. Ox. III.* 635. f. 15. t. 7. f. 14.
- hirtus.* β Lichen filamentosus ramosissimus erectus, tuberculis farinaceis sparsis. *Sp. pl.* 1623. *Weis. crypt.* 101.
Lichen barbatus, teretibus cauliculis, ramis simplicibus parallelis. *Hall. hist.* 1973.
Usnea vulgatissima tenuior et brevior sine orbiculis. *Dill. musc.* 67. t. 13. f. 12.
Muscus ramosus. *Tab. ic.* 807. *Ger.* 1372. *R. syn.* 65.
Anglis, flowering Lichen.
Habitat in sylvis et sepibus frequens. 4. I—XII.
Obs. variat colore fulvo et luteo.
- plicatus.* 96. LICHEN filamentosus pendulus, ramis implexis, scutellis radiatis. *Sp. pl.* 1622. *Mat. med.* 495.
Lichen teres pendulus exasperatus ramis descendens, scutellis radiatis. *Hall. hist.* 1971.
Usnea vulgaris, loris longis implexis. *Dill. musc.* 56. t. 11. f. 1.
Muscus arboreus, Usnea officinarum. *B. pin.* 361. *R. syn.* 64.
Muscus quercinus. *Ger. em.* 1558.
Muscus arboreus vulgaris et quercinus. *Park.* 1311.
Anglis, tree Lichen, or Mols.
Habitat in ramis arborum; in sylvis densis passim. 4. I—XII.

97. LICHEN filamentosus pendulus subarticulatus, *ta-barbatus*.
 mis patentibus. *Sp. pl.* 1622. *Weis. crypt.* 100.
 Lichen barbatus, trunco articulato lævi. *Hall. hist.*
 1970.
 Usnea barbata, loris tenuibus fibrosis. *Dill. musc.* 63.
t. 12. f. 6.
 Muscus capillaceus longissimus. *B. pin.* 361.
Anglis, bearded Lichen.
 Habitat in ramis arborum, in sylva dicta Forest of Dean
 in Glocestriensi comitatu, et in sylva prope Bishops-
 castle in Salopia. *Dill. musc.* 4. I—XII.
98. LICHEN filamentosus articulatus, ramis tenuissi-*articula-*
 mis punctatis. *Sp. pl.* 1623. *Roy. lugdb.* 513. *tus.*
 Usnea capillacea et nodosa. *Dill. musc.* 60. *t. 11. f. 4.*
 Muscus arboreus nodosus. *B. pin.* 361. *R. syn.* 65.
f. geniculatus. Park. 1311.
Anglis, jointed Lichen.
 Habitat in ramis arborum in sylvis passim; in sylva ad
 villam Stokeſ-church; prope Burnley in Lancastria.
Dill. musc. 4. I—XII.
99. LICHEN filamentosus ramosus subpendulus impli-*jubatus.*
 cato-flexuosus, axillis compressis.
 Lichen filamentosus pendulus, axillis compressis. *Sp.*
pl. 1622. *Weis. crypt.* 101.
 Lichen barbatus pendulus, filamentis nigris prolixis.
Hall. hist. 1968.
 Usnea jubata nigricans. *Dill. musc.* 67. *t. 12. f. 7.*
 Muscus corallinus saxatilis fœniculaceus. *R. syn.* 65.
 Musco-fungus trichoides e nigridine cinereus jubæ ad
 instar e saxis et arboribus dependens. *Hist. Ox.* III.
 633.
- β Lichen filamentosus ramosus divaricatus decumbens chalybei-
 implicato-flexuosus. *Sp. pl.* 1623. *Fl. dan.* 261. *formis.*
 Lichen fruticosus diffusus, caulibus vagis flexuosis ri-
 gidis teretibus. *Hall. hist.* 1969.
 Usnea rigida horsum versus extensa. *Dill. hist.* 66.
t. 13. f. 10.
 Muscus caule rigido instar fili chalybei. *R. syn.* 65.
Anglis, wiry Lichen.

Habitat in rupibus et arborum truncis; in Wallia, et in boreali parte Britanniae passim. 4. I—XII.

scaber. . 100. LICHEN filamentosus ramosissimus decumbens implexus scaber, scutellis concavis integerrimis. Usnea cæspitosa exilis capillacea atra. *Dill. musc.* 66. t. 13. f. 9.

Anglis, scabrous Lichen.

Habitat in rupibus et saxis grandioribus circa Llanberris copiose. in boreali parte Angliæ passim. 4. I—XII.

Obs. Scutellæ magnitudo fere seminis Papaveris albi, hemisphæricæ, truncatæ, marginatæ, atra fundo atrofusco, margine integerrima.

lanatus. 101. LICHEN filamentosus ramosissimus decumbens implicatus, ramulis brevissimis divaricatis.

Lichen fruticosus minimus flexuosus, cauliculis nigris capillaceis. *Hall. hist.* 1967.

Usnea lanæ nigræ instar saxi adhærens. *Dill. hist.* 66. t. 13. f. 8.

Muscus coralloides lanæ nigræ instar saxi adhærens. *R. syn.* 65.

Anglis, woolly Lichen.

Habitat in rupibus et saxosis, in Cornubia D. Stephens, in rupibus Cambriæ circa Borth, miliaris distantia a Bangor S. Brewer; in septentrionali parte Britanniae passim. 4. I—XII.

pubescens. 102. LICHEN filamentosus ramosissimus decumbens implexus nitidus. *Sp. pl.* 1623.

Anglis, pubescent Lichen.

Habitat in rupibus et saxosis in Westmorlandia passim. 4. I—XII.

Obs. affinis præcedenti sed minor.

TREMELLA. 1204.

Fructificationes vix manifestæ, in corpore gelatinoso.

juniperi- 1. TREMELLA sessilis membranacea auriformis fulva. *Sp. pl.* 1625.

na.

Agaricus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Lichen gelatinosus niger undosus et crispus. *Hall. hist.* 2038.

Lichen arboreus membranaceus nigricans. *Buxb. comm. petrop. III. 273. t. 14. f. 3.*

Anglis, tree Tremella, or Witches Butter.

Habitat in truncis arborum. ☉. IX—V.

Nostoc. 5. TREMELLA sessilis subrotunda plicata undulata fusco-virens.

Tremella sessilis plicata undulata. Sp. pl. 1625. Weis. crypt. 29.

Tremella terrestris sinuosa pinguis et fugax. Dill. musc. 52. t. 10. f. 14.

Linkia terrestris gelatinosa membranacea vulgatissima. Mich. gen. 126. t. 67. f. 1.

Lichen gelatinosus plicatus undulatus, laciniis crispis granulosis. Hall. hist. 2041.

Ulva terrestris pinguis et fugax. R. syn. 64.

Nostoc paracelsi. Act. paris. 1708. p. 228.

Anglis, jelly Tremella.

Habitat in pratis et pascuis post pluvias. ☉. I—XII.

utricula- 6. TREMELLA sessilis tuberculosa gelatinosa, tuberculis cavis.

Tremella fluviatilis gelatinosa et uterculosa. Dill. musc. 14. t. 10. f. 16?

Linkia palustris gelatinosa saxis adnascens ex obscuro fulvo et concava vesicam referens. Mich. gen. 126. t. 67. f. 2.

Anglis, bladder Tremella.

Habitat in rivulis supra lapides, circa Pentir et Llanberris, Dill. in rivulis montosis in comitatibus Westmorlandico et Cumberlandica. Prope Tidewell in comitatu Derbiensi. 4. I—XII.

Desc. Rupes et lapides sub aqua late tegit, hæc Tremella, substantia gelatinosa, viridia, rigidiuscula, fragilis, nunc. semi, nunc ad duas uncias crassa; superficies nitidiuscula, tuberculosa, subinde lævis, tuberculis intus cavis, variæ magnitudinis Pisi vel nucis Avellanæ.

7. TREMELLA

7. TREMELLA tuberculosa solida rugosa. *Sp. pl. verrucosa.*
1625.
Tremella subrotunda sinuosa difformis gelatinosa. *Lin.*
It. w-goth. 193.
Anglis, warty Tremella.
Habitat in rivulis et fontibus limpidis supra lapides. 24.
I—XII.
Obs. Substantia valde tenerrima lacte viridis.
8. TREMELLA hemisphærica sparsa. *Sp. pl. 1626. hemis-*
Anglis, hemisphæric Tremella. phærica.
Habitat in confervis, Fucis et rupibus submarinis. 24.
I—XII.
9. TREMELLA subrotunda sinuosa difformis gelatino-difformis.
sa. Sp. pl. 1626. It. w-goth. 193.
Anglis, sea Tremella.
Habitat in confervis et Fucis et rupibus submarinis. 24.
I—XII.
10. TREMELLA rotunda imbricata livida. *Sp. pl. adnata.*
1627. *It. scan. 323. Fl. suec. 1140.*
Anglis, livid Tremella.
Habitat in rupibus et lapidibus submarinis. 24. I—XII.
11. TREMELLA subglobosa sessilis solitaria glabra. *purpurea.*
Sp. pl. 1626. Fl. suec. 1142.
Sphæria convexa miniata hirsuta. Hall. hist. 2189.
Lichenoides tuberculosum amæne purpureum. Dill.
giff. 206. musc. 127. t. 18. f. 6. R. syn. 71.
Lycogala globosum rubrum, grani panici magnitudine
et forma. Mich. gen. 216. t. 95. f. 3.
Noctoc granulorum coccineum arboribus adnascens.
Vaill. paris. 144.
Anglis, purple Tremella.
Habitat in arborum ramis moribundis et emortuis. ☉.
IX—V.
12. TREMELLA sessilis gelatinosa difformis albida. *albica.*
Echinus gelatinosus crytallinus. Hall. hist. 2319?
Anglis, whitish Tremella.

Habitat in arborum ramis dejectis semiputridis, in sylvis densis. ☉. IX—V.

Desc. Frons gelatinosa, difformis, albida subinde glaucescens et flavescens, subrugosa, solida diaphana.

granulata.

13. TREMELLA gelatinosa sphærica aggregata viridis.

Tremella pellustris, vesiculis sphæricis fungiformibus. Dill. musc. 51. t. 10. f. 17.

Ulva sphærica aggregata. Sp. pl. 1633.

Lichenoides fungiforme, capitulis vel vesiculis sphæricis, aqueo humore repletis. R. syn. 70.

Anglis, granulated Tremella.

Habitat ad margines fossarum et fluvium et in sylvis arenosis humidis, inter Newington et Hackney, prope Londinum, Dill. circa Charlton in Cantia copiose.

☉. X—V.

U L V A. 1206.

Fruclificationes in membrana diaphana.

pavonia. 1. ULVA plana reniformis decussatim striata sessilis Syst. nat. 719.

Fucus fronde sessili reniformi decussatim striata. Sp. pl. 1630. Gmel. fuc. 169.

Fucus maritimus gallopavonis pennas referens. B. pin. 155. Hist. Ox. III. 645. t. 8. f. 7. R. syn. 43. Ell. cor. 88. t. 33. f. c.

Corallina foliacea, frondibus proliferus reniformibus striatis transversimque fasciatis. Pall. Zooph. 419.

Anglis, striated Laver, or Turkey feather.

Habitat in rupibus et saxis submarinis passim. I—XII.

Obs. margines foliæ et fossarum ciliatæ, pilis tenuissimis.

Lactuca. 2. ULVA palmata prolifera membranacea: ramentis inferne angustatis. Sp. pl. 1632. Gmel. fuc. 215.

Ulva marina lactucæ similis. R. syn. 62.

Tremella marina vulgaris lactucæ similis. Dill. musc. 42. t. 8. f. 1.

Fucus marinus, lactuca marina dictus. Park: 1293.

Muscus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Habitat in rupibus, saxis et muris, in rupibus meridiem spectantibus, subter arcem Keven-lees-castle dictam, in comitatu Radnoriensi. Littl. Brown; prope Halifax, D. Bolton; in montibus Scotiis passim. 4. I—XII.

- deustus.* 81. LICHEN umbilicatus undique lævis. *Sp. pl.* 1618. Lichen fronde rotunde lobata inferne nigerrima glebosa. *Hall. hist.* 1998. Lichen pulmonarius saxatilis e cinereo fuscus minimus. *Vaill. paris.* 116. t. 21. f. 14. Lichenoides coriaceum cinereum, peltis atris compressis. *Dill. musc.* 218. t. 29. f. 117. Lichenoides saxatile, foliis minus divisis cinereo-fuscis. *R. syn.* 73. *Anglis,* sooty Lichen. Habitat in rupibus et saxis; in rupe Sancti Vincentii prope Bristolium. D. Dare. circa Llanberris. D. Davies. 4. I—XII.

- polyrrhizos.* 82. LICHEN umbilicatus polyphyllus utrinque lævis polyrrhizos. *Sp. pl.* 1618. *Fl. dan.* 471. f. 3? Lichenoides rugosum durum pullum, peltis atris verrucosis. *Dill. musc.* 220. t. 30. f. 118. *Anglis,* finged Lichen. Habitat in rupibus et saxis, in rupe Sancti Vincentii prope Bristolium; circa Llanberris. D. Davies, et in montibus Scotiis. 4. I—XII.

- vellens.* 83. Lichen umbilicatus subtus hirsutissimus. *Sp. pl.* 1617. Lichen fronde coriacea inferne pustulata villosa, scutellis atris. *Hall. hist.* 1997. Lichenoides pullum superne glabrum, inferne nigrum et cirrosum. *Dill. musc.* 226. t. 30. f. 130. β Lichenoides coriaceum, latissimo folio umbilicato et verrucoso. *Dill. musc.* 545. t. 82. f. 5. *Anglis,* fleecy Lichen. Habitat in rupibus et saxis circa Llanberris, Dill. in Clark's Park et Paradise prope Money-musk, Aberdeenshire. 4. I—XII.

84. LICHEN

84. LICHEN umbilicatus peltis turbinatis truncatis *probofci-*
perforatis. *Sp. pl.* 1617. *deus.*

Lichen foliaceus ciliatus, peltis cylindricis rectis spar-
sis perforatis truncatis circulo duplici. *Amœn.*
acad. II. 264.

Lichenoides corneum, marginibus eleganter firmbria-
tis. *Dill. musc.* 218. t. 29. f. 116.

Anglis, button Lichen.

Habitat in rupibus et saxis circa Llanberris in Arvoniam,
S. Brewer; et in monte Berwyn in comitatu Der-
biensi. *Littl. Brown*; in montibus Scotiis passim.
ꝛ. I—XII.

85. LICHEN umbilicatus polyphyllis undique lævis *pollyphyl-*
crenatus. *Sp. pl.* 1618. *lus.*

Lichenoides tenue pullum, foliis utrinque glabris.

Dill. musc. 225. t. 30. f. 129.

Anglis, smooth Lichen.

Habitat in rupibus et saxis passim. ꝛ. I—XII.

***** Scyphiferi.

86. LICHEN stipite erecto cylindrico cyphifero, tu- *pyxida-*
berculis marginalibus subrotundis. *tus.*

Lichen scyphifer simplex crenulatus, tuberculis fuscis. *Sp. pl.* 1619. *Weis. crypt.* 85.

Lichen scypho infundibuliformi simplici. *Hall. hist.*
1912.

Lichen pyxidatus major. *Vaill. paris.* 115. t. 21. f.
8. *Mich. gen.* 82. t. 41. f. 1.

Coralloides scyphiforme, tuberculis fuscis. *Dill. musc.*
79. t. 14. f. 6.

Lichenoides tubulosum pyxidatum cinereum. *Dill.*
gifs. 204. *R. syn.* 68.

Muscus pyxidatus. *Ger. em.* 1560. *Park.* 1308.

Muscus pyxioides terrestris. *B. pin.* 361.

β. Lichen caule simplici, calyce turbinato, centro sim-
pliciter prolifero. *Fl. lapp.* 431.

Lichen infundibulis glabris proliferis. *Hall. hist.* 1923.

Lichen pyxidatus prolifer. *Vaill. paris.* 115. t. 21.
f. 5.

- Lichenoides tubulosum pyxidatum proliferum. *Dill. giss.* 204. *musc.* 80. t. 14. f. 6. D—H. *R. syn.* 60.
- γ Lichen scyphiger infundibulorum fungis fuscis. *Hall. hist.* 1919.
- Lichen pyxidatus major, acetabulo fimbriato et tuberculoso. *Vaill. par.* 115. t. 21. f. 11.
- Lichenoides tubulosum pyxidatum, tuberculis fusconigris. *Dill. giss.* 204. *musc.* 81. t. 14. f. 6. c. proliferum. *Dill. giss.* 204. *musc.* 81. t. 14. f. 6. K—M. *R. syn.* 69.
- fimbriatus. δ Lichen scyphifer simplex denticulatus, stipite cylindrico. *Sp. pl.* 1619.
- Lichen scyphiger infundibulis ferratis. *Hall. hist.* 1915. coronatis. *Hall. hist.* 1916.
- Lichen pyxidatus minor. *Vaill. paris.* 115. t. 21. f. 6. *Mich. gen.* 82. t. 41. f. 4, 5.
- Lichenoides tubulosum pyxidatum, marginibus serratis. *Dill. giss.* 204. *R. syn.* 69.
- Coralloides scyphiforme gracile, marginibus ferratis. *Dill. musc.* 84. t. 14. f. 8.
- Lichen infundibulis ferratis proliferis. *Hall. hist.* 1924.
- Lichen infundibulis proliferis ferratis foliosis, et verrucosis. *Hall. hist.* 1925.
- Lichenoides pyxidatum proliferum, marginibus serratis. *R. syn.* 69.
- filiformis. ζ Lichen scyphifer simplex integerrimis filiformis, tuberculis fuscis. *Fl. angl.* 456.
- Lichen corniculo simplici apice depresso scutellato. *Hall. hist.* 1908.
- Lichenoides tubulosum pyxidatum exiguum, fuscovirens. *Dill. giss.* 206. *Ry. syn.* 70.
- Coralloides scyphis gracilibus tubiformibus, pedicularis folio. *Dill. musc.* 85. t. 14. f. 10.
- exiguus. η Lichen stipite brevissimo scyphifero.
- Coralloides scyphis humilibus intus fuscis. *Dill. musc.* 86. t. 14. f. 11.
- foliaceus. θ Lichen foliis suberectis laciniatis crispis scyphiferis, scyphis brevissimis conicis. *Fl. angl.* 457.
- Lichen infundibulis explanatis fungiferis. *Hall. hist.* 1934.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- Lichen cornubus acutis farinosis subramosis. *Hall. hist.* 1903.
- Coralloides vix ramosum, scyphis obscuris. *Dill. musc.* 90. t. 15. f. 14.
- Lichenoides tubulosum cinereum, non ramosum. *R. syn.* 68
- deformis. μ Lichen scyphifer simpliciusculus subventricosus, calycibus dentatis. *Sp. pl.* 1620. *Fl. lapp.* t. 11. f. 5.
- Lichen pyxidatus difformis, acetabulis cornigeris. *Hall. hist.* 1938.
- Lichen pyxidatus teres, acetabulis minoribus repandis. *Mich. gen.* 80. t. 41. f. 1.
- Lichenoides tubulosum minus ramosum cauliculis crassioribus. *R. syn.* 68.
- Lichenoides tubulosum magis ramosum, maxime difforme. *R. syn.* 68.
- Coralloides crassius subincanum, calycibus dentatis. *Dill. hist.* 95. t. 15. f. 18.
- ventrico-
sus. Lichen scyphifer ramosissimus ventricosus, scyphis dentatis, tuberculis fuscis. *Fl. angl.* 457.
- Lichen infundibulis proliferis palmatis serratis et punctatis. *Hall. hist.* 1937.
- Lichenoides tubulosum cinereum, valde crustaceum, ramulis brevioribus et crebrioribus ex acetabulis enascentibus. *Dill. giss.* 203. *R. syn.* 68.
- Coralloides cornucopioides incanum, scyphis cristatis. *Dill. musc.* 94. t. 15. f. 17.
- Muscus cornucopioides cristatus alpinus saxatilis. *Bocc. musc.* 11. 21. t. 8.
- digitatus. ξ Lichen scyphifer ramosissimus, ramis cylindricis, calycibus integris nodosis. *Sp. pl.* 1620. *Weis. crypt.* 91.
- Lichen infundibulis palmatis, fungulis coccineis. *Hall. hist.* 1939.
- Lichenoides coralliforme, apicibus coccineis. *R. syn.* 68.
- Coralloides ramulosum, tuberculis coccineis. *Dill. musc.* 96. t. 15. f. 19.
- Coralloides parum ramosum, tuberculis fuscis. *Dill. musc.* 97. t. 13. f. 20.

- Lichen verrucosus bicornis et tricornis, fungulis fuscis terminatis. *Hall. hist.* 1941.
- ☞ Lichen scyphifer ramosus, denticulatus filiformis. *gracilis.*
Sp. pl. 1619. *Weis. crypt.* 93.
- Lichen infundibulis proliferis et corniculatis. *Hall, hist.* 1940.
- Lichen pyxidatus et corniculatus ramosus alpinus e fusco-cinereus, pyxidulis crenatis. *Mich. gen.* 81. t. 41. f. 5.
- Lichenoides pyxidatum cinereum elatius, ramulis pyxidatim desinentibus. *R. syn.* 69.
- Coralloides scyphiforme ferratum elatius, caulibus gracilibus glabris. *Dill. musc.* 88. t. 14. f. 13.
- Musco-fungus pyxidatus gracilior ramosus, calycibus ferratis. *Hist. Ox. III.* 632. f. 15. t. 7. f. 6.
- ☞ Lichen corniculis erectis ramosis, infundibulis gracilibus. *Hall. hist.* 1945.
- Lichen pyxidatus et corniculatus cinereus angustior et elatior, corniculis longioribus. *Mich. gen.* 80. t. 41. f. 3, 4.
- Lichenoides tubulosum cinereum pulverulentum et crustaceum, ramulis ex acetabulis enascentibus minus copiosis. *Dill. giss.* 203. *R. syn.* 68.
- Coralloides scyphiforme cornutum. *Dill. musc.* 92. t. 15. f. 16.
- Anglis,* cup Lichen.
- Habitat in ericetis, pascuis montosis et sylvis. 4. I—
XII.

***** Fruticulosi.

87. LICHEN fruticulosus perforatus, ramulis brevissimis autis. *Sp. pl.* 1621. *uncialis.*
- Lichen fruticosus levis perforatus cauliculis palmatis brevissimis. *Hall. hist.* 1960.
- Lichenoides tubulosum, cauliculis mollioribus et crassioribus, minus. *R. syn.* 67.
- Coralloides perforatum minus molle et tenue. *Dill. musc.* 99. t. 16. f. 22.
- ☞ Lichen fruticosus levis perforatus spinosus, ramulis punctiferis. *Hall. hist.* 98. t. 16. f. 21.

Lichen •

- Lichen cornubus acutis farinosis subramosis. *Hall. hist.* 1903.
- Coralloides vix ramosum, scyphis obscuris. *Dill. musc.* 90. t. 15. f. 14.
- Lichenoides tubulosum cinereum, non ramosum. *R. syn.* 68
- deformis. μ Lichen scyphifer simpliciusculus subventricosus, calycibus dentatis. *Sp. pl.* 1620. *Fl. lapp.* t. 11. f. 5.
- Lichen pyxidatus difformis, acetabulis cornigeris. *Hall. hist.* 1938.
- Lichen pyxidatus teres, acetabulis minoribus repandis. *Mich. gen.* 80. t. 41. f. 1.
- Lichenoides tubulosum minus ramosum cauliculis crassioribus. *R. syn.* 68.
- Lichenoides tubulosum magis ramosum, maxime difforme. *R. syn.* 68.
- Coralloides crassius subincanum, calycibus dentatis. *Dill. hist.* 95. t. 15. f. 18.
- ventricosus. ν Lichen scyphifer ramosissimus ventricosus, scyphis dentatis, tuberculis fuscis. *Fl. angl.* 457.
- Lichen infundibulis proliferis palmatis serratis et punctatis. *Hall. hist.* 1937.
- Lichenoides tubulosum cinereum, valde crustaceum, ramulis brevioribus et crebrioribus ex acetabulis enascentibus. *Dill. giss.* 203. *R. syn.* 68.
- Coralloides cornucopioides incanum, scyphis cristatis. *Dill. musc.* 94. t. 15. f. 17.
- Muscus cornucopioides cristatus alpinus saxatilis. *Bocc. musc.* 11. 21. t. 8.
- digitatus. ξ Lichen scyphifer ramosissimus, ramis cylindricis, calycibus integris nodosis. *Sp. pl.* 1620. *Weis. crypt.* 91.
- Lichen infundibulis palmatis, fungulis coccineis. *Hall. hist.* 1939.
- Lichenoides coralliforme, apicibus coccineis. *R. syn.* 68.
- Coralloides ramulosum, tuberculis coccineis. *Dill. musc.* 96. t. 15. f. 19.
- Coralloides parum ramosum, tuberculis fuscis. *Dill. musc.* 97. t. 13. f. 20.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Lichen coralloides tubulosus albidus major et mollior, caulibus crassioribus minus ramosis, receptaculis florum perexiguis rufescentibus. *Mich. gen.* 79. t. 40. f. 2.

Lichenoides tubulosum, cauliculis mollioribus et crassioribus, majus. *R. syn.* 67.

Coralloides perforatum majus, molle et crassum. *Dill. musc.* 98. t. 16. 5. 21.

Musco-fungus cetanoides mollior et elatior albidus tubulosus. *Hist. Ox.* III. 633. f. 15. t. 7. f. 7.

Spinosus. γ Lichen fruticulosus tubulosus ramosissimus spinosus, ramulis tuberculatis subdigitatis. *Fl. angl.* 459.

Lichen fruticosus vage ramosus foliosus, extremis ramis palmatis fungiferis. *Hall. hist.* 1952.

Lichen coralloides tubulosus major, ex cinereo fusco-virens caulibus crassioribus et durioribus, minus ramosis, extremitatibus palmatis, receptaculis florum exiguis rufescentibus. *Mich. gen.* 79. t. 40. f. 3, 5.

Lichenoides tubulosum ramosum rigidius, majus et crassius cinereo-fuscum. *R. syn.* 67.

Coralloides sparsum, caulibus tortuosis et spinosis. *Dill. musc.* 101. t. 16. f. 25.

Anglis, perforated Lichen.

Habitat in ericetis et saxosis, montosis. 4. I—XII.

Obl. medius inter uncialem et subulatum est.

subulatus. 88. LICHEN fruticulosus subdichotomus, ramis simplicibus subulatis. *Sp. pl.* 1620. *Weis. crypt.* 97.

Lichen caule fruticoso erecto, ramis adscendentibus dichotomis. *Hall. hist.* 1955.

Lichenoides tubulosum cinereum minus crustaceum minusque ramosum. *R. syn.* 67.

Coralloides corniculis longioribus et rarioribus. *Dill. musc.* 102. f. 26.

Muscus corniculatus. *Tab. ic.* 800. *Ger. em.* 1560. *Park.* 1308.

furcatus. β Lichen fruticulosus tubulosus ramosus, ramis erectis furcatis. *Fl. angl.* 458.

Lichen caule fruticoso subfolioso ramis adscendentibus repetito bifurcatis punctiferis. *Hall. hist.* 1954.

Lichen

Lichen coralloides tubulosus medius e cinereo viridans ramulis erectis nonnihil foliosis, extremitatibus apertis et palmatis, receptaculis florum rufescentibus exiguis. *Mich. gen.* 80. t. 40. f. 4. et *D.*

Lichenoides tubulosum cinereum, ramosius et crustaceum. *R. syn.* 67.

Coralloides corniculis brevioribus et crebrioribus. *Dill. musc.* 104. t. 16. f. 27.

Musco-fungus montanus corniculatus. *Hist. Ox. III.* 632. f. 15. t. 7. f. 1. *Vaill. paris.* t. 26. f. 7.

Anglis, horned, β forked Lichen.

Habitat in ericetis et pascuis montosis. \mathcal{U} . I—XII.

89. LICHEN fruticulosus perforatus ramosissimus, ramangifemulis nutantibus. *Sp. pl.* 1620. *Weis. crypt.* 95. *rinus.* *Fl. dan.* 180.

Lichen fruticosus farinosus, furculis innumeris nutantibus. *Hall. hist.* 1957. *alopestris.*

Lichen coralloides tubulosus major ramosissimus receptaculis florum rufescentibus perexiguis. *Mich. gen.* 79. t. 40. f. 1.

Lichenoides tubulosum ramosissimum, fruticuli specie, candicans. *R. syn.* 66.

Coralloides montanum fruticuli specie, ubique candicans. *Dill. musc.* 107. t. 16. f. 29.

Musco-fungus coralloides montanus ramosissimus cinereus vulgaris. *Hist. Ox. III.* 633. f. 15. t. 7. f. 9.

8 Lichen caule ramoso, alis perforatis, ramis capitatis. *sylvaticus.* *Roy. lugdb.* 511.

Lichen caule fruticoso folioso, ramis subnutantibus, minute multifidis fungiferis. *Hall. hist.* 1956.

Lichenoides tubulosum ramosissimum, fruticuli specie, candicans, corniculis rufescentibus. *R. syn.* 67.

Coralloides fruticuli specie candicans, corniculis rufescentibus. *Dill. musc.* 110. t. 16. f. 30.

Muscus terrestris coralloides erectus, cornibus rufescentibus. *B. pin.* 361. *Park.* 1310.

Anglis, Rhendeer Lichen.

Habitat in ericetis, pascuis montosis et sylvis. \mathcal{U} . I—XII.

90. LICHEN

- pascalis.* 90. LICHEN fruticulosus solidus, tectus foliolis crustaceis. *Sp. pl.* 1621. *Weis. crypt.* 98. *Fl. dan.* 151. Lichen caule fruticoso, fungifero, verrucis calcariis tecto. *Hall. hist.* 1963.
Lichen alpinus glaucus ramosus botryoides. *Sch. it. alp. II.* 137. t. 19. f. 4. *Mich. gen.* 78. t. 35. f. 5—8.
Lichenoides non tubulosum cinereum ramosum, totum crustaceum. *R. syn.* 66.
Coralloides crispum et botryforme alpinum. *Dill. musc.* 114. t. 17. f. 33.
Musco-fungus coralloides fruticosior et lignosior. *Hist. Ox. III.* 635. f. 15. t. 7. f. 12.
Anglis, crisp Lichen.
Habitat in rupibus et saxosis montosis. 4. I—XII.

- fragilis.* 91. LICHEN fruticulosus solidus lævis, ramulis teretiusculis obtusis, tuberculis terminalibus globosis cevis.
Lichen fruticulosus solidus ramulis teretibus obtusis. *Sp. pl.* 1621. *Fl. lapp.* t. 11. f. 1.
Lichen caule fruticoso ramoso erecto densissime cespitoso. *Hall. hist.* 1962.
Lichenoides non tubulosum ramosissimum, fruticuli specie, cinereo-fuscum. *R. syn.* 65.
Coralloides alpinum corallinæ minoris facie. *Dill. musc.* 116. t. 17. f. 34.
globiferus. β Lichen fruticulosus solidus ramosissimus ramulis teretibus acutis furcatis, tuberculis terminalibus globosis. *Fl. angl.* 460. *Lin. Mantiss.* 133.
Lichen fruticosus coralloides non tubulosus cinereus ramosissimus, receptaculis florum sphericis concoloribus. *Mich. gen.* 103. t. 39. f. 6.
Lichenoides non tubulosum, ramulis scutellis nigris terminatis. *R. syn.* 66.
Coralloides cupressiforme, capitulis globosis. *Dill. musc.* 117. t. 17. f. 35.
Anglis, brittle, β globe Lichen.
Habitat in rupibus, saxis et ericetis montosis. 4. I—XII.

92. LICHEN

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- floridus.* 95. LICHEN filamentosus ramosissimus erectus, tuberculis farinaceis sparsis, scutellis radiatis.
Lichen filamentosus ramosus erectus, scutellis radiatis. *Sp. pl.* 1624. *Weis. crypt.* 102.
Lichen barbatus, ramis teretibus, furculis parallelis simplicibus, scutellis radiatis. *Hall. hist.* 1972.
Usnea vulgarissima tenuior et brevior cum orbiculis. *Dill. musc.* 69. t. 13. f. 13.
Muscus ramosus floridus. *Tab. ic.* 808. *Ger. em.* 1560.
Muscus arboreus cum orbiculis. *B. pin.* 361. *R. syn.* 65.
Musco-fungus arborum capillaceus, scutis amplis, per ambitum pilis radiatis. *Hist. Ox.* III. 635. f. 15. t. 7. f. 14.
- hirtus.* β Lichen filamentosus ramosissimus erectus, tuberculis farinaceis sparsis. *Sp. pl.* 1623. *Weis. crypt.* 101.
Lichen barbatus, teretibus cauliculis, ramis simplicibus parallelis. *Hall. hist.* 1973.
Usnea vulgarissima tenuior et brevior sine orbiculis. *Dill. musc.* 67. t. 13. f. 12.
Muscus ramosus. *Tab. ic.* 807. *Ger.* 1372. *R. syn.* 65.
Anglis, flowering Lichen.
Habitat in sylvis et sepibus frequens. 4. I—XII.
Obs. variat colore fulvo et luteo.
- picatus.* 96. LICHEN filamentosus pendulus, ramis implexis, scutellis radiatis. *Sp. pl.* 1622. *Mat. med.* 495.
Lichen teres pendulus exasperatus ramis descendens, scutellis radiatis. *Hall. hist.* 1971.
Usnea vulgaris, loris longis implexis. *Dill. musc.* 56. t. 11. f. 1.
Muscus arboreus, Usnea officinarum. *B. pin.* 361. *R. syn.* 64.
Muscus quercinus. *Ger. em.* 1558.
Muscus arboreus vulgaris et quercinus. *Park.* 1311.
Anglis, tree Lichen, or Mols.
Habitat in ramis arborum; in sylvis densis passim. 4. I—XII.

97. LICHEN filamentosus pendulus subarticulatus, ramis patentibus. *Sp. pl.* 1622. *Weis. crypt.* 100.

Lichen barbatus, trunco articulato lævi. *Hall. hist.* 1970.

Usnea barbata, loris tenuibus fibrosis. *Dill. musc.* 63. t. 12. f. 6.

Muscus capillaceus longissimus. *B. pin.* 361.

Anglis, bearded Lichen.

Habitat in ramis arborum, in sylva dicta Forest of Dean in Glocestriensi comitatu, et in sylva prope Bishopscastle in Salopia. *Dill. musc.* 4. I—XII.

98. LICHEN filamentosus articulatus, ramis tenuissimis punctatis. *Sp. pl.* 1623. *Roy. lugdb.* 513. *art.*

Usnea capillacea et nodosa. *Dill. musc.* 60. t. 11. f. 4.

Muscus arboreus nodosus. *B. pin.* 361. *R. syn.* 65. f. geniculatus. *Park.* 1311.

Anglis, jointed Lichen.

Habitat in ramis arborum in sylvis passim; in sylva ad villam Stokeſ-church; prope Burnley in Lancastria. *Dill. musc.* 4. I—XII.

99. LICHEN filamentosus ramosus subpendulus implicato-flexuosus, axillis compressis.

Lichen filamentosus pendulus, axillis compressis. *Sp. pl.* 1622. *Weis. crypt.* 101.

Lichen barbatus pendulus, filamentis nigris prolixis. *Hall. hist.* 1968.

Usnea jubata nigricans. *Dill. musc.* 67. t. 12. f. 7.

Muscus corallinus saxatilis fœniculaceus. *R. syn.* 65.

Musco-fungus trichoides e nigridine cinereus jubæ ad instar e saxis et arboribus dependens. *Hist. Ox.* III. 633.

β Lichen filamentosus ramosus divaricatus decumbens chalybeiformis. *Sp. pl.* 1623. *Fl. dan.* 261. *formis.*

Lichen fruticosus diffusus, caulibus vagis flexuosis rigidis teretibus. *Hall. hist.* 1969.

Usnea rigida horsum versus extensa. *Dill. hist.* 66. t. 13. f. 10.

Muscus caule rigido instar fili chalybei. *R. syn.* 65.

Anglis, wiry Lichen.

Habitat in rupibus et arborum truncis; in Wallia, et in boreali parte Britanniaë passim. 4. I—XII.

scaber. . 100. LICHEN filamentosus ramosissimus decumbens implexus scaber, scutellis concavis integerrimis. Usnea cæspitosa exilis capillacea atra. *Dill. musc.* 66. t. 13. f. 9.

Anglis, scabrous Lichen.

Habitat in rupibus et saxis grandioribus circa Llanberris copiose. in boreali parte Angliæ passim. 4. I—XII.

Obs. Scutellæ magnitudo fere seminis *Papaveris albi*, hemisphæricæ, truncatæ, marginatæ, atra fundo atrofusco, margine integerrima.

lanatus. 101. LICHEN filamentosus ramosissimus decumbens implicatus, ramulis brevissimis divaricatis.

Lichen fruticosus minimus flexuosus, cauliculis nigris capillaceis. *Hall. hist.* 1967.

Uinea lanæ nigræ instar saxi adhærens. *Dill. hist.* 66. t. 13. f. 8.

Musculus coralloides lanæ nigræ instar saxi adhærens. *R. syn.* 65.

Anglis, woolly Lichen.

Habitat in rupibus et saxosis, in Cornubia D. Stephens, in rupibus Cambriæ circa Borth, miliaris distantia a Bangor S. Brewer; in septentrionali parte Britanniaë passim. 4. I—XII.

pubescens. 102. LICHEN filamentosus ramosissimus decumbens implexus nitidus. *Sp. pl.* 1623.

Anglis, pubescent Lichen.

Habitat in rupibus et saxosis in Westmorlandia passim. 4. I—XII.

Obs. affinis præcedenti sed minor.

TREMELLA. 1204.

Frustrificationes vix manifestæ, in corpore gelatinoso.

juniperi- 1. TREMELLA sessilis membranacea auriformis fulva. *Sp. pl.* 1625.

na.

Agaricus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Lichen gelatinosus niger undosus et crispus. *Hall. hist.* 2038.

Lichen arboreus membranaceus nigricans. *Buxb. comm. petrop. III. 273. t. 14. f. 3.*

Anglis, tree Tremella, or Witches Butter.

Habitat in truncis arborum. ☉. IX—V.

Nostoc. 5. TREMELLA sessilis subrotunda plicata undulata fusco-virens.

Tremella sessilis plicata undulata. Sp. pl. 1625. Weis. crypt. 29.

Tremella terrestris sinuosa pinguis et fugax. Dill. musc. 52. t. 10. f. 14.

Linkia terrestris gelatinosa membranacea vulgatissima. Mich. gen. 126. t. 67. f. 1.

Lichen gelatinosus plicatus undulatus, laciniis crispis granulosis. Hall. hist. 2041.

Ulva terrestris pinguis et fugax. R. syn. 64.

Nostoc paracelsi. Act. paris. 1708. p. 228.

Anglis, jelly Tremella.

Habitat in pratis et pascuis post pluvias. ☉. I—XII.

utricula- 6. TREMELLA sessilis tuberculosa gelatinosa, tuberculis cavis.

Tremella fluviatilis gelatinosa et uterculosa. Dill. musc. 14. t. 10. f. 16?

Linkia palustris gelatinosa saxis adnascens ex obscuro fulvo et concava vesicam referens. Mich. gen. 126. t. 67. f. 2.

Anglis, bladder Tremella.

Habitat in rivulis supra lapides, circa Pentir et Llanberris, Dill. in rivulis montosis in comitatibus Westmorlandico et Cumberlandica. Prope Tidewell in comitatu Derbiensi. 4. I—XII.

Desc. Rupes et lapides sub aqua late tegit, hæc Tremella, substantia gelatinosa, viridia, rigidiuscula, fragilis, nunc. semi, nunc ad duas uncias crassa; superficies nitidiuscula, tuberculosa, subinde lævis, tuberculis intus cavis, variæ magnitudinis Pisi vel nucis Avellanæ.

7. TREMELLA

7. TREMELLA tuberculosa solida rugosa. *Sp. pl. verrucosa.*
1625.

Tremella subrotunda sinuosa difformis gelatinosa. *Lin.*

It. w-goth. 193.

Anglis, warty Tremella.

Habitat in rivulis et fontibus limpidis supra lapides. 4.

I—XII.

Obs. Substantia valde tenerrima lacte viridis.

8. TREMELLA hemisphærica sparsa. *Sp. pl.* 1626. *hemis-*
Anglis, hemisphæric Tremella. *sphærica.*

Habitat in confervis, Fucis et rupibus submarinis. 4.

I—XII.

9. TREMELLA subrotunda sinuosa difformis gelatino-*difformis.*
sa. *Sp. pl.* 1626. *It. w-goth.* 193.

Anglis, sea Tremella.

Habitat in confervis et Fucis et rupibus submarinis. 4.

I—XII.

10. TREMELLA rotunda imbricata livida. *Sp. pl. adnata.*
1627. *It. scan.* 323. *Fl. suec.* 1140.

Anglis, livid Tremella.

Habitat in rupibus et lapidibus submarinis. 4. I—XII.

11. TREMELLA subglobosa sessilis solitaria glabra. *purpurea.*
Sp. pl. 1626. *Fl. suec.* 1142.

Sphæria convexa miniata hirsuta. *Hall. hist.* 2189.

Lichenoides tuberculosum amæne purpureum. *Dill.*

giff. 206. *musc.* 127. t. 18. f. 6. *R. syn.* 71.

Lycogala globosum rubrum, grani panici magnitudine
et forma. *Mich. gen.* 216. t. 95. f. 3.

Noctoc granulorum coccineum arboribus adnascens.

Vaill. paris. 144.

Anglis, purple Tremella.

Habitat in arborum ramis moribundis et emortuis. ☉.

IX—V.

12. TREMELLA sessilis gelatinosa difformis albida. *albica.*

Echinus gelatinosus crytallinus. *Hall. hist.* 2319?

Anglis, whitish Tremella.

Habitat in arborum ramis dejectis semiputridis, in sylvis densis. ☉. IX—V.

Desc. Frons gelatinosa, difformis, albida subinde glaucescens et flavescens, subrugosa, solida diaphana.

granulata.

13. TREMELLA gelatinosa sphærica aggregata viridis.

Tremella pellustris, vesiculis sphæricis fungiformibus.

Dill. musc. 51. t. 10. f. 17.

Ulva sphærica aggregata. Sp. pl. 1633.

Lichenoides fungiforme, capitulis vel vesiculis sphæricis, aqueo humore repletis. R. syn. 70.

Anglis, granulated Tremella.

Habitat ad margines fossarum et fluvium et in sylvis arenosis humidis, inter Newington et Hackney, prope Londinum, Dill. circa Charlton in Cantia copiose.

☉. X—V.

U L V A. 1206.

Frustrificationes in membrana diaphana.

pavonia. 1. ULVA plana reniformis decussatim striata sessilis
Syst. nat. 719.

Fucus fronde sessili reniformi decussatim striata. Sp. pl. 1630. Gmel. fuc. 169.

Fucus maritimus gallopavonis pennas referens. B. pin. 155. Hist. Ox. III. 645. t. 8. f. 7. R. syn. 43. Ell. cor. 88. t. 33. f. c.

Corallina foliacea, frondibus proliferus reniformibus striatis transversimque fasciatis. Pall. Zooph. 419.

Anglis, striated Laver, or Turkey feather.

Habitat in rupibus et saxis submarinis passim. I—XII.

Obs. margines foliæ et fossarum ciliatæ, pilis tenuissimis.

Lactuca. 2. ULVA palmata prolifera membranacea: ramentis inferne angustatis. Sp. pl. 1632. Gmel. fuc. 215.

Ulva marina lactucæ similis. R. syn. 62.

Tremella marina vulgaris lactucæ similis. Dill. musc. 42. t. 8. f. 1.

Fucus marinus, lactuca marina dictus. Park: 1293.

Muscus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Alga longissimo lato tenuique folio. *B. pin.* 364.
bist. III. 801.

Anglis, brown Laver.

Habitat in rupibus et saxis submarinis, prope Sheerness
in insula Shepey copiose. ☉. V—X.

Desc. Frons pedatis et tripedatis, a duabus et octo uncias
late, criformis, membranacea, tenuis, undulata, nitida,
fusca.

lanceolata. 6. ULVA ovato-lanceolata plana. *Syst. nat.* 719.

Tremella marina, porri folio. *Dill. musc.* 46. t. 5.

f. 5.

Anglis, spear Laver.

Habitat in scopulis marinis, in insula Mona. D. Brewer.

4. I—XII.

linza. 7. ULVA fronde oblonga bilata. *Sp. pl.* 1633.

Ulva marina fasciata. *R. syn.* 62.

Tremella mariana fasciata. *Dill. musc.* 46. t. 9. f. 6.

Fucus linza dictus, lactucæ marinæ similis. *B. bist.*
III. 801.

Muscus lactucæ marinæ similis. *B. pin.* 364.

Anglis, ribbin Laver.

Habitat in saxis et rupibus submarinis; in fossis prope
Sheerness. 4. I—XII.

dichotoma. 8. ULVA plana dichotoma viridis. *Fl. angl.* 476.

Fucus membranaceus dichotomus gramineus. *R. syn.*

45.

Anglis, dichotomous Laver.

Habitat in rupibus et saxis submarinis in insula Walney
et in Devonis, Cornubia et Suffexia. 4. I—XII.

intestinalis. 9. ULVA tubulosa simplex membranacea viridis.

Ulva tubulosa simplex. *Sp. pl.* 1632. *Weis. crypt.*

271.

Ulva marina tubulosa, intestinorum figuram referens.

R. syn. 62.

Tremella marina tubulosa intestinorum figuram. *Dill.*

musc. 42. t. 9. f. 7.

Fucus cavius. *B. pin.* 364. *bist.* III. 803.

Anglis,

Anglis, gut Laver.

Habitat in fossis et stagnis marinis. ☉. III—X.

10. ULVA tubulosa subramosa compressa viridis, ramis compressa alternis longissimis.

Ulva tubulosa ramosa compressa. *Sp. pl.* 1632.

Ulva marina tenuissima et compressa. *R. syn.* 63.

Tremella marina tenuissima et compressa. *Dill. musc.* 48. t. 9, 10. f. 8.

Alga tubulosa albida, navibus adhærens. *Pet. gaz. t.* 9. f. 6.

Anglis, compressed Laver.

Habitat in rupibus, saxis et fossis marinis. ☉. I—XII.

11. ULVA tubulosa ramosissima teretiusecula; ramis oppositis acutis. *purpurascens.*

Anglis, purplish Laver.

Habitat in rupibus et saxis submarinis; prope Christchurch in comitatu Hantonienti. ☉. III—X.

Desc. Frons semipedalis, crassitie fili emporetici, tubulosa, teretiusecula, purpurascens, diaphana, ramosissima; Rami oppositi, subdistichi, teretes, acuti.

12. ULVA tubulosa articulata ramosissima, ramis oppositis dichotomis. *articulata.*

Corallina lenta purpurea compressa. *R. syn.* 34.

Fucus purpureus humilis tenuiter divisus geniculatus.

Hist. Ox. III. 646. f. 15. t. 8. f. 14.

- β Tremella marina cæspitosa, segmentis tenuibus. *Dill. musc.* 40. t. 10. f. 9.

Anglis, jointed Laver.

Habitat in rupibus et saxis submarinis, in Cornubia, Devoniam, Dorsetiam, Sufflexiam, &c. frequens. ☉. III—XI.

13. ULVA gelatinosa tubulosa uniformis simplex. *fistulosa.*

Anglis, pipe Laver.

Habitat in saxis et fucis marinis. ☉. V—IX.

Desc. Radix repens, Frondes numerosæ, gelatinosæ, uniformes, fistulosæ, apice clausa, fuscescentes, triunciales,

ciales, crassitie fili emporetici. Semina numerosa, parva, rotunda, fusca.

diaphana. 14. ULVA gelatinosa pallida diaphana teretiuscula ramosa, ramis brevissimis acutiusculis.

Alcionum polymorphum gelatinosum. *Faun. suec.* 2228. *Syst. nat.* 1295.

Alcionum cinerescence-hyalinum ramosissimum, ramis teretibus subacutis. *Pall. Zooph.* 207.

Spongia ramosa altera anglica. *Park.* 1304.

Fucus spongiosus nodosus. *Johns. it. cant.* 3. f. 3. *Ger. em.* 1570. *R. syn.* 49. *Ellis. cor.* 87. t. 32. f. d. D.

Urtica marina nodosa. *Bocc. musc. fic.* 209. t. 5. f. 13. *Anglis*, diaphanous Laver.

Habitat in littore maris; prope Sheerness copiose. 4, I—XII.

Desc. Frons gelatinosa, solida, teretiuscula, pallida, diaphana, ramosa; ramis brevissimis teretibus acutiusculis. Semina numerosissima, minuta, fusca, variat longitudine a quatuor ad sedecim uncias, crassitudine a digito fere ad carpum.

flavescens. 15. ULVA gelatinosa subteretiuscula subramosa flavescens, ramis brevissimis obtusis.

Alga minor flavescens varie divisa. *Mart. cent.* 32.

Anglis, yellowish Laver.

Habitat in rupibus et saxis marinis; in insulis Anglesea et Walney. 4. I—XII.

Desc. Frons gelatinosa, solida, teres, et subinde compressa et dilatata, tum margine sinuato-dentata, subramosa, pallide flavescens; ramis brevissimis, obtusis. Semina numerosa, parva, rotunda, fusca, longitudine trium et quatuor unciarum, crassitudine fere digiti minoris. An varietas præcedentis?

filiformis. 16. ULVA gelatinosa filiformis ramosissima purpurascens, ramis sparsis distantibus longissimis.

Anglis, filiform Laver.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

incrassata. 21. ULVA gelatinosa plana sinuato-dentata viridis, margine incrassato.

Tremella viridissima corniculis palmatis. *Hall. bist.* 2125.

Tremella palustris gelatinosa, Damæ cornuum facie. *Dill. musc.* 51. t. 10. f. 10.

Conferva gelatinosa, damæ cornua repræsentans. *R. syn.* 60.

Fucus fontanus pinguis, corniculatus viridis. *Vaill. paris.* 56. t. 10. f. 3.

Anglis, thick Laver.

Habitat in stagnis et fossis, prope Chichester, D. Manningham, in insula Selsey in Suffexia copiose, inter Greenwich et Woolwich; prope Doncaster, circa Spalding, et alibi in comitatu Lincolnensi. ☉. III—X.

prunifor- 22. ULVA subglobosa solitaria. *Sp. pl.* 1633. *Fl. suec.* 1159. intus succulenta. *Syst. nat.* 1633.

Fucus subglobosus sessilis mollis. *Gledit. monogr.* *Anglis*, plumb Laver.

Habitat in fossis et stagnis, in lacubus alpinis in Westmorlandia. ☉. V—X.

Obs. magnitudo pruni spinosæ s. insititiæ.

pisiformis. 23. ULVA globosa solitaria spongiosa fuscescens, *Anglis*, pea Laver.

Habitat in fossis inter Greenwich et Woolwich. ☉. III—VIII.

Desc. Frons variæ magnitudinis a seminis rapi ad pisum, spherica, reticulato-contexta, pulposa, fusco virens.

F U C U S. 1205.

Masc. *Vesiculæ* villis intertextæ.

Fem. *Vesiculæ* adpersæ granis immerfis apice prominulis. *Sem.* Solitaria.

* *Foliis distinctis.*

natans. 1. FUCUS caule tereti ramosissimo, foliis lanceolatis ferratis,

- ferratis, fructificationibus globosis pedunculatis.
Sp. pl. 1628. *Gmel. fuc.* 92.
Fucus folliculaceus, ferrato folio. *B. pin.* 365. *Barr. ic.* 1122.
Lenticula marina ferratis foliis. *Lob. obs.* 256. *Dalech. hist.* 1397. *Ger. em.* 1615. *Park.* 1281.
Anglis, flote Fucus, or Gulf-weed.
 Habitat in littoribus maritimis; in insula Shepey D. Banks; ex Northumbria recepti; inter rejeclamenta maris prope Falmouth. 4. I—XII.
2. **FUCUS caule compresso ramoso, foliis ovalibus integerrimis.**
Fucus vermicularis. *Gmel. fuc.* 162. t. 18. f. 4.
Fucus polypodioides. *Gmel. fuc.* 186.
 Alga minor caulifera, foliis parvis oblongis ex viridifuscentibus. *Mart. cent.* 32.
Anglis, ovate Fucus.
 Habitat in rupibus et saxis submarinis, prope Scarborough in comitatu Eboracensi, et Christ-church in comitatu Hantonensi. 5. V—X.
3. **FUCUS caule tereti ramoso, foliis ovato-oblongis sanguine-undulatis integerrimis.** *Fl. angl.* 475. *Gmel. fuc. us.* 185. t. 24. f. 2. *Fl. dan.* 349.
Fucus frondibus membranaceis ovato-oblongis integerrimis petiolatis, caule tereti ramoso. *Syst. nat.* 718.
Fucus sive alga folio membranaceo purpureo, lapathi sanguinei figura et magnitudine. *Hist. Ox. III.* 645. f. 15. t. 8. f. 6. *R. syn.* 49.
Anglis, dock-leaved Fucus.
 Habitat in rupibus et saxis submarinis passim; prope Falmouth copiose. 4. I—XII.
Obs. Fructificationes subrotundæ, pedunculatæ, atro-rubentes, ad latera ramorum et nervorum; magnitudine seminis rapi.
4. **FUCUS caule tereti ramoso, foliis oblongis undulatis sinuatis dentatis.**
Fucus frondibus membranaceis oblongis undulatis sinuatis,

Habitat in rupibus et arborum truncis; in Wallia, et in boreali parte Britanniaë passim. 4. I—XII.

scaber. . 100. LICHEN filamentosus ramosissimus decumbens implexus scaber, scutellis concavis integerrimis. Usnea cæspitosa exilis capillacea atra. *Dill. musc.* 66. t. 13. f. 9.

Anglis, scabrous Lichen.

Habitat in rupibus et saxis grandioribus circa Llanberris copiose. in boreali parte Angliæ passim. 4. I—XII.

Obs. Scutellæ magnitudo fere seminis Papaveris albi, hemisphæricæ, concavæ, marginatæ, atra fundo atrofusco, margine integerrima.

lanatus. 101. LICHEN filamentosus ramosissimus decumbens implicatus, ramulis brevissimis divaricatis.

Lichen fruticosus minimus flexuosus, cauliculis nigris capillaceis. *Hall. hist.* 1967.

Uinea lanæ nigræ instar saxis adhærens. *Dill. hist.* 66. t. 13. f. 8.

Musculus coralloides lanæ nigræ instar saxis adhærens. *R. syn.* 65.

Anglis, woolly Lichen.

Habitat in rupibus et saxosis, in Cornubia D. Stephens, in rupibus Cambriæ circa Borth, miliaris distantia a Bangor S. Brewer; in septentrionali parte Britanniaë passim. 4. I—XII.

pubescens. 102. LICHEN filamentosus ramosissimus decumbens implexus nitidus. *Sp. pl.* 1623.

Anglis, pubescent Lichen.

Habitat in rupibus et saxosis in Westmorlandia passim. 4. I—XII.

Obs. affinis præcedenti sed minor.

TREMELLA. 1204.

Fruclificationes vix manifestæ, in corpore gelatinoso.

juniperi- 1. TREMELLA sessilis membranacea auriformis fulva. *Sp. pl.* 1625.

na.

Agaricus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Lichen gelatinosus niger undosus et crispus. *Hall. hist.* 2038.

Lichen arboreus membranaceus nigricans. *Buxb. comm. petrop. III. 273. t. 14. f. 3.*

Anglis, tree Tremella, or Witches Butter.

Habitat in truncis arborum. ☉. IX—V.

Nostoc. 5. TREMELLA sessilis subrotunda plicata undulata fusco-virens.

Tremella sessilis plicata undulata. Sp. pl. 1625. Weis. crypt. 29.

Tremella terrestris sinuosa pinguis et fugax. Dill. musc. 52. t. 10. f. 14.

Linkia terrestris gelatinosa membranacea vulgatissima. Mich. gen. 126. t. 67. f. 1.

Lichen gelatinosus plicatus undulatus, laciniis crispis granulosis. *Hall. hist. 2041.*

Ulva terrestris pinguis et fugax. R. syn. 64.

Nostoc paracelsi. Aët. paris. 1708. p. 228.

Anglis, jelly Tremella.

Habitat in pratis et pascuis post pluvias. ☉. I—XII.

utricula- 6. TREMELLA sessilis tuberculosa gelatinosa, tuberculis cavis.

Tremella fluviatilis gelatinosa et uterculosa. Dill. musc. 14. t. 10. f. 16?

Linkia palustris gelatinosa saxis adnascens ex obscuro fulvo et concava vesicam referens. Mich. gen. 126. t. 67. f. 2.

Anglis, bladder Tremella.

Habitat in rivulis supra lapides, circa Pentir et Ilanberris, *Dill. in rivulis montosis in comitatibus Westmorlandico et Cumberlandica. Prope Tidewell in comitatu Derbiensi. 4. I—XII.*

Desc. Rupes et lapides sub aqua late tegit, hæc Tremella, substantia gelatinosa, viridia, rigidiuscula, fragilis, nunc. semi, nunc ad duas uncias crassa; superficies nitidiuscula, tuberculosa, subinde lævis, tuberculis intus cavis, variæ magnitudinis Pisi vel nucis Avellanæ.

• 7. TREMELLA

7. TREMELLA tuberculosa solida rugosa. *Sp. pl. verrucosa.*
1625.

Tremella subrotunda sinuosa difformis gelatinosa. *Lin.*

It. w-goth. 193.

Anglis, warty Tremella.

Habitat in rivulis et fontibus limpidis supra lapides. 4.

I—XII.

Obs. Substantia valde tenerrima lacte viridis.

8. TREMELLA hemisphærica sparsa. *Sp. pl. 1626. hemis-*
Anglis, hemisphæric Tremella. phærica.

Habitat in confervis, Fucis et rupibus submarinis. 4.

I—XII.

9. TREMELLA subrotunda sinuosa difformis gelatino-*difformis.*
sa. Sp. pl. 1626. It. w-goth. 193.

Anglis, sea Tremella.

Habitat in confervis et Fucis et rupibus submarinis. 4.

I—XII.

10. TREMELLA rotunda imbricata livida. *Sp. pl. adnata.*
1627. *It. scan. 323. Fl. suec. 1140.*

Anglis, livid Tremella.

Habitat in rupibus et lapidibus submarinis. 4. I—XII.

11. TREMELLA subglobosa sessilis solitaria glabra. *purpurea.*
Sp. pl. 1626. Fl. suec. 1142.

Sphæria convexa miniata hirsuta. Hall. hist. 2189.

Lichenoides tuberculosum amæne purpureum. Dill.

giff. 206. musc. 127. t. 18. f. 6. R. syn. 71.

Lycogala globosum rubrum, grani panici magnitudine
et forma. Mich. gen. 216. t. 95. f. 3.

Noctoc granulorum coccineum arboribus adnascens.

Vaill. paris. 144.

Anglis, purple Tremella.

Habitat in arborum ramis moribundis et emortuis. 0.

IX—V.

12. TREMELLA sessilis gelatinosa difformis albida. *albica.*

Echinus gelatinosus crytallinus. Hall. hist. 2319?

Anglis, whitish Tremella.

Habitat in arborum ramis dejectis semiputridis, in sylvis densis. ☉. IX—V.

Desc. Frons gelatinosa, difformis, albida subinde glaucescens et flavescens, subrugosa, solida diaphana.

granu-
lata.

13. TREMELLA gelatinosa sphærica aggregata viridis.

Tremella pellustris, vesiculis sphæricis fungiformibus.

Dill. musc. 51. t. 10. f. 17.

Ulva sphærica aggregata. Sp. pl. 1633.

Lichenoides fungiforme, capitulis vel vesiculis sphæricis, aqueo humore repletis. R. syn. 70.

Anglis, granulated Tremella.

Habitat ad margines fossarum et fluvium et in sylvis arenosis humidis, inter Newington et Hackney, prope Londinum, Dill. circa Charlton in Cantio copiose.

☉. X—V.

U L V A. 1206.

Frustrificationes in membrana diaphana.

pavonia. 1. ULVA plana reniformis decussatim striata sessilis
Syst. nat. 719.

Fucus fronde sessili reniformi decussatim striata. Sp. pl. 1630. Gmel. fuc. 169.

Fucus maritimus gallopavonis pennas referens. B. pin. 155. Hist. Ox. III. 645. t. 8. f. 7. R. syn. 43. Ell. cor. 88. t. 33. f. c.

Corallina foliacea, frondibus proliferus reniformibus striatis transversimque fasciatis. Pall. Zooph. 419.

Anglis, striated Laver, or Turkey feather.

Habitat in rupibus et saxis submarinis passim. I—XII.

Obs. margines foliæ et fossarum ciliatæ, pilis tenuissimis.

Lactuca. 2. ULVA palmata prolifera membranacea: ramentis inferne angustatis. Sp. pl. 1632. Gmel. fuc. 215.

Ulva marina lactucæ similis. R. syn. 62.

Tremella marina vulgaris lactucæ similis. Dill. musc. 42. t. 8. f. 1.

Fucus marinus, lactuca marina dictus. Park: 1293.

Muscus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Alga longissimo lato tenuique folio. *B. pin.* 364.
hist. III. 801.

Anglis, brown Laver.

Habitat in rupibus et saxis submarinis, prope Sheerness
in insula Shepey copiose. ☉. V—X.

Desc. Frons pedalis et tripedalis, a duabus et octo uncias
late, ensiformis, membranacea, tenuis, undulata, nitida,
fusca.

lanceolata. 6. ULVA ovato-lanceolata plana. *Syst. nat.* 719.

Tremella marina, porri folio. *Dill. musc.* 46. t. 5.
f. 5.

Anglis, spear Laver.

Habitat in scopulis marinis, in insula Mona. *D. Brewer.*
4. I—XII.

linza. 7. ULVA fronde oblonga bullata. *Sp. pl.* 1633.

Ulva marina fasciata. *R. syn.* 62.

Tremella mariana fasciata. *Dill. musc.* 46. t. 9. f. 6.

Fucus linza dictus, lactucæ marinæ similis. *B. hist.*
III. 801.

Muscus lactucæ marinæ similis. *B. pin.* 364.

Anglis, ribbin Laver.

Habitat in saxis et rupibus submarinis; in fossis prope
Sheerness. 4. I—XII.

dichotoma. 8. ULVA plana dichotoma viridis. *Fl. angl.* 476.

Fucus membranaceus dichotomus gramineus. *R. syn.*
45.

Anglis, dichotomous Laver.

Habitat in rupibus et saxis submarinis in insula Walney
et in Devoniam, Cornubiam et Suffexiam. 4. I—XII.

intestinalis. 9. ULVA tubulosa simplex membranacea viridis.

Ulva tubulosa simplex. *Sp. pl.* 1632. *Weis. crypt.*
271.

Ulva marina tubulosa, intestinorum figuram referens.
R. syn. 62.

Tremella marina tubulosa intestinorum figuram. *Dill.*
musc. 42. t. 9. f. 7.

Fucus cavus. *B. pin.* 364. *hist.* III. 803.

Anglis,

Anglis, gut Laver.

Habitat in fossis et stagnis marinis. ☉. III—X.

10. ULVA tubulosa subramosa compressa viridis, ramis compressa alternis longissimis.

Ulva tubulosa ramosa compressa. *Sp. pl.* 1632.

Ulva marina tenuissima et compressa. *R. syn.* 63.

Tremella marina tenuissima et compressa. *Dill. musc.* 48. t. 9, 10. f. 8.

Alga tubulosa albida, navibus adhærens. *Pet. gaz. t.* 9. f. 6.

Anglis, compressed Laver.

Habitat in rupibus, saxis et fossis marinis. ☉. I—XII.

11. ULVA tubulosa ramosissima teretiusecula; ramis oppositis acutis. *purpurascens.*

Anglis, purplish Laver.

Habitat in rupibus et saxis submarinis; prope Christchurch in comitatu Hantonienfi. ☉. III—X.

Desc. Frons semipedalis, crassitie fili emporetici, tubulosa, teretiusecula, purpurascens, diaphana, ramosissima; Rami oppositi, subdistichi, teretes, acuti.

12. ULVA tubulosa articulata ramosissima, ramis oppositis dichotomis. *articulata.*

Corallina lenta purpurea compressa. *R. syn.* 34.

Fucus purpureus humilis tenuiter divisus geniculatus.

Hist. Ox. III. 646. f. 15. t. 8. f. 14.

- β Tremella marina cæspitosa, segmentis tenuibus. *Dill. musc.* 40. t. 10. f. 9.

Anglis, jointed Laver.

Habitat in rupibus et saxis submarinis, in Cornubia, Devoniam, Dorsetiam, Sufflexiam, &c. frequens. ☉. III—XI.

13. ULVA gelatinosa tubulosa uniformis simplex. *fistulosa.*

Anglis, pipe Laver.

Habitat in saxis et fucis marinis. ☉. V—IX.

Desc. Radix repens, Frondes numerosæ, gelatinosæ, uniformes, fistulosæ, apice clausa, fuscescentes, triunciales,

ciales, crassitie fili emporetici. Semina numerosa, parva, rotunda, fusca.

- diaphana.* 14. **ULVA** gelatinosa pallida diaphana teretiuscula ramosa, ramis brevissimis acutiusculis.
 Alcionum polymorphum gelatinosum. *Faun. suec.* 2228. *Syst. nat.* 1295.
 Alcionum cinerescence-hyalinum ramosissimum, ramis teretibus subacutis. *Pall. Zooph.* 207.
 Spongia ramosa altera anglica. *Park.* 1304.
 Fucus spongiosus nodosus. *Johns. it. cant.* 3. f. 3. *Ger. em.* 1570. *R. syn.* 49. *Ellis. cor.* 87. t. 32. f. d. D.
 Urtica marina nodosa. *Bocc. musc. fic.* 209. t. 5. f. 13. *Anglis*, diaphanous Laver.
 Habitat in littore maris; prope Sheerness copiose. 4; I—XII.
 Desc. Frons gelatinosa, solida, teretiuscula, pallida, diaphana, ramosa; ramis brevissimis teretibus acutiusculis. Semina numerosissima, minuta, fusca, variat longitudine a quatuor ad sedecim uncias, crassitudine a digito fere ad carpum.
- flavescens.* 15. **ULVA** gelatinosa subteretiuscula subramosa flavescens, ramis brevissimis obtusis.
 Alga minor flavescens varie divisa. *Mart. cent.* 32. *Anglis*, yellowish Laver.
 Habitat in rupibus et saxis marinis; in insulis Anglesea et Walney. 4. I—XII.
 Desc. Frons gelatinosa, solida, teres, et subinde compressa et dilatata, tum margine sinuato-dentata, subramosa, pallide flavescens; ramis brevissimis, obtusis. Semina numerosa, parva, rotunda, fusca, longitudine trium et quatuor unciarum, crassitudine fere digiti minoris. An varietas præcedentis?
- filiformis.* 16. **ULVA** gelatinosa filiformis ramosissima purpurascens, ramis sparsis distantibus longissimis.
Anglis, filiform Laver.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

incrassata. 21. ULVA gelatinosa plana sinuato-dentata viridis, margine incrassato.
 Tremella viridissima corniculis palmatis. *Hall. bist.* 2125.
 Tremella palustris gelatinosa, Damæ cornuum facie. *Dill. musc.* 51. t. 10. f. 10.
 Conferva gelatinosa, damæ cornua repræsentans. *R. syn.* 60.
 Fucus fontanus pinguis, corniculatus viridis. *Vaill. paris.* 56. t. 10. f. 3.
Anglis, thick Laver.
 Habitat in stagnis et fossis, prope Chichester, D. Manningham, in insula Selsey in Sufflexia copiose, inter Greenwich et Woolwich; prope Doncaster, circa Spalding, et alibi in comitatu Lincolnensi. ☉. III—X.

prunifor- 22. ULVA subglobosa solitaria. *Sp. pl.* 1633. *Fl. suec.* 1159. intus succulenta. *Syst. nat.* 1633.
mis. Fucus subglobosus sessilis mollis. *Gledit. monogr.*
Anglis, plumb Laver.
 Habitat in fossis et stagnis, in lacubus alpinis in Westmorlandia. ☉. V—X.
 Obs. magnitudo pruni spinosæ s. insititiæ.

pisiformis, 23. ULVA globosa solitaria spongiosa fuscescens, *Anglis*, pea Laver.
 Habitat in fossis inter Greenwich et Woolwich. ☉. III—VII.
 Desc. Frons variæ magnitudinis a seminis rapi ad pisum, spherica, reticulato-contexta, pulposa, fusco virens.

F U C U S. 1205.

Masc. *Vesiculæ* villis intertextæ.

Fem. *Vesiculæ* adpersæ granis immerfis apice prominulis. *Sem.* Solitaria.

* *Foliis distinctis.*

natans. 1. FUCUS caule tereti ramosissimo, foliis lanceolatis ferratis,

- ferratis, fructificationibus globosis pedunculatis.
Sp. pl. 1628. *Gmel. fuc.* 92.
Fucus folliculaceus, serrato folio. *B. pin.* 365. *Barr. ic.* 1122.
Lenticula marina ferratis foliis. *Lob. obs.* 256. *Dalech. hist.* 1397. *Ger. em.* 1615. *Park.* 1281.
Anglis, flote Fucus, or Gulf-weed.
Habitat in littoribus maritimis; in insula Shepey D. Banks; ex Northumbria recepi; inter rejectamenta maris prope Falmouth. 4. I—XII.
2. **FUCUS caule compresso ramoso, foliis ovalibus integerrimis.**
Fucus vermicularis. *Gmel. fuc.* 162. t. 18. f. 4.
Fucus polypodioides. *Gmel. fuc.* 186.
Alga minor caulifera, foliis parvis oblongis ex viridifuscentibus. *Mart. cent.* 32.
Anglis, ovate Fucus.
Habitat in rupibus et saxis submarinis, prope Scarborough in comitatu Eboracensi, et Christ-church in comitatu Hantonensi. ☉. V—X.
3. **FUCUS caule tereti ramoso, foliis ovato-oblongis sanguine-undulatis integerrimis.** *Fl. angl.* 475. *Gmel. fuc. us.* 185. t. 24. f. 2. *Fl. dan.* 349.
Fucus frondibus membranaceis ovato-oblongis integerrimis petiolatis, caule tereti ramoso. *Syst. nat.* 718.
Fucus sive alga folio membranaceo purpureo, lapathi sanguinei figura et magnitudine. *Hist. Ox. III.* 645. f. 15. t. 8. f. 6. *R. syn.* 49.
Anglis, dock-leaved Fucus.
Habitat in rupibus et saxis submarinis passim; prope Falmouth copiose. 4. I—XII.
Obs. Fructificationes subrotundæ, pedunculatæ, atro-rubentes, ad latera ramorum et nervorum; magnitudine seminis rapi.
4. **FUCUS caule tereti ramoso, foliis oblongis undulatis sinuatis dentatis.**
Fucus frondibus membranaceis oblongis undulatis sinuatis,

sinuatis, stipite tereti ramoſo. *Syst. nat.* 718. *Sp. pl.* 1630. *Fl. dan.* 1652.

Fucus crenatus. *Gmel. fuc.* 184. t. 24. f. 1.

Fucus membranaceus purpureus latifolius pinnatus.

R. fyn. 47.

Anglis, reddish Fucus.

Habitat in rupibus et saxis marinis. 4. I—XII.

fimbriatus.

5. **FUCUS** caule compresso subsimplici, foliis distichis linearibus ciliatis, fructificationibus marginalibus oblongis.

Fucus caulescens. *Gmel. fuc.* 173. t. 20. f. 2.

Anglis, fringed Fucus.

Habitat in littoribus marinis, in insula Portlandica sed rarius. 4. I—XII.

** *Ramosi fronde unita.*

filiquosus.

6. **FUCUS** fronde compressa ramoſa, foliis distichis alternis integerrimis, fructificationibus pedunculatis oblongis mucronatis. *Syst. nat.* 716. *Sp. pl.* 1629. *Fl. dan.* 106. *Gmel. fuc.* 81. t. 2. B.

Fucus maritimus alter tuberculis paucissimis. *B. pin.* 365.

Fucus angustifolius vesiculis oblongis siliquarum æmulis. *R. fyn.* 48. *Hist. Ox.* III. 647.

Quercus marina quarta. *Ger. em.* 1569.

Anglis, polded Fucus.

Habitat in rupibus et saxis marinis. 4. I—XII.

concatenatus.

7. **FUCUS** fronde filiformi ramoſissima, ramulis dichotomis vesiculis moniliformibus distantibus innatis, foliis subulatis. *Syst. nat.* 717?

Fucus caule tereti ramoſissimo, fructificationibus oblongis concatenatis. *Roy. lugd.* 514. *Sp. pl.* 1628.

Fucus folio tenuissimo diviso, siliquatus. *R. fyn.* 48.

Fucus foliis ericæ Raii. *Reaum. act. gall.* 1712. 31. t. 3. f. 5.

Fucus marinus foliis abrotani maris. *Lob. ic.* 2. p. 254.

Fucus foliculaceus foliis abrotani. *B. pin.* 365.

Muscus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Fucus abrotanoides vesiculis parvis rotundis ornatus.

Hist. Ox. III. 648. t. 8. f. 17.

Fucus radicibus arborum fibrosis similis. *R. syn. 49.*

Fucus folliculaceus : foeniculi folio. *B. pin. 365?*

Anglis, fibrous Fucus.

Habitat in littoribus Eboracensi, Lancastrensi et Han-
toniensi. 4. I—XII.

tamarisci- 12. FUCUS fronde filiformi ramosissima, ramis alter-
folius. nis vesiculis subrotundis innatis, foliis alternis ge-
minatis subulatis basi tuberculatis.

selaginoi- Fucus fronde filiformi ramosissima, ramis dichotomis,
des. foliis subulatis alternis basi vesiculosis. *Syst. nat.*
717. *Mant. 134.*

Fucus abies marina. *Gmel. fuc. 83. t. 11. A. f. 1.*

Fucus tamarisco similis maritima. *B. pin. 365.*

Fucus foliis ericæ aut tamarisci *R. syn. 49.*

Erica marina quibusdam. *B. hist. III. 799.*

Anglis, tamarisk Fucus.

Habitat in rupibus et saxis submarinis. Prope Mar-
ketjew copiose et alibi in Cornubia et Devoniam; in
littore Eboracensi. 4. I—XII.

*** *Alati f. Fronde plana, stipite medium folium percur-
rente.*

ferratus. 13. FUCUS fronde plana dichotoma ferrato-dentata,
fructificationibus terminalibus tuberculosis. *Syst.*
nat. 715. Sp. pl. 1626. Gmel. fuc. 57.

Fucus f. Alga latifolia major dentata. *R. syn. 42.*

Hist. Ox. III. 648. f. 15. t. 9. f. 1. Aët. gall. 1711.
285. t. 9. f. 1.

β Fucus marina humilis latifoliæ, ferratæ similis. *R.*
syn. 52.

Anglis, serrated Fucus, or sea Wrack.

Habitat in rupibus et saxis marinis. 4. I—XII.

vesiculo- 14. FUCUS fronde plana dichotoma integerrima, ve-
lus. siculis axillaribus geminis, terminalibus tubercula-
tis. *Syst. nat. 715. Sp. pl. 1626. Gmel. fuc. 60.*

Fucus

Fucus maritimus vel Quercus maritima vesiculas habens. *B. pin.* 363.

Fucus s. Alga marina latifolia vulgatissima. *R. syn.* 40.

Quercus marina. *Lob. ic. 2rp.* 255. *Ger. em.* 1567. herbacea et varietas. *Park.* 1293.

β Fucus maritimus s. Quercus maritima foliorum extremis tumidis. *B. pin.* 365.

Fucus palmaris latioribus foliis in binas ternasve vesiculas verrucosas terminatis. *Hist. Ox. III.* 647. *f.* 15. *t.* 8. *f.* 10.

γ Fucus fronde plana dichotoma integerrima axillis didivari-
varicatis, vesiculis axillaribus geminis. *Sp. pl.* 1627. *tus.*

Fucus bullatus fruticescens, caule nudo, foliis rectis compressis bifidis. *Hist. Ox. III.* 647. *f.* 15. *t.* 8. *f.* 5.

Anglis, bladder Fucus, or Sea Oak.

Habitat in rupibus et saxis marinis. 4. I—XII.

15. FUCUS fronde plana dichotoma integerrima punctata: inferne lineari canaliculata, fructificationibus tuberculatis geminis. *Syst. nat.* 715. *Sp. pl.* 1627. *Fl. dan.* 286.

Fucus spiralis maritimus major. *R. syn.* 497.

Alga spiralis maritima major. *Hist. Ox. III.* 646.

Anglis, spiral Fucus.

Habitat in saxis et rupibus, submarinis in littore Cantiano, Suffexiano, et Essexiano frequens. 4. I—XII.

16. FUCUS fronde plana spirali' perfoliata repando-volubilis, dentata. *Syst. nat.* 715. *Sp. pl.* 1627. *Gmel: fuc.* 180?

Fucus spiralis ferratus. *Bar. ic.* 1303?

Fucus spiralis maritima minor. *R. syn.* 42. *Tourn.* 568.

Alga spiralis maritima. *Bocc. sic.* 70. *t.* 38. *f.* 2?

Anglis, twisted Fucus.

Habitat in saxis et rupibus marinis in estuario Cestriensi. *R. syn.* 4. I—XII.

linearis. 17. FUCUS alatus dichotomus integerrimus acutus, fructificationibus tuberculatis terminalibus.

Fucus filiformis. *Gmel. fuc. 72. t. 1. f. 1.*

Fucus s. Quercus marina minima angustifolia. *Doody.*

R. syn. ed. 11. 328.

Anglis, linear Fucus.

Habitat in rupibus et saxis submarinis. 4. I—XII.

alatus. 18. FUCUS alatus membranaceus subdichotomus integerrimus coloratus, laciniis alternis bifidis.

Fucus frondibus membranaceis subdichotomis, nervosis, laciniis alternis decurrentibus bifidis. *Lin.*

Mant. 135. Fl. dan. 352. Gmel. fuc. 187. t. 25.

f. 1.

Fucus dichotomus parvus costatus et membranaceus.

R. syn. 44.

Fucus purpureus tenuiter divisus non geniculatus.

Hist. Ox. III. 646.

Anglis, winged Fucus.

Habitat in rupibus et saxis submarinis. 4. V—X.

esculentus. 19. FUCUS alatus simplicissimus ensiformis, stipite pinnata.

Fucus fronde simplici indivia ensiformi, stipite pinnata folium percurrent. *Mant. 135. Fl. dan. 200.*

t. 29. f. 1. Gunner. act Nidros. 4. t. 8. f. 1.

Fucus alatus sive phasganoides. *B. pin. 364. pr. 154.*

Fucus scoticus latissimus edulis dulcis. *Sibb. pr. 26.*

R. syn. 46.

Anglis, esculent Fucus.

Habitat in rupibus et saxis submarinis, in littore Cumberlandico, et Scotico passim. 4. I—XII.

*** Fronde plana avenia.

saccharinus. 20. FUCUS fronde plana avenia simplici ensiformi, stipite tereti brevissima.

Fucus fronde simplici ensiformi, stipite brevissima.

Syst. nat. 718. Sp. pl. 1630. Fl. dan. 416. Gmel. fuc. 194. t. 27 et 28.

Fucus folio singulari, longissimo lato, in medio rugoso,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Fucus ciliatus. *Gmel. fuc. t. 21. f. 1.*
 Alga minor suave rubens varie divisa. *Mart. cent.*
 32. t. 33.

Anglis, jagged Fucus.

Habitat in rupibus et saxis submarinis. ☉. IV—X.

Obs. variat marginibus integris et dentato-crispatis.

ciliatus. 25. **FUCUS** frondibus planis aveniis ramosis membranaceis acutis ciliatis.

Fucus frondibus membranaceis lanceolatis proliferis ciliatis. *Mant. 136. Syst. 718. Fl. dan. 353.*

Fucus holosetaceus. *Gmel. fuc. 177. t. 21. f. 2.*

Fucus ligulatus. *Gmel. fuc. 178. t. 21. f. 3.*

Fucus membranaceus rubens angustifolius, marginibus ligulis armatis. *R. syn. 47.*

Fucus humilis membranaceus acaulos elegantissimus ruber, capillis longis fimbriatis. *Hist. Ox. III. 646.*

Anglis, ciliated Fucus.

Habitat in rupibus et saxis marinis. ☉. IV—IX.

Obs. variat longitudine ciliarum a linea ad unciam. Frustrificationes ad apicem ciliarum, rotundæ, magnitudine seminis papaveris albi.

crispus. 26. **FUCUS** frondibus submembranaceis planis aveniis dichotomis, apicibus dilatatis bifidis.

Fucus frondibus submembranaceis dichotomis, laciniis dilatatis. *Syst. uat. 718. Mant. 134.*

Fucus membranaceus purpureus varie ramosus. *R. syn. 47.*

Alga crispa scabiosa rubra et pallida. *B. hist. III. 795.*

Anglis, curled Fucus.

Habitat in rupibus et saxis submarinis. ☉. I—XII.

crispatus. 27. **FUCUS** frondibus membranaceis planis aveniis linearibus ramosis, marginibus crispo-lacinulatis.

Fucus frondibus membranaceis sublinearibus ramosissimis crispis coloratis. *Syst. nat. 718?*

Fucus laceratus. *Gmel. fuc. 179. t. 21. f. 4?*

Ulva fronde ramosa plana undulata purpurea. *Fl. angl. 476.*

Anglis,

Anglis, crisped Fucus.

Habitat in rupibus et saxis submarinis, in comitatibus Cornubiensi, Devoniensi et Hantoniensi passim. ☉. V—X.

Desc. Frons palmaris membranaceus, tenerrimus, rubens, ramosus, linearis, planus, subundulatus, avenius, crispolacinulatus, obtusus, lacinulis fructiferis, fructificationes subrotundæ, parvæ, atro-rubescens.

28. FUCUS frondibus membranaceis planis aveniis bifidus. dilatatis bifidis.

Anglis, bifid Fucus.

Habitat in saxis et rupibus submarinis, in littore Hantoniensi. ☉. V—X.

Desc. Radix ramosus compressus repens. Frons uncialis et sesuncialis, membranacea, semel et subinde bis bifida, cuneiformia s. versus apicem dilatata, plana, avenia, purpurea, diaphana.

29. FUCUS frondibus cartilagineis planis aveniis rappinnatifidus. ramis dentato-pinnatifidis, dentibus callosis. dus.

Gmel. fuc. 156. t. 16. f. 3. Fl. dan. 354.

Fucus Osmunda. Gmel. fuc. 155. t. 16. f. 2.

Fucus Dealensis pedicularis rubriæ folio. R. syn. 48.

Pet mus. 405.

Alga cervi cornu divisura. B. hist. III. 797?

Anglis, pinnatifid Fucus.

Habitat in rupibus et saxis marinis. ☉. IV—X.

30. FUCUS frondibus cartilagineis planis aveniis ramulifidus. ramis alternis distichis subduplicato-pinnatifidis obtusis.

Fucus membranaceus acaulos angustior, foliis palmæ in modum divisis, marginibus laciniatis et veluti crispis. *Hist. Ox. III. 646. f. 15. t. 8. f. 2. Act. gal. 1712. t. 4. f. 6.*

Anglis, multifid Fucus.

Habitat in rupibus et saxis submarinis in comitatibus Devoniensi et Hantoniensi. ☉? IV—X.

Desc. Frons tri et sexuncialis cartilaginea, plana avenia, linearis, inferne angustior et crassior, fusco-rubra, sed

versus apicem pallidior et flavescens, ramosa; ramis alternis distichis subduplicato-pinnatifidis, laciniis obtusis.

dentatus. 31. FUCUS frondibus membranaceis aveniis alternatim pinnatifidis: sinibus obtusis, laciniis apice erosis. *Syst. nat.* 719. *Fl. dan.* 354.

Fucus corymbiferus. *Gmel. fuc.* 124. t. 9.

Fucus membranaceus rubens, foliolis latiusculis ad extrema dentatis. *Hist. Ox.* III. 646. f. 15. t. 8. f. 5.

Anglis, dentated Fucus.

Habitat in rupibus et saxis marinis, in comitatibus Devonienſi et Eboraceniſi. ☉. V—X.

herbaceus. 32. FUCUS frondibus subcartilagineis planis aveniis duplicato-pinnatis, laciniis suboppositis lineari-lanceolatis serratis.

Anglis, herbaceous Fucus.

Habitat in rupibus et saxis submarinis, inter reſectamenta maris prope Hastings in Suffexia, et in Northumbria. 4. I—XII.

Desc. Frons ſeſquipedalis et bipedalis circiter duas lineas lata, subcartilaginea, ovata, plana, avenia, herbacea, duplicato pinnata; laciniis oppositis lineari-lanceolatis, serratis, ferraturis ſubinde ſetaceis.

ceranoides. 33. FUCUS fronde cartilaginea plana avenia dichotoma integerrima punctata lanceolata, fructificationibus tuberculatis bifidis terminalibus. *Syst. nat.* 715.

Fucus fronde dichotoma plana integra, apicibus bifidis vesiculosis. *Sp. pl.* 1626. *Gmel. fuc.* 115. t. 7. f. 1, 2, 3.

Fucus membranaceus ceranoides varie dissectus. *R. ſyn.* 44.

Fucus dichotomus ceranoides angulis rotundiusculos efformans. *Hist. Ox.* III. 646. f. 15. t. 8. f. 11?

β Fucus membranaceus ceranoides ramosus, per ſiccitatem obsolete virens. *Pluk. alm.* 161. *R. ſyn.* 44.

δ Fucus ceranoides albidus, ramulorum apicibus stellatis. *Pluk. alm.* 161. *R. ſyn.* 44.

γ Fucus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Fucus longo angusto crassoque folio. *B. pin.* 364.
pr. 155. *R. syn.* 43.

Fucus marinus secundus. *Dod. pempt.* 479. *Park.*
1293.

Quercus marina secunda. *Ger. em.* 1568.

Anglis, thong Fucus, or sea Thongs vulgo.

Habitat in rupibus et saxis submarinis. 4. VI--IX.

tomento-
sus.
elonga-
tus,

36. FUCUS fronde compressa dichotoma tomentosa ob-
tusa.

Fucus fronde filiformi compressa dichotoma articu-
lata, geniculis tumidiusculis. *Syst. nat.* 716. *Sp.*
pl. 1627.

Fucus spongiosus teres ramosior viridis erectis.
Hist. Ox. III. 647. *f.* 15. *t.* 8. *f.* 7.

Spongia dichotomos teretifolia viridis. *R. syn.* 29.
Anglis, woolly Fucus.

Habitat in rupibus et saxis submarinis; in Cornubia.

R. syn. prope Exmouth in Devoniam, in littore Ebo-
racensi. 4. V—X.

nodosus.

37. FUCUS fronde compressa dichotoma, foliis di-
stichis integerrimis, vesiculis innatis solitariis di-
latatis. *Syst. nat.* 716. *Gmel. fuc.* 78. *t.* 1. *B. f.* 1.
Fl. dan. 146.

Fucus caule compresso dichotomo medio ramorum in
vesiculam dilatato. *Sp. pl.* 1628.

Fucus maritimus nodosus. *B. pin.* 365. *R. syn.* 48.
Aët. gall. 1712. 26. *t.* 2. *f.* 3.

Fucus marinus tertius. *Dod. pempt.* 480.

Fucus maritimus, vesiculis majoribus singularibus per
intervalla dispositis. *Hist. Ox.* III. 647. *t.* 8. *f.* 2.

Quercus marina tertia. *Ger. em.* 1568.

Anglis, knobbed Fucus.

Habitat in rupibus et saxis submarinis. 4. V—X.

pumilus.

38. FUCUS fronde compressa dichotoma, axillis diva-
ricatis, vesiculis subsolitariis axillis ultimis innatis
fructificationibus globosis terminalibus.

Anglis, dwarf Fucus.

Habitat

Habitat in rupibus et saxis inter fluxum et refluxum maris frequens. 4. VI—X.

Desc. Frons fere semiuncialis, cartilaginea, dichotoma, compressa, fere plana, inferne angustior, atro-olivacea et subinde purpuro-cærulea; Vesiculæ solitariae, et subinde geminae, rotundæ, inflatæ, infra ultimas axillas innatæ. Fructificationes globosæ, apicibus ramorum terminantes, parvæ, glabræ.

39. FUCUS fronde cartilaginea filiformis compressa dichotoma acuta.

Anglis, filiform Fucus.

Habitat in rupibus et saxis prope insula Walney in comitatu Lancastriensi. 4. V—X.

Desc. Frons semipedalis, cartilaginea, filiformis, compressa, dichotoma, diaphana, rubescens.

40. FUCUS fronde filiformi compressa ramosissima, aculeatus. dentibus marginalibus subulatis alternis erectis.

Syst. nat. 717. Fl. dan. 355.

Fucus muscoides. Gmel. fuc. 130. t. 12.

Fucus angustifolius, foliis dentatis. R. syn. 48.

Fucus tenuifolius, foliis dentatis. Hist. Ox. III. 648.

f. 15. t. 9. f. 4.

Anglis, prickly Fucus.

Habitat in rupibus et saxis marinis, in Devoniam, Cornubiam et Northumbriam frequens. 4. V—X.

41. FUCUS fronde cartilaginea compressa ramosissima, corneus, ramis alternis distichis pinnatis, laciniis oppositis setaceis, fructificationibus subrotundis pedunculatis.

Fucus sericeus. Gmel. fuc. 149. t. 15. f. 3?

Fucus flavicans teretifoliis ramulis ennatim enascentibus. R. syn. 50.

Fruticulus parvus pennatus flavicans, corneus et tenax. Buddl. hort. succ. II. 2.

Anglis, horney Fucus.

Habitat in rupibus submarinis in Cornubia et Devoniam passim. 4. V—X.

42. FUCUS

- pinnatus.* 42. FUCUS fronde cartilaginea filiformi compressa subtriplicato-pinnata, laciniis subulatis erectiusculis.
Anglis, pinnated Fucus.
 Habitat in rupibus et saxis submarinis in Cornubia, Devoniam et Suffexia passim; prope Scarborough in comitatu Eboracensi. ☉. V—X.
 Desc. Frons triuncialis, filiformis, compressa, cartilaginea, rubra, duplicato vel triplicato-pinnata acuta; laciniis oppositis subulatis, brevissimis, erectiusculis.
- obtusus.* 43. FUCUS fronde cartilaginea filiformi compressa subduplicato-pinnata, laciniis obovatis apice tuberculatis.
Anglis, obtuse Fucus.
 Habitat in saxis et rupibus marinis prope Hastings in Suffexia et in Devoniam passim. ☉. V—X.
 Desc. Frons quadipollicaris, filiformis, compressa, cartilaginea, diaphana, purpurascens, subduplicato-pinnata; laciniis oppositis, obovatis, patentibus, apice tuberculatis.
- filicinus.* 44. FUCUS fronde cartilaginea compressa obtusa subdecomposito-pinnata, laciniis horizontalibus obtusis.
Anglis, fern Fucus.
 Habitat in rupibus et saxis submarinis prope Walney in comitatu Lancastriensi. ☉. V—X.
- cartilagineus.* 45. FUCUS fronde cartilaginea compressa decomposito-pinnata, laciniis linearibus. *Syst. nat.* 719.
Fucus vesicolor. Gmel. fuc. 158. t. 17. f. 2.
Fucus coronopi facie. Buddl. hort. sic. 1. 12. R. syn. 45.
 Frutices marini arbuscularum forma muticolores.
Seb. mus. III. 191. t. 102. f. 1, 2.
Anglis, cape Fucus, or Weed.
 Habitat in rupibus et saxis; in Cornubia D. Stevens. 4. I—XII.
- coccineus.* 46. FUCUS fronde cartilaginea compressa ramosissima, ramulis alternatim secundis, fructificationibus globosis lateralibus. *Fl. angl.* 473.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

*verruco-
sus.*

50. **Fucus** fronde cartilaginea setacea ramosa, ramis longissimis, fructificationibus lateralibus globosis sessilibus.

Fucus longissimus. Gmel. fuc. 134. t. 13.

Fucus marinus purpurascens parvus, caule et ramulis s. foliolis teretibus. R. syn. 50.

Fucus teres ramosissimus, Reaum. act. gall. 1712. t. 5. f. 9.

Fucus teres rubens minus ramosus in longum protensus. R. syn. 51.

Anglis, warty Fucus.

Habitat in rupibus et saxis submarinis passim. 4. VI—X.

albidus.

51. **FUCUS** fronde cartilaginea filiformi teretiüscula subdichotoma; ramis setaceis distantibus subsecundis, fructificationibus lateralibus subrotundis sessilibus. *Fl. angl. 470. Gmel. fuc. 138.*

Fucus verrucosus. Gmel. fuc. 136. t. 14. f. 1.

Fucus teres albus tenuissime divilus. R. syn. 50.

Anglis, whitish Fucus.

Habitat in rupibus et saxis submarinis; in comitatibus Suffexiano, Hantonienſi et Corbunienſi. 4. VI—X.

Obs. variat albido et fuscente.

*tubercu-
latus.*

52. **FUCUS** fronde filiformi subdichotoma, ramis obtusis tuberculatis.

Fucus kali geniculato similis, non tamen geniculatus. R. syn. 43.

Anglis, tuberculated Fucus.

Habitat in rupibus et saxis submarinis prope St. Ives in Cornubia. 4. VI—X.

Obs. affinis F. fastigiato, sed differt non uniformiter dichotomus nec fastigiatus sed ramis inæqualibus, ultimis fructificationibus tuberculatis aspersis.

*fastigia-
tus.*

53. **FUCUS** fronde filiformi dichotoma fastigiata obtusa, vesiculis lanceolatis terminalibus.

Fucus fronde filiformi dichotoma ramosissima fastigiata obtusa. Syst. nat. 718. Sp. pl. 1631. Fl. dan. 393. Gmel. fuc. 106. t. 6. f. 1.

Fucus

Fucus palmaris tenuis in orbem expansis in segmenta bifida vel trifida breviora tertia divisus. *Hist. Ox. III. 649. f. 15. t. 9. f. 19.*

Fucus s. alga exigua dichotoma foliorum segmentis longiusculis, crassis et subrotundis. *R. syn. 45.*

Anglis, fastigiated Fucus.

Habitat in rupibus et saxis submarinis. 4. VI—X.

54. **FUCUS** fronde filiformi dichotoma fastigiata *acufurcellatus.*
minata. *Sp. pl. 1631.*

Fucus lumbricatis. *Fl. angl. 471. Gmel. fuc. 108. t. 6. f. 2.*

Fucus parvus segmentis prælongis teretibus acutis.

R. syn. 45. Hist. Ox. III. 648. f. 15. t. 9. f. 4.

Fucus forcellata, lumbricalis species. *B. pin. 366.*

Furcellaria. *Imp. hist. 842.*

Anglis, forked Fucus.

Habitat in rupibus et saxis submarinis. 4. I—XII.

55. **FUCUS** fronde cartilaginea filiformi dichotoma *diffusus.*
divaricata diffusa.

Anglis, diffuse Fucus.

Habitat in rupibus submarinis in Cornubia et Devoniam passim, et in insula Portlandica. 4. I—X.

Desc. Frons semipedalis fili emporetici crassitudine, cartilaginea, flavescens s. purpurascens, filiformis, diaphana, dichotoma valde divaricata, diffusa, apicibus acutis.

56. **FUCUS** fronde cartilaginea filiformi implicata *diplicatus.*
aphana. *Fl. angl. 470. Gmel. fuc. 142. t. 14. f. 2.*

Fucus trichoides nostras aurei coloris ramulorum apicibus furcatis. *Pluk. ph. 184. f. 2. R. syn. 45.*

Fucus coralloides erectus. *R. syn. 51.*

Alga exigua dichotoma arenacei coloris. *R. hist. 51.*

Anglis, matted Fucus.

Habitat in rupibus et saxis submarinis. 0. V—XI.

57. **FUCUS** fronde filiformi ramosissima, ramulis *purpurafertis* setaceis fructificationibus globosis innatis. *scens.*
Fl. angl. 471. Gmel. fuc. 139.

Fucus

Fucus teres rubens ramosissimus. *R. syn.* 51.

Anglis, purplish Fucus.

Habitat in rupibus et saxis submarinis. 4. V—X.

*incur-
vus.*

58. **FUCUS** fronde filiformi ramosissima, ramis imbricatis apicibus involutis, ramulis subulatis subsecundis.

Fucus pinnastroides. *Gmel. fuc.* 127. t. 11. f. 1.

Pinus maritima s. *Fucus teres*, cujus ramuli setis sursum tendentibus sunt obliti. *R. syn.* 50.

Anglis, pine Fucus.

Habitat in rupibus et saxis submarinis in Suffexia copiose. I—XII.

*amphi-
bious.*

59. **FUCUS** fronde filiformi ramosissima, ramis alternis involutis, ramulis brevissimis multifidis, fructificationibus oblongis pedunculatis. *Gmel. fuc.* 135.

Fucoides erectum fruticuli specie, summitatibus inflexis. *R. syn.* 38. t. 2. f. 6.

Anglis, amphibious Fucus.

Habitat in rupibus, saxis submarinis et ad radices plantarum in fossis et palustribus maritimis frequens. 4. VII—X.

muscoides. 60. **FUCUS** fronde filiformi ramosissima, ramis sparsis divaricatis flexuosis, spinis mollibus alternis.

Fucus fronde tereti ramosissima, ramis sparsis spinis mollibus alternis. *Sp. pl.* 1630.

Fucus folliculaceus fuscus s. *nigricans* ex insula cypro. *R. hist.* III. 13. *Pet. musc.* 219.

Anglis, thorny Fucus.

Habitat in rupibus submarinis, in littoribus Eboracensi et Northumbrienti sed rarius. 4. V—X.

lanosus.

61. **FUCUS** frondibus capillaceis dichotomis scabris fructificationibus tuberculatis lateralibus.

Fucus frondibus capillaceis dichotomis ramosissimis scabris. *Syst. nat.* 718.

Anglis, wool Fucus.

Habitat in rupibus et saxis submarinis; in insula Walney, in comitatu Lancastriensi. 4. VII—X.

62. **FUCUS**

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Byssus palustris confervoides non ramosa viridis, filamentis crassioribus, setas aprinas æmulantibus.

Mich. gen. 210. t. 89. f. 6.

Anglis, river Conferva, or Crow-silk.

Habitat in rivulis leniter fluentibus: β in fossis. 4.
I—XII.

fontinalis. 2. CONFERVA filamentis simplicissimis æqualibus digito brevioribus. *Sp. pl.* 1633.

Conferva cespitosa, filis rectis, undique divergentibus.

Hall. hist. 2114.

Conferva minima byssi facie. *Dill. musc.* 14. t. 12. f. 3.

Conferva fontalis fusca, omnium minima mollis. *R. syn.* 58.

Byssus minima palustris obscura, crassioribus filamentis plerumque simplicibus. *Mich. gen.* 211. t. 89. f. 8.

Anglis, Spring Conferva.

Habitat in saxis, in rivulis et fontibus; in rivo novo versus Hornsey. *Dill.* ☉. III—VI.

violacea. 3. CONFERVA filamentis simplicissimis æqualibus digito brevioribus violaceis.

Conferva mucosa confragosis rivulis innascens. *Dill. musc.* 15. t. 2. f. 4.

Anglis, violet Conferva.

Habitat in rivulis alpinis prope Llandberris. *Dill.* ☉. V—X.

• • • Filamentis ramosis æqualibus.

furcata. 4. CONFERVA filamentis æqualibus extremitatibus ramosis, ramis simplicibus. *Fl. angl.* 478.

Conferva capillacea, filamentis bifidis et trifidis. *Dill. musc.* 16. t. 12. f. 6.

Conferva rivulorum bicornis, filamentis tenuissimis. *R. syn.* 58.

β Conferva palustris, filamentis brevioribus et crassioribus. *Dill. musc.* 17. t. 3. f. 10.

Conferva palustris subhirsuta filamentis brevioribus
et crassioribus. *R. syn.* 477.

Anglis, forked Conferva.

Habitat in rivulis leniter fluentibus. β in fossis. \odot .
X—V.

5. CONFERVA filamentis æqualibus dichotomis. *Sp. dichotoma.*
pl. 1635. *Fl. dan.* 358.

Conferva dichotoma, setis porcinis similis. *Dill. musc.*
17. t. 3. f. 9. *Hall. hist.* 2117.

Conferva Plinii setis porcinis similis. *Merr. pin.*
R. syn. 58.

Anglis, dichotomous Conferva.

Habitat in fossis, inter Greenwich et Woolwich copiose,
prope Gravesend. \odot . III—X.

6. CONFERVA filamentis æqualibus ramosis implica-*bullosa.*
tis, bullas aëreas includentibus. *Sp. pl.* 1634.

Conferva pilis tenerrimis, vellere gossypiaceo. *Hall.*
hist. 2120.

Conferva palustris bombycina. *R. syn.* 60. *Dill. musc.*
18. t. 3. f. 11.

Alga bombycina. *B. pr.* 155. *pin.* 363. *Park.* 1261.

Musculus aquaticus bombycinus, tenuissimis filamen-
tis. *Loef. pruss.* 173. f. 55.

Anglis, cotton Conferva.

Habitat in fossis et stagnis passim. \odot . III—VI.

7. CONFERVA filamentis æqualibus versus basin ra-*canalicu-*
mosioribus. *Sp. pl.* 1634. ramis longis. *laris.*

Conferva filamentis tenuibus ramosis intricatis fuscis,
Hall. hist. 2118.

Conferva rivulorum capillacea densissime congestis ra-
mulis. *R. syn.* 58. *Dill. hist.* 21. t. 5. f. 15.

Alga in tubulis aquam fontanam deducuntibus. *B.*
pin. 364.

Anglis, mill Conferva.

Habitat in rivulis purioribus, inque tubulis et assaribus ad
molas deducuntibus. \mathcal{U} . I—XII.

- amphibia.* 8. **CONFERVA** filamentis æqualibus ramosis, exsiccatione coeuntibus in aculeos. *Sp. pl.* 479.
Conferva amphibia, fibrillosa et spongiosa. Dill. musc. 22. t. 5. f. 17.
Conferva terrestris exilis fibrillosa. R. syn. 59.
Anglis, amphibious Conferva.
Habitat ad ripas fluviorum, fossarum et paludium frequens. ꝥ. I—XII.
- rigida.* 9. **CONFERVA** filamentis æqualibus ramosissimis rigidiusculis, ramulis alternis brevissimis. *Fl. angl.* 479.
Conferva filis crassiusculis rigidioribus, subarticulatis. Hall. hist. 2122.
Conferva fluviatilis fibrillosa subrigida. Dill. musc. 21. t. 4. f. 16.
Anglis, rigid Conferva.
Habitat in fluvii; in ericeto Hounflejano, in fluvio Lug. prope Mortimer's Cross in comitatu Herefordiensi. Dill. ꝥ. I—X.
- foeniculacea.* 10. **CONFERVA** filamentis æqualibus ramosissimis, ramis ramulisque longissimis sparsis. *Fl. angl.* 479.
Conferva marina foeniculacea. Dill. musc. 16. t. 2. f. 8.
Anglis, fennel Conferva.
Habitat in rupibus et saxis submarinis in Mona insula. Dill. in Cornubia. ☉. VI—X.
- littoralis.* 11. **CONFERVA** filamentis æqualibus ramosissimis elongatis scabriusculis. *Sp. pl.* 1635.
Conferva maritima filiformis longissima fusca, ramis alternis simplicibus brevioribus pallidis. Fl. succ. 1170.
Conferva marina capillacea longa ramosissima. R. syn. 50. *Dill. musc.* 23. t. 6. f. 19.
Anglis, soft Conferva.
Habitat in rupibus et saxis submarinis passim. ☉. V—X.
- tomentosa.* 12. **CONFERVA** filamentis æqualibus tenuissimis ramosissimis, ramis simplicibus longis confertis fuscis. *Fl. angl.* 480.

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

cancellata. 17. CONFERVA filis ramosis, filamentis alternis brevibus multipartitis digitatis. *Sp. pl.* 1635.

Conferva marina cancellata. *R. syn.* 59. *Dill. musc.* 24. t. 4. f. 22.

Anglis, latticed Conferva.

Habitat in saxis et rupibus submarinis passim. 4. I—XII.

multifida. 18. CONFERVA filamentis æqualibus ramosissimis, ramulis oppositis brevissimis multifidis.

Anglis, tufted Conferva.

Habitat in saxis et rupibus submarinis in comitatibus Hantoniensi et Dorsetiensi. ☉. V—X.

Desc. Fila quadripollicaria, æqualia subgelatinosa, rubra ramosissima; Rami oppositi, longissimi; Ramuli oppositi, brevissimi, multifidi, tenuissimi, remoti, ut verticillati apparent.

spongiosa. 19. CONFERVA filamentis æqualibus subdichotomis, ramulis brevissimis simplicibus undique imbricatis.

Fucus hirsutus, fronde filiformi tereti dichotoma obtecta undique hirsutie. *Mant.* 134.

Fucus teretifolius spongiosus pilosissimus. *R. syn.* 46.

Musculus marinus hirsutius, flagellis longis ramosis subviridibus. *Hist. Ox. III.* 650. f. 15. t. 9. f. 6.

Anglis, sponge Conferva.

Habitat in rupibus et saxis submarinis. 4. I—XII.

** Filamentis anastomosantibus.

reticulata. 20. CONFERVA filamentis reticulato - coadunatis. *Sp. pl.* 1635.

Conferva reticulata. *R. syn.* 59. *Dill. musc.* 20. t. 4. f. 14. *Hall. hist.* 2119.

Conferva reticulata crispa. *Pluk. ph.* t. 24. f. 2. *Hist. Ox. III.* 644. f. 15. t. 4. f. 4.

Anglis, net Conferva.

Habitat in fossis et stagnis, circa Hounslow. ☉. V—X.

*** Filamentis

*** *Filamentis nodosis.*

21. CONFERVA filis simplicissimis setiformibus rec-^{fluviati-}
tis, geniculis crassioribus angulatis. *Sp. pl.* 1635. ^{lis.}

Conferva fluviatilis lubrica setosa, equiseti facie. *Dill.*
musc. 39. t. 7. f. 47.

Corallina fluviatilis non ramosa. *Vaill. paris.* 40.
t. 4. f. 5.

β Conferva fluviatilis nodosa, fucum æmulans. *Dill.*
musc. 39. t. 7. f. 38.

Anglis, horse-tail Conferva.

Habitat in fluviis, prope Bangoriam in rivo dicto Afon
Cegin, inter molendinum vocatum Pandu maes y Geir-
chan et Malun Escob. *Sam. Brewer. prope Ludlow*
D. Littl. Brown. in comitatibus Eboracensi, Cum-
berlandico et Westmorlandico. 4. I—XII.

22. CONFERVA filamentis moniliformibus setaceis *atra.*
ramosissimis atris, articulis globosis subgelatinosis.

Conferva fontana nodosa lubrica, filamentis tenuissi-
mis nigris. *Dill. musc.* 39. t. 7. f. 46.

Anglis, bead Conferva.

Habitat in fontibus et rivulis; in fonte quodam parvo
in lacum dictum Gors Velen prope Llanfaithly, se
exonerante, et in scaturigine quadam campi Gors
Back, dicta Ffynnon bach y Lufg, media via inter
ædes parochiales et Trefadog, in Mona insula D.
Brewer. prope Martin in comitatu Surriensi. 4.
V—XII.

23. CONFERVA filamentis moniliformibus filiformi-^{gelatino-}
bus ramosis, articulis globosis gelatinosis. ^{sa.}

Conferva filis ramosis moniliformibus, articulis glo-
bosis gelatinosis. *Sp. pl.* 1635.

Conferva gelatinosa, globulis punctatis concatenatis.
Hall. hist. 2124.

Conferva fontana nodosa, spermatis ranarum instar
lubrica, major et fusca. *R. syn.* 62. *Dill. musc.*
36. t. 7. f. 42.

CRYPTOGAMIA ALGÆ.

Chara batrachosperma, caule ramoso filamentoso; ramis foliaceis capillaceis, in glomeres lubricos gemiculatos articulatum circa caulem dispositis. *Weis. crypt.* 33. t. 1.

β *Conferva fontana nodosa*, spermatis ranarum instar lubrica, minor et viridis. *Dill. musc.* 37. t. 7. f. 43.

Corallina pinguis ramosa viridis. *Vaill. paris.* 40. t. 7. f. 6.

γ *Conferva stagnalis*, globulis virescentibus mucosis. *Dill. musc.* 38. t. 7. f. 44.

δ *Conferva alpina lubrica*, filamentis nodosis cæruleis. *Dill. musc.* 38. t. 7. f. 45.

Anglis, gelly Conferva.

Habitat in fontibus et rivulis purissimis passim; inter Greenwich et Woolwich; β in rivulo quodam saltus Enfieldensis in Middlesexia. γ; δ. prope Mancunium in Lancastria. D. Harrison. ©. I—XII.

*** *Filamentis geniculatis.*

capillaris. 24. CONFERVA filis geniculatis simplicibus articulis alternatim compressis. *Sp. pl.* 1636.

Conferva filamentis longis geniculatis simplicibus. *Dill. musc.* 25. t. 5. f. 25.

Conferva palustris, sive Filum marinum anglicum. *R. syn.* 60.

Conferva geniculata minima nostras. *Pluk. pb.* t. 84. f. 9. *Hist. Ox.* III. 644. f. 15. t. 4. f. 4.

Anglis, thread Conferva.

Habitat in fossis et stagnis marinis et rivulis. ©. II—IX.

corallinoides. 25. CONFERVA filamentis geniculatis dichotomis; articulis apice incrassatis,

Conferva filamentis geniculatis dichotomis. *Sp. pl.* 1636.

Conferva marina gelatinosa, corallinae instar geniculata crassior. *Dill. musc.* 33. t. 5. f. 3.

Corallina confervoides geniculata alba, geniculis crassiusculis pellucidis. *R. syn.* 34.

Anglis, coralline Conferva.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

- Muscus capillaceus multifidus niger. *B. pin.* 363.
Anglis, palmated Conferva.
 Habitat in fucis, præcipue in fuco nodoso. 4. I—XII.

- tubulosa.* 30. CONFERVA filamentis geniculatis ramosissimis, articulis ovalibus alternatim compressis.
 Conferva marina fistulosa. *Dill. musc.* 34. t. 6. f. 39.
 Conferva marina geniculata albicans, diaphragmatis distinctis. *R. syn.* 60.
 Conferva marina nodosa ex albo rubescens mollis, sed minus lubrica. *R. syn.* 61.
 Ulva filiformis articulata, articulis alternatim compressis. *Sp. pl.* 1632.
Anglis, hollow Conferva.
 Habitat in saxis, rupibus et fucis marinis passim. ☉. IV—X.

- rubra.* 31. CONFERVA filamentis geniculatis ramosissimis, ramis setaceis, articulis cylindricis brevibus.
 Conferva flosculosa. *Ellis. philos. transact.* LVII. 425. t. 18. f. e, E.
 Conferva geniculata ramosissima lubrica longis sparse ramulis. *R. syn.* 61. *Dill. musc.* 35. t. 6. f. 38.
Anglis, red Conferva.
 Habitat in rupibus, saxis et fucis marinis. ☉. V—X.
 Obs. fructificationes sessiles solitariae subrotundæ, filamentis subulato subjectæ.

- purpurascens.* 32. CONFERVA filamentis geniculatis ramosissimis, ramis confertis, articulis cylindricis longis.
 Conferva marina nodosa, Coralloides montani instar ramosa. *Dill. musc.* 36. t. 7. 41.
Anglis, purple Conferva.
 Habitat in saxis et fucis submarinis passim. ☉. V—X.

- nodulosa.* 33. CONFERVA filamentis geniculatis ramosissimis, articulis oblongis, ramulorum subrotundis moniliformibus.
 Conferva marina nodosa lubrica ramosissima et elegantissima

- tissima rubens. *R. fyn.* 62. t. 2. f. 3. *Dill. musc.*
 35. t. 7. f. 40.
Anglis, knotted Conferva.
 Habitat in *saxis, rupibus et fucis*. ☉. V—X.
 Obs. Fructificationes laterales, sessiles, subrotundæ, glomeratæ.
34. CONFERVA filamentis geniculatis ramosissimis, *pellucida*.
 ramis oppositis, articulis cylindricis longissimis.
Anglis, pellucid Conferva.
 Habitat in *rupibus et saxis submarinis*, in *Devonia, Cornubia, Hantonia, et Suffexia*. ☉. V—X.
 Desc. Fila fere semipedalia, geniculata, nitida, pellucida, viridi-purpurea, ramosissima, ramis oppositis, fere triebotomis; articulis cylindricis, æqualibus, longissimis.
35. CONFERVA filamentis geniculatis flexuosis ramosissimis, ramulis brevissimis horizontalibus. *vagabunda*.
 Conferva filamentis flexuosis, ramis ramulisque brevioribus. *Sp. pl.* 1637.
 Conferva marina trichoides, lanæ instar expansa.
R. fyn. 60. *Dill. musc.* 30. t. 5. f. 32.
Anglis, floating Conferva.
 Habitat in *paludibus et fossis maritimis*. ☉. IV—X.
36. CONFERVA filamentis geniculatis ramosissimis *rupestris*.
 viridibus, ramis ramulisque brevioribus confertis.
 Conferva filamentis geniculatis ramosissimis viridibus.
Sp. pl. 1637.
 Conferva marina trichoides ramosior. *Dill. musc.* 28.
 t. 5. f. 29.
 Conferva marina trichoides s. muscus marinus virens tenuifolius. *Pluk. mant.* 53. t. 182. f. 6. *R. fyn.* 60.
 β Conferva fluviatilis trichoides, extremitatibus ramosis.
Dill. giss. musc. 27. t. 25. f. 28.
Anglis, rock Conferva or Sea-beard.
 Habitat in *rupibus et saxis submarinis et in fluviiis passim*.
 4. I—XII.
37. CONFERVA filamentis geniculatis ramosissimis *sericca*.
 elongatis viridibus, ramulis confertis tenuissimis.
 Conferva

Conferva trichoides virgata sericea. Dill. musc. 31.
t. 4. f. 33.

Conferva marina trichoides, ramulis virgatis longioribus glabris. R. syn. 60.

Anglis, silky Conferva.

Habitat in rupibus et saxis submarinis; in insula Shepey copiose; in Rico novo prope Londinum, et ericeto Hounsteano. u. I—XII.

glomerata.

38. CONFERVA filamentis geniculatis ramosissimis, ramulis brevissimis multifidis.

Conferva filamentis geniculatis, ramulis brevioribus multifidis. Sp. pl. 1637.

Conferva fontinalis ramosissima glomeratim congesta.

R. syn. 59. Dill. musc. 28. t. 5. f. 31.

Alga fontalis trichoides. B. pin. 364.

Alga s. Conferva fontalis trichoides. Park. 1260.

Anglis, cluster Conferva.

Habitat in rivulis et fontibus, passim. ☉. IV—X.

fulva.

39. CONFERVA filamentis geniculatis ramosis, ramis ramulisque brevissimis fulvis.

Anglis, tawny Conferva.

Habitat in saxis et facis marinis, in littore Eboracensi.

☉. V—IX.

nigrescens.

40. CONFERVA filamentis geniculatis ramosissimis, ramis longissimis, ramulis subsfasciculatis brevissimis subulatis.

Anglis, blackish Conferva.

Habitat in rupibus et saxis submarinis, prope St. Ives in Cornubia et Exmouth in Devoniam. u. I—XII.

Desc. Fila semipedalia, caespitosa, geniculata, tenuè atro-rubescens, (in sicca nigra) levia, ramosissima; Rami alterni, longi, ramuli brevissimi, subsfasciculati, subulati.

fusca.

41. CONFERVA filamentis geniculatis ramosissimis, ramulis alternis simplicibus.

Anglis, brown Conferva.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Muscus pelagicus pinnatus rubens, ramulis numerosis mollibus latius se spargentibus. *Pluk. ph. t. 48. f. 2.*

Muscus marinus purpureus parvus, foliis oblongis millifolii fere divisura. *R. hist. 79.*

Anglis, scarlet Conferva.

Habitat in saxis, rupibus et fucis marinis. 4. I—XII.

pennata. 46. CONFERVA filamentis geniculatis ramosis, ramis confertis triplicato-pinnatis, laciniis subulatis.

Conferva marina pennata. *Pet. hort. succ. R. syn. 59.*

Anglis, pinnated Conferva.

Habitat in saxis et rupibus submarinis, inter Dover et Margate, in insula Walney in comitatu Lancastriensi. 4. I—XII.

parasitica. 47. CONFERVA filamentis geniculatis ramosis, ramis duplicato-pinnatis.

Anglis, feathered Conferva.

Habitat in fucis, in littoribus Eboracensi, Cornubiensi, et Dorsetiensi passim.

Desc. Fila pollicaria, geniculata, fusca, ramosa, Rami fere pollicares duplicato-pinnati; pinnulis acutis.

ægagropila. 48. CONFERVA filamentis geniculatis ramosissimis e centro confertissimis globum constituentibus. *Sp. pl. 1637.*

Conferva globosa. *Philos. transact. XLI. 498.*

Anglis, globe Conferva, or Moor-balls.

Habitat in lacubus montosis; in Wallingfen-moor, Yorkshire. *Mr. Tho. Knowlton. 4. I—XII.*

BYSSUS. 1208.

Lanugo aut Pulvis simplicissimus.

* *Filamentosa.*

Flos aquæ. 1. BYSSUS filamentis plumosis natantibus. *Sp. pl. 1637.*

Byssus

Byssus latissima papiri instar super aquam sparsa. *R. syn.* 57. *Dill. musc.* 21.

Conferva crustacea tenerrima viridis. *Hall. hist.* 2113. *Anglis*, paper Byssus.

Habitat in aquis stagnantibus. ☉. V—VIII.

2. BYSSUS lanuginosa violacea lignis adnascens. *R. phosphorea.*
syn. 56. *Dill. musc.* 54. t. 1. 6. *Sp. pl.* 1638.

Byssus cærulea cespitosa crispa. *Hall. hist.* 2102.

Byssus purpurea delicatissima, arborum corticibus adnascens, brevissimis et tenuissimis filamentis. *Mich. gen.* 211. t. 9. f. 3.

Fungus violaceus herpatis modo lignis irrepens. *R. hist.* III. 23.

Anglis, violet Byssus.

Habitat in corticibus arborum et lignis putrescentibus. ☉. X—V.

3. BYSSUS filamentis simplicibus tenuissimis æruginosa. *æruginosa.*

Byssus lanuginosa æruginosa. *Dill. musc.* 4. t. f. 7. *Weis. crypt.* 18.

Lichen crusta viridissima. *Hall. hist.* 2097.

Anglis, verdigris Byssus.

Habitat in stipitibus, filicis, et lignis putrescentibus. ☉. VIII—IV.

4. BYSSUS capillacea viridis, filamentis ramosis. *Syst. velutina.*
nat. 721. *Sp. pl.* 1638. *Weis. crypt.* 12.

Byssus viridis, filamentis ramosis. *Hall. hist.* 2101.

Byssus tenerrima viridis velutum referens. *R. syn.* 36. *Dill. musc.* 7. t. 1. f. 14.

Byssus terrestris viridis herbacea et mollissima, filamentis ramosis et non ramosis. *Mich. gen.* 211. t. 89. f. 5.

Anglis, velvet Byssus.

Habitat in terris umbrosis et corticibus arborum humidiusculis. ☉. X—VI.

5. BYSSUS filamentis simplicibus ramosisque rubris. *rubra.*
Anglis, red Byssus.

Habitat

Habitat in saxis et rupibus, præcipue marinis. 4.
I—XII.

nigra.

6. BYSSUS filamentis pulverulentis ramosis atris.

Byssus nigra velutina. *Hall. hist.* 2104.

Byssus petræa nigerrima fibrosa. *R. syn.* 57. *Dill. musc.*
9. t. 1. f. 18.

Byssus minima saxatilis nigra ramosissima, viticulis
nudis. *Mich. gen.* 212. t. 90. f. 5.

Anglis, black Byssus.

Habitat in rupibus et saxis grandioribus, in boreali parte
Angliæ et in Wallia. 4. I—XII.

aurea.

7. BYSSUS filamentis pulverulentis simplicibus ramo-
sisque.

Byssus capillacea pulverulenta, fructificationibus spar-
sis filamentis simplicibus ramosisque. *Gledit. fung.*
18. *Sp. pl.* 1638. *Weis. crypt.* 13.

Byssus cespitosa pulvinata crocea. *Hall. hist.* 2106.

Byssus petræa crocea glomerulis lanuginosis. *Dill.*
musc. 8. t. 1. f. 16.

Byssus aureus derbiensis humifusus. *Pet. gaz.* 1. t. 13.
f. 3. *R. syn.* 56.

Byssus minima saxatilis aurea inodora, filamentis
partim simplicibus, partim ramosis. *Mich. gen.*
210. t. 89. f. 2.

Anglis, golden Byssus.

Habitat in rupibus cameratis passim. 4. I—XII.

fulva.

8. BYSSUS filamentis ramosis fulvis.

Byssus arborea crocea fibrosa. *R. syn.* 57. *Dill.*
musc. *Dill. musc.* 9. t. 1. f. 17.

Anglis, tawny Byssus.

Habitat in lignis putrescentibus humidiusculis. 4.
I—XII.

barbata.

9. BYSSUS filamentis erectiusculis summitatibus ramo-
sis subfastigiatis fulvis.

Byssus arborea barbata fulvi coloris. *R. syn.* 57.
Dill. musc. 9. t. 1. f. 19.

Pseudo-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Chara batrachosperma, caule ramoso filamentoso; ramis foliaceis capillaceis, in glomeres lubricos gemiculatos articulatum circa caulem dispositis. *Weis. crypt. 33. t. 1.*

β *Conferva fontana nodosa*, spermatis ranarum instar lubrica, minor et viridis. *Dill. musc. 37. t. 7. f. 43.*
Corallina pinguis ramosa viridis. Vaill. paris. 40. t. 7. f. 6.

γ *Conferva stagnalis*, globulis virescentibus mucosis. *Dill. musc. 38. t. 7. f. 44.*

δ *Conferva alpina lubrica*, filamentis nodosis cæruleis. *Dill. musc. 38. t. 7. f. 45.*

Anglis, gelly Conferva.

Habitat in fontibus et rivulis purissimis passim; inter Greenwich et Woolwich; β in rivulo quodam saltus Enfieldensis in Middlesexia. γ; δ. prope Mancunium in Lancastria. D. Harrison. ☉. I—XII.

*** *Filamentis geniculatis.*

capillaris. 24. CONFERVA filis geniculatis simplicibus articulis alternatim compressis. *Sp. pl. 1636.*

Conferva filamentis longis geniculatis simplicibus. Dill. musc. 25. t. 5. f. 25.

Conferva palustris, sive Filum marinum anglicum. R. syn. 60.

Conferva geniculata minima nostras. Pluk. ph. t. 84. f. 9. Hist. Ox. III. 644. f. 15. t. 4. f. 4.

Anglis, thread Conferva.

Habitat in fossis et stagnis marinis et rivulis. ☉. II—IX.

corallinoides. 25. CONFERVA filamentis geniculatis dichotomis; articulis apice incrassatis.

Conferva filamentis geniculatis dichotomis. Sp. pl. 1636.

Conferva marina gelatinosa, corallinae instar geniculata crassior. Dill. musc. 33. t. 5. f. 3.

Corallina confervoides geniculata alba, geniculis crassiusculis pellucidis. R. syn. 34.

Anglis, coralline Conferva.

Habitat

Habitat in saxis et rupibus submarinis passim. ☉. V—X.

26. CONFERVA filamentis geniculatus dichotomis, *setacea*.
ramis longissimis setaceis, articulis cylindricis.

Conferva marina gelatinosa, corallinæ instar geniculata tenuior. *Dill. musc.* 33. t. 6. f. 37.

Corallina confervoides gelatinosa rubens, ramulis et geniculis perangustis. *R. syn.* 34.

Anglis, setaceous Conferva.

Habitat in saxis et rupibus submarinis passim. ☉. V—X.

Obf. ramis longis setaceis.

27. CONFERVA filamentis geniculatis ramosis, ramis *elongata*.
dichotomis longis setaceis, articulis brevissimis.

Anglis, pointed Conferva.

Habitat in saxis et rupibus submarinis, in Devoniam, Cornubiam, Suffexiam, et in insula Mona, passim. ☉. IV—X.

Desc. Fila *dedrantal* et *pedalia*, crassitie fili *emporetici* tenuioris, articulata, *lævia*, *fusco-purpurea*, basi *ramosa*; Rami *dichotomi*, *longissimi*, *setacei*; articulis *brevissimis*.

28. CONFERVA filamentis geniculatis dichotomis, *ciliata*.
apicibus conniventibus, articulis apice ciliatis.

Conferva ciliata. *Ellis. philos. transact.* LVII. 425. t. 18. f. h. H.

Anglis, ciliated Conferva.

Habitat in fucis tenuioribus passim. ☉. V—IX.

29. CONFERVA filamentis geniculatis subdichotomis *polymorpha*.
fastigiatis, ramis fasciculatis.

Conferva filamentis geniculatis, ramis fasciculatis.

Sp. pl. 163. *Fl. dan.* 395. *Ellis. philos. transact.* LVII. 424. t. 18. f. a, A, b, B.

Conferva marina geniculata nigra palmata. *Dill. musc.* 32. t. 6. f. 35.

Conferva marina geniculata ramosissima lubrica, brevibus et palmatim congestis ramulis. *R. syn.* 61.

- Muscus capillaceus multifidus niger. *B. pin.* 363.
Anglis, palmated Conferva.
 Habitat in fucis, præcipue in fuco nodoso. \mathcal{C} . I—XII.
- tubulosa.* 30. CONFERVA filamentis geniculatis ramosissimis, articulis ovalibus alternatim compressis.
 Conferva marina fistulosa. *Dill. musc.* 34. t. 6. f. 39.
 Conferva marina geniculata albicans, diaphragmatis distinctis. *R. syn.* 60.
 Conferva marina nodosa ex albo rubescens mollis, sed minus lubrica. *R. syn.* 61.
 Ulva filiformis articulata, articulis alternatim compressis. *Sp. pl.* 1632.
Anglis, hollow Conferva.
 Habitat in saxis, rupibus et fucis marinis passim. \odot . IV—X.
- rubra.* 31. CONFERVA filamentis geniculatis ramosissimis, ramis setaceis, articulis cylindricis brevibus.
 Conferva flosculosa. *Ellis. philos. transact.* LVII. 425. t. 18. f. e, E.
 Conferva geniculata ramosissima lubrica longis sparse ramulis. *R. syn.* 61. *Dill. musc.* 35. t. 6. f. 38.
Anglis, red Conferva.
 Habitat in rupibus, saxis et fucis marinis. \odot . V—X.
 Obs. fructificationes sessiles solitariae subrotundæ, filamentis subulato subjectæ.
- purpurascens.* 32. CONFERVA filamentis geniculatis ramosissimis, ramis confertis, articulis cylindricis longis.
 Conferva marina nodosa, Coralloides montani instar ramosa. *Dill. musc.* 36. t. 7. 41.
Anglis, purple Conferva.
 Habitat in saxis et fucis submarinis passim. \odot . V—X.
- nodulosa.* 33. CONFERVA filamentis geniculatis ramosissimis, articulis oblongis, ramulorum subrotundis moniliformibus.
 Conferva marina nodosa lubrica ramosissima et elegantissima

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Conferva trichoides virgata sericea. *Dill. musc.* 31.
t. 4. f. 33.

Conferva marina trichoides, ramulis virgatis longioribus glabris. *R. syn.* 60.

Anglis, silky Conferva.

Habitat in rupibus et saxis submarinis; in insula Shepey copiose; in Rivo novo prope Londinum, et ericeto Hounsteano. 4. I—XII.

*glomera-
ta.*

38. **CONFERVA** filamentis geniculatis ramosissimis, ramulis brevissimis multifidis.

Conferva filamentis geniculatis, ramulis brevioribus multifidis. *Sp. pl.* 1637.

Conferva fontinalis ramosissima glomeratim congesta.

R. syn. 59. *Dill. musc.* 28. t. 5. f. 31.

Alga fontalis trichoides. *B. pin.* 364.

Alga f. Conferva fontalis trichoides. *Park.* 1260.

Anglis, cluster Conferva.

Habitat in rivulis et fontibus, passim. ☉. IV—X.

fulva.

39. **CONFERVA** filamentis geniculatis ramosis, ramis ramulisque brevissimis fulvis.

Anglis, tawny Conferva.

Habitat in saxis et fucis marinis, in littore Eboracensi.

☉. V—IX.

nigrescens.

40. **CONFERVA** filamentis geniculatis ramosissimis, ramis longissimis, ramulis subfasciculatis brevissimis subulatis.

Anglis, blackish Conferva.

Habitat in rupibus et saxis submarinis, prope St. Ives in Cornubia et Exmouth in Devoniam. 4. I—XII.

Desc. Fila semipedalia, caespitosa, geniculata, tenue atro-rubescens, (in sicca nigra) lævia, ramosissima; Rami alterni, longi, ramuli brevissimi, subfasciculati, subulati.

fusca.

41. **CONFERVA** filamentis geniculatis ramosissimis, ramulis alternis simplicibus.

Anglis, brown Conferva.

Habitat

Habitat in saxis et rupibus submarinis. ☉. VI—X.

Desc. Fila tri et quadripolicaria, geniculata, atro-fusca aut rubescentia, lævia, ramosissima, Rami alterni, longi, ramuli, alterni simplices, brevi, distantes; Fructificationes ramuli terminatæ et laterales sessiles, parvæ, subrotundæ, glomeratæ.

42. CONFERVA filamentis geniculatis ramosissimis *fucoides*. ramulis multifidis, inferioribus fasciculatis fructiferis.

Anglis, fucus Conferva.

Habitat in rupibus, saxis et fucis marinis passim. 4. I—XII.

Desc. Fila pedalia, geniculata, lævia atro-rubescens, ramosissima, Rami alterni; ramuli multifidi, subdichotomi, inferiores fasciculati, fructiferi. Fructificationes terminales, radiatæ, parvæ.

43. CONFERVA filamentis geniculatis ramosis, geniculis villosis, ramis subsimplicibus distantibus.

Anglis, hairy Conferva.

Habitat in saxis et rupibus submarinis, in Cornubia, sed rarius. ☉. V—IX.

Desc. Fila semipedalia, geniculata, viridia, diaphana, ramosa; Rami longi, setacei, subsimplices; Geniculis villosis s. ramulis brevissimis, tenuissimis, verticillatis.

44. CONFERVA filamentis geniculatis ramosissimis, *imbricata*. ramis acutis, ramulis verticillatis imbricatis dichotomis.

Muscus marinus hirsutus, flagellis longioribus, rarius divisus, ruber. *Hist. Ox. III. 650. f. 15. t. 9. f. 7.*

Anglis, imbricated Conferva.

Habitat in rupibus saxis et fucis marinis. 4. I—XII.

45. CONFERVA filamentis geniculatis ramosis, ramis *coccinea*, alternis duplicato-pinnatis, laciniis multifidis.

Conferva plumosa. *Ellis. philos. transact. LVII. 425. t. 18. f. c; c; d, D.*

Muscus marinus folio multifido. *Johns. merc. bot. II.*

Muscus pelagicus pinnatus rubens, ramulis numerosis mollibus latius se spargentibus. *Pluk. ph. t. 48. f. 2.*

Muscus marinus purpureus parvus, foliis oblongis millifolii fere divisura. *R. hist. 79.*

Anglis, scarlet Conferva.

Habitat in saxis, rupibus et fucis marinis. 4. I—XII.

pennata. 46. CONFERVA filamentis geniculatis ramosis, ramis confertis triplicato-pinnatis, laciniis subulatis.

Conferva marina pennata. *Pet. hort. sicc. R. syn. 59.*

Anglis, pinnated Conferva.

Habitat in saxis et rupibus submarinis, inter Dover et Margate, in insula Walney in comitatu Lancastriensi. 4. I—XII.

parasitica. 47. CONFERVA filamentis geniculatis ramosis, ramis duplicato-pinnatis.

Anglis, feathered Conferva.

Habitat in fucis, in littoribus Eboracensi, Cornubiensi, et Dorsetiensi passim.

Desc. Fila pollicaria, geniculata, fusca, ramosa, Rami fere pollicares duplicato-pinnati; pinnulis acutis.

*ægagropi-
la.* 48. CONFERVA filamentis geniculatis ramosissimis e centro confertissimis globum constituentibus. *Sp. pl. 1637.*

Conferva globosa. *Philos. transact. XLI. 498.*

Anglis, globe Conferva, or Moor-balls.

Habitat in lacubus montosis; in Wallingfen-moor, Yorkshire. *Mr. Tho. Knowlton. 4. I—XII.*

BYSSUS. 1208.

Lanugo aut Pulvis simplicissimus.

* *Filamentosa.*

Flos aquæ. 1. BYSSUS filamentis plumosis natantibus. *Sp. pl. 1637.*

Byssus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Habitat in saxis et rupibus, præcipue marinis. 4.
I—XII.

nigra.

6. BYSSUS filamentis pulverulentis ramosis atris.

Byssus nigra velutina. *Hall. hist.* 2104.

Byssus petræa nigerrima fibrosa. *R. syn.* 57. *Dill. musc.*
9. t. 1. f. 18.

Byssus minima saxatilis nigra ramosissima, viticulis
nudis. *Mich. gen.* 212. t. 90. f. 5.

Anglis, black Byssus.

Habitat in rupibus et saxis grandioribus, in boreali parte
Angliæ et in *Wallia*. 4. I—XII.

aurea.

7. BYSSUS filamentis pulverulentis simplicibus ramo-
sisque.

Byssus capillacea pulverulenta, fructificationibus spar-
sis filamentis simplicibus ramosisque. *Gledit. fung.*
18. *Sp. pl.* 1638. *Weis. crypt.* 13.

Byssus cespitosa pulvinata crocea. *Hall. hist.* 2106.

Byssus petræa crocea glomerulis lanuginosis. *Dill.*
musc. 8. t. 1. f. 16.

Byssus aureus derbiensis humifusus. *Pet. gaz.* 1. t. 13.
f. 3. *R. syn.* 56.

Byssus minima saxatilis aurea inodora, filamentis
partim simplicibus, partim ramosis. *Mich. gen.*
210. t. 89. f. 2.

Anglis, golden Byssus.

Habitat in rupibus cameratis passim. 4. I—XII.

fulva.

8. BYSSUS filamentis ramosis fulvis.

Byssus arborea crocea fibrosa. *R. syn.* 57. *Dill.*
musc. *Dill. musc.* 9. t. 1. f. 17.

Anglis, tawny Byssus.

Habitat in lignis putrescentibus humidiusculis. 4.
I—XII.

barbata.

9. BYSSUS filamentis erectiusculis summitatibus ramo-
sis subfastigiatis fulvis.

Byssus arborea barbata fulvi coloris. *R. syn.* 57.
Dill. musc. 9. t. 1. f. 19.

Pseudo-

Pseudo-spongia fungoides fulva lignis adnascens. R.
hist. III. 23.

Anglis, bearded Byssus.

Habitat in lignis et truncis arborum putrescentibus. 4.
I—XII.

10. BYSSUS filamentis ramosissimis, ramulis fascicu-*candida*.
latis albidis.

Byssus tenerrima villosa et elegantissime ramulosa. R.

syn. 476. t. 23. Dill. musc. 7. t. 1. f. 15.

Anglis, white Byssus.

Habitat in foliis et lignis putrescentibus. ☉. IX—IV.

11. BYSSUS filamentis longissimis tenuissimis ramosis *septica*.
implexis.

Byssus capillacea molissima parallela fragillissima pal-
lida. *Syst. nat. III. 235.*

Byssus longissima tenerrima nivea. *Hall. hist. 2108.*

Byssus major speluncis et cellis vinariis innascens,
filtrum vel pannum laneum simulans, primum al-
ba, dein nigra, filamentis tenuissimis non ramosis.

Mich. gen. 211. t. 89. f. 9.

Byssus evanida floccosa nivea. *Dill. musc. 5. t. 1.*
f. 9.

Fungus niveus aqueus, lignis cellarum vinariarum
adhærens. R. *syn. 26. hist. III. 23.*

Fungus fimosus niveus, ramosissimus mollis. R. *syn.*
13.

■ *Byssus tenerrima murina doliaris.* *Dill. musc. 6. t. 1.*
f. 12.

Fungus spongiosus niger reticulatus, doliolis vinosis
adnascens. R. *syn. 57.*

Anglis, flaky, β mouse-skin Byssus.

Habitat in cellis et cryptis subterraneis humidiusculis.
β supra dolia vinosa. 4. V—XII.

12. BYSSUS capillacea perennis cinerea tenax rupi *crypta-*
innata. *Sp. pl. 1638.* *rum.*

Byssus albida brevis setacea. *Dill. musc. 10. t. 1.*
f. 20.

Anglis, cave Byssus.

Habitat

Habitat in rupibus calcareis cameratis passim. 4.
I—XII.

** Pulverulentæ.

antiquita- 13. BYSSUS pulverulenta atra. *Sp. pl.* 1638. *It. gotb.*
tis. 298.

Anglis, foot Byssus.

Habitat in muris antiquis, rupibus et saxis grandioribus,
passim. 4. I—XII.

saxatilis. 14. BYSSUS pulverulenta cinerea rupes operiens. *Sp.*
pl. 1638. *Fl. suec.* 1179.

Anglis, rock Byssus.

Habitat in rupibus et saxis calcareis in comitatibus West-
morlandico, Cumberlandico, Eboracensi et Derbi-
ensi. 4. I—XII.

Jolithus. 15. BYSSUS pulverulenta sanguinea saxis innascens.
Sp. pl. 1638.

Byssus germanica minima-saxatilis aurea violæ odorem
spirans. *Mich. gen.* 210. t. 89. f. 3.

Jolithus f. Lapis violaceus. *Swenck foss.* 382.

Lichen crustaceus, iridis odore, ruberrimus. *Hall. hist.*
2090.

Anglis, red Byssus.

Habitat in rupibus et saxis quartzosis, in umbris hu-
midis. 4. I—XII.

botryoides. 16. BYSSUS pulverulenta viridis. *Sp. pl.* 1639. *Weis.*
crypt. 15.

Byssus botryoides saturate virens. *R. syn.* 56. *Dill.*
musc. 3. t. 1. f. 5.

Lichen crustaceus, verrucosus viridis. *Hall. hist.* 2089.

Anglis, green Byssus.

Habitat in cortice arborum, parietibus et terra humida et
umbrosa. 4. I—XII.

candela- 17. BYSSUS pulverulenta flava, lignis adnascens. *R.*
ris. *syn.* 56. *Dill. musc.* 3. t. 1. f. 4. *Sp. pl.* 1639.
Weis. crypt. 16.

Lichen

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Alectorolophoides fulcis crispis. *Battar. fung.* 39.
t. 14.

Fungus angulosus et velut in lacinias sectus. *B. pin.*
371. *Vaill. paris. t. II. f. 9—15.*

Fungus luteus s. pallidus, Chanterelle dictus se con-
torquens esculentus. *B. hist. III. 832. R. syn. 2.*

Fungi lethales. *Ger. em.* 1380.

Anglis, chantarelle Agaric.

Habitat in sylvis et sepibus. VIII—X.

quinque- 2. AGARICUS stipitatus, pileo subflavescente partito,
partitus. lamellis albidioribus interne dentato-connexis. *Sp.*
pl. 1640.

Agaricus caulescens pileo cinereo quinquepartito, la-
mellis albis. *Fl. suec.* 1222.

Anglis, lobed Agaric.

Habitat in pratis, pascuis et nemorosis. IX. X.

campe- 3. AGARICUS stipitatus, pileo convexo squamato al-
stris. bido, lamellis rufis. *Sp. pl.* 1641. *Schæff. fung.*
t. 33.

Amanita petiolo annulato, pileo campaniformi albido,
lamellis roseis. *Hall. hist.* 2443.

Fungus pileolo lato et rotundo. *B. pin.* 370. *Battar.*
29. t. 7. f. A.

Fungus campestris albus superne, inferne rubens. *B.*
hist. III. 824. R. syn. 2. Mich. gen. 174.

Fungus esculentus. *Park.* 1317.

georgii. β Agaricus stipitatus pileo flavo convexo, lamellis albis.
Sp. pl. 1642.

Amanita pileo flavo oris striatis et lanuginosis, lamellis
albis. *Hall. hist.* 2367.

Fungus esculentus pileo et lamellis albis. *R. syn.* 2.

Fungus croceus magnus. *Sterb. fung.* 63. t. II. f. 3.

Anglis, common Agaric or Mushroom.

Habitat in pratis et pascuis. VIII—X.

Obs. in β, succus luteus est.

integer. 4. AGARICUS stipitatus, pileo convexiusculo viscido
purpureo, lamellis omnibus æqualibus albis.

Agaricus stipitatus, lamellis omnibus magnitudine
æqualibus

æqualibus. *Sp. pl.* 1640. *Schæf. fung. t.* 15, 16, 58, 75.

Amanita petiolo farto, pileolo convexo umbilicato, oris rugosis, lamellis crassis et albis. *Hall. hist.* 2373.

Omphalomyces coccineus. *Battar. fung. 37. t.* 15. *f. E.*

Fungus major rubescens, pediculo brevi crasso, lamellis crebris albentibus.

Anglis, red Agaric.

Habitat in sylvis et nemorosis. VIII—X.

5. AGARICUS stipitatus, pileo convexiusculo viscido luteus. luteo, lamellis omnibus æqualibus flavis.

Fungus glutine flavo limacino resplendens. *Vaill. par.* 62?

Anglis, yellow Agaric.

Habitat in pratis et pascuis sub arboribus, et nemorosis. VIII—X.

Obs. affinis præcedenti.

6. AGARICUS stipitatus, pileo convexo-fuscescenti, bulbosus. lamellis cærulescentibus, stipite brevi basi bulboso violaceo.

Agaricus cærulescens. *Schæff. fung. t.* 34?

Fungus pileo magno orbiculari sublivido, lamellis albis, pediculo brevi bulbiformi violaceo. *R. syn.* 3.

Anglis, bulbous Agaric.

Habitat in pratis et pascuis, sed rarius. Prope Newington. D. Dillenius. IX—XII.

7. AGARICUS stipitatus, pileo rimoso : margine vio-violaceus. laceo tomentoso, stipite cærulescente, lana ferruginea. *Sp. pl.* 1641. *Schæff. fung. t.* 3. 56.

Amanita violaceus, petiolo bulboso annulato, pileo plano. *Hall. hist.* 2455.

Fungus cæruleus major. *Buxb. cent. IV. t.* 9.

Fungus esculentus bulbosus dilute purpureus. *Mich. gen.* 149. *t.* 74. *f.* 1.

Fungus major violaceus. *Vaill. paris.* 67.

Anglis, violet Agaric.

Habitat in sylvis et sepibus. VIII—XI.

amethysti- 8. AGARICUS stipitatus, pileo subcampanulato lævi,
nus. lamellis stipiteque violaceis.

Amanita obscure violaceus, pileolo conico. *Hall. hist.*
2458?

Fungus minor amethystinus. *Vaill. paris.* 67.

Fungus parvus violaceus marmoreus perniciosus. *Sterb.*
72. t. 23.

Anglis, amethyst Agaric.

Habitat in sylvis. VIII—X.

muscari- 9. AGARICUS stipitatus, lamellis dimidiatis solitariis,
us. stipite volvato: apice dilatato, basi ovato. *Sp. pl.*
1640. *Schæff. t.* 27, 28.

Amanita petiolo annulato, pileo sanguineo, lamellis
albis. *Hall. hist.* 2373.

Fungus, &c. *Mich. gen.* 187, 188. t. 78. f. 1, 2.

Fungi muscas interficientis III. species aut varietas.
B. pin. 373.

Mell muscarum venenosum. *Sterb. fung.* 214. t. 22.
f. d.

Anglis, fly Agaric.

Habitat in sylvis et sepibus solo arenoso. VIII—X.

procerus. 10. AGARICUS stipitatus, pileo hemisphærico lace-
rato-squamoso rufescente-cinereo, lamellis albis,
stipite longo cylindrico acetabulo inserto.

Agaricus procerus. *Scop. carn.* 1465. *Schæff. fung.*
t. 22, 23.

Amanita petiolo procero squamoso annulato, pileo
conico squamoso subspadiceo, laminis albis. *Hall.*
hist. 190. et *nomencl.* 2366*.

Agaricus petiolo procero annulato, in acetabulum
pilei inmisso, pileo squamoso et maculato, lamellis
albis. *Hall. hist.* 2371.

Fungus major, pediculo longo, modice crasso, lamel-
lis albis creberrimis, superne ad margines apparen-
tibus. *R. syn.* 4?

Fungus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

- lactifluus.* 14. AGARICUS stipitatus, pileo plano carneo lactescente, lamellis rufis, stipite longo carneo. *Sp. pl.* 1641. *Schæff. fung. t. 5.*
 Agaricus rubescens. *Schæff. fung. t. 73?*
 Fungus pileolo lato pumiceo lacteum et dulcem succum fundens. *B. pin. 371.*
 Fungus lactescens piperatus rufus. *Vaill. paris. 61.*
Anglis, milky Agaric.
 Habitat in sylvis. IX. X.
- dulcis.* 15. AGARICUS stipitatus, pileo convexo albo lactescente, lamellis albidis, stipite longo albo.
 Fungus lactescens non acris. *R. syn. 4.*
Anglis, mild Agaric.
 Habitat in sylvis. VII—X.
- viridis.* 16. AGARICUS stipitatus, pileo convexo viridi, lamellis albidis, stipite longo virescente.
 Agaricus virens. *Scop. carn. 1570. Schæff. fung. t. 1.*
 Amanita annulatus pileo convexo cæruleo-viridi lamellis roseo cæruleis. *Hall. hist. 2444.*
 Fungus esculentus, pileolo pulvinato viridi, inferne cum pediculo albo. *Mich. gen. 152.*
 Fungus magnus viridis. *Sterb. fung. 67. t. 5. f. c.*
R. syn. 4.
Anglis, green Agaric.
 Habitat in sylvis. VIII—X.
- squamosus.* 17. AGARICUS stipitatus, pileo convexiusculo flavescente lacerato-squamoso, lamellis albidis crenatis, stipite brevi solido pallido. *Schæff. fung. t. 29, 30.*
 Amanita pileo flavescente squamoso, lamellis terreis obscuris, annulo araneoso. *Hall. hist. 2412.*
Anglis, squamous Agaric.
 Habitat in lignis putrescentibus, præsertim salicis. VIII, —X.
- viscidus.* 18. AGARICUS stipitatus pileo purpurascente-fusco viscido, lamellis fusco-purpurascensibus. *Sp. pl.* 1642.

Agaricus truncatus. Schæf. fung. t. 251.

Agaricus pileo convexo subrubro viscido, lamellis una cum bulbofo nudo et pleno stipite albis. Scop. carn. 1477.

Amanita dura ex fusco-rubens quercina. R. syn. 5.
Anglis, viscid Agaric.

Habitat in nemorosis et sylvis. IX. X.

19. **AGARICUS** stipitatus, pileo convexo rufo, lamellis flavescentibus, stipite longo flavo. *cinnamomeus.*

Agaricus stipitatus, pileo sordide flavo, lamellis luteo-rufis. Sp. pl. 1642. Fl. suec. 1205?

Agaricus caulescens flavus, pileo plano. Fl. lapp. 594.

Agaricus croceus. Schæff. fung. t. 4.

Anglis, brown Agaric.

Habitat in sylvis passim. VIII—X.

20. **AGARICUS** stipitatus, pileo convexo luteo, lamellis virescentibus, stipite flavo. *fascicularis.*

Agaricus lateritius. Schæff. fung. t. 49.

Amanita annulo araneoso, pileo flavo, lamellis ex flavo fuliginosis. Hall. hist. 2482.

Fungus major subluteus, lamellis luridis. Buxb. cent. iv. t. 4?

Fungus mediæ magnitudinis, pileolo superne e rufo flavicante, lamellis subtus sordide virentibus. R. syn. 10.

Anglis, clustered Agaric.

Habitat in lignis putrescentibus. VIII—X.

21. **AGARICUS** stipitatus, pileo convexo flavescente, lamellis testaceis, stipite flavo. *testaceus.*

Agaricus incurvus. Schæff. fung. t. 65.

Amanita leoninus annulo arenoso. Hall. hist. 243.

Anglis, testaceous Agaric.

Habitat in lignis putrescentibus et ad radices arborum. VIII—X.

22. **AGARICUS** stipitatus, pileo hemisphærico fusco-flavescente, lamellis rufis, stipite longo tomentoso. *mutabilis.*

Agaricus mutabilis. Scop. carn. 1515. Schæff. fung. t. 9.

Fungus fasciculosus, pileo orbiculari lutescente, pediculo fusco tenerrime villosa, lamellis ex flavo candidantibus. R. syn. 9.

Anglis, variable Agaric.

Habitat in sylvis, sepibus et lignis putrescentibus. VIII.—X.

lividus. 23. AGARICUS stipitatus, pileo convexo livido, lamellis albidis, stipite longo albo.

Agaricus multiformis. Schæff. fung. t. 14.

Amanita orbicularis sublivida lamellis subfuscis, pilei oris leviter purpurascens. R. syn. 5?

Anglis, livid Agaric.

Habitat in sylvis et ad sepes. IX—V.

equestris. 24. AGARICUS stipitatus, pileo pallido; disco stellatim luteo, lamellis sulphureis. Sp. pl. 1643.

Agaricus tuberculatus. Schæff. fung. t. 79?

Fungus parvus galeriformam exprimens rufus. R. syn. 7?

Fungus vaccinus. Sterb. fung. 210. t. 21. f. c.

Anglis, starry Agaric.

Habitat in pascuis et nemorosis. IX. X.

pratensis. 25. AGARICUS stipitatus, pileo convexiusculo pallide lutescente lamellis stipiteque concoloribus.

Agaricus collinus. Schæff. fung. t. 220?

Anglis, field Agaric, or Champignon.

Habitat in pratis pascuis et ericetis. VIII—X.

farinaceus. 26. AGARICUS stipitatus, pileo convexiusculo purpurascens-fusco, lamellis convexis distantibus incarnatis pulverulentis, stipite longo.

Agaricus laccatus. Scop. carn. 1530. Schæff. fung. t. 13.

Amanita petiolo fistuloso lineato, pileo convexo flavo-rufo. Hall. hist. 2436.

Fungus multiplex sordide carneus. Vaill. paris. 66?

Anglis, powdered Agaric.

Habitat

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Agaricus fugax. *Schæff. fung. t. 67, 68.*

Agaricus fulcescens. *Schæff. fung. t. 17.*

Agaricus rufocandidus. *Schæff. fung. t. 201.*

Agaricus margaritaceus. *Schæff. fung. t. 216.*

Amanita pileo ovato striato cinereo, anulo fugaci.
Hall. hist. 2479.

Fungus multiplex ovatus cinereus. *Vaill. paris. 73.*
t. 12. f. 10, 11.

Hydrophorus oris laceris alter. *Battar. 54. t. 27.*
f. G.

Anglis, striated Agaric.

Habitat in ruderatis et ad sepes. IX. X.

campanu- 31. AGARICUS stipitatus, pileo campanulato striato
latus. pellucido, lamellis adscendentibus, stipite nudo.

Sp. pl. 1643.

Agaricus plicatus, bruneus et pallefcens. *Schæff.*
fung. t. 31. 22, et 211.

Amanita albus, petiolo gracile prælongo, pileolo stri-
ato conico. *Hall. hist. 2345.*

Fungus minimus fimetarius pileo undique et utrinque
striato cinereo aut murino, pediculo lanuginoso et
fistuloso. *Mich, gen. 167. t. 75. f. 6.*

Fungus multiplex obtuse conicus, colore griseo mu-
rino. *Vaill. paris. 71. t. 12. f. 12.*

Fungus parvus, pediculo oblongo firmo lento, pileolo
in medio fastigiato, striis exterius apparentibus.
R. syn. 8.

β Amanita petiolo procero gracilissimo, pileolo conico
striato murino inferne albo. *Hall. hist. 2391.*

Amanita parva verna utrinque striata fusca, pileo ob-
tuse coniformi. *Dill. giss. 184. R. syn. 8.*

Anglis, bell Agaric.

Habitat in pratis, pascuis et sylvis. VIII.—X.

separatus. 32. AGARICUS stipitatus, pileo lævi livido, lamellis
nigricantibus separatis, stipite bulboso volvato. *Sp.*
pl. 1643. Fl. succ. 1220.

Agaricus candicans. *Schæff. fung. t. 217?*

Anglis, separated Agaric.

Habitat in pratis et pascuis. IX. X.

33. AGA-

33. AGARICUS stipitatus, pileo hemisphærico livido-*adnatus*,
 flavescente, lamellis adnatis atro-griseis, stipite lon-
 gissimo nudo livido.
 Agaricus helvolus. *Schæff. fung.* 210.
 Fungus parvus pediculo oblongo pileolo hemisphærico,
 ex albido subluteus. *R. syn.* 7.
Anglis, long-stalked Agaric.
 Habitat in pratis et pascuis e fimo bubulo. IX. X.
34. AGARICUS stipitatus, pileo convexo acuto gri-*mammo-*
 seo, lamellis convexis griseis crenatis, stipite nudo. *sus.*
 Fungus minor capitulo pileum nauticum referente,
 pediculo longiore. *Buxb. cent. IV.* 13. t. 21. f. 1.
Anglis, crenated Agaric.
 Habitat in sylvis et nemorosis. VIII—XII.
35. AGARICUS stipitatus, pileo hemisphærico viscido *chypeatus*.
 acuminato, lamellis albis, stipite longo cylindrico
 albo. *Schæff. fung. t.* 52. f. 7, 8, 9.
 Amanita radice caudata, petiolo fistuloso procero, pileo
 umbonato striato cinereo, lamellis albis. *Hall.*
hist. 2388.
 Fungus clypeatus in medio protuberans. *Vaill. paris.*
 68.
Anglis, shield Agaric.
 Habitat in sylvis, nemorosis et lignis putrescentibus. IX.
 X.
36. AGARICUS stipitatus, pileo campanulato obtuso *aquosus*.
 fusco-flavescente, lamellis atro-griseis, stipite fistu-
 loso griseo villosa.
 Agaricus truncorum. *Scop. carn.* 1480. *Schæff. fung.*
 t. 6.
 Fungus multiplex campaniformis colore fusco. *Vaill.*
paris. 73. t. 12. f. 5, 6.
 Fungus parvus lethalis galericulatus. *Lob. ic.* 272.
R. syn. 7.
 Fungus parvus galericulatus alter flavus. *B. pin.* 373.
Anglis, watery Agaric.
 Habitat in lignis putridis humidis. VIII—X.
37. AGARI-

Agaricus fugax. *Schæff. fung. t. 67, 68.*

Agaricus fulcescens. *Schæff. fung. t. 17.*

Agaricus rufocandidus. *Schæff. fung. t. 201.*

Agaricus margaritaceus. *Schæff. fung. t. 216.*

Amanita pileo ovato striato cinereo, anulo fugaci.
Hall. hist. 2479.

Fungus multiplex ovatus cinereus. *Vaill. paris. 73.
t. 12. f. 10, 11.*

Hydrophorus oris laceris alter. *Battar. 54. t. 27.
f. G.*

Anglis, striated Agaric.

Habitat in ruderatis et ad sepes. IX. X.

campanu- 31. AGARICUS stipitatus, pileo campanulato striato
latus. pellucido, lamellis adscendentibus, stipite nudo.

Sp. pl. 1643.

Agaricus plicatus, bruneus et pallefcens. *Schæff.
fung. t. 31. 22, et 211.*

Amanita albus, petiolo gracile prælongo, pileolo stri-
ato conico. *Hall. hist. 2345.*

Fungus minimus fimetarius pileo undique et utrinque
striato cinereo aut murino, pediculo lanuginoso et
fistuloso. *Mich, gen. 167. t. 75. f. 6.*

Fungus multiplex obtuse conicus, colore griseo mu-
rino. *Vaill. paris. 71. t. 12. f. 12.*

Fungus parvus, pediculo oblongo firmo lento, pileolo
in medio fastigiato, striis exterius apparentibus.
R. syn. 8.

β Amanita petiolo procero gracilissimo, pileolo conico
striato murino inferne albo. *Hall. hist. 2391.*

Amanita parva verna utrinque striata fusca, pileo ob-
tuse coniformi. *Dill. giss. 184. R. syn. 8.*

Anglis, bell Agaric.

Habitat in pratis, pascuis et sylvis. VIII.—X.

separatus. 32. AGARICUS stipitatus, pileo lævi livido, lamellis
nigricantibus separatis, stipite bulboso volvato. *Sp.
pl. 1643. Fl. succ. 1220.*

Agaricus candicans. *Schæff. fung. t. 217?*

Anglis, separated Agaric.

Habitat in pratis et pascuis. IX. X.

33. AGA-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

- canicus.* 37. AGARICUS stipitatus, pileo campanulato acuto striato livido nitido, lamellis albis, stipite fistuloso cinereo.
 Agaricus galericulatus. Scop. carn. 1564. Schæff. fung. t. 52.
 Amanita pileolo conico spadiceo, lamellis albis. Hall. hist. 2393.
 Fungus multiplex campaniformis, colore castaneo. Vaill. paris. 73. t. 12. f. 3, 4.
 Anglis, cone Agaric.
 Habitat in lignis putrescentibus. IX. X.
- fuliginosus.* 38. AGARICUS stipitatus, pileo hemisphærico cinereo striato, lamellis griseis pulverulentis, stipite fistuloso griseo. Schæff. fung. t. 6. f. 2?
 Bulla campaniformis utrinque sulcata. Battar. fung. 56. t. 27. f. E?
 Fungus minor tenerrimus farina respersus, pileolo superne cinereo, lamellis subtus tenuissimis creberrimis nigris. R. syn. 9?
 Fungi minores plurimi simul nascentes turbinati, exterius cinerei aut subfulvi, striis nigricantibus. R. syn. 10.
 Anglis, sooty Agaric.
 Habitat in pratis et pascuis ad sepes. IX. X.
- candidus.* 39. AGARICUS stipitatus, pileo hemisphærico, lamellis stipiteque albis.
 Aminata albus, pileo hemisphærico excarni, oris striatis. Hall. hist. 2348.
 Fungus plures juxta se nascentes, parvi turbinati candidi ubivis coloris. R. syn. 10. t. 1. f. 2. a a.
 Fungus minimus albus, pileo hemisphærico in arborum caudicibus. Mich. gen. 146. t. 74. f. 7.
 Anglis, white Agaric.
 Habitat in sylvis, ramis arborum dejectis putridis. VIII—X.
- fragilis.* 40. AGARICUS stipitatus, pileo convexo viscido pelucido, lamellis que luteis, stipite nudo. Sp. pl. 1643. Schæff. fung. t. 230.
- Amanita

Amanita flavo-cinnamomeus viscidus striatus papillatus. *Hall. hist.* 2425.

Fungus pediculo croceo splendoris particeps. *Vaill. paris.* 69. t. 11. f. 16, 17, 18.

Fungus pratensis minor externe viscidus, striis subtus fulvis seu croceis. *R. syn.* 8?

Anglis, brittle Agaric.

Habitat in sylvis et sepibus inter dejecta folia et muscos. VIII—XI.

41. AGARICUS stipitatus, pileo plano plicato pellucido lamellisque lividis brevissimis, stipite longissimo griseo nudo.

Fungus capillaceus, capitulo rorido, nigro punctulo in summitate notato. *Pluk. ph.* t. 116. f. 7. *R. syn.* 13.

Anglis, tender Agaric.

Habitat in pratis et pascuis, in stercore equino. IX—X.

42. AGARICUS stipitatus, pileo plicato membranaceo, umbelliformis lamellis basi latioribus. *Sp. pl.* 1643. *Fl. suec.* 1192. *serus.* *Scop. carn.* 1571.

Amanita albus tener pileolo campanulato striato. *Hall. hist.* 2346.

Fungus minimus totus albus, pileo hemisphærico, utrinque striato lamellis rarioribus. *Mich. gen.* 166. t. 80. f. 11.

Fungus parvus candidissimus lamellatus, pediculo longo gracili. *R. syn.* 9.

Anglis, umbrella Agaric.

Habitat in sylvis inter folia dejecta semiputrida. IX—XII.

43. AGARICUS stipitatus, pileo planiusculo margine plicato plicato luteo-rufo, lamellis pallidioribus, stipite longo rufo.

Agaricus alliatus. *Schæff. fung.* t. 99?

Anglis, plicated Agaric.

Habitat in sylvis et sepibus. IX. X.

44. AGARICUS stipitatus albus, pileo plicato membranaceo, *androsaceus.*

branaceo, stipite nigro. *Sp. pl.* 1644. *Fl. suec.* 1193. *Schæff. fung.* 239.

Amanita petiolo nigro, pileolo albo striato ex carni.
Hall. hist. 2351.

Fungus caule nigro capillari androsaces capitulo. *Bacc. musc.* 143. t. 104.

Fungus pileo cardicante lamellis paucis, pediculo fulco splendente. *Vaill. paris.* 69. t. 11. f. 21, 22, 23.

Fungus minimus adianti aurei capitulis. *R. syn.* 9.
Anglis, black-stalked Agaric.

Habitat in foliis et ramulis dejectis putrescentibus. VIII
—XII.

clavus. 45. AGARICUS stipitatus, pileo luteo convexo striato, lamellis stipiteque albis. *Sp. pl.* 1644. *Fl. suec.* 1212.

Agaricus acicula. *Schæff. fung.* t. 222.

Amanita minimus, oris adtractis flavus lamellis albis.
Hall. hist. 2370.

Fungus minimus aurantius mammillaris. *Vaill. paris.* 76. t. 11. f. 19, 20.

Fungus minimus, capitulo conico, rufescens, lamellis subtus paucis. *R. syn.* 9?

Anglis, dwarf Agaric.

Habitat in sylvis et sepibus. IX—III.

pilosus. 46. AGARICUS stipitatus, pileo convexo albo piloso, pilis rubris, stipite longo.

Anglis, hairy Agaric.

Habitat in sylvis, in foliis putrescentibus, præcipue illicis.
IX—XI.

Desc. Stipes fere uncialis, filiformis, albidus aut rufus, aut rubescens, ad basin subpilosus. Pileus magnitudine seminis cannabis, convexus, fere hemisphæricus, albus, pilosus; pilis erectis, rigidiusculis rubris; Lamellæ albide.

** Parasitici, subacaules, dimidiati.

quercinus. 47. AGARICUS acaulis, lamellis labyrinthiformibus.
Sp. pl. 1644. *Schæff. fung.* 57.

Agaricus

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Amanati sessilis, lamellis pulverulentis bifidis. *Hall. hist.* 2333.

Fungus arboreus albus durus, lamellis instar lapidis hæmatis. *R. syn.* 25.

Anglis, alder Agaric.

Habitat in truncis arborum. 4. I—XII.

lateralis. 51. AGARICUS substipitatus, pileo albido lævi, lamellis subsimplicibus pallidis, stipite laterali.

Agaricus semipetiatus. *Schæff. fung. t.* 208?

Agaricus parvus lamellatus pectunculi forma elegans.

Dill. giss. 192. *R. syn.* 25.

Agaricum squamosum rufescens, pediculo donatum, subtus lamellis densis. *Mich. gen.* 123. t. 65. f.

71

Amanita semipetiatus albidus, subtus terreus. *Hall. hist.* 2337.

Fungus parvus, lamellatus, pectunculi forma, albo adnescens. *Vaill. paris.* 70. t. 10. f. 1, 2.

Anglis, lateral Agaric.

Habitat in sylvis et sepibus, in ramis arborum dejectis putrescentibus. VIII—XII.

BOLETUS. 1210.

Fungus horizontalis; subtus porosus.

* *Parasitici acaules.*

suberosus. 1. BOLETUS acaulis pulvinatus albus lævis, poris acutis difformibus. *Sp. pl.* 1645.

Boletus acaulis coriaceo-convexus villosus albus, poris difformibus rotundis flexuosisque. *Fl. suec.* 1253.

Anglis, cork Boletus.

Habitat in truncis arborum, in fraxinis in comitatu Westmorlandico passim. 4. I—XII.

fomentarius. 2. BOLETUS acaulis pulvinatus inæqualis obtusus, poris teretibus æqualibus glaucis. *Sp. pl.* 1645.

Fl. suec. 1252.

Agaricus porosus ignarius Fagi, superne candicans, inferne fuscus. *Dill. gif.* 193. *R. syn.* 24.

β Agaricus

- β Agaricus porosus igniarius Carpini. *Dill. gifs.* 193.
R. syn. 24.
Anglis, tinder Agaric.
 Habitat in truncis arborum. 4. I—XII.
3. BOLETUS acaulis pulvinatus lævis, poris tenuissi- igniarius:
 mis. *Sp. pl.* 1646. *Fl. suec.* 1256.
 Boletus unguulatus, fulvus et variegatus. *Schæff.*
fung. t. 137, 138. 262, 263.
 Polyporus convexo-planus durissimus cinereus, inferne
 albidus. *Hall. hist.* 2288.
 Agaricus pedis equini facie. *Tourn.* 562. *R. syn.* 22.
Battar. 71. t. 37. f. E.
 Agaricum igniarium, &c. *Mich. gen.* 118. i. 61.
 Fungus in caudicibus nascens, unguis equini figura.
B. pin. 372.
 Fungus durus igniarius. *Park.* 1323.
- β Boletus annulatus. *Schæff. fung. t.* 106.
 Polyporus sessilis convexo-planus, anulis versicolori-
 bus, poris albis tenuissimis. *Hall. hist.* 2283.
Anglis, touchwood Boletus.
 Habitat in arborum truncis. 4. I—XII.
4. BOLETUS acaulis coriaceus convexus lobatus fla- coriaceus
 vus lævis, poris tenuissimis.
 Boletus caudicinus. *Scop. carn.* 1596. *Schæff. fung.*
t. 131, 132.
 Agaricus speciosus. *Battar. fung.* 68. t. 34. f. B.
Anglis, coriaceous Boletus.
 Habitat in truncis arborum. 4. I—XII.
5. BOLETUS acaulis convexus sublobatus rufus lævis hepaticus:
 poris teretibus flavis. *Schæff. fung. t.* 116—120.
 Polyporus sessilis sanguineus crassus mollis obscure lo-
 batus. *Hall. hist.* 2315.
 Agaricus porosus rubens carnosus, hepatis facie. *Dill.*
gifs. 192. *R. syn.* 23.
 Agaricum esculentum, castanæ adnascens, latissimum,
 hepatis facie, superne ex rubro ferrugineum, interne
 sanguineum, inferne ochroleucum. *Mich. gen.* 117.
t. 60.
- S f
- Anglis,*

Anglis, liver Boletus.

Habitat in truncis arborum. 4. I—XII.

versicolor. 6. BOLETUS acaulis, fasciis discoloribus, poris albis.

Sp. pl. 1645. *Schæff. fung. t.* 136.

Boletus variegatus, mesentericus, atrofuscus et multicolor. *Schæff. fung. t.* 263, 267, 268, et 269.

Polyporus sessilis cespitosus planus hirsutus, anulis versicoloribus. *Hall hist.* 2282.

Agaricus varii coloris squamosus. *Tourn.* 562. *R. syn* 24.

Agaricus squamis iridiformibus. *Battar. fung.* 69. *t.* 35. *f. A.*

Anglis, striped Boletus.

Habitat in truncis arborum. 4. I—XII.

villosus. 7. BOLETUS acaulis convexus villosus fuscus, poris angularibus flavescens.

Agaricus villosus et porosus substantiæ coriacæ. *Dill. giss.* 193. *R. syn.* 24.

Anglis, villous Boletus.

Habitat in truncis arborum. 4, I—XII.

squamosus. 8. BOLETUS subacaulis planiusculus lacerato-squamosus flavescens, poris tetragonis.

Boletus juglandis. *Schæff. fung. t.* 101, 102.

Polyporus sessilis, convexo-planus, poris amplissimis albidis. *Hall. hist.* 2278.

Fungus angulosus pediculo exiguo. *B. pin.* 370. *R. syn.* 11.

Auricula flammea Malchi. *Sterb. fung.* 105. *t.* 13.

Anglis, scaly Boletus.

Habitat in truncis arborum præcipue Fraxinis et Betulis; in boreali parte Angliæ frequens. V—X.

albus. 9. BOLETUS acaulis pulvinatus lævis, poris oblongis.

Fl. angl. 496.

Polyporus sessilis convexo-planus farinosus. *Hall. hist.* 2287?

Agaricus officinali similis. *Dill. giss.* 192. *R. syn.* 23.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Fungus porosus tenuis coriaceis. *Buxb. cent. 5. t. 15.*
f. 1.
Anglis, woolly Boletus.
Habitat in sylvis. Prope Esher in comitatu Surriensi.
 IX. X.

- perennis.* 13. BOLETUS stipitatus perennis, pileo utrinque planiusculo. *Sp. pl. 1646. Fl. suec. 1245.*
 Boletus coriaceus. *Scop. carn. 1588. Schæff. fung. t. 125.*
 Polyporus petiolatus infundibuliforma cespitosus intus nigricans, extus albus. *Hall. hist. 2298.*
 Polyporus lignosus et cespitosus infundibulum imitans, superne nigricans, inferne cum pediculo albus, areolis carbonariis innascens. *Mich. gen. 131. t. 70. f. 6.*
 Fungus lignosus fasciatus. *Vaill. paris. 61. t. 12. f. 7.*
Anglis, perennial Boletus.
Habitat in sylvis, ad ligna putrida. 4. I—XII.

H Y D N U M. 1211.

Fungus horizontalis, subtus echinatus.

- imbricat-* 1. HYDNUM stipitatum, pileo convexo imbricatum.
tum. *Sp. pl. 1647. Schæff. fung. t. 140. 273. Fl. dan. 176.*
 Echinus petiolatus albicans, superne squamosus. *Hall. hist. 2324.*
 Erinaceus esculentus albus crassus. *Mich. gen. 132. t. 72. f. 2.*
 Fungus pœne albus, prona parte erinaceus. *B. hist. III. 828. R. syn. 11.*
Anglis, imbricated Hydnum.
Habitat in sylvis; prope Maidston in Cantio. IX. X.

- repan-* 2. HYDNUM stipitatum, pileo convexo lævi flexuoso.
dum. *Sp. pl. 1647. Fl. suec. 1258. Fl. dan. 310. Schæff. fung. t. 141, 318.*

Echinus

- Echinus petiolatus subfuscus, petiolo lævi. *Hall. hist.* 2325.
 Erinaceus esculentus pallide luteus. *Mich. gen.* 132. t. 72. f. 3.
Anglis, smooth Hydnum.
 Habitat in sylvis. IX—XI.
3. HYDNUM stipitatum, pileo dimidiato. *Sp. pl.* 1648. *auriscalpium.*
Schæff. fung. t. 143.
 Echinus petiolo gracili, laterali, pileolo plano obscuro. *Hall. hist.* 2321.
 Erinaceus parvus hirsutus ex fusco fulvus, pileolo femiorbiculari, pediculo tenuiore. *Mich. gen.* 132. t. 78. f. 8.
 Fungus erinaceus parvus, in conis abietis nascens. *Buxb. cent. 1. t.* 57. f. 1.
Anglis, fir Hydnum.
 Habitat in ramis foliis et conicis dejectis putrescentibus Pini et Abietis, passim. IX. X.

P H A L L U S. 1212.

Fungus supra reticulatus, subtus lævis.

1. PHALLUS pileo ovato celluloso, stipite nudo rugoso. *Sp. pl.* 1648. *Fl. dan.* 53. *Schæff. fung. t.* 199, 298—300. *esculentus.*
 Boletus capite tereti reticulato. *Hall. hist.* 2247.
 Boletus esculentus rugosus. *Mich. gen.* 203. t. 85. f. 1, 2. albicans, quasi fuligine infectus. *Tourn.* 561. t. 329. f. *A. Battar. fung.* 24. t. 2. f. *F.*
 Fungus porosus rugosus albicans, quasi fuligine infectus. *B. pin.* 370.
 Fungus favaginosus. *Park.* 1317. *R. fyn.* 11.
Anglis, esculent Morel.
 Habitat in sylvis et sepibus.
2. PHALLUS volvatus stipitatus, pileo albo celluloso impudicus.
 apice pervio.

Phallus volvatus stipitatus, pileo celluloso. *Sp. pl.* 1648. *Fl. dan.* 175. *Schæff. fung. t.* 196—198.

Phallus volva exceptus, capituli apice patulo. *Gled. fung.* 55. *Hall. hist.* 2248.

Phallus vulgaris totus albus, volva rotunda, pileolo cellulato, ac summa parte umbilicato pervio orniato. *Mich. gen.* 201. t. 83.

Boletus phalloides. *Tourn.* 562. *Battar. fung.* 23. t. 2. f. A.

Fungus phalloides. *B. hist. III.* 843. *R. syn.* 12. batavicus pileo laciniato et veluti crispo. *R. hist. III.* 25. *Sterb. fung.* 277. t. 30. f. A—D.

Fungus foetidus penis imaginem referens. *B. pin.* 374. *Anglis*, stinking Morel.

Habitat in sylvis et ad sepes. VII—IX.

caninus. 3. PHALLUS volvatus stipitatus, pileo rubro celluloso acuto, apice clauso.

Phallus caninus. *Schæff. fung. t.* 330?

Phallus exilis marattæ. *Battar. fung.* 76. desc? *Anglis*, red-headed Morel.

Habitat in sylvis et numerosis, prope Shrewsbury. *D. Ehret.* VII—IX.

CLATHRUS. 1213.

Fungus subrotundus subcancellatus, villo repletus.

* *Stipitati.*

denudatus. 1. CLATHRUS stipitatus, capitulo oblongo volvato. *Sp. pl.* 1649.

Clathroides purpureum, pediculo donatum. *Mich. gen.* 214. t. 94. f. 1.

Trichia gregaria purpurea petiolata, capitulo sphaerico. *Hall. hist.* 2164. t. 48. f. 6.

Anglis, purple Clathrus.

Habitat in lignis putridis. VI—X.

nudus. 2. CLATHRUS stipitatus, capitulo oblongo axi longitudinali adnato. *Sp. pl.* 1649. *Fl. dan.* 216.

Clathroi-

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Mucor perennis pallidus, pileo fulvo. *Sp. pl.* 1655.
Trichia petiolata rufa, capitulo sphaerico. *Hall. hist.*
 2163. t. 48. f. 4.
Anglis, brown Clathrus.
Habitat in lignis putridis. V—X.

virescens, 7. **CLATHRUS** stipitatus viridis, capitulo globofo fa-
 rinaceo.

Mucor perennis viridis, foliis furfuraceis, stipite fili-
 formi, capitulo globofo. *Sp. pl.* 1655.
Anglis, greenish Clathrus.
Habitat in terra umbrosa ad latera viarum et fossarum
et in lignis putridis. V—X.

** *Acaulis*,

turbina- 8. **CLATHRUS** acaulis turbinatus, villo flavescente-
tus. Clathroides purpureus, pediculo carens. *Mich. gen.*
 214. t. 94. f. 2.
Trichia gregaria sessilis piriformis, flava. *Hall. hist.*
 2168. t. 48. f. 7.
Anglis, turbinated Clathrus.
Habitat in lignis putridis. VIII—IV.

HELVELLA. 1214.

Fungus horizontalis, utrinque lævis.

• *Stipitatae.*

Mitra. 1. **HELVELLA** pileo deflexo adnato lobato difformi.
Sp. pl. 1649.
Elvela mitra. *Schæf. fung.* 159—162, 154, 282,
 320, et 322.
Boletus capitulo explanato laciniato. *Hall. hist.* 2246.
Boletus leucophæus. *Battar. fung.* 25. t. 3. f. B.
Boletus pileolo spiralibus plicis contorto. *Battar.*
fung. 24. t. 2. f. G.
Fungoides fungiforme crispum laciniatum et compli-
catum, pediculo crasso striato rimoso et fistuloso.
Mich. gen. 204. t. 86. f. 7, 8, 9.

Fungus

Fungus terrestris pediculo striato et cavernoso, capitulo plicatili, subtus plano. *R. syn.* 8.

Anglis, curled or mitred Helvella.

Habitat in sylvis. VIII—X.

Obs. variat stipite lævi, et striato; Pileo flavescente, fusco et livido.

2. HELVELLA pileo plano livido, stipite lævi. *planus.*

Anglis, plain Helvella.

Habitat in sylvis. VIII—X.

* *Acaulis.*

3. HELVELLA acaulis. *acaulis.*

Agaricus acaulis utrinque planiusculus. *Fl. lapp.* 517.

suec. 1243.

Agaricus villosus tenuis, inferne lævis. *Dill. giss.*

193. *R. syn.* 21?

Anglis, sessile Helvella.

Habitat in lignis et arborum truncis putrescentibus. 4.

I—XII.

PEZIZA. 1215.

Fungus campanulatus.

1. PEZIZA campanulata lentifera villosa interne lævis. *lentifera.*

Peziza campanulata lentifera. *Sp. pl.* 1649. *Fl. dan.*

469.

Peziza sericea. *Schæff. fung. t.* 180.

Cyathus sericeus intus lævis. *Hall. hist.* 2215.

Cyathoides cyathiforme, cinereum et veluti sericeum.

Mich. gen. 222. t. 102. f. 1.

Fungoides infundibuliforma, semine foetum. *Tourn.*

560. *Vaill. paris.* 56. t. 11. f. 677.

Fungus pyxoides seminifer. *Fl. pruss.* 90. t. 16. f.

sup.

Fungus minimus ligneis tabulis areolarum hortorum

adnascens. *B. pin.* 514.

Fungi calyciformes seminiferi. *Mentz. pug. t.* 6.

R. syn. 20.

Anglis,

Anglis, seeding Peziza.

Habitat in sylvis, agris, arborum et lignis putridis. I—XII.

Ariata. 2. PEZIZA campanulata lentifera villosa, interne striata. *Fl. dan.* 105.

Peziza hirsuta. Schæff. fung. t. 178.

Cyathus hirsutus, intus striatus. Hall. hist 2214.

Cyathoides cyathiforme, obscurum, externe hirsutum interne plumbeum glabrum et striatum. Mich. gen. 222. t. 182. f. 2.

Fungoides infundibuliforma, semine foetum, interne striatum, externe hirsutum. Vaill. paris. 57. t. 11. f. 4, 5.

Fungus seminifer, externe hirsutus, interne striatus. R. syn. 21.

Fungus non vescus XLI. Fl. pruss. 90. t. 16. f. inf.

Anglis, striated Peziza.

Habitat in sylvis, agris et lignis putridis. I—XII.

laevis. 3. PEZIZA campanulata lentifera ventricosa glabra.

Peziza crucibuliformis. Scop. carn. 1630. Schæff. fung. t. 179. 181.

Cyathoides luteum crucibuliforme. Mich. gen. 222. t. 102. f. 3.

Fungus seminifer minor, fere hemisphaericus. R. syn. 20. t. 1. f. 2. b, c.

Anglis, smooth Peziza,

Habitat in lignis et putridis. IX—V.

punctata. 4. PEZIZA turbinata truncata, disco punctata. *Sp. pl.* 1650. *Fl. succ.* 1275. *dan.* 288.

Fungus minimus infundibuliformis, superne nigris punctis notatus. R. syn. 12.

Fungus minimus lignosus disco punctato. Bocc. musc. II. 149. t. 107.

Anglis, dotted Peziza.

Habitat in stercore equino. V—X.

cornucopioides. 5. PEZIZA infundibuliformis, disco patente sinuato punctato. *Sp. pl.* 1650. *Fl. dan.* 268. *Schæff. fung. t. 165, 166.*

Peziza

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Elvela hemisphærica excavata et sessilis, intus candida glabra extus pilosa et subobscura. *Gled. fung.* 46.

Fungoides scutellatum, interne album, externe obscurum et subhirsutum. *Mich. gen.* 206. t. 86. f. 4.

Anglis, hairy Peziza.

Habitat in sylvis et sepibus humidiusculis et lignis putridis humidis. IX. X.

stipitata. 9. PEZIZA stipitata, hemisphærica, externe fusca hispida, stipite cylindrico.

Peziza hispida. *Schæff. fung.* t. 167.

Peziza terrea petiolo longo gracili, acetabulo extus subaspero, plano aut concavo. *Hall. hist.* 2227.

Fungoides fuscum, pediculo longiore donatum. *Buxb. comm. Aët. petrop.* IV. 282. t. 29. f. 3.

Anglis, stalked Peziza.

Habitat in sylvis; prope Guildford. IX. X.

coccinea. 10. PEZIZA subgloboso-campanulata interne coccinea, margine integro. *Schæff. fung.* t. 148.

Peziza cava albida, intus coccinea. *Hall. hist.* 2222.

Peziza miniata major. *Dill. giss.* 194. *R. syn.* 17.

Fungus membranaceus seu coriaceus acetabuli modo concavus, colore intus coccineo seu cremesino facturo. *R. syn.* 18.

Fungoides coccineum, acetabuli forma. *Mich. gen.* 207.

Fungi perniciosi gen. xxiv. spec. 2. *Clus. hist.* 287.

Anglis, scarlet Peziza.

Habitat in lignis putridis in sylvis et sepibus. VIII—III.

cyathoides. 11. PEZIZA cyathiformis, margine obtusa erecto. *Sp. pl.* 1651.

Peziza lutea parva, marginibus lævibus. *R. syn.* 18, t. 24. f. 4.

Anglis, smooth Peziza.

Habitat in lignis putridis. VIII—IV.

fulva. 12. PEZIZA plana fulva, margine convexo glabro. Peziza

Peziza lenticulatis parva miniata. *R. syn.* 18. t. 24.
f. 2.

Peziza sessilis aurantia plana annulata. *Hall. hist.*
2224.

Fungus fimosus bracteolatus croceus. *Pet. Vaill.*
paris. t. 13. f. 14.

Anglis, orange Peziza.

Habitat in stercore equino et bubulo, et solo glareoso.
VIII—X.

Obs. variat colore flavo et fusco.

13. PEZIZA concava atra.

Peziza planissima, ex glauco nigricans. *Hall. hist. atra.*
2243.

Anglis, black Peziza.

Habitat in fimo vaccino. IX—V.

14. PEZIZA plana, margine convexo piloso. *Sp. pl. scutellata.*
1651.

Peziza sessilis miniata plana, ora ciliata. *Hall. hist.*
2245.

Peziza lutea parva, marginibus pilosis. *R. syn.* 18.
t. 24. f. 3.

Elvela ciliata. *Scop. carn.* 1620. *Schæff. fung.* 284.

Anglis, ciliated Peziza.

*Habitat in stercore equino et bubulo et lignis putridis in
sylvis et sepibus.* VIII—X.

15. PEZIZA turbinata cochleata. *Sp. pl.* 1657. *cochleata.*

Elvelæ. *Schæff. fung.* t. 150, 155, 156, 274, 280.

*Fungoides auriculam judæ referens, intus rufescens
extus candicans et quasi farinosum.* *Vaill. paris.*
57. t. 11. f. 8.

Anglis, wreathed Peziza.

Habitat in sylvis. IX. X.

CLAVARIA. 1216.

Fungus, lævis, oblongus.

* *Indivisa.*

- pistillaris.* 1. CLAVARIA clavæformis simplicissima. *Sp. pl.* 1651.
 Clavaria alba pistilli forma. *Vaill. paris.* 39. t. 7.
 f. 5.
 Clavaria lutea minima. *Mich. gen.* 208. t. 87. f. 5.
 Clavaria cespitosa media lutea. *Mich. gen.* 209. t. 87.
 f. 10, 11.
 Clavaria ligula. *Schæff. fung.* t. 171.
 Clavaria alba longa petiolata. *Hall. hist.* 2203.
 Fungoides clavatum minus. *R. syn.* 14.
 β Clavaria minima nivea. *Mich. gen.* 208. t. 87. f. 6, et
 9. *Hall. hist.* 2207.
 Fungus clavatus minimus. *R. syn.* 14.
 γ Clavaria vermiculata, simplex lumbriciformis ochro-
 leuca. *Fl. carn.* 1623.
 Clavaria vermiculata fistulosa et non fistulosa candi-
 da. *Mich. gen.* 209. t. 87. f. 12, 13.
 Fungoides clavatum incurvum in acutum mucronem
 productum. *R. syn.* 14.
 δ Clavaria major lutea. *Mich. gen.* 208. t. 87. f. 1,
 2, 3.
 Clavaria pistillaris. *Schæff. fung.* t. 169.
Anglis, club Clavaria.
Habitat in sylvis et ericetis. IX—XI.
- militaris.* 2. CLAVARIA clavata integerrima, capite squamoso.
Sp. pl. 1652.
 Clavaria militaris crocea. *Vaill. paris.* 39. t. 7. f. 4.
 Fungus fontanus purpureus elegans. *R. syn.* 12.
Anglis, red Clavaria.
Habitat in sylvis humidis et paludibus. IX. X.
- ophioglossoides.* 3. CLAVARIA clavata integerrima compressa obtusa.
Sp. pl. 1652. *Schæff. fung.* t. 327.
 Clavaria indivisa nigra compressa. *Hall. hist.* 2206.
 Clavaria ophioglossoides. *R. syn.* 14.

Clavaria

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

Habitat in lignis putrescentibus. I—XII.

coralloi-
des. 6. CLAVARIA ramis confertis ramosissimis inæqualibus. *Sp. pl.* 1652. *Schæff. fung. t.* 175, 176, 177, 285—288.

Clavaria caule crassissimo, ramis innumerabilibus cespitosis, breviter dentatis. *Hall. hist.* 2201.

Corallo-fungus flavus. *Vaill. paris.* 41. t. 8. f. 4.

Coralloides flava albida. *Tourn.* 564. t. 332. f. B.

Fungus ramosus flavus et albidus. *B. hist. III.* 837. *R. syn.* 16.

Fungoides ramosum maximum, brassicæ caulifloræ facie et magnitudine. *R. syn.* 16.

Barba caprina. *Sterb. fung.* 69. t. 11. f. A—D.

Anglis, coral Clavaria.

Habitat in ericetis, nemorosis et pascuis. VIII—X.

fastigiata. 7. CLAVARIA ramis confertis ramosissimis fastigiatis obtusis luteis. *Sp. pl.* 1652.

Clavaria albida, purpurea, flammea. *Schæff. fung. t.* 170, 172, 174.

Clavaria caule plano sulcato ramis ferratis. *Hall. hist.* 2200.

Coralloides minor cristata. *Buxb. cent. IV.* 39. t. 66. f. 1.

Fungoides fungiforme luteum foetidum et minus ramosum. *R. syn.* 479. t. 24. f. 5.

Anglis, yellow Clavaria.

Habitat in sylvis. VIII—X.

muscoides. 8. CLAVARIA ramis ramosis acuminatis inæqualibus luteis. *Sp. pl.* 1652. *Schæff. fung. t.* 173.

Clavaria caule ramoso, ramis incurvis acutis. *Hall. hist.* 2199.

Fungus parvus ramosus luteus. *R. syn.* 16. t. 24. f. 7.

Fungus ramosus minimus, coloris aurantii. *R. syn.* 16.

Anglis, dwarf Clavaria.

Habitat in ericetis et sylvis. IX. X.

LYCOPERDON.

LYCOPERDON. 1712.

Fungus subrotundus, Seminibus farinaceis repletus.

* *Solida, subterranea absque radice.*

1. LYCOPERDON globosum solidum muricatum, *tuberis*
radice destitutum. *Sp. pl.* 1653.

Lycoperdon subterraneum conglomeratum, cortice
nigro exasperato. *Hall. hist.* 2177.

Tubera. *Math.* 544. *B. pin.* 376. *hist. III.* 849.
Tourn. t. 333. *R. syn.* 28. *terræ. Ger. em.* 1583.
edulia. Park. 1319.

Tuber brumale, pulpa obscura odorata. *Mich. gen.*
221. *t.* 164.

Anglis, esculent Puff-ball or Truffles.

Habitat *sub terra; in sylvis et pascuis elatis.* Downs
dictis in comitatibus Wiltoniensi, Hantonensi, Can-
tiano, &c. passim. IX—V.

2. LYCOPERDON globosum solidiusculum lacerum *cervinum*,
centro farinifero, radice instructum. *Sp. pl.* 1653.

Sphæria subrotunda, tuberculis minimis, medulla
atra pulverulenta. *Hall. hist.* 2191.

Tubera cervina. *Lob. ic.* 276. *B. pin.* 376.

Tubera perniciosâ terrestria s. cervina. *Sterb. fung.*
215. *t.* 32. *f. B. B. B. R. syn.* 28.

Lycoperdaltrum tuberosum arrhizon fulvum cortice
duriose crasso et granulato, medulla ex albo pur-
purascente, semine nigro crassiore. *Mich. gen.* 320.
t. 99. *f.* 4.

Anglis, subterraneous Puff-ball.

Habitat *in sylvis et ad sepes.* In Canewood, Hamp-
stead. *R. syn. in Devon. IX. X.*

** *Solidum parasiticum.*

3. LYCOPERDON subrotundum solidum tubercula-*fraxinea*
tum nigrum. *um.*

Sphæria maxima convexa nigerrima. *Hall. hist.* 2192.

Lichen-agaricus crustaceus crassus, bovillum quasi

renem repræsentans, niger et veluti deustus. *Mich. gen.* 104. t. 54. f. 1.
 Fungus fraxineus niger durus orbiculatus. *R. syn.* 16.
hist. 109.
Anglis, ash Puff-ball.
 Habitat in putrescentibus fraxini truncis. I—XII.

*** Pulverulenta, radicata supra terram.

Bovista. 4. LYCOPERDON subrotundum, lacerato-dehiscens.
Sp. pl. 1656.

Lycoperdon faccatum cinereum, polline obscure viridi. *Hall. hist.* 2172.

vulgare. α Lycoperdon vulgare. *Tourn.* 563.

Lycoperdon areolatum. *Schæff. fung. t.* 190.

Crepitus Lupi s. Fungus ovatus. *Park.* 1323. *R. syn.* 26.

Fungus rotundus orbicularis. *B. pin.* 374.

Fungus tertius s. orbicularis. *Ger. em.* 1582.

β Fungus orbicularis per totum pulvere repletus. *R. syn.* 26.

γ Lycoperdon furfuraceum. *Schæff. fung. t.* 294.

Fungus Lupi crepitus dictus in summitate solum pulverulentus. *R. syn.* 26.

Fungus niger calycis figuram referens. *B. pin.* 375.

δ Lycoperdon maximum. *Schæff. fung. t.* 191.

Fungus maximus rotundus pulverulentus dictus germanis Bofst. *B. hist. III.* 848. *R. syn.* 26.

Fungus cucurbitiformis magnitudine capitis humani et ponderosus candiusque instar nivis. *Merr. pin.*

echinatum.

Lycoperdon echinatum. *Schæf. fung. t.* 186.

Lycoperdon esculentum obscurum, echinis quadrangularibus et quadripartitis insignitis. *Mich. gen.* 218. t. 97. f. 5.

Lycoperdon gemmatum. *Schæff. fung. t.* 189.

Lycoperdon niveum sphaericum, superficie in areolas adamantis instar dissecta ac distributa. *Tourn.* 564. *Mich. gen.* 218.

excipuliforme.

Lycoperdon excipuliforme. *Schæff. fung.* 187, 292, 295.

Lycoperdon

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

*fornica-
tum.*

7. LYCOPERDON volva quadrifida fornicata, capitulo glabro: ore obtuso ciliato.

Lycoperdon coronatum. Schæff. fung. t. 183.

Geaster volvæ radiis et operculo elevatis. *D. Watson. Act. angl. No. 474. p. 234.*

Geasteroides phragmites fuscum e volva erumpens quatuor radiis elevatum, curiosum. *Battar. fung. 74. t. 39. f. 1—5.*

Fungus pulverulentus turriculam fornicatam referens. *D. Rand. Blacks. spec. bot. 24. t. 2.*

Anglis, turret Puff-ball.

Habitat in pratis et pascuis; at Bucklebury, ten miles from Reading, Berkshire, and about Wickham near Bromley in Kent. *Blacks. spec. bot. prope Doncaster. D. Tofield. IX. X.*

Carpobolus.

8. LYCOPERDON volva multifida, fructu globoso ex seminibus combinatis. *Sp. pl. 1654.*

Lycoperdon corticibus revolutis stellatis, globo projectili. Hall. hist. 2175.

Caropobolus. Mich. gen. 221. t. 101. f. 1, 2.

Anglis, projectile Puff-ball.

Habitat in lignis putridis, in sylvis et sepibus. VIII—X.

*** Parasitica, in farinam fatescentia.

piliformis.

9. LYCOPERDON globosum scabrum, ore perforato. *Syst. nat. 726.*

Sphæria solitaria, scutellis nudis atris obtuse conicis. Hall. hist. 2180.

Lichen crustaceus tartareus, arboribus adnascens, niger et veluti deustus, receptaculis florum rotundis per siccitatem nonnihil adpressis et umbilicatis. *Mich. gen. 104. t. 54. f. 5.*

Anglis, pea Puff-ball.

Habitat in arborum truncis putrescentibus. IX—XI.

nigrum.

10. LYCOPERDON parasiticum sessile subrotundum nigrum farina concolori.

Sphæria convexa carne nigra. Hall. hist. 2184.

Lichenoides

Lichenoides tuberculosum compressum nigrum lignis putridis adnascens. *R. syn.* 51. *Dill. musc.* 126. *t.* 18. *f.* 7.

Lichen-agaricus bullatus parvus, ex obscure nigricans, sub cute arborum exsiccatarum erumpens. *Mich. gen.* 105. *t.* 54. *ord.* 11. *f.* 2.

Anglis, black Puff-ball.

Habitat in lignis et truncis arborum putridis. IX—IV.

11. LYCOPERDON parasiticum sessile subrotundum varioloscortice exteriori fulvo deciduo, farina atra compacta. *Syst. nat.* III. 234.

Lycogala globosum grani pisi magnitudine, æris recti colore. *Mich. gen.* 216. *t.* 95. *f.* 2.

Lycogala sessile fulvum. *Hall. hist.* 2142.

Anglis, orange Puff-ball.

Habitat in lignis et truncis arborum putridis. IX—IV.

Obs. affinis Tremellæ purpuræ, sed major et caro in senescente atra est.

12. LYCOPERDON parasiticum globosum fuscum farina concolori.

Lycogala griseum majus. *Mich. gen.* 216. *t.* 95. *f.* 1?

Anglis, brown Puff-ball.

Habitat in arborum truncis putridis. IX—V.

13. LYCOPERDON cortice farinaque purpurea. *Sp. epidendrum* *pl.* 1654.

Lycoperdon sanguineum sphaericum. *Buxb. cent.* V. *t.* 29. *f.* 2.

Lycoperdon leve miniatum sphaericum. *Hall. hist.* 2173.

Anglis, purple Puff-ball.

Habitat in lignis putridis. IX—V.

14. LYCOPERDON aggregatum parasiticum ore multifido lacero; pulvere fulvo. *Sp. pl.* 1655. *epiphyl- lum.*

Mucor fragiformis. *Schæff. fung.* *t.* 193.

Anglis, yellow Puff-ball.

Habitat in lignis et foliis semiputridis et hypnis. VIII
—V.

MUCOR. 1218.

Fungus vesiculis stipitatis.

* *Perennis.*

- Embolus.* 1. MUCOR seta nigra villo fusco. *Sp. pl.* 1655.
Embolus nigerrimus, villo albo adpersus. *Hall. hist.*
2137.
Anglis, globe Mould.
Habitat in lignis putridis. I—XII.

** *Fugaces.*

- Mucedo.* 2. MUCOR stipitatus, capsula globosa. *Sp. pl.* 1655.
Fl. dan. 467. f. 4.
Mucor petiolatus cinereus. *Hall. hist.* 2147.
Mucor vulgaris, capitulo lucido per maturitatem nigro, pediculo griseo. *Mich. gen.* 215. t. 95. f. 1.
Mucedo capitulo diaphano, deinde viridi et postremo nigro. *Malp. veget.* t. 25. f. 108.
Fungus bombycinus murini coloris e fimo felino tenuissimis capillis: *Pluk. alm.* 164. *R. syn.* 13.
Anglis, common Mould.
Habitat in variis putridis: pane, plantis, &c. I—XII.
- leprosus.* 3. MUCOR fetaceus, seminibus radicalibus. *Sp. pl.* 1656.
Aspergillus cespitosus densissimus, initio niveus deinde aureus, seminibus ovatis. *Mich. gen.* 213. t. 91. f. 5.
Anglis, leprous Mould.
Habitat in cavernulis et cellulis fornicatis. IX—IV.
- glaucus.* 4. MUCOR stipitatus, capitulo subrotundo aggregato.
Sp. pl. 1656.
Aspergillus capitatus, capitulo glauco, seminibus rotundis. *Mich. gen.* 212. t. 91. f. 1.
Anglis, glaucous Mould.
Habitat in Pomis, Melonibus aliisque corruptis. I—XII.

5. MUCOR

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

A P P E N D I X.

- pauciflora.* 2—3. **SCIRPUS** culmo striato nudo, spica terminali pauciflora valvis calycinis longiore. *Lightf. Scot. 1078. Fl. dan. 167?*
Scirpus caule tereti, spica nuda pauciflora, glumis calycinis longiore. Hall. hist. 1335.
Scirpus minimus, spica brevior squamosa spadicea. Sch. agrost. 364. t. 7, f. 19.
Anglis, chocolate-head Club-rush.
Habitat in montosis. Upon Malghyrdy in Breadalbane.
Mr. Stuart. Lightf. Scot. 4. VIII.
Obs. varietas scirpi cæspitosi videtur.
- glabra.* 4—5. **FESTUCA** panicula ramosa erecta coarctata, spiculis subulatis trifloris aristatis glabris. *Lightf. Scot. 1085.*
Anglis, smooth Fescue-grass.
Habitat in pascuis. Found at Ardbigland in Galloway.
Lightf. Scot. 4. VII.
Obs. affinis festucae cambricæ, an eadem.
- tuberosum.* 2. **SYMPHYTUM** foliis semidecurrentibus, summis oppositis. *Syst. nat. 147. Sp. pl. 195. Jacq. obs. III. 12. t. 63.*
Symphytum tuberosum. Ger. em. 806. majus. Clus. hist. 166.
Anglis, tuberous-rooted Camfrey.
Habitat in Scotia. Mr. Yalden found it sparingly opposite the new well at the water of Leith, but more plentifully

plentifully in Dr. Robertson's walks at North Marchiston. 4. VII.

IMPERATORIA. 359.

Fruetus subrotundus, compressus, medio gibbus, margine cinctus. *Petala* inflexo-emarginata.

125. 1. IMPERATORIA. *B. pin.* 156. *Ger. em.* 1001. *ostruthi-Sp. pl.* 371. *Lightf. scot.* 169. major. *Hist. Ox. III. um.* 278. *f.* 9. *t.* 4. *f.* 1.

Anglis, Masterwort.

Habitat in pascuis et ruderatis. Upon the banks of the Clyde in several places, particularly about Ardencaple, and in the isle of Bute, near Mount Stewart; on the banks of the Tweed, about half a mile above the old castle of Drummelzier. *Dr. Burges's. Lightf. scot.* 4. VI.

7. ALLIUM scapo nudo adæquante folia teretia sub-schœno-
lato-filiformia. *Syst. nat.* 140. *Sp. pl.* 432. prafum.

Allium radice multiplici, foliis radicalibus teretibus, scapi longitudine, floribus umbellatis cylindricis. *Hall. hist.* 1226.

Allium montanum capite rotundo. *Hist. Ox. III.* 386. *f.* 4. *t.* 14. *f.* 4.

Porrum sectivum juncifolium. *B. pin.* 72.

Schœnoprasum. *Dod. pempt.* 639. *Ger. em.* 176.

Anglis, chived or cives Garlick.

Habitat in pratis et pascuis; by Fast-castle on the borders of Berwickshire. *Dr. Parsons. Fl. scot. in comitatu Westmorlandico passim.* 4. VI.

9—10. JUNCUS folio subulato, gluma biflora terminali. *Sp. pl.* 467. *Amœn. acad. II.* 266 *t.* 3. *f.* 3. *Lightf. scot.* 1200?

Anglis, two-flowered Rush.

Habitat in alpibus Scoticis. Upon the top of Malghyrdy, and upon Ben-Teskerny, in Breadalbane. *Mr. Stuart. Lightf. scot.* 4. VIII.

- spicatus.* 14. JUNCUS foliis planis, spica racemosa nutante.
Sp. pl. 469. *Fl. lapp.* 125. t. 10. f. 4. *Lightf. scot.* 187.
Anglis, spiked Rush.
 Habitat in alpidibus Scoticis. Upon Ben-na-scree, on the north side of Loch-Urn, on the western coast of Inverness-shire, &c. 4. VII.

DORONICUM. 959.

Recept. nudum. Pappus simplex. Calycis squamæ duplicis ordinis æquales, disco longiores. Sem. radii nuda pappoque destituta.

- pardati-
anches.* *370. 1. DORONICUM foliis cordatis obtusis denticulatis, radicalibus petiolatis caulinis amplexicaulibus. *Sp. pl.* 1247. *Mill. ic.* 128. *Lightf. scot.* 485.
 Doronicum foliis cordatis obtusis, imis longis petiolatis, superioribus amplexicaulibus. *Hall. hist.* 88.
 Doronicum maximum, foliis caulem amplexantibus. *B. pin.* 185. *Carn. epit.* 823.
 Doronicum maximum. *Ger. em.* 726.
Anglis, great Leopard's or Wolf's Bane.
 Habitat in montosis. In several places in the Lowlands, as about Hoddam-castle in Annandale, &c. 4. V. VI.

- incurva.* 7—8. CAREX spica conica, e spiculis congestis androgynis sessilibus composita, involucro nullo, culmo incurvo. *Lightf. scot.* 544. t. 24.
Anglis, curved Carex.
 Habitat in arenosis. It was discovered in deep loose sea-sand, at the mouth of the water of Naver, and near Shelberry, in Dunrossness, in Shetland. *Dr. Hope. Lightf. Scot.* 4. VIII.

- tomentosa.* 17—18. CAREX spicis femineis subpedunculatis, capulis subglobosis tomentosis. *Syst. nat.* 617. *Lightf. scot.* 553.
 Carex spicis sessilibus approximatis teretibus folio infidentibus. *Hall. hist.* 1383.
 Cyperoides sylvaticum angustifolium, spicis parvis tenuibus spadiceo viridibus. *Sch. agr.* 425. t. 10. f. 11.

Anglis,

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

Habitat in rupibus. Upon the rocks on the Highland mountains frequent. Lightf. scot. IX, X.
 Obs. an varietas Jungermanniaë julaceæ.

cæruleo-nigricans. 5—6. LICHEN leprofus cærulescens, tuberculis nigricantibus. Lightf. scot. 805.

Lichenoides granosum subglaucum, tuberculis planis nigricantibus. Dill. musc. 544. s. 82. f. 2.

Anglis, bluish Lichen.

Habitat in rupibus. Upon the Highland rocks, but not common. Lightf. scot. 4. I—XII.

cornuta. 8—9. ULVA subrigida cornuta terrestris. Lightf. scot. 972.

Tremella terrestris cornuta. Dill. musc. 52. t. 10. f. 13.

Anglis, horned Laver.

Habitat in arenosis humidis; near Leith. Mr. Yalden, Lightf. scot.

Obs. an varietas Jungermanniaë pinguidis.

montana. 7—8. ULVA plana coriacea terrestris sanguinea. Lightf. scot. 973.

Anglis, mountain Laver.

Habitat in montanis. Amongst the grass and moss on the sides of the mountains in the isle of Skye, and in Rosshire, on the side of a mountain called Cail-veg, between Loch-broom and Lead-beg, on the western coast. Lightf. scot. VIII.

laciniata. 3—4. ULVA frondibus planis purpureis, apice dilatatis laciniatis undulatis. Lightf. scot. 974.

Anglis, lacinated Laver.

Habitat in littoribus marinis; upon the coast of Jura. Lightf. scot. VIII.

Obs. an varietas Ulvæ umbilicalis.

endivia-folius. 24—25. FUCUS fronde membranacea laciniata, laciniis dilatatis undulatis, marginibus crispis tuberculato-punctatis. Lightf. scot. 948. t. 32.

Anglis, endive-leaved Fucus.

Habitat

Habitat in littoribus. In the Frith of Forth, and on the coast of Jona. VIII.

19—20. CONFERVA filamentis ramosis geniculatis, verticillatis ramulis verticillatis furcatis incurvis. *Lightf. scot. lata.* 984.

Anglis, verticillate Conferva.

Habitat in rupibus et saxis submarinis. In the Frith of Forth, and many other places not unfrequent. *Lightf. scot.*

27—26. CONFERVA filamentis geniculatis ramosissimi-*diaphana.* mis, ramis dichotomis apice forcipatis, septis ruberrimis, articulis diaphanis. *Lightf. scot. 998.*

Anglis, diaphanous Conferva.

Habitat in rupibus et saxis submarinis in littoribus Scotiæ. *Lightf. scot.*

22—23. AGARICUS stipitatus, pileo conico viscido *aurantio,* lamellis luteis, stipite nudo. *Lightf. scot. tius.* 1025.

Agaricus conicus. Schæff fung. t. 2.

Agaricus palustris fanguineo colore. Dill. giss. 180?

Amanita glutinosus flavus pileo umbonato. Hall. hist. 2420.

Anglis, orange Agaric.

Habitat in pascuis siccis, in Scotia. *Lightf. scot. IX.* X.

10—11. LYCOPERDON *nigrum,* aggregatum parasiticum sphaericum, ore integro. *Lightf. scot. 1069. tum.* t. 31.

Anglis, clustered Puff-ball.

Habitat in arborum truncis putridis. *Lightf. scot. XI.* —V.

PINGUICULA *villosa.* *Lightf. scot. 77. t. 6.* In Strathwardale in the Isle of Skye, in a bog just by Mr. M'Kennon's, of Coryattacha, and on the side of a mountain there called Benna-Caillick. *Lightf. scot.*

CIRCÆA alpina. *At the foot of the mountains about Loch-Broom in Ross-shire. Under the rocks on the east-side of St. Mary's Loch, at the head of Yarrow-water, in the county of Peebles. Dr. Burgess. Lightf. scot. 77. 1115.*

SCHŒNUS rufus.

Schœnus compressus varietas. Lightf. scot. t. 24.

AGROSTIS pumilla. *Vide figura in fronte tom. II: Lightf. scot.*

FESTUCA rubra. *In the Isle of Canay plentifully. Lightf. scot. 102.*

CORNUS herbacea. *On the sides of the Highland mountains not unfrequent. Lightf. scot. 119.*

ATROPA Belladonna. *Jacq. austr. IV. 5. t. 309.*

CAUCALIS arvensis.

Caucalis helvetica. Jacq. hort. III. 12. t. 16.

SISON Amomum. *Jacq. hort. III. 13. t. 17.*

SISON verticillatum. *Lightf. scot. 1095. t. 35.*

OENANTHE pimpinelloides. *Jacq. austr. IV. 49. t. 395:*

CARUM Carui. *Jacq. austr. IV. 49. t. 393.*

PIMPINELLA major. *Jacq. austr. IV. 50. t. 396.*

PIMPINELLA saxifraga. *Jacq. austr. IV. 50. t. 395.*

VIBURNUM Lantana. *Jacq. austr. IV. 21. t. 341.*

GALANTHUS nivalis. *Jacq. austr. IV. 7. t. 313.*

ORNITHOGALUM umbellatum. *Jacq. austr. IV. 22. t. 340.*

JUNCUS trifidus. *Lightf. scot. 183. t. 9. Upon the summits of the Highland mountains to the south of Little-Lock-Broom, in Ross-shire, and on Ben-nascree above Arnesdale, on the side of Loch-Urn, in Inverness-shire. Upon Ben-nevis in Lochabar. Lightf. scot. 184, 1120.*

JUNCUS triglumis. *Upon Craigneulicht near the head of Loch-Tay. Upon Ben-Lomond, and in mossy ground at the top of Carndearg, one of the lower heads of Ben-Squilert, in Glen-creran, in Upper-Lorn. Upon Malghyrdy and Malhan-tarmonach, on the east side of*

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

- RESEDA *lutea*. *Jacq. austr. IV. 28. t. 353.*
 EUPHORBIA *platyphyllos*. *Jacq. austr. IV. 40. t. 376.*
 POTENTILLA *verna*. *Upon the top of Craig-chailleach, near Finlarig in Breadalbane. Fl. scot. 270.*
 CLEMATIS *Vitalba*. *Jacq. austr. IV. 4. t. 308.*
 THALICTRUM *alpinum*. *Lightf. scot. 286. t. 13.*
 STACHYS *germanica*. *Jacq. austr. IV. 10. t. 319.*
 BARTSIA *viscosa*. *Lightf. scot. 321. t. 14.*
 HESPERIS *inodora*. *Jacq. austr. IV. 25. t. 347.*
- CARDAMINE *petræa*. *Lightf. scot. 347. t. 15. Upon Craig-chailleach in Breadalbane; upon Baikeval in the island of Rum, and upon Ben-na-caillich in the isle of Skye. Lightf. scot.*
 BRASSICA *monensis*.
Silybrium monense. Lightf. scot. 353. t. 15.
 SISYMBRIUM *Iris*. *Jacq. austr. IV. 11. t. 322.*
 GERANIUM *pyrenaicum*. *In mountainous pastures and waste places near Edinburgh. Dr. Parsons. Lightf. scot. 367.*
- OROBUS *sylvaticus*. *Lightf. scot. 390. t. 16.*
 HEDYSARUM *onobrychis*. *Jacq. austr. IV. 27. t. 352.*
 ASTRAGALUS *arenarius*. *Upon the hills in the King's Park at Edinburgh; upon the sands of Musselburgh, upon the hill of Moncreis, near Perth; and in dry ground near York-Cascade at the seat of the Duke of Argyle, at Blair. Light. scot. 400.*
- ASTRAGALUS *uralensis*. *Light. scot. 401. t. 17. Upon Carn-Dearg one of the lower heads of Ben-Squilt, a high mountain in Glen-cieran, in Upper Lorn. Mr. Stuart.*
- HIERACIUM *alpinum*. *Light. scot. 43. t. 18. Upon Ben mor, and Craig-vore just above Loch-Laraigan-Lochain in Breadalbane; upon Cory-yaal and near the top of Bedannam-bian, the highest mountain in Glenco. On the high mountains to the south of Little-Loch-Broom, in Ross-shire, also on Ben-nagron, two miles above M'Kinnon's Castle in Strath, in the isle of Skye. Lightf. scot.*
- SERRATULA

SERRATULA alpina. Lightf. scot. 448. t. 19. *Among the rocks on the eastern side of Malghyrdy, and near the top of Ben-Cruipen in Breadalbane; on the north side of Stuic-an-lochain in Glen-lyon; upon Ben-achalader and Ben-dotha, in Glenorchy; and upon Malmore in Glenco, in Upper-Lorn; upon a Brae a mile from Talisker in the isle of Skye.* Lightf. scot.

GNAPHALIUM supinum.

Gnaphalium alpinum. Lightf. scot. 470. t. 20.

ORCHIS militaris β.

Orchis fusca. Jacq. austr. IV. 4. t. 307.

SATYRIUM hircinum. Jacq. austr. IV. 35. t. 367.

SATYRIUM albidum. *In the dry mountainous pastures not very uncommon.* Lightf. scot. 520.

SATYRIUM repens. Jacq. austr. 36. t. 369. Lightf. scot. 550. t. 22. *In a birch wood called Ca-bue or Yellow-hill, facing the house of Mr. McKenzie of Dundonald, two miles from the head of Little-Loch-Broom on the western coast of Ross-shire.* Lightf. scot.

OPHRYS corallorrhiza. Lightf. scot. 523. t. 23. *In a moist hanging-wood, on the south side near the head of Little-Loch-Broom, Ross-shire.* Lightf. scot.

CAREX patula.

Carex pauciflora, spica simplici androgyna; floribus femineis subternis remotiusculis patentibus, masculo unico terminali. Lightf. scot. 543. t. 6. *In a boggy soil about half-way up the mountains of Goatfield, in the isle of Arran, near the place where peat is dug, in the ascent from Brodwick-castle.* Lightf. scot.

CAREX atrata. *Upon the Highland mountains frequent, as upon Benteskerney, Mal-ghyrdy, Mal-nan-tarmonach, &c. in Breadalbane.* Lightf. scot. 555.

CAREX inflata. *In bogs and marshes as about Finlarig, in Breadalbane.* Mr. Stuart. Lightf. scot. 567.

BETULA nana. Lightf. scot. 575. t. 25. *This grows in great abundance in Corry-yeallin, on the west side of a hill directly north from Miggerny, in Glenlyon, in Breadalbane; on some of the moors of Loch-glass, in Ross-shire. Lightf.*

SALIX myrsinitis. *Upon the Highland mountains, as upon Ben-Achulader, in Glenurchy, Mal-ghyrdy, in Breadalbane; and on Craig-vore, a high projecting rock on the west side of Loch-laraig-an-lochain, a small lake between Loch-tay and Glen-lyon. Mr. Stuart. Lightf. scot. 599.*

SALIX reticulata. *It grows upon several of the Highland mountains, in a talky soil, as upon Cleg-chaillech, Mal-ghyrdy, and Mal-grea, mountains in Breadalbane, &c. Lightf. scot. 601.*

ACROSTICUM septentrionale. *This grows out of the clefts of the rocks about Arthur's Seat, at Edinburgh. Lightf. scot. 656.*

ISOETES lacustris. *In some small locks near the foot of the Paps of Jura, and in others above Great-Breeze-Hill, near Talisker, in the isle of Skye. Lightf. scot. 684.*

LYCOPODIUM annotinum. *In rough stony places on the sides of the Highland mountains. Light. scot. 689.*

MNIUM trichomanes.

Jungermannia frondibus simpliciter pinnatis, foliolis ovatis planis integerrimis, caule apice femineo. Lightf. scot. 799.

MNIUM fissum.

Jungermannia frondibus simpliciter pinnatis, foliolis bifidis, caule apice femineo. Lightf. scot. 770.

MNIUM Jungermannia.

Jungermannia purpurea pinnis subtus auriculatis, foliolis subrotundis convexo-concavis amplexicaulis. Lightf. scot. 778.

BRYUM

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

LICHEN *islandicus* γ.

Lichen hispidus. fruticulosus solidus ramosissimus, ramis divaricatis subcompressis glabris spinulosis, angulis obtusis. *Lightf. scot.* 883.

LICHEN *laciniatus*.

Lichen glomuliferis, coriaceus repens laciniatus glaber, laciniarum axillis orbicularibus, scutellis fulvis et glomulis atroviridibus. *Lightf. scot.* 853.

LICHEN *herbaceus*.

Lichen læte-virens, coriaceus repens lobatus obtusus glaber læte-virens, subtus-avenius, scutellis fulvis. *Lightf. scot.* 852.

LICHEN *verrucosus*.

Lichen scrobiculatus, coriaceus repens lobatus obtusus, superne scrobiculatus verrucosus, inferne villosus tumoribus albidis. *Lightf. scot.* 850.

LICHEN *croceus*. Upon a rock near the top of Bentekerney, in Breadalbane. Mr. Stuart. *Lightf. scot.* 856.

LICHEN *polyrrhizos*.

Lichen torrefactus, umbilicatus pullus, supra rugosus subtus-reticulato-fibrosus, verrucis nigris crispis. *Lightf. scot.* 862.

LICHEN *velleus*.

Lichen polyrrhizus. Lin. secundum. *Lightf. in Fl. scot.* 864.

LICHEN *proboscideus*.

Lichen crinitus, umbilicatus margine crinitus, peltis nigris concinnatis subpedunculatis. *Lightf. scot.* 860. On Goatfield in the isle of Arran, Ben-Nevis in Lochaber, and on the mountains of Breadalbane. *Lightf. scot.*

LICHEN *radiatus*.

Lichen corniculatus, fruticulosus solidus ramosus, ramis declinatis compressis dichotomis, scutellis plano-convexis fungiformibus. *Lightf. scot.* 885. On Creg-chaillech, in Breadalbane, on Ben-Lomond, on the mountains in Ross-shire, and in the isle of Skye. *Lightf. scot.*

LICHEN

LICHEN *scaber.*

Lichen *exilis*, filamentosus ramosissimus caespitosus, filamentis capillaceis implexis opacis scabris. *Lightf. Scot.* 895.

TREMELLA *auricula.*

Peziza concava rugosa auriformis. *Lightf. Scot.* 1055.

TREMELLA *turbinata.*

Peziza *polymorpha*, turbinata, primum cava, deindè plana seu convexa, externe rugosa. *Lightf. Scot.* 1055.

ULVA *laetuca* γ .

Ulva *crispa*, tenera crispa terrestris. *Lightf. Scot.* 972.

ULVA *dichotoma.*

Ulva fronde plana dichotoma reticulata viridis. *Lightf. Scot.* 975. t. 34.

ULVA *purpurascens.*

Fucus *verticillatus*, frondibus tubulosis subarticulatis ramosis setaceo-ligulatis. *Lightf. Scot.* 962. t. 31.
Upon the rocks of the little isles of Jura. Lightf. Scot.

ULVA *articulata* α et β .

α Fucus fronde articulata ramosissima, articulis ovato-cylindricis tubulosis, ramis oppositis et verticillatis. *Lightf. Scot.* 959.

β Fucus frondibus caespitosis repentibus ramosis articulatis, articulis planis. *Lightf. Scot.* 961.

FUCUS *ovalis.*

Fucus *vermicularis*, caule filiformi ramoso, ramis dichotomis, foliis oblongis teretibus petiolatis. *Lightf. Scot.* 958.

FUCUS *esculentus.* *Lightf. Scot.* 938. t. 28.

FUCUS *bulbosus.*

Fucus *polychides*, fronde palmata, foliolis ensiformibus, radice cava, stirpe plana marginibus plicatis. *Lightf. Scot.* 936.

FUCUS *palmatus*. *Lightf. scot.* 933. t. 27.

FUCUS *crispus*.

Fucus prolifer, frondibus membranaceis dichotomis catenato-proliferis, apicibus dilatatis bifidis. *Lightf. scot.* 949. t. 30.

FUCUS *herbaceus*.

Fucus ligulatus, fronde membranacea lineari bipinnato-ligulata, ligulis ensiformibus ciliatis. *Light. scot.* 964. t. 29. *In the Frith of Forth about Newhaven.* *Lightf. scot.*

FUCUS *pumilus*.

Fucus pygmaeus fronde cartilaginea compressa apice dilatata palmata, fructificationibus terminalibus subrotundis apice perforatis. *Lightf. scot.* 964. t. 32.

FUCUS *corneus*.

Fucus nereideus, fronde cartilaginea filiformi-compressa ramosa, ramulis setaceo-pinnatis. *Lightf. scot.* 656. *In the Frith of Forth.*

FUCUS *verrucosus*.

Fucus flagelliformis, fronde filiformi tereti ramosa, ramis alternis subdistichis longissimis uniformibus. *Lightf. scot.* 928.

FUCUS *purpurascens*.

Fucus tuberculatus, fronde filiformi tereti ramosissima; ramis alternis ramulis confertis tuberculatis, tuberculis subrotundis distantibus innatis. *Lightf. scot.* 927.

CONFERVA *violacea*.

Conferva confragosa, filamentis mucosis simplicibus æqualibus violaceis. *Lightf. scot.* 976. *Upon the rocks in the water-falls on the mountain of Goatfield in the isle of Arran.* *Lightf. scot.*

CONFERVA *ciliata*.

Conferva ciliata, filamentis geniculatis dichotomis, apice forcipatis, articulis verticillatim ciliatis. *Lightf. scot.* 998.

CONFERVA

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

BOLETUS villosus.

Boletus spongiosus, acaulis villosus fuscus, poris albidis ciliatis angularibus. *Lightf. scot.* 1033.

BOLETUS hepaticus.

Boletus hepaticus, acaulis lævis carnosus sanguineus mollis, obscure lobatus. *Lightf. scot.* 1034.

PEZIZA striata.

Peziza striata campanulata lentifera extus hirsuta. intus striata. *Lightf. scot.* 1049.

PEZIZA lævis.

Peziza crucibuliformis lentifera glabra sublutea. *Lightf. scot.* 1049.

CLAVARIA pistillaris. γ . δ .

γ *Clavaria vermiculata*, lumbriciformis ochroleuca. *Lightf. scot.* 1057.

δ *Clavaria herculeana*, clavæformis simplicissima crassissima luteo-fusca rugosa, vertice depresso. *Lightf. scot.* 1056.

LYCOPERDON *nigrum*.

Sphæria tuberculosa, atra convexa carne nigra. *Lightf. scot.* 1113.

F I N I S,

I N D E X

N O M I N U M · L A T I N O R U M .

Generica nomina Romana Characterè traduntur :
 Cursivis Literis Synonyma.
 Primus numerus Genus designat ; cæteri Synonyma.

| A. | | Page | | Page |
|-------------------|------|----------------------------|------------------------|---------------------------------|
| <i>Abies</i> | | 424 | <i>Ægilops</i> | 441, 52 |
| <i>Abrotanum</i> | | 358 | <i>Ægopodium</i> | 129 |
| <i>Abſinthium</i> | 358, | 359 | <i>Æthusa</i> | 123 |
| <i>Acanthium</i> | | 354 | <i>Agaricus</i> | 609, 626, 633, 639, 653, 663 |
| <i>Acer</i> | | 445 | <i>Agaricum</i> | 563, 625 |
| <i>Acetosa</i> | | 156 | <i>Agrifolium</i> | 446 |
| <i>Acetofella</i> | | 156 | <i>Agrimonia</i> | 206 |
| <i>Achillea</i> | 374, | 375 | <i>Agrostemma</i> | 198 |
| <i>Acinos</i> | 261, | 263 | <i>Agrostis</i> | 30—32, 654 |
| <i>Aconitum</i> | | 228 | <i>Aira</i> | 33, 37 |
| <i>Acorus</i> | | 147, 14 | <i>Ajuga</i> | 248 |
| <i>Acrosticum</i> | 450, | 451, 457, 658 | <i>Aizoon</i> | 198, 197 |
| <i>Actæa</i> | | 228 | <i>Alcea</i> | 207, 208 |
| <i>Adiantum</i> | 460, | 461, 450, 454, 456, 459 | <i>Alchemilla</i> | 71, 72, 178 |
| <i>Adonis</i> | | 239 | <i>Alcionum</i> | 570 |
| <i>Adoxa</i> | | 172 | <i>Aleærolophis</i> | 267 |
| | | | <i>Aleærolophoides</i> | 610 |
| | | | <i>Aleærolophus</i> | |

666 INDEX NOMINUM LATINORUM.

| | Page | | Page |
|---------------------|--|-------------------------|------------------------------|
| <i>Aleurolophus</i> | 268 | <i>Aphanes</i> | 72 |
| <i>Alga</i> | 395, 396, 568—570, 577, 580, 581, 589, 593, 602 | <i>Apium</i> | 128, 129, 115 |
| <i>Alisma</i> | 158, 159 | <i>Aponogiton</i> | 397 |
| <i>Alliaria</i> | 287 | <i>Aquifolium</i> | 446 |
| <i>Allium</i> | 138—140, 649 | <i>Aquilegia</i> | 235 |
| <i>Alni</i> | 214 | <i>Arabis</i> | 292, 293 |
| <i>Alnus</i> | 98, 416 | <i>Aracus</i> | 319, 321 |
| <i>Aloe</i> | 236 | <i>Arbutus</i> | 177, 655 |
| <i>Alopecurus</i> | 27, 28 | <i>Arctium</i> | 348 |
| <i>Alphanthemis</i> | 160 | <i>Arctia</i> | 84 |
| <i>Alfinastrum</i> | 173 | <i>Arenaria</i> | 191—193, 655 |
| <i>Alfne</i> | 131, 6, 7, 60, 63, 72, 73, 138, 147, 160, 186, 190—195, 200—203, 255, 256, 276 | <i>Argemone</i> | 230, 231 |
| <i>Alfinella</i> | 73 | <i>Argentina</i> | 223 |
| <i>Althæa</i> | 306 | <i>Argyrocome</i> | 360 |
| <i>Alyssum</i> | 277 | <i>Aria</i> | 214 |
| <i>Amanita</i> | 610—624 | <i>Aristolochia</i> | 394 |
| <i>Amara dulcis</i> | 93 | <i>Armeria</i> | 184, 185 |
| <i>Amaranthus</i> | 418 | <i>Armerius</i> | 185, 189 |
| <i>Ambrosia</i> | 284 | <i>Artemisia</i> | 357—359 |
| <i>Amomum</i> | 119 | <i>Arum</i> | 395 |
| <i>Ampeloprasum</i> | 139 | <i>Arundo</i> | 53, 54, 23 |
| <i>Anacampteros</i> | 195, 434 | <i>Atarum</i> | 205 |
| <i>Anagallis</i> | 87, 5, 63, 86, 94 | <i>Ascyron</i> | 334 |
| <i>Anblatum</i> | 266 | <i>Asparagus</i> | 145 |
| <i>Anchusa</i> | 80, 79 | <i>Aspergillus</i> | 646, 647 |
| <i>Andromeda</i> | 176, 166 | <i>Asperugo</i> | 82 |
| <i>Androsæmum</i> | 332—334 | <i>Asperula</i> | 66, 67 |
| <i>Anemone</i> | 236, 237 | <i>Asphodelus</i> | 143, 145 |
| <i>Anethum</i> | 126 | <i>Asplenium</i> | 452—454 |
| <i>Angelica</i> | 118, 129 | <i>Aster</i> | 368, 360, 363 |
| <i>Anonis</i> | 312, 313 | <i>Astragalus</i> | 322, 323, 314, 650 |
| <i>Anthemis</i> | 373, 374 | <i>Athamanta</i> | 115, 116 |
| <i>Antnericum</i> | 144, 145, 151 | <i>Athanasia</i> | 357 |
| <i>Anthoceros</i> | 522, 523 | <i>Atioplex</i> | 442—444, 105—107 |
| <i>Antboxanthon</i> | 155 | <i>Atropa</i> | 93, 654 |
| <i>Anthoxanthum</i> | 11 | <i>Avena</i> | 51—53, 30, 31, 34—36, 440 |
| <i>Anthyllis</i> | 313, 60, 138 | <i>Auricula flammea</i> | 626 |
| <i>Antirrhinum</i> | 271, 274 | <i>———— muris</i> | 192 |
| <i>Aparine</i> | 68—70, 82 | <i>Azalea</i> | 88 |
| <i>Aphaca</i> | 315 | | |
| | | B. | |
| | | <i>Bacharis</i> | 363 |
| | | <i>Bella-</i> | |

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

668 INDEX NOMINUM LATINORUM.

| | Page | | Page |
|---------------------------|-------------------|-----------------------|--------------------------|
| <i>Cauda equina</i> | 2 | <i>Cineraria</i> | 369, 370 |
| — <i>muris</i> | 136 | <i>Circaea</i> | 10, 11, 654 |
| <i>Centaureo</i> | 357—377 | <i>Cirsium</i> | 349, 350, 352—354 |
| <i>Centaarium</i> | 102, 103, 168 | <i>Cistus</i> | 232—234 |
| <i>Centunculus</i> | 63 | <i>Clabroidastrum</i> | 631 |
| <i>Cerastium</i> | 200—202, 655 | <i>Clabroides</i> | 630, 632 |
| <i>Cerasus</i> | 213 | <i>Clathrus</i> | 630—632 |
| <i>Ceratophyllum</i> | 419 | <i>Clavaria</i> | 638—640, 654 |
| <i>Cerefolium</i> | 124, 125 | <i>Clematis</i> | 238, 656, 91 |
| <i>Cerintbe</i> | 81 | <i>Clinopodium</i> | 261, 263, 267 |
| <i>Ceryomices</i> | 627 | <i>Clymenum</i> | 320, 332 |
| <i>Ceterach</i> | 452 | <i>Cnicus</i> | 355 |
| <i>Chætophylla</i> | 72 | <i>Cochlearia</i> | 283, 284 |
| <i>Chætophyllum</i> | 124, 125 | <i>Colchicum</i> | 157 |
| <i>Chamaeclema</i> | 254 | <i>Comarum</i> | 227 |
| <i>Chamaecistus</i> | 88, 181, 233, 234 | <i>Conferva</i> | 591—605, 572, 653, 662 |
| <i>Chamaedrys</i> | 6, 226, 237, 248 | | 663 |
| <i>Chamaefilix</i> | 453 | <i>Conium</i> | 114, 115 |
| <i>Chamaegenista</i> | 311 | <i>Confiligo</i> | 245 |
| <i>Chamaelea</i> | 167 | <i>Consolida</i> | 235, 248, 249 |
| <i>Chamaelinum</i> | 134 | <i>Convallaria</i> | 146, 147 |
| <i>Chamaemelum</i> | 372, 374 | <i>Convolvulus</i> | 88, 89, 171 |
| <i>Chamaemorus</i> | 221 | <i>Conyza</i> | 362, 363, 356, 366, 369, |
| <i>Chamaenerion</i> | 161—163 | | 370 |
| <i>Chamaeorchis</i> | 390 | <i>Corallina</i> | 566, 569, 598, 599 |
| <i>Chamaepericlymenum</i> | 71 | <i>Coralloides</i> | 286, 527, 328, 539, |
| <i>Chamaepitys</i> | 248 | | 551—359, 631 |
| <i>Chamaepuce</i> | 464 | <i>Corallorbiza</i> | 388 |
| <i>Chamaeitea</i> | 429 | <i>Coreopsis</i> | 356 |
| <i>Chamaerubus</i> | 221 | <i>Coriandrum</i> | 123 |
| <i>Chamaeschænus</i> | 20 | <i>Cornu cervinum</i> | 65 |
| <i>Chara</i> | 297, 998 | <i>Cornus</i> | 70, 71, 654 |
| <i>Chelidonium</i> | 228, 229, 244 | <i>Coronopus</i> | 64, 65, 284 |
| <i>Cheiranthus</i> | 287, 288 | <i>Corrigeola</i> | 100 |
| <i>Chenopodium</i> | 104—108 | <i>Corylus</i> | 423 |
| <i>Cherleria</i> | 193 | <i>Cotyledon</i> | 194, 110 |
| <i>Chlora</i> | 168 | <i>Cracca</i> | 318, 321 |
| <i>Chondrilla</i> | 338 | <i>Crambe</i> | 299, 298 |
| <i>Chrysiophoriana</i> | 228 | <i>Crassula</i> | 73, 195 |
| <i>Chrysanthemum</i> | 370 | <i>Cratægus</i> | 214 |
| <i>Chrytosplenium</i> | 178, 179, 268 | <i>Cratægonon</i> | 268, 270 |
| <i>Cichoreum</i> | 348, 341, 342 | <i>Crepis</i> | 339, 341, 342 |
| <i>Cicuta</i> | 122, 123, 115 | <i>Crepitus lupi</i> | 642 |
| <i>Cicutaria</i> | 123—125 | <i>Critimum</i> | |

INDEX NOMINUM LATINORUM. 669

| | Page |
|---------------------|---------------------------------|
| <i>Crithmum</i> | 117, 112, 369 |
| <i>Crista galli</i> | 268 |
| <i>Crocus</i> | 13 |
| <i>Cruciata</i> | 441 |
| <i>Cucubalus</i> | 185—187 |
| <i>Cuminum</i> | 126 |
| <i>Cuscuta</i> | 104 |
| <i>Cyanus</i> | 375, 376 |
| <i>Cyatocides</i> | 633, 634 |
| <i>Cyatbus</i> | 633, 634 |
| <i>Cymbalaria</i> | 271 |
| <i>Cynoscrambe</i> | 435 |
| <i>Cynoglossum</i> | 80, 81 |
| <i>Cynoforbis</i> | 383 |
| <i>Cynofurus</i> | 59 |
| <i>Cyperoides</i> | 400, 401, 407, 409, 410, 412 |
| <i>Cyperus</i> | 17, 14—17, 21, 403 |
| <i>Cyperella</i> | 16 |
| <i>Cypripedium</i> | 392 |

D.

| | |
|--------------------|--------------------|
| <i>Dactylis</i> | 43 |
| <i>Damaconium</i> | 158, 159 |
| <i>Daphne</i> | 167 |
| <i>Darea</i> | 461 |
| <i>Datura</i> | 92 |
| <i>Daucus</i> | 114, 115, 120, 128 |
| <i>Delphinium</i> | 235 |
| <i>Dens leonis</i> | 239 |
| <i>Dentaria</i> | 285, 286, 388 |
| <i>Dianthus</i> | 183—185 |
| <i>Diapensia</i> | 111 |
| <i>Digitalis</i> | 275, 276 |
| <i>Digitaria</i> | 25 |
| <i>Dipsacus</i> | —61 |
| <i>Doronicum</i> | —650 |
| <i>Dortmanna</i> | 377 |
| <i>Draba</i> | 278, 279, 378 |
| <i>Dracunculus</i> | 375 |
| <i>Drosera</i> | 155 |
| <i>Dryas</i> | 226 |

| | Page |
|-------------------|----------|
| <i>Dryopteris</i> | 457, 460 |

É.

| | |
|------------------------|--------------------------------------|
| <i>Ebulus</i> | 130 |
| <i>Echium</i> | 82, 83, 78, 81 |
| <i>Echinophora</i> | 112, 113 |
| <i>Echinus</i> | 565, 628, 629 |
| <i>Elatine</i> | 173, 271, 272 |
| <i>Elleborine</i> | 392 |
| <i>Eleoselinum</i> | 129 |
| <i>Eliechrysum</i> | 359 |
| <i>Elymus</i> | 56—58 |
| <i>Embolus</i> | 646 |
| <i>Elvela</i> | 563, 632, 636, 637 |
| <i>Empetrum</i> | 430, 431 |
| <i>Epilobium</i> | 161—163, 655 |
| <i>Epipactis</i> | 386—390, 393 |
| <i>Equisetum</i> | 447—449, 2, 173, 398 |
| <i>Erica</i> | 165—167, 177, 431 |
| <i>Erigeron</i> | 363, 365 |
| <i>Eriocaulon</i> | 415 |
| <i>Eriophorum</i> | 21, 22 |
| <i>Eruca</i> | 137, 278, 286, 290, 291, 296, 298 |
| <i>Ervum</i> | 320, 321, 315 |
| <i>Eryngium</i> | 109, 110 |
| <i>Eryfimo similis</i> | —292 |
| <i>Erysimum</i> | 286, 297, 172 |
| <i>Esula</i> | 208 |
| <i>Eudnymus</i> | 98 |
| <i>Eupatorium</i> | 356, 206, 355 |
| <i>Euphorbia</i> | 207—211, 656 |
| <i>Euphrasia</i> | 269, 267, 268 |

F.

| | |
|-----------------------|----------------------------|
| <i>Fagus</i> | 422 |
| <i>Fagopyrum</i> | 171, 172 |
| <i>Festuca</i> | 44—52, 33, 55, 648, 654 |
| <i>Ferrum equinum</i> | 321, 322 |
| <i>Ficaria</i> | 244 |
| <i>Filago</i> | |

670 INDEX NOMINUM LATINORUM.

| | Page | | Page |
|------------------------|---|-----------------------|---|
| <i>Filago</i> | 357, 359—362 | <i>Geum</i> | 226, 179, 181 |
| <i>Filipendula</i> | 121, 217 | <i>Gladiolus</i> | 14—16, 174, 378 |
| <i>Filix</i> | 449—461 | <i>Glaſtum</i> | 299 |
| <i>Filicula</i> | 451, 453, 456 | <i>Gluacrum</i> | 229 |
| <i>Filum marinum</i> | 587 | <i>Glaux</i> | 101, 322, 323 |
| <i>Flos adonis</i> | 239 | <i>Glecoma</i> | 254, 260 |
| — <i> cuculi</i> | 294 | <i>Glycyrrhiza</i> | 322 |
| <i>Fœniculum</i> | 117, 126, 128 | <i>Gnaphalium</i> | 359—362, 357, 657 |
| <i>Fœnum græcum</i> | 324 | <i>Gramen</i> | 11, 18, 20—59, 132, 149—153, 190, 277, 400— 415, 440, 441 |
| — <i> burgundiacum</i> | 330 | <i>Graminifolia</i> | 277, 462 |
| <i>Fontales</i> | 169 | <i>Grossularia</i> | 99, 100 |
| <i>Fontinalis</i> | 467, 468, 74 | <i>Gratiola</i> | 206, 265 |
| <i>Fragaria</i> | 221—225, 136 | | H. |
| <i>Franca</i> | 137, 138 | <i>Halimus</i> | 442 |
| <i>Frangula</i> | 98 | <i>Hedera</i> | 100 |
| <i>Frankenia</i> | 137, 138 | — <i> terreſtris</i> | 254 |
| <i>Fraxinus</i> | 445, 446, 215 | <i>Hederula</i> | 340 |
| <i>Fritillaria</i> | 144 | <i>Hedypnois</i> | 340—342 |
| <i>Frutices</i> | 586 | <i>Hedysarum</i> | 322, 656 |
| <i>Fruticulus</i> | 585 | <i>Helianium</i> | 368 |
| <i>Fucus</i> | 572—591, 395, 566—570, 595, 596, 652, 661, 662 | <i>Helianthemum</i> | 332—334 |
| <i>Fucoides</i> | 587, 590 | <i>Heliotropium</i> | 79 |
| <i>Fumaria</i> | 309, 310 | <i>Helleborine</i> | 390, 392—394 |
| <i>Fungoidaſter</i> | 635 | <i>Helleboraster</i> | 245, 246 |
| <i>Fungoides</i> | 563, 632—640 | <i>Helleborus</i> | 245 |
| <i>Fungus</i> | 528, 563, 605, 607, 610 —630, 633—640, 642—644, 646 | <i>Helvella</i> | 632, 633 |
| <i>Furcellaria</i> | 589 | <i>Hepatica</i> | 512, 520—522 |
| | G. | <i>H. paticoides</i> | 509, 511, 513—515 |
| <i>Galanthus</i> | 140, 654 | <i>Heraeleum</i> | 117 |
| <i>Galeobdolon</i> | 257, 258 | <i>Herba</i> | 202 |
| <i>Galeopsis</i> | 256—259 | — <i> dysenterica</i> | 369 |
| <i>Galium</i> | 67—70, 441 | — <i> Gerardi</i> | 129 |
| <i>Geaſter</i> | 643, 644 | — <i> impia</i> | 362 |
| <i>Geaſteroides</i> | 644 | — <i> Paris</i> | 172 |
| <i>Geniſta</i> | 310 | — <i> trientalis</i> | 160 |
| <i>Geiſtelka</i> | 311, 312 | <i>Herniaria</i> | 108, 109, 72 |
| <i>Gentiana</i> | 102, 103, 168 | <i>Heſperis</i> | 288, 287, 292, 656 |
| <i>Gentianella</i> | 103 | <i>Hieracium</i> | 343—346, 337, 339 347—656 |
| <i>Geranium</i> | 300—305, 656 | <i>Hippophaë</i> | |

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

672 INDEX NOMINUM LATINORUM.

| | Page | | Page. |
|--------------------------|--|-----------------------|---|
| <i>Lepidium</i> | 279, 280, 282 | <i>Lythrum</i> | 205, 206 |
| <i>Leucoium</i> | 140, 287, 378 | | |
| <i>Leucanthemum</i> | 371 | M. | |
| <i>Lichen</i> | 523—562, 517, 519— 522, 564, 565, 605, 608, 609, 644, 652, 659—661 | <i>Malva</i> | 306—308 |
| <i>Lichen-agaricus</i> | 639, 641 | <i>Malus</i> | 216, 217 |
| <i>Lichenastrum</i> | 473, 509—519, 522, 523 | <i>Marchantia</i> | 519—521 |
| <i>Lichenoides</i> | 523—562, 565, 566, 645, 652 | <i>Majorana</i> | 262 |
| <i>Ligusticum</i> | 117, 118 | <i>Marsilea</i> | 517, 518 |
| <i>Ligustrum</i> | 3 | <i>Marrubium</i> | 260, 261, 9 |
| <i>Ligula marina</i> | 587 | <i>Matricaria</i> | 371—374 |
| <i>Lilium convallium</i> | 146 | <i>Medica</i> | 330, 331 |
| <i>Limnopeuce</i> | 2 | <i>Medicago</i> | 330, 331 |
| <i>Limodorum</i> | 387 | <i>Melampyrum</i> | 269, 270 |
| <i>Limonium</i> | 132, 133 | <i>Melica</i> | 37 |
| <i>Limofella</i> | 276 | <i>Melilotus</i> | 324 |
| <i>Lincbia</i> | 564 | <i>Melissa</i> | 263, 264 |
| <i>Lithophytoides</i> | 639 | <i>Melissophyllon</i> | 264 |
| <i>Lithospermum</i> | 78, 79 | <i>Melittis</i> | 264 |
| <i>Littorella</i> | 415 | <i>Mell-muscarum</i> | 624 |
| <i>Lobelia</i> | 377, 378 | <i>Mentha</i> | 249—254 |
| <i>Lolium</i> | 55 | <i>Menthastrum</i> | 251 |
| <i>Lonchitis</i> | 450, 456 | <i>Menyanthes</i> | 85 |
| <i>Lonicera</i> | 95 | <i>Mercurialis</i> | 435, 381 |
| <i>Lotus</i> | 329 | <i>Mezerion</i> | 167 |
| <i>Loto affinis</i> | 313 | <i>Mespilus</i> | 217, 214, 215 |
| <i>Lunaria</i> | 279, 449, 520 | <i>Meum</i> | 116 |
| <i>Lupulus</i> | 433 | <i>Milium</i> | 29, 28 |
| <i>Lutcola</i> | 207 | <i>Millefolium</i> | 9, 76, 85, 375, 419, 420 |
| <i>Lychnis</i> | 198, 199, 137, 183, 186—189, 193 | <i>Millegrana</i> | 135 |
| <i>Lychnoides</i> | 192, 198 | <i>Militaris</i> | 236 |
| <i>Lycogala</i> | 565, 645 | <i>Mnium</i> | 472, 473, 469, 470, 475, 482, 486, 487, 489—492, 519, 658 |
| <i>Lycoperdon</i> | 641—645, 664 | <i>Mollugo</i> | 60, 67, 68, 70 |
| <i>Lycoperdastrum</i> | 645 | <i>Molle</i> | 139 |
| <i>Lycopodium</i> | 462—464, 658 | <i>Monotropa</i> | 175 |
| <i>Lycopsis</i> | 82, 83 | <i>Montia</i> | 60 |
| <i>Lycopus</i> | 9 | <i>Mursus diaboli</i> | 62 |
| <i>Lyfimachia</i> | 85—87, 161—163, 206, 265 | <i>-----ranae</i> | 436 |
| | | <i>Moscatellina</i> | 171 |
| | | <i>Mucedo</i> | 646 |
| | | <i>Mucor</i> | 646, 647, 631, 632, 645 |
| | | <i>Muscipula</i> | |

INDEX NOMINUM LATINORUM. 673

| | Page |
|-----------------------------|---------------------|
| <i>Muscipula</i> | 186, 199 |
| 486—509, 511—515, 540, | |
| —562, 567, 568, 587, 593, | 55 |
| 596, 603, 604 | |
| <i>Muscoides</i> | |
| <i>Musco fungus</i> | 531, 537, 538, 541, |
| 545, 553, 555—559, 560, 561 | |
| <i>Myagre affinis</i> | 287 |
| <i>Myagrurn</i> | 277 |
| <i>Myolotis</i> | 78, 190, 200—202 |
| <i>Myosurus</i> | 136 |
| <i>Myrica</i> | 432 |
| <i>Myriophyllum</i> | 419, 420 |
| <i>Myrrhis</i> | 124, 125 |
| <i>Myrtus</i> | 432 |

N.

| | |
|--------------------------|-------------------|
| <i>Napus</i> | 290 |
| <i>Narcissus</i> | 141 |
| <i>Nardus</i> | 22, 32 |
| <i>Narthesium</i> | 145 |
| <i>Nasturtium</i> | 278, 280—284, 293 |
| | —297 |
| <i>Nasmythia</i> | 414, 415 |
| <i>Nepeta</i> | 249 |
| <i>Nidus avis</i> | 386, 388 |
| <i>Noli me tangere</i> | 381 |
| <i>Nosloc</i> | 363—365 |
| <i>Nucula terrestris</i> | 122 |
| <i>Nummularia</i> | 87 |
| <i>Nux vesicaria</i> | 131 |
| <i>Nymphaea</i> | 334, 85, 436 |

O.

| | |
|---------------------|----------|
| <i>Ocymum</i> | 236 |
| <i>Ocymoides</i> | 187 |
| <i>Odontites</i> | 268 |
| <i>Oenanthe</i> | 121, 654 |
| <i>Omphalomyces</i> | 611, 613 |

| | Page |
|----------------------|--------------------|
| <i>Onobrychis</i> | 322 |
| <i>Ononis</i> | 312, 313 |
| <i>Onopordon</i> | 354 |
| <i>Onopteris</i> | 454 |
| <i>Ophyoglossum</i> | 449 |
| <i>Ophrys</i> | 387—392, 657 |
| <i>Opulus</i> | 130 |
| <i>Orchis</i> | 387, 392, 657 |
| <i>Oreofelinum</i> | 115 |
| <i>Origanum</i> | 262 |
| <i>Ornithopus</i> | 321 |
| <i>Ornithogalum</i> | 143, 142, 144, 654 |
| <i>Ornithopodium</i> | 321 |
| <i>Ornus</i> | 215 |
| <i>Orobanche</i> | 265—267, 175, 388 |
| <i>Orobus</i> | 314, 656 |
| <i>Osmunda</i> | 449, 450 |
| <i>Ostrya</i> | 422 |
| <i>Ostrya</i> | 422 |
| <i>Othonna</i> | 369, 370 |
| <i>Oxalis</i> | 198, 156 |
| <i>Oxyacantha</i> | 215 |
| <i>Oxycoccus</i> | 165 |
| <i>Oxys</i> | 198 |

P.

| | |
|------------------------|---------------------|
| <i>Padus</i> | 313 |
| <i>Palma christi</i> | 385 |
| <i>Panicum</i> | 24, 25, 29 |
| <i>Panax coloni</i> | 259 |
| <i>Papaver</i> | 230, 231, 186, 229 |
| <i>Paralyfis</i> | 84 |
| <i>Parietaria</i> | 442 |
| <i>Paris</i> | 172 |
| <i>Parnassia</i> | 131, 132 |
| <i>Paronicbia</i> | 101, 182, 278, 279, |
| | 292 |
| <i>Pastinaca</i> | 125, 119 |
| <i>Pecten veneris</i> | 123 |
| <i>Pedicularis</i> | 270, 271, 268 |
| <i>Peloria</i> | 273 |
| <i>Pentapbylloides</i> | 136, 222, 223, |
| | 227 |
| X x | <i>Pentapbyllum</i> |

674 INDEX NOMINUM LATINORUM.

| | Page |
|--------------------------|------------------------|
| <i>Pentaptyllum</i> | 71, 223—225 |
| <i>Pentopteris</i> | 419, 420 |
| <i>Pentapterisptylum</i> | 420 |
| <i>Peplis</i> | 147, 207, 208 |
| <i>Peplus</i> | 208 |
| <i>Percepius</i> | 72 |
| <i>Perfoliata</i> | 111, 290 |
| <i>Perichlymenum</i> | 71, 94 |
| <i>Periscaria</i> | 169, 170, 381 |
| <i>Peruvia</i> | 91 |
| <i>Pes anserinus</i> | 105 |
| — <i>columbinum</i> | 307 |
| <i>Petasites</i> | 364 |
| <i>Peucedanum</i> | 116, 128 |
| <i>Peziza</i> | 633—637, 563, 661, 669 |
| <i>Phalangium</i> | 143—145 |
| <i>Phalaris</i> | 23, 26 |
| <i>Phallus</i> | 629, 630 |
| <i>Phoenix</i> | 441 |
| <i>Phascum</i> | 466, 467, 469 |
| <i>Phellandrium</i> | 122 |
| <i>Phleum</i> | 25, 26, 23, 28 |
| <i>Phyllitis</i> | 452 |
| <i>Phytuma</i> | 97 |
| <i>Picea</i> | 424 |
| <i>Picris</i> | 342, 341 |
| <i>Pilularia</i> | 462 |
| <i>Pilosella</i> | 343—345 |
| <i>Pimpinella</i> | 127, 128, 65, 421, 654 |
| <i>Pinguicula</i> | 8, 653 |
| <i>Pinus</i> | 423, 424, 590 |
| <i>Piperitis</i> | 279 |
| <i>Pistacea</i> | 132 |
| <i>Pisum</i> | 313, 314 |
| <i>Plantaginella</i> | 276 |
| <i>Plantago</i> | 63—65, 158, 159, 415 |
| <i>Pneumonanthe</i> | 162 |
| <i>Poa</i> | 38—43, 31—34, 46, 47 |
| <i>Podagraria</i> | 129 |
| <i>Polemonium</i> | 89 |
| <i>Polycarpon</i> | 60 |
| <i>Polygalon</i> | 322 |

| | Page |
|--------------------------|--|
| <i>Polygonatum</i> | 140, 147 |
| <i>Polygonum</i> | 168—172, 72, 74, 100, 108, 135, 137, 178 |
| <i>Polypodium</i> | 455—460 |
| <i>Polyperus</i> | 625—628 |
| <i>Polytrichon</i> | 489 |
| <i>Polytrichum</i> | 469—472 |
| <i>Populago</i> | 245 |
| <i>Populus</i> | 433, 434 |
| <i>Portulaca</i> | 60, 147, 442 |
| <i>Porrum</i> | 139, 649 |
| <i>Potamogeton</i> | 74—77, 169, 395, 397 |
| <i>Potentilla</i> | 222—225, 656 |
| <i>Poterium</i> | 421 |
| <i>Prenanthes</i> | 338 |
| <i>Primula</i> | 83, 84 |
| <i>Prunella</i> | 264 |
| <i>Prunus</i> | 212, 213 |
| <i>Pseudo-asphodelus</i> | 145, 147 |
| — <i>cyperus</i> | 14, 21, 416 |
| — <i>iris</i> | 14 |
| — <i>linarea</i> | 101 |
| — <i>melantheum</i> | 198 |
| — <i>Narcissus</i> | 141, 144 |
| — <i>Orcus</i> | 387, 390 |
| <i>Psarmica</i> | 375 |
| <i>Pteris</i> | 451 |
| <i>Pulegium</i> | 254 |
| <i>Pulmonaria</i> | 80, 81, 344, 345 |
| <i>Pulsatilla</i> | 237 |
| <i>Pyrola</i> | 175, 176, 160, 387, 655 |
| <i>Pyrus</i> | 216 |
| <i>Pyraister</i> | 216 |

Q.

| | |
|----------------------|---------------|
| <i>Quercus</i> | 421 |
| — <i>marina</i> | 577, 584 |
| <i>Quinquefolium</i> | 223, 225, 227 |

R.

| | |
|-----------------|-----|
| <i>Radiola</i> | 194 |
| <i>Radicula</i> | |

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

676 INDEX NOMINUM LATINORUM.

| | Page | | Pa |
|-----------------------|--------------------------------|-----------------------|--------------------|
| <i>Sherardia</i> | 66 | <i>Stramonium</i> | |
| <i>Sibbaldia</i> | 136 | <i>Stratiotes</i> | 236, 4 |
| <i>Sibthorpia</i> | 276 | <i>Stratbiopteris</i> | 4 |
| <i>Sideritis</i> | 256, 260 | <i>Subularia</i> | 277, |
| <i>Silene</i> | 187—189 | <i>Succisa</i> | |
| <i>Silibum</i> | 353 | <i>Suillus</i> | 6 |
| <i>Sinapi</i> | 297 | <i>Swertia</i> | 101, 1 |
| <i>Sinapis</i> | 297, 298 | <i>Symphytum</i> | 81, 6 |
| <i>Sifon</i> | 119, 120, 654 | <i>Sycemorus</i> | |
| <i>Sisymbrium</i> | 295—297, 259, 290, 291, 656 | T. | |
| <i>Sium</i> | 118—120, 122, 295 | <i>Tamus</i> | 43 |
| <i>Smilax</i> | 88, 89 | <i>Tanacetum</i> | 3 |
| <i>Smyrniun</i> | 126, 118 | <i>Taraxacum</i> | 3 |
| <i>Solanum</i> | 22, 93, 172 | <i>Targionia</i> | 5 |
| <i>Solanifolia</i> | 10, 11 | <i>Taxus</i> | 436, 4 |
| <i>Solidago</i> | 367 | <i>Telebium</i> | 1 |
| <i>Soldanella</i> | 89 | <i>Testiculus</i> | 3 |
| <i>Sonchus</i> | 336—338 | <i>Teucrium</i> | 247, 248, 2 |
| <i>Soncho-affinis</i> | 348 | <i>Thalictrum</i> | 238, 239, 6 |
| <i>Sophia</i> | 297 | <i>Tbapsus</i> | |
| <i>Sorbus</i> | 215, 216, 214 | <i>Thesium</i> | 1 |
| <i>Sparganium</i> | 400—401 | <i>Thlaspi</i> | 281—285, 2 |
| <i>Spartium</i> | 310 | <i>Thymus</i> | 262, 2 |
| <i>Spartum</i> | 22, 29, 43, 54, 56 | <i>Thysselinum</i> | 1 |
| <i>Spatula</i> | 14 | <i>Tilia</i> | 231, 2 |
| <i>Spergula</i> | 202, 203, 193, 276 | <i>Tillæa</i> | |
| <i>Sphagnum</i> | 465—468, 474 | <i>Titbymalus</i> | 207, 2 |
| <i>Sphæria</i> | 565, 639, 641, 644, 665 | <i>Tofieldia</i> | 1 |
| <i>Spondylium</i> | 117 | <i>Tordylium</i> | 112, 1 |
| <i>Spina alba</i> | 354 | <i>Tormentilla</i> | 225, |
| — <i>acida</i> | 137 | <i>Trachelium</i> | |
| — <i>Appendix</i> | 215 | <i>Tragopogon</i> | 3 |
| <i>Spirea</i> | 217 | <i>Tragoselinum</i> | 1 |
| <i>Splachnum</i> | 468, 469 | <i>Tragorchis</i> | 383, 3 |
| <i>Spongia</i> | 570, 584, 647 | <i>Tragum</i> | 1 |
| <i>Stachys</i> | 259, 260, 656 | <i>Tremella</i> | 562—569, 537, 572, |
| <i>Stæbelina</i> | 267 | <i>Tribulus</i> | |
| <i>Stapyledendron</i> | 131 | <i>Trichia</i> | 630, 6 |
| <i>Staphylea</i> | 131 | <i>Trichomanes</i> | 461, 452, 4 |
| <i>Statyce</i> | 132, 133 | <i>Trientalis</i> | 1 |
| <i>Stellaria</i> | 189, 190, 439, 440 | <i>Trifolium</i> | 323—331, 85, 1 |
| <i>Stipa</i> | 29 | <i>Trigloch</i> | |

INDEX NOMINUM LATINORUM. 677

| | Page | | Page |
|---------------------|------------------|--------------------------|---|
| <i>Triglochin</i> | 152, 461 | <i>Virga-aurea</i> | 366—368 |
| <i>Triorchis</i> | 389 | — <i>pastoris</i> | 61 |
| <i>Tripolium</i> | 368 | <i>Viscago</i> | 186—189 |
| <i>Triticum</i> | 57, 58 | <i>Viscum</i> | 431 |
| <i>Trixago</i> | 260 | <i>Vitis idæa</i> | 163—165, 177 |
| <i>Trollius</i> | 244 | <i>Volubilis</i> | 171 |
| <i>Tuber</i> | 641 | <i>Valneraria</i> | 313 |
| <i>Tubera</i> | 641 | <i>Vulvaria</i> | 107 |
| <i>Tunica</i> | 183, 185 | | |
| <i>Tu.ritis</i> | 291—293 | U. | |
| <i>Tussilago</i> | 364 | | |
| <i>Typha</i> | 340 | <i>Ulex</i> | 312 |
| | | <i>Ulmaria</i> | 218 |
| V. | | <i>Ulmus</i> | 109 |
| <i>Vaccina</i> | 163—165, 177 | <i>Ulva</i> | 566—572, 518, 522, 564, 581, 600, 652, 661 |
| <i>Vaccinium</i> | 163—165 | <i>Umbelicus veneris</i> | 194 |
| <i>Valantia</i> | 441 | <i>Urtica</i> | 417, 418, 257 |
| <i>Valeriana</i> | 12, 13, 89 | <i>Usnea</i> | 559—562, 570 |
| <i>Valerianella</i> | 13 | <i>Utricularia</i> | 8, 9 |
| <i>Veila</i> | 278 | <i>Uva ursi</i> | 177 |
| <i>Verbascum</i> | 85—61 | | |
| <i>Verbasculum</i> | 84 | X. | |
| <i>Verbena</i> | 249 | <i>Xanthium</i> | 418 |
| <i>Verbena</i> | 355, 356 | <i>Xyris</i> | 14 |
| <i>Vermicularis</i> | 108, 195, 196 | | |
| <i>Veronica</i> | 3—8, 272 | Z. | |
| <i>Viburnum</i> | 129, 130, 654 | | |
| <i>Vicia</i> | 317—321 | <i>Zannichellia</i> | 397 |
| <i>Vinca</i> | 91 | <i>Zosteria</i> | 395 |
| <i>Viola</i> | 378—380, 85, 288 | | |
| <i>Viorna</i> | 283 | | |

I N D E X

NOMINUM ANGLICANORUM.

| A. | | Page | | | Page |
|-----------------|------|------|---------------|------|------|
| A CORUS | | 147 | Arrowhead | | 420 |
| Acrostichum | 450, | 451 | Arsmart | 170, | 171 |
| Adders tongue | | 449 | Afarabacca | | 205 |
| Adonis | | 239 | Ash-tree | | 446 |
| Agaric | 610, | 624, | — mountain | | 215 |
| Agrimony | | 206 | — weed | | 129 |
| — hemp | | 355 | Asphodel | 145, | 157 |
| Alder | | 416 | Astragalus | | 323 |
| — berry-bearing | | 98 | Avens | | 226 |
| Alexanders | | 126 | Azalea | | 88 |
| Alkanet | | 80 | | | |
| — bastard | | 79 | B. | | |
| Alheal | | 259 | Ballote | | 260 |
| — feed | | 135 | Balm, bastard | | 264 |
| Althæa | | 306 | Balsam | | 381 |
| Amaranth | | 419 | Bane-berries | | 228 |
| Angelica | | 118 | Barberry | | 137 |
| Anthericum | | 144 | Barley-grass | | 56 |
| Anthoceros | | 523 | Bartsia | 267, | 268 |
| Anemony | 236, | 237 | Basil | 261, | 263 |
| Apple-tree | | 217 | Bawd-money | | 116 |
| — thorny | | 92 | Beam-tree | | 214 |
| Arbutus | | 177 | Bears-foot | | 245 |
| Archangel | 255, | 256 | — berry | | 177 |
| | | | Bedstraw | | |

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

680 INDEX NOMINUM ANGLICANORUM.

| | Page |
|--------------------|---------------|
| Champignon | 616 |
| Charlock | 298 |
| Chara | 398 |
| Cheese-renning | 69 |
| | 193 |
| Cherry-tree | 213 |
| Chervil | 125 |
| | 228 |
| | 422 |
| Chickweed | 131 |
| ——— germander | 7 |
| ——— mouse-ear | 200, 201 |
| ——— water | 60 |
| ——— winter-green | 161 |
| Chives | 649 |
| Cicely | 124 |
| Cicuta | 122 |
| Cineraria | 370 |
| Cinquefoil | 222—225 |
| ——— marsh | 227 |
| Cistus | 232—234 |
| Cives | 649 |
| Clary | 10 |
| Clathrus | 630—632 |
| Clavaria | 638—640 |
| Cleavers | 70 |
| Climber | 238 |
| Ciot bur | 348 |
| Cloud-berry | 221 |
| Clover | 325, 326, 331 |
| Cockle | 198 |
| Cock's comb | 268 |
| ——— head | 322 |
| Codlings and Cream | 162 |
| Colchicum | 155 |
| Colewort | 299 |
| Coltsfoot | 364 |
| Columbine | 235, 236 |
| Comarum | 227 |
| Comfrey | 81, 648 |
| Conferva | 592—604, 653 |
| Conyza | 363 |
| Coral-wort | 286 |

| | Page |
|------------------|----------|
| Coriander | 123 |
| Cornel-tree | 79, 71 |
| Corn-fallet | 13 |
| Cow-parsnep | 117 |
| ——— quakes | 38 |
| ——— weed | 125 |
| ——— wheat | 269, 270 |
| Cowslips | 84 |
| Crab-tree | 216 |
| Crake-berries | 165 |
| Crepis | 339 |
| Cress, bastard | 281, 283 |
| ——— bitter | 294 |
| ——— rocket | 278 |
| ——— rock | 285 |
| ——— swines | 284 |
| ——— wall | 292, 293 |
| ——— water | 296, 297 |
| ——— winter | 286 |
| Crocus | 13 |
| Crosswort | 441 |
| Crowfoot | 240—243 |
| ——— berries | 431 |
| ——— silk | 592 |
| Cuckow-flower | 199, 294 |
| ——— pint | 395 |
| Cudweed | 359—362 |
| Currants | 99 |
| Cyperus, bastard | 21 |
| ——— millet | 21 |
| ——— sweet | 17 |

D.

| | |
|---------------|----------|
| Daffodil | 141 |
| ——— chequered | 144 |
| Daisy | 370, 371 |
| Dandelion | 339 |
| Danewort | 130 |
| Daphne | 167 |
| Datura | 92 |
| Devil's bit | 62, 342 |
| Dewberry | 220 |
| Diers- | |

INDEX NOMINUM ANGLICANORUM. 681

| | |
|-------------|----------|
| | Page |
| Diers-weed | 220 |
| Dittander | 279 |
| Dock | 153—156 |
| Dodder | 204 |
| Dogberry | 70 |
| Dropwort | 217 |
| ———— water | 121 |
| Dryas | 227 |
| Duck's meat | 399, 340 |
| Dwale | 93 |

E.

| | |
|-------------|----------|
| Echinophora | 112 |
| Elder | 130 |
| ———— water | 130 |
| Elecampane | 368, 369 |
| Elm | 109 |
| Epilobium | 161—163 |
| Erigeron | 363 |
| Eryngo | 109, 110 |
| Kyebright | 269 |
| Eupatorium | 356 |

F.

| | |
|------------------|----------|
| Fennel | 126 |
| ———— hogs | 116 |
| Fern, female | 451 |
| ———— flowering | 450 |
| ———— male | 458 |
| ———— stone | 450 |
| Feverfew | 372, 373 |
| Figwort | 275 |
| Pinkle | 126 |
| Fir | 423, 424 |
| Fiveleaved-grass | 225 |
| Flax | 133—135 |
| Pleabane | 369 |
| Flixweed | 297 |
| Flos adonis | 239 |
| Flower-de luce | 14 |
| ———— gentle | 419 |

| | |
|------------|--------------|
| | Page |
| Fluellin | 272 |
| Fontinalis | 467, 468 |
| Foxgloves | 276 |
| Frankenia | 137, 138 |
| Fritillary | 144 |
| Frog-bit | 436 |
| Fucus | 573—592, 653 |
| Fumitory | 309, 310 |
| Furze | 312 |

G.

| | |
|---------------------|--------------|
| Gale | 432 |
| Galeobdolon | 258 |
| Galingale | 17 |
| Garlick | 138—140, 649 |
| Gentian | 102, 103 |
| Gatter-tree | 70 |
| Germander | 6, 247, 248 |
| Gilly-flower, sea | 130 |
| Gladiole | 578 |
| Gladwyn | 14 |
| Glasswort | 1 |
| Globeflower | 244 |
| Goats-beard | 335 |
| Goldilocks | 242 |
| Gold of pleasure | 277 |
| Golden-rod | 367, 368 |
| Goose-corn | 144 |
| ———— foot | 104—107 |
| ———— grass | 69, 70, 82 |
| ———— tongue | 375 |
| Gout-weed | 129 |
| Grass, arrow-headed | 152 |
| ———— Barley | 56, 57 |
| ———— Dent | 30—32 |
| ———— Brome | 49—51 |
| ———— Canary | 23, 24 |
| ———— Cat's tails | 26 |
| ———— Cook's foot | 43, 44 |
| ———— Cotton | 22 |
| ———— Couch | 58 |
| ———— Darnel | 55 |
| Grass, | |

682 INDEX NOMINUM ANGLICANORUM.

| | Page | | Page |
|--------------|------------|------------------|----------|
| Grass, Dog's | 58 | Hafel, wych | 109 |
| — — tail | 59 | Hawke-weed | 343—346 |
| — Feather | 29 | Haw-tree. | 214, 215 |
| — Fescue | 44—48, 648 | Heath | 165, 167 |
| — Fiveleaved | 225 | — berried | 431 |
| — Fox-tail | 27—29 | — sea | 137, 138 |
| — Hard | 441 | Hedge-hyssop | 206 |
| — Hair | 33—37 | Hedypnois | 341, 342 |
| — Knot | 171 | Hellebore | 245, 246 |
| — Lime | 56 | Helleborine | 392, 393 |
| — Marle | 326 | Hellweed | 104 |
| — Mat | 22 | Helvella | 633 |
| — Meadow | 38—43 | Hemlock | 115 |
| — Melic | 37 | — water | 123 |
| — Millet | 30 | Henbane | 92 |
| — Oat | 52, 53 | Henbit | 7, 256 |
| — Panic | 24, 25 | Hep-tree | 220 |
| — Parnassus | 132 | Herb-christopher | 228 |
| — Pepper | 462 | — bennet | 226 |
| — Quaking | 38 | — Gerard | 129 |
| — Quick | 58 | — Paris | 172 |
| — Reed | 54 | — Robert | 305 |
| — Soft | 440, 441 | High-taper | 98 |
| — Spring | 11 | Holly-tree | 446 |
| — Scurvy | 283, 284 | — Knee | 437 |
| — Spiked | 153 | — Sea | 109 |
| — Vernal | 11 | Honey-wort | 120 |
| — Wheat | 58 | Honey-suckle | 94 |
| — Wrack | 396 | Horehound, black | 260 |
| Green-weed | 311, 312 | — water | 9 |
| Greymill | 79 | — white | 261 |
| Gromwell | 79 | Hornbeam | 423 |
| Ground-ivy | 254 | Hornwort | 419 |
| — pine | 248 | Horse-beach | 425 |
| Groundsel | 365, 367 | — radish | 284 |
| | | — shoe-vetch | 322 |
| | | — tail | 447—449 |
| | | Houseleek | 211 |
| | | Hounds-tongue | 80 |
| | | Hyacinth | 142 |
| | | Hydnum | 628, 629 |
| | | Hypnum | 493, 509 |
| | | Hyoseris | 346 |
| | | Hypochæris | 347 |
| | | I. Ille- | |

H,

| | |
|----------------|-----|
| Hardbeam | 423 |
| Harebells | 142 |
| Hare's-ear | 111 |
| — tail | 22 |
| Hart's-tongue | 452 |
| Hafel-nut tree | 423 |

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page



HISTORY

Tens of thousands of important historical sources, many previously unobtainable, are now available for the first time with a Forgotten Books Full Membership.

Unlimited Access
\$8.99/month

Continue

*Fair usage policy applies

684 INDEX, NOMINUM ANGLICANORUM.

| | Page | | Page |
|--------------------|---------------|----------------|----------|
| Mniem | 472—474 | Nettle, dead | 255, 256 |
| Mo. — wort | 87 | —— hemp | 256, 257 |
| —— Cornish | 276 | —— hedge | 259 |
| Monotropa | 175 | —— roman | 417 |
| Moonwort | 449 | Nightshade | 93 |
| Moorballs | 604 | —— deadly | 93 |
| Morel | 629, 630 | —— enchanter's | 10, 11 |
| Morocco | 239 | Nipplewort | 348 |
| Moscatel | 172 | Nonfuch | 331 |
| Moss, bog | 465 | Nut, earth | 122 |
| —— cup | 555 | —— bladder | 131 |
| —— Club | 463, 464 | —— Kipper | 122 |
| —— tree | 566 | —— Hawke | 122 |
| —— water | 567, 568 | —— Pig | 122 |
| Mother-wort | 261 | | |
| Mould | 646, 647 | O. | |
| Mouse-ear | 343 | Oak-tree | 421 |
| —— scorpion-grafs | 78 | —— sea | 577 |
| —— tail | 136 | Oneberry | 172 |
| Mudwort | 276 | Onopordum | 354 |
| Mugwort | 276 | Ophrys | 388—392 |
| Mullein | 90, 91 | Orache | 442—444 |
| Mushroom | 610 | —— stinking | 107 |
| Muscovy | 301 | Orchis | 382—387 |
| Mustard | 298 | Orobus | 314 |
| —— hedge | 286, 287, 297 | Orpine | 195 |
| —— Mithridate | 281 | Osier | 430 |
| —— tower | 291, 292 | Osmund-royal | 450 |
| —— treacle | 282 | Ox-eye | 371, 374 |
| Myagrum | 277 | — slips | 84 |
| Myrtle | 432 | Oyster-green | 567 |
| N. | | P. | |
| Nape | 290 | Pagils | 84 |
| Narcissus | 141 | Panick-grafs | 24, 25 |
| Narthefium | 145 | Parsley | 129 |
| Nasmythia | 415 | —— fools | 123 |
| Navew | 290 | —— stone | 120 |
| Navel-wort | 194 | Parsnep, cows | 117 |
| Needle, shepherd's | 124 | —— water | 119 |
| Nep-mint | 249 | —— wild | 125 |
| Nettle | 417, 418 | Pasque- | |

INDEX NOMINUM ANGLICANORUM. 685

| | Page | | Page |
|-------------------|---------------|-------------------|-------------------|
| Pasque-flower | 237 | Prickmadman | 195 |
| Pear-tree | 216 | — wood | 70, 99 |
| Pearlwort | 73 | Prim | 3 |
| Peas, everlasting | 237 | Primrose | 84 |
| — heath | 314 | Privet | 9 |
| — sea | 314 | Prunus | 212, 213 |
| Pellitory, wall | 442 | Puff-balls | 641—645, 653 |
| Penny-royal | 254 | Purslain-water | 147 |
| Pennywort, marsh | 110 | — sea | 444 |
| — wall | 194 | Purley-piert | 72 |
| Replis | 147 | | |
| Pepper-grass | 462 | Q. | |
| — wall | 196 | Quicken-tree | 215 |
| — wort | 279, 280 | Quick in the hand | 388 |
| Periwinkle | 91 | Quil-wort | 462 |
| Perficaria | 169, 171 | | |
| Peziza | 634—637 | R. | |
| Pnascum | 466, 467 | Radish, horse | 284 |
| Pheasant's eye | 239 | — water | 297 |
| Picris | 343 | Ragwort | 366, 370 |
| Pilcorn | 52 | Rampions | 95 |
| Pilewort | 244 | Ramsons | 146 |
| Pimpernel, male | 87 | Rape | 296 |
| — water | 94 | Raspberry | 220 |
| — yellow | 86 | Rattle, yellow | 268 |
| Pimpinell | 127, 128 | Reed | 54 |
| Pink | 184, 185, 199 | — mace | 400 |
| Pipperidge-bush | 127 | Reseda | 207 |
| Pitch tree | 424 | Rest-harrow | 312, 313 |
| Plantain | 63—65 | Rhamnus | 98 |
| — water | 158, 159 | Ribwort | 64 |
| Plumb | 212 | Riccia | 521, 522 |
| Plowman spikenard | 363 | Rocket, base | 207 |
| Polycarpon | 60 | — sea | 298 |
| Polymonium | 89 | — wild | 288, 291, 296 |
| Polypody | 455—460 | Rose-tree | 218—220 |
| Polytrichum | 470—472 | — wort | 434 |
| Pondweed | 74—77 | Rue, meadow | 238, 239 |
| Poplar | 434 | Rupture-wort | 108, 109 |
| Poppy | 230—231 | Ruppia | 77 |
| — horned | 229 | Ruth | 148—152, 649, 650 |
| — spatling | 186 | Ruth, | |
| Prenanthes | 339 | | |

686 INDEX NOMINUM ANGLICANORUM.

| | Page | | Page |
|-----------------|-------------------|----------------------|----------|
| Rash, bog | 15, 16 | Setterwort | 245 |
| — bull | 19, 20 | Shepherd's needle | 124 |
| — club | 17—21, 648 | — purple | 283 |
| — flowering | 174 | Sherardia | 66 |
| — hares-tail | 22 | Sibbaldia | 136 |
| | | Sibthorpia | 276 |
| | | Silverweed | 223 |
| S. | | Simpson | 365 |
| Sage | 10 | Skullcap | 265 |
| — wood | 248 | Sloe-tree | 212 |
| Saffron | 13, 157 | Smallage | 129 |
| Saintfoin | 322 | Snake's head | 144 |
| St. John's wort | 332—334 | Snakeweed | 168 |
| — Peter's wort | 334 | Snapdragon | 274 |
| Sallow | 430 | Sneezewort | 375 |
| — thorn | 432 | Snowdrop | 140 |
| Saltwort | 1, 107 | Soap-wort | 183 |
| Samolus | 94 | Soldier, fresh-water | 236 |
| Sampire | 11, 112, 117, 369 | Solomon's seal | 146, 147 |
| Sandwort | 192, 193 | Sorb | 215 |
| Sanicle | 111 | Sorrel | 156 |
| Santolina | 357 | — wood | 198 |
| Satyrion | 386, 387 | Southernwood | 358 |
| Sauce-alone | 287 | Sow-thistle | 336, 337 |
| Saw-wort | 349, 350 | — bane | 105 |
| Saufrage | 179—183 | Sparagus | 146 |
| — burnet | 65, 127, 128 | Spearwort | 240 |
| — English | 203 | Sphagnum | 465 |
| — golden | 179 | Spignel | 115, 116 |
| — meadow | 116 | Spikeard | 363 |
| Scabious | 62, 63 | Spindle-tree | 99 |
| — sheeps | 377 | Spiraea | 217, 218 |
| Scorpion gras | 78 | Splachnum | 468, 469 |
| Scurvy-gras | 283, 284 | Spleenwort | 452—456 |
| — Scottish | 89 | Spurge | 202—211 |
| Self heal | 264 | — laurel | 167 |
| Sea-beard | 601 | — olive | 167 |
| — belts | 579 | Squill | 142 |
| — Buck-thorn | 432 | Squinancy berries | 100 |
| — laes | 567 | — wort | 67 |
| Selinum | 115 | Stachys | 259, 260 |
| Septfoil | 225 | Star of Bethlahem | 143, 144 |
| Service | 214—216 | — wort | 368 |
| | | Stitchwort | |

THIS PAGE IS LOCKED TO FREE MEMBERS

Purchase full membership to immediately unlock this page

SAVE \$3,999,994

Did you know we sell
paperback books too?

To buy our entire catalog
in paperback would cost
over \$4,000,000

Access it all now for
\$8.99/month

*Fair usage policy applies

Continue

688 INDEX NOMINUM ANGLICANORUM.

| | Page | | Page |
|-----------------|---------------|--------------|------|
| White-rot | 110 | Wormwood | 358 |
| Whitlow-grass | 182, 278, 279 | Wrack, grass | 396 |
| Whortle-berries | 164 | ——— sea | 576 |
| Wilding | 216 | | |
| Winter-green | 176 | X. | |
| Whorts | 164 | | |
| Williams, wild | 199 | Xanthium | 418 |
| Willow | 425—430, 651 | | |
| ——— herb | 161—163 | Y. | |
| ——— ——— hooded | 265 | | |
| ——— rosebay | 161 | Yarrow | 375 |
| ——— sweet | 432 | Yellow wort | 168 |
| Woad, wild | 299 | Yew-tree | 437 |
| Wolfs-bane | 650 | | |
| Woodbine | 294 | Z. | |
| ——— roof | 66, 67 | | |
| ——— waxen | 311 | Zannichellia | 397 |
| Wormseed | 88 | | |