

Flora of Minnesota

Authors: Anita Cholewa

Locality: Minnesota

Abstract:

A comprehensive, annotated list of the vascular plants known, or reported, to occur in Minnesota (native, naturalized, and adventive species).

Family delineations are based on the Angiosperm Phylogeny Group arrangement (Bot. J. Linnean Soc. 141: 399-436, 2003) and species names are based on published volumes of the Flora of North America (FNA), recent journals for changes post FNA (if widely accepted by the major taxonomic indices), and Gleason and Cronquist 1991 for others. Although not all taxonomists may agree with FNA, a wide array of experts from many areas have been consulted and this consequently represents the most up-to-date understanding of our flora.

Common names as given in the USDA-NRCS Plants database (2008) are also provided, followed by other common names widely used, and by Indian names where known.

County-level distribution maps are available at www.bonap.org

Families: 158

Genera: 806

Species: 2647 (species rank)

Total Taxa: 2832 (including subsp. and var.)

ACANTHACEAE

Ruellia humilis Nuttall - fringeleaf wild petunia

reported for Minnesota (Washington Co.) by MN-DNR but no specimens at MIN; known from NE, IA, WI, south- and eastward; special concern

ACORACEAE

Acorus americanus (Rafinesque) Rafinesque - sweetflag

root extract reportedly repels some insects

Acorus calamus L. - calamus

native to Europe; sterile polyploid species

ADOXACEAE

Adoxa moschatellina L. - muskroot

Sambucus nigra subsp. ***canadensis*** (L.) Bolli - black elderberry

Sambucus racemosa var. ***racemosa*** - red elderberry

Viburnum acerifolium L. - mapleleaf viburnum

erroneous report

Viburnum edule (Michaux) Rafinesque - squashberry

Viburnum lantana L. - wayfaring tree

native to Eurasia

Viburnum lentago L. - nannyberry

Viburnum opulus var. ***americanum*** Aiton - American cranberrybush

Viburnum opulus var. ***opulus*** - European cranberrybush

native to Eurasia

Viburnum rafinesquianum var. ***affine*** (Bush ex C.K. Schneider) House - downy arrowwood

Viburnum rafinesquianum var. ***rafinesquianum*** - downy arrowwood

ALISMATACEAE

Alisma gramineum Lej. - narrowleaf water plantain

Alisma subcordatum Rafinesque - American water plantain

Alisma triviale Pursh - northern water plantain

Sagittaria brevirostra Mackenzie & Bush - shortbeak arrowhead

Sagittaria cristata Engelmann - crested arrowhead

Sagittaria cuneata E. Sheldon - arumleaf arrowhead

Sagittaria graminea subsp. ***graminea*** - grassy arrowhead

reported for Minnesota by FNA but no specimens at MIN; known from Ontario, WI, IA

Sagittaria latifolia Willdenow - broadleaf arrowhead

corms are edible

Sagittaria montevidensis subsp. ***calycina*** (Engelmann) Begon - hooded arrowhead

Sagittaria rigida Pursh - sessilefruit arrowhead

ALLIACEAE

Allium canadense var. ***canadense*** - meadow garlic

Allium cernuum Roth - nodding onion

Allium hollandicum R.M. Fritsch - purple sensation onion

hybrid ornamental rarely escaping; known only from St. Louis Co. (one roadside population, collected 2005);

Allium schoenoprasum L. - wild chives

vars. not recognized; endangered

Allium stellatum Ker-Gawler - autumn onion

Allium textile A. Nelson & J.F. Macbride - textile onion

Allium tricoccum var. ***burdickii*** Hanes - narrowleaf wild leek

Allium tricoccum var. *tricoccum* - ramp

Allium tuberosum Rottler ex Spreng. - Chinese chives

Allium x proliferum (Moench) Schrad. ex Willd. - garden onion

AMARANTHACEAE

Amaranthus albus L. - prostrate pigweed

native to Eurasia; often confused with *A. blitoides*

Amaranthus blitoides S. Watson - mat amaranth

considered native to the central US by FNA; state DOA noxious (CN) weed, 1st collected 1886, Winona Co.

Amaranthus caudatus L. - love-lies-bleeding

report unverified

Amaranthus graecizans L.

misapplied to North American flora, see *A. albus* or *A. blitoides*; misapplied name

Amaranthus hybridus L. - slim amaranth

native to eastern US; highly variable and often confused with *A. powellii* or *A. retroflexus*;

Amaranthus hypochondriacus L. - Prince-of-Wales feather

native to southwestern US; adventive near cultivated fields but unlikely to be persisting;

Amaranthus palmeri S. Watson - carelessweed

native to southern North America

Amaranthus powellii S. Watson - Powell's amaranth

native to southwestern US; often hybridizes with *A. hybridus*

Amaranthus retroflexus var. *retroflexus* - redroot amaranth

native to subtropical America; state DOA noxious (CN) weed, 1st collected 1880, Wabasha Co.; highly variable and poisonous to livestock

Amaranthus spinosus L. - spiny amaranth

native to tropical America; reported for Minnesota by FNA but no specimens at MIN

Amaranthus tuberculatus (Moquin-Tandon) J.D. Sauer - roughfruit amaranth

Atriplex argentea var. *argentea* - silverscale saltbush

expected in Minnesota

Atriplex dioica (Nuttall) J.F. Macbride - Suckley's endolepis

reported for Minnesota by FNA but no specimens at MIN; known from Manitoba, ND, SD on saline soils

Atriplex glabriuscula var. *glabriuscula* - Scotland orache

erroneous report

Atriplex heterosperma Bunge - twoscale saltbush

western species reported for Minnesota by the Atlas of the Great Plains but no specimens at MIN

Atriplex hortensis L. - garden orach

native to Asia; known only from Pipestone Co. (one urban population, collected 1925)

Atriplex patula L. - spear saltbush

native to the Mediterranean region

Atriplex prostrata Boucher ex de Candolle - triangle orache

native to Eurasia

Axyris amaranthoides L. - Russian pigweed

native to Siberia

Beta vulgaris subsp. *vulgaris* - sugar beet

native to Europe; known only from Roseau Co. (one riverside population, collected 1939) but doubtfully persisting

Chenopodium album L. - lamb's-quarters

Chenopodium berlandieri var. *bushianum* (Aellen) Cronquist - Bush's goosefoot

Chenopodium berlandieri var. *zschackei* (Murr) Murr ex Asch. - Zschack's goosefoot

Chenopodium capitatum var. *capitatum* (L.) Ambrosi - blite goosefoot

Chenopodium desiccatum A. Nelson - aridland goosefoot

native to western US; known only from Wabasha Co. (collected 1983)

Chenopodium foggii Wahl - Fogg's goosefoot

eastern species erroneously reported for Minnesota by USDA-NRCS; probably misidentified *C. standleyanum* or *C. pratericola*; erroneous record

Chenopodium glaucum var. *glaucum* - oakleaf goosefoot

Chenopodium glaucum var. *salinum* (Standley) B. Boivin - Rocky Mountain goosefoot

Chenopodium leptophyllum (Moquin-Tandon) Nuttall ex S. Watson - narrowleaf goosefoot

often confused with *C. pratericola*

Chenopodium pratericola Rydberg - desert goosefoot

Chenopodium rubrum var. *humile* (Hooker) S. Watson - marshland goosefoot

Chenopodium rubrum var. *rubrum* - red goosefoot

Chenopodium simplex (Torrey) Rafinesque - mapleleaf goosefoot

Chenopodium standleyanum Aellen - Standley's goosefoot

Chenopodium strictum Roth - lateflowering goosefoot

known only from Mille Lacs Co. (collected 1892); native and introduced forms exist but are difficult to distinguish

Chenopodium urbicum L. - city goosefoot

reported for Minnesota by USDA-NRCS but no specimens at MIN

Corispermum americanum var. *americanum* - American bugseed

Corispermum nitidum Kitaibel ex Schultes

misapplied to our flora, see *C. americanum*; misapplied name

Corispermum orientale L.

misapplied to our flora, see *C. villosum*; misapplied name

Corispermum pallasii Steven - Siberian bugseed

Corispermum villosum Rydberg - hairy bugseed

Cycloloma atriplicifolium (Sprengel) J.M. Coulter - winged pigweed

Dysphania ambrosioides (L.) Mosyakin & Clements - Mexican-tea

to be expected in the far southeast corner (known from IA, WI), on drying river beds and lake bottoms

Dysphania botrys (L.) Mosyakin & Clements - Jerusalem oak goosefoot

native to Eurasia

Froelichia floridana (Nuttall) Moquin-Tandon - plains snakecotton

Kochia scoparia subsp. ***scoparia*** - burningbush

native to Eurasia; state DOA noxious (CN) weed, 1st collected 1932 in both Dakota and Hennepin counties

Monolepis nuttalliana (Schultes) Greene - Nuttall's povertyweed

Salicornia rubra A. Nelson - red swampfire

eastern limit of distribution; threatened

Salsola collina Pallas - slender Russian thistle

native to Siberia

Salsola kali L. - prickly Russian-thistle

misapplied to our flora, see *S. tragus*

Salsola tragus L. - prickly Russian-thistle

native to Eurasia; state DOA listed weed, 1st collected 1890, Hennepin Co.

Suaeda calceoliformis (Hooker) Moquin-Tandon - push seepweed

ANACARDIACEAE

Rhus aromatica var. ***aromatica*** - fragrant sumac

native to eastern and southern US

Rhus glabra L. - northern smooth sumac

Rhus typhina L. - staghorn sumac

Rhus x borealis Greene - northern smooth

(*R. glabra* x *typhina*)

Rhus x pulvinata Greene - northern smooth sumac

uncertain status; see *Rhus glabra*

Toxicodendron radicans var. ***negundo*** (Greene) Gillis - eastern poison-ivy

state DOA noxious weed; can cause severe skin irritation

Toxicodendron rydbergii (Small ex Rydberg) Greene - western poison-ivy

state DOA noxious weed; can cause severe skin irritation

Toxicodendron vernix (L.) Kuntze - poison sumac

can cause severe skin irritation

APIACEAE

Aegopodium podagraria L. - bishop's goutweed

native to Eurasia; known to be an aggressive invader in other regions

Aethusa cynapium L. - fool's parsley

native to Eurasia

Anethum graveolens L. - dill

native to southern Europe

Angelica atropurpurea L. - purplestem angelica

Angelica venenosa (Greenway) Fernald - hairy angelica

reported for Minnesota by Gleason & Cronquist 1991 but no specimens at MIN; known from MO, IL, MI in dry woods

Anthriscus sylvestris (L.) Hoffman - wild chervil

native to Europe

Berula erecta (W. Hudson) Coville - cutleaf waterparsnip

Carum carvi L. - caraway

native to Eurasia

Cicuta bulbifera L. - bulblet-bearing water hemlock

Cicuta maculata var. ***angustifolia*** Hook. - spotted water hemlock

Cicuta maculata var. ***bolanderi*** (S. Wats.) G. Mulligan - spotted water hemlock

Cicuta maculata var. ***maculata*** - spotted water hemlock

one of the deadliest poisonous plants in the US, to humans and animals

Conioselinum chinense (L.) Britton, Sterns & Poggenburg - eastern hemlockparsley

reported for Minnesota by Gleason & Cronquist 1991 but no specimens at MIN; known from Ontario, WI, IA in wet meadows and swamps

Conium maculatum L. - poison hemlock

native to Eurasia; plants deadly poisonous to humans and animals (the poison of Socrates); first collected 1992

Coriandrum sativum L. - coriander

Cryptotaenia canadensis (L.) de Candolle - Canadian honewort

Cymopterus acaulis var. ***acaulis*** - plains springparsley

eastern limit of distribution; known only from Clay Co. (one roadside population, collected 1960)

Daucus carota L. - Queen Anne's lace

native to Eurasia; state DOA noxious (CN) weed, 1st collected 1916, Ramsey Co.

Eryngium yuccifolium Michaux - button eryngo

Foeniculum vulgare Mill. - sweet fennel

Heracleum maximum W. Bartram - common cow-parsnip

Levisticum officinale W.D.J. Koch - garden lovage

native to Mediterranean region; reported for Minnesota as a garden escape by UDSA-NRCS but no specimens at MIN

Lomatium orientale J.M. Coulter & Rose - northern Idaho biscuitroot

Oenanthe javanica (Blume) DC. - Java waterdropwort

native to Europe

Osmorrhiza berteroii DC. - sweetcicely

Osmorrhiza claytonii (Michaux) C.B. Clarke - Clayton's sweetroot

Osmorrhiza depauperata R.A. Philippi - bluntseed sweetroot

known only from Cook Co. (last collected 2003); special concern

Osmorrhiza longistylis (Torrey) de Candolle - longstyle sweetroot

Oxypolis rigidior (L.) Rafinesque - stiff cowbane

***Pastinaca sativa* L. - wild parsnip**

native to Eurasia; can cause severe skin irritation

***Pimpinella saxifraga* L. - solidstem burnet saxifrage**

native to Eurasia; known only from Pine Co. (one field population, collected 1982)

***Polytaenia nuttallii* de Candolle - Nuttall's prairie parsley**

known only from Fillmore Co. (last collected 1915); special concern

***Sanicula canadensis* var. *canadensis* - Canadian black snakeroot**

***Sanicula canadensis* var. *grandis* Fernald - Canadian black snakeroot**

***Sanicula gregaria* E.P. Bicknell - clustered black snakeroot**

***Sanicula marilandica* L. - Maryland black snakeroot**

***Sanicula odorata* (Raf.) Pryer & Phillippe - clustered black snakeroot**

***Sanicula trifoliata* E.P. Bicknell - largefruit black snakeroot**

***Sium carsonii* Durand ex A. Gray - Carson's waterparsnip**

erroneous report

***Sium suave* Walter - hemlock waterparsnip**

***Spermolepis inermis* (Nuttall ex de Candolle) Mathias & Constance - Red River scaleseed**

native to southern Great Plains; known only from Fillmore Co. (one trailside population, collected 1994) but doubtfully persisting

***Taenidia integrerrima* (L.) Drude - yellow pimpernel**

***Thaspium barbinode* (Michaux) Nuttall - hairyjoint meadow parsnip**

***Thaspium trifoliatum* var. *aureum* (Nutt.) Britton - purple meadow parsnip**

reported for Minnesota by Gleason & Cronquist 1991 but no specimens at MIN; known from Ontario, WI, IL

***Torilis japonica* de Candolle - erect hedge parsley**

native to Eurasia and northern Africa; known only from Ramsey Co. (one park population, collected 2009)

***Zizia aptera* (A. Gray) Fernald - meadow zizia**

***Zizia aurea* (L.) W.D.J. Koch - golden zizia**

APOCYNACEAE

***Apocynum androsaemifolium* L. - spreading dogbane**

plants poisonous

***Apocynum cannabinum* L. - Indian hemp**

plants poisonous

***Apocynum sibiricum* Jacquin - Indian hemp**

included in *A. cannabinum* by some

***Apocynum x floribundum* Greene - intermediate dogbane**

(*A. androsaemifolium* x *cannabinum*)

***Asclepias amplexicaulis* J.E. Smith - clasping milkweed**

***Asclepias exaltata* L. - poke milkweed**

***Asclepias hirtella* (Pennell) Woodson - green milkweed**

northern limit of distribution; known only from Mower Co. (one population, collected 1982); threatened

***Asclepias incarnata* subsp. *incarnata* - swamp milkweed**

***Asclepias lanuginosa* Nuttall - sidecluster milkweed**

***Asclepias ovalifolia* J. Decaisne - oval-leaf milkweed**

***Asclepias purpurascens* L. - purple milkweed**

known only from Wabasha Co. (historical collection from 1883); rare historic record

***Asclepias quadrifolia* Jacq. - fourleaf milkweed**

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from IA, IL in dry woods

***Asclepias speciosa* Torrey - showy milkweed**

***Asclepias stenophylla* A. Gray - slimleaf milkweed**

***Asclepias sullivantii* Engelmann ex A. Gray - prairie milkweed**

northern limit of distribution; threatened

***Asclepias syriaca* L. - common milkweed**

state DOA noxious (CN) weed

***Asclepias tuberosa* subsp. *interior* Woodson - butterfly milkweed**

***Asclepias verticillata* L. - whorled milkweed**

***Asclepias viridiflora* Rafinesque - green comet milkweed**

***Cynanchum louiseae* Kartesz & Gandhi - Louise**

***Vinca minor* L. - common periwinkle**

native to southern Europe; occasionally escaping from cultivation and persisting but no specimens at MIN

***Vincetoxicum nigrum* (L.) Moench - Louise**

native to Europe; known only from Ramsey Co. (last collected 1997)

AQUIFOLIACEAE

***Ilex mucronata* (L.) M. Powell, Savol. & S. Andrews - catberry**

***Ilex verticillata* (L.) A. Gray - common winterberry**

ARACEAE

***Arisaema dracontium* (L.) Schott - green-dragon**

may cause skin irritation, contains calcium-oxalate crystals; special concern

***Arisaema triphyllum* (L.) Schott - jack in the pulpit**

may cause skin irritation, contains calcium-oxalate crystals; subsp. not formally recognized by FNA since numerous intermediate forms exist with much overlap in character expression

***Calla palustris* L. - water arum**

may cause skin irritation, contains calcium-oxalate crystals

***Lemna minor* L. - common duckweed**

most of our specimens were misidentified and instead are *L. turionifera*

***Lemna obscura* (Austin) Daubs - little duckweed**

northern limit of distribution known only from Washington Co. (one population, collected 1988); DNR watch list

Lemna perpusilla* Torrey - minute duckweed**Lemna trisulca* L. - ivy leaf duckweed*****Lemna turionifera* Landolt - turion duckweed*****Peltandra virginica* (L.) Schott - green arrow arum**

disjunct from central Great Lakes region; known only from Cook Co. (one population, collected 1995)

***Pistia stratiotes* L. - water lettuce**

native to tropics; known only from Winona Co. (collected 2000), doubtfully persisting

Spirodela polyrrhiza* (L.) Schleiden - common duckmeat**Symplocarpus foetidus* (L.) Salisbiry ex W. P. C. Barton - skunk cabbage**

may cause skin irritation, contains calcium-oxalate crystals; roots toxic

Wolffia borealis* (Engelmann) Landolt - northern watermeal**Wolffia brasiliensis* Weddell - northern watermeal**

misapplied; see Wolffia borealis

Wolffia columbiiana* H. Karsten - Columbian watermeal**Wolffia punctata* Griseb. - northern watermeal**

misapplied to our flora, see W. borealis

ARALIACEAE***Aralia hispida* Ventenan - bristly sarsaparilla*****Aralia nudicaulis* L. - wild sarsaparilla*****Aralia racemosa* subsp. *racemosa* - American spikenard*****Hydrocotyle americana* L. - American marsh pennywort*****Hydrocotyle ranunculoides* L. f. - floating marsh pennywort**

possibly introduced, disjunct from eastern OH and TN; known only from Scott Co. (Shakopee-Mdewakanton tribal lands, collected 2007)

***Hydrocotyle umbellata* L. - many-flower marsh pennywort**

reported for Minnesota by Gleason & Cronquist 1991 but no specimens at MIN; known from MI, IL

Panax quinquefolius* L. - American ginseng**Panax trifolius* L. - dwarf ginseng****ARISTOLOCHIACEAE*****Asarum canadense* L. - Canadian wildginger****ASPARAGACEAE*****Asparagus officinalis* L. - common asparagus**

native to Eurasia and north Africa, but evidence suggests that American Indians traded this in pre-European times

***Muscari botryoides* (L.) Mill. - common grape hyacinth**

native to Europe; reported for Minnesota by USDA-NRCS but no specimens at MIN

ASPLENIACEAE***Asplenium platyneuron* (L.) Britton, Stearns & Poggenburg - ebony spleenwort*****Asplenium rhizophyllum* L. - walking fern*****Asplenium trichomanes* subsp. *trichomanes* - maindenhair spleenwort****ASTERACEAE*****Achillea alpina* L. - Chinese yarrow**

southern limit of distribution; known only from Roseau Co. (last collected 1983); threatened

***Achillea millefolium* L. - common yarrow**

both native and introduced forms exist but are not distinguishable except on a molecular level; although much morphological variability occurs subsp. and vars. not recognized, the differences are not considered taxonomically important

***Achillea nobilis* L. - noble yarrow**

native to Europe; apparently introduced with wildlife planting

***Achillea ptarmica* L. - sneezeweed yarrow**

native to Eurasia

***Acropiton repens* (L.) de Candolle - hardheads**

native to Asia; state DOA noxious (CN) weed, known only from Lac Qui Parle Co. (one crop field population, collected 1981)

***Adenocaulon bicolor* Hook. - American trailplant**

erroneous report

***Ageratina altissima* var. *altissima* - white snakeroot**

roots and leaves toxic to humans, deadly poisonous to livestock

Agoseris glauca* var. *glauca* - pale agoseris**Agoseris parviflora* (Nuttall) Greene - false agoseris**

erroneous report

***Agoseris x agrestis* - pale agoseris**

erroneous report

***Ambrosia acanthicarpa* Hooker - flatspine burr ragweed**

native to western US, known only from Hennepin Co. (historical collections to 1896)

***Ambrosia artemisiifolia* L. - annual ragweed**

state DOA noxious (CN) weed; pollen causes allergies

***Ambrosia bidentata* Michaux - lanceleaf ragweed**

known only from Ramsey Co. (one gravel pit population, collected 1914)

Ambrosia psilostachya* de Candolle - cuman ragweed**Ambrosia tomentosa* Nutt. - skeleton-leaf burr ragweed**

to be expected on disturbed wet clays or sands (known from ND, SD, IA, WI)

Ambrosia trifida L. - great ragweed
state DOA noxious (CN) weed; pollen causes allergies

Anaphalis margaritacea (L.) Bentham & Hooker f. - western pearly everlasting

Antennaria howellii subsp. *canadensis* (Greene) R.J. Bayer - Canadian pussytoes

Antennaria howellii subsp. *howellii* - Howell's pussytoes
reported for Minnesota by FNA but no specimens at MIN; known from Ontario, WI, SD and westward

Antennaria howellii subsp. *neodioica* (Greene) R.J. Bayer - Howell's pussytoes

Antennaria howellii subsp. *petaloidea* (Fernald) R.J. Bayer - small pussytoes

Antennaria microphylla Rydberg - littleleaf pussytoes

Antennaria neglecta Greene - field pussytoes

Antennaria parlinii subsp. *fallax* (Greene) R.J. Bayer - Parlin's pussytoes

Antennaria parlinii subsp. *parlinii* - Parlin's pussytoes
known only from Chisago Co. (red-cedar glades; collected 1990)

Antennaria parvifolia Nuttall - small-leaf pussytoes

Antennaria plantaginifolia (L.) Richardson - woman's tobacco

Antennaria rosea subsp. *arida* (E.E. Nelson) R.J. Bayer - rosy pussytoes
erroneous report

Antennaria rosea subsp. *rosea* - rosy pussytoes
reported for Minnesota by FNA but no specimens at MIN; known from Ontario, Manitoba, ND

Anthemis cotula L. - stinking chamomile
native to Eurasia; pollen may cause allergies

Arctium lappa L. - greater burdock
native to Eurasia; reported for Minnesota by FNA but no specimens at MIN

Arctium minus (Hill) Bernhardi - lesser burdock
native to Eurasia; state DOA noxious (CN) weed, 1st collected 1877, Hennepin Co.

Arctium tomentosum Miller - woolly burdock
native to Eurasia; known only from Kittson Co. (last collected 1958)

Arnica lonchophylla Greene - longleaf arnica
southern limit of distribution; vars. Not recognized; threatened

Arnoglossum atriplicifolium (L.) H. Robinson - pale Indian plantain
erroneous report

Arnoglossum plantagineum Rafinesque - groovestem Indian plantain
northern limit of distribution; threatened

Arnoglossum reniforme (Hooker) H. Robinson - great Indian plantain

Artemisia abrotanum L. - southernwood
native to Eurasia and north Africa

Artemisia absinthium L. - absinthium
native to Europe; state DOA noxious (CN) weed, 1st collected 1892, Otter Tail Co.; pollen may cause allergies

Artemisia annua L. - annual wormwood
erroneous report

Artemisia biennis Willdenow - biennial wormwood
native to northwestern US

Artemisia campestris subsp. *borealis* (Pall.) Hall & Clem.
misapplied to the regional flora, see subsp. *caudata*; misapplied name

Artemisia campestris subsp. *caudata* (Michaux) H.M. Hall & Clements - field sagewort

Artemisia cana subsp. *cana* - silver sagebrush
report unverified

Artemisia dracunculus L. - tarragon

Artemisia frigida Willdenow - prairie sagewort

Artemisia longifolia Nuttall - longleaf wormwood
reported for Minnesota by FNA but no specimens at MIN; known from Manitoba, ND, SD on alkaline soils

Artemisia ludoviciana subsp. *ludoviciana* - white sagebrush
pollen may cause allergies

Artemisia pontica L. - Roman wormwood
native to Eurasia; known only from Isanti Co. (one old homestead population, collected 1963)

Artemisia serrata - sawtooth wormwood

Artemisia stelleriana Besser - old woman
native to eastern Asia; known only from St. Louis Co. (Duluth area; last collected 2009)

Artemisia vulgaris L. - common wormwood
native to Eurasia

Bellis perennis L. - lawndaisy
native to Eurasia and Pacific Islands, occasional escape from gardens

Bidens aristosa Britton - bearded beggarticks
report unverified

Bidens beckii Torrey ex Sprengel - Beck's water-marigold

Bidens cernua L. - nodding beggartick

Bidens connata Muhlenberg ex Willdenow - purplestem beggarticks

Bidens discoidea Britton - small beggarticks

Bidens frondosa L. - devil's beggartick

Bidens trichosperma (Michaux) Britton - crowned beggarticks

Bidens tripartita L. - threelobe beggarticks

Bidens vulgata Greene - big devils beggartick

Boltonia asteroides var. *latisquama* (A. Gray) Cronquist - white doll's daisy

Boltonia asteroides var. *recognita* (Fernald & Griscom) Cronquist - white doll's daisy

Brickellia eupatorioides var. *corymbulosa* (Torrey & A. Gray) Shinners - false boneset

Canadanthus modestus (Lindley) G.L. Nesom - **giant mountain aster**
Carduus acanthoides subsp. ***acanthoides*** - **spiny plumeless thistle**
native to Eurasia; state DOA noxious weed, 1st collected 1949, Becker & Rock counties

Carduus crispus L. - **curly plumeless thistle**
native to Eurasia; reported for Minnesota by USDA-NRCS but no specimens at MIN

Carduus nutans L. - **nodding plumeless thistle**
native to Eurasia; state DOA noxious weed, 1st collected 1939, Goodhue Co.

Centaurea cyanus L. - **garden cornflower**
native to Mediterranean region

Centaurea diffusa Lam. - **diffuse knapweed**

Centaurea jacea L. - **brownray knapweed**
native to Europe; known only from St. Louis Co. (one roadside population, collected 1953)

Centaurea montana L. - **perennial cornflower**
native to montane Europe; known only from St. Louis Co. (Duluth area; one population, collected 1952), doubtfully persisting

Centaurea nigrescens Willdenow - **Tyrol knapweed**
erroneous report

Centaurea phrygia L. - **wig knapweed**
native to Europe; known only from St. Louis Co. (last collected 2006)

Centaurea solstitialis L. - **yellow star-thistle**
native to southern Europe; reported for Minnesota by FNA but no specimens at MIN; known from all surrounding states and provinces

Centaurea stoebe L.

Centaurea stoebe subsp. ***micranthos*** (S.G. Gmelin ex Gugler) Hayek - **spotted knapweed**
native to Europe; state DOA noxious (CN) weed, 1st collected 1918, Dakota Co.

Centaurea x moncktonii C.E. Britton - **meadow knapweed**
(C. jacea x nigra x nigrescens); native to Europe; known only from St. Louis Co. (last collected 2006)

Chrysopsis villosa var. ***angustifolia*** (Rydb.) Cronquist
misapplied name; see *Heterotheca stenophylla*

Cichorium intybus L. - **chicory**
native to Eurasia

Cirsium altissimum (L.) Sprengel - **tall thistle**

Cirsium arvense (L.) Scopoli - **Canada thistle**
native to Eurasia; state DOA noxious weed, 1st collected 1878, Hennepin Co.; vars. not recognized

Cirsium discolor (Muhlenberg ex Willdenow) Sprengel - **field thistle**

Cirsium flodmanii (Rydb.) Arthur - **Flodman's thistle**

Cirsium muticum Michaux - **swamp thistle**

Cirsium palustre (L.) Scop. - **marsh thistle**
native to Eurasia; reported for Minnesota (Houston Co.) by U.S. Forest Service but no specimens at MIN

Cirsium pumilum var. ***hillii*** (Canby) B. Boivin - **Hill's thistle**

Cirsium undulatum (Nuttall) Sprengel - **wavyleaf thistle**
reported for Minnesota by FNA but no specimens at MIN; easily confused with *C. flodmanii*

Cirsium vulgare (Savi) Tenore - **bull thistle**
native to Eurasia; state DOA noxious weed, 1st collected 1883, Wabasha Co.

Cirsium x iowense (Pammel) Fernald - **Iowa thistle**
(*C. altissimum* x *discolor*); progeny of this hybrid have low fertility but occur in stable populations; FNA recognizes this taxon but recommends further genetic study before a proper classification can be made

Conyza canadensis (L.) Cronquist - **Canadian horseweed**
vars. not recognized

Conyza ramosissima Cronquist - **dwarf horseweed**

Coreopsis lanceolata L. - **lanceleaf tickseed**
native to eastern US

Coreopsis palmata Nuttall - **stiff tickseed**

Coreopsis tinctoria Nuttall - **golden tickseed**
native to southern US; vars. not recognized

Cosmos bipinnatus Cavendilles - **garden cosmos**
native to Mexico and southwestern U.S.; reported as naturalized for Minnesota by FNA but no specimens at MIN

Cosmos sulphureus Cavendilles - **sulphur cosmos**
native to Mexico; known only from Olmsted Co. (collected 1994), apparently introduced as part of prairie planting but doubtfully persisting

Cota tinctoria (L.) J. Gay ex Gussone - **golden chamomile**
native to Eurasia

Crepis capillaris (L.) Wallroth - **smooth hawksbeard**
native to Europe; state DOA noxious (CN) weed, but not known from Minnesota

Crepis runcinata subsp. ***runcinata*** - **fiddleleaf hawksbeard**

Crepis tectorum L. - **narrowleaf hawksbeard**
native to Europe; state DOA noxious (CN) weed, 1st collected 1934, Becker Co.

Cyclachaena xanthiiifolia (Nuttall) Fresenius - **giant sumpweed**
state DOA noxious (CN) weed

Dieteria canescens (Pursh) Nutt. - **hoary tansyaster**

Doellingeria umbellata var. ***pubens*** (A. Gray) Britton - **parasol whitetop**

Doellingeria umbellata var. ***umbellata*** - **parasol whitetop**

Dyssodia papposa (Venteran) A. Hitchcock - **field marigold**

Echinacea angustifolia de Candolle - **narrow leaved purple coneflower**

Echinacea pallida (Nuttall) Nuttall - **pale purple coneflower**

Echinops exaltatus Schrader - **tall globethistle**
native to Siberia; known only from St. Louis Co. (one population from former agricultural test plots, collected 2004)

Echinops sphaerocephalus - **great globethistle**
native to Eurasia; reported for Minnesota by USDA-NRCS but no specimens at MIN

***Eclipta prostrata* (L.) L. - false daisy**

known only from Washington Co. (one riverside population, collected 1976), probably introduced from further east

***Erechtites hieraciifolius* var. *hieraciifolius* - American burnweed**

***Erigeron acris* L. - bitter fleabane**

Erigeron acris* var. *kamtschaticus (de Candolle) Herder

***Erigeron annuus* (Linnaeus) Persoon - eastern daisy fleabane**

Erigeron glabellus* var. *pubescens Hooker - streamside fleabane

Erigeron lonchophyllum Hooker - shortray fleabane

Erigeron philadelphicus* var. *philadelphicus - Philadelphia fleabane

Erigeron pulchellus* var. *pulchellus

Erigeron pulchellus* var. *tolsteadii Cronquist - Tolstead's erigeron

native to Minnesota; DNR watch list

Erigeron strigosus* var. *septentrionalis (Fernald & Wiegand) Fernald

Erigeron strigosus* var. *strigosus

***Eupatorium altissimum* L.** - tall boneset

***Eupatorium perfoliatum* L.** - common boneset

vars. not recognized

***Eupatorium serotinum* Michaux - lateflowering throughwort**

reported for Minnesota by FNA but no specimens at MIN; known from WI, IA, NE

***Eupatorium sessilifolium* L.** - upland boneset

northwest limit of distribution; threatened

***Eurybia macrophylla* (L.) Cassini - bigleaf aster**

***Euthamia graminifolia* (L.) Nuttall - flat-top goldentop**

vars. not recognized

***Euthamia gymnospermoides* Greene - Texas goldentop**

***Eutrochium maculatum* (L.) E.E. Lamont - spotted joe pye weed**

Eutrochium maculatum* var. *bruneri (A. Gray) E.E. Lamont

Eutrochium maculatum* var. *foliosum (Fernald) E.E. Lamont

known only from Cook Co. (last collected 1945); rare historic record

Eutrochium maculatum* var. *maculatum

***Eutrochium purpureum* (L.) E.E. Lamont - sweet scented joe pye weed**

Eutrochium purpureum* var. *holzingeri (Rydberg) E.E. Lamont - Holzinger's eupatorium

***Eutrochium purpureum* var. *purpureum* - sweetscented joe pye weed**

reported for Minnesota by FNA but no specimens at MIN; known from WI, IA, NE

***Gaillardia aristata* Pursh - common gaillardia**

***Gaillardia pulchella* Fougeroux - Indian blanket**

native to southern US

***Galinsoga parviflora* var. *parviflora* - gallant soldier**

native from southwestern US to South America

***Galinsoga quadriradiata* Ruiz & Pav - shaggy soldier**

native from Central America to South America

***Gnaphalium uliginosum* L.** - marsh cudweed

native to Europe

***Grindelia hirsutula* Hooker & Arnott - hairy gumweed**

***Grindelia squarrosa* (Pursh) Dunal - curlycup gumweed**

state DOA noxious (CN) weed; vars. not recognized

***Gutierrezia sarothrae* (Pursh) Britton & Rusby - broom snakeweed**

reported for Minnesota by FNA but no specimens at MIN; known from Manitoba, ND, SD on rocky soils

***Hasteola suaveolens* (L.) Pojarkova - false Indian plantain**

northwest limit of distribution; endangered

***Helenium autumnale* L.** - common sneezeweed

poisonous to cattle; vars. not recognized

***Helenium flexuosum* Rafinesque - purplehead sneezeweed**

native to eastern US; known only from Carlton Co. (one swampy roadside population, collected 1944)

***Helianthus annuus* L.** - common sunflower

state DOA noxious (CN) weed, named cultivars exempt

***Helianthus giganteus* L.** - giant sunflower

***Helianthus grosseserratus* M. Martens - sawtooth sunflower**

***Helianthus hirsutus* Rafinesque - hairy sunflower**

***Helianthus maximiliani* Schrader - Maximilian's sunflower**

***Helianthus microcephalus* Torrey & A. Gray - small woodland sunflower**

reported for Minnesota by FNA but no specimens at MIN; known from IA, IL

Helianthus nuttallii* subsp. *rydbergii (Britton) R.W. Long - Rydberg's sunflower

Helianthus occidentalis* subsp. *occidentalis - fewleaf sunflower

Helianthus pauciflorus* subsp. *pauciflorus Nutt. - stiff sunflower

Helianthus pauciflorus* subsp. *subrhomboideus (Rydberg) O. Spring & E.E. Schilling - stiff sunflower

Helianthus petiolaris* subsp. *petiolaris - prairie sunflower

***Helianthus strumosus* L.** - paleleaf woodland sunflower

***Helianthus tuberosus* L.** - Jerusalem artichoke

state DOA noxious (CN) weed

Helianthus x laetiflorus Persoon - cheerful sunflower

(*H. pauciflorus* x *tuberous*)

Helianthus helianthoides* var. *scabra (Dunal) Fernald - smooth oxeye

Heterotheca stenophylla* var. *angustifolia (Rydberg) Semple - stiffleaf false goldenaster

known only from Pipestone Co. (collected 1963)

Heterotheca villosa var. *ballardii* (Rydberg) Semple - hairy false goldenaster
Heterotheca villosa var. *foliosa* (Nuttall) V.L. Harms - hairy false goldenaster
known only from Traverse Co. (collected 1938)
Heterotheca villosa var. *minor* (Hooker) Semple - hairy false goldenaster
Heterotheca villosa var. *villosa* - hairy false goldenaster
Hieracium aurantiacum L. - orange hawkweed
native to Europe; state DOA noxious (CN) weed, 1st collected 1947, St. Louis Co.
Hieracium caespitosum Dumortier - meadow hawkweed
native to Europe
Hieracium gronovii L. - queendevil
reported for Minnesota by FNA but no specimens at MIN; known from MI, IN, IL, MO
Hieracium longipilum Torrey ex Hooker - hairy hawkweed
Hieracium pilosella L. - mouse-ear hawkweed
native to Europe
Hieracium piloselloides Villars - tall hawkweed
native to Europe
Hieracium scabrum Michaux - rough hawkweed
Hieracium umbellatum L. - narrowleaf hawkweed
Hieracium vulgatum Fries - common hawkweed
native to Europe; correct name for this taxon is in some doubt fide FNA
Hieracium x fernaldii Lepage
(*H. scabrum* x *umbellatum*); reported for Minnesota by USDA-NRCS but no specimens at MIN; FNA makes no mention of this hybrid
Hieracium x floribundum Wimmer & Grabowski - smooth king devil
(*H. caespitosum* x *lactucella*); native to Europe
Inula britannica L. - British yellowhead
native to Europe; known only from Dakota Co. (nursery holding area, collected 2004)
Inula helenium L. - elecampane
native to Europe
Ionactis linariifolia (L.) Greene - flaxleaf whitetop aster
report unverified
Iva annua L. - annual marsh elder
to be expected on moist disturbed sites; known from ND, SD, IA, WI
Iva axillaris Pursh - povertyweed
reported for Minnesota by FNA but no specimens at MIN; known from WI, Manitoba, ND, SD on wet saline soils
Krigia biflora (Walter) S.F. Blake - twoflower dwarfdandelion
Krigia virginica (L.) Willdenow - Virginia dwarfdandelion
reported for Minnesota by FNA but no specimens at MIN; known from WI, IL, IA
Lactuca biennis (Moench) Fernald - tall blue lettuce
Lactuca canadensis L. - Canada lettuce
Lactuca floridana (L.) Gaertner - woodland lettuce
Lactuca hirsuta Muhlenberg ex Nuttall - hairy lettuce
erroneous report
Lactuca ludoviciana (Nuttall) Ridell - biannual lettuce
Lactuca serriola L. - prickly lettuce
native to Europe; toxic to livestock
Lapsana communis L. - comon nipplewort
native to Eurasia
Leontodon autumnalis L. - fall dandelion
native to Eurasia; known only from St. Louis Co. (one turf-planting population, collected 1995); vars. not recognized
Leucanthemella serotina (L.) Tzvelev - giant daisy
native to eastern Europe
Leucanthemum lacustre (Brot.) Samp. - giant daisy
misapplied to US flora (except CA), see *L. serotina*
Leucanthemum vulgare Lamarck - oxeye daisy
native to Europe; state DOA noxious (CN) weed, 1st collected 1878, Hennepin Co.
Liatris aspera Michaux - tall blazing star
Liatris cylindracea Michaux - Ontario blazing star
Liatris ligulistylis (A. Nelson) K. Schumann - Rocky Mountain blazing star
Liatris punctata var. *punctata* - dotted blazing star
Liatris pycnostachya var. *pycnostachya* NULL - prairie blazing star
Liatris squarrosa var. *glabrata* (Rydb.) Gaiser - scaly blazing star
report unverified
Liatris x frostii Gaiser
(*L. aspera* x *pycnostachya*); reported as endemic to Minnesota by USDA-NRCS but no specimens at MIN
Liatris x sphaeroidea Michaux - spherical blazing star
(*L. aspera* x *ligulistylis*); known only from Marshall Co. (collected 1900)
Logfia arvensis (L.) Holub - field cottonrose
native to Eurasia and northwest Africa; known only from St. Louis Co. (last collected 2006)
Lygodesmia juncea (Pursh) D. Don ex Hooker - rush skeletonplant
Madia glomerata Hooker - mountain tarweed
native to western US
Matricaria chamomilla L. - German chamomile
native to Eurasia; pollen causes allergies
Matricaria discoidea de Candolle - disc mayweed
native to northwestern US
Matricaria maritima L. - scentless false mayweed

misapplied; see *Tripleurospermum inodorum*

***Matricaria matricarioides* (Less.) Porter - disc mayweed**
misapplied; see *M. discoidea*

***Mulgedium pulchellum* (Pursh) G. Don - blue lettuce**
native to Europe; reported for Minnesota by FNA but no specimens at MIN

***Nothocalais cuspidata* (Pursh) Greene - prairie false dandelion**
native to Eurasia; known only from Pipestone Co. (one population, collected 1954), doubtfully persisting

***Onopordum acanthium* var. *acanthium* - Scotch cottontistle**
native to Eurasia; known only from Pipestone Co. (one population, collected 1954), doubtfully persisting

***Packera aurea* (L.) A. Löve & D. Löve - golden ragwort**
leaves and roots possibly toxic to humans, deadly poisonous to horses and cattle

***Packera cana* (Hooker) W.A. Weber & Á. Löve - woolly groundsel**
southeast limit of distribution; known only from Polk Co. (last collected 1993); endangered

***Packera indecora* (Greene) Á. Löve & D. Löve - elegant groundsel**

***Packera pauciflora* (Pursh) Á. Löve & D. Löve - alpine groundsel**
disjunct from western North America and northern Canada; only known specimens at DUL

***Packera paupercula* (Michaux) Á. Löve & D. Löve - balsam groundsel**
leaves and roots possibly toxic to humans, deadly poisonous to horses and cattle

***Packera pseudaurea* var. *semicordata* (Mackenzie & Bush) Trock & T.M. Barkley - falsegold groundsel**

***Parthenium integrifolium* L. - wild quinine**
northwest limit of distribution; endangered

***Petasites frigidus* var. *palmatus* (Aiton) Cronquist - arctic sweet coltsfoot**

***Petasites frigidus* var. *sagittatus* (Banks ex Pursh) Cherniawsky - arrowleaf sweet coltsfoot**

***Petasites frigidus* var. *x vitifolius* (Greene) Cherniawsky - arctic sweet coltsfoot**

***Petasites hybridus* (L.) G. Gaertn., B. Mey. & Scherb. - pestilence wort**

***Polymnia canadensis* L. - whiteflower leafcup**

***Prenanthes alba* L. - white rattlesnakeroot**

***Prenanthes aspera* Michaux - rough rattlesnakeroot**

***Prenanthes crepidinea* Michaux - nodding rattlesnakeroot**
known only from Houston Co. (collected 1900); rare historic record

***Prenanthes racemosa* Michaux - purple rattlesnakeroot**
vars./subsp. not recognized

***Pseudognaphalium macounii* (Greene) Kartesz - clammy cudweed**

***Pseudognaphalium micradenium* (Weatherby) G.L. Nesom - Heller's cudweed**
reported for Minnesota by FNA but no specimens at MIN; known from WI, MI

***Pseudognaphalium obtusifolium* (L.) Hilliard & B.L. Burtt - rabbit-tobacco**

***Pseudognaphalium viscosum* (Kunth) W.A. Weber - clammy cudweed**
misapplied to the regional flora, see *P. macounii*

***Ratibida columnifera* (Nuttall) Wooton & Standley - upright prairie coneflower**

***Ratibida pinnata* (Venterat) Barnhart - pinnate prairie coneflower**

***Rudbeckia hirta* var. *pulcherrima* Farwell - blackeyed Susan**

***Rudbeckia laciniata* var. *laciniata* - cutleaf coneflower**

***Rudbeckia subtomentosa* Pursh - sweet coneflower**
possibly introduced or possibly a range extension from southern IA and WI; known only from Mower Co. (one railroad population, collected 2001); DNR watch list

***Rudbeckia triloba* var. *triloba* - brownneyed Susan**

***Senecio eremophilus* var. *eremophilus* - desert ragwort**
to be expected in far northeast corner (known from adjacent Ontario), on grassy, rocky outcroppings and cliffs

***Senecio integerimus* var. *integerimus* - lambstongue ragwort**
leaves and roots possibly toxic

***Senecio viscosus* L. - sticky ragwort**
native to Eurasia; known only from St. Louis Co. (Duluth area; last collected 2006)

***Senecio vulgaris* L. - old-man-in-the-spring**
native to Eurasia; leaves and roots possibly toxic to humans, deadly poisonous to horses and cattle

***Shinnersoseris rostrata* (A. Gray) Tomb - beaked skeletonweed**
eastern limit of distribution; threatened

***Silphium integrifolium* var. *integrifolium* - wholeleaf rosinweed**
possibly to be expected along southern border (known from SD, IA, WI), in road/railside prairies

***Silphium laciniatum* L. - compass plant**

***Silphium perfoliatum* var. *perfoliatum* - cup plant**

***Solidago altissima* subsp. *altissima* - Canada goldenrod**

***Solidago altissima* subsp. *gilvocanescens* (Rydberg) Semple - Canada goldenrod**

***Solidago canadensis* var. *canadensis* - Canada goldenrod**

***Solidago canadensis* var. *hargeri* Fernald - Canada goldenrod**

***Solidago canadensis* var. *salebrosa* (Piper) M.E. Jones - late goldenrod**
misapplied to our flora, see *S. canadensis* or possibly *S. altissima*

***Solidago flexicaulis* L. - zigzag goldenrod**

***Solidago gigantea* Aiton - giant goldenrod**

***Solidago hispida* Muhlenberg ex Willdenow - hairy goldenrod**
vars. not recognized

***Solidago jejunifolia* Steele - showy goldenrod**

***Solidago juncea* Aiton - early goldenrod**

***Solidago lepida* var. *molina* Fernald**

***Solidago missouriensis* Nutt. - Missouri goldenrod**
vars. not recognized

Solidago mollis Bartling - **velvety goldenrod**
Solidago nemoralis subsp. *decemflora* (de Candolle) Brammall ex Semple - **gray goldenrod**
Solidago nemoralis subsp. *nemoralis* - **gray goldenrod**
Solidago ptarmicoides (Torr. & A. Gray) B. Boivin - **prairie goldenrod**
Solidago riddellii Frank - **Riddell's goldenrod**
Solidago rigida L. - **stiff goldenrod**
Solidago rigida subsp. *humilis* (Porter) S.B. Heard & Semple - **stiff goldenrod**
Solidago rigida subsp. *rigida* - **stiff goldenrod**
Solidago rigidiuscula (Torr. & A. Gray) Porter - **showy goldenrod**
Solidago sciaphila E.S. Steele - **shadowy goldenrod**
Solidago simplex var. *simplex* - **Mt. Albert goldenrod**
 reported for Minnesota by FNA but no specimens at MIN; known from Ontario, Manitoba, ND, SD in rocky seepages
Solidago speciosa var. *speciosa*
Solidago uliginosa Nuttall - **bog goldenrod**
Solidago ulmifolia Muhlenberg ex Willdenow
Solidago ulmifolia var. *ulmifolia* - **elmleaf goldenrod**
Solidago x bernardii B. Boivin - **yellow stiff-aster**
(*S. ptarmicoides* x *riddellii*)
Sonchus arvensis subsp. *arvensis* - **field sowthistle**
Sonchus arvensis subsp. *uliginosus* (M. Bieberstein) Nyman - **field sowthistle**
Sonchus asper (L.) Hill - **spiny sowthistle**
 native to Mediterranean region
Sonchus oleraceus L. - **common sowthistle**
 native to Europe
Symphyotrichum boreale (Torrey & A. Gray) Å. Løve & D. Løve - **northern bog aster**
Symphyotrichum ciliatum (Ledebour) G.L. Nesom - **rayless alkali aster**
 native to western US, Canada, and Siberia
Symphyotrichum ciliolatum - **Lindley's aster**
Symphyotrichum cordifolium - **common blue wood aster**
Symphyotrichum drummondii (Lindley) G.L. Nesom
Symphyotrichum drummondii var. *drummondii* - **Drummond's aster**
Symphyotrichum ericoides (L.) G.L. Nesom - **white heath aster**
Symphyotrichum ericoides var. *ericoides*
Symphyotrichum ericoides var. *pansum* (S.F. Blake) G.L. Nesom - **manyflowered aster**
 known only from Wilkin Co. (one prairie population, collected 1981)
Symphyotrichum falcatum var. *commutatum* (Torrey & A. Gray) G.L. Nesom
Symphyotrichum falcatum var. *falcatum* - **white prairie aster**
 report unverified
Symphyotrichum firmum (Nees) G.L. Nesom - **purplestem aster**
Symphyotrichum laeve (L.) Å. Løve & D. Løve - **smooth blue aster**
Symphyotrichum laeve var. *geyeri* (A. Gray) G.L. Nesom - **Geyer's aster**
 known only from type specimen (as *A. laeviformis*) collected at Lake Benton, Lincoln Co., 1891; rare historic record
Symphyotrichum laeve var. *laeve*
Symphyotrichum lanceolatum (Willd.) G.L. Nesom - **white panicled aster**
Symphyotrichum lanceolatum var. *hesperium* (A. Gray) G.L. Nesom
Symphyotrichum lanceolatum var. *hirsuticaule* (Semple & Chmielewski) G.L. Nesom
 known only from Itasca Co. (collected 2008)
Symphyotrichum lanceolatum var. *interior* (Wiegand) G. L. Nesom
Symphyotrichum lanceolatum var. *lanceolatum* - **white panicle aster**
Symphyotrichum lanceolatum var. *latifolium* (Semple & Chmielewski) G.L. Nesom - **white panicle aster**
 reported for Minnesota by FNA but no specimens at MIN; known from adjacent states and provinces
Symphyotrichum lateriflorum (L.) Å. Løve & D. Løve - **calico aster**
Symphyotrichum novae-angliae (L.) G.L. Nesom - **New England aster**
Symphyotrichum novi-belgii (L.) G.L. Nesom - **New York aster**
Symphyotrichum novi-belgii var. *novi-belgii*
 native to far eastern US; known only from Lake Co. (one population, collected 1952, consists of single poor specimen)
Symphyotrichum oblongifolium (Nuttall) G.L. Nesom - **aromatic aster**
Symphyotrichum ontarionis (Wiegand) G.L. Nesom
Symphyotrichum ontarionis var. *ontarionis* - **bottomland aster**
Symphyotrichum oolentangiense - **skyblue aster**
 vars. not recognized
Symphyotrichum patens var. *patens* - **late purple aster**
 erroneous report
Symphyotrichum pilosum (Willdenow) G.L. Nesom - **hairy white old field aster**
Symphyotrichum pilosum var. *pilosum* - **hairy white oldfield aster**
Symphyotrichum pilosum var. *pringlei* (A. Gray) G.L. Nesom - **Pringle's aster**
 known only from Winona Co. (one park population, collected 1914)
Symphyotrichum praealtum (Poiret) G.L. Nesom - **willowleaf aster**
 several varieties have been proposed but more work is needed to establish their validity
Symphyotrichum prenanthoides (Muhlenberg ex Willdenow) G.L. Nesom - **crookedstem aster**
Symphyotrichum puniceum (L.) Å. Løve & D. Løve
Symphyotrichum puniceum var. *puniceum* - **purplestem aster**
Symphyotrichum robynsonianum (J. Rousseau) Brouillet & Labrecque - **longleaf aster**
 erroneous report

- Symphytum sericeum* (Venterat) G.L. Nesom - western silky aster**
***Symphytum shortii* (Lindley) G.L. Nesom - Short's aster**
 northwest limit of distribution; special concern
- Symphytum urophyllum* (Lindley ex de Candolle) G.L. Nesom - white arrowleaf aster**
- Symphytum x amethystinum* (Nuttall) G.L. Nesom - amethyst aster**
 (S. ericoides x novae-angliae)
- Symphytum x longulum* (E. Sheldon) G.L. Nesom - long aster**
 (S. boreale x puniceum) known only from MN and WI
- Tanacetum parthenium* (L.) Schultz-Bipontinus - feverfew**
 native to Eurasia and north Africa
- Tanacetum vulgare* L. - common tansy**
 native to Eurasia; state DOA noxious (CN) weed, 1st collected 1875, Hennepin Co.; plant oil poisonous
- Taraxacum erythrospermum* Andrzejowski ex Besser - rock dandelion**
 native to Europe
- Taraxacum officinale* F.H. Wiggers - common dandelion**
 native to Europe
- Tephroseris palustris* (L.) Reichenbach - marsh fleabane**
- Thelesperma megapotamicum* (Sprengel) Kuntze - Hopi tea greenthread**
 native to western Great Plains and southwestern US; known only from Sherburne Co., Sherburne National Wildlife Refuge (last collected 2001), introduced with wildlife planting
- Tragopogon dubius* Scopoli - yellow salsify**
 native to Europe
- Tragopogon porrifolius* L. - salsify**
 to be expected on heavily disturbed sites (naturalized across much of North America and known from surrounding states and provinces); native to Europe and north Africa; known to hybridize with both T. dubius and T. pratensis
- Tragopogon pratensis* L. - jack-go-to-bed-at-noon**
 native to Europe
- Tripleurospermum inodorum* (L.) Schultz-Bipontinus - scentless false mayweed**
 native to Europe
- Tripleurospermum maritimum* (L.) W.D.J. Koch**
 misapplied to the regional flora, see T. inodorum; misapplied name
- Tripleurospermum perforatum* (Mérat) M. Lainz - scentless false mayweed**
 misapplied; see T. inodorum
- Tussilago farfara* L. - coltsfoot**
 native to Eurasia; known only from Wright Co. (one agricultural-field population, collected 1913)
- Vernonia baldwinii* Torrey - Baldwin's ironweed**
 native from central to southwestern Great Plains; vars. not recognized; rare historic record
- Vernonia fasciculata* Michaux - prairie ironweed**
 vars. not recognized
- Xanthista spinulosum* var. *spinulosum* - lacy tansyaster**
- Xanthium strumarium* L. - rough cocklebur**
 state DOA noxious (CN) weed, 1st collected 1886; highly toxic to livestock; vars. Not recognized

ATHYRIACEAE

- Diplazium pycnocarpon* (Sprengel) M. Broun - glade fern**
 northwest limit of distribution; threatened

AZOLLACEAE

- Azolla caroliniana* Willdenow - Carolina mosquitofern**
 reported for Minnesota by FNA but no specimens at MIN; widespread in eastern US and more cold tolerant than A. mexicana, from which it differs in megasporangium traits; unfortunately sporocarps are rarely collected but necessary for identification
- Azolla mexicana* C. Presl - Mexican mosquitofern**
 some of our specimens probably A. caroliniana (see comments under Azolla caroliniana); special concern

BALSAMINACEAE

- Impatiens balfourii* Hook.f. - Balfour's touch-me-not**
- Impatiens capensis* Meerburgh - jewelweed**
- Impatiens glandulifera* Royle - ornamental jewelweed**
 native to Himalayan region; known only from the Caribou River in Lake Co. (collected 2008); can become extremely invasive in milder environments

BERBERIDACEAE

- Berberis repens* Lindley - creeping barberry**
 native to western US; reported for Minnesota by MN-DNR and FNA but no specimens at MIN
- Berberis thunbergii* de Candolle - Japanese barberry**
 native to Japan
- Berberis vulgaris* L. - common barberry**
 native to Europe
- Caulophyllum thalictroides* (L.) Michaux - blue cohosh**
- Jeffersonia diphylla* (L.) Persoon - twinleaf**
- Podophyllum peltatum* L. - mayapple**
 roots and leaves poisonous

BETULACEAE

- Alnus glutinosa* (L.) Gaertn. - European alder**
 native to Europe; reported as naturalized in east-central Minnesota by FNA but no specimens at MIN
- Alnus incana* subsp. *rugosa* (Du Roi) R.T. Clausen - speckled alder**

Alnus viridis subsp. *crispa* (Aiton) Turrill - mountain alder

Betula alleghaniensis - yellow birch

Betula cordifolia Regel - mountain paper birch

Betula glandulosa Michaux

misapplied to our flora; see *B. pumila*; misapplied name

Betula nana L. - dwarf birch

erroneous report

Betula nigra L. - river birch

Betula papyrifera Marshall - paper birch

Betula pubescens Ehrhart - downy birch

native to Europe; known only from Houston Co. (one roadside population, collected 1920)

Betula pumila L. - bog birch

Betula x purpurea Schneider - Purpus' birch

(*B. alleghaniensis* x *pumila*)

Betula x rosendahlii Butters & Abbe - Rosendahl's birch

(*B. cordifolia* x *papyrifera*); taxonomic status uncertain, not included in FNA or USDA-NRCS; known only from Cook Co. (last collected 2000)

Betula x sandbergii Britton - Sandberg's birch

(*B. papyrifera* x *pumila*)

Carpinus caroliniana subsp. *virginiana* (Marshall) Furlow - American hornbeam

Corylus americana Walter - American hazelnut

Corylus cornuta subsp. *cornuta* - beaked hazelnut

Ostrya virginiana (Miller) K. Koch - hophornbeam

BIGNONIACEAE

Catalpa ovata G. Don - Chinese catalpa

native to China; known only from Houston Co. (one population, collected 1979, no further data, probably cultivated and not escaping)

Catalpa speciosa (Warder) Warder ex Engelmann - northern catalpa

native to southern US

BORAGINACEAE

Anchusa arvensis (L.) M. von Bieberstein - small bugloss

native to Europe

Asperugo procumbens L. - German-madwort

native to Eurasia; reported for Minnesota by USDA-NRCS but no specimens at MIN

Borago officinalis L. - common borage

native to Europe; doubtfully persisting

Buglossoides arvensis (L.) I.M. Johnst. - corn gromwell

report unverified

Cynoglossum officinale L. - gypsyflower

native to Europe; leaves and roots potentially toxic and may cause skin irritation

Cynoglossum virginianum var. *boreale* (Fernald) Cooperriider - wild comfrey

Echium vulgare L. - common viper's bugloss

native to southern Europe; plant toxic

Ellisia nyctelea (L.) L. - aunt lucy

Eriodictyon angustifolium Nutt. - narrowleaf yerba santa

Hackelia deflexa var. *americana* (A. Gray) gernald & I.M. Johnston - nodding stickseed

Hackelia virginiana (L.) I.M. Johnston - beggarslice

Hydrophyllum appendiculatum Michaux - great waterleaf

Hydrophyllum virginianum var. *virginianum* - eastern waterleaf

Lappula occidentalis var. *occidentalis* - flatspine stickseed

Lappula squarrosa (Retzius) Dumortier - European stickseed

native to Eurasia

Lithospermum canescens (Michaux) J.G.C. Lehmann - hoary puccoon

Lithospermum carolinense var. *croceum* (Fernald) Cusick - Carolina puccoon

Lithospermum incisum J.G.C. Lehmann - narrowleaf stoneseed

Lithospermum latifolium Michaux - American stoneseed

Lithospermum officinale L. - European stoneseed

native to Eurasia; reported for Minnesota by USDA-NRCS but no specimens at MIN

Mertensia paniculata var. *paniculata* - tall bluebells

Mertensia virginica (L.) Persoon ex Link - Virginia bluebells

Myosotis arvensis (L.) Hill - field forget-me-not

native to Eurasia

Myosotis laxa J.G.C. Lehmann - bay forget-me-not

Myosotis scorpioides L. - true forget-me-not

native to Europe

Myosotis sylvatica J.F. Ehrhart ex Hoffmann - woodland forget-me-not

native to Eurasia

Myosotis verna Nuttall - spring forget-me-not

Onosmodium bejariense DC.

Onosmodium bejariense var. *hispidissimum* (Mack.) B.L. Turner - softhair marbleseed

Onosmodium bejariense var. *occidentale* (Mack.) B.L. Turner - western marbleseed

Phacelia franklinii (R. Brown) A. Gray - Franklin's phacelia

Plagiobothrys scouleri var. *hispidulus* (Greene) Dorn - sleeping popcorn flower

Symphytum asperum Lepechin - prickly comfrey

native to Caucasus Mountains; known only from Goodhue Co. (historical collection from 1881)

Symphytum officinale L. - common comfrey
native to Eurasia

BRASSICACEAE

Alliaria petiolata (M. Bieberstein) Cavara & Grande - **garlic mustard**
native to Europe; state DOA noxious weed, 1st collected 1933, Hennepin Co.

Alyssum alyssoides (L.) L. - **pale madwort**
native to Europe

Alyssum desertorum Stapf - **desert madwort**

Arabidopsis lyrata (L.) O

Arabidopsis lyrata subsp. *lyrata* - **lyrate rockcress**

Arabidopsis thaliana (L.) Heynh. - **mouseear cress**

native to Eurasia; reported for Minnesota by USDA-NRCS but no specimens at MIN

Arabis divaricarpa A. Nelson - **spreadingpod rockcress**

misapplied; see Boechera grahamii

Arabis perstellata E.L. Braun - **Short's rockcress**

misapplied; see Boechera dentata

Arabis pycnocarpa M. Hopkins - **hairy rockcress**

Arabis pycnocarpa var. *adpressipilis* M. Hopkins

known only from Waseca Co. (historical collection from 1891), probably a waif, natural distribution is IA to OH and southward

Armoracia rusticana P. Gaertner, B. Meyer & Scherbius - **horseradish**

native to Eurasia; poisonous to livestock

Barbarea orthoceras Ledebour - **American yellowrocket**

Barbarea vulgaris W.T. Aiton - **garden yellowrocket**

native to Eurasia; state DOA noxious (CN) weed, 1st collected 1877, Hennepin Co.; vars. not recognized

Berteroa incana (L.) de Candolle - **hoary alyssum**

native to Europe; state DOA noxious (CN) weed, 1st collected 1894, Ramsey Co.

Boechera canadensis (L.) Al-Shehbaz - **sicklepod**

Boechera collinsii (Fernald) Å. Løve & D. Løve - **Collins' rockcress**

known only from Cook Co. (one cliffside population, collected 1938); rare historic record

Boechera dentata (Rafinesque) Al-Shehbaz & Zarucchi - **Short's rockcress**

Boechera divaricarpa (A. Nelson) Å. Løve & D. Løve

misapplied to our flora, see B. grahamii; misapplied name

Boechera grahamii (Lehmann) Windham & Al-Shehbaz - **spreadingpod rockcress**

Boechera laevigata (Muhlenberg ex Willdenow) Al-Shehbaz - **smooth rockcress**

Boechera missouriensis (Greene) Al-Shehbaz - **green rockcress**

known only from Pine Co. (one roadside population, collected 1980)

Boechera retrofracta (Graham) Å. Løve & D. Løve - **second rockcress**

Boechera stricta (Graham) Al-Shehbaz - **Drummond's rockcress**

Brassica juncea (L.) Czernajew - **brown mustard**

native to Asia; raw seeds somewhat toxic to livestock but cultivated as an oil crop and for condiments

Brassica napus L. - **rape**

native to Mediterranean; known only from Pennington and St. Louis counties (city weeds, last collected 2009), doubtfully persisting; seeds somewhat toxic to livestock; supsp. not recognized

Brassica nigra (L.) W.D.J. Koch - **black mustard**

native to Europe; seeds somewhat toxic to livestock

Brassica oleracea L. - **cabbage**

native to Eurasia; occasional escape from cultivation; cultivars sometimes recognized as vars.

Brassica rapa L. - **field mustard**

native to Europe; raw seeds poisonous but commonly cultivated as a oil crop

Camelina alyssum (Miller) Thellung - **gold-of-pleasure**

native to Europe; known only from Chisago Co. (historical collections from 1885)

Camelina microcarpa de Candolle - **littlepod false flax**

native to Europe

Camelina rumelica Velen. - **graceful false flax**

native to Eurasia; known only from St. Louis Co., doubtfully persisting; only known specimen at DUL

Camelina sativa (L.) Crantz - **gold-of-pleasure**

native to Eurasia

Capsella bursa-pastoris (L.) Medikus - **shepherd's purse**

native to southern Europe

Cardamine bulbosa (Schreber ex Muhlenberg) Britton, Sterns & Poggenburg - **bulbous bittercress**

Cardamine concatenata (Michaux) O. Schwarz - **cutleaf toothwort**

Cardamine diphylla (Michx.) Alph. Wood - **crinkleroot**

reported for Minnesota by USDA-NRCS and FNA but no specimens at MIN; known from Ontario, IL, WI

Cardamine douglassii Britt. - **limestone bittercress**

reported for Minnesota by FNA but no specimens at MIN; known from Ontario, WI, IA

Cardamine flexuosa Withering - **woodland bittercress**

native to Europe; known only from Hennepin Co. (garden weed, collected 2007)

Cardamine impatiens L. - **narrowleaf bittercress**

native to Eurasia; invasive with explosively dehiscent fruits; known only from Ramsey Co. (along Mississippi River in St. Paul, first collected 2008); vars. not recognized

Cardamine parviflora L. - **sand bittercress**

vars. not recognized

Cardamine pensylvanica Muhlenberg ex Willdenow - **Pennsylvania bittercress**

Cardamine pratensis L. - **cuckoo flower**

disjunct from MI, and OH eastward; however, most populations in US introduced from Europe; vars. not recognized; threatened

Chorispora tenella (Pallas) de Candolle - **crossflower**

native to Eurasia; known only from Ramsey Co. (one lawn population, collected 1988), doubtfully persisting

***Conringia orientalis* (L.) Dumortier - hare's ear mustard**

native to Eurasia

***Descurainia incana* (Bernhardi ex Fischer & C.A. Meyer) Dorn - mountain tansymustard**

known only from St. Louis Co. (last collected 1936); rare historic record

***Descurainia pinnata subsp. brachycarpa* (Richardson) Detling - western tansymustard**

***Descurainia sophia* (L.) Webb ex Prantl - herb sophia**

native to Eurasia; state DOA noxious (CN) weed, 1st collected 1932, Dakota Co.

***Diplotaxis muralis* (L.) de Candolle - annual wallrocket**

native to Europe; known only from Martin Co. (one lawn population, collected 1957), doubtfully persisting

***Draba arabisans* Michaux - rock draba**

vars. not recognized; special concern

***Draba cana* Rydberg - cushion draba**

***Draba incana* L. - twisted draba**

reported for Minnesota by FNA but no specimens at MIN; known from Manitoba and Ontario on rock outcrops and gravelly lakeshores

***Draba nemorosa* L. - woodland draba**

vars. not recognized

***Draba norvegica* Gunnerus - Norwegian draba**

disjunct from the high-arctic of Canada; known only from Cook Co. (last collected 1998); vars. not recognized; endangered

***Draba reptans* (Lamarck) Fernald - Carolina draba**

vars. not recognized

***Draba verna* L. - spring draba**

native to Eurasia; known only from St. Louis Co. (Duluth area; one campground population, collected 2001), doubtfully persisting

***Eruca vesicaria* subsp. *sativa* (Mill.) Thell. - rocketsalad**

***Erucastrum gallicum* (Willdenow) O.E. Schultz - common dogmustard**

native to Europe

***Erysimum asperum* (Nuttall) de Candolle - western wallflower**

***Erysimum cheiranthoides* L. - wormseed wallflower**

native to Europe

***Erysimum inconspicuum* (S. Watson) MacMillan - shy wallflower**

***Hesperis matronalis* L. - dames rocket**

native to Eurasia

***Iberis umbellata* L. - globe candytuft**

native to Europe; reported for Minnesota (occasional garden escape near Ely) by Lakela 1965 but no specimens at MIN; doubtfully persisting

***Iodanthus pinnatifidus* (Michaux) Steudel - purple rocket**

northwest limit of distribution; endangered

***Lepidium campestre* (L.) W.T. Aiton - field pepperweed**

native to Eurasia; seeds somewhat toxic to livestock

***Lepidium densiflorum* Schrader - common pepperweed**

seeds somewhat toxic to livestock

***Lepidium draba* L. - whitetop**

native to Mediterranean region

***Lepidium perfoliatum* L. - clasping pepperweed**

native to Eurasia; known only from Clay Co. (one railroad population, collected 1958), doubtfully persisting; seeds somewhat toxic to livestock

***Lepidium ramosissimum* A. Nelson - manybranched pepperweed**

reported for Minnesota by FNA but no specimens at MIN

***Lepidium sativum* L. - gardencress pepperweed**

native to Eurasia; seeds somewhat toxic to livestock

***Lepidium virginicum* subsp. *virginicum* - Virginia pepperweed**

seeds somewhat toxic to livestock

***Lobularia maritima* (L.) Desvaux - sweet alyssum**

native to Mediterranean region

***Lunaria annua* L. - annual honesty**

native to southeastern Europe; known only from Hennepin Co. (one population, collected 1941), doubtfully persisting

***Nasturtium microphyllum* Boenninghausen ex Reichenbach - onerow yellowcress**

native to Europe

***Nasturtium officinale* W.T. Aiton - watercress**

native to Eurasia

***Neslia paniculata* (L.) Desvaux - ballmustard**

native to western Asia

***Physaria ludoviciana* (Nuttall) O'Kane & Al-Shehbaz - foothill bladderpod**

disjunct from Great Plains westward; endangered

***Raphanus raphanistrum* subsp. *raphanistrum* - wild radish**

native to Eurasia; state DOA noxious (CN) weed, 1st collected 1888, Anoka Co.

***Raphanus sativus* L. - cultivated radish**

native to Eurasia; doubtfully persisting

***Rorippa aquatica* (Eaton) E.J. Palmer & Steyermark - lakecress**

known only from Wilkin Co. (one population, collected 1981)

***Rorippa austriaca* (Crantz) Besser - Austrian yellowcress**

native to Europe; known only from Pipestone Co. (one riverside population, collected 1964)

***Rorippa curvipes* Greene - bluntleaf yellowcress**

***Rorippa nasturtium-aquaticum* (L.) Hayek - watercress**

misapplied; see Nasturtium officinale

***Rorippa palustris* subsp. *hispida* (Desvaux) Jonsell - hispid yellowcress**

***Rorippa palustris* subsp. *palustris* - bog yellowcress**

***Rorippa sessiliflora* (Nuttall) Hitchcock - stalkless yellowcress**

known only from historical collections to 1891; special concern

Rorippa sinuata (Nuttall) Hitchcock - **spreading yellowcress**

Rorippa sphaerocarpa (A.Gray) Britton - **roundfruit yellowcress**

no specimens deposited at MIN; perhaps a significant range extension from the southwest and western US; DNR watch list

Rorippa sylvestris (L.) Besser - **creeping yellowcress**

native to Eurasia; DNR watch list

Sinapis alba L. - **white mustard**

native to Europe; known only from Ramsey Co. (one agricultural population, collected 1981), doubtfully persisting

Sinapis arvensis L. - **charlock mustard**

native to Europe; state DOA noxious (CN) weed, 1st collected 1877, Hennepin Co.

Sisymbrium altissimum L. - **tall tumblemustard**

native to Eurasia

Sisymbrium loeselii L. - **small tumbleweed mustard**

native to eastern Mediterranean region

Sisymbrium officinale (L.) Scopoli - **hedgemustard**

native to Mediterranean region

Subularia aquatica subsp. ***americana*** (G.A. Mulligan & Calder) B. Boivin - **American waterawlwort**

Thlaspi arvense L. - **field pennycress**

native to Eurasia; state DOA restricted weed, 1st collected 1883, Goodhue Co.; seeds somewhat toxic to livestock

Turritis glabra L. - **tower rockcress**

BUTOMACEAE

Butomus umbellatus L. - **flowering rush**

native to Eurasia; on MN-DNR Prohibited Invasive Species list; 1st collected 1972, Rice Co.

BUXACEAE

Pachysandra terminalis Siebold & Zuccarini - **Japanese pachysandra**

native to China and Japan; occasionally escaping from cultivation but no specimens at MIN

CABOMBACEAE

Brasenia schreberi J.F. Gmelin - **watershield**

CACTACEAE

Coryphantha missouriensis (Sweet) Britton & Rose - **Missouri foxtail cactus**

erroneous report

Coryphantha vivipara (Nuttall) Britton & Rose - **spinystar**

eastern limit of distribution; endangered

Opuntia fragilis (Nuttall) Haworth - **brittle pricklypear**

Opuntia humifusa (Rafinesque) Rafinesque

Opuntia humifusa var. ***humifusa*** - **devil's-tongue**

Opuntia macrorhiza Engelmann

misapplied to our flora fide FNA, see O. humifusa; some taxonomists (B. Parfitt, pers. comm.) do not believe this is a separate species from O. humifusa but instead represents a color phase but this is unresolved; plants with flower color of O. macrorhiza; misapplied name

CAMPANULACEAE

Campanula aparinoides Pursh - **marsh bellflower**

Campanula cervicaria L. - **bristly bluebells**

native to Europe

Campanula glomerata L. - **Dane's blood**

native to Europe; known only from St. Louis Co. (one population in former agricultural test plots, collected 2004)

Campanula persicifolia L. - **peachleaf bellflower**

native to Asia; known only from St. Louis Co. (Duluth area; collected 2009)

Campanula rapunculoides L. - **rampion bellflower**

native to Eurasia; known to be an aggressive invader in other regions

Campanula rotundifolia L. - **bluebell bellflower**

Campanulastrum americanum (L.) Small - **American bellflower**

Triodanis leptocarpa (Nuttall) Nieuwland - **slimpod Venus' looking-glass**

Triodanis perfoliata (L.) Nieuwland - **clasping Venus' looking-glass**

CANNABACEAE

Cannabis sativa L. - **marijuana**

both native and alien forms occur; state DOA noxious weed; contains intoxicating compounds (federally regulated)

Celtis occidentalis L. - **common hackberry**

Humulus japonicus Siebold & Zuccarini - **Japanese hop**

native to eastern Asia; known only from Fillmore Co. (one riverside population, collected 1992)

Humulus lupulus var. ***lupuloides*** E. Small - **common hop**

Humulus lupulus var. ***lupulus*** - **common hop**

Humulus lupulus var. ***pubescens*** E. Small - **common hop**

report unverified

CAPRIFOLIACEAE

Lonicera caerulea var. ***edulis*** Turczaninow ex Herder - **mountain fly honeysuckle**

native to eastern Siberia; known only from St. Louis Co. (Duluth area; one population collected 2009)

Lonicera caerulea var. ***villosa*** (Michaux) Torrey & A. Gray

Lonicera canadensis W. Bartram ex Marshall - **American fly honeysuckle**

Lonicera dioica L. - **limber honeysuckle**

erroneously called Japanese honeysuckle by some

Lonicera hirsuta Eaton - hairy honeysuckle

Lonicera morrowii A. Gray - Morrow's honeysuckle

native to Japan

Lonicera oblongifolia (Goldie) Hooker - swamp fly honeysuckle

Lonicera reticulata Raf. - grape honeysuckle

Lonicera sempervirens L. - trumpet honeysuckle

Lonicera tatarica L. - Tatarian honeysuckle

native to Eurasia

Lonicera villosa var. *solonis* (Eaton) Fernald - mountain fly honeysuckle

Lonicera villosa var. *tonsa* Fernald - mountain fly honeysuckle

Lonicera x bella Zabel - fly honeysuckle

(*L. morrowii* x *tatarica*); parents native to Eurasia

Lonicera x salicifolia Dieck ex Zabel - willow-leaf honeysuckle

(*L. ruprechtiana* x *xylosteoides*) native to Eurasia; known only from St. Louis Co. (Duluth area; one population collected 2010); only specimen at DUL

Lonicera xylosteum L. - dwarf fly-honeysuckle

native to Europe

Symporicarpos albus (L.) S.F. Blake - common snowberry

Symporicarpos occidentalis Hooker - western snowberry

Symporicarpos orbiculatus Moench - coralberry

Triosteum aurantiacum var. *aurantiacum* - orangefruit horse-gentian

Triosteum perfoliatum L. - feverwort

CARYOPHYLLACEAE

Agrostemma githago var. *githago* - common corncockle

native to Eurasia; (last collected 1912) becoming less common in North America due to herbicide and seed screening protocols; seeds toxic, contain saponin

Arenaria serpyllifolia var. *serpyllifolia* - thymeleaf sandwort

native to Eurasia and Africa

Cerastium arvense subsp. *arvense*

native to western Europe; known only from St. Louis Co. (Duluth area; last collected 2009)

Cerastium arvense subsp. *strictum* Gaudin - field chickweed

Cerastium brachypodium (Engelmann ex A. Gray) B.L. Robinson - shortstalk chickweed

Cerastium fontanum Baumgarten

Cerastium fontanum subsp. *vulgare* (Hartman) Greuter & Burdet - common mouse-ear chickweed

native to Europe

Cerastium nutans var. *nutans* - nodding chickweed

Cerastium velutinum var. *velutinum* - large field mouse ear chickweed

reported for Minnesota by FNA but no specimens at MIN; known from Ontario, IA, MI

Cerastium vulgatum L.

rejected name; see *Cerastium fontanum*

Dianthus armeria subsp. *armeria* - Deptford pink

native to Eurasia

Dianthus barbatus subsp. *barbatus* - sweet william

native to Eurasia; doubtfully persisting

Dianthus deltoides subsp. *deltoides* - maiden pink

native to Europe; doubtfully persisting

Gypsophila elegans M. Bieberstein - showy baby's-breath

native to Eurasia; doubtfully persisting

Gypsophila muralis L. - low baby's-breath

native to Eurasia; doubtfully persisting

Gypsophila paniculata L. - baby's-breath

native to Eurasia

Minuartia dawsonensis (Britton) House - rock stitchwort

Minuartia michauxii (Fenzl) Farw. - Michaux's stitchwort

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from WI, IA, SD; to be expected along the southern border on calcareous gravel and ledges; erroneous record

Minuartia patula (Michx.) Mattfeld - pitcher's stitchwort

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from WI, IL

Moehringia lateriflora (L.) Fenzl - bluntleaf sandwort

Moehringia macrophylla (Hooker) Fenzl - largeleaf sandwort

Myosoton aquaticum (L.) Moench - giant chickweed

native to Eurasia

Paronychia canadensis (L.) Alph. Wood - smooth forked nailwort

northwest limit of distribution; endangered

Paronychia fastigiata var. *fastigiata* - hairy forked nailwort

northwest limit of distribution; known only from Washington Co. (last collected 1981); endangered

Petrorhagia saxifraga var. *saxifraga* - saxifrage pink

native to Eurasia; known only from St. Louis Co. (historical collections to 1940)

Sagina nodosa subsp. *borealis* G.E. Crow - knotty pearlwort

disjunct from arctic Canada; endangered

Sagina procumbens L. - birdeye pearlwort

native to Europe; known only from St. Louis Co. (Duluth area; last collected 2003)

Saponaria officinalis L. - bouncingbet

native to Eurasia; plants used as a liquid soap but toxic as contain saponin

Scleranthus annuus subsp. *annuus* - German knotgrass

native to Eurasia

Silene antirrhina L. - sleepy silene
Silene armeria L. - sweet william silene
native to Europe
Silene chalcedonica (L.) E.H.L. Krause - Maltese cross
native to Asia; although perennial doubtfully persisting more than a few years
Silene csereii Baumgarten - Balkan catchfly
native to southeastern Europe
Silene dichotoma subsp. *dichotoma* Ehrhart - forked catchfly
Silene dioica (L.) Clairville - red catchfly
native to Europe
Silene drummondii subsp. *drummondii* - Drummond's campion
eastern limit of distribution; special concern
Silene flos-cuculi subsp. *flos-cuculi* - ragged robin
native to Europe; known only from Aitkin Co. (one old-field population, collected 2005)
Silene latifolia Poiret - bladder campion
native to Eurasia; state DOA noxious (CN) weed, 1st collected 1878, Martin Co.; subsp. not recognized for North America material
Silene nivea (Nuttall) Muhlenberg ex Otth - evening campion
northwest limit of distribution; threatened
Silene noctiflora L. - nightflowering silene
native to Europe; state DOA noxious weed, 1st collected 1878, Hennepin Co.
Silene stellata (L.) W.T. Aiton - widowstrill
vars. not recognized
Silene vulgaris subsp. *vulgaris* - maidens tears
Spergula arvensis L. - corn spurry
native to Eurasia
Spergularia rubra (L.) J. Presl & C. Presl - red sandspurry
native to Eurasia
Spergularia salina J. Presl & C. Presl - salt sandspurry
to be expected as a widespread weed (known from nearly all states and provinces); native to Eurasia
Stellaria alsine Grimm - bog stitchwort
native to eastern Canada, US
Stellaria borealis subsp. *borealis* - boreal starwort
frequently hybridizes with S. longifolia, though no hybrids reported for Minnesota
Stellaria crassifolia Ehrhart - fleshy starwort
Stellaria graminea L. - grass-like starwort
native to Europe
Stellaria longifolia Muhlenberg ex Willdenow - longleaf starwort
frequently hybridizes with S. borealis, though no hybrids reported for Minnesota
Stellaria longipes subsp. *longipes* - longstalk starwort
Stellaria media (L.) Villars - common chickweed
native to Europe
Stellaria pubera Michx. - star chickweed
erroneous report
Vaccaria hispanica (Miller) Rauschert - cow soapwort
native to Eurasia

CELASTRACEAE

Celastrus orbiculatus Thunberg - Oriental bittersweet
native to Asia; known only from Hennepin Co. (one city-park population, collected 2002)
Celastrus scandens L. - American bittersweet
fruit toxic, possibly other parts
Euonymus alatus (Thunberg) Siebold - burningbush
native to eastern Asia; known only from St. Louis Co. (collected 2007)
Euonymus atropurpureus Jacquin - eastern wahoo
bark, fruits, seeds poisonous
Parnassia glauca Raf. - fen grass of parnassus
Parnassia palustris L. - marsh grass of parnassus
Parnassia parviflora DC. - smallflower grass of Parnassus

CERATOPHYLLACEAE

Ceratophyllum demersum L. - coon's tail
Ceratophyllum echinatum A. Gray - spineless hornwort

CISTACEAE

Crocanthemum bicknellii (Fernald) Janchen - hoary frostweed
Crocanthemum canadense (L.) Britton - longbranch frostweed
Hudsonia tomentosa Nuttall - woolly beachheather
Lechea intermedia var. *intermedia* Legg. ex Britton - largepod pinweed
Lechea stricta Leggett ex Britton - prairie pinweed
Lechea tenuifolia Michaux - narrow leaf pinweed

CLEOMACEAE

Peritoma serrulata (Pursh) de Candolle - Rocky Mountain beechplant
Polanisia dodecandra var. *dodecandra* - redwhisker clammyweed
Polanisia dodecandra var. *trachysperma* (Torrey & A. Gray) H.H. Illis - sandyseed clammyweed

Polanisia jamesii (Torrey & A. Gray) H.H. Iltis - **James' clammyweed**
northern limit of distribution; endangered

COLCHICACEAE

Clintonia borealis (Aiton) Rafinesque - **bluebead**
Uvularia grandiflora Smith - **largeflower bellwort**
Uvularia sessilifolia L. - **sessileleaf bellwort**

COMANDRACEAE

Comandra umbellata subsp. *pallida* (A. DC.) Piehl - **bastard toadflax**
erroneous report
Comandra umbellata subsp. *umbellata*
Geocaulon lividum (Richardson) Fernald - **false toadflax**

COMMELINACEAE

Commelina communis L. - **Asiatic dayflower**
native to Asia
Commelina erecta L. - **whitemouth dayflower**
known only from Wabasha Co. (last collected 1996); endangered
Tradescantia bracteata Small - **longbract spiderwort**
Tradescantia occidentalis var. *occidentalis* - **prairie spiderwort**
Tradescantia ohiensis Rafinesque - **bluejacket**
Tradescantia virginiana L. - **Virginia spiderwort**
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from one disjunct population in WI, otherwise from southeastern IA, IL south- and eastward;
erroneous record

CONVOLVULACEAE

Calystegia macounii (Greene) Brummitt - **Macoun's false bindweed**
Calystegia sepium (L.) R. Brown - **hedge false bindweed**
Calystegia sepium subsp. *americana* (Sims) Brummitt - **hedge false bindweed**
Calystegia sepium subsp. *angulata* Brummitt - **hedge false bindweed**
Calystegia sepium subsp. *appalachiana* Brummitt - **Appalachia false bindweed**
report unverified
Calystegia spithamea (L.) Pursh
Calystegia spithamea subsp. *stans* (Michx.) Brunnitt - **low false bindweed**
Convolvulus arvensis L. - **field bindweed**
native to Europe; state DOA noxious weed, 1st collected 1902, Jackson Co.
Cuscuta cephalanthi Englemann - **buttonbush dodder**
Cuscuta coryli Engelmann - **hazel dodder**
Cuscuta cuspidata Engelm. - **cusp dodder**
Cuscuta glomerata Choisy - **rope dodder**
Cuscuta gronovii Willdenow ex Roemer & Schultes - **scaldweed**
Cuscuta gronovii var. *gronovii*
Cuscuta indecora var. *indecora* - **bigseed alfalfa dodder**
Cuscuta megalocarpa Rydberg - **bigfruit dodder**
known only from St. Louis Co. (one lake-side population, collected 1951); rare historic record
Cuscuta obtusiflora var. *glandulosa* Engelmann - **Peruvian dodder**
mostly found in the southern US & southward; known only from Winona Co. (historical collection from 1897); rare historic record
Cuscuta pentagona Engelmann - **five angle dodder**
Cuscuta pentagona var. *pentagona* - **fiveangled dodder**
Cuscuta polygonorum Engelmann - **smartweed dodder**
Ipomoea cristulata Hallier f. - **Trans-Pecos morning-glory**
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from IA
Ipomoea hederacea Jacquin - **ivyleaf morning-glory**
native to warm temperate Americas; known only from Winona Co. (historical collections from 1886)
Ipomoea purpurea (L.) Roth - **tall morning-glory**
native to tropical Americas; occasionally escaping from cultivation but unlikely to be persisting

CORNACEAE

Cornus alternifolia L. f. - **alternateleaf dogwood**
Cornus amomum Miller - **silky dogwood**
Cornus canadensis L. - **bunchberry dogwood**
Cornus foemina Mill. - **gray dogwood**
see C. racemosa; misapplied name
Cornus obliqua Raf. - **silky dogwood**
Cornus racemosa Lamarck - **gray dogwood**
Cornus rugosa Lamarck - **roundleaf dogwood**
Cornus sericea subsp. *sericea* - **red osier dogwood**

CRASSULACEAE

Crassula aquatica (L.) Schenck - **water pygmyweed**
Hylotelephium telephoides (Michaux) H. Ohba - **Alleghany stonecrop**
native to eastern US; known only from Lake Co. (one Lake Superior island population, collected 1999)
Hylotelephium telephium (L.) H. Ohba - **witch's moneybags**

native to Europe

***Phedimus aizoon* (L.) 't Hart - fei cai**

native to Asia; rarely escapes from cultivation in the Lake Superior area

***Phedimus spurius* (M. Bieberstein) 't Hart - Caucasian stonecrop**

native to the Caucasus Mountains; known only from St. Louis Co. (Duluth area; last collected 2009)

***Rhodiola integrifolia* subsp. *leedyi* (Rosendahl & J.W. Moore) H. Ohba - Leedy's stonecrop**

federal Threatened list / disjunct from western North America; endangered

***Sedum acre* L. - goldmoss stonecrop**

native to Greenland, Eurasia, and north Africa

***Sedum rupestre* L. - Jenny's stonecrop**

native to Europe; known only from St. Louis Co. (Duluth area; one population collected 2001)

CUCURBITACEAE

***Citrullus lanatus* subsp. *lanatus* - watermelon**

***Echinocystis lobata* (Michaux) Torrey & A. Gray - wild cucumber**

***Sicyos angulatus* L. - bur cucumber**

***Thladiantha dubia* Bunge**

native to Asia; reported for Minnesota by USDA-NRCS but based on cultivated specimens

CUPRESSACEAE

***Juniperus communis* var. *depressa* Pursh - common juniper**

***Juniperus horizontalis* Moench - creeping juniper**

***Juniperus virginiana* var. *virginiana* - eastern red-cedar**

***Thuja occidentalis* subsp. *occidentalis* - arborvitae**

CYPERACEAE

***Bolboschoenus fluviatilis* (Torrey) Soj - river bulrush**

***Bolboschoenus maritimus* subsp. *paludosus* (A. Nelson) T. Koyama - cosmopolitan bulrush**

***Bulbostylis capillaris* (L.) C.B. Clarke - densetuft hairsedge**

vars. not recognized

***Carex adusta* Boott - lesser brown sedge**

***Carex aggregata* Mack. - glomerate sedge**

reported for Minnesota by FNA but no specimens at MIN; known from SD, IA, WI

***Carex albicans* Willdenow ex Sprengel - whitetinge sedge**

to be expected in the southeastern corner (known from IA, WI), in open forests on dry acidic soils over sandstone and granite or calcareous soils

***Carex albursina* E. Sheldon - white bear sedge**

***Carex alopecoidea* Tuckerman - foxtail sedge**

***Carex annectens* (E.P. Bicknell) E.P. Bicknell - yellow fruit sedge**

vars. not recognized; special concern

***Carex aquatilis* var. *altior* (Rydb.) Fernald**

misapplied; see *C. aquatilis* var. *substricta*

***Carex aquatilis* var. *aquatilis* - water sedge**

known only from the North Shore (fide G. Wheeler)

***Carex aquatilis* var. *substricta* Kükenthal**

***Carex arcta* Boott - northern cluster sedge**

***Carex arctata* Boott - drooping woodland sedge**

***Carex assiniboinensis* W. Boott - Assiniboa sedge**

***Carex atherodes* Sprengel - wheat sedge**

***Carex aurea* Nuttall - golden sedge**

Gleason & Cronquist (1991) included *C. garberi* in this taxon

***Carex backii* W. Boott - Back's sedge**

Gleason & Cronquist (1991) included *C. saximontana* in this taxon

***Carex bebbii* (L.H. Bailey) Olney ex Fernald - Bebb's sedge**

***Carex bicknellii* Britton - Bicknell's sedge**

***Carex blanda* Dewey - eastern woodland sedge**

***Carex brevior* (Dewey) Mackenzie ex Lunell - shortbeak sedge**

Gleason & Cronquist (1991) included *C. merritt-fernaldii* and *C. molesta* in this taxon

***Carex bromoides* subsp. *bromoides* - brome-like sedge**

Carex brunneoscens* subsp. *brunneoscens

***Carex brunneoscens* subsp. *sphaerostachya* (Tuckerman) Kalela**

***Carex bushii* Mackenzie - Bush's sedge**

known only from Ramsey Co. (one population from a "storm water wetland", collected 1995)

***Carex buxbaumii* Wahlenberg - Buxbaum's sedge**

***Carex canescens* subsp. *canescens* - silvery sedge**

***Carex canescens* subsp. *dis juncta* (Fernald) Toivonen - silvery sedge**

***Carex canescens* var. *suboliacea* Laestad. - Lapland sedge**

misapplied to our flora, see subsp. *canescens*

***Carex capillaris* L. - hair-like sedge**

vars. not recognized; DNR watch list

***Carex careyana* Torrey ex Dewey - Carey's sedge**

northwest limit of distribution; endangered

***Carex castanea* Wahlenberg - chestnut sedge**

***Carex cephaloidea* (Dewey) Dewey - thinleaf sedge**

***Carex cephalophora* Muhlenberg ex Willdenow - oval-leaf sedge**

Carex chordorrhiza Ehrhart ex L. f. - **creeping sedge**
Carex communis var. *communis* - **fibrousroot sedge**
Carex comosa Boott - **long-hair sedge**
Carex conjuncta Boott - **soft fox sedge**
disjunct from southern IA and IL; known only from Rice Co. (last collected 1980) and restored wetland in Ramsey Co. (collected 1998); threatened
Carex conoidea Willdenow - **open field sedge**
this includes *C. katahdinensis*, which was on the state Threatened list
Carex crawei Dewey - **Crawe's sedge**
Carex crawfordii Fernald - **Crawford's sedge**
Carex crinita var. *crinita* - **fringed sedge**
Carex cristatella Britton - **crested sedge**
Carex crus-corvi Shuttleworth - **ravenfoot sedge**
known only from Goodhue Co. (historical collection from 1885); Gleason & Cronquist (1991) included *C. flava* var. *fertilis* in this taxon; rare historic record
Carex cryptolepis Mackenzie - **northeastern sedge**
*Carex davisi*ii Schweinitz & Torrey - **Davis' sedge**
northwest limit of distribution; threatened
Carex debilis var. *rudgei* L.H. Bailey - **white edge sedge**
Carex deflexa var. *deflexa* - **northern sedge**
Carex deweyana var. *deweyana* - **Dewey sedge**
Carex diandra Schrank - **lesser paniced sedge**
Carex disperma Dewey - **softleaf sedge**
Carex divulsa Stokes - **grassland sedge**
erroneous report
Carex duriuscula C.A. Meyer - **needleleaf sedge**
Carex eburnea Boott - **bristleleaf sedge**
Carex echinata Murray
Carex echinata subsp. *echinata* - **star sedge**
Carex echinodes (Fernald) Rothrock, Reznicek & Hipp - **quill sedge**
recently (2009, Systematic Botany 34(2)) elevated from a variety of *C. tenera* to the species level
Carex emoryi Dewey - **Emory's sedge**
Carex exilis Dewey - **coastal sedge**
Carex festucacea Schkuhr ex Willdenow - **fescue sedge**
northwest limit of distribution; known only from Jackson Co. (one population, collected 1982); threatened
Carex filifolia var. *filifolia* - **threadleaf sedge**
Carex flava L. - **yellow sedge**
Carex foenea Willdenow - **dryspike sedge**
(of Gleason & Cronquist)
Carex formosa Dewey - **handsome sedge**
Carex garberi Fernald - **elk sedge**
Carex gracilescens Steudel
misapplied to our flora, see *C. ormostachya*; misapplied name
Carex gracillima Schweinitz - **graceful sedge**
Carex granularis Muhlenberg ex Willdenow - **limestone meadow sedge**
vars. not recognized
Carex gravida L.H. Bailey - **heavy sedge**
Carex grayi J. Carey - **Gray's sedge**
Carex grisea Wahlenberg - **inflated narrow-leaf sedge**
Carex gynandra Schweinitz - **nodding sedge**
Carex gynocrates Wormskjld ex Drejer - **northern bog sedge**
Carex hallii Olney - **deer sedge**
Carex haydenii Dewey - **Hayden's sedge**
Carex hirtifolia Mackenzie - **pubescent sedge**
Carex hitchcockiana Dewey - **Hitchcock's sedge**
Carex hookeriana Dewey - **Hooker's sedge**
native from SD to MT and northward; known only from Becker Co. (one farm-road population, collected 2005, probably introduced); special concern
Carex hystericina Muhlenberg ex Willdenow - **bottlebrush sedge**
Carex inops subsp. *heliophila* (Mackenzie) Crins - **long-stolon sedge**
Carex interior L.H. Bailey - **inland sedge**
Carex intumescens Rudge - **greater bladder sedge**
Carex jamesii Schweinitz - **James' sedge**
northwest limit of distribution; threatened
Carex lacustris Willdenow - **hairy sedge**
Carex laeviconica Dewey - **smoothcone sedge**
Carex laeviginata (K - smoothsheath sedge
northwest limit of distribution; threatened
Carex lanuginosa
misapplied to our flora, see *C. pellita*; misapplied name
Carex lapponica O. Lang - **Lapland sedge**
erroneous report
Carex lasiocarpa subsp. *americana* (Fernald) Hultén - **American woollyfruit sedge**
Carex laxiculmis var. *copulata* (L.H. Bailey) Fernald
Carex laxiculmis var. *laxiculmis* - **spreading sedge**
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from IA and WI
Carex lenticularis Michaux

Carex lenticularis var. *lenticularis* - lakeshore sedge
Carex leptalea Wahlenberg - bristlystalked sedge
Carex leptonervia (Fernald) Fernald - nerveless woodland sedge
Carex limosa L. - mud sedge
Carex livida (Wahlenberg) Willdenow - vivid sedge
Carex lucorum var. *lucorum* - Blue Ridge sedge
 reported for Minnesota by FNA but no specimens at MIN; known from WI and MI; DNR watch list
Carex lupulina Willdenow - hop sedge
Carex lurida Wahlenberg - shallow sedge
 known only from Mille Lacs Co.; only known specimen deposited at NY; DNR watch list
Carex magellanica Lamarck
Carex magellanica subsp. *irrigua* (Wahlenburg) Hiitonen - boreal bog sedge
Carex meadii Dewey - Mead's sedge
Carex media R. Brown ex Richardson - closed-head sedge
Carex merritt-fernaldii Mackenzie - Fernald's sedge
Carex michauxiana Boeckeler - Michaux' sedge
Carex molesta Mackenzie ex Bright - troublesome sedge
Carex muehlenbergii var. *enervis* Boott - Muehlenberg's sedge
 report unverified
Carex muehlenbergii var. *muehlenbergii* - Muehlenberg's sedge
Carex muricata L.
 misapplied to our flora, see C. echinata; misapplied name
Carex muskingumensis Schweinitz - Muskingum sedge
Carex normalis Mackenzie - greater straw sedge
Carex norvegica Retz. - closed-head sedge
 misapplied to our flora, see C. media
Carex novae-angliae Schweinitz - New England sedge
 range extension from WI; known only from Lake Co. (last collected 2006); threatened
Carex obtusata Liljeblad - obtuse sedge
Carex oligocarpa Willdenow - rich woods sedge
Carex oligosperma Michaux - fewseed sedge
Carex ormostachya Wiegand - necklace spike sedge
Carex pallescens L. - pale sedge
 western limit of distribution; endangered
Carex parryana Dewey
 misapplied to our flora, see C. hallii; misapplied name
Carex pauciflora Lightfoot - fewflower sedge
Carex peckii Howe - Peck's sedge
Carex pedunculata Muhlenberg ex Willdenow - longstalk sedge
Carex pellita Willdenow - woolly sedge
 Gleason & Cronquist (1991) included C. lanuginosa in this taxon
Carex pensylvanica Lamarck - Pennsylvania sedge
Carex pinguis L.H. Bailey
Carex plantaginea Lamarck - plantainleaf sedge
 western limit of distribution; endangered
Carex praegracilis W. Boott - clustered field sedge
Carex prairea Dewey - prairie sedge
Carex prasina Wahlenberg - drooping sedge
 to be expected on forested seeps in the St. Croix Valley; known from the WI side in Burnett Co.
Carex praticola Rydberg - meadow sedge
Carex projecta Mackenzie - necklace sedge
Carex pseudocyperus L. - cypress-like sedge
Carex radiata (Wahlenberg) Small - eastern star sedge
Carex retrorsa Schweinitz - knotsheath sedge
Carex richardsonii R. Brown - Richardson's sedge
Carex rosea Schkuhr ex Willdenow - rosy sedge
Carex rossii Boott - Ross's sedge
Carex rostrata Stokes - beaked sedge
 most of our specimens identified as C. rostrata in the past are actually C. utriculata fide G. Wheeler; in Minnesota this species appears restricted to the North Shore and vicinity
Carex sartwellii Dewey - Sartwell's sedge
Carex saximontana Mackenzie - Rocky Mountain sedge
Carex scabrata Schweinitz - eastern rough sedge
 to be expected on forested seeps in the St. Croix Valley; known from the WI side in Burnett Co.
Carex scirpoidea subsp. *scirpoidea* - northern singlespike sedge
Carex scoparia var. *scoparia* - broom sedge
Carex siccata Dewey - dryspike sedge
Carex sparganioides Muhlenberg ex Willdenow - bur-reed sedge
Carex sprengelii Dewey ex Sprengel - Sprengel's sedge
Carex squarrosa L. - squarrose sedge
Carex stenophylla Wahlenb. - needleleaf sedge
 misapplied to our flora, see C. duriuscula; misapplied name
Carex sterilis Willdenow - dioecious sedge
Carex stipata var. *stipata* - awlfruit sedge

Carex stricta Lamarck - **upright sedge**
Carex suberecta (Olney) Britton - **prairie straw sedge**
reported for Minnesota by FNA but no specimens at MIN; known from IA, WI
Carex supina subsp. *spaniocarpa* (Steudel) Hultén - **weak arctic sedge**
southern limit of distribution; known only from Cook Co. (one cliffside population, collected 1936); endangered
Carex sychnocephala J. Carey - **manyhead sedge**
Carex tenera var. *tenera* - **quill sedge**
Carex tenuiflora Wahlenberg - **sparseflower sedge**
Carex tetanica Schkuhr - **rigid sedge**
Carex tincta (Fernald) Fernald - **tinged sedge**
erroneous report
Carex tonsa var. *rugosperma* (Mackenzie) Crins
Carex tonsa var. *tonsa*
range extension from WI; known only from Lake Co. (collected 2003)
Carex torreyi Tuckerman - **Torrey's sedge**
Carex torta Boott in E. Tuckerman - **twisted sedge**
erroneous report
Carex tribuloides var. *tribuloides* - **blunt broom sedge**
Carex trichocarpa Willdenow - **hairyfruit sedge**
Carex trisperma Dewey - **threeseeded sedge**
Carex tuckermanii Dewey - **Tuckerman's sedge**
Carex typhina Michaux - **cattail sedge**
Carex umbellata Schkuhr ex Willdenow - **parasol sedge**
Gleason & Cronquist (1991) included C. tonsa in this taxon
Carex utriculata Boott in W.J. Hooker - **Northwest Territories sedge**
Carex vaginata Tausch - **sheathed sedge**
Carex vesicaria L. - **blister sedge**
Carex viridula subsp. *viridula* - **little green sedge**
Carex vulpinoidea Michaux - **fox sedge**
Carex woodii Dewey - **pretty sedge**
Carex x knieskernii Dewey - **sedge**
(C. arctata x castanea) reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, MI
Carex xerantica L.H. Bailey - **whitescale sedge**
Cladium mariscoides (Muhlenberg) Torrey - **smooth sawgrass**
Cyperus acuminatus Torrey & Hooker - **tapertip flatsedge**
Cyperus bipartitus Torrey - **slender flatsedge**
Cyperus diandrus Torrey - **umbrella flatsedge**
Cyperus erythrorhizos Muhlenberg - **redroot flatsedge**
Cyperus esculentus var. *leptostachyus* Boeckeler - **yellow nutsedge**
native to tropical & subtropical regions but persisting; state DOA noxious (CN) weed, 1st collected 1887, Winona Co.
Cyperus filiculmis Vahl - **fern flatsedge**
native to eastern coastal states; known only from Goodhue Co. (one railroad population, collected 2010)
Cyperus fuscus L. - **brown flatsedge**
native to temperate Eurasia; known only from Itasca Co. (one park population, collected 2007)
Cyperus houghtonii Torrey - **Houghton's flatsedge**
Cyperus lupulinus subsp. *lupulinus* - **Great Plains flatsedge**
Cyperus lupulinus subsp. *macilentus* (Fernald) Marcks - **Great Plains flatsedge**
Cyperus odoratus L. - **fragrant nutsedge**
Cyperus rotundus L. - **nutgrass**
native to subtropical regions; known only from Stearns Co. (historical collection from 1896)
Cyperus schweinitzii Torrey - **Schweinitz's flatsedge**
Cyperus squarrosus L. - **bearded flatsedge**
Cyperus strigosus L. - **strawcolored flatsedge**
Cyperus x mesochorus Geise - **Schweinitz's flatsedge**
(C. lupulinus x schweinitzii)
Dulichium arundinaceum var. *arundinaceum* - **threeway sedge**
Eleocharis acicularis (Leiberg, J.B.) - **needle spikerush**
Eleocharis coloradoensis (Britton) Gilly - **dwarf spikerush**
last collected 1939; special concern
Eleocharis compressa var. *acutisquamata* (Buckley) S.G. Smith - **sharpscale spikerush**
Eleocharis compressa var. *compressa* - **flatstem spikerush**
Eleocharis elliptica Kunth - **elliptic spikerush**
Eleocharis engelmannii Steudel - **Engelmann's spikerush**
Eleocharis erythropoda Steudel - **bald spikerush**
Eleocharis flavescens var. *olivacea* (Torrey) Gleason - **bright green spikerush**
northwest limit of distribution; threatened
Eleocharis intermedia Schultes - **matted spikerush**
Eleocharis macrostachya Britton - **large spike spikerush**
Eleocharis mamillata subsp. *mamillata* - **soft-stem spikerush**
reported for Minnesota by FNA but no specimens at MIN; known from Canada and WI; often mistaken for E. macrostachya or E. palustris
Eleocharis nitida Fernald - **quill spikerush**
Eleocharis obtusa (Willdenow) Schultes - **blunt spikerush**
Eleocharis ovata (Roth) Roemer & Schultes - **ovate spikerush**
Gleason & Cronquist (1991) included E. engelmannii and E. obtusa in this taxon

***Eleocharis palustris* (L.) Roemer & Schultes - common spikerush**
Gleason & Cronquist (1991) included *E. erythropoda*, *E. macrostachya*, and *E. mamillata* in this taxon

***Eleocharis parvula* (Roemer & Schultes) Link ex Bluff Nees, & Schauer**

***Eleocharis quinqueflora* (Hartmann) O. Schwarz - fewflower spikerush**

***Eleocharis robbinsii* Oakes - Robbins' spikerush**
range extension from WI; threatened

***Eleocharis rostellata* (Torrey) Torrey - beaked spikerush**

***Eleocharis tenuis* var. *verrucosa* (Svenson) Svenson - slender spikerush**
erroneous report

***Eleocharis wolfii* (A. Gray) A. Gray ex Britton - Wolf's spikerush**

***Eriophorum angustifolium* Honckeny**

***Eriophorum angustifolium* subsp. *angustifolium* - tall cottongrass**

***Eriophorum chamissonis* C.A. Meyer - Chamisso's cottongrass**
this is part of the circumboreal *E. chamissonis*-*scheuchzeri* complex; in the strict sense (see J. Cayouette, DAO, 2004) this taxon is restricted to AK and BC and ours would then be called *E. russeum* Fries subsp. *leiocarpum* Novoselova

***Eriophorum gracile* W.D.J. Koch ex Roth - slender cottongrass**

***Eriophorum tenellum* Nuttall - fewnerved cottongrass**

***Eriophorum vaginatum* L. - tufted cottongrass**
vars. not recognized

***Eriophorum virginicum* L. - tawny cottongrass**

***Eriophorum viridicarinatum* (Engelmann) Fernald - thinleaf cottonsedge**

***Fimbristylis autumnalis* (L.) Roemer & Schultes - slender fimbry**

***Fimbristylis puberula* var. *interior* (Britton) Kral - hairy fimbry**
range extension from NE; endangered

***Lipocarpha aristulata* (Coville) G.C. Tucker - awned halfchaff sedge**
reported for Minnesota by FNA but no specimens at MIN; known from NE, west- and southward

***Lipocarpha micrantha* (Vahl) G.C. Tucker - smallflower halfchaff sedge**

***Rhynchospora alba* (L.) Vahl - white beaksedge**

***Rhynchospora capillacea* Torrey - needle beaksedge**

***Rhynchospora capitellata* (Michaux) Vahl - brownish beaksedge**
reported for Minnesota by FNA but no specimens at MIN; known from WI (Burnett and Douglas counties); DNR watch list

***Rhynchospora fusca* (L.) W.T. Aiton - brown beaksedge**

***Schoenoplectella purshiana* var. *purshiana* (Fernald) Lye - weakstem bulrush**

***Schoenoplectella smithii* var. *setosa* (Fernald) Shiels & Monfils - Smith's bulrush**

***Schoenoplectella smithii* var. *smithii* (A. Gray) Shiels & Monfils - Smith's bulrush**

***Schoenoplectus acutus* (Muhlenberg ex Bigelow) Å. Løve & D. Løve**

***Schoenoplectus acutus* var. *acutus* - hardstem bulrush**

***Schoenoplectus heterochaetus* (Chase) Soj**

***Schoenoplectus pungens* (Vahl) Palla - common threesquare**

***Schoenoplectus subterminalis* (Torrey) Soj - swaying bulrush**

***Schoenoplectus tabernaemontani* (C.C. Gmelin) Palla - softstem bulrush**

***Schoenoplectus torreyi* (Olney) Palla - Torrey's bulrush**

***Schoenoplectus x kuekenthalianus* (Junge) D.H. Kent -**
erroneous report; no USDA name

***Schoenoplectus x oblongus* (T. Koyama) Soj - hybrid bulrush**
(S. acutus x heterochaetus)

***Scirpus americanus* Pers.**
misapplied; see *Schoenoplectus pungens*

***Scirpus atrocinctus* - blackgirdle bulrush**
hybridizes with *S. cyperinus* and *S. pedicellatus*

***Scirpus atrovirens* Willdenow - green bulrush**
highly variable taxon often confused with closely related *S. georgianus*, *S. hattorianus*, and *S. pallidus*; also hybridizes with *S. pallidus*

***Scirpus cyperinus* - woolgrass**
Gleason & Cronquist (1991) included *S. atrocinctus* and *S. pedicellatus* in this taxon; also hybridizes with *S. pedicellatus*

***Scirpus georgianus* R.M. Harper - Georgia bulrush**
known only from Chisago Co. (one road-ditch population collected 1973); DNR watch list

***Scirpus hattorianus* Makino - mosquito bulrush**

***Scirpus microcarpus* J. Presl & C. Presl - panicled bulrush**

***Scirpus pallidus* (Britton) Fernald - cloaked bulrush**

***Scirpus pedicellatus* Fernald - stalked bulrush**

***Scirpus pendulus* Muhlenberg - rufous bulrush**

***Scleria triglomerata* Michaux - whip nutrush**
northwest limit of distribution; endangered

***Scleria verticillata* Muhlenberg ex Willdenow - low nutrush**
northwest limit of distribution; threatened

***Trichophorum alpinum* (L.) Persoon - alpine bulrush**

***Trichophorum cespitosum* (L.) Schur - tufted bulrush**

***Trichophorum clintonii* (A. Gray) S.G. Smith - Clinton's bulrush**

DENNSTAEDTIACEAE

***Pteridium aquilinum* var. *latiusculum* (Desvaux) L. Underwood ex A. Heller - western bracken fern**
state DOA noxious (CN) weed; moderately toxic to livestock

DIERVILLACEAE

Diervilla lonicera Miller - northern bush honeysuckle

DIOSCOREACEAE

Dioscorea villosa L. - wild yam

DIPSACACEAE

Cephalaria gigantea (Lebed.) Bobr. - Tatarian cephalaria

native to Eurasia; known only from Hennepin Co. (Minneapolis city park; collected 1941)

Dipsacus fullonum L. - fuller's teasel

native to Eurasia; known only from Winona Co. (collected 2004)

Dipsacus laciniatus L. - cutleaf teasel

native to Europe

Knautia arvensis (L.) J.M. Coulter - field scabiosa

native to Europe

DROSERACEAE

Drosera anglica Hudson - English sundew

Drosera intermedia Hayne - spoonleaf sundew

Drosera linearis Goldie - slenderleaf sundew

Drosera rotundifolia L. - roundleaf sundew

Drosera x obovata Mert. & Koch - obovate-leaved sundew

(*D. anglica* x *rotundifolia*) disjunct from Pacific Northwest; known only from Beltrami Co. peatland (collected 1978)

DRYOPTERIDACEAE

Athyrium filix-femina var. *angustum* (Willdenow) G. Lawson - subarctic lady fern

note that the epithet is "filix-femina" not "felix-femina" as often misspelled

Cystopteris bulbifera (L.) Bernhardi - bulblet bladder fern

Cystopteris fragilis (L.) Bernhardi - brittle bladder fern

Cystopteris laurentiana (Weatherby) Blasdell - St. Lawrence bladder fern

Cystopteris protrusa (Wealthery) Blasdell - lowland bladder fern

Cystopteris tennesseensis Shaver - Tennessee bladder fern

Cystopteris tenuis (Michaux) Desvaux - upland brittle bladder fern

Cystopteris x illinoensis R.C. Moran - Illinois bladder fern

Deparia acrostichoides (Swartz) M. Kato - silver false spleenwort

Dryopteris campyloptera (Kunze) Clarkson - mountain wood fern

erroneous report

Dryopteris carthusiana (Villars) H.P. Fuchs - spinulose wood fern

Dryopteris carthusiana x *cristata*

Dryopteris cristata (L.) A. Gray - crested wood fern

Dryopteris expansa (C. Presl) Fraser-Jenkins & Jermy - spreading wood fern

Dryopteris filix-mas (L.) Schott - male fern

report unverified

Dryopteris fragrans (L.) Schott - fragrant wood fern

Dryopteris goldiana (Hooker ex Goldie) A. Gray - Goldie's wood fern

Dryopteris intermedia (Muhlenberg ex Willdenow) A. Gray - intermediate wood fern

Dryopteris marginalis (L.) A. Gray - marginal wood fern

northwest limit of distribution; endangered

Dryopteris x boottii (Tuckerman) L. Underwood

(*D. cristata* x *intermedia*) disposition uncertain (FNA does not recognize names for hybrids involving *D. cristata*); known only from Winona Co. (last collected 1980); uncertain status

Dryopteris x triploidea Wherry - triploid shield fern

(*D. carthusiana* x *intermedia*)

Dryopteris x uliginosa (A. Braun ex Dowell) Druce

(*D. carthusiana* x *cristata*) not *D. uliginosa* (Kunze) C. Christensen; an accepted name for the hybrid (fide ITIS & IPNI) but not included in FNA; uncertain status

Gymnocarpium dryopteris (L.) Newman - western oak fern

Gymnocarpium jessoense subsp. *parvulum* Sarvela - Asian oak fern

Gymnocarpium robertianum (Hoffmann) Newman - scented oak fern

Gymnocarpium x brittonianum (Sarvela) Pryer & Haufler - Britton's oak fern

(*G. disjunctum* x *dryopteris*)

Gymnocarpium x intermedium Sarvela - intermediate oak fern

(*G. jessoense* x *parvulum*)

Matteuccia struthiopteris var. *pensylvanica* (Willdenow) C.V. Morton - ostrich fern

Onoclea sensibilis L. - sensitive fern

Polystichum acrostichoides (Michaux) Schott - christmas fern

northwest limit of distribution; endangered

Polystichum braunii (Spenn) Fe - Braun's holly fern

Polystichum lonchitis (L.) Roth - northern holly fern

reported for Minnesota (northeast corner) by FNA but no specimens at MIN; known from Ontario and WI

Woodsia alpina (Bolton) A. Gray - alpine woodsia

Woodsia glabella R. Brown ex Richardson - smooth woodsia

Woodsia ilvensis (L.) R. Brown - rusty woodsia

Woodsia obtusa subsp. *obtusa* - bluntlobe cliff fern

Woodsia oregana subsp. *cathcartiana* (B.L. Robinson) Windham - Oregon cliff fern

Woodsia scopulina subsp. *laurentiana* Windham - Laurent's cliff fern

known only from Cook Co.; threatened

- Woodsia x abbeae*** Butters - **Abbe's woodsia**
(*W. ilvensis* x *oregana cathcartiana*) sterile hybrid
- Woodsia x gracilis*** (Lawson) Butters - **slender woodsia**
(*W. alpina* x *ilvensis*) sterile hybrid
- Woodsia x maxonii*** R.M. Tryon - **Maxon's woodsia**
(*W. oregana* x *scopulina*) known only from Cook Co. (one population, collected 1947); rare historic record

ELAEAGNACEAE

- Elaeagnus angustifolia*** L. - **Russian-olive**
native to Eurasia
- Elaeagnus commutata*** Bernhardi ex Rudberg - **silverberry**
- Elaeagnus umbellata*** Thunb. - **autumn olive**
native to eastern Asia; reported for Minnesota (Brown Co.) by U.S. Forest Service but no specimens at MIN; known from IA and WI
- Shepherdia argentea*** (Pursh) Nuttall - **silver buffaloberry**
- Shepherdia canadensis* (L.)** Nuttall - **russet buffaloberry**

ELATINACEAE

- Elatine americana*** (Pursh) Arnott - **American waterwort**
- Elatine minima*** (Nuttall) von Fischer & C.A. Meyer - **small waterwort**
- Elatine rubella*** Rydb. - **southwestern waterwort**
- Elatine triandra*** Schkuhr - **threestem waterwort**

EQUISETACEAE

- Equisetum arvense*** L. - **field horsetail**
toxic to livestock, especially horses
- Equisetum fluviatile*** L. - **water horsetail**
- Equisetum hyemale subsp. affine*** (Engelmann) Calder & Roy L. Taylor - **scouringrush horsetail**
toxic to livestock, especially horses; considered a weed by some as can be aggressive
- Equisetum laevigatum*** A. Braun - **smooth horsetail**
- Equisetum palustre*** L. - **marsh horsetail**
toxic to livestock
- Equisetum pratense*** Ehrhart - **meadow horsetail**
- Equisetum scirpoideum*** Michaux - **dwarf scouringrush**
- Equisetum sylvaticum*** L. - **woodland horsetail**
- Equisetum variegatum subsp. variegatum*** - **variegated scouringrush**
- Equisetum x ferrissii*** Clute - **Ferriss' scouring-rush**
(*E. hyemale* x *laevigatum*)
- Equisetum x litorale*** K. - **shallow-water horsetail**
(*E. arvense* x *fluviatile*)
- Equisetum x mackaii*** (Newman) Brichan - **Mack's scouring-rush**
(*E. hyemale* x *variegatum*)
- Equisetum x nelsonii*** (A.A. Eaton) J.H. Schaffner - **Nelson's horsetail**

ERICACEAE

- Andromeda polifolia*** L. - **bog rosemary**
- Andromeda polifolia*** var. *latifolia* Aiton - **bog rosemary**
- Arctostaphylos uva-ursi* (L.)** Sprengel - **kinnikinnick (Chippewa)**
subsp. not recognized; leaves contain toxic compounds
- Chamaedaphne calyculata* (L.)** Moench - **leatherleaf**
- Chimaphila umbellata subsp. umbellata***
- Empetrum atropurpureum*** Fernald & Wiegand - **purple crowberry**
disjunct from eastern Canada; known only from Cook Co. (last collected 1980); endangered
- Empetrum nigrum*** L. - **black crowberry**
known only from Cook Co. (last collected in 1929); var./subsp. not recognized; endangered
- Epigaea repens*** L. - **trailing arbutus**
leaves contain toxic compounds
- Gaultheria hispida* (L.)** Muhlenberg ex Bigelow - **creeping snowberry**
- Gaultheria procumbens* L.** - **eastern teaberry**
leaves contain toxic compounds
- Gaylussacia baccata*** (Wangenheim) K. Koch - **black huckleberry**
- Kalmia polifolia*** Wangenheim - **bog laurel**
- Moneses uniflora* (L.)** A. Gray - **single delight**
- Monotropa hypopitys* L.** - **pinesap**
(note change in spelling)
- Monotropa uniflora* L.** - **Indianpipe**
- Orthilia secunda* (L.)** House - **sidebells wintergreen**
- Pyrola americana*** Sweet - **American wintergreen**
leaves contain toxic compounds
- Pyrola asarifolia subsp. asarifolia*** - **liverleaf wintergreen**
- Pyrola chlorantha*** Swartz - **greenflowered wintergreen**
- Pyrola elliptica*** Nuttall - **waxflower shinleaf**
- Pyrola minor* L.** - **snowline wintergreen**
- Rhododendron groenlandicum*** (Oeder) Kron & Judd - **bog Labrador tea**

Vaccinium angustifolium Aiton - lowbush blueberry
Vaccinium caespitosum Michaux - dwarf bilberry
(note change in spelling)
Vaccinium macrocarpon Aiton - cranberry
Vaccinium membranaceum Douglas ex Torrey - thinleaf blueberry
erroneous report
Vaccinium myrtilloides Michaux - velvetleaf blueberry
Vaccinium oxycoccus L. - small cranberry
Vaccinium pallidum Aiton - Blue Ridge blueberry
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from WI, IL south-and eastward
Vaccinium uliginosum L. - bog blueberry
known only from Cook Co. (last collected 2002); vars. not recognized; endangered
Vaccinium vitis-idaea L. - lingonberry

ERIOCAULACEAE

Eriocaulon aquaticum (Hill) Druce - sevenangle pipewort

EUPHORBIACEAE

Acalypha rhomboidea Rafinesque - common threeseed mercury
Croton glandulosus var. *septentrionalis* Müll. Arg. - vente-conmigo
Euphorbia agraria M. Bieb. - urban spurge
erroneous report
Euphorbia brachycera Engelm. - horned spurge
native to western Great Plains and mountain states; reported for Minnesota by USDA-NRCS but no specimens at MIN; erroneous record
Euphorbia commutata Engelm. ex A. Gray - tinted woodland spurge
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IA
Euphorbia corollata L. - flowering spurge
Euphorbia cyathophora Murray - fire on the mountain
Euphorbia cyparissias L. - cypress spurge
native to Eurasia; may cause severe skin irritation, poisonous to cattle
Euphorbia davidii Subils - David's spurge
Euphorbia dentata var. *dentata* - toothed spurge
Euphorbia esula L. - leafy spurge
misapplied; see E. virgata
Euphorbia geyeri var. *geyeri* - Geyer's sandmat
Euphorbia glyptosperma Engelmann - ribseed sandmat
Euphorbia helioscopia L. - madwoman's milk
native to Europe; known only from Winona Co. (historical collection from 1885)
Euphorbia hexagona Nuttall ex Sprengel - sixangle spurge
Euphorbia humistrata Engelm. ex A. Gray - spreading sandmat
native to southern Great Plains; reported for Minnesota by USDA-NRCS but no specimens at MIN
Euphorbia maculata L. - spotted sandmat
Euphorbia marginata Pursh - snow on the mountain
may cause severe skin irritation
Euphorbia missurica Rafinesque - prairie sandmat
Euphorbia nutans Lagasca y Segura - eyebane
Euphorbia prostrata Aiton - thymeleaf sandmat
Euphorbia serpillifolia Pers. - thymeleaf sandmat
Euphorbia spathulata Lamarck - warty spurge
Euphorbia vermiculata Raf. - wormseed sandmat
Euphorbia virgata Waldst. & Kit. - leafy spurge

FABACEAE

Amorpha canescens Pursh - leadplant
Amorpha fruticosa L. - false indigo bush
Amorpha nana Nuttall - dwarf false indigo
Amphicarpaea bracteata var. *bracteata* - American hogpeanut
Amphicarpaea bracteata var. *comosa* (L.) Fernald - American hogpeanut
Apios americana Medikus - groundnut
Astragalus adsurgens var. *robustior* Hook. - prairie milk vetch
Astragalus agrestis Douglas ex G. Don - purple milkvetch
Astragalus alpinus var. *alpinus* - alpine milkvetch
disjunct from Canada and western mountains of US; first collected 1991; endangered
Astragalus canadensis var. *canadensis* - Canadian milkvetch
Astragalus cicer L. - chickpea milkvetch
native to Europe; known only from St. Louis Co. (one mining road population, collected 2001)
Astragalus crassicarpus var. *crassicarpus* - groundplum milkvetch
Astragalus flexuosus var. *flexuosus* - flexible milkvetch
Astragalus laxmannii var. *robustior* (Hook.) Barneby & S.L. Welsh - prairie milkvetch
Astragalus lotiflorus Hooker - lotus milkvetch
Astragalus missouriensis var. *missouriensis* - Missouri milkvetch
Astragalus neglectus (Torrey & A. Gray) E. Sheldon - Cooper's milkvetch
Astragalus racemosus var. *racemosus* - cream milkvetch
known only from Pipestone Co. (historical collection from 1895); rare historic record

Astragalus tenellus Pursh - looseflower milkvetch

eastern limit of distribution; known only from Otter Tail Co. (historical collections to 1892); rare historic record

Baptisia alba (L.) Vent - long leaf wild indigo***Baptisia alba*** var. ***macrophylla*** (Larisey) Isley - white wild indigo***Baptisia australis*** var. ***australis*** - blue wild indigo

native to eastern and southeastern US; known only from St. Louis Co. (Duluth area; one roadside population, collected 1956), probably garden escapee; roots possibly toxic

Baptisia bracteata var. ***leucophaea*** (Nutt.) Kartesz & Gandhi - longbract wild indigo***Baptisia tinctoria*** (L.) R. Brown ex W.T. Aiton - horseflyweed

native to eastern US and Canada; known only from Lake City area (historical collection from 1882)

Caragana arborescens Lamarck - Siberian peashrub

native to Siberia

Caragana frutex (L.) K.Koch - Russian peashrub***Chamaecrista fasciculata*** (Michaux) Greene - partridge pea***Chamaecrista fasciculata*** var. ***fasciculata*** - partridge pea***Citroria mariana*** L. - Atlantic pigeonwings

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from WI, IA, NE

Coronilla varia L. - crown vetch

native to Mediterranean region

Crotalaria sagittalis L. - arrowhead rattlebox***Dalea candida*** var. ***candida*** - white prairie clover***Dalea candida*** var. ***oligophylla*** (Torrey) Shinners - white prairie clover***Dalea leporina*** (Aiton) Bullock - foxtail prairie clover***Dalea purpurea*** Venterat***Dalea purpurea*** var. ***purpurea*** - purple prairie clover***Dalea villosa*** var. ***villosa*** - silky prairie clover***Desmanthus illinoensis*** (Michaux) MacMillan ex B.L. Robinson & Fernald - Illinois bundleflower***Desmodium canadense*** (L.) DeCandolle - showy ticktrefoil***Desmodium canescens*** (L.) DC. - hoary ticktrefoil

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IA, NE

Desmodium cuspidatum var. ***longifolium*** (Torrey & A. Gray) B.G. Schubert - largebract ticktrefoil***Desmodium glutinosum*** (Muhlenberg ex Willdenow) Wood - pointedleaf ticktrefoil***Desmodium illinoense*** A. Gray - Illinois ticktrefoil***Desmodium nudiflorum*** (L.) de Candolle - nakedflower ticktrefoil***Gleditsia triacanthos*** L. - honeylocust***Glycine max*** (L.) Merrill - soybean

native to China and Japan; occasionally reported along farm roadsides but not persisting

Glycyrrhiza lepidota Pursh - American licorice***Gymnocladus dioicus*** - Kentucky coffeetree

seeds toxic to livestock; special concern

Lathyrus japonicus var. ***maritimus*** (L.) Kartesz & Gandhi - beach pea***Lathyrus latifolius*** L. - perennial pea

native to southern Europe; toxic to animals

Lathyrus ochroleucus Hooker - cream pea***Lathyrus palustris*** L. - marsh pea***Lathyrus sylvestris*** L. - flat pea

native to Europe; reported for Minnesota by USDA-NRCS but no specimens at MIN

Lathyrus tuberosus L. - tuberous sweetpea

native to Eurasia

Lathyrus venosus Muhlenberg ex Willdenow***Lespedeza capitata*** Michx. - roundhead lespedeza***Lespedeza cuneata*** G. Don - sericea lespedeza

native to eastern Asia; known only from Sherburne Co. (nursery holding area, collected 2005)

Lespedeza frutescens (L.) Hornem. - shrubby lespedeza***Lespedeza leptostachya*** Engelmann ex A. Gray - prairie lespedeza

federal Threatened list; northern limit of distribution; threatened

Lespedeza violacea (L.) Pers. - violet lespedeza

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IA, NE

Lespedeza virginica (L.) Britton - slender lespedeza

native to eastern US; known only from Winona Co. (one revegetated-field population, collected 1997)

Lotus corniculatus var. ***corniculatus*** - bird's-foot trefoil

native to Europe

Lotus unifoliolatus var. ***unifoliolatus*** - American bird's-foot trefoil***Lupinus perennis*** var. ***occidentalis*** S. Watson - sundial lupine***Lupinus perennis*** var. ***perennis*** - sundial lupine***Lupinus polyphyllus*** var. ***polyphyllus*** - bigleaf lupine***Medicago lupulina*** L. - black medick

native to Eurasia

Medicago sativa subsp. ***falcata*** (L.) Arcangeli - alfalfa

native to Eurasia

Medicago sativa subsp. ***sativa*** - alfalfa

native to Eurasia

Melilotus albus Medik. - white sweet clover

native to Eurasia; moderately toxic to livestock; considered by some to be a color variant of *M. officinalis*

Melilotus indicus (L.) All. - annual yellow sweetclover

native to Mediterranean region; reported for Minnesota by USDA-NRCS but no specimens at MIN

***Melilotus officinalis* (L.) Lamarck - white sweet clover**

native to Eurasia; moderately toxic to livestock

***Onobrychis viciifolia* Scopoli - sainfoin**

native to southern Europe; known only from Otter Tail Co. (historical collection of 1911; population soon destroyed after discovery)

***Oxytropis borealis* var. *viscida* (Nutt.) S.L. Welsh - viscid locoweed**

***Oxytropis campestris* var. *chartacea* (Fassett) Barneby - field locoweed**

report unverified

***Oxytropis campestris* var. *dispar* (A. Nelson) Barneby - field locoweed**

***Oxytropis lambertii* var. *lambertii* - purple locoweed**

poisonous to livestock

***Oxytropis splendens* Douglas ex Hooker - showy locoweed**

only known specimen from Minnesota ("Chippewa River") deposited at GH; poisonous to livestock

***Oxytropis viscida* Nuttall - viscid locoweed**

disjunct from western North America and arctic regions, known only from Cook Co., (last collected 1997); poisonous to livestock; endangered

***Pediomelum argophyllum* (Pursh) J.W.Grimes - silverleaf Indian breadroot**

***Pediomelum esculentum* (Pursh) Rydberg - large Indian breadroot**

***Psoralidium tenuiflorum* (Pursh) Rydberg - slimflower scurfpea**

***Robinia hispida* var. *hispida* - bristly locust**

native to southeastern US; flowers possibly toxic

***Robinia pseudoacacia* L. - black locust**

native to southeastern US; potentially invasive; toxic, especially to children and horses

***Robinia viscosa* var. *viscosa* - clammy locust**

***Securigera varia* - crown vetch**

***Strophostyles helvola* (L.) Elliott - amberique-bean**

***Strophostyles leiosperma* (Torrey & A. Gray) Piper - slickseed fuzzybean**

***Tephrosia virginiana* (L.) Persoon - Virginia tephrosia**

may cause skin irritation, seeds toxic; special concern

***Trifolium arvense* L. - rabbitfoot clover**

native to Eurasia and northern Africa

***Trifolium aureum* Pollich - golden clover**

native to Eurasia

***Trifolium campestre* Schreber - field clover**

native to Eurasia and northern Africa

***Trifolium dubium* Sibth. - suckling clover**

native to Europe; reported for Minnesota by USDA-NRCS but no specimens at MIN

***Trifolium hybridum* L. - alsike clover**

native to Eurasia

***Trifolium incarnatum* L. - crimson clover**

native to Europe

***Trifolium pratense* L. - red clover**

native to Europe

***Trifolium repens* L. - white clover**

native to Eurasia

***Vicia americana* subsp. *americana* - American vetch**

***Vicia americana* subsp. *minor* (Hook.) Piper & Beattie - American vetch**

***Vicia caroliniana* Walt. - Carolina vetch**

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IL

***Vicia cracca* subsp. *cracca* - bird vetch**

native to Eurasia

***Vicia sativa* subsp. *nigra* (L.) Ehrh. - garden vetch**

***Vicia sativa* subsp. *sativa* - garden vetch**

native to southern Europe

***Vicia tetrasperma* (L.) Schreber - lentil vetch**

native to Eurasia; known only from St. Louis Co. (Duluth area; one population collected 2009)

***Vicia villosa* subsp. *varia* (Host) Corbière - winter vetch**

erroneous report

***Vicia villosa* subsp. *villosa* - winter vetch**

FAGACEAE

***Fagus grandifolia* Ehrhart - American beech**

***Quercus alba* L. - white oak**

***Quercus bicolor* Willdenow - swamp white oak**

***Quercus coccinea* M - scarlet oak**

erroneous report

***Quercus ellipsoidalis* - northern pin oak**

hybrids with Q. rubra especially common in Minnesota

***Quercus macrocarpa* Michaux - bur oak**

***Quercus muehlenbergii* Engelmann - chinkapin oak**

known only from Houston Co. (historical collection from 1899); rare historic record

***Quercus prinoides* Willd. - dwarf chinkapin oak**

erroneous report

***Quercus rubra* L. - northern red oak**

buds, young shoots, and acorns toxic, especially to cattle

***Quercus velutina* Lamarck - black oak**

***Quercus x bebbiana* C.K. Schneider - Bebb's hybrid oak**

(Q. alba x macrocarpa)

***Quercus x hawkinsiae* Sudworth - Hawkins' hybrid oak**

(Q. rubra x velutina) known only from Houston Co. (last collected 1982)

***Quercus x paleolithicola* Trelease - fossil oak**

(Q. ellipsoidalis x velutina) known only from Houston (last collected 1915) and Wabash (collected 2000) counties

***Quercus x schuettei* Trelease - Schuette's hybrid oak**

(Q. bicolor x macrocarpa) known in Minnesota only from urban street trees (historical collections to 1918); this hybrid not named in FNA

FUMARIACEAE

***Adlumia fungosa* (Aiton) Greene ex Britton, Sterns & Poggenberg - Allegheny vine**

***Corydalis aurea var. aurea* - scrambled eggs**

plant potentially toxic

***Corydalis micrantha* (Engelmann ex A. Gray) A. Gray - smallflower fumewort**

***Corydalis sempervirens* (L.) Persoon - rock harlequin**

***Dicentra canadensis* (Goldie) Walpers - squirrel corn**

***Dicentra cucullaria* (L.) Bernhardi - dutchman's breeches**

may cause skin irritation, leaves and roots potentially poisonous

***Fumaria officinalis* L. - drug fumitory**

native to Europe and northern Africa; known only from Winona Co. (one population, collected 1901)

GENTIANACEAE

***Bartonia virginica* (L.) Britton, Sterns & Poggenburg - yellow screwstem**

western limit of distribution; endangered

***Centaurium pulchellum* (Swartz) Druce - branched centaury**

native to Europe; doubtfully persisting annual

***Gentiana affinis* Grisebach - pleated gentian**

***Gentiana alba* Muhl. ex Nutt. - plain gentian**

***Gentiana andrewsii var. andrewsii* - closed bottle gentian**

***Gentiana andrewsii var. dakotica* A. Nelson - Dakota gentian**

***Gentiana flava* A. Gray**

***Gentiana puberulenta* J.S. Pringle - downy gentian**

***Gentiana rubricaulis* Schweinitz - closed gentian**

***Gentiana x billingtonii* Farwell - Billington's gentian**

(G. andrewsii x puberulenta) known only from Dakota Co. (one railroad population, collected 1956)

***Gentiana x curtisiae* J.S. Pringle - Curtis' gentian**

(Gentiana alba x puberulenta)

***Gentiana x grandilacustris* J.S. Pringle - big lake gentian**

(G. andrewsii x rubricaulis) reported as endemic to MN and WI by USDA-NRCS; in MN known only from Clearwater County (Itasca SP, collected 1935)

***Gentiana x pallidocyannea* J.S. Pringle**

(G. alba x andrewsii) reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IA

***Gentianella amarella* subsp. *acuta* (Michaux) J.M. Gillett - autumn dwarf gentian**

***Gentianella quinquefolia* subsp. *occidentalis* (A. Gray) Small - agueweed**

***Gentianopsis crinita* (Froelich) Ma - greater fringed gentian**

***Gentianopsis detonsa* (Rottbøll) Ma - Macoun's gentian**

misapplied name; see Gentianopsis virgata subsp. macounii

***Gentianopsis virgata* subsp. *macounii* (T. Holm) J.S. Pringle - Macoun's gentian**

***Gentianopsis virgata* subsp. *virgata* - lesser fringed gentian**

***Halenia deflexa* var. *deflexa* - American spurred gentian**

GERANIACEAE

***Erodium cicutarium* (L.) L'Héritier ex Aiton - redstem stork's bill**

native to Mediterranean region

***Geranium bicknellii* Britton - Bicknell's cranesbill**

***Geranium carolinianum* var. *carolinianum* - Carolina geranium**

***Geranium carolinianum* var. *sphaerospermum* (Fernald) Breitung - Carolina geranium**

***Geranium maculatum* L. - spotted geranium**

***Geranium pratense* L. - meadow geranium**

native to Europe; known only from St. Louis Co. (last collected 2004)

***Geranium pusillum* L. - small geranium**

native to Europe; reported for Minnesota by USDA-NRCS but no specimens at MIN

***Geranium robertianum* subsp. *robertianum* - Robert geranium**

***Geranium sanguineum* L. - bloody geranium**

native to Europe; only known collection at DUL

***Geranium sibiricum* L. - Siberian geranium**

native to Eurasia

GROSSULARIACEAE

***Ribes americanum* Miller - American black currant**

***Ribes aureum* var. *villosum* de Candolle - golden currant**

considered native in the Great Plains (west of the Mississippi River) but widely spreading elsewhere

***Ribes cynosbati* L. - eastern prickly gooseberry**

***Ribes glandulosum* Grauer - skunk currant**

***Ribes hirtellum* Michaux - hairy stem gooseberry**

***Ribes hudsonianum* Richardson - northern black currant**

vars. not recognized, although in some regions distinctions occur

Ribes lacustre (Persoon) Poirer - prickly currant

Ribes missouriense Nuttall - Missouri gooseberry

Ribes nigrum L. - European black currant

native to Eurasia; known only from Ramsey Co. (one agricultural population, collected 1924)

Ribes oxyacanthoides var. *oxyacanthoides* - Canadian gooseberry

Ribes rubrum L. - cultivated currant

native to Eurasia

Ribes triste Pallas - red currant

Ribes uva-crispa L. - European gooseberry

erroneous report

HALORAGACEAE

Myriophyllum alterniflorum de Candolle - alternate flower watermilfoil

Myriophyllum farwellii Morong - Farwell's watermilfoil

Myriophyllum heterophyllum Michaux - twoleaf watermilfoil

known only from St. Louis Co. (one population, collected 1992); special concern

Myriophyllum humile (Rafinesque) Morong - low watermilfoil

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from WI, IL

Myriophyllum sibiricum Komarov - shortspike watermilfoil

Myriophyllum spicatum L. - Eurasian watermilfoil

native to Eurasia; state DOA noxious weed and MN-DNR Prohibited Invasive Species list; recent research suggests that this is problematical only when forming dense surface mats that shade out native vegetation below

Myriophyllum tenellum Bigelow - slender watermilfoil

Myriophyllum verticillatum L. - whorl-leaf watermilfoil

Proserpinaca palustris var. *crebra* Fernald & Grisc. - marsh mermaidweed

report unverified

HAMAMELIDACEAE

Hamamelis virginiana L. - American witchhazel

HEMEROCALLIDACEAE

Hemerocallis fulva (L.) L. - orange daylily

native to Asia

Hemerocallis lilioasphodelus L. - yellow daylily

native to Asia; reported by Ownbey & Morley 1991 as occasionally escaping from cultivation but no specimens at MIN

HYACINTHEACEAE

Ornithogalum umbellatum L. - sleepy-dick

native to Eurasia and north Africa; rarely escaping cultivation

Scilla siberica Haworth - Siberian squill

native to Russia; rarely escaping cultivation

HYDRANGEACEAE

Philadelphus coronarius L. - sweet mock orange

erroneous report

Philadelphus pubescens var. *pubescens* - hoary mock orange

HYDROCHARITACEAE

Egeria densa Planch. - Brazilian waterweed

native to S. America; unverified report from Hennepin Co.; probably not long-persisting (introduced with aquarium trade)

Elodea bifoliata H. St. John - twoleaf waterweed

Elodea canadensis Michaux - Canadian waterweed

Elodea nuttallii (Planchon) H. St. John - western waterweed

Najas flexilis (Willdenow) Rostkovius & W.L.E. Schmidt - nodding waternymph

Najas gracillima (A. Braun ex Engelmann) Magnus - slender waternymph

Najas guadalupensis subsp. *guadalupensis* - southern waternymph

Najas guadalupensis subsp. *olivacea* (Rosendahl & Butters) R.R. Haynes & Hellquist - Guadalupe waternymph

Najas marina L. - spiny naiad

Najas minor Allioni - brittle waternymph

on MN-DNR Prohibited Invasive Species list; disjunct from IL, IN, MI, probably introduced; known only from Dakota Co. (one population, collected 2003)

Vallisneria americana Michaux - American eelgrass

vars. not recognized

HYPERICACEAE

Hypericum ascyron subsp. *pyramidalatum* (Aiton) N. Robson - great St. Johnswort

Hypericum boreale (Britton) E.P. Bicknell - northern St. Johnswort

Hypericum canadense L. - lesser Canadian St. Johnswort

erroneous report

Hypericum ellipticum Hooker - pale St. Johnswort

Hypericum gentianoides (L.) Britton, Sterns & Poggenb. - orangegrass

reported for Minnesota by Gleason & Cronquist 1991 but no specimens at MIN; known from Ontario, WI, IA

Hypericum kalmianum L. - Kalm's St. Johnswort

disjunct from central and eastern Great Lakes, known only from Lake Co. (one population at wilderness entry point, 2005)

Hypericum maculatum subsp. *obtusiusculum* (Tourlet) Hayek - spotted St. Johnswort

native to Europe; known only from St. Louis Co. (Duluth area; collected 2007)

Hypericum majus (A.Gray) Britton - **large St. Johnswort**

Hypericum mutilem subsp. mutilem L. - **dwarf St. Johnswort**

Hypericum perforatum L. - **common Saint John**

native to Europe

Hypericum perforatum var. *perforatum* L. - **common St. Johnswort**

Hypericum prolificum L. - **shrubby St. Johnswort**

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IA

Hypericum punctatum Lamarck - **spotted St. Johnswort**

Hypericum pyramidatum Aiton - **great St. Johnswort**

Triadenium fraseri (Spach) Gleason - **Fraser's marsh St. Johnswort**

HYPOXIDACEAE

Hypoxis hirsuta (L.) Coville - **common goldstar**

IRIDACEAE

Belamcanda chinensis (L.) DC. - **blackberry lily**

native to eastern Asia; reported by Ownbey & Morley 1991 as escaping cultivation in Waseca Co. but no specimens at MIN

Iris germanica L. - **German iris**

native to Europe; known only from Sherburne Co. (Sherburne National Wildlife Refuge, one hillside population, collected 2001)

Iris missouriensis Nuttall - **Rocky Mountain iris**

erroneous report

Iris pallida Lamarck - **sweet iris**

native to Mediterranean region; known only from Douglas Co. (one roadside population, collected 1940)

Iris pseudacorus L. - **paleyellow iris**

native to Europe

Iris versicolor L. - **harlequin blue flag**

roots poisonous

Iris virginica L. - **Virginia iris**

vars. not recognized

Sisyrinchium angustifolium Miller - **narrowleaf blue-eyed grass**

native of eastern US and Canada; known only from St. Louis Co. (Duluth area; last collected 1950), probably introduced initially with ballast; DNR watch list

Sisyrinchium campestre E.P. Bicknell - **prairie blue-eyed grass**

Sisyrinchium montanum var. *cerebrum* Fernald - **strict blue-eyed grass**

erroneous report

Sisyrinchium montanum var. *montanum* - **strict blue-eyed grass**

Sisyrinchium mucronatum Michaux - **needletip blue-eyed grass**

ISOETACEAE

Isoetes echinospora Durieu - **spiny-spore quillwort**

vars. not recognized

Isoetes lacustris L. - **lake quillwort**

Isoetes melanopoda Gay & Durieu - **blackfoot quillwort**

northern limit of distribution; endangered

Isoetes tenella L. - **spiny-spore quillwort**

misapplied to our flora, see I. echinospora

Isoetes x hickeyi Taylor & Luebke - **quillwort**

JUGLANDACEAE

Carya cordiformis (Wangenheim) K. Koch - **bitternut hickory**

Carya ovata var. *ovata* - **shagbark hickory**

Juglans cinerea L. - **butternut**

Juglans nigra L. - **black walnut**

moderately toxic to dogs and horses (particularly bark/wood shavings for bedding)

JUNCACEAE

Juncus acuminatus Michaux - **tapertip rush**

reported for Minnesota by FNA and MN-DNR but no specimens at MIN; known from IA, WI

Juncus alpinoarticulatus Chaix - **northern green rush**

Juncus anthelatus (Wiegand) R.E. Brooks - **greater poverty rush**

reported for Minnesota by FNA and MN-DNR but no specimens at MIN; known from Ontario, IA, WI

Juncus arcticus var. *arcticus* - **arctic rush**

erroneous report

Juncus arcticus var. *balticus* (Willdenow) Trautvetter - **arctic rush**

Juncus articulatus L. - **jointleaf rush**

previously known only from Ramsey Co. (collected 1922) but recently (1998) collected from Aitkin Co.; endangered

Juncus brachycarpus Engelmann - **whiteroot rush**

disjunct from IL; known only from Blue Earth Co. (collected 1948)

Juncus brachycephalus (Engelmann) Buchenau - **smallhead rush**

erroneous report

Juncus brevicaudatus (Engelmann) Fernald - **narrowpanicle rush**

Juncus bufonius L. - **toad rush**

vars. not recognized

Juncus canadensis J. Gay - **Canadian rush**

Juncus compressus Jacquin - **roundfruit rush**

Juncus dudleyi Wiegand - **Dudley's rush**

Juncus effusus L. - common rush
vars. not recognized

Juncus filiformis L. - thread rush

Juncus gerardii Loiseleur-Deslongchamps - saltmeadow rush

Juncus greenei Oakes & Tuckerman - Greene's rush

Juncus interior Wiegand - inland rush

Juncus longistylis Torrey - longstyle rush

Juncus marginatus Rostkovius - grassleaf rush

Juncus nodosus L. - knotted rush

Juncus pelocarpus E. Meyer - brownfruit rush

Juncus stygius var. *americanus* Buchenau - moor rush

Juncus subtilis E. Meyer - greater creeping rush
disjunct from eastern Canada; known only from Cook Co. (one wilderness population, collected 1998); endangered

Juncus tenuis Willdenow - poverty rush

Juncus torreyi Coville - Torrey's rush

Juncus vaseyi Engelmann - Vasey's rush

Juncus x gracilescens J. Hermann - slender rush
(*J. alpinoarticulatus* x *brevicaudatus*)

Luzula acuminata var. *acuminata* - hairy woodrush

Luzula campestris (L.) DC.
misapplied to our flora, see *L. multiflora*

Luzula luzuloides subsp. *luzuloides* - oakforest woodrush
native to Europe; known only from St. Louis Co. (Duluth area; last collected 1941)

Luzula multiflora (Ehrhart) Lejeune

Luzula multiflora subsp. *multiflora* - common woodrush
native to Eurasia

Luzula parviflora (Ehrhart) Desvaux - smallflowered woodrush
vars. not recognized; threatened

JUNCAGINACEAE

Triglochin maritima L. - seaside arrowgrass

Triglochin palustris L. - marsh arrowgrass

LAMIACEAE

Agastache foeniculum (Pursh) Kuntze - blue giant hyssop

Agastache nepetoides (L.) Kuntze - yellow giant hyssop
known only from Chippewa Co. (one population, collected 1938); rare historic record

Agastache scrophulariifolia (Willdenow) Kuntze - purple giant hyssop

Ajuga genevensis L. - blue bugle
native to Eurasia; known only from Lake and St. Louis counties (last collected 2009)

Blephilia hirsuta (Pursh) Bentham - hairy pagoda-plant

Blephilia subnuda R. Simmers & Kraul - Cumberland pagoda-plant
native to southern US

Chaiturus marrubiastrum (L.) Spenn. - lion's tail

Clinopodium arkansanum (Nutt.) House - Ozark calamint
report unverified

Clinopodium vulgare L. - wild basil

Dracocephalum parviflorum Nuttall - American dragonhead

Elsholtzia ciliata (Thunberg) Hylander - crested latesummer mint
native to Asia; known only from Anoka Co. (one collection, 1959) as escape from birdseed

Galeopsis bifida Boenn. - splitlip hempnettle

Galeopsis tetrahit var. *tetrahit* - brittlestem hempnettle

Glechoma hederacea L. - ground ivy
native to Eurasia; plant oil may be irritating

Hedeoma drummondii Benth. - Drummond's false pennyroyal
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from ND, SD, NE

Hedeoma hispida Pursh - rough false pennyroyal

Hedeoma pulegioides (L.) Persoon - American false pennyroyal
native to eastern and southeastern US; known only from Houston Co. (historical collection from 1899); rare historic record

Hyssopus officinalis L. - hyssop
native to Eurasia; known only from Chisago Co. (one riverbank population, collected 1938)

Lamiastrum galeobdolon (L.) Ehrend. & Polatschek - yellow archangel
native to Europe; known only from St. Louis Co. (Duluth area; collected 2008)

Lamium album L. - snowflake
native to Eurasia; known only from Ramsey Co. (last collected 1973)

Lamium amplexicaule L. - henbit deadnettle
native to Eurasia and northern Africa

Lamium galeobdolon (L.) L.

Lamium maculatum L. - spotted henbit
native to eastern Asia; known only from St. Louis Co. (one cemetery population, collected 2007)

Leonurus cardiaca subsp. *cardiaca* - common motherwort

Leonurus marrubiastrum L. - lion
native to Europe and northern Asia

Leonurus sibiricus L. - honeyweed
native to Asia

Lycopus americanus Muhlenberg ex W.P.C. Barton - American water horehound
Lycopus asper Greene - rough bugleweed
Lycopus uniflorus Michaux - northern bugleweed
Lycopus virginicus L. - Virginia water horehound
Lycopus x sherardii Steele - Sherard's bugleweed
(*L. uniflorus* x *virginicus*)
Marrubium vulgare L. - horehound
native to Eurasia; known only from Wabasha Co. (historical collections to 1890)
Mentha arvensis L. - wild mint
Mentha spicata L. - spearmint
native to Europe
Mentha x gracilis Sole - ginger mint
Mentha x piperita L. - peppermint
(*M. aquatica* x *spicata*) cultigen of Europe; reported for Minnesota by USDA-NRCS but no specimens at MIN
Moluccella laevis L. - shellflower
native to Mediterranean region; known only from Isanti Co. (one homestead population, collected 1963), doubtfully persisting
Monarda didyma L. - scarlet beebealm
native to eastern US
Monarda fistulosa var. *fistulosa* - wild bergamot
Monarda fistulosa var. *menthaefolia* (Graham) Fernald - mintleaf bergamot
Monarda punctata var. *villicaulis* (Pennell) E.J. Palmer & Steyermark - spotted beebealm
Nepeta cataria L. - catnip
native to Eurasia
Nepeta racemosa Lam. - raceme catnip
Origanum vulgare L. - oregano
Perilla frutescens var. *frutescens* - beefsteak plant
Physostegia parviflora Nuttall ex A. Gray - western false dragonhead
erroneous report
Physostegia virginiana subsp. *praemorsa* (Shinners) Cantino
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from NE, IA, IL
Physostegia virginiana subsp. *virginiana* - obedient plant
Prunella vulgaris subsp. *lanceolata* (W.C.P. Barton) Hultén - lance selfheal
Prunella vulgaris subsp. *vulgaris* - common selfheal
native to Europe
Pycnanthemum flexuosum (Walt.) Britton, Sterns & Poggenburg - Appalachian mountainmint
native to southeastern US; reported for Minnesota by MN-DNR and NatureServe but no specimens at MIN
Pycnanthemum tenuifolium Schrad. - narrowleaf mountainmint
report unverified
Pycnanthemum virginianum (L.) T. Durand & B.D. Jacks ex B.L. Robinson & Fernald - Virginia mountain mint
Salvia azurea var. *grandiflora* Benth. - pitcher sage
report unverified
Salvia nutans L. - nodding sage
native to southeastern Europe; known only from Clay Co. (one riverbank population, collected 1937)
Salvia pratensis L.
erroneous report
Salvia reflexa Hornemann - lanceleaf sage
Salvia x sylvestris L. - woodland sage
Satureja arkansana (Nutt.) Briq. - Ozark calamint
Satureja glabella var. *angustifolia* (Torr.) Svenson - Ozark calamint
reported for Minnesota by USDA-NRCS and NatureServe but no specimens at MIN; known from Ontario, WI, IL
Scutellaria galericulata L. - marsh skullcap
Scutellaria lateriflora var. *lateriflora* - blue skullcap
Scutellaria leonardii Epling - Leonard
included in *S. parvula* by some
Scutellaria ovata subsp. *ovata* - heartleaf skullcap
Scutellaria ovata var. *versicolor* (Nuttall) Fernald - heartleaf skullcap
northern limit of distribution; threatened
Scutellaria parvula var. *missouriensis* (Torr.) Goodman & C.A. Lawson - Leonard's skullcap
Scutellaria parvula var. *parvula* - small skullcap
Stachys hispida Pursh - hairy hedge nettle
included in *S. tenuifolia* by some including USDA-NRCS
Stachys palustris L. - marsh hedge-nettle
Stachys palustris var. *pilosa* (Nutt.) Fernald - hairy hedge-nettle
Stachys pilosa var. *arenicola* (Britton) Mulligan & Munro - hairy hedge-nettle
Stachys pilosa var. *pilosa* - hairy hedge-nettle
Stachys tenuifolia Willdenow - smooth hedge-nettle
Teucrium canadense var. *canadense* - Canada germander
Teucrium canadense var. *occidentale* (A. Gray) E.M. McClintock & Epling - western germander
Teucrium canadense var. *virginicum* (L.) Eaton
Thymus praecox subsp. *arcticus* (Durand) Jalas - creeping thyme
native to Eurasia; known only from Lake Co. (lakeshore population collected 1980); only specimen at DUL
Thymus pulegioides L. - lemon thyme
native to Europe
Thymus serpyllum L. - Breckland thyme
erroneous report

Trichostema brachiatum L. - fluxweed

LENTIBULARIACEAE

- Pinguicula vulgaris* L. - common butterwort
Utricularia cornuta Michaux - horned bladderwort
Utricularia geminiscapa L. Benjamin - hiddenfruit bladderwort
range extension from IA and WI; threatened
Utricularia gibba L. - humped bladderwort
Utricularia intermedia Hayne - flatleaf bladderwort
Utricularia macrorhiza Leconte - common bladderwort
Utricularia minor L. - lesser bladderwort
Utricularia ochroleuca R.W. Hartm. - yellowishwhite bladderwort
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario and Manitoba
Utricularia purpurea Walter - eastern purple bladderwort
Utricularia resupinata B.D. Greene ex Bigelow - lavender bladderwort

LILIACEAE

- Aletris farinosa* L. - white colicroot
reported for Minnesota by USDA-NRCS but no specimens at MIN; known east of the Mississippi River
Erythronium albidum Nuttall - white fawnlily
vars. not recognized
Erythronium americanum subsp. *americanum* - dogtooth violet
Erythronium propullans A. Gray - Minnesota fawnlily
federal and endemic to Minnesota; endangered
Lilium lancifolium Thunberg - tiger lily
native to China
Lilium michiganense Farwell - Michigan lily
Lilium philadelphicum L. - wood lily
vars. not recognized
Medeola virginiana L. - Indian-cucumber
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IL
Prosartes trachycarpa S. Watson - roughfruit fairybells
disjunct from western US and Canada; known only from Cook Co. (last collected 2000); endangered
Streptopus amplexifolius (L.) de Candolle - claspleaf twistedstalk
vars. not recognized
Streptopus lanceolatus (Aiton) Reveal - twisted stalk

LIMNANTHACEAE

- Floerkea proserpinacoides* Willdenow - false mermaidweed

LINACEAE

- Linum lewisii* var. *lewisii* - prairie flax
Linum perenne L. - blue flax
native to Europe; known only from Goodhue Co. (Frontenac SP), introduced with roadside plantings (collected 1999)
Linum rigidum var. *rigidum* - stiffstem flax
Linum rigidum var. *simulans* C.M. Rogers - stiffstem flax
Linum sulcatum var. *sulcatum* - grooved flax
Linum usitatissimum L. - common flax
native to Europe; seeds possibly toxic

LINNAEACEAE

- Linnaea borealis* subsp. *americana* (Forbes) Hultén ex Clausen

LOASACEAE

- Mentzelia decapetala* (Pursh) Urban - tenpetal blazingstar
Mentzelia nuda (Pursh) Torrey & A. Gray - bractless blazingstar

LOBELIACEAE

- Lobelia cardinalis* L. - cardinalflower
Lobelia dortmanna L. - Dortmann's cardinalflower
Lobelia inflata L. - Indian-tobacco
plant potentially toxic
Lobelia kalmii L. - Ontario lobelia
Lobelia siphilitica L. - great blue lobelia
leaves and roots potentially toxic
Lobelia siphilitica var. *ludoviciana* A.L.P.P. de Candolle - Ontario lobelia
Lobelia siphilitica var. *siphilitica* - Ontario lobelia
Lobelia spicata var. *campanulata* McVaugh - palespike lobelia
reported by USDA-NRCS for Minnesota but no specimens at MIN; known from Manitoba, Ontario, WI, IA
Lobelia spicata var. *hirtella* A. Gray - palespike lobelia
Lobelia spicata var. *spicata* - palespike lobelia

LYCOPODIACEAE

- Diphasiastrum complanatum* L. - groundcedar

Diphasiastrum digitatum (Dillenius ex A. Braun) Holub - **fan clubmoss**
Diphasiastrum tristachyum (Pursh) Holub - **deeproot clubmoss**
Diphasiastrum x habereri (House) Holub - **Haberer's groundcedar**
(D. digitatum x tristachyum) easily confused with both parents
Diphasiastrum x zeilleri (Rouy) Holub - **Zeiller's groundcedar**
(D. complanatum x tristachyum)
Huperzia appalachiana Beitel & Mickel - **Appalachian clubmoss**
hybridizes frequently with H. selago and rarely (one population, collected 2001 from Cook Co.) with H. lucidula; special concern
Huperzia appressa (Desv.) Å. Löve & D. Löve
misapplied to our flora, see H. appalachiana; misapplied name
Huperzia lucidula (Michaux) Trevisan - **shining clubmoss**
Huperzia porophila (F.E. Lloyd & L. Underwood) Holub - **rock clubmoss**
disjunct from IL and IN; threatened
Huperzia selago (L.) Bernhardi ex Schrank & Martius - **fir clubmoss**
vars./subsp. not recognized
Huperzia x bartleyi (Cusick) Kartesz & Gandhi - **Bartley's clubmoss**
(H. lucidula x porophila) sterile hybrid; range extension from Great Lakes states; known only from Lake Co. (Tettegouche State Park, collected 2001)
Huperzia x buttersii (Abbe) Kartesz & Gandhi - **Butters' fir-moss**
(H. lucidula x selago)
Huperzia x josephbeitelli A. Haines
(H. appalachiana x selago)
Huperzia x protoporophila A. Haines
(H. appalachiana x lucidula)
Lycopodiella inundata (L.) Holub - **inundated clubmoss**
Lycopodium annotinum L. - **stiff clubmoss**
Lycopodium canadense Nessel
Lycopodium clavatum L. - **running clubmoss**
potentially toxic
Lycopodium dendroideum Michaux - **tree groundpine**
Lycopodium hickeyi W.H. Wagner, Beitel & R.C. Moran - **Pennsylvania clubmoss**
Lycopodium lagopus (Laestadius ex C. Hartman) G. Zinserling ex Kuzeneva-Prochorova - **one-cone clubmoss**
Lycopodium obscurum L. - **rare clubmoss**
to be expected in rich hardwoods in the St. Croix Valley; known from adjacent WI eastward

LYTHRACEAE

Ammannia coccinea C.F. Rottb - **valley redstem**
Ammannia robusta Heer & Regel - **grand redstem**
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from all adjacent states
Decodon verticillatus (L.) Elliott
Didiplis diandra (Nuttall ex de Candolle) A.W. Wood - **waterpurslane**
only known specimen from Minnesota (Mower Co.) deposited at MO; rare historic record
Lythrum alatum var. **alatum** - **winged lythrum**
Lythrum salicaria L. - **purple loosestrife**
native to Eurasia; state DOA noxious weed and MN-DNR Prohibited Invasive Species list, 1st collected 1924, Ramsey Co.
Rotala ramosior (L.) Koehne - **lowland rotala**

MALVACEAE

Abelmoschus esculentus (L.) Moench - **okra**
Abutilon theophrasti Medikus - **velvetleaf**
native to southern Asia; state DOA noxious (CN) weed, 1st collected 1876, Hennepin Co.
Alcea rosea L. - **hollyhock**
Althaea officinalis L. - **common marshmallow**
native to Europe; known only from Winona Co. (collected 1994)
Althaea rosea (L.) Cavanilles - **hollyhock**
native to Europe; known only from Fillmore Co. (one parking-lot population, collected 1982), doubtfully persisting
Callirhoe involucrata var. **involucrata** - **purple poppymallow**
report unverified
Callirhoe triangulata (Leavenworth) A. Gray - **clustered poppymallow**
native from WI, IA and southward; known only from Wabasha Co. (one population, collected 2008)
Hibiscus esculentus L. - **okra**
native to Asian tropics; known only from Sherburne Co., Sherburne National Wildlife Refuge (collected 1972), doubtfully persisting
Hibiscus laevis Allioni - **halberdleaf rosemallow**
range extension from IA; known only from Ramsey Co. (one railroad population, collected 2000)
Hibiscus trionum L. - **flower of an hour**
native to southern Europe; state DOA noxious (CN) weed, 1st collected 1878, Hennepin Co.
Lavatera thuringiaca L. - **tree lavatera**
native to southern Europe; reported for Minnesota by USDA-NRCS but no specimens at MIN; doubtfully persisting annual
Malva moschata L. - **musk mallow**
native to Europe; known only from St. Louis Co. (lawn weed, collected 2007)
Malva neglecta Wallroth - **common mallow**
native from Eurasia and northern Africa
Malva parviflora L. - **cheeseweed mallow**
Malva pusilla Sm. - **low mallow**
Malva rotundifolia L. - **low mallow**
native to Europe
Malva sylvestris L. - **high mallow**

native to Eurasia; yard weed, doubtfully persisting

***Malva verticillata* L. - cluster mallow**

native to Asia; known only from Kittson Co. (one riverbank population, collected 1958), doubtfully persisting

***Napaea dioica* L. - glademallow**

northwest limit of distribution; threatened

***Sphaeralcea coccinea* var. *coccinea* (Nutt.) Rydb. - scarlet globemallow**

report unverified

***Tilia americana* L. - American basswood**

***Tilia x europaea* L. - common linden**

(*Tilia cordata* x *platyphyllos*)

MARSILEACEAE

***Marsilea vestita* Hooker & Greville - hairy waterclover**

northeast limit of distribution; endangered

MELANTHIACEAE

***Trillium cernuum* L. - whip-poor-will flower**

vars. not recognized

***Trillium flexipes* Rafinesque - nodding wakerobin**

***Trillium grandiflorum* (Michaux) Salisbury - white trillium**

***Trillium nivale* Riddell - snow trillium**

***Zigadenus elegans* Pursh - mountain deathcamas**

plants poisonous; vars. not recognized

MENISPERMACEAE

***Menispermum canadense* L. - common moonseed**

seeds possibly poisonous

MENYANTHACEAE

***Menyanthes trifoliata* L. - buckbean**

MOLLUGINACEAE

***Mollugo verticillata* L. - green carpet weed**

native to tropical America, now widely established (archeological evidence indicates a pre-Columbian presence at least as far north as TN)

MORACEAE

***Fatoua villosa* (Thunberg) Nakai - hairy crabweed**

native to Asia; known only from Anoka Co. (one nursery population, collected 1998)

***Morus alba* L. - white mulberry**

native to eastern Asia; hybridizes with *M. rubra* and the hybrids spreading rapidly in the state (M. Nepal, pers. comm.) and frequently mistaken for this parent

***Morus rubra* L. - red mulberry**

known only from SE counties; rare historic record

MYRICACEAE

***Comptonia peregrina* (L.) J.M. Coulter - sweet fern**

***Myrica gale* L. - sweetgale**

MYRSINACEAE

***Lysimachia arvensis* (L.) U. Manns & Anderberg - scarlet pimpernel**

native to Europe (last collected 1902); (nomenclatural change was made after FNA went to press; see *Willdenowia* 39: 51. 2009)

***Lysimachia borealis* (Rafinesque) U. Manns & Anderberg - starflower**

(nomenclatural change was made after FNA went to press; see *Willdenowia* 39: 51. 2009)

***Lysimachia ciliata* L. - fringed loosestrife**

***Lysimachia hybrida* Michaux - Mississippi loosestrife**

***Lysimachia lanceolata* Walter - lanceleaf loosestrife**

to be expected especially in southeastern Minnesota on moist sites; known from IA, WI, Ontario, and Manitoba

***Lysimachia maritima* (L.) Galasso, Banfi & Soldano - sea milkwort**

known only from Kittson Co. (last collected 1992); endangered

***Lysimachia minima* (L.) U. Manns & Anderberg - chaffweed**

known only from Big Stone Co. (collected 1901); (nomenclatural change made after FNA went to press (see *Willdenowia* 39: 52. 2009); rare historic record

***Lysimachia nummularia* L. - creeping jenny**

native to Europe; rarely producing seed

***Lysimachia quadriflora* Sims - four flowered yellow loosestrife**

***Lysimachia quadrifolia* L. - fourflower yellow loosestrife**

northwest limit of distribution; known only from Pine Co. (last collected 2000); special concern

***Lysimachia terrestris* (L.) Britton, Sterns & Poggenburg - earth loosestrife**

***Lysimachia thrysiflora* L. - tufted loosestrife**

***Lysimachia vulgaris* L. - garden yellow loosestrife**

native to Eurasia; known only from St. Louis Co. (Duluth area; last collected 1993)

***Lysimachia x commixta* Fernald - mixed loosestrife**

(*L. terrestris* x *thrysiflora*) known only from Washington Co. (one riverbank population, collected 1961); rarely producing seed

NELUMBONACEAE

***Nelumbo lutea* Willdenow - American lotus**

NYCTAGINACEAE

Mirabilis albida (Walter) Heimerl - **white four o'clock**

Mirabilis linearis var. *linearis* - **narrowleaf four o'clock**

known only from Houston Co. (historical collections to 1899) but to be expected along the Dakota borders in disturbed rocky or gravelly sites

Mirabilis nyctaginea (Michaux) MacMillan - **heartleaf four o'clock**

NYMPHAEACEAE

Nuphar lutea subsp. *pumila* (Timm) E.O. Beal - **yellow pond-lily**

misapplied to our flora, see *N. microphylla*

Nuphar microphylla (Persoon) Fernald - **yellow pond-lily**

Nuphar rubrodisca Morong - **yellow pond-lily**

possibly a hybrid (sometimes sterile) of *N. microphylla* and *N. variegata*

Nuphar variegata Durand - **variegated yellow pond-lily**

subsp. not recognized

Nymphaea leibergii Morong - **Leiberg's waterlily**

Nymphaea odorata Aiton - **American white waterlily**

Nymphaea odorata subsp. *odorata*

Nymphaea odorata subsp. *tuberosa* (Paine) Wiersema & Hellquist

Nymphaea tetragona Georgi - **Leiberg's waterlily**

misapplied to our flora, see *N. leibergii*

OLEACEAE

Fraxinus americana L. - **white ash**

Fraxinus nigra Marshall - **black ash**

Fraxinus pennsylvanica Marshall - **green ash**

vars. not recognized

Fraxinus quadrangulata Michx. - **blue ash**

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IA

Syringa reticulata subsp. *amurensis* (F. Ruprecht) P.S. Greene & M.C. Chang - **Amur lilac**

naturalized only from Ramsey Co. (city park, collected 2008)

Syringa reticulata subsp. *reticulata* - **Japanese tree lilac**

naturalized only in St. Louis Co. (Duluth area; last collected 2009)

Syringa villosa Vahl - **villous lilac**

native to China

Syringa vulgaris L. - **common lilac**

native to southern Europe

Syringa x persica L. - **Persian lilac**

(*S. afghanica* x *sibirica*) native to Asia; known only from Isanti Co. (one population, collected 1971), probable garden escapee

ONAGRACEAE

Calylophus serrulatus (Nuttall) P.H. Raven - **yellow sundrops**

Chamerion angustifolium subsp. *angustifolium* (L.) Holub

Chamerion angustifolium subsp. *circumvagum* (Mosquin) Hoch

Circaeа alpina subsp. *alpina* - **small enchanter's nightshade**

Circaeа lutetiana var. *canadensis* L. - **broadleaf enchanter's nightshade**

Circaeа x intermedia J.F. Ehrlhart - **hybrid enchanter's nightshade**

(*C. alpina* x *lutetiana*)

Epilobium brachycarpum C. Presl - **tall annual willowherb**

reported for Minnesota by USDA-NRCS but no specimens at MIN; probably based on a misidentified specimen; known from Canada, ND, SD, WI

Epilobium ciliatum subsp. *ciliatum* - **fringed willowherb**

Epilobium ciliatum subsp. *glandulosum* (Lehm.) Hoch & P.H. Raven - **fringed willowherb**

Epilobium coloratum Biehler - **purpleleaf willowherb**

Epilobium glandulosum Lehman - **fringed willowherb**

included in *E. ciliatum* by some as a subspecies or variety

Epilobium leptophyllum Rafinesque - **bog willowherb**

Epilobium palustre L. - **marsh willowherb**

Epilobium strictum Muhlenberg ex Sprengel - **downy willowherb**

Epilobium x wisconsinense Ugent - **Wisconsin willowherb**

(*E. ciliatum* x *coloratum*)

Gaura biennis L. - **biennial beeblissom**

Gaura coccinea Nuttall ex Pursh - **scarlet beeblissom**

Gaura parviflora Douglas ex Lehm. - **velvetweed**

Ludwigia palustris (L.) Elliott - **marsh seedbox**

Ludwigia peploides (Kunth) P.H. Raven - **floating primrose-willow**

Ludwigia polycarpa Short & R. Peter - **manyfruit primrose-willow**

Oenothera biennis var. *biennis* - **common evening primrose**

Oenothera clelandii W. Dietrich, P.H. Raven & W.L. Wagner - **Cleland's evening primrose**

Oenothera laciniata Hill - **cutleaf evening primrose**

Oenothera nuttallii Sweet - **Nuttall's evening primrose**

Oenothera oakesiana (A. Gray) Robbins ex. S. Wats. & J.M. Coulter. - **Oak's evening primrose**

Oenothera parviflora L. - **northern evening primrose**

Oenothera parviflora var. *oakesiana* (A. Gray) Fernald

Oenothera perennis L. - **little evening primrose**

Oenothera pilosella Raf. - **meadow evening primrose**

Oenothera rhombipetala Nuttall ex Torrey & A. Gray - fourpoint evening primrose

Oenothera villosa subsp. *strigosa* (Rydb.) W. Dietr. & P.H. Raven

Oenothera villosa subsp. *villosa*

OPHIOGLOSSACEAE

Botrychium acuminatum W.H. Wagner - tailed grape fern

restricted to Lake Superior region; known only from Cook Co. (last collected 1999); special concern

Botrychium ascendens W.H. Wagner - trianglelobe moonwort

known only from mine dumps in Crow Wing Co. (and one location in St. Louis Co.); disjunct from the western montane region and Hudson Bay; endangered

Botrychium campestre W.H. Wagner & Farrar - Iowa moonwort

Botrychium crenulatum W. H. Wagner - scalloped moonwort

reported but no specimens deposited at MIN; DNR watch list

Botrychium dissectum - cutleaf grape fern

Botrychium gallicomontanum Farrar & Johnson-Groh - Frenchman's Bluff grape fern

"previously thought to be endemic to Minnesota (fide D. Farrar a population also exists in MT); endangered"

Botrychium hesperium (Maxon & R.T. Clausen) W.H. Wagner & Lellinger - western moonwort

one specimen previously identified by W.H. Wagner as possibly this species, was later annotated as coming from a population of *B. matricariifolium*. *Botrychium hesperium* is known from the Thunder Bay area of Ontario and Upper Peninsula MI; erroneous record

Botrychium lanceolatum subsp. *angustisegmentum* (Pease & A.H. Moore) R.T. Clausen - lanceleaf grape fern

Botrychium lineare W.H. Wagner - narrowleaf grape fern

Wagner & Wagner did not include this species in their treatment for FNA; known only from mine dumps in St. Louis Co. (collected 2007); disjunct from MT; endangered

Botrychium lunaria (L.) Swartz - common moonwort

Botrychium matricariifolium (Dill) A. Braun ex W.D.J. Koch - matricary grape fern

Botrychium michiganense W.H. Wagner ex A.V. Gilman, Farrar & Zika - Michigan moonwort

Botrychium minganense Victorin - Mingan moonwort

Botrychium mormo W.H. Wagner - little goblin moonwort

Botrychium multifidum (S.G. Gmelin) Ruprecht - leathery grape fern

Botrychium neolonaria

unpublished

Botrychium oneidense (Gilbert) House - bluntlobe grape fern

western limit of distribution; threatened

Botrychium pallidum W.H. Wagner - pale botrychium

mostly limited to upper Great Lakes region; special concern

Botrychium pedunculosum W.H. Wagner - stalked moonwort

Botrychium pseudopinnatum W.H. Wagner - false daisyleaf moonwort

range extension from Ontario; known only from St. Louis County (only known specimen at DUL)

Botrychium rugulosum W.H. Wagner - ternate grape fern

western limit of distribution; special concern

Botrychium simplex E. Hitchcock - little grape fern

Botrychium spathulatum W.H. Wagner - spatulate botrychium

known only from Crow Wing and Cook counties; endangered

Botrychium virginianum (L.) Swartz - rattlesnake fern

Ophioglossum pusillum Rafinesque - northern adder's tongue

ORCHIDACEAE

Amerorchis rotundifolia (Banks ex Pursh) Hultn - roundleaf orchid

Aplectrum hyemale (Muhlenberg ex Willdenow) Torrey - adam and eve

Arethusa bulbosa L. - dragon's-mouth

Calopogon oklahomensis D.H. Goldman - Oklahoma grasspink

Calopogon tuberosus var. *tuberosus* - tuberous grasspink

Calypso bulbosa var. *americana* (R. Brown) Luer - fairy slipper

Coeloglossum viride (L.) Hartman - longbract frog orchid

vars. not recognized

Corallorrhiza maculata var. *maculata*

Corallorrhiza maculata var. *occidentalis* (Lindley) Ames

Corallorrhiza odontorhiza (Willdenow) Poiret

Corallorrhiza odontorhiza var. *odontorhiza* - autumn coralroot

Corallorrhiza striata var. *striata* - hooded coralroot

Corallorrhiza trifida Chtelain - yellow coralroot

Cypripedium acaule Aiton - moccasin flower

may cause skin irritation

Cypripedium arietinum R. Brown - ram's head lady's slipper

may cause skin irritation; threatened

Cypripedium candidum Muhlenberg ex Willdenow - white lady's slipper

may cause skin irritation; special concern

Cypripedium parviflorum Salisbury - greater yellow lady

may cause skin irritation

Cypripedium parviflorum var. *makasin* (Farwell) Sheviak - greater yellow lady's slipper

Cypripedium parviflorum var. *parviflorum* - greater yellow lady's slipper

misapplied; see *Cypripedium parviflorum* var. *makasin*

Cypripedium parviflorum var. *pubescens* (Willdenow) O.W. Knight - greater yellow lady's slipper

Cypripedium reginae Walter - showy lady's slipper

may cause skin irritation

Cypripedium x andrewsii var. *andrewsii* - Andrews's lady-slipper

report unverified

Cypripedium x andrewsii var. *favillianum* (J. T. Curtis) B. Boivin - Andrews's lady-slipper
Epipactis helleborine (L.) Crantz - broadleaf helleborine
Galearis spectabilis (L.) Rafinesque - showy orchid
Goodyera pubescens (Willdenow) R. Brown - downy rattlesnake plantain
Goodyera repens (L.) R. Brown - lesser rattlesnake plantain
vars. not recognized
Goodyera tesselata Loddiges - checkered rattlesnake plantain
Habenaria hyperborea (L.) R. Br. - northern green orchid
misapplied; see *Platanthera aquilonis* or *Platanthera huronensis*
Liparis liliifolia (L.) Richard ex Lindley - brown widelip orchid
Liparis loeselii (L.) Richard - yellow widelip orchid
Listera auriculata Wiegand - auricled twayblade
western limit of distribution; endangered
Listera convallarioides (Swartz) Nuttall ex Elliott - broadlipped twayblade
known only from Cook Co. (one population, collected 1924); special concern
Listera cordata var. *cordata* - heartleaf twayblade
Malaxis monophyllos (L.) Swartz
Malaxis monophyllos var. *brachypoda* (A. Gray) F. Morris & E.A. Eames - white adder's-mouth orchid
Malaxis paludosa (L.) Swartz - bog adder's-mouth orchid
southern limit of distribution; endangered
Malaxis unifolia Michaux - green adder's-mouth orchid
Platanthera aquilonis Sheviak - northern green orchid
Platanthera clavellata (Michaux) Luer - small green wood orchid
Platanthera dilatata (Pursh) Lindley ex L.C. Beck
Platanthera dilatata var. *dilatata* - scentbottle
Platanthera flava var. *herbiola* (R. Brown) Luer - palegreen orchid
northwest limit of distribution; threatened
Platanthera hookeri (Torrey ex A. Gray) Lindley - Hooker's orchid
Platanthera huronensis (Nuttall) Lindley - Huron green orchid
Platanthera hyperborea (L.) Lindl. - northern green orchid
misapplied; see *Platanthera huronensis* or *Platanthera aquilonis*
Platanthera lacera (Michaux) G. Don - green fringed orchid
Platanthera obtusata (Banks ex Pursh) Lindley - bluntleaved orchid
Platanthera orbiculata (Pursh) Lindley - lesser roundleaved orchid
Platanthera praecalaria Sheviak & M.L. Bowles - Great Plains white fringed orchid
federal Threatened list; endangered
Platanthera psycodes (L.) Lindley - lesser purple fringed orchid
vars. not recognized
Pogonia ophioglossoides (L.) Ker Gawler - snakemouth orchid
Spiranthes casei var. *casei* - Case's lady's tresses
Spiranthes cernua (L.) Richard - nodding lady's tresses
Spiranthes lacera var. *gracilis* (Bigelow) Luer - northern slender lady's tresses
known only from Hennepin Co. (historical collection from 1889)
Spiranthes lacera var. *lacera* - northern slender lady's tresses
Spiranthes magnicamporum Sheviak - Great Plains lady's tresses
Spiranthes romanzoffiana Chamisso - hooded lady's tresses
Spiranthes x simpsonii Catling & Sheviak - Simpson's lady's-tresses
report unverified; see *S. lacera* x *ramanzoffiana*

OROBANCHACEAE

Agalinis aspera (Douglas ex Bentham) Britton - tall false foxglove
Agalinis auriculata (Michaux) S.F. Blake - earleaf false foxglove
northwest limit of distribution; endangered
Agalinis gattingeri (Small) Small - roundstem false foxglove
northwest limit of distribution; endangered
Agalinis paupercula var. *borealis* Pennell - smallflower false foxglove
Agalinis paupercula var. *paupercula* - smallflower false foxglove
Agalinis purpurea (L.) Pennell - purple false foxglove
Agalinis tenuifolia var. *macrophylla* (Benth.) S.F. Blake - slenderleaf false foxglove
Agalinis tenuifolia var. *parviflora* (Nutt.) Pennell - slenderleaf false foxglove
Aureolaria grandiflora var. *pulchra* Pennell - largeflower yellow false foxglove
northern limit of distribution; known only from Houston Co. (historical collection from 1899); rare historic record
Aureolaria pedicularia var. *ambigens* (Fernald) Farwell - fernleaf yellow false foxglove
Aureolaria pedicularia var. *intercedens* Pennell
known only from Hennepin Co. (historical collection from 1878)
Aureolaria pedicularia var. *pedicularia* - fernleaf yellow false foxglove
report unverified
Castilleja coccinea (L.) Sprengel - scarlet Indian paintbrush
Castilleja septentrionalis Lindley - Labrador Indian paintbrush
southwest limit of distribution; endangered
Castilleja sessiliflora Pursh - downy paintedcup
Euphrasia hudsoniana var. *contracta* P.D. Sell & Yeo - Hudson Bay eyebright
report unverified
Euphrasia hudsoniana var. *ramosior* Sell & Yeo - Hudson Bay eyebright

Euphrasia micrantha - northern eyebright

Euphrasia nemorosa (Pers.) Wallr.

Euphrasia officinalis L. - eyebright

native to Europe; DNR watch list

Euphrasia stricta J.F. Lehmann - drug eyebright

Melampyrum lineare var. *latifolium* (Muhl. ex Britton) Beauverd - narrowleaf cowwheat

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IL

Melampyrum lineare var. *lineare* - narrowleaf cowwheat

Melampyrum lineare var. *pectinatum* (Pennell) Fernald - narrowleaf cowwheat

Orobanche fasciculata Nuttall - clustered broomrape

federal noxious weed; threatened

Orobanche ludoviciana subsp. *ludoviciana* - Louisiana broomrape

federal noxious weed; threatened

Orobanche uniflora L. - oneflowered broomrape

federal noxious weed; threatened

Orthocarpus luteus Nuttall - yellow owl's-clover

Pedicularis canadensis L. - Canadian lousewort

Pedicularis lanceolata Michaux - swamp lousewort

OSMUNDACEAE

Osmunda claytoniana L. - interrupted fern

Osmunda regalis var. *spectabilis* (Willdenow) A. Gray - royal fern

Osmundastrum cinnamomeum (L.) C. Presl - cinnamon fern

(see J.S. Metzgar et al., 2008, Systematic Botany 33(1): 31-36 for discussion of the name change)

OXALIDACEAE

Oxalis dillenii N.J. Jacquin - slender yellow wood sorrel

Oxalis montana Raf. - mountain wood sorrel

Oxalis stricta L. - common yellow oxalis

Oxalis violacea L. - violet wood sorrel

PAPAVERACEAE

Chelidonium majus L. - celandine

native to Eurasia

Eschscholzia californica subsp. *californica* - California poppy

Papaver rhoeas L. - corn poppy

native from Eurasia and west Africa; occasionally found as annual escape but generally not persisting

Papaver somniferum L. - opium poppy

native to Mediterranean region; known only from St. Louis Co. (Duluth area; collected 1978)

Sanguinaria canadensis L. - bloodroot

plant toxic

PEDALIACEAE

Proboscidea louisianica (Mill.) Thell. - ram's-horn

report unverified

PENTHORACEAE

Penthorum sedoides L. - ditch stonecrop

PHRYMACEAE

Mimulus glabratus var. *jamesii* A. Gray - James' monkey flower

Mimulus ringens var. *ringens* - Allegheny monkey flower

Phryma leptostachya L. - American lopseed

PHYTOLACCACEAE

Phytolacca americana var. *americana* - American pokeweed

reported for Minnesota by FNA but no specimens at MIN; known from Ontario, WI, IA

PINACEAE

Abies balsamea (L.) Miller - balsam fir

may cause skin irritation

Abies fraseri (Pursh) Poiret - Fraser fir

native to southeastern US; known only from Kanabec Co. (collected 1999), probable escapee from tree farm

Larix decidua Miller - European larch

native to Eurasia

Larix laricina (Du Roi) K. Koch - tamarack

Picea abies (L.) H. Karsten - Norway spruce

native to Europe

Picea glauca (Moench) Voss - white spruce

Picea glauca x *marianna*

Picea mariana (Miller) Britton, Sterns & Poggenburg - black spruce

Picea pungens Engelmann - blue spruce

native to western US; known only from Lake of the Woods Co. (collected 1981), probable escapee from plantation

Picea x rosendahlii Little & Pauley

invalid name; see *P. glauca* x *marianna*

Pinus banksiana Lambert - jack pine
Pinus resinosa Aiton - red pine
Pinus rigida Miller - pitch pine
native to eastern US and Canada; known only from Washington Co. (collected 1960), escapee from plantation
Pinus strobus L. - eastern white pine
Pinus sylvestris L. - Scots pine
native to Europe
Pseudotsuga menziesii (Mirbel) Franco - douglas-fir
native to the Pacific Northwest; known only from Winona Co. (collected 1967), escapee from plantation
Tsuga canadensis (L.) Carrige - eastern hemlock

PLANTAGINACEAE

Bacopa rotundifolia (Michaux) R. Wetstein - disk waterhyssop
Besseyea bullii (Eaton) Rydberg - Bull's coraldrops
northwest limit of distribution; threatened
Callitricha hermaphroditica L. - northern water-starwort
Callitricha heterophylla subsp. *heterophylla* - twoheaded water-starwort
Callitricha palustris L. - vernal water-starwort
Chaenorhinum minus (L.) Lange - dwarf snapdragon
native to Europe
Chelone glabra L. - white turtlehead
Chelone obliqua var. *speciosa* Pennell & Wherry - red turtlehead
Digitalis grandiflora Lamarck - yellow foxglove
native to Europe and western Asia; known only from St. Louis Co. (Duluth area; one weedy population, collected 1997)
Digitalis lanata J.F. Ehrhart - Grecian foxglove
native to southern Europe; known only from Washington Co. (last collected 1999); known to be invasive
Gratiola neglecta Torrey - clammy hedgehyssop
Hippuris vulgaris L. - clammy hedgehyssop
Leucospora multifida (Michx.) Nutt. - narrowleaf paleseed
report unverified
Limosella aquatica L. - water mudwort
Linaria dalmatica subsp. *dalmatica*
native to eastern Mediterranean region
Linaria genistifolia (L.) Miller - broomleaf toadflax
native to Europe; known only from Pope Co. (one roadside population, collected 1938)
Linaria texana Scheele - Canada toadflax
reported for Minnesota by the Atlas of the Flora of the Great Plains 1986 but no specimens at MIN; known from Saskatchewan, ND, SD, NE
Linaria vulgaris Miller - butter and eggs
native to Europe
Lindernia dubia var. *anagallidea* (Michaux) T.S. Cooperider - yellowseed false pimpernel
Lindernia dubia var. *dubia* - yellowseed false pimpernel
Littorella uniflora (L.) Ascherson
Nuttallanthus canadensis (L.) D.A. Sutton - Canada toadflax
Odontites vernus subsp. *serotinus* Corb. - red bartsia
Penstemon albidus Nuttall - white penstemon
Penstemon digitalis Nuttall ex Sims - tall slope penstemon
Penstemon gracilis var. *gracilis* - lilac penstemon
Penstemon grandiflorus Nuttall - large beardtongue
Penstemon laevigatus Aiton - eastern smooth beardtongue
native to southeastern US; known only from Stearns Co. (collected 1971), possible garden escape; erroneous record
Penstemon pallidus Small - pale beardtongue
Penstemon smallii A. Heller - Small's beardtongue
Plantago aristata Michaux - largebracted plantain
Plantago elongata subsp. *elongata* - prairie plantain
Plantago eriopoda Torrey - redwool plantain
Plantago lanceolata L. - narrowleaf plantain
native to Eurasia; state DOA restricted weed, 1st collected 1891, Brown and Cook counties
Plantago major L. - common plantain
native to Eurasia
Plantago patagonica N.J. Jacquin - woolly plantain
Plantago psyllium L. - sand plantain
native to eastern Mediterranean region
Plantago rugelii var. *rugelii* - blackseed plantain
Plantago virginica L. - Virginia plantain
known only from Cook Co. (one population, collected 1944); rare historic record
Pseudolysimachion longifolium (L.) Opiz - longleaf speedwell
Veronica americana Schweinitz ex Bentham - American speedwell
Veronica anagallis-aquatica L. - water speedwell
Veronica arvensis L. - corn speedwell
native to Eurasia
Veronica austriaca subsp. *teucrium* (L.) D.A. Webb - broadleaf speedwell
native to Eurasia; known only from Lake Co. (one lakeshore population, collected 1999)
Veronica longifolia L. - longleaf speedwell
native to Europe

Veronica officinalis var. *officinalis* - common gypsyweed
Veronica peregrina var. *peregrina* - neckweed
Veronica peregrina var. *xalapensis* (Kunth) Pennell - hairy purslane speedwell
Veronica persica Poiret - birdeye speedwell
native to southwestern Asia
Veronica prostrata L. - prostrate speedwell
erroneous report
Veronica satureifolia Poiteau & Turpe - basil-leaved speedwell
native to Europe; known only from St. Louis Co. (Duluth area; lawn weed, collected 2002)
Veronica scutellata L. - skullcap speedwell
Veronica serpyllifolia var. *humifusa* (J.J. Dickson) Syme - brightblue speedwell
Veronica serpyllifolia var. *serpyllifolia* - thymeleaf speedwell
native to Europe
Veronica verna L. - spring speedwell
native to Europe
Veronicastrum virginicum (L.) Farwell - culver's root

POACEAE

Achnatherum hymenoides (Roemer & Schultes) Barkworth - Indian ricegrass
eastern limit of distribution; endangered
Agropyron cristatum (L.) Gaertner - crested wheatgrass
native to Eurasia
Agrostis alba L. - redtop
misapplied to our flora, see *A. gigantea* or *A. stolonifera*
Agrostis canina L. - velvet bentgrass
erroneous report
Agrostis gigantea Roth - redtop
native to Eurasia
Agrostis hyemalis (Walter) Britton, Sterns & Poggenburg - winter bentgrass
northwestern edge of range; known only from southeastern Minnesota (Anoka Co. populations prob. introduced); previous widespread reports were based on previous inclusion of *A. scabra* in this taxon; endangered
Agrostis perennans (Walter) E. Tuckerman - upland bentgrass
Agrostis scabra Willdenow - rough bentgrass
native to Pacific coastal countries, introduced elsewhere; includes *A. gigantea* which was on the state Special Concern list
Agrostis stolonifera L. - creeping bentgrass
both native and introduced (from Eurasia) forms may occur but vars. not recognized
Alopecurus aequalis var. *aequalis* - shortawn foxtail
Alopecurus carolinianus Walter - Carolina foxtail
Alopecurus geniculatus L. - water foxtail
native to Eurasia; reported for Minnesota (southwestern counties) by FNA but no specimens at MIN
Alopecurus pratensis L. - meadow foxtail
native to Eurasia
Ammophila breviligulata subsp. *breviligulata* - American beachgrass
western limit of distribution; known only from St. Louis Co. (Duluth area; last collected 1983); threatened
Andropogon gerardii Vitman - big bluestem
Andropogon hallii Hackel - sand bluestem
native to the Great Plains; introduced with roadside/wildlife plantings
Anthoxanthum aristatum Boiss. - annual vernalgrass
erroneous report
Anthoxanthum hirtum (Schrank) Y. Schouten & Veldkamp - northern sweetgrass
Aristida basiramea Engelmann ex Vasey - forked threeawn
Aristida dichotoma var. *curtissii* A. Gray - Curtis's threeawn
Aristida dichotoma - churchmouse threeawn
possibly also occurring but no specimens at MIN
Aristida longespica var. *geniculata* (Rafinesque) Fernald - slimspike threeawn
known only from Anoka Co. (last collected 2005) but probably introduced; endangered
Aristida longespica var. *longespica* - slimspike threeawn
erroneous report
Aristida oligantha Michaux - prairie threeawn
Aristida purpurea var. *longiseta* (Steudel) Vasey - Fendler threeawn
Aristida purpurea var. *purpurea* - purple threeawn
possibly also occurring but no specimens at MIN; special concern
Aristida tuberculosa Nuttall - seaside threeawn
Arrhenatherum elatius subsp. *bulbosum* (Willdenow) Schübeler & G. Martens - tall oatgrass
known only from Ramsey Co. (collected 1891)
Arrhenatherum elatius subsp. *elatius* - tall oatgrass
known only from Hennepin Co. (collected 1943)
Avena fatua L. - wild oat
native to Eurasia; occasionally persisting near cultivated fields; state DOA noxious (CN) weed
Avena sativa L. - common oat
native to Eurasia; occasionally persisting near cultivated fields; moderately toxic to livestock
Avenula hookeri (Scribnier) Holub - spikeoat
Avenula pubescens (Huds.) Dumort. - downy alpine oatgrass
erroneous report
Beckmannia syzigachne (Steudel) Fernald - American sloughgrass
Bouteloua curtipendula var. *curtipendula* - sideoats gramma

Bouteloua gracilis (Kunth) Lagasca ex Griffiths - **blue grama**
Bouteloua hirsuta var. hirsuta - **hairy grama**
Brachyelytrum aristosum (Michaux) P. Beauvois - **northern shorthusk**
Brachyelytrum erectum (Schreber) P. Beauvois - **bearded shorthusk**
Bromus catharticus var. catharticus - **rescuegrass**
Bromus ciliatus L. - **fringed brome**
Bromus hordeaceus subsp. hordeaceus - **soft brome**
Bromus inermis Leysser - **smooth brome**
native to Eurasia
Bromus japonicus Thunberg - **field brome**
native to Eurasia
Bromus kalmii A. Gray - **arctic brome**
Bromus latiglumis (Scribnier ex Shear) A.S. Hitchcock - **earlyleaf brome**
Bromus pubescens Muhlenberg ex Willdenow - **hairy woodland brome**
Bromus pumpellianus subsp. pumpellianus - **Pumpelly's brome**
native to Canada and western US; difficult to distinguish from *B. inermis* (especially) or *B. latiglumis*; previous report from Minnesota based on misidentified specimen, however an unverified report from Norman Co. may be this species
Bromus secalinus L. - **chess**
native to Europe; known only from historical collections (to 1899)
Bromus tectorum L. - **cheatgrass**
native to Eurasia
Buchloe dactyloides (Nuttall) Engelman - **buffalograss**
Calamagrostis canadensis var. canadensis - **bluejoint**
Calamagrostis canadensis var. langsdorffii (Link) Inman - **bluejoint**
Calamagrostis canadensis var. macouniana (Vasey) Stebbins - **Macoun's reedgrass**
Calamagrostis montanensis (Scribnier) Vasey - **plains reedgrass**
eastern limit of distribution; special concern
Calamagrostis purpurascens R. Brown - **purple reedgrass**
disjunct from western and northern North America; known only from Cook Co. (last collected 2005); endangered
Calamagrostis stricta (Timm) Koeler - **slimstem reedgrass**
Calamagrostis stricta subsp. inexpectata (A. Gray) C.W. Greene - **northern reedgrass**
Calamovilfa longifolia var. longifolia - **prairie sandreed**
Catabrosa aquatica (L.) P. Beauvois - **water whorlgrass**
to be expected along the North Dakota border, in wet meadows and along streams
Cenchrus longispinus (Hackel) Fernald - **mat sandbur**
state DOA noxious (CN) weed
Cenchrus spinifex Cavanilles - **coastal sandbur**
disjunct from IL and IA
Cinna arundinacea L. - **sweet woodreed**
Cinna latifolia (Treviranus ex Gppert) Grisebach - **drooping woodreed**
Dactylis glomerata L. - **orchardgrass**
native to Eurasia and Africa
Danthonia spicata (L.) P. Beauvois ex Roemer & Schultes - **poverty oatgrass**
Deschampsia cespitosa subsp. cespitosa - **tufted hairgrass**
Deschampsia flexuosa (L.) Trinius - **wavy hairgrass**
Diarrhena obovata (Gleason) Brandenburg - **obovate beakgrain**
known only from Fillmore Co. (last collected 1999); endangered
Dichanthelium aciculare subsp. angustifolium (Elliott) Freckmann & Lelong - **needleleaf rosette grass**
disjunct from southern and eastern US; known only from Clearwater Co. (Itasca State Park (collected 1962), probably an introduced waif
Dichanthelium acuminatum subsp. acuminatum - **tapered rosette grass**
Dichanthelium acuminatum subsp. columbianum (Scribnier) Freckmann & Lelong - **tapered rosette grass**
Dichanthelium acuminatum subsp. fasciculatum (Torrey) Freckmann & Lelong - **western panicgrass**
Dichanthelium acuminatum subsp. implicatum (Scribnier) Freckmann & Lelong - **western panicgrass**
Dichanthelium acuminatum subsp. lindheimeri (Nash) Freckmann & Lelong - **Lindheimer panicgrass**
Dichanthelium boreale (Nash) Freckmann - **northern panicgrass**
Dichanthelium depauperatum (Muhlenberg) Gould - **starved panicgrass**
Dichanthelium dichotomum (L.) Gould - **cypress panicgrass**
reported for Minnesota by MN-DNR but no specimens at MIN; known from southern WI south- and eastward
Dichanthelium latifolium (L.) Harvill - **broadleaf rosette grass**
Dichanthelium leibergii (Vasey) Freckmann - **Leiberg's panicgrass**
Dichanthelium linearifolium (Scribnier) Gould - **slimleaf panicgrass**
Dichanthelium oligosanthes subsp. oligosanthes - **Heller's rosette grass**
Dichanthelium oligosanthes subsp. scribnerianum (Nash) Freckmann & Lelong - **Scribner's rosette grass**
Dichanthelium ovale subsp. praecocius (Hitchcock & Chase) Freckmann & Lelong - **egg-leaf rosette grass**
Dichanthelium ovale subsp. pseudopubescens (Nash) Freckmann & Lelong - **whitehair rosette grass**
Dichanthelium ovale subsp. villosissimum (Nash) Freckmann & Lelong - **whitehair rosette grass**
Dichanthelium perlóngum (Nash) Freckmann - **slimleaf panicgrass**
Dichanthelium sabulorum (Lam.) Gould & C.A. Clark - **western panicgrass**
misapplied to our flora, see *D. acuminatum* subsp. *Implicatum*
Dichanthelium wilcoxianum (Vasey) Freckmann - **fall rosette grass**
Dichanthelium x scopariooides (Ashe) Mohlenbr. - **Shasta panicgrass**
(*D. acuminatum* x *oligosanthes*) reported for Minnesota by USDA-NRCS but no specimens at MIN; this hybrid not reported in FNA
Dichanthelium xanthophysum (A. Gray) Freckmann - **slender rosette grass**
Digitaria cognata (Schultes) Pilger - **fall witchgrass**

Digitaria ischaemum (Schreber) Muhlenberg - **smooth crabgrass**

native to Eurasia

Digitaria sanguinalis (L.) Scopoli - **hairy crabgrass**

native to Eurasia

Distichlis spicata (L.) Greene - **saltgrass**

vars. not recognized

Echinochloa crus-galli (L.) P. Beauvois - **barnyard grass**

native to Eurasia

Echinochloa frumentacea Link - **billion-dollar grass**

native to India; reported from experimental plots in Minnesota, unknown if spreading or persisting

Echinochloa muricata var. *microstachya* Wiegand - **rough barnyard grass**

Echinochloa muricata var. *muricata* - **rough barnyard grass**

Echinochloa walteri (Pursh) A. Heller - **coast cockspur grass**

Eleusine coracana subsp. *coracana* - **finger millet**

native to Africa; known only from Ramsey Co. (one agricultural population, collected 2003), doubtfully persisting

Eleusine indica (L.) Gaertner - **Indian goosegrass**

native to Africa; occasional weed near cultivated fields

Elyhordeum macounii (Vasey) Barkworth & Dewey - **Macoun's barley**

(Elymus trachycaulis x Hordeum jubatum)

Elyhordeum montanense (Scrib. ex Beal) Bowden - **mountain barley**

(Elymus virginicus x Hordeum jubatum) to be expected on disturbed sites if both parents nearby

Elymus canadensis f. *crescendus* Ramaley

Elymus canadensis var. *canadensis* - **Canada wildrye**

Elymus canadensis var. *robustus* (Scribner & J.G. Smith) Mackenzie & Bush - **Canada wildrye**

known only from Chippewa Co. (historical collection from 1894)

Elymus curvatus Piper - **Virginia wildrye**

Elymus diversiglumis - **diverseglume wild rye**

Elymus hystrix L. - **eastern bottlebrush grass**

Elymus interruptus Buckley - **diverseglume wild rye**

misapplied to our flora, see E. diversiglumis

Elymus repens (L.) Gould - **quackgrass**

native to Eurasia; state DOA noxious (CN) weed, 1st collected 1886, St. Louis Co.

Elymus riparius Wiegand - **riverbank wildrye**

easily confused with E. villosus and possibly E. diversiglumis

Elymus trachycaulus subsp. *subsecundus* (Link) A. Löve & D. Löve - **slender wheatgrass**

Elymus trachycaulus subsp. *trachycaulus* - **slender wheatgrass**

Elymus trachycaulus unnamed polymorphic Group II - **slender wheatgrasse**

see FNA for discussion

Elymus villosus Muhlenberg ex Willdenow - **hairy wildrye**

Elymus virginicus var. *halophilus* (E.P. Bicknell) Wiegand

presently known only from Cottonwood Co. (however, most specimens of the species not identified to variety)

Elymus virginicus var. *jejonus* (Ramaley) Bush

Elymus virginicus var. *virginicus*

Elymus wiegandii Fernald - **Wiegand's wildrye**

Eragrostis capillaris (L.) Nees - **lace grass**

to be expected in the southeastern corner along the lower Mississippi River (known from IA, WI), on dry sandy riverbanks

Eragrostis cilianensis (Allioni) Vignolo ex Janchen - **stinkgrass**

native to Europe

Eragrostis frankii C.A. Meyer ex Steudel - **sandbar lovegrass**

Eragrostis hypnoides (Lamarck) Britton - **teal lovegrass**

Eragrostis minor Host - **little lovegrass**

native to Europe

Eragrostis pectinacea var. *pectinacea* - **tufted lovegrass**

Eragrostis pilosa var. *perplexa* (L.H. Harvey) S.D. Koch

known only from Pipestone Co. (collected 1931)

Eragrostis pilosa var. *pilosa*

known only from Pipestone Co. (historical collection from 1883)

Eragrostis spectabilis (Pursh) Steudel - **purple lovegrass**

Eragrostis trichodes (Nuttall) Alph. Wood - **sand lovegrass**

disjunct from central Great Plains

Eriochloa contracta Hitchc. - **prairie cupgrass**

reported for Minnesota (Mankato region) by FNA but no specimens at MIN; known from MO and NE

Eriochloa villosa (Thunberg) Kumth - **Chinese cupgrass**

native to eastern Asia; state DOA noxious (CN) weed, 1st collected 1967, Blue Earth Co.

Festuca brachyphylla Schultes & Schultes f. - **meadow ryegrass**

misapplied to our flora; see Schedonorus arundinaceus; previous reports of this species in Minnesota were based on a broader interpretation of the taxon; ours now to be considered the narrower F. saximontana

Festuca longifolia Thuijl. - **hard fescue**

misapplied to our flora, see F. trachyphylla

Festuca ovina var. *ovina*

misapplied; see F. saximontana

Festuca paradoxa Desvaux - **clustered fescue**

known only from Goodhue Co. (historical collection from 1886)

Festuca rubra subsp. *commutata* Gauden - **red fescue**

native to Europe, generally found in lawns but occasionally escaping; possibly occurring in Minnesota

Festuca rubra subsp. *fallax* (Thuillier) Nyman - **red fescue**

native to Europe; possibly occurring in Minnesota
Festuca rubra subsp. *rubra* - red fescue
native to Eurasia
Festuca saximontana Rydberg
Festuca saximontana var. *saximontana* - Rocky Mountain fescue
Festuca subverticillata (Persoon) E.B. Alexeev - nodding fescue
Festuca trachyphylla (Hackel) Krajina - hard fescue
native to Europe
Glyceria borealis (Nash) Batchelder - small floating mannagrass
Glyceria canadensis var. *canadensis* - rattlesnake mannagrass
Glyceria grandis var. *grandis* - American mannagrass
Glyceria septentrionalis var. *septentrionalis* - floating mannagrass
reported for Minnesota by USDA-NRCS and NatureServe but no specimens at MIN; known from adjacent La Crosse Co. WI south- and eastward FNA
Glyceria striata (Lamarck) Hitchcock - fowl mannagrass
vars. not recognized
Hesperostipa comata (Trinius & Ruprecht) Barkworth
Hesperostipa comata subsp. *comata* - needle and thread
Hesperostipa curtiseta (Hitchc.) Barkworth - shortbristle needle and thread
to be expected in the far northwest corner (known from ND, Manitoba), in prairies; easily confused with *H. comata*
Hesperostipa sparteae (Trinius) Barkworth - porcupinegrass
Hordeum jubatum subsp. *jubatum* - foxtail barley
Hordeum pusillum Nuttall - little barley
Hordeum vulgare L. - common barley
native to Europe
Koeleria macrantha (Ledebour) Schultes - prairie junegrass
Koeleria pyramidata (Lam.) P. Beauv. - prairie junegrass
misapplied to North American flora, see *K. macrantha*
Leersia lenticularis Michaux - catchfly grass
Leersia oryzoides (L.) Swartz - rice cutgrass
vars. not recognized
Leersia virginica Willdenow - whitegrass
Leptochloa fusca subsp. *fascicularis* (Lamarck) N. Snow - bearded sprangletop
Leymus arenarius (L.) Hochstetter - sand ryegrass
Leymus cinereus (Scribnier & Merrill) Å. Löve - basin wildrye
reported for Minnesota by MN-DNR and USDA-NRCS but no specimens at MIN; known only from western Great Plains westward
Leymus racemosus (Lamarck) Tzvelev - mammoth wildrye
native to Eurasia; known only from Houston Co. (one riverbank population, collected 1980)
Lolium multiflorum Lamarck - Italian ryegrass
Lolium perenne L. - perennial ryegrass
native to Europe
Lolium temulentum subsp. *temulentum* - darnel ryegrass
native to Mediterranean region; known only from Blue Earth Co. (historical collection from 1883)
Melica nitens (Scribnier) Nuttall ex Piper - threeflower melicgrass
northwest limit of distribution; threatened
Milium effusum var. *cisatlanticum* Fernald - American milletgrass
Misanthus sacchariflorus (Maximowicz) Bentham - amur silvergrass
native to eastern Asia; potentially invasive (first recorded naturalized population collected in 1977 in Le Sueur Co.)
Muhlenbergia asperifolia (Nees & Meyen ex Trinius) Parodi - scratchgrass
Muhlenbergia cuspidata (Torrey) Rydberg - plains muhly
Muhlenbergia frondosa (Poirer) Fernald - wirestem muhly
state DOA noxious (CN) weed
Muhlenbergia glomerata (Willdenow) Trinius - spiked muhly
Muhlenbergia mexicana var. *filiformis* (Torrey) Scribnier - Mexican muhly
Muhlenbergia mexicana var. *mexicana* - Mexican muhly
Muhlenbergia racemosa (Michaux) Britton, Sterns & Poggenburg - satingrass
Muhlenbergia richardsonis (Trinius) Rydberg - mat muhly
Muhlenbergia schreberi J.F. Gmelin - nimblewill
Muhlenbergia sobolifera (Muhl. ex Willd.) Trin. - rock muhly
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from WI, IL, IA; perhaps to be expected on limestone outcrops in the southeast corner
Muhlenbergia sylvatica (Torrey) Torrey ex A. Gray - woodland muhly
Muhlenbergia uniflora (Muhlenberg) Fernald - bog muhly
Munroa squarrosa (Nuttall) Torrey - false buffalograss
erroneous report
Nassella viridula (Trinius) Barkworth - green needlegrass
Oryzopsis asperifolia Michaux - roughleaf ricegrass
Panicum capillare subsp. *capillare* - witchgrass
Panicum dichotomiflorum subsp. *dichotomiflorum* - fall panicgrass
state DOA noxious (CN) weed
Panicum flexile (Gattinger) Scribn. - wiry panicgrass
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from adjacent states
Panicum leucothrix Nash - western panicgrass
misapplied; see *Dichanthelium acuminatum* subsp. *implicatum*
Panicum miliaceum subsp. *miliaceum* - broomcorn millet
Panicum miliaceum subsp. *ruderale* (Kitagawa) Tzvelev - broomcorn millet
reported for Minnesota and Wisconsin by USDA-NRCS but no specimens at MIN

Panicum philadelphicum subsp. *gattingeri* (Nash) Freckmann & Lelong - **Philadelphia panicgrass**

erroneous report

Panicum philadelphicum subsp. *philadelphicum* - **Philadelphia panicgrass**

Panicum virgatum L. - **switchgrass**

vars. not recognized

Pascopyrum smithii (Ryderberg) Barkworth & D.R. Dewey - **western wheatgrass**

probably not native as known to have originated in sagebrush deserts of western North America; considered a possible hybrid of *Leymus triticoides* and *Elymus lanceolatus* fide M. Barkworth

Paspalum setaceum var. *ciliatifolium* (Michx.) Vasey

misapplied to our flora, see var. *muhlenbergii*

Paspalum setaceum var. *muhlenbergii* (Nash) D.J. Banks

Paspalum setaceum var. *stramineum* (Nash) D.J. Banks

possibly occurring in Minnesota, known from WI

Pennisetum glaucum (L.) R. Br. - **yellow foxtail**

misapplied; see *S. pumila*

Phalaris arundinacea L. - **reed canarygrass**

disagreement exists as to whether native or alien forms occur; USDA-NRCS considers this a native

Phalaris canariensis L. - **annual canarygrass**

native to southern Europe; introduced with bird seed but doubtfully persisting

Phleum pratense subsp. *pratense* - **timothy**

native to Europe

Phragmites australis subsp. *americanus* Saltonstall - **American common reed**

Phragmites australis subsp. *australis* - **European common reed**

native to Europe

Piptatherum canadense (Poiret) Dorn - **Canadian ricegrass**

range extension from Manitoba and WI; threatened

Piptatherum pungens (Torrey) Dorn - **mountain ricegrass**

Piptatherum racemosum (Smith) Eaton - **blackseed ricegrass**

Poa alpina subsp. *alpina* - **alpine bluegrass**

report from Lake Co. not yet verified

Poa alsodes A. Gray - **grove bluegrass**

Poa annua L. - **annual bluegrass**

native to Eurasia

Poa arida Vasey - **plains bluegrass**

Poa bulbosa subsp. *vivipara* (Koeler) Arcangeli - **bulbous bluegrass**

native to Europe

Poa chaixii Villars - **broadleaf bluegrass**

native to northern Europe; known only from St. Louis Co. (Duluth area; last collected 1949)

Poa compressa L. - **Canada bluegrass**

native to Europe

Poa glauca Vahl - **glaucous bluegrass**

Poa glauca subsp. *glauca*

Poa interior Rydberg - **inland bluegrass**

Poa nemoralis L. - **wood bluegrass**

native to Europe; known only from St. Louis Co. (last collected 1945)

Poa paludigena Fernald & Wiegand - **bog bluegrass**

northwest limit of distribution; threatened

Poa palustris L. - **waterfowl meadow grass**

Poa pratensis subsp. *agassizensis* (B. Boivin & D. Löve) Roy L. Taylor & MacBryde - **Kentucky bluegrass**

habitat: prairies; possibly native or a derivative of cultivated forms

Poa pratensis subsp. *angustifolia* (L.) Lejeune - **Kentucky bluegrass**

native to western Eurasia

Poa pratensis subsp. *pratensis* - **Kentucky bluegrass**

widespread; probably not native

Poa saltuensis subsp. *languida* (Hitchcock) A. Haines - **oldpasture bluegrass**

Poa saltuensis subsp. *saltuensis* - **oldpasture bluegrass**

Poa secunda subsp. *juncifolia* (Scribnier) Soreng - **Sandberg bluegrass**

reported for Minnesota by USDA-NRCS and NatureServe but no specimens at MIN; known from ND, SD, and Canada

Poa sylvestris A. Gray - **woodland bluegrass**

Poa trivialis subsp. *trivialis* - **rough bluegrass**

native to Europe

Poa wolfii Scribnier - **Wolf's bluegrass**

Polypogon monspeliensis (L.) Desfontaines - **annual rabbitsfoot grass**

native of Europe; known only from Polk Co. (one agricultural population, collected 1993), doubtfully persisting annual

Psathyrostachys juncea (Fischer) Nevski - **Russian wildrye**

native to Asia; known only from Goodhue Co. (one roadside population, collected 1977); reports in FNA from southwestern counties not verified

Puccinellia distans (Jacquin) Parlato - **weeping alkaligrass**

native to Eurasia

Puccinellia nuttalliana (Schultes) Hitchcock - **Nuttall's alkaligrass**

Schedonnardus paniculatus (Nuttall) Trelease - **tumblegrass**

Schedonorus arundinaceus (Schreber) Dumortier - **meadow ryegrass**

native to Eurasia

Schedonorus pratensis (Hudson) P. Beauvois - **meadow fescue**

native to Europe

Schizachne purpurascens (Torrey) Swallen - **false melic**

vars. not recognized

Schizachyrium scoparium (Michaux) Nash
Schizachyrium scoparium var. *scoparium* - little bluestem
Scolochloa festucacea (Willdenow) Link - common rivergrass
Secale cereale L. - cereal rye
native to Europe

Setaria faberii R.A.W. Herrmann - Japanese bristlegrass
native to China; state DOA noxious (CN) weed, 1st collected 1957, Blue Earth Co.

Setaria italica (L.) P. Beauvois - foxtail bristlegrass
native to Eurasia

Setaria lutescens (Weigel) Hubbard - yellow foxtail
misapplied to North American flora, see *S. pumila*

Setaria pumila (Poiret) Roemer & Schultes - yellow foxtail
native to Europe

Setaria pumila subsp. *pallidefusca* (Schumacher) B.K. Simon - cattail grass
known only from Anoka Co. (one road edge population, collected 2010)

Setaria pumila subsp. *pumila*

Setaria verticillata (L.) P. Beauvois - hooked bristlegrass
native to Europe

Setaria viridis var. *major* (Gaudin) Petermann - green bristlegrass

Setaria viridis var. *viridis* - green bristlegrass

Sorghastrum nutans (L.) Nash - Indiangrass

Sorghum bicolor (L.) Moench - Sudan grass

Sorghum bicolor subsp. *drummondii* (Nees ex Steud.) de Wet & Harlan - Sudan grass

Sorghum x alnum Parodi - Columbus grass
(*S. bicolor* x *halepense*) native to Argentina; state DOA listed weed; reported for Minnesota by MN-DNR but no specimens at MIN; known from Dane Co. WI

Spartina gracilis Trinius - alkali cordgrass

Spartina pectinata Link - prairie cordgrass

Sphenopholis intermedia (Rydberg) Rydberg - slender wedgescale

Sphenopholis obtusata (Michaux) Scribner - prairie wedgescale

Sporobolus compositus var. *compositus* - composite dropseed

Sporobolus cryptandrus (Torrey) A. Gray - sand dropseed
vars. not recognized

Sporobolus heterolepis (A. Gray) A. Gray - prairie dropseed

Sporobolus neglectus Nash - puffsheat dropseed

Sporobolus vaginiflorus var. *vaginiflorus* - poverty dropseed

Torreyochloa pallida (Torrey) G.L. Church

Torreyochloa pallida subsp. *fernaldii* (Hitchcock) Dore - Fernald's false mannagrass

Torreyochloa pallida subsp. *pallida* - pale false mannagrass

Tridens flavus var. *flavus* - purpletop tridens
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from IA, WI; perhaps to be expected in the far southeastern corner

Triplasis purpurea var. *purpurea* - purple sandgrass

Trisetum spicatum (L.) K. Richter - spike trisetum

Triticum aestivum L. - common wheat
native to Eurasia; occasionally persisting near cultivated fields

Vulpia octoflora var. *glauca* (Nuttall) Fernald - sixweeks fescue

Vulpia octoflora var. *octoflora* - sixweeks fescue
widespread in US and reported for Minnesota by Flora of the Great Plains 1986 (one specimen from Anoka Co., collected 1915, exhibits some of the characteristics of this variety)

Zea mays L. - corn
occasionally found along roadsides in farming areas; not persisting

Zizania aquatica var. *aquatica* - annual wildrice
reported for western counties by FNA but probably based on misidentified specimens; no specimens at MIN; DNR watch list

Zizania palustris var. *interior* (Fassett) Dore - northern wildrice

Zizania palustris var. *palustris* - northern wildrice

PODOSTEMACEAE

Podostemum ceratophyllum Michaux - hornleaf riverweed
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from ND, Ontario, MI

POLEMONIACEAE

Collomia linearis Nuttall - tiny trumpet

Phlox divaricata subsp. *laphamii* (A.W. Wood) Wherry - Lapham's phlox

Phlox drummondii Hooker - annual phlox
reported for Minnesota by Ownbey & Morley 1991 as a rare garden escape; doubtfully persisting annual

Phlox drummondii subsp. *drummondii* - annual phlox
report unverified

Phlox maculata subsp. *maculata* - wild sweetwilliam

Phlox paniculata L. - fall phlox
native to eastern US

Phlox pilosa subsp. *fulgida* (Wherry) Wherry - downy phlox

Phlox subulata var. *subulata* - moss phlox
native to eastern US

Polemonium caeruleum L. - charity
native to Europe; reported for Minnesota by Lakela 1965 but no specimens at MIN; erroneous record

Polemonium occidentale subsp. *lacustre* Wherry - western polemonium

Polemonium occidentale subsp. *occidentale* - western polemonium

Polemonium reptans var. *reptans* - Greek valerian

POLYGALACEAE

Polygala cruciata var. *aquilonia* Fernald & B.G. Schub. - drumheads

Polygala paucifolia Willdenow - gaywings

Polygala polygama Walter

Polygala sanguinea L. - purple milkwort

Polygala senega L. - Seneca snakeroot

Polygala verticillata var. *isocycla* Fernald - whorled milkwort

POLYGONACEAE

Bistorta vivipara (L.) Delarbre - alpine bistort

reproduces mostly asexually; threatened

Eriogonum annuum Nuttall - annual buckwheat

native to western Great Plains; known only from Sherburne Co. (Sherburne National Wildlife Refuge, last collected 2002), apparently introduced with roadside plantings; vars. not recognized

Fagopyrum esculentum Moench - buckwheat

native to China

Fallopia cilinodis (Michaux) Holub - fringed black bindweed

Fallopia convolvulus (L.) Å. Löve - black-bindweed

native to Eurasia; state DOA noxious (CN) weed, 1st collected 1879, Cook Co.

Fallopia dumetorum (L.) Holub - corpse bindweed

native to Eurasia; to be expected on disturbed sites along forest edges (known from IA, WI and MI); easily and often confused with *F. scandens*

Fallopia japonica var. *japonica* - Japanese knotweed

native to southeast Asia

Fallopia sachalinensis (F. Schmidt) Ronse Decr. - giant knotweed

erroneous report

Fallopia scandens (L.) Holub - climbing false buckwheat

Fallopia x bohemica (Chrtkov & Chrtkov - Bohemian knotweed

(*F. japonica* x *sachalinensis*) native to Europe; known only from St. Louis Co. (Duluth area; collected 2000); easily confused with *Fallopia sachalinensis* and only recently was the hybrid distinguished from the parents

Persicaria amphibia (L.) Delarbre

vars. Not formally recognized in FNA but acknowledged as being somewhat distinctive

Persicaria arifolia (L.) K. Haraldson - halberdleaf tearthumb

Persicaria bungeana (Turczaninow) Nakai - Bunge's smartweed

native to Eurasia

Persicaria careyi (Olney) Greene - Carey's smartweed

western limit of distribution; known only from Carlton Co. (one roadside population, collected 1940); special concern

Persicaria hydropiper (L.) Spach - marshpepper knotweed

native to Europe; oily glands can cause skin irritation; sometimes confused with *P. punctata*

Persicaria hydropiperoides (Michaux) Small - swamp smartweed

highly variable in appearance; rare historic record

Persicaria lapathifolia (L.) A. Gray - curlytop knotweed

highly variable in appearance

Persicaria longiseta (Bruijn) Kitagawa - Oriental lady's thumb

native to eastern Asia; reported for Minnesota by Ownbey & Morley 1991 and FNA but no specimens at MIN

Persicaria maculosa A. Gray - spotted ladysthumb

native to Eurasia, Africa and Oceania; state DOA noxious (CN) weed, 1st collected 1877, Hennepin Co.; may cause skin irritation

Persicaria minor (Huds.) Opiz - pygmy smartweed

Persicaria orientalis (L.) Spach - kiss me over the garden gate

native to India

Persicaria pensylvanica (L.) M. G - Pennsylvania smartweed

state DOA noxious (CN) weed; highly variable in appearance; may cause skin irritation

Persicaria punctata (Elliott) Small - dotted smartweed

vars. not recognized

Persicaria sagittata (L.) H. Gross - arrowleaf tearthumb

highly variable in appearance

Persicaria virginiana (L.) Gaertner - jumpseed

Polygonella articulata (L.) Meisner - coastal jointweed

Polygonum achoreum S.F. Blake - leathery knotweed

Polygonum aviculare subsp. *aviculare* - prostrate knotweed

native to Eurasia

Polygonum aviculare subsp. *buxiforme* (Small) Costea & Tardif - box knotweed

Polygonum aviculare subsp. *depressum* (Meisner) Arcangeli - prostrate knotweed

native to Europe

Polygonum aviculare subsp. *neglectum* (Besser) Arcangeli - narrowleaf knotweed

native to Europe

Polygonum aviculare subsp. *rurivagum* (Jord. ex Boreau) Berher

native to Europe; possibly occurring in Minnesota, known from IA, SD, and Manitoba, in disturbed places

Polygonum cespitosum Blume - Oriental lady's thumb

misapplied; see *Persicaria longiseta*

Polygonum douglasii Greene - Douglas' knotweed

Polygonum erectum L. - erect knotweed

Polygonum ramosissimum subsp. *prolificum* (Small) Costea & Tardif - bushy knotweed

known only from Lac Qui Parle Co. (collected 1988)

Polygonum ramosissimum subsp. *ramosissimum* - bushy knotweed

Polygonum tenuie Michaux - pleatleaf knotweed

Rheum rhabarbarum L. - garden rhubarb

native to Siberia; leaf blades toxic to humans and animals

Rumex acetosa L. - garden sorrel

native to Eurasia and northwest Africa; occasionally misidentified as *R. acetosella*

Rumex acetosella L. - common sheep sorrel

native to Eurasia; polyploid complex highly variable in appearance

Rumex alpinus L. - Munk's rhubarb

native to Eurasia; known only from Cook Co. (collected 1977); only known specimen at UWSP

Rumex altissimus Alph. Wood - pale dock

Rumex britannica L. - greater water dock

Rumex crispus L. - curly dock

native to Eurasia; state DOA noxious (CN) weed, 1st collected 1877, Hennepin Co.

Rumex fueginus Philippi - golden dock

frequently misidentified as *R. maritimus*, which is native to Eurasia and known from Alaska and Yukon and rarely appearing elsewhere

Rumex longifolius de Candolle - dooryard dock

native to Eurasia

Rumex mexicanus

misapplied to our flora, see *R. triangulivalvis*; misapplied name

Rumex obtusifolius L. - bitter dock

native to Eurasia; vars. not recognized

Rumex occidentalis S. Watson - western dock

reported for Minnesota by FNA but no specimens at MIN; known from Canada, ND, SD, IA; weedy in wet meadows, bogs, on riverbanks, etc.; frequently misidentified as *R. longifolius*

Rumex patientia L. - patience dock

native to Eurasia

Rumex pseudonatronatus (Borb - field dock

native to Eurasia; frequently misidentified as *R. longifolius* and sometimes *R. crispus*

Rumex salicifolius Weinmann

misapplied to our flora, see *R. triangulivalvis*; misapplied name

Rumex sanguineus L. - redvein dock

native to Africa and Eurasia; reported for Minnesota by USDA-NRCS but no specimens at MIN

Rumex stenophyllus Ledebour - narrowleaf dock

native to Eurasia

Rumex thrysiflorus Fingerhuth - garden sorrel

Rumex triangulivalvis (Danser) Rechinger f. - Mexican dock

Rumex venosus Pursh - veiny dock

reported for Minnesota by FNA but no specimens at MIN, known from Manitoba, ND, SD, IA and WI; weedy on sandy soils

Rumex verticillatus L. - swamp dock

Rumex x pratensis Mert. & W.D.J. Koch - dock

(*R. crispus* x *obtusifolius*) common sterile hybrid to be expected on disturbed sites if parents nearby

POLYPODIACEAE

Polypodium virginianum L. - rock polypody

PONTEDERIACEAE

Eichhornia crassipes (Mart.) Solms - common water hyacinth

native to tropical Americas; reported from Winona Co. but no specimens at MIN; unlikely to be persisting

Heteranthera dubia (Jacquin) MacMillan - grassleaf mudplantain

Heteranthera limosa (Swartz) Willdenow - blue mudplantain

northeast limit of distribution; threatened

Pontederia cordata L. - pickerelweed

PORTULACACEAE

Claytonia caroliniana Michaux - Carolina springbeauty

Claytonia virginica L. - Virginia springbeauty

Montia chamissoi (Ledebour ex Sprengel) Greene - water minerslettuce

disjunct from Rocky Mountains; known only from Winona Co. (last collected 1985); endangered

Phemeranthus parviflorus (Nuttall) Kiger - sunbright

Phemeranthus rugospermus (Holzinger) Kiger - prairie fameflower

northern limit of distribution; threatened

Portulaca grandiflora Hooker - rose moss

native to South America; reported for Minnesota by FNA as escaping and naturalizing near gardens but no specimens at MIN

Portulaca oleracea L. - little hogweed

native to southern Eurasia; some ethnobotanical evidence suggests this was introduced to the western hemisphere in pre-Columbian times; known to have the highest antioxidant content of any leafy plant examined

POTAMOGETONACEAE

Potamogeton alpinus Balbis - alpine pondweed

vars. not recognized

Potamogeton amplifolius Tuckerman - largeleaf pondweed

Potamogeton bicupulatus Fernald - snailseed pondweed

disjunct from eastern US; endangered

Potamogeton confervoides Reichenbach - Tuckerman's pondweed

westernmost limit of distribution, known only from St. Louis Co. (one population, collected 2008); DNR watch list

Potamogeton crispus L. - curly pondweed

native to Eurasia; on MN-DNR Prohibited Invasive Species list; 1st collected 1929, Hennepin Co.

Potamogeton diversifolius Rafinesque - waterthread pondweed

Potamogeton epihydrus - ribbonleaf pondweed

Potamogeton foliosus subsp. foliosus - leafy pondweed

Potamogeton friesii Ruprecht - Fries' pondweed

Potamogeton gramineus L. - variableleaf pondweed

Potamogeton illinoensis Morong - Illinois pondweed

Potamogeton lateralis

invalid name

Potamogeton natans L. - floating leaf pondweed

Potamogeton nodosus Poiret - longleaf pondweed

Potamogeton oakesianus J.W. Robbins - Oakes' pondweed

Potamogeton obtusifolius Mertens & W.D.J. Koch - bluntleaf pondweed

Potamogeton praelongus Wulfen - whitestem pondweed

Potamogeton pulcher Tuckerman - spotted pondweed

disjunct from eastern US; known only from Clearwater (collected 2007) and Cass (collected 2008) counties; endangered

Potamogeton pusillus subsp. pusillus - small pondweed

Potamogeton pusillus subsp. tenuissimus (Mertens & W.D.J. Koch) R.R. Haynes & Hellquist - small pondweed

Potamogeton richardsonii (A. Bennett) Rydberg - Richardson's pondweed

Potamogeton robbinsii Oakes - Robbins' pondweed

Potamogeton spirillus Tuckerman - spiral pondweed

Potamogeton strictifolius A. Bennett - narrowleaf pondweed

Potamogeton vaseyi J.W. Robbins - Vasey's pondweed

Potamogeton x haynesii Hellquist & G.E. Crow - Haynes' hybrid pondweed

(P. strictifolius x zosteriformis) only known specimen from Minnesota deposited at GH; DNR watch list

Potamogeton x spathuliformis (J.W. Robbins) Morong - spoon pondweed

(P. gramineus x illinoensis) known only from St. Louis Co. (collected 1940)

Potamogeton zosteriformis Fernald - flatstem pondweed

Stuckenia filiformis (Persoon) Brner - fine leaf pondweed

Stuckenia filiformis subsp. alpina (Blytt) R.R. Haynes, Les, & M. Král - fineleaf pondweed

Stuckenia filiformis subsp. occidentalis (J.W. Robbins) R.R. Haynes, Les, & M. Král - western fineleaf pondweed

Stuckenia pectinata (L.) Brner - sago pondweed

Stuckenia vaginata (Turczaninow) Holub - sheathed pondweed

last collected 1956; endangered

Zannichellia palustris L. - horned pondweed

PRIMULACEAE

Androsace occidentalis Pursh - western rockjasmine

Androsace septentrionalis L. - pygmyflower rockjasmine

Dodecatheon amethystinum (Fassett) Fassett - jeweled shooting star

Dodecatheon meadia L. - pride of Ohio

Dodecatheon radicum Greene - jeweled shooting star

misapplied; see D. amethystinum

Primula fassettii Mast & Reveal - jeweled shooting star

known only from the southeastern corner of the state; DNR watch list

Primula meadia (L.) Mast & Reveal - pride of Ohio

known only from Mower Co. (last collected 2008); endangered

Primula mistassinica Michaux - Mistassini primrose

vars. not recognized

PTERIDACEAE

Adiantum pedatum L. - northern maidenhair

Cheilanthes feei T. Moore - slender lip fern

Cheilanthes lanosa (Michaux) D.C. Eaton - hairy lip fern

erroneous report

Cryptogramma stelleri (S.G. Gmelin) Prantl - fragile rockbrake

Pellaea atropurpurea (L.) Link - purple cliffbrake

Pellaea glabella subsp. glabella - smooth cliffbrake

RANUNCULACEAE

Aconitum napellus L. - Venus' chariot

native to Europe; known only from Cook Co. (one roadside population, collected 1975)

Actaea pachypoda Elliott - doll

plants poisonous

Actaea rubra (Aiton) Willdenow - red baneberry

plants poisonous

Anemone acutiloba (de Candolle) G. Lawson - sharplobe hepatica

Anemone americana (de Candolle) H. Hara - roundlobe hepatica

Anemone canadensis L. - Canadian anemone

Anemone caroliniana Walter - Carolina anemone

Anemone cylindrica A. Gray - candle anemone

Anemone halleri Allioni - glacial anemone

native to Europe; known only from Freeborn Co. (collected 1995), introduced with prairie planting

***Anemone multifida* var. *multifida* - Pacific anemone**

known only from Mahnomen Co. (one roadside population, collected 1941); rare historic record

Anemone patens* L. - cutleaf anemone**Anemone patens* var. *multifida* Pritzel - eastern pasque flower**

plants poisonous

Anemone quinquefolia* var. *quinquefolia* - two leaf anemone**Anemone virginiana* var. *alba* (Oakes) A.W. Wood - tall thimbleweed*****Anemone virginiana* var. *cylindroidea* B. Boivin - tall thimbleweed**

reported for northern Minnesota by FNA but no specimens at MIN; known from Saskatchewan and Manitoba

Anemone virginiana* var. *virginiana* - tall thimbleweed**Aquilegia brevistyla* Hooker - smallflower columbine**

erroneous report

Aquilegia canadensis* L. - red columbine**Aquilegia vulgaris* L. - European columbine**

native to Europe; reported as a garden escape for Minnesota by FNA but no specimens at MIN

***Caltha natans* Pallas - floating marsh marigold**

southern limit of distribution; known only from St. Louis Co.; endangered

Caltha palustris* L. - yellow marsh marigold**Ceratocephala testiculata* (Crantz) Roth - curveseed buttercup**

invalid name; see *Ranunculus testiculatus*

Clematis occidentalis* (Hornem.) DC.**Clematis occidentalis* var. *occidentalis* - western blue virginsbower*****Clematis recta* L. - ground virginsbower*****Clematis terniflora* de Candolle - sweet autumn virginsbower**

native to eastern Asia; known only from St. Louis Co. (Duluth area; one population, collected 1995)

***Clematis virginiana* L. - devil's darning needles**

may cause severe skin irritation

***Consolida ajacis* (L.) Schur - doubtful knight's-spur**

native to Europe and Africa; known only from Polk Co. (historical collection from 1882), probably escaped from cultivation

Coptis trifolia* (L.) Salisbury - threeleaf goldthread**Delphinium carolinianum* Walter*****Delphinium carolinianum* subsp. *virescens* (Nuttall) R.E. Brooks - Carolina larkspur**

plants toxic

***Delphinium tricorne* Michx. - dwarf larkspur**

erroneous report

Enemion biternatum* Rafinesque - eastern false rue anemone**Hydrastis canadensis* L. - goldenseal**

northwest limit of distribution; endangered

Myosurus minimus* L. - tiny mousetail**Nigella damascena* L. - devil in the bush**

native to Eurasia; known only from Otter Tail Co. (one weedy population, collected 1975), doubtfully persisting

***Ranunculus abortivus* L. - littleleaf buttercup**

toxic to livestock

***Ranunculus acris* L. - tall buttercup**

native to Eurasia; state DOA noxious (CN) weed, 1st collected 1878, St. Louis Co.; can cause severe skin irritation, toxic to livestock

***Ranunculus ambigens* S. Wats. - waterplantain spearwort**

erroneous report

Ranunculus aquatilis* L.**Ranunculus aquatilis* var. *diffusus* Withering - white water crowfoot*****Ranunculus cymbalaria* Pursh - alkali buttercup**

plant highly toxic

Ranunculus fascicularis* Muhlenberg ex J.M. Bigelow - early buttercup**Ranunculus flabellaris* Rafinesque - yellow water buttercup*****Ranunculus flammula* var. *flammula* - greater creeping spearwort**

erroneous report

Ranunculus flammula* var. *ovalis* (J.M. Bigelow) L.D. Benson - greater creeping spearwort**Ranunculus flammula* var. *reptans* (L.) E. Meyer - greater creeping spearwort*****Ranunculus gmelinii* de Candolle - Gmelin's buttercup**

vars. not recognized

Ranunculus hispidus* var. *caricetorum* (Greene) T. Duncan - bristly buttercup**Ranunculus hispidus* var. *nitidus* (Chapman) T. Duncan - bristly buttercup*****Ranunculus lapponicus* L. - Lapland buttercup*****Ranunculus macounii* Britton - Macoun's buttercup*****Ranunculus pedatifidus* Sm. - surefoot buttercup**

erroneous report

Ranunculus pensylvanicus* L. f. - Pennsylvania buttercup**Ranunculus recurvatus* var. *recurvatus* - blisterwort*****Ranunculus repens* L. - creeping buttercup**

native to Eurasia

Ranunculus rhomboideus* Goldie - Labrador buttercup**Ranunculus sceleratus* var. *multifidus* Nuttall - cursed buttercup*****Ranunculus sceleratus* var. *sceleratus* - cursed buttercup*****Ranunculus testiculatus* Crantz - curveseed buttercup**

native to Eurasia; known only from Rock Co. (collected 1981), doubtfully persisting

***Thalictrum confine* Fernald**

misapplied to our flora, see *T. venulosum*; misapplied name

Thalictrum dasycarpum Fischer & Ay Lallemand - **purple meadow-rue**

Thalictrum dioicum L. - **early meadow-rue**

Thalictrum revolutum de Candolle - **waxyleaf meadow-rue**

although FNA indicates this as widespread in northeastern Minnesota, it is known only from two populations in Lake (collected 1914) and St. Louis (collected 2008) counties; DNR watch list

Thalictrum thalictroides (L.) A.J. Eames & B. Boivin - **rue anemone**

roots possibly toxic

Thalictrum venulosum Trelease - **veiny meadow-rue**

RESEDAEAE

Reseda lutea L. - **yellow mignonette**

native to Mediterranean; invasive and possibly to be expected on disturbed sites; known from Saskatchewan, Manitoba, Ontario, IA, KS, NE, WI

RHAMNACEAE

Ceanothus americanus L.

Ceanothus herbaceus Rafinesque - **Jersey tea**

Ceanothus masonii McMinn - **Mason's ceanothus**
erroneous report

Frangula alnus Mill. - **glossy buckthorn**

Rhamnus alnifolia L'Hritier - **alderleaf buckthorn**

Rhamnus cathartica L. - **common buckthorn**

native to Eurasia; state DOA restricted noxious weed, 1st collected 1937, Hennepin Co.

Rhamnus davurica Pall. - **Dahurian buckthorn**

Rhamnus frangula L. - **glossy buckthorn**

native to Europe; state DOA restricted noxious weed, 1st collected 1938, Ramsey Co.

ROSACEAE

Agrimonia eupatoria L. - **churchsteeple**

native of Eurasia; reported for Minnesota by USDA-NRCS but no specimens at MIN; erroneous record

Agrimonia gryposepala Wallroth - **tall hairy agrimony**

Agrimonia pubescens Wallroth - **soft agrimony**

Agrimonia rostellata Wallr. - **beaked agrimony**

erroneous report

Agrimonia striata Michaux - **roadside agrimony**

Amelanchier alnifolia var. ***alnifolia*** (Nutt.) Nutt. ex M. Roem. - **Saskatoon serviceberry**

Amelanchier arborea (Michaux f.) Fernald - **common serviceberry**

Amelanchier bartramiana (Tausch) M. Roemer - **oblongfruit serviceberry**

Amelanchier bartramiana x ***laevis***

Amelanchier humilis Wiegand - **low serviceberry**

Amelanchier interior E.L. Nielsen - **Pacific serviceberry**

Amelanchier intermedia Spach - **serviceberry**

Amelanchier laevis Wiegand - **Allegheny serviceberry**

Amelanchier sanguinea (Pursh) DC. - **roundleaf serviceberry**

Amelanchier spicata (Lamarck) K. Koch - **running serviceberry**

Amelanchier x neglecta Eggleston ex G.N. Jones

(*A. bartramiana* x *laevis*); invalid name

Aronia melanocarpa (Michaux) Elliott - **black chokeberry**

Chamaerhodos erecta (L.) Bunge - **Nuttall's little rose**

Chamaerhodos erecta subsp. ***nuttallii*** (Pickering ex Rydb.) Hultén - **Nuttall's little rose**

Comarum palustre L. - **purple marshlocks**

Cotoneaster lucidus Schlechtendal - **shiny cotoneaster**

Crataegus calpodendron (Ehrhart) Medikus - **pear hawthorn**

all sources - recognized taxon for Minnesota; special concern

Crataegus chrysocarpa var. ***blanchardii*** (Sarg.) J.B. Phipps - **fireberry hawthorn**

Crataegus chrysocarpa var. ***chrysocarpa*** - **red haw**

All sources - recognized taxon for Minnesota

Crataegus chrysocarpa var. ***faxonii*** (Sargent) Eggleston - **fireberry hawthorn**

Crataegus chrysocarpa var. ***praecox*** (Sarg.) J.B. Phipps - **fireberry hawthorn**

Crataegus chrysocarpa var. ***vigintistamina*** J.B. Phipps - **fireberry hawthorn**

Crataegus coccinea var. ***pringlei*** - **red haw**

Crataegus crus-galli L. - **cockspur hawthorn**

USDA-NRCS 2009 - recognized taxon for Minnesota; NatureServe 2009 - recognized taxon for Minnesota; ITIS 2009 - recognized taxon; Smith 2008 - not included; erroneous record

Crataegus douglasii Lindley - **black hawthorn**

disjunct from western North America; All sources - recognized taxon for Minnesota; special concern

Crataegus flabellata (Bosc ex Spach) K. Koch - **fanleaf hawthorn**

USDA-NRCS 2009 - recognized taxon for Minnesota; NatureServe 2009 - recognized taxon but not reported for Minnesota; ITIS 2009 - recognized taxon; Smith 2008 - not included; erroneous record

Crataegus fluviatilis Sargent - **bigfruit hawthorn**

Crataegus holmesiana Ashe - **Holme's hawthorn**

USDA-NRCS 2009 - recognized taxon for Minnesota; NatureServe 2009 - recognized taxon for Minnesota; ITIS 2009 - recognized taxon; Smith 2008 - not included

Crataegus irrasa Sarg. - **Blanchard's hawthorn**

USDA-NRCS 2009 - recognized taxon for Minnesota - erroneously (based on misidentified specimens); NatureServe 2009 - recognized taxon for Minnesota; ITIS 2009 - recognized taxon; Smith 2008 - not included

Crataegus laurentiana Sarg. - Laurentian hawthorn
Crataegus macracantha Loddiges - fleshy hawthorn
USDA-NRCS 2009 – synonym for *C. succulenta*; NatureServe 2009 – not included; ITIS 2009 – synonym for *C. succulenta*; Smith 2008 – recognized taxon for Minnesota

Crataegus macrosperma Ashe - big fruit hawthorn
All sources - recognized taxon for Minnesota

Crataegus mollis* var. *mollis - downy hawthorn
All sources - recognized taxon for Minnesota

Crataegus monogyna* var. *monogyna - oneseed hawthorn

Crataegus punctata Jacquin - dotted hawthorn
All sources - recognized taxon for Minnesota

Crataegus scabrida Sargent - rough hawthorn
known only from Houston Co. (Spring Grove area, historical collections from 1902); USDA-NRCS 2009 – recognized taxon for Minnesota; NatureServe 2009 – recognized taxon for Minnesota; ITIS 2009 – recognized taxon; Smith 2008 – recognized taxon for Minnesota; rare historic record

Crataegus schuettei* var. *cuneata Kruschke ex J.B. Phipps - Schuette's hawthorn

Crataegus sheridana A. Nels. - Great Plains fireberry hawthorn
USDA-NRCS 2009 – not included; NatureServe 2009 – not included; ITIS 2009 – not included; Smith 2008 – recognized taxon for Minnesota; known only from Big Stone Co. (collected 1997) but no specimens at MIN

Crataegus submollis Sargent - Quebec hawthorn
All sources – recognized taxon for Minnesota

Crataegus succulenta* var. *succulenta
All sources – recognized taxon for Minnesota

Dasiphora fruticosa (L.) Rydb. - shrubby cinquefoil

Drymocallis arguta (Pursh) Rydb. - tall cinquefoil

Filipendula rubra (Hill) B.L. Robinson - queen of the prairie
native from IA, IL, WI and eastward; known only from St. Louis Co. (Duluth area; one population collected 2009); only specimen at DUL

Filipendula ulmaria (L.) Maximowicz - queen of the meadow
native to Eurasia; known only from St. Louis Co.

Fragaria vesca* subsp. *americana Porter - woodland strawberry

Fragaria virginiana* subsp. *glaucia (S. Watson) Staudt - Virginia strawberry

Fragaria virginiana* subsp. *virginiana - Virginia strawberry

Geum aleppicum Jacquin - yellow avens

Geum canadense* var. *canadense - white avens

Geum laciniatum Murray - rough avens

Geum macrophyllum* var. *macrophyllum - largeleaf avens

Geum macrophyllum* var. *perincisum (Rydberg) Raup - largeleaf avens

Geum rivale L. - water avens

Geum triflorum* var. *triflorum - old man's whiskers

Geum virginianum L. - cream avens
reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IL

Malus baccata* var. *baccata - Siberian crab apple

Malus floribunda Siebold ex Van Houtte - Japanese flowering crab apple

Malus ioensis* var. *ioensis - prairie crab apple

Malus prunifolia (Willd.) Borkh. - plumleaf crab apple

Malus pumila Mill. - paradise apple

Malus x soulardii (L.H. Bailey) Britton - Soulard crab
(*Malus ioensis* × *pumila*); erroneous record

Physocarpus intermedius (Rydb.) C.K. Schneid. - Atlantic ninebark

Physocarpus opulifolius (L.) Maximowicz - common ninebark

Potentilla anserina L. - silverweed cinquefoil

Potentilla anserina* subsp. *anserina - silverweed cinquefoil

Potentilla argentea L. - silver cinquefoil
native to Europe

Potentilla bipinnatifida Douglas - tansy cinquefoil

Potentilla effusa* var. *effusa - branched cinquefoil
report unverified

Potentilla fruticosa L. - shrubby cinquefoil
invalid name; see *Dasiphora fruticosa*

Potentilla gracilis* var. *fastigiata (Nuttall) S. Watson - slender cinquefoil

Potentilla gracilis* var. *flabelliformis (Lehm.) Nutt. - slender cinquefoil

Potentilla hippiana* var. *hippiana - woolly cinquefoil

Potentilla inclinata Vill. - ashy cinquefoil
report unverified

Potentilla intermedia L. - downy cinquefoil
native to Eurasia; known only from St. Louis Co. (one resort-grounds population, collected 1950)

Potentilla lasiodonta Rydberg - sandhills cinquefoil
known only from Polk Co. (collected 1940)

Potentilla littoralis Rydberg - coast cinquefoil

Potentilla norvegica L. - Norwegian cinquefoil

Potentilla pensylvanica L. - Pennsylvania cinquefoil

Potentilla pensylvanica* var. *arida B. Boivin - Pennsylvania cinquefoil
some authors include this in var. *bipinnatifida*

Potentilla pulcherrima Lehmann - beautiful cinquefoil
erroneous report

Potentilla recta L. - sulphur cinquefoil

native to Europe

Potentilla reptans L. - creeping cinquefoil

report unverified

Potentilla rivalis Nuttall - brook cinquefoil

Potentilla simplex Michaux - common cinquefoil

Potentilla supina subsp. paradoxoa (Nutt.) Soják - paradox cinquefoil

Potentilla verna L. - spring cinquefoil

native to Europe; known only from Hennepin Co. (roadside ground cover, collected 2009)

Prunus americana Marshall - American plum

seeds contain cyanide compounds

Prunus domestica L. - European plum

seeds contain cyanide compounds; cultigen from Europe reported as naturalized by MN-DNR (no further data) and USDA-NRCS but no specimens at MIN

Prunus nigra Aiton - Canadian plum

seeds contain cyanide compounds

Prunus pensylvanica L. f. - pin cherry

seeds contain cyanide compounds

Prunus pumila var. besseyi (L.H. Bailey) Gleason - western sandcherry

Prunus pumila var. pumila - Great Lakes sandcherry

Prunus pumila var. susquehanae (Willd.) H. Jaeger - Susquehanna sandcherry

Prunus serotina var. serotina - black cherry

bark, leaves, seeds highly toxic to humans, pets, and especially ruminant livestock

Prunus tomentosa Thunberg - Nanking cherry

seeds contain cyanide compounds; native to Asia; naturalized only in Washington Co. (one population, collected 1990)

Prunus triloba Lindley - flowering plum

erroneous report

Prunus virginiana var. virginiana - choke cherry

bark, leaves, seeds highly toxic to humans and livestock

Pyrus ussuriensis Maxim. - Chinese pear

Rosa acicularis subsp. *sayi* (Schweinitz) W.H. Lewis - prickly rose

Rosa arkansana Porter - prairie rose

Rosa blanda Aiton - smooth rose

Rosa canina L. - dog rose

Rosa carolina var. *carolina* - Carolina rose

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IA, NE

Rosa eglanteria L.

rejected name; see Rosa rubiginosa; native to Europe; known only from Dakota Co. (one population, collected 1943)

Rosa ferruginea Vill. - redleaf rose

see Rosa glauca; uncertain status

Rosa glauca Pourr. - redleaf rose

Rosa multiflora Thunberg - multiflora rose

native to eastern Asia; known only from Washington Co., reports from national forests unconfirmed

Rosa rubiginosa L.

Rosa rubiginosa var. *rubiginosa* - sweetbrier rose

Rosa rugosa Thunberg - rugosa rose

native to eastern Asia; known only from Itasca Co. (one roadside population, collected 1958)

Rosa setigera Michx. - climbing rose

Rosa spinosissima L. - Scotch rose

Rosa woodsii Lindl.

Rosa woodsii var. *woodsii* - Wood's rose

Rosa x dulcissima Lunell - hybrid rose

(R. acicularis x Woodsii)

Rosa x housei (Erlanson) - House's hybrid rose

(R. acicularis x blanda)

Rubus acridens L.H. Bailey -

FNA (in press) – not included; USDA-NRCS 2009 – synonym for R. kennedyanus; NatureServe 2009 – not recognized; ITIS 2009 – synonym for R. kennedyanus; Smith 2008 – recognized taxon for Minnesota (reported only from northern MN)

Rubus allegheniensis Porter - Allegheny blackberry

All sources – recognized taxon for Minnesota (widespread in eastern US)

Rubus arcticus subsp. *acaulis* (Michaux) Fock - dwarf raspberry

All sources – recognized taxon for Minnesota (widespread across Canada with extensions into US)

Rubus canadensis L. - smooth blackberry

All sources – recognized taxon for Minnesota (common in eastern US and Canada)

Rubus canadensis x *pensylvanicus* - showy blackberry

Rubus chamaemorus L. - cloudberry

common across Canada with few extensions into US; All sources – recognized taxon for Minnesota; reported only from the northeastern counties; threatened

Rubus flagellaris Willdenow - northern dewberry

All sources – recognized taxon for Minnesota; but species limits defined differently among authors. The common view of this taxon (Alice et al. in press) is of an extremely polymorphic taxon when considered across its range in the US and Canada, with some characters subject to environmental variation and intergradation common among the variants. Some authors (e.g., Smith 2008) take a restricted view and separate the variants as species.

Rubus hispida L. - bristly dewberry

FNA (in press) – recognized taxon for Minnesota; known only from along one portage trail in St. Louis Co. (collected 1952); USDA-NRCS 2009 – recognized taxon but not reported for Minnesota; NatureServe 2009 – recognized taxon but not reported for Minnesota; erroneous record

Rubus idaeus subsp. *idaeus* - American red raspberry

FNA (in press) – recognized taxon for Minnesota (as garden escape; no specimens at MIN); USDA-NRCS 2009 – recognized taxon for Minnesota (common across much of North America); NatureServe 2009 – recognized taxon for Minnesota; ITIS 2009 – recognized taxon; Smith 2008 – not included

Rubus idaeus subsp. *strigosus* (Michx.) Focke - grayleaf red raspberry

All sources – recognized taxon for Minnesota (Smith 2008 – synonym for var. *strigosus* (Michx.) Maxim.)

***Rubus illecebrosus* Focke - strawberry raspberry**

native to Japan; FNA (in press) – recognized taxon (but known only from Nova Scotia and Quebec south to WV); USDA-NRCS 2009 – recognized taxon for Minnesota – probably erroneously (no specimens at MIN); NatureServe 2009 – recognized taxon for Minnesota; ITIS 2009 – recognized taxon; Smith 2008 – not included

***Rubus occidentalis* L. - black raspberry**

All sources – recognized taxon for Minnesota (common in eastern half of US and Canada)

***Rubus parviflorus* Nuttall - thimbleberry**

All sources – recognized taxon for Minnesota (common in western US and Canada; disjunct in central states and provinces)

***Rubus pensylvanicus* Poiret - Pennsylvania blackberry**

FNA (in press) – recognized taxon for Minnesota; USDA-NRCS 2009 – recognized taxon for Minnesota (common in eastern US and Canada); NatureServe 2009 – recognized taxon for Minnesota; ITIS 2009 – recognized taxon for Minnesota; Smith 2008 – not included

***Rubus pergratus* Blanch. - smooth blackberry**

FNA (in press) – synonym for *R. canadensis*; USDA-NRCS 2009 – recognized taxon for Minnesota (common from MN and IA to ME and VA); NatureServe 2009 – recognized taxon for Minnesota; ITIS 2009 – recognized taxon; Smith 2008 – not included

***Rubus pubescens* Rafinesque - dwarf red blackberry**

All sources – recognized taxon for Minnesota (widespread in northern US and Canada)

***Rubus quaeasitus* L.H. Bailey**

FNA (in press) – not included; USDA-NRCS 2009 – not included; NatureServe 2009 – not included; ITIS 2009 – recognized taxon; Smith 2008 – recognized taxon for Minnesota (known only from Carlton Co. and disjunct with WI populations from eastern provinces of Prince Edward Island and New Brunswick; erroneous record

***Rubus repens* L. - robin runaway**

report unverified

***Rubus setosus* Bigelow - setose blackberry**

All sources – recognized taxon for Minnesota (Smith 2008 recognizes *R. setosus* var. *rotundior* Bailey as a synonym for *R. fulleri*)

***Rubus superioris* L.H. Bailey - Vermont blackberry**

FNA (in press) – synonym for *R. setosus*; USDA-NRCS 2009 – synonym for *R. vermontanus*; NatureServe 2009 – not included; ITIS 2009 – synonym for *R. vermontanus*; Smith 2008 – recognized taxon for Minnesota

***Rubus x elegantulus* Blanchard - showy blackberry**

report unverified; (*Rubus canadensis* x *pensylvanicus*)

***Rubus x multiflora* Blanch. - variable blackberry**

(*Rubus flagellaris* x *setosus*)

***Rubus x neglectus* Peck**

(*Rubus idaeus* *strigosus* x *occidentalis*)

***Rubus x paracaulis* L.H. Bailey - blackberry**

(*Rubus arcticus* *acaulis* x *pubescens*)

***Rubus x rosendahlii* - northern dewberry**

(*Rubus flagellaris* x *hispidus*)

***Rubus x wisconsinensis* L.H. Bailey - Wisconsin blackberry**

(*Rubus pensylvanicus* x *setosus*)

***Sanguisorba minor* Scop. - small burnet**

***Sanguisorba officinalis* L. - great burnet**

native to the west coast; reported for Minnesota by USDA but no specimens at MIN

***Sibbaldiopsis tridentata* (Aiton) Rydb. - shrubby five-fingers**

***Sorbaria sorbifolia* (L.) A. Braun - false spirea**

***Sorbus americana* Marshall - American mountain ash**

***Sorbus aucuparia* L. - European mountain ash**

native to Europe

***Sorbus decora* (Sargent) C.K. Schneider - northern mountain ash**

***Spiraea alba* var. *alba* - white meadowsweet**

***Spiraea alba* var. *latifolia* (Aiton) Dippel - white meadowsweet**

***Spiraea lucida* Douglas ex Greene - shinyleaf spirea**

report unverified

***Spiraea tomentosa* var. *rosea* (Rafinesque) Fernald - steeplebush**

***Spiraea x billiardii* Hort. ex K. Koch - triumphans bilerd spirea**

(*S. douglasii* x *salicifolia*) native to Allegheny region; known only from St. Louis Co. (Duluth area; collected 2008)

***Spiraea x vanhouttei* (Briot) Carré - spiraea**

(*S. cantoniensis* x *trilobata*) native to Europe

***Waldsteinia fragarioides* (Michaux) Trattinnick - Appalachian barren strawberry**

RUBIACEAE

***Cephalanthus occidentalis* L. - common buttonbush**

poisonous to livestock

***Galium aparine* L. - stickywilly**

***Galium asprellum* Michaux - rough bedstraw**

***Galium boreale* L. - northern bedstraw**

***Galium brevipes* - limestone swamp bedstraw**

***Galium circaeans* var. *hypomalacum* Fernald - licorice bedstraw**

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IA, NE; DNR watch list

***Galium concinnum* Torrey & A. Gray - shining bedstraw**

***Galium labradoricum* (Wiegand) Wiegand - northern bog bedstraw**

***Galium lanceolatum* Torr. - lanceleaf wild licorice**

reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, Barron and Washburn counties of WI and IL

***Galium mollugo* L. - false baby's breath**

native to Europe

***Galium obtusum* var. *obtusum* - bluntnleaf bedstraw**

***Galium odoratum* (L.) Scopoli - sweetscented bedstraw**

native to Europe; known only from Washington Co. (one population, collected 1983)

Galium tinctorium L. - stiff marsh bedstraw
Galium trifidum subsp. *trifidum* L. - threepetal bedstraw
Galium triflorum Michaux - fragrant bedstraw
Galium verum var. *verum* - yellow spring bedstraw
native to Eurasia
Houstonia canadensis Willd. ex Roem. & Schult. - Canadian summer bluet
erroneous report
Houstonia longifolia Gaertn. - longleaf summer bluet
Mitchella repens L. - partridgeberry

RUPPIACEAE

Ruppia cirrhosa (Petagna) Grande - spiral ditchgrass
Ruppia maritima L. - spiral ditchgrass
misapplied to our flora, see R. cirrhosa

RUSCACEAE

Convallaria majalis var. *majalis* - European lily of the valley
native to Eurasia; roots and flowers toxic to pets, potentially toxic to humans
Maianthemum canadense Desfontaines - Canada mayflower
vars. Not recognized
Maianthemum canadense (L.) Link
Maianthemum canadense subsp. *racemosum* - feathery false lily of the valley
Maianthemum stellatum (L.) Link - starry false lily of the valley
Maianthemum trifolium (L.) Sloboda - threeleaf false lily of the valley
Polygonatum biflorum (Walter) Elliott - smooth solomon's seal
Polygonatum pubescens (Willdenow) Pursh - hairy solomon's seal

RUTACEAE

Phellodendron amurense Rupr. - amur corktree
erroneous report
Phellodendron chinense C.K. Schneider - Chinese corktree
native to China; known only from Winona Co. (collected 2001), probably naturalized from nearby ornamental plantings; can become invasive
Ptelea trifoliata var. *trifoliata* - common hoptree
native to south and eastern US; known only from Hennepin Co. (last collected 1951), possibly planted
Zanthoxylum americanum Miller - common pricklyash

SALICACEAE

Populus alba L. - white poplar
native to Eurasia; possibly can become invasive as spreads by root suckering
Populus balsamifera L. - balsam poplar
Populus deltoides subsp. *monilifera* (Aiton) Eckenwalder - plains cottonwood
Populus grandidentata Michaux - big-tooth aspen
Populus nigra L. - Lombardy poplar
native to Eurasia; probably planted
Populus tremuloides Michaux - quaking aspen
Populus x canadensis Moench - Carolina poplar
(*P. deltoides* x *nigra*) reported for Minnesota by Flora of the Great Plains 1986 but no specimens at MIN
Populus x canescens (Aiton) Smith - gray poplar
(*P. alba* x *tremula*) native to Eurasia; reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IL
Populus x jackii Sargent - balm-of-Gilead
(*P. balsamifera* x *deltoides*) rarely produces viable seed
Populus x smithii B. Boivin - cottonwood
(*P. grandidentata* x *tremuloides*) reported for Minnesota by Ownbey and Morley 1991 and MN-DNR but no specimens at MIN; known from Ontario, NE, IL
Salix alba L. - white willow
native to Europe; only two specimens (verified by G. Argus, 2010) indicate possible naturalization in Minnesota
Salix amygdaloides Andersson - peachleaf willow
Salix babylonica L.
misapplied; see *Salix* x *sepulcralis*
Salix bebbiana - Bebb willow
Salix candida Flugge ex Willd. - sageleaf willow
Salix cordata Michaux - heartleaf willow
to be expected in eastern counties on sandy soil; known from WI and eastward; considered to be a synonym of *S. eriocephala* in Smith 2008
Salix daphnoides Villars - daphne willow
native to Eurasia; known only from St. Louis Co. (Duluth area; one population collected 1980)
Salix discolor - pussy willow
Salix eriocephala Michx. - diamond willow
Salix eriocephala var. *famelica* (C.R Ball) Dorn - yellow willow
Salix exigua Nuttall
misapplied to eastern US flora see *S. interior*; misapplied name
Salix famelica (C. R. Ball) Argus - yellow willow
fide G. Argus (pers. comm.) populations of this western species are quite distinct from the eastern *S. eriocephala*; in Minnesota, however, there exists a narrow band of hybrids
Salix humilis var. *humilis* - prairie willow
Salix humilis var. *tristis* (Aiton) Griggs - prairie willow
Salix interior Rowlee - sandbar willow

Salix lucida Muhlenberg - **shining willow**
vars. not recognized

Salix lutea Nuttall
misapplied to our flora (G. Argus, pers. comm.) see *S. famelica*; misapplied name

Salix maccalliana Rowlee - **McCalla's willow**

Salix myricoides Muhl. - **bayberry willow**
erroneous report

Salix nigra Marshall - **black willow**
vars. not recognized

Salix pedicellaris Pursh - **bog willow**

Salix pellita (Andersson) Bebb - **satiny willow**
state Special Concern list; threatened

Salix pentandra L. - **laural willow**
native to Europe; fide G. Argus 2010, only carpellate plants are found in the US

Salix petiolaris Smith - **meadow willow**

Salix planifolia Pursh - **diamondleaf willow**

Salix pseudomonticola C.R. Ball - **false mountain willow**
vegetative specimens may be easily confused with *S. famelica*; special concern

Salix purpurea L. - **purpleosier willow**
native to Europe; reported for Minnesota by FNA but no specimens at MIN

Salix pyrifolia Andersson - **balsam willow**

Salix sericea Marshall - **silky willow**
vegetative material may be difficult to distinguish from *S. petiolaris*; known only from Fillmore Co. (collected 1982)

Salix serissima (L.H. Bailey) Fernald - **autumn willow**
potentially invasive (fide G. Argus 2010), unlike most willows, seeds of this species experience winter dormancy and germinate very early in the spring before the seedlings can be outcompeted by others, and Smith 2008 indicates few Minnesota plants are long-lived

Salix x beschellii B. Boivin - **hybrid willow**
(*S. bebbiana* x *discolor*) only one specimen identified as "possibly *S. bebbiana* x *discolor*" suggested by the presence of ferruginous hairs but otherwise characters indicate *S. bebbiana* (fide G. Argus, 2010); Isanti County (collected 1926)

Salix x confinera Wangenb.
(*Salix discolor* x *humilis*)

Salix x fragilis L. - **crack willow**
(*S. alba* x *euxina*) native to Europe; viable seed rarely produced

Salix x rubella Bebb ex C.K. Schneider - **hybrid willow**
(*S. candida* x *eriocephala*) – erroneously reported for Minnesota by USDA-NRCS; currently known only from Newfoundland and NY

Salix x sepulcralis Simonkai

Salix x sepulcralis* var. *chrysocoma (Dode) Meikle - **weeping willow**
(*S. alba* x *babylonica*) native to Europe; previously all weeping willows were called *S. babylonica*, which is not cold hardy; strongly pendulous and golden branches; known only from Anoka Co. (collected 2006)

***Salix x sepulcralis* var. *sepulcralis* - weeping willow**
(*S. alba* x *babylonica*) native to Europe; previously all weeping willows were called *S. babylonica*, which is not cold hardy; possibly var. *sepulcralis* (fide G. Argus 2010) -- known only from Cass Co. (collected 1992)

SALVINIACEAE

Salvinia molesta D.S. Mitch. - **kariba weed**
Federally listed Noxious Weed and on MN-DNR's Prohibited Invasive Species list; common aquarium plant native to tropical Asia and known only from southernmost states; unlikely to be overwintering in Minnesota; reports from Winona Co. unverified

SAPINDACEAE

Acer ginnala Maximowicz - **amur maple**
native to Asia

Acer negundo* var. *interius (Britton) Sargent - **box elder**

***Acer negundo* var. *negundo* - box elder**

Acer negundo* var. *violaceum Booth ex G. Kirchn. - **boxelder**
report unverified

Acer nigrum Michaux f. - **black maple**

Acer pensylvanicum L. - **striped maple**
report unverified

Acer platanoides L. - **Norway maple**
native to Europe; naturalized only in St. Louis Co. (Duluth area; collected 2003)

***Acer rubrum* var. *rubrum* - red maple**
fall leaves deadly poisonous to horses

Acer saccharinum L. - **silver maple**

Acer saccharum* var. *saccharum Marshall - **sugar maple**

Acer spicatum Lamarck - **mountain maple**

Aesculus glabra Willdenow - **Ohio buckeye**
native to eastern US; toxic to pets

Aesculus hippocastanum L.

SARRACENIACEAE

Sarracenia purpurea* subsp. *purpurea - **purple pitcher plant**

Sarracenia purpurea* var. *gibbosa - **purple pitcher plant**
erroneous report

SAXIFRAGACEAE

Chrysosplenium alternifolium L. - **Iowa golden saxifrage**

misapplied to US flora see *C. iowense*; misapplied name

Chrysosplenium americanum Schweinitz ex Hooker - **American golden saxifrage**

Chrysosplenium iowense Rydberg - **Iowa golden saxifrage**

disjunct from Canadian prairies in the Driftless Area of MN and IA; endangered

Heuchera richardsonii R. Brown - **Richardson's alumroot**

Micranthes pensylvanica (L.) Haworth - **eastern swamp saxifrage**

Micranthes virginiana (Michaux) Small - **early saxifrage**

Mitella diphylla L. - **twoleaf miterwort**

Mitella nuda L. - **naked miterwort**

Saxifraga cernua L. - **nodding saxifrage**

disjunct from arctic Canada and Rocky Mountains, known only from Cook Co. (last collected 1980); rarely producing seed, reproduction primarily from bulblets; endangered

Saxifraga paniculata Miller - **white mountain saxifrage**

Lake Superior populations disjunct from eastern Canada; special concern

Sullivantia sullivantii (Torrey & A. Gray) Britton - **Sullivant's coolwort**

northern limit of distribution; threatened

Tiarella cordifolia L. - **heartleaf foamflower**

western limit of distribution; known only from Stearns Co. (one population, collected 1998)

SCHEUCHZERIACEAE

Scheuchzeria palustris L. - **rannoch-rush**

SCROPHULARIACEAE

Scrophularia lanceolata Pursh - **lanceleaf figwort**

Scrophularia marilandica L. - **carpenter's square**

Scrophularia nodosa L. - **woodland figwort**

native to Europe; known only from St. Louis Co. (Duluth area; one population, collected 1997)

Verbascum blattaria L. - **moth mullein**

native to Eurasia; known only from Beltrami Co. (one roadside population, collected 1999)

Verbascum chaixii Villars - **nettle-leaf mullein**

native to Europe; known only from St. Louis Co. (Duluth area; one lawn-weed population, collected 2002)

Verbascum epixanthemum Boissier & Heldreich - **Grecian mullein**

Verbascum nigrum L. - **black mullein**

native to Eurasia; known only from Minneapolis area (one population, collected 1933)

Verbascum phlomoides L. - **orange mullein**

native to Europe; reported for Minnesota by USDA-NRCS but no specimens at MIN

Verbascum thapsus L. - **flannel mullein**

native to Europe; leaves and inflorescence possibly toxic

SELAGINELLACEAE

Selaginella rupestris (L.) Spring - **northern selaginella**

Selaginella selaginoides (L.) Palisot de Beauvois ex Martius & Schrank - **club spikemoss**

known only from Cook Co. (last collected 1998); endangered

SMILACACEAE

Smilax ecirrhata S. Watson - **upright Carrionflower**

Smilax herbacea L. - **smooth Carrionflower**

Smilax illinoensis Mangaly - **Illinois greenbrier**

Smilax lasioneura Hooker - **Blue Ridge Carrionflower**

Smilax pulverulenta Michaux - **downy Carrionflower**

reported for Minnesota (far southeast corner) by FNA but no specimens at MIN (probably based on misidentified specimens)

Smilax rotundifolia L. - **roundleaf greenbrier**

Smilax tamnoides L. - **bristly greenbrier**

SOLANACEAE

Datura innoxia Mill. - **white lady's slipper**

see *D. wrightii*; misapplied name

Datura stramonium L. - **jimsonweed**

native from tropical US to Central America; state DOA noxious (CN) weed, 1st collected 1878, St. Louis Co.; contains hallucinogenic compounds, deadly poisonous to pets and livestock

Datura wrightii Regel - **white lady's slipper**

native from southwestern US to Mexico; occasionally escaping, rarely persisting

Leucophysalis grandiflora (Hooker) Rydberg - **large false groundcherry**

Lycium barbarum L. - **matrimony vine**

native to Eurasia

Nicandra physalodes (L.) Gaertner - **apple of Peru**

native to Peru; known only from Winona Co. (last collected 1901)

Nicotiana rustica L. - **Aztec tobacco**

native to tropical America; reported for Minnesota by USDA-NRCS but no specimens at MIN; annual unlikely to be persisting

Petunia axillaris (Lam.) Britton, Sterns & Poggenb. - **large white petunia**

Petunia x atkinsiana (Sweet) D. Don ex W.H. Baxter - **petunia**

(*Petunia axillaris* x *integrifolia*)

Petunia x axillaris (Lam.) Britton, Sterns & Poggenb.

native to Argentina; reported for Minnesota by USDA-NRCS but no specimens at MIN; doubtfully persisting annual

Physalis alkekengi L. - **strawberry groundcherry**

native to Eurasia; reported for Minnesota by Morley 1969 as a rare garden escape but no specimens at MIN

Physalis grisea (Waterf.) M. Martínez - **strawberry-tomato**

Physalis heterophylla var. *heterophylla* - clammy groundcherry

Physalis hispida (Waterf.) Cronquist - prairie groundcherry

report unverified

Physalis longifolia var. *longifolia* - longleaf groundcherry

reported for Minnesota by Gleason & Cronquist 1991 but no specimens at MIN; known throughout the US and eastern Canada

Physalis longifolia var. *subglabrata* (Mack. & Bush) Cronquist - longleaf groundcherry

report unverified

Physalis philadelphica var. *immaculata* Waterf. - **Mexican groundcherry**

report unverified

Physalis pubescens var. *integrifolia* (Dunal) Waterf. - **husk tomato**

Physalis virginiana var. *virginiana* (Dunal) Waterf. - **Virginia groundcherry**

Solanum americanum Mill. - **West Indian black nightshade**

misapplied; see *S. ptycanthemum*

Solanum carolinense var. *carolinense* - **Carolina horseradish**

Solanum dulcamara var. *dulcamara* - climbing nightshade

Solanum nigrum L. - **black nightshade**

see *Solanum ptycanthemum*; uncertain status

Solanum physalifolium Rusby - **hoe nightshade**

Solanum ptycanthemum Dunal - **West Indian black nightshade**

Solanum rostratum Dunal - **buffalobur nightshade**

native to central US; state DOA noxious (CN) weed, 1st collected 1890, Hennepin Co.

Solanum triflorum Nuttall - **cutleaf nightshade**

native to western US

Solanum tuberosum L. - **Irish potato**

native to Andes Mountains of South America; occasionally escaping from cultivation, rarely persisting more than 1 or 2 years

SPARGANIACEAE

Sparganium americanum Nuttall - **American bur-reed**

Sparganium androcladum (Engelmann) Morong - **branched bur-reed**

Sparganium angustifolium Michaux - **narrowleaf bur-reed**

Sparganium emersum Rehmann - **European bur-reed**

Sparganium erectum L. - **broadfruit bur-reed**

misapplied to North American flora, see *S. eurycarpum*

Sparganium eurycarpum Engelmann - **broadfruit bur-reed**

Sparganium fluctuans (Engelmann ex Morong) B.L. Robinson - **floating bur-reed**

Sparganium glomeratum (Beurling ex Laestadius) L.M. Newman - **clustered bur-reed**

Sparganium natans L. - **small bur-reed**

STAPHYLEACEAE

Staphylea trifolia L. - **American bladdernut**

TAMARICACEAE

Tamarix gallica L. - **French tamarisk**

Tamarix ramosissima Ledeb. - **saltcedar**

TAXACEAE

Taxus canadensis Marshall - **Canada yew**

deadly poisonous to humans, pets, and livestock

THELYPTERIDACEAE

Phegopteris connectilis (Michaux) Watt - **long beech fern**

Phegopteris hexagonoptera (Michaux) Fe - **broad beech fern**

northwest limit of distribution; endangered

Thelypteris palustris var. *pubescens* (Lawson) Fernald - **eastern marsh fern**

THYMELAEACEAE

Daphne mezereum L. - **paradise plant**

native to Eurasia; known to be invasive in the eastern US and Canada and poisonous to humans; known only from Lake Co. (only known specimen at DUL) though scheduled to be extirpated by state Dept. of Agriculture

Dirca palustris L. - **eastern leatherwood**

mildly toxic to mammals and may cause dermatitis

TOFIELDIACEAE

Tofieldia pusilla (Michaux) Persoon - **scotch false asphodel**

disjunct from arctic regions and Canadian Rocky Mountains; endangered

Triantha glutinosa (Michaux) Baker - **sticky tofieldia**

TYPHACEAE

Typha angustifolia L. - **narrowleaf cattail**

native to eastern seaboard states and provinces

Typha latifolia L. - **broadleaf cattail**

Typha x glauca Godron - **hybrid cattail**

(*T. angustifolia* x *latifolia*) although reported from only 20 of the 87 counties, fide Lee Frelich, UM research associate, forest ecology (pers. comm.), this hybrid is much

more widespread, often out-competing both parents

ULMACEAE

Ulmus americana K. - American elm

Ulmus glabra Huds. - Wych elm

native to Eurasia; erroneously reported for Minnesota by USDA-NRCS

Ulmus pumila - Siberian elm

native to eastern Asia

Ulmus rubra Muhlenberg - slippery elm

Ulmus thomasii Sargent - rock elm

URTICACEAE

Boehmeria cylindrica (L.) Swartz - smallspike false nettle

Laportea canadensis (L.) Weddell - Canadian woodnettle

Parietaria pensylvanica Muhlenberg ex Willdenow - Pennsylvania pellitory

Pilea fontana (Lunell) Rydberg - lesser clearweed

Pilea pumila (L.) A. Gray - Canadian clearweed

Urtica dioica subsp. *dioica* - stinging nettle

misapplied; see *Urtica dioica* subsp. *gracilis*

Urtica dioica subsp. *gracilis* (Aiton) Selander - stinging nettle

can cause skin irritation

VALERIANACEAE

Valeriana edulis var. *ciliata* (Torrey & A. Gray) Cronquist - tobacco root

northwest limit of distribution; threatened

Valeriana officinalis L. - garden valerian

native to Eurasia

VERBENACEAE

Glandularia canadensis (L.) Small - rose mock vervain

Phyla lanceolata (Michaux) Greene - lanceleaf fogfruit

Verbena bracteata Lagasca y Segura & Rodr - bigbract verbena

Verbena hastata var. *hastata* - swamp verbena

Verbena hastata var. *scabra* Moldenke - swamp verbena

Verbena officinalis var. *officinalis*

native to Europe; known only from St. Louis Co. (one garden plot population, collected 2005)

Verbena simplex Lehmann - narrowleaf vervain

Verbena stricta Ventenat - hoary vervain

Verbena urticifolia var. *leiocarpa* L.M. Perry & Fernald - white vervain

Verbena urticifolia var. *urticifolia* - white vervain

Verbena x deamii Moldenke - vervain

(*V. bracteata* x *stricta*) known only from New Ulm area (historical collection from 1891); rare historic record

Verbena x perriana Moldenke - vervain

(*V. bracteata* x *urticifolia*) known only from Minneapolis area (historical collections to 1877); rare historic record

Verbena x rydbergii Moldenke - vervain

(*V. hastata* x *stricta*) reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, WI, IA, NE

VIOLACEAE

Hybanthus concolor Sprengel - eastern greenviolet

range extension from IA and WI; known only from Winona Co. (one population, collected 1999); endangered

Viola adunca var. *adunca* - hookedspur violet

Viola affinis LeConte - sand violet

Viola arvensis Murray - European field pansy

native to Europe; known only from Cass Co. (last collected 1939), doubtfully persisting

Viola bicolor Pursh - field pansy

Viola blanda Willdenow - sweet white violet

Viola canadensis var. *canadensis* - creepingroot violet

Viola canadensis var. *rugulosa* (Greene) C.L. Hitchcock - Canadian white violet

Viola cucullata Aiton - marsh blue violet

Viola labradorica Schrank - alpine violet

Viola lanceolata L. - bog white violet

Viola macloskeyi subsp. *pallens* (Banks ex Ging.) M.S. Baker - small white violet

Viola missouriensis Greene - Missouri violet

Viola nephrophylla Greene - northern bog violet

Viola novae-angliae House - New England blue violet

Viola nuttallii Pursh - Nuttall's violet

eastern limit of distribution; threatened

Viola palustris L. - marsh violet

Viola pedata var. *pedata* L. - birdfoot violet

Viola pedatifida G. Don - prairie violet

Viola primulifolia var. *primulifolia* L. - primrose-leaved violet

Viola pubescens var. *pubescens* - downy yellow violet

Viola pubescens var. *scabriuscula* Torrey & A. Gray - downy yellow violet

Viola renifolia A. Gray - white violet

Viola sagittata var. *ovata* - arrowleaf violet

Viola sagittata var. *sagittata* - arrowleaf violet

Viola selkirkii Pursh ex Goldie - Selkirk's violet

Viola sororia Willdenow - common blue violet

Viola tricolor L. - johnny jumpup

introduced from Europe

Viola x bernardii (Greene) Greene - Bernard violet

(*V. pedatifida* x *sororia*) reported for Minnesota by USDA-NRCS but no specimens at MIN

Viola x bisselli House - hybrid violet

(*V. cucullata* x *sororia*) known only from St. Louis Co.

Viola x conjugens Greene - hybrid violet

(*V. sagittata* x *sororia*)

Viola x mistura House - hybrid violet

(*V. palmata* x *sagittata*)

Viola x napae House

Viola x populifolia Greene - Peck's violet

(*V. sororia* x *triloba*) reported for Minnesota by USDA-NRCS but no specimens at MIN

Viola x sublanceolata House - lanceleaf violet

(*Viola lanceolata* x *macloskeyi*); reported for Minnesota by USDA-NRCS but no specimens at MIN; known from Ontario, IA; report unverified

VISCACEAE

Arceuthobium pusillum Peck - eastern dwarf mistletoe

VITACEAE

Parthenocissus quinquefolia (L.) Planchon - Virginia creeper

leaves and fruit toxic

Parthenocissus tricuspidata (Siebold & Zuccarini) Planchon - boston ivy

native to China and Japan; known only from Pipestone Co. (one population, collected 1961)

Parthenocissus vitacea (Knerr) Hitchcock - woodbine

Vitis aestivalis var. *argentifolia* (Munson) Fernald - summer grape

see *Vitis aestivalis* var. *bicolor*; threatened

Vitis aestivalis var. *bicolor* Michx. - summer grape

Vitis riparia Michaux - frost grape

XYRIDACEAE

Xyris montana Ries - northern yelloweyed grass

Xyris torta Smith - slender yelloweyed grass

northwest limit of distribution; endangered

ZYGOPHYLLACEAE

Tribulus terrestris L. - puncturevine

introduced from Mediterranean region