Endangered Karst Invertebrate Taxonomy of Central Texas

U.S. Fish and Wildlife Service Austin Ecological Services Field Office 10711 Burnet Rd. Suite #200 Austin, TX 78758

> Original date: July 28, 2011 Revised on: April 4, 2019

TABLE OF CONTENTS

1.0	INTRODUCTION	1
2.0	ENDANGERED KARST INVERTEBRATE TAXONOMY	1
2.1	Batrisodes texanus (Coffin Cave mold beetle)	2
2.2	Batrisodes venyivi (Helotes mold beetle)	3
2.3	Cicurina baronia (Robber Baron Cave meshweaver)	4
2.4	Cicurina madla (Madla Cave meshweaver)	5
2.5	Cicurina venii (Braken Bat Cave meshweaver)	6
2.6	Cicurina vespera (Government Canyon Bat Cave meshweaver)	7
2.7	Neoleptoneta microps (Government Canyon Bat Cave spider)	8
2.8	Neoleptoneta myopica (Tooth Cave spider)	9
2.9	Rhadine exilis (no common name)	10
2.10	0 Rhadine infernalis (no common name)	11
2.1	1 Rhadine persephone (Tooth Cave ground beetle)	12
	2 Tartarocreagris texana (Tooth Cave pseudoscorpion)	
2.13	3 Texamaurops reddelli (Kretschmarr Cave mold beetle)	14
2.14	4 Texella cokendolpheri (Cokendolpher cave harvestman)	15
2.15	5 Texella reddelli (Bee Creek Cave Harvestman)	16
2.16	6 Texella reyesi (Bone Cave Harvestman)	17
3.0	KARST INVERTEBRATE COLLECTION AND IDENTIFICATION	18
4.0	LITERATURE CITED	20

LIST OF FIGURES

Table 1. Taxonomists	19
Figure 10. Texella reyesi	17
Figure 9. Texella tuberculata showing general Texella morphology	15
Figure 8. Tartarocreagris attenuata	13
Figure 7. Rhadine persephone	12
Figure 6. Rhadine infernalis	
Figure 5. Rhadine exilis (on right)	10
Figure 4. Neoleptoneta myopica	
Figure 3. Cicurina madla from a cave in Government Canyon State Natural Area	5
Figure 2. Cicurina baronia from Robber Baron Cave	4
Figure 1. Batrisodes texanus	2

1.0 INTRODUCTION

The purpose of this document is to provide up-to date taxonomic information for the sixteen endangered karst invertebrates that occupy caves and karst areas in Bexar, Travis, and Williamson Counties, Texas. It covers six insects (three ground beetles and three mold beetles) and ten arachnids (three harvestmen, six spiders, and one pseudoscorpion). Methods to collect specimens for taxonomic identification are also provided as well as a list of taxonomists that are qualified to provide identifications. This document will be updated periodically as we receive more information.

2.0 ENDANGERED KARST INVERTEBRATE TAXONOMY

Taxonomic verification of these species is usually not possible in the field and typically requires examination of adult specimens under a microscope, dissection of the genitalia by a taxonomic expert, and/or for the *Cicurina* species, DNA identification. The following are the species covered in this document:

Insects:

- Rhadine exilis, R. infernalis, and R. persephone (small, essentially eyeless ground beetles)
- Batrisodes venyivi, B. texanus, and Texamaurops reddelli (small, eyeless mold beetles)

Arachnids:

- *Texella cokendolpheri. T. reddelli*, and *T. reyesi* (small, eyeless harvestmen (daddylonglegs))
- Neoleptoneta microps, N. myopica, Cicurina baronia, C. madla, C. venii, and C. vespera (small, eyeless, or essentially eyeless spiders)
- Tartarocreagris texana (small, eyeless pseudoscorpion)

2.1 Batrisodes texanus (Coffin Cave mold beetle)

Scientific Name: Batrisodes (Excavodes) texanus

Common Name: Coffin Cave mold beetle

<u>Description</u>: This species was listed as the Kretschmarr Cave mold beetle (*Texamaurops reddelli*) (Barr 1974a) then later split into two species: the Kretschmarr Cave mold beetle and Coffin Cave mold beetle (*Batrisodes texamus*) (Chandler 1992). Chandler and Reddell (2001) further split *B. texamus* into *B. texamus* and *B. cryptotexanus*. Chandler et al. (2009) again stated that these are two distinct species described by Chandler (1992). The Service has not yet officially recognized this taxonomic revision and considers everything identified as either *B. texanus* or *B. cryptotexanus* to be in the entity listed as endangered.

<u>Type Specimen</u>: The male holotype was collected from Inner Space Cavern, Williamson County, Texas, on May 23, 1965 by William Russell. The holotype is curated at the Field Museum of Natural History (Chandler 1992).

<u>Selected Characteristics</u>: Length 0.10-0.11 inches (in) (2.60-2.88 millimeter (mm)). This is a tiny, reddish-brown beetle that superficially resembles an ant (Figure 1).

<u>Taxonomic Classification</u>: Class Insecta (insects), Order Coleoptera (beetles), Suborder Polyphaga, Family Pselaphidae (mold beetles), Tribe Batrisini, Genus *Batrisodes*, Subgenus *Excavodes*.

<u>Distribution</u>: Williamson County

Figure 1. Batrisodes texanus



Photo by Dr. Kemble White

2.2 Batrisodes venyivi (Helotes mold beetle)

Scientific Name: Batrisodes (Excavodes) venyivi

Common Name: Helotes mold beetle

<u>Description</u>: This species was described by Chandler (1992).

<u>Type Specimen</u>: The male holotype was collected from Helotes Hilltop Cave, Bexar County, Texas, on September 29, 1984 by J. Ivy and G. Veni (Chandler 1992).

<u>Selected Characteristics</u>: Length 0.08 in (2.24 mm) (Chandler 1992). This is a tiny, reddish- brown beetle that superficially resembles an ant. A congener (member of the same genus) is shown to illustrate the general morphology of the genus (Figure 1).

<u>Taxonomic Classification</u>: Class Insecta (insects), Order Coleoptera (beetles), Suborder Polyphaga, Family Pselaphidae (mold beetles), Tribe Batrisini, Genus *Batrisodes*, Subgenus *Excavodes*.

2.3 Cicurina baronia (Robber Baron Cave meshweaver)

Scientific Name: Cicurina (Cicurella) baronia

Common Name: This species has been referred to by two common names, the Robber Baron Cave spider (Service 2000) and the Robber Baron Cave meshweaver (Breene et al. 2003). The latter name is now accepted as the official common name (Breene et al. 2003).

<u>Description</u>: The species was described by Gertsch (1992).


<u>Type Specimen</u>: The female holotype was collected by R. Bartholomew from Robber BaronCave, Bexar County, Texas, in April 1969 (Gertsch 1992).

<u>Selected Characteristics</u>: This species is eyeless and has reduced pigment (Figure 2). Additional descriptive information on this species is found in Cokendolpher (2004), Reddell and Cokendolpher (2004), Paquin and Hedin (2004), and Paquin and Dupérré (2009).

<u>Taxonomic Classification</u>: Class Arachnida (arachnids), Order Araneae (spiders), Family Dictynidae, Genus *Cicurina*, Subgenus *Cicurella*.

Distribution: Bexar County

Figure 2. Cicurina baronia from Robber Baron Cave



2.4 Cicurina madla (Madla Cave meshweaver)

Scientific Name: Cicurina (Cicurella) madla

Common Name: This species has been referred to by two common names, Madla's Cave Spider (Service 2000) and Madla Cave meshweaver (Breene et al. 2003). The latter name is now accepted as the official common name (Breene et al. 2003).

<u>Description</u>: The species was described by Gertsch (1992).

<u>Type Specimen</u>: The female holotype was collected by D. McKenzie and J. Reddell in Madla's Cave, 3.1 miles (mi) (5 kilometers (km)) north of Helotes, Bexar County, Texas, on October 4, 1963 (Gertsch 1992).

<u>Selected Characteristics</u>: This species is eyeless and has reduced pigment (Figure 3). Additional descriptive information on this species is found in Cokendolpher (2004), Reddell and Cokendolpher (2004), Paquin and Hedin (2004), and Paquin and Dupérré (2009).

<u>Taxonomic Classification</u>: Class Arachnida (arachnids), Order Araneae (spiders), Family Dictynidae, Genus *Cicurina*, Subgenus *Cicurella*.

Distribution: Bexar County

Figure 3. Cicurina madla from a cave in Government Canyon State Natural Area





2.5 Cicurina venii (Braken Bat Cave meshweaver)

Scientific Name: Cicurina (Cicurella) venii

Common Name: The Service (2000) listed no common name for this species. The Committee on Common Names of Arachnids (Breene et al. 2003) listed the official common name of this species as the Braken Bat Cave meshweaver, which is now accepted as the official common name (Breene et al. 2003).

<u>Description</u>: The species was described by Gertsch (1992).

<u>Type Specimen</u>: The female holotype was collected in Braken Bat Cave on November 22, 1980, by G. Veni (Gertsch 1992, Paquin and Dupérré 2009).

<u>Selected Characteristics</u>: This species is eyeless and has reduced pigment. A congener is shown to illustrate the general morphology of the genus (Figure 3). Additional descriptive information on this species is found in Cokendolpher (2004), Reddell and Cokendolpher (2004), Paquin and Hedin (2004), and Paquin and Dupérré (2009).

<u>Taxonomic Classification</u>: Class Arachnida (arachnids), Order Araneae (spiders), Family Dictynidae, Genus *Cicurina*, Subgenus *Cicurella*.

2.6 Cicurina vespera (Government Canyon Bat Cave meshweaver)

Scientific Name: Cicurina (Cicurella) vespera (Gertsch).

Common Name: This species has been referred to by two common names, the vesper cave spider (Service 2000) and the Government Canyon Bat Cave meshweaver (Breene et al. 2003). The latter name has been accepted as the official common name (Breene et al. 2003).

<u>Description</u>: The species was described by Gertsch (1992).

<u>Type Specimen</u>: The female holotype was collected from Government Canyon Bat Cave on August 11, 1965, by J. Fish and J. Reddell (Gertsch 1992).

<u>Selected Characteristics</u>: This species is eyeless and has reduced pigment. A congener is shown to illustrate the general morphology of the genus (Figure 3). Additional descriptive information on this species is found in Cokendolpher (2004), Reddell and Cokendolpher (2004), Paquin and Hedin (2004), and Paquin and Dupérré (2009).

<u>Taxonomic Classification</u>: Class Arachnida (arachnids), Order Araneae (spiders), Family Dictynidae, Genus *Cicurina*, Subgenus *Cicurella*. A possible synonymy between *C. vespera* and *C. madla* was suggested by the molecular analysis of Paquin and Hedin (2004); however, no formal synonymy was set forth in their work.

2.7 Neoleptoneta microps (Government Canyon Bat Cave spider)

Scientific Name: Neoleptoneta microps

Common Name: This species has been referred to by two common names, the Government Canyon cave spider (Service 2000) and the Government Canyon Bat Cave spider (Breene et al. 2003). The latter name is now accepted as the official common name (Breene et al. 2003).

<u>Description</u>: *Neoleptoneta microps* was first collected in 1965 and described by Gertsch (1974) as *Leptoneta microps*. The species was reassigned to *Neoleptoneta* following Brignoli (1977) and Platnick (1986). A review of the taxonomic history of nearctic leptonetids is available in Ubick et al. (2005). Ledford et al. (2011), reassigned the species to *Tayshaneta microps*. The Service has not officially recognized this taxonomic revision.

<u>Type Specimen</u>: The female holotype was collected from Government Canyon Bat Cave, Bexar County, Texas on August 11, 1965, by J. Fish and J. Reddell (Gertsch 1974).

<u>Selected Characteristics</u>: This is a small, yellowish, essentially eyeless troglobite. A congener is shown to illustrate the general morphology of the genus (Gertsch 1974) (Figure 4).

<u>Taxonomic Classification</u>: Class Arachnida (arachnids), Order Araneae (spiders), Infraorder Araneomorphae (true spiders), Family Leptonetidae.

2.8 Neoleptoneta myopica (Tooth Cave spider)

Scientific Name: Neoleptoneta myopica

<u>Common Name</u>: Tooth Cave spider

<u>Description</u>: Neoleptoneta myopica was described by Gertsch (1974) as Leptoneta myopica. The species was reassigned to Neoleptoneta following Brignoli (1977) and Platnick (1986). A review of the taxonomic history of nearctic leptonids is available in Ubick et al. (2005). Ledford et al. 2011, reassigned the species to Tayshaneta myopica. The Service has not officially recognized this taxonomic revision.

<u>Type Specimen</u>: The female holotype was collected from Tooth Cave, Travis County, Texas, on March 30, 1965, by J. Reddell (Gertsch 1974). The holotype is curated at the American Museum of Natural History (Ledford 2011).

<u>Selected Characteristics</u>: Body length is 0.06 in (1.6 mm). This is a small, yellowish/whitish, essentially eyeless troglobite. A congener is shown to illustrate the general morphology of the genus (Gertsch 1974) (Figure 4).

<u>Taxonomic Classification</u>: Class Arachnida (arachnids), Order Araneae (spiders), Infraorder Araneomorphae (true spiders), Family Leptonetidae.

<u>Distribution</u>: Travis and Williamson Counties

Figure 4. Neoleptoneta myopica



2.9 Rhadine exilis (no common name)

Scientific Name: Rhadine exilis

Common Name: This species has no common name.


<u>Description</u>: This species was originally described as *Agonum exile* by Barr and Lawrence (1960). Later this species was referred to as *Rhadine exilis* by Reddell (1966) in a checklist. Barr (1974b) formally reassigned the species to the genus *Rhadine*.

<u>Type Specimen</u>: The male holotype (the only specimen collected) was collected from Marnock Cave, 0.99 mi (1.6 km) north of Helotes, Bexar County, Texas, on July 2, 1959 by J. F. Lawrence and F. Moore (Barr 1974b).

<u>Selected Characteristics</u>: Mean length is 0.29 in (7.4 mm). Body is extremely slender (Figure 5).

<u>Taxonomic Classification</u>: Class Insecta (insects), Order Coleoptera (beetles), Suborder Adephaga, Family Carabidae (ground beetles), Tribe Agonini (agonines).

Distribution: Bexar County





2.10 Rhadine infernalis (no common name)

Scientific Name: Rhadine infernalis

<u>Common Name</u>: This species has no common name.

<u>Description</u>: This species was originally described as *Agonum infernale* by Barr

and

Lawrence (1960). Barr (1974b) reassigned the species to the genus *Rhadine*.

<u>Type Specimen</u>: The male holotype was collected from Madla's Cave, 3.1 mi (5 km) north of Helotes, Bexar County, Texas, on July 6-7, 1959, by J. F. Lawrence and J. R. Reid (Barr 1974b).

<u>Selected Characteristics</u>: Mean length is 0.28 in (7.2 mm). Body is robust (Figure 6).

<u>Intraspecific Variation</u>: Taxonomists have delineated three subspecies (*R. infernalis ewersi*, *R. infernalis infernalis*, and *R. infernalis* ssp.). Two have been formally described (Barr 1960). In a more recent report, the third subspecies was characterized as valid, but was not formally described (Reddell 1998).

<u>Taxonomic Classification</u>: Class Insecta (insects), Order Coleoptera (beetles), Suborder Adephaga, Family Carabidae (ground beetles), Tribe Agonini (agonines).

Distribution: Bexar County

Figure 6. Rhadine infernalis



2.11 Rhadine persephone (Tooth Cave ground beetle)

Scientific Name: Rhadine persephone

<u>Common Name</u>: Tooth Cave ground beetle

<u>Description</u>: This species was originally described by Barr (1974b).

<u>Type Specimen</u>: The male holotype was collected from Tooth Cave, Travis County, Texas, on May 16, 1965 by R. W. Mitchell, T.C. Barr, and W.M. Andrews (Barr 1974b). The holotype is curated at the Museum of Natural History (Barr 1974b).

<u>Selected Characteristics</u>: Body length is 0.28-0.34 in (7.2-8.7 mm). Body is robust (Barr 1974b) (Figure 7).

<u>Taxonomic Classification</u>: Class Insecta (insects), Order Coleoptera (beetles), Suborder Adephaga, Family Caribidae (ground beetles), Tribe Agonini (agonines)

Distribution: Travis and Williamson Counties



Figure 7. Rhadine persephone

2.12 Tartarocreagris texana (Tooth Cave pseudoscorpion)

Scientific Name: Tartarocreagris texana

Common Name: Tooth Cave pseudoscorpion

<u>Description</u>: This species was originally described as *Microcreagris texana* by Muchmore (1969). The genus *Microcreagris* was reassigned to *Tartarocreagris* following Muchmore (1992).

<u>Type Specimen</u>: The female holotype was collected from Tooth Cave, Travis County, Texas, on May 16, 1965, by James Reddell. The holotype is curated at the American Museum of Natural History (Muchmore 1992).

<u>Selected Characteristics</u>: Body 0.16 in (4.1 mm). This is an eyeless pseudoscorpion. The upper portion of the body (carapace, chelicerae and palps) are golden brown, body and legs are light tan (Muchmore 1969). A congener is shown to illustrate the general morphology of the genus (Figure 8).

<u>Taxonomic Classification</u>: Class Arachnida (arachnids), Order Pseudoscorpiones, family Neobisiidae.

Distribution: Travis County






Photo by Mark Sanders

2.13 Texamaurops reddelli (Kretschmarr Cave mold beetle)

Scientific Name: Texamaurops reddelli

Common Name: Kretschmarr Cave mold beetle

<u>Description</u>: This species was originally described by Barr and Steeves (1963)

<u>Type Specimen</u>: The female holotype was collected from Kretschmarr Cave, Travis County, Texas on March 2, 1963, by James Reddell and Davis MacKenzie (Barr and Steeves 1963). The holotype is curated at the Field Museum of Natural History Chicago.

<u>Selected Characteristics</u>: Body length is 0.10-0.15 in (2.72-3.96 mm). A small reddish-brown beetle with attenuated (long) legs (Barr and Steeves 1963).

<u>Taxonomic Classification</u>: Class Insecta (insects), Order Coleoptera (beetles), Suborder Polyphaga, Family Psephidae (mold beetles), Tribe Batrisini.

Distribution: Travis County

2.14 Texella cokendolpheri (Cokendolpher cave harvestman)

Scientific Name: Texella cokendolpheri

<u>Common Name</u>: This species has been referred to by two common names, the Robber Baron Cave harvestman (Service 2000) and the Cokendolpher cave harvestman (Breene et al. 2003). The latter name is now accepted as the official common name (Breene et al. 2003).

<u>Description</u>: This species was described by Ubick and Briggs (1992).

<u>Type Specimen</u>: The male holotype was collected from Robber Baron Cave, Bexar County, Texas, on April 3, 1982, by A. Grubbs (Ubick and Briggs 1992).

<u>Selected Characteristics</u>: Pale orange. A congener is shown to illustrate the general morphology of the genus (Figure 9).

<u>Taxonomic Classification</u>: Class Arachnida (arachnids), Order Opiliones (opilionids, or harvestmen), Suborder Laniatores, Family Phalangodidae.

Distribution: Bexar County

Figure 9. Texella tuberculata showing general Texella morphology





2.15 Texella reddelli (Bee Creek Cave Harvestman)

Scientific Name: Texella reddelli

Common Name: Bee Creek cave harvestman (Goodnight and Goodnight 1967)

<u>Description</u>: This species was described by Goodnight and Goodnight (1967).

<u>Type Specimen</u>: The male holotype was collected from Bee Creek Cave (also referred to as Pine Creek Cave), Travis County, Texas, on October 2, 1963, by James Reddell and David MacKenzie (Goodnight and Goodnight 1967).

<u>Selected Characteristics</u>: Pale orange. Body length 0.07-0.08 in (1.90-2.18 mm) (Ubick and Briggs 1992). A congener is shown to illustrate the general morphology of the genus (Figure 10 and 11).

<u>Taxonomic Classification</u>: Class Arachnida (arachnids), Order Opiliones (opilionids, or harvestmen), Suborder Laniatores, Family Phalangodidae.

Distribution: Travis County

2.16 Texella reyesi (Bone Cave Harvestman)

Scientific Name: Texella reyesi

Common Name: Bone Cave harvestman (Ubick and Briggs 1992)

<u>Description:</u> This species was described by Ubick and Briggs (1992).

<u>Type Specimen:</u> The male holotype was collected from Bone Cave, Travis County, Texas, on June 4, 1989, by William Elliott, James Reddell, and Marcelino Reyes. The holotype is curated at the California Academy of Science (Ubick and Briggs 1992).

<u>Selected Characteristics:</u> Pale orange. Body length 0.05-0.10 in (1.41-2.67 mm) (Ubick and Briggs 1992) (Figure 10).

<u>Taxonomic Classification</u>: Class Arachnida (arachnids), Order Opiliones (opilionids, or harvestmen), Suborder Laniatores, Family Phalangodidae.

Distribution: Travis and Williamson Counties



Figure 10. Texella reyesi

Photo by Mark Sanders

3.0 KARST INVERTEBRATE COLLECTION AND IDENTIFICATION

Endangered karst invertebrates may not be possible to distinguish in the field from closely related species. Therefore, specimens should be collected from locations not known to contain endangered karst invertebrates by an individual that has a valid section 10(a)(1)(A) permit (or Enhancement of Survival permit) from the Service or be accompanied by someone that has such a permit. These collections should be identified as specifically as possible and sent to an appropriate university or museum in the list in Table 1 or to a taxonomist and collection that is approved in writing by the Austin Ecological Services Office (Recovery and Candidate Conservation branch) for taxonomic determination and curation. Please call the Austin Ecological Services Office (512-490-0057 ext. 231) for guidance if you have any questions as to which taxonomist a specimen should be sent to for identification and curation. Documentation from taxonomists regarding specimen identification (including accession or collection numbers) should be provided in all 10(a)(1)(A) permitees' annual reports.

Guidelines to follow while collecting are found in the USFWS Section 10(a)(1)(A) Permit Requirements for Conducting Presence/Absence Surveys for Endangered Karst Invertebrates. The most recent version can be found at:

 $https://www.fws.gov/southwest/es/AustinTexas/ESA_Sci_permits.html \#Presence_absence_karst.$

Table 1. Taxonomists

Species	Taxonomists	Contact Information
Texella species	Darrel Ubick	Department of Entomology California Academy of Science 55 Music Concourse Drive Golden Gate park San Francisco, CA 94128 (415) 379-5308 dubick@calacademy.org
Cicurina species	Dr. Marshal Hedin	Department of Biology Life Sciences North, Room 204E San Diego State University 5500 Campanile Blvd. San Diego, CA 92182-4614 mhedin@mail.sdsu.edu
Rhadine species	James Reddell	Texas Memorial Museum 3001 Lake Austin Blvd., Suite 1.314 University of Texas at Austin Austin, TX 78703 (512) 366-2038 jreddell.caves@mail.utexas.edu
Tartarocreagris species	Charles Stephen	Auburn University Department of Biological Sciences 3331 Funchess Hall Auburn, AL 36849 (334) 734-9021 czs0021@auburn.edu
Batrisodes and Texamaurops species	Dr. Don Chandler	University of New Hampshire Department of Zoology Durham, NC 03824 (603) 862-1735
Neoleptoneta or Texella species	Dr. Joel Ledford	University of California-Davis Department of Plant Biology Sciences Lab Building Davis, CA 95616 (530) 400-4187

4.0 LITERATURE CITED

- Barr, T.C., Jr. 1960. The Cavernicolus beetle of the subgenus *Rhadine*, genus *Agonum* (Coleoptera: Carabidae). American Midland Naturalist 64(1): 45-65.
- Barr, T.C., Jr. 1974a. The eyeless beetles of the genus Arianops Brendel (Coleoptera, Pselaphidae), Bulletin of the American Museum of Natural History 154: 1-52.
- Barr, T.C., Jr. 1974b. Revision of *Rhadine* LeConte (Coleoptera, Carabidae) I. The *subterranea* group. American Museum Novitates 2539: 30.
- Barr, T.C., Jr. and J.F. Lawrence. 1960. New cavernicolous species of *Agonum (Rhadine)* from Texas (Coleoptera: Carabidae). The Wasmann Journal of Biology 18: 137-145.
- Barr, T.D. and H.R. Steeves. 1963. Texamaurops a new genus of pselaphids from caves in central Texas. (Coleptera, Pselaphidae). Colepterist's Bulletin 17: 117-120.
- Breene, R.G., D.A. Dean, G.B. Edwards, B. Hebert, H.W. Levi, G. Manning, K. McWest, and L. Sorkin. 2003. Common names of Arachnids 2003. 5th edition. The American Arachnological Society Committee on Common Names of Arachnids. American Tarantula Society.
- Brignoli, P.M. 1977. Spiders from Mexico, III. A new leptonetid from Oaxaca (Araneae, Leptonetidae). Accademia Nazionale dei Lincei 171 (3): 213-218.
- Chandler, D.S. 1992. The Pselaphidae (Coleoptera) of Texas caves. Texas Memorial Museum Speleological Monographs 3: 241-254.
- Chandler, D.S. and J.R. Reddell. 2001. A review of the ant-like litter beetles found in Texas caves (Coleoptera: Staphylinidae: Pselaphinae), pp. 115-128. Texas Memorial Museum, Speleological Monographs 5: 115-128.
- Chandler, D.S., J.R. Reddell, and P. Paquin. 2009. New cave Pselaphinae and records from Texas with a discussion of the relationships and distributions of the Texas troglobitic Pselaphinae (Coleptera: Staphlinidae: Pselaphinae). Texas Memorial Museum, Speleological Monograph 7: 125-140.
- Cokendolpher, J.C. 2004. Cicurina spiders from caves in Bexar County, Texas (Araneae: Dictynidae). Texas Memorial Museum, Speleological Monographs 6: 13-58.
- Gertsch, W.J. 1974. The spider family Leptonetidae in North America. The Journal of Arachnology 1: 145-203.

- Gertsch, W.J. 1992. Distribution patterns and speciation in North American cave spiders with a list of the troglobites and revision of the Cicurinas of the subgenus *Cicurella*. Texas Memorial Museum Speleological Monographs 3: 75-122.
- Goodnight, C.J. and M.L. Goodnight. 1967. Opilionida from Texas caves (Opiliones, Phalangodidae). American Museum Novitates, No. 2301, 8 pp.
- Ledford, J. 2001. Studies on the cave-spider family Leptonetidae in North America. PhD dissertation. University of California-Berkely. Spring 2011. 254 pp.
- Ledford, J., P. Paquin, J. Cokendolpher, J. Campbell, and C. Griswold. 2011. Systematics of the spider genus Neoleptoneta Brignoli, 1972 (Araneae: Leptonetidae) with a discussion of the morphology and relationships for North American Leptonetidae. Invertebrate Systematics 25: 334-388.
- Muchmore, W.B. 1969. New species and records of cavernicolous pseudoscorpions of the genus Microcreagris (Arachnida, Chelonethida, Neobisiidae, Ideobisiinae). American Museum Novitates, No. 2932, 21, pp.
- Muchmore, W.B. 1992. Cavernicolous pseudoscorpions from Texas and New Mexico (Arachnida, Pseudoscorpionida). Texas Memorial Museum Speleological Monographs 3: 127-154.
- Paquin, P. and M. Hedin. 2004. Genetic and morphological analysis of species limits in *Cicurina* spiders (Araneae, Dictynidae) from southern Travis and northern Hays counties (TX), with emphasis on *Cicurina cueva* Gertsch and relatives. Special report for the Department of Interior United States Fish & Wildlife Service Contract No. 201814G959. Revised version 10 May 2005.
- Paquin, P. and N. Dupérré. 2009. A first step towards the revision of *Cicurina*: redescription of type specimens of 60 troglobitic species of the subgenus *Cicurella* (Araneae: Dictynidae), a first visual assessment of their distribution. Zootaxa 2002: 1-67.
- Platnick, N.I. 1986. On the tibial and patellar glands, relationships, and American genera of the spider family Leptonetidae (Arachnida, Araneae). American Museum Novitates 2855: 1-16. Reddell, J.R. 1966. A checklist of the cave fauna of Texas. II. Insecta. The Texas Journal of Science 18: 25-56.
- Reddell, J.R. 1966. A checklist of the cave fauna of Texas. II. Insecta. The Texas Journal of Science 18: 25-26.
- Reddell, J.R. 1998. Troglobitic ground beetles of the genus *Rhadine* from Bexar County, Texas. A report prepared for the Texas Parks and Wildlife Foundation.
- Reddell, J.R. and Cokendolpher, J.C., 2004. The cave spiders of Bexar and Comal counties, Texas. Texas Memorial Museum, Speleological Monographs, 6:75-94.

- Service (U.S. Fish and Wildlife Service). 2000. Endangered and threatened wildlife and plants; final rule to list nine Bexar County, Texas invertebrate species as endangered. Federal Register 65: 81419-81433.
- Ubick, D. and T.S. Briggs. 1992. The harvestman family Phalangodidae. 3. Revision of *Texella* Goodnight and Goodnight (Opiliones: Laniatores). Pages 155-240 *in* J. R. Reddell, editor. Texas Memorial Museum Speleological Monographs, 3, Studies on the Cave and Endogean Fauna of North America IV. Texas Memorial Museum, Austin, Texas.
- Ubick, D., P. Paquin, P.E. Cushing, and V. Roth editors. 2005. Spiders of North America: an identification manual. American Arachnological Society, New York.