

# Croaking Gourami (*Trichopsis vittata*)

## Ecological Risk Screening Summary

U.S. Fish & Wildlife Service, February 2011  
Revised, April 2019  
Web Version, 9/10/2019



Photo: BEDO. Licensed under the Creative Commons Attribution-Share Alike 4.0 International.  
Available: [https://commons.wikimedia.org/wiki/File:Trichopsis\\_vittata.jpg](https://commons.wikimedia.org/wiki/File:Trichopsis_vittata.jpg).

## 1 Native Range and Status in the United States

---

### Native Range

From Froese and Pauly (2019):

“Asia: Thailand to Viet Nam and the islands of Sumatra, Borneo and Java.”

From Low (2019):

“The species occurs widely in Java, Sumatra, Borneo (Kalimantan, Sarawak), the Malay Peninsula (Singapore, Peninsular Malaysia, southern Thailand), and the Mekong and Chao Phraya River drainages of Indochina.”

## **Status in the United States**

From Nico et al. (2019):

“Locally established in Palm Beach County, Florida, from at least the late 1970s (Lee et al. 1980 et seq., Courtenay and Stauffer 1990). Thought to be extirpated (Shafland et al. 2008a, b); however, recent collections (October 2012-July 2014) reveal it is in fact established (Schofield and Pecora 2013).”

*Trichopsis vittata* is in trade in the United States.

From TRiN’s Tropical Fish (2019):

“Croaking Gourami *Trichopsis vittata*  
\$3.00”

## **Means of Introductions in the United States**

From Froese and Pauly (2019):

“Accidentally released from aquarium fish farms and became established since at least in the late 1970s [Lever 1996] in a localized area on the south side of Lake Worth Drainage District canal L-36, Delray Beach, Palm Beach County, Florida.”

## **Remarks**

From Low (2019):

“Recent genetic evidence has suggested that what is currently considered *Trichopsis vittata* may in fact comprise a complex of four ecologically- and morphologically-similar, but genetically distinct species (Noren et al. 2017). Nonetheless, until a comprehensive revisionary study of this taxon has been carried out, all known populations are tentatively considered conspecific.”

From Panijpan et al. (2015):

“Our DNA analyses confirmed that some wild-caught fish from these places were hybrids as a result of crossing between female *T. vittata* and male *T. pumila*: the fish inherited the COI gene from maternal *T. vittata* and the RAG1 gene from paternal *T. pumila*.”

## 2 Biology and Ecology

---

### Taxonomic Hierarchy and Taxonomic Standing

From Fricke et al. (2019):

“**Current status:** Valid as *Trichopsis vittata* (Cuvier 1831).”

From ITIS (2019):

“Kingdom Animalia  
Subkingdom Bilateria  
Infrakingdom Deuterostomia  
Phylum Chordata  
Subphylum Vertebrata  
Infraphylum Gnathostomata  
Superclass Actinopterygii  
Class Teleostei  
Superorder Acanthopterygii  
Order Perciformes  
Suborder Anabantoidei  
Family Osphronemidae  
Subfamily Macropodinae  
Genus *Trichopsis*  
Species *Trichopsis vittata* (Cuvier in Cuvier and Valenciennes, 1831)”

### Size, Weight, and Age Range

From Froese and Pauly (2019):

“Max length : 7.0 cm SL male/unsexed; [Baird et al. 1999]; common length : 3.9 cm TL male/unsexed; [Hugg 1996]”

### Environment

From Froese and Pauly (2019):

“Freshwater; demersal; pH range: 6.0 - 8.0; dH range: 5 - 19. [...]; 22°C - 28°C [Riehl and Baensch 1991; assumed to be recommended aquarium temperature]”

From Noren et al. (2017):

“It is able to survive in brackish water up to 20 psu salinity, and in temperatures down to 7.2 °C (Schofield and Schulte 2016).”

## Climate/Range

From Froese and Pauly (2019):

“Tropical”

## Distribution Outside the United States

Native

From Froese and Pauly (2019):

“Asia: Thailand to Viet Nam and the islands of Sumatra, Borneo and Java.”

From Low (2019):

“The species occurs widely in Java, Sumatra, Borneo (Kalimantan, Sarawak), the Malay Peninsula (Singapore, Peninsular Malaysia, southern Thailand), and the Mekong and Chao Phraya River drainages of Indochina.”

Introduced

From Akash and Hossain (2017):

“The species [*Trichopsis vittata*] was first found in Bangladesh in 2012 (Hossain et al. 2012). Morphometric and morphomeristic attributes of feral specimens collected from Bangladesh have been documented (Islam 2014) and it was again reported by Hossain (2014).”

“Figure 2 and Table S1 [in source material] clearly denote the spread of *T. vittata* in Bangladesh over the last four years with a north-eastwardly colonization pattern.”

From Knight and Balasubramanian (2015):

“During recent (April to June 2014) surveys of the water bodies around Chennai [India], we recorded two species, *Trichopsis vittata* and *Macropodus opercularis* belonging to the family Osphronemidae. Specimens of *Trichopsis vittata* [...] were collected from an irrigation channel that conveys water from the Chembarampakkam Lake [...].”

It is unknown if the collection in India represents an established population.

According to Froese and Pauly (2019), *T. vittata* has been reported as introduced to the Philippines but it is unknown if the population has become established.

## Means of Introduction Outside the United States

From Akash and Hossain (2017):

“We opine that the establishment of *T. vittata* in the country is likely owing to aquaculture acting as an invasion vector.”

From Knight and Balasubramanian (2015):

“There is an unconfirmed report of a well established non-native population in Dhaka, Bangladesh [has been confirmed since this publication] (Andrew Rao pers. com. 2014). The role of inter-basin water transfer playing a role in introducing non-native species into an ecosystem cannot be ruled out.”

## Short Description

From Froese and Pauly (2019):

“Anal spines: 6-8. Has 24-28 branched anal-fin rays; 13 transverse scale rows; and 2 or more stripes along the body [Rainboth 1996]. Anal fin with a few elongate, filament-like rays extending backwards almost to tip of caudal fin; black blotch above pectoral base [Kottelat 2001].”

## Biology

From Froese and Pauly (2019):

“Found in flooded fields in the middle Mekong [Taki 1978]. Feed on zooplankton, crustaceans and insect larvae. Eggs sink to the bottom and were then collected by one or both parents and were embedded among the foam bubbles in the nest [Britz and Cambray 2001]. An air-breathing species [Huang et al. 2011].”

From Low (2019):

“The species occurs in a wide range of habitats, including freshwater marshes, ponds, lakes, peatlands, slow-flowing streams, floodplains, rice fields, irrigation canals and roadside ditches. It is partial to sluggish, open habitats with dense growths of submerged vegetation, and can tolerate a degree of anthropogenic disturbance and pollution (Linke, 1991, Ng and Tan 2005, Noren et al. 2017, pers. obs. B.W. Low 2012–2018).”

From Noren et al. (2017):

“The male is territorial and builds a small inconspicuous bubble nest among the vegetation, in which the eggs are deposited and guarded by the male until the larvae become free-swimming (Britz et al. 2001, Liengpornpan et al. 2006).”

From Ladich and Schleinzer (2015):

“[...] *Trichopsis vittata*, a freshwater fish from Southeast Asia possessing a highly specialized sound-generating mechanism found only in a single genus. The croaking gourami produces pulsed sounds by stretching and plucking two enhanced pectoral fin tendons during rapid pectoral fin beating. Croaking sounds typically consist of a series of double-pulsed bursts with main energies between 1 and 1.5 kHz.”

From Bischof (1996):

“[...] *T. vittata* was the only species to utter croaks in series [Ladich et al., 1992a].”

## Human Uses

From Low (2019):

“There is no targeted fishery for the species, but it is occasionally sold in markets as part of mixed catches. It is commonly seen in the aquarium trade (Noren et al. 2017).”

## Diseases

**No records of OIE-reportable diseases (OIE 2019) were found for *Trichopsis vittata*.**

According to Poelen (2014), *Trichopsis vittata* is a host for *Heteronchocleidus bushkielli*, and *Paracapillaria philippinensis*.

According to Senapin et al. (2014), *T. vittata* is a host for *Euclinostomum* sp. infection.

## Threat to Humans

From Froese and Pauly (2019):

“Harmless”

## 3 Impacts of Introductions

---

From Nico et al. (2019):

“The impacts of this species are currently unknown, as no studies have been done to determine how it has affected ecosystems in the invaded range. The absence of data does not equate to lack of effects. It does, however, mean that research is required to evaluate effects before conclusions can be made.”

The following concerns *potential* impacts from introduction for *Trichopsis vittata*:

From Knight and Balasubramanian (2015):


“Though the ecological impacts of neither *Trichopsis vittata* or *Macropodus opercularis* have been documented, it is quite obvious that like *T. trichopterus*, they would also compete for niche space with native species such as *Trichogaster lalius* and *Pseudosphromenus cupanus* (Knight 2010b). As both *T. vittata* and *M. opercularis* are air-breathers and exhibit parental care, they will be able to ensure better survival of their young, even in stagnant and polluted water systems with high biological oxygen demand (BOD) where other fish species may not be able to survive and reproduce.”

## 4 Global Distribution

---



**Figure 1.** Known global distribution of *Trichopsis vittata*. Locations are in Bangladesh, Myanmar, Thailand, Laos, Cambodia, Vietnam, Malaysia, Indonesia, Sweden, and the United States Map from GBIF Secretariat (2019). The location in Sweden does not represent an established population, the specimen was purchased from a pet shop, and was not used to select source points for the climate match.



**Figure 2.** Known distribution of *Trichopsis vittata* in Bangladesh. Map from Akash and Hossain (2017; Figure 2), licensed under Creative Commons BY 4.0 International.



## 5 Distribution Within the United States

---




**Figure 3.** Known distribution of *Trichopsis vittata* in the United States. Map from Nico et al. (2019).


## 6 Climate Matching

### Summary of Climate Matching Analysis

The climate match was low for most of the contiguous United States. Peninsular Florida has a high match. The southern Atlantic Coast had a medium match along with the Texas Gulf Coast and small areas of the Gulf Coast in Louisiana and Georgia. The Climate 6 score (Sanders et al. 2018; 16 climate variables; Euclidean distance) for contiguous United States was 0.018, medium (scores greater than 0.005, but less than 0.103, are classified as medium). The only State to receive a high individual Climate 6 score is Florida. Georgia, South Carolina, and Texas all received medium individual Climate 6 scores while all remaining States received low individual Climate 6 scores.



**Figure 4.** RAMP (Sanders et al. 2018) source map showing weather stations in Southeast Asia and in Florida in the United States selected as source locations (red; Bangladesh, India, Myanmar, Laos, Thailand, Cambodia, Vietnam, Malaysia, Indonesia, United States) and non-source locations (gray) for *Trichopsis vittata* climate matching. Source locations from Akash and Hossain (2017) and GBIF Secretariat (2019). Selected source locations are within 100 km of one or more species occurrences, and do not necessarily represent the locations of occurrences themselves.



**Figure 5.** Map of RAMP (Sanders et al. 2018) climate matches for *Trichopsis vittata* in the contiguous United States based on source locations reported by Akash and Hossain (2017) and GBIF Secretariat (2019). 0 = Lowest match, 10 = Highest match.

The High, Medium, and Low Climate match Categories are based on the following table:

Climate 6: Proportion of (Sum of Climate Scores 6-10) / (Sum of total Climate Scores)	Climate Match Category
$0.000 \leq X \leq 0.005$	Low
$0.005 < X < 0.103$	Medium
$\geq 0.103$	High

## 7 Certainty of Assessment

The certainty of this assessment is low. There is an abundance of information available on the biology, environment and distribution of *Trichopsis vittata*. There are records of introduction and at least one established population. No information was found regarding actual impacts of introduction.

## 8 Risk Assessment

---

### Summary of Risk to the Contiguous United States

The Croaking Gourami (*Trichopsis vittata*) is a freshwater fish native to Southeast Asia. Native to Cambodia, Laos, Malaysia, Myanmar, Singapore, Thailand, Vietnam, and Indonesia. *T. vittata* is a unique air-breathing fish that produces pulsed sounds by stretching and plucking their two enhanced pectoral fin tendons. This species is regularly found in the aquarium trade. Croaking Gourami have been introduced outside of their native range. *T. vittata* has become established in Florida in the United States where they were accidentally released from an aquarium fish farm. A non-native, established population is also present in Bangladesh and *T. vittata* has been reported as introduced in India and the Philippines but it is uncertain if these populations have become established. The overall climate match for the contiguous United States is medium. Florida and coastal Texas had a high match, much of the South Atlantic and Gulf Coasts had a medium match, and all remaining areas had a low match. The certainty of assessment is low. The overall risk assessment category of *Trichopsis vittata* is uncertain.

### Assessment Elements

- **History of Invasiveness (Sec. 3): None Documented**
- **Climate Match (Sec. 6): Medium**
- **Certainty of Assessment (Sec. 7): Low**
- **Remarks/Important additional information:** No additional information.
- **Overall Risk Assessment Category: Uncertain**

## 9 References

---

**Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.**

- Akash, M., and M. A. R. Hossain. 2018. A Southeast Asian species in the Ganges Delta: on spreading extent of non-native Croaking Gourami *Trichopsis vittata* (Cuvier, 1831) in Bangladesh. *BioInvasions Records* 7(4):447–450.
- Bischof, C. 1996. Diversity in the agonistic behavior of Croaking Gourami (*Trichopsis vittata*, *T. schalleri*, and *T. pumila*; Anabantoidei) and the Paradise Fish (*Macropodus opercularis*; Anabantoidei). *Aggressive Behavior* 22:447–455.
- Fricke, R., Eschmeyer, W. N., and R. van der Laan, editors. 2019. Eschmeyer's catalog of fishes: genera, species, references. Available: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. (April 2019).
- Froese, R., and D. Pauly, editors. 2019. *Trichopsis vittata* (Cuvier, 1831). FishBase. Available: <http://www.fishbase.org/summary/Trichopsis-vittata.html>. (April 2019).

- GBIF Secretariat. 2019. GBIF backbone taxonomy: *Trichopsis vittata* (Cuvier, 1831). Global Biodiversity Information Facility, Copenhagen. Available: <https://www.gbif.org/species/5211201>. (September 2019).
- ITIS (Integrated Taxonomic Information System). 2019. *Trichopsis vittata* (Cuvier in Cuvier and Valenciennes, 1831). Integrated Taxonomic Information System. Reston, Virginia. Available: [https://www.itis.gov/servlet/SingleRpt/SingleRpt?search\\_topic=TSN&search\\_value=553301#null](https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=553301#null). (April 2019).
- Knight, J. D. M., and S. Balasubramanian. 2015. On a record of two alien fish species (Teleostei: Osphronemidae) from the natural waters of Chennai, Tamil Nadu, India. *Journal of Threatened Taxa* 7(3):7044–7046.
- Ladich, F., and G. Schleinzer. 2015. Effect of temperature on acoustic communication: sound production in the Croaking Gourami (labyrinth fishes). *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology* 182:8–13.
- Low, B. W. 2019. *Trichopsis vittata*. The IUCN Red List of Threatened Species 2019: e.T187906A89805994. Available: <https://www.iucnredlist.org/species/187906/89805994>. (September 2019).
- Nico, L., P. Fuller, and P. Schofield. 2019. *Trichopsis vittata* (Cuvier in Cuvier and Valenciennes, 1831). U.S. Geological Survey, Nonindigenous Aquatic Species Database, Gainesville, Florida. Available: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=333>.
- Noren, M., S. O. Kullander, M. M. Rahman, and A. R. Mollah. 2017. First records of Croaking Gourami, *Trichopsis vittata* (Cuvier, 1831) (Teleostei: Osphronemidae), from Myanmar and Bangladesh. *Check List* 13(4):81–85.
- OIE (World Organisation for Animal Health). 2019. OIE-listed diseases, infections and infestations in force in 2019. Available: <http://www.oie.int/animal-health-in-the-world/oie-listed-diseases-2019/>. (August 2019).
- Poelen, J. H., J. D. Simons, and C. J. Mungall. 2014. Global Biotic Interactions: an open infrastructure to share and analyze species-interaction datasets. *Ecological Informatics* 24:148–159.
- Sanders, S., C. Castiglione, and M. Hoff. 2018. Risk assessment mapping program: RAMP, version 3.1. U.S. Fish and Wildlife Service.
- Senapin, S., K. Phiwaiya, P. Laosinchai, C. Kowasupat, P. Ruenwongsa, and B. Panijpan. 2014. Phylogenetic analysis of parasitic trematodes of the genus *Euclinostomum* found in *Trichopsis* and betta fish. *Journal of Parasitology* 100(3):368–371.



TRiN's Tropical Fish. 2019. Croaking Gourami *Trichopsis vittata*. Available: <http://trinsfish.com/croaking-gourami-trichopsis-vittata/>. (August 2019).

## 10 References Quoted But Not Accessed

---

**Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.**

- Baird, I. G., V. Inthaphaisy, P. Kisouvannalath, B. Phylavanh, and B. Mounsouphom. 1999. The fishes of southern Lao. Lao Community Fisheries and Dolphin Protection Project. Ministry of Agriculture and Forestry, Lao PDR.
- Britz, R., and J. A. Cambray. 2001. Structure of egg surfaces and attachment organs in anabantoids. *Ichthyological Explorations of Freshwaters* 12(3):267–288.
- Courtenay, W. R., Jr., and J. R. Stauffer, Jr. 1990. The introduced fish problem and the aquarium fish industry. *Journal of the World Aquaculture Society* 21(3):145–159.
- Hossain, M. A. R. 2014. Habitat and fish diversity: Bangladesh perspective. Pages 1–26 in M. A. Wahab, M. S. Shah, M. A. R. Hossain, B. K. Barman, and M. E. Hoq, editors. *Advances in fisheries research in Bangladesh: I. Proceedings of the 5th Fisheries Conference & Research Fair 2012, Dhaka, Bangladesh*. Bangladesh Agricultural Research Council, Dhaka, Bangladesh Fisheries Research Forum, Dhaka.
- Hossain, M. A. R., M. A. Wahab, and B. Belton. 2012. The checklist of the riverine fishes of Bangladesh. *FAN-Bangladesh* 2:29–39.
- Huang, C. Y., C. P. Lin, and H. C. Lin. 2011. Morphological and biochemical variations in the gills of 12 aquatic air-breathing anabantoid fish. *Physiological and Biochemical Zoology* 84(2):125–134.
- Hugg, D. O. 1996. MAPFISH georeferenced mapping database. *Freshwater and estuarine fishes of North America*. Life Science Software. Dennis O. and Steven Hugg, Edgewater, Maryland.
- Islam, M. S. 2014. Morphometric, meristic and landmark-based analysis of a rare indigenous Croaking gourami, *Trichopsis vittata*. Master's thesis. Bangladesh Agricultural University, Mymensingh, Bangladesh.
- Knight, J. D. M. 2010b. Invasive ornamental fish: a potential threat to aquatic biodiversity in peninsular India. *Journal of Threatened Taxa* 2(2):700–704.
- Kottelat, M. 1985. Fresh-water fishes of Kampuchea. *Hydrobiologia* 121:249–279.
- Kottelat, M. 2001. [no title given] Page 198 in *Fishes of Laos*. WHT Publications, Colombo 5, Sri Lanka.

- Ladich, F., C. Bischof, G. Schleinzer, and A. Fuchs. 1992a. Intra- and interspecific differences in agonistic vocalization in Croaking Gouramis (Genus *Trichopsis*, Anabantoidei, Teleostei). *Bioacoustics* 4:131–141.
- Lee, D. S., C. R. Gilbert, C. H. Hocutt, R. E. Jenkins, D. E. McAllister, and J. R. Stauffer, Jr. 1980. Atlas of North American freshwater fishes. North Carolina State Museum of Natural History, Raleigh.
- Lever, C. 1996. [no title given] Page 408 in Naturalized fishes of the world. Academic Press, California.
- Liengpornpan, S., M. Jaroensutasinee, and K. Jaroensutasinee. 2006. Mating habits and nesting habitats of the Croaking Gourami *Trichopsis vittata*. *Acta Zoologica Sinica* 52(5):846–853.
- Rainboth, W. J. 1996. [no title given] Page 265 in *Fishes of the Cambodian Mekong*. FAO species identification field guide for fishery purposes. FAO, Rome.
- Riehl, R., and H. A. Baensch. 1991. [no title given] Page 992 in *Aquarien Atlas*. Band. 1. Melle: Mergus, Verlag für Natur-und Heimtierkunde, Germany.
- Schofield, P. J., and D. J. Pecora. 2013. Croaking Gourami, *Trichopsis vittata* (Cuvier, 1831), in Florida, USA. *BioInvasions Records* 2:247–251.
- Schofield, P. J., and J. M. Schulte. 2016. Small but tough: what can ecophysiology of Croaking Gourami *Trichopsis vittata* (Cuvier, 1831) tell us about invasiveness of non-native fishes in Florida? *NeoBiota* 28:51–65.
- Taki, Y. 1978. An analytical study of the fish fauna of the Mekong basin as a biological production system in nature. Research Institute of Evolutionary Biology, Special Publications 1, Tokyo, Japan.