

Variable Platyfish (*Xiphophorus variatus*)

Ecological Risk Screening Summary

U.S. Fish & Wildlife Service, January 2017
Revised, December 2017
Web Version, 10/23/2019



Photo: Marrabbio2. Licensed under Creative Commons Attribution-Share Alike 3.0 Unported License. Available: https://commons.wikimedia.org/wiki/File:Xipho_variatus_maschio.jpg. (January 1, 2017).

1 Native Range and Status in the United States

Native Range

From Froese and Pauly (2017):

“North America: endemic to Mexico from southern Tamaulipas to northern Veracruz.”

From Culumber and Rosenthal (2013):

“*Xiphophorus variatus* is a species of platyfish inhabiting streams of north- and east-central Mexico from the coastal plain to the foothills of the Sierra Madre Oriental.”

Status in the United States

From Froese and Pauly (2017):

“Introduced in the 1960s. Suspected self-sustaining populations reported to exist on the island of Oahu. However, the species is not contained in the 1992 list of introduced and established freshwater fishes of Hawaii. Also [Maciolek 1984].”

From NatureServe (2016):

“Established in springs in Beaverhead, Granite, and Madison counties, Montana; in canals along the eastern shore of Tampa Bay, Hillsborough County, Florida; and in Gainesville, Alachua County, Florida; uncommon (Page and Burr 1991). Populations reported from Arizona and California apparently are no longer extant.”

From Nico (2017):

“This species has been recorded from Yuma and Tempe, Arizona (Minckley 1973); several sites in southern California (St. Amant and Hoover 1969; St. Amant and Sharp 1971; Moyle 1976; Hubbs et al. 1979; Shapovalov et al. 1981; Courtenay et al. 1984, 1991; Swift et al. 1993); Colorado (Zuckerman and Behnke 1986); as many as six counties in Florida (Courtenay et al. 1974; Burgess et al. 1977; J. D. Williams, personal observation; museum specimens); Oahu, Hawaii (Maciolek 1984; Mundy 2005); and Beaverhead, Broadwater, Granite, and Madison counties, Montana (Brown 1971; Courtenay 1985; Holton 1990).”

“Locally established in Florida (Courtenay and Meffe 1989; museum specimens): a population in Alachua County, Florida, first recorded in the 1970s (Burgess et al. 1977), still survives. Locally established in Montana (Courtenay and Meffe 1989); possibly established locally in California (Swift et al. 1993). Possibly established on Oahu, Hawaii: listed as established by Maciolek (1984), but this species is not listed in the state by recent reports (Devick 1991; Yamamoto and Tagawa 2000; Mundy 2005). It was established locally in Arizona from 1963 to 1965, but that population was presumably destroyed by a flood in late 1965 (Minckley 1973). The species was reported [as extirpated] from Colorado (Zuckerman and Behnke 1986).”

“It has been popular as an ornamental fish for many years.”

From Courtenay and Stauffer (1990):

“As with other species of *Xiphophorus*, variable platyfish are known to have escaped from aquarium fish farms in Hillsborough County, Florida, and in Orange and Riverside counties, California (St. Amant and Hoover 1969; St. Amant and Sharp 1971; Courtenay et al. 1974). None of these introductions results in permanent establishment. The species was established in canals in Tempe, Arizona, but was extirpated by a flood; individuals have been collected periodically from drains near Yuma, Yuma County, Arizona, without evidence of establishment (Minckley 1973). The species is established in Gainesville, Alachua County, Florida (Burgess et al. 1977), probably from introductions by university students (Courtenay et al. 1986). Brown

(1971) reported established populations, released by aquarists, in thermal outflows in Beaverhead, Granite, and Madison counties, Montana; the population in Beaverhead County remains extant (Courtenay et al. 1986).”

From Cohen et al. (2014):

“Here we provide the first report of establishment of this species in Texas, where it has persisted for at least a decade in Waller Creek in the city of Austin (Travis County).”

Means of Introductions in the United States

From Courtenay and Stauffer (1990):

“As with other species of *Xiphophorus*, variable platyfish are known to have escaped from aquarium fish farms in Hillsborough County, Florida, and in Orange and Riverside counties, California (St. Amant and Hoover 1969; St. Amant and Sharp 1971; Courtenay et al. 1974). [...] The species is established in Gainesville, Alachua County, Florida (Burgess et al. 1977), probably from introductions by university students (Courtenay et al. 1986). Brown (1971) reported established populations, released by aquarists, in thermal outflows in Beaverhead, Granite, and Madison counties, Montana; the population in Beaverhead County remains extant (Courtenay et al. 1986).”

From Nico (2017):

“Probably due to aquarium releases.”

From Froese and Pauly (2017):

“Accidentally released from aquaria and populations were found to be reproducing but it is uncertain whether it will become established in Florida [Shafland 1979].”

Remarks

From Magalhães and Jacobi (2017):

“This situation is already occurring in the area (Magalhães et al., 2011), as well as the probable intra-specific hybridization between *X. maculatus* and *X. variatus* (A.L.B. Magalhães, pers. obs.).”

From Cohen et al. (2014):

“This population likely includes genes from *X. maculatus* and/or *X. hellerii* that may enhance cold tolerance. *Xiphophorus hellerii* occurs in the wild at higher altitudes than *X. variatus* (Miller et al., 2005) and may have greater cold tolerance.”

2 Biology and Ecology

Taxonomic Hierarchy and Taxonomic Standing

From Eschmeyer et al. (2017):

“**Current status:** Valid as *Xiphophorus variatus* (Meek 1904).”

From ITIS (2017):

“Kingdom Animalia
Subkingdom Bilateria
Infrakingdom Deuterostomia
Phylum Chordata
Subphylum Vertebrata
Infraphylum Gnathostomata
Superclass Osteichthyes
Class Actinopterygii
Subclass Neopterygii
Infraclass Teleostei
Superorder Acanthopterygii
Order Cyprinodontiformes
Suborder Cyprinodontoidei
Family Poeciliidae
Subfamily Poecilinae
Genus *Xiphophorus*
Species *Xiphophorus variatus*”

Size, Weight, and Age Range

From Froese and Pauly (2017):

“Max length : 7.0 cm TL male/unsexed; [Page and Burr 1991]; common length : 3.9 cm TL male/unsexed; [Hugg 1996]”

From NatureServe (2016):

“Length: 6 centimeters”

Environment

From Froese and Pauly (2017):

“Freshwater; benthopelagic; pH range: 7.0 - 8.0; dH range: 9 - 19; non-migratory.”

“15°C - 25°C [Riehl and Baensch 1991] [assumed to be recommended aquarium temperature range]”

From Cohen et al. (2014):

“Borowsky reports them occurring in water temperatures of 17°C (Borowsky, 1984) and 18–24°C (Borowsky, 1990) in winter and spring at a spring dominated site in the species’ native range, but presumably the species can tolerate lower temperatures. Dawes (2001) puts the species’ lower limit at 16°C, and Hoedman (1974) reports it tolerates temperatures as low as 15°C.”

“The population had apparently survived a period of at least 4 days at or near 7°C, and their improved equilibrium suggests that they acclimated to the lower temperature.”

Climate/Range

From Froese and Pauly (2017):

“Subtropical; [...]; 25°N - 20°N”

Distribution Outside the United States

Native

From Froese and Pauly (2017):

“North America: endemic to Mexico from southern Tamaulipas to northern Veracruz.”

From Culumber and Rosenthal (2013):

“*Xiphophorus variatus* is a species of platyfish inhabiting streams of north- and east-central Mexico from the coastal plain to the foothills of the Sierra Madre Oriental.”

Introduced

Established populations of *Xiphophorus variatus* resulting from introductions have been documented in Hong Kong, Costa Rica, and Colombia. Probably established populations of *X. variatus* resulting from introductions have been documented in Singapore. An introduction to Spain was listed but no further information was available (Froese and Pauly 2017).

From Froese and Pauly (2017):

“Established in the Magdalena and Orinoco watersheds [in Colombia].”

“Recorded from several tributaries on the Río Reventazón drainage [in Costa Rica] [Bussing 1998].”

FAO (2017) lists *Xiphophorus variatus* as introduced to and established in Colombia, Puerto Rico, Hong Kong, and Costa Rica; introduced and probably established in Singapore.

From Magalhães and Jacobi (2017):

“We found females of *P. reticulata*, *P. sphenops*, *X. hellerii*, *X. maculatus*, and *X. variatus* in reproductive activity during the 12 months of the year in almost all creeks [in Paraíba do Sul River basin, southeastern Brazil], [...]”

From Lee et al. (2006):

“Eight of the ten most common species recorded [in 220 study sites in Hong Kong, China] were native species. Two exotics were *Gambusia affinis* and *Xiphophorus variatus*.”

Means of Introduction Outside the United States

FAO (2017) listed reasons for introductions of *Xiphophorus variatus* include aquaculture, ornamental, and accidental.

From Froese and Pauly (2017):

“Widespread in fish rearing facilities and has presumably escaped into local waters [in Colombia].”

Short Description

From Nico (2017):

“Similar to some of the other poeciliids, this species exhibits considerable natural variation (e.g., Wischnath 1993).”

From Culumber and Rosenthal (2013):

“This species exhibits high polymorphism at the autosomal tailspot locus, with at least 6 known patterns that are exhibited in both sexes (Borowsky 1984).”

Biology

From Froese and Pauly (2017):

“[...] non-migratory.”

“Occurs in warm springs, weedy canals and ditches [Page and Burr 1991]. Feeds on worms, crustaceans, insects and plant matter [Mills and Vevers 1989].”

“After 24 days gestation, up to 100 young are born.”

“Reproductive guild: bearers, internal live bearers”

From NatureServe (2016):

“Usually lentic habitats in native range. Has been collected from habitats with low oxygen levels. Apparently has little tolerance for brackish water. North America: warm springs, weedy canals and ditches (Page and Burr 1991).”

“Food Comments: Feeds during day primarily on mud or bottom ooze.”

Human Uses

From Froese and Pauly (2017):

“aquarium: commercial”

“One of the most frequently found species in the pet and aquarium stores [in Spain] [Maceda-Veiga et al. 2013].

“Used for genetic research [Robins et al. 1991].”

From Nico (2017):

“It has been popular as an ornamental fish for many years.”

Diseases

No records of OIE-reportable diseases (OIE 2019) were found for *Xiphophorus variatus*.

From Froese and Pauly (2017):

“Fin-rot Disease (late stage), Bacterial diseases
White spot Disease, Parasitic infestations (protozoa, worms, etc.)
Fin Rot (early stage), Bacterial diseases
Skin Fungi (*Saprolegnia* sp.), Fungal diseases
Turbidity of the Skin (Freshwater fish),
Parasitic infestations (protozoa, worms, etc.)
Bacterial Infections (general), Bacterial diseases
Fin-rot Disease (late stage), Bacterial diseases
Costia Disease, Parasitic infestations (protozoa, worms, etc.)
Ichthyobodo Infection, Parasitic infestations (protozoa, worms, etc.)
Fish tuberculosis (FishMB), Bacterial diseases”

Poelen et al. (2014) list *Bothriocephalus acheilognathi*, *Centrocestus formosanus*, *Eustrongylides ignotus*, *Posthodiplostomum minimum*, and *Valipora campylancristrota* as parasites of *Xiphophorus variatus*.

Mendoza et al. (2015) report that *Xiphophorus variatus* could be a vector for introduction of the pathogen *Mycobacterium marium*.

Magalhães (2006) reports infection with *Lernaea* sp.

Threat to Humans

From Froese and Pauly (2017):

“Potential pest [FAO 1997]”

3 Impacts of Introductions

The follow information concerns *potential* impacts of introductions or broad generalizations.

From Froese and Pauly (2017):

“At least one country reports adverse ecological impact after introduction.”

“The introduction of these two species (*Xiphophorus helleri* and *X. variatus*) is likely to have affected native cyprinid species such as *Aphyocypris lini*, the native endemic Hong Kong minnow.”

From Nico (2017):

“Impact of Introduction [in the United States]: Unknown.”

From Magalhães and Jacobi (2017):

“The presence of reproductive adults and juveniles of non-native poeciliids in these creeks, besides characterizing establishment by recruitment, may lead to biotic homogenization, i.e. the establishment of the same non-native species at two or more locations, decreasing beta-diversity over time (Olden et al., 2011). This situation is already occurring in the area (Magalhães et al., 2011), as well as the probable intra-specific hybridization between *X. maculatus* and *X. variatus* (A.L.B. Magalhães, pers. obs.)”

From Cohen et al. (2014):

“However, in some areas of Waller Creek, it [*X. variatus*] has become the dominant fish species, suggesting that it may be strongly impacting the community. It has replaced native fishes near Monterrey, Nuevo León, in Mexico (S. Contreras, in litt., 1976), but in Hawaii may have negligible ecological impact compared to *X. maculatus* and especially *X. hellerii* (Maciolek, 1984), [...]”

4 Global Distribution



Figure 1. Known global distribution of *Xiphophorus variatus*. Locations are in the United States, Mexico, Costa Rica, Colombia, Dominican Republic, Mauritius, and China. Map from GBIF Secretariat (2017). The location in Mauritius was not used to select source points for the climate matching. The record information listed this observation as from a farm pond, therefore it was not representative of an established wild population.

An additional georeferenced locations in southern Brazil are provided in Magalhães (2006) and Magalhães and Jacobi (2017).

5 Distribution Within the United States




Figure 2. Known distribution of *Xiphophorus variatus* in the contiguous United States. Map from BISON (2017). The locations in Montana were not used to select source points for the climate match. Those populations only persist in thermal outflows (Courtenay and Stauffer 1990). The conditions in a thermal outflow would not be reflected accurately in the climate variables used in RAMP for those locations therefore including them as source points would result in an incorrect climate match. The locations in Arizona, California, and Colorado were not used to select source points for the climate match; they either did not establish populations or were extirpated (Courtenay and Stauffer 1990; NatureServe 2016; Nico 2017).




Figure 3. Known distribution of *Xiphophorus variatus* in Hawai'i. Map from BISON (2017).

6 Climate Matching

Summary of Climate Matching Analysis

The climate match for *Xiphophorus variatus* was high along the southern Atlantic Coast, the eastern and western ends of the Gulf Coast, most of Texas, and in some areas in the southwest. The rest of the United States was low to medium match. The Climate 6 score (Sanders et al. 2018; 16 climate variables; Euclidean distance) for the contiguous United States was 0.161, high (scores 0.103 and greater are classified as high). Most States had low individual Climate 6 scores, except for Arizona, Florida, Georgia, New Mexico, North Carolina, Oklahoma, South Carolina, and Texas, which had high individual scores, and Alabama, Arizona, California, Kansas, and Louisiana, which had medium individual scores.




Figure 4. RAMP (Sanders et al. 2018) source map showing weather stations selected as source locations (red; China, United States (Florida, Texas, Hawaii), Mexico, Costa Rica, Colombia, Dominican Republic) and non-source locations (gray) for *Xiphophorus variatus* climate matching. Source locations from Courtenay and Stauffer (1990), Magalhães (2006), NatureServe (2016), BISON (2017), GBIF Secretariat (2017), and Magalhães and Jacobi (2017). Selected source locations are within 100 km of one or more species occurrences, and do not necessarily represent the locations of occurrences themselves.




Figure 5. Map of RAMP (Sanders et al. 2018) climate matches for *Xiphophorus variatus* in the contiguous United States based on source locations reported by Courtenay and Stauffer (1990), Magalhães (2006), NatureServe (2016), BISON (2017), GBIF Secretariat (2017), Magalhães and Jacobi (2017). Counts of climate match scores are tabulated on the left. 0 = Lowest match, 10 = Highest match.

The High, Medium, and Low Climate match Categories are based on the following table:

Climate 6: Proportion of (Sum of Climate Scores 6-10) / (Sum of total Climate Scores)	Climate Match Category
$0.000 \leq X \leq 0.005$	Low
$0.005 < X < 0.103$	Medium
≥ 0.103	High

7 Certainty of Assessment

The certainty of this assessment is low. There was information regarding the ecology, biology, and distribution of *Xiphophorus variatus*. Many records of introductions were found that resulted in established populations. Many records of potential impacts were found but no scientifically defensible records of documented impacts.

8 Risk Assessment

Summary of Risk to the Contiguous United States

Variable Platyfish (*Xiphophorus variatus*) is a freshwater fish native to Mexico. The species is popular in the aquarium industry and has been used as a research subject. *X. variatus* is in trade within the United States. The history of invasiveness for *X. variatus* is None Documented. There were numerous records of introductions resulting in established populations. There were many records of potential impacts but no scientifically defensible information on documented impacts. The overall climate match is high. Most of the contiguous United States had a low of medium match with the areas of high match in the Southeast and Gulf. The certainty of assessment is low due to lack of information on impacts. The overall risk assessment category is uncertain.

Assessment Elements

- **History of Invasiveness (Sec. 3): None Documented**
- **Climate Match (Sec. 6): High**
- **Certainty of Assessment (Sec. 7): Low**
- **Remarks/Important additional information** No additional information.
- **Overall Risk Assessment Category: Uncertain**

9 References

Note: The following references were accessed for this ERSS. References cited within quoted text but not accessed are included below in Section 10.

BISON. 2017. Biodiversity Information Serving Our Nation (BISON). U.S. Geological Survey. Available: <https://bison.usgs.gov>. (December 2017).

Cohen, A. E., L. E. Dugan, D. A. Hendrickson, F. D. Martin, J. Juynh, B. J. Labay, and M. J. Casarez. 2014. Population of Variable Platyfish (*Xiphophorus variatus*) established in Waller Creek, Travis County, Texas. *The Southwestern Naturalist* 59(3):413-419.

Courtenay, W. R., and J. R. Stauffer. 1990. The introduced fish problem and the aquarium fish industry. *Journal of the World Aquaculture Society* 21(3):145-159.

Culumber, Z. W., and G. G. Rosenthal. 2013. Mating preferences do not maintain the tailspot polymorphism in the platyfish, *Xiphophorus variatus*. *Behavioral Ecology* 24(6):1286-1291.

Eschmeyer, W. N., R. Fricke, and R. van der Laan, editors. 2017. *Catalog of fishes: genera, species, references*. Available: <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.as> (December 2017).

- FAO (Fisheries and Agriculture Organization of the United Nations). 2017. Database on introductions of aquatic species. FAO, Rome. Available: <http://www.fao.org/fishery/introsp/search/en>. (December 2017).
- Froese, R., and D. Pauly, editors. 2017. *Xiphophorus variatus* (Meek, 1904). FishBase. Available: <http://www.fishbase.org/summary/Xiphophorus-variatus.html>. (December 2017).
- GBIF Secretariat. 2017. GBIF backbone taxonomy: *Xiphophorus variatus* (Meek, 1904). Global Biodiversity Information Facility, Copenhagen. Available: <https://www.gbif.org/species/2350220>. (December 2017).
- ITIS (Integrated Taxonomic Information System). 2017. *Xiphophorus variatus* (Meek, 1904). Integrated Taxonomic Information System, Reston, Virginia. Available: https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=165925. (December 2017).
- Lee, V. L., M. L. Young, T. K. Chan, S. K. Lam, F. K. Ng, and J. W. Chan. 2006. Biodiversity and conservation of freshwater fish in Hong Kong, China. *Acta Hydrobiologica Sinica* 30(6):660–666.
- Magalhães, A. L. B. 2006. First record of lernaeciosis in a native fish species from a natural environment in Minas Gerais state, Brazil. *Pan American Journal of Aquatic Sciences* 1(1):8–10.
- Magalhães, A. L. B., and C. M. Jacobi. 2010. E-commerce of freshwater aquarium fishes: potential disseminator of exotic species in Brazil. *Acta Scientiarum. Biological Sciences* 32(2):243–248.
- Magalhães, A. L. B., and C. M. Jacobi. 2017. Colorful invasion in permissive Neotropical ecosystems: establishment of ornamental non-native poeciliids of the genera *Poecilia/Xiphophorus* (Cyprinodontiformes: Poeciliidae) and management alternatives. *Neotropical Ichthyology* 15(1):e160094.
- Mendoza, R., S. Luna, and C. Aguilera. 2015. Risk assessment of the ornamental fish trade in Mexico: analysis of freshwater species and effectiveness of the FISK (Fish Invasiveness Screening Kit). *Biological Invasions* 17:3491–3502.
- NatureServe. 2016. NatureServe Explorer: An online encyclopedia of life, version 7.1. NatureServe, Arlington, Virginia. Available: <http://explorer.natureserve.org>. (December 2017).
- Nico, L. 2017. *Xiphophorus variatus* (Meek, 1904). U.S. Geological Survey, Nonindigenous Aquatic Species Database, Gainesville, Florida. Available: <https://nas.er.usgs.gov/queries/FactSheet.aspx?SpeciesID=874>. (December 2017).

OIE (World Organisation for Animal Health). 2019. OIE-listed diseases, infections and infestations in force in 2019. Available: <http://www.oie.int/animal-health-in-the-world/oie-listed-diseases-2019/>. (October 2019).

Poelen, J. H., J. D. Simons, and C. J. Mungall. 2014. Global Biotic Interactions: an open infrastructure to share and analyze species-interaction datasets. *Ecological Informatics* 24:148–159.

Sanders, S., C. Castiglione, and M. Hoff. 2018. Risk Assessment Mapping Program: RAMP, version 3.1. U.S. Fish and Wildlife Service.

10 References Quoted But Not Accessed

Note: The following references are cited within quoted text within this ERSS, but were not accessed for its preparation. They are included here to provide the reader with more information.

Borowsky, R. 1984. The evolutionary genetics of *Xiphophorus*. Pages 235–310 in B. J. Turner, editor. *Evolutionary genetics of fishes*. Plenum Press, New York.

Borowsky, R. 1990. Habitat choice by allelic variants in *Xiphophorus variatus* (Pisces; Poeciliidae) and implications for maintenance of genetic polymorphism. *Evolution* 44:1338–1345.

Brown, C. J. D. 1971. *Fishes of Montana*. Montana State University, Bozeman.

Burgess, G. H., C. R. Gilbert, V. Guillory, and D. C. Taphorn. 1977. Distributional notes on some north Florida freshwater fishes. *Florida Scientist* 40(1):33–41.

Bussing, W. A. 1998. *Peces de las aguas continentales de Costa Rica*, 2nd edition. Editorial de la Universidad de Costa Rica, San José.

Contreras, S. 1976. [Source material did not give full citation for this reference.]

Courtenay, W. R., Jr. 1985. Florida Atlantic University quarterly reports for 1985 to the U.S. Fish and Wildlife Service, Gainesville, Florida.

Courtenay, W. R., Jr., D. A. Hensley, J. N. Taylor, and J. A. McCann. 1984. Distribution of exotic fishes in the continental United States. Pages 41–77 in W. R. Courtenay, Jr. and J. R. Stauffer, Jr., editors. *Distribution, biology and management of exotic fishes*. Johns Hopkins University Press, Maryland.

Courtenay, W. R., Jr., D. A. Hensley, J. N. Taylor, and J. A. McCann. 1986. Distribution of exotic fishes in North America. Pages 675–698 in C. H. Hocutt, and E. O. Wiley, editors. *Zoogeography of North American freshwater fishes*. John Wiley & Sons, New York.

- Courtenay, W. R., Jr., D. P. Jennings, and J. D. Williams. 1991. Appendix 2: exotic fishes. Pages 97–107 in C. R. Robins, R. M. Bailey, C. E. Bond, J. R. Brooker, E. A. Lachner, R. N. Lea, and W. B. Scott. Common and scientific names of fishes from the United States and Canada, 5th edition. American Fisheries Society, Special Publication 20, Bethesda, Maryland.
- Courtenay, W. R., Jr., and G. K. Meffe. 1989. Small fishes in strange places: a review of introduced poeciliids. Pages 319–331 in G. K. Meffe and F. F. Snelson, Jr., editors. Ecology and evolution of livebearing fishes (Poecillidae). Prentice Hall, Englewood Cliffs, New Jersey.
- Courtenay, W. R., Jr., H. F. Sahlman, W. W. Miley, II, and D. J. Herrema. 1974. Exotic fishes in fresh and brackish waters of Florida. *Biological Conservation* 6(4):292–302.
- Dawes, J. 2001. Complete encyclopedia of the freshwater aquarium. Firefly Books, Buffalo, New York.
- Devick, W. S. 1991. Pattern of introductions of aquatic organisms to Hawaii freshwater habitats. *In* New directions in research, management and conservations of Hawaiian freshwater stream ecosystems. Proceedings of 1990 symposium on freshwater stream biology and fisheries management, Hawaii Department of Land and Natural Resources, Division of Aquatic Resources, Honolulu, Hawaii.
- FAO. 1997. FAO database on introduced aquatic species. FAO Database on Introduced Aquatic Species, FAO, Rome.
- Hoedman, J. J. 1974. Naturalist's guide to freshwater aquarium fish. M. Banister (translator). Sterling Publishing, New York.
- Holton, G. D. 1990. A field guide to Montana fishes. Montana Department of Fish, Wildlife and Parks, Helena.
- Hubbs, C. L., W. I. Follett, and L. J. Dempster. 1979. List of the fishes of California. California Academy Science, Occasional Papers 133.
- Hugg, D. O. 1996. MAPFISH georeferenced mapping database. Freshwater and estuarine fishes of North America. Life Science Software, Edgewater, Maryland.
- Maceda-Veiga, A., de Sostoa, A. Escribano-Alacid, and E. García-Berthou. 2013. The aquarium trade as a potential source of fish introductions in southwestern Europe. *Biological Invasions* 15:2707–2716.
- Maciolek, J. A. 1984. Exotic fishes in Hawaii and other islands of Oceania. Pages 131–161 in W. R. Courtenay, Jr., and J. R. Stauffer, Jr., editors. Distribution, biology, and management of exotic fishes. The Johns Hopkins University Press, Baltimore, Maryland.

- Magalhães, A. L. B., L. Casatti, and J. R. S. Vitule. 2011. Alterações no Código Florestal Brasileiro favorecerão espécies não-nativas de peixes de água doce. *Nat Conserv.* 9(1):121–24.
- Meek, S. E. 1904. The fresh-water fishes of Mexico north of the Isthmus of Tehuantepec. *Field Columbian Museum, Zoological Series* 5:1–17.
- Miller, R. R., W. L. Minckley, and S. M. Norris. 2005. *Freshwater fishes of Mexico*, 1st edition. University of Chicago Press, Chicago, Illinois.
- Mills, D., and G. Vevers. 1989. *The Tetra encyclopedia of freshwater tropical aquarium fishes*. Tetra Press, New Jersey.
- Minckley, W. L. 1973. *Fishes of Arizona*. Arizona Fish and Game Department. Sims Printing, Phoenix, Arizona.
- Moyle, P. B. 1976. *Inland fishes of California*. University of California Press, Berkeley.
- Mundy, B. C. 2005. Fishes of the Hawaiian Archipelago. *Bishop Museum Bulletins in Zoology* 6.
- Olden, J. D., J. L. Lockwood, and C. L. Parr. 2011. Biological invasions and the homogenization of faunas and floras. Pages 224–243 *in* R. J. Ladle, and R. J. Whittaker, editors. *Conservation biogeography*. Blackwell Publishing, Oxford, UK.
- Page, L. M., and B. M. Burr. 1991. *A field guide to freshwater fishes of North America north of Mexico*. Houghton Mifflin, Boston.
- Robins, C. R., R. M. Bailey, C. E. Bond, J. R. Brooker, E. A. Lachner, R. N. Lea, and W. B. Scott. 1991. *World fishes important to North Americans. Exclusive of species from the continental waters of the United States and Canada*. American Fisheries Society, Special Publication 21, Bethesda, Maryland.
- Shafland, P. L. 1979. Non-native fish introductions with special reference to Florida. *Fisheries* 4(3):18–23.
- Shapovalov, L., A. J. Cordone, and W. A. Dill. 1981. A list of freshwater and anadromous fishes of California. *California Fish and Game* 67(1):4–38.
- St. Amant, J. A., and F. G. Hoover. 1969. Addition of *Misgurnus anguillicaudatus* (Cantor) to the California fauna. *California Fish and Game* 57(2):330–331.
- St. Amant, J. A., and I. Sharp. 1971. Addition of *Xiphophorus variatus* (Meek) to the California fauna. *California Fish and Game* 57:128–129.

- Swift, C., T. R. Haglund, M. Ruiz, and R. N. Fisher. 1993. The status and distribution of the freshwater fishes of southern California. *Bulletin of the Southern California Academy of Science* 92(3):101–167.
- Wischnath, L. 1993. *Atlas of livebearers of the world*. TFH Publication, Neptune City, New Jersey.
- Yamamoto, M. N., and A. W. Tagawa. 2000. *Hawaii's native and exotic freshwater animals*. Mutual Publishing, Honolulu, Hawaii.
- Zuckerman, L. D., and R. J. Behnke. 1986. Introduced fishes in the San Luis Valley, Colorado. Pages 435–452 *in* R. H. Stroud, editor. *Fish culture in fisheries management. Proceedings of a symposium on the role of fish culture in fisheries management at Lake Ozark, Missouri, March 31–April 3, 1985*. American Fisheries Society, Bethesda, Maryland.