

# FEDERAL AND STATE ENDANGERED AND THREATENED SPECIES EXPENDITURES



*Fiscal Year 2018*

## TABLE OF CONTENTS

<b>ENDANGERED AND THREATENED SPECIES EXPENDITURES FISCAL YEAR 2018</b>	
PURPOSE .....	3
BACKGROUND .....	4
SECTION 18(1), FEDERAL EXPENDITURES .....	5
SECTION 18(2), STATE EXPENDITURES .....	5
SUMMARY OF DATA.....	5
COMPARISON TO PREVIOUS REPORTS.....	6
<b>Table 1.</b> FY 2018 Reported Expenditures for Endangered and Threatened Species.....	7
<b>Table 2.</b> Species Ranked in Descending Order of Total FY 2018 Reported Expenditures.....	105
<b>Table 3.</b> FY 2018 Reported Land Acquisition Expenditures for Endangered and Threatened Species.....	165
<b>Table 4.</b> FY 2018 Expenditures Reported by Individual Federal Agencies and the States.....	176
<b>Appendix A.</b> Supplemental Guidance for Reporting Expenditures for the Conservation of Endangered and Threatened Species.....	179
<b>Appendix B.</b> Reports Provided by Other Federal Agencies.....	184
<b>Appendix C.</b> Expenditures Reported by the States.....	343

## **ENDANGERED SPECIES EXPENDITURES FISCAL YEAR 2018**

### **PURPOSE**

The Endangered Species Act of 1973 (16 U.S.C. 1531 *et seq.*) (Act) was reauthorized and amended on October 7, 1988, under Public Law 100-478. A new section 18 was added to the Act that requires the U.S. Fish and Wildlife Service (Service) on behalf of the Secretary of the Interior to report annually on certain expenditures on a species-by-species basis. The section reads:

SEC. 18. On or before January 15, 1990, and each January 15 thereafter, the Secretary of the Interior, acting through the Fish and Wildlife Service, shall submit to the Congress an annual report covering the preceding fiscal year which shall contain

(1) the accounting on a species-by-species basis of all reasonably identifiable Federal expenditures made primarily for the conservation of endangered or threatened species pursuant to this Act; and

(2) an accounting on a species-by-species basis of all reasonably identifiable expenditures made primarily for the conservation of endangered or threatened species pursuant to this Act by States receiving grants under section 6.

This document constitutes the 30th report by the Service pursuant to the requirements of section 18 of the Act. The report covers fiscal year 2018 (October 1, 2017 - September 30, 2018), although some States may be reporting for a different fiscal year period (without duplicating a previous year's reported data). To assist Federal and State agencies with more accurate accounting of expenditures, a multi-species reporting category was added to the FY12 and subsequent reports. This option gives agencies another way to report expenditures when it is not possible to determine individual species costs for those species that are part of a multi-species conservation activity.

The term "State" as used in this report is defined in section 3(17) of the Act to mean any of the several States, the District of Columbia, the Commonwealth of Puerto Rico, American Samoa, the Virgin Islands, Guam, and the Trust Territory of the Pacific Islands.

The term "State agency" as used in this report is defined in section 3(18) of the Act to mean any State agency, department, board, commission, or other governmental entity which is responsible for the management and conservation of fish, plant, or wildlife resources within a State.

The term "conservation," as defined in section 3(3) of the Act, means the use of all methods and procedures necessary to bring any endangered or threatened species to the point where the measures provided for by the Act are no longer necessary--including, but not limited to, research, census, law enforcement, habitat acquisition and maintenance, propagation, live trapping, and transplantation. For the purposes of this report, the definition of conservation is further expanded to include any and all actions taken by Federal and State agencies on behalf of threatened or endangered species listed pursuant to the Act.

## BACKGROUND

At the time the reporting requirement was enacted by Congress, the Service determined that a good faith effort was expected to develop methods to track and obtain data that are "reasonably identifiable." No extensive or extraordinary measures to develop exceptionally precise data were sought, nor were the reporting agencies expected to prorate staff salaries and other normal operational and maintenance costs not directed toward a particular species. The expenditures reported in this and the preceding reports do not necessarily constitute all expenditures made for each species.

This report represents a compilation of reasonably identifiable expenditures for the conservation of listed species reported independently by Federal and State agencies. Expenditures for FY 2018 are reported for 1,719 domestic threatened and endangered species of the 1,912 domestic species that were listed under United States jurisdiction in 50 CFR Part 17 during FY 2018 up through September 30, 2018 (Tables 1 and 3).

Expenditures are identified by listed entity, which may be a species<sup>1</sup>, subspecies<sup>2</sup>, Distinct Population Segment (DPS)<sup>3</sup>, or Evolutionarily Significant Unit (ESU)<sup>4</sup>, as identified in the Act's implementing regulations at 50 CFR 17.11 and 17.12.

These expenditures are reported separately for the Service and in aggregate for all other Federal agencies and the States (Tables 1 and 3).

For these annual reports, the Service requests that all reporting agencies submit their data directly into our Web-based data entry system to ensure accuracy, consistency, and efficiency. At the time of submission, the reporting agencies certify that their information is accurate. The information provided in this report, pursuant to section 18 of the Endangered Species Act, is presented as it was submitted by these agencies. We do not verify an agency's data after the data are submitted to us. In order to improve the Service's efficiency in producing future editions of this report, the Service will no longer enter a reporting agency's expenditures data into our Web-based system. All agencies must enter their own expenditures data into our Web-based system.

Expenditures for candidate species or other species not listed in 50 CFR Part 17 as of the end of each fiscal year are not included in this report. In addition, other types of expenses that are not incurred primarily for the conservation of listed species, such as those for litigation, are not reported.

---

<sup>1</sup> **Species** For purposes of the ESA, this term includes any species or subspecies of fish or wildlife or plants, and any Distinct Population Segment of any species of vertebrate fish or wildlife that interbreeds when mature.

<sup>2</sup> **Subspecies** A taxonomic rank below that of species, usually recognizing individuals that have certain heritable characteristics distinct from other subspecies of a species.

<sup>3</sup> **Distinct Population Segment** A subdivision of a vertebrate species that is treated as a species for purposes of listing under the Endangered Species Act.

<sup>4</sup> **Evolutionarily Significant Unit** A Pacific salmonid stock that is substantially reproductively isolated from other stocks of the same species and which represents an important part of the evolutionary legacy of the species; functionally equivalent to DPS for the purpose of listing under ESA.

## **SECTION 18(1), FEDERAL EXPENDITURES**

### **Expenditures Reported by the Fish and Wildlife Service**

The Service's operational categories of expenditures include fisheries, refuges, land acquisition, law enforcement, research, international conservation, and regional and field office operations for listing, recovery, consultation, migratory birds, and grants. All reasonably identifiable expenditures that could be attributed to individual species within these activities have been included.

The Federal assistance portion under section 6 of the Act for each species is included in the FWS Total in Table 1. The State share for each species for which a Federal contribution was made is included in the States Total column of Table 1.

A portion of the Service's funds for consultation, listing, and recovery are incorporated into salaries, operations, and maintenance costs. If these costs contributed to the conservation of listed species and are reasonably identifiable to specific species, they are included in this report.

### **Expenditures Reported by other Federal Agencies**

The Director of the Service sends a request to other Federal agencies identified as likely to have species-specific expenditures under the Act to report for each fiscal year. The Service provides guidance identifying what constitutes reasonably identifiable expenditures and how these are to be reported (Appendix A). For FY 2018, the Service and 22 other Federal agencies identified expenditures for the conservation of individual threatened and endangered species. Results from all responding Federal agencies are included in Appendix B. The data identified by those responding Federal agencies for individual species and multi-species conservation activities are tabulated in the "Other Fed" columns of this report (Tables 1 and 3).

## **SECTION 18(2), STATE EXPENDITURES**

The assistance of the Association of Fish and Wildlife Agencies is sought by the Service to help coordinate the reporting of reasonably identifiable expenditures made by the States. States responding to this data call do so by reporting their data by species directly to the Service via our Web-based system. Collective results from all responding State agencies are included in Appendix C. For FY 2018, 32 State agencies identified expenditures for the conservation of individual threatened and endangered species.

## **SUMMARY OF DATA**

**Table 1** reports the activity-related expenditures that are reasonably identifiable to specific individual species as well as multi-species activity costs by the Service, other Federal agencies collectively, and the States collectively. Expenditures for land acquisition (Table 3) are not included in this tabulation. Species are grouped in the same general order as they appear in 50 CFR Part 17. Table 1 also provides the rank for each species from Table 2.

**Table 2** ranks each species in descending order of the species total from Table 1. A cumulative total of the activity-related, species-specific expenditures is also provided.

**Table 3** displays land acquisition expenditures, reasonably identifiable to specific individual species, by the Service, other Federal agencies collectively, and the States collectively. Expenditures for land purchased for multi-species conservation activities are listed at the end of the table. Table 3 also provides the rank for land expenditures by species.

**Table 4** shows the combined total expenditures for species and land acquisition by reporting Federal agency, the Service, and the States collectively. Expenditures for foreign species and multi-species conservation activities are included.

**Total expenditures** reported for domestic and foreign species in **FY 2018** were **\$ 1,519,328,304**, of which **\$ 1,387,579,479** was reported by Federal agencies and **\$ 131,748,825** was reported by the States (Table 4).

## **COMPARISON TO PREVIOUS REPORTS**

Over the years, these reports have been refined as our experience in planning and coordinating with the reporting agencies to track such expenditures has improved. A good faith effort is made to develop species-by-species expenditures for these reports, but due to differences in reporting methods, this report cannot be easily compared to prior expenditures reports. These year-to-year differences include changes in the number of Federal and State agencies reporting data, changes in how expenditures are calculated, changes in the agencies' abilities to track expenditures, and changes in the number of listed species. Therefore, real dollar increases or decreases cannot accurately be calculated or inferred from the overall or individual totals.

## **TABLE 1. FY 2018 REPORTED EXPENDITURES FOR ENDANGERED AND THREATENED SPECIES**

The list of species in Table 1 essentially follows the taxonomic and nomenclatural conventions used in 50 CFR Part 17 for the identification and sequence of species' names. For ease of reference, the alphabetized, inverted common name is provided within each group (usually Class) of animals and plants. Thus, the gray bat is found with other bats under "Bat, gray" within mammals, and Tar River spiny mussel is found under "Spiny mussel, Tar River" within the clams. Groups of animals follow the sequence: Mammals, Birds, Reptiles, Amphibians, Fishes, and Invertebrates, which include Clams, Snails, Insects, Arachnids, and Crustaceans. Groups of plants follow the sequence: Flowering Plants, Conifers and Cycads, Ferns and Allies, and Lichens.

For additional information about the names, historical range, or status of an endangered or threatened species, the species can be found in the Code of Federal Regulations at 50 CFR 17.11 (list of wildlife) or 17.12 (list of plants), or through our Web site at <http://www.fws.gov/endangered>. The STATUS column indicates whether the species is endangered (E), threatened (T), experimental (EXPN), or threatened by similarity of appearance (SAT).

Fish and Wildlife Service expenditures (FWS Total) and other Federal agency expenditures (Other Fed) are totaled in the "Federal Total" column. The States Total is the amount reported to the Service directly by the States. None of these reported totals includes costs for land acquisition. (See Table 3 for reported land acquisition expenditures.)

The RANK is from Table 2 and indicates each species' rank by identifiable expenditures, where #1 is the highest expenditure.

Multi-species conservation activity-related expenditures are tabulated at the end of Table 1.

**TABLE 1. REPORTED EXPENDITURES FOR FY 2018 FOR ENDANGERED AND THREATENED SPECIES,  
NOT INCLUDING LAND ACQUISITION COSTS**

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
Mammals	158	Bat, Florida bonneted ( <i>Eumops floridanus</i> ) - Wherever found	E	\$424,000	\$366,607	\$790,607	\$0	\$790,607
	143	Bat, gray ( <i>Myotis grisescens</i> ) - Wherever found	E	\$397,924	\$522,119	\$920,043	\$25,800	\$945,843
	93	Bat, Hawaiian hoary ( <i>Lasiurus cinereus semotus</i> ) - Wherever found	E	\$616,841	\$1,121,167	\$1,738,008	\$0	\$1,738,008
	50	Bat, Indiana ( <i>Myotis sodalis</i> ) - Wherever found	E	\$1,820,373	\$3,471,152	\$5,291,525	\$191,727	\$5,483,252
	606	Bat, little Mariana fruit ( <i>Pteropus tokudae</i> ) - Wherever found	E	\$42,842	\$0	\$42,842	\$0	\$42,842
	398	Bat, Mariana fruit (=Mariana flying fox) ( <i>Pteropus mariannus mariannus</i> ) - Wherever found	T	\$59,402	\$65,302	\$124,704	\$0	\$124,704
	626	Bat, Mexican long-nosed ( <i>Leptonycteris nivalis</i> ) - Wherever found	E	\$5,000	\$21,300	\$26,300	\$13,576	\$39,876
	43	Bat, Northern long-eared ( <i>Myotis septentrionalis</i> ) - Wherever found	T	\$1,538,481	\$4,934,333	\$6,472,814	\$361,930	\$6,834,744
	431	Bat, Ozark big-eared ( <i>Corynorhinus (=Plecotus) townsendii ingens</i> ) - Wherever found	E	\$74,204	\$20,347	\$94,551	\$4,000	\$98,551
	636	Bat, Pacific sheath-tailed ( <i>Emballonura semicaudata rotensis</i> ) - Wherever found	E	\$38,684	\$0	\$38,684	\$0	\$38,684
	897	Bat, Pacific sheath-tailed ( <i>Emballonura semicaudata semicaudata</i> ) - American Samoa	E	\$14,500	\$0	\$14,500	\$0	\$14,500
	274	Bat, Virginia big-eared ( <i>Corynorhinus (=Plecotus) townsendii virginianus</i> ) - Wherever found	E	\$132,285	\$116,603	\$248,888	\$37,182	\$286,070
	1181	Bear, brown ( <i>Ursus arctos pruinosus</i> ) - Wherever found	E	\$0	\$4,073	\$4,073	\$0	\$4,073
	33	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - U.S.A., conterminous (lower 48) States, except where listed as an experimental population	T	\$2,394,362	\$3,760,428	\$6,154,790	\$3,099,083	\$9,253,873



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	170	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - Bitterroot Grizzly Bear Nonessential Experimental Population Area	EXPN	\$590,096	\$40,000	\$630,096	\$59,500	\$689,596
	808	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - Cabinet-Yaak Recovery Zone Population	UR	\$0	\$0	\$0	\$19,500	\$19,500
	587	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - Greater Yellowstone Ecosystem DPS	UR	\$0	\$0	\$0	\$47,000	\$47,000
	75	Bear, polar ( <i>Ursus maritimus</i> ) - Wherever found	T	\$872,000	\$1,827,190	\$2,699,190	\$0	\$2,699,190
	292	Bison, wood ( <i>Bison bison athabasca</i> ) - Wherever found	T	\$186,000	\$0	\$186,000	\$62,017	\$248,017
	300	Caribou DPS, Southern Mountain ( <i>Rangifer tarandus ssp. caribou</i> ) - Southern Mountain DPS	E	\$219,132	\$3,900	\$223,032	\$10,000	\$233,032
	490	Chimpanzee ( <i>Pan troglodytes</i> ) - Wherever found	E	\$74,339	\$0	\$74,339	\$0	\$74,339
	232	Deer, Columbian white-tailed ( <i>Odocoileus virginianus leucurus</i> ) - Columbia River DPS	T	\$424,023	\$146	\$424,169	\$10,000	\$434,169
	394	Deer, key ( <i>Odocoileus virginianus clavium</i> ) - Wherever found	E	\$121,000	\$5,231	\$126,231	\$0	\$126,231
	348	Dolphin, Maui ( <i>Cephalorhynchus hectori maui</i> )	E	\$0	\$158,636	\$158,636	\$0	\$158,636
	72	Elephant, African ( <i>Loxodonta africana</i> ) - Wherever found	T	\$3,083,083	\$0	\$3,083,083	\$0	\$3,083,083
	112	Elephant, Asian ( <i>Elephas maximus</i> ) - Wherever found	E	\$1,296,246	\$0	\$1,296,246	\$0	\$1,296,246
	116	Ferret, black-footed ( <i>Mustela nigripes</i> ) - Wherever found, except where listed as an experimental population	E	\$407,929	\$775,500	\$1,183,429	\$0	\$1,183,429
	142	Ferret, black-footed ( <i>Mustela nigripes</i> ) - U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	EXPN	\$18,500	\$534,876	\$553,376	\$396,627	\$950,003
	119	Fox, San Joaquin kit ( <i>Vulpes macrotis mutica</i> ) - wherever found	E	\$798,000	\$358,730	\$1,156,730	\$0	\$1,156,730
	1197	Fox, Santa Catalina Island ( <i>Urocyon littoralis catalinae</i> ) - Wherever found	T	\$3,350	\$0	\$3,350	\$0	\$3,350
	154	Gibbons ( <i>Hylobates</i> spp. (including <i>Nomascus</i> )) - Wherever found	E	\$843,638	\$0	\$843,638	\$0	\$843,638
	260	Gorilla ( <i>Gorilla gorilla</i> ) - Wherever found	E	\$316,116	\$3,150	\$319,266	\$0	\$319,266

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	263	Jaguar ( <i>Panthera onca</i> ) - Wherever found	E	\$301,182	\$16,300	\$317,482	\$0	\$317,482
	494	Jaguarundi, Gulf Coast ( <i>Herpailurus (=Felis) yagouaroundi cacomitli</i> ) - Wherever found	E	\$72,300	\$210	\$72,510	\$0	\$72,510
	616	Kangaroo rat, Fresno ( <i>Dipodomys nitratoides exilis</i> ) - Wherever found	E	\$27,000	\$14,438	\$41,438	\$0	\$41,438
	320	Kangaroo rat, giant ( <i>Dipodomys ingens</i> ) - Wherever found	E	\$124,000	\$74,458	\$198,458	\$0	\$198,458
	649	Kangaroo rat, Morro Bay ( <i>Dipodomys heermanni morroensis</i> ) - Wherever found	E	\$36,000	\$0	\$36,000	\$0	\$36,000
	221	Kangaroo rat, San Bernardino Merriam's ( <i>Dipodomys merriami parvus</i> ) - Wherever found	E	\$406,000	\$67,138	\$473,138	\$0	\$473,138
	184	Kangaroo rat, Stephens' ( <i>Dipodomys stephensi</i> (incl. <i>D. cascus</i> )) - Wherever found	E	\$57,000	\$539,888	\$596,888	\$0	\$596,888
	345	Kangaroo rat, Tipton ( <i>Dipodomys nitratoides nitratoides</i> ) - Wherever found	E	\$103,000	\$57,300	\$160,300	\$0	\$160,300
	1170	Leopard ( <i>Panthera pardus</i> ) - except where threatened	E	\$4,620	\$0	\$4,620	\$0	\$4,620
	436	Lion ( <i>Panthera leo</i> ssp. <i>leo</i> ) - Wherever found	E	\$97,076	\$0	\$97,076	\$0	\$97,076
	436	Lion ( <i>Panthera leo</i> ssp. <i>melanochaita</i> ) - Wherever found	T	\$97,076	\$0	\$97,076	\$0	\$97,076
	70	Lynx, Canada ( <i>Lynx canadensis</i> ) - Wherever Found in Contiguous U.S.	T	\$771,967	\$1,226,188	\$1,998,155	\$1,240,002	\$3,238,157
	36	Manatee, West Indian ( <i>Trichechus manatus</i> ) - Wherever found	T	\$2,171,500	\$6,479,480	\$8,650,980	\$30,100	\$8,681,080
	150	Markhor, chiltan (=wild goat) ( <i>Capra falconeri (=aegrargus) chiltanensis</i> ) - Wherever found	E	\$889,996	\$0	\$889,996	\$0	\$889,996
	516	Mountain beaver, Point Arena ( <i>Aplodontia rufa nigra</i> ) - Wherever found	E	\$59,000	\$5,500	\$64,500	\$0	\$64,500
	420	Mouse, Alabama beach ( <i>Peromyscus polionotus ammobates</i> ) - Wherever found	E	\$96,500	\$6,094	\$102,594	\$0	\$102,594
	748	Mouse, Anastasia Island beach ( <i>Peromyscus polionotus phasma</i> ) - Wherever found	E	\$21,000	\$3,220	\$24,220	\$0	\$24,220

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	470	Mouse, Choctawhatchee beach ( <i>Peromyscus polionotus allophrys</i> ) - Wherever found	E	\$37,500	\$45,781	\$83,281	\$0	\$83,281
	547	Mouse, Key Largo cotton ( <i>Peromyscus gossypinus allapaticola</i> ) - Wherever found	E	\$55,500	\$1,561	\$57,061	\$0	\$57,061
	254	Mouse, New Mexico meadow jumping ( <i>Zapus hudsonius luteus</i> ) - Wherever found	E	\$103,865	\$241,341	\$345,206	\$0	\$345,206
	230	Mouse, Pacific pocket ( <i>Perognathus longimembris pacificus</i> ) - Wherever found	E	\$74,000	\$362,400	\$436,400	\$0	\$436,400
	399	Mouse, Perdido Key beach ( <i>Peromyscus polionotus trissyllepsis</i> ) - Wherever found	E	\$78,000	\$46,134	\$124,134	\$0	\$124,134
	370	Mouse, Preble's meadow jumping ( <i>Zapus hudsonius preblei</i> ) - wherever found	T	\$57,400	\$65,439	\$122,839	\$15,745	\$138,584
	135	Mouse, salt marsh harvest ( <i>Reithrodontomys raviventris</i> ) - wherever found	E	\$723,000	\$266,500	\$989,500	\$0	\$989,500
	334	Mouse, southeastern beach ( <i>Peromyscus polionotus niveiventris</i> ) - wherever found	T	\$135,500	\$35,820	\$171,320	\$0	\$171,320
	553	Mouse, St. Andrew beach ( <i>Peromyscus polionotus peninsularis</i> ) - Wherever found	E	\$18,000	\$37,660	\$55,660	\$0	\$55,660
	120	Ocelot ( <i>Leopardus (=Felis) pardalis</i> ) - wherever found	E	\$1,066,400	\$24,910	\$1,091,310	\$0	\$1,091,310
	918	Orangutan ( <i>Pongo pygmaeus</i> ) - Wherever found	E	\$13,585	\$0	\$13,585	\$0	\$13,585
	237	Otter, Northern Sea ( <i>Enhydra lutris kenyoni</i> ) - Southwest Alaska DPS	T	\$273,810	\$138,751	\$412,561	\$4,000	\$416,561
	231	Otter, southern sea ( <i>Enhydra lutris nereis</i> ) - Wherever found	T	\$0	\$434,499	\$434,499	\$0	\$434,499
	739	Otter, southern sea ( <i>Enhydra lutris nereis</i> ) - Wherever found	NS	\$25,000	\$0	\$25,000	\$0	\$25,000
	321	Pangolin, Temnick's ground ( <i>Manis temminckii</i> ) - Wherever found	E	\$193,532	\$0	\$193,532	\$0	\$193,532
	83	Panther, Florida ( <i>Puma (=Felis) concolor coryi</i> ) - Wherever found	E	\$1,154,500	\$1,033,376	\$2,187,876	\$0	\$2,187,876
	216	Pocket gopher, Olympia ( <i>Thomomys mazama pugetensis</i> ) - Wherever found	T	\$452,868	\$2,000	\$454,868	\$35,000	\$489,868
	270	Pocket gopher, Roy Prairie ( <i>Thomomys mazama glacialis</i> ) - Wherever found	T	\$261,401	\$0	\$261,401	\$35,000	\$296,401

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	215	Pocket gopher, Tenino ( <i>Thomomys mazama tumuli</i> ) - Wherever found	T	\$324,991	\$130,000	\$454,991	\$35,000	\$489,991
	220	Pocket gopher, Yelm ( <i>Thomomys mazama yelmensis</i> ) - Wherever found	T	\$325,801	\$75,000	\$400,801	\$75,000	\$475,801
	168	Prairie dog, Utah ( <i>Cynomys parvidens</i> ) - Wherever found	T	\$0	\$702,600	\$702,600	\$0	\$702,600
	972	Pronghorn, Sonoran ( <i>Antilocapra americana sonoriensis</i> ) - Wherever found, except where listed as an experimental population	E	\$0	\$11,430	\$11,430	\$0	\$11,430
	185	Pronghorn, Sonoran ( <i>Antilocapra americana sonoriensis</i> ) - U.S.A. (AZ), Mexico	EXPN	\$412,000	\$184,780	\$596,780	\$0	\$596,780
	229	Puma (=mountain lion) ( <i>Puma (=Felis) concolor</i> (all subsp. except <i>coryi</i> )) - FL	SAT	\$200	\$437,370	\$437,570	\$0	\$437,570
	191	Rabbit, Columbia Basin Pygmy ( <i>Brachylagus idahoensis</i> ) - Columbia Basin DPS	E	\$331,520	\$18,600	\$350,120	\$229,600	\$579,720
	433	Rabbit, Lower Keys marsh ( <i>Sylvilagus palustris hefneri</i> ) - Wherever found	E	\$70,500	\$27,293	\$97,793	\$0	\$97,793
	534	Rabbit, riparian brush ( <i>Sylvilagus bachmani riparius</i> ) - Wherever found	E	\$53,000	\$8,300	\$61,300	\$0	\$61,300
	138	Rhinoceros, black ( <i>Diceros bicornis</i> ) - Africa	E	\$969,295	\$0	\$969,295	\$0	\$969,295
	423	Rhinoceros, great Indian ( <i>Rhinoceros unicornis</i> ) - Nepal, India	E	\$100,672	\$0	\$100,672	\$0	\$100,672
	303	Rhinoceros, Javan ( <i>Rhinoceros sondaicus</i> ) - Sikkim, Thailand, Malaysia, Indonesia, Indochina, Burma, Bangladesh	E	\$225,852	\$0	\$225,852	\$0	\$225,852
	362	Rhinoceros, Southern white ( <i>Ceratotherium simum ssp. simum</i> ) - Wherever found	SAT	\$145,451	\$0	\$145,451	\$0	\$145,451
	211	Rhinoceros, Sumatran ( <i>Dicerorhinus sumatrensis</i> ) - Wherever found	E	\$505,326	\$0	\$505,326	\$0	\$505,326
	758	Rice rat, Silver ( <i>Oryzomys palustris natator</i> ) - lower FL Keys	E	\$18,000	\$5,303	\$23,303	\$0	\$23,303
	94	Seal, bearded ( <i>Erignathus barbatus nauticus</i> ) - Beringia DPS	T	\$0	\$1,705,000	\$1,705,000	\$0	\$1,705,000
	249	Seal, Guadalupe fur ( <i>Arctocephalus townsendi</i> ) - Wherever found	T	\$0	\$366,042	\$366,042	\$0	\$366,042

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	58	Seal, Hawaiian monk ( <i>Monachus schauinslandi</i> ) - Wherever found	E	\$92,020	\$4,181,114	\$4,273,134	\$0	\$4,273,134
	49	Sea lion, Steller ( <i>Eumetopias jubatus</i> ) - Western DPS	E	\$33,000	\$5,531,634	\$5,564,634	\$125,520	\$5,690,154
	1182	Seal, Ringed ( <i>Phoca (=Pusa) hispida botnica</i> ) - Wherever found	T	\$0	\$4,000	\$4,000	\$0	\$4,000
	955	Seal, Ringed ( <i>Phoca (=Pusa) hispida hispida</i> ) - Wherever found	T	\$0	\$11,998	\$11,998	\$0	\$11,998
	330	Sheep, Peninsular bighorn ( <i>Ovis canadensis nelsoni</i> ) - Peninsular CA pop.	E	\$116,000	\$64,500	\$180,500	\$0	\$180,500
	236	Sheep, Sierra Nevada bighorn ( <i>Ovis canadensis sierrae</i> ) - Sierra Nevada	E	\$150,000	\$267,980	\$417,980	\$0	\$417,980
	456	Shrew, Buena Vista Lake ornate ( <i>Sorex ornatus relictus</i> ) - Wherever found	E	\$73,000	\$15,390	\$88,390	\$0	\$88,390
	572	Squirrel, Carolina northern flying ( <i>Glaucomys sabrinus coloratus</i> ) - Wherever found	E	\$16,300	\$25,824	\$42,124	\$7,500	\$49,624
	747	Squirrel, Mount Graham red ( <i>Tamiasciurus hudsonicus grahamensis</i> ) - Wherever found	E	\$24,500	\$0	\$24,500	\$0	\$24,500
	266	Squirrel, Northern Idaho Ground ( <i>Urocitellus brunneus</i> ) - Wherever found	T	\$282,892	\$3,100	\$285,992	\$24,000	\$309,992
	106	Tiger ( <i>Panthera tigris</i> ) - Wherever found	E	\$1,409,159	\$0	\$1,409,159	\$0	\$1,409,159
	207	Vole, Amargosa ( <i>Microtus californicus scirpensis</i> ) - Wherever found	E	\$396,800	\$117,000	\$513,800	\$0	\$513,800
	583	Vole, Florida salt marsh ( <i>Microtus pennsylvanicus dukecampbelli</i> ) - Wherever found	E	\$47,500	\$0	\$47,500	\$0	\$47,500
	69	Whale, beluga ( <i>Delphinapterus leucas</i> ) - Cook Inlet DPS	E	\$0	\$3,335,312	\$3,335,312	\$194,968	\$3,530,280
	91	Whale, blue ( <i>Balaenoptera musculus</i> ) - Wherever found	E	\$0	\$1,898,201	\$1,898,201	\$0	\$1,898,201
	132	Whale, bowhead ( <i>Balaena mysticetus</i> ) - Wherever found	E	\$0	\$743,134	\$743,134	\$261,737	\$1,004,871
	155	Whale, false killer ( <i>Pseudorca crassidens</i> ) - Main Hawaiian Islands Insular DPS	E	\$122,078	\$700,065	\$822,143	\$0	\$822,143
	84	Whale, finback ( <i>Balaenoptera physalus</i> ) - Wherever found	E	\$0	\$2,167,630	\$2,167,630	\$0	\$2,167,630

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	336	Whale, gray ( <i>Eschrichtius robustus</i> ) - Western North Pacific DPS	E	\$0	\$165,135	\$165,135	\$3,000	\$168,135
	151	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Western North Pacific	E	\$0	\$871,481	\$871,481	\$3,000	\$874,481
	108	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Mexico DPS	T	\$0	\$1,385,855	\$1,385,855	\$0	\$1,385,855
	228	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Central America DPS	E	\$0	\$440,142	\$440,142	\$0	\$440,142
	87	Whale, killer ( <i>Orcinus orca</i> ) - Southern Resident DPS	E	\$0	\$1,689,236	\$1,689,236	\$388,600	\$2,077,836
	30	Whale, North Atlantic Right ( <i>Eubalaena glacialis</i> ) - Wherever found	E	\$0	\$9,817,274	\$9,817,274	\$125,000	\$9,942,274
	186	Whale, North Pacific Right ( <i>Eubalaena japonica</i> ) - Wherever found	E	\$0	\$595,341	\$595,341	\$0	\$595,341
	98	Whale, Sei ( <i>Balaenoptera borealis</i> ) - Wherever found	E	\$0	\$1,613,657	\$1,613,657	\$0	\$1,613,657
	1259	Whale, Southern Right ( <i>Eubalaena australis</i> ) - Wherever found	E	\$0	\$476	\$476	\$0	\$476
	88	Whale, sperm ( <i>Physeter catodon</i> (=macrocephalus)) - Wherever found	E	\$310	\$2,031,421	\$2,031,731	\$0	\$2,031,731
	57	Wolf, gray ( <i>Canis lupus</i> ) - U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	E	\$974,620	\$1,522,863	\$2,497,483	\$2,116,640	\$4,614,123
	107	Wolf, gray ( <i>Canis lupus</i> ) - MN	T	\$0	\$1,239,292	\$1,239,292	\$154,900	\$1,394,192
	1003	Wolf, gray ( <i>Canis lupus</i> ) - Western Great Lakes DPS	RT	\$10,000	\$0	\$10,000	\$0	\$10,000
	622	Wolf, Mexican ( <i>Canis lupus baileyi</i> ) - Wherever found, except where listed as an experimental population	E	\$0	\$40,230	\$40,230	\$0	\$40,230
	73	Wolf, Mexican ( <i>Canis lupus baileyi</i> ) - U.S.A. (portions of AZ and NM)see 17.84(k)	EXPN	\$2,463,677	\$355,600	\$2,819,277	\$0	\$2,819,277
	1253	Wolf, red ( <i>Canis rufus</i> ) - Wherever found, except where listed as an experimental population	E	\$0	\$0	\$0	\$1,000	\$1,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	164	Wolf, red ( <i>Canis rufus</i> ) - U.S.A. (portions of NC and TN)	EXPN	\$728,400	\$0	\$728,400	\$0	\$728,400
	503	Woodrat, Key Largo ( <i>Neotoma floridana smalli</i> ) - Wherever found	E	\$68,500	\$1,048	\$69,548	\$0	\$69,548
	850	Woodrat, riparian (=San Joaquin Valley) ( <i>Neotoma fuscipes riparia</i> ) - Wherever found	E	\$8,700	\$8,300	\$17,000	\$0	\$17,000
<b>Mammals subtotal</b>				<b>\$37,371,492</b>	<b>\$73,853,127</b>	<b>\$111,224,619</b>	<b>\$9,443,254</b>	<b>\$120,667,873</b>
Birds	299	Akekee ( <i>Loxops caeruleirostris</i> ) - Wherever found	E	\$158,581	\$74,489	\$233,070	\$0	\$233,070
	337	Akepa, Hawaii ( <i>Loxops coccineus</i> ) - Wherever found	E	\$142,869	\$22,610	\$165,479	\$0	\$165,479
	313	Akepa, Maui ( <i>Loxops ochraceus</i> ) - Wherever found	E	\$24,696	\$186,400	\$211,096	\$0	\$211,096
	815	Akialoa, Kauai (honeycreeper) ( <i>Akialoa stejnegeri</i> ) - Wherever found	E	\$19,012	\$0	\$19,012	\$0	\$19,012
	396	Akiapolaau ( <i>Hemignathus wilsoni</i> ) - Wherever found	E	\$21,610	\$104,009	\$125,619	\$0	\$125,619
	393	Akikiki ( <i>Oreomystis bairdi</i> ) - Wherever found	E	\$42,587	\$84,489	\$127,076	\$0	\$127,076
	354	Albatross, short-tailed ( <i>Phoebastria (=Diomedea) albatrus</i> ) - Wherever found	E	\$149,311	\$1,958	\$151,269	\$2,000	\$153,269
	830	Blackbird, yellow-shouldered ( <i>Agelaius xanthomus</i> ) - Wherever found	E	\$18,000	\$250	\$18,250	\$0	\$18,250
	245	Bobwhite, masked (quail) ( <i>Colinus virginianus ridgwayi</i> ) - Wherever found	E	\$353,726	\$21,400	\$375,126	\$0	\$375,126
	31	Caracara, Audubon's crested ( <i>Polyborus plancus audubonii</i> ) - FL pop.	T	\$247,500	\$9,407,913	\$9,655,413	\$0	\$9,655,413
	61	Condor, California ( <i>Gymnogyps californianus</i> ) - U.S.A. only, except where listed as an experimental population	E	\$3,346,481	\$741,632	\$4,088,113	\$0	\$4,088,113
	386	Condor, California ( <i>Gymnogyps californianus</i> ) - U.S.A. (specific portions of Arizona, Nevada, and Utah)	EXPN	\$86,097	\$43,130	\$129,227	\$0	\$129,227
	481	Condor, California ( <i>Gymnogyps californianus</i> ) - Pacific Northwest NEP	PEXP	\$78,318	\$0	\$78,318	\$0	\$78,318
	156	Coot, Hawaiian ( <i>Fulica americana alai</i> ) - Wherever found	E	\$113,113	\$698,397	\$811,510	\$0	\$811,510

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1271	Crane, hooded ( <i>Grus monacha</i> ) - Wherever found	E	\$0	\$20	\$20	\$0	\$20
	289	Crane, Mississippi sandhill ( <i>Grus canadensis pulla</i> ) - Wherever found	E	\$250,000	\$4,626	\$254,626	\$500	\$255,126
	1271	Crane, Siberian white ( <i>Grus leucogeranus</i> ) - Wherever found	E	\$0	\$20	\$20	\$0	\$20
	176	Crane, whooping ( <i>Grus americana</i> ) - Wherever found, except where listed as an experimental population	E	\$412,802	\$222,811	\$635,613	\$9,400	\$645,013
	675	Crane, whooping ( <i>Grus americana</i> ) - U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	EXPN	\$0	\$31,688	\$31,688	\$0	\$31,688
	189	Crane, whooping ( <i>Grus americana</i> ) - U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	EXPN	\$241,500	\$256,523	\$498,023	\$92,000	\$590,023
	660	Crane, whooping ( <i>Grus americana</i> ) - U.S.A (Southwestern Louisiana)	EXPN	\$33,720	\$0	\$33,720	\$0	\$33,720
	293	Creeper, Hawaii ( <i>Oreomystis mana</i> ) - Wherever found	E	\$143,883	\$104,010	\$247,893	\$0	\$247,893
	738	Creeper, Molokai ( <i>Paroreomyza flammea</i> ) - Wherever found	E	\$25,335	\$0	\$25,335	\$0	\$25,335
	957	Creeper, Oahu ( <i>Paroreomyza maculata</i> ) - Wherever found	E	\$11,914	\$0	\$11,914	\$0	\$11,914
	296	Crow, Hawaiian (=alala) ( <i>Corvus hawaiiensis</i> ) - Wherever found	E	\$132,467	\$110,400	\$242,867	\$0	\$242,867
	146	Crow, Mariana (=aga) ( <i>Corvus kubaryi</i> ) - Wherever found	E	\$861,882	\$58,049	\$919,931	\$0	\$919,931
	1232	Crow, white-necked ( <i>Corvus leucognaphalus</i> ) - Wherever found	E	\$2,000	\$0	\$2,000	\$0	\$2,000
	48	Cuckoo, yellow-billed ( <i>Coccyzus americanus</i> ) - Western U.S. DPS	T	\$499,562	\$5,274,443	\$5,774,005	\$16,500	\$5,790,505
	1255	Curlew, Eskimo ( <i>Numenius borealis</i> ) - Wherever found	E	\$629	\$100	\$729	\$0	\$729
	152	Duck, Hawaiian (=koloa) ( <i>Anas wyvilliana</i> ) - Wherever found	E	\$88,860	\$760,984	\$849,844	\$0	\$849,844
	530	Duck, Laysan ( <i>Anas laysanensis</i> ) - Wherever found	E	\$61,773	\$300	\$62,073	\$0	\$62,073



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	63	Eider, spectacled ( <i>Somateria fischeri</i> ) - Wherever found	T	\$759,000	\$3,088,136	\$3,847,136	\$0	\$3,847,136
	67	Eider, Steller's ( <i>Polysticta stelleri</i> ) - AK breeding pop.	T	\$574,000	\$3,113,916	\$3,687,916	\$0	\$3,687,916
	770	Elepaio, Oahu ( <i>Chasiempis ibidis</i> ) - Wherever found	E	\$22,655	\$0	\$22,655	\$0	\$22,655
	350	Falcon, Northern Aplomado ( <i>Falco femoralis septentrionalis</i> ) - Wherever found, except where listed as an experimental population	E	\$0	\$157,326	\$157,326	\$0	\$157,326
	380	Falcon, Northern Aplomado ( <i>Falco femoralis septentrionalis</i> ) - U.S.A (AZ, NM)	EXPN	\$19,700	\$114,500	\$134,200	\$0	\$134,200
	836	Finch, Laysan (honeycreeper) ( <i>Telespyza cantans</i> ) - Wherever found	E	\$17,713	\$0	\$17,713	\$0	\$17,713
	813	Finch, Nihoa (honeycreeper) ( <i>Telespyza ultima</i> ) - Wherever found	E	\$19,133	\$0	\$19,133	\$0	\$19,133
	42	Flycatcher, southwestern willow ( <i>Empidonax traillii extimus</i> ) - Wherever found	E	\$438,021	\$6,850,696	\$7,288,717	\$1,830	\$7,290,547
	165	Gallinule, Hawaiian common ( <i>Gallinula galeata sandvicensis</i> ) - Wherever found	E	\$122,845	\$599,145	\$721,990	\$0	\$721,990
	71	Gnatcatcher, coastal California ( <i>Polioptila californica californica</i> ) - Wherever found	T	\$884,960	\$2,242,596	\$3,127,556	\$0	\$3,127,556
	102	Goose, Hawaiian ( <i>Branta</i> (=Nesochen) <i>sandvicensis</i> ) - Wherever found	E	\$331,937	\$1,167,348	\$1,499,285	\$0	\$1,499,285
	869	Ground-Dove, Friendly ( <i>Gallicolumba stairi</i> ) - American Samoa DPS	E	\$15,758	\$0	\$15,758	\$0	\$15,758
	268	Hawk, Hawaiian (=lo) ( <i>Buteo solitarius</i> ) - Wherever found	E	\$104,674	\$198,200	\$302,874	\$0	\$302,874
	870	Hawk, Puerto Rican broad-winged ( <i>Buteo platypterus brunnescens</i> ) - Wherever found	E	\$15,500	\$250	\$15,750	\$0	\$15,750
	368	Hawk, Puerto Rican sharp-shinned ( <i>Accipiter striatus venator</i> ) - Wherever found	E	\$117,000	\$22,750	\$139,750	\$0	\$139,750
	403	Honeycreeper, (Akohekohe) crested ( <i>Palmeria dolei</i> ) - Wherever found	E	\$18,135	\$105,000	\$123,135	\$0	\$123,135

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	932	(honeyeater), Mao (= maomao) ( <i>Gymnomyza samoensis</i> ) - Wherever found	E	\$12,929	\$0	\$12,929	\$0	\$12,929
	121	Horned lark, streaked ( <i>Eremophila alpestris strigata</i> ) - Wherever found	T	\$724,564	\$330,709	\$1,055,273	\$31,000	\$1,086,273
	356	ʻIiwi ( <i>Drepanis coccinea</i> ) - Wherever found	T	\$44,660	\$107,341	\$152,001	\$0	\$152,001
	401	Kingfisher, Guam Micronesian ( <i>Halcyon cinnamomina cinnamomina</i> ) - Wherever found	E	\$64,511	\$59,002	\$123,513	\$0	\$123,513
	111	Kite, Everglade snail ( <i>Rostrhamus sociabilis plumbeus</i> ) - Wherever found	E	\$448,000	\$884,647	\$1,332,647	\$0	\$1,332,647
	144	Knot, red ( <i>Calidris canutus rufa</i> ) - Wherever found	T	\$504,095	\$372,730	\$876,825	\$62,360	\$939,185
	483	Megapode, Micronesian ( <i>Megapodius laperouse</i> ) - Wherever found	E	\$30,656	\$47,617	\$78,273	\$0	\$78,273
	689	Millerbird, Nihoa (old world warbler) ( <i>Acrocephalus familiaris kingi</i> ) - Wherever found	E	\$30,135	\$0	\$30,135	\$0	\$30,135
	623	Moorhen, Mariana common ( <i>Gallinula chloropus guami</i> ) - Wherever found	E	\$34,001	\$6,221	\$40,222	\$0	\$40,222
	46	Murrelet, marbled ( <i>Brachyramphus marmoratus</i> ) - U.S.A. (CA, OR, WA)	T	\$2,128,419	\$2,145,462	\$4,273,881	\$1,618,600	\$5,892,481
	999	Nightjar, Puerto Rican ( <i>Caprimulgus noctitherus</i> ) - Wherever found	E	\$10,000	\$250	\$10,250	\$0	\$10,250
	834	Nukupuu, Kauai ( <i>Hemignathus hanapepe</i> ) - Wherever found	E	\$17,828	\$0	\$17,828	\$0	\$17,828
	771	Nukupuu, Maui ( <i>Hemignathus affinis</i> ) - Wherever found	E	\$22,357	\$0	\$22,357	\$0	\$22,357
	968	ʻOʻo, Kauai (honeyeater) ( <i>Moho braccatus</i> ) - Wherever found	E	\$11,663	\$0	\$11,663	\$0	\$11,663
	941	ʻOʻu (honeycreeper) ( <i>Psittirostra psittacea</i> ) - Wherever found	E	\$12,588	\$0	\$12,588	\$0	\$12,588
	153	Owl, Mexican spotted ( <i>Strix occidentalis lucida</i> ) - Wherever found	T	\$84,800	\$762,156	\$846,956	\$0	\$846,956
	5	Owl, northern spotted ( <i>Strix occidentalis caurina</i> ) - Wherever found	T	\$4,072,138	\$40,446,261	\$44,518,399	\$914,600	\$45,432,999
	411	Palila (honeycreeper) ( <i>Loxioides bailleui</i> ) - Wherever found	E	\$98,027	\$11,150	\$109,177	\$0	\$109,177

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	366	Parrotbill, Maui (Kiwikiu) ( <i>Pseudonestor xanthophrys</i> ) - Wherever found	E	\$36,533	\$105,000	\$141,533	\$0	\$141,533
	97	Parrot, Puerto Rican ( <i>Amazona vittata</i> ) - Wherever found	E	\$1,512,500	\$130,000	\$1,642,500	\$0	\$1,642,500
	1267	Parrot, thick-billed ( <i>Rhynchopsitta pachyrhyncha</i> ) - Wherever found	E	\$200	\$0	\$200	\$0	\$200
	1077	Petrel, Bermuda ( <i>Pterodroma cahow</i> ) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	163	Petrel, Hawaiian ( <i>Pterodroma sandwichensis</i> ) - Wherever found	E	\$255,211	\$492,954	\$748,165	\$0	\$748,165
	902	Pigeon, Puerto Rican plain ( <i>Columba inornata wetmorei</i> ) - Wherever found	E	\$12,500	\$1,707	\$14,207	\$0	\$14,207
	118	Plover, piping ( <i>Charadrius melodus</i> ) - [Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	E	\$831,029	\$290,654	\$1,121,683	\$35,099	\$1,156,782
	38	Plover, piping ( <i>Charadrius melodus</i> ) - [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	T	\$1,336,288	\$6,426,543	\$7,762,831	\$368,502	\$8,131,333
	60	Plover, western snowy ( <i>Charadrius nivosus nivosus</i> ) - Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	T	\$1,496,588	\$2,499,382	\$3,995,970	\$92,400	\$4,088,370
	469	Po'ouli (honeycreeper) ( <i>Melamprosops phaeosoma</i> ) - Wherever found	E	\$84,263	\$0	\$84,263	\$0	\$84,263
	227	Prairie-chicken, Attwater's greater ( <i>Tympanuchus cupido attwateri</i> ) - Wherever found	E	\$432,835	\$10,500	\$443,335	\$0	\$443,335
	110	Rail, California clapper ( <i>Rallus longirostris obsoletus</i> ) - Wherever found	E	\$617,416	\$724,927	\$1,342,343	\$0	\$1,342,343
	479	Rail, Guam ( <i>Rallus owstoni</i> ) - Wherever found, except where listed as an experimental population	E	\$34,171	\$44,861	\$79,032	\$0	\$79,032
	717	Rail, Guam ( <i>Rallus owstoni</i> ) - Rota	EXPN	\$27,125	\$0	\$27,125	\$0	\$27,125
	247	Rail, light-footed clapper ( <i>Rallus longirostris levipes</i> ) - Wherever found	E	\$169,833	\$200,325	\$370,158	\$0	\$370,158

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	77	Rail, Yuma Ridgways (Rallus obsoletus [=longirostris] yumanensis) - Wherever found	E	\$554,371	\$2,024,772	\$2,579,143	\$0	\$2,579,143
	103	Sage-grouse, Gunnison (Centrocercus minimus) - Wherever found	T	\$0	\$1,030,379	\$1,030,379	\$464,417	\$1,494,796
	117	Scrub-jay, Florida (Aphelocoma coerulescens) - Wherever found	T	\$480,000	\$685,757	\$1,165,757	\$0	\$1,165,757
	304	Shearwater, Newell's Townsend's (Puffinus auricularis newelli) - Wherever found	T	\$214,506	\$8,954	\$223,460	\$0	\$223,460
	76	Shrike, San Clemente loggerhead (Lanius ludovicianus mearnsi) - Wherever found	E	\$30,050	\$2,597,414	\$2,627,464	\$0	\$2,627,464
	113	Sparrow, Cape Sable seaside (Ammodramus maritimus mirabilis) - Wherever found	E	\$507,000	\$741,425	\$1,248,425	\$0	\$1,248,425
	252	Sparrow, Florida grasshopper (Ammodramus savannarum floridanus) - Wherever found	E	\$230,000	\$128,094	\$358,094	\$0	\$358,094
	137	Sparrow, San Clemente sage (Amphispiza belli clementeae) - Wherever found	T	\$21,849	\$967,094	\$988,943	\$0	\$988,943
	147	Stilt, Hawaiian (Himantopus mexicanus knudseni) - Wherever found	E	\$136,581	\$775,396	\$911,977	\$0	\$911,977
	104	Stork, wood (Mycteria americana) - AL, FL, GA, MS, NC, SC	T	\$225,000	\$1,151,257	\$1,376,257	\$75,400	\$1,451,657
	442	Storm-petrel, band-rumped (Oceanodroma castro) - USA (HI)	E	\$86,983	\$8,954	\$95,937	\$0	\$95,937
	262	Swiftlet, Mariana gray (Aerodramus vanikorensis bartschi) - Wherever found	E	\$49,858	\$267,965	\$317,823	\$0	\$317,823
	86	Tern, California least (Sterna antillarum browni) - Wherever found	E	\$758,334	\$1,378,417	\$2,136,751	\$0	\$2,136,751
	45	Tern, least (Sterna antillarum) - interior pop.	E	\$453,472	\$5,500,502	\$5,953,974	\$56,801	\$6,010,775
	325	Tern, roseate (Sterna dougallii dougallii) - Northeast U.S. nesting population	E	\$52,483	\$103,558	\$156,041	\$32,000	\$188,041
	459	Tern, roseate (Sterna dougallii dougallii) - Western Hemisphere except NE U.S.	T	\$22,000	\$65,000	\$87,000	\$0	\$87,000
	416	Thrush, large Kauai (=kamao) (Myadestes myadestinus) - Wherever found	E	\$104,441	\$0	\$104,441	\$0	\$104,441
	911	Thrush, Molokai (Myadestes lanaiensis rutha) - Wherever found	E	\$13,878	\$0	\$13,878	\$0	\$13,878

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	706	Thrush, small Kauai (=puaiohi) ( <i>Myadestes palmeri</i> ) - Wherever found	E	\$28,501	\$0	\$28,501	\$0	\$28,501
	863	Towhee, Inyo California ( <i>Pipilo crissalis eremophilus</i> ) - Wherever found	T	\$9,453	\$6,702	\$16,155	\$0	\$16,155
	62	Vireo, least Bell's ( <i>Vireo bellii pusillus</i> ) - Wherever found	E	\$864,338	\$3,160,664	\$4,025,002	\$0	\$4,025,002
	913	Warbler, elfin-woods ( <i>Setophaga angelae</i> ) - Wherever found	T	\$13,500	\$250	\$13,750	\$0	\$13,750
	217	Warbler, Kirtland's ( <i>Setophaga kirtlandii</i> (= <i>Dendroica kirtlandii</i> )) - Wherever found	E	\$130,600	\$301,346	\$431,946	\$56,366	\$488,312
	718	Warbler, nightingale reed (old world warbler) ( <i>Acrocephalus luscini</i> a) - Wherever found	E	\$27,107	\$0	\$27,107	\$0	\$27,107
	1100	Warbler (=wood), Bachman's ( <i>Vermivora bachmanii</i> ) - Wherever found	E	\$7,000	\$0	\$7,000	\$100	\$7,100
	244	Warbler (=wood), golden-cheeked ( <i>Dendroica chrysoparia</i> ) - Wherever found	E	\$253,420	\$127,020	\$380,440	\$0	\$380,440
	1054	White-eye, bridled ( <i>Zosterops conspicillatus conspicillatus</i> ) - Wherever found	E	\$8,444	\$0	\$8,444	\$0	\$8,444
	646	White-eye, Rota bridled ( <i>Zosterops rotensis</i> ) - Wherever found	E	\$36,706	\$0	\$36,706	\$0	\$36,706
	690	Woodpecker, ivory-billed ( <i>Campephilus principalis</i> ) - Wherever found	E	\$25,029	\$5,000	\$30,029	\$0	\$30,029
	19	Woodpecker, red-cockaded ( <i>Picoides borealis</i> ) - Wherever found	E	\$1,631,589	\$14,849,299	\$16,480,888	\$195,200	\$16,676,088
<b>Birds subtotal</b>				<b>\$33,211,615</b>	<b>\$127,856,132</b>	<b>\$161,067,747</b>	<b>\$4,125,075</b>	<b>\$165,192,822</b>
Reptiles	92	Alligator, American ( <i>Alligator mississippiensis</i> ) - Wherever found	SAT	\$156,500	\$1,470,725	\$1,627,225	\$135,000	\$1,762,225
	1088	Anole, Culebra Island giant ( <i>Anolis roosevelti</i> ) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1210	Boa, Mona ( <i>Epicrates monensis monensis</i> ) - Wherever found	T	\$3,000	\$0	\$3,000	\$0	\$3,000
	621	Boa, Puerto Rican ( <i>Epicrates inornatus</i> ) - Wherever found	E	\$35,500	\$5,413	\$40,913	\$0	\$40,913
	724	Boa, Virgin Islands tree ( <i>Chilabothrus granti</i> ) - Wherever found	E	\$12,000	\$0	\$12,000	\$14,556	\$26,556
	240	Crocodile, American ( <i>Crocodylus acutus</i> ) - U.S.A. (FL)	T	\$39,000	\$372,809	\$411,809	\$0	\$411,809

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	414	Gartersnake, narrow-headed ( <i>Thamnophis rufipunctatus</i> ) - Wherever found	T	\$61,500	\$46,260	\$107,760	\$0	\$107,760
	328	Gartersnake, northern Mexican ( <i>Thamnophis eques megalops</i> ) - Wherever found	T	\$67,000	\$116,975	\$183,975	\$0	\$183,975
	1163	Gecko, Monito ( <i>Sphaerodactylus micropithecus</i> ) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	659	Iguana, Cuban ground ( <i>Cyclura nubila nubila</i> ) - ( <i>Cyclura nubila nubila</i> )	T	\$0	\$33,800	\$33,800	\$0	\$33,800
	1163	Iguana, Mona ground ( <i>Cyclura stejnegeri</i> ) - Wherever found	T	\$5,000	\$0	\$5,000	\$0	\$5,000
	235	Lizard, blunt-nosed leopard ( <i>Gambelia silus</i> ) - Wherever found	E	\$252,000	\$172,335	\$424,335	\$0	\$424,335
	347	Lizard, Coachella Valley fringe-toed ( <i>Uma inornata</i> ) - Wherever found	T	\$48,000	\$110,929	\$158,929	\$0	\$158,929
	763	Lizard, St. Croix ground ( <i>Ameiva polops</i> ) - Wherever found	E	\$20,500	\$2,500	\$23,000	\$0	\$23,000
	145	Massasauga (=rattlesnake), eastern ( <i>Sistrurus catenatus</i> ) - Wherever found	T	\$542,520	\$309,195	\$851,715	\$78,607	\$930,322
	633	Pinesnake, black ( <i>Pituophis melanoleucus lodingi</i> ) - Wherever found	T	\$0	\$38,556	\$38,556	\$400	\$38,956
	105	Pinesnake, Louisiana ( <i>Pituophis ruthveni</i> ) - Wherever found	T	\$54,600	\$1,362,475	\$1,417,075	\$7,399	\$1,424,474
	844	Rattlesnake, New Mexican ridge-nosed ( <i>Crotalus willardi obscurus</i> ) - Wherever found	T	\$16,000	\$1,250	\$17,250	\$0	\$17,250
	109	Sea turtle, green ( <i>Chelonia mydas</i> ) - Central North Pacific DPS	T	\$130,231	\$1,251,640	\$1,381,871	\$0	\$1,381,871
	166	Sea turtle, green ( <i>Chelonia mydas</i> ) - Central South Pacific DPS	E	\$19,532	\$699,570	\$719,102	\$0	\$719,102
	96	Sea turtle, green ( <i>Chelonia mydas</i> ) - Central West Pacific DPS	E	\$88,920	\$1,557,800	\$1,646,720	\$0	\$1,646,720
	467	Sea turtle, green ( <i>Chelonia mydas</i> ) - East Indian-West Pacific DPS	T	\$84,784	\$0	\$84,784	\$0	\$84,784
	887	Sea turtle, green ( <i>Chelonia mydas</i> ) - Mediterranean DPS	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	99	Sea turtle, green ( <i>Chelonia mydas</i> ) - East Pacific DPS	T	\$66,628	\$1,474,462	\$1,541,090	\$0	\$1,541,090

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	80	Sea turtle, green ( <i>Chelonia mydas</i> ) - North Atlantic DPS	T	\$147,500	\$2,321,002	\$2,468,502	\$3,000	\$2,471,502
	125	Sea turtle, green ( <i>Chelonia mydas</i> ) - South Atlantic DPS	T	\$50,800	\$997,330	\$1,048,130	\$12,000	\$1,060,130
	65	Sea turtle, hawksbill ( <i>Eretmochelys imbricata</i> ) - Wherever found	E	\$525,783	\$3,220,650	\$3,746,433	\$1,400	\$3,747,833
	51	Sea turtle, Kemp's ridley ( <i>Lepidochelys kempii</i> ) - Wherever found	E	\$160,500	\$5,275,332	\$5,435,832	\$12,900	\$5,448,732
	56	Sea turtle, leatherback ( <i>Dermochelys coriacea</i> ) - Wherever found	E	\$815,320	\$3,893,215	\$4,708,535	\$6,800	\$4,715,335
	678	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - Wherever found	RT	\$0	\$0	\$0	\$31,300	\$31,300
	53	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - Northwest Atlantic Ocean DPS	T	\$568,206	\$4,228,283	\$4,796,489	\$150,000	\$4,946,489
	90	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - North Pacific Ocean DPS	E	\$10,907	\$1,894,139	\$1,905,046	\$0	\$1,905,046
	887	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - Mediterranean Sea DPS	E	\$15,000	\$0	\$15,000	\$0	\$15,000
	430	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - North Indian Ocean DPS	E	\$98,641	\$0	\$98,641	\$0	\$98,641
	261	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - Northeast Atlantic Ocean DPS	E	\$73,800	\$244,853	\$318,653	\$0	\$318,653
	242	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - South Atlantic Ocean DPS	T	\$0	\$390,716	\$390,716	\$0	\$390,716
	183	Sea turtle, olive ridley ( <i>Lepidochelys olivacea</i> ) - Wherever found, except when listed as endangered under 50 CFR 224.101	T	\$22,679	\$576,537	\$599,216	\$0	\$599,216
	264	Sea turtle, olive ridley ( <i>Lepidochelys olivacea</i> ) - Breeding colony populations on Pacific coast of Mexico	E	\$316,689	\$0	\$316,689	\$0	\$316,689
	367	Skink, bluetail mole ( <i>Eumeces egregius lividus</i> ) - Wherever found	T	\$83,000	\$58,106	\$141,106	\$0	\$141,106
	243	Skink, sand ( <i>Neoseps reynoldsi</i> ) - Wherever found	T	\$277,500	\$110,725	\$388,225	\$0	\$388,225
	544	Skink, Slevin's ( <i>Emoia slevini</i> ) - Wherever found	E	\$58,305	\$0	\$58,305	\$0	\$58,305
	929	Snake, Atlantic salt marsh ( <i>Nerodia clarkii taeniata</i> ) - Wherever found	T	\$12,500	\$569	\$13,069	\$0	\$13,069

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1090	Snake, copperbelly water ( <i>Nerodia erythrogaster neglecta</i> ) - Indiana north of 40 degrees north latitude, Michigan, Ohio	T	\$0	\$1,066	\$1,066	\$6,366	\$7,432
	1163	Snake, copperbelly water ( <i>Nerodia erythrogaster neglecta</i> ) - Illinois, Kentucky, and Indiana south of 40 degrees north latitude	RT	\$5,000	\$0	\$5,000	\$0	\$5,000
	101	Snake, eastern indigo ( <i>Drymarchon corais couperi</i> ) - Wherever found	T	\$426,000	\$891,654	\$1,317,654	\$200,000	\$1,517,654
	131	Snake, giant garter ( <i>Thamnophis gigas</i> ) - Wherever found	T	\$589,100	\$422,718	\$1,011,818	\$0	\$1,011,818
	257	Snake, San Francisco garter ( <i>Thamnophis sirtalis tetrataenia</i> ) - Wherever found	E	\$277,300	\$54,594	\$331,894	\$0	\$331,894
	664	Tortoise, desert ( <i>Gopherus agassizii</i> ) - AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	SAT	\$33,100	\$100	\$33,200	\$0	\$33,200
	26	Tortoise, desert ( <i>Gopherus agassizii</i> ) - Wherever found, except AZ south and east of Colorado R., and Mexico	T	\$1,805,700	\$10,854,890	\$12,660,590	\$0	\$12,660,590
	78	Tortoise, gopher ( <i>Gopherus polyphemus</i> ) - West of Mobile and Tombigbee Rivers	T	\$48,500	\$2,499,279	\$2,547,779	\$6,000	\$2,553,779
	426	Tracaja ( <i>Podocnemis unifilis</i> ) - Wherever found	E	\$99,909	\$0	\$99,909	\$0	\$99,909
	1059	Turtle, Alabama red-bellied ( <i>Pseudemys alabamensis</i> ) - Wherever found	E	\$200	\$8,058	\$8,258	\$100	\$8,358
	181	Turtle, bog ( <i>Clemmys muhlenbergii</i> ) - Wherever found, except GA, NC, SC, TN, VA	T	\$285,076	\$284,330	\$569,406	\$38,000	\$607,406
	307	Turtle, bog ( <i>Clemmys muhlenbergii</i> ) - U.S.A. (GA, NC, SC, TN, VA)	SAT	\$99,943	\$4,500	\$104,443	\$115,100	\$219,543
	1227	Turtle, flattened musk ( <i>Sternotherus depressus</i> ) - Black Warrior R. system upstream from Bankhead Dam	T	\$900	\$1,337	\$2,237	\$0	\$2,237
	835	Turtle, Plymouth Redbelly ( <i>Pseudemys rubriventris bangsi</i> ) - Wherever found	E	\$17,340	\$100	\$17,440	\$300	\$17,740
	687	Turtle, ringed map ( <i>Graptemys oculifera</i> ) - Wherever found	T	\$24,000	\$6,169	\$30,169	\$400	\$30,569



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	901	Turtle, Sonoyta mud ( <i>Kinosternon sonoriense longifemorale</i> ) - Wherever found	E	\$13,500	\$756	\$14,256	\$0	\$14,256
	1199	Turtle, yellow-blotched map ( <i>Graptemys flavimaculata</i> ) - Wherever found	T	\$0	\$2,913	\$2,913	\$400	\$3,313
	290	Whipsnake (=striped racer), Alameda ( <i>Masticophis lateralis euryxanthus</i> ) - Wherever found	T	\$96,600	\$153,929	\$250,529	\$0	\$250,529
<b>Reptiles subtotal</b>				<b>\$8,790,013</b>	<b>\$47,424,544</b>	<b>\$56,214,557</b>	<b>\$820,028</b>	<b>\$57,034,585</b>
Amphibians	907	Coqui, golden ( <i>Eleutherodactylus jasperi</i> ) - Wherever found	T	\$14,000	\$0	\$14,000	\$0	\$14,000
	175	Coqui, Llanero ( <i>Eleutherodactylus juanariveroi</i> ) - Wherever found	E	\$4,500	\$652,524	\$657,024	\$0	\$657,024
	82	Frog, California red-legged ( <i>Rana draytonii</i> ) - Wherever found	T	\$1,409,605	\$992,043	\$2,401,648	\$0	\$2,401,648
	312	Frog, Chiricahua leopard ( <i>Rana chiricahuensis</i> ) - Wherever found	T	\$75,489	\$139,494	\$214,983	\$0	\$214,983
	222	Frog, dusky gopher ( <i>Rana sevosa</i> ) - Wherever found	E	\$169,000	\$271,568	\$440,568	\$22,000	\$462,568
	1210	Frog, Goliath ( <i>Conraua goliath</i> ) - Wherever found	T	\$0	\$3,000	\$3,000	\$0	\$3,000
	374	Frog, mountain yellow-legged ( <i>Rana muscosa</i> ) - Southern California DPS	E	\$76,115	\$60,706	\$136,821	\$0	\$136,821
	205	Frog, mountain yellow-legged ( <i>Rana muscosa</i> ) - Northern California DPS	E	\$216,242	\$311,600	\$527,842	\$0	\$527,842
	95	Frog, Oregon spotted ( <i>Rana pretiosa</i> ) - Wherever found	T	\$982,887	\$618,808	\$1,601,695	\$52,800	\$1,654,495
	141	Frog, Sierra Nevada Yellow-legged ( <i>Rana sierrae</i> ) - Wherever found	E	\$247,124	\$703,200	\$950,324	\$0	\$950,324
	1115	Guajon ( <i>Eleutherodactylus cooki</i> ) - Wherever found	T	\$6,500	\$0	\$6,500	\$0	\$6,500
	265	Hellbender, Ozark ( <i>Cryptobranchus alleganiensis bishopi</i> ) - Wherever found	E	\$196,000	\$15,887	\$211,887	\$100,900	\$312,787
	1087	Salamander, Austin blind ( <i>Eurycea waterlooensis</i> ) - Wherever found	E	\$5,000	\$0	\$5,000	\$2,502	\$7,502
	253	Salamander, Barton Springs ( <i>Eurycea sosorum</i> ) - Wherever found	E	\$10,000	\$336,224	\$346,224	\$2,502	\$348,726

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	193	Salamander, California tiger ( <i>Ambystoma californiense</i> ) - U.S.A. (CA - Santa Barbara County)	E	\$568,688	\$400	\$569,088	\$0	\$569,088
	182	Salamander, California tiger ( <i>Ambystoma californiense</i> ) - U.S.A. (CA - Sonoma County)	E	\$568,969	\$36,000	\$604,969	\$0	\$604,969
	128	Salamander, California tiger ( <i>Ambystoma californiense</i> ) - U.S.A. (Central CA DPS)	T	\$774,855	\$255,617	\$1,030,472	\$0	\$1,030,472
	233	Salamander, Cheat Mountain ( <i>Plethodon nettingi</i> ) - Wherever found	T	\$28,471	\$404,239	\$432,710	\$0	\$432,710
	985	Salamander, desert slender ( <i>Batrachoseps aridus</i> ) - Wherever found	E	\$578	\$10,200	\$10,778	\$0	\$10,778
	190	Salamander, frosted flatwoods ( <i>Ambystoma cingulatum</i> ) - Wherever found	T	\$232,500	\$319,432	\$551,932	\$30,000	\$581,932
	1162	Salamander, Georgetown ( <i>Eurycea naufragia</i> ) - Wherever found	T	\$2,500	\$0	\$2,500	\$2,502	\$5,002
	511	Salamander, Jemez Mountains ( <i>Plethodon neomexicanus</i> ) - Wherever found	E	\$17,000	\$50,000	\$67,000	\$0	\$67,000
	1003	Salamander, Jollyville Plateau ( <i>Eurycea tonkawae</i> ) - Wherever found	T	\$10,000	\$0	\$10,000	\$0	\$10,000
	1064	Salamander, Red Hills ( <i>Phaeognathus hubrichti</i> ) - Wherever found	T	\$7,000	\$1,248	\$8,248	\$0	\$8,248
	364	Salamander, Reticulated flatwoods ( <i>Ambystoma bishopi</i> ) - Wherever found	E	\$30,000	\$84,369	\$114,369	\$30,000	\$144,369
	448	Salamander, Salado ( <i>Eurycea chisholmensis</i> ) - Wherever found	T	\$5,000	\$89,216	\$94,216	\$0	\$94,216
	983	Salamander, San Marcos ( <i>Eurycea nana</i> ) - Wherever found	T	\$10,000	\$840	\$10,840	\$0	\$10,840
	1220	Salamander, San Marcos ( <i>Eurycea nana</i> ) - Spring Lake and nearby habitats immediately downstream of the dam forming Spring Lake	RT	\$0	\$0	\$0	\$2,502	\$2,502
	485	Salamander, Santa Cruz long-toed ( <i>Ambystoma macrodactylum croceum</i> ) - Wherever found	E	\$75,963	\$1,600	\$77,563	\$0	\$77,563
	486	Salamander, Shenandoah ( <i>Plethodon shenandoah</i> ) - Wherever found	E	\$2,000	\$75,421	\$77,421	\$0	\$77,421

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	945	Salamander, Sonora tiger ( <i>Ambystoma tigrinum stebbinsi</i> ) - Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	983	Salamander, Texas blind ( <i>Eurycea [=Typhlomolge] rathbuni</i> ) - Wherever found	E	\$10,000	\$840	\$10,840	\$0	\$10,840
	134	Toad, arroyo (=arroyo southwestern) ( <i>Anaxyrus californicus</i> ) - Wherever found	E	\$257,341	\$737,609	\$994,950	\$0	\$994,950
	294	Toad, Houston ( <i>Bufo houstonensis</i> ) - Wherever found	E	\$111,000	\$132,007	\$243,007	\$4,387	\$247,394
	359	Toad, Puerto Rican crested ( <i>Peltophryne lemur</i> ) - Wherever found	T	\$5,000	\$142,491	\$147,491	\$0	\$147,491
	250	Toad, Wyoming ( <i>Bufo hemiophrys baxteri</i> ) - Wherever found	E	\$331,300	\$14,500	\$345,800	\$15,251	\$361,051
	271	Toad, Yosemite ( <i>Anaxyrus canorus</i> ) - Wherever found	T	\$221,760	\$67,606	\$289,366	\$0	\$289,366
	1130	Waterdog, black warrior (=Sipsey Fork) ( <i>Necturus alabamensis</i> ) - Wherever found	E	\$4,500	\$1,637	\$6,137	\$0	\$6,137
<b>Amphibians subtotal</b>				<b>\$6,699,387</b>	<b>\$6,530,326</b>	<b>\$13,229,713</b>	<b>\$265,346</b>	<b>\$13,495,059</b>
Fishes	620	Blindcat, Mexican (catfish) ( <i>Prietella phreatophila</i> ) - Wherever found	E	\$0	\$41,000	\$41,000	\$0	\$41,000
	130	Bocaccio ( <i>Sebastes paucispinis</i> ) - Puget Sound - Georgia Basin DPS	E	\$0	\$776,899	\$776,899	\$235,600	\$1,012,499
	32	Bonytail ( <i>Gila elegans</i> ) - Wherever found	E	\$365,325	\$8,919,502	\$9,284,827	\$217,932	\$9,502,759
	841	Bonytongue, Asian ( <i>Scleropages formosus</i> ) - Wherever found	E	\$17,325	\$0	\$17,325	\$0	\$17,325
	537	Catfish, Yaqui ( <i>Ictalurus pricei</i> ) - Wherever found	T	\$43,700	\$16,000	\$59,700	\$0	\$59,700
	1256	Cavefish, Alabama ( <i>Speoplatyrhinus poulsoni</i> ) - Wherever found	E	\$700	\$0	\$700	\$0	\$700
	464	Cavefish, Ozark ( <i>Amblyopsis rosae</i> ) - Wherever found	T	\$57,609	\$27,960	\$85,569	\$500	\$86,069
	353	Chub, Borax Lake ( <i>Gila boraxobius</i> ) - Wherever found	E	\$115,100	\$1,300	\$116,400	\$37,000	\$153,400
	1238	Chub, Chihuahua ( <i>Gila nigrescens</i> ) - Wherever found	T	\$0	\$1,600	\$1,600	\$0	\$1,600
	178	Chub, Gila ( <i>Gila intermedia</i> ) - Wherever found	E	\$48,000	\$589,200	\$637,200	\$0	\$637,200
	41	Chub, humpback ( <i>Gila cypha</i> ) - Wherever found	E	\$31,300	\$7,188,803	\$7,220,103	\$83,248	\$7,303,351

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	921	Chub, Hutton tui ( <i>Gila bicolor</i> ssp.) - Wherever found	T	\$13,500	\$0	\$13,500	\$0	\$13,500
	618	Chub, Owens Tui ( <i>Gila bicolor</i> ssp. <i>snyderi</i> ) - Wherever found	E	\$24,024	\$17,000	\$41,024	\$0	\$41,024
	282	Chub, Pahrnagat roundtail ( <i>Gila robusta jordani</i> ) - Wherever found	E	\$195,617	\$73,300	\$268,917	\$0	\$268,917
	1234	Chub, slender ( <i>Erimystax cahni</i> ) - Wherever found	T	\$1,500	\$370	\$1,870	\$0	\$1,870
	1221	Chub, Sonora ( <i>Gila ditaenia</i> ) - Wherever found	T	\$2,500	\$0	\$2,500	\$0	\$2,500
	512	Chub, spotfin ( <i>Erimonax monachus</i> ) - Wherever found, except where listed as an experimental population	T	\$60,000	\$2,666	\$62,666	\$4,000	\$66,666
	1248	Chub, spotfin ( <i>Erimonax monachus</i> ) - U.S.A. (TN-specified portions of the Tellico River; see 17.84(m)(1)(i))	EXPN	\$100	\$1,056	\$1,156	\$0	\$1,156
	1254	Chub, spotfin ( <i>Erimonax monachus</i> ) - U.S.A. (AL, TN-specified portions of Shoal Creek; see 17.84(m)(1)(ii))	EXPN	\$800	\$0	\$800	\$0	\$800
	1270	Chub, spotfin ( <i>Erimonax monachus</i> ) - U.S.A. (TN-specified portions of the French Broad and Holston Rivers; see 17.84(m)(1)(iii))	EXPN	\$100	\$0	\$100	\$0	\$100
	308	Chub, Virgin River ( <i>Gila seminuda</i> (=robusta)) - Wherever found	E	\$24,072	\$195,300	\$219,372	\$0	\$219,372
	1098	Chub, Yaqui ( <i>Gila purpurea</i> ) - Wherever found	E	\$7,200	\$0	\$7,200	\$0	\$7,200
	226	Cui-ui ( <i>Chasmistes cujus</i> ) - Wherever found	E	\$383,229	\$61,640	\$444,869	\$0	\$444,869
	990	Dace, Ash Meadows speckled ( <i>Rhinichthys osculus nevadensis</i> ) - Wherever found	E	\$10,582	\$0	\$10,582	\$0	\$10,582
	388	Dace, blackside ( <i>Phoxinus cumberlandensis</i> ) - Wherever found	T	\$123,000	\$5,929	\$128,929	\$0	\$128,929
	849	Dace, Clover Valley speckled ( <i>Rhinichthys osculus oligoporus</i> ) - Wherever found	E	\$17,039	\$0	\$17,039	\$0	\$17,039
	281	Dace, desert ( <i>Eremichthys acros</i> ) - Wherever found	T	\$41,063	\$229,750	\$270,813	\$0	\$270,813

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	545	Dace, Foscett speckled ( <i>Rhinichthys osculus</i> ssp.) - Wherever found	T	\$0	\$15,000	\$15,000	\$43,000	\$58,000
	1223	Dace, Independence Valley speckled ( <i>Rhinichthys osculus lethoporus</i> ) - Wherever found	E	\$2,402	\$0	\$2,402	\$0	\$2,402
	508	Dace, Kendall Warm Springs ( <i>Rhinichthys osculus thermalis</i> ) - Wherever found	E	\$66,500	\$1,500	\$68,000	\$0	\$68,000
	1156	Dace, Laurel ( <i>Chrosomus saylori</i> ) - Wherever found	E	\$5,000	\$250	\$5,250	\$0	\$5,250
	301	Dace, Moapa ( <i>Moapa coriacea</i> ) - Wherever found	E	\$174,898	\$56,000	\$230,898	\$0	\$230,898
	1045	Darter, amber ( <i>Percina antesella</i> ) - Wherever found	E	\$0	\$3,732	\$3,732	\$5,000	\$8,732
	1260	Darter, bayou ( <i>Etheostoma rubrum</i> ) - Wherever found	T	\$0	\$0	\$0	\$400	\$400
	1138	Darter, bluemask ( <i>Etheostoma akatulo</i> ) - Wherever found	E	\$4,000	\$1,926	\$5,926	\$0	\$5,926
	484	Darter, boulder ( <i>Etheostoma wapiti</i> ) - Wherever found	E	\$18,500	\$59,500	\$78,000	\$0	\$78,000
	1153	Darter, Cherokee ( <i>Etheostoma scotti</i> ) - Wherever found	T	\$0	\$330	\$330	\$5,000	\$5,330
	666	Darter, Cumberland ( <i>Etheostoma susanae</i> ) - Wherever found	E	\$31,000	\$748	\$31,748	\$1,000	\$32,748
	1143	Darter, diamond ( <i>Crystallaria cincotta</i> ) - Wherever found	E	\$4,744	\$1,030	\$5,774	\$0	\$5,774
	310	Darter, duskytail ( <i>Etheostoma percnum</i> ) - Wherever found	E	\$34,500	\$181,841	\$216,341	\$0	\$216,341
	715	Darter, Etowah ( <i>Etheostoma etowahae</i> ) - Wherever found	E	\$22,000	\$330	\$22,330	\$5,000	\$27,330
	983	Darter, fountain ( <i>Etheostoma fonticola</i> ) - Wherever found	E	\$10,000	\$840	\$10,840	\$0	\$10,840
	1063	Darter, goldline ( <i>Percina aurolineata</i> ) - Wherever found	T	\$2,500	\$772	\$3,272	\$5,000	\$8,272
	778	Darter, Kentucky arrow ( <i>Etheostoma spilotum</i> ) - Wherever found	T	\$17,500	\$3,182	\$20,682	\$500	\$21,182
	451	Darter, leopard ( <i>Percina pantherina</i> ) - Wherever found	T	\$57,674	\$0	\$57,674	\$32,500	\$90,174
	1266	Darter, Maryland ( <i>Etheostoma sellare</i> ) - Wherever found	E	\$0	\$240	\$240	\$0	\$240

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	331	Darter, Niangua ( <i>Etheostoma nianguae</i> ) - Wherever found	T	\$145,000	\$13,565	\$158,565	\$18,500	\$177,065
	272	Darter, Okaloosa ( <i>Etheostoma okaloosae</i> ) - Wherever found	T	\$17,000	\$270,000	\$287,000	\$0	\$287,000
	1257	Darter, Pearl ( <i>Percina aurora</i> ) - Wherever found	T	\$0	\$678	\$678	\$0	\$678
	798	Darter, relict ( <i>Etheostoma chienense</i> ) - Wherever found	E	\$19,500	\$0	\$19,500	\$500	\$20,000
	1211	Darter, rush ( <i>Etheostoma phytophilum</i> ) - Wherever found	E	\$2,500	\$418	\$2,918	\$0	\$2,918
	1210	Darter, slackwater ( <i>Etheostoma boschungii</i> ) - Wherever found	T	\$3,000	\$0	\$3,000	\$0	\$3,000
	940	Darter, snail ( <i>Percina tanasi</i> ) - Wherever found	T	\$8,500	\$1,127	\$9,627	\$3,000	\$12,627
	1221	Darter, vermilion ( <i>Etheostoma chermocki</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	378	Darter, watercress ( <i>Etheostoma nuchale</i> ) - Wherever found	E	\$134,500	\$1,120	\$135,620	\$0	\$135,620
	884	Darter, yellowcheek ( <i>Etheostoma moorei</i> ) - Wherever found	E	\$14,500	\$0	\$14,500	\$800	\$15,300
	66	Eulachon ( <i>Thaleichthys pacificus</i> ) - Southern DPS	T	\$0	\$3,435,561	\$3,435,561	\$282,600	\$3,718,161
	1232	Gambusia, Big Bend ( <i>Gambusia gaigei</i> ) - Wherever found	E	\$1,000	\$1,000	\$2,000	\$0	\$2,000
	1253	Gambusia, Clear Creek ( <i>Gambusia heterochir</i> ) - Wherever found	E	\$1,000	\$0	\$1,000	\$0	\$1,000
	886	Gambusia, Pecos ( <i>Gambusia nobilis</i> ) - Wherever found	E	\$5,250	\$10,000	\$15,250	\$0	\$15,250
	1163	Gambusia, San Marcos ( <i>Gambusia georgei</i> ) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	161	Goby, tidewater ( <i>Eucyclogobius newberryi</i> ) - Wherever found	E	\$455,214	\$311,972	\$767,186	\$0	\$767,186
	209	Grouper, Nassau ( <i>Epinephelus striatus</i> ) -	T	\$0	\$510,951	\$510,951	\$0	\$510,951
	1154	Logperch, Conasauga ( <i>Percina jenkinsi</i> ) - Wherever found	E	\$300	\$0	\$300	\$5,000	\$5,300
	338	Logperch, Roanoke ( <i>Percina rex</i> ) - Wherever found	E	\$142,000	\$20,140	\$162,140	\$3,000	\$165,140
	1262	Madtom, chucky ( <i>Noturus crypticus</i> ) - Wherever found	E	\$300	\$0	\$300	\$0	\$300

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	527	Madtom, Neosho ( <i>Noturus placidus</i> ) - Wherever found	T	\$61,158	\$1,440	\$62,598	\$0	\$62,598
	685	Madtom, pygmy ( <i>Noturus stanauli</i> ) - Wherever found	E	\$30,500	\$311	\$30,811	\$0	\$30,811
	1251	Madtom, Scioto ( <i>Noturus trautmani</i> ) - Wherever found	E	\$0	\$1,030	\$1,030	\$0	\$1,030
	1065	Madtom, smoky ( <i>Noturus baileyi</i> ) - Wherever found	E	\$300	\$7,947	\$8,247	\$0	\$8,247
	1253	Madtom, smoky ( <i>Noturus baileyi</i> ) - Tellico River, TN	EXPN	\$1,000	\$0	\$1,000	\$0	\$1,000
	624	Madtom, yellowfin ( <i>Noturus flavipinnis</i> ) - Wherever found, except where listed as an experimental population	T	\$32,000	\$8,157	\$40,157	\$0	\$40,157
	1210	Madtom, yellowfin ( <i>Noturus flavipinnis</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(e)(1)(iii))	EXPN	\$3,000	\$0	\$3,000	\$0	\$3,000
	842	Minnow, Devils River ( <i>Dionda diaboli</i> ) - Wherever found	T	\$2,500	\$100	\$2,600	\$14,705	\$17,305
	172	Minnow, loach ( <i>Tiaroga cobitis</i> ) - Wherever found	E	\$48,000	\$625,507	\$673,507	\$0	\$673,507
	44	Minnow, Rio Grande Silvery ( <i>Hybognathus amarus</i> ) - Wherever found, except where listed as an experimental population	E	\$97,790	\$6,731,846	\$6,829,636	\$0	\$6,829,636
	1262	Minnow, Rio Grande Silvery ( <i>Hybognathus amarus</i> ) - Rio Grande, from Little Box Canyon to Amistad Dam	EXPN	\$0	\$300	\$300	\$0	\$300
	59	Pikeminnow (=squawfish), Colorado ( <i>Ptychocheilus lucius</i> ) - Wherever found, except where listed as an experimental population	E	\$108,600	\$3,925,294	\$4,033,894	\$83,248	\$4,117,142
	980	Pikeminnow (=squawfish), Colorado ( <i>Ptychocheilus lucius</i> ) - Salt and Verde R. drainages, AZ	EXPN	\$11,000	\$0	\$11,000	\$0	\$11,000
	427	Poolfish, Pahump ( <i>Empetrichthys latos</i> ) - Wherever found	E	\$94,894	\$5,000	\$99,894	\$0	\$99,894
	381	Pupfish, Ash Meadows Amargosa ( <i>Cyprinodon nevadensis mionectes</i> ) - Wherever found	E	\$133,581	\$0	\$133,581	\$0	\$133,581

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1088	Pupfish, Comanche Springs (Cyprinodon elegans) - Wherever found	E	\$2,500	\$5,000	\$7,500	\$0	\$7,500
	437	Pupfish, desert (Cyprinodon macularius) - Wherever found	E	\$23,000	\$74,055	\$97,055	\$0	\$97,055
	149	Pupfish, Devils Hole (Cyprinodon diabolis) - Wherever found	E	\$506,463	\$400,000	\$906,463	\$0	\$906,463
	1221	Pupfish, Leon Springs (Cyprinodon bovinus) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	501	Pupfish, Owens (Cyprinodon radiosus) - Wherever found	E	\$24,024	\$46,000	\$70,024	\$0	\$70,024
	558	Pupfish, Warm Springs (Cyprinodon nevadensis pectoralis) - Wherever found	E	\$53,479	\$0	\$53,479	\$0	\$53,479
	267	Rockfish, canary (Sebastes pinniger) - Puget Sound - Georgia Basin DPS	T	\$0	\$5,330	\$5,330	\$303,200	\$308,530
	127	Rockfish, yelloweye (Sebastes ruberrimus) - Puget Sound - Georgia Basin DPS	T	\$0	\$727,751	\$727,751	\$303,200	\$1,030,951
	37	Salmon, Atlantic (Salmo salar) - Gulf of Maine DPS	E	\$2,434,963	\$6,209,208	\$8,644,171	\$0	\$8,644,171
	8	Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Snake River fall-run ESU	T	\$391,488	\$35,468,592	\$35,860,080	\$408,900	\$36,268,980
	4	Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Snake River spring/summer-run ESU	T	\$302,988	\$55,493,134	\$55,796,122	\$1,175,000	\$56,971,122
	13	Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Sacramento River winter-run ESU	E	\$102,441	\$21,878,718	\$21,981,159	\$0	\$21,981,159
	1	Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Puget Sound ESU	T	\$213,546	\$10,543,952	\$10,757,498	\$63,067,800	\$73,825,298
	10	Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Lower Columbia River ESU	T	\$754,988	\$19,752,046	\$20,507,034	\$9,555,200	\$30,062,234
	20	Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Upper Willamette River ESU	T	\$80,988	\$14,702,592	\$14,783,580	\$407,000	\$15,190,580
	6	Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Upper Columbia spring-run ESU	E	\$157,909	\$36,685,972	\$36,843,881	\$5,163,800	\$42,007,681
	17	Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Central Valley spring-run ESU	T	\$264,200	\$18,172,587	\$18,436,787	\$0	\$18,436,787



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	64	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) tshawytscha) - California Coastal ESU	T	\$0	\$3,760,184	\$3,760,184	\$0	\$3,760,184
	1015	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) tshawytscha) - Central Valley spring-run ESU	EXPN	\$0	\$9,700	\$9,700	\$0	\$9,700
	25	Salmon, chum ( <i>Oncorhynchus keta</i> ) - Columbia River ESU	T	\$800	\$12,015,363	\$12,016,163	\$706,000	\$12,722,163
	40	Salmon, chum ( <i>Oncorhynchus keta</i> ) - Hood Canal summer-run ESU	T	\$27,507	\$4,593,564	\$4,621,071	\$2,996,400	\$7,617,471
	29	Salmon, coho ( <i>Oncorhynchus</i> (=Salmo) kisutch) - Central California Coast ESU	E	\$5,123	\$10,750,215	\$10,755,338	\$0	\$10,755,338
	21	Salmon, coho ( <i>Oncorhynchus</i> (=Salmo) kisutch) - Southern Oregon - Northern California Coast ESU	T	\$319,725	\$13,359,410	\$13,679,135	\$110,000	\$13,789,135
	34	Salmon, coho ( <i>Oncorhynchus</i> (=Salmo) kisutch) - Oregon Coast ESU	T	\$268,728	\$6,967,459	\$7,236,187	\$1,912,000	\$9,148,187
	18	Salmon, coho ( <i>Oncorhynchus</i> (=Salmo) kisutch) - Lower Columbia River ESU	T	\$181,628	\$14,454,702	\$14,636,330	\$3,268,600	\$17,904,930
	11	Salmon, sockeye ( <i>Oncorhynchus</i> (=Salmo) nerka) - Snake River ESU	E	\$578,722	\$24,052,804	\$24,631,526	\$223,300	\$24,854,826
	54	Salmon, sockeye ( <i>Oncorhynchus</i> (=Salmo) nerka) - Ozette Lake ESU	T	\$0	\$4,669,228	\$4,669,228	\$230,300	\$4,899,528
	1256	Sawfish, largetooth ( <i>Pristis pristis</i> ) - Wherever found	E	\$0	\$700	\$700	\$0	\$700
	157	Sawfish, smalltooth ( <i>Pristis pectinata</i> ) - US DPS - Smalltooth sawfish originating from U.S. waters.	E	\$0	\$793,678	\$793,678	\$0	\$793,678
	502	Sculpin, Grotto ( <i>Cottus specus</i> ) - Wherever found	E	\$27,900	\$5,000	\$32,900	\$37,000	\$69,900
	1190	Sculpin, pygmy ( <i>Cottus paulus</i> (=pygmaeus)) - Wherever found	T	\$3,500	\$0	\$3,500	\$0	\$3,500
	808	Shark, Scalloped Hammerhead ( <i>Sphyrna lewini</i> ) - Wherever found	RT	\$0	\$0	\$0	\$19,500	\$19,500
	1006	Shark, Scalloped Hammerhead ( <i>Sphyrna lewini</i> ) - Eastern Pacific DPS	E	\$0	\$9,902	\$9,902	\$0	\$9,902
	1242	Shark, Scalloped Hammerhead ( <i>Sphyrna lewini</i> ) - Central and Southwest Atlantic DPS	T	\$0	\$1,407	\$1,407	\$0	\$1,407

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1206	Shark, Scalloped Hammerhead ( <i>Sphyrna lewini</i> ) - Indo-West Pacific DPS	T	\$0	\$3,127	\$3,127	\$0	\$3,127
	317	Shiner, Arkansas River ( <i>Notropis girardi</i> ) - Arkansas River Basin (AR, KS, NM, OK, TX)	T	\$140,986	\$61,920	\$202,906	\$0	\$202,906
	980	Shiner, beautiful ( <i>Cyprinella formosa</i> ) - Wherever found	T	\$11,000	\$0	\$11,000	\$0	\$11,000
	1101	Shiner, blue ( <i>Cyprinella caerulea</i> ) - Wherever found	T	\$2,000	\$0	\$2,000	\$5,000	\$7,000
	1200	Shiner, Cahaba ( <i>Notropis cahabae</i> ) - Wherever found	E	\$2,500	\$805	\$3,305	\$0	\$3,305
	755	Shiner, Cape Fear ( <i>Notropis mekistocholas</i> ) - Wherever found	E	\$16,000	\$3,640	\$19,640	\$3,800	\$23,440
	1151	Shiner, palezone ( <i>Notropis albizonatus</i> ) - Wherever found	E	\$5,500	\$0	\$5,500	\$0	\$5,500
	89	Shiner, Pecos bluntnose ( <i>Notropis simus pecosensis</i> ) - Wherever found	T	\$10,000	\$2,010,750	\$2,020,750	\$0	\$2,020,750
	628	Shiner, sharpnose ( <i>Notropis oxyrhynchus</i> ) - Wherever found	E	\$25,500	\$0	\$25,500	\$14,165	\$39,665
	628	Shiner, smalleye ( <i>Notropis buccula</i> ) - Wherever found	E	\$25,500	\$0	\$25,500	\$14,165	\$39,665
	139	Shiner, Topeka ( <i>Notropis topeka</i> (=tristis)) - Wherever found, except where listed as an experimental population	E	\$827,700	\$116,432	\$944,132	\$10,300	\$954,432
	1122	Silverside, Waccamaw ( <i>Menidia extensa</i> ) - Wherever found	T	\$3,000	\$0	\$3,000	\$3,400	\$6,400
	14	Smelt, delta ( <i>Hypomesus transpacificus</i> ) - Wherever found	T	\$1,081,441	\$19,724,428	\$20,805,869	\$0	\$20,805,869
	173	Spikedace ( <i>Meda fulgida</i> ) - Wherever found	E	\$46,000	\$625,507	\$671,507	\$0	\$671,507
	1001	Spinedace, Big Spring ( <i>Lepidomeda mollispinis pratensis</i> ) - Wherever found	T	\$3,803	\$6,300	\$10,103	\$0	\$10,103
	612	Spinedace, Little Colorado ( <i>Lepidomeda vittata</i> ) - Wherever found	T	\$13,500	\$28,575	\$42,075	\$0	\$42,075
	491	Spinedace, White River ( <i>Lepidomeda albivallis</i> ) - Wherever found	E	\$72,416	\$1,500	\$73,916	\$0	\$73,916
	585	Springfish, Hiko White River ( <i>Crenichthys baileyi grandis</i> ) - Wherever found	E	\$24,295	\$22,886	\$47,181	\$0	\$47,181

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	463	Springfish, Railroad Valley ( <i>Crenichthys nevadae</i> ) - Wherever found	T	\$74,697	\$11,500	\$86,197	\$0	\$86,197
	601	Springfish, White River ( <i>Crenichthys baileyi baileyi</i> ) - Wherever found	E	\$29,116	\$15,000	\$44,116	\$0	\$44,116
	16	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - California Central Valley DPS	T	\$65,024	\$18,899,463	\$18,964,487	\$0	\$18,964,487
	3	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Snake River Basin DPS	T	\$20,903	\$57,148,406	\$57,169,309	\$1,056,000	\$58,225,309
	47	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Central California Coast DPS	T	\$0	\$5,835,104	\$5,835,104	\$0	\$5,835,104
	15	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Lower Columbia River DPS	T	\$182,360	\$16,154,762	\$16,337,122	\$2,704,400	\$19,041,522
	68	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - South-Central California Coast DPS	T	\$0	\$3,653,700	\$3,653,700	\$0	\$3,653,700
	52	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Southern California DPS	E	\$0	\$5,396,190	\$5,396,190	\$0	\$5,396,190
	9	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Upper Columbia River DPS	T	\$139,229	\$29,601,351	\$29,740,580	\$2,741,600	\$32,482,180
	27	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Upper Willamette River DPS	T	\$78,700	\$11,984,400	\$12,063,100	\$421,000	\$12,484,100
	2	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Middle Columbia River DPS	T	\$115,300	\$59,631,969	\$59,747,269	\$1,865,400	\$61,612,669
	39	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Northern California DPS	T	\$0	\$7,689,200	\$7,689,200	\$0	\$7,689,200
	22	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Puget Sound DPS	T	\$66,001	\$10,679,817	\$10,745,818	\$2,549,000	\$13,294,818
	395	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Middle Columbia River DPS	EXPN	\$95,000	\$4,550	\$99,550	\$26,200	\$125,750
	460	Stickleback, unarmored threespine ( <i>Gasterosteus aculeatus williamsoni</i> ) - Wherever found	E	\$78,425	\$8,500	\$86,925	\$0	\$86,925
	639	Sturgeon, Alabama ( <i>Scaphirhynchus suttkusi</i> ) - Wherever found	E	\$12,000	\$26,022	\$38,022	\$0	\$38,022
	122	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - Gulf of Maine DPS	T	\$0	\$1,072,614	\$1,072,614	\$0	\$1,072,614
	136	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - New York Bight DPS	E	\$0	\$960,133	\$960,133	\$29,000	\$989,133

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	167	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - Chesapeake Bay DPS	E	\$110,000	\$601,497	\$711,497	\$0	\$711,497
	213	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - Carolina DPS	E	\$0	\$500,494	\$500,494	\$0	\$500,494
	192	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - South Atlantic DPS	E	\$0	\$501,030	\$501,030	\$75,000	\$576,030
	28	Sturgeon, green ( <i>Acipenser medirostris</i> ) - Southern DPS	T	\$2,100	\$10,831,222	\$10,833,322	\$29,600	\$10,862,922
	55	Sturgeon, Gulf ( <i>Acipenser oxyrinchus (=oxyrhynchus) desotoi</i> ) - Wherever found	T	\$279,500	\$4,527,886	\$4,807,386	\$20,000	\$4,827,386
	12	Sturgeon, pallid ( <i>Scaphirhynchus albus</i> ) - Wherever found	E	\$1,416,200	\$20,679,532	\$22,095,732	\$69,900	\$22,165,632
	100	Sturgeon, shortnose ( <i>Acipenser brevirostrum</i> ) - Wherever found	E	\$7,500	\$1,373,084	\$1,380,584	\$137,500	\$1,518,084
	552	Sturgeon, Shovelnose ( <i>Scaphirhynchus platorynchus</i> ) - Wherever found	SAT	\$529	\$31,635	\$32,164	\$23,700	\$55,864
	24	Sturgeon, white ( <i>Acipenser transmontanus</i> ) - U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	E	\$278,112	\$12,369,651	\$12,647,763	\$341,730	\$12,989,493
	518	Sucker, June ( <i>Chasmistes liorus</i> ) - Wherever found	E	\$14,000	\$50,000	\$64,000	\$0	\$64,000
	81	Sucker, Lost River ( <i>Deltistes luxatus</i> ) - Wherever found	E	\$620,776	\$1,788,900	\$2,409,676	\$16,000	\$2,425,676
	23	Sucker, razorback ( <i>Xyrauchen texanus</i> ) - Wherever found	E	\$480,118	\$12,311,856	\$12,791,974	\$217,932	\$13,009,906
	302	Sucker, Santa Ana ( <i>Catostomus santaanae</i> ) - 3 CA river basins	T	\$163,606	\$66,155	\$229,761	\$0	\$229,761
	79	Sucker, shortnose ( <i>Chasmistes brevirostris</i> ) - Wherever found	E	\$615,679	\$1,900,900	\$2,516,579	\$16,000	\$2,532,579
	223	Sucker, Warner ( <i>Catostomus warnerensis</i> ) - Wherever found	T	\$222,500	\$156,780	\$379,280	\$82,000	\$461,280
	649	Sucker, Zuni bluehead ( <i>Catostomus discobolus yarrowi</i> ) - Wherever found	E	\$2,500	\$33,500	\$36,000	\$0	\$36,000
	1258	Sunfish, spring pygmy ( <i>Elassoma alabamae</i> ) - Wherever found	T	\$500	\$0	\$500	\$0	\$500
	171	Topminnow, Gila (incl. Yaqui) ( <i>Poeciliopsis occidentalis</i> ) - Wherever found	E	\$56,000	\$623,580	\$679,580	\$0	\$679,580
	592	Totoaba (seatrout or weakfish) ( <i>Cynoscion macdonaldi</i> ) - Wherever found	E	\$46,200	\$0	\$46,200	\$0	\$46,200

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	583	Trout, Apache ( <i>Oncorhynchus apache</i> ) - Wherever found	T	\$21,500	\$26,000	\$47,500	\$0	\$47,500
	7	Trout, bull ( <i>Salvelinus confluentus</i> ) - U.S.A., conterminous, lower 48 states	T	\$7,335,866	\$29,273,292	\$36,609,158	\$3,655,500	\$40,264,658
	126	Trout, bull ( <i>Salvelinus confluentus</i> ) - Clackamas River subbasin experimental population	EXPN	\$606,500	\$0	\$606,500	\$447,600	\$1,054,100
	322	Trout, Gila ( <i>Oncorhynchus gilae</i> ) - Wherever found	T	\$99,000	\$93,957	\$192,957	\$0	\$192,957
	115	Trout, Greenback Cutthroat ( <i>Oncorhynchus clarkii stomias</i> ) - Wherever found	T	\$441,500	\$653,238	\$1,094,738	\$99,302	\$1,194,040
	35	Trout, Lahontan cutthroat ( <i>Oncorhynchus clarkii henshawi</i> ) - Wherever found	T	\$5,946,849	\$2,855,658	\$8,802,507	\$0	\$8,802,507
	631	Trout, Little Kern golden ( <i>Oncorhynchus aguabonita whitei</i> ) - Wherever found	T	\$1,210	\$37,813	\$39,023	\$0	\$39,023
	206	Trout, Paiute cutthroat ( <i>Oncorhynchus clarkii seleniris</i> ) - Wherever found	T	\$505,327	\$12,000	\$517,327	\$0	\$517,327
	493	Tui chub, Mohave ( <i>Gila bicolor</i> ssp. <i>mohavensis</i> ) - Wherever found	E	\$726	\$72,063	\$72,789	\$0	\$72,789
	287	Woundfin ( <i>Plagopterus argentissimus</i> ) - Wherever found, except where listed as an experimental population	E	\$65,317	\$195,300	\$260,617	\$0	\$260,617
<b>Fishes subtotal</b>				<b>\$32,952,611</b>	<b>\$702,913,787</b>	<b>\$735,866,398</b>	<b>\$107,649,427</b>	<b>\$843,515,825</b>
Clams	1231	Acornshell, southern ( <i>Epioblasma othcaloogensis</i> ) - Wherever found	E	\$1,500	\$542	\$2,042	\$0	\$2,042
	285	Bankclimber, purple (mussel) ( <i>Elliptioideus sloatianus</i> ) - Wherever found	T	\$162,000	\$95,240	\$257,240	\$5,000	\$262,240
	629	Bean, Choctaw ( <i>Villosa choctawensis</i> ) - Wherever found	E	\$4,000	\$35,499	\$39,499	\$0	\$39,499
	555	Bean, Cumberland (pearlymussel) ( <i>Villosa trabalis</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$32,500	\$1,443	\$33,943	\$20,000	\$53,943
	1222	Bean, Cumberland (pearlymussel) ( <i>Villosa trabalis</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$2,409	\$2,409	\$0	\$2,409
	1091	Bean, purple ( <i>Villosa perpurpurea</i> ) - Wherever found	E	\$3,600	\$3,816	\$7,416	\$0	\$7,416

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	539	Bean, rayed ( <i>Villosa fabalis</i> ) - Wherever found	E	\$23,300	\$29,888	\$53,188	\$6,166	\$59,354
	1258	Blossom, green (pearlymussel) ( <i>Epioblasma torulosa gubernaculum</i> ) - Wherever found	E	\$400	\$100	\$500	\$0	\$500
	1205	Blossom, tubercled (pearlymussel) ( <i>Epioblasma torulosa torulosa</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$1,500	\$1,631	\$3,131	\$0	\$3,131
	1256	Blossom, turgid (pearlymussel) ( <i>Epioblasma turgidula</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$700	\$0	\$700	\$0	\$700
	1265	Blossom, turgid (pearlymussel) ( <i>Epioblasma turgidula</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$250	\$250	\$0	\$250
	1086	Catspaw, white (pearlymussel) ( <i>Epioblasma obliquata perobliqua</i> ) - Wherever found	E	\$500	\$846	\$1,346	\$6,166	\$7,512
	404	Clubshell ( <i>Pleurobema clava</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$44,340	\$30,683	\$75,023	\$45,391	\$120,414
	747	Clubshell ( <i>Pleurobema clava</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$24,500	\$0	\$24,500	\$0	\$24,500
	1258	Clubshell, black ( <i>Pleurobema curtum</i> ) - Wherever found	E	\$0	\$100	\$100	\$400	\$500
	1048	Clubshell, ovate ( <i>Pleurobema perovatum</i> ) - Wherever found	E	\$3,000	\$2,711	\$5,711	\$2,900	\$8,611
	905	Clubshell, southern ( <i>Pleurobema decisum</i> ) - Wherever found	E	\$2,500	\$8,165	\$10,665	\$3,400	\$14,065
	550	Combshell, Cumberlandian ( <i>Epioblasma brevidens</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$34,500	\$1,417	\$35,917	\$20,000	\$55,917
	1252	Combshell, Cumberlandian ( <i>Epioblasma brevidens</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$1,002	\$1,002	\$0	\$1,002

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1239	Combshell, southern ( <i>Epioblasma penita</i> ) - Wherever found	E	\$0	\$100	\$100	\$1,400	\$1,500
	1174	Combshell, upland ( <i>Epioblasma metastrata</i> ) - Wherever found	E	\$1,500	\$1,896	\$3,396	\$1,000	\$4,396
	775	Ebonysell, round ( <i>Fusconaia rotulata</i> ) - Wherever found	E	\$700	\$20,648	\$21,348	\$0	\$21,348
	279	Elktoe, Appalachian ( <i>Alasmidonta raveneliana</i> ) - Wherever found	E	\$69,500	\$196,814	\$266,314	\$9,900	\$276,214
	786	Elktoe, Cumberland ( <i>Alasmidonta atropurpurea</i> ) - Wherever found	E	\$13,000	\$2,618	\$15,618	\$5,000	\$20,618
	382	Fanshell ( <i>Cyprogenia stegaria</i> ) - Wherever found	E	\$60,900	\$62,274	\$123,174	\$10,166	\$133,340
	1193	Fanshell ( <i>Cyprogenia stegaria</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$3,379	\$3,379	\$0	\$3,379
	883	Fatmucket, Arkansas ( <i>Lampsilis powellii</i> ) - Wherever found	T	\$8,000	\$3,312	\$11,312	\$4,000	\$15,312
	258	Heelsplitter, Carolina ( <i>Lasmigona decorata</i> ) - Wherever found	E	\$258,000	\$6,280	\$264,280	\$67,600	\$331,880
	607	Heelsplitter, Inflated ( <i>Potamilus inflatus</i> ) - Wherever found	T	\$36,500	\$5,934	\$42,434	\$400	\$42,834
	140	Higgins eye (pearlymussel) ( <i>Lampsilis higginsii</i> ) - Wherever found	E	\$290,200	\$645,500	\$935,700	\$15,800	\$951,500
	713	Hornshell, Texas ( <i>Popenaias popeii</i> ) - Wherever found	E	\$4,500	\$23,100	\$27,600	\$0	\$27,600
	487	Kidneyshell, fluted ( <i>Ptychobranhus subtentus</i> ) - Wherever found	E	\$15,000	\$61,953	\$76,953	\$0	\$76,953
	624	Kidneyshell, southern ( <i>Ptychobranhus jonesi</i> ) - Wherever found	E	\$5,000	\$35,157	\$40,157	\$0	\$40,157
	1055	Kidneyshell, triangular ( <i>Ptychobranhus greenii</i> ) - Wherever found	E	\$5,000	\$1,430	\$6,430	\$2,000	\$8,430
	1221	Lampmussel, Alabama ( <i>Lampsilis virescens</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1225	Lampmussel, Alabama ( <i>Lampsilis virescens</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$2,380	\$2,380	\$0	\$2,380

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	412	Lance, yellow ( <i>Elliptio lanceolata</i> ) - Wherever found	T	\$91,418	\$2,018	\$93,436	\$15,310	\$108,746
	1189	Lilliput, pale (pearlymussel) ( <i>Toxolasma cylindrellus</i> ) - Wherever found	E	\$3,500	\$100	\$3,600	\$0	\$3,600
	589	Mapleleaf, winged ( <i>Quadrula fragosa</i> ) - Wherever found, except where listed as an experimental population	E	\$25,447	\$21,188	\$46,635	\$0	\$46,635
	1052	Moccasinshell, Alabama ( <i>Medionidus acutissimus</i> ) - Wherever found	T	\$4,000	\$1,056	\$5,056	\$3,400	\$8,456
	1179	Moccasinshell, Coosa ( <i>Medionidus parvulus</i> ) - Wherever found	E	\$3,000	\$120	\$3,120	\$1,000	\$4,120
	632	Moccasinshell, Gulf ( <i>Medionidus penicillatus</i> ) - Wherever found	E	\$16,500	\$20,000	\$36,500	\$2,500	\$39,000
	668	Moccasinshell, Ochlockonee ( <i>Medionidus simpsonianus</i> ) - Wherever found	E	\$11,500	\$20,000	\$31,500	\$1,000	\$32,500
	876	Moccasinshell, Suwannee ( <i>Medionidus walkeri</i> ) - Wherever found	T	\$9,000	\$1,536	\$10,536	\$5,000	\$15,536
	1249	Monkeyface, Appalachian (pearlymussel) ( <i>Quadrula sparsa</i> ) - Wherever found	E	\$1,000	\$100	\$1,100	\$0	\$1,100
	1164	Monkeyface, Appalachian (pearlymussel) ( <i>Quadrula sparsa</i> ) - USA (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$4,500	\$425	\$4,925	\$0	\$4,925
	474	Monkeyface, Cumberland (pearlymussel) ( <i>Quadrula intermedia</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$21,000	\$60,010	\$81,010	\$0	\$81,010
	1262	Monkeyface, Cumberland (pearlymussel) ( <i>Quadrula intermedia</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$300	\$0	\$300	\$0	\$300
	1262	Monkeyface, Cumberland (pearlymussel) ( <i>Quadrula intermedia</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$300	\$0	\$300	\$0	\$300
	291	Mucket, Neosho ( <i>Lampsilis rafinesqueana</i> ) - Wherever found	E	\$118,804	\$121,842	\$240,646	\$8,800	\$249,446



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1160	Mucket, orangenacre ( <i>Lampsilis perovalis</i> ) - Wherever found	T	\$3,500	\$1,576	\$5,076	\$0	\$5,076
	283	Mucket, pink (pearlymussel) ( <i>Lampsilis abrupta</i> ) - Wherever found	E	\$191,500	\$59,204	\$250,704	\$16,166	\$266,870
	848	Mussel, oyster ( <i>Epioblasma capsaeformis</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$15,500	\$1,557	\$17,057	\$0	\$17,057
	1246	Mussel, oyster ( <i>Epioblasma capsaeformis</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$1,253	\$1,253	\$0	\$1,253
	478	Mussel, scaleshell ( <i>Leptodea leptodon</i> ) - Wherever found	E	\$61,847	\$15,683	\$77,530	\$1,700	\$79,230
	298	Mussel, sheepnose ( <i>Plethobasus cyphus</i> ) - Wherever found	E	\$146,560	\$77,404	\$223,964	\$13,666	\$237,630
	225	Mussel, snuffbox ( <i>Epioblasma triquetra</i> ) - Wherever found	E	\$332,810	\$86,656	\$419,466	\$26,266	\$445,732
	1260	Mussel, snuffbox ( <i>Epioblasma triquetra</i> ) - Wherever found	NS	\$0	\$0	\$0	\$400	\$400
	1142	Pearlshell, Alabama ( <i>Margaritifera marrianae</i> ) - Wherever found	E	\$5,500	\$290	\$5,790	\$0	\$5,790
	255	Pearlshell, Louisiana ( <i>Margaritifera hembeli</i> ) - Wherever found	T	\$329,000	\$13,600	\$342,600	\$1,000	\$343,600
	519	Pearlymussel, birdwing ( <i>Lemiox rimosus</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$4,000	\$59,900	\$63,900	\$0	\$63,900
	1261	Pearlymussel, birdwing ( <i>Lemiox rimosus</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert, and Lauderdale Counties, AL	EXPN	\$0	\$350	\$350	\$0	\$350
	514	Pearlymussel, cracking ( <i>Hemistena lata</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$5,500	\$60,210	\$65,710	\$0	\$65,710
	1264	Pearlymussel, cracking ( <i>Hemistena lata</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$251	\$251	\$0	\$251
	1195	Pearlymussel, Curtis ( <i>Epioblasma florentina curtisii</i> ) - Wherever found	E	\$2,700	\$666	\$3,366	\$0	\$3,366

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	966	Pearlymussel, dromedary ( <i>Dromus dromas</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$5,500	\$1,177	\$6,677	\$5,000	\$11,677
	1192	Pearlymussel, dromedary ( <i>Dromus dromas</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$3,382	\$3,382	\$0	\$3,382
	543	Pearlymussel, littlewing ( <i>Pegias fabula</i> ) - Wherever found	E	\$16,000	\$32,364	\$48,364	\$10,000	\$58,364
	510	Pearlymussel, slabside ( <i>Pleuroaia dolabelloides</i> ) - Wherever found	E	\$5,500	\$61,127	\$66,627	\$400	\$67,027
	1202	Pigtoe, Cumberland ( <i>Pleurobema gibberum</i> ) - Wherever found	E	\$2,500	\$726	\$3,226	\$0	\$3,226
	1209	Pigtoe, dark ( <i>Pleurobema furvum</i> ) - Wherever found	E	\$2,500	\$573	\$3,073	\$0	\$3,073
	471	Pigtoe, finerayed ( <i>Fusconaia cuneolus</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$22,500	\$60,150	\$82,650	\$0	\$82,650
	1185	Pigtoe, finerayed ( <i>Fusconaia cuneolus</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$300	\$3,508	\$3,808	\$0	\$3,808
	1262	Pigtoe, finerayed ( <i>Fusconaia cuneolus</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$300	\$0	\$300	\$0	\$300
	1260	Pigtoe, flat ( <i>Pleurobema marshalli</i> ) - Wherever found	E	\$0	\$0	\$0	\$400	\$400
	611	Pigtoe, fuzzy ( <i>Pleurobema strodeanum</i> ) - Wherever found	T	\$7,000	\$35,499	\$42,499	\$0	\$42,499
	1172	Pigtoe, Georgia ( <i>Pleurobema hanleyianum</i> ) - Wherever found	E	\$1,500	\$0	\$1,500	\$3,000	\$4,500
	1240	Pigtoe, heavy ( <i>Pleurobema taitianum</i> ) - Wherever found	E	\$700	\$764	\$1,464	\$0	\$1,464
	588	Pigtoe, narrow ( <i>Fusconaia escambia</i> ) - Wherever found	T	\$26,000	\$20,989	\$46,989	\$0	\$46,989
	363	Pigtoe, oval ( <i>Pleurobema pyriforme</i> ) - Wherever found	E	\$44,000	\$96,152	\$140,152	\$5,000	\$145,152

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	574	Pigtoe, rough ( <i>Pleurobema plenum</i> ) - Wherever found	E	\$28,500	\$19,018	\$47,518	\$2,000	\$49,518
	1161	Pigtoe, rough ( <i>Pleurobema plenum</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$2,000	\$3,006	\$5,006	\$0	\$5,006
	472	Pigtoe, shiny ( <i>Fusconaia cor</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$22,000	\$60,250	\$82,250	\$0	\$82,250
	1204	Pigtoe, shiny ( <i>Fusconaia cor</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$300	\$2,881	\$3,181	\$0	\$3,181
	1262	Pigtoe, shiny ( <i>Fusconaia cor</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$300	\$0	\$300	\$0	\$300
	1029	Pigtoe, southern ( <i>Pleurobema georgianum</i> ) - Wherever found	E	\$3,500	\$829	\$4,329	\$5,000	\$9,329
	780	Pigtoe, tapered ( <i>Fusconaia burkei</i> ) - Wherever found	T	\$6,500	\$14,510	\$21,010	\$0	\$21,010
	538	Pimpleback, orangefoot (pearlymussel) ( <i>Plethobasus cooperianus</i> ) - Wherever found	E	\$33,000	\$23,170	\$56,170	\$3,529	\$59,699
	1163	Pimpleback, orangefoot (pearlymussel) ( <i>Plethobasus cooperianus</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$0	\$0	\$5,000	\$5,000
	309	Pocketbook, fat ( <i>Potamilus capax</i> ) - Wherever found	E	\$74,500	\$144,504	\$219,004	\$0	\$219,004
	868	Pocketbook, finelined ( <i>Lampsilis altilis</i> ) - Wherever found	T	\$4,500	\$6,263	\$10,763	\$5,000	\$15,763
	476	Pocketbook, Ouachita rock ( <i>Arkansia wheeleri</i> ) - Wherever found	E	\$72,963	\$5,588	\$78,551	\$1,600	\$80,151
	610	Pocketbook, shinyrayed ( <i>Lampsilis subangulata</i> ) - Wherever found	E	\$17,500	\$20,000	\$37,500	\$5,000	\$42,500
	1201	Pocketbook, speckled ( <i>Lampsilis streckeri</i> ) - Wherever found	E	\$0	\$100	\$100	\$3,200	\$3,300

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	704	Purple Cat's paw (=Purple Cat's paw pearlymussel) ( <i>Epioblasma obliquata obliquata</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$10,000	\$12,654	\$22,654	\$6,166	\$28,820
	1264	Purple Cat's paw (=Purple Cat's paw pearlymussel) ( <i>Epioblasma obliquata obliquata</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$251	\$251	\$0	\$251
	286	Rabbitsfoot ( <i>Quadrula cylindrica cylindrica</i> ) - Wherever found	T	\$160,716	\$93,147	\$253,863	\$7,866	\$261,729
	1260	Rabbitsfoot ( <i>Quadrula cylindrica cylindrica</i> ) -	NS	\$0	\$0	\$0	\$400	\$400
	1099	Rabbitsfoot, rough ( <i>Quadrula cylindrica strigillata</i> ) - Wherever found	E	\$4,000	\$3,106	\$7,106	\$0	\$7,106
	343	Riffleshell, northern ( <i>Epioblasma torulosa rangiana</i> ) - Wherever found	E	\$120,555	\$33,865	\$154,420	\$7,166	\$161,586
	541	Riffleshell, tan ( <i>Epioblasma florentina walkeri</i> (=E. walkeri)) - Wherever found	E	\$47,600	\$1,183	\$48,783	\$10,000	\$58,783
	580	Ring pink (mussel) ( <i>Obovaria retusa</i> ) - Wherever found	E	\$26,000	\$20,180	\$46,180	\$2,000	\$48,180
	602	Sandshell, Southern ( <i>Hamiota australis</i> ) - Wherever found	T	\$8,000	\$35,499	\$43,499	\$0	\$43,499
	392	Slabshell, Chipola ( <i>Elliptio chipolaensis</i> ) - Wherever found	T	\$32,000	\$95,240	\$127,240	\$0	\$127,240
	339	Spectaclecase (mussel) ( <i>Cumberlandia monodonta</i> ) - Wherever found	E	\$80,310	\$55,956	\$136,266	\$28,800	\$165,066
	1232	Spinymussel, Altamaha ( <i>Elliptio spinosa</i> ) - Wherever found	E	\$0	\$0	\$0	\$2,000	\$2,000
	200	Spinymussel, James ( <i>Pleurobema collina</i> ) - Wherever found	E	\$535,120	\$2,010	\$537,130	\$2,100	\$539,230
	750	Spinymussel, Tar River ( <i>Elliptio steinstansana</i> ) - Wherever found	E	\$13,000	\$6,125	\$19,125	\$4,900	\$24,025
	1260	Stirrupshell ( <i>Quadrula stapes</i> ) - Wherever found	E	\$0	\$0	\$0	\$400	\$400
	346	Threeridge, fat (mussel) ( <i>Amblyma neislerii</i> ) - Wherever found	E	\$56,500	\$95,240	\$151,740	\$7,529	\$159,269

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1196	Wartyback, white (pearlymussel) ( <i>Plethobasus cicatricosus</i> ) - Wherever found	E	\$3,000	\$351	\$3,351	\$0	\$3,351
	180	Wedgemussel, dwarf ( <i>Alasmidonta heterodon</i> ) - Wherever found	E	\$577,835	\$34,864	\$612,699	\$7,400	\$620,099
<b>Clams subtotal</b>				<b>\$4,592,325</b>	<b>\$3,021,713</b>	<b>\$7,614,038</b>	<b>\$463,753</b>	<b>\$8,077,791</b>
Snails	74	Abalone, Black ( <i>Haliotis cracherodii</i> ) - Wherever found	E	\$0	\$2,699,460	\$2,699,460	\$0	\$2,699,460
	133	Abalone, White ( <i>Haliotis sorenseni</i> ) - North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	E	\$0	\$1,003,000	\$1,003,000	\$0	\$1,003,000
	1198	Ambersnail, Kanab ( <i>Oxyloma haydeni kanabensis</i> ) - Wherever found	E	\$0	\$3,333	\$3,333	\$0	\$3,333
	1183	Campeloma, slender ( <i>Campeloma decampi</i> ) - Wherever found	E	\$3,000	\$997	\$3,997	\$0	\$3,997
	691	Cavesnail, Tumbling Creek ( <i>Antrobia culveri</i> ) - Wherever found	E	\$0	\$24,000	\$24,000	\$6,000	\$30,000
	856	Elimia, lacy (snail) ( <i>Elimia crenatella</i> ) - Wherever found	T	\$16,500	\$0	\$16,500	\$0	\$16,500
	862	Hornsnail, rough ( <i>Pleurocera foremani</i> ) - Wherever found	E	\$15,000	\$1,196	\$16,196	\$0	\$16,196
	385	Limpet, Banbury Springs ( <i>Lanx</i> sp.) - Wherever found	E	\$120,345	\$9,100	\$129,445	\$0	\$129,445
	1207	Lioplax, cylindrical (snail) ( <i>Lioplax cyclostomaformis</i> ) - Wherever found	E	\$2,500	\$622	\$3,122	\$0	\$3,122
	951	Marstonia, royal (snail) ( <i>Pyrgulopsis ogmorhapse</i> ) - Wherever found	E	\$12,000	\$125	\$12,125	\$0	\$12,125
	927	[no common name] Snail ( <i>Eua zebrina</i> ) - Wherever found	E	\$13,223	\$0	\$13,223	\$0	\$13,223
	927	[no common name] Snail ( <i>Ostodes strigatus</i> ) - Wherever found	E	\$13,223	\$0	\$13,223	\$0	\$13,223
	1268	Pebblesnail, flat ( <i>Lepyrium showalteri</i> ) - Wherever found	E	\$0	\$149	\$149	\$0	\$149
	735	Riversnail, Anthony's ( <i>Athearnia anthonyi</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$25,500	\$0	\$25,500	\$0	\$25,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1177	Riversnail, Anthony's ( <i>Athearnia anthonyi</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$4,254	\$4,254	\$0	\$4,254
	1077	Rocksnailed, interrupted (=Georgia) ( <i>Leptoxis foremani</i> ) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	947	Rocksnailed, painted ( <i>Leptoxis taeniata</i> ) - Wherever found	T	\$12,000	\$396	\$12,396	\$0	\$12,396
	1003	Rocksnailed, plicate ( <i>Leptoxis plicata</i> ) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	546	Rocksnailed, round ( <i>Leptoxis ampla</i> ) - Wherever found	T	\$56,500	\$597	\$57,097	\$0	\$57,097
	1254	Snail, armored ( <i>Pyrgulopsis</i> (=Marstonia) <i>pachyta</i> ) - Wherever found	E	\$800	\$0	\$800	\$0	\$800
	429	Snail, Bliss Rapids ( <i>Taylorconcha serpenticola</i> ) - Wherever found	T	\$85,066	\$13,650	\$98,716	\$0	\$98,716
	422	Snail, Chittenango ovate amber ( <i>Succinea chittenangoensis</i> ) - Wherever found	T	\$100,000	\$0	\$100,000	\$800	\$100,800
	480	Snail, fragile tree ( <i>Samoana fragilis</i> ) - Wherever found	E	\$60,797	\$18,107	\$78,904	\$0	\$78,904
	413	Snail, Guam tree ( <i>Partula radiolata</i> ) - Wherever found	E	\$60,797	\$47,654	\$108,451	\$0	\$108,451
	415	Snail, Humped tree ( <i>Partula gibba</i> ) - Wherever found	E	\$62,825	\$44,094	\$106,919	\$0	\$106,919
	587	Snail, Iowa Pleistocene ( <i>Discus macclintocki</i> ) - Wherever found	E	\$47,000	\$0	\$47,000	\$0	\$47,000
	693	Snail, Lanai tree ( <i>Partulina semicarinata</i> ) - Wherever found	E	\$29,956	\$0	\$29,956	\$0	\$29,956
	693	Snail, Lanai tree ( <i>Partulina variabilis</i> ) - Wherever found	E	\$29,956	\$0	\$29,956	\$0	\$29,956
	634	Snail, Langford's tree ( <i>Partula langfordi</i> ) - Wherever found	E	\$38,938	\$0	\$38,938	\$0	\$38,938
	358	Snail, Morro shoulderband (=Banded dune) ( <i>Helminthoglypta walkeriana</i> ) - Wherever found	E	\$146,137	\$2,818	\$148,955	\$0	\$148,955
	936	Snail, Newcomb's ( <i>Erinna newcombi</i> ) - Wherever found	T	\$12,797	\$0	\$12,797	\$0	\$12,797
	645	Snail, nooday ( <i>Mesodon clarki nantahala</i> ) - Wherever found	T	\$37,000	\$0	\$37,000	\$0	\$37,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1212	Snail, painted snake coiled forest ( <i>Anguispira picta</i> ) - Wherever found	T	\$2,500	\$400	\$2,900	\$0	\$2,900
	866	Snail, Pecos assiminea ( <i>Assiminea pecos</i> ) - Wherever found	E	\$13,250	\$2,600	\$15,850	\$0	\$15,850
	315	Snail, Snake River physa ( <i>Physa natricina</i> ) - Wherever found	E	\$128,877	\$76,461	\$205,338	\$0	\$205,338
	554	Snails, Oahu tree ( <i>Achatinella</i> spp.) - Wherever found	E*	\$55,191	\$0	\$55,191	\$0	\$55,191
	692	Snail, Stock Island tree ( <i>Orthalicus reses</i> (not incl. <i>nesodryas</i> )) - Wherever found	T	\$28,500	\$1,484	\$29,984	\$0	\$29,984
	1219	Snail, tulotoma ( <i>Tulotoma magnifica</i> ) - Wherever found	T	\$2,000	\$545	\$2,545	\$0	\$2,545
	1232	Snail, Virginia fringed mountain ( <i>Polygyris virginiensis</i> ) - Wherever found	E	\$2,000	\$0	\$2,000	\$0	\$2,000
	1224	Springsnail, Alamosa ( <i>Tryonia alamosae</i> ) - Wherever found	E	\$2,400	\$0	\$2,400	\$0	\$2,400
	457	Springsnail, Bruneau Hot ( <i>Pyrgulopsis bruneauensis</i> ) - Wherever found	E	\$83,919	\$2,600	\$86,519	\$815	\$87,334
	1269	Springsnail, Chupadera ( <i>Pyrgulopsis chupaderae</i> ) - Wherever found	E	\$125	\$0	\$125	\$0	\$125
	891	Springsnail, Koster's ( <i>Juturnia kosteri</i> ) - Wherever found	E	\$12,250	\$2,500	\$14,750	\$0	\$14,750
	1221	Springsnail, Phantom ( <i>Pyrgulopsis texana</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	890	Springsnail, Roswell ( <i>Pyrgulopsis roswellensis</i> ) - Wherever found	E	\$12,250	\$2,600	\$14,850	\$0	\$14,850
	1243	Springsnail, San Bernardino ( <i>Pyrgulopsis bernardina</i> ) - Wherever found	T	\$1,300	\$100	\$1,400	\$0	\$1,400
	1270	Springsnail, Socorro ( <i>Pyrgulopsis neomexicana</i> ) - Wherever found	E	\$0	\$100	\$100	\$0	\$100
	1101	Springsnail, Three Forks ( <i>Pyrgulopsis trivialis</i> ) - Wherever found	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	744	Tree snail, Newcomb's ( <i>Newcombia cumingi</i> ) - Wherever found	E	\$24,659	\$0	\$24,659	\$0	\$24,659
	1253	Tryonia, Diamond ( <i>Pseudotryonia adamantina</i> ) - Wherever found	E	\$1,000	\$0	\$1,000	\$0	\$1,000
<b>Snails subtotal</b>				<b>\$1,399,581</b>	<b>\$3,962,942</b>	<b>\$5,362,523</b>	<b>\$7,615</b>	<b>\$5,370,138</b>

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
Insects	123	Beetle, American burying ( <i>Nicrophorus americanus</i> ) - Wherever found, except where listed as an experimental population	E	\$556,984	\$346,663	\$903,647	\$164,466	\$1,068,113
	548	Beetle, American burying ( <i>Nicrophorus americanus</i> ) - Ex Pop, SW Missouri	EXPN	\$47,500	\$9,200	\$56,700	\$0	\$56,700
	977	Beetle, American burying ( <i>Nicrophorus americanus</i> ) - Ex-Pop, SW Missouri	NS	\$11,100	\$0	\$11,100	\$0	\$11,100
	248	Beetle, Casey's June ( <i>Dinacoma caseyi</i> ) - Wherever found	E	\$360,014	\$8,052	\$368,066	\$0	\$368,066
	1221	Beetle, Coffin Cave mold ( <i>Batrisodes texanus</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	865	Beetle, Comal Springs dryopid ( <i>Stygoparnus comalensis</i> ) - Wherever found	E	\$10,000	\$0	\$10,000	\$5,972	\$15,972
	983	Beetle, Comal Springs riffle ( <i>Heterelmis comalensis</i> ) - Wherever found	E	\$10,000	\$840	\$10,840	\$0	\$10,840
	812	Beetle, delta green ground ( <i>Elaphrus viridis</i> ) - Wherever found	T	\$15,061	\$4,100	\$19,161	\$0	\$19,161
	1221	Beetle, Helotes mold ( <i>Batrisodes venyivi</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	798	Beetle, Hungerford's crawling water ( <i>Brychius hungerfordi</i> ) - Wherever found	E	\$20,000	\$0	\$20,000	\$0	\$20,000
	1221	Beetle, Kretschmarr Cave mold ( <i>Texamaurops reddelli</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	506	Beetle, Mount Hermon June ( <i>Polyphylla barbata</i> ) - Wherever found	E	\$68,697	\$0	\$68,697	\$0	\$68,697
	1163	Beetle, [no common name] ( <i>Rhadine exilis</i> ) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1163	Beetle, [no common name] ( <i>Rhadine infernalis</i> ) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1221	Beetle, Tooth Cave ground ( <i>Rhadine persephone</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	162	Beetle, valley elderberry longhorn ( <i>Desmocerus californicus dimorphus</i> ) - Wherever found	T	\$468,658	\$281,970	\$750,628	\$0	\$750,628
	218	Bumble bee, Rusty patched ( <i>Bombus affinis</i> ) - Wherever found	E	\$419,154	\$32,100	\$451,254	\$30,000	\$481,254
	507	Butterfly, Bartram's hairstreak ( <i>Strymon acis bartrami</i> ) - Wherever found	E	\$38,500	\$30,000	\$68,500	\$0	\$68,500



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	295	Butterfly, bay checkerspot ( <i>Euphydryas editha bayensis</i> ) - Wherever found	T	\$40,425	\$203,500	\$243,925	\$0	\$243,925
	594	Butterfly, Behren's silverspot ( <i>Speyeria zerene behrensii</i> ) - Wherever found	E	\$39,039	\$6,500	\$45,539	\$0	\$45,539
	513	Butterfly, callippe silverspot ( <i>Speyeria callippe callippe</i> ) - Wherever found	E	\$61,000	\$5,600	\$66,600	\$0	\$66,600
	613	Butterfly, El Segundo blue ( <i>Euphilotes battoides allyni</i> ) - Wherever found	E	\$42,042	\$0	\$42,042	\$0	\$42,042
	148	Butterfly, Fender's blue ( <i>Icaricia icarioides fenderi</i> ) - Wherever found	E	\$642,560	\$265,169	\$907,729	\$0	\$907,729
	557	Butterfly, Florida leafwing ( <i>Anaea troglodyta floralis</i> ) - Wherever found	E	\$23,500	\$30,000	\$53,500	\$0	\$53,500
	159	Butterfly, Karner blue ( <i>Lycacides melissa samuelis</i> ) - Wherever found	E	\$131,400	\$425,528	\$556,928	\$217,654	\$774,582
	327	Butterfly, Lange's metalmark ( <i>Apodemia mormo langei</i> ) - Wherever found	E	\$184,819	\$900	\$185,719	\$0	\$185,719
	1221	Butterfly, lotis blue ( <i>Lycacides argyrognomon lotis</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	389	Butterfly, Mariana eight-spot ( <i>Hypolimnias octocula marianensis</i> ) - Wherever found	E	\$67,945	\$59,749	\$127,694	\$0	\$127,694
	361	Butterfly, Mariana wandering ( <i>Vagrans egistina</i> ) - Wherever found	E	\$140,965	\$4,927	\$145,892	\$0	\$145,892
	614	Butterfly, Miami Blue ( <i>Cyclargus (=Hemiargus) thomasi bethunebakeri</i> ) - Wherever found	E	\$41,500	\$100	\$41,600	\$0	\$41,600
	340	Butterfly, mission blue ( <i>Icaricia icarioides missionensis</i> ) - Wherever found	E	\$36,036	\$126,600	\$162,636	\$0	\$162,636
	323	Butterfly, Mitchell's satyr ( <i>Neonympha mitchellii mitchellii</i> ) - Wherever found	E	\$179,000	\$9,755	\$188,755	\$2,300	\$191,055
	489	Butterfly, Mount Charleston blue ( <i>Icaricia (Plebejus) shasta charlestonensis</i> ) - Wherever found	E	\$74,725	\$0	\$74,725	\$0	\$74,725
	954	Butterfly, Myrtle's silverspot ( <i>Speyeria zerene myrtleae</i> ) - Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	204	Butterfly, Oregon silverspot ( <i>Speyeria zerene hippolyta</i> ) - Wherever found	T	\$462,240	\$66,671	\$528,911	\$0	\$528,911
	1037	Butterfly, Oregon silverspot ( <i>Speyeria zerene hippolyta</i> ) -	EXPN	\$4,903	\$0	\$4,903	\$4,000	\$8,903

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	529	Butterfly, Palos Verdes blue ( <i>Glaucopsyche lygdamus palosverdesensis</i> ) - Wherever found	E	\$1,623	\$60,522	\$62,145	\$0	\$62,145
	129	Butterfly, Quino checkerspot ( <i>Euphydryas editha quino</i> (=E. e. <i>wrighti</i> )) - Wherever found	E	\$152,000	\$860,532	\$1,012,532	\$0	\$1,012,532
	269	Butterfly, Saint Francis' satyr ( <i>Neonympha mitchellii francisci</i> ) - Wherever found	E	\$8,000	\$288,585	\$296,585	\$0	\$296,585
	700	Butterfly, San Bruno elfin ( <i>Callophrys mossii bayensis</i> ) - Wherever found	E	\$21,714	\$7,500	\$29,214	\$0	\$29,214
	619	Butterfly, Schaus swallowtail ( <i>Heraclides aristodemus ponceanus</i> ) - Wherever found	E	\$29,000	\$12,004	\$41,004	\$0	\$41,004
	462	Butterfly, Smith's blue ( <i>Euphilotes enoptes smithi</i> ) - Wherever found	E	\$79,800	\$6,461	\$86,261	\$0	\$86,261
	797	Butterfly, Uncompahgre fritillary ( <i>Boloria acrocneuma</i> ) - Wherever found	E	\$8,600	\$11,600	\$20,200	\$0	\$20,200
	208	Checkerspot, Taylor's (=whulge) ( <i>Euphydryas editha taylori</i> ) - Wherever found	E	\$431,220	\$16,682	\$447,902	\$64,000	\$511,902
	1096	Damselfly, blackline Hawaiian ( <i>Megalagrion nigrohamatum nigrolineatum</i> ) - Wherever found	E	\$7,270	\$0	\$7,270	\$0	\$7,270
	1121	Damselfly, crimson Hawaiian ( <i>Megalagrion leptodemas</i> ) - Wherever found	E	\$6,409	\$0	\$6,409	\$0	\$6,409
	1108	Damselfly, flying earwig Hawaiian ( <i>Megalagrion nesiotes</i> ) - Wherever found	E	\$6,692	\$0	\$6,692	\$0	\$6,692
	1127	Damselfly, oceanic Hawaiian ( <i>Megalagrion oceanicum</i> ) - Wherever found	E	\$6,256	\$0	\$6,256	\$0	\$6,256
	767	Damselfly, orangeblack Hawaiian ( <i>Megalagrion xanthomelas</i> ) - Wherever found	E	\$22,576	\$200	\$22,776	\$0	\$22,776
	881	Damselfly, Pacific Hawaiian ( <i>Megalagrion pacificum</i> ) - Wherever found	E	\$10,475	\$5,000	\$15,475	\$0	\$15,475
	532	Damselfly, Rota blue ( <i>Ischnura luta</i> ) - Wherever found	E	\$61,836	\$0	\$61,836	\$0	\$61,836
	234	Dragonfly, Hine's emerald ( <i>Somatochlora hineana</i> ) - Wherever found	E	\$411,400	\$20,150	\$431,550	\$0	\$431,550

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	85	Fly, Delhi Sands flower-loving ( <i>Rhaphiomidas terminatus abdominalis</i> ) - Wherever found	E	\$2,163,525	\$0	\$2,163,525	\$0	\$2,163,525
	1125	Fly, Hawaiian picture-wing ( <i>Drosophila aglaia</i> ) - Wherever found	E	\$6,320	\$0	\$6,320	\$0	\$6,320
	944	Fly, Hawaiian picture-wing ( <i>Drosophila differens</i> ) - Wherever found	E	\$12,503	\$0	\$12,503	\$0	\$12,503
	1119	Fly, Hawaiian picture-wing ( <i>Drosophila digressa</i> ) - Wherever found	E	\$6,419	\$0	\$6,419	\$0	\$6,419
	447	Fly, Hawaiian picture-wing ( <i>Drosophila hemipeza</i> ) - Wherever found	E	\$94,284	\$0	\$94,284	\$0	\$94,284
	1089	Fly, Hawaiian picture-wing ( <i>Drosophila heteroneura</i> ) - Wherever found	E	\$7,433	\$0	\$7,433	\$0	\$7,433
	974	Fly, Hawaiian picture-wing ( <i>Drosophila mulli</i> ) - Wherever found	T	\$11,322	\$0	\$11,322	\$0	\$11,322
	1110	Fly, Hawaiian picture-wing ( <i>Drosophila musaphilia</i> ) - Wherever found	E	\$5,983	\$632	\$6,615	\$0	\$6,615
	1125	Fly, Hawaiian picture-wing ( <i>Drosophila obatai</i> ) - Wherever found	E	\$6,320	\$0	\$6,320	\$0	\$6,320
	1119	Fly, Hawaiian picture-wing ( <i>Drosophila ochrobasis</i> ) - Wherever found	E	\$6,419	\$0	\$6,419	\$0	\$6,419
	1107	Fly, Hawaiian picture-wing ( <i>Drosophila sharpi</i> ) - Wherever found	E	\$6,062	\$632	\$6,694	\$0	\$6,694
	1125	Fly, Hawaiian picture-wing ( <i>Drosophila substenoptera</i> ) - Wherever found	E	\$6,320	\$0	\$6,320	\$0	\$6,320
	1125	Fly, Hawaiian picture-wing ( <i>Drosophila tarphytrichia</i> ) - Wherever found	E	\$6,320	\$0	\$6,320	\$0	\$6,320
	1228	Grasshopper, Zayante band-winged ( <i>Trimerotropis infantilis</i> ) - Wherever found	E	\$2,200	\$0	\$2,200	\$0	\$2,200
	326	Moth, Blackburn's sphinx ( <i>Manduca blackburni</i> ) - Wherever found	E	\$64,920	\$121,700	\$186,620	\$0	\$186,620
	838	Moth, Kern primrose sphinx ( <i>Euproserpinus euterpe</i> ) - Wherever found	T	\$2,200	\$15,368	\$17,568	\$0	\$17,568
	757	Naucorid, Ash Meadows ( <i>Ambrysus amargosus</i> ) - Wherever found	T	\$23,400	\$0	\$23,400	\$0	\$23,400
	1125	Picture-wing fly, Hawaiian ( <i>Drosophila montgomeryi</i> ) - Wherever found	E	\$6,320	\$0	\$6,320	\$0	\$6,320
	971	Picture-wing fly, Hawaiian ( <i>Drosophila neoclavisetae</i> ) - Wherever found	E	\$11,489	\$0	\$11,489	\$0	\$11,489

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	492	Skipper, Carson wandering (Pseudocopacodes eunus obscurus) - Wherever found	E	\$28,000	\$45,075	\$73,075	\$0	\$73,075
	280	Skipper, Dakota (Hesperia dacotae) - Wherever found	T	\$91,200	\$122,350	\$213,550	\$62,100	\$275,650
	1003	Skipper, Laguna Mountains (Pyrgus ruralis lagunae) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	219	Skipperling, Poweshiek (Oarisma poweshiek) - Wherever found	E	\$427,700	\$34,300	\$462,000	\$14,980	\$476,980
	1166	Skipper, Pawnee montane (Hesperia leonardus montana) - Wherever found	T	\$4,700	\$110	\$4,810	\$0	\$4,810
	1190	Tiger beetle, Miami (Cicindelidia floridana) - Wherever found	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	466	Tiger beetle, Northeastern beach (Cicindela dorsalis dorsalis) - Wherever found	T	\$81,432	\$3,868	\$85,300	\$0	\$85,300
	571	Tiger beetle, Ohlone (Cicindela ohlone) - Wherever found	E	\$49,890	\$0	\$49,890	\$0	\$49,890
	406	Tiger beetle, Puritan (Cicindela puritana) - Wherever found	T	\$105,349	\$7,240	\$112,589	\$0	\$112,589
	408	Tiger beetle, Salt Creek (Cicindela nevadica lincolniana) - Wherever found	E	\$55,300	\$30,617	\$85,917	\$25,000	\$110,917
	526	Yellow-faced bee, anthricinan (Hylaeus anthracinus) - Wherever found	E	\$62,420	\$200	\$62,620	\$0	\$62,620
	714	Yellow-faced bee, assimulans (Hylaeus assimulans) - Wherever found	E	\$27,592	\$0	\$27,592	\$0	\$27,592
	1042	Yellow-faced bee, easy (Hylaeus facilis) - Wherever found	E	\$8,784	\$0	\$8,784	\$0	\$8,784
	1125	Yellow-faced bee, Hawaiian (Hylaeus kuakea) - Wherever found	E	\$6,320	\$0	\$6,320	\$0	\$6,320
	1042	Yellow-faced bee, Hawaiian (Hylaeus longiceps) - Wherever found	E	\$8,784	\$0	\$8,784	\$0	\$8,784
	1125	Yellow-faced bee, Hawaiian (Hylaeus mana) - Wherever found	E	\$6,320	\$0	\$6,320	\$0	\$6,320
	1119	Yellow-faced bee, hilaris (Hylaeus hilaris) - Wherever found	E	\$6,419	\$0	\$6,419	\$0	\$6,419
<b>Insects subtotal</b>				<b>\$8,930,607</b>	<b>\$3,589,252</b>	<b>\$12,519,859</b>	<b>\$590,472</b>	<b>\$13,110,331</b>
Arachnids	1221	Harvestman, Bee Creek Cave (Texella reddelli) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1003	Harvestman, Bone Cave ( <i>Texella reyesi</i> ) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1221	Meshweaver, Braken Bat Cave ( <i>Cicurina venii</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1221	Meshweaver, Government Canyon Bat Cave ( <i>Cicurina vespera</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1221	Meshweaver, Madla Cave ( <i>Cicurina madla</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1221	Meshweaver, Robber Baron Cave ( <i>Cicurina baronia</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1221	Pseudoscorpion, Tooth Cave ( <i>Tartarocreagris texana</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1221	Spider, Government Canyon Bat Cave ( <i>Neoleptoneta microps</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	871	Spider, Kauai cave wolf or pe'e pe'e maka 'ole ( <i>Adelocosa anops</i> ) - Wherever found	E	\$15,737	\$0	\$15,737	\$0	\$15,737
	582	Spider, spruce-fir moss ( <i>Microhexura montivaga</i> ) - Wherever found	E	\$43,500	\$4,265	\$47,765	\$0	\$47,765
	1221	Spider, Tooth Cave ( <i>Neoleptoneta myopica</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
<b>Arachnids subtotal</b>				<b>\$89,237</b>	<b>\$4,265</b>	<b>\$93,502</b>	<b>\$0</b>	<b>\$93,502</b>
Crustaceans	1221	Amphipod, diminutive ( <i>Gammarus hyalleloides</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1266	Amphipod, Hay's Spring ( <i>Stygobromus hayi</i> ) - Wherever found	E	\$0	\$240	\$240	\$0	\$240
	1003	Amphipod, Illinois cave ( <i>Gammarus acherondytes</i> ) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	826	Amphipod, Kauai cave ( <i>Spelaeorchestia koloana</i> ) - Wherever found	E	\$18,360	\$0	\$18,360	\$0	\$18,360
	886	Amphipod, Noel's ( <i>Gammarus desperatus</i> ) - Wherever found	E	\$12,250	\$3,000	\$15,250	\$0	\$15,250
	1221	Amphipod, Peck's cave ( <i>Stygobromus (=Stygonectes) pecki</i> ) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1163	Crayfish, Benton County cave ( <i>Cambarus aculabrum</i> ) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	402	Crayfish, Big Sandy ( <i>Cambarus callainus</i> ) - Wherever found	T	\$116,012	\$7,486	\$123,498	\$0	\$123,498
	525	Crayfish, Guyandotte River ( <i>Cambarus veteranus</i> ) - Wherever found	E	\$62,648	\$0	\$62,648	\$0	\$62,648

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	656	Crayfish, Hell Creek Cave ( <i>Cambarus zophonastes</i> ) - Wherever found	E	\$5,000	\$30,000	\$35,000	\$0	\$35,000
	799	Crayfish, Nashville ( <i>Orconectes shoupi</i> ) - Wherever found	E	\$16,500	\$3,442	\$19,942	\$0	\$19,942
	879	Crayfish, Shasta ( <i>Pacifastacus fortis</i> ) - Wherever found	E	\$10,395	\$5,100	\$15,495	\$0	\$15,495
	278	Fairy shrimp, Conservancy ( <i>Branchinecta conservatio</i> ) - Wherever found	E	\$161,855	\$114,899	\$276,754	\$0	\$276,754
	581	Fairy shrimp, longhorn ( <i>Branchinecta longiantenna</i> ) - Wherever found	E	\$29,856	\$17,959	\$47,815	\$0	\$47,815
	311	Fairy shrimp, Riverside ( <i>Streptocephalus woottoni</i> ) - Wherever found	E	\$23,095	\$192,500	\$215,595	\$0	\$215,595
	239	Fairy shrimp, San Diego ( <i>Branchinecta sandiegonensis</i> ) - Wherever found	E	\$87,170	\$325,223	\$412,393	\$0	\$412,393
	124	Fairy shrimp, vernal pool ( <i>Branchinecta lynchi</i> ) - Wherever found	T	\$652,792	\$410,109	\$1,062,901	\$0	\$1,062,901
	970	Isopod, Lee County cave ( <i>Lirceus usdagalun</i> ) - Wherever found	E	\$6,500	\$5,000	\$11,500	\$0	\$11,500
	522	Isopod, Madison Cave ( <i>Antrolana lira</i> ) - Wherever found	T	\$62,843	\$240	\$63,083	\$0	\$63,083
	952	Isopod, Socorro ( <i>Thermosphaeroma thermophilus</i> ) - Wherever found	E	\$12,000	\$100	\$12,100	\$0	\$12,100
	577	Shrimp, Alabama cave ( <i>Palaemonias alabamae</i> ) - Wherever found	E	\$8,000	\$41,000	\$49,000	\$0	\$49,000
	852	Shrimp, anchialine pool ( <i>Procaris hawaiana</i> ) - Wherever found	E	\$16,474	\$200	\$16,674	\$0	\$16,674
	861	Shrimp, anchialine pool ( <i>Vetericaris chaceorum</i> ) - Wherever found	E	\$16,254	\$0	\$16,254	\$0	\$16,254
	722	Shrimp, California freshwater ( <i>Syncaris pacifica</i> ) - Wherever found	E	\$12,012	\$14,700	\$26,712	\$0	\$26,712
	661	Shrimp, Kentucky cave ( <i>Palaemonias ganteri</i> ) - Wherever found	E	\$3,500	\$30,000	\$33,500	\$0	\$33,500
	160	Tadpole shrimp, vernal pool ( <i>Lepidurus packardi</i> ) - Wherever found	E	\$532,516	\$237,999	\$770,515	\$0	\$770,515
<b>Crustaceans subtotal</b>				<b>\$1,886,032</b>	<b>\$1,439,197</b>	<b>\$3,325,229</b>	<b>\$0</b>	<b>\$3,325,229</b>
Corals	203	Coral, Boulder star ( <i>Orbicella franksi</i> ) - Wherever found	T	\$0	\$528,942	\$528,942	\$0	\$528,942
	188	Coral, elkhorn ( <i>Acropora palmata</i> ) - Wherever found	T	\$0	\$592,097	\$592,097	\$0	\$592,097

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	197	Coral, Lobed Star ( <i>Orbicella annularis</i> ) - Wherever found	T	\$0	\$554,942	\$554,942	\$0	\$554,942
	196	Coral, Mountainous Star ( <i>Orbicella faveolata</i> ) - Wherever found	T	\$0	\$558,642	\$558,642	\$0	\$558,642
	524	Coral, Pillar ( <i>Dendrogyra cylindrus</i> ) - Wherever found	T	\$0	\$62,942	\$62,942	\$0	\$62,942
	214	Coral, Pillar ( <i>Dendrogyra cylindrus</i> ) - Wherever found	NS	\$0	\$497,000	\$497,000	\$0	\$497,000
	197	Coral, Rough Cactus ( <i>Mycetophyllia ferox</i> ) - Wherever found	T	\$0	\$554,942	\$554,942	\$0	\$554,942
	179	Coral, staghorn ( <i>Acropora cervicornis</i> ) - Wherever found	T	\$0	\$630,397	\$630,397	\$0	\$630,397
	187	No Common Name ( <i>Acropora globiceps</i> ) - Wherever found	T	\$0	\$594,055	\$594,055	\$0	\$594,055
	201	No Common Name ( <i>Acropora jacquelinae</i> ) - Wherever found	T	\$0	\$538,000	\$538,000	\$0	\$538,000
	195	No Common Name ( <i>Acropora retusa</i> ) - Wherever found	T	\$0	\$561,338	\$561,338	\$0	\$561,338
	199	No Common Name ( <i>Acropora speciosa</i> ) - Wherever found	T	\$0	\$545,217	\$545,217	\$0	\$545,217
	201	No Common Name ( <i>Euphyllia paradivisa</i> ) - Wherever found	T	\$0	\$538,000	\$538,000	\$0	\$538,000
	201	No Common Name ( <i>Isopora crateriformis</i> ) - Wherever found	T	\$0	\$538,000	\$538,000	\$0	\$538,000
	677	No Common Name ( <i>Pavona diffluens</i> ) - Wherever found	T	\$0	\$31,467	\$31,467	\$0	\$31,467
	297	No Common Name ( <i>Seriatopora aculeata</i> ) - Wherever found	T	\$0	\$242,055	\$242,055	\$0	\$242,055
<b>Corals subtotal</b>				<b>\$0</b>	<b>\$7,568,036</b>	<b>\$7,568,036</b>	<b>\$0</b>	<b>\$7,568,036</b>
Flowering Plants	1078	A`e ( <i>Zanthoxylum dipetalum</i> var. <i>tomentosum</i> ) - Wherever found	E	\$7,989	\$0	\$7,989	\$0	\$7,989
	853	A`e ( <i>Zanthoxylum hawaiiense</i> ) - Wherever found	E	\$16,593	\$0	\$16,593	\$0	\$16,593
	1152	A`e ( <i>Zanthoxylum oahuense</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	375	`Ahinahina ( <i>Argyroxiphium sandwicense</i> ssp. <i>macrocephalum</i> ) - Wherever found	T	\$11,705	\$125,000	\$136,705	\$0	\$136,705
	1116	`Ahinahina ( <i>Argyroxiphium sandwicense</i> ssp. <i>sandwicense</i> ) - Wherever found	E	\$6,480	\$0	\$6,480	\$0	\$6,480

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	943	`Aiakeakua, popolo ( <i>Solanum sandwicense</i> ) - Wherever found	E	\$12,515	\$0	\$12,515	\$0	\$12,515
	734	`Aiea ( <i>Nothoecstrum breviflorum</i> ) - Wherever found	E	\$15,509	\$10,000	\$25,509	\$0	\$25,509
	540	`Aiea ( <i>Nothoecstrum latifolium</i> ) - Wherever found	E	\$47,238	\$12,000	\$59,238	\$0	\$59,238
	1083	`Aiea ( <i>Nothoecstrum peltatum</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1120	`Akoko ( <i>Euphorbia celastroides</i> var. <i>kaenana</i> ) - Wherever found	E	\$6,412	\$0	\$6,412	\$0	\$6,412
	1152	`Akoko ( <i>Euphorbia deppeana</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1083	`Akoko ( <i>Euphorbia eleanoriae</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1074	`Akoko ( <i>Euphorbia haelealeana</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1144	`Akoko ( <i>Euphorbia herbstii</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1021	`Akoko ( <i>Euphorbia kuwaleana</i> ) - Wherever found	E	\$7,736	\$1,792	\$9,528	\$0	\$9,528
	1152	`Akoko ( <i>Euphorbia remyi</i> var. <i>kauaiensis</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1073	`Akoko ( <i>Euphorbia remyi</i> var. <i>remyi</i> ) - Wherever found	E	\$8,083	\$0	\$8,083	\$0	\$8,083
	1144	`Akoko ( <i>Euphorbia rockii</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1083	"Akoko ( <i>Euphorbia halemanui</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	432	`Akoko, Ewa Plains ( <i>Euphorbia skottsbergii</i> var. <i>skottsbergii</i> ) - Wherever found	E	\$96,575	\$1,561	\$98,136	\$0	\$98,136
	880	`aku ( <i>Cyanea tritomantha</i> ) - Wherever found	E	\$13,520	\$1,973	\$15,493	\$0	\$15,493
	1060	`aku`aku ( <i>Cyanea platyphylla</i> ) - Wherever found	E	\$8,326	\$0	\$8,326	\$0	\$8,326
	1150	`Ala `ala wai nui ( <i>Peperomia subpetiolata</i> ) - Wherever found	E	\$5,599	\$0	\$5,599	\$0	\$5,599
	801	Alani ( <i>Melicope adscendens</i> ) - Wherever found	E	\$19,768	\$0	\$19,768	\$0	\$19,768
	926	Alani ( <i>Melicope balloui</i> ) - Wherever found	E	\$13,284	\$0	\$13,284	\$0	\$13,284



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1008	Alani (Melicope christophersenii) - Wherever found	E	\$8,073	\$1,792	\$9,865	\$0	\$9,865
	1152	Alani (Melicope degeneri) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1083	Alani (Melicope haupuensis) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1083	Alani (Melicope hiiakae) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	814	Alani (Melicope knudsenii) - Wherever found	E	\$19,044	\$0	\$19,044	\$0	\$19,044
	1074	Alani (Melicope lydgatei) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1074	Alani (Melicope makahae) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1051	Alani (Melicope mucronulata) - Wherever found	E	\$8,459	\$0	\$8,459	\$0	\$8,459
	434	Alani (Melicope munroi) - Wherever found	E	\$97,728	\$0	\$97,728	\$0	\$97,728
	570	Alani (Melicope ovalis) - Wherever found	E	\$14,036	\$36,000	\$50,036	\$0	\$50,036
	1152	Alani (Melicope pallida) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1152	Alani (Melicope paniculata) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1016	Alani (Melicope puberula) - Wherever found	E	\$9,669	\$0	\$9,669	\$0	\$9,669
	1051	Alani (Melicope quadrangularis) - Wherever found	E	\$8,459	\$0	\$8,459	\$0	\$8,459
	1083	Alani (Melicope reflexa) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1097	Alani (Melicope saint-johnii) - Wherever found	E	\$5,417	\$1,792	\$7,209	\$0	\$7,209
	1082	Alani (Melicope zahlbruckneri) - Wherever found	E	\$5,833	\$1,973	\$7,806	\$0	\$7,806
	997	Allocarya, Calistoga (Plagiobothrys strictus) - Wherever found	E	\$10,000	\$300	\$10,300	\$0	\$10,300
	1076	Alopecurus, Sonoma (Alopecurus aequalis var. sonomensis) - Wherever found	E	\$4,620	\$3,400	\$8,020	\$0	\$8,020
	275	Amaranth, seabeach (Amaranthus pumilus) - Wherever found	T	\$119,133	\$162,660	\$281,793	\$0	\$281,793
	324	Ambrosia, San Diego (Ambrosia pumila) - Wherever found	E	\$26,000	\$163,000	\$189,000	\$0	\$189,000
	725	Ambrosia, south Texas (Ambrosia cheiranthifolia) - Wherever found	E	\$26,500	\$0	\$26,500	\$0	\$26,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	397	Amole, purple ( <i>Chlorogalum purpureum</i> ) - Wherever found	T	\$19,000	\$106,097	\$125,097	\$0	\$125,097
	1232	Amphianthus, little ( <i>Amphianthus pusillus</i> ) - Wherever found	T	\$0	\$0	\$0	\$2,000	\$2,000
	1093	`Anaunau ( <i>Lepidium arbuscula</i> ) - Wherever found	E	\$5,602	\$1,792	\$7,394	\$0	\$7,394
	782	`Anunu ( <i>Sicyos albus</i> ) - Wherever found	E	\$16,007	\$4,973	\$20,980	\$0	\$20,980
	896	`Anunu ( <i>Sicyos macrophyllus</i> ) - Wherever found	E	\$8,509	\$6,000	\$14,509	\$0	\$14,509
	707	Aplokating-palaoan ( <i>Psychotria malaspiniae</i> ) - Wherever found	E	\$28,470	\$0	\$28,470	\$0	\$28,470
	772	Arrowhead, bunched ( <i>Sagittaria fasciculata</i> ) - Wherever found	E	\$18,000	\$0	\$18,000	\$4,100	\$22,100
	498	Aster, decurrent false ( <i>Boltonia decurrens</i> ) - Wherever found	T	\$15,200	\$56,044	\$71,244	\$0	\$71,244
	645	Aster, Florida golden ( <i>Chrysopsis floridana</i> ) - Wherever found	E	\$8,000	\$0	\$8,000	\$29,000	\$37,000
	781	Aster, Ruth's golden ( <i>Pityopsis ruthii</i> ) - Wherever found	E	\$14,000	\$7,000	\$21,000	\$0	\$21,000
	1109	Aupaka ( <i>Isodendrion hosakae</i> ) - Wherever found	E	\$6,665	\$0	\$6,665	\$0	\$6,665
	943	Aupaka ( <i>Isodendrion laurifolium</i> ) - Wherever found	E	\$12,515	\$0	\$12,515	\$0	\$12,515
	829	Aupaka ( <i>Isodendrion longifolium</i> ) - Wherever found	T	\$18,257	\$0	\$18,257	\$0	\$18,257
	783	Avens, spreading ( <i>Geum radiatum</i> ) - Wherever found	E	\$6,000	\$14,475	\$20,475	\$400	\$20,875
	915	`Awikiwiki ( <i>Canavalia molokaiensis</i> ) - Wherever found	E	\$8,712	\$5,000	\$13,712	\$0	\$13,712
	1152	`Awikiwiki ( <i>Canavalia napaliensis</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1032	`Awikiwiki ( <i>Canavalia pubescens</i> ) - Wherever found	E	\$9,148	\$0	\$9,148	\$0	\$9,148
	1083	Awiwi ( <i>Schenkia sebaeoides</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1041	'Awiwi ( <i>Kadua cookiana</i> ) - Wherever found	E	\$8,796	\$0	\$8,796	\$0	\$8,796
	668	Ayenia, Texas ( <i>Ayenia limitaris</i> ) - Wherever found	E	\$32,500	\$0	\$32,500	\$0	\$32,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	305	Baccharis, Encinitas ( <i>Baccharis vanessae</i> ) - Wherever found	T	\$2,460	\$220,300	\$222,760	\$0	\$222,760
	1136	Barbara's Buttons, Mohr's ( <i>Marshallia mohrii</i> ) - Wherever found	T	\$300	\$1,677	\$1,977	\$4,000	\$5,977
	669	Barberry, island ( <i>Berberis pinnata</i> ssp. <i>insularis</i> ) - Wherever found	E	\$22,100	\$10,399	\$32,499	\$0	\$32,499
	1106	Barberry, Nevin's ( <i>Berberis nevinii</i> ) - Wherever found	E	\$0	\$6,700	\$6,700	\$0	\$6,700
	1088	Bariaco ( <i>Trichilia triacantha</i> ) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	597	Beaked-rush, Knieskern's ( <i>Rhynchospora knieskernii</i> ) - Wherever found	T	\$29,987	\$14,874	\$44,861	\$0	\$44,861
	1072	Bean, sea ( <i>Mucuna sloanei</i> var. <i>persericea</i> ) - Wherever found	E	\$8,104	\$0	\$8,104	\$0	\$8,104
	889	Beardtongue, Parachute ( <i>Penstemon debilis</i> ) - Wherever found	T	\$6,100	\$8,800	\$14,900	\$0	\$14,900
	1092	Beardtongue, Penland ( <i>Penstemon penlandii</i> ) - Wherever found	E	\$400	\$7,000	\$7,400	\$0	\$7,400
	674	Beargrass, Britton's ( <i>Nolina brittoniana</i> ) - Wherever found	E	\$1,200	\$21,700	\$22,900	\$8,900	\$31,800
	683	Bear-poppy, Dwarf ( <i>Arctomecon humilis</i> ) - Wherever found	E	\$0	\$30,903	\$30,903	\$0	\$30,903
	672	Beauty, Harper's ( <i>Harperocallis flava</i> ) - Wherever found	E	\$2,000	\$29,700	\$31,700	\$300	\$32,000
	903	Bedstraw, El Dorado ( <i>Galium californicum</i> ssp. <i>sierrae</i> ) - Wherever found	E	\$1,700	\$12,450	\$14,150	\$0	\$14,150
	1175	Bedstraw, island ( <i>Galium buxifolium</i> ) - Wherever found	E	\$0	\$4,356	\$4,356	\$0	\$4,356
	1135	Bellflower, Brooksville ( <i>Campanula robinsiae</i> ) - Wherever found	E	\$0	\$0	\$0	\$6,000	\$6,000
	1260	Birch, Virginia round-leaf ( <i>Betula uber</i> ) - Wherever found	T	\$400	\$0	\$400	\$0	\$400
	665	Bird's beak, palmate-bracted ( <i>Cordylanthus palmatus</i> ) - Wherever found	E	\$10,000	\$22,992	\$32,992	\$0	\$32,992
	1232	Bird's-beak, Pennell's ( <i>Cordylanthus tenuis</i> ssp. <i>capillaris</i> ) - Wherever found	E	\$1,700	\$300	\$2,000	\$0	\$2,000
	535	Bird's-beak, salt marsh ( <i>Cordylanthus maritimus</i> ssp. <i>maritimus</i> ) - Wherever found	E	\$27,900	\$32,884	\$60,784	\$0	\$60,784

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	536	Bird's-beak, soft ( <i>Cordylanthus mollis</i> ssp. <i>mollis</i> ) - Wherever found	E	\$34,400	\$25,800	\$60,200	\$0	\$60,200
	1178	Birds-in-a-nest, white ( <i>Macbridea alba</i> ) - Wherever found	T	\$1,200	\$1,100	\$2,300	\$1,900	\$4,200
	1002	Bittercress, small-anthered ( <i>Cardamine micranthera</i> ) - Wherever found	E	\$9,700	\$0	\$9,700	\$400	\$10,100
	768	Bladderpod, Dudley Bluffs ( <i>Lesquerella congesta</i> ) - Wherever found	T	\$5,300	\$16,000	\$21,300	\$1,450	\$22,750
	1237	Bladderpod, kodachrome ( <i>Lesquerella tumulosa</i> ) - Wherever found	E	\$0	\$1,650	\$1,650	\$0	\$1,650
	1221	Bladderpod, lyrate ( <i>Lesquerella lyrata</i> ) - Wherever found	T	\$2,500	\$0	\$2,500	\$0	\$2,500
	562	Bladderpod, Missouri ( <i>Physaria filiformis</i> ) - Wherever found	T	\$5,000	\$45,142	\$50,142	\$2,000	\$52,142
	1270	Bladderpod, San Bernardino Mountains ( <i>Lesquerella kingii</i> ssp. <i>bernardina</i> ) - Wherever found	E	\$0	\$100	\$100	\$0	\$100
	599	Bladderpod, Short's ( <i>Physaria globosa</i> ) - Wherever found	E	\$17,500	\$25,843	\$43,343	\$1,300	\$44,643
	1187	Bladderpod, Spring Creek ( <i>Lesquerella perforata</i> ) - Wherever found	E	\$2,000	\$1,660	\$3,660	\$0	\$3,660
	1210	Bladderpod, white ( <i>Lesquerella pallida</i> ) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	387	Bladderpod, White Bluffs ( <i>Physaria douglasii</i> ssp. <i>tuplashensis</i> ) - Wherever found	T	\$129,037	\$0	\$129,037	\$0	\$129,037
	1077	Bladderpod, Zapata ( <i>Lesquerella thamnophila</i> ) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	874	Blazingstar, Ash Meadows ( <i>Mentzelia leucophylla</i> ) - Wherever found	T	\$15,600	\$0	\$15,600	\$0	\$15,600
	933	Blazingstar, Heller's ( <i>Liatris helleri</i> ) - Wherever found	T	\$4,500	\$5,200	\$9,700	\$3,200	\$12,900
	1159	Blazingstar, scrub ( <i>Liatris ohlingerae</i> ) - Wherever found	E	\$0	\$0	\$0	\$5,100	\$5,100
	1152	Bluegrass, Hawaiian ( <i>Poa sandvicensis</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1158	Bluegrass, Mann's ( <i>Poa mannii</i> ) - Wherever found	E	\$5,143	\$0	\$5,143	\$0	\$5,143

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	984	Bluegrass, Napa ( <i>Poa napensis</i> ) - Wherever found	E	\$10,600	\$200	\$10,800	\$0	\$10,800
	1159	Bluegrass, San Bernardino ( <i>Poa atropurpurea</i> ) - Wherever found	E	\$4,600	\$500	\$5,100	\$0	\$5,100
	774	Blue-star, Kearney's ( <i>Amsonia kearneyana</i> ) - Wherever found	E	\$11,000	\$10,900	\$21,900	\$0	\$21,900
	1132	Bluet, Roan Mountain ( <i>Hedyotis purpurea</i> var. <i>montana</i> ) - Wherever found	E	\$2,900	\$2,700	\$5,600	\$500	\$6,100
	679	Bonamia, Florida ( <i>Bonamia grandiflora</i> ) - Wherever found	T	\$6,000	\$21,700	\$27,700	\$3,500	\$31,200
	1025	Boxwood, Vahl's ( <i>Buxus vahlii</i> ) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	1244	Brickell-bush, Florida ( <i>Brickellia mosieri</i> ) - Wherever found	E	\$0	\$0	\$0	\$1,300	\$1,300
	556	Brodiaea, Chinese Camp ( <i>Brodiaea pallida</i> ) - Wherever found	T	\$49,900	\$3,800	\$53,700	\$0	\$53,700
	238	Brodiaea, thread-leaved ( <i>Brodiaea filifolia</i> ) - Wherever found	T	\$245,600	\$166,868	\$412,468	\$0	\$412,468
	1151	Buckwheat, cushenbury ( <i>Eriogonum ovalifolium</i> var. <i>vineum</i> ) - Wherever found	E	\$200	\$5,300	\$5,500	\$0	\$5,500
	708	Buckwheat, Ione (incl. Irish Hill) ( <i>Eriogonum apricum</i> (incl. var. <i>prostratum</i> )) - Wherever found	E	\$25,700	\$2,700	\$28,400	\$0	\$28,400
	625	Buckwheat, scrub ( <i>Eriogonum longifolium</i> var. <i>gnaphalifolium</i> ) - Wherever found	T	\$5,500	\$26,426	\$31,926	\$8,000	\$39,926
	528	Buckwheat, steamboat ( <i>Eriogonum ovalifolium</i> var. <i>williamsiae</i> ) - Wherever found	E	\$38,400	\$24,193	\$62,593	\$0	\$62,593
	1262	Bully, Everglades ( <i>Sideroxylon reclinatum</i> ssp. <i>austrofloridense</i> ) - Wherever found	T	\$0	\$100	\$100	\$200	\$300
	438	Bulrush, Northeastern ( <i>Scirpus ancistrochaetus</i> ) -	E	\$75,395	\$13,440	\$88,835	\$8,038	\$96,873
	907	Bush-clover, prairie ( <i>Lespedeza leptostachya</i> ) - Wherever found	T	\$9,000	\$3,500	\$12,500	\$1,500	\$14,000
	342	Bush-mallow, San Clemente Island ( <i>Malacothamnus clementinus</i> ) - Wherever found	E	\$3,500	\$158,417	\$161,917	\$0	\$161,917

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1113	Bush-mallow, Santa Cruz Island (Malacothamnus fasciculatus var. nesioticus) - Wherever found	E	\$0	\$6,534	\$6,534	\$0	\$6,534
	1235	Buttercup, autumn (Ranunculus aestivalis (=acriformis)) - Wherever found	E	\$0	\$1,800	\$1,800	\$0	\$1,800
	468	Butterfly plant, Colorado (Gaura neomexicana var. coloradensis) - Wherever found	T	\$45,000	\$39,174	\$84,174	\$100	\$84,274
	647	Butterweed, Layne's (Senecio layneae) - Wherever found	T	\$17,300	\$19,250	\$36,550	\$0	\$36,550
	1049	Butterwort, Godfrey's (Pinguicula ionantha) - Wherever found	T	\$1,200	\$5,600	\$6,800	\$1,800	\$8,600
	314	Button-celery, San Diego (Eryngium aristulatum var. parishii) - Wherever found	E	\$8,300	\$198,390	\$206,690	\$0	\$206,690
	832	Cactus, Acuna (Echinomastus erectocentrus var. acunensis) - Wherever found	E	\$2,000	\$15,908	\$17,908	\$0	\$17,908
	695	Cactus, Arizona hedgehog (Echinocereus triglochidiatus var. arizonicus) - Wherever found	E	\$12,000	\$17,704	\$29,704	\$0	\$29,704
	737	Cactus, Bakersfield (Opuntia treleasei) - Wherever found	E	\$23,000	\$2,400	\$25,400	\$0	\$25,400
	864	Cactus, black lace (Echinocereus reichenbachii var. albertii) - Wherever found	E	\$8,000	\$0	\$8,000	\$8,089	\$16,089
	1135	Cactus, Brady pincushion (Pediocactus bradyi) - Wherever found	E	\$4,500	\$1,500	\$6,000	\$0	\$6,000
	1239	Cactus, Chisos Mountain hedgehog (Echinocereus chisoensis var. chisoensis) - Wherever found	T	\$1,000	\$500	\$1,500	\$0	\$1,500
	1182	Cactus, Cochise pincushion (Coryphantha robbinsorum) - Wherever found	T	\$4,000	\$0	\$4,000	\$0	\$4,000
	360	Cactus, Colorado hookless (Sclerocactus glaucus) - Wherever found	T	\$13,100	\$131,633	\$144,733	\$1,200	\$145,933
	743	Cactus, Fickeisen plains (Pediocactus peeblesianus fickeiseniae) - Wherever found	E	\$800	\$24,000	\$24,800	\$0	\$24,800
	1092	Cactus, Florida semaphore (Consolea corallicola) - Wherever found	E	\$4,500	\$2,300	\$6,800	\$600	\$7,400

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	882	Cactus, Key tree ( <i>Pilosocereus robinii</i> ) - Wherever found	E	\$3,500	\$1,342	\$4,842	\$10,500	\$15,342
	671	Cactus, Knowlton's ( <i>Pediocactus knowltonii</i> ) - Wherever found	E	\$15,300	\$17,000	\$32,300	\$0	\$32,300
	390	Cactus, Kuenzler hedgehog ( <i>Echinocereus fendleri</i> var. <i>kuenzleri</i> ) - Wherever found	T	\$27,000	\$100,386	\$127,386	\$0	\$127,386
	831	Cactus, Lee pincushion ( <i>Coryphantha sneedii</i> var. <i>leei</i> ) - Wherever found	T	\$15,000	\$3,000	\$18,000	\$0	\$18,000
	1239	Cactus, Lloyd's Mariposa ( <i>Echinomastus mariposensis</i> ) - Wherever found	T	\$1,000	\$500	\$1,500	\$0	\$1,500
	593	Cactus, Mesa Verde ( <i>Sclerocactus mesae-verdae</i> ) - Wherever found	T	\$15,300	\$30,500	\$45,800	\$0	\$45,800
	1253	Cactus, Nellie cory ( <i>Coryphantha minima</i> ) - Wherever found	E	\$1,000	\$0	\$1,000	\$0	\$1,000
	1217	Cactus, Nichol's Turk's head ( <i>Echinocactus horizontalonius</i> var. <i>nicholii</i> ) - Wherever found	E	\$500	\$2,100	\$2,600	\$0	\$2,600
	919	Cactus, Pariette ( <i>Sclerocactus brevispinus</i> ) - Wherever found	T	\$0	\$13,579	\$13,579	\$0	\$13,579
	1216	Cactus, Peebles Navajo ( <i>Pediocactus peeblesianus</i> var. <i>peeblesianus</i> ) - Wherever found	E	\$300	\$2,350	\$2,650	\$0	\$2,650
	424	Cactus, Pima pineapple ( <i>Coryphantha scheeri</i> var. <i>robustispina</i> ) - Wherever found	E	\$49,845	\$50,200	\$100,045	\$0	\$100,045
	785	Cactus, San Rafael ( <i>Pediocactus despainii</i> ) - Wherever found	E	\$0	\$20,750	\$20,750	\$0	\$20,750
	716	Cactus, Siler pincushion ( <i>Pediocactus</i> (= <i>Echinocactus</i> ,= <i>Utahia</i> ) <i>sileri</i> ) - Wherever found	T	\$6,500	\$20,700	\$27,200	\$0	\$27,200
	1050	Cactus, Sneed pincushion ( <i>Coryphantha sneedii</i> var. <i>sneedii</i> ) - Wherever found	E	\$7,000	\$1,500	\$8,500	\$0	\$8,500
	773	Cactus, star ( <i>Astrophytum asterias</i> ) - Wherever found	E	\$22,000	\$0	\$22,000	\$0	\$22,000
	1100	Cactus, Tobusch fishhook ( <i>Sclerocactus brevihamatus</i> ssp. <i>tobuschii</i> ) - Wherever found	T	\$5,000	\$2,100	\$7,100	\$0	\$7,100
	846	Cactus, Uinta Basin hookless ( <i>Sclerocactus wetlandicus</i> ) - Wherever found	T	\$1,100	\$15,992	\$17,092	\$0	\$17,092

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	615	Cactus, Winkler ( <i>Pediocactus winkleri</i> ) - Wherever found	T	\$0	\$41,500	\$41,500	\$0	\$41,500
	410	Cactus, Wright fishhook ( <i>Sclerocactus wrightiae</i> ) - Wherever found	E	\$0	\$109,750	\$109,750	\$0	\$109,750
	1047	Campion, fringed ( <i>Silene polypetala</i> ) - Wherever found	E	\$0	\$1,912	\$1,912	\$6,700	\$8,612
	1135	Capa rosa ( <i>Callicarpa ampla</i> ) - Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	256	Catchfly, Spalding's ( <i>Silene spaldingii</i> ) - Wherever found	T	\$270,645	\$64,605	\$335,250	\$0	\$335,250
	1258	Cat's-eye, Terlingua Creek ( <i>Cryptantha crassipes</i> ) - Wherever found	E	\$500	\$0	\$500	\$0	\$500
	843	Ceanothus, coyote ( <i>Ceanothus ferrisae</i> ) - Wherever found	E	\$17,200	\$100	\$17,300	\$0	\$17,300
	899	Ceanothus, Pine Hill ( <i>Ceanothus roderickii</i> ) - Wherever found	E	\$4,358	\$10,050	\$14,408	\$0	\$14,408
	723	Centauray, spring-loving ( <i>Centaureum namophilum</i> ) - Wherever found	T	\$20,600	\$6,000	\$26,600	\$0	\$26,600
	439	Chaff-flower, round-leaved ( <i>Achyranthes splendens</i> var. <i>rotundata</i> ) - Wherever found	E	\$96,856	\$0	\$96,856	\$0	\$96,856
	174	Chaffseed, American ( <i>Schwalbea americana</i> ) - Wherever found	E	\$127,749	\$531,795	\$659,544	\$5,600	\$665,144
	858	Checker-mallow, Keck's ( <i>Sidalcea keckii</i> ) - Wherever found	E	\$8,500	\$7,893	\$16,393	\$0	\$16,393
	1233	Checker-mallow, Kenwood Marsh ( <i>Sidalcea oregana</i> ssp. <i>valida</i> ) - Wherever found	E	\$1,700	\$200	\$1,900	\$0	\$1,900
	246	Checker-mallow, Nelson's ( <i>Sidalcea nelsoniana</i> ) - Wherever found	T	\$218,450	\$152,500	\$370,950	\$0	\$370,950
	1262	Checker-mallow, pedate ( <i>Sidalcea pedata</i> ) - Wherever found	E	\$0	\$300	\$300	\$0	\$300
	319	Checkermallow, Wenatchee Mountains ( <i>Sidalcea oregana</i> var. <i>calva</i> ) - Wherever found	E	\$193,000	\$6,000	\$199,000	\$0	\$199,000
	970	Chumbo, Higo ( <i>Harrisia portoricensis</i> ) - Wherever found	T	\$11,500	\$0	\$11,500	\$0	\$11,500
	1115	Chupacallos ( <i>Pleodendron macranthum</i> ) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1262	Clarkia, Pismo (Clarkia speciosa ssp. immaculata) - Wherever found	E	\$0	\$300	\$300	\$0	\$300
	969	Clarkia, Presidio (Clarkia franciscana) - Wherever found	E	\$9,400	\$2,200	\$11,600	\$0	\$11,600
	794	Clarkia, Springville (Clarkia springvillensis) - Wherever found	T	\$11,500	\$8,819	\$20,319	\$0	\$20,319
	850	Cliffrose, Arizona (Purshia (=Cowania) subintegra) - Wherever found	E	\$4,000	\$13,000	\$17,000	\$0	\$17,000
	1229	Clover, Monterey (Trifolium trichocalyx) - Wherever found	E	\$1,750	\$400	\$2,150	\$0	\$2,150
	332	Clover, running buffalo (Trifolium stoloniferum) -	E	\$95,842	\$72,218	\$168,060	\$8,858	\$176,918
	788	Clover, showy Indian (Trifolium amoenum) - Wherever found	E	\$17,300	\$3,200	\$20,500	\$0	\$20,500
	878	Cobana negra (Stahlia monosperma) - Wherever found	T	\$15,500	\$0	\$15,500	\$0	\$15,500
	318	Coneflower, smooth (Echinacea laevigata) - Wherever found	E	\$30,500	\$158,366	\$188,866	\$11,400	\$200,266
	1239	Cory cactus, bunched (Coryphantha ramillosa) - Wherever found	T	\$1,000	\$500	\$1,500	\$0	\$1,500
	1244	Crabgrass, Florida pineland (Digitaria pauciflora) - Wherever found	T	\$0	\$1,100	\$1,100	\$200	\$1,300
	1262	Crownbeard, big-leaved (Verbesina dissita) - Wherever found	T	\$0	\$300	\$300	\$0	\$300
	1062	Crownscale, San Jacinto Valley (Atriplex coronata var. notatior) - Wherever found	E	\$8,100	\$200	\$8,300	\$0	\$8,300
	651	Cycladenia, Jones (Cycladenia humilis var. jonesii) - Wherever found	T	\$800	\$35,030	\$35,830	\$0	\$35,830
	1168	Daisy, Lakeside (Hymenoxys herbacea) - Wherever found	T	\$1,600	\$3,076	\$4,676	\$0	\$4,676
	897	Daisy, Parish's (Erigeron parishii) - Wherever found	T	\$300	\$14,200	\$14,500	\$0	\$14,500
	224	Daisy, Willamette (Erigeron decumbens) - Wherever found	E	\$112,310	\$341,517	\$453,827	\$0	\$453,827
	564	Dawn-flower, Texas prairie (Hymenoxys texana) - Wherever found	E	\$47,000	\$5,000	\$52,000	\$0	\$52,000
	387	Desert buckwheat, Umtanum (Eriogonum codium) - Wherever found	T	\$129,037	\$0	\$129,037	\$0	\$129,037

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1077	Dogweed, ashy ( <i>Thymophylla tephroleuca</i> ) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	644	Dropwort, Canby's ( <i>Oxypolis canbyi</i> ) - Wherever found	E	\$4,514	\$5,240	\$9,754	\$27,400	\$37,154
	895	Dudleya, Conejo ( <i>Dudleya abramsii</i> ssp. <i>parva</i> ) - Wherever found	T	\$14,300	\$300	\$14,600	\$0	\$14,600
	892	Dudleya, marcescent ( <i>Dudleya cymosa</i> ssp. <i>marcescens</i> ) - Wherever found	T	\$14,300	\$400	\$14,700	\$0	\$14,700
	1243	Dudleya, Santa Clara Valley ( <i>Dudleya setchellii</i> ) - Wherever found	E	\$0	\$1,400	\$1,400	\$0	\$1,400
	845	Dudleya, Santa Monica Mountains ( <i>Dudleya cymosa</i> ssp. <i>ovatifolia</i> ) - Wherever found	T	\$17,200	\$0	\$17,200	\$0	\$17,200
	641	Dudleya, Verity's ( <i>Dudleya verityi</i> ) - Wherever found	T	\$37,400	\$0	\$37,400	\$0	\$37,400
	924	Dwarf-flax, Marin ( <i>Hesperolinon congestum</i> ) - Wherever found	T	\$11,200	\$2,200	\$13,400	\$0	\$13,400
	1128	'Ena`ena ( <i>Pseudognaphalium sandwicensium</i> var. <i>molokaiense</i> ) - Wherever found	E	\$5,701	\$500	\$6,201	\$0	\$6,201
	1095	Erubia ( <i>Solanum drymophilum</i> ) - Wherever found	E	\$7,000	\$300	\$7,300	\$0	\$7,300
	383	Evening-primrose, Antioch Dunes ( <i>Oenothera deltooides</i> ssp. <i>howellii</i> ) - Wherever found	E	\$129,600	\$1,700	\$131,300	\$0	\$131,300
	680	Evening-primrose, San Benito ( <i>Camissonia benitensis</i> ) - Wherever found	T	\$0	\$31,000	\$31,000	\$0	\$31,000
	1151	Fescue, Guadalupe ( <i>Festuca ligulata</i> ) - Wherever found	E	\$5,000	\$500	\$5,500	\$0	\$5,500
	306	Fiddleneck, large-flowered ( <i>Amsinckia grandiflora</i> ) - Wherever found	E	\$16,600	\$203,163	\$219,763	\$0	\$219,763
	720	Flannelbush, Mexican ( <i>Fremontodendron mexicanum</i> ) - Wherever found	E	\$300	\$26,600	\$26,900	\$0	\$26,900
	875	Flannelbush, Pine Hill ( <i>Fremontodendron californicum</i> ssp. <i>decumbens</i> ) - Wherever found	E	\$0	\$15,550	\$15,550	\$0	\$15,550
	1213	Flax, Carter's small-flowered ( <i>Linum carteri carteri</i> ) - Wherever found	E	\$1,500	\$0	\$1,500	\$1,300	\$2,800

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1235	Flax, sand ( <i>Linum arenicola</i> ) - Wherever found	E	\$0	\$0	\$0	\$1,800	\$1,800
	791	Fleabane, Zuni ( <i>Erigeron rhizomatus</i> ) - Wherever found	T	\$15,500	\$4,907	\$20,407	\$0	\$20,407
	435	Four-o'clock, MacFarlane's ( <i>Mirabilis macfarlanei</i> ) - Wherever found	T	\$86,414	\$10,757	\$97,171	\$0	\$97,171
	1217	Fringepod, Santa Cruz Island ( <i>Thysanocarpus conchuliferus</i> ) - Wherever found	E	\$2,600	\$0	\$2,600	\$0	\$2,600
	1178	Fringe-tree, pygmy ( <i>Chionanthus pygmaeus</i> ) - Wherever found	E	\$1,200	\$0	\$1,200	\$3,000	\$4,200
	259	Fritillary, Gentner's ( <i>Fritillaria gentneri</i> ) - Wherever found	E	\$100,000	\$225,500	\$325,500	\$0	\$325,500
	1094	Gardenia (=Na`u), Hawaiian ( <i>Gardenia brighamii</i> ) - Wherever found	E	\$7,319	\$0	\$7,319	\$0	\$7,319
	595	Geranium, Hawaiian red-flowered ( <i>Geranium arboreum</i> ) - Wherever found	E	\$9,386	\$36,000	\$45,386	\$0	\$45,386
	746	Gerardia, sandplain ( <i>Agalinis acuta</i> ) - Wherever found	E	\$22,680	\$1,240	\$23,920	\$600	\$24,520
	1009	Gilia, Hoffmann's slender-flowered ( <i>Gilia tenuiflora</i> ssp. <i>hoffmannii</i> ) - Wherever found	E	\$2,600	\$7,258	\$9,858	\$0	\$9,858
	793	Gilia, Monterey ( <i>Gilia tenuiflora</i> ssp. <i>arenaria</i> ) - Wherever found	E	\$15,300	\$5,044	\$20,344	\$0	\$20,344
	1139	Glade cress, Kentucky ( <i>Leavenworthia exigua laciniata</i> ) - Wherever found	T	\$0	\$5,920	\$5,920	\$0	\$5,920
	1208	Gladecress, Texas golden ( <i>Leavenworthia texana</i> ) - Wherever found	E	\$3,100	\$0	\$3,100	\$0	\$3,100
	954	Goetzea, beautiful ( <i>Goetzea elegans</i> ) - Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	1226	Goldenrod, Blue Ridge ( <i>Solidago spithamea</i> ) - Wherever found	T	\$2,000	\$0	\$2,000	\$300	\$2,300
	568	Goldenrod, Houghton's ( <i>Solidago houghtonii</i> ) - Wherever found	T	\$49,975	\$513	\$50,488	\$0	\$50,488
	904	Goldenrod, Short's ( <i>Solidago shortii</i> ) - Wherever found	E	\$11,000	\$1,332	\$12,332	\$1,800	\$14,132
	672	Goldfields, Burke's ( <i>Lasthenia burkei</i> ) - Wherever found	E	\$10,300	\$21,700	\$32,000	\$0	\$32,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	371	Goldfields, Contra Costa ( <i>Lasthenia conjugens</i> ) - Wherever found	E	\$111,100	\$26,763	\$137,863	\$0	\$137,863
	973	Gooseberry, Miccosukee ( <i>Ribes echinellum</i> ) - Wherever found	T	\$1,200	\$10,000	\$11,200	\$200	\$11,400
	627	Gourd, Okeechobee ( <i>Cucurbita okeechobeensis</i> ssp. <i>okeechobeensis</i> ) - Wherever found	E	\$39,000	\$500	\$39,500	\$200	\$39,700
	712	Grass, Colusa ( <i>Neostapfia colusana</i> ) - Wherever found	T	\$22,800	\$4,990	\$27,790	\$0	\$27,790
	477	Grass, Eureka Dune ( <i>Swallenia alexandrae</i> ) -	T	\$29,700	\$50,000	\$79,700	\$0	\$79,700
	1031	Grass, Solano ( <i>Tuctoria mucronata</i> ) - Wherever found	E	\$8,500	\$800	\$9,300	\$0	\$9,300
	251	Grass, Tennessee yellow-eyed ( <i>Xyris</i> <i>tennesseensis</i> ) - Wherever found	E	\$1,700	\$353,780	\$355,480	\$5,000	\$360,480
	1188	Ground-plum, Guthrie's (=Pyne's) ( <i>Astragalus</i> <i>bibullatus</i> ) - Wherever found	E	\$3,000	\$650	\$3,650	\$0	\$3,650
	723	Gumplant, Ash Meadows ( <i>Grindelia</i> <i>fraxinipratensis</i> ) - Wherever found	T	\$20,600	\$6,000	\$26,600	\$0	\$26,600
	1074	Haha ( <i>Cyanea acuminata</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1152	Haha ( <i>Cyanea asarifolia</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	598	Haha ( <i>Cyanea asplenifolia</i> ) - Wherever found	E	\$8,846	\$36,000	\$44,846	\$0	\$44,846
	909	Haha ( <i>Cyanea calycina</i> ) - Wherever found	E	\$9,087	\$4,865	\$13,952	\$0	\$13,952
	1152	Haha ( <i>Cyanea copelandii</i> ssp. <i>copelandii</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	573	Haha ( <i>Cyanea copelandii</i> ssp. <i>haleakalaensis</i> ) - Wherever found	E	\$13,537	\$36,000	\$49,537	\$0	\$49,537
	1074	Haha ( <i>Cyanea crispa</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1083	Haha ( <i>Cyanea dolichopoda</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	979	Haha ( <i>Cyanea dunbariae</i> ) - Wherever found	E	\$11,031	\$0	\$11,031	\$0	\$11,031
	1034	Haha ( <i>Cyanea duvalliorum</i> ) - Wherever found	E	\$9,099	\$0	\$9,099	\$0	\$9,099
	1083	Haha ( <i>Cyanea eleeleensis</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	985	Haha ( <i>Cyanea gibsonii</i> ) - Wherever found	E	\$10,778	\$0	\$10,778	\$0	\$10,778
	920	Haha ( <i>Cyanea glabra</i> ) - Wherever found	E	\$13,537	\$0	\$13,537	\$0	\$13,537

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1013	Haha ( <i>Cyanea grimesiana</i> ssp. <i>grimesiana</i> ) - Wherever found	E	\$9,722	\$0	\$9,722	\$0	\$9,722
	1060	Haha ( <i>Cyanea grimesiana</i> ssp. <i>obatae</i> ) - Wherever found	E	\$8,326	\$0	\$8,326	\$0	\$8,326
	1060	Haha ( <i>Cyanea hamatiflora</i> ssp. <i>carlsonii</i> ) - Wherever found	E	\$8,326	\$0	\$8,326	\$0	\$8,326
	733	Haha ( <i>Cyanea hamatiflora</i> ssp. <i>hamatiflora</i> ) - Wherever found	E	\$13,537	\$12,000	\$25,537	\$0	\$25,537
	1078	Haha ( <i>Cyanea humboldtiana</i> ) - Wherever found	E	\$7,989	\$0	\$7,989	\$0	\$7,989
	1083	Haha ( <i>Cyanea kolekoleensis</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1074	Haha ( <i>Cyanea koolauensis</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1152	Haha ( <i>Cyanea kuhihewa</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	769	Haha ( <i>Cyanea kunthiana</i> ) - Wherever found	E	\$10,678	\$12,000	\$22,678	\$0	\$22,678
	1152	Haha ( <i>Cyanea lanceolata</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	963	Haha ( <i>Cyanea lobata</i> ) - Wherever found	E	\$11,713	\$0	\$11,713	\$0	\$11,713
	1074	Haha ( <i>Cyanea longiflora</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1007	Haha ( <i>Cyanea magnicalyx</i> ) - Wherever found	E	\$9,868	\$0	\$9,868	\$0	\$9,868
	985	Haha ( <i>Cyanea manni</i> ) - Wherever found	E	\$10,778	\$0	\$10,778	\$0	\$10,778
	686	Haha ( <i>Cyanea maritae</i> ) - Wherever found	E	\$6,780	\$23,999	\$30,779	\$0	\$30,779
	1056	Haha ( <i>Cyanea marksii</i> ) - Wherever found	E	\$8,425	\$0	\$8,425	\$0	\$8,425
	1033	Haha ( <i>Cyanea mauiensis</i> ) - Wherever found	E	\$9,145	\$0	\$9,145	\$0	\$9,145
	935	Haha ( <i>Cyanea mceldowneyi</i> ) - Wherever found	E	\$12,863	\$0	\$12,863	\$0	\$12,863
	989	Haha ( <i>Cyanea munroi</i> ) - Wherever found	E	\$10,629	\$0	\$10,629	\$0	\$10,629
	1040	Haha ( <i>Cyanea obtusa</i> ) - Wherever found	E	\$8,892	\$0	\$8,892	\$0	\$8,892
	1083	Haha ( <i>Cyanea pinnatifida</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	979	Haha ( <i>Cyanea procera</i> ) - Wherever found	E	\$11,031	\$0	\$11,031	\$0	\$11,031
	1033	Haha ( <i>Cyanea profuga</i> ) - Wherever found	E	\$9,145	\$0	\$9,145	\$0	\$9,145
	1083	Haha ( <i>Cyanea purpurellifolia</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1083	Haha ( <i>Cyanea recta</i> ) - Wherever found	T	\$7,736	\$0	\$7,736	\$0	\$7,736

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1147	Haha ( <i>Cyanea remyi</i> ) - Wherever found	E	\$5,670	\$0	\$5,670	\$0	\$5,670
	1083	Haha ( <i>Cyanea rivularis</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	425	Haha ( <i>Cyanea shipmanii</i> ) - Wherever found	E	\$93,037	\$6,973	\$100,010	\$0	\$100,010
	421	Haha ( <i>Cyanea stictophylla</i> ) - Wherever found	E	\$95,356	\$6,973	\$102,329	\$0	\$102,329
	1060	Haha ( <i>Cyanea st.-johnii</i> ) - Wherever found	E	\$8,326	\$0	\$8,326	\$0	\$8,326
	444	Haha ( <i>Cyanea superba</i> ) - Wherever found	E	\$95,023	\$0	\$95,023	\$0	\$95,023
	1152	Haha ( <i>Cyanea truncata</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1083	Haha ( <i>Cyanea undulata</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	586	Haiwale ( <i>Cyrtandra ferripilosa</i> ) - Wherever found	E	\$6,780	\$40,300	\$47,080	\$0	\$47,080
	1004	Haiwale ( <i>Cyrtandra gracilis</i> ) - Wherever found	E	\$5,670	\$4,300	\$9,970	\$0	\$9,970
	1014	Haiwale ( <i>Cyrtandra paliku</i> ) - Wherever found	E	\$5,417	\$4,300	\$9,717	\$0	\$9,717
	953	Haiwale ( <i>Cyrtandra waiolani</i> ) - Wherever found	E	\$7,736	\$4,300	\$12,036	\$0	\$12,036
	1152	Ha`iwale ( <i>Cyrtandra crenata</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1144	Ha`iwale ( <i>Cyrtandra dentata</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	818	Ha`iwale ( <i>Cyrtandra filipes</i> ) - Wherever found	E	\$18,662	\$0	\$18,662	\$0	\$18,662
	998	Ha`iwale ( <i>Cyrtandra giffardii</i> ) - Wherever found	E	\$8,326	\$1,973	\$10,299	\$0	\$10,299
	1056	Ha`iwale ( <i>Cyrtandra hematos</i> ) - Wherever found	E	\$8,425	\$0	\$8,425	\$0	\$8,425
	1078	Ha`iwale ( <i>Cyrtandra kaulantha</i> ) - Wherever found	E	\$7,989	\$0	\$7,989	\$0	\$7,989
	939	Ha`iwale ( <i>Cyrtandra limahuliensis</i> ) - Wherever found	T	\$12,639	\$0	\$12,639	\$0	\$12,639
	877	Ha`iwale ( <i>Cyrtandra munroi</i> ) - Wherever found	E	\$15,516	\$0	\$15,516	\$0	\$15,516
	1083	Ha`iwale ( <i>Cyrtandra oenobarba</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1112	Ha`iwale ( <i>Cyrtandra oxybapha</i> ) - Wherever found	E	\$6,573	\$0	\$6,573	\$0	\$6,573
	1083	Ha`iwale ( <i>Cyrtandra polyantha</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1083	Ha`iwale ( <i>Cyrtandra sessilis</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1144	Ha`iwale ( <i>Cyrtandra subumbellata</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1079	Ha`iwale ( <i>Cyrtandra tintinnabula</i> ) - Wherever found	E	\$6,007	\$1,973	\$7,980	\$0	\$7,980
	1144	Ha`iwale ( <i>Cyrtandra viridiflora</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	976	Hala pepe ( <i>Pleomele fernaldii</i> ) - Wherever found	E	\$11,214	\$0	\$11,214	\$0	\$11,214
	1060	Hala pepe ( <i>Pleomele forbesii</i> ) - Wherever found	E	\$5,754	\$2,572	\$8,326	\$0	\$8,326
	736	Hala pepe ( <i>Pleomele hawaiiensis</i> ) - Wherever found	E	\$10,487	\$15,000	\$25,487	\$0	\$25,487
	1067	Halomtano, Berenghenas ( <i>Solanum guamense</i> ) - Wherever found	E	\$8,238	\$0	\$8,238	\$0	\$8,238
	496	Halumtano, Cebello ( <i>Bulbophyllum guamense</i> ) - Wherever found	T	\$72,018	\$0	\$72,018	\$0	\$72,018
	900	Harebells, Avon Park ( <i>Crotalaria avonensis</i> ) - Wherever found	E	\$2,000	\$0	\$2,000	\$12,400	\$14,400
	454	Harperella ( <i>Ptilimnium nodosum</i> ) - Wherever found	E	\$29,511	\$48,740	\$78,251	\$10,862	\$89,113
	938	Hau kuahiwi ( <i>Hibiscadelphus giffardianus</i> ) - Wherever found	E	\$7,736	\$5,000	\$12,736	\$0	\$12,736
	1078	Hau kuahiwi ( <i>Hibiscadelphus hualalaiensis</i> ) - Wherever found	E	\$7,989	\$0	\$7,989	\$0	\$7,989
	1083	Hau kuahiwi ( <i>Hibiscadelphus woodii</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	682	Heartleaf, dwarf-flowered ( <i>Hexastylis naniflora</i> ) - Wherever found	T	\$4,000	\$23,568	\$27,568	\$3,400	\$30,968
	684	Heather, mountain golden ( <i>Hudsonia montana</i> ) - Wherever found	T	\$4,500	\$26,400	\$30,900	\$0	\$30,900
	1149	Heau ( <i>Exocarpos luteolus</i> ) - Wherever found	E	\$5,602	\$0	\$5,602	\$0	\$5,602
	964	Heau ( <i>Exocarpos menziesii</i> ) -	E	\$10,695	\$1,000	\$11,695	\$0	\$11,695
	1083	Hibiscus, Clay's ( <i>Hibiscus clayi</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	751	Higuero de sierra ( <i>Crescentia portoricensis</i> ) - Wherever found	E	\$24,000	\$0	\$24,000	\$0	\$24,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	800	Hoawa ( <i>Pittosporum halophilum</i> ) - Wherever found	E	\$14,873	\$5,000	\$19,873	\$0	\$19,873
	824	Hoawa ( <i>Pittosporum hawaiiense</i> ) - Wherever found	E	\$13,520	\$5,000	\$18,520	\$0	\$18,520
	1026	Ho`awa ( <i>Pittosporum napaliense</i> ) - Wherever found	E	\$9,473	\$0	\$9,473	\$0	\$9,473
	630	Holei ( <i>Ochrosia haleakalae</i> ) - Wherever found	E	\$37,989	\$1,500	\$39,489	\$0	\$39,489
	1152	Holei ( <i>Ochrosia kilauaeensis</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1135	Holly, Cook's ( <i>Ilex cookii</i> ) - Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	475	Honohono ( <i>Haplostachys haplostachya</i> ) - Wherever found	E	\$50,158	\$30,000	\$80,158	\$0	\$80,158
	349	Howellia, Water ( <i>Howellia aquatilis</i> ) -	T	\$79,549	\$10,244	\$89,793	\$68,000	\$157,793
	1074	Hulumoa ( <i>Korthalsella degeneri</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1038	Hypericum, highlands scrub ( <i>Hypericum cumulicola</i> ) - Wherever found	E	\$900	\$0	\$900	\$8,000	\$8,900
	417	Iagu, Hayun (=Guam), Tronkon guafi ( <i>Rota</i> ) ( <i>Serianthes nelsonii</i> ) - Wherever found	E	\$43,183	\$60,869	\$104,052	\$0	\$104,052
	949	Ihi ( <i>Portulaca villosa</i> ) - Wherever found	E	\$12,274	\$0	\$12,274	\$0	\$12,274
	816	Iliau, dwarf ( <i>Wilkesia hobdyi</i> ) - Wherever found	E	\$7,736	\$11,229	\$18,965	\$0	\$18,965
	596	Ipomopsis, Holy Ghost ( <i>Ipomopsis sancti-spiritus</i> ) - Wherever found	E	\$5,000	\$40,000	\$45,000	\$0	\$45,000
	377	Iris, dwarf lake ( <i>Iris lacustris</i> ) - Wherever found	T	\$129,600	\$342	\$129,942	\$6,500	\$136,442
	1176	Irisette, white ( <i>Sisyrinchium dichotomum</i> ) - Wherever found	E	\$2,000	\$0	\$2,000	\$2,300	\$4,300
	681	Ischaemum, Hilo ( <i>Ischaemum byrone</i> ) - Wherever found	E	\$6,983	\$24,000	\$30,983	\$0	\$30,983
	874	Ivesia, Ash Meadows ( <i>Ivesia kingii</i> var. <i>eremica</i> ) - Wherever found	T	\$15,600	\$0	\$15,600	\$0	\$15,600
	379	Ivesia, Webber's ( <i>Ivesia webberi</i> ) - Wherever found	T	\$60,100	\$75,404	\$135,504	\$0	\$135,504
	1122	Jacquemontia, beach ( <i>Jacquemontia reclinata</i> ) - Wherever found	E	\$900	\$0	\$900	\$5,500	\$6,400



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	405	Jewelflower, California ( <i>Caulanthus californicus</i> ) - Wherever found	E	\$15,000	\$105,004	\$120,004	\$0	\$120,004
	1254	Jewelflower, Metcalf Canyon ( <i>Streptanthus albidus</i> ssp. <i>albidus</i> ) - Wherever found	E	\$0	\$800	\$800	\$0	\$800
	702	Joint-vetch, Sensitive ( <i>Aeschynomene virginica</i> ) - Wherever found	T	\$28,408	\$530	\$28,938	\$0	\$28,938
	1144	Kamakahala ( <i>Labordia cyrtandrae</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1083	Kamakahala ( <i>Labordia helleri</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	961	Kamakahala ( <i>Labordia lydgatei</i> ) - Wherever found	E	\$11,792	\$0	\$11,792	\$0	\$11,792
	1083	Kamakahala ( <i>Labordia pumila</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1051	Kamakahala ( <i>Labordia tinifolia</i> var. <i>lanaiensis</i> ) - Wherever found	E	\$8,459	\$0	\$8,459	\$0	\$8,459
	1083	Kamakahala ( <i>Labordia tinifolia</i> var. <i>wahiawaensis</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	985	Kamakahala ( <i>Labordia triflora</i> ) - Wherever found	E	\$10,778	\$0	\$10,778	\$0	\$10,778
	991	Kamanomano ( <i>Cenchrus agrimonioides</i> ) - Wherever found	E	\$10,505	\$0	\$10,505	\$0	\$10,505
	942	Kamapua`a ( <i>Kadua fluviatilis</i> ) - Wherever found	E	\$12,575	\$0	\$12,575	\$0	\$12,575
	1083	Kauai hau kuahiwi ( <i>Hibiscadelphus distans</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	796	Kauila ( <i>Colubrina oppositifolia</i> ) - Wherever found	E	\$10,294	\$10,000	\$20,294	\$0	\$20,294
	1152	Kaulu ( <i>Pteralyxia kauaiensis</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1144	Kaulu ( <i>Pteralyxia macrocarpa</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1083	Kio`ele ( <i>Kadua coriacea</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1074	Kiponapona ( <i>Phyllostegia racemosa</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1066	Kohe malama malama o kanaloa ( <i>Kanaloa kahoolawensis</i> ) - Wherever found	E	\$8,243	\$0	\$8,243	\$0	\$8,243
	1083	Koholapehu ( <i>Dubautia latifolia</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	978	Koki'o (Kokia drynarioides) - Wherever found	E	\$10,574	\$500	\$11,074	\$0	\$11,074
	1083	Koki'o (Kokia kauaiensis) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1051	Koki'o, Cooke's (Kokia cookei) - Wherever found	E	\$8,459	\$0	\$8,459	\$0	\$8,459
	867	Koki'o ke'oke'o (Hibiscus arnottianus ssp. immaculatus) - Wherever found	E	\$10,778	\$5,000	\$15,778	\$0	\$15,778
	1026	Koki'o ke'oke'o (Hibiscus waimeae ssp. hanneriae) - Wherever found	E	\$9,473	\$0	\$9,473	\$0	\$9,473
	1145	Kolea (Myrsine fosbergii) - Wherever found	E	\$5,701	\$0	\$5,701	\$0	\$5,701
	1144	Kolea (Myrsine juddii) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1152	Kolea (Myrsine knudsenii) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1073	Kolea (Myrsine linearifolia) - Wherever found	T	\$8,083	\$0	\$8,083	\$0	\$8,083
	1152	Kolea (Myrsine mezii) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	959	Kolea (Myrsine vaccinioides) - Wherever found	E	\$11,893	\$0	\$11,893	\$0	\$11,893
	721	Ko'oko'olau (Bidens amplexens) - Wherever found	E	\$26,896	\$0	\$26,896	\$0	\$26,896
	608	Ko'oko'olau (Bidens campylothea ssp. pentamera) - Wherever found	E	\$6,780	\$36,000	\$42,780	\$0	\$42,780
	608	Ko'oko'olau (Bidens campylothea ssp. waihoiensis) - Wherever found	E	\$6,780	\$36,000	\$42,780	\$0	\$42,780
	441	Ko'oko'olau (Bidens conjuncta) - Wherever found	E	\$96,095	\$0	\$96,095	\$0	\$96,095
	604	Ko'oko'olau (Bidens micrantha ssp. ctenophylla) - Wherever found	E	\$42,870	\$500	\$43,370	\$0	\$43,370
	549	Ko'oko'olau (Bidens micrantha ssp. kalealaha) - Wherever found	E	\$20,442	\$36,000	\$56,442	\$0	\$56,442
	1069	Ko'oko'olau (Bidens wiebkei) - Wherever found	E	\$8,188	\$0	\$8,188	\$0	\$8,188
	822	Ko'oloa'ula (Abutilon menziesii) - Wherever found	E	\$16,862	\$1,734	\$18,596	\$0	\$18,596
	1051	Kopa (Kadua cordata remyi) - Wherever found	E	\$8,459	\$0	\$8,459	\$0	\$8,459

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1155	Kopiko ( <i>Psychotria grandiflora</i> ) - Wherever found	E	\$5,265	\$0	\$5,265	\$0	\$5,265
	1083	Kopiko ( <i>Psychotria hexandra</i> ssp. <i>oahuensis</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1083	Kopiko ( <i>Psychotria hobbyi</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1133	Kuahiwi laukahi ( <i>Plantago hawaiiensis</i> ) - Wherever found	E	\$5,598	\$500	\$6,098	\$0	\$6,098
	575	Kuahiwi laukahi ( <i>Plantago princeps</i> ) - Wherever found	E	\$11,713	\$37,792	\$49,505	\$0	\$49,505
	956	Kuawawaenuhu ( <i>Schiedea lychnoides</i> ) - Wherever found	E	\$11,988	\$0	\$11,988	\$0	\$11,988
	654	Kula wahine noho ( <i>Isodendron pyrifolium</i> ) - Wherever found	E	\$35,045	\$500	\$35,545	\$0	\$35,545
	1008	Kulu`i ( <i>Nototrichium humile</i> ) - Wherever found	E	\$8,073	\$1,792	\$9,865	\$0	\$9,865
	1254	Ladies-tresses, Canelo Hills ( <i>Spiranthes delitescens</i> ) - Wherever found	E	\$800	\$0	\$800	\$0	\$800
	699	Ladies-tresses, Navasota ( <i>Spiranthes parksii</i> ) - Wherever found	E	\$19,000	\$10,265	\$29,265	\$0	\$29,265
	241	Ladies'-tresses, Ute ( <i>Spiranthes diluvialis</i> ) - Wherever found	T	\$75,339	\$320,813	\$396,152	\$100	\$396,252
	987	Larkspur, Baker's ( <i>Delphinium bakeri</i> ) - Wherever found	E	\$10,600	\$100	\$10,700	\$0	\$10,700
	357	Larkspur, San Clemente Island ( <i>Delphinium variegatum</i> ssp. <i>kinkiense</i> ) - Wherever found	E	\$3,500	\$145,951	\$149,451	\$0	\$149,451
	994	Larkspur, yellow ( <i>Delphinium luteum</i> ) - Wherever found	E	\$10,400	\$0	\$10,400	\$0	\$10,400
	906	Lau `ehu ( <i>Panicum niihauense</i> ) - Wherever found	E	\$7,736	\$6,300	\$14,036	\$0	\$14,036
	1051	Laulihilihi ( <i>Schiedea stellarioides</i> ) - Wherever found	E	\$8,459	\$0	\$8,459	\$0	\$8,459
	212	Layia, beach ( <i>Layia carnosa</i> ) - Wherever found	E	\$365,105	\$136,000	\$501,105	\$0	\$501,105
	655	Lead-plant, Crenulate ( <i>Amorpha crenulata</i> ) - Wherever found	E	\$7,500	\$21,700	\$29,200	\$6,000	\$35,200
	1210	Leather flower, Alabama ( <i>Clematis socialis</i> ) - Wherever found	E	\$0	\$0	\$0	\$3,000	\$3,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1023	Lehua makanoe ( <i>Lysimachia daphnoides</i> ) - Wherever found	E	\$9,517	\$0	\$9,517	\$0	\$9,517
	1210	Lessingia, San Francisco ( <i>Lessingia germanorum</i> (=L.g. var. <i>germanorum</i> )) - Wherever found	E	\$0	\$3,000	\$3,000	\$0	\$3,000
	1152	Liliwai ( <i>Acaena exigua</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1203	Lily, Minnesota dwarf trout ( <i>Erythronium propullans</i> ) - Wherever found	E	\$0	\$3,200	\$3,200	\$0	\$3,200
	1232	Lily, Pitkin Marsh ( <i>Lilium pardalinum</i> ssp. <i>pitkinense</i> ) - Wherever found	E	\$1,700	\$300	\$2,000	\$0	\$2,000
	673	Lily, Western ( <i>Lilium occidentale</i> ) - Wherever found	E	\$5,810	\$26,000	\$31,810	\$0	\$31,810
	1267	Liveforever, Laguna Beach ( <i>Dudleya stolonifera</i> ) - Wherever found	T	\$0	\$200	\$200	\$0	\$200
	1184	Liveforever, Santa Barbara Island ( <i>Dudleya traskiae</i> ) - Wherever found	E	\$2,700	\$1,139	\$3,839	\$0	\$3,839
	1221	Locoweed, Fassett's ( <i>Oxytropis campestris</i> var. <i>chartacea</i> ) - Wherever found	T	\$0	\$0	\$0	\$2,500	\$2,500
	210	Lomatium, Bradshaw's ( <i>Lomatium bradshawii</i> ) - Wherever found	E	\$110,550	\$399,619	\$510,169	\$0	\$510,169
	482	Lomatium, Cook's ( <i>Lomatium cookii</i> ) - Wherever found	E	\$14,310	\$64,000	\$78,310	\$0	\$78,310
	288	Loosestrife, rough-leaved ( <i>Lysimachia asperulaefolia</i> ) - Wherever found	E	\$23,000	\$232,640	\$255,640	\$2,600	\$258,240
	357	Lotus, San Clemente Island ( <i>Acmispon dendroideus</i> var. <i>traskiae</i> ) - Wherever found	T	\$3,500	\$145,951	\$149,451	\$0	\$149,451
	1051	Loulu ( <i>Pritchardia aylmer-robinsonii</i> ) - Wherever found	E	\$8,459	\$0	\$8,459	\$0	\$8,459
	1152	Loulu ( <i>Pritchardia bakeri</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1083	Loulu ( <i>Pritchardia hardyi</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1144	Loulu ( <i>Pritchardia kaalae</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	967	Loulu ( <i>Pritchardia lanigera</i> ) - Wherever found	E	\$11,676	\$0	\$11,676	\$0	\$11,676
	1129	Loulu ( <i>Pritchardia maideniana</i> ) - Wherever found	E	\$5,670	\$500	\$6,170	\$0	\$6,170

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	985	Loulu ( <i>Pritchardia munroi</i> ) - Wherever found	E	\$10,778	\$0	\$10,778	\$0	\$10,778
	985	Loulu ( <i>Pritchardia napaliensis</i> ) - Wherever found	E	\$10,778	\$0	\$10,778	\$0	\$10,778
	888	Loulu ( <i>Pritchardia remota</i> ) - Wherever found	E	\$14,948	\$0	\$14,948	\$0	\$14,948
	1147	Loulu ( <i>Pritchardia schattaueri</i> ) - Wherever found	E	\$5,670	\$0	\$5,670	\$0	\$5,670
	1083	Loulu ( <i>Pritchardia viscosa</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	440	Lousewort, Furbish ( <i>Pedicularis furbishiae</i> ) - Wherever found	E	\$96,220	\$0	\$96,220	\$0	\$96,220
	1215	Love grass, Fosberg's ( <i>Eragrostis fosbergii</i> ) - Wherever found	E	\$2,689	\$0	\$2,689	\$0	\$2,689
	1208	Lupine, clover (Tidestrom's) ( <i>Lupinus tidestromii</i> ) - Wherever found	E	\$0	\$3,100	\$3,100	\$0	\$3,100
	177	Lupine, Kincaid's ( <i>Lupinus sulphureus</i> ssp. <i>kincaidii</i> ) - Wherever found	T	\$268,930	\$371,519	\$640,449	\$0	\$640,449
	828	Lupine, Nipomo Mesa ( <i>Lupinus nipomensis</i> ) - Wherever found	E	\$18,000	\$300	\$18,300	\$0	\$18,300
	1239	Lupine, scrub ( <i>Lupinus aridorum</i> ) - Wherever found	E	\$0	\$0	\$0	\$1,500	\$1,500
	820	Mahoe ( <i>Alectryon macrococcus</i> ) - Wherever found	E	\$16,873	\$1,734	\$18,607	\$0	\$18,607
	934	Makou ( <i>Peucedanum sandwicense</i> ) - Wherever found	T	\$7,888	\$5,000	\$12,888	\$0	\$12,888
	776	Makou ( <i>Ranunculus hawaiiensis</i> ) - Wherever found	E	\$7,305	\$13,973	\$21,278	\$0	\$21,278
	719	Makou ( <i>Ranunculus mauiensis</i> ) - Wherever found	E	\$14,946	\$12,000	\$26,946	\$0	\$26,946
	1270	Malacothrix, island ( <i>Malacothrix squalida</i> ) - Wherever found	E	\$0	\$100	\$100	\$0	\$100
	913	Malacothrix, Santa Cruz Island ( <i>Malacothrix indecora</i> ) - Wherever found	E	\$2,700	\$11,050	\$13,750	\$0	\$13,750
	749	Mallow, Gierisch ( <i>Sphaeralcea gierischii</i> ) - Wherever found	E	\$4,500	\$19,700	\$24,200	\$0	\$24,200
	563	Mallow, Kern ( <i>Eremalche kernensis</i> ) - Wherever found	E	\$35,900	\$16,200	\$52,100	\$0	\$52,100

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1151	Manaca, palma de ( <i>Calyptronoma rivalis</i> ) - Wherever found	T	\$5,500	\$0	\$5,500	\$0	\$5,500
	747	Manioc, Walker's ( <i>Manihot walkerae</i> ) - Wherever found	E	\$24,500	\$0	\$24,500	\$0	\$24,500
	1226	Manzanita, Del Mar ( <i>Arctostaphylos glandulosa</i> <i>ssp. crassifolia</i> ) - Wherever found	E	\$2,100	\$200	\$2,300	\$0	\$2,300
	533	Manzanita, Franciscan ( <i>Arctostaphylos</i> <i>franciscana</i> ) - Wherever found	E	\$17,300	\$44,200	\$61,500	\$0	\$61,500
	653	Manzanita, Ione ( <i>Arctostaphylos myrtifolia</i> ) - Wherever found	T	\$28,900	\$6,703	\$35,603	\$0	\$35,603
	960	Manzanita, Morro ( <i>Arctostaphylos morroensis</i> ) - Wherever found	T	\$11,000	\$800	\$11,800	\$0	\$11,800
	1244	Manzanita, pallid ( <i>Arctostaphylos pallida</i> ) - Wherever found	T	\$0	\$1,300	\$1,300	\$0	\$1,300
	1092	Manzanita, Presidio ( <i>Arctostaphylos hookeri</i> <i>var. ravenii</i> ) - Wherever found	E	\$3,200	\$4,200	\$7,400	\$0	\$7,400
	847	Manzanita, Santa Rosa Island ( <i>Arctostaphylos</i> <i>confertiflora</i> ) - Wherever found	E	\$2,700	\$14,368	\$17,068	\$0	\$17,068
	1114	Ma`o hau hele, (=native yellow hibiscus) ( <i>Hibiscus brackenridgei</i> ) - Wherever found	E	\$6,007	\$500	\$6,507	\$0	\$6,507
	1083	Ma`oli`oli ( <i>Schiedea apokremnos</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1043	Ma`oli`oli ( <i>Schiedea hawaiiensis</i> ) - Wherever found	E	\$8,762	\$0	\$8,762	\$0	\$8,762
	1074	Ma`oli`oli ( <i>Schiedea kealiae</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1070	Ma`oli`oli ( <i>Schiedea pubescens</i> ) - Wherever found	E	\$8,172	\$0	\$8,172	\$0	\$8,172
	1152	Mapele ( <i>Cyrtandra cyaneoides</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1236	Mariposa lily, Tiburon ( <i>Calochortus</i> <i>tiburonensis</i> ) - Wherever found	T	\$1,700	\$0	\$1,700	\$0	\$1,700
	1159	Meadowfoam, Butte County ( <i>Limnanthes</i> <i>floccosa ssp. californica</i> ) - Wherever found	E	\$2,600	\$2,500	\$5,100	\$0	\$5,100
	859	Meadowfoam, large-flowered woolly ( <i>Limnanthes pumila ssp. grandiflora</i> ) - Wherever found	E	\$16,310	\$0	\$16,310	\$0	\$16,310

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	656	Meadowfoam, Sebastopol ( <i>Limnanthes vinculans</i> ) - Wherever found	E	\$11,200	\$23,800	\$35,000	\$0	\$35,000
	854	Meadowrue, Cooley's ( <i>Thalictrum cooleyi</i> ) - Wherever found	E	\$13,000	\$0	\$13,000	\$3,540	\$16,540
	837	Mehamehame ( <i>Flueggea neowawraea</i> ) - Wherever found	E	\$15,799	\$1,792	\$17,591	\$0	\$17,591
	810	Mesa-mint, Otay ( <i>Pogogyne nudiuscula</i> ) - Wherever found	E	\$7,300	\$12,100	\$19,400	\$0	\$19,400
	667	Mesa-mint, San Diego ( <i>Pogogyne abramsii</i> ) - Wherever found	E	\$7,700	\$24,890	\$32,590	\$0	\$32,590
	1154	Milkpea, Small's ( <i>Galactia smallii</i> ) - Wherever found	E	\$5,000	\$0	\$5,000	\$300	\$5,300
	561	Milk-vetch, Applegate's ( <i>Astragalus applegatei</i> ) - Wherever found	E	\$53,010	\$0	\$53,010	\$0	\$53,010
	874	Milk-vetch, Ash meadows ( <i>Astragalus phoenix</i> ) - Wherever found	T	\$15,600	\$0	\$15,600	\$0	\$15,600
	449	Milk-vetch, Braunton's ( <i>Astragalus brauntonii</i> ) - Wherever found	E	\$88,900	\$2,984	\$91,884	\$0	\$91,884
	1230	Milk-vetch, Clara Hunt's ( <i>Astragalus clarianus</i> ) - Wherever found	E	\$1,700	\$400	\$2,100	\$0	\$2,100
	792	Milk-vetch, Coachella Valley ( <i>Astragalus lentiginosus</i> var. <i>coachellae</i> ) - Wherever found	E	\$19,300	\$1,100	\$20,400	\$0	\$20,400
	1262	Milk-vetch, coastal dunes ( <i>Astragalus tener</i> var. <i>titi</i> ) - Wherever found	E	\$0	\$300	\$300	\$0	\$300
	1035	Milk-vetch, Cushenbury ( <i>Astragalus albens</i> ) - Wherever found	E	\$200	\$8,800	\$9,000	\$0	\$9,000
	564	Milk-vetch, Fish Slough ( <i>Astragalus lentiginosus</i> var. <i>piscinensis</i> ) - Wherever found	T	\$36,000	\$16,000	\$52,000	\$0	\$52,000
	1182	Milk-vetch, heliotrope ( <i>Astragalus montii</i> ) - Wherever found	T	\$0	\$4,000	\$4,000	\$0	\$4,000
	703	Milk-vetch, Holmgren ( <i>Astragalus holmgreniorum</i> ) - Wherever found	E	\$800	\$28,100	\$28,900	\$0	\$28,900
	560	Milk-vetch, Jesup's ( <i>Astragalus robbinsii</i> var. <i>jesupi</i> ) - Wherever found	E	\$40,070	\$0	\$40,070	\$13,000	\$53,070
	643	Milk-vetch, Lane Mountain ( <i>Astragalus jaegerianus</i> ) - Wherever found	E	\$0	\$37,200	\$37,200	\$0	\$37,200

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	705	Milk-vetch, Mancos ( <i>Astragalus humillimus</i> ) - Wherever found	E	\$15,300	\$13,300	\$28,600	\$0	\$28,600
	1000	Milkvetch, Osterhout ( <i>Astragalus osterhoutii</i> ) - Wherever found	E	\$400	\$9,000	\$9,400	\$715	\$10,115
	276	Milk-vetch, Peirson's ( <i>Astragalus magdalenae</i> var. <i>peirsonii</i> ) - Wherever found	T	\$18,100	\$261,200	\$279,300	\$0	\$279,300
	452	Milk-vetch, Sentry ( <i>Astragalus cremnophylax</i> var. <i>cremnophylax</i> ) - Wherever found	E	\$4,000	\$86,000	\$90,000	\$0	\$90,000
	742	Milk-vetch, Shivwits ( <i>Astragalus ampullarioides</i> ) - Wherever found	E	\$0	\$24,925	\$24,925	\$0	\$24,925
	517	Milk-vetch, triple-ribbed ( <i>Astragalus tricarinatus</i> ) - Wherever found	E	\$4,600	\$59,500	\$64,100	\$0	\$64,100
	810	Milk-vetch, Ventura Marsh ( <i>Astragalus pycnostachyus</i> var. <i>lanosissimus</i> ) - Wherever found	E	\$19,400	\$0	\$19,400	\$0	\$19,400
	352	Milkweed, Mead's ( <i>Asclepias meadii</i> ) - Wherever found	T	\$65,300	\$43,541	\$108,841	\$45,000	\$153,841
	688	Milkweed, Welsh's ( <i>Asclepias welshii</i> ) - Wherever found	T	\$300	\$30,200	\$30,500	\$0	\$30,500
	1126	Mint, Garrett's ( <i>Dicerandra christmanii</i> ) - Wherever found	E	\$5,500	\$0	\$5,500	\$800	\$6,300
	921	Mint, Lakela's ( <i>Dicerandra immaculata</i> ) - Wherever found	E	\$1,000	\$0	\$1,000	\$12,500	\$13,500
	1267	Mint, longspurred ( <i>Dicerandra cornutissima</i> ) - Wherever found	E	\$0	\$0	\$0	\$200	\$200
	710	Mint, scrub ( <i>Dicerandra frutescens</i> ) - Wherever found	E	\$900	\$21,700	\$22,600	\$5,500	\$28,100
	973	Monardella, willowy ( <i>Monardella viminea</i> ) - Wherever found	E	\$4,400	\$7,000	\$11,400	\$0	\$11,400
	650	Monkey-flower, Michigan ( <i>Mimulus michiganensis</i> ) - Wherever found	E	\$34,500	\$1,365	\$35,865	\$0	\$35,865
	840	Monkshood, northern wild ( <i>Aconitum noveboracense</i> ) - Wherever found	T	\$14,700	\$1,779	\$16,479	\$1,000	\$17,479
	762	Morning-glory, Stebbins' ( <i>Calystegia stebbinsii</i> ) - Wherever found	E	\$4,400	\$18,700	\$23,100	\$0	\$23,100
	542	Mountainbalm, Indian Knob ( <i>Eriodictyon altissimum</i> ) - Wherever found	E	\$58,400	\$200	\$58,600	\$0	\$58,600



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1270	Mountain-mahogany, Catalina Island (Cercocarpus traskiae) - Wherever found	E	\$0	\$100	\$100	\$0	\$100
	1025	Mustard, Carter's (Warea carteri) - Wherever found	E	\$2,000	\$0	\$2,000	\$7,500	\$9,500
	860	Mustard, Penland alpine fen (Eutrema penlandii) - Wherever found	T	\$7,300	\$9,000	\$16,300	\$0	\$16,300
	1214	Mustard, slender-petaled (Thelypodium stenopetalum) - Wherever found	E	\$2,700	\$0	\$2,700	\$0	\$2,700
	1152	Naenae (Dubautia kalalaensis) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1083	Naenae (Dubautia kenwoodii) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1144	Na`ena`e (Dubautia herbstobatae) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1152	Na`ena`e (Dubautia imbricata ssp. imbricata) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1152	Na`ena`e (Dubautia pauciflorula) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	985	Na`ena`e (Dubautia plantaginea ssp. humilis) - Wherever found	E	\$10,778	\$0	\$10,778	\$0	\$10,778
	1152	Na`ena`e (Dubautia plantaginea ssp. magnifolia) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1152	Na`ena`e (Dubautia waialealae) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1083	Nani wai`ale`ale (Viola kauaiensis var. wahiawaensis) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1074	Nanu (Gardenia manni) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	787	Nanu (Gardenia remyi) - Wherever found	E	\$8,555	\$12,000	\$20,555	\$0	\$20,555
	566	Naupaka, dwarf (Scaevola coriacea) - Wherever found	E	\$46,230	\$5,000	\$51,230	\$0	\$51,230
	1247	Navarretia, few-flowered (Navarretia leucocephala ssp. pauciflora (=N. pauciflora)) - Wherever found	E	\$0	\$1,200	\$1,200	\$0	\$1,200
	973	Navarretia, many-flowered (Navarretia leucocephala ssp. pliantha) - Wherever found	E	\$2,700	\$8,700	\$11,400	\$0	\$11,400
	329	Navarretia, spreading (Navarretia fossalis) - Wherever found	T	\$8,100	\$173,600	\$181,700	\$0	\$181,700

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1155	Nehe ( <i>Lipochaeta fauriei</i> ) - Wherever found	E	\$5,265	\$0	\$5,265	\$0	\$5,265
	1097	Nehe ( <i>Lipochaeta lobata</i> var. <i>leptophylla</i> ) - Wherever found	E	\$5,417	\$1,792	\$7,209	\$0	\$7,209
	1152	Nehe ( <i>Lipochaeta micrantha</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	908	Nehe ( <i>Lipochaeta venosa</i> ) - Wherever found	E	\$13,953	\$0	\$13,953	\$0	\$13,953
	1152	Nehe ( <i>Lipochaeta waimeaensis</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1061	Nehe ( <i>Melanthera kamolensis</i> ) - Wherever found	E	\$8,307	\$0	\$8,307	\$0	\$8,307
	1074	Nehe ( <i>Melanthera tenuifolia</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1149	Nioi ( <i>Eugenia koolauensis</i> ) - Wherever found	E	\$5,602	\$0	\$5,602	\$0	\$5,602
	578	Niterwort, Amargosa ( <i>Nitrophila mohavensis</i> ) - Wherever found	E	\$12,400	\$36,200	\$48,600	\$0	\$48,600
	1030	No Common Name ( <i>Abutilon eremitopetalum</i> ) - Wherever found	E	\$9,321	\$0	\$9,321	\$0	\$9,321
	1012	No Common Name ( <i>Abutilon sandwicense</i> ) - Wherever found	E	\$8,073	\$1,734	\$9,807	\$0	\$9,807
	400	No Common Name ( <i>Achyranthes mutica</i> ) - Wherever found	E	\$123,569	\$0	\$123,569	\$0	\$123,569
	1210	No Common Name ( <i>Agave eggersiana</i> ) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1080	No Common Name ( <i>Amaranthus brownii</i> ) - Wherever found	E	\$7,952	\$0	\$7,952	\$0	\$7,952
	921	No Common Name ( <i>Aristida chaseae</i> ) - Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	1182	No Common Name ( <i>Auerodendron pauciflorum</i> ) - Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	1070	No Common Name ( <i>Bidens hillebrandiana</i> ssp. <i>hillebrandiana</i> ) - Wherever found	E	\$8,172	\$0	\$8,172	\$0	\$8,172
	817	No Common Name ( <i>Bonamia menziesii</i> ) - Wherever found	E	\$17,120	\$1,761	\$18,881	\$0	\$18,881
	1151	No Common Name ( <i>Calypttranthes thomasiana</i> ) - Wherever found	E	\$5,500	\$0	\$5,500	\$0	\$5,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1077	No Common Name ( <i>Catesbaea melanocarpa</i> ) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	1210	No Common Name ( <i>Chamaecrista glandulosa</i> var. <i>mirabilis</i> ) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	545	No Common Name ( <i>Cordia bellonis</i> ) - Wherever found	E	\$58,000	\$0	\$58,000	\$0	\$58,000
	1151	No Common Name ( <i>Cranichis ricartii</i> ) - Wherever found	E	\$5,500	\$0	\$5,500	\$0	\$5,500
	1131	No Common Name ( <i>Cyanea kauaulaensis</i> ) - Wherever found	E	\$6,106	\$0	\$6,106	\$0	\$6,106
	1078	No Common Name ( <i>Cyperus fauriei</i> ) - Wherever found	E	\$7,989	\$0	\$7,989	\$0	\$7,989
	1141	No Common Name ( <i>Cyperus neokunthianus</i> ) - Wherever found	E	\$5,853	\$0	\$5,853	\$0	\$5,853
	1083	No Common Name ( <i>Cyperus pennatiformis</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	569	No Common Name ( <i>Cyrtandra nanawaleensis</i> ) - Wherever found	E	\$6,106	\$44,254	\$50,360	\$0	\$50,360
	1056	No Common Name ( <i>Cyrtandra wagneri</i> ) - Wherever found	E	\$8,425	\$0	\$8,425	\$0	\$8,425
	1190	No Common Name ( <i>Daphnopsis helleriana</i> ) - Wherever found	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	912	No Common Name ( <i>Delissea rhytidosperma</i> ) - Wherever found	E	\$13,820	\$0	\$13,820	\$0	\$13,820
	1078	No Common Name ( <i>Delissea undulata</i> ) - Wherever found	E	\$7,989	\$0	\$7,989	\$0	\$7,989
	461	No Common Name ( <i>Dendrobium guamense</i> ) - Wherever found	T	\$52,917	\$33,773	\$86,690	\$0	\$86,690
	600	No Common Name ( <i>Eugenia bryanii</i> ) - Wherever found	E	\$32,840	\$11,386	\$44,226	\$0	\$44,226
	851	No Common Name ( <i>Eugenia woodburyana</i> ) - Wherever found	E	\$16,500	\$300	\$16,800	\$0	\$16,800
	1117	No Common Name ( <i>Festuca hawaiiensis</i> ) - Wherever found	E	\$6,443	\$0	\$6,443	\$0	\$6,443
	989	No Common Name ( <i>Festuca molokaiensis</i> ) - Wherever found	E	\$10,629	\$0	\$10,629	\$0	\$10,629
	523	No Common Name ( <i>Geocarpon minimum</i> ) - Wherever found	T	\$44,811	\$16,587	\$61,398	\$1,600	\$62,998

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1115	No Common Name ( <i>Gesneria pauciflora</i> ) - Wherever found	T	\$6,500	\$0	\$6,500	\$0	\$6,500
	1232	No Common Name ( <i>Gonocalyx concolor</i> ) - Wherever found	E	\$2,000	\$0	\$2,000	\$0	\$2,000
	1068	No Common Name ( <i>Gouania hillebrandii</i> ) - Wherever found	E	\$8,205	\$0	\$8,205	\$0	\$8,205
	1148	No Common Name ( <i>Gouania meyenii</i> ) - Wherever found	E	\$5,652	\$0	\$5,652	\$0	\$5,652
	1084	No Common Name ( <i>Gouania vitifolia</i> ) - Wherever found	E	\$7,634	\$0	\$7,634	\$0	\$7,634
	1149	No Common Name ( <i>Hesperomannia arborescens</i> ) - Wherever found	E	\$5,602	\$0	\$5,602	\$0	\$5,602
	1144	No Common Name ( <i>Hesperomannia arbuscula</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1083	No Common Name ( <i>Hesperomannia lydgatei</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1135	No Common Name ( <i>Ilex sintenisii</i> ) - Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	1144	No Common Name ( <i>Kadua degeneri</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1005	No Common Name ( <i>Kadua hauptuensis</i> ) - Wherever found	E	\$9,909	\$0	\$9,909	\$0	\$9,909
	1008	No Common Name ( <i>Kadua parvula</i> ) - Wherever found	E	\$8,073	\$1,792	\$9,865	\$0	\$9,865
	1152	No Common Name ( <i>Kadua st.-johnii</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1165	No Common Name ( <i>Keysseria (=Lagenifera) erici</i> ) - Wherever found	E	\$4,907	\$0	\$4,907	\$0	\$4,907
	1218	No Common Name ( <i>Keysseria (=Lagenifera) helenae</i> ) - Wherever found	E	\$2,588	\$0	\$2,588	\$0	\$2,588
	1141	No Common Name ( <i>Labordia lorenciana</i> ) - Wherever found	E	\$5,853	\$0	\$5,853	\$0	\$5,853
	632	No Common Name ( <i>Lepanthes eltoroensis</i> ) - Wherever found	E	\$39,000	\$0	\$39,000	\$0	\$39,000
	1141	No Common Name ( <i>Lepidium orbiculare</i> ) - Wherever found	E	\$5,853	\$0	\$5,853	\$0	\$5,853
	1163	No Common Name ( <i>Leptocereus grantianus</i> ) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1074	No Common Name ( <i>Lobelia koolauensis</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1152	No Common Name ( <i>Lobelia monostachya</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1008	No Common Name ( <i>Lobelia niihauensis</i> ) - Wherever found	E	\$8,073	\$1,792	\$9,865	\$0	\$9,865
	1144	No Common Name ( <i>Lobelia oahuensis</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	931	No Common Name ( <i>Lyonia truncata</i> var. <i>proctorii</i> ) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1152	No Common Name ( <i>Lysimachia filifolia</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1152	No Common Name ( <i>Lysimachia iniki</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	985	No Common Name ( <i>Lysimachia lydgatei</i> ) - Wherever found	E	\$10,778	\$0	\$10,778	\$0	\$10,778
	985	No Common Name ( <i>Lysimachia maxima</i> ) - Wherever found	E	\$10,778	\$0	\$10,778	\$0	\$10,778
	1083	No Common Name ( <i>Lysimachia pendens</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1083	No Common Name ( <i>Lysimachia scopulensis</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1083	No Common Name ( <i>Lysimachia venosa</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	521	No Common Name ( <i>Maesa walkeri</i> ) - Wherever found	T	\$41,129	\$22,270	\$63,399	\$0	\$63,399
	1050	No Common Name ( <i>Mitracarpus maxwelliae</i> ) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1050	No Common Name ( <i>Mitracarpus polycladus</i> ) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	1210	No Common Name ( <i>Myrcia paganii</i> ) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1085	No Common Name ( <i>Neraudia angulata</i> ) - Wherever found	E	\$5,754	\$1,792	\$7,546	\$0	\$7,546
	732	No Common Name ( <i>Neraudia ovata</i> ) - Wherever found	E	\$13,222	\$12,435	\$25,657	\$0	\$25,657
	985	No Common Name ( <i>Neraudia sericea</i> ) - Wherever found	E	\$10,778	\$0	\$10,778	\$0	\$10,778

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	520	No Common Name ( <i>Nervilia jacksoniae</i> ) - Wherever found	T	\$41,466	\$22,270	\$63,736	\$0	\$63,736
	657	No Common Name ( <i>Nesogenes rotensis</i> ) - Wherever found	E	\$34,671	\$0	\$34,671	\$0	\$34,671
	726	No Common Name ( <i>Osmoxylon mariannense</i> ) - Wherever found	E	\$26,488	\$0	\$26,488	\$0	\$26,488
	565	No Common Name ( <i>Phyllanthus saffordii</i> ) - Wherever found	E	\$30,883	\$21,020	\$51,903	\$0	\$51,903
	488	No Common Name ( <i>Phyllostegia bracteata</i> ) - Wherever found	E	\$14,290	\$61,000	\$75,290	\$0	\$75,290
	499	No Common Name ( <i>Phyllostegia brevidens</i> ) - Wherever found	E	\$10,196	\$60,000	\$70,196	\$0	\$70,196
	1022	No Common Name ( <i>Phyllostegia floribunda</i> ) - Wherever found	E	\$9,523	\$0	\$9,523	\$0	\$9,523
	979	No Common Name ( <i>Phyllostegia glabra</i> var. <i>lanaiensis</i> ) - Wherever found	E	\$11,031	\$0	\$11,031	\$0	\$11,031
	500	No Common Name ( <i>Phyllostegia haliakalae</i> ) - Wherever found	E	\$9,099	\$61,000	\$70,099	\$0	\$70,099
	1070	No Common Name ( <i>Phyllostegia helleri</i> ) - Wherever found	E	\$8,172	\$0	\$8,172	\$0	\$8,172
	1074	No Common Name ( <i>Phyllostegia hirsuta</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	979	No Common Name ( <i>Phyllostegia hispida</i> ) - Wherever found	E	\$11,031	\$0	\$11,031	\$0	\$11,031
	1074	No Common Name ( <i>Phyllostegia kaalaensis</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1083	No Common Name ( <i>Phyllostegia knudsenii</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	979	No Common Name ( <i>Phyllostegia mannii</i> ) - Wherever found	E	\$11,031	\$0	\$11,031	\$0	\$11,031
	1074	No Common Name ( <i>Phyllostegia mollis</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1058	No Common Name ( <i>Phyllostegia parviflora</i> ) - Wherever found	E	\$8,410	\$0	\$8,410	\$0	\$8,410
	1056	No Common Name ( <i>Phyllostegia pilosa</i> ) - Wherever found	E	\$8,425	\$0	\$8,425	\$0	\$8,425
	1083	No Common Name ( <i>Phyllostegia renovans</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1111	No Common Name ( <i>Phyllostegia stachyoides</i> ) - Wherever found	E	\$6,106	\$500	\$6,606	\$0	\$6,606

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1074	No Common Name ( <i>Phyllostegia velutina</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1152	No Common Name ( <i>Phyllostegia waimeae</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1060	No Common Name ( <i>Phyllostegia warshaueri</i> ) - Wherever found	E	\$8,326	\$0	\$8,326	\$0	\$8,326
	939	No Common Name ( <i>Phyllostegia wawrana</i> ) - Wherever found	E	\$12,639	\$0	\$12,639	\$0	\$12,639
	756	No Common Name ( <i>Platanthera holochila</i> ) - Wherever found	E	\$11,411	\$12,000	\$23,411	\$0	\$23,411
	1083	No Common Name ( <i>Platydesma cornuta</i> var. <i>cornuta</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	988	No Common Name ( <i>Platydesma cornuta</i> var. <i>decurrens</i> ) - Wherever found	E	\$8,073	\$2,572	\$10,645	\$0	\$10,645
	1117	No Common Name ( <i>Platydesma remyi</i> ) - Wherever found	E	\$6,443	\$0	\$6,443	\$0	\$6,443
	1152	No Common Name ( <i>Poa siphonoglossa</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1011	No Common Name ( <i>Polyscias bisattenuata</i> ) - Wherever found	E	\$9,820	\$0	\$9,820	\$0	\$9,820
	1083	No Common Name ( <i>Polyscias flynnii</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1152	No Common Name ( <i>Polyscias lydgatei</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1083	No Common Name ( <i>Polyscias racemosa</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1044	No Common Name ( <i>Remya kauaiensis</i> ) - Wherever found	E	\$8,750	\$0	\$8,750	\$0	\$8,750
	1118	No Common Name ( <i>Remya montgomeryi</i> ) - Wherever found	E	\$6,431	\$0	\$6,431	\$0	\$6,431
	1149	No Common Name ( <i>Sanicula mariversa</i> ) - Wherever found	E	\$5,602	\$0	\$5,602	\$0	\$5,602
	1041	No Common Name ( <i>Sanicula purpurea</i> ) - Wherever found	E	\$8,796	\$0	\$8,796	\$0	\$8,796
	591	No Common Name ( <i>Sanicula sandwicensis</i> ) - Wherever found	E	\$8,359	\$37,973	\$46,332	\$0	\$46,332
	1141	No Common Name ( <i>Santalum involutum</i> ) - Wherever found	E	\$5,853	\$0	\$5,853	\$0	\$5,853

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1152	No Common Name ( <i>Schiedea attenuata</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	923	No Common Name ( <i>Schiedea diffusa</i> ssp. <i>macraei</i> ) - Wherever found	E	\$8,425	\$5,000	\$13,425	\$0	\$13,425
	766	No Common Name ( <i>Schiedea diffusa</i> subsp. <i>diffusa</i> ) - Wherever found	E	\$10,931	\$12,000	\$22,931	\$0	\$22,931
	495	No Common Name ( <i>Schiedea haleakalensis</i> ) - Wherever found	E	\$11,452	\$61,000	\$72,452	\$0	\$72,452
	1083	No Common Name ( <i>Schiedea helleri</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1036	No Common Name ( <i>Schiedea hookeri</i> ) - Wherever found	E	\$6,427	\$2,572	\$8,999	\$0	\$8,999
	1141	No Common Name ( <i>Schiedea jacobii</i> ) - Wherever found	E	\$5,853	\$0	\$5,853	\$0	\$5,853
	1074	No Common Name ( <i>Schiedea kaalae</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	912	No Common Name ( <i>Schiedea kauaiensis</i> ) - Wherever found	E	\$13,820	\$0	\$13,820	\$0	\$13,820
	1018	No Common Name ( <i>Schiedea laui</i> ) - Wherever found	E	\$9,615	\$0	\$9,615	\$0	\$9,615
	1051	No Common Name ( <i>Schiedea lydgatei</i> ) - Wherever found	E	\$8,459	\$0	\$8,459	\$0	\$8,459
	1051	No Common Name ( <i>Schiedea membranacea</i> ) - Wherever found	E	\$8,459	\$0	\$8,459	\$0	\$8,459
	1057	No Common Name ( <i>Schiedea nuttallii</i> ) - Wherever found	E	\$8,420	\$0	\$8,420	\$0	\$8,420
	1074	No Common Name ( <i>Schiedea obovata</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1039	No Common Name ( <i>Schiedea salicaria</i> ) - Wherever found	E	\$8,895	\$0	\$8,895	\$0	\$8,895
	1051	No Common Name ( <i>Schiedea sarmentosa</i> ) - Wherever found	E	\$8,459	\$0	\$8,459	\$0	\$8,459
	1152	No Common Name ( <i>Schiedea spergulina</i> var. <i>leiopoda</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	996	No Common Name ( <i>Schiedea spergulina</i> var. <i>spergulina</i> ) - Wherever found	T	\$10,320	\$0	\$10,320	\$0	\$10,320
	1144	No Common Name ( <i>Schiedea trinervis</i> ) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	958	No Common Name ( <i>Schiedea verticillata</i> ) - Wherever found	E	\$11,906	\$0	\$11,906	\$0	\$11,906
	1152	No Common Name ( <i>Schiedea viscosa</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1115	No Common Name ( <i>Schoepfia arenaria</i> ) - Wherever found	T	\$6,500	\$0	\$6,500	\$0	\$6,500
	1024	No Common Name ( <i>Sicyos lanceoloideus</i> ) - Wherever found	E	\$8,509	\$1,000	\$9,509	\$0	\$9,509
	1061	No Common Name ( <i>Silene alexandri</i> ) - Wherever found	E	\$8,307	\$0	\$8,307	\$0	\$8,307
	803	No Common Name ( <i>Silene hawaiiensis</i> ) - Wherever found	T	\$8,743	\$11,000	\$19,743	\$0	\$19,743
	729	No Common Name ( <i>Silene lanceolata</i> ) - Wherever found	E	\$6,091	\$20,000	\$26,091	\$0	\$26,091
	1078	No Common Name ( <i>Silene perlmanii</i> ) - Wherever found	E	\$5,417	\$2,572	\$7,989	\$0	\$7,989
	697	No Common Name ( <i>Spermolepis hawaiiensis</i> ) - Wherever found	E	\$5,827	\$23,801	\$29,628	\$0	\$29,628
	384	No Common Name ( <i>Stenogyne angustifolia</i> var. <i>angustifolia</i> ) - Wherever found	E	\$111,067	\$20,000	\$131,067	\$0	\$131,067
	443	No Common Name ( <i>Stenogyne bifida</i> ) - Wherever found	E	\$95,662	\$0	\$95,662	\$0	\$95,662
	1083	No Common Name ( <i>Stenogyne campanulata</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1043	No Common Name ( <i>Stenogyne cranwelliae</i> ) - Wherever found	E	\$8,762	\$0	\$8,762	\$0	\$8,762
	1083	No Common Name ( <i>Stenogyne kaalae</i> ssp. <i>sherffii</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	1074	No Common Name ( <i>Stenogyne kanehoana</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	893	No Common Name ( <i>Stenogyne kauaulaensis</i> ) - Wherever found	E	\$14,647	\$0	\$14,647	\$0	\$14,647
	446	No Common Name ( <i>Stenogyne kealiae</i> ) - Wherever found	E	\$94,686	\$0	\$94,686	\$0	\$94,686
	409	No Common Name ( <i>Tabernaemontana rotensis</i> ) - Wherever found	T	\$51,225	\$59,071	\$110,296	\$0	\$110,296
	1151	No Common Name ( <i>Ternstroemia subsessilis</i> ) - Wherever found	E	\$5,500	\$0	\$5,500	\$0	\$5,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	731	No Common Name (Tetramolopium arenarium) - Wherever found	E	\$5,855	\$20,000	\$25,855	\$0	\$25,855
	1074	No Common Name (Tetramolopium filiforme) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1060	No Common Name (Tetramolopium lepidotum ssp. lepidotum) - Wherever found	E	\$5,754	\$2,572	\$8,326	\$0	\$8,326
	930	No Common Name (Tetramolopium remyi) - Wherever found	E	\$13,059	\$0	\$13,059	\$0	\$13,059
	419	No Common Name (Tetramolopium rockii) - Wherever found	T	\$97,981	\$5,000	\$102,981	\$0	\$102,981
	1081	No Common Name (Tinospora homosepala) - Wherever found	E	\$7,936	\$0	\$7,936	\$0	\$7,936
	1169	No Common Name (Trematolobelia singularis) - Wherever found	E	\$4,626	\$0	\$4,626	\$0	\$4,626
	418	No Common Name (Tuberolabium guamense) - Wherever found	T	\$58,746	\$44,680	\$103,426	\$0	\$103,426
	1101	No Common Name (Varronia rupicola) - Wherever found	T	\$7,000	\$0	\$7,000	\$0	\$7,000
	931	No Common Name (Vernonia proctorii) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	811	No Common Name (Vigna o-wahuensis) - Wherever found	E	\$19,381	\$0	\$19,381	\$0	\$19,381
	1152	No Common Name (Viola helenae) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	979	No Common Name (Viola lanaiensis) - Wherever found	E	\$11,031	\$0	\$11,031	\$0	\$11,031
	1144	No Common Name (Viola oahuensis) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1141	No Common Name (Wikstroemia skottsbergiana) -	E	\$5,853	\$0	\$5,853	\$0	\$5,853
	766	No Common Name (Wikstroemia villosa) - Wherever found	E	\$10,931	\$12,000	\$22,931	\$0	\$22,931
	1152	No Common Name (Xylosma crenatum) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	779	Nohoanu (Geranium hanaense) - Wherever found	E	\$9,099	\$12,000	\$21,099	\$0	\$21,099
	1010	Nohoanu (Geranium hillebrandii) - Wherever found	E	\$9,827	\$0	\$9,827	\$0	\$9,827

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1016	Nohoanu ( <i>Geranium kauaiense</i> ) - Wherever found	E	\$9,669	\$0	\$9,669	\$0	\$9,669
	765	Nohoanu ( <i>Geranium multiflorum</i> ) - Wherever found	E	\$10,965	\$12,000	\$22,965	\$0	\$22,965
	766	Nui, haha ( <i>Cyanea horrida</i> ) - Wherever found	E	\$10,931	\$12,000	\$22,931	\$0	\$22,931
	925	Oha ( <i>Delissea subcordata</i> ) - Wherever found	E	\$13,313	\$0	\$13,313	\$0	\$13,313
	465	Ohai ( <i>Sesbania tomentosa</i> ) - Wherever found	E	\$82,734	\$2,700	\$85,434	\$0	\$85,434
	1155	`Oha wai ( <i>Clermontia drepanomorpha</i> ) - Wherever found	E	\$5,265	\$0	\$5,265	\$0	\$5,265
	914	`Oha wai ( <i>Clermontia lindseyana</i> ) - Wherever found	E	\$6,760	\$6,973	\$13,733	\$0	\$13,733
	1046	`Oha wai ( <i>Clermontia oblongifolia</i> ssp. <i>brevipes</i> ) - Wherever found	E	\$8,712	\$0	\$8,712	\$0	\$8,712
	855	`Oha wai ( <i>Clermontia oblongifolia</i> ssp. <i>mauiensis</i> ) - Wherever found	E	\$16,538	\$0	\$16,538	\$0	\$16,538
	965	`Oha wai ( <i>Clermontia peleana</i> ) - Wherever found	E	\$6,680	\$5,000	\$11,680	\$0	\$11,680
	1060	`Oha wai ( <i>Clermontia pyrularia</i> ) - Wherever found	E	\$8,326	\$0	\$8,326	\$0	\$8,326
	584	`Oha wai ( <i>Clermontia samuelii</i> ) - Wherever found	E	\$11,452	\$36,000	\$47,452	\$0	\$47,452
	579	`Ohe ( <i>Joinvillea ascendens ascendens</i> ) - Wherever found	E	\$12,281	\$36,000	\$48,281	\$0	\$48,281
	1074	`Ohe`ohe ( <i>Polyscias gymnocarpa</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1155	Olulu ( <i>Brighamia insignis</i> ) - Wherever found	E	\$5,265	\$0	\$5,265	\$0	\$5,265
	1244	Onion, Munz's ( <i>Allium munzii</i> ) - Wherever found	E	\$1,200	\$100	\$1,300	\$0	\$1,300
	1149	Opuhe ( <i>Urera kaalae</i> ) - Wherever found	E	\$5,602	\$0	\$5,602	\$0	\$5,602
	445	Orchid, eastern prairie fringed ( <i>Platanthera leucophaea</i> ) - Wherever found	T	\$28,941	\$27,076	\$56,017	\$39,000	\$95,017
	391	Orchid, western prairie fringed ( <i>Platanthera praeclara</i> ) - Wherever found	T	\$42,229	\$62,656	\$104,885	\$22,415	\$127,300
	576	Orchid, white fringeless ( <i>Platanthera integrilabia</i> ) -	T	\$29,000	\$14,156	\$43,156	\$6,000	\$49,156

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	798	Orcutt grass, California ( <i>Orcuttia californica</i> ) - Wherever found	E	\$7,200	\$12,800	\$20,000	\$0	\$20,000
	885	Orcutt grass, hairy ( <i>Orcuttia pilosa</i> ) - Wherever found	E	\$11,200	\$4,056	\$15,256	\$0	\$15,256
	792	Orcutt grass, Sacramento ( <i>Orcuttia viscida</i> ) - Wherever found	E	\$15,900	\$4,500	\$20,400	\$0	\$20,400
	764	Orcutt grass, San Joaquin ( <i>Orcuttia inaequalis</i> ) - Wherever found	T	\$16,200	\$6,778	\$22,978	\$0	\$22,978
	753	Orcutt grass, slender ( <i>Orcuttia tenuis</i> ) - Wherever found	T	\$12,800	\$10,800	\$23,600	\$0	\$23,600
	741	Owl's-clover, fleshy ( <i>Castilleja campestris</i> ssp. <i>succulenta</i> ) - Wherever found	T	\$20,000	\$4,930	\$24,930	\$0	\$24,930
	531	<i>Oxytheca</i> , Cushenbury ( <i>Oxytheca parishii</i> var. <i>goodmaniana</i> ) - Wherever found	E	\$230	\$61,800	\$62,030	\$0	\$62,030
	1152	Pa`iniu ( <i>Astelia waialealae</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1262	Paintbrush, ash-grey ( <i>Castilleja cinerea</i> ) - Wherever found	T	\$0	\$300	\$300	\$0	\$300
	169	Paintbrush, golden ( <i>Castilleja levisecta</i> ) - Wherever found	T	\$505,413	\$191,346	\$696,759	\$2,600	\$699,359
	341	Paintbrush, San Clemente Island ( <i>Castilleja</i> <i>grisea</i> ) - Wherever found	T	\$3,800	\$158,417	\$162,217	\$0	\$162,217
	995	Paintbrush, soft-leaved ( <i>Castilleja mollis</i> ) - Wherever found	E	\$2,700	\$7,674	\$10,374	\$0	\$10,374
	948	Paintbrush, Tiburon ( <i>Castilleja affinis</i> ssp. <i>neglecta</i> ) - Wherever found	E	\$11,200	\$1,100	\$12,300	\$0	\$12,300
	1115	Palo colorado ( <i>Ternstroemia luquillensis</i> ) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1115	Palo de jazmin ( <i>Styrax portoricensis</i> ) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1088	Palo de nigua ( <i>Cornutia obovata</i> ) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1115	Palo de ramon ( <i>Banara vanderbiltii</i> ) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1101	Palo de rosa ( <i>Ottoschulzia rhodoxylon</i> ) - Wherever found	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	1046	Pamakani ( <i>Tetramolopium capillare</i> ) - Wherever found	E	\$8,712	\$0	\$8,712	\$0	\$8,712

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1060	Pamakani ( <i>Viola chamissoniana</i> ssp. <i>chamissoniana</i> ) - Wherever found	E	\$5,754	\$2,572	\$8,326	\$0	\$8,326
	986	Panicgrass, Carter's ( <i>Panicum fauriei</i> var. <i>carteri</i> ) - Wherever found	E	\$5,701	\$5,000	\$10,701	\$0	\$10,701
	1030	Papala ( <i>Charpentiera densiflora</i> ) - Wherever found	E	\$9,321	\$0	\$9,321	\$0	\$9,321
	658	Paudedo ( <i>Hedyotis megalantha</i> ) - Wherever found	E	\$34,368	\$0	\$34,368	\$0	\$34,368
	1221	Pawpaw, beautiful ( <i>Deeringothamnus pulchellus</i> ) - Wherever found	E	\$900	\$100	\$1,000	\$1,500	\$2,500
	1050	Pawpaw, four-petal ( <i>Asimina tetramera</i> ) - Wherever found	E	\$900	\$6,500	\$7,400	\$1,100	\$8,500
	1003	Pawpaw, Rugel's ( <i>Deeringothamnus rugelii</i> ) - Wherever found	E	\$4,500	\$0	\$4,500	\$5,500	\$10,000
	1224	Pea, Big Pine partridge ( <i>Chamaecrista lineata keyensis</i> ) - Wherever found	E	\$700	\$0	\$700	\$1,700	\$2,400
	921	Pelos del diablo ( <i>Aristida portoricensis</i> ) - Wherever found	E	\$13,500	\$0	\$13,500	\$0	\$13,500
	703	Penny-cress, Kneeland Prairie ( <i>Thlaspi californicum</i> ) - Wherever found	E	\$28,900	\$0	\$28,900	\$0	\$28,900
	676	Pennyroyal, Todsens's ( <i>Hedeoma todsenii</i> ) - Wherever found	E	\$0	\$31,500	\$31,500	\$0	\$31,500
	335	Penstemon, blowout ( <i>Penstemon haydenii</i> ) - Wherever found	E	\$43,200	\$121,049	\$164,249	\$4,800	\$169,049
	671	Pentachaeta, Lyon's ( <i>Pentachaeta lyonii</i> ) - Wherever found	E	\$32,200	\$100	\$32,300	\$0	\$32,300
	1224	Pentachaeta, white-rayed ( <i>Pentachaeta bellidiflora</i> ) - Wherever found	E	\$0	\$2,400	\$2,400	\$0	\$2,400
	1163	Peperomia, Wheeler's ( <i>Peperomia wheeleri</i> ) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	194	Peppergrass, Slickspot ( <i>Lepidium papilliferum</i> )	T	\$153,508	\$408,500	\$562,008	\$0	\$562,008
	795	Phacelia, clay ( <i>Phacelia argillacea</i> ) - Wherever found	E	\$0	\$20,300	\$20,300	\$0	\$20,300
	806	Phacelia, DeBeque ( <i>Phacelia submutica</i> ) - Wherever found	T	\$3,100	\$15,200	\$18,300	\$1,340	\$19,640
	962	Phacelia, island ( <i>Phacelia insularis</i> ssp. <i>insularis</i> ) - Wherever found	E	\$2,700	\$9,079	\$11,779	\$0	\$11,779

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	898	Phacelia, North Park ( <i>Phacelia formosula</i> ) - Wherever found	E	\$800	\$11,500	\$12,300	\$2,165	\$14,465
	850	Phlox, Texas trailing ( <i>Phlox nivalis</i> ssp. <i>texensis</i> ) - Wherever found	E	\$12,000	\$5,000	\$17,000	\$0	\$17,000
	202	Phlox, Yreka ( <i>Phlox hirsuta</i> ) - Wherever found	E	\$531,200	\$2,000	\$533,200	\$0	\$533,200
	638	Pigeon wings ( <i>Clitoria fragrans</i> ) - Wherever found	T	\$2,000	\$30,365	\$32,365	\$6,000	\$38,365
	922	Pilo ( <i>Kadua laxiflora</i> ) - Wherever found	E	\$13,491	\$0	\$13,491	\$0	\$13,491
	1083	Pilo kea lau li'i ( <i>Platydesma rostrata</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	648	Pinkroot, gentian ( <i>Spigelia gentianoides</i> ) - Wherever found	E	\$12,500	\$23,819	\$36,319	\$200	\$36,519
	351	Pink, swamp ( <i>Helonias bullata</i> ) - Wherever found	T	\$124,271	\$24,495	\$148,766	\$6,700	\$155,466
	365	Piperia, Yadon's ( <i>Piperia yadonii</i> ) - Wherever found	E	\$34,500	\$109,105	\$143,605	\$0	\$143,605
	1258	Pitaya, Davis' green ( <i>Echinocereus viridiflorus</i> var. <i>davisii</i> ) - Wherever found	E	\$500	\$0	\$500	\$0	\$500
	1123	Pitcher-plant, Alabama canebrake ( <i>Sarracenia rubra</i> ssp. <i>alabamensis</i> ) - Wherever found	E	\$6,000	\$399	\$6,399	\$0	\$6,399
	1077	Pitcher-plant, green ( <i>Sarracenia oreophila</i> ) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	730	Pitcher-plant, mountain sweet ( <i>Sarracenia rubra</i> ssp. <i>jonesii</i> ) - Wherever found	E	\$20,000	\$2,867	\$22,867	\$3,200	\$26,067
	652	Plum, scrub ( <i>Prunus geniculata</i> ) - Wherever found	E	\$8,000	\$21,700	\$29,700	\$6,000	\$35,700
	807	Po'e ( <i>Portulaca sclerocarpa</i> ) - Wherever found	E	\$7,105	\$12,500	\$19,605	\$0	\$19,605
	284	Pogonia, small whorled ( <i>Isotria medeoloides</i> ) -	T	\$187,009	\$68,798	\$255,807	\$6,681	\$262,488
	609	Polygala, Lewton's ( <i>Polygala lewtonii</i> ) - Wherever found	E	\$2,000	\$26,426	\$28,426	\$14,200	\$42,626
	1126	Polygala, tiny ( <i>Polygala smallii</i> ) - Wherever found	E	\$6,000	\$0	\$6,000	\$300	\$6,300
	509	Polygonum, Scotts Valley ( <i>Polygonum hickmanii</i> ) - Wherever found	E	\$67,300	\$0	\$67,300	\$0	\$67,300

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	114	Pondberry ( <i>Lindera melissifolia</i> ) - Wherever found	E	\$29,500	\$1,165,048	\$1,194,548	\$12,400	\$1,206,948
	1265	Pondweed, Little Aguja (=Creek) ( <i>Potamogeton clystocarpus</i> ) - Wherever found	E	\$250	\$0	\$250	\$0	\$250
	333	Popcornflower, rough ( <i>Plagiobothrys hirtus</i> ) - Wherever found	E	\$171,500	\$200	\$171,700	\$0	\$171,700
	1033	Popolo ( <i>Cyanea solanacea</i> ) - Wherever found	E	\$9,145	\$0	\$9,145	\$0	\$9,145
	497	Popolo ( <i>Solanum nelsonii</i> ) - Wherever found	E	\$66,338	\$5,000	\$71,338	\$0	\$71,338
	515	Popolo ku mai ( <i>Solanum incompletum</i> ) - Wherever found	E	\$52,865	\$12,000	\$64,865	\$0	\$64,865
	1241	Poppy-mallow, Texas ( <i>Callirhoe scabriuscula</i> ) - Wherever found	E	\$1,000	\$420	\$1,420	\$0	\$1,420
	754	Poppy, Sacramento prickly ( <i>Argemone pleiacantha</i> ssp. <i>pinnatisecta</i> ) - Wherever found	E	\$5,000	\$18,550	\$23,550	\$0	\$23,550
	823	Potato-bean, Prices ( <i>Apios priceana</i> ) -	T	\$14,000	\$4,058	\$18,058	\$500	\$18,558
	603	Potentilla, Hickman's ( <i>Potentilla hickmanii</i> ) - Wherever found	E	\$6,800	\$36,600	\$43,400	\$0	\$43,400
	1132	Prairie-clover, Florida ( <i>Dalea carthagenensis floridana</i> ) - Wherever found	E	\$0	\$1,100	\$1,100	\$5,000	\$6,100
	819	Prairie-clover, leafy ( <i>Dalea foliosa</i> ) - Wherever found	E	\$17,000	\$1,626	\$18,626	\$0	\$18,626
	1249	Prickly-apple, aboriginal ( <i>Harrisia</i> (=Cereus) <i>aboriginum</i> (=gracilis)) - Wherever found	E	\$300	\$0	\$300	\$800	\$1,100
	1028	Prickly-apple, fragrant ( <i>Cereus eriophorus</i> var. <i>fragrans</i> ) - Wherever found	E	\$900	\$2,000	\$2,900	\$6,500	\$9,400
	1135	Prickly-ash, St. Thomas ( <i>Zanthoxylum thomasianum</i> ) - Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	1232	Primrose, Maguire ( <i>Primula maguirei</i> ) - Wherever found	T	\$0	\$2,000	\$2,000	\$0	\$2,000
	975	Pua `ala ( <i>Brighamia rockii</i> ) - Wherever found	E	\$10,778	\$500	\$11,278	\$0	\$11,278
	982	Pussypaws, Mariposa ( <i>Calyptridium pulchellum</i> ) - Wherever found	T	\$9,400	\$1,500	\$10,900	\$0	\$10,900

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1053	Pu`uka`a (Cyperus trachysanthos) - Wherever found	E	\$6,684	\$1,761	\$8,445	\$0	\$8,445
	1180	Ragwort, San Francisco Peaks (Packera franciscana) - Wherever found	T	\$1,700	\$2,400	\$4,100	\$0	\$4,100
	567	Rattleweed, hairy (Baptisia arachnifera) - Wherever found	E	\$11,000	\$0	\$11,000	\$40,000	\$51,000
	1186	Reedgrass, Hillebrand's (Calamagrostis hillebrandii) - Wherever found	E	\$3,744	\$0	\$3,744	\$0	\$3,744
	760	Reedgrass, Maui (Calamagrostis expansa) - Wherever found	E	\$11,208	\$12,000	\$23,208	\$0	\$23,208
	1250	Reed-mustard, Barneby (Schoenocrambe barnebyi) - Wherever found	E	\$0	\$1,050	\$1,050	\$0	\$1,050
	1194	Reed-mustard, clay (Schoenocrambe argillacea) - Wherever found	T	\$0	\$3,369	\$3,369	\$0	\$3,369
	1167	Reed-mustard, shrubby (Schoenocrambe suffrutescens) - Wherever found	E	\$0	\$4,712	\$4,712	\$0	\$4,712
	1026	Remya, Maui (Remya mauiensis) - Wherever found	E	\$9,473	\$0	\$9,473	\$0	\$9,473
	740	Rhododendron, Chapman (Rhododendron chapmanii) - Wherever found	E	\$2,500	\$22,000	\$24,500	\$450	\$24,950
	1270	Ridge-cress, Barneby (Lepidium barnebyanum) - Wherever found	E	\$0	\$100	\$100	\$0	\$100
	937	Rock-cress, Braun's (Arabis perstellata) - Wherever found	E	\$10,000	\$2,749	\$12,749	\$0	\$12,749
	1103	Rockcress, Georgia (Arabis georgiana) - Wherever found	T	\$0	\$1,826	\$1,826	\$5,000	\$6,826
	872	Rock-cress, Hoffmann's (Arabis hoffmannii) - Wherever found	E	\$2,700	\$13,025	\$15,725	\$0	\$15,725
	910	Rock-cress, McDonald's (Arabis macdonaldiana) - Wherever found	E	\$2,500	\$11,400	\$13,900	\$0	\$13,900
	355	Rockcress, Santa Cruz Island (Sibara filifolia) - Wherever found	E	\$3,800	\$149,163	\$152,963	\$0	\$152,963
	789	Rock cress, Shale barren (Arabis serotina) - Wherever found	E	\$14,680	\$240	\$14,920	\$5,560	\$20,480
	992	Rose-mallow, Neches River (Hibiscus dasycalyx) - Wherever found	T	\$3,500	\$7,000	\$10,500	\$0	\$10,500
	617	Rosemary, Apalachicola (Conradina glabra) - Wherever found	E	\$19,500	\$21,700	\$41,200	\$200	\$41,400



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	790	Rosemary, Cumberland ( <i>Conradina verticillata</i> ) - Wherever found	T	\$19,000	\$1,476	\$20,476	\$0	\$20,476
	917	Rosemary, Etonia ( <i>Conradina etonia</i> ) - Wherever found	E	\$4,500	\$125	\$4,625	\$9,000	\$13,625
	1092	Rosemary, short-leaved ( <i>Conradina brevifolia</i> ) - Wherever found	E	\$900	\$0	\$900	\$6,500	\$7,400
	473	Roseroot, Leedy's ( <i>Rhodiola integrifolia</i> ssp. <i>leedyi</i> ) - Wherever found	T	\$81,200	\$723	\$81,923	\$0	\$81,923
	945	Rush-pea, slender ( <i>Hoffmannseggia tenella</i> ) - Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1017	Rush-rose, island ( <i>Helianthemum greenei</i> ) - Wherever found	T	\$2,900	\$6,732	\$9,632	\$0	\$9,632
	801	Sandalwood, Lanai (= `iliah) ( <i>Santalum</i> <i>haleakalae</i> var. <i>lanaiense</i> ) - Wherever found	E	\$19,768	\$0	\$19,768	\$0	\$19,768
	709	Sandlace ( <i>Polygonella myriophylla</i> ) - Wherever found	E	\$2,000	\$21,700	\$23,700	\$4,500	\$28,200
	1244	Sandmat, pineland ( <i>Chamaesyce deltoidea</i> <i>pinetorum</i> ) - Wherever found	T	\$0	\$100	\$100	\$1,200	\$1,300
	1002	Sand-verbena, large-fruited ( <i>Abronia</i> <i>macrocarpa</i> ) - Wherever found	E	\$8,000	\$2,100	\$10,100	\$0	\$10,100
	1270	Sandwort, Bear Valley ( <i>Arenaria ursina</i> ) - Wherever found	T	\$0	\$100	\$100	\$0	\$100
	728	Sandwort, Cumberland ( <i>Arenaria</i> <i>cumberlandensis</i> ) - Wherever found	E	\$23,500	\$2,851	\$26,351	\$0	\$26,351
	821	Sandwort, Marsh ( <i>Arenaria paludicola</i> ) - Wherever found	E	\$17,300	\$1,300	\$18,600	\$0	\$18,600
	1147	Schiedea, Diamond Head ( <i>Schiedea adamantis</i> ) - Wherever found	E	\$5,670	\$0	\$5,670	\$0	\$5,670
	1171	Seablite, California ( <i>Suaeda californica</i> ) -	E	\$2,710	\$1,900	\$4,610	\$0	\$4,610
	198	Seagrass, Johnson's ( <i>Halophila johnsonii</i> ) - Wherever found	T	\$0	\$548,488	\$548,488	\$700	\$549,188
	1035	Sedge, golden ( <i>Carex lutea</i> ) - Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1134	Sedge, Navajo ( <i>Carex specuicola</i> ) - Wherever found	T	\$4,000	\$2,010	\$6,010	\$0	\$6,010
	1191	Silverbush, Blodgett's ( <i>Argythamnia blodgettii</i> ) - Wherever found	T	\$0	\$942	\$942	\$2,500	\$3,442

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	827	Silversword, Mauna Loa (=Ka'u) (Argyroxiphium kauense) - Wherever found	E	\$8,356	\$10,000	\$18,356	\$0	\$18,356
	1172	Skullcap, Florida (Scutellaria floridana) - Wherever found	T	\$3,500	\$0	\$3,500	\$1,000	\$4,500
	663	Skullcap, large-flowered (Scutellaria montana) - Wherever found	T	\$8,000	\$10,276	\$18,276	\$15,000	\$33,276
	559	Skyrocket, Pagosa (Ipomopsis polyantha) - Wherever found	E	\$16,700	\$25,000	\$41,700	\$11,465	\$53,165
	701	Snakeroot (Eryngium cuneifolium) - Wherever found	E	\$2,000	\$21,700	\$23,700	\$5,500	\$29,200
	916	Sneezeweed, Virginia (Helenium virginicum) - Wherever found	T	\$12,200	\$240	\$12,440	\$1,200	\$13,640
	1253	Snowbells, Texas (Styrax texanus) - Wherever found	E	\$1,000	\$0	\$1,000	\$0	\$1,000
	804	Spineflower, Ben Lomond (Chorizanthe pungens var. hartwegiana) - Wherever found	E	\$19,600	\$100	\$19,700	\$0	\$19,700
	950	Spineflower, Howell's (Chorizanthe howellii) - Wherever found	E	\$12,000	\$200	\$12,200	\$0	\$12,200
	453	Spineflower, Monterey (Chorizanthe pungens var. pungens) - Wherever found	T	\$81,900	\$7,244	\$89,144	\$0	\$89,144
	316	Spineflower, Orcutt's (Chorizanthe orcuttiana) - Wherever found	E	\$65,300	\$139,184	\$204,484	\$0	\$204,484
	1140	Spineflower, Robust (Chorizanthe robusta var. robusta) - Wherever found	E	\$4,100	\$1,800	\$5,900	\$0	\$5,900
	797	Spineflower, Scotts Valley (Chorizanthe robusta var. hartwegii) - Wherever found	E	\$20,200	\$0	\$20,200	\$0	\$20,200
	1235	Spineflower, slender-horned (Dodecahema leptoceras) - Wherever found	E	\$1,800	\$0	\$1,800	\$0	\$1,800
	1214	Spineflower, Sonoma (Chorizanthe valida) - Wherever found	E	\$1,700	\$1,000	\$2,700	\$0	\$2,700
	551	Spiraea, Virginia (Spiraea virginiana) - Wherever found	T	\$39,775	\$5,294	\$45,069	\$10,839	\$55,908
	1146	Spurge, deltoid (Chamaesyce deltoidea ssp. deltoidea) - Wherever found	E	\$4,500	\$0	\$4,500	\$1,200	\$5,700
	1071	Spurge, Garber's (Chamaesyce garberi) - Wherever found	T	\$6,000	\$942	\$6,942	\$1,200	\$8,142

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	977	Spurge, Hoover's ( <i>Chamaesyce hooveri</i> ) - Wherever found	T	\$7,700	\$3,400	\$11,100	\$0	\$11,100
	839	Spurge, telephus ( <i>Euphorbia telephioides</i> ) - Wherever found	T	\$11,000	\$0	\$11,000	\$6,500	\$17,500
	1230	Spurge, wedge ( <i>Chamaesyce deltoidea</i> <i>serpyllum</i> ) - Wherever found	E	\$900	\$0	\$900	\$1,200	\$2,100
	1104	Stickseed, showy ( <i>Hackelia venusta</i> ) - Wherever found	E	\$810	\$6,000	\$6,810	\$0	\$6,810
	1267	Stonecrop, Lake County ( <i>Parvisedum</i> <i>leiocarpum</i> ) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	277	Sumac, Michaux's ( <i>Rhus michauxii</i> ) - Wherever found	E	\$60,500	\$212,415	\$272,915	\$5,000	\$277,915
	1124	Sunburst, Hartweg's golden ( <i>Pseudobahia</i> <i>bahiifolia</i> ) - Wherever found	E	\$0	\$6,364	\$6,364	\$0	\$6,364
	809	Sunburst, San Joaquin adobe ( <i>Pseudobahia</i> <i>peirsonii</i> ) - Wherever found	T	\$13,500	\$5,916	\$19,416	\$0	\$19,416
	696	Sunflower, Pecos (=puzzle, =paradox) ( <i>Helianthus paradoxus</i> ) - Wherever found	T	\$6,000	\$23,650	\$29,650	\$0	\$29,650
	1028	Sunflower, San Mateo woolly ( <i>Eriophyllum</i> <i>latilobum</i> ) - Wherever found	E	\$9,400	\$0	\$9,400	\$0	\$9,400
	458	Sunflower, Schweinitz's ( <i>Helianthus</i> <i>schweinitzii</i> ) - Wherever found	E	\$41,500	\$38,825	\$80,325	\$6,800	\$87,125
	831	Sunflower, whorled ( <i>Helianthus verticillatus</i> ) -	E	\$13,000	\$0	\$13,000	\$5,000	\$18,000
	874	Sunray, Ash Meadows ( <i>Enceliopsis nudicaulis</i> var. <i>corrugata</i> ) - Wherever found	T	\$15,600	\$0	\$15,600	\$0	\$15,600
	662	Sunshine, Sonoma ( <i>Blennosperma bakeri</i> ) - Wherever found	E	\$11,200	\$22,200	\$33,400	\$0	\$33,400
	1270	Taraxacum, California ( <i>Taraxacum</i> <i>californicum</i> ) - Wherever found	E	\$0	\$100	\$100	\$0	\$100
	1203	Tarplant, Gaviota ( <i>Deinandra increscens</i> ssp. <i>villosa</i> ) - Wherever found	E	\$3,000	\$200	\$3,200	\$0	\$3,200
	505	Tarplant, Otay ( <i>Deinandra</i> (=Hemizonia) <i>conjugens</i> ) - Wherever found	T	\$10,500	\$58,200	\$68,700	\$0	\$68,700
	698	Tarplant, Santa Cruz ( <i>Holocarpha macradenia</i> ) - Wherever found	T	\$23,900	\$5,400	\$29,300	\$0	\$29,300

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	670	Thelypody, Howell's spectacular ( <i>Thelypodium howellii</i> ssp. <i>spectabilis</i> ) - Wherever found	T	\$32,310	\$0	\$32,310	\$0	\$32,310
	1189	Thistle, Chorro Creek bog ( <i>Cirsium fontinale</i> var. <i>obispoense</i> ) - Wherever found	E	\$3,300	\$300	\$3,600	\$0	\$3,600
	1019	Thistle, fountain ( <i>Cirsium fontinale</i> var. <i>fontinale</i> ) - Wherever found	E	\$9,400	\$200	\$9,600	\$0	\$9,600
	450	Thistle, La Graciosa ( <i>Cirsium loncholepis</i> ) - Wherever found	E	\$86,100	\$5,568	\$91,668	\$0	\$91,668
	1267	Thistle, Loch Lomond coyote ( <i>Eryngium constancei</i> ) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	455	Thistle, Pitcher's ( <i>Cirsium pitcheri</i> ) - Wherever found	T	\$41,000	\$42,126	\$83,126	\$5,600	\$88,726
	1163	Thistle, Sacramento Mountains ( <i>Cirsium vinaceum</i> ) - Wherever found	T	\$5,000	\$0	\$5,000	\$0	\$5,000
	642	Thistle, Suisun ( <i>Cirsium hydrophilum</i> var. <i>hydrophilum</i> ) - Wherever found	E	\$29,100	\$8,200	\$37,300	\$0	\$37,300
	610	Thornmint, San Diego ( <i>Acanthomintha ilicifolia</i> ) - Wherever found	T	\$13,500	\$29,000	\$42,500	\$0	\$42,500
	1270	Thornmint, San Mateo ( <i>Acanthomintha obovata</i> ssp. <i>duttonii</i> ) - Wherever found	E	\$0	\$100	\$100	\$0	\$100
	761	Thoroughwort, Cape Sable ( <i>Chromolaena frustrata</i> ) - Wherever found	E	\$21,500	\$0	\$21,500	\$1,700	\$23,200
	1173	Townsendia, Last Chance ( <i>Townsendia aprica</i> ) - Wherever found	T	\$0	\$4,450	\$4,450	\$0	\$4,450
	1077	Trillium, persistent ( <i>Trillium persistens</i> ) - Wherever found	E	\$0	\$0	\$0	\$8,000	\$8,000
	1020	Trillium, relict ( <i>Trillium reliquum</i> ) - Wherever found	E	\$0	\$5,538	\$5,538	\$4,000	\$9,538
	857	Tuctoria, Greene's ( <i>Tuctoria greenei</i> ) - Wherever found	E	\$12,400	\$4,024	\$16,424	\$0	\$16,424
	873	Twinpod, Dudley Bluffs ( <i>Physaria obcordata</i> ) - Wherever found	T	\$5,200	\$10,500	\$15,700	\$0	\$15,700
	504	Ufa-halomtano ( <i>Heritiera longipetiolata</i> ) - Wherever found	E	\$47,170	\$22,314	\$69,484	\$0	\$69,484
	635	Uhi uhi ( <i>Mezoneuron kawaiense</i> ) - Wherever found	E	\$28,826	\$10,000	\$38,826	\$0	\$38,826

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1062	Uvillo ( <i>Eugenia haematocarpa</i> ) - Wherever found	E	\$8,000	\$300	\$8,300	\$0	\$8,300
	833	Vervain, Red Hills ( <i>Verbena californica</i> ) - Wherever found	T	\$1,200	\$16,700	\$17,900	\$0	\$17,900
	946	Vetch, Hawaiian ( <i>Vicia menziesii</i> ) - Wherever found	E	\$12,496	\$0	\$12,496	\$0	\$12,496
	273	Wallflower, Ben Lomond ( <i>Erysimum teretifolium</i> ) - Wherever found	E	\$283,900	\$2,500	\$286,400	\$0	\$286,400
	369	Wallflower, Contra Costa ( <i>Erysimum capitatum</i> var. <i>angustatum</i> ) - Wherever found	E	\$139,000	\$700	\$139,700	\$0	\$139,700
	372	Wallflower, Menzies' ( <i>Erysimum menziesii</i> ) - Wherever found	E	\$6,600	\$131,200	\$137,800	\$0	\$137,800
	1172	Walnut (=Nogal), West Indian ( <i>Juglans jamaicensis</i> ) - Wherever found	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	590	Warea, wide-leaf ( <i>Warea amplexifolia</i> ) - Wherever found	E	\$0	\$0	\$0	\$46,500	\$46,500
	759	Watercress, Gambel's ( <i>Rorippa gambellii</i> ) - Wherever found	E	\$23,100	\$200	\$23,300	\$0	\$23,300
	1245	Water-plantain, Kral's ( <i>Sagittaria secundifolia</i> ) - Wherever found	T	\$0	\$299	\$299	\$1,000	\$1,299
	694	Water-umbel, Huachuca ( <i>Lilaeopsis schaffneriana</i> var. <i>recurva</i> ) - Wherever found	E	\$13,500	\$16,300	\$29,800	\$0	\$29,800
	1267	Water-willow, Cooley's ( <i>Justicia cooleyi</i> ) - Wherever found	E	\$0	\$0	\$0	\$200	\$200
	1077	Whitlow-wort, papery ( <i>Paronychia chartacea</i> ) - Wherever found	T	\$2,000	\$0	\$2,000	\$6,000	\$8,000
	805	Wild buckwheat, clay-loving ( <i>Eriogonum pelinophilum</i> ) - Wherever found	E	\$500	\$19,150	\$19,650	\$0	\$19,650
	564	Wild-buckwheat, gypsum ( <i>Eriogonum gypsophilum</i> ) - Wherever found	T	\$0	\$52,000	\$52,000	\$0	\$52,000
	1236	Wild-buckwheat, southern mountain ( <i>Eriogonum kennedyi</i> var. <i>austromontanum</i> ) - Wherever found	T	\$1,700	\$0	\$1,700	\$0	\$1,700
	983	Wild-rice, Texas ( <i>Zizania texana</i> ) - Wherever found	E	\$10,000	\$840	\$10,840	\$0	\$10,840
	894	Wire-lettuce, Malheur ( <i>Stephanomeria malheurensis</i> ) - Wherever found	E	\$12,620	\$2,000	\$14,620	\$0	\$14,620

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	727	Wireweed ( <i>Polygonella basiramia</i> ) - Wherever found	E	\$2,000	\$21,700	\$23,700	\$2,700	\$26,400
	357	Woodland-star, San Clemente Island ( <i>Lithophragma maximum</i> ) - Wherever found	E	\$3,500	\$145,951	\$149,451	\$0	\$149,451
	784	Woolly-star, Santa Ana River ( <i>Eriastrum densifolium</i> ssp. <i>sanctorum</i> ) - Wherever found	E	\$12,500	\$8,318	\$20,818	\$0	\$20,818
	373	Woolly-threads, San Joaquin ( <i>Monolopia</i> (=Lembertia) <i>congdonii</i> ) - Wherever found	E	\$58,600	\$78,904	\$137,504	\$0	\$137,504
	407	Yellowhead, desert ( <i>Yermo xanthocephalus</i> ) - Wherever found	T	\$61,500	\$50,000	\$111,500	\$0	\$111,500
	928	Yerba santa, Lompoc ( <i>Eriodictyon capitatum</i> ) - Wherever found	E	\$12,900	\$200	\$13,100	\$0	\$13,100
	856	Ziziphus, Florida ( <i>Ziziphus celata</i> ) - Wherever found	E	\$3,000	\$0	\$3,000	\$13,500	\$16,500
<b>Flowering Plants subtotal</b>				<b>\$15,725,173</b>	<b>\$14,779,388</b>	<b>\$30,504,561</b>	<b>\$807,367</b>	<b>\$31,311,928</b>
Conifers and Cycads	1102	Cypress, Gowen ( <i>Cupressus goveniana</i> ssp. <i>goveniana</i> ) - Wherever found	T	\$6,832	\$0	\$6,832	\$0	\$6,832
	981	Cypress, Santa Cruz ( <i>Cupressus abramsiana</i> ) - Wherever found	T	\$10,931	\$0	\$10,931	\$0	\$10,931
	376	Fadang ( <i>Cycas micronesica</i> ) - Wherever found	T	\$65,865	\$70,664	\$136,529	\$0	\$136,529
	428	Torreya, Florida ( <i>Torreya taxifolia</i> ) - Wherever found	E	\$23,500	\$73,932	\$97,432	\$1,500	\$98,932
<b>Conifers and Cycads subtotal</b>				<b>\$107,128</b>	<b>\$144,596</b>	<b>\$251,724</b>	<b>\$1,500</b>	<b>\$253,224</b>
Ferns and Allies	1155	Aumakua, Palapalai ( <i>Dryopteris crinalis</i> var. <i>podosorus</i> ) - Wherever found	E	\$5,265	\$0	\$5,265	\$0	\$5,265
	993	Diellia, Asplenium-leaved ( <i>Asplenium dielirectum</i> ) - Wherever found	E	\$10,449	\$0	\$10,449	\$0	\$10,449
	1263	Fern, Alabama streak-sorus ( <i>Thelypteris pilosa</i> var. <i>alabamensis</i> ) - Wherever found	T	\$0	\$299	\$299	\$0	\$299
	711	Fern, Aleutian shield ( <i>Polystichum aleuticum</i> ) - Wherever found	E	\$18,000	\$10,000	\$28,000	\$0	\$28,000
	605	Fern, American hart's-tongue ( <i>Asplenium scolopendrium</i> var. <i>americanum</i> ) - Wherever found	T	\$43,100	\$125	\$43,225	\$0	\$43,225
	1115	Fern, Elfin tree ( <i>Cyathea dryopteroides</i> ) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1203	Fern, Florida bristle ( <i>Trichomanes punctatum</i> ssp. <i>floridanum</i> ) - Wherever found	E	\$2,000	\$0	\$2,000	\$1,200	\$3,200
	1157	Fern, pendant kihi ( <i>Adenophorus periens</i> ) - Wherever found	E	\$5,244	\$0	\$5,244	\$0	\$5,244
	1083	Hohiu ( <i>Dryopteris glabra</i> var. <i>pusilla</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	640	Ihi`ihi ( <i>Marsilea villosa</i> ) - Wherever found	E	\$36,220	\$1,792	\$38,012	\$0	\$38,012
	825	Kupukupu makalii ( <i>Cyclosorus boydiae</i> ) - Wherever found	E	\$6,375	\$12,000	\$18,375	\$0	\$18,375
	1151	No Common Name ( <i>Adiantum vivesii</i> ) - Wherever found	E	\$5,500	\$0	\$5,500	\$0	\$5,500
	1074	No Common Name ( <i>Asplenium dielfalcatum</i> ) - Wherever found	E	\$8,073	\$0	\$8,073	\$0	\$8,073
	1141	No Common Name ( <i>Asplenium diellaciniatum</i> ) - Wherever found	E	\$5,853	\$0	\$5,853	\$0	\$5,853
	985	No Common Name ( <i>Asplenium dielmannii</i> ) - Wherever found	E	\$10,778	\$0	\$10,778	\$0	\$10,778
	1083	No Common Name ( <i>Asplenium dielpallidum</i> ) - Wherever found	E	\$7,736	\$0	\$7,736	\$0	\$7,736
	745	No Common Name ( <i>Asplenium peruvianum</i> var. <i>insulare</i> ) - Wherever found	E	\$10,574	\$13,973	\$24,547	\$0	\$24,547
	1027	No Common Name ( <i>Asplenium unisorum</i> ) - Wherever found	E	\$7,736	\$1,734	\$9,470	\$0	\$9,470
	1145	No Common Name ( <i>Deparia kaalaana</i> ) - Wherever found	E	\$5,701	\$0	\$5,701	\$0	\$5,701
	1075	No Common Name ( <i>Diplazium molokaiense</i> ) - Wherever found	E	\$8,024	\$0	\$8,024	\$0	\$8,024
	961	No Common Name ( <i>Doryopteris angelica</i> ) - Wherever found	E	\$11,792	\$0	\$11,792	\$0	\$11,792
	1152	No Common Name ( <i>Doryopteris takeuchii</i> ) - Wherever found	E	\$5,417	\$0	\$5,417	\$0	\$5,417
	1135	No Common Name ( <i>Elaphoglossum serpens</i> ) - Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	1034	No Common Name ( <i>Huperzia stemmermanniae</i> ) - Wherever found	E	\$9,099	\$0	\$9,099	\$0	\$9,099
	752	No Common Name ( <i>Microlepia strigosa</i> var. <i>mauiensis</i> ) - Wherever found	E	\$11,888	\$12,000	\$23,888	\$0	\$23,888

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1163	No Common Name (Polystichum calderonense) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1144	No Common Name (Pteris lidgatei) - Wherever found	E	\$5,754	\$0	\$5,754	\$0	\$5,754
	1135	No Common Name (Tectaria estremerana) - Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	1088	No Common Name (Thelypteris inabonensis) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1135	No Common Name (Thelypteris verecunda) - Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	1163	No Common Name (Thelypteris yaucoensis) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1137	Olua (Hypolepis hawaiiensis var. mauiensis) - Wherever found	E	\$5,954	\$0	\$5,954	\$0	\$5,954
	777	Pauoa (Ctenitis squamigera) - Wherever found	E	\$9,276	\$12,000	\$21,276	\$0	\$21,276
	1232	Quillwort, black spored (Isoetes melanospora) - Wherever found	E	\$0	\$0	\$0	\$2,000	\$2,000
	344	Quillwort, Louisiana (Isoetes louisianensis) - Wherever found	E	\$28,000	\$133,116	\$161,116	\$0	\$161,116
	1210	Quillwort, mat-forming (Isoetes tegetiformans) - Wherever found	E	\$0	\$0	\$0	\$3,000	\$3,000
	802	Wawae`iole (Huperzia mannii) - Wherever found	E	\$7,765	\$12,000	\$19,765	\$0	\$19,765
	1105	Wawae`iole (Huperzia nutans) - Wherever found	E	\$6,764	\$0	\$6,764	\$0	\$6,764
<b>Ferns and Allies subtotal</b>				<b>\$348,073</b>	<b>\$209,039</b>	<b>\$557,112</b>	<b>\$6,200</b>	<b>\$563,312</b>
Lichens	637	Cladonia, Florida perforate (Cladonia perforata) - Wherever found	E	\$3,200	\$28,200	\$31,400	\$7,000	\$38,400
	1210	Lichen, rock gnome (Gymnoderma lineare) - Wherever found	E	\$2,400	\$0	\$2,400	\$600	\$3,000
<b>Lichens subtotal</b>				<b>\$5,600</b>	<b>\$28,200</b>	<b>\$33,800</b>	<b>\$7,600</b>	<b>\$41,400</b>
Multi-species								
<b>Multi-species subtotal</b>				<b>\$2,682,138</b>	<b>\$133,994,427</b>	<b>\$136,676,565</b>	<b>\$1,559,778</b>	<b>\$138,223,343</b>
<b>Total Species Expenditures</b>				<b>\$154,791,012</b>	<b>\$1,127,318,971</b>	<b>\$1,282,109,983</b>	<b>\$125,747,415</b>	<b>\$1,407,857,398</b>



**TABLE 2. SPECIES RANKED IN DESCENDING ORDER OF TOTAL FY 2018  
REPORTED EXPENDITURES**

Species are arranged in descending rank order of total identifiable expenditures (Table 1). Cumulative totals of the expenditures totals are provided. The cumulative total column is the sum of all higher ranked species totals (from Table 1) and the selected species entry. This table does not include costs for land acquisition or multi-species conservation activities.

The species list in the report follows the same breakdown as in 50 CFR 17.11 and 17.12. All populations are itemized separately, not just Evolutionarily Significant Units (ESUs). Thus, it appears as if there are more listed species. Separate populations of the same species may be ranked differently. The STATUS column indicates whether the species is endangered (E), threatened (T), experimental (EXPN), or threatened by similarity of appearance (SAT).

**TABLE 2. SPECIES RANKED IN DESCENDING ORDER OF TOTAL FY 2018 REPORTED EXPENDITURES, NOT INCLUDING LAND ACQUISITION COSTS**

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) tshawytscha) - Puget Sound ESU	T	\$73,825,298	\$73,825,298
2	Steelhead ( <i>Oncorhynchus</i> (=Salmo) mykiss) - Middle Columbia River DPS	T	\$61,612,669	\$135,437,967
3	Steelhead ( <i>Oncorhynchus</i> (=Salmo) mykiss) - Snake River Basin DPS	T	\$58,225,309	\$193,663,276
4	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) tshawytscha) - Snake River spring/summer-run ESU	T	\$56,971,122	\$250,634,398
5	Owl, northern spotted ( <i>Strix occidentalis caurina</i> ) - Wherever found	T	\$45,432,999	\$296,067,397
6	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) tshawytscha) - Upper Columbia spring-run ESU	E	\$42,007,681	\$338,075,078
7	Trout, bull ( <i>Salvelinus confluentus</i> ) - U.S.A., conterminous, lower 48 states	T	\$40,264,658	\$378,339,736
8	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) tshawytscha) - Snake River fall-run ESU	T	\$36,268,980	\$414,608,716
9	Steelhead ( <i>Oncorhynchus</i> (=Salmo) mykiss) - Upper Columbia River DPS	T	\$32,482,180	\$447,090,896
10	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) tshawytscha) - Lower Columbia River ESU	T	\$30,062,234	\$477,153,130
11	Salmon, sockeye ( <i>Oncorhynchus</i> (=Salmo) nerka) - Snake River ESU	E	\$24,854,826	\$502,007,956
12	Sturgeon, pallid ( <i>Scaphirhynchus albus</i> ) - Wherever found	E	\$22,165,632	\$524,173,588
13	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) tshawytscha) - Sacramento River winter-run ESU	E	\$21,981,159	\$546,154,747
14	Smelt, delta ( <i>Hypomesus transpacificus</i> ) - Wherever found	T	\$20,805,869	\$566,960,616
15	Steelhead ( <i>Oncorhynchus</i> (=Salmo) mykiss) - Lower Columbia River DPS	T	\$19,041,522	\$586,002,138
16	Steelhead ( <i>Oncorhynchus</i> (=Salmo) mykiss) - California Central Valley DPS	T	\$18,964,487	\$604,966,625
17	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) tshawytscha) - Central Valley spring-run ESU	T	\$18,436,787	\$623,403,412
18	Salmon, coho ( <i>Oncorhynchus</i> (=Salmo) kisutch) - Lower Columbia River ESU	T	\$17,904,930	\$641,308,342
19	Woodpecker, red-cockaded ( <i>Picoides borealis</i> ) - Wherever found	E	\$16,676,088	\$657,984,430
20	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) tshawytscha) - Upper Willamette River ESU	T	\$15,190,580	\$673,175,010
21	Salmon, coho ( <i>Oncorhynchus</i> (=Salmo) kisutch) - Southern Oregon - Northern California Coast ESU	T	\$13,789,135	\$686,964,145
22	Steelhead ( <i>Oncorhynchus</i> (=Salmo) mykiss) - Puget Sound DPS	T	\$13,294,818	\$700,258,963
23	Sucker, razorback ( <i>Xyrauchen texanus</i> ) - Wherever found	E	\$13,009,906	\$713,268,869
24	Sturgeon, white ( <i>Acipenser transmontanus</i> ) - U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	E	\$12,989,493	\$726,258,362
25	Salmon, chum ( <i>Oncorhynchus keta</i> ) - Columbia River ESU	T	\$12,722,163	\$738,980,525
26	Tortoise, desert ( <i>Gopherus agassizii</i> ) - Wherever found, except AZ south and east of Colorado R., and Mexico	T	\$12,660,590	\$751,641,115
27	Steelhead ( <i>Oncorhynchus</i> (=Salmo) mykiss) - Upper Willamette River DPS	T	\$12,484,100	\$764,125,215

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
28	Sturgeon, green ( <i>Acipenser medirostris</i> ) - Southern DPS	T	\$10,862,922	\$774,988,137
29	Salmon, coho ( <i>Oncorhynchus</i> (=Salmo) <i>kisutch</i> ) - Central California Coast ESU	E	\$10,755,338	\$785,743,475
30	Whale, North Atlantic Right ( <i>Eubalaena glacialis</i> ) - Wherever found	E	\$9,942,274	\$795,685,749
31	Caracara, Audubon's crested ( <i>Polyborus plancus audubonii</i> ) - FL pop.	T	\$9,655,413	\$805,341,162
32	Bonytail ( <i>Gila elegans</i> ) - Wherever found	E	\$9,502,759	\$814,843,921
33	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - U.S.A., conterminous (lower 48) States, except where listed as an experimental population	T	\$9,253,873	\$824,097,794
34	Salmon, coho ( <i>Oncorhynchus</i> (=Salmo) <i>kisutch</i> ) - Oregon Coast ESU	T	\$9,148,187	\$833,245,981
35	Trout, Lahontan cutthroat ( <i>Oncorhynchus clarkii henshawi</i> ) - Wherever found	T	\$8,802,507	\$842,048,488
36	Manatee, West Indian ( <i>Trichechus manatus</i> ) - Wherever found	T	\$8,681,080	\$850,729,568
37	Salmon, Atlantic ( <i>Salmo salar</i> ) - Gulf of Maine DPS	E	\$8,644,171	\$859,373,739
38	Plover, piping ( <i>Charadrius melodus</i> ) - [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	T	\$8,131,333	\$867,505,072
39	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Northern California DPS	T	\$7,689,200	\$875,194,272
40	Salmon, chum ( <i>Oncorhynchus keta</i> ) - Hood Canal summer-run ESU	T	\$7,617,471	\$882,811,743
41	Chub, humpback ( <i>Gila cypha</i> ) - Wherever found	E	\$7,303,351	\$890,115,094
42	Flycatcher, southwestern willow ( <i>Empidonax traillii extimus</i> ) - Wherever found	E	\$7,290,547	\$897,405,641
43	Bat, Northern long-eared ( <i>Myotis septentrionalis</i> ) - Wherever found	T	\$6,834,744	\$904,240,385
44	Minnnow, Rio Grande Silvery ( <i>Hybognathus amarus</i> ) - Wherever found, except where listed as an experimental population	E	\$6,829,636	\$911,070,021
45	Tern, least ( <i>Sterna antillarum</i> ) - interior pop.	E	\$6,010,775	\$917,080,796
46	Murrelet, marbled ( <i>Brachyramphus marmoratus</i> ) - U.S.A. (CA, OR, WA)	T	\$5,892,481	\$922,973,277
47	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Central California Coast DPS	T	\$5,835,104	\$928,808,381
48	Cuckoo, yellow-billed ( <i>Coccyzus americanus</i> ) - Western U.S. DPS	T	\$5,790,505	\$934,598,886
49	Sea lion, Steller ( <i>Eumetopias jubatus</i> ) - Western DPS	E	\$5,690,154	\$940,289,040
50	Bat, Indiana ( <i>Myotis sodalis</i> ) - Wherever found	E	\$5,483,252	\$945,772,292
51	Sea turtle, Kemp's ridley ( <i>Lepidochelys kempii</i> ) - Wherever found	E	\$5,448,732	\$951,221,024
52	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Southern California DPS	E	\$5,396,190	\$956,617,214
53	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - Northwest Atlantic Ocean DPS	T	\$4,946,489	\$961,563,703
54	Salmon, sockeye ( <i>Oncorhynchus</i> (=Salmo) <i>nerka</i> ) - Ozette Lake ESU	T	\$4,899,528	\$966,463,231
55	Sturgeon, Gulf ( <i>Acipenser oxyrinchus</i> (=oxyrinchus) <i>desotoi</i> ) - Wherever found	T	\$4,827,386	\$971,290,617
56	Sea turtle, leatherback ( <i>Dermochelys coriacea</i> ) - Wherever found	E	\$4,715,335	\$976,005,952

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
57	Wolf, gray ( <i>Canis lupus</i> ) - U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	E	\$4,614,123	\$980,620,075
58	Seal, Hawaiian monk ( <i>Monachus schauinslandi</i> ) - Wherever found	E	\$4,273,134	\$984,893,209
59	Pikeminnow (=squawfish), Colorado ( <i>Ptychocheilus lucius</i> ) - Wherever found, except where listed as an experimental population	E	\$4,117,142	\$989,010,351
60	Plover, western snowy ( <i>Charadrius nivosus nivosus</i> ) - Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	T	\$4,088,370	\$993,098,721
61	Condor, California ( <i>Gymnogyps californianus</i> ) - U.S.A. only, except where listed as an experimental population	E	\$4,088,113	\$997,186,834
62	Vireo, least Bell's ( <i>Vireo bellii pusillus</i> ) - Wherever found	E	\$4,025,002	\$1,001,211,836
63	Eider, spectacled ( <i>Somateria fischeri</i> ) - Wherever found	T	\$3,847,136	\$1,005,058,972
64	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i> ) - California Coastal ESU	T	\$3,760,184	\$1,008,819,156
65	Sea turtle, hawksbill ( <i>Eretmochelys imbricata</i> ) - Wherever found	E	\$3,747,833	\$1,012,566,989
66	Eulachon ( <i>Thaleichthys pacificus</i> ) - Southern DPS	T	\$3,718,161	\$1,016,285,150
67	Eider, Steller's ( <i>Polysticta stelleri</i> ) - AK breeding pop.	T	\$3,687,916	\$1,019,973,066
68	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - South-Central California Coast DPS	T	\$3,653,700	\$1,023,626,766
69	Whale, beluga ( <i>Delphinapterus leucas</i> ) - Cook Inlet DPS	E	\$3,530,280	\$1,027,157,046
70	Lynx, Canada ( <i>Lynx canadensis</i> ) - Wherever Found in Contiguous U.S.	T	\$3,238,157	\$1,030,395,203
71	Gnatcatcher, coastal California ( <i>Polioptila californica californica</i> ) - Wherever found	T	\$3,127,556	\$1,033,522,759
72	Elephant, African ( <i>Loxodonta africana</i> ) - Wherever found	T	\$3,083,083	\$1,036,605,842
73	Wolf, Mexican ( <i>Canis lupus baileyi</i> ) - U.S.A. (portions of AZ and NM)see 17.84(k)	EXPN	\$2,819,277	\$1,039,425,119
74	Abalone, Black ( <i>Haliotis cracherodii</i> ) - Wherever found	E	\$2,699,460	\$1,042,124,579
75	Bear, polar ( <i>Ursus maritimus</i> ) - Wherever found	T	\$2,699,190	\$1,044,823,769
76	Shrike, San Clemente loggerhead ( <i>Lanius ludovicianus mearnsi</i> ) - Wherever found	E	\$2,627,464	\$1,047,451,233
77	Rail, Yuma Ridgways ( <i>Rallus obsoletus</i> [=longirostris] <i>yumanensis</i> ) - Wherever found	E	\$2,579,143	\$1,050,030,376
78	Tortoise, gopher ( <i>Gopherus polyphemus</i> ) - West of Mobile and Tombigbee Rivers	T	\$2,553,779	\$1,052,584,155
79	Sucker, shortnose ( <i>Chasmistes brevirostris</i> ) - Wherever found	E	\$2,532,579	\$1,055,116,734
80	Sea turtle, green ( <i>Chelonia mydas</i> ) - North Atlantic DPS	T	\$2,471,502	\$1,057,588,236
81	Sucker, Lost River ( <i>Deltistes luxatus</i> ) - Wherever found	E	\$2,425,676	\$1,060,013,912
82	Frog, California red-legged ( <i>Rana draytonii</i> ) - Wherever found	T	\$2,401,648	\$1,062,415,560
83	Panther, Florida ( <i>Puma</i> (=Felis) <i>concolor coryi</i> ) - Wherever found	E	\$2,187,876	\$1,064,603,436

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
84	Whale, finback ( <i>Balaenoptera physalus</i> ) - Wherever found	E	\$2,167,630	\$1,066,771,066
85	Fly, Delhi Sands flower-loving ( <i>Rhaphiomidas terminatus abdominalis</i> ) - Wherever found	E	\$2,163,525	\$1,068,934,591
86	Tern, California least ( <i>Sterna antillarum browni</i> ) - Wherever found	E	\$2,136,751	\$1,071,071,342
87	Whale, killer ( <i>Orcinus orca</i> ) - Southern Resident DPS	E	\$2,077,836	\$1,073,149,178
88	Whale, sperm ( <i>Physeter catodon</i> (=macrocephalus)) - Wherever found	E	\$2,031,731	\$1,075,180,909
89	Shiner, Pecos bluntnose ( <i>Notropis simus pecosensis</i> ) - Wherever found	T	\$2,020,750	\$1,077,201,659
90	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - North Pacific Ocean DPS	E	\$1,905,046	\$1,079,106,705
91	Whale, blue ( <i>Balaenoptera musculus</i> ) - Wherever found	E	\$1,898,201	\$1,081,004,906
92	Alligator, American ( <i>Alligator mississippiensis</i> ) - Wherever found	SAT	\$1,762,225	\$1,082,767,131
93	Bat, Hawaiian hoary ( <i>Lasiurus cinereus semotus</i> ) - Wherever found	E	\$1,738,008	\$1,084,505,139
94	Seal, bearded ( <i>Erignathus barbatus nauticus</i> ) - Beringia DPS	T	\$1,705,000	\$1,086,210,139
95	Frog, Oregon spotted ( <i>Rana pretiosa</i> ) - Wherever found	T	\$1,654,495	\$1,087,864,634
96	Sea turtle, green ( <i>Chelonia mydas</i> ) - Central West Pacific DPS	E	\$1,646,720	\$1,089,511,354
97	Parrot, Puerto Rican ( <i>Amazona vittata</i> ) - Wherever found	E	\$1,642,500	\$1,091,153,854
98	Whale, Sei ( <i>Balaenoptera borealis</i> ) - Wherever found	E	\$1,613,657	\$1,092,767,511
99	Sea turtle, green ( <i>Chelonia mydas</i> ) - East Pacific DPS	T	\$1,541,090	\$1,094,308,601
100	Sturgeon, shortnose ( <i>Acipenser brevirostrum</i> ) - Wherever found	E	\$1,518,084	\$1,095,826,685
101	Snake, eastern indigo ( <i>Drymarchon corais couperi</i> ) - Wherever found	T	\$1,517,654	\$1,097,344,339
102	Goose, Hawaiian ( <i>Branta</i> (=Nesochen) <i>sandvicensis</i> ) - Wherever found	E	\$1,499,285	\$1,098,843,624
103	Sage-grouse, Gunnison ( <i>Centrocercus minimus</i> ) - Wherever found	T	\$1,494,796	\$1,100,338,420
104	Stork, wood ( <i>Mycteria americana</i> ) - AL, FL, GA, MS, NC, SC	T	\$1,451,657	\$1,101,790,077
105	Pinesnake, Louisiana ( <i>Pituophis ruthveni</i> ) - Wherever found	T	\$1,424,474	\$1,103,214,551
106	Tiger ( <i>Panthera tigris</i> ) - Wherever found	E	\$1,409,159	\$1,104,623,710
107	Wolf, gray ( <i>Canis lupus</i> ) - MN	T	\$1,394,192	\$1,106,017,902
108	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Mexico DPS	T	\$1,385,855	\$1,107,403,757
109	Sea turtle, green ( <i>Chelonia mydas</i> ) - Central North Pacific DPS	T	\$1,381,871	\$1,108,785,628
110	Rail, California clapper ( <i>Rallus longirostris obsoletus</i> ) - Wherever found	E	\$1,342,343	\$1,110,127,971
111	Kite, Everglade snail ( <i>Rostrhamus sociabilis plumbeus</i> ) - Wherever found	E	\$1,332,647	\$1,111,460,618
112	Elephant, Asian ( <i>Elephas maximus</i> ) - Wherever found	E	\$1,296,246	\$1,112,756,864
113	Sparrow, Cape Sable seaside ( <i>Ammodramus maritimus mirabilis</i> ) - Wherever found	E	\$1,248,425	\$1,114,005,289
114	Pondberry ( <i>Lindera melissifolia</i> ) - Wherever found	E	\$1,206,948	\$1,115,212,237
115	Trout, Greenback Cutthroat ( <i>Oncorhynchus clarkii stomias</i> ) - Wherever found	T	\$1,194,040	\$1,116,406,277

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
116	Ferret, black-footed ( <i>Mustela nigripes</i> ) - Wherever found, except where listed as an experimental population	E	\$1,183,429	\$1,117,589,706
117	Scrub-jay, Florida ( <i>Aphelocoma coerulescens</i> ) - Wherever found	T	\$1,165,757	\$1,118,755,463
118	Plover, piping ( <i>Charadrius melodus</i> ) - [Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	E	\$1,156,782	\$1,119,912,245
119	Fox, San Joaquin kit ( <i>Vulpes macrotis mutica</i> ) - wherever found	E	\$1,156,730	\$1,121,068,975
120	Ocelot ( <i>Leopardus</i> (=Felis) <i>pardalis</i> ) - wherever found	E	\$1,091,310	\$1,122,160,285
121	Horned lark, streaked ( <i>Eremophila alpestris strigata</i> ) - Wherever found	T	\$1,086,273	\$1,123,246,558
122	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - Gulf of Maine DPS	T	\$1,072,614	\$1,124,319,172
123	Beetle, American burying ( <i>Nicrophorus americanus</i> ) - Wherever found, except where listed as an experimental population	E	\$1,068,113	\$1,125,387,285
124	Fairy shrimp, vernal pool ( <i>Branchinecta lynchi</i> ) - Wherever found	T	\$1,062,901	\$1,126,450,186
125	Sea turtle, green ( <i>Chelonia mydas</i> ) - South Atlantic DPS	T	\$1,060,130	\$1,127,510,316
126	Trout, bull ( <i>Salvelinus confluentus</i> ) - Clackamas River subbasin experimental population	EXPN	\$1,054,100	\$1,128,564,416
127	Rockfish, yelloweye ( <i>Sebastes ruberrimus</i> ) - Puget Sound - Georgia Basin DPS	T	\$1,030,951	\$1,129,595,367
128	Salamander, California tiger ( <i>Ambystoma californiense</i> ) - U.S.A. (Central CA DPS)	T	\$1,030,472	\$1,130,625,839
129	Butterfly, Quino checkerspot ( <i>Euphydryas editha quino</i> (=E. e. <i>wrighti</i> )) - Wherever found	E	\$1,012,532	\$1,131,638,371
130	Bocaccio ( <i>Sebastes paucispinis</i> ) - Puget Sound - Georgia Basin DPS	E	\$1,012,499	\$1,132,650,870
131	Snake, giant garter ( <i>Thamnophis gigas</i> ) - Wherever found	T	\$1,011,818	\$1,133,662,688
132	Whale, bowhead ( <i>Balaena mysticetus</i> ) - Wherever found	E	\$1,004,871	\$1,134,667,559
133	Abalone, White ( <i>Haliotis sorenseni</i> ) - North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	E	\$1,003,000	\$1,135,670,559
134	Toad, arroyo (=arroyo southwestern) ( <i>Anaxyrus californicus</i> ) - Wherever found	E	\$994,950	\$1,136,665,509
135	Mouse, salt marsh harvest ( <i>Reithrodontomys raviventris</i> ) - wherever found	E	\$989,500	\$1,137,655,009
136	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - New York Bight DPS	E	\$989,133	\$1,138,644,142
137	Sparrow, San Clemente sage ( <i>Amphispiza belli clementeae</i> ) - Wherever found	T	\$988,943	\$1,139,633,085
138	Rhinoceros, black ( <i>Diceros bicornis</i> ) - Africa	E	\$969,295	\$1,140,602,380
139	Shiner, Topeka ( <i>Notropis topeka</i> (=tristis)) - Wherever found, except where listed as an experimental population	E	\$954,432	\$1,141,556,812
140	Higgins eye (pearlymussel) ( <i>Lampsilis higginsii</i> ) - Wherever found	E	\$951,500	\$1,142,508,312
141	Frog, Sierra Nevada Yellow-legged ( <i>Rana sierrae</i> ) - Wherever found	E	\$950,324	\$1,143,458,636
142	Ferret, black-footed ( <i>Mustela nigripes</i> ) - U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	EXPN	\$950,003	\$1,144,408,639

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
143	Bat, gray ( <i>Myotis grisescens</i> ) - Wherever found	E	\$945,843	\$1,145,354,482
144	Knot, red ( <i>Calidris canutus rufa</i> ) - Wherever found	T	\$939,185	\$1,146,293,667
145	Massasauga (=rattlesnake), eastern ( <i>Sistrurus catenatus</i> ) - Wherever found	T	\$930,322	\$1,147,223,989
146	Crow, Mariana (=aga) ( <i>Corvus kubaryi</i> ) - Wherever found	E	\$919,931	\$1,148,143,920
147	Stilt, Hawaiian ( <i>Himantopus mexicanus knudseni</i> ) - Wherever found	E	\$911,977	\$1,149,055,897
148	Butterfly, Fender's blue ( <i>Icaricia icarioides fenderi</i> ) - Wherever found	E	\$907,729	\$1,149,963,626
149	Pupfish, Devils Hole ( <i>Cyprinodon diabolis</i> ) - Wherever found	E	\$906,463	\$1,150,870,089
150	Markhor, chiltan (=wild goat) ( <i>Capra falconeri</i> (=aegrugus) chiltanensis) - Wherever found	E	\$889,996	\$1,151,760,085
151	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Western North Pacific	E	\$874,481	\$1,152,634,566
152	Duck, Hawaiian (=koloa) ( <i>Anas wyvilliana</i> ) - Wherever found	E	\$849,844	\$1,153,484,410
153	Owl, Mexican spotted ( <i>Strix occidentalis lucida</i> ) - Wherever found	T	\$846,956	\$1,154,331,366
154	Gibbons ( <i>Hylobates</i> spp. (including <i>Nomascus</i> )) - Wherever found	E	\$843,638	\$1,155,175,004
155	Whale, false killer ( <i>Pseudorca crassidens</i> ) - Main Hawaiian Islands Insular DPS	E	\$822,143	\$1,155,997,147
156	Coot, Hawaiian ( <i>Fulica americana alai</i> ) - Wherever found	E	\$811,510	\$1,156,808,657
157	Sawfish, smalltooth ( <i>Pristis pectinata</i> ) - US DPS - Smalltooth sawfish originating from U.S. waters.	E	\$793,678	\$1,157,602,335
158	Bat, Florida bonneted ( <i>Eumops floridanus</i> ) - Wherever found	E	\$790,607	\$1,158,392,942
159	Butterfly, Karner blue ( <i>Lycæides melissa samuelis</i> ) - Wherever found	E	\$774,582	\$1,159,167,524
160	Tadpole shrimp, vernal pool ( <i>Lepidurus packardii</i> ) - Wherever found	E	\$770,515	\$1,159,938,039
161	Goby, tidewater ( <i>Eucyclogobius newberryi</i> ) - Wherever found	E	\$767,186	\$1,160,705,225
162	Beetle, valley elderberry longhorn ( <i>Desmocerus californicus dimorphus</i> ) - Wherever found	T	\$750,628	\$1,161,455,853
163	Petrel, Hawaiian ( <i>Pterodroma sandwichensis</i> ) - Wherever found	E	\$748,165	\$1,162,204,018
164	Wolf, red ( <i>Canis rufus</i> ) - U.S.A. (portions of NC and TN)	EXPN	\$728,400	\$1,162,932,418
165	Gallinule, Hawaiian common ( <i>Gallinula galeata sandvicensis</i> ) - Wherever found	E	\$721,990	\$1,163,654,408
166	Sea turtle, green ( <i>Chelonia mydas</i> ) - Central South Pacific DPS	E	\$719,102	\$1,164,373,510
167	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - Chesapeake Bay DPS	E	\$711,497	\$1,165,085,007
168	Prairie dog, Utah ( <i>Cynomys parvidens</i> ) - Wherever found	T	\$702,600	\$1,165,787,607
169	Paintbrush, golden ( <i>Castilleja levisecta</i> ) - Wherever found	T	\$699,359	\$1,166,486,966
170	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - Bitterroot Grizzly Bear Nonessential Experimental Population Area	EXPN	\$689,596	\$1,167,176,562
171	Topminnow, Gila (incl. Yaqui) ( <i>Poeciliopsis occidentalis</i> ) - Wherever found	E	\$679,580	\$1,167,856,142
172	Minnnow, loach ( <i>Tiaroga cobitis</i> ) - Wherever found	E	\$673,507	\$1,168,529,649
173	Spikedace ( <i>Meda fulgida</i> ) - Wherever found	E	\$671,507	\$1,169,201,156
174	Chaffseed, American ( <i>Schwalbea americana</i> ) - Wherever found	E	\$665,144	\$1,169,866,300

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
175	Coqui, Llanero ( <i>Eleutherodactylus juanariveroi</i> ) - Wherever found	E	\$657,024	\$1,170,523,324
176	Crane, whooping ( <i>Grus americana</i> ) - Wherever found, except where listed as an experimental	E	\$645,013	\$1,171,168,337
177	Lupine, Kincaid's ( <i>Lupinus sulphureus</i> ssp. <i>kincaidii</i> ) - Wherever found	T	\$640,449	\$1,171,808,786
178	Chub, Gila ( <i>Gila intermedia</i> ) - Wherever found	E	\$637,200	\$1,172,445,986
179	Coral, staghorn ( <i>Acropora cervicornis</i> ) - Wherever found	T	\$630,397	\$1,173,076,383
180	Wedgemussel, dwarf ( <i>Alasmidonta heterodon</i> ) - Wherever found	E	\$620,099	\$1,173,696,482
181	Turtle, bog ( <i>Clemmys muhlenbergii</i> ) - Wherever found, except GA, NC, SC, TN, VA	T	\$607,406	\$1,174,303,888
182	Salamander, California tiger ( <i>Ambystoma californiense</i> ) - U.S.A. (CA - Sonoma County)	E	\$604,969	\$1,174,908,857
183	Sea turtle, olive ridley ( <i>Lepidochelys olivacea</i> ) - Wherever found, except when listed as endangered under 50 CFR 224.101	T	\$599,216	\$1,175,508,073
184	Kangaroo rat, Stephens' ( <i>Dipodomys stephensi</i> (incl. <i>D. cascus</i> )) - Wherever found	E	\$596,888	\$1,176,104,961
185	Pronghorn, Sonoran ( <i>Antilocapra americana sonoriensis</i> ) - U.S.A. (AZ), Mexico	EXPN	\$596,780	\$1,176,701,741
186	Whale, North Pacific Right ( <i>Eubalaena japonica</i> ) - Wherever found	E	\$595,341	\$1,177,297,082
187	No Common Name ( <i>Acropora globiceps</i> ) - Wherever found	T	\$594,055	\$1,177,891,137
188	Coral, elkhorn ( <i>Acropora palmata</i> ) - Wherever found	T	\$592,097	\$1,178,483,234
189	Crane, whooping ( <i>Grus americana</i> ) - U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	EXPN	\$590,023	\$1,179,073,257
190	Salamander, frosted flatwoods ( <i>Ambystoma cingulatum</i> ) - Wherever found	T	\$581,932	\$1,179,655,189
191	Rabbit, Columbia Basin Pygmy ( <i>Brachylagus idahoensis</i> ) - Columbia Basin DPS	E	\$579,720	\$1,180,234,909
192	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - South Atlantic DPS	E	\$576,030	\$1,180,810,939
193	Salamander, California tiger ( <i>Ambystoma californiense</i> ) - U.S.A. (CA - Santa Barbara County)	E	\$569,088	\$1,181,380,027
194	Peppergrass, Slickspot ( <i>Lepidium papilliferum</i> ) -	T	\$562,008	\$1,181,942,035
195	No Common Name ( <i>Acropora retusa</i> ) - Wherever found	T	\$561,338	\$1,182,503,373
196	Coral, Mountainous Star ( <i>Orbicella faveolata</i> ) - Wherever found	T	\$558,642	\$1,183,062,015
197	Coral, Lobed Star ( <i>Orbicella annularis</i> ) - Wherever found	T	\$554,942	\$1,183,616,957
197	Coral, Rough Cactus ( <i>Mycetophyllia ferox</i> ) - Wherever found	T	\$554,942	\$1,184,171,899
198	Seagrass, Johnson's ( <i>Halophila johnsonii</i> ) - Wherever found	T	\$549,188	\$1,184,721,087
199	No Common Name ( <i>Acropora speciosa</i> ) - Wherever found	T	\$545,217	\$1,185,266,304
200	Spinymussel, James ( <i>Pleurobema collina</i> ) - Wherever found	E	\$539,230	\$1,185,805,534
201	No Common Name ( <i>Acropora jacquelineae</i> ) - Wherever found	T	\$538,000	\$1,186,343,534
201	No Common Name ( <i>Euphyllia paradivisa</i> ) - Wherever found	T	\$538,000	\$1,186,881,534
201	No Common Name ( <i>Isopora crateriformis</i> ) - Wherever found	T	\$538,000	\$1,187,419,534



Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
202	Phlox, Yreka ( <i>Phlox hirsuta</i> ) - Wherever found	E	\$533,200	\$1,187,952,734
203	Coral, Boulder star ( <i>Orbicella franksi</i> ) - Wherever found	T	\$528,942	\$1,188,481,676
204	Butterfly, Oregon silverspot ( <i>Speyeria zerene hippolyta</i> ) - Wherever found	T	\$528,911	\$1,189,010,587
205	Frog, mountain yellow-legged ( <i>Rana muscosa</i> ) - Northern California DPS	E	\$527,842	\$1,189,538,429
206	Trout, Paiute cutthroat ( <i>Oncorhynchus clarkii seleniris</i> ) - Wherever found	T	\$517,327	\$1,190,055,756
207	Vole, Amargosa ( <i>Microtus californicus scirpensis</i> ) - Wherever found	E	\$513,800	\$1,190,569,556
208	Checkerspot, Taylor's (=whulge) ( <i>Euphydryas editha taylori</i> ) - Wherever found	E	\$511,902	\$1,191,081,458
209	Grouper, Nassau ( <i>Epinephelus striatus</i> ) -	T	\$510,951	\$1,191,592,409
210	Lomatium, Bradshaw's ( <i>Lomatium bradshawii</i> ) - Wherever found	E	\$510,169	\$1,192,102,578
211	Rhinoceros, Sumatran ( <i>Dicerorhinus sumatrensis</i> ) - Wherever found	E	\$505,326	\$1,192,607,904
212	Layia, beach ( <i>Layia carnosa</i> ) - Wherever found	E	\$501,105	\$1,193,109,009
213	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - Carolina DPS	E	\$500,494	\$1,193,609,503
214	Coral, Pillar ( <i>Dendrogyra cylindrus</i> ) - Wherever found	NS	\$497,000	\$1,194,106,503
215	Pocket gopher, Tenino ( <i>Thomomys mazama tumuli</i> ) - Wherever found	T	\$489,991	\$1,194,596,494
216	Pocket gopher, Olympia ( <i>Thomomys mazama pugetensis</i> ) - Wherever found	T	\$489,868	\$1,195,086,362
217	Warbler, Kirtland's ( <i>Setophaga kirtlandii</i> (= <i>Dendroica kirtlandii</i> )) - Wherever found	E	\$488,312	\$1,195,574,674
218	Bumble bee, Rusty patched ( <i>Bombus affinis</i> ) - Wherever found	E	\$481,254	\$1,196,055,928
219	Skipperling, Poweshiek ( <i>Oarisma poweshiek</i> ) - Wherever found	E	\$476,980	\$1,196,532,908
220	Pocket gopher, Yelm ( <i>Thomomys mazama yelmensis</i> ) - Wherever found	T	\$475,801	\$1,197,008,709
221	Kangaroo rat, San Bernardino Merriam's ( <i>Dipodomys merriami parvus</i> ) - Wherever found	E	\$473,138	\$1,197,481,847
222	Frog, dusky gopher ( <i>Rana sevosia</i> ) - Wherever found	E	\$462,568	\$1,197,944,415
223	Sucker, Warner ( <i>Catostomus warnerensis</i> ) - Wherever found	T	\$461,280	\$1,198,405,695
224	Daisy, Willamette ( <i>Erigeron decumbens</i> ) - Wherever found	E	\$453,827	\$1,198,859,522
225	Mussel, snuffbox ( <i>Epioblasma triquetra</i> ) - Wherever found	E	\$445,732	\$1,199,305,254
226	Cui-ui ( <i>Chasmistes cujus</i> ) - Wherever found	E	\$444,869	\$1,199,750,123
227	Prairie-chicken, Attwater's greater ( <i>Tympanuchus cupido attwateri</i> ) - Wherever found	E	\$443,335	\$1,200,193,458
228	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Central America DPS	E	\$440,142	\$1,200,633,600
229	Puma (=mountain lion) ( <i>Puma</i> (= <i>Felis</i> ) <i>concolor</i> (all subsp. except <i>coryi</i> )) - FL	SAT	\$437,570	\$1,201,071,170
230	Mouse, Pacific pocket ( <i>Perognathus longimembris pacificus</i> ) - Wherever found	E	\$436,400	\$1,201,507,570
231	Otter, southern sea ( <i>Enhydra lutris nereis</i> ) - Wherever found	T	\$434,499	\$1,201,942,069
232	Deer, Columbian white-tailed ( <i>Odocoileus virginianus leucurus</i> ) - Columbia River DPS	T	\$434,169	\$1,202,376,238
233	Salamander, Cheat Mountain ( <i>Plethodon nettingi</i> ) - Wherever found	T	\$432,710	\$1,202,808,948
234	Dragonfly, Hine's emerald ( <i>Somatochlora hineana</i> ) - Wherever found	E	\$431,550	\$1,203,240,498

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
235	Lizard, blunt-nosed leopard ( <i>Gambelia silus</i> ) - Wherever found	E	\$424,335	\$1,203,664,833
236	Sheep, Sierra Nevada bighorn ( <i>Ovis canadensis sierrae</i> ) - Sierra Nevada	E	\$417,980	\$1,204,082,813
237	Otter, Northern Sea ( <i>Enhydra lutris kenyoni</i> ) - Southwest Alaska DPS	T	\$416,561	\$1,204,499,374
238	Brodiaea, thread-leaved ( <i>Brodiaea filifolia</i> ) - Wherever found	T	\$412,468	\$1,204,911,842
239	Fairy shrimp, San Diego ( <i>Branchinecta sandiegonensis</i> ) - Wherever found	E	\$412,393	\$1,205,324,235
240	Crocodile, American ( <i>Crocodylus acutus</i> ) - U.S.A. (FL)	T	\$411,809	\$1,205,736,044
241	Ladies'-tresses, Ute ( <i>Spiranthes diluvialis</i> ) - Wherever found	T	\$396,252	\$1,206,132,296
242	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - South Atlantic Ocean DPS	T	\$390,716	\$1,206,523,012
243	Skink, sand ( <i>Neoseps reynoldsi</i> ) - Wherever found	T	\$388,225	\$1,206,911,237
244	Warbler (=wood), golden-cheeked ( <i>Dendroica chrysoparia</i> ) - Wherever found	E	\$380,440	\$1,207,291,677
245	Bobwhite, masked (quail) ( <i>Colinus virginianus ridgwayi</i> ) - Wherever found	E	\$375,126	\$1,207,666,803
246	Checker-mallow, Nelson's ( <i>Sidalcea nelsoniana</i> ) - Wherever found	T	\$370,950	\$1,208,037,753
247	Rail, light-footed clapper ( <i>Rallus longirostris levipes</i> ) - Wherever found	E	\$370,158	\$1,208,407,911
248	Beetle, Casey's June ( <i>Dinacoma caseyi</i> ) - Wherever found	E	\$368,066	\$1,208,775,977
249	Seal, Guadalupe fur ( <i>Arctocephalus townsendi</i> ) - Wherever found	T	\$366,042	\$1,209,142,019
250	Toad, Wyoming ( <i>Bufo hemiophrys baxteri</i> ) - Wherever found	E	\$361,051	\$1,209,503,070
251	Grass, Tennessee yellow-eyed ( <i>Xyris tennesseensis</i> ) - Wherever found	E	\$360,480	\$1,209,863,550
252	Sparrow, Florida grasshopper ( <i>Ammodramus savannarum floridanus</i> ) - Wherever found	E	\$358,094	\$1,210,221,644
253	Salamander, Barton Springs ( <i>Eurycea sosorum</i> ) - Wherever found	E	\$348,726	\$1,210,570,370
254	Mouse, New Mexico meadow jumping ( <i>Zapus hudsonius luteus</i> ) - Wherever found	E	\$345,206	\$1,210,915,576
255	Pearlshell, Louisiana ( <i>Margaritifera hembeli</i> ) - Wherever found	T	\$343,600	\$1,211,259,176
256	Catchfly, Spalding's ( <i>Silene spaldingii</i> ) - Wherever found	T	\$335,250	\$1,211,594,426
257	Snake, San Francisco garter ( <i>Thamnophis sirtalis tetrataenia</i> ) - Wherever found	E	\$331,894	\$1,211,926,320
258	Heelsplitter, Carolina ( <i>Lasmigona decorata</i> ) - Wherever found	E	\$331,880	\$1,212,258,200
259	Fritillary, Gentner's ( <i>Fritillaria gentneri</i> ) - Wherever found	E	\$325,500	\$1,212,583,700
260	Gorilla ( <i>Gorilla gorilla</i> ) - Wherever found	E	\$319,266	\$1,212,902,966
261	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - Northeast Atlantic Ocean DPS	E	\$318,653	\$1,213,221,619
262	Swiftlet, Mariana gray ( <i>Aerodramus vanikorensis bartschi</i> ) - Wherever found	E	\$317,823	\$1,213,539,442
263	Jaguar ( <i>Panthera onca</i> ) - Wherever found	E	\$317,482	\$1,213,856,924
264	Sea turtle, olive ridley ( <i>Lepidochelys olivacea</i> ) - Breeding colony populations on Pacific coast of	E	\$316,689	\$1,214,173,613
265	Hellbender, Ozark ( <i>Cryptobranchus alleganiensis bishopi</i> ) - Wherever found	E	\$312,787	\$1,214,486,400
266	Squirrel, Northern Idaho Ground ( <i>Uroditellus brunneus</i> ) - Wherever found	T	\$309,992	\$1,214,796,392
267	Rockfish, canary ( <i>Sebastes pinniger</i> ) - Puget Sound - Georgia Basin DPS	T	\$308,530	\$1,215,104,922

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
268	Hawk, Hawaiian (=lo) ( <i>Buteo solitarius</i> ) - Wherever found	E	\$302,874	\$1,215,407,796
269	Butterfly, Saint Francis' satyr ( <i>Neonympha mitchellii francisci</i> ) - Wherever found	E	\$296,585	\$1,215,704,381
270	Pocket gopher, Roy Prairie ( <i>Thomomys mazama glacialis</i> ) - Wherever found	T	\$296,401	\$1,216,000,782
271	Toad, Yosemite ( <i>Anaxyrus canorus</i> ) - Wherever found	T	\$289,366	\$1,216,290,148
272	Darter, Okaloosa ( <i>Etheostoma okaloosae</i> ) - Wherever found	T	\$287,000	\$1,216,577,148
273	Wallflower, Ben Lomond ( <i>Erysimum teretifolium</i> ) - Wherever found	E	\$286,400	\$1,216,863,548
274	Bat, Virginia big-eared ( <i>Corynorhinus (=Plecotus) townsendii virginianus</i> ) - Wherever found	E	\$286,070	\$1,217,149,618
275	Amaranth, seabeach ( <i>Amaranthus pumilus</i> ) - Wherever found	T	\$281,793	\$1,217,431,411
276	Milk-vetch, Peirson's ( <i>Astragalus magdalenae</i> var. <i>peirsonii</i> ) - Wherever found	T	\$279,300	\$1,217,710,711
277	Sumac, Michaux's ( <i>Rhus michauxii</i> ) - Wherever found	E	\$277,915	\$1,217,988,626
278	Fairy shrimp, Conservancy ( <i>Branchinecta conservatio</i> ) - Wherever found	E	\$276,754	\$1,218,265,380
279	Elktoe, Appalachian ( <i>Alasmidonta raveneliana</i> ) - Wherever found	E	\$276,214	\$1,218,541,594
280	Skipper, Dakota ( <i>Hesperia dacotae</i> ) - Wherever found	T	\$275,650	\$1,218,817,244
281	Dace, desert ( <i>Eremichthys acros</i> ) - Wherever found	T	\$270,813	\$1,219,088,057
282	Chub, Pahrnagat roundtail ( <i>Gila robusta jordani</i> ) - Wherever found	E	\$268,917	\$1,219,356,974
283	Mucket, pink (pearlymussel) ( <i>Lampsilis abrupta</i> ) - Wherever found	E	\$266,870	\$1,219,623,844
284	Pogonia, small whorled ( <i>Isotria medeoloides</i> ) -	T	\$262,488	\$1,219,886,332
285	Bankclimber, purple (mussel) ( <i>Elliptoideus sloatianus</i> ) - Wherever found	T	\$262,240	\$1,220,148,572
286	Rabbitsfoot ( <i>Quadrula cylindrica cylindrica</i> ) - Wherever found	T	\$261,729	\$1,220,410,301
287	Woundfin ( <i>Plagopterus argentissimus</i> ) - Wherever found, except where listed as an experimental population	E	\$260,617	\$1,220,670,918
288	Loosestrife, rough-leaved ( <i>Lysimachia asperulaefolia</i> ) - Wherever found	E	\$258,240	\$1,220,929,158
289	Crane, Mississippi sandhill ( <i>Grus canadensis pulla</i> ) - Wherever found	E	\$255,126	\$1,221,184,284
290	Whipsnake (=striped racer), Alameda ( <i>Masticophis lateralis euryxanthus</i> ) - Wherever found	T	\$250,529	\$1,221,434,813
291	Mucket, Neosho ( <i>Lampsilis rafinesqueana</i> ) - Wherever found	E	\$249,446	\$1,221,684,259
292	Bison, wood ( <i>Bison bison athabascaae</i> ) - Wherever found	T	\$248,017	\$1,221,932,276
293	Creeper, Hawaii ( <i>Oreomystis mana</i> ) - Wherever found	E	\$247,893	\$1,222,180,169
294	Toad, Houston ( <i>Bufo houstonensis</i> ) - Wherever found	E	\$247,394	\$1,222,427,563
295	Butterfly, bay checkerspot ( <i>Euphydryas editha bayensis</i> ) - Wherever found	T	\$243,925	\$1,222,671,488
296	Crow, Hawaiian (=alala) ( <i>Corvus hawaiiensis</i> ) - Wherever found	E	\$242,867	\$1,222,914,355
297	No Common Name ( <i>Seriatopora aculeata</i> ) - Wherever found	T	\$242,055	\$1,223,156,410
298	Mussel, sheepnose ( <i>Plethobasus cyphus</i> ) - Wherever found	E	\$237,630	\$1,223,394,040
299	Akekee ( <i>Loxops caeruleirostris</i> ) - Wherever found	E	\$233,070	\$1,223,627,110

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
300	Caribou DPS, Southern Mountain ( <i>Rangifer tarandus</i> ssp. <i>caribou</i> ) - Southern Mountain DPS	E	\$233,032	\$1,223,860,142
301	Dace, Moapa ( <i>Moapa coriacea</i> ) - Wherever found	E	\$230,898	\$1,224,091,040
302	Sucker, Santa Ana ( <i>Catostomus santaanae</i> ) - 3 CA river basins	T	\$229,761	\$1,224,320,801
303	Rhinoceros, Javan ( <i>Rhinoceros sondaicus</i> ) - Sikkim, Thailand, Malaysia, Indonesia, Indochina, Burma, Bangladesh	E	\$225,852	\$1,224,546,653
304	Shearwater, Newell's Townsend's ( <i>Puffinus auricularis newelli</i> ) - Wherever found	T	\$223,460	\$1,224,770,113
305	Baccharis, Encinitas ( <i>Baccharis vanessae</i> ) - Wherever found	T	\$222,760	\$1,224,992,873
306	Fiddleneck, large-flowered ( <i>Amsinckia grandiflora</i> ) - Wherever found	E	\$219,763	\$1,225,212,636
307	Turtle, bog ( <i>Clemmys muhlenbergii</i> ) - U.S.A. (GA, NC, SC, TN, VA)	SAT	\$219,543	\$1,225,432,179
308	Chub, Virgin River ( <i>Gila seminuda</i> (=robusta)) - Wherever found	E	\$219,372	\$1,225,651,551
309	Pocketbook, fat ( <i>Potamilus capax</i> ) - Wherever found	E	\$219,004	\$1,225,870,555
310	Darter, duskytail ( <i>Etheostoma percnurum</i> ) - Wherever found	E	\$216,341	\$1,226,086,896
311	Fairy shrimp, Riverside ( <i>Streptocephalus woottoni</i> ) - Wherever found	E	\$215,595	\$1,226,302,491
312	Frog, Chiricahua leopard ( <i>Rana chiricahuensis</i> ) - Wherever found	T	\$214,983	\$1,226,517,474
313	Akepa, Maui ( <i>Loxops ochraceus</i> ) - Wherever found	E	\$211,096	\$1,226,728,570
314	Button-celery, San Diego ( <i>Eryngium aristulatum</i> var. <i>parishii</i> ) - Wherever found	E	\$206,690	\$1,226,935,260
315	Snail, Snake River physa ( <i>Physa natricina</i> ) - Wherever found	E	\$205,338	\$1,227,140,598
316	Spineflower, Orcutt's ( <i>Chorizanthe orcuttiana</i> ) - Wherever found	E	\$204,484	\$1,227,345,082
317	Shiner, Arkansas River ( <i>Notropis girardi</i> ) - Arkansas River Basin (AR, KS, NM, OK, TX)	T	\$202,906	\$1,227,547,988
318	Coneflower, smooth ( <i>Echinacea laevigata</i> ) - Wherever found	E	\$200,266	\$1,227,748,254
319	Checkermallow, Wenatchee Mountains ( <i>Sidalcea oregana</i> var. <i>calva</i> ) - Wherever found	E	\$199,000	\$1,227,947,254
320	Kangaroo rat, giant ( <i>Dipodomys ingens</i> ) - Wherever found	E	\$198,458	\$1,228,145,712
321	Pangolin, Temnick's ground ( <i>Manis temminckii</i> ) - Wherever found	E	\$193,532	\$1,228,339,244
322	Trout, Gila ( <i>Oncorhynchus gilae</i> ) - Wherever found	T	\$192,957	\$1,228,532,201
323	Butterfly, Mitchell's satyr ( <i>Neonympha mitchellii mitchellii</i> ) - Wherever found	E	\$191,055	\$1,228,723,256
324	Ambrosia, San Diego ( <i>Ambrosia pumila</i> ) - Wherever found	E	\$189,000	\$1,228,912,256
325	Tern, roseate ( <i>Sterna dougallii dougallii</i> ) - Northeast U.S. nesting population	E	\$188,041	\$1,229,100,297
326	Moth, Blackburn's sphinx ( <i>Manduca blackburni</i> ) - Wherever found	E	\$186,620	\$1,229,286,917
327	Butterfly, Lange's metalmark ( <i>Apodemia mormo langei</i> ) - Wherever found	E	\$185,719	\$1,229,472,636
328	Gartersnake, northern Mexican ( <i>Thamnophis eques megalops</i> ) - Wherever found	T	\$183,975	\$1,229,656,611
329	Navarretia, spreading ( <i>Navarretia fossalis</i> ) - Wherever found	T	\$181,700	\$1,229,838,311
330	Sheep, Peninsular bighorn ( <i>Ovis canadensis nelsoni</i> ) - Peninsular CA pop.	E	\$180,500	\$1,230,018,811
331	Darter, Niangua ( <i>Etheostoma nianguae</i> ) - Wherever found	T	\$177,065	\$1,230,195,876

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
332	Clover, running buffalo ( <i>Trifolium stoloniferum</i> ) -	E	\$176,918	\$1,230,372,794
333	Popcornflower, rough ( <i>Plagiobothrys hirtus</i> ) - Wherever found	E	\$171,700	\$1,230,544,494
334	Mouse, southeastern beach ( <i>Peromyscus polionotus niveiventris</i> ) - wherever found	T	\$171,320	\$1,230,715,814
335	Penstemon, blowout ( <i>Penstemon haydenii</i> ) - Wherever found	E	\$169,049	\$1,230,884,863
336	Whale, gray ( <i>Eschrichtius robustus</i> ) - Western North Pacific DPS	E	\$168,135	\$1,231,052,998
337	Akepa, Hawaii ( <i>Loxops coccineus</i> ) - Wherever found	E	\$165,479	\$1,231,218,477
338	Logperch, Roanoke ( <i>Percina rex</i> ) - Wherever found	E	\$165,140	\$1,231,383,617
339	Spectaclecase (mussel) ( <i>Cumberlandia monodonta</i> ) - Wherever found	E	\$165,066	\$1,231,548,683
340	Butterfly, mission blue ( <i>Icaricia icarioides missionensis</i> ) - Wherever found	E	\$162,636	\$1,231,711,319
341	Paintbrush, San Clemente Island ( <i>Castilleja grisea</i> ) - Wherever found	T	\$162,217	\$1,231,873,536
342	Bush-mallow, San Clemente Island ( <i>Malacothamnus clementinus</i> ) - Wherever found	E	\$161,917	\$1,232,035,453
343	Riffleshell, northern ( <i>Epioblasma torulosa rangiana</i> ) - Wherever found	E	\$161,586	\$1,232,197,039
344	Quillwort, Louisiana ( <i>Isoetes louisianensis</i> ) - Wherever found	E	\$161,116	\$1,232,358,155
345	Kangaroo rat, Tipton ( <i>Dipodomys nitratoides nitratoides</i> ) - Wherever found	E	\$160,300	\$1,232,518,455
346	Threeridge, fat (mussel) ( <i>Amblema neislerii</i> ) - Wherever found	E	\$159,269	\$1,232,677,724
347	Lizard, Coachella Valley fringe-toed ( <i>Uma inornata</i> ) - Wherever found	T	\$158,929	\$1,232,836,653
348	Dolphin, Maui ( <i>Cephalorhynchus hectori maui</i> )	E	\$158,636	\$1,232,995,289
349	Howellia, Water ( <i>Howellia aquatilis</i> ) -	T	\$157,793	\$1,233,153,082
350	Falcon, Northern Aplomado ( <i>Falco femoralis septentrionalis</i> ) - Wherever found, except where listed as an experimental population	E	\$157,326	\$1,233,310,408
351	Pink, swamp ( <i>Helonias bullata</i> ) - Wherever found	T	\$155,466	\$1,233,465,874
352	Milkweed, Mead's ( <i>Asclepias meadii</i> ) - Wherever found	T	\$153,841	\$1,233,619,715
353	Chub, Borax Lake ( <i>Gila boraxobius</i> ) - Wherever found	E	\$153,400	\$1,233,773,115
354	Albatross, short-tailed ( <i>Phoebastria (=Diomedea) albatrus</i> ) - Wherever found	E	\$153,269	\$1,233,926,384
355	Rockcress, Santa Cruz Island ( <i>Sibara filifolia</i> ) - Wherever found	E	\$152,963	\$1,234,079,347
356	'I'iwi ( <i>Drepanis coccinea</i> ) - Wherever found	T	\$152,001	\$1,234,231,348
357	Woodland-star, San Clemente Island ( <i>Lithophragma maximum</i> ) - Wherever found	E	\$149,451	\$1,234,380,799
357	Larkspur, San Clemente Island ( <i>Delphinium variegatum</i> ssp. <i>kinkiense</i> ) - Wherever found	E	\$149,451	\$1,234,530,250
357	Lotus, San Clemente Island ( <i>Acemison dendroideus</i> var. <i>traskiae</i> ) - Wherever found	T	\$149,451	\$1,234,679,701
358	Snail, Morro shoulderband (=Banded dune) ( <i>Helminthoglypta walkeriana</i> ) - Wherever found	E	\$148,955	\$1,234,828,656
359	Toad, Puerto Rican crested ( <i>Peltophryne lemur</i> ) - Wherever found	T	\$147,491	\$1,234,976,147
360	Cactus, Colorado hookless ( <i>Sclerocactus glaucus</i> ) - Wherever found	T	\$145,933	\$1,235,122,080
361	Butterfly, Mariana wandering ( <i>Vagrans egistina</i> ) - Wherever found	E	\$145,892	\$1,235,267,972

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
362	Rhinoceros, Southern white ( <i>Ceratotherium simum</i> ssp. <i>simum</i> ) - Wherever found	SAT	\$145,451	\$1,235,413,423
363	Pigtoe, oval ( <i>Pleurobema pyriforme</i> ) - Wherever found	E	\$145,152	\$1,235,558,575
364	Salamander, Reticulated flatwoods ( <i>Ambystoma bishopi</i> ) - Wherever found	E	\$144,369	\$1,235,702,944
365	Piperia, Yadon's ( <i>Piperia yadonii</i> ) - Wherever found	E	\$143,605	\$1,235,846,549
366	Parrotbill, Maui (Kiwikiu) ( <i>Pseudonestor xanthophrys</i> ) - Wherever found	E	\$141,533	\$1,235,988,082
367	Skink, bluetail mole ( <i>Eumeces egregius lividus</i> ) - Wherever found	T	\$141,106	\$1,236,129,188
368	Hawk, Puerto Rican sharp-shinned ( <i>Accipiter striatus venator</i> ) - Wherever found	E	\$139,750	\$1,236,268,938
369	Wallflower, Contra Costa ( <i>Erysimum capitatum</i> var. <i>angustatum</i> ) - Wherever found	E	\$139,700	\$1,236,408,638
370	Mouse, Preble's meadow jumping ( <i>Zapus hudsonius preblei</i> ) - wherever found	T	\$138,584	\$1,236,547,222
371	Goldfields, Contra Costa ( <i>Lasthenia conjugens</i> ) - Wherever found	E	\$137,863	\$1,236,685,085
372	Wallflower, Menzies' ( <i>Erysimum menziesii</i> ) - Wherever found	E	\$137,800	\$1,236,822,885
373	Wooly-threads, San Joaquin ( <i>Monolopia (=Lembertia) congdonii</i> ) - Wherever found	E	\$137,504	\$1,236,960,389
374	Frog, mountain yellow-legged ( <i>Rana muscosa</i> ) - Southern California DPS	E	\$136,821	\$1,237,097,210
375	'Ahinahina ( <i>Argyroxiphium sandwicense</i> ssp. <i>macrocephalum</i> ) - Wherever found	T	\$136,705	\$1,237,233,915
376	Fadang ( <i>Cycas micronesica</i> ) - Wherever found	T	\$136,529	\$1,237,370,444
377	Iris, dwarf lake ( <i>Iris lacustris</i> ) - Wherever found	T	\$136,442	\$1,237,506,886
378	Darter, watercress ( <i>Etheostoma nuchale</i> ) - Wherever found	E	\$135,620	\$1,237,642,506
379	Ivesia, Webber's ( <i>Ivesia webberi</i> ) - Wherever found	T	\$135,504	\$1,237,778,010
380	Falcon, Northern Aplomado ( <i>Falco femoralis septentrionalis</i> ) - U.S.A (AZ, NM)	EXPN	\$134,200	\$1,237,912,210
381	Pupfish, Ash Meadows Amargosa ( <i>Cyprinodon nevadensis mionectes</i> ) - Wherever found	E	\$133,581	\$1,238,045,791
382	Fanshell ( <i>Cyprogenia stegaria</i> ) - Wherever found	E	\$133,340	\$1,238,179,131
383	Evening-primrose, Antioch Dunes ( <i>Oenothera deltoides</i> ssp. <i>howellii</i> ) - Wherever found	E	\$131,300	\$1,238,310,431
384	No Common Name ( <i>Stenogyne angustifolia</i> var. <i>angustifolia</i> ) - Wherever found	E	\$131,067	\$1,238,441,498
385	Limpet, Banbury Springs ( <i>Lanx</i> sp.) - Wherever found	E	\$129,445	\$1,238,570,943
386	Condor, California ( <i>Gymnogyps californianus</i> ) - U.S.A. (specific portions of Arizona, Nevada, and Utah)	EXPN	\$129,227	\$1,238,700,170
387	Desert buckwheat, Umtanum ( <i>Eriogonum codium</i> ) - Wherever found	T	\$129,037	\$1,238,829,207
387	Bladderpod, White Bluffs ( <i>Physaria douglasii</i> ssp. <i>tuplashensis</i> ) - Wherever found	T	\$129,037	\$1,238,958,244
388	Dace, blackside ( <i>Phoxinus cumberlandensis</i> ) - Wherever found	T	\$128,929	\$1,239,087,173
389	Butterfly, Mariana eight-spot ( <i>Hypolimnas octocula marianensis</i> ) - Wherever found	E	\$127,694	\$1,239,214,867
390	Cactus, Kuenzler hedgehog ( <i>Echinocereus fendleri</i> var. <i>kuenzleri</i> ) - Wherever found	T	\$127,386	\$1,239,342,253
391	Orchid, western prairie fringed ( <i>Platanthera praeclara</i> ) - Wherever found	T	\$127,300	\$1,239,469,553
392	Slabshell, Chipola ( <i>Elliptio chipolaensis</i> ) - Wherever found	T	\$127,240	\$1,239,596,793
393	Akikiki ( <i>Oreomystis bairdi</i> ) - Wherever found	E	\$127,076	\$1,239,723,869

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
394	Deer, key ( <i>Odocoileus virginianus clavium</i> ) - Wherever found	E	\$126,231	\$1,239,850,100
395	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Middle Columbia River DPS	EXPN	\$125,750	\$1,239,975,850
396	Akiapolaau ( <i>Hemignathus wilsoni</i> ) - Wherever found	E	\$125,619	\$1,240,101,469
397	Amole, purple ( <i>Chlorogalum purpureum</i> ) - Wherever found	T	\$125,097	\$1,240,226,566
398	Bat, Mariana fruit (=Mariana flying fox) ( <i>Pteropus mariannus mariannus</i> ) - Wherever found	T	\$124,704	\$1,240,351,270
399	Mouse, Perdido Key beach ( <i>Peromyscus polionotus trissyllepsis</i> ) - Wherever found	E	\$124,134	\$1,240,475,404
400	No Common Name ( <i>Achyranthes mutica</i> ) - Wherever found	E	\$123,569	\$1,240,598,973
401	Kingfisher, Guam Micronesian ( <i>Halcyon cinnamomina cinnamomina</i> ) - Wherever found	E	\$123,513	\$1,240,722,486
402	Crayfish, Big Sandy ( <i>Cambarus callainus</i> ) - Wherever found	T	\$123,498	\$1,240,845,984
403	Honeycreeper, (Akohekohe) crested ( <i>Palmeria dolei</i> ) - Wherever found	E	\$123,135	\$1,240,969,119
404	Clubshell ( <i>Pleurobema clava</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$120,414	\$1,241,089,533
405	Jewelflower, California ( <i>Caulanthus californicus</i> ) - Wherever found	E	\$120,004	\$1,241,209,537
406	Tiger beetle, Puritan ( <i>Cicindela puritana</i> ) - Wherever found	T	\$112,589	\$1,241,322,126
407	Yellowhead, desert ( <i>Yermo xanthocephalus</i> ) - Wherever found	T	\$111,500	\$1,241,433,626
408	Tiger beetle, Salt Creek ( <i>Cicindela nevadica lincolniana</i> ) - Wherever found	E	\$110,917	\$1,241,544,543
409	No Common Name ( <i>Tabernaemontana rotensis</i> ) - Wherever found	T	\$110,296	\$1,241,654,839
410	Cactus, Wright fishhook ( <i>Sclerocactus wrightiae</i> ) - Wherever found	E	\$109,750	\$1,241,764,589
411	Palila (honeycreeper) ( <i>Loxioides bailleui</i> ) - Wherever found	E	\$109,177	\$1,241,873,766
412	Lance, yellow ( <i>Elliptio lanceolata</i> ) - Wherever found	T	\$108,746	\$1,241,982,512
413	Snail, Guam tree ( <i>Partula radiolata</i> ) - Wherever found	E	\$108,451	\$1,242,090,963
414	Gartersnake, narrow-headed ( <i>Thamnophis rufipunctatus</i> ) - Wherever found	T	\$107,760	\$1,242,198,723
415	Snail, Humped tree ( <i>Partula gibba</i> ) - Wherever found	E	\$106,919	\$1,242,305,642
416	Thrush, large Kauai (=kamao) ( <i>Myadestes myadestinus</i> ) - Wherever found	E	\$104,441	\$1,242,410,083
417	Iagu, Hayun (=Guam), Tronkon guafi (Rota) ( <i>Serianthes nelsonii</i> ) - Wherever found	E	\$104,052	\$1,242,514,135
418	No Common Name ( <i>Tuberolabium guamense</i> ) - Wherever found	T	\$103,426	\$1,242,617,561
419	No Common Name ( <i>Tetramolopium rockii</i> ) - Wherever found	T	\$102,981	\$1,242,720,542
420	Mouse, Alabama beach ( <i>Peromyscus polionotus ammobates</i> ) - Wherever found	E	\$102,594	\$1,242,823,136
421	Haha ( <i>Cyanea stictophylla</i> ) - Wherever found	E	\$102,329	\$1,242,925,465
422	Snail, Chittenango ovate amber ( <i>Succinea chittenangoensis</i> ) - Wherever found	T	\$100,800	\$1,243,026,265
423	Rhinoceros, great Indian ( <i>Rhinoceros unicornis</i> ) - Nepal, India	E	\$100,672	\$1,243,126,937
424	Cactus, Pima pineapple ( <i>Coryphantha scheeri</i> var. <i>robustispina</i> ) - Wherever found	E	\$100,045	\$1,243,226,982
425	Haha ( <i>Cyanea shipmanii</i> ) - Wherever found	E	\$100,010	\$1,243,326,992

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
426	Tracaja (Podocnemis unifilis) - Wherever found	E	\$99,909	\$1,243,426,901
427	Poolfish, Pahump (Empetrichthys latos) - Wherever found	E	\$99,894	\$1,243,526,795
428	Torreya, Florida (Torreya taxifolia) - Wherever found	E	\$98,932	\$1,243,625,727
429	Snail, Bliss Rapids (Taylorconcha serpenticola) - Wherever found	T	\$98,716	\$1,243,724,443
430	Sea turtle, loggerhead (Caretta caretta) - North Indian Ocean DPS	E	\$98,641	\$1,243,823,084
431	Bat, Ozark big-eared (Corynorhinus (=Plecotus) townsendii ingens) - Wherever found	E	\$98,551	\$1,243,921,635
432	`Akoko, Ewa Plains (Euphorbia skottsbergii var. skottsbergii) - Wherever found	E	\$98,136	\$1,244,019,771
433	Rabbit, Lower Keys marsh (Sylvilagus palustris hefneri) - Wherever found	E	\$97,793	\$1,244,117,564
434	Alani (Melicope munroi) - Wherever found	E	\$97,728	\$1,244,215,292
435	Four-o'clock, MacFarlane's (Mirabilis macfarlanei) - Wherever found	T	\$97,171	\$1,244,312,463
436	Lion (Panthera leo ssp. leo) - Wherever found	E	\$97,076	\$1,244,409,539
436	Lion (Panthera leo ssp. melanochaita) - Wherever found	T	\$97,076	\$1,244,506,615
437	Pupfish, desert (Cyprinodon macularius) - Wherever found	E	\$97,055	\$1,244,603,670
438	Bulrush, Northeastern (Scirpus ancistrochaetus) -	E	\$96,873	\$1,244,700,543
439	Chaff-flower, round-leaved (Achyranthes splendens var. rotundata) - Wherever found	E	\$96,856	\$1,244,797,399
440	Lousewort, Furbish (Pedicularis furbishiae) - Wherever found	E	\$96,220	\$1,244,893,619
441	Ko'oko'olau (Bidens conjuncta) - Wherever found	E	\$96,095	\$1,244,989,714
442	Storm-petrel, band-rumped (Oceanodroma castro) - USA (HI)	E	\$95,937	\$1,245,085,651
443	No Common Name (Stenogyne bifida) - Wherever found	E	\$95,662	\$1,245,181,313
444	Haha (Cyanea superba) - Wherever found	E	\$95,023	\$1,245,276,336
445	Orchid, eastern prairie fringed (Platanthera leucophaea) - Wherever found	T	\$95,017	\$1,245,371,353
446	No Common Name (Stenogyne kealiae) - Wherever found	E	\$94,686	\$1,245,466,039
447	Fly, Hawaiian picture-wing (Drosophila hemipeza) - Wherever found	E	\$94,284	\$1,245,560,323
448	Salamander, Salado (Eurycea chisholmensis) - Wherever found	T	\$94,216	\$1,245,654,539
449	Milk-vetch, Braunton's (Astragalus brauntonii) - Wherever found	E	\$91,884	\$1,245,746,423
450	Thistle, La Graciosa (Cirsium loncholepis) - Wherever found	E	\$91,668	\$1,245,838,091
451	Darter, leopard (Percina pantherina) - Wherever found	T	\$90,174	\$1,245,928,265
452	Milk-vetch, Sentry (Astragalus cremnophylax var. cremnophylax) - Wherever found	E	\$90,000	\$1,246,018,265
453	Spineflower, Monterey (Chorizanthe pungens var. pungens) - Wherever found	T	\$89,144	\$1,246,107,409
454	Harperella (Ptilimnium nodosum) - Wherever found	E	\$89,113	\$1,246,196,522
455	Thistle, Pitcher's (Cirsium pitcheri) - Wherever found	T	\$88,726	\$1,246,285,248
456	Shrew, Buena Vista Lake ornate (Sorex ornatus relictus) - Wherever found	E	\$88,390	\$1,246,373,638
457	Springsnail, Bruneau Hot (Pyrgulopsis bruneauensis) - Wherever found	E	\$87,334	\$1,246,460,972



Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
458	Sunflower, Schweinitz's ( <i>Helianthus schweinitzii</i> ) - Wherever found	E	\$87,125	\$1,246,548,097
459	Tern, roseate ( <i>Sterna dougallii dougallii</i> ) - Western Hemisphere except NE U.S.	T	\$87,000	\$1,246,635,097
460	Stickleback, unarmored threespine ( <i>Gasterosteus aculeatus williamsoni</i> ) - Wherever found	E	\$86,925	\$1,246,722,022
461	No Common Name ( <i>Dendrobium guamense</i> ) - Wherever found	T	\$86,690	\$1,246,808,712
462	Butterfly, Smith's blue ( <i>Euphilotes enoptes smithi</i> ) - Wherever found	E	\$86,261	\$1,246,894,973
463	Springfish, Railroad Valley ( <i>Crenichthys nevadae</i> ) - Wherever found	T	\$86,197	\$1,246,981,170
464	Cavefish, Ozark ( <i>Amblyopsis rosae</i> ) - Wherever found	T	\$86,069	\$1,247,067,239
465	Ohai ( <i>Sesbania tomentosa</i> ) - Wherever found	E	\$85,434	\$1,247,152,673
466	Tiger beetle, Northeastern beach ( <i>Cicindela dorsalis dorsalis</i> ) - Wherever found	T	\$85,300	\$1,247,237,973
467	Sea turtle, green ( <i>Chelonia mydas</i> ) - East Indian-West Pacific DPS	T	\$84,784	\$1,247,322,757
468	Butterfly plant, Colorado ( <i>Gaura neomexicana</i> var. <i>coloradensis</i> ) - Wherever found	T	\$84,274	\$1,247,407,031
469	Po`ouli (honeycreeper) ( <i>Melamprosops phaeosoma</i> ) - Wherever found	E	\$84,263	\$1,247,491,294
470	Mouse, Choctawhatchee beach ( <i>Peromyscus polionotus allopshys</i> ) - Wherever found	E	\$83,281	\$1,247,574,575
471	Pigtoe, finerayed ( <i>Fusconaia cuneolus</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$82,650	\$1,247,657,225
472	Pigtoe, shiny ( <i>Fusconaia cor</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$82,250	\$1,247,739,475
473	Roseroot, Leedy's ( <i>Rhodiola integrifolia</i> ssp. <i>leedyi</i> ) - Wherever found	T	\$81,923	\$1,247,821,398
474	Monkeyface, Cumberland (pearlymussel) ( <i>Quadrula intermedia</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$81,010	\$1,247,902,408
475	Honohono ( <i>Haplostachys haplostachya</i> ) - Wherever found	E	\$80,158	\$1,247,982,566
476	Pocketbook, Ouachita rock ( <i>Arkansia wheeleri</i> ) - Wherever found	E	\$80,151	\$1,248,062,717
477	Grass, Eureka Dune ( <i>Swallenia alexandrae</i> ) -	T	\$79,700	\$1,248,142,417
478	Mussel, scaleshell ( <i>Leptodea leptodon</i> ) - Wherever found	E	\$79,230	\$1,248,221,647
479	Rail, Guam ( <i>Rallus owstoni</i> ) - Wherever found, except where listed as an experimental population	E	\$79,032	\$1,248,300,679
480	Snail, fragile tree ( <i>Samoana fragilis</i> ) - Wherever found	E	\$78,904	\$1,248,379,583
481	Condor, California ( <i>Gymnogyps californianus</i> ) - Pacific Northwest NEP	PEXP	\$78,318	\$1,248,457,901
482	Lomatium, Cook's ( <i>Lomatium cookii</i> ) - Wherever found	E	\$78,310	\$1,248,536,211
483	Megapode, Micronesian ( <i>Megapodius laperouse</i> ) - Wherever found	E	\$78,273	\$1,248,614,484
484	Darter, boulder ( <i>Etheostoma wapiti</i> ) - Wherever found	E	\$78,000	\$1,248,692,484
485	Salamander, Santa Cruz long-toed ( <i>Ambystoma macrodactylum croceum</i> ) - Wherever found	E	\$77,563	\$1,248,770,047
486	Salamander, Shenandoah ( <i>Plethodon shenandoah</i> ) - Wherever found	E	\$77,421	\$1,248,847,468

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
487	Kidneyshell, fluted ( <i>Ptychobranthus subtentus</i> ) - Wherever found	E	\$76,953	\$1,248,924,421
488	No Common Name ( <i>Phyllostegia bracteata</i> ) - Wherever found	E	\$75,290	\$1,248,999,711
489	Butterfly, Mount Charleston blue ( <i>Icaricia (Plebejus) shasta charlestonensis</i> ) - Wherever found	E	\$74,725	\$1,249,074,436
490	Chimpanzee ( <i>Pan troglodytes</i> ) - Wherever found	E	\$74,339	\$1,249,148,775
491	Spinedace, White River ( <i>Lepidomeda albivallis</i> ) - Wherever found	E	\$73,916	\$1,249,222,691
492	Skipper, Carson wandering ( <i>Pseudocopa eodes eunus obscurus</i> ) - Wherever found	E	\$73,075	\$1,249,295,766
493	Tui chub, Mohave ( <i>Gila bicolor ssp. mohavensis</i> ) - Wherever found	E	\$72,789	\$1,249,368,555
494	Jaguarundi, Gulf Coast ( <i>Herpailurus (=Felis) yagouaroundi cacomitli</i> ) - Wherever found	E	\$72,510	\$1,249,441,065
495	No Common Name ( <i>Schiedea haleakalensis</i> ) - Wherever found	E	\$72,452	\$1,249,513,517
496	Halumtano, Cebello ( <i>Bulbophyllum guamense</i> ) - Wherever found	T	\$72,018	\$1,249,585,535
497	Popolo ( <i>Solanum nelsonii</i> ) - Wherever found	E	\$71,338	\$1,249,656,873
498	Aster, decurrent false ( <i>Boltonia decurrens</i> ) - Wherever found	T	\$71,244	\$1,249,728,117
499	No Common Name ( <i>Phyllostegia brevidens</i> ) - Wherever found	E	\$70,196	\$1,249,798,313
500	No Common Name ( <i>Phyllostegia haliakalae</i> ) - Wherever found	E	\$70,099	\$1,249,868,412
501	Pupfish, Owens ( <i>Cyprinodon radiosus</i> ) - Wherever found	E	\$70,024	\$1,249,938,436
502	Sculpin, Grotto ( <i>Cottus specus</i> ) - Wherever found	E	\$69,900	\$1,250,008,336
503	Woodrat, Key Largo ( <i>Neotoma floridana smalli</i> ) - Wherever found	E	\$69,548	\$1,250,077,884
504	Ufa-halomtano ( <i>Heritiera longipetiolata</i> ) - Wherever found	E	\$69,484	\$1,250,147,368
505	Tarplant, Otay ( <i>Deinandra (=Hemizonia) conjugens</i> ) - Wherever found	T	\$68,700	\$1,250,216,068
506	Beetle, Mount Hermon June ( <i>Polyphylla barbata</i> ) - Wherever found	E	\$68,697	\$1,250,284,765
507	Butterfly, Bartram's hairstreak ( <i>Strymon acis bartrami</i> ) - Wherever found	E	\$68,500	\$1,250,353,265
508	Dace, Kendall Warm Springs ( <i>Rhinichthys osculus thermalis</i> ) - Wherever found	E	\$68,000	\$1,250,421,265
509	Polygonum, Scotts Valley ( <i>Polygonum hickmanii</i> ) - Wherever found	E	\$67,300	\$1,250,488,565
510	Pearlymussel, slabside ( <i>Pleurotaia dolabelloides</i> ) - Wherever found	E	\$67,027	\$1,250,555,592
511	Salamander, Jemez Mountains ( <i>Plethodon neomexicanus</i> ) - Wherever found	E	\$67,000	\$1,250,622,592
512	Chub, spotfin ( <i>Erimonax monachus</i> ) - Wherever found, except where listed as an experimental	T	\$66,666	\$1,250,689,258
513	Butterfly, callippe silverspot ( <i>Speyeria callippe callippe</i> ) - Wherever found	E	\$66,600	\$1,250,755,858
514	Pearlymussel, cracking ( <i>Hemistena lata</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$65,710	\$1,250,821,568
515	Popolo ku mai ( <i>Solanum incompletum</i> ) - Wherever found	E	\$64,865	\$1,250,886,433
516	Mountain beaver, Point Arena ( <i>Aplodontia rufa nigra</i> ) - Wherever found	E	\$64,500	\$1,250,950,933
517	Milk-vetch, triple-ribbed ( <i>Astragalus tricarinatus</i> ) - Wherever found	E	\$64,100	\$1,251,015,033
518	Sucker, June ( <i>Chasmistes liorus</i> ) - Wherever found	E	\$64,000	\$1,251,079,033

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
519	Pearlymussel, birdwing ( <i>Lemiox rimosus</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$63,900	\$1,251,142,933
520	No Common Name ( <i>Nervilia jacksoniae</i> ) - Wherever found	T	\$63,736	\$1,251,206,669
521	No Common Name ( <i>Maesa walkeri</i> ) - Wherever found	T	\$63,399	\$1,251,270,068
522	Isopod, Madison Cave ( <i>Antrolana lira</i> ) - Wherever found	T	\$63,083	\$1,251,333,151
523	No Common Name ( <i>Geocarpon minimum</i> ) - Wherever found	T	\$62,998	\$1,251,396,149
524	Coral, Pillar ( <i>Dendrogyra cylindrus</i> ) - Wherever found	T	\$62,942	\$1,251,459,091
525	Crayfish, Guyandotte River ( <i>Cambarus veteranus</i> ) - Wherever found	E	\$62,648	\$1,251,521,739
526	Yellow-faced bee, anthracinan ( <i>Hylaeus anthracinus</i> ) - Wherever found	E	\$62,620	\$1,251,584,359
527	Madtom, Neosho ( <i>Noturus placidus</i> ) - Wherever found	T	\$62,598	\$1,251,646,957
528	Buckwheat, steamboat ( <i>Eriogonum ovalifolium</i> var. <i>williamsiae</i> ) - Wherever found	E	\$62,593	\$1,251,709,550
529	Butterfly, Palos Verdes blue ( <i>Glaucopsyche lygdamus palosverdesensis</i> ) - Wherever found	E	\$62,145	\$1,251,771,695
530	Duck, Laysan ( <i>Anas laysanensis</i> ) - Wherever found	E	\$62,073	\$1,251,833,768
531	Oxytheca, Cushenbury ( <i>Oxytheca parishii</i> var. <i>goodmaniana</i> ) - Wherever found	E	\$62,030	\$1,251,895,798
532	Damselfly, Rota blue ( <i>Ischnura luta</i> ) - Wherever found	E	\$61,836	\$1,251,957,634
533	Manzanita, Franciscan ( <i>Arctostaphylos franciscana</i> ) - Wherever found	E	\$61,500	\$1,252,019,134
534	Rabbit, riparian brush ( <i>Sylvilagus bachmani riparius</i> ) - Wherever found	E	\$61,300	\$1,252,080,434
535	Bird's-beak, salt marsh ( <i>Cordylanthus maritimus</i> ssp. <i>maritimus</i> ) - Wherever found	E	\$60,784	\$1,252,141,218
536	Bird's-beak, soft ( <i>Cordylanthus mollis</i> ssp. <i>mollis</i> ) - Wherever found	E	\$60,200	\$1,252,201,418
537	Catfish, Yaqui ( <i>Ictalurus pricei</i> ) - Wherever found	T	\$59,700	\$1,252,261,118
538	Pimpleback, orangefoot (pearlymussel) ( <i>Plethobasus cooperianus</i> ) - Wherever found	E	\$59,699	\$1,252,320,817
539	Bean, rayed ( <i>Villosa fabalis</i> ) - Wherever found	E	\$59,354	\$1,252,380,171
540	`Aiea ( <i>Nothoecstrum latifolium</i> ) - Wherever found	E	\$59,238	\$1,252,439,409
541	Riffleshell, tan ( <i>Epioblasma florentina walkeri</i> (=E. <i>walkeri</i> )) - Wherever found	E	\$58,783	\$1,252,498,192
542	Mountainbalm, Indian Knob ( <i>Eriodictyon altissimum</i> ) - Wherever found	E	\$58,600	\$1,252,556,792
543	Pearlymussel, littlewing ( <i>Pegias fabula</i> ) - Wherever found	E	\$58,364	\$1,252,615,156
544	Skink, Slevin's ( <i>Emoia slevini</i> ) - Wherever found	E	\$58,305	\$1,252,673,461
545	Dace, Foskett speckled ( <i>Rhinichthys osculus</i> ssp.) - Wherever found	T	\$58,000	\$1,252,731,461
545	No Common Name ( <i>Cordia bellonis</i> ) - Wherever found	E	\$58,000	\$1,252,789,461
546	Rocksnailed, round ( <i>Leptoxis ampla</i> ) - Wherever found	T	\$57,097	\$1,252,846,558
547	Mouse, Key Largo cotton ( <i>Peromyscus gossypinus allapaticola</i> ) - Wherever found	E	\$57,061	\$1,252,903,619
548	Beetle, American burying ( <i>Nicrophorus americanus</i> ) - Ex Pop, SW Missouri	EXPN	\$56,700	\$1,252,960,319
549	Ko`oko`olau ( <i>Bidens micrantha</i> ssp. <i>kalealaha</i> ) - Wherever found	E	\$56,442	\$1,253,016,761

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
550	Combshell, Cumberlandian ( <i>Epioblasma brevidens</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$55,917	\$1,253,072,678
551	Spiraea, Virginia ( <i>Spiraea virginiana</i> ) - Wherever found	T	\$55,908	\$1,253,128,586
552	Sturgeon, Shovelnose ( <i>Scaphirhynchus platyrhynchus</i> ) - Wherever found	SAT	\$55,864	\$1,253,184,450
553	Mouse, St. Andrew beach ( <i>Peromyscus polionotus peninsularis</i> ) - Wherever found	E	\$55,660	\$1,253,240,110
554	Snails, Oahu tree ( <i>Achatinella</i> spp.) - Wherever found	E*	\$55,191	\$1,253,295,301
555	Bean, Cumberland (pearlymussel) ( <i>Villosa trabalis</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$53,943	\$1,253,349,244
556	Brodiaea, Chinese Camp ( <i>Brodiaea pallida</i> ) - Wherever found	T	\$53,700	\$1,253,402,944
557	Butterfly, Florida leafwing ( <i>Anaea troglodyta floridalis</i> ) - Wherever found	E	\$53,500	\$1,253,456,444
558	Pupfish, Warm Springs ( <i>Cyprinodon nevadensis pectoralis</i> ) - Wherever found	E	\$53,479	\$1,253,509,923
559	Skyrocket, Pagosa ( <i>Ipomopsis polyantha</i> ) - Wherever found	E	\$53,165	\$1,253,563,088
560	Milk-vetch, Jesup's ( <i>Astragalus robbinsii</i> var. <i>jesupi</i> ) - Wherever found	E	\$53,070	\$1,253,616,158
561	Milk-vetch, Applegate's ( <i>Astragalus applegatei</i> ) - Wherever found	E	\$53,010	\$1,253,669,168
562	Bladderpod, Missouri ( <i>Physaria filiformis</i> ) - Wherever found	T	\$52,142	\$1,253,721,310
563	Mallow, Kern ( <i>Eremalche kernensis</i> ) - Wherever found	E	\$52,100	\$1,253,773,410
564	Wild-buckwheat, gypsum ( <i>Eriogonum gypsophilum</i> ) - Wherever found	T	\$52,000	\$1,253,825,410
564	Milk-vetch, Fish Slough ( <i>Astragalus lentiginosus</i> var. <i>piscinensis</i> ) - Wherever found	T	\$52,000	\$1,253,877,410
564	Dawn-flower, Texas prairie ( <i>Hymenoxys texana</i> ) - Wherever found	E	\$52,000	\$1,253,929,410
565	No Common Name ( <i>Phyllanthus saffordii</i> ) - Wherever found	E	\$51,903	\$1,253,981,313
566	Naupaka, dwarf ( <i>Scaevola coriacea</i> ) - Wherever found	E	\$51,230	\$1,254,032,543
567	Rattleweed, hairy ( <i>Baptisia arachnifera</i> ) - Wherever found	E	\$51,000	\$1,254,083,543
568	Goldenrod, Houghton's ( <i>Solidago houghtonii</i> ) - Wherever found	T	\$50,488	\$1,254,134,031
569	No Common Name ( <i>Cyrtandra nanawaleensis</i> ) - Wherever found	E	\$50,360	\$1,254,184,391
570	Alani ( <i>Melicope ovalis</i> ) - Wherever found	E	\$50,036	\$1,254,234,427
571	Tiger beetle, Ohlone ( <i>Cicindela ohlone</i> ) - Wherever found	E	\$49,890	\$1,254,284,317
572	Squirrel, Carolina northern flying ( <i>Glaucomys sabrinus coloratus</i> ) - Wherever found	E	\$49,624	\$1,254,333,941
573	Haha ( <i>Cyanea copelandii</i> ssp. <i>haleakalaensis</i> ) - Wherever found	E	\$49,537	\$1,254,383,478
574	Pigtoe, rough ( <i>Pleurobema plenum</i> ) - Wherever found	E	\$49,518	\$1,254,432,996
575	Kuahiwi laukahi ( <i>Plantago princeps</i> ) - Wherever found	E	\$49,505	\$1,254,482,501
576	Orchid, white fringeless ( <i>Platanthera integrilabia</i> ) -	T	\$49,156	\$1,254,531,657
577	Shrimp, Alabama cave ( <i>Palaemonias alabamae</i> ) - Wherever found	E	\$49,000	\$1,254,580,657
578	Niterwort, Amargosa ( <i>Nitrophila mohavensis</i> ) - Wherever found	E	\$48,600	\$1,254,629,257

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
579	`Ohe ( <i>Joinvillea ascendens ascendens</i> ) - Wherever found	E	\$48,281	\$1,254,677,538
580	Ring pink (mussel) ( <i>Obovaria retusa</i> ) - Wherever found	E	\$48,180	\$1,254,725,718
581	Fairy shrimp, longhorn ( <i>Branchinecta longiantenna</i> ) - Wherever found	E	\$47,815	\$1,254,773,533
582	Spider, spruce-fir moss ( <i>Microhexura montivaga</i> ) - Wherever found	E	\$47,765	\$1,254,821,298
583	Vole, Florida salt marsh ( <i>Microtus pennsylvanicus dukecampbelli</i> ) - Wherever found	E	\$47,500	\$1,254,868,798
583	Trout, Apache ( <i>Oncorhynchus apache</i> ) - Wherever found	T	\$47,500	\$1,254,916,298
584	`Oha wai ( <i>Clermontia samuelii</i> ) - Wherever found	E	\$47,452	\$1,254,963,750
585	Springfish, Hiko White River ( <i>Crenichthys baileyi grandis</i> ) - Wherever found	E	\$47,181	\$1,255,010,931
586	Haiwale ( <i>Cyrtandra ferripilosa</i> ) - Wherever found	E	\$47,080	\$1,255,058,011
587	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - Greater Yellowstone Ecosystem DPS	UR	\$47,000	\$1,255,105,011
587	Snail, Iowa Pleistocene ( <i>Discus macclintocki</i> ) - Wherever found	E	\$47,000	\$1,255,152,011
588	Pigtoe, narrow ( <i>Fusconaia escambia</i> ) - Wherever found	T	\$46,989	\$1,255,199,000
589	Mapleleaf, winged ( <i>Quadrula fragosa</i> ) - Wherever found, except where listed as an experimental population	E	\$46,635	\$1,255,245,635
590	Warea, wide-leaf ( <i>Warea amplexifolia</i> ) - Wherever found	E	\$46,500	\$1,255,292,135
591	No Common Name ( <i>Sanicula sandwicensis</i> ) - Wherever found	E	\$46,332	\$1,255,338,467
592	Totoaba (seatrout or weakfish) ( <i>Cynoscion macdonaldi</i> ) - Wherever found	E	\$46,200	\$1,255,384,667
593	Cactus, Mesa Verde ( <i>Sclerocactus mesae-verdae</i> ) - Wherever found	T	\$45,800	\$1,255,430,467
594	Butterfly, Behren's silverspot ( <i>Speyeria zerene behrensii</i> ) - Wherever found	E	\$45,539	\$1,255,476,006
595	Geranium, Hawaiian red-flowered ( <i>Geranium arboreum</i> ) - Wherever found	E	\$45,386	\$1,255,521,392
596	Ipomopsis, Holy Ghost ( <i>Ipomopsis sancti-spiritus</i> ) - Wherever found	E	\$45,000	\$1,255,566,392
597	Beaked-rush, Knieskern's ( <i>Rhynchospora knieskernii</i> ) - Wherever found	T	\$44,861	\$1,255,611,253
598	Haha ( <i>Cyanea asplenifolia</i> ) - Wherever found	E	\$44,846	\$1,255,656,099
599	Bladderpod, Short's ( <i>Physaria globosa</i> ) - Wherever found	E	\$44,643	\$1,255,700,742
600	No Common Name ( <i>Eugenia bryanii</i> ) - Wherever found	E	\$44,226	\$1,255,744,968
601	Springfish, White River ( <i>Crenichthys baileyi baileyi</i> ) - Wherever found	E	\$44,116	\$1,255,789,084
602	Sandshell, Southern ( <i>Hamiota australis</i> ) - Wherever found	T	\$43,499	\$1,255,832,583
603	Potentilla, Hickman's ( <i>Potentilla hickmanii</i> ) - Wherever found	E	\$43,400	\$1,255,875,983
604	Ko`oko`olau ( <i>Bidens micrantha</i> ssp. <i>ctenophylla</i> ) - Wherever found	E	\$43,370	\$1,255,919,353
605	Fern, American hart's-tongue ( <i>Asplenium scolopendrium</i> var. <i>americanum</i> ) - Wherever found	T	\$43,225	\$1,255,962,578
606	Bat, little Mariana fruit ( <i>Pteropus tokudae</i> ) - Wherever found	E	\$42,842	\$1,256,005,420
607	Heelsplitter, Inflated ( <i>Potamilus inflatus</i> ) - Wherever found	T	\$42,834	\$1,256,048,254
608	Ko`oko`olau ( <i>Bidens campylotheca</i> ssp. <i>waihoiensis</i> ) - Wherever found	E	\$42,780	\$1,256,091,034

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
608	Ko'oko'olau ( <i>Bidens campylotheca</i> ssp. <i>pentamera</i> ) - Wherever found	E	\$42,780	\$1,256,133,814
609	Polygala, Lewton's ( <i>Polygala lewtonii</i> ) - Wherever found	E	\$42,626	\$1,256,176,440
610	Pocketbook, shinyrayed ( <i>Lampsilis subangulata</i> ) - Wherever found	E	\$42,500	\$1,256,218,940
610	Thornmint, San Diego ( <i>Acanthomintha ilicifolia</i> ) - Wherever found	T	\$42,500	\$1,256,261,440
611	Pigtoe, fuzzy ( <i>Pleurobema strodeanum</i> ) - Wherever found	T	\$42,499	\$1,256,303,939
612	Spinedace, Little Colorado ( <i>Lepidomeda vittata</i> ) - Wherever found	T	\$42,075	\$1,256,346,014
613	Butterfly, El Segundo blue ( <i>Euphilotes battoides allyni</i> ) - Wherever found	E	\$42,042	\$1,256,388,056
614	Butterfly, Miami Blue ( <i>Cyclargus</i> (=Hemiargus) <i>thomasi bethunebakeri</i> ) - Wherever found	E	\$41,600	\$1,256,429,656
615	Cactus, Winkler ( <i>Pediocactus winkleri</i> ) - Wherever found	T	\$41,500	\$1,256,471,156
616	Kangaroo rat, Fresno ( <i>Dipodomys nitratoideis exilis</i> ) - Wherever found	E	\$41,438	\$1,256,512,594
617	Rosemary, Apalachicola ( <i>Conradina glabra</i> ) - Wherever found	E	\$41,400	\$1,256,553,994
618	Chub, Owens Tui ( <i>Gila bicolor</i> ssp. <i>snyderi</i> ) - Wherever found	E	\$41,024	\$1,256,595,018
619	Butterfly, Schaus swallowtail ( <i>Heraclides aristodemus ponceanus</i> ) - Wherever found	E	\$41,004	\$1,256,636,022
620	Blindcat, Mexican (catfish) ( <i>Prietella phreatophila</i> ) - Wherever found	E	\$41,000	\$1,256,677,022
621	Boa, Puerto Rican ( <i>Epicrates inornatus</i> ) - Wherever found	E	\$40,913	\$1,256,717,935
622	Wolf, Mexican ( <i>Canis lupus baileyi</i> ) - Wherever found, except where listed as an experimental population	E	\$40,230	\$1,256,758,165
623	Moorhen, Mariana common ( <i>Gallinula chloropus guami</i> ) - Wherever found	E	\$40,222	\$1,256,798,387
624	Madtom, yellowfin ( <i>Noturus flavipinnis</i> ) - Wherever found, except where listed as an experimental population	T	\$40,157	\$1,256,838,544
624	Kidneyshell, southern ( <i>Ptychobranhus jonesi</i> ) - Wherever found	E	\$40,157	\$1,256,878,701
625	Buckwheat, scrub ( <i>Eriogonum longifolium</i> var. <i>gnaphalifolium</i> ) - Wherever found	T	\$39,926	\$1,256,918,627
626	Bat, Mexican long-nosed ( <i>Leptonycteris nivalis</i> ) - Wherever found	E	\$39,876	\$1,256,958,503
627	Gourd, Okeechobee ( <i>Cucurbita okeechobeensis</i> ssp. <i>okeechobeensis</i> ) - Wherever found	E	\$39,700	\$1,256,998,203
628	Shiner, sharpnose ( <i>Notropis oxyrhynchus</i> ) - Wherever found	E	\$39,665	\$1,257,037,868
628	Shiner, smalleye ( <i>Notropis buccula</i> ) - Wherever found	E	\$39,665	\$1,257,077,533
629	Bean, Choctaw ( <i>Villosa choctawensis</i> ) - Wherever found	E	\$39,499	\$1,257,117,032
630	Holei ( <i>Ochrosia haleakalae</i> ) - Wherever found	E	\$39,489	\$1,257,156,521
631	Trout, Little Kern golden ( <i>Oncorhynchus aguabonita whitei</i> ) - Wherever found	T	\$39,023	\$1,257,195,544
632	Moccasinshell, Gulf ( <i>Medionidus penicillatus</i> ) - Wherever found	E	\$39,000	\$1,257,234,544
632	No Common Name ( <i>Lepanthes eltoroensis</i> ) - Wherever found	E	\$39,000	\$1,257,273,544
633	Pinesnake, black ( <i>Pituophis melanoleucus lodingi</i> ) - Wherever found	T	\$38,956	\$1,257,312,500
634	Snail, Langford's tree ( <i>Partula langfordi</i> ) - Wherever found	E	\$38,938	\$1,257,351,438

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
635	Uhi uhi ( <i>Mezoneuron kawaiense</i> ) - Wherever found	E	\$38,826	\$1,257,390,264
636	Bat, Pacific sheath-tailed ( <i>Emballonura semicaudata rotensis</i> ) - Wherever found	E	\$38,684	\$1,257,428,948
637	Cladonia, Florida perforate ( <i>Cladonia perforata</i> ) - Wherever found	E	\$38,400	\$1,257,467,348
638	Pigeon wings ( <i>Clitoria fragrans</i> ) - Wherever found	T	\$38,365	\$1,257,505,713
639	Sturgeon, Alabama ( <i>Scaphirhynchus suttkusi</i> ) - Wherever found	E	\$38,022	\$1,257,543,735
640	Ihi`ihi ( <i>Marsilea villosa</i> ) - Wherever found	E	\$38,012	\$1,257,581,747
641	Dudleya, Verity's ( <i>Dudleya verityi</i> ) - Wherever found	T	\$37,400	\$1,257,619,147
642	Thistle, Suisun ( <i>Cirsium hydrophilum</i> var. <i>hydrophilum</i> ) - Wherever found	E	\$37,300	\$1,257,656,447
643	Milk-vetch, Lane Mountain ( <i>Astragalus jaegerianus</i> ) - Wherever found	E	\$37,200	\$1,257,693,647
644	Dropwort, Canby's ( <i>Oxypolis canbyi</i> ) - Wherever found	E	\$37,154	\$1,257,730,801
645	Snail, noonday ( <i>Mesodon clarki nantahala</i> ) - Wherever found	T	\$37,000	\$1,257,767,801
645	Aster, Florida golden ( <i>Chrysopsis floridana</i> ) - Wherever found	E	\$37,000	\$1,257,804,801
646	White-eye, Rota bridled ( <i>Zosterops rotensis</i> ) - Wherever found	E	\$36,706	\$1,257,841,507
647	Butterweed, Layne's ( <i>Senecio layneae</i> ) - Wherever found	T	\$36,550	\$1,257,878,057
648	Pinkroot, gentian ( <i>Spigelia gentianoides</i> ) - Wherever found	E	\$36,519	\$1,257,914,576
649	Kangaroo rat, Morro Bay ( <i>Dipodomys heermanni morroensis</i> ) - Wherever found	E	\$36,000	\$1,257,950,576
649	Sucker, Zuni bluehead ( <i>Catostomus discobolus yarrowi</i> ) - Wherever found	E	\$36,000	\$1,257,986,576
650	Monkey-flower, Michigan ( <i>Mimulus michiganensis</i> ) - Wherever found	E	\$35,865	\$1,258,022,441
651	Cycladenia, Jones ( <i>Cycladenia humilis</i> var. <i>jonesii</i> ) - Wherever found	T	\$35,830	\$1,258,058,271
652	Plum, scrub ( <i>Prunus geniculata</i> ) - Wherever found	E	\$35,700	\$1,258,093,971
653	Manzanita, Ione ( <i>Arctostaphylos myrtifolia</i> ) - Wherever found	T	\$35,603	\$1,258,129,574
654	Kula wahine noho ( <i>Isodendrion pyrifolium</i> ) - Wherever found	E	\$35,545	\$1,258,165,119
655	Lead-plant, Crenulate ( <i>Amorpha crenulata</i> ) - Wherever found	E	\$35,200	\$1,258,200,319
656	Crayfish, Hell Creek Cave ( <i>Cambarus zophonastes</i> ) - Wherever found	E	\$35,000	\$1,258,235,319
656	Meadowfoam, Sebastopol ( <i>Limnanthes vinculans</i> ) - Wherever found	E	\$35,000	\$1,258,270,319
657	No Common Name ( <i>Nesogenes rotensis</i> ) - Wherever found	E	\$34,671	\$1,258,304,990
658	Paudedo ( <i>Hedyotis megalantha</i> ) - Wherever found	E	\$34,368	\$1,258,339,358
659	Iguana, Cuban ground ( <i>Cyclura nubila nubila</i> ) - ( <i>Cyclura nubila nubila</i> )	T	\$33,800	\$1,258,373,158
660	Crane, whooping ( <i>Grus americana</i> ) - U.S.A (Southwestern Louisiana)	EXPN	\$33,720	\$1,258,406,878
661	Shrimp, Kentucky cave ( <i>Palaemonias ganteri</i> ) - Wherever found	E	\$33,500	\$1,258,440,378
662	Sunshine, Sonoma ( <i>Blennosperma bakeri</i> ) - Wherever found	E	\$33,400	\$1,258,473,778
663	Skullcap, large-flowered ( <i>Scutellaria montana</i> ) - Wherever found	T	\$33,276	\$1,258,507,054

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
664	Tortoise, desert ( <i>Gopherus agassizii</i> ) - AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	SAT	\$33,200	\$1,258,540,254
665	Bird's beak, palmate-bracted ( <i>Cordylanthus palmatus</i> ) - Wherever found	E	\$32,992	\$1,258,573,246
666	Darter, Cumberland ( <i>Etheostoma susanae</i> ) - Wherever found	E	\$32,748	\$1,258,605,994
667	Mesa-mint, San Diego ( <i>Pogogyne abramsii</i> ) - Wherever found	E	\$32,590	\$1,258,638,584
668	Moccasinshell, Ochlockonee ( <i>Medionidus simpsonianus</i> ) - Wherever found	E	\$32,500	\$1,258,671,084
668	Ayenia, Texas ( <i>Ayenia limitaris</i> ) - Wherever found	E	\$32,500	\$1,258,703,584
669	Barberry, island ( <i>Berberis pinnata</i> ssp. <i>insularis</i> ) - Wherever found	E	\$32,499	\$1,258,736,083
670	Thelypody, Howell's spectacular ( <i>Thelypodium howellii</i> ssp. <i>spectabilis</i> ) - Wherever found	T	\$32,310	\$1,258,768,393
671	Pentachaeta, Lyon's ( <i>Pentachaeta lyonii</i> ) - Wherever found	E	\$32,300	\$1,258,800,693
671	Cactus, Knowlton's ( <i>Pediocactus knowltonii</i> ) - Wherever found	E	\$32,300	\$1,258,832,993
672	Beauty, Harper's ( <i>Harperocallis flava</i> ) - Wherever found	E	\$32,000	\$1,258,864,993
672	Goldfields, Burke's ( <i>Lasthenia burkei</i> ) - Wherever found	E	\$32,000	\$1,258,896,993
673	Lily, Western ( <i>Lilium occidentale</i> ) - Wherever found	E	\$31,810	\$1,258,928,803
674	Beargrass, Britton's ( <i>Nolina brittoniana</i> ) - Wherever found	E	\$31,800	\$1,258,960,603
675	Crane, whooping ( <i>Grus americana</i> ) - U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	EXPN	\$31,688	\$1,258,992,291
676	Pennyroyal, Todsens' ( <i>Hedeoma todsenii</i> ) - Wherever found	E	\$31,500	\$1,259,023,791
677	No Common Name ( <i>Pavona diffluens</i> ) - Wherever found	T	\$31,467	\$1,259,055,258
678	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - Wherever found	RT	\$31,300	\$1,259,086,558
679	Bonamia, Florida ( <i>Bonamia grandiflora</i> ) - Wherever found	T	\$31,200	\$1,259,117,758
680	Evening-primrose, San Benito ( <i>Camissonia benitensis</i> ) - Wherever found	T	\$31,000	\$1,259,148,758
681	Ischaemum, Hilo ( <i>Ischaemum byrone</i> ) - Wherever found	E	\$30,983	\$1,259,179,741
682	Heartleaf, dwarf-flowered ( <i>Hexastylis naniflora</i> ) - Wherever found	T	\$30,968	\$1,259,210,709
683	Bear-poppy, Dwarf ( <i>Arctomecon humilis</i> ) - Wherever found	E	\$30,903	\$1,259,241,612
684	Heather, mountain golden ( <i>Hudsonia montana</i> ) - Wherever found	T	\$30,900	\$1,259,272,512
685	Madtom, pygmy ( <i>Noturus stanauli</i> ) - Wherever found	E	\$30,811	\$1,259,303,323
686	Haha ( <i>Cyanea maritae</i> ) - Wherever found	E	\$30,779	\$1,259,334,102
687	Turtle, ringed map ( <i>Graptemys oculifera</i> ) - Wherever found	T	\$30,569	\$1,259,364,671
688	Milkweed, Welsh's ( <i>Asclepias welshii</i> ) - Wherever found	T	\$30,500	\$1,259,395,171
689	Millerbird, Nihoa (old world warbler) ( <i>Acrocephalus familiaris kingi</i> ) - Wherever found	E	\$30,135	\$1,259,425,306
690	Woodpecker, ivory-billed ( <i>Campephilus principalis</i> ) - Wherever found	E	\$30,029	\$1,259,455,335
691	Cavesnail, Tumbling Creek ( <i>Antrobia culveri</i> ) - Wherever found	E	\$30,000	\$1,259,485,335



Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
692	Snail, Stock Island tree ( <i>Orthalicus reses</i> (not incl. <i>nesodryas</i> )) - Wherever found	T	\$29,984	\$1,259,515,319
693	Snail, Lanai tree ( <i>Partulina semicarinata</i> ) - Wherever found	E	\$29,956	\$1,259,545,275
693	Snail, Lanai tree ( <i>Partulina variabilis</i> ) - Wherever found	E	\$29,956	\$1,259,575,231
694	Water-umbel, Huachuca ( <i>Lilaeopsis schaffneriana</i> var. <i>recurva</i> ) - Wherever found	E	\$29,800	\$1,259,605,031
695	Cactus, Arizona hedgehog ( <i>Echinocereus triglochidiatus</i> var. <i>arizonicus</i> ) - Wherever found	E	\$29,704	\$1,259,634,735
696	Sunflower, Pecos (=puzzle, =paradox) ( <i>Helianthus paradoxus</i> ) - Wherever found	T	\$29,650	\$1,259,664,385
697	No Common Name ( <i>Spermolepis hawaiiensis</i> ) - Wherever found	E	\$29,628	\$1,259,694,013
698	Tarplant, Santa Cruz ( <i>Holocarpha macradenia</i> ) - Wherever found	T	\$29,300	\$1,259,723,313
699	Ladies-tresses, Navasota ( <i>Spiranthes parksii</i> ) - Wherever found	E	\$29,265	\$1,259,752,578
700	Butterfly, San Bruno elfin ( <i>Callophrys mossii bayensis</i> ) - Wherever found	E	\$29,214	\$1,259,781,792
701	Snakeroot ( <i>Eryngium cuneifolium</i> ) - Wherever found	E	\$29,200	\$1,259,810,992
702	Joint-vetch, Sensitive ( <i>Aeschynomene virginica</i> ) - Wherever found	T	\$28,938	\$1,259,839,930
703	Penny-cress, Kneeland Prairie ( <i>Thlaspi californicum</i> ) - Wherever found	E	\$28,900	\$1,259,868,830
703	Milk-vetch, Holmgren ( <i>Astragalus holmgreniorum</i> ) - Wherever found	E	\$28,900	\$1,259,897,730
704	Purple Cat's paw (=Purple Cat's paw pearlymussel) ( <i>Epioblasma obliquata obliquata</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$28,820	\$1,259,926,550
705	Milk-vetch, Mancos ( <i>Astragalus humillimus</i> ) - Wherever found	E	\$28,600	\$1,259,955,150
706	Thrush, small Kauai (=puaiohi) ( <i>Myadestes palmeri</i> ) - Wherever found	E	\$28,501	\$1,259,983,651
707	Aplokating-palaoan ( <i>Psychotria malaspiniae</i> ) - Wherever found	E	\$28,470	\$1,260,012,121
708	Buckwheat, Ione (incl. Irish Hill) ( <i>Eriogonum apricum</i> (incl. var. <i>prostratum</i> )) - Wherever found	E	\$28,400	\$1,260,040,521
709	Sandlace ( <i>Polygonella myriophylla</i> ) - Wherever found	E	\$28,200	\$1,260,068,721
710	Mint, scrub ( <i>Dicerandra frutescens</i> ) - Wherever found	E	\$28,100	\$1,260,096,821
711	Fern, Aleutian shield ( <i>Polystichum aleuticum</i> ) - Wherever found	E	\$28,000	\$1,260,124,821
712	Grass, Colusa ( <i>Neostapfia colusana</i> ) - Wherever found	T	\$27,790	\$1,260,152,611
713	Hornshell, Texas ( <i>Popenaias popeii</i> ) - Wherever found	E	\$27,600	\$1,260,180,211
714	Yellow-faced bee, assimilans ( <i>Hylaeus assimilans</i> ) - Wherever found	E	\$27,592	\$1,260,207,803
715	Darter, Etowah ( <i>Etheostoma etowahae</i> ) - Wherever found	E	\$27,330	\$1,260,235,133
716	Cactus, Siler pincushion ( <i>Pediocactus</i> (=Echinocactus,=Utahia) <i>sileri</i> ) - Wherever found	T	\$27,200	\$1,260,262,333
717	Rail, Guam ( <i>Rallus owstoni</i> ) - Rota	EXPN	\$27,125	\$1,260,289,458
718	Warbler, nightingale reed (old world warbler) ( <i>Acrocephalus luscinia</i> ) - Wherever found	E	\$27,107	\$1,260,316,565
719	Makou ( <i>Ranunculus mauiensis</i> ) - Wherever found	E	\$26,946	\$1,260,343,511
720	Flannelbush, Mexican ( <i>Fremontodendron mexicanum</i> ) - Wherever found	E	\$26,900	\$1,260,370,411

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
721	Ko'oko'olau ( <i>Bidens amplexans</i> ) - Wherever found	E	\$26,896	\$1,260,397,307
722	Shrimp, California freshwater ( <i>Syncaris pacifica</i> ) - Wherever found	E	\$26,712	\$1,260,424,019
723	Centaury, spring-loving ( <i>Centaureum namophilum</i> ) - Wherever found	T	\$26,600	\$1,260,450,619
723	Gumplant, Ash Meadows ( <i>Grindelia fraxinipratensis</i> ) - Wherever found	T	\$26,600	\$1,260,477,219
724	Boa, Virgin Islands tree ( <i>Chilabothrus granti</i> ) - Wherever found	E	\$26,556	\$1,260,503,775
725	Ambrosia, south Texas ( <i>Ambrosia cheiranthifolia</i> ) - Wherever found	E	\$26,500	\$1,260,530,275
726	No Common Name ( <i>Osmoxylon mariannense</i> ) - Wherever found	E	\$26,488	\$1,260,556,763
727	Wireweed ( <i>Polygonella basiramia</i> ) - Wherever found	E	\$26,400	\$1,260,583,163
728	Sandwort, Cumberland ( <i>Arenaria cumberlandensis</i> ) - Wherever found	E	\$26,351	\$1,260,609,514
729	No Common Name ( <i>Silene lanceolata</i> ) - Wherever found	E	\$26,091	\$1,260,635,605
730	Pitcher-plant, mountain sweet ( <i>Sarracenia rubra</i> ssp. <i>jonesii</i> ) - Wherever found	E	\$26,067	\$1,260,661,672
731	No Common Name ( <i>Tetramolopium arenarium</i> ) - Wherever found	E	\$25,855	\$1,260,687,527
732	No Common Name ( <i>Neraudia ovata</i> ) - Wherever found	E	\$25,657	\$1,260,713,184
733	Haha ( <i>Cyanea hamatiflora</i> ssp. <i>hamatiflora</i> ) - Wherever found	E	\$25,537	\$1,260,738,721
734	`Aiea ( <i>Nothocestrum breviflorum</i> ) - Wherever found	E	\$25,509	\$1,260,764,230
735	Riversnail, Anthony's ( <i>Athearnia anthonyi</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$25,500	\$1,260,789,730
736	Hala pepe ( <i>Pleomele hawaiiensis</i> ) - Wherever found	E	\$25,487	\$1,260,815,217
737	Cactus, Bakersfield ( <i>Opuntia treleasei</i> ) - Wherever found	E	\$25,400	\$1,260,840,617
738	Creeper, Molokai ( <i>Paroreomyza flammea</i> ) - Wherever found	E	\$25,335	\$1,260,865,952
739	Otter, southern sea ( <i>Enhydra lutris nereis</i> ) - Wherever found	NS	\$25,000	\$1,260,890,952
740	Rhododendron, Chapman ( <i>Rhododendron chapmanii</i> ) - Wherever found	E	\$24,950	\$1,260,915,902
741	Owl's-clover, fleshy ( <i>Castilleja campestris</i> ssp. <i>succulenta</i> ) - Wherever found	T	\$24,930	\$1,260,940,832
742	Milk-vetch, Shivwits ( <i>Astragalus ampullarioides</i> ) - Wherever found	E	\$24,925	\$1,260,965,757
743	Cactus, Fickeisen plains ( <i>Pediocactus peeblesianus fickeiseniae</i> ) - Wherever found	E	\$24,800	\$1,260,990,557
744	Tree snail, Newcomb's ( <i>Newcombia cumingi</i> ) - Wherever found	E	\$24,659	\$1,261,015,216
745	No Common Name ( <i>Asplenium peruvianum</i> var. <i>insulare</i> ) - Wherever found	E	\$24,547	\$1,261,039,763
746	Gerardia, sandplain ( <i>Agalinis acuta</i> ) - Wherever found	E	\$24,520	\$1,261,064,283
747	Squirrel, Mount Graham red ( <i>Tamiasciurus hudsonicus grahamensis</i> ) - Wherever found	E	\$24,500	\$1,261,088,783
747	Clubshell ( <i>Pleurobema clava</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$24,500	\$1,261,113,283
747	Manioc, Walker's ( <i>Manihot walkerae</i> ) - Wherever found	E	\$24,500	\$1,261,137,783
748	Mouse, Anastasia Island beach ( <i>Peromyscus polionotus phasma</i> ) - Wherever found	E	\$24,220	\$1,261,162,003

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
749	Mallow, Gierisch ( <i>Sphaeralcea gierischii</i> ) - Wherever found	E	\$24,200	\$1,261,186,203
750	Spinymussel, Tar River ( <i>Elliptio steinstansana</i> ) - Wherever found	E	\$24,025	\$1,261,210,228
751	Higuero de sierra ( <i>Crescentia portoricensis</i> ) - Wherever found	E	\$24,000	\$1,261,234,228
752	No Common Name ( <i>Microlepidia strigosa</i> var. <i>mauiensis</i> ) - Wherever found	E	\$23,888	\$1,261,258,116
753	Orcutt grass, slender ( <i>Orcuttia tenuis</i> ) - Wherever found	T	\$23,600	\$1,261,281,716
754	Poppy, Sacramento prickly ( <i>Argemone pleiacantha</i> ssp. <i>pinnatisecta</i> ) - Wherever found	E	\$23,550	\$1,261,305,266
755	Shiner, Cape Fear ( <i>Notropis mekistocholas</i> ) - Wherever found	E	\$23,440	\$1,261,328,706
756	No Common Name ( <i>Platanthera holochila</i> ) - Wherever found	E	\$23,411	\$1,261,352,117
757	Naucorid, Ash Meadows ( <i>Ambrysus amargosus</i> ) - Wherever found	T	\$23,400	\$1,261,375,517
758	Rice rat, Silver ( <i>Oryzomys palustris natator</i> ) - lower FL Keys	E	\$23,303	\$1,261,398,820
759	Watercress, Gambel's ( <i>Rorippa gambellii</i> ) - Wherever found	E	\$23,300	\$1,261,422,120
760	Reedgrass, Maui ( <i>Calamagrostis expansa</i> ) - Wherever found	E	\$23,208	\$1,261,445,328
761	Thoroughwort, Cape Sable ( <i>Chromolaena frustrata</i> ) - Wherever found	E	\$23,200	\$1,261,468,528
762	Morning-glory, Stebbins' ( <i>Calystegia stebbinsii</i> ) - Wherever found	E	\$23,100	\$1,261,491,628
763	Lizard, St. Croix ground ( <i>Ameiva polops</i> ) - Wherever found	E	\$23,000	\$1,261,514,628
764	Orcutt grass, San Joaquin ( <i>Orcuttia inaequalis</i> ) - Wherever found	T	\$22,978	\$1,261,537,606
765	Nohoanu ( <i>Geranium multiflorum</i> ) - Wherever found	E	\$22,965	\$1,261,560,571
766	No Common Name ( <i>Wikstroemia villosa</i> ) - Wherever found	E	\$22,931	\$1,261,583,502
766	Nui, haha ( <i>Cyanea horrida</i> ) - Wherever found	E	\$22,931	\$1,261,606,433
766	No Common Name ( <i>Schiedea diffusa</i> subsp. <i>diffusa</i> ) - Wherever found	E	\$22,931	\$1,261,629,364
767	Damselfly, orangeblack Hawaiian ( <i>Megalagrion xanthomelas</i> ) - Wherever found	E	\$22,776	\$1,261,652,140
768	Bladderpod, Dudley Bluffs ( <i>Lesquerella congesta</i> ) - Wherever found	T	\$22,750	\$1,261,674,890
769	Haha ( <i>Cyanea kunthiana</i> ) - Wherever found	E	\$22,678	\$1,261,697,568
770	Elepaio, Oahu ( <i>Chasiempis ibidis</i> ) - Wherever found	E	\$22,655	\$1,261,720,223
771	Nukupuu, Maui ( <i>Hemignathus affinis</i> ) - Wherever found	E	\$22,357	\$1,261,742,580
772	Arrowhead, bunched ( <i>Sagittaria fasciculata</i> ) - Wherever found	E	\$22,100	\$1,261,764,680
773	Cactus, star ( <i>Astrophytum asterias</i> ) - Wherever found	E	\$22,000	\$1,261,786,680
774	Blue-star, Kearney's ( <i>Amsonia kearneyana</i> ) - Wherever found	E	\$21,900	\$1,261,808,580
775	Ebonysheal, round ( <i>Fusconaia rotulata</i> ) - Wherever found	E	\$21,348	\$1,261,829,928
776	Makou ( <i>Ranunculus hawaiiensis</i> ) - Wherever found	E	\$21,278	\$1,261,851,206
777	Pauoa ( <i>Ctenitis squamigera</i> ) - Wherever found	E	\$21,276	\$1,261,872,482
778	Darter, Kentucky arrow ( <i>Etheostoma spilotum</i> ) - Wherever found	T	\$21,182	\$1,261,893,664
779	Nohoanu ( <i>Geranium hanaense</i> ) - Wherever found	E	\$21,099	\$1,261,914,763

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
780	Pigtoe, tapered ( <i>Fusconaia burkei</i> ) - Wherever found	T	\$21,010	\$1,261,935,773
781	Aster, Ruth's golden ( <i>Pityopsis ruthii</i> ) - Wherever found	E	\$21,000	\$1,261,956,773
782	`Anunu ( <i>Sicyos albus</i> ) - Wherever found	E	\$20,980	\$1,261,977,753
783	Avens, spreading ( <i>Geum radiatum</i> ) - Wherever found	E	\$20,875	\$1,261,998,628
784	Woolly-star, Santa Ana River ( <i>Eriastrum densifolium</i> ssp. <i>sanctorum</i> ) - Wherever found	E	\$20,818	\$1,262,019,446
785	Cactus, San Rafael ( <i>Pediocactus despainii</i> ) - Wherever found	E	\$20,750	\$1,262,040,196
786	Elktoe, Cumberland ( <i>Alasmidonta atropurpurea</i> ) - Wherever found	E	\$20,618	\$1,262,060,814
787	Nanu ( <i>Gardenia remyi</i> ) - Wherever found	E	\$20,555	\$1,262,081,369
788	Clover, showy Indian ( <i>Trifolium amoenum</i> ) - Wherever found	E	\$20,500	\$1,262,101,869
789	Rock cress, Shale barren ( <i>Arabis serotina</i> ) - Wherever found	E	\$20,480	\$1,262,122,349
790	Rosemary, Cumberland ( <i>Conradina verticillata</i> ) - Wherever found	T	\$20,476	\$1,262,142,825
791	Fleabane, Zuni ( <i>Erigeron rhizomatus</i> ) - Wherever found	T	\$20,407	\$1,262,163,232
792	Orcutt grass, Sacramento ( <i>Orcuttia viscida</i> ) - Wherever found	E	\$20,400	\$1,262,183,632
792	Milk-vetch, Coachella Valley ( <i>Astragalus lentiginosus</i> var. <i>coachellae</i> ) - Wherever found	E	\$20,400	\$1,262,204,032
793	Gilia, Monterey ( <i>Gilia tenuiflora</i> ssp. <i>arenaria</i> ) - Wherever found	E	\$20,344	\$1,262,224,376
794	Clarkia, Springville ( <i>Clarkia springvillensis</i> ) - Wherever found	T	\$20,319	\$1,262,244,695
795	Phacelia, clay ( <i>Phacelia argillacea</i> ) - Wherever found	E	\$20,300	\$1,262,264,995
796	Kaulia ( <i>Colubrina oppositifolia</i> ) - Wherever found	E	\$20,294	\$1,262,285,289
797	Butterfly, Uncompahgre fritillary ( <i>Boloria acrocneuma</i> ) - Wherever found	E	\$20,200	\$1,262,305,489
797	Spineflower, Scotts Valley ( <i>Chorizanthe robusta</i> var. <i>hartwegii</i> ) - Wherever found	E	\$20,200	\$1,262,325,689
798	Darter, relict ( <i>Etheostoma chienense</i> ) - Wherever found	E	\$20,000	\$1,262,345,689
798	Beetle, Hungerford's crawling water ( <i>Brychius hungerfordi</i> ) - Wherever found	E	\$20,000	\$1,262,365,689
798	Orcutt grass, California ( <i>Orcuttia californica</i> ) - Wherever found	E	\$20,000	\$1,262,385,689
799	Crayfish, Nashville ( <i>Orconectes shoupi</i> ) - Wherever found	E	\$19,942	\$1,262,405,631
800	Hoawa ( <i>Pittosporum halophilum</i> ) - Wherever found	E	\$19,873	\$1,262,425,504
801	Sandalwood, Lanai (= `iliahi) ( <i>Santalum haleakalae</i> var. <i>lanaiense</i> ) - Wherever found	E	\$19,768	\$1,262,445,272
801	Alani ( <i>Melicope adscendens</i> ) - Wherever found	E	\$19,768	\$1,262,465,040
802	Wawae `iole ( <i>Huperzia mannii</i> ) - Wherever found	E	\$19,765	\$1,262,484,805
803	No Common Name ( <i>Silene hawaiiensis</i> ) - Wherever found	T	\$19,743	\$1,262,504,548
804	Spineflower, Ben Lomond ( <i>Chorizanthe pungens</i> var. <i>hartwegiana</i> ) - Wherever found	E	\$19,700	\$1,262,524,248
805	Wild buckwheat, clay-loving ( <i>Eriogonum pelinophilum</i> ) - Wherever found	E	\$19,650	\$1,262,543,898
806	Phacelia, DeBeque ( <i>Phacelia submutica</i> ) - Wherever found	T	\$19,640	\$1,262,563,538
807	Po`e ( <i>Portulaca sclerocarpa</i> ) - Wherever found	E	\$19,605	\$1,262,583,143

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
808	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - Cabinet-Yaak Recovery Zone Population	UR	\$19,500	\$1,262,602,643
808	Shark, Scalloped Hammerhead ( <i>Sphyrna lewini</i> ) - Wherever found	RT	\$19,500	\$1,262,622,143
809	Sunburst, San Joaquin adobe ( <i>Pseudobahia peirsonii</i> ) - Wherever found	T	\$19,416	\$1,262,641,559
810	Milk-vetch, Ventura Marsh ( <i>Astragalus pycnostachyus</i> var. <i>lanosissimus</i> ) - Wherever found	E	\$19,400	\$1,262,660,959
810	Mesa-mint, Otay ( <i>Pogogyne nudiuscula</i> ) - Wherever found	E	\$19,400	\$1,262,680,359
811	No Common Name ( <i>Vigna o-wahuensis</i> ) - Wherever found	E	\$19,381	\$1,262,699,740
812	Beetle, delta green ground ( <i>Elaphrus viridis</i> ) - Wherever found	T	\$19,161	\$1,262,718,901
813	Finch, Nihoa (honeycreeper) ( <i>Telespyza ultima</i> ) - Wherever found	E	\$19,133	\$1,262,738,034
814	Alani ( <i>Melicope knudsenii</i> ) - Wherever found	E	\$19,044	\$1,262,757,078
815	Akialoa, Kauai (honeycreeper) ( <i>Akialoa stejnegeri</i> ) - Wherever found	E	\$19,012	\$1,262,776,090
816	Iliau, dwarf ( <i>Wilkesia hobdyi</i> ) - Wherever found	E	\$18,965	\$1,262,795,055
817	No Common Name ( <i>Bonamia menziesii</i> ) - Wherever found	E	\$18,881	\$1,262,813,936
818	Ha`iwale ( <i>Cyrtandra filipes</i> ) - Wherever found	E	\$18,662	\$1,262,832,598
819	Prairie-clover, leafy ( <i>Dalea foliosa</i> ) - Wherever found	E	\$18,626	\$1,262,851,224
820	Mahoe ( <i>Alectryon macrococcus</i> ) - Wherever found	E	\$18,607	\$1,262,869,831
821	Sandwort, Marsh ( <i>Arenaria paludicola</i> ) - Wherever found	E	\$18,600	\$1,262,888,431
822	Ko`olua`ula ( <i>Abutilon menziesii</i> ) - Wherever found	E	\$18,596	\$1,262,907,027
823	Potato-bean, Prices ( <i>Apios priceana</i> ) -	T	\$18,558	\$1,262,925,585
824	Hoawa ( <i>Pittosporum hawaiiense</i> ) - Wherever found	E	\$18,520	\$1,262,944,105
825	Kupukupu makalii ( <i>Cyclosorus boydiae</i> ) - Wherever found	E	\$18,375	\$1,262,962,480
826	Amphipod, Kauai cave ( <i>Spelaeorchestia koloana</i> ) - Wherever found	E	\$18,360	\$1,262,980,840
827	Silversword, Mauna Loa (=Ka'u) ( <i>Argyroxiphium kauense</i> ) - Wherever found	E	\$18,356	\$1,262,999,196
828	Lupine, Nipomo Mesa ( <i>Lupinus nipomensis</i> ) - Wherever found	E	\$18,300	\$1,263,017,496
829	Aupaka ( <i>Isodendrion longifolium</i> ) - Wherever found	T	\$18,257	\$1,263,035,753
830	Blackbird, yellow-shouldered ( <i>Agelaius xanthomus</i> ) - Wherever found	E	\$18,250	\$1,263,054,003
831	Cactus, Lee pincushion ( <i>Coryphantha sneedii</i> var. <i>leei</i> ) - Wherever found	T	\$18,000	\$1,263,072,003
831	Sunflower, whorled ( <i>Helianthus verticillatus</i> ) -	E	\$18,000	\$1,263,090,003
832	Cactus, Acuna ( <i>Echinomastus erectocentrus</i> var. <i>acunensis</i> ) - Wherever found	E	\$17,908	\$1,263,107,911
833	Vervain, Red Hills ( <i>Verbena californica</i> ) - Wherever found	T	\$17,900	\$1,263,125,811
834	Nukupuu, Kauai ( <i>Hemignathus hanapepe</i> ) - Wherever found	E	\$17,828	\$1,263,143,639
835	Turtle, Plymouth Redbelly ( <i>Pseudemys rubriventris bangsi</i> ) - Wherever found	E	\$17,740	\$1,263,161,379
836	Finch, Laysan (honeycreeper) ( <i>Telespyza cantans</i> ) - Wherever found	E	\$17,713	\$1,263,179,092
837	Mehamehame ( <i>Flueggea neowawraea</i> ) - Wherever found	E	\$17,591	\$1,263,196,683

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
838	Moth, Kern primrose sphinx ( <i>Euproserpinus euterpe</i> ) - Wherever found	T	\$17,568	\$1,263,214,251
839	Spurge, telephus ( <i>Euphorbia telephoides</i> ) - Wherever found	T	\$17,500	\$1,263,231,751
840	Monkshood, northern wild ( <i>Aconitum noveboracense</i> ) - Wherever found	T	\$17,479	\$1,263,249,230
841	Bonytongue, Asian ( <i>Scleropages formosus</i> ) - Wherever found	E	\$17,325	\$1,263,266,555
842	Minnow, Devils River ( <i>Dionda diaboli</i> ) - Wherever found	T	\$17,305	\$1,263,283,860
843	Ceanothus, coyote ( <i>Ceanothus ferrisiae</i> ) - Wherever found	E	\$17,300	\$1,263,301,160
844	Rattlesnake, New Mexican ridge-nosed ( <i>Crotalus willardi obscurus</i> ) - Wherever found	T	\$17,250	\$1,263,318,410
845	Dudleya, Santa Monica Mountains ( <i>Dudleya cymosa</i> ssp. <i>ovatifolia</i> ) - Wherever found	T	\$17,200	\$1,263,335,610
846	Cactus, Uinta Basin hookless ( <i>Sclerocactus wetlandicus</i> ) - Wherever found	T	\$17,092	\$1,263,352,702
847	Manzanita, Santa Rosa Island ( <i>Arctostaphylos confertiflora</i> ) - Wherever found	E	\$17,068	\$1,263,369,770
848	Mussel, oyster ( <i>Epioblasma capsaeformis</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$17,057	\$1,263,386,827
849	Dace, Clover Valley speckled ( <i>Rhinichthys osculus oligoporus</i> ) - Wherever found	E	\$17,039	\$1,263,403,866
850	Woodrat, riparian (=San Joaquin Valley) ( <i>Neotoma fuscipes riparia</i> ) - Wherever found	E	\$17,000	\$1,263,420,866
850	Phlox, Texas trailing ( <i>Phlox nivalis</i> ssp. <i>texensis</i> ) - Wherever found	E	\$17,000	\$1,263,437,866
850	Cliffrose, Arizona ( <i>Purshia</i> (=Cowania) <i>subintegra</i> ) - Wherever found	E	\$17,000	\$1,263,454,866
851	No Common Name ( <i>Eugenia woodburyana</i> ) - Wherever found	E	\$16,800	\$1,263,471,666
852	Shrimp, anchialine pool ( <i>Procaris hawaiana</i> ) - Wherever found	E	\$16,674	\$1,263,488,340
853	A`e ( <i>Zanthoxylum hawaiiense</i> ) - Wherever found	E	\$16,593	\$1,263,504,933
854	Meadowrue, Cooley's ( <i>Thalictrum cooleyi</i> ) - Wherever found	E	\$16,540	\$1,263,521,473
855	`Oha wai ( <i>Clermontia oblongifolia</i> ssp. <i>mauiensis</i> ) - Wherever found	E	\$16,538	\$1,263,538,011
856	Elimia, lacy (snail) ( <i>Elimia crenatella</i> ) - Wherever found	T	\$16,500	\$1,263,554,511
856	Ziziphus, Florida ( <i>Ziziphus celata</i> ) - Wherever found	E	\$16,500	\$1,263,571,011
857	Tuctoria, Greene's ( <i>Tuctoria greenei</i> ) - Wherever found	E	\$16,424	\$1,263,587,435
858	Checker-mallow, Keck's ( <i>Sidalcea keckii</i> ) - Wherever found	E	\$16,393	\$1,263,603,828
859	Meadowfoam, large-flowered woolly ( <i>Limnanthes pumila</i> ssp. <i>grandiflora</i> ) - Wherever found	E	\$16,310	\$1,263,620,138
860	Mustard, Penland alpine fen ( <i>Eutrema penlandii</i> ) - Wherever found	T	\$16,300	\$1,263,636,438
861	Shrimp, anchialine pool ( <i>Vetericaris chaceorum</i> ) - Wherever found	E	\$16,254	\$1,263,652,692
862	Hornsnail, rough ( <i>Pleurocera foremani</i> ) - Wherever found	E	\$16,196	\$1,263,668,888
863	Towhee, Inyo California ( <i>Pipilo crissalis eremophilus</i> ) - Wherever found	T	\$16,155	\$1,263,685,043
864	Cactus, black lace ( <i>Echinocereus reichenbachii</i> var. <i>albertii</i> ) - Wherever found	E	\$16,089	\$1,263,701,132
865	Beetle, Comal Springs dryopid ( <i>Stygoparnus comalensis</i> ) - Wherever found	E	\$15,972	\$1,263,717,104
866	Snail, Pecos assiminea ( <i>Assimineia pecos</i> ) - Wherever found	E	\$15,850	\$1,263,732,954

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
867	Koki`o ke`oke`o ( <i>Hibiscus arnottianus</i> ssp. <i>immaculatus</i> ) - Wherever found	E	\$15,778	\$1,263,748,732
868	Pocketbook, finelined ( <i>Lampsilis altilis</i> ) - Wherever found	T	\$15,763	\$1,263,764,495
869	Ground-Dove, Friendly ( <i>Gallicolumba stairi</i> ) - American Samoa DPS	E	\$15,758	\$1,263,780,253
870	Hawk, Puerto Rican broad-winged ( <i>Buteo platypterus brunnescens</i> ) - Wherever found	E	\$15,750	\$1,263,796,003
871	Spider, Kauai cave wolf or pe'e pe'e maka 'ole ( <i>Adelocosa anops</i> ) - Wherever found	E	\$15,737	\$1,263,811,740
872	Rock-cress, Hoffmann's ( <i>Arabis hoffmannii</i> ) - Wherever found	E	\$15,725	\$1,263,827,465
873	Twinpod, Dudley Bluffs ( <i>Physaria obcordata</i> ) - Wherever found	T	\$15,700	\$1,263,843,165
874	Milk-vetch, Ash meadows ( <i>Astragalus phoenix</i> ) - Wherever found	T	\$15,600	\$1,263,858,765
874	Ivesia, Ash Meadows ( <i>Ivesia kingii</i> var. <i>eremica</i> ) - Wherever found	T	\$15,600	\$1,263,874,365
874	Blazingstar, Ash Meadows ( <i>Mentzelia leucophylla</i> ) - Wherever found	T	\$15,600	\$1,263,889,965
874	Sunray, Ash Meadows ( <i>Enceliopsis nudicaulis</i> var. <i>corrugata</i> ) - Wherever found	T	\$15,600	\$1,263,905,565
875	Flannelbush, Pine Hill ( <i>Fremontodendron californicum</i> ssp. <i>decumbens</i> ) - Wherever found	E	\$15,550	\$1,263,921,115
876	Moccasinshell, Suwannee ( <i>Medionidus walkeri</i> ) - Wherever found	T	\$15,536	\$1,263,936,651
877	Ha`iwale ( <i>Cyrtandra munroi</i> ) - Wherever found	E	\$15,516	\$1,263,952,167
878	Cobana negra ( <i>Stahlia monosperma</i> ) - Wherever found	T	\$15,500	\$1,263,967,667
879	Crayfish, Shasta ( <i>Pacifastacus fortis</i> ) - Wherever found	E	\$15,495	\$1,263,983,162
880	`aku ( <i>Cyanea tritomantha</i> ) - Wherever found	E	\$15,493	\$1,263,998,655
881	Damselfly, Pacific Hawaiian ( <i>Megalagrion pacificum</i> ) - Wherever found	E	\$15,475	\$1,264,014,130
882	Cactus, Key tree ( <i>Pilosocereus robinii</i> ) - Wherever found	E	\$15,342	\$1,264,029,472
883	Fatmucket, Arkansas ( <i>Lampsilis powellii</i> ) - Wherever found	T	\$15,312	\$1,264,044,784
884	Darter, yellowcheek ( <i>Etheostoma moorei</i> ) - Wherever found	E	\$15,300	\$1,264,060,084
885	Orcutt grass, hairy ( <i>Orcuttia pilosa</i> ) - Wherever found	E	\$15,256	\$1,264,075,340
886	Gambusia, Pecos ( <i>Gambusia nobilis</i> ) - Wherever found	E	\$15,250	\$1,264,090,590
886	Amphipod, Noel's ( <i>Gammarus desperatus</i> ) - Wherever found	E	\$15,250	\$1,264,105,840
887	Sea turtle, green ( <i>Chelonia mydas</i> ) - Mediterranean DPS	E	\$15,000	\$1,264,120,840
887	Sea turtle, loggerhead ( <i>Caretta caretta</i> ) - Mediterranean Sea DPS	E	\$15,000	\$1,264,135,840
888	Loulu ( <i>Pritchardia remota</i> ) - Wherever found	E	\$14,948	\$1,264,150,788
889	Beardtongue, Parachute ( <i>Penstemon debilis</i> ) - Wherever found	T	\$14,900	\$1,264,165,688
890	Springsnail, Roswell ( <i>Pyrgulopsis roswellensis</i> ) - Wherever found	E	\$14,850	\$1,264,180,538
891	Springsnail, Koster's ( <i>Juturnia kosteri</i> ) - Wherever found	E	\$14,750	\$1,264,195,288
892	Dudleya, marcescent ( <i>Dudleya cymosa</i> ssp. <i>marcescens</i> ) - Wherever found	T	\$14,700	\$1,264,209,988
893	No Common Name ( <i>Stenogyne kauaulaensis</i> ) - Wherever found	E	\$14,647	\$1,264,224,635
894	Wire-lettuce, Malheur ( <i>Stephanomeria malheurensis</i> ) - Wherever found	E	\$14,620	\$1,264,239,255

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
895	Dudleya, Conejo ( <i>Dudleya abramsii</i> ssp. <i>parva</i> ) - Wherever found	T	\$14,600	\$1,264,253,855
896	`Anunu ( <i>Sicyos macrophyllus</i> ) - Wherever found	E	\$14,509	\$1,264,268,364
897	Bat, Pacific sheath-tailed ( <i>Emballonura semicaudata semicaudata</i> ) - American Samoa	E	\$14,500	\$1,264,282,864
897	Daisy, Parish's ( <i>Erigeron parishii</i> ) - Wherever found	T	\$14,500	\$1,264,297,364
898	Phacelia, North Park ( <i>Phacelia formosula</i> ) - Wherever found	E	\$14,465	\$1,264,311,829
899	Ceanothus, Pine Hill ( <i>Ceanothus roderickii</i> ) - Wherever found	E	\$14,408	\$1,264,326,237
900	Harebells, Avon Park ( <i>Crotalaria avonensis</i> ) - Wherever found	E	\$14,400	\$1,264,340,637
901	Turtle, Sonoyta mud ( <i>Kinosternon sonoriense longifemorale</i> ) - Wherever found	E	\$14,256	\$1,264,354,893
902	Pigeon, Puerto Rican plain ( <i>Columba inornata wetmorei</i> ) - Wherever found	E	\$14,207	\$1,264,369,100
903	Bedstraw, El Dorado ( <i>Galium californicum</i> ssp. <i>sierrae</i> ) - Wherever found	E	\$14,150	\$1,264,383,250
904	Goldenrod, Short's ( <i>Solidago shortii</i> ) - Wherever found	E	\$14,132	\$1,264,397,382
905	Clubshell, southern ( <i>Pleurobema decisum</i> ) - Wherever found	E	\$14,065	\$1,264,411,447
906	Lau `ehu ( <i>Panicum niihauense</i> ) - Wherever found	E	\$14,036	\$1,264,425,483
907	Coqui, golden ( <i>Eleutherodactylus jasperii</i> ) - Wherever found	T	\$14,000	\$1,264,439,483
907	Bush-clover, prairie ( <i>Lespedeza leptostachya</i> ) - Wherever found	T	\$14,000	\$1,264,453,483
908	Nehe ( <i>Lipochaeta venosa</i> ) - Wherever found	E	\$13,953	\$1,264,467,436
909	Haha ( <i>Cyanea calycina</i> ) - Wherever found	E	\$13,952	\$1,264,481,388
910	Rock-cress, McDonald's ( <i>Arabis macdonaldiana</i> ) - Wherever found	E	\$13,900	\$1,264,495,288
911	Thrush, Molokai ( <i>Myadestes lanaiensis rutha</i> ) - Wherever found	E	\$13,878	\$1,264,509,166
912	No Common Name ( <i>Delissea rhytidosperma</i> ) - Wherever found	E	\$13,820	\$1,264,522,986
912	No Common Name ( <i>Schiedea kauaiensis</i> ) - Wherever found	E	\$13,820	\$1,264,536,806
913	Warbler, elfin-woods ( <i>Setophaga angelae</i> ) - Wherever found	T	\$13,750	\$1,264,550,556
913	Malacothrix, Santa Cruz Island ( <i>Malacothrix indecora</i> ) - Wherever found	E	\$13,750	\$1,264,564,306
914	`Oha wai ( <i>Clermontia lindseyana</i> ) - Wherever found	E	\$13,733	\$1,264,578,039
915	`Awikiwiki ( <i>Canavalia molokaiensis</i> ) - Wherever found	E	\$13,712	\$1,264,591,751
916	Sneezeweed, Virginia ( <i>Helenium virginicum</i> ) - Wherever found	T	\$13,640	\$1,264,605,391
917	Rosemary, Etonia ( <i>Conradina etonia</i> ) - Wherever found	E	\$13,625	\$1,264,619,016
918	Orangutan ( <i>Pongo pygmaeus</i> ) - Wherever found	E	\$13,585	\$1,264,632,601
919	Cactus, Pariette ( <i>Sclerocactus brevispinus</i> ) - Wherever found	T	\$13,579	\$1,264,646,180
920	Haha ( <i>Cyanea glabra</i> ) - Wherever found	E	\$13,537	\$1,264,659,717
921	Chub, Hutton tui ( <i>Gila bicolor</i> ssp.) - Wherever found	T	\$13,500	\$1,264,673,217
921	Mint, Lakela's ( <i>Dicerandra immaculata</i> ) - Wherever found	E	\$13,500	\$1,264,686,717
921	Pelos del diablo ( <i>Aristida portoricensis</i> ) - Wherever found	E	\$13,500	\$1,264,700,217



Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
921	No Common Name ( <i>Aristida chaseae</i> ) - Wherever found	E	\$13,500	\$1,264,713,717
922	Pilo ( <i>Kadua laxiflora</i> ) - Wherever found	E	\$13,491	\$1,264,727,208
923	No Common Name ( <i>Schiedea diffusa</i> ssp. <i>macraei</i> ) - Wherever found	E	\$13,425	\$1,264,740,633
924	Dwarf-flax, Marin ( <i>Hesperolinon congestum</i> ) - Wherever found	T	\$13,400	\$1,264,754,033
925	Oha ( <i>Delissea subcordata</i> ) - Wherever found	E	\$13,313	\$1,264,767,346
926	Alani ( <i>Melicope balloui</i> ) - Wherever found	E	\$13,284	\$1,264,780,630
927	[no common name] Snail ( <i>Ostodes strigatus</i> ) - Wherever found	E	\$13,223	\$1,264,793,853
927	[no common name] Snail ( <i>Eua zebrina</i> ) - Wherever found	E	\$13,223	\$1,264,807,076
928	Yerba santa, Lompoc ( <i>Eriodictyon capitatum</i> ) - Wherever found	E	\$13,100	\$1,264,820,176
929	Snake, Atlantic salt marsh ( <i>Nerodia clarkii taeniata</i> ) - Wherever found	T	\$13,069	\$1,264,833,245
930	No Common Name ( <i>Tetramolopium remyi</i> ) - Wherever found	E	\$13,059	\$1,264,846,304
931	No Common Name ( <i>Lyonia truncata</i> var. <i>proctorii</i> ) - Wherever found	E	\$13,000	\$1,264,859,304
931	No Common Name ( <i>Vernonia proctorii</i> ) - Wherever found	E	\$13,000	\$1,264,872,304
932	(honeyeater), Mao (= maomao) ( <i>Gymnomyza samoensis</i> ) - Wherever found	E	\$12,929	\$1,264,885,233
933	Blazingstar, Heller's ( <i>Liatris helleri</i> ) - Wherever found	T	\$12,900	\$1,264,898,133
934	Makou ( <i>Peucedanum sandwicense</i> ) - Wherever found	T	\$12,888	\$1,264,911,021
935	Haha ( <i>Cyanea mceldowneyi</i> ) - Wherever found	E	\$12,863	\$1,264,923,884
936	Snail, Newcomb's ( <i>Erinna newcombi</i> ) - Wherever found	T	\$12,797	\$1,264,936,681
937	Rock-cress, Braun's ( <i>Arabis perstellata</i> ) - Wherever found	E	\$12,749	\$1,264,949,430
938	Hau kuahiwi ( <i>Hibiscadelphus giffardianus</i> ) - Wherever found	E	\$12,736	\$1,264,962,166
939	Ha`iwale ( <i>Cyrtandra limahuliensis</i> ) - Wherever found	T	\$12,639	\$1,264,974,805
939	No Common Name ( <i>Phyllostegia wawrana</i> ) - Wherever found	E	\$12,639	\$1,264,987,444
940	Darter, snail ( <i>Percina tanasi</i> ) - Wherever found	T	\$12,627	\$1,265,000,071
941	`O`u (honeycreeper) ( <i>Psittirostra psittacea</i> ) - Wherever found	E	\$12,588	\$1,265,012,659
942	Kamapua`a ( <i>Kadua fluviatilis</i> ) - Wherever found	E	\$12,575	\$1,265,025,234
943	Aupaka ( <i>Isodendrion laurifolium</i> ) - Wherever found	E	\$12,515	\$1,265,037,749
943	`Aiakeakua, popolo ( <i>Solanum sandwicense</i> ) - Wherever found	E	\$12,515	\$1,265,050,264
944	Fly, Hawaiian picture-wing ( <i>Drosophila differens</i> ) - Wherever found	E	\$12,503	\$1,265,062,767
945	Salamander, Sonora tiger ( <i>Ambystoma tigrinum stebbinsi</i> ) - Wherever found	E	\$12,500	\$1,265,075,267
945	Rush-pea, slender ( <i>Hoffmannseggia tenella</i> ) - Wherever found	E	\$12,500	\$1,265,087,767
946	Vetch, Hawaiian ( <i>Vicia menziesii</i> ) - Wherever found	E	\$12,496	\$1,265,100,263
947	Rocksail, painted ( <i>Leptoxis taeniata</i> ) - Wherever found	T	\$12,396	\$1,265,112,659
948	Paintbrush, Tiburon ( <i>Castilleja affinis</i> ssp. <i>neglecta</i> ) - Wherever found	E	\$12,300	\$1,265,124,959

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
949	Ihi ( <i>Portulaca villosa</i> ) - Wherever found	E	\$12,274	\$1,265,137,233
950	Spineflower, Howell's ( <i>Chorizanthe howellii</i> ) - Wherever found	E	\$12,200	\$1,265,149,433
951	Marstonia, royal (snail) ( <i>Pyrgulopsis ogmorhapha</i> ) - Wherever found	E	\$12,125	\$1,265,161,558
952	Isopod, Socorro ( <i>Thermosphaeroma thermophilus</i> ) - Wherever found	E	\$12,100	\$1,265,173,658
953	Haiwale ( <i>Cyrtandra waiolani</i> ) - Wherever found	E	\$12,036	\$1,265,185,694
954	Butterfly, Myrtle's silverspot ( <i>Speyeria zerene myrtleae</i> ) - Wherever found	E	\$12,000	\$1,265,197,694
954	Goetzea, beautiful ( <i>Goetzea elegans</i> ) - Wherever found	E	\$12,000	\$1,265,209,694
955	Seal, Ringed ( <i>Phoca (=Pusa) hispida hispida</i> ) - Wherever found	T	\$11,998	\$1,265,221,692
956	Kuawawaenuhu ( <i>Schiedea lychnoides</i> ) - Wherever found	E	\$11,988	\$1,265,233,680
957	Creeper, Oahu ( <i>Paroreomyza maculata</i> ) - Wherever found	E	\$11,914	\$1,265,245,594
958	No Common Name ( <i>Schiedea verticillata</i> ) - Wherever found	E	\$11,906	\$1,265,257,500
959	Kolea ( <i>Myrsine vaccinioides</i> ) - Wherever found	E	\$11,893	\$1,265,269,393
960	Manzanita, Morro ( <i>Arctostaphylos morroensis</i> ) - Wherever found	T	\$11,800	\$1,265,281,193
961	Kamakahala ( <i>Labordia lydgatei</i> ) - Wherever found	E	\$11,792	\$1,265,292,985
961	No Common Name ( <i>Doryopteris angelica</i> ) - Wherever found	E	\$11,792	\$1,265,304,777
962	Phacelia, island ( <i>Phacelia insularis</i> ssp. <i>insularis</i> ) - Wherever found	E	\$11,779	\$1,265,316,556
963	Haha ( <i>Cyanea lobata</i> ) - Wherever found	E	\$11,713	\$1,265,328,269
964	Heau ( <i>Exocarpos menziesii</i> ) -	E	\$11,695	\$1,265,339,964
965	`Oha wai ( <i>Clermontia peleana</i> ) - Wherever found	E	\$11,680	\$1,265,351,644
966	Pearlymussel, dromedary ( <i>Dromus dromas</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$11,677	\$1,265,363,321
967	Loulu ( <i>Pritchardia lanigera</i> ) - Wherever found	E	\$11,676	\$1,265,374,997
968	`O`o, Kauai (honeyeater) ( <i>Moho braccatus</i> ) - Wherever found	E	\$11,663	\$1,265,386,660
969	Clarkia, Presidio ( <i>Clarkia franciscana</i> ) - Wherever found	E	\$11,600	\$1,265,398,260
970	Isopod, Lee County cave ( <i>Lirceus usdagalun</i> ) - Wherever found	E	\$11,500	\$1,265,409,760
970	Chumbo, Higo ( <i>Harrisia portoricensis</i> ) - Wherever found	T	\$11,500	\$1,265,421,260
971	Picture-wing fly, Hawaiian ( <i>Drosophila neoclavisetae</i> ) - Wherever found	E	\$11,489	\$1,265,432,749
972	Pronghorn, Sonoran ( <i>Antilocapra americana sonoriensis</i> ) - Wherever found, except where listed as an experimental population	E	\$11,430	\$1,265,444,179
973	Monardella, willowy ( <i>Monardella viminea</i> ) - Wherever found	E	\$11,400	\$1,265,455,579
973	Navarretia, many-flowered ( <i>Navarretia leucocephala</i> ssp. <i>plieantha</i> ) - Wherever found	E	\$11,400	\$1,265,466,979
973	Gooseberry, Miccosukee ( <i>Ribes echinellum</i> ) - Wherever found	T	\$11,400	\$1,265,478,379
974	Fly, Hawaiian picture-wing ( <i>Drosophila mulli</i> ) - Wherever found	T	\$11,322	\$1,265,489,701

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
975	Pua `ala ( <i>Brighamia rockii</i> ) - Wherever found	E	\$11,278	\$1,265,500,979
976	Hala pepe ( <i>Pleomele fernaldii</i> ) - Wherever found	E	\$11,214	\$1,265,512,193
977	Beetle, American burying ( <i>Nicrophorus americanus</i> ) - Ex-Pop, SW Missouri	NS	\$11,100	\$1,265,523,293
977	Spurge, Hoover's ( <i>Chamaesyce hooveri</i> ) - Wherever found	T	\$11,100	\$1,265,534,393
978	Koki`o ( <i>Kokia drynarioides</i> ) - Wherever found	E	\$11,074	\$1,265,545,467
979	No Common Name ( <i>Phyllostegia glabra</i> var. <i>lanaiensis</i> ) - Wherever found	E	\$11,031	\$1,265,556,498
979	No Common Name ( <i>Viola lanaiensis</i> ) - Wherever found	E	\$11,031	\$1,265,567,529
979	Haha ( <i>Cyanea dunbariae</i> ) - Wherever found	E	\$11,031	\$1,265,578,560
979	Haha ( <i>Cyanea procera</i> ) - Wherever found	E	\$11,031	\$1,265,589,591
979	No Common Name ( <i>Phyllostegia mannii</i> ) - Wherever found	E	\$11,031	\$1,265,600,622
979	No Common Name ( <i>Phyllostegia hispida</i> ) - Wherever found	E	\$11,031	\$1,265,611,653
980	Pikeminnow (=squawfish), Colorado ( <i>Ptychocheilus lucius</i> ) - Salt and Verde R. drainages, AZ	EXPN	\$11,000	\$1,265,622,653
980	Shiner, beautiful ( <i>Cyprinella formosa</i> ) - Wherever found	T	\$11,000	\$1,265,633,653
981	Cypress, Santa Cruz ( <i>Cupressus abramsiana</i> ) - Wherever found	T	\$10,931	\$1,265,644,584
982	Pussypaws, Mariposa ( <i>Calyptridium pulchellum</i> ) - Wherever found	T	\$10,900	\$1,265,655,484
983	Salamander, Texas blind ( <i>Eurycea</i> [=Typhlomolge] <i>rathbuni</i> ) - Wherever found	E	\$10,840	\$1,265,666,324
983	Salamander, San Marcos ( <i>Eurycea nana</i> ) - Wherever found	T	\$10,840	\$1,265,677,164
983	Darter, fountain ( <i>Etheostoma fonticola</i> ) - Wherever found	E	\$10,840	\$1,265,688,004
983	Beetle, Comal Springs riffle ( <i>Heterelmis comalensis</i> ) - Wherever found	E	\$10,840	\$1,265,698,844
983	Wild-rice, Texas ( <i>Zizania texana</i> ) - Wherever found	E	\$10,840	\$1,265,709,684
984	Bluegrass, Napa ( <i>Poa napensis</i> ) - Wherever found	E	\$10,800	\$1,265,720,484
985	Salamander, desert slender ( <i>Batrachoseps aridus</i> ) - Wherever found	E	\$10,778	\$1,265,731,262
985	Kamakahala ( <i>Labordia triflora</i> ) - Wherever found	E	\$10,778	\$1,265,742,040
985	No Common Name ( <i>Neraudia sericea</i> ) - Wherever found	E	\$10,778	\$1,265,752,818
985	Loulu ( <i>Pritchardia munroi</i> ) - Wherever found	E	\$10,778	\$1,265,763,596
985	Haha ( <i>Cyanea gibsonii</i> ) - Wherever found	E	\$10,778	\$1,265,774,374
985	Haha ( <i>Cyanea mannii</i> ) - Wherever found	E	\$10,778	\$1,265,785,152
985	Na`ena`e ( <i>Dubautia plantaginea</i> ssp. <i>humilis</i> ) - Wherever found	E	\$10,778	\$1,265,795,930
985	No Common Name ( <i>Lysimachia lydgatei</i> ) - Wherever found	E	\$10,778	\$1,265,806,708
985	No Common Name ( <i>Lysimachia maxima</i> ) - Wherever found	E	\$10,778	\$1,265,817,486
985	Loulu ( <i>Pritchardia napaliensis</i> ) - Wherever found	E	\$10,778	\$1,265,828,264
985	No Common Name ( <i>Asplenium dielmannii</i> ) - Wherever found	E	\$10,778	\$1,265,839,042
986	Panicgrass, Carter's ( <i>Panicum fauriei</i> var. <i>carteri</i> ) - Wherever found	E	\$10,701	\$1,265,849,743

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
987	Larkspur, Baker's ( <i>Delphinium bakeri</i> ) - Wherever found	E	\$10,700	\$1,265,860,443
988	No Common Name ( <i>Platydesma cornuta</i> var. <i>decurrens</i> ) - Wherever found	E	\$10,645	\$1,265,871,088
989	Haha ( <i>Cyanea munroi</i> ) - Wherever found	E	\$10,629	\$1,265,881,717
989	No Common Name ( <i>Festuca molokaiensis</i> ) - Wherever found	E	\$10,629	\$1,265,892,346
990	Dace, Ash Meadows speckled ( <i>Rhinichthys osculus nevadensis</i> ) - Wherever found	E	\$10,582	\$1,265,902,928
991	Kamanomano ( <i>Cenchrus agrimonoides</i> ) - Wherever found	E	\$10,505	\$1,265,913,433
992	Rose-mallow, Neches River ( <i>Hibiscus dasycalyx</i> ) - Wherever found	T	\$10,500	\$1,265,923,933
993	Diellia, Asplenium-leaved ( <i>Asplenium dielirectum</i> ) - Wherever found	E	\$10,449	\$1,265,934,382
994	Larkspur, yellow ( <i>Delphinium luteum</i> ) - Wherever found	E	\$10,400	\$1,265,944,782
995	Paintbrush, soft-leaved ( <i>Castilleja mollis</i> ) - Wherever found	E	\$10,374	\$1,265,955,156
996	No Common Name ( <i>Schiedea spergulina</i> var. <i>spergulina</i> ) - Wherever found	T	\$10,320	\$1,265,965,476
997	Allocarya, Calistoga ( <i>Plagiobothrys strictus</i> ) - Wherever found	E	\$10,300	\$1,265,975,776
998	Ha`iwale ( <i>Cyrtandra giffardii</i> ) - Wherever found	E	\$10,299	\$1,265,986,075
999	Nightjar, Puerto Rican ( <i>Caprimulgus noctitherus</i> ) - Wherever found	E	\$10,250	\$1,265,996,325
1000	Milkvetch, Osterhout ( <i>Astragalus osterhoutii</i> ) - Wherever found	E	\$10,115	\$1,266,006,440
1001	Spinedace, Big Spring ( <i>Lepidomeda mollispinis pratensis</i> ) - Wherever found	T	\$10,103	\$1,266,016,543
1002	Bittercress, small-anthered ( <i>Cardamine micranthera</i> ) - Wherever found	E	\$10,100	\$1,266,026,643
1002	Sand-verbena, large-fruited ( <i>Abronia macrocarpa</i> ) - Wherever found	E	\$10,100	\$1,266,036,743
1003	Wolf, gray ( <i>Canis lupus</i> ) - Western Great Lakes DPS	RT	\$10,000	\$1,266,046,743
1003	Salamander, Jollyville Plateau ( <i>Eurycea tonkawae</i> ) - Wherever found	T	\$10,000	\$1,266,056,743
1003	Rocksnailed, plicate ( <i>Leptoxis plicata</i> ) - Wherever found	E	\$10,000	\$1,266,066,743
1003	Skipper, Laguna Mountains ( <i>Pyrgus ruralis lagunae</i> ) - Wherever found	E	\$10,000	\$1,266,076,743
1003	Harvestman, Bone Cave ( <i>Texella reyesi</i> ) - Wherever found	E	\$10,000	\$1,266,086,743
1003	Amphipod, Illinois cave ( <i>Gammarus acherondytes</i> ) - Wherever found	E	\$10,000	\$1,266,096,743
1003	Pawpaw, Rugel's ( <i>Deeringothamnus rugelii</i> ) - Wherever found	E	\$10,000	\$1,266,106,743
1004	Haiwale ( <i>Cyrtandra gracilis</i> ) - Wherever found	E	\$9,970	\$1,266,116,713
1005	No Common Name ( <i>Kadua haupuensis</i> ) - Wherever found	E	\$9,909	\$1,266,126,622
1006	Shark, Scalloped Hammerhead ( <i>Sphyrna lewini</i> ) - Eastern Pacific DPS	E	\$9,902	\$1,266,136,524
1007	Haha ( <i>Cyanea magnicalyx</i> ) - Wherever found	E	\$9,868	\$1,266,146,392
1008	No Common Name ( <i>Kadua parvula</i> ) - Wherever found	E	\$9,865	\$1,266,156,257
1008	No Common Name ( <i>Lobelia niihauensis</i> ) - Wherever found	E	\$9,865	\$1,266,166,122
1008	Alani ( <i>Melicope christophersenii</i> ) - Wherever found	E	\$9,865	\$1,266,175,987
1008	Kulu`i ( <i>Nototrichium humile</i> ) - Wherever found	E	\$9,865	\$1,266,185,852

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1009	Gilia, Hoffmann's slender-flowered ( <i>Gilia tenuiflora</i> ssp. <i>hoffmannii</i> ) - Wherever found	E	\$9,858	\$1,266,195,710
1010	Nohoanu ( <i>Geranium hillebrandii</i> ) - Wherever found	E	\$9,827	\$1,266,205,537
1011	No Common Name ( <i>Polyscias bisattenuata</i> ) - Wherever found	E	\$9,820	\$1,266,215,357
1012	No Common Name ( <i>Abutilon sandwicense</i> ) - Wherever found	E	\$9,807	\$1,266,225,164
1013	Haha ( <i>Cyanea grimesiana</i> ssp. <i>grimesiana</i> ) - Wherever found	E	\$9,722	\$1,266,234,886
1014	Haiwale ( <i>Cyrtandra paliku</i> ) - Wherever found	E	\$9,717	\$1,266,244,603
1015	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i> ) - Central Valley spring-run ESU	EXPN	\$9,700	\$1,266,254,303
1016	Nohoanu ( <i>Geranium kauaiense</i> ) - Wherever found	E	\$9,669	\$1,266,263,972
1016	Alani ( <i>Melicope puberula</i> ) - Wherever found	E	\$9,669	\$1,266,273,641
1017	Rush-rose, island ( <i>Helianthemum greenei</i> ) - Wherever found	T	\$9,632	\$1,266,283,273
1018	No Common Name ( <i>Schiedea laui</i> ) - Wherever found	E	\$9,615	\$1,266,292,888
1019	Thistle, fountain ( <i>Cirsium fontinale</i> var. <i>fontinale</i> ) - Wherever found	E	\$9,600	\$1,266,302,488
1020	Trillium, relict ( <i>Trillium reliquum</i> ) - Wherever found	E	\$9,538	\$1,266,312,026
1021	`Akoko ( <i>Euphorbia kuwaleana</i> ) - Wherever found	E	\$9,528	\$1,266,321,554
1022	No Common Name ( <i>Phyllostegia floribunda</i> ) - Wherever found	E	\$9,523	\$1,266,331,077
1023	Lehua makanoë ( <i>Lysimachia daphnoides</i> ) - Wherever found	E	\$9,517	\$1,266,340,594
1024	No Common Name ( <i>Sicyos lanceoloideus</i> ) - Wherever found	E	\$9,509	\$1,266,350,103
1025	Boxwood, Vahl's ( <i>Buxus vahlii</i> ) - Wherever found	E	\$9,500	\$1,266,359,603
1025	Mustard, Carter's ( <i>Warea carteri</i> ) - Wherever found	E	\$9,500	\$1,266,369,103
1026	Koki`o ke`oke`o ( <i>Hibiscus waimeae</i> ssp. <i>hannerae</i> ) - Wherever found	E	\$9,473	\$1,266,378,576
1026	Remya, Maui ( <i>Remya mauiensis</i> ) - Wherever found	E	\$9,473	\$1,266,388,049
1026	Ho`awa ( <i>Pittosporum napaliense</i> ) - Wherever found	E	\$9,473	\$1,266,397,522
1027	No Common Name ( <i>Asplenium unisorum</i> ) - Wherever found	E	\$9,470	\$1,266,406,992
1028	Prickly-apple, fragrant ( <i>Cereus eriophorus</i> var. <i>fragrans</i> ) - Wherever found	E	\$9,400	\$1,266,416,392
1028	Sunflower, San Mateo woolly ( <i>Eriophyllum latilobum</i> ) - Wherever found	E	\$9,400	\$1,266,425,792
1029	Pigtoe, southern ( <i>Pleurobema georgianum</i> ) - Wherever found	E	\$9,329	\$1,266,435,121
1030	Papala ( <i>Charpentiera densiflora</i> ) - Wherever found	E	\$9,321	\$1,266,444,442
1030	No Common Name ( <i>Abutilon eremitopetalum</i> ) - Wherever found	E	\$9,321	\$1,266,453,763
1031	Grass, Solano ( <i>Tuctoria mucronata</i> ) - Wherever found	E	\$9,300	\$1,266,463,063
1032	`Awikiwiki ( <i>Canavalia pubescens</i> ) - Wherever found	E	\$9,148	\$1,266,472,211
1033	Popolo ( <i>Cyanea solanacea</i> ) - Wherever found	E	\$9,145	\$1,266,481,356
1033	Haha ( <i>Cyanea profuga</i> ) - Wherever found	E	\$9,145	\$1,266,490,501
1033	Haha ( <i>Cyanea mauiensis</i> ) - Wherever found	E	\$9,145	\$1,266,499,646

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1034	Haha ( <i>Cyanea duvalliorum</i> ) - Wherever found	E	\$9,099	\$1,266,508,745
1034	No Common Name ( <i>Huperzia stemmermanniae</i> ) - Wherever found	E	\$9,099	\$1,266,517,844
1035	Milk-vetch, Cushenbury ( <i>Astragalus albens</i> ) - Wherever found	E	\$9,000	\$1,266,526,844
1035	Sedge, golden ( <i>Carex lutea</i> ) - Wherever found	E	\$9,000	\$1,266,535,844
1036	No Common Name ( <i>Schiedea hookeri</i> ) - Wherever found	E	\$8,999	\$1,266,544,843
1037	Butterfly, Oregon silverspot ( <i>Speyeria zerene hippolyta</i> ) -	EXPN	\$8,903	\$1,266,553,746
1038	Hypericum, highlands scrub ( <i>Hypericum cumulicola</i> ) - Wherever found	E	\$8,900	\$1,266,562,646
1039	No Common Name ( <i>Schiedea salicaria</i> ) - Wherever found	E	\$8,895	\$1,266,571,541
1040	Haha ( <i>Cyanea obtusa</i> ) - Wherever found	E	\$8,892	\$1,266,580,433
1041	No Common Name ( <i>Sanicula purpurea</i> ) - Wherever found	E	\$8,796	\$1,266,589,229
1041	'Awiwi ( <i>Kadua cookiana</i> ) - Wherever found	E	\$8,796	\$1,266,598,025
1042	Yellow-faced bee, easy ( <i>Hylaeus facilis</i> ) - Wherever found	E	\$8,784	\$1,266,606,809
1042	Yellow-faced bee, Hawaiian ( <i>Hylaeus longiceps</i> ) - Wherever found	E	\$8,784	\$1,266,615,593
1043	Ma`oli`oli ( <i>Schiedea hawaiiensis</i> ) - Wherever found	E	\$8,762	\$1,266,624,355
1043	No Common Name ( <i>Stenogyne cranwelliae</i> ) - Wherever found	E	\$8,762	\$1,266,633,117
1044	No Common Name ( <i>Remya kauaiensis</i> ) - Wherever found	E	\$8,750	\$1,266,641,867
1045	Darter, amber ( <i>Percina antesella</i> ) - Wherever found	E	\$8,732	\$1,266,650,599
1046	Pamakani ( <i>Tetramolopium capillare</i> ) - Wherever found	E	\$8,712	\$1,266,659,311
1046	'Oha wai ( <i>Clermontia oblongifolia</i> ssp. <i>brevipes</i> ) - Wherever found	E	\$8,712	\$1,266,668,023
1047	Campion, fringed ( <i>Silene polypetala</i> ) - Wherever found	E	\$8,612	\$1,266,676,635
1048	Clubshell, ovate ( <i>Pleurobema perovatum</i> ) - Wherever found	E	\$8,611	\$1,266,685,246
1049	Butterwort, Godfrey's ( <i>Pinguicula ionantha</i> ) - Wherever found	T	\$8,600	\$1,266,693,846
1050	Pawpaw, four-petal ( <i>Asimina tetramera</i> ) - Wherever found	E	\$8,500	\$1,266,702,346
1050	Cactus, Sneed pincushion ( <i>Coryphantha sneedii</i> var. <i>sneedii</i> ) - Wherever found	E	\$8,500	\$1,266,710,846
1050	No Common Name ( <i>Mitracarpus maxwelliae</i> ) - Wherever found	E	\$8,500	\$1,266,719,346
1050	No Common Name ( <i>Mitracarpus polycladus</i> ) - Wherever found	E	\$8,500	\$1,266,727,846
1051	Loulu ( <i>Pritchardia aylmer-robinsonii</i> ) - Wherever found	E	\$8,459	\$1,266,736,305
1051	No Common Name ( <i>Schiedea membranacea</i> ) - Wherever found	E	\$8,459	\$1,266,744,764
1051	No Common Name ( <i>Schiedea sarmentosa</i> ) - Wherever found	E	\$8,459	\$1,266,753,223
1051	Koki`o, Cooke's ( <i>Kokia cookei</i> ) - Wherever found	E	\$8,459	\$1,266,761,682
1051	Alani ( <i>Melicope mucronulata</i> ) - Wherever found	E	\$8,459	\$1,266,770,141
1051	Alani ( <i>Melicope quadrangularis</i> ) - Wherever found	E	\$8,459	\$1,266,778,600
1051	Kamakahala ( <i>Labordia tinifolia</i> var. <i>lanaiensis</i> ) - Wherever found	E	\$8,459	\$1,266,787,059

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1051	No Common Name ( <i>Schiedea lydgatei</i> ) - Wherever found	E	\$8,459	\$1,266,795,518
1051	Laulihilihi ( <i>Schiedea stellarioides</i> ) - Wherever found	E	\$8,459	\$1,266,803,977
1051	Kopa ( <i>Kadua cordata remyi</i> ) - Wherever found	E	\$8,459	\$1,266,812,436
1052	Moccasinshell, Alabama ( <i>Medionidus acutissimus</i> ) - Wherever found	T	\$8,456	\$1,266,820,892
1053	Pu`uka`a ( <i>Cyperus trachysanthos</i> ) - Wherever found	E	\$8,445	\$1,266,829,337
1054	White-eye, bridled ( <i>Zosterops conspicillatus conspicillatus</i> ) - Wherever found	E	\$8,444	\$1,266,837,781
1055	Kidneyshell, triangular ( <i>Ptychobranthus greenii</i> ) - Wherever found	E	\$8,430	\$1,266,846,211
1056	Haha ( <i>Cyanea marksii</i> ) - Wherever found	E	\$8,425	\$1,266,854,636
1056	Ha`iwale ( <i>Cyrtandra hematos</i> ) - Wherever found	E	\$8,425	\$1,266,863,061
1056	No Common Name ( <i>Phyllostegia pilosa</i> ) - Wherever found	E	\$8,425	\$1,266,871,486
1056	No Common Name ( <i>Cyrtandra wagneri</i> ) - Wherever found	E	\$8,425	\$1,266,879,911
1057	No Common Name ( <i>Schiedea nuttallii</i> ) - Wherever found	E	\$8,420	\$1,266,888,331
1058	No Common Name ( <i>Phyllostegia parviflora</i> ) - Wherever found	E	\$8,410	\$1,266,896,741
1059	Turtle, Alabama red-bellied ( <i>Pseudemys alabamensis</i> ) - Wherever found	E	\$8,358	\$1,266,905,099
1060	`Oha wai ( <i>Clermontia pyralaria</i> ) - Wherever found	E	\$8,326	\$1,266,913,425
1060	Haha ( <i>Cyanea st.-johnii</i> ) - Wherever found	E	\$8,326	\$1,266,921,751
1060	No Common Name ( <i>Tetramolopium lepidotum</i> ssp. <i>lepidotum</i> ) - Wherever found	E	\$8,326	\$1,266,930,077
1060	Pamakani ( <i>Viola chamissoniana</i> ssp. <i>chamissoniana</i> ) - Wherever found	E	\$8,326	\$1,266,938,403
1060	`aku`aku ( <i>Cyanea platyphylla</i> ) - Wherever found	E	\$8,326	\$1,266,946,729
1060	Hala pepe ( <i>Pleomele forbesii</i> ) - Wherever found	E	\$8,326	\$1,266,955,055
1060	Haha ( <i>Cyanea grimesiana</i> ssp. <i>obatae</i> ) - Wherever found	E	\$8,326	\$1,266,963,381
1060	Haha ( <i>Cyanea hamatiflora</i> ssp. <i>carlsonii</i> ) - Wherever found	E	\$8,326	\$1,266,971,707
1060	No Common Name ( <i>Phyllostegia warshaueri</i> ) - Wherever found	E	\$8,326	\$1,266,980,033
1061	No Common Name ( <i>Silene alexandri</i> ) - Wherever found	E	\$8,307	\$1,266,988,340
1061	Nehe ( <i>Melanthera kamolensis</i> ) - Wherever found	E	\$8,307	\$1,266,996,647
1062	Uvillo ( <i>Eugenia haematocarpa</i> ) - Wherever found	E	\$8,300	\$1,267,004,947
1062	Crownscale, San Jacinto Valley ( <i>Atriplex coronata</i> var. <i>notatior</i> ) - Wherever found	E	\$8,300	\$1,267,013,247
1063	Darter, goldline ( <i>Percina aurolineata</i> ) - Wherever found	T	\$8,272	\$1,267,021,519
1064	Salamander, Red Hills ( <i>Phaeognathus hubrichti</i> ) - Wherever found	T	\$8,248	\$1,267,029,767
1065	Madtom, smoky ( <i>Noturus baileyi</i> ) - Wherever found	E	\$8,247	\$1,267,038,014
1066	Kohe malama malama o kanaloa ( <i>Kanaloa kahoolawensis</i> ) - Wherever found	E	\$8,243	\$1,267,046,257
1067	Halomtano, Berenghenas ( <i>Solanum guamense</i> ) - Wherever found	E	\$8,238	\$1,267,054,495
1068	No Common Name ( <i>Gouania hillebrandii</i> ) - Wherever found	E	\$8,205	\$1,267,062,700

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1069	Ko`oko`olau ( <i>Bidens wiebkei</i> ) - Wherever found	E	\$8,188	\$1,267,070,888
1070	No Common Name ( <i>Phyllostegia helleri</i> ) -	E	\$8,172	\$1,267,079,060
1070	Ma`oli`oli ( <i>Schiedea pubescens</i> ) - Wherever found	E	\$8,172	\$1,267,087,232
1070	No Common Name ( <i>Bidens hillebrandiana</i> ssp. <i>hillebrandiana</i> ) - Wherever found	E	\$8,172	\$1,267,095,404
1071	Spurge, Garber's ( <i>Chamaesyce garberi</i> ) - Wherever found	T	\$8,142	\$1,267,103,546
1072	Bean, sea ( <i>Mucuna sloanei</i> var. <i>persericea</i> ) - Wherever found	E	\$8,104	\$1,267,111,650
1073	Kolea ( <i>Myrsine linearifolia</i> ) - Wherever found	T	\$8,083	\$1,267,119,733
1073	`Akoko ( <i>Euphorbia remyi</i> var. <i>remyi</i> ) - Wherever found	E	\$8,083	\$1,267,127,816
1074	`Akoko ( <i>Euphorbia haeleleana</i> ) - Wherever found	E	\$8,073	\$1,267,135,889
1074	No Common Name ( <i>Lobelia koolauensis</i> ) - Wherever found	E	\$8,073	\$1,267,143,962
1074	No Common Name ( <i>Phyllostegia hirsuta</i> ) - Wherever found	E	\$8,073	\$1,267,152,035
1074	Ma`oli`oli ( <i>Schiedea kealiae</i> ) - Wherever found	E	\$8,073	\$1,267,160,108
1074	No Common Name ( <i>Schiedea obovata</i> ) - Wherever found	E	\$8,073	\$1,267,168,181
1074	Haha ( <i>Cyanea crispa</i> ) - Wherever found	E	\$8,073	\$1,267,176,254
1074	Alani ( <i>Melicope lydgatei</i> ) - Wherever found	E	\$8,073	\$1,267,184,327
1074	Alani ( <i>Melicope makahae</i> ) - Wherever found	E	\$8,073	\$1,267,192,400
1074	No Common Name ( <i>Schiedea kaalae</i> ) - Wherever found	E	\$8,073	\$1,267,200,473
1074	No Common Name ( <i>Stenogyne kanehoana</i> ) - Wherever found	E	\$8,073	\$1,267,208,546
1074	No Common Name ( <i>Tetramolopium filiforme</i> ) - Wherever found	E	\$8,073	\$1,267,216,619
1074	`Ohe`ohe ( <i>Polyscias gymnocarpa</i> ) - Wherever found	E	\$8,073	\$1,267,224,692
1074	Nehe ( <i>Melanthera tenuifolia</i> ) - Wherever found	E	\$8,073	\$1,267,232,765
1074	No Common Name ( <i>Phyllostegia mollis</i> ) - Wherever found	E	\$8,073	\$1,267,240,838
1074	Hulumoa ( <i>Korthalsella degeneri</i> ) - Wherever found	E	\$8,073	\$1,267,248,911
1074	Kiponapona ( <i>Phyllostegia racemosa</i> ) - Wherever found	E	\$8,073	\$1,267,256,984
1074	No Common Name ( <i>Phyllostegia velutina</i> ) - Wherever found	E	\$8,073	\$1,267,265,057
1074	Haha ( <i>Cyanea acuminata</i> ) - Wherever found	E	\$8,073	\$1,267,273,130
1074	Haha ( <i>Cyanea koolauensis</i> ) - Wherever found	E	\$8,073	\$1,267,281,203
1074	Haha ( <i>Cyanea longiflora</i> ) - Wherever found	E	\$8,073	\$1,267,289,276
1074	Nanu ( <i>Gardenia mannii</i> ) - Wherever found	E	\$8,073	\$1,267,297,349
1074	No Common Name ( <i>Phyllostegia kaalaensis</i> ) - Wherever found	E	\$8,073	\$1,267,305,422
1074	No Common Name ( <i>Asplenium dielfalcatum</i> ) - Wherever found	E	\$8,073	\$1,267,313,495
1075	No Common Name ( <i>Diplazium molokaiense</i> ) - Wherever found	E	\$8,024	\$1,267,321,519
1076	Alopecurus, Sonoma ( <i>Alopecurus aequalis</i> var. <i>sonomensis</i> ) - Wherever found	E	\$8,020	\$1,267,329,539



Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1077	Petrel, Bermuda ( <i>Pterodroma cahow</i> ) - Wherever found	E	\$8,000	\$1,267,337,539
1077	Rocksail, interrupted (=Georgia) ( <i>Leptoxis foremani</i> ) - Wherever found	E	\$8,000	\$1,267,345,539
1077	Bladderpod, Zapata ( <i>Lesquerella thamnophila</i> ) - Wherever found	E	\$8,000	\$1,267,353,539
1077	Dogweed, ashy ( <i>Thymophylla tephroleuca</i> ) - Wherever found	E	\$8,000	\$1,267,361,539
1077	Whitlow-wort, papery ( <i>Paronychia chartacea</i> ) - Wherever found	T	\$8,000	\$1,267,369,539
1077	Pitcher-plant, green ( <i>Sarracenia oreophila</i> ) - Wherever found	E	\$8,000	\$1,267,377,539
1077	Trillium, persistent ( <i>Trillium persistens</i> ) - Wherever found	E	\$8,000	\$1,267,385,539
1077	No Common Name ( <i>Catesbaea melanocarpa</i> ) - Wherever found	E	\$8,000	\$1,267,393,539
1078	Haha ( <i>Cyanea humboldtiana</i> ) - Wherever found	E	\$7,989	\$1,267,401,528
1078	Ha`iwale ( <i>Cyrtandra kaulantha</i> ) - Wherever found	E	\$7,989	\$1,267,409,517
1078	No Common Name ( <i>Delissea undulata</i> ) - Wherever found	E	\$7,989	\$1,267,417,506
1078	Hau kuahiwi ( <i>Hibiscadelphus hualalaiensis</i> ) - Wherever found	E	\$7,989	\$1,267,425,495
1078	No Common Name ( <i>Cyperus fauriei</i> ) - Wherever found	E	\$7,989	\$1,267,433,484
1078	No Common Name ( <i>Silene perlmanii</i> ) - Wherever found	E	\$7,989	\$1,267,441,473
1078	A`e ( <i>Zanthoxylum dipetalum</i> var. <i>tomentosum</i> ) - Wherever found	E	\$7,989	\$1,267,449,462
1079	Ha`iwale ( <i>Cyrtandra tintinnabula</i> ) - Wherever found	E	\$7,980	\$1,267,457,442
1080	No Common Name ( <i>Amaranthus brownii</i> ) - Wherever found	E	\$7,952	\$1,267,465,394
1081	No Common Name ( <i>Tinospora homosepala</i> ) - Wherever found	E	\$7,936	\$1,267,473,330
1082	Alani ( <i>Melicope zahlbruckneri</i> ) - Wherever found	E	\$7,806	\$1,267,481,136
1083	Ha`iwale ( <i>Cyrtandra oenobarba</i> ) - Wherever found	E	\$7,736	\$1,267,488,872
1083	Haha ( <i>Cyanea rivularis</i> ) - Wherever found	E	\$7,736	\$1,267,496,608
1083	No Common Name ( <i>Phyllostegia knudsenii</i> ) - Wherever found	E	\$7,736	\$1,267,504,344
1083	Haha ( <i>Cyanea purpurellifolia</i> ) - Wherever found	E	\$7,736	\$1,267,512,080
1083	No Common Name ( <i>Stenogyne kaalae</i> ssp. <i>sherffii</i> ) - Wherever found	E	\$7,736	\$1,267,519,816
1083	"Akoko ( <i>Euphorbia halemanui</i> ) - Wherever found	E	\$7,736	\$1,267,527,552
1083	Ha`iwale ( <i>Cyrtandra polyantha</i> ) - Wherever found	E	\$7,736	\$1,267,535,288
1083	Koholapehu ( <i>Dubautia latifolia</i> ) - Wherever found	E	\$7,736	\$1,267,543,024
1083	Kio`ele ( <i>Kadua coriacea</i> ) - Wherever found	E	\$7,736	\$1,267,550,760
1083	No Common Name ( <i>Hesperomannia lydgatei</i> ) - Wherever found	E	\$7,736	\$1,267,558,496
1083	Kauai hau kuahiwi ( <i>Hibiscadelphus distans</i> ) - Wherever found	E	\$7,736	\$1,267,566,232
1083	Hibiscus, Clay's ( <i>Hibiscus clayi</i> ) - Wherever found	E	\$7,736	\$1,267,573,968
1083	Koki`o ( <i>Kokia kauaiensis</i> ) - Wherever found	E	\$7,736	\$1,267,581,704
1083	Alani ( <i>Melicope haupeensis</i> ) - Wherever found	E	\$7,736	\$1,267,589,440

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1083	Alani ( <i>Melicope reflexa</i> ) - Wherever found	E	\$7,736	\$1,267,597,176
1083	No Common Name ( <i>Polyscias racemosa</i> ) - Wherever found	E	\$7,736	\$1,267,604,912
1083	`Aiea ( <i>Nothocestrum peltatum</i> ) - Wherever found	E	\$7,736	\$1,267,612,648
1083	Nani wai`ale`ale ( <i>Viola kauaiensis</i> var. <i>wahiawaensis</i> ) - Wherever found	E	\$7,736	\$1,267,620,384
1083	Haha ( <i>Cyanea pinnatifida</i> ) - Wherever found	E	\$7,736	\$1,267,628,120
1083	Alani ( <i>Melicope hiiakae</i> ) - Wherever found	E	\$7,736	\$1,267,635,856
1083	No Common Name ( <i>Cyperus pennatifolius</i> ) - Wherever found	E	\$7,736	\$1,267,643,592
1083	Ma`oli`oli ( <i>Schiedea apokremnos</i> ) - Wherever found	E	\$7,736	\$1,267,651,328
1083	No Common Name ( <i>Schiedea helleri</i> ) - Wherever found	E	\$7,736	\$1,267,659,064
1083	Awiji ( <i>Schenkia sebaeoides</i> ) - Wherever found	E	\$7,736	\$1,267,666,800
1083	Haha ( <i>Cyanea recta</i> ) - Wherever found	T	\$7,736	\$1,267,674,536
1083	Haha ( <i>Cyanea undulata</i> ) - Wherever found	E	\$7,736	\$1,267,682,272
1083	Ha`iwale ( <i>Cyrtandra sessilis</i> ) - Wherever found	E	\$7,736	\$1,267,690,008
1083	Kamakahala ( <i>Labordia helleri</i> ) - Wherever found	E	\$7,736	\$1,267,697,744
1083	Kamakahala ( <i>Labordia pumila</i> ) - Wherever found	E	\$7,736	\$1,267,705,480
1083	No Common Name ( <i>Platydesma cornuta</i> var. <i>cornuta</i> ) - Wherever found	E	\$7,736	\$1,267,713,216
1083	Pilo kea lau li`i ( <i>Platydesma rostrata</i> ) - Wherever found	E	\$7,736	\$1,267,720,952
1083	Loulu ( <i>Pritchardia hardyi</i> ) - Wherever found	E	\$7,736	\$1,267,728,688
1083	Loulu ( <i>Pritchardia viscosa</i> ) - Wherever found	E	\$7,736	\$1,267,736,424
1083	Kopiko ( <i>Psychotria hobdyi</i> ) - Wherever found	E	\$7,736	\$1,267,744,160
1083	No Common Name ( <i>Stenogyne campanulata</i> ) - Wherever found	E	\$7,736	\$1,267,751,896
1083	Hau kuahiwi ( <i>Hibiscadelphus woodii</i> ) - Wherever found	E	\$7,736	\$1,267,759,632
1083	Kamakahala ( <i>Labordia tinifolia</i> var. <i>wahiawaensis</i> ) - Wherever found	E	\$7,736	\$1,267,767,368
1083	No Common Name ( <i>Lysimachia venosa</i> ) - Wherever found	E	\$7,736	\$1,267,775,104
1083	Kopiko ( <i>Psychotria hexandra</i> ssp. <i>oahuensis</i> ) - Wherever found	E	\$7,736	\$1,267,782,840
1083	Haha ( <i>Cyanea eleeleensis</i> ) - Wherever found	E	\$7,736	\$1,267,790,576
1083	`Akoko ( <i>Euphorbia eleanoriae</i> ) - Wherever found	E	\$7,736	\$1,267,798,312
1083	Haha ( <i>Cyanea dolichopoda</i> ) - Wherever found	E	\$7,736	\$1,267,806,048
1083	Haha ( <i>Cyanea kolekoleensis</i> ) - Wherever found	E	\$7,736	\$1,267,813,784
1083	Naenae ( <i>Dubautia kenwoodii</i> ) - Wherever found	E	\$7,736	\$1,267,821,520
1083	No Common Name ( <i>Lysimachia pendens</i> ) - Wherever found	E	\$7,736	\$1,267,829,256
1083	No Common Name ( <i>Lysimachia scopulensis</i> ) - Wherever found	E	\$7,736	\$1,267,836,992
1083	No Common Name ( <i>Phyllostegia renovans</i> ) - Wherever found	E	\$7,736	\$1,267,844,728

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1083	No Common Name ( <i>Polyscias flynnii</i> ) - Wherever found	E	\$7,736	\$1,267,852,464
1083	No Common Name ( <i>Asplenium dielpallidum</i> ) - Wherever found	E	\$7,736	\$1,267,860,200
1083	Hohiu ( <i>Dryopteris glabra</i> var. <i>pusilla</i> ) - Wherever found	E	\$7,736	\$1,267,867,936
1084	No Common Name ( <i>Gouania vitifolia</i> ) - Wherever found	E	\$7,634	\$1,267,875,570
1085	No Common Name ( <i>Neraudia angulata</i> ) - Wherever found	E	\$7,546	\$1,267,883,116
1086	Catspaw, white (pearlymussel) ( <i>Epioblasma obliquata perobliqua</i> ) - Wherever found	E	\$7,512	\$1,267,890,628
1087	Salamander, Austin blind ( <i>Eurycea waterlooensis</i> ) - Wherever found	E	\$7,502	\$1,267,898,130
1088	Anole, Culebra Island giant ( <i>Anolis roosevelti</i> ) - Wherever found	E	\$7,500	\$1,267,905,630
1088	Pupfish, Comanche Springs ( <i>Cyprinodon elegans</i> ) - Wherever found	E	\$7,500	\$1,267,913,130
1088	Palo de nigua ( <i>Cornutia obovata</i> ) - Wherever found	E	\$7,500	\$1,267,920,630
1088	Bariaco ( <i>Trichilia triacantha</i> ) - Wherever found	E	\$7,500	\$1,267,928,130
1088	No Common Name ( <i>Thelypteris inabonensis</i> ) - Wherever found	E	\$7,500	\$1,267,935,630
1089	Fly, Hawaiian picture-wing ( <i>Drosophila heteroneura</i> ) - Wherever found	E	\$7,433	\$1,267,943,063
1090	Snake, copperbelly water ( <i>Nerodia erythrogaster neglecta</i> ) - Indiana north of 40 degrees north latitude, Michigan, Ohio	T	\$7,432	\$1,267,950,495
1091	Bean, purple ( <i>Villosa perpurpurea</i> ) - Wherever found	E	\$7,416	\$1,267,957,911
1092	Manzanita, Presidio ( <i>Arctostaphylos hookeri</i> var. <i>ravenii</i> ) - Wherever found	E	\$7,400	\$1,267,965,311
1092	Rosemary, short-leaved ( <i>Conradina brevifolia</i> ) - Wherever found	E	\$7,400	\$1,267,972,711
1092	Beardtongue, Penland ( <i>Penstemon penlandii</i> ) - Wherever found	E	\$7,400	\$1,267,980,111
1092	Cactus, Florida semaphore ( <i>Consolea corallicola</i> ) - Wherever found	E	\$7,400	\$1,267,987,511
1093	`Anaunau ( <i>Lepidium arbuscula</i> ) - Wherever found	E	\$7,394	\$1,267,994,905
1094	Gardenia (=Na`u), Hawaiian ( <i>Gardenia brighamii</i> ) - Wherever found	E	\$7,319	\$1,268,002,224
1095	Erubia ( <i>Solanum drymophilum</i> ) - Wherever found	E	\$7,300	\$1,268,009,524
1096	Damselfly, blackline Hawaiian ( <i>Megalagrion nigrohamatum nigrolineatum</i> ) - Wherever found	E	\$7,270	\$1,268,016,794
1097	Alani ( <i>Melicope saint-johnii</i> ) - Wherever found	E	\$7,209	\$1,268,024,003
1097	Nehe ( <i>Lipochaeta lobata</i> var. <i>leptophylla</i> ) - Wherever found	E	\$7,209	\$1,268,031,212
1098	Chub, Yaqui ( <i>Gila purpurea</i> ) - Wherever found	E	\$7,200	\$1,268,038,412
1099	Rabbitsfoot, rough ( <i>Quadrula cylindrica strigillata</i> ) - Wherever found	E	\$7,106	\$1,268,045,518
1100	Warbler (=wood), Bachman's ( <i>Vermivora bachmanii</i> ) - Wherever found	E	\$7,100	\$1,268,052,618
1100	Cactus, Tobusch fishhook ( <i>Sclerocactus brevihamatus</i> ssp. <i>tobuschii</i> ) - Wherever found	T	\$7,100	\$1,268,059,718
1101	Shiner, blue ( <i>Cyprinella caerulea</i> ) - Wherever found	T	\$7,000	\$1,268,066,718
1101	Springsnail, Three Forks ( <i>Pyrgulopsis trivialis</i> ) - Wherever found	E	\$7,000	\$1,268,073,718
1101	No Common Name ( <i>Varronia rupicola</i> ) - Wherever found	T	\$7,000	\$1,268,080,718

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1101	Palo de rosa ( <i>Ottoschulzia rhodoxylon</i> ) - Wherever found	E	\$7,000	\$1,268,087,718
1102	Cypress, Gowen ( <i>Cupressus goveniana</i> ssp. <i>goveniana</i> ) - Wherever found	T	\$6,832	\$1,268,094,550
1103	Rockcress, Georgia ( <i>Arabis georgiana</i> ) - Wherever found	T	\$6,826	\$1,268,101,376
1104	Stickseed, showy ( <i>Hackelia venusta</i> ) - Wherever found	E	\$6,810	\$1,268,108,186
1105	Wawae`iole ( <i>Huperzia nutans</i> ) - Wherever found	E	\$6,764	\$1,268,114,950
1106	Barberry, Nevin's ( <i>Berberis nevinii</i> ) - Wherever found	E	\$6,700	\$1,268,121,650
1107	Fly, Hawaiian picture-wing ( <i>Drosophila sharpi</i> ) - Wherever found	E	\$6,694	\$1,268,128,344
1108	Damselfly, flying earwig Hawaiian ( <i>Megalagrion nesiotes</i> ) - Wherever found	E	\$6,692	\$1,268,135,036
1109	Aupaka ( <i>Isodendrion hosakae</i> ) - Wherever found	E	\$6,665	\$1,268,141,701
1110	Fly, Hawaiian picture-wing ( <i>Drosophila musaphilia</i> ) - Wherever found	E	\$6,615	\$1,268,148,316
1111	No Common Name ( <i>Phyllostegia stachyoides</i> ) - Wherever found	E	\$6,606	\$1,268,154,922
1112	Ha`iwale ( <i>Cyrtandra oxybapha</i> ) - Wherever found	E	\$6,573	\$1,268,161,495
1113	Bush-mallow, Santa Cruz Island ( <i>Malacothamnus fasciculatus</i> var. <i>nesioticus</i> ) - Wherever found	E	\$6,534	\$1,268,168,029
1114	Ma`o hau hele, (=native yellow hibiscus) ( <i>Hibiscus brackenridgei</i> ) - Wherever found	E	\$6,507	\$1,268,174,536
1115	Guajon ( <i>Eleutherodactylus cooki</i> ) - Wherever found	T	\$6,500	\$1,268,181,036
1115	Palo de ramon ( <i>Banara vanderbiltii</i> ) - Wherever found	E	\$6,500	\$1,268,187,536
1115	Chupacallos ( <i>Pleodendron macranthum</i> ) - Wherever found	E	\$6,500	\$1,268,194,036
1115	Palo colorado ( <i>Ternstroemia luquillensis</i> ) - Wherever found	E	\$6,500	\$1,268,200,536
1115	Palo de jazmin ( <i>Styrax portoricensis</i> ) - Wherever found	E	\$6,500	\$1,268,207,036
1115	No Common Name ( <i>Gesneria pauciflora</i> ) - Wherever found	T	\$6,500	\$1,268,213,536
1115	No Common Name ( <i>Schoepfia arenaria</i> ) - Wherever found	T	\$6,500	\$1,268,220,036
1115	Fern, Elfin tree ( <i>Cyathea dryopteroides</i> ) - Wherever found	E	\$6,500	\$1,268,226,536
1116	`Ahinahina ( <i>Argyroxiphium sandwicense</i> ssp. <i>sandwicense</i> ) - Wherever found	E	\$6,480	\$1,268,233,016
1117	No Common Name ( <i>Platydesma remyi</i> ) - Wherever found	E	\$6,443	\$1,268,239,459
1117	No Common Name ( <i>Festuca hawaiiensis</i> ) - Wherever found	E	\$6,443	\$1,268,245,902
1118	No Common Name ( <i>Remya montgomeryi</i> ) - Wherever found	E	\$6,431	\$1,268,252,333
1119	Yellow-faced bee, hilaris ( <i>Hylaeus hilaris</i> ) - Wherever found	E	\$6,419	\$1,268,258,752
1119	Fly, Hawaiian picture-wing ( <i>Drosophila digressa</i> ) - Wherever found	E	\$6,419	\$1,268,265,171
1119	Fly, Hawaiian picture-wing ( <i>Drosophila ochrobasis</i> ) - Wherever found	E	\$6,419	\$1,268,271,590
1120	`Akoko ( <i>Euphorbia celastroides</i> var. <i>kaenana</i> ) - Wherever found	E	\$6,412	\$1,268,278,002
1121	Damselfly, crimson Hawaiian ( <i>Megalagrion leptodemas</i> ) - Wherever found	E	\$6,409	\$1,268,284,411
1122	Silverside, Waccamaw ( <i>Menidia extensa</i> ) - Wherever found	T	\$6,400	\$1,268,290,811

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1122	Jacquemontia, beach ( <i>Jacquemontia reclinata</i> ) - Wherever found	E	\$6,400	\$1,268,297,211
1123	Pitcher-plant, Alabama canebrake ( <i>Sarracenia rubra</i> ssp. <i>alabamensis</i> ) - Wherever found	E	\$6,399	\$1,268,303,610
1124	Sunburst, Hartweg's golden ( <i>Pseudobahia bahiifolia</i> ) - Wherever found	E	\$6,364	\$1,268,309,974
1125	Fly, Hawaiian picture-wing ( <i>Drosophila aglaia</i> ) - Wherever found	E	\$6,320	\$1,268,316,294
1125	Picture-wing fly, Hawaiian ( <i>Drosophila montgomeryi</i> ) - Wherever found	E	\$6,320	\$1,268,322,614
1125	Fly, Hawaiian picture-wing ( <i>Drosophila obatai</i> ) - Wherever found	E	\$6,320	\$1,268,328,934
1125	Fly, Hawaiian picture-wing ( <i>Drosophila substenoptera</i> ) - Wherever found	E	\$6,320	\$1,268,335,254
1125	Fly, Hawaiian picture-wing ( <i>Drosophila tarphytrichia</i> ) - Wherever found	E	\$6,320	\$1,268,341,574
1125	Yellow-faced bee, Hawaiian ( <i>Hylaeus mana</i> ) - Wherever found	E	\$6,320	\$1,268,347,894
1125	Yellow-faced bee, Hawaiian ( <i>Hylaeus kuakea</i> ) - Wherever found	E	\$6,320	\$1,268,354,214
1126	Polygala, tiny ( <i>Polygala smallii</i> ) - Wherever found	E	\$6,300	\$1,268,360,514
1126	Mint, Garrett's ( <i>Dicerandra christmanii</i> ) - Wherever found	E	\$6,300	\$1,268,366,814
1127	Damselfly, oceanic Hawaiian ( <i>Megalagrion oceanicum</i> ) - Wherever found	E	\$6,256	\$1,268,373,070
1128	`Ena`ena ( <i>Pseudognaphalium sandwicensium</i> var. <i>molokaiense</i> ) - Wherever found	E	\$6,201	\$1,268,379,271
1129	Loulu ( <i>Pritchardia maideniana</i> ) - Wherever found	E	\$6,170	\$1,268,385,441
1130	Waterdog, black warrior (=Sipsey Fork) ( <i>Necturus alabamensis</i> ) - Wherever found	E	\$6,137	\$1,268,391,578
1131	No Common Name ( <i>Cyanea kauaulaensis</i> ) - Wherever found	E	\$6,106	\$1,268,397,684
1132	Bluet, Roan Mountain ( <i>Hedyotis purpurea</i> var. <i>montana</i> ) - Wherever found	E	\$6,100	\$1,268,403,784
1132	Prairie-clover, Florida ( <i>Dalea carthagenensis floridana</i> ) - Wherever found	E	\$6,100	\$1,268,409,884
1133	Kuahiwi laukahi ( <i>Plantago hawaiiensis</i> ) - Wherever found	E	\$6,098	\$1,268,415,982
1134	Sedge, Navajo ( <i>Carex specuicola</i> ) - Wherever found	T	\$6,010	\$1,268,421,992
1135	Bellflower, Brooksville ( <i>Campanula robinsiae</i> ) - Wherever found	E	\$6,000	\$1,268,427,992
1135	Cactus, Brady pincushion ( <i>Pediocactus bradyi</i> ) - Wherever found	E	\$6,000	\$1,268,433,992
1135	Capa rosa ( <i>Callicarpa ampla</i> ) - Wherever found	E	\$6,000	\$1,268,439,992
1135	Holly, Cook's ( <i>Ilex cookii</i> ) - Wherever found	E	\$6,000	\$1,268,445,992
1135	Prickly-ash, St. Thomas ( <i>Zanthoxylum thomasianum</i> ) - Wherever found	E	\$6,000	\$1,268,451,992
1135	No Common Name ( <i>Ilex sintenisii</i> ) - Wherever found	E	\$6,000	\$1,268,457,992
1135	No Common Name ( <i>Elaphoglossum serpens</i> ) - Wherever found	E	\$6,000	\$1,268,463,992
1135	No Common Name ( <i>Tectaria estremerana</i> ) - Wherever found	E	\$6,000	\$1,268,469,992
1135	No Common Name ( <i>Thelypteris verecunda</i> ) - Wherever found	E	\$6,000	\$1,268,475,992
1136	Barbara's Buttons, Mohr's ( <i>Marshallia mohrii</i> ) - Wherever found	T	\$5,977	\$1,268,481,969
1137	Olua ( <i>Hypolepis hawaiiensis</i> var. <i>mauiensis</i> ) - Wherever found	E	\$5,954	\$1,268,487,923
1138	Darter, bluemask ( <i>Etheostoma akatulo</i> ) - Wherever found	E	\$5,926	\$1,268,493,849

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1139	Glade cress, Kentucky ( <i>Leavenworthia exigua laciniata</i> ) - Wherever found	T	\$5,920	\$1,268,499,769
1140	Spineflower, Robust ( <i>Chorizanthe robusta</i> var. <i>robusta</i> ) - Wherever found	E	\$5,900	\$1,268,505,669
1141	No Common Name ( <i>Wikstroemia skottsbergiana</i> ) -	E	\$5,853	\$1,268,511,522
1141	No Common Name ( <i>Cyperus neokunthianus</i> ) - Wherever found	E	\$5,853	\$1,268,517,375
1141	No Common Name ( <i>Schiedea jacobii</i> ) - Wherever found	E	\$5,853	\$1,268,523,228
1141	No Common Name ( <i>Lepidium orbiculare</i> ) - Wherever found	E	\$5,853	\$1,268,529,081
1141	No Common Name ( <i>Santalum involutum</i> ) - Wherever found	E	\$5,853	\$1,268,534,934
1141	No Common Name ( <i>Labordia lorenciana</i> ) - Wherever found	E	\$5,853	\$1,268,540,787
1141	No Common Name ( <i>Asplenium diellaciniatum</i> ) - Wherever found	E	\$5,853	\$1,268,546,640
1142	Pearlshell, Alabama ( <i>Margaritifera marrianae</i> ) - Wherever found	E	\$5,790	\$1,268,552,430
1143	Darter, diamond ( <i>Crystallaria cincotta</i> ) - Wherever found	E	\$5,774	\$1,268,558,204
1144	Ha`iwale ( <i>Cyrtandra dentata</i> ) - Wherever found	E	\$5,754	\$1,268,563,958
1144	Kaulu ( <i>Pteralyxia macrocarpa</i> ) - Wherever found	E	\$5,754	\$1,268,569,712
1144	No Common Name ( <i>Schiedea trinervis</i> ) - Wherever found	E	\$5,754	\$1,268,575,466
1144	Ha`iwale ( <i>Cyrtandra subumbellata</i> ) - Wherever found	E	\$5,754	\$1,268,581,220
1144	No Common Name ( <i>Kadua degeneri</i> ) - Wherever found	E	\$5,754	\$1,268,586,974
1144	No Common Name ( <i>Hesperomannia arbuscula</i> ) - Wherever found	E	\$5,754	\$1,268,592,728
1144	No Common Name ( <i>Lobelia oahuensis</i> ) - Wherever found	E	\$5,754	\$1,268,598,482
1144	No Common Name ( <i>Viola oahuensis</i> ) - Wherever found	E	\$5,754	\$1,268,604,236
1144	Kamakahala ( <i>Labordia cyrtandrae</i> ) - Wherever found	E	\$5,754	\$1,268,609,990
1144	Na`ena`e ( <i>Dubautia herbstobatae</i> ) - Wherever found	E	\$5,754	\$1,268,615,744
1144	Loulu ( <i>Pritchardia kaalae</i> ) - Wherever found	E	\$5,754	\$1,268,621,498
1144	Ha`iwale ( <i>Cyrtandra viridiflora</i> ) - Wherever found	E	\$5,754	\$1,268,627,252
1144	Kolea ( <i>Myrsine juddii</i> ) - Wherever found	E	\$5,754	\$1,268,633,006
1144	`Akoko ( <i>Euphorbia herbstii</i> ) - Wherever found	E	\$5,754	\$1,268,638,760
1144	`Akoko ( <i>Euphorbia rockii</i> ) - Wherever found	E	\$5,754	\$1,268,644,514
1144	No Common Name ( <i>Pteris lidgatei</i> ) - Wherever found	E	\$5,754	\$1,268,650,268
1145	Kolea ( <i>Myrsine fosbergii</i> ) - Wherever found	E	\$5,701	\$1,268,655,969
1145	No Common Name ( <i>Deparia kaalaana</i> ) - Wherever found	E	\$5,701	\$1,268,661,670
1146	Spurge, deltoid ( <i>Chamaesyce deltoidea</i> ssp. <i>deltoidea</i> ) - Wherever found	E	\$5,700	\$1,268,667,370
1147	Schiedea, Diamond Head ( <i>Schiedea adamantis</i> ) - Wherever found	E	\$5,670	\$1,268,673,040
1147	Loulu ( <i>Pritchardia schattaueri</i> ) - Wherever found	E	\$5,670	\$1,268,678,710
1147	Haha ( <i>Cyanea remyi</i> ) - Wherever found	E	\$5,670	\$1,268,684,380

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1148	No Common Name ( <i>Gouania meyenii</i> ) - Wherever found	E	\$5,652	\$1,268,690,032
1149	No Common Name ( <i>Hesperomannia arborescens</i> ) - Wherever found	E	\$5,602	\$1,268,695,634
1149	Opuhe ( <i>Urera kaalae</i> ) - Wherever found	E	\$5,602	\$1,268,701,236
1149	Heau ( <i>Exocarpos luteolus</i> ) - Wherever found	E	\$5,602	\$1,268,706,838
1149	Nioi ( <i>Eugenia koolauensis</i> ) - Wherever found	E	\$5,602	\$1,268,712,440
1149	No Common Name ( <i>Sanicula mariversa</i> ) - Wherever found	E	\$5,602	\$1,268,718,042
1150	`Ala `ala wai nui ( <i>Peperomia subpetiolata</i> ) - Wherever found	E	\$5,599	\$1,268,723,641
1151	Shiner, palezone ( <i>Notropis albizonatus</i> ) - Wherever found	E	\$5,500	\$1,268,729,141
1151	Fescue, Guadalupe ( <i>Festuca ligulata</i> ) - Wherever found	E	\$5,500	\$1,268,734,641
1151	Buckwheat, cushenbury ( <i>Eriogonum ovalifolium</i> var. <i>vineum</i> ) - Wherever found	E	\$5,500	\$1,268,740,141
1151	No Common Name ( <i>Calyptanthus thomasiana</i> ) - Wherever found	E	\$5,500	\$1,268,745,641
1151	Manaca, palma de ( <i>Calyptronoma rivalis</i> ) - Wherever found	T	\$5,500	\$1,268,751,141
1151	No Common Name ( <i>Ternstroemia subsessilis</i> ) - Wherever found	E	\$5,500	\$1,268,756,641
1151	No Common Name ( <i>Cranichis ricartii</i> ) - Wherever found	E	\$5,500	\$1,268,762,141
1151	No Common Name ( <i>Adiantum vivesii</i> ) - Wherever found	E	\$5,500	\$1,268,767,641
1152	Haha ( <i>Cyanea lanceolata</i> ) - Wherever found	E	\$5,417	\$1,268,773,058
1152	Na`ena`e ( <i>Dubautia waialealae</i> ) - Wherever found	E	\$5,417	\$1,268,778,475
1152	Liliwai ( <i>Acaena exigua</i> ) - Wherever found	E	\$5,417	\$1,268,783,892
1152	`Awikiwiki ( <i>Canavalia napaliensis</i> ) - Wherever found	E	\$5,417	\$1,268,789,309
1152	`Akoko ( <i>Euphorbia deppeana</i> ) - Wherever found	E	\$5,417	\$1,268,794,726
1152	`Akoko ( <i>Euphorbia remyi</i> var. <i>kauaiensis</i> ) - Wherever found	E	\$5,417	\$1,268,800,143
1152	No Common Name ( <i>Kadua st.-johnii</i> ) - Wherever found	E	\$5,417	\$1,268,805,560
1152	Alani ( <i>Melicope degeneri</i> ) - Wherever found	E	\$5,417	\$1,268,810,977
1152	Alani ( <i>Melicope pallida</i> ) - Wherever found	E	\$5,417	\$1,268,816,394
1152	Alani ( <i>Melicope paniculata</i> ) - Wherever found	E	\$5,417	\$1,268,821,811
1152	Bluegrass, Hawaiian ( <i>Poa sandvicensis</i> ) - Wherever found	E	\$5,417	\$1,268,827,228
1152	Kaulu ( <i>Pteralyxia kauaiensis</i> ) - Wherever found	E	\$5,417	\$1,268,832,645
1152	No Common Name ( <i>Polyscias lydgatei</i> ) - Wherever found	E	\$5,417	\$1,268,838,062
1152	No Common Name ( <i>Viola helenae</i> ) - Wherever found	E	\$5,417	\$1,268,843,479
1152	Ha`iwale ( <i>Cyrtandra crenata</i> ) - Wherever found	E	\$5,417	\$1,268,848,896
1152	Nehe ( <i>Lipochaeta micrantha</i> ) - Wherever found	E	\$5,417	\$1,268,854,313
1152	Nehe ( <i>Lipochaeta waimeaensis</i> ) - Wherever found	E	\$5,417	\$1,268,859,730
1152	No Common Name ( <i>Lobelia monostachya</i> ) - Wherever found	E	\$5,417	\$1,268,865,147

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1152	No Common Name ( <i>Lysimachia filifolia</i> ) - Wherever found	E	\$5,417	\$1,268,870,564
1152	<i>Kolea</i> ( <i>Myrsine mezii</i> ) - Wherever found	E	\$5,417	\$1,268,875,981
1152	No Common Name ( <i>Poa siphonoglossa</i> ) - Wherever found	E	\$5,417	\$1,268,881,398
1152	No Common Name ( <i>Xylosma crenatum</i> ) - Wherever found	E	\$5,417	\$1,268,886,815
1152	Holei ( <i>Ochrosia kilauaeaensis</i> ) - Wherever found	E	\$5,417	\$1,268,892,232
1152	No Common Name ( <i>Schiedea spergulina</i> var. <i>leiopoda</i> ) - Wherever found	E	\$5,417	\$1,268,897,649
1152	No Common Name ( <i>Schiedea viscosa</i> ) - Wherever found	E	\$5,417	\$1,268,903,066
1152	Haha ( <i>Cyanea asarifolia</i> ) - Wherever found	E	\$5,417	\$1,268,908,483
1152	Haha ( <i>Cyanea copelandii</i> ssp. <i>copelandii</i> ) - Wherever found	E	\$5,417	\$1,268,913,900
1152	Haha ( <i>Cyanea truncata</i> ) - Wherever found	E	\$5,417	\$1,268,919,317
1152	Mapele ( <i>Cyrtandra cyaneoides</i> ) - Wherever found	E	\$5,417	\$1,268,924,734
1152	Na'ena'e ( <i>Dubautia imbricata</i> ssp. <i>imbricata</i> ) - Wherever found	E	\$5,417	\$1,268,930,151
1152	Na'ena'e ( <i>Dubautia pauciflorula</i> ) - Wherever found	E	\$5,417	\$1,268,935,568
1152	No Common Name ( <i>Phyllostegia waimeae</i> ) - Wherever found	E	\$5,417	\$1,268,940,985
1152	Pa'iniu ( <i>Astelia waialealae</i> ) - Wherever found	E	\$5,417	\$1,268,946,402
1152	Haha ( <i>Cyanea kuhihewa</i> ) - Wherever found	E	\$5,417	\$1,268,951,819
1152	Na'ena'e ( <i>Dubautia plantaginea</i> ssp. <i>magnifolia</i> ) - Wherever found	E	\$5,417	\$1,268,957,236
1152	No Common Name ( <i>Schiedea attenuata</i> ) - Wherever found	E	\$5,417	\$1,268,962,653
1152	A'e ( <i>Zanthoxylum oahuense</i> ) - Wherever found	E	\$5,417	\$1,268,968,070
1152	Naenae ( <i>Dubautia kalalauensis</i> ) - Wherever found	E	\$5,417	\$1,268,973,487
1152	No Common Name ( <i>Lysimachia iniki</i> ) - Wherever found	E	\$5,417	\$1,268,978,904
1152	<i>Kolea</i> ( <i>Myrsine knudsenii</i> ) - Wherever found	E	\$5,417	\$1,268,984,321
1152	Loulu ( <i>Pritchardia bakeri</i> ) - Wherever found	E	\$5,417	\$1,268,989,738
1152	No Common Name ( <i>Doryopteris takeuchii</i> ) - Wherever found	E	\$5,417	\$1,268,995,155
1153	Darter, Cherokee ( <i>Etheostoma scotti</i> ) - Wherever found	T	\$5,330	\$1,269,000,485
1154	Logperch, Conasauga ( <i>Percina jenkinsi</i> ) - Wherever found	E	\$5,300	\$1,269,005,785
1154	Milkpea, Small's ( <i>Galactia smallii</i> ) - Wherever found	E	\$5,300	\$1,269,011,085
1155	'Oha wai ( <i>Clermontia drepanomorpha</i> ) - Wherever found	E	\$5,265	\$1,269,016,350
1155	Kopiko ( <i>Psychotria grandiflora</i> ) - Wherever found	E	\$5,265	\$1,269,021,615
1155	Olulu ( <i>Brighamia insignis</i> ) - Wherever found	E	\$5,265	\$1,269,026,880
1155	Nehe ( <i>Lipochaeta fauriei</i> ) - Wherever found	E	\$5,265	\$1,269,032,145
1155	Aumakua, Palapalai ( <i>Dryopteris crinalis</i> var. <i>podosorus</i> ) - Wherever found	E	\$5,265	\$1,269,037,410
1156	Dace, Laurel ( <i>Chrosomus saylora</i> ) - Wherever found	E	\$5,250	\$1,269,042,660



Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1157	Fern, pendant kiki ( <i>Adenophorus periens</i> ) - Wherever found	E	\$5,244	\$1,269,047,904
1158	Bluegrass, Mann's ( <i>Poa mannii</i> ) - Wherever found	E	\$5,143	\$1,269,053,047
1159	Bluegrass, San Bernardino ( <i>Poa atropurpurea</i> ) - Wherever found	E	\$5,100	\$1,269,058,147
1159	Blazingstar, scrub ( <i>Liatris ohlingerae</i> ) - Wherever found	E	\$5,100	\$1,269,063,247
1159	Meadowfoam, Butte County ( <i>Limnanthes floccosa</i> ssp. <i>californica</i> ) - Wherever found	E	\$5,100	\$1,269,068,347
1160	Mucket, orangenacre ( <i>Lampsilis perovalis</i> ) - Wherever found	T	\$5,076	\$1,269,073,423
1161	Pigtoe, rough ( <i>Pleurobema plenum</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$5,006	\$1,269,078,429
1162	Salamander, Georgetown ( <i>Eurycea naufragia</i> ) - Wherever found	T	\$5,002	\$1,269,083,431
1163	Iguana, Mona ground ( <i>Cyclura stejnegeri</i> ) - Wherever found	T	\$5,000	\$1,269,088,431
1163	Gecko, Monito ( <i>Sphaerodactylus micropithecus</i> ) - Wherever found	E	\$5,000	\$1,269,093,431
1163	Snake, copperbelly water ( <i>Nerodia erythrogaster neglecta</i> ) - Illinois, Kentucky, and Indiana south of 40 degrees north latitude	RT	\$5,000	\$1,269,098,431
1163	Gambusia, San Marcos ( <i>Gambusia georgei</i> ) - Wherever found	E	\$5,000	\$1,269,103,431
1163	Pimpleback, orangefoot (pearlymussel) ( <i>Plethobasus cooperianus</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$5,000	\$1,269,108,431
1163	Beetle, [no common name] ( <i>Rhadine infernalis</i> ) - Wherever found	E	\$5,000	\$1,269,113,431
1163	Beetle, [no common name] ( <i>Rhadine exilis</i> ) - Wherever found	E	\$5,000	\$1,269,118,431
1163	Crayfish, Benton County cave ( <i>Cambarus aculabrum</i> ) - Wherever found	E	\$5,000	\$1,269,123,431
1163	Thistle, Sacramento Mountains ( <i>Cirsium vinaceum</i> ) - Wherever found	T	\$5,000	\$1,269,128,431
1163	Peperomia, Wheeler's ( <i>Peperomia wheeleri</i> ) - Wherever found	E	\$5,000	\$1,269,133,431
1163	No Common Name ( <i>Leptocereus grantianus</i> ) - Wherever found	E	\$5,000	\$1,269,138,431
1163	No Common Name ( <i>Polystichum calderonense</i> ) - Wherever found	E	\$5,000	\$1,269,143,431
1163	No Common Name ( <i>Thelypteris yaucoensis</i> ) - Wherever found	E	\$5,000	\$1,269,148,431
1164	Monkeyface, Appalachian (pearlymussel) ( <i>Quadrula sparsa</i> ) - USA (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$4,925	\$1,269,153,356
1165	No Common Name ( <i>Keysseria</i> (=Lagenifera) <i>erici</i> ) - Wherever found	E	\$4,907	\$1,269,158,263
1166	Skipper, Pawnee montane ( <i>Hesperia leonardus montana</i> ) - Wherever found	T	\$4,810	\$1,269,163,073
1167	Reed-mustard, shrubby ( <i>Schoenocrambe suffrutescens</i> ) - Wherever found	E	\$4,712	\$1,269,167,785
1168	Daisy, Lakeside ( <i>Hymenoxys herbacea</i> ) - Wherever found	T	\$4,676	\$1,269,172,461
1169	No Common Name ( <i>Trematolobelia singularis</i> ) - Wherever found	E	\$4,626	\$1,269,177,087
1170	Leopard ( <i>Panthera pardus</i> ) - except where threatened	E	\$4,620	\$1,269,181,707
1171	Seablite, California ( <i>Suaeda californica</i> ) -	E	\$4,610	\$1,269,186,317

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1172	Pigtoe, Georgia ( <i>Pleurobema hanleyianum</i> ) - Wherever found	E	\$4,500	\$1,269,190,817
1172	Skullcap, Florida ( <i>Scutellaria floridana</i> ) - Wherever found	T	\$4,500	\$1,269,195,317
1172	Walnut (=Nogal), West Indian ( <i>Juglans jamaicensis</i> ) - Wherever found	E	\$4,500	\$1,269,199,817
1173	Townsendia, Last Chance ( <i>Townsendia aprica</i> ) - Wherever found	T	\$4,450	\$1,269,204,267
1174	Combshell, upland ( <i>Epioblasma metastrata</i> ) - Wherever found	E	\$4,396	\$1,269,208,663
1175	Bedstraw, island ( <i>Galium buxifolium</i> ) - Wherever found	E	\$4,356	\$1,269,213,019
1176	Irisette, white ( <i>Sisyrinchium dichotomum</i> ) - Wherever found	E	\$4,300	\$1,269,217,319
1177	Riversnail, Anthony's ( <i>Athearnia anthonyi</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$4,254	\$1,269,221,573
1178	Birds-in-a-nest, white ( <i>Macbridea alba</i> ) - Wherever found	T	\$4,200	\$1,269,225,773
1178	Fringe-tree, pygmy ( <i>Chionanthus pygmaeus</i> ) - Wherever found	E	\$4,200	\$1,269,229,973
1179	Moccasinshell, Coosa ( <i>Medionidus parvulus</i> ) - Wherever found	E	\$4,120	\$1,269,234,093
1180	Ragwort, San Francisco Peaks ( <i>Packera franciscana</i> ) - Wherever found	T	\$4,100	\$1,269,238,193
1181	Bear, brown ( <i>Ursus arctos pruinosus</i> ) - Wherever found	E	\$4,073	\$1,269,242,266
1182	Seal, Ringed ( <i>Phoca (=Pusa) hispida botnica</i> ) - Wherever found	T	\$4,000	\$1,269,246,266
1182	Milk-vetch, heliotrope ( <i>Astragalus montii</i> ) - Wherever found	T	\$4,000	\$1,269,250,266
1182	Cactus, Cochise pincushion ( <i>Coryphantha robbinsorum</i> ) - Wherever found	T	\$4,000	\$1,269,254,266
1182	No Common Name ( <i>Auerodendron pauciflorum</i> ) - Wherever found	E	\$4,000	\$1,269,258,266
1183	Campeloma, slender ( <i>Campeloma decampi</i> ) - Wherever found	E	\$3,997	\$1,269,262,263
1184	Liveforever, Santa Barbara Island ( <i>Dudleya traskiae</i> ) - Wherever found	E	\$3,839	\$1,269,266,102
1185	Pigtoe, finerayed ( <i>Fusconaia cuneolus</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$3,808	\$1,269,269,910
1186	Reedgrass, Hillebrand's ( <i>Calamagrostis hillebrandii</i> ) - Wherever found	E	\$3,744	\$1,269,273,654
1187	Bladderpod, Spring Creek ( <i>Lesquerella perforata</i> ) - Wherever found	E	\$3,660	\$1,269,277,314
1188	Ground-plum, Guthrie's (=Pyne's) ( <i>Astragalus bibullatus</i> ) - Wherever found	E	\$3,650	\$1,269,280,964
1189	Lilliput, pale (pearlymussel) ( <i>Toxolasma cylindrellus</i> ) - Wherever found	E	\$3,600	\$1,269,284,564
1189	Thistle, Chorro Creek bog ( <i>Cirsium fontinale</i> var. <i>obispoense</i> ) - Wherever found	E	\$3,600	\$1,269,288,164
1190	Sculpin, pygmy ( <i>Cottus paulus</i> (=pygmaeus)) - Wherever found	T	\$3,500	\$1,269,291,664
1190	Tiger beetle, Miami ( <i>Cicindelidia floridana</i> ) - Wherever found	E	\$3,500	\$1,269,295,164
1190	No Common Name ( <i>Daphnopsis helleriana</i> ) - Wherever found	E	\$3,500	\$1,269,298,664
1191	Silverbush, Blodgett's ( <i>Argythamnia blodgettii</i> ) - Wherever found	T	\$3,442	\$1,269,302,106
1192	Pearlymussel, dromedary ( <i>Dromus dromas</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$3,382	\$1,269,305,488

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1193	Fanshell ( <i>Cyprogenia stegaria</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$3,379	\$1,269,308,867
1194	Reed-mustard, clay ( <i>Schoenocrambe argillacea</i> ) - Wherever found	T	\$3,369	\$1,269,312,236
1195	Pearlymussel, Curtis ( <i>Epioblasma florentina curtisii</i> ) - Wherever found	E	\$3,366	\$1,269,315,602
1196	Wartyback, white (pearlymussel) ( <i>Plethobasus cicatricosus</i> ) - Wherever found	E	\$3,351	\$1,269,318,953
1197	Fox, Santa Catalina Island ( <i>Urocyon littoralis catalinae</i> ) - Wherever found	T	\$3,350	\$1,269,322,303
1198	Ambersnail, Kanab ( <i>Oxyloma haydeni kanabensis</i> ) - Wherever found	E	\$3,333	\$1,269,325,636
1199	Turtle, yellow-blotched map ( <i>Graptemys flavimaculata</i> ) - Wherever found	T	\$3,313	\$1,269,328,949
1200	Shiner, Cahaba ( <i>Notropis cahabae</i> ) - Wherever found	E	\$3,305	\$1,269,332,254
1201	Pocketbook, speckled ( <i>Lampsilis streckeri</i> ) - Wherever found	E	\$3,300	\$1,269,335,554
1202	Pigtoe, Cumberland ( <i>Pleurobema gibberum</i> ) - Wherever found	E	\$3,226	\$1,269,338,780
1203	Lily, Minnesota dwarf trout ( <i>Erythronium propullans</i> ) - Wherever found	E	\$3,200	\$1,269,341,980
1203	Tarplant, Gaviota ( <i>Deinandra increscens</i> ssp. <i>villosa</i> ) - Wherever found	E	\$3,200	\$1,269,345,180
1203	Fern, Florida bristle ( <i>Trichomanes punctatum</i> ssp. <i>floridanum</i> ) - Wherever found	E	\$3,200	\$1,269,348,380
1204	Pigtoe, shiny ( <i>Fusconaia cor</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$3,181	\$1,269,351,561
1205	Blossom, tubercled (pearlymussel) ( <i>Epioblasma torulosa torulosa</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$3,131	\$1,269,354,692
1206	Shark, Scalloped Hammerhead ( <i>Sphyrna lewini</i> ) - Indo-West Pacific DPS	T	\$3,127	\$1,269,357,819
1207	Lioplax, cylindrical (snail) ( <i>Lioplax cyclostomaformis</i> ) - Wherever found	E	\$3,122	\$1,269,360,941
1208	Lupine, clover (Tidestrom"s) ( <i>Lupinus tidestromii</i> ) - Wherever found	E	\$3,100	\$1,269,364,041
1208	Gladecress, Texas golden ( <i>Leavenworthia texana</i> ) - Wherever found	E	\$3,100	\$1,269,367,141
1209	Pigtoe, dark ( <i>Pleurobema furvum</i> ) - Wherever found	E	\$3,073	\$1,269,370,214
1210	Boa, Mona ( <i>Epicrates monensis monensis</i> ) - Wherever found	T	\$3,000	\$1,269,373,214
1210	Frog, Goliath ( <i>Conraua goliath</i> ) - Wherever found	T	\$3,000	\$1,269,376,214
1210	Darter, slackwater ( <i>Etheostoma boschungii</i> ) - Wherever found	T	\$3,000	\$1,269,379,214
1210	Madtom, yellowfin ( <i>Noturus flavipinnis</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(e)(1)(iii))	EXPN	\$3,000	\$1,269,382,214
1210	No Common Name ( <i>Agave eggersiana</i> ) - Wherever found	E	\$3,000	\$1,269,385,214
1210	No Common Name ( <i>Chamaecrista glandulosa</i> var. <i>mirabilis</i> ) - Wherever found	E	\$3,000	\$1,269,388,214
1210	Bladderpod, white ( <i>Lesquerella pallida</i> ) - Wherever found	E	\$3,000	\$1,269,391,214
1210	No Common Name ( <i>Myrcia paganii</i> ) - Wherever found	E	\$3,000	\$1,269,394,214
1210	Leather flower, Alabama ( <i>Clematis socialis</i> ) - Wherever found	E	\$3,000	\$1,269,397,214

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1210	Lessingia, San Francisco ( <i>Lessingia germanorum</i> (=L.g. var. <i>germanorum</i> )) - Wherever found	E	\$3,000	\$1,269,400,214
1210	Quillwort, mat-forming ( <i>Isoetes tegetiformans</i> ) - Wherever found	E	\$3,000	\$1,269,403,214
1210	Lichen, rock gnome ( <i>Gymnoderma lineare</i> ) - Wherever found	E	\$3,000	\$1,269,406,214
1211	Darter, rush ( <i>Etheostoma phytophilum</i> ) - Wherever found	E	\$2,918	\$1,269,409,132
1212	Snail, painted snake coiled forest ( <i>Anguispira picta</i> ) - Wherever found	T	\$2,900	\$1,269,412,032
1213	Flax, Carter's small-flowered ( <i>Linum carteri carteri</i> ) - Wherever found	E	\$2,800	\$1,269,414,832
1214	Spineflower, Sonoma ( <i>Chorizanthe valida</i> ) - Wherever found	E	\$2,700	\$1,269,417,532
1214	Mustard, slender-petaled ( <i>Thelypodium stenopetalum</i> ) - Wherever found	E	\$2,700	\$1,269,420,232
1215	Love grass, Fosberg's ( <i>Eragrostis fosbergii</i> ) - Wherever found	E	\$2,689	\$1,269,422,921
1216	Cactus, Peebles Navajo ( <i>Pediocactus peeblesianus</i> var. <i>peeblesianus</i> ) - Wherever found	E	\$2,650	\$1,269,425,571
1217	Cactus, Nichol's Turk's head ( <i>Echinocactus horzonthalonius</i> var. <i>nicholii</i> ) - Wherever found	E	\$2,600	\$1,269,428,171
1217	Fringepod, Santa Cruz Island ( <i>Thysanocarpus conchuliferus</i> ) - Wherever found	E	\$2,600	\$1,269,430,771
1218	No Common Name ( <i>Keysseria</i> (=Lagenifera) <i>helenae</i> ) - Wherever found	E	\$2,588	\$1,269,433,359
1219	Snail, tulotoma ( <i>Tulotoma magnifica</i> ) - Wherever found	T	\$2,545	\$1,269,435,904
1220	Salamander, San Marcos ( <i>Eurycea nana</i> ) - Spring Lake and nearby habitats immediately downstream of the dam forming Spring Lake	RT	\$2,502	\$1,269,438,406
1221	Pupfish, Leon Springs ( <i>Cyprinodon bovinus</i> ) - Wherever found	E	\$2,500	\$1,269,440,906
1221	Chub, Sonora ( <i>Gila ditaenia</i> ) - Wherever found	T	\$2,500	\$1,269,443,406
1221	Darter, vermilion ( <i>Etheostoma chermocki</i> ) - Wherever found	E	\$2,500	\$1,269,445,906
1221	Lampmussel, Alabama ( <i>Lampsilis virescens</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$2,500	\$1,269,448,406
1221	Springsnail, Phantom ( <i>Pyrgulopsis texana</i> ) - Wherever found	E	\$2,500	\$1,269,450,906
1221	Butterfly, lotis blue ( <i>Lycæides argyrognomon lotis</i> ) - Wherever found	E	\$2,500	\$1,269,453,406
1221	Beetle, Coffin Cave mold ( <i>Batrisodes texanus</i> ) - Wherever found	E	\$2,500	\$1,269,455,906
1221	Beetle, Kretschmarr Cave mold ( <i>Texamaurops reddelli</i> ) - Wherever found	E	\$2,500	\$1,269,458,406
1221	Beetle, Tooth Cave ground ( <i>Rhadine persephone</i> ) - Wherever found	E	\$2,500	\$1,269,460,906
1221	Beetle, Helotes mold ( <i>Batrisodes venyivi</i> ) - Wherever found	E	\$2,500	\$1,269,463,406
1221	Harvestman, Bee Creek Cave ( <i>Texella reddelli</i> ) - Wherever found	E	\$2,500	\$1,269,465,906
1221	Pseudoscorpion, Tooth Cave ( <i>Tartarocreagris texana</i> ) - Wherever found	E	\$2,500	\$1,269,468,406
1221	Spider, Tooth Cave ( <i>Neoleptoneta myopica</i> ) - Wherever found	E	\$2,500	\$1,269,470,906
1221	Spider, Government Canyon Bat Cave ( <i>Neoleptoneta microps</i> ) - Wherever found	E	\$2,500	\$1,269,473,406
1221	Meshweaver, Madla Cave ( <i>Cicurina madla</i> ) - Wherever found	E	\$2,500	\$1,269,475,906

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1221	Meshweaver, Robber Baron Cave ( <i>Cicurina baronia</i> ) - Wherever found	E	\$2,500	\$1,269,478,406
1221	Meshweaver, Government Canyon Bat Cave ( <i>Cicurina vespera</i> ) - Wherever found	E	\$2,500	\$1,269,480,906
1221	Meshweaver, Braken Bat Cave ( <i>Cicurina venii</i> ) - Wherever found	E	\$2,500	\$1,269,483,406
1221	Amphipod, Peck's cave ( <i>Stygobromus (=Stygonectes) pecki</i> ) - Wherever found	E	\$2,500	\$1,269,485,906
1221	Amphipod, diminutive ( <i>Gammarus hyalleloides</i> ) - Wherever found	E	\$2,500	\$1,269,488,406
1221	Bladderpod, lyrate ( <i>Lesquerella lyrata</i> ) - Wherever found	T	\$2,500	\$1,269,490,906
1221	Pawpaw, beautiful ( <i>Deeringothamnus pulchellus</i> ) - Wherever found	E	\$2,500	\$1,269,493,406
1221	Locoweed, Fassett's ( <i>Oxytropis campestris</i> var. <i>chartacea</i> ) - Wherever found	T	\$2,500	\$1,269,495,906
1222	Bean, Cumberland (pearlymussel) ( <i>Villosa trabalis</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$2,409	\$1,269,498,315
1223	Dace, Independence Valley speckled ( <i>Rhinichthys osculus lethoporus</i> ) - Wherever found	E	\$2,402	\$1,269,500,717
1224	Springsnail, Alamosa ( <i>Tryonia alamosae</i> ) - Wherever found	E	\$2,400	\$1,269,503,117
1224	Pea, Big Pine partridge ( <i>Chamaecrista lineata keyensis</i> ) - Wherever found	E	\$2,400	\$1,269,505,517
1224	Pentachaeta, white-rayed ( <i>Pentachaeta bellidiflora</i> ) - Wherever found	E	\$2,400	\$1,269,507,917
1225	Lampmussel, Alabama ( <i>Lampsilis virescens</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$2,380	\$1,269,510,297
1226	Manzanita, Del Mar ( <i>Arctostaphylos glandulosa</i> ssp. <i>crassifolia</i> ) - Wherever found	E	\$2,300	\$1,269,512,597
1226	Goldenrod, Blue Ridge ( <i>Solidago spithamea</i> ) - Wherever found	T	\$2,300	\$1,269,514,897
1227	Turtle, flattened musk ( <i>Sternotherus depressus</i> ) - Black Warrior R. system upstream from Bankhead Dam	T	\$2,237	\$1,269,517,134
1228	Grasshopper, Zayante band-winged ( <i>Trimerotropis infantilis</i> ) - Wherever found	E	\$2,200	\$1,269,519,334
1229	Clover, Monterey ( <i>Trifolium trichocalyx</i> ) - Wherever found	E	\$2,150	\$1,269,521,484
1230	Milk-vetch, Clara Hunt's ( <i>Astragalus clarianus</i> ) - Wherever found	E	\$2,100	\$1,269,523,584
1230	Spurge, wedge ( <i>Chamaesyce deltoidea serpyllum</i> ) - Wherever found	E	\$2,100	\$1,269,525,684
1231	Acornshell, southern ( <i>Epioblasma othcaloogensis</i> ) - Wherever found	E	\$2,042	\$1,269,527,726
1232	Crow, white-necked ( <i>Corvus leucognaphalus</i> ) - Wherever found	E	\$2,000	\$1,269,529,726
1232	Gambusia, Big Bend ( <i>Gambusia gaigei</i> ) - Wherever found	E	\$2,000	\$1,269,531,726
1232	Spinymussel, Altamaha ( <i>Elliptio spinosa</i> ) - Wherever found	E	\$2,000	\$1,269,533,726
1232	Snail, Virginia fringed mountain ( <i>Polygyriscus virginianus</i> ) - Wherever found	E	\$2,000	\$1,269,535,726
1232	Lily, Pitkin Marsh ( <i>Lilium pardalinum</i> ssp. <i>pitkinense</i> ) - Wherever found	E	\$2,000	\$1,269,537,726
1232	Amphianthus, little ( <i>Amphianthus pusillus</i> ) - Wherever found	T	\$2,000	\$1,269,539,726
1232	Primrose, Maguire ( <i>Primula maguirei</i> ) - Wherever found	T	\$2,000	\$1,269,541,726
1232	Bird's-beak, Pennell's ( <i>Cordylanthus tenuis</i> ssp. <i>capillaris</i> ) - Wherever found	E	\$2,000	\$1,269,543,726

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1232	No Common Name ( <i>Gonocalyx concolor</i> ) - Wherever found	E	\$2,000	\$1,269,545,726
1232	Quillwort, black spored ( <i>Isoetes melanospora</i> ) - Wherever found	E	\$2,000	\$1,269,547,726
1233	Checker-mallow, Kenwood Marsh ( <i>Sidalcea oregana</i> ssp. <i>valida</i> ) - Wherever found	E	\$1,900	\$1,269,549,626
1234	Chub, slender ( <i>Erimystax cahni</i> ) - Wherever found	T	\$1,870	\$1,269,551,496
1235	Flax, sand ( <i>Linum arenicola</i> ) - Wherever found	E	\$1,800	\$1,269,553,296
1235	Buttercup, autumn ( <i>Ranunculus aestivalis</i> (=acriformis)) - Wherever found	E	\$1,800	\$1,269,555,096
1235	Spineflower, slender-horned ( <i>Dodecahema leptoceras</i> ) - Wherever found	E	\$1,800	\$1,269,556,896
1236	Wild-buckwheat, southern mountain ( <i>Eriogonum kennedyi</i> var. <i>austromontanum</i> ) - Wherever found	T	\$1,700	\$1,269,558,596
1236	Mariposa lily, Tiburon ( <i>Calochortus tiburonensis</i> ) - Wherever found	T	\$1,700	\$1,269,560,296
1237	Bladderpod, kodachrome ( <i>Lesquerella tumulosa</i> ) - Wherever found	E	\$1,650	\$1,269,561,946
1238	Chub, Chihuahua ( <i>Gila nigrescens</i> ) - Wherever found	T	\$1,600	\$1,269,563,546
1239	Combshell, southern ( <i>Epioblasma penita</i> ) - Wherever found	E	\$1,500	\$1,269,565,046
1239	Cory cactus, bunched ( <i>Coryphantha ramillosa</i> ) - Wherever found	T	\$1,500	\$1,269,566,546
1239	Cactus, Lloyd's Mariposa ( <i>Echinomastus mariposensis</i> ) - Wherever found	T	\$1,500	\$1,269,568,046
1239	Cactus, Chisos Mountain hedgehog ( <i>Echinocereus chisoensis</i> var. <i>chisoensis</i> ) - Wherever found	T	\$1,500	\$1,269,569,546
1239	Lupine, scrub ( <i>Lupinus aridorum</i> ) - Wherever found	E	\$1,500	\$1,269,571,046
1240	Pigtoe, heavy ( <i>Pleurobema taitianum</i> ) - Wherever found	E	\$1,464	\$1,269,572,510
1241	Poppy-mallow, Texas ( <i>Callirhoe scabriuscula</i> ) - Wherever found	E	\$1,420	\$1,269,573,930
1242	Shark, Scalloped Hammerhead ( <i>Sphyrna lewini</i> ) - Central and Southwest Atlantic DPS	T	\$1,407	\$1,269,575,337
1243	Springsnail, San Bernardino ( <i>Pyrgulopsis bernardina</i> ) - Wherever found	T	\$1,400	\$1,269,576,737
1243	Dudleya, Santa Clara Valley ( <i>Dudleya setchellii</i> ) - Wherever found	E	\$1,400	\$1,269,578,137
1244	Manzanita, pallid ( <i>Arctostaphylos pallida</i> ) - Wherever found	T	\$1,300	\$1,269,579,437
1244	Brickell-bush, Florida ( <i>Brickellia mosieri</i> ) - Wherever found	E	\$1,300	\$1,269,580,737
1244	Crabgrass, Florida pineland ( <i>Digitaria pauciflora</i> ) - Wherever found	T	\$1,300	\$1,269,582,037
1244	Onion, Munz's ( <i>Allium munzii</i> ) - Wherever found	E	\$1,300	\$1,269,583,337
1244	Sandmat, pineland ( <i>Chamaesyce deltoidea pinetorum</i> ) - Wherever found	T	\$1,300	\$1,269,584,637
1245	Water-plantain, Kral's ( <i>Sagittaria secundifolia</i> ) - Wherever found	T	\$1,299	\$1,269,585,936
1246	Mussel, oyster ( <i>Epioblasma capsaeformis</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$1,253	\$1,269,587,189
1247	Navarretia, few-flowered ( <i>Navarretia leucocephala</i> ssp. <i>pauciflora</i> (=N. <i>pauciflora</i> )) - Wherever found	E	\$1,200	\$1,269,588,389

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1248	Chub, spotfin ( <i>Erimonax monachus</i> ) - U.S.A. (TN-specified portions of the Tellico River; see 17.84(m)(1)(i))	EXPN	\$1,156	\$1,269,589,545
1249	Monkeyface, Appalachian (pearlymussel) ( <i>Quadrula sparsa</i> ) - Wherever found	E	\$1,100	\$1,269,590,645
1249	Prickly-apple, aboriginal ( <i>Harrisia (=Cereus) aboriginum (=gracilis)</i> ) - Wherever found	E	\$1,100	\$1,269,591,745
1250	Reed-mustard, Barneby ( <i>Schoenocrambe barnebyi</i> ) - Wherever found	E	\$1,050	\$1,269,592,795
1251	Madtom, Scioto ( <i>Noturus trautmani</i> ) - Wherever found	E	\$1,030	\$1,269,593,825
1252	Combshell, Cumberlandian ( <i>Epioblasma brevidens</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$1,002	\$1,269,594,827
1253	Wolf, red ( <i>Canis rufus</i> ) - Wherever found, except where listed as an experimental population	E	\$1,000	\$1,269,595,827
1253	Gambusia, Clear Creek ( <i>Gambusia heterochir</i> ) - Wherever found	E	\$1,000	\$1,269,596,827
1253	Madtom, smoky ( <i>Noturus baileyi</i> ) - Tellico River, TN	EXPN	\$1,000	\$1,269,597,827
1253	Tryonia, Diamond ( <i>Pseudotryonia adamantina</i> ) - Wherever found	E	\$1,000	\$1,269,598,827
1253	Cactus, Nellie cory ( <i>Coryphantha minima</i> ) - Wherever found	E	\$1,000	\$1,269,599,827
1253	Snowbells, Texas ( <i>Styrax texanus</i> ) - Wherever found	E	\$1,000	\$1,269,600,827
1254	Chub, spotfin ( <i>Erimonax monachus</i> ) - U.S.A. (AL, TN-specified portions of Shoal Creek; see 17.84(m)(1)(ii))	EXPN	\$800	\$1,269,601,627
1254	Snail, armored ( <i>Pyrgulopsis (=Marstonia) pachyta</i> ) - Wherever found	E	\$800	\$1,269,602,427
1254	Jewelflower, Metcalf Canyon ( <i>Streptanthus albidus</i> ssp. <i>albidus</i> ) - Wherever found	E	\$800	\$1,269,603,227
1254	Ladies-tresses, Canelo Hills ( <i>Spiranthes delitescens</i> ) - Wherever found	E	\$800	\$1,269,604,027
1255	Curlew, Eskimo ( <i>Numenius borealis</i> ) - Wherever found	E	\$729	\$1,269,604,756
1256	Cavefish, Alabama ( <i>Speoplatyrhinus poulsoni</i> ) - Wherever found	E	\$700	\$1,269,605,456
1256	Sawfish, largetooth ( <i>Pristis pristis</i> ) - Wherever found	E	\$700	\$1,269,606,156
1256	Blossom, turgid (pearlymussel) ( <i>Epioblasma turgidula</i> ) - Wherever found; Except where listed as Experimental Populations	E	\$700	\$1,269,606,856
1257	Darter, Pearl ( <i>Percina aurora</i> ) - Wherever found	T	\$678	\$1,269,607,534
1258	Sunfish, spring pygmy ( <i>Elassoma alabamae</i> ) - Wherever found	T	\$500	\$1,269,608,034
1258	Blossom, green (pearlymussel) ( <i>Epioblasma torulosa gubernaculum</i> ) - Wherever found	E	\$500	\$1,269,608,534
1258	Clubshell, black ( <i>Pleurobema curtum</i> ) - Wherever found	E	\$500	\$1,269,609,034
1258	Pitaya, Davis' green ( <i>Echinocereus viridiflorus</i> var. <i>davisii</i> ) - Wherever found	E	\$500	\$1,269,609,534
1258	Cat's-eye, Terlingua Creek ( <i>Cryptantha crassipes</i> ) - Wherever found	E	\$500	\$1,269,610,034
1259	Whale, Southern Right ( <i>Eubalaena australis</i> ) - Wherever found	E	\$476	\$1,269,610,510
1260	Darter, bayou ( <i>Etheostoma rubrum</i> ) - Wherever found	T	\$400	\$1,269,610,910
1260	Pigtoe, flat ( <i>Pleurobema marshalli</i> ) - Wherever found	E	\$400	\$1,269,611,310

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1260	Stirrupshell ( <i>Quadrula stapes</i> ) - Wherever found	E	\$400	\$1,269,611,710
1260	Mussel, snuffbox ( <i>Epioblasma triquetra</i> ) - Wherever found	NS	\$400	\$1,269,612,110
1260	Rabbitsfoot ( <i>Quadrula cylindrica cylindrica</i> ) -	NS	\$400	\$1,269,612,510
1260	Birch, Virginia round-leaf ( <i>Betula uber</i> ) - Wherever found	T	\$400	\$1,269,612,910
1261	Pearlymussel, birdwing ( <i>Lemiox rimosus</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert, and Lauderdale Counties, AL	EXPN	\$350	\$1,269,613,260
1262	Madtom, chunky ( <i>Noturus crypticus</i> ) - Wherever found	E	\$300	\$1,269,613,560
1262	Minnow, Rio Grande Shiner ( <i>Hypognathus amarus</i> ) - Rio Grande, from Little Box Canyon to Assisted Dam	EXPN	\$300	\$1,269,613,860
1262	Monkeyface, Cumberland (pearlymussel) ( <i>Quadrula intermedia</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$300	\$1,269,614,160
1262	Monkeyface, Cumberland (pearlymussel) ( <i>Quadrula intermedia</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$300	\$1,269,614,460
1262	Pigtoe, finereyed ( <i>Fusconaia cuneolus</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$300	\$1,269,614,760
1262	Pigtoe, shiny ( <i>Fusconaia cor</i> ) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$300	\$1,269,615,060
1262	Milk-vetch, coastal dunes ( <i>Astragalus tener</i> var. <i>titi</i> ) - Wherever found	E	\$300	\$1,269,615,360
1262	Paintbrush, ash-grey ( <i>Castilleja cinerea</i> ) - Wherever found	T	\$300	\$1,269,615,660
1262	Clarkia, Pismo ( <i>Clarkia speciosa</i> ssp. <i>immaculata</i> ) - Wherever found	E	\$300	\$1,269,615,960
1262	Checker-mallow, pedate ( <i>Sidalcea pedata</i> ) - Wherever found	E	\$300	\$1,269,616,260
1262	Crownbeard, big-leaved ( <i>Verbesina dissita</i> ) - Wherever found	T	\$300	\$1,269,616,560
1262	Bully, Everglades ( <i>Sideroxylon reclinatum</i> ssp. <i>austrofloridense</i> ) - Wherever found	T	\$300	\$1,269,616,860
1263	Fern, Alabama streak-sorus ( <i>Thelypteris pilosa</i> var. <i>alabamensis</i> ) - Wherever found	T	\$299	\$1,269,617,159
1264	Purple Cat's paw (=Purple Cat's paw pearlymussel) ( <i>Epioblasma obliquata obliquata</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$251	\$1,269,617,410
1264	Pearlymussel, cracking ( <i>Hemistena lata</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$251	\$1,269,617,661
1265	Blossom, turgid (pearlymussel) ( <i>Epioblasma turgidula</i> ) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$250	\$1,269,617,911
1265	Pondweed, Little Aguja (=Creek) ( <i>Potamogeton clystocarpus</i> ) - Wherever found	E	\$250	\$1,269,618,161
1266	Darter, Maryland ( <i>Etheostoma sellare</i> ) - Wherever found	E	\$240	\$1,269,618,401
1266	Amphipod, Hay's Spring ( <i>Stygobromus hayi</i> ) - Wherever found	E	\$240	\$1,269,618,641



Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1267	Parrot, thick-billed ( <i>Rhynchopsitta pachyrhyncha</i> ) - Wherever found	E	\$200	\$1,269,618,841
1267	Liveforever, Laguna Beach ( <i>Dudleya stolonifera</i> ) - Wherever found	T	\$200	\$1,269,619,041
1267	Stonecrop, Lake County ( <i>Parvisedum leiocarpum</i> ) - Wherever found	E	\$200	\$1,269,619,241
1267	Water-willow, Cooley's ( <i>Justicia cooleyi</i> ) - Wherever found	E	\$200	\$1,269,619,441
1267	Thistle, Loch Lomond coyote ( <i>Eryngium constancei</i> ) - Wherever found	E	\$200	\$1,269,619,641
1267	Mint, longspurred ( <i>Dicerandra cornutissima</i> ) - Wherever found	E	\$200	\$1,269,619,841
1268	Pebblesnail, flat ( <i>Lepyrium showalteri</i> ) - Wherever found	E	\$149	\$1,269,619,990
1269	Springsnail, Chupadera ( <i>Pyrgulopsis chupadera</i> ) - Wherever found	E	\$125	\$1,269,620,115
1270	Chub, spotfin ( <i>Erimonax monachus</i> ) - U.S.A. (TN-specified portions of the French Broad and Holston Rivers; see 17.84(m)(1)(iii))	EXPN	\$100	\$1,269,620,215
1270	Springsnail, Socorro ( <i>Pyrgulopsis neomexicana</i> ) - Wherever found	E	\$100	\$1,269,620,315
1270	Sandwort, Bear Valley ( <i>Arenaria ursina</i> ) - Wherever found	T	\$100	\$1,269,620,415
1270	Mountain-mahogany, Catalina Island ( <i>Cercocarpus traskiae</i> ) - Wherever found	E	\$100	\$1,269,620,515
1270	Taraxacum, California ( <i>Taraxacum californicum</i> ) - Wherever found	E	\$100	\$1,269,620,615
1270	Ridge-cress, Barneby ( <i>Lepidium barnebyanum</i> ) - Wherever found	E	\$100	\$1,269,620,715
1270	Thornmint, San Mateo ( <i>Acanthomintha obovata</i> ssp. <i>duttonii</i> ) - Wherever found	E	\$100	\$1,269,620,815
1270	Bladderpod, San Bernardino Mountains ( <i>Lesquerella kingii</i> ssp. <i>bernardina</i> ) - Wherever found	E	\$100	\$1,269,620,915
1270	Malacothrix, island ( <i>Malacothrix squalida</i> ) - Wherever found	E	\$100	\$1,269,621,015
1271	Crane, Siberian white ( <i>Grus leucogeranus</i> ) - Wherever found	E	\$20	\$1,269,621,035
1271	Crane, hooded ( <i>Grus monacha</i> ) - Wherever found	E	\$20	\$1,269,621,055
1272	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - North Cascades Ecosystem Recovery Zone	UR	\$0	\$1,269,621,055
1272	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - Alaska	RT	\$0	\$1,269,621,055
1272	Deer, Columbian white-tailed ( <i>Odocoileus virginianus leucurus</i> ) - Douglas County DPS	DM	\$0	\$1,269,621,055
1272	Wolf, gray ( <i>Canis lupus</i> ) - U.S.A. (portions of AZ, NM, TX), Mexico, except where listed as an experimental population, captive wolves who were, or whose ancestors were, removed from the wild in this area.	SU	\$0	\$1,269,621,055
1272	Wolf, gray ( <i>Canis lupus</i> ) - U.S.A. (MI, MN, ND, SD, WI); captive wolves who were, or whose ancestors were, removed from the wild in this area.	SU	\$0	\$1,269,621,055
1272	Wolf, gray ( <i>Canis lupus</i> ) - U.S.A. (ME, NH, NY, VT); captive wolves who were, or whose ancestors were, removed from the wild in this area.	SU	\$0	\$1,269,621,055
1272	Wolf, gray ( <i>Canis lupus</i> ) - U.S.A. (CO, ID, MT, OR, UT, WA, WY; portions of AZ and NM), except where listed as an experimental population captive wolves who were, or whose ancestors were, removed from the wild in this area	SU	\$0	\$1,269,621,055

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1272	Wolf, gray ( <i>Canis lupus</i> ) - WI, MN, MI	SU	\$0	\$1,269,621,055
1272	Wolf, gray ( <i>Canis lupus</i> ) - Alaska population	SU	\$0	\$1,269,621,055
1272	Wolf, Mexican ( <i>Canis lupus baileyi</i> ) -	SU	\$0	\$1,269,621,055
1272	Whale, gray ( <i>Eschrichtius robustus</i> ) - except where listed	DM	\$0	\$1,269,621,055
1272	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - West Indies DPS	DM	\$0	\$1,269,621,055
1272	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Hawaii DPS	DM	\$0	\$1,269,621,055
1272	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Brazil DPS	DM	\$0	\$1,269,621,055
1272	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Gabon/Southwest Africa DPS	DM	\$0	\$1,269,621,055
1272	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Southeast Africa/Madagascar DPS	DM	\$0	\$1,269,621,055
1272	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - West Australia DPS	DM	\$0	\$1,269,621,055
1272	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - East Australia DPS	DM	\$0	\$1,269,621,055
1272	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Oceania DPS	DM	\$0	\$1,269,621,055
1272	Whale, humpback ( <i>Megaptera novaeangliae</i> ) - Southeastern Pacific DPS	DM	\$0	\$1,269,621,055
1272	Bear, brown ( <i>Ursus arctos arctos</i> ) - Kenai Pop.	SC	\$0	\$1,269,621,055
1272	Lynx, Canada ( <i>Lynx canadensis</i> ) - NM population	RT	\$0	\$1,269,621,055
1272	Bat, Mariana fruit (=Mariana flying fox) ( <i>Pteropus mariannus mariannus</i> ) - Aguijan, etc.	RT	\$0	\$1,269,621,055
1272	Bat, Mariana fruit (=Mariana flying fox) ( <i>Pteropus mariannus mariannus</i> ) - Rota, etc.	RT	\$0	\$1,269,621,055
1272	Caribou DPS, Southern Mountain ( <i>Rangifer tarandus ssp. caribou</i> ) - MT pop.	SU	\$0	\$1,269,621,055
1272	Bat, Pacific sheath-tailed ( <i>Emballonura semicaudata semicaudata</i> ) - Caroline Islands	RT	\$0	\$1,269,621,055
1272	Rabbit, Columbia Basin Pygmy ( <i>Brachylagus idahoensis</i> ) - Wherever found	RT	\$0	\$1,269,621,055
1272	Otter, Northern Sea ( <i>Enhydra lutris kenyoni</i> ) - NSO (WA)	SU	\$0	\$1,269,621,055
1272	Otter, Northern Sea ( <i>Enhydra lutris kenyoni</i> ) - SCAK NSO	SU	\$0	\$1,269,621,055
1272	Otter, Northern Sea ( <i>Enhydra lutris kenyoni</i> ) - SEAK NSO	SU	\$0	\$1,269,621,055
1272	Seal, bearded ( <i>Erignathus barbatus nauticus</i> ) - Wherever found	SU	\$0	\$1,269,621,055
1272	Dolphin, Taiwanese humpback ( <i>Sousa chinensis taiwanensis</i> )	PE	\$0	\$1,269,621,055
1272	Dolphin, South Island Hector's ( <i>Cephalorhynchus hectori hectori</i> )	T	\$0	\$1,269,621,055
1272	Akialoa, Kauai (honeycreeper) ( <i>Akialoa stejnegeri</i> ) -	NS	\$0	\$1,269,621,055
1272	Akialoa, Kauai (honeycreeper) ( <i>Akialoa stejnegeri</i> ) -	NS	\$0	\$1,269,621,055
1272	Plover, western snowy ( <i>Charadrius nivosus nivosus</i> ) - Interior pop.	RT	\$0	\$1,269,621,055
1272	Tern, least ( <i>Sterna antillarum</i> ) - Atlantic	SC	\$0	\$1,269,621,055
1272	Murrelet, marbled ( <i>Brachyramphus marmoratus</i> ) - except CA, OR, WA	SC	\$0	\$1,269,621,055
1272	Storm-petrel, band-rumped ( <i>Oceanodroma castro</i> ) - Wherever found	SC	\$0	\$1,269,621,055
1272	Goshawk, Queen Charlotte ( <i>Accipiter gentilis laingi</i> ) - Wherever found	RT	\$0	\$1,269,621,055

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1272	Tortoise, gopher ( <i>Gopherus polyphemus</i> ) - eastern	C	\$0	\$1,269,621,055
1272	Tortoise, gopher ( <i>Gopherus polyphemus</i> ) - Florida	RT	\$0	\$1,269,621,055
1272	Tortoise, gopher ( <i>Gopherus polyphemus</i> ) - Wherever found	UR	\$0	\$1,269,621,055
1272	Tortoise, desert ( <i>Gopherus agassizii</i> ) - Sonoran	RT	\$0	\$1,269,621,055
1272	Salamander, Texas blind ( <i>Eurycea [=Typhlomolge] rathbuni</i> ) -	SU	\$0	\$1,269,621,055
1272	Salamander, California tiger ( <i>Ambystoma californiense</i> ) - Wherever found	RT	\$0	\$1,269,621,055
1272	Salamander, Jollyville Plateau ( <i>Eurycea tonkawae</i> ) - Buttercup Creek	SC	\$0	\$1,269,621,055
1272	Tui chub, Mohave ( <i>Gila bicolor</i> ssp. <i>mohavensis</i> ) - Lewis Center	NS	\$0	\$1,269,621,055
1272	Tui chub, Mohave ( <i>Gila bicolor</i> ssp. <i>mohavensis</i> ) - Morning Star Mine	NS	\$0	\$1,269,621,055
1272	Chub, spotfin ( <i>Erimonax monachus</i> ) - ERP	SU	\$0	\$1,269,621,055
1272	Chub, spotfin ( <i>Erimonax monachus</i> ) - BRP	SU	\$0	\$1,269,621,055
1272	Chub, spotfin ( <i>Erimonax monachus</i> ) - HRP	SU	\$0	\$1,269,621,055
1272	Chub, spotfin ( <i>Erimonax monachus</i> ) - LTRP	SU	\$0	\$1,269,621,055
1272	Darter, candy ( <i>Etheostoma osburni</i> ) - Wherever found	E	\$0	\$1,269,621,055
1272	Pupfish, desert ( <i>Cyprinodon macularius</i> ) -	SU	\$0	\$1,269,621,055
1272	Salmon, sockeye ( <i>Oncorhynchus (=Salmo) nerka</i> ) - Kokanee - Lake Sammamish and Lake Washinton	RT	\$0	\$1,269,621,055
1272	Salmon, sockeye ( <i>Oncorhynchus (=Salmo) nerka</i> ) - Issaquah Creek population	RT	\$0	\$1,269,621,055
1272	Goby, tidewater ( <i>Eucyclogobius newberryi</i> ) - Populations north of Orange County, CA	RT	\$0	\$1,269,621,055
1272	Salmon, Atlantic ( <i>Salmo salar</i> ) - Kennebec River DPS	RT	\$0	\$1,269,621,055
1272	Steelhead ( <i>Oncorhynchus (=Salmo) mykiss</i> ) - Klamath Mountains Province	UR	\$0	\$1,269,621,055
1272	Sturgeon, Atlantic ( <i>Acipenser oxyrinchus oxyrinchus</i> ) - Wherever found	RT	\$0	\$1,269,621,055
1272	Darter, rush ( <i>Etheostoma phytophilum</i> ) -	NS	\$0	\$1,269,621,055
1272	Coelacanth, African ( <i>Latimeria chalumnae</i> )	T	\$0	\$1,269,621,055
1272	Angelshark, Argentine ( <i>Squatina argentina</i> )	E	\$0	\$1,269,621,055
1272	Guitarfish, Blackchin ( <i>Rhinobatos cemiculus</i> )	T	\$0	\$1,269,621,055
1272	Guitarfish Brazilian ( <i>Rhinobatos horkelii</i> )	E	\$0	\$1,269,621,055
1272	Guitarfish Common ( <i>Rhinobatos rhinobatos</i> )	T	\$0	\$1,269,621,055
1272	Ray, giant manta ( <i>Manta birostris</i> )	T	\$0	\$1,269,621,055
1272	Grouper, Gulf ( <i>Mycteroperca jordani</i> )	E	\$0	\$1,269,621,055
1272	Grouper, Island ( <i>Mycteroperca fusca</i> )	T	\$0	\$1,269,621,055
1272	Shark, Narrownose smoothhound ( <i>Mustelus schmitti</i> )	T	\$0	\$1,269,621,055
1272	Shark, Oceanic whitetip ( <i>Carcharhinus longimanus</i> )	T	\$0	\$1,269,621,055

<b>Rank</b>	<b>Species (50 CFR Part 17)</b>	<b>Status</b>	<b>Species Total</b>	<b>Cumulatative Total</b>
1272	Angelshark, spiny ( <i>Squatina guggenheim</i> )	E	\$0	\$1,269,621,055
1272	Shark, Striped smoothhound ( <i>Mustelus fasciatus</i> )	E	\$0	\$1,269,621,055
1272	Mucket, Neosho ( <i>Lampsilis rafinesqueana</i> ) - Wherever found	NS	\$0	\$1,269,621,055
1272	Mucket, Neosho ( <i>Lampsilis rafinesqueana</i> ) -	NS	\$0	\$1,269,621,055
1272	Riffleshell, tan ( <i>Epioblasma florentina walkeri</i> (=E. walkeri)) - BSFP	SU	\$0	\$1,269,621,055
1272	Riffleshell, tan ( <i>Epioblasma florentina walkeri</i> (=E. walkeri)) - ICP	SU	\$0	\$1,269,621,055
1272	Pigtoe, heavy ( <i>Pleurobema taitianum</i> ) - Wherever found	NS	\$0	\$1,269,621,055
1272	Pigtoe, heavy ( <i>Pleurobema taitianum</i> ) - Wherever found	NS	\$0	\$1,269,621,055
1272	Bean, rayed ( <i>Villosa fabalis</i> ) - Wherever found	NS	\$0	\$1,269,621,055
1272	Heelsplitter, Inflated ( <i>Potamilus inflatus</i> ) -	NS	\$0	\$1,269,621,055
1272	Hornsnail, rough ( <i>Pleurocera foremani</i> ) -	SU	\$0	\$1,269,621,055
1272	Hornsnail, rough ( <i>Pleurocera foremani</i> ) -	SU	\$0	\$1,269,621,055
1272	Hornsnail, rough ( <i>Pleurocera foremani</i> ) -	SU	\$0	\$1,269,621,055
1272	Beetle, American burying ( <i>Nicrophorus americanus</i> ) - Wherever found	NS	\$0	\$1,269,621,055
1272	Sturgeon, Sakhalin ( <i>Acipenser mikadoi</i> ) - Wherever found	NS	\$0	\$1,269,621,055
1272	Shark, Daggernose ( <i>Isogomphodon oxyrinchus</i> )	E	\$0	\$1,269,621,055
<b>Total Ranked Species Expenditures</b>				<b>\$1,269,621,055</b>

**TABLE 3. FY 2018 REPORTED LAND ACQUISITION EXPENDITURES FOR  
ENDANGERED AND THREATENED SPECIES**

Species are arranged in the same sequence as Table 1. These costs are not included in the expenditures displayed in Tables 1 and 2. Only those species that had land acquisition costs reported for FY 2018 are included here. Land rank is included to show the relative order of the species for which land expenditures are reported. Land acquisition costs for multi-species conservation activities are tabulated at the end of Table 3. The STATUS column indicates whether the species is endangered (E), threatened (T), experimental (EXPN), or threatened by similarity of appearance (SAT).

**TABLE 3. REPORTED LAND ACQUISITION EXPENDITURES FOR FY 2018 FOR ENDANGERED AND THREATENED SPECIES**

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
Mammals	55	Bat, Florida bonneted ( <i>Eumops floridanus</i> ) - Wherever found	E	\$0	\$250,000	\$250,000	\$0	\$250,000
	35	Bat, Hawaiian hoary ( <i>Lasiurus cinereus semotus</i> ) - Wherever found	E	\$620,476	\$0	\$620,476	\$0	\$620,476
	34	Bat, Indiana ( <i>Myotis sodalis</i> ) - Wherever found	E	\$285,667	\$0	\$285,667	\$339,643	\$625,310
	2	Bat, Northern long-eared ( <i>Myotis septentrionalis</i> ) - Wherever found	T	\$0	\$11,947,000	\$11,947,000	\$391,500	\$12,338,500
	36	Bear, grizzly ( <i>Ursus arctos horribilis</i> ) - U.S.A., conterminous (lower 48) States, except where listed as an experimental population	T	\$605,167	\$0	\$605,167	\$0	\$605,167
	90	Deer, Columbian white-tailed ( <i>Odocoileus virginianus leucurus</i> ) - Columbia River DPS	T	\$7,500	\$0	\$7,500	\$0	\$7,500
	39	Fox, San Joaquin kit ( <i>Vulpes macrotis mutica</i> ) - wherever found	E	\$300,000	\$207,500	\$507,500	\$0	\$507,500
	74	Kangaroo rat, giant ( <i>Dipodomys ingens</i> ) - Wherever found	E	\$0	\$75,000	\$75,000	\$0	\$75,000
	68	Kangaroo rat, Tipton ( <i>Dipodomys nitratoides nitratoides</i> ) - Wherever found	E	\$0	\$132,500	\$132,500	\$0	\$132,500
	22	Lynx, Canada ( <i>Lynx canadensis</i> ) - Wherever Found in Contiguous U.S.	T	\$861,917	\$0	\$861,917	\$90,000	\$951,917
	24	Manatee, West Indian ( <i>Trichechus manatus</i> ) - Wherever found	T	\$167,356	\$600,058	\$767,414	\$0	\$767,414
	63	Otter, Northern Sea ( <i>Enhydra lutris kenyoni</i> ) - Southwest Alaska DPS	T	\$190,000	\$0	\$190,000	\$0	\$190,000
	95	Panther, Florida ( <i>Puma (=Felis) concolor coryi</i> ) - Wherever found	E	\$0	\$2,151	\$2,151	\$0	\$2,151
	86	Pocket gopher, Yelm ( <i>Thomomys mazama yelmensis</i> ) - Wherever found	T	\$0	\$0	\$0	\$26,000	\$26,000
	84	Pronghorn, Sonoran ( <i>Antilocapra americana sonoriensis</i> ) - Wherever found, except where listed as an experimental population	E	\$0	\$27,000	\$27,000	\$0	\$27,000
	5	Puma (=mountain lion) ( <i>Puma (=Felis) concolor</i> (all subsp. except <i>coryi</i> )) - FL	SAT	\$0	\$4,000,000	\$4,000,000	\$0	\$4,000,000
	23	Rabbit, Columbia Basin Pygmy ( <i>Brachylagus idahoensis</i> ) - Columbia Basin DPS	E	\$0	\$0	\$0	\$940,000	\$940,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	87	Seal, Hawaiian monk ( <i>Monachus schauinslandi</i> ) - Wherever found	E	\$25,995	\$0	\$25,995	\$0	\$25,995
	62	Sea lion, Steller ( <i>Eumetopias jubatus</i> ) - Western DPS	E	\$194,771	\$0	\$194,771	\$0	\$194,771
	28	Wolf, gray ( <i>Canis lupus</i> ) - U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	E	\$91,000	\$0	\$91,000	\$590,000	\$681,000
	71	Wolf, red ( <i>Canis rufus</i> ) - Wherever found, except where listed as an experimental population	E	\$110,000	\$0	\$110,000	\$0	\$110,000
<b>Mammals subtotal</b>				<b>\$3,459,849</b>	<b>\$17,241,209</b>	<b>\$20,701,058</b>	<b>\$2,377,143</b>	<b>\$23,078,201</b>
Birds	92	Albatross, short-tailed ( <i>Phoebastria</i> (=Diomedea) albatrus) - Wherever found	E	\$4,771	\$0	\$4,771	\$0	\$4,771
	83	Bobwhite, masked (quail) ( <i>Colinus virginianus ridgwayi</i> ) - Wherever found	E	\$30,000	\$0	\$30,000	\$0	\$30,000
	17	Caracara, Audubon's crested ( <i>Polyborus plancus audubonii</i> ) - FL pop.	T	\$3,913	\$1,500,000	\$1,503,913	\$0	\$1,503,913
	51	Condor, California ( <i>Gymnogyps californianus</i> ) - U.S.A. only, except where listed as an experimental population	E	\$300,000	\$0	\$300,000	\$0	\$300,000
	26	Condor, California ( <i>Gymnogyps californianus</i> ) - U.S.A. (specific portions of Arizona, Nevada, and Utah)	EXPN	\$0	\$728,000	\$728,000	\$0	\$728,000
	32	Coot, Hawaiian ( <i>Fulica americana alai</i> ) - Wherever found	E	\$646,471	\$0	\$646,471	\$0	\$646,471
	40	Crane, whooping ( <i>Grus americana</i> ) - Wherever found, except where listed as an experimental population	E	\$445,395	\$0	\$445,395	\$57,702	\$503,097
	35	Creepers, Hawaii ( <i>Oreomystis mana</i> ) - Wherever found	E	\$620,476	\$0	\$620,476	\$0	\$620,476
	50	Cuckoo, yellow-billed ( <i>Coccyzus americanus</i> ) - Western U.S. DPS	T	\$0	\$331,608	\$331,608	\$0	\$331,608
	32	Duck, Hawaiian (=koloa) ( <i>Anas wyvilliana</i> ) - Wherever found	E	\$646,471	\$0	\$646,471	\$0	\$646,471
	76	Eider, spectacled ( <i>Somateria fischeri</i> ) - Wherever found	T	\$64,000	\$0	\$64,000	\$0	\$64,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	25	Eider, Steller's ( <i>Polysticta stelleri</i> ) - AK breeding pop.	T	\$729,000	\$0	\$729,000	\$0	\$729,000
	41	Falcon, Northern Aplomado ( <i>Falco femoralis septentrionalis</i> ) - Wherever found, except where listed as an experimental population	E	\$426,293	\$0	\$426,293	\$0	\$426,293
	85	Gnatcatcher, coastal California ( <i>Poliophtila californica californica</i> ) - Wherever found	T	\$26,385	\$0	\$26,385	\$0	\$26,385
	35	Goose, Hawaiian ( <i>Branta (=Nesochen) sandvicensis</i> ) - Wherever found	E	\$620,476	\$0	\$620,476	\$0	\$620,476
	35	Hawk, Hawaiian (=Io) ( <i>Buteo solitarius</i> ) - Wherever found	E	\$620,476	\$0	\$620,476	\$0	\$620,476
	14	Kite, Everglade snail ( <i>Rostrhamus sociabilis plumbeus</i> ) - Wherever found	E	\$0	\$2,000,531	\$2,000,531	\$0	\$2,000,531
	92	Murrelet, marbled ( <i>Brachyramphus marmoratus</i> ) - U.S.A. (CA, OR, WA)	T	\$4,771	\$0	\$4,771	\$0	\$4,771
	7	Plover, piping ( <i>Charadrius melodus</i> ) - [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	T	\$3,264,343	\$6,530	\$3,270,873	\$55,500	\$3,326,373
	80	Scrub-jay, Florida ( <i>Aphelocoma coerulescens</i> ) - Wherever found	T	\$3,913	\$42,203	\$46,116	\$0	\$46,116
	21	Sparrow, Florida grasshopper ( <i>Ammodramus savannarum floridanus</i> ) - Wherever found	E	\$0	\$1,036,255	\$1,036,255	\$0	\$1,036,255
	87	Stilt, Hawaiian ( <i>Himantopus mexicanus knudseni</i> ) - Wherever found	E	\$25,995	\$0	\$25,995	\$0	\$25,995
	3	Stork, wood ( <i>Mycteria americana</i> ) - AL, FL, GA, MS, NC, SC	T	\$171,269	\$4,000,000	\$4,171,269	\$0	\$4,171,269
	30	Tern, least ( <i>Sterna antillarum</i> ) - interior pop.	E	\$623,243	\$41,530	\$664,773	\$0	\$664,773
	57	Tern, roseate ( <i>Sterna dougallii dougallii</i> ) - Northeast U.S. nesting population	E	\$226,500	\$0	\$226,500	\$0	\$226,500
	85	Vireo, least Bell's ( <i>Vireo bellii pusillus</i> ) - Wherever found	E	\$26,385	\$0	\$26,385	\$0	\$26,385
	44	Woodpecker, red-cockaded ( <i>Picoides borealis</i> ) - Wherever found	E	\$57,356	\$353,328	\$410,684	\$0	\$410,684
<b>Birds subtotal</b>				<b>\$9,587,902</b>	<b>\$10,039,985</b>	<b>\$19,627,887</b>	<b>\$113,202</b>	<b>\$19,741,089</b>
Reptiles	94	Alligator, American ( <i>Alligator mississippiensis</i> ) - Wherever found	SAT	\$0	\$2,758	\$2,758	\$0	\$2,758
	60	Lizard, blunt-nosed leopard ( <i>Gambelia silus</i> ) - Wherever found	E	\$0	\$207,500	\$207,500	\$0	\$207,500



Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	92	Sea turtle, green ( <i>Chelonia mydas</i> ) - East Pacific DPS	T	\$4,771	\$0	\$4,771	\$0	\$4,771
	11	Sea turtle, green ( <i>Chelonia mydas</i> ) - North Atlantic DPS	T	\$2,165,556	\$0	\$2,165,556	\$0	\$2,165,556
	15	Sea turtle, hawksbill ( <i>Eretmochelys imbricata</i> ) - Wherever found	E	\$2,000,000	\$0	\$2,000,000	\$0	\$2,000,000
	9	Sea turtle, Kemp's ridley ( <i>Lepidochelys kempii</i> ) - Wherever found	E	\$2,481,848	\$0	\$2,481,848	\$0	\$2,481,848
	10	Sea turtle, leatherback ( <i>Dermochelys coriacea</i> ) - Wherever found	E	\$2,170,327	\$0	\$2,170,327	\$0	\$2,170,327
	93	Skink, bluetail mole ( <i>Eumeces egregius lividus</i> ) - Wherever found	T	\$3,913	\$0	\$3,913	\$0	\$3,913
	93	Skink, sand ( <i>Neoseps reynoldsi</i> ) - Wherever found	T	\$3,913	\$0	\$3,913	\$0	\$3,913
	89	Snake, copperbelly water ( <i>Nerodia erythrogaster neglecta</i> ) - Indiana north of 40 degrees north latitude, Michigan, Ohio	T	\$11,667	\$0	\$11,667	\$0	\$11,667
	4	Snake, eastern indigo ( <i>Drymarchon corais couperi</i> ) - Wherever found	T	\$3,913	\$2,000,000	\$2,003,913	\$2,000,000	\$4,003,913
	47	Tortoise, desert ( <i>Gopherus agassizii</i> ) - Wherever found, except AZ south and east of Colorado R., and Mexico	T	\$0	\$365,725	\$365,725	\$0	\$365,725
	81	Turtle, bog ( <i>Clemmys muhlenbergii</i> ) - Wherever found, except GA, NC, SC, TN, VA	T	\$17,500	\$22,000	\$39,500	\$0	\$39,500
<b>Reptiles subtotal</b>				<b>\$8,863,408</b>	<b>\$2,597,983</b>	<b>\$11,461,391</b>	<b>\$2,000,000</b>	<b>\$13,461,391</b>
Amphibians	83	Frog, Chiricahua leopard ( <i>Rana chiricahuensis</i> ) - Wherever found	T	\$30,000	\$0	\$30,000	\$0	\$30,000
	26	Salamander, California tiger ( <i>Ambystoma californiense</i> ) - U.S.A. (Central CA DPS)	T	\$0	\$728,000	\$728,000	\$0	\$728,000
	66	Salamander, Cheat Mountain ( <i>Plethodon nettingi</i> ) - Wherever found	T	\$149,000	\$0	\$149,000	\$0	\$149,000
	65	Salamander, frosted flatwoods ( <i>Ambystoma cingulatum</i> ) - Wherever found	T	\$55,556	\$109,028	\$164,584	\$0	\$164,584
	98	Salamander, Reticulated flatwoods ( <i>Ambystoma bishopi</i> ) - Wherever found	E	\$0	\$1,101	\$1,101	\$0	\$1,101
<b>Amphibians subtotal</b>				<b>\$234,556</b>	<b>\$838,129</b>	<b>\$1,072,685</b>	<b>\$0</b>	<b>\$1,072,685</b>
Fishes	92	Eulachon ( <i>Thaleichthys pacificus</i> ) - Southern DPS	T	\$4,771	\$0	\$4,771	\$0	\$4,771

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	83	Gambusia, Pecos ( <i>Gambusia nobilis</i> ) - Wherever found	E	\$0	\$30,000	\$30,000	\$0	\$30,000
	20	Minnow, Rio Grande Silvery ( <i>Hybognathus amarus</i> ) - Wherever found, except where listed as an experimental population	E	\$0	\$1,200,000	\$1,200,000	\$0	\$1,200,000
	83	Pupfish, Comanche Springs ( <i>Cyprinodon elegans</i> ) - Wherever found	E	\$0	\$30,000	\$30,000	\$0	\$30,000
	38	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i> ) - Snake River fall-run ESU	T	\$0	\$582,500	\$582,500	\$0	\$582,500
	38	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i> ) - Snake River spring/summer-run ESU	T	\$0	\$582,500	\$582,500	\$0	\$582,500
	75	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i> ) - Sacramento River winter-run ESU	E	\$0	\$66,000	\$66,000	\$0	\$66,000
	45	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i> ) - Puget Sound ESU	T	\$0	\$387,341	\$387,341	\$0	\$387,341
	31	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i> ) - Lower Columbia River ESU	T	\$61,250	\$586,632	\$647,882	\$0	\$647,882
	18	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i> ) - Upper Willamette River ESU	T	\$0	\$1,459,803	\$1,459,803	\$0	\$1,459,803
	37	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i> ) - Upper Columbia spring-run ESU	E	\$0	\$602,978	\$602,978	\$0	\$602,978
	59	Salmon, Chinook ( <i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i> ) - Central Valley spring-run ESU	T	\$0	\$217,500	\$217,500	\$0	\$217,500
	13	Salmon, coho ( <i>Oncorhynchus</i> (=Salmo) <i>kisutch</i> ) - Lower Columbia River ESU	T	\$61,250	\$2,029,372	\$2,090,622	\$0	\$2,090,622
	38	Salmon, sockeye ( <i>Oncorhynchus</i> (=Salmo) <i>nerka</i> ) - Snake River ESU	E	\$0	\$582,500	\$582,500	\$0	\$582,500
	61	Shiner, Pecos bluntnose ( <i>Notropis simus pecosensis</i> ) - Wherever found	T	\$0	\$200,000	\$200,000	\$0	\$200,000
	29	Shiner, Topeka ( <i>Notropis topeka</i> (=tristis)) - Wherever found, except where listed as an experimental population	E	\$674,733	\$0	\$674,733	\$0	\$674,733
	59	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - California Central Valley DPS	T	\$0	\$217,500	\$217,500	\$0	\$217,500
	38	Steelhead ( <i>Oncorhynchus</i> (=Salmo) <i>mykiss</i> ) - Snake River Basin DPS	T	\$0	\$582,500	\$582,500	\$0	\$582,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	33	Steelhead (Oncorhynchus (=Salmo) mykiss) - Lower Columbia River DPS	T	\$61,250	\$582,500	\$643,750	\$0	\$643,750
	38	Steelhead (Oncorhynchus (=Salmo) mykiss) - Upper Columbia River DPS	T	\$0	\$582,500	\$582,500	\$0	\$582,500
	12	Steelhead (Oncorhynchus (=Salmo) mykiss) - Upper Willamette River DPS	T	\$0	\$2,133,205	\$2,133,205	\$0	\$2,133,205
	38	Steelhead (Oncorhynchus (=Salmo) mykiss) - Middle Columbia River DPS	T	\$0	\$582,500	\$582,500	\$0	\$582,500
	45	Steelhead (Oncorhynchus (=Salmo) mykiss) - Puget Sound DPS	T	\$0	\$387,341	\$387,341	\$0	\$387,341
	42	Sturgeon, Atlantic (Acipenser oxyrinchus oxyrinchus) - Gulf of Maine DPS	T	\$415,300	\$0	\$415,300	\$0	\$415,300
	92	Sturgeon, green (Acipenser medirostris) - Southern DPS	T	\$4,771	\$0	\$4,771	\$0	\$4,771
	27	Sturgeon, pallid (Scaphirhynchus albus) - Wherever found	E	\$531,977	\$165,990	\$697,967	\$0	\$697,967
	70	Sturgeon, shortnose (Acipenser brevirostrum) - Wherever found	E	\$111,800	\$0	\$111,800	\$0	\$111,800
	6	Sucker, Lost River (Deltistes luxatus) - Wherever found	E	\$3,557,000	\$0	\$3,557,000	\$0	\$3,557,000
	6	Sucker, shortnose (Chasmistes brevirostris) - Wherever found	E	\$3,557,000	\$0	\$3,557,000	\$0	\$3,557,000
	1	Trout, bull (Salvelinus confluentus) - U.S.A., conterminous, lower 48 states	T	\$522,667	\$13,000,509	\$13,523,176	\$590,000	\$14,113,176
<b>Fishes subtotal</b>				<b>\$9,563,769</b>	<b>\$26,791,671</b>	<b>\$36,355,440</b>	<b>\$590,000</b>	<b>\$36,945,440</b>
Clams	79	Bankclimber, purple (mussel) (Elliptioideus sloatianus) - Wherever found	T	\$55,556	\$0	\$55,556	\$0	\$55,556
	64	Fatmucket, Arkansas (Lampsilis powellii) - Wherever found	T	\$0	\$0	\$0	\$174,000	\$174,000
	48	Higgins eye (pearlymussel) (Lampsilis higginsii) - Wherever found	E	\$362,683	\$0	\$362,683	\$0	\$362,683
	73	Lance, yellow (Elliptio lanceolata) - Wherever found	T	\$0	\$0	\$0	\$77,600	\$77,600
	72	Mucket, pink (pearlymussel) (Lampsilis abrupta) - Wherever found	E	\$79,000	\$0	\$79,000	\$0	\$79,000
	48	Mussel, sheepnose (Plethobasus cyphus) - Wherever found	E	\$362,683	\$0	\$362,683	\$0	\$362,683
	82	Mussel, snuffbox (Epioblasma triquetra) - Wherever found	E	\$0	\$0	\$0	\$38,500	\$38,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	72	Pocketbook, fat (Potamilus capax) - Wherever found	E	\$79,000	\$0	\$79,000	\$0	\$79,000
	53	Pocketbook, Ouachita rock (Arkansia wheeleri) - Wherever found	E	\$270,200	\$0	\$270,200	\$0	\$270,200
	48	Spectaclecase (mussel) (Cumberlandia monodonta) - Wherever found	E	\$362,683	\$0	\$362,683	\$0	\$362,683
	96	Spinymussel, James (Pleurobema collina) - Wherever found	E	\$0	\$0	\$0	\$2,100	\$2,100
	73	Spinymussel, Tar River (Elliptio steinstansana) - Wherever found	E	\$0	\$0	\$0	\$77,600	\$77,600
	49	Wedgemussel, dwarf (Alasmidonta heterodon) - Wherever found	E	\$256,750	\$0	\$256,750	\$77,665	\$334,415
<b>Clams subtotal</b>				<b>\$1,828,555</b>	<b>\$0</b>	<b>\$1,828,555</b>	<b>\$447,465</b>	<b>\$2,276,020</b>
Insects	78	Beetle, American burying (Nicrophorus americanus) - Wherever found, except where listed as an experimental population	E	\$0	\$0	\$0	\$60,000	\$60,000
	85	Butterfly, Quino checkerspot (Euphydryas editha quino (=E. e. wrighti)) - Wherever found	E	\$26,385	\$0	\$26,385	\$0	\$26,385
	35	Damselfly, Pacific Hawaiian (Megalagrion pacificum) - Wherever found	E	\$620,476	\$0	\$620,476	\$0	\$620,476
	35	Fly, Hawaiian picture-wing (Drosophila heteroneura) - Wherever found	E	\$620,476	\$0	\$620,476	\$0	\$620,476
	74	Moth, Kern primrose sphinx (Euproserpinus euterpe) - Wherever found	T	\$0	\$75,000	\$75,000	\$0	\$75,000
	54	Tiger beetle, Puritan (Cicindela puritana) - Wherever found	T	\$256,750	\$0	\$256,750	\$0	\$256,750
	69	Tiger beetle, Salt Creek (Cicindela nevadica lincolniana) - Wherever found	E	\$0	\$0	\$0	\$128,300	\$128,300
<b>Insects subtotal</b>				<b>\$1,524,087</b>	<b>\$75,000</b>	<b>\$1,599,087</b>	<b>\$188,300</b>	<b>\$1,787,387</b>
Crustaceans	85	Fairy shrimp, San Diego (Branchinecta sandiegonensis) - Wherever found	E	\$26,385	\$0	\$26,385	\$0	\$26,385
<b>Crustaceans subtotal</b>				<b>\$26,385</b>	<b>\$0</b>	<b>\$26,385</b>	<b>\$0</b>	<b>\$26,385</b>
Flowering Plants	35	'Aiea (Nothoecstrum breviflorum) - Wherever found	E	\$620,476	\$0	\$620,476	\$0	\$620,476
	85	Ambrosia, San Diego (Ambrosia pumila) - Wherever found	E	\$26,385	\$0	\$26,385	\$0	\$26,385
	93	Beargrass, Britton's (Nolina brittoniana) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	93	Blazingstar, scrub ( <i>Liatris ohlingerae</i> ) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
	83	Blue-star, Kearney's ( <i>Amsonia kearneyana</i> ) - Wherever found	E	\$30,000	\$0	\$30,000	\$0	\$30,000
	93	Bonamia, Florida ( <i>Bonamia grandiflora</i> ) - Wherever found	T	\$3,913	\$0	\$3,913	\$0	\$3,913
	97	Brodiaea, Chinese Camp ( <i>Brodiaea pallida</i> ) - Wherever found	T	\$0	\$2,000	\$2,000	\$0	\$2,000
	93	Buckwheat, scrub ( <i>Eriogonum longifolium</i> var. <i>gnaphalifolium</i> ) - Wherever found	T	\$3,913	\$0	\$3,913	\$0	\$3,913
	52	Bulrush, Northeastern ( <i>Scirpus ancistrochaetus</i> ) -	E	\$274,250	\$0	\$274,250	\$0	\$274,250
	85	Button-celery, San Diego ( <i>Eryngium aristulatum</i> var. <i>parishii</i> ) - Wherever found	E	\$26,385	\$0	\$26,385	\$0	\$26,385
	83	Cactus, Pima pineapple ( <i>Coryphantha scheeri</i> var. <i>robustispina</i> ) - Wherever found	E	\$30,000	\$0	\$30,000	\$0	\$30,000
	58	Catchfly, Spalding's ( <i>Silene spaldingii</i> ) - Wherever found	T	\$0	\$0	\$0	\$221,000	\$221,000
	77	Checker-mallow, Nelson's ( <i>Sidalcea nelsoniana</i> ) - Wherever found	T	\$61,250	\$0	\$61,250	\$0	\$61,250
	88	Dropwort, Canby's ( <i>Oxypolis canbyi</i> ) - Wherever found	E	\$0	\$0	\$0	\$20,000	\$20,000
	85	Flannelbush, Mexican ( <i>Fremontodendron mexicanum</i> ) - Wherever found	E	\$26,385	\$0	\$26,385	\$0	\$26,385
	93	Fringe-tree, pygmy ( <i>Chionanthus pygmaeus</i> ) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
	8	Haha ( <i>Cyanea hamatiflora</i> ssp. <i>carlsonii</i> ) - Wherever found	E	\$2,481,905	\$0	\$2,481,905	\$0	\$2,481,905
	35	Ha'iwale ( <i>Cyrtandra tintinnabula</i> ) - Wherever found	E	\$620,476	\$0	\$620,476	\$0	\$620,476
	90	Howellia, Water ( <i>Howellia aquatilis</i> ) -	T	\$7,500	\$0	\$7,500	\$0	\$7,500
	93	Hypericum, highlands scrub ( <i>Hypericum cumulicola</i> ) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
	67	Iris, dwarf lake ( <i>Iris lacustris</i> ) - Wherever found	T	\$147,000	\$0	\$147,000	\$0	\$147,000
	43	Joint-vetch, Sensitive ( <i>Aeschynomene virginica</i> ) - Wherever found	T	\$413,500	\$0	\$413,500	\$0	\$413,500
	35	Kiponapona ( <i>Phyllostegia racemosa</i> ) - Wherever found	E	\$620,476	\$0	\$620,476	\$0	\$620,476

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	93	Lupine, scrub ( <i>Lupinus aridorum</i> ) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
	46	Mallow, Kern ( <i>Eremalche kernensis</i> ) - Wherever found	E	\$300,000	\$75,000	\$375,000	\$0	\$375,000
	85	Mesa-mint, Otay ( <i>Pogogyne nudiuscula</i> ) - Wherever found	E	\$26,385	\$0	\$26,385	\$0	\$26,385
	85	Mesa-mint, San Diego ( <i>Pogogyne abramsii</i> ) - Wherever found	E	\$26,385	\$0	\$26,385	\$0	\$26,385
	93	Mint, Garrett's ( <i>Dicerandra christmanii</i> ) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
	93	Mustard, Carter's ( <i>Warea carteri</i> ) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
	85	Navarretia, spreading ( <i>Navarretia fossalis</i> ) - Wherever found	T	\$26,385	\$0	\$26,385	\$0	\$26,385
	19	No Common Name ( <i>Phyllostegia velutina</i> ) - Wherever found	E	\$1,240,952	\$0	\$1,240,952	\$0	\$1,240,952
	87	Ohai ( <i>Sesbania tomentosa</i> ) - Wherever found	E	\$25,995	\$0	\$25,995	\$0	\$25,995
	16	'Oha wai ( <i>Clermontia lindseyana</i> ) - Wherever found	E	\$1,861,429	\$0	\$1,861,429	\$0	\$1,861,429
	56	Orchid, western prairie fringed ( <i>Platanthera praeclara</i> ) - Wherever found	T	\$229,338	\$0	\$229,338	\$0	\$229,338
	85	Orcutt grass, California ( <i>Orcuttia californica</i> ) - Wherever found	E	\$26,385	\$0	\$26,385	\$0	\$26,385
	93	Pigeon wings ( <i>Clitoria fragrans</i> ) - Wherever found	T	\$3,913	\$0	\$3,913	\$0	\$3,913
	93	Plum, scrub ( <i>Prunus geniculata</i> ) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
	35	Po'e ( <i>Portulaca sclerocarpa</i> ) - Wherever found	E	\$620,476	\$0	\$620,476	\$0	\$620,476
	93	Polygala, Lewton's ( <i>Polygala lewtonii</i> ) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
	93	Sandlace ( <i>Polygonella myriophylla</i> ) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
	85	Tarplant, Otay ( <i>Deinandra</i> (=Hemizonia) <i>conjugens</i> ) - Wherever found	T	\$26,385	\$0	\$26,385	\$0	\$26,385
	67	Thistle, Pitcher's ( <i>Cirsium pitcheri</i> ) - Wherever found	T	\$147,000	\$0	\$147,000	\$0	\$147,000
	85	Thornmint, San Diego ( <i>Acanthomintha ilicifolia</i> ) - Wherever found	T	\$26,385	\$0	\$26,385	\$0	\$26,385

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	93	Whitlow-wort, papery (Paronychia chartacea) - Wherever found	T	\$3,913	\$0	\$3,913	\$0	\$3,913
	91	Wild-buckwheat, gypsum (Eriogonum gypsophilum) - Wherever found	T	\$0	\$7,000	\$7,000	\$0	\$7,000
	93	Wireweed (Polygonella basiramia) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
	93	Ziziphus, Florida (Ziziphus celata) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
<b>Flowering Plants subtotal</b>				<b>\$10,032,096</b>	<b>\$84,000</b>	<b>\$10,116,096</b>	<b>\$241,000</b>	<b>\$10,357,096</b>
Lichens	93	Cladonia, Florida perforate (Cladonia perforata) - Wherever found	E	\$3,913	\$0	\$3,913	\$0	\$3,913
<b>Lichens subtotal</b>				<b>\$3,913</b>	<b>\$0</b>	<b>\$3,913</b>	<b>\$0</b>	<b>\$3,913</b>
Multi-species				\$0	\$2,676,999	\$2,676,999	\$44,300	\$2,721,299
<b>Total Land Expenditures</b>				<b>\$45,124,520</b>	<b>\$60,344,976</b>	<b>\$105,469,496</b>	<b>\$6,001,410</b>	<b>\$111,470,906</b>

**TABLE 4. FY 2018 EXPENDITURES REPORTED BY INDIVIDUAL FEDERAL AGENCIES AND THE STATES**

This table displays combined species expenditures and land acquisition costs by each reporting Federal agency, the Service, and the States collectively. Expenditures for foreign species are included, as well as costs for multi-species conservation activities.



**TABLE 4. FY 2018 EXPENDITURES REPORTED BY INDIVIDUAL FEDERAL AGENCIES  
AND THE STATES**


<b>Agency Type</b>	<b>Expense Type</b>	<b>Agency</b>	<b>Species Total</b>	<b>Land Total</b>	<b>Cumulative Total</b>
Federal	Federal Domestic Expenditures	DHS Coast Guard	\$382,362	\$0	\$382,362
		DHS Customs and Border Protection	\$323,300	\$0	\$323,300
		DOC National Oceanic and Atmospheric Administration	\$197,913,000	\$0	\$197,913,000
		DOD Army	\$10,482,107	\$0	\$10,482,107
		DOD Army Corps of Engineers	\$190,660,844	\$1,341,073	\$192,001,917
		DOD Marine Corps	\$13,407,400	\$0	\$13,407,400
		DOD Navy	\$17,643,011	\$0	\$17,643,011
		DOE Bonneville Power Administration	\$239,774,729	\$18,260,158	\$258,034,887
		DOE Southwestern Power Administration	\$39,649	\$35,000	\$74,649
		DOE Western Area Power Administration	\$2,439,667	\$0	\$2,439,667
		DOI Bureau of Indian Affairs	\$2,302,875	\$0	\$2,302,875
		DOI Bureau of Land Management	\$23,226,857	\$433,333	\$23,660,190
		DOI Bureau of Reclamation	\$160,049,000	\$9,732,000	\$169,781,000
		DOI National Park Service	\$17,982,068	\$0	\$17,982,068
		DOI US Geological Survey	\$10,361,301	\$0	\$10,361,301
		DOT Federal Aviation Administration	\$176,000	\$0	\$176,000
		Florida Fish & Wildlife Conservation Commission	\$8,830,282	\$547,413	\$9,377,695
		State Expenditures Data Entry	\$33,000	\$0	\$33,000
		Tennessee Valley Authority	\$1,663,700	\$0	\$1,663,700
		USDA Animal and Plant Health Inspection Service	\$10,236,200	\$0	\$10,236,200
		USDA Forest Service	\$62,746,067	\$0	\$62,746,067
		USDA Natural Resources Conservation Service	\$21,963,493	\$27,319,000	\$49,282,493
		US Fish and Wildlife Service	\$141,076,464	\$45,124,520	\$186,200,984
		<i>Federal Domestic Expenditures subtotal</i>	<i>\$1,133,713,376</i>	<i>\$102,792,497</i>	<i>\$1,236,505,873</i>
	Federal Foreign Expenditures	USDA Animal and Plant Health Inspection Service	\$33,800	\$0	\$33,800
		DOI National Park Service	\$41,000	\$0	\$41,000
		DOI US Geological Survey	\$55,200	\$0	\$55,200
		DOD Army Corps of Engineers	\$590,632	\$0	\$590,632
		US Fish and Wildlife Service	\$11,032,410	\$0	\$11,032,410
		<i>Federal Foreign Expenditures subtotal</i>	<i>\$11,753,042</i>	<i>\$0</i>	<i>\$11,753,042</i>
		<b><i>Federal Domestic and Foreign Expenditures Subtotal</i></b>	<b><i>\$1,145,466,418</i></b>	<b><i>\$102,792,497</i></b>	<b><i>\$1,248,258,915</i></b>

Multi-Agency	Federal Multi-species Expenditures	DOI Bureau of Reclamation	\$65,007,500	\$2,602,500	\$67,610,000
		DOI National Park Service	\$11,536	\$0	\$11,536
		USDA Animal and Plant Health Inspection Service	\$365,480	\$0	\$365,480
		DOD Army	\$30,781,382	\$0	\$30,781,382
		DOD Army Corps of Engineers	\$1,165,366	\$0	\$1,165,366
		DOT Federal Aviation Administration	\$89,800	\$0	\$89,800
		DOD Navy	\$507,786	\$0	\$507,786
		USDA Natural Resources Conservation Service	\$2,403,828	\$0	\$2,403,828
		DHS Coast Guard	\$32,289,839	\$0	\$32,289,839
		DOE Southwestern Power Administration	\$17,900	\$0	\$17,900
		Federal Communications Commission	\$127,900	\$0	\$127,900
		Florida Fish & Wildlife Conservation Commission	\$1,193,110	\$74,499	\$1,267,609
		US Fish and Wildlife Service	\$2,682,138	\$0	\$2,682,138
	<i>Federal Multi-species Expenditures subtotal</i>		<i>\$136,643,565</i>	<i>\$2,676,999</i>	<i>\$139,320,564</i>
	<b><i>Federal Domestic and Foreign Expenditures subtotal</i></b>		<b><i>\$1,282,109,983</i></b>	<b><i>\$105,469,496</i></b>	<b><i>\$1,387,579,479</i></b>
State	State Domestic Expenditures	State Governments	\$124,187,637	\$5,957,110	\$130,144,747
	State Multi-species Expenditures	State Governments	\$1,559,778	\$44,300	\$1,604,078
	<b><i>State Domestic and Foreign Expenditures Subtotal</i></b>		<b><i>\$125,747,415</i></b>	<b><i>\$6,001,410</i></b>	<b><i>\$131,748,825</i></b>
<b>Total Federal and State Domestic and Foreign Expenditures</b>			<b>\$1,407,857,398</b>	<b>\$111,470,906</b>	<b>\$1,519,328,304</b>


## Appendix A. Supplemental Guidance for Reporting Expenditures for the Conservation of Endangered and Threatened Species -- FY 2018

This reporting guidance provides information on how to report expenditures attributable to individual species to the extent this information is reasonably available. The Congress requested information for this report on a species-by-species basis, but to provide the most accurate account of expenditures for the conservation of species listed under the Endangered Species Act, a new reporting feature was added in FY 2012. This “multi-species” option allows the reporting of lump sum costs for multi-species conservation activities **when it is not possible to account accurately for individual species.**

Each agency must enter data directly into our Web-based system and will need to designate a contact person (more than one person is permissible) to enter the data. The contact person(s) will need to obtain a username and password before entering the data. Credentials can be obtained by accessing our Environmental Conservation Online System (ECOS) home page at <http://ecos.fws.gov/> and clicking on “Contact Us” located at the bottom of the ECOS home page. This brings you the Help Desk where you will click on “help desk web form.”



Fill in the requested information along with your affiliated agency and request access to the “Expenditures” application.



The screenshot shows a web browser window with the URL <https://ecos.fws.gov/ecp/helpTickets/create>. The page header includes the U.S. Fish & Wildlife Service logo and the text "ECOS Environmental Conservation Online System" and "Conserving the Nature of America". The main content area is titled "ECOS Help Desk" and contains a form with the following fields:

- Name: \* John Doe
- Email: \* john\_doe@usgs.gov
- Phone: \* 555-555-5555
- Issue Title: \* Need to enter expenditures
- Issue Description: \* I am John Doe with the U.S. Geological Survey and I need access to the Expenditures module.

To the right of the form is a box titled "Are you contacting the correct help desk?" with the following text: "Note that this form will contact the *web page technical and program staff*. If you have a question about the FWS or Wildlife in general, [contact the FWS help desk](#)."

Please allow 2 working days for access approval (an email confirmation will be sent to you).

Once you receive your log-in credentials and access approval email, log in at

<https://ecos.fws.gov> to begin entering expenditures data. Contact the ECOS Help Desk (970-266-2999) if you experience any problems.

The following information should be used to gather the data.

## BACKGROUND

- Section 18 of the Endangered Species Act of 1973 (16 U.S. 1531 *et seq.*) reads, in part, “...the Fish and Wildlife Service, shall submit to the Congress an annual report covering the preceding fiscal year which shall contain--
  - (1) an accounting on a species-by-species basis of all reasonably identifiable Federal expenditures made primarily for the conservation of endangered and threatened species pursuant to this Act; and
  - (2) an accounting on a species-by-species basis for all reasonably identifiable expenditures made primarily for the conservation of endangered and threatened species pursuant to this Act by States receiving grants under section 6.”
- The conference report for the 1988 amendments made some of the following points concerning section 18 of the Act:
  - Provision should not become unduly burdensome or result in diversion of funds from the operation of the program.

- The Secretary of the Interior is to make a good faith effort to develop and obtain data but is not to undertake extensive or extraordinary measures.
- Generalized dollar estimates will suffice.
- Amendment does not apply to expenditures for delisted, proposed, and candidate species.
- Amendment is not to require new species-specific time sheets for the Service or other reporting agency.
- Listing, consultation, and law enforcement are to be included to the extent they are "reasonably identifiable."
- Only salary costs that can be reasonably identified as applicable to listed species need be reported.
- For other Federal and State agencies, a good faith effort to collect information is expected.
- Submission of data from any agency is not to be a precondition to receiving any contracts or grants or establishing other arrangements with the Fish and Wildlife Service.
- Only one data set per agency should be submitted for any given fiscal year.

#### DATA FORMAT

- Species should be identified by the same scientific name as found in the most current list of endangered and threatened species.
- Amounts should be reported in whole dollars. Amounts above \$2,000 may be rounded to the nearest \$500 or \$1,000; smaller amounts may be rounded to the nearest \$100. **Do not report amounts lower than \$100.**

#### EXPENDITURES TO BE REPORTED

- Only species on the Federal list of Endangered and Threatened Wildlife and Plants (50 CFR Part 17) at the end of Federal fiscal year 2018 (October 1, 2017 to September 30, 2018) are to be reported. Expenditures made prior to the actual listing date of a species, but still within the same year, may be reported (e.g., costs of public meetings, notices, surveys, initial recovery efforts). Funds expended for unlisted, separate populations of listed vertebrates cannot be included in the report (e.g., Atlantic coast least terns, Alaska gray wolves). Expenditures for State-listed species are not reportable unless they are also federally listed. Amounts for foreign species on the list (e.g., grants or contracts carried

out in another country) are reportable in the same manner as domestic species, but again, to the extent they are reasonably identifiable.

- Only reasonably identifiable expenditures for listed species will be compiled for this report. Extraordinary accounting to track funds expended on listed species is not expected. **Funds reported should be those actually expended, not obligated.**
- Examples of reportable expenditures include status surveys, habitat management, habitat acquisition, research, propagation (including surrogate species), recovery plan development or implementation, and mitigation. **Projects must primarily benefit listed species** and not other conservation goals, although if a readily identifiable portion of the expenditure is for the benefit of listed species (either individually or collectively), that portion of the expenditure may be reported.
- All habitat (land or water) acquisition costs are to be reported separately from all other identifiable expenditures. Such acquisitions must be primarily for the purpose of conserving federally listed species. As with other projects, those portions of habitat acquisition costs that are specifically for a listed species may be prorated out of the total acquisition costs when the other habitat is being acquired for other purposes (e.g., 25 acres of a 250-acre purchase will be set aside).
- Expenditures in a single project devoted to a number of listed species should be prorated by species, if possible. **When it is not possible to determine individual listed species costs in a multi-species project by prorating, those costs should be entered, in total, in the multi-species category option. The comment box, which is part of this feature, should be used to include information about the project.**
- General surveys or projects that cover a large number of species, some of which may not be listed, are reportable only to extent that the agency can reasonably identify the portion of costs that benefit listed species.
- Expenditures associated with consultations pursuant to section 7 of the Act should be reported as for the benefit of a specific species to the extent they are readily identifiable to that particular species. Thus, a formal consultation dealing with a single species, or up to several species where the relative costs are easily divisible, should be reported under that or those species. **Consultation costs not readily attributable to individual species can be reported in the multi-species category.**
- Monitoring and survey costs should be reported when: (1) the cost of the monitoring or survey was incurred during the period the species was under a proposed listing and the species was actually listed in the same fiscal year that the monitoring and surveys were conducted; (2) the monitoring or survey covered both candidates and listed species, and the costs can be prorated to include only the listed species; or (3) the cost was readily identifiable to individual species. **Costs meeting the criteria in (1) and (2), but not readily identifiable to individual species can be reported in the multi-species category.**

- Salary and benefits of an employee working full-time on a single species or whose time is devoted to particular species that can be readily identified should be reported for those species. Travel costs should be treated similarly. **Staff costs for work on multi-species activities that cannot be identified to particular species can be reported in the multi-species category.**
- Any State or Federal project that incurs increased costs related directly to mitigation or other conservation efforts on behalf of federally listed species should report that added cost, by species, if readily identifiable, or **may report such costs in the multi-species category.**
- All State agencies (e.g., fish and wildlife, parks, heritage programs, forests, highways) should report their expenditures, but only a single, collective report from each State should be submitted. **Expenditures should include conservation costs for fish, wildlife, and plants.** For ESA section 6 grants or other Federal grants, States should report **only the State cost-share portion** of each grant. States should not include other sources of non-Federal cost share, e.g., third-party contributions, unless those contributions are donated to the State as a general donation.
- For reimbursable funds, the paying or disbursing agency should report the expenditure.
- **Litigation costs should not be reported.**

## **Appendix B. Reports Provided by Other Federal Agencies for Fiscal Year 2018**

### Department of Agriculture:

- Animal and Plant Health Inspection Service
- Farm Service Agency\*
- Forest Service
- Natural Resources Conservation Service

### Department of Commerce:

- National Oceanic and Atmospheric Administration

### Department of Defense:

- Office of the Under Secretary
  - Representing: Air Force\*, Army (includes Defense Logistics Agency), Marine Corps, Navy, and Army Corps of Engineers

### Department of Energy:

- Bonneville Power Administration
- Southeastern Power Administration\*
- Southwestern Power Administration
- Western Area Power Administration

### Department of Homeland Security

- Coast Guard
- Customs and Border Protection
- Federal Emergency Management Agency\*
- Office of Science and Technology\*

### Department of the Interior:

- Bureau of Indian Affairs
- Bureau of Land Management
- Bureau of Ocean Energy Management\*
- Bureau of Reclamation
- Bureau of Safety and Environmental Enforcement\*
- National Park Service
- Office of Surface Mining\*
- U.S. Geological Survey

### Department of Transportation:

- Federal Aviation Administration
- Federal Highway Administration\*

### Independent Agencies:



Environmental Protection Agency\*  
Federal Communications Commission  
Federal Energy Regulatory Commission\*  
Nuclear Regulatory Commission  
Smithsonian Institution\*  
Tennessee Valley Authority

\* Agencies with no expenditures data submitted for FY 2018

## DHS Coast Guard - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Manatee, West Indian	Trichechus manatus	Wherever found	\$10,000	\$0	\$10,000
		Sea lion, Steller	Eumetopias jubatus	Western DPS	\$900	\$0	\$900
		Whale, North Atlantic Right	Eubalaena glacialis	Wherever found	\$357,469	\$0	\$357,469
	<i>Mammals Subtotal</i>				<i>\$368,369</i>	<i>\$0</i>	<i>\$368,369</i>
	Reptiles						
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$7,996	\$0	\$7,996
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$5,997	\$0	\$5,997
	<i>Reptiles Total</i>				<i>\$13,993</i>	<i>\$0</i>	<i>\$13,993</i>
	Multi-Species						\$32,289,839
	<i>Multi-Species Subtotal</i>				<i>\$32,289,839</i>	<i>\$0</i>	<i>\$32,289,839</i>
	<i>Domestic Total</i>				<i>\$32,672,201</i>	<i>\$0</i>	<i>\$32,672,201</i>
	<b>Grand Total</b>				<b>\$32,672,201</b>	<b>\$0</b>	<b>\$32,672,201</b>

## DHS Customs and Border Protection - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Birds	Gnatcatcher, coastal California	Polioptila californica californica	Wherever found	\$12,000	\$0	\$12,000
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$19,000	\$0	\$19,000
		<i>Birds Subtotal</i>			<i>\$31,000</i>	<i>\$0</i>	<i>\$31,000</i>
	Insects	Butterfly, Quino checkerspot	Euphydryas editha quino (=E. e. wrighti)	Wherever found	\$90,000	\$0	\$90,000
		<i>Insects Subtotal</i>					<i>\$90,000</i>
	Crustaceans	Fairy shrimp, Riverside	Streptocephalus woottoni	Wherever found	\$17,500	\$0	\$17,500
		Fairy shrimp, San Diego	Branchinecta sandiegonensis	Wherever found	\$17,500	\$0	\$17,500
		<i>Crustaceans Subtotal</i>			<i>\$35,000</i>	<i>\$0</i>	<i>\$35,000</i>
	Flowering Plants	Ambrosia, San Diego	Ambrosia pumila	Wherever found	\$12,000	\$0	\$12,000
		Baccharis, Encinitas	Baccharis vanessae	Wherever found	\$12,000	\$0	\$12,000
		Button-celery, San Diego	Eryngium aristulatum var. parishii	Wherever found	\$12,000	\$0	\$12,000
		Cactus, Pima pineapple	Coryphantha scheeri var. robustispina	Wherever found	\$20,200	\$0	\$20,200
		Flannelbush, Mexican	Fremontodendron mexicanum	Wherever found	\$12,000	\$0	\$12,000
		Mesa-mint, Otay	Pogogyne nudiuscula	Wherever found	\$12,000	\$0	\$12,000
		Navarretia, spreading	Navarretia fossalis	Wherever found	\$12,000	\$0	\$12,000
		Orcutt grass, California	Orcuttia californica	Wherever found	\$12,000	\$0	\$12,000
		Tarplant, Otay	Deinandra (=Hemizonia) conjugens	Wherever found	\$51,100	\$0	\$51,100
		Thornmint, San Diego	Acanthomintha ilicifolia	Wherever found	\$12,000	\$0	\$12,000
		<i>Flowering Plants Total</i>			<i>\$167,300</i>	<i>\$0</i>	<i>\$167,300</i>
		<i>Domestic Subtotal</i>			<i>\$323,300</i>	<i>\$0</i>	<i>\$323,300</i>
		<b>Total</b>			<b>\$323,300</b>	<b>\$0</b>	<b>\$323,300</b>

## DOC National Oceanic and Atmospheric Administration - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Sea lion, Steller	<i>Eumetopias jubatus</i>	Western DPS	\$5,413,000	\$0	\$5,413,000
		Seal, bearded	<i>Erignathus barbatus nauticus</i>	Beringia DPS	\$1,630,000	\$0	\$1,630,000
		Seal, Guadalupe fur	<i>Arctocephalus townsendi</i>	Wherever found	\$200,000	\$0	\$200,000
		Seal, Hawaiian monk	<i>Monachus schauinslandi</i>	Wherever found	\$3,833,000	\$0	\$3,833,000
		whale, beluga	<i>Delphinapterus leucas</i>	Cook Inlet DPS	\$3,142,000	\$0	\$3,142,000
		Whale, blue	<i>Balaenoptera musculus</i>	Wherever found	\$1,333,000	\$0	\$1,333,000
		Whale, bowhead	<i>Balaena mysticetus</i>	Wherever found	\$572,000	\$0	\$572,000
		whale, false killer	<i>Pseudorca crassidens</i>	Main Hawaiian Islands Insular DPS	\$419,000	\$0	\$419,000
		Whale, finback	<i>Balaenoptera physalus</i>	Wherever found	\$1,333,000	\$0	\$1,333,000
		Whale, humpback	<i>Megaptera novaeangliae</i>	Central America DPS	\$427,000	\$0	\$427,000
		Whale, humpback	<i>Megaptera novaeangliae</i>	Mexico DPS	\$804,000	\$0	\$804,000
		Whale, humpback	<i>Megaptera novaeangliae</i>	Western North Pacific	\$572,000	\$0	\$572,000
		Whale, killer	<i>Orcinus orca</i>	Southern Resident DPS	\$1,443,000	\$0	\$1,443,000
		Whale, North Atlantic Right	<i>Eubalaena glacialis</i>	Wherever found	\$8,591,000	\$0	\$8,591,000
		Whale, North Pacific Right	<i>Eubalaena japonica</i>	Wherever found	\$572,000	\$0	\$572,000
		Whale, Sei	<i>Balaenoptera borealis</i>	Wherever found	\$1,333,000	\$0	\$1,333,000
		Whale, sperm	<i>Physeter catodon</i> (= <i>macrocephalus</i> )	Wherever found	\$1,333,000	\$0	\$1,333,000

	<i>Mammals Subtotal</i>				\$32,950,000	\$0	\$32,950,000
	Reptiles	Sea turtle, green	Chelonia mydas	Central North Pacific DPS	\$926,000	\$0	\$926,000
		Sea turtle, green	Chelonia mydas	Central South Pacific DPS	\$692,000	\$0	\$692,000
		Sea turtle, green	Chelonia mydas	Central West Pacific DPS	\$1,371,000	\$0	\$1,371,000
		Sea turtle, green	Chelonia mydas	East Pacific DPS	\$1,313,000	\$0	\$1,313,000
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$1,135,000	\$0	\$1,135,000
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$2,221,000	\$0	\$2,221,000
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$2,330,000	\$0	\$2,330,000
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$2,826,000	\$0	\$2,826,000
		Sea turtle, loggerhead	Caretta caretta	North Pacific Ocean DPS	\$1,885,000	\$0	\$1,885,000
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$2,156,000	\$0	\$2,156,000
		Sea turtle, olive ridley	Lepidochelys olivacea	Wherever found, except when listed as endangered under 50 CFR 224.101	\$558,000	\$0	\$558,000
	<i>Reptiles Subtotal</i>				\$17,413,000	\$0	\$17,413,000
	Fishes	Bocaccio	Sebastes paucispinis	Puget Sound - Georgia Basin DPS	\$728,000	\$0	\$728,000
		Eulachon	Thaleichthys pacificus	Southern DPS	\$866,000	\$0	\$866,000
		grouper, Nassau	Epinephelus striatus		\$497,000	\$0	\$497,000

		rockfish, yelloweye	Sebastes ruberrimus	Puget Sound - Georgia Basin DPS	\$700,000	\$0	\$700,000
		Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$5,955,000	\$0	\$5,955,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$3,627,000	\$0	\$3,627,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$3,627,000	\$0	\$3,627,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$5,416,000	\$0	\$5,416,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$4,549,000	\$0	\$4,549,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter-run ESU	\$3,627,000	\$0	\$3,627,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$5,006,000	\$0	\$5,006,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$6,387,000	\$0	\$6,387,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$3,599,000	\$0	\$3,599,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$3,638,000	\$0	\$3,638,000
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$5,130,000	\$0	\$5,130,000
		Salmon, chum	Oncorhynchus keta	Hood Canal summer-run ESU	\$4,549,000	\$0	\$4,549,000

		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$3,627,000	\$0	\$3,627,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$5,416,000	\$0	\$5,416,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$4,445,000	\$0	\$4,445,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$5,334,000	\$0	\$5,334,000
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU	\$4,549,000	\$0	\$4,549,000
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$3,377,000	\$0	\$3,377,000
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$497,000	\$0	\$497,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$3,627,000	\$0	\$3,627,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Central California Coast DPS	\$3,627,000	\$0	\$3,627,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$5,416,000	\$0	\$5,416,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$5,266,000	\$0	\$5,266,000

		Steelhead	Oncorhynchus (=Salmo) mykiss	Northern California DPS	\$3,952,000	\$0	\$3,952,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$4,549,000	\$0	\$4,549,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$6,387,000	\$0	\$6,387,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$3,627,000	\$0	\$3,627,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$3,627,000	\$0	\$3,627,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$3,599,000	\$0	\$3,599,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$3,638,000	\$0	\$3,638,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Carolina DPS	\$497,000	\$0	\$497,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Chesapeake Bay DPS	\$597,000	\$0	\$597,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Gulf of Maine DPS	\$597,000	\$0	\$597,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	New York Bight DPS	\$597,000	\$0	\$597,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	South Atlantic DPS	\$497,000	\$0	\$497,000
		sturgeon, green	Acipenser medirostris	Southern DPS	\$777,000	\$0	\$777,000
		Sturgeon, Gulf	Acipenser oxyrinchus (=oxyrhynchus) desotoi	Wherever found	\$497,000	\$0	\$497,000
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$880,000	\$0	\$880,000
		<i>Fishes Subtotal</i>			<i>\$137,400,000</i>	<i>\$0</i>	<i>\$137,400,000</i>



	Snails	Abalone, Black	Haliotis cracherodii	Wherever found	\$1,987,000	\$0	\$1,987,000
		Abalone, White	Haliotis sorenseni	North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	\$777,000	\$0	\$777,000
	<i>Snails Subtotal</i>				<i>\$2,764,000</i>	<i>\$0</i>	<i>\$2,764,000</i>
	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$511,000	\$0	\$511,000
		Coral, elkhorn	Acropora palmata	Wherever found	\$511,000	\$0	\$511,000
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$497,000	\$0	\$497,000
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$497,000	\$0	\$497,000
		coral, Pillar	Dendrogyra cylindrus	Wherever found	\$497,000	\$0	\$497,000
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$497,000	\$0	\$497,000
		Coral, staghorn	Acropora cervicornis	Wherever found	\$497,000	\$0	\$497,000
		No common name	Acropora globiceps	Wherever found	\$528,000	\$0	\$528,000
		No common name	Acropora jacquelinae	Wherever found	\$528,000	\$0	\$528,000
		No common name	Acropora retusa	Wherever found	\$528,000	\$0	\$528,000
		No common name	Acropora speciosa	Wherever found	\$528,000	\$0	\$528,000

		No common name	Euphyllia paradivisa	Wherever found	\$528,000	\$0	\$528,000
		No common name	Isopora crateriformis	Wherever found	\$528,000	\$0	\$528,000
		No common name	Seriatopora aculeata	Wherever found	\$214,000	\$0	\$214,000
	<i>Corals Subtotal</i>				<i>\$6,889,000</i>	<i>\$0</i>	<i>\$6,889,000</i>
	Flowering Plants	Seagrass, Johnson's	Halophila johnsonii	Wherever found	\$497,000	\$0	\$497,000
	<i>Flowering Plants Subtotal</i>						<i>\$497,000</i>
<i>Domestic Subtotal</i>					<i>\$197,913,000</i>	<i>\$0</i>	<i>\$197,913,000</i>
Grand Total					\$197,913,000	\$0	\$197,913,000

## DOD Army - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$16,061	\$0	\$16,061
		Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$155,164	\$0	\$155,164
		Bat, Indiana	Myotis sodalis	Wherever found	\$367,111	\$0	\$367,111
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$269,514	\$0	\$269,514
		Ferret, black-footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$1,000	\$0	\$1,000
		Fox, San Joaquin kit	Vulpes macrotis mutica	wherever found	\$82,506	\$0	\$82,506
		Wolf, gray	Canis lupus	MN	\$10,560	\$0	\$10,560
	<i>Mammals Subtotal</i>				<i>\$901,916</i>	<i>\$0</i>	<i>\$901,916</i>
	Birds	Crane, whooping	Grus americana	Wherever found, except where	\$125,271	\$0	\$125,271
		Falcon, Northern Aplomado	Falco femoralis septentrionalis	Wherever found, except where	\$143,000	\$0	\$143,000
		Owl, Mexican spotted	Strix occidentalis lucida	Wherever found	\$124,702	\$0	\$124,702
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern	\$2,000	\$0	\$2,000
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-	\$42,000	\$0	\$42,000
		Rail, California clapper	Rallus longirostris obsoletus	Wherever found	\$159,000	\$0	\$159,000
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$793	\$0	\$793
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$6,072	\$0	\$6,072
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$54,894	\$0	\$54,894
		Warbler (=wood), golden-cheeked	Dendroica chrysoparia	Wherever found	\$24,500	\$0	\$24,500
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$5,741,231	\$0	\$5,741,231
	<i>Birds Subtotal</i>				<i>\$6,423,463</i>	<i>\$0</i>	<i>\$6,423,463</i>

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Reptiles	Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$131,247	\$0	\$131,247
		Pinesnake, Louisiana	Pituophis ruthveni	Wherever found	\$120,000	\$0	\$120,000
		Sea turtle, green	Chelonia mydas	South Atlantic DPS	\$400	\$0	\$400
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$400	\$0	\$400
		Sea turtle, leatherback	Dermodochelys coriacea	Wherever found	\$400	\$0	\$400
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean	\$400	\$0	\$400
		Snake, black pine	Pituophis melanoleucus lodingi	Wherever found	\$28,942	\$0	\$28,942
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$107,160	\$0	\$107,160
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$251,800	\$0	\$251,800
		Tortoise, gopher	Gopherus polyphemus	West of Mobile and Tombigbee Rivers	\$104,950	\$0	\$104,950
		turtle, bog	Clemmys muhlenbergii	Wherever found, except GA, NC, SC, TN, VA	\$15,000	\$0	\$15,000
	<i>Reptiles Subtotal</i>				<i>\$760,699</i>	<i>\$0</i>	<i>\$760,699</i>
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$21,886	\$0	\$21,886
		Salamander, California tiger	Ambystoma californiense	U.S.A. (Central CA DPS)	\$12,931	\$0	\$12,931
		Salamander, frosted flatwoods	Ambystoma cingulatum	Wherever found	\$53,580	\$0	\$53,580
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$66,320	\$0	\$66,320
		Toad, Houston	Bufo houstonensis	Wherever found	\$35,581	\$0	\$35,581

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	<i>Amphibians Subtotal</i>				\$190,298	\$0	\$190,298
	Fishes	Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$25,000	\$0	\$25,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	New York Bight DPS	\$2,000	\$0	\$2,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	South Atlantic DPS	\$536	\$0	\$536
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$55,580	\$0	\$55,580
	<i>Fishes Subtotal</i>				\$83,116	\$0	\$83,116
	Clams	Bean, rayed	Villosa fabalis	Wherever found	\$11,777	\$0	\$11,777
	<i>Clams Subtotal</i>				\$11,777	\$0	\$11,777
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$88,378	\$0	\$88,378
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$24,000	\$0	\$24,000
		Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$158,000	\$0	\$158,000
		Butterfly, Mariana eight-spot	Hypolimnas octocula marianensis	Wherever found	\$25,000	\$0	\$25,000
		Butterfly, Oregon silverspot	Speyeria zerene hippolyta	Wherever found	\$38,000	\$0	\$38,000
		Butterfly, Saint Francis' satyr	Neonympha mitchellii francisci	Wherever found	\$260,785	\$0	\$260,785
		Skipper, Dakota	Hesperia dacotae	Wherever found	\$11,807	\$0	\$11,807
		Tiger beetle, Salt Creek	Cicindela nevadica lincolniana	Wherever found	\$11,617	\$0	\$11,617
	<i>Insects Subtotal</i>				\$617,587	\$0	\$617,587
	Crustaceans	Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$107,258	\$0	\$107,258
		Shrimp, Alabama cave	Palaemonias alabamae	Wherever found	\$41,000	\$0	\$41,000
	<i>Crustaceans Subtotal</i>				\$148,258	\$0	\$148,258
	Flowering Plants	Amaranth, seabeach	Amaranthus pumilus	Wherever found	\$2,000	\$0	\$2,000
		amole, purple	Chlorogalum purpureum	Wherever found	\$105,897	\$0	\$105,897
		Barbara's Buttons, Mohr's	Marshallia mohrii	Wherever found	\$300	\$0	\$300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Bear-poppy, Dwarf	Arctomecon humilis	Wherever found	\$903	\$0	\$903
		Bulrush, Northeastern	Scirpus ancistrochaetus	Wherever found	\$2,500	\$0	\$2,500
		Chaffseed, American	Schwalbea americana	Wherever found	\$131,585	\$0	\$131,585
		Clover, running buffalo	Trifolium stoloniferum		\$18,500	\$0	\$18,500
		Coneflower, smooth	Echinacea laevigata	Wherever found	\$16,716	\$0	\$16,716
		Grass, Tennessee yellow-eyed	Xyris tennesseensis	Wherever found	\$352,732	\$0	\$352,732
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$200	\$0	\$200
		Ladies'-tresses, Navasota	Spiranthes parksii	Wherever found	\$9,425	\$0	\$9,425
		Loosestrife, rough-leaved	Lysimachia asperulaefolia	Wherever found	\$157,390	\$0	\$157,390
		Lupine, Kincaid's	Lupinus sulphureus ssp. kincaidii	Wherever found	\$600	\$0	\$600
		Milkweed, Mead's	Asclepias meadii	Wherever found	\$12,460	\$0	\$12,460
		No common name	Cyrtandra nanawaleensis	Wherever found	\$44,254	\$0	\$44,254
		No common name	Geocarpon minimum	Wherever found	\$1,000	\$0	\$1,000
		Orchid, western prairie fringed	Platanthera praeclara	Wherever found	\$11,267	\$0	\$11,267
		Peppergrass, Slickspot	Lepidium papilliferum		\$85,000	\$0	\$85,000
		Piperia, Yadon's	Piperia yadonii	Wherever found	\$107,505	\$0	\$107,505
		Pogonia, small whorled	Isotria medeoloides	Wherever found	\$6,320	\$0	\$6,320
		Rhododendron, Chapman	Rhododendron chapmanii	Wherever found	\$300	\$0	\$300
		Skullcap, large-flowered	Scutellaria montana	Wherever found	\$3,000	\$0	\$3,000
		Sumac, Michaux's	Rhus michauxii	Wherever found	\$179,355	\$0	\$179,355
		<i>Flowering Plants Subtotal</i>			<i>\$1,249,209</i>	<i>\$0</i>	<i>\$1,249,209</i>
	Ferns and Allies	Quillwort, Louisiana	Isoetes louisianensis	Wherever found	\$95,784	\$0	\$95,784
		<i>Ferns and Allies Subtotal</i>			<i>\$95,784</i>	<i>\$0</i>	<i>\$95,784</i>
	Multi-Species				\$30,781,382	\$0	\$30,781,382
		<i>Multi-Species Subtotal</i>			<i>\$30,781,382</i>	<i>\$0</i>	<i>\$30,781,382</i>
	<i>Domestic Subtotal</i>				<i>\$41,263,489</i>	<i>\$0</i>	<i>\$41,263,489</i>
	Grand Total				\$41,263,489	\$0	\$41,263,489

## DOD Army Corps of Engineers - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	bat, Florida bonneted	Eumops floridanus	Wherever found	\$202,729	\$0	\$202,729
		Bat, gray	Myotis grisescens	Wherever found	\$174,891	\$0	\$174,891
		Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$3,196	\$0	\$3,196
		Bat, Indiana	Myotis sodalis	Wherever found	\$804,943	\$0	\$804,943
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$1,376,120	\$0	\$1,376,120
		Bat, Ozark big-eared	Corynorhinus (=Plecotus) townsendii ingens	Wherever found	\$10,080	\$0	\$10,080
		Bat, Virginia big-eared	Corynorhinus (=Plecotus) townsendii virginianus	Wherever found	\$5,103	\$0	\$5,103
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$3,293	\$0	\$3,293
		Bear, polar	Ursus maritimus	Wherever found	\$103,306	\$0	\$103,306
		Deer, Columbian white-tailed	Odocoileus virginianus leucurus	Columbia River DPS	\$146	\$0	\$146
		Deer, key	Odocoileus virginianus	Wherever found	\$3,989	\$0	\$3,989
		Ferret, black-footed	Mustela nigripes	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$800	\$0	\$800
		Fox, San Joaquin kit	Vulpes macrotis mutica	wherever found	\$35,022	\$0	\$35,022
		Jaguarundi, Gulf Coast	Herpailurus (=Felis) yagouaroundi cacomitli	Wherever found	\$210	\$0	\$210
		Kangaroo rat, Fresno	Dipodomys nitratoides exilis	Wherever found	\$2,934	\$0	\$2,934

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Kangaroo rat, San Bernardino Merriam's	Dipodomys merriami parvus	Wherever found	\$67,138	\$0	\$67,138
		Kangaroo rat, Stephens'	Dipodomys stephensi (incl. D. cascus)	Wherever found	\$2,684	\$0	\$2,684
		Kangaroo rat, Tipton	Dipodomys nitratooides nitratooides	Wherever found	\$800	\$0	\$800
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$13,030	\$0	\$13,030
		Manatee, West Indian	Trichechus manatus	Wherever found	\$504,886	\$0	\$504,886
		Mouse, Alabama beach	Peromyscus polionotus ammobates	Wherever found	\$1,594	\$0	\$1,594
		Mouse, Choctawhatchee beach	Peromyscus polionotus allophrys	Wherever found	\$2,021	\$0	\$2,021
		Mouse, Key Largo cotton	Peromyscus gossypinus allapaticola	Wherever found	\$1,561	\$0	\$1,561
		Mouse, New Mexico meadow jumping	Zapus hudsonius luteus	Wherever found	\$12,107	\$0	\$12,107
		Mouse, Perdido Key beach	Peromyscus polionotus trissyllepsis	Wherever found	\$1,594	\$0	\$1,594
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	wherever found	\$8,438	\$0	\$8,438
		Mouse, salt marsh harvest	Reithrodontomys raviventris	wherever found	\$93,200	\$0	\$93,200
		Mouse, southeastern beach	Peromyscus polionotus niveiventris	wherever found	\$100	\$0	\$100
		Ocelot	Leopardus (=Felis) pardalis	wherever found	\$210	\$0	\$210
		Otter, Northern Sea	Enhydra lutris kenyoni	Southwest Alaska DPS	\$83,501	\$0	\$83,501
		Otter, southern sea	Enhydra lutris nereis	Wherever found	\$15,754	\$0	\$15,754
		Panther, Florida	Puma (=Felis) concolor coryi	Wherever found	\$85,307	\$0	\$85,307
		Rabbit, Lower Keys marsh	Sylvilagus palustris hefneri	Wherever found	\$1,351	\$0	\$1,351
		Rabbit, riparian brush	Sylvilagus bachmani riparius	Wherever found	\$800	\$0	\$800



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Rice rat, Silver	<i>Oryzomys palustris natator</i>	lower FL Keys	\$4,361	\$0	\$4,361
		Sea lion, Steller	<i>Eumetopias jubatus</i>	Western DPS	\$117,234	\$0	\$117,234
		Seal, Hawaiian monk	<i>Monachus schauinslandi</i>	Wherever found	\$5,025	\$0	\$5,025
		Seal, Ringed	<i>Phoca (=Pusa) hispida hispida</i>	Wherever found	\$11,998	\$0	\$11,998
		Shrew, Buena Vista Lake ornate	<i>Sorex ornatus relictus</i>	Wherever found	\$100	\$0	\$100
		Squirrel, Carolina northern flying	<i>Glaucomys sabrinus coloratus</i>	Wherever found	\$1,424	\$0	\$1,424
		whale, beluga	<i>Delphinapterus leucas</i>	Cook Inlet DPS	\$34,676	\$0	\$34,676
		Whale, blue	<i>Balaenoptera musculus</i>	Wherever found	\$8,074	\$0	\$8,074
		Whale, bowhead	<i>Balaena mysticetus</i>	Wherever found	\$11,999	\$0	\$11,999
		whale, false killer	<i>Pseudorca crassidens</i>	Main Hawaiian Islands Insular DPS	\$2,882	\$0	\$2,882
		Whale, finback	<i>Balaenoptera physalus</i>	Wherever found	\$77,496	\$0	\$77,496
		Whale, gray	<i>Eschrichtius robustus</i>	Western North Pacific DPS	\$3,000	\$0	\$3,000
		Whale, humpback	<i>Megaptera novaeangliae</i>	Central America DPS	\$3,773	\$0	\$3,773
		Whale, humpback	<i>Megaptera novaeangliae</i>	Mexico DPS	\$110,407	\$0	\$110,407
		Whale, humpback	<i>Megaptera novaeangliae</i>	Western North Pacific	\$40,750	\$0	\$40,750
		Whale, killer	<i>Orcinus orca</i>	Southern Resident DPS	\$39,506	\$0	\$39,506
		Whale, North Atlantic Right	<i>Eubalaena glacialis</i>	Wherever found	\$224,284	\$0	\$224,284
		Whale, North Pacific Right	<i>Eubalaena japonica</i>	Wherever found	\$22,841	\$0	\$22,841
		Whale, Sei	<i>Balaenoptera borealis</i>	Wherever found	\$6,194	\$0	\$6,194
		Whale, Southern Right	<i>Eubalaena australis</i>	Wherever found	\$476	\$0	\$476
		Whale, sperm	<i>Physeter catodon</i> (=macrocephalus)	Wherever found	\$11,644	\$0	\$11,644

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$38,029	\$0	\$38,029
		Woodrat, Key Largo	Neotoma floridana smalli	Wherever found	\$1,048	\$0	\$1,048
		Woodrat, riparian (=San Joaquin Valley)	Neotoma fuscipes riparia	Wherever found	\$800	\$0	\$800
	<i>Mammals Subtotal</i>				<i>\$4,400,859</i>	<i>\$0</i>	<i>\$4,400,859</i>
	Birds	Albatross, short-tailed	Phoebastria (=Diomedea) albatrus	Wherever found	\$1,200	\$0	\$1,200
		Blackbird, yellow-shouldered	Agelaius xanthomus	Wherever found	\$250	\$0	\$250
		Caracara, Audubon's crested	Polyborus plancus audubonii	FL pop.	\$8,907,913	\$0	\$8,907,913
		Condor, California	Gymnogyps californianus	U.S.A. only, except where listed as an experimental population	\$352	\$0	\$352
		Crane, Mississippi sandhill	Grus canadensis pulla	Wherever found	\$3,926	\$0	\$3,926

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Crane, whooping	<i>Grus americana</i>	U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	\$4,083	\$0	\$4,083
		Crane, whooping	<i>Grus americana</i>	U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	\$17,000	\$0	\$17,000
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$71,160	\$0	\$71,160
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$404,050	\$0	\$404,050
		Duck, Hawaiian (=koloa)	<i>Anas wyvilliana</i>	Wherever found	\$3,196	\$0	\$3,196
		Eider, spectacled	<i>Somateria fischeri</i>	Wherever found	\$86,450	\$0	\$86,450
		Eider, Steller's	<i>Polysticta stelleri</i>	AK breeding pop.	\$103,230	\$0	\$103,230
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$552,588	\$0	\$552,588
		gallinule, Hawaiian common	<i>Gallinula galeata sandvicensis</i>	Wherever found	\$3,196	\$0	\$3,196
		Gnatcatcher, coastal California	<i>Polioptila californica californica</i>	Wherever found	\$147,014	\$0	\$147,014
		Hawk, Puerto Rican broad-winged	<i>Buteo platypterus brunnescens</i>	Wherever found	\$250	\$0	\$250

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Hawk, Puerto Rican sharp-shinned	Accipiter striatus venator	Wherever found	\$250	\$0	\$250
		Horned lark, streaked	Eremophila alpestris strigata	Wherever found	\$210,709	\$0	\$210,709
		Kite, Everglade snail	Rostrhamus sociabilis plumbeus	Wherever found	\$626,626	\$0	\$626,626
		Knot, red	Calidris canutus rufa	Wherever found	\$327,857	\$0	\$327,857
		Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR, WA)	\$47,323	\$0	\$47,323
		Nightjar, Puerto Rican	Caprimulgus noctitherus	Wherever found	\$250	\$0	\$250
		Owl, Mexican spotted	Strix occidentalis lucida	Wherever found	\$2,707	\$0	\$2,707
		Owl, northern spotted	Strix occidentalis caurina	Wherever found	\$14,111	\$0	\$14,111
		petrel, Hawaiian	Pterodroma sandwichensis	Wherever found	\$3,196	\$0	\$3,196
		Pigeon, Puerto Rican plain	Columba inornata wetmorei	Wherever found	\$1,707	\$0	\$1,707
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] Wherever found, except those areas where listed as endangered.	\$4,955,622	\$6,530	\$4,962,152
		Plover, piping	Charadrius melodus	[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont)	\$38,351	\$0	\$38,351

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, western snowy	<i>Charadrius nivosus nivosus</i>	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$346,134	\$0	\$346,134
		Rail, California clapper	<i>Rallus longirostris obsoletus</i>	Wherever found	\$116,132	\$0	\$116,132
		Rail, light-footed clapper	<i>Rallus longirostris levipes</i>	Wherever found	\$4,600	\$0	\$4,600
		Rail, Yuma Ridgways	<i>Rallus obsoletus</i> [= <i>longirostris</i> ] <i>yumanensis</i>	Wherever found	\$9,272	\$0	\$9,272
		sage-grouse, Gunnison	<i>Centrocercus minimus</i>	Wherever found	\$2,400	\$0	\$2,400
		scrub-jay, Florida	<i>Aphelocoma coerulescens</i>	Wherever found	\$25,922	\$0	\$25,922
		Shearwater, Newell's Townsend's	<i>Puffinus auricularis newelli</i>	Wherever found	\$3,196	\$0	\$3,196
		Sparrow, Cape Sable seaside	<i>Ammodramus maritimus mirabilis</i>	Wherever found	\$267,425	\$0	\$267,425
		Sparrow, Florida grasshopper	<i>Ammodramus savannarum floridanus</i>	Wherever found	\$5,052	\$0	\$5,052
		Stilt, Hawaiian	<i>Himantopus mexicanus knudseni</i>	Wherever found	\$3,196	\$0	\$3,196
		Stork, wood	<i>Mycteria americana</i>	AL, FL, GA, MS, NC, SC	\$448,075	\$0	\$448,075
		Storm-petrel, band-rumped	<i>Oceanodroma castro</i>	USA (HI)	\$3,196	\$0	\$3,196
		Tern, California least	<i>Sterna antillarum browni</i>	Wherever found	\$62,064	\$0	\$62,064
		Tern, least	<i>Sterna antillarum</i>	interior pop.	\$5,253,914	\$6,530	\$5,260,444
		Tern, roseate	<i>Sterna dougallii dougallii</i>	Northeast U.S. nesting population	\$54,032	\$0	\$54,032
		Vireo, least Bell's	<i>Vireo bellii pusillus</i>	Wherever found	\$2,482,720	\$0	\$2,482,720
		Warbler (=wood), golden-cheeked	<i>Dendroica chrysoparia</i>	Wherever found	\$32,520	\$0	\$32,520
		Warbler, elfin-woods	<i>Setophaga angelae</i>	Wherever found	\$250	\$0	\$250

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Warbler, Kirtland's	Setophaga kirtlandii (= Dendroica kirtlandii)	Wherever found	\$14,246	\$0	\$14,246
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$95,587	\$0	\$95,587
	<i>Birds Subtotal</i>				<i>\$25,764,500</i>	<i>\$13,060</i>	<i>\$25,777,560</i>
	Reptiles	Alligator, American	Alligator mississippiensis	Wherever found	\$537	\$0	\$537
		Boa, Puerto Rican	Epicrates inornatus	Wherever found	\$5,413	\$0	\$5,413
		Crocodile, American	Crocodylus acutus	U.S.A. (FL)	\$33,325	\$0	\$33,325
		gartersnake, northern Mexican	Thamnophis eques megalops	Wherever found	\$4,225	\$0	\$4,225
		Lizard, blunt-nosed leopard	Gambelia silus	Wherever found	\$3,421	\$0	\$3,421
		Lizard, Coachella Valley fringe-toed	Uma inornata	Wherever found	\$2,684	\$0	\$2,684
		Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$44,422	\$0	\$44,422
		Pinesnake, Louisiana	Pituophis ruthveni	Wherever found	\$467	\$0	\$467
		Sea turtle, green	Chelonia mydas	Central North Pacific DPS	\$8,222	\$0	\$8,222
		Sea turtle, green	Chelonia mydas	Central South Pacific DPS	\$7,320	\$0	\$7,320
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$270,417	\$0	\$270,417
		Sea turtle, green	Chelonia mydas	South Atlantic DPS	\$996,930	\$0	\$996,930
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$656,467	\$0	\$656,467
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$1,620,555	\$0	\$1,620,555
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$996,391	\$0	\$996,391
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$708,250	\$0	\$708,250

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, olive ridley	Lepidochelys olivacea	Wherever found, except when listed as endangered under 50 CFR 224.101	\$3,664	\$0	\$3,664
		Skink, bluetail mole	Eumeces egregius lividus	Wherever found	\$8,106	\$0	\$8,106
		Skink, sand	Neoseps reynoldsi	Wherever found	\$11,476	\$0	\$11,476
		Snake, Atlantic salt marsh	Nerodia clarkii taeniata	Wherever found	\$569	\$0	\$569
		Snake, black pine	Pituophis melanoleucus lodingi	Wherever found	\$5,614	\$0	\$5,614
		Snake, copperbelly water	Nerodia erythrogaster neglecta	Indiana north of 40 degrees north latitude, Michigan, Ohio	\$1,066	\$0	\$1,066
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$203,392	\$0	\$203,392
		Snake, giant garter	Thamnophis gigas	Wherever found	\$239,948	\$0	\$239,948
		Snake, San Francisco garter	Thamnophis sirtalis tetrataenia	Wherever found	\$17,600	\$0	\$17,600
		Tortoise, desert	Gopherus agassizii	AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	\$100	\$0	\$100
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$8,486	\$0	\$8,486
		Tortoise, gopher	Gopherus polyphemus	West of Mobile and Tombigbee Rivers	\$40,379	\$0	\$40,379

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Turtle, Alabama red-bellied	<i>Pseudemys alabamensis</i>	Wherever found	\$7,958	\$0	\$7,958
		turtle, bog	<i>Clemmys muhlenbergii</i>	Wherever found, except GA, NC, SC, TN, VA	\$103,883	\$0	\$103,883
		Turtle, flattened musk	<i>Sternotherus depressus</i>	Black Warrior R. system upstream from Bankhead Dam	\$1,137	\$0	\$1,137
		Turtle, Plymouth Redbelly	<i>Pseudemys rubriventris bangsi</i>	Wherever found	\$100	\$0	\$100
		Turtle, ringed map	<i>Gratemys oculifera</i>	Wherever found	\$5,969	\$0	\$5,969
		Turtle, yellow-blotched map	<i>Gratemys flavimaculata</i>	Wherever found	\$2,813	\$0	\$2,813
		Whipsnake (=striped racer), Alameda	<i>Masticophis lateralis euryxanthus</i>	Wherever found	\$25,200	\$0	\$25,200
	<i>Reptiles Subtotal</i>				<i>\$6,046,506</i>	<i>\$0</i>	<i>\$6,046,506</i>
	Amphibians	Frog, California red-legged	<i>Rana draytonii</i>	Wherever found	\$226,176	\$0	\$226,176
		Frog, Chiricahua leopard	<i>Rana chiricahuensis</i>	Wherever found	\$1,907	\$0	\$1,907
		Frog, dusky gopher	<i>Rana sevosia</i>	Wherever found	\$9,593	\$0	\$9,593
		Frog, mountain yellow-legged	<i>Rana muscosa</i>	Northern California DPS	\$800	\$0	\$800
		Frog, Oregon spotted	<i>Rana pretiosa</i>	Wherever found	\$219	\$0	\$219
		Frog, Sierra Nevada Yellow-legged	<i>Rana sierrae</i>	Wherever found	\$4,800	\$0	\$4,800
		Hellbender, Ozark	<i>Cryptobranchus alleganiensis bishopi</i>	Wherever found	\$4,000	\$0	\$4,000
		Salamander, California tiger	<i>Ambystoma californiense</i>	U.S.A. (CA - Sonoma County)	\$30,400	\$0	\$30,400
		Salamander, California tiger	<i>Ambystoma californiense</i>	U.S.A. (Central CA DPS)	\$73,963	\$0	\$73,963
		Salamander, Cheat Mountain	<i>Plethodon nettingi</i>	Wherever found	\$103	\$0	\$103
		Salamander, Red Hills	<i>Phaeognathus hubrichti</i>	Wherever found	\$1,248	\$0	\$1,248



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		salamander, Reticulated flatwoods	Ambystoma bishopi	Wherever found	\$3,715	\$0	\$3,715
		Salamander, San Marcos	Eurycea nana	Wherever found	\$840	\$0	\$840
		Salamander, Santa Cruz long-toed	Ambystoma macrodactylum croceum	Wherever found	\$1,600	\$0	\$1,600
		Salamander, Shenandoah	Plethodon shenandoah	Wherever found	\$240	\$0	\$240
		Salamander, Texas blind	Eurycea [=Typhlomolge] rathbuni	Wherever found	\$840	\$0	\$840
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$20,740	\$0	\$20,740
		Toad, Houston	Bufo houstonensis	Wherever found	\$840	\$0	\$840
		toad, Yosemite	Anaxyrus canorus	Wherever found	\$2,400	\$0	\$2,400
		Waterdog, black warrior (=Sipsey Fork)	Necturus alabamensis	Wherever found	\$1,137	\$0	\$1,137
	<i>Amphibians Subtotal</i>				<i>\$385,561</i>	<i>\$0</i>	<i>\$385,561</i>
	Fishes	Bocaccio	Sebastes paucispinis	Puget Sound - Georgia Basin DPS	\$28,223	\$0	\$28,223
		Bonytail	Gila elegans	Wherever found	\$17,784	\$0	\$17,784
		Cavefish, Ozark	Amblyopsis rosae	Wherever found	\$12,960	\$0	\$12,960
		Chub, humpback	Gila cypha	Wherever found	\$14,300	\$0	\$14,300
		Chub, slender	Erimystax cahni	Wherever found	\$370	\$0	\$370
		Chub, spotfin	Erimonax monachus	U.S.A. (TN-specified portions of the Tellico River; see 17.84(m)(1)(i))	\$1,056	\$0	\$1,056
		Chub, spotfin	Erimonax monachus	Wherever found, except where listed as an experimental population	\$2,666	\$0	\$2,666

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Dace, blackside	<i>Phoxinus cumberlandensis</i>	Wherever found	\$3,044	\$0	\$3,044
		dace, Laurel	<i>Chrosomus saylori</i>	Wherever found	\$250	\$0	\$250
		Darter, amber	<i>Percina antesella</i>	Wherever found	\$330	\$0	\$330
		darter, bluemask	<i>Etheostoma akatulo</i>	Wherever found	\$626	\$0	\$626
		Darter, Cherokee	<i>Etheostoma scotti</i>	Wherever found	\$330	\$0	\$330
		Darter, Cumberland	<i>Etheostoma susanae</i>	Wherever found	\$748	\$0	\$748
		Darter, diamond	<i>Crystallaria cincotta</i>	Wherever found	\$1,030	\$0	\$1,030
		Darter, duskytail	<i>Etheostoma percnum</i>	Wherever found	\$173,894	\$0	\$173,894
		Darter, Etowah	<i>Etheostoma etowahae</i>	Wherever found	\$330	\$0	\$330
		Darter, fountain	<i>Etheostoma fonticola</i>	Wherever found	\$840	\$0	\$840
		Darter, goldline	<i>Percina aurolineata</i>	Wherever found	\$772	\$0	\$772
		Darter, Kentucky arrow	<i>Etheostoma spilotum</i>	Wherever found	\$3,182	\$0	\$3,182
		Darter, Maryland	<i>Etheostoma sellare</i>	Wherever found	\$240	\$0	\$240
		Darter, Niangua	<i>Etheostoma nianguae</i>	Wherever found	\$3,565	\$0	\$3,565
		Darter, Pearl	<i>Percina aurora</i>	Wherever found	\$678	\$0	\$678
		Darter, rush	<i>Etheostoma phytophilum</i>	Wherever found	\$418	\$0	\$418
		Darter, snail	<i>Percina tanasi</i>	Wherever found	\$1,127	\$0	\$1,127
		Darter, watercress	<i>Etheostoma nuchale</i>	Wherever found	\$120	\$0	\$120
		Eulachon	<i>Thaleichthys pacificus</i>	Southern DPS	\$2,561,561	\$0	\$2,561,561
		Goby, tidewater	<i>Eucyclogobius newberryi</i>	Wherever found	\$40,922	\$0	\$40,922
		grouper, Nassau	<i>Epinephelus striatus</i>		\$13,951	\$0	\$13,951
		Logperch, Roanoke	<i>Percina rex</i>	Wherever found	\$2,140	\$0	\$2,140
		Madtom, Neosho	<i>Noturus placidus</i>	Wherever found	\$1,440	\$0	\$1,440
		Madtom, pygmy	<i>Noturus stanauli</i>	Wherever found	\$251	\$0	\$251
		Madtom, Scioto	<i>Noturus trautmani</i>	Wherever found	\$1,030	\$0	\$1,030
		Madtom, yellowfin	<i>Noturus flavipinnis</i>	Wherever found, except where listed as an experimental population	\$210	\$0	\$210
		Minnow, loach	<i>Tiaroga cobitis</i>	Wherever found	\$1,907	\$0	\$1,907

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Minnow, Rio Grande Silvery	Hybognathus amarus	Wherever found, except where listed as an experimental population	\$641,846	\$0	\$641,846
		Pikeminnow (=squawfish), Colorado	Ptychocheilus lucius	Wherever found, except where listed as an experimental population	\$15,100	\$0	\$15,100
		Pupfish, desert	Cyprinodon macularius	Wherever found	\$2,318	\$0	\$2,318
		rockfish, canary	Sebastes pinniger	Puget Sound - Georgia Basin DPS	\$4,630	\$0	\$4,630
		rockfish, yelloweye	Sebastes ruberrimus	Puget Sound - Georgia Basin DPS	\$7,075	\$0	\$7,075
		Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$135,040	\$0	\$135,040
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$38,684	\$0	\$38,684
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$1,144,131	\$0	\$1,144,131
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$3,395,368	\$0	\$3,395,368
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$4,751,252	\$387,341	\$5,138,593
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter-run ESU	\$1,974,422	\$0	\$1,974,422

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$11,499,900	\$0	\$11,499,900
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$11,538,365	\$0	\$11,538,365
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$7,370,997	\$0	\$7,370,997
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$4,023,581	\$0	\$4,023,581
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$3,404,415	\$0	\$3,404,415
		Salmon, chum	Oncorhynchus keta	Hood Canal summer-run ESU	\$30,564	\$0	\$30,564
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$1,266,465	\$0	\$1,266,465
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$3,392,435	\$0	\$3,392,435
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$2,585,653	\$0	\$2,585,653
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$11,714,521	\$0	\$11,714,521
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$177,654	\$0	\$177,654

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Shark, Scalloped Hammerhead	Sphyrna lewini	Central and Southwest Atlantic DPS	\$1,407	\$0	\$1,407
		Shiner, Arkansas River	Notropis girardi	Arkansas River Basin (AR, KS, NM, OK, TX)	\$10,080	\$0	\$10,080
		Shiner, Cahaba	Notropis cahabae	Wherever found	\$805	\$0	\$805
		Shiner, Cape Fear	Notropis mekistocholas	Wherever found	\$3,640	\$0	\$3,640
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$28,932	\$0	\$28,932
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$195,471	\$0	\$195,471
		Spikedace	Meda fulgida	Wherever found	\$1,907	\$0	\$1,907
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$1,778,163	\$0	\$1,778,163
		Steelhead	Oncorhynchus (=Salmo) mykiss	Central California Coast DPS	\$2,040,154	\$0	\$2,040,154
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$3,391,502	\$0	\$3,391,502
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$7,837,651	\$0	\$7,837,651
		Steelhead	Oncorhynchus (=Salmo) mykiss	Northern California DPS	\$25,000	\$0	\$25,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$3,986,462	\$387,341	\$4,373,803
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$11,600,561	\$0	\$11,600,561
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$23,600	\$0	\$23,600

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$49,370	\$0	\$49,370
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$7,402,870	\$0	\$7,402,870
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$4,023,216	\$0	\$4,023,216
		Sturgeon, Alabama	Scaphirhynchus suttkusi	Wherever found	\$6,387	\$0	\$6,387
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Gulf of Maine DPS	\$424,230	\$0	\$424,230
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	New York Bight DPS	\$358,039	\$0	\$358,039
		sturgeon, green	Acipenser medirostris	Southern DPS	\$3,642,922	\$0	\$3,642,922
		Sturgeon, Gulf	Acipenser oxyrinchus (=oxyrhynchus) desotoi	Wherever found	\$3,911,206	\$0	\$3,911,206
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$19,484,326	\$165,990	\$19,650,316
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$375,615	\$0	\$375,615
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$39,853	\$0	\$39,853
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$18,584	\$0	\$18,584
		Sucker, Santa Ana	Catostomus santaanae	3 CA river basins	\$44,504	\$0	\$44,504
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$7,411,051	\$387,341	\$7,798,392
		Trout, Gila	Oncorhynchus gilae	Wherever found	\$1,907	\$0	\$1,907
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$4,038	\$0	\$4,038
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$15,204	\$0	\$15,204
	<i>Fishes Subtotal</i>				<i>\$150,145,368</i>	<i>\$1,328,013</i>	<i>\$151,473,381</i>

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Clams	Acornshell, southern	<i>Epioblasma othcaloogensis</i>	Wherever found	\$542	\$0	\$542
		Bankclimber, purple (mussel)	<i>Elliptoideus sloatianus</i>	Wherever found	\$75,240	\$0	\$75,240
		Bean, Choctaw	<i>Villosa choctawensis</i>	Wherever found	\$1,632	\$0	\$1,632
		Bean, Cumberland (pearlymussel)	<i>Villosa trabalis</i>	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$2,409	\$0	\$2,409
		Bean, Cumberland (pearlymussel)	<i>Villosa trabalis</i>	Wherever found; Except where listed as Experimental Populations	\$736	\$0	\$736
		Bean, purple	<i>Villosa perpurpurea</i>	Wherever found	\$3,816	\$0	\$3,816
		Bean, rayed	<i>Villosa fabalis</i>	Wherever found	\$18,111	\$0	\$18,111
		Blossom, green (pearlymussel)	<i>Epioblasma torulosa gubernaculum</i>	Wherever found	\$100	\$0	\$100
		Blossom, tubercled (pearlymussel)	<i>Epioblasma torulosa torulosa</i>	Wherever found; Except where listed as Experimental Populations	\$1,631	\$0	\$1,631
		Blossom, turgid (pearlymussel)	<i>Epioblasma turgidula</i>	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$250	\$0	\$250

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Catspaw, white (pearlymussel)	Epioblasma obliquata perobliqua	Wherever found	\$846	\$0	\$846
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$30,483	\$0	\$30,483
		Clubshell, black	Pleurobema curtum	Wherever found	\$100	\$0	\$100
		Clubshell, ovate	Pleurobema perovatum	Wherever found	\$2,711	\$0	\$2,711
		Clubshell, southern	Pleurobema decisum	Wherever found	\$8,165	\$0	\$8,165
		Combshell, Cumberlandian	Epioblasma brevidens	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$1,002	\$0	\$1,002
		Combshell, Cumberlandian	Epioblasma brevidens	Wherever found; Except where listed as Experimental Populations	\$710	\$0	\$710
		Combshell, southern	Epioblasma penita	Wherever found	\$100	\$0	\$100
		Combshell, upland	Epioblasma metastriata	Wherever found	\$1,896	\$0	\$1,896
		Ebonys shell, round	Fusconaia rotulata	Wherever found	\$648	\$0	\$648
		Elktoe, Appalachian	Alasmidonta raveneliana	Wherever found	\$6,814	\$0	\$6,814
		Elktoe, Cumberland	Alasmidonta atropurpurea	Wherever found	\$626	\$0	\$626



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Fanshell	Cyprogenia stegaria	U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$3,379	\$0	\$3,379
		Fanshell	Cyprogenia stegaria	Wherever found	\$62,074	\$0	\$62,074
		fatmucket, Arkansas	Lampsilis powellii	Wherever found	\$3,312	\$0	\$3,312
		Heelsplitter, Carolina	Lasmigona decorata	Wherever found	\$6,280	\$0	\$6,280
		heelsplitter, Inflated	Potamilus inflatus	Wherever found	\$5,934	\$0	\$5,934
		Higgins eye (pearlymussel)	Lampsilis higginsii	Wherever found	\$624,400	\$0	\$624,400
		Kidneyshell, fluted	Ptychobranthus subtentus	Wherever found	\$1,746	\$0	\$1,746
		Kidneyshell, southern	Ptychobranthus jonesi	Wherever found	\$1,290	\$0	\$1,290
		Kidneyshell, triangular	Ptychobranthus greenii	Wherever found	\$1,430	\$0	\$1,430
		Lampmussel, Alabama	Lampsilis virescens	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$2,380	\$0	\$2,380
		Lance, yellow	Elliptio lanceolata	Wherever found	\$2,018	\$0	\$2,018
		Lilliput, pale (pearlymussel)	Toxolasma cylindrellus	Wherever found	\$100	\$0	\$100
		Mapleleaf, winged	Quadrula fragosa	Wherever found, except where listed as an experimental population	\$3,088	\$0	\$3,088
		Moccasinshell, Alabama	Medionidus acutissimus	Wherever found	\$1,056	\$0	\$1,056
		Moccasinshell, Coosa	Medionidus parvulus	Wherever found	\$120	\$0	\$120
		Moccasinshell, Suwannee	Medionidus walkeri	Wherever found	\$1,536	\$0	\$1,536

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Monkeyface, Appalachian (pearlymussel)	Quadrula sparsa	USA (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$425	\$0	\$425
		Monkeyface, Appalachian (pearlymussel)	Quadrula sparsa	Wherever found	\$100	\$0	\$100
		Monkeyface, Cumberland (pearlymussel)	Quadrula intermedia	Wherever found; Except where listed as Experimental Populations	\$510	\$0	\$510
		Mucket, Neosho	Lampsilis rafinesqueana	Wherever found	\$2,042	\$0	\$2,042
		Mucket, orangenacre	Lampsilis perovalis	Wherever found	\$1,576	\$0	\$1,576
		Mucket, pink (pearlymussel)	Lampsilis abrupta	Wherever found	\$58,297	\$0	\$58,297
		Mussel, oyster	Epioblasma capsaeformis	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$1,253	\$0	\$1,253
		Mussel, oyster	Epioblasma capsaeformis	Wherever found; Except where listed as Experimental Populations	\$850	\$0	\$850
		Mussel, scaleshell	Leptodea leptodon	Wherever found	\$12,083	\$0	\$12,083
		Mussel, sheepnose	Plethobasus cyphus	Wherever found	\$77,204	\$0	\$77,204
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$26,956	\$0	\$26,956
		Pearlshell, Louisiana	Margaritifera hembeli	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pearlymussel, birdwing	Lemiox rimosus	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert, and Lauderdale Counties, AL	\$350	\$0	\$350
		Pearlymussel, birdwing	Lemiox rimosus	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Pearlymussel, cracking	Hemistena lata	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$251	\$0	\$251
		Pearlymussel, cracking	Hemistena lata	Wherever found; Except where listed as Experimental Populations	\$710	\$0	\$710
		Pearlymussel, Curtis	Epioblasma florentina curtisii	Wherever found	\$666	\$0	\$666

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pearlymussel, dromedary	<i>Dromus dromas</i>	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$3,382	\$0	\$3,382
		Pearlymussel, dromedary	<i>Dromus dromas</i>	Wherever found; Except where listed as Experimental Populations	\$470	\$0	\$470
		Pearlymussel, littlewing	<i>Pegias fabula</i>	Wherever found	\$1,657	\$0	\$1,657
		Pearlymussel, slabside	<i>Pleuroaia dolabelloides</i>	Wherever found	\$1,627	\$0	\$1,627
		Pigtoe, Cumberland	<i>Pleurobema gibberum</i>	Wherever found	\$726	\$0	\$726
		Pigtoe, dark	<i>Pleurobema furvum</i>	Wherever found	\$573	\$0	\$573
		Pigtoe, finerayed	<i>Fusconaia cuneolus</i>	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$3,508	\$0	\$3,508
		Pigtoe, finerayed	<i>Fusconaia cuneolus</i>	Wherever found; Except where listed as Experimental Populations	\$650	\$0	\$650
		Pigtoe, fuzzy	<i>Pleurobema strodeanum</i>	Wherever found	\$1,632	\$0	\$1,632
		Pigtoe, heavy	<i>Pleurobema taitianum</i>	Wherever found	\$764	\$0	\$764
		Pigtoe, narrow	<i>Fusconaia escambia</i>	Wherever found	\$989	\$0	\$989
		Pigtoe, oval	<i>Pleurobema pyriforme</i>	Wherever found	\$76,152	\$0	\$76,152

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pigtoe, rough	Pleurobema plenum	U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$3,006	\$0	\$3,006
		Pigtoe, rough	Pleurobema plenum	Wherever found	\$18,818	\$0	\$18,818
		Pigtoe, shiny	Fusconaia cor	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$2,881	\$0	\$2,881
		Pigtoe, shiny	Fusconaia cor	Wherever found; Except where listed as Experimental Populations	\$750	\$0	\$750
		Pigtoe, southern	Pleurobema georgianum	Wherever found	\$829	\$0	\$829
		Pigtoe, tapered	Fusconaia burkei	Wherever found	\$643	\$0	\$643
		Pimpleback, orangefoot	Plethobasus cooperianus	Wherever found	\$23,170	\$0	\$23,170
		Pocketbook, fat	Potamilus capax	Wherever found	\$144,404	\$0	\$144,404
		Pocketbook, finelined	Lampsilis altilis	Wherever found	\$2,263	\$0	\$2,263
		Pocketbook, Ouachita rock	Arkansia wheeleri	Wherever found	\$5,488	\$0	\$5,488
		Purple Cat's paw (=Purple Cat's paw pearl mussel)	Epioblasma obliquata obliquata	AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$251	\$0	\$251

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Purple Cat's paw (=Purple Cat's paw pearlymussel)	Epioblasma obliquata obliquata	Wherever found; Except where listed as Experimental Populations	\$12,654	\$0	\$12,654
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$33,347	\$0	\$33,347
		Rabbitsfoot, rough	Quadrula cylindrica strigillata	Wherever found	\$3,106	\$0	\$3,106
		Riffleshell, northern	Epioblasma torulosa rangiana	Wherever found	\$33,865	\$0	\$33,865
		Riffleshell, tan	Epioblasma florentina walkeri	Wherever found	\$476	\$0	\$476
		Ring pink (mussel)	Obovaria retusa	Wherever found	\$19,980	\$0	\$19,980
		Sandshell, Southern	Hamiota australis	Wherever found	\$1,632	\$0	\$1,632
		Slabshell, Chipola	Elliptio chipolaensis	Wherever found	\$75,240	\$0	\$75,240
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$50,949	\$0	\$50,949
		Spinymussel, James	Pleurobema collina	Wherever found	\$310	\$0	\$310
		Spinymussel, Tar River	Elliptio steinstansana	Wherever found	\$6,125	\$0	\$6,125
		Threeridge, fat (mussel)	Amblema neislerii	Wherever found	\$75,240	\$0	\$75,240
		Wartyback, white (pearlymussel)	Plethobasus cicatricosus	Wherever found	\$351	\$0	\$351
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$29,264	\$0	\$29,264
	<i>Clams Subtotal</i>				<i>\$1,699,656</i>	<i>\$0</i>	<i>\$1,699,656</i>
	Snails						
		Abalone, Black	Haliotis cracherodii	Wherever found	\$68,000	\$0	\$68,000
		Campeloma, slender	Campeloma decampi	Wherever found	\$997	\$0	\$997
		Cavesnail, Tumbling Creek	Antrobia culveri	Wherever found	\$9,000	\$0	\$9,000
		Hornsnail, rough	Pleurocera foremani	Wherever found	\$1,196	\$0	\$1,196
		Lioplax, cylindrical (snail)	Lioplax cyclostomaformis	Wherever found	\$622	\$0	\$622
		Marstonia, royal (snail)	Pyrgulopsis ogmorhappe	Wherever found	\$125	\$0	\$125
		Pebblesnail, flat	Lepyrium showalteri	Wherever found	\$149	\$0	\$149

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Riversnail, Anthony's	Athearnia anthonyi	U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$4,254	\$0	\$4,254
		Rocksail, painted	Leptoxis taeniata	Wherever found	\$396	\$0	\$396
		Rocksail, round	Leptoxis ampla	Wherever found	\$597	\$0	\$597
		Snail, Bliss Rapids	Taylorconcha serpenticola	Wherever found	\$2,550	\$0	\$2,550
		Snail, Morro shoulderband (=Banded dune)	Helminthoglypta walkeriana	Wherever found	\$2,318	\$0	\$2,318
		Snail, Snake River physa	Physa natricina	Wherever found	\$8,261	\$0	\$8,261
		Snail, Stock Island tree	Orthalicus reses (not incl.	Wherever found	\$1,484	\$0	\$1,484
		Snail, tulotoma	Tulotoma magnifica	Wherever found	\$545	\$0	\$545
	<i>Snails Subtotal</i>				<i>\$100,494</i>	<i>\$0</i>	<i>\$100,494</i>
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$103,585	\$0	\$103,585
		Beetle, Casey's June	Dinacoma caseyi	Wherever found	\$8,052	\$0	\$8,052
		Beetle, Comal Springs riffle	Heterelmis comalensis	Wherever found	\$840	\$0	\$840
		Beetle, delta green ground	Elaphrus viridis	Wherever found	\$1,600	\$0	\$1,600
		Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$54,920	\$0	\$54,920
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$6,200	\$0	\$6,200
		Butterfly, callippe silverspot	Speyeria callippe callippe	Wherever found	\$5,000	\$0	\$5,000
		Butterfly, Fender's blue	Icaricia icarioides fenderi	Wherever found	\$211,169	\$0	\$211,169
		Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$19,280	\$0	\$19,280
		Butterfly, Miami Blue	Cyclargus (=Hemiargus) thomasi bethunebakeri	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Butterfly, mission blue	Icaricia icarioides missionensis	Wherever found	\$1,600	\$0	\$1,600
		Butterfly, Mitchell's satyr	Neonympha mitchellii mitchellii	Wherever found	\$1,055	\$0	\$1,055
		Butterfly, Quino checkerspot	Euphydryas editha quino (=E. e. wrighti)	Wherever found	\$2,684	\$0	\$2,684
		Butterfly, San Bruno elfin	Callophrys mossii bayensis	Wherever found	\$1,600	\$0	\$1,600
		Butterfly, Schaus swallowtail	Heraclides aristodemus ponceanus	Wherever found	\$2,004	\$0	\$2,004
		Butterfly, Smith's blue	Euphilotes enoptes smithi	Wherever found	\$5,600	\$0	\$5,600
		Moth, Kern primrose sphinx	Euproserpinus euterpe	Wherever found	\$5,368	\$0	\$5,368
		Tiger beetle, Northeastern beach	Cicindela dorsalis dorsalis	Wherever found	\$2,730	\$0	\$2,730
		Tiger beetle, Puritan	Cicindela puritana	Wherever found	\$7,240	\$0	\$7,240
		Tiger beetle, Salt Creek	Cicindela nevadica lincolniana	Wherever found	\$2,600	\$0	\$2,600
		<i>Insects Subtotal</i>			\$443,227	\$0	\$443,227
	Arachnids	Spider, spruce-fir moss	Microhexura montivaga	Wherever found	\$265	\$0	\$265
		<i>Arachnids Subtotal</i>					\$265
	Crustaceans	Amphipod, Hay's Spring	Stygobromus hayi	Wherever found	\$240	\$0	\$240
		Crayfish, Big Sandy	Cambarus callainus	Wherever found	\$7,486	\$0	\$7,486
		Crayfish, Nashville	Orconectes shoupi	Wherever found	\$3,442	\$0	\$3,442
		Fairy shrimp, Conservancy	Branchinecta conservatio	Wherever found	\$15,265	\$0	\$15,265
		Fairy shrimp, longhorn	Branchinecta longiantenna	Wherever found	\$3,400	\$0	\$3,400
		Fairy shrimp, San Diego	Branchinecta sandiegonensis	Wherever found	\$5,002	\$0	\$5,002
		Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$68,596	\$0	\$68,596
		Isopod, Madison Cave	Antrolana lira	Wherever found	\$240	\$0	\$240
		Shrimp, California freshwater	Syncaris pacifica	Wherever found	\$9,600	\$0	\$9,600
		Tadpole shrimp, vernal pool	Lepidurus packardi	Wherever found	\$61,931	\$0	\$61,931
		<i>Crustaceans Subtotal</i>			\$175,202	\$0	\$175,202
	Flowering Plants	Alopecurus, Sonoma	Alopecurus aequalis var. sonomensis	Wherever found	\$2,400	\$0	\$2,400



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Amaranth, seabeach	<i>Amaranthus pumilus</i>	Wherever found	\$106,773	\$0	\$106,773
		Aster, decurrent false	<i>Boltonia decurrens</i>	Wherever found	\$56,044	\$0	\$56,044
		Avens, spreading	<i>Geum radiatum</i>	Wherever found	\$125	\$0	\$125
		Barbara's Buttons, Mohr's	<i>Marshallia mohrii</i>	Wherever found	\$377	\$0	\$377
		Bird's beak, palmate-bracted	<i>Cordylanthus palmatus</i>	Wherever found	\$1,800	\$0	\$1,800
		Bird's-beak, soft	<i>Cordylanthus mollis</i> ssp.	Wherever found	\$25,800	\$0	\$25,800
		bladderpod, Missouri	<i>Physaria filiformis</i>	Wherever found	\$142	\$0	\$142
		Bladderpod, Short's	<i>Physaria globosa</i>	Wherever found	\$25,783	\$0	\$25,783
		Bladderpod, Spring Creek	<i>Lesquerella perforata</i>	Wherever found	\$1,600	\$0	\$1,600
		Brodiaea, thread-leaved	<i>Brodiaea filifolia</i>	Wherever found	\$5,368	\$0	\$5,368
		Bulrush, Northeastern	<i>Scirpus ancistrochaetus</i>	Wherever found	\$1,840	\$0	\$1,840
		Butterfly plant, Colorado	<i>Gaura neomexicana</i> var.	Wherever found	\$1,800	\$0	\$1,800
		Cactus, Key tree	<i>Pilosocereus robinii</i>	Wherever found	\$1,342	\$0	\$1,342
		cactus, Tobusch fishhook	<i>Sclerocactus brevihamatus</i> ssp. <i>tobuschii</i>	Wherever found	\$2,100	\$0	\$2,100
		Campion, fringed	<i>Silene polypetala</i>	Wherever found	\$1,912	\$0	\$1,912
		Catchfly, Spalding's	<i>Silene spaldingii</i>	Wherever found	\$4,480	\$0	\$4,480
		Chaffseed, American	<i>Schwalbea americana</i>	Wherever found	\$410	\$0	\$410
		Checker-mallow, Keck's	<i>Sidalcea keckii</i>	Wherever found	\$2,593	\$0	\$2,593
		Clarkia, Springville	<i>Clarkia springvillensis</i>	Wherever found	\$4,019	\$0	\$4,019
		Clover, running buffalo	<i>Trifolium stoloniferum</i>		\$43,718	\$0	\$43,718
		Clover, showy Indian	<i>Trifolium amoenum</i>	Wherever found	\$3,200	\$0	\$3,200
		Coneflower, smooth	<i>Echinacea laevigata</i>	Wherever found	\$650	\$0	\$650
		Daisy, Lakeside	<i>Hymenoxys herbacea</i>	Wherever found	\$3,076	\$0	\$3,076
		Daisy, Willamette	<i>Erigeron decumbens</i>	Wherever found	\$204,219	\$0	\$204,219
		Dawn-flower, Texas prairie	<i>Hymenoxys texana</i>	Wherever found	\$5,000	\$0	\$5,000
		Dropwort, Canby's	<i>Oxypolis canbyi</i>	Wherever found	\$240	\$0	\$240
		Fiddleneck, large-flowered	<i>Amsinckia grandiflora</i>	Wherever found	\$1,600	\$0	\$1,600
		Fleabane, Zuni	<i>Erigeron rhizomatus</i>	Wherever found	\$1,907	\$0	\$1,907
		Four-o'clock, MacFarlane's	<i>Mirabilis macfarlanei</i>	Wherever found	\$1,157	\$0	\$1,157
		Gerardia, sandplain	<i>Agalinis acuta</i>	Wherever found	\$1,240	\$0	\$1,240
		Gilia, Monterey	<i>Gilia tenuiflora</i> ssp. <i>arenaria</i>	Wherever found	\$1,600	\$0	\$1,600

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Glade cress, Kentucky	<i>Leavenworthia exigua laciniata</i>	Wherever found	\$5,920	\$0	\$5,920
		Goldenrod, Houghton's	<i>Solidago houghtonii</i>	Wherever found	\$513	\$0	\$513
		Goldenrod, Short's	<i>Solidago shortii</i>	Wherever found	\$1,332	\$0	\$1,332
		Goldfields, Burke's	<i>Lasthenia burkei</i>	Wherever found	\$21,400	\$0	\$21,400
		Goldfields, Contra Costa	<i>Lasthenia conjugens</i>	Wherever found	\$10,400	\$0	\$10,400
		Gourd, Okeechobee	<i>Cucurbita okeechobeensis</i> ssp. <i>okeechobeensis</i>	Wherever found	\$500	\$0	\$500
		Grass, Colusa	<i>Neostapfia colusana</i>	Wherever found	\$4,690	\$0	\$4,690
		Grass, Solano	<i>Tuctoria mucronata</i>	Wherever found	\$800	\$0	\$800
		Grass, Tennessee yellow-eyed	<i>Xyris tennesseensis</i>	Wherever found	\$1,048	\$0	\$1,048
		Ground-plum, Guthrie's (=Pyne's)	<i>Astragalus bibullatus</i>	Wherever found	\$650	\$0	\$650
		Harperella	<i>Ptilimnium nodosum</i>	Wherever found	\$2,313	\$0	\$2,313
		Heartleaf, dwarf-flowered	<i>Hexastylis naniflora</i>	Wherever found	\$22,568	\$0	\$22,568
		Iris, dwarf lake	<i>Iris lacustris</i>	Wherever found	\$342	\$0	\$342
		Joint-vetch, Sensitive	<i>Aeschynomene virginica</i>	Wherever found	\$530	\$0	\$530
		Ladies'-tresses, Ute	<i>Spiranthes diluvialis</i>	Wherever found	\$37,085	\$0	\$37,085
		Ladies-tresses, Navasota	<i>Spiranthes parksii</i>	Wherever found	\$840	\$0	\$840
		Layia, beach	<i>Layia carnosa</i>	Wherever found	\$2,400	\$0	\$2,400
		Lily, Minnesota dwarf trout	<i>Erythronium propullans</i>	Wherever found	\$3,200	\$0	\$3,200
		Lomatium, Bradshaw's	<i>Lomatium bradshawii</i>	Wherever found	\$190,619	\$0	\$190,619
		Loosestrife, rough-leaved	<i>Lysimachia asperulaefolia</i>	Wherever found	\$25,250	\$0	\$25,250
		Lupine, Kincaid's	<i>Lupinus sulphureus</i> ssp. <i>kincaidii</i>	Wherever found	\$196,919	\$0	\$196,919
		Manzanita, Ione	<i>Arctostaphylos myrtifolia</i>	Wherever found	\$1,803	\$0	\$1,803
		Manzanita, pallid	<i>Arctostaphylos pallida</i>	Wherever found	\$200	\$0	\$200
		Meadowfoam, Butte County	<i>Limnanthes floccosa</i> ssp. <i>californica</i>	Wherever found	\$2,400	\$0	\$2,400
		Meadowfoam, Sebastopol	<i>Limnanthes vinculans</i>	Wherever found	\$23,800	\$0	\$23,800
		Milk-vetch, Braunton's	<i>Astragalus brauntonii</i>	Wherever found	\$2,684	\$0	\$2,684

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Milkweed, Mead's	<i>Asclepias meadii</i>	Wherever found	\$6,081	\$0	\$6,081
		Monkshood, northern wild	<i>Aconitum noveboracense</i>	Wherever found	\$1,779	\$0	\$1,779
		Navarretia, many-flowered	<i>Navarretia leucocephala</i> ssp. <i>plieantha</i>	Wherever found	\$2,400	\$0	\$2,400
		No common name	<i>Geocarpon minimum</i>	Wherever found	\$5,587	\$0	\$5,587
		Orchid, eastern prairie fringed	<i>Platanthera leucophaea</i>	Wherever found	\$22,376	\$0	\$22,376
		Orchid, western prairie fringed	<i>Platanthera praeclara</i>	Wherever found	\$3,500	\$0	\$3,500
		Orchid, white fringeless	<i>Platanthera integrilabia</i>		\$656	\$0	\$656
		Orcutt grass, hairy	<i>Orcuttia pilosa</i>	Wherever found	\$1,956	\$0	\$1,956
		Orcutt grass, Sacramento	<i>Orcuttia viscida</i>	Wherever found	\$3,200	\$0	\$3,200
		Orcutt grass, San Joaquin	<i>Orcuttia inaequalis</i>	Wherever found	\$2,378	\$0	\$2,378
		Orcutt grass, slender	<i>Orcuttia tenuis</i>	Wherever found	\$3,200	\$0	\$3,200
		Owl's-clover, fleshy	<i>Castilleja campestris</i> ssp. <i>succulenta</i>	Wherever found	\$3,130	\$0	\$3,130
		paintbrush, golden	<i>Castilleja levisecta</i>	Wherever found	\$162,119	\$0	\$162,119
		Pawpaw, beautiful	<i>Deeringothamnus pulchellus</i>	Wherever found	\$100	\$0	\$100
		Pentachaeta, white-rayed	<i>Pentachaeta bellidiflora</i>	Wherever found	\$2,400	\$0	\$2,400
		Pink, swamp	<i>Helonias bullata</i>	Wherever found	\$650	\$0	\$650
		Pinkroot, gentian	<i>Spigelia gentianoides</i>	Wherever found	\$2,119	\$0	\$2,119
		Piperia, Yadon's	<i>Piperia yadonii</i>	Wherever found	\$1,600	\$0	\$1,600
		Pitcher-plant, Alabama canebrake	<i>Sarracenia rubra</i> ssp. <i>alabamensis</i>	Wherever found	\$399	\$0	\$399
		Pitcher-plant, mountain sweet	<i>Sarracenia rubra</i> ssp. <i>jonesii</i>	Wherever found	\$2,867	\$0	\$2,867
		Pogonia, small whorled	<i>Isotria medeoloides</i>	Wherever found	\$26,777	\$0	\$26,777
		Pondberry	<i>Lindera melissifolia</i>	Wherever found	\$1,048	\$0	\$1,048
		Poppy-mallow, Texas	<i>Callirhoe scabriuscula</i>	Wherever found	\$420	\$0	\$420
		potato-bean, Prices	<i>Apios priceana</i>		\$3,058	\$0	\$3,058
		Potentilla, Hickman's	<i>Potentilla hickmanii</i>	Wherever found	\$2,400	\$0	\$2,400
		Prairie-clover, leafy	<i>Dalea foliosa</i>	Wherever found	\$826	\$0	\$826
		rock cress, Shale barren	<i>Arabis serotina</i>	Wherever found	\$240	\$0	\$240

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Rock-cress, Braun's	<i>Arabis perstellata</i>	Wherever found	\$2,349	\$0	\$2,349
		Rockcress, Georgia	<i>Arabis georgiana</i>	Wherever found	\$1,826	\$0	\$1,826
		Rosemary, Cumberland	<i>Conradina verticillata</i>	Wherever found	\$376	\$0	\$376
		Rosemary, Etonia	<i>Conradina etonia</i>	Wherever found	\$125	\$0	\$125
		roseroot, Leedy's	<i>Rhodiola integrifolia</i> ssp.	Wherever found	\$723	\$0	\$723
		Sand-verbena, large-fruited	<i>Abronia macrocarpa</i>	Wherever found	\$2,100	\$0	\$2,100
		Sandwort, Cumberland	<i>Arenaria cumberlandensis</i>	Wherever found	\$251	\$0	\$251
		Seablite, California	<i>Suaeda californica</i>		\$1,600	\$0	\$1,600
		Seagrass, Johnson's	<i>Halophila johnsonii</i>	Wherever found	\$50,488	\$0	\$50,488
		Skullcap, large-flowered	<i>Scutellaria montana</i>	Wherever found	\$1,276	\$0	\$1,276
		Sneezeweed, Virginia	<i>Helenium virginicum</i>	Wherever found	\$240	\$0	\$240
		Spineflower, Monterey	<i>Chorizanthe pungens</i> var. <i>pungens</i>	Wherever found	\$4,000	\$0	\$4,000
		spineflower, Robust	<i>Chorizanthe robusta</i> var. <i>robusta</i>	Wherever found	\$1,600	\$0	\$1,600
		Spiraea, Virginia	<i>Spiraea virginiana</i>	Wherever found	\$3,994	\$0	\$3,994
		Spurge, Hoover's	<i>Chamaesyce hooveri</i>	Wherever found	\$100	\$0	\$100
		Sumac, Michaux's	<i>Rhus michauxii</i>	Wherever found	\$5,260	\$0	\$5,260
		Sunburst, Hartweg's golden	<i>Pseudobahia bahiifolia</i>	Wherever found	\$3,564	\$0	\$3,564
		Sunburst, San Joaquin adobe	<i>Pseudobahia peirsonii</i>	Wherever found	\$4,616	\$0	\$4,616
		Sunflower, Pecos (=puzzle, =paradox)	<i>Helianthus paradoxus</i>	Wherever found	\$1,650	\$0	\$1,650
		Sunflower, Schweinitz's	<i>Helianthus schweinitzii</i>	Wherever found	\$15,425	\$0	\$15,425
		Sunshine, Sonoma	<i>Blennosperma bakeri</i>	Wherever found	\$22,200	\$0	\$22,200
		Tarplant, Santa Cruz	<i>Holocarpha macradenia</i>	Wherever found	\$4,000	\$0	\$4,000
		Thistle, La Graciosa	<i>Cirsium loncholepis</i>	Wherever found	\$5,368	\$0	\$5,368
		Thistle, Pitcher's	<i>Cirsium pitcheri</i>	Wherever found	\$2,545	\$0	\$2,545
		Thistle, Suisun	<i>Cirsium hydrophilum</i> var. <i>hydrophilum</i>	Wherever found	\$8,200	\$0	\$8,200
		Thornmint, San Mateo	<i>Acanthomintha obovata</i> ssp. <i>duttonii</i>	Wherever found	\$100	\$0	\$100
		Trillium, relict	<i>Trillium reliquum</i>	Wherever found	\$5,538	\$0	\$5,538

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Tuctoria, Greene's	Tuctoria greenei	Wherever found	\$424	\$0	\$424
		Wallflower, Ben Lomond	Erysimum teretifolium	Wherever found	\$2,400	\$0	\$2,400
		Water-plantain, Kral's	Sagittaria secundifolia	Wherever found	\$299	\$0	\$299
		Wild-rice, Texas	Zizania texana	Wherever found	\$840	\$0	\$840
		Woolly-star, Santa Ana River	Eriastrum densifolium ssp. sanctorum	Wherever found	\$2,318	\$0	\$2,318
	<i>Flowering Plants Subtotal</i>				\$1,493,582	\$0	\$1,493,582
	Conifers and Cycads	Torreya, Florida	Torreya taxifolia	Wherever found	\$2,232	\$0	\$2,232
	<i>Conifers and Cycads Subtotal</i>						
	Ferns and Allies	Fern, Alabama streak-sorus	Thelypteris pilosa var. alabamensis	Wherever found	\$299	\$0	\$299
		Fern, American hart's-tongue	Asplenium scolopendrium var. americanum	Wherever found	\$125	\$0	\$125
		Quillwort, Louisiana	Isoetes louisianensis	Wherever found	\$2,968	\$0	\$2,968
	<i>Ferns and Allies Subtotal</i>				\$3,392	\$0	\$3,392
	<i>Domestic Subtotal</i>				\$190,660,844	\$1,341,073	\$192,001,917
Foreign	Mammals	Bear, brown	Ursus arctos pruinosus	Wherever found	\$4,073	\$0	\$4,073
	<i>Mammals Subtotal</i>				\$4,073		\$4,073
	Reptiles	Sea turtle, loggerhead	Caretta caretta	Northeast Atlantic Ocean DPS	\$195,843	\$0	\$195,843
		Sea turtle, loggerhead	Caretta caretta	South Atlantic	\$390,716	\$0	\$390,716
	<i>Reptiles Subtotal</i>				\$586,559	\$0	\$586,559
	<i>Foreign Subtotal</i>				\$590,632	\$0	\$590,632
	Multi-Species				\$1,165,366	\$0	\$1,165,366
	<i>Multi-Species Subtotal</i>				\$1,165,366	\$0	\$1,165,366
<b>Grand Total</b>					\$192,416,842	\$1,341,073	\$193,757,915

## DOD Marine Corps - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$231,000	\$0	\$231,000
		Bat, Indiana	Myotis sodalis	Wherever found	\$40,000	\$0	\$40,000
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$40,000	\$0	\$40,000
		Kangaroo rat, Stephens'	Dipodomys stephensi (incl. D. cascus)	Wherever found	\$331,000	\$0	\$331,000
		Manatee, West Indian	Trichechus manatus	Wherever found	\$5,000	\$0	\$5,000
		Mouse, Pacific pocket	Perognathus longimembris pacificus	Wherever found	\$347,200	\$0	\$347,200
		Pronghorn, Sonoran	Antilocapra americana sonoriensis	U.S.A. (AZ), Mexico	\$93,100	\$0	\$93,100
		Seal, Hawaiian monk	Monachus schauinslandi	Wherever found	\$1,000	\$0	\$1,000
	<i>Mammals Subtotal</i>				<i>\$1,088,300</i>	<i>\$0</i>	<i>\$1,088,300</i>
	Birds	Coot, Hawaiian	Fulica americana alai	Wherever found	\$13,000	\$0	\$13,000
		Duck, Hawaiian (=koloa)	Anas wyvilliana	Wherever found	\$13,000	\$0	\$13,000
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$248,200	\$0	\$248,200
		gallinule, Hawaiian common	Gallinula galeata sandvicensis	Wherever found	\$13,000	\$0	\$13,000
		Gnatcatcher, coastal California	Polioptila californica californica	Wherever found	\$1,928,600	\$0	\$1,928,600
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$356,200	\$0	\$356,200
		Stilt, Hawaiian	Himantopus mexicanus	Wherever found	\$89,000	\$0	\$89,000
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$18,500	\$0	\$18,500
		Tern, California least	Sterna antillarum browni	Wherever found	\$409,600	\$0	\$409,600
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$275,600	\$0	\$275,600

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$394,500	\$0	\$394,500
	<i>Birds Subtotal</i>				\$3,759,200	\$0	\$3,759,200
	Reptiles	Alligator, American	Alligator mississippiensis	Wherever found	\$10,000	\$0	\$10,000
		Sea turtle, green	Chelonia mydas	Central North Pacific DPS	\$1,000	\$0	\$1,000
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$15,800	\$0	\$15,800
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$6,720,500	\$0	\$6,720,500
	<i>Reptiles Subtotal</i>				\$6,747,300	\$0	\$6,747,300
	Amphibians	Salamander, frosted flatwoods	Ambystoma cingulatum	Wherever found	\$8,000	\$0	\$8,000
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$192,100	\$0	\$192,100
	<i>Amphibians Subtotal</i>				\$200,100	\$0	\$200,100
	Fishes	Goby, tidewater	Eucyclogobius newberryi	Wherever found	\$252,800	\$0	\$252,800
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$216,300	\$0	\$216,300
	<i>Fishes Subtotal</i>				\$469,100	\$0	\$469,100
	Insects	Yellow-faced bee, anthricinan	Hylaeus anthracinus	Wherever found	\$200	\$0	\$200
	<i>Insects Subtotal</i>				\$200	\$0	\$200
	Crustaceans	Fairy shrimp, Riverside	Streptocephalus woottoni	Wherever found	\$175,000	\$0	\$175,000
		Fairy shrimp, San Diego	Branchinecta sandiegonensis	Wherever found	\$268,900	\$0	\$268,900
	<i>Crustaceans Subtotal</i>				\$443,900	\$0	\$443,900
	Flowering Plants	Baccharis, Encinitas	Baccharis vanessae	Wherever found	\$200,500	\$0	\$200,500
		Brodiaea, thread-leaved	Brodiaea filifolia	Wherever found	\$161,500	\$0	\$161,500
		Button-celery, San Diego	Eryngium aristulatum var. parishii	Wherever found	\$161,500	\$0	\$161,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Navarretia, spreading	Navarretia fossalis	Wherever found	\$161,600	\$0	\$161,600
		Ohai	Sesbania tomentosa	Wherever found	\$200	\$0	\$200
		Pondberry	Lindera melissifolia	Wherever found	\$14,000	\$0	\$14,000
	<i>Flowering Plants Subtotal</i>				<i>\$699,300</i>	<i>\$0</i>	<i>\$699,300</i>
<i>Domestic Subtotal</i>					<i>\$13,407,400</i>	<i>\$0</i>	<i>\$13,407,400</i>
Total					\$13,407,400	\$0	\$13,407,400



## DOD Navy - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$3,455	\$0	\$3,455
		Bat, Indiana	Myotis sodalis	Wherever found	\$238	\$0	\$238
		Bat, Mariana fruit (=Mariana flying fox)	Pteropus mariannus mariannus	Wherever found	\$65,302	\$0	\$65,302
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$357,972	\$0	\$357,972
		Bear, polar	Ursus maritimus	Wherever found	\$4,234	\$0	\$4,234
		Deer, key	Odocoileus virginianus clavium	Wherever found	\$942	\$0	\$942
		dolphin, Maui	Cephalorhynchus hectori maui		\$158,636	\$0	\$158,636
		Fox, San Joaquin kit	Vulpes macrotis mutica	wherever found	\$2,519	\$0	\$2,519
		Kangaroo rat, Fresno	Dipodomys nitratooides exilis	Wherever found	\$3,704	\$0	\$3,704
		Kangaroo rat, Stephens'	Dipodomys stephensi (incl. D. cascus)	Wherever found	\$58,204	\$0	\$58,204
		Manatee, West Indian	Trichechus manatus	Wherever found	\$25,283	\$0	\$25,283
		Otter, southern sea	Enhydra lutris nereis	Wherever found	\$97,622	\$0	\$97,622
		Rabbit, Lower Keys marsh	Sylvilagus palustris hefneri	Wherever found	\$942	\$0	\$942
		Rice rat, Silver	Oryzomys palustris natator	lower FL Keys	\$942	\$0	\$942
		Seal, bearded	Erignathus barbatus nauticus	Beringia DPS	\$60,000	\$0	\$60,000
		Seal, Guadalupe fur	Arctocephalus townsendi	Wherever found	\$165,842	\$0	\$165,842
		Seal, Hawaiian monk	Monachus schauinslandi	Wherever found	\$308,589	\$0	\$308,589
		Shrew, Buena Vista Lake ornate	Sorex ornatus relictus	Wherever found	\$11,290	\$0	\$11,290
		whale, beluga	Delphinapterus leucas	Cook Inlet DPS	\$158,636	\$0	\$158,636
		Whale, blue	Balaenoptera musculus	Wherever found	\$556,627	\$0	\$556,627
		Whale, bowhead	Balaena mysticetus	Wherever found	\$158,635	\$0	\$158,635
		whale, false killer	Pseudorca crassidens	Main Hawaiian Is	\$278,183	\$0	\$278,183
		Whale, finback	Balaenoptera physalus	Wherever found	\$756,534	\$0	\$756,534
		Whale, gray	Eschrichtius robustus	Western North Pa	\$161,635	\$0	\$161,635
		Whale, humpback	Megaptera novaeangliae	Central America I	\$9,369	\$0	\$9,369
		Whale, humpback	Megaptera novaeangliae	Mexico DPS	\$152,144	\$0	\$152,144

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Whale, humpback	Megaptera novaeangliae	Western North Pa	\$258,231	\$0	\$258,231
		Whale, killer	Orcinus orca	Southern Resident	\$178,730	\$0	\$178,730
		Whale, North Atlantic Right	Eubalaena glacialis	Wherever found	\$582,232	\$0	\$582,232
		Whale, Sei	Balaenoptera borealis	Wherever found	\$274,413	\$0	\$274,413
		Whale, sperm	Physeter catodon (=macrocephalus)	Wherever found	\$686,227	\$0	\$686,227
	<i>Mammals Subtotal</i>				<i>\$5,537,312</i>	<i>\$0</i>	<i>\$5,537,312</i>
	Birds	Albatross, short-tailed	Phoebastria (=Diomedea) albatrus	Wherever found	\$758	\$0	\$758
		Coot, Hawaiian	Fulica americana alai	Wherever found	\$7,349	\$0	\$7,349
		Crow, Mariana (=aga)	Corvus kubaryi	Wherever found	\$49,610	\$0	\$49,610
		Duck, Hawaiian (=koloa)	Anas wyvilliana	Wherever found	\$6,900	\$0	\$6,900
		Eider, spectacled	Somateria fischeri	Wherever found	\$3,686	\$0	\$3,686
		Eider, Steller's	Polysticta stelleri	AK breeding pop.	\$3,686	\$0	\$3,686
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$1,992	\$0	\$1,992
		gallinule, Hawaiian common	Gallinula galeata sandvicensis	Wherever found	\$7,349	\$0	\$7,349
		Gnatcatcher, coastal California	Polioptila californica californica	Wherever found	\$74,982	\$0	\$74,982
		Goose, Hawaiian	Branta (=Nesochen) sandvicensis	Wherever found	\$11,448	\$0	\$11,448
		Kingfisher, Guam Micronesian	Halcyon cinnamomina cinnamomina	Wherever found	\$59,002	\$0	\$59,002
		Knot, red	Calidris canutus rufa	Wherever found	\$11,268	\$0	\$11,268
		Megapode, Micronesian	Megapodius laperouse	Wherever found	\$47,617	\$0	\$47,617
		Moorhen, Mariana common	Gallinula chloropus guami	Wherever found	\$6,221	\$0	\$6,221
		Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR,	\$318,587	\$0	\$318,587
		petrel, Hawaiian	Pterodroma sandwichensis	Wherever found	\$758	\$0	\$758
		Plover, piping	Charadrius melodus	[Atlantic Coast an	\$7,926	\$0	\$7,926
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast popu	\$527,144	\$0	\$527,144
		Rail, Guam	Rallus owstoni	Wherever found, e	\$44,861	\$0	\$44,861

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Rail, light-footed clapper	Rallus longirostris levipes	Wherever found	\$152,909	\$0	\$152,909
		Shearwater, Newell's Townsend's	Puffinus auricularis newelli	Wherever found	\$758	\$0	\$758
		Shrike, San Clemente loggerhead	Lanius ludovicianus mearnsi	Wherever found	\$2,597,414	\$0	\$2,597,414
		Sparrow, San Clemente sage	Amphispiza belli clementeae	Wherever found	\$967,094	\$0	\$967,094
		Stilt, Hawaiian	Himantopus mexicanus knudseni	Wherever found	\$15,452	\$0	\$15,452
		Storm-petrel, band-rumped	Oceanodroma castro	USA (HI)	\$758	\$0	\$758
		Swiftlet, Mariana gray	Aerodramus vanikorensis bartschi	Wherever found	\$115,296	\$0	\$115,296
		Tern, California least	Sterna antillarum browni	Wherever found	\$530,253	\$0	\$530,253
		Tern, least	Sterna antillarum	interior pop.	\$4,000	\$0	\$4,000
		Tern, roseate	Sterna dougallii dougallii	Northeast U.S. nes	\$3,926	\$0	\$3,926
		Towhee, Inyo California	Pipilo crissalis eremophilus	Wherever found	\$202	\$0	\$202
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$31,375	\$0	\$31,375
		'Iwi	Drepanis coccinea	Wherever found	\$2,341	\$0	\$2,341
	<i>Birds Subtotal</i>				<i>\$5,612,922</i>	<i>\$0</i>	<i>\$5,612,922</i>
	Reptiles	Alligator, American	Alligator mississippiensis	Wherever found	\$2,568	\$0	\$2,568
		Crocodile, American	Crocodylus acutus	U.S.A. (FL)	\$942	\$0	\$942
		Sea turtle, green	Chelonia mydas	Central North Pac	\$286,718	\$0	\$286,718
		Sea turtle, green	Chelonia mydas	East Pacific DPS	\$159,662	\$0	\$159,662
		Sea turtle, green	Chelonia mydas	North Atlantic DP	\$20,399	\$0	\$20,399
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$132,831	\$0	\$132,831
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$23,399	\$0	\$23,399
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$22,974	\$0	\$22,974
		Sea turtle, loggerhead	Caretta caretta	North Pacific Oce	\$1,000	\$0	\$1,000
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlanti	\$27,179	\$0	\$27,179
		Sea turtle, olive ridley	Lepidochelys olivacea	Wherever found, e	\$14,873	\$0	\$14,873
	<i>Reptiles Subtotal</i>				<i>\$692,545</i>	<i>\$0</i>	<i>\$692,545</i>

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Amphibians	Coqui, Llanero	Eleutherodactylus juanariveroi	Wherever found	\$652,524	\$0	\$652,524
		Frog, California red-legged	Rana draytonii	Wherever found	\$232,009	\$0	\$232,009
		Frog, mountain yellow-legged	Rana muscosa	Southern Californ	\$5,608	\$0	\$5,608
		Salamander, Barton Springs	Eurycea sosorum	Wherever found	\$336,224	\$0	\$336,224
		Salamander, Cheat Mountain	Plethodon nettingi	Wherever found	\$388,136	\$0	\$388,136
		salamander, Reticulated flatwoods	Ambystoma bishopi	Wherever found	\$290	\$0	\$290
		Salamander, Salado	Eurycea chisholmensis	Wherever found	\$89,216	\$0	\$89,216
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$188,909	\$0	\$188,909
		Toad, Houston	Bufo houstonensis	Wherever found	\$70,786	\$0	\$70,786
		Toad, Puerto Rican crested	Peltophryne lemur	Wherever found	\$142,491	\$0	\$142,491
	<i>Amphibians Subtotal</i>				<i>\$2,106,193</i>	<i>\$0</i>	<i>\$2,106,193</i>
	Fishes	Bocaccio	Sebastes paucispinis	Puget Sound - Ge	\$20,676	\$0	\$20,676
		Goby, tidewater	Eucyclogobius newberryi	Wherever found	\$2,250	\$0	\$2,250
		rockfish, canary	Sebastes pinniger	Puget Sound - Ge	\$700	\$0	\$700
		rockfish, yelloweye	Sebastes ruberrimus	Puget Sound - Ge	\$20,676	\$0	\$20,676
		Salmon, Atlantic	Salmo salar	Gulf of Maine DP	\$14,168	\$0	\$14,168
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal	\$700	\$0	\$700
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spr	\$700	\$0	\$700
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spr	\$700	\$0	\$700
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$700	\$0	\$700
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River	\$700	\$0	\$700

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-r	\$700	\$0	\$700
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring	\$700	\$0	\$700
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia s	\$111,999	\$0	\$111,999
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette	\$700	\$0	\$700
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU	\$700	\$0	\$700
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$700	\$0	\$700
		Sawfish, largetooth	Pristis pristis	Wherever found	\$700	\$0	\$700
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smallto	\$10,597	\$0	\$10,597
		Shark, Scalloped Hammerhead	Sphyrna lewini	Eastern Pacific Df	\$9,902	\$0	\$9,902
		Shark, Scalloped Hammerhead	Sphyrna lewini	Indo-West Pacific	\$3,127	\$0	\$3,127
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central	\$700	\$0	\$700
		Steelhead	Oncorhynchus (=Salmo) mykiss	Central California	\$700	\$0	\$700
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia I	\$99,461	\$0	\$99,461
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia	\$700	\$0	\$700
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia	\$700	\$0	\$700
		Steelhead	Oncorhynchus (=Salmo) mykiss	Northern Californ	\$700	\$0	\$700
		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$63,655	\$0	\$63,655

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin	\$700	\$0	\$700
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California	\$700	\$0	\$700
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California	\$3,720	\$0	\$3,720
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River	\$700	\$0	\$700
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette	\$700	\$0	\$700
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Carolina DPS	\$3,494	\$0	\$3,494
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Chesapeake Bay I	\$4,497	\$0	\$4,497
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Gulf of Maine DP	\$51,384	\$0	\$51,384
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	New York Bight I	\$2,594	\$0	\$2,594
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	South Atlantic DP	\$3,494	\$0	\$3,494
		sturgeon, green	Acipenser medirostris	Southern DPS	\$700	\$0	\$700
		Sturgeon, Gulf	Acipenser oxyrinchus (=oxyrhynchus) desotoi	Wherever found	\$6,329	\$0	\$6,329
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$11,489	\$0	\$11,489
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous	\$121,218	\$0	\$121,218
		Tui chub, Mohave	Gila bicolor ssp. mohavensis	Wherever found	\$202	\$0	\$202
		<i>Fishes Subtotal</i>			<i>\$580,332</i>	<i>\$0</i>	<i>\$580,332</i>
	Clams	Pearlshell, Alabama	Margaritifera marrianae	Wherever found	\$290	\$0	\$290
		<i>Clams Subtotal</i>			<i>\$290</i>	<i>\$0</i>	<i>\$290</i>
	Snails	Abalone, Black	Haliotis cracherodii	Wherever found	\$633,460	\$0	\$633,460
		Abalone, White	Haliotis sorenseni	North America (W)	\$225,000	\$0	\$225,000
		Snail, fragile tree	Samoana fragilis	Wherever found	\$18,107	\$0	\$18,107

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Snail, Guam tree	Partula radiolata	Wherever found	\$47,454	\$0	\$47,454
		Snail, Humped tree	Partula gibba	Wherever found	\$43,794	\$0	\$43,794
	<i>Snails Subtotal</i>				<i>\$967,815</i>	<i>\$0</i>	<i>\$967,815</i>
	Insects	Butterfly, Mariana eight-spot	Hypolimnas octocula marianensis	Wherever found	\$33,749	\$0	\$33,749
		Butterfly, Mariana wandering	Vagrans egistina	Wherever found	\$4,927	\$0	\$4,927
		Butterfly, Palos Verdes blue	Glaucopsyche lygdamus palosverdesensis	Wherever found	\$60,522	\$0	\$60,522
		Butterfly, Quino checkerspot	Euphydryas editha quino (=E. e. wrighti)	Wherever found	\$178,848	\$0	\$178,848
		Butterfly, Smith's blue	Euphilotes enoptes smithi	Wherever found	\$761	\$0	\$761
		Checkerspot, Taylor's (=whulge)	Euphydryas editha taylori	Wherever found	\$12,182	\$0	\$12,182
		fly, Hawaiian picture-wing	Drosophila musaphilia	Wherever found	\$632	\$0	\$632
		Fly, Hawaiian picture-wing	Drosophila sharpi	Wherever found	\$632	\$0	\$632
		Tiger beetle, Northeastern beach	Cicindela dorsalis dorsalis	Wherever found	\$238	\$0	\$238
	<i>Insects Subtotal</i>				<i>\$292,491</i>	<i>\$0</i>	<i>\$292,491</i>
	Crustaceans	Fairy shrimp, San Diego	Branchinecta sandiegonensis	Wherever found	\$33,721	\$0	\$33,721
		Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$11,993	\$0	\$11,993
		Tadpole shrimp, vernal pool	Lepidurus packardi	Wherever found	\$4,881	\$0	\$4,881
	<i>Crustaceans Subtotal</i>				<i>\$50,595</i>	<i>\$0</i>	<i>\$50,595</i>
	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$1,642	\$0	\$1,642
		Coral, elkhorn	Acropora palmata	Wherever found	\$10,597	\$0	\$10,597
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$1,642	\$0	\$1,642
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$1,642	\$0	\$1,642
		Coral, Pillar	Dendrogyra cylindrus	Wherever found	\$1,642	\$0	\$1,642
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$1,642	\$0	\$1,642
		Coral, staghorn	Acropora cervicornis	Wherever found	\$10,597	\$0	\$10,597
		No common name	Acropora globiceps	Wherever found	\$56,055	\$0	\$56,055
		No common name	Acropora retusa	Wherever found	\$23,338	\$0	\$23,338

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		No common name	Acropora speciosa	Wherever found	\$7,217	\$0	\$7,217
		No common name	Pavona diffluens	Wherever found	\$31,467	\$0	\$31,467
		No common name	Seriatopora aculeata	Wherever found	\$28,055	\$0	\$28,055
	<i>Corals Subtotal</i>				<i>\$175,536</i>	<i>\$0</i>	<i>\$175,536</i>
	Flowering Plants	Alani	Melicope christophersenii	Wherever found	\$1,792	\$0	\$1,792
		Alani	Melicope saint-johnii	Wherever found	\$1,792	\$0	\$1,792
		Beaked-rush, Knieskern's	Rhynchospora knieskernii	Wherever found	\$14,874	\$0	\$14,874
		Bird's-beak, salt marsh	Cordylanthus maritimus ssp. maritimus	Wherever found	\$25,584	\$0	\$25,584
		Bush-mallow, San Clemente Island	Malacothamnus clementinus	Wherever found	\$158,417	\$0	\$158,417
		Button-celery, San Diego	Eryngium aristulatum var. parishii	Wherever found	\$24,890	\$0	\$24,890
		Gilia, Monterey	Gilia tenuiflora ssp. arenaria	Wherever found	\$3,044	\$0	\$3,044
		Haha	Cyanea calycina	Wherever found	\$1,761	\$0	\$1,761
		Hala pepe	Pleomele forbesii	Wherever found	\$2,572	\$0	\$2,572
		Iagu, Hayun (=Guam), Tronkon guafi (Rota))	Serianthes nelsonii	Wherever found	\$60,869	\$0	\$60,869
		Iliau, dwarf	Wilkesia hobdyi	Wherever found	\$11,229	\$0	\$11,229
		Ko`oloa`ula	Abutilon menziesii	Wherever found	\$1,734	\$0	\$1,734
		Kuahiwi laukahi	Plantago princeps	Wherever found	\$1,792	\$0	\$1,792
		Kulu`i	Nototrichium humile	Wherever found	\$1,792	\$0	\$1,792
		Larkspur, San Clemente Island	Delphinium variegatum ssp. kinkiense	Wherever found	\$145,951	\$0	\$145,951
		Lau `ehu	Panicum niihauense	Wherever found	\$6,300	\$0	\$6,300
		Lotus, San Clemente Island	Acmispon dendroideus var. traskiae (=Lotus d. ssp. traskiae)	Wherever found	\$145,951	\$0	\$145,951
		Mahoe	Alectryon macrococcus	Wherever found	\$1,734	\$0	\$1,734
		Mehamehame	Flueggea neowawraea	Wherever found	\$1,792	\$0	\$1,792
		Mesa-mint, San Diego	Pogogyne abramsii	Wherever found	\$24,890	\$0	\$24,890



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Nehe	Lipochaeta lobata var. leptophylla	Wherever found	\$1,792	\$0	\$1,792
		No common name	Abutilon sandwicense	Wherever found	\$1,734	\$0	\$1,734
		No common name	Bonamia menziesii	Wherever found	\$1,761	\$0	\$1,761
		No common name	Dendrobium guamense	Wherever found	\$33,473	\$0	\$33,473
		No common name	Eugenia bryanii	Wherever found	\$11,086	\$0	\$11,086
		No common name	Kadua parvula	Wherever found	\$1,792	\$0	\$1,792
		No common name	Lobelia niihauensis	Wherever found	\$1,792	\$0	\$1,792
		No common name	Maesa walkeri	Wherever found	\$22,270	\$0	\$22,270
		No common name	Neraudia angulata	Wherever found	\$1,792	\$0	\$1,792
		No common name	Nervilia jacksoniae	Wherever found	\$22,270	\$0	\$22,270
		No common name	Phyllanthus saffordii	Wherever found	\$21,020	\$0	\$21,020
		No common name	Platydesma cornuta var. decurrens	Wherever found	\$2,572	\$0	\$2,572
		No common name	Schiedea hookeri	Wherever found	\$2,572	\$0	\$2,572
		No common name	Silene perlmanii	Wherever found	\$2,572	\$0	\$2,572
		No common name	Spermolepis hawaiiensis	Wherever found	\$13,801	\$0	\$13,801
		No common name	Tabernaemontana rotensis	Wherever found	\$59,071	\$0	\$59,071
		No common name	Tetramolopium lepidotum ssp. lepidotum	Wherever found	\$2,572	\$0	\$2,572
		No common name	Tuberolabium guamense	Wherever found	\$44,680	\$0	\$44,680
		paintbrush, golden	Castilleja levisecta	Wherever found	\$14,227	\$0	\$14,227
		Paintbrush, San Clemente Island	Castilleja grisea	Wherever found	\$158,417	\$0	\$158,417
		Pamakani	Viola chamissoniana ssp. chamissoniana	Wherever found	\$2,572	\$0	\$2,572
		Pink, swamp	Helonias bullata	Wherever found	\$13,745	\$0	\$13,745
		Pogonia, small whorled	Isotria medeoloides	Wherever found	\$8,101	\$0	\$8,101
		Pu'uka`a	Cyperus trachysanthos	Wherever found	\$1,761	\$0	\$1,761
		Rockcress, Santa Cruz Island	Sibara filifolia	Wherever found	\$149,163	\$0	\$149,163
		silverbush, Blodgett's	Argythamnia blodgettii	Wherever found	\$942	\$0	\$942

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Spineflower, Monterey	Chorizanthe pungens var. pungens	Wherever found	\$3,044	\$0	\$3,044
		Spineflower, Orcutt's	Chorizanthe orcuttiana	Wherever found	\$139,084	\$0	\$139,084
		Spurge, Garber's	Chamaesyce garberi	Wherever found	\$942	\$0	\$942
		Ufa-halomtano	Heritiera longipetiolata	Wherever found	\$22,314	\$0	\$22,314
		Woodland-star, San Clemente Island	Lithophragma maximum	Wherever found	\$145,951	\$0	\$145,951
		`Akoko	Euphorbia kuwaleana	Wherever found	\$1,792	\$0	\$1,792
		`Akoko, Ewa Plains	Euphorbia skottsbergii var. skottsbergii	Wherever found	\$1,561	\$0	\$1,561
		`Anaunau	Lepidium arbuscula	Wherever found	\$1,792	\$0	\$1,792
	<i>Flowering Plants Subtotal</i>				<i>\$1,552,790</i>	<i>\$0</i>	<i>\$1,552,790</i>
	Conifers and Cycads	Fadang	Cycas micronesica	Wherever found	\$70,664	\$0	\$70,664
	<i>Conifers and Cycads Subtotal</i>				<i>\$70,664</i>	<i>\$0</i>	<i>\$70,664</i>
	Ferns and Allies	Ihi`ihi	Marsilea villosa	Wherever found	\$1,792	\$0	\$1,792
		No common name	Asplenium unisorum	Wherever found	\$1,734	\$0	\$1,734
	<i>Ferns and Allies Subtotal</i>				<i>\$3,526</i>	<i>\$0</i>	<i>\$3,526</i>
	Multi-Species				\$507,786	\$0	\$507,786
	<i>Multi-Species Subtotal</i>				<i>\$507,786</i>	<i>\$0</i>	<i>\$507,786</i>
<i>Domestic Total</i>					<i>\$18,150,797</i>	<i>\$0</i>	<i>\$18,150,797</i>
<b>Grand Total</b>					<b>\$18,150,797</b>	<b>\$0</b>	<b>\$18,150,797</b>

## DOE Bonneville Power Administration - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Fishes	Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$10,122,893	\$4,132	\$10,127,025
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$18,282,642	\$0	\$18,282,642
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$33,477,114	\$0	\$33,477,114
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$21,057,907	\$20,478	\$21,078,385
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$7,011,061	\$1,459,803	\$8,470,864
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$3,434,098	\$0	\$3,434,098
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$4,529,332	\$2,029,372	\$6,558,704
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$186,019	\$0	\$186,019
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$8,063,783	\$0	\$8,063,783
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$6,153,819	\$0	\$6,153,819
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$43,281,600	\$0	\$43,281,600
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$34,502,590	\$0	\$34,502,590
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$14,687,663	\$0	\$14,687,663
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$4,267,009	\$2,133,205	\$6,400,214

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$12,321,598	\$0	\$12,321,598
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$16,264,037	\$12,613,168	\$28,877,205
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$2,131,564	\$0	\$2,131,564
	<i>Fishes Subtotal</i>				<i>\$239,774,729</i>	<i>\$18,260,158</i>	<i>\$258,034,887</i>
	<i>Domestic Total</i>				<i>\$239,774,729</i>	<i>\$18,260,158</i>	<i>\$258,034,887</i>
	<b>Grand Total</b>				<b>\$239,774,729</b>	<b>\$18,260,158</b>	<b>\$258,034,887</b>

## DOE Southwestern Power Administration - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$5,237	\$0	\$5,237
		Bat, Indiana	Myotis sodalis	Wherever found	\$5,238	\$0	\$5,238
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$5,237	\$0	\$5,237
		Bat, Ozark big-eared	Corynorhinus (=Plecotus) townsendii ingens	Wherever found	\$5,237	\$0	\$5,237
	<i>Mammals Subtotal</i>				<i>\$20,949</i>	<i>\$0</i>	<i>\$20,949</i>
	Birds	Tern, least	Sterna antillarum	interior pop.	\$9,500	\$35,000	\$44,500
	<i>Birds Subtotal</i>				<i>\$9,500</i>	<i>\$35,000</i>	<i>\$44,500</i>
	Insects	Beetle, American burying	Nicrophorus americanus	Ex Pop, SW Missouri	\$9,200	\$0	\$9,200
	<i>Insects Subtotal</i>				<i>\$9,200</i>	<i>\$0</i>	<i>\$9,200</i>
	Multi-Species				\$17,900	\$0	\$17,900
	<i>Multi-Species Subtotal</i>				<i>\$17,900</i>	<i>\$0</i>	<i>\$17,900</i>
	<i>Domestic Total</i>				<i>\$57,549</i>	<i>\$35,000</i>	<i>\$92,549</i>
	<b>Grand Total</b>				<b>\$57,549</b>	<b>\$35,000</b>	<b>\$92,549</b>

## DOI Bureau of Indian Affairs - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Indiana	<i>Myotis sodalis</i>	Wherever found	\$55,300	\$0	\$55,300
		Bear, grizzly	<i>Ursus arctos horribilis</i>	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$189,150	\$0	\$189,150
		Ferret, black-footed	<i>Mustela nigripes</i>	Wherever found, except where listed as an experimental population	\$32,300	\$0	\$32,300
		Mouse, New Mexico meadow jumping	<i>Zapus hudsonius luteus</i>	Wherever found	\$65,000	\$0	\$65,000
		Panther, Florida	<i>Puma (=Felis) concolor coryi</i>	Wherever found	\$100,000	\$0	\$100,000
		Wolf, gray	<i>Canis lupus</i>	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$319,600	\$0	\$319,600
		Wolf, Mexican	<i>Canis lupus baileyi</i>	Wherever found, except where listed as an experimental population	\$28,200	\$0	\$28,200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	<i>Mammals Subtotal</i>				\$789,550	\$0	\$789,550
	Birds	Condor, California	Gymnogyps californianus	U.S.A. only, except where listed as an experimental population	\$146,500	\$0	\$146,500
		Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR, WA)	\$50,000	\$0	\$50,000
		Owl, northern spotted	Strix occidentalis caurina	Wherever found	\$404,000	\$0	\$404,000
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$85,000	\$0	\$85,000
	<i>Birds Subtotal</i>				\$685,500	\$0	\$685,500
	Amphibians						
		Frog, Oregon spotted	Rana pretiosa	Wherever found	\$18,000	\$0	\$18,000
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$90,000	\$0	\$90,000
	<i>Amphibians Subtotal</i>				\$108,000	\$0	\$108,000
	Fishes	Chub, Pahrnagat roundtail	Gila robusta jordani	Wherever found	\$64,800	\$0	\$64,800
		Minnow, Rio Grande Silvery	Hybognathus amarus	Wherever found, except where listed as an experimental population	\$62,000	\$0	\$62,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$119,300	\$0	\$119,300
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$99,500	\$0	\$99,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$79,700	\$0	\$79,700

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$13,625	\$0	\$13,625
		Sucker, Zuni bluehead	Catostomus discobolus yarrowi	Wherever found	\$17,500	\$0	\$17,500
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$24,600	\$0	\$24,600
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$105,000	\$0	\$105,000
	Fishes Subtotal				\$586,025	\$0	\$586,025
	Clams	Mucket, Neosho	Lampsilis rafinesqueana	Wherever found	\$119,800	\$0	\$119,800
	<i>Clams Subtotal</i>				<i>\$119,800</i>	<i>\$0</i>	<i>\$119,800</i>
	Insects	Skipper, Dakota	Hesperia dacotae	Wherever found	\$14,000	\$0	\$14,000
	<i>Insects Subtotal</i>				<i>\$14,000</i>	<i>\$0</i>	<i>\$14,000</i>
	<i>Domestic Subtotal</i>				<i>\$2,302,875</i>	<i>\$0</i>	<i>\$2,302,875</i>
	Total				\$2,302,875	\$0	\$2,302,875



## DOI Bureau of Land Management - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	<i>Myotis grisescens</i>	Wherever found	\$600	\$0	\$600
		Bat, Indiana	<i>Myotis sodalis</i>	Wherever found	\$600	\$0	\$600
		Bat, Mexican long-nosed	<i>Leptonycteris nivalis</i>	Wherever found	\$19,000	\$0	\$19,000
		Bat, Northern long-eared	<i>Myotis septentrionalis</i>	Wherever found	\$37,369	\$0	\$37,369
		Bear, grizzly	<i>Ursus arctos horribilis</i>	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$154,000	\$0	\$154,000
		Bear, polar	<i>Ursus maritimus</i>	Wherever found	\$226,000	\$0	\$226,000
		Caribou DPS, Southern Mountain	<i>Rangifer tarandus ssp. caribou</i>	Southern Mountain DPS	\$2,500	\$0	\$2,500
		Ferret, black-footed	<i>Mustela nigripes</i>	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$255,011	\$0	\$255,011
		Ferret, black-footed	<i>Mustela nigripes</i>	Wherever found, except where listed as an experimental population	\$557,000	\$0	\$557,000
		Fox, San Joaquin kit	<i>Vulpes macrotis mutica</i>	wherever found	\$91,000	\$0	\$91,000
		Jaguar	<i>Panthera onca</i>	Wherever found	\$4,800	\$0	\$4,800
		Kangaroo rat, giant	<i>Dipodomys ingens</i>	Wherever found	\$34,000	\$0	\$34,000
		Kangaroo rat, Stephens'	<i>Dipodomys stephensi</i> (incl. <i>D. cascus</i> )	Wherever found	\$145,000	\$0	\$145,000
		Kangaroo rat, Tipton	<i>Dipodomys nitratooides nitratooides</i>	Wherever found	\$10,000	\$0	\$10,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$305,303	\$0	\$305,303
		Manatee, West Indian	Trichechus manatus	Wherever found	\$1,100	\$0	\$1,100
		Mountain beaver, Point Arena	Aplodontia rufa nigra	Wherever found	\$5,500	\$0	\$5,500
		Mouse, Alabama beach	Peromyscus polionotus	Wherever found	\$500	\$0	\$500
		Mouse, New Mexico meadow jumping	Zapus hudsonius luteus	Wherever found	\$54,134	\$0	\$54,134
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	wherever found	\$9,466	\$0	\$9,466
		Ocelot	Leopardus (=Felis) pardalis	wherever found	\$4,800	\$0	\$4,800
		Prairie dog, Utah	Cynomys parvidens	Wherever found	\$222,400	\$0	\$222,400
		Pronghorn, Sonoran	Antilocapra americana sonoriensis	U.S.A. (AZ), Mexico	\$85,000	\$0	\$85,000
		Pronghorn, Sonoran	Antilocapra americana sonoriensis	Wherever found, except where listed as an experimental population	\$4,750	\$27,000	\$31,750
		Rabbit, Columbia Basin Pygmy	Brachylagus idahoensis	Columbia Basin DPS	\$18,600	\$0	\$18,600
		Sea lion, Steller	Eumetopias jubatus	Western DPS	\$500	\$0	\$500
		Seal, bearded	Erignathus barbatus nauticus	Beringia DPS	\$15,000	\$0	\$15,000
		Seal, Ringed	Phoca (=Pusa) hispida botnica	Wherever found	\$4,000	\$0	\$4,000
		Sheep, Peninsular bighorn	Ovis canadensis nelsoni	Peninsular CA pop.	\$15,500	\$0	\$15,500
		Sheep, Sierra Nevada bighorn	Ovis canadensis sierrae	Sierra Nevada	\$17,000	\$0	\$17,000
		Shrew, Buena Vista Lake ornate	Sorex ornatus relictus	Wherever found	\$1,000	\$0	\$1,000
		Squirrel, Northern Idaho Ground	Urocitellus brunneus	Wherever found	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Vole, Amargosa	Microtus californicus scirpensis	Wherever found	\$117,000	\$0	\$117,000
		Whale, blue	Balaenoptera musculus	Wherever found	\$500	\$0	\$500
		Whale, bowhead	Balaena mysticetus	Wherever found	\$500	\$0	\$500
		Whale, finback	Balaenoptera physalus	Wherever found	\$500	\$0	\$500
		Whale, gray	Eschrichtius robustus	Western North Pacific DPS	\$500	\$0	\$500
		Whale, humpback	Megaptera novaeangliae	Mexico DPS	\$500	\$0	\$500
		Whale, humpback	Megaptera novaeangliae	Western North Pacific	\$500	\$0	\$500
		Whale, North Pacific Right	Eubalaena japonica	Wherever found	\$500	\$0	\$500
		Whale, sperm	Physeter catodon (=macrocephalus)	Wherever found	\$500	\$0	\$500
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$39,700	\$0	\$39,700
		Wolf, Mexican	Canis lupus baileyi	U.S.A. (portions of AZ and NM)see 17.84(k)	\$290,500	\$0	\$290,500
	<i>Mammals Subtotal</i>				\$2,753,633	\$27,000	\$2,780,633
	Birds	Bobwhite, masked (quail)	Colinus virginianus ridgwayi	Wherever found	\$10,400	\$0	\$10,400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Condor, California	<i>Gymnogyps californianus</i>	U.S.A. (specific portions of Arizona, Nevada, and Utah)	\$18,230	\$0	\$18,230
		Condor, California	<i>Gymnogyps californianus</i>	U.S.A. only, except where listed as an experimental population	\$34,130	\$0	\$34,130
		Crane, Mississippi sandhill	<i>Grus canadensis pulla</i>	Wherever found	\$100	\$0	\$100
		Crane, whooping	<i>Grus americana</i>	U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	\$14,088	\$0	\$14,088
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$4,500	\$0	\$4,500
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$844,276	\$31,608	\$875,884
		Eider, spectacled	<i>Somateria fischeri</i>	Wherever found	\$48,000	\$0	\$48,000
		Eider, Steller's	<i>Polysticta stelleri</i>	AK breeding pop.	\$57,000	\$0	\$57,000
		Falcon, Northern Aplomado	<i>Falco femoralis septentrionalis</i>	U.S.A (AZ, NM)	\$114,500	\$0	\$114,500
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$933,749	\$0	\$933,749
		Gnatcatcher, coastal California	<i>Polioptila californica californica</i>	Wherever found	\$52,000	\$0	\$52,000
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$7,800	\$0	\$7,800
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$978,288	\$0	\$978,288

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$427,413	\$0	\$427,413
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$2,774,148	\$0	\$2,774,148
		Plover, piping	<i>Charadrius melodus</i>	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$36,488	\$0	\$36,488
		Plover, western snowy	<i>Charadrius nivosus nivosus</i>	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$388,000	\$0	\$388,000
		Rail, Yuma Ridgways	<i>Rallus obsoletus</i> [=longirostris] yumanensis	Wherever found	\$409,000	\$0	\$409,000
		sage-grouse, Gunnison	<i>Centrocercus minimus</i>	Wherever found	\$723,868	\$0	\$723,868
		Tern, California least	<i>Sterna antillarum browni</i>	Wherever found	\$1,500	\$0	\$1,500
		Tern, least	<i>Sterna antillarum</i>	interior pop.	\$30,488	\$0	\$30,488
		Towhee, Inyo California	<i>Pipilo crissalis eremophilus</i>	Wherever found	\$6,500	\$0	\$6,500
		Vireo, least Bell's	<i>Vireo bellii pusillus</i>	Wherever found	\$136,500	\$0	\$136,500
		Woodpecker, red-cockaded	<i>Picoides borealis</i>	Wherever found	\$10,100	\$0	\$10,100
	<i>Birds Subtotal</i>				<i>\$8,061,066</i>	<i>\$31,608</i>	<i>\$8,092,674</i>
	Reptiles	gartersnake, narrow-headed	<i>Thamnophis rufipunctatus</i>	Wherever found	\$13,750	\$0	\$13,750
		gartersnake, northern Mexican	<i>Thamnophis eques megalops</i>	Wherever found	\$64,750	\$0	\$64,750
		Lizard, blunt-nosed leopard	<i>Gambelia silus</i>	Wherever found	\$61,000	\$0	\$61,000
		Lizard, Coachella Valley fringe-toed	<i>Uma inornata</i>	Wherever found	\$69,000	\$0	\$69,000
		Pinesnake, Louisiana	<i>Pituophis ruthveni</i>	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Rattlesnake, New Mexican ridge-nosed	Crotalus willardi obscurus	Wherever found	\$1,250	\$0	\$1,250
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$100	\$0	\$100
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$100	\$0	\$100
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$400	\$0	\$400
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$100	\$0	\$100
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$600	\$0	\$600
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$2,766,600	\$365,725	\$3,132,325
		Tortoise, gopher	Gopherus polyphemus	West of Mobile and Tombigbee Rivers	\$100	\$0	\$100
		Turtle, Alabama red-bellied	Pseudemys alabamensis	Wherever found	\$100	\$0	\$100
		Turtle, yellow-blotched map	Graptemys flavimaculata	Wherever found	\$100	\$0	\$100
		<i>Reptiles Subtotal</i>			<i>\$2,978,050</i>	<i>\$365,725</i>	<i>\$3,343,775</i>
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$164,500	\$0	\$164,500
		Frog, Chiricahua leopard	Rana chiricahuensis	Wherever found	\$60,100	\$0	\$60,100
		Frog, dusky gopher	Rana sevosa	Wherever found	\$100	\$0	\$100
		Frog, Oregon spotted	Rana pretiosa	Wherever found	\$26,800	\$0	\$26,800
		Salamander, California tiger	Ambystoma californiense	U.S.A. (Central CA DPS)	\$14,000	\$0	\$14,000
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$1,500	\$0	\$1,500
		Toad, Wyoming	Bufo hemiophrys baxteri	Wherever found	\$11,000	\$0	\$11,000
		<i>Amphibians Subtotal</i>			<i>\$278,000</i>	<i>\$0</i>	<i>\$278,000</i>
	Fishes	Bonytail	Gila elegans	Wherever found	\$45,550	\$0	\$45,550
		Chub, Borax Lake	Gila boraxobius	Wherever found	\$1,300	\$0	\$1,300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Chub, Chihuahua	<i>Gila nigrescens</i>	Wherever found	\$1,500	\$0	\$1,500
		Chub, Gila	<i>Gila intermedia</i>	Wherever found	\$30,100	\$0	\$30,100
		Chub, humpback	<i>Gila cypha</i>	Wherever found	\$53,075	\$0	\$53,075
		Chub, Owens Tui	<i>Gila bicolor</i> ssp. <i>snyderi</i>	Wherever found	\$17,000	\$0	\$17,000
		Chub, Pahrnagat roundtail	<i>Gila robusta jordani</i>	Wherever found	\$3,500	\$0	\$3,500
		Chub, Virgin River	<i>Gila seminuda</i> (=robusta)	Wherever found	\$106,300	\$0	\$106,300
		Cui-ui	<i>Chasmistes cujus</i>	Wherever found	\$11,640	\$0	\$11,640
		Dace, desert	<i>Eremichthys acros</i>	Wherever found	\$229,750	\$0	\$229,750
		Dace, Foskett speckled	<i>Rhinichthys osculus</i> ssp.	Wherever found	\$15,000	\$0	\$15,000
		Dace, Kendall Warm Springs	<i>Rhinichthys osculus thermalis</i>	Wherever found	\$1,500	\$0	\$1,500
		Dace, Moapa	<i>Moapa coriacea</i>	Wherever found	\$56,000	\$0	\$56,000
		Gambusia, Pecos	<i>Gambusia nobilis</i>	Wherever found	\$5,000	\$0	\$5,000
		Minnow, loach	<i>Tiaroga cobitis</i>	Wherever found	\$18,600	\$0	\$18,600
		Minnow, Rio Grande Silvery	<i>Hybognathus amarus</i>	Wherever found, except where listed as an experimental population	\$4,000	\$0	\$4,000
		Pikeminnow (=squawfish), Colorado	<i>Ptychocheilus lucius</i>	Wherever found, except where listed as an experimental population	\$117,276	\$0	\$117,276
		Poolfish, Pahrump	<i>Empetrichthys latos</i>	Wherever found	\$5,000	\$0	\$5,000
		Pupfish, desert	<i>Cyprinodon macularius</i>	Wherever found	\$65,737	\$0	\$65,737
		Pupfish, Owens	<i>Cyprinodon radiosus</i>	Wherever found	\$46,000	\$0	\$46,000
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	California Coastal ESU	\$27,500	\$0	\$27,500
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	Central Valley spring-run ESU	\$7,500	\$0	\$7,500
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	Central Valley spring-run ESU	\$9,000	\$0	\$9,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter-run ESU	\$10,500	\$0	\$10,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$168,000	\$0	\$168,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$87,000	\$0	\$87,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$11,000	\$0	\$11,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$3,000	\$0	\$3,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$1,422,500	\$0	\$1,422,500
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$235,600	\$0	\$235,600
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$111,400	\$0	\$111,400
		Shiner, Arkansas River	Notropis girardi	Arkansas River Basin (AR, KS, NM, OK, TX)	\$500	\$0	\$500
		Shiner, Pecos bluntnose	Notropis simus pecosensis	Wherever found	\$10,750	\$0	\$10,750
		Spinedace	Meda fulgida	Wherever found	\$18,600	\$0	\$18,600
		Spinedace, Big Spring	Lepidomeda mollispinis pratensis	Wherever found	\$6,300	\$0	\$6,300
		Spinedace, Little Colorado	Lepidomeda vittata	Wherever found	\$3,375	\$0	\$3,375
		Spinedace, White River	Lepidomeda albivallis	Wherever found	\$1,500	\$0	\$1,500
		Springfish, Hiko White River	Crenichthys baileyi grandis	Wherever found	\$22,886	\$0	\$22,886
		Springfish, Railroad Valley	Crenichthys nevadae	Wherever found	\$11,500	\$0	\$11,500
		Springfish, White River	Crenichthys baileyi baileyi	Wherever found	\$15,000	\$0	\$15,000



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	California Central Valley DPS	\$8,500	\$0	\$8,500
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Central California Coast DPS	\$1,500	\$0	\$1,500
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Lower Columbia River DPS	\$4,800	\$0	\$4,800
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Middle Columbia River DPS	\$36,960	\$0	\$36,960
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Northern California DPS	\$26,500	\$0	\$26,500
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Snake River Basin DPS	\$200,500	\$0	\$200,500
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	South-Central California Coast DPS	\$1,500	\$0	\$1,500
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Southern California DPS	\$3,000	\$0	\$3,000
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Upper Columbia River DPS	\$13,720	\$0	\$13,720
		Sturgeon, Gulf	<i>Acipenser oxyrinchus (=oxyrinchus) desotoi</i>	Wherever found	\$100	\$0	\$100
		Sturgeon, pallid	<i>Scaphirhynchus albus</i>	Wherever found	\$17,288	\$0	\$17,288
		Sturgeon, white	<i>Acipenser transmontanus</i>	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$3,000	\$0	\$3,000
		Sucker, Lost River	<i>Deltistes luxatus</i>	Wherever found	\$10,900	\$0	\$10,900
		Sucker, razorback	<i>Xyrauchen texanus</i>	Wherever found	\$131,577	\$0	\$131,577
		Sucker, shortnose	<i>Chasmistes brevirostris</i>	Wherever found	\$14,900	\$0	\$14,900
		Sucker, Warner	<i>Catostomus warnerensis</i>	Wherever found	\$146,780	\$0	\$146,780
		Topminnow, Gila (incl. Yaqui)	<i>Poeciliopsis occidentalis</i>	Wherever found	\$34,780	\$0	\$34,780

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$752,866	\$0	\$752,866
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$33,250	\$0	\$33,250
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$395,603	\$0	\$395,603
		Woundfin	Plagopterus argentissimus	Wherever found, except where listed as an experimental population	\$106,300	\$0	\$106,300
	<i>Fishes Subtotal</i>				<i>\$4,961,563</i>	<i>\$0</i>	<i>\$4,961,563</i>
	Clams	Hornshell, Texas	Popenaias popeii	Wherever found	\$23,000	\$0	\$23,000
		Mussel, scaleshell	Leptodea leptodon	Wherever found	\$100	\$0	\$100
		Pocketbook, fat	Potamilus capax	Wherever found	\$100	\$0	\$100
		Pocketbook, speckled	Lampsilis streckeri	Wherever found	\$100	\$0	\$100
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$100	\$0	\$100
	<i>Clams Subtotal</i>				<i>\$23,400</i>	<i>\$0</i>	<i>\$23,400</i>
	Snails	Ambersnail, Kanab	Oxyloma haydeni kanabensis	Wherever found	\$3,333	\$0	\$3,333
		Limpet, Banbury Springs	Lanx sp.	Wherever found	\$1,000	\$0	\$1,000
		Snail, Bliss Rapids	Taylorconcha serpenticola	Wherever found	\$3,000	\$0	\$3,000
		Snail, Morro shoulderband (=Banded dune)	Helminthoglypta walkeriana	Wherever found	\$500	\$0	\$500
		Snail, Pecos assiminea	Assiminea pecos	Wherever found	\$2,500	\$0	\$2,500
		Snail, Snake River physa	Physa natricina	Wherever found	\$3,000	\$0	\$3,000
		Springsnail, Bruneau Hot	Pyrgulopsis bruneauensis	Wherever found	\$2,500	\$0	\$2,500
		Springsnail, Koster's	Juturnia kosteri	Wherever found	\$2,500	\$0	\$2,500
		Springsnail, Roswell	Pyrgulopsis roswellensis	Wherever found	\$2,500	\$0	\$2,500
	<i>Snails Subtotal</i>				<i>\$20,833</i>	<i>\$0</i>	<i>\$20,833</i>

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$5,500	\$0	\$5,500
		Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$18,000	\$0	\$18,000
		Butterfly, Behren's silverspot	Speyeria zerene behrensii	Wherever found	\$6,500	\$0	\$6,500
		Butterfly, Quino checkerspot	Euphydryas editha quino (=E. e. wrighti)	Wherever found	\$589,000	\$0	\$589,000
		Butterfly, Uncompahgre fritillary	Boloria acrocneuma	Wherever found	\$8,000	\$0	\$8,000
		Moth, Kern primrose sphinx	Euproserpinus euterpe	Wherever found	\$10,000	\$0	\$10,000
		Skipper, Carson wandering	Pseudocopaeodes eunus obscurus	Wherever found	\$44,975	\$0	\$44,975
		Skipper, Dakota	Hesperia dacotae	Wherever found	\$19,000	\$0	\$19,000
	<i>Insects Subtotal</i>				<i>\$700,975</i>	<i>\$0</i>	<i>\$700,975</i>
	Crustaceans	Amphipod, Noel's	Gammarus desperatus	Wherever found	\$3,000	\$0	\$3,000
		Crayfish, Shasta	Pacifastacus fortis	Wherever found	\$5,000	\$0	\$5,000
		Fairy shrimp, longhorn	Branchinecta longiantenna	Wherever found	\$2,400	\$0	\$2,400
		Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$41,900	\$0	\$41,900
		Tadpole shrimp, vernal pool	Lepidurus packardii	Wherever found	\$30,500	\$0	\$30,500
	<i>Crustaceans Subtotal</i>				<i>\$82,800</i>	<i>\$0</i>	<i>\$82,800</i>
	Flowering Plants	Ambrosia, San Diego	Ambrosia pumila	Wherever found	\$151,000	\$0	\$151,000
		Baccharis, Encinitas	Baccharis vanessae	Wherever found	\$500	\$0	\$500
		Barberry, Nevin's	Berberis nevinii	Wherever found	\$6,500	\$0	\$6,500
		Bear-poppy, Dwarf	Arctomecon humilis	Wherever found	\$30,000	\$0	\$30,000
		beardtongue, Parachute	Penstemon debilis	Wherever found	\$8,800	\$0	\$8,800
		Beardtongue, Penland	Penstemon penlandii	Wherever found	\$7,000	\$0	\$7,000
		Bedstraw, El Dorado	Galium californicum ssp. sierrae	Wherever found	\$12,250	\$0	\$12,250
		Bladderpod, Dudley Bluffs	Lesquerella congesta	Wherever found	\$16,000	\$0	\$16,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Bladderpod, kodachrome	<i>Lesquerella tumulosa</i>	Wherever found	\$1,650	\$0	\$1,650
		Blue-star, Kearney's	<i>Amsonia kearneyana</i>	Wherever found	\$10,900	\$0	\$10,900
		Bluegrass, San Bernardino	<i>Poa atropurpurea</i>	Wherever found	\$500	\$0	\$500
		Brodiaea, Chinese Camp	<i>Brodiaea pallida</i>	Wherever found	\$3,500	\$2,000	\$5,500
		Buckwheat, cushenbury	<i>Eriogonum ovalifolium</i> var. <i>vineum</i>	Wherever found	\$5,000	\$0	\$5,000
		Buckwheat, Ione (incl. Irish Hill)	<i>Eriogonum apricum</i> (incl. var. <i>prostratum</i> )	Wherever found	\$2,500	\$0	\$2,500
		Buckwheat, steamboat	<i>Eriogonum ovalifolium</i> var. <i>williamsiae</i>	Wherever found	\$24,193	\$0	\$24,193
		Buttercup, autumn	<i>Ranunculus aestivalis</i> (= <i>acriiformis</i> )	Wherever found	\$1,800	\$0	\$1,800
		Butterfly plant, Colorado	<i>Gaura neomexicana</i> var. <i>coloradensis</i>	Wherever found	\$31,874	\$0	\$31,874
		Butterweed, Layne's	<i>Senecio layneae</i>	Wherever found	\$13,250	\$0	\$13,250
		Cactus, Acuna	<i>Echinomastus erectocentrus</i> var. <i>acunensis</i>	Wherever found	\$15,300	\$0	\$15,300
		Cactus, Arizona hedgehog	<i>Echinocereus triglochidiatus</i> var. <i>arizonicus</i>	Wherever found	\$500	\$0	\$500
		Cactus, Bakersfield	<i>Opuntia treleasei</i>	Wherever found	\$1,500	\$0	\$1,500
		Cactus, Brady pincushion	<i>Pediocactus bradyi</i>	Wherever found	\$1,000	\$0	\$1,000
		Cactus, Colorado hookless	<i>Sclerocactus glaucus</i>	Wherever found	\$131,633	\$0	\$131,633
		Cactus, Fickeisen plains	<i>Pediocactus peeblesianus</i> <i>fickeiseniae</i>	Wherever found	\$18,000	\$0	\$18,000
		Cactus, Knowlton's	<i>Pediocactus knowltonii</i>	Wherever found	\$17,000	\$0	\$17,000
		Cactus, Kuenzler hedgehog	<i>Echinocereus fendleri</i> var. <i>kuenzleri</i>	Wherever found	\$83,500	\$0	\$83,500
		Cactus, Lee pincushion	<i>Coryphantha sneedii</i> var. <i>leei</i>	Wherever found	\$3,000	\$0	\$3,000
		Cactus, Mesa Verde	<i>Sclerocactus mesae-verdae</i>	Wherever found	\$17,000	\$0	\$17,000
		Cactus, Nichol's Turk's head	<i>Echinocactus horizontalionius</i> var. <i>nicholii</i>	Wherever found	\$2,100	\$0	\$2,100
		cactus, Pariette	<i>Sclerocactus brevispinus</i>	Wherever found	\$13,579	\$0	\$13,579

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Cactus, Peebles Navajo	<i>Pediocactus peeblesianus</i> var. <i>peeblesianus</i>	Wherever found	\$2,350	\$0	\$2,350
		Cactus, San Rafael	<i>Pediocactus despainii</i>	Wherever found	\$8,250	\$0	\$8,250
		Cactus, Siler pincushion	<i>Pediocactus</i> (=Echinocactus, =Utahia) <i>sileri</i>	Wherever found	\$20,700	\$0	\$20,700
		Cactus, Sneed pincushion	<i>Coryphantha sneedii</i> var. <i>sneedii</i>	Wherever found	\$1,500	\$0	\$1,500
		cactus, Uinta Basin hookless	<i>Sclerocactus wetlandicus</i>	Wherever found	\$15,992	\$0	\$15,992
		Cactus, Winkler	<i>Pediocactus winkleri</i>	Wherever found	\$6,000	\$0	\$6,000
		Cactus, Wright fishhook	<i>Sclerocactus wrightiae</i>	Wherever found	\$74,250	\$0	\$74,250
		Catchfly, Spalding's	<i>Silene spaldingii</i>	Wherever found	\$26,425	\$0	\$26,425
		Ceanothus, Pine Hill	<i>Ceanothus roderickii</i>	Wherever found	\$9,750	\$0	\$9,750
		Centaury, spring-loving	<i>Centaureum namophilum</i>	Wherever found	\$6,000	\$0	\$6,000
		Checker-mallow, Keck's	<i>Sidalcea keckii</i>	Wherever found	\$1,000	\$0	\$1,000
		Checker-mallow, Nelson's	<i>Sidalcea nelsoniana</i>	Wherever found	\$2,500	\$0	\$2,500
		Clarkia, Springville	<i>Clarkia springvillensis</i>	Wherever found	\$3,500	\$0	\$3,500
		Cliffrose, Arizona	<i>Purshia</i> (=Cowania) <i>subintegra</i>	Wherever found	\$13,000	\$0	\$13,000
		Cycladenia, Jones	<i>Cycladenia humilis</i> var. <i>jonesii</i>	Wherever found	\$33,030	\$0	\$33,030
		Daisy, Parish's	<i>Erigeron parishii</i>	Wherever found	\$10,500	\$0	\$10,500
		Daisy, Willamette	<i>Erigeron decumbens</i>	Wherever found	\$75,000	\$0	\$75,000
		Evening-primrose, San Benito	<i>Camissonia benitensis</i>	Wherever found	\$31,000	\$0	\$31,000
		Flannelbush, Mexican	<i>Fremontodendron mexicanum</i>	Wherever found	\$500	\$0	\$500
		Flannelbush, Pine Hill	<i>Fremontodendron californicum</i> ssp. <i>decumbens</i>	Wherever found	\$15,250	\$0	\$15,250
		Fleabane, Zuni	<i>Erigeron rhizomatus</i>	Wherever found	\$3,000	\$0	\$3,000
		Four-o'clock, MacFarlane's	<i>Mirabilis macfarlanei</i>	Wherever found	\$9,500	\$0	\$9,500
		Fritillary, Gentner's	<i>Fritillaria gentneri</i>	Wherever found	\$212,000	\$0	\$212,000
		Goldfields, Contra Costa	<i>Lasthenia conjugens</i>	Wherever found	\$1,000	\$0	\$1,000
		Gumplant, Ash Meadows	<i>Grindelia fraxinipratensis</i>	Wherever found	\$6,000	\$0	\$6,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Howellia, Water	Howellia aquatilis		\$4,300	\$0	\$4,300
		ivesia, Webber's	Ivesia webberi	Wherever found	\$75,404	\$0	\$75,404
		Jewelflower, California	Caulanthus californicus	Wherever found	\$13,500	\$0	\$13,500
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$212,628	\$0	\$212,628
		Layia, beach	Layia carnosa	Wherever found	\$131,000	\$0	\$131,000
		Lily, Western	Lilium occidentale	Wherever found	\$26,000	\$0	\$26,000
		lomatium, Bradshaw's	Lomatium bradshawii	Wherever found	\$59,000	\$0	\$59,000
		Lomatium, Cook's	Lomatium cookii	Wherever found	\$64,000	\$0	\$64,000
		Lupine, Kincaid's	Lupinus sulphureus ssp. kincaidii	Wherever found	\$113,000	\$0	\$113,000
		mallow, Gierisch	Sphaeralcea gierischii	Wherever found	\$19,700	\$0	\$19,700
		Mallow, Kern	Eremalche kernensis	Wherever found	\$9,000	\$0	\$9,000
		Manzanita, Ione	Arctostaphylos myrtifolia	Wherever found	\$4,700	\$0	\$4,700
		Manzanita, Morro	Arctostaphylos morroensis	Wherever found	\$500	\$0	\$500
		Milk-vetch, Coachella Valley	Astragalus lentiginosus var. coachellae	Wherever found	\$1,000	\$0	\$1,000
		Milk-vetch, Cushenbury	Astragalus albens	Wherever found	\$8,500	\$0	\$8,500
		Milk-vetch, Fish Slough	Astragalus lentiginosus var. piscinensis	Wherever found	\$16,000	\$0	\$16,000
		Milk-vetch, Holmgren	Astragalus holmgreniorum	Wherever found	\$28,100	\$0	\$28,100
		Milk-vetch, Lane Mountain	Astragalus jaegerianus	Wherever found	\$37,000	\$0	\$37,000
		Milk-vetch, Mancos	Astragalus humillimus	Wherever found	\$4,800	\$0	\$4,800
		Milk-vetch, Peirson's	Astragalus magdalenae var. peirsonii	Wherever found	\$261,000	\$0	\$261,000
		Milk-vetch, Shivwits	Astragalus ampullarioides	Wherever found	\$19,000	\$0	\$19,000
		Milk-vetch, triple-ribbed	Astragalus tricarinatus	Wherever found	\$56,000	\$0	\$56,000
		milkvetch, Osterhout	Astragalus osterhoutii	Wherever found	\$9,000	\$0	\$9,000
		Milkweed, Welsh's	Asclepias welshii	Wherever found	\$30,200	\$0	\$30,200
		Morning-glory, Stebbins'	Calystegia stebbinsii	Wherever found	\$18,700	\$0	\$18,700
		Mustard, Penland alpine fen	Eutrema penlandii	Wherever found	\$9,000	\$0	\$9,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Navarretia, few-flowered	Navarretia leucocephala ssp. pauciflora (=N. pauciflora)	Wherever found	\$1,000	\$0	\$1,000
		Navarretia, many-flowered	Navarretia leucocephala ssp. plieantha	Wherever found	\$6,000	\$0	\$6,000
		Niterwort, Amargosa	Nitrophila mohavensis	Wherever found	\$36,000	\$0	\$36,000
		Orchid, western prairie fringed	Platanthera praeclara	Wherever found	\$11,189	\$0	\$11,189
		Orcutt grass, California	Orcuttia californica	Wherever found	\$500	\$0	\$500
		Orcutt grass, San Joaquin	Orcuttia inaequalis	Wherever found	\$300	\$0	\$300
		Orcutt grass, slender	Orcuttia tenuis	Wherever found	\$4,000	\$0	\$4,000
		Owl's-clover, fleshy	Castilleja campestris ssp. succulenta	Wherever found	\$1,500	\$0	\$1,500
		oxytheca, Cushenbury	Oxytheca parishii var. goodmaniana	Wherever found	\$61,500	\$0	\$61,500
		Pawpaw, four-petal	Asimina tetramera	Wherever found	\$6,500	\$0	\$6,500
		Pennyroyal, Todsens	Hedeoma todsenii	Wherever found	\$31,500	\$0	\$31,500
		Penstemon, blowout	Penstemon haydenii	Wherever found	\$28,849	\$0	\$28,849
		Peppergrass, Slickspot	Lepidium papilliferum		\$269,400	\$0	\$269,400
		Phacelia, clay	Phacelia argillacea	Wherever found	\$300	\$0	\$300
		Phacelia, DeBeque	Phacelia submutica	Wherever found	\$15,200	\$0	\$15,200
		Phacelia, North Park	Phacelia formosula	Wherever found	\$8,000	\$0	\$8,000
		popcornflower, rough	Plagiobothrys hirtus	Wherever found	\$200	\$0	\$200
		Reed-mustard, Barneby	Schoenocrambe barnebyi	Wherever found	\$1,050	\$0	\$1,050
		Reed-mustard, clay	Schoenocrambe argillacea	Wherever found	\$3,369	\$0	\$3,369
		Reed-mustard, shrubby	Schoenocrambe suffrutescens	Wherever found	\$4,712	\$0	\$4,712
		Ridge-cress, Barneby	Lepidium barnebyanum	Wherever found	\$100	\$0	\$100
		Rock-cress, McDonald's	Arabis macdonaldiana	Wherever found	\$1,000	\$0	\$1,000
		Seagrass, Johnson's	Halophila johnsonii	Wherever found	\$1,000	\$0	\$1,000
		Sedge, Navajo	Carex specuicola	Wherever found	\$2,010	\$0	\$2,010
		Skyrocket, Pagosa	Ipomopsis polyantha	Wherever found	\$25,000	\$0	\$25,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sunflower, Pecos (=puzzle, =paradox)	Helianthus paradoxus	Wherever found	\$22,000	\$0	\$22,000
		Townsendia, Last Chance	Townsendia aprica	Wherever found	\$4,050	\$0	\$4,050
		Twinpod, Dudley Bluffs	Physaria obcordata	Wherever found	\$10,500	\$0	\$10,500
		Vervain, Red Hills	Verbena californica	Wherever found	\$16,400	\$0	\$16,400
		Wallflower, Menzies'	Erysimum menziesii	Wherever found	\$131,000	\$0	\$131,000
		Water-umbel, Huachuca	Lilaeopsis schaffneriana var. recurva	Wherever found	\$12,300	\$0	\$12,300
		wild buckwheat, clay-loving	Eriogonum pelinophilum	Wherever found	\$19,150	\$0	\$19,150
		Wild-buckwheat, gypsum	Eriogonum gypsophilum	Wherever found	\$52,000	\$7,000	\$59,000
		Wire-lettuce, Malheur	Stephanomeria malheurensis	Wherever found	\$2,000	\$0	\$2,000
		Woolly-star, Santa Ana River	Eriastrum densifolium ssp. sanctorum	Wherever found	\$6,000	\$0	\$6,000
		Wooly-threads, San Joaquin	Monolopia (=Lembertia) congdonii	Wherever found	\$64,000	\$0	\$64,000
		Yellowhead, desert	Yermo xanthocephalus	Wherever found	\$50,000	\$0	\$50,000
	<i>Flowering Plants Subtotal</i>				\$3,359,937	\$9,000	\$3,368,937
	Ferns and Allies	Quillwort, Louisiana	Isoetes louisianensis	Wherever found	\$100	\$0	\$100
	<i>Ferns and Allies Subtotal</i>				\$100	\$0	\$100
	Lichens	Cladonia, Florida perforate	Cladonia perforata	Wherever found	\$6,500	\$0	\$6,500
	<i>Lichens Subtotal</i>				\$6,500	\$0	\$6,500
	<i>Domestic Subtotal</i>				\$23,226,857	\$433,333	\$23,660,190
	<b>Total</b>				\$23,226,857	\$433,333	\$23,660,190



## DOI Bureau of Reclamation - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Northern long-eared	<i>Myotis septentrionalis</i>	Wherever found	\$6,500	\$0	\$6,500
		Fox, San Joaquin kit	<i>Vulpes macrotis mutica</i>	wherever found	\$91,500	\$207,500	\$299,000
		Kangaroo rat, Fresno	<i>Dipodomys nitratoides exilis</i>	Wherever found	\$6,500	\$0	\$6,500
		Kangaroo rat, giant	<i>Dipodomys ingens</i>	Wherever found	\$1,800	\$75,000	\$76,800
		Kangaroo rat, Tipton	<i>Dipodomys nitratoides nitratoides</i>	Wherever found	\$46,500	\$132,500	\$179,000
		Mouse, New Mexico meadow jumping	<i>Zapus hudsonius luteus</i>	Wherever found	\$100,000	\$0	\$100,000
		Mouse, Preble's meadow jumping	<i>Zapus hudsonius preblei</i>	wherever found	\$1,000	\$0	\$1,000
		Mouse, salt marsh harvest	<i>Reithrodontomys raviventris</i>	wherever found	\$2,500	\$0	\$2,500
		Rabbit, riparian brush	<i>Sylvilagus bachmani riparius</i>	Wherever found	\$7,500	\$0	\$7,500
		Sheep, Peninsular bighorn	<i>Ovis canadensis nelsoni</i>	Peninsular CA pop.	\$49,000	\$0	\$49,000
		Shrew, Buena Vista Lake ornate	<i>Sorex ornatus relictus</i>	Wherever found	\$3,000	\$0	\$3,000
		Whale, killer	<i>Orcinus orca</i>	Southern Resident DPS	\$28,000	\$0	\$28,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$200	\$0	\$200
		Woodrat, riparian (=San Joaquin Valley)	Neotoma fuscipes riparia	Wherever found	\$7,500	\$0	\$7,500
	<i>Mammals Subtotal</i>				<i>\$351,500</i>	<i>\$415,000</i>	<i>\$766,500</i>
	Birds	Condor, California	Gymnogyps californianus	U.S.A. (specific portions of Arizona, Nevada, and Utah)	\$900	\$728,000	\$728,900
		Crane, whooping	Grus americana	U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	\$600	\$0	\$600
		Cuckoo, yellow-billed	Coccyzus americanus	Western U.S. DPS	\$3,930,500	\$300,000	\$4,230,500
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$4,741,500	\$0	\$4,741,500
		Knot, red	Calidris canutus rufa	Wherever found	\$200	\$0	\$200
		Owl, northern spotted	Strix occidentalis caurina	Wherever found	\$3,000	\$0	\$3,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$8,000	\$0	\$8,000
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$1,800	\$0	\$1,800
		Rail, Yuma Ridgways	Rallus obsoletus [=longirostris] yumanensis	Wherever found	\$1,594,000	\$0	\$1,594,000
		Tern, California least	Sterna antillarum browni	Wherever found	\$1,300	\$0	\$1,300
		Tern, least	Sterna antillarum	interior pop.	\$28,000	\$0	\$28,000
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$3,500	\$0	\$3,500
		<i>Birds Subtotal</i>			<i>\$10,313,300</i>	<i>\$1,028,000</i>	<i>\$11,341,300</i>
	Reptiles	Lizard, blunt-nosed leopard	Gambelia silus	Wherever found	\$64,000	\$207,500	\$271,500
		Sea turtle, green	Chelonia mydas	East Pacific DPS	\$1,800	\$0	\$1,800
		Snake, giant garter	Thamnophis gigas	Wherever found	\$98,000	\$0	\$98,000
		Snake, San Francisco garter	Thamnophis sirtalis tetrataenia	Wherever found	\$900	\$0	\$900
		Whipsnake (=striped racer), Alameda	Masticophis lateralis euryxanthus	Wherever found	\$1,100	\$0	\$1,100
		<i>Reptiles Subtotal</i>			<i>\$165,800</i>	<i>\$207,500</i>	<i>\$373,300</i>
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$51,500	\$0	\$51,500
		Frog, Oregon spotted	Rana pretiosa	Wherever found	\$119,000	\$0	\$119,000
		Frog, Sierra Nevada Yellow-legged	Rana sierrae	Wherever found	\$800	\$0	\$800

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salamander, California tiger	Ambystoma californiense	U.S.A. (CA - Santa Barbara County)	\$300	\$0	\$300
		Salamander, California tiger	Ambystoma californiense	U.S.A. (CA - Sonoma County)	\$5,500	\$0	\$5,500
		Salamander, California tiger	Ambystoma californiense	U.S.A. (Central CA DPS)	\$105,500	\$728,000	\$833,500
	<i>Amphibians Subtotal</i>				<i>\$282,600</i>	<i>\$728,000</i>	<i>\$1,010,600</i>
	Fishes	Bonytail	Gila elegans	Wherever found	\$8,582,500	\$0	\$8,582,500
		Chub, Gila	Gila intermedia	Wherever found	\$559,000	\$0	\$559,000
		Chub, humpback	Gila cypha	Wherever found	\$6,500,500	\$0	\$6,500,500
		Cui-ui	Chasmistes cujus	Wherever found	\$50,000	\$0	\$50,000
		Eulachon	Thaleichthys pacificus	Southern DPS	\$8,000	\$0	\$8,000
		Gambusia, Pecos	Gambusia nobilis	Wherever found	\$5,000	\$30,000	\$35,000
		Minnow, loach	Tiaroga cobitis	Wherever found	\$559,000	\$0	\$559,000
		Minnow, Rio Grande Silvery	Hybognathus amarus	Wherever found, except where listed as an experimental population	\$6,000,000	\$1,200,000	\$7,200,000
		Pikeminnow (=squawfish), Colorado	Ptychocheilus lucius	Wherever found, except where listed as an experimental population	\$3,500,500	\$0	\$3,500,500
		Pupfish, Comanche Springs	Cyprinodon elegans	Wherever found	\$5,000	\$30,000	\$35,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$13,372,500	\$217,500	\$13,590,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$410,000	\$582,500	\$992,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter-run ESU	\$16,121,000	\$66,000	\$16,187,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$337,000	\$582,500	\$919,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$2,874,500	\$582,500	\$3,457,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$3,750,000	\$582,500	\$4,332,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$17,500	\$0	\$17,500
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$5,690,000	\$0	\$5,690,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$75,000	\$0	\$75,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$5,000	\$0	\$5,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$4,660,500	\$0	\$4,660,500
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$337,000	\$582,500	\$919,500
		Shiner, Pecos bluntnose	Notropis simus pecosensis	Wherever found	\$2,000,000	\$200,000	\$2,200,000
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$900	\$0	\$900
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$19,459,000	\$0	\$19,459,000
		Spikedace	Meda fulgida	Wherever found	\$559,000	\$0	\$559,000
		Spinedace, Little Colorado	Lepidomeda vittata	Wherever found	\$25,000	\$0	\$25,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$13,479,500	\$217,500	\$13,697,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$470,500	\$582,500	\$1,053,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$1,461,000	\$582,500	\$2,043,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Northern California DPS	\$3,568,500	\$0	\$3,568,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$3,885,500	\$582,500	\$4,468,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$500	\$0	\$500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$1,495,500	\$0	\$1,495,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$3,244,500	\$582,500	\$3,827,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$20,500	\$0	\$20,500
		sturgeon, green	Acipenser medirostris	Southern DPS	\$6,407,000	\$0	\$6,407,000
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$587,500	\$0	\$587,500
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$5,000	\$0	\$5,000
		Sucker, June	Chasmistes liorus	Wherever found	\$50,000	\$0	\$50,000
		Sucker, Lost River	Deltistes luxatus	Wherever found	\$1,768,000	\$0	\$1,768,000
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$11,768,000	\$0	\$11,768,000
		Sucker, shortnose	Chasmistes brevirostris	Wherever found	\$1,768,000	\$0	\$1,768,000
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	Wherever found	\$559,000	\$0	\$559,000
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$1,692,500	\$0	\$1,692,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$50,500	\$0	\$50,500
	<i>Fishes Subtotal</i>				<i>\$147,744,900</i>	<i>\$7,203,500</i>	<i>\$154,948,400</i>
	Snails	Limpet, Banbury Springs	Lanx sp.	Wherever found	\$8,000	\$0	\$8,000
		Snail, Bliss Rapids	Taylorconcha serpenticola	Wherever found	\$8,000	\$0	\$8,000
		Snail, Snake River physa	Physa natricina	Wherever found	\$61,000	\$0	\$61,000
	<i>Snails Subtotal</i>				<i>\$77,000</i>	<i>\$0</i>	<i>\$77,000</i>
	Insects	Beetle, delta green ground	Elaphrus viridis	Wherever found	\$2,500	\$0	\$2,500
		Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$140,000	\$0	\$140,000
		Butterfly, bay checkerspot	Euphydryas editha bayensis	Wherever found	\$203,500	\$0	\$203,500
		Butterfly, callippe silverspot	Speyeria callippe callippe	Wherever found	\$600	\$0	\$600
		Butterfly, Fender's blue	Icaricia icarioides fenderi	Wherever found	\$54,000	\$0	\$54,000
		Butterfly, Lange's metalmark	Apodemia mormo langei	Wherever found	\$900	\$0	\$900
		Butterfly, San Bruno elfin	Callophrys mossii bayensis	Wherever found	\$900	\$0	\$900
		Moth, Kern primrose sphinx	Euproserpinus euterpe	Wherever found	\$0	\$75,000	\$75,000
		Skipper, Dakota	Hesperia dacotae	Wherever found	\$700	\$0	\$700
		skipperling, Poweshiek	Oarisma poweshiek	Wherever found	\$200	\$0	\$200
	<i>Insects Subtotal</i>				<i>\$403,300</i>	<i>\$75,000</i>	<i>\$478,300</i>
	Crustaceans	Fairy shrimp, Conservancy	Branchinecta conservatio	Wherever found	\$86,000	\$0	\$86,000
		Fairy shrimp, longhorn	Branchinecta longiantenna	Wherever found	\$1,600	\$0	\$1,600
		Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$126,000	\$0	\$126,000
		Tadpole shrimp, vernal pool	Lepidurus packardi	Wherever found	\$89,500	\$0	\$89,500
	<i>Crustaceans Subtotal</i>				<i>\$303,100</i>	<i>\$0</i>	<i>\$303,100</i>
	Flowering Plants	Bird's beak, palmate-bracted	Cordylanthus palmatus	Wherever found	\$1,500	\$0	\$1,500
		Cactus, Bakersfield	Opuntia treleasei	Wherever found	\$600	\$0	\$600
		Cactus, Mesa Verde	Sclerocactus mesae-verdae	Wherever found	\$5,000	\$0	\$5,000
		Checker-mallow, Keck's	Sidalcea keckii	Wherever found	\$600	\$0	\$600
		Clarkia, Springville	Clarkia springvillensis	Wherever found	\$1,300	\$0	\$1,300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Dudleya, Santa Clara Valley	Dudleya setchellii	Wherever found	\$1,400	\$0	\$1,400
		Evening-primrose, Antioch Dunes	Oenothera deltooides ssp. howellii	Wherever found	\$1,500	\$0	\$1,500
		Fiddleneck, large-flowered	Amsinckia grandiflora	Wherever found	\$187,000	\$0	\$187,000
		Goldfields, Contra Costa	Lasthenia conjugens	Wherever found	\$900	\$0	\$900
		Jewelflower, California	Caulanthus californicus	Wherever found	\$80,500	\$0	\$80,500
		Jewelflower, Metcalf Canyon	Streptanthus albidus ssp. albidus	Wherever found	\$600	\$0	\$600
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$50,000	\$0	\$50,000
		Lupine, Kincaid's	Lupinus sulphureus ssp. kincaidii	Wherever found	\$49,000	\$0	\$49,000
		Mallow, Kern	Eremalche kernensis	Wherever found	\$7,000	\$75,000	\$82,000
		Manzanita, pallid	Arctostaphylos pallida	Wherever found	\$900	\$0	\$900
		Orchid, western prairie fringed	Platanthera praeclara	Wherever found	\$200	\$0	\$200
		Orcutt grass, hairy	Orcuttia pilosa	Wherever found	\$1,500	\$0	\$1,500
		Orcutt grass, San Joaquin	Orcuttia inaequalis	Wherever found	\$2,500	\$0	\$2,500
		Paintbrush, Tiburon	Castilleja affinis ssp. neglecta	Wherever found	\$100	\$0	\$100
		Pussypaws, Mariposa	Calyptridium pulchellum	Wherever found	\$1,300	\$0	\$1,300
		Spurge, Hoover's	Chamaesyce hooveri	Wherever found	\$3,000	\$0	\$3,000
		Sunburst, Hartweg's golden	Pseudobahia bahiifolia	Wherever found	\$1,300	\$0	\$1,300
		Sunburst, San Joaquin adobe	Pseudobahia peirsonii	Wherever found	\$1,300	\$0	\$1,300
		Tarplant, Santa Cruz	Holocarpha macradenia	Wherever found	\$1,400	\$0	\$1,400
		Tuctoria, Greene's	Tuctoria greenei	Wherever found	\$2,500	\$0	\$2,500
		Wallflower, Contra Costa	Erysimum capitatum var. angustatum	Wherever found	\$600	\$0	\$600
		Woolly-threads, San Joaquin	Monolopia (=Lembertia) congdonii	Wherever found	\$4,000	\$0	\$4,000
	<i>Flowering Plants Subtotal</i>				\$407,500	\$75,000	\$482,500
	Multi-Species				\$65,007,500	\$2,602,500	\$67,610,000



<b>Locality</b>	<b>Group Name</b>	<b>Common Name</b>	<b>Scientific Name</b>	<b>Population</b>	<b>General Expenditures</b>	<b>Land Expenditures</b>	<b>Grand Total</b>
	<i>Multi-Species Subtotal</i>				\$65,007,500	\$2,602,500	\$67,610,000
<i>Domestic Total</i>					\$225,056,500	\$12,334,500	\$237,391,000
<b>Total</b>					<b>\$225,056,500</b>	<b>\$12,334,500</b>	<b>\$237,391,000</b>

## DOI National Parks Service - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
Domestic	Mammals	bat, Florida bonneted	Eumops floridanus	Wherever found	\$33,000	\$0	\$33,000
		Bat, gray	Myotis grisescens	Wherever found	\$88,000	\$0	\$88,000
		Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$15,000	\$0	\$15,000
		Bat, Indiana	Myotis sodalis	Wherever found	\$291,514	\$0	\$291,514
		Bat, Mexican long-nosed	Leptonycteris nivalis	Wherever found	\$2,300	\$0	\$2,300
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$762,211	\$0	\$762,211
		Bat, Ozark big-eared	Corynorhinus (=Plecotus) townsendii ingens	Wherever found	\$5,000	\$0	\$5,000
		Bat, Virginia big-eared	Corynorhinus (=Plecotus) townsendii virginianus	Wherever found	\$11,000	\$0	\$11,000
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$2,063,708	\$0	\$2,063,708
		Ferret, black-footed	Mustela nigripes	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$101,000	\$0	\$101,000
		Ferret, black-footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$40,000	\$0	\$40,000
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$24,600	\$0	\$24,600
		Manatee, West Indian	Trichechus manatus	Wherever found	\$19,000	\$0	\$19,000
		Mouse, Anastasia Island beach	Peromyscus polionotus phasma	Wherever found	\$800	\$0	\$800

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Mouse, New Mexico meadow jumping	<i>Zapus hudsonius luteus</i>	Wherever found	\$10,000	\$0	\$10,000
		Mouse, Perdido Key beach	<i>Peromyscus polionotus trissyllepsis</i>	Wherever found	\$2,000	\$0	\$2,000
		Mouse, southeastern beach	<i>Peromyscus polionotus</i>	wherever found	\$2,000	\$0	\$2,000
		Panther, Florida	<i>Puma (=Felis) concolor coryi</i>	Wherever found	\$50,000	\$0	\$50,000
		Prairie dog, Utah	<i>Cynomys parvidens</i>	Wherever found	\$139,200	\$0	\$139,200
		Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	U.S.A. (AZ), Mexico	\$3,680	\$0	\$3,680
		Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	Wherever found, except where listed as an experimental population	\$3,680	\$0	\$3,680
		Seal, Hawaiian monk	<i>Monachus schauinslandi</i>	Wherever found	\$6,500	\$0	\$6,500
		Sheep, Sierra Nevada bighorn	<i>Ovis canadensis sierrae</i>	Sierra Nevada	\$185,723	\$0	\$185,723
		Squirrel, Carolina northern flying	<i>Glaucomys sabrinus coloratus</i>	Wherever found	\$7,700	\$0	\$7,700
		Whale, finback	<i>Balaenoptera physalus</i>	Wherever found	\$100	\$0	\$100
		Whale, humpback	<i>Megaptera novaeangliae</i>	Mexico DPS	\$318,804	\$0	\$318,804
		Whale, North Atlantic Right	<i>Eubalaena glacialis</i>	Wherever found	\$600	\$0	\$600
		Whale, Sei	<i>Balaenoptera borealis</i>	Wherever found	\$50	\$0	\$50
		Whale, sperm	<i>Physeter catodon</i> (=macrocephalus)	Wherever found	\$50	\$0	\$50
		Wolf, gray	<i>Canis lupus</i>	MN	\$40,000	\$0	\$40,000

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$544,101	\$0	\$544,101
		Wolf, Mexican	Canis lupus baileyi	U.S.A. (portions of AZ and NM)see 17.84(k)	\$100	\$0	\$100
	<i>Mammals Subtotal</i>				<i>\$4,771,421</i>	<i>\$0</i>	<i>\$4,771,421</i>
	Birds	akepa, Maui	Loxops ochraceus	Wherever found	\$105,000	\$0	\$105,000
		Condor, California	Gymnogyps californianus	U.S.A. (specific portions of Arizona, Nevada, and Utah)	\$17,000	\$0	\$17,000
		Condor, California	Gymnogyps californianus	U.S.A. only, except where listed as an experimental population	\$400,000	\$0	\$400,000
		Coot, Hawaiian	Fulica americana alai	Wherever found	\$71,000	\$0	\$71,000
		Cuckoo, yellow-billed	Coccyzus americanus	Western U.S. DPS	\$14,225	\$0	\$14,225

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Falcon, Northern Aplomado	Falco femoralis septentrionalis	Wherever found, except where listed as an experimental population	\$14,326	\$0	\$14,326
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$7,500	\$0	\$7,500
		Goose, Hawaiian	Branta (=Nesochen) sandvicensis	Wherever found	\$515,000	\$0	\$515,000
		honeycreeper, (Akohekohe) crested	Palmeria dolei	Wherever found	\$105,000	\$0	\$105,000
		Kite, Everglade snail	Rostrhamus sociabilis plumbeus	Wherever found	\$5,000	\$0	\$5,000
		Knot, red	Calidris canutus rufa	Wherever found	\$12,405	\$0	\$12,405
		Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR, WA)	\$169,700	\$0	\$169,700
		Owl, Mexican spotted	Strix occidentalis lucida	Wherever found	\$157,334	\$0	\$157,334
		Owl, northern spotted	Strix occidentalis caurina	Wherever found	\$409,906	\$0	\$409,906
		parrotbill, Maui (Kiwikiu)	Pseudonestor xanthophrys	Wherever found	\$105,000	\$0	\$105,000
		petrel, Hawaiian	Pterodroma sandwichensis	Wherever found	\$489,000	\$0	\$489,000
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$1,004,207	\$0	\$1,004,207

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Plover, piping	Charadrius melodus	[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$141,102	\$0	\$141,102
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS- U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$70,000	\$0	\$70,000
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$22,000	\$0	\$22,000
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$1,000	\$0	\$1,000
		Shearwater, Newell's Townsend's	Puffinus auricularis newelli	Wherever found	\$5,000	\$0	\$5,000
		Sparrow, Cape Sable seaside	Ammodramus maritimus mirabilis	Wherever found	\$400,000	\$0	\$400,000
		Stilt, Hawaiian	Himantopus mexicanus knudseni	Wherever found	\$71,000	\$0	\$71,000
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$51,100	\$0	\$51,100
		Storm-petrel, band-rumped	Oceanodroma castro	USA (HI)	\$5,000	\$0	\$5,000
		Swiftlet, Mariana gray	Aerodramus vanikorensis bartschi	Wherever found	\$200	\$0	\$200
		Tern, least	Sterna antillarum	interior pop.	\$37,300	\$0	\$37,300
		Tern, roseate	Sterna dougallii dougallii	Northeast U.S. nesting population	\$11,600	\$0	\$11,600
		Tern, roseate	Sterna dougallii dougallii	Western Hemisphere except NE U.S.	\$2,500	\$0	\$2,500

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Woodpecker, ivory-billed	Campephilus principalis	Wherever found	\$5,000	\$0	\$5,000
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$11,000	\$0	\$11,000
		'Iwi	Drepanis coccinea	Wherever found	\$105,000	\$0	\$105,000
	<i>Birds Subtotal</i>				<i>\$4,540,405</i>	<i>\$0</i>	<i>\$4,540,405</i>
	Reptiles	Alligator, American	Alligator mississippiensis	Wherever found	\$95,000	\$0	\$95,000
		Crocodile, American	Crocodylus acutus	U.S.A. (FL)	\$142,000	\$0	\$142,000
		gartersnake, northern Mexican	Thamnophis eques megalops	Wherever found	\$4,000	\$0	\$4,000
		Lizard, St. Croix ground	Ameiva polops	Wherever found	\$2,500	\$0	\$2,500
		Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$2,228	\$0	\$2,228
		Sea turtle, green	Chelonia mydas	Central North Pacific DPS	\$2,700	\$0	\$2,700
		Sea turtle, green	Chelonia mydas	Central South Pacific DPS	\$250	\$0	\$250
		Sea turtle, green	Chelonia mydas	Central West Pacific DPS	\$200	\$0	\$200
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$431,239	\$0	\$431,239
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$133,902	\$0	\$133,902
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$1,059,819	\$0	\$1,059,819
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$4,500	\$0	\$4,500
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$576,664	\$0	\$576,664
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$5,000	\$0	\$5,000
		Snake, San Francisco garter	Thamnophis sirtalis tetrataenia	Wherever found	\$7,500	\$0	\$7,500
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$279,352	\$0	\$279,352

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Tortoise, gopher	Gopherus polyphemus	West of Mobile and Tombigbee Rivers	\$100	\$0	\$100
		turtle, bog	Clemmys muhlenbergii	U.S.A. (GA, NC, SC, TN, VA)	\$4,000	\$0	\$4,000
		Turtle, Sonoyta mud	Kinosternon sonoriense longifemorale	Wherever found	\$756	\$0	\$756
		Whipsnake (=striped racer), Alameda	Masticophis lateralis euryxanthus	Wherever found	\$75,000	\$0	\$75,000
	<i>Reptiles Subtotal</i>				\$2,826,710	\$0	\$2,826,710
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$101,000	\$0	\$101,000
		Frog, mountain yellow-legged	Rana muscosa	Northern California DPS	\$310,700	\$0	\$310,700
		Frog, Sierra Nevada Yellow-legged	Rana sierrae	Wherever found	\$505,100	\$0	\$505,100
		Hellbender, Ozark	Cryptobranchus alleganiensis bishopi	Wherever found	\$7,887	\$0	\$7,887
		Salamander, California tiger	Ambystoma californiense	U.S.A. (Central CA DPS)	\$2,500	\$0	\$2,500
		Salamander, Jemez Mountains	Plethodon neomexicanus	Wherever found	\$50,000	\$0	\$50,000
		Salamander, Shenandoah	Plethodon shenandoah	Wherever found	\$15,000	\$0	\$15,000
	<i>Amphibians Subtotal</i>				\$992,187	\$0	\$992,187
	Fishes	Bonytail	Gila elegans	Wherever found	\$500	\$0	\$500
		Chub, humpback	Gila cypha	Wherever found	\$314,760	\$0	\$314,760
		Dace, blackside	Phoxinus cumberlandensis	Wherever found	\$1,600	\$0	\$1,600
		Darter, duskytail	Etheostoma percnurum	Wherever found	\$7,947	\$0	\$7,947
		Gambusia, Big Bend	Gambusia gaigei	Wherever found	\$1,000	\$0	\$1,000
		Goby, tidewater	Eucyclogobius newberryi	Wherever found	\$10,000	\$0	\$10,000
		Madtom, smoky	Noturus baileyi	Wherever found	\$7,947	\$0	\$7,947



Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Madtom, yellowfin	Noturus flavipinnis	Wherever found, except where listed as an experimental population	\$7,947	\$0	\$7,947
		Minnow, Rio Grande Silvery	Hybognathus amarus	Rio Grande, from Little Box Canyon to Amistad Dam	\$300	\$0	\$300
		Pikeminnow (=squawfish), Colorado	Ptychocheilus lucius	Wherever found, except where listed as an experimental population	\$750	\$0	\$750
		Pupfish, desert	Cyprinodon macularius	Wherever found	\$3,000	\$0	\$3,000
		Pupfish, Devils Hole	Cyprinodon diabolis	Wherever found	\$400,000	\$0	\$400,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$54,000	\$0	\$54,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$1,500	\$0	\$1,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$3,850	\$0	\$3,850
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$352,000	\$0	\$352,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$3,850	\$0	\$3,850
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$3,850	\$0	\$3,850
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$3,850	\$0	\$3,850
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$163,250	\$0	\$163,250
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$3,850	\$0	\$3,850

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$134,000	\$0	\$134,000
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU	\$10,000	\$0	\$10,000
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$15,250	\$0	\$15,250
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$9,900	\$0	\$9,900
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$1,500	\$0	\$1,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Central California Coast DPS	\$163,250	\$0	\$163,250
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$3,850	\$0	\$3,850
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$3,850	\$0	\$3,850
		Steelhead	Oncorhynchus (=Salmo) mykiss	Northern California DPS	\$56,000	\$0	\$56,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$356,000	\$0	\$356,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$3,850	\$0	\$3,850
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	New York Bight DPS	\$500	\$0	\$500
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$400	\$0	\$400
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$45,750	\$0	\$45,750

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	Wherever found	\$1,300	\$0	\$1,300
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$286,000	\$0	\$286,000
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$45,000	\$0	\$45,000
		Tui chub, Mohave	Gila bicolor ssp. mohavensis	Wherever found	\$32,045	\$0	\$32,045
	<i>Fishes Subtotal</i>				<i>\$2,514,196</i>	<i>\$0</i>	<i>\$2,514,196</i>
	Clams	Bean, Cumberland (pearlymussel)	Villosa trabalis	Wherever found; Except where listed as Experimental Populations	\$707	\$0	\$707
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$200	\$0	\$200
		Combshell, Cumberlandian	Epioblasma brevidens	Wherever found; Except where listed as Experimental Populations	\$707	\$0	\$707
		Elktoe, Cumberland	Alasmidonta atropurpurea	Wherever found	\$707	\$0	\$707
		Fanshell	Cyprogenia stegaria	Wherever found	\$200	\$0	\$200
		Higgins eye (pearlymussel)	Lampsilis higginsii	Wherever found	\$18,000	\$0	\$18,000
		Kidneyshell, fluted	Ptychobranhus subtentus	Wherever found	\$707	\$0	\$707
		Mapleleaf, winged	Quadrula fragosa	Wherever found, except where listed as an experimental population	\$18,000	\$0	\$18,000
		Mucket, pink (pearlymussel)	Lampsilis abrupta	Wherever found	\$907	\$0	\$907

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Mussel, oyster	Epioblasma capsaeformis	Wherever found; Except where listed as Experimental Populations	\$707	\$0	\$707
		Mussel, sheepnose	Plethobasus cyphus	Wherever found	\$200	\$0	\$200
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$200	\$0	\$200
		Pearlymussel, dromedary	Dromus dromas	Wherever found; Except where listed as Experimental Populations	\$707	\$0	\$707
		Pearlymussel, littlewing	Pegias fabula	Wherever found	\$707	\$0	\$707
		Pigtoe, rough	Pleurobema plenum	Wherever found	\$200	\$0	\$200
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$200	\$0	\$200
		Riffleshell, tan	Epioblasma florentina walkeri (=E. walkeri)	Wherever found	\$707	\$0	\$707
		Ring pink (mussel)	Obovaria retusa	Wherever found	\$200	\$0	\$200
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$4,907	\$0	\$4,907
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$2,500	\$0	\$2,500
	<i>Clams Subtotal</i>				<i>\$51,370</i>	<i>\$0</i>	<i>\$51,370</i>
	Snails	Abalone, Black	Haliotis cracherodii	Wherever found	\$11,000	\$0	\$11,000
		Abalone, White	Haliotis sorenseni	North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	\$1,000	\$0	\$1,000
		Snail, Guam tree	Partula radiolata	Wherever found	\$200	\$0	\$200
		Snail, Humped tree	Partula gibba	Wherever found	\$300	\$0	\$300
	<i>Snails Subtotal</i>				<i>\$12,500</i>	<i>\$0</i>	<i>\$12,500</i>

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
	Insects	Butterfly, Bartram's hairstreak	Strymon acis bartrami	Wherever found	\$30,000	\$0	\$30,000
		Butterfly, Florida leafwing	Anaea troglodyta floridalis	Wherever found	\$30,000	\$0	\$30,000
		Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$500	\$0	\$500
		Butterfly, Mariana eight-spot	Hypolimnas octocula marianensis	Wherever found	\$1,000	\$0	\$1,000
		Butterfly, mission blue	Icaricia icarioides missionensis	Wherever found	\$125,000	\$0	\$125,000
		Butterfly, Mitchell's satyr	Neonympha mitchellii mitchellii	Wherever found	\$3,000	\$0	\$3,000
		Butterfly, Oregon silverspot	Speyeria zerene hippolyta	Wherever found	\$3,800	\$0	\$3,800
		Butterfly, San Bruno elfin	Callophrys mossii bayensis	Wherever found	\$5,000	\$0	\$5,000
		Butterfly, Schaus swallowtail	Heraclides aristodemus ponceanus	Wherever found	\$10,000	\$0	\$10,000
		Damselfly, orangeblack Hawaiian	Megalagrion xanthomelas	Wherever found	\$200	\$0	\$200
		Damselfly, Pacific Hawaiian	Megalagrion pacificum	Wherever found	\$5,000	\$0	\$5,000
		Dragonfly, Hine's emerald	Somatochlora hineana	Wherever found	\$50	\$0	\$50
		Tiger beetle, Northeastern beach	Cicindela dorsalis dorsalis	Wherever found	\$900	\$0	\$900
	<i>Insects Subtotal</i>				<i>\$214,450</i>	<i>\$0</i>	<i>\$214,450</i>
	Crustaceans	Shrimp, anchialine pool	Procaris hawaiiiana	Wherever found	\$200	\$0	\$200
		Shrimp, California freshwater	Syncaris pacifica	Wherever found	\$5,000	\$0	\$5,000
		Shrimp, Kentucky cave	Palaemonias ganteri	Wherever found	\$30,000	\$0	\$30,000
	<i>Crustaceans Subtotal</i>				<i>\$35,200</i>	<i>\$0</i>	<i>\$35,200</i>
	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$16,300	\$0	\$16,300
		Coral, elkhorn	Acropora palmata	Wherever found	\$30,500	\$0	\$30,500
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$16,300	\$0	\$16,300
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$20,000	\$0	\$20,000
		Coral, Pillar	Dendrogyra cylindrus	Wherever found	\$21,300	\$0	\$21,300
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$16,300	\$0	\$16,300
		Coral, staghorn	Acropora cervicornis	Wherever found	\$82,800	\$0	\$82,800
		No common name	Acropora globiceps	Wherever found	\$10,000	\$0	\$10,000

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		No common name	Acropora jacquelineae	Wherever found	\$10,000	\$0	\$10,000
		No common name	Acropora retusa	Wherever found	\$10,000	\$0	\$10,000
		No common name	Acropora speciosa	Wherever found	\$10,000	\$0	\$10,000
		No common name	Euphyllia paradivisa	Wherever found	\$10,000	\$0	\$10,000
		No common name	Isopora crateriformis	Wherever found	\$10,000	\$0	\$10,000
	<i>Corals Subtotal</i>				\$263,500	\$0	\$263,500
	Flowering Plants	Alani	Melicope ovalis	Wherever found	\$36,000	\$0	\$36,000
		Alani	Melicope zahlbruckneri	Wherever found	\$1,973	\$0	\$1,973
		Alopecurus, Sonoma	Alopecurus aequalis var. sonomensis	Wherever found	\$1,000	\$0	\$1,000
		Amaranth, seabeach	Amaranthus pumilus	Wherever found	\$53,887	\$0	\$53,887
		Avens, spreading	Geum radiatum	Wherever found	\$3,350	\$0	\$3,350
		Barberry, island	Berberis pinnata ssp. insularis	Wherever found	\$2,000	\$0	\$2,000
		bladderpod, Missouri	Physaria filiformis	Wherever found	\$40,000	\$0	\$40,000
		Blazingstar, Heller's	Liatris helleri	Wherever found	\$3,500	\$0	\$3,500
		Cactus, Acuna	Echinomastus erectocentrus var. acunensis	Wherever found	\$608	\$0	\$608
		Cactus, Brady pincushion	Pediocactus bradyi	Wherever found	\$500	\$0	\$500
		Cactus, Chisos Mountain hedgehog	Echinocereus chisoensis var. chisoensis	Wherever found	\$500	\$0	\$500
		Cactus, Florida semaphore	Consolea corallicola	Wherever found	\$2,300	\$0	\$2,300
		Cactus, Lloyd's Mariposa	Echinomastus mariposensis	Wherever found	\$500	\$0	\$500
		Cactus, Winkler	Pediocactus winkleri	Wherever found	\$35,500	\$0	\$35,500
		Cactus, Wright fishhook	Sclerocactus wrightiae	Wherever found	\$35,500	\$0	\$35,500
		Catchfly, Spalding's	Silene spaldingii	Wherever found	\$1,500	\$0	\$1,500
		Clarkia, Presidio	Clarkia franciscana	Wherever found	\$2,000	\$0	\$2,000
		Cory cactus, bunched	Coryphantha ramillosa	Wherever found	\$500	\$0	\$500
		Crabgrass, Florida pineland	Digitaria pauciflora	Wherever found	\$1,000	\$0	\$1,000
		Cycladenia, Jones	Cycladenia humilis var. jonesii	Wherever found	\$2,000	\$0	\$2,000
		Daisy, Parish's	Erigeron parishii	Wherever found	\$3,500	\$0	\$3,500
		Dwarf-flax, Marin	Hesperolinon congestum	Wherever found	\$2,000	\$0	\$2,000
		Fescue, Guadalupe	Festuca ligulata	Wherever found	\$500	\$0	\$500

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Geranium, Hawaiian red-flowered	<i>Geranium arboreum</i>	Wherever found	\$36,000	\$0	\$36,000
		Grass, Eureka Dune	<i>Swallenia alexandrae</i>		\$50,000	\$0	\$50,000
		Haha	<i>Cyanea asplenifolia</i>	Wherever found	\$36,000	\$0	\$36,000
		Haha	<i>Cyanea copelandii</i> ssp. <i>haleakalaensis</i>	Wherever found	\$36,000	\$0	\$36,000
		Haha	<i>Cyanea hamatiflora</i> ssp. <i>hamatiflora</i>	Wherever found	\$12,000	\$0	\$12,000
		Haha	<i>Cyanea kunthiana</i>	Wherever found	\$12,000	\$0	\$12,000
		haha	<i>Cyanea maritae</i>	Wherever found	\$23,999	\$0	\$23,999
		Haha	<i>Cyanea shipmanii</i>	Wherever found	\$6,973	\$0	\$6,973
		Haha	<i>Cyanea stictophylla</i>	Wherever found	\$6,973	\$0	\$6,973
		haiwale	<i>Cyrtandra ferripilosa</i>	Wherever found	\$36,000	\$0	\$36,000
		Hala pepe	<i>Pleomele hawaiiensis</i>	Wherever found	\$5,000	\$0	\$5,000
		Harperella	<i>Ptilimnium nodosum</i>	Wherever found	\$46,427	\$0	\$46,427
		Hau kuahiwi	<i>Hibiscadelphus giffardianus</i>	Wherever found	\$5,000	\$0	\$5,000
		Ha`iwale	<i>Cyrtandra giffardii</i>	Wherever found	\$1,973	\$0	\$1,973
		Ha`iwale	<i>Cyrtandra tintinnabula</i>	Wherever found	\$1,973	\$0	\$1,973
		Heau	<i>Exocarpos menziesii</i>		\$1,000	\$0	\$1,000
		Hoawa	<i>Pittosporum halophilum</i>	Wherever found	\$5,000	\$0	\$5,000
		Hoawa	<i>Pittosporum hawaiiense</i>	Wherever found	\$5,000	\$0	\$5,000
		Holei	<i>Ochrosia haleakalae</i>	Wherever found	\$1,500	\$0	\$1,500
		Ischaemum, Hilo	<i>Ischaemum byrone</i>	Wherever found	\$24,000	\$0	\$24,000
		Koki`o	<i>Kokia drynarioides</i>	Wherever found	\$500	\$0	\$500
		Koki`o ke`oke`o	<i>Hibiscus arnottianus</i> ssp. <i>immaculatus</i>	Wherever found	\$5,000	\$0	\$5,000
		Ko`oko`olau	<i>Bidens campylotheca</i> ssp. <i>pentamera</i>	Wherever found	\$36,000	\$0	\$36,000
		Ko`oko`olau	<i>Bidens campylotheca</i> ssp. <i>waihoiense</i>	Wherever found	\$36,000	\$0	\$36,000
		Ko`oko`olau	<i>Bidens micrantha</i> ssp. <i>ctenophylla</i>	Wherever found	\$500	\$0	\$500
		Ko`oko`olau	<i>Bidens micrantha</i> ssp. <i>kalealaha</i>	Wherever found	\$36,000	\$0	\$36,000

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Kuahiwi laukahi	<i>Plantago hawaiiensis</i>	Wherever found	\$500	\$0	\$500
		Kuahiwi laukahi	<i>Plantago princeps</i>	Wherever found	\$36,000	\$0	\$36,000
		Kula wahine noho	<i>Isodendron pyrifolium</i>	Wherever found	\$500	\$0	\$500
		Layia, beach	<i>Layia carnosa</i>	Wherever found	\$2,500	\$0	\$2,500
		Lessingia, San Francisco	<i>Lessingia germanorum</i> (=L.g. var. <i>germanorum</i> )	Wherever found	\$3,000	\$0	\$3,000
		Lo`ulu	<i>Pritchardia maideniana</i>	Wherever found	\$500	\$0	\$500
		Lupine, clover	<i>Lupinus tidestromii</i>	Wherever found	\$3,000	\$0	\$3,000
		Makou	<i>Peucedanum sandwicense</i>	Wherever found	\$5,000	\$0	\$5,000
		Makou	<i>Ranunculus hawaiiensis</i>	Wherever found	\$13,973	\$0	\$13,973
		Makou	<i>Ranunculus mauiensis</i>	Wherever found	\$12,000	\$0	\$12,000
		Manzanita, Franciscan	<i>Arctostaphylos franciscana</i>	Wherever found	\$44,000	\$0	\$44,000
		Manzanita, Presidio	<i>Arctostaphylos hookeri</i> var. <i>ravenii</i>	Wherever found	\$4,000	\$0	\$4,000
		Manzanita, Santa Rosa Island	<i>Arctostaphylos confertiflora</i>	Wherever found	\$1,600	\$0	\$1,600
		Ma`o hau hele, (=native yellow hibiscus)	<i>Hibiscus brackenridgei</i>	Wherever found	\$500	\$0	\$500
		Milk-vetch, Sentry	<i>Astragalus cremnophylax</i> var. <i>cremnophylax</i>	Wherever found	\$86,000	\$0	\$86,000
		Milk-vetch, Shivwits	<i>Astragalus ampullarioides</i>	Wherever found	\$5,925	\$0	\$5,925
		Milk-vetch, triple-ribbed	<i>Astragalus tricarinatus</i>	Wherever found	\$3,500	\$0	\$3,500
		monkey-flower, Michigan	<i>Mimulus michiganensis</i>	Wherever found	\$1,365	\$0	\$1,365
		Nanu	<i>Gardenia remyi</i>	Wherever found	\$12,000	\$0	\$12,000
		Naupaka, dwarf	<i>Scaevola coriacea</i>	Wherever found	\$5,000	\$0	\$5,000
		No common name	<i>Neraudia ovata</i>	Wherever found	\$2,435	\$0	\$2,435
		No common name	<i>Phyllostegia bracteata</i>	Wherever found	\$61,000	\$0	\$61,000
		No common name	<i>Phyllostegia brevidens</i>	Wherever found	\$60,000	\$0	\$60,000
		No common name	<i>Phyllostegia haliakalae</i>	Wherever found	\$61,000	\$0	\$61,000
		No common name	<i>Phyllostegia stachyoides</i>	Wherever found	\$500	\$0	\$500
		No common name	<i>Platanthera holochila</i>	Wherever found	\$12,000	\$0	\$12,000
		No common name	<i>Sanicula sandwicensis</i>	Wherever found	\$37,973	\$0	\$37,973
		No common name	<i>Schiedea diffusa</i> ssp. <i>macraei</i>	Wherever found	\$5,000	\$0	\$5,000



Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		No common name	<i>Schiedea diffusa</i> subsp. <i>diffusa</i>	Wherever found	\$12,000	\$0	\$12,000
		No common name	<i>Schiedea haleakalensis</i>	Wherever found	\$61,000	\$0	\$61,000
		No common name	<i>Silene hawaiiensis</i>	Wherever found	\$1,000	\$0	\$1,000
		No common name	<i>Tetramolopium rockii</i>	Wherever found	\$5,000	\$0	\$5,000
		No common name	<i>Wikstroemia villosa</i>	Wherever found	\$12,000	\$0	\$12,000
		Nohoanu	<i>Geranium hanaense</i>	Wherever found	\$12,000	\$0	\$12,000
		Nohoanu	<i>Geranium multiflorum</i>	Wherever found	\$12,000	\$0	\$12,000
		nui, haha	<i>Cyanea horrida</i>	Wherever found	\$12,000	\$0	\$12,000
		Ohai	<i>Sesbania tomentosa</i>	Wherever found	\$2,500	\$0	\$2,500
		Orchid, western prairie fringed	<i>Platanthera praeclara</i>	Wherever found	\$8,000	\$0	\$8,000
		Orchid, white fringeless	<i>Platanthera integrilabia</i>		\$1,000	\$0	\$1,000
		paintbrush, golden	<i>Castilleja levisecta</i>	Wherever found	\$15,000	\$0	\$15,000
		Paintbrush, Tiburon	<i>Castilleja affinis</i> ssp. <i>neglecta</i>	Wherever found	\$1,000	\$0	\$1,000
		Panicgrass, Carter's	<i>Panicum fauriei</i> var. <i>carteri</i>	Wherever found	\$5,000	\$0	\$5,000
		Phlox, Texas trailing	<i>Phlox nivalis</i> ssp. <i>texensis</i>	Wherever found	\$5,000	\$0	\$5,000
		Pink, swamp	<i>Helonias bullata</i>	Wherever found	\$3,500	\$0	\$3,500
		Pogonia, small whorled	<i>Isotria medeoloides</i>	Wherever found	\$5,600	\$0	\$5,600
		Popolo	<i>Solanum nelsonii</i>	Wherever found	\$5,000	\$0	\$5,000
		Popolo ku mai	<i>Solanum incompletum</i>	Wherever found	\$12,000	\$0	\$12,000
		potato-bean, Prices	<i>Apios priceana</i>		\$1,000	\$0	\$1,000
		Potentilla, Hickman's	<i>Potentilla hickmanii</i>	Wherever found	\$34,000	\$0	\$34,000
		Po`e	<i>Portulaca sclerocarpa</i>	Wherever found	\$2,500	\$0	\$2,500
		Prairie-clover, Florida	<i>Dalea carthagenensis floridana</i>	Wherever found	\$1,000	\$0	\$1,000
		Prickly-apple, fragrant	<i>Cereus eriophorus</i> var. <i>fragrans</i>	Wherever found	\$2,000	\$0	\$2,000
		Pua `ala	<i>Brighamia rockii</i>	Wherever found	\$500	\$0	\$500
		Reedgrass, Maui	<i>Calamagrostis expansa</i>	Wherever found	\$12,000	\$0	\$12,000
		Rosemary, Cumberland	<i>Conradina verticillata</i>	Wherever found	\$1,100	\$0	\$1,100
		Sandwort, Cumberland	<i>Arenaria cumberlandensis</i>	Wherever found	\$1,000	\$0	\$1,000
		Sandwort, Marsh	<i>Arenaria paludicola</i>	Wherever found	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General	Land	Grand Total
		Silversword, Mauna Loa (=Ka'u)	Argyroxiphium kauense	Wherever found	\$10,000	\$0	\$10,000
		Skullcap, large-flowered	Scutellaria montana	Wherever found	\$1,000	\$0	\$1,000
		Spineflower, Sonoma	Chorizanthe valida	Wherever found	\$1,000	\$0	\$1,000
		Thistle, Pitcher's	Cirsium pitcheri	Wherever found	\$21,930	\$0	\$21,930
		Townsendia, Last Chance	Townsendia aprica	Wherever found	\$400	\$0	\$400
		`Ahinahina	Argyroxiphium sandwicense ssp. macrocephalum	Wherever found	\$125,000	\$0	\$125,000
		`Aiea	Nothoestrum latifolium	Wherever found	\$12,000	\$0	\$12,000
		`aku	Cyanea tritomantha	Wherever found	\$1,973	\$0	\$1,973
		`Anunu	Sicyos albus	Wherever found	\$4,973	\$0	\$4,973
		`Anunu	Sicyos macrophyllus	Wherever found	\$6,000	\$0	\$6,000
		`Awikiwiki	Canavalia molokaiensis	Wherever found	\$5,000	\$0	\$5,000
		`Ena`ena	Pseudognaphalium sandwicense var. molokaiense	Wherever found	\$500	\$0	\$500
		`Oha wai	Clermontia lindseyana	Wherever found	\$6,973	\$0	\$6,973
		`Oha wai	Clermontia peleana	Wherever found	\$5,000	\$0	\$5,000
		`Oha wai	Clermontia samuelii	Wherever found	\$36,000	\$0	\$36,000
		`Ohe	Joinvillea ascendens ascendens	Wherever found	\$36,000	\$0	\$36,000
	<i>Flowering Plants Subtotal</i>				<i>\$1,698,156</i>	<i>\$0</i>	<i>\$1,698,156</i>
	Ferns and Allies	Kupukupu makalii	Cyclosorus boydiae	Wherever found	\$12,000	\$0	\$12,000
		No common name	Asplenium peruvianum var. insulare	Wherever found	\$13,973	\$0	\$13,973
		No common name	Microlepia strigosa var. mauiensis	Wherever found	\$12,000	\$0	\$12,000
		Pauoa	Ctenitis squamigera	Wherever found	\$12,000	\$0	\$12,000
		Wawae`iole	Huperzia mannii	Wherever found	\$12,000	\$0	\$12,000
	<i>Ferns and Allies Subtotal</i>				<i>\$61,973</i>	<i>\$0</i>	<i>\$61,973</i>
<i>Domestic Subtotal</i>					<i>\$17,982,068</i>	<i>\$0</i>	<i>\$17,982,068</i>
Foreign	Fishes	Blindcat, Mexican (catfish)	Prietella phreatophila	Wherever found	\$41,000	\$0	\$41,000
<i>Foreign Subtotal</i>					<i>\$41,000</i>	<i>\$0</i>	<i>\$41,000</i>
	Multi-Species				\$11,536	\$0	\$11,536

<b>Locality</b>	<b>Group Name</b>	<b>Common Name</b>	<b>Scientific Name</b>	<b>Population</b>	<b>General</b>	<b>Land</b>	<b>Grand Total</b>
<i>Multi-Species Subtotal</i>					<i>\$11,536</i>	<i>\$0</i>	<i>\$11,536</i>
<b>Total</b>					<b>\$18,034,604</b>	<b>\$0</b>	<b>\$18,034,604</b>

## DOI US Geological Survey - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$9,330	\$0	\$9,330
		Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$168,352	\$0	\$168,352
		Bat, Indiana	Myotis sodalis	Wherever found	\$30,000	\$0	\$30,000
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$90	\$0	\$90
		Bat, Ozark big-eared	Corynorhinus (=Plecotus) townsendii ingens	Wherever found	\$30	\$0	\$30
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$678,176	\$0	\$678,176
		Bear, polar	Ursus maritimus	Wherever found	\$1,493,650	\$0	\$1,493,650
		Ferret, black-footed	Mustela nigripes	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$66,901	\$0	\$66,901
		Jaguar	Panthera onca	Wherever found	\$6,000	\$0	\$6,000
		Kangaroo rat, giant	Dipodomys ingens	Wherever found	\$3,000	\$0	\$3,000
		Kangaroo rat, Stephens'	Dipodomys stephensi (incl. D. cactus)	Wherever found	\$3,000	\$0	\$3,000
		Manatee, West Indian	Trichechus manatus	Wherever found	\$841,839	\$0	\$841,839
		Mouse, Anastasia Island beach	Peromyscus polionotus phasma	Wherever found	\$1,200	\$0	\$1,200
		Mouse, Choctawhatchee beach	Peromyscus polionotus allophrys	Wherever found	\$1,500	\$0	\$1,500
		Mouse, Pacific pocket	Perognathus longimembris pacificus	Wherever found	\$15,000	\$0	\$15,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Mouse, Perdido Key beach	<i>Peromyscus polionotus trissyllepsis</i>	Wherever found	\$1,500	\$0	\$1,500
		Mouse, St. Andrew beach	<i>Peromyscus polionotus peninsularis</i>	Wherever found	\$1,500	\$0	\$1,500
		Ocelot	<i>Leopardus (=Felis) pardalis</i>	wherever found	\$6,000	\$0	\$6,000
		Otter, Northern Sea	<i>Enhydra lutris kenyoni</i>	Southwest Alaska DPS	\$55,250	\$0	\$55,250
		Otter, southern sea	<i>Enhydra lutris nereis</i>	Wherever found	\$321,123	\$0	\$321,123
		Panther, Florida	<i>Puma (=Felis) concolor coryi</i>	Wherever found	\$3,000	\$0	\$3,000
		Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	U.S.A. (AZ), Mexico	\$3,000	\$0	\$3,000
		Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	Wherever found, except where listed as an experimental population	\$3,000	\$0	\$3,000
		Puma (=mountain lion)	<i>Puma (=Felis) concolor</i> (all subsp. except <i>coryi</i> )	FL	\$3,000	\$0	\$3,000
		Seal, Guadalupe fur	<i>Arctocephalus townsendi</i>	Wherever found	\$200	\$0	\$200
		Sheep, Sierra Nevada bighorn	<i>Ovis canadensis sierrae</i>	Sierra Nevada	\$56,977	\$0	\$56,977
		Squirrel, Carolina northern flying	<i>Glaucomys sabrinus coloratus</i>	Wherever found	\$15,000	\$0	\$15,000
		Whale, North Atlantic Right	<i>Eubalaena glacialis</i>	Wherever found	\$8,250	\$0	\$8,250
		Wolf, gray	<i>Canis lupus</i>	MN	\$300	\$0	\$300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$7,500	\$0	\$7,500
		Wolf, Mexican	Canis lupus baileyi	Wherever found, except where listed as an experimental population	\$12,030	\$0	\$12,030
	<i>Mammals Subtotal</i>				<i>\$3,815,698</i>	<i>\$0</i>	<i>\$3,815,698</i>
	Birds	Akekee	Loxops caeruleirostris	Wherever found	\$74,489	\$0	\$74,489
		akepa, Hawaii	Loxops coccineus	Wherever found	\$2,610	\$0	\$2,610
		akiapolaau	Hemignathus wilsoni	Wherever found	\$2,609	\$0	\$2,609
		Akikiki	Oreomystis bairdi	Wherever found	\$74,489	\$0	\$74,489
		Bobwhite, masked (quail)	Colinus virginianus ridgwayi	Wherever found	\$11,000	\$0	\$11,000
		Condor, California	Gymnogyps californianus	U.S.A. (specific portions of Arizona, Nevada, and Utah)	\$7,000	\$0	\$7,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Condor, California	<i>Gymnogyps californianus</i>	U.S.A. only, except where listed as an experimental population	\$155,903	\$0	\$155,903
		Coot, Hawaiian	<i>Fulica americana alai</i>	Wherever found	\$41,448	\$0	\$41,448
		Crane, Mississippi sandhill	<i>Grus canadensis pulla</i>	Wherever found	\$600	\$0	\$600
		Crane, whooping	<i>Grus americana</i>	U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	\$252,440	\$0	\$252,440
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$280	\$0	\$280
		Creeper, Hawaii	<i>Oreomystis mana</i>	Wherever found	\$2,610	\$0	\$2,610
		Crow, Hawaiian (= 'alala)	<i>Corvus hawaiiensis</i>	Wherever found	\$25,000	\$0	\$25,000
		Crow, Mariana (=aga)	<i>Corvus kubaryi</i>	Wherever found	\$8,439	\$0	\$8,439
		Duck, Hawaiian (=koloa)	<i>Anas wyvilliana</i>	Wherever found	\$172,288	\$0	\$172,288
		Duck, Laysan	<i>Anas laysanensis</i>	Wherever found	\$300	\$0	\$300
		Eider, spectacled	<i>Somateria fischeri</i>	Wherever found	\$20,000	\$0	\$20,000
		Eider, Steller's	<i>Polysticta stelleri</i>	AK breeding pop.	\$20,000	\$0	\$20,000
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$33,267	\$0	\$33,267
		gallinule, Hawaiian common	<i>Gallinula galeata sandvicensis</i>	Wherever found	\$10,000	\$0	\$10,000
		Gnatcatcher, coastal California	<i>Polioptila californica californica</i>	Wherever found	\$28,000	\$0	\$28,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Goose, Hawaiian	Branta (=Nesochen) sandvicensis	Wherever found	\$1,500	\$0	\$1,500
		Hawk, Puerto Rican sharp-shinned	Accipiter striatus venator	Wherever found	\$22,500	\$0	\$22,500
		Kite, Everglade snail	Rostrhamus sociabilis plumbeus	Wherever found	\$30,000	\$0	\$30,000
		Knot, red	Calidris canutus rufa	Wherever found	\$4,800	\$0	\$4,800
		Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR, WA)	\$37,255	\$0	\$37,255
		Owl, northern spotted	Strix occidentalis caurina	Wherever found	\$244,191	\$0	\$244,191
		Palila (honeycreeper)	Loxioides bailleui	Wherever found	\$1,150	\$0	\$1,150
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$2,400	\$0	\$2,400
		Plover, piping	Charadrius melodus	[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$44,201	\$0	\$44,201



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$30,390	\$0	\$30,390
		Rail, California clapper	Rallus longirostris obsoletus	Wherever found	\$245,495	\$0	\$245,495
		Rail, light-footed clapper	Rallus longirostris levipes	Wherever found	\$5,816	\$0	\$5,816
		Sparrow, Cape Sable seaside	Ammodramus maritimus mirabilis	Wherever found	\$74,000	\$0	\$74,000
		Stilt, Hawaiian	Himantopus mexicanus knudseni	Wherever found	\$31,148	\$0	\$31,148
		Swiftlet, Mariana gray	Aerodramus vanikorensis bartschi	Wherever found	\$42,169	\$0	\$42,169
		Tern, California least	Sterna antillarum browni	Wherever found	\$1,000	\$0	\$1,000
		Tern, least	Sterna antillarum	interior pop.	\$1,200	\$0	\$1,200
		Tern, roseate	Sterna dougallii dougallii	Northeast U.S. nesting population	\$30,000	\$0	\$30,000
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$100,000	\$0	\$100,000
		Warbler, Kirtland's	Setophaga kirtlandii (= Dendroica kirtlandii)	Wherever found	\$26,000	\$0	\$26,000
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$109,442	\$0	\$109,442
	<i>Birds Subtotal</i>				<i>\$2,027,429</i>	<i>\$0</i>	<i>\$2,027,429</i>
	Reptiles	Crocodile, American	Crocodylus acutus	U.S.A. (FL)	\$120,000	\$0	\$120,000
		gartersnake, narrow-headed	Thamnophis rufipunctatus	Wherever found	\$29,010	\$0	\$29,010
		gartersnake, northern Mexican	Thamnophis eques megalops	Wherever found	\$28,000	\$0	\$28,000
		Lizard, blunt-nosed leopard	Gambelia silus	Wherever found	\$10,000	\$0	\$10,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Lizard, Coachella Valley fringe-toed	Uma inornata	Wherever found	\$39,245	\$0	\$39,245
		Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$70,000	\$0	\$70,000
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$95,410	\$0	\$95,410
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$65,000	\$0	\$65,000
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$150,010	\$0	\$150,010
		Sea turtle, loggerhead	Caretta caretta	North Pacific Ocean DPS	\$8,139	\$0	\$8,139
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$98,410	\$0	\$98,410
		Snake, black pine	Pituophis melanoleucus lodingi	Wherever found	\$4,000	\$0	\$4,000
		Snake, giant garter	Thamnophis gigas	Wherever found	\$28,494	\$0	\$28,494
		Snake, San Francisco garter	Thamnophis sirtalis tetrataenia	Wherever found	\$28,494	\$0	\$28,494
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$545,912	\$0	\$545,912
		Whipsnake (=striped racer), Alameda	Masticophis lateralis euryxanthus	Wherever found	\$38,495	\$0	\$38,495
	<i>Reptiles Subtotal</i>				<i>\$1,358,619</i>	<i>\$0</i>	<i>\$1,358,619</i>
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$82,207	\$0	\$82,207
		Frog, Chiricahua leopard	Rana chiricahuensis	Wherever found	\$59,287	\$0	\$59,287
		Frog, mountain yellow-legged	Rana muscosa	Southern California DPS	\$52,671	\$0	\$52,671
		Frog, Oregon spotted	Rana pretiosa	Wherever found	\$312,685	\$0	\$312,685
		Frog, Sierra Nevada Yellow-legged	Rana sierrae	Wherever found	\$81,379	\$0	\$81,379

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salamander, desert slender	Batrachoseps aridus	Wherever found	\$10,000	\$0	\$10,000
		Salamander, frosted flatwoods	Ambystoma cingulatum	Wherever found	\$46,800	\$0	\$46,800
		Salamander, Shenandoah	Plethodon shenandoah	Wherever found	\$60,181	\$0	\$60,181
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$12,260	\$0	\$12,260
		Toad, Houston	Bufo houstonensis	Wherever found	\$4,800	\$0	\$4,800
		Toad, Wyoming	Bufo hemiophrys baxteri	Wherever found	\$3,000	\$0	\$3,000
		toad, Yosemite	Anaxyrus canorus	Wherever found	\$29,006	\$0	\$29,006
	<i>Amphibians Subtotal</i>				\$754,276	\$0	\$754,276
	Fishes	Cavefish, Ozark	Amblyopsis rosae	Wherever found	\$15,000	\$0	\$15,000
		Chub, humpback	Gila cypha	Wherever found	\$33,000	\$0	\$33,000
		Darter, amber	Percina antesella	Wherever found	\$3,402	\$0	\$3,402
		Darter, Okaloosa	Etheostoma okaloosae	Wherever found	\$95,000	\$0	\$95,000
		Goby, tidewater	Eucyclogobius newberryi	Wherever found	\$6,000	\$0	\$6,000
		Logperch, Roanoke	Percina rex	Wherever found	\$18,000	\$0	\$18,000
		Minnow, Rio Grande Silvery	Hybognathus amarus	Wherever found, except where listed as an experimental population	\$24,000	\$0	\$24,000
		Pikeminnow (=squawfish), Colorado	Ptychocheilus lucius	Wherever found, except where listed as an experimental population	\$6,000	\$0	\$6,000
		Pupfish, desert	Cyprinodon macularius	Wherever found	\$3,000	\$0	\$3,000
		Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$60,000	\$0	\$60,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$12,000	\$0	\$12,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$15,256	\$0	\$15,256
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$181,000	\$0	\$181,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter-run ESU	\$141,196	\$0	\$141,196
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$43,369	\$0	\$43,369
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$31,896	\$0	\$31,896
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU	\$95,528	\$0	\$95,528
		Shiner, Arkansas River	Notropis girardi	Arkansas River Basin (AR, KS, NM, OK, TX)	\$51,340	\$0	\$51,340
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$58,282	\$0	\$58,282
		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$45,000	\$0	\$45,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$15,000	\$0	\$15,000
		Stickleback, unarmored threespine	Gasterosteus aculeatus williamsoni	Wherever found	\$8,000	\$0	\$8,000
		Sturgeon, Alabama	Scaphirhynchus suttkusi	Wherever found	\$19,635	\$0	\$19,635
		Sturgeon, Gulf	Acipenser oxyrinchus (=oxyrhynchus) desotoi	Wherever found	\$50,800	\$0	\$50,800
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$388,818	\$0	\$388,818
		Sturgeon, Shovelnose	Scaphirhynchus platyrhynchus	Wherever found	\$31,635	\$0	\$31,635
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$62,077	\$0	\$62,077
		Sucker, Santa Ana	Catostomus santaanae	3 CA river basins	\$21,651	\$0	\$21,651

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	Wherever found	\$10,500	\$0	\$10,500
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$338,172	\$0	\$338,172
		Trout, Gila	Oncorhynchus gilae	Wherever found	\$48,000	\$0	\$48,000
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$134,027	\$0	\$134,027
		Tui chub, Mohave	Gila bicolor ssp. mohavensis	Wherever found	\$39,816	\$0	\$39,816
	<i>Fishes Subtotal</i>				<i>\$2,106,400</i>	<i>\$0</i>	<i>\$2,106,400</i>
	Clams	Bean, Choctaw	Villosa choctawensis	Wherever found	\$13,867	\$0	\$13,867
		Kidneyshell, southern	Ptychobranhus jonesi	Wherever found	\$13,867	\$0	\$13,867
		Pigtoe, fuzzy	Pleurobema strodeanum	Wherever found	\$13,867	\$0	\$13,867
		Pigtoe, tapered	Fusconaia burkei	Wherever found	\$13,867	\$0	\$13,867
		Sandshell, Southern	Hamiota australis	Wherever found	\$13,867	\$0	\$13,867
	<i>Clams Subtotal</i>				<i>\$69,335</i>	<i>\$0</i>	<i>\$69,335</i>
	Insects	Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$7,500	\$0	\$7,500
		Butterfly, Mitchell's satyr	Neonympha mitchellii mitchellii	Wherever found	\$5,000	\$0	\$5,000
		Butterfly, Oregon silverspot	Speyeria zerene hippolyta	Wherever found	\$24,871	\$0	\$24,871
	<i>Insects Subtotal</i>				<i>\$37,371</i>	<i>\$0</i>	<i>\$37,371</i>
	Arachnids	Spider, spruce-fir moss	Microhexura montivaga	Wherever found	\$3,000	\$0	\$3,000
	<i>Arachnids Subtotal</i>				<i>\$3,000</i>	<i>\$0</i>	<i>\$3,000</i>
	Crustaceans	Crayfish, Hell Creek Cave	Cambarus zophonastes	Wherever found	\$30,000	\$0	\$30,000
	<i>Crustaceans Subtotal</i>				<i>\$30,000</i>	<i>\$0</i>	<i>\$30,000</i>
	Flowering Plants	Baccharis, Encinitas	Baccharis vanessae	Wherever found	\$7,000	\$0	\$7,000
		Barberry, island	Berberis pinnata ssp. insularis	Wherever found	\$8,399	\$0	\$8,399
		Bedstraw, island	Galium buxifolium	Wherever found	\$4,256	\$0	\$4,256
		Bird's-beak, salt marsh	Cordylanthus maritimus ssp. maritimus	Wherever found	\$7,000	\$0	\$7,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Bush-mallow, Santa Cruz Island	Malacothamnus fasciculatus var. nesioticus	Wherever found	\$6,534	\$0	\$6,534
		Cactus, Arizona hedgehog	Echinocereus triglochidiatus var. arizonicus	Wherever found	\$14,004	\$0	\$14,004
		Gilia, Hoffmann's slender-flowered	Gilia tenuiflora ssp. hoffmannii	Wherever found	\$7,058	\$0	\$7,058
		Haha	Cyanea calycina	Wherever found	\$3,104	\$0	\$3,104
		Liveforever, Santa Barbara Island	Dudleya traskiae	Wherever found	\$1,139	\$0	\$1,139
		Malacothrix, Santa Cruz Island	Malacothrix indecora	Wherever found	\$11,050	\$0	\$11,050
		Manzanita, Santa Rosa Island	Arctostaphylos confertiflora	Wherever found	\$12,768	\$0	\$12,768
		Monardella, willowy	Monardella viminea	Wherever found	\$7,000	\$0	\$7,000
		Paintbrush, soft-leaved	Castilleja mollis	Wherever found	\$7,674	\$0	\$7,674
		Phacelia, island	Phacelia insularis ssp. insularis	Wherever found	\$8,979	\$0	\$8,979
		Rock-cress, Hoffmann's	Arabis hoffmannii	Wherever found	\$12,925	\$0	\$12,925
		Rush-rose, island	Helianthemum greenei	Wherever found	\$6,632	\$0	\$6,632
		Tarplant, Otay	Deinandra (=Hemizonia) conjugens	Wherever found	\$7,000	\$0	\$7,000
		Thistle, Pitcher's	Cirsium pitcheri	Wherever found	\$9,651	\$0	\$9,651
		Thornmint, San Diego	Acanthomintha ilicifolia	Wherever found	\$7,000	\$0	\$7,000
		<i>Flowering Plants Subtotal</i>			\$149,173	\$0	\$149,173
	Ferns and Allies	Fern, Aleutian shield	Polystichum aleuticum	Wherever found	\$10,000	\$0	\$10,000
		<i>Ferns and Allies Subtotal</i>			\$10,000	\$0	\$10,000
	<i>Domestic Subtotal</i>				\$10,361,301	\$0	\$10,361,301
Foreign	Mammals	Gorilla	Gorilla gorilla	Wherever found	\$3,150	\$0	\$3,150
	Birds	Crane, hooded	Grus monacha	Wherever found	\$20	\$0	\$20
		Crane, Siberian white	Grus leucogeranus	Wherever found	\$20	\$0	\$20
	<i>Birds Subtotal</i>				\$40	\$0	\$40

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Reptiles	Sea turtle, loggerhead	Caretta caretta	Northeast Atlantic Ocean DPS	\$49,010	\$0	\$49,010
	Amphibians	Frog, Goliath	Conraua goliath	Wherever found	\$3,000	\$0	\$3,000
<i>Foreign Subtotal</i>					<i>\$55,200</i>	<i>\$0</i>	<i>\$55,200</i>
<b>Total</b>					<b>\$10,416,501</b>	<b>\$0</b>	<b>\$10,416,501</b>

## DOT Federal Aviation Administration - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	bat, Florida bonneted	Eumops floridanus	Wherever found	\$16,000	\$0	\$16,000
		jumping	Zapus hudsonius preblei	wherever found	\$8,000	\$0	\$8,000
		Prairie dog, Utah	Cynomys parvidens	Wherever found	\$10,000	\$0	\$10,000
	<i>Mammals Subtotal</i>				<i>\$34,000</i>	<i>\$0</i>	<i>\$34,000</i>
	Birds	Crane, whooping	Grus americana	except where	\$5,000	\$0	\$5,000
		willow	Empidonax traillii extimus	Wherever found	\$2,500	\$0	\$2,500
		Horned lark, streaked	Eremophila alpestris strigata	Wherever found	\$120,000	\$0	\$120,000
		Knot, red	Calidris canutus rufa	Wherever found	\$1,000	\$0	\$1,000
		Stork, wood	Mycteria americana	NC, SC	\$10,000	\$0	\$10,000
	<i>Birds Subtotal</i>				<i>\$138,500</i>	<i>\$0</i>	<i>\$138,500</i>
	Reptiles	Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$3,500	\$0	\$3,500
	<i>Reptiles Subtotal</i>				<i>\$3,500</i>	<i>\$0</i>	<i>\$3,500</i>
	Multi-Species				\$89,800	\$0	\$89,800
	<i>Multi-Species Subtotal</i>				<i>\$89,800</i>	<i>\$0</i>	<i>\$89,800</i>
	<i>Domestic Subtotal</i>				<i>\$265,800</i>	<i>\$0</i>	<i>\$265,800</i>
Total					\$265,800	\$0	\$265,800



### Federal Communications Commission - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Multi-Species				\$127,900	\$0	\$127,900
Total					\$127,900	\$0	\$127,900

## Tennessee Valley Authority - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$168,000	\$0	\$168,000
		Bat, Indiana	Myotis sodalis	Wherever found	\$633,000	\$0	\$633,000
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$142,000	\$0	\$142,000
		Bat, Virginia big-eared	Corynorhinus (=Plecotus) townsendii virginianus	Wherever found	\$100,500	\$0	\$100,500
		Squirrel, Carolina northern flying	Glaucomys sabrinus coloratus	Wherever found	\$1,700	\$0	\$1,700
	<i>Mammals Subtotal</i>				<i>\$1,045,200</i>	<i>\$0</i>	<i>\$1,045,200</i>
	Birds	Knot, red	Calidris canutus rufa	Wherever found	\$200	\$0	\$200
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$1,000	\$0	\$1,000
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$2,000	\$0	\$2,000
		Tern, least	Sterna antillarum	interior pop.	\$200	\$0	\$200
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$2,500	\$0	\$2,500
	<i>Birds Subtotal</i>				<i>\$5,900</i>	<i>\$0</i>	<i>\$5,900</i>
	Reptiles	turtle, bog	Clemmys muhlenbergii	U.S.A. (GA, NC, SC, TN, VA)	\$500	\$0	\$500
		Turtle, flattened musk	Sternotherus depressus	Black Warrior R. system upstream from Bankhead Dam	\$200	\$0	\$200
		Turtle, ringed map	Graptemys oculifera	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	<i>Reptiles Subtotal</i>				\$900	\$0	\$900
	Amphibians	Waterdog, black warrior (=Sipsey Fork)	Necturus alabamensis	Wherever found	\$500	\$0	\$500
	<i>Amphibians Subtotal</i>				\$500	\$0	\$500
	Fishes	Darter, boulder	Etheostoma wapiti	Wherever found	\$59,500	\$0	\$59,500
	<i>Fishes Subtotal</i>				\$59,500	\$0	\$59,500
	Clams	Kidneyshell, fluted	Ptychobranthus subtentus	Wherever found	\$59,500	\$0	\$59,500
		Monkeyface, Cumberland (pearlymussel)	Quadrula intermedia	Wherever found; Except where listed as Experimental Populations	\$59,500	\$0	\$59,500
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$59,500	\$0	\$59,500
		Pearlymussel, birdwing	Lemiox rimosus	Wherever found; Except where listed as Experimental Populations	\$59,500	\$0	\$59,500
		Pearlymussel, cracking	Hemistena lata	Wherever found; Except where listed as Experimental Populations	\$59,500	\$0	\$59,500
		Pearlymussel, slabside	Pleuroaia dolabelloides	Wherever found	\$59,500	\$0	\$59,500
		Pigtoe, finerayed	Fusconaia cuneolus	Wherever found; Except where listed as Experimental Populations	\$59,500	\$0	\$59,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pigtoe, shiny	Fusconaia cor	Wherever found; Except where listed as Experimental Populations	\$59,500	\$0	\$59,500
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$59,500	\$0	\$59,500
	<i>Clams Subtotal</i>				\$535,500	\$0	\$535,500
	Snails	Snail, painted snake coiled forest	Anguispira picta	Wherever found	\$400	\$0	\$400
	Snails Subtotal						
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$200	\$0	\$200
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$200	\$0	\$200
		Butterfly, Mitchell's satyr	Neonympha mitchellii mitchellii	Wherever found	\$700	\$0	\$700
	<i>Insects Subtotal</i>				\$1,100	\$0	\$1,100
	Arachnids	Spider, spruce-fir moss	Microhexura montivaga	Wherever found	\$1,000	\$0	\$1,000
	<i>Arachnids Subtotal</i>				\$1,000	\$0	\$1,000
	Flowering Plants	Aster, Ruth's golden	Pityopsis ruthii	Wherever found	\$5,000	\$0	\$5,000
		Barbara's Buttons, Mohr's	Marshallia mohrii	Wherever found	\$1,000	\$0	\$1,000
		Orchid, white fringeless	Platanthera integrilabia		\$1,500	\$0	\$1,500
		Prairie-clover, leafy	Dalea foliosa	Wherever found	\$800	\$0	\$800
		Rock-cress, Braun's	Arabis perstellata	Wherever found	\$400	\$0	\$400
		Skullcap, large-flowered	Scutellaria montana	Wherever found	\$5,000	\$0	\$5,000
	<i>Flowering Plants Subtotal</i>				\$13,700	\$0	\$13,700
<i>Domestic Total</i>					\$1,663,700	\$0	\$1,663,700
Total					\$1,663,700	\$0	\$1,663,700

## USDA Animal and Plant Health Inspection Service - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$515,700	\$0	\$515,700
		Bat, Indiana	Myotis sodalis	Wherever found	\$9,200	\$0	\$9,200
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$2,000	\$0	\$2,000
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$227,000	\$0	\$227,000
		Caribou DPS, Southern Mountain	Rangifer tarandus ssp. caribou	Southern Mountain DPS	\$1,400	\$0	\$1,400
		Deer, key	Odocoileus virginianus clavium	Wherever found	\$300	\$0	\$300
		Ferret, black-footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$134,000	\$0	\$134,000
		Jaguar	Panthera onca	Wherever found	\$1,500	\$0	\$1,500
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$11,200	\$0	\$11,200
		Mouse, Alabama beach	Peromyscus polionotus ammobates	Wherever found	\$4,000	\$0	\$4,000
		Mouse, Choctawhatchee beach	Peromyscus polionotus allophrys	Wherever found	\$32,500	\$0	\$32,500
		Mouse, New Mexico meadow jumping	Zapus hudsonius luteus	Wherever found	\$100	\$0	\$100
		Mouse, Pacific pocket	Perognathus longimembris pacificus	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Mouse, Perdido Key beach	<i>Peromyscus polionotus trissyllepsis</i>	Wherever found	\$32,500	\$0	\$32,500
		Mouse, salt marsh harvest	<i>Reithrodontomys raviventris</i>	wherever found	\$170,800	\$0	\$170,800
		Mouse, southeastern beach	<i>Peromyscus polionotus niveiventris</i>	wherever found	\$32,500	\$0	\$32,500
		Mouse, St. Andrew beach	<i>Peromyscus polionotus peninsularis</i>	Wherever found	\$32,500	\$0	\$32,500
		Ocelot	<i>Leopardus (=Felis) pardalis</i>	wherever found	\$9,900	\$0	\$9,900
		Prairie dog, Utah	<i>Cynomys parvidens</i>	Wherever found	\$2,000	\$0	\$2,000
		Rabbit, Lower Keys marsh	<i>Sylvilagus palustris hefneri</i>	Wherever found	\$25,000	\$0	\$25,000
		Seal, Hawaiian monk	<i>Monachus schauinslandi</i>	Wherever found	\$27,000	\$0	\$27,000
		Squirrel, Northern Idaho Ground	<i>Uroditellus brunneus</i>	Wherever found	\$2,100	\$0	\$2,100
		Wolf, gray	<i>Canis lupus</i>	MN	\$295,000	\$0	\$295,000
		Wolf, gray	<i>Canis lupus</i>	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$420,000	\$0	\$420,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, Mexican	Canis lupus baileyi	U.S.A. (portions of AZ and NM)see 17.84(k)	\$65,000	\$0	\$65,000
	<i>Mammals Subtotal</i>				\$2,053,400	\$0	\$2,053,400
	Birds	Coot, Hawaiian	Fulica americana alai	Wherever found	\$565,600	\$0	\$565,600
		Cuckoo, yellow-billed	Coccyzus americanus	Western U.S. DPS	\$13,400	\$0	\$13,400
		Duck, Hawaiian (=koloa)	Anas wyvilliana	Wherever found	\$565,600	\$0	\$565,600
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$81,700	\$0	\$81,700
		gallinule, Hawaiian common	Gallinula galeata sandvicensis	Wherever found	\$565,600	\$0	\$565,600
		Goose, Hawaiian	Branta (=Nesochen) sandvicensis	Wherever found	\$515,000	\$0	\$515,000
		Hawk, Hawaiian (=lo)	Buteo solitarius	Wherever found	\$72,600	\$0	\$72,600
		Knot, red	Calidris canutus rufa	Wherever found	\$100	\$0	\$100
		Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR, WA)	\$100	\$0	\$100
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$328,300	\$0	\$328,300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, piping	Charadrius melodus	[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$67,000	\$0	\$67,000
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS- U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$584,700	\$0	\$584,700
		Prairie-chicken, Attwater's greater	Tympanuchus cupido attwateri	Wherever found	\$10,500	\$0	\$10,500
		Rail, California clapper	Rallus longirostris obsoletus	Wherever found	\$204,300	\$0	\$204,300
		Rail, light-footed clapper	Rallus longirostris levipes	Wherever found	\$37,000	\$0	\$37,000
		Rail, Yuma Ridgways	Rallus obsoletus [=longirostris] yumanensis	Wherever found	\$12,500	\$0	\$12,500
		Sparrow, Florida grasshopper	Ammodramus savannarum floridanus	Wherever found	\$200	\$0	\$200
		Stilt, Hawaiian	Himantopus mexicanus knudseni	Wherever found	\$565,600	\$0	\$565,600
		Swiftlet, Mariana gray	Aerodramus vanikorensis bartschi	Wherever found	\$110,300	\$0	\$110,300
		Tern, California least	Sterna antillarum browni	Wherever found	\$372,700	\$0	\$372,700
		Tern, least	Sterna antillarum	interior pop.	\$57,000	\$0	\$57,000
		Tern, roseate	Sterna dougallii dougallii	Northeast U.S. nesting population	\$4,000	\$0	\$4,000



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Tern, roseate	<i>Sterna dougallii dougallii</i>	Western Hemisphere except NE U.S.	\$62,500	\$0	\$62,500
		Vireo, least Bell's	<i>Vireo bellii pusillus</i>	Wherever found	\$400	\$0	\$400
		Warbler, Kirtland's	<i>Setophaga kirtlandii</i> (= <i>Dendroica kirtlandii</i> )	Wherever found	\$33,000	\$0	\$33,000
	<i>Birds Subtotal</i>				<i>\$4,829,700</i>	<i>\$0</i>	<i>\$4,829,700</i>
	Reptiles	gartersnake, northern Mexican	<i>Thamnophis eques megalops</i>	Wherever found	\$12,500	\$0	\$12,500
		Massasauga (=rattlesnake), eastern	<i>Sistrurus catenatus</i>	Wherever found	\$8,000	\$0	\$8,000
		Sea turtle, green	<i>Chelonia mydas</i>	Central North Pacific DPS	\$27,000	\$0	\$27,000
		Sea turtle, green	<i>Chelonia mydas</i>	Central West Pacific DPS	\$186,600	\$0	\$186,600
		Sea turtle, green	<i>Chelonia mydas</i>	North Atlantic DPS	\$35,400	\$0	\$35,400
		Sea turtle, hawksbill	<i>Eretmochelys imbricata</i>	Wherever found	\$1,500	\$0	\$1,500
		Sea turtle, Kemp's ridley	<i>Lepidochelys kempii</i>	Wherever found	\$41,500	\$0	\$41,500
		Sea turtle, leatherback	<i>Dermochelys coriacea</i>	Wherever found	\$33,000	\$0	\$33,000
		Sea turtle, loggerhead	<i>Caretta caretta</i>	Northwest Atlantic Ocean DPS	\$48,000	\$0	\$48,000
		Snake, eastern indigo	<i>Drymarchon corais couperi</i>	Wherever found	\$15,000	\$0	\$15,000
		Snake, San Francisco garter	<i>Thamnophis sirtalis tetrataenia</i>	Wherever found	\$100	\$0	\$100
		Tortoise, desert	<i>Gopherus agassizii</i>	Wherever found, except AZ south and east of Colorado R., and Mexico	\$95,800	\$0	\$95,800

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		turtle, bog	Clemmys muhlenbergii	Wherever found, except GA, NC, SC, TN, VA	\$9,100	\$0	\$9,100
	<i>Reptiles Subtotal</i>				<i>\$513,500</i>	<i>\$0</i>	<i>\$513,500</i>
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$15,400	\$0	\$15,400
		Frog, Chiricahua leopard	Rana chiricahuensis	Wherever found	\$12,600	\$0	\$12,600
		Frog, mountain yellow-legged	Rana muscosa	Northern California DPS	\$100	\$0	\$100
		Frog, mountain yellow-legged	Rana muscosa	Southern California DPS	\$100	\$0	\$100
		Frog, Sierra Nevada Yellow-legged	Rana sierrae	Wherever found	\$200	\$0	\$200
		Salamander, California tiger	Ambystoma californiense	U.S.A. (CA - Santa Barbara County)	\$100	\$0	\$100
		Salamander, California tiger	Ambystoma californiense	U.S.A. (CA - Sonoma County)	\$100	\$0	\$100
		Salamander, desert slender	Batrachoseps aridus	Wherever found	\$200	\$0	\$200
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$300	\$0	\$300
		toad, Yosemite	Anaxyrus canorus	Wherever found	\$200	\$0	\$200
	<i>Amphibians Subtotal</i>				<i>\$29,300</i>	<i>\$0</i>	<i>\$29,300</i>
	Fishes	Chub, Chihuahua	Gila nigrescens	Wherever found	\$100	\$0	\$100
		Chub, Gila	Gila intermedia	Wherever found	\$100	\$0	\$100
		darter, bluemask	Etheostoma akatulo	Wherever found	\$1,300	\$0	\$1,300
		Darter, Niangua	Etheostoma nianguae	Wherever found	\$10,000	\$0	\$10,000
		Darter, Okaloosa	Etheostoma okaloosae	Wherever found	\$150,000	\$0	\$150,000
		Darter, watercress	Etheostoma nuchale	Wherever found	\$1,000	\$0	\$1,000
		Minnow, Devils River	Dionda diaboli	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pikeminnow (=squawfish), Colorado	Ptychocheilus lucius	Wherever found, except where listed as an experimental population	\$12,500	\$0	\$12,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$300	\$0	\$300
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$500	\$0	\$500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$17,500	\$0	\$17,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter-run ESU	\$400	\$0	\$400
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$17,500	\$0	\$17,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$17,500	\$0	\$17,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$466,000	\$0	\$466,000
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$14,000	\$0	\$14,000
		Salmon, chum	Oncorhynchus keta	Hood Canal summer-run ESU	\$14,000	\$0	\$14,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$500	\$0	\$500
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$466,000	\$0	\$466,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$17,500	\$0	\$17,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$17,900	\$0	\$17,900
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU	\$14,000	\$0	\$14,000
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$448,400	\$0	\$448,400
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$600	\$0	\$600
		Steelhead	Oncorhynchus (=Salmo) mykiss	Central California Coast DPS	\$2,500	\$0	\$2,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$126,000	\$0	\$126,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$126,000	\$0	\$126,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Northern California DPS	\$500	\$0	\$500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$126,000	\$0	\$126,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$400	\$0	\$400
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$400	\$0	\$400
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$126,000	\$0	\$126,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$17,500	\$0	\$17,500
		sturgeon, green	Acipenser medirostris	Southern DPS	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$100	\$0	\$100
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$12,700	\$0	\$12,700
		Trout, Apache	Oncorhynchus apache	Wherever found	\$1,000	\$0	\$1,000
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$1,200	\$0	\$1,200
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$100	\$0	\$100
	<i>Fishes Subtotal</i>				<i>\$2,228,200</i>	<i>\$0</i>	<i>\$2,228,200</i>
	Clams	Higgins eye (pearlymussel)	Lampsilis higginsii	Wherever found	\$100	\$0	\$100
		Hornshell, Texas	Popenaias popeii	Wherever found	\$100	\$0	\$100
		Mapleleaf, winged	Quadrula fragosa	Wherever found, except where listed as an experimental population	\$100	\$0	\$100
		Mussel, scaleshell	Leptodea leptodon	Wherever found	\$100	\$0	\$100
		Pearlshell, Louisiana	Margaritifera hembeli	Wherever found	\$13,500	\$0	\$13,500
		Pocketbook, Ouachita rock	Arkansia wheeleri	Wherever found	\$100	\$0	\$100
		Spinymussel, James	Pleurobema collina	Wherever found	\$1,700	\$0	\$1,700
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$600	\$0	\$600
	<i>Clams Subtotal</i>				<i>\$16,300</i>	<i>\$0</i>	<i>\$16,300</i>
	Snails	Cavesnail, Tumbling Creek	Antrobia culveri	Wherever found	\$15,000	\$0	\$15,000
		Limpet, Banbury Springs	Lanx sp.	Wherever found	\$100	\$0	\$100
		Snail, Bliss Rapids	Taylorconcha serpenticola	Wherever found	\$100	\$0	\$100
		Snail, Pecos assiminea	Assiminea pecos	Wherever found	\$100	\$0	\$100
		Snail, Snake River physa	Physa natricina	Wherever found	\$200	\$0	\$200
		Springsnail, Bruneau Hot	Pyrgulopsis bruneauensis	Wherever found	\$100	\$0	\$100
		Springsnail, Roswell	Pyrgulopsis roswellensis	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		springsnail, San Bernardino	Pyrgulopsis bernardina	Wherever found	\$100	\$0	\$100
		Springsnail, Socorro	Pyrgulopsis neomexicana	Wherever found	\$100	\$0	\$100
	<i>Snails Subtotal</i>				<i>\$15,900</i>	<i>\$0</i>	<i>\$15,900</i>
	Insects	Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$400	\$0	\$400
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$300	\$0	\$300
		Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$8,000	\$0	\$8,000
		Butterfly, Saint Francis' satyr	Neonympha mitchellii francisci	Wherever found	\$27,800	\$0	\$27,800
		Butterfly, Smith's blue	Euphilotes enoptes smithi	Wherever found	\$100	\$0	\$100
		Butterfly, Uncompahgre fritillary	Boloria acrocneuma	Wherever found	\$100	\$0	\$100
		Dragonfly, Hine's emerald	Somatochlora hineana	Wherever found	\$20,000	\$0	\$20,000
		Skipper, Carson wandering	Pseudocopaeodes eunus obscurus	Wherever found	\$100	\$0	\$100
		Skipper, Pawnee montane	Hesperia leonardus montana	Wherever found	\$100	\$0	\$100
		Tiger beetle, Salt Creek	Cicindela nevadica lincolniana	Wherever found	\$400	\$0	\$400
	<i>Insects Subtotal</i>				<i>\$57,300</i>	<i>\$0</i>	<i>\$57,300</i>
	Crustaceans	Crayfish, Shasta	Pacifastacus fortis	Wherever found	\$100	\$0	\$100
		Fairy shrimp, longhorn	Branchinecta longiantenna	Wherever found	\$100	\$0	\$100
		Fairy shrimp, San Diego	Branchinecta sandiegonensis	Wherever found	\$100	\$0	\$100
		Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$600	\$0	\$600
		Isopod, Lee County cave	Lirceus usdagalun	Wherever found	\$5,000	\$0	\$5,000
		Isopod, Socorro	Thermosphaeroma thermophilus	Wherever found	\$100	\$0	\$100
		Shrimp, California freshwater	Syncaris pacifica	Wherever found	\$100	\$0	\$100
		Tadpole shrimp, vernal pool	Lepidurus packardi	Wherever found	\$100	\$0	\$100
	<i>Crustaceans Subtotal</i>				<i>\$6,200</i>	<i>\$0</i>	<i>\$6,200</i>
	Flowering Plants	Allocarya, Calistoga	Plagiobothrys strictus	Wherever found	\$300	\$0	\$300
		amole, purple	Chlorogalum purpureum	Wherever found	\$200	\$0	\$200
		Baccharis, Encinitas	Baccharis vanessae	Wherever found	\$300	\$0	\$300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Barberry, Nevin's	<i>Berberis nevinii</i>	Wherever found	\$200	\$0	\$200
		Beargrass, Britton's	<i>Nolina brittoniana</i>	Wherever found	\$21,700	\$0	\$21,700
		Beauty, Harper's	<i>Harperocallis flava</i>	Wherever found	\$21,700	\$0	\$21,700
		Bedstraw, El Dorado	<i>Galium californicum</i> ssp. <i>sierrae</i>	Wherever found	\$200	\$0	\$200
		Bedstraw, island	<i>Galium buxifolium</i>	Wherever found	\$100	\$0	\$100
		Bird's beak, palmate-bracted	<i>Cordylanthus palmatus</i>	Wherever found	\$100	\$0	\$100
		Bird's-beak, Pennell's	<i>Cordylanthus tenuis</i> ssp. <i>capillaris</i>	Wherever found	\$300	\$0	\$300
		Bird's-beak, salt marsh	<i>Cordylanthus maritimus</i> ssp. <i>maritimus</i>	Wherever found	\$300	\$0	\$300
		bladderpod, Missouri	<i>Physaria filiformis</i>	Wherever found	\$5,000	\$0	\$5,000
		Bladderpod, San Bernardino Mountains	<i>Lesquerella kingii</i> ssp. <i>bernardina</i>	Wherever found	\$100	\$0	\$100
		Bluegrass, Napa	<i>Poa napensis</i>	Wherever found	\$200	\$0	\$200
		Bonamia, Florida	<i>Bonamia grandiflora</i>	Wherever found	\$21,700	\$0	\$21,700
		Brodiaea, Chinese Camp	<i>Brodiaea pallida</i>	Wherever found	\$300	\$0	\$300
		Buckwheat, cushenbury	<i>Eriogonum ovalifolium</i> var. <i>vineum</i>	Wherever found	\$300	\$0	\$300
		Buckwheat, Ione (incl. Irish Hill)	<i>Eriogonum apricum</i> (incl. var. <i>prostratum</i> )	Wherever found	\$200	\$0	\$200
		Bully, Everglades	<i>Sideroxylon reclinatum</i> ssp. <i>austrofloridense</i>	Wherever found	\$100	\$0	\$100
		Bulrush, Northeastern	<i>Scirpus ancistrochaetus</i>	Wherever found	\$9,000	\$0	\$9,000
		Butterweed, Layne's	<i>Senecio layneae</i>	Wherever found	\$200	\$0	\$200
		Cactus, Bakersfield	<i>Opuntia treleasei</i>	Wherever found	\$300	\$0	\$300
		Catchfly, Spalding's	<i>Silene spaldingii</i>	Wherever found	\$100	\$0	\$100
		Ceanothus, coyote	<i>Ceanothus ferrisae</i>	Wherever found	\$100	\$0	\$100
		Ceanothus, Pine Hill	<i>Ceanothus roderickii</i>	Wherever found	\$300	\$0	\$300
		Chaffseed, American	<i>Schwalbea americana</i>	Wherever found	\$27,800	\$0	\$27,800
		Checker-mallow, Keck's	<i>Sidalcea keckii</i>	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Checker-mallow, Kenwood Marsh	<i>Sidalcea oregana</i> ssp. <i>valida</i>	Wherever found	\$200	\$0	\$200
		Checker-mallow, pedate	<i>Sidalcea pedata</i>	Wherever found	\$300	\$0	\$300
		Clarkia, Pismo	<i>Clarkia speciosa</i> ssp. <i>immaculata</i>	Wherever found	\$300	\$0	\$300
		Clarkia, Presidio	<i>Clarkia franciscana</i>	Wherever found	\$200	\$0	\$200
		Clover, Monterey	<i>Trifolium trichocalyx</i>	Wherever found	\$400	\$0	\$400
		Crabgrass, Florida pineland	<i>Digitaria pauciflora</i>	Wherever found	\$100	\$0	\$100
		Crownbeard, big-leaved	<i>Verbesina dissita</i>	Wherever found	\$300	\$0	\$300
		Crownscale, San Jacinto Valley	<i>Atriplex coronata</i> var. <i>notatior</i>	Wherever found	\$200	\$0	\$200
		Daisy, Parish's	<i>Erigeron parishii</i>	Wherever found	\$200	\$0	\$200
		Dudleya, Conejo	<i>Dudleya abramsii</i> ssp. <i>parva</i>	Wherever found	\$300	\$0	\$300
		Dudleya, marcescent	<i>Dudleya cymosa</i> ssp. <i>marcescens</i>	Wherever found	\$400	\$0	\$400
		Dwarf-flax, Marin	<i>Hesperolinon congestum</i>	Wherever found	\$200	\$0	\$200
		Erubia	<i>Solanum drymophilum</i>	Wherever found	\$300	\$0	\$300
		Evening-primrose, Antioch Dunes	<i>Oenothera deltoides</i> ssp. <i>howellii</i>	Wherever found	\$200	\$0	\$200
		Fiddleneck, large-flowered	<i>Amsinckia grandiflora</i>	Wherever found	\$200	\$0	\$200
		Flannelbush, Mexican	<i>Fremontodendron mexicanum</i>	Wherever found	\$100	\$0	\$100
		Flannelbush, Pine Hill	<i>Fremontodendron californicum</i> ssp. <i>decumbens</i>	Wherever found	\$300	\$0	\$300
		Four-o'clock, MacFarlane's	<i>Mirabilis macfarlanei</i>	Wherever found	\$100	\$0	\$100
		Gilia, Hoffmann's slender-flowered	<i>Gilia tenuiflora</i> ssp. <i>hoffmannii</i>	Wherever found	\$200	\$0	\$200
		Gilia, Monterey	<i>Gilia tenuiflora</i> ssp. <i>arenaria</i>	Wherever found	\$400	\$0	\$400
		Goldfields, Burke's	<i>Lasthenia burkei</i>	Wherever found	\$300	\$0	\$300
		Goldfields, Contra Costa	<i>Lasthenia conjugens</i>	Wherever found	\$100	\$0	\$100
		Grass, Colusa	<i>Neostapfia colusana</i>	Wherever found	\$300	\$0	\$300
		haiwale	<i>Cyrtandra ferripilosa</i>	Wherever found	\$4,300	\$0	\$4,300



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Haiwale	<i>Cyrtandra gracilis</i>	Wherever found	\$4,300	\$0	\$4,300
		Haiwale	<i>Cyrtandra paliku</i>	Wherever found	\$4,300	\$0	\$4,300
		Haiwale	<i>Cyrtandra waiolani</i>	Wherever found	\$4,300	\$0	\$4,300
		Howellia, Water	<i>Howellia aquatilis</i>		\$100	\$0	\$100
		Jewelflower, California	<i>Caulanthus californicus</i>	Wherever found	\$100	\$0	\$100
		Jewelflower, Metcalf Canyon	<i>Streptanthus albidus</i> ssp. <i>albidus</i>	Wherever found	\$200	\$0	\$200
		Ladies'-tresses, Ute	<i>Spiranthes diluvialis</i>	Wherever found	\$100	\$0	\$100
		Larkspur, Baker's	<i>Delphinium bakeri</i>	Wherever found	\$100	\$0	\$100
		Layia, beach	<i>Layia carnosa</i>	Wherever found	\$100	\$0	\$100
		Lead-plant, Crenulate	<i>Amorpha crenulata</i>	Wherever found	\$21,700	\$0	\$21,700
		Lily, Pitkin Marsh	<i>Lilium pardalinum</i> ssp. <i>pitkinense</i>	Wherever found	\$300	\$0	\$300
		Liveforever, Laguna Beach	<i>Dudleya stolonifera</i>	Wherever found	\$200	\$0	\$200
		Loosestrife, rough-leaved	<i>Lysimachia asperulaefolia</i>	Wherever found	\$27,800	\$0	\$27,800
		Lupine, clover	<i>Lupinus tidestromii</i>	Wherever found	\$100	\$0	\$100
		Lupine, Nipomo Mesa	<i>Lupinus nipomensis</i>	Wherever found	\$300	\$0	\$300
		Malacothrix, island	<i>Malacothrix squalida</i>	Wherever found	\$100	\$0	\$100
		Mallow, Kern	<i>Eremalche kernensis</i>	Wherever found	\$200	\$0	\$200
		Manzanita, Del Mar	<i>Arctostaphylos glandulosa</i> ssp. <i>crassifolia</i>	Wherever found	\$200	\$0	\$200
		Manzanita, Franciscan	<i>Arctostaphylos franciscana</i>	Wherever found	\$200	\$0	\$200
		Manzanita, Ione	<i>Arctostaphylos myrtifolia</i>	Wherever found	\$200	\$0	\$200
		Manzanita, Morro	<i>Arctostaphylos morroensis</i>	Wherever found	\$300	\$0	\$300
		Manzanita, pallid	<i>Arctostaphylos pallida</i>	Wherever found	\$200	\$0	\$200
		Manzanita, Presidio	<i>Arctostaphylos hookeri</i> var. <i>ravenii</i>	Wherever found	\$200	\$0	\$200
		Meadowfoam, Butte County	<i>Limnanthes floccosa</i> ssp. <i>californica</i>	Wherever found	\$100	\$0	\$100
		Mesa-mint, Otay	<i>Pogogyne nudiuscula</i>	Wherever found	\$100	\$0	\$100
		Milk-vetch, Braunton's	<i>Astragalus brauntonii</i>	Wherever found	\$300	\$0	\$300
		Milk-vetch, Clara Hunt's	<i>Astragalus clarianus</i>	Wherever found	\$400	\$0	\$400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Milk-vetch, Coachella Valley	<i>Astragalus lentiginosus</i> var. <i>coachellae</i>	Wherever found	\$100	\$0	\$100
		Milk-vetch, coastal dunes	<i>Astragalus tener</i> var. <i>titi</i>	Wherever found	\$300	\$0	\$300
		Milk-vetch, Cushenbury	<i>Astragalus albens</i>	Wherever found	\$300	\$0	\$300
		Milk-vetch, Lane Mountain	<i>Astragalus jaegerianus</i>	Wherever found	\$200	\$0	\$200
		Milk-vetch, Peirson's	<i>Astragalus magdalenae</i> var. <i>peirsonii</i>	Wherever found	\$200	\$0	\$200
		Milkweed, Mead's	<i>Asclepias meadii</i>	Wherever found	\$20,000	\$0	\$20,000
		Mint, scrub	<i>Dicerandra frutescens</i>	Wherever found	\$21,700	\$0	\$21,700
		Mountain-mahogany, Catalina Island	<i>Cercocarpus traskiae</i>	Wherever found	\$100	\$0	\$100
		mountainbalm, Indian Knob	<i>Eriodictyon altissimum</i>	Wherever found	\$200	\$0	\$200
		Navarretia, few-flowered	<i>Navarretia leucocephala</i> ssp. <i>pauciflora</i> (=N. <i>pauciflora</i> )	Wherever found	\$200	\$0	\$200
		Navarretia, many-flowered	<i>Navarretia leucocephala</i> ssp. <i>plieantha</i>	Wherever found	\$300	\$0	\$300
		Niterwort, Amargosa	<i>Nitrophila mohavensis</i>	Wherever found	\$200	\$0	\$200
		No common name	<i>Dendrobium guamense</i>	Wherever found	\$300	\$0	\$300
		No common name	<i>Eugenia bryanii</i>	Wherever found	\$300	\$0	\$300
		No common name	<i>Eugenia woodburyana</i>	Wherever found	\$300	\$0	\$300
		No common name	<i>Geocarpon minimum</i>	Wherever found	\$10,000	\$0	\$10,000
		No common name	<i>Sicyos lanceoloideus</i>	Wherever found	\$1,000	\$0	\$1,000
		Onion, Munz's	<i>Allium munzii</i>	Wherever found	\$100	\$0	\$100
		Orchid, eastern prairie fringed	<i>Platanthera leucophaea</i>	Wherever found	\$1,200	\$0	\$1,200
		Orcutt grass, California	<i>Orcuttia californica</i>	Wherever found	\$300	\$0	\$300
		Orcutt grass, hairy	<i>Orcuttia pilosa</i>	Wherever found	\$600	\$0	\$600
		Orcutt grass, San Joaquin	<i>Orcuttia inaequalis</i>	Wherever found	\$300	\$0	\$300
		Owl's-clover, fleshy	<i>Castilleja campestris</i> ssp. <i>succulenta</i>	Wherever found	\$300	\$0	\$300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		oxythecha, Cushenbury	Oxythecha parishii var. goodmaniana	Wherever found	\$300	\$0	\$300
		Paintbrush, ash-grey	Castilleja cinerea	Wherever found	\$300	\$0	\$300
		Pentachaeta, Lyon's	Pentachaeta lyonii	Wherever found	\$100	\$0	\$100
		Peppergrass, Slickspot	Lepidium papilliferum		\$100	\$0	\$100
		Phacelia, island	Phacelia insularis ssp. insularis	Wherever found	\$100	\$0	\$100
		Pinkroot, gentian	Spigelia gentianoides	Wherever found	\$21,700	\$0	\$21,700
		Plum, scrub	Prunus geniculata	Wherever found	\$21,700	\$0	\$21,700
		Pogonia, small whorled	Isotria medeoloides	Wherever found	\$500	\$0	\$500
		Potentilla, Hickman's	Potentilla hickmanii	Wherever found	\$200	\$0	\$200
		Prairie-clover, Florida	Dalea carthagenensis floridana	Wherever found	\$100	\$0	\$100
		Pussypaws, Mariposa	Calyptridium pulchellum	Wherever found	\$200	\$0	\$200
		Rhododendron, Chapman	Rhododendron chapmanii	Wherever found	\$21,700	\$0	\$21,700
		Rock-cress, Hoffmann's	Arabis hoffmannii	Wherever found	\$100	\$0	\$100
		Rock-cress, McDonald's	Arabis macdonaldiana	Wherever found	\$400	\$0	\$400
		Rosemary, Apalachicola	Conradina glabra	Wherever found	\$21,700	\$0	\$21,700
		Rush-rose, island	Helianthemum greenei	Wherever found	\$100	\$0	\$100
		Sandlace	Polygonella myriophylla	Wherever found	\$21,700	\$0	\$21,700
		Sandmat, pineland	Chamaesyce deltoidea pinetorum	Wherever found	\$100	\$0	\$100
		Sandwort, Bear Valley	Arenaria ursina	Wherever found	\$100	\$0	\$100
		Sandwort, Marsh	Arenaria paludicola	Wherever found	\$300	\$0	\$300
		Seablite, California	Suaeda californica		\$300	\$0	\$300
		Snakeroot	Eryngium cuneifolium	Wherever found	\$21,700	\$0	\$21,700
		Spineflower, Ben Lomond	Chorizanthe pungens var. hartwegiana	Wherever found	\$100	\$0	\$100
		Spineflower, Howell's	Chorizanthe howellii	Wherever found	\$200	\$0	\$200
		Spineflower, Monterey	Chorizanthe pungens var. pungens	Wherever found	\$200	\$0	\$200
		Spineflower, Orcutt's	Chorizanthe orcuttiana	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		spineflower, Robust	Chorizanthe robusta var. robusta	Wherever found	\$200	\$0	\$200
		Spiraea, Virginia	Spiraea virginiana	Wherever found	\$1,300	\$0	\$1,300
		Spurge, Hoover's	Chamaesyce hooveri	Wherever found	\$300	\$0	\$300
		Stonecrop, Lake County	Parvisedum leiocarpum	Wherever found	\$200	\$0	\$200
		Sumac, Michaux's	Rhus michauxii	Wherever found	\$27,800	\$0	\$27,800
		Sunburst, Hartweg's golden	Pseudobahia bahiifolia	Wherever found	\$200	\$0	\$200
		Taraxacum, California	Taraxacum californicum	Wherever found	\$100	\$0	\$100
		Tarplant, Gaviota	Deinandra increscens ssp. villosa	Wherever found	\$200	\$0	\$200
		Tarplant, Otay	Deinandra (=Hemizonia) conjugens	Wherever found	\$100	\$0	\$100
		Thistle, Chorro Creek bog	Cirsium fontinale var. obispoense	Wherever found	\$300	\$0	\$300
		Thistle, fountain	Cirsium fontinale var. fontinale	Wherever found	\$200	\$0	\$200
		Thistle, La Graciosa	Cirsium loncholepis	Wherever found	\$200	\$0	\$200
		Thistle, Loch Lomond coyote	Eryngium constancei	Wherever found	\$200	\$0	\$200
		Thistle, Pitcher's	Cirsium pitcheri	Wherever found	\$8,000	\$0	\$8,000
		Tuctoria, Greene's	Tuctoria greenei	Wherever found	\$300	\$0	\$300
		Uvillo	Eugenia haematocarpa	Wherever found	\$300	\$0	\$300
		Vervain, Red Hills	Verbena californica	Wherever found	\$300	\$0	\$300
		Wallflower, Ben Lomond	Erysimum teretifolium	Wherever found	\$100	\$0	\$100
		Wallflower, Contra Costa	Erysimum capitatum var. angustatum	Wherever found	\$100	\$0	\$100
		Wallflower, Menzies'	Erysimum menziesii	Wherever found	\$200	\$0	\$200
		Watercress, Gambel's	Rorippa gambellii	Wherever found	\$200	\$0	\$200
		Wireweed	Polygonella basiramia	Wherever found	\$21,700	\$0	\$21,700
		Yerba santa, Lompoc	Eriodictyon capitatum	Wherever found	\$200	\$0	\$200
		<i>Flowering Plants Subtotal</i>			<i>\$443,000</i>	<i>\$0</i>	<i>\$443,000</i>

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Conifers and Cycads	Torreya, Florida	Torreya taxifolia	Wherever found	\$21,700	\$0	\$21,700
	<i>Conifers and Cycads Subtotal</i>				\$21,700	\$0	\$21,700
	Lichens	Cladonia, Florida perforate	Cladonia perforata	Wherever found	\$21,700	\$0	\$21,700
	<i>Lichens Subtotal</i>				\$21,700	\$0	\$21,700
<i>Domestic Total</i>					\$10,236,200	\$0	\$10,236,200
<i>Multi-Species Subtotal</i>		<i>Multi-Species</i>			\$365,480	\$0	\$365,480
Foreign	Reptiles	Iguana, Cuban ground	Cyclura nubila nubila	(Cyclura nubila nubila)	\$33,800	\$0	\$33,800
<i>Foreign Subtotal</i>					\$33,800	\$0	\$33,800
<b>Total</b>					<b>\$10,635,480</b>	<b>\$0</b>	<b>\$10,635,480</b>

## USDA Forest Service - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	<i>Myotis grisescens</i>	Wherever found	\$60,000	\$0	\$60,000
		Bat, Hawaiian hoary	<i>Lasiurus cinereus semotus</i>	Wherever found	\$2,000	\$0	\$2,000
		Bat, Indiana	<i>Myotis sodalis</i>	Wherever found	\$1,198,458	\$0	\$1,198,458
		Bat, Northern long-eared	<i>Myotis septentrionalis</i>	Wherever found	\$176,770	\$0	\$176,770
		Bear, grizzly	<i>Ursus arctos horribilis</i>	Bitterroot Grizzly Bear Nonessential Experimental Population Area	\$40,000	\$0	\$40,000
		Bear, grizzly	<i>Ursus arctos horribilis</i>	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$417,101	\$0	\$417,101
		Ferret, black-footed	<i>Mustela nigripes</i>	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$111,164	\$0	\$111,164
		Jaguar	<i>Panthera onca</i>	Wherever found	\$4,000	\$0	\$4,000
		Lynx, Canada	<i>Lynx canadensis</i>	Wherever Found in Contiguous U.S.	\$856,055	\$0	\$856,055
		Mouse, Preble's meadow jumping	<i>Zapus hudsonius preblei</i>	wherever found	\$32,035	\$0	\$32,035
		Ocelot	<i>Leopardus (=Felis) pardalis</i>	wherever found	\$4,000	\$0	\$4,000
		Panther, Florida	<i>Puma (=Felis) concolor coryi</i>	Wherever found	\$5,000	\$0	\$5,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Prairie dog, Utah	<i>Cynomys parvidens</i>	Wherever found	\$320,000	\$0	\$320,000
		Puma (=mountain lion)	<i>Puma (=Felis) concolor</i> (all subsp. except <i>coryi</i> )	FL	\$34,370	\$0	\$34,370
		Sheep, Sierra Nevada bighorn	<i>Ovis canadensis sierrae</i>	Sierra Nevada	\$8,280	\$0	\$8,280
		Wolf, gray	<i>Canis lupus</i>	MN	\$893,432	\$0	\$893,432
		Wolf, gray	<i>Canis lupus</i>	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$153,233	\$0	\$153,233
	<i>Mammals Total</i>				<i>\$4,315,898</i>	<i>\$0</i>	<i>\$4,315,898</i>
	Birds	akepa, Hawaii	<i>Loxops coccineus</i>	Wherever found	\$20,000	\$0	\$20,000
		akiapolaau	<i>Hemignathus wilsoni</i>	Wherever found	\$20,000	\$0	\$20,000
		Akikiki	<i>Oreomystis bairdi</i>	Wherever found	\$10,000	\$0	\$10,000
		Creeper, Hawaii	<i>Oreomystis mana</i>	Wherever found	\$20,000	\$0	\$20,000
		Crow, Hawaiian (=alala)	<i>Corvus hawaiiensis</i>	Wherever found	\$4,000	\$0	\$4,000
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$16,000	\$0	\$16,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$136,120	\$0	\$136,120
		Hawk, Hawaiian (=lo)	Buteo solitarius	Wherever found	\$10,000	\$0	\$10,000
		Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR, WA)	\$544,209	\$0	\$544,209
		Owl, northern spotted	Strix occidentalis caurina	Wherever found	\$36,494,350	\$0	\$36,494,350
		Palila (honeycreeper)	Loxioides bailleui	Wherever found	\$10,000	\$0	\$10,000
		Parrot, Puerto Rican	Amazona vittata	Wherever found	\$130,000	\$0	\$130,000
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$153,014	\$0	\$153,014
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$138,511	\$0	\$138,511
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$514,345	\$0	\$514,345
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$15,410	\$0	\$15,410
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$45,000	\$0	\$45,000
		Warbler (=wood), golden-cheeked	Dendroica chrysoparia	Wherever found	\$70,000	\$0	\$70,000
		Warbler, Kirtland's	Setophaga kirtlandii (=Dendroica kirtlandii)	Wherever found	\$228,100	\$0	\$228,100
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$5,697,654	\$0	\$5,697,654
	<i>Birds Total</i>				<i>\$44,276,713</i>	<i>\$0</i>	<i>\$44,276,713</i>
	Reptiles	Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$49,798	\$0	\$49,798
		Pinesnake, Louisiana	Pituophis ruthveni	Wherever found	\$745,908	\$0	\$745,908
		Skink, sand	Neoseps reynoldsi	Wherever found	\$49,249	\$0	\$49,249
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$257,602	\$0	\$257,602



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$98,635	\$0	\$98,635
		Tortoise, gopher	Gopherus polyphemus	West of Mobile and Tombigbee Rivers	\$2,037,150	\$0	\$2,037,150
		<i>Reptiles Total</i>			<i>\$3,238,342</i>	<i>\$0</i>	<i>\$3,238,342</i>
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$47,425	\$0	\$47,425
		Frog, Chiricahua leopard	Rana chiricahuensis	Wherever found	\$5,600	\$0	\$5,600
		Frog, dusky gopher	Rana sevosa	Wherever found	\$261,875	\$0	\$261,875
		Frog, mountain yellow-legged	Rana muscosa	Southern California DPS	\$2,327	\$0	\$2,327
		Frog, Oregon spotted	Rana pretiosa	Wherever found	\$111,104	\$0	\$111,104
		Frog, Sierra Nevada Yellow-legged	Rana sierrae	Wherever found	\$110,921	\$0	\$110,921
		Salamander, Cheat Mountain	Plethodon nettingi	Wherever found	\$16,000	\$0	\$16,000
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$165,480	\$0	\$165,480
		Toad, Houston	Bufo houstonensis	Wherever found	\$20,000	\$0	\$20,000
		toad, Yosemite	Anaxyrus canorus	Wherever found	\$36,000	\$0	\$36,000
		<i>Amphibians Total</i>			<i>\$776,732</i>	<i>\$0</i>	<i>\$776,732</i>
	Fishes	Catfish, Yaqui	Ictalurus pricei	Wherever found	\$16,000	\$0	\$16,000
		Dace, blackside	Phoxinus cumberlandensis	Wherever found	\$1,285	\$0	\$1,285
		Minnow, loach	Tiaroga cobitis	Wherever found	\$46,000	\$0	\$46,000
		Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$20,000	\$0	\$20,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$384,435	\$0	\$384,435
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$10,000	\$0	\$10,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$834,805	\$0	\$834,805
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$187,200	\$0	\$187,200
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$750	\$0	\$750
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$27,000	\$0	\$27,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$570,085	\$0	\$570,085
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$856,771	\$0	\$856,771
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$173,842	\$0	\$173,842
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$10,000	\$0	\$10,000
		Spikedace	Meda fulgida	Wherever found	\$46,000	\$0	\$46,000
		Spinedace, Little Colorado	Lepidomeda vittata	Wherever found	\$200	\$0	\$200
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$486,830	\$0	\$486,830
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$322,058	\$0	\$322,058
		Steelhead	Oncorhynchus (=Salmo) mykiss	Northern California DPS	\$60,000	\$0	\$60,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$245,555	\$0	\$245,555

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$900	\$0	\$900
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$509,898	\$0	\$509,898
		Stickleback, unarmored threespine	Gasterosteus aculeatus williamsoni	Wherever found	\$500	\$0	\$500
		Sturgeon, Gulf	Acipenser oxyrinchus (=oxyrinchus) desotoi	Wherever found	\$1,000	\$0	\$1,000
		Sucker, Lost River	Deltistes luxatus	Wherever found	\$10,000	\$0	\$10,000
		Sucker, shortnose	Chasmistes brevirostris	Wherever found	\$118,000	\$0	\$118,000
		Sucker, Warner	Catostomus warnerensis	Wherever found	\$10,000	\$0	\$10,000
		Sucker, Zuni bluehead	Catostomus discobolus yarrowi	Wherever found	\$16,000	\$0	\$16,000
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	Wherever found	\$18,000	\$0	\$18,000
		Trout, Apache	Oncorhynchus apache	Wherever found	\$25,000	\$0	\$25,000
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$1,777,148	\$0	\$1,777,148
		Trout, Gila	Oncorhynchus gilae	Wherever found	\$44,050	\$0	\$44,050
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$565,950	\$0	\$565,950
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$23,660	\$0	\$23,660
		Trout, Little Kern golden	Oncorhynchus aguabonita whitei	Wherever found	\$37,813	\$0	\$37,813
		Trout, Paiute cutthroat	Oncorhynchus clarkii seleniris	Wherever found	\$12,000	\$0	\$12,000
		<i>Fishes Total</i>			<i>\$7,468,735</i>	<i>\$0</i>	<i>\$7,468,735</i>
	Clams	Elktoe, Cumberland	Alasmidonta atropurpurea	Wherever found	\$1,285	\$0	\$1,285
		Pearlymussel, littlewing	Pegias fabula	Wherever found	\$30,000	\$0	\$30,000
		Pocketbook, finlined	Lampsilis altilis	Wherever found	\$4,000	\$0	\$4,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$100	\$0	\$100
	<i>Clams Total</i>				\$35,385	\$0	\$35,385
	Snails	Snail, Snake River physa	Physa natricina	Wherever found	\$4,000	\$0	\$4,000
	<i>Snails Subtotal</i>				\$4,000	\$0	\$4,000
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$77,000	\$0	\$77,000
		Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$232,248	\$0	\$232,248
		Checkerspot, Taylor's (=whulge)	Euphydryas editha taylori	Wherever found	\$4,500	\$0	\$4,500
		Dragonfly, Hine's emerald	Somatochlora hineana	Wherever found	\$100	\$0	\$100
		Moth, Blackburn's sphinx	Manduca blackburni	Wherever found	\$6,000	\$0	\$6,000
		Skipper, Dakota	Hesperia dacotae	Wherever found	\$40,843	\$0	\$40,843
		Skipper, Pawnee montane	Hesperia leonardus montana	Wherever found	\$10	\$0	\$10
	<i>Insects Total</i>				\$360,701	\$0	\$360,701
	Flowering Plants	Aster, Ruth's golden	Pityopsis ruthii	Wherever found	\$2,000	\$0	\$2,000
		Avens, spreading	Geum radiatum	Wherever found	\$11,000	\$0	\$11,000
		Beauty, Harper's	Harperocallis flava	Wherever found	\$8,000	\$0	\$8,000
		Birds-in-a-nest, white	Macbridea alba	Wherever found	\$1,100	\$0	\$1,100
		Blazingstar, Heller's	Liatris helleri	Wherever found	\$1,700	\$0	\$1,700
		Bluet, Roan Mountain	Hedyotis purpurea var. montana	Wherever found	\$2,700	\$0	\$2,700
		Buckwheat, scrub	Eriogonum longifolium var. gnaphalifolium	Wherever found	\$26,426	\$0	\$26,426
		Butterweed, Layne's	Senecio layneae	Wherever found	\$5,800	\$0	\$5,800
		Butterwort, Godfrey's	Pinguicula ionantha	Wherever found	\$5,600	\$0	\$5,600
		Cactus, Arizona hedgehog	Echinocereus triglochidiatus var. arizonicus	Wherever found	\$3,200	\$0	\$3,200
		Cactus, Fickeisen plains	Pediocactus peeblesianus fickeiseniae	Wherever found	\$6,000	\$0	\$6,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Cactus, Kuenzler hedgehog	Echinocereus fendleri var. kuenzleri	Wherever found	\$16,886	\$0	\$16,886
		Cactus, Pima pineapple	Coryphantha scheeri var. robustispina	Wherever found	\$5,000	\$0	\$5,000
		Catchfly, Spalding's	Silene spaldingii	Wherever found	\$32,000	\$0	\$32,000
		Chaffseed, American	Schwalbea americana	Wherever found	\$70,000	\$0	\$70,000
		Checkermallow, Wenatchee Mountains	Sidalcea oregana var. calva	Wherever found	\$6,000	\$0	\$6,000
		Clover, running buffalo	Trifolium stoloniferum		\$10,000	\$0	\$10,000
		Coneflower, smooth	Echinacea laevigata	Wherever found	\$141,000	\$0	\$141,000
		Dropwort, Canby's	Oxypolis canbyi	Wherever found	\$5,000	\$0	\$5,000
		Flannelbush, Mexican	Fremontodendron mexicanum	Wherever found	\$14,000	\$0	\$14,000
		Fritillary, Gentner's	Fritillaria gentneri	Wherever found	\$10,000	\$0	\$10,000
		Gooseberry, Miccosukee	Ribes echinellum	Wherever found	\$10,000	\$0	\$10,000
		Hala pepe	Pleomele hawaiiensis	Wherever found	\$10,000	\$0	\$10,000
		Heartleaf, dwarf-flowered	Hexastylis naniflora	Wherever found	\$1,000	\$0	\$1,000
		Heather, mountain golden	Hudsonia montana	Wherever found	\$26,400	\$0	\$26,400
		Honohono	Haplostachys haplostachya	Wherever found	\$30,000	\$0	\$30,000
		Howellia, Water	Howellia aquatilis		\$5,744	\$0	\$5,744
		Ipomopsis, Holy Ghost	Ipomopsis sancti-spiritus	Wherever found	\$40,000	\$0	\$40,000
		Kauila	Colubrina oppositifolia	Wherever found	\$10,000	\$0	\$10,000
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$2,000	\$0	\$2,000
		Loosestrife, rough-leaved	Lysimachia asperulaefolia	Wherever found	\$22,200	\$0	\$22,200
		Lupine, Kincaid's	Lupinus sulphureus ssp. kincaidii	Wherever found	\$12,000	\$0	\$12,000
		Milk-vetch, heliotrope	Astragalus montii	Wherever found	\$4,000	\$0	\$4,000
		No common name	Neraudia ovata	Wherever found	\$10,000	\$0	\$10,000
		No common name	Silene hawaiiensis	Wherever found	\$10,000	\$0	\$10,000
		No common name	Silene lanceolata	Wherever found	\$20,000	\$0	\$20,000
		No common name	Spermolepis hawaiiensis	Wherever found	\$10,000	\$0	\$10,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		No common name	<i>Stenogyne angustifolia</i> var. <i>angustifolia</i>	Wherever found	\$20,000	\$0	\$20,000
		No common name	<i>Tetramolopium arenarium</i>	Wherever found	\$20,000	\$0	\$20,000
		Orchid, western prairie fringed	<i>Platanthera praeclara</i>	Wherever found	\$12,000	\$0	\$12,000
		Orchid, white fringeless	<i>Platanthera integrilabia</i>		\$11,000	\$0	\$11,000
		Orcutt grass, slender	<i>Orcuttia tenuis</i>	Wherever found	\$3,600	\$0	\$3,600
		Penstemon, blowout	<i>Penstemon haydenii</i>	Wherever found	\$91,500	\$0	\$91,500
		Peppergrass, Slickspot	<i>Lepidium papilliferum</i>		\$54,000	\$0	\$54,000
		Phacelia, clay	<i>Phacelia argillacea</i>	Wherever found	\$20,000	\$0	\$20,000
		Phlox, Yreka	<i>Phlox hirsuta</i>	Wherever found	\$2,000	\$0	\$2,000
		Pigeon wings	<i>Clitoria fragrans</i>	Wherever found	\$30,365	\$0	\$30,365
		Pink, swamp	<i>Helonias bullata</i>	Wherever found	\$6,600	\$0	\$6,600
		Pogonia, small whorled	<i>Isotria medeoloides</i>	Wherever found	\$15,300	\$0	\$15,300
		Polygala, Lewton's	<i>Polygala lewtonii</i>	Wherever found	\$26,426	\$0	\$26,426
		Pondberry	<i>Lindera melissifolia</i>	Wherever found	\$1,150,000	\$0	\$1,150,000
		Poppy, Sacramento prickly	<i>Argemone pleiakantha</i> ssp. <i>pinnatisecta</i>	Wherever found	\$18,550	\$0	\$18,550
		Po`e	<i>Portulaca sclerocarpa</i>	Wherever found	\$10,000	\$0	\$10,000
		Primrose, Maguire	<i>Primula maguirei</i>	Wherever found	\$2,000	\$0	\$2,000
		Ragwort, San Francisco Peaks	<i>Packera franciscana</i>	Wherever found	\$2,400	\$0	\$2,400
		Rock-cress, McDonald's	<i>Arabis macdonaldiana</i>	Wherever found	\$10,000	\$0	\$10,000
		Rose-mallow, Neches River	<i>Hibiscus dasycalyx</i>	Wherever found	\$7,000	\$0	\$7,000
		Sandwort, Cumberland	<i>Arenaria cumberlandensis</i>	Wherever found	\$1,600	\$0	\$1,600
		Stickseed, showy	<i>Hackelia venusta</i>	Wherever found	\$6,000	\$0	\$6,000
		Sunflower, Schweinitz's	<i>Helianthus schweinitzii</i>	Wherever found	\$23,400	\$0	\$23,400
		Thornmint, San Diego	<i>Acanthomintha ilicifolia</i>	Wherever found	\$10,000	\$0	\$10,000
		Tuctoria, Greene's	<i>Tuctoria greenei</i>	Wherever found	\$800	\$0	\$800
		Uhi uhi	<i>Mezoneuron kavaiense</i>	Wherever found	\$10,000	\$0	\$10,000
		Water-umbel, Huachuca	<i>Lilaeopsis schaffneriana</i> var. <i>recurva</i>	Wherever found	\$4,000	\$0	\$4,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		'Aiea	Nothocestrum breviflorum	Wherever found	\$10,000	\$0	\$10,000
	<i>Flowering Plants Total</i>				\$2,185,297	\$0	\$2,185,297
	Conifers and Cycads	Torreya, Florida	Torreya taxifolia	Wherever found	\$50,000	\$0	\$50,000
	<i>Conifers and Cycads Subtotal</i>				\$50,000	\$0	\$50,000
	Ferns and Allies	Quillwort, Louisiana	Isoetes louisianensis	Wherever found	\$34,264	\$0	\$34,264
	<i>Ferns and Allies Subtotal</i>				\$34,264	\$0	\$34,264
<i>Domestic Total</i>					\$62,746,067	\$0	\$62,746,067
<b>Total</b>					<b>\$62,746,067</b>	<b>\$0</b>	<b>\$62,746,067</b>

## USDA Natural Resources Conservation Service - FY 2018

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	bat, Florida bonneted	Eumops floridanus	Wherever found	\$8,000	\$250,000	\$258,000
		Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$27,300	\$0	\$27,300
		Bat, Indiana	Myotis sodalis	Wherever found	\$32,050	\$0	\$32,050
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$1,755,050	\$11,947,000	\$13,702,050
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population	\$24,500	\$0	\$24,500
		Ferret, black-footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$7,700	\$0	\$7,700
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$12,500	\$0	\$12,500
		Manatee, West Indian	Trichechus manatus	Wherever found	\$100,000	\$600,000	\$700,000
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	wherever found	\$3,000	\$0	\$3,000
		Pocket gopher, Olympia	Thomomys mazama pugetensis	Wherever found	\$2,000	\$0	\$2,000
		Pocket gopher, Tenino	Thomomys mazama tumuli	Wherever found	\$130,000	\$0	\$130,000
		Pocket gopher, Yelm	Thomomys mazama yelmensis	Wherever found	\$75,000	\$0	\$75,000
		Prairie dog, Utah	Cynomys parvidens	Wherever found	\$9,000	\$0	\$9,000
		Puma (=mountain lion)	Puma (=Felis) concolor (all subsp. except coryi)	FL	\$400,000	\$4,000,000	\$4,400,000
	<i>Mammals Subtotal</i>				<i>\$2,586,100</i>	<i>\$16,797,000</i>	<i>\$19,383,100</i>



Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Birds	akepa, Maui	<i>Loxops ochraceus</i>	Wherever found	\$81,400	\$0	\$81,400
		akiapolaau	<i>Hemignathus wilsoni</i>	Wherever found	\$81,400	\$0	\$81,400
		Caracara, Audubon's crested	<i>Polyborus plancus audubonii</i>	FL pop.	\$500,000	\$1,500,000	\$2,000,000
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$13,600	\$0	\$13,600
		Creeper, Hawaii	<i>Oreomystis mana</i>	Wherever found	\$81,400	\$0	\$81,400
		Crow, Hawaiian (= 'alala)	<i>Corvus hawaiiensis</i>	Wherever found	\$81,400	\$0	\$81,400
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$1,900	\$0	\$1,900
		Curlew, Eskimo	<i>Numenius borealis</i>	Wherever found	\$100	\$0	\$100
		Eider, spectacled	<i>Somateria fischeri</i>	Wherever found	\$2,930,000	\$0	\$2,930,000
		Eider, Steller's	<i>Polysticta stelleri</i>	AK breeding pop.	\$2,930,000	\$0	\$2,930,000
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$106,580	\$0	\$106,580
		Goose, Hawaiian	<i>Branta</i> (=Nesochen) <i>sandvicensis</i>	Wherever found	\$124,400	\$0	\$124,400
		Hawk, Hawaiian (= 'lo)	<i>Buteo solitarius</i>	Wherever found	\$115,600	\$0	\$115,600
		Kite, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	Wherever found	\$150,000	\$2,000,000	\$2,150,000
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$4,100	\$0	\$4,100
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$92,300	\$0	\$92,300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$5,600	\$0	\$5,600
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$33,600	\$0	\$33,600
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$40,000	\$0	\$40,000
		Sparrow, Florida grasshopper	Ammodramus savannarum floridanus	Wherever found	\$50,000	\$1,000,000	\$1,050,000
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$600,000	\$4,000,000	\$4,600,000
		Tern, least	Sterna antillarum	interior pop.	\$3,900	\$0	\$3,900
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$2,694,000	\$0	\$2,694,000
		<i>Birds Subtotal</i>			<i>\$10,721,280</i>	<i>\$8,500,000</i>	<i>\$19,221,280</i>
	Reptiles	Pinesnake, Louisiana	Pituophis ruthveni	Wherever found	\$496,000	\$0	\$496,000
		Skink, bluetail mole	Eumeces egregius lividus	Wherever found	\$50,000	\$0	\$50,000
		Skink, sand	Neoseps reynoldsi	Wherever found	\$50,000	\$0	\$50,000
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$300,000	\$2,000,000	\$2,300,000
		Tortoise, gopher	Gopherus polyphemus	West of Mobile and Tombigbee Rivers	\$316,600	\$0	\$316,600
		turtle, bog	Clemmys muhlenbergii	Wherever found, except GA, NC, SC, TN, VA	\$156,347	\$22,000	\$178,347
		<i>Reptiles Subtotal</i>			<i>\$1,368,947</i>	<i>\$2,022,000</i>	<i>\$3,390,947</i>
	Amphibians	Frog, Oregon spotted	Rana pretiosa	Wherever found	\$31,000	\$0	\$31,000
		Hellbender, Ozark	Cryptobranchus alleganiensis bishopi	Wherever found	\$4,000	\$0	\$4,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salamander, frosted flatwoods	Ambystoma cingulatum	Wherever found	\$175,000	\$0	\$175,000
		salamander, Reticulated flatwoods	Ambystoma bishopi	Wherever found	\$80,000	\$0	\$80,000
		Toad, Wyoming	Bufo hemiophrys baxteri	Wherever found	\$500	\$0	\$500
	<i>Amphibians Subtotal</i>				<i>\$290,500</i>	<i>\$0</i>	<i>\$290,500</i>
	Fishes	Chub, Virgin River	Gila seminuda (=robusta)	Wherever found	\$89,000	\$0	\$89,000
		Darter, Okaloosa	Etheostoma okaloosae	Wherever found	\$25,000	\$0	\$25,000
		Madtom, pygmy	Noturus stanauli	Wherever found	\$60	\$0	\$60
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$2,000	\$0	\$2,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$700,000	\$0	\$700,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$153,000	\$0	\$153,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$153,000	\$0	\$153,000
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$2,000	\$0	\$2,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$2,000	\$0	\$2,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$220,688	\$0	\$220,688
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$25,000	\$0	\$25,000
		Sculpin, Grotto	Cottus specus	Wherever found	\$5,000	\$0	\$5,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$61,700	\$0	\$61,700
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$2,000	\$0	\$2,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$1,300,000	\$0	\$1,300,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$1,600,000	\$0	\$1,600,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$185,000	\$0	\$185,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$17,000	\$0	\$17,000
		Sturgeon, Gulf	Acipenser oxyrinchus (=oxyrinchus) desotoi	Wherever found	\$50,000	\$0	\$50,000
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$3,600	\$0	\$3,600
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$50,000	\$0	\$50,000
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$100	\$0	\$100
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$599,500	\$0	\$599,500
		Woundfin	Plagopterus argentissimus	Wherever found, except where listed as an experimental population	\$89,000	\$0	\$89,000
	<i>Fishes Subtotal</i>				<i>\$5,334,648</i>	<i>\$0</i>	<i>\$5,334,648</i>
	Clams	Bankclimber, purple (mussel)	Elliptoideus sloatianus	Wherever found	\$20,000	\$0	\$20,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Bean, Choctaw	Villosa choctawensis	Wherever found	\$20,000	\$0	\$20,000
		Ebonyshell, round	Fusconaia rotulata	Wherever found	\$20,000	\$0	\$20,000
		Elktoe, Appalachian	Alasmidonta raveneliana	Wherever found	\$190,000	\$0	\$190,000
		Kidneyshell, southern	Ptychobranthus jonesi	Wherever found	\$20,000	\$0	\$20,000
		Moccasinshell, Gulf	Medionidus penicillatus	Wherever found	\$20,000	\$0	\$20,000
		Moccasinshell, Ochlockonee	Medionidus simpsonianus	Wherever found	\$20,000	\$0	\$20,000
		Mussel, scaleshell	Leptodea leptodon	Wherever found	\$400	\$0	\$400
		Pigtoe, fuzzy	Pleurobema strodeanum	Wherever found	\$20,000	\$0	\$20,000
		Pigtoe, narrow	Fusconaia escambia	Wherever found	\$20,000	\$0	\$20,000
		Pigtoe, oval	Pleurobema pyriforme	Wherever found	\$20,000	\$0	\$20,000
		Pocketbook, shinyrayed	Lampsilis subangulata	Wherever found	\$20,000	\$0	\$20,000
		Sandshell, Southern	Hamiota australis	Wherever found	\$20,000	\$0	\$20,000
		Slabshell, Chipola	Elliptio chipolaensis	Wherever found	\$20,000	\$0	\$20,000
		Threeridge, fat (mussel)	Amblema neislerii	Wherever found	\$20,000	\$0	\$20,000
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$2,500	\$0	\$2,500
	<i>Clams Subtotal</i>				<i>\$452,900</i>	<i>\$0</i>	<i>\$452,900</i>
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$68,500	\$0	\$68,500
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$1,400	\$0	\$1,400
		Moth, Blackburn's sphinx	Manduca blackburni	Wherever found	\$115,700	\$0	\$115,700
		Skipper, Dakota	Hesperia dacotae	Wherever found	\$32,500	\$0	\$32,500
		skipperling, Poweshiek	Oarisma poweshiek	Wherever found	\$30,600	\$0	\$30,600
		Tiger beetle, Salt Creek	Cicindela nevadica lincolniana	Wherever found	\$12,500	\$0	\$12,500
	<i>Insects Subtotal</i>				<i>\$261,200</i>	<i>\$0</i>	<i>\$261,200</i>
	Corals	Coral, elkhorn	Acropora palmata	Wherever found	\$40,000	\$0	\$40,000
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$40,000	\$0	\$40,000
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$40,000	\$0	\$40,000
		Coral, Pillar	Dendrogyra cylindrus	Wherever found	\$40,000	\$0	\$40,000
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$40,000	\$0	\$40,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Coral, staghorn	Acropora cervicornis	Wherever found	\$40,000	\$0	\$40,000
	<i>Corals Subtotal</i>				\$240,000	\$0	\$240,000
	Flowering Plants	Bladderpod, Short's	Physaria globosa	Wherever found	\$60	\$0	\$60
		Bladderpod, Spring Creek	Lesquerella perforata	Wherever found	\$60	\$0	\$60
		Bulrush, Northeastern	Scirpus ancistrochaetus	Wherever found	\$100	\$0	\$100
		Butterfly plant, Colorado	Gaura neomexicana var. coloradensis	Wherever found	\$500	\$0	\$500
		Cactus, San Rafael	Pediocactus despainii	Wherever found	\$12,500	\$0	\$12,500
		Catchfly, Spalding's	Silene spaldingii	Wherever found	\$100	\$0	\$100
		Chaffseed, American	Schwalbea americana	Wherever found	\$302,000	\$0	\$302,000
		Checker-mallow, Nelson's	Sidalcea nelsoniana	Wherever found	\$150,000	\$0	\$150,000
		Daisy, Willamette	Erigeron decumbens	Wherever found	\$62,298	\$0	\$62,298
		Howellia, Water	Howellia aquatilis		\$100	\$0	\$100
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$8,800	\$0	\$8,800
		lomatium, Bradshaw's	Lomatium bradshawii	Wherever found	\$150,000	\$0	\$150,000
		Milkweed, Mead's	Asclepias meadii	Wherever found	\$1,500	\$0	\$1,500
		Orchid, western prairie fringed	Platanthera praeclara	Wherever found	\$13,000	\$0	\$13,000
		Penstemon, blowout	Penstemon haydenii	Wherever found	\$700	\$0	\$700
		Pogonia, small whorled	Isotria medeoloides	Wherever found	\$6,200	\$0	\$6,200
	<i>Flowering Plants Subtotal</i>				\$707,918	\$0	\$707,918
	<i>Multi-Species Subtotal</i>				\$2,403,828	\$0	\$2,403,828
	<i>Domestic Total</i>				\$24,367,321	\$27,319,000	\$51,686,321
	<b>Total</b>				\$24,367,321	\$27,319,000	\$51,686,321

## Appendix C. Expenditures Reported by States for FY 2018

Through the Association of Fish and Wildlife Agencies, States voluntarily provided annual ESA expenditures directly into the Service's Web-based system. This report includes collective expenditure data from all responding State agencies.

<b>States Provided Expenditure Data for FY 2018</b>	<b>States Did Not Provided Expenditure Data for FY 2018</b>
Alaska	Alabama
Arkansas	American Samoa
Colorado	Arizona
Florida	California
Georgia	Connecticut
Idaho	Delaware
Indiana	District of Columbia
Iowa	Guam
Kentucky	Hawaii
Massachusetts	Illinois
Minnesota	Kansas
Mississippi	Louisiana
Missouri	Maine
Montana	Maryland
Nebraska	Michigan
New Hampshire	Nevada
New Jersey	New Mexico
New York	Northern Mariana Islands
North Carolina	Oklahoma
North Dakota	Puerto Rico
Ohio	Rhode Island
Oregon	Tennessee
Pennsylvania	Utah
South Carolina	Virginia
South Dakota	
Texas	
Vermont	
U.S. Virgin Islands	
Washington	
West Virginia	
Wisconsin	
Wyoming	

### Appendix C. Expenditures Reported by the States for FY 2018

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
Mammals	Bat, gray -- Wherever found	Myotis grisescens	\$25,800	\$0
	Bat, Indiana -- Wherever found	Myotis sodalis	\$191,727	\$339,643
	Bat, Mexican long-nosed -- Wherever found	Leptonycteris nivalis	\$13,576	\$0
	Bat, Northern long-eared -- Wherever found	Myotis septentrionalis	\$361,930	\$391,500
	Bat, Ozark big-eared -- Wherever found	Corynorhinus (=Plecotus) townsendii ingens	\$4,000	\$0
	Bat, Virginia big-eared -- Wherever found	Corynorhinus (=Plecotus) townsendii	\$37,182	\$0
	Bear, grizzly -- U.S.A., conterminous (lower 48) States, except where listed as an experimental population	Ursus arctos horribilis	\$3,099,083	\$0
	Bear, grizzly -- Bitterroot Grizzly Bear Nonessential Experimental Population Area	Ursus arctos horribilis	\$59,500	\$0
	Bear, grizzly -- Cabinet-Yaak Recovery Zone	Ursus arctos horribilis	\$19,500	\$0
	Bear, grizzly -- Greater Yellowstone Ecosystem DPS	Ursus arctos horribilis	\$47,000	\$0
	Bison, wood -- Wherever found	Bison bison athabascae	\$62,017	\$0
	Caribou DPS, Southern Mountain -- Southern Mountain DPS	Rangifer tarandus ssp. caribou	\$10,000	\$0
	Deer, Columbian white-tailed -- Columbia River DPS	Odocoileus virginianus leucurus	\$10,000	\$0
	Ferret, black-footed -- U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	Mustela nigripes	\$396,627	\$0
	Lynx, Canada -- Wherever Found in Contiguous U.S.	Lynx canadensis	\$1,240,002	\$90,000
	Manatee, West Indian -- Wherever found	Trichechus manatus	\$30,100	\$0
	Mouse, Preble's meadow jumping -- wherever found	Zapus hudsonius preblei	\$15,745	\$0
	Otter, Northern Sea -- Southwest Alaska DPS	Enhydra lutris kenyoni	\$4,000	\$0
	Pocket gopher, Olympia -- Wherever found	Thomomys mazama pugetensis	\$35,000	\$0
	Pocket gopher, Roy Prairie -- Wherever found	Thomomys mazama glacialis	\$35,000	\$0
	Pocket gopher, Tenino -- Wherever found	Thomomys mazama tumuli	\$35,000	\$0
	Pocket gopher, Yelm -- Wherever found	Thomomys mazama yelmensis	\$75,000	\$26,000
	Rabbit, Columbia Basin Pygmy -- Columbia Basin	Brachylagus idahoensis	\$229,600	\$940,000



Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Sea lion, Steller -- Western DPS	Eumetopias jubatus	\$125,520	\$0
	Squirrel, Carolina northern flying -- Wherever found	Glaucomys sabrinus coloratus	\$7,500	\$0
	Squirrel, Northern Idaho Ground -- Wherever found	Urocitellus brunneus	\$24,000	\$0
	whale, beluga -- Cook Inlet DPS	Delphinapterus leucas	\$194,968	\$0
	Whale, bowhead -- Wherever found	Balaena mysticetus	\$261,737	\$0
	Whale, gray -- Western North Pacific DPS	Eschrichtius robustus	\$3,000	\$0
	Whale, humpback -- Western North Pacific	Megaptera novaeangliae	\$3,000	\$0
	Whale, killer -- Southern Resident DPS	Orcinus orca	\$388,600	\$0
	Whale, North Atlantic Right -- Wherever found	Eubalaena glacialis	\$125,000	\$0
	Wolf, gray -- U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	Canis lupus	\$2,116,640	\$590,000
	Wolf, gray -- MN	Canis lupus	\$154,900	\$0
	Wolf, red -- Wherever found, except where listed as an experimental population	Canis rufus	\$1,000	\$0
<i>Mammals subtotal</i>			<i>\$9,443,254</i>	<i>\$2,377,143</i>
Birds	Albatross, short-tailed -- Wherever found	Phoebastria (=Diomedea) albatrus	\$2,000	\$0
	Crane, Mississippi sandhill -- Wherever found	Grus canadensis pulla	\$500	\$0
	Crane, whooping -- Wherever found, except where listed as an experimental population	Grus americana	\$9,400	\$57,702
	Crane, whooping -- U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	Grus americana	\$92,000	\$0
	Cuckoo, yellow-billed -- Western U.S. DPS	Coccyzus americanus	\$16,500	\$0
	Flycatcher, southwestern willow -- Wherever found	Empidonax traillii extimus	\$1,830	\$0
	Horned lark, streaked -- Wherever found	Eremophila alpestris strigata	\$31,000	\$0
	Knot, red -- Wherever found	Calidris canutus rufa	\$62,360	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Murrelet, marbled -- U.S.A. (CA, OR, WA)	Brachyramphus marmoratus	\$1,618,600	\$0
	Owl, northern spotted -- Wherever found	Strix occidentalis caurina	\$914,600	\$0
	Plover, piping -- [Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	Charadrius melodus	\$35,099	\$0
	Plover, piping -- [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	Charadrius melodus	\$368,502	\$55,500
	Plover, western snowy -- Pacific Coast population DPS- U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	Charadrius nivosus nivosus	\$92,400	\$0
	sage-grouse, Gunnison -- Wherever found	Centrocercus minimus	\$464,417	\$0
	Stork, wood -- AL, FL, GA, MS, NC, SC	Mycteria americana	\$75,400	\$0
	Tern, least -- interior pop.	Sterna antillarum	\$56,801	\$0
	Tern, roseate -- Northeast U.S. nesting population	Sterna dougallii dougallii	\$32,000	\$0
	Warbler, Kirtland's -- Wherever found	Setophaga kirtlandii (= Dendroica kirtlandii)	\$56,366	\$0
	Warbler (=wood), Bachman's -- Wherever found	Vermivora bachmanii	\$100	\$0
	Woodpecker, red-cockaded -- Wherever found	Picoides borealis	\$195,200	\$0
<i>Birds subtotal</i>			<i>\$4,125,075</i>	<i>\$113,202</i>
Reptiles	Alligator, American -- Wherever found	Alligator mississippiensis	\$135,000	\$0
	Boa, Virgin Islands tree -- Wherever found	Chilabothrus granti	\$14,556	\$0
	Massasauga (=rattlesnake), eastern -- Wherever found	Sistrurus catenatus	\$78,607	\$0
	piresnake, black -- Wherever found	Pituophis melanoleucus lodingi	\$400	\$0
	Pinesnake, Louisiana -- Wherever found	Pituophis ruthveni	\$7,399	\$0
	Sea turtle, green -- North Atlantic DPS	Chelonia mydas	\$3,000	\$0
	Sea turtle, green -- South Atlantic DPS	Chelonia mydas	\$12,000	\$0
	Sea turtle, hawksbill -- Wherever found	Eretmochelys imbricata	\$1,400	\$0
	Sea turtle, Kemp's ridley -- Wherever found	Lepidochelys kempii	\$12,900	\$0
	Sea turtle, leatherback -- Wherever found	Dermochelys coriacea	\$6,800	\$0
	Sea turtle, loggerhead -- Wherever found	Caretta caretta	\$31,300	\$0
	Sea turtle, loggerhead -- Northwest Atlantic Ocean	Caretta caretta	\$150,000	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Snake, copperbelly water -- Indiana north of 40 degrees north latitude, Michigan, Ohio	Nerodia erythrogaster neglecta	\$6,366	\$0
	Snake, eastern indigo -- Wherever found	Drymarchon corais couperi	\$200,000	\$2,000,000
	Tortoise, gopher -- West of Mobile and Tombigbee Rivers	Gopherus polyphemus	\$6,000	\$0
	Turtle, Alabama red-bellied -- Wherever found	Pseudemys alabamensis	\$100	\$0
	turtle, bog -- Wherever found, except GA, NC, SC, TN, VA	Clemmys muhlenbergii	\$38,000	\$0
	turtle, bog -- U.S.A. (GA, NC, SC, TN, VA)	Clemmys muhlenbergii	\$115,100	\$0
	Turtle, Plymouth Redbelly -- Wherever found	Pseudemys rubriventris bangsi	\$300	\$0
	Turtle, ringed map -- Wherever found	Graptemys oculifera	\$400	\$0
	Turtle, yellow-blotched map -- Wherever found	Graptemys flavimaculata	\$400	\$0
<i>Reptiles subtotal</i>			<i>\$820,028</i>	<i>\$2,000,000</i>
Amphibians	Frog, dusky gopher -- Wherever found	Rana sevosia	\$22,000	\$0
	Frog, Oregon spotted -- Wherever found	Rana pretiosa	\$52,800	\$0
	Hellbender, Ozark -- Wherever found	Cryptobranchus alleganiensis bishopi	\$100,900	\$0
	Salamander, Austin blind -- Wherever found	Eurycea waterlooensis	\$2,502	\$0
	Salamander, Barton Springs -- Wherever found	Eurycea sosorum	\$2,502	\$0
	Salamander, frosted flatwoods -- Wherever found	Ambystoma cingulatum	\$30,000	\$0
	Salamander, Georgetown -- Wherever found	Eurycea naufragia	\$2,502	\$0
	salamander, Reticulated flatwoods -- Wherever found	Ambystoma bishopi	\$30,000	\$0
	Salamander, San Marcos -- Spring Lake and nearby habitats immediately downstream of the dam forming Spring Lake	Eurycea nana	\$2,502	\$0
	Toad, Houston -- Wherever found	Bufo houstonensis	\$4,387	\$0
	Toad, Wyoming -- Wherever found	Bufo hemiophrys baxteri	\$15,251	\$0
<i>Amphibians subtotal</i>			<i>\$265,346</i>	<i>\$0</i>
Fishes	Bocaccio -- Puget Sound - Georgia Basin DPS	Sebastes paucispinis	\$235,600	\$0
	Bonytail -- Wherever found	Gila elegans	\$217,932	\$0
	Cavefish, Ozark -- Wherever found	Amblyopsis rosae	\$500	\$0
	Chub, Borax Lake -- Wherever found	Gila boraxobius	\$37,000	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Chub, humpback -- Wherever found	<i>Gila cypha</i>	\$83,248	\$0
	Chub, spotfin -- Wherever found, except where listed as an experimental population	<i>Erimonax monachus</i>	\$4,000	\$0
	Dace, Foskett speckled -- Wherever found	<i>Rhinichthys osculus</i> ssp.	\$43,000	\$0
	Darter, amber -- Wherever found	<i>Percina antesella</i>	\$5,000	\$0
	Darter, bayou -- Wherever found	<i>Etheostoma rubrum</i>	\$400	\$0
	Darter, Cherokee -- Wherever found	<i>Etheostoma scotti</i>	\$5,000	\$0
	Darter, Cumberland -- Wherever found	<i>Etheostoma susanae</i>	\$1,000	\$0
	Darter, Etowah -- Wherever found	<i>Etheostoma etowahae</i>	\$5,000	\$0
	Darter, goldline -- Wherever found	<i>Percina aurolineata</i>	\$5,000	\$0
	Darter, Kentucky arrow -- Wherever found	<i>Etheostoma spilotum</i>	\$500	\$0
	Darter, leopard -- Wherever found	<i>Percina pantherina</i>	\$32,500	\$0
	Darter, Niangua -- Wherever found	<i>Etheostoma nianguae</i>	\$18,500	\$0
	Darter, relict -- Wherever found	<i>Etheostoma chienense</i>	\$500	\$0
	Darter, snail -- Wherever found	<i>Percina tanasi</i>	\$3,000	\$0
	Darter, yellowcheek -- Wherever found	<i>Etheostoma moorei</i>	\$800	\$0
	Eulachon -- Southern DPS	<i>Thaleichthys pacificus</i>	\$282,600	\$0
	Logperch, Conasauga -- Wherever found	<i>Percina jenkinsi</i>	\$5,000	\$0
	Logperch, Roanoke -- Wherever found	<i>Percina rex</i>	\$3,000	\$0
	Minnow, Devils River -- Wherever found	<i>Dionda diaboli</i>	\$14,705	\$0
	Pikeminnow (=squawfish), Colorado -- Wherever found, except where listed as an experimental	<i>Ptychocheilus lucius</i>	\$83,248	\$0
	rockfish, canary -- Puget Sound - Georgia Basin DPS	<i>Sebastes pinniger</i>	\$303,200	\$0
	rockfish, yelloweye -- Puget Sound - Georgia Basin	<i>Sebastes ruberrimus</i>	\$303,200	\$0
	Salmon, Chinook -- Snake River fall-run ESU	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	\$408,900	\$0
	Salmon, Chinook -- Snake River spring/summer-run	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	\$1,175,000	\$0
	Salmon, Chinook -- Puget Sound ESU	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	\$63,067,800	\$0
	Salmon, Chinook -- Lower Columbia River ESU	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	\$9,555,200	\$0
	Salmon, Chinook -- Upper Willamette River ESU	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	\$407,000	\$0
	Salmon, Chinook -- Upper Columbia spring-run ESU	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	\$5,163,800	\$0
	Salmon, chum -- Columbia River ESU	<i>Oncorhynchus keta</i>	\$706,000	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Salmon, chum -- Hood Canal summer-run ESU	Oncorhynchus keta	\$2,996,400	\$0
	Salmon, coho -- Southern Oregon - Northern California Coast ESU	Oncorhynchus (=Salmo) kisutch	\$110,000	\$0
	Salmon, coho -- Oregon Coast ESU	Oncorhynchus (=Salmo) kisutch	\$1,912,000	\$0
	Salmon, coho -- Lower Columbia River ESU	Oncorhynchus (=Salmo) kisutch	\$3,268,600	\$0
	Salmon, sockeye -- Snake River ESU	Oncorhynchus (=Salmo) nerka	\$223,300	\$0
	Salmon, sockeye -- Ozette Lake ESU	Oncorhynchus (=Salmo) nerka	\$230,300	\$0
	Sculpin, Grotto -- Wherever found	Cottus specus	\$37,000	\$0
	Shark, Scalloped Hammerhead -- Wherever found	Sphyrna lewini	\$19,500	\$0
	Shiner, blue -- Wherever found	Cyprinella caerulea	\$5,000	\$0
	Shiner, Cape Fear -- Wherever found	Notropis mekistocholas	\$3,800	\$0
	Shiner, sharpnose -- Wherever found	Notropis oxyrinchus	\$14,165	\$0
	Shiner, smalleye -- Wherever found	Notropis buccula	\$14,165	\$0
	Shiner, Topeka -- Wherever found, except where listed as an experimental population	Notropis topeka (=tristis)	\$10,300	\$0
	Silverside, Waccamaw -- Wherever found	Menidia extensa	\$3,400	\$0
	Steelhead -- Snake River Basin DPS	Oncorhynchus (=Salmo) mykiss	\$1,056,000	\$0
	Steelhead -- Lower Columbia River DPS	Oncorhynchus (=Salmo) mykiss	\$2,704,400	\$0
	Steelhead -- Upper Columbia River DPS	Oncorhynchus (=Salmo) mykiss	\$2,741,600	\$0
	Steelhead -- Upper Willamette River DPS	Oncorhynchus (=Salmo) mykiss	\$421,000	\$0
	Steelhead -- Middle Columbia River DPS	Oncorhynchus (=Salmo) mykiss	\$1,865,400	\$0
	Steelhead -- Puget Sound DPS	Oncorhynchus (=Salmo) mykiss	\$2,549,000	\$0
	Steelhead -- Middle Columbia River DPS	Oncorhynchus (=Salmo) mykiss	\$26,200	\$0
	Sturgeon, Atlantic -- New York Bight DPS	Acipenser oxyrinchus oxyrinchus	\$29,000	\$0
	Sturgeon, Atlantic -- South Atlantic DPS	Acipenser oxyrinchus oxyrinchus	\$75,000	\$0
	sturgeon, green -- Southern DPS	Acipenser medirostris	\$29,600	\$0
	Sturgeon, Gulf -- Wherever found	Acipenser oxyrinchus (=oxyrinchus) desotoi	\$20,000	\$0
	Sturgeon, pallid -- Wherever found	Scaphirhynchus albus	\$69,900	\$0
	Sturgeon, shortnose -- Wherever found	Acipenser brevirostrum	\$137,500	\$0
	Sturgeon, Shovelnose -- Wherever found	Scaphirhynchus platyrhynchus	\$23,700	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Sturgeon, white -- U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	Acipenser transmontanus	\$341,730	\$0
	Sucker, Lost River -- Wherever found	Deltistes luxatus	\$16,000	\$0
	Sucker, razorback -- Wherever found	Xyrauchen texanus	\$217,932	\$0
	Sucker, shortnose -- Wherever found	Chasmistes brevirostris	\$16,000	\$0
	Sucker, Warner -- Wherever found	Catostomus warnerensis	\$82,000	\$0
	Trout, bull -- U.S.A., conterminous, lower 48 states	Salvelinus confluentus	\$3,655,500	\$590,000
	Trout, bull -- Clackamas River subbasin experimental population	Salvelinus confluentus	\$447,600	\$0
	trout, Greenback Cutthroat -- Wherever found	Oncorhynchus clarkii stomias	\$99,302	\$0
<i>Fishes subtotal</i>			<i>\$107,649,427</i>	<i>\$590,000</i>
Clams	Bankclimber, purple (mussel) -- Wherever found	Elliptoideus sloatianus	\$5,000	\$0
	Bean, Cumberland (pearlymussel) -- Wherever found; Except where listed as Experimental Populations	Villosa trabalis	\$20,000	\$0
	Bean, rayed -- Wherever found	Villosa fabalis	\$6,166	\$0
	Catspaw, white (pearlymussel) -- Wherever found	Epioblasma obliquata perobliqua	\$6,166	\$0
	Clubshell -- Wherever found; Except where listed as Experimental Populations	Pleurobema clava	\$45,391	\$0
	Clubshell, black -- Wherever found	Pleurobema curtum	\$400	\$0
	Clubshell, ovate -- Wherever found	Pleurobema perovatum	\$2,900	\$0
	Clubshell, southern -- Wherever found	Pleurobema decisum	\$3,400	\$0
	Combshell, Cumberlandian -- Wherever found; Except where listed as Experimental Populations	Epioblasma brevidens	\$20,000	\$0
	Combshell, southern -- Wherever found	Epioblasma penita	\$1,400	\$0
	Combshell, upland -- Wherever found	Epioblasma metastriata	\$1,000	\$0
	Elktoe, Appalachian -- Wherever found	Alasmidonta raveneliana	\$9,900	\$0
	Elktoe, Cumberland -- Wherever found	Alasmidonta atropurpurea	\$5,000	\$0
	Fanshell -- Wherever found	Cyprogenia stegaria	\$10,166	\$0
	fatmucket, Arkansas -- Wherever found	Lampsilis powellii	\$4,000	\$174,000
	Heelsplitter, Carolina -- Wherever found	Lasmigona decorata	\$67,600	\$0
	heelsplitter, Inflated -- Wherever found	Potamilus inflatus	\$400	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Higgins eye (pearlymussel) -- Wherever found	Lampsilis higginsii	\$15,800	\$0
	Kidneyshell, triangular -- Wherever found	Ptychobranthus greenii	\$2,000	\$0
	Lance, yellow -- Wherever found	Elliptio lanceolata	\$15,310	\$77,600
	Moccasinshell, Alabama -- Wherever found	Medionidus acutissimus	\$3,400	\$0
	Moccasinshell, Coosa -- Wherever found	Medionidus parvulus	\$1,000	\$0
	Moccasinshell, Gulf -- Wherever found	Medionidus penicillatus	\$2,500	\$0
	Moccasinshell, Ochlockonee -- Wherever found	Medionidus simpsonianus	\$1,000	\$0
	Moccasinshell, Suwannee -- Wherever found	Medionidus walkeri	\$5,000	\$0
	Mucket, Neosho -- Wherever found	Lampsilis rafinesqueana	\$8,800	\$0
	Mucket, pink (pearlymussel) -- Wherever found	Lampsilis abrupta	\$16,166	\$0
	Mussel, scaleshell -- Wherever found	Leptodea leptodon	\$1,700	\$0
	Mussel, sheepnose -- Wherever found	Plethobasus cyphus	\$13,666	\$0
	Mussel, snuffbox -- Wherever found	Epioblasma triquetra	\$26,266	\$38,500
	Mussel, snuffbox -- Wherever found	Epioblasma triquetra	\$400	\$0
	Pearlshell, Louisiana -- Wherever found	Margaritifera hembeli	\$1,000	\$0
	Pearlymussel, dromedary -- Wherever found; Except where listed as Experimental Populations	Dromus dromas	\$5,000	\$0
	Pearlymussel, littlewing -- Wherever found	Pegias fabula	\$10,000	\$0
	Pearlymussel, slabside -- Wherever found	Pleuroaia dolabelloides	\$400	\$0
	Pigtoe, flat -- Wherever found	Pleurobema marshalli	\$400	\$0
	Pigtoe, Georgia -- Wherever found	Pleurobema hanleyianum	\$3,000	\$0
	Pigtoe, oval -- Wherever found	Pleurobema pyriforme	\$5,000	\$0
	Pigtoe, rough -- Wherever found	Pleurobema plenum	\$2,000	\$0
	Pigtoe, southern -- Wherever found	Pleurobema georgianum	\$5,000	\$0
	Pimpleback, orangefoot (pearlymussel) -- Wherever found	Plethobasus cooperianus	\$3,529	\$0
	Pimpleback, orangefoot (pearlymussel) -- U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	Plethobasus cooperianus	\$5,000	\$0
	Pocketbook, finelined -- Wherever found	Lampsilis altilis	\$5,000	\$0
	Pocketbook, Ouachita rock -- Wherever found	Arkansia wheeleri	\$1,600	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Pocketbook, shinyrayed -- Wherever found	Lampsilis subangulata	\$5,000	\$0
	Pocketbook, speckled -- Wherever found	Lampsilis streckeri	\$3,200	\$0
	Purple Cat's paw (=Purple Cat's paw pearlymussel) -- Wherever found; Except where listed as Experimental Populations	Epioblasma obliquata obliquata	\$6,166	\$0
	Rabbitsfoot -- Wherever found	Quadrula cylindrica cylindrica	\$7,866	\$0
	Rabbitsfoot --	Quadrula cylindrica cylindrica	\$400	\$0
	Riffleshell, northern -- Wherever found	Epioblasma torulosa rangiana	\$7,166	\$0
	Riffleshell, tan -- Wherever found	Epioblasma florentina walkeri (=E. walkeri)	\$10,000	\$0
	Ring pink (mussel) -- Wherever found	Obovaria retusa	\$2,000	\$0
	Spectaclecase (mussel) -- Wherever found	Cumberlandia monodonta	\$28,800	\$0
	Spinymussel, Altamaha -- Wherever found	Elliptio spinosa	\$2,000	\$0
	Spinymussel, James -- Wherever found	Pleurobema collina	\$2,100	\$2,100
	Spinymussel, Tar River -- Wherever found	Elliptio steinstansana	\$4,900	\$77,600
	Stirrupshell -- Wherever found	Quadrula stapes	\$400	\$0
	Threeridge, fat (mussel) -- Wherever found	Amblema neislerii	\$7,529	\$0
	Wedgemussel, dwarf -- Wherever found	Alasmidonta heterodon	\$7,400	\$77,665
<i>Clams subtotal</i>			<i>\$463,753</i>	<i>\$447,465</i>
Snails	Cavesnail, Tumbling Creek -- Wherever found	Antrobia culveri	\$6,000	\$0
	Snail, Chittenango ovate amber -- Wherever found	Succinea chittenangoensis	\$800	\$0
	Springsnail, Bruneau Hot -- Wherever found	Pyrgulopsis bruneauensis	\$815	\$0
<i>Snails subtotal</i>			<i>\$7,615</i>	<i>\$0</i>
Insects	Beetle, American burying -- Wherever found, except where listed as an experimental population	Nicrophorus americanus	\$164,466	\$60,000
	Beetle, Comal Springs dryopid -- Wherever found	Stygoparnus comalensis	\$5,972	\$0
	Bumble bee, Rusty patched -- Wherever found	Bombus affinis	\$30,000	\$0
	Butterfly, Karner blue -- Wherever found	Lycaeides melissa samuelis	\$217,654	\$0
	Butterfly, Mitchell's satyr -- Wherever found	Neonympha mitchellii mitchellii	\$2,300	\$0
	Butterfly, Oregon silverspot --	Speyeria zerene hippolyta	\$4,000	\$0
	Checkerspot, Taylor's (=whulge) -- Wherever found	Euphydryas editha taylori	\$64,000	\$0
	Skipper, Dakota -- Wherever found	Hesperia dacotae	\$62,100	\$0



Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	skipperling, Poweshiek -- Wherever found	Oarisma poweshiek	\$14,980	\$0
	Tiger beetle, Salt Creek -- Wherever found	Cicindela nevadica lincolniana	\$25,000	\$128,300
<i>Insects subtotal</i>			<i>\$590,472</i>	<i>\$188,300</i>
Flowering Plants	Amphianthus, little -- Wherever found	Amphianthus pusillus	\$2,000	\$0
	Arrowhead, bunched -- Wherever found	Sagittaria fasciculata	\$4,100	\$0
	Aster, Florida golden -- Wherever found	Chrysopsis floridana	\$29,000	\$0
	Avens, spreading -- Wherever found	Geum radiatum	\$400	\$0
	Barbara's Buttons, Mohr's -- Wherever found	Marshallia mohrii	\$4,000	\$0
	Beargrass, Britton's -- Wherever found	Nolina brittoniana	\$8,900	\$0
	Beauty, Harper's -- Wherever found	Harperocallis flava	\$300	\$0
	Bellflower, Brooksville -- Wherever found	Campanula robinsiae	\$6,000	\$0
	Birds-in-a-nest, white -- Wherever found	Macbridea alba	\$1,900	\$0
	Bittercress, small-anthered -- Wherever found	Cardamine micranthera	\$400	\$0
	Bladderpod, Dudley Bluffs -- Wherever found	Lesquerella congesta	\$1,450	\$0
	bladderpod, Missouri -- Wherever found	Physaria filiformis	\$2,000	\$0
	Bladderpod, Short's -- Wherever found	Physaria globosa	\$1,300	\$0
	Blazingstar, Heller's -- Wherever found	Liatris helleri	\$3,200	\$0
	Blazingstar, scrub -- Wherever found	Liatris ohlingerae	\$5,100	\$0
	Bluet, Roan Mountain -- Wherever found	Hedyotis purpurea var. montana	\$500	\$0
	Bonamia, Florida -- Wherever found	Bonamia grandiflora	\$3,500	\$0
	Brickell-bush, Florida -- Wherever found	Brickellia mosieri	\$1,300	\$0
	Buckwheat, scrub -- Wherever found	Eriogonum longifolium var. gnaphalifolium	\$8,000	\$0
	Bully, Everglades -- Wherever found	Sideroxylon reclinatum ssp. austrofloridense	\$200	\$0
	Bulrush, Northeastern --	Scirpus ancistrochaetus	\$8,038	\$0
	Bush-clover, prairie -- Wherever found	Lespedeza leptostachya	\$1,500	\$0
	Butterfly plant, Colorado -- Wherever found	Gaura neomexicana var. coloradensis	\$100	\$0
	Butterwort, Godfrey's -- Wherever found	Pinguicula ionantha	\$1,800	\$0
	Cactus, black lace -- Wherever found	Echinocereus reichenbachii var. albertii	\$8,089	\$0
	Cactus, Colorado hookless -- Wherever found	Sclerocactus glaucus	\$1,200	\$0
	Cactus, Florida semaphore -- Wherever found	Consolea corallicola	\$600	\$0
	Cactus, Key tree -- Wherever found	Pilosocereus robinii	\$10,500	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Campion, fringed -- Wherever found	<i>Silene polypetala</i>	\$6,700	\$0
	Catchfly, Spalding's -- Wherever found	<i>Silene spaldingii</i>	\$0	\$221,000
	Chaffseed, American -- Wherever found	<i>Schwalbea americana</i>	\$5,600	\$0
	Clover, running buffalo --	<i>Trifolium stoloniferum</i>	\$8,858	\$0
	Coneflower, smooth -- Wherever found	<i>Echinacea laevigata</i>	\$11,400	\$0
	Crabgrass, Florida pineland -- Wherever found	<i>Digitaria pauciflora</i>	\$200	\$0
	Dropwort, Canby's -- Wherever found	<i>Oxypolis canbyi</i>	\$27,400	\$20,000
	Flax, Carter's small-flowered -- Wherever found	<i>Linum carteri carteri</i>	\$1,300	\$0
	Flax, sand -- Wherever found	<i>Linum arenicola</i>	\$1,800	\$0
	Fringe-tree, pygmy -- Wherever found	<i>Chionanthus pygmaeus</i>	\$3,000	\$0
	Gerardia, sandplain -- Wherever found	<i>Agalinis acuta</i>	\$600	\$0
	Goldenrod, Blue Ridge -- Wherever found	<i>Solidago spithamaea</i>	\$300	\$0
	Goldenrod, Short's -- Wherever found	<i>Solidago shortii</i>	\$1,800	\$0
	Gooseberry, Miccosukee -- Wherever found	<i>Ribes echinellum</i>	\$200	\$0
	Gourd, Okeechobee -- Wherever found	<i>Cucurbita okeechobeensis</i> ssp. <i>okeechobeensis</i>	\$200	\$0
	Grass, Tennessee yellow-eyed -- Wherever found	<i>Xyris tennesseensis</i>	\$5,000	\$0
	Harebells, Avon Park -- Wherever found	<i>Crotalaria avonensis</i>	\$12,400	\$0
	Harperella -- Wherever found	<i>Ptilimnium nodosum</i>	\$10,862	\$0
	Heartleaf, dwarf-flowered -- Wherever found	<i>Hexastylis naniflora</i>	\$3,400	\$0
	Howellia, Water --	<i>Howellia aquatilis</i>	\$68,000	\$0
	Hypericum, highlands scrub -- Wherever found	<i>Hypericum cumulicola</i>	\$8,000	\$0
	Iris, dwarf lake -- Wherever found	<i>Iris lacustris</i>	\$6,500	\$0
	Irisette, white -- Wherever found	<i>Sisyrinchium dichotomum</i>	\$2,300	\$0
	Jacquemontia, beach -- Wherever found	<i>Jacquemontia reclinata</i>	\$5,500	\$0
	Ladies'-tresses, Ute -- Wherever found	<i>Spiranthes diluvialis</i>	\$100	\$0
	Lead-plant, Crenulate -- Wherever found	<i>Amorpha crenulata</i>	\$6,000	\$0
	Leather flower, Alabama -- Wherever found	<i>Clematis socialis</i>	\$3,000	\$0
	Locoweed, Fassett's -- Wherever found	<i>Oxytropis campestris</i> var. <i>chartacea</i>	\$2,500	\$0
	Loosestrife, rough-leaved -- Wherever found	<i>Lysimachia asperulaefolia</i>	\$2,600	\$0
	Lupine, scrub -- Wherever found	<i>Lupinus aridorum</i>	\$1,500	\$0
	Meadowrue, Cooley's -- Wherever found	<i>Thalictrum cooleyi</i>	\$3,540	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Milkpea, Small's -- Wherever found	Galactia smallii	\$300	\$0
	Milk-vetch, Jesup's -- Wherever found	Astragalus robbinsii var. jesupi	\$13,000	\$0
	milkvetch, Osterhout -- Wherever found	Astragalus osterhoutii	\$715	\$0
	Milkweed, Mead's -- Wherever found	Asclepias meadii	\$45,000	\$0
	Mint, Garrett's -- Wherever found	Dicerandra christmanii	\$800	\$0
	Mint, Lakela's -- Wherever found	Dicerandra immaculata	\$12,500	\$0
	Mint, longspurred -- Wherever found	Dicerandra cornutissima	\$200	\$0
	Mint, scrub -- Wherever found	Dicerandra frutescens	\$5,500	\$0
	Monkshood, northern wild -- Wherever found	Aconitum noveboracense	\$1,000	\$0
	Mustard, Carter's -- Wherever found	Warea carteri	\$7,500	\$0
	No common name -- Wherever found	Geocarpon minimum	\$1,600	\$0
	Orchid, eastern prairie fringed -- Wherever found	Platanthera leucophaea	\$39,000	\$0
	Orchid, western prairie fringed -- Wherever found	Platanthera praeclara	\$22,415	\$0
	Orchid, white fringeless --	Platanthera integrilabia	\$6,000	\$0
	paintbrush, golden -- Wherever found	Castilleja levisecta	\$2,600	\$0
	Pawpaw, beautiful -- Wherever found	Deeringothamnus pulchellus	\$1,500	\$0
	Pawpaw, four-petal -- Wherever found	Asimina tetramera	\$1,100	\$0
	Pawpaw, Rugel's -- Wherever found	Deeringothamnus rugelii	\$5,500	\$0
	Pea, Big Pine partridge -- Wherever found	Chamaecrista lineata keyensis	\$1,700	\$0
	Penstemon, blowout -- Wherever found	Penstemon haydenii	\$4,800	\$0
	Phacelia, DeBeque -- Wherever found	Phacelia submutica	\$1,340	\$0
	Phacelia, North Park -- Wherever found	Phacelia formosula	\$2,165	\$0
	Pigeon wings -- Wherever found	Clitoria fragrans	\$6,000	\$0
	Pinkroot, gentian -- Wherever found	Spigelia gentianoides	\$200	\$0
	Pink, swamp -- Wherever found	Helonias bullata	\$6,700	\$0
	Pitcher-plant, mountain sweet -- Wherever found	Sarracenia rubra ssp. jonesii	\$3,200	\$0
	Plum, scrub -- Wherever found	Prunus geniculata	\$6,000	\$0
	Pogonia, small whorled --	Isotria medeoloides	\$6,681	\$0
	Polygala, Lewton's -- Wherever found	Polygala lewtonii	\$14,200	\$0
	Polygala, tiny -- Wherever found	Polygala smallii	\$300	\$0
	Pondberry -- Wherever found	Lindera melissifolia	\$12,400	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	potato-bean, Prices --	<i>Apios priceana</i>	\$500	\$0
	Prairie-clover, Florida -- Wherever found	<i>Dalea carthagenensis floridana</i>	\$5,000	\$0
	Prickly-apple, aboriginal -- Wherever found	<i>Harrisia (=Cereus) aboriginum (=gracilis)</i>	\$800	\$0
	Prickly-apple, fragrant -- Wherever found	<i>Cereus eriophorus</i> var. <i>fragrans</i>	\$6,500	\$0
	Rattleweed, hairy -- Wherever found	<i>Baptisia arachnifera</i>	\$40,000	\$0
	Rhododendron, Chapman -- Wherever found	<i>Rhododendron chapmanii</i>	\$450	\$0
	Rockcress, Georgia -- Wherever found	<i>Arabis georgiana</i>	\$5,000	\$0
	rock cress, Shale barren -- Wherever found	<i>Arabis serotina</i>	\$5,560	\$0
	Rosemary, Apalachicola -- Wherever found	<i>Conradina glabra</i>	\$200	\$0
	Rosemary, Etonia -- Wherever found	<i>Conradina etonia</i>	\$9,000	\$0
	Rosemary, short-leaved -- Wherever found	<i>Conradina brevifolia</i>	\$6,500	\$0
	Sandlace -- Wherever found	<i>Polygonella myriophylla</i>	\$4,500	\$0
	Sandmat, pineland -- Wherever found	<i>Chamaesyce deltoidea pinetorum</i>	\$1,200	\$0
	Seagrass, Johnson's -- Wherever found	<i>Halophila johnsonii</i>	\$700	\$0
	silverbush, Blodgett's -- Wherever found	<i>Argythamnia blodgettii</i>	\$2,500	\$0
	Skullcap, Florida -- Wherever found	<i>Scutellaria floridana</i>	\$1,000	\$0
	Skullcap, large-flowered -- Wherever found	<i>Scutellaria montana</i>	\$15,000	\$0
	Skyrocket, Pagosa -- Wherever found	<i>Ipomopsis polyantha</i>	\$11,465	\$0
	Snakeroot -- Wherever found	<i>Eryngium cuneifolium</i>	\$5,500	\$0
	Sneezeweed, Virginia -- Wherever found	<i>Helenium virginicum</i>	\$1,200	\$0
	Spiraea, Virginia -- Wherever found	<i>Spiraea virginiana</i>	\$10,839	\$0
	Spurge, deltoid -- Wherever found	<i>Chamaesyce deltoidea</i> ssp. <i>deltoidea</i>	\$1,200	\$0
	Spurge, Garber's -- Wherever found	<i>Chamaesyce garberi</i>	\$1,200	\$0
	Spurge, telephus -- Wherever found	<i>Euphorbia telephioides</i>	\$6,500	\$0
	Spurge, wedge -- Wherever found	<i>Chamaesyce deltoidea serpyllum</i>	\$1,200	\$0
	Sumac, Michaux's -- Wherever found	<i>Rhus michauxii</i>	\$5,000	\$0
	Sunflower, Schweinitz's -- Wherever found	<i>Helianthus schweinitzii</i>	\$6,800	\$0
	Sunflower, whorled --	<i>Helianthus verticillatus</i>	\$5,000	\$0
	Thistle, Pitcher's -- Wherever found	<i>Cirsium pitcheri</i>	\$5,600	\$0
	Thoroughwort, Cape Sable -- Wherever found	<i>Chromolaena frustrata</i>	\$1,700	\$0
	Trillium, persistent -- Wherever found	<i>Trillium persistens</i>	\$8,000	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Trillium, relict -- Wherever found	Trillium reliquum	\$4,000	\$0
	Warea, wide-leaf -- Wherever found	Warea amplexifolia	\$46,500	\$0
	Water-plantain, Kral's -- Wherever found	Sagittaria secundifolia	\$1,000	\$0
	Water-willow, Cooley's -- Wherever found	Justicia cooleyi	\$200	\$0
	Whitlow-wort, papery -- Wherever found	Paronychia chartacea	\$6,000	\$0
	Wireweed -- Wherever found	Polygonella basiramia	\$2,700	\$0
	Ziziphus, Florida -- Wherever found	Ziziphus celata	\$13,500	\$0
<i>Flowering Plants subtotal</i>			<i>\$807,367</i>	<i>\$241,000</i>
Conifers and Cycads	Torreya, Florida -- Wherever found	Torreya taxifolia	\$1,500	\$0
<i>Conifers and Cycads subtotal</i>			<i>\$1,500</i>	<i>\$0</i>
Ferns and Allies	fern, Florida bristle -- Wherever found	Trichomanes punctatum ssp. floridanum	\$1,200	\$0
	Quillwort, black spored -- Wherever found	Isoetes melanospora	\$2,000	\$0
	Quillwort, mat-forming -- Wherever found	Isoetes tegetiformans	\$3,000	\$0
<i>Ferns and Allies subtotal</i>			<i>\$6,200</i>	<i>\$0</i>
Lichens	Cladonia, Florida perforate -- Wherever found	Cladonia perforata	\$7,000	\$0
	Lichen, rock gnome -- Wherever found	Gymnoderma lineare	\$600	\$0
<i>Lichens subtotal</i>			<i>\$7,600</i>	<i>\$0</i>
<i>Multi-Species Expenditures subtotal</i>			<i>\$1,559,778</i>	<i>\$44,300</i>
<b>Total</b>			<b>\$125,747,415</b>	<b>\$6,001,410</b>