


FEDERAL AND STATE ENDANGERED AND THREATENED SPECIES EXPENDITURES



Fiscal Year 2017

TABLE OF CONTENTS

ENDANGERED AND THREATENED SPECIES EXPENDITURES FISCAL YEAR 2017	
PURPOSE	3
BACKGROUND	4
SECTION 18(1), FEDERAL EXPENDITURES	5
SECTION 18(2), STATE EXPENDITURES	5
SUMMARY OF DATA	5
COMPARISON TO PREVIOUS REPORTS	6
Table 1. FY 2017 Reported Expenditures for Endangered and Threatened Species.....	7
Table 2. Species Ranked in Descending Order of Total FY 2017 Reported Expenditures.....	58
Table 3. FY 2017 Reported Land Acquisition Expenditures for Endangered and Threatened Species.....	121
Table 4. FY 2017 Expenditures Reported by Individual Federal Agencies and the States.....	133
Appendix A. Supplemental Guidance for Reporting Expenditures for the Conservation of Endangered and Threatened Species.....	136
Appendix B. Reports Provided by Other Federal Agencies.....	141
Appendix C. Expenditures Reported by the States.....	421

ENDANGERED SPECIES EXPENDITURES FISCAL YEAR 2017

PURPOSE

The Endangered Species Act of 1973 (16 U.S.C. 1531 *et seq.*) (Act) was reauthorized and amended on October 7, 1988, under Public Law 100-478. A new section 18 was added to the Act that requires the U.S. Fish and Wildlife Service (Service) on behalf of the Secretary of the Interior to report annually on certain expenditures on a species-by-species basis. The section reads:

SEC. 18. On or before January 15, 1990, and each January 15 thereafter, the Secretary of the Interior, acting through the Fish and Wildlife Service, shall submit to the Congress an annual report covering the preceding fiscal year which shall contain

(1) the accounting on a species-by-species basis of all reasonably identifiable Federal expenditures made primarily for the conservation of endangered or threatened species pursuant to this Act; and

(2) an accounting on a species-by-species basis of all reasonably identifiable expenditures made primarily for the conservation of endangered or threatened species pursuant to this Act by States receiving grants under section 6.

This document constitutes the 28th report by the Service pursuant to the requirements of section 18 of the Act. The report covers fiscal year 2017 (October 1, 2016 - September 30, 2017), although some States may be reporting for a different fiscal year period (without duplicating a previous year's reported data). To assist Federal and State agencies with more accurate accounting of expenditures, a multi-species reporting category was added to the FY12 and subsequent reports. This option gives agencies another way to report expenditures when it is not possible to determine individual species costs for those species that are part of a multi-species conservation activity.

The term "State" as used in this report is defined in section 3(17) of the Act to mean any of the several States, the District of Columbia, the Commonwealth of Puerto Rico, American Samoa, the Virgin Islands, Guam, and the Trust Territory of the Pacific Islands.

The term "State agency" as used in this report is defined in section 3(18) of the Act to mean any State agency, department, board, commission, or other governmental entity which is responsible for the management and conservation of fish, plant, or wildlife resources within a State.

The term "conservation," as defined in section 3(3) of the Act, means the use of all methods and procedures necessary to bring any endangered or threatened species to the point where the measures provided for by the Act are no longer necessary--including, but not limited to, research, census, law enforcement, habitat acquisition and maintenance, propagation, live trapping, and transplantation. For the purposes of this report, the definition of conservation is further expanded to include any and all actions taken by Federal and State agencies on behalf of threatened or endangered species listed pursuant to the Act.

BACKGROUND

At the time the reporting requirement was enacted by Congress, the Service determined that a good faith effort was expected to develop methods to track and obtain data that are "reasonably identifiable." No extensive or extraordinary measures to develop exceptionally precise data were sought, nor were the reporting agencies expected to prorate staff salaries and other normal operational and maintenance costs not directed toward a particular species. The expenditures reported in this and the preceding reports do not necessarily constitute all expenditures made for each species.

This report represents a compilation of reasonably identifiable expenditures for the conservation of listed species reported independently by Federal and State agencies. Expenditures for FY 2017 are reported for 1,685 domestic threatened and endangered species of the 1,772 domestic species that were listed under United States jurisdiction in 50 CFR Part 17 during FY 2017 up through September 30, 2017 (Tables 1 and 3).

Expenditures are identified by listed entity, which may be a species¹, subspecies², Distinct Population Segment (DPS)³, or Evolutionarily Significant Unit (ESU)⁴, as identified in the Act's implementing regulations at 50 CFR 17.11 and 17.12.

These expenditures are reported separately for the Service and in aggregate for all other Federal agencies and the States (Tables 1 and 3).

For these annual reports, the Service requests that all reporting agencies submit their data directly into our Web-based data entry system to ensure accuracy, consistency, and efficiency. At the time of submission, the reporting agencies certify that their information is accurate. The information provided in this report, pursuant to section 18 of the Endangered Species Act, is presented as it was submitted by these agencies. We do not verify an agency's data after the data are submitted to us. In order to improve the Service's efficiency in producing future editions of this report, the Service will no longer enter a reporting agency's expenditures data into our Web-based system. All agencies must enter their own expenditures data into our Web-based system.

Expenditures for candidate species or other species not listed in 50 CFR Part 17 as of the end of each fiscal year are not included in this report. In addition, other types of expenses that are not incurred primarily for the conservation of listed species, such as those for litigation, are not reported.

¹ **Species** For purposes of the ESA, this term includes any species or subspecies of fish or wildlife or plants, and any Distinct Population Segment of any species of vertebrate fish or wildlife that interbreeds when mature.

² **Subspecies** A taxonomic rank below that of species, usually recognizing individuals that have certain heritable characteristics distinct from other subspecies of a species.

³ **Distinct Population Segment** A subdivision of a vertebrate species that is treated as a species for purposes of listing under the Endangered Species Act.

⁴ **Evolutionarily Significant Unit** A Pacific salmonid stock that is substantially reproductively isolated from other stocks of the same species and which represents an important part of the evolutionary legacy of the species; functionally equivalent to DPS for the purpose of listing under ESA.

SECTION 18(1), FEDERAL EXPENDITURES

Expenditures Reported by the Fish and Wildlife Service

The Service's operational categories of expenditures include fisheries, refuges, land acquisition, law enforcement, research, international conservation, and regional and field office operations for listing, recovery, consultation, migratory birds, and grants. All reasonably identifiable expenditures that could be attributed to individual species within these activities have been included.

The Federal assistance portion under section 6 of the Act for each species is included in the FWS Total in Table 1. The State share for each species for which a Federal contribution was made is included in the States Total column of Table 1.

A portion of the Service's funds for consultation, listing, and recovery are incorporated into salaries, operations, and maintenance costs. If these costs contributed to the conservation of listed species and are reasonably identifiable to specific species, they are included in this report.

Expenditures Reported by other Federal Agencies

The Director of the Service sends a request to other Federal agencies identified as likely to have species-specific expenditures under the Act to report for each fiscal year. The Service provides guidance identifying what constitutes reasonably identifiable expenditures and how these are to be reported (Appendix A). For FY 2017, the Service and 27 other Federal agencies identified expenditures for the conservation of individual threatened and endangered species. Results from all responding Federal agencies are included in Appendix B. The data identified by those responding Federal agencies for individual species and multi-species conservation activities are tabulated in the "Other Fed" columns of this report (Tables 1 and 3).

SECTION 18(2), STATE EXPENDITURES

The assistance of the Association of Fish and Wildlife Agencies is sought by the Service to help coordinate the reporting of reasonably identifiable expenditures made by the States. States report their data by species directly to the Service via our Web-based system. Collective results from all responding State agencies are included in Appendix C. For FY 2017, 36 State agencies identified expenditures for the conservation of individual threatened and endangered species.

SUMMARY OF DATA

Table 1 reports the activity-related expenditures that are reasonably identifiable to specific individual species as well as multi-species activity costs by the Service, other Federal agencies collectively, and the States collectively. Expenditures for land acquisition (Table 3) are not included in this tabulation. Species are grouped in the same general order as they appear in 50 CFR Part 17. Table 1 also provides the rank for each species from Table 2.

Table 2 ranks each species in descending order of the species total from Table 1. A cumulative total of the activity-related, species-specific expenditures is also provided.

Table 3 displays land acquisition expenditures, reasonably identifiable to specific individual species, by the Service, other Federal agencies collectively, and the States collectively. Expenditures for land purchased for multi-species conservation activities are listed at the end of the table. Table 3 also provides the rank for land expenditures by species.

Table 4 shows the combined total expenditures for species and land acquisition by reporting Federal agency, the Service, and the States collectively. Expenditures for foreign species and multi-species conservation activities are included.

Total expenditures reported for domestic and foreign species in **FY 2017** were **\$1,346,111,912** of which **\$1,287,980,311** was reported by Federal agencies and **\$58,131,601** was reported by the States (Table 4).

COMPARISON TO PREVIOUS REPORTS

Over the years, these reports have been refined as our experience in planning and coordinating with the reporting agencies to track such expenditures has improved. A good faith effort is made to develop species-by-species expenditures for these reports, but due to differences in reporting methods, this report cannot be easily compared to prior expenditures reports. These year-to-year differences include changes in the number of Federal and State agencies reporting data, changes in how expenditures are calculated, changes in the agencies' abilities to track expenditures, and changes in the number of listed species. Therefore, real dollar increases or decreases cannot accurately be calculated or inferred from the overall or individual totals.

TABLE 1. FY 2017 REPORTED EXPENDITURES FOR ENDANGERED AND THREATENED SPECIES

The list of species in Table 1 essentially follows the taxonomic and nomenclatural conventions used in 50 CFR Part 17 for the identification and sequence of species' names. For ease of reference, the alphabetized, inverted common name is provided within each group (usually Class) of animals and plants. Thus, the gray bat is found with other bats under "Bat, gray" within mammals, and Tar River spiny mussel is found under "Spiny mussel, Tar River" within the clams. Groups of animals follow the sequence: Mammals, Birds, Reptiles, Amphibians, Fishes, and Invertebrates, which include Clams, Snails, Insects, Arachnids, and Crustaceans. Groups of plants follow the sequence: Flowering Plants, Conifers and Cycads, Ferns and Allies, and Lichens.

For additional information about the names, historical range, or status of an endangered or threatened species, the species can be found in the Code of Federal Regulations at [50 CFR 17.11 \(list of wildlife\)](#) or [17.12 \(list of plants\)](#), or through our Web site at <http://www.fws.gov/angered>. The STATUS column indicates whether the species is endangered (E), threatened (T), experimental (EXPN), or threatened by similarity of appearance (SAT).

Fish and Wildlife Service expenditures (FWS Total) and other Federal agency expenditures (Other Fed) are totaled in the "Federal Total" column. The States Total is the amount reported to the Service directly by the States. None of these reported totals includes costs for land acquisition. (See Table 3 for reported land acquisition expenditures.)

The RANK is from Table 2 and indicates each species' rank by identifiable expenditures, where #1 is the highest expenditure.

Multi-species conservation activity-related expenditures are tabulated at the end of Table 1.

**TABLE 1. REPORTED EXPENDITURES FOR FY 2017 FOR ENDANGERED AND THREATENED SPECIES,
NOT INCLUDING LAND ACQUISITION COSTS**

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
Mammals	1426	Addax (<i>Addax nasomaculatus</i>) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	1040	Ass, Asian wild (<i>Equus hemionus</i>) - Wherever found	E	\$0	\$12,500	\$12,500	\$0	\$12,500
	1423	Babirusa (<i>Babirusa babirusa</i>) - Wherever found	E	\$0	\$300	\$300	\$0	\$300
	165	Bat, Florida bonneted (<i>Eumops floridanus</i>) - Wherever found	E	\$369,500	\$399,334	\$768,834	\$33,965	\$802,799
	133	Bat, gray (<i>Myotis grisescens</i>) - Wherever found	E	\$300,780	\$646,723	\$947,503	\$127,190	\$1,074,693
	84	Bat, Hawaiian hoary (<i>Lasiurus cinereus semotus</i>) - Wherever found	E	\$1,199,808	\$863,036	\$2,062,844	\$0	\$2,062,844
	42	Bat, Indiana (<i>Myotis sodalis</i>) - Wherever found	E	\$2,038,556	\$3,489,421	\$5,527,977	\$254,874	\$5,782,851
	326	Bat, lesser long-nosed (<i>Leptonycteris curasoae yerbabuena</i>) - Wherever found	E	\$40,500	\$152,674	\$193,174	\$19,855	\$213,029
	1380	Bat, little Mariana fruit (<i>Pteropus tokudae</i>) - Wherever found	E	\$2,104	\$0	\$2,104	\$0	\$2,104
	316	Bat, Mariana fruit (=Mariana flying fox) (<i>Pteropus mariannus mariannus</i>) - Wherever found	T	\$95,171	\$143,282	\$238,453	\$0	\$238,453
	509	Bat, Mexican long-nosed (<i>Leptonycteris nivalis</i>) - Wherever found	E	\$7,500	\$41,843	\$49,343	\$41,155	\$90,498
	39	Bat, Northern long-eared (<i>Myotis septentrionalis</i>) - Wherever found	T	\$1,641,693	\$5,028,226	\$6,669,919	\$295,751	\$6,965,670
	468	Bat, Ozark big-eared (<i>Corynorhinus</i> (=Plecotus) <i>townsendii ingens</i>) - Wherever found	E	\$98,500	\$24,883	\$123,383	\$3,780	\$127,163
	1161	Bat, Pacific sheath-tailed (<i>Emballonura semicaudata rotensis</i>) - Wherever found	E	\$9,573	\$0	\$9,573	\$0	\$9,573
	301	Bat, Virginia big-eared (<i>Corynorhinus</i> (=Plecotus) <i>townsendii virginianus</i>) - Wherever found	E	\$66,111	\$194,512	\$260,623	\$1,500	\$262,123
	1333	Bear, brown (<i>Ursus arctos pruinosus</i>) - Wherever found	E	\$0	\$4,800	\$4,800	\$0	\$4,800
	40	Bear, grizzly (<i>Ursus arctos horribilis</i>) - U.S.A., conterminous (lower 48) States, except where listed as an experimental population	T	\$3,658,509	\$2,627,323	\$6,285,832	\$629,465	\$6,915,297
	1141	Bear, grizzly (<i>Ursus arctos horribilis</i>) - Bitterroot Grizzly Bear Nonessential Experimental Population Area	EXPN	\$0	\$10,000	\$10,000	\$0	\$10,000
	60	Bear, polar (<i>Ursus maritimus</i>) - Wherever found	T	\$1,577,500	\$2,349,146	\$3,926,646	\$73,270	\$3,999,916
	311	Bison, wood (<i>Bison bison athabasca</i>) - Wherever found	T	\$8,500	\$200	\$8,700	\$232,863	\$241,563
	202	Caribou, Woodland (<i>Rangifer tarandus caribou</i>) - Selkirk Mountain population	E	\$592,969	\$2,400	\$595,369	\$20,901	\$616,270
	1394	Cat, leopard (<i>Prionailurus</i> (=Felis) <i>bengalensis bengalensis</i>) - Wherever found	E	\$1,400	\$200	\$1,600	\$0	\$1,600
	1426	Cat, tiger (<i>Leopardus</i> (=Felis) <i>tigrinus</i>) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	352	Cheetah (<i>Acinonyx jubatus</i>) - Wherever found	E	\$0	\$186,050	\$186,050	\$0	\$186,050
	120	Chimpanzee (<i>Pan troglodytes</i>) - Wherever found	E	\$1,285,751	\$0	\$1,285,751	\$0	\$1,285,751
	378	Chimpanzee, pygmy (<i>Pan paniscus</i>) - Wherever found	E	\$166,409	\$0	\$166,409	\$0	\$166,409
	273	Deer, Columbian white-tailed (<i>Odocoileus virginianus leucurus</i>) - Columbia River DPS	E	\$334,036	\$6,700	\$340,736	\$5,000	\$345,736
	387	Deer, Eld's brow-antlered (<i>Cervus eldi</i>) - Wherever found	E	\$98,200	\$59,200	\$157,400	\$0	\$157,400
	103	Deer, key (<i>Odocoileus virginianus clavium</i>) - Wherever found	E	\$666,000	\$1,007,635	\$1,673,635	\$0	\$1,673,635
	1343	Deer, musk (<i>Moschus</i> spp. (all species)) - Afghanistan, Bhutan, Burma, China [Tibet, Yunnan], India, Nepal, Pakistan, Sikkim	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	1232	Dog, African wild (<i>Lycan pictus</i>) - Wherever found	E	\$0	\$8,000	\$8,000	\$0	\$8,000
	1040	Drill (<i>Mandrillus</i> (=Papio) <i>leucophaeus</i>) - Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1427	Dugong (<i>Dugong dugon</i>) - Wherever found	E	\$0	\$100	\$100	\$0	\$100

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	51	Elephant, African (<i>Loxodonta africana</i>) - Wherever found	T	\$4,674,200	\$0	\$4,674,200	\$0	\$4,674,200
	83	Elephant, Asian (<i>Elephas maximus</i>) - Wherever found	E	\$1,517,593	\$592,300	\$2,109,893	\$0	\$2,109,893
	66	Ferret, black-footed (<i>Mustela nigripes</i>) - Wherever found, except where listed as an experimental population	E	\$2,300,641	\$763,262	\$3,063,903	\$188,125	\$3,252,028
	186	Ferret, black-footed (<i>Mustela nigripes</i>) - U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	EXPN	\$10,500	\$179,178	\$189,678	\$526,370	\$716,048
	122	Fox, San Joaquin kit (<i>Vulpes macrotis mutica</i>) - wherever found	E	\$793,705	\$444,463	\$1,238,168	\$0	\$1,238,168
	757	Fox, Santa Catalina Island (<i>Urocyon littoralis catalinae</i>) - Wherever found	T	\$3,190	\$29,680	\$32,870	\$0	\$32,870
	529	Gazelle, dama (<i>Gazella dama</i>) - Wherever found	E	\$0	\$79,450	\$79,450	\$0	\$79,450
	1423	Gazelle, slender-horned (<i>Gazella leptoceros</i>) - Wherever found	E	\$0	\$300	\$300	\$0	\$300
	199	Gibbons (<i>Hylobates</i> spp. (including <i>Nomascus</i>)) - Wherever found	E	\$621,049	\$700	\$621,749	\$0	\$621,749
	131	Gorilla (<i>Gorilla gorilla</i>) - Wherever found	E	\$1,123,231	\$300	\$1,123,531	\$0	\$1,123,531
	586	Horse, Przewalski's (<i>Equus przewalskii</i>) - Wherever found	E	\$0	\$61,950	\$61,950	\$0	\$61,950
	547	Jaguar (<i>Panthera onca</i>) - Wherever found	E	\$30,000	\$32,600	\$62,600	\$11,702	\$74,302
	517	Jaguarundi, Guatemalan (<i>Herpailurus</i> (=Felis) <i>yagouaroundi fossata</i>) - Mexico, Nicaragua	E	\$85,729	\$0	\$85,729	\$0	\$85,729
	1046	Jaguarundi, Gulf Coast (<i>Herpailurus</i> (=Felis) <i>yagouaroundi cacomitli</i>) - Wherever found	E	\$6,100	\$6,200	\$12,300	\$0	\$12,300
	226	Jaguarundi, Panamanian (<i>Herpailurus</i> (=Felis) <i>yagouaroundi panamensis</i>) - Wherever found	E	\$490,600	\$0	\$490,600	\$0	\$490,600
	260	Kangaroo rat, Fresno (<i>Dipodomys nitratoides exilis</i>) - Wherever found	E	\$25,740	\$366,002	\$391,742	\$0	\$391,742
	223	Kangaroo rat, giant (<i>Dipodomys ingens</i>) - Wherever found	E	\$118,258	\$392,340	\$510,598	\$0	\$510,598
	739	Kangaroo rat, Morro Bay (<i>Dipodomys heermanni morroensis</i>) - Wherever found	E	\$34,632	\$0	\$34,632	\$0	\$34,632
	220	Kangaroo rat, San Bernardino Merriam's (<i>Dipodomys merriami parvus</i>) - Wherever found	E	\$386,870	\$134,602	\$521,472	\$0	\$521,472
	251	Kangaroo rat, Stephens' (<i>Dipodomys stephensi</i> (incl. <i>D. cascus</i>)) - Wherever found	E	\$54,340	\$363,274	\$417,614	\$0	\$417,614
	467	Kangaroo rat, Tipton (<i>Dipodomys nitratoides nitratoides</i>) - Wherever found	E	\$97,900	\$29,396	\$127,296	\$0	\$127,296
	1426	Koala (<i>Phascolarctos cinereus</i>) - Australia	T	\$0	\$200	\$200	\$0	\$200
	1426	Langur, Francois' (<i>Trachypithecus</i> (=Presbytis) <i>francoisi</i>) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	1426	Lechwe, red (<i>Kobus leche</i>) - Wherever found	T	\$0	\$200	\$200	\$0	\$200
	1403	Lemurs (<i>Lemuridae</i> (incl. genera <i>Lemur</i> , <i>Phaner</i> , <i>Hapalemur</i> , <i>Lepilemur</i> , <i>Microcebus</i> , <i>Allocebus</i> , <i>Cheirogaleus</i> , <i>Varecia</i>)) - Wherever found	E	\$0	\$1,100	\$1,100	\$0	\$1,100
	1232	Leopard (<i>Panthera pardus</i>) - except where threatened	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	395	Leopard (<i>Panthera pardus</i>) - Gabon to Kenya and southward	T	\$152,177	\$0	\$152,177	\$0	\$152,177
	565	Leopard, clouded (<i>Neofelis nebulosa</i>) - Wherever found	E	\$0	\$69,800	\$69,800	\$0	\$69,800
	498	Leopard, snow (<i>Uncia</i> (=Panthera) <i>uncia</i>) - Wherever found	E	\$99,200	\$300	\$99,500	\$0	\$99,500
	242	Lion (<i>Panthera leo</i> ssp. <i>leo</i>) - Wherever found	E	\$457,071	\$200	\$457,271	\$0	\$457,271
	72	Lynx, Canada (<i>Lynx canadensis</i>) - Wherever Found in Contiguous U.S.	T	\$1,899,398	\$625,500	\$2,524,898	\$247,135	\$2,772,033
	1194	Manatee, West African (<i>Trichechus senegalensis</i>) - Wherever found	T	\$0	\$9,000	\$9,000	\$0	\$9,000
	25	Manatee, West Indian (<i>Trichechus manatus</i>) - Wherever found	E	\$2,555,100	\$2,557,439	\$5,112,539	\$5,032,819	\$10,145,358

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1020	Mandrill (<i>Mandrillus</i> (=Papio) sphinx) - Wherever found	E	\$12,500	\$500	\$13,000	\$0	\$13,000
	1040	Mangabey, white-collared (<i>Cercocebus torquatus</i>) - Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1040	Monkey, black colobus (<i>Colobus satanas</i>) - Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1426	Monkey, Diana (<i>Cercopithecus diana</i>) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	1040	Monkey, Preuss' red colobus (<i>Procolobus</i> (=Colobus) preussi (=badius p.)) - Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1040	Monkey, red-eared nose-spotted (<i>Cercopithecus erythrotis</i>) - Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	593	Mountain beaver, Point Arena (<i>Aplodontia rufa nigra</i>) - Wherever found	E	\$56,100	\$5,000	\$61,100	\$0	\$61,100
	449	Mouse, Alabama beach (<i>Peromyscus polionotus ammobates</i>) - Wherever found	E	\$120,500	\$13,907	\$134,407	\$0	\$134,407
	499	Mouse, Anastasia Island beach (<i>Peromyscus polionotus phasma</i>) - Wherever found	E	\$93,500	\$700	\$94,200	\$1,115	\$95,315
	485	Mouse, Choctawhatchee beach (<i>Peromyscus polionotus allophrys</i>) - Wherever found	E	\$15,500	\$81,474	\$96,974	\$8,916	\$105,890
	943	Mouse, Key Largo cotton (<i>Peromyscus gossypinus allapaticola</i>) - Wherever found	E	\$15,000	\$1,829	\$16,829	\$0	\$16,829
	281	Mouse, New Mexico meadow jumping (<i>Zapus hudsonius luteus</i>) - Wherever found	E	\$90,813	\$200,510	\$291,323	\$41,372	\$332,695
	239	Mouse, Pacific pocket (<i>Perognathus longimembris pacificus</i>) - Wherever found	E	\$70,840	\$391,424	\$462,264	\$0	\$462,264
	640	Mouse, Perdido Key beach (<i>Peromyscus polionotus trissyllepsis</i>) - Wherever found	E	\$5,000	\$39,420	\$44,420	\$7,802	\$52,222
	235	Mouse, Preble's meadow jumping (<i>Zapus hudsonius preblei</i>) - wherever found	T	\$51,658	\$392,011	\$443,669	\$31,263	\$474,932
	144	Mouse, salt marsh harvest (<i>Reithrodontomys raviventris</i>) - wherever found	E	\$688,161	\$284,900	\$973,061	\$0	\$973,061
	404	Mouse, southeastern beach (<i>Peromyscus polionotus niveiventris</i>) - wherever found	T	\$99,500	\$48,600	\$148,100	\$1,115	\$149,215
	495	Mouse, St. Andrew beach (<i>Peromyscus polionotus peninsularis</i>) - Wherever found	E	\$13,000	\$84,343	\$97,343	\$3,344	\$100,687
	155	Ocelot (<i>Leopardus</i> (=Felis) pardalis) - wherever found	E	\$788,200	\$43,800	\$832,000	\$8,609	\$840,609
	298	Orangutan (<i>Pongo pygmaeus</i>) - Wherever found	E	\$275,335	\$200	\$275,535	\$0	\$275,535
	346	Oryx, scimitar-horned (<i>Oryx dammah</i>) - Wherever found	E	\$0	\$189,750	\$189,750	\$0	\$189,750
	823	Otter, giant (<i>Pteronura brasiliensis</i>) - South America	E	\$24,745	\$0	\$24,745	\$0	\$24,745
	126	Otter, Northern Sea (<i>Enhydra lutris kenyonii</i>) - Southwest Alaska DPS	T	\$1,062,000	\$124,982	\$1,186,982	\$0	\$1,186,982
	151	Otter, southern sea (<i>Enhydra lutris nereis</i>) - Wherever found	T	\$23,636	\$847,359	\$870,995	\$0	\$870,995
	234	Panda, giant (<i>Ailuropoda melanoleuca</i>) - Wherever found	E	\$0	\$478,300	\$478,300	\$0	\$478,300
	339	Pangolin, Temnick's ground (<i>Manis temminckii</i>) - Wherever found	E	\$199,797	\$0	\$199,797	\$0	\$199,797
	68	Panther, Florida (<i>Puma</i> (=Felis) concolor coryi) - Wherever found	E	\$1,058,500	\$1,138,685	\$2,197,185	\$804,101	\$3,001,286
	266	Pocket gopher, Olympia (<i>Thomomys mazama pugetensis</i>) - Wherever found	T	\$272,401	\$83,000	\$355,401	\$5,000	\$360,401
	246	Pocket gopher, Roy Prairie (<i>Thomomys mazama glacialis</i>) - Wherever found	T	\$169,531	\$266,666	\$436,197	\$5,000	\$441,197
	312	Pocket gopher, Tenino (<i>Thomomys mazama tumuli</i>) - Wherever found	T	\$236,431	\$0	\$236,431	\$5,000	\$241,431

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	205	Pocket gopher, Yelm (<i>Thomomys mazama yelmensis</i>) - Wherever found	T	\$576,531	\$0	\$576,531	\$20,000	\$596,531
	74	Prairie dog, Utah (<i>Cynomys parvidens</i>) - Wherever found	T	\$876,000	\$1,630,948	\$2,506,948	\$70,000	\$2,576,948
	1359	Pronghorn, peninsular (<i>Antilocapra americana peninsularis</i>) - Mexico (Baja California)	E	\$0	\$3,000	\$3,000	\$0	\$3,000
	321	Pronghorn, Sonoran (<i>Antilocapra americana sonoriensis</i>) - Wherever found, except where listed as an experimental population	E	\$0	\$232,500	\$232,500	\$0	\$232,500
	123	Pronghorn, Sonoran (<i>Antilocapra americana sonoriensis</i>) - U.S.A. (AZ), Mexico	EXPN	\$306,500	\$110,000	\$416,500	\$816,763	\$1,233,263
	929	Puma (=cougar), eastern (<i>Puma (=Felis concolor cougar)</i>) - Wherever found	E	\$10,391	\$6,361	\$16,752	\$900	\$17,652
	1359	Puma (=mountain lion) (<i>Puma (=Felis concolor</i> (all subsp. except <i>coryi</i>)) - FL	SAT	\$0	\$3,000	\$3,000	\$0	\$3,000
	265	Rabbit, Columbia Basin Pygmy (<i>Brachylagus idahoensis</i>) - Columbia Basin DPS	E	\$261,912	\$6,200	\$268,112	\$100,689	\$368,801
	371	Rabbit, Lower Keys marsh (<i>Sylvilagus palustris hefneri</i>) - Wherever found	E	\$22,500	\$147,405	\$169,905	\$0	\$169,905
	631	Rabbit, riparian brush (<i>Sylvilagus bachmani riparius</i>) - Wherever found	E	\$50,410	\$2,688	\$53,098	\$0	\$53,098
	129	Rhinoceros, black (<i>Diceros bicornis</i>) - Africa	E	\$1,143,586	\$300	\$1,143,886	\$0	\$1,143,886
	465	Rhinoceros, great Indian (<i>Rhinoceros unicornis</i>) - Nepal, India	E	\$127,581	\$0	\$127,581	\$0	\$127,581
	368	Rhinoceros, Javan (<i>Rhinoceros sondaicus</i>) - Sikkim, Thailand, Malaysia, Indonesia, Indochina, Burma, Bangladesh	E	\$170,916	\$0	\$170,916	\$0	\$170,916
	257	Rhinoceros, Southern white (<i>Ceratotherium simum ssp. simum</i>) - Wherever found	SAT	\$405,775	\$300	\$406,075	\$0	\$406,075
	254	Rhinoceros, Sumatran (<i>Dicerorhinus sumatrensis</i>) - Wherever found	E	\$410,106	\$0	\$410,106	\$0	\$410,106
	808	Rice rat (<i>Oryzomys palustris natator</i>) - lower FL Keys	E	\$15,000	\$11,420	\$26,420	\$0	\$26,420
	854	Seal, bearded (<i>Erignathus barbatus nauticus</i>) - Okhotsk DPS	T	\$0	\$22,186	\$22,186	\$0	\$22,186
	343	Seal, Guadalupe fur (<i>Arctocephalus townsendi</i>) - Wherever found	T	\$0	\$191,706	\$191,706	\$0	\$191,706
	56	Seal, Hawaiian monk (<i>Monachus schauinslandi</i>) - Wherever found	E	\$81,770	\$4,101,134	\$4,182,904	\$0	\$4,182,904
	65	Sea lion, Steller (<i>Eumetopias jubatus</i>) - Western DPS	E	\$31,400	\$2,400,258	\$2,431,658	\$1,139,184	\$3,570,842
	1423	Seal, Ringed (<i>Phoca (=Pusa) hispida botnica</i>) - Wherever found	T	\$0	\$300	\$300	\$0	\$300
	320	Seal, Ringed (<i>Phoca (=Pusa) hispida hispida</i>) - Wherever found	T	\$0	\$60	\$60	\$234,029	\$234,089
	1406	Seal, spotted (<i>Phoca largha</i>) - Southern DPS	T	\$0	\$1,000	\$1,000	\$0	\$1,000
	473	Sheep, Peninsular bighorn (<i>Ovis canadensis nelsoni</i>) - Peninsular CA pop.	E	\$110,220	\$10,500	\$120,720	\$0	\$120,720
	277	Sheep, Sierra Nevada bighorn (<i>Ovis canadensis sierrae</i>) - Sierra Nevada	E	\$142,340	\$197,128	\$339,468	\$0	\$339,468
	503	Shrew, Buena Vista Lake ornate (<i>Sorex ornatus relictus</i>) - Wherever found	E	\$69,300	\$23,516	\$92,816	\$0	\$92,816
	1423	Siamang (<i>Symphalangus syndactylus</i>) - Wherever found	E	\$0	\$300	\$300	\$0	\$300
	677	Squirrel, Carolina northern flying (<i>Glaucomys sabrinus coloratus</i>) - Wherever found	E	\$30,500	\$16,104	\$46,604	\$0	\$46,604
	694	Squirrel, Mount Graham red (<i>Tamiasciurus hudsonicus grahamensis</i>) - Wherever found	E	\$14,000	\$100	\$14,100	\$28,246	\$42,346
	261	Squirrel, Northern Idaho Ground (<i>Uroditellus brunneus</i>) - Wherever found	T	\$303,528	\$30,692	\$334,220	\$54,449	\$388,669
	1418	Tamarin, cotton-top (<i>Saguinus oedipus</i>) - Wherever found	E	\$0	\$500	\$500	\$0	\$500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	710	Tamarin, golden-rumped (<i>Leontopithecus</i> spp.) - Wherever found	E	\$0	\$39,900	\$39,900	\$0	\$39,900
	450	Tapir, Central American (<i>Tapirus bairdii</i>) - Wherever found	E	\$133,574	\$200	\$133,774	\$0	\$133,774
	1426	Tapir, South American (=Brazilian) (<i>Tapirus terrestris</i>) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	90	Tiger (<i>Panthera tigris</i>) - Wherever found	E	\$1,717,273	\$295,500	\$2,012,773	\$0	\$2,012,773
	222	Vole, Amargosa (<i>Microtus californicus scirpensis</i>) - Wherever found	E	\$331,117	\$180,000	\$511,117	\$0	\$511,117
	1115	Vole, Florida salt marsh (<i>Microtus pennsylvanicus dukecampbelli</i>) - Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	1348	Vole, Hualapai Mexican (<i>Microtus mexicanus hualpaiensis</i>) - Wherever found	E	\$3,000	\$100	\$3,100	\$1,000	\$4,100
	63	Whale, beluga (<i>Delphinapterus leucas</i>) - Cook Inlet DPS	E	\$0	\$3,568,804	\$3,568,804	\$34,548	\$3,603,352
	71	Whale, blue (<i>Balaenoptera musculus</i>) - Wherever found	E	\$800	\$2,780,024	\$2,780,824	\$0	\$2,780,824
	117	Whale, bowhead (<i>Balaena mysticetus</i>) - Wherever found	E	\$0	\$901,111	\$901,111	\$401,045	\$1,302,156
	167	Whale, false killer (<i>Pseudorca crassidens</i>) - Main Hawaiian Islands Insular DPS	E	\$0	\$793,495	\$793,495	\$0	\$793,495
	75	Whale, finback (<i>Balaenoptera physalus</i>) - Wherever found	E	\$0	\$2,541,447	\$2,541,447	\$0	\$2,541,447
	290	Whale, gray (<i>Eschrichtius robustus</i>) - Western North Pacific DPS	E	\$0	\$291,166	\$291,166	\$0	\$291,166
	147	Whale, humpback (<i>Megaptera novaeangliae</i>) - Western North Pacific	E	\$0	\$956,621	\$956,621	\$0	\$956,621
	108	Whale, humpback (<i>Megaptera novaeangliae</i>) - Mexico DPS	T	\$48,663	\$1,509,421	\$1,558,084	\$0	\$1,558,084
	114	Whale, humpback (<i>Megaptera novaeangliae</i>) - Central America DPS	E	\$0	\$1,411,338	\$1,411,338	\$0	\$1,411,338
	79	Whale, killer (<i>Orcinus orca</i>) - Southern Resident DPS	E	\$0	\$2,329,103	\$2,329,103	\$8,385	\$2,337,488
	29	Whale, North Atlantic Right (<i>Eubalaena glacialis</i>) - Wherever found	E	\$0	\$8,848,633	\$8,848,633	\$94,564	\$8,943,197
	158	Whale, North Pacific Right (<i>Eubalaena japonica</i>) - Wherever found	E	\$0	\$832,185	\$832,185	\$0	\$832,185
	101	Whale, Sei (<i>Balaenoptera borealis</i>) - Wherever found	E	\$0	\$1,752,980	\$1,752,980	\$0	\$1,752,980
	94	Whale, sperm (<i>Physeter catodon</i> (=macrocephalus)) - Wherever found	E	\$3,600	\$1,962,323	\$1,965,923	\$0	\$1,965,923
	49	Wolf, gray (<i>Canis lupus</i>) - U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	E	\$3,377,433	\$1,468,952	\$4,846,385	\$10,770	\$4,857,155
	208	Wolf, gray (<i>Canis lupus</i>) - MN	T	\$20,000	\$556,700	\$576,700	\$0	\$576,700
	308	Wolf, gray (<i>Canis lupus</i>) - WY, EXPN population	EXPN	\$1,871	\$246,000	\$247,871	\$0	\$247,871
	284	Wolf, maned (<i>Chrysocyon brachyurus</i>) - Wherever found	E	\$0	\$302,000	\$302,000	\$0	\$302,000
	575	Wolf, Mexican (<i>Canis lupus baileyi</i>) - Wherever found, except where listed as an experimental population	E	\$0	\$65,700	\$65,700	\$0	\$65,700
	69	Wolf, Mexican (<i>Canis lupus baileyi</i>) - U.S.A. (portions of AZ and NM)see 17.84(k)	EXPN	\$2,525,888	\$80,200	\$2,606,088	\$368,975	\$2,975,063
	314	Wolf, red (<i>Canis rufus</i>) - Wherever found, except where listed as an experimental population	E	\$240,000	\$0	\$240,000	\$0	\$240,000
	170	Wolf, red (<i>Canis rufus</i>) - U.S.A. (portions of NC and TN)	EXPN	\$772,500	\$3,600	\$776,100	\$0	\$776,100
	789	Woodrat, Key Largo (<i>Neotoma floridana smalli</i>) - Wherever found	E	\$23,000	\$5,339	\$28,339	\$0	\$28,339
	1092	Woodrat, riparian (=San Joaquin Valley) (<i>Neotoma fuscipes riparia</i>) - Wherever found	E	\$8,350	\$2,520	\$10,870	\$0	\$10,870
Mammals subtotal				\$51,691,777	\$71,591,552	\$123,283,329	\$12,051,904	\$135,335,233
Birds	285	Akekee (<i>Loxops caeruleirostris</i>) - Wherever found	E	\$236,484	\$64,215	\$300,699	\$0	\$300,699
	536	Akepa, Hawaii (<i>Loxops coccineus</i>) - Wherever found	E	\$50,844	\$26,635	\$77,479	\$0	\$77,479

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	620	Akepa, Maui (<i>Loxops ochraceus</i>) - Wherever found	E	\$19,290	\$36,250	\$55,540	\$0	\$55,540
	1399	Akialoa, Kauai (honeycreeper) (<i>Akialoa stejnegeri</i>) - Wherever found	E	\$1,362	\$0	\$1,362	\$0	\$1,362
	568	Akiapolaau (<i>Hemignathus wilsoni</i>) - Wherever found	E	\$42,585	\$26,635	\$69,220	\$0	\$69,220
	259	Akikiki (<i>Oreomystis bairdi</i>) - Wherever found	E	\$328,831	\$64,215	\$393,046	\$0	\$393,046
	353	Albatross, short-tailed (<i>Phoebastria</i> (=Diomedea) <i>albatrus</i>) - Wherever found	E	\$141,525	\$43,386	\$184,911	\$0	\$184,911
	483	Blackbird, yellow-shouldered (<i>Agelaius xanthomus</i>) - Wherever found	E	\$84,500	\$25,000	\$109,500	\$0	\$109,500
	304	Bobwhite, masked (quail) (<i>Colinus virginianus ridgwayi</i>) - Wherever found	E	\$245,164	\$10,600	\$255,764	\$2,687	\$258,451
	148	Caracara, Audubon's crested (<i>Polyborus plancus audubonii</i>) - FL pop.	T	\$212,500	\$685,745	\$898,245	\$0	\$898,245
	62	Condor, California (<i>Gymnogyps californianus</i>) - U.S.A. only, except where listed as an experimental population	E	\$3,140,711	\$708,674	\$3,849,385	\$0	\$3,849,385
	249	Condor, California (<i>Gymnogyps californianus</i>) - U.S.A. (specific portions of Arizona, Nevada, and Utah)	EXPN	\$77,830	\$88,335	\$166,165	\$252,240	\$418,405
	116	Coot, Hawaiian (<i>Fulica americana alai</i>) - Wherever found	E	\$675,699	\$682,064	\$1,357,763	\$0	\$1,357,763
	1428	Crane, black-necked (<i>Grus nigricollis</i>) - Wherever found	E	\$0	\$40	\$40	\$0	\$40
	1428	Crane, hooded (<i>Grus monacha</i>) - Wherever found	E	\$0	\$40	\$40	\$0	\$40
	1429	Crane, Japanese (<i>Grus japonensis</i>) - Wherever found	E	\$0	\$20	\$20	\$0	\$20
	190	Crane, Mississippi sandhill (<i>Grus canadensis pulla</i>) - Wherever found	E	\$686,000	\$16,013	\$702,013	\$0	\$702,013
	1428	Crane, white-naped (<i>Grus vipio</i>) - Wherever found	E	\$0	\$40	\$40	\$0	\$40
	132	Crane, whooping (<i>Grus americana</i>) - Wherever found, except where listed as an experimental population	E	\$857,025	\$224,274	\$1,081,299	\$12,700	\$1,093,999
	291	Crane, whooping (<i>Grus americana</i>) - U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	EXPN	\$46,500	\$227,850	\$274,350	\$14,666	\$289,016
	118	Crane, whooping (<i>Grus americana</i>) - U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	EXPN	\$900	\$1,292,304	\$1,293,204	\$0	\$1,293,204
	289	Crane, whooping (<i>Grus americana</i>) - U.S.A (Southwestern Louisiana)	EXPN	\$76,000	\$215,250	\$291,250	\$0	\$291,250
	531	Creeper, Hawaii (<i>Oreomystis mana</i>) - Wherever found	E	\$51,858	\$26,635	\$78,493	\$0	\$78,493
	1355	Creeper, Molokai (<i>Paroreomyza flammea</i>) - Wherever found	E	\$3,390	\$0	\$3,390	\$0	\$3,390
	991	Creeper, Oahu (<i>Paroreomyza maculata</i>) - Wherever found	E	\$14,220	\$0	\$14,220	\$0	\$14,220
	121	Crow, Hawaiian (=alala) (<i>Corvus hawaiiensis</i>) - Wherever found	E	\$1,259,686	\$25,000	\$1,284,686	\$0	\$1,284,686
	125	Crow, Mariana (=aga) (<i>Corvus kubaryi</i>) - Wherever found	E	\$956,075	\$240,825	\$1,196,900	\$0	\$1,196,900
	1396	Crow, white-necked (<i>Corvus leucognaphalus</i>) - Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	44	Cuckoo, yellow-billed (<i>Coccyzus americanus</i>) - Western U.S. DPS	T	\$561,378	\$4,785,947	\$5,347,325	\$34,262	\$5,381,587
	1187	Curassow, blue-billed (<i>Crax alberti</i>) - Wherever found	E	\$0	\$9,100	\$9,100	\$0	\$9,100
	1391	Curlew, Eskimo (<i>Numenius borealis</i>) - Wherever found	E	\$500	\$1,200	\$1,700	\$0	\$1,700
	173	Duck, Hawaiian (=koloa) (<i>Anas wyvilliana</i>) - Wherever found	E	\$90,507	\$681,330	\$771,837	\$0	\$771,837
	390	Duck, Laysan (<i>Anas laysanensis</i>) - Wherever found	E	\$43,705	\$111,178	\$154,883	\$0	\$154,883
	458	Eagle, harpy (<i>Harpia harpyja</i>) - Wherever found	E	\$130,000	\$0	\$130,000	\$0	\$130,000
	142	Eider, spectacled (<i>Somateria fischeri</i>) - Wherever found	T	\$849,500	\$130,058	\$979,558	\$0	\$979,558
	179	Eider, Steller's (<i>Polysticta stelleri</i>) - AK breeding pop.	T	\$604,000	\$154,512	\$758,512	\$0	\$758,512
	362	Elepaio, Oahu (<i>Chasiempis ibidis</i>) - Wherever found	E	\$35,657	\$140,567	\$176,224	\$0	\$176,224
	324	Falcon, Northern Aplomado (<i>Falco femoralis septentrionalis</i>) - Wherever found, except where listed as an experimental population	E	\$0	\$215,059	\$215,059	\$0	\$215,059

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	384	Falcon, Northern Aplomado (<i>Falco femoralis septentrionalis</i>) - U.S.A (AZ, NM)	EXPN	\$118,700	\$37,900	\$156,600	\$1,500	\$158,100
	1047	Finch, Laysan (honeycreeper) (<i>Telespyza cantans</i>) - Wherever found	E	\$12,271	\$0	\$12,271	\$0	\$12,271
	949	Finch, Nihoa (honeycreeper) (<i>Telespyza ultima</i>) - Wherever found	E	\$16,327	\$0	\$16,327	\$0	\$16,327
	30	Flycatcher, southwestern willow (<i>Empidonax traillii extimus</i>) - Wherever found	E	\$461,246	\$8,442,027	\$8,903,273	\$35,691	\$8,938,964
	197	Gallinule, Hawaiian common (<i>Gallinula galeata sandvicensis</i>) - Wherever found	E	\$118,020	\$521,869	\$639,889	\$0	\$639,889
	91	Gnatcatcher, coastal California (<i>Polioptila californica californica</i>) - Wherever found	T	\$804,724	\$1,187,536	\$1,992,260	\$0	\$1,992,260
	87	Goose, Hawaiian (<i>Branta (=Nesochen) sandvicensis</i>) - Wherever found	E	\$1,166,799	\$872,539	\$2,039,338	\$0	\$2,039,338
	1411	Ground-Dove, Friendly (<i>Gallicolumba stairi</i>) - American Samoa DPS	E	\$226	\$500	\$726	\$0	\$726
	478	Hawk, Hawaiian (=Io) (<i>Buteo solitarius</i>) - Wherever found	E	\$72,900	\$44,600	\$117,500	\$0	\$117,500
	651	Hawk, Puerto Rican broad-winged (<i>Buteo platypterus brunnescens</i>) - Wherever found	E	\$35,500	\$15,000	\$50,500	\$0	\$50,500
	574	Hawk, Puerto Rican sharp-shinned (<i>Accipiter striatus venator</i>) - Wherever found	E	\$36,000	\$30,000	\$66,000	\$0	\$66,000
	545	Honeycreeper, (Akohekohe) crested (<i>Palmeria dolei</i>) - Wherever found	E	\$39,896	\$35,251	\$75,147	\$0	\$75,147
	104	Horned lark, streaked (<i>Eremophila alpestris strigata</i>) - Wherever found	T	\$581,345	\$1,085,496	\$1,666,841	\$2,000	\$1,668,841
	256	Kingfisher, Guam Micronesian (<i>Halcyon cinnamomina cinnamomina</i>) - Wherever found	E	\$200,467	\$209,074	\$409,541	\$0	\$409,541
	110	Kite, Everglade snail (<i>Rostrhamus sociabilis plumbeus</i>) - Wherever found	E	\$420,000	\$982,696	\$1,402,696	\$74,362	\$1,477,058
	135	Knot, red (<i>Calidris canutus rufa</i>) - Wherever found	T	\$465,290	\$507,323	\$972,613	\$50,060	\$1,022,673
	641	Megapode, Micronesian (<i>Megapodius laperouse</i>) - Wherever found	E	\$36,574	\$15,179	\$51,753	\$0	\$51,753
	675	Millerbird, Nihoa (old world warbler) (<i>Acrocephalus familiaris kingi</i>) - Wherever found	E	\$46,747	\$0	\$46,747	\$0	\$46,747
	391	Moorhen, Mariana common (<i>Gallinula chloropus guami</i>) - Wherever found	E	\$49,346	\$105,342	\$154,688	\$0	\$154,688
	47	Murrelet, marbled (<i>Brachyramphus marmoratus</i>) - U.S.A. (CA, OR, WA)	T	\$2,514,505	\$2,586,499	\$5,101,004	\$109,100	\$5,210,104
	1206	Nightjar, Puerto Rican (<i>Caprimulgus noctitherus</i>) - Wherever found	E	\$8,500	\$0	\$8,500	\$0	\$8,500
	954	Nukupuu, Kauai (<i>Hemignathus hanapepe</i>) - Wherever found	E	\$16,248	\$0	\$16,248	\$0	\$16,248
	905	Nukupuu, Maui (<i>Hemignathus affinis</i>) - Wherever found	E	\$19,019	\$0	\$19,019	\$0	\$19,019
	1399	'O'o, Kauai (honeyeater) (<i>Moho braccatus</i>) - Wherever found	E	\$1,362	\$0	\$1,362	\$0	\$1,362
	1299	'O'u (honeycreeper) (<i>Psittirostra psittacea</i>) - Wherever found	E	\$6,309	\$0	\$6,309	\$0	\$6,309
	184	Owl, Mexican spotted (<i>Strix occidentalis lucida</i>) - Wherever found	T	\$95,604	\$637,823	\$733,427	\$500	\$733,927
	23	Owl, northern spotted (<i>Strix occidentalis caurina</i>) - Wherever found	T	\$6,610,558	\$5,241,917	\$11,852,475	\$2,200	\$11,854,675
	98	Palila (honeycreeper) (<i>Loxioides bailleui</i>) - Wherever found	E	\$1,798,316	\$10,000	\$1,808,316	\$0	\$1,808,316
	382	Parrotbill, Maui (Kiwikiu) (<i>Pseudonestor xanthophrys</i>) - Wherever found	E	\$105,782	\$54,851	\$160,633	\$0	\$160,633
	112	Parrot, Puerto Rican (<i>Amazona vittata</i>) - Wherever found	E	\$1,445,000	\$0	\$1,445,000	\$0	\$1,445,000
	1406	Parrot, thick-billed (<i>Rhynchopsitta pachyrhyncha</i>) - Wherever found	E	\$1,000	\$0	\$1,000	\$0	\$1,000
	1133	Petrel, Bermuda (<i>Pterodroma cahow</i>) - Wherever found	E	\$0	\$10,118	\$10,118	\$0	\$10,118

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	96	Petrel, Hawaiian (<i>Pterodroma sandwichensis</i>) - Wherever found	E	\$936,144	\$930,578	\$1,866,722	\$0	\$1,866,722
	1164	Pigeon, Puerto Rican plain (<i>Columba inornata wetmorei</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	193	Plover, piping (<i>Charadrius melodus</i>) - [Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	E	\$522,200	\$123,158	\$645,358	\$9,947	\$655,305
	28	Plover, piping (<i>Charadrius melodus</i>) - [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	T	\$1,473,747	\$7,618,744	\$9,092,491	\$432,949	\$9,525,440
	52	Plover, western snowy (<i>Charadrius nivosus nivosus</i>) - Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	T	\$1,689,993	\$2,893,235	\$4,583,228	\$36,200	\$4,619,428
	1373	Po'ouli (honeycreeper) (<i>Melamprosops phaeosoma</i>) - Wherever found	E	\$2,376	\$0	\$2,376	\$0	\$2,376
	119	Prairie-chicken, Attwater's greater (<i>Tympanuchus cupido attwateri</i>) - Wherever found	E	\$1,278,788	\$10,700	\$1,289,488	\$0	\$1,289,488
	143	Rail, California clapper (<i>Rallus longirostris obsoletus</i>) - Wherever found	E	\$588,015	\$391,145	\$979,160	\$0	\$979,160
	279	Rail, Guam (<i>Rallus owstoni</i>) - Wherever found, except where listed as an experimental population	E	\$274,813	\$59,026	\$333,839	\$0	\$333,839
	623	Rail, Guam (<i>Rallus owstoni</i>) - Rota	EXPN	\$54,784	\$0	\$54,784	\$0	\$54,784
	255	Rail, light-footed clapper (<i>Rallus longirostris levipes</i>) - Wherever found	E	\$161,746	\$247,919	\$409,665	\$0	\$409,665
	81	Rail, Yuma clapper (<i>Rallus longirostris yumanensis</i>) - Wherever found	E	\$351,077	\$1,796,350	\$2,147,427	\$56,771	\$2,204,198
	64	Sage-grouse, Gunnison (<i>Centrocercus minimus</i>) - Wherever found	T	\$429,873	\$2,722,479	\$3,152,352	\$430,274	\$3,582,626
	99	Scrub-jay, Florida (<i>Aphelocoma coerulescens</i>) - Wherever found	T	\$524,000	\$1,158,600	\$1,682,600	\$124,496	\$1,807,096
	217	Shearwater, Newell's Townsend's (<i>Puffinus auricularis newelli</i>) - Wherever found	T	\$394,341	\$142,305	\$536,646	\$0	\$536,646
	76	Shrike, San Clemente loggerhead (<i>Lanius ludovicianus mearnsi</i>) - Wherever found	E	\$28,619	\$2,383,719	\$2,412,338	\$0	\$2,412,338
	130	Sparrow, Cape Sable seaside (<i>Ammodramus maritimus mirabilis</i>) - Wherever found	E	\$478,000	\$658,840	\$1,136,840	\$0	\$1,136,840
	211	Sparrow, Florida grasshopper (<i>Ammodramus savannarum floridanus</i>) - Wherever found	E	\$244,000	\$234,624	\$478,624	\$79,954	\$558,578
	161	Sparrow, San Clemente sage (<i>Amphispiza belli clementeae</i>) - Wherever found	T	\$20,809	\$804,173	\$824,982	\$0	\$824,982
	113	Stilt, Hawaiian (<i>Himantopus mexicanus knudseni</i>) - Wherever found	E	\$670,212	\$759,100	\$1,429,312	\$0	\$1,429,312
	93	Stork, wood (<i>Mycteria americana</i>) - AL, FL, GA, MS, NC, SC	T	\$330,500	\$1,612,131	\$1,942,631	\$30,000	\$1,972,631
	318	Storm-petrel, band-rumped (<i>Oceanodroma castro</i>) - USA (HI)	E	\$16,180	\$219,431	\$235,611	\$0	\$235,611
	206	Swiftlet, Mariana gray (<i>Aerodramus vanikorensis bartschi</i>) - Wherever found	E	\$28,912	\$563,489	\$592,401	\$0	\$592,401
	73	Tern, California least (<i>Sterna antillarum browni</i>) - Wherever found	E	\$722,133	\$1,996,203	\$2,718,336	\$0	\$2,718,336
	43	Tern, least (<i>Sterna antillarum</i>) - interior pop.	E	\$344,321	\$5,367,349	\$5,711,670	\$55,701	\$5,767,371
	192	Tern, roseate (<i>Sterna dougallii dougallii</i>) - northeast U.S. nesting pop.	E	\$503,720	\$113,006	\$616,726	\$42,000	\$658,726
	310	Tern, roseate (<i>Sterna dougallii dougallii</i>) - Western Hemisphere except NE U.S.	T	\$25,500	\$216,343	\$241,843	\$0	\$241,843
	1421	Thrush, large Kauai (=kamao) (<i>Myadestes myadestinus</i>) - Wherever found	E	\$348	\$0	\$348	\$0	\$348

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	962	Thrush, Molokai (<i>Myadestes lanaiensis rutha</i>) - Wherever found	E	\$15,680	\$0	\$15,680	\$0	\$15,680
	489	Thrush, small Kauai (=puaiohi) (<i>Myadestes palmeri</i>) - Wherever found	E	\$98,886	\$5,345	\$104,231	\$0	\$104,231
	428	Towhee, Inyo California (<i>Pipilo crissalis eremophilus</i>) - Wherever found	T	\$9,003	\$129,515	\$138,518	\$0	\$138,518
	241	Vireo, Black-capped (<i>Vireo atricapilla</i>) - Wherever found	E	\$156,600	\$296,440	\$453,040	\$4,500	\$457,540
	77	Vireo, least Bell's (<i>Vireo bellii pusillus</i>) - Wherever found	E	\$823,179	\$1,567,201	\$2,390,380	\$0	\$2,390,380
	1349	Warbler, elfin-woods (<i>Setophaga angelae</i>) - Wherever found	T	\$3,000	\$1,000	\$4,000	\$0	\$4,000
	296	Warbler, Kirtland's (<i>Setophaga kirtlandii</i> (= <i>Dendroica kirtlandii</i>)) - Wherever found	E	\$177,000	\$99,446	\$276,446	\$300	\$276,746
	431	Warbler, nightingale reed (old world warbler) (<i>Acrocephalus lusciniia</i>) - Wherever found	E	\$137,864	\$0	\$137,864	\$0	\$137,864
	1396	Warbler (=wood), Bachman's (<i>Vermivora bachmanii</i>) - Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	160	Warbler (=wood), golden-cheeked (<i>Dendroica chrysoparia</i>) - Wherever found	E	\$232,500	\$597,200	\$829,700	\$0	\$829,700
	1422	White-eye, bridled (<i>Zosterops conspicillatus conspicillatus</i>) - Wherever found	E	\$304	\$0	\$304	\$0	\$304
	952	White-eye, Rota bridled (<i>Zosterops rotensis</i>) - Wherever found	E	\$16,272	\$0	\$16,272	\$0	\$16,272
	1210	Woodpecker, ivory-billed (<i>Campephilus principalis</i>) - Wherever found	E	\$3,000	\$5,200	\$8,200	\$200	\$8,400
	21	Woodpecker, red-cockaded (<i>Picoides borealis</i>) - Wherever found	E	\$1,718,800	\$10,866,578	\$12,585,378	\$181,805	\$12,767,183
Birds subtotal				\$44,328,546	\$80,183,108	\$124,511,654	\$2,077,065	\$126,588,719
Reptiles	97	Alligator, American (<i>Alligator mississippiensis</i>) - Wherever found	SAT	\$125,500	\$268,648	\$394,148	\$1,421,352	\$1,815,500
	619	Alligator, Chinese (<i>Alligator sinensis</i>) - Wherever found	E	\$0	\$55,750	\$55,750	\$0	\$55,750
	1396	Anole, Culebra Island giant (<i>Anolis roosevelti</i>) - Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	1396	Boa, Mona (<i>Epicrates monensis monensis</i>) - Wherever found	T	\$1,500	\$0	\$1,500	\$0	\$1,500
	436	Boa, Puerto Rican (<i>Epicrates inornatus</i>) - Wherever found	E	\$27,000	\$110,068	\$137,068	\$0	\$137,068
	995	Boa, Virgin Islands tree (<i>Epicrates monensis granti</i>) - Wherever found	E	\$14,000	\$0	\$14,000	\$0	\$14,000
	567	Crocodile, African slender-snouted (<i>Crocodylus cataphractus</i>) - Wherever found	E	\$69,420	\$0	\$69,420	\$0	\$69,420
	244	Crocodile, American (<i>Crocodylus acutus</i>) - U.S.A. (FL)	T	\$38,500	\$390,877	\$429,377	\$19,522	\$448,899
	619	Crocodile, Cuban (<i>Crocodylus rhombifer</i>) - Wherever found	E	\$0	\$55,750	\$55,750	\$0	\$55,750
	619	Crocodile, Philippine (<i>Crocodylus novaeguineae mindorensis</i>) - Wherever found	E	\$0	\$55,750	\$55,750	\$0	\$55,750
	313	Gartersnake, narrow-headed (<i>Thamnophis rufipunctatus</i>) - Wherever found	T	\$70,500	\$44,340	\$114,840	\$125,480	\$240,320
	286	Gartersnake, northern Mexican (<i>Thamnophis eques megalops</i>) - Wherever found	T	\$57,000	\$123,034	\$180,034	\$118,798	\$298,832
	619	Gavial (<i>Gavialis gangeticus</i>) - Wherever found	E	\$0	\$55,750	\$55,750	\$0	\$55,750
	1141	Gecko, Monito (<i>Sphaerodactylus micropithecus</i>) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	776	Iguana, Cuban ground (<i>Cyclura nubila nubila</i>) - (<i>Cyclura nubila nubila</i>)	T	\$0	\$30,000	\$30,000	\$0	\$30,000
	619	Iguana, Fiji banded (<i>Brachylophus fasciatus</i>) - Wherever found	E	\$0	\$55,750	\$55,750	\$0	\$55,750
	860	Iguana, Mona ground (<i>Cyclura stejnegeri</i>) - Wherever found	T	\$22,000	\$0	\$22,000	\$0	\$22,000
	276	Lizard, blunt-nosed leopard (<i>Gambelia silus</i>) - Wherever found	E	\$240,008	\$101,107	\$341,115	\$0	\$341,115
	409	Lizard, Coachella Valley fringe-toed (<i>Uma inornata</i>) - Wherever found	T	\$45,980	\$101,001	\$146,981	\$0	\$146,981
	947	Lizard, St. Croix ground (<i>Ameiva polops</i>) - Wherever found	E	\$14,000	\$2,500	\$16,500	\$0	\$16,500
	262	Massasauga (=rattlesnake), eastern (<i>Sistrurus catenatus</i>) - Wherever found	T	\$269,200	\$103,107	\$372,307	\$11,500	\$383,807

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	619	Monitor, Komodo Island (<i>Varanus komodoensis</i>) - Wherever found	E	\$0	\$55,750	\$55,750	\$0	\$55,750
	948	Rattlesnake, New Mexican ridge-nosed (<i>Crotalus willardi obscurus</i>) - Wherever found	T	\$3,500	\$9,000	\$12,500	\$3,950	\$16,450
	128	Sea turtle, green (<i>Chelonia mydas</i>) - Central North Pacific DPS	T	\$94,359	\$1,071,806	\$1,166,165	\$2,500	\$1,168,665
	169	Sea turtle, green (<i>Chelonia mydas</i>) - Central South Pacific DPS	E	\$31,134	\$748,827	\$779,961	\$0	\$779,961
	102	Sea turtle, green (<i>Chelonia mydas</i>) - Central West Pacific DPS	E	\$39,749	\$1,653,627	\$1,693,376	\$2,500	\$1,695,876
	369	Sea turtle, green (<i>Chelonia mydas</i>) - East Indian-West Pacific DPS	T	\$0	\$170,521	\$170,521	\$0	\$170,521
	618	Sea turtle, green (<i>Chelonia mydas</i>) - North Indian DPS	T	\$55,286	\$521	\$55,807	\$0	\$55,807
	1417	Sea turtle, green (<i>Chelonia mydas</i>) - Southwest Indian DPS	T	\$0	\$521	\$521	\$0	\$521
	1052	Sea turtle, green (<i>Chelonia mydas</i>) - Southwest Pacific DPS	T	\$0	\$12,111	\$12,111	\$0	\$12,111
	106	Sea turtle, green (<i>Chelonia mydas</i>) - East Pacific DPS	T	\$61,173	\$1,521,277	\$1,582,450	\$0	\$1,582,450
	89	Sea turtle, green (<i>Chelonia mydas</i>) - North Atlantic DPS	T	\$204,000	\$1,814,826	\$2,018,826	\$2,110	\$2,020,936
	136	Sea turtle, green (<i>Chelonia mydas</i>) - South Atlantic DPS	T	\$2,700	\$670,237	\$672,937	\$327,838	\$1,000,775
	53	Sea turtle, hawksbill (<i>Eretmochelys imbricata</i>) - Wherever found	E	\$1,247,284	\$3,194,486	\$4,441,770	\$9,935	\$4,451,705
	57	Sea turtle, Kemp's ridley (<i>Lepidochelys kempii</i>) - Wherever found	E	\$165,706	\$3,861,552	\$4,027,258	\$73,280	\$4,100,538
	54	Sea turtle, leatherback (<i>Dermochelys coriacea</i>) - Wherever found	E	\$692,049	\$3,631,226	\$4,323,275	\$14,934	\$4,338,209
	45	Sea turtle, loggerhead (<i>Caretta caretta</i>) - Northwest Atlantic Ocean DPS	T	\$860,800	\$3,644,152	\$4,504,952	\$874,115	\$5,379,067
	95	Sea turtle, loggerhead (<i>Caretta caretta</i>) - North Pacific Ocean DPS	E	\$2,891	\$1,917,457	\$1,920,348	\$0	\$1,920,348
	654	Sea turtle, loggerhead (<i>Caretta caretta</i>) - Mediterranean Sea DPS	E	\$49,497	\$0	\$49,497	\$0	\$49,497
	238	Sea turtle, loggerhead (<i>Caretta caretta</i>) - Northeast Atlantic Ocean DPS	E	\$287,639	\$177,166	\$464,805	\$0	\$464,805
	272	Sea turtle, loggerhead (<i>Caretta caretta</i>) - South Atlantic Ocean DPS	T	\$0	\$349,235	\$349,235	\$0	\$349,235
	1397	Sea turtle, loggerhead (<i>Caretta caretta</i>) - South Pacific Ocean DPS	E	\$0	\$1,406	\$1,406	\$0	\$1,406
	600	Sea turtle, loggerhead (<i>Caretta caretta</i>) - Southeast Indo-Pacific Ocean DPS	T	\$58,930	\$0	\$58,930	\$0	\$58,930
	153	Sea turtle, olive ridley (<i>Lepidochelys olivacea</i>) - Wherever found, except when listed as endangered under 50 CFR 224.101	T	\$221,388	\$636,363	\$857,751	\$0	\$857,751
	809	Sea turtle, olive ridley (<i>Lepidochelys olivacea</i>) - Breeding colony populations on Pacific coast of Mexico	E	\$25,000	\$1,406	\$26,406	\$0	\$26,406
	159	Skink, bluetail mole (<i>Eumeces egregius lividus</i>) - Wherever found	T	\$75,000	\$755,028	\$830,028	\$0	\$830,028
	263	Skink, sand (<i>Neoseps reynoldsi</i>) - Wherever found	T	\$310,000	\$66,267	\$376,267	\$0	\$376,267
	858	Skink, Slevin's (<i>Emoia slevini</i>) - Wherever found	E	\$22,141	\$0	\$22,141	\$0	\$22,141
	978	Snake, Atlantic salt marsh (<i>Nerodia clarkii taeniata</i>) - Wherever found	T	\$13,000	\$1,871	\$14,871	\$0	\$14,871
	584	Snake, black pine (<i>Pituophis melanoleucus lodingi</i>) - Wherever found	T	\$700	\$61,808	\$62,508	\$0	\$62,508
	1044	Snake, copperbelly water (<i>Nerodia erythrogaster neglecta</i>) - Indiana north of 40 degrees north latitude, Michigan, Ohio	T	\$9,500	\$2,886	\$12,386	\$0	\$12,386
	80	Snake, eastern indigo (<i>Drymarchon corais couperi</i>) - Wherever found	T	\$508,500	\$1,568,422	\$2,076,922	\$150,000	\$2,226,922
	149	Snake, giant garter (<i>Thamnophis gigas</i>) - Wherever found	T	\$561,008	\$323,507	\$884,515	\$0	\$884,515
	268	Snake, San Francisco garter (<i>Thamnophis sirtalis tetrataenia</i>) - Wherever found	E	\$264,155	\$88,194	\$352,349	\$0	\$352,349
	619	Tartaruga (<i>Podocnemis expansa</i>) - Wherever found	E	\$0	\$55,750	\$55,750	\$0	\$55,750
	341	Tortoise, angulated (<i>Geochelone yniphora</i>) - Wherever found	E	\$196,521	\$0	\$196,521	\$0	\$196,521
	323	Tortoise, desert (<i>Gopherus agassizii</i>) - AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	SAT	\$64,046	\$159,378	\$223,424	\$0	\$223,424
	13	Tortoise, desert (<i>Gopherus agassizii</i>) - Wherever found, except AZ south and east of Colorado R., and Mexico	T	\$1,756,368	\$14,079,956	\$15,836,324	\$138,056	\$15,974,380

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	178	Tortoise, gopher (<i>Gopherus polyphemus</i>) - West of Mobile and Tombigbee Rivers	T	\$105,000	\$651,670	\$756,670	\$6,500	\$763,170
	389	Tortoise, Madagascar radiated (<i>Geochelone radiata</i>) - Wherever found	E	\$100,000	\$55,750	\$155,750	\$0	\$155,750
	619	Tracaja (<i>Podocnemis unifilis</i>) - Wherever found	E	\$0	\$55,750	\$55,750	\$0	\$55,750
	921	Turtle, Alabama red-bellied (<i>Pseudemys alabamensis</i>) - Wherever found	E	\$562	\$17,488	\$18,050	\$0	\$18,050
	162	Turtle, bog (<i>Clemmys muhlenbergii</i>) - Wherever found, except GA, NC, SC, TN, VA	T	\$416,966	\$297,916	\$714,882	\$103,258	\$818,140
	670	Turtle, bog (<i>Clemmys muhlenbergii</i>) - U.S.A. (GA, NC, SC, TN, VA)	SAT	\$7,000	\$5,456	\$12,456	\$35,000	\$47,456
	756	Turtle, Central American river (<i>Dermatemys mawii</i>) - Wherever found	E	\$0	\$32,902	\$32,902	\$0	\$32,902
	1264	Turtle, flattened musk (<i>Sternotherus depressus</i>) - Black Warrior R. system upstream from Bankhead Dam	T	\$3,500	\$3,856	\$7,356	\$0	\$7,356
	788	Turtle, Plymouth Redbelly (<i>Pseudemys rubriventris bangsi</i>) - Wherever found	E	\$25,100	\$771	\$25,871	\$2,500	\$28,371
	1034	Turtle, ringed map (<i>Graptemys oculifera</i>) - Wherever found	T	\$0	\$12,648	\$12,648	\$0	\$12,648
	1205	Turtle, yellow-blotched map (<i>Graptemys flavimaculata</i>) - Wherever found	T	\$0	\$8,549	\$8,549	\$0	\$8,549
	303	Whipsnake (=striped racer), Alameda (<i>Masticophis lateralis euryxanthus</i>) - Wherever found	T	\$92,070	\$166,431	\$258,501	\$0	\$258,501
Reptiles subtotal				\$9,640,330	\$45,169,159	\$54,809,489	\$3,443,128	\$58,252,617
Amphibians	701	Coqui, golden (<i>Eleutherodactylus jasperi</i>) - Wherever found	T	\$13,000	\$28,250	\$41,250	\$0	\$41,250
	1164	Coqui, Llanero (<i>Eleutherodactylus juanariveroi</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	67	Frog, California red-legged (<i>Rana draytonii</i>) - Wherever found	T	\$1,342,481	\$1,781,829	\$3,124,310	\$0	\$3,124,310
	210	Frog, Chiricahua leopard (<i>Rana chiricahuensis</i>) - Wherever found	T	\$88,488	\$188,396	\$276,884	\$291,140	\$568,024
	280	Frog, dusky gopher (<i>Rana sevosia</i>) - Wherever found	E	\$316,000	\$17,734	\$333,734	\$0	\$333,734
	507	Frog, mountain yellow-legged (<i>Rana muscosa</i>) - Southern California DPS	E	\$72,490	\$18,578	\$91,068	\$0	\$91,068
	250	Frog, mountain yellow-legged (<i>Rana muscosa</i>) - Northern California DPS	E	\$205,945	\$212,008	\$417,953	\$0	\$417,953
	107	Frog, Oregon spotted (<i>Rana pretiosa</i>) - Wherever found	T	\$1,021,785	\$529,726	\$1,551,511	\$24,480	\$1,575,991
	700	Frog, Panamanian golden (<i>Atelopus varius zeteki</i>) - Wherever found	E	\$0	\$41,750	\$41,750	\$0	\$41,750
	187	Frog, Sierra Nevada Yellow-legged (<i>Rana sierrae</i>) - Wherever found	E	\$235,356	\$470,500	\$705,856	\$0	\$705,856
	1311	Guajon (<i>Eleutherodactylus cooki</i>) - Wherever found	T	\$6,000	\$0	\$6,000	\$0	\$6,000
	546	Hellbender, Ozark (<i>Cryptobranchus alleganiensis bishopi</i>) - Wherever found	E	\$46,500	\$10,000	\$56,500	\$17,900	\$74,400
	1317	Salamander, Austin blind (<i>Eurycea waterlooensis</i>) - Wherever found	E	\$5,500	\$200	\$5,700	\$0	\$5,700
	969	Salamander, Barton Springs (<i>Eurycea sosorum</i>) - Wherever found	E	\$15,000	\$300	\$15,300	\$0	\$15,300
	196	Salamander, California tiger (<i>Ambystoma californiense</i>) - U.S.A. (CA - Santa Barbara County)	E	\$541,608	\$100,600	\$642,208	\$0	\$642,208
	198	Salamander, California tiger (<i>Ambystoma californiense</i>) - U.S.A. (CA - Sonoma County)	E	\$570,447	\$52,156	\$622,603	\$0	\$622,603
	100	Salamander, California tiger (<i>Ambystoma californiense</i>) - U.S.A. (Central CA DPS)	T	\$737,957	\$1,052,249	\$1,790,206	\$0	\$1,790,206
	895	Salamander, Cheat Mountain (<i>Plethodon nettingi</i>) - Wherever found	T	\$12,180	\$200	\$12,380	\$7,336	\$19,716
	1112	Salamander, desert slender (<i>Batrachoseps aridus</i>) - Wherever found	E	\$550	\$10,000	\$10,550	\$0	\$10,550

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	157	Salamander, frosted flatwoods (<i>Ambystoma cingulatum</i>) - Wherever found	T	\$442,000	\$350,681	\$792,681	\$46,933	\$839,614
	1321	Salamander, Georgetown (<i>Eurycea naufragia</i>) - Wherever found	T	\$5,000	\$400	\$5,400	\$0	\$5,400
	619	Salamander, Japanese giant (<i>Andrias japonicus</i> (=davidianus j.)) - Wherever found	E	\$0	\$55,750	\$55,750	\$0	\$55,750
	383	Salamander, Jemez Mountains (<i>Plethodon neomexicanus</i>) - Wherever found	E	\$10,000	\$150,500	\$160,500	\$0	\$160,500
	1306	Salamander, Jollyville Plateau (<i>Eurycea tonkawae</i>) - Wherever found	T	\$5,500	\$600	\$6,100	\$0	\$6,100
	1358	Salamander, Red Hills (<i>Phaeognathus hubrichti</i>) - Wherever found	T	\$1,500	\$1,700	\$3,200	\$0	\$3,200
	195	Salamander, Reticulated flatwoods (<i>Ambystoma bishopi</i>) - Wherever found	E	\$32,500	\$588,754	\$621,254	\$30,171	\$651,425
	1321	Salamander, Salado (<i>Eurycea chisholmensis</i>) - Wherever found	T	\$5,000	\$400	\$5,400	\$0	\$5,400
	1141	Salamander, San Marcos (<i>Eurycea nana</i>) - Wherever found	T	\$10,000	\$0	\$10,000	\$0	\$10,000
	530	Salamander, Santa Cruz long-toed (<i>Ambystoma macrodactylum croceum</i>) - Wherever found	E	\$72,345	\$6,400	\$78,745	\$0	\$78,745
	780	Salamander, Shenandoah (<i>Plethodon shenandoah</i>) - Wherever found	E	\$1,800	\$27,800	\$29,600	\$0	\$29,600
	624	Salamander, Sonora tiger (<i>Ambystoma tigrinum stebbinsi</i>) - Wherever found	E	\$4,000	\$11,380	\$15,380	\$39,010	\$54,390
	888	Salamander, Texas blind (<i>Typhlomolge rathbuni</i>) - Wherever found	E	\$20,000	\$0	\$20,000	\$0	\$20,000
	200	Toad, arroyo (=arroyo southwestern) (<i>Anaxyrus californicus</i>) - Wherever found	E	\$245,087	\$375,272	\$620,359	\$0	\$620,359
	491	Toad, Houston (<i>Bufo houstonensis</i>) - Wherever found	E	\$95,000	\$7,400	\$102,400	\$1,093	\$103,493
	1206	Toad, Puerto Rican crested (<i>Peltophryne lemur</i>) - Wherever found	T	\$8,500	\$0	\$8,500	\$0	\$8,500
	201	Toad, Wyoming (<i>Bufo hemiophrys baxteri</i>) - Wherever found	E	\$578,592	\$33,020	\$611,612	\$6,778	\$618,390
	247	Toad, Yosemite (<i>Anaxyrus canorus</i>) - Wherever found	T	\$211,200	\$226,522	\$437,722	\$0	\$437,722
Amphibians subtotal				\$6,988,811	\$6,351,055	\$13,339,866	\$464,841	\$13,804,707
Fishes	1232	Blindcat, Mexican (catfish) (<i>Prietella phreatophila</i>) - Wherever found	E	\$0	\$8,000	\$8,000	\$0	\$8,000
	180	Bocaccio (<i>Sebastes paucispinis</i>) - Puget Sound - Georgia Basin DPS	E	\$0	\$610,290	\$610,290	\$139,822	\$750,112
	569	Catfish, Yaqui (<i>Ictalurus pricei</i>) - Wherever found	T	\$66,500	\$100	\$66,600	\$2,000	\$68,600
	1349	Cavefish, Alabama (<i>Speoplatyrhinus poulsoni</i>) - Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	476	Cavefish, Ozark (<i>Amblyopsis rosae</i>) - Wherever found	T	\$111,500	\$7,840	\$119,340	\$850	\$120,190
	20	Chub, bonytail (<i>Gila elegans</i>) - Wherever found	E	\$381,734	\$12,406,267	\$12,788,001	\$373,075	\$13,161,076
	443	Chub, Borax Lake (<i>Gila boraxobius</i>) - Wherever found	E	\$119,822	\$5,500	\$125,322	\$10,000	\$135,322
	1121	Chub, Chihuahua (<i>Gila nigrescens</i>) - Wherever found	T	\$5,000	\$5,390	\$10,390	\$0	\$10,390
	194	Chub, Gila (<i>Gila intermedia</i>) - Wherever found	E	\$47,000	\$590,218	\$637,218	\$14,944	\$652,162
	31	Chub, humpback (<i>Gila cypha</i>) - Wherever found	E	\$54,292	\$8,355,972	\$8,410,264	\$383,921	\$8,794,185
	555	Chub, Hutton tui (<i>Gila bicolor</i> ssp.) - Wherever found	T	\$2,664	\$68,709	\$71,373	\$0	\$71,373
	628	Chub, Owens Tui (<i>Gila bicolor</i> ssp. <i>snyderi</i>) - Wherever found	E	\$22,880	\$31,000	\$53,880	\$0	\$53,880
	297	Chub, Pahrangat roundtail (<i>Gila robusta jordani</i>) - Wherever found	E	\$186,302	\$90,362	\$276,664	\$0	\$276,664
	595	Chub, slender (<i>Erimystax cahni</i>) - Wherever found	T	\$59,000	\$1,522	\$60,522	\$0	\$60,522
	1362	Chub, Sonora (<i>Gila ditaenia</i>) - Wherever found	T	\$700	\$100	\$800	\$2,000	\$2,800
	515	Chub, spotfin (<i>Erimonax monachus</i>) - Wherever found, except where listed as an experimental population	T	\$39,000	\$47,118	\$86,118	\$0	\$86,118
	1384	Chub, spotfin (<i>Erimonax monachus</i>) - U.S.A. (TN-specified portions of the Tellico River; see 17.84(m)(1)(i))	EXPN	\$0	\$2,020	\$2,020	\$0	\$2,020
	974	Chub, spotfin (<i>Erimonax monachus</i>) - U.S.A. (AL, TN-specified portions of Shoal Creek; see 17.84(m)(1)(ii))	EXPN	\$0	\$0	\$0	\$15,000	\$15,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1385	Chub, spotfin (Erimonax monachus) - U.S.A. (TN-specified portions of the French Broad and Holston Rivers; see 17.84(m)(1)(iii))	EXPN	\$2,000	\$0	\$2,000	\$0	\$2,000
	182	Chub, Virgin River (<i>Gila seminuda</i> (=robusta)) - Wherever found	E	\$232,759	\$36,700	\$269,459	\$478,500	\$747,959
	1126	Chub, Yaqui (<i>Gila purpurea</i>) - Wherever found	E	\$7,600	\$100	\$7,700	\$2,600	\$10,300
	236	Cui-ui (<i>Chasmistes cujus</i>) - Wherever found	E	\$364,980	\$102,300	\$467,280	\$0	\$467,280
	1070	Dace, Ash Meadows speckled (<i>Rhinichthys osculus nevadensis</i>) - Wherever found	E	\$10,078	\$1,500	\$11,578	\$0	\$11,578
	454	Dace, blackside (<i>Phoxinus cumberlandensis</i>) - Wherever found	T	\$127,500	\$4,832	\$132,332	\$500	\$132,832
	955	Dace, Clover Valley speckled (<i>Rhinichthys osculus oligoporus</i>) - Wherever found	E	\$16,228	\$0	\$16,228	\$0	\$16,228
	417	Dace, desert (<i>Eremichthys acros</i>) - Wherever found	T	\$39,108	\$103,000	\$142,108	\$0	\$142,108
	329	Dace, Foskett speckled (<i>Rhinichthys osculus</i> ssp.) - Wherever found	T	\$162,169	\$24,200	\$186,369	\$21,000	\$207,369
	1377	Dace, Independence Valley speckled (<i>Rhinichthys osculus lethoporus</i>) - Wherever found	E	\$2,288	\$0	\$2,288	\$0	\$2,288
	760	Dace, Kendall Warm Springs (<i>Rhinichthys osculus thermalis</i>) - Wherever found	E	\$30,013	\$2,500	\$32,513	\$0	\$32,513
	753	Dace, Laurel (<i>Chrosomus saylori</i>) - Wherever found	E	\$32,000	\$1,100	\$33,100	\$0	\$33,100
	366	Dace, Moapa (<i>Moapa coriacea</i>) - Wherever found	E	\$166,570	\$6,500	\$173,070	\$0	\$173,070
	722	Darter, amber (<i>Percina antesella</i>) - Wherever found	E	\$2,000	\$30,600	\$32,600	\$5,000	\$37,600
	1157	Darter, bayou (<i>Etheostoma rubrum</i>) - Wherever found	T	\$0	\$9,685	\$9,685	\$0	\$9,685
	1336	Darter, bluemark (<i>Etheostoma akatulo</i>) - Wherever found	E	\$3,000	\$1,704	\$4,704	\$0	\$4,704
	399	Darter, boulder (<i>Etheostoma wapiti</i>) - Wherever found	E	\$4,500	\$131,600	\$136,100	\$15,000	\$151,100
	642	Darter, Cherokee (<i>Etheostoma scotti</i>) - Wherever found	T	\$9,500	\$37,100	\$46,600	\$5,000	\$51,600
	723	Darter, Cumberland (<i>Etheostoma susanae</i>) - Wherever found	E	\$34,000	\$1,504	\$35,504	\$2,000	\$37,504
	1171	Darter, diamond (<i>Crystallaria cincotta</i>) - Wherever found	E	\$8,582	\$853	\$9,435	\$0	\$9,435
	520	Darter, duskytail (<i>Etheostoma percunum</i>) - Wherever found	E	\$59,000	\$9,275	\$68,275	\$15,000	\$83,275
	415	Darter, Etowah (<i>Etheostoma etowahae</i>) - Wherever found	E	\$111,500	\$26,100	\$137,600	\$5,000	\$142,600
	731	Darter, fountain (<i>Etheostoma fonticola</i>) - Wherever found	E	\$25,000	\$11,074	\$36,074	\$0	\$36,074
	985	Darter, goldline (<i>Percina aurolineata</i>) - Wherever found	T	\$5,500	\$4,070	\$9,570	\$5,000	\$14,570
	510	Darter, Kentucky arrow (<i>Etheostoma spilotum</i>) - Wherever found	T	\$88,500	\$1,100	\$89,600	\$500	\$90,100
	471	Darter, leopard (<i>Percina pantherina</i>) - Wherever found	T	\$64,000	\$29,880	\$93,880	\$27,960	\$121,840
	602	Darter, Niangua (<i>Etheostoma nianguae</i>) - Wherever found	T	\$42,500	\$15,906	\$58,406	\$0	\$58,406
	278	Darter, Okaloosa (<i>Etheostoma okaloosae</i>) - Wherever found	T	\$24,500	\$309,939	\$334,439	\$0	\$334,439
	724	Darter, relict (<i>Etheostoma chienense</i>) - Wherever found	E	\$37,000	\$0	\$37,000	\$0	\$37,000
	506	Darter, rush (<i>Etheostoma phytophilum</i>) - Wherever found	E	\$91,500	\$0	\$91,500	\$0	\$91,500
	839	Darter, slackwater (<i>Etheostoma boschungii</i>) - Wherever found	T	\$6,500	\$0	\$6,500	\$17,000	\$23,500
	309	Darter, snail (<i>Percina tanasi</i>) - Wherever found	T	\$87,500	\$140,867	\$228,367	\$14,000	\$242,367
	1418	Darter, vermilion (<i>Etheostoma chermocki</i>) - Wherever found	E	\$500	\$0	\$500	\$0	\$500
	1164	Darter, watercress (<i>Etheostoma nuchale</i>) - Wherever found	E	\$8,500	\$1,000	\$9,500	\$0	\$9,500
	945	Darter, yellowcheek (<i>Etheostoma moorei</i>) - Wherever found	E	\$11,000	\$100	\$11,100	\$5,600	\$16,700
	105	Eulachon (<i>Thaleichthys pacificus</i>) - Southern DPS	T	\$0	\$1,592,017	\$1,592,017	\$62,187	\$1,654,204
	1289	Gambusia, Big Bend (<i>Gambusia gaigei</i>) - Wherever found	E	\$1,500	\$5,000	\$6,500	\$0	\$6,500
	1406	Gambusia, Clear Creek (<i>Gambusia heterochir</i>) - Wherever found	E	\$1,000	\$0	\$1,000	\$0	\$1,000
	965	Gambusia, Pecos (<i>Gambusia nobilis</i>) - Wherever found	E	\$500	\$15,000	\$15,500	\$0	\$15,500
	1418	Gambusia, San Marcos (<i>Gambusia georgei</i>) - Wherever found	E	\$500	\$0	\$500	\$0	\$500
	177	Goby, tidewater (<i>Eucyclogobius newberryi</i>) - Wherever found	E	\$433,537	\$331,864	\$765,401	\$0	\$765,401
	237	Grouper, Nassau (<i>Epinephelus striatus</i>)	T	\$0	\$464,945	\$464,945	\$0	\$464,945

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	607	Logperch, Conasauga (<i>Percina jenkinsi</i>) - Wherever found	E	\$27,000	\$25,800	\$52,800	\$5,000	\$57,800
	379	Logperch, Roanoke (<i>Percina rex</i>) - Wherever found	E	\$109,000	\$53,549	\$162,549	\$0	\$162,549
	1053	Madtom, chunky (<i>Noturus crypticus</i>) - Wherever found	E	\$11,000	\$1,100	\$12,100	\$0	\$12,100
	749	Madtom, Neosho (<i>Noturus placidus</i>) - Wherever found	T	\$12,000	\$19,822	\$31,822	\$1,500	\$33,322
	599	Madtom, pygmy (<i>Noturus stanauli</i>) - Wherever found	E	\$57,633	\$1,302	\$58,935	\$0	\$58,935
	869	Madtom, pygmy (<i>Noturus stanauli</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(t)(1)(i))	EXPN	\$0	\$0	\$0	\$21,000	\$21,000
	1353	Madtom, Scioto (<i>Noturus trautmansi</i>) - Wherever found	E	\$0	\$3,559	\$3,559	\$0	\$3,559
	836	Madtom, smoky (<i>Noturus baileyi</i>) - Wherever found	E	\$1,000	\$7,569	\$8,569	\$15,000	\$23,569
	824	Madtom, yellowfin (<i>Noturus flavipinnis</i>) - Wherever found, except where listed as an experimental population	T	\$15,500	\$9,091	\$24,591	\$0	\$24,591
	974	Madtom, yellowfin (<i>Noturus flavipinnis</i>) - U.S.A. (TN-specified portions of the Tellico River; see 17.84(e)(1)(ii))	EXPN	\$0	\$0	\$0	\$15,000	\$15,000
	1359	Madtom, yellowfin (<i>Noturus flavipinnis</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(e)(1)(iii))	EXPN	\$3,000	\$0	\$3,000	\$0	\$3,000
	659	Minnow, Devils River (<i>Dionda diaboli</i>) - Wherever found	T	\$2,500	\$800	\$3,300	\$45,304	\$48,604
	172	Minnow, loach (<i>Tiaroga cobitis</i>) - Wherever found	E	\$33,000	\$571,790	\$604,790	\$168,074	\$772,864
	33	Minnow, Rio Grande Silvery (<i>Hybognathus amarus</i>) - Wherever found, except where listed as an experimental population	E	\$43,765	\$8,311,083	\$8,354,848	\$14,785	\$8,369,633
	1406	Minnow, Rio Grande Silvery (<i>Hybognathus amarus</i>) - Rio Grande, from Little Box Canyon to Amistad Dam	EXPN	\$0	\$1,000	\$1,000	\$0	\$1,000
	36	Pikeminnow (=squawfish), Colorado (<i>Ptychocheilus lucius</i>) - Wherever found, except where listed as an experimental population	E	\$46,760	\$7,721,934	\$7,768,694	\$113,317	\$7,882,011
	380	Pikeminnow (=squawfish), Colorado (<i>Ptychocheilus lucius</i>) - Salt and Verde R. drainages, AZ	EXPN	\$0	\$0	\$0	\$162,000	\$162,000
	504	Poolfish, Pahrump (<i>Empetrichthys latos</i>) - Wherever found	E	\$90,375	\$2,300	\$92,675	\$0	\$92,675
	463	Pupfish, Ash Meadows Amargosa (<i>Cyprinodon nevadensis mionectes</i>) - Wherever found	E	\$127,220	\$1,500	\$128,720	\$0	\$128,720
	740	Pupfish, Comanche Springs (<i>Cyprinodon elegans</i>) - Wherever found	E	\$0	\$13,000	\$13,000	\$21,559	\$34,559
	243	Pupfish, desert (<i>Cyprinodon macularius</i>) - Wherever found	E	\$8,653	\$397,400	\$406,053	\$48,918	\$454,971
	140	Pupfish, Devils Hole (<i>Cyprinodon diabolis</i>) - Wherever found	E	\$482,345	\$501,500	\$983,845	\$0	\$983,845
	558	Pupfish, Owens (<i>Cyprinodon radiosus</i>) - Wherever found	E	\$22,880	\$48,000	\$70,880	\$0	\$70,880
	647	Pupfish, Warm Springs (<i>Cyprinodon nevadensis pectoralis</i>) - Wherever found	E	\$50,933	\$0	\$50,933	\$0	\$50,933
	185	Rockfish, canary (<i>Sebastes pinniger</i>) - Puget Sound - Georgia Basin DPS	T	\$0	\$559,859	\$559,859	\$162,910	\$722,769
	176	Rockfish, yelloweye (<i>Sebastes ruberrimus</i>) - Puget Sound - Georgia Basin DPS	T	\$0	\$602,793	\$602,793	\$162,910	\$765,703
	26	Salmon, Atlantic (<i>Salmo salar</i>) - Gulf of Maine DPS	E	\$3,065,541	\$6,842,367	\$9,907,908	\$0	\$9,907,908
	7	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Snake River fall-run ESU	T	\$376,500	\$32,264,433	\$32,640,933	\$195,502	\$32,836,435
	1	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Snake River spring/summer-run ESU	T	\$273,000	\$117,927,481	\$118,200,481	\$1,026,624	\$119,227,105
	18	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Sacramento River winter-run ESU	E	\$86,146	\$13,920,299	\$14,006,445	\$0	\$14,006,445
	14	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Puget Sound ESU	T	\$132,631	\$8,996,940	\$9,129,571	\$5,428,723	\$14,558,294

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	10	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Lower Columbia River ESU	T	\$715,062	\$20,966,939	\$21,682,001	\$862,859	\$22,544,860
	4	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Upper Willamette River ESU	T	\$22,123	\$48,745,659	\$48,767,782	\$267,583	\$49,035,365
	8	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Upper Columbia spring-run ESU	E	\$142,000	\$31,390,212	\$31,532,212	\$263,029	\$31,795,241
	37	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Central Valley spring-run ESU	T	\$166,120	\$7,360,796	\$7,526,916	\$0	\$7,526,916
	59	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - California Coastal ESU	T	\$0	\$4,017,394	\$4,017,394	\$0	\$4,017,394
	32	Salmon, chum (<i>Oncorhynchus keta</i>) - Columbia River ESU	T	\$3,473	\$8,432,570	\$8,436,043	\$208,782	\$8,644,825
	48	Salmon, chum (<i>Oncorhynchus keta</i>) - Hood Canal summer-run ESU	T	\$69,848	\$4,599,648	\$4,669,496	\$212,667	\$4,882,163
	46	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) - Central California Coast ESU	E	\$22,028	\$5,305,767	\$5,327,795	\$0	\$5,327,795
	12	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) - Southern Oregon - Northern California Coast ESU	T	\$271,929	\$17,959,142	\$18,231,071	\$24,000	\$18,255,071
	38	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) - Oregon Coast ESU	T	\$211,107	\$5,867,756	\$6,078,863	\$1,275,000	\$7,353,863
	24	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) - Lower Columbia River ESU	T	\$105,488	\$10,313,380	\$10,418,868	\$977,280	\$11,396,148
	11	Salmon, sockeye (<i>Oncorhynchus</i> (=Salmo) nerka) - Snake River ESU	E	\$551,164	\$19,646,847	\$20,198,011	\$1,284,828	\$21,482,839
	50	Salmon, sockeye (<i>Oncorhynchus</i> (=Salmo) nerka) - Ozette Lake ESU	T	\$1,263	\$4,616,628	\$4,617,891	\$72,232	\$4,690,123
	1387	Sawfish, largetooth (<i>Pristis pristis</i>) - Wherever found	E	\$0	\$1,853	\$1,853	\$0	\$1,853
	146	Sawfish, smalltooth (<i>Pristis pectinata</i>) - US DPS - Smalltooth sawfish originating from U.S. waters.	E	\$0	\$877,947	\$877,947	\$89,298	\$967,245
	875	Sculpin, Grotto (<i>Cottus specus</i>) - Wherever found	E	\$6,500	\$4,000	\$10,500	\$10,000	\$20,500
	959	Sculpin, pygmy (<i>Cottus paulus</i> (=pygmaeus)) - Wherever found	T	\$14,000	\$0	\$14,000	\$2,000	\$16,000
	1023	Shark, Scalloped Hammerhead (<i>Sphyrna lewini</i>) - Eastern Pacific DPS	E	\$0	\$12,907	\$12,907	\$0	\$12,907
	1054	Shark, Scalloped Hammerhead (<i>Sphyrna lewini</i>) - Central and Southwest Atlantic DPS	T	\$0	\$12,078	\$12,078	\$0	\$12,078
	1374	Shark, Scalloped Hammerhead (<i>Sphyrna lewini</i>) - Indo-West Pacific DPS	T	\$0	\$2,355	\$2,355	\$0	\$2,355
	342	Shiner, Arkansas River (<i>Notropis girardi</i>) - Arkansas River Basin (AR, KS, NM, OK, TX)	T	\$110,598	\$82,880	\$193,478	\$3,000	\$196,478
	1226	Shiner, beautiful (<i>Cyprinella formosa</i>) - Wherever found	T	\$7,500	\$100	\$7,600	\$500	\$8,100
	732	Shiner, blue (<i>Cyprinella caerulea</i>) - Wherever found	T	\$5,000	\$25,800	\$30,800	\$5,000	\$35,800
	1295	Shiner, Cahaba (<i>Notropis cahabae</i>) - Wherever found	E	\$5,000	\$1,386	\$6,386	\$0	\$6,386
	490	Shiner, Cape Fear (<i>Notropis mekistocholas</i>) - Wherever found	E	\$103,500	\$0	\$103,500	\$0	\$103,500
	1140	Shiner, palezone (<i>Notropis albizonatus</i>) - Wherever found	E	\$8,500	\$1,504	\$10,004	\$0	\$10,004
	209	Shiner, Pecos bluntnose (<i>Notropis simus pecosensis</i>) - Wherever found	T	\$65,000	\$507,000	\$572,000	\$0	\$572,000
	843	Shiner, sharpnose (<i>Notropis oxyrhynchus</i>) - Wherever found	E	\$23,000	\$100	\$23,100	\$0	\$23,100
	844	Shiner, smalleye (<i>Notropis buccula</i>) - Wherever found	E	\$23,000	\$0	\$23,000	\$0	\$23,000
	139	Shiner, Topeka (<i>Notropis topeka</i> (=tristis)) - Wherever found, except where listed as an experimental population	E	\$579,782	\$388,491	\$968,273	\$15,900	\$984,173

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	124	Shiner, Topeka (<i>Notropis topeka</i> (=tristis)) - U.S.A. (MO-specified portions of Little Creek, Big Muddy Creek, and Spring Creek watersheds in Adair, Gentry, Harrison, Putnam, Sullivan, and Worth Counties; see 17.84(d)(1)(i))	EXPN	\$1,185,500	\$17,600	\$1,203,100	\$0	\$1,203,100
	1276	Silverside, Waccamaw (<i>Menidia extensa</i>) - Wherever found	T	\$7,000	\$0	\$7,000	\$0	\$7,000
	34	Smelt, delta (<i>Hypomesus transpacificus</i>) - Wherever found	T	\$1,029,944	\$7,322,932	\$8,352,876	\$0	\$8,352,876
	175	Spinedace (<i>Meda fulgida</i>) - Wherever found	E	\$27,000	\$571,790	\$598,790	\$170,562	\$769,352
	1238	Spinedace, Big Spring (<i>Lepidomeda mollispinis pratensis</i>) - Wherever found	T	\$3,622	\$4,300	\$7,922	\$0	\$7,922
	502	Spinedace, Little Colorado (<i>Lepidomeda vittata</i>) - Wherever found	T	\$5,500	\$78,500	\$84,000	\$8,980	\$92,980
	519	Spinedace, White River (<i>Lepidomeda albivallis</i>) - Wherever found	E	\$68,968	\$15,300	\$84,268	\$0	\$84,268
	741	Springfish, Hiko White River (<i>Crenichthys baileyi grandis</i>) - Wherever found	E	\$23,138	\$11,366	\$34,504	\$0	\$34,504
	548	Springfish, Railroad Valley (<i>Crenichthys nevadae</i>) - Wherever found	T	\$71,140	\$2,800	\$73,940	\$0	\$73,940
	759	Springfish, White River (<i>Crenichthys baileyi baileyi</i>) - Wherever found	E	\$27,730	\$4,800	\$32,530	\$0	\$32,530
	17	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - California Central Valley DPS	T	\$61,928	\$14,100,714	\$14,162,642	\$0	\$14,162,642
	2	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Snake River Basin DPS	T	\$92,000	\$58,291,541	\$58,383,541	\$1,767,150	\$60,150,691
	41	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Central California Coast DPS	T	\$0	\$6,270,991	\$6,270,991	\$0	\$6,270,991
	16	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Lower Columbia River DPS	T	\$47,000	\$12,898,097	\$12,945,097	\$1,228,560	\$14,173,657
	61	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - South-Central California Coast DPS	T	\$0	\$3,980,889	\$3,980,889	\$0	\$3,980,889
	58	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Southern California DPS	E	\$0	\$4,100,474	\$4,100,474	\$0	\$4,100,474
	9	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Upper Columbia River DPS	T	\$217,500	\$26,595,918	\$26,813,418	\$310,630	\$27,124,048
	19	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Upper Willamette River DPS	T	\$0	\$13,093,034	\$13,093,034	\$268,995	\$13,362,029
	3	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Middle Columbia River DPS	T	\$398,185	\$53,950,695	\$54,348,880	\$874,763	\$55,223,643
	55	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Northern California DPS	T	\$0	\$4,226,406	\$4,226,406	\$0	\$4,226,406
	27	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Puget Sound DPS	T	\$57,883	\$8,255,588	\$8,313,471	\$1,301,804	\$9,615,275
	661	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Middle Columbia River DPS	EXPN	\$23,000	\$3,850	\$26,850	\$21,694	\$48,544
	370	Stickleback, unarmored threespine (<i>Gasterosteus aculeatus williamsoni</i>) - Wherever found	E	\$74,690	\$95,684	\$170,374	\$0	\$170,374
	961	Sturgeon, Alabama (<i>Scaphirhynchus suttkusi</i>) - Wherever found	E	\$4,500	\$199	\$4,699	\$11,000	\$15,699
	127	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) - Gulf of Maine DPS	T	\$64,667	\$1,120,223	\$1,184,890	\$0	\$1,184,890
	164	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) - New York Bight DPS	E	\$64,667	\$688,830	\$753,497	\$50,500	\$803,997
	115	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) - Chesapeake Bay DPS	E	\$64,667	\$1,315,229	\$1,379,896	\$0	\$1,379,896
	150	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) - Carolina DPS	E	\$341,500	\$537,530	\$879,030	\$0	\$879,030

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	183	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) - South Atlantic DPS	E	\$0	\$714,886	\$714,886	\$25,000	\$739,886
	734	Sturgeon, Beluga (<i>Huso huso</i>) - Wherever found	T	\$35,403	\$0	\$35,403	\$0	\$35,403
	70	Sturgeon, green (<i>Acipenser medirostris</i>) - Southern DPS	T	\$4,500	\$2,814,767	\$2,819,267	\$20,747	\$2,840,014
	78	Sturgeon (Gulf subspecies), Atlantic (<i>Acipenser oxyrinchus</i> (=oxyrhynchus) <i>desotoi</i>) - Wherever found	T	\$499,000	\$1,875,346	\$2,374,346	\$14,716	\$2,389,062
	5	Sturgeon, pallid (<i>Scaphirhynchus albus</i>) - Wherever found	E	\$2,089,909	\$32,033,233	\$34,123,142	\$78,000	\$34,201,142
	92	Sturgeon, shortnose (<i>Acipenser brevirostrum</i>) - Wherever found	E	\$50,749	\$1,784,252	\$1,835,001	\$146,000	\$1,981,001
	737	Sturgeon, Shovelnose (<i>Scaphirhynchus platyrhynchus</i>) - Wherever found	SAT	\$1,500	\$33,350	\$34,850	\$0	\$34,850
	22	Sturgeon, white (<i>Acipenser transmontanus</i>) - U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	E	\$499,868	\$11,181,030	\$11,680,898	\$279,417	\$11,960,315
	138	Sucker, June (<i>Chasmistes liorus</i>) - Wherever found	E	\$105,000	\$301,600	\$406,600	\$585,000	\$991,600
	86	Sucker, Lost River (<i>Deltistes luxatus</i>) - Wherever found	E	\$601,982	\$1,406,400	\$2,008,382	\$35,000	\$2,043,382
	15	Sucker, razorback (<i>Xyrauchen texanus</i>) - Wherever found	E	\$455,046	\$13,481,966	\$13,937,012	\$617,257	\$14,554,269
	163	Sucker, Santa Ana (<i>Catostomus santaanae</i>) - 3 CA river basins	T	\$155,815	\$653,803	\$809,618	\$0	\$809,618
	88	Sucker, shortnose (<i>Chasmistes brevirostris</i>) - Wherever found	E	\$591,982	\$1,408,400	\$2,000,382	\$35,000	\$2,035,382
	293	Sucker, Warner (<i>Catostomus warnerensis</i>) - Wherever found	T	\$102,383	\$120,340	\$222,723	\$61,000	\$283,723
	1135	Sucker, Zuni bluehead (<i>Catostomus discobolus yarrowi</i>) - Wherever found	E	\$7,900	\$500	\$8,400	\$1,700	\$10,100
	1349	Sunfish, spring pygmy (<i>Elassoma alabamae</i>) - Wherever found	T	\$4,000	\$0	\$4,000	\$0	\$4,000
	166	Topminnow, Gila (incl. Yaqui) (<i>Poeciliopsis occidentalis</i>) - Wherever found	E	\$32,500	\$615,300	\$647,800	\$145,744	\$793,544
	576	Trout, Apache (<i>Oncorhynchus apache</i>) - Wherever found	T	\$8,000	\$500	\$8,500	\$56,900	\$65,400
	6	Trout, bull (<i>Salvelinus confluentus</i>) - U.S.A., conterminous, lower 48 states	T	\$9,021,195	\$23,318,508	\$32,339,703	\$1,295,760	\$33,635,463
	506	Trout, bull (<i>Salvelinus confluentus</i>) - Clackamas River subbasin experimental population	EXPN	\$66,500	\$0	\$66,500	\$25,000	\$91,500
	385	Trout, Gila (<i>Oncorhynchus gilae</i>) - Wherever found	T	\$55,000	\$17,000	\$72,000	\$86,060	\$158,060
	215	Trout, Greenback Cutthroat (<i>Oncorhynchus clarkii stomias</i>) - Wherever found	T	\$366,825	\$76,805	\$443,630	\$97,649	\$541,279
	35	Trout, Lahontan cutthroat (<i>Oncorhynchus clarkii henshawi</i>) - Wherever found	T	\$5,665,881	\$2,164,363	\$7,830,244	\$84,000	\$7,914,244
	1403	Trout, Little Kern golden (<i>Oncorhynchus aguabonita whitei</i>) - Wherever found	T	\$1,100	\$0	\$1,100	\$0	\$1,100
	229	Trout, Paiute cutthroat (<i>Oncorhynchus clarkii seleniris</i>) - Wherever found	T	\$481,264	\$5,000	\$486,264	\$0	\$486,264
	899	Tui chub, Mohave (<i>Gila bicolor</i> ssp. <i>mohavensis</i>) - Wherever found	E	\$660	\$18,680	\$19,340	\$0	\$19,340
	181	Woundfin (<i>Plagopterus argentissimus</i>) - Wherever found, except where listed as an experimental population	E	\$281,183	\$30,900	\$312,083	\$437,842	\$749,925
Fishes subtotal				\$36,913,547	\$727,243,942	\$764,157,489	\$24,947,973	\$789,105,462
Clams	1183	Acornshell, southern (<i>Epioblasma othcaloogensis</i>) - Wherever found	E	\$1,500	\$7,742	\$9,242	\$0	\$9,242
	283	Bankclimber, purple (mussel) (<i>Elliptoideus sloatianus</i>) - Wherever found	T	\$168,500	\$132,521	\$301,021	\$5,000	\$306,021
	699	Bean, Choctaw (<i>Villosa choctawensis</i>) - Wherever found	E	\$5,000	\$36,860	\$41,860	\$0	\$41,860
	616	Bean, Cumberland (pearlymussel) (<i>Villosa trabalis</i>) - Wherever found; Except where listed as Experimental Populations	E	\$34,000	\$2,371	\$36,371	\$20,000	\$56,371

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1370	Bean, Cumberland (pearlymussel) (<i>Villosa trabalis</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$2,423	\$2,423	\$0	\$2,423
	588	Bean, purple (<i>Villosa perpurpurea</i>) - Wherever found	E	\$61,000	\$620	\$61,620	\$0	\$61,620
	663	Bean, rayed (<i>Villosa fabalis</i>) - Wherever found	E	\$25,815	\$16,583	\$42,398	\$6,000	\$48,398
	1381	Blossom, green (pearlymussel) (<i>Epioblasma torulosa gubernaculum</i>) - Wherever found	E	\$1,700	\$403	\$2,103	\$0	\$2,103
	1266	Blossom, tubercled (pearlymussel) (<i>Epioblasma torulosa torulosa</i>) - Wherever found; Except where listed as Experimental Populations	E	\$3,315	\$4,014	\$7,329	\$0	\$7,329
	1372	Blossom, turgid (pearlymussel) (<i>Epioblasma turgidula</i>) - Wherever found; Except where listed as Experimental Populations	E	\$2,000	\$400	\$2,400	\$0	\$2,400
	1420	Blossom, turgid (pearlymussel) (<i>Epioblasma turgidula</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$404	\$404	\$0	\$404
	1425	Blossom, yellow (pearlymussel) (<i>Epioblasma florentina florentina</i>) - Wherever found; Except where listed as Experimental Populations	E	\$0	\$202	\$202	\$0	\$202
	1406	Blossom, yellow (pearlymussel) (<i>Epioblasma florentina florentina</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$1,000	\$0	\$1,000	\$0	\$1,000
	1350	Catspaw, white (pearlymussel) (<i>Epioblasma obliquata perobliqua</i>) - Wherever found	E	\$1,500	\$2,478	\$3,978	\$0	\$3,978
	496	Clubshell (<i>Pleurobema clava</i>) - Wherever found; Except where listed as Experimental Populations	E	\$37,970	\$51,336	\$89,306	\$11,000	\$100,306
	861	Clubshell (<i>Pleurobema clava</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$21,500	\$404	\$21,904	\$0	\$21,904
	1259	Clubshell, black (<i>Pleurobema curtum</i>) - Wherever found	E	\$0	\$7,425	\$7,425	\$0	\$7,425
	928	Clubshell, ovate (<i>Pleurobema perovatum</i>) - Wherever found	E	\$4,500	\$10,802	\$15,302	\$2,500	\$17,802
	831	Clubshell, southern (<i>Pleurobema decisum</i>) - Wherever found	E	\$6,000	\$15,066	\$21,066	\$3,000	\$24,066
	601	Combshell, Cumberlandian (<i>Epioblasma brevidens</i>) - Wherever found; Except where listed as Experimental Populations	E	\$30,500	\$8,390	\$38,890	\$20,000	\$58,890
	1378	Combshell, Cumberlandian (<i>Epioblasma brevidens</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$2,222	\$2,222	\$0	\$2,222
	1317	Combshell, Cumberlandian (<i>Epioblasma brevidens</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$0	\$0	\$5,700	\$5,700
	713	Combshell, southern (<i>Epioblasma penita</i>) - Wherever found	E	\$31,500	\$7,125	\$38,625	\$1,000	\$39,625
	1008	Combshell, upland (<i>Epioblasma metastrata</i>) - Wherever found	E	\$9,500	\$3,019	\$12,519	\$1,000	\$13,519
	790	Ebonyshell, round (<i>Fusconaia rotulata</i>) - Wherever found	E	\$6,500	\$21,800	\$28,300	\$0	\$28,300
	305	Elktoe, Appalachian (<i>Alasmidonta raveneliana</i>) - Wherever found	E	\$40,000	\$217,462	\$257,462	\$0	\$257,462
	761	Elktoe, Cumberland (<i>Alasmidonta atropurpurea</i>) - Wherever found	E	\$17,500	\$2,780	\$20,280	\$12,000	\$32,280
	481	Fanshell (<i>Cyprogenia stegaria</i>) - Wherever found	E	\$64,815	\$42,585	\$107,400	\$5,800	\$113,200
	1388	Fanshell (<i>Cyprogenia stegaria</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$1,808	\$1,808	\$0	\$1,808
	998	Fatmucket, Arkansas (<i>Lampsilis powellii</i>) - Wherever found	T	\$8,500	\$2,545	\$11,045	\$2,900	\$13,945
	240	Heelsplitter, Carolina (<i>Lasmigona decorata</i>) - Wherever found	E	\$406,000	\$53,489	\$459,489	\$0	\$459,489

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	622	Heelsplitter, Inflated (<i>Potamilus inflatus</i>) - Wherever found	T	\$31,500	\$23,437	\$54,937	\$0	\$54,937
	191	Higgins eye (pearlymussel) (<i>Lampsilis higginsii</i>) - Wherever found	E	\$245,000	\$446,011	\$691,011	\$5,000	\$696,011
	733	Kidneyshell, fluted (<i>Ptychobranhus subtentum</i>) - Wherever found	E	\$24,500	\$5,307	\$29,807	\$5,700	\$35,507
	678	Kidneyshell, southern (<i>Ptychobranhus jonesi</i>) - Wherever found	E	\$8,500	\$37,843	\$46,343	\$0	\$46,343
	649	Kidneyshell, triangular (<i>Ptychobranhus greenii</i>) - Wherever found	E	\$44,000	\$4,682	\$48,682	\$2,000	\$50,682
	1242	Lampmussel, Alabama (<i>Lampsilis virescens</i>) - Wherever found; Except where listed as Experimental Populations	E	\$7,500	\$400	\$7,900	\$0	\$7,900
	1317	Lampmussel, Alabama (<i>Lampsilis virescens</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$0	\$0	\$5,700	\$5,700
	1331	Lilliput, pale (pearlymussel) (<i>Toxolasma cylindrellus</i>) - Wherever found	E	\$4,500	\$400	\$4,900	\$0	\$4,900
	292	Mapleleaf, winged (<i>Quadrula fragosa</i>) - Wherever found, except where listed as an experimental population	E	\$188,000	\$100,507	\$288,507	\$0	\$288,507
	686	Mapleleaf, winged (<i>Quadrula fragosa</i>) - U.S.A. (AL-specified portions of the Tennessee River; see 17.85(a)(1))	EXPN	\$44,000	\$0	\$44,000	\$0	\$44,000
	665	Moccasinshell, Alabama (<i>Medionidus acutissimus</i>) - Wherever found	T	\$42,500	\$2,382	\$44,882	\$3,000	\$47,882
	687	Moccasinshell, Coosa (<i>Medionidus parvulus</i>) - Wherever found	E	\$42,500	\$0	\$42,500	\$1,000	\$43,500
	614	Moccasinshell, Gulf (<i>Medionidus penicillatus</i>) - Wherever found	E	\$26,500	\$27,400	\$53,900	\$2,500	\$56,400
	744	Moccasinshell, Ochlockonee (<i>Medionidus simpsonianus</i>) - Wherever found	E	\$11,500	\$21,800	\$33,300	\$1,000	\$34,300
	407	Moccasinshell, Suwannee (<i>Medionidus walkeri</i>) - Wherever found	T	\$122,000	\$21,700	\$143,700	\$5,000	\$148,700
	915	Monkeyface, Appalachian (pearlymussel) (<i>Quadrula sparsa</i>) - Wherever found	E	\$18,000	\$400	\$18,400	\$0	\$18,400
	1419	Monkeyface, Appalachian (pearlymussel) (<i>Quadrula sparsa</i>) - USA (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$422	\$422	\$0	\$422
	452	Monkeyface, Cumberland (pearlymussel) (<i>Quadrula intermedia</i>) - Wherever found; Except where listed as Experimental Populations	E	\$2,000	\$131,322	\$133,322	\$0	\$133,322
	995	Monkeyface, Cumberland (pearlymussel) (<i>Quadrula intermedia</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$14,000	\$0	\$14,000	\$0	\$14,000
	300	Mucket, Neosho (<i>Lampsilis rafinesqueana</i>) - Wherever found	E	\$120,928	\$144,011	\$264,939	\$100	\$265,039
	967	Mucket, orangenacre (<i>Lampsilis perovalis</i>) - Wherever found	T	\$5,000	\$10,403	\$15,403	\$0	\$15,403
	207	Mucket, pink (pearlymussel) (<i>Lampsilis abrupta</i>) - Wherever found	E	\$419,415	\$132,979	\$552,394	\$32,900	\$585,294
	736	Mussel, oyster (<i>Epioblasma capsaeformis</i>) - Wherever found; Except where listed as Experimental Populations	E	\$32,500	\$2,591	\$35,091	\$0	\$35,091
	1384	Mussel, oyster (<i>Epioblasma capsaeformis</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$2,020	\$2,020	\$0	\$2,020
	1317	Mussel, oyster (<i>Epioblasma capsaeformis</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$0	\$0	\$5,700	\$5,700
	534	Mussel, scaleshell (<i>Leptodea leptodon</i>) - Wherever found	E	\$50,000	\$27,158	\$77,158	\$1,000	\$78,158
	336	Mussel, sheepsnose (<i>Plethobasus cyphus</i>) - Wherever found	E	\$61,015	\$134,459	\$195,474	\$6,800	\$202,274
	430	Mussel, snuffbox (<i>Epioblasma triquetra</i>) - Wherever found	E	\$79,937	\$50,712	\$130,649	\$7,600	\$138,249
	1164	Pearlshell, Alabama (<i>Margaritifera marrianae</i>) - Wherever found	E	\$9,500	\$0	\$9,500	\$0	\$9,500
	287	Pearlshell, Louisiana (<i>Margaritifera hembeli</i>) - Wherever found	T	\$279,000	\$13,320	\$292,320	\$900	\$293,220

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	990	Pearlymussel, birdwing (<i>Lemiox rimosus</i>) - Wherever found; Except where listed as Experimental Populations	E	\$8,000	\$620	\$8,620	\$5,700	\$14,320
	1425	Pearlymussel, birdwing (<i>Lemiox rimosus</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert, and Lauderdale Counties, AL	EXPN	\$0	\$202	\$202	\$0	\$202
	401	Pearlymussel, cracking (<i>Hemistena lata</i>) - Wherever found; Except where listed as Experimental Populations	E	\$19,000	\$131,120	\$150,120	\$0	\$150,120
	1420	Pearlymussel, cracking (<i>Hemistena lata</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$404	\$404	\$0	\$404
	1426	Pearlymussel, Curtis (<i>Epioblasma florentina curtisii</i>) - Wherever found	E	\$200	\$0	\$200	\$0	\$200
	909	Pearlymussel, dromedary (<i>Dromus dromas</i>) - Wherever found; Except where listed as Experimental Populations	E	\$8,500	\$2,591	\$11,091	\$7,700	\$18,791
	1400	Pearlymussel, dromedary (<i>Dromus dromas</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$1,212	\$1,212	\$0	\$1,212
	814	Pearlymussel, littlewing (<i>Pegias fabula</i>) - Wherever found	E	\$18,500	\$2,275	\$20,775	\$5,000	\$25,775
	396	Pearlymussel, slabside (<i>Pleuroaia dolabelloides</i>) - Wherever found	E	\$13,500	\$138,649	\$152,149	\$0	\$152,149
	1351	Pigtoe, Cumberland (<i>Pleurobema gibberum</i>) - Wherever found	E	\$2,000	\$1,814	\$3,814	\$0	\$3,814
	1339	Pigtoe, dark (<i>Pleurobema furvum</i>) - Wherever found	E	\$4,500	\$120	\$4,620	\$0	\$4,620
	412	Pigtoe, finereyed (<i>Fusconaia cuneolus</i>) - Wherever found; Except where listed as Experimental Populations	E	\$13,500	\$131,120	\$144,620	\$0	\$144,620
	1405	Pigtoe, finereyed (<i>Fusconaia cuneolus</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$1,010	\$1,010	\$0	\$1,010
	625	Pigtoe, fuzzy (<i>Pleurobema strodeanum</i>) - Wherever found	T	\$17,000	\$37,059	\$54,059	\$0	\$54,059
	705	Pigtoe, Georgia (<i>Pleurobema hanleyianum</i>) - Wherever found	E	\$37,500	\$199	\$37,699	\$3,000	\$40,699
	1096	Pigtoe, heavy (<i>Pleurobema taitianum</i>) - Wherever found	E	\$3,500	\$7,324	\$10,824	\$0	\$10,824
	698	Pigtoe, narrow (<i>Fusconaia escambia</i>) - Wherever found	T	\$20,000	\$21,900	\$41,900	\$0	\$41,900
	552	Pigtoe, oval (<i>Pleurobema pyriforme</i>) - Wherever found	E	\$18,000	\$50,294	\$68,294	\$5,000	\$73,294
	528	Pigtoe, rough (<i>Pleurobema plenum</i>) - Wherever found	E	\$48,000	\$29,147	\$77,147	\$2,800	\$79,947
	1405	Pigtoe, rough (<i>Pleurobema plenum</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$1,010	\$1,010	\$0	\$1,010
	1106	Pigtoe, shiny (<i>Fusconaia cor</i>) - Wherever found; Except where listed as Experimental Populations	E	\$10,000	\$620	\$10,620	\$0	\$10,620
	1415	Pigtoe, shiny (<i>Fusconaia cor</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$0	\$606	\$606	\$0	\$606
	457	Pigtoe, shiny (<i>Fusconaia cor</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$130,500	\$130,500	\$0	\$130,500
	668	Pigtoe, southern (<i>Pleurobema georgianum</i>) - Wherever found	E	\$42,500	\$0	\$42,500	\$5,000	\$47,500
	865	Pigtoe, tapered (<i>Fusconaia burkei</i>) - Wherever found	T	\$5,000	\$16,361	\$21,361	\$0	\$21,361
	553	Pimpleback, orangefoot (pearlymussel) (<i>Plethobasus cooperianus</i>) - Wherever found	E	\$42,000	\$28,733	\$70,733	\$2,000	\$72,733
	365	Pocketbook, fat (<i>Potamilus capax</i>) - Wherever found	E	\$90,500	\$66,066	\$156,566	\$17,350	\$173,916
	644	Pocketbook, finelined (<i>Lampsilis altilis</i>) - Wherever found	T	\$43,500	\$2,961	\$46,461	\$5,000	\$51,461
	589	Pocketbook, Ouachita rock (<i>Arkansia wheeleri</i>) - Wherever found	E	\$58,000	\$3,430	\$61,430	\$0	\$61,430

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	554	Pocketbook, shinyrayed (<i>Lampsilis subangulata</i>) - Wherever found	E	\$24,000	\$43,499	\$67,499	\$5,000	\$72,499
	1126	Pocketbook, speckled (<i>Lampsilis streckeri</i>) - Wherever found	E	\$9,000	\$100	\$9,100	\$1,200	\$10,300
	801	Purple Cat's paw (=Purple Cat's paw pearlymussel) (<i>Epioblasma obliquata obliquata</i>) - Wherever found; Except where listed as Experimental Populations	E	\$3,000	\$3,903	\$6,903	\$20,000	\$26,903
	1014	Purple cat's paw (=purple cat's paw pearlymussel) (<i>Epioblasma obliquata obliquata</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$10,000	\$3,202	\$13,202	\$0	\$13,202
	386	Rabbitsfoot (<i>Quadrula cylindrica cylindrica</i>) - Wherever found	T	\$103,375	\$50,381	\$153,756	\$3,810	\$157,566
	881	Rabbitsfoot, rough (<i>Quadrula cylindrica strigillata</i>) - Wherever found	E	\$19,500	\$822	\$20,322	\$0	\$20,322
	479	Riffleshell, northern (<i>Epioblasma torulosa rangiana</i>) - Wherever found	E	\$94,015	\$17,650	\$111,665	\$5,000	\$116,665
	581	Riffleshell, tan (<i>Epioblasma florentina walkeri</i> (=E. walkeri)) - Wherever found	E	\$42,500	\$1,449	\$43,949	\$20,000	\$63,949
	594	Ring pink (mussel) (<i>Obovaria retusa</i>) - Wherever found	E	\$32,500	\$26,473	\$58,973	\$2,000	\$60,973
	804	Sandshell, Southern (<i>Hamiota australis</i>) - Wherever found	T	\$4,500	\$22,161	\$26,661	\$0	\$26,661
	440	Slabshell, Chipola (<i>Elliptio chipolaensis</i>) - Wherever found	T	\$8,000	\$128,508	\$136,508	\$0	\$136,508
	299	Spectaclecase (mussel) (<i>Cumberlandia monodonta</i>) - Wherever found	E	\$116,815	\$134,748	\$251,563	\$17,700	\$269,263
	1352	Spinymussel, Altamaha (<i>Elliptio spinosa</i>) - Wherever found	E	\$250	\$1,500	\$1,750	\$2,000	\$3,750
	410	Spinymussel, James (<i>Pleurobema collina</i>) - Wherever found	E	\$144,000	\$1,700	\$145,700	\$0	\$145,700
	872	Spinymussel, Tar River (<i>Elliptio steinstansana</i>) - Wherever found	E	\$19,000	\$1,819	\$20,819	\$0	\$20,819
	327	Threeridge, fat (mussel) (<i>Amblema neislerii</i>) - Wherever found	E	\$74,500	\$131,221	\$205,721	\$4,000	\$209,721
	1256	Wartyback, white (pearlymussel) (<i>Plethobasus cicatricosus</i>) - Wherever found	E	\$5,500	\$1,208	\$6,708	\$800	\$7,508
	230	Wedgemussel, dwarf (<i>Alasmidonta heterodon</i>) - Wherever found	E	\$345,172	\$131,067	\$476,239	\$8,441	\$484,680
Clams subtotal				\$4,502,737	\$3,505,703	\$8,008,440	\$330,301	\$8,338,741
Snails	137	Abalone, Black (<i>Haliotis cracherodii</i>) - Wherever found	E	\$0	\$997,866	\$997,866	\$0	\$997,866
	82	Abalone, White (<i>Haliotis sorenseni</i>) - North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	E	\$0	\$2,167,091	\$2,167,091	\$0	\$2,167,091
	859	Ambersnail, Kanab (<i>Oxyloma haydeni kanabensis</i>) - Wherever found	E	\$11,500	\$500	\$12,000	\$10,004	\$22,004
	1328	Campeloma, slender (<i>Campeloma decampi</i>) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1354	Cavesnail, Tumbling Creek (<i>Antrobia culveri</i>) - Wherever found	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	974	Elimia, lacy (snail) (<i>Elimia crenatella</i>) - Wherever found	T	\$15,000	\$0	\$15,000	\$0	\$15,000
	1325	Hornsnail, rough (<i>Pleurocera foremani</i>) - Wherever found	E	\$4,000	\$1,196	\$5,196	\$0	\$5,196
	541	Limpet, Banbury Springs (<i>Lanx</i> sp.) - Wherever found	E	\$70,839	\$5,550	\$76,389	\$0	\$76,389
	1365	Lioplax, cylindrical (snail) (<i>Lioplax cyclostomaformis</i>) - Wherever found	E	\$2,500	\$100	\$2,600	\$0	\$2,600
	1367	Marstonia, royal (snail) (<i>Pyrgulopsis ogmorhappe</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	989	[no common name] Snail (<i>Eua zebrina</i>) - Wherever found	E	\$6,424	\$8,000	\$14,424	\$0	\$14,424
	1290	[no common name] Snail (<i>Ostodes strigatus</i>) - Wherever found	E	\$6,424	\$0	\$6,424	\$0	\$6,424
	1043	Riversnail, Anthony's (<i>Athearnia anthonyi</i>) - Wherever found; Except where listed as Experimental Populations	E	\$12,000	\$400	\$12,400	\$0	\$12,400
	1386	Riversnail, Anthony's (<i>Athearnia anthonyi</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$0	\$1,866	\$1,866	\$0	\$1,866
	1324	Rocksnaail, interrupted (=Georgia) (<i>Leptoxis foremani</i>) - Wherever found	E	\$5,000	\$199	\$5,199	\$0	\$5,199
	1398	Rocksnaail, painted (<i>Leptoxis taeniata</i>) - Wherever found	T	\$1,000	\$399	\$1,399	\$0	\$1,399

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1272	Rocksnailed, plicate (<i>Leptoxis plicata</i>) - Wherever found	E	\$7,000	\$120	\$7,120	\$0	\$7,120
	707	Rocksnailed, round (<i>Leptoxis ampla</i>) - Wherever found	T	\$32,000	\$0	\$32,000	\$8,500	\$40,500
	1396	Snail, armored (<i>Pyrgulopsis (=Marstonia) pachyta</i>) - Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	331	Snail, Bliss Rapids (<i>Taylorconcha serpenticola</i>) - Wherever found	T	\$195,046	\$10,680	\$205,726	\$0	\$205,726
	674	Snail, Chittanooga ovate amber (<i>Succinea chittengoensis</i>) - Wherever found	T	\$45,000	\$0	\$45,000	\$2,000	\$47,000
	1366	Snail, flat-spined three-toothed (<i>Triodopsis platysayoides</i>) - Wherever found	T	\$228	\$0	\$228	\$2,275	\$2,503
	578	Snail, fragile tree (<i>Samoana fragilis</i>) - Wherever found	E	\$64,886	\$0	\$64,886	\$0	\$64,886
	248	Snail, Guam tree (<i>Partula radiolata</i>) - Wherever found	E	\$79,482	\$355,793	\$435,275	\$0	\$435,275
	459	Snail, Humped tree (<i>Partula gibba</i>) - Wherever found	E	\$76,040	\$53,315	\$129,355	\$0	\$129,355
	696	Snail, Iowa Pleistocene (<i>Discus macclintocki</i>) - Wherever found	E	\$25,000	\$17,200	\$42,200	\$0	\$42,200
	755	Snail, Lanai tree (<i>Partulina semicarinata</i>) - Wherever found	E	\$33,016	\$0	\$33,016	\$0	\$33,016
	813	Snail, Lanai tree (<i>Partulina variabilis</i>) - Wherever found	E	\$25,918	\$0	\$25,918	\$0	\$25,918
	1275	Snail, Langford's tree (<i>Partula langfordi</i>) - Wherever found	E	\$7,038	\$0	\$7,038	\$0	\$7,038
	425	Snail, Morro shoulderband (=Banded dune) (<i>Helminthoglypta walkeriana</i>) - Wherever found	E	\$138,926	\$500	\$139,426	\$0	\$139,426
	927	Snail, Newcomb's (<i>Erinna newcombi</i>) - Wherever found	T	\$17,834	\$0	\$17,834	\$0	\$17,834
	860	Snail, noonday (<i>Mesodon clarki nantahala</i>) - Wherever found	T	\$22,000	\$0	\$22,000	\$0	\$22,000
	1362	Snail, painted snake coiled forest (<i>Anguispira picta</i>) - Wherever found	T	\$2,000	\$800	\$2,800	\$0	\$2,800
	1232	Snail, Pecos assiminea (<i>Assiminea pecos</i>) - Wherever found	E	\$5,500	\$2,500	\$8,000	\$0	\$8,000
	264	Snail, Snake River physa (<i>Physa natricina</i>) - Wherever found	E	\$195,046	\$177,602	\$372,648	\$0	\$372,648
	134	Snails, Oahu tree (<i>Achatinella</i> spp.) - Wherever found	E	\$673,040	\$370,769	\$1,043,809	\$0	\$1,043,809
	925	Snail, Stock Island tree (<i>Orthalicus reses</i> (not incl. <i>nesodryas</i>)) - Wherever found	T	\$16,500	\$1,341	\$17,841	\$0	\$17,841
	1360	Snail, tulotoma (<i>Tulotoma magnifica</i>) - Wherever found	T	\$1,500	\$1,399	\$2,899	\$0	\$2,899
	1389	Snail, Virginia fringed mountain (<i>Polygyriscus virginianus</i>) - Wherever found	E	\$1,800	\$0	\$1,800	\$0	\$1,800
	1327	Springsnail, Alamosa (<i>Tryonia alamosae</i>) - Wherever found	E	\$5,000	\$100	\$5,100	\$0	\$5,100
	497	Springsnail, Bruneau Hot (<i>Pyrgulopsis bruneauensis</i>) - Wherever found	E	\$90,732	\$9,050	\$99,782	\$0	\$99,782
	1327	Springsnail, Chupadera (<i>Pyrgulopsis chupaderae</i>) - Wherever found	E	\$5,000	\$100	\$5,100	\$0	\$5,100
	1232	Springsnail, Koster's (<i>Juturnia kosteri</i>) - Wherever found	E	\$5,500	\$2,500	\$8,000	\$0	\$8,000
	1194	Springsnail, Phantom (<i>Pyrgulopsis texana</i>) - Wherever found	E	\$9,000	\$0	\$9,000	\$0	\$9,000
	1232	Springsnail, Roswell (<i>Pyrgulopsis roswellensis</i>) - Wherever found	E	\$5,500	\$2,500	\$8,000	\$0	\$8,000
	1382	Springsnail, San Bernardino (<i>Pyrgulopsis bernardina</i>) - Wherever found	T	\$1,000	\$100	\$1,100	\$1,000	\$2,100
	1390	Springsnail, Socorro (<i>Pyrgulopsis neomexicana</i>) - Wherever found	E	\$1,660	\$100	\$1,760	\$0	\$1,760
	829	Springsnail, Three Forks (<i>Pyrgulopsis trivialis</i>) - Wherever found	E	\$16,500	\$100	\$16,600	\$7,513	\$24,113
	746	Tree snail, Newcomb's (<i>Newcombia cumingi</i>) - Wherever found	E	\$34,030	\$0	\$34,030	\$0	\$34,030
Snails subtotal				\$1,985,909	\$4,189,736	\$6,175,645	\$31,292	\$6,206,937
Insects	145	Beetle, American burying (<i>Nicrophorus americanus</i>) - Wherever found, except where listed as an experimental population	E	\$591,647	\$336,730	\$928,377	\$39,306	\$967,683
	934	Beetle, American burying (<i>Nicrophorus americanus</i>) - Ex Pop, SW Missouri	EXPN	\$17,500	\$0	\$17,500	\$0	\$17,500
	274	Beetle, Casey's June (<i>Dinacoma caseyi</i>) - Wherever found	E	\$342,870	\$2,684	\$345,554	\$0	\$345,554
	1218	Beetle, Coffin Cave mold (<i>Batrisodes texanus</i>) - Wherever found	E	\$5,500	\$2,750	\$8,250	\$0	\$8,250

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1130	Beetle, Comal Springs dryopid (<i>Stygoparnus comalensis</i>) - Wherever found	E	\$10,000	\$200	\$10,200	\$0	\$10,200
	817	Beetle, Comal Springs riffle (<i>Heterelmis comalensis</i>) - Wherever found	E	\$25,000	\$400	\$25,400	\$0	\$25,400
	972	Beetle, delta green ground (<i>Elaphrus viridis</i>) - Wherever found	T	\$14,344	\$800	\$15,144	\$0	\$15,144
	1367	Beetle, Helotes mold (<i>Batrisodes venyivi</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1141	Beetle, Hungerford's crawling water (<i>Brychius hungerfordi</i>) - Wherever found	E	\$9,000	\$1,000	\$10,000	\$0	\$10,000
	1364	Beetle, Kretschmarr Cave mold (<i>Texamaurops reddelli</i>) - Wherever found	E	\$2,500	\$200	\$2,700	\$0	\$2,700
	538	Beetle, Mount Hermon June (<i>Polyphylla barbata</i>) - Wherever found	E	\$65,426	\$11,200	\$76,626	\$0	\$76,626
	643	Beetle, [no common name] (<i>Rhadine exilis</i>) - Wherever found	E	\$1,500	\$50,000	\$51,500	\$0	\$51,500
	906	Beetle, [no common name] (<i>Rhadine infernalis</i>) - Wherever found	E	\$9,000	\$10,000	\$19,000	\$0	\$19,000
	1391	Beetle, Tooth Cave ground (<i>Rhadine persephone</i>) - Wherever found	E	\$1,500	\$200	\$1,700	\$0	\$1,700
	174	Beetle, valley elderberry longhorn (<i>Desmocerus californicus dimorphus</i>) - Wherever found	T	\$446,341	\$324,747	\$771,088	\$0	\$771,088
	252	Bumble bee, Rusty patched (<i>Bombus affinis</i>) - Wherever found	E	\$372,574	\$42,311	\$414,885	\$1,300	\$416,185
	684	Butterfly, Bartram's hairstreak (<i>Strymon acis bartrami</i>) - Wherever found	E	\$44,000	\$900	\$44,900	\$0	\$44,900
	692	Butterfly, bay checkerspot (<i>Euphydryas editha bayensis</i>) - Wherever found	T	\$38,500	\$4,000	\$42,500	\$0	\$42,500
	689	Butterfly, Behren's silverspot (<i>Speyeria zerene behrensii</i>) - Wherever found	E	\$37,180	\$6,000	\$43,180	\$0	\$43,180
	500	Butterfly, callippe silverspot (<i>Speyeria callippe callippe</i>) - Wherever found	E	\$58,520	\$36,200	\$94,720	\$0	\$94,720
	288	Butterfly, El Segundo blue (<i>Euphilotes battoides allyni</i>) - Wherever found	E	\$40,040	\$253,000	\$293,040	\$0	\$293,040
	141	Butterfly, Fender's blue (<i>Icaricia icarioides fenderi</i>) - Wherever found	E	\$695,510	\$286,196	\$981,706	\$0	\$981,706
	764	Butterfly, Florida leafwing (<i>Anaea troglodyta floralis</i>) - Wherever found	E	\$31,000	\$900	\$31,900	\$0	\$31,900
	156	Butterfly, Karner blue (<i>Lycaeides melissa samuelis</i>) - Wherever found	E	\$634,950	\$121,004	\$755,954	\$83,700	\$839,654
	363	Butterfly, Lange's metalmark (<i>Apodemia mormo langei</i>) - Wherever found	E	\$176,018	\$0	\$176,018	\$0	\$176,018
	1375	Butterfly, lotis blue (<i>Lycaeides argyrognomon lotis</i>) - Wherever found	E	\$2,310	\$0	\$2,310	\$0	\$2,310
	361	Butterfly, Mariana eight-spot (<i>Hypolimnys octocula marianensis</i>) - Wherever found	E	\$96,868	\$79,840	\$176,708	\$0	\$176,708
	851	Butterfly, Mariana wandering (<i>Vagrans egistina</i>) - Wherever found	E	\$19,545	\$2,905	\$22,450	\$0	\$22,450
	803	Butterfly, Miami Blue (<i>Cyclargus (=Hemiargus) thomasi bethunebakeri</i>) - Wherever found	E	\$25,500	\$1,200	\$26,700	\$0	\$26,700
	579	Butterfly, mission blue (<i>Icaricia icarioides missionensis</i>) - Wherever found	E	\$34,320	\$30,000	\$64,320	\$0	\$64,320
	505	Butterfly, Mitchell's satyr (<i>Neonympha mitchellii mitchellii</i>) - Wherever found	E	\$82,500	\$8,805	\$91,305	\$800	\$92,105
	556	Butterfly, Mount Charleston blue (<i>Icaricia (Plebejus) shasta charlestonensis</i>) - Wherever found	E	\$71,167	\$0	\$71,167	\$0	\$71,167
	1063	Butterfly, Myrtle's silverspot (<i>Speyeria zerene myrtleae</i>) - Wherever found	E	\$11,770	\$0	\$11,770	\$0	\$11,770

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	245	Butterfly, Oregon silverspot (<i>Speyeria zerene hippolyta</i>) - Wherever found	T	\$408,613	\$38,382	\$446,995	\$0	\$446,995
	325	Butterfly, Palos Verdes blue (<i>Glaucopteryx lygdamus palosverdesensis</i>) - Wherever found	E	\$1,546	\$213,136	\$214,682	\$0	\$214,682
	168	Butterfly, Quino checkerspot (<i>Euphydryas editha quino</i> (=E. e. wrighti)) - Wherever found	E	\$145,643	\$645,628	\$791,271	\$0	\$791,271
	306	Butterfly, Saint Francis' satyr (<i>Neonympha mitchellii francisci</i>) - Wherever found	E	\$5,000	\$244,481	\$249,481	\$0	\$249,481
	819	Butterfly, San Bruno elfin (<i>Callophrys mossii bayensis</i>) - Wherever found	E	\$20,680	\$4,500	\$25,180	\$0	\$25,180
	777	Butterfly, Schaus swallowtail (<i>Heraclides aristodemus ponceanus</i>) - Wherever found	E	\$18,500	\$11,418	\$29,918	\$0	\$29,918
	522	Butterfly, Smith's blue (<i>Euphilotes enoptes smithi</i>) - Wherever found	E	\$76,000	\$6,400	\$82,400	\$0	\$82,400
	693	Butterfly, Uncompahgre fritillary (<i>Boloria acrocnema</i>) - Wherever found	E	\$22,013	\$20,379	\$42,392	\$0	\$42,392
	109	Checkerspot, Taylor's (=whulge) (<i>Euphydryas editha taylori</i>) - Wherever found	E	\$909,538	\$546,666	\$1,456,204	\$22,000	\$1,478,204
	944	Damselfly, blackline Hawaiian (<i>Megalagrion nigrohamatum nigrolineatum</i>) - Wherever found	E	\$16,812	\$0	\$16,812	\$0	\$16,812
	1163	Damselfly, crimson Hawaiian (<i>Megalagrion leptodemas</i>) - Wherever found	E	\$9,510	\$0	\$9,510	\$0	\$9,510
	745	Damselfly, flying earwig Hawaiian (<i>Megalagrion nesiotis</i>) - Wherever found	E	\$34,067	\$0	\$34,067	\$0	\$34,067
	970	Damselfly, oceanic Hawaiian (<i>Megalagrion oceanicum</i>) - Wherever found	E	\$15,291	\$0	\$15,291	\$0	\$15,291
	527	Damselfly, orangeblack Hawaiian (<i>Megalagrion xanthomelas</i>) - Wherever found	E	\$42,691	\$37,928	\$80,619	\$0	\$80,619
	542	Damselfly, Pacific Hawaiian (<i>Megalagrion pacificum</i>) - Wherever found	E	\$42,642	\$33,600	\$76,242	\$0	\$76,242
	984	Damselfly, Rota blue (<i>Ischnura luta</i>) - Wherever found	E	\$14,580	\$0	\$14,580	\$0	\$14,580
	282	Dragonfly, Hine's emerald (<i>Somatochlora hineana</i>) - Wherever found	E	\$189,500	\$124,625	\$314,125	\$0	\$314,125
	85	Fly, Delhi Sands flower-loving (<i>Rhaphiomidas terminatus abdominalis</i>) - Wherever found	E	\$2,060,500	\$0	\$2,060,500	\$0	\$2,060,500
	1207	Fly, Hawaiian picture-wing (<i>Drosophila aglaia</i>) - Wherever found	E	\$8,496	\$0	\$8,496	\$0	\$8,496
	356	Fly, Hawaiian picture-wing (<i>Drosophila differens</i>) - Wherever found	E	\$180,211	\$0	\$180,211	\$0	\$180,211
	1007	Fly, Hawaiian picture-wing (<i>Drosophila digressa</i>) - Wherever found	E	\$13,566	\$0	\$13,566	\$0	\$13,566
	973	Fly, Hawaiian picture-wing (<i>Drosophila hemipeza</i>) - Wherever found	E	\$15,131	\$0	\$15,131	\$0	\$15,131
	1003	Fly, Hawaiian picture-wing (<i>Drosophila heteroneura</i>) - Wherever found	E	\$13,610	\$0	\$13,610	\$0	\$13,610
	1338	Fly, Hawaiian picture-wing (<i>Drosophila mulli</i>) - Wherever found	T	\$4,642	\$0	\$4,642	\$0	\$4,642
	1065	Fly, Hawaiian picture-wing (<i>Drosophila musaphilia</i>) - Wherever found	E	\$11,740	\$0	\$11,740	\$0	\$11,740
	1101	Fly, Hawaiian picture-wing (<i>Drosophila obatai</i>) - Wherever found	E	\$10,726	\$0	\$10,726	\$0	\$10,726
	1101	Fly, Hawaiian picture-wing (<i>Drosophila ochrobasis</i>) - Wherever found	E	\$10,726	\$0	\$10,726	\$0	\$10,726
	1065	Fly, Hawaiian picture-wing (<i>Drosophila sharpi</i>) - Wherever found	E	\$11,740	\$0	\$11,740	\$0	\$11,740
	420	Fly, Hawaiian picture-wing (<i>Drosophila substenoptera</i>) - Wherever found	E	\$12,247	\$128,415	\$140,662	\$0	\$140,662

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1101	Fly, Hawaiian picture-wing (<i>Drosophila tarphytrichia</i>) - Wherever found	E	\$10,726	\$0	\$10,726	\$0	\$10,726
	958	Grasshopper, Zayante band-winged (<i>Trimerotropis infantilis</i>) - Wherever found	E	\$1,650	\$14,400	\$16,050	\$0	\$16,050
	221	Moth, Blackburn's sphinx (<i>Manduca blackburni</i>) - Wherever found	E	\$506,670	\$12,300	\$518,970	\$0	\$518,970
	1341	Moth, Kern primrose sphinx (<i>Euproserpinus euterpe</i>) - Wherever found	T	\$2,090	\$2,500	\$4,590	\$0	\$4,590
	866	Naucorid, Ash Meadows (<i>Ambrysus amargosus</i>) - Wherever found	T	\$21,324	\$0	\$21,324	\$0	\$21,324
	453	Picture-wing fly, Hawaiian (<i>Drosophila montgomeryi</i>) - Wherever found	E	\$4,642	\$128,415	\$133,057	\$0	\$133,057
	621	Skipper, Carson wandering (<i>Pseudocopaeodes eunus obscurus</i>) - Wherever found	E	\$27,456	\$27,920	\$55,376	\$0	\$55,376
	307	Skipper, Dakota (<i>Hesperia dacotae</i>) - Wherever found	T	\$149,129	\$68,975	\$218,104	\$30,505	\$248,609
	1089	Skipper, Laguna Mountains (<i>Pyrgus ruralis lagunae</i>) - Wherever found	E	\$9,240	\$1,680	\$10,920	\$0	\$10,920
	228	Skipperling, Poweshiek (<i>Oarisma poweshiek</i>) - Wherever found	E	\$478,547	\$6,225	\$484,772	\$2,600	\$487,372
	1427	Skipper, Pawnee montane (<i>Hesperia leonardus montana</i>) - Wherever found	T	\$0	\$100	\$100	\$0	\$100
	394	Tiger beetle, Miami (<i>Cicindelia floridana</i>) - Wherever found	E	\$152,500	\$0	\$152,500	\$0	\$152,500
	598	Tiger beetle, Northeastern beach (<i>Cicindela dorsalis dorsalis</i>) - Wherever found	T	\$49,456	\$10,395	\$59,851	\$0	\$59,851
	638	Tiger beetle, Ohlone (<i>Cicindela ohlone</i>) - Wherever found	E	\$47,515	\$4,800	\$52,315	\$0	\$52,315
	582	Tiger beetle, Puritan (<i>Cicindela puritana</i>) - Wherever found	T	\$53,869	\$9,765	\$63,634	\$0	\$63,634
	493	Tiger beetle, Salt Creek (<i>Cicindela nevadica lincolniiana</i>) - Wherever found	E	\$69,125	\$15,575	\$84,700	\$17,600	\$102,300
	727	Yellow-faced bee, anthracinan (<i>Hylaeus anthracinus</i>) - Wherever found	E	\$36,390	\$200	\$36,590	\$0	\$36,590
	1189	Yellow-faced bee, assimilans (<i>Hylaeus assimilans</i>) - Wherever found	E	\$9,066	\$0	\$9,066	\$0	\$9,066
	1189	Yellow-faced bee, easy (<i>Hylaeus facilis</i>) - Wherever found	E	\$9,066	\$0	\$9,066	\$0	\$9,066
	721	Yellow-faced bee, Hawaiian (<i>Hylaeus kuakea</i>) - Wherever found	E	\$733	\$37,075	\$37,808	\$0	\$37,808
	1069	Yellow-faced bee, Hawaiian (<i>Hylaeus longiceps</i>) - Wherever found	E	\$11,601	\$0	\$11,601	\$0	\$11,601
	1410	Yellow-faced bee, Hawaiian (<i>Hylaeus mana</i>) - Wherever found	E	\$733	\$0	\$733	\$0	\$733
	1410	Yellow-faced bee, hiliaris (<i>Hylaeus hiliaris</i>) - Wherever found	E	\$733	\$0	\$733	\$0	\$733
Insects subtotal				\$9,995,122	\$4,010,055	\$14,005,177	\$197,811	\$14,202,988
Arachnids	1364	Harvestman, Bee Creek Cave (<i>Texella reddelli</i>) - Wherever found	E	\$2,500	\$200	\$2,700	\$0	\$2,700
	359	Harvestman, Bone Cave (<i>Texella reyesi</i>) - Wherever found	E	\$175,000	\$2,950	\$177,950	\$0	\$177,950
	1367	Harvestman, Cokendolpher Cave (<i>Texella cokendolpheri</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1367	Meshweaver, Braken Bat Cave (<i>Cicurina venii</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1359	Meshweaver, Government Canyon Bat Cave (<i>Cicurina vespera</i>) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1358	Meshweaver, Madla's Cave (<i>Cicurina madla</i>) - Wherever found	E	\$3,000	\$200	\$3,200	\$0	\$3,200
	1359	Meshweaver, Robber Baron Cave (<i>Cicurina baronia</i>) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1391	Pseudoscorpion, Tooth Cave (<i>Tartarocreagris texana</i>) - Wherever found	E	\$1,500	\$200	\$1,700	\$0	\$1,700
	1359	Spider, Government Canyon Bat Cave (<i>Neoleptoneta microps</i>) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	920	Spider, Kauai cave wolf or pe'e pe'e maka 'ole (Adelocosa anops) - Wherever found	E	\$18,129	\$0	\$18,129	\$0	\$18,129
	673	Spider, spruce-fir moss (Microhexura montivaga) - Wherever found	E	\$43,000	\$4,004	\$47,004	\$0	\$47,004
	1391	Spider, Tooth Cave (Neoleptoneta myopica) - Wherever found	E	\$1,500	\$200	\$1,700	\$0	\$1,700
Arachnids subtotal				\$258,629	\$7,754	\$266,383	\$0	\$266,383
Crustaceans	1359	Amphipod, diminutive (Gammarus hyalleloides) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1342	Amphipod, Hay's Spring (Stygobromus hayi) - Wherever found	E	\$4,532	\$0	\$4,532	\$0	\$4,532
	1359	Amphipod, Illinois cave (Gammarus acherondytes) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	920	Amphipod, Kauai cave (Spelaorchestia koloana) - Wherever found	E	\$18,129	\$0	\$18,129	\$0	\$18,129
	1232	Amphipod, Noel's (Gammarus desperatus) - Wherever found	E	\$5,500	\$2,500	\$8,000	\$0	\$8,000
	1130	Amphipod, Peck's cave (Stygobromus (=Stygonectes) pecki) - Wherever found	E	\$10,000	\$200	\$10,200	\$0	\$10,200
	1359	Amphipod, Pecos (Gammarus pecos) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1141	Crayfish, Benton County cave (Cambarus aculabrum) - Wherever found	E	\$3,900	\$6,000	\$9,900	\$100	\$10,000
	494	Crayfish, Big Sandy (Cambarus callainus) - Wherever found	T	\$100,756	\$140	\$100,896	\$0	\$100,896
	627	Crayfish, Guyandotte River (Cambarus veteranus) - Wherever found	E	\$53,982	\$0	\$53,982	\$0	\$53,982
	1347	Crayfish, Hell Creek Cave (Cambarus zophonastes) - Wherever found	E	\$3,000	\$0	\$3,000	\$1,200	\$4,200
	525	Crayfish, Nashville (Orconectes shoupi) - Wherever found	E	\$79,000	\$2,730	\$81,730	\$0	\$81,730
	890	Crayfish, Shasta (Pacifastacus fortis) - Wherever found	E	\$9,900	\$10,000	\$19,900	\$0	\$19,900
	344	Fairy shrimp, Conservancy (Branchinecta conservatio) - Wherever found	E	\$154,148	\$37,420	\$191,568	\$0	\$191,568
	348	Fairy shrimp, longhorn (Branchinecta longiantenna) - Wherever found	E	\$28,434	\$160,600	\$189,034	\$0	\$189,034
	797	Fairy shrimp, Riverside (Streptocephalus woottoni) - Wherever found	E	\$21,996	\$5,468	\$27,464	\$0	\$27,464
	345	Fairy shrimp, San Diego (Branchinecta sandiegonensis) - Wherever found	E	\$83,019	\$108,529	\$191,548	\$0	\$191,548
	111	Fairy shrimp, vernal pool (Branchinecta lynchi) - Wherever found	T	\$641,441	\$807,350	\$1,448,791	\$0	\$1,448,791
	942	Isopod, Lee County cave (Lirceus usdagalun) - Wherever found	E	\$12,000	\$5,000	\$17,000	\$0	\$17,000
	650	Isopod, Madison Cave (Antrolana lira) - Wherever found	T	\$50,581	\$0	\$50,581	\$0	\$50,581
	1326	Isopod, Socorro (Thermosphaeroma thermophilus) - Wherever found	E	\$1,660	\$3,500	\$5,160	\$0	\$5,160
	1406	Shrimp, Alabama cave (Palaemonias alabamae) - Wherever found	E	\$1,000	\$0	\$1,000	\$0	\$1,000
	957	Shrimp, anchialine pool (Procaris hawaiiiana) - Wherever found	E	\$16,101	\$0	\$16,101	\$0	\$16,101
	1018	Shrimp, anchialine pool (Vetericaris chaceorum) - Wherever found	E	\$13,059	\$0	\$13,059	\$0	\$13,059
	923	Shrimp, California freshwater (Syncaris pacifica) - Wherever found	E	\$11,440	\$6,556	\$17,996	\$0	\$17,996
	779	Shrimp, Kentucky cave (Palaemonias ganteri) - Wherever found	E	\$9,000	\$20,802	\$29,802	\$0	\$29,802
	152	Tadpole shrimp, vernal pool (Lepidurus packardii) - Wherever found	E	\$507,158	\$363,293	\$870,451	\$0	\$870,451
Crustaceans subtotal				\$1,848,736	\$1,540,088	\$3,388,824	\$1,300	\$3,390,124
Corals	232	Coral, Boulder star (Orbicella franksi) - Wherever found	T	\$0	\$483,618	\$483,618	\$0	\$483,618
	213	Coral, elkhorn (Acropora palmata) - Wherever found	T	\$0	\$549,410	\$549,410	\$0	\$549,410
	218	Coral, Lobed Star (Orbicella annularis) - Wherever found	T	\$0	\$523,628	\$523,628	\$0	\$523,628
	231	Coral, Mountainous Star (Orbicella faveolata) - Wherever found	T	\$0	\$484,138	\$484,138	\$0	\$484,138
	216	Coral, Pillar (Dendrogyra cylindrus) - Wherever found	T	\$0	\$537,504	\$537,504	\$0	\$537,504
	219	Coral, Rough Cactus (Mycetophyllia ferox) - Wherever found	T	\$0	\$523,385	\$523,385	\$0	\$523,385
	203	Coral, staghorn (Acropora cervicornis) - Wherever found	T	\$0	\$601,866	\$601,866	\$0	\$601,866
	189	No Common Name (Acropora globiceps) - Wherever found	T	\$0	\$702,853	\$702,853	\$0	\$702,853
	269	No Common Name (Acropora jacquelinae) - Wherever found	T	\$0	\$350,300	\$350,300	\$0	\$350,300
	275	No Common Name (Acropora retusa) - Wherever found	T	\$0	\$344,279	\$344,279	\$0	\$344,279

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1141	No Common Name (Acropora rudis) - Wherever found	T	\$0	\$10,000	\$10,000	\$0	\$10,000
	269	No Common Name (Acropora speciosa) - Wherever found	T	\$0	\$350,300	\$350,300	\$0	\$350,300
	269	No Common Name (Euphyllia paradivisa) - Wherever found	T	\$0	\$350,300	\$350,300	\$0	\$350,300
	269	No Common Name (Isopora crateriformis) - Wherever found	T	\$0	\$350,300	\$350,300	\$0	\$350,300
	253	No Common Name (Seriatopora aculeata) - Wherever found	T	\$0	\$410,419	\$410,419	\$0	\$410,419
Corals subtotal				\$0	\$6,572,300	\$6,572,300	\$0	\$6,572,300
Flowering Plants	1037	A`e (Zanthoxylum dipetalum var. tomentosum) - Wherever found	E	\$12,545	\$0	\$12,545	\$0	\$12,545
	666	A`e (Zanthoxylum hawaiiense) - Wherever found	E	\$15,634	\$32,000	\$47,634	\$0	\$47,634
	999	A`e (Zanthoxylum oahuense) - Wherever found	E	\$13,925	\$0	\$13,925	\$0	\$13,925
	418	`Ahinahina (Argyroxiphium sandwicense ssp. macrocephalum) - Wherever found	T	\$6,492	\$135,420	\$141,912	\$0	\$141,912
	1270	`Ahinahina (Argyroxiphium sandwicense ssp. sandwicense) - Wherever found	E	\$7,253	\$0	\$7,253	\$0	\$7,253
	798	`Aiakeakua, popolo (Solanum sandwicense) - Wherever found	E	\$27,450	\$0	\$27,450	\$0	\$27,450
	877	`Aiea (Nothocestrum breviflorum) - Wherever found	E	\$20,439	\$0	\$20,439	\$0	\$20,439
	806	`Aiea (Nothocestrum latifolium) - Wherever found	E	\$5,914	\$20,700	\$26,614	\$0	\$26,614
	1081	`Aiea (Nothocestrum peltatum) - Wherever found	E	\$11,044	\$0	\$11,044	\$0	\$11,044
	441	`Akoko (Euphorbia celastroides var. kaenana) - Wherever found	E	\$7,700	\$128,415	\$136,115	\$0	\$136,115
	1221	`Akoko (Euphorbia deppeana) - Wherever found	E	\$8,164	\$0	\$8,164	\$0	\$8,164
	1185	`Akoko (Euphorbia eleanoriae) - Wherever found	E	\$9,186	\$0	\$9,186	\$0	\$9,186
	1004	`Akoko (Euphorbia haeleleana) - Wherever found	E	\$13,594	\$0	\$13,594	\$0	\$13,594
	426	`Akoko (Euphorbia herbstii) - Wherever found	E	\$10,463	\$128,415	\$138,878	\$0	\$138,878
	941	`Akoko (Euphorbia kuwaleana) - Wherever found	E	\$6,208	\$10,817	\$17,025	\$0	\$17,025
	1188	`Akoko (Euphorbia remyi var. kauaiensis) - Wherever found	E	\$9,087	\$0	\$9,087	\$0	\$9,087
	1042	`Akoko (Euphorbia remyi var. remyi) - Wherever found	E	\$12,490	\$0	\$12,490	\$0	\$12,490
	1136	`Akoko (Euphorbia rockii) - Wherever found	E	\$10,093	\$0	\$10,093	\$0	\$10,093
	1269	`Akoko (Euphorbia halemanui) - Wherever found	E	\$7,288	\$0	\$7,288	\$0	\$7,288
	258	`Akoko, Ewa Plains (Euphorbia skottsbergii var. skottsbergii) - Wherever found	E	\$394,960	\$4,434	\$399,394	\$0	\$399,394
	903	`aku (Cyanea tritomantha) - Wherever found	E	\$16,711	\$2,500	\$19,211	\$0	\$19,211
	1146	`aku`aku (Cyanea platyphylla) - Wherever found	E	\$9,941	\$0	\$9,941	\$0	\$9,941
	1212	`Ala`ala wai nui (Peperomia subpetiolata) - Wherever found	E	\$8,376	\$0	\$8,376	\$0	\$8,376
	856	Alani (Melicope adscendens) - Wherever found	E	\$20,157	\$2,000	\$22,157	\$0	\$22,157
	1208	Alani (Melicope balloui) - Wherever found	E	\$8,478	\$0	\$8,478	\$0	\$8,478
	893	Alani (Melicope christophersenii) - Wherever found	E	\$5,914	\$13,895	\$19,809	\$0	\$19,809
	1011	Alani (Melicope degeneri) - Wherever found	E	\$13,451	\$0	\$13,451	\$0	\$13,451
	1090	Alani (Melicope haupuensis) - Wherever found	E	\$10,913	\$0	\$10,913	\$0	\$10,913
	1143	Alani (Melicope hiiakae) - Wherever found	E	\$9,995	\$0	\$9,995	\$0	\$9,995
	901	Alani (Melicope knudsenii) - Wherever found	E	\$17,275	\$2,000	\$19,275	\$0	\$19,275
	615	Alani (Melicope lydgatei) - Wherever found	E	\$56,389	\$0	\$56,389	\$0	\$56,389
	1310	Alani (Melicope makahae) - Wherever found	E	\$6,034	\$0	\$6,034	\$0	\$6,034
	1039	Alani (Melicope mucronulata) - Wherever found	E	\$10,502	\$2,000	\$12,502	\$0	\$12,502
	1236	Alani (Melicope munroi) - Wherever found	E	\$7,942	\$0	\$7,942	\$0	\$7,942
	664	Alani (Melicope ovalis) - Wherever found	E	\$6,928	\$41,400	\$48,328	\$0	\$48,328
	1253	Alani (Melicope pallida) - Wherever found	E	\$7,582	\$0	\$7,582	\$0	\$7,582
	1261	Alani (Melicope paniculata) - Wherever found	E	\$7,408	\$0	\$7,408	\$0	\$7,408
	889	Alani (Melicope puberula) - Wherever found	E	\$19,994	\$0	\$19,994	\$0	\$19,994

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1076	Alani (Melicope quadrangularis) - Wherever found	E	\$11,372	\$0	\$11,372	\$0	\$11,372
	1308	Alani (Melicope reflexa) - Wherever found	E	\$6,067	\$0	\$6,067	\$0	\$6,067
	886	Alani (Melicope saint-johnii) - Wherever found	E	\$6,208	\$13,895	\$20,103	\$0	\$20,103
	1029	Alani (Melicope zahlbruckneri) - Wherever found	E	\$10,728	\$2,000	\$12,728	\$0	\$12,728
	1132	Allocarya, Calistoga (Plagiobothrys strictus) - Wherever found	E	\$10,120	\$0	\$10,120	\$0	\$10,120
	1321	Alopecurus, Sonoma (Alopecurus aequalis var. sonomensis) - Wherever found	E	\$4,400	\$1,000	\$5,400	\$0	\$5,400
	302	Amaranth, seabeach (Amaranthus pumilus) - Wherever found	T	\$133,237	\$125,904	\$259,141	\$0	\$259,141
	816	Ambrosia, San Diego (Ambrosia pumila) - Wherever found	E	\$24,792	\$771	\$25,563	\$0	\$25,563
	735	Ambrosia, south Texas (Ambrosia cheiranthifolia) - Wherever found	E	\$35,000	\$400	\$35,400	\$0	\$35,400
	487	Amole, purple (Chlorogalum purpureum) - Wherever found	T	\$17,520	\$87,693	\$105,213	\$0	\$105,213
	1232	Amphianthus, little (Amphianthus pusillus) - Wherever found	T	\$5,500	\$500	\$6,000	\$2,000	\$8,000
	886	ʻAnaunau (Lepidium arbuscula) - Wherever found	E	\$6,208	\$13,895	\$20,103	\$0	\$20,103
	1184	ʻAnunu (Sicyos albus) - Wherever found	E	\$9,231	\$0	\$9,231	\$0	\$9,231
	718	ʻAnunu (Sicyos macrophyllus) - Wherever found	E	\$6,140	\$32,000	\$38,140	\$0	\$38,140
	845	Aplokating-palaoan (Psychotria malaspinae) - Wherever found	E	\$22,987	\$0	\$22,987	\$0	\$22,987
	793	Arrowhead, bunched (Sagittaria fasciculata) - Wherever found	E	\$21,500	\$6,439	\$27,939	\$0	\$27,939
	521	Aster, decurrent false (Boltonia decurrens) - Wherever found	T	\$8,000	\$75,100	\$83,100	\$0	\$83,100
	769	Aster, Florida golden (Chrysopsis floridana) - Wherever found	E	\$27,000	\$200	\$27,200	\$3,500	\$30,700
	1179	Aster, Ruth's golden (Pityopsis ruthii) - Wherever found	E	\$3,500	\$5,802	\$9,302	\$0	\$9,302
	704	Aupaka (Isodendron hosakae) - Wherever found	E	\$9,109	\$32,000	\$41,109	\$0	\$41,109
	1111	Aupaka (Isodendron laurifolium) - Wherever found	E	\$10,552	\$0	\$10,552	\$0	\$10,552
	1111	Aupaka (Isodendron longifolium) - Wherever found	T	\$10,552	\$0	\$10,552	\$0	\$10,552
	1169	Avens, spreading (Geum radiatum) - Wherever found	E	\$5,000	\$4,450	\$9,450	\$0	\$9,450
	1059	ʻAwikiwiki (Canavalia molokaiensis) - Wherever found	E	\$6,928	\$5,000	\$11,928	\$0	\$11,928
	1269	ʻAwikiwiki (Canavalia napaliensis) - Wherever found	E	\$7,288	\$0	\$7,288	\$0	\$7,288
	940	ʻAwikiwiki (Canavalia pubescens) - Wherever found	E	\$17,194	\$0	\$17,194	\$0	\$17,194
	1313	Awiwi (Schenkia sebaeoides) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1303	ʻAwiwi (Kadua cookiana) - Wherever found	E	\$6,203	\$0	\$6,203	\$0	\$6,203
	676	Ayenia, Texas (Ayenia limitaris) - Wherever found	E	\$46,500	\$200	\$46,700	\$0	\$46,700
	1369	Baccharis, Encinitas (Baccharis vanessae) - Wherever found	T	\$2,343	\$100	\$2,443	\$0	\$2,443
	1291	Barbara's Buttons, Mohr's (Marshallia mohrii) - Wherever found	T	\$1,500	\$922	\$2,422	\$4,000	\$6,422
	800	Barberry, island (Berberis pinnata ssp. insularis) - Wherever found	E	\$21,067	\$5,878	\$26,945	\$0	\$26,945
	1412	Barberry, Nevin's (Berberis nevinii) - Wherever found	E	\$0	\$700	\$700	\$0	\$700
	1257	Bariaco (Trichilia triacantha) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	492	Beaked-rush, Knieskem's (Rhynchospora knieskemii) - Wherever found	T	\$22,150	\$80,560	\$102,710	\$0	\$102,710
	1252	Bean, sea (Mucuna sloanei var. persericea) - Wherever found	E	\$7,607	\$0	\$7,607	\$0	\$7,607
	767	Beardtongue, Parachute (Penstemon debilis) - Wherever found	T	\$1,541	\$10,206	\$11,747	\$19,550	\$31,297
	1079	Beardtongue, Penland (Penstemon penlandii) - Wherever found	E	\$0	\$11,056	\$11,056	\$0	\$11,056
	758	Beargrass, Britton's (Nolina brittoniana) - Wherever found	E	\$0	\$21,700	\$21,700	\$11,000	\$32,700
	837	Bear-poppy, Dwarf (Arctomecon humilis) - Wherever found	E	\$10,000	\$13,516	\$23,516	\$0	\$23,516
	672	Beauty, Harper's (Harperocallis flava) - Wherever found	E	\$7,500	\$21,700	\$29,200	\$18,000	\$47,200
	908	Bedstraw, El Dorado (Galium californicum ssp. sierrae) - Wherever found	E	\$1,650	\$17,300	\$18,950	\$0	\$18,950
	1229	Bedstraw, island (Galium buxifolium) - Wherever found	E	\$0	\$8,059	\$8,059	\$0	\$8,059
	1354	Bellflower, Brooksville (Campanula robinsiae) - Wherever found	E	\$0	\$0	\$0	\$3,500	\$3,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1412	Birch, Virginia round-leaf (<i>Betula uber</i>) - Wherever found	T	\$700	\$0	\$700	\$0	\$700
	830	Bird's beak, palmate-bracted (<i>Cordylanthus palmatus</i>) - Wherever found	E	\$9,790	\$14,300	\$24,090	\$0	\$24,090
	1393	Bird's-beak, Pennell's (<i>Cordylanthus tenuis</i> ssp. <i>capillaris</i>) - Wherever found	E	\$1,650	\$0	\$1,650	\$0	\$1,650
	451	Bird's-beak, salt marsh (<i>Cordylanthus maritimus</i> ssp. <i>maritimus</i>) - Wherever found	E	\$26,542	\$106,996	\$133,538	\$0	\$133,538
	402	Bird's-beak, soft (<i>Cordylanthus mollis</i> ssp. <i>mollis</i>) - Wherever found	E	\$32,778	\$117,200	\$149,978	\$0	\$149,978
	1319	Birds-in-a-nest, white (<i>Macbridea alba</i>) - Wherever found	T	\$3,500	\$1,100	\$4,600	\$900	\$5,500
	1408	Bittercress, small-anthered (<i>Cardamine micranthera</i>) - Wherever found	E	\$700	\$200	\$900	\$0	\$900
	912	Bladderpod, Dudley Bluffs (<i>Lesquerella congesta</i>) - Wherever found	T	\$4,403	\$12,956	\$17,359	\$1,108	\$18,467
	1312	Bladderpod, kodachrome (<i>Lesquerella tumulosa</i>) - Wherever found	E	\$2,000	\$3,950	\$5,950	\$0	\$5,950
	1426	Bladderpod, lyrate (<i>Lesquerella lyrata</i>) - Wherever found	T	\$0	\$200	\$200	\$0	\$200
	679	Bladderpod, Missouri (<i>Physaria filiformis</i>) - Wherever found	T	\$9,500	\$35,840	\$45,340	\$500	\$45,840
	1426	Bladderpod, San Bernardino Mountains (<i>Lesquerella kingii</i> ssp. <i>bernardina</i>) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	630	Bladderpod, Short's (<i>Physaria globosa</i>) - Wherever found	E	\$42,500	\$9,727	\$52,227	\$1,200	\$53,427
	1294	Bladderpod, Spring Creek (<i>Lesquerella perforata</i>) - Wherever found	E	\$4,000	\$2,400	\$6,400	\$0	\$6,400
	1323	Bladderpod, white (<i>Lesquerella pallida</i>) - Wherever found	E	\$5,000	\$200	\$5,200	\$0	\$5,200
	726	Bladderpod, White Bluffs (<i>Physaria douglasii</i> ssp. <i>tuplashensis</i>) - Wherever found	T	\$36,200	\$400	\$36,600	\$0	\$36,600
	1358	Bladderpod, Zapata (<i>Lesquerella thamnophila</i>) - Wherever found	E	\$2,800	\$400	\$3,200	\$0	\$3,200
	979	Blazingstar, Ash Meadows (<i>Mentzelia leucophylla</i>) - Wherever found	T	\$14,848	\$0	\$14,848	\$0	\$14,848
	1204	Blazingstar, Heller's (<i>Liatris helleri</i>) - Wherever found	T	\$5,000	\$3,600	\$8,600	\$0	\$8,600
	1318	Blazingstar, scrub (<i>Liatris ohlingerae</i>) - Wherever found	E	\$0	\$100	\$100	\$5,500	\$5,600
	776	Bluecurls, Hidden Lake (<i>Trichostema austromontanum</i> ssp. <i>compactum</i>) - Wherever found	T	\$30,000	\$0	\$30,000	\$0	\$30,000
	1268	Bluegrass, Hawaiian (<i>Poa sandvicensis</i>) - Wherever found	E	\$7,309	\$0	\$7,309	\$0	\$7,309
	1268	Bluegrass, Mann's (<i>Poa mannii</i>) - Wherever found	E	\$7,309	\$0	\$7,309	\$0	\$7,309
	1132	Bluegrass, Napa (<i>Poa napensis</i>) - Wherever found	E	\$10,120	\$0	\$10,120	\$0	\$10,120
	1330	Bluegrass, San Bernardino (<i>Poa atropurpurea</i>) - Wherever found	E	\$4,433	\$500	\$4,933	\$0	\$4,933
	1318	Blue-star, Kearney's (<i>Amsonia kearneyana</i>) - Wherever found	E	\$500	\$4,600	\$5,100	\$500	\$5,600
	1364	Bluet, Roan Mountain (<i>Hedyotis purpurea</i> var. <i>montana</i>) - Wherever found	E	\$2,700	\$0	\$2,700	\$0	\$2,700
	762	Bonamia, Florida (<i>Bonamia grandiflora</i>) - Wherever found	T	\$7,500	\$21,700	\$29,200	\$3,000	\$32,200
	1311	Boxwood, Vahl's (<i>Buxus vahlII</i>) - Wherever found	E	\$6,000	\$0	\$6,000	\$0	\$6,000
	1362	Brickell-bush, Florida (<i>Brickellia mosieri</i>) - Wherever found	E	\$1,500	\$100	\$1,600	\$1,200	\$2,800
	658	Brodiaea, Chinese Camp (<i>Brodiaea pallida</i>) - Wherever found	T	\$47,520	\$1,200	\$48,720	\$0	\$48,720
	319	Brodiaea, thread-leaved (<i>Brodiaea filifolia</i>) - Wherever found	T	\$233,899	\$1,342	\$235,241	\$0	\$235,241
	1424	Buckwheat, cushenbury (<i>Eriogonum ovalifolium</i> var. <i>vineum</i>) - Wherever found	E	\$220	\$0	\$220	\$0	\$220
	826	Buckwheat, Ione (incl. Irish Hill) (<i>Eriogonum apricum</i> (incl. var. <i>prostratum</i>)) - Wherever found	E	\$24,500	\$0	\$24,500	\$0	\$24,500
	1057	Buckwheat, scrub (<i>Eriogonum longifolium</i> var. <i>gnaphalifolium</i>) - Wherever found	T	\$0	\$0	\$0	\$12,000	\$12,000
	577	Buckwheat, steamboat (<i>Eriogonum ovalifolium</i> var. <i>williamsiae</i>) - Wherever found	E	\$36,608	\$28,407	\$65,015	\$0	\$65,015

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1404	Buckwheat, Umtanum Desert (<i>Eriogonum codium</i>) - Wherever found	T	\$0	\$1,050	\$1,050	\$0	\$1,050
	486	Bulrush, Northeastern (<i>Scirpus ancistrochaetus</i>) - Wherever found	E	\$70,214	\$34,400	\$104,614	\$852	\$105,466
	1075	Bush-clover, prairie (<i>Lespedeza leptostachya</i>) - Wherever found	T	\$3,500	\$6,585	\$10,085	\$1,300	\$11,385
	315	Bush-mallow, San Clemente Island (<i>Malacothamnus clementinus</i>) - Wherever found	E	\$3,319	\$236,089	\$239,408	\$0	\$239,408
	1058	Bush-mallow, Santa Cruz Island (<i>Malacothamnus fasciculatus</i> var. <i>nesioticus</i>) - Wherever found	E	\$0	\$11,985	\$11,985	\$0	\$11,985
	1073	Buttercup, autumn (<i>Ranunculus aestivalis</i> (= <i>acriformis</i>)) - Wherever found	E	\$5,500	\$5,950	\$11,450	\$0	\$11,450
	397	Butterfly plant, Colorado (<i>Gaura neomexicana</i> var. <i>coloradensis</i>) - Wherever found	T	\$84,244	\$67,313	\$151,557	\$100	\$151,657
	818	Butterweed, Layne's (<i>Senecio layneae</i>) - Wherever found	T	\$16,500	\$8,700	\$25,200	\$0	\$25,200
	1257	Butterwort, Godfrey's (<i>Pinguicula ionantha</i>) - Wherever found	T	\$5,000	\$1,100	\$6,100	\$1,400	\$7,500
	544	Button-celery, San Diego (<i>Eryngium aristulatum</i> var. <i>parishii</i>) - Wherever found	E	\$7,862	\$68,048	\$75,910	\$0	\$75,910
	719	Cactus, Acuna (<i>Echinomastus erectocentrus</i> var. <i>acunensis</i>) - Wherever found	E	\$2,500	\$33,000	\$35,500	\$2,500	\$38,000
	864	Cactus, Arizona hedgehog (<i>Echinocereus triglochidiatus</i> var. <i>arizonicus</i>) - Wherever found	E	\$10,000	\$9,402	\$19,402	\$2,000	\$21,402
	832	Cactus, Bakersfield (<i>Opuntia treleasei</i>) - Wherever found	E	\$21,780	\$2,220	\$24,000	\$0	\$24,000
	751	Cactus, black lace (<i>Echinocereus reichenbachii</i> var. <i>albertii</i>) - Wherever found	E	\$7,200	\$100	\$7,300	\$25,936	\$33,236
	1276	Cactus, Brady pincushion (<i>Pediocactus bradyi</i>) - Wherever found	E	\$2,000	\$2,500	\$4,500	\$2,500	\$7,000
	1385	Cactus, Chisos Mountain hedgehog (<i>Echinocereus chisoensis</i> var. <i>chisoensis</i>) - Wherever found	T	\$1,000	\$1,000	\$2,000	\$0	\$2,000
	1328	Cactus, Cochise pincushion (<i>Coryphantha robbinsiorum</i>) - Wherever found	T	\$4,400	\$100	\$4,500	\$500	\$5,000
	400	Cactus, Colorado hookless (<i>Sclerocactus glaucus</i>) - Wherever found	T	\$18,924	\$130,596	\$149,520	\$1,518	\$151,038
	844	Cactus, Fickeisen plains (<i>Pediocactus peeblesianus fickeiseniae</i>) - Wherever found	E	\$4,000	\$17,000	\$21,000	\$2,000	\$23,000
	1201	Cactus, Florida semaphore (<i>Consolea corallicola</i>) - Wherever found	E	\$4,500	\$3,600	\$8,100	\$600	\$8,700
	932	Cactus, Key tree (<i>Pilosocereus robinii</i>) - Wherever found	E	\$4,500	\$1,053	\$5,553	\$12,000	\$17,553
	892	Cactus, Knowlton's (<i>Pediocactus knowltonii</i>) - Wherever found	E	\$10,921	\$5,800	\$16,721	\$3,125	\$19,846
	466	Cactus, Kuenzler hedgehog (<i>Echinocereus fendleri</i> var. <i>kuenzleri</i>) - Wherever found	E	\$32,000	\$95,300	\$127,300	\$0	\$127,300
	924	Cactus, Lee pincushion (<i>Coryphantha sneedii</i> var. <i>leei</i>) - Wherever found	T	\$17,959	\$0	\$17,959	\$0	\$17,959
	1385	Cactus, Lloyd's Mariposa (<i>Echinomastus mariposensis</i>) - Wherever found	T	\$1,000	\$1,000	\$2,000	\$0	\$2,000
	840	Cactus, Mesa Verde (<i>Sclerocactus mesae-verdae</i>) - Wherever found	T	\$7,420	\$15,875	\$23,295	\$0	\$23,295
	1406	Cactus, Nellie cory (<i>Coryphantha minima</i>) - Wherever found	E	\$1,000	\$0	\$1,000	\$0	\$1,000
	1359	Cactus, Nichol's Turk's head (<i>Echinocactus horizontalonius</i> var. <i>nicholii</i>) - Wherever found	E	\$300	\$2,200	\$2,500	\$500	\$3,000
	690	Cactus, Pariette (<i>Sclerocactus brevispinus</i>) - Wherever found	T	\$36,000	\$7,076	\$43,076	\$0	\$43,076
	911	Cactus, Peebles Navajo (<i>Pediocactus peeblesianus</i> var. <i>peeblesianus</i>) - Wherever found	E	\$200	\$16,300	\$16,500	\$2,000	\$18,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	501	Cactus, Pima pineapple (<i>Coryphantha scheeri</i> var. <i>robustispina</i>) - Wherever found	E	\$12,500	\$77,765	\$90,265	\$3,500	\$93,765
	717	Cactus, San Rafael (<i>Pediocactus despainii</i>) - Wherever found	E	\$16,500	\$21,900	\$38,400	\$0	\$38,400
	835	Cactus, Siler pincushion (<i>Pediocactus</i> (=Echinocactus,=Utahia) <i>sileri</i>) - Wherever found	T	\$4,000	\$18,150	\$22,150	\$1,500	\$23,650
	911	Cactus, Sneed pincushion (<i>Coryphantha sneedii</i> var. <i>sneedii</i>) - Wherever found	E	\$500	\$18,000	\$18,500	\$0	\$18,500
	1286	Cactus, star (<i>Astrophytum asterias</i>) - Wherever found	E	\$6,400	\$200	\$6,600	\$0	\$6,600
	469	Cactus, Tobusch fishhook (<i>Sclerocactus brevihamatus</i> ssp. <i>tobuschii</i>) - Wherever found	E	\$125,000	\$500	\$125,500	\$0	\$125,500
	587	Cactus, Uinta Basin hookless (<i>Sclerocactus wetlandicus</i>) - Wherever found	T	\$47,500	\$14,186	\$61,686	\$0	\$61,686
	657	Cactus, Winkler (<i>Pediocactus winkleri</i>) - Wherever found	T	\$15,500	\$33,400	\$48,900	\$0	\$48,900
	524	Cactus, Wright fishhook (<i>Sclerocactus wrightiae</i>) - Wherever found	E	\$7,500	\$74,400	\$81,900	\$0	\$81,900
	1251	Campion, fringed (<i>Silene polypetala</i>) - Wherever found	E	\$1,000	\$0	\$1,000	\$6,700	\$7,700
	1367	Capa rosa (<i>Callicarpa ampla</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	212	Catchfly, Spalding's (<i>Silene spaldingii</i>) - Wherever found	T	\$476,313	\$28,922	\$505,235	\$46,970	\$552,205
	1418	Cat's-eye, Terlingua Creek (<i>Cryptantha crassipes</i>) - Wherever found	E	\$500	\$0	\$500	\$0	\$500
	878	Ceanothus, coyote (<i>Ceanothus ferrisiae</i>) - Wherever found	E	\$16,390	\$4,000	\$20,390	\$0	\$20,390
	1050	Ceanothus, Pine Hill (<i>Ceanothus roderickii</i>) - Wherever found	E	\$4,150	\$8,000	\$12,150	\$0	\$12,150
	897	Centauray, spring-loving (<i>Centaureum namophilum</i>) - Wherever found	T	\$19,636	\$0	\$19,636	\$0	\$19,636
	374	Chaff-flower, round-leaved (<i>Achyranthes splendens</i> var. <i>rotundata</i>) - Wherever found	E	\$167,952	\$0	\$167,952	\$0	\$167,952
	335	Chaffseed, American (<i>Schwalbea americana</i>) - Wherever found	E	\$92,244	\$108,674	\$200,918	\$2,500	\$203,418
	1200	Checker-mallow, Keck's (<i>Sidalcea keckii</i>) - Wherever found	E	\$8,140	\$629	\$8,769	\$0	\$8,769
	1393	Checker-mallow, Kenwood Marsh (<i>Sidalcea oregana</i> ssp. <i>valida</i>) - Wherever found	E	\$1,650	\$0	\$1,650	\$0	\$1,650
	475	Checker-mallow, Nelson's (<i>Sidalcea nelsoniana</i>) - Wherever found	T	\$119,954	\$400	\$120,354	\$0	\$120,354
	1413	Checkermallow, Wenatchee Mountains (<i>Sidalcea oregana</i> var. <i>calva</i>) - Local Endemic	E	\$452	\$200	\$652	\$0	\$652
	1232	Chumbo, Higo (<i>Harrisia portoricensis</i>) - Wherever found	T	\$8,000	\$0	\$8,000	\$0	\$8,000
	1328	Chupacallos (<i>Pleodendron macranthum</i>) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1084	Clarkia, Presidio (<i>Clarkia franciscana</i>) - Wherever found	E	\$9,020	\$2,000	\$11,020	\$0	\$11,020
	916	Clarkia, Springville (<i>Clarkia springvillensis</i>) - Wherever found	T	\$11,000	\$7,361	\$18,361	\$0	\$18,361
	1367	Cliffrose, Arizona (<i>Purshia</i> (=Cowania) <i>subintegra</i>) - Wherever found	E	\$300	\$1,700	\$2,000	\$500	\$2,500
	1392	Clover, Monterey (<i>Trifolium trichocalyx</i>) - Wherever found	E	\$1,666	\$0	\$1,666	\$0	\$1,666
	340	Clover, running buffalo (<i>Trifolium stoloniferum</i>) - Wherever found	E	\$100,998	\$67,260	\$168,258	\$30,784	\$199,042
	919	Clover, showy Indian (<i>Trifolium amoenum</i>) - Wherever found	E	\$16,500	\$1,700	\$18,200	\$0	\$18,200
	911	Cobana negra (<i>Stahlia monosperma</i>) - Wherever found	T	\$18,500	\$0	\$18,500	\$0	\$18,500
	636	Coneflower, smooth (<i>Echinacea laevigata</i>) - Wherever found	E	\$26,500	\$20,970	\$47,470	\$5,000	\$52,470
	1406	Cory cactus, bunched (<i>Coryphantha ramillosa</i>) - Wherever found	T	\$0	\$1,000	\$1,000	\$0	\$1,000
	1427	Crownbeard, big-leaved (<i>Verbesina dissita</i>) - Wherever found	T	\$0	\$100	\$100	\$0	\$100
	1249	Crownscale, San Jacinto Valley (<i>Atriplex coronata</i> var. <i>notatior</i>) - Wherever found	E	\$7,733	\$0	\$7,733	\$0	\$7,733
	812	Cycladenia, Jones (<i>Cycladenia humilis</i> var. <i>jonesii</i>) - Wherever found	T	\$3,000	\$22,550	\$25,550	\$500	\$26,050
	1332	Daisy, Lakeside (<i>Hymenoxys herbacea</i>) - Wherever found	T	\$3,000	\$1,859	\$4,859	\$0	\$4,859
	1245	Daisy, Parish's (<i>Erigeron parishii</i>) - Wherever found	T	\$220	\$7,600	\$7,820	\$0	\$7,820

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	271	Daisy, Willamette (<i>Erigeron decumbens</i>) - Wherever found	E	\$92,173	\$257,696	\$349,869	\$0	\$349,869
	1273	Dawn-flower, Texas prairie (<i>Hymenoxys texana</i>) - Wherever found	E	\$2,000	\$5,100	\$7,100	\$0	\$7,100
	227	Desert-parsley, Bradshaw's (<i>Lomatium bradshawii</i>) - Wherever found	E	\$260,092	\$229,796	\$489,888	\$0	\$489,888
	1391	Dogweed, ashy (<i>Thymophylla tephroleuca</i>) - Wherever found	E	\$1,400	\$300	\$1,700	\$0	\$1,700
	482	Dropwort, Canby's (<i>Oxypolis canbyi</i>) - Wherever found	E	\$87,000	\$10,410	\$97,410	\$15,000	\$112,410
	1002	Dudleya, Conejo (<i>Dudleya abramsii</i> ssp. <i>parva</i>) - Wherever found	T	\$13,651	\$0	\$13,651	\$0	\$13,651
	1001	Dudleya, marcescent (<i>Dudleya cymosa</i> ssp. <i>marcescens</i>) - Wherever found	T	\$13,684	\$0	\$13,684	\$0	\$13,684
	878	Dudleya, Santa Clara Valley (<i>Dudleya setchellii</i>) - Wherever found	E	\$16,390	\$4,000	\$20,390	\$0	\$20,390
	956	Dudleya, Santa Cruz Island (<i>Dudleya nesiotica</i>) - Wherever found	T	\$16,151	\$0	\$16,151	\$0	\$16,151
	725	Dudleya, Verity's (<i>Dudleya verityi</i>) - Wherever found	T	\$35,614	\$1,000	\$36,614	\$0	\$36,614
	1031	Dwarf-flax, Marin (<i>Hesperolinon congestum</i>) - Wherever found	T	\$10,670	\$2,000	\$12,670	\$0	\$12,670
	1293	'Ena'ena (<i>Pseudognaphalium sandwicensium</i> var. <i>molokaiense</i>) - Wherever found	E	\$5,914	\$500	\$6,414	\$0	\$6,414
	1328	Erubia (<i>Solanum drymophilum</i>) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	470	Evening-primrose, Antioch Dunes (<i>Oenothera deltooides</i> ssp. <i>howellii</i>) - Wherever found	E	\$123,450	\$0	\$123,450	\$0	\$123,450
	532	Evening-primrose, Eureka Valley (<i>Oenothera avita</i> ssp. <i>eurekensis</i>) -	E	\$28,350	\$50,000	\$78,350	\$0	\$78,350
	936	Evening-primrose, San Benito (<i>Camissonia benitensis</i>) - Wherever found	T	\$0	\$17,400	\$17,400	\$0	\$17,400
	421	Fiddleneck, large-flowered (<i>Amsinckia grandiflora</i>) - Wherever found	E	\$15,840	\$124,800	\$140,640	\$0	\$140,640
	1409	Flannelbush, Mexican (<i>Fremontodendron mexicanum</i>) - Wherever found	E	\$250	\$500	\$750	\$0	\$750
	1164	Flannelbush, Pine Hill (<i>Fremontodendron californicum</i> ssp. <i>decumbens</i>) - Wherever found	E	\$0	\$9,500	\$9,500	\$0	\$9,500
	1391	Flax, Carter's small-flowered (<i>Linum carteri carteri</i>) - Wherever found	E	\$1,000	\$0	\$1,000	\$700	\$1,700
	1343	Flax, sand (<i>Linum arenicola</i>) - Wherever found	E	\$2,500	\$700	\$3,200	\$1,300	\$4,500
	1317	Fleabane, Zuni (<i>Erigeron rhizomatus</i>) - Wherever found	T	\$1,000	\$4,200	\$5,200	\$500	\$5,700
	388	Four-o'clock, MacFarlane's (<i>Mirabilis macfarlanei</i>) - Wherever found	T	\$118,648	\$33,600	\$152,248	\$4,983	\$157,231
	1364	Fringepod, Santa Cruz Island (<i>Thysanocarpus conchuliferus</i>) - Wherever found	E	\$2,500	\$200	\$2,700	\$0	\$2,700
	1359	Fringe-tree, pygmy (<i>Chionanthus pygmaeus</i>) - Wherever found	E	\$0	\$0	\$0	\$3,000	\$3,000
	270	Fritillary, Gentner's (<i>Fritillaria gentneri</i>) - Wherever found	E	\$125,262	\$225,000	\$350,262	\$0	\$350,262
	1048	Gardenia (=Na'u), Hawaiian (<i>Gardenia brighamii</i>) - Wherever found	E	\$12,216	\$0	\$12,216	\$0	\$12,216
	537	Geranium, Hawaiian red-flowered (<i>Geranium arboreum</i>) - Wherever found	E	\$14,691	\$62,000	\$76,691	\$0	\$76,691
	1024	Gerardia, sandplain (<i>Agalinis acuta</i>) - Wherever found	E	\$9,600	\$2,640	\$12,240	\$650	\$12,890
	1083	Gilia, Hoffmann's slender-flowered (<i>Gilia tenuiflora</i> ssp. <i>hoffmannii</i>) - Wherever found	E	\$2,500	\$8,530	\$11,030	\$0	\$11,030
	730	Gilia, Monterey (<i>Gilia tenuiflora</i> ssp. <i>arenaria</i>) - Wherever found	E	\$14,560	\$21,635	\$36,195	\$0	\$36,195
	1426	Gladecress, Fleshy-fruit (<i>Leavenworthia crassa</i>) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	853	Glade cress, Kentucky (<i>Leavenworthia exigua laciniata</i>) - Wherever found	T	\$18,500	\$3,710	\$22,210	\$0	\$22,210
	1401	Gladecress, Texas golden (<i>Leavenworthia texana</i>) - Wherever found	E	\$1,000	\$200	\$1,200	\$0	\$1,200
	959	Goetzea, beautiful (<i>Goetzea elegans</i>) - Wherever found	E	\$16,000	\$0	\$16,000	\$0	\$16,000
	1382	Goldenrod, Blue Ridge (<i>Solidago spithamea</i>) - Wherever found	T	\$2,000	\$100	\$2,100	\$0	\$2,100
	671	Goldenrod, Houghton's (<i>Solidago houghtonii</i>) - Wherever found	T	\$47,200	\$100	\$47,300	\$0	\$47,300

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1016	Goldenrod, Short's (<i>Solidago shortii</i>) - Wherever found	E	\$12,000	\$100	\$12,100	\$1,000	\$13,100
	1141	Goldenrod, white-haired (<i>Solidago albopilosa</i>) - Wherever found	T	\$10,000	\$0	\$10,000	\$0	\$10,000
	635	Goldfields, Burke's (<i>Lasthenia burkei</i>) - Wherever found	E	\$9,790	\$42,700	\$52,490	\$0	\$52,490
	332	Goldfields, Contra Costa (<i>Lasthenia conjugens</i>) - Wherever found	E	\$105,778	\$98,780	\$204,558	\$0	\$204,558
	1130	Gooseberry, Miccosukee (<i>Ribes echinellum</i>) - Wherever found	T	\$10,000	\$0	\$10,000	\$200	\$10,200
	572	Gourd, Okeechobee (<i>Cucurbita okeechobeensis</i> ssp. <i>okeechobeensis</i>) - Wherever found	E	\$66,500	\$500	\$67,000	\$200	\$67,200
	821	Grass, Colusa (<i>Neostapfia colusana</i>) - Wherever found	T	\$21,684	\$3,259	\$24,943	\$0	\$24,943
	533	Grass, Eureka Dune (<i>Swallenia alexandrae</i>) -	E	\$28,240	\$50,000	\$78,240	\$0	\$78,240
	1228	Grass, Solano (<i>Tuctoria mucronata</i>) - Wherever found	E	\$8,074	\$0	\$8,074	\$0	\$8,074
	926	Grass, Tennessee yellow-eyed (<i>Xyris tennesseensis</i>) - Wherever found	E	\$8,500	\$4,339	\$12,839	\$5,000	\$17,839
	1038	Ground-plum, Guthrie's (=Pyne's) (<i>Astragalus bibullatus</i>) - Wherever found	E	\$11,500	\$1,028	\$12,528	\$0	\$12,528
	896	Gumplant, Ash Meadows (<i>Grindelia fraxinipratensis</i>) - Wherever found	T	\$19,548	\$100	\$19,648	\$0	\$19,648
	611	Haha (<i>Cyanea acuminata</i>) - Wherever found	E	\$57,087	\$0	\$57,087	\$0	\$57,087
	1265	Haha (<i>Cyanea asarifolia</i>) - Wherever found	E	\$7,339	\$0	\$7,339	\$0	\$7,339
	571	Haha (<i>Cyanea asplenifolia</i>) - Wherever found	E	\$6,067	\$62,000	\$68,067	\$0	\$68,067
	882	Haha (<i>Cyanea calycina</i>) - Wherever found	E	\$6,421	\$13,895	\$20,316	\$0	\$20,316
	1219	Haha (<i>Cyanea copelandii</i> ssp. <i>copelandii</i>) - Wherever found	E	\$8,223	\$0	\$8,223	\$0	\$8,223
	653	Haha (<i>Cyanea copelandii</i> ssp. <i>haleakalaensis</i>) - Wherever found	E	\$8,124	\$41,400	\$49,524	\$0	\$49,524
	592	Haha (<i>Cyanea crispa</i>) - Wherever found	E	\$61,153	\$0	\$61,153	\$0	\$61,153
	1176	Haha (<i>Cyanea dolichopoda</i>) - Wherever found	E	\$9,344	\$0	\$9,344	\$0	\$9,344
	1153	Haha (<i>Cyanea dunbariae</i>) - Wherever found	E	\$9,792	\$0	\$9,792	\$0	\$9,792
	1292	Haha (<i>Cyanea duvalliorum</i>) - Wherever found	E	\$6,420	\$0	\$6,420	\$0	\$6,420
	1176	Haha (<i>Cyanea eleleensis</i>) - Wherever found	E	\$9,344	\$0	\$9,344	\$0	\$9,344
	1258	Haha (<i>Cyanea gibsonii</i>) - Wherever found	E	\$7,432	\$0	\$7,432	\$0	\$7,432
	1281	Haha (<i>Cyanea glabra</i>) - Wherever found	E	\$6,934	\$0	\$6,934	\$0	\$6,934
	1144	Haha (<i>Cyanea grimesiana</i> ssp. <i>grimesiana</i>) - Wherever found	E	\$9,984	\$0	\$9,984	\$0	\$9,984
	433	Haha (<i>Cyanea grimesiana</i> ssp. <i>obatae</i>) - Wherever found	E	\$8,945	\$128,415	\$137,360	\$0	\$137,360
	902	Haha (<i>Cyanea hamatiflora</i> ssp. <i>carlsonii</i>) - Wherever found	E	\$19,248	\$0	\$19,248	\$0	\$19,248
	783	Haha (<i>Cyanea hamatiflora</i> ssp. <i>hamatiflora</i>) - Wherever found	E	\$8,376	\$20,700	\$29,076	\$0	\$29,076
	605	Haha (<i>Cyanea humboldtiana</i>) - Wherever found	E	\$58,087	\$0	\$58,087	\$0	\$58,087
	1180	Haha (<i>Cyanea kolekoleensis</i>) - Wherever found	E	\$9,295	\$0	\$9,295	\$0	\$9,295
	350	Haha (<i>Cyanea koolauensis</i>) - Wherever found	E	\$58,186	\$128,415	\$186,601	\$0	\$186,601
	1176	Haha (<i>Cyanea kuhihewa</i>) - Wherever found	E	\$9,344	\$0	\$9,344	\$0	\$9,344
	799	Haha (<i>Cyanea kunthiana</i>) - Wherever found	E	\$6,519	\$20,700	\$27,219	\$0	\$27,219
	1313	Haha (<i>Cyanea lanceolata</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1066	Haha (<i>Cyanea lobata</i>) - Wherever found	E	\$11,739	\$0	\$11,739	\$0	\$11,739
	444	Haha (<i>Cyanea longiflora</i>) - Wherever found	E	\$6,890	\$128,415	\$135,305	\$0	\$135,305
	1313	Haha (<i>Cyanea magnicalyx</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1308	Haha (<i>Cyanea manni</i>) - Wherever found	E	\$6,067	\$0	\$6,067	\$0	\$6,067
	773	Haha (<i>Cyanea maritae</i>) - Wherever found	E	\$9,640	\$20,700	\$30,340	\$0	\$30,340
	933	Haha (<i>Cyanea marksii</i>) - Wherever found	E	\$17,527	\$0	\$17,527	\$0	\$17,527
	1308	Haha (<i>Cyanea mauiensis</i>) - Wherever found	E	\$6,067	\$0	\$6,067	\$0	\$6,067
	1305	Haha (<i>Cyanea mcelandowneyi</i>) - Wherever found	E	\$6,168	\$0	\$6,168	\$0	\$6,168
	1220	Haha (<i>Cyanea munroi</i>) - Wherever found	E	\$8,173	\$0	\$8,173	\$0	\$8,173

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1158	Haha (Cyanea obtusa) - Wherever found	E	\$9,640	\$0	\$9,640	\$0	\$9,640
	1214	Haha (Cyanea pinnatifida) - Wherever found	E	\$8,314	\$0	\$8,314	\$0	\$8,314
	1110	Haha (Cyanea procera) - Wherever found	E	\$10,553	\$0	\$10,553	\$0	\$10,553
	1158	Haha (Cyanea profuga) - Wherever found	E	\$9,640	\$0	\$9,640	\$0	\$9,640
	1162	Haha (Cyanea purpurellifolia) - Wherever found	E	\$9,539	\$0	\$9,539	\$0	\$9,539
	1269	Haha (Cyanea recta) - Wherever found	T	\$7,288	\$0	\$7,288	\$0	\$7,288
	1103	Haha (Cyanea remyi) - Wherever found	E	\$10,704	\$0	\$10,704	\$0	\$10,704
	1090	Haha (Cyanea rivularis) - Wherever found	E	\$10,913	\$0	\$10,913	\$0	\$10,913
	743	Haha (Cyanea shipmanii) - Wherever found	E	\$21,385	\$13,000	\$34,385	\$0	\$34,385
	822	Haha (Cyanea stictophylla) - Wherever found	E	\$21,844	\$3,000	\$24,844	\$0	\$24,844
	983	Haha (Cyanea st.-johnii) - Wherever found	E	\$14,599	\$0	\$14,599	\$0	\$14,599
	423	Haha (Cyanea superba) - Wherever found	E	\$11,625	\$128,415	\$140,040	\$0	\$140,040
	1151	Haha (Cyanea truncata) - Wherever found	E	\$9,832	\$0	\$9,832	\$0	\$9,832
	1090	Haha (Cyanea undulata) - Wherever found	E	\$10,913	\$0	\$10,913	\$0	\$10,913
	562	Haiwale (Cyrtandra ferripilosa) - Wherever found	E	\$8,122	\$62,000	\$70,122	\$0	\$70,122
	1165	Haiwale (Cyrtandra gracilis) - Wherever found	E	\$9,488	\$0	\$9,488	\$0	\$9,488
	1090	Haiwale (Cyrtandra paliku) - Wherever found	E	\$10,913	\$0	\$10,913	\$0	\$10,913
	1234	Haiwale (Cyrtandra waiolani) - Wherever found	E	\$7,970	\$0	\$7,970	\$0	\$7,970
	1302	Ha`iwale (Cyrtandra crenata) - Wherever found	E	\$6,208	\$0	\$6,208	\$0	\$6,208
	444	Ha`iwale (Cyrtandra dentata) - Wherever found	E	\$6,890	\$128,415	\$135,305	\$0	\$135,305
	1051	Ha`iwale (Cyrtandra filipes) - Wherever found	E	\$12,140	\$0	\$12,140	\$0	\$12,140
	1093	Ha`iwale (Cyrtandra giffardii) - Wherever found	E	\$8,364	\$2,500	\$10,864	\$0	\$10,864
	1313	Ha`iwale (Cyrtandra hematos) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1134	Ha`iwale (Cyrtandra kaulantha) - Wherever found	E	\$10,115	\$0	\$10,115	\$0	\$10,115
	1269	Ha`iwale (Cyrtandra limahuliensis) - Wherever found	T	\$7,288	\$0	\$7,288	\$0	\$7,288
	1202	Ha`iwale (Cyrtandra munroi) - Wherever found	E	\$8,662	\$0	\$8,662	\$0	\$8,662
	1269	Ha`iwale (Cyrtandra oenobarba) - Wherever found	E	\$7,288	\$0	\$7,288	\$0	\$7,288
	1308	Ha`iwale (Cyrtandra oxybapha) - Wherever found	E	\$6,067	\$0	\$6,067	\$0	\$6,067
	1214	Ha`iwale (Cyrtandra polyantha) - Wherever found	E	\$8,314	\$0	\$8,314	\$0	\$8,314
	1211	Ha`iwale (Cyrtandra sessilis) - Wherever found	E	\$8,377	\$0	\$8,377	\$0	\$8,377
	1155	Ha`iwale (Cyrtandra subumbellata) - Wherever found	E	\$9,757	\$0	\$9,757	\$0	\$9,757
	1125	Ha`iwale (Cyrtandra tintinnabula) - Wherever found	E	\$10,320	\$0	\$10,320	\$0	\$10,320
	596	Ha`iwale (Cyrtandra viridiflora) - Wherever found	E	\$60,274	\$0	\$60,274	\$0	\$60,274
	1309	Hala pepe (Pleomele fernaldii) - Wherever found	E	\$6,055	\$0	\$6,055	\$0	\$6,055
	876	Hala pepe (Pleomele forbesii) - Wherever found	E	\$6,592	\$13,895	\$20,487	\$0	\$20,487
	971	Hala pepe (Pleomele hawaiiensis) - Wherever found	E	\$10,260	\$5,000	\$15,260	\$0	\$15,260
	966	Halomtano, Berenghenas (Solanum guamense) - Wherever found	E	\$15,489	\$0	\$15,489	\$0	\$15,489
	580	Halumtano, Cebello (Bulbophyllum guamense) - Wherever found	T	\$64,191	\$100	\$64,291	\$0	\$64,291
	1349	Harebells, Avon Park (Crotalaria avonensis) - Wherever found	E	\$2,000	\$0	\$2,000	\$2,000	\$4,000
	564	Harperella (Ptilimnium nodosum) - Wherever found	E	\$41,501	\$5,910	\$47,411	\$22,417	\$69,828
	1095	Hau kuahiwi (Hibiscadelphus giffardianus) - Wherever found	E	\$8,327	\$2,500	\$10,827	\$0	\$10,827
	1120	Hau kuahiwi (Hibiscadelphus hualalaiensis) - Wherever found	E	\$10,398	\$0	\$10,398	\$0	\$10,398
	1166	Hau kuahiwi (Hibiscadelphus woodii) - Wherever found	E	\$9,475	\$0	\$9,475	\$0	\$9,475
	935	Heartleaf, dwarf-flowered (Hexastylis naniflora) - Wherever found	T	\$4,000	\$13,463	\$17,463	\$0	\$17,463
	1343	Heather, mountain golden (Hudsonia montana) - Wherever found	T	\$4,500	\$0	\$4,500	\$0	\$4,500
	828	Heau (Exocarpos luteolus) - Wherever found	E	\$24,276	\$0	\$24,276	\$0	\$24,276
	1223	Heau (Exocarpos menziesii) -	E	\$6,140	\$2,000	\$8,140	\$0	\$8,140

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1260	Hibiscus, Clay's (<i>Hibiscus clayi</i>) - Wherever found	E	\$7,420	\$0	\$7,420	\$0	\$7,420
	1020	Higuero de sierra (<i>Crescentia portoricensis</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1000	Hoawa (<i>Pittosporum halophilum</i>) - Wherever found	E	\$11,700	\$2,000	\$13,700	\$0	\$13,700
	778	Hoawa (<i>Pittosporum hawaiiense</i>) - Wherever found	E	\$16,891	\$13,000	\$29,891	\$0	\$29,891
	1061	Ho`awa (<i>Pittosporum napaliense</i>) - Wherever found	E	\$11,845	\$0	\$11,845	\$0	\$11,845
	1102	Holei (<i>Ochrosia haleakalae</i>) - Wherever found	E	\$8,211	\$2,500	\$10,711	\$0	\$10,711
	1230	Holei (<i>Ochrosia kilauaeensis</i>) - Wherever found	E	\$8,036	\$0	\$8,036	\$0	\$8,036
	1343	Holly, Cook's (<i>Ilex cookii</i>) - Wherever found	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	681	Honohono (<i>Haplostachys haplostachya</i>) - Wherever found	E	\$12,441	\$33,000	\$45,441	\$0	\$45,441
	424	Howellia, water (<i>Howellia aquatilis</i>) - Wherever found	T	\$127,421	\$7,400	\$134,821	\$5,169	\$139,990
	1231	Hulumoa (<i>Korthalsella degeneri</i>) - Wherever found	E	\$8,002	\$0	\$8,002	\$0	\$8,002
	1257	Hypericum, highlands scrub (<i>Hypericum cumulicola</i>) - Wherever found	E	\$0	\$0	\$0	\$7,500	\$7,500
	484	Iagu, Hayun (=Guam, Tronkon guafi (Rota)) (<i>Serianthes nelsonii</i>) - Wherever found	E	\$109,285	\$0	\$109,285	\$0	\$109,285
	1282	Ihi (<i>Portulaca villosa</i>) - Wherever found	E	\$6,140	\$750	\$6,890	\$0	\$6,890
	1068	Iliau, dwarf (<i>Wilkesia hobbdi</i>) - Wherever found	E	\$7,309	\$4,418	\$11,727	\$0	\$11,727
	776	Ipomopsis, Holy Ghost (<i>Ipomopsis sancti-spiritus</i>) - Wherever found	E	\$30,000	\$0	\$30,000	\$0	\$30,000
	942	Iris, dwarf lake (<i>Iris lacustris</i>) - Wherever found	T	\$16,500	\$500	\$17,000	\$0	\$17,000
	1385	Irisette, white (<i>Sisyrinchium dichotomum</i>) - Wherever found	E	\$2,000	\$0	\$2,000	\$0	\$2,000
	772	Ischaemum, Hilo (<i>Ischaemum byrone</i>) - Wherever found	E	\$9,693	\$20,700	\$30,393	\$0	\$30,393
	979	Ivesia, Ash Meadows (<i>Ivesia kingii</i> var. <i>eremica</i>) - Wherever found	T	\$14,848	\$0	\$14,848	\$0	\$14,848
	512	Ivesia, Webber's (<i>Ivesia webberi</i>) - Wherever found	T	\$57,200	\$30,620	\$87,820	\$0	\$87,820
	1319	Jacquemontia, beach (<i>Jacquemontia reclinata</i>) - Wherever found	E	\$0	\$500	\$500	\$5,000	\$5,500
	680	Jewelflower, California (<i>Caulanthus californicus</i>) - Wherever found	E	\$14,306	\$31,300	\$45,606	\$0	\$45,606
	907	Jewelflower, Metcalf Canyon (<i>Streptanthus albidus</i> ssp. <i>albidus</i>) - Wherever found	E	\$16,390	\$2,600	\$18,990	\$0	\$18,990
	1426	Jewelflower, Tiburon (<i>Streptanthus niger</i>) - Wherever found	E	\$0	\$200	\$200	\$0	\$200
	703	Joint-vetch, Sensitive (<i>Aeschynomene virginica</i>) - Wherever found	T	\$39,626	\$1,620	\$41,246	\$0	\$41,246
	414	Kamakahala (<i>Labordia cyrtandrae</i>) - Wherever found	E	\$14,224	\$128,415	\$142,639	\$0	\$142,639
	1269	Kamakahala (<i>Labordia helleri</i>) - Wherever found	E	\$7,288	\$0	\$7,288	\$0	\$7,288
	910	Kamakahala (<i>Labordia lydgatei</i>) - Wherever found	E	\$18,521	\$0	\$18,521	\$0	\$18,521
	1209	Kamakahala (<i>Labordia pumila</i>) - Wherever found	E	\$8,434	\$0	\$8,434	\$0	\$8,434
	1077	Kamakahala (<i>Labordia tinifolia</i> var. <i>lanaiensis</i>) - Wherever found	E	\$11,143	\$0	\$11,143	\$0	\$11,143
	1055	Kamakahala (<i>Labordia tinifolia</i> var. <i>wahiawaensis</i>) - Wherever found	E	\$12,058	\$0	\$12,058	\$0	\$12,058
	1110	Kamakahala (<i>Labordia triflora</i>) - Wherever found	E	\$10,553	\$0	\$10,553	\$0	\$10,553
	434	Kamanomano (<i>Cenchrus agrimonioides</i>) - Wherever found	E	\$8,878	\$128,415	\$137,293	\$0	\$137,293
	1407	Kamapua`a (<i>Kadua fluviatilis</i>) - Wherever found	E	\$943	\$0	\$943	\$0	\$943
	1166	Kauai hau kuahiwi (<i>Hibiscadelphus distans</i>) - Wherever found	E	\$9,475	\$0	\$9,475	\$0	\$9,475
	1056	Kauila (<i>Colubrina oppositifolia</i>) - Wherever found	E	\$12,041	\$0	\$12,041	\$0	\$12,041
	1123	Kaulu (<i>Pteralyxia kauaiensis</i>) - Wherever found	E	\$10,351	\$0	\$10,351	\$0	\$10,351
	1235	Kaulu (<i>Pteralyxia macrocarpa</i>) - Wherever found	E	\$7,963	\$0	\$7,963	\$0	\$7,963
	688	Kio`ele (<i>Kadua coriacea</i>) - Wherever found	E	\$11,481	\$32,000	\$43,481	\$0	\$43,481
	1041	Kiponapona (<i>Phyllostegia racemosa</i>) - Wherever found	E	\$12,491	\$0	\$12,491	\$0	\$12,491
	1142	Kohe malama malama o kanaloa (<i>Kanaloa kahoolawensis</i>) - Wherever found	E	\$9,998	\$0	\$9,998	\$0	\$9,998
	1124	Koholapehu (<i>Dubautia latifolia</i>) - Wherever found	E	\$10,330	\$0	\$10,330	\$0	\$10,330
	986	Koki`o (<i>Kokia drynarioides</i>) - Wherever found	E	\$14,545	\$0	\$14,545	\$0	\$14,545

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	977	Koki'o (Kokia kauaiensis) - Wherever found	E	\$14,918	\$0	\$14,918	\$0	\$14,918
	1114	Koki'o, Cooke's (Kokia cookei) - Wherever found	E	\$10,502	\$0	\$10,502	\$0	\$10,502
	1254	Koki'o ke'oke'o (Hibiscus arnottianus ssp. immaculatus) - Wherever found	E	\$7,060	\$500	\$7,560	\$0	\$7,560
	1269	Koki'o ke'oke'o (Hibiscus waimeae ssp. hanneriae) - Wherever found	E	\$7,288	\$0	\$7,288	\$0	\$7,288
	1313	Kolea (Myrsine fosbergii) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1137	Kolea (Myrsine juddii) - Wherever found	E	\$10,081	\$0	\$10,081	\$0	\$10,081
	992	Kolea (Myrsine knudsenii) - Wherever found	E	\$14,075	\$0	\$14,075	\$0	\$14,075
	1178	Kolea (Myrsine linearifolia) - Wherever found	T	\$9,316	\$0	\$9,316	\$0	\$9,316
	1082	Kolea (Myrsine mezii) - Wherever found	E	\$11,033	\$0	\$11,033	\$0	\$11,033
	1239	Kolea (Myrsine vaccinioides) - Wherever found	E	\$7,921	\$0	\$7,921	\$0	\$7,921
	1313	Ko'oko'olau (Bidens amplexens) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	660	Ko'oko'olau (Bidens campylotheca ssp. pentamera) - Wherever found	E	\$7,187	\$41,400	\$48,587	\$0	\$48,587
	669	Ko'oko'olau (Bidens campylotheca ssp. waihoiensis) - Wherever found	E	\$6,067	\$41,400	\$47,467	\$0	\$47,467
	1308	Ko'oko'olau (Bidens conjuncta) - Wherever found	E	\$6,067	\$0	\$6,067	\$0	\$6,067
	795	Ko'oko'olau (Bidens micrantha ssp. ctenophylla) - Wherever found	E	\$26,659	\$1,000	\$27,659	\$0	\$27,659
	608	Ko'oko'olau (Bidens micrantha ssp. kalealaha) - Wherever found	E	\$14,251	\$43,400	\$57,651	\$0	\$57,651
	1313	Ko'oko'olau (Bidens wiebkei) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	405	Ko'oloa'ula (Abutilon menziesii) - Wherever found	E	\$14,320	\$134,808	\$149,128	\$0	\$149,128
	1313	Kopa (Kadua cordata remyi) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1017	Kopiko (Psychotria grandiflora) - Wherever found	E	\$13,081	\$0	\$13,081	\$0	\$13,081
	1248	Kopiko (Psychotria hexandra ssp. oahuensis) - Wherever found	E	\$7,777	\$0	\$7,777	\$0	\$7,777
	1268	Kopiko (Psychotria hobdyi) - Wherever found	E	\$7,309	\$0	\$7,309	\$0	\$7,309
	1012	Kuahiwi laukahi (Plantago hawaiiensis) - Wherever found	E	\$11,926	\$1,500	\$13,426	\$0	\$13,426
	355	Kuahiwi laukahi (Plantago princeps) - Wherever found	E	\$18,086	\$163,769	\$181,855	\$0	\$181,855
	1122	Kuawawaenuhu (Schiedea lychnoides) - Wherever found	E	\$10,361	\$0	\$10,361	\$0	\$10,361
	648	Kula wahine noho (Isodendron pyrifolium) - Wherever found	E	\$49,744	\$1,000	\$50,744	\$0	\$50,744
	398	Kulu'i (Nototrichium humile) - Wherever found	E	\$8,036	\$143,069	\$151,105	\$0	\$151,105
	1257	Ladies-tresses, Canelo Hills (Spiranthes delitescens) - Wherever found	E	\$2,000	\$4,000	\$6,000	\$1,500	\$7,500
	744	Ladies-tresses, Navasota (Spiranthes parksii) - Wherever found	E	\$15,000	\$19,300	\$34,300	\$0	\$34,300
	204	Ladies'-tresses, Ute (Spiranthes diluvialis) - Wherever found	T	\$141,712	\$458,843	\$600,555	\$1,208	\$601,763
	1132	Larkspur, Baker's (Delphinium bakeri) - Wherever found	E	\$10,120	\$0	\$10,120	\$0	\$10,120
	357	Larkspur, San Clemente Island (Delphinium variegatum ssp. kinkiense) - Wherever found	E	\$3,319	\$174,927	\$178,246	\$0	\$178,246
	1147	Larkspur, yellow (Delphinium luteum) - Wherever found	E	\$9,900	\$0	\$9,900	\$0	\$9,900
	597	Lau`ehu (Panicum niuhauense) - Wherever found	E	\$11,044	\$49,092	\$60,136	\$0	\$60,136
	1045	Lauhilihi (Schiedea stellarioides) - Wherever found	E	\$12,317	\$0	\$12,317	\$0	\$12,317
	188	Layia, beach (Layia carnosae) - Wherever found	E	\$347,719	\$356,500	\$704,219	\$0	\$704,219
	729	Lead-plant, Crenulate (Amorpha crenulata) - Wherever found	E	\$6,500	\$21,700	\$28,200	\$8,000	\$36,200
	1085	Leather flower, Alabama (Clematis socialis) - Wherever found	E	\$9,000	\$0	\$9,000	\$2,000	\$11,000
	1346	Leather flower, Morefield's (Clematis morefieldii) -	E	\$3,000	\$1,302	\$4,302	\$0	\$4,302
	868	Lehua makanoe (Lysimachia daphnoides) - Wherever found	E	\$21,008	\$0	\$21,008	\$0	\$21,008
	1382	Lessingia, San Francisco (Lessingia germanorum (=L.g. var. germanorum)) - Wherever found	E	\$0	\$2,100	\$2,100	\$0	\$2,100
	1383	Liliwai (Acaena exigua) - Wherever found	E	\$2,074	\$0	\$2,074	\$0	\$2,074

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1406	Lily, Minnesota dwarf trout (<i>Erythronium propullans</i>) - Wherever found	E	\$1,000	\$0	\$1,000	\$0	\$1,000
	1393	Lily, Pitkin Marsh (<i>Lilium pardalinum</i> ssp. <i>pitkinense</i>) - Wherever found	E	\$1,650	\$0	\$1,650	\$0	\$1,650
	480	Lily, Western (<i>Lilium occidentale</i>) - Wherever found	E	\$65,562	\$51,000	\$116,562	\$0	\$116,562
	1354	Liveforever, Santa Barbara Island (<i>Dudleya traskiae</i>) - Wherever found	E	\$2,500	\$1,000	\$3,500	\$0	\$3,500
	508	Lomatium, Cook's (<i>Lomatium cookii</i>) - Wherever found	E	\$32,463	\$58,500	\$90,963	\$0	\$90,963
	373	Loosestrife, rough-leaved (<i>Lysimachia asperulaefolia</i>) - Wherever found	E	\$21,000	\$147,696	\$168,696	\$0	\$168,696
	364	Lotus, San Clemente Island (<i>Acmispon dendroideus</i> var. <i>traskiae</i> (=Lotus d. ssp. <i>traskiae</i>)) - Wherever found	T	\$3,319	\$171,430	\$174,749	\$0	\$174,749
	1268	Lo'ulu (<i>Pritchardia hardyi</i>) - Wherever found	E	\$7,309	\$0	\$7,309	\$0	\$7,309
	446	Lo'ulu (<i>Pritchardia kaalae</i>) - Wherever found	E	\$6,373	\$128,415	\$134,788	\$0	\$134,788
	1247	Lo'ulu (<i>Pritchardia lanigera</i>) - Wherever found	E	\$7,792	\$0	\$7,792	\$0	\$7,792
	950	Lo'ulu (<i>Pritchardia maideniana</i>) - Wherever found	E	\$14,300	\$2,000	\$16,300	\$0	\$16,300
	1215	Lo'ulu (<i>Pritchardia munroi</i>) - Wherever found	E	\$8,313	\$0	\$8,313	\$0	\$8,313
	1177	Lo'ulu (<i>Pritchardia napaliensis</i>) - Wherever found	E	\$9,337	\$0	\$9,337	\$0	\$9,337
	1009	Lo'ulu (<i>Pritchardia remota</i>) - Wherever found	E	\$13,468	\$0	\$13,468	\$0	\$13,468
	1116	Lo'ulu (<i>Pritchardia schattaueri</i>) - Wherever found	E	\$10,483	\$0	\$10,483	\$0	\$10,483
	1080	Lo'ulu (<i>Pritchardia viscosa</i>) - Wherever found	E	\$11,053	\$0	\$11,053	\$0	\$11,053
	1313	Loulu, Baker's (<i>Pritchardia bakeri</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	632	Lousewort, Furbish (<i>Pedicularis furbishiae</i>) - Wherever found	E	\$46,345	\$0	\$46,345	\$6,500	\$52,845
	1216	Love grass, Fosberg's (<i>Eragrostis fosbergii</i>) - Wherever found	E	\$8,263	\$0	\$8,263	\$0	\$8,263
	1359	Lupine, clover (<i>Lupinus tidestromii</i>) - Wherever found	E	\$0	\$3,000	\$3,000	\$0	\$3,000
	233	Lupine, Kincaid's (<i>Lupinus sulphureus</i> ssp. <i>kincaidii</i>) - Wherever found	T	\$209,299	\$271,496	\$480,795	\$0	\$480,795
	939	Lupine, Nipomo Mesa (<i>Lupinus nipomensis</i>) - Wherever found	E	\$17,210	\$0	\$17,210	\$0	\$17,210
	869	Lupine, scrub (<i>Lupinus aridorum</i>) - Wherever found	E	\$0	\$0	\$0	\$21,000	\$21,000
	381	Mahoe (<i>Alectryon macrococcus</i>) - Wherever found	E	\$16,034	\$145,069	\$161,103	\$0	\$161,103
	1006	Makou (<i>Peucedanum sandwicense</i>) - Wherever found	T	\$8,574	\$5,000	\$13,574	\$0	\$13,574
	1285	Makou (<i>Ranunculus hawaiiensis</i>) - Wherever found	E	\$6,605	\$0	\$6,605	\$0	\$6,605
	1313	Makou (<i>Ranunculus mauiensis</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1427	Malacothrix, island (<i>Malacothrix squalida</i>) - Wherever found	E	\$0	\$100	\$100	\$0	\$100
	1287	Malacothrix, Santa Cruz Island (<i>Malacothrix indecora</i>) - Wherever found	E	\$2,500	\$4,027	\$6,527	\$0	\$6,527
	846	Mallow, Gierisch (<i>Sphaeralcea gierischii</i>) - Wherever found	E	\$2,000	\$19,450	\$21,450	\$1,500	\$22,950
	604	Mallow, Kern (<i>Eremalche kernensis</i>) - Wherever found	E	\$34,140	\$24,000	\$58,140	\$0	\$58,140
	1343	Manaca, palma de (<i>Calyptronoma rivalis</i>) - Wherever found	T	\$4,500	\$0	\$4,500	\$0	\$4,500
	1244	Manioc, Walker's (<i>Manihot walkerae</i>) - Wherever found	E	\$7,300	\$560	\$7,860	\$0	\$7,860
	1371	Manzanita, Del Mar (<i>Arctostaphylos glandulosa</i> ssp. <i>crassifolia</i>) - Wherever found	E	\$2,000	\$420	\$2,420	\$0	\$2,420
	540	Manzanita, Franciscan (<i>Arctostaphylos franciscana</i>) - Wherever found	E	\$16,500	\$60,000	\$76,500	\$0	\$76,500
	771	Manzanita, Ione (<i>Arctostaphylos myrtifolia</i>) - Wherever found	T	\$27,500	\$2,931	\$30,431	\$0	\$30,431
	1087	Manzanita, Morro (<i>Arctostaphylos morroensis</i>) - Wherever found	T	\$10,461	\$500	\$10,961	\$0	\$10,961
	1365	Manzanita, pallid (<i>Arctostaphylos pallida</i>) - Wherever found	T	\$0	\$2,600	\$2,600	\$0	\$2,600
	1347	Manzanita, Presidio (<i>Arctostaphylos hookeri</i> var. <i>ravenii</i>) - Wherever found	E	\$2,200	\$2,000	\$4,200	\$0	\$4,200

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1049	Manzanita, Santa Rosa Island (<i>Arctostaphylos confertiflora</i>) - Wherever found	E	\$2,500	\$9,659	\$12,159	\$0	\$12,159
	411	Ma`o hau hele, (=native yellow hibiscus) (<i>Hibiscus brackenridgei</i>) - Wherever found	E	\$15,869	\$129,415	\$145,284	\$0	\$145,284
	1267	Ma`oli`oli (<i>Schiedea apokremnos</i>) - Wherever found	E	\$7,319	\$0	\$7,319	\$0	\$7,319
	695	Ma`oli`oli (<i>Schiedea hawaiiensis</i>) - Wherever found	E	\$10,254	\$32,000	\$42,254	\$0	\$42,254
	1297	Ma`oli`oli (<i>Schiedea kealiae</i>) - Wherever found	E	\$6,337	\$0	\$6,337	\$0	\$6,337
	1313	Ma`oli`oli (<i>Schiedea pubescens</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1269	Mapele (<i>Cyrtandra cyaneoides</i>) - Wherever found	E	\$7,288	\$0	\$7,288	\$0	\$7,288
	1393	Mariposa lily, Tiburon (<i>Calochortus tiburonensis</i>) - Wherever found	T	\$1,650	\$0	\$1,650	\$0	\$1,650
	1356	Meadowfoam, Butte County (<i>Limnanthes floccosa</i> ssp. <i>californica</i>) - Wherever found	E	\$2,500	\$800	\$3,300	\$0	\$3,300
	775	Meadowfoam, large-flowered woolly (<i>Limnanthes pumila</i> ssp. <i>grandiflora</i>) - Wherever found	E	\$30,022	\$0	\$30,022	\$0	\$30,022
	636	Meadowfoam, Sebastopol (<i>Limnanthes vinculans</i>) - Wherever found	E	\$10,670	\$41,800	\$52,470	\$0	\$52,470
	1015	Meadowrue, Cooley's (<i>Thalictrum cooleyi</i>) - Wherever found	E	\$11,000	\$0	\$11,000	\$2,200	\$13,200
	337	Mehamehame (<i>Flueggea neowawraea</i>) - Wherever found	E	\$57,165	\$145,069	\$202,234	\$0	\$202,234
	1159	Mesa-mint, Otay (<i>Pogogyne nudiuscula</i>) - Wherever found	E	\$6,952	\$2,684	\$9,636	\$0	\$9,636
	549	Mesa-mint, San Diego (<i>Pogogyne abramsii</i>) - Wherever found	E	\$7,359	\$66,580	\$73,939	\$0	\$73,939
	1323	Milkpea, Small's (<i>Galactia smallii</i>) - Wherever found	E	\$4,500	\$0	\$4,500	\$700	\$5,200
	637	Milk-vetch, Applegate's (<i>Astragalus applegatei</i>) - Wherever found	E	\$52,364	\$0	\$52,364	\$0	\$52,364
	979	Milk-vetch, Ash meadows (<i>Astragalus phoenix</i>) - Wherever found	T	\$14,848	\$0	\$14,848	\$0	\$14,848
	513	Milk-vetch, Braunton's (<i>Astragalus brauntonii</i>) - Wherever found	E	\$84,668	\$2,684	\$87,352	\$0	\$87,352
	1393	Milk-vetch, Clara Hunt's (<i>Astragalus clarianus</i>) - Wherever found	E	\$1,650	\$0	\$1,650	\$0	\$1,650
	862	Milk-vetch, Coachella Valley (<i>Astragalus lentiginosus</i> var. <i>coachellae</i>) - Wherever found	E	\$18,370	\$3,500	\$21,870	\$0	\$21,870
	1335	Milk-vetch, Cushenbury (<i>Astragalus albens</i>) - Wherever found	E	\$220	\$4,500	\$4,720	\$0	\$4,720
	1316	Milkvetch, Deseret (<i>Astragalus desereticus</i>) - Wherever found	T	\$3,500	\$2,250	\$5,750	\$0	\$5,750
	682	Milk-vetch, Fish Slough (<i>Astragalus lentiginosus</i> var. <i>piscinensis</i>) - Wherever found	T	\$34,320	\$11,000	\$45,320	\$0	\$45,320
	815	Milk-vetch, heliotrope (<i>Astragalus montii</i>) - Wherever found	T	\$23,500	\$2,250	\$25,750	\$0	\$25,750
	904	Milk-vetch, Holmgren (<i>Astragalus holmgreniorum</i>) - Wherever found	E	\$5,567	\$12,000	\$17,567	\$1,500	\$19,067
	754	Milk-vetch, Jesup's (<i>Astragalus robbinsii</i> var. <i>jesupi</i>) - Wherever found	E	\$20,053	\$0	\$20,053	\$13,000	\$33,053
	367	Milk-vetch, Lane Mountain (<i>Astragalus jaegerianus</i>) - Wherever found	E	\$0	\$171,853	\$171,853	\$0	\$171,853
	960	Milk-vetch, Mancos (<i>Astragalus humillimus</i>) - Wherever found	E	\$7,200	\$8,600	\$15,800	\$0	\$15,800
	900	Milkvetch, Osterhout (<i>Astragalus osterhoutii</i>) - Wherever found	E	\$0	\$19,312	\$19,312	\$0	\$19,312
	295	Milk-vetch, Peirson's (<i>Astragalus magdalenae</i> var. <i>peirsonii</i>) - Wherever found	T	\$17,270	\$260,000	\$277,270	\$0	\$277,270
	521	Milk-vetch, Sentry (<i>Astragalus cremnophylax</i> var. <i>cremnophylax</i>) - Wherever found	E	\$2,000	\$80,100	\$82,100	\$1,000	\$83,100
	911	Milk-vetch, Shivwits (<i>Astragalus ampullarioides</i>) - Wherever found	E	\$6,500	\$12,000	\$18,500	\$0	\$18,500
	1262	Milk-vetch, triple-ribbed (<i>Astragalus tricarinatus</i>) - Wherever found	E	\$4,400	\$3,000	\$7,400	\$0	\$7,400
	841	Milk-vetch, Ventura Marsh (<i>Astragalus pycnostachyus</i> var. <i>lanosissimus</i>) - Wherever found	E	\$18,517	\$4,636	\$23,153	\$0	\$23,153
	328	Milkweed, Mead's (<i>Asclepias meadii</i>) - Wherever found	T	\$16,963	\$181,903	\$198,866	\$10,800	\$209,666
	774	Milkweed, Welsh's (<i>Asclepias welshii</i>) - Wherever found	T	\$0	\$29,100	\$29,100	\$1,000	\$30,100

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1284	Mint, Garrett's (<i>Dicerandra christmanii</i>) - Wherever found	E	\$6,000	\$0	\$6,000	\$700	\$6,700
	995	Mint, Lakela's (<i>Dicerandra immaculata</i>) - Wherever found	E	\$2,000	\$0	\$2,000	\$12,000	\$14,000
	1311	Mint, longspurred (<i>Dicerandra cornutissima</i>) - Wherever found	E	\$0	\$0	\$0	\$6,000	\$6,000
	848	Mint, scrub (<i>Dicerandra frutescens</i>) - Wherever found	E	\$0	\$21,700	\$21,700	\$1,000	\$22,700
	360	Monardella, willowly (<i>Monardella viminea</i>) - Wherever found	E	\$4,252	\$172,780	\$177,032	\$0	\$177,032
	964	Monkey-flower, Michigan (<i>Mimulus michiganensis</i>) - Wherever found	E	\$14,000	\$1,540	\$15,540	\$0	\$15,540
	535	Monkeyflower, Vandenberg (<i>Diplacus vanderbergensis</i>) - Wherever found	E	\$0	\$77,500	\$77,500	\$0	\$77,500
	1263	Monkshood, northern wild (<i>Aconitum noveboracense</i>) - Wherever found	T	\$5,100	\$1,769	\$6,869	\$500	\$7,369
	1033	Morning-glory, Stebbins' (<i>Calystegia stebbinsii</i>) - Wherever found	E	\$4,150	\$8,500	\$12,650	\$0	\$12,650
	606	Mountainbalm, Indian Knob (<i>Eriodictyon altissimum</i>) - Wherever found	E	\$55,579	\$2,318	\$57,897	\$0	\$57,897
	1337	Mustard, Carter's (<i>Warea carteri</i>) - Wherever found	E	\$0	\$200	\$200	\$4,500	\$4,700
	766	Mustard, Penland alpine fen (<i>Eutrema penlandii</i>) - Wherever found	T	\$15,409	\$16,056	\$31,465	\$0	\$31,465
	1364	Mustard, slender-petaled (<i>Thelypodium stenopetalum</i>) - Wherever found	E	\$2,500	\$200	\$2,700	\$0	\$2,700
	1173	Naenae (<i>Dubautia kalalauensis</i>) - Wherever found	E	\$9,395	\$0	\$9,395	\$0	\$9,395
	1176	Naenae (<i>Dubautia kenwoodii</i>) - Wherever found	E	\$9,344	\$0	\$9,344	\$0	\$9,344
	444	Na'ena'e (<i>Dubautia herbstobatae</i>) - Wherever found	E	\$6,890	\$128,415	\$135,305	\$0	\$135,305
	1269	Na'ena'e (<i>Dubautia imbricata</i> ssp. <i>imbricata</i>) - Wherever found	E	\$7,288	\$0	\$7,288	\$0	\$7,288
	1060	Na'ena'e (<i>Dubautia pauciflorula</i>) - Wherever found	E	\$11,927	\$0	\$11,927	\$0	\$11,927
	1114	Na'ena'e (<i>Dubautia plantaginea</i> ssp. <i>humilis</i>) - Wherever found	E	\$10,502	\$0	\$10,502	\$0	\$10,502
	1269	Na'ena'e (<i>Dubautia plantaginea</i> ssp. <i>magnifolia</i>) - Wherever found	E	\$7,288	\$0	\$7,288	\$0	\$7,288
	1178	Na'ena'e (<i>Dubautia waialealae</i>) - Wherever found	E	\$9,316	\$0	\$9,316	\$0	\$9,316
	1022	Nani wai'ale'ale (<i>Viola kauaiensis</i> var. <i>wahiawaensis</i>) - Wherever found	E	\$12,961	\$0	\$12,961	\$0	\$12,961
	351	Nanu (<i>Gardenia manni</i>) - Wherever found	E	\$58,087	\$128,415	\$186,502	\$0	\$186,502
	786	Nanu (<i>Gardenia remyi</i>) - Wherever found	E	\$7,985	\$20,700	\$28,685	\$0	\$28,685
	988	Naupaka, dwarf (<i>Scaevola coriacea</i>) - Wherever found	E	\$9,454	\$5,000	\$14,454	\$0	\$14,454
	1331	Navarretia, many-flowered (<i>Navarretia leucocephala</i> ssp. <i>plieantha</i>) - Wherever found	E	\$2,500	\$2,400	\$4,900	\$0	\$4,900
	656	Navarretia, spreading (<i>Navarretia fossalis</i>) - Wherever found	T	\$7,722	\$41,188	\$48,910	\$0	\$48,910
	1269	Nehe (<i>Lipochaeta fauriei</i>) - Wherever found	E	\$7,288	\$0	\$7,288	\$0	\$7,288
	873	Nehe (<i>Lipochaeta lobata</i> var. <i>leptophylla</i>) - Wherever found	E	\$6,208	\$14,604	\$20,812	\$0	\$20,812
	1078	Nehe (<i>Lipochaeta micrantha</i>) - Wherever found	E	\$11,133	\$0	\$11,133	\$0	\$11,133
	1100	Nehe (<i>Lipochaeta waimeaensis</i>) - Wherever found	E	\$10,762	\$0	\$10,762	\$0	\$10,762
	1237	Nehe (<i>Melanthera kamolensis</i>) - Wherever found	E	\$7,941	\$0	\$7,941	\$0	\$7,941
	447	Nehe (<i>Melanthera tenuifolia</i>) - Wherever found	E	\$6,353	\$128,415	\$134,768	\$0	\$134,768
	354	Nioi (<i>Eugenia koolauensis</i>) - Wherever found	E	\$54,099	\$128,415	\$182,514	\$0	\$182,514
	711	Niterwort, Amargosa (<i>Nitrophila mohavensis</i>) - Wherever found	E	\$11,848	\$28,000	\$39,848	\$0	\$39,848
	1191	No Common Name (<i>Abutilon eremitopetalum</i>) - Wherever found	E	\$9,034	\$0	\$9,034	\$0	\$9,034
	317	No Common Name (<i>Abutilon sandwicense</i>) - Wherever found	E	\$6,208	\$229,522	\$235,730	\$0	\$235,730
	1298	No Common Name (<i>Achyranthes mutica</i>) - Wherever found	E	\$6,323	\$0	\$6,323	\$0	\$6,323
	1349	No Common Name (<i>Agave eggersiana</i>) - Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	1167	No Common Name (<i>Amaranthus brownii</i>) - Wherever found	E	\$9,462	\$0	\$9,462	\$0	\$9,462
	1020	No Common Name (<i>Aristida chaseae</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1257	No Common Name (Auerodendron pauciflorum) - Wherever found	E	\$7,500	\$0	\$7,500	\$0	\$7,500
	1243	No Common Name (Bidens hillebrandiana ssp. hillebrandiana) - Wherever found	E	\$7,870	\$0	\$7,870	\$0	\$7,870
	781	No Common Name (Bonamia menziesii) - Wherever found	E	\$14,769	\$14,604	\$29,373	\$0	\$29,373
	1354	No Common Name (Calyptanthus thomasiana) - Wherever found	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	1328	No Common Name (Catesbaea melanocarpa) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1385	No Common Name (Chamaecrista glandulosa var. mirabilis) - Wherever found	E	\$2,000	\$0	\$2,000	\$0	\$2,000
	1232	No Common Name (Cordia bellonis) - Wherever found	E	\$8,000	\$0	\$8,000	\$0	\$8,000
	1382	No Common Name (Cranichis ricartii) - Wherever found	E	\$2,000	\$100	\$2,100	\$0	\$2,100
	1258	No Common Name (Cyanea kauaulaensis) - Wherever found	E	\$7,432	\$0	\$7,432	\$0	\$7,432
	1308	No Common Name (Cyperus fauriei) - Wherever found	E	\$6,067	\$0	\$6,067	\$0	\$6,067
	1313	No Common Name (Cyperus neokunthianus) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1128	No Common Name (Cyperus pennatiformis) - Wherever found	E	\$10,207	\$0	\$10,207	\$0	\$10,207
	833	No Common Name (Cyrandra nanawaleensis) - Wherever found	E	\$6,067	\$17,766	\$23,833	\$0	\$23,833
	1062	No Common Name (Cyrandra wagneri) - Wherever found	E	\$11,843	\$0	\$11,843	\$0	\$11,843
	1343	No Common Name (Daphnopsis hellerana) - Wherever found	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	1094	No Common Name (Delissea rhytidosperra) - Wherever found	E	\$10,862	\$0	\$10,862	\$0	\$10,862
	1098	No Common Name (Delissea undulata) - Wherever found	E	\$10,789	\$0	\$10,789	\$0	\$10,789
	474	No Common Name (Dendrobium guamense) - Wherever found	T	\$62,275	\$58,105	\$120,380	\$0	\$120,380
	931	No Common Name (Eugenia bryanii) - Wherever found	E	\$17,582	\$0	\$17,582	\$0	\$17,582
	1115	No Common Name (Eugenia woodburyana) - Wherever found	E	\$10,500	\$0	\$10,500	\$0	\$10,500
	720	No Common Name (Festuca hawaiiensis) - Wherever found	E	\$5,914	\$32,000	\$37,914	\$0	\$37,914
	1304	No Common Name (Festuca molokaiensis) - Wherever found	E	\$6,198	\$0	\$6,198	\$0	\$6,198
	946	No Common Name (Geocarpon minimum) - Wherever found	T	\$10,000	\$6,273	\$16,273	\$400	\$16,673
	1385	No Common Name (Gesneria pauciflora) - Wherever found	T	\$2,000	\$0	\$2,000	\$0	\$2,000
	1396	No Common Name (Gonocalyx concolor) - Wherever found	E	\$1,500	\$0	\$1,500	\$0	\$1,500
	1274	No Common Name (Gouania hillebrandii) - Wherever found	E	\$7,060	\$0	\$7,060	\$0	\$7,060
	1172	No Common Name (Gouania meyenii) - Wherever found	E	\$9,432	\$0	\$9,432	\$0	\$9,432
	416	No Common Name (Gouania vitifolia) - Wherever found	E	\$14,122	\$128,415	\$142,537	\$0	\$142,537
	347	No Common Name (Hesperomannia arborescens) - Wherever found	E	\$61,239	\$128,415	\$189,654	\$0	\$189,654
	427	No Common Name (Hesperomannia arbuscula) - Wherever found	E	\$10,162	\$128,415	\$138,577	\$0	\$138,577
	1260	No Common Name (Hesperomannia lydgatei) - Wherever found	E	\$7,420	\$0	\$7,420	\$0	\$7,420
	1367	No Common Name (Ilex sintenisii) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	413	No Common Name (Kadua degeneri) - Wherever found	E	\$14,667	\$128,415	\$143,082	\$0	\$143,082
	1313	No Common Name (Kadua hauptuensis) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	406	No Common Name (Kadua parvula) - Wherever found	E	\$7,022	\$141,690	\$148,712	\$0	\$148,712
	1176	No Common Name (Kadua st. johnii) - Wherever found	E	\$9,344	\$0	\$9,344	\$0	\$9,344
	1170	No Common Name (Keysseria (=Lagenifera) erici) - Wherever found	E	\$9,448	\$0	\$9,448	\$0	\$9,448
	1170	No Common Name (Keysseria (=Lagenifera) helenae) - Wherever found	E	\$9,448	\$0	\$9,448	\$0	\$9,448
	1258	No Common Name (Labordia lorenciana) - Wherever found	E	\$7,432	\$0	\$7,432	\$0	\$7,432
	1226	No Common Name (Lepanthes eltoroensis) - Wherever found	E	\$8,000	\$100	\$8,100	\$0	\$8,100
	1313	No Common Name (Lepidium orbiculare) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1359	No Common Name (Leptocereus grantianus) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	708	No Common Name (Lipochaeta venosa) - Wherever found	E	\$8,407	\$32,000	\$40,407	\$0	\$40,407
	1145	No Common Name (Lobelia koolauensis) - Wherever found	E	\$9,961	\$0	\$9,961	\$0	\$9,961

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1151	No Common Name (<i>Lobelia monostachya</i>) - Wherever found	E	\$9,832	\$0	\$9,832	\$0	\$9,832
	871	No Common Name (<i>Lobelia niihauensis</i>) - Wherever found	E	\$7,582	\$13,275	\$20,857	\$0	\$20,857
	994	No Common Name (<i>Lobelia oahuensis</i>) - Wherever found	E	\$14,041	\$0	\$14,041	\$0	\$14,041
	1020	No Common Name (<i>Lyonia truncata</i> var. <i>proctorii</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1227	No Common Name (<i>Lysimachia filifolia</i>) - Wherever found	E	\$8,089	\$0	\$8,089	\$0	\$8,089
	1173	No Common Name (<i>Lysimachia iniki</i>) - Wherever found	E	\$9,395	\$0	\$9,395	\$0	\$9,395
	1196	No Common Name (<i>Lysimachia lydgatei</i>) - Wherever found	E	\$8,984	\$0	\$8,984	\$0	\$8,984
	1110	No Common Name (<i>Lysimachia maxima</i>) - Wherever found	E	\$10,553	\$0	\$10,553	\$0	\$10,553
	1097	No Common Name (<i>Lysimachia pendens</i>) - Wherever found	E	\$10,813	\$0	\$10,813	\$0	\$10,813
	1182	No Common Name (<i>Lysimachia scopulensis</i>) - Wherever found	E	\$9,244	\$0	\$9,244	\$0	\$9,244
	1076	No Common Name (<i>Lysimachia venosa</i>) - Wherever found	E	\$11,372	\$0	\$11,372	\$0	\$11,372
	1414	No Common Name (<i>Maesa walkeri</i>) - Wherever found	T	\$614	\$0	\$614	\$0	\$614
	1354	No Common Name (<i>Mitracarpus maxwelliae</i>) - Wherever found	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	1359	No Common Name (<i>Mitracarpus polycladus</i>) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1343	No Common Name (<i>Myrcia paganii</i>) - Wherever found	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	393	No Common Name (<i>Neraudia angulata</i>) - Wherever found	E	\$11,118	\$141,690	\$152,808	\$0	\$152,808
	634	No Common Name (<i>Neraudia ovata</i>) - Wherever found	E	\$18,000	\$34,500	\$52,500	\$0	\$52,500
	1105	No Common Name (<i>Neraudia sericea</i>) - Wherever found	E	\$10,633	\$0	\$10,633	\$0	\$10,633
	796	No Common Name (<i>Nervilia jacksoniae</i>) - Wherever found	T	\$27,454	\$100	\$27,554	\$0	\$27,554
	842	No Common Name (<i>Nesogenes rotensis</i>) - Wherever found	E	\$23,142	\$0	\$23,142	\$0	\$23,142
	747	No Common Name (<i>Osmoxylon mariannense</i>) - Wherever found	E	\$33,720	\$0	\$33,720	\$0	\$33,720
	1027	No Common Name (<i>Phyllanthus saffordii</i>) - Wherever found	E	\$12,826	\$0	\$12,826	\$0	\$12,826
	559	No Common Name (<i>Phyllostegia bracteata</i>) - Wherever found	E	\$8,475	\$62,000	\$70,475	\$0	\$70,475
	566	No Common Name (<i>Phyllostegia brevidens</i>) - Wherever found	E	\$7,432	\$62,000	\$69,432	\$0	\$69,432
	1117	No Common Name (<i>Phyllostegia floribunda</i>) - Wherever found	E	\$10,432	\$0	\$10,432	\$0	\$10,432
	1193	No Common Name (<i>Phyllostegia glabra</i> var. <i>lanaiensis</i>) - Wherever found	E	\$9,016	\$0	\$9,016	\$0	\$9,016
	561	No Common Name (<i>Phyllostegia haliakalae</i>) - Wherever found	E	\$8,254	\$62,000	\$70,254	\$0	\$70,254
	1313	No Common Name (<i>Phyllostegia helleri</i>) -	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	349	No Common Name (<i>Phyllostegia hirsuta</i>) - Wherever found	E	\$60,274	\$128,415	\$188,689	\$0	\$188,689
	1104	No Common Name (<i>Phyllostegia hispida</i>) - Wherever found	E	\$10,685	\$0	\$10,685	\$0	\$10,685
	435	No Common Name (<i>Phyllostegia kaalaensis</i>) - Wherever found	E	\$8,784	\$128,415	\$137,199	\$0	\$137,199
	1166	No Common Name (<i>Phyllostegia knudsenii</i>) - Wherever found	E	\$9,475	\$0	\$9,475	\$0	\$9,475
	1152	No Common Name (<i>Phyllostegia manni</i>) - Wherever found	E	\$9,823	\$0	\$9,823	\$0	\$9,823
	429	No Common Name (<i>Phyllostegia mollis</i>) - Wherever found	E	\$9,916	\$128,415	\$138,331	\$0	\$138,331
	922	No Common Name (<i>Phyllostegia parviflora</i>) - Wherever found	E	\$18,038	\$0	\$18,038	\$0	\$18,038
	1131	No Common Name (<i>Phyllostegia pilosa</i>) - Wherever found	E	\$10,176	\$0	\$10,176	\$0	\$10,176
	1072	No Common Name (<i>Phyllostegia renovans</i>) - Wherever found	E	\$11,476	\$0	\$11,476	\$0	\$11,476
	1064	No Common Name (<i>Phyllostegia stachyoides</i>) - Wherever found	E	\$9,264	\$2,500	\$11,764	\$0	\$11,764
	1199	No Common Name (<i>Phyllostegia velutina</i>) - Wherever found	E	\$8,793	\$0	\$8,793	\$0	\$8,793
	1081	No Common Name (<i>Phyllostegia waimeae</i>) - Wherever found	E	\$11,044	\$0	\$11,044	\$0	\$11,044
	1030	No Common Name (<i>Phyllostegia warshaueri</i>) - Wherever found	E	\$12,704	\$0	\$12,704	\$0	\$12,704
	1166	No Common Name (<i>Phyllostegia wawrana</i>) - Wherever found	E	\$9,475	\$0	\$9,475	\$0	\$9,475
	1010	No Common Name (<i>Platanthera holochila</i>) - Wherever found	E	\$13,363	\$100	\$13,463	\$0	\$13,463
	617	No Common Name (<i>Platydesma cornuta</i> var. <i>cornuta</i>) - Wherever found	E	\$56,227	\$0	\$56,227	\$0	\$56,227

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	880	No Common Name (<i>Platydesma cornuta</i> var. <i>decurrens</i>) - Wherever found	E	\$7,456	\$12,875	\$20,331	\$0	\$20,331
	1067	No Common Name (<i>Platydesma remyi</i>) - Wherever found	E	\$11,731	\$0	\$11,731	\$0	\$11,731
	1268	No Common Name (<i>Poa siphonoglossa</i>) - Wherever found	E	\$7,309	\$0	\$7,309	\$0	\$7,309
	980	No Common Name (<i>Polyscias bisattenuata</i>) - Wherever found	E	\$14,839	\$0	\$14,839	\$0	\$14,839
	1025	No Common Name (<i>Polyscias flynnii</i>) - Wherever found	E	\$12,862	\$0	\$12,862	\$0	\$12,862
	1180	No Common Name (<i>Polyscias lydgatei</i>) - Wherever found	E	\$9,295	\$0	\$9,295	\$0	\$9,295
	612	No Common Name (<i>Polyscias racemosa</i>) - Wherever found	E	\$56,956	\$0	\$56,956	\$0	\$56,956
	1225	No Common Name (<i>Remya kauaiensis</i>) - Wherever found	E	\$8,120	\$0	\$8,120	\$0	\$8,120
	1088	No Common Name (<i>Remya montgomeryi</i>) - Wherever found	E	\$10,933	\$0	\$10,933	\$0	\$10,933
	445	No Common Name (<i>Sanicula mariversa</i>) - Wherever found	E	\$6,766	\$128,415	\$135,181	\$0	\$135,181
	1036	No Common Name (<i>Sanicula purpurea</i>) - Wherever found	E	\$12,596	\$0	\$12,596	\$0	\$12,596
	662	No Common Name (<i>Sanicula sandwicensis</i>) - Wherever found	E	\$6,605	\$41,900	\$48,505	\$0	\$48,505
	1313	No Common Name (<i>Santalum involutum</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1174	No Common Name (<i>Schiedea attenuata</i>) - Wherever found	E	\$9,374	\$0	\$9,374	\$0	\$9,374
	1148	No Common Name (<i>Schiedea diffusa</i> ssp. <i>macraei</i>) - Wherever found	E	\$9,896	\$0	\$9,896	\$0	\$9,896
	1296	No Common Name (<i>Schiedea diffusa</i> subsp. <i>diffusa</i>) - Wherever found	E	\$6,379	\$0	\$6,379	\$0	\$6,379
	557	No Common Name (<i>Schiedea haleakalensis</i>) - Wherever found	E	\$9,065	\$62,000	\$71,065	\$0	\$71,065
	1267	No Common Name (<i>Schiedea helleri</i>) - Wherever found	E	\$7,319	\$0	\$7,319	\$0	\$7,319
	898	No Common Name (<i>Schiedea hookeri</i>) - Wherever found	E	\$6,390	\$13,075	\$19,465	\$0	\$19,465
	1222	No Common Name (<i>Schiedea jacobii</i>) - Wherever found	E	\$8,152	\$0	\$8,152	\$0	\$8,152
	422	No Common Name (<i>Schiedea kaalae</i>) - Wherever found	E	\$11,921	\$128,415	\$140,336	\$0	\$140,336
	975	No Common Name (<i>Schiedea kauaiensis</i>) - Wherever found	E	\$14,999	\$0	\$14,999	\$0	\$14,999
	1203	No Common Name (<i>Schiedea laui</i>) - Wherever found	E	\$8,641	\$0	\$8,641	\$0	\$8,641
	1278	No Common Name (<i>Schiedea lydgatei</i>) - Wherever found	E	\$6,959	\$0	\$6,959	\$0	\$6,959
	1021	No Common Name (<i>Schiedea membranacea</i>) - Wherever found	E	\$12,971	\$0	\$12,971	\$0	\$12,971
	437	No Common Name (<i>Schiedea nuttallii</i>) - Wherever found	E	\$8,537	\$128,415	\$136,952	\$0	\$136,952
	438	No Common Name (<i>Schiedea obovata</i>) - Wherever found	E	\$8,438	\$128,415	\$136,853	\$0	\$136,853
	1307	No Common Name (<i>Schiedea salicaria</i>) - Wherever found	E	\$6,097	\$0	\$6,097	\$0	\$6,097
	1307	No Common Name (<i>Schiedea sarmentosa</i>) - Wherever found	E	\$6,097	\$0	\$6,097	\$0	\$6,097
	1181	No Common Name (<i>Schiedea spergulina</i> var. <i>leiopoda</i>) - Wherever found	E	\$9,275	\$0	\$9,275	\$0	\$9,275
	1224	No Common Name (<i>Schiedea spergulina</i> var. <i>spergulina</i>) - Wherever found	T	\$8,124	\$0	\$8,124	\$0	\$8,124
	448	No Common Name (<i>Schiedea trinervis</i>) - Wherever found	E	\$6,238	\$128,415	\$134,653	\$0	\$134,653
	1129	No Common Name (<i>Schiedea verticillata</i>) - Wherever found	E	\$10,204	\$0	\$10,204	\$0	\$10,204
	1118	No Common Name (<i>Schiedea viscosa</i>) - Wherever found	E	\$10,412	\$0	\$10,412	\$0	\$10,412
	1194	No Common Name (<i>Schoepfia arenaria</i>) - Wherever found	T	\$9,000	\$0	\$9,000	\$0	\$9,000
	1258	No Common Name (<i>Sicyos lanceoloideus</i>) - Wherever found	E	\$7,432	\$0	\$7,432	\$0	\$7,432
	1109	No Common Name (<i>Silene alexandri</i>) - Wherever found	E	\$10,573	\$0	\$10,573	\$0	\$10,573
	716	No Common Name (<i>Silene hawaiiensis</i>) - Wherever found	T	\$7,175	\$32,000	\$39,175	\$0	\$39,175
	714	No Common Name (<i>Silene lanceolata</i>) - Wherever found	E	\$7,399	\$32,000	\$39,399	\$0	\$39,399
	847	No Common Name (<i>Silene perlmanii</i>) - Wherever found	E	\$9,801	\$13,075	\$22,876	\$0	\$22,876
	613	No Common Name (<i>Spermolepis hawaiiensis</i>) - Wherever found	E	\$8,132	\$48,530	\$56,662	\$0	\$56,662
	706	No Common Name (<i>Stenogyne angustifolia</i> var. <i>angustifolia</i>) - Wherever found	E	\$8,654	\$32,000	\$40,654	\$0	\$40,654

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1190	No Common Name (<i>Stenogyne bifida</i>) - Wherever found	E	\$9,055	\$0	\$9,055	\$0	\$9,055
	1091	No Common Name (<i>Stenogyne campanulata</i>) - Wherever found	E	\$10,882	\$0	\$10,882	\$0	\$10,882
	1233	No Common Name (<i>Stenogyne cranwelliae</i>) - Wherever found	E	\$7,990	\$0	\$7,990	\$0	\$7,990
	1271	No Common Name (<i>Stenogyne kaalae</i> ssp. <i>sherrfii</i>) - Wherever found	E	\$7,181	\$0	\$7,181	\$0	\$7,181
	439	No Common Name (<i>Stenogyne kanehoana</i>) - Wherever found	E	\$8,283	\$128,415	\$136,698	\$0	\$136,698
	1138	No Common Name (<i>Stenogyne kauaulaensis</i>) - Wherever found	E	\$10,015	\$0	\$10,015	\$0	\$10,015
	1091	No Common Name (<i>Stenogyne kealiae</i>) - Wherever found	E	\$10,882	\$0	\$10,882	\$0	\$10,882
	511	No Common Name (<i>Tabernaemontana rotensis</i>) - Wherever found	T	\$36,833	\$53,215	\$90,048	\$0	\$90,048
	1367	No Common Name (<i>Ternstroemia subsessilis</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	715	No Common Name (<i>Tetramolopium arenarium</i>) - Wherever found	E	\$7,375	\$32,000	\$39,375	\$0	\$39,375
	442	No Common Name (<i>Tetramolopium filiforme</i>) - Wherever found	E	\$7,562	\$128,415	\$135,977	\$0	\$135,977
	884	No Common Name (<i>Tetramolopium lepidotum</i> ssp. <i>lepidotum</i>) - Wherever found	E	\$8,486	\$11,750	\$20,236	\$0	\$20,236
	1190	No Common Name (<i>Tetramolopium remyi</i>) - Wherever found	E	\$9,055	\$0	\$9,055	\$0	\$9,055
	938	No Common Name (<i>Tetramolopium rockii</i>) - Wherever found	T	\$12,309	\$5,000	\$17,309	\$0	\$17,309
	1026	No Common Name (<i>Tinospora homosepala</i>) - Wherever found	E	\$12,847	\$0	\$12,847	\$0	\$12,847
	1283	No Common Name (<i>Trematolobelia singularis</i>) - Wherever found	E	\$6,735	\$0	\$6,735	\$0	\$6,735
	330	No Common Name (<i>Tuberolabium guamense</i>) - Wherever found	T	\$153,535	\$53,315	\$206,850	\$0	\$206,850
	1289	No Common Name (<i>Varronia rupicola</i>) - Wherever found	T	\$6,500	\$0	\$6,500	\$0	\$6,500
	1057	No Common Name (<i>Vernonia proctorii</i>) - Wherever found	E	\$12,000	\$0	\$12,000	\$0	\$12,000
	855	No Common Name (<i>Vigna o-wahuensis</i>) - Wherever found	E	\$22,163	\$0	\$22,163	\$0	\$22,163
	1088	No Common Name (<i>Viola helenae</i>) - Wherever found	E	\$10,933	\$0	\$10,933	\$0	\$10,933
	1197	No Common Name (<i>Viola lanaiensis</i>) - Wherever found	E	\$8,956	\$0	\$8,956	\$0	\$8,956
	629	No Common Name (<i>Viola oahuensis</i>) - Wherever found	E	\$53,847	\$0	\$53,847	\$0	\$53,847
	1313	No Common Name (<i>Wikstroemia skottsbergiana</i>) -	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	806	No Common Name (<i>Wikstroemia villosa</i>) - Wherever found	E	\$5,914	\$20,700	\$26,614	\$0	\$26,614
	997	No Common Name (<i>Xylosma crenatum</i>) - Wherever found	E	\$13,975	\$0	\$13,975	\$0	\$13,975
	802	Nohoanu (<i>Geranium hanaense</i>) - Wherever found	E	\$6,198	\$20,700	\$26,898	\$0	\$26,898
	1246	Nohoanu (<i>Geranium hillebrandii</i>) - Wherever found	E	\$7,801	\$0	\$7,801	\$0	\$7,801
	857	Nohoanu (<i>Geranium kauaiense</i>) - Wherever found	E	\$22,154	\$0	\$22,154	\$0	\$22,154
	792	Nohoanu (<i>Geranium multiflorum</i>) - Wherever found	E	\$7,414	\$20,700	\$28,114	\$0	\$28,114
	770	Nui, haha (<i>Cyanea horrida</i>) - Wherever found	E	\$9,945	\$20,700	\$30,645	\$0	\$30,645
	1418	Oak, Hinckley (<i>Quercus hinckleyi</i>) - Wherever found	T	\$500	\$0	\$500	\$0	\$500
	419	Oha (<i>Delissea subcordata</i>) - Wherever found	E	\$12,321	\$128,415	\$140,736	\$0	\$140,736
	787	Ohai (<i>Sesbania tomentosa</i>) - Wherever found	E	\$21,981	\$6,552	\$28,533	\$0	\$28,533
	1296	`Oha wai (<i>Clermontia drepanomorpha</i>) - Wherever found	E	\$6,379	\$0	\$6,379	\$0	\$6,379
	867	`Oha wai (<i>Clermontia lindseyana</i>) - Wherever found	E	\$18,739	\$2,500	\$21,239	\$0	\$21,239
	1156	`Oha wai (<i>Clermontia oblongifolia</i> ssp. <i>brevipes</i>) - Wherever found	E	\$9,741	\$0	\$9,741	\$0	\$9,741
	1241	`Oha wai (<i>Clermontia oblongifolia</i> ssp. <i>mauiensis</i>) - Wherever found	E	\$7,902	\$0	\$7,902	\$0	\$7,902
	834	`Oha wai (<i>Clermontia peleana</i>) - Wherever found	E	\$19,821	\$4,000	\$23,821	\$0	\$23,821
	885	`Oha wai (<i>Clermontia pyrularia</i>) - Wherever found	E	\$20,110	\$0	\$20,110	\$0	\$20,110
	560	`Oha wai (<i>Clermontia samuelii</i>) - Wherever found	E	\$8,436	\$62,000	\$70,436	\$0	\$70,436
	539	`Ohe (<i>Joinvillea ascendens ascendens</i>) - Wherever found	E	\$55,830	\$20,700	\$76,530	\$0	\$76,530
	609	`Ohe`ohe (<i>Polyscias gymnocarpa</i>) - Wherever found	E	\$57,600	\$0	\$57,600	\$0	\$57,600
	1094	Olulu (<i>Brighamia insignis</i>) - Wherever found	E	\$10,862	\$0	\$10,862	\$0	\$10,862
	1402	Onion, Munz's (<i>Allium munzii</i>) - Wherever found	E	\$1,133	\$0	\$1,133	\$0	\$1,133
	1150	Opuhe (<i>Urera kaalae</i>) - Wherever found	E	\$9,852	\$0	\$9,852	\$0	\$9,852

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	460	Orchid, eastern prairie fringed (<i>Platanthera leucophaea</i>) - Wherever found	T	\$110,202	\$16,043	\$126,245	\$3,100	\$129,345
	523	Orchid, western prairie fringed (<i>Platanthera praeclara</i>) - Wherever found	T	\$39,572	\$40,819	\$80,391	\$1,800	\$82,191
	645	Orchid, white fringeless (<i>Platanthera integrilabia</i>) -	T	\$15,000	\$32,400	\$47,400	\$4,000	\$51,400
	685	Orcutt grass, California (<i>Orcuttia californica</i>) - Wherever found	E	\$6,809	\$37,454	\$44,263	\$0	\$44,263
	1019	Orcutt grass, hairy (<i>Orcuttia pilosa</i>) - Wherever found	E	\$10,670	\$2,373	\$13,043	\$0	\$13,043
	917	Orcutt grass, Sacramento (<i>Orcuttia viscida</i>) - Wherever found	E	\$15,150	\$3,200	\$18,350	\$0	\$18,350
	914	Orcutt grass, San Joaquin (<i>Orcuttia inaequalis</i>) - Wherever found	T	\$15,400	\$3,018	\$18,418	\$0	\$18,418
	874	Orcutt grass, slender (<i>Orcuttia tenuis</i>) - Wherever found	T	\$12,210	\$8,400	\$20,610	\$0	\$20,610
	838	Owl's-clover, fleshy (<i>Castilleja campestris</i> ssp. <i>succulenta</i>) - Wherever found	T	\$19,030	\$4,473	\$23,503	\$0	\$23,503
	1334	Oxytheca, Cushenbury (<i>Oxytheca parishii</i> var. <i>goodmaniana</i>) - Wherever found	E	\$250	\$4,500	\$4,750	\$0	\$4,750
	785	Pa'iniu (<i>Astelia waialealae</i>) - Wherever found	E	\$28,689	\$0	\$28,689	\$0	\$28,689
	225	Paintbrush, golden (<i>Castilleja levisecta</i>) - Wherever found	T	\$320,918	\$174,774	\$495,692	\$0	\$495,692
	377	Paintbrush, San Clemente Island (<i>Castilleja grisea</i>) - Wherever found	T	\$3,319	\$163,511	\$166,830	\$0	\$166,830
	1032	Paintbrush, soft-leaved (<i>Castilleja mollis</i>) - Wherever found	E	\$2,500	\$10,159	\$12,659	\$0	\$12,659
	993	Paintbrush, Tiburon (<i>Castilleja affinis</i> ssp. <i>neglecta</i>) - Wherever found	E	\$10,670	\$3,400	\$14,070	\$0	\$14,070
	1359	Palo colorado (<i>Ternstroemia luquillensis</i>) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1359	Palo de jazmin (<i>Styrax portoricensis</i>) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1141	Palo de nigua (<i>Cornutia obovata</i>) - Wherever found	E	\$10,000	\$0	\$10,000	\$0	\$10,000
	1289	Palo de ramon (<i>Banara vanderbiltii</i>) - Wherever found	E	\$6,500	\$0	\$6,500	\$0	\$6,500
	1020	Palo de rosa (<i>Ottoschulzia rhodoxylon</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1154	Pamakani (<i>Tetramolopium capillare</i>) - Wherever found	E	\$9,770	\$0	\$9,770	\$0	\$9,770
	408	Pamakani (<i>Viola chamissoniana</i> ssp. <i>chamissoniana</i>) - Wherever found	E	\$6,910	\$140,165	\$147,075	\$0	\$147,075
	1178	Panicgrass, Carter's (<i>Panicum fauriei</i> var. <i>carteri</i>) - Wherever found	E	\$8,316	\$1,000	\$9,316	\$0	\$9,316
	1314	Papala (<i>Charpentiera densiflora</i>) - Wherever found	E	\$5,853	\$0	\$5,853	\$0	\$5,853
	1258	Paudedo (<i>Hedyotis megalantha</i>) - Wherever found	E	\$7,432	\$0	\$7,432	\$0	\$7,432
	1327	Pawpaw, beautiful (<i>Deeringothamnus pulchellus</i>) - Wherever found	E	\$0	\$100	\$100	\$5,000	\$5,100
	1147	Pawpaw, four-petal (<i>Asimina tetramera</i>) - Wherever found	E	\$0	\$9,000	\$9,000	\$900	\$9,900
	1311	Pawpaw, Rugel's (<i>Deeringothamnus rugelii</i>) - Wherever found	E	\$4,500	\$0	\$4,500	\$1,500	\$6,000
	1340	Pea, Big Pine partridge (<i>Chamaecrista lineata keyensis</i>) - Wherever found	E	\$3,500	\$400	\$3,900	\$700	\$4,600
	1020	Pelos del diablo (<i>Aristida portoricensis</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	794	Penny-cress, Kneeland Prairie (<i>Thlaspi californicum</i>) - Wherever found	E	\$27,500	\$200	\$27,700	\$0	\$27,700
	526	Pennyroyal, Todsens's (<i>Hedeoma todsenii</i>) - Wherever found	E	\$3,000	\$78,100	\$81,100	\$0	\$81,100
	392	Penstemon, blowout (<i>Penstemon haydenii</i>) - Wherever found	E	\$49,516	\$93,934	\$143,450	\$10,100	\$153,550
	768	Pentachaeta, Lyon's (<i>Pentachaeta lyonii</i>) - Wherever found	E	\$30,692	\$100	\$30,792	\$0	\$30,792
	1391	Pentachaeta, white-rayed (<i>Pentachaeta bellidiflora</i>) - Wherever found	E	\$0	\$1,700	\$1,700	\$0	\$1,700
	1367	Peperomia, Wheeler's (<i>Peperomia wheeleri</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	154	Peppergrass, Slickspot (<i>Lepidium papilliferum</i>) -	T	\$332,116	\$473,975	\$806,091	\$45,040	\$851,131
	1040	Phacelia, clay (<i>Phacelia argillacea</i>) - Wherever found	E	\$12,500	\$0	\$12,500	\$0	\$12,500
	1119	Phacelia, DeBeque (<i>Phacelia submutica</i>) - Wherever found	T	\$2,201	\$7,100	\$9,301	\$1,108	\$10,409
	987	Phacelia, island (<i>Phacelia insularis</i> ssp. <i>insularis</i>) - Wherever found	E	\$2,500	\$11,985	\$14,485	\$0	\$14,485
	937	Phacelia, North Park (<i>Phacelia formosula</i>) - Wherever found	E	\$1,541	\$15,806	\$17,347	\$0	\$17,347

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	959	Phlox, Texas trailing (<i>Phlox nivalis</i> ssp. <i>texensis</i>) - Wherever found	E	\$11,000	\$5,000	\$16,000	\$0	\$16,000
	224	Phlox, Yreka (<i>Phlox hirsuta</i>) - Wherever found	E	\$506,550	\$0	\$506,550	\$0	\$506,550
	334	Pigeon wings (<i>Clitoria fragrans</i>) - Wherever found	T	\$0	\$198,008	\$198,008	\$6,000	\$204,008
	1258	Pilo (<i>Kadua laxiflora</i>) - Wherever found	E	\$7,432	\$0	\$7,432	\$0	\$7,432
	1268	Pilo kea lau li'i (<i>Platydesma rostrata</i>) - Wherever found	E	\$7,309	\$0	\$7,309	\$0	\$7,309
	750	Pinkroot, gentian (<i>Spigelia gentianoides</i>) - Wherever found	E	\$3,500	\$22,270	\$25,770	\$7,500	\$33,270
	338	Pink, swamp (<i>Helonias bullata</i>) - Wherever found	T	\$153,030	\$41,395	\$194,425	\$6,000	\$200,425
	432	Piperia, Yadon's (<i>Piperia yadonii</i>) - Wherever found	E	\$32,841	\$104,535	\$137,376	\$0	\$137,376
	1418	Pitaya, Davis' green (<i>Echinocereus viridiflorus</i> var. <i>davisii</i>) - Wherever found	E	\$500	\$0	\$500	\$0	\$500
	1349	Pitcher-plant, Alabama canebrake (<i>Sarracenia rubra</i> ssp. <i>alabamensis</i>) - Wherever found	E	\$4,000	\$0	\$4,000	\$0	\$4,000
	1198	Pitcher-plant, green (<i>Sarracenia oreophila</i>) - Wherever found	E	\$4,500	\$4,420	\$8,920	\$0	\$8,920
	782	Pitcher-plant, mountain sweet (<i>Sarracenia rubra</i> ssp. <i>jonesii</i>) - Wherever found	E	\$20,000	\$9,254	\$29,254	\$0	\$29,254
	752	Plum, scrub (<i>Prunus geniculata</i>) - Wherever found	E	\$6,500	\$21,700	\$28,200	\$5,000	\$33,200
	683	Po'e (<i>Portulaca sclerocarpa</i>) - Wherever found	E	\$8,931	\$36,000	\$44,931	\$0	\$44,931
	333	Pogonia, small whorled (<i>Isotria medeoloides</i>) - Wherever found	T	\$175,375	\$22,928	\$198,303	\$6,200	\$204,503
	934	Polygala, Lewton's (<i>Polygala lewtonii</i>) - Wherever found	E	\$0	\$0	\$0	\$17,500	\$17,500
	1323	Polygala, tiny (<i>Polygala smallii</i>) - Wherever found	E	\$5,000	\$0	\$5,000	\$200	\$5,200
	551	Polygonum, Scotts Valley (<i>Polygonum hickmanii</i>) - Wherever found	E	\$64,085	\$9,600	\$73,685	\$0	\$73,685
	563	Pondberry (<i>Lindera melissifolia</i>) - Wherever found	E	\$30,500	\$34,102	\$64,602	\$5,300	\$69,902
	1418	Pondweed, Little Aguja (=Creek) (<i>Potamogeton clystocarpus</i>) - Wherever found	E	\$500	\$0	\$500	\$0	\$500
	477	Popcornflower, rough (<i>Plagiobothrys hirtus</i>) - Wherever found	E	\$119,021	\$0	\$119,021	\$0	\$119,021
	1158	Popolo (<i>Cyanea solanacea</i>) - Wherever found	E	\$9,640	\$0	\$9,640	\$0	\$9,640
	1195	Popolo (<i>Solanum nelsonii</i>) - Wherever found	E	\$3,985	\$5,000	\$8,985	\$0	\$8,985
	697	Popolo ku mai (<i>Solanum incompletum</i>) - Wherever found	E	\$10,076	\$32,000	\$42,076	\$0	\$42,076
	1416	Poppy-mallow, Texas (<i>Callirhoe scabruscula</i>) - Wherever found	E	\$500	\$100	\$600	\$0	\$600
	805	Poppy, Sacramento prickly (<i>Argemone pleiacantha</i> ssp. <i>pinnatisecta</i>) - Wherever found	E	\$1,000	\$25,620	\$26,620	\$0	\$26,620
	516	Potato-bean, Price's (<i>Apios priceana</i>) -	T	\$32,500	\$53,326	\$85,826	\$0	\$85,826
	472	Potentilla, Hickman's (<i>Potentilla hickmanii</i>) - Wherever found	E	\$6,507	\$115,000	\$121,507	\$0	\$121,507
	913	Prairie-clover, leafy (<i>Dalea foliosa</i>) - Wherever found	E	\$16,000	\$2,434	\$18,434	\$0	\$18,434
	1426	Prickly-apple, aboriginal (<i>Harrisia</i> (=Cereus) aboriginum (=gracilis)) - Wherever found	E	\$0	\$0	\$0	\$200	\$200
	1391	Prickly-apple, fragrant (<i>Cereus eriophorus</i> var. <i>fragrans</i>) - Wherever found	E	\$0	\$1,000	\$1,000	\$700	\$1,700
	1328	Prickly-ash, St. Thomas (<i>Zanthoxylum thomasianum</i>) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	1363	Primrose, Maguire (<i>Primula maguirei</i>) - Wherever found	T	\$500	\$2,250	\$2,750	\$0	\$2,750
	1127	Pua`ala (<i>Brighamia rockii</i>) - Wherever found	E	\$9,792	\$500	\$10,292	\$0	\$10,292
	1192	Pussypaws, Mariposa (<i>Calyptidium pulchellum</i>) - Wherever found	T	\$9,020	\$0	\$9,020	\$0	\$9,020
	825	Pu`uka`a (<i>Cyperus trachysanthos</i>) - Wherever found	E	\$6,224	\$18,350	\$24,574	\$0	\$24,574
	1379	Ragwort, San Francisco Peaks (<i>Packera franciscana</i>) - Wherever found	T	\$1,500	\$200	\$1,700	\$500	\$2,200
	728	Rattleweed, hairy (<i>Baptisia arachnifera</i>) - Wherever found	E	\$16,500	\$0	\$16,500	\$20,000	\$36,500

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1314	Reedgrass, Hillebrand's (<i>Calamagrostis hillebrandii</i>) - Wherever found	E	\$5,853	\$0	\$5,853	\$0	\$5,853
	807	Reedgrass, Maui (<i>Calamagrostis expansa</i>) - Wherever found	E	\$5,853	\$20,700	\$26,553	\$0	\$26,553
	1320	Reed-mustard, Barneby (<i>Schoenocrambe barnebyi</i>) - Wherever found	E	\$2,000	\$3,450	\$5,450	\$0	\$5,450
	951	Reed-mustard, clay (<i>Schoenocrambe argillacea</i>) - Wherever found	T	\$2,000	\$14,292	\$16,292	\$0	\$16,292
	996	Reed-mustard, shrubby (<i>Schoenocrambe suffrutescens</i>) - Wherever found	E	\$2,000	\$11,993	\$13,993	\$0	\$13,993
	1279	Remya, Maui (<i>Remya mauiensis</i>) - Wherever found	E	\$6,949	\$0	\$6,949	\$0	\$6,949
	849	Rhododendron, Chapman (<i>Rhododendron chapmanii</i>) - Wherever found	E	\$0	\$22,200	\$22,200	\$400	\$22,600
	1368	Ridge-cress, Barneby (<i>Lepidium barnebyanum</i>) - Wherever found	E	\$0	\$2,450	\$2,450	\$0	\$2,450
	879	Rock-cress, Braun's (<i>Arabis perstellata</i>) - Wherever found	E	\$17,000	\$3,334	\$20,334	\$0	\$20,334
	1046	Rockcress, Georgia (<i>Arabis georgiana</i>) - Wherever found	T	\$4,000	\$6,300	\$10,300	\$2,000	\$12,300
	982	Rock-cress, Hoffmann's (<i>Arabis hoffmannii</i>) - Wherever found	E	\$2,500	\$12,185	\$14,685	\$0	\$14,685
	1277	Rock-cress, McDonald's (<i>Arabis macdonaldiana</i>) - Wherever found	E	\$2,762	\$4,200	\$6,962	\$0	\$6,962
	376	Rockcress, Santa Cruz Island (<i>Sibara filifolia</i>) - Wherever found	E	\$3,649	\$163,314	\$166,963	\$0	\$166,963
	887	Rock cress, Shale barren (<i>Arabis serotina</i>) - Wherever found	E	\$13,213	\$200	\$13,413	\$6,683	\$20,096
	810	Rose-mallow, Neches River (<i>Hibiscus dasycalyx</i>) - Wherever found	T	\$7,600	\$0	\$7,600	\$18,671	\$26,271
	696	Rosemary, Apalachicola (<i>Conradina glabra</i>) - Wherever found	E	\$8,000	\$21,700	\$29,700	\$12,500	\$42,200
	1250	Rosemary, Cumberland (<i>Conradina verticillata</i>) - Wherever found	T	\$5,000	\$2,704	\$7,704	\$0	\$7,704
	995	Rosemary, Etonia (<i>Conradina etonia</i>) - Wherever found	E	\$7,500	\$0	\$7,500	\$6,500	\$14,000
	1311	Rosemary, short-leaved (<i>Conradina brevifolia</i>) - Wherever found	E	\$0	\$0	\$0	\$6,000	\$6,000
	1315	Roseroot, Leedy's (<i>Rhodiola integrifolia</i> ssp. <i>leedyi</i>) - Wherever found	T	\$5,000	\$808	\$5,808	\$0	\$5,808
	742	Rush-pea, slender (<i>Hoffmannseggia tenella</i>) - Wherever found	E	\$34,500	\$0	\$34,500	\$0	\$34,500
	1099	Rush-rose, island (<i>Helianthemum greenei</i>) - Wherever found	T	\$2,720	\$8,059	\$10,779	\$0	\$10,779
	894	Sandalwood, Lanai (= `iliahi) (<i>Santalum haleakalae</i> var. <i>lanaiense</i>) - Wherever found	E	\$17,803	\$2,000	\$19,803	\$0	\$19,803
	818	Sandlance (<i>Polygonella myriophylla</i>) - Wherever found	E	\$0	\$21,700	\$21,700	\$3,500	\$25,200
	1130	Sand-verbena, large-fruited (<i>Abronia macrocarpa</i>) - Wherever found	E	\$10,100	\$100	\$10,200	\$0	\$10,200
	1139	Sandwort, Cumberland (<i>Arenaria cumberlandensis</i>) - Wherever found	E	\$7,500	\$2,506	\$10,006	\$0	\$10,006
	827	Sandwort, Marsh (<i>Arenaria paludicola</i>) - Wherever found	E	\$16,494	\$7,954	\$24,448	\$0	\$24,448
	1149	Schiedea, Diamond Head (<i>Schiedea adamantis</i>) - Wherever found	E	\$9,863	\$0	\$9,863	\$0	\$9,863
	883	Seablite, California (<i>Suaeda californica</i>) -	E	\$16,261	\$4,000	\$20,261	\$0	\$20,261
	214	Seagrass, Johnson's (<i>Halophila johnsonii</i>) - Wherever found	T	\$0	\$547,624	\$547,624	\$200	\$547,824
	1020	Sedge, golden (<i>Carex lutea</i>) - Wherever found	E	\$13,000	\$0	\$13,000	\$0	\$13,000
	1300	Sedge, Navajo (<i>Carex specuicola</i>) - Wherever found	T	\$1,100	\$4,700	\$5,800	\$500	\$6,300
	981	Silverbush, Blodgett's (<i>Argythamnia blodgettii</i>) - Wherever found	T	\$200	\$9,605	\$9,805	\$5,000	\$14,805
	953	Silversword, Mauna Loa (=Ka'u) (<i>Argyroxiphium kauense</i>) - Wherever found	E	\$11,253	\$5,000	\$16,253	\$0	\$16,253
	1257	Skullcap, Florida (<i>Scutellaria floridana</i>) - Wherever found	T	\$6,000	\$1,100	\$7,100	\$400	\$7,500
	652	Skullcap, large-flowered (<i>Scutellaria montana</i>) - Wherever found	T	\$7,500	\$32,141	\$39,641	\$10,000	\$49,641
	610	Skyrocket, Pagosa (<i>Ipomopsis polyantha</i>) - Wherever found	E	\$13,575	\$12,456	\$26,031	\$31,530	\$57,561
	849	Snakeroot (<i>Eryngium cuneifolium</i>) - Wherever found	E	\$0	\$21,700	\$21,700	\$900	\$22,600
	1376	Sneezeweed, Virginia (<i>Helenium virginicum</i>) - Wherever found	T	\$1,800	\$500	\$2,300	\$0	\$2,300
	1160	Snowbells, Texas (<i>Styrax texanus</i>) - Wherever found	E	\$9,500	\$100	\$9,600	\$0	\$9,600
	791	Spineflower, Ben Lomond (<i>Chorizanthe pungens</i> var. <i>hartwegiana</i>) - Wherever found	E	\$18,629	\$9,600	\$28,229	\$0	\$28,229
	1074	Spineflower, Howell's (<i>Chorizanthe howellii</i>) - Wherever found	E	\$11,440	\$0	\$11,440	\$0	\$11,440

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	358	Spineflower, Monterey (Chorizanthe pungens var. pungens) - Wherever found	T	\$78,040	\$100,033	\$178,073	\$0	\$178,073
	372	Spineflower, Orcutt's (Chorizanthe orcuttiana) - Wherever found	E	\$62,232	\$107,398	\$169,630	\$0	\$169,630
	1240	Spineflower, Robust (Chorizanthe robusta var. robusta) - Wherever found	E	\$3,904	\$4,000	\$7,904	\$0	\$7,904
	852	Spineflower, Scotts Valley (Chorizanthe robusta var. hartwegii) - Wherever found	E	\$19,242	\$3,200	\$22,442	\$0	\$22,442
	1345	Spineflower, slender-horned (Dodecema leptoceras) - Wherever found	E	\$1,683	\$2,684	\$4,367	\$0	\$4,367
	1361	Spineflower, Sonoma (Chorizanthe valida) - Wherever found	E	\$1,650	\$1,200	\$2,850	\$0	\$2,850
	639	Spiraea, Virginia (Spiraea virginiana) - Wherever found	T	\$36,767	\$5,131	\$41,898	\$10,325	\$52,223
	1344	Spurge, deltoid (Chamaesyce deltoidea ssp. deltoidea) - Wherever found	E	\$4,000	\$200	\$4,200	\$200	\$4,400
	1005	Spurge, Garber's (Chamaesyce garberi) - Wherever found	T	\$6,000	\$4,090	\$10,090	\$3,500	\$13,590
	1213	Spurge, Hoover's (Chamaesyce hooveri) - Wherever found	T	\$7,370	\$945	\$8,315	\$0	\$8,315
	514	Spurge, telephus (Euphorbia telephioides) - Wherever found	T	\$6,000	\$73,331	\$79,331	\$7,000	\$86,331
	1344	Spurge, wedge (Chamaesyce deltoidea serpyllum) - Wherever found	E	\$3,000	\$700	\$3,700	\$700	\$4,400
	1413	Stickseed, showy (Hackelia venusta) - Wherever found	E	\$452	\$200	\$652	\$0	\$652
	322	Sumac, Michaux's (Rhus michauxii) - Wherever found	E	\$55,500	\$169,838	\$225,338	\$3,000	\$228,338
	1329	Sunburst, Hartweg's golden (Pseudobahia bahiifolia) - Wherever found	E	\$0	\$4,959	\$4,959	\$0	\$4,959
	811	Sunburst, San Joaquin adobe (Pseudobahia peirsonii) - Wherever found	T	\$12,870	\$13,373	\$26,243	\$0	\$26,243
	974	Sunflower, Pecos (=puzzle, =paradox) (Helianthus paradoxus) - Wherever found	T	\$5,500	\$9,500	\$15,000	\$0	\$15,000
	1071	Sunflower, San Mateo woolly (Eriophyllum latilobum) - Wherever found	E	\$9,020	\$2,500	\$11,520	\$0	\$11,520
	667	Sunflower, Schweinitz's (Helianthus schweinitzii) - Wherever found	E	\$35,500	\$12,062	\$47,562	\$0	\$47,562
	1108	Sunflower, whorled (Helianthus verticillatus) -	E	\$5,500	\$100	\$5,600	\$5,000	\$10,600
	976	Sunray, Ash Meadows (Enceliopsis nudicaulis var. corrugata) - Wherever found	T	\$14,856	\$100	\$14,956	\$0	\$14,956
	646	Sunshine, Sonoma (Blennosperma bakeri) - Wherever found	E	\$10,670	\$40,300	\$50,970	\$0	\$50,970
	1427	Taraxacum, California (Taraxacum californicum) - Wherever found	E	\$0	\$100	\$100	\$0	\$100
	633	Tarplant, Gaviota (Deinandra increscens ssp. villosa) - Wherever found	E	\$2,605	\$50,100	\$52,705	\$0	\$52,705
	850	Tarplant, Otay (Deinandra (=Hemizonia) conjugens) - Wherever found	T	\$10,038	\$12,468	\$22,506	\$0	\$22,506
	765	Tarplant, Santa Cruz (Holocarpha macradenia) - Wherever found	T	\$22,720	\$8,900	\$31,620	\$0	\$31,620
	550	Thelypody, Howell's spectacular (Thelypodium howellii ssp. spectabilis) - Wherever found	T	\$73,503	\$200	\$73,703	\$0	\$73,703
	1357	Thistle, Chorro Creek bog (Cirsium fontinale var. obispoense) - Wherever found	E	\$3,188	\$100	\$3,288	\$0	\$3,288
	1186	Thistle, fountain (Cirsium fontinale var. fontinale) - Wherever found	E	\$9,020	\$100	\$9,120	\$0	\$9,120
	403	Thistle, La Graciosa (Cirsium loncholepis) - Wherever found	E	\$81,949	\$67,784	\$149,733	\$0	\$149,733
	462	Thistle, Pitcher's (Cirsium pitcheri) - Wherever found	T	\$27,500	\$100,756	\$128,256	\$500	\$128,756
	1403	Thistle, Sacramento Mountains (Cirsium vinaceum) - Wherever found	T	\$1,000	\$100	\$1,100	\$0	\$1,100
	748	Thistle, Suisun (Cirsium hydrophilum var. hydrophilum) - Wherever found	E	\$27,648	\$5,700	\$33,348	\$0	\$33,348
	963	Thornmint, San Diego (Acanthomintha ilicifolia) - Wherever found	T	\$12,900	\$2,684	\$15,584	\$0	\$15,584

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	1395	Thormint, San Mateo (<i>Acanthomintha obovata</i> ssp. <i>duttonii</i>) - Wherever found	E	\$0	\$1,547	\$1,547	\$0	\$1,547
	1367	Thoroughwort, Cape Sable (<i>Chromolaena frustrata</i>) - Wherever found	E	\$400	\$100	\$500	\$2,000	\$2,500
	1322	Townsendia, Last Chance (<i>Townsendia aprica</i>) - Wherever found	T	\$1,000	\$4,350	\$5,350	\$0	\$5,350
	668	Trillium, persistent (<i>Trillium persistens</i>) - Wherever found	E	\$41,500	\$0	\$41,500	\$6,000	\$47,500
	712	Trillium, relict (<i>Trillium reliquum</i>) - Wherever found	E	\$1,500	\$35,210	\$36,710	\$3,000	\$39,710
	1035	Tuctoria, Greene's (<i>Tuctoria greenei</i>) - Wherever found	E	\$11,880	\$745	\$12,625	\$0	\$12,625
	918	Twinpod, Dudley Bluffs (<i>Physaria obcordata</i>) - Wherever found	T	\$4,256	\$12,956	\$17,212	\$1,108	\$18,320
	573	Ufa-halomtano (<i>Heritiera longipetiolata</i>) - Wherever found	E	\$66,219	\$0	\$66,219	\$0	\$66,219
	709	Uhi uhi (<i>Mezoneuron kavaense</i>) - Wherever found	E	\$40,347	\$0	\$40,347	\$0	\$40,347
	1343	Uvillo (<i>Eugenia haematocarpa</i>) - Wherever found	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	1365	Vervain, Red Hills (<i>Verbena californica</i>) - Wherever found	T	\$1,100	\$1,500	\$2,600	\$0	\$2,600
	1113	Vetch, Hawaiian (<i>Vicia menziesii</i>) - Wherever found	E	\$10,534	\$0	\$10,534	\$0	\$10,534
	1280	Wahane (<i>Pritchardia aylmer-robinsonii</i>) - Wherever found	E	\$6,948	\$0	\$6,948	\$0	\$6,948
	294	Wallflower, Ben Lomond (<i>Erysimum teretifolium</i>) - Wherever found	E	\$270,309	\$9,800	\$280,109	\$0	\$280,109
	455	Wallflower, Contra Costa (<i>Erysimum capitatum</i> var. <i>angustatum</i>) - Wherever found	E	\$132,460	\$200	\$132,660	\$0	\$132,660
	691	Wallflower, Menzies' (<i>Erysimum menziesii</i>) - Wherever found	E	\$6,310	\$36,200	\$42,510	\$0	\$42,510
	1367	Walnut (=Nogal), West Indian (<i>Juglans jamaicensis</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	583	Warea, wide-leaf (<i>Warea amplexifolia</i>) - Wherever found	E	\$12,000	\$200	\$12,200	\$50,500	\$62,700
	488	Watercress, Gambel's (<i>Rorippa gambellii</i>) - Wherever found	E	\$21,955	\$82,336	\$104,291	\$0	\$104,291
	1408	Water-plantain, Kral's (<i>Sagittaria secundifolia</i>) - Wherever found	T	\$400	\$0	\$400	\$500	\$900
	543	Water-umbel, Huachuca (<i>Lilaeopsis schaffneriana</i> var. <i>recurva</i>) - Wherever found	E	\$11,500	\$62,100	\$73,600	\$2,500	\$76,100
	1385	Water-willow, Cooley's (<i>Justicia cooleyi</i>) - Wherever found	E	\$0	\$0	\$0	\$2,000	\$2,000
	1385	Whitlow-wort, papery (<i>Paronychia chartacea</i>) - Wherever found	T	\$0	\$0	\$0	\$2,000	\$2,000
	738	Wild buckwheat, clay-loving (<i>Eriogonum pelinophilum</i>) - Wherever found	E	\$0	\$28,006	\$28,006	\$6,808	\$34,814
	461	Wild-buckwheat, gypsum (<i>Eriogonum gypsophilum</i>) - Wherever found	T	\$32,000	\$97,000	\$129,000	\$0	\$129,000
	1393	Wild-buckwheat, southern mountain (<i>Eriogonum kennedyi</i> var. <i>austromontanum</i>) - Wherever found	T	\$1,650	\$0	\$1,650	\$0	\$1,650
	456	Wild-rice, Texas (<i>Zizania texana</i>) - Wherever found	E	\$130,000	\$600	\$130,600	\$0	\$130,600
	891	Wire-lettuce, Malheur (<i>Stephanomeria malheurensis</i>) - Wherever found	E	\$18,151	\$1,700	\$19,851	\$0	\$19,851
	702	Wireweed (<i>Polygonella basiramia</i>) - Wherever found	E	\$0	\$39,548	\$39,548	\$1,700	\$41,248
	377	Woodland-star, San Clemente Island (<i>Lithophragma maximum</i>) - Wherever found	E	\$3,319	\$163,511	\$166,830	\$0	\$166,830
	863	Woolly-star, Santa Ana River (<i>Eriastrum densifolium</i> ssp. <i>sanctorum</i>) - Wherever found	E	\$11,935	\$9,918	\$21,853	\$0	\$21,853
	375	Wooly-threads, San Joaquin (<i>Monolopia</i> (=Lembertia) <i>congdonii</i>) - Wherever found	E	\$55,810	\$111,360	\$167,170	\$0	\$167,170
	655	Yellowhead, desert (<i>Yermo xanthocephalus</i>) - Wherever found	T	\$5,068	\$44,100	\$49,168	\$0	\$49,168
	267	Yerba santa, Lompoc (<i>Eriodictyon capitatum</i>) - Wherever found	E	\$12,312	\$346,500	\$358,812	\$0	\$358,812
	626	Ziziphus, Florida (<i>Ziziphus celata</i>) - Wherever found	E	\$23,000	\$0	\$23,000	\$31,000	\$54,000
Flowering Plants subtotal				\$16,152,353	\$18,842,257	\$34,994,610	\$808,343	\$35,802,953
Conifers and Cycads	1288	Cypress, Gowen (<i>Cupressus goveniana</i> ssp. <i>goveniana</i>) - Wherever found	T	\$6,507	\$0	\$6,507	\$0	\$6,507

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	930	Cypress, Santa Cruz (<i>Cupressus abramsiana</i>) - Wherever found	T	\$10,410	\$7,200	\$17,610	\$0	\$17,610
	171	Fadang (<i>Cycas micronesica</i>) - Wherever found	T	\$76,379	\$697,550	\$773,929	\$0	\$773,929
	729	Torreya, Florida (<i>Torreya taxifolia</i>) - Wherever found	E	\$6,000	\$21,700	\$27,700	\$8,500	\$36,200
Conifers and Cycads subtotal				\$99,296	\$726,450	\$825,746	\$8,500	\$834,246
Ferns and Allies	1175	Aumakua, Palapalai (<i>Dryopteris crinalis</i> var. <i>podosorus</i>) - Wherever found	E	\$9,364	\$0	\$9,364	\$0	\$9,364
	1258	Diellia, Asplenium-leaved (<i>Asplenium dielirectum</i>) - Wherever found	E	\$7,432	\$0	\$7,432	\$0	\$7,432
	763	Fern, Aleutian shield (<i>Polystichum aleuticum</i>) - Wherever found	E	\$22,000	\$10,000	\$32,000	\$0	\$32,000
	820	Fern, American hart's-tongue (<i>Asplenium scolopendrium</i> var. <i>americanum</i>) - Wherever found	T	\$22,600	\$2,379	\$24,979	\$0	\$24,979
	1343	Fern, Elfin tree (<i>Cyathea dryopteroides</i>) - Wherever found	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	1364	Fern, Florida bristle (<i>Trichomanes punctatum</i> ssp. <i>floridanum</i>) - Wherever found	E	\$2,000	\$0	\$2,000	\$700	\$2,700
	1028	Fern, pendant kihi (<i>Adenophorus periens</i>) - Wherever found	E	\$12,820	\$0	\$12,820	\$0	\$12,820
	1313	Hohiu (<i>Dryopteris glabra</i> var. <i>pusilla</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	570	Ihi' ihi (<i>Marsilea villosa</i>) - Wherever found	E	\$12,580	\$55,821	\$68,401	\$0	\$68,401
	1313	Kupukupu makalii (<i>Cyclosorus boydiae</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1359	No Common Name (<i>Adiantum vivesii</i>) - Wherever found	E	\$3,000	\$0	\$3,000	\$0	\$3,000
	1217	No Common Name (<i>Asplenium dielfalcatum</i>) - Wherever found	E	\$8,256	\$0	\$8,256	\$0	\$8,256
	1313	No Common Name (<i>Asplenium diellaciniatum</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	968	No Common Name (<i>Asplenium dielmannii</i>) - Wherever found	E	\$15,396	\$0	\$15,396	\$0	\$15,396
	968	No Common Name (<i>Asplenium dielpallidum</i>) - Wherever found	E	\$15,396	\$0	\$15,396	\$0	\$15,396
	591	No Common Name (<i>Asplenium peruvianum</i> var. <i>insulare</i>) - Wherever found	E	\$8,542	\$52,700	\$61,242	\$0	\$61,242
	870	No Common Name (<i>Asplenium unisorum</i>) - Wherever found	E	\$6,348	\$14,604	\$20,952	\$0	\$20,952
	1313	No Common Name (<i>Deparia kaalaana</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1086	No Common Name (<i>Diplazium molokaiense</i>) - Wherever found	E	\$10,993	\$0	\$10,993	\$0	\$10,993
	1168	No Common Name (<i>Doryopteris angelica</i>) - Wherever found	E	\$9,457	\$0	\$9,457	\$0	\$9,457
	1301	No Common Name (<i>Doryopteris takeuchii</i>) - Wherever found	E	\$6,212	\$0	\$6,212	\$0	\$6,212
	1276	No Common Name (<i>Elaphoglossum serpens</i>) - Wherever found	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	1255	No Common Name (<i>Huperzia stemmermanniae</i>) - Wherever found	E	\$7,520	\$0	\$7,520	\$0	\$7,520
	806	No Common Name (<i>Microlepia strigosa</i> var. <i>mauiensis</i>) - Wherever found	E	\$5,914	\$20,700	\$26,614	\$0	\$26,614
	1328	No Common Name (<i>Polystichum calderonense</i>) - Wherever found	E	\$5,000	\$0	\$5,000	\$0	\$5,000
	585	No Common Name (<i>Pteris lidgatei</i>) - Wherever found	E	\$62,446	\$0	\$62,446	\$0	\$62,446
	1276	No Common Name (<i>Tectaria estremerana</i>) - Wherever found	E	\$7,000	\$0	\$7,000	\$0	\$7,000
	1343	No Common Name (<i>Thelypteris inabonensis</i>) - Wherever found	E	\$4,500	\$0	\$4,500	\$0	\$4,500
	1354	No Common Name (<i>Thelypteris verecunda</i>) - Wherever found	E	\$3,500	\$0	\$3,500	\$0	\$3,500
	1367	No Common Name (<i>Thelypteris yaucoensis</i>) - Wherever found	E	\$2,500	\$0	\$2,500	\$0	\$2,500
	1313	Olua (<i>Hypolepis hawaiiensis</i> var. <i>mauiensis</i>) - Wherever found	E	\$5,914	\$0	\$5,914	\$0	\$5,914
	1107	Pauoa (<i>Ctenitis squamigera</i>) - Wherever found	E	\$10,618	\$0	\$10,618	\$0	\$10,618
	603	Quillwort, black spored (<i>Isoetes melanospora</i>) - Wherever found	E	\$54,000	\$2,200	\$56,200	\$2,000	\$58,200
	464	Quillwort, Louisiana (<i>Isoetes louisianensis</i>) - Wherever found	E	\$1,500	\$127,082	\$128,582	\$0	\$128,582
	995	Quillwort, mat-forming (<i>Isoetes tegetiformans</i>) - Wherever found	E	\$11,000	\$0	\$11,000	\$3,000	\$14,000
	784	Wawae'iole (<i>Huperzia mannii</i>) - Wherever found	E	\$8,120	\$20,700	\$28,820	\$0	\$28,820
	590	Wawae'iole (<i>Huperzia nutans</i>) - Wherever found	E	\$61,363	\$0	\$61,363	\$0	\$61,363

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
Ferns and Allies subtotal				\$458,447	\$306,186	\$764,633	\$5,700	\$770,333
Lichens	518	Cladonia, Florida perforate (Cladonia perforata) - Wherever found	E	\$2,000	\$81,800	\$83,800	\$1,900	\$85,700
	1013	Lichen, rock gnome (Gymnoderma lineare) - Wherever found	E	\$12,200	\$1,100	\$13,300	\$0	\$13,300
Lichens subtotal				\$14,200	\$82,900	\$97,100	\$1,900	\$99,000
Multi-species				\$3,625,640	\$77,472,044	\$81,097,684	\$8,704,696	\$89,802,380
Multi-species subtotal				\$3,625,640	\$77,472,044	\$81,097,684	\$8,704,696	\$89,802,380
Total Species Expenditures				\$188,504,080	\$1,047,794,289	\$1,236,298,369	\$53,074,754	\$1,289,373,123

**TABLE 2. SPECIES RANKED IN DESCENDING ORDER OF TOTAL FY 2017
REPORTED EXPENDITURES**

Species are arranged in descending rank order of total identifiable expenditures (Table 1). Cumulative totals of the expenditures totals are provided. The cumulative total column is the sum of all higher ranked species totals (from Table 1) and the selected species entry. This table does not include costs for land acquisition or multi-species conservation activities.

The species list in the report follows the same breakdown as in 50 CFR 17.11 and 17.12. All populations are itemized separately, not just Evolutionarily Significant Units (ESUs). Thus, it appears as if there are more listed species. Separate populations of the same species may be ranked differently. The STATUS column indicates whether the species is endangered (E), threatened (T), experimental (EXPN), or threatened by similarity of appearance (SAT).

TABLE 2. SPECIES RANKED IN DESCENDING ORDER OF TOTAL FY 2017 REPORTED EXPENDITURES, NOT INCLUDING LAND ACQUISITION COSTS

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Snake River spring/summer-run ESU	T	\$119,227,105	\$119,227,105
2	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Snake River Basin DPS	T	\$60,150,691	\$179,377,796
3	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Middle Columbia River DPS	T	\$55,223,643	\$234,601,439
4	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Upper Willamette River ESU	T	\$49,035,365	\$283,636,804
5	Sturgeon, pallid (<i>Scaphirhynchus albus</i>) - Wherever found	E	\$34,201,142	\$317,837,946
6	Trout, bull (<i>Salvelinus confluentus</i>) - U.S.A., conterminous, lower 48 states	T	\$33,635,463	\$351,473,409
7	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Snake River fall-run ESU	T	\$32,836,435	\$384,309,844
8	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Upper Columbia spring-run ESU	E	\$31,795,241	\$416,105,085
9	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Upper Columbia River DPS	T	\$27,124,048	\$443,229,133
10	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Lower Columbia River ESU	T	\$22,544,860	\$465,773,993
11	Salmon, sockeye (<i>Oncorhynchus</i> (=Salmo) nerka) - Snake River ESU	E	\$21,482,839	\$487,256,832
12	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) - Southern Oregon - Northern California Coast ESU	T	\$18,255,071	\$505,511,903
13	Tortoise, desert (<i>Gopherus agassizii</i>) - Wherever found, except AZ south and east of Colorado R., and Mexico	T	\$15,974,380	\$521,486,283
14	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Puget Sound ESU	T	\$14,558,294	\$536,044,577
15	Sucker, razorback (<i>Xyrauchen texanus</i>) - Wherever found	E	\$14,554,269	\$550,598,846
16	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Lower Columbia River DPS	T	\$14,173,657	\$564,772,503
17	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - California Central Valley DPS	T	\$14,162,642	\$578,935,145
18	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Sacramento River winter-run ESU	E	\$14,006,445	\$592,941,590
19	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Upper Willamette River DPS	T	\$13,362,029	\$606,303,619
20	Chub, bonytail (<i>Gila elegans</i>) - Wherever found	E	\$13,161,076	\$619,464,695

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
21	Woodpecker, red-cockaded (Picoides borealis) - Wherever found	E	\$12,767,183	\$632,231,878
22	Sturgeon, white (Acipenser transmontanus) - U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	E	\$11,960,315	\$644,192,193
23	Owl, northern spotted (Strix occidentalis caurina) - Wherever found	T	\$11,854,675	\$656,046,868
24	Salmon, coho (Oncorhynchus (=Salmo) kisutch) - Lower Columbia River ESU	T	\$11,396,148	\$667,443,016
25	Manatee, West Indian (Trichechus manatus) - Wherever found	E	\$10,145,358	\$677,588,374
26	Salmon, Atlantic (Salmo salar) - Gulf of Maine DPS	E	\$9,907,908	\$687,496,282
27	Steelhead (Oncorhynchus (=Salmo) mykiss) - Puget Sound DPS	T	\$9,615,275	\$697,111,557
28	Plover, piping (Charadrius melodus) - [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	T	\$9,525,440	\$706,636,997
29	Whale, North Atlantic Right (Eubalaena glacialis) - Wherever found	E	\$8,943,197	\$715,580,194
30	Flycatcher, southwestern willow (Empidonax traillii extimus) - Wherever found	E	\$8,938,964	\$724,519,158
31	Chub, humpback (Gila cypha) - Wherever found	E	\$8,794,185	\$733,313,343
32	Salmon, chum (Oncorhynchus keta) - Columbia River ESU	T	\$8,644,825	\$741,958,168
33	Minnnow, Rio Grande Silvery (Hybognathus amarus) - Wherever found, except where listed as an experimental population	E	\$8,369,633	\$750,327,801
34	Smelt, delta (Hypomesus transpacificus) - Wherever found	T	\$8,352,876	\$758,680,677
35	Trout, Lahontan cutthroat (Oncorhynchus clarkii henshawi) - Wherever found	T	\$7,914,244	\$766,594,921
36	Pikeminnow (=squawfish), Colorado (Ptychocheilus lucius) - Wherever found, except where listed as an experimental population	E	\$7,882,011	\$774,476,932
37	Salmon, Chinook (Oncorhynchus (=Salmo) tshawytscha) - Central Valley spring-run ESU	T	\$7,526,916	\$782,003,848
38	Salmon, coho (Oncorhynchus (=Salmo) kisutch) - Oregon Coast ESU	T	\$7,353,863	\$789,357,711
39	Bat, Northern long-eared (Myotis septentrionalis) - Wherever found	T	\$6,965,670	\$796,323,381
40	Bear, grizzly (Ursus arctos horribilis) - U.S.A., conterminous (lower 48) States, except where listed as an experimental population	T	\$6,915,297	\$803,238,678
41	Steelhead (Oncorhynchus (=Salmo) mykiss) - Central California Coast DPS	T	\$6,270,991	\$809,509,669
42	Bat, Indiana (Myotis sodalis) - Wherever found	E	\$5,782,851	\$815,292,520
43	Tern, least (Sterna antillarum) - interior pop.	E	\$5,767,371	\$821,059,891
44	Cuckoo, yellow-billed (Coccyzus americanus) - Western U.S. DPS	T	\$5,381,587	\$826,441,478
45	Sea turtle, loggerhead (Caretta caretta) - Northwest Atlantic Ocean DPS	T	\$5,379,067	\$831,820,545
46	Salmon, coho (Oncorhynchus (=Salmo) kisutch) - Central California Coast ESU	E	\$5,327,795	\$837,148,340

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
47	Murrelet, marbled (<i>Brachyramphus marmoratus</i>) - U.S.A. (CA, OR, WA)	T	\$5,210,104	\$842,358,444
48	Salmon, chum (<i>Oncorhynchus keta</i>) - Hood Canal summer-run ESU	T	\$4,882,163	\$847,240,607
49	Wolf, gray (<i>Canis lupus</i>) - U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	E	\$4,857,155	\$852,097,762
50	Salmon, sockeye (<i>Oncorhynchus (=Salmo) nerka</i>) - Ozette Lake ESU	T	\$4,690,123	\$856,787,885
51	Elephant, African (<i>Loxodonta africana</i>) - Wherever found	T	\$4,674,200	\$861,462,085
52	Plover, western snowy (<i>Charadrius nivosus nivosus</i>) - Pacific Coast population DPS- U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	T	\$4,619,428	\$866,081,513
53	Sea turtle, hawksbill (<i>Eretmochelys imbricata</i>) - Wherever found	E	\$4,451,705	\$870,533,218
54	Sea turtle, leatherback (<i>Dermochelys coriacea</i>) - Wherever found	E	\$4,338,209	\$874,871,427
55	Steelhead (<i>Oncorhynchus (=Salmo) mykiss</i>) - Northern California DPS	T	\$4,226,406	\$879,097,833
56	Seal, Hawaiian monk (<i>Monachus schauinslandi</i>) - Wherever found	E	\$4,182,904	\$883,280,737
57	Sea turtle, Kemp's ridley (<i>Lepidochelys kempii</i>) - Wherever found	E	\$4,100,538	\$887,381,275
58	Steelhead (<i>Oncorhynchus (=Salmo) mykiss</i>) - Southern California DPS	E	\$4,100,474	\$891,481,749
59	Salmon, Chinook (<i>Oncorhynchus (=Salmo) tshawytscha</i>) - California Coastal ESU	T	\$4,017,394	\$895,499,143
60	Bear, polar (<i>Ursus maritimus</i>) - Wherever found	T	\$3,999,916	\$899,499,059
61	Steelhead (<i>Oncorhynchus (=Salmo) mykiss</i>) - South-Central California Coast DPS	T	\$3,980,889	\$903,479,948
62	Condor, California (<i>Gymnogyps californianus</i>) - U.S.A. only, except where listed as an experimental population	E	\$3,849,385	\$907,329,333
63	Whale, beluga (<i>Delphinapterus leucas</i>) - Cook Inlet DPS	E	\$3,603,352	\$910,932,685
64	Sage-grouse, Gunnison (<i>Centrocercus minimus</i>) - Wherever found	T	\$3,582,626	\$914,515,311
65	Sea lion, Steller (<i>Eumetopias jubatus</i>) - Western DPS	E	\$3,570,842	\$918,086,153
66	Ferret, black-footed (<i>Mustela nigripes</i>) - Wherever found, except where listed as an experimental population	E	\$3,252,028	\$921,338,181
67	Frog, California red-legged (<i>Rana draytonii</i>) - Wherever found	T	\$3,124,310	\$924,462,491
68	Panther, Florida (<i>Puma (=Felis) concolor coryi</i>) - Wherever found	E	\$3,001,286	\$927,463,777
69	Wolf, Mexican (<i>Canis lupus baileyi</i>) - U.S.A. (portions of AZ and NM)see 17.84(k)	EXPN	\$2,975,063	\$930,438,840
70	Sturgeon, green (<i>Acipenser medirostris</i>) - Southern DPS	T	\$2,840,014	\$933,278,854
71	Whale, blue (<i>Balaenoptera musculus</i>) - Wherever found	E	\$2,780,824	\$936,059,678

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
72	Lynx, Canada (<i>Lynx canadensis</i>) - Wherever Found in Contiguous U.S.	T	\$2,772,033	\$938,831,711
73	Tern, California least (<i>Sterna antillarum browni</i>) - Wherever found	E	\$2,718,336	\$941,550,047
74	Prairie dog, Utah (<i>Cynomys parvidens</i>) - Wherever found	T	\$2,576,948	\$944,126,995
75	Whale, finback (<i>Balaenoptera physalus</i>) - Wherever found	E	\$2,541,447	\$946,668,442
76	Shrike, San Clemente loggerhead (<i>Lanius ludovicianus mearnsi</i>) - Wherever found	E	\$2,412,338	\$949,080,780
77	Vireo, least Bell's (<i>Vireo bellii pusillus</i>) - Wherever found	E	\$2,390,380	\$951,471,160
78	Sturgeon (Gulf subspecies), Atlantic (<i>Acipenser oxyrinchus (=oxyrynchus) desotoi</i>) - Wherever found	T	\$2,389,062	\$953,860,222
79	Whale, killer (<i>Orcinus orca</i>) - Southern Resident DPS	E	\$2,337,488	\$956,197,710
80	Snake, eastern indigo (<i>Drymarchon corais couperi</i>) - Wherever found	T	\$2,226,922	\$958,424,632
81	Rail, Yuma clapper (<i>Rallus longirostris yumanensis</i>) - Wherever found	E	\$2,204,198	\$960,628,830
82	Abalone, White (<i>Haliotis sorenseni</i>) - North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	E	\$2,167,091	\$962,795,921
83	Elephant, Asian (<i>Elephas maximus</i>) - Wherever found	E	\$2,109,893	\$964,905,814
84	Bat, Hawaiian hoary (<i>Lasiurus cinereus semotus</i>) - Wherever found	E	\$2,062,844	\$966,968,658
85	Fly, Delhi Sands flower-loving (<i>Rhaphiomidas terminatus abdominalis</i>) - Wherever found	E	\$2,060,500	\$969,029,158
86	Sucker, Lost River (<i>Deltistes luxatus</i>) - Wherever found	E	\$2,043,382	\$971,072,540
87	Goose, Hawaiian (<i>Branta (=Nesochen) sandvicensis</i>) - Wherever found	E	\$2,039,338	\$973,111,878
88	Sucker, shortnose (<i>Chasmistes brevirostris</i>) - Wherever found	E	\$2,035,382	\$975,147,260
89	Sea turtle, green (<i>Chelonia mydas</i>) - North Atlantic DPS	T	\$2,020,936	\$977,168,196
90	Tiger (<i>Panthera tigris</i>) - Wherever found	E	\$2,012,773	\$979,180,969
91	Gnatcatcher, coastal California (<i>Polioptila californica californica</i>) - Wherever found	T	\$1,992,260	\$981,173,229
92	Sturgeon, shortnose (<i>Acipenser brevirostrum</i>) - Wherever found	E	\$1,981,001	\$983,154,230
93	Stork, wood (<i>Mycteria americana</i>) - AL, FL, GA, MS, NC, SC	T	\$1,972,631	\$985,126,861
94	Whale, sperm (<i>Physeter catodon (=macrocephalus)</i>) - Wherever found	E	\$1,965,923	\$987,092,784
95	Sea turtle, loggerhead (<i>Caretta caretta</i>) - North Pacific Ocean DPS	E	\$1,920,348	\$989,013,132
96	Petrel, Hawaiian (<i>Pterodroma sandwichensis</i>) - Wherever found	E	\$1,866,722	\$990,879,854
97	Alligator, American (<i>Alligator mississippiensis</i>) - Wherever found	SAT	\$1,815,500	\$992,695,354
98	Palila (honeycreeper) (<i>Loxioides bailleui</i>) - Wherever found	E	\$1,808,316	\$994,503,670
99	Scrub-jay, Florida (<i>Aphelocoma coerulescens</i>) - Wherever found	T	\$1,807,096	\$996,310,766
100	Salamander, California tiger (<i>Ambystoma californiense</i>) - U.S.A. (Central CA DPS)	T	\$1,790,206	\$998,100,972

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
101	Whale, Sei (<i>Balaenoptera borealis</i>) - Wherever found	E	\$1,752,980	\$999,853,952
102	Sea turtle, green (<i>Chelonia mydas</i>) - Central West Pacific DPS	E	\$1,695,876	\$1,001,549,828
103	Deer, key (<i>Odocoileus virginianus clavium</i>) - Wherever found	E	\$1,673,635	\$1,003,223,463
104	Horned lark, streaked (<i>Eremophila alpestris strigata</i>) - Wherever found	T	\$1,668,841	\$1,004,892,304
105	Eulachon (<i>Thaleichthys pacificus</i>) - Southern DPS	T	\$1,654,204	\$1,006,546,508
106	Sea turtle, green (<i>Chelonia mydas</i>) - East Pacific DPS	T	\$1,582,450	\$1,008,128,958
107	Frog, Oregon spotted (<i>Rana pretiosa</i>) - Wherever found	T	\$1,575,991	\$1,009,704,949
108	Whale, humpback (<i>Megaptera novaeangliae</i>) - Mexico DPS	T	\$1,558,084	\$1,011,263,033
109	Checkerspot, Taylor's (=whulge) (<i>Euphydryas editha taylori</i>) - Wherever found	E	\$1,478,204	\$1,012,741,237
110	Kite, Everglade snail (<i>Rostrhamus sociabilis plumbeus</i>) - Wherever found	E	\$1,477,058	\$1,014,218,295
111	Fairy shrimp, vernal pool (<i>Branchinecta lynchi</i>) - Wherever found	T	\$1,448,791	\$1,015,667,086
112	Parrot, Puerto Rican (<i>Amazona vittata</i>) - Wherever found	E	\$1,445,000	\$1,017,112,086
113	Stilt, Hawaiian (<i>Himantopus mexicanus knudseni</i>) - Wherever found	E	\$1,429,312	\$1,018,541,398
114	Whale, humpback (<i>Megaptera novaeangliae</i>) - Central America DPS	E	\$1,411,338	\$1,019,952,736
115	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) - Chesapeake Bay DPS	E	\$1,379,896	\$1,021,332,632
116	Coot, Hawaiian (<i>Fulica americana alai</i>) - Wherever found	E	\$1,357,763	\$1,022,690,395
117	Whale, bowhead (<i>Balaena mysticetus</i>) - Wherever found	E	\$1,302,156	\$1,023,992,551
118	Crane, whooping (<i>Grus americana</i>) - U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	EXPN	\$1,293,204	\$1,025,285,755
119	Prairie-chicken, Attwater's greater (<i>Tympanuchus cupido attwateri</i>) - Wherever found	E	\$1,289,488	\$1,026,575,243
120	Chimpanzee (<i>Pan troglodytes</i>) - Wherever found	E	\$1,285,751	\$1,027,860,994
121	Crow, Hawaiian (=alala) (<i>Corvus hawaiiensis</i>) - Wherever found	E	\$1,284,686	\$1,029,145,680
122	Fox, San Joaquin kit (<i>Vulpes macrotis mutica</i>) - wherever found	E	\$1,238,168	\$1,030,383,848
123	Pronghorn, Sonoran (<i>Antilocapra americana sonoriensis</i>) - U.S.A. (AZ), Mexico	EXPN	\$1,233,263	\$1,031,617,111
124	Shiner, Topeka (<i>Notropis topeka</i> (=tristis)) - U.S.A. (MO-specified portions of Little Creek, Big Muddy Creek, and Spring Creek watersheds in Adair, Gentry, Harrison, Putnam, Sullivan, and Worth Counties; see 17.84(d)(1)(i))	EXPN	\$1,203,100	\$1,032,820,211
125	Crow, Mariana (=aga) (<i>Corvus kubaryi</i>) - Wherever found	E	\$1,196,900	\$1,034,017,111
126	Otter, Northern Sea (<i>Enhydra lutris kenyoni</i>) - Southwest Alaska DPS	T	\$1,186,982	\$1,035,204,093
127	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) - Gulf of Maine DPS	T	\$1,184,890	\$1,036,388,983

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
128	Sea turtle, green (<i>Chelonia mydas</i>) - Central North Pacific DPS	T	\$1,168,665	\$1,037,557,648
129	Rhinoceros, black (<i>Diceros bicornis</i>) - Africa	E	\$1,143,886	\$1,038,701,534
130	Sparrow, Cape Sable seaside (<i>Ammodramus maritimus mirabilis</i>) - Wherever found	E	\$1,136,840	\$1,039,838,374
131	Gorilla (<i>Gorilla gorilla</i>) - Wherever found	E	\$1,123,531	\$1,040,961,905
132	Crane, whooping (<i>Grus americana</i>) - Wherever found, except where listed as an experimental population	E	\$1,093,999	\$1,042,055,904
133	Bat, gray (<i>Myotis grisescens</i>) - Wherever found	E	\$1,074,693	\$1,043,130,597
134	Snails, Oahu tree (<i>Achatinella</i> spp.) - Wherever found	E	\$1,043,809	\$1,044,174,406
135	Knot, red (<i>Calidris canutus rufa</i>) - Wherever found	T	\$1,022,673	\$1,045,197,079
136	Sea turtle, green (<i>Chelonia mydas</i>) - South Atlantic DPS	T	\$1,000,775	\$1,046,197,854
137	Abalone, Black (<i>Haliotis cracherodii</i>) - Wherever found	E	\$997,866	\$1,047,195,720
138	Sucker, June (<i>Chasmistes liorus</i>) - Wherever found	E	\$991,600	\$1,048,187,320
139	Shiner, Topeka (<i>Notropis topeka</i> (=tristis)) - Wherever found, except where listed as an experimental population	E	\$984,173	\$1,049,171,493
140	Pupfish, Devils Hole (<i>Cyprinodon diabolis</i>) - Wherever found	E	\$983,845	\$1,050,155,338
141	Butterfly, Fender's blue (<i>Icaricia icarioides fenderi</i>) - Wherever found	E	\$981,706	\$1,051,137,044
142	Eider, spectacled (<i>Somateria fischeri</i>) - Wherever found	T	\$979,558	\$1,052,116,602
143	Rail, California clapper (<i>Rallus longirostris obsoletus</i>) - Wherever found	E	\$979,160	\$1,053,095,762
144	Mouse, salt marsh harvest (<i>Reithrodontomys raviventris</i>) - wherever found	E	\$973,061	\$1,054,068,823
145	Beetle, American burying (<i>Nicrophorus americanus</i>) - Wherever found, except where listed as an experimental population	E	\$967,683	\$1,055,036,506
146	Sawfish, smalltooth (<i>Pristis pectinata</i>) - US DPS - Smalltooth sawfish originating from U.S. waters.	E	\$967,245	\$1,056,003,751
147	Whale, humpback (<i>Megaptera novaeangliae</i>) - Western North Pacific	E	\$956,621	\$1,056,960,372
148	Caracara, Audubon's crested (<i>Polyborus plancus audubonii</i>) - FL pop.	T	\$898,245	\$1,057,858,617
149	Snake, giant garter (<i>Thamnophis gigas</i>) - Wherever found	T	\$884,515	\$1,058,743,132
150	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) - Carolina DPS	E	\$879,030	\$1,059,622,162
151	Otter, southern sea (<i>Enhydra lutris nereis</i>) - Wherever found	T	\$870,995	\$1,060,493,157
152	Tadpole shrimp, vernal pool (<i>Lepidurus packardi</i>) - Wherever found	E	\$870,451	\$1,061,363,608
153	Sea turtle, olive ridley (<i>Lepidochelys olivacea</i>) - Wherever found, except when listed as endangered under 50 CFR 224.101	T	\$857,751	\$1,062,221,359
154	Peppergrass, Slickspot (<i>Lepidium papilliferum</i>) -	T	\$851,131	\$1,063,072,490

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
155	Ocelot (<i>Leopardus</i> (=Felis) <i>pardalis</i>) - wherever found	E	\$840,609	\$1,063,913,099
156	Butterfly, Karner blue (<i>Lycaeides melissa samuelis</i>) - Wherever found	E	\$839,654	\$1,064,752,753
157	Salamander, frosted flatwoods (<i>Ambystoma cingulatum</i>) - Wherever found	T	\$839,614	\$1,065,592,367
158	Whale, North Pacific Right (<i>Eubalaena japonica</i>) - Wherever found	E	\$832,185	\$1,066,424,552
159	Skink, bluetail mole (<i>Eumeces egregius lividus</i>) - Wherever found	T	\$830,028	\$1,067,254,580
160	Warbler (=wood), golden-cheeked (<i>Dendroica chrysoparia</i>) - Wherever found	E	\$829,700	\$1,068,084,280
161	Sparrow, San Clemente sage (<i>Amphispiza belli clementeae</i>) - Wherever found	T	\$824,982	\$1,068,909,262
162	Turtle, bog (<i>Clemmys muhlenbergii</i>) - Wherever found, except GA, NC, SC, TN, VA	T	\$818,140	\$1,069,727,402
163	Sucker, Santa Ana (<i>Catostomus santaanae</i>) - 3 CA river basins	T	\$809,618	\$1,070,537,020
164	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) - New York Bight DPS	E	\$803,997	\$1,071,341,017
165	Bat, Florida bonneted (<i>Eumops floridanus</i>) - Wherever found	E	\$802,799	\$1,072,143,816
166	Topminnow, Gila (incl. Yaqui) (<i>Poeciliopsis occidentalis</i>) - Wherever found	E	\$793,544	\$1,072,937,360
167	Whale, false killer (<i>Pseudorca crassidens</i>) - Main Hawaiian Islands Insular DPS	E	\$793,495	\$1,073,730,855
168	Butterfly, Quino checkerspot (<i>Euphydryas editha quino</i> (=E. e. wrighti)) - Wherever found	E	\$791,271	\$1,074,522,126
169	Sea turtle, green (<i>Chelonia mydas</i>) - Central South Pacific DPS	E	\$779,961	\$1,075,302,087
170	Wolf, red (<i>Canis rufus</i>) - U.S.A. (portions of NC and TN)	EXPN	\$776,100	\$1,076,078,187
171	Fadang (<i>Cycas micronesica</i>) - Wherever found	T	\$773,929	\$1,076,852,116
172	Minnow, loach (<i>Tiaroga cobitis</i>) - Wherever found	E	\$772,864	\$1,077,624,980
173	Duck, Hawaiian (=koloa) (<i>Anas wyvilliana</i>) - Wherever found	E	\$771,837	\$1,078,396,817
174	Beetle, valley elderberry longhorn (<i>Desmocerus californicus dimorphus</i>) - Wherever found	T	\$771,088	\$1,079,167,905
175	Spikedace (<i>Meda fulgida</i>) - Wherever found	E	\$769,352	\$1,079,937,257
176	Rockfish, yelloweye (<i>Sebastes ruberrimus</i>) - Puget Sound - Georgia Basin DPS	T	\$765,703	\$1,080,702,960
177	Goby, tidewater (<i>Eucyclogobius newberryi</i>) - Wherever found	E	\$765,401	\$1,081,468,361
178	Tortoise, gopher (<i>Gopherus polyphemus</i>) - West of Mobile and Tombigbee Rivers	T	\$763,170	\$1,082,231,531
179	Eider, Steller's (<i>Polysticta stelleri</i>) - AK breeding pop.	T	\$758,512	\$1,082,990,043
180	Bocaccio (<i>Sebastes paucispinis</i>) - Puget Sound - Georgia Basin DPS	E	\$750,112	\$1,083,740,155
181	Woundfin (<i>Plagopterus argentissimus</i>) - Wherever found, except where listed as an experimental population	E	\$749,925	\$1,084,490,080
182	Chub, Virgin River (<i>Gila seminuda</i> (=robusta)) - Wherever found	E	\$747,959	\$1,085,238,039

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
183	Sturgeon, Atlantic (<i>Acipenser oxyrinchus oxyrinchus</i>) - South Atlantic DPS	E	\$739,886	\$1,085,977,925
184	Owl, Mexican spotted (<i>Strix occidentalis lucida</i>) - Wherever found	T	\$733,927	\$1,086,711,852
185	Rockfish, canary (<i>Sebastes pinniger</i>) - Puget Sound - Georgia Basin DPS	T	\$722,769	\$1,087,434,621
186	Ferret, black-footed (<i>Mustela nigripes</i>) - U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	EXPN	\$716,048	\$1,088,150,669
187	Frog, Sierra Nevada Yellow-legged (<i>Rana sierrae</i>) - Wherever found	E	\$705,856	\$1,088,856,525
188	Layia, beach (<i>Layia carnosa</i>) - Wherever found	E	\$704,219	\$1,089,560,744
189	No Common Name (<i>Acropora globiceps</i>) - Wherever found	T	\$702,853	\$1,090,263,597
190	Crane, Mississippi sandhill (<i>Grus canadensis pulla</i>) - Wherever found	E	\$702,013	\$1,090,965,610
191	Higgins eye (pearlymussel) (<i>Lampsilis higginsii</i>) - Wherever found	E	\$696,011	\$1,091,661,621
192	Tern, roseate (<i>Sterna dougallii dougallii</i>) - northeast U.S. nesting pop.	E	\$658,726	\$1,092,320,347
193	Plover, piping (<i>Charadrius melodus</i>) - [Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	E	\$655,305	\$1,092,975,652
194	Chub, Gila (<i>Gila intermedia</i>) - Wherever found	E	\$652,162	\$1,093,627,814
195	Salamander, Reticulated flatwoods (<i>Ambystoma bishopi</i>) - Wherever found	E	\$651,425	\$1,094,279,239
196	Salamander, California tiger (<i>Ambystoma californiense</i>) - U.S.A. (CA - Santa Barbara County)	E	\$642,208	\$1,094,921,447
197	Gallinule, Hawaiian common (<i>Gallinula galeata sandvicensis</i>) - Wherever found	E	\$639,889	\$1,095,561,336
198	Salamander, California tiger (<i>Ambystoma californiense</i>) - U.S.A. (CA - Sonoma County)	E	\$622,603	\$1,096,183,939
199	Gibbons (<i>Hylobates</i> spp. (including <i>Nomascus</i>)) - Wherever found	E	\$621,749	\$1,096,805,688
200	Toad, arroyo (=arroyo southwestern) (<i>Anaxyrus californicus</i>) - Wherever found	E	\$620,359	\$1,097,426,047
201	Toad, Wyoming (<i>Bufo hemiophrys baxteri</i>) - Wherever found	E	\$618,390	\$1,098,044,437
202	Caribou, Woodland (<i>Rangifer tarandus caribou</i>) - Selkirk Mountain population	E	\$616,270	\$1,098,660,707
203	Coral, staghorn (<i>Acropora cervicornis</i>) - Wherever found	T	\$601,866	\$1,099,262,573
204	Ladies'-tresses, Ute (<i>Spiranthes diluvialis</i>) - Wherever found	T	\$601,763	\$1,099,864,336
205	Pocket gopher, Yelm (<i>Thomomys mazama yelmensis</i>) - Wherever found	T	\$596,531	\$1,100,460,867
206	Swiftlet, Mariana gray (<i>Aerodramus vanikorensis bartschi</i>) - Wherever found	E	\$592,401	\$1,101,053,268
207	Mucket, pink (pearlymussel) (<i>Lampsilis abrupta</i>) - Wherever found	E	\$585,294	\$1,101,638,562
208	Wolf, gray (<i>Canis lupus</i>) - MN	T	\$576,700	\$1,102,215,262
209	Shiner, Pecos bluntnose (<i>Notropis simus pecosensis</i>) - Wherever found	T	\$572,000	\$1,102,787,262
210	Frog, Chiricahua leopard (<i>Rana chiricahuensis</i>) - Wherever found	T	\$568,024	\$1,103,355,286

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
211	Sparrow, Florida grasshopper (<i>Ammodramus savannarum floridanus</i>) - Wherever found	E	\$558,578	\$1,103,913,864
212	Catchfly, Spalding's (<i>Silene spaldingii</i>) - Wherever found	T	\$552,205	\$1,104,466,069
213	Coral, elkhorn (<i>Acropora palmata</i>) - Wherever found	T	\$549,410	\$1,105,015,479
214	Seagrass, Johnson's (<i>Halophila johnsonii</i>) - Wherever found	T	\$547,824	\$1,105,563,303
215	Trout, Greenback Cutthroat (<i>Oncorhynchus clarkii stomias</i>) - Wherever found	T	\$541,279	\$1,106,104,582
216	Coral, Pillar (<i>Dendrogyra cylindrus</i>) - Wherever found	T	\$537,504	\$1,106,642,086
217	Shearwater, Newell's Townsend's (<i>Puffinus auricularis newelli</i>) - Wherever found	T	\$536,646	\$1,107,178,732
218	Coral, Lobed Star (<i>Orbicella annularis</i>) - Wherever found	T	\$523,628	\$1,107,702,360
219	Coral, Rough Cactus (<i>Mycetophyllia ferox</i>) - Wherever found	T	\$523,385	\$1,108,225,745
220	Kangaroo rat, San Bernardino Merriam's (<i>Dipodomys merriami parvus</i>) - Wherever found	E	\$521,472	\$1,108,747,217
221	Moth, Blackburn's sphinx (<i>Manduca blackburni</i>) - Wherever found	E	\$518,970	\$1,109,266,187
222	Vole, Amargosa (<i>Microtus californicus scirpensis</i>) - Wherever found	E	\$511,117	\$1,109,777,304
223	Kangaroo rat, giant (<i>Dipodomys ingens</i>) - Wherever found	E	\$510,598	\$1,110,287,902
224	Phlox, Yreka (<i>Phlox hirsuta</i>) - Wherever found	E	\$506,550	\$1,110,794,452
225	Paintbrush, golden (<i>Castilleja levisecta</i>) - Wherever found	T	\$495,692	\$1,111,290,144
226	Jaguarundi, Panamanian (<i>Herpailurus (=Felis) yagouaroundi panamensis</i>) - Wherever found	E	\$490,600	\$1,111,780,744
227	Desert-parsley, Bradshaw's (<i>Lomatium bradshawii</i>) - Wherever found	E	\$489,888	\$1,112,270,632
228	Skipperling, Poweshiek (<i>Oarisma poweshiek</i>) - Wherever found	E	\$487,372	\$1,112,758,004
229	Trout, Paiute cutthroat (<i>Oncorhynchus clarkii seleniris</i>) - Wherever found	T	\$486,264	\$1,113,244,268
230	Wedgemussel, dwarf (<i>Alasmidonta heterodon</i>) - Wherever found	E	\$484,680	\$1,113,728,948
231	Coral, Mountainous Star (<i>Orbicella faveolata</i>) - Wherever found	T	\$484,138	\$1,114,213,086
232	Coral, Boulder star (<i>Orbicella franksi</i>) - Wherever found	T	\$483,618	\$1,114,696,704
233	Lupine, Kincaid's (<i>Lupinus sulphureus</i> ssp. <i>kincaidii</i>) - Wherever found	T	\$480,795	\$1,115,177,499
234	Panda, giant (<i>Ailuropoda melanoleuca</i>) - Wherever found	E	\$478,300	\$1,115,655,799
235	Mouse, Preble's meadow jumping (<i>Zapus hudsonius preblei</i>) - wherever found	T	\$474,932	\$1,116,130,731
236	Cui-ui (<i>Chasmistes cujus</i>) - Wherever found	E	\$467,280	\$1,116,598,011
237	Grouper, Nassau (<i>Epinephelus striatus</i>)	T	\$464,945	\$1,117,062,956
238	Sea turtle, loggerhead (<i>Caretta caretta</i>) - Northeast Atlantic Ocean DPS	E	\$464,805	\$1,117,527,761
239	Mouse, Pacific pocket (<i>Perognathus longimembris pacificus</i>) - Wherever found	E	\$462,264	\$1,117,990,025

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
240	Heelsplitter, Carolina (<i>Lasmigona decorata</i>) - Wherever found	E	\$459,489	\$1,118,449,514
241	Vireo, Black-capped (<i>Vireo atricapilla</i>) - Wherever found	E	\$457,540	\$1,118,907,054
242	Lion (<i>Panthera leo</i> ssp. <i>leo</i>) - Wherever found	E	\$457,271	\$1,119,364,325
243	Pupfish, desert (<i>Cyprinodon macularius</i>) - Wherever found	E	\$454,971	\$1,119,819,296
244	Crocodile, American (<i>Crocodylus acutus</i>) - U.S.A. (FL)	T	\$448,899	\$1,120,268,195
245	Butterfly, Oregon silverspot (<i>Speyeria zerene hippolyta</i>) - Wherever found	T	\$446,995	\$1,120,715,190
246	Pocket gopher, Roy Prairie (<i>Thomomys mazama glacialis</i>) - Wherever found	T	\$441,197	\$1,121,156,387
247	Toad, Yosemite (<i>Anaxyrus canorus</i>) - Wherever found	T	\$437,722	\$1,121,594,109
248	Snail, Guam tree (<i>Partula radiolata</i>) - Wherever found	E	\$435,275	\$1,122,029,384
249	Condor, California (<i>Gymnogyps californianus</i>) - U.S.A. (specific portions of Arizona, Nevada, and Utah)	EXPN	\$418,405	\$1,122,447,789
250	Frog, mountain yellow-legged (<i>Rana muscosa</i>) - Northern California DPS	E	\$417,953	\$1,122,865,742
251	Kangaroo rat, Stephens' (<i>Dipodomys stephensi</i> (incl. <i>D. cascus</i>)) - Wherever found	E	\$417,614	\$1,123,283,356
252	Bumble bee, Rusty patched (<i>Bombus affinis</i>) - Wherever found	E	\$416,185	\$1,123,699,541
253	No Common Name (<i>Seriatopora aculeata</i>) - Wherever found	T	\$410,419	\$1,124,109,960
254	Rhinoceros, Sumatran (<i>Dicerorhinus sumatrensis</i>) - Wherever found	E	\$410,106	\$1,124,520,066
255	Rail, light-footed clapper (<i>Rallus longirostris levipes</i>) - Wherever found	E	\$409,665	\$1,124,929,731
256	Kingfisher, Guam Micronesian (<i>Halcyon cinnamomina cinnamomina</i>) - Wherever found	E	\$409,541	\$1,125,339,272
257	Rhinoceros, Southern white (<i>Ceratotherium simum</i> ssp. <i>simum</i>) - Wherever found	SAT	\$406,075	\$1,125,745,347
258	`Akoko, Ewa Plains (<i>Euphorbia skottsbergii</i> var. <i>skottsbergii</i>) - Wherever found	E	\$399,394	\$1,126,144,741
259	Akikiki (<i>Oreomystis bairdi</i>) - Wherever found	E	\$393,046	\$1,126,537,787
260	Kangaroo rat, Fresno (<i>Dipodomys nitratoideis exilis</i>) - Wherever found	E	\$391,742	\$1,126,929,529
261	Squirrel, Northern Idaho Ground (<i>Uroditellus brunneus</i>) - Wherever found	T	\$388,669	\$1,127,318,198
262	Massasauga (=rattlesnake), eastern (<i>Sistrurus catenatus</i>) - Wherever found	T	\$383,807	\$1,127,702,005
263	Skink, sand (<i>Neoseps reynoldsi</i>) - Wherever found	T	\$376,267	\$1,128,078,272
264	Snail, Snake River physa (<i>Physa natricina</i>) - Wherever found	E	\$372,648	\$1,128,450,920
265	Rabbit, Columbia Basin Pygmy (<i>Brachylagus idahoensis</i>) - Columbia Basin DPS	E	\$368,801	\$1,128,819,721
266	Pocket gopher, Olympia (<i>Thomomys mazama pugetensis</i>) - Wherever found	T	\$360,401	\$1,129,180,122
267	Yerba santa, Lompoc (<i>Eriodictyon capitatum</i>) - Wherever found	E	\$358,812	\$1,129,538,934
268	Snake, San Francisco garter (<i>Thamnophis sirtalis tetrataenia</i>) - Wherever found	E	\$352,349	\$1,129,891,283
269	No Common Name (<i>Acropora jacquelineae</i>) - Wherever found	T	\$350,300	\$1,130,241,583

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
269	No Common Name (<i>Acropora speciosa</i>) - Wherever found	T	\$350,300	\$1,130,591,883
269	No Common Name (<i>Euphyllia paradivisa</i>) - Wherever found	T	\$350,300	\$1,130,942,183
269	No Common Name (<i>Isopora crateriformis</i>) - Wherever found	T	\$350,300	\$1,131,292,483
270	Fritillary, Gentner's (<i>Fritillaria gentneri</i>) - Wherever found	E	\$350,262	\$1,131,642,745
271	Daisy, Willamette (<i>Erigeron decumbens</i>) - Wherever found	E	\$349,869	\$1,131,992,614
272	Sea turtle, loggerhead (<i>Caretta caretta</i>) - South Atlantic Ocean DPS	T	\$349,235	\$1,132,341,849
273	Deer, Columbian white-tailed (<i>Odocoileus virginianus leucurus</i>) - Columbia River DPS	E	\$345,736	\$1,132,687,585
274	Beetle, Casey's June (<i>Dinacoma caseyi</i>) - Wherever found	E	\$345,554	\$1,133,033,139
275	No Common Name (<i>Acropora retusa</i>) - Wherever found	T	\$344,279	\$1,133,377,418
276	Lizard, blunt-nosed leopard (<i>Gambelia silus</i>) - Wherever found	E	\$341,115	\$1,133,718,533
277	Sheep, Sierra Nevada bighorn (<i>Ovis canadensis sierrae</i>) - Sierra Nevada	E	\$339,468	\$1,134,058,001
278	Darter, Okaloosa (<i>Etheostoma okaloosae</i>) - Wherever found	T	\$334,439	\$1,134,392,440
279	Rail, Guam (<i>Rallus owstoni</i>) - Wherever found, except where listed as an experimental population	E	\$333,839	\$1,134,726,279
280	Frog, dusky gopher (<i>Rana sevosa</i>) - Wherever found	E	\$333,734	\$1,135,060,013
281	Mouse, New Mexico meadow jumping (<i>Zapus hudsonius luteus</i>) - Wherever found	E	\$332,695	\$1,135,392,708
282	Dragonfly, Hine's emerald (<i>Somatochlora hineana</i>) - Wherever found	E	\$314,125	\$1,135,706,833
283	Bankclimber, purple (mussel) (<i>Elliptoideus sloatianus</i>) - Wherever found	T	\$306,021	\$1,136,012,854
284	Wolf, maned (<i>Chrysocyon brachyurus</i>) - Wherever found	E	\$302,000	\$1,136,314,854
285	Akekee (<i>Loxops caeruleirostris</i>) - Wherever found	E	\$300,699	\$1,136,615,553
286	Gartersnake, northern Mexican (<i>Thamnophis eques megalops</i>) - Wherever found	T	\$298,832	\$1,136,914,385
287	Pearlshell, Louisiana (<i>Margaritifera hembeli</i>) - Wherever found	T	\$293,220	\$1,137,207,605
288	Butterfly, El Segundo blue (<i>Euphilotes battoides allyni</i>) - Wherever found	E	\$293,040	\$1,137,500,645
289	Crane, whooping (<i>Grus americana</i>) - U.S.A (Southwestern Louisiana)	EXPN	\$291,250	\$1,137,791,895
290	Whale, gray (<i>Eschrichtius robustus</i>) - Western North Pacific DPS	E	\$291,166	\$1,138,083,061
291	Crane, whooping (<i>Grus americana</i>) - U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	EXPN	\$289,016	\$1,138,372,077
292	Mapleleaf, winged (<i>Quadrula fragosa</i>) - Wherever found, except where listed as an experimental population	E	\$288,507	\$1,138,660,584
293	Sucker, Warner (<i>Catostomus warnerensis</i>) - Wherever found	T	\$283,723	\$1,138,944,307
294	Wallflower, Ben Lomond (<i>Erysimum teretifolium</i>) - Wherever found	E	\$280,109	\$1,139,224,416

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
295	Milk-vetch, Peirson's (<i>Astragalus magdalenae</i> var. <i>peirsonii</i>) - Wherever found	T	\$277,270	\$1,139,501,686
296	Warbler, Kirtland's (<i>Setophaga kirtlandii</i> (= <i>Dendroica kirtlandii</i>)) - Wherever found	E	\$276,746	\$1,139,778,432
297	Chub, Pahrnagat roundtail (<i>Gila robusta jordani</i>) - Wherever found	E	\$276,664	\$1,140,055,096
298	Orangutan (<i>Pongo pygmaeus</i>) - Wherever found	E	\$275,535	\$1,140,330,631
299	Spectaclecase (mussel) (<i>Cumberlandia monodonta</i>) - Wherever found	E	\$269,263	\$1,140,599,894
300	Mucket, Neosho (<i>Lampsilis rafinesqueana</i>) - Wherever found	E	\$265,039	\$1,140,864,933
301	Bat, Virginia big-eared (<i>Corynorhinus</i> (= <i>Plecotus</i>) <i>townsendii virginianus</i>) - Wherever found	E	\$262,123	\$1,141,127,056
302	Amaranth, seabeach (<i>Amaranthus pumilus</i>) - Wherever found	T	\$259,141	\$1,141,386,197
303	Whipsnake (=striped racer), Alameda (<i>Masticophis lateralis euryxanthus</i>) - Wherever found	T	\$258,501	\$1,141,644,698
304	Bobwhite, masked (quail) (<i>Colinus virginianus ridgwayi</i>) - Wherever found	E	\$258,451	\$1,141,903,149
305	Elktoe, Appalachian (<i>Alasmidonta raveneliana</i>) - Wherever found	E	\$257,462	\$1,142,160,611
306	Butterfly, Saint Francis' satyr (<i>Neonympha mitchellii francisci</i>) - Wherever found	E	\$249,481	\$1,142,410,092
307	Skipper, Dakota (<i>Hesperia dacotae</i>) - Wherever found	T	\$248,609	\$1,142,658,701
308	Wolf, gray (<i>Canis lupus</i>) - WY, EXPN population	EXPN	\$247,871	\$1,142,906,572
309	Darter, snail (<i>Percina tanasi</i>) - Wherever found	T	\$242,367	\$1,143,148,939
310	Tern, roseate (<i>Sterna dougallii dougallii</i>) - Western Hemisphere except NE U.S.	T	\$241,843	\$1,143,390,782
311	Bison, wood (<i>Bison bison athabascaae</i>) - Wherever found	T	\$241,563	\$1,143,632,345
312	Pocket gopher, Tenino (<i>Thomomys mazama tumuli</i>) - Wherever found	T	\$241,431	\$1,143,873,776
313	Gartersnake, narrow-headed (<i>Thamnophis rufipunctatus</i>) - Wherever found	T	\$240,320	\$1,144,114,096
314	Wolf, red (<i>Canis rufus</i>) - Wherever found, except where listed as an experimental population	E	\$240,000	\$1,144,354,096
315	Bush-mallow, San Clemente Island (<i>Malacothamnus clementinus</i>) - Wherever found	E	\$239,408	\$1,144,593,504
316	Bat, Mariana fruit (=Mariana flying fox) (<i>Pteropus mariannus mariannus</i>) - Wherever found	T	\$238,453	\$1,144,831,957
317	No Common Name (<i>Abutilon sandwicense</i>) - Wherever found	E	\$235,730	\$1,145,067,687
318	Storm-petrel, band-rumped (<i>Oceanodroma castro</i>) - USA (HI)	E	\$235,611	\$1,145,303,298
319	Brodiaea, thread-leaved (<i>Brodiaea filifolia</i>) - Wherever found	T	\$235,241	\$1,145,538,539
320	Seal, Ringed (<i>Phoca</i> (= <i>Pusa</i>) <i>hispida hispida</i>) - Wherever found	T	\$234,089	\$1,145,772,628
321	Pronghorn, Sonoran (<i>Antilocapra americana sonoriensis</i>) - Wherever found, except where listed as an experimental population	E	\$232,500	\$1,146,005,128

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
322	Sumac, Michaux's (<i>Rhus michauxii</i>) - Wherever found	E	\$228,338	\$1,146,233,466
323	Tortoise, desert (<i>Gopherus agassizii</i>) - AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	SAT	\$223,424	\$1,146,456,890
324	Falcon, Northern Aplomado (<i>Falco femoralis septentrionalis</i>) - Wherever found, except where listed as an experimental population	E	\$215,059	\$1,146,671,949
325	Butterfly, Palos Verdes blue (<i>Glaucopsyche lygdamus palosverdesensis</i>) - Wherever found	E	\$214,682	\$1,146,886,631
326	Bat, lesser long-nosed (<i>Leptonycteris curasoae yerbabuenae</i>) - Wherever found	E	\$213,029	\$1,147,099,660
327	Threeridge, fat (mussel) (<i>Amblema neislerii</i>) - Wherever found	E	\$209,721	\$1,147,309,381
328	Milkweed, Mead's (<i>Asclepias meadii</i>) - Wherever found	T	\$209,666	\$1,147,519,047
329	Dace, Foscett speckled (<i>Rhinichthys osculus</i> ssp.) - Wherever found	T	\$207,369	\$1,147,726,416
330	No Common Name (<i>Tuberolabium guamense</i>) - Wherever found	T	\$206,850	\$1,147,933,266
331	Snail, Bliss Rapids (<i>Taylorconcha serpenticola</i>) - Wherever found	T	\$205,726	\$1,148,138,992
332	Goldfields, Contra Costa (<i>Lasthenia conjugens</i>) - Wherever found	E	\$204,558	\$1,148,343,550
333	Pogonia, small whorled (<i>Isotria medeoloides</i>) - Wherever found	T	\$204,503	\$1,148,548,053
334	Pigeon wings (<i>Clitoria fragrans</i>) - Wherever found	T	\$204,008	\$1,148,752,061
335	Chaffseed, American (<i>Schwalbea americana</i>) - Wherever found	E	\$203,418	\$1,148,955,479
336	Mussel, sheepsnose (<i>Plethobasus cyphus</i>) - Wherever found	E	\$202,274	\$1,149,157,753
337	Mehamehame (<i>Flueggea neowawraea</i>) - Wherever found	E	\$202,234	\$1,149,359,987
338	Pink, swamp (<i>Helonias bullata</i>) - Wherever found	T	\$200,425	\$1,149,560,412
339	Pangolin, Temnick's ground (<i>Manis temminckii</i>) - Wherever found	E	\$199,797	\$1,149,760,209
340	Clover, running buffalo (<i>Trifolium stoloniferum</i>) - Wherever found	E	\$199,042	\$1,149,959,251
341	Tortoise, angulated (<i>Geochelone yniphora</i>) - Wherever found	E	\$196,521	\$1,150,155,772
342	Shiner, Arkansas River (<i>Notropis girardi</i>) - Arkansas River Basin (AR, KS, NM, OK, TX)	T	\$196,478	\$1,150,352,250
343	Seal, Guadalupe fur (<i>Arctocephalus townsendi</i>) - Wherever found	T	\$191,706	\$1,150,543,956
344	Fairy shrimp, Conservancy (<i>Branchinecta conservatio</i>) - Wherever found	E	\$191,568	\$1,150,735,524
345	Fairy shrimp, San Diego (<i>Branchinecta sandiegonensis</i>) - Wherever found	E	\$191,548	\$1,150,927,072
346	Oryx, scimitar-horned (<i>Oryx dammah</i>) - Wherever found	E	\$189,750	\$1,151,116,822
347	No Common Name (<i>Hesperomannia arborescens</i>) - Wherever found	E	\$189,654	\$1,151,306,476
348	Fairy shrimp, longhorn (<i>Branchinecta longiantenna</i>) - Wherever found	E	\$189,034	\$1,151,495,510
349	No Common Name (<i>Phyllostegia hirsuta</i>) - Wherever found	E	\$188,689	\$1,151,684,199

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
350	Haha (<i>Cyanea koolauensis</i>) - Wherever found	E	\$186,601	\$1,151,870,800
351	Nanu (<i>Gardenia mannii</i>) - Wherever found	E	\$186,502	\$1,152,057,302
352	Cheetah (<i>Acinonyx jubatus</i>) - Wherever found	E	\$186,050	\$1,152,243,352
353	Albatross, short-tailed (<i>Phoebastria (=Diomedea) albatrus</i>) - Wherever found	E	\$184,911	\$1,152,428,263
354	Nioi (<i>Eugenia koolauensis</i>) - Wherever found	E	\$182,514	\$1,152,610,777
355	Kuahiwi laukahi (<i>Plantago princeps</i>) - Wherever found	E	\$181,855	\$1,152,792,632
356	Fly, Hawaiian picture-wing (<i>Drosophila differens</i>) - Wherever found	E	\$180,211	\$1,152,972,843
357	Larkspur, San Clemente Island (<i>Delphinium variegatum</i> ssp. <i>kinkiense</i>) - Wherever found	E	\$178,246	\$1,153,151,089
358	Spineflower, Monterey (<i>Chorizanthe pungens</i> var. <i>pungens</i>) - Wherever found	T	\$178,073	\$1,153,329,162
359	Harvestman, Bone Cave (<i>Texella reyesi</i>) - Wherever found	E	\$177,950	\$1,153,507,112
360	Monardella, willowy (<i>Monardella viminea</i>) - Wherever found	E	\$177,032	\$1,153,684,144
361	Butterfly, Mariana eight-spot (<i>Hypolimnys octocula marianensis</i>) - Wherever found	E	\$176,708	\$1,153,860,852
362	Elepaio, Oahu (<i>Chasiempis ibidis</i>) - Wherever found	E	\$176,224	\$1,154,037,076
363	Butterfly, Lange's metalmark (<i>Apodemia mormo langei</i>) - Wherever found	E	\$176,018	\$1,154,213,094
364	Lotus, San Clemente Island (<i>Acmispon dendroideus</i> var. <i>traskiae (=Lotus d. ssp. traskiae)</i>) - Wherever found	T	\$174,749	\$1,154,387,843
365	Pocketbook, fat (<i>Potamilus capax</i>) - Wherever found	E	\$173,916	\$1,154,561,759
366	Dace, Moapa (<i>Moapa coriacea</i>) - Wherever found	E	\$173,070	\$1,154,734,829
367	Milk-vetch, Lane Mountain (<i>Astragalus jaegerianus</i>) - Wherever found	E	\$171,853	\$1,154,906,682
368	Rhinoceros, Javan (<i>Rhinoceros sondaicus</i>) - Sikkim, Thailand, Malaysia, Indonesia, Indochina, Burma, Bangladesh	E	\$170,916	\$1,155,077,598
369	Sea turtle, green (<i>Chelonia mydas</i>) - East Indian-West Pacific DPS	T	\$170,521	\$1,155,248,119
370	Stickleback, unarmored threespine (<i>Gasterosteus aculeatus williamsoni</i>) - Wherever found	E	\$170,374	\$1,155,418,493
371	Rabbit, Lower Keys marsh (<i>Sylvilagus palustris hefneri</i>) - Wherever found	E	\$169,905	\$1,155,588,398
372	Spineflower, Orcutt's (<i>Chorizanthe orcuttiana</i>) - Wherever found	E	\$169,630	\$1,155,758,028
373	Loosestrife, rough-leaved (<i>Lysimachia asperulaefolia</i>) - Wherever found	E	\$168,696	\$1,155,926,724
374	Chaff-flower, round-leaved (<i>Achyranthes splendens</i> var. <i>rotundata</i>) - Wherever found	E	\$167,952	\$1,156,094,676
375	Wooly-threads, San Joaquin (<i>Monolopia (=Lembertia) congdonii</i>) - Wherever found	E	\$167,170	\$1,156,261,846

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
376	Rockcress, Santa Cruz Island (<i>Sibara filifolia</i>) - Wherever found	E	\$166,963	\$1,156,428,809
377	Paintbrush, San Clemente Island (<i>Castilleja grisea</i>) - Wherever found	T	\$166,830	\$1,156,595,639
377	Woodland-star, San Clemente Island (<i>Lithophragma maximum</i>) - Wherever found	E	\$166,830	\$1,156,762,469
378	Chimpanzee, pygmy (<i>Pan paniscus</i>) - Wherever found	E	\$166,409	\$1,156,928,878
379	Logperch, Roanoke (<i>Percina rex</i>) - Wherever found	E	\$162,549	\$1,157,091,427
380	Pikeminnow (=squawfish), Colorado (<i>Ptychocheilus lucius</i>) - Salt and Verde R. drainages, AZ	EXPN	\$162,000	\$1,157,253,427
381	Mahoe (<i>Alectryon macrococcus</i>) - Wherever found	E	\$161,103	\$1,157,414,530
382	Parrotbill, Maui (<i>Kiwikiu</i>) (<i>Pseudonestor xanthophrys</i>) - Wherever found	E	\$160,633	\$1,157,575,163
383	Salamander, Jemez Mountains (<i>Plethodon neomexicanus</i>) - Wherever found	E	\$160,500	\$1,157,735,663
384	Falcon, Northern Aplomado (<i>Falco femoralis septentrionalis</i>) - U.S.A (AZ, NM)	EXPN	\$158,100	\$1,157,893,763
385	Trout, Gila (<i>Oncorhynchus gilae</i>) - Wherever found	T	\$158,060	\$1,158,051,823
386	Rabbitsfoot (<i>Quadrula cylindrica cylindrica</i>) - Wherever found	T	\$157,566	\$1,158,209,389
387	Deer, Eld's brow-antlered (<i>Cervus eldi</i>) - Wherever found	E	\$157,400	\$1,158,366,789
388	Four-o'clock, MacFarlane's (<i>Mirabilis macfarlanei</i>) - Wherever found	T	\$157,231	\$1,158,524,020
389	Tortoise, Madagascar radiated (<i>Geochelone radiata</i>) - Wherever found	E	\$155,750	\$1,158,679,770
390	Duck, Laysan (<i>Anas laysanensis</i>) - Wherever found	E	\$154,883	\$1,158,834,653
391	Moorhen, Mariana common (<i>Gallinula chloropus guami</i>) - Wherever found	E	\$154,688	\$1,158,989,341
392	Penstemon, blowout (<i>Penstemon haydenii</i>) - Wherever found	E	\$153,550	\$1,159,142,891
393	No Common Name (<i>Neraudia angulata</i>) - Wherever found	E	\$152,808	\$1,159,295,699
394	Tiger beetle, Miami (<i>Cicindelidia floridana</i>) - Wherever found	E	\$152,500	\$1,159,448,199
395	Leopard (<i>Panthera pardus</i>) - Gabon to Kenya and southward	T	\$152,177	\$1,159,600,376
396	Pearlymussel, slabside (<i>Pleuronaia dolabelloides</i>) - Wherever found	E	\$152,149	\$1,159,752,525
397	Butterfly plant, Colorado (<i>Gaura neomexicana</i> var. <i>coloradensis</i>) - Wherever found	T	\$151,657	\$1,159,904,182
398	Kulu'i (<i>Nototrichium humile</i>) - Wherever found	E	\$151,105	\$1,160,055,287
399	Darter, boulder (<i>Etheostoma wapiti</i>) - Wherever found	E	\$151,100	\$1,160,206,387
400	Cactus, Colorado hookless (<i>Sclerocactus glaucus</i>) - Wherever found	T	\$151,038	\$1,160,357,425
401	Pearlymussel, cracking (<i>Hemistena lata</i>) - Wherever found; Except where listed as Experimental Populations	E	\$150,120	\$1,160,507,545
402	Bird's-beak, soft (<i>Cordylanthus mollis</i> ssp. <i>mollis</i>) - Wherever found	E	\$149,978	\$1,160,657,523
403	Thistle, La Graciosa (<i>Cirsium loncholepis</i>) - Wherever found	E	\$149,733	\$1,160,807,256
404	Mouse, southeastern beach (<i>Peromyscus polionotus niveiventris</i>) - wherever found	T	\$149,215	\$1,160,956,471

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
405	Ko'oloa'ula (<i>Abutilon menziesii</i>) - Wherever found	E	\$149,128	\$1,161,105,599
406	No Common Name (<i>Kadua parvula</i>) - Wherever found	E	\$148,712	\$1,161,254,311
407	Moccasinshell, Suwannee (<i>Medionidus walkeri</i>) - Wherever found	T	\$148,700	\$1,161,403,011
408	Pamakani (<i>Viola chamissoniana</i> ssp. <i>chamissoniana</i>) - Wherever found	E	\$147,075	\$1,161,550,086
409	Lizard, Coachella Valley fringe-toed (<i>Uma inornata</i>) - Wherever found	T	\$146,981	\$1,161,697,067
410	Spinymussel, James (<i>Pleurobema collina</i>) - Wherever found	E	\$145,700	\$1,161,842,767
411	Ma'o hau hele, (=native yellow hibiscus) (<i>Hibiscus brackenridgei</i>) - Wherever found	E	\$145,284	\$1,161,988,051
412	Pigtoe, finerayed (<i>Fusconaia cuneolus</i>) - Wherever found; Except where listed as Experimental Populations	E	\$144,620	\$1,162,132,671
413	No Common Name (<i>Kadua degeneri</i>) - Wherever found	E	\$143,082	\$1,162,275,753
414	Kamakahala (<i>Labordia cyrtandrae</i>) - Wherever found	E	\$142,639	\$1,162,418,392
415	Darter, Etowah (<i>Etheostoma etowahae</i>) - Wherever found	E	\$142,600	\$1,162,560,992
416	No Common Name (<i>Gouania vitifolia</i>) - Wherever found	E	\$142,537	\$1,162,703,529
417	Dace, desert (<i>Eremichthys acros</i>) - Wherever found	T	\$142,108	\$1,162,845,637
418	`Ahinahina (<i>Argyroxiphium sandwicense</i> ssp. <i>macrocephalum</i>) - Wherever found	T	\$141,912	\$1,162,987,549
419	Oha (<i>Delissea subcordata</i>) - Wherever found	E	\$140,736	\$1,163,128,285
420	Fly, Hawaiian picture-wing (<i>Drosophila substenoptera</i>) - Wherever found	E	\$140,662	\$1,163,268,947
421	Fiddleneck, large-flowered (<i>Amsinckia grandiflora</i>) - Wherever found	E	\$140,640	\$1,163,409,587
422	No Common Name (<i>Schiedea kaalae</i>) - Wherever found	E	\$140,336	\$1,163,549,923
423	Haha (<i>Cyanea superba</i>) - Wherever found	E	\$140,040	\$1,163,689,963
424	Howellia, water (<i>Howellia aquatilis</i>) - Wherever found	T	\$139,990	\$1,163,829,953
425	Snail, Morro shoulderband (=Banded dune) (<i>Helminthoglypta walkeriana</i>) - Wherever found	E	\$139,426	\$1,163,969,379
426	`Akoko (<i>Euphorbia herbstii</i>) - Wherever found	E	\$138,878	\$1,164,108,257
427	No Common Name (<i>Hesperomannia arbuscula</i>) - Wherever found	E	\$138,577	\$1,164,246,834
428	Towhee, Inyo California (<i>Pipilo crissalis eremophilus</i>) - Wherever found	T	\$138,518	\$1,164,385,352
429	No Common Name (<i>Phyllostegia mollis</i>) - Wherever found	E	\$138,331	\$1,164,523,683
430	Mussel, snuffbox (<i>Epioblasma triquetra</i>) - Wherever found	E	\$138,249	\$1,164,661,932
431	Warbler, nightingale reed (old world warbler) (<i>Acrocephalus luscini</i> a) - Wherever found	E	\$137,864	\$1,164,799,796
432	Piperia, Yadon's (<i>Piperia yadonii</i>) - Wherever found	E	\$137,376	\$1,164,937,172
433	Haha (<i>Cyanea grimesiana</i> ssp. <i>obatae</i>) - Wherever found	E	\$137,360	\$1,165,074,532

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
434	Kamanomano (<i>Cenchrus agrimonioides</i>) - Wherever found	E	\$137,293	\$1,165,211,825
435	No Common Name (<i>Phyllostegia kaalaensis</i>) - Wherever found	E	\$137,199	\$1,165,349,024
436	Boa, Puerto Rican (<i>Epicrates inornatus</i>) - Wherever found	E	\$137,068	\$1,165,486,092
437	No Common Name (<i>Schiedea nuttallii</i>) - Wherever found	E	\$136,952	\$1,165,623,044
438	No Common Name (<i>Schiedea obovata</i>) - Wherever found	E	\$136,853	\$1,165,759,897
439	No Common Name (<i>Stenogyne kanehoana</i>) - Wherever found	E	\$136,698	\$1,165,896,595
440	Slabshell, Chipola (<i>Elliptio chipolaensis</i>) - Wherever found	T	\$136,508	\$1,166,033,103
441	`Akoko (<i>Euphorbia celastroides</i> var. <i>kaenana</i>) - Wherever found	E	\$136,115	\$1,166,169,218
442	No Common Name (<i>Tetramolopium filiforme</i>) - Wherever found	E	\$135,977	\$1,166,305,195
443	Chub, Borax Lake (<i>Gila boraxobius</i>) - Wherever found	E	\$135,322	\$1,166,440,517
444	Ha`iwale (<i>Cyrtandra dentata</i>) - Wherever found	E	\$135,305	\$1,166,575,822
444	Haha (<i>Cyanea longiflora</i>) - Wherever found	E	\$135,305	\$1,166,711,127
444	Na`ena`e (<i>Dubautia herbstobatae</i>) - Wherever found	E	\$135,305	\$1,166,846,432
445	No Common Name (<i>Sanicula mariversa</i>) - Wherever found	E	\$135,181	\$1,166,981,613
446	Lo`ulu (<i>Pritchardia kaalae</i>) - Wherever found	E	\$134,788	\$1,167,116,401
447	Nehe (<i>Melanthera tenuifolia</i>) - Wherever found	E	\$134,768	\$1,167,251,169
448	No Common Name (<i>Schiedea trinervis</i>) - Wherever found	E	\$134,653	\$1,167,385,822
449	Mouse, Alabama beach (<i>Peromyscus polionotus ammobates</i>) - Wherever found	E	\$134,407	\$1,167,520,229
450	Tapir, Central American (<i>Tapirus bairdii</i>) - Wherever found	E	\$133,774	\$1,167,654,003
451	Bird's-beak, salt marsh (<i>Cordylanthus maritimus</i> ssp. <i>maritimus</i>) - Wherever found	E	\$133,538	\$1,167,787,541
452	Monkeyface, Cumberland (pearlymussel) (<i>Quadrula intermedia</i>) - Wherever found; Except where listed as Experimental Populations	E	\$133,322	\$1,167,920,863
453	Picture-wing fly, Hawaiian (<i>Drosophila montgomeryi</i>) - Wherever found	E	\$133,057	\$1,168,053,920
454	Dace, blackside (<i>Phoxinus cumberlandensis</i>) - Wherever found	T	\$132,832	\$1,168,186,752
455	Wallflower, Contra Costa (<i>Erysimum capitatum</i> var. <i>angustatum</i>) - Wherever found	E	\$132,660	\$1,168,319,412
456	Wild-rice, Texas (<i>Zizania texana</i>) - Wherever found	E	\$130,600	\$1,168,450,012
457	Pigtoe, shiny (<i>Fusconaia cor</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$130,500	\$1,168,580,512
458	Eagle, harpy (<i>Harpia harpyja</i>) - Wherever found	E	\$130,000	\$1,168,710,512
459	Snail, Humped tree (<i>Partula gibba</i>) - Wherever found	E	\$129,355	\$1,168,839,867
460	Orchid, eastern prairie fringed (<i>Platanthera leucophaea</i>) - Wherever found	T	\$129,345	\$1,168,969,212
461	Wild-buckwheat, gypsum (<i>Eriogonum gypsophilum</i>) - Wherever found	T	\$129,000	\$1,169,098,212

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
462	Thistle, Pitcher's (<i>Cirsium pitcheri</i>) - Wherever found	T	\$128,756	\$1,169,226,968
463	Pupfish, Ash Meadows Amargosa (<i>Cyprinodon nevadensis mionectes</i>) - Wherever found	E	\$128,720	\$1,169,355,688
464	Quillwort, Louisiana (<i>Isoetes louisianensis</i>) - Wherever found	E	128582	\$1,169,484,270
465	Rhinoceros, great Indian (<i>Rhinoceros unicornis</i>) - Nepal, India	E	\$127,581	\$1,169,611,851
466	Cactus, Kuenzler hedgehog (<i>Echinocereus fendleri</i> var. <i>kuenzleri</i>) - Wherever found	E	\$127,300	\$1,169,739,151
467	Kangaroo rat, Tipton (<i>Dipodomys nitratoideus nitratoideus</i>) - Wherever found	E	\$127,296	\$1,169,866,447
468	Bat, Ozark big-eared (<i>Corynorhinus</i> (=Plecotus) <i>townsendii ingens</i>) - Wherever found	E	\$127,163	\$1,169,993,610
469	Cactus, Tobusch fishhook (<i>Sclerocactus brevihamatus</i> ssp. <i>tobuschii</i>) - Wherever found	E	\$125,500	\$1,170,119,110
470	Evening-primrose, Antioch Dunes (<i>Oenothera deltoidea</i> ssp. <i>howellii</i>) - Wherever found	E	\$123,450	\$1,170,242,560
471	Darter, leopard (<i>Percina pantherina</i>) - Wherever found	T	\$121,840	\$1,170,364,400
472	Potentilla, Hickman's (<i>Potentilla hickmanii</i>) - Wherever found	E	\$121,507	\$1,170,485,907
473	Sheep, Peninsular bighorn (<i>Ovis canadensis nelsoni</i>) - Peninsular CA pop.	E	\$120,720	\$1,170,606,627
474	No Common Name (<i>Dendrobium guamense</i>) - Wherever found	T	\$120,380	\$1,170,727,007
475	Checker-mallow, Nelson's (<i>Sidalcea nelsoniana</i>) - Wherever found	T	\$120,354	\$1,170,847,361
476	Cavefish, Ozark (<i>Amblyopsis rosae</i>) - Wherever found	T	\$120,190	\$1,170,967,551
477	Popcornflower, rough (<i>Plagiobothrys hirtus</i>) - Wherever found	E	\$119,021	\$1,171,086,572
478	Hawk, Hawaiian (=ʻIo) (<i>Buteo solitarius</i>) - Wherever found	E	\$117,500	\$1,171,204,072
479	Riffleshell, northern (<i>Epioblasma torulosa rangiana</i>) - Wherever found	E	\$116,665	\$1,171,320,737
480	Lily, Western (<i>Lilium occidentale</i>) - Wherever found	E	\$116,562	\$1,171,437,299
481	Fanshell (<i>Cyprogenia stegaria</i>) - Wherever found	E	\$113,200	\$1,171,550,499
482	Dropwort, Canby's (<i>Oxypolis canbyi</i>) - Wherever found	E	\$112,410	\$1,171,662,909
483	Blackbird, yellow-shouldered (<i>Agelaius xanthomus</i>) - Wherever found	E	\$109,500	\$1,171,772,409
484	Iagu, Hayun (=Guam), Tronkon guafi (Rota) (<i>Serianthes nelsonii</i>) - Wherever found	E	\$109,285	\$1,171,881,694
485	Mouse, Choctawhatchee beach (<i>Peromyscus polionotus allophrys</i>) - Wherever found	E	\$105,890	\$1,171,987,584
486	Bulrush, Northeastern (<i>Scirpus ancistrochaetus</i>) - Wherever found	E	\$105,466	\$1,172,093,050
487	Amole, purple (<i>Chlorogalum purpureum</i>) - Wherever found	T	\$105,213	\$1,172,198,263

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
488	Watercress, Gambel's (<i>Rorippa gambellii</i>) - Wherever found	E	\$104,291	\$1,172,302,554
489	Thrush, small Kauai (=puaiohi) (<i>Myadestes palmeri</i>) - Wherever found	E	\$104,231	\$1,172,406,785
490	Shiner, Cape Fear (<i>Notropis mekistocholas</i>) - Wherever found	E	\$103,500	\$1,172,510,285
491	Toad, Houston (<i>Bufo houstonensis</i>) - Wherever found	E	\$103,493	\$1,172,613,778
492	Beaked-rush, Knieskern's (<i>Rhynchospora knieskernii</i>) - Wherever found	T	\$102,710	\$1,172,716,488
493	Tiger beetle, Salt Creek (<i>Cicindela nevadica lincolniana</i>) - Wherever found	E	\$102,300	\$1,172,818,788
494	Crayfish, Big Sandy (<i>Cambarus callainus</i>) - Wherever found	T	\$100,896	\$1,172,919,684
495	Mouse, St. Andrew beach (<i>Peromyscus polionotus peninsularis</i>) - Wherever found	E	\$100,687	\$1,173,020,371
496	Clubshell (<i>Pleurobema clava</i>) - Wherever found; Except where listed as Experimental Populations	E	\$100,306	\$1,173,120,677
497	Springsnail, Bruneau Hot (<i>Pyrgulopsis bruneauensis</i>) - Wherever found	E	\$99,782	\$1,173,220,459
498	Leopard, snow (<i>Uncia (=Panthera) uncia</i>) - Wherever found	E	\$99,500	\$1,173,319,959
499	Mouse, Anastasia Island beach (<i>Peromyscus polionotus phasma</i>) - Wherever found	E	\$95,315	\$1,173,415,274
500	Butterfly, callippe silverspot (<i>Speyeria callippe callippe</i>) - Wherever found	E	\$94,720	\$1,173,509,994
501	Cactus, Pima pineapple (<i>Coryphantha scheeri</i> var. <i>robustispina</i>) - Wherever found	E	\$93,765	\$1,173,603,759
502	Spinedace, Little Colorado (<i>Lepidomeda vittata</i>) - Wherever found	T	\$92,980	\$1,173,696,739
503	Shrew, Buena Vista Lake ornate (<i>Sorex ornatus relictus</i>) - Wherever found	E	\$92,816	\$1,173,789,555
504	Poolfish, Pahrump (<i>Empetrichthys latos</i>) - Wherever found	E	\$92,675	\$1,173,882,230
505	Butterfly, Mitchell's satyr (<i>Neonympha mitchellii mitchellii</i>) - Wherever found	E	\$92,105	\$1,173,974,335
506	Darter, rush (<i>Etheostoma phytophilum</i>) - Wherever found	E	\$91,500	\$1,174,065,835
506	Trout, bull (<i>Salvelinus confluentus</i>) - Clackamas River subbasin experimental population	EXPN	\$91,500	\$1,174,157,335
507	Frog, mountain yellow-legged (<i>Rana muscosa</i>) - Southern California DPS	E	\$91,068	\$1,174,248,403
508	Lomatium, Cook's (<i>Lomatium cookii</i>) - Wherever found	E	\$90,963	\$1,174,339,366
509	Bat, Mexican long-nosed (<i>Leptonycteris nivalis</i>) - Wherever found	E	\$90,498	\$1,174,429,864
510	Darter, Kentucky arrow (<i>Etheostoma spilotum</i>) - Wherever found	T	\$90,100	\$1,174,519,964
511	No Common Name (<i>Tabernaemontana rotensis</i>) - Wherever found	T	\$90,048	\$1,174,610,012
512	Ivesia, Webber's (<i>Ivesia webberi</i>) - Wherever found	T	\$87,820	\$1,174,697,832
513	Milk-vetch, Braunton's (<i>Astragalus brauntonii</i>) - Wherever found	E	\$87,352	\$1,174,785,184
514	Spurge, telephus (<i>Euphorbia telephioides</i>) - Wherever found	T	\$86,331	\$1,174,871,515
515	Chub, spotfin (<i>Erimonax monachus</i>) - Wherever found, except where listed as an experimental population	T	\$86,118	\$1,174,957,633

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
516	Potato-bean, Price's (<i>Apios priceana</i>) -	T	\$85,826	\$1,175,043,459
517	Jaguarundi, Guatemalan (<i>Herpailurus (=Felis) yagouaroundi fossata</i>) - Mexico, Nicaragua	E	\$85,729	\$1,175,129,188
518	Cladonia, Florida perforate (<i>Cladonia perforata</i>) - Wherever found	E	85700	\$1,175,214,888
519	Spinedace, White River (<i>Lepidomeda albivallis</i>) - Wherever found	E	\$84,268	\$1,175,299,156
520	Darter, duskytail (<i>Etheostoma percnurum</i>) - Wherever found	E	\$83,275	\$1,175,382,431
521	Aster, decurrent false (<i>Boltonia decurrens</i>) - Wherever found	T	\$83,100	\$1,175,465,531
521	Milk-vetch, Sentry (<i>Astragalus cremnophylax</i> var. <i>cremnophylax</i>) - Wherever found	E	\$83,100	\$1,175,548,631
522	Butterfly, Smith's blue (<i>Euphilotes enoptes smithi</i>) - Wherever found	E	\$82,400	\$1,175,631,031
523	Orchid, western prairie fringed (<i>Platanthera praeclara</i>) - Wherever found	T	\$82,191	\$1,175,713,222
524	Cactus, Wright fishhook (<i>Sclerocactus wrightiae</i>) - Wherever found	E	\$81,900	\$1,175,795,122
525	Crayfish, Nashville (<i>Orconectes shoupi</i>) - Wherever found	E	\$81,730	\$1,175,876,852
526	Pennyroyal, Todsens' (<i>Hedeoma todsenii</i>) - Wherever found	E	\$81,100	\$1,175,957,952
527	Damselfly, orangeblack Hawaiian (<i>Megalagrion xanthomelas</i>) - Wherever found	E	\$80,619	\$1,176,038,571
528	Pigtoe, rough (<i>Pleurobema plenum</i>) - Wherever found	E	\$79,947	\$1,176,118,518
529	Gazelle, dama (<i>Gazella dama</i>) - Wherever found	E	\$79,450	\$1,176,197,968
530	Salamander, Santa Cruz long-toed (<i>Ambystoma macrodactylum croceum</i>) - Wherever found	E	\$78,745	\$1,176,276,713
531	Creeper, Hawaii (<i>Oreomystis mana</i>) - Wherever found	E	\$78,493	\$1,176,355,206
532	Evening-primrose, Eureka Valley (<i>Oenothera avita</i> ssp. <i>eurekensis</i>) -	E	\$78,350	\$1,176,433,556
533	Grass, Eureka Dune (<i>Swallenia alexandrae</i>) -	E	\$78,240	\$1,176,511,796
534	Mussel, scaleshell (<i>Leptodea leptodon</i>) - Wherever found	E	\$78,158	\$1,176,589,954
535	Monkeyflower, Vandenberg (<i>Diplacus vandenbergensis</i>) - Wherever found	E	\$77,500	\$1,176,667,454
536	Akepa, Hawaii (<i>Loxops coccineus</i>) - Wherever found	E	\$77,479	\$1,176,744,933
537	Geranium, Hawaiian red-flowered (<i>Geranium arboreum</i>) - Wherever found	E	\$76,691	\$1,176,821,624
538	Beetle, Mount Hermon June (<i>Polyphylla barbata</i>) - Wherever found	E	\$76,626	\$1,176,898,250
539	`Ohe (<i>Joinvillea ascendens ascendens</i>) - Wherever found	E	\$76,530	\$1,176,974,780
540	Manzanita, Franciscan (<i>Arctostaphylos franciscana</i>) - Wherever found	E	\$76,500	\$1,177,051,280
541	Limpet, Banbury Springs (<i>Lanx</i> sp.) - Wherever found	E	\$76,389	\$1,177,127,669
542	Damselfly, Pacific Hawaiian (<i>Megalagrion pacificum</i>) - Wherever found	E	\$76,242	\$1,177,203,911
543	Water-umbel, Huachuca (<i>Lilaeopsis schaffneriana</i> var. <i>recurva</i>) - Wherever found	E	\$76,100	\$1,177,280,011
544	Button-celery, San Diego (<i>Eryngium aristulatum</i> var. <i>parishii</i>) - Wherever found	E	\$75,910	\$1,177,355,921

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
545	Honeycreeper, (Akohekohe) crested (<i>Palmeria dolei</i>) - Wherever found	E	\$75,147	\$1,177,431,068
546	Hellbender, Ozark (<i>Cryptobranchus alleganiensis bishopi</i>) - Wherever found	E	\$74,400	\$1,177,505,468
547	Jaguar (<i>Panthera onca</i>) - Wherever found	E	\$74,302	\$1,177,579,770
548	Springfish, Railroad Valley (<i>Crenichthys nevadae</i>) - Wherever found	T	\$73,940	\$1,177,653,710
549	Mesa-mint, San Diego (<i>Pogogyne abramsii</i>) - Wherever found	E	\$73,939	\$1,177,727,649
550	Thelypody, Howell's spectacular (<i>Thelypodium howellii</i> ssp. <i>spectabilis</i>) - Wherever found	T	\$73,703	\$1,177,801,352
551	Polygonum, Scotts Valley (<i>Polygonum hickmanii</i>) - Wherever found	E	\$73,685	\$1,177,875,037
552	Pigtoe, oval (<i>Pleurobema pyriforme</i>) - Wherever found	E	\$73,294	\$1,177,948,331
553	Pimpleback, orangefoot (pearlymussel) (<i>Plethobasus cooperianus</i>) - Wherever found	E	\$72,733	\$1,178,021,064
554	Pocketbook, shinyrayed (<i>Lampsilis subangulata</i>) - Wherever found	E	\$72,499	\$1,178,093,563
555	Chub, Hutton tui (<i>Gila bicolor</i> ssp.) - Wherever found	T	\$71,373	\$1,178,164,936
556	Butterfly, Mount Charleston blue (<i>Icaricia (Plebejus) shasta charlestonensis</i>) - Wherever found	E	\$71,167	\$1,178,236,103
557	No Common Name (<i>Schiedea haleakalensis</i>) - Wherever found	E	\$71,065	\$1,178,307,168
558	Pupfish, Owens (<i>Cyprinodon radiosus</i>) - Wherever found	E	\$70,880	\$1,178,378,048
559	No Common Name (<i>Phyllostegia bracteata</i>) - Wherever found	E	\$70,475	\$1,178,448,523
560	`Oha wai (<i>Clermontia samuelii</i>) - Wherever found	E	\$70,436	\$1,178,518,959
561	No Common Name (<i>Phyllostegia haliakalae</i>) - Wherever found	E	\$70,254	\$1,178,589,213
562	Haiwale (<i>Cyrtandra ferripilosa</i>) - Wherever found	E	\$70,122	\$1,178,659,335
563	Pondberry (<i>Lindera melissifolia</i>) - Wherever found	E	\$69,902	\$1,178,729,237
564	Harperella (<i>Ptilimnium nodosum</i>) - Wherever found	E	\$69,828	\$1,178,799,065
565	Leopard, clouded (<i>Neofelis nebulosa</i>) - Wherever found	E	\$69,800	\$1,178,868,865
566	No Common Name (<i>Phyllostegia brevidens</i>) - Wherever found	E	\$69,432	\$1,178,938,297
567	Crocodile, African slender-snouted (<i>Crocodylus cataphractus</i>) - Wherever found	E	\$69,420	\$1,179,007,717
568	Akiapolaau (<i>Hemignathus wilsoni</i>) - Wherever found	E	\$69,220	\$1,179,076,937
569	Catfish, Yaqui (<i>Ictalurus pricei</i>) - Wherever found	T	\$68,600	\$1,179,145,537
570	Ihi`ihi (<i>Marsilea villosa</i>) - Wherever found	E	\$68,401	\$1,179,213,938
571	Haha (<i>Cyanea asplenifolia</i>) - Wherever found	E	\$68,067	\$1,179,282,005
572	Gourd, Okeechobee (<i>Cucurbita okeechobeensis</i> ssp. <i>okeechobeensis</i>) - Wherever found	E	\$67,200	\$1,179,349,205

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
573	Ufa-halomtano (<i>Heritiera longipetiolata</i>) - Wherever found	E	\$66,219	\$1,179,415,424
574	Hawk, Puerto Rican sharp-shinned (<i>Accipiter striatus venator</i>) - Wherever found	E	\$66,000	\$1,179,481,424
575	Wolf, Mexican (<i>Canis lupus baileyi</i>) - Wherever found, except where listed as an experimental population	E	\$65,700	\$1,179,547,124
576	Trout, Apache (<i>Oncorhynchus apache</i>) - Wherever found	T	\$65,400	\$1,179,612,524
577	Buckwheat, steamboat (<i>Eriogonum ovalifolium</i> var. <i>williamsiae</i>) - Wherever found	E	\$65,015	\$1,179,677,539
578	Snail, fragile tree (<i>Samoana fragilis</i>) - Wherever found	E	\$64,886	\$1,179,742,425
579	Butterfly, mission blue (<i>Icaricia icarioides missionensis</i>) - Wherever found	E	\$64,320	\$1,179,806,745
580	Halumtano, Cebello (<i>Bulbophyllum guamense</i>) - Wherever found	T	\$64,291	\$1,179,871,036
581	Riffleshell, tan (<i>Epioblasma florentina walkeri</i> (=E. <i>walkeri</i>)) - Wherever found	E	\$63,949	\$1,179,934,985
582	Tiger beetle, Puritan (<i>Cicindela puritana</i>) - Wherever found	T	\$63,634	\$1,179,998,619
583	Warea, wide-leaf (<i>Warea amplexifolia</i>) - Wherever found	E	\$62,700	\$1,180,061,319
584	Snake, black pine (<i>Pituophis melanoleucus lodingi</i>) - Wherever found	T	\$62,508	\$1,180,123,827
585	No Common Name (<i>Pteris lidgatei</i>) - Wherever found	E	\$62,446	\$1,180,186,273
586	Horse, Przewalski's (<i>Equus przewalskii</i>) - Wherever found	E	\$61,950	\$1,180,248,223
587	Cactus, Uinta Basin hookless (<i>Sclerocactus wetlandicus</i>) - Wherever found	T	\$61,686	\$1,180,309,909
588	Bean, purple (<i>Villosa perpurpurea</i>) - Wherever found	E	\$61,620	\$1,180,371,529
589	Pocketbook, Ouachita rock (<i>Arkansia wheeleri</i>) - Wherever found	E	\$61,430	\$1,180,432,959
590	Wawae`iole (<i>Huperzia nutans</i>) - Wherever found	E	\$61,363	\$1,180,494,322
591	No Common Name (<i>Asplenium peruvianum</i> var. <i>insulare</i>) - Wherever found	E	61242	\$1,180,555,564
592	Haha (<i>Cyanea crispa</i>) - Wherever found	E	\$61,153	\$1,180,616,717
593	Mountain beaver, Point Arena (<i>Aplodontia rufa nigra</i>) - Wherever found	E	\$61,100	\$1,180,677,817
594	Ring pink (mussel) (<i>Obovaria retusa</i>) - Wherever found	E	\$60,973	\$1,180,738,790
595	Chub, slender (<i>Erimystax cahni</i>) - Wherever found	T	\$60,522	\$1,180,799,312
596	Ha`iwale (<i>Cyrtandra viridiflora</i>) - Wherever found	E	\$60,274	\$1,180,859,586
597	Lau `ehu (<i>Panicum niihauense</i>) - Wherever found	E	\$60,136	\$1,180,919,722
598	Tiger beetle, Northeastern beach (<i>Cicindela dorsalis dorsalis</i>) - Wherever found	T	\$59,851	\$1,180,979,573
599	Madtom, pygmy (<i>Noturus stanauli</i>) - Wherever found	E	\$58,935	\$1,181,038,508
600	Sea turtle, loggerhead (<i>Caretta caretta</i>) - Southeast Indo-Pacific Ocean DPS	T	\$58,930	\$1,181,097,438
601	Combshell, Cumberlandian (<i>Epioblasma brevidens</i>) - Wherever found; Except where listed as Experimental Populations	E	\$58,890	\$1,181,156,328
602	Darter, Niangua (<i>Etheostoma nianguae</i>) - Wherever found	T	\$58,406	\$1,181,214,734

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
603	Quillwort, black spored (<i>Isoetes melanospora</i>) - Wherever found	E	\$58,200	\$1,181,272,934
604	Mallow, Kern (<i>Eremalche kernensis</i>) - Wherever found	E	58140	\$1,181,331,074
605	Haha (<i>Cyanea humboldtiana</i>) - Wherever found	E	\$58,087	\$1,181,389,161
606	Mountainbalm, Indian Knob (<i>Eriodictyon altissimum</i>) - Wherever found	E	\$57,897	\$1,181,447,058
607	Logperch, Conasauga (<i>Percina jenkinsi</i>) - Wherever found	E	\$57,800	\$1,181,504,858
608	Ko'oko'olau (<i>Bidens micrantha</i> ssp. <i>kalealaha</i>) - Wherever found	E	\$57,651	\$1,181,562,509
609	'Ohe'ohe (<i>Polyscias gymnocarpa</i>) - Wherever found	E	\$57,600	\$1,181,620,109
610	Skyrocket, Pagosa (<i>Ipomopsis polyantha</i>) - Wherever found	E	\$57,561	\$1,181,677,670
611	Haha (<i>Cyanea acuminata</i>) - Wherever found	E	\$57,087	\$1,181,734,757
612	No Common Name (<i>Polyscias racemosa</i>) - Wherever found	E	\$56,956	\$1,181,791,713
613	No Common Name (<i>Spermolepis hawaiiensis</i>) - Wherever found	E	\$56,662	\$1,181,848,375
614	Moccasinshell, Gulf (<i>Medionidus penicillatus</i>) - Wherever found	E	\$56,400	\$1,181,904,775
615	Alani (<i>Melicope lydgatei</i>) - Wherever found	E	\$56,389	\$1,181,961,164
616	Bean, Cumberland (pearlymussel) (<i>Villosa trabalis</i>) - Wherever found; Except where listed as Experimental Populations	E	\$56,371	\$1,182,017,535
617	No Common Name (<i>Platydesma cornuta</i> var. <i>cornuta</i>) - Wherever found	E	\$56,227	\$1,182,073,762
618	Sea turtle, green (<i>Chelonia mydas</i>) - North Indian DPS	T	\$55,807	\$1,182,129,569
619	Alligator, Chinese (<i>Alligator sinensis</i>) - Wherever found	E	\$55,750	\$1,182,185,319
619	Crocodile, Cuban (<i>Crocodylus rhombifer</i>) - Wherever found	E	\$55,750	\$1,182,241,069
619	Crocodile, Philippine (<i>Crocodylus novaeguineae mindorensis</i>) - Wherever found	E	\$55,750	\$1,182,296,819
619	Gavial (<i>Gavialis gangeticus</i>) - Wherever found	E	\$55,750	\$1,182,352,569
619	Iguana, Fiji banded (<i>Brachylophus fasciatus</i>) - Wherever found	E	\$55,750	\$1,182,408,319
619	Monitor, Komodo Island (<i>Varanus komodoensis</i>) - Wherever found	E	\$55,750	\$1,182,464,069
619	Salamander, Japanese giant (<i>Andrias japonicus</i> (= <i>daavidianus</i> j.)) - Wherever found	E	\$55,750	\$1,182,519,819
619	Tartaruga (<i>Podocnemis expansa</i>) - Wherever found	E	\$55,750	\$1,182,575,569
619	Tracaja (<i>Podocnemis unifilis</i>) - Wherever found	E	\$55,750	\$1,182,631,319
620	Akepa, Maui (<i>Loxops ochraceus</i>) - Wherever found	E	\$55,540	\$1,182,686,859
621	Skipper, Carson wandering (<i>Pseudocopaodes eunus obscurus</i>) - Wherever found	E	\$55,376	\$1,182,742,235
622	Heelsplitter, Inflated (<i>Potamilus inflatus</i>) - Wherever found	T	\$54,937	\$1,182,797,172
623	Rail, Guam (<i>Rallus owstoni</i>) - Rota	EXPN	\$54,784	\$1,182,851,956
624	Salamander, Sonora tiger (<i>Ambystoma tigrinum stebbinsi</i>) - Wherever found	E	\$54,390	\$1,182,906,346
625	Pigtoe, fuzzy (<i>Pleurobema strodeanum</i>) - Wherever found	T	\$54,059	\$1,182,960,405

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
626	Ziziphus, Florida (<i>Ziziphus celata</i>) - Wherever found	E	\$54,000	\$1,183,014,405
627	Crayfish, Guyandotte River (<i>Cambarus veteranus</i>) - Wherever found	E	\$53,982	\$1,183,068,387
628	Chub, Owens Tui (<i>Gila bicolor</i> ssp. <i>snyderi</i>) - Wherever found	E	\$53,880	\$1,183,122,267
629	No Common Name (<i>Viola oahuensis</i>) - Wherever found	E	\$53,847	\$1,183,176,114
630	Bladderpod, Short's (<i>Physaria globosa</i>) - Wherever found	E	\$53,427	\$1,183,229,541
631	Rabbit, riparian brush (<i>Sylvilagus bachmani riparius</i>) - Wherever found	E	\$53,098	\$1,183,282,639
632	Lousewort, Furbish (<i>Pedicularis furbishiae</i>) - Wherever found	E	\$52,845	\$1,183,335,484
633	Tarplant, Gaviota (<i>Deinandra increscens</i> ssp. <i>villosa</i>) - Wherever found	E	\$52,705	\$1,183,388,189
634	No Common Name (<i>Neraudia ovata</i>) - Wherever found	E	\$52,500	\$1,183,440,689
635	Goldfields, Burke's (<i>Lasthenia burkei</i>) - Wherever found	E	\$52,490	\$1,183,493,179
636	Coneflower, smooth (<i>Echinacea laevigata</i>) - Wherever found	E	\$52,470	\$1,183,545,649
636	Meadowfoam, Sebastopol (<i>Limnanthes vinculans</i>) - Wherever found	E	\$52,470	\$1,183,598,119
637	Milk-vetch, Applegate's (<i>Astragalus applegatei</i>) - Wherever found	E	\$52,364	\$1,183,650,483
638	Tiger beetle, Ohlone (<i>Cicindela ohlone</i>) - Wherever found	E	\$52,315	\$1,183,702,798
639	Spiraea, Virginia (<i>Spiraea virginiana</i>) - Wherever found	T	\$52,223	\$1,183,755,021
640	Mouse, Perdido Key beach (<i>Peromyscus polionotus trissyllepsis</i>) - Wherever found	E	\$52,222	\$1,183,807,243
641	Megapode, Micronesian (<i>Megapodius laperouse</i>) - Wherever found	E	\$51,753	\$1,183,858,996
642	Darter, Cherokee (<i>Etheostoma scotti</i>) - Wherever found	T	\$51,600	\$1,183,910,596
643	Beetle, [no common name] (<i>Rhadine exilis</i>) - Wherever found	E	\$51,500	\$1,183,962,096
644	Pocketbook, finelined (<i>Lampsilis altilis</i>) - Wherever found	T	\$51,461	\$1,184,013,557
645	Orchid, white fringeless (<i>Platanthera integrilabia</i>) -	T	\$51,400	\$1,184,064,957
646	Sunshine, Sonoma (<i>Blennosperma bakeri</i>) - Wherever found	E	\$50,970	\$1,184,115,927
647	Pupfish, Warm Springs (<i>Cyprinodon nevadensis pectoralis</i>) - Wherever found	E	\$50,933	\$1,184,166,860
648	Kula wahine noho (<i>Isodendron pyrifolium</i>) - Wherever found	E	\$50,744	\$1,184,217,604
649	Kidneyshell, triangular (<i>Ptychobranchus greenii</i>) - Wherever found	E	\$50,682	\$1,184,268,286
650	Isopod, Madison Cave (<i>Antrolana lira</i>) - Wherever found	T	\$50,581	\$1,184,318,867
651	Hawk, Puerto Rican broad-winged (<i>Buteo platypterus brunescens</i>) - Wherever found	E	\$50,500	\$1,184,369,367
652	Skullcap, large-flowered (<i>Scutellaria montana</i>) - Wherever found	T	\$49,641	\$1,184,419,008
653	Haha (<i>Cyanea copelandii</i> ssp. <i>haleakalaensis</i>) - Wherever found	E	\$49,524	\$1,184,468,532
654	Sea turtle, loggerhead (<i>Caretta caretta</i>) - Mediterranean Sea DPS	E	\$49,497	\$1,184,518,029
655	Yellowhead, desert (<i>Yermo xanthocephalus</i>) - Wherever found	T	\$49,168	\$1,184,567,197

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
656	Navarretia, spreading (<i>Navarretia fossalis</i>) - Wherever found	T	\$48,910	\$1,184,616,107
657	Cactus, Winkler (<i>Pediocactus winkleri</i>) - Wherever found	T	\$48,900	\$1,184,665,007
658	Brodiaea, Chinese Camp (<i>Brodiaea pallida</i>) - Wherever found	T	\$48,720	\$1,184,713,727
659	Minnow, Devils River (<i>Dionda diaboli</i>) - Wherever found	T	\$48,604	\$1,184,762,331
660	Ko`oko`olau (<i>Bidens campylotheca</i> ssp. <i>pentamera</i>) - Wherever found	E	\$48,587	\$1,184,810,918
661	Steelhead (<i>Oncorhynchus</i> (=Salmo) <i>mykiss</i>) - Middle Columbia River DPS	EXPN	\$48,544	\$1,184,859,462
662	No Common Name (<i>Sanicula sandwicensis</i>) - Wherever found	E	\$48,505	\$1,184,907,967
663	Bean, rayed (<i>Villosa fabalis</i>) - Wherever found	E	\$48,398	\$1,184,956,365
664	Alani (<i>Melicope ovalis</i>) - Wherever found	E	\$48,328	\$1,185,004,693
665	Moccasinshell, Alabama (<i>Medionidus acutissimus</i>) - Wherever found	T	\$47,882	\$1,185,052,575
666	A`e (<i>Zanthoxylum hawaiiense</i>) - Wherever found	E	\$47,634	\$1,185,100,209
667	Sunflower, Schweinitz's (<i>Helianthus schweinitzii</i>) - Wherever found	E	\$47,562	\$1,185,147,771
668	Pigtoe, southern (<i>Pleurobema georgianum</i>) - Wherever found	E	\$47,500	\$1,185,195,271
668	Trillium, persistent (<i>Trillium persistens</i>) - Wherever found	E	\$47,500	\$1,185,242,771
669	Ko`oko`olau (<i>Bidens campylotheca</i> ssp. <i>waihoiensis</i>) - Wherever found	E	\$47,467	\$1,185,290,238
670	Turtle, bog (<i>Clemmys muhlenbergii</i>) - U.S.A. (GA, NC, SC, TN, VA)	SAT	\$47,456	\$1,185,337,694
671	Goldenrod, Houghton's (<i>Solidago houghtonii</i>) - Wherever found	T	\$47,300	\$1,185,384,994
672	Beauty, Harper's (<i>Harperocallis flava</i>) - Wherever found	E	\$47,200	\$1,185,432,194
673	Spider, spruce-fir moss (<i>Microhexura montivaga</i>) - Wherever found	E	\$47,004	\$1,185,479,198
674	Snail, Chittenango ovate amber (<i>Succinea chittenangoensis</i>) - Wherever found	T	\$47,000	\$1,185,526,198
675	Millerbird, Nihoa (old world warbler) (<i>Acrocephalus familiaris kingi</i>) - Wherever found	E	\$46,747	\$1,185,572,945
676	Ayenia, Texas (<i>Ayenia limitaris</i>) - Wherever found	E	\$46,700	\$1,185,619,645
677	Squirrel, Carolina northern flying (<i>Glaucomys sabrinus coloratus</i>) - Wherever found	E	\$46,604	\$1,185,666,249
678	Kidneyshell, southern (<i>Ptychobranhus jonesi</i>) - Wherever found	E	\$46,343	\$1,185,712,592
679	Bladderpod, Missouri (<i>Physaria filiformis</i>) - Wherever found	T	\$45,840	\$1,185,758,432
680	Jewelflower, California (<i>Caulanthus californicus</i>) - Wherever found	E	\$45,606	\$1,185,804,038
681	Honohono (<i>Haplostachys haplostachya</i>) - Wherever found	E	\$45,441	\$1,185,849,479
682	Milk-vetch, Fish Slough (<i>Astragalus lentiginosus</i> var. <i>piscinensis</i>) - Wherever found	T	\$45,320	\$1,185,894,799
683	Po`e (<i>Portulaca sclerocarpa</i>) - Wherever found	E	\$44,931	\$1,185,939,730
684	Butterfly, Bartram's hairstreak (<i>Strymon acis bartrami</i>) - Wherever found	E	\$44,900	\$1,185,984,630
685	Orcutt grass, California (<i>Orcuttia californica</i>) - Wherever found	E	\$44,263	\$1,186,028,893

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
686	Mapleleaf, winged (<i>Quadrula fragosa</i>) - U.S.A. (AL-specified portions of the Tennessee River; see 17.85(a)(1))	EXPN	\$44,000	\$1,186,072,893
687	Moccasinshell, Coosa (<i>Medionidus parvulus</i>) - Wherever found	E	\$43,500	\$1,186,116,393
688	Kio'ele (<i>Kadua coriacea</i>) - Wherever found	E	\$43,481	\$1,186,159,874
689	Butterfly, Behren's silverspot (<i>Speyeria zerene behrensii</i>) - Wherever found	E	\$43,180	\$1,186,203,054
690	Cactus, Pariette (<i>Sclerocactus brevispinus</i>) - Wherever found	T	\$43,076	\$1,186,246,130
691	Wallflower, Menzies' (<i>Erysimum menziesii</i>) - Wherever found	E	\$42,510	\$1,186,288,640
692	Butterfly, bay checkerspot (<i>Euphydryas editha bayensis</i>) - Wherever found	T	\$42,500	\$1,186,331,140
693	Butterfly, Uncompahgre fritillary (<i>Boloria acrocneuma</i>) - Wherever found	E	\$42,392	\$1,186,373,532
694	Squirrel, Mount Graham red (<i>Tamiasciurus hudsonicus grahamensis</i>) - Wherever found	E	\$42,346	\$1,186,415,878
695	Ma'oli'oli (<i>Schiedea hawaiiensis</i>) - Wherever found	E	\$42,254	\$1,186,458,132
696	Rosemary, Apalachicola (<i>Conradina glabra</i>) - Wherever found	E	\$42,200	\$1,186,500,332
696	Snail, Iowa Pleistocene (<i>Discus macclintocki</i>) - Wherever found	E	\$42,200	\$1,186,542,532
697	Popolo ku mai (<i>Solanum incompletum</i>) - Wherever found	E	\$42,076	\$1,186,584,608
698	Pigtoe, narrow (<i>Fusconaia escambia</i>) - Wherever found	T	\$41,900	\$1,186,626,508
699	Bean, Choctaw (<i>Villosa choctawensis</i>) - Wherever found	E	\$41,860	\$1,186,668,368
700	Frog, Panamanian golden (<i>Atelopus varius zeteki</i>) - Wherever found	E	\$41,750	\$1,186,710,118
701	Coqui, golden (<i>Eleutherodactylus jasper</i>) - Wherever found	T	\$41,250	\$1,186,751,368
702	Wireweed (<i>Polygonella basiramia</i>) - Wherever found	E	\$41,248	\$1,186,792,616
703	Joint-vetch, Sensitive (<i>Aeschynomene virginica</i>) - Wherever found	T	\$41,246	\$1,186,833,862
704	Aupaka (<i>Isodendrion hosakae</i>) - Wherever found	E	\$41,109	\$1,186,874,971
705	Pigtoe, Georgia (<i>Pleurobema hanleyianum</i>) - Wherever found	E	\$40,699	\$1,186,915,670
706	No Common Name (<i>Stenogyne angustifolia</i> var. <i>angustifolia</i>) - Wherever found	E	\$40,654	\$1,186,956,324
707	Rocksnail, round (<i>Leptoxis ampla</i>) - Wherever found	T	\$40,500	\$1,186,996,824
708	No Common Name (<i>Lipochaeta venosa</i>) - Wherever found	E	\$40,407	\$1,187,037,231
709	Uhi uhi (<i>Mezoneuron kavaiense</i>) - Wherever found	E	\$40,347	\$1,187,077,578
710	Tamarin, golden-rumped (<i>Leontopithecus</i> spp.) - Wherever found	E	\$39,900	\$1,187,117,478
711	Niterwort, Amargosa (<i>Nitrophila mohavensis</i>) - Wherever found	E	\$39,848	\$1,187,157,326
712	Trillium, relict (<i>Trillium reliquum</i>) - Wherever found	E	\$39,710	\$1,187,197,036
713	Combshell, southern (<i>Epioblasma penita</i>) - Wherever found	E	\$39,625	\$1,187,236,661
714	No Common Name (<i>Silene lanceolata</i>) - Wherever found	E	\$39,399	\$1,187,276,060

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
715	No Common Name (<i>Tetramolopium arenarium</i>) - Wherever found	E	\$39,375	\$1,187,315,435
716	No Common Name (<i>Silene hawaiiensis</i>) - Wherever found	T	\$39,175	\$1,187,354,610
717	Cactus, San Rafael (<i>Pediocactus despainii</i>) - Wherever found	E	\$38,400	\$1,187,393,010
718	`Anunu (<i>Sicyos macrophyllus</i>) - Wherever found	E	\$38,140	\$1,187,431,150
719	Cactus, Acuna (<i>Echinomastus erectocentrus</i> var. <i>acunensis</i>) - Wherever found	E	\$38,000	\$1,187,469,150
720	No Common Name (<i>Festuca hawaiiensis</i>) - Wherever found	E	\$37,914	\$1,187,507,064
721	Yellow-faced bee, Hawaiian (<i>Hylaeus kuakea</i>) - Wherever found	E	\$37,808	\$1,187,544,872
722	Darter, amber (<i>Percina antesella</i>) - Wherever found	E	\$37,600	\$1,187,582,472
723	Darter, Cumberland (<i>Etheostoma susanae</i>) - Wherever found	E	\$37,504	\$1,187,619,976
724	Darter, relict (<i>Etheostoma chienense</i>) - Wherever found	E	\$37,000	\$1,187,656,976
725	Dudleya, Verity's (<i>Dudleya verityi</i>) - Wherever found	T	\$36,614	\$1,187,693,590
726	Bladderpod, White Bluffs (<i>Physaria douglasii</i> ssp. <i>tuplashensis</i>) - Wherever found	T	\$36,600	\$1,187,730,190
727	Yellow-faced bee, anthricinan (<i>Hylaeus anthracinus</i>) - Wherever found	E	\$36,590	\$1,187,766,780
728	Rattleweed, hairy (<i>Baptisia arachnifera</i>) - Wherever found	E	\$36,500	\$1,187,803,280
729	Lead-plant, Crenulate (<i>Amorpha crenulata</i>) - Wherever found	E	\$36,200	\$1,187,839,480
729	Torreya, Florida (<i>Torreya taxifolia</i>) - Wherever found	E	\$36,200	\$1,187,875,680
730	Gilia, Monterey (<i>Gilia tenuiflora</i> ssp. <i>arenaria</i>) - Wherever found	E	\$36,195	\$1,187,911,875
731	Darter, fountain (<i>Etheostoma fonticola</i>) - Wherever found	E	\$36,074	\$1,187,947,949
732	Shiner, blue (<i>Cyprinella caerulea</i>) - Wherever found	T	\$35,800	\$1,187,983,749
733	Kidneyshell, fluted (<i>Ptychobranchus subtentum</i>) - Wherever found	E	\$35,507	\$1,188,019,256
734	Sturgeon, Beluga (<i>Huso huso</i>) - Wherever found	T	\$35,403	\$1,188,054,659
735	Ambrosia, south Texas (<i>Ambrosia cheiranthifolia</i>) - Wherever found	E	\$35,400	\$1,188,090,059
736	Mussel, oyster (<i>Epioblasma capsaeformis</i>) - Wherever found; Except where listed as Experimental Populations	E	\$35,091	\$1,188,125,150
737	Sturgeon, Shovelnose (<i>Scaphirhynchus platyrhynchus</i>) - Wherever found	SAT	\$34,850	\$1,188,160,000
738	Wild buckwheat, clay-loving (<i>Eriogonum pelinophilum</i>) - Wherever found	E	\$34,814	\$1,188,194,814
739	Kangaroo rat, Morro Bay (<i>Dipodomys heermanni morroensis</i>) - Wherever found	E	\$34,632	\$1,188,229,446
740	Pupfish, Comanche Springs (<i>Cyprinodon elegans</i>) - Wherever found	E	\$34,559	\$1,188,264,005
741	Springfish, Hiko White River (<i>Crenichthys baileyi grandis</i>) - Wherever found	E	\$34,504	\$1,188,298,509
742	Rush-pea, slender (<i>Hoffmannseggia tenella</i>) - Wherever found	E	\$34,500	\$1,188,333,009
743	Haha (<i>Cyanea shipmanii</i>) - Wherever found	E	\$34,385	\$1,188,367,394
744	Ladies-tresses, Navasota (<i>Spiranthes parksii</i>) - Wherever found	E	\$34,300	\$1,188,401,694

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
744	Moccasinshell, Ochlockonee (<i>Medionidus simpsonianus</i>) - Wherever found	E	\$34,300	\$1,188,435,994
745	Damselfly, flying earwig Hawaiian (<i>Megalagrion nesiotes</i>) - Wherever found	E	\$34,067	\$1,188,470,061
746	Tree snail, Newcomb's (<i>Newcombia cumingi</i>) - Wherever found	E	\$34,030	\$1,188,504,091
747	No Common Name (<i>Osmoxylon mariannense</i>) - Wherever found	E	\$33,720	\$1,188,537,811
748	Thistle, Suisun (<i>Cirsium hydrophilum</i> var. <i>hydrophilum</i>) - Wherever found	E	\$33,348	\$1,188,571,159
749	Madtom, Neosho (<i>Noturus placidus</i>) - Wherever found	T	\$33,322	\$1,188,604,481
750	Pinkroot, gentian (<i>Spigelia gentianoides</i>) - Wherever found	E	\$33,270	\$1,188,637,751
751	Cactus, black lace (<i>Echinocereus reichenbachii</i> var. <i>albertii</i>) - Wherever found	E	\$33,236	\$1,188,670,987
752	Plum, scrub (<i>Prunus geniculata</i>) - Wherever found	E	\$33,200	\$1,188,704,187
753	Dace, Laurel (<i>Chrosomus saylori</i>) - Wherever found	E	\$33,100	\$1,188,737,287
754	Milk-vetch, Jesup's (<i>Astragalus robbinsii</i> var. <i>jesupi</i>) - Wherever found	E	\$33,053	\$1,188,770,340
755	Snail, Lanai tree (<i>Partulina semicarinata</i>) - Wherever found	E	\$33,016	\$1,188,803,356
756	Turtle, Central American river (<i>Dermatemys mawii</i>) - Wherever found	E	\$32,902	\$1,188,836,258
757	Fox, Santa Catalina Island (<i>Urocyon littoralis catalinae</i>) - Wherever found	T	\$32,870	\$1,188,869,128
758	Beargrass, Britton's (<i>Nolina brittoniana</i>) - Wherever found	E	\$32,700	\$1,188,901,828
759	Springfish, White River (<i>Crenichthys baileyi baileyi</i>) - Wherever found	E	\$32,530	\$1,188,934,358
760	Dace, Kendall Warm Springs (<i>Rhinichthys osculus thermalis</i>) - Wherever found	E	\$32,513	\$1,188,966,871
761	Elktoe, Cumberland (<i>Alasmidonta atropurpurea</i>) - Wherever found	E	\$32,280	\$1,188,999,151
762	Bonamia, Florida (<i>Bonamia grandiflora</i>) - Wherever found	T	\$32,200	\$1,189,031,351
763	Fern, Aleutian shield (<i>Polystichum aleuticum</i>) - Wherever found	E	\$32,000	\$1,189,063,351
764	Butterfly, Florida leafwing (<i>Anaea troglodyta floridalis</i>) - Wherever found	E	\$31,900	\$1,189,095,251
765	Tarplant, Santa Cruz (<i>Holocarpha macradenia</i>) - Wherever found	T	\$31,620	\$1,189,126,871
766	Mustard, Penland alpine fen (<i>Eutrema penlandii</i>) - Wherever found	T	\$31,465	\$1,189,158,336
767	Beardtongue, Parachute (<i>Penstemon debilis</i>) - Wherever found	T	\$31,297	\$1,189,189,633
768	Pentachaeta, Lyon's (<i>Pentachaeta lyonii</i>) - Wherever found	E	\$30,792	\$1,189,220,425
769	Aster, Florida golden (<i>Chrysopsis floridana</i>) - Wherever found	E	\$30,700	\$1,189,251,125
770	Nui, haha (<i>Cyanea horrida</i>) - Wherever found	E	\$30,645	\$1,189,281,770
771	Manzanita, Ione (<i>Arctostaphylos myrtifolia</i>) - Wherever found	T	\$30,431	\$1,189,312,201
772	Ischaemum, Hilo (<i>Ischaemum byrone</i>) - Wherever found	E	\$30,393	\$1,189,342,594
773	Haha (<i>Cyanea maritae</i>) - Wherever found	E	\$30,340	\$1,189,372,934
774	Milkweed, Welsh's (<i>Asclepias welshii</i>) - Wherever found	T	\$30,100	\$1,189,403,034

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
775	Meadowfoam, large-flowered woolly (<i>Limnanthes pumila</i> ssp. <i>grandiflora</i>) - Wherever found	E	\$30,022	\$1,189,433,056
776	Bluecurls, Hidden Lake (<i>Trichostema austromontanum</i> ssp. <i>compactum</i>) - Wherever found	T	\$30,000	\$1,189,463,056
776	Iguana, Cuban ground (<i>Cyclura nubila nubila</i>) - (<i>Cyclura nubila nubila</i>)	T	\$30,000	\$1,189,493,056
776	Ipomopsis, Holy Ghost (<i>Ipomopsis sancti-spiritus</i>) - Wherever found	E	\$30,000	\$1,189,523,056
777	Butterfly, Schaus swallowtail (<i>Heraclides aristodemus ponceanus</i>) - Wherever found	E	\$29,918	\$1,189,552,974
778	Hoawa (<i>Pittosporum hawaiiense</i>) - Wherever found	E	\$29,891	\$1,189,582,865
779	Shrimp, Kentucky cave (<i>Palaemonias ganteri</i>) - Wherever found	E	\$29,802	\$1,189,612,667
780	Salamander, Shenandoah (<i>Plethodon shenandoah</i>) - Wherever found	E	\$29,600	\$1,189,642,267
781	No Common Name (<i>Bonamia menziesii</i>) - Wherever found	E	\$29,373	\$1,189,671,640
782	Pitcher-plant, mountain sweet (<i>Sarracenia rubra</i> ssp. <i>jonesii</i>) - Wherever found	E	\$29,254	\$1,189,700,894
783	Haha (<i>Cyanea hamatiflora</i> ssp. <i>hamatiflora</i>) - Wherever found	E	\$29,076	\$1,189,729,970
784	Wawae`iole (<i>Huperzia mannii</i>) - Wherever found	E	28820	\$1,189,758,790
785	Pa`iniu (<i>Astelia waialealae</i>) - Wherever found	E	\$28,689	\$1,189,787,479
786	Nanu (<i>Gardenia remyi</i>) - Wherever found	E	\$28,685	\$1,189,816,164
787	Ohai (<i>Sesbania tomentosa</i>) - Wherever found	E	\$28,533	\$1,189,844,697
788	Turtle, Plymouth Redbelly (<i>Pseudemys rubriventris bangsi</i>) - Wherever found	E	\$28,371	\$1,189,873,068
789	Woodrat, Key Largo (<i>Neotoma floridana smalli</i>) - Wherever found	E	\$28,339	\$1,189,901,407
790	Ebonyshell, round (<i>Fusconaia rotulata</i>) - Wherever found	E	\$28,300	\$1,189,929,707
791	Spineflower, Ben Lomond (<i>Chorizanthe pungens</i> var. <i>hartwegiana</i>) - Wherever found	E	\$28,229	\$1,189,957,936
792	Nohoanu (<i>Geranium multiflorum</i>) - Wherever found	E	\$28,114	\$1,189,986,050
793	Arrowhead, bunched (<i>Sagittaria fasciculata</i>) - Wherever found	E	\$27,939	\$1,190,013,989
794	Penny-cress, Kneeland Prairie (<i>Thlaspi californicum</i>) - Wherever found	E	\$27,700	\$1,190,041,689
795	Ko`oko`olau (<i>Bidens micrantha</i> ssp. <i>ctenophylla</i>) - Wherever found	E	\$27,659	\$1,190,069,348
796	No Common Name (<i>Nervilia jacksoniae</i>) - Wherever found	T	\$27,554	\$1,190,096,902
797	Fairy shrimp, Riverside (<i>Streptocephalus woottoni</i>) - Wherever found	E	\$27,464	\$1,190,124,366
798	`Aiakeakua, popolo (<i>Solanum sandwicense</i>) - Wherever found	E	\$27,450	\$1,190,151,816
799	Haha (<i>Cyanea kunthiana</i>) - Wherever found	E	\$27,219	\$1,190,179,035
800	Barberry, island (<i>Berberis pinnata</i> ssp. <i>insularis</i>) - Wherever found	E	\$26,945	\$1,190,205,980

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
801	Purple Cat's paw (=Purple Cat's paw pearlymussel) (<i>Epioblasma obliquata obliquata</i>) - Wherever found; Except where listed as Experimental Populations	E	\$26,903	\$1,190,232,883
802	Nohoanu (<i>Geranium hanaense</i>) - Wherever found	E	\$26,898	\$1,190,259,781
803	Butterfly, Miami Blue (<i>Cyclargus</i> (=Hemiargus) <i>thomasi bethunebakeri</i>) - Wherever found	E	\$26,700	\$1,190,286,481
804	Sandshell, Southern (<i>Hamiota australis</i>) - Wherever found	T	\$26,661	\$1,190,313,142
805	Poppy, Sacramento prickly (<i>Argemone pleiakantha</i> ssp. <i>pinnatisecta</i>) - Wherever found	E	\$26,620	\$1,190,339,762
806	`Aiea (<i>Nothocestrum latifolium</i>) - Wherever found	E	\$26,614	\$1,190,366,376
806	No Common Name (<i>Microlepidia strigosa</i> var. <i>mauiensis</i>) - Wherever found	E	\$26,614	\$1,190,392,990
806	No Common Name (<i>Wikstroemia villosa</i>) - Wherever found	E	\$26,614	\$1,190,419,604
807	Reedgrass, Maui (<i>Calamagrostis expansa</i>) - Wherever found	E	\$26,553	\$1,190,446,157
808	Rice rat (<i>Oryzomys palustris natator</i>) - lower FL Keys	E	\$26,420	\$1,190,472,577
809	Sea turtle, olive ridley (<i>Lepidochelys olivacea</i>) - Breeding colony populations on Pacific coast of Mexico	E	\$26,406	\$1,190,498,983
810	Rose-mallow, Neches River (<i>Hibiscus dasycalyx</i>) - Wherever found	T	\$26,271	\$1,190,525,254
811	Sunburst, San Joaquin adobe (<i>Pseudobahia peirsonii</i>) - Wherever found	T	\$26,243	\$1,190,551,497
812	Cycladenia, Jones (<i>Cycladenia humilis</i> var. <i>jonesii</i>) - Wherever found	T	\$26,050	\$1,190,577,547
813	Snail, Lanai tree (<i>Partulina variabilis</i>) - Wherever found	E	\$25,918	\$1,190,603,465
814	Pearlymussel, littlewing (<i>Pegias fabula</i>) - Wherever found	E	\$25,775	\$1,190,629,240
815	Milk-vetch, heliotrope (<i>Astragalus montii</i>) - Wherever found	T	\$25,750	\$1,190,654,990
816	Ambrosia, San Diego (<i>Ambrosia pumila</i>) - Wherever found	E	\$25,563	\$1,190,680,553
817	Beetle, Comal Springs riffle (<i>Heterelmis comalensis</i>) - Wherever found	E	\$25,400	\$1,190,705,953
818	Butterweed, Layne's (<i>Senecio layneae</i>) - Wherever found	T	\$25,200	\$1,190,731,153
818	Sandlace (<i>Polygonella myriophylla</i>) - Wherever found	E	\$25,200	\$1,190,756,353
819	Butterfly, San Bruno elfin (<i>Callophrys mossii bayensis</i>) - Wherever found	E	\$25,180	\$1,190,781,533
820	Fern, American hart's-tongue (<i>Asplenium scolopendrium</i> var. <i>americanum</i>) - Wherever found	T	\$24,979	\$1,190,806,512
821	Grass, Colusa (<i>Neostapfia colusana</i>) - Wherever found	T	\$24,943	\$1,190,831,455
822	Haha (<i>Cyanea stictophylla</i>) - Wherever found	E	\$24,844	\$1,190,856,299
823	Otter, giant (<i>Pteronura brasiliensis</i>) - South America	E	\$24,745	\$1,190,881,044

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
824	Madtom, yellowfin (<i>Noturus flavipinnis</i>) - Wherever found, except where listed as an experimental population	T	\$24,591	\$1,190,905,635
825	Pu`uka`a (<i>Cyperus trachysanthos</i>) - Wherever found	E	\$24,574	\$1,190,930,209
826	Buckwheat, Ione (incl. Irish Hill) (<i>Eriogonum apricum</i> (incl. var. <i>prostratum</i>)) - Wherever found	E	\$24,500	\$1,190,954,709
827	Sandwort, Marsh (<i>Arenaria paludicola</i>) - Wherever found	E	\$24,448	\$1,190,979,157
828	Heau (<i>Exocarpos luteolus</i>) - Wherever found	E	\$24,276	\$1,191,003,433
829	Springsnail, Three Forks (<i>Pyrgulopsis trivialis</i>) - Wherever found	E	\$24,113	\$1,191,027,546
830	Bird's beak, palmate-bracted (<i>Cordylanthus palmatus</i>) - Wherever found	E	\$24,090	\$1,191,051,636
831	Clubshell, southern (<i>Pleurobema decisum</i>) - Wherever found	E	\$24,066	\$1,191,075,702
832	Cactus, Bakersfield (<i>Opuntia treleasei</i>) - Wherever found	E	\$24,000	\$1,191,099,702
833	No Common Name (<i>Cyrtandra nanawaleensis</i>) - Wherever found	E	\$23,833	\$1,191,123,535
834	`Oha wai (<i>Clermontia peleana</i>) - Wherever found	E	\$23,821	\$1,191,147,356
835	Cactus, Siler pincushion (<i>Pediocactus</i> (=Echinocactus,=Utahia) <i>sileri</i>) - Wherever found	T	\$23,650	\$1,191,171,006
836	Madtom, smoky (<i>Noturus baileyi</i>) - Wherever found	E	\$23,569	\$1,191,194,575
837	Bear-poppy, Dwarf (<i>Arctomecon humilis</i>) - Wherever found	E	\$23,516	\$1,191,218,091
838	Owl's-clover, fleshy (<i>Castilleja campestris</i> ssp. <i>succulenta</i>) - Wherever found	T	\$23,503	\$1,191,241,594
839	Darter, slackwater (<i>Etheostoma boschungii</i>) - Wherever found	T	\$23,500	\$1,191,265,094
840	Cactus, Mesa Verde (<i>Sclerocactus mesae-verdae</i>) - Wherever found	T	\$23,295	\$1,191,288,389
841	Milk-vetch, Ventura Marsh (<i>Astragalus pycnostachyus</i> var. <i>lanosissimus</i>) - Wherever found	E	\$23,153	\$1,191,311,542
842	No Common Name (<i>Nesogenes rotensis</i>) - Wherever found	E	\$23,142	\$1,191,334,684
843	Shiner, sharpnose (<i>Notropis oxyrhynchus</i>) - Wherever found	E	\$23,100	\$1,191,357,784
844	Cactus, Fickeisen plains (<i>Pediocactus peeblesianus fickeiseniae</i>) - Wherever found	E	\$23,000	\$1,191,380,784
844	Shiner, smalleye (<i>Notropis buccula</i>) - Wherever found	E	\$23,000	\$1,191,403,784
845	Aplokating-palaoan (<i>Psychotria malaspinae</i>) - Wherever found	E	\$22,987	\$1,191,426,771
846	Mallow, Gierisch (<i>Sphaeralcea gierischii</i>) - Wherever found	E	\$22,950	\$1,191,449,721
847	No Common Name (<i>Silene perlmanii</i>) - Wherever found	E	\$22,876	\$1,191,472,597
848	Mint, scrub (<i>Dicerandra frutescens</i>) - Wherever found	E	\$22,700	\$1,191,495,297
849	Rhododendron, Chapman (<i>Rhododendron chapmanii</i>) - Wherever found	E	\$22,600	\$1,191,517,897
849	Snakeroot (<i>Eryngium cuneifolium</i>) - Wherever found	E	\$22,600	\$1,191,540,497

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
850	Tarplant, Otay (<i>Deinandra</i> (=Hemizonia) <i>conjugens</i>) - Wherever found	T	\$22,506	\$1,191,563,003
851	Butterfly, Mariana wandering (<i>Vagrans egistina</i>) - Wherever found	E	\$22,450	\$1,191,585,453
852	Spineflower, Scotts Valley (<i>Chorizanthe robusta</i> var. <i>hartwegii</i>) - Wherever found	E	\$22,442	\$1,191,607,895
853	Glade cress, Kentucky (<i>Leavenworthia exigua laciniata</i>) - Wherever found	T	\$22,210	\$1,191,630,105
854	Seal, bearded (<i>Erignathus barbatus nauticus</i>) - Okhotsk DPS	T	\$22,186	\$1,191,652,291
855	No Common Name (<i>Vigna o-wahuensis</i>) - Wherever found	E	\$22,163	\$1,191,674,454
856	Alani (<i>Melicope adscendens</i>) - Wherever found	E	\$22,157	\$1,191,696,611
857	Nohoanu (<i>Geranium kauaiense</i>) - Wherever found	E	\$22,154	\$1,191,718,765
858	Skink, Slevin's (<i>Emoia slevini</i>) - Wherever found	E	\$22,141	\$1,191,740,906
859	Ambersnail, Kanab (<i>Oxyloma haydeni kanabensis</i>) - Wherever found	E	\$22,004	\$1,191,762,910
860	Iguana, Mona ground (<i>Cyclura stejnegeri</i>) - Wherever found	T	\$22,000	\$1,191,784,910
860	Snail, noonday (<i>Mesodon clarki nantahala</i>) - Wherever found	T	\$22,000	\$1,191,806,910
861	Clubshell (<i>Pleurobema clava</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$21,904	\$1,191,828,814
862	Milk-vetch, Coachella Valley (<i>Astragalus lentiginosus</i> var. <i>coachellae</i>) - Wherever found	E	\$21,870	\$1,191,850,684
863	Woolly-star, Santa Ana River (<i>Eriastrum densifolium</i> ssp. <i>sanctorum</i>) - Wherever found	E	\$21,853	\$1,191,872,537
864	Cactus, Arizona hedgehog (<i>Echinocereus triglochidiatus</i> var. <i>arizonicus</i>) - Wherever found	E	\$21,402	\$1,191,893,939
865	Pigtoe, tapered (<i>Fusconaia burkei</i>) - Wherever found	T	\$21,361	\$1,191,915,300
866	Naucorid, Ash Meadows (<i>Ambrysus amargosus</i>) - Wherever found	T	\$21,324	\$1,191,936,624
867	Oha wai (<i>Clermontia lindseyana</i>) - Wherever found	E	\$21,239	\$1,191,957,863
868	Lehua makanoë (<i>Lysimachia daphnoides</i>) - Wherever found	E	\$21,008	\$1,191,978,871
869	Lupine, scrub (<i>Lupinus aridorum</i>) - Wherever found	E	\$21,000	\$1,191,999,871
869	Madtom, pygmy (<i>Noturus stanauli</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(t)(1)(i))	EXPN	\$21,000	\$1,192,020,871
870	No Common Name (<i>Asplenium unisorum</i>) - Wherever found	E	\$20,952	\$1,192,041,823
871	No Common Name (<i>Lobelia niihauensis</i>) - Wherever found	E	\$20,857	\$1,192,062,680
872	Spinymussel, Tar River (<i>Elliptio steinstansana</i>) - Wherever found	E	\$20,819	\$1,192,083,499
873	Nehe (<i>Lipochaeta lobata</i> var. <i>leptophylla</i>) - Wherever found	E	\$20,812	\$1,192,104,311
874	Orcutt grass, slender (<i>Orcuttia tenuis</i>) - Wherever found	T	\$20,610	\$1,192,124,921

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
875	Sculpin, Grotto (<i>Cottus specus</i>) - Wherever found	E	\$20,500	\$1,192,145,421
876	Hala pepe (<i>Pleomele forbesii</i>) - Wherever found	E	\$20,487	\$1,192,165,908
877	`Aiea (<i>Nothocestrum breviflorum</i>) - Wherever found	E	\$20,439	\$1,192,186,347
878	Ceanothus, coyote (<i>Ceanothus ferrisae</i>) - Wherever found	E	\$20,390	\$1,192,206,737
878	Dudleya, Santa Clara Valley (<i>Dudleya setchellii</i>) - Wherever found	E	\$20,390	\$1,192,227,127
879	Rock-cress, Braun's (<i>Arabis perstellata</i>) - Wherever found	E	\$20,334	\$1,192,247,461
880	No Common Name (<i>Platydesma cornuta</i> var. <i>decurrens</i>) - Wherever found	E	\$20,331	\$1,192,267,792
881	Rabbitsfoot, rough (<i>Quadrula cylindrica strigillata</i>) - Wherever found	E	\$20,322	\$1,192,288,114
882	Haha (<i>Cyanea calycina</i>) - Wherever found	E	\$20,316	\$1,192,308,430
883	Seablite, California (<i>Suaeda californica</i>) -	E	\$20,261	\$1,192,328,691
884	No Common Name (<i>Tetramolopium lepidotum</i> ssp. <i>lepidotum</i>) - Wherever found	E	\$20,236	\$1,192,348,927
885	`Oha wai (<i>Clermontia pyrularia</i>) - Wherever found	E	\$20,110	\$1,192,369,037
886	`Anaunau (<i>Lepidium arbuscula</i>) - Wherever found	E	\$20,103	\$1,192,389,140
886	Alani (<i>Melicope saint-johnii</i>) - Wherever found	E	\$20,103	\$1,192,409,243
887	Rock cress, Shale barren (<i>Arabis serotina</i>) - Wherever found	E	\$20,096	\$1,192,429,339
888	Salamander, Texas blind (<i>Typhlomolge rathbuni</i>) - Wherever found	E	\$20,000	\$1,192,449,339
889	Alani (<i>Melicope puberula</i>) - Wherever found	E	\$19,994	\$1,192,469,333
890	Crayfish, Shasta (<i>Pacifastacus fortis</i>) - Wherever found	E	\$19,900	\$1,192,489,233
891	Wire-lettuce, Malheur (<i>Stephanomeria malheurensis</i>) - Wherever found	E	\$19,851	\$1,192,509,084
892	Cactus, Knowlton's (<i>Pediocactus knowltonii</i>) - Wherever found	E	\$19,846	\$1,192,528,930
893	Alani (<i>Melicope christophersenii</i>) - Wherever found	E	\$19,809	\$1,192,548,739
894	Sandalwood, Lanai (= `iliahii) (<i>Santalum haleakalae</i> var. <i>lanaiense</i>) - Wherever found	E	\$19,803	\$1,192,568,542
895	Salamander, Cheat Mountain (<i>Plethodon nettingi</i>) - Wherever found	T	\$19,716	\$1,192,588,258
896	Gumplant, Ash Meadows (<i>Grindelia fraxinipratensis</i>) - Wherever found	T	\$19,648	\$1,192,607,906
897	Centaury, spring-loving (<i>Centaurium namophilum</i>) - Wherever found	T	\$19,636	\$1,192,627,542
898	No Common Name (<i>Schiedea hookeri</i>) - Wherever found	E	\$19,465	\$1,192,647,007
899	Tui chub, Mohave (<i>Gila bicolor</i> ssp. <i>mohavensis</i>) - Wherever found	E	\$19,340	\$1,192,666,347
900	Milkvetch, Osterhout (<i>Astragalus osterhoutii</i>) - Wherever found	E	\$19,312	\$1,192,685,659
901	Alani (<i>Melicope knudsenii</i>) - Wherever found	E	\$19,275	\$1,192,704,934
902	Haha (<i>Cyanea hamatiflora</i> ssp. <i>carlsonii</i>) - Wherever found	E	\$19,248	\$1,192,724,182
903	`aku (<i>Cyanea tritomantha</i>) - Wherever found	E	\$19,211	\$1,192,743,393
904	Milk-vetch, Holmgren (<i>Astragalus holmgreniorum</i>) - Wherever found	E	\$19,067	\$1,192,762,460

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
905	Nukupuu, Maui (<i>Hemignathus affinis</i>) - Wherever found	E	\$19,019	\$1,192,781,479
906	Beetle, [no common name] (<i>Rhadine infernalis</i>) - Wherever found	E	\$19,000	\$1,192,800,479
907	Jewelflower, Metcalf Canyon (<i>Streptanthus albidus</i> ssp. <i>albidus</i>) - Wherever found	E	\$18,990	\$1,192,819,469
908	Bedstraw, El Dorado (<i>Galium californicum</i> ssp. <i>sierrae</i>) - Wherever found	E	\$18,950	\$1,192,838,419
909	Pearlymussel, dromedary (<i>Dromus dromas</i>) - Wherever found; Except where listed as Experimental Populations	E	\$18,791	\$1,192,857,210
910	Kamakahala (<i>Labordia lydgatei</i>) - Wherever found	E	\$18,521	\$1,192,875,731
911	Cactus, Peebles Navajo (<i>Pediocactus peeblesianus</i> var. <i>peeblesianus</i>) - Wherever found	E	\$18,500	\$1,192,894,231
911	Cactus, Sneed pincushion (<i>Coryphantha sneedii</i> var. <i>sneedii</i>) - Wherever found	E	\$18,500	\$1,192,912,731
911	Cobana negra (<i>Stahlia monosperma</i>) - Wherever found	T	\$18,500	\$1,192,931,231
911	Milk-vetch, Shivwits (<i>Astragalus ampullarioides</i>) - Wherever found	E	\$18,500	\$1,192,949,731
912	Bladderpod, Dudley Bluffs (<i>Lesquerella congesta</i>) - Wherever found	T	\$18,467	\$1,192,968,198
913	Prairie-clover, leafy (<i>Dalea foliosa</i>) - Wherever found	E	\$18,434	\$1,192,986,632
914	Orcutt grass, San Joaquin (<i>Orcuttia inaequalis</i>) - Wherever found	T	\$18,418	\$1,193,005,050
915	Monkeyface, Appalachian (pearlymussel) (<i>Quadrula sparsa</i>) - Wherever found	E	\$18,400	\$1,193,023,450
916	Clarkia, Springville (<i>Clarkia springvillensis</i>) - Wherever found	T	\$18,361	\$1,193,041,811
917	Orcutt grass, Sacramento (<i>Orcuttia viscida</i>) - Wherever found	E	\$18,350	\$1,193,060,161
918	Twinpod, Dudley Bluffs (<i>Physaria obcordata</i>) - Wherever found	T	\$18,320	\$1,193,078,481
919	Clover, showy Indian (<i>Trifolium amoenum</i>) - Wherever found	E	\$18,200	\$1,193,096,681
920	Amphipod, Kauai cave (<i>Spelaeorchestia koloana</i>) - Wherever found	E	\$18,129	\$1,193,114,810
920	Spider, Kauai cave wolf or pe'e pe'e maka 'ole (<i>Adelocosa anops</i>) - Wherever found	E	\$18,129	\$1,193,132,939
921	Turtle, Alabama red-bellied (<i>Pseudemys alabamensis</i>) - Wherever found	E	\$18,050	\$1,193,150,989
922	No Common Name (<i>Phyllostegia parviflora</i>) - Wherever found	E	\$18,038	\$1,193,169,027
923	Shrimp, California freshwater (<i>Syncaris pacifica</i>) - Wherever found	E	\$17,996	\$1,193,187,023
924	Cactus, Lee pincushion (<i>Coryphantha sneedii</i> var. <i>leei</i>) - Wherever found	T	\$17,959	\$1,193,204,982
925	Snail, Stock Island tree (<i>Orthalicus reses</i> (not incl. <i>nesodryas</i>)) - Wherever found	T	\$17,841	\$1,193,222,823
926	Grass, Tennessee yellow-eyed (<i>Xyris tennesseensis</i>) - Wherever found	E	\$17,839	\$1,193,240,662
927	Snail, Newcomb's (<i>Erinna newcombi</i>) - Wherever found	T	\$17,834	\$1,193,258,496
928	Clubshell, ovate (<i>Pleurobema perovatum</i>) - Wherever found	E	\$17,802	\$1,193,276,298
929	Puma (=cougar), eastern (<i>Puma</i> (=Felis) <i>concolor</i> cougar) - Wherever found	E	\$17,652	\$1,193,293,950

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
930	Cypress, Santa Cruz (<i>Cupressus abramsiana</i>) - Wherever found	T	\$17,610	\$1,193,311,560
931	No Common Name (<i>Eugenia bryanii</i>) - Wherever found	E	\$17,582	\$1,193,329,142
932	Cactus, Key tree (<i>Pilosocereus robinii</i>) - Wherever found	E	\$17,553	\$1,193,346,695
933	Haha (<i>Cyanea marksii</i>) - Wherever found	E	\$17,527	\$1,193,364,222
934	Beetle, American burying (<i>Nicrophorus americanus</i>) - Ex Pop, SW Missouri	EXPN	\$17,500	\$1,193,381,722
934	Polygala, Lewton's (<i>Polygala lewtonii</i>) - Wherever found	E	\$17,500	\$1,193,399,222
935	Heartleaf, dwarf-flowered (<i>Hexastylis naniflora</i>) - Wherever found	T	\$17,463	\$1,193,416,685
936	Evening-primrose, San Benito (<i>Camissonia benitensis</i>) - Wherever found	T	\$17,400	\$1,193,434,085
937	Phacelia, North Park (<i>Phacelia formosula</i>) - Wherever found	E	\$17,347	\$1,193,451,432
938	No Common Name (<i>Tetramolopium rockii</i>) - Wherever found	T	\$17,309	\$1,193,468,741
939	Lupine, Nipomo Mesa (<i>Lupinus nipomensis</i>) - Wherever found	E	\$17,210	\$1,193,485,951
940	`Awikiwiki (<i>Canavalia pubescens</i>) - Wherever found	E	\$17,194	\$1,193,503,145
941	`Akoko (<i>Euphorbia kuwaleana</i>) - Wherever found	E	\$17,025	\$1,193,520,170
942	Iris, dwarf lake (<i>Iris lacustris</i>) - Wherever found	T	\$17,000	\$1,193,537,170
942	Isopod, Lee County cave (<i>Lirceus usdagalun</i>) - Wherever found	E	\$17,000	\$1,193,554,170
943	Mouse, Key Largo cotton (<i>Peromyscus gossypinus allapaticola</i>) - Wherever found	E	\$16,829	\$1,193,570,999
944	Damselfly, blackline Hawaiian (<i>Megalagrion nigrohamatum nigrolineatum</i>) - Wherever found	E	\$16,812	\$1,193,587,811
945	Darter, yellowcheek (<i>Etheostoma moorei</i>) - Wherever found	E	\$16,700	\$1,193,604,511
946	No Common Name (<i>Geocarpon minimum</i>) - Wherever found	T	\$16,673	\$1,193,621,184
947	Lizard, St. Croix ground (<i>Ameiva polops</i>) - Wherever found	E	\$16,500	\$1,193,637,684
948	Rattlesnake, New Mexican ridge-nosed (<i>Crotalus willardi obscurus</i>) - Wherever found	T	\$16,450	\$1,193,654,134
949	Finch, Nihoa (honeycreeper) (<i>Telespyza ultima</i>) - Wherever found	E	\$16,327	\$1,193,670,461
950	Lo`ulu (<i>Pritchardia maideniana</i>) - Wherever found	E	\$16,300	\$1,193,686,761
951	Reed-mustard, clay (<i>Schoenocrambe argillacea</i>) - Wherever found	T	\$16,292	\$1,193,703,053
952	White-eye, Rota bridled (<i>Zosterops rotensis</i>) - Wherever found	E	\$16,272	\$1,193,719,325
953	Silversword, Mauna Loa (=Ka'u) (<i>Argyroxiphium kauense</i>) - Wherever found	E	\$16,253	\$1,193,735,578
954	Nukupuu, Kauai (<i>Hemignathus hanapepe</i>) - Wherever found	E	\$16,248	\$1,193,751,826
955	Dace, Clover Valley speckled (<i>Rhinichthys osculus oligoporus</i>) - Wherever found	E	\$16,228	\$1,193,768,054
956	Dudleya, Santa Cruz Island (<i>Dudleya nesiotica</i>) - Wherever found	T	\$16,151	\$1,193,784,205
957	Shrimp, anchialine pool (<i>Procaris hawaiana</i>) - Wherever found	E	\$16,101	\$1,193,800,306

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
958	Grasshopper, Zayante band-winged (<i>Trimerotropis infantilis</i>) - Wherever found	E	\$16,050	\$1,193,816,356
959	Goetzea, beautiful (<i>Goetzea elegans</i>) - Wherever found	E	\$16,000	\$1,193,832,356
959	Phlox, Texas trailing (<i>Phlox nivalis</i> ssp. <i>texensis</i>) - Wherever found	E	\$16,000	\$1,193,848,356
959	Sculpin, pygmy (<i>Cottus paulus</i> (=pygmaeus)) - Wherever found	T	\$16,000	\$1,193,864,356
960	Milk-vetch, Mancos (<i>Astragalus humillimus</i>) - Wherever found	E	\$15,800	\$1,193,880,156
961	Sturgeon, Alabama (<i>Scaphirhynchus suttkusi</i>) - Wherever found	E	\$15,699	\$1,193,895,855
962	Thrush, Molokai (<i>Myadestes lanaiensis</i> <i>rutha</i>) - Wherever found	E	\$15,680	\$1,193,911,535
963	Thornmint, San Diego (<i>Acanthomintha ilicifolia</i>) - Wherever found	T	\$15,584	\$1,193,927,119
964	Monkey-flower, Michigan (<i>Mimulus michiganensis</i>) - Wherever found	E	\$15,540	\$1,193,942,659
965	Gambusia, Pecos (<i>Gambusia nobilis</i>) - Wherever found	E	\$15,500	\$1,193,958,159
966	Halomtano, Berenghenas (<i>Solanum guamense</i>) - Wherever found	E	\$15,489	\$1,193,973,648
967	Mucket, orangenacre (<i>Lampsilis perovalis</i>) - Wherever found	T	\$15,403	\$1,193,989,051
968	No Common Name (<i>Asplenium dielmannii</i>) - Wherever found	E	\$15,396	\$1,194,004,447
968	No Common Name (<i>Asplenium dielpallidum</i>) - Wherever found	E	\$15,396	\$1,194,019,843
969	Salamander, Barton Springs (<i>Eurycea sosorum</i>) - Wherever found	E	\$15,300	\$1,194,035,143
970	Damselfly, oceanic Hawaiian (<i>Megalagrion oceanicum</i>) - Wherever found	E	\$15,291	\$1,194,050,434
971	Hala pepe (<i>Pleomele hawaiiensis</i>) - Wherever found	E	\$15,260	\$1,194,065,694
972	Beetle, delta green ground (<i>Elaphrus viridis</i>) - Wherever found	T	\$15,144	\$1,194,080,838
973	Fly, Hawaiian picture-wing (<i>Drosophila hemipeza</i>) - Wherever found	E	\$15,131	\$1,194,095,969
974	Chub, spotfin (<i>Erimonax monachus</i>) - U.S.A. (AL, TN-specified portions of Shoal Creek; see 17.84(m)(1)(ii))	EXPN	\$15,000	\$1,194,110,969
974	Elimia, lacy (snail) (<i>Elimia crenatella</i>) - Wherever found	T	\$15,000	\$1,194,125,969
974	Madtom, yellowfin (<i>Noturus flavipinnis</i>) - U.S.A. (TN-specified portions of the Tellico River; see 17.84(e)(1)(ii))	EXPN	\$15,000	\$1,194,140,969
974	Sunflower, Pecos (=puzzle, =paradox) (<i>Helianthus paradoxus</i>) - Wherever found	T	\$15,000	\$1,194,155,969
975	No Common Name (<i>Schiedea kauaiensis</i>) - Wherever found	E	\$14,999	\$1,194,170,968
976	Sunray, Ash Meadows (<i>Enceliopsis nudicaulis</i> var. <i>corrugata</i>) - Wherever found	T	\$14,956	\$1,194,185,924
977	Koki'o (<i>Kokia kauaiensis</i>) - Wherever found	E	\$14,918	\$1,194,200,842
978	Snake, Atlantic salt marsh (<i>Nerodia clarkii</i> <i>taeniata</i>) - Wherever found	T	\$14,871	\$1,194,215,713
979	Blazingstar, Ash Meadows (<i>Mentzelia leucophylla</i>) - Wherever found	T	\$14,848	\$1,194,230,561
979	Ivesia, Ash Meadows (<i>Ivesia kingii</i> var. <i>eremica</i>) - Wherever found	T	\$14,848	\$1,194,245,409
979	Milk-vetch, Ash meadows (<i>Astragalus phoenix</i>) - Wherever found	T	\$14,848	\$1,194,260,257

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
980	No Common Name (<i>Polyscias bisattenuata</i>) - Wherever found	E	\$14,839	\$1,194,275,096
981	Silverbush, Blodgett's (<i>Argythamnia blodgettii</i>) - Wherever found	T	\$14,805	\$1,194,289,901
982	Rock-cress, Hoffmann's (<i>Arabis hoffmannii</i>) - Wherever found	E	\$14,685	\$1,194,304,586
983	Haha (<i>Cyanea st.-johnii</i>) - Wherever found	E	\$14,599	\$1,194,319,185
984	Damselfly, Rota blue (<i>Ischnura luta</i>) - Wherever found	E	\$14,580	\$1,194,333,765
985	Darter, goldline (<i>Percina aurolineata</i>) - Wherever found	T	\$14,570	\$1,194,348,335
986	Koki`o (<i>Kokia drynarioides</i>) - Wherever found	E	\$14,545	\$1,194,362,880
987	Phacelia, island (<i>Phacelia insularis</i> ssp. <i>insularis</i>) - Wherever found	E	\$14,485	\$1,194,377,365
988	Naupaka, dwarf (<i>Scaevola coriacea</i>) - Wherever found	E	\$14,454	\$1,194,391,819
989	[no common name] Snail (<i>Eua zebrina</i>) - Wherever found	E	\$14,424	\$1,194,406,243
990	Pearlymussel, birdwing (<i>Lemiox rimosus</i>) - Wherever found; Except where listed as Experimental Populations	E	\$14,320	\$1,194,420,563
991	Creeper, Oahu (<i>Paroreomyza maculata</i>) - Wherever found	E	\$14,220	\$1,194,434,783
992	Kolea (<i>Myrsine knudsenii</i>) - Wherever found	E	\$14,075	\$1,194,448,858
993	Paintbrush, Tiburon (<i>Castilleja affinis</i> ssp. <i>neglecta</i>) - Wherever found	E	\$14,070	\$1,194,462,928
994	No Common Name (<i>Lobelia oahuensis</i>) - Wherever found	E	\$14,041	\$1,194,476,969
995	Boa, Virgin Islands tree (<i>Epicrates monensis granti</i>) - Wherever found	E	\$14,000	\$1,194,490,969
995	Mint, Lakela's (<i>Dicerandra immaculata</i>) - Wherever found	E	\$14,000	\$1,194,504,969
995	Monkeyface, Cumberland (pearlymussel) (<i>Quadrula intermedia</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$14,000	\$1,194,518,969
995	Quillwort, mat-forming (<i>Isoetes tegetiformans</i>) - Wherever found	E	14000	\$1,194,532,969
995	Rosemary, Etonia (<i>Conradina etonia</i>) - Wherever found	E	\$14,000	\$1,194,546,969
996	Reed-mustard, shrubby (<i>Schoenocrambe suffrutescens</i>) - Wherever found	E	\$13,993	\$1,194,560,962
997	No Common Name (<i>Xylosma crenatum</i>) - Wherever found	E	\$13,975	\$1,194,574,937
998	Fatmucket, Arkansas (<i>Lampsilis powellii</i>) - Wherever found	T	\$13,945	\$1,194,588,882
999	A`e (<i>Zanthoxylum oahuense</i>) - Wherever found	E	\$13,925	\$1,194,602,807
1000	Hoawa (<i>Pittosporum halophilum</i>) - Wherever found	E	\$13,700	\$1,194,616,507
1001	Dudleya, marcescent (<i>Dudleya cymosa</i> ssp. <i>marcescens</i>) - Wherever found	T	\$13,684	\$1,194,630,191
1002	Dudleya, Conejo (<i>Dudleya abramsii</i> ssp. <i>parva</i>) - Wherever found	T	\$13,651	\$1,194,643,842
1003	Fly, Hawaiian picture-wing (<i>Drosophila heteroneura</i>) - Wherever found	E	\$13,610	\$1,194,657,452
1004	`Akoko (<i>Euphorbia haelecleana</i>) - Wherever found	E	\$13,594	\$1,194,671,046
1005	Spurge, Garber's (<i>Chamaesyce garberi</i>) - Wherever found	T	\$13,590	\$1,194,684,636

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1006	Makou (<i>Peucedanum sandwicense</i>) - Wherever found	T	\$13,574	\$1,194,698,210
1007	Fly, Hawaiian picture-wing (<i>Drosophila digressa</i>) - Wherever found	E	\$13,566	\$1,194,711,776
1008	Combshell, upland (<i>Epioblasma metastriata</i>) - Wherever found	E	\$13,519	\$1,194,725,295
1009	Lo`ulu (<i>Pritchardia remota</i>) - Wherever found	E	\$13,468	\$1,194,738,763
1010	No Common Name (<i>Platanthera holochila</i>) - Wherever found	E	\$13,463	\$1,194,752,226
1011	Alani (<i>Melicope degeneri</i>) - Wherever found	E	\$13,451	\$1,194,765,677
1012	Kuahiwi laukahi (<i>Plantago hawaiiensis</i>) - Wherever found	E	\$13,426	\$1,194,779,103
1013	Lichen, rock gnome (<i>Gymnoderma lineare</i>) - Wherever found	E	13300	\$1,194,792,403
1014	Purple cat's paw (=purple cat's paw pearlymussel) (<i>Epioblasma obliquata obliquata</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$13,202	\$1,194,805,605
1015	Meadowrue, Cooley's (<i>Thalictrum cooleyi</i>) - Wherever found	E	\$13,200	\$1,194,818,805
1016	Goldenrod, Short's (<i>Solidago shortii</i>) - Wherever found	E	\$13,100	\$1,194,831,905
1017	Kopiko (<i>Psychotria grandiflora</i>) - Wherever found	E	\$13,081	\$1,194,844,986
1018	Shrimp, anchialine pool (<i>Vetericaris chaceorum</i>) - Wherever found	E	\$13,059	\$1,194,858,045
1019	Orcutt grass, hairy (<i>Orcuttia pilosa</i>) - Wherever found	E	\$13,043	\$1,194,871,088
1020	Higuero de sierra (<i>Crescentia portoricensis</i>) - Wherever found	E	\$13,000	\$1,194,884,088
1020	Mandrill (<i>Mandrillus</i> (=Papio) sphinx) - Wherever found	E	\$13,000	\$1,194,897,088
1020	No Common Name (<i>Aristida chaseae</i>) - Wherever found	E	\$13,000	\$1,194,910,088
1020	No Common Name (<i>Lyonia truncata</i> var. proctorii) - Wherever found	E	\$13,000	\$1,194,923,088
1020	Palo de rosa (<i>Ottoschulzia rhodoxylon</i>) - Wherever found	E	\$13,000	\$1,194,936,088
1020	Pelos del diablo (<i>Aristida portoricensis</i>) - Wherever found	E	\$13,000	\$1,194,949,088
1020	Sedge, golden (<i>Carex lutea</i>) - Wherever found	E	\$13,000	\$1,194,962,088
1021	No Common Name (<i>Schiedea membranacea</i>) - Wherever found	E	\$12,971	\$1,194,975,059
1022	Nani wai`ale`ale (<i>Viola kauaiensis</i> var. wahiawaensis) - Wherever found	E	\$12,961	\$1,194,988,020
1023	Shark, Scalloped Hammerhead (<i>Sphyrna lewini</i>) - Eastern Pacific DPS	E	\$12,907	\$1,195,000,927
1024	Gerardia, sandplain (<i>Agalinis acuta</i>) - Wherever found	E	\$12,890	\$1,195,013,817
1025	No Common Name (<i>Polyscias flynnii</i>) - Wherever found	E	\$12,862	\$1,195,026,679
1026	No Common Name (<i>Tinospora homosepala</i>) - Wherever found	E	\$12,847	\$1,195,039,526
1027	No Common Name (<i>Phyllanthus saffordii</i>) - Wherever found	E	\$12,826	\$1,195,052,352
1028	Fern, pendant kihi (<i>Adenophorus periens</i>) - Wherever found	E	\$12,820	\$1,195,065,172
1029	Alani (<i>Melicope zahlbruckneri</i>) - Wherever found	E	\$12,728	\$1,195,077,900

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1030	No Common Name (<i>Phyllostegia warshaueri</i>) - Wherever found	E	\$12,704	\$1,195,090,604
1031	Dwarf-flax, Marin (<i>Hesperolinon congestum</i>) - Wherever found	T	\$12,670	\$1,195,103,274
1032	Paintbrush, soft-leaved (<i>Castilleja mollis</i>) - Wherever found	E	\$12,659	\$1,195,115,933
1033	Morning-glory, Stebbins' (<i>Calystegia stebbinsii</i>) - Wherever found	E	\$12,650	\$1,195,128,583
1034	Turtle, ringed map (<i>Graptemys oculifera</i>) - Wherever found	T	\$12,648	\$1,195,141,231
1035	Tuctoria, Greene's (<i>Tuctoria greenei</i>) - Wherever found	E	\$12,625	\$1,195,153,856
1036	No Common Name (<i>Sanicula purpurea</i>) - Wherever found	E	\$12,596	\$1,195,166,452
1037	A`e (<i>Zanthoxylum dipetalum</i> var. <i>tomentosum</i>) - Wherever found	E	\$12,545	\$1,195,178,997
1038	Ground-plum, Guthrie's (=Pyne's) (<i>Astragalus bibullatus</i>) - Wherever found	E	\$12,528	\$1,195,191,525
1039	Alani (<i>Melicope mucronulata</i>) - Wherever found	E	\$12,502	\$1,195,204,027
1040	Ass, Asian wild (<i>Equus hemionus</i>) - Wherever found	E	\$12,500	\$1,195,216,527
1040	Drill (<i>Mandrillus</i> (=Papio) <i>leucophaeus</i>) - Wherever found	E	\$12,500	\$1,195,229,027
1040	Mangabey, white-collared (<i>Cercocebus torquatus</i>) - Wherever found	E	\$12,500	\$1,195,241,527
1040	Monkey, black colobus (<i>Colobus satanas</i>) - Wherever found	E	\$12,500	\$1,195,254,027
1040	Monkey, Preuss' red colobus (<i>Procolobus</i> (=Colobus) <i>preussi</i> (=badius p.)) - Wherever found	E	\$12,500	\$1,195,266,527
1040	Monkey, red-eared nose-spotted (<i>Cercopithecus erythrotis</i>) - Wherever found	E	\$12,500	\$1,195,279,027
1040	Phacelia, clay (<i>Phacelia argillacea</i>) - Wherever found	E	\$12,500	\$1,195,291,527
1041	Kiponapona (<i>Phyllostegia racemosa</i>) - Wherever found	E	\$12,491	\$1,195,304,018
1042	`Akoko (<i>Euphorbia remyi</i> var. <i>remyi</i>) - Wherever found	E	\$12,490	\$1,195,316,508
1043	Riversnail, Anthony's (<i>Athearnia anthonyi</i>) - Wherever found; Except where listed as Experimental Populations	E	\$12,400	\$1,195,328,908
1044	Snake, copperbelly water (<i>Nerodia erythrogaster neglecta</i>) - Indiana north of 40 degrees north latitude, Michigan, Ohio	T	\$12,386	\$1,195,341,294
1045	Lauhilihi (<i>Schiedea stellarioides</i>) - Wherever found	E	\$12,317	\$1,195,353,611
1046	Jaguarundi, Gulf Coast (<i>Herpailurus</i> (=Felis) <i>yagouaroundi cacomitli</i>) - Wherever found	E	\$12,300	\$1,195,365,911
1046	Rockcress, Georgia (<i>Arabis georgiana</i>) - Wherever found	T	\$12,300	\$1,195,378,211
1047	Finch, Laysan (honeycreeper) (<i>Telespyza cantans</i>) - Wherever found	E	\$12,271	\$1,195,390,482
1048	Gardenia (=Na`u), Hawaiian (<i>Gardenia brighamii</i>) - Wherever found	E	\$12,216	\$1,195,402,698
1049	Manzanita, Santa Rosa Island (<i>Arctostaphylos confertiflora</i>) - Wherever found	E	\$12,159	\$1,195,414,857
1050	Ceanothus, Pine Hill (<i>Ceanothus roderickii</i>) - Wherever found	E	\$12,150	\$1,195,427,007

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1051	Ha`iwale (<i>Cyrtandra filipes</i>) - Wherever found	E	\$12,140	\$1,195,439,147
1052	Sea turtle, green (<i>Chelonia mydas</i>) - Southwest Pacific DPS	T	\$12,111	\$1,195,451,258
1053	Madtom, chunky (<i>Noturus crypticus</i>) - Wherever found	E	\$12,100	\$1,195,463,358
1054	Shark, Scalloped Hammerhead (<i>Sphyrna lewini</i>) - Central and Southwest Atlantic DPS	T	\$12,078	\$1,195,475,436
1055	Kamakahala (<i>Labordia tinifolia</i> var. <i>wahiawaensis</i>) - Wherever found	E	\$12,058	\$1,195,487,494
1056	Kauila (<i>Colubrina oppositifolia</i>) - Wherever found	E	\$12,041	\$1,195,499,535
1057	Buckwheat, scrub (<i>Eriogonum longifolium</i> var. <i>gnaphalifolium</i>) - Wherever found	T	\$12,000	\$1,195,511,535
1057	No Common Name (<i>Vernonia proctorii</i>) - Wherever found	E	\$12,000	\$1,195,523,535
1058	Bush-mallow, Santa Cruz Island (<i>Malacothamnus fasciculatus</i> var. <i>nesioticus</i>) - Wherever found	E	\$11,985	\$1,195,535,520
1059	`Awikiwiki (<i>Canavalia molokaiensis</i>) - Wherever found	E	\$11,928	\$1,195,547,448
1060	Na`ena`e (<i>Dubautia pauciflorula</i>) - Wherever found	E	\$11,927	\$1,195,559,375
1061	Ho`awa (<i>Pittosporum napaliense</i>) - Wherever found	E	\$11,845	\$1,195,571,220
1062	No Common Name (<i>Cyrtandra wagneri</i>) - Wherever found	E	\$11,843	\$1,195,583,063
1063	Butterfly, Myrtle's silverspot (<i>Speyeria zerene myrtleae</i>) - Wherever found	E	\$11,770	\$1,195,594,833
1064	No Common Name (<i>Phyllostegia stachyoides</i>) - Wherever found	E	\$11,764	\$1,195,606,597
1065	Fly, Hawaiian picture-wing (<i>Drosophila musaphilia</i>) - Wherever found	E	\$11,740	\$1,195,618,337
1065	Fly, Hawaiian picture-wing (<i>Drosophila sharpi</i>) - Wherever found	E	\$11,740	\$1,195,630,077
1066	Haha (<i>Cyanea lobata</i>) - Wherever found	E	\$11,739	\$1,195,641,816
1067	No Common Name (<i>Platydesma remyi</i>) - Wherever found	E	\$11,731	\$1,195,653,547
1068	Iliau, dwarf (<i>Wilkesia hobdyi</i>) - Wherever found	E	\$11,727	\$1,195,665,274
1069	Yellow-faced bee, Hawaiian (<i>Hylaeus longiceps</i>) - Wherever found	E	\$11,601	\$1,195,676,875
1070	Dace, Ash Meadows speckled (<i>Rhinichthys osculus nevadensis</i>) - Wherever found	E	\$11,578	\$1,195,688,453
1071	Sunflower, San Mateo woolly (<i>Eriophyllum latilobum</i>) - Wherever found	E	\$11,520	\$1,195,699,973
1072	No Common Name (<i>Phyllostegia renovans</i>) - Wherever found	E	\$11,476	\$1,195,711,449
1073	Buttercup, autumn (<i>Ranunculus aestivalis</i> (= <i>acriiformis</i>)) - Wherever found	E	\$11,450	\$1,195,722,899
1074	Spineflower, Howell's (<i>Chorizanthe howellii</i>) - Wherever found	E	\$11,440	\$1,195,734,339
1075	Bush-clover, prairie (<i>Lespedeza leptostachya</i>) - Wherever found	T	\$11,385	\$1,195,745,724
1076	Alani (<i>Melicope quadrangularis</i>) - Wherever found	E	\$11,372	\$1,195,757,096
1076	No Common Name (<i>Lysimachia venosa</i>) - Wherever found	E	\$11,372	\$1,195,768,468
1077	Kamakahala (<i>Labordia tinifolia</i> var. <i>lanaiensis</i>) - Wherever found	E	\$11,143	\$1,195,779,611

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1078	Nehe (<i>Lipochaeta micrantha</i>) - Wherever found	E	\$11,133	\$1,195,790,744
1079	Beardtongue, Penland (<i>Penstemon penlandii</i>) - Wherever found	E	\$11,056	\$1,195,801,800
1080	Lo`ulu (<i>Pritchardia viscosa</i>) - Wherever found	E	\$11,053	\$1,195,812,853
1081	`Aiea (<i>Nothocestrum peltatum</i>) - Wherever found	E	\$11,044	\$1,195,823,897
1081	No Common Name (<i>Phyllostegia waimeae</i>) - Wherever found	E	\$11,044	\$1,195,834,941
1082	Kolea (<i>Myrsine mezii</i>) - Wherever found	E	\$11,033	\$1,195,845,974
1083	Gilia, Hoffmann's slender-flowered (<i>Gilia tenuiflora</i> ssp. <i>hoffmannii</i>) - Wherever found	E	\$11,030	\$1,195,857,004
1084	Clarkia, Presidio (<i>Clarkia franciscana</i>) - Wherever found	E	\$11,020	\$1,195,868,024
1085	Leather flower, Alabama (<i>Clematis socialis</i>) - Wherever found	E	\$11,000	\$1,195,879,024
1086	No Common Name (<i>Diplazium molokaiense</i>) - Wherever found	E	\$10,993	\$1,195,890,017
1087	Manzanita, Morro (<i>Arctostaphylos morroensis</i>) - Wherever found	T	\$10,961	\$1,195,900,978
1088	No Common Name (<i>Remya montgomeryi</i>) - Wherever found	E	\$10,933	\$1,195,911,911
1088	No Common Name (<i>Viola helenae</i>) - Wherever found	E	\$10,933	\$1,195,922,844
1089	Skipper, Laguna Mountains (<i>Pyrgus ruralis lagunae</i>) - Wherever found	E	\$10,920	\$1,195,933,764
1090	Alani (<i>Melicope haupeensis</i>) - Wherever found	E	\$10,913	\$1,195,944,677
1090	Haha (<i>Cyanea rivularis</i>) - Wherever found	E	\$10,913	\$1,195,955,590
1090	Haha (<i>Cyanea undulata</i>) - Wherever found	E	\$10,913	\$1,195,966,503
1090	Haiwale (<i>Cyrtandra paliku</i>) - Wherever found	E	\$10,913	\$1,195,977,416
1091	No Common Name (<i>Stenogyne campanulata</i>) - Wherever found	E	\$10,882	\$1,195,988,298
1091	No Common Name (<i>Stenogyne kealiae</i>) - Wherever found	E	\$10,882	\$1,195,999,180
1092	Woodrat, riparian (=San Joaquin Valley) (<i>Neotoma fuscipes riparia</i>) - Wherever found	E	\$10,870	\$1,196,010,050
1093	Ha`iwale (<i>Cyrtandra giffardii</i>) - Wherever found	E	\$10,864	\$1,196,020,914
1094	No Common Name (<i>Delissea rhytidosperma</i>) - Wherever found	E	\$10,862	\$1,196,031,776
1094	Olulu (<i>Brighamia insignis</i>) - Wherever found	E	\$10,862	\$1,196,042,638
1095	Hau kuahiwi (<i>Hibiscadelphus giffardianus</i>) - Wherever found	E	\$10,827	\$1,196,053,465
1096	Pigtoe, heavy (<i>Pleurobema taitianum</i>) - Wherever found	E	\$10,824	\$1,196,064,289
1097	No Common Name (<i>Lysimachia pendens</i>) - Wherever found	E	\$10,813	\$1,196,075,102
1098	No Common Name (<i>Delissea undulata</i>) - Wherever found	E	\$10,789	\$1,196,085,891
1099	Rush-rose, island (<i>Helianthemum greenei</i>) - Wherever found	T	\$10,779	\$1,196,096,670
1100	Nehe (<i>Lipochaeta waimeaensis</i>) - Wherever found	E	\$10,762	\$1,196,107,432

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1101	Fly, Hawaiian picture-wing (<i>Drosophila obatai</i>) - Wherever found	E	\$10,726	\$1,196,118,158
1101	Fly, Hawaiian picture-wing (<i>Drosophila ochrobasis</i>) - Wherever found	E	\$10,726	\$1,196,128,884
1101	Fly, Hawaiian picture-wing (<i>Drosophila tarphytrichia</i>) - Wherever found	E	\$10,726	\$1,196,139,610
1102	Holei (<i>Ochrosia haleakalae</i>) - Wherever found	E	\$10,711	\$1,196,150,321
1103	Haha (<i>Cyanea remyi</i>) - Wherever found	E	\$10,704	\$1,196,161,025
1104	No Common Name (<i>Phyllostegia hispida</i>) - Wherever found	E	\$10,685	\$1,196,171,710
1105	No Common Name (<i>Neraudia sericea</i>) - Wherever found	E	\$10,633	\$1,196,182,343
1106	Pigtoe, shiny (<i>Fusconaia cor</i>) - Wherever found; Except where listed as Experimental Populations	E	\$10,620	\$1,196,192,963
1107	Pauoa (<i>Ctenitis squamigera</i>) - Wherever found	E	10618	\$1,196,203,581
1108	Sunflower, whorled (<i>Helianthus verticillatus</i>) -	E	\$10,600	\$1,196,214,181
1109	No Common Name (<i>Silene alexandri</i>) - Wherever found	E	\$10,573	\$1,196,224,754
1110	Haha (<i>Cyanea procera</i>) - Wherever found	E	\$10,553	\$1,196,235,307
1110	Kamakahala (<i>Labordia triflora</i>) - Wherever found	E	\$10,553	\$1,196,245,860
1110	No Common Name (<i>Lysimachia maxima</i>) - Wherever found	E	\$10,553	\$1,196,256,413
1111	Aupaka (<i>Isodendrion laurifolium</i>) - Wherever found	E	\$10,552	\$1,196,266,965
1111	Aupaka (<i>Isodendrion longifolium</i>) - Wherever found	T	\$10,552	\$1,196,277,517
1112	Salamander, desert slender (<i>Batrachoseps aridus</i>) - Wherever found	E	\$10,550	\$1,196,288,067
1113	Vetch, Hawaiian (<i>Vicia menziesii</i>) - Wherever found	E	\$10,534	\$1,196,298,601
1114	Koki'o, Cooke's (<i>Kokia cookei</i>) - Wherever found	E	\$10,502	\$1,196,309,103
1114	Na'ena'e (<i>Dubautia plantaginea</i> ssp. <i>humilis</i>) - Wherever found	E	\$10,502	\$1,196,319,605
1115	No Common Name (<i>Eugenia woodburyana</i>) - Wherever found	E	\$10,500	\$1,196,330,105
1115	Vole, Florida salt marsh (<i>Microtus pennsylvanicus</i> <i>dukecampbelli</i>) - Wherever found	E	\$10,500	\$1,196,340,605
1116	Lo'ulu (<i>Pritchardia schattaueri</i>) - Wherever found	E	\$10,483	\$1,196,351,088
1117	No Common Name (<i>Phyllostegia floribunda</i>) - Wherever found	E	\$10,432	\$1,196,361,520
1118	No Common Name (<i>Schiedea viscosa</i>) - Wherever found	E	\$10,412	\$1,196,371,932
1119	Phacelia, DeBeque (<i>Phacelia submutica</i>) - Wherever found	T	\$10,409	\$1,196,382,341
1120	Hau kuahiwi (<i>Hibiscadelphus hualalaiensis</i>) - Wherever found	E	\$10,398	\$1,196,392,739
1121	Chub, Chihuahuahua (<i>Gila nigrescens</i>) - Wherever found	T	\$10,390	\$1,196,403,129
1122	Kuawawaenuhu (<i>Schiedea lychnoides</i>) - Wherever found	E	\$10,361	\$1,196,413,490
1123	Kaulu (<i>Pteralyxia kauaiensis</i>) - Wherever found	E	\$10,351	\$1,196,423,841
1124	Koholapehu (<i>Dubautia latifolia</i>) - Wherever found	E	\$10,330	\$1,196,434,171

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1125	Ha`iwale (Cyrtandra tintinnabula) - Wherever found	E	\$10,320	\$1,196,444,491
1126	Chub, Yaqui (Gila purpurea) - Wherever found	E	\$10,300	\$1,196,454,791
1126	Pocketbook, speckled (Lampsilis streckeri) - Wherever found	E	\$10,300	\$1,196,465,091
1127	Pua `ala (Brighamia rockii) - Wherever found	E	\$10,292	\$1,196,475,383
1128	No Common Name (Cyperus pennatiformis) - Wherever found	E	\$10,207	\$1,196,485,590
1129	No Common Name (Schiedea verticillata) - Wherever found	E	\$10,204	\$1,196,495,794
1130	Amphipod, Peck's cave (Stygobromus (=Stygonectes) pecki) - Wherever found	E	\$10,200	\$1,196,505,994
1130	Beetle, Comal Springs dryopid (Stygoparnus comalensis) - Wherever found	E	\$10,200	\$1,196,516,194
1130	Gooseberry, Miccosukee (Ribes echinellum) - Wherever found	T	\$10,200	\$1,196,526,394
1130	Sand-verbena, large-fruited (Abronia macrocarpa) - Wherever found	E	\$10,200	\$1,196,536,594
1131	No Common Name (Phyllostegia pilosa) - Wherever found	E	\$10,176	\$1,196,546,770
1132	Allocarya, Calistoga (Plagiobothrys strictus) - Wherever found	E	\$10,120	\$1,196,556,890
1132	Bluegrass, Napa (Poa napensis) - Wherever found	E	\$10,120	\$1,196,567,010
1132	Larkspur, Baker's (Delphinium bakeri) - Wherever found	E	\$10,120	\$1,196,577,130
1133	Petrel, Bermuda (Pterodroma cahow) - Wherever found	E	\$10,118	\$1,196,587,248
1134	Ha`iwale (Cyrtandra kaulantha) - Wherever found	E	\$10,115	\$1,196,597,363
1135	Sucker, Zuni bluehead (Catostomus discobolus yarrowi) - Wherever found	E	\$10,100	\$1,196,607,463
1136	`Akoko (Euphorbia rockii) - Wherever found	E	\$10,093	\$1,196,617,556
1137	Kolea (Myrsine juddii) - Wherever found	E	\$10,081	\$1,196,627,637
1138	No Common Name (Stenogyne kauaulaensis) - Wherever found	E	\$10,015	\$1,196,637,652
1139	Sandwort, Cumberland (Arenaria cumberlandensis) - Wherever found	E	\$10,006	\$1,196,647,658
1140	Shiner, palezone (Notropis albizonatus) - Wherever found	E	\$10,004	\$1,196,657,662
1141	Bear, grizzly (Ursus arctos horribilis) - Bitterroot Grizzly Bear Nonessential Experimental Population Area	EXPN	\$10,000	\$1,196,667,662
1141	Beetle, Hungerford's crawling water (Brychius hungerfordi) - Wherever found	E	\$10,000	\$1,196,677,662
1141	Crayfish, Benton County cave (Cambarus aculabrum) - Wherever found	E	\$10,000	\$1,196,687,662
1141	Gecko, Monito (Sphaerodactylus micropithecus) - Wherever found	E	\$10,000	\$1,196,697,662
1141	Goldenrod, white-haired (Solidago albopilosa) - Wherever found	T	\$10,000	\$1,196,707,662
1141	No Common Name (Acropora rudis) - Wherever found	T	\$10,000	\$1,196,717,662
1141	Palo de nigua (Cornutia obovata) - Wherever found	E	\$10,000	\$1,196,727,662
1141	Salamander, San Marcos (Eurycea nana) - Wherever found	T	\$10,000	\$1,196,737,662
1142	Kohe malama malama o kanaloa (Kanaloa kahoolawensis) - Wherever found	E	\$9,998	\$1,196,747,660

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1143	Alani (<i>Melicope hiiakae</i>) - Wherever found	E	\$9,995	\$1,196,757,655
1144	Haha (<i>Cyanea grimesiana</i> ssp. <i>grimesiana</i>) - Wherever found	E	\$9,984	\$1,196,767,639
1145	No Common Name (<i>Lobelia koolauensis</i>) - Wherever found	E	\$9,961	\$1,196,777,600
1146	`aku`aku (<i>Cyanea platyphylla</i>) - Wherever found	E	\$9,941	\$1,196,787,541
1147	Larkspur, yellow (<i>Delphinium luteum</i>) - Wherever found	E	\$9,900	\$1,196,797,441
1147	Pawpaw, four-petal (<i>Asimina tetramera</i>) - Wherever found	E	\$9,900	\$1,196,807,341
1148	No Common Name (<i>Schiedea diffusa</i> ssp. <i>macraei</i>) - Wherever found	E	\$9,896	\$1,196,817,237
1149	Schiedea, Diamond Head (<i>Schiedea adamantis</i>) - Wherever found	E	\$9,863	\$1,196,827,100
1150	Opuhe (<i>Urera kaalae</i>) - Wherever found	E	\$9,852	\$1,196,836,952
1151	Haha (<i>Cyanea truncata</i>) - Wherever found	E	\$9,832	\$1,196,846,784
1151	No Common Name (<i>Lobelia monostachya</i>) - Wherever found	E	\$9,832	\$1,196,856,616
1152	No Common Name (<i>Phyllostegia mannii</i>) - Wherever found	E	\$9,823	\$1,196,866,439
1153	Haha (<i>Cyanea dunbariae</i>) - Wherever found	E	\$9,792	\$1,196,876,231
1154	Pamakani (<i>Tetramolopium capillare</i>) - Wherever found	E	\$9,770	\$1,196,886,001
1155	Ha`iwale (<i>Cyrtandra subumbellata</i>) - Wherever found	E	\$9,757	\$1,196,895,758
1156	`Oha wai (<i>Clermontia oblongifolia</i> ssp. <i>brevipes</i>) - Wherever found	E	\$9,741	\$1,196,905,499
1157	Darter, bayou (<i>Etheostoma rubrum</i>) - Wherever found	T	\$9,685	\$1,196,915,184
1158	Haha (<i>Cyanea obtusa</i>) - Wherever found	E	\$9,640	\$1,196,924,824
1158	Haha (<i>Cyanea profuga</i>) - Wherever found	E	\$9,640	\$1,196,934,464
1158	Popolo (<i>Cyanea solanacea</i>) - Wherever found	E	\$9,640	\$1,196,944,104
1159	Mesa-mint, Otay (<i>Pogogyne nudiuscula</i>) - Wherever found	E	\$9,636	\$1,196,953,740
1160	Snowbells, Texas (<i>Styrax texanus</i>) - Wherever found	E	\$9,600	\$1,196,963,340
1161	Bat, Pacific sheath-tailed (<i>Emballonura semicaudata rotensis</i>) - Wherever found	E	\$9,573	\$1,196,972,913
1162	Haha (<i>Cyanea purpurellifolia</i>) - Wherever found	E	\$9,539	\$1,196,982,452
1163	Damselfly, crimson Hawaiian (<i>Megalagrion leptodemas</i>) - Wherever found	E	\$9,510	\$1,196,991,962
1164	Coqui, Llanero (<i>Eleutherodactylus juanariveroi</i>) - Wherever found	E	\$9,500	\$1,197,001,462
1164	Darter, watercress (<i>Etheostoma nuchale</i>) - Wherever found	E	\$9,500	\$1,197,010,962
1164	Flannelbush, Pine Hill (<i>Fremontodendron californicum</i> ssp. <i>decumbens</i>) - Wherever found	E	\$9,500	\$1,197,020,462
1164	Pearlshell, Alabama (<i>Margaritifera marrianae</i>) - Wherever found	E	\$9,500	\$1,197,029,962
1164	Pigeon, Puerto Rican plain (<i>Columba inornata wetmorei</i>) - Wherever found	E	\$9,500	\$1,197,039,462
1165	Haiwale (<i>Cyrtandra gracilis</i>) - Wherever found	E	\$9,488	\$1,197,048,950

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1166	Hau kuahiwi (<i>Hibiscadelphus woodii</i>) - Wherever found	E	\$9,475	\$1,197,058,425
1166	Kauai hau kuahiwi (<i>Hibiscadelphus distans</i>) - Wherever found	E	\$9,475	\$1,197,067,900
1166	No Common Name (<i>Phyllostegia knudsenii</i>) - Wherever found	E	\$9,475	\$1,197,077,375
1166	No Common Name (<i>Phyllostegia wawrana</i>) - Wherever found	E	\$9,475	\$1,197,086,850
1167	No Common Name (<i>Amaranthus brownii</i>) - Wherever found	E	\$9,462	\$1,197,096,312
1168	No Common Name (<i>Doryopteris angelica</i>) - Wherever found	E	\$9,457	\$1,197,105,769
1169	Avens, spreading (<i>Geum radiatum</i>) - Wherever found	E	\$9,450	\$1,197,115,219
1170	No Common Name (<i>Keysseria (=Lagenifera) erici</i>) - Wherever found	E	\$9,448	\$1,197,124,667
1170	No Common Name (<i>Keysseria (=Lagenifera) helenae</i>) - Wherever found	E	\$9,448	\$1,197,134,115
1171	Darter, diamond (<i>Crystallaria cincotta</i>) - Wherever found	E	\$9,435	\$1,197,143,550
1172	No Common Name (<i>Gouania meyenii</i>) - Wherever found	E	\$9,432	\$1,197,152,982
1173	Naenae (<i>Dubautia kalalauensis</i>) - Wherever found	E	\$9,395	\$1,197,162,377
1173	No Common Name (<i>Lysimachia iniki</i>) - Wherever found	E	\$9,395	\$1,197,171,772
1174	No Common Name (<i>Schiedea attenuata</i>) - Wherever found	E	\$9,374	\$1,197,181,146
1175	Aumakua, Palapalai (<i>Dryopteris crinalis</i> var. <i>podosorus</i>) - Wherever found	E	\$9,364	\$1,197,190,510
1176	Haha (<i>Cyanea dolichopoda</i>) - Wherever found	E	\$9,344	\$1,197,199,854
1176	Haha (<i>Cyanea eleeleensis</i>) - Wherever found	E	\$9,344	\$1,197,209,198
1176	Haha (<i>Cyanea kuhihewa</i>) - Wherever found	E	\$9,344	\$1,197,218,542
1176	Naenae (<i>Dubautia kenwoodii</i>) - Wherever found	E	\$9,344	\$1,197,227,886
1176	No Common Name (<i>Kadua st.-johnii</i>) - Wherever found	E	\$9,344	\$1,197,237,230
1177	Lo`ulu (<i>Pritchardia napaliensis</i>) - Wherever found	E	\$9,337	\$1,197,246,567
1178	Kolea (<i>Myrsine linearifolia</i>) - Wherever found	T	\$9,316	\$1,197,255,883
1178	Na`ena`e (<i>Dubautia waialealae</i>) - Wherever found	E	\$9,316	\$1,197,265,199
1178	Panicgrass, Carter's (<i>Panicum fauriei</i> var. <i>carteri</i>) - Wherever found	E	\$9,316	\$1,197,274,515
1179	Aster, Ruth's golden (<i>Pityopsis ruthii</i>) - Wherever found	E	\$9,302	\$1,197,283,817
1180	Haha (<i>Cyanea kolekoleensis</i>) - Wherever found	E	\$9,295	\$1,197,293,112
1180	No Common Name (<i>Polyscias lydgatei</i>) - Wherever found	E	\$9,295	\$1,197,302,407
1181	No Common Name (<i>Schiedea spergulina</i> var. <i>leiopoda</i>) - Wherever found	E	\$9,275	\$1,197,311,682
1182	No Common Name (<i>Lysimachia scopulensis</i>) - Wherever found	E	\$9,244	\$1,197,320,926
1183	Acornshell, southern (<i>Epioblasma othcaloogensis</i>) - Wherever found	E	\$9,242	\$1,197,330,168
1184	`Anunu (<i>Sicyos albus</i>) - Wherever found	E	\$9,231	\$1,197,339,399
1185	`Akoko (<i>Euphorbia eleanoriae</i>) - Wherever found	E	\$9,186	\$1,197,348,585

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1186	Thistle, fountain (<i>Cirsium fontinale</i> var. <i>fontinale</i>) - Wherever found	E	\$9,120	\$1,197,357,705
1187	Curassow, blue-billed (<i>Crax alberti</i>) - Wherever found	E	\$9,100	\$1,197,366,805
1188	`Akoko (<i>Euphorbia remyi</i> var. <i>kauaiensis</i>) - Wherever found	E	\$9,087	\$1,197,375,892
1189	Yellow-faced bee, assimulans (<i>Hylaeus assimulans</i>) - Wherever found	E	\$9,066	\$1,197,384,958
1189	Yellow-faced bee, easy (<i>Hylaeus facilis</i>) - Wherever found	E	\$9,066	\$1,197,394,024
1190	No Common Name (<i>Stenogyne bifida</i>) - Wherever found	E	\$9,055	\$1,197,403,079
1190	No Common Name (<i>Tetramolopium remyi</i>) - Wherever found	E	\$9,055	\$1,197,412,134
1191	No Common Name (<i>Abutilon eremitopetalum</i>) - Wherever found	E	\$9,034	\$1,197,421,168
1192	Pussypaws, Mariposa (<i>Calyptridium pulchellum</i>) - Wherever found	T	\$9,020	\$1,197,430,188
1193	No Common Name (<i>Phyllostegia glabra</i> var. <i>lanaiensis</i>) - Wherever found	E	\$9,016	\$1,197,439,204
1194	Manatee, West African (<i>Trichechus senegalensis</i>) - Wherever found	T	\$9,000	\$1,197,448,204
1194	No Common Name (<i>Schoepfia arenaria</i>) - Wherever found	T	\$9,000	\$1,197,457,204
1194	Springsnail, Phantom (<i>Pyrgulopsis texana</i>) - Wherever found	E	\$9,000	\$1,197,466,204
1195	Popolo (<i>Solanum nelsonii</i>) - Wherever found	E	\$8,985	\$1,197,475,189
1196	No Common Name (<i>Lysimachia lydgatei</i>) - Wherever found	E	\$8,984	\$1,197,484,173
1197	No Common Name (<i>Viola lanaiensis</i>) - Wherever found	E	\$8,956	\$1,197,493,129
1198	Pitcher-plant, green (<i>Sarracenia oreophila</i>) - Wherever found	E	\$8,920	\$1,197,502,049
1199	No Common Name (<i>Phyllostegia velutina</i>) - Wherever found	E	\$8,793	\$1,197,510,842
1200	Checker-mallow, Keck's (<i>Sidalcea keckii</i>) - Wherever found	E	\$8,769	\$1,197,519,611
1201	Cactus, Florida semaphore (<i>Consolea corallicola</i>) - Wherever found	E	\$8,700	\$1,197,528,311
1202	Ha`iwale (<i>Cyrtandra munroi</i>) - Wherever found	E	\$8,662	\$1,197,536,973
1203	No Common Name (<i>Schiedea laui</i>) - Wherever found	E	\$8,641	\$1,197,545,614
1204	Blazingstar, Heller's (<i>Liatris helleri</i>) - Wherever found	T	\$8,600	\$1,197,554,214
1205	Turtle, yellow-blotched map (<i>Graptemys flavimaculata</i>) - Wherever found	T	\$8,549	\$1,197,562,763
1206	Nightjar, Puerto Rican (<i>Caprimulgus noctitherus</i>) - Wherever found	E	\$8,500	\$1,197,571,263
1206	Toad, Puerto Rican crested (<i>Peltophryne lemur</i>) - Wherever found	T	\$8,500	\$1,197,579,763
1207	Fly, Hawaiian picture-wing (<i>Drosophila aglaia</i>) - Wherever found	E	\$8,496	\$1,197,588,259
1208	Alani (<i>Melicope balloui</i>) - Wherever found	E	\$8,478	\$1,197,596,737
1209	Kamakahala (<i>Labordia pumila</i>) - Wherever found	E	\$8,434	\$1,197,605,171
1210	Woodpecker, ivory-billed (<i>Campephilus principalis</i>) - Wherever found	E	\$8,400	\$1,197,613,571
1211	Ha`iwale (<i>Cyrtandra sessilis</i>) - Wherever found	E	\$8,377	\$1,197,621,948
1212	`Ala `ala wai nui (<i>Peperomia subpetiolata</i>) - Wherever found	E	\$8,376	\$1,197,630,324

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1213	Spurge, Hoover's (<i>Chamaesyce hooveri</i>) - Wherever found	T	\$8,315	\$1,197,638,639
1214	Ha`iwale (<i>Cyrtandra polyantha</i>) - Wherever found	E	\$8,314	\$1,197,646,953
1214	Haha (<i>Cyanea pinnatifida</i>) - Wherever found	E	\$8,314	\$1,197,655,267
1215	Lo`ulu (<i>Pritchardia munroi</i>) - Wherever found	E	\$8,313	\$1,197,663,580
1216	Love grass, Fosberg's (<i>Eragrostis fosbergii</i>) - Wherever found	E	\$8,263	\$1,197,671,843
1217	No Common Name (<i>Asplenium dielfalcatum</i>) - Wherever found	E	\$8,256	\$1,197,680,099
1218	Beetle, Coffin Cave mold (<i>Batrisodes texanus</i>) - Wherever found	E	\$8,250	\$1,197,688,349
1219	Haha (<i>Cyanea copelandii</i> ssp. <i>copelandii</i>) - Wherever found	E	\$8,223	\$1,197,696,572
1220	Haha (<i>Cyanea munroi</i>) - Wherever found	E	\$8,173	\$1,197,704,745
1221	`Akoko (<i>Euphorbia deppeana</i>) - Wherever found	E	\$8,164	\$1,197,712,909
1222	No Common Name (<i>Schiedea jacobii</i>) - Wherever found	E	\$8,152	\$1,197,721,061
1223	Heau (<i>Exocarpos menziesii</i>) -	E	\$8,140	\$1,197,729,201
1224	No Common Name (<i>Schiedea spergulina</i> var. <i>spergulina</i>) - Wherever found	T	\$8,124	\$1,197,737,325
1225	No Common Name (<i>Remya kauaiensis</i>) - Wherever found	E	\$8,120	\$1,197,745,445
1226	No Common Name (<i>Lepanthes eltoroensis</i>) - Wherever found	E	\$8,100	\$1,197,753,545
1226	Shiner, beautiful (<i>Cyprinella formosa</i>) - Wherever found	T	\$8,100	\$1,197,761,645
1227	No Common Name (<i>Lysimachia filifolia</i>) - Wherever found	E	\$8,089	\$1,197,769,734
1228	Grass, Solano (<i>Tuctoria mucronata</i>) - Wherever found	E	\$8,074	\$1,197,777,808
1229	Bedstraw, island (<i>Galium buxifolium</i>) - Wherever found	E	\$8,059	\$1,197,785,867
1230	Holei (<i>Ochrosia kilaueaensis</i>) - Wherever found	E	\$8,036	\$1,197,793,903
1231	Hulumoa (<i>Korthalsella degeneri</i>) - Wherever found	E	\$8,002	\$1,197,801,905
1232	Amphianthus, little (<i>Amphianthus pusillus</i>) - Wherever found	T	\$8,000	\$1,197,809,905
1232	Amphipod, Noel's (<i>Gammarus desperatus</i>) - Wherever found	E	\$8,000	\$1,197,817,905
1232	Blindcat, Mexican (catfish) (<i>Prietella phreatophila</i>) - Wherever found	E	\$8,000	\$1,197,825,905
1232	Chumbo, Higo (<i>Harrisia portoricensis</i>) - Wherever found	T	\$8,000	\$1,197,833,905
1232	Dog, African wild (<i>Lycaon pictus</i>) - Wherever found	E	\$8,000	\$1,197,841,905
1232	Leopard (<i>Panthera pardus</i>) - except where threatened	E	\$8,000	\$1,197,849,905
1232	No Common Name (<i>Cordia bellonis</i>) - Wherever found	E	\$8,000	\$1,197,857,905
1232	Snail, Pecos <i>Assiminea pecos</i>) - Wherever found	E	\$8,000	\$1,197,865,905
1232	Springsnail, Koster's (<i>Juturnia kosteri</i>) - Wherever found	E	\$8,000	\$1,197,873,905
1232	Springsnail, Roswell (<i>Pyrgulopsis roswellensis</i>) - Wherever found	E	\$8,000	\$1,197,881,905
1233	No Common Name (<i>Stenogyne cranwelliae</i>) - Wherever found	E	\$7,990	\$1,197,889,895

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1234	Haiwale (<i>Cyrtandra waiolani</i>) - Wherever found	E	\$7,970	\$1,197,897,865
1235	Kaulu (<i>Pteralyxia macrocarpa</i>) - Wherever found	E	\$7,963	\$1,197,905,828
1236	Alani (<i>Melicope munroi</i>) - Wherever found	E	\$7,942	\$1,197,913,770
1237	Nehe (<i>Melanthera kamolensis</i>) - Wherever found	E	\$7,941	\$1,197,921,711
1238	Spinedace, Big Spring (<i>Lepidomeda mollispinis pratensis</i>) - Wherever found	T	\$7,922	\$1,197,929,633
1239	Kolea (<i>Myrsine vaccinioides</i>) - Wherever found	E	\$7,921	\$1,197,937,554
1240	Spineflower, Robust (<i>Chorizanthe robusta</i> var. <i>robusta</i>) - Wherever found	E	\$7,904	\$1,197,945,458
1241	`Oha wai (<i>Clermontia oblongifolia</i> ssp. <i>mauiensis</i>) - Wherever found	E	\$7,902	\$1,197,953,360
1242	Lampmussel, Alabama (<i>Lampsilis virescens</i>) - Wherever found; Except where listed as Experimental Populations	E	\$7,900	\$1,197,961,260
1243	No Common Name (<i>Bidens hillebrandiana</i> ssp. <i>hillebrandiana</i>) - Wherever found	E	\$7,870	\$1,197,969,130
1244	Manioc, Walker's (<i>Manihot walkerae</i>) - Wherever found	E	\$7,860	\$1,197,976,990
1245	Daisy, Parish's (<i>Erigeron parishii</i>) - Wherever found	T	\$7,820	\$1,197,984,810
1246	Nohoanu (<i>Geranium hillebrandii</i>) - Wherever found	E	\$7,801	\$1,197,992,611
1247	Lo`ulu (<i>Pritchardia lanigera</i>) - Wherever found	E	\$7,792	\$1,198,000,403
1248	Kopiko (<i>Psychotria hexandra</i> ssp. <i>oahuensis</i>) - Wherever found	E	\$7,777	\$1,198,008,180
1249	Crownscale, San Jacinto Valley (<i>Atriplex coronata</i> var. <i>notatior</i>) - Wherever found	E	\$7,733	\$1,198,015,913
1250	Rosemary, Cumberland (<i>Conradina verticillata</i>) - Wherever found	T	\$7,704	\$1,198,023,617
1251	Campion, fringed (<i>Silene polypetala</i>) - Wherever found	E	\$7,700	\$1,198,031,317
1252	Bean, sea (<i>Mucuna sloanei</i> var. <i>persericea</i>) - Wherever found	E	\$7,607	\$1,198,038,924
1253	Alani (<i>Melicope pallida</i>) - Wherever found	E	\$7,582	\$1,198,046,506
1254	Koki`o ke`oke`o (<i>Hibiscus arnottianus</i> ssp. <i>immaculatus</i>) - Wherever found	E	\$7,560	\$1,198,054,066
1255	No Common Name (<i>Huperzia stemmermanniae</i>) - Wherever found	E	\$7,520	\$1,198,061,586
1256	Wartyback, white (pearlymussel) (<i>Plethobasus cicatricosus</i>) - Wherever found	E	\$7,508	\$1,198,069,094
1257	Bariaco (<i>Trichilia triacantha</i>) - Wherever found	E	\$7,500	\$1,198,076,594
1257	Butterwort, Godfrey's (<i>Pinguicula ionantha</i>) - Wherever found	T	\$7,500	\$1,198,084,094
1257	Hypericum, highlands scrub (<i>Hypericum cumulicola</i>) - Wherever found	E	\$7,500	\$1,198,091,594
1257	Ladies-tresses, Canelo Hills (<i>Spiranthes delitescens</i>) - Wherever found	E	\$7,500	\$1,198,099,094
1257	No Common Name (<i>Auerodendron pauciflorum</i>) - Wherever found	E	\$7,500	\$1,198,106,594
1257	Skullcap, Florida (<i>Scutellaria floridana</i>) - Wherever found	T	\$7,500	\$1,198,114,094
1258	Diellia, Asplenium-leaved (<i>Asplenium dielirectum</i>) - Wherever found	E	\$7,432	\$1,198,121,526
1258	Haha (<i>Cyanea gibsonii</i>) - Wherever found	E	\$7,432	\$1,198,128,958

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1258	No Common Name (<i>Cyanea kauaulaensis</i>) - Wherever found	E	\$7,432	\$1,198,136,390
1258	No Common Name (<i>Labordia lorenciana</i>) - Wherever found	E	\$7,432	\$1,198,143,822
1258	No Common Name (<i>Sicyos lanceoloideus</i>) - Wherever found	E	\$7,432	\$1,198,151,254
1258	Paudedo (<i>Hedyotis megalantha</i>) - Wherever found	E	\$7,432	\$1,198,158,686
1258	Pilo (<i>Kadua laxiflora</i>) - Wherever found	E	\$7,432	\$1,198,166,118
1259	Clubshell, black (<i>Pleurobema curtum</i>) - Wherever found	E	\$7,425	\$1,198,173,543
1260	Hibiscus, Clay's (<i>Hibiscus clayi</i>) - Wherever found	E	\$7,420	\$1,198,180,963
1260	No Common Name (<i>Hesperomannia lydgatei</i>) - Wherever found	E	\$7,420	\$1,198,188,383
1261	Alani (<i>Melicope paniculata</i>) - Wherever found	E	\$7,408	\$1,198,195,791
1262	Milk-vetch, triple-ribbed (<i>Astragalus tricarinatus</i>) - Wherever found	E	\$7,400	\$1,198,203,191
1263	Monkshood, northern wild (<i>Aconitum noveboracense</i>) - Wherever found	T	\$7,369	\$1,198,210,560
1264	Turtle, flattened musk (<i>Sternotherus depressus</i>) - Black Warrior R. system upstream from Bankhead Dam	T	\$7,356	\$1,198,217,916
1265	Haha (<i>Cyanea asarifolia</i>) - Wherever found	E	\$7,339	\$1,198,225,255
1266	Blossom, tubercled (pearlymussel) (<i>Epioblasma torulosa torulosa</i>) - Wherever found; Except where listed as Experimental Populations	E	\$7,329	\$1,198,232,584
1267	Ma`oli`oli (<i>Schiedea apokremnos</i>) - Wherever found	E	\$7,319	\$1,198,239,903
1267	No Common Name (<i>Schiedea helleri</i>) - Wherever found	E	\$7,319	\$1,198,247,222
1268	Bluegrass, Hawaiian (<i>Poa sandvicensis</i>) - Wherever found	E	\$7,309	\$1,198,254,531
1268	Bluegrass, Mann's (<i>Poa mannii</i>) - Wherever found	E	\$7,309	\$1,198,261,840
1268	Kopiko (<i>Psychotria hobdyi</i>) - Wherever found	E	\$7,309	\$1,198,269,149
1268	Lo`ulu (<i>Pritchardia hardyi</i>) - Wherever found	E	\$7,309	\$1,198,276,458
1268	No Common Name (<i>Poa siphonoglossa</i>) - Wherever found	E	\$7,309	\$1,198,283,767
1268	Pilo kea lau li`i (<i>Platydesma rostrata</i>) - Wherever found	E	\$7,309	\$1,198,291,076
1269	`Awikiwiki (<i>Canavalia napaliensis</i>) - Wherever found	E	\$7,288	\$1,198,298,364
1269	'Akoko (<i>Euphorbia halemanui</i>) - Wherever found	E	\$7,288	\$1,198,305,652
1269	Ha`iwale (<i>Cyrtandra limahuliensis</i>) - Wherever found	T	\$7,288	\$1,198,312,940
1269	Ha`iwale (<i>Cyrtandra oenobarba</i>) - Wherever found	E	\$7,288	\$1,198,320,228
1269	Haha (<i>Cyanea recta</i>) - Wherever found	T	\$7,288	\$1,198,327,516
1269	Kamakahala (<i>Labordia helleri</i>) - Wherever found	E	\$7,288	\$1,198,334,804
1269	Koki`o ke`oke`o (<i>Hibiscus waimeae</i> ssp. <i>hannerae</i>) - Wherever found	E	\$7,288	\$1,198,342,092
1269	Mapele (<i>Cyrtandra cyaneoides</i>) - Wherever found	E	\$7,288	\$1,198,349,380

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1269	Na`ena`e (<i>Dubautia imbricata</i> ssp. <i>imbricata</i>) - Wherever found	E	\$7,288	\$1,198,356,668
1269	Na`ena`e (<i>Dubautia plantaginea</i> ssp. <i>magnifolia</i>) - Wherever found	E	\$7,288	\$1,198,363,956
1269	Nehe (<i>Lipochaeta fauriei</i>) - Wherever found	E	\$7,288	\$1,198,371,244
1270	`Ahinahina (<i>Argyroxiphium sandwicense</i> ssp. <i>sandwicense</i>) - Wherever found	E	\$7,253	\$1,198,378,497
1271	No Common Name (<i>Stenogyne kaalae</i> ssp. <i>sherffii</i>) - Wherever found	E	\$7,181	\$1,198,385,678
1272	Rocksnaail, plicate (<i>Leptoxis plicata</i>) - Wherever found	E	\$7,120	\$1,198,392,798
1273	Dawn-flower, Texas prairie (<i>Hymenoxys texana</i>) - Wherever found	E	\$7,100	\$1,198,399,898
1274	No Common Name (<i>Gouania hillebrandii</i>) - Wherever found	E	\$7,060	\$1,198,406,958
1275	Snail, Langford's tree (<i>Partula langfordi</i>) - Wherever found	E	\$7,038	\$1,198,413,996
1276	Cactus, Brady pincushion (<i>Pediocactus bradyi</i>) - Wherever found	E	\$7,000	\$1,198,420,996
1276	No Common Name (<i>Elaphoglossum serpens</i>) - Wherever found	E	\$7,000	\$1,198,427,996
1276	No Common Name (<i>Tectaria estremerana</i>) - Wherever found	E	7000	\$1,198,434,996
1276	Silverside, Waccamaw (<i>Menidia extensa</i>) - Wherever found	T	\$7,000	\$1,198,441,996
1277	Rock-cress, McDonald's (<i>Arabis macdonaldiana</i>) - Wherever found	E	\$6,962	\$1,198,448,958
1278	No Common Name (<i>Schiedea lydgatei</i>) - Wherever found	E	\$6,959	\$1,198,455,917
1279	Remya, Maui (<i>Remya mauiensis</i>) - Wherever found	E	\$6,949	\$1,198,462,866
1280	Wahane (<i>Pritchardia aylmer-robinsonii</i>) - Wherever found	E	\$6,948	\$1,198,469,814
1281	Haha (<i>Cyanea glabra</i>) - Wherever found	E	\$6,934	\$1,198,476,748
1282	Ihi (<i>Portulaca villosa</i>) - Wherever found	E	\$6,890	\$1,198,483,638
1283	No Common Name (<i>Trematolobelia singularis</i>) - Wherever found	E	\$6,735	\$1,198,490,373
1284	Mint, Garrett's (<i>Dicerandra christmanii</i>) - Wherever found	E	\$6,700	\$1,198,497,073
1285	Makou (<i>Ranunculus hawaiiensis</i>) - Wherever found	E	\$6,605	\$1,198,503,678
1286	Cactus, star (<i>Astrophytum asterias</i>) - Wherever found	E	\$6,600	\$1,198,510,278
1287	Malacothrix, Santa Cruz Island (<i>Malacothrix indecora</i>) - Wherever found	E	\$6,527	\$1,198,516,805
1288	Cypress, Gowen (<i>Cupressus goveniana</i> ssp. <i>goveniana</i>) - Wherever found	T	\$6,507	\$1,198,523,312
1289	Gambusia, Big Bend (<i>Gambusia gaigei</i>) - Wherever found	E	\$6,500	\$1,198,529,812
1289	No Common Name (<i>Varronia rupicola</i>) - Wherever found	T	\$6,500	\$1,198,536,312
1289	Palo de ramon (<i>Banara vanderbiltii</i>) - Wherever found	E	\$6,500	\$1,198,542,812
1290	[no common name] Snail (<i>Ostodes strigatus</i>) - Wherever found	E	\$6,424	\$1,198,549,236
1291	Barbara's Buttons, Mohr's (<i>Marshallia mohrii</i>) - Wherever found	T	\$6,422	\$1,198,555,658
1292	Haha (<i>Cyanea duvalliorum</i>) - Wherever found	E	\$6,420	\$1,198,562,078
1293	`Ena`ena (<i>Pseudognaphalium sandwicense</i> var. <i>molokaiense</i>) - Wherever found	E	\$6,414	\$1,198,568,492

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1294	Bladderpod, Spring Creek (<i>Lesquerella perforata</i>) - Wherever found	E	\$6,400	\$1,198,574,892
1295	Shiner, Cahaba (<i>Notropis cahabae</i>) - Wherever found	E	\$6,386	\$1,198,581,278
1296	'Oha wai (<i>Clermontia drepanomorpha</i>) - Wherever found	E	\$6,379	\$1,198,587,657
1296	No Common Name (<i>Schiedea diffusa</i> subsp. <i>diffusa</i>) - Wherever found	E	\$6,379	\$1,198,594,036
1297	Ma`oli`oli (<i>Schiedea kealiae</i>) - Wherever found	E	\$6,337	\$1,198,600,373
1298	No Common Name (<i>Achyranthes mutica</i>) - Wherever found	E	\$6,323	\$1,198,606,696
1299	'O`u (honeycreeper) (<i>Psittirostra psittacea</i>) - Wherever found	E	\$6,309	\$1,198,613,005
1300	Sedge, Navajo (<i>Carex specuicola</i>) - Wherever found	T	\$6,300	\$1,198,619,305
1301	No Common Name (<i>Doryopteris takeuchii</i>) - Wherever found	E	\$6,212	\$1,198,625,517
1302	Ha`iwale (<i>Cyrtandra crenata</i>) - Wherever found	E	\$6,208	\$1,198,631,725
1303	'Awiwi (<i>Kadua cookiana</i>) - Wherever found	E	\$6,203	\$1,198,637,928
1304	No Common Name (<i>Festuca molokaiensis</i>) - Wherever found	E	\$6,198	\$1,198,644,126
1305	Haha (<i>Cyanea mceldowneyi</i>) - Wherever found	E	\$6,168	\$1,198,650,294
1306	Salamander, Jollyville Plateau (<i>Eurycea tonkawae</i>) - Wherever found	T	\$6,100	\$1,198,656,394
1307	No Common Name (<i>Schiedea salicaria</i>) - Wherever found	E	\$6,097	\$1,198,662,491
1307	No Common Name (<i>Schiedea sarmentosa</i>) - Wherever found	E	\$6,097	\$1,198,668,588
1308	Alani (<i>Melicope reflexa</i>) - Wherever found	E	\$6,067	\$1,198,674,655
1308	Ha`iwale (<i>Cyrtandra oxybapha</i>) - Wherever found	E	\$6,067	\$1,198,680,722
1308	Haha (<i>Cyanea mannii</i>) - Wherever found	E	\$6,067	\$1,198,686,789
1308	Haha (<i>Cyanea mauiensis</i>) - Wherever found	E	\$6,067	\$1,198,692,856
1308	Ko`oko`olau (<i>Bidens conjuncta</i>) - Wherever found	E	\$6,067	\$1,198,698,923
1308	No Common Name (<i>Cyperus fauriei</i>) - Wherever found	E	\$6,067	\$1,198,704,990
1309	Hala pepe (<i>Pleomele fernaldii</i>) - Wherever found	E	\$6,055	\$1,198,711,045
1310	Alani (<i>Melicope makahae</i>) - Wherever found	E	\$6,034	\$1,198,717,079
1311	Boxwood, Vahl's (<i>Buxus vahlii</i>) - Wherever found	E	\$6,000	\$1,198,723,079
1311	Guajon (<i>Eleutherodactylus cooki</i>) - Wherever found	T	\$6,000	\$1,198,729,079
1311	Mint, longspurred (<i>Dicerandra cornutissima</i>) - Wherever found	E	\$6,000	\$1,198,735,079
1311	Pawpaw, Rugel's (<i>Deeringothamnus rugelii</i>) - Wherever found	E	\$6,000	\$1,198,741,079
1311	Rosemary, short-leaved (<i>Conradina brevifolia</i>) - Wherever found	E	\$6,000	\$1,198,747,079
1312	Bladderpod, kodachrome (<i>Lesquerella tumulosa</i>) - Wherever found	E	\$5,950	\$1,198,753,029
1313	Awiwi (<i>Schenkia sebaeoides</i>) - Wherever found	E	\$5,914	\$1,198,758,943
1313	Ha`iwale (<i>Cyrtandra hematos</i>) - Wherever found	E	\$5,914	\$1,198,764,857

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1313	Haha (<i>Cyanea lanceolata</i>) - Wherever found	E	\$5,914	\$1,198,770,771
1313	Haha (<i>Cyanea magnicalyx</i>) - Wherever found	E	\$5,914	\$1,198,776,685
1313	Hohiu (<i>Dryopteris glabra</i> var. <i>pusilla</i>) - Wherever found	E	\$5,914	\$1,198,782,599
1313	Ko`oko`olau (<i>Bidens amplexans</i>) - Wherever found	E	\$5,914	\$1,198,788,513
1313	Ko`oko`olau (<i>Bidens wiebkei</i>) - Wherever found	E	\$5,914	\$1,198,794,427
1313	Kolea (<i>Myrsine fosbergii</i>) - Wherever found	E	\$5,914	\$1,198,800,341
1313	Kopa (<i>Kadua cordata remyi</i>) - Wherever found	E	\$5,914	\$1,198,806,255
1313	Kupukupu makalii (<i>Cyclosorus boydiae</i>) - Wherever found	E	\$5,914	\$1,198,812,169
1313	Loulu, Baker's (<i>Pritchardia bakeri</i>) - Wherever found	E	\$5,914	\$1,198,818,083
1313	Ma`oli`oli (<i>Schiedea pubescens</i>) - Wherever found	E	\$5,914	\$1,198,823,997
1313	Makou (<i>Ranunculus mauiensis</i>) - Wherever found	E	\$5,914	\$1,198,829,911
1313	No Common Name (<i>Asplenium diellaciniatum</i>) - Wherever found	E	\$5,914	\$1,198,835,825
1313	No Common Name (<i>Cyperus neokunthianus</i>) - Wherever found	E	\$5,914	\$1,198,841,739
1313	No Common Name (<i>Deparia kaalaana</i>) - Wherever found	E	\$5,914	\$1,198,847,653
1313	No Common Name (<i>Kadua haupuensis</i>) - Wherever found	E	\$5,914	\$1,198,853,567
1313	No Common Name (<i>Lepidium orbiculare</i>) - Wherever found	E	\$5,914	\$1,198,859,481
1313	No Common Name (<i>Phyllostegia helleri</i>) -	E	\$5,914	\$1,198,865,395
1313	No Common Name (<i>Santalum involutum</i>) - Wherever found	E	\$5,914	\$1,198,871,309
1313	No Common Name (<i>Wikstroemia skottsbergiana</i>) -	E	\$5,914	\$1,198,877,223
1313	Olua (<i>Hypolepis hawaiiensis</i> var. <i>mauiensis</i>) - Wherever found	E	5914	\$1,198,883,137
1314	Papala (<i>Charpentiera densiflora</i>) - Wherever found	E	\$5,853	\$1,198,888,990
1314	Reedgrass, Hillebrand's (<i>Calamagrostis hillebrandii</i>) - Wherever found	E	\$5,853	\$1,198,894,843
1315	Roseroot, Leedy's (<i>Rhodiola integrifolia</i> ssp. <i>leedyi</i>) - Wherever found	T	\$5,808	\$1,198,900,651
1316	Milkvetch, Deseret (<i>Astragalus desereticus</i>) - Wherever found	T	\$5,750	\$1,198,906,401
1317	Combshell, Cumberlandian (<i>Epioblasma brevidens</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$5,700	\$1,198,912,101
1317	Fleabane, Zuni (<i>Erigeron rhizomatus</i>) - Wherever found	T	\$5,700	\$1,198,917,801
1317	Lampmussel, Alabama (<i>Lampsilis virescens</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$5,700	\$1,198,923,501
1317	Mussel, oyster (<i>Epioblasma capsaeformis</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$5,700	\$1,198,929,201
1317	Salamander, Austin blind (<i>Eurycea waterlooensis</i>) - Wherever found	E	\$5,700	\$1,198,934,901

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1318	Blazingstar, scrub (<i>Liatris ohlingerae</i>) - Wherever found	E	\$5,600	\$1,198,940,501
1318	Blue-star, Kearney's (<i>Amsonia kearneyana</i>) - Wherever found	E	\$5,600	\$1,198,946,101
1319	Birds-in-a-nest, white (<i>Macbridea alba</i>) - Wherever found	T	\$5,500	\$1,198,951,601
1319	Jacquemontia, beach (<i>Jacquemontia reclinata</i>) - Wherever found	E	\$5,500	\$1,198,957,101
1320	Reed-mustard, Barneby (<i>Schoenocrambe barnebyi</i>) - Wherever found	E	\$5,450	\$1,198,962,551
1321	Alopecurus, Sonoma (<i>Alopecurus aequalis</i> var. <i>sonomensis</i>) - Wherever found	E	\$5,400	\$1,198,967,951
1321	Salamander, Georgetown (<i>Eurycea naufragia</i>) - Wherever found	T	\$5,400	\$1,198,973,351
1321	Salamander, Salado (<i>Eurycea chisholmensis</i>) - Wherever found	T	\$5,400	\$1,198,978,751
1322	Townsendia, Last Chance (<i>Townsendia aprica</i>) - Wherever found	T	\$5,350	\$1,198,984,101
1323	Bladderpod, white (<i>Lesquerella pallida</i>) - Wherever found	E	\$5,200	\$1,198,989,301
1323	Milkpea, Small's (<i>Galactia smallii</i>) - Wherever found	E	\$5,200	\$1,198,994,501
1323	Polygala, tiny (<i>Polygala smallii</i>) - Wherever found	E	\$5,200	\$1,198,999,701
1324	Rocksnailed, interrupted (=Georgia) (<i>Leptoxis foremani</i>) - Wherever found	E	\$5,199	\$1,199,004,900
1325	Hornsnail, rough (<i>Pleurocera foremani</i>) - Wherever found	E	\$5,196	\$1,199,010,096
1326	Isopod, Socorro (<i>Thermosphaeroma thermophilus</i>) - Wherever found	E	\$5,160	\$1,199,015,256
1327	Pawpaw, beautiful (<i>Deeringothamnus pulchellus</i>) - Wherever found	E	\$5,100	\$1,199,020,356
1327	Springsnail, Alamosa (<i>Tryonia alamosae</i>) - Wherever found	E	\$5,100	\$1,199,025,456
1327	Springsnail, Chupadera (<i>Pyrgulopsis chupadera</i>) - Wherever found	E	\$5,100	\$1,199,030,556
1328	Cactus, Cochise pincushion (<i>Coryphantha robbinsiorum</i>) - Wherever found	T	\$5,000	\$1,199,035,556
1328	Campeloma, slender (<i>Campeloma decampi</i>) - Wherever found	E	\$5,000	\$1,199,040,556
1328	Chupacallos (<i>Pleodendron macranthum</i>) - Wherever found	E	\$5,000	\$1,199,045,556
1328	Erubia (<i>Solanum drymophilum</i>) - Wherever found	E	\$5,000	\$1,199,050,556
1328	No Common Name (<i>Catesbaea melanocarpa</i>) - Wherever found	E	\$5,000	\$1,199,055,556
1328	No Common Name (<i>Polystichum calderonense</i>) - Wherever found	E	\$5,000	\$1,199,060,556
1328	Prickly-ash, St. Thomas (<i>Zanthoxylum thomasianum</i>) - Wherever found	E	\$5,000	\$1,199,065,556
1329	Sunburst, Hartweg's golden (<i>Pseudobahia bahiifolia</i>) - Wherever found	E	\$4,959	\$1,199,070,515
1330	Bluegrass, San Bernardino (<i>Poa atropurpurea</i>) - Wherever found	E	\$4,933	\$1,199,075,448
1331	Lilliput, pale (pearlymussel) (<i>Toxolasma cylindrellus</i>) - Wherever found	E	\$4,900	\$1,199,080,348
1331	Navarretia, many-flowered (<i>Navarretia leucocephala</i> ssp. <i>plieantha</i>) - Wherever found	E	\$4,900	\$1,199,085,248
1332	Daisy, Lakeside (<i>Hymenoxys herbacea</i>) - Wherever found	T	\$4,859	\$1,199,090,107
1333	Bear, brown (<i>Ursus arctos pruinosus</i>) - Wherever found	E	\$4,800	\$1,199,094,907

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1334	Oxytheca, Cushenbury (<i>Oxytheca parishii</i> var. <i>goodmaniana</i>) - Wherever found	E	\$4,750	\$1,199,099,657
1335	Milk-vetch, Cushenbury (<i>Astragalus albens</i>) - Wherever found	E	\$4,720	\$1,199,104,377
1336	Darter, bluemask (<i>Etheostoma akatulo</i>) - Wherever found	E	\$4,704	\$1,199,109,081
1337	Mustard, Carter's (<i>Warea carteri</i>) - Wherever found	E	\$4,700	\$1,199,113,781
1338	Fly, Hawaiian picture-wing (<i>Drosophila mulli</i>) - Wherever found	T	\$4,642	\$1,199,118,423
1339	Pigtoe, dark (<i>Pleurobema furvum</i>) - Wherever found	E	\$4,620	\$1,199,123,043
1340	Pea, Big Pine partridge (<i>Chamaecrista lineata keyensis</i>) - Wherever found	E	\$4,600	\$1,199,127,643
1341	Moth, Kern primrose sphinx (<i>Euproserpinus euterpe</i>) - Wherever found	T	\$4,590	\$1,199,132,233
1342	Amphipod, Hay's Spring (<i>Stygobromus hayi</i>) - Wherever found	E	\$4,532	\$1,199,136,765
1343	Deer, musk (<i>Moschus</i> spp. (all species)) - Afghanistan, Bhutan, Burma, China [Tibet, Yunnan], India, Nepal, Pakistan, Sikkim	E	\$4,500	\$1,199,141,265
1343	Fern, Elfin tree (<i>Cyathea dryopteroides</i>) - Wherever found	E	\$4,500	\$1,199,145,765
1343	Flax, sand (<i>Linum arenicola</i>) - Wherever found	E	\$4,500	\$1,199,150,265
1343	Heather, mountain golden (<i>Hudsonia montana</i>) - Wherever found	T	\$4,500	\$1,199,154,765
1343	Holly, Cook's (<i>Ilex cookii</i>) - Wherever found	E	\$4,500	\$1,199,159,265
1343	Manaca, palma de (<i>Calyptronoma rivalis</i>) - Wherever found	T	\$4,500	\$1,199,163,765
1343	No Common Name (<i>Daphnopsis hellerana</i>) - Wherever found	E	\$4,500	\$1,199,168,265
1343	No Common Name (<i>Myrcia paganii</i>) - Wherever found	E	\$4,500	\$1,199,172,765
1343	No Common Name (<i>Thelypteris inabonensis</i>) - Wherever found	E	4500	\$1,199,177,265
1343	Uvillo (<i>Eugenia haematocarpa</i>) - Wherever found	E	\$4,500	\$1,199,181,765
1344	Spurge, deltoid (<i>Chamaesyce deltoidea</i> ssp. <i>deltoidea</i>) - Wherever found	E	\$4,400	\$1,199,186,165
1344	Spurge, wedge (<i>Chamaesyce deltoidea serpyllum</i>) - Wherever found	E	\$4,400	\$1,199,190,565
1345	Spineflower, slender-horned (<i>Dodecahema leptoceras</i>) - Wherever found	E	\$4,367	\$1,199,194,932
1346	Leather flower, Morefield's (<i>Clematis morefieldii</i>) -	E	\$4,302	\$1,199,199,234
1347	Crayfish, Hell Creek Cave (<i>Cambarus zophonastes</i>) - Wherever found	E	\$4,200	\$1,199,203,434
1347	Manzanita, Presidio (<i>Arctostaphylos hookeri</i> var. <i>ravenii</i>) - Wherever found	E	\$4,200	\$1,199,207,634
1348	Vole, Hualapai Mexican (<i>Microtus mexicanus hualpaiensis</i>) - Wherever found	E	\$4,100	\$1,199,211,734
1349	Cavefish, Alabama (<i>Speoplatyrhinus poulsoni</i>) - Wherever found	E	\$4,000	\$1,199,215,734
1349	Harebells, Avon Park (<i>Crotalaria avonensis</i>) - Wherever found	E	\$4,000	\$1,199,219,734
1349	No Common Name (<i>Agave eggersiana</i>) - Wherever found	E	\$4,000	\$1,199,223,734
1349	Pitcher-plant, Alabama canebrake (<i>Sarracenia rubra</i> ssp. <i>alabamensis</i>) - Wherever found	E	\$4,000	\$1,199,227,734

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1349	Sunfish, spring pygmy (<i>Elassoma alabamae</i>) - Wherever found	T	\$4,000	\$1,199,231,734
1349	Warbler, elfin-woods (<i>Setophaga angelae</i>) - Wherever found	T	\$4,000	\$1,199,235,734
1350	Catspaw, white (pearlymussel) (<i>Epioblasma obliquata perobliqua</i>) - Wherever found	E	\$3,978	\$1,199,239,712
1351	Pigtoe, Cumberland (<i>Pleurobema gibberum</i>) - Wherever found	E	\$3,814	\$1,199,243,526
1352	Spinymussel, Altamaha (<i>Elliptio spinosa</i>) - Wherever found	E	\$3,750	\$1,199,247,276
1353	Madtom, Scioto (<i>Noturus trautmani</i>) - Wherever found	E	\$3,559	\$1,199,250,835
1354	Bellflower, Brooksville (<i>Campanula robinsiae</i>) - Wherever found	E	\$3,500	\$1,199,254,335
1354	Cavesnail, Tumbling Creek (<i>Antrobia culveri</i>) - Wherever found	E	\$3,500	\$1,199,257,835
1354	Liveforever, Santa Barbara Island (<i>Dudleya traskiae</i>) - Wherever found	E	\$3,500	\$1,199,261,335
1354	No Common Name (<i>Calyptanthes thomasi</i>) - Wherever found	E	\$3,500	\$1,199,264,835
1354	No Common Name (<i>Mitracarpus maxwelliae</i>) - Wherever found	E	\$3,500	\$1,199,268,335
1354	No Common Name (<i>Thelypteris verecunda</i>) - Wherever found	E	3500	\$1,199,271,835
1355	Creeper, Molokai (<i>Paroreomyza flammea</i>) - Wherever found	E	\$3,390	\$1,199,275,225
1356	Meadowfoam, Butte County (<i>Limnanthes floccosa</i> ssp. <i>californica</i>) - Wherever found	E	\$3,300	\$1,199,278,525
1357	Thistle, Chorro Creek bog (<i>Cirsium fontinale</i> var. <i>obispoense</i>) - Wherever found	E	\$3,288	\$1,199,281,813
1358	Bladderpod, Zapata (<i>Lesquerella thamnophila</i>) - Wherever found	E	\$3,200	\$1,199,285,013
1358	Meshweaver, Madla's Cave (<i>Cicurina madla</i>) - Wherever found	E	\$3,200	\$1,199,288,213
1358	Salamander, Red Hills (<i>Phaeognathus hubrichti</i>) - Wherever found	T	\$3,200	\$1,199,291,413
1359	Amphipod, diminutive (<i>Gammarus hyalleoides</i>) - Wherever found	E	\$3,000	\$1,199,294,413
1359	Amphipod, Illinois cave (<i>Gammarus acherondytes</i>) - Wherever found	E	\$3,000	\$1,199,297,413
1359	Amphipod, Pecos (<i>Gammarus pecos</i>) - Wherever found	E	\$3,000	\$1,199,300,413
1359	Cactus, Nichol's Turk's head (<i>Echinocactus horizontalonius</i> var. <i>nicholii</i>) - Wherever found	E	\$3,000	\$1,199,303,413
1359	Fringe-tree, pygmy (<i>Chionanthus pygmaeus</i>) - Wherever found	E	\$3,000	\$1,199,306,413
1359	Lupine, clover (<i>Lupinus tidestromii</i>) - Wherever found	E	\$3,000	\$1,199,309,413
1359	Madtom, yellowfin (<i>Noturus flavipinnis</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(e)(1)(iii))	EXPN	\$3,000	\$1,199,312,413
1359	Meshweaver, Government Canyon Bat Cave (<i>Cicurina vespera</i>) - Wherever found	E	\$3,000	\$1,199,315,413
1359	Meshweaver, Robber Baron Cave (<i>Cicurina baronia</i>) - Wherever found	E	\$3,000	\$1,199,318,413
1359	No Common Name (<i>Adiantum vivesii</i>) - Wherever found	E	\$3,000	\$1,199,321,413
1359	No Common Name (<i>Leptocereus grantianus</i>) - Wherever found	E	\$3,000	\$1,199,324,413

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1359	No Common Name (<i>Mitracarpus polycladus</i>) - Wherever found	E	\$3,000	\$1,199,327,413
1359	Palo colorado (<i>Ternstroemia luquillensis</i>) - Wherever found	E	\$3,000	\$1,199,330,413
1359	Palo de jazmin (<i>Styrax portoricensis</i>) - Wherever found	E	\$3,000	\$1,199,333,413
1359	Pronghorn, peninsular (<i>Antilocapra americana peninsularis</i>) - Mexico (Baja California)	E	\$3,000	\$1,199,336,413
1359	Puma (=mountain lion) (<i>Puma</i> (=Felis) <i>concolor</i> (all subsp. except <i>coryi</i>)) - FL	SAT	\$3,000	\$1,199,339,413
1359	Spider, Government Canyon Bat Cave (<i>Neoleptoneta microps</i>) - Wherever found	E	\$3,000	\$1,199,342,413
1360	Snail, tulotoma (<i>Tulotoma magnifica</i>) - Wherever found	T	\$2,899	\$1,199,345,312
1361	Spineflower, Sonoma (<i>Chorizanthe valida</i>) - Wherever found	E	\$2,850	\$1,199,348,162
1362	Brickell-bush, Florida (<i>Brickellia mosieri</i>) - Wherever found	E	\$2,800	\$1,199,350,962
1362	Chub, Sonora (<i>Gila ditaenia</i>) - Wherever found	T	\$2,800	\$1,199,353,762
1362	Snail, painted snake coiled forest (<i>Anguispira picta</i>) - Wherever found	T	\$2,800	\$1,199,356,562
1363	Primrose, Maguire (<i>Primula maguirei</i>) - Wherever found	T	\$2,750	\$1,199,359,312
1364	Beetle, Kretschmarr Cave mold (<i>Texamaurops reddelli</i>) - Wherever found	E	\$2,700	\$1,199,362,012
1364	Bluet, Roan Mountain (<i>Hedyotis purpurea</i> var. <i>montana</i>) - Wherever found	E	\$2,700	\$1,199,364,712
1364	Fern, Florida bristle (<i>Trichomanes punctatum</i> ssp. <i>floridanum</i>) - Wherever found	E	\$2,700	\$1,199,367,412
1364	Fringepod, Santa Cruz Island (<i>Thysanocarpus conchuliferus</i>) - Wherever found	E	\$2,700	\$1,199,370,112
1364	Harvestman, Bee Creek Cave (<i>Texella reddelli</i>) - Wherever found	E	\$2,700	\$1,199,372,812
1364	Mustard, slender-petaled (<i>Thelypodium stenopetalum</i>) - Wherever found	E	\$2,700	\$1,199,375,512
1365	Lioplax, cylindrical (snail) (<i>Lioplax cyclostomaformis</i>) - Wherever found	E	\$2,600	\$1,199,378,112
1365	Manzanita, pallid (<i>Arctostaphylos pallida</i>) - Wherever found	T	\$2,600	\$1,199,380,712
1365	Vervain, Red Hills (<i>Verbena californica</i>) - Wherever found	T	\$2,600	\$1,199,383,312
1366	Snail, flat-spined three-toothed (<i>Triodopsis platysayoides</i>) - Wherever found	T	\$2,503	\$1,199,385,815
1367	Beetle, Helotes mold (<i>Batrisodes venyivi</i>) - Wherever found	E	\$2,500	\$1,199,388,315
1367	Capa rosa (<i>Callicarpa ampla</i>) - Wherever found	E	\$2,500	\$1,199,390,815
1367	Cliffrose, Arizona (<i>Purshia</i> (=Cowania) <i>subintegra</i>) - Wherever found	E	\$2,500	\$1,199,393,315
1367	Harvestman, Cokendolpher Cave (<i>Texella cokendolpheri</i>) - Wherever found	E	\$2,500	\$1,199,395,815
1367	Marstonia, royal (snail) (<i>Pyrgulopsis ogmorhapse</i>) - Wherever found	E	\$2,500	\$1,199,398,315
1367	Meshweaver, Braken Bat Cave (<i>Cicurina venii</i>) - Wherever found	E	\$2,500	\$1,199,400,815
1367	No Common Name (<i>Ilex sintenisii</i>) - Wherever found	E	\$2,500	\$1,199,403,315
1367	No Common Name (<i>Ternstroemia subsessilis</i>) - Wherever found	E	\$2,500	\$1,199,405,815
1367	No Common Name (<i>Thelypteris yaucoensis</i>) - Wherever found	E	2500	\$1,199,408,315

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1367	Peperomia, Wheeler's (<i>Peperomia wheeleri</i>) - Wherever found	E	\$2,500	\$1,199,410,815
1367	Thoroughwort, Cape Sable (<i>Chromolaena frustrata</i>) - Wherever found	E	\$2,500	\$1,199,413,315
1367	Walnut (=Nogal), West Indian (<i>Juglans jamaicensis</i>) - Wherever found	E	\$2,500	\$1,199,415,815
1368	Ridge-cress, Barneby (<i>Lepidium barnebyanum</i>) - Wherever found	E	\$2,450	\$1,199,418,265
1369	Baccharis, Encinitas (<i>Baccharis vanessae</i>) - Wherever found	T	\$2,443	\$1,199,420,708
1370	Bean, Cumberland (pearlymussel) (<i>Villosa trabalis</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$2,423	\$1,199,423,131
1371	Manzanita, Del Mar (<i>Arctostaphylos glandulosa</i> ssp. <i>crassifolia</i>) - Wherever found	E	\$2,420	\$1,199,425,551
1372	Blossom, turgid (pearlymussel) (<i>Epioblasma turgidula</i>) - Wherever found; Except where listed as Experimental Populations	E	\$2,400	\$1,199,427,951
1373	Po`ouli (honeycreeper) (<i>Melamprosops phaeosoma</i>) - Wherever found	E	\$2,376	\$1,199,430,327
1374	Shark, Scalloped Hammerhead (<i>Sphyrna lewini</i>) - Indo-West Pacific DPS	T	\$2,355	\$1,199,432,682
1375	Butterfly, lotis blue (<i>Lycaeides argyrognomon lotis</i>) - Wherever found	E	\$2,310	\$1,199,434,992
1376	Sneezeweed, Virginia (<i>Helenium virginicum</i>) - Wherever found	T	\$2,300	\$1,199,437,292
1377	Dace, Independence Valley speckled (<i>Rhinichthys osculus lethoporus</i>) - Wherever found	E	\$2,288	\$1,199,439,580
1378	Combshell, Cumberlandian (<i>Epioblasma brevidens</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$2,222	\$1,199,441,802
1379	Ragwort, San Francisco Peaks (<i>Packera franciscana</i>) - Wherever found	T	\$2,200	\$1,199,444,002
1380	Bat, little Mariana fruit (<i>Pteropus tokudae</i>) - Wherever found	E	\$2,104	\$1,199,446,106
1381	Blossom, green (pearlymussel) (<i>Epioblasma torulosa gubernaculum</i>) - Wherever found	E	\$2,103	\$1,199,448,209
1382	Goldenrod, Blue Ridge (<i>Solidago spithamea</i>) - Wherever found	T	\$2,100	\$1,199,450,309
1382	Lessingia, San Francisco (<i>Lessingia germanorum</i> (=L.g. var. <i>germanorum</i>)) - Wherever found	E	\$2,100	\$1,199,452,409
1382	No Common Name (<i>Cranichis ricartii</i>) - Wherever found	E	\$2,100	\$1,199,454,509
1382	Springsnail, San Bernardino (<i>Pyrgulopsis bernardina</i>) - Wherever found	T	\$2,100	\$1,199,456,609
1383	Liliwai (<i>Acaena exigua</i>) - Wherever found	E	\$2,074	\$1,199,458,683
1384	Chub, spotfin (<i>Erimonax monachus</i>) - U.S.A. (TN-specified portions of the Tellico River; see 17.84(m)(1)(i))	EXPN	\$2,020	\$1,199,460,703
1384	Mussel, oyster (<i>Epioblasma capsaeformis</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$2,020	\$1,199,462,723

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1385	Cactus, Chisos Mountain hedgehog (<i>Echinocereus chisoensis</i> var. <i>chisoensis</i>) - Wherever found	T	\$2,000	\$1,199,464,723
1385	Cactus, Lloyd's Mariposa (<i>Echinomastus mariposensis</i>) - Wherever found	T	\$2,000	\$1,199,466,723
1385	Chub, spotfin (<i>Erimonax monachus</i>) - U.S.A. (TN-specified portions of the French Broad and Holston Rivers; see 17.84(m)(1)(iii))	EXPN	\$2,000	\$1,199,468,723
1385	Irisette, white (<i>Sisyrinchium dichotomum</i>) - Wherever found	E	\$2,000	\$1,199,470,723
1385	No Common Name (<i>Chamaecrista glandulosa</i> var. <i>mirabilis</i>) - Wherever found	E	\$2,000	\$1,199,472,723
1385	No Common Name (<i>Gesneria pauciflora</i>) - Wherever found	T	\$2,000	\$1,199,474,723
1385	Water-willow, Cooley's (<i>Justicia cooleyi</i>) - Wherever found	E	\$2,000	\$1,199,476,723
1385	Whitlow-wort, papery (<i>Paronychia chartacea</i>) - Wherever found	T	\$2,000	\$1,199,478,723
1386	Riversnail, Anthony's (<i>Athearnia anthonyi</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$1,866	\$1,199,480,589
1387	Sawfish, largetooth (<i>Pristis pristis</i>) - Wherever found	E	\$1,853	\$1,199,482,442
1388	Fanshell (<i>Cyprogenia stegaria</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$1,808	\$1,199,484,250
1389	Snail, Virginia fringed mountain (<i>Polygyriscus virginianus</i>) - Wherever found	E	\$1,800	\$1,199,486,050
1390	Springsnail, Socorro (<i>Pyrgulopsis neomexicana</i>) - Wherever found	E	\$1,760	\$1,199,487,810
1391	Beetle, Tooth Cave ground (<i>Rhadine persephone</i>) - Wherever found	E	\$1,700	\$1,199,489,510
1391	Curlew, Eskimo (<i>Numenius borealis</i>) - Wherever found	E	\$1,700	\$1,199,491,210
1391	Dogweed, ashy (<i>Thymophylla tephroleuca</i>) - Wherever found	E	\$1,700	\$1,199,492,910
1391	Flax, Carter's small-flowered (<i>Linum carteri carteri</i>) - Wherever found	E	\$1,700	\$1,199,494,610
1391	Pentachaeta, white-rayed (<i>Pentachaeta bellidiflora</i>) - Wherever found	E	\$1,700	\$1,199,496,310
1391	Prickly-apple, fragrant (<i>Cereus eriophorus</i> var. <i>fragrans</i>) - Wherever found	E	\$1,700	\$1,199,498,010
1391	Pseudoscorpion, Tooth Cave (<i>Tartarocreagris texana</i>) - Wherever found	E	\$1,700	\$1,199,499,710
1391	Spider, Tooth Cave (<i>Neoleptoneta myopica</i>) - Wherever found	E	\$1,700	\$1,199,501,410
1392	Clover, Monterey (<i>Trifolium trichocalyx</i>) - Wherever found	E	\$1,666	\$1,199,503,076
1393	Bird's-beak, Pennell's (<i>Cordylanthus tenuis</i> ssp. <i>capillaris</i>) - Wherever found	E	\$1,650	\$1,199,504,726
1393	Checker-mallow, Kenwood Marsh (<i>Sidalcea oregana</i> ssp. <i>valida</i>) - Wherever found	E	\$1,650	\$1,199,506,376
1393	Lily, Pitkin Marsh (<i>Lilium pardalinum</i> ssp. <i>pitkinense</i>) - Wherever found	E	\$1,650	\$1,199,508,026
1393	Mariposa lily, Tiburon (<i>Calochortus tiburonensis</i>) - Wherever found	T	\$1,650	\$1,199,509,676
1393	Milk-vetch, Clara Hunt's (<i>Astragalus clarianus</i>) - Wherever found	E	\$1,650	\$1,199,511,326

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1393	Wild-buckwheat, southern mountain (<i>Eriogonum kennedyi</i> var. <i>austromontanum</i>) - Wherever found	T	\$1,650	\$1,199,512,976
1394	Cat, leopard (<i>Prionailurus</i> (=Felis) <i>bengalensis bengalensis</i>) - Wherever found	E	\$1,600	\$1,199,514,576
1395	Thornmint, San Mateo (<i>Acanthomintha obovata</i> ssp. <i>duttonii</i>) - Wherever found	E	\$1,547	\$1,199,516,123
1396	Anole, Culebra Island giant (<i>Anolis roosevelti</i>) - Wherever found	E	\$1,500	\$1,199,517,623
1396	Boa, Mona (<i>Epicrates monensis monensis</i>) - Wherever found	T	\$1,500	\$1,199,519,123
1396	Crow, white-necked (<i>Corvus leucognaphalus</i>) - Wherever found	E	\$1,500	\$1,199,520,623
1396	No Common Name (<i>Gonocalyx concolor</i>) - Wherever found	E	\$1,500	\$1,199,522,123
1396	Snail, armored (<i>Pyrgulopsis</i> (=Marstonia) <i>pachyta</i>) - Wherever found	E	\$1,500	\$1,199,523,623
1396	Warbler (=wood), Bachman's (<i>Vermivora bachmanii</i>) - Wherever found	E	\$1,500	\$1,199,525,123
1397	Sea turtle, loggerhead (<i>Caretta caretta</i>) - South Pacific Ocean DPS	E	\$1,406	\$1,199,526,529
1398	Rocksnaail, painted (<i>Leptoxis taeniata</i>) - Wherever found	T	\$1,399	\$1,199,527,928
1399	`O`o, Kauai (honeyeater) (<i>Moho braccatus</i>) - Wherever found	E	\$1,362	\$1,199,529,290
1399	Akialoa, Kauai (honeycreeper) (<i>Akialoa stejnegeri</i>) - Wherever found	E	\$1,362	\$1,199,530,652
1400	Pearlymussel, dromedary (<i>Dromus dromas</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$1,212	\$1,199,531,864
1401	Gladecress, Texas golden (<i>Leavenworthia texana</i>) - Wherever found	E	\$1,200	\$1,199,533,064
1402	Onion, Munz's (<i>Allium munzii</i>) - Wherever found	E	\$1,133	\$1,199,534,197
1403	Lemurs (<i>Lemuridae</i> (incl. genera <i>Lemur</i> , <i>Phaner</i> , <i>Hapalemur</i> , <i>Lepilemur</i> , <i>Microcebus</i> , <i>Allocebus</i> , <i>Cheirogaleus</i> , <i>Varecia</i>)) - Wherever found	E	\$1,100	\$1,199,535,297
1403	Thistle, Sacramento Mountains (<i>Cirsium vinaceum</i>) - Wherever found	T	\$1,100	\$1,199,536,397
1403	Trout, Little Kern golden (<i>Oncorhynchus aguabonita whitei</i>) - Wherever found	T	\$1,100	\$1,199,537,497
1404	Buckwheat, Umtanum Desert (<i>Eriogonum codium</i>) - Wherever found	T	\$1,050	\$1,199,538,547
1405	Pigtoe, finerayed (<i>Fusconaia cuneolus</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$1,010	\$1,199,539,557
1405	Pigtoe, rough (<i>Pleurobema plenum</i>) - U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$1,010	\$1,199,540,567
1406	Blossom, yellow (pearlymussel) (<i>Epioblasma florentina florentina</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$1,000	\$1,199,541,567
1406	Cactus, Nellie cory (<i>Coryphantha minima</i>) - Wherever found	E	\$1,000	\$1,199,542,567
1406	Cory cactus, bunched (<i>Coryphantha ramillosa</i>) - Wherever found	T	\$1,000	\$1,199,543,567

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1406	Gambusia, Clear Creek (<i>Gambusia heterochir</i>) - Wherever found	E	\$1,000	\$1,199,544,567
1406	Lily, Minnesota dwarf trout (<i>Erythronium propullans</i>) - Wherever found	E	\$1,000	\$1,199,545,567
1406	Minnow, Rio Grande Silvery (<i>Hybognathus amarus</i>) - Rio Grande, from Little Box Canyon to Amistad Dam	EXPN	\$1,000	\$1,199,546,567
1406	Parrot, thick-billed (<i>Rhynchopsitta pachyrhyncha</i>) - Wherever found	E	\$1,000	\$1,199,547,567
1406	Seal, spotted (<i>Phoca largha</i>) - Southern DPS	T	\$1,000	\$1,199,548,567
1406	Shrimp, Alabama cave (<i>Palaemonias alabamae</i>) - Wherever found	E	\$1,000	\$1,199,549,567
1407	Kamapua`a (<i>Kadua fluviatilis</i>) - Wherever found	E	\$943	\$1,199,550,510
1408	Bittercress, small-anthered (<i>Cardamine micranthera</i>) - Wherever found	E	\$900	\$1,199,551,410
1408	Water-plantain, Kral's (<i>Sagittaria secundifolia</i>) - Wherever found	T	\$900	\$1,199,552,310
1409	Flannelbush, Mexican (<i>Fremontodendron mexicanum</i>) - Wherever found	E	\$750	\$1,199,553,060
1410	Yellow-faced bee, Hawaiian (<i>Hylaeus mana</i>) - Wherever found	E	\$733	\$1,199,553,793
1410	Yellow-faced bee, hiliaris (<i>Hylaeus hiliaris</i>) - Wherever found	E	\$733	\$1,199,554,526
1411	Ground-Dove, Friendly (<i>Gallucolumba stairi</i>) - American Samoa DPS	E	\$726	\$1,199,555,252
1412	Barberry, Nevin's (<i>Berberis nevinii</i>) - Wherever found	E	\$700	\$1,199,555,952
1412	Birch, Virginia round-leaf (<i>Betula uber</i>) - Wherever found	T	\$700	\$1,199,556,652
1413	Checkermallow, Wenatchee Mountains (<i>Sidalcea oregana</i> var. <i>calva</i>) - Local Endemic	E	\$652	\$1,199,557,304
1413	Stickseed, showy (<i>Hackelia venusta</i>) - Wherever found	E	\$652	\$1,199,557,956
1414	No Common Name (<i>Maesa walkeri</i>) - Wherever found	T	\$614	\$1,199,558,570
1415	Pigtoe, shiny (<i>Fusconaia cor</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$606	\$1,199,559,176
1416	Poppy-mallow, Texas (<i>Callirhoe scabriuscula</i>) - Wherever found	E	\$600	\$1,199,559,776
1417	Sea turtle, green (<i>Chelonia mydas</i>) - Southwest Indian DPS	T	\$521	\$1,199,560,297
1418	Cat's-eye, Terlingua Creek (<i>Cryptantha crassipes</i>) - Wherever found	E	\$500	\$1,199,560,797
1418	Darter, vermilion (<i>Etheostoma chermocki</i>) - Wherever found	E	\$500	\$1,199,561,297
1418	Gambusia, San Marcos (<i>Gambusia georgei</i>) - Wherever found	E	\$500	\$1,199,561,797
1418	Oak, Hinckley (<i>Quercus hinckleyi</i>) - Wherever found	T	\$500	\$1,199,562,297
1418	Pitaya, Davis' green (<i>Echinocereus viridiflorus</i> var. <i>davisii</i>) - Wherever found	E	\$500	\$1,199,562,797
1418	Pondweed, Little Aguja (=Creek) (<i>Potamogeton clystocarpus</i>) - Wherever found	E	\$500	\$1,199,563,297
1418	Tamarin, cotton-top (<i>Saguinus oedipus</i>) - Wherever found	E	\$500	\$1,199,563,797

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulatative Total
1419	Monkeyface, Appalachian (pearlymussel) (<i>Quadrula sparsa</i>) - USA (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	EXPN	\$422	\$1,199,564,219
1420	Blossom, turgid (pearlymussel) (<i>Epioblasma turgidula</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$404	\$1,199,564,623
1420	Pearlymussel, cracking (<i>Hemistena lata</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	EXPN	\$404	\$1,199,565,027
1421	Thrush, large Kauai (=kamao) (<i>Myadestes myadestinus</i>) - Wherever found	E	\$348	\$1,199,565,375
1422	White-eye, bridled (<i>Zosterops conspicillatus conspicillatus</i>) - Wherever found	E	\$304	\$1,199,565,679
1423	Babirusa (<i>Babirusa babirusa</i>) - Wherever found	E	\$300	\$1,199,565,979
1423	Gazelle, slender-horned (<i>Gazella leptoceros</i>) - Wherever found	E	\$300	\$1,199,566,279
1423	Seal, Ringed (<i>Phoca (=Pusa) hispida botnica</i>) - Wherever found	T	\$300	\$1,199,566,579
1423	Siamang (<i>Symphalangus syndactylus</i>) - Wherever found	E	\$300	\$1,199,566,879
1424	Buckwheat, cushenbury (<i>Eriogonum ovalifolium</i> var. <i>vineum</i>) - Wherever found	E	\$220	\$1,199,567,099
1425	Blossom, yellow (pearlymussel) (<i>Epioblasma florentina florentina</i>) - Wherever found; Except where listed as Experimental Populations	E	\$202	\$1,199,567,301
1425	Pearlymussel, birdwing (<i>Lemiox rimosus</i>) - AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert, and Lauderdale Counties, AL	EXPN	\$202	\$1,199,567,503
1426	Addax (<i>Addax nasomaculatus</i>) - Wherever found	E	\$200	\$1,199,567,703
1426	Bladderpod, lyrate (<i>Lesquerella lyrata</i>) - Wherever found	T	\$200	\$1,199,567,903
1426	Bladderpod, San Bernardino Mountains (<i>Lesquerella kingii</i> ssp. <i>bernardina</i>) - Wherever found	E	\$200	\$1,199,568,103
1426	Cat, tiger (<i>Leopardus (=Felis) tigrinus</i>) - Wherever found	E	\$200	\$1,199,568,303
1426	Gladecress, Fleshy-fruit (<i>Leavenworthia crassa</i>) - Wherever found	E	\$200	\$1,199,568,503
1426	Jewelflower, Tiburon (<i>Streptanthus niger</i>) - Wherever found	E	\$200	\$1,199,568,703
1426	Koala (<i>Phascolarctos cinereus</i>) - Australia	T	\$200	\$1,199,568,903
1426	Langur, Francois' (<i>Trachypithecus (=Presbytis) francoisi</i>) - Wherever found	E	\$200	\$1,199,569,103
1426	Lechwe, red (<i>Kobus leche</i>) - Wherever found	T	\$200	\$1,199,569,303
1426	Monkey, Diana (<i>Cercopithecus diana</i>) - Wherever found	E	\$200	\$1,199,569,503
1426	Pearlymussel, Curtis (<i>Epioblasma florentina curtisii</i>) - Wherever found	E	\$200	\$1,199,569,703
1426	Prickly-apple, aboriginal (<i>Harrisia (=Cereus) aboriginum (=gracilis)</i>) - Wherever found	E	\$200	\$1,199,569,903
1426	Tapir, South American (=Brazilian) (<i>Tapirus terrestris</i>) - Wherever found	E	\$200	\$1,199,570,103

Rank	Species (50 CFR Part 17)	Status	Species Total	Cumulative Total
1427	Crownbeard, big-leaved (<i>Verbesina dissita</i>) - Wherever found	T	\$100	\$1,199,570,203
1427	Dugong (<i>Dugong dugon</i>) - Wherever found	E	\$100	\$1,199,570,303
1427	Malacothrix, island (<i>Malacothrix squalida</i>) - Wherever found	E	\$100	\$1,199,570,403
1427	Skipper, Pawnee montane (<i>Hesperia leonardus montana</i>) - Wherever found	T	\$100	\$1,199,570,503
1427	Taraxacum, California (<i>Taraxacum californicum</i>) - Wherever found	E	\$100	\$1,199,570,603
1428	Crane, black-necked (<i>Grus nigricollis</i>) - Wherever found	E	\$40	\$1,199,570,643
1428	Crane, hooded (<i>Grus monacha</i>) - Wherever found	E	\$40	\$1,199,570,683
1428	Crane, white-naped (<i>Grus vipio</i>) - Wherever found	E	\$40	\$1,199,570,723
1429	Crane, Japanese (<i>Grus japonensis</i>) - Wherever found	E	\$20	\$1,199,570,743
1430	Whale, humpback (<i>Megaptera novaeangliae</i>) - Cape Verde Island/Northwest Africa DPS	E	\$0	\$1,199,570,743
1430	Whale, humpback (<i>Megaptera novaeangliae</i>) - Arabian Sea DPS	E	\$0	\$1,199,570,743
1430	Bat, Pacific sheath-tailed (<i>Emballonura semicaudata semicaudata</i>) - American Samoa	E	\$0	\$1,199,570,743
1430	(honeyeater), Mao (= maomao) (<i>Gymnomyza samoensis</i>) - Wherever found	E	\$0	\$1,199,570,743
1430	Snake, Dusky sea (<i>Aipysurus fuscus</i>)	E	\$0	\$1,199,570,743
1430	Cardinalfish, Banggai (<i>Pterapogon kauderni</i>)	T	\$0	\$1,199,570,743
1430	Coelacanth, African (<i>Latimeria chalumnae</i>) - Tanzanian DPS	T	\$0	\$1,199,570,743
1430	Angelshark, Sawback (<i>Squatina aculeata</i>)	E	\$0	\$1,199,570,743
1430	Angelshark, Smoothback (<i>Squatina oculata</i>)	E	\$0	\$1,199,570,743
1430	Shark, Angel (<i>Squatina squatina</i>)	E	\$0	\$1,199,570,743
1430	Butterfly, Oregon silverspot (<i>Speyeria zerene hippolyta</i>) -	EXPN	\$0	\$1,199,570,743
1430	No Common Name (<i>Cantharellus noumeae</i>) -	E	\$0	\$1,199,570,743
1430	No Common Name (<i>Siderastrea glynni</i>)	E	\$0	\$1,199,570,743
1430	No Common Name (<i>Tubastraea floreana</i>) -	E	\$0	\$1,199,570,743
Total Ranked Species Expenditures				\$1,199,570,743

**TABLE 3. FY 2017 REPORTED LAND ACQUISITION EXPENDITURES FOR
ENDANGERED AND THREATENED SPECIES**

Species are arranged in the same sequence as Table 1. These costs are not included in the expenditures displayed in Tables 1 and 2. Only those species that had land acquisition costs reported for FY 2017 are included here. Land rank is included to show the relative order of the species for which land expenditures are reported. Land acquisition costs for multi-species conservation activities are tabulated at the end of Table 3. The STATUS column indicates whether the species is endangered (E), threatened (T), experimental (EXPN), or threatened by similarity of appearance (SAT).

TABLE 3. REPORTED LAND ACQUISITION EXPENDITURES FOR FY 2017 FOR ENDANGERED AND THREATENED SPECIES

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
Mammals	47	Bat, gray (<i>Myotis grisescens</i>) - Wherever found	E	\$0	\$0	\$0	\$100,000	\$100,000
	28	Bat, Hawaiian hoary (<i>Lasiurus cinereus semotus</i>) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	30	Bat, Indiana (<i>Myotis sodalis</i>) - Wherever found	E	\$250,000	\$0	\$250,000	\$100,000	\$350,000
	27	Bat, Northern long-eared (<i>Myotis septentrionalis</i>) - Wherever found	T	\$0	\$250,000	\$250,000	\$189,400	\$439,400
	61	Bear, polar (<i>Ursus maritimus</i>) - Wherever found	T	\$16,000	\$0	\$16,000	\$0	\$16,000
	35	Lynx, Canada (<i>Lynx canadensis</i>) - Wherever Found in Contiguous U.S.	T	\$275,250	\$0	\$275,250	\$0	\$275,250
	36	Manatee, West Indian (<i>Trichechus manatus</i>) - Wherever found	E	\$270,336	\$0	\$270,336	\$0	\$270,336
	62	Ocelot (<i>Leopardus (=Felis) pardalis</i>) - wherever found	E	\$7,067	\$0	\$7,067	\$0	\$7,067
	3	Panther, Florida (<i>Puma (=Felis) concolor coryi</i>) - Wherever found	E	\$0	\$3,000,000	\$3,000,000	\$0	\$3,000,000
	20	Prairie dog, Utah (<i>Cynomys parvidens</i>) - Wherever found	T	\$0	\$0	\$0	\$530,000	\$530,000
	51	Pronghorn, Sonoran (<i>Antilocapra americana sonoriensis</i>) - Wherever found, except where listed as an experimental population	E	\$0	\$48,970	\$48,970	\$0	\$48,970
	61	Whale, bowhead (<i>Balaena mysticetus</i>) - Wherever found	E	\$16,000	\$0	\$16,000	\$0	\$16,000
	43	Wolf, red (<i>Canis rufus</i>) - Wherever found, except where listed as an experimental population	E	\$137,500	\$0	\$137,500	\$0	\$137,500
Mammals subtotal				\$1,370,248	\$3,298,970	\$4,669,218	\$919,400	\$5,588,618

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
Birds	4	Caracara, Audubon's crested (Polyborus plancus audubonii) - FL pop.	T	\$783	\$2,500,000	\$2,500,783	\$0	\$2,500,783
	28	Coot, Hawaiian (Fulica americana alai) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	11	Crane, whooping (Grus americana) - Wherever found, except where listed as an experimental population	E	\$1,069,573	\$0	\$1,069,573	\$0	\$1,069,573
	28	Creep, Hawaii (Oreomystis mana) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	37	Cuckoo, yellow-billed (Coccyzus americanus) - Western U.S. DPS	T	\$0	\$250,000	\$250,000	\$0	\$250,000
	28	Duck, Hawaiian (=koloa) (Anas wyvilliana) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	61	Eider, spectacled (Somateria fischeri) - Wherever found	T	\$16,000	\$0	\$16,000	\$0	\$16,000
	40	Eider, Steller's (Polysticta stelleri) - AK breeding pop.	T	\$216,000	\$0	\$216,000	\$0	\$216,000
	48	Falcon, Northern Aplomado (Falco femoralis septentrionalis) - Wherever found, except where listed as an experimental population	E	\$63,200	\$0	\$63,200	\$0	\$63,200
	60	Gnatcatcher, coastal California (Polioptila californica californica) - Wherever found	T	\$22,077	\$0	\$22,077	\$0	\$22,077
	28	Goose, Hawaiian (Branta (=Nesochen) sandvicensis) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	28	Hawk, Hawaiian (=lo) (Buteo solitarius) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	8	Plover, piping (Charadrius melodus) - [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	T	\$1,388,269	\$0	\$1,388,269	\$0	\$1,388,269
	56	Scrub-jay, Florida (Aphelocoma coerulescens) - Wherever found	T	\$783	\$0	\$783	\$27,990	\$28,773

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	6	Sparrow, Florida grasshopper (Ammodramus savannarum floridanus) - Wherever found	E	\$0	\$1,500,000	\$1,500,000	\$16,310	\$1,516,310
	1	Stork, wood (Mycteria americana) - AL, FL, GA, MS, NC, SC	T	\$223,227	\$10,798,527	\$11,021,754	\$0	\$11,021,754
	15	Tern, least (Sterna antillarum) - interior pop.	E	\$868,542	\$70,000	\$938,542	\$0	\$938,542
	26	Tern, roseate (Sterna dougallii dougallii) - northeast U.S. nesting pop.	E	\$440,000	\$0	\$440,000	\$0	\$440,000
	25	Vireo, Black-capped (Vireo atricapilla) - Wherever found	E	\$440,748	\$0	\$440,748	\$0	\$440,748
	60	Vireo, least Bell's (Vireo bellii pusillus) - Wherever found	E	\$22,077	\$0	\$22,077	\$0	\$22,077
	24	Warbler (=wood), golden-cheeked (Dendroica chrysoparia) - Wherever found	E	\$447,815	\$0	\$447,815	\$0	\$447,815
	12	Woodpecker, red-cockaded (Picoides borealis) - Wherever found	E	\$705,069	\$0	\$705,069	\$344,598	\$1,049,667
Birds subtotal				\$7,914,638	\$15,118,527	\$23,033,165	\$388,898	\$23,422,063
Reptiles	64	Alligator, American (Alligator mississippiensis) - Wherever found	SAT	\$0	\$0	\$0	\$2,429	\$2,429
	38	Sea turtle, green (Chelonia mydas) - North Atlantic DPS	T	\$229,511	\$0	\$229,511	\$0	\$229,511
	62	Sea turtle, hawksbill (Eretmochelys imbricata) - Wherever found	E	\$7,067	\$0	\$7,067	\$0	\$7,067
	34	Sea turtle, Kemp's ridley (Lepidochelys kempii) - Wherever found	E	\$285,644	\$0	\$285,644	\$0	\$285,644
	38	Sea turtle, leatherback (Dermochelys coriacea) - Wherever found	E	\$229,511	\$0	\$229,511	\$0	\$229,511
	13	Skink, bluetail mole (Eumeces egregius lividus) - Wherever found	T	\$783	\$1,000,000	\$1,000,783	\$0	\$1,000,783
	67	Skink, sand (Neoseps reynoldsi) - Wherever found	T	\$783	\$0	\$783	\$0	\$783

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	2	Snake, eastern indigo (<i>Drymarchon corais couperi</i>) - Wherever found	T	\$783	\$0	\$783	\$3,000,000	\$3,000,783
	57	Snake, giant garter (<i>Thamnophis gigas</i>) - Wherever found	T	\$28,167	\$0	\$28,167	\$0	\$28,167
	29	Tortoise, desert (<i>Gopherus agassizii</i>) - Wherever found, except AZ south and east of Colorado R., and Mexico	T	\$0	\$357,600	\$357,600	\$0	\$357,600
	49	Tortoise, gopher (<i>Gopherus polyphemus</i>) - West of Mobile and Tombigbee Rivers	T	\$0	\$61,000	\$61,000	\$0	\$61,000
	17	Turtle, bog (<i>Clemmys muhlenbergii</i>) - Wherever found, except GA, NC, SC, TN, VA	T	\$216,250	\$497,767	\$714,017	\$0	\$714,017
Reptiles subtotal				\$998,499	\$1,916,367	\$2,914,866	\$3,002,429	\$5,917,295
Amphibians	57	Salamander, California tiger (<i>Ambystoma californiense</i>) - U.S.A. (Central CA DPS)	T	\$28,167	\$0	\$28,167	\$0	\$28,167
	31	Salamander, frosted flatwoods (<i>Ambystoma cingulatum</i>) - Wherever found	T	\$222,444	\$0	\$222,444	\$118,599	\$341,043
	66	Salamander, Reticulated flatwoods (<i>Ambystoma bishopi</i>) - Wherever found	E	\$0	\$0	\$0	\$1,198	\$1,198
Amphibians subtotal				\$250,611	\$0	\$250,611	\$119,797	\$370,408
Fishes	58	Cui-ui (<i>Chasmistes cujus</i>) - Wherever found	E	\$26,250	\$0	\$26,250	\$0	\$26,250
	53	Darter, Cherokee (<i>Etheostoma scotti</i>) - Wherever found	T	\$0	\$0	\$0	\$40,000	\$40,000
	53	Darter, Etowah (<i>Etheostoma etowahae</i>) - Wherever found	E	\$0	\$0	\$0	\$40,000	\$40,000
	54	Eulachon (<i>Thaleichthys pacificus</i>) - Southern DPS	T	\$35,667	\$0	\$35,667	\$0	\$35,667
	3	Minnow, Rio Grande Silvery (<i>Hybognathus amarus</i>) - Wherever found, except where listed as an experimental population	E	\$0	\$3,000,000	\$3,000,000	\$0	\$3,000,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	22	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Snake River fall-run ESU	T	\$0	\$506,000	\$506,000	\$0	\$506,000
	22	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Snake River spring/summer-run ESU	T	\$0	\$506,000	\$506,000	\$0	\$506,000
	49	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Sacramento River winter-run ESU	E	\$0	\$61,000	\$61,000	\$0	\$61,000
	14	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Lower Columbia River ESU	T	\$0	\$972,462	\$972,462	\$0	\$972,462
	22	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Upper Columbia spring-run ESU	E	\$0	\$506,000	\$506,000	\$0	\$506,000
	49	Salmon, Chinook (<i>Oncorhynchus</i> (=Salmo) tshawytscha) - Central Valley spring-run ESU	T	\$0	\$61,000	\$61,000	\$0	\$61,000
	54	Salmon, coho (<i>Oncorhynchus</i> (=Salmo) kisutch) - Oregon Coast ESU	T	\$35,667	\$0	\$35,667	\$0	\$35,667
	22	Salmon, sockeye (<i>Oncorhynchus</i> (=Salmo) nerka) - Snake River ESU	E	\$0	\$506,000	\$506,000	\$0	\$506,000
	10	Shiner, Pecos bluntnose (<i>Notropis simus pecosensis</i>) - Wherever found	T	\$0	\$1,100,000	\$1,100,000	\$0	\$1,100,000
	18	Shiner, Topeka (<i>Notropis topeka</i> (=tristis)) - Wherever found, except where listed as an experimental population	E	\$650,163	\$0	\$650,163	\$0	\$650,163
	49	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - California Central Valley DPS	T	\$0	\$61,000	\$61,000	\$0	\$61,000
	22	Steelhead (<i>Oncorhynchus</i> (=Salmo) mykiss) - Snake River Basin DPS	T	\$0	\$506,000	\$506,000	\$0	\$506,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	19	Steelhead (Oncorhynchus (=Salmo) mykiss) - Lower Columbia River DPS	T	\$0	\$532,763	\$532,763	\$0	\$532,763
	21	Steelhead (Oncorhynchus (=Salmo) mykiss) - Upper Columbia River DPS	T	\$0	\$512,254	\$512,254	\$0	\$512,254
	22	Steelhead (Oncorhynchus (=Salmo) mykiss) - Middle Columbia River DPS	T	\$0	\$506,000	\$506,000	\$0	\$506,000
	54	Sturgeon, green (Acipenser medirostris) - Southern DPS	T	\$35,667	\$0	\$35,667	\$0	\$35,667
	7	Sturgeon, pallid (Scaphirhynchus albus) - Wherever found	E	\$861,475	\$556,043	\$1,417,518	\$0	\$1,417,518
	52	Trout, bull (Salvelinus confluentus) - U.S.A., conterminous, lower 48 states	T	\$0	\$46,000	\$46,000	\$0	\$46,000
	58	Trout, Lahontan cutthroat (Oncorhynchus clarkii henshawi) - Wherever found	T	\$26,250	\$0	\$26,250	\$0	\$26,250
Fishes subtotal				\$1,671,139	\$9,938,522	\$11,609,661	\$80,000	\$11,689,661
Clams	39	Bankclimber, purple (mussel) (Elliptioideus sloatianus) - Wherever found	T	\$222,444	\$0	\$222,444	\$0	\$222,444
	41	Higgins eye (pearlymussel) (Lampsilis higginsii) - Wherever found	E	\$185,667	\$0	\$185,667	\$0	\$185,667
	44	Mussel, sheepnose (Plethobasus cyphus) - Wherever found	E	\$134,667	\$0	\$134,667	\$0	\$134,667
	44	Spectaclecase (mussel) (Cumberlandia monodonta) - Wherever found	E	\$134,667	\$0	\$134,667	\$0	\$134,667
	33	Wedgemussel, dwarf (Alasmidonta heterodon) - Wherever found	E	\$275,250	\$35,000	\$310,250	\$0	\$310,250
Clams subtotal				\$952,695	\$35,000	\$987,695	\$0	\$987,695
Insects	45	Beetle, valley elderberry longhorn (Desmocerus californicus dimorphus) - Wherever found	T	\$0	\$110,000	\$110,000	\$0	\$110,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	50	Butterfly, Karner blue (<i>Lycaeides melissa samuelis</i>) - Wherever found	E	\$0	\$0	\$0	\$50,000	\$50,000
	60	Butterfly, Quino checkerspot (<i>Euphydryas editha quino</i> (=E. e. wrighti)) - Wherever found	E	\$22,077	\$0	\$22,077	\$0	\$22,077
	28	Damselfly, Pacific Hawaiian (<i>Megalagrion pacificum</i>) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	28	Fly, Hawaiian picture-wing (<i>Drosophila heteroneura</i>) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	35	Tiger beetle, Puritan (<i>Cicindela puritana</i>) - Wherever found	T	\$275,250	\$0	\$275,250	\$0	\$275,250
Insects subtotal				\$1,093,517	\$110,000	\$1,203,517	\$50,000	\$1,253,517
Crustaceans	57	Fairy shrimp, Conservancy (<i>Branchinecta conservatio</i>) - Wherever found	E	\$28,167	\$0	\$28,167	\$0	\$28,167
	57	Fairy shrimp, longhorn (<i>Branchinecta longiantenna</i>) - Wherever found	E	\$28,167	\$0	\$28,167	\$0	\$28,167
	60	Fairy shrimp, San Diego (<i>Branchinecta sandiegonensis</i>) - Wherever found	E	\$22,077	\$0	\$22,077	\$0	\$22,077
	42	Fairy shrimp, vernal pool (<i>Branchinecta lynchi</i>) - Wherever found	T	\$28,167	\$152,720	\$180,887	\$0	\$180,887
	57	Tadpole shrimp, vernal pool (<i>Lepidurus packardi</i>) - Wherever found	E	\$28,167	\$0	\$28,167	\$0	\$28,167
Crustaceans subtotal				\$134,745	\$152,720	\$287,465	\$0	\$287,465
Flowering Plants	28	`Aiea (<i>Nothocestrum breviflorum</i>) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	55	Amaranth, seabeach (<i>Amaranthus pumilus</i>) - Wherever found	T	\$33,800	\$0	\$33,800	\$0	\$33,800
	60	Ambrosia, San Diego (<i>Ambrosia pumila</i>) - Wherever found	E	\$22,077	\$0	\$22,077	\$0	\$22,077

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	62	Ayenia, Texas (<i>Ayenia limitaris</i>) - Wherever found	E	\$7,067	\$0	\$7,067	\$0	\$7,067
	67	Beargrass, Britton's (<i>Nolina brittoniana</i>) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	65	Bedstraw, El Dorado (<i>Galium californicum</i> ssp. <i>sierrae</i>) - Wherever found	E	\$0	\$1,500	\$1,500	\$0	\$1,500
	63	Bladderpod, Dudley Bluffs (<i>Lesquerella congesta</i>) - Wherever found	T	\$0	\$5,000	\$5,000	\$0	\$5,000
	62	Bladderpod, Zapata (<i>Lesquerella thamnophila</i>) - Wherever found	E	\$7,067	\$0	\$7,067	\$0	\$7,067
	67	Blazingstar, scrub (<i>Liatris ohlingerae</i>) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	67	Bonamia, Florida (<i>Bonamia grandiflora</i>) - Wherever found	T	\$783	\$0	\$783	\$0	\$783
	67	Buckwheat, scrub (<i>Eriogonum longifolium</i> var. <i>gnaphalifolium</i>) - Wherever found	T	\$783	\$0	\$783	\$0	\$783
	23	Bulrush, Northeastern (<i>Scirpus ancistrochaetus</i>) - Wherever found	E	\$491,500	\$0	\$491,500	\$0	\$491,500
	70	Butterweed, Layne's (<i>Senecio layneae</i>) - Wherever found	T	\$0	\$600	\$600	\$0	\$600
	60	Button-celery, San Diego (<i>Eryngium aristulatum</i> var. <i>parishii</i>) - Wherever found	E	\$22,077	\$0	\$22,077	\$0	\$22,077
	62	Cactus, star (<i>Astrophytum asterias</i>) - Wherever found	E	\$7,067	\$0	\$7,067	\$0	\$7,067
	32	Catchfly, Spalding's (<i>Silene spaldingii</i>) - Wherever found	T	\$310,440	\$0	\$310,440	\$0	\$310,440
	71	Ceanothus, Pine Hill (<i>Ceanothus roderickii</i>) - Wherever found	E	\$0	\$400	\$400	\$0	\$400
	62	Dogweed, ashy (<i>Thymophylla tephroleuca</i>) - Wherever found	E	\$7,067	\$0	\$7,067	\$0	\$7,067
	59	Dropwort, Canby's (<i>Oxypolis canbyi</i>) - Wherever found	E	\$0	\$0	\$0	\$25,000	\$25,000

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	60	Flannelbush, Mexican (Fremontodendron mexicanum) - Wherever found	E	\$22,077	\$0	\$22,077	\$0	\$22,077
	68	Flannelbush, Pine Hill (Fremontodendron californicum ssp. decumbens) - Wherever found	E	\$0	\$700	\$700	\$0	\$700
	67	Fringe-tree, pygmy (Chionanthus pygmaeus) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	5	Haha (Cyanea hamatiflora ssp. carlsonii) - Wherever found	E	\$1,592,381	\$0	\$1,592,381	\$0	\$1,592,381
	28	Ha'iwale (Cyrtandra tintinnabula) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	32	Howellia, water (Howellia aquatilis) - Wherever found	T	\$310,440	\$0	\$310,440	\$0	\$310,440
	67	Hypericum, highlands scrub (Hypericum cumulicola) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	28	Kiponapona (Phyllostegia racemosa) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	67	Lupine, scrub (Lupinus aridorum) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	62	Manioc, Walker's (Manihot walkerae) - Wherever found	E	\$7,067	\$0	\$7,067	\$0	\$7,067
	60	Mesa-mint, Otay (Pogogyne nudiuscula) - Wherever found	E	\$22,077	\$0	\$22,077	\$0	\$22,077
	60	Mesa-mint, San Diego (Pogogyne abramsii) - Wherever found	E	\$22,077	\$0	\$22,077	\$0	\$22,077
	67	Mint, Garrett's (Dicerandra christmanii) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	69	Morning-glory, Stebbins' (Calystegia stebbinsii) - Wherever found	E	\$0	\$650	\$650	\$0	\$650
	67	Mustard, Carter's (Warea carteri) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	60	Navarretia, spreading (Navarretia fossalis) - Wherever found	T	\$22,077	\$0	\$22,077	\$0	\$22,077

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
	16	No Common Name (Phyllostegia velutina) - Wherever found	E	\$796,190	\$0	\$796,190	\$0	\$796,190
	9	`Oha wai (Clermontia lindseyana) - Wherever found	E	\$1,194,286	\$0	\$1,194,286	\$0	\$1,194,286
	46	Orchid, western prairie fringed (Platanthera praeclara) - Wherever found	T	\$109,338	\$0	\$109,338	\$0	\$109,338
	60	Orcutt grass, California (Orcuttia californica) - Wherever found	E	\$22,077	\$0	\$22,077	\$0	\$22,077
	67	Pigeon wings (Clitoria fragrans) - Wherever found	T	\$783	\$0	\$783	\$0	\$783
	55	Pink, swamp (Helonias bullata) - Wherever found	T	\$33,800	\$0	\$33,800	\$0	\$33,800
	67	Plum, scrub (Prunus geniculata) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	28	Po`e (Portulaca sclerocarpa) - Wherever found	E	\$398,095	\$0	\$398,095	\$0	\$398,095
	67	Polygala, Lewton's (Polygala lewtonii) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	67	Sandlace (Polygonella myriophylla) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	60	Tarplant, Otay (Deinandra (=Hemizonia) conjugens) - Wherever found	T	\$22,077	\$0	\$22,077	\$0	\$22,077
	60	Thornmint, San Diego (Acanthomintha ilicifolia) - Wherever found	T	\$22,077	\$0	\$22,077	\$0	\$22,077
	63	Twinpod, Dudley Bluffs (Physaria obcordata) - Wherever found	T	\$0	\$5,000	\$5,000	\$0	\$5,000
	67	Whitlow-wort, papery (Paronychia chartacea) - Wherever found	T	\$783	\$0	\$783	\$0	\$783
	67	Wireweed (Polygonella basiramia) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
	67	Ziziphus, Florida (Ziziphus celata) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
Flowering Plants subtotal				\$6,711,111	\$13,850	\$6,724,961	\$25,000	\$6,749,961

Group Name	Rank	Species (50 CFR Part 17)	Status	FWS Total	Other Fed	Federal Total	States Total	Species Total
Lichens	67	Cladonia, Florida perforate (Cladonia perforata) - Wherever found	E	\$783	\$0	\$783	\$0	\$783
Lichens subtotal				\$783	\$0	\$783	\$0	\$783
Multi-species				\$0	\$0	\$0	\$471,323	\$471,323
Multi-species subtotal				\$0	\$0	\$0	\$471,323	\$471,323
Total Land Expenditures				\$21,097,986	\$30,583,956	\$51,681,942	\$5,056,847	\$56,738,789

TABLE 4. FY 2017 EXPENDITURES REPORTED BY INDIVIDUAL FEDERAL AGENCIES AND THE STATES

This table displays combined species expenditures and land acquisition costs by each reporting Federal agency, the Service, and the States collectively. Expenditures for foreign species are included, as well as costs for multi-species conservation activities.

Table 4. FY2017 EXPENDITURES REPORTED BY INDIVIDUAL FEDERAL AGENCIES AND THE STATES


Agency Type	Expense Type	Agency	Species Total	Land Total	Cumulative Total
Agency	Federal Domestic Expenditures	DHS Coast Guard	\$795,000	\$0	\$795,000
		DHS Customs and Border Protection	\$150,085	\$0	\$150,085
		DHS Federal Emergency Management Agency	\$40,010	\$0	\$40,010
		DOC National Oceanic and Atmospheric Administration	\$189,687,000	\$0	\$189,687,000
		DOD Air Force	\$15,119,880	\$0	\$15,119,880
		DOD Army	\$21,747,662	\$0	\$21,747,662
		DOD Army Corps of Engineers	\$181,136,616	\$556,043	\$181,692,659
		DOD Marine Corps	\$14,937,430	\$0	\$14,937,430
		DOD Navy	\$27,212,222	\$0	\$27,212,222
		DOE Bonneville Power Administration	\$302,777,939	\$499,479	\$303,277,418
		DOE Southwestern Power Administration	\$19,200	\$70,000	\$89,200
		DOE Western Area Power Administration	\$13,581,207	\$0	\$13,581,207
		DOI Bureau of Indian Affairs	\$2,427,793	\$0	\$2,427,793
		DOI Bureau of Land Management	\$21,755,835	\$420,420	\$22,176,255
		DOI Bureau of Reclamation	\$112,694,700	\$9,133,000	\$121,827,700
		DOI National Park Service	\$18,087,172	\$0	\$18,087,172
		DOI US Geological Survey	\$17,122,605	\$0	\$17,122,605
		DOT Federal Aviation Administration	\$1,502,500	\$0	\$1,502,500
		Environmental Protection Agency	\$179,456	\$0	\$179,456
		Federal Energy Regulatory Commission	\$284,600	\$0	\$284,600
		Nuclear Regulatory Commission	\$221,100	\$0	\$221,100
		Smithsonian Institution	\$478,400	\$0	\$478,400
		Tennessee Valley Authority	\$2,035,300	\$0	\$2,035,300
		USDA Animal and Plant Health Inspection Service	\$10,177,000	\$0	\$10,177,000
		USDA Forest Service	\$475,425	\$0	\$475,425
		USDA Natural Resources Conservation Service	\$11,885,129	\$19,905,014	\$31,790,143
		US Fish and Wildlife Service	\$168,401,946	\$21,097,986	\$189,499,932
	<i>Federal Domestic Expenditures subtotal</i>		<i>\$1,134,933,212</i>	<i>\$51,681,942</i>	<i>\$1,186,615,154</i>
	Federal Foreign Expenditures	Environmental Protection Agency	\$100	\$0	\$100
		DOD Navy	\$3,772	\$0	\$3,772
		DOI National Park Service	\$8,000	\$0	\$8,000
		USDA Animal and Plant Health Inspection Service	\$40,100	\$0	\$40,100
		DOI US Geological Survey	\$46,042	\$0	\$46,042
		DOC National Oceanic and Atmospheric Administration	\$170,000	\$0	\$170,000
		DOD Army Corps of Engineers	\$543,915	\$0	\$543,915
		Smithsonian Institution	\$2,979,050	\$0	\$2,979,050
		US Fish and Wildlife Service	\$16,476,494	\$0	\$16,476,494
	<i>Federal Foreign Expenditures subtotal</i>		<i>\$20,267,473</i>	<i>\$0</i>	<i>\$20,267,473</i>
	<i>Federal Domestic and Foreign Expenditures Subtotal</i>		<i>\$1,155,200,685</i>	<i>\$51,681,942</i>	<i>\$1,206,882,627</i>
MultiAgency	Federal Multi-species Expenditures	DOI Bureau of Reclamation	\$9,677,500	\$0	\$9,677,500

Agency Type	Expense Type	Agency	Species Total	Land Total	Cumulative Total
		DOI US Geological Survey	\$22,809	\$0	\$22,809
		USDA Animal and Plant Health Inspection Service	\$256,150	\$0	\$256,150
		DOD Army	\$25,602,301	\$0	\$25,602,301
		DOD Army Corps of Engineers	\$1,297,157	\$0	\$1,297,157
		Environmental Protection Agency	\$4,289,552	\$0	\$4,289,552
		DOT Federal Aviation Administration	\$17,500	\$0	\$17,500
		DOD Navy	\$288,622	\$0	\$288,622
		Nuclear Regulatory Commission	\$29,500	\$0	\$29,500
		USDA Natural Resources Conservation Service	\$1,903,515	\$0	\$1,903,515
		DHS Coast Guard	\$33,847,738	\$0	\$33,847,738
		DOE Southwestern Power Administration	\$9,700	\$0	\$9,700
		Federal Communications Commission	\$155,000	\$0	\$155,000
		DHS Federal Emergency Management Agency	\$75,000	\$0	\$75,000
		US Fish and Wildlife Service	\$3,625,640	\$0	\$3,625,640
		<i>Federal Multi-species Expenditures subtotal</i>	<i>\$81,097,684</i>	<i>\$0</i>	<i>\$81,097,684</i>
		Federal Domestic and Foreign Expenditures Subtotal	\$1,236,298,369	\$51,681,942	\$1,287,980,311
State	State Domestic Expenditures	State Governments	\$44,370,058	\$4,585,524	\$48,955,582
	State Multi-species Expenditures	State Governments	\$8,704,696	\$471,323	\$9,176,019
		State Domestic and Foreign Expenditures Subtotal	\$53,074,754	\$5,056,847	\$58,131,601
		Total Federal and State Domestic and Foreign Expenditures	\$1,289,373,123	\$56,738,789	\$1,346,111,912


Appendix A. Supplemental Guidance for Reporting Expenditures for the Conservation of Endangered and Threatened Species -- FY 2017

This reporting guidance provides information on how to report expenditures attributable to individual species to the extent this information is reasonably available. The Congress requested information for this report on a species-by-species basis, but to provide the most accurate account of expenditures for the conservation of species listed under the Endangered Species Act, a new reporting feature was added in FY 2012. This “multi-species” option allows the reporting of lump sum costs for multi-species conservation activities **when it is not possible to account accurately for individual species.**

Each agency must enter data directly into our Web-based system and will need to designate a contact person (more than one person is permissible) to enter the data. The contact person(s) will need to obtain a username and password before entering the data. Credentials can be obtained by accessing our Environmental Conservation Online System (ECOS) home page at <http://ecos.fws.gov/> and clicking on “Contact Us” located at the bottom of the ECOS home page. This brings you the Help Desk where you will click on “help desk web form.”



Fill in the requested information along with your affiliated agency and request access to the “Expenditures” application.



The screenshot shows a web browser window with the URL <https://ecos.fws.gov/ecp/helpTickets/create>. The page header includes the U.S. Fish & Wildlife Service logo and the text "ECOS Environmental Conservation Online System" and "Conserving the Nature of America". The main content area is titled "ECOS Help Desk" and contains a form with the following fields:

- Name: * John Doe
- Email: * john_doe@usgs.gov
- Phone: * 555-555-5555
- Issue Title: * Need to enter expenditures
- Issue Description: * I am John Doe with the U.S. Geological Survey and I need access to the Expenditures module.

To the right of the form is a box titled "Are you contacting the correct help desk?" with the following text: "Note that this form will contact the *web page technical and program staff*. If you have a question about the FWS or Wildlife in general, [contact the FWS help desk](#)."

Please allow 2 working days for access approval (an email confirmation will be sent to you).

Once you receive your log-in credentials and access approval email, log in at

<https://ecos.fws.gov> to begin entering expenditures data. Contact the ECOS Help Desk (970-266-2999) if you experience any problems.

The following information should be used to gather the data.

BACKGROUND

- Section 18 of the Endangered Species Act of 1973 (16 U.S. 1531 *et seq.*) reads, in part, “...the Fish and Wildlife Service, shall submit to the Congress an annual report covering the preceding fiscal year which shall contain--
 - (1) an accounting on a species-by-species basis of all reasonably identifiable Federal expenditures made primarily for the conservation of endangered and threatened species pursuant to this Act; and
 - (2) an accounting on a species-by-species basis for all reasonably identifiable expenditures made primarily for the conservation of endangered and threatened species pursuant to this Act by States receiving grants under section 6.”
- The conference report for the 1988 amendments made some of the following points concerning section 18 of the Act:
 - Provision should not become unduly burdensome or result in diversion of funds from the operation of the program.

- The Secretary of the Interior is to make a good faith effort to develop and obtain data but is not to undertake extensive or extraordinary measures.
- Generalized dollar estimates will suffice.
- Amendment does not apply to expenditures for delisted, proposed, and candidate species.
- Amendment is not to require new species-specific time sheets for the Service or other reporting agency.
- Listing, consultation, and law enforcement are to be included to the extent they are "reasonably identifiable."
- Only salary costs that can be reasonably identified as applicable to listed species need be reported.
- For other Federal and State agencies, a good faith effort to collect information is expected.
- Submission of data from any agency is not to be a precondition to receiving any contracts or grants or establishing other arrangements with the Fish and Wildlife Service.
- Only one data set per agency should be submitted for any given fiscal year.

DATA FORMAT

- Species should be identified by the same scientific name as found in the most current list of endangered and threatened species.
- Amounts should be reported in whole dollars. Amounts above \$2,000 may be rounded to the nearest \$500 or \$1,000; smaller amounts may be rounded to the nearest \$100. **Do not report amounts lower than \$100.**

EXPENDITURES TO BE REPORTED

- Only species on the Federal list of Endangered and Threatened Wildlife and Plants (50 CFR Part 17) at the end of Federal fiscal year 2017 (October 1, 2016 to September 30, 2017) are to be reported. Expenditures made prior to the actual listing date of a species, but still within the same year, may be reported (e.g., costs of public meetings, notices, surveys, initial recovery efforts). Funds expended for unlisted, separate populations of listed vertebrates cannot be included in the report (e.g., Atlantic coast least terns, Alaska gray wolves). Expenditures for State-listed species are not reportable unless they are also federally listed. Amounts for foreign species on the list (e.g., grants or contracts carried

out in another country) are reportable in the same manner as domestic species, but again, to the extent they are reasonably identifiable.

- Only reasonably identifiable expenditures for listed species will be compiled for this report. Extraordinary accounting to track funds expended on listed species is not expected. **Funds reported should be those actually expended, not obligated.**
- Examples of reportable expenditures include status surveys, habitat management, habitat acquisition, research, propagation (including surrogate species), recovery plan development or implementation, and mitigation. **Projects must primarily benefit listed species** and not other conservation goals, although if a readily identifiable portion of the expenditure is for the benefit of listed species (either individually or collectively), that portion of the expenditure may be reported.
- All habitat (land or water) acquisition costs are to be reported separately from all other identifiable expenditures. Such acquisitions must be primarily for the purpose of conserving federally listed species. As with other projects, those portions of habitat acquisition costs that are specifically for a listed species may be prorated out of the total acquisition costs when the other habitat is being acquired for other purposes (e.g., 25 acres of a 250-acre purchase will be set aside).
- Expenditures in a single project devoted to a number of listed species should be prorated by species, if possible. **When it is not possible to determine individual listed species costs in a multi-species project by prorating, those costs should be entered, in total, in the multi-species category option. The comment box, which is part of this feature, should be used to include information about the project.**
- General surveys or projects that cover a large number of species, some of which may not be listed, are reportable only to extent that the agency can reasonably identify the portion of costs that benefit listed species.
- Expenditures associated with consultations pursuant to section 7 of the Act should be reported as for the benefit of a specific species to the extent they are readily identifiable to that particular species. Thus, a formal consultation dealing with a single species, or up to several species where the relative costs are easily divisible, should be reported under that or those species. **Consultation costs not readily attributable to individual species can be reported in the multi-species category.**
- Monitoring and survey costs should be reported when: (1) the cost of the monitoring or survey was incurred during the period the species was under a proposed listing and the species was actually listed in the same fiscal year that the monitoring and surveys were conducted; (2) the monitoring or survey covered both candidates and listed species, and the costs can be prorated to include only the listed species; or (3) the cost was readily identifiable to individual species. **Costs meeting the criteria in (1) and (2), but not readily identifiable to individual species can be reported in the multi-species category.**

- Salary and benefits of an employee working full-time on a single species or whose time is devoted to particular species that can be readily identified should be reported for those species. Travel costs should be treated similarly. **Staff costs for work on multi-species activities that cannot be identified to particular species can be reported in the multi-species category.**
- Any State or Federal project that incurs increased costs related directly to mitigation or other conservation efforts on behalf of federally listed species should report that added cost, by species, if readily identifiable, or **may report such costs in the multi-species category.**
- All State agencies (e.g., fish and wildlife, parks, heritage programs, forests, highways) should report their expenditures, but only a single, collective report from each State should be submitted. **Expenditures should include conservation costs for fish, wildlife, and plants.** For ESA section 6 grants or other Federal grants, States should report **only the State cost-share portion** of each grant. States should not include other sources of non-Federal cost share, e.g., third-party contributions, unless those contributions are donated to the State as a general donation.
- For reimbursable funds, the paying or disbursing agency should report the expenditure.
- **Litigation costs should not be reported.**

Appendix B. Reports Provided by Other Federal Agencies for Fiscal Year 2017

Department of Agriculture:

- Animal and Plant Health Inspection Service
- Farm Service Agency*
- Forest Service
- Natural Resources Conservation Service

Department of Commerce:

- National Oceanic and Atmospheric Administration

Department of Defense:

- Office of the Under Secretary
 - Representing: Air Force, Army (includes Defense Logistics Agency), Marine Corps, Navy, and Army Corps of Engineers

Department of Energy:

- Bonneville Power Administration
- Southeastern Power Administration*
- Southwestern Power Administration
- Western Area Power Administration

Department of Homeland Security

- Coast Guard
- Customs and Border Protection
- Federal Emergency Management Agency
- Office of Science and Technology*

Department of the Interior:

- Bureau of Indian Affairs
- Bureau of Land Management
- Bureau of Ocean Energy Management*
- Bureau of Reclamation
- Bureau of Safety and Environmental Enforcement*
- National Park Service
- Office of Surface Mining*
- U.S. Geological Survey

Department of Transportation:

- Federal Aviation Administration
- Federal Highway Administration*

Independent Agencies:

Environmental Protection Agency
Federal Communications Commission
Federal Energy Regulatory Commission
Nuclear Regulatory Commission
Smithsonian Institution
Tennessee Valley Authority

* Agencies reporting no expenditures for FY 2017

DHS Coast Guard - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Otter, southern sea	Enhydra lutris nereis	Wherever found	\$150,000	\$0	\$150,000
		Sea lion, Steller	Eumetopias jubatus	Western DPS	\$150,000	\$0	\$150,000
		Whale, killer	Orcinus orca	Southern Resident DPS	\$60,000	\$0	\$60,000
		Whale, North Atlantic Right	Eubalaena glacialis	Wherever found	\$20,000	\$0	\$20,000
	Mammals subtotal				\$380,000	\$0	\$380,000
	Birds	Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR, WA)	\$5,000	\$0	\$5,000
		Tern, California least	Sterna antillarum browni	Wherever found	\$15,000	\$0	\$15,000
	Birds subtotal				\$20,000	\$0	\$20,000
	Reptiles	Sea turtle, green	Chelonia mydas	Central South Pacific DPS	\$2,500	\$0	\$2,500
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$2,500	\$0	\$2,500
		Whipsnake (=striped racer), Alameda	Masticophis lateralis euryxanthus	Wherever found	\$5,000	\$0	\$5,000
	Reptiles subtotal				\$10,000	\$0	\$10,000
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$7,500	\$0	\$7,500
		Salamander, California tiger	Ambystoma californiense	U.S.A. (CA - Santa Barbara County)	\$7,500	\$0	\$7,500
	Amphibians subtotal				\$15,000	\$0	\$15,000
	Fishes	Eulachon	Thaleichthys pacificus	Southern DPS	\$37,000	\$0	\$37,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$37,000	\$0	\$37,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$37,000	\$0	\$37,000
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$37,000	\$0	\$37,000
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$100,000	\$0	\$100,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$40,000	\$0	\$40,000
		sturgeon, green	Acipenser medirostris	Southern DPS	\$37,000	\$0	\$37,000
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$40,000	\$0	\$40,000
		Trout, Paiute cutthroat	Oncorhynchus clarkii seleniris	Wherever found	\$5,000	\$0	\$5,000
	Fishes subtotal				\$370,000	\$0	\$370,000
	Multi-Species				\$33,847,738	\$0	\$33,847,738
	Multi-Species subtotal				\$33,847,738	\$0	\$33,847,738
	Domestic subtotal				\$34,642,738	\$0	\$34,642,738
Total					\$34,642,738	\$0	\$34,642,738

DHS Customs and Border Protection - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, lesser long-nosed	Leptonycteris curasoae yerbabuenae	Wherever found	\$3,139	\$0	\$3,139
	Mammals subtotal				\$3,139	\$0	\$3,139
	Birds	Gnatcatcher, coastal California	Polioptila californica californica	Wherever found	\$54,000	\$0	\$54,000
		Rail, Yuma clapper	Rallus longirostris yumanensis	Wherever found	\$19,000	\$0	\$19,000
	Birds subtotal				\$73,000	\$0	\$73,000
	Amphibians	Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$13,000	\$0	\$13,000
	Amphibians subtotal				\$13,000	\$0	\$13,000
	Insects	Butterfly, Quino checkerspot	Euphydryas editha quino (=E. e. wrighti)	Wherever found	\$33,000	\$0	\$33,000
	Insects subtotal				\$33,000	\$0	\$33,000
	Flowering Plants	Cactus, Pima pineapple	Coryphantha scheeri var. robustispina	Wherever found	\$27,946	\$0	\$27,946
	Flowering Plants subtotal				\$27,946	\$0	\$27,946
Domestic subtotal					\$150,085	\$0	\$150,085
Total					\$150,085	\$0	\$150,085

DHS Federal Emergency Management Agency - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Birds	Cuckoo, yellow-billed	Coccyzus americanus	Western U.S. DPS	\$1,110	\$0	\$1,110
		Warbler (=wood), golden-cheeked	Dendroica chrysoparia	Wherever found	\$29,000	\$0	\$29,000
	Birds subtotal				\$30,110	\$0	\$30,110
	Fishes	Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$2,000	\$0	\$2,000
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$3,000	\$0	\$3,000
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$3,000	\$0	\$3,000
	Fishes subtotal				\$8,000	\$0	\$8,000
	Clams	Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$400	\$0	\$400
	Clams subtotal				\$400	\$0	\$400
	Flowering Plants	Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$1,500	\$0	\$1,500
	Flowering Plants subtotal				\$1,500	\$0	\$1,500
	Multi-Species				\$75,000	\$0	\$75,000
	Multi-Species subtotal				\$75,000	\$0	\$75,000
	Domestic subtotal				\$115,010	\$0	\$115,010
Total					\$115,010	\$0	\$115,010

DOC National Oceanic and Atmospheric Administration - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Sea lion, Steller	<i>Eumetopias jubatus</i>	Western DPS	\$2,058,000	\$0	\$2,058,000
		Seal, Guadalupe fur	<i>Arctocephalus townsendi</i>	Wherever found	\$190,000	\$0	\$190,000
		Seal, Hawaiian monk	<i>Monachus schauinslandi</i>	Wherever found	\$3,896,000	\$0	\$3,896,000
		whale, beluga	<i>Delphinapterus leucas</i>	Cook Inlet DPS	\$2,772,000	\$0	\$2,772,000
		Whale, blue	<i>Balaenoptera musculus</i>	Wherever found	\$1,089,000	\$0	\$1,089,000
		Whale, bowhead	<i>Balaena mysticetus</i>	Wherever found	\$505,000	\$0	\$505,000
		whale, false killer	<i>Pseudorca crassidens</i>	Main Hawaiian Islands Insular DPS	\$366,000	\$0	\$366,000
		Whale, finback	<i>Balaenoptera physalus</i>	Wherever found	\$1,089,000	\$0	\$1,089,000
		Whale, humpback	<i>Megaptera novaeangliae</i>	Central America DPS	\$845,000	\$0	\$845,000
		Whale, humpback	<i>Megaptera novaeangliae</i>	Mexico DPS	\$480,000	\$0	\$480,000
		Whale, humpback	<i>Megaptera novaeangliae</i>	Western North Pacific	\$505,000	\$0	\$505,000
		Whale, killer	<i>Orcinus orca</i>	Southern Resident DPS	\$1,528,000	\$0	\$1,528,000
		Whale, North Atlantic Right	<i>Eubalaena glacialis</i>	Wherever found	\$7,322,000	\$0	\$7,322,000
		Whale, North Pacific Right	<i>Eubalaena japonica</i>	Wherever found	\$505,000	\$0	\$505,000
		Whale, Sei	<i>Balaenoptera borealis</i>	Wherever found	\$1,122,000	\$0	\$1,122,000
		Whale, sperm	<i>Physeter catodon</i> (= <i>macrocephalus</i>)	Wherever found	\$1,267,000	\$0	\$1,267,000
	Mammals subtotal				\$25,539,000	\$0	\$25,539,000
	Reptiles	Sea turtle, green	<i>Chelonia mydas</i>	Central North Pacific DPS	\$960,000	\$0	\$960,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, green	Chelonia mydas	Central South Pacific DPS	\$728,000	\$0	\$728,000
Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, green	Chelonia mydas	Central West Pacific DPS	\$1,285,000	\$0	\$1,285,000
		Sea turtle, green	Chelonia mydas	East Pacific DPS	\$1,349,000	\$0	\$1,349,000
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$1,033,000	\$0	\$1,033,000
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$2,096,000	\$0	\$2,096,000
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$1,290,000	\$0	\$1,290,000
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$2,872,000	\$0	\$2,872,000
		Sea turtle, loggerhead	Caretta caretta	North Pacific Ocean DPS	\$1,914,000	\$0	\$1,914,000
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$2,087,000	\$0	\$2,087,000
		Sea turtle, olive ridley	Lepidochelys olivacea	Wherever found, except when listed as endangered under 50 CFR 224.101	\$603,000	\$0	\$603,000
	Reptiles subtotal				\$16,217,000	\$0	\$16,217,000
	Fishes	Bocaccio	Sebastes paucispinis	Puget Sound - Georgia Basin DPS	\$556,000	\$0	\$556,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Eulachon	Thaleichthys pacificus	Southern DPS	\$556,000	\$0	\$556,000
		grouper, Nassau	Epinephelus striatus		\$452,000	\$0	\$452,000
		rockfish, canary	Sebastes pinniger	Puget Sound - Georgia Basin DPS	\$556,000	\$0	\$556,000
		rockfish, yelloweye	Sebastes ruberrimus	Puget Sound - Georgia Basin DPS	\$556,000	\$0	\$556,000
		Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$6,186,000	\$0	\$6,186,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$3,747,000	\$0	\$3,747,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$3,747,000	\$0	\$3,747,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$5,379,000	\$0	\$5,379,000
Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$4,530,000	\$0	\$4,530,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter- run ESU	\$3,747,000	\$0	\$3,747,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$4,988,000	\$0	\$4,988,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer -run ESU	\$6,320,000	\$0	\$6,320,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$3,755,000	\$0	\$3,755,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$3,644,000	\$0	\$3,644,000
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$5,136,000	\$0	\$5,136,000
		Salmon, chum	Oncorhynchus keta	Hood Canal summer-run ESU	\$4,530,000	\$0	\$4,530,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$3,747,000	\$0	\$3,747,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$5,379,000	\$0	\$5,379,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$4,839,000	\$0	\$4,839,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$5,419,000	\$0	\$5,419,000
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU	\$4,530,000	\$0	\$4,530,000
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$3,349,000	\$0	\$3,349,000
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$651,000	\$0	\$651,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$3,747,000	\$0	\$3,747,000
Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	Central California Coast DPS	\$3,747,000	\$0	\$3,747,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$5,379,000	\$0	\$5,379,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$5,229,000	\$0	\$5,229,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Northern California DPS	\$4,072,000	\$0	\$4,072,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$4,530,000	\$0	\$4,530,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$6,320,000	\$0	\$6,320,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$3,747,000	\$0	\$3,747,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$3,747,000	\$0	\$3,747,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$3,644,000	\$0	\$3,644,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$3,644,000	\$0	\$3,644,000
		Sturgeon (Gulf subspecies), Atlantic	Acipenser oxyrinchus (=oxyrinchus) desotoi	Wherever found	\$518,000	\$0	\$518,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Carolina DPS	\$452,000	\$0	\$452,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Chesapeake Bay DPS	\$1,015,000	\$0	\$1,015,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Gulf of Maine DPS	\$530,000	\$0	\$530,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	New York Bight DPS	\$672,000	\$0	\$672,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	South Atlantic DPS	\$697,000	\$0	\$697,000
		sturgeon, green	Acipenser medirostris	Southern DPS	\$658,000	\$0	\$658,000
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$918,000	\$0	\$918,000
	Fishes subtotal				\$139,565,000	\$0	\$139,565,000
	Snails	Abalone, Black	Haliotis cracherodii	Wherever found	\$658,000	\$0	\$658,000
Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Abalone, White	Haliotis sorenseni	North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	\$1,862,000	\$0	\$1,862,000
	Snails subtotal				\$2,520,000	\$0	\$2,520,000
	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$452,000	\$0	\$452,000
		Coral, elkhorn	Acropora palmata	Wherever found	\$452,000	\$0	\$452,000
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$452,000	\$0	\$452,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$452,000	\$0	\$452,000
		Coral, Pillar	Dendrogyra cylindrus	Wherever found	\$452,000	\$0	\$452,000
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$452,000	\$0	\$452,000
		Coral, staghorn	Acropora cervicornis	Wherever found	\$452,000	\$0	\$452,000
		No common name	Acropora globiceps	Wherever found	\$340,000	\$0	\$340,000
		No common name	Acropora jacquelineae	Wherever found	\$340,000	\$0	\$340,000
		No common name	Acropora retusa	Wherever found	\$190,000	\$0	\$190,000
		No common name	Acropora speciosa	Wherever found	\$340,000	\$0	\$340,000
		No common name	Euphyllia paradivisa	Wherever found	\$340,000	\$0	\$340,000
		No common name	Isopora crateriformis	Wherever found	\$340,000	\$0	\$340,000
		No common name	Seriatopora aculeata	Wherever found	\$340,000	\$0	\$340,000
	Corals subtotal				\$5,394,000	\$0	\$5,394,000
	Flowering Plants	Seagrass, Johnson's	Halophila johnsonii	Wherever found	\$452,000	\$0	\$452,000
	Flowering Plants subtotal				\$452,000	\$0	\$452,000
Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic subtotal					\$189,687,000	\$0	\$189,687,000
Foreign	Reptiles	Sea turtle, green	Chelonia mydas	East Indian- West Pacific DPS	\$170,000	\$0	\$170,000
	Reptiles subtotal				\$170,000	\$0	\$170,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Foreign subtotal					\$170,000	\$0	\$170,000
Total					\$189,857,000	\$0	\$189,857,000

DOD Air Force - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	bat, Florida bonneted	Eumops floridanus	Wherever found	\$200,567	\$0	\$200,567
		Bat, gray	Myotis grisescens	Wherever found	\$79,666	\$0	\$79,666
		Bat, Indiana	Myotis sodalis	Wherever found	\$222,116	\$0	\$222,116
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$79,666	\$0	\$79,666
		Bear, polar	Ursus maritimus	Wherever found	\$280,000	\$0	\$280,000
		Manatee, West Indian	Trichechus manatus	Wherever found	\$2,000	\$0	\$2,000
		Mouse, Choctawhatchee beach	Peromyscus polionotus allophrys	Wherever found	\$47,250	\$0	\$47,250
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	wherever found	\$334,000	\$0	\$334,000
		Mouse, southeastern beach	Peromyscus polionotus niveiventris	wherever found	\$15,000	\$0	\$15,000
		Mouse, St. Andrew beach	Peromyscus polionotus peninsularis	Wherever found	\$47,250	\$0	\$47,250
		Otter, southern sea	Enhydra lutris nereis	Wherever found	\$22,500	\$0	\$22,500
		Panther, Florida	Puma (=Felis) concolor coryi	Wherever found	\$98,000	\$0	\$98,000
		Sea lion, Steller	Eumetopias jubatus	Western DPS	\$58,771	\$0	\$58,771
		whale, beluga	Delphinapterus leucas	Cook Inlet DPS	\$423,673	\$0	\$423,673
	Mammals subtotal				\$1,910,459	\$0	\$1,910,459
	Birds	Caracara, Audubon's crested	Polyborus plancus audubonii	FL pop.	\$90,994	\$0	\$90,994

		Condor, California	<i>Gymnogyps californianus</i>	U.S.A. only, except where listed as an experimental population	\$5,000	\$0	\$5,000
		Coot, Hawaiian	<i>Fulica americana alai</i>	Wherever found	\$52,200	\$0	\$52,200
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$28,646	\$0	\$28,646
		Duck, Hawaiian (=koloa)	<i>Anas wyvilliana</i>	Wherever found	\$52,200	\$0	\$52,200
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$192,500	\$0	\$192,500
		Kite, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	Wherever found	\$210,516	\$0	\$210,516
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$57,500	\$0	\$57,500
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$3,000	\$0	\$3,000
		petrel, Hawaiian	<i>Pterodroma sandwichensis</i>	Wherever found	\$120,000	\$0	\$120,000
		Plover, piping	<i>Charadrius melodus</i>	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$57,500	\$0	\$57,500

		Plover, western snowy	<i>Charadrius nivosus nivosus</i>	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$529,000	\$0	\$529,000
		scrub-jay, Florida	<i>Aphelocoma coerulescens</i>	Wherever found	\$1,067,509	\$0	\$1,067,509
		Shearwater, Newell's Townsend's	<i>Puffinus auricularis newelli</i>	Wherever found	\$33,333	\$0	\$33,333
		Sparrow, Florida grasshopper	<i>Ammodramus savannarum floridanus</i>	Wherever found	\$180,589	\$0	\$180,589
		Stilt, Hawaiian	<i>Himantopus mexicanus knudseni</i>	Wherever found	\$52,200	\$0	\$52,200
		Stork, wood	<i>Mycteria americana</i>	AL, FL, GA, MS, NC, SC	\$312,017	\$0	\$312,017
		Tern, California least	<i>Sterna antillarum browni</i>	Wherever found	\$321,000	\$0	\$321,000
		Tern, least	<i>Sterna antillarum</i>	interior pop.	\$975	\$0	\$975
		Warbler (=wood), golden-cheeked	<i>Dendroica chrysoparia</i>	Wherever found	\$200,000	\$0	\$200,000
		Woodpecker, red-cockaded	<i>Picoides borealis</i>	Wherever found	\$1,347,489	\$0	\$1,347,489
	Birds subtotal				\$4,914,168	\$0	\$4,914,168
	Reptiles	Alligator, American	<i>Alligator mississippiensis</i>	Wherever found	\$144,161	\$0	\$144,161
		Boa, Puerto Rican	<i>Epicrates inornatus</i>	Wherever found	\$70,000	\$0	\$70,000
		Sea turtle, green	<i>Chelonia mydas</i>	South Atlantic DPS	\$152,210	\$0	\$152,210

		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$2,500	\$0	\$2,500
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$70,150	\$0	\$70,150
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$152,210	\$0	\$152,210
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$184,710	\$0	\$184,710
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$724,264	\$0	\$724,264
		Snake, giant garter	Thamnophis gigas	Wherever found	\$39,211	\$0	\$39,211
		Snake, San Francisco garter	Thamnophis sirtalis tetrataenia	Wherever found	\$16,967	\$0	\$16,967
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$669,325	\$0	\$669,325
	Reptiles subtotal				\$2,225,708	\$0	\$2,225,708
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$767,466	\$0	\$767,466
		Salamander, California tiger	Ambystoma californiense	U.S.A. (Central CA DPS)	\$469,965	\$0	\$469,965
		Salamander, frosted flatwoods	Ambystoma cingulatum	Wherever found	\$136,000	\$0	\$136,000
		salamander, Reticulated flatwoods	Ambystoma bishopi	Wherever found	\$463,664	\$0	\$463,664
	Amphibians subtotal				\$1,837,095	\$0	\$1,837,095
	Fishes	Darter, Okaloosa	Etheostoma okaloosae	Wherever found	\$208,372	\$0	\$208,372

		Goby, tidewater	<i>Eucyclogobius newberryi</i>	Wherever found	\$50,000	\$0	\$50,000
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Central California Coast DPS	\$1,100,000	\$0	\$1,100,000
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	South-Central California Coast DPS	\$75,000	\$0	\$75,000
		Stickleback, unarmored threespine	<i>Gasterosteus aculeatus williamsoni</i>	Wherever found	\$85,000	\$0	\$85,000
		Sturgeon (Gulf subspecies), Atlantic	<i>Acipenser oxyrinchus (=oxyrinchus) desotoi</i>	Wherever found	\$67,900	\$0	\$67,900
		Sturgeon, Atlantic	<i>Acipenser oxyrinchus oxyrinchus</i>	Carolina DPS	\$60,000	\$0	\$60,000
	Fishes subtotal				\$1,646,272	\$0	\$1,646,272
	Snails	Abalone, Black	<i>Haliotis cracherodii</i>	Wherever found	\$6,976	\$0	\$6,976
	Snails subtotal				\$6,976	\$0	\$6,976
	Insects	Beetle, valley elderberry longhorn	<i>Desmocerus californicus dimorphus</i>	Wherever found	\$23,693	\$0	\$23,693
		Beetle, [no common name]	<i>Rhadine exilis</i>	Wherever found	\$50,000	\$0	\$50,000
		Beetle, [no common name]	<i>Rhadine infernalis</i>	Wherever found	\$10,000	\$0	\$10,000
		Butterfly, El Segundo blue	<i>Euphilotes battoides allyni</i>	Wherever found	\$253,000	\$0	\$253,000
	Insects subtotal				\$336,693	\$0	\$336,693
	Crustaceans	Fairy shrimp, Conservancy	<i>Branchinecta conservatio</i>	Wherever found	\$13,807	\$0	\$13,807
		Fairy shrimp, vernal pool	<i>Branchinecta lynchi</i>	Wherever found	\$515,283	\$0	\$515,283

		Tadpole shrimp, vernal pool	<i>Lepidurus packardii</i>	Wherever found	\$250,283	\$0	\$250,283
	Crustaceans subtotal				\$779,373	\$0	\$779,373
	Flowering Plants	Beaked-rush, Knieskern's	<i>Rhynchospora knieskernii</i>	Wherever found	\$20,000	\$0	\$20,000
		Butterfly plant, Colorado	<i>Gaura neomexicana</i> var. <i>coloradensis</i>	Wherever found	\$27,000	\$0	\$27,000
		Gerardia, sandplain	<i>Agalinis acuta</i>	Wherever found	\$2,000	\$0	\$2,000
		Goldfields, Contra Costa	<i>Lasthenia conjugens</i>	Wherever found	\$80,780	\$0	\$80,780
		Layia, beach	<i>Layia carnosae</i>	Wherever found	\$192,500	\$0	\$192,500
		Monkeyflower, Vandenberg	<i>Diplacus vandenbergensis</i>	Wherever found	\$77,500	\$0	\$77,500
		Peppergrass, Slickspot	<i>Lepidium papilliferum</i>		\$181,500	\$0	\$181,500
		Pigeon wings	<i>Clitoria fragrans</i>	Wherever found	\$198,008	\$0	\$198,008
		Spurge, telephus	<i>Euphorbia telephioides</i>	Wherever found	\$70,000	\$0	\$70,000
		Tarplant, Gaviota	<i>Deinandra increscens</i> ssp. <i>villosa</i>	Wherever found	\$50,000	\$0	\$50,000
		Tarplant, Otay	<i>Deinandra (=Hemizonia)</i> <i>conjugens</i>	Wherever found	\$7,000	\$0	\$7,000
		Thistle, La Graciosa	<i>Cirsium loncholepis</i>	Wherever found	\$65,000	\$0	\$65,000
		Watercress, Gambel's	<i>Rorippa gambellii</i>	Wherever found	\$77,500	\$0	\$77,500
		Wireweed	<i>Polygonella basiramia</i>	Wherever found	\$17,848	\$0	\$17,848
		Yerba santa, Lompoc	<i>Eriodictyon capitatum</i>	Wherever found	\$346,500	\$0	\$346,500
	Flowering Plants subtotal				\$1,413,136	\$0	\$1,413,136
	Lichens	Cladonia, Florida perforate	<i>Cladonia perforata</i>	Wherever found	\$50,000	\$0	\$50,000

	Lichens subtotal				\$50,000	\$0	\$50,000
Domestic subtotal					\$15,119,880	\$0	\$15,119,880
Total					\$15,119,880	\$0	\$15,119,880

DOD Army Corps of Engineers - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	bat, Florida bonneted	<i>Eumops floridanus</i>	Wherever found	\$156,624	\$0	\$156,624
		Bat, gray	<i>Myotis grisescens</i>	Wherever found	\$241,164	\$0	\$241,164
		Bat, Hawaiian hoary	<i>Lasiurus cinereus semotus</i>	Wherever found	\$1,446	\$0	\$1,446
		Bat, Indiana	<i>Myotis sodalis</i>	Wherever found	\$703,676	\$0	\$703,676
		Bat, lesser long-nosed	<i>Leptonycteris curasoae yerbabuena</i>	Wherever found	\$9,272	\$0	\$9,272
		Bat, Northern long-eared	<i>Myotis septentrionalis</i>	Wherever found	\$1,480,229	\$0	\$1,480,229
		Bat, Virginia big-eared	<i>Corynorhinus (=Plecotus) townsendii virginianus</i>	Wherever found	\$4,826	\$0	\$4,826
		Bear, polar	<i>Ursus maritimus</i>	Wherever found	\$65,411	\$0	\$65,411
		Deer, key	<i>Odocoileus virginianus clavium</i>	Wherever found	\$3,345	\$0	\$3,345
		Ferret, black-footed	<i>Mustela nigripes</i>	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$1,300	\$0	\$1,300
		Fox, San Joaquin kit	<i>Vulpes macrotis mutica</i>	wherever found	\$84,472	\$0	\$84,472
		Jaguarundi, Gulf Coast	<i>Herpailurus (=Felis) yagouaroundi cacomitli</i>	Wherever found	\$3,700	\$0	\$3,700
		Kangaroo rat, Fresno	<i>Dipodomys nitratoide exilis</i>	Wherever found	\$2,502	\$0	\$2,502

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Kangaroo rat, San Bernardino Merriam's	Dipodomys merriami parvus	Wherever found	\$103,602	\$0	\$103,602
		Kangaroo rat, Stephens'	Dipodomys stephensi (incl. D. cactus)	Wherever found	\$5,917	\$0	\$5,917
		Kangaroo rat, Tipton	Dipodomys nitratoides nitratoides	Wherever found	\$800	\$0	\$800
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$23,247	\$0	\$23,247
		Manatee, West Indian	Trichechus manatus	Wherever found	\$677,891	\$0	\$677,891
		Mouse, Alabama beach	Peromyscus polionotus ammobates	Wherever found	\$3,907	\$0	\$3,907
		Mouse, Choctawhatchee beach	Peromyscus polionotus allophrys	Wherever found	\$224	\$0	\$224
		Mouse, Key Largo cotton	Peromyscus gossypinus allapaticola	Wherever found	\$1,729	\$0	\$1,729
		Mouse, New Mexico meadow jumping	Zapus hudsonius luteus	Wherever found	\$4,310	\$0	\$4,310
		Mouse, Perdido Key beach	Peromyscus polionotus trissyllepsis	Wherever found	\$3,320	\$0	\$3,320
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	wherever found	\$3,600	\$0	\$3,600
		Mouse, salt marsh harvest	Reithrodontomys raviventris	wherever found	\$86,200	\$0	\$86,200
		Mouse, St. Andrew beach	Peromyscus polionotus peninsularis	Wherever found	\$3,093	\$0	\$3,093

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Ocelot	Leopardus (=Felis) pardalis	wherever found	\$3,700	\$0	\$3,700
		Otter, Northern Sea	Enhydra lutris kenyoni	Southwest Alaska DPS	\$23,496	\$0	\$23,496
		Otter, southern sea	Enhydra lutris nereis	Wherever found	\$8,000	\$0	\$8,000
		Panther, Florida	Puma (=Felis) concolor coryi	Wherever found	\$42,185	\$0	\$42,185
		Puma (=cougar), eastern	Puma (=Felis) concolor couguar	Wherever found	\$4,861	\$0	\$4,861
		Rabbit, Lower Keys marsh	Sylvilagus palustris hefneri	Wherever found	\$7,715	\$0	\$7,715
		Rice rat	Oryzomys palustris natator	lower FL Keys	\$7,530	\$0	\$7,530
		Sea lion, Steller	Eumetopias jubatus	Western DPS	\$124,485	\$0	\$124,485
		Seal, Hawaiian monk	Monachus schauinslandi	Wherever found	\$12,695	\$0	\$12,695
		Shrew, Buena Vista Lake ornate	Sorex ornatus relictus	Wherever found	\$1,547	\$0	\$1,547
		Squirrel, Carolina northern flying	Glaucomys sabrinus coloratus	Wherever found	\$404	\$0	\$404
		whale, beluga	Delphinapterus leucas	Cook Inlet DPS	\$84,131	\$0	\$84,131
		Whale, blue	Balaenoptera musculus	Wherever found	\$5,288	\$0	\$5,288
		whale, false killer	Pseudorca crassidens	Main Hawaiian Islands Insular DPS	\$4,415	\$0	\$4,415
		Whale, finback	Balaenoptera physalus	Wherever found	\$113,783	\$0	\$113,783
		Whale, humpback	Megaptera novaeangliae	Central America DPS	\$2,017	\$0	\$2,017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Whale, humpback	Megaptera novaeangliae	Mexico DPS	\$138,633	\$0	\$138,633
		Whale, humpback	Megaptera novaeangliae	Western North Pacific	\$250	\$0	\$250
		Whale, killer	Orcinus orca	Southern Resident DPS	\$56,525	\$0	\$56,525
		Whale, North Atlantic Right	Eubalaena glacialis	Wherever found	\$336,499	\$0	\$336,499
		Whale, North Pacific Right	Eubalaena japonica	Wherever found	\$5,479	\$0	\$5,479
		Whale, Sei	Balaenoptera borealis	Wherever found	\$9,244	\$0	\$9,244
		Whale, sperm	Physeter catodon (=macrocephalus)	Wherever found	\$32,098	\$0	\$32,098
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$34,725	\$0	\$34,725

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Woodrat, Key Largo	Neotoma floridana smalli	Wherever found	\$1,788	\$0	\$1,788
	Mammals subtotal				\$4,737,300	\$0	\$4,737,300
	Birds	Albatross, short-tailed	Phoebastria (=Diomedea) albatrus	Wherever found	\$1,057	\$0	\$1,057
		Caracara, Audubon's crested	Polyborus plancus audubonii	FL pop.	\$94,651	\$0	\$94,651
		Condor, California	Gymnogyps californianus	U.S.A. only, except where listed as an experimental population	\$11,482	\$0	\$11,482
		Coot, Hawaiian	Fulica americana alai	Wherever found	\$1,446	\$0	\$1,446
		Crane, Mississippi sandhill	Grus canadensis pulla	Wherever found	\$15,953	\$0	\$15,953
		Crane, whooping	Grus americana	U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	\$14,752	\$0	\$14,752

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Crane, whooping	<i>Grus americana</i>	U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	\$23,000	\$0	\$23,000
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$36,720	\$0	\$36,720
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$358,912	\$0	\$358,912
		Duck, Hawaiian (=koloa)	<i>Anas wyvilliana</i>	Wherever found	\$1,446	\$0	\$1,446
		Eider, spectacled	<i>Somateria fischeri</i>	Wherever found	\$67,933	\$0	\$67,933
		Eider, Steller's	<i>Polysticta stelleri</i>	AK breeding pop.	\$98,887	\$0	\$98,887
		elepaio, Oahu	<i>Chasiempis ibidis</i>	Wherever found	\$366	\$0	\$366
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$1,213,333	\$0	\$1,213,333
		gallinule, Hawaiian common	<i>Gallinula galeata sandvicensis</i>	Wherever found	\$1,446	\$0	\$1,446
		Gnatcatcher, coastal California	<i>Polioptila californica californica</i>	Wherever found	\$286,543	\$0	\$286,543
		Goose, Hawaiian	<i>Branta (=Nesochen) sandvicensis</i>	Wherever found	\$400	\$0	\$400
		Horned lark, streaked	<i>Eremophila alpestris strigata</i>	Wherever found	\$564,830	\$0	\$564,830

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Kite, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	Wherever found	\$656,380	\$0	\$656,380
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$228,507	\$0	\$228,507
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$80,512	\$0	\$80,512
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$11,124	\$0	\$11,124
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$4,341	\$0	\$4,341
		Plover, piping	<i>Charadrius melodus</i>	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$5,151,397	\$0	\$5,151,397
		Plover, piping	<i>Charadrius melodus</i>	[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$32,855	\$0	\$32,855

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$257,137	\$0	\$257,137
		Rail, California clapper	Rallus longirostris obsoletus	Wherever found	\$89,602	\$0	\$89,602
		Rail, light-footed clapper	Rallus longirostris levipes	Wherever found	\$8,554	\$0	\$8,554
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$1,600	\$0	\$1,600
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$38,991	\$0	\$38,991
		Shearwater, Newell's Townsend's	Puffinus auricularis newelli	Wherever found	\$200	\$0	\$200
		Sparrow, Cape Sable seaside	Ammodramus maritimus mirabilis	Wherever found	\$210,840	\$0	\$210,840
		Sparrow, Florida grasshopper	Ammodramus savannarum floridanus	Wherever found	\$4,035	\$0	\$4,035
		Stilt, Hawaiian	Himantopus mexicanus knudseni	Wherever found	\$3,646	\$0	\$3,646
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$423,222	\$0	\$423,222
		Storm-petrel, band-rumped	Oceanodroma castro	USA (HI)	\$200	\$0	\$200
		Tern, California least	Sterna antillarum browni	Wherever found	\$50,826	\$0	\$50,826
		Tern, least	Sterna antillarum	interior pop.	\$5,127,853	\$0	\$5,127,853

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Tern, roseate	<i>Sterna dougallii dougallii</i>	northeast U.S. nesting pop.	\$31,340	\$0	\$31,340
		Vireo, Black-capped	<i>Vireo atricapilla</i>	Wherever found	\$31,840	\$0	\$31,840
		Vireo, least Bell's	<i>Vireo bellii pusillus</i>	Wherever found	\$789,231	\$0	\$789,231
		Warbler (=wood), golden-cheeked	<i>Dendroica chrysoparia</i>	Wherever found	\$45,800	\$0	\$45,800
		Warbler, elfin-woods	<i>Setophaga angelae</i>	Wherever found	\$1,000	\$0	\$1,000
		Warbler, Kirtland's	<i>Setophaga kirtlandii</i> (= <i>Dendroica kirtlandii</i>)	Wherever found	\$8,311	\$0	\$8,311
		Woodpecker, ivory-billed	<i>Campephilus principalis</i>	Wherever found	\$100	\$0	\$100
		Woodpecker, red-cockaded	<i>Picoides borealis</i>	Wherever found	\$213,198	\$0	\$213,198
	Birds subtotal				\$16,295,799	\$0	\$16,295,799
	Reptiles	Alligator, American	<i>Alligator mississippiensis</i>	Wherever found	\$1,075	\$0	\$1,075
		Boa, Puerto Rican	<i>Epicrates inornatus</i>	Wherever found	\$3,068	\$0	\$3,068
		Crocodile, American	<i>Crocodylus acutus</i>	U.S.A. (FL)	\$44,372	\$0	\$44,372
		gartersnake, northern Mexican	<i>Thamnophis eques megalops</i>	Wherever found	\$6,954	\$0	\$6,954
		Lizard, blunt-nosed leopard	<i>Gambelia silus</i>	Wherever found	\$13,591	\$0	\$13,591
		Sea turtle, green	<i>Chelonia mydas</i>	Central North Pacific DPS	\$521	\$0	\$521
		Sea turtle, green	<i>Chelonia mydas</i>	East Pacific DPS	\$16,070	\$0	\$16,070

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$232,368	\$0	\$232,368
		Sea turtle, green	Chelonia mydas	South Atlantic DPS	\$491,104	\$0	\$491,104
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$524,549	\$0	\$524,549
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$1,117,652	\$0	\$1,117,652
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$497,177	\$0	\$497,177
		Sea turtle, loggerhead	Caretta caretta	North Pacific Ocean DPS	\$237	\$0	\$237
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$388,161	\$0	\$388,161
		Sea turtle, olive ridley	Lepidochelys olivacea	Wherever found, except when listed as endangered under 50 CFR 224.101	\$6,981	\$0	\$6,981
		Skink, bluetail mole	Eumeces egregius lividus	Wherever found	\$5,028	\$0	\$5,028
		Skink, sand	Neoseps reynoldsi	Wherever found	\$14,887	\$0	\$14,887
		Snake, Atlantic salt marsh	Nerodia clarkii taeniata	Wherever found	\$1,771	\$0	\$1,771
		Snake, black pine	Pituophis melanoleucus lodingi	Wherever found	\$12,861	\$0	\$12,861

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Snake, copperbelly water	Nerodia erythrogaster neglecta	Indiana north of 40 degrees north latitude, Michigan, Ohio	\$2,234	\$0	\$2,234
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$154,983	\$0	\$154,983
		Snake, giant garter	Thamnophis gigas	Wherever found	\$75,027	\$0	\$75,027
		Snake, San Francisco garter	Thamnophis sirtalis tetrataenia	Wherever found	\$57,800	\$0	\$57,800
		Tortoise, desert	Gopherus agassizii	AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	\$1,547	\$0	\$1,547
		Tortoise, gopher	Gopherus polyphemus	West of Mobile and Tombigbee Rivers	\$50,786	\$0	\$50,786
		Turtle, Alabama red-bellied	Pseudemys alabamensis	Wherever found	\$17,488	\$0	\$17,488
		turtle, bog	Clemmys muhlenbergii	U.S.A. (GA, NC, SC, TN, VA)	\$956	\$0	\$956
		turtle, bog	Clemmys muhlenbergii	Wherever found, except GA, NC, SC, TN, VA	\$92,754	\$0	\$92,754

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Turtle, flattened musk	<i>Sternotherus depressus</i>	Black Warrior R. system upstream from Bankhead Dam	\$3,656	\$0	\$3,656
		Turtle, Plymouth Redbelly	<i>Pseudemys rubriventris bangsi</i>	Wherever found	\$171	\$0	\$171
		Turtle, ringed map	<i>Graptemys oculifera</i>	Wherever found	\$12,448	\$0	\$12,448
		Turtle, yellow-blotched map	<i>Graptemys flavimaculata</i>	Wherever found	\$8,549	\$0	\$8,549
		Whipsnake (=striped racer), Alameda	<i>Masticophis lateralis euryxanthus</i>	Wherever found	\$49,800	\$0	\$49,800
	Reptiles subtotal				\$3,906,626	\$0	\$3,906,626
	Amphibians	Frog, California red-legged	<i>Rana draytonii</i>	Wherever found	\$360,632	\$0	\$360,632
		Frog, Chiricahua leopard	<i>Rana chiricahuensis</i>	Wherever found	\$9,158	\$0	\$9,158
		Frog, dusky gopher	<i>Rana sevosa</i>	Wherever found	\$13,734	\$0	\$13,734
		Frog, mountain yellow-legged	<i>Rana muscosa</i>	Northern California DPS	\$945	\$0	\$945
		Frog, Oregon spotted	<i>Rana pretiosa</i>	Wherever found	\$100	\$0	\$100
		Salamander, Barton Springs	<i>Eurycea sosorum</i>	Wherever found	\$100	\$0	\$100
		Salamander, California tiger	<i>Ambystoma californiense</i>	U.S.A. (CA - Sonoma County)	\$45,000	\$0	\$45,000
		Salamander, California tiger	<i>Ambystoma californiense</i>	U.S.A. (Central CA DPS)	\$124,940	\$0	\$124,940

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salamander, frosted flatwoods	Ambystoma cingulatum	Wherever found	\$7,210	\$0	\$7,210
		Salamander, Georgetown	Eurycea naufragia	Wherever found	\$400	\$0	\$400
		Salamander, Jollyville Plateau	Eurycea tonkawae	Wherever found	\$400	\$0	\$400
		Salamander, Red Hills	Phaeognathus hubrichti	Wherever found	\$1,700	\$0	\$1,700
		salamander, Reticulated flatwoods	Ambystoma bishopi	Wherever found	\$1,436	\$0	\$1,436
		Salamander, Salado	Eurycea chisholmensis	Wherever found	\$400	\$0	\$400
		Salamander, Santa Cruz long-toed	Ambystoma macrodactylum croceum	Wherever found	\$6,400	\$0	\$6,400
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$20,374	\$0	\$20,374
		Toad, Houston	Bufo houstonensis	Wherever found	\$2,200	\$0	\$2,200
		toad, Yosemite	Anaxyrus canorus	Wherever found	\$4,800	\$0	\$4,800
	Amphibians subtotal				\$599,929	\$0	\$599,929
	Fishes	Bocaccio	Sebastes paucispinis	Puget Sound - Georgia Basin DPS	\$33,489	\$0	\$33,489
		Cavefish, Ozark	Amblyopsis rosae	Wherever found	\$1,440	\$0	\$1,440
		Chub, bonytail	Gila elegans	Wherever found	\$10,600	\$0	\$10,600
		Chub, Chihuahua	Gila nigrescens	Wherever found	\$3,790	\$0	\$3,790
		Chub, Gila	Gila intermedia	Wherever found	\$2,318	\$0	\$2,318
		Chub, humpback	Gila cypha	Wherever found	\$10,600	\$0	\$10,600
		Chub, slender	Erimystax cahni	Wherever found	\$422	\$0	\$422

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Chub, spotfin	<i>Erimonax monachus</i>	U.S.A. (TN-specified portions of the Tellico River; see 17.84(m)(1)(i))	\$2,020	\$0	\$2,020
		Chub, spotfin	<i>Erimonax monachus</i>	Wherever found, except where listed as an experimental population	\$2,457	\$0	\$2,457
		Chub, Virgin River	<i>Gila seminuda (=robusta)</i>	Wherever found	\$2,400	\$0	\$2,400
		Cui-ui	<i>Chasmistes cujus</i>	Wherever found	\$800	\$0	\$800
		Dace, blackside	<i>Phoxinus cumberlandensis</i>	Wherever found	\$1,432	\$0	\$1,432
		Darter, amber	<i>Percina antesella</i>	Wherever found	\$4,800	\$0	\$4,800
		Darter, bayou	<i>Etheostoma rubrum</i>	Wherever found	\$9,685	\$0	\$9,685
		darter, bluemask	<i>Etheostoma akatulo</i>	Wherever found	\$404	\$0	\$404
		Darter, Cherokee	<i>Etheostoma scotti</i>	Wherever found	\$11,300	\$0	\$11,300
		Darter, Cumberland	<i>Etheostoma susanae</i>	Wherever found	\$404	\$0	\$404
		Darter, diamond	<i>Crystallaria cincotta</i>	Wherever found	\$853	\$0	\$853
		Darter, duskytail	<i>Etheostoma percnum</i>	Wherever found	\$606	\$0	\$606
		Darter, Etowah	<i>Etheostoma etowahae</i>	Wherever found	\$300	\$0	\$300
		Darter, fountain	<i>Etheostoma fonticola</i>	Wherever found	\$600	\$0	\$600
		Darter, goldline	<i>Percina aurolineata</i>	Wherever found	\$4,070	\$0	\$4,070
		Darter, leopard	<i>Percina pantherina</i>	Wherever found	\$29,780	\$0	\$29,780
		Darter, Niangua	<i>Etheostoma nianguae</i>	Wherever found	\$15,906	\$0	\$15,906
		Darter, Okaloosa	<i>Etheostoma okaloosae</i>	Wherever found	\$1,567	\$0	\$1,567
		Darter, snail	<i>Percina tanasi</i>	Wherever found	\$8,967	\$0	\$8,967

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Eulachon	Thaleichthys pacificus	Southern DPS	\$993,761	\$0	\$993,761
		Goby, tidewater	Eucyclogobius newberryi	Wherever found	\$27,280	\$0	\$27,280
		grouper, Nassau	Epinephelus striatus		\$1,867	\$0	\$1,867
		Logperch, Roanoke	Percina rex	Wherever found	\$2,149	\$0	\$2,149
		Madtom, pygmy	Noturus stanauli	Wherever found	\$202	\$0	\$202
		Madtom, Scioto	Noturus trautmani	Wherever found	\$3,412	\$0	\$3,412
		Madtom, yellowfin	Noturus flavipinnis	Wherever found, except where listed as an experimental population	\$422	\$0	\$422
		Minnow, Devils River	Dionda diaboli	Wherever found	\$400	\$0	\$400
		Minnow, loach	Tiaroga cobitis	Wherever found	\$3,790	\$0	\$3,790
		Minnow, Rio Grande Silvery	Hybognathus amarus	Wherever found, except where listed as an experimental population	\$1,242,712	\$0	\$1,242,712
		Pikeminnow (=squawfish), Colorado	Ptychocheilus lucius	Wherever found, except where listed as an experimental population	\$12,200	\$0	\$12,200
		rockfish, canary	Sebastes pinniger	Puget Sound - Georgia Basin DPS	\$1,266	\$0	\$1,266

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		rockfish, yelloweye	Sebastes ruberrimus	Puget Sound - Georgia Basin DPS	\$18,628	\$0	\$18,628
		Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$89,040	\$0	\$89,040
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$117,888	\$0	\$117,888
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$1,034,290	\$0	\$1,034,290
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$5,941,616	\$0	\$5,941,616
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$2,941,154	\$0	\$2,941,154
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter- run ESU	\$417,106	\$0	\$417,106
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$6,842,414	\$0	\$6,842,414
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer -run ESU	\$6,859,336	\$0	\$6,859,336
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$4,390,271	\$0	\$4,390,271
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$38,292,457	\$0	\$38,292,457
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$389,161	\$0	\$389,161
		Salmon, chum	Oncorhynchus keta	Hood Canal summer-run ESU	\$22,198	\$0	\$22,198

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$1,156,421	\$0	\$1,156,421
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$949,408	\$0	\$949,408
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$17,100	\$0	\$17,100
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$2,742,900	\$0	\$2,742,900
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$7,505,859	\$0	\$7,505,859
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$166,886	\$0	\$166,886
		Shark, Scalloped Hammerhead	Sphyrna lewini	Central and Southwest Atlantic DPS	\$1,000	\$0	\$1,000
		Shark, Scalloped Hammerhead	Sphyrna lewini	Indo-West Pacific DPS	\$1,744	\$0	\$1,744
		Shiner, Arkansas River	Notropis girardi	Arkansas River Basin (AR, KS, NM, OK, TX)	\$2,880	\$0	\$2,880
		Shiner, Cahaba	Notropis cahabae	Wherever found	\$1,386	\$0	\$1,386
		Shiner, palezone	Notropis albizonatus	Wherever found	\$404	\$0	\$404
		Shiner, sharpnose	Notropis oxyrhynchus	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$59,816	\$0	\$59,816
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$875,554	\$0	\$875,554
		Spikedace	Meda fulgida	Wherever found	\$3,790	\$0	\$3,790
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$1,425,196	\$0	\$1,425,196
		Steelhead	Oncorhynchus (=Salmo) mykiss	Central California Coast DPS	\$1,110,117	\$0	\$1,110,117
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$949,729	\$0	\$949,729
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$4,888,781	\$0	\$4,888,781
		Steelhead	Oncorhynchus (=Salmo) mykiss	Northern California DPS	\$37,000	\$0	\$37,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$2,725,420	\$0	\$2,725,420
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$7,541,851	\$0	\$7,541,851
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$42,400	\$0	\$42,400
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$18,558	\$0	\$18,558
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$4,393,331	\$0	\$4,393,331

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$5,292,417	\$0	\$5,292,417
		Stickleback, unarmored threespine	Gasterosteus aculeatus williamsoni	Wherever found	\$2,684	\$0	\$2,684
		Sturgeon (Gulf subspecies), Atlantic	Acipenser oxyrinchus (=oxyrinchus) desotoi	Wherever found	\$1,237,895	\$0	\$1,237,895
		Sturgeon, Alabama	Scaphirhynchus suttkusi	Wherever found	\$199	\$0	\$199
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Gulf of Maine DPS	\$455,823	\$0	\$455,823
		sturgeon, green	Acipenser medirostris	Southern DPS	\$1,951,461	\$0	\$1,951,461
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$30,859,511	\$556,043	\$31,415,554
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$393,197	\$0	\$393,197
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$108,894	\$0	\$108,894
		Sucker, June	Chasmistes liorus	Wherever found	\$1,600	\$0	\$1,600
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$12,200	\$0	\$12,200
		Sucker, Santa Ana	Catostomus santaanae	3 CA river basins	\$568,818	\$0	\$568,818
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$3,988,164	\$0	\$3,988,164
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Trout, Lahontan cutthroat	<i>Oncorhynchus clarkii henshawi</i>	Wherever found	\$16,804	\$0	\$16,804
		Woundfin	<i>Plagopterus argentissimus</i>	Wherever found, except where listed as an experimental population	\$1,600	\$0	\$1,600
	Fishes subtotal				\$151,315,928	\$556,043	\$151,871,971
	Clams	Acornshell, southern	<i>Epioblasma othcaloogensis</i>	Wherever found	\$7,742	\$0	\$7,742
		Bankclimber, purple (mussel)	<i>Elliptoideus sloatianus</i>	Wherever found	\$110,721	\$0	\$110,721
		Bean, Choctaw	<i>Villosa choctawensis</i>	Wherever found	\$660	\$0	\$660
		Bean, Cumberland (pearlymussel)	<i>Villosa trabalis</i>	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$2,423	\$0	\$2,423
		Bean, Cumberland (pearlymussel)	<i>Villosa trabalis</i>	Wherever found; Except where listed as Experimental Populations	\$1,326	\$0	\$1,326
		Bean, purple	<i>Villosa perpurpurea</i>	Wherever found	\$220	\$0	\$220
		Bean, rayed	<i>Villosa fabalis</i>	Wherever found	\$11,198	\$0	\$11,198
		Blossom, green (pearlymussel)	<i>Epioblasma torulosa gubernaculum</i>	Wherever found	\$403	\$0	\$403

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Blossom, tubercled (pearlymussel)	Epioblasma torulosa torulosa	Wherever found; Except where listed as Experimental Populations	\$3,614	\$0	\$3,614
		Blossom, turgid (pearlymussel)	Epioblasma turgidula	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$404	\$0	\$404
		Blossom, yellow (pearlymussel)	Epioblasma florentina florentina	Wherever found; Except where listed as Experimental Populations	\$202	\$0	\$202
		Catspaw, white (pearlymussel)	Epioblasma obliquata perobliqua	Wherever found	\$2,078	\$0	\$2,078
		Clubshell	Pleurobema clava	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$404	\$0	\$404

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$39,029	\$0	\$39,029
		Clubshell, black	Pleurobema curtum	Wherever found	\$7,125	\$0	\$7,125
		Clubshell, ovate	Pleurobema perovatum	Wherever found	\$10,802	\$0	\$10,802
		Clubshell, southern	Pleurobema decisum	Wherever found	\$15,066	\$0	\$15,066
		Combshell, Cumberlandian	Epioblasma brevidens	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$2,222	\$0	\$2,222
		Combshell, Cumberlandian	Epioblasma brevidens	Wherever found; Except where listed as Experimental Populations	\$7,345	\$0	\$7,345
		Combshell, southern	Epioblasma penita	Wherever found	\$7,125	\$0	\$7,125
		Combshell, upland	Epioblasma metastrata	Wherever found	\$3,019	\$0	\$3,019
		Elktoe, Appalachian	Alasmidonta raveneliana	Wherever found	\$14,762	\$0	\$14,762
		Elktoe, Cumberland	Alasmidonta atropurpurea	Wherever found	\$1,235	\$0	\$1,235

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Fanshell	<i>Cyprogenia stegaria</i>	U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$808	\$0	\$808
		Fanshell	<i>Cyprogenia stegaria</i>	Wherever found	\$38,707	\$0	\$38,707
		fatmucket, Arkansas	<i>Lampsilis powellii</i>	Wherever found	\$2,445	\$0	\$2,445
		Heelsplitter, Alabama (=inflated)	<i>Potamilus inflatus</i>	Wherever found	\$18,337	\$0	\$18,337
		Heelsplitter, Carolina	<i>Lasmigona decorata</i>	Wherever found	\$52,889	\$0	\$52,889
		Higgins eye (pearlymussel)	<i>Lampsilis higginsii</i>	Wherever found	\$438,320	\$0	\$438,320
		Kidneyshell, fluted	<i>Ptychobranthus subtentum</i>	Wherever found	\$3,662	\$0	\$3,662
		Kidneyshell, southern	<i>Ptychobranthus jonesi</i>	Wherever found	\$1,643	\$0	\$1,643
		Kidneyshell, triangular	<i>Ptychobranthus greenii</i>	Wherever found	\$4,682	\$0	\$4,682
		Mapleleaf, winged	<i>Quadrula fragosa</i>	Wherever found, except where listed as an experimental population	\$25,850	\$0	\$25,850
		Moccasinshell, Alabama	<i>Medionidus acutissimus</i>	Wherever found	\$2,282	\$0	\$2,282

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Moccasinshell, Gulf	Medionidus penicillatus	Wherever found	\$5,500	\$0	\$5,500
		Monkeyface, Appalachian (pearlymussel)	Quadrula sparsa	USA (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$422	\$0	\$422
		Monkeyface, Cumberland (pearlymussel)	Quadrula intermedia	Wherever found; Except where listed as Experimental Populations	\$422	\$0	\$422
		Mucket, Neosho	Lampsilis rafinesqueana	Wherever found	\$7,011	\$0	\$7,011
		Mucket, orangenacre	Lampsilis perovalis	Wherever found	\$10,403	\$0	\$10,403
		Mucket, pink (pearlymussel)	Lampsilis abrupta	Wherever found	\$125,756	\$0	\$125,756
		Mussel, oyster	Epioblasma capsaeformis	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$2,020	\$0	\$2,020

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Mussel, oyster	Epioblasma capsaeformis	Wherever found; Except where listed as Experimental Populations	\$1,546	\$0	\$1,546
		Mussel, scaleshell	Leptodea leptodon	Wherever found	\$21,483	\$0	\$21,483
		Mussel, sheepnose	Plethobasus cyphus	Wherever found	\$124,045	\$0	\$124,045
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$34,827	\$0	\$34,827
		Pearlshell, Louisiana	Margaritifera hembeli	Wherever found	\$1,920	\$0	\$1,920
		Pearlymussel, birdwing	Lemiox rimosus	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert, and Lauderdale Counties, AL	\$202	\$0	\$202
		Pearlymussel, birdwing	Lemiox rimosus	Wherever found; Except where listed as Experimental Populations	\$220	\$0	\$220

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pearlymussel, cracking	Hemistena lata	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$404	\$0	\$404
		Pearlymussel, cracking	Hemistena lata	Wherever found; Except where listed as Experimental Populations	\$220	\$0	\$220
		Pearlymussel, dromedary	Dromus dromas	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$1,212	\$0	\$1,212
		Pearlymussel, dromedary	Dromus dromas	Wherever found; Except where listed as Experimental Populations	\$1,546	\$0	\$1,546
		Pearlymussel, littlewing	Pegias fabula	Wherever found	\$1,230	\$0	\$1,230

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pearlymussel, slabside	<i>Pleuonaia dolabelloides</i>	Wherever found	\$7,749	\$0	\$7,749
		Pigtoe, Cumberland	<i>Pleurobema gibberum</i>	Wherever found	\$1,414	\$0	\$1,414
		Pigtoe, dark	<i>Pleurobema furvum</i>	Wherever found	\$120	\$0	\$120
		Pigtoe, finerayed	<i>Fusconaia cuneolus</i>	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$1,010	\$0	\$1,010
		Pigtoe, finerayed	<i>Fusconaia cuneolus</i>	Wherever found; Except where listed as Experimental Populations	\$220	\$0	\$220
		Pigtoe, fuzzy	<i>Pleurobema strodeanum</i>	Wherever found	\$859	\$0	\$859
		Pigtoe, Georgia	<i>Pleurobema hanleyianum</i>	Wherever found	\$199	\$0	\$199
		Pigtoe, heavy	<i>Pleurobema taitianum</i>	Wherever found	\$7,324	\$0	\$7,324
		Pigtoe, oval	<i>Pleurobema pyriforme</i>	Wherever found	\$8,594	\$0	\$8,594

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pigtoe, rough	Pleurobema plenum	U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$1,010	\$0	\$1,010
		Pigtoe, rough	Pleurobema plenum	Wherever found	\$26,647	\$0	\$26,647
		Pigtoe, shiny	Fusconaia cor	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$606	\$0	\$606
		Pigtoe, shiny	Fusconaia cor	Wherever found; Except where listed as Experimental Populations	\$220	\$0	\$220
		Pigtoe, tapered	Fusconaia burkei	Wherever found	\$261	\$0	\$261
		Pimpleback, orangefoot (pearlymussel)	Plethobasus cooperianus	Wherever found	\$28,333	\$0	\$28,333
		Pocketbook, fat	Potamilus capax	Wherever found	\$65,366	\$0	\$65,366
		Pocketbook, finelined	Lampsilis altilis	Wherever found	\$2,961	\$0	\$2,961
		Pocketbook, Ouachita rock	Arkansia wheeleri	Wherever found	\$3,330	\$0	\$3,330

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pocketbook, shinyrayed	Lampsilis subangulata	Wherever found	\$5,399	\$0	\$5,399
		Purple cat's paw (=purple cat's paw pearlymussel)	Epioblasma obliquata obliquata	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$202	\$0	\$202
		Purple Cat's paw (=Purple Cat's paw pearlymussel)	Epioblasma obliquata obliquata	Wherever found; Except where listed as Experimental Populations	\$2,730	\$0	\$2,730
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$46,655	\$0	\$46,655
		Rabbitsfoot, rough	Quadrula cylindrica strigillata	Wherever found	\$422	\$0	\$422
		Riffleshell, northern	Epioblasma torulosa rangiana	Wherever found	\$12,210	\$0	\$12,210
		Riffleshell, tan	Epioblasma florentina walkeri (=E. walkeri)	Wherever found	\$404	\$0	\$404
		Ring pink (mussel)	Obovaria retusa	Wherever found	\$25,873	\$0	\$25,873
		sandshell, Southern	Hamiota australis	Wherever found	\$261	\$0	\$261
		Slabshell, Chipola	Elliptio chipolaensis	Wherever found	\$106,708	\$0	\$106,708
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$57,542	\$0	\$57,542

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Spnymussel, Altamaha	<i>Elliptio spinosa</i>	Wherever found	\$1,500	\$0	\$1,500
		Spnymussel, Tar River	<i>Elliptio steinstansana</i>	Wherever found	\$1,819	\$0	\$1,819
		Threeridge, fat (mussel)	<i>Amblema neislerii</i>	Wherever found	\$109,221	\$0	\$109,221
		Wartyback, white (pearlymussel)	<i>Plethobasus cicatricosus</i>	Wherever found	\$808	\$0	\$808
		Wedgemussel, dwarf	<i>Alasmidonta heterodon</i>	Wherever found	\$32,754	\$0	\$32,754
	Clams subtotal				\$1,739,870	\$0	\$1,739,870
	Snails	Hornsnail, rough	<i>Pleurocera foremani</i>	Wherever found	\$1,196	\$0	\$1,196
		Lioplax, cylindrical (snail)	<i>Lioplax cyclostomaformis</i>	Wherever found	\$100	\$0	\$100
		Riversnail, Anthony's	<i>Athearnia anthonyi</i>	U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$1,866	\$0	\$1,866
		Rocksnail, interrupted (=Georgia)	<i>Leptoxis foremani</i>	Wherever found	\$199	\$0	\$199
		Rocksnail, painted	<i>Leptoxis taeniata</i>	Wherever found	\$399	\$0	\$399
		Rocksnail, plicate	<i>Leptoxis plicata</i>	Wherever found	\$120	\$0	\$120
		Snail, Bliss Rapids	<i>Taylorconcha serpenticola</i>	Wherever found	\$4,130	\$0	\$4,130
		Snail, Snake River physa	<i>Physa natricina</i>	Wherever found	\$11,052	\$0	\$11,052

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Snail, Stock Island tree	Orthalicus reses (not incl. nesodryas)	Wherever found	\$941	\$0	\$941
		Snail, tulotoma	Tulotoma magnifica	Wherever found	\$399	\$0	\$399
	Snails subtotal				\$20,402	\$0	\$20,402
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$107,065	\$0	\$107,065
		Beetle, Casey's June	Dinacoma caseyi	Wherever found	\$2,684	\$0	\$2,684
		Beetle, Coffin Cave mold	Batrisodes texanus	Wherever found	\$2,750	\$0	\$2,750
		Beetle, Comal Springs dryopid	Stygoparnus comalensis	Wherever found	\$200	\$0	\$200
		Beetle, Comal Springs riffle	Heterelmis comalensis	Wherever found	\$400	\$0	\$400
		Beetle, delta green ground	Elaphrus viridis	Wherever found	\$800	\$0	\$800
		Beetle, Mount Hermon June	Polyphylla barbata	Wherever found	\$11,200	\$0	\$11,200
		Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$173,942	\$0	\$173,942
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$3,760	\$0	\$3,760
		Butterfly, bay checkerspot	Euphydryas editha bayensis	Wherever found	\$4,000	\$0	\$4,000
		Butterfly, callippe silverspot	Speyeria callippe callippe	Wherever found	\$36,200	\$0	\$36,200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Butterfly, Fender's blue	<i>Icaricia icarioides fenderi</i>	Wherever found	\$198,796	\$0	\$198,796
		Butterfly, Karner blue	<i>Lycaeides melissa samuelis</i>	Wherever found	\$1,204	\$0	\$1,204
		Butterfly, Mitchell's satyr	<i>Neonympha mitchellii mitchellii</i>	Wherever found	\$7,455	\$0	\$7,455
		Butterfly, Quino checkerspot	<i>Euphydryas editha quino</i> (=E. e. <i>wrighti</i>)	Wherever found	\$5,002	\$0	\$5,002
		Butterfly, Schaus swallowtail	<i>Heraclides aristodemus ponceanus</i>	Wherever found	\$2,518	\$0	\$2,518
		Butterfly, Smith's blue	<i>Euphilotes enoptes smithi</i>	Wherever found	\$6,400	\$0	\$6,400
		Dragonfly, Hine's emerald	<i>Somatochlora hineana</i>	Wherever found	\$28,000	\$0	\$28,000
		Grasshopper, Zayante band-winged	<i>Trimerotropis infantilis</i>	Wherever found	\$14,400	\$0	\$14,400
		Skipper, Dakota	<i>Hesperia dacotae</i>	Wherever found	\$640	\$0	\$640
		Tiger beetle, Northeastern beach	<i>Cicindela dorsalis dorsalis</i>	Wherever found	\$350	\$0	\$350
		Tiger beetle, Ohlone	<i>Cicindela ohlone</i>	Wherever found	\$4,800	\$0	\$4,800
		Tiger beetle, Puritan	<i>Cicindela puritana</i>	Wherever found	\$1,750	\$0	\$1,750
		Tiger beetle, Salt Creek	<i>Cicindela nevadica lincolniana</i>	Wherever found	\$1,600	\$0	\$1,600
	Insects subtotal				\$615,916	\$0	\$615,916
	Arachnids	Harvestman, Bone Cave	<i>Texella reyesi</i>	Wherever found	\$2,750	\$0	\$2,750

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Meshweaver, Madla's Cave	Cicurina madla	Wherever found	\$200	\$0	\$200
		Spider, spruce- fir moss	Microhexura montivaga	Wherever found	\$404	\$0	\$404
	Arachnids subtotal				\$3,354	\$0	\$3,354
	Crustaceans	Amphipod, Peck's cave	Stygbromus (=Stygonectes) pecki	Wherever found	\$200	\$0	\$200
		Crayfish, Big Sandy	Cambarus callainus	Wherever found	\$140	\$0	\$140
		Crayfish, Nashville	Orconectes shoupi	Wherever found	\$2,330	\$0	\$2,330
		Fairy shrimp, Conservancy	Branchinecta conservatio	Wherever found	\$15,613	\$0	\$15,613
		Fairy shrimp, longhorn	Branchinecta longiantenna	Wherever found	\$35,600	\$0	\$35,600
		Fairy shrimp, Riverside	Streptocephalus woottoni	Wherever found	\$5,368	\$0	\$5,368
		Fairy shrimp, San Diego	Branchinecta sandiegonensis	Wherever found	\$10,370	\$0	\$10,370
		Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$119,993	\$0	\$119,993
		Shrimp, California freshwater	Syncaris pacifica	Wherever found	\$1,600	\$0	\$1,600
		Shrimp, Kentucky cave	Palaemonias ganteri	Wherever found	\$702	\$0	\$702
		Tadpole shrimp, vernal pool	Lepidurus packardi	Wherever found	\$47,890	\$0	\$47,890
	Crustaceans subtotal				\$239,806	\$0	\$239,806
	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$1,614	\$0	\$1,614
		Coral, elkhorn	Acropora palmata	Wherever found	\$2,595	\$0	\$2,595

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Coral, Lobed Star	<i>Orbicella annularis</i>	Wherever found	\$1,625	\$0	\$1,625
		Coral, Mountainous Star	<i>Orbicella faveolata</i>	Wherever found	\$2,134	\$0	\$2,134
		Coral, Pillar	<i>Dendrogyra cylindrus</i>	Wherever found	\$1,601	\$0	\$1,601
		Coral, Rough Cactus	<i>Mycetophyllia ferox</i>	Wherever found	\$1,480	\$0	\$1,480
		Coral, staghorn	<i>Acropora cervicornis</i>	Wherever found	\$2,551	\$0	\$2,551
		No common name	<i>Acropora globiceps</i>	Wherever found	\$1,744	\$0	\$1,744
		No common name	<i>Acropora jacquelineae</i>	Wherever found	\$200	\$0	\$200
		No common name	<i>Acropora retusa</i>	Wherever found	\$200	\$0	\$200
		No common name	<i>Acropora speciosa</i>	Wherever found	\$200	\$0	\$200
		No common name	<i>Euphyllia paradivisa</i>	Wherever found	\$200	\$0	\$200
		No common name	<i>Isopora crateriformis</i>	Wherever found	\$200	\$0	\$200
		No common name	<i>Seriatopora aculeata</i>	Wherever found	\$1,544	\$0	\$1,544
	Corals subtotal				\$17,888	\$0	\$17,888
	Flowering Plants	Amaranth, seabeach	<i>Amaranthus pumilus</i>	Wherever found	\$73,269	\$0	\$73,269
		Ambrosia, San Diego	<i>Ambrosia pumila</i>	Wherever found	\$671	\$0	\$671
		Amphianthus, little	<i>Amphianthus pusillus</i>	Wherever found	\$500	\$0	\$500
		Arrowhead, bunched	<i>Sagittaria fasciculata</i>	Wherever found	\$6,439	\$0	\$6,439

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Aster, decurrent false	<i>Boltonia decurrens</i>	Wherever found	\$75,000	\$0	\$75,000
		Aster, Ruth's golden	<i>Pityopsis ruthii</i>	Wherever found	\$202	\$0	\$202
		Bird's beak, palmate-bracted	<i>Cordylanthus palmatus</i>	Wherever found	\$5,800	\$0	\$5,800
		Bird's-beak, soft	<i>Cordylanthus mollis</i> ssp. <i>mollis</i>	Wherever found	\$27,200	\$0	\$27,200
		bladderpod, Missouri	<i>Physaria filiformis</i>	Wherever found	\$640	\$0	\$640
		Bladderpod, Short's	<i>Physaria globosa</i>	Wherever found	\$9,527	\$0	\$9,527
		Bladderpod, Spring Creek	<i>Lesquerella perforata</i>	Wherever found	\$2,200	\$0	\$2,200
		Brodiaea, thread-leaved	<i>Brodiaea filifolia</i>	Wherever found	\$1,342	\$0	\$1,342
		Bulrush, Northeastern	<i>Scirpus ancistrochaetus</i>	Wherever found	\$3,600	\$0	\$3,600
		Bush-clover, prairie	<i>Lespedeza leptostachya</i>	Wherever found	\$160	\$0	\$160
		Butterweed, Layne's	<i>Senecio layneae</i>	Wherever found	\$800	\$0	\$800
		Button-celery, San Diego	<i>Eryngium aristulatum</i> var. <i>parishii</i>	Wherever found	\$5,368	\$0	\$5,368
		Cactus, Bakersfield	<i>Opuntia treleasei</i>	Wherever found	\$800	\$0	\$800
		Cactus, Key tree	<i>Pilosocereus robinii</i>	Wherever found	\$653	\$0	\$653
		Cactus, Pima pineapple	<i>Coryphantha scheeri</i> var. <i>robustispina</i>	Wherever found	\$5,368	\$0	\$5,368
		Cactus, San Rafael	<i>Pediocactus despainii</i>	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		cactus, Tobusch fishhook	<i>Sclerocactus brevihamatus</i> ssp. <i>tobuschii</i>	Wherever found	\$400	\$0	\$400
		Catchfly, Spalding's	<i>Silene spaldingii</i>	Wherever found	\$3,122	\$0	\$3,122
		Ceanothus, coyote	<i>Ceanothus ferrisiae</i>	Wherever found	\$4,000	\$0	\$4,000
		Chaffseed, American	<i>Schwalbea americana</i>	Wherever found	\$1,000	\$0	\$1,000
		Checker- mallow, Keck's	<i>Sidalcea keckii</i>	Wherever found	\$129	\$0	\$129
		Clarkia, Springville	<i>Clarkia springvillensis</i>	Wherever found	\$4,861	\$0	\$4,861
		Clover, running buffalo	<i>Trifolium stoloniferum</i>	Wherever found	\$48,060	\$0	\$48,060
		Clover, showy Indian	<i>Trifolium amoenum</i>	Wherever found	\$1,600	\$0	\$1,600
		Coneflower, smooth	<i>Echinacea laevigata</i>	Wherever found	\$546	\$0	\$546
		Daisy, Lakeside	<i>Hymenoxys herbacea</i>	Wherever found	\$1,759	\$0	\$1,759
		Daisy, Willamette	<i>Erigeron decumbens</i>	Wherever found	\$207,396	\$0	\$207,396
		Dawn-flower, Texas prairie	<i>Hymenoxys texana</i>	Wherever found	\$5,000	\$0	\$5,000
		Desert-parsley, Bradshaw's	<i>Lomatium bradshawii</i>	Wherever found	\$195,596	\$0	\$195,596
		Dropwort, Canby's	<i>Oxypolis canbyi</i>	Wherever found	\$5,210	\$0	\$5,210
		Dudleya, Santa Clara Valley	<i>Dudleya setchellii</i>	Wherever found	\$4,000	\$0	\$4,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Evening- primrose, San Benito	Camissonia benitensis	Wherever found	\$2,400	\$0	\$2,400
		Fiddleneck, large-flowered	Amsinckia grandiflora	Wherever found	\$1,800	\$0	\$1,800
		Gerardia, sandplain	Agalinis acuta	Wherever found	\$240	\$0	\$240
		Glade cress, Kentucky	Leavenworthia exigua laciniata	Wherever found	\$3,510	\$0	\$3,510
		Goldfields, Burke's	Lasthenia burkei	Wherever found	\$42,600	\$0	\$42,600
		Goldfields, Contra Costa	Lasthenia conjugens	Wherever found	\$9,800	\$0	\$9,800
		Gourd, Okeechobee	Cucurbita okeechobeensis ssp. okeechobeensis	Wherever found	\$500	\$0	\$500
		Grass, Colusa	Neostapfia colusana	Wherever found	\$3,259	\$0	\$3,259
		Grass, Tennessee yellow-eyed	Xyris tennesseensis	Wherever found	\$272	\$0	\$272
		Ground-plum, Guthrie's (=Pyne's)	Astragalus bibullatus	Wherever found	\$1,028	\$0	\$1,028
		Harperella	Ptilimnium nodosum	Wherever found	\$5,710	\$0	\$5,710
		Heartleaf, dwarf-flowered	Hexastylis naniflora	Wherever found	\$13,263	\$0	\$13,263
		Jewelflower, California	Caulanthus californicus	Wherever found	\$1,600	\$0	\$1,600
		Jewelflower, Metcalf Canyon	Streptanthus albidus ssp. albidus	Wherever found	\$2,400	\$0	\$2,400
		Joint-vetch, Sensitive	Aeschynomene virginica	Wherever found	\$220	\$0	\$220

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Ladies'-tresses, Navasota	<i>Spiranthes parksii</i>	Wherever found	\$1,300	\$0	\$1,300
		Ladies'-tresses, Ute	<i>Spiranthes diluvialis</i>	Wherever found	\$35,225	\$0	\$35,225
		Leather flower, Morefield's	<i>Clematis morefieldii</i>		\$202	\$0	\$202
		Loosestrife, rough-leaved	<i>Lysimachia asperulaefolia</i>	Wherever found	\$15,250	\$0	\$15,250
		Lupine, Kincaid's	<i>Lupinus sulphureus</i> ssp. <i>kincaidii</i>	Wherever found	\$162,896	\$0	\$162,896
		Manioc, Walker's	<i>Manihot walkerae</i>	Wherever found	\$160	\$0	\$160
		Manzanita, Ione	<i>Arctostaphylos myrtifolia</i>	Wherever found	\$1,431	\$0	\$1,431
		Manzanita, pallid	<i>Arctostaphylos pallida</i>	Wherever found	\$2,600	\$0	\$2,600
		Meadowfoam, Butte County	<i>Limnanthes floccosa</i> ssp. <i>californica</i>	Wherever found	\$800	\$0	\$800
		Meadowfoam, Sebastopol	<i>Limnanthes vinculans</i>	Wherever found	\$41,800	\$0	\$41,800
		Mesa-mint, Otay	<i>Pogogyne nudiuscula</i>	Wherever found	\$2,684	\$0	\$2,684
		Milk-vetch, Braunton's	<i>Astragalus brauntonii</i>	Wherever found	\$2,684	\$0	\$2,684
		Milk-vetch, Ventura Marsh	<i>Astragalus pycnostachyus</i> var. <i>lanosissimus</i>	Wherever found	\$4,636	\$0	\$4,636
		Milkweed, Mead's	<i>Asclepias meadii</i>	Wherever found	\$5,835	\$0	\$5,835
		Monkshood, northern wild	<i>Aconitum noveboracense</i>	Wherever found	\$969	\$0	\$969
		mountainbalm, Indian Knob	<i>Eriodictyon altissimum</i>	Wherever found	\$2,318	\$0	\$2,318

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Navarretia, many-flowered	Navarretia leucocephala ssp. plieantha	Wherever found	\$2,400	\$0	\$2,400
		Navarretia, spreading	Navarretia fossalis	Wherever found	\$5,368	\$0	\$5,368
		No common name	Geocarpon minimum	Wherever found	\$5,173	\$0	\$5,173
		Orchid, eastern prairie fringed	Platanthera leucophaea	Wherever found	\$8,416	\$0	\$8,416
		Orchid, western prairie fringed	Platanthera praeclara	Wherever found	\$7,319	\$0	\$7,319
		Orcutt grass, California	Orcuttia californica	Wherever found	\$2,684	\$0	\$2,684
		Orcutt grass, hairy	Orcuttia pilosa	Wherever found	\$2,373	\$0	\$2,373
		Orcutt grass, Sacramento	Orcuttia viscida	Wherever found	\$3,200	\$0	\$3,200
		Orcutt grass, San Joaquin	Orcuttia inaequalis	Wherever found	\$2,518	\$0	\$2,518
		Orcutt grass, slender	Orcuttia tenuis	Wherever found	\$2,400	\$0	\$2,400
		Owl's-clover, fleshy	Castilleja campestris ssp. succulenta	Wherever found	\$3,973	\$0	\$3,973
		paintbrush, golden	Castilleja levisecta	Wherever found	\$127,396	\$0	\$127,396
		Paintbrush, Tiburon	Castilleja affinis ssp. neglecta	Wherever found	\$2,400	\$0	\$2,400
		Pentachaeta, white-rayed	Pentachaeta bellidiflora	Wherever found	\$1,600	\$0	\$1,600
		Pink, swamp	Helonias bullata	Wherever found	\$1,228	\$0	\$1,228
		Pinkroot, gentian	Spigelia gentianoides	Wherever found	\$570	\$0	\$570
		Piperia, Yadon's	Piperia yadonii	Wherever found	\$3,200	\$0	\$3,200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pitcher-plant, green	<i>Sarracenia oreophila</i>	Wherever found	\$500	\$0	\$500
		Pitcher-plant, mountain sweet	<i>Sarracenia rubra</i> ssp. <i>jonesii</i>	Wherever found	\$9,254	\$0	\$9,254
		Pogonia, small whorled	<i>Isotria medeoloides</i>	Wherever found	\$13,280	\$0	\$13,280
		Polygonum, Scotts Valley	<i>Polygonum hickmanii</i>	Wherever found	\$9,600	\$0	\$9,600
		Pondberry	<i>Lindera melissifolia</i>	Wherever found	\$19,252	\$0	\$19,252
		Poppy-mallow, Texas	<i>Callirhoe scabriuscula</i>	Wherever found	\$100	\$0	\$100
		Potato-bean, Price's	<i>Apios priceana</i>		\$12,126	\$0	\$12,126
		Prairie-clover, leafy	<i>Dalea foliosa</i>	Wherever found	\$2,434	\$0	\$2,434
		Rock-cress, Braun's	<i>Arabis perstellata</i>	Wherever found	\$3,134	\$0	\$3,134
		Rockcress, Georgia	<i>Arabis georgiana</i>	Wherever found	\$100	\$0	\$100
		Rosemary, Cumberland	<i>Conradina verticillata</i>	Wherever found	\$404	\$0	\$404
		roseroot, Leedy's	<i>Rhodiola integrifolia</i> ssp. <i>leedyi</i>	Wherever found	\$652	\$0	\$652
		Sand-verbena, large-fruited	<i>Abronia macrocarpa</i>	Wherever found	\$100	\$0	\$100
		Sandwort, Cumberland	<i>Arenaria cumberlandensis</i>	Wherever found	\$606	\$0	\$606
		Sandwort, Marsh	<i>Arenaria paludicola</i>	Wherever found	\$6,954	\$0	\$6,954
		Seablite, California	<i>Suaeda californica</i>		\$4,000	\$0	\$4,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Seagrass, Johnson's	<i>Halophila johnsonii</i>	Wherever found	\$90,524	\$0	\$90,524
		Skullcap, large-flowered	<i>Scutellaria montana</i>	Wherever found	\$202	\$0	\$202
		Sneezeweed, Virginia	<i>Helenium virginicum</i>	Wherever found	\$400	\$0	\$400
		Spineflower, Ben Lomond	<i>Chorizanthe pungens</i> var. <i>hartwegiana</i>	Wherever found	\$9,600	\$0	\$9,600
		Spineflower, Monterey	<i>Chorizanthe pungens</i> var. <i>pungens</i>	Wherever found	\$9,600	\$0	\$9,600
		spineflower, Robust	<i>Chorizanthe robusta</i> var. <i>robusta</i>	Wherever found	\$4,000	\$0	\$4,000
		spineflower, Scotts Valley	<i>Chorizanthe robusta</i> var. <i>hartwegii</i>	Wherever found	\$3,200	\$0	\$3,200
		Spineflower, slender-horned	<i>Dodecahema leptoceras</i>	Wherever found	\$2,684	\$0	\$2,684
		Spiraea, Virginia	<i>Spiraea virginiana</i>	Wherever found	\$2,884	\$0	\$2,884
		Spurge, Hoover's	<i>Chamaesyce hooveri</i>	Wherever found	\$145	\$0	\$145
		Spurge, telephus	<i>Euphorbia telephioides</i>	Wherever found	\$3,131	\$0	\$3,131
		Sumac, Michaux's	<i>Rhus michauxii</i>	Wherever found	\$4,564	\$0	\$4,564
		Sunburst, Hartweg's golden	<i>Pseudobahia bahiifolia</i>	Wherever found	\$4,859	\$0	\$4,859
		Sunburst, San Joaquin adobe	<i>Pseudobahia peirsonii</i>	Wherever found	\$12,853	\$0	\$12,853
		Sunflower, San Mateo woolly	<i>Eriophyllum latilobum</i>	Wherever found	\$2,400	\$0	\$2,400
		Sunflower, Schweinitz's	<i>Helianthus schweinitzii</i>	Wherever found	\$11,762	\$0	\$11,762
		Sunshine, Sonoma	<i>Blennosperma bakeri</i>	Wherever found	\$40,200	\$0	\$40,200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Tarplant, Otay	Deinandra (=Hemizonia) conjugens	Wherever found	\$5,368	\$0	\$5,368
		Tarplant, Santa Cruz	Holocarpha macradenia	Wherever found	\$8,800	\$0	\$8,800
		Thistle, La Graciosa	Cirsium loncholepis	Wherever found	\$2,684	\$0	\$2,684
		Thistle, Pitcher's	Cirsium pitcheri	Wherever found	\$2,111	\$0	\$2,111
		Thistle, Suisun	Cirsium hydrophilum var. hydrophilum	Wherever found	\$5,600	\$0	\$5,600
		Thornmint, San Diego	Acanthomintha ilicifolia	Wherever found	\$2,684	\$0	\$2,684
		Thornmint, San Mateo	Acanthomintha obovata ssp. duttonii	Wherever found	\$1,547	\$0	\$1,547
		Trillium, relict	Trillium reliquum	Wherever found	\$15,210	\$0	\$15,210
		Tuctoria, Greene's	Tuctoria greenei	Wherever found	\$145	\$0	\$145
		Wallflower, Ben Lomond	Erysimum teretifolium	Wherever found	\$9,600	\$0	\$9,600
		Watercress, Gambel's	Rorippa gambellii	Wherever found	\$4,636	\$0	\$4,636
		Wild-rice, Texas	Zizania texana	Wherever found	\$600	\$0	\$600
		Woolly-star, Santa Ana River	Eriastrum densifolium ssp. sanctorum	Wherever found	\$9,818	\$0	\$9,818
		Woolly-threads, San Joaquin	Monolopia (=Lembertia) congdonii	Wherever found	\$4,000	\$0	\$4,000
		Flowering Plants subtotal			\$1,612,235	\$0	\$1,612,235
	Conifers and Cycads	Cypress, Santa Cruz	Cupressus abramsiana	Wherever found	\$7,200	\$0	\$7,200
		Conifers and Cycads subtotal			\$7,200	\$0	\$7,200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Ferns and Allies	Fern, American hart's-tongue	Asplenium scolopendrium var. americanum	Wherever found	\$1,279	\$0	\$1,279
		Quillwort, black spored	Isoetes melanospora	Wherever found	\$2,000	\$0	\$2,000
		Quillwort, Louisiana	Isoetes louisianensis	Wherever found	\$19,084	\$0	\$19,084
	Ferns and Allies subtotal				\$22,363	\$0	\$22,363
	Lichens	Cladonia, Florida perforate	Cladonia perforata	Wherever found	\$2,000	\$0	\$2,000
	Lichens subtotal				\$2,000	\$0	\$2,000
Domestic subtotal					\$181,136,616	\$556,043	\$181,692,659
Foreign	Mammals	Bear, brown	Ursus arctos pruinosus	Wherever found	\$4,800	\$0	\$4,800
	Mammals subtotal				\$4,800	\$0	\$4,800
	Reptiles	Sea turtle, green	Chelonia mydas	East Indian- West Pacific DPS	\$521	\$0	\$521
		Sea turtle, green	Chelonia mydas	North Indian DPS	\$521	\$0	\$521
		Sea turtle, green	Chelonia mydas	Southwest Indian DPS	\$521	\$0	\$521
		Sea turtle, green	Chelonia mydas	Southwest Pacific DPS	\$12,111	\$0	\$12,111
		Sea turtle, loggerhead	Caretta caretta	Northeast Atlantic Ocean DPS	\$177,166	\$0	\$177,166
		Sea turtle, loggerhead	Caretta caretta	South Atlantic Ocean DPS	\$348,275	\$0	\$348,275
	Reptiles subtotal				\$539,115	\$0	\$539,115

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Foreign subtotal					\$543,915	\$0	\$543,915
Domestic	Multi-Species				\$1,297,157	\$0	\$1,297,157
	Multi-Species subtotal				\$1,297,157	\$0	\$1,297,157
Domestic subtotal					\$1,297,157	\$0	\$1,297,157
Total					\$182,977,688	\$556,043	\$183,533,731

DOD Army - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$115,000	\$0	\$115,000
		Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$237,634	\$0	\$237,634
		Bat, Indiana	Myotis sodalis	Wherever found	\$708,649	\$0	\$708,649
		Bat, lesser long-nosed	Leptonycteris curasoae yerbabuena	Wherever found	\$48,000	\$0	\$48,000
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$1,222,830	\$0	\$1,222,830
		Fox, San Joaquin kit	Vulpes macrotis mutica	wherever found	\$96,223	\$0	\$96,223
		Jaguar	Panthera onca	Wherever found	\$7,000	\$0	\$7,000
		Ocelot	Leopardus (=Felis) pardalis	wherever found	\$7,000	\$0	\$7,000
		Pocket gopher, Roy Prairie	Thomomys mazama glacialis	Wherever found	\$266,666	\$0	\$266,666
		Pronghorn, Sonoran	Antilocapra americana sonoriensis	Wherever found, except where listed as an experimental population	\$49,200	\$0	\$49,200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$36,190	\$0	\$36,190
	Mammals subtotal				\$2,794,392	\$0	\$2,794,392
	Birds	Blackbird, yellow-shouldered	Agelaius xanthomus	Wherever found	\$25,000	\$0	\$25,000
		Condor, California	Gymnogyps californianus	U.S.A. (specific portions of Arizona, Nevada, and Utah)	\$12,000	\$0	\$12,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Crane, whooping	<i>Grus americana</i>	U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	\$120,822	\$0	\$120,822
		elepaio, Oahu	<i>Chasiempis ibidis</i>	Wherever found	\$100,000	\$0	\$100,000
		Falcon, Northern Aplomado	<i>Falco femoralis septentrionalis</i>	Wherever found, except where listed as an experimental population	\$175,259	\$0	\$175,259
		Goose, Hawaiian	<i>Branta (=Nesochen) sandvicensis</i>	Wherever found	\$160,000	\$0	\$160,000
		Horned lark, streaked	<i>Eremophila alpestris strigata</i>	Wherever found	\$466,666	\$0	\$466,666
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$151,277	\$0	\$151,277
		Palila (honeycreeper)	<i>Loxioides bailleui</i>	Wherever found	\$10,000	\$0	\$10,000
		petrel, Hawaiian	<i>Pterodroma sandwichensis</i>	Wherever found	\$47,050	\$0	\$47,050

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$2,000	\$0	\$2,000
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$25,000	\$0	\$25,000
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$300	\$0	\$300
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$51,812	\$0	\$51,812
		Storm-petrel, band-rumped	Oceanodroma castro	USA (HI)	\$47,050	\$0	\$47,050
		Tern, least	Sterna antillarum	interior pop.	\$1,000	\$0	\$1,000
		Vireo, Black-capped	Vireo atricapilla	Wherever found	\$255,600	\$0	\$255,600
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$4,671	\$0	\$4,671
		Warbler (=wood), golden-cheeked	Dendroica chrysoparia	Wherever found	\$318,400	\$0	\$318,400
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$7,041,042	\$0	\$7,041,042
	Birds subtotal				\$9,014,949	\$0	\$9,014,949

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Reptiles	Alligator, American	Alligator mississippiensis	Wherever found	\$3,000	\$0	\$3,000
		Boa, Puerto Rican	Epicrates inornatus	Wherever found	\$25,000	\$0	\$25,000
		gartersnake, northern Mexican	Thamnophis eques megalops	Wherever found	\$700	\$0	\$700
		Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$87,935	\$0	\$87,935
		Snake, black pine	Pituophis melanoleucus lodingi	Wherever found	\$48,947	\$0	\$48,947
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$183,092	\$0	\$183,092
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$355,853	\$0	\$355,853
		Tortoise, gopher	Gopherus polyphemus	West of Mobile and Tombigbee Rivers	\$108,584	\$0	\$108,584
	Reptiles subtotal				\$813,111	\$0	\$813,111
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$49,329	\$0	\$49,329
		Frog, Chiricahua leopard	Rana chiricahuensis	Wherever found	\$2,000	\$0	\$2,000
		Salamander, California tiger	Ambystoma californiense	U.S.A. (Central CA DPS)	\$21,456	\$0	\$21,456
		Salamander, frosted flatwoods	Ambystoma cingulatum	Wherever found	\$90,621	\$0	\$90,621

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salamander, Sonora tiger	Ambystoma tigrinum stebbinsi	Wherever found	\$1,500	\$0	\$1,500
		Toad, Houston	Bufo houstonensis	Wherever found	\$5,000	\$0	\$5,000
	Amphibians subtotal				\$169,906	\$0	\$169,906
	Fishes	Logperch, Roanoke	Percina rex	Wherever found	\$45,000	\$0	\$45,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$16,488	\$0	\$16,488
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$250,000	\$0	\$250,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$16,448	\$0	\$16,448
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$71,983	\$0	\$71,983
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$16,448	\$0	\$16,448
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Chesapeake Bay DPS	\$3,273	\$0	\$3,273
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	South Atlantic DPS	\$906	\$0	\$906
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$9,156	\$0	\$9,156
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$16,448	\$0	\$16,448
	Fishes subtotal				\$446,150	\$0	\$446,150

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Clams	Pocketbook, shinyrayed	Lampsilis subangulata	Wherever found	\$2,000	\$0	\$2,000
	Clams subtotal				\$2,000	\$0	\$2,000
	Snails	Snails, Oahu tree	Achatinella spp.	Wherever found	\$300,000	\$0	\$300,000
	Snails subtotal				\$300,000	\$0	\$300,000
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$191,360	\$0	\$191,360
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$17,178	\$0	\$17,178
		Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$82,000	\$0	\$82,000
		Butterfly, Oregon silverspot	Speyeria zerene hippolyta	Wherever found	\$10,700	\$0	\$10,700
		Butterfly, Saint Francis' satyr	Neonympha mitchellii francisci	Wherever found	\$244,481	\$0	\$244,481
		Checkerspot, Taylor's (=whulge)	Euphydryas editha taylori	Wherever found	\$541,666	\$0	\$541,666
		fly, Hawaiian picture-wing	Drosophila substenoptera	Wherever found	\$128,415	\$0	\$128,415
		picture-wing fly, Hawaiian	Drosophila montgomeryi	Wherever found	\$128,415	\$0	\$128,415
	Insects subtotal				\$1,344,215	\$0	\$1,344,215
	Crustaceans	Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$61,136	\$0	\$61,136
	Crustaceans subtotal				\$61,136	\$0	\$61,136
	Flowering Plants	Amaranth, seabeach	Amaranthus pumilus	Wherever found	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		amole, purple	<i>Chlorogalum purpureum</i>	Wherever found	\$87,693	\$0	\$87,693
		Aupaka	<i>Isodendrion hosakae</i>	Wherever found	\$32,000	\$0	\$32,000
		A`e	<i>Zanthoxylum hawaiiense</i>	Wherever found	\$32,000	\$0	\$32,000
		Barbara's Buttons, Mohr's	<i>Marshallia mohrii</i>	Wherever found	\$822	\$0	\$822
		Bear-poppy, Dwarf	<i>Arctomecon humilis</i>	Wherever found	\$266	\$0	\$266
		Bird's-beak, soft	<i>Cordylanthus mollis</i> ssp. <i>mollis</i>	Wherever found	\$90,000	\$0	\$90,000
		Chaffseed, American	<i>Schwalbea americana</i>	Wherever found	\$105,574	\$0	\$105,574
		Clover, running buffalo	<i>Trifolium stoloniferum</i>	Wherever found	\$19,000	\$0	\$19,000
		Coneflower, smooth	<i>Echinacea laevigata</i>	Wherever found	\$20,124	\$0	\$20,124
		Grass, Tennessee yellow-eyed	<i>Xyris tennesseensis</i>	Wherever found	\$4,067	\$0	\$4,067
		Haha	<i>Cyanea grimesiana</i> ssp. <i>obatae</i>	Wherever found	\$128,415	\$0	\$128,415
		Haha	<i>Cyanea koolauensis</i>	Wherever found	\$128,415	\$0	\$128,415
		Haha	<i>Cyanea longiflora</i>	Wherever found	\$128,415	\$0	\$128,415
		Haha	<i>Cyanea superba</i>	Wherever found	\$128,415	\$0	\$128,415
		Ha`iwale	<i>Cyrtandra dentata</i>	Wherever found	\$128,415	\$0	\$128,415
		Honohono	<i>Haplostachys</i> <i>haplostachya</i>	Wherever found	\$32,000	\$0	\$32,000
		Kamakahala	<i>Labordia cyrtandrae</i>	Wherever found	\$128,415	\$0	\$128,415
		Kamanomano	<i>Cenchrus agrimonioides</i>	Wherever found	\$128,415	\$0	\$128,415
		Kio`ele	<i>Kadua coriacea</i>	Wherever found	\$32,000	\$0	\$32,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Ko`oloa`ula	<i>Abutilon menziesii</i>	Wherever found	\$32,000	\$0	\$32,000
		Kuahiwi laukahi	<i>Plantago princeps</i>	Wherever found	\$128,415	\$0	\$128,415
		Kulu`i	<i>Nototrichium humile</i>	Wherever found	\$128,415	\$0	\$128,415
		Ladies'-tresses, Navasota	<i>Spiranthes parksii</i>	Wherever found	\$18,000	\$0	\$18,000
		Lau `ehu	<i>Panicum niuhauense</i>	Wherever found	\$47,865	\$0	\$47,865
		Loosestrife, rough- leaved	<i>Lysimachia asperulaefolia</i>	Wherever found	\$132,246	\$0	\$132,246
		Lo`ulu	<i>Pritchardia kaalae</i>	Wherever found	\$128,415	\$0	\$128,415
		Mahoe	<i>Alectryon macrococcus</i>	Wherever found	\$128,415	\$0	\$128,415
		Ma`o hau hele, (=native yellow hibiscus)	<i>Hibiscus brackenridgei</i>	Wherever found	\$128,415	\$0	\$128,415
		Ma`oli`oli	<i>Schiedea hawaiiensis</i>	Wherever found	\$32,000	\$0	\$32,000
		Mehamehame	<i>Flueggea neowawraea</i>	Wherever found	\$128,415	\$0	\$128,415
		Milk-vetch, Lane Mountain	<i>Astragalus jaegerianus</i>	Wherever found	\$105,853	\$0	\$105,853
		Nanu	<i>Gardenia mannii</i>	Wherever found	\$128,415	\$0	\$128,415
		Na`ena`e	<i>Dubautia herbstobatae</i>	Wherever found	\$128,415	\$0	\$128,415
		Nehe	<i>Melanthera tenuifolia</i>	Wherever found	\$128,415	\$0	\$128,415
		Nioi	<i>Eugenia koolauensis</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Abutilon sandwicense</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Cyrtandra nanawaleensis</i>	Wherever found	\$17,766	\$0	\$17,766
		No common name	<i>Festuca hawaiiensis</i>	Wherever found	\$32,000	\$0	\$32,000
		No common name	<i>Geocarpon minimum</i>	Wherever found	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		No common name	<i>Gouania vitifolia</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Hesperomannia arborescens</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Hesperomannia arbuscula</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Kadua degeneri</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Kadua parvula</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Lipochaeta venosa</i>	Wherever found	\$32,000	\$0	\$32,000
		No common name	<i>Neraudia angulata</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Neraudia ovata</i>	Wherever found	\$32,000	\$0	\$32,000
		No common name	<i>Phyllostegia hirsuta</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Phyllostegia kaalaensis</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Phyllostegia mollis</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Sanicula mariversa</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Schiedea kaalae</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Schiedea nuttallii</i>	Wherever found	\$128,415	\$0	\$128,415
		No common name	<i>Schiedea obovata</i>	Wherever found	\$128,415	\$0	\$128,415

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		No common name	Schiedea trinervis	Wherever found	\$128,415	\$0	\$128,415
		No common name	Silene hawaiiensis	Wherever found	\$32,000	\$0	\$32,000
		No common name	Silene lanceolata	Wherever found	\$32,000	\$0	\$32,000
		No common name	Spermolepis hawaiiensis	Wherever found	\$32,000	\$0	\$32,000
		No common name	Stenogyne angustifolia angustifolia	Wherever found	\$32,000	\$0	\$32,000
		No common name	Stenogyne kanehoana	Wherever found	\$128,415	\$0	\$128,415
		No common name	Tetramolopium arenarium	Wherever found	\$32,000	\$0	\$32,000
		No common name	Tetramolopium filiforme	Wherever found	\$128,415	\$0	\$128,415
		Oha	Delissea subcordata	Wherever found	\$128,415	\$0	\$128,415
		Pamakani	Viola chamissoniana ssp. chamissoniana	Wherever found	\$128,415	\$0	\$128,415
		Pennyroyal, Todsens's	Hedeoma todsenii	Wherever found	\$47,000	\$0	\$47,000
		Peppergrass, Slickspot	Lepidium papilliferum		\$80,000	\$0	\$80,000
		Piperia, Yadon's	Piperia yadonii	Wherever found	\$79,700	\$0	\$79,700
		Pogonia, small whorled	Isotria medeoloides	Wherever found	\$2,348	\$0	\$2,348
		Popolo ku mai	Solanum incompletum	Wherever found	\$32,000	\$0	\$32,000
		Potato-bean, Price's	Apios priceana		\$40,000	\$0	\$40,000
		Po`e	Portulaca sclerocarpa	Wherever found	\$32,000	\$0	\$32,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Rhododendron, Chapman	Rhododendron chapmanii	Wherever found	\$500	\$0	\$500
		Rockcress, Georgia	Arabis georgiana	Wherever found	\$5,000	\$0	\$5,000
		Skullcap, large-flowered	Scutellaria montana	Wherever found	\$3,839	\$0	\$3,839
		Spineflower, Monterey	Chorizanthe pungens var. pungens	Wherever found	\$68,798	\$0	\$68,798
		Sumac, Michaux's	Rhus michauxii	Wherever found	\$164,574	\$0	\$164,574
		Trillium, relict	Trillium reliquum	Wherever found	\$20,000	\$0	\$20,000
		Water-umbel, Huachuca	Lilaeopsis schaffneriana var. recurva	Wherever found	\$43,000	\$0	\$43,000
		`Akoko	Euphorbia celastroides var. kaenana	Wherever found	\$128,415	\$0	\$128,415
		`Akoko	Euphorbia herbstii	Wherever found	\$128,415	\$0	\$128,415
		`Anunu	Sicyos macrophyllus	Wherever found	\$32,000	\$0	\$32,000
		Flowering Plants subtotal			\$6,629,805	\$0	\$6,629,805
	Ferns and Allies	Ihi`ihi	Marsilea villosa	Wherever found	\$32,000	\$0	\$32,000
		No common name	Asplenium peruvianum var. insulare	Wherever found	\$32,000	\$0	\$32,000
		Quillwort, Louisiana	Isoetes louisianensis	Wherever found	\$107,998	\$0	\$107,998
		Ferns and Allies subtotal			\$171,998	\$0	\$171,998
		Multi-Species			\$25,602,301	\$0	\$25,602,301
		Multi-Species subtotal			\$25,602,301	\$0	\$25,602,301
	Domestic subtotal				\$47,349,963	\$0	\$47,349,963
	Total				\$47,349,963	\$0	\$47,349,963

DOD Marine Corps - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Indiana	Myotis sodalis	Wherever found	\$32,500	\$0	\$32,500
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$91,980	\$0	\$91,980
		Kangaroo rat, giant	Dipodomys ingens	Wherever found	\$322,000	\$0	\$322,000
		Manatee, West Indian	Trichechus manatus	Wherever found	\$5,000	\$0	\$5,000
		Mouse, Pacific pocket	Perognathus longimembris pacificus	Wherever found	\$346,000	\$0	\$346,000
		Pronghorn, Sonoran	Antilocapra americana sonoriensis	Wherever found, except where listed as an experimental population	\$90,250	\$0	\$90,250
		Seal, Hawaiian monk	Monachus schauinslandi	Wherever found	\$1,000	\$0	\$1,000
	Mammals subtotal				\$888,730	\$0	\$888,730
	Birds	Coot, Hawaiian	Fulica americana alai	Wherever found	\$60,000	\$0	\$60,000
		Duck, Hawaiian (=koloa)	Anas wyvilliana	Wherever found	\$60,000	\$0	\$60,000
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$235,810	\$0	\$235,810
		gallinule, Hawaiian common	Gallinula galeata sandvicensis	Wherever found	\$60,000	\$0	\$60,000
		Gnatcatcher, coastal California	Polioptila californica californica	Wherever found	\$409,000	\$0	\$409,000

		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$255,000	\$0	\$255,000
		Stilt, Hawaiian	Himantopus mexicanus knudseni	Wherever found	\$150,000	\$0	\$150,000
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$25,000	\$0	\$25,000
		Tern, California least	Sterna antillarum browni	Wherever found	\$267,000	\$0	\$267,000
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$258,810	\$0	\$258,810
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$441,530	\$0	\$441,530
	Birds subtotal				\$2,222,150	\$0	\$2,222,150
	Reptiles	Alligator, American	Alligator mississippiensis	Wherever found	\$30,000	\$0	\$30,000
		Sea turtle, green	Chelonia mydas	Central North Pacific DPS	\$1,000	\$0	\$1,000
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$164,990	\$0	\$164,990
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$10,295,330	\$0	\$10,295,330
	Reptiles subtotal				\$10,491,320	\$0	\$10,491,320
	Amphibians	salamander, Reticulated flatwoods	Ambystoma bishopi	Wherever found	\$11,240	\$0	\$11,240

		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$178,000	\$0	\$178,000
		toad, Yosemite	Anaxyrus canorus	Wherever found	\$150,000	\$0	\$150,000
	Amphibians subtotal				\$339,240	\$0	\$339,240
	Fishes	Goby, tidewater	Eucyclogobius newberryi	Wherever found	\$244,000	\$0	\$244,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$48,000	\$0	\$48,000
	Fishes subtotal				\$292,000	\$0	\$292,000
	Insects	Butterfly, Quino checkerspot	Euphydryas editha quino (=E. e. wrighti)	Wherever found	\$338,850	\$0	\$338,850
		Yellow-faced bee, anthricinan	Hylaeus anthracinus	Wherever found	\$200	\$0	\$200
	Insects subtotal				\$339,050	\$0	\$339,050
	Crustaceans	Fairy shrimp, San Diego	Branchinecta sandiegonensis	Wherever found	\$39,080	\$0	\$39,080
	Crustaceans subtotal				\$39,080	\$0	\$39,080
	Flowering Plants	Button-celery, San Diego	Eryngium aristulatum var. parishii	Wherever found	\$34,770	\$0	\$34,770
		Mesa-mint, San Diego	Pogogyne abramsii	Wherever found	\$34,770	\$0	\$34,770
		Monardella, willowy	Monardella viminea	Wherever found	\$172,780	\$0	\$172,780
		Navarretia, spreading	Navarretia fossalis	Wherever found	\$34,770	\$0	\$34,770
		Orcutt grass, California	Orcuttia californica	Wherever found	\$34,770	\$0	\$34,770
		Pondberry	Lindera melissifolia	Wherever found	\$14,000	\$0	\$14,000
	Flowering Plants subtotal				\$325,860	\$0	\$325,860
Domestic subtotal					\$14,937,430	\$0	\$14,937,430
Total					\$14,937,430	\$0	\$14,937,430

DOD Navy - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$21,362	\$0	\$21,362
		Bat, Indiana	Myotis sodalis	Wherever found	\$28,502	\$0	\$28,502
		Bat, Mariana fruit (=Mariana flying fox)	Pteropus mariannus mariannus	Wherever found	\$143,282	\$0	\$143,282
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$175,913	\$0	\$175,913
		Bear, polar	Ursus maritimus	Wherever found	\$35,966	\$0	\$35,966
		Deer, key	Odocoileus virginianus clavium	Wherever found	\$3,390	\$0	\$3,390
		Kangaroo rat, Fresno	Dipodomys nitratoides exilis	Wherever found	\$7,500	\$0	\$7,500
		Kangaroo rat, Stephens'	Dipodomys stephensi (incl. D. cascus)	Wherever found	\$318,357	\$0	\$318,357
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$2,477	\$0	\$2,477
		Manatee, West Indian	Trichechus manatus	Wherever found	\$104,757	\$0	\$104,757
		Otter, Northern Sea	Enhydra lutris kenyoni	Southwest Alaska DPS	\$12,186	\$0	\$12,186
		Otter, southern sea	Enhydra lutris nereis	Wherever found	\$188,417	\$0	\$188,417
		Rabbit, Lower Keys marsh	Sylvilagus palustris hefneri	Wherever found	\$112,190	\$0	\$112,190
		Rice rat	Oryzomys palustris natator	lower FL Keys	\$3,390	\$0	\$3,390
		Seal, bearded	Erignathus barbatus nauticus	Okhotsk DPS	\$22,186	\$0	\$22,186

		Seal, Guadalupe fur	<i>Arctocephalus townsendi</i>	Wherever found	\$1,406	\$0	\$1,406
		Seal, Hawaiian monk	<i>Monachus schauinslandi</i>	Wherever found	\$175,139	\$0	\$175,139
		Shrew, Buena Vista Lake ornate	<i>Sorex ornatus relictus</i>	Wherever found	\$12,469	\$0	\$12,469
		whale, beluga	<i>Delphinapterus leucas</i>	Cook Inlet DPS	\$288,000	\$0	\$288,000
		Whale, blue	<i>Balaenoptera musculus</i>	Wherever found	\$1,683,529	\$0	\$1,683,529
		Whale, bowhead	<i>Balaena mysticetus</i>	Wherever found	\$395,111	\$0	\$395,111
		whale, false killer	<i>Pseudorca crassidens</i>	Main Hawaiian Islands Insular DPS	\$423,080	\$0	\$423,080
		Whale, finback	<i>Balaenoptera physalus</i>	Wherever found	\$1,335,618	\$0	\$1,335,618
		Whale, gray	<i>Eschrichtius robustus</i>	Western North Pacific DPS	\$288,446	\$0	\$288,446
		Whale, humpback	<i>Megaptera novaeangliae</i>	Central America DPS	\$563,657	\$0	\$563,657
		Whale, humpback	<i>Megaptera novaeangliae</i>	Mexico DPS	\$611,066	\$0	\$611,066
		Whale, humpback	<i>Megaptera novaeangliae</i>	Western North Pacific	\$447,651	\$0	\$447,651
		Whale, killer	<i>Orcinus orca</i>	Southern Resident DPS	\$682,528	\$0	\$682,528
		Whale, North Atlantic Right	<i>Eubalaena glacialis</i>	Wherever found	\$1,165,732	\$0	\$1,165,732
		Whale, North Pacific Right	<i>Eubalaena japonica</i>	Wherever found	\$320,656	\$0	\$320,656
		Whale, Sei	<i>Balaenoptera borealis</i>	Wherever found	\$619,382	\$0	\$619,382
		Whale, sperm	<i>Physeter catodon</i> (= <i>macrocephalus</i>)	Wherever found	\$660,958	\$0	\$660,958
		Mammals subtotal			\$10,854,303	\$0	\$10,854,303

	Birds	Albatross, short-tailed	Phoebastria (=Diomedea) albatrus	Wherever found	\$40,373	\$0	\$40,373
		Coot, Hawaiian	Fulica americana alai	Wherever found	\$119,161	\$0	\$119,161
		Crow, Mariana (=aga)	Corvus kubaryi	Wherever found	\$170,825	\$0	\$170,825
		Duck, Hawaiian (=koloa)	Anas wyvilliana	Wherever found	\$107,615	\$0	\$107,615
		Eider, spectacled	Somateria fischeri	Wherever found	\$125	\$0	\$125
		Eider, Steller's	Polysticta stelleri	AK breeding pop.	\$125	\$0	\$125
		elepaio, Oahu	Chasiempis ibidis	Wherever found	\$40,201	\$0	\$40,201
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$33,548	\$0	\$33,548
		gallinule, Hawaiian common	Gallinula galeata sandvicensis	Wherever found	\$119,903	\$0	\$119,903
		Gnatcatcher, coastal California	Polioptila californica californica	Wherever found	\$374,213	\$0	\$374,213
		Goose, Hawaiian	Branta (=Nesochen) sandvicensis	Wherever found	\$27,892	\$0	\$27,892
		Kingfisher, Guam Micronesian	Halcyon cinnamomina cinnamomina	Wherever found	\$170,825	\$0	\$170,825
		Knot, red	Calidris canutus rufa	Wherever found	\$72,419	\$0	\$72,419
		Megapode, Micronesian	Megapodius laperouse	Wherever found	\$15,179	\$0	\$15,179
		Moorhen, Mariana common	Gallinula chloropus guami	Wherever found	\$104,242	\$0	\$104,242
		Murrelet, marbled	Brachyramphus marmoratus	U.S.A. (CA, OR, WA)	\$726,505	\$0	\$726,505
		Owl, Mexican spotted	Strix occidentalis lucida	Wherever found	\$10,000	\$0	\$10,000

		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$2,216	\$0	\$2,216
		petrel, Bermuda	<i>Pterodroma cahow</i>	Wherever found	\$10,118	\$0	\$10,118
		petrel, Hawaiian	<i>Pterodroma sandwichensis</i>	Wherever found	\$35,681	\$0	\$35,681
		Plover, piping	<i>Charadrius melodus</i>	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$52,167	\$0	\$52,167
		Plover, western snowy	<i>Charadrius nivosus nivosus</i>	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$812,150	\$0	\$812,150
		Rail, Guam	<i>Rallus owstoni</i>	Wherever found, except where listed as an experimental population	\$59,026	\$0	\$59,026
		Rail, light-footed clapper	<i>Rallus longirostris levipes</i>	Wherever found	\$162,765	\$0	\$162,765
		Shearwater, Newell's Townsend's	<i>Puffinus auricularis newelli</i>	Wherever found	\$41,652	\$0	\$41,652

		Shrike, San Clemente loggerhead	<i>Lanius ludovicianus mearnsi</i>	Wherever found	\$2,383,719	\$0	\$2,383,719
		Sparrow, San Clemente sage	<i>Amphispiza belli clementeae</i>	Wherever found	\$804,173	\$0	\$804,173
		Stilt, Hawaiian	<i>Himantopus mexicanus knudseni</i>	Wherever found	\$124,897	\$0	\$124,897
		Stork, wood	<i>Mycteria americana</i>	AL, FL, GA, MS, NC, SC	\$2,980	\$0	\$2,980
		Storm-petrel, band-rumped	<i>Oceanodroma castro</i>	USA (HI)	\$35,681	\$0	\$35,681
		Swiftlet, Mariana gray	<i>Aerodramus vanikorensis bartschi</i>	Wherever found	\$445,976	\$0	\$445,976
		Tern, California least	<i>Sterna antillarum browni</i>	Wherever found	\$873,297	\$0	\$873,297
		Tern, least	<i>Sterna antillarum</i>	interior pop.	\$1,420	\$0	\$1,420
		Tern, roseate	<i>Sterna dougallii dougallii</i>	Western Hemisphere except NE U.S.	\$17,843	\$0	\$17,843
		Towhee, Inyo California	<i>Pipilo crissalis eremophilus</i>	Wherever found	\$94,515	\$0	\$94,515
		Vireo, least Bell's	<i>Vireo bellii pusillus</i>	Wherever found	\$361,983	\$0	\$361,983
	Birds subtotal				\$8,455,410	\$0	\$8,455,410
	Reptiles	Alligator, American	<i>Alligator mississippiensis</i>	Wherever found	\$1,092	\$0	\$1,092
		Crocodile, American	<i>Crocodylus acutus</i>	U.S.A. (FL)	\$8,605	\$0	\$8,605
		Sea turtle, green	<i>Chelonia mydas</i>	Central North Pacific DPS	\$85,778	\$0	\$85,778
		Sea turtle, green	<i>Chelonia mydas</i>	Central West Pacific DPS	\$90,000	\$0	\$90,000
		Sea turtle, green	<i>Chelonia mydas</i>	East Pacific DPS	\$156,207	\$0	\$156,207

		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$108,035	\$0	\$108,035
		Sea turtle, green	Chelonia mydas	South Atlantic DPS	\$5,243	\$0	\$5,243
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$278,654	\$0	\$278,654
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$77,551	\$0	\$77,551
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$57,555	\$0	\$57,555
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$74,893	\$0	\$74,893
		Sea turtle, olive ridley	Lepidochelys olivacea	Wherever found, except when listed as endangered under 50 CFR 224.101	\$24,962	\$0	\$24,962
		Skink, sand	Neoseps reynoldsi	Wherever found	\$1,380	\$0	\$1,380
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$9,083	\$0	\$9,083
		Snake, giant garter	Thamnophis gigas	Wherever found	\$13,546	\$0	\$13,546
		Tortoise, desert	Gopherus agassizii	AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	\$150,831	\$0	\$150,831
	Reptiles subtotal				\$1,143,415	\$0	\$1,143,415

	Amphibians	Frog, California red-legged	<i>Rana draytonii</i>	Wherever found	\$21,635	\$0	\$21,635
		Salamander, frosted flatwoods	<i>Ambystoma cingulatum</i>	Wherever found	\$750	\$0	\$750
		salamander, Reticulated flatwoods	<i>Ambystoma bishopi</i>	Wherever found	\$12,414	\$0	\$12,414
		Toad, arroyo (=arroyo southwestern)	<i>Anaxyrus californicus</i>	Wherever found	\$150,980	\$0	\$150,980
	Amphibians subtotal				\$185,779	\$0	\$185,779
	Fishes	Bocaccio	<i>Sebastes paucispinis</i>	Puget Sound - Georgia Basin DPS	\$20,801	\$0	\$20,801
		Chub, humpback	<i>Gila cypha</i>	Wherever found	\$960	\$0	\$960
		Chub, Hutton tui	<i>Gila bicolor</i> ssp.	Wherever found	\$68,709	\$0	\$68,709
		Eulachon	<i>Thaleichthys pacificus</i>	Southern DPS	\$1,406	\$0	\$1,406
		grouper, Nassau	<i>Epinephelus striatus</i>		\$11,078	\$0	\$11,078
		rockfish, canary	<i>Sebastes pinniger</i>	Puget Sound - Georgia Basin DPS	\$2,593	\$0	\$2,593
		rockfish, yelloweye	<i>Sebastes ruberrimus</i>	Puget Sound - Georgia Basin DPS	\$28,165	\$0	\$28,165
		Salmon, Atlantic	<i>Salmo salar</i>	Gulf of Maine DPS	\$12,590	\$0	\$12,590
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	California Coastal ESU	\$1,406	\$0	\$1,406
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Central Valley spring-run ESU	\$1,406	\$0	\$1,406
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Lower Columbia River ESU	\$1,406	\$0	\$1,406

		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$75,345	\$0	\$75,345
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter- run ESU	\$1,406	\$0	\$1,406
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$1,406	\$0	\$1,406
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer -run ESU	\$1,406	\$0	\$1,406
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$1,406	\$0	\$1,406
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$1,406	\$0	\$1,406
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$1,406	\$0	\$1,406
		Salmon, chum	Oncorhynchus keta	Hood Canal summer-run ESU	\$42,450	\$0	\$42,450
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$1,406	\$0	\$1,406
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$1,406	\$0	\$1,406
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$1,406	\$0	\$1,406
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$1,406	\$0	\$1,406

		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU	\$1,406	\$0	\$1,406
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$1,406	\$0	\$1,406
		Sawfish, largetooth	Pristis pristis	Wherever found	\$1,853	\$0	\$1,853
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$15,561	\$0	\$15,561
		Shark, Scalloped Hammerhead	Sphyrna lewini	Central and Southwest Atlantic DPS	\$11,078	\$0	\$11,078
		Shark, Scalloped Hammerhead	Sphyrna lewini	Eastern Pacific DPS	\$12,907	\$0	\$12,907
		Shark, Scalloped Hammerhead	Sphyrna lewini	Indo-West Pacific DPS	\$511	\$0	\$511
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$1,500	\$0	\$1,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$1,406	\$0	\$1,406
		Steelhead	Oncorhynchus (=Salmo) mykiss	Central California Coast DPS	\$1,406	\$0	\$1,406
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$1,406	\$0	\$1,406
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$1,406	\$0	\$1,406
		Steelhead	Oncorhynchus (=Salmo) mykiss	Northern California DPS	\$1,406	\$0	\$1,406

		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$75,168	\$0	\$75,168
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$1,406	\$0	\$1,406
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$1,406	\$0	\$1,406
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$15,166	\$0	\$15,166
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$1,406	\$0	\$1,406
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$1,406	\$0	\$1,406
		Sturgeon (Gulf subspecies), Atlantic	Acipenser oxyrinchus (=oxyrinchus) desotoi	Wherever found	\$2,651	\$0	\$2,651
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Carolina DPS	\$12,530	\$0	\$12,530
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Chesapeake Bay DPS	\$292,856	\$0	\$292,856
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Gulf of Maine DPS	\$13,149	\$0	\$13,149
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	New York Bight DPS	\$12,530	\$0	\$12,530
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	South Atlantic DPS	\$12,980	\$0	\$12,980
		sturgeon, green	Acipenser medirostris	Southern DPS	\$1,406	\$0	\$1,406
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$287,784	\$0	\$287,784
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$86,105	\$0	\$86,105

	Fishes subtotal				\$1,153,576	\$0	\$1,153,576
	Snails	Abalone, Black	<i>Haliotis cracherodii</i>	Wherever found	\$319,690	\$0	\$319,690
		Abalone, White	<i>Haliotis sorenseni</i>	North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	\$303,891	\$0	\$303,891
		Snail, Guam tree	<i>Partula radiolata</i>	Wherever found	\$355,193	\$0	\$355,193
		Snail, Humped tree	<i>Partula gibba</i>	Wherever found	\$53,215	\$0	\$53,215
		Snails, Oahu tree	<i>Achatinella</i> spp.	Wherever found	\$70,769	\$0	\$70,769
	Snails subtotal				\$1,102,758	\$0	\$1,102,758
	Insects	Bumble bee, Rusty patched	<i>Bombus affinis</i>	Wherever found	\$8,473	\$0	\$8,473
		Butterfly, Mariana eight- spot	<i>Hypolimnas octocula marianensis</i>	Wherever found	\$59,026	\$0	\$59,026
		Butterfly, Mariana wandering	<i>Vagrans egistina</i>	Wherever found	\$2,905	\$0	\$2,905
		Butterfly, Palos Verdes blue	<i>Glaucopsyche lygdamus palosverdesensis</i>	Wherever found	\$213,136	\$0	\$213,136
		Butterfly, Quino checkerspot	<i>Euphydryas editha quino</i> (= <i>E. e. wrighti</i>)	Wherever found	\$256,666	\$0	\$256,666
		Checkerspot, Taylor's (=whulge)	<i>Euphydryas editha taylori</i>	Wherever found	\$500	\$0	\$500

		Damselfly, orangeblack Hawaiian	Megalagrion xanthomelas	Wherever found	\$6,162	\$0	\$6,162
		yellow-faced bee, Hawaiian	Hylaeus kuakea	Wherever found	\$37,075	\$0	\$37,075
	Insects subtotal				\$583,943	\$0	\$583,943
	Crustaceans	Fairy shrimp, San Diego	Branchinecta sandiegonensis	Wherever found	\$59,079	\$0	\$59,079
		Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$17,804	\$0	\$17,804
		Tadpole shrimp, vernal pool	Lepidurus packardi	Wherever found	\$17,804	\$0	\$17,804
	Crustaceans subtotal				\$94,687	\$0	\$94,687
	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$14,468	\$0	\$14,468
		Coral, elkhorn	Acropora palmata	Wherever found	\$15,361	\$0	\$15,361
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$14,468	\$0	\$14,468
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$14,468	\$0	\$14,468
		Coral, Pillar	Dendrogyra cylindrus	Wherever found	\$14,468	\$0	\$14,468
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$14,468	\$0	\$14,468
		Coral, staghorn	Acropora cervicornis	Wherever found	\$15,361	\$0	\$15,361
		No common name	Acropora globiceps	Wherever found	\$351,009	\$0	\$351,009
		No common name	Acropora retusa	Wherever found	\$143,979	\$0	\$143,979
		No common name	Seriatopora aculeata	Wherever found	\$68,875	\$0	\$68,875
	Corals subtotal				\$666,925	\$0	\$666,925
	Flowering Plants	Alani	Melicope christophersenii	Wherever found	\$13,895	\$0	\$13,895
		Alani	Melicope saint-johnii	Wherever found	\$13,895	\$0	\$13,895

		Beaked-rush, Knieskern's	<i>Rhynchospora knieskernii</i>	Wherever found	\$60,560	\$0	\$60,560
		Bird's-beak, salt marsh	<i>Cordylanthus maritimus</i> ssp. <i>maritimus</i>	Wherever found	\$106,996	\$0	\$106,996
		Bush-mallow, San Clemente Island	<i>Malacothamnus</i> <i>clementinus</i>	Wherever found	\$236,089	\$0	\$236,089
		Button-celery, San Diego	<i>Eryngium aristulatum</i> var. <i>parishii</i>	Wherever found	\$27,910	\$0	\$27,910
		Gilia, Monterey	<i>Gilia tenuiflora</i> ssp. <i>arenaria</i>	Wherever found	\$21,635	\$0	\$21,635
		Haha	<i>Cyanea calycina</i>	Wherever found	\$13,895	\$0	\$13,895
		Hala pepe	<i>Pleomele forbesii</i>	Wherever found	\$13,895	\$0	\$13,895
		Iliau, dwarf	<i>Wilkesia hobdyi</i>	Wherever found	\$4,418	\$0	\$4,418
		Indian paintbrush, San Clemente Island	<i>Castilleja grisea</i>	Wherever found	\$163,511	\$0	\$163,511
		Ko`oloa`ula	<i>Abutilon menziesii</i>	Wherever found	\$102,808	\$0	\$102,808
		Kuahiwi laukahi	<i>Plantago princeps</i>	Wherever found	\$14,654	\$0	\$14,654
		Kulu`i	<i>Nototrichium humile</i>	Wherever found	\$14,654	\$0	\$14,654
		Larkspur, San Clemente Island	<i>Delphinium variegatum</i> ssp. <i>kinkiense</i>	Wherever found	\$174,927	\$0	\$174,927
		Lau `ehu	<i>Panicum niihauense</i>	Wherever found	\$1,227	\$0	\$1,227
		Lotus, San Clemente Island	<i>Acmispon dendroideus</i> var. <i>traskiae</i> (=Lotus d. ssp. <i>traskiae</i>)	Wherever found	\$171,430	\$0	\$171,430
		Mahoe	<i>Alectryon macrococcus</i>	Wherever found	\$14,654	\$0	\$14,654
		Mehamehame	<i>Flueggea neowawraea</i>	Wherever found	\$14,654	\$0	\$14,654
		Mesa-mint, San Diego	<i>Pogogyne abramsii</i>	Wherever found	\$31,810	\$0	\$31,810
		Nehe	<i>Lipochaeta lobata</i> var. <i>leptophylla</i>	Wherever found	\$14,604	\$0	\$14,604

		No common name	<i>Abutilon sandwicense</i>	Wherever found	\$101,107	\$0	\$101,107
		No common name	<i>Bonamia menziesii</i>	Wherever found	\$14,604	\$0	\$14,604
		No common name	<i>Dendrobium guamense</i>	Wherever found	\$58,005	\$0	\$58,005
		No common name	<i>Kadua parvula</i>	Wherever found	\$13,275	\$0	\$13,275
		No common name	<i>Lobelia niihauensis</i>	Wherever found	\$13,275	\$0	\$13,275
		No common name	<i>Neraudia angulata</i>	Wherever found	\$13,275	\$0	\$13,275
		No common name	<i>Platydesma cornuta decurrens</i>	Wherever found	\$12,875	\$0	\$12,875
		No common name	<i>Schiedea hookeri</i>	Wherever found	\$13,075	\$0	\$13,075
		No common name	<i>Silene perlmanii</i>	Wherever found	\$13,075	\$0	\$13,075
		No common name	<i>Spermolepis hawaiiensis</i>	Wherever found	\$16,530	\$0	\$16,530
		No common name	<i>Tabernaemontana rotensis</i>	Wherever found	\$53,215	\$0	\$53,215
		No common name	<i>Tetramolopium lepidotum</i> ssp. <i>lepidotum</i>	Wherever found	\$11,750	\$0	\$11,750
		No common name	<i>Tuberolabium guamense</i>	Wherever found	\$53,215	\$0	\$53,215
		paintbrush, golden	<i>Castilleja levisecta</i>	Wherever found	\$33,278	\$0	\$33,278
		Pamakani	<i>Viola chamissoniana</i> ssp. <i>chamissoniana</i>	Wherever found	\$11,750	\$0	\$11,750
		Pink, swamp	<i>Helonias bullata</i>	Wherever found	\$35,492	\$0	\$35,492
		Piperia, Yadon's	<i>Piperia yadonii</i>	Wherever found	\$21,635	\$0	\$21,635
		Pondberry	<i>Lindera melissifolia</i>	Wherever found	\$750	\$0	\$750

		Pu`uka`a	Cyperus trachysanthos	Wherever found	\$18,350	\$0	\$18,350	
		Rockcress, Santa Cruz Island	Sibara filifolia	Wherever found	\$163,114	\$0	\$163,114	
		silverbush, Blodgett's	Argythamnia blodgettii	Wherever found	\$8,605	\$0	\$8,605	
		Spineflower, Monterey	Chorizanthe pungens var. pungens	Wherever found	\$21,635	\$0	\$21,635	
		Spineflower, Orcutt's	Chorizanthe orcuttiana	Wherever found	\$107,398	\$0	\$107,398	
		Spurge, Garber's	Chamaesyce garberi	Wherever found	\$3,390	\$0	\$3,390	
		Woodland-star, San Clemente Island	Lithophragma maximum	Wherever found	\$163,511	\$0	\$163,511	
		`Akoko	Euphorbia kuwaleana	Wherever found	\$10,817	\$0	\$10,817	
		`Akoko, Ewa Plains	Euphorbia skottsbergii var. skottsbergii	Wherever found	\$4,434	\$0	\$4,434	
		`Anaunau	Lepidium arbuscula	Wherever found	\$13,895	\$0	\$13,895	
		Flowering Plants subtotal			\$2,237,451	\$0	\$2,237,451	
		Conifers and Cycads	Fadang	Cycas micronesica	Wherever found	\$697,550	\$0	\$697,550
		Conifers and Cycads subtotal			\$697,550	\$0	\$697,550	
		Ferns and Allies	Ihi`ihi	Marsilea villosa	Wherever found	\$21,821	\$0	\$21,821
		No common name	Asplenium unisorum	Wherever found	\$14,604	\$0	\$14,604	
		Ferns and Allies subtotal			\$36,425	\$0	\$36,425	
Domestic subtotal					\$27,212,222	\$0	\$27,212,222	
Foreign	Reptiles	Sea turtle, loggerhead	Caretta caretta	South Atlantic Ocean DPS	\$960	\$0	\$960	
		Sea turtle, loggerhead	Caretta caretta	South Pacific Ocean DPS	\$1,406	\$0	\$1,406	

		Sea turtle, olive ridley	Lepidochelys olivacea	Breeding colony populations on Pacific coast of Mexico	\$1,406	\$0	\$1,406
	Reptiles subtotal				\$3,772	\$0	\$3,772
Foreign subtotal					\$3,772	\$0	\$3,772
Domestic	Multi-Species				\$288,622	\$0	\$288,622
	Multi-Species subtotal				\$288,622	\$0	\$288,622
Domestic subtotal					\$288,622	\$0	\$288,622
Total					\$27,504,616	\$0	\$27,504,616

DOE Bonneville Power Administration - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Fishes	Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$9,069,893	\$466,462	\$9,536,355
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$19,933,663	\$0	\$19,933,663
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$101,785,289	\$0	\$101,785,289
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$19,706,573	\$0	\$19,706,573
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$6,711,196	\$0	\$6,711,196
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$2,857,453	\$0	\$2,857,453
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$3,452,242	\$0	\$3,452,242
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$213,216	\$0	\$213,216
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$7,855,782	\$0	\$7,855,782
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$5,851,112	\$26,763	\$5,877,875
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$42,000,260	\$0	\$42,000,260
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$34,456,984	\$0	\$34,456,984

		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$15,875,259	\$6,254	\$15,881,513
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$4,099,361	\$0	\$4,099,361
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$10,965,527	\$0	\$10,965,527
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$16,128,452	\$0	\$16,128,452
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$1,815,677	\$0	\$1,815,677
	Fishes subtotal				\$302,777,939	\$499,479	\$303,277,418
Domestic subtotal					\$302,777,939	\$499,479	\$303,277,418
Total					\$302,777,939	\$499,479	\$303,277,418

DOE Southwestern Power Administration - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Birds	Tern, least	Sterna antillarum	interior pop.	\$7,000	\$70,000	\$77,000
	Birds subtotal				\$7,000	\$70,000	\$77,000
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$12,200	\$0	\$12,200
	Insects subtotal				\$12,200	\$0	\$12,200
	Multi-Species				\$9,700	\$0	\$9,700
	Multi-Species subtotal				\$9,700	\$0	\$9,700
Domestic subtotal					\$28,900	\$70,000	\$98,900
Total					\$28,900	\$70,000	\$98,900

DOE Western Area Power Administration - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Indiana	Myotis sodalis	Wherever found	\$4,775	\$0	\$4,775
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$4,775	\$0	\$4,775
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population or delisted	\$4,775	\$0	\$4,775
		Ferret, black- footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$9,800	\$0	\$9,800
		Fox, San Joaquin kit	Vulpes macrotis mutica	wherever found	\$35,500	\$0	\$35,500
		Kangaroo rat, giant	Dipodomys ingens	Wherever found	\$23,500	\$0	\$23,500
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$15,182	\$0	\$15,182
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	wherever found	\$26,252	\$0	\$26,252

		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$8,437	\$0	\$8,437
	Mammals subtotal				\$132,996	\$0	\$132,996
	Birds	Condor, California	Gymnogyps californianus	U.S.A. only, except where listed as an experimental population	\$5,000	\$0	\$5,000
		Crane, whooping	Grus americana	Wherever found, except where listed as an experimental population	\$4,775	\$0	\$4,775
		Cuckoo, yellow-billed	Coccyzus americanus	Western U.S. DPS	\$53,368	\$0	\$53,368

		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$10,000	\$0	\$10,000
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$4,775	\$0	\$4,775
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$96,408	\$0	\$96,408
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$9,000	\$0	\$9,000
		Plover, piping	<i>Charadrius melodus</i>	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$4,775	\$0	\$4,775
		sage-grouse, Gunnison	<i>Centrocercus minimus</i>	Wherever found	\$67,671	\$0	\$67,671
		Tern, least	<i>Sterna antillarum</i>	interior pop.	\$4,775	\$0	\$4,775
		Vireo, least Bell's	<i>Vireo bellii pusillus</i>	Wherever found	\$8,000	\$0	\$8,000
	Birds subtotal				\$268,547	\$0	\$268,547
	Reptiles	gartersnake, narrow-headed	<i>Thamnophis rufipunctatus</i>	Wherever found	\$5,000	\$0	\$5,000
		gartersnake, northern Mexican	<i>Thamnophis eques megalops</i>	Wherever found	\$6,800	\$0	\$6,800
		Lizard, blunt-nosed leopard	<i>Gambelia silus</i>	Wherever found	\$36,000	\$0	\$36,000
		Massasauga (=rattlesnake), eastern	<i>Sistrurus catenatus</i>	Wherever found	\$4,775	\$0	\$4,775
		Snake, giant garter	<i>Thamnophis gigas</i>	Wherever found	\$23,500	\$0	\$23,500

		Tortoise, desert	<i>Gopherus agassizii</i>	Wherever found, except AZ south and east of Colorado R., and Mexico	\$153,700	\$0	\$153,700
		Whipsnake (=striped racer), Alameda	<i>Masticophis lateralis euryxanthus</i>	Wherever found	\$8,000	\$0	\$8,000
	Reptiles subtotal				\$237,775	\$0	\$237,775
	Amphibians	Frog, California red-legged	<i>Rana draytonii</i>	Wherever found	\$11,500	\$0	\$11,500
		Salamander, California tiger	<i>Ambystoma californiense</i>	U.S.A. (Central CA DPS)	\$10,500	\$0	\$10,500
	Amphibians subtotal				\$22,000	\$0	\$22,000
	Fishes	Chub, bonytail	<i>Gila elegans</i>	Wherever found	\$3,041,917	\$0	\$3,041,917
		Chub, humpback	<i>Gila cypha</i>	Wherever found	\$3,041,917	\$0	\$3,041,917
		Chub, Pahrnagat roundtail	<i>Gila robusta jordani</i>	Wherever found	\$3,462	\$0	\$3,462
		Pikeminnow (=squawfish), Colorado	<i>Ptychocheilus lucius</i>	Wherever found, except where listed as an experimental population	\$3,041,917	\$0	\$3,041,917
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	Central Valley spring-run ESU	\$22,000	\$0	\$22,000
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) <i>tshawytscha</i>	Sacramento River winter- run ESU	\$22,000	\$0	\$22,000

		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$4,775	\$0	\$4,775
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$8,000	\$0	\$8,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$22,000	\$0	\$22,000
		sturgeon, green	Acipenser medirostris	Southern DPS	\$22,000	\$0	\$22,000
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$300,000	\$0	\$300,000
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$3,046,918	\$0	\$3,046,918
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$4,775	\$0	\$4,775
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$4,915	\$0	\$4,915
		Fishes subtotal			\$12,586,596	\$0	\$12,586,596
	Clams	Higgins eye (pearlymussel)	Lampsilis higginsii	Wherever found	\$4,775	\$0	\$4,775
		Mussel, scaleshell	Leptodea leptodon	Wherever found	\$4,775	\$0	\$4,775
		Clams subtotal			\$9,550	\$0	\$9,550
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$4,775	\$0	\$4,775
		Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$48,500	\$0	\$48,500

		Butterfly, Uncompahgre fritillary	<i>Boloria acrocne</i>	Wherever found	\$10,279	\$0	\$10,279
		Skipper, Dakota	<i>Hesperia dacotae</i>	Wherever found	\$4,775	\$0	\$4,775
		skipperling, Poweshiek	<i>Oarisma poweshiek</i>	Wherever found	\$4,775	\$0	\$4,775
		Tiger beetle, Salt Creek	<i>Cicindela nevadica lincolniana</i>	Wherever found	\$4,775	\$0	\$4,775
	Insects subtotal				\$77,879	\$0	\$77,879
	Crustaceans	Fairy shrimp, Conservancy	<i>Branchinecta conservatio</i>	Wherever found	\$8,000	\$0	\$8,000
		Fairy shrimp, vernal pool	<i>Branchinecta lynchi</i>	Wherever found	\$22,000	\$0	\$22,000
		Tadpole shrimp, vernal pool	<i>Lepidurus packardii</i>	Wherever found	\$9,000	\$0	\$9,000
	Crustaceans subtotal				\$39,000	\$0	\$39,000
	Flowering Plants	Bird's beak, palmate- bracted	<i>Cordylanthus palmatus</i>	Wherever found	\$8,500	\$0	\$8,500
		Bush-clover, prairie	<i>Lespedeza leptostachya</i>	Wherever found	\$4,775	\$0	\$4,775
		Butterfly plant, Colorado	<i>Gaura neomexicana</i> var. <i>coloradensis</i>	Wherever found	\$13,090	\$0	\$13,090
		Cactus, Mesa Verde	<i>Sclerocactus mesae- verdae</i>	Wherever found	\$4,775	\$0	\$4,775
		Cactus, Pima pineapple	<i>Coryphantha scheeri</i> var. <i>robustispina</i>	Wherever found	\$39,100	\$0	\$39,100
		Checker- mallow, Keck's	<i>Sidalcea keckii</i>	Wherever found	\$500	\$0	\$500
		Fiddleneck, large- flowered	<i>Amsinckia grandiflora</i>	Wherever found	\$8,000	\$0	\$8,000
		Fritillary, Gentner's	<i>Fritillaria gentneri</i>	Wherever found	\$500	\$0	\$500

		Goldfields, Contra Costa	Lasthenia conjugens	Wherever found	\$8,000	\$0	\$8,000
		Jewelflower, California	Caulanthus californicus	Wherever found	\$24,000	\$0	\$24,000
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$44,124	\$0	\$44,124
		Milk-vetch, Mancos	Astragalus humillimus	Wherever found	\$6,000	\$0	\$6,000
		Milkweed, Mead's	Asclepias meadii	Wherever found	\$4,775	\$0	\$4,775
		Orchid, eastern prairie fringed	Platanthera leucophaea	Wherever found	\$4,775	\$0	\$4,775
		Orchid, western prairie fringed	Platanthera praeclara	Wherever found	\$7,000	\$0	\$7,000
		Phacelia, North Park	Phacelia formosula	Wherever found	\$5,450	\$0	\$5,450
		Wooly-threads, San Joaquin	Monolopia (=Lembertia) congdonii	Wherever found	\$23,500	\$0	\$23,500
		Flowering Plants subtotal			\$206,864	\$0	\$206,864
Domestic subtotal					\$13,581,207	\$0	\$13,581,207
Total					\$13,581,207	\$0	\$13,581,207

DOI Bureau of Indian Affairs - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$83,000	\$0	\$83,000
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population or delisted	\$40,235	\$0	\$40,235
		Ferret, black- footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$100,000	\$0	\$100,000
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$40,235	\$0	\$40,235
		Panther, Florida	Puma (=Felis) concolor coryi	Wherever found	\$88,000	\$0	\$88,000
		Wolf, gray	Canis lupus	MN	\$80,000	\$0	\$80,000

		Wolf, gray	<i>Canis lupus</i>	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$76,000	\$0	\$76,000
	Mammals subtotal				\$507,470	\$0	\$507,470
	Birds	Condor, California	<i>Gymnogyps californianus</i>	U.S.A. only, except where listed as an experimental population	\$80,000	\$0	\$80,000
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$44,771	\$0	\$44,771
		Horned lark, streaked	<i>Eremophila alpestris strigata</i>	Wherever found	\$30,000	\$0	\$30,000
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$40,336	\$0	\$40,336
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$482,864	\$0	\$482,864

		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$30,000	\$0	\$30,000
	Birds subtotal				\$707,971	\$0	\$707,971
	Fishes	Chub, Pahrnagat roundtail	Gila robusta jordani	Wherever found	\$83,000	\$0	\$83,000
		Minnow, Rio Grande Silvery	Hybognathus amarus	Wherever found, except where listed as an experimental population	\$44,771	\$0	\$44,771
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$659,617	\$0	\$659,617
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$35,000	\$0	\$35,000
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$11,409	\$0	\$11,409
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$37,825	\$0	\$37,825
	Fishes subtotal				\$871,622	\$0	\$871,622
	Clams	Mucket, Neosho	Lampsilis rafinesqueana	Wherever found	\$137,000	\$0	\$137,000
	Clams subtotal				\$137,000	\$0	\$137,000
	Insects	Skipper, Dakota	Hesperia dacotae	Wherever found	\$22,730	\$0	\$22,730
	Insects subtotal				\$22,730	\$0	\$22,730
	Flowering Plants	Lily, Western	Lilium occidentale	Wherever found	\$31,000	\$0	\$31,000

		Milkweed, Mead's	Asclepias meadii	Wherever found	\$150,000	\$0	\$150,000
	Flowering Plants subtotal				\$181,000	\$0	\$181,000
Domestic subtotal					\$2,427,793	\$0	\$2,427,793
Total					\$2,427,793	\$0	\$2,427,793

DOI Bureau of Land Management - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	bat, Florida bonneted	Eumops floridanus	Wherever found	\$100	\$0	\$100
		Bat, gray	Myotis grisescens	Wherever found	\$1,700	\$0	\$1,700
		Bat, Indiana	Myotis sodalis	Wherever found	\$1,700	\$0	\$1,700
		Bat, lesser long-nosed	Leptonycteris curasoae yerbabuenae	Wherever found	\$51,950	\$0	\$51,950
		Bat, Mexican long-nosed	Leptonycteris nivalis	Wherever found	\$25,000	\$0	\$25,000
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$134,380	\$0	\$134,380
		Bat, Ozark big-eared	Corynorhinus (=Plecotus) townsendii ingens	Wherever found	\$1,600	\$0	\$1,600
		Bat, Virginia big-eared	Corynorhinus (=Plecotus) townsendii virginianus	Wherever found	\$100	\$0	\$100
		Bear, grizzly	Ursus arctos horribilis	Bitterroot Ecosystem Recovery Zone Population	\$10,000	\$0	\$10,000
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population or delisted	\$170,090	\$0	\$170,090
		Bear, polar	Ursus maritimus	Wherever found	\$95,000	\$0	\$95,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Caribou, Woodland	Rangifer tarandus caribou	Selkirk Mountain population	\$1,000	\$0	\$1,000
		Ferret, black- footed	Mustela nigripes	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$156,178	\$0	\$156,178
		Ferret, black- footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$97,000	\$0	\$97,000
		Fox, San Joaquin kit	Vulpes macrotis mutica	wherever found	\$75,500	\$0	\$75,500
		Jaguar	Panthera onca	Wherever found	\$16,300	\$0	\$16,300
		Kangaroo rat, giant	Dipodomys ingens	Wherever found	\$42,500	\$0	\$42,500
		Kangaroo rat, San Bernardino Merriam's	Dipodomys merriami parvus	Wherever found	\$31,000	\$0	\$31,000
		Kangaroo rat, Stephens'	Dipodomys stephensi (incl. D. cascus)	Wherever found	\$36,000	\$0	\$36,000
		Kangaroo rat, Tipton	Dipodomys nitratoides nitratoides	Wherever found	\$11,500	\$0	\$11,500
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$252,164	\$0	\$252,164
		Manatee, West Indian	Trichechus manatus	Wherever found	\$2,100	\$0	\$2,100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Mountain beaver, Point Arena	<i>Aplodontia rufa nigra</i>	Wherever found	\$5,000	\$0	\$5,000
		Mouse, Alabama beach	<i>Peromyscus polionotus ammobates</i>	Wherever found	\$6,000	\$0	\$6,000
		Mouse, Key Largo cotton	<i>Peromyscus gossypinus allapaticola</i>	Wherever found	\$100	\$0	\$100
		Mouse, New Mexico meadow jumping	<i>Zapus hudsonius luteus</i>	Wherever found	\$95,800	\$0	\$95,800
		Mouse, Perdido Key beach	<i>Peromyscus polionotus trissyllepsis</i>	Wherever found	\$100	\$0	\$100
		Mouse, Preble's meadow jumping	<i>Zapus hudsonius preblei</i>	wherever found	\$13,878	\$0	\$13,878
		Mouse, southeastern beach	<i>Peromyscus polionotus niveiventris</i>	wherever found	\$100	\$0	\$100
		Ocelot	<i>Leopardus (=Felis) pardalis</i>	wherever found	\$14,300	\$0	\$14,300
		Panther, Florida	<i>Puma (=Felis) concolor coryi</i>	Wherever found	\$100	\$0	\$100
		Prairie dog, Utah	<i>Cynomys parvidens</i>	Wherever found	\$191,734	\$0	\$191,734
		Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	U.S.A. (AZ), Mexico	\$109,000	\$0	\$109,000
		Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	Wherever found, except where listed as an experimental population	\$4,250	\$48,970	\$53,220
		Rabbit, Columbia Basin Pygmy	<i>Brachylagus idahoensis</i>	Columbia Basin DPS	\$6,000	\$0	\$6,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Rabbit, Lower Keys marsh	<i>Sylvilagus palustris hefneri</i>	Wherever found	\$100	\$0	\$100
		Rice rat	<i>Oryzomys palustris natator</i>	lower FL Keys	\$100	\$0	\$100
		Sheep, Peninsular bighorn	<i>Ovis canadensis nelsoni</i>	Peninsular CA pop.	\$10,500	\$0	\$10,500
		Sheep, Sierra Nevada bighorn	<i>Ovis canadensis sierrae</i>	Sierra Nevada	\$12,000	\$0	\$12,000
		Shrew, Buena Vista Lake ornate	<i>Sorex ornatus relictus</i>	Wherever found	\$2,500	\$0	\$2,500
		Squirrel, Northern Idaho Ground	<i>Uroditellus brunneus</i>	Wherever found	\$2,000	\$0	\$2,000
		Vole, Amargosa	<i>Microtus californicus scirpensis</i>	Wherever found	\$180,000	\$0	\$180,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	<i>Canis lupus</i>	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$38,650	\$0	\$38,650
		Wolf, Mexican	<i>Canis lupus baileyi</i>	U.S.A. (portions of AZ and NM)see 17.84(k)	\$18,500	\$0	\$18,500
		Woodrat, Key Largo	<i>Neotoma floridana smalli</i>	Wherever found	\$100	\$0	\$100
	Mammals subtotal				\$1,923,674	\$48,970	\$1,972,644
	Birds	Bobwhite, masked (quail)	<i>Colinus virginianus ridgwayi</i>	Wherever found	\$4,500	\$0	\$4,500
		Caracara, Audubon's crested	<i>Polyborus plancus audubonii</i>	FL pop.	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Condor, California	<i>Gymnogyps californianus</i>	U.S.A. (specific portions of Arizona, Nevada, and Utah)	\$43,625	\$0	\$43,625
		Condor, California	<i>Gymnogyps californianus</i>	U.S.A. only, except where listed as an experimental population	\$57,525	\$0	\$57,525
		Crane, whooping	<i>Grus americana</i>	U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	\$4,600	\$0	\$4,600
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$1,000	\$0	\$1,000
		Cuckoo, yellow- billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$1,076,675	\$0	\$1,076,675
		Eider, spectacled	<i>Somateria fischeri</i>	Wherever found	\$33,000	\$0	\$33,000
		Eider, Steller's	<i>Polysticta stelleri</i>	AK breeding pop.	\$34,000	\$0	\$34,000
		Falcon, Northern Aplomado	<i>Falco femoralis</i> <i>septentrionalis</i>	U.S.A (AZ, NM)	\$37,400	\$0	\$37,400
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$900,116	\$0	\$900,116

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Gnatcatcher, coastal California	<i>Polioptila californica californica</i>	Wherever found	\$25,000	\$0	\$25,000
		Horned lark, streaked	<i>Eremophila alpestris strigata</i>	Wherever found	\$500	\$0	\$500
		Kite, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	Wherever found	\$100	\$0	\$100
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$4,950	\$0	\$4,950
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$1,534,100	\$0	\$1,534,100
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$210,164	\$0	\$210,164
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$4,078,100	\$0	\$4,078,100
		Plover, piping	<i>Charadrius melodus</i>	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$14,750	\$0	\$14,750
		Plover, western snowy	<i>Charadrius nivosus nivosus</i>	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$326,000	\$0	\$326,000
		Rail, Yuma clapper	<i>Rallus longirostris yumanensis</i>	Wherever found	\$398,350	\$0	\$398,350

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		sage-grouse, Gunnison	<i>Centrocercus minimus</i>	Wherever found	\$1,180,800	\$0	\$1,180,800
		scrub-jay, Florida	<i>Aphelocoma coerulescens</i>	Wherever found	\$1,100	\$0	\$1,100
		Stork, wood	<i>Mycteria americana</i>	AL, FL, GA, MS, NC, SC	\$100	\$0	\$100
		Tern, least	<i>Sterna antillarum</i>	interior pop.	\$14,950	\$0	\$14,950
		Towhee, Inyo California	<i>Pipilo crissalis eremophilus</i>	Wherever found	\$35,000	\$0	\$35,000
		Vireo, Black-capped	<i>Vireo atricapilla</i>	Wherever found	\$3,000	\$0	\$3,000
		Vireo, least Bell's	<i>Vireo bellii pusillus</i>	Wherever found	\$39,500	\$0	\$39,500
		Warbler, Kirtland's	<i>Setophaga kirtlandii</i> (= <i>Dendroica kirtlandii</i>)	Wherever found	\$100	\$0	\$100
		Woodpecker, ivory-billed	<i>Campephilus principalis</i>	Wherever found	\$100	\$0	\$100
		Woodpecker, red-cockaded	<i>Picoides borealis</i>	Wherever found	\$7,200	\$0	\$7,200
	Birds subtotal				\$10,066,405	\$0	\$10,066,405
	Reptiles	Alligator, American	<i>Alligator mississippiensis</i>	Wherever found	\$100	\$0	\$100
		gartersnake, narrow-headed	<i>Thamnophis rufipunctatus</i>	Wherever found	\$14,750	\$0	\$14,750
		gartersnake, northern Mexican	<i>Thamnophis eques megalops</i>	Wherever found	\$62,150	\$0	\$62,150
		Lizard, blunt-nosed leopard	<i>Gambelia silus</i>	Wherever found	\$34,500	\$0	\$34,500
		Lizard, Coachella Valley fringe-toed	<i>Uma inornata</i>	Wherever found	\$4,000	\$0	\$4,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Rattlesnake, New Mexican ridge-nosed	<i>Crotalus willardi obscurus</i>	Wherever found	\$8,000	\$0	\$8,000
		Sea turtle, green	<i>Chelonia mydas</i>	North Atlantic DPS	\$100	\$0	\$100
		Sea turtle, hawksbill	<i>Eretmochelys imbricata</i>	Wherever found	\$100	\$0	\$100
		Sea turtle, Kemp's ridley	<i>Lepidochelys kempii</i>	Wherever found	\$1,600	\$0	\$1,600
		Sea turtle, leatherback	<i>Dermochelys coriacea</i>	Wherever found	\$100	\$0	\$100
		Sea turtle, loggerhead	<i>Caretta caretta</i>	Northwest Atlantic Ocean DPS	\$1,600	\$0	\$1,600
		Snake, Atlantic salt marsh	<i>Nerodia clarkii taeniata</i>	Wherever found	\$100	\$0	\$100
		Snake, eastern indigo	<i>Drymarchon corais couperi</i>	Wherever found	\$100	\$0	\$100
		Tortoise, desert	<i>Gopherus agassizii</i>	Wherever found, except AZ south and east of Colorado R., and Mexico	\$1,711,647	\$357,600	\$2,069,247
		Tortoise, gopher	<i>Gopherus polyphemus</i>	West of Mobile and Tombigbee Rivers	\$100	\$0	\$100
	Reptiles subtotal				\$1,838,947	\$357,600	\$2,196,547
	Amphibians	Frog, California red-legged	<i>Rana draytonii</i>	Wherever found	\$72,500	\$0	\$72,500
		Frog, Chiricahua leopard	<i>Rana chiricahuensis</i>	Wherever found	\$68,500	\$0	\$68,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Frog, Oregon spotted	<i>Rana pretiosa</i>	Wherever found	\$101,300	\$0	\$101,300
		Hellbender, Ozark	<i>Cryptobranchus alleganiensis bishopi</i>	Wherever found	\$100	\$0	\$100
		Salamander, California tiger	<i>Ambystoma californiense</i>	U.S.A. (Central CA DPS)	\$10,000	\$0	\$10,000
		Salamander, desert slender	<i>Batrachoseps aridus</i>	Wherever found	\$10,000	\$0	\$10,000
		Toad, arroyo (=arroyo southwestern)	<i>Anaxyrus californicus</i>	Wherever found	\$500	\$0	\$500
		Toad, Wyoming	<i>Bufo hemiophrys baxteri</i>	Wherever found	\$29,000	\$0	\$29,000
	Amphibians subtotal				\$291,900	\$0	\$291,900
	Fishes	Chub, bonytail	<i>Gila elegans</i>	Wherever found	\$28,700	\$0	\$28,700
		Chub, Borax Lake	<i>Gila boraxobius</i>	Wherever found	\$2,500	\$0	\$2,500
		Chub, Chihuahua	<i>Gila nigrescens</i>	Wherever found	\$1,500	\$0	\$1,500
		Chub, Gila	<i>Gila intermedia</i>	Wherever found	\$30,300	\$0	\$30,300
		Chub, humpback	<i>Gila cypha</i>	Wherever found	\$22,700	\$0	\$22,700
		Chub, Owens Tui	<i>Gila bicolor ssp. snyderi</i>	Wherever found	\$31,000	\$0	\$31,000
		Chub, Pahrnagat roundtail	<i>Gila robusta jordani</i>	Wherever found	\$2,300	\$0	\$2,300
		Chub, Virgin River	<i>Gila seminuda</i> (=robusta)	Wherever found	\$25,800	\$0	\$25,800
		Dace, desert	<i>Eremichthys acros</i>	Wherever found	\$94,000	\$0	\$94,000
		Dace, Foskett speckled	<i>Rhinichthys osculus ssp.</i>	Wherever found	\$21,200	\$0	\$21,200
		Dace, Moapa	<i>Moapa coriacea</i>	Wherever found	\$5,000	\$0	\$5,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Darter, leopard	<i>Percina pantherina</i>	Wherever found	\$100	\$0	\$100
		Darter, snail	<i>Percina tanasi</i>	Wherever found	\$100	\$0	\$100
		Darter, yellowcheek	<i>Etheostoma moorei</i>	Wherever found	\$100	\$0	\$100
		Gambusia, Pecos	<i>Gambusia nobilis</i>	Wherever found	\$2,000	\$0	\$2,000
		Minnow, loach	<i>Tiaroga cobitis</i>	Wherever found	\$10,500	\$0	\$10,500
		Minnow, Rio Grande Silvery	<i>Hybognathus amarus</i>	Wherever found, except where listed as an experimental population	\$1,000	\$0	\$1,000
		Pikeminnow (=squawfish), Colorado	<i>Ptychocheilus lucius</i>	Wherever found, except where listed as an experimental population	\$44,867	\$0	\$44,867
		Poolfish, Pahrump	<i>Empetrichthys latos</i>	Wherever found	\$2,300	\$0	\$2,300
		Pupfish, desert	<i>Cyprinodon macularius</i>	Wherever found	\$355,900	\$0	\$355,900
		Pupfish, Owens	<i>Cyprinodon radiosus</i>	Wherever found	\$48,000	\$0	\$48,000
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	California Coastal ESU	\$45,000	\$0	\$45,000
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Central Valley spring-run ESU	\$8,000	\$0	\$8,000
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Lower Columbia River ESU	\$27,500	\$0	\$27,500
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Sacramento River winter- run ESU	\$8,000	\$0	\$8,000
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Snake River fall-run ESU	\$49,400	\$0	\$49,400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$232,400	\$0	\$232,400
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$64,600	\$0	\$64,600
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$55,000	\$0	\$55,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$777,150	\$0	\$777,150
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$851,500	\$0	\$851,500
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$111,300	\$0	\$111,300
		Shiner, Arkansas River	Notropis girardi	Arkansas River Basin (AR, KS, NM, OK, TX)	\$1,000	\$0	\$1,000
		Shiner, Pecos bluntnose	Notropis simus pecosensis	Wherever found	\$7,000	\$0	\$7,000
		Spikedace	Meda fulgida	Wherever found	\$10,500	\$0	\$10,500
		Spinedace, Big Spring	Lepidomeda mollispinis pratensis	Wherever found	\$4,300	\$0	\$4,300
		Spinedace, White River	Lepidomeda albivallis	Wherever found	\$13,700	\$0	\$13,700
		Springfish, Hiko White River	Crenichthys baileyi grandis	Wherever found	\$11,266	\$0	\$11,266
		Springfish, Railroad Valley	Crenichthys nevadae	Wherever found	\$2,300	\$0	\$2,300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Springfish, White River	<i>Crenichthys baileyi baileyi</i>	Wherever found	\$3,300	\$0	\$3,300
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	California Central Valley DPS	\$8,000	\$0	\$8,000
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Lower Columbia River DPS	\$69,000	\$0	\$69,000
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Middle Columbia River DPS	\$131,300	\$0	\$131,300
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Northern California DPS	\$60,000	\$0	\$60,000
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Snake River Basin DPS	\$245,800	\$0	\$245,800
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Upper Columbia River DPS	\$17,000	\$0	\$17,000
		Sturgeon (Gulf subspecies), Atlantic	<i>Acipenser oxyrinchus (=oxyrinchus) desotoi</i>	Wherever found	\$100	\$0	\$100
		Sturgeon, pallid	<i>Scaphirhynchus albus</i>	Wherever found	\$9,200	\$0	\$9,200
		Sturgeon, white	<i>Acipenser transmontanus</i>	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$1,000	\$0	\$1,000
		Sucker, Lost River	<i>Deltistes luxatus</i>	Wherever found	\$7,400	\$0	\$7,400
		Sucker, razorback	<i>Xyrauchen texanus</i>	Wherever found	\$39,700	\$0	\$39,700
		Sucker, shortnose	<i>Chasmistes brevirostris</i>	Wherever found	\$9,400	\$0	\$9,400
		Sucker, Warner	<i>Catostomus warnerensis</i>	Wherever found	\$115,040	\$0	\$115,040

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	Wherever found	\$50,000	\$0	\$50,000
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$559,972	\$0	\$559,972
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$15,490	\$0	\$15,490
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$188,178	\$0	\$188,178
		Woundfin	Plagopterus argentissimus	Wherever found, except where listed as an experimental population	\$20,800	\$0	\$20,800
	Fishes subtotal				\$4,560,463	\$0	\$4,560,463
	Clams	Bean, Choctaw	Villosa choctawensis	Wherever found	\$100	\$0	\$100
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$100	\$0	\$100
		Fanshell	Cyprogenia stegaria	Wherever found	\$200	\$0	\$200
		fatmucket, Arkansas	Lampsilis powellii	Wherever found	\$100	\$0	\$100
		Heelsplitter, Alabama (=inflated)	Potamilus inflatus	Wherever found	\$100	\$0	\$100
		Higgins eye (pearlymussel)	Lampsilis higginsii	Wherever found	\$150	\$0	\$150
		Kidneyshell, fluted	Ptychobranthus subtentum	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Kidneyshell, southern	<i>Ptychobranchnus jonesi</i>	Wherever found	\$100	\$0	\$100
		Moccasinshell, Alabama	<i>Medionidus acutissimus</i>	Wherever found	\$100	\$0	\$100
		Moccasinshell, Gulf	<i>Medionidus penicillatus</i>	Wherever found	\$100	\$0	\$100
		Mucket, pink (pearlymussel)	<i>Lampsilis abrupta</i>	Wherever found	\$100	\$0	\$100
		Mussel, scaleshell	<i>Leptodea leptodon</i>	Wherever found	\$100	\$0	\$100
		Mussel, snuffbox	<i>Epioblasma triquetra</i>	Wherever found	\$150	\$0	\$150
		Pearlshell, Louisiana	<i>Margaritifera hembeli</i>	Wherever found	\$100	\$0	\$100
		Pigtoe, fuzzy	<i>Pleurobema strodeanum</i>	Wherever found	\$100	\$0	\$100
		Pigtoe, narrow	<i>Fusconaia escambia</i>	Wherever found	\$100	\$0	\$100
		Pigtoe, rough	<i>Pleurobema plenum</i>	Wherever found	\$100	\$0	\$100
		Pocketbook, fat	<i>Potamilus capax</i>	Wherever found	\$200	\$0	\$200
		Pocketbook, Ouachita rock	<i>Arkansia wheeleri</i>	Wherever found	\$100	\$0	\$100
		Pocketbook, speckled	<i>Lampsilis streckeri</i>	Wherever found	\$100	\$0	\$100
		Rabbitsfoot	<i>Quadrula cylindrica cylindrica</i>	Wherever found	\$200	\$0	\$200
		sandshell, Southern	<i>Hamiota australis</i>	Wherever found	\$100	\$0	\$100
		Spectaclecase (mussel)	<i>Cumberlandia monodonta</i>	Wherever found	\$100	\$0	\$100
	Clams subtotal				\$2,700	\$0	\$2,700
	Snails	Limpet, Banbury Springs	<i>Lanx</i> sp.	Wherever found	\$3,750	\$0	\$3,750

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Snail, Bliss Rapids	Taylorconcha serpenticola	Wherever found	\$3,750	\$0	\$3,750
		Snail, Morro shoulderband (=Banded dune)	Helminthoglypta walkeriana	Wherever found	\$500	\$0	\$500
		Snail, Pecos assiminea	Assiminea pecos	Wherever found	\$2,500	\$0	\$2,500
		Snail, Snake River physa	Physa natricina	Wherever found	\$4,250	\$0	\$4,250
		Springsnail, Bruneau Hot	Pyrgulopsis bruneauensis	Wherever found	\$3,250	\$0	\$3,250
		Springsnail, Koster's	Juturnia kosteri	Wherever found	\$2,500	\$0	\$2,500
		Springsnail, Roswell	Pyrgulopsis roswellensis	Wherever found	\$2,500	\$0	\$2,500
	Snails subtotal				\$23,000	\$0	\$23,000
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$7,350	\$0	\$7,350
		Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$32,300	\$0	\$32,300
		Butterfly, Behren's silverspot	Speyeria zerene behrensii	Wherever found	\$6,000	\$0	\$6,000
		Butterfly, Fender's blue	Icaricia icarioides fenderi	Wherever found	\$44,100	\$0	\$44,100
		Butterfly, Quino checkerspot	Euphydryas editha quino (=E. e. wrighti)	Wherever found	\$6,500	\$0	\$6,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Butterfly, Uncompahgre fritillary	<i>Boloria acrocne</i>	Wherever found	\$10,100	\$0	\$10,100
		Checkerspot, Taylor's (=whulge)	<i>Euphydryas editha taylori</i>	Wherever found	\$500	\$0	\$500
		Moth, Kern primrose sphinx	<i>Euproserpinus euterpe</i>	Wherever found	\$2,500	\$0	\$2,500
		Skipper, Carson wandering	<i>Pseudocopaeodes eunus obscurus</i>	Wherever found	\$27,820	\$0	\$27,820
		Skipper, Dakota	<i>Hesperia dacotae</i>	Wherever found	\$15,150	\$0	\$15,150
		Skipper, Pawnee montane	<i>Hesperia leonardus montana</i>	Wherever found	\$100	\$0	\$100
		skipperling, Poweshiek	<i>Oarisma poweshiek</i>	Wherever found	\$150	\$0	\$150
	Insects subtotal				\$152,570	\$0	\$152,570
	Crustaceans	Amphipod, Noel's	<i>Gammarus desperatus</i>	Wherever found	\$2,500	\$0	\$2,500
		Crayfish, Shasta	<i>Pacifastacus fortis</i>	Wherever found	\$10,000	\$0	\$10,000
		Fairy shrimp, longhorn	<i>Branchinecta longiantenna</i>	Wherever found	\$3,500	\$0	\$3,500
		Fairy shrimp, vernal pool	<i>Branchinecta lynchi</i>	Wherever found	\$34,500	\$0	\$34,500
		Isopod, Socorro	<i>Thermosphaeroma thermophilus</i>	Wherever found	\$3,500	\$0	\$3,500
		Shrimp, Kentucky cave	<i>Palaemonias ganteri</i>	Wherever found	\$100	\$0	\$100
		Tadpole shrimp, vernal pool	<i>Lepidurus packardi</i>	Wherever found	\$30,000	\$0	\$30,000
	Crustaceans subtotal				\$84,100	\$0	\$84,100
	Flowering Plants	Aster, Florida golden	<i>Chrysopsis floridana</i>	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Barberry, Nevin's	<i>Berberis nevinii</i>	Wherever found	\$500	\$0	\$500
		Bear-poppy, Dwarf	<i>Arctomecon humilis</i>	Wherever found	\$11,000	\$0	\$11,000
		beardtongue, Parachute	<i>Penstemon debilis</i>	Wherever found	\$10,206	\$0	\$10,206
		Beardtongue, Penland	<i>Penstemon penlandii</i>	Wherever found	\$11,056	\$0	\$11,056
		Bedstraw, El Dorado	<i>Galium californicum</i> ssp. <i>sierrae</i>	Wherever found	\$17,300	\$1,500	\$18,800
		Birds-in-a-nest, white	<i>Macbridea alba</i>	Wherever found	\$1,100	\$0	\$1,100
		Bladderpod, Dudley Bluffs	<i>Lesquerella congesta</i>	Wherever found	\$12,756	\$5,000	\$17,756
		Bladderpod, kodachrome	<i>Lesquerella tumulosa</i>	Wherever found	\$1,500	\$0	\$1,500
		Blue-star, Kearney's	<i>Amsonia kearneyana</i>	Wherever found	\$4,500	\$0	\$4,500
		Bluegrass, San Bernardino	<i>Poa atropurpurea</i>	Wherever found	\$500	\$0	\$500
		Brodiaea, Chinese Camp	<i>Brodiaea pallida</i>	Wherever found	\$1,200	\$0	\$1,200
		Buckwheat, steamboat	<i>Eriogonum ovalifolium</i> var. <i>williamsiae</i>	Wherever found	\$28,407	\$0	\$28,407
		Buckwheat, Umtanum Desert	<i>Eriogonum codium</i>	Wherever found	\$850	\$0	\$850
		Bulrush, Northeastern	<i>Scirpus ancistrochaetus</i>	Wherever found	\$100	\$0	\$100
		Bush-clover, prairie	<i>Lespedeza leptostachya</i>	Wherever found	\$150	\$0	\$150
		Buttercup, autumn	<i>Ranunculus aestivalis</i> (= <i>acrifomis</i>)	Wherever found	\$3,700	\$0	\$3,700

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Butterfly plant, Colorado	<i>Gaura neomexicana</i> var. <i>coloradensis</i>	Wherever found	\$25,723	\$0	\$25,723
		Butterweed, Layne's	<i>Senecio layneae</i>	Wherever found	\$7,800	\$600	\$8,400
		Butterwort, Godfrey's	<i>Pinguicula ionantha</i>	Wherever found	\$1,100	\$0	\$1,100
		Cactus, Acuna	<i>Echinomastus erectocentrus</i> var. <i>acunensis</i>	Wherever found	\$8,900	\$0	\$8,900
		Cactus, Bakersfield	<i>Opuntia treleasei</i>	Wherever found	\$1,000	\$0	\$1,000
		Cactus, Brady pincushion	<i>Pediocactus bradyi</i>	Wherever found	\$500	\$0	\$500
		Cactus, Colorado hookless	<i>Sclerocactus glaucus</i>	Wherever found	\$130,596	\$0	\$130,596
		Cactus, Fickeisen plains	<i>Pediocactus peeblesianus fickeiseniae</i>	Wherever found	\$16,500	\$0	\$16,500
		Cactus, Knowlton's	<i>Pediocactus knowltonii</i>	Wherever found	\$5,700	\$0	\$5,700
		Cactus, Kuenzler hedgehog	<i>Echinocereus fendleri</i> var. <i>kuenzleri</i>	Wherever found	\$95,300	\$0	\$95,300
		Cactus, Mesa Verde	<i>Sclerocactus mesae-verdae</i>	Wherever found	\$6,000	\$0	\$6,000
		Cactus, Nichol's Turk's head	<i>Echinocactus horizonthalonius</i> var. <i>nicholii</i>	Wherever found	\$2,100	\$0	\$2,100
		cactus, Pariette	<i>Sclerocactus brevispinus</i>	Wherever found	\$4,826	\$0	\$4,826

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Cactus, Peebles Navajo	<i>Pediocactus peeblesianus</i> var. <i>peeblesianus</i>	Wherever found	\$16,000	\$0	\$16,000
		Cactus, Pima pineapple	<i>Coryphantha scheeri</i> var. <i>robustispina</i>	Wherever found	\$800	\$0	\$800
		Cactus, San Rafael	<i>Pediocactus despainii</i>	Wherever found	\$8,300	\$0	\$8,300
		Cactus, Siler pincushion	<i>Pediocactus</i> (=Echinocactus,=Utahia) <i>sileri</i>	Wherever found	\$12,800	\$0	\$12,800
		Cactus, Sneed pincushion	<i>Coryphantha sneedii</i> var. <i>sneedii</i>	Wherever found	\$18,000	\$0	\$18,000
		cactus, Uinta Basin hookless	<i>Sclerocactus wetlandicus</i>	Wherever found	\$11,936	\$0	\$11,936
		Cactus, Winkler	<i>Pediocactus winkleri</i>	Wherever found	\$1,400	\$0	\$1,400
		Cactus, Wright fishhook	<i>Sclerocactus wrightiae</i>	Wherever found	\$38,400	\$0	\$38,400
		Catchfly, Spalding's	<i>Silene spaldingii</i>	Wherever found	\$23,500	\$0	\$23,500
		Ceanothus, Pine Hill	<i>Ceanothus roderickii</i>	Wherever found	\$8,000	\$400	\$8,400
		Chaffseed, American	<i>Schwalbea americana</i>	Wherever found	\$100	\$0	\$100
		Clarkia, Springville	<i>Clarkia springvillensis</i>	Wherever found	\$2,500	\$0	\$2,500
		Cliffrose, Arizona	<i>Purshia</i> (=Cowania) <i>subintegra</i>	Wherever found	\$1,200	\$0	\$1,200
		Cycladenia, Jones	<i>Cycladenia humilis</i> var. <i>jonesii</i>	Wherever found	\$20,100	\$0	\$20,100
		Daisy, Parish's	<i>Erigeron parishii</i>	Wherever found	\$4,500	\$0	\$4,500
		Daisy, Willamette	<i>Erigeron decumbens</i>	Wherever found	\$50,200	\$0	\$50,200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Desert-parsley, Bradshaw's	Lomatium bradshawii	Wherever found	\$33,200	\$0	\$33,200
		Evening- primrose, San Benito	Camissonia benitensis	Wherever found	\$15,000	\$0	\$15,000
		Flannelbush, Mexican	Fremontodendron mexicanum	Wherever found	\$500	\$0	\$500
		Flannelbush, Pine Hill	Fremontodendron californicum ssp. decumbens	Wherever found	\$9,500	\$700	\$10,200
		Fleabane, Zuni	Erigeron rhizomatus	Wherever found	\$4,000	\$0	\$4,000
		Four-o'clock, MacFarlane's	Mirabilis macfarlanei	Wherever found	\$33,500	\$0	\$33,500
		Fritillary, Gentner's	Fritillaria gentneri	Wherever found	\$224,500	\$0	\$224,500
		Howellia, water	Howellia aquatilis	Wherever found	\$7,000	\$0	\$7,000
		ivesia, Webber's	Ivesia webberi	Wherever found	\$30,620	\$0	\$30,620
		Jewelflower, California	Caulanthus californicus	Wherever found	\$5,500	\$0	\$5,500
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$213,444	\$0	\$213,444
		ladies-tresses, Canelo Hills	Spiranthes delitescens	Wherever found	\$3,900	\$0	\$3,900
		Layia, beach	Layia carnosa	Wherever found	\$162,000	\$0	\$162,000
		Lily, Western	Lilium occidentale	Wherever found	\$20,000	\$0	\$20,000
		Lomatium, Cook's	Lomatium cookii	Wherever found	\$58,500	\$0	\$58,500
		Lupine, Kincaid's	Lupinus sulphureus ssp. kincaidii	Wherever found	\$60,200	\$0	\$60,200
		mallow, Gierisch	Sphaeralcea gierischii	Wherever found	\$16,700	\$0	\$16,700
		Mallow, Kern	Eremalche kernensis	Wherever found	\$24,000	\$0	\$24,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Manzanita, Ione	<i>Arctostaphylos myrtifolia</i>	Wherever found	\$1,500	\$0	\$1,500
		Manzanita, Morro	<i>Arctostaphylos morroensis</i>	Wherever found	\$500	\$0	\$500
		Milk-vetch, Coachella Valley	<i>Astragalus lentiginosus</i> var. <i>coachellae</i>	Wherever found	\$3,500	\$0	\$3,500
		Milk-vetch, Cushenbury	<i>Astragalus albens</i>	Wherever found	\$4,500	\$0	\$4,500
		Milk-vetch, Fish Slough	<i>Astragalus lentiginosus</i> var. <i>piscinensis</i>	Wherever found	\$11,000	\$0	\$11,000
		Milk-vetch, Holmgren	<i>Astragalus holmgreniorum</i>	Wherever found	\$11,500	\$0	\$11,500
		Milk-vetch, Lane Mountain	<i>Astragalus jaegerianus</i>	Wherever found	\$66,000	\$0	\$66,000
		Milk-vetch, Mancos	<i>Astragalus humillimus</i>	Wherever found	\$2,500	\$0	\$2,500
		Milk-vetch, Peirson's	<i>Astragalus magdalenae</i> var. <i>peirsonii</i>	Wherever found	\$260,000	\$0	\$260,000
		Milk-vetch, Shivwits	<i>Astragalus ampullarioides</i>	Wherever found	\$5,500	\$0	\$5,500
		milkvetch, Osterhout	<i>Astragalus osterhoutii</i>	Wherever found	\$19,312	\$0	\$19,312
		Milkweed, Welsh's	<i>Asclepias welshii</i>	Wherever found	\$29,000	\$0	\$29,000
		Morning-glory, Stebbins'	<i>Calystegia stebbinsii</i>	Wherever found	\$8,500	\$650	\$9,150
		Mustard, Penland alpine fen	<i>Eutrema penlandii</i>	Wherever found	\$15,856	\$0	\$15,856
		Niterwort, Amargosa	<i>Nitrophila mohavensis</i>	Wherever found	\$28,000	\$0	\$28,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		No common name	<i>Geocarpon minimum</i>	Wherever found	\$100	\$0	\$100
		Orchid, western prairie fringed	<i>Platanthera praeclara</i>	Wherever found	\$1,000	\$0	\$1,000
		Orchid, white fringeless	<i>Platanthera integrilabia</i>		\$100	\$0	\$100
		Orcutt grass, San Joaquin	<i>Orcuttia inaequalis</i>	Wherever found	\$500	\$0	\$500
		Orcutt grass, slender	<i>Orcuttia tenuis</i>	Wherever found	\$6,000	\$0	\$6,000
		Owl's-clover, fleshy	<i>Castilleja campestris</i> ssp. <i>succulenta</i>	Wherever found	\$500	\$0	\$500
		oxytheca, Cushenbury	<i>Oxytheca parishii</i> var. <i>goodmaniana</i>	Wherever found	\$4,500	\$0	\$4,500
		Pawpaw, beautiful	<i>Deeringothamnus pulchellus</i>	Wherever found	\$100	\$0	\$100
		Pawpaw, four- petal	<i>Asimina tetramera</i>	Wherever found	\$9,000	\$0	\$9,000
		Pennyroyal, Todsens	<i>Hedeoma todsenii</i>	Wherever found	\$31,000	\$0	\$31,000
		Penstemon, blowout	<i>Penstemon haydenii</i>	Wherever found	\$92,934	\$0	\$92,934
		Peppergrass, Slickspot	<i>Lepidium papilliferum</i>		\$211,875	\$0	\$211,875
		Phacelia, DeBeque	<i>Phacelia submutica</i>	Wherever found	\$7,100	\$0	\$7,100
		Phacelia, North Park	<i>Phacelia formosula</i>	Wherever found	\$10,356	\$0	\$10,356
		Pogonia, small whorled	<i>Isotria medeoloides</i>	Wherever found	\$100	\$0	\$100
		Pondberry	<i>Lindera melissifolia</i>	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Poppy, Sacramento prickly	<i>Argemone pleiacantha</i> ssp. <i>pinnatisecta</i>	Wherever found	\$25,620	\$0	\$25,620
		Potato-bean, Price's	<i>Apios priceana</i>		\$100	\$0	\$100
		Reed-mustard, Barneby	<i>Schoenocrambe barnebyi</i>	Wherever found	\$1,000	\$0	\$1,000
		Reed-mustard, clay	<i>Schoenocrambe argillacea</i>	Wherever found	\$11,842	\$0	\$11,842
		Reed-mustard, shrubby	<i>Schoenocrambe suffrutescens</i>	Wherever found	\$9,543	\$0	\$9,543
		Rock-cress, McDonald's	<i>Arabis macdonaldiana</i>	Wherever found	\$4,000	\$0	\$4,000
		Seagrass, Johnson's	<i>Halophila johnsonii</i>	Wherever found	\$5,100	\$0	\$5,100
		Sedge, Navajo	<i>Carex specuicola</i>	Wherever found	\$4,600	\$0	\$4,600
		Skullcap, Florida	<i>Scutellaria floridana</i>	Wherever found	\$1,100	\$0	\$1,100
		Skyrocket, Pagosa	<i>Ipomopsis polyantha</i>	Wherever found	\$12,456	\$0	\$12,456
		Sunflower, Pecos (=puzzle, =paradox)	<i>Helianthus paradoxus</i>	Wherever found	\$9,300	\$0	\$9,300
		Townsendia, Last Chance	<i>Townsendia aprica</i>	Wherever found	\$3,850	\$0	\$3,850
		Twinpod, Dudley Bluffs	<i>Physaria obcordata</i>	Wherever found	\$12,756	\$5,000	\$17,756
		Vervain, Red Hills	<i>Verbena californica</i>	Wherever found	\$1,500	\$0	\$1,500
		Wallflower, Menzies'	<i>Erysimum menziesii</i>	Wherever found	\$36,000	\$0	\$36,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Water-umbel, Huachuca	Lilaeopsis schaffneriana var. recurva	Wherever found	\$19,000	\$0	\$19,000
		wild buckwheat, clay-loving	Eriogonum pelinophilum	Wherever found	\$28,006	\$0	\$28,006
		Wild-buckwheat, gypsum	Eriogonum gypsophilum	Wherever found	\$97,000	\$0	\$97,000
		Wire-lettuce, Malheur	Stephanomeria malheurensis	Wherever found	\$1,500	\$0	\$1,500
		Wooly-threads, San Joaquin	Monolopia (=Lembertia) congdonii	Wherever found	\$82,500	\$0	\$82,500
		Yellowhead, desert	Yermo xanthocephalus	Wherever found	\$44,000	\$0	\$44,000
	Flowering Plants subtotal				\$2,803,976	\$13,850	\$2,817,826
	Lichens	Cladonia, Florida perforate	Cladonia perforata	Wherever found	\$8,100	\$0	\$8,100
	Lichens subtotal				\$8,100	\$0	\$8,100
Domestic subtotal					\$21,755,835	\$420,420	\$22,176,255
Total					\$21,755,835	\$420,420	\$22,176,255

DOI Bureau of Reclamation - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Northern long-eared	<i>Myotis septentrionalis</i>	Wherever found	\$4,500	\$0	\$4,500
		Ferret, black- footed	<i>Mustela nigripes</i>	Wherever found, except where listed as an experimental population	\$1,800	\$0	\$1,800
		Fox, San Joaquin kit	<i>Vulpes macrotis mutica</i>	wherever found	\$76,000	\$0	\$76,000
		Kangaroo rat, Fresno	<i>Dipodomys nitratoides exilis</i>	Wherever found	\$356,000	\$0	\$356,000
		Kangaroo rat, giant	<i>Dipodomys ingens</i>	Wherever found	\$500	\$0	\$500
		Kangaroo rat, Tipton	<i>Dipodomys nitratoides nitratoides</i>	Wherever found	\$5,000	\$0	\$5,000
		Mouse, New Mexico meadow jumping	<i>Zapus hudsonius luteus</i>	Wherever found	\$100,000	\$0	\$100,000
		Mouse, Preble's meadow jumping	<i>Zapus hudsonius preblei</i>	wherever found	\$2,000	\$0	\$2,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	<i>Canis lupus</i>	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$2,000	\$0	\$2,000
	Mammals subtotal				\$547,800	\$0	\$547,800
	Birds	Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$2,500	\$0	\$2,500
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$3,225,500	\$250,000	\$3,475,500
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$5,717,500	\$0	\$5,717,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$8,500	\$0	\$8,500
		Rail, Yuma clapper	Rallus longirostris yumanensis	Wherever found	\$1,364,000	\$0	\$1,364,000
		Tern, least	Sterna antillarum	interior pop.	\$26,500	\$0	\$26,500
	Birds subtotal				\$10,344,500	\$250,000	\$10,594,500
	Reptiles	Lizard, blunt-nosed leopard	Gambelia silus	Wherever found	\$800	\$0	\$800
		Snake, giant garter	Thamnophis gigas	Wherever found	\$42,500	\$0	\$42,500
	Reptiles subtotal				\$43,300	\$0	\$43,300
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$128,500	\$0	\$128,500
		Frog, Oregon spotted	Rana pretiosa	Wherever found	\$145,000	\$0	\$145,000
		Salamander, California tiger	Ambystoma californiense	U.S.A. (CA - Santa Barbara County)	\$93,000	\$0	\$93,000
		Salamander, California tiger	Ambystoma californiense	U.S.A. (Central CA DPS)	\$76,000	\$0	\$76,000
	Amphibians subtotal				\$442,500	\$0	\$442,500
	Fishes	Chub, bonytail	Gila elegans	Wherever found	\$9,322,500	\$0	\$9,322,500
		Chub, Gila	Gila intermedia	Wherever found	\$557,000	\$0	\$557,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Chub, humpback	<i>Gila cypha</i>	Wherever found	\$4,605,000	\$0	\$4,605,000
		Chub, Virgin River	<i>Gila seminuda</i> (=robusta)	Wherever found	\$8,000	\$0	\$8,000
		Cui-ui	<i>Chasmistes cujus</i>	Wherever found	\$101,500	\$0	\$101,500
		Gambusia, Pecos	<i>Gambusia nobilis</i>	Wherever found	\$13,000	\$0	\$13,000
		Minnow, loach	<i>Tiaroga cobitis</i>	Wherever found	\$557,000	\$0	\$557,000
		Minnow, Rio Grande Silvery	<i>Hybognathus amarus</i>	Wherever found, except where listed as an experimental population	\$7,000,000	\$3,000,000	\$10,000,000
		Pikeminnow (=squawfish), Colorado	<i>Ptychocheilus lucius</i>	Wherever found, except where listed as an experimental population	\$4,608,000	\$0	\$4,608,000
		Pupfish, Comanche Springs	<i>Cyprinodon elegans</i>	Wherever found	\$13,000	\$0	\$13,000
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Central Valley spring-run ESU	\$2,524,500	\$61,000	\$2,585,500
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Lower Columbia River ESU	\$524,000	\$506,000	\$1,030,000
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Sacramento River winter- run ESU	\$9,346,000	\$61,000	\$9,407,000
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Snake River fall-run ESU	\$364,000	\$506,000	\$870,000
		Salmon, Chinook	<i>Oncorhynchus</i> (=Salmo) tshawytscha	Snake River spring/summer -run ESU	\$2,643,500	\$506,000	\$3,149,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$3,074,500	\$506,000	\$3,580,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$8,000	\$0	\$8,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$8,648,500	\$0	\$8,648,500
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$354,000	\$506,000	\$860,000
		Shiner, Pecos bluntnose	Notropis simus pecosensis	Wherever found	\$500,000	\$1,100,000	\$1,600,000
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$1,400	\$0	\$1,400
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$6,230,500	\$0	\$6,230,500
		Spikedace	Meda fulgida	Wherever found	\$557,000	\$0	\$557,000
		Spinedace, Little Colorado	Lepidomeda vittata	Wherever found	\$78,000	\$0	\$78,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$8,869,000	\$61,000	\$8,930,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Central California Coast DPS	\$20,000	\$0	\$20,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$524,000	\$506,000	\$1,030,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$1,566,000	\$506,000	\$2,072,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$9,398,000	\$506,000	\$9,904,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$3,000	\$0	\$3,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$270,500	\$0	\$270,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$2,522,500	\$506,000	\$3,028,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$26,000	\$0	\$26,000
		sturgeon, green	Acipenser medirostris	Southern DPS	\$138,500	\$0	\$138,500
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$490,500	\$0	\$490,500
		Sucker, June	Chasmistes liorus	Wherever found	\$300,000	\$0	\$300,000
		Sucker, Lost River	Deltistes luxatus	Wherever found	\$1,398,000	\$0	\$1,398,000
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$10,248,000	\$0	\$10,248,000
		Sucker, shortnose	Chasmistes brevirostris	Wherever found	\$1,398,000	\$0	\$1,398,000
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	Wherever found	\$557,000	\$0	\$557,000
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$1,126,000	\$46,000	\$1,172,000
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$101,500	\$0	\$101,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Woundfin	Plagopterus argentissimus	Wherever found, except where listed as an experimental population	\$8,000	\$0	\$8,000
	Fishes subtotal				\$100,603,400	\$8,883,000	\$109,486,400
	Snails	Snail, Snake River physa	Physa natricina	Wherever found	\$159,500	\$0	\$159,500
	Snails subtotal				\$159,500	\$0	\$159,500
	Insects	Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$38,000	\$0	\$38,000
		Butterfly, Fender's blue	Icaricia icarioides fenderi	Wherever found	\$43,000	\$0	\$43,000
		Skipper, Dakota	Hesperia dacotae	Wherever found	\$1,100	\$0	\$1,100
		skipperling, Poweshiek	Oarisma poweshiek	Wherever found	\$400	\$0	\$400
	Insects subtotal				\$82,500	\$0	\$82,500
	Crustaceans	Fairy shrimp, longhorn	Branchinecta longiantenna	Wherever found	\$121,500	\$0	\$121,500
		Fairy shrimp, vernal pool	Branchinecta lynchi	Wherever found	\$27,000	\$0	\$27,000
	Crustaceans subtotal				\$148,500	\$0	\$148,500
	Flowering Plants	Cactus, Mesa Verde	Sclerocactus mesae-verdae	Wherever found	\$5,000	\$0	\$5,000
		Fiddleneck, large-flowered	Amsinckia grandiflora	Wherever found	\$115,000	\$0	\$115,000
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$150,000	\$0	\$150,000
		Lupine, Kincaid's	Lupinus sulphureus ssp. kincaidii	Wherever found	\$48,000	\$0	\$48,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Orchid, western prairie fringed	Platanthera praeclara	Wherever found	\$3,500	\$0	\$3,500
		Spurge, Hoover's	Chamaesyce hooveri	Wherever found	\$600	\$0	\$600
		Tuctoria, Greene's	Tuctoria greenei	Wherever found	\$600	\$0	\$600
	Flowering Plants subtotal				\$322,700	\$0	\$322,700
	Multi-Species				\$9,677,500	\$0	\$9,677,500
	Multi-Species subtotal				\$9,677,500	\$0	\$9,677,500
Domestic subtotal					\$122,372,200	\$9,133,000	\$131,505,200
Total					\$122,372,200	\$9,133,000	\$131,505,200

DOI National Park Service - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	bat, Florida bonneted	<i>Eumops floridanus</i>	Wherever found	\$20,000	\$0	\$20,000
		Bat, gray	<i>Myotis grisescens</i>	Wherever found	\$65,200	\$0	\$65,200
		Bat, Hawaiian hoary	<i>Lasiurus cinereus semotus</i>	Wherever found	\$86,600	\$0	\$86,600
		Bat, Indiana	<i>Myotis sodalis</i>	Wherever found	\$406,723	\$0	\$406,723
		Bat, lesser long-nosed	<i>Leptonycteris curasoae yerbabuena</i>	Wherever found	\$25,800	\$0	\$25,800
		Bat, Mexican long-nosed	<i>Leptonycteris nivalis</i>	Wherever found	\$3,500	\$0	\$3,500
		Bat, Northern long-eared	<i>Myotis septentrionalis</i>	Wherever found	\$1,023,619	\$0	\$1,023,619
		Bat, Ozark big-eared	<i>Corynorhinus (=Plecotus) townsendii ingens</i>	Wherever found	\$13,100	\$0	\$13,100
		Bat, Virginia big-eared	<i>Corynorhinus (=Plecotus) townsendii virginianus</i>	Wherever found	\$77,843	\$0	\$77,843
		Bear, grizzly	<i>Ursus arctos horribilis</i>	U.S.A., conterminous (lower 48) States, except where listed as an experimental population or delisted	\$1,589,000	\$0	\$1,589,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Ferret, black- footed	<i>Mustela nigripes</i>	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$15,000	\$0	\$15,000
		Ferret, black- footed	<i>Mustela nigripes</i>	Wherever found, except where listed as an experimental population	\$35,000	\$0	\$35,000
		Lynx, Canada	<i>Lynx canadensis</i>	Wherever Found in Contiguous U.S.	\$12,720	\$0	\$12,720
		Manatee, West Indian	<i>Trichechus manatus</i>	Wherever found	\$14,500	\$0	\$14,500
		Mouse, Anastasia Island beach	<i>Peromyscus polionotus phasma</i>	Wherever found	\$700	\$0	\$700
		Mouse, Perdido Key beach	<i>Peromyscus polionotus trissyllepsis</i>	Wherever found	\$2,000	\$0	\$2,000
		Mouse, southeastern beach	<i>Peromyscus polionotus niveiventris</i>	wherever found	\$1,000	\$0	\$1,000
		Panther, Florida	<i>Puma (=Felis) concolor coryi</i>	Wherever found	\$185,000	\$0	\$185,000
		Prairie dog, Utah	<i>Cynomys parvidens</i>	Wherever found	\$68,900	\$0	\$68,900
		Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	U.S.A. (AZ), Mexico	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	Wherever found, except where listed as an experimental population	\$88,700	\$0	\$88,700
		Seal, Hawaiian monk	<i>Monachus schauinslandi</i>	Wherever found	\$6,200	\$0	\$6,200
		Sheep, Sierra Nevada bighorn	<i>Ovis canadensis sierrae</i>	Sierra Nevada	\$50,500	\$0	\$50,500
		Squirrel, Carolina northern flying	<i>Glaucomys sabrinus coloratus</i>	Wherever found	\$5,400	\$0	\$5,400
		Whale, finback	<i>Balaenoptera physalus</i>	Wherever found	\$100	\$0	\$100
		Whale, humpback	<i>Megaptera novaeangliae</i>	Mexico DPS	\$279,422	\$0	\$279,422
		Whale, killer	<i>Orcinus orca</i>	Southern Resident DPS	\$50	\$0	\$50
		Whale, North Atlantic Right	<i>Eubalaena glacialis</i>	Wherever found	\$500	\$0	\$500
		Whale, North Pacific Right	<i>Eubalaena japonica</i>	Wherever found	\$50	\$0	\$50
		Whale, Sei	<i>Balaenoptera borealis</i>	Wherever found	\$100	\$0	\$100
		Whale, sperm	<i>Physeter catodon</i> (=macrocephalus)	Wherever found	\$100	\$0	\$100
		Wolf, gray	<i>Canis lupus</i>	MN	\$40,000	\$0	\$40,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$369,850	\$0	\$369,850
	Mammals subtotal				\$4,488,177	\$0	\$4,488,177
	Birds	akepa, Maui	Loxops ochraceus	Wherever found	\$36,250	\$0	\$36,250
		Condor, California	Gymnogyps californianus	U.S.A. (specific portions of Arizona, Nevada, and Utah)	\$32,250	\$0	\$32,250
		Condor, California	Gymnogyps californianus	U.S.A. only, except where listed as an experimental population	\$370,000	\$0	\$370,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Coot, Hawaiian	<i>Fulica americana alai</i>	Wherever found	\$78,500	\$0	\$78,500
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$13,800	\$0	\$13,800
		Falcon, Northern Aplomado	<i>Falco femoralis septentrionalis</i>	Wherever found, except where listed as an experimental population	\$37,500	\$0	\$37,500
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$1,500	\$0	\$1,500
		Goose, Hawaiian	<i>Branta (=Nesochen) sandvicensis</i>	Wherever found	\$370,927	\$0	\$370,927
		Ground-Dove, Friendly	<i>Gallicolumba stairi</i>	American Samoa DPS	\$500	\$0	\$500
		honeycreeper, (Akohekohe) crested	<i>Palmeria dolei</i>	Wherever found	\$35,000	\$0	\$35,000
		Kite, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	Wherever found	\$5,000	\$0	\$5,000
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$31,992	\$0	\$31,992
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$113,500	\$0	\$113,500
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$153,350	\$0	\$153,350
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$399,655	\$0	\$399,655
		parrotbill, Maui (Kiwikiu)	<i>Pseudonestor xanthophrys</i>	Wherever found	\$54,600	\$0	\$54,600
		petrel, Hawaiian	<i>Pterodroma sandwichensis</i>	Wherever found	\$703,327	\$0	\$703,327

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$967,422	\$0	\$967,422
		Plover, piping	Charadrius melodus	[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$24,943	\$0	\$24,943
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$69,000	\$0	\$69,000
		Rail, Yuma clapper	Rallus longirostris yumanensis	Wherever found	\$2,000	\$0	\$2,000
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$8,000	\$0	\$8,000
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Shearwater, Newell's Townsend's	<i>Puffinus auricularis newelli</i>	Wherever found	\$66,100	\$0	\$66,100
		Sparrow, Cape Sable seaside	<i>Ammodramus maritimus mirabilis</i>	Wherever found	\$390,000	\$0	\$390,000
		Stilt, Hawaiian	<i>Himantopus mexicanus knudseni</i>	Wherever found	\$78,500	\$0	\$78,500
		Stork, wood	<i>Mycteria americana</i>	AL, FL, GA, MS, NC, SC	\$41,200	\$0	\$41,200
		Storm-petrel, band-rumped	<i>Oceanodroma castro</i>	USA (HI)	\$136,500	\$0	\$136,500
		Tern, least	<i>Sterna antillarum</i>	interior pop.	\$35,000	\$0	\$35,000
		Tern, roseate	<i>Sterna dougallii dougallii</i>	Western Hemisphere except NE U.S.	\$2,500	\$0	\$2,500
		Tern, roseate	<i>Sterna dougallii dougallii</i>	northeast U.S. nesting pop.	\$40,200	\$0	\$40,200
		Vireo, Black-capped	<i>Vireo atricapilla</i>	Wherever found	\$2,500	\$0	\$2,500
		Woodpecker, ivory-billed	<i>Campephilus principalis</i>	Wherever found	\$5,000	\$0	\$5,000
		Woodpecker, red-cockaded	<i>Picoides borealis</i>	Wherever found	\$15,000	\$0	\$15,000
	Birds subtotal				\$4,321,716	\$0	\$4,321,716
	Reptiles	Alligator, American	<i>Alligator mississippiensis</i>	Wherever found	\$78,600	\$0	\$78,600
		Crocodile, American	<i>Crocodylus acutus</i>	U.S.A. (FL)	\$128,000	\$0	\$128,000
		gartersnake, northern Mexican	<i>Thamnophis eques megalops</i>	Wherever found	\$13,000	\$0	\$13,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Lizard, St. Croix ground	Ameiva polops	Wherever found	\$2,500	\$0	\$2,500
		Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$940	\$0	\$940
		Sea turtle, green	Chelonia mydas	Central North Pacific DPS	\$2,650	\$0	\$2,650
		Sea turtle, green	Chelonia mydas	Central South Pacific DPS	\$10,000	\$0	\$10,000
		Sea turtle, green	Chelonia mydas	Central West Pacific DPS	\$200	\$0	\$200
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$352,500	\$0	\$352,500
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$135,829	\$0	\$135,829
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$1,109,970	\$0	\$1,109,970
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$3,600	\$0	\$3,600
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$545,301	\$0	\$545,301
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$2,500	\$0	\$2,500
		Snake, San Francisco garter	Thamnophis sirtalis tetrataenia	Wherever found	\$6,500	\$0	\$6,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Tortoise, desert	<i>Gopherus agassizii</i>	AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	\$6,500	\$0	\$6,500
		Tortoise, desert	<i>Gopherus agassizii</i>	Wherever found, except AZ south and east of Colorado R., and Mexico	\$247,400	\$0	\$247,400
		turtle, bog	<i>Clemmys muhlenbergii</i>	U.S.A. (GA, NC, SC, TN, VA)	\$4,000	\$0	\$4,000
		Whipsnake (=striped racer), Alameda	<i>Masticophis lateralis euryxanthus</i>	Wherever found	\$75,000	\$0	\$75,000
	Reptiles subtotal				\$2,724,990	\$0	\$2,724,990
	Amphibians	Frog, California red-legged	<i>Rana draytonii</i>	Wherever found	\$102,000	\$0	\$102,000
		Frog, mountain yellow-legged	<i>Rana muscosa</i>	Northern California DPS	\$178,600	\$0	\$178,600
		Frog, Sierra Nevada Yellow-legged	<i>Rana sierrae</i>	Wherever found	\$389,200	\$0	\$389,200
		Hellbender, Ozark	<i>Cryptobranchus alleganiensis bishopi</i>	Wherever found	\$6,900	\$0	\$6,900
		Salamander, California tiger	<i>Ambystoma californiense</i>	U.S.A. (Central CA DPS)	\$2,500	\$0	\$2,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salamander, Jemez Mountains	<i>Plethodon neomexicanus</i>	Wherever found	\$150,500	\$0	\$150,500
		Salamander, Shenandoah	<i>Plethodon shenandoah</i>	Wherever found	\$15,000	\$0	\$15,000
		toad, Yosemite	<i>Anaxyrus canorus</i>	Wherever found	\$10,000	\$0	\$10,000
	Amphibians subtotal				\$854,700	\$0	\$854,700
	Fishes	Chub, bonytail	<i>Gila elegans</i>	Wherever found	\$500	\$0	\$500
		Chub, humpback	<i>Gila cypha</i>	Wherever found	\$605,290	\$0	\$605,290
		Dace, blackside	<i>Phoxinus cumberlandensis</i>	Wherever found	\$2,300	\$0	\$2,300
		Darter, duskytail	<i>Etheostoma percnurum</i>	Wherever found	\$7,569	\$0	\$7,569
		Eulachon	<i>Thaleichthys pacificus</i>	Southern DPS	\$3,850	\$0	\$3,850
		Gambusia, Big Bend	<i>Gambusia gaigei</i>	Wherever found	\$5,000	\$0	\$5,000
		Goby, tidewater	<i>Eucyclogobius newberryi</i>	Wherever found	\$4,500	\$0	\$4,500
		Madtom, smoky	<i>Noturus baileyi</i>	Wherever found	\$7,569	\$0	\$7,569
		Madtom, yellowfin	<i>Noturus flavipinnis</i>	Wherever found, except where listed as an experimental population	\$7,569	\$0	\$7,569
		Minnow, Rio Grande Silvery	<i>Hybognathus amarus</i>	Rio Grande, from Little Box Canyon to Amistad Dam	\$1,000	\$0	\$1,000
		Pikeminnow (=squawfish), Colorado	<i>Ptychocheilus lucius</i>	Wherever found, except where listed as an experimental population	\$500	\$0	\$500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pupfish, desert	Cyprinodon macularius	Wherever found	\$41,000	\$0	\$41,000
		Pupfish, Devils Hole	Cyprinodon diabolis	Wherever found	\$500,000	\$0	\$500,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$54,000	\$0	\$54,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$1,500	\$0	\$1,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$3,850	\$0	\$3,850
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$353,500	\$0	\$353,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$3,850	\$0	\$3,850
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer-run ESU	\$3,850	\$0	\$3,850
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$3,850	\$0	\$3,850
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$396,000	\$0	\$396,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$3,850	\$0	\$3,850
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$56,500	\$0	\$56,500
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU	\$10,000	\$0	\$10,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sawfish, smalltooth	<i>Pristis pectinata</i>	US DPS - Smalltooth sawfish originating from U.S. waters.	\$15,000	\$0	\$15,000
		Shiner, Topeka	<i>Notropis topeka</i> (=tristis)	Wherever found, except where listed as an experimental population	\$8,700	\$0	\$8,700
		Steelhead	<i>Oncorhynchus</i> (=Salmo) mykiss	California Central Valley DPS	\$6,500	\$0	\$6,500
		Steelhead	<i>Oncorhynchus</i> (=Salmo) mykiss	Central California Coast DPS	\$288,000	\$0	\$288,000
		Steelhead	<i>Oncorhynchus</i> (=Salmo) mykiss	Lower Columbia River DPS	\$3,850	\$0	\$3,850
		Steelhead	<i>Oncorhynchus</i> (=Salmo) mykiss	Middle Columbia River DPS	\$3,850	\$0	\$3,850
		Steelhead	<i>Oncorhynchus</i> (=Salmo) mykiss	Northern California DPS	\$56,000	\$0	\$56,000
		Steelhead	<i>Oncorhynchus</i> (=Salmo) mykiss	Puget Sound DPS	\$353,000	\$0	\$353,000
		Steelhead	<i>Oncorhynchus</i> (=Salmo) mykiss	Upper Willamette River DPS	\$3,850	\$0	\$3,850
		Sturgeon, Atlantic	<i>Acipenser oxyrinchus oxyrinchus</i>	New York Bight DPS	\$300	\$0	\$300
		Sturgeon, shortnose	<i>Acipenser brevirostrum</i>	Wherever found	\$300	\$0	\$300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$77,873	\$0	\$77,873
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	Wherever found	\$300	\$0	\$300
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$267,000	\$0	\$267,000
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$55,000	\$0	\$55,000
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$500	\$0	\$500
		Tui chub, Mohave	Gila bicolor ssp. mohavensis	Wherever found	\$18,680	\$0	\$18,680
	Fishes subtotal				\$3,236,100	\$0	\$3,236,100
	Clams	Bean, Cumberland (pearlymussel)	Villosa trabalis	Wherever found; Except where listed as Experimental Populations	\$645	\$0	\$645
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$200	\$0	\$200
		Combshell, Cumberlandian	Epioblasma brevidens	Wherever found; Except where listed as Experimental Populations	\$645	\$0	\$645
		Elktoe, Cumberland	Alasmidonta atropurpurea	Wherever found	\$645	\$0	\$645

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Fanshell	<i>Cyprogenia stegaria</i>	Wherever found	\$200	\$0	\$200
		Kidneyshell, fluted	<i>Ptychobranthus subtentum</i>	Wherever found	\$645	\$0	\$645
		Mapleleaf, winged	<i>Quadrula fragosa</i>	Wherever found, except where listed as an experimental population	\$2,600	\$0	\$2,600
		Mucket, pink (pearlymussel)	<i>Lampsilis abrupta</i>	Wherever found	\$845	\$0	\$845
		Mussel, oyster	<i>Epioblasma capsaeformis</i>	Wherever found; Except where listed as Experimental Populations	\$645	\$0	\$645
		Mussel, sheepnose	<i>Plethobasus cyphus</i>	Wherever found	\$200	\$0	\$200
		Mussel, snuffbox	<i>Epioblasma triquetra</i>	Wherever found	\$200	\$0	\$200
		Pearlymussel, dromedary	<i>Dromus dromas</i>	Wherever found; Except where listed as Experimental Populations	\$645	\$0	\$645
		Pearlymussel, littlewing	<i>Pegias fabula</i>	Wherever found	\$645	\$0	\$645
		Pigtoe, rough	<i>Pleurobema plenum</i>	Wherever found	\$200	\$0	\$200
		Rabbitsfoot	<i>Quadrula cylindrica cylindrica</i>	Wherever found	\$200	\$0	\$200
		Riffleshell, tan	<i>Epioblasma florentina walkeri</i> (=E. walkeri)	Wherever found	\$645	\$0	\$645

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Ring pink (mussel)	Obovaria retusa	Wherever found	\$200	\$0	\$200
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$845	\$0	\$845
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$4,000	\$0	\$4,000
	Clams subtotal				\$14,850	\$0	\$14,850
	Snails	Abalone, Black	Haliotis cracherodii	Wherever found	\$12,000	\$0	\$12,000
		Snail, Guam tree	Partula radiolata	Wherever found	\$100	\$0	\$100
		Snail, Humped tree	Partula gibba	Wherever found	\$100	\$0	\$100
		[no common name] Snail	Eua zebrina	Wherever found	\$8,000	\$0	\$8,000
	Snails subtotal				\$20,200	\$0	\$20,200
	Insects	Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$1,300	\$0	\$1,300
		Butterfly, Mariana eight- spot	Hypolimnna octocula marianensis	Wherever found	\$20,814	\$0	\$20,814
		Butterfly, mission blue	Icaricia icarioides missionensis	Wherever found	\$30,000	\$0	\$30,000
		Butterfly, Mitchell's satyr	Neonympha mitchellii mitchellii	Wherever found	\$250	\$0	\$250
		Butterfly, Oregon silverspot	Speyeria zerene hippolyta	Wherever found	\$3,718	\$0	\$3,718
		Butterfly, San Bruno elfin	Callophrys mossii bayensis	Wherever found	\$4,500	\$0	\$4,500
		Butterfly, Schaus swallowtail	Heraclides aristodemus ponceanus	Wherever found	\$8,000	\$0	\$8,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Checkerspot, Taylor's (=whulge)	<i>Euphydryas editha taylori</i>	Wherever found	\$500	\$0	\$500
		Damselfly, Pacific Hawaiian	<i>Megalagrion pacificum</i>	Wherever found	\$33,600	\$0	\$33,600
		Dragonfly, Hine's emerald	<i>Somatochlora hineana</i>	Wherever found	\$50	\$0	\$50
		Tiger beetle, Northeastern beach	<i>Cicindela dorsalis dorsalis</i>	Wherever found	\$1,100	\$0	\$1,100
	Insects subtotal				\$103,832	\$0	\$103,832
	Crustaceans	Shrimp, Kentucky cave	<i>Palaemonias ganteri</i>	Wherever found	\$20,000	\$0	\$20,000
	Crustaceans subtotal				\$20,000	\$0	\$20,000
	Corals	coral, Boulder star	<i>Orbicella franksi</i>	Wherever found	\$12,500	\$0	\$12,500
		Coral, elkhorn	<i>Acropora palmata</i>	Wherever found	\$36,000	\$0	\$36,000
		Coral, Lobed Star	<i>Orbicella annularis</i>	Wherever found	\$12,500	\$0	\$12,500
		Coral, Mountainous Star	<i>Orbicella faveolata</i>	Wherever found	\$12,500	\$0	\$12,500
		Coral, Pillar	<i>Dendrogyra cylindrus</i>	Wherever found	\$26,500	\$0	\$26,500
		Coral, Rough Cactus	<i>Mycetophyllia ferox</i>	Wherever found	\$12,500	\$0	\$12,500
		Coral, staghorn	<i>Acropora cervicornis</i>	Wherever found	\$88,500	\$0	\$88,500
		No common name	<i>Acropora globiceps</i>	Wherever found	\$10,000	\$0	\$10,000
		No common name	<i>Acropora jacquelinae</i>	Wherever found	\$10,000	\$0	\$10,000
		No common name	<i>Acropora retusa</i>	Wherever found	\$10,000	\$0	\$10,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		No common name	Acropora rudis	Wherever found	\$10,000	\$0	\$10,000
		No common name	Acropora speciosa	Wherever found	\$10,000	\$0	\$10,000
		No common name	Euphyllia paradivisa	Wherever found	\$10,000	\$0	\$10,000
		No common name	Isopora crateriformis	Wherever found	\$10,000	\$0	\$10,000
	Corals subtotal				\$271,000	\$0	\$271,000
	Flowering Plants	Alani	Melicope ovalis	Wherever found	\$41,400	\$0	\$41,400
		Alani	Melicope zahlbruckneri	Wherever found	\$2,000	\$0	\$2,000
		Alopecurus, Sonoma	Alopecurus aequalis var. sonomensis	Wherever found	\$1,000	\$0	\$1,000
		Amaranth, seabeach	Amaranthus pumilus	Wherever found	\$51,635	\$0	\$51,635
		Avens, spreading	Geum radiatum	Wherever found	\$3,350	\$0	\$3,350
		ballart, Menzies	Exocarpos menziesii		\$2,000	\$0	\$2,000
		Barberry, island	Berberis pinnata ssp. insularis	Wherever found	\$2,000	\$0	\$2,000
		bladderpod, Missouri	Physaria filiformis	Wherever found	\$35,000	\$0	\$35,000
		Blazingstar, Heller's	Liatris helleri	Wherever found	\$3,500	\$0	\$3,500
		Cactus, Acuna	Echinomastus erectocentrus var. acunensis	Wherever found	\$24,000	\$0	\$24,000
		Cactus, Brady pincushion	Pediocactus bradyi	Wherever found	\$1,500	\$0	\$1,500
		Cactus, Chisos Mountain hedgehog	Echinocereus chisoensis var. chisoensis	Wherever found	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Cactus, Florida semaphore	<i>Consolea corallicola</i>	Wherever found	\$3,200	\$0	\$3,200
		Cactus, Lloyd's Mariposa	<i>Echinomastus mariposensis</i>	Wherever found	\$1,000	\$0	\$1,000
		Cactus, Peebles Navajo	<i>Pediocactus peeblesianus</i> var. <i>peeblesianus</i>	Wherever found	\$200	\$0	\$200
		Cactus, Winkler	<i>Pediocactus winkleri</i>	Wherever found	\$32,000	\$0	\$32,000
		Cactus, Wright fishhook	<i>Sclerocactus wrightiae</i>	Wherever found	\$32,000	\$0	\$32,000
		Catchfly, Spalding's	<i>Silene spaldingii</i>	Wherever found	\$1,500	\$0	\$1,500
		Clarkia, Presidio	<i>Clarkia franciscana</i>	Wherever found	\$2,000	\$0	\$2,000
		Cory cactus, bunched	<i>Coryphantha ramillosa</i>	Wherever found	\$1,000	\$0	\$1,000
		Cycladenia, Jones	<i>Cycladenia humilis</i> var. <i>jonesii</i>	Wherever found	\$100	\$0	\$100
		Daisy, Parish's	<i>Erigeron parishii</i>	Wherever found	\$3,000	\$0	\$3,000
		Dudleya, Verity's	<i>Dudleya verityi</i>	Wherever found	\$1,000	\$0	\$1,000
		Dwarf-flax, Marin	<i>Hesperolinon congestum</i>	Wherever found	\$2,000	\$0	\$2,000
		Evening- primrose, Eureka Valley	<i>Oenothera avita</i> ssp. <i>eurekensis</i>		\$50,000	\$0	\$50,000
		Geranium, Hawaiian red-flowered	<i>Geranium arboreum</i>	Wherever found	\$62,000	\$0	\$62,000
		Grass, Eureka Dune	<i>Swallenia alexandrae</i>		\$50,000	\$0	\$50,000
		Haha	<i>Cyanea asplenifolia</i>	Wherever found	\$62,000	\$0	\$62,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Haha	<i>Cyanea copelandii</i> ssp. <i>haleakalaensis</i>	Wherever found	\$41,400	\$0	\$41,400
		Haha	<i>Cyanea hamatiflora</i> ssp. <i>hamatiflora</i>	Wherever found	\$20,700	\$0	\$20,700
		Haha	<i>Cyanea kunthiana</i>	Wherever found	\$20,700	\$0	\$20,700
		haha	<i>Cyanea maritae</i>	Wherever found	\$20,700	\$0	\$20,700
		Haha	<i>Cyanea shipmanii</i>	Wherever found	\$13,000	\$0	\$13,000
		Haha	<i>Cyanea stictophylla</i>	Wherever found	\$3,000	\$0	\$3,000
		haiwale	<i>Cyrtandra ferripilosa</i>	Wherever found	\$62,000	\$0	\$62,000
		Hala pepe	<i>Pleomele hawaiiensis</i>	Wherever found	\$5,000	\$0	\$5,000
		Hau kuahiwi	<i>Hibiscadelphus giffardianus</i>	Wherever found	\$2,500	\$0	\$2,500
		Ha`iwale	<i>Cyrtandra giffardii</i>	Wherever found	\$2,500	\$0	\$2,500
		Holei	<i>Ochrosia haleakalae</i>	Wherever found	\$2,500	\$0	\$2,500
		Honohono	<i>Haplostachys haplostachya</i>	Wherever found	\$1,000	\$0	\$1,000
		Ihi	<i>Portulaca villosa</i>	Wherever found	\$750	\$0	\$750
		Ischaemum, Hilo	<i>Ischaemum byrone</i>	Wherever found	\$20,700	\$0	\$20,700
		Koki`o ke`oke`o	<i>Hibiscus arnottianus</i> ssp. <i>immaculatus</i>	Wherever found	\$500	\$0	\$500
		Ko`oko`olau	<i>Bidens campylothea pentamera</i>	Wherever found	\$41,400	\$0	\$41,400
		Ko`oko`olau	<i>Bidens campylothea waihoiensis</i>	Wherever found	\$41,400	\$0	\$41,400
		Ko`oko`olau	<i>Bidens micrantha ctenophylla</i>	Wherever found	\$1,000	\$0	\$1,000
		Ko`oko`olau	<i>Bidens micrantha</i> ssp. <i>kalealaha</i>	Wherever found	\$41,400	\$0	\$41,400
		Kuahiwi laukahi	<i>Plantago hawaiiensis</i>	Wherever found	\$1,500	\$0	\$1,500
		Kuahiwi laukahi	<i>Plantago princeps</i>	Wherever found	\$20,700	\$0	\$20,700

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Kula wahine noho	Isodendrion pyrifolium	Wherever found	\$1,000	\$0	\$1,000
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$200	\$0	\$200
		Layia, beach	Layia carnosa	Wherever found	\$1,900	\$0	\$1,900
		Lessingia, San Francisco	Lessingia germanorum (=L.g. var. germanorum)	Wherever found	\$2,000	\$0	\$2,000
		Liveforever, Santa Barbara Island	Dudleya traskiae	Wherever found	\$1,000	\$0	\$1,000
		Lo`ulu	Pritchardia maideniana	Wherever found	\$2,000	\$0	\$2,000
		Lupine, clover	Lupinus tidestromii	Wherever found	\$3,000	\$0	\$3,000
		Makou	Peucedanum sandwicense	Wherever found	\$5,000	\$0	\$5,000
		Manzanita, Franciscan	Arctostaphylos franciscana	Wherever found	\$60,000	\$0	\$60,000
		Manzanita, Presidio	Arctostaphylos hookeri var. ravenii	Wherever found	\$2,000	\$0	\$2,000
		Manzanita, Santa Rosa Island	Arctostaphylos confertiflora	Wherever found	\$1,600	\$0	\$1,600
		Ma`o hau hele, (=native yellow hibiscus)	Hibiscus brackenridgei	Wherever found	\$1,000	\$0	\$1,000
		Milk-vetch, Sentry	Astragalus cremnophylax var. cremnophylax	Wherever found	\$80,000	\$0	\$80,000
		Milk-vetch, Shivwits	Astragalus ampullarioides	Wherever found	\$5,500	\$0	\$5,500
		Milk-vetch, triple-ribbed	Astragalus tricarinatus	Wherever found	\$3,000	\$0	\$3,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		monkey-flower, Michigan	Mimulus michiganensis	Wherever found	\$1,540	\$0	\$1,540
		Nanu	Gardenia remyi	Wherever found	\$20,700	\$0	\$20,700
		Naupaka, dwarf	Scaevola coriacea	Wherever found	\$5,000	\$0	\$5,000
		No common name	Neraudia ovata	Wherever found	\$2,500	\$0	\$2,500
		No common name	Phyllostegia bracteata	Wherever found	\$62,000	\$0	\$62,000
		No common name	Phyllostegia brevidens	Wherever found	\$62,000	\$0	\$62,000
		No common name	Phyllostegia haliakalae	Wherever found	\$62,000	\$0	\$62,000
		No common name	Phyllostegia stachyoides	Wherever found	\$2,500	\$0	\$2,500
		No common name	Pittosporum halophilum	Wherever found	\$2,000	\$0	\$2,000
		No common name	Pittosporum hawaiiense	Wherever found	\$13,000	\$0	\$13,000
		No common name	Sanicula sandwicensis	Wherever found	\$41,900	\$0	\$41,900
		No common name	Schiedea haleakalensis	Wherever found	\$62,000	\$0	\$62,000
		No common name	Tetramolopium rockii	Wherever found	\$5,000	\$0	\$5,000
		No common name	Wikstroemia villosa	Wherever found	\$20,700	\$0	\$20,700
		Nohoanu	Geranium hanaense	Wherever found	\$20,700	\$0	\$20,700
		Nohoanu	Geranium multiflorum	Wherever found	\$20,700	\$0	\$20,700
		nui, haha	Cyanea horrida	Wherever found	\$20,700	\$0	\$20,700
		Ohai	Sesbania tomentosa	Wherever found	\$4,552	\$0	\$4,552

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Orchid, western prairie fringed	<i>Platanthera praeclara</i>	Wherever found	\$8,000	\$0	\$8,000
		Orchid, white fringeless	<i>Platanthera integrilabia</i>		\$28,000	\$0	\$28,000
		paintbrush, golden	<i>Castilleja levisecta</i>	Wherever found	\$14,000	\$0	\$14,000
		Paintbrush, Tiburon	<i>Castilleja affinis</i> ssp. <i>neglecta</i>	Wherever found	\$1,000	\$0	\$1,000
		Panicgrass, Carter's	<i>Panicum fauriei</i> var. <i>carteri</i>	Wherever found	\$1,000	\$0	\$1,000
		Phlox, Texas trailing	<i>Phlox nivalis</i> ssp. <i>texensis</i>	Wherever found	\$5,000	\$0	\$5,000
		Pink, swamp	<i>Helonias bullata</i>	Wherever found	\$3,500	\$0	\$3,500
		Pitcher-plant, green	<i>Sarracenia oreophila</i>	Wherever found	\$3,920	\$0	\$3,920
		Pogonia, small whorled	<i>Isotria medeoloides</i>	Wherever found	\$5,500	\$0	\$5,500
		Popolo	<i>Solanum nelsonii</i>	Wherever found	\$5,000	\$0	\$5,000
		Potentilla, Hickman's	<i>Potentilla hickmanii</i>	Wherever found	\$115,000	\$0	\$115,000
		Po`e	<i>Portulaca sclerocarpa</i>	Wherever found	\$4,000	\$0	\$4,000
		Prickly-apple, fragrant	<i>Cereus eriophorus</i> var. <i>fragrans</i>	Wherever found	\$1,000	\$0	\$1,000
		Pua `ala	<i>Brighamia rockii</i>	Wherever found	\$500	\$0	\$500
		Reedgrass, Maui	<i>Calamagrostis expansa</i>	Wherever found	\$20,700	\$0	\$20,700
		Rosemary, Cumberland	<i>Conradina verticillata</i>	Wherever found	\$1,200	\$0	\$1,200
		Sandwort, Cumberland	<i>Arenaria cumberlandensis</i>	Wherever found	\$800	\$0	\$800
		Sandwort, Marsh	<i>Arenaria paludicola</i>	Wherever found	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Silversword, Mauna Loa (=Ka'u)	Argyroxiphium kauense	Wherever found	\$5,000	\$0	\$5,000
		Skullcap, large-flowered	Scutellaria montana	Wherever found	\$1,500	\$0	\$1,500
		Spineflower, Sonoma	Chorizanthe valida	Wherever found	\$1,200	\$0	\$1,200
		Thistle, Pitcher's	Cirsium pitcheri	Wherever found	\$65,860	\$0	\$65,860
		Townsendia, Last Chance	Townsendia aprica	Wherever found	\$400	\$0	\$400
		`Ahinahina	Argyroxiphium sandwicense ssp. macrocephalum	Wherever found	\$124,100	\$0	\$124,100
		`Aiea	Nothoestrum latifolium	Wherever found	\$20,700	\$0	\$20,700
		`aku	Cyanea tritomantha	Wherever found	\$2,500	\$0	\$2,500
		`Awikiwiki	Canavalia molokaiensis	Wherever found	\$5,000	\$0	\$5,000
		`Ena`ena	Pseudognaphalium (=Gnaphalium) sandwicense var. molokaiense	Wherever found	\$500	\$0	\$500
		`Oha wai	Clermontia lindseyana	Wherever found	\$2,500	\$0	\$2,500
		`Oha wai	Clermontia peleana	Wherever found	\$4,000	\$0	\$4,000
		`Oha wai	Clermontia samuelii	Wherever found	\$62,000	\$0	\$62,000
		`Ohe	Joinvillea ascendens ascendens	Wherever found	\$20,700	\$0	\$20,700
	Flowering Plants subtotal				\$1,969,507	\$0	\$1,969,507
	Ferns and Allies	fern, Maui	Microlepidia strigosa var. mauiensis	Wherever found	\$20,700	\$0	\$20,700
		No common name	Asplenium peruvianum var. insulare	Wherever found	\$20,700	\$0	\$20,700

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wawae`iole	Huperzia mannii	Wherever found	\$20,700	\$0	\$20,700
	Ferns and Allies subtotal				\$62,100	\$0	\$62,100
Domestic subtotal					\$18,087,172	\$0	\$18,087,172
Foreign	Fishes	Blindcat, Mexican (catfish)	Prietella phreatophila	Wherever found	\$8,000	\$0	\$8,000
	Fishes subtotal				\$8,000	\$0	\$8,000
Foreign subtotal					\$8,000	\$0	\$8,000
Total					\$18,095,172	\$0	\$18,095,172

DOI US Geological Survey - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	bat, Florida bonneted	<i>Eumops floridanus</i>	Wherever found	\$12,043	\$0	\$12,043
		Bat, gray	<i>Myotis grisescens</i>	Wherever found	\$11,803	\$0	\$11,803
		Bat, Hawaiian hoary	<i>Lasiurus cinereus semotus</i>	Wherever found	\$187,974	\$0	\$187,974
		Bat, Indiana	<i>Myotis sodalis</i>	Wherever found	\$147,443	\$0	\$147,443
		Bat, lesser long-nosed	<i>Leptonycteris curasoae yerbabuena</i>	Wherever found	\$14,013	\$0	\$14,013
		Bat, Mexican long-nosed	<i>Leptonycteris nivalis</i>	Wherever found	\$13,243	\$0	\$13,243
		Bat, Northern long-eared	<i>Myotis septentrionalis</i>	Wherever found	\$275,192	\$0	\$275,192
		Bat, Ozark big-eared	<i>Corynorhinus (=Plecotus) townsendii ingens</i>	Wherever found	\$10,183	\$0	\$10,183
		Bat, Virginia big-eared	<i>Corynorhinus (=Plecotus) townsendii virginianus</i>	Wherever found	\$9,943	\$0	\$9,943
		Bear, grizzly	<i>Ursus arctos horribilis</i>	U.S.A., conterminous (lower 48) States, except where listed as an experimental population or delisted	\$669,923	\$0	\$669,923
		Bear, polar	<i>Ursus maritimus</i>	Wherever found	\$1,869,069	\$0	\$1,869,069

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Ferret, black- footed	<i>Mustela nigripes</i>	Wherever found, except where listed as an experimental population	\$144,162	\$0	\$144,162
		Jaguar	<i>Panthera onca</i>	Wherever found	\$6,000	\$0	\$6,000
		Kangaroo rat, giant	<i>Dipodomys ingens</i>	Wherever found	\$3,000	\$0	\$3,000
		Kangaroo rat, Stephens'	<i>Dipodomys stephensi</i> (incl. <i>D. cascus</i>)	Wherever found	\$3,000	\$0	\$3,000
		Lynx, Canada	<i>Lynx canadensis</i>	Wherever Found in Contiguous U.S.	\$204,665	\$0	\$204,665
		Manatee, West Indian	<i>Trichechus manatus</i>	Wherever found	\$1,671,700	\$0	\$1,671,700
		Mouse, Choctawhatche beach	<i>Peromyscus polionotus allophrys</i>	Wherever found	\$1,500	\$0	\$1,500
		Mouse, Pacific pocket	<i>Perognathus longimembris pacificus</i>	Wherever found	\$45,424	\$0	\$45,424
		Mouse, Perdido Key beach	<i>Peromyscus polionotus trissyllepsis</i>	Wherever found	\$1,500	\$0	\$1,500
		Mouse, Preble's meadow jumping	<i>Zapus hudsonius preblei</i>	wherever found	\$8,281	\$0	\$8,281
		Mouse, St. Andrew beach	<i>Peromyscus polionotus peninsularis</i>	Wherever found	\$1,500	\$0	\$1,500
		Ocelot	<i>Leopardus (=Felis) pardalis</i>	wherever found	\$6,000	\$0	\$6,000
		Otter, Northern Sea	<i>Enhydra lutris kenyoni</i>	Southwest Alaska DPS	\$60,300	\$0	\$60,300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Otter, southern sea	Enhydra lutris nereis	Wherever found	\$478,442	\$0	\$478,442
		Panther, Florida	Puma (=Felis) concolor coryi	Wherever found	\$24,600	\$0	\$24,600
		Prairie dog, Utah	Cynomys parvidens	Wherever found	\$16,814	\$0	\$16,814
		Puma (=cougar), eastern	Puma (=Felis) concolor cougar	Wherever found	\$1,500	\$0	\$1,500
		Puma (=mountain lion)	Puma (=Felis) concolor (all subsp. except coryi)	FL	\$3,000	\$0	\$3,000
		Rabbit, Lower Keys marsh	Sylvilagus palustris hefneri	Wherever found	\$2,000	\$0	\$2,000
		Sea lion, Steller	Eumetopias jubatus	Western DPS	\$7,002	\$0	\$7,002
		Seal, Ringed	Phoca (=Pusa) hispida hispida	Wherever found	\$60	\$0	\$60
		Sheep, Sierra Nevada bighorn	Ovis canadensis sierrae	Sierra Nevada	\$133,928	\$0	\$133,928
		Squirrel, Carolina northern flying	Glaucomys sabrinus coloratus	Wherever found	\$7,500	\$0	\$7,500
		Squirrel, Northern Idaho Ground	Uroditellus brunneus	Wherever found	\$23,892	\$0	\$23,892
		Wolf, gray	Canis lupus	MN	\$436,500	\$0	\$436,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	<i>Canis lupus</i>	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$33,700	\$0	\$33,700
		Wolf, gray	<i>Canis lupus</i>	WY, EXPN population	\$24,400	\$0	\$24,400
		Wolf, Mexican	<i>Canis lupus baileyi</i>	U.S.A. (portions of AZ and NM)see 17.84(k)	\$1,500	\$0	\$1,500
		Wolf, red	<i>Canis rufus</i>	U.S.A. (portions of NC and TN)	\$3,600	\$0	\$3,600
		Woodrat, Key Largo	<i>Neotoma floridana smalli</i>	Wherever found	\$3,051	\$0	\$3,051
	Mammals subtotal				\$6,579,350	\$0	\$6,579,350
	Birds	Akekee	<i>Loxops caeruleirostris</i>	Wherever found	\$64,215	\$0	\$64,215

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		akepa, Hawaii	<i>Loxops coccineus</i>	Wherever found	\$26,635	\$0	\$26,635
		akiapolaau	<i>Hemignathus wilsoni</i>	Wherever found	\$26,635	\$0	\$26,635
		Akikiki	<i>Oreomystis bairdi</i>	Wherever found	\$64,215	\$0	\$64,215
		Albatross, short-tailed	<i>Phoebastria (=Diomedea) albatrus</i>	Wherever found	\$1,956	\$0	\$1,956
		Bobwhite, masked (quail)	<i>Colinus virginianus ridgwayi</i>	Wherever found	\$6,000	\$0	\$6,000
		Condor, California	<i>Gymnogyps californianus</i>	U.S.A. (specific portions of Arizona, Nevada, and Utah)	\$360	\$0	\$360
		Condor, California	<i>Gymnogyps californianus</i>	U.S.A. only, except where listed as an experimental population	\$165,267	\$0	\$165,267
		Coot, Hawaiian	<i>Fulica americana alai</i>	Wherever found	\$50,237	\$0	\$50,237
		Crane, Mississippi sandhill	<i>Grus canadensis pulla</i>	Wherever found	\$60	\$0	\$60
		Crane, whooping	<i>Grus americana</i>	U.S.A (Southwestern Louisiana)	\$215,000	\$0	\$215,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Crane, whooping	<i>Grus americana</i>	U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	\$1,156,480	\$0	\$1,156,480
		Crane, whooping	<i>Grus americana</i>	U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	\$200,000	\$0	\$200,000
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$142,829	\$0	\$142,829
		Creepers, Hawaii	<i>Oreomystis mana</i>	Wherever found	\$26,635	\$0	\$26,635
		Crow, Hawaiian (=alala)	<i>Corvus hawaiiensis</i>	Wherever found	\$25,000	\$0	\$25,000
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$4,500	\$0	\$4,500
		Duck, Hawaiian (=koloa)	<i>Anas wyvilliana</i>	Wherever found	\$139,549	\$0	\$139,549
		Duck, Laysan	<i>Anas laysanensis</i>	Wherever found	\$111,178	\$0	\$111,178
		Eider, spectacled	<i>Somateria fischeri</i>	Wherever found	\$27,500	\$0	\$27,500
		Eider, Steller's	<i>Polysticta stelleri</i>	AK breeding pop.	\$20,000	\$0	\$20,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$33,267	\$0	\$33,267
		gallinule, Hawaiian common	<i>Gallinula galeata sandvicensis</i>	Wherever found	\$20,000	\$0	\$20,000
		Gnatcatcher, coastal California	<i>Polioptila californica californica</i>	Wherever found	\$26,000	\$0	\$26,000
		Goose, Hawaiian	<i>Branta (=Nesochen) sandvicensis</i>	Wherever found	\$6,600	\$0	\$6,600
		Hawk, Puerto Rican broad- winged	<i>Buteo platypterus brunnescens</i>	Wherever found	\$15,000	\$0	\$15,000
		Hawk, Puerto Rican sharp- shinned	<i>Accipiter striatus venator</i>	Wherever found	\$30,000	\$0	\$30,000
		honeycreeper, (Akohekohe) crested	<i>Palmeria dolei</i>	Wherever found	\$251	\$0	\$251
		Kingfisher, Guam Micronesian	<i>Halcyon cinnamomina cinnamomina</i>	Wherever found	\$38,249	\$0	\$38,249
		Kite, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	Wherever found	\$35,000	\$0	\$35,000
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$34,400	\$0	\$34,400
		Moorhen, Mariana common	<i>Gallinula chloropus guami</i>	Wherever found	\$600	\$0	\$600
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$69,646	\$0	\$69,646
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$248,529	\$0	\$248,529

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		parrotbill, Maui (Kiwikiu)	Pseudonestor xanthophrys	Wherever found	\$251	\$0	\$251
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$1,015,930	\$0	\$1,015,930
		Plover, piping	Charadrius melodus	[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$15,080	\$0	\$15,080
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$19,480	\$0	\$19,480
		Rail, California clapper	Rallus longirostris obsoletus	Wherever found	\$97,143	\$0	\$97,143
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$278,408	\$0	\$278,408

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Shearwater, Newell's Townsend's	<i>Puffinus auricularis newelli</i>	Wherever found	\$300	\$0	\$300
		Sparrow, Cape Sable seaside	<i>Ammodramus maritimus mirabilis</i>	Wherever found	\$58,000	\$0	\$58,000
		Stilt, Hawaiian	<i>Himantopus mexicanus knudseni</i>	Wherever found	\$29,037	\$0	\$29,037
		Swiftlet, Mariana gray	<i>Aerodramus vanikorensis bartschi</i>	Wherever found	\$10,713	\$0	\$10,713
		Tern, California least	<i>Sterna antillarum browni</i>	Wherever found	\$18,680	\$0	\$18,680
		Tern, least	<i>Sterna antillarum</i>	interior pop.	\$97,926	\$0	\$97,926
		Tern, roseate	<i>Sterna dougallii dougallii</i>	Western Hemisphere except NE U.S.	\$133,000	\$0	\$133,000
		Tern, roseate	<i>Sterna dougallii dougallii</i>	northeast U.S. nesting pop.	\$3,000	\$0	\$3,000
		Thrush, small Kauai (=puaiohi)	<i>Myadestes palmeri</i>	Wherever found	\$5,345	\$0	\$5,345
		Vireo, least Bell's	<i>Vireo bellii pusillus</i>	Wherever found	\$100,000	\$0	\$100,000
		Warbler, Kirtland's	<i>Setophaga kirtlandii</i> (= <i>Dendroica kirtlandii</i>)	Wherever found	\$24,000	\$0	\$24,000
		Woodpecker, red-cockaded	<i>Picoides borealis</i>	Wherever found	\$147,919	\$0	\$147,919
	Birds subtotal				\$5,086,005	\$0	\$5,086,005
	Reptiles	Alligator, American	<i>Alligator mississippiensis</i>	Wherever found	\$10,620	\$0	\$10,620
		Boa, Puerto Rican	<i>Epicrates inornatus</i>	Wherever found	\$12,000	\$0	\$12,000
		Crocodile, American	<i>Crocodylus acutus</i>	U.S.A. (FL)	\$193,000	\$0	\$193,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		gartersnake, narrow-headed	Thamnophis rufipunctatus	Wherever found	\$24,090	\$0	\$24,090
		gartersnake, northern Mexican	Thamnophis eques megalops	Wherever found	\$20,430	\$0	\$20,430
		Lizard, blunt-nosed leopard	Gambelia silus	Wherever found	\$10,000	\$0	\$10,000
		Lizard, Coachella Valley fringe-toed	Uma inornata	Wherever found	\$97,001	\$0	\$97,001
		Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$5,000	\$0	\$5,000
		Sea turtle, green	Chelonia mydas	Central North Pacific DPS	\$10,737	\$0	\$10,737
		Sea turtle, green	Chelonia mydas	Central South Pacific DPS	\$8,327	\$0	\$8,327
		Sea turtle, green	Chelonia mydas	Central West Pacific DPS	\$6,227	\$0	\$6,227
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$44,000	\$0	\$44,000
		Sea turtle, green	Chelonia mydas	South Atlantic DPS	\$18,680	\$0	\$18,680
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$146,800	\$0	\$146,800
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$139,000	\$0	\$139,000
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$141,133	\$0	\$141,133

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, olive ridley	<i>Lepidochelys olivacea</i>	Wherever found, except when listed as endangered under 50 CFR 224.101	\$300	\$0	\$300
		Snake, eastern indigo	<i>Drymarchon corais couperi</i>	Wherever found	\$12,000	\$0	\$12,000
		Snake, giant garter	<i>Thamnophis gigas</i>	Wherever found	\$127,035	\$0	\$127,035
		Snake, San Francisco garter	<i>Thamnophis sirtalis tetrataenia</i>	Wherever found	\$6,927	\$0	\$6,927
		Tortoise, desert	<i>Gopherus agassizii</i>	Wherever found, except AZ south and east of Colorado R., and Mexico	\$534,231	\$0	\$534,231
		Turtle, Plymouth Redbelly	<i>Pseudemys rubriventris bangsi</i>	Wherever found	\$600	\$0	\$600
		Whipsnake (=striped racer), Alameda	<i>Masticophis lateralis euryxanthus</i>	Wherever found	\$16,927	\$0	\$16,927
	Reptiles subtotal				\$1,585,065	\$0	\$1,585,065
	Amphibians	Frog, California red-legged	<i>Rana draytonii</i>	Wherever found	\$130,601	\$0	\$130,601
		Frog, Chiricahua leopard	<i>Rana chiricahuensis</i>	Wherever found	\$94,738	\$0	\$94,738
		Frog, dusky gopher	<i>Rana sevosa</i>	Wherever found	\$4,000	\$0	\$4,000
		Frog, mountain yellow-legged	<i>Rana muscosa</i>	Northern California DPS	\$30,951	\$0	\$30,951

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Frog, mountain yellow-legged	<i>Rana muscosa</i>	Southern California DPS	\$18,578	\$0	\$18,578
		Frog, Oregon spotted	<i>Rana pretiosa</i>	Wherever found	\$227,826	\$0	\$227,826
		Frog, Sierra Nevada Yellow-legged	<i>Rana sierrae</i>	Wherever found	\$80,300	\$0	\$80,300
		Salamander, frosted flatwoods	<i>Ambystoma cingulatum</i>	Wherever found	\$15,900	\$0	\$15,900
		Salamander, Shenandoah	<i>Plethodon shenandoah</i>	Wherever found	\$12,800	\$0	\$12,800
		Salamander, Sonora tiger	<i>Ambystoma tigrinum stebbinsi</i>	Wherever found	\$9,780	\$0	\$9,780
		Toad, arroyo (=arroyo southwestern)	<i>Anaxyrus californicus</i>	Wherever found	\$11,000	\$0	\$11,000
		Toad, Wyoming	<i>Bufo hemiophrys baxteri</i>	Wherever found	\$2,520	\$0	\$2,520
		toad, Yosemite	<i>Anaxyrus canorus</i>	Wherever found	\$61,722	\$0	\$61,722
	Amphibians subtotal				\$700,716	\$0	\$700,716
	Fishes	Cavefish, Ozark	<i>Amblyopsis rosae</i>	Wherever found	\$6,000	\$0	\$6,000
		Chub, bonytail	<i>Gila elegans</i>	Wherever found	\$1,750	\$0	\$1,750
		Chub, Borax Lake	<i>Gila boraxobius</i>	Wherever found	\$3,000	\$0	\$3,000
		Chub, humpback	<i>Gila cypha</i>	Wherever found	\$68,805	\$0	\$68,805
		Chub, Pahrnagat roundtail	<i>Gila robusta jordani</i>	Wherever found	\$1,500	\$0	\$1,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Chub, spotfin	Erimonax monachus	Wherever found, except where listed as an experimental population	\$44,261	\$0	\$44,261
		Dace, Ash Meadows speckled	Rhinichthys osculus nevadensis	Wherever found	\$1,500	\$0	\$1,500
		Dace, desert	Eremichthys acros	Wherever found	\$9,000	\$0	\$9,000
		Dace, Foskett speckled	Rhinichthys osculus ssp.	Wherever found	\$3,000	\$0	\$3,000
		Dace, Kendall Warm Springs	Rhinichthys osculus thermalis	Wherever found	\$2,500	\$0	\$2,500
		Dace, Moapa	Moapa coriacea	Wherever found	\$1,500	\$0	\$1,500
		Darter, amber	Percina antesella	Wherever found	\$25,800	\$0	\$25,800
		Darter, Cherokee	Etheostoma scotti	Wherever found	\$25,800	\$0	\$25,800
		Darter, Etowah	Etheostoma etowahae	Wherever found	\$25,800	\$0	\$25,800
		Darter, fountain	Etheostoma fonticola	Wherever found	\$10,474	\$0	\$10,474
		Goby, tidewater	Eucyclogobius newberryi	Wherever found	\$6,000	\$0	\$6,000
		Logperch, Conasauga	Percina jenkinsi	Wherever found	\$25,800	\$0	\$25,800
		Logperch, Roanoke	Percina rex	Wherever found	\$6,000	\$0	\$6,000
		Madtom, Neosho	Noturus placidus	Wherever found	\$19,422	\$0	\$19,422
		Minnow, Rio Grande Silvery	Hybognathus amarus	Wherever found, except where listed as an experimental population	\$22,500	\$0	\$22,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pikeminnow (=squawfish), Colorado	Ptychocheilus lucius	Wherever found, except where listed as an experimental population	\$13,750	\$0	\$13,750
		Pupfish, Ash Meadows Amargosa	Cyprinodon nevadensis mionectes	Wherever found	\$1,500	\$0	\$1,500
		Pupfish, Devils Hole	Cyprinodon diabolis	Wherever found	\$1,500	\$0	\$1,500
		Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$144,479	\$0	\$144,479
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$15,000	\$0	\$15,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$3,000	\$0	\$3,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$7,474	\$0	\$7,474
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$114,324	\$0	\$114,324
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter- run ESU	\$378,687	\$0	\$378,687
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$3,000	\$0	\$3,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer -run ESU	\$3,000	\$0	\$3,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$7,474	\$0	\$7,474

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$9,000	\$0	\$9,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$7,974	\$0	\$7,974
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$19,020	\$0	\$19,020
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU	\$70,222	\$0	\$70,222
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$4,500	\$0	\$4,500
		Shiner, Arkansas River	Notropis girardi	Arkansas River Basin (AR, KS, NM, OK, TX)	\$75,000	\$0	\$75,000
		Shiner, blue	Cyprinella caerulea	Wherever found	\$25,800	\$0	\$25,800

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Shiner, Topeka	<i>Notropis topeka</i> (=tristis)	U.S.A. (MO-specified portions of Little Creek, Big Muddy Creek, and Spring Creek watersheds in Adair, Gentry, Harrison, Putnam, Sullivan, and Worth Counties; see 17.84(d)(1)(i))	\$15,000	\$0	\$15,000
		Shiner, Topeka	<i>Notropis topeka</i> (=tristis)	Wherever found, except where listed as an experimental population	\$60,000	\$0	\$60,000
		Smelt, delta	<i>Hypomesus transpacificus</i>	Wherever found	\$105,658	\$0	\$105,658
		Spinedace, White River	<i>Lepidomeda albivallis</i>	Wherever found	\$1,500	\$0	\$1,500
		Springfish, White River	<i>Crenichthys baileyi baileyi</i>	Wherever found	\$1,500	\$0	\$1,500
		Steelhead	<i>Oncorhynchus</i> (=Salmo) mykiss	California Central Valley DPS	\$9,000	\$0	\$9,000
		Steelhead	<i>Oncorhynchus</i> (=Salmo) mykiss	Lower Columbia River DPS	\$5,500	\$0	\$5,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$5,500	\$0	\$5,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Puget Sound DPS	\$45,000	\$0	\$45,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$16,500	\$0	\$16,500
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$7,474	\$0	\$7,474
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$6,000	\$0	\$6,000
		Stickleback, unarmored threespine	Gasterosteus aculeatus williamsoni	Wherever found	\$8,000	\$0	\$8,000
		Sturgeon (Gulf subspecies), Atlantic	Acipenser oxyrinchus (=oxyrinchus) desotoi	Wherever found	\$47,300	\$0	\$47,300
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Gulf of Maine DPS	\$5,000	\$0	\$5,000
		sturgeon, green	Acipenser medirostris	Southern DPS	\$6,000	\$0	\$6,000
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$319,872	\$0	\$319,872
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$76,572	\$0	\$76,572
		Sturgeon, Shovelnose	Scaphirhynchus platyrhynchus	Wherever found	\$33,350	\$0	\$33,350
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$92,200	\$0	\$92,200
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$6,250	\$0	\$6,250

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sucker, Santa Ana	Catostomus santaanae	3 CA river basins	\$84,605	\$0	\$84,605
		Sucker, Warner	Catostomus warnerensis	Wherever found	\$3,000	\$0	\$3,000
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	Wherever found	\$7,500	\$0	\$7,500
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$185,392	\$0	\$185,392
		Trout, Gila	Oncorhynchus gilae	Wherever found	\$16,500	\$0	\$16,500
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$40,204	\$0	\$40,204
	Fishes subtotal				\$2,425,493	\$0	\$2,425,493
	Clams	Bankclimber, purple (mussel)	Elliptoideus sloatianus	Wherever found	\$1,800	\$0	\$1,800
		Bean, Choctaw	Villosa choctawensis	Wherever found	\$16,100	\$0	\$16,100
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$5,354	\$0	\$5,354
		Ebonysell, round	Fusconaia rotulata	Wherever found	\$1,800	\$0	\$1,800
		Kidneyshell, southern	Ptychobranchnus jonesi	Wherever found	\$16,100	\$0	\$16,100
		Mapleleaf, winged	Quadrula fragosa	Wherever found, except where listed as an experimental population	\$71,881	\$0	\$71,881
		Moccasinshell, Gulf	Medionidus penicillatus	Wherever found	\$1,800	\$0	\$1,800

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Moccasinshell, Ochlockonee	Medionidus simpsonianus	Wherever found	\$1,800	\$0	\$1,800
		Moccasinshell, Suwannee	Medionidus walkeri	Wherever found	\$21,700	\$0	\$21,700
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$3,000	\$0	\$3,000
		Pearlshell, Louisiana	Margaritifera hembeli	Wherever found	\$1,500	\$0	\$1,500
		Pigtoe, fuzzy	Pleurobema strodeanum	Wherever found	\$16,100	\$0	\$16,100
		Pigtoe, narrow	Fusconaia escambia	Wherever found	\$1,800	\$0	\$1,800
		Pigtoe, oval	Pleurobema pyriforme	Wherever found	\$21,700	\$0	\$21,700
		Pigtoe, tapered	Fusconaia burkei	Wherever found	\$16,100	\$0	\$16,100
		Pocketbook, shinyrayed	Lampsilis subangulata	Wherever found	\$16,100	\$0	\$16,100
		Purple cat's paw (=purple cat's paw pearlymussel)	Epioblasma obliquata	AL; Free- Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	\$3,000	\$0	\$3,000
		sandshell, Southern	Hamiota australis	Wherever found	\$1,800	\$0	\$1,800
		Slabshell, Chipola	Elliptio chipolaensis	Wherever found	\$1,800	\$0	\$1,800
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$75,185	\$0	\$75,185
		Threeridge, fat (mussel)	Amblema neislerii	Wherever found	\$1,800	\$0	\$1,800

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$62,471	\$0	\$62,471
	Clams subtotal				\$360,691	\$0	\$360,691
	Snails	Snail, Iowa Pleistocene	Discus macclintocki	Wherever found	\$17,200	\$0	\$17,200
	Snails subtotal				\$17,200	\$0	\$17,200
	Insects	Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$5,000	\$0	\$5,000
		Butterfly, Oregon silverspot	Speyeria zerene hippolyta	Wherever found	\$23,964	\$0	\$23,964
		Butterfly, Quino checkerspot	Euphydryas editha quino (=E. e. wrighti)	Wherever found	\$5,610	\$0	\$5,610
		Damselfly, orangeblack Hawaiian	Megalagrion xanthomelas	Wherever found	\$31,766	\$0	\$31,766
		Dragonfly, Hine's emerald	Somatochlora hineana	Wherever found	\$73,221	\$0	\$73,221
		Skipper, Dakota	Hesperia dacotae	Wherever found	\$22,780	\$0	\$22,780
		Tiger beetle, Northeastern beach	Cicindela dorsalis dorsalis	Wherever found	\$8,945	\$0	\$8,945
		Tiger beetle, Puritan	Cicindela puritana	Wherever found	\$8,015	\$0	\$8,015
	Insects subtotal				\$179,301	\$0	\$179,301
	Arachnids	Spider, spruce- fir moss	Microhexura montivaga	Wherever found	\$1,500	\$0	\$1,500
	Arachnids subtotal				\$1,500	\$0	\$1,500
	Crustaceans	Crayfish, cave	Cambarus aculabrum	Wherever found	\$6,000	\$0	\$6,000
	Crustaceans subtotal				\$6,000	\$0	\$6,000
	Flowering Plants	Barberry, island	Berberis pinnata ssp. insularis	Wherever found	\$3,878	\$0	\$3,878

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Bear-poppy, Dwarf	<i>Arctomecon humilis</i>	Wherever found	\$2,250	\$0	\$2,250
		Bedstraw, island	<i>Galium buxifolium</i>	Wherever found	\$8,059	\$0	\$8,059
		Bladderpod, kodachrome	<i>Lesquerella tumulosa</i>	Wherever found	\$2,250	\$0	\$2,250
		Bush-mallow, Santa Cruz Island	<i>Malacothamnus fasciculatus</i> var. <i>nesioticus</i>	Wherever found	\$11,985	\$0	\$11,985
		Buttercup, autumn	<i>Ranunculus aestivalis</i> (= <i>acriformis</i>)	Wherever found	\$2,250	\$0	\$2,250
		Cactus, Arizona hedgehog	<i>Echinocereus triglochidiatus</i> var. <i>arizonicus</i>	Wherever found	\$8,902	\$0	\$8,902
		cactus, Pariette	<i>Sclerocactus brevispinus</i>	Wherever found	\$2,250	\$0	\$2,250
		Cactus, Pima pineapple	<i>Coryphantha scheeri</i> var. <i>robustispina</i>	Wherever found	\$4,451	\$0	\$4,451
		Cactus, Siler pincushion	<i>Pediocactus</i> (= <i>Echinocactus</i> , = <i>Utahia</i>) <i>sileri</i>	Wherever found	\$2,250	\$0	\$2,250
		cactus, Uinta Basin hookless	<i>Sclerocactus wetlandicus</i>	Wherever found	\$2,250	\$0	\$2,250
		Cycladenia, Jones	<i>Cycladenia humilis</i> var. <i>jonesii</i>	Wherever found	\$2,250	\$0	\$2,250
		Gilia, Hoffmann's slender-flowered	<i>Gilia tenuiflora</i> ssp. <i>hoffmannii</i>	Wherever found	\$8,530	\$0	\$8,530
		Ladies'-tresses, Ute	<i>Spiranthes diluvialis</i>	Wherever found	\$2,250	\$0	\$2,250
		Malacothrix, Santa Cruz Island	<i>Malacothrix indecora</i>	Wherever found	\$3,927	\$0	\$3,927
		mallow, Gierisch	<i>Sphaeralcea gierischii</i>	Wherever found	\$2,250	\$0	\$2,250

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Manzanita, Santa Rosa Island	Arctostaphylos confertiflora	Wherever found	\$8,059	\$0	\$8,059
		Milk-vetch, heliotrope	Astragalus montii	Wherever found	\$2,250	\$0	\$2,250
		milkvetch, Deseret	Astragalus desereticus	Wherever found	\$2,250	\$0	\$2,250
		Paintbrush, soft-leaved	Castilleja mollis	Wherever found	\$10,159	\$0	\$10,159
		Phacelia, island	Phacelia insularis ssp. insularis	Wherever found	\$11,985	\$0	\$11,985
		Primrose, Maguire	Primula maguirei	Wherever found	\$2,250	\$0	\$2,250
		Reed-mustard, Barneby	Schoenocrambe barnebyi	Wherever found	\$2,250	\$0	\$2,250
		Reed-mustard, clay	Schoenocrambe argillacea	Wherever found	\$2,250	\$0	\$2,250
		Reed-mustard, shrubby	Schoenocrambe suffrutescens	Wherever found	\$2,250	\$0	\$2,250
		Ridge-cress, Barneby	Lepidium barnebyanum	Wherever found	\$2,250	\$0	\$2,250
		Rock-cress, Hoffmann's	Arabis hoffmannii	Wherever found	\$11,985	\$0	\$11,985
		Rush-rose, island	Helianthemum greenei	Wherever found	\$8,059	\$0	\$8,059
		Thistle, Pitcher's	Cirsium pitcheri	Wherever found	\$23,985	\$0	\$23,985
		`Ahinahina	Argyroxiphium sandwicense ssp. macrocephalum	Wherever found	\$11,320	\$0	\$11,320
	Flowering Plants subtotal				\$171,284	\$0	\$171,284
	Ferns and Allies	Fern, Aleutian shield	Polystichum aleuticum	Wherever found	\$10,000	\$0	\$10,000
	Ferns and Allies subtotal				\$10,000	\$0	\$10,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic subtotal					\$17,122,605	\$0	\$17,122,605
Foreign	Mammals	Manatee, West African	Trichechus senegalensis	Wherever found	\$9,000	\$0	\$9,000
		Pronghorn, peninsular	Antilocapra americana peninsularis	Mexico (Baja California)	\$3,000	\$0	\$3,000
		Tiger	Panthera tigris	Wherever found	\$1,000	\$0	\$1,000
	Mammals subtotal				\$13,000	\$0	\$13,000
	Birds	Crane, black-necked	Grus nigricollis	Wherever found	\$40	\$0	\$40
		Crane, hooded	Grus monacha	Wherever found	\$40	\$0	\$40
		Crane, Japanese	Grus japonensis	Wherever found	\$20	\$0	\$20
		Crane, white-naped	Grus vipio	Wherever found	\$40	\$0	\$40
	Birds subtotal				\$140	\$0	\$140
	Reptiles	Turtle, Central American river	Dermatemys mawii	Wherever found	\$32,902	\$0	\$32,902
	Reptiles subtotal				\$32,902	\$0	\$32,902
Foreign subtotal					\$46,042	\$0	\$46,042
Domestic	Multi-Species				\$22,809	\$0	\$22,809
	Multi-Species subtotal				\$22,809	\$0	\$22,809
Domestic subtotal					\$22,809	\$0	\$22,809
Total					\$17,191,456	\$0	\$17,191,456

DOT Federal Aviation Administration - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Indiana	Myotis sodalis	Wherever found	\$26,500	\$0	\$26,500
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$23,000	\$0	\$23,000
		Bear, polar	Ursus maritimus	Wherever found	\$2,500	\$0	\$2,500
		Prairie dog, Utah	Cynomys parvidens	Wherever found	\$1,350,000	\$0	\$1,350,000
	Mammals subtotal				\$1,402,000	\$0	\$1,402,000
	Birds	Eider, spectacled	Somateria fischeri	Wherever found	\$1,500	\$0	\$1,500
		Eider, Steller's	Polysticta stelleri	AK breeding pop.	\$1,500	\$0	\$1,500
		Horned lark, streaked	Eremophila alpestris strigata	Wherever found	\$11,000	\$0	\$11,000
		Knot, red	Calidris canutus rufa	Wherever found	\$3,000	\$0	\$3,000
	Birds subtotal				\$17,000	\$0	\$17,000
	Reptiles	Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$1,500	\$0	\$1,500
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$29,500	\$0	\$29,500
	Reptiles subtotal				\$31,000	\$0	\$31,000
	Fishes	Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Willamette River ESU	\$15,000	\$0	\$15,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$15,000	\$0	\$15,000
	Fishes subtotal				\$30,000	\$0	\$30,000
	Insects	Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$22,500	\$0	\$22,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Insects subtotal				\$22,500	\$0	\$22,500
	Multi-Species				\$17,500	\$0	\$17,500
	Multi-Species subtotal				\$17,500	\$0	\$17,500
Domestic subtotal					\$1,520,000	\$0	\$1,520,000
Total					\$1,520,000	\$0	\$1,520,000

Environmental Protection Agency - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$190	\$0	\$190
		Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$720	\$0	\$720
		Bat, Indiana	Myotis sodalis	Wherever found	\$5,762	\$0	\$5,762
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$9,092	\$0	\$9,092
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population or delisted	\$1,000	\$0	\$1,000
		Ferret, black- footed	Mustela nigripes	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$500	\$0	\$500
		Ferret, black- footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$500	\$0	\$500
		Fox, San Joaquin kit	Vulpes macrotis mutica	wherever found	\$720	\$0	\$720
		Jaguar	Panthera onca	Wherever found	\$1,000	\$0	\$1,000
		Jaguarundi, Gulf Coast	Herpailurus (=Felis) yagouarundi cacomitli	Wherever found	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$1,010	\$0	\$1,010
		Manatee, West Indian	Trichechus manatus	Wherever found	\$291	\$0	\$291
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	wherever found	\$1,000	\$0	\$1,000
		Ocelot	Leopardus (=Felis) pardalis	wherever found	\$1,000	\$0	\$1,000
		Seal, Guadalupe fur	Arctocephalus townsendi	Wherever found	\$300	\$0	\$300
		Whale, blue	Balaenoptera musculus	Wherever found	\$1,207	\$0	\$1,207
		Whale, finback	Balaenoptera physalus	Wherever found	\$1,646	\$0	\$1,646
		Whale, gray	Eschrichtius robustus	Western North Pacific DPS	\$720	\$0	\$720
		Whale, humpback	Megaptera novaeangliae	Central America DPS	\$664	\$0	\$664
		Whale, humpback	Megaptera novaeangliae	Western North Pacific	\$720	\$0	\$720
		Whale, North Atlantic Right	Eubalaena glacialis	Wherever found	\$602	\$0	\$602
		Whale, Sei	Balaenoptera borealis	Wherever found	\$1,254	\$0	\$1,254
		Whale, sperm	Physeter catodon (=macrocephalus)	Wherever found	\$1,167	\$0	\$1,167
		Wolf, gray	Canis lupus	MN	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	Canis lupus	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$200	\$0	\$200
		Wolf, gray	Canis lupus	WY, EXPN population	\$200	\$0	\$200
		Wolf, Mexican	Canis lupus baileyi	U.S.A. (portions of AZ and NM)see 17.84(k)	\$200	\$0	\$200
		Wolf, Mexican	Canis lupus baileyi	Wherever found, except where listed as an experimental population	\$200	\$0	\$200
	Mammals subtotal				\$33,065	\$0	\$33,065
	Birds	Coot, Hawaiian	Fulica americana alai	Wherever found	\$720	\$0	\$720

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Crane, whooping	<i>Grus americana</i>	U.S.A (Southwestern Louisiana)	\$250	\$0	\$250
		Crane, whooping	<i>Grus americana</i>	U.S.A. (AL, AR, CO, FL, GA, ID, IL, IN, IA, KY, LA, MI, MN, MS, MO, NC, NM, OH, SC, TN, UT, VA, WI, WV, western half of WY)	\$250	\$0	\$250
		Crane, whooping	<i>Grus americana</i>	U.S.A. (CO, ID, FL, NM, UT, and the western half of Wyoming)	\$250	\$0	\$250
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$250	\$0	\$250
		Curlew, Eskimo	<i>Numenius borealis</i>	Wherever found	\$1,000	\$0	\$1,000
		Duck, Hawaiian (=koloa)	<i>Anas wyvilliana</i>	Wherever found	\$720	\$0	\$720
		Falcon, Northern Aplomado	<i>Falco femoralis septentrionalis</i>	U.S.A (AZ, NM)	\$500	\$0	\$500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Falcon, Northern Aplomado	Falco femoralis septentrionalis	Wherever found, except where listed as an experimental population	\$500	\$0	\$500
		Flycatcher, southwestern willow	Empidonax traillii extimus	Wherever found	\$100	\$0	\$100
		gallinule, Hawaiian common	Gallinula galeata sandvicensis	Wherever found	\$720	\$0	\$720
		Gnatcatcher, coastal California	Polioptila californica californica	Wherever found	\$900	\$0	\$900
		Goose, Hawaiian	Branta (=Nesochen) sandvicensis	Wherever found	\$720	\$0	\$720
		Knot, red	Calidris canutus rufa	Wherever found	\$1,030	\$0	\$1,030
		Owl, Mexican spotted	Strix occidentalis lucida	Wherever found	\$1,000	\$0	\$1,000
		petrel, Hawaiian	Pterodroma sandwichensis	Wherever found	\$720	\$0	\$720
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$1,803	\$0	\$1,803

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, piping	Charadrius melodus	[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$2,280	\$0	\$2,280
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$100	\$0	\$100
		Shearwater, Newell's Townsend's	Puffinus auricularis newelli	Wherever found	\$720	\$0	\$720
		Stilt, Hawaiian	Himantopus mexicanus knudseni	Wherever found	\$720	\$0	\$720
		Tern, California least	Sterna antillarum browni	Wherever found	\$100	\$0	\$100
		Tern, least	Sterna antillarum	interior pop.	\$950	\$0	\$950
		Tern, roseate	Sterna dougallii dougallii	northeast U.S. nesting pop.	\$166	\$0	\$166
		Vireo, Black-capped	Vireo atricapilla	Wherever found	\$1,000	\$0	\$1,000
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$900	\$0	\$900
		Warbler (=wood), golden-cheeked	Dendroica chrysoparia	Wherever found	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Warbler, Kirtland's	Setophaga kirtlandii (= Dendroica kirtlandii)	Wherever found	\$235	\$0	\$235
	Birds subtotal				\$19,604	\$0	\$19,604
	Reptiles	Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$557	\$0	\$557
		Rattlesnake, New Mexican ridge-nosed	Crotalus willardi obscurus	Wherever found	\$1,000	\$0	\$1,000
		Sea turtle, green	Chelonia mydas	Central North Pacific DPS	\$720	\$0	\$720
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$3,223	\$0	\$3,223
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$3,454	\$0	\$3,454
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$529	\$0	\$529
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$3,984	\$0	\$3,984
		Sea turtle, loggerhead	Caretta caretta	North Pacific Ocean DPS	\$720	\$0	\$720
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$3,264	\$0	\$3,264
		Sea turtle, olive ridley	Lepidochelys olivacea	Wherever found, except when listed as endangered under 50 CFR 224.101	\$720	\$0	\$720

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Snake, copperbelly water	Nerodia erythrogaster neglecta	Indiana north of 40 degrees north latitude, Michigan, Ohio	\$352	\$0	\$352
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$3,720	\$0	\$3,720
		turtle, bog	Clemmys muhlenbergii	Wherever found, except GA, NC, SC, TN, VA	\$249	\$0	\$249
	Reptiles subtotal				\$22,492	\$0	\$22,492
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$720	\$0	\$720
		Frog, Chiricahua leopard	Rana chiricahuensis	Wherever found	\$1,000	\$0	\$1,000
	Amphibians subtotal				\$1,720	\$0	\$1,720
	Fishes	Madtom, Scioto	Noturus trautmani	Wherever found	\$147	\$0	\$147
		Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$3,258	\$0	\$3,258
		Shark, Scalloped Hammerhead	Sphyrna lewini	Indo-West Pacific DPS	\$100	\$0	\$100
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$500	\$0	\$500
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$720	\$0	\$720

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$840	\$0	\$840
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$100	\$0	\$100
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Gulf of Maine DPS	\$7,251	\$0	\$7,251
		sturgeon, green	Acipenser medirostris	Southern DPS	\$100	\$0	\$100
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$950	\$0	\$950
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$7,743	\$0	\$7,743
		Sucker, Santa Ana	Catostomus santaanae	3 CA river basins	\$180	\$0	\$180
	Fishes subtotal				\$21,889	\$0	\$21,889
	Clams	Bean, rayed	Villosa fabalis	Wherever found	\$4,485	\$0	\$4,485
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$6,153	\$0	\$6,153
		Fanshell	Cyprogenia stegaria	Wherever found	\$3,078	\$0	\$3,078
		Higgins eye (pearlymussel)	Lampsilis higginsii	Wherever found	\$1,466	\$0	\$1,466
		Mapleleaf, winged	Quadrula fragosa	Wherever found, except where listed as an experimental population	\$176	\$0	\$176
		Mucket, pink (pearlymussel)	Lampsilis abrupta	Wherever found	\$3,078	\$0	\$3,078

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Mussel, sheepnose	Plethobasus cyphus	Wherever found	\$7,514	\$0	\$7,514
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$8,335	\$0	\$8,335
		Purple Cat's paw (=Purple Cat's paw pearlymussel)	Epioblasma obliquata obliquata	Wherever found; Except where listed as Experimental Populations	\$1,173	\$0	\$1,173
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$2,726	\$0	\$2,726
		Riffleshell, northern	Epioblasma torulosa rangiana	Wherever found	\$5,140	\$0	\$5,140
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$176	\$0	\$176
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$7,742	\$0	\$7,742
	Clams subtotal				\$51,242	\$0	\$51,242
	Snails	Abalone, Black	Haliotis cracherodii	Wherever found	\$1,200	\$0	\$1,200
		Abalone, White	Haliotis sorenseni	North America (West Coast from Point Conception, CA, U.S.A., to Punta Abreojos, Baja California, Mexico)	\$1,200	\$0	\$1,200
	Snails subtotal				\$2,400	\$0	\$2,400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$1,580	\$0	\$1,580
		Dragonfly, Hine's emerald	Somatochlora hineana	Wherever found	\$2,354	\$0	\$2,354
	Insects subtotal				\$3,934	\$0	\$3,934
	Crustaceans	Fairy shrimp, Riverside	Streptocephalus woottoni	Wherever found	\$100	\$0	\$100
	Crustaceans subtotal				\$100	\$0	\$100
	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$2,736	\$0	\$2,736
		Coral, elkhorn	Acropora palmata	Wherever found	\$3,254	\$0	\$3,254
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$2,735	\$0	\$2,735
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$2,736	\$0	\$2,736
		Coral, Pillar	Dendrogyra cylindrus	Wherever found	\$2,735	\$0	\$2,735
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$2,737	\$0	\$2,737
		Coral, staghorn	Acropora cervicornis	Wherever found	\$3,254	\$0	\$3,254
		No common name	Acropora globiceps	Wherever found	\$100	\$0	\$100
		No common name	Acropora jacquelineae	Wherever found	\$100	\$0	\$100
		No common name	Acropora retusa	Wherever found	\$100	\$0	\$100
		No common name	Acropora speciosa	Wherever found	\$100	\$0	\$100
		No common name	Euphyllia paradivisa	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		No common name	<i>Isopora crateriformis</i>	Wherever found	\$100	\$0	\$100
	Corals subtotal				\$20,787	\$0	\$20,787
	Flowering Plants	Milkweed, Mead's	<i>Asclepias meadii</i>	Wherever found	\$293	\$0	\$293
		Orchid, eastern prairie fringed	<i>Platanthera leucophaea</i>	Wherever found	\$352	\$0	\$352
		Pink, swamp	<i>Helonias bullata</i>	Wherever found	\$1,175	\$0	\$1,175
		roseroot, Leedy's	<i>Rhodiola integrifolia</i> ssp. <i>leedyi</i>	Wherever found	\$156	\$0	\$156
		Spiraea, Virginia	<i>Spiraea virginiana</i>	Wherever found	\$147	\$0	\$147
		Woolly-star, Santa Ana River	<i>Eriastrum densifolium</i> ssp. <i>sanctorum</i>	Wherever found	\$100	\$0	\$100
	Flowering Plants subtotal				\$2,223	\$0	\$2,223
Domestic subtotal					\$179,456	\$0	\$179,456
Foreign	Mammals	Dugong	<i>Dugong dugon</i>	Wherever found	\$100	\$0	\$100
	Mammals subtotal				\$100	\$0	\$100
Foreign subtotal					\$100	\$0	\$100
Domestic	Multi-Species				\$4,289,552	\$0	\$4,289,552
	Multi-Species subtotal				\$4,289,552	\$0	\$4,289,552
Domestic subtotal					\$4,289,552	\$0	\$4,289,552
Total					\$4,469,108	\$0	\$4,469,108

Federal Communications Commission - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Multi-Species				\$155,000	\$0	\$155,000
	Multi-Species subtotal				\$155,000	\$0	\$155,000
Domestic subtotal					\$155,000	\$0	\$155,000
Total					\$155,000	\$0	\$155,000

Federal Energy Regulatory Commission - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$1,000	\$0	\$1,000
		Bat, Indiana	Myotis sodalis	Wherever found	\$19,000	\$0	\$19,000
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$30,000	\$0	\$30,000
		Bat, Virginia big-eared	Corynorhinus (=Plecotus) townsendii virginianus	Wherever found	\$300	\$0	\$300
		Bear, polar	Ursus maritimus	Wherever found	\$1,000	\$0	\$1,000
		Bison, wood	Bison bison athabascae	Wherever found	\$200	\$0	\$200
		Jaguarundi, Gulf Coast	Herpailurus (=Felis) yagouaroundi cacomitli	Wherever found	\$1,000	\$0	\$1,000
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$1,000	\$0	\$1,000
		Manatee, West Indian	Trichechus manatus	Wherever found	\$2,000	\$0	\$2,000
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	wherever found	\$1,000	\$0	\$1,000
		Ocelot	Leopardus (=Felis) pardalis	wherever found	\$1,000	\$0	\$1,000
		Otter, Northern Sea	Enhydra lutris kenyoni	Southwest Alaska DPS	\$1,000	\$0	\$1,000
		Sea lion, Steller	Eumetopias jubatus	Western DPS	\$2,000	\$0	\$2,000
		Seal, Ringed	Phoca (=Pusa) hispida botnica	Wherever found	\$300	\$0	\$300
		Seal, spotted	Phoca largha	Southern DPS	\$1,000	\$0	\$1,000
		whale, beluga	Delphinapterus leucas	Cook Inlet DPS	\$1,000	\$0	\$1,000
		Whale, blue	Balaenoptera musculus	Wherever found	\$1,000	\$0	\$1,000
		Whale, bowhead	Balaena mysticetus	Wherever found	\$1,000	\$0	\$1,000
		Whale, finback	Balaenoptera physalus	Wherever found	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Whale, gray	<i>Eschrichtius robustus</i>	Western North Pacific DPS	\$2,000	\$0	\$2,000
		Whale, humpback	<i>Megaptera novaeangliae</i>	Western North Pacific	\$3,000	\$0	\$3,000
		Whale, killer	<i>Orcinus orca</i>	Southern Resident DPS	\$2,000	\$0	\$2,000
		Whale, North Atlantic Right	<i>Eubalaena glacialis</i>	Wherever found	\$3,000	\$0	\$3,000
		Whale, North Pacific Right	<i>Eubalaena japonica</i>	Wherever found	\$1,000	\$0	\$1,000
		Whale, Sei	<i>Balaenoptera borealis</i>	Wherever found	\$1,000	\$0	\$1,000
		Whale, sperm	<i>Physeter catodon</i> (= <i>macrocephalus</i>)	Wherever found	\$1,000	\$0	\$1,000
		Wolf, gray	<i>Canis lupus</i>	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$8,000	\$0	\$8,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Mammals subtotal				\$86,800	\$0	\$86,800
	Birds	Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$3,000	\$0	\$3,000
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$1,000	\$0	\$1,000
		Falcon, Northern Aplomado	<i>Falco femoralis septentrionalis</i>	Wherever found, except where listed as an experimental population	\$1,000	\$0	\$1,000
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$3,000	\$0	\$3,000
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$1,000	\$0	\$1,000
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$1,000	\$0	\$1,000
		Plover, piping	<i>Charadrius melodus</i>	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$11,000	\$0	\$11,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$200	\$0	\$200
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$3,000	\$0	\$3,000
		Tern, least	Sterna antillarum	interior pop.	\$11,000	\$0	\$11,000
		Vireo, Black-capped	Vireo atricapilla	Wherever found	\$2,000	\$0	\$2,000
		Warbler, Kirtland's	Setophaga kirtlandii (= Dendroica kirtlandii)	Wherever found	\$1,000	\$0	\$1,000
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$1,000	\$0	\$1,000
	Birds subtotal				\$39,200	\$0	\$39,200
	Reptiles	Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$1,000	\$0	\$1,000
		Sea turtle, green	Chelonia mydas	Central North Pacific DPS	\$300	\$0	\$300
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$500	\$0	\$500
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$1,000	\$0	\$1,000
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$1,000	\$0	\$1,000
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, olive ridley	<i>Lepidochelys olivacea</i>	Wherever found, except when listed as endangered under 50 CFR 224.101	\$200	\$0	\$200
		Snake, eastern indigo	<i>Drymarchon corais couperi</i>	Wherever found	\$300	\$0	\$300
		Tortoise, gopher	<i>Gopherus polyphemus</i>	West of Mobile and Tombigbee Rivers	\$2,000	\$0	\$2,000
		turtle, bog	<i>Clemmys muhlenbergii</i>	Wherever found, except GA, NC, SC, TN, VA	\$3,000	\$0	\$3,000
	Reptiles subtotal				\$10,300	\$0	\$10,300
	Amphibians	Frog, California red-legged	<i>Rana draytonii</i>	Wherever found	\$4,000	\$0	\$4,000
		Frog, Oregon spotted	<i>Rana pretiosa</i>	Wherever found	\$22,000	\$0	\$22,000
		Frog, Sierra Nevada Yellow-legged	<i>Rana sierrae</i>	Wherever found	\$1,000	\$0	\$1,000
		Salamander, Cheat Mountain	<i>Plethodon nettingi</i>	Wherever found	\$200	\$0	\$200
		Salamander, frosted flatwoods	<i>Ambystoma cingulatum</i>	Wherever found	\$200	\$0	\$200
		Toad, arroyo (=arroyo southwestern)	<i>Anaxyrus californicus</i>	Wherever found	\$200	\$0	\$200
	Amphibians subtotal				\$27,600	\$0	\$27,600
	Fishes	Cavefish, Ozark	<i>Amblyopsis rosae</i>	Wherever found	\$400	\$0	\$400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Logperch, Roanoke	Percina rex	Wherever found	\$400	\$0	\$400
		Madtom, Neosho	Noturus placidus	Wherever found	\$400	\$0	\$400
		Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$21,000	\$0	\$21,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$19,000	\$0	\$19,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$2,000	\$0	\$2,000
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$1,000	\$0	\$1,000
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$1,000	\$0	\$1,000
		Shiner, Arkansas River	Notropis girardi	Arkansas River Basin (AR, KS, NM, OK, TX)	\$2,000	\$0	\$2,000
		Smelt, delta	Hypomesus transpacificus	Wherever found	\$1,000	\$0	\$1,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$11,000	\$0	\$11,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Carolina DPS	\$9,000	\$0	\$9,000
		sturgeon, green	Acipenser medirostris	Southern DPS	\$300	\$0	\$300
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$1,000	\$0	\$1,000
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$3,000	\$0	\$3,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$300	\$0	\$300
		Sucker, Lost River	Deltistes luxatus	Wherever found	\$1,000	\$0	\$1,000
		Sucker, shortnose	Chasmistes brevirostris	Wherever found	\$1,000	\$0	\$1,000
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$14,000	\$0	\$14,000
		trout, Greenback Cutthroat	Oncorhynchus clarkii stomias	Wherever found	\$400	\$0	\$400
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$1,000	\$0	\$1,000
	Fishes subtotal				\$90,200	\$0	\$90,200
	Clams	Bean, rayed	Villosa fabalis	Wherever found	\$500	\$0	\$500
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$100	\$0	\$100
		Clubshell, black	Pleurobema curtum	Wherever found	\$300	\$0	\$300
		Elktoe, Appalachian	Alasmidonta raveneliana	Wherever found	\$300	\$0	\$300
		Fanshell	Cyprogenia stegaria	U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$1,000	\$0	\$1,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Heelsplitter, Alabama (=inflated)	Potamilus inflatus	Wherever found	\$5,000	\$0	\$5,000
		Mucket, pink (pearlymussel)	Lampsilis abrupta	Wherever found	\$1,000	\$0	\$1,000
		Mussel, sheepnose	Plethobasus cyphus	Wherever found	\$1,000	\$0	\$1,000
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$2,000	\$0	\$2,000
		Pocketbook, fat	Potamilus capax	Wherever found	\$500	\$0	\$500
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$200	\$0	\$200
		Riffleshell, northern	Epioblasma torulosa rangiana	Wherever found	\$300	\$0	\$300
		Threeridge, fat (mussel)	Amblema neislerii	Wherever found	\$200	\$0	\$200
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$1,000	\$0	\$1,000
	Clams subtotal				\$13,400	\$0	\$13,400
	Snails	Snail, Bliss Rapids	Taylorconcha serpenticola	Wherever found	\$1,000	\$0	\$1,000
		Snail, Snake River physa	Physa natricina	Wherever found	\$1,000	\$0	\$1,000
		Snail, tulotoma	Tulotoma magnifica	Wherever found	\$1,000	\$0	\$1,000
	Snails subtotal				\$3,000	\$0	\$3,000
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$2,000	\$0	\$2,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Beetle, valley elderberry longhorn	Desmocerus californicus dimorphus	Wherever found	\$500	\$0	\$500
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$1,000	\$0	\$1,000
		Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$200	\$0	\$200
		Dragonfly, Hine's emerald	Somatochlora hineana	Wherever found	\$1,000	\$0	\$1,000
	Insects subtotal				\$4,700	\$0	\$4,700
	Crustaceans	Crayfish, Nashville	Orconectes shoupi	Wherever found	\$400	\$0	\$400
	Crustaceans subtotal				\$400	\$0	\$400
	Corals	coral, Boulder star	Orbicella franksi	Wherever found	\$300	\$0	\$300
		Coral, elkhorn	Acropora palmata	Wherever found	\$200	\$0	\$200
		Coral, Lobed Star	Orbicella annularis	Wherever found	\$300	\$0	\$300
		Coral, Mountainous Star	Orbicella faveolata	Wherever found	\$300	\$0	\$300
		Coral, Pillar	Dendrogyra cylindrus	Wherever found	\$200	\$0	\$200
		Coral, Rough Cactus	Mycetophyllia ferox	Wherever found	\$200	\$0	\$200
		Coral, staghorn	Acropora cervicornis	Wherever found	\$200	\$0	\$200
	Corals subtotal				\$1,700	\$0	\$1,700
	Flowering Plants	Ambrosia, south Texas	Ambrosia cheiranthifolia	Wherever found	\$200	\$0	\$200
		Bulrush, Northeastern	Scirpus ancistrochaetus	Wherever found	\$2,000	\$0	\$2,000
		Clover, running buffalo	Trifolium stoloniferum	Wherever found	\$200	\$0	\$200
		Harperella	Ptilimnium nodosum	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Heartleaf, dwarf-flowered	Hexastylis naniflora	Wherever found	\$200	\$0	\$200
		Iris, dwarf lake	Iris lacustris	Wherever found	\$500	\$0	\$500
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$300	\$0	\$300
		Orchid, eastern prairie fringed	Platanthera leucophaea	Wherever found	\$1,000	\$0	\$1,000
		Pogonia, small whorled	Isotria medeoloides	Wherever found	\$300	\$0	\$300
		Rockcress, Georgia	Arabis georgiana	Wherever found	\$1,000	\$0	\$1,000
		Spiraea, Virginia	Spiraea virginiana	Wherever found	\$1,000	\$0	\$1,000
		Sunflower, Schweinitz's	Helianthus schweinitzii	Wherever found	\$200	\$0	\$200
	Flowering Plants subtotal				\$7,100	\$0	\$7,100
	Ferns and Allies	Quillwort, black spored	Isoetes melanospora	Wherever found	\$200	\$0	\$200
	Ferns and Allies subtotal				\$200	\$0	\$200
Domestic subtotal					\$284,600	\$0	\$284,600
Total					\$284,600	\$0	\$284,600

Nuclear Regulatory Commission - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$3,500	\$0	\$3,500
		Bat, Indiana	Myotis sodalis	Wherever found	\$3,500	\$0	\$3,500
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$9,400	\$0	\$9,400
		Bat, Virginia big-eared	Corynorhinus (=Plecotus) townsendii virginianus	Wherever found	\$1,500	\$0	\$1,500
		Ferret, black- footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$200	\$0	\$200
		Manatee, West Indian	Trichechus manatus	Wherever found	\$1,500	\$0	\$1,500
		Ocelot	Leopardus (=Felis) pardalis	wherever found	\$100	\$0	\$100
		Panther, Florida	Puma (=Felis) concolor coryi	Wherever found	\$200	\$0	\$200
		Squirrel, Carolina northern flying	Glaucomys sabrinus coloratus	Wherever found	\$1,100	\$0	\$1,100
		Whale, finback	Balaenoptera physalus	Wherever found	\$300	\$0	\$300
		Whale, humpback	Megaptera novaeangliae	Mexico DPS	\$300	\$0	\$300
		Whale, North Atlantic Right	Eubalaena glacialis	Wherever found	\$300	\$0	\$300
	Mammals subtotal				\$21,900	\$0	\$21,900

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Birds	Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$1,300	\$0	\$1,300
		Kite, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	Wherever found	\$200	\$0	\$200
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$2,300	\$0	\$2,300
		Plover, piping	<i>Charadrius melodus</i>	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$2,000	\$0	\$2,000
		Stork, wood	<i>Mycteria americana</i>	AL, FL, GA, MS, NC, SC	\$600	\$0	\$600
		Tern, least	<i>Sterna antillarum</i>	interior pop.	\$300	\$0	\$300
		Tern, roseate	<i>Sterna dougallii dougallii</i>	northeast U.S. nesting pop.	\$200	\$0	\$200
		Vireo, Black-capped	<i>Vireo atricapilla</i>	Wherever found	\$200	\$0	\$200
		Warbler (=wood), golden-cheeked	<i>Dendroica chrysoparia</i>	Wherever found	\$200	\$0	\$200
		Woodpecker, red-cockaded	<i>Picoides borealis</i>	Wherever found	\$800	\$0	\$800
	Birds subtotal				\$8,100	\$0	\$8,100
	Reptiles	Crocodile, American	<i>Crocodylus acutus</i>	U.S.A. (FL)	\$16,500	\$0	\$16,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$700	\$0	\$700
		Sea turtle, green	Chelonia mydas	North Atlantic DPS	\$5,000	\$0	\$5,000
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$4,000	\$0	\$4,000
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$4,500	\$0	\$4,500
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$4,500	\$0	\$4,500
		Sea turtle, loggerhead	Caretta caretta	North Pacific Ocean DPS	\$2,500	\$0	\$2,500
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$2,500	\$0	\$2,500
		Sea turtle, olive ridley	Lepidochelys olivacea	Wherever found, except when listed as endangered under 50 CFR 224.101	\$200	\$0	\$200
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$200	\$0	\$200
	Reptiles subtotal				\$40,600	\$0	\$40,600
	Amphibians	Salamander, Austin blind	Eurycea waterlooensis	Wherever found	\$200	\$0	\$200
		Salamander, Barton Springs	Eurycea sosorum	Wherever found	\$200	\$0	\$200
		Salamander, Jollyville Plateau	Eurycea tonkawae	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Amphibians subtotal				\$600	\$0	\$600
	Fishes	Chub, slender	Erimystax cahni	Wherever found	\$1,100	\$0	\$1,100
		Chub, spotfin	Erimonax monachus	Wherever found, except where listed as an experimental population	\$400	\$0	\$400
		Dace, blackside	Phoxinus cumberlandensis	Wherever found	\$1,100	\$0	\$1,100
		dace, Laurel	Chrosomus saylori	Wherever found	\$1,100	\$0	\$1,100
		darer, bluemask	Etheostoma akatulo	Wherever found	\$1,100	\$0	\$1,100
		Darter, boulder	Etheostoma wapiti	Wherever found	\$1,100	\$0	\$1,100
		Darter, Cumberland	Etheostoma susanae	Wherever found	\$1,100	\$0	\$1,100
		Darter, duskytail	Etheostoma percnum	Wherever found	\$1,100	\$0	\$1,100
		Darter, Kentucky arrow	Etheostoma spilotum	Wherever found	\$1,100	\$0	\$1,100
		Darter, snail	Percina tanasi	Wherever found	\$1,100	\$0	\$1,100
		Madtom, chunky	Noturus crypticus	Wherever found	\$1,100	\$0	\$1,100
		Madtom, pygmy	Noturus stanauli	Wherever found	\$1,100	\$0	\$1,100
		Madtom, yellowfin	Noturus flavipinnis	Wherever found, except where listed as an experimental population	\$1,100	\$0	\$1,100
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$6,500	\$0	\$6,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$3,500	\$0	\$3,500
		Shiner, palezone	Notropis albizonatus	Wherever found	\$1,100	\$0	\$1,100
		Shiner, Topeka	Notropis topeka (=tristis)	Wherever found, except where listed as an experimental population	\$300	\$0	\$300
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$6,500	\$0	\$6,500
		Sturgeon (Gulf subspecies), Atlantic	Acipenser oxyrinchus (=oxyrinchus) desotoi	Wherever found	\$1,500	\$0	\$1,500
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Carolina DPS	\$4,000	\$0	\$4,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Chesapeake Bay DPS	\$4,100	\$0	\$4,100
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	Gulf of Maine DPS	\$4,000	\$0	\$4,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	New York Bight DPS	\$4,000	\$0	\$4,000
		Sturgeon, Atlantic	Acipenser oxyrinchus oxyrinchus	South Atlantic DPS	\$4,000	\$0	\$4,000
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$42,500	\$0	\$42,500
		Sturgeon, shortnose	Acipenser brevirostrum	Wherever found	\$8,500	\$0	\$8,500
	Fishes subtotal				\$104,100	\$0	\$104,100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Clams	Bean, Cumberland (pearlymussel)	Villosa trabalis	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Bean, purple	Villosa perpurpurea	Wherever found	\$400	\$0	\$400
		Bean, rayed	Villosa fabalis	Wherever found	\$400	\$0	\$400
		Blossom, tubercled (pearlymussel)	Epioblasma torulosa torulosa	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Blossom, turgid (pearlymussel)	Epioblasma turgidula	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Catspaw, white (pearlymussel)	Epioblasma obliquata perobliqua	Wherever found	\$400	\$0	\$400
		Clubshell	Pleurobema clava	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Combshell, Cumberlandian	Epioblasma brevidens	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Elktoe, Appalachian	Alasmidonta raveneliana	Wherever found	\$400	\$0	\$400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Elktoe, Cumberland	<i>Alasmidonta atropurpurea</i>	Wherever found	\$400	\$0	\$400
		Fanshell	<i>Cyprogenia stegaria</i>	Wherever found	\$400	\$0	\$400
		Heelsplitter, Carolina	<i>Lasmigona decorata</i>	Wherever found	\$600	\$0	\$600
		Higgins eye (pearlymussel)	<i>Lampsilis higginsii</i>	Wherever found	\$1,300	\$0	\$1,300
		Kidneyshell, fluted	<i>Ptychobranchus subtentum</i>	Wherever found	\$400	\$0	\$400
		Lampmussel, Alabama	<i>Lampsilis virescens</i>	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Lilliput, pale (pearlymussel)	<i>Toxolasma cylindrellus</i>	Wherever found	\$400	\$0	\$400
		Monkeyface, Appalachian (pearlymussel)	<i>Quadrula sparsa</i>	Wherever found	\$400	\$0	\$400
		Monkeyface, Cumberland (pearlymussel)	<i>Quadrula intermedia</i>	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Mucket, pink (pearlymussel)	<i>Lampsilis abrupta</i>	Wherever found	\$2,200	\$0	\$2,200
		Mussel, oyster	<i>Epioblasma capsaeformis</i>	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Mussel, sheepnose	Plethobasus cyphus	Wherever found	\$1,700	\$0	\$1,700
		Mussel, snuffbox	Epioblasma triquetra	Wherever found	\$2,200	\$0	\$2,200
		Pearlymussel, birdwing	Lemiox rimosus	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Pearlymussel, cracking	Hemistena lata	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Pearlymussel, dromedary	Dromus dromas	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Pearlymussel, littlewing	Pegias fabula	Wherever found	\$400	\$0	\$400
		Pearlymussel, slabside	Pleuonaia dolabelloides	Wherever found	\$400	\$0	\$400
		Pigtoe, Cumberland	Pleurobema gibberum	Wherever found	\$400	\$0	\$400
		Pigtoe, finerayed	Fusconaia cuneolus	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Pigtoe, rough	Pleurobema plenum	Wherever found	\$2,200	\$0	\$2,200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pigtoe, shiny	Fusconaia cor	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Pimpleback, orangefoot (pearlymussel)	Plethobasus cooperianus	Wherever found	\$400	\$0	\$400
		Rabbitsfoot	Quadrula cylindrica cylindrica	Wherever found	\$400	\$0	\$400
		Rabbitsfoot, rough	Quadrula cylindrica strigillata	Wherever found	\$400	\$0	\$400
		Riffleshell, tan	Epioblasma florentina walkeri (=E. walker)	Wherever found	\$400	\$0	\$400
		Ring pink (mussel)	Obovaria retusa	Wherever found	\$400	\$0	\$400
		Spectaclecase (mussel)	Cumberlandia monodonta	Wherever found	\$400	\$0	\$400
		Wartyback, white (pearlymussel)	Plethobasus cicatricosus	Wherever found	\$400	\$0	\$400
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$400	\$0	\$400
	Clams subtotal				\$23,400	\$0	\$23,400
	Snails	Riversnail, Anthony's	Athearnia anthonyi	Wherever found; Except where listed as Experimental Populations	\$400	\$0	\$400
		Snail, painted snake coiled forest	Anguispira picta	Wherever found	\$400	\$0	\$400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Snails subtotal				\$800	\$0	\$800
	Insects	Beetle, American burying	Nicrophorus americanus	Wherever found, except where listed as an experimental population	\$200	\$0	\$200
		Beetle, Kretschmarr Cave mold	Texamaurops reddelli	Wherever found	\$200	\$0	\$200
		Beetle, Tooth Cave ground	Rhadine persephone	Wherever found	\$200	\$0	\$200
		Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$1,100	\$0	\$1,100
		Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$700	\$0	\$700
	Insects subtotal				\$2,400	\$0	\$2,400
	Arachnids	Harvestman, Bee Creek Cave	Texella reddelli	Wherever found	\$200	\$0	\$200
		Harvestman, Bone Cave	Texella reyesi	Wherever found	\$200	\$0	\$200
		Pseudoscorpion, Tooth Cave	Tartarocreagris texana	Wherever found	\$200	\$0	\$200
		Spider, spruce- fir moss	Microhexura montivaga	Wherever found	\$1,100	\$0	\$1,100
		Spider, Tooth Cave	Neoleptoneta myopica	Wherever found	\$200	\$0	\$200
	Arachnids subtotal				\$1,900	\$0	\$1,900
	Flowering Plants	Avens, spreading	Geum radiatum	Wherever found	\$1,100	\$0	\$1,100
		Bulrush, Northeastern	Scirpus ancistrochaetus	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Bush-clover, prairie	<i>Lespedeza leptostachya</i>	Wherever found	\$1,500	\$0	\$1,500
		Coneflower, smooth	<i>Echinacea laevigata</i>	Wherever found	\$200	\$0	\$200
		Dropwort, Canby's	<i>Oxypolis canbyi</i>	Wherever found	\$200	\$0	\$200
		Gerardia, sandplain	<i>Agalinis acuta</i>	Wherever found	\$400	\$0	\$400
		Leather flower, Morefield's	<i>Clematis morefieldii</i>		\$1,100	\$0	\$1,100
		Loosestrife, rough-leaved	<i>Lysimachia asperulaefolia</i>	Wherever found	\$200	\$0	\$200
		Monkshood, northern wild	<i>Aconitum noveboracense</i>	Wherever found	\$700	\$0	\$700
		Orchid, eastern prairie fringed	<i>Platanthera leucophaea</i>	Wherever found	\$700	\$0	\$700
		Orchid, white fringeless	<i>Platanthera integrilabia</i>		\$1,100	\$0	\$1,100
		Pogonia, small whorled	<i>Isotria medeoloides</i>	Wherever found	\$1,300	\$0	\$1,300
		Potato-bean, Price's	<i>Apios priceana</i>		\$1,100	\$0	\$1,100
		Rosemary, Cumberland	<i>Conradina verticillata</i>	Wherever found	\$1,100	\$0	\$1,100
		Sandwort, Cumberland	<i>Arenaria cumberlandensis</i>	Wherever found	\$1,100	\$0	\$1,100
		Skullcap, large-flowered	<i>Scutellaria montana</i>	Wherever found	\$1,100	\$0	\$1,100
		Spiraea, Virginia	<i>Spiraea virginiana</i>	Wherever found	\$1,100	\$0	\$1,100
		Sumac, Michaux's	<i>Rhus michauxii</i>	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Thistle, Pitcher's	Cirsium pitcheri	Wherever found	\$700	\$0	\$700
	Flowering Plants subtotal				\$15,100	\$0	\$15,100
	Ferns and Allies	Fern, American hart's-tongue	Asplenium scolopendrium var. americanum	Wherever found	\$1,100	\$0	\$1,100
	Ferns and Allies subtotal				\$1,100	\$0	\$1,100
	Lichens	Lichen, rock gnome	Gymnoderma lineare	Wherever found	\$1,100	\$0	\$1,100
	Lichens subtotal				\$1,100	\$0	\$1,100
	Multi-Species				\$29,500	\$0	\$29,500
	Multi-Species subtotal				\$29,500	\$0	\$29,500
Domestic subtotal					\$250,600	\$0	\$250,600
Total					\$250,600	\$0	\$250,600

Smithsonian Institution - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Ferret, black- footed	<i>Mustela nigripes</i>	Wherever found, except where listed as an experimental population	\$173,100	\$0	\$173,100
		Fox, San Joaquin kit	<i>Vulpes macrotis mutica</i>	wherever found	\$70,000	\$0	\$70,000
		Fox, Santa Catalina Island	<i>Urocyon littoralis catalinae</i>	Wherever found	\$28,000	\$0	\$28,000
		Otter, Northern Sea	<i>Enhydra lutris kenyon</i>	Southwest Alaska DPS	\$28,000	\$0	\$28,000
		Shrew, Buena Vista Lake ornate	<i>Sorex ornatus relictus</i>	Wherever found	\$7,000	\$0	\$7,000
	Mammals subtotal				\$306,100	\$0	\$306,100
	Birds	Condor, California	<i>Gymnogyps californianus</i>	U.S.A. only, except where listed as an experimental population	\$14,000	\$0	\$14,000
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$22,250	\$0	\$22,250
		Crow, Mariana (=aga)	<i>Corvus kubaryi</i>	Wherever found	\$70,000	\$0	\$70,000
		petrel, Hawaiian	<i>Pterodroma sandwichensis</i>	Wherever found	\$23,800	\$0	\$23,800

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Warbler, Kirtland's	Setophaga kirtlandii (= Dendroica kirtlandii)	Wherever found	\$14,000	\$0	\$14,000
	Birds subtotal				\$144,050	\$0	\$144,050
	Amphibians	Coqui, golden	Eleutherodactylus jasperii	Wherever found	\$28,250	\$0	\$28,250
	Amphibians subtotal				\$28,250	\$0	\$28,250
Domestic subtotal					\$478,400	\$0	\$478,400
Foreign	Mammals	Ass, Asian wild	Equus hemionus	Wherever found	\$12,500	\$0	\$12,500
		Cheetah	Acinonyx jubatus	Wherever found	\$185,250	\$0	\$185,250
		Deer, Eld's brow-antlered	Cervus eldi	Wherever found	\$59,000	\$0	\$59,000
		Dog, African wild	Lycaon pictus	Wherever found	\$8,000	\$0	\$8,000
		Elephant, Asian	Elephas maximus	Wherever found	\$591,300	\$0	\$591,300
		Gazelle, dama	Gazella dama	Wherever found	\$79,250	\$0	\$79,250
		Horse, Przewalski's	Equus przewalskii	Wherever found	\$61,950	\$0	\$61,950
		Leopard, clouded	Neofelis nebulosa	Wherever found	\$69,500	\$0	\$69,500
		Oryx, scimitar-horned	Oryx dammah	Wherever found	\$189,550	\$0	\$189,550
		Panda, giant	Ailuropoda melanoleuca	Wherever found	\$478,000	\$0	\$478,000
		Tamarin, golden-rumped	Leontopithecus spp.	Wherever found	\$39,900	\$0	\$39,900
		Tiger	Panthera tigris	Wherever found	\$294,500	\$0	\$294,500
		Wolf, maned	Chrysocyon brachyurus	Wherever found	\$302,000	\$0	\$302,000
	Mammals subtotal				\$2,370,700	\$0	\$2,370,700
	Birds	Curassow, blue-billed	Crax alberti	Wherever found	\$9,100	\$0	\$9,100
	Birds subtotal				\$9,100	\$0	\$9,100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Reptiles	Alligator, Chinese	Alligator sinensis	Wherever found	\$55,750	\$0	\$55,750
		Crocodile, Cuban	Crocodylus rhombifer	Wherever found	\$55,750	\$0	\$55,750
		Crocodile, Philippine	Crocodylus novaeguineae mindorensis	Wherever found	\$55,750	\$0	\$55,750
		Gavial	Gavialis gangeticus	Wherever found	\$55,750	\$0	\$55,750
		Iguana, Fiji banded	Brachylophus fasciatus	Wherever found	\$55,750	\$0	\$55,750
		Monitor, Komodo Island	Varanus komodoensis	Wherever found	\$55,750	\$0	\$55,750
		Tartaruga	Podocnemis expansa	Wherever found	\$55,750	\$0	\$55,750
		Tortoise, Madagascar radiated	Geochelone radiata	Wherever found	\$55,750	\$0	\$55,750
		Tracaja	Podocnemis unifilis	Wherever found	\$55,750	\$0	\$55,750
	Reptiles subtotal				\$501,750	\$0	\$501,750
	Amphibians	Frog, Panamanian golden	Atelopus varius zeteki	Wherever found	\$41,750	\$0	\$41,750
		Salamander, Japanese giant	Andrias japonicus (=davidianus j.)	Wherever found	\$55,750	\$0	\$55,750
	Amphibians subtotal				\$97,500	\$0	\$97,500
Foreign subtotal					\$2,979,050	\$0	\$2,979,050
Total					\$3,457,450	\$0	\$3,457,450

Tennessee Valley Authority - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, gray	Myotis grisescens	Wherever found	\$127,000	\$0	\$127,000
		Bat, Indiana	Myotis sodalis	Wherever found	\$665,500	\$0	\$665,500
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$184,000	\$0	\$184,000
		Bat, Virginia big-eared	Corynorhinus (=Plecotus) townsendii virginianus	Wherever found	\$100,000	\$0	\$100,000
		Squirrel, Carolina northern flying	Glaucomys sabrinus coloratus	Wherever found	\$1,700	\$0	\$1,700
	Mammals subtotal				\$1,078,200	\$0	\$1,078,200
	Birds	Knot, red	Calidris canutus rufa	Wherever found	\$200	\$0	\$200
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$1,000	\$0	\$1,000
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$2,200	\$0	\$2,200
		Tern, least	Sterna antillarum	interior pop.	\$200	\$0	\$200
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$2,400	\$0	\$2,400
	Birds subtotal				\$6,000	\$0	\$6,000
	Reptiles	turtle, bog	Clemmys muhlenbergii	U.S.A. (GA, NC, SC, TN, VA)	\$500	\$0	\$500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Turtle, flattened musk	<i>Sternotherus depressus</i>	Black Warrior R. system upstream from Bankhead Dam	\$200	\$0	\$200
		Turtle, ringed map	<i>Graptemys oculifera</i>	Wherever found	\$200	\$0	\$200
	Reptiles subtotal				\$900	\$0	\$900
	Fishes	Darter, boulder	<i>Etheostoma wapiti</i>	Wherever found	\$130,500	\$0	\$130,500
		Darter, snail	<i>Percina tanasi</i>	Wherever found	\$130,500	\$0	\$130,500
	Fishes subtotal				\$261,000	\$0	\$261,000
	Clams	Monkeyface, Cumberland (pearlymussel)	<i>Quadrula intermedia</i>	Wherever found; Except where listed as Experimental Populations	\$130,500	\$0	\$130,500
		Pearlymussel, cracking	<i>Hemistena lata</i>	Wherever found; Except where listed as Experimental Populations	\$130,500	\$0	\$130,500
		Pearlymussel, slabside	<i>Pleuronaia dolabelloides</i>	Wherever found	\$130,500	\$0	\$130,500
		Pigtoe, finerayed	<i>Fusconaia cuneolus</i>	Wherever found; Except where listed as Experimental Populations	\$130,500	\$0	\$130,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pigtoe, shiny	<i>Fusconaia cor</i>	U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	\$130,500	\$0	\$130,500
	Clams subtotal				\$652,500	\$0	\$652,500
	Snails	Snail, painted snake coiled forest	<i>Anguispira picta</i>	Wherever found	\$400	\$0	\$400
	Snails subtotal				\$400	\$0	\$400
	Insects	Beetle, American burying	<i>Nicrophorus americanus</i>	Wherever found, except where listed as an experimental population	\$200	\$0	\$200
		Bumble bee, Rusty patched	<i>Bombus affinis</i>	Wherever found	\$200	\$0	\$200
		Butterfly, Mitchell's satyr	<i>Neonympha mitchellii mitchellii</i>	Wherever found	\$700	\$0	\$700
	Insects subtotal				\$1,100	\$0	\$1,100
	Arachnids	Spider, spruce- fir moss	<i>Microhexura montivaga</i>	Wherever found	\$1,000	\$0	\$1,000
	Arachnids subtotal				\$1,000	\$0	\$1,000
	Flowering Plants	Aster, Ruth's golden	<i>Pityopsis ruthii</i>	Wherever found	\$5,500	\$0	\$5,500
		Orchid, white fringeless	<i>Platanthera integrilabia</i>		\$3,200	\$0	\$3,200
		Skullcap, large-flowered	<i>Scutellaria montana</i>	Wherever found	\$25,500	\$0	\$25,500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Flowering Plants subtotal				\$34,200	\$0	\$34,200
Domestic subtotal					\$2,035,300	\$0	\$2,035,300
Total					\$2,035,300	\$0	\$2,035,300

USDA Animal and Plant Health Inspection Service - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Hawaiian hoary	Lasiurus cinereus semotus	Wherever found	\$324,300	\$0	\$324,300
		Bat, Indiana	Myotis sodalis	Wherever found	\$13,900	\$0	\$13,900
		Bat, lesser long-nosed	Leptonycteris curasoae yerbabuenae	Wherever found	\$500	\$0	\$500
		Bat, Mexican long-nosed	Leptonycteris nivalis	Wherever found	\$100	\$0	\$100
		Bat, Northern long-eared	Myotis septentrionalis	Wherever found	\$17,000	\$0	\$17,000
		Bear, grizzly	Ursus arctos horribilis	U.S.A., conterminous (lower 48) States, except where listed as an experimental population or delisted	\$152,200	\$0	\$152,200
		Bear, polar	Ursus maritimus	Wherever found	\$200	\$0	\$200
		Caribou, Woodland	Rangifer tarandus caribou	Selkirk Mountain population	\$1,400	\$0	\$1,400
		Deer, key	Odocoileus virginianus clavium	Wherever found	\$1,000,900	\$0	\$1,000,900
		Ferret, black- footed	Mustela nigripes	Wherever found, except where listed as an experimental population	\$201,700	\$0	\$201,700
		Jaguar	Panthera onca	Wherever found	\$2,300	\$0	\$2,300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Jaguarundi, Gulf Coast	Herpailurus (=Felis) yagouaroundi cacomitli	Wherever found	\$500	\$0	\$500
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$17,200	\$0	\$17,200
		Manatee, West Indian	Trichechus manatus	Wherever found	\$700	\$0	\$700
		Mouse, Alabama beach	Peromyscus polionotus ammobates	Wherever found	\$4,000	\$0	\$4,000
		Mouse, Choctawhatche beach	Peromyscus polionotus allophrys	Wherever found	\$32,500	\$0	\$32,500
		Mouse, New Mexico meadow jumping	Zapus hudsonius luteus	Wherever found	\$400	\$0	\$400
		Mouse, Perdido Key beach	Peromyscus polionotus trissyllepsis	Wherever found	\$32,500	\$0	\$32,500
		Mouse, salt marsh harvest	Reithrodontomys raviventris	wherever found	\$194,500	\$0	\$194,500
		Mouse, southeastern beach	Peromyscus polionotus niveiventris	wherever found	\$32,500	\$0	\$32,500
		Mouse, St. Andrew beach	Peromyscus polionotus peninsularis	Wherever found	\$32,500	\$0	\$32,500
		Ocelot	Leopardus (=Felis) pardalis	wherever found	\$10,700	\$0	\$10,700
		Panther, Florida	Puma (=Felis) concolor coryi	Wherever found	\$600	\$0	\$600
		Prairie dog, Utah	Cynomys parvidens	Wherever found	\$2,000	\$0	\$2,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pronghorn, Sonoran	<i>Antilocapra americana sonoriensis</i>	Wherever found, except where listed as an experimental population	\$100	\$0	\$100
		Rabbit, Columbia Basin Pygmy	<i>Brachylagus idahoensis</i>	Columbia Basin DPS	\$200	\$0	\$200
		Rabbit, Lower Keys marsh	<i>Sylvilagus palustris hefneri</i>	Wherever found	\$25,400	\$0	\$25,400
		Rice rat	<i>Oryzomys palustris natator</i>	lower FL Keys	\$400	\$0	\$400
		Seal, Hawaiian monk	<i>Monachus schauinslandi</i>	Wherever found	\$10,100	\$0	\$10,100
		Sheep, Sierra Nevada bighorn	<i>Ovis canadensis sierrae</i>	Sierra Nevada	\$700	\$0	\$700
		Squirrel, Mount Graham red	<i>Tamiasciurus hudsonicus grahamensis</i>	Wherever found	\$100	\$0	\$100
		Squirrel, Northern Idaho Ground	<i>Urocitellus brunneus</i>	Wherever found	\$4,800	\$0	\$4,800
		Vole, Hualapai Mexican	<i>Microtus mexicanus hualpaiensis</i>	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	<i>Canis lupus</i>	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$860,200	\$0	\$860,200
		Wolf, gray	<i>Canis lupus</i>	WY, EXPN population	\$221,400	\$0	\$221,400
		Wolf, Mexican	<i>Canis lupus baileyi</i>	U.S.A. (portions of AZ and NM)see 17.84(k)	\$60,000	\$0	\$60,000
		Wolf, Mexican	<i>Canis lupus baileyi</i>	Wherever found, except where listed as an experimental population	\$65,500	\$0	\$65,500
		Woodrat, Key Largo	<i>Neotoma floridana smalli</i>	Wherever found	\$400	\$0	\$400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Mammals subtotal				\$3,324,500	\$0	\$3,324,500
	Birds	Bobwhite, masked (quail)	<i>Colinus virginianus ridgwayi</i>	Wherever found	\$100	\$0	\$100
		Condor, California	<i>Gymnogyps californianus</i>	U.S.A. (specific portions of Arizona, Nevada, and Utah)	\$100	\$0	\$100
		Condor, California	<i>Gymnogyps californianus</i>	U.S.A. only, except where listed as an experimental population	\$400	\$0	\$400
		Coot, Hawaiian	<i>Fulica americana alai</i>	Wherever found	\$319,300	\$0	\$319,300
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$1,000	\$0	\$1,000
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$15,500	\$0	\$15,500
		Duck, Hawaiian (=koloa)	<i>Anas wyvilliana</i>	Wherever found	\$319,300	\$0	\$319,300
		Falcon, Northern Aplomado	<i>Falco femoralis septentrionalis</i>	Wherever found, except where listed as an experimental population	\$800	\$0	\$800

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$36,100	\$0	\$36,100
		gallinule, Hawaiian common	<i>Gallinula galeata sandvicensis</i>	Wherever found	\$319,300	\$0	\$319,300
		Gnatcatcher, coastal California	<i>Polioptila californica californica</i>	Wherever found	\$200	\$0	\$200
		Goose, Hawaiian	<i>Branta (=Nesochen) sandvicensis</i>	Wherever found	\$293,500	\$0	\$293,500
		Hawk, Hawaiian (=ʻIo)	<i>Buteo solitarius</i>	Wherever found	\$43,400	\$0	\$43,400
		Kite, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	Wherever found	\$500	\$0	\$500
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$3,000	\$0	\$3,000
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$12,400	\$0	\$12,400
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$500	\$0	\$500
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$200	\$0	\$200
		Plover, piping	<i>Charadrius melodus</i>	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$320,000	\$0	\$320,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, piping	Charadrius melodus	[Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	\$48,000	\$0	\$48,000
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$570,000	\$0	\$570,000
		Prairie-chicken, Attwater's greater	Tympanuchus cupido attwateri	Wherever found	\$10,700	\$0	\$10,700
		Rail, California clapper	Rallus longirostris obsoletus	Wherever found	\$204,400	\$0	\$204,400
		Rail, light-footed clapper	Rallus longirostris levipes	Wherever found	\$76,600	\$0	\$76,600
		Rail, Yuma clapper	Rallus longirostris yumanensis	Wherever found	\$13,000	\$0	\$13,000
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$500	\$0	\$500
		Stilt, Hawaiian	Himantopus mexicanus knudseni	Wherever found	\$319,300	\$0	\$319,300
		Swiftlet, Mariana gray	Aerodramus vanikorensis bartschi	Wherever found	\$106,800	\$0	\$106,800

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Tern, California least	<i>Sterna antillarum browni</i>	Wherever found	\$443,000	\$0	\$443,000
		Tern, least	<i>Sterna antillarum</i>	interior pop.	\$37,500	\$0	\$37,500
		Tern, roseate	<i>Sterna dougallii dougallii</i>	Western Hemisphere except NE U.S.	\$63,000	\$0	\$63,000
		Tern, roseate	<i>Sterna dougallii dougallii</i>	northeast U.S. nesting pop.	\$38,100	\$0	\$38,100
		Vireo, Black-capped	<i>Vireo atricapilla</i>	Wherever found	\$300	\$0	\$300
		Vireo, least Bell's	<i>Vireo bellii pusillus</i>	Wherever found	\$200	\$0	\$200
		Warbler (=wood), golden-cheeked	<i>Dendroica chrysoparia</i>	Wherever found	\$2,800	\$0	\$2,800
		Warbler, Kirtland's	<i>Setophaga kirtlandii</i> (= <i>Dendroica kirtlandii</i>)	Wherever found	\$51,800	\$0	\$51,800
	Birds subtotal				\$3,671,600	\$0	\$3,671,600
	Reptiles	Crocodile, American	<i>Crocodylus acutus</i>	U.S.A. (FL)	\$400	\$0	\$400
		gartersnake, narrow-headed	<i>Thamnophis rufipunctatus</i>	Wherever found	\$500	\$0	\$500
		gartersnake, northern Mexican	<i>Thamnophis eques megalops</i>	Wherever found	\$13,000	\$0	\$13,000
		Sea turtle, green	<i>Chelonia mydas</i>	Central North Pacific DPS	\$10,100	\$0	\$10,100
		Sea turtle, green	<i>Chelonia mydas</i>	Central West Pacific DPS	\$272,200	\$0	\$272,200
		Sea turtle, green	<i>Chelonia mydas</i>	North Atlantic DPS	\$36,600	\$0	\$36,600
		Sea turtle, green	<i>Chelonia mydas</i>	South Atlantic DPS	\$3,000	\$0	\$3,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sea turtle, hawksbill	Eretmochelys imbricata	Wherever found	\$2,100	\$0	\$2,100
		Sea turtle, Kemp's ridley	Lepidochelys kempii	Wherever found	\$49,600	\$0	\$49,600
		Sea turtle, leatherback	Dermochelys coriacea	Wherever found	\$36,600	\$0	\$36,600
		Sea turtle, loggerhead	Caretta caretta	Northwest Atlantic Ocean DPS	\$49,600	\$0	\$49,600
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$400	\$0	\$400
		Tortoise, desert	Gopherus agassizii	AZ south and east of Colorado R., and Mexico, when found outside of Mexico or said range in AZ	\$500	\$0	\$500
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$69,100	\$0	\$69,100
		Tortoise, gopher	Gopherus polyphemus	West of Mobile and Tombigbee Rivers	\$10,700	\$0	\$10,700
		turtle, bog	Clemmys muhlenbergii	Wherever found, except GA, NC, SC, TN, VA	\$28,700	\$0	\$28,700
	Reptiles subtotal				\$583,100	\$0	\$583,100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Amphibians	Frog, California red-legged	<i>Rana draytonii</i>	Wherever found	\$15,100	\$0	\$15,100
		Frog, Chiricahua leopard	<i>Rana chiricahuensis</i>	Wherever found	\$13,000	\$0	\$13,000
		Salamander, California tiger	<i>Ambystoma californiense</i>	U.S.A. (CA - Santa Barbara County)	\$100	\$0	\$100
		Salamander, California tiger	<i>Ambystoma californiense</i>	U.S.A. (CA - Sonoma County)	\$100	\$0	\$100
		Salamander, California tiger	<i>Ambystoma californiense</i>	U.S.A. (Central CA DPS)	\$100	\$0	\$100
		Salamander, Sonora tiger	<i>Ambystoma tigrinum stebbinsi</i>	Wherever found	\$100	\$0	\$100
		Toad, Houston	<i>Bufo houstonensis</i>	Wherever found	\$200	\$0	\$200
	Amphibians subtotal				\$28,700	\$0	\$28,700
	Fishes	Catfish, Yaqui	<i>Ictalurus pricei</i>	Wherever found	\$100	\$0	\$100
		Chub, bonytail	<i>Gila elegans</i>	Wherever found	\$100	\$0	\$100
		Chub, Chihuahua	<i>Gila nigrescens</i>	Wherever found	\$100	\$0	\$100
		Chub, Gila	<i>Gila intermedia</i>	Wherever found	\$600	\$0	\$600
		Chub, humpback	<i>Gila cypha</i>	Wherever found	\$500	\$0	\$500
		Chub, Sonora	<i>Gila ditaenia</i>	Wherever found	\$100	\$0	\$100
		Chub, Virgin River	<i>Gila seminuda (=robusta)</i>	Wherever found	\$500	\$0	\$500
		Chub, Yaqui	<i>Gila purpurea</i>	Wherever found	\$100	\$0	\$100
		Darter, Okaloosa	<i>Etheostoma okaloosae</i>	Wherever found	\$75,000	\$0	\$75,000
		Darter, watercress	<i>Etheostoma nuchale</i>	Wherever found	\$1,000	\$0	\$1,000
		Minnow, Devils River	<i>Dionda diaboli</i>	Wherever found	\$400	\$0	\$400
		Minnow, loach	<i>Tiaroga cobitis</i>	Wherever found	\$500	\$0	\$500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Minnow, Rio Grande Silvery	Hybognathus amarus	Wherever found, except where listed as an experimental population	\$100	\$0	\$100
		Pikeminnow (=squawfish), Colorado	Ptychocheilus lucius	Wherever found, except where listed as an experimental population	\$500	\$0	\$500
		Pupfish, desert	Cyprinodon macularius	Wherever found	\$500	\$0	\$500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	California Coastal ESU	\$100	\$0	\$100
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Central Valley spring-run ESU	\$100	\$0	\$100
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$9,700	\$0	\$9,700
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Sacramento River winter- run ESU	\$100	\$0	\$100
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall-run ESU	\$5,200	\$0	\$5,200
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer -run ESU	\$5,200	\$0	\$5,200
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Upper Columbia spring-run ESU	\$432,000	\$0	\$432,000
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$5,200	\$0	\$5,200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, chum	Oncorhynchus keta	Hood Canal summer-run ESU	\$5,000	\$0	\$5,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$100	\$0	\$100
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$427,000	\$0	\$427,000
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$218,400	\$0	\$218,400
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$5,100	\$0	\$5,100
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Ozette Lake ESU	\$5,000	\$0	\$5,000
		Salmon, sockeye	Oncorhynchus (=Salmo) nerka	Snake River ESU	\$427,000	\$0	\$427,000
		Shiner, Arkansas River	Notropis girardi	Arkansas River Basin (AR, KS, NM, OK, TX)	\$2,000	\$0	\$2,000
		Shiner, beautiful	Cyprinella formosa	Wherever found	\$100	\$0	\$100
		Spikedace	Meda fulgida	Wherever found	\$500	\$0	\$500
		Spinedace, Little Colorado	Lepidomeda vittata	Wherever found	\$500	\$0	\$500
		Steelhead	Oncorhynchus (=Salmo) mykiss	California Central Valley DPS	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Steelhead	Oncorhynchus (=Salmo) mykiss	Central California Coast DPS	\$100	\$0	\$100
		Steelhead	Oncorhynchus (=Salmo) mykiss	Lower Columbia River DPS	\$112,000	\$0	\$112,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Middle Columbia River DPS	\$112,000	\$0	\$112,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Snake River Basin DPS	\$112,000	\$0	\$112,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	South-Central California Coast DPS	\$100	\$0	\$100
		Steelhead	Oncorhynchus (=Salmo) mykiss	Southern California DPS	\$100	\$0	\$100
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Columbia River DPS	\$112,000	\$0	\$112,000
		Steelhead	Oncorhynchus (=Salmo) mykiss	Upper Willamette River DPS	\$5,000	\$0	\$5,000
		Sturgeon, pallid	Scaphirhynchus albus	Wherever found	\$300	\$0	\$300
		Sturgeon, white	Acipenser transmontanus	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$100	\$0	\$100
		Sucker, razorback	Xyrauchen texanus	Wherever found	\$13,000	\$0	\$13,000
		Sucker, Santa Ana	Catostomus santaanae	3 CA river basins	\$200	\$0	\$200
		Sucker, Zuni bluehead	Catostomus discobolus yarrowi	Wherever found	\$500	\$0	\$500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Topminnow, Gila (incl. Yaqui)	Poeciliopsis occidentalis	Wherever found	\$500	\$0	\$500
		Trout, Apache	Oncorhynchus apache	Wherever found	\$500	\$0	\$500
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$3,000	\$0	\$3,000
		Trout, Gila	Oncorhynchus gilae	Wherever found	\$500	\$0	\$500
		Woundfin	Plagopterus argentissimus	Wherever found, except where listed as an experimental population	\$500	\$0	\$500
	Fishes subtotal				\$2,100,900	\$0	\$2,100,900
	Clams	Pearlshell, Louisiana	Margaritifera hembeli	Wherever found	\$9,800	\$0	\$9,800
		Spiny mussel, James	Pleurobema collina	Wherever found	\$1,700	\$0	\$1,700
		Wedgemussel, dwarf	Alasmidonta heterodon	Wherever found	\$16,200	\$0	\$16,200
	Clams subtotal				\$27,700	\$0	\$27,700
	Snails	Ambersnail, Kanab	Oxyloma haydeni kanabensis	Wherever found	\$500	\$0	\$500
		Limpet, Banbury Springs	Lanx sp.	Wherever found	\$1,800	\$0	\$1,800
		Snail, Bliss Rapids	Taylorconcha serpenticola	Wherever found	\$1,800	\$0	\$1,800
		Snail, Snake River physa	Physa natricina	Wherever found	\$1,800	\$0	\$1,800
		Snail, Stock Island tree	Orthalicus reses (not incl. nesodryas)	Wherever found	\$400	\$0	\$400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Springsnail, Alamosa	Tryonia alamosae	Wherever found	\$100	\$0	\$100
		Springsnail, Bruneau Hot	Pyrgulopsis bruneauensis	Wherever found	\$1,800	\$0	\$1,800
		Springsnail, Chupadera	Pyrgulopsis chupaderae	Wherever found	\$100	\$0	\$100
		springsnail, San Bernardino	Pyrgulopsis bernardina	Wherever found	\$100	\$0	\$100
		Springsnail, Socorro	Pyrgulopsis neomexicana	Wherever found	\$100	\$0	\$100
		Springsnail, Three Forks	Pyrgulopsis trivialis	Wherever found	\$100	\$0	\$100
	Snails subtotal				\$8,600	\$0	\$8,600
	Insects	Bumble bee, Rusty patched	Bombus affinis	Wherever found	\$1,300	\$0	\$1,300
		Butterfly, Bartram's hairstreak	Strymon acis bartrami	Wherever found	\$900	\$0	\$900
		Butterfly, Florida leafwing	Anaea troglodyta floridalis	Wherever found	\$900	\$0	\$900
		Butterfly, Karner blue	Lycaeides melissa samuelis	Wherever found	\$8,100	\$0	\$8,100
		Butterfly, Miami Blue	Cyclargus (=Hemiargus) thomasi bethunebakeri	Wherever found	\$1,200	\$0	\$1,200
		Butterfly, Schaus swallowtail	Heraclides aristodemus ponceanus	Wherever found	\$900	\$0	\$900
		Dragonfly, Hine's emerald	Somatochlora hineana	Wherever found	\$20,000	\$0	\$20,000
		Skipper, Carson wandering	Pseudocopaeodes eunus obscurus	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Skipper, Dakota	<i>Hesperia dacotae</i>	Wherever found	\$300	\$0	\$300
		skipperling, Poweshiek	<i>Oarisma poweshiek</i>	Wherever found	\$300	\$0	\$300
	Insects subtotal				\$34,000	\$0	\$34,000
	Crustaceans	Fairy shrimp, vernal pool	<i>Branchinecta lynchi</i>	Wherever found	\$100	\$0	\$100
		Isopod, Lee County cave	<i>Lirceus usdagalun</i>	Wherever found	\$5,000	\$0	\$5,000
	Crustaceans subtotal				\$5,100	\$0	\$5,100
	Flowering Plants	Ambrosia, San Diego	<i>Ambrosia pumila</i>	Wherever found	\$100	\$0	\$100
		Ambrosia, south Texas	<i>Ambrosia cheiranthifolia</i>	Wherever found	\$200	\$0	\$200
		Aster, decurrent false	<i>Boltonia decurrens</i>	Wherever found	\$100	\$0	\$100
		Aster, Florida golden	<i>Chrysopsis floridana</i>	Wherever found	\$100	\$0	\$100
		Aster, Ruth's golden	<i>Pityopsis ruthii</i>	Wherever found	\$100	\$0	\$100
		Ayenia, Texas	<i>Ayenia limitaris</i>	Wherever found	\$200	\$0	\$200
		Baccharis, Encinitas	<i>Baccharis vanessae</i>	Wherever found	\$100	\$0	\$100
		Barbara's Buttons, Mohr's	<i>Marshallia mohrii</i>	Wherever found	\$100	\$0	\$100
		Barberry, Nevin's	<i>Berberis nevinii</i>	Wherever found	\$200	\$0	\$200
		Beargrass, Britton's	<i>Nolina brittoniana</i>	Wherever found	\$21,700	\$0	\$21,700
		Beauty, Harper's	<i>Harperocallis flava</i>	Wherever found	\$21,700	\$0	\$21,700
		Bittercress, small-anthered	<i>Cardamine micranthera</i>	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Bladderpod, Dudley Bluffs	<i>Lesquerella congesta</i>	Wherever found	\$200	\$0	\$200
		Bladderpod, kodachrome	<i>Lesquerella tumulosa</i>	Wherever found	\$200	\$0	\$200
		Bladderpod, lyrata	<i>Lesquerella lyrata</i>	Wherever found	\$200	\$0	\$200
		bladderpod, Missouri	<i>Physaria filiformis</i>	Wherever found	\$200	\$0	\$200
		Bladderpod, San Bernardino Mountains	<i>Lesquerella kingii</i> ssp. <i>bernardina</i>	Wherever found	\$200	\$0	\$200
		Bladderpod, Short's	<i>Physaria globosa</i>	Wherever found	\$200	\$0	\$200
		Bladderpod, Spring Creek	<i>Lesquerella perforata</i>	Wherever found	\$200	\$0	\$200
		Bladderpod, white	<i>Lesquerella pallida</i>	Wherever found	\$200	\$0	\$200
		Bladderpod, White Bluffs	<i>Physaria douglasii</i> ssp. <i>tuplashensis</i>	Wherever found	\$400	\$0	\$400
		Bladderpod, Zapata	<i>Lesquerella thamnophila</i>	Wherever found	\$400	\$0	\$400
		Blazingstar, Heller's	<i>Liatris helleri</i>	Wherever found	\$100	\$0	\$100
		Blazingstar, scrub	<i>Liatris ohlingerae</i>	Wherever found	\$100	\$0	\$100
		Blue-star, Kearney's	<i>Amsonia kearneyana</i>	Wherever found	\$100	\$0	\$100
		Bonamia, Florida	<i>Bonamia grandiflora</i>	Wherever found	\$21,700	\$0	\$21,700
		Brickell-bush, Florida	<i>Brickellia mosieri</i>	Wherever found	\$100	\$0	\$100
		Buckwheat, Umtanum Desert	<i>Eriogonum codium</i>	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Bulrush, Northeastern	Scirpus ancistrochaetus	Wherever found	\$28,500	\$0	\$28,500
		Butterweed, Layne's	Senecio layneae	Wherever found	\$100	\$0	\$100
		Cactus, Acuna	Echinomastus erectocentrus var. acunensis	Wherever found	\$100	\$0	\$100
		Cactus, Arizona hedgehog	Echinocereus triglochidiatus var. arizonicus	Wherever found	\$500	\$0	\$500
		Cactus, black lace	Echinocereus reichenbachii var. albertii	Wherever found	\$100	\$0	\$100
		Cactus, Brady pincushion	Pediocactus bradyi	Wherever found	\$500	\$0	\$500
		Cactus, Cochise pincushion	Coryphantha robbinsiorum	Wherever found	\$100	\$0	\$100
		Cactus, Fickeisen plains	Pediocactus peeblesianus fickeiseniae	Wherever found	\$500	\$0	\$500
		Cactus, Florida semaphore	Consolea corallicola	Wherever found	\$400	\$0	\$400
		Cactus, Key tree	Pilosocereus robinii	Wherever found	\$400	\$0	\$400
		Cactus, Knowlton's	Pediocactus knowltonii	Wherever found	\$100	\$0	\$100
		Cactus, Mesa Verde	Sclerocactus mesae-verdae	Wherever found	\$100	\$0	\$100
		Cactus, Nichol's Turk's head	Echinocactus horizontalonius var. nicholii	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Cactus, Peebles Navajo	<i>Pediocactus peeblesianus</i> var. <i>peeblesianus</i>	Wherever found	\$100	\$0	\$100
		Cactus, Pima pineapple	<i>Coryphantha scheeri</i> var. <i>robustispina</i>	Wherever found	\$100	\$0	\$100
		Cactus, Siler pincushion	<i>Pediocactus</i> (=Echinocactus,=Utahia) <i>sileri</i>	Wherever found	\$100	\$0	\$100
		Cactus, star	<i>Astrophytum asterias</i>	Wherever found	\$200	\$0	\$200
		cactus, Tobusch fishhook	<i>Sclerocactus brevihamatus</i> ssp. <i>tobuschii</i>	Wherever found	\$100	\$0	\$100
		Catchfly, Spalding's	<i>Silene spaldingii</i>	Wherever found	\$700	\$0	\$700
		Checkermallow, Wenatchee Mountains	<i>Sidalcea oregana</i> var. <i>calva</i>	Local Endemic	\$200	\$0	\$200
		Cliffrose, Arizona	<i>Purshia</i> (=Cowania) <i>subintegra</i>	Wherever found	\$500	\$0	\$500
		Clover, showy Indian	<i>Trifolium amoenum</i>	Wherever found	\$100	\$0	\$100
		Coneflower, smooth	<i>Echinacea laevigata</i>	Wherever found	\$100	\$0	\$100
		Crownbeard, big-leaved	<i>Verbesina dissita</i>	Wherever found	\$100	\$0	\$100
		Cycladenia, Jones	<i>Cycladenia humilis</i> var. <i>jonesii</i>	Wherever found	\$100	\$0	\$100
		Daisy, Lakeside	<i>Hymenoxys herbacea</i>	Wherever found	\$100	\$0	\$100
		Daisy, Parish's	<i>Erigeron parishii</i>	Wherever found	\$100	\$0	\$100
		Daisy, Willamette	<i>Erigeron decumbens</i>	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Dawn-flower, Texas prairie	Hymenoxys texana	Wherever found	\$100	\$0	\$100
		Dogweed, ashy	Thymophylla tephroleuca	Wherever found	\$300	\$0	\$300
		Flax, sand	Linum arenicola	Wherever found	\$700	\$0	\$700
		Fleabane, Zuni	Erigeron rhizomatus	Wherever found	\$200	\$0	\$200
		Four-o'clock, MacFarlane's	Mirabilis macfarlanei	Wherever found	\$100	\$0	\$100
		Fringepod, Santa Cruz Island	Thysanocarpus conchuliferus	Wherever found	\$200	\$0	\$200
		Glade cress, Kentucky	Leavenworthia exigua laciniata	Wherever found	\$200	\$0	\$200
		Gladecress, Fleshy-fruit	Leavenworthia crassa	Wherever found	\$200	\$0	\$200
		Gladecress, Texas golden	Leavenworthia texana	Wherever found	\$200	\$0	\$200
		Goldenrod, Blue Ridge	Solidago spithamaea	Wherever found	\$100	\$0	\$100
		Goldenrod, Houghton's	Solidago houghtonii	Wherever found	\$100	\$0	\$100
		Goldenrod, Short's	Solidago shortii	Wherever found	\$100	\$0	\$100
		Goldfields, Burke's	Lasthenia burkei	Wherever found	\$100	\$0	\$100
		Goldfields, Contra Costa	Lasthenia conjugens	Wherever found	\$200	\$0	\$200
		Gumplant, Ash Meadows	Grindelia fraxinipratensis	Wherever found	\$100	\$0	\$100
		halumtano, Cebello	Bulbophyllum guamense	Wherever found	\$100	\$0	\$100
		Howellia, water	Howellia aquatilis	Wherever found	\$300	\$0	\$300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Jacquemontia, beach	Jacquemontia reclinata	Wherever found	\$500	\$0	\$500
		Jewelflower, California	Caulanthus californicus	Wherever found	\$200	\$0	\$200
		Jewelflower, Metcalf Canyon	Streptanthus albidus ssp. albidus	Wherever found	\$200	\$0	\$200
		Jewelflower, Tiburon	Streptanthus niger	Wherever found	\$200	\$0	\$200
		Joint-vetch, Sensitive	Aeschynomene virginica	Wherever found	\$1,400	\$0	\$1,400
		Ladies'-tresses, Ute	Spiranthes diluvialis	Wherever found	\$800	\$0	\$800
		ladies-tresses, Canelo Hills	Spiranthes delitescens	Wherever found	\$100	\$0	\$100
		Layia, beach	Layia carnosa	Wherever found	\$100	\$0	\$100
		Lead-plant, Crenulate	Amorpha crenulata	Wherever found	\$21,700	\$0	\$21,700
		Lessingia, San Francisco	Lessingia germanorum (=L.g. var. germanorum)	Wherever found	\$100	\$0	\$100
		Malacothrix, island	Malacothrix squalida	Wherever found	\$100	\$0	\$100
		Malacothrix, Santa Cruz Island	Malacothrix indecora	Wherever found	\$100	\$0	\$100
		mallow, Gierisch	Sphaeralcea gierischii	Wherever found	\$500	\$0	\$500
		Manioc, Walker's	Manihot walkerae	Wherever found	\$400	\$0	\$400
		Milk-vetch, Holmgren	Astragalus holmgreniorum	Wherever found	\$500	\$0	\$500
		Milk-vetch, Mancos	Astragalus humillimus	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Milk-vetch, Sentry	<i>Astragalus cremnophylax</i> var. <i>cremnophylax</i>	Wherever found	\$100	\$0	\$100
		Milkweed, Mead's	<i>Asclepias meadii</i>	Wherever found	\$20,000	\$0	\$20,000
		Milkweed, Welsh's	<i>Asclepias welshii</i>	Wherever found	\$100	\$0	\$100
		Mint, scrub	<i>Dicerandra frutescens</i>	Wherever found	\$21,700	\$0	\$21,700
		Monkshood, northern wild	<i>Aconitum noveboracense</i>	Wherever found	\$100	\$0	\$100
		Mustard, Carter's	<i>Warea carteri</i>	Wherever found	\$200	\$0	\$200
		Mustard, Penland alpine fen	<i>Eutrema penlandii</i>	Wherever found	\$200	\$0	\$200
		Mustard, slender- petaled	<i>Thelypodium stenopetalum</i>	Wherever found	\$200	\$0	\$200
		No common name	<i>Cranichis ricartii</i>	Wherever found	\$100	\$0	\$100
		No common name	<i>Dendrobium guamense</i>	Wherever found	\$100	\$0	\$100
		No common name	<i>Lepanthes eltoroensis</i>	Wherever found	\$100	\$0	\$100
		No common name	<i>Nervilia jacksoniae</i>	Wherever found	\$100	\$0	\$100
		No common name	<i>Platanthera holochila</i>	Wherever found	\$100	\$0	\$100
		No common name	<i>Tuberolabium guamense</i>	Wherever found	\$100	\$0	\$100
		Orchid, eastern prairie fringed	<i>Platanthera leucophaea</i>	Wherever found	\$800	\$0	\$800
		paintbrush, golden	<i>Castilleja levisecta</i>	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Pea, Big Pine partridge	<i>Chamaecrista lineata keyensis</i>	Wherever found	\$400	\$0	\$400
		Penny-cress, Kneeland Prairie	<i>Thlaspi californicum</i>	Wherever found	\$200	\$0	\$200
		Pennyroyal, Todsens's	<i>Hedeoma todsenii</i>	Wherever found	\$100	\$0	\$100
		Pentachaeta, Lyon's	<i>Pentachaeta lyonii</i>	Wherever found	\$100	\$0	\$100
		Pentachaeta, white-rayed	<i>Pentachaeta bellidiflora</i>	Wherever found	\$100	\$0	\$100
		Peppergrass, Slickspot	<i>Lepidium papilliferum</i>		\$300	\$0	\$300
		Pinkroot, gentian	<i>Spigelia gentianoides</i>	Wherever found	\$21,700	\$0	\$21,700
		Plum, scrub	<i>Prunus geniculata</i>	Wherever found	\$21,700	\$0	\$21,700
		Pogonia, small whorled	<i>Isotria medeoloides</i>	Wherever found	\$100	\$0	\$100
		Ragwort, San Francisco Peaks	<i>Packera franciscana</i>	Wherever found	\$200	\$0	\$200
		Reed-mustard, Barneby	<i>Schoenocrambe barnebyi</i>	Wherever found	\$200	\$0	\$200
		Reed-mustard, clay	<i>Schoenocrambe argillacea</i>	Wherever found	\$200	\$0	\$200
		Reed-mustard, shrubby	<i>Schoenocrambe suffrutescens</i>	Wherever found	\$200	\$0	\$200
		Rhododendron, Chapman	<i>Rhododendron chapmanii</i>	Wherever found	\$21,700	\$0	\$21,700
		Ridge-cress, Barneby	<i>Lepidium barnebyanum</i>	Wherever found	\$200	\$0	\$200
		rock cress, Shale barren	<i>Arabis serotina</i>	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Rock-cress, Braun's	<i>Arabis perstellata</i>	Wherever found	\$200	\$0	\$200
		Rock-cress, Hoffmann's	<i>Arabis hoffmannii</i>	Wherever found	\$200	\$0	\$200
		Rock-cress, McDonald's	<i>Arabis macdonaldiana</i>	Wherever found	\$200	\$0	\$200
		Rockcross, Georgia	<i>Arabis georgiana</i>	Wherever found	\$200	\$0	\$200
		Rockcross, Santa Cruz Island	<i>Sibara filifolia</i>	Wherever found	\$200	\$0	\$200
		Rosemary, Apalachicola	<i>Conradina glabra</i>	Wherever found	\$21,700	\$0	\$21,700
		Sandlace	<i>Polygonella myriophylla</i>	Wherever found	\$21,700	\$0	\$21,700
		Sedge, Navajo	<i>Carex specuicola</i>	Wherever found	\$100	\$0	\$100
		silverbush, Blodgett's	<i>Argythamnia blodgettii</i>	Wherever found	\$1,000	\$0	\$1,000
		Snakeroot	<i>Eryngium cuneifolium</i>	Wherever found	\$21,700	\$0	\$21,700
		Sneezeweed, Virginia	<i>Helenium virginicum</i>	Wherever found	\$100	\$0	\$100
		Snowbells, Texas	<i>Styrax texanus</i>	Wherever found	\$100	\$0	\$100
		Spurge, deltoid	<i>Chamaesyce deltoidea</i> ssp. <i>deltoidea</i>	Wherever found	\$200	\$0	\$200
		Spurge, Garber's	<i>Chamaesyce garberi</i>	Wherever found	\$700	\$0	\$700
		Spurge, Hoover's	<i>Chamaesyce hooveri</i>	Wherever found	\$200	\$0	\$200
		Spurge, telephus	<i>Euphorbia telephioides</i>	Wherever found	\$200	\$0	\$200
		Spurge, wedge	<i>Chamaesyce deltoidea serpyllum</i>	Wherever found	\$700	\$0	\$700
		Stickseed, showy	<i>Hackelia venusta</i>	Wherever found	\$200	\$0	\$200
		Sumac, Michaux's	<i>Rhus michauxii</i>	Wherever found	\$500	\$0	\$500

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Sunburst, Hartweg's golden	<i>Pseudobahia bahiifolia</i>	Wherever found	\$100	\$0	\$100
		Sunburst, San Joaquin adobe	<i>Pseudobahia peirsonii</i>	Wherever found	\$100	\$0	\$100
		Sunflower, Pecos (=puzzle, =paradox)	<i>Helianthus paradoxus</i>	Wherever found	\$200	\$0	\$200
		Sunflower, San Mateo woolly	<i>Eriophyllum latilobum</i>	Wherever found	\$100	\$0	\$100
		Sunflower, Schweinitz's	<i>Helianthus schweinitzii</i>	Wherever found	\$100	\$0	\$100
		Sunflower, whorled	<i>Helianthus verticillatus</i>		\$100	\$0	\$100
		Sunray, Ash Meadows	<i>Enceliopsis nudicaulis</i> var. <i>corrugata</i>	Wherever found	\$100	\$0	\$100
		Sunshine, Sonoma	<i>Blennosperma bakeri</i>	Wherever found	\$100	\$0	\$100
		Taraxacum, California	<i>Taraxacum californicum</i>	Wherever found	\$100	\$0	\$100
		Tarplant, Gaviota	<i>Deinandra increscens</i> ssp. <i>villosa</i>	Wherever found	\$100	\$0	\$100
		Tarplant, Otay	<i>Deinandra</i> (=Hemizonia) <i>conjugens</i>	Wherever found	\$100	\$0	\$100
		Tarplant, Santa Cruz	<i>Holocarpha macradenia</i>	Wherever found	\$100	\$0	\$100
		Thelypody, Howell's spectacular	<i>Thelypodium howellii</i> <i>spectabilis</i>	Wherever found	\$200	\$0	\$200
		Thistle, Chorro Creek bog	<i>Cirsium fontinale</i> var. <i>obispoense</i>	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Thistle, fountain	<i>Cirsium fontinale</i> var. <i>fontinale</i>	Wherever found	\$100	\$0	\$100
		Thistle, La Graciosa	<i>Cirsium loncholepis</i>	Wherever found	\$100	\$0	\$100
		Thistle, Pitcher's	<i>Cirsium pitcheri</i>	Wherever found	\$8,100	\$0	\$8,100
		Thistle, Sacramento Mountains	<i>Cirsium vinaceum</i>	Wherever found	\$100	\$0	\$100
		Thistle, Suisun	<i>Cirsium hydrophilum</i> var. <i>hydrophilum</i>	Wherever found	\$100	\$0	\$100
		Thoroughwort, Cape Sable	<i>Chromolaena frustrata</i>	Wherever found	\$100	\$0	\$100
		Townsendia, Last Chance	<i>Townsendia aprica</i>	Wherever found	\$100	\$0	\$100
		Twinpod, Dudley Bluffs	<i>Physaria obcordata</i>	Wherever found	\$200	\$0	\$200
		Wallflower, Ben Lomond	<i>Erysimum teretifolium</i>	Wherever found	\$200	\$0	\$200
		Wallflower, Contra Costa	<i>Erysimum capitatum</i> var. <i>angustatum</i>	Wherever found	\$200	\$0	\$200
		Wallflower, Menzies'	<i>Erysimum menziesii</i>	Wherever found	\$200	\$0	\$200
		Warea, wide- leaf	<i>Warea amplexifolia</i>	Wherever found	\$200	\$0	\$200
		Water-umbel, Huachuca	<i>Lilaeopsis schaffneriana</i> var. <i>recurva</i>	Wherever found	\$100	\$0	\$100
		Watercress, Gambel's	<i>Rorippa gambellii</i>	Wherever found	\$200	\$0	\$200
		Wire-lettuce, Malheur	<i>Stephanomeria malheurensis</i>	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wireweed	Polygonella basiramia	Wherever found	\$21,700	\$0	\$21,700
		Wooly-threads, San Joaquin	Monolopia (=Lembertia) congdonii	Wherever found	\$100	\$0	\$100
		Yellowhead, desert	Yermo xanthocephalus	Wherever found	\$100	\$0	\$100
	Flowering Plants subtotal				\$349,400	\$0	\$349,400
	Conifers and Cycads	Torreya, Florida	Torreya taxifolia	Wherever found	\$21,700	\$0	\$21,700
	Conifers and Cycads subtotal				\$21,700	\$0	\$21,700
	Lichens	Cladonia, Florida perforate	Cladonia perforata	Wherever found	\$21,700	\$0	\$21,700
	Lichens subtotal				\$21,700	\$0	\$21,700
Domestic subtotal					\$10,177,000	\$0	\$10,177,000
Foreign	Mammals	Addax	Addax nasomaculatus	Wherever found	\$200	\$0	\$200
		Babirusa	Babyrousa babyrussa	Wherever found	\$300	\$0	\$300
		Cat, leopard	Prionailurus (=Felis) bengalensis bengalensis	Wherever found	\$200	\$0	\$200
		Cat, tiger	Leopardus (=Felis) tigrinus	Wherever found	\$200	\$0	\$200
		Cheetah	Acinonyx jubatus	Wherever found	\$800	\$0	\$800
		Deer, Eld's brow-antlered	Cervus eldi	Wherever found	\$200	\$0	\$200
		Elephant, Asian	Elephas maximus	Wherever found	\$1,000	\$0	\$1,000
		Gazelle, dama	Gazella dama	Wherever found	\$200	\$0	\$200
		Gazelle, slender-horned	Gazella leptoceros	Wherever found	\$300	\$0	\$300
		Gibbons	Hylobates spp. (including Nomascus)	Wherever found	\$700	\$0	\$700
		Gorilla	Gorilla gorilla	Wherever found	\$300	\$0	\$300
		Koala	Phascolarctos cinereus	Australia	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Langur, Francois'	Trachypithecus (=Presbytis) francoisi	Wherever found	\$200	\$0	\$200
		Lechwe, red	Kobus leche	Wherever found	\$200	\$0	\$200
		Lemurs	Lemuridae (incl. genera Lemur, Phaner, Hapalemur, Lepilemur, Microcebus, Allocebus, Cheirogaleus, Varecia)	Wherever found	\$1,100	\$0	\$1,100
		Leopard, clouded	Neofelis nebulosa	Wherever found	\$300	\$0	\$300
		Leopard, snow	Uncia (=Panthera) uncia	Wherever found	\$300	\$0	\$300
		Lion	Panthera leo ssp. leo	Wherever found	\$200	\$0	\$200
		Mandrill	Mandrillus (=Papio) sphinx	Wherever found	\$500	\$0	\$500
		Monkey, Diana	Cercopithecus diana	Wherever found	\$200	\$0	\$200
		Orangutan	Pongo pygmaeus	Wherever found	\$200	\$0	\$200
		Oryx, scimitar-horned	Oryx dammah	Wherever found	\$200	\$0	\$200
		Panda, giant	Ailuropoda melanoleuca	Wherever found	\$300	\$0	\$300
		Rhinoceros, black	Diceros bicornis	Africa	\$300	\$0	\$300
		Rhinoceros, Southern white	Ceratotherium simum ssp. simum	Wherever found	\$300	\$0	\$300
		Siamang	Symphalangus syndactylus	Wherever found	\$300	\$0	\$300
		Tamarin, cotton- top	Saguinus oedipus	Wherever found	\$500	\$0	\$500
		Tapir, Central American	Tapirus bairdii	Wherever found	\$200	\$0	\$200

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Tapir, South American (=Brazilian)	Tapirus terrestris	Wherever found	\$200	\$0	\$200
	Mammals subtotal				\$10,100	\$0	\$10,100
	Reptiles	Iguana, Cuban ground	Cyclura nubila nubila	(Cyclura nubila nubila)	\$30,000	\$0	\$30,000
	Reptiles subtotal				\$30,000	\$0	\$30,000
Foreign subtotal					\$40,100	\$0	\$40,100
Domestic	Multi-Species				\$256,150	\$0	\$256,150
	Multi-Species subtotal				\$256,150	\$0	\$256,150
Domestic subtotal					\$256,150	\$0	\$256,150
Total					\$10,473,250	\$0	\$10,473,250

USDA Forest Service - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	Bat, Indiana	Myotis sodalis	Wherever found	\$475,425	\$0	\$475,425
	Mammals subtotal				\$475,425	\$0	\$475,425
Domestic subtotal					\$475,425	\$0	\$475,425
Total					\$475,425	\$0	\$475,425

USDA Natural Resources Conservation Service - FY 2017

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic	Mammals	bat, Florida bonneted	<i>Eumops floridanus</i>	Wherever found	\$10,000	\$0	\$10,000
		Bat, gray	<i>Myotis grisescens</i>	Wherever found	\$500	\$0	\$500
		Bat, Hawaiian hoary	<i>Lasiurus cinereus semotus</i>	Wherever found	\$3,000	\$0	\$3,000
		Bat, Indiana	<i>Myotis sodalis</i>	Wherever found	\$23,750	\$0	\$23,750
		Bat, Northern long-eared	<i>Myotis septentrionalis</i>	Wherever found	\$179,650	\$250,000	\$429,650
		Bear, grizzly	<i>Ursus arctos horribilis</i>	U.S.A., conterminous (lower 48) States, except where listed as an experimental population or delisted	\$100	\$0	\$100
		Deer, Columbian white-tailed	<i>Odocoileus virginianus leucurus</i>	Columbia River DPS	\$6,700	\$0	\$6,700
		Ferret, black-footed	<i>Mustela nigripes</i>	U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	\$6,200	\$0	\$6,200
		Fox, San Joaquin kit	<i>Vulpes macrotis mutica</i>	wherever found	\$6,048	\$0	\$6,048
		Fox, Santa Catalina Island	<i>Urocyon littoralis catalinae</i>	Wherever found	\$1,680	\$0	\$1,680
		Kangaroo rat, giant	<i>Dipodomys ingens</i>	Wherever found	\$840	\$0	\$840

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Kangaroo rat, Tipton	Dipodomys nitratoides nitratoides	Wherever found	\$12,096	\$0	\$12,096
		Lynx, Canada	Lynx canadensis	Wherever Found in Contiguous U.S.	\$55,600	\$0	\$55,600
		Manatee, West Indian	Trichechus manatus	Wherever found	\$75,000	\$0	\$75,000
		Mouse, Preble's meadow jumping	Zapus hudsonius preblei	wherever found	\$2,000	\$0	\$2,000
		Mouse, salt marsh harvest	Reithrodontomys raviventris	wherever found	\$4,200	\$0	\$4,200
		Panther, Florida	Puma (=Felis) concolor coryi	Wherever found	\$700,000	\$3,000,000	\$3,700,000
		Pocket gopher, Olympia	Thomomys mazama pugetensis	Wherever found	\$83,000	\$0	\$83,000
		Prairie dog, Utah	Cynomys parvidens	Wherever found	\$1,500	\$0	\$1,500
		Rabbit, riparian brush	Sylvilagus bachmani riparius	Wherever found	\$2,688	\$0	\$2,688

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Wolf, gray	<i>Canis lupus</i>	U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	\$1,000	\$0	\$1,000
		Woodrat, riparian (=San Joaquin Valley)	<i>Neotoma fuscipes riparia</i>	Wherever found	\$2,520	\$0	\$2,520
	Mammals subtotal				\$1,178,072	\$3,250,000	\$4,428,072
	Birds	Caracara, Audubon's crested	<i>Polyborus plancus audubonii</i>	FL pop.	\$500,000	\$2,500,000	\$3,000,000
		Coot, Hawaiian	<i>Fulica americana alai</i>	Wherever found	\$500	\$0	\$500
		Crane, whooping	<i>Grus americana</i>	Wherever found, except where listed as an experimental population	\$8,650	\$0	\$8,650

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Cuckoo, yellow-billed	<i>Coccyzus americanus</i>	Western U.S. DPS	\$6,936	\$0	\$6,936
		Curlew, Eskimo	<i>Numenius borealis</i>	Wherever found	\$200	\$0	\$200
		Duck, Hawaiian (=koloa)	<i>Anas wyvilliana</i>	Wherever found	\$500	\$0	\$500
		Flycatcher, southwestern willow	<i>Empidonax traillii extimus</i>	Wherever found	\$23,482	\$0	\$23,482
		gallinule, Hawaiian common	<i>Gallinula galeata sandvicensis</i>	Wherever found	\$500	\$0	\$500
		Gnatcatcher, coastal California	<i>Polioptila californica californica</i>	Wherever found	\$11,680	\$0	\$11,680
		Goose, Hawaiian	<i>Branta (=Nesochen) sandvicensis</i>	Wherever found	\$12,500	\$0	\$12,500
		Hawk, Hawaiian (=ʻIo)	<i>Buteo solitarius</i>	Wherever found	\$1,200	\$0	\$1,200
		Horned lark, streaked	<i>Eremophila alpestris strigata</i>	Wherever found	\$12,500	\$0	\$12,500
		Kite, Everglade snail	<i>Rostrhamus sociabilis plumbeus</i>	Wherever found	\$75,000	\$0	\$75,000
		Knot, red	<i>Calidris canutus rufa</i>	Wherever found	\$60,250	\$0	\$60,250
		Moorhen, Mariana common	<i>Gallinula chloropus guami</i>	Wherever found	\$500	\$0	\$500
		Murrelet, marbled	<i>Brachyramphus marmoratus</i>	U.S.A. (CA, OR, WA)	\$500	\$0	\$500
		Owl, Mexican spotted	<i>Strix occidentalis lucida</i>	Wherever found	\$4,000	\$0	\$4,000
		Owl, northern spotted	<i>Strix occidentalis caurina</i>	Wherever found	\$16,012	\$0	\$16,012

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Plover, piping	Charadrius melodus	[Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	\$8,500	\$0	\$8,500
		Plover, western snowy	Charadrius nivosus nivosus	Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	\$168	\$0	\$168
		sage-grouse, Gunnison	Centrocercus minimus	Wherever found	\$1,186,000	\$0	\$1,186,000
		scrub-jay, Florida	Aphelocoma coerulescens	Wherever found	\$50,000	\$0	\$50,000
		Sparrow, Florida grasshopper	Ammodramus savannarum floridanus	Wherever found	\$50,000	\$1,500,000	\$1,550,000
		Stilt, Hawaiian	Himantopus mexicanus knudseni	Wherever found	\$800	\$0	\$800
		Stork, wood	Mycteria americana	AL, FL, GA, MS, NC, SC	\$750,000	\$10,798,527	\$11,548,527
		Tern, California least	Sterna antillarum browni	Wherever found	\$7,300	\$0	\$7,300
		Vireo, least Bell's	Vireo bellii pusillus	Wherever found	\$3,906	\$0	\$3,906
		Woodpecker, red-cockaded	Picoides borealis	Wherever found	\$1,649,000	\$0	\$1,649,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Birds subtotal				\$4,440,584	\$14,798,527	\$19,239,111
	Reptiles	Lizard, blunt-nosed leopard	Gambelia silus	Wherever found	\$6,216	\$0	\$6,216
		Massasauga (=rattlesnake), eastern	Sistrurus catenatus	Wherever found	\$2,200	\$0	\$2,200
		Skink, bluetail mole	Eumeces egregius lividus	Wherever found	\$750,000	\$1,000,000	\$1,750,000
		Skink, sand	Neoseps reynoldsi	Wherever found	\$50,000	\$0	\$50,000
		Snake, copperbelly water	Nerodia erythrogaster neglecta	Indiana north of 40 degrees north latitude, Michigan, Ohio	\$300	\$0	\$300
		Snake, eastern indigo	Drymarchon corais couperi	Wherever found	\$480,000	\$0	\$480,000
		Snake, giant garter	Thamnophis gigas	Wherever found	\$2,688	\$0	\$2,688
		Tortoise, desert	Gopherus agassizii	Wherever found, except AZ south and east of Colorado R., and Mexico	\$10,150	\$0	\$10,150
		Tortoise, gopher	Gopherus polyphemus	West of Mobile and Tombigbee Rivers	\$479,500	\$61,000	\$540,500
		turtle, bog	Clemmys muhlenbergii	Wherever found, except GA, NC, SC, TN, VA	\$173,213	\$497,767	\$670,980
		Whipsnake (=striped racer), Alameda	Masticophis lateralis euryxanthus	Wherever found	\$11,704	\$0	\$11,704

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
	Reptiles subtotal				\$1,965,971	\$1,558,767	\$3,524,738
	Amphibians	Frog, California red-legged	Rana draytonii	Wherever found	\$110,346	\$0	\$110,346
		Frog, mountain yellow-legged	Rana muscosa	Northern California DPS	\$1,512	\$0	\$1,512
		Frog, Oregon spotted	Rana pretiosa	Wherever found	\$33,500	\$0	\$33,500
		Hellbender, Ozark	Cryptobranchus alleganiensis bishopi	Wherever found	\$3,000	\$0	\$3,000
		Salamander, California tiger	Ambystoma californiense	U.S.A. (CA - Sonoma County)	\$7,056	\$0	\$7,056
		Salamander, California tiger	Ambystoma californiense	U.S.A. (Central CA DPS)	\$336,788	\$0	\$336,788
		Salamander, frosted flatwoods	Ambystoma cingulatum	Wherever found	\$100,000	\$0	\$100,000
		salamander, Reticulated flatwoods	Ambystoma bishopi	Wherever found	\$100,000	\$0	\$100,000
		Toad, arroyo (=arroyo southwestern)	Anaxyrus californicus	Wherever found	\$1,218	\$0	\$1,218
		Toad, Wyoming	Bufo hemiophrys baxteri	Wherever found	\$1,500	\$0	\$1,500
	Amphibians subtotal				\$694,920	\$0	\$694,920
	Fishes	Chub, bonytail	Gila elegans	Wherever found	\$200	\$0	\$200
		Chub, humpback	Gila cypha	Wherever found	\$200	\$0	\$200
		Chub, Pahrnagat roundtail	Gila robusta jordani	Wherever found	\$100	\$0	\$100
		darter, bluemask	Etheostoma akatulo	Wherever found	\$200	\$0	\$200
		Darter, Okaloosa	Etheostoma okaloosae	Wherever found	\$25,000	\$0	\$25,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Darter, snail	Percina tanasi	Wherever found	\$200	\$0	\$200
		Goby, tidewater	Eucyclogobius newberryi	Wherever found	\$84	\$0	\$84
		Pikeminnow (=squawfish), Colorado	Ptychocheilus lucius	Wherever found, except where listed as an experimental population	\$200	\$0	\$200
		Salmon, Atlantic	Salmo salar	Gulf of Maine DPS	\$386,000	\$0	\$386,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Lower Columbia River ESU	\$2,500	\$0	\$2,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Puget Sound ESU	\$323,000	\$0	\$323,000
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River fall- run ESU	\$73,500	\$0	\$73,500
		Salmon, Chinook	Oncorhynchus (=Salmo) tshawytscha	Snake River spring/summer -run ESU	\$73,500	\$0	\$73,500
		Salmon, chum	Oncorhynchus keta	Columbia River ESU	\$2,500	\$0	\$2,500
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Central California Coast ESU	\$840	\$0	\$840
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Lower Columbia River ESU	\$2,500	\$0	\$2,500
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Oregon Coast ESU	\$14,700	\$0	\$14,700

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Salmon, coho	Oncorhynchus (=Salmo) kisutch	Southern Oregon - Northern California Coast ESU	\$2,000	\$0	\$2,000
		Sawfish, smalltooth	Pristis pectinata	US DPS - Smalltooth sawfish originating from U.S. waters.	\$25,000	\$0	\$25,000
		Sculpin, Grotto	Cottus specus	Wherever found	\$4,000	\$0	\$4,000
		Shiner, Topeka	Notropis topeka (=tristis)	U.S.A. (MO-specified portions of Little Creek, Big Muddy Creek, and Spring Creek watersheds in Adair, Gentry, Harrison, Putnam, Sullivan, and Worth Counties; see 17.84(d)(1)(i))	\$2,600	\$0	\$2,600
		Spinedace, White River	Lepidomeda albivallis	Wherever found	\$100	\$0	\$100
		Springfish, Hiko White River	Crenichthys baileyi grandis	Wherever found	\$100	\$0	\$100

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Springfish, Railroad Valley	<i>Crenichthys nevadae</i>	Wherever found	\$500	\$0	\$500
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	California Central Valley DPS	\$672	\$0	\$672
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Central California Coast DPS	\$4,368	\$0	\$4,368
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Lower Columbia River DPS	\$2,500	\$0	\$2,500
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Puget Sound DPS	\$527,000	\$0	\$527,000
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Snake River Basin DPS	\$199,000	\$0	\$199,000
		Steelhead	<i>Oncorhynchus (=Salmo) mykiss</i>	Southern California DPS	\$1,050	\$0	\$1,050
		Sturgeon, Atlantic	<i>Acipenser oxyrinchus oxyrinchus</i>	Gulf of Maine DPS	\$105,000	\$0	\$105,000
		Sturgeon, pallid	<i>Scaphirhynchus albus</i>	Wherever found	\$9,400	\$0	\$9,400
		Sturgeon, shortnose	<i>Acipenser brevirostrum</i>	Wherever found	\$80,000	\$0	\$80,000
		Sturgeon, white	<i>Acipenser transmontanus</i>	U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	\$1,600	\$0	\$1,600
		Sucker, razorback	<i>Xyrauchen texanus</i>	Wherever found	\$200	\$0	\$200
		Sucker, Warner	<i>Catostomus warnerensis</i>	Wherever found	\$2,300	\$0	\$2,300

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Trout, bull	Salvelinus confluentus	U.S.A., conterminous, lower 48 states	\$896,200	\$0	\$896,200
		Trout, Lahontan cutthroat	Oncorhynchus clarkii henshawi	Wherever found	\$500	\$0	\$500
	Fishes subtotal				\$2,769,314	\$0	\$2,769,314
	Clams	Bankclimber, purple (mussel)	Elliptoideus sloatianus	Wherever found	\$20,000	\$0	\$20,000
		Bean, Choctaw	Villosa choctawensis	Wherever found	\$20,000	\$0	\$20,000
		Ebonysell, round	Fusconaia rotulata	Wherever found	\$20,000	\$0	\$20,000
		Elktoe, Appalachian	Alasmidonta raveneliana	Wherever found	\$202,000	\$0	\$202,000
		Elktoe, Cumberland	Alasmidonta atropurpurea	Wherever found	\$500	\$0	\$500
		Kidneyshell, fluted	Ptychobranthus subtentum	Wherever found	\$500	\$0	\$500
		Kidneyshell, southern	Ptychobranthus jonesi	Wherever found	\$20,000	\$0	\$20,000
		Moccasinshell, Gulf	Medionidus penicillatus	Wherever found	\$20,000	\$0	\$20,000
		Moccasinshell, Ochlockonee	Medionidus simpsonianus	Wherever found	\$20,000	\$0	\$20,000
		Mussel, scaleshell	Leptodea leptodon	Wherever found	\$800	\$0	\$800
		Pigtoe, fuzzy	Pleurobema strodeanum	Wherever found	\$20,000	\$0	\$20,000
		Pigtoe, narrow	Fusconaia escambia	Wherever found	\$20,000	\$0	\$20,000
		Pigtoe, oval	Pleurobema pyriforme	Wherever found	\$20,000	\$0	\$20,000
		Pocketbook, shinyrayed	Lampsilis subangulata	Wherever found	\$20,000	\$0	\$20,000
		sandshell, Southern	Hamiota australis	Wherever found	\$20,000	\$0	\$20,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Slabshell, Chipola	<i>Elliptio chipolaensis</i>	Wherever found	\$20,000	\$0	\$20,000
		Spectaclecase (mussel)	<i>Cumberlandia monodonta</i>	Wherever found	\$500	\$0	\$500
		Threeridge, fat (mussel)	<i>Amblema neislerii</i>	Wherever found	\$20,000	\$0	\$20,000
		Wedgemussel, dwarf	<i>Alasmidonta heterodon</i>	Wherever found	\$6,100	\$35,000	\$41,100
	Clams subtotal				\$470,400	\$35,000	\$505,400
	Snails	Snail, Guam tree	<i>Partula radiolata</i>	Wherever found	\$500	\$0	\$500
		Springsnail, Bruneau Hot	<i>Pyrgulopsis bruneauensis</i>	Wherever found	\$4,000	\$0	\$4,000
	Snails subtotal				\$4,500	\$0	\$4,500
	Insects	Beetle, American burying	<i>Nicrophorus americanus</i>	Wherever found, except where listed as an experimental population	\$10,000	\$0	\$10,000
		Beetle, Hungerford's crawling water	<i>Brychius hungerfordi</i>	Wherever found	\$1,000	\$0	\$1,000
		Beetle, valley elderberry longhorn	<i>Desmocerus californicus dimorphus</i>	Wherever found	\$7,812	\$110,000	\$117,812
		Bumble bee, Rusty patched	<i>Bombus affinis</i>	Wherever found	\$9,300	\$0	\$9,300
		Butterfly, Fender's blue	<i>Icaricia icarioides fenderi</i>	Wherever found	\$300	\$0	\$300
		Butterfly, Mitchell's satyr	<i>Neonympha mitchellii mitchellii</i>	Wherever found	\$400	\$0	\$400

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Checkerspot, Taylor's (=whulge)	<i>Euphydryas editha taylori</i>	Wherever found	\$3,500	\$0	\$3,500
		Moth, Blackburn's sphinx	<i>Manduca blackburni</i>	Wherever found	\$12,300	\$0	\$12,300
		Skipper, Dakota	<i>Hesperia dacetoe</i>	Wherever found	\$1,500	\$0	\$1,500
		Skipper, Laguna Mountains	<i>Pyrgus ruralis lagunae</i>	Wherever found	\$1,680	\$0	\$1,680
		skipperling, Poweshiek	<i>Oarisma poweshiek</i>	Wherever found	\$600	\$0	\$600
		Tiger beetle, Salt Creek	<i>Cicindela nevadica lincolniana</i>	Wherever found	\$9,200	\$0	\$9,200
	Insects subtotal				\$57,592	\$110,000	\$167,592
	Crustaceans	Fairy shrimp, vernal pool	<i>Branchinecta lynchi</i>	Wherever found	\$9,534	\$152,720	\$162,254
		Shrimp, California freshwater	<i>Syncaris pacifica</i>	Wherever found	\$4,956	\$0	\$4,956
		Tadpole shrimp, vernal pool	<i>Lepidurus packardi</i>	Wherever found	\$8,316	\$0	\$8,316
	Crustaceans subtotal				\$22,806	\$152,720	\$175,526
	Corals	Coral, elkhorn	<i>Acropora palmata</i>	Wherever found	\$40,000	\$0	\$40,000
		Coral, Lobed Star	<i>Orbicella annularis</i>	Wherever found	\$40,000	\$0	\$40,000
		Coral, Pillar	<i>Dendrogyra cylindrus</i>	Wherever found	\$40,000	\$0	\$40,000
		Coral, Rough Cactus	<i>Mycetophyllia ferox</i>	Wherever found	\$40,000	\$0	\$40,000
		Coral, staghorn	<i>Acropora cervicornis</i>	Wherever found	\$40,000	\$0	\$40,000
	Corals subtotal				\$200,000	\$0	\$200,000
	Flowering Plants	Alani	<i>Melicope adscendens</i>	Wherever found	\$2,000	\$0	\$2,000
		Alani	<i>Melicope knudsenii</i>	Wherever found	\$2,000	\$0	\$2,000
		Alani	<i>Melicope mucronulata</i>	Wherever found	\$2,000	\$0	\$2,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Butterfly plant, Colorado	<i>Gaura neomexicana</i> var. <i>coloradensis</i>	Wherever found	\$1,500	\$0	\$1,500
		Cactus, Bakersfield	<i>Opuntia treleasei</i>	Wherever found	\$420	\$0	\$420
		Cactus, San Rafael	<i>Pediocactus despainii</i>	Wherever found	\$13,500	\$0	\$13,500
		Cactus, Siler pincushion	<i>Pediocactus</i> (=Echinocactus,=Utahia) <i>sileri</i>	Wherever found	\$3,000	\$0	\$3,000
		Cactus, Wright fishhook	<i>Sclerocactus wrightiae</i>	Wherever found	\$4,000	\$0	\$4,000
		Catchfly, Spalding's	<i>Silene spaldingii</i>	Wherever found	\$100	\$0	\$100
		Chaffseed, American	<i>Schwalbea americana</i>	Wherever found	\$2,000	\$0	\$2,000
		Checker- mallow, Nelson's	<i>Sidalcea nelsoniana</i>	Wherever found	\$400	\$0	\$400
		Desert-parsley, Bradshaw's	<i>Lomatium bradshawii</i>	Wherever found	\$1,000	\$0	\$1,000
		Dropwort, Canby's	<i>Oxypolis canbyi</i>	Wherever found	\$5,000	\$0	\$5,000
		Howellia, water	<i>Howellia aquatilis</i>	Wherever found	\$100	\$0	\$100
		Ko`oko`olau	<i>Bidens micrantha</i> ssp. <i>kalealaha</i>	Wherever found	\$2,000	\$0	\$2,000
		Ladies'-tresses, Ute	<i>Spiranthes diluvialis</i>	Wherever found	\$11,000	\$0	\$11,000
		Lupine, Kincaid's	<i>Lupinus sulphureus</i> ssp. <i>kincaidii</i>	Wherever found	\$400	\$0	\$400
		Mahoe	<i>Alectryon macrococcus</i>	Wherever found	\$2,000	\$0	\$2,000

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
		Manzanita, Del Mar	Arctostaphylos glandulosa ssp. crassifolia	Wherever found	\$420	\$0	\$420
		Mehamehame	Flueggea neowawraea	Wherever found	\$2,000	\$0	\$2,000
		Milk-vetch, Shivwits	Astragalus ampullarioides	Wherever found	\$1,000	\$0	\$1,000
		Milkweed, Mead's	Asclepias meadii	Wherever found	\$1,000	\$0	\$1,000
		Navarretia, spreading	Navarretia fossalis	Wherever found	\$1,050	\$0	\$1,050
		Ohai	Sesbania tomentosa	Wherever found	\$2,000	\$0	\$2,000
		Orchid, western prairie fringed	Platanthera praeclara	Wherever found	\$14,000	\$0	\$14,000
		Penstemon, blowout	Penstemon haydenii	Wherever found	\$1,000	\$0	\$1,000
		Peppergrass, Slickspot	Lepidium papilliferum		\$300	\$0	\$300
		Sandalwood, Lanai (= `iliahi)	Santalum haleakalae var. lanaiense	Wherever found	\$2,000	\$0	\$2,000
		Sunburst, San Joaquin adobe	Pseudobahia peirsonii	Wherever found	\$420	\$0	\$420
		Wire-lettuce, Malheur	Stephanomeria malheurensis	Wherever found	\$100	\$0	\$100
		Woolly-threads, San Joaquin	Monolopia (=Lembertia) congdonii	Wherever found	\$1,260	\$0	\$1,260
	Flowering Plants subtotal				\$78,970	\$0	\$78,970
	Ferns and Allies	Ihi`ihi	Marsilea villosa	Wherever found	\$2,000	\$0	\$2,000
	Ferns and Allies subtotal				\$2,000	\$0	\$2,000
	Multi-Species				\$1,903,515	\$0	\$1,903,515
	Multi-Species subtotal				\$1,903,515	\$0	\$1,903,515

Locality	Group Name	Common Name	Scientific Name	Population	General Expenditures	Land Expenditures	Grand Total
Domestic subtotal					\$13,788,644	\$19,905,014	\$33,693,658
Total					\$13,788,644	\$19,905,014	\$33,693,658

Appendix C. Expenditures Reported by States for FY 2017

Through the Association of Fish and Wildlife Agencies, States voluntarily provided annual ESA expenditures directly into the Service's Web-based system. This report includes collective expenditure data from all responding State agencies.

States Provided Expenditure Data for FY 2017	States Did Not Provide Expenditure Data for FY 2017
Alabama Alaska Arizona Arkansas Colorado Florida Georgia Idaho Indiana Iowa Kansas Kentucky Louisiana Maine Massachusetts Minnesota Missouri Montana Nebraska New Hampshire New Jersey New York North Dakota Ohio Oklahoma Oregon Pennsylvania South Carolina South Dakota Tennessee Texas Utah Vermont Washington West Virginia Wyoming	American Samoa California Connecticut Delaware District of Columbia Guam Hawaii Illinois Maryland Michigan Mississippi Nevada New Mexico North Carolina Northern Mariana Islands Puerto Rico Rhode Island Virgin Islands Virginia Wisconsin

Appendix C. Expenditures Reported by the States for FY 2017

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
Mammals	Bat, gray -- Wherever found	<i>Myotis grisescens</i>	\$127,190	\$100,000
	Bat, Indiana -- Wherever found	<i>Myotis sodalis</i>	\$254,874	\$100,000
	Bat, lesser long-nosed -- Wherever found	<i>Leptonycteris curasoae yerbabuena</i>	\$19,855	\$0
	Bat, Mexican long-nosed -- Wherever found	<i>Leptonycteris nivalis</i>	\$41,155	\$0
	Bat, Northern long-eared -- Wherever found	<i>Myotis septentrionalis</i>	\$295,751	\$189,400
	Bat, Ozark big-eared -- Wherever found	<i>Corynorhinus (=Plecotus) townsendii ingens</i>	\$3,780	\$0
	Bat, Virginia big-eared -- Wherever found	<i>Corynorhinus (=Plecotus) townsendii virginianus</i>	\$1,500	\$0
	Bear, grizzly -- U.S.A., conterminous (lower 48) States, except where listed as an experimental population	<i>Ursus arctos horribilis</i>	\$629,465	\$0
	Bear, polar -- Wherever found	<i>Ursus maritimus</i>	\$73,270	\$0
	Bison, wood -- Wherever found	<i>Bison bison athabasca</i>	\$232,863	\$0
	Caribou, Woodland -- Selkirk Mountain population	<i>Rangifer tarandus caribou</i>	\$20,901	\$0
	Deer, Columbian white-tailed -- Columbia River DPS	<i>Odocoileus virginianus leucurus</i>	\$5,000	\$0
	Ferret, black-footed -- Wherever found, except where listed as an experimental population	<i>Mustela nigripes</i>	\$188,125	\$0
	Ferret, black-footed -- U.S.A. (WY and specified portions of AZ, CO, MT, SD, and UT, see 17.84(g)(9))	<i>Mustela nigripes</i>	\$526,370	\$0
	Jaguar -- Wherever found	<i>Panthera onca</i>	\$11,702	\$0
	Lynx, Canada -- Wherever Found in Contiguous U.S.	<i>Lynx canadensis</i>	\$247,135	\$0
	Manatee, West Indian -- Wherever found	<i>Trichechus manatus</i>	\$42,500	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Mouse, New Mexico meadow jumping -- Wherever found	<i>Zapus hudsonius luteus</i>	\$41,372	\$0
	Mouse, Preble's meadow jumping -- wherever found	<i>Zapus hudsonius preblei</i>	\$31,263	\$0
	Ocelot -- wherever found	<i>Leopardus (=Felis) pardalis</i>	\$8,609	\$0
	Pocket gopher, Olympia -- Wherever found	<i>Thomomys mazama pugetensis</i>	\$5,000	\$0
	Pocket gopher, Roy Prairie -- Wherever found	<i>Thomomys mazama glacialis</i>	\$5,000	\$0
	Pocket gopher, Tenino -- Wherever found	<i>Thomomys mazama tumuli</i>	\$5,000	\$0
	Pocket gopher, Yelm -- Wherever found	<i>Thomomys mazama yelmensis</i>	\$20,000	\$0
	Prairie dog, Utah -- Wherever found	<i>Cynomys parvidens</i>	\$70,000	\$530,000
	Pronghorn, Sonoran -- U.S.A. (AZ), Mexico	<i>Antilocapra americana sonoriensis</i>	\$816,763	\$0
	Puma (=cougar), eastern -- Wherever found	<i>Puma (=Felis) concolor cougar</i>	\$900	\$0
	Rabbit, Columbia Basin Pygmy -- Columbia Basin DPS	<i>Brachylagus idahoensis</i>	\$100,689	\$0
	Sea lion, Steller -- Western DPS	<i>Eumetopias jubatus</i>	\$1,139,184	\$0
	Seal, Ringed -- Wherever found	<i>Phoca (=Pusa) hispida hispida</i>	\$234,029	\$0
	Squirrel, Mount Graham red -- Wherever found	<i>Tamiasciurus hudsonicus grahamensis</i>	\$28,246	\$0
	Squirrel, Northern Idaho Ground -- Wherever found	<i>Uroditellus brunneus</i>	\$54,449	\$0
	Vole, Hualapai Mexican -- Wherever found	<i>Microtus mexicanus hualpaiensis</i>	\$1,000	\$0
	whale, beluga -- Cook Inlet DPS	<i>Delphinapterus leucas</i>	\$34,548	\$0
	Whale, bowhead -- Wherever found	<i>Balaena mysticetus</i>	\$401,045	\$0
	Whale, killer -- Southern Resident DPS	<i>Orcinus orca</i>	\$8,385	\$0
	Whale, North Atlantic Right -- Wherever found	<i>Eubalaena glacialis</i>	\$55,000	\$0
	Wolf, gray -- U.S.A.: All of AL, AR, CA, CO, CT, DE, FL, GA, IA, IN, IL, KS, KY, LA, MA, MD, ME, MI, MO, MS, NC, ND, NE, NH, NJ, NV, NY, OH, OK, PA, RI, SC, SD, TN, TX, VA, VT, WI, and WV; and portions of AZ, NM, OR, UT, and WA. Mexico.	<i>Canis lupus</i>	\$18,970	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Wolf, Mexican -- U.S.A. (portions of AZ and NM)see 17.84(k)	<i>Canis lupus baileyi</i>	\$360,775	\$0
Mammals subtotal			\$6,161,663	\$919,400
Birds	Bobwhite, masked (quail) -- Wherever found	<i>Colinus virginianus ridgwayi</i>	\$2,687	\$0
	Condor, California -- U.S.A. (specific portions of Arizona, Nevada, and Utah)	<i>Gymnogyps californianus</i>	\$252,240	\$0
	Crane, whooping -- Wherever found, except where listed as an experimental population	<i>Grus americana</i>	\$12,700	\$0
	Cuckoo, yellow-billed -- Western U.S. DPS	<i>Coccyzus americanus</i>	\$34,262	\$0
	Falcon, Northern Aplomado -- U.S.A (AZ, NM)	<i>Falco femoralis septentrionalis</i>	\$1,500	\$0
	Flycatcher, southwestern willow -- Wherever found	<i>Empidonax traillii extimus</i>	\$35,691	\$0
	Horned lark, streaked -- Wherever found	<i>Eremophila alpestris strigata</i>	\$2,000	\$0
	Knot, red -- Wherever found	<i>Calidris canutus rufa</i>	\$50,060	\$0
	Murrelet, marbled -- U.S.A. (CA, OR, WA)	<i>Brachyramphus marmoratus</i>	\$109,100	\$0
	Owl, Mexican spotted -- Wherever found	<i>Strix occidentalis lucida</i>	\$500	\$0
	Owl, northern spotted -- Wherever found	<i>Strix occidentalis caurina</i>	\$2,200	\$0
	Plover, piping -- [Great Lakes watershed DPS] - Great Lakes, watershed in States of IL, IN, MI, MN, NY, OH, PA, and WI and Canada (Ont.)	<i>Charadrius melodus</i>	\$9,947	\$0
	Plover, piping -- [Atlantic Coast and Northern Great Plains populations] - Wherever found, except those areas where listed as endangered.	<i>Charadrius melodus</i>	\$432,949	\$0
	Plover, western snowy -- Pacific Coast population DPS-U.S.A. (CA, OR, WA), Mexico (within 50 miles of Pacific coast)	<i>Charadrius nivosus nivosus</i>	\$36,200	\$0
	Rail, Yuma clapper -- Wherever found	<i>Rallus longirostris yumanensis</i>	\$56,771	\$0
	sage-grouse, Gunnison -- Wherever found	<i>Centrocercus minimus</i>	\$430,274	\$0
	Stork, wood -- AL, FL, GA, MS, NC, SC	<i>Mycteria americana</i>	\$30,000	\$0
	Tern, least -- interior pop.	<i>Sterna antillarum</i>	\$55,701	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Tern, roseate -- northeast U.S. nesting pop.	<i>Sterna dougallii dougallii</i>	\$42,000	\$0
	Vireo, Black-capped -- Wherever found	<i>Vireo atricapilla</i>	\$4,500	\$0
	Warbler, Kirtland's -- Wherever found	<i>Setophaga kirtlandii</i> (= <i>Dendroica kirtlandii</i>)	\$300	\$0
	Woodpecker, ivory-billed -- Wherever found	<i>Campephilus principalis</i>	\$200	\$0
	Woodpecker, red-cockaded -- Wherever found	<i>Picoides borealis</i>	\$142,100	\$0
Birds subtotal			\$1,743,882	\$0
Reptiles	Alligator, American -- Wherever found	<i>Alligator mississippiensis</i>	\$35,000	\$0
	gartersnake, narrow-headed -- Wherever found	<i>Thamnophis rufipunctatus</i>	\$125,480	\$0
	gartersnake, northern Mexican -- Wherever found	<i>Thamnophis eques megalops</i>	\$118,798	\$0
	Massasauga (=rattlesnake), eastern -- Wherever found	<i>Sistrurus catenatus</i>	\$11,500	\$0
	Rattlesnake, New Mexican ridge-nosed -- Wherever found	<i>Crotalus willardi obscurus</i>	\$3,950	\$0
	Sea turtle, green -- Central North Pacific DPS	<i>Chelonia mydas</i>	\$2,500	\$0
	Sea turtle, green -- Central West Pacific DPS	<i>Chelonia mydas</i>	\$5,000	\$0
	Sea turtle, green -- North Atlantic DPS	<i>Chelonia mydas</i>	\$2,110	\$0
	Sea turtle, Kemp's ridley -- Wherever found	<i>Lepidochelys kempii</i>	\$23,608	\$0
	Sea turtle, leatherback -- Wherever found	<i>Dermodochelys coriacea</i>	\$5,000	\$0
	Sea turtle, loggerhead -- Northwest Atlantic Ocean DPS	<i>Caretta caretta</i>	\$275,545	\$0
	Snake, eastern indigo -- Wherever found	<i>Drymarchon corais couperi</i>	\$150,000	\$3,000,000
	Tortoise, desert -- Wherever found, except AZ south and east of Colorado R., and Mexico	<i>Gopherus agassizii</i>	\$138,056	\$0
	Tortoise, gopher -- West of Mobile and Tombigbee Rivers	<i>Gopherus polyphemus</i>	\$6,500	\$0
	turtle, bog -- Wherever found, except GA, NC, SC, TN, VA	<i>Clemmys muhlenbergii</i>	\$103,258	\$0
	turtle, bog -- U.S.A. (GA, NC, SC, TN, VA)	<i>Clemmys muhlenbergii</i>	\$35,000	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Turtle, Plymouth Redbelly -- Wherever found	<i>Pseudemys rubriventris bangsi</i>	\$2,500	\$0
Reptiles subtotal			\$1,043,805	\$3,000,000
Amphibians	Frog, Chiricahua leopard -- Wherever found	<i>Rana chiricahuensis</i>	\$291,140	\$0
	Frog, Oregon spotted -- Wherever found	<i>Rana pretiosa</i>	\$24,480	\$0
	Hellbender, Ozark -- Wherever found	<i>Cryptobranchus alleganiensis bishopi</i>	\$17,900	\$0
	Salamander, Cheat Mountain -- Wherever found	<i>Plethodon nettingi</i>	\$7,336	\$0
	Salamander, frosted flatwoods -- Wherever found	<i>Ambystoma cingulatum</i>	\$30,000	\$0
	salamander, Reticulated flatwoods -- Wherever found	<i>Ambystoma bishopi</i>	\$30,000	\$0
	Salamander, Sonora tiger -- Wherever found	<i>Ambystoma tigrinum stebbinsi</i>	\$39,010	\$0
	Toad, Houston -- Wherever found	<i>Bufo houstonensis</i>	\$1,093	\$0
	Toad, Wyoming -- Wherever found	<i>Bufo hemiophrys baxteri</i>	\$6,778	\$0
Amphibians subtotal			\$447,737	\$0
Fishes	Bocaccio -- Puget Sound - Georgia Basin DPS	<i>Sebastes paucispinis</i>	\$139,822	\$0
	Catfish, Yaqui -- Wherever found	<i>Ictalurus pricei</i>	\$2,000	\$0
	Cavefish, Ozark -- Wherever found	<i>Amblyopsis rosae</i>	\$850	\$0
	Chub, bonytail -- Wherever found	<i>Gila elegans</i>	\$373,075	\$0
	Chub, Borax Lake -- Wherever found	<i>Gila boraxobius</i>	\$10,000	\$0
	Chub, Gila -- Wherever found	<i>Gila intermedia</i>	\$14,944	\$0
	Chub, humpback -- Wherever found	<i>Gila cypha</i>	\$383,921	\$0
	Chub, Sonora -- Wherever found	<i>Gila ditaenia</i>	\$2,000	\$0
	Chub, spotfin -- U.S.A. (AL, TN-specified portions of Shoal Creek; see 17.84(m)(1)(ii))	<i>Erimonax monachus</i>	\$15,000	\$0
	Chub, Virgin River -- Wherever found	<i>Gila seminuda (=robusta)</i>	\$478,500	\$0
	Chub, Yaqui -- Wherever found	<i>Gila purpurea</i>	\$2,600	\$0
	Dace, blackside -- Wherever found	<i>Phoxinus cumberlandensis</i>	\$500	\$0
	Dace, Foskett speckled -- Wherever found	<i>Rhinichthys osculus</i> ssp.	\$21,000	\$0
	Darter, amber -- Wherever found	<i>Percina antesella</i>	\$5,000	\$0
	Darter, boulder -- Wherever found	<i>Etheostoma wapiti</i>	\$15,000	\$0
	Darter, Cherokee -- Wherever found	<i>Etheostoma scotti</i>	\$5,000	\$40,000

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Darter, Cumberland -- Wherever found	Etheostoma susanae	\$2,000	\$0
	Darter, duskytail -- Wherever found	Etheostoma percnum	\$15,000	\$0
	Darter, Etowah -- Wherever found	Etheostoma etowahae	\$5,000	\$40,000
	Darter, goldline -- Wherever found	Percina aurolineata	\$5,000	\$0
	Darter, Kentucky arrow -- Wherever found	Etheostoma spilotum	\$500	\$0
	Darter, leopard -- Wherever found	Percina pantherina	\$27,960	\$0
	Darter, slackwater -- Wherever found	Etheostoma boschungii	\$17,000	\$0
	Darter, snail -- Wherever found	Percina tanasi	\$14,000	\$0
	Darter, yellowcheek -- Wherever found	Etheostoma moorei	\$5,600	\$0
	Eulachon -- Southern DPS	Thaleichthys pacificus	\$62,189	\$0
	Logperch, Conasauga -- Wherever found	Percina jenkinsi	\$5,000	\$0
	Madtom, Neosho -- Wherever found	Noturus placidus	\$1,500	\$0
	Madtom, pygmy -- U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.84(t)(1)(i))	Noturus stanauli	\$21,000	\$0
	Madtom, smoky -- Wherever found	Noturus baileyi	\$15,000	\$0
	Madtom, yellowfin -- U.S.A. (TN-specified portions of the Tellico River; see 17.84(e)(1)(ii))	Noturus flavipinnis	\$15,000	\$0
	Minnow, Devils River -- Wherever found	Dionda diaboli	\$45,304	\$0
	Minnow, loach -- Wherever found	Tiaroga cobitis	\$168,074	\$0
	Minnow, Rio Grande Silvery -- Wherever found, except where listed as an experimental population	Hybognathus amarus	\$14,785	\$0
	Pikeminnow (=squawfish), Colorado -- Wherever found, except where listed as an experimental population	Ptychocheilus lucius	\$113,317	\$0
	Pikeminnow (=squawfish), Colorado -- Salt and Verde R. drainages, AZ	Ptychocheilus lucius	\$162,000	\$0
	Pupfish, Comanche Springs -- Wherever found	Cyprinodon elegans	\$21,559	\$0
	Pupfish, desert -- Wherever found	Cyprinodon macularius	\$48,918	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	rockfish, canary -- Puget Sound - Georgia Basin DPS	Sebastes pinniger	\$162,910	\$0
	rockfish, yelloweye -- Puget Sound - Georgia Basin DPS	Sebastes ruberrimus	\$162,910	\$0
	Salmon, Chinook -- Snake River fall-run ESU	Oncorhynchus (=Salmo) tshawytscha	\$214,053	\$0
	Salmon, Chinook -- Snake River spring/summer-run ESU	Oncorhynchus (=Salmo) tshawytscha	\$1,203,575	\$0
	Salmon, Chinook -- Puget Sound ESU	Oncorhynchus (=Salmo) tshawytscha	\$5,428,723	\$0
	Salmon, Chinook -- Lower Columbia River ESU	Oncorhynchus (=Salmo) tshawytscha	\$862,859	\$0
	Salmon, Chinook -- Upper Willamette River ESU	Oncorhynchus (=Salmo) tshawytscha	\$267,583	\$0
	Salmon, Chinook -- Upper Columbia spring-run ESU	Oncorhynchus (=Salmo) tshawytscha	\$263,029	\$0
	Salmon, chum -- Columbia River ESU	Oncorhynchus keta	\$208,782	\$0
	Salmon, chum -- Hood Canal summer-run ESU	Oncorhynchus keta	\$212,667	\$0
	Salmon, coho -- Southern Oregon - Northern California Coast ESU	Oncorhynchus (=Salmo) kisutch	\$24,000	\$0
	Salmon, coho -- Oregon Coast ESU	Oncorhynchus (=Salmo) kisutch	\$1,275,000	\$0
	Salmon, coho -- Lower Columbia River ESU	Oncorhynchus (=Salmo) kisutch	\$977,280	\$0
	Salmon, sockeye -- Snake River ESU	Oncorhynchus (=Salmo) nerka	\$1,284,828	\$0
	Salmon, sockeye -- Ozette Lake ESU	Oncorhynchus (=Salmo) nerka	\$72,232	\$0
	Sculpin, Grotto -- Wherever found	Cottus specus	\$10,000	\$0
	Sculpin, pygmy -- Wherever found	Cottus paulus (=pygmaeus)	\$2,000	\$0
	Shiner, Arkansas River -- Arkansas River Basin (AR, KS, NM, OK, TX)	Notropis girardi	\$3,000	\$0
	Shiner, beautiful -- Wherever found	Cyprinella formosa	\$500	\$0
	Shiner, blue -- Wherever found	Cyprinella caerulea	\$5,000	\$0
	Shiner, Topeka -- Wherever found, except where listed as an experimental population	Notropis topeka (=tristis)	\$15,900	\$0
	Spikedace -- Wherever found	Meda fulgida	\$170,562	\$0
	Spinedace, Little Colorado -- Wherever found	Lepidomeda vittata	\$8,980	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Steelhead -- Snake River Basin DPS	<i>Oncorhynchus (=Salmo) mykiss</i>	\$1,767,150	\$0
	Steelhead -- Lower Columbia River DPS	<i>Oncorhynchus (=Salmo) mykiss</i>	\$571,280	\$0
	Steelhead -- Upper Columbia River DPS	<i>Oncorhynchus (=Salmo) mykiss</i>	\$310,630	\$0
	Steelhead -- Upper Willamette River DPS	<i>Oncorhynchus (=Salmo) mykiss</i>	\$262,000	\$0
	Steelhead -- Middle Columbia River DPS	<i>Oncorhynchus (=Salmo) mykiss</i>	\$853,069	\$0
	Steelhead -- Puget Sound DPS	<i>Oncorhynchus (=Salmo) mykiss</i>	\$1,301,804	\$0
	Steelhead -- Middle Columbia River DPS	<i>Oncorhynchus (=Salmo) mykiss</i>	\$21,694	\$0
	Sturgeon, Alabama -- Wherever found	<i>Scaphirhynchus suttkusi</i>	\$11,000	\$0
	Sturgeon, Atlantic -- New York Bight DPS	<i>Acipenser oxyrinchus oxyrinchus</i>	\$50,500	\$0
	Sturgeon, Atlantic -- South Atlantic DPS	<i>Acipenser oxyrinchus oxyrinchus</i>	\$25,000	\$0
	sturgeon, green -- Southern DPS	<i>Acipenser medirostris</i>	\$20,747	\$0
	Sturgeon (Gulf subspecies), Atlantic -- Wherever found	<i>Acipenser oxyrinchus (=oxyrhynchus) desotoi</i>	\$11,000	\$0
	Sturgeon, pallid -- Wherever found	<i>Scaphirhynchus albus</i>	\$78,000	\$0
	Sturgeon, shortnose -- Wherever found	<i>Acipenser brevirostrum</i>	\$146,000	\$0
	Sturgeon, white -- U.S.A. (ID, MT), Canada (B.C.), Kootenai R. system	<i>Acipenser transmontanus</i>	\$213,542	\$0
	Sucker, June -- Wherever found	<i>Chasmistes liorus</i>	\$585,000	\$0
	Sucker, Lost River -- Wherever found	<i>Deltistes luxatus</i>	\$35,000	\$0
	Sucker, razorback -- Wherever found	<i>Xyrauchen texanus</i>	\$617,257	\$0
	Sucker, shortnose -- Wherever found	<i>Chasmistes brevirostris</i>	\$35,000	\$0
	Sucker, Warner -- Wherever found	<i>Catostomus warnerensis</i>	\$61,000	\$0
	Sucker, Zuni bluehead -- Wherever found	<i>Catostomus discobolus yarrowi</i>	\$1,700	\$0
	Topminnow, Gila (incl. Yaqui) -- Wherever found	<i>Poeciliopsis occidentalis</i>	\$145,744	\$0
	Trout, Apache -- Wherever found	<i>Oncorhynchus apache</i>	\$56,900	\$0
	Trout, bull -- U.S.A., conterminous, lower 48 states	<i>Salvelinus confluentus</i>	\$1,100,258	\$0
	Trout, bull -- Clackamas River subbasin experimental population	<i>Salvelinus confluentus</i>	\$25,000	\$0
	Trout, Gila -- Wherever found	<i>Oncorhynchus gilae</i>	\$86,060	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	trout, Greenback Cutthroat -- Wherever found	<i>Oncorhynchus clarkii stomias</i>	\$97,649	\$0
	Trout, Lahontan cutthroat -- Wherever found	<i>Oncorhynchus clarkii henshawi</i>	\$84,000	\$0
	Woundfin -- Wherever found, except where listed as an experimental population	<i>Plagopterus argentissimus</i>	\$437,842	\$0
Fishes subtotal			\$24,854,959	\$80,000
Clams	Bankclimber, purple (mussel) -- Wherever found	<i>Elliptoideus sloatianus</i>	\$5,000	\$0
	Bean, Cumberland (pearlymussel) -- Wherever found; Except where listed as Experimental Populations	<i>Villosa trabalis</i>	\$20,000	\$0
	Bean, rayed -- Wherever found	<i>Villosa fabalis</i>	\$6,000	\$0
	Clubshell -- Wherever found; Except where listed as Experimental Populations	<i>Pleurobema clava</i>	\$11,000	\$0
	Clubshell, ovate -- Wherever found	<i>Pleurobema perovatum</i>	\$2,500	\$0
	Clubshell, southern -- Wherever found	<i>Pleurobema decisum</i>	\$3,000	\$0
	Combshell, Cumberlandian -- Wherever found; Except where listed as Experimental Populations	<i>Epioblasma brevidens</i>	\$20,000	\$0
	Combshell, Cumberlandian -- U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	<i>Epioblasma brevidens</i>	\$5,700	\$0
	Combshell, southern -- Wherever found	<i>Epioblasma penita</i>	\$1,000	\$0
	Combshell, upland -- Wherever found	<i>Epioblasma metastrata</i>	\$1,000	\$0
	Elktoe, Cumberland -- Wherever found	<i>Alasmidonta atropurpurea</i>	\$12,000	\$0
	Fanshell -- Wherever found	<i>Cyprogenia stegaria</i>	\$5,800	\$0
	fatmucket, Arkansas -- Wherever found	<i>Lampsilis powellii</i>	\$2,900	\$0
	Higgins eye (pearlymussel) -- Wherever found	<i>Lampsilis higginsii</i>	\$5,000	\$0
	Kidneyshell, fluted -- Wherever found	<i>Ptychobranhus subtentum</i>	\$5,700	\$0
	Kidneyshell, triangular -- Wherever found	<i>Ptychobranhus greenii</i>	\$2,000	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Lampmussel, Alabama -- AL; Free-Flowing Reach of the Tennessee River below the Wilson Dam, Colbert and Lauderdale Counties, AL	Lampsilis virescens	\$5,700	\$0
	Moccasinshell, Alabama -- Wherever found	Medionidus acutissimus	\$3,000	\$0
	Moccasinshell, Coosa -- Wherever found	Medionidus parvulus	\$1,000	\$0
	Moccasinshell, Gulf -- Wherever found	Medionidus penicillatus	\$2,500	\$0
	Moccasinshell, Ochlockonee -- Wherever found	Medionidus simpsonianus	\$1,000	\$0
	Moccasinshell, Suwannee -- Wherever found	Medionidus walkeri	\$5,000	\$0
	Mucket, Neosho -- Wherever found	Lampsilis rafinesqueana	\$100	\$0
	Mucket, pink (pearlymussel) -- Wherever found	Lampsilis abrupta	\$32,900	\$0
	Mussel, oyster -- U.S.A. (TN - specified portions of the French Broad and Holston Rivers; see 17.85(b)(1))	Epioblasma capsaeformis	\$5,700	\$0
	Mussel, scaleshell -- Wherever found	Leptodea leptodon	\$1,000	\$0
	Mussel, sheepsnose -- Wherever found	Plethobasus cyphus	\$6,800	\$0
	Mussel, snuffbox -- Wherever found	Epioblasma triquetra	\$7,600	\$0
	Pearlshell, Louisiana -- Wherever found	Margaritifera hembeli	\$900	\$0
	Pearlymussel, birdwing -- Wherever found; Except where listed as Experimental Populations	Lemiox rimosus	\$5,700	\$0
	Pearlymussel, dromedary -- Wherever found; Except where listed as Experimental Populations	Dromus dromas	\$7,700	\$0
	Pearlymussel, littlewing -- Wherever found	Pegias fabula	\$5,000	\$0
	Pigtoe, Georgia -- Wherever found	Pleurobema hanleyianum	\$3,000	\$0
	Pigtoe, oval -- Wherever found	Pleurobema pyriforme	\$5,000	\$0
	Pigtoe, rough -- Wherever found	Pleurobema plenum	\$2,800	\$0
	Pigtoe, southern -- Wherever found	Pleurobema georgianum	\$5,000	\$0
	Pimpleback, orangefoot (pearlymussel) -- Wherever found	Plethobasus cooperianus	\$2,000	\$0
	Pocketbook, fat -- Wherever found	Potamilus capax	\$17,350	\$0
	Pocketbook, finelined -- Wherever found	Lampsilis altilis	\$5,000	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Pocketbook, shinyrayed -- Wherever found	Lampsilis subangulata	\$5,000	\$0
	Pocketbook, speckled -- Wherever found	Lampsilis streckeri	\$1,200	\$0
	Purple Cat's paw (=Purple Cat's paw pearlymussel) -- Wherever found; Except where listed as Experimental Populations	Epioblasma obliquata obliquata	\$20,000	\$0
	Rabbitsfoot -- Wherever found	Quadrula cylindrica cylindrica	\$3,810	\$0
	Riffleshell, northern -- Wherever found	Epioblasma torulosa rangiana	\$5,000	\$0
	Riffleshell, tan -- Wherever found	Epioblasma florentina walkeri (=E. walkeri)	\$20,000	\$0
	Ring pink (mussel) -- Wherever found	Obovaria retusa	\$2,000	\$0
	Spectaclecase (mussel) -- Wherever found	Cumberlandia monodonta	\$17,700	\$0
	Spinymussel, Altamaha -- Wherever found	Elliptio spinosa	\$2,000	\$0
	Threeridge, fat (mussel) -- Wherever found	Amblema neislerii	\$4,000	\$0
	Wartyback, white (pearlymussel) -- Wherever found	Plethobasus cicatricosus	\$800	\$0
	Wedgemussel, dwarf -- Wherever found	Alasmidonta heterodon	\$8,441	\$0
Clams subtotal			\$330,301	\$0
Snails	Ambersnail, Kanab -- Wherever found	Oxyloma haydeni kanabensis	\$10,004	\$0
	Rocksnailed, round -- Wherever found	Leptoxis ampla	\$8,500	\$0
	Snail, Chittenango ovate amber -- Wherever found	Succinea chittenangoensis	\$2,000	\$0
	Snail, flat-spired three-toothed -- Wherever found	Triodopsis platysayoides	\$2,275	\$0
	springsnail, San Bernardino -- Wherever found	Pyrgulopsis bernardina	\$1,000	\$0
	Springsnail, Three Forks -- Wherever found	Pyrgulopsis trivialis	\$7,513	\$0
Snails subtotal			\$31,292	\$0
Insects	Beetle, American burying -- Wherever found, except where listed as an experimental population	Nicrophorus americanus	\$39,306	\$0
	Bumble bee, Rusty patched -- Wherever found	Bombus affinis	\$1,300	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Butterfly, Karner blue -- Wherever found	<i>Lycaeides melissa samuelis</i>	\$83,700	\$50,000
	Butterfly, Mitchell's satyr -- Wherever found	<i>Neonympha mitchellii mitchellii</i>	\$800	\$0
	Checkerspot, Taylor's (=whulge) -- Wherever found	<i>Euphydryas editha taylori</i>	\$22,000	\$0
	Skipper, Dakota -- Wherever found	<i>Hesperia dacotae</i>	\$30,505	\$0
	skipperling, Poweshiek -- Wherever found	<i>Oarisma poweshiek</i>	\$2,600	\$0
	Tiger beetle, Salt Creek -- Wherever found	<i>Cicindela nevadica lincolniana</i>	\$17,600	\$0
Insects subtotal			\$197,811	\$50,000
Crustaceans	Crayfish, cave -- Wherever found	<i>Cambarus aculabrum</i>	\$100	\$0
	Crayfish, cave -- Wherever found	<i>Cambarus zophonastes</i>	\$1,200	\$0
Crustaceans subtotal			\$1,300	\$0
Flowering Plants	Amphianthus, little -- Wherever found	<i>Amphianthus pusillus</i>	\$2,000	\$0
	Aster, Florida golden -- Wherever found	<i>Chrysopsis floridana</i>	\$3,500	\$0
	Barbara's Buttons, Mohr's -- Wherever found	<i>Marshallia mohrii</i>	\$4,000	\$0
	beardtongue, Parachute -- Wherever found	<i>Penstemon debilis</i>	\$19,550	\$0
	Beargrass, Britton's -- Wherever found	<i>Nolina brittoniana</i>	\$11,000	\$0
	Beauty, Harper's -- Wherever found	<i>Harperocallis flava</i>	\$18,000	\$0
	Bellflower, Brooksville -- Wherever found	<i>Campanula robinsiae</i>	\$3,500	\$0
	Birds-in-a-nest, white -- Wherever found	<i>Macbridea alba</i>	\$900	\$0
	Bladderpod, Dudley Bluffs -- Wherever found	<i>Lesquerella congesta</i>	\$1,108	\$0
	bladderpod, Missouri -- Wherever found	<i>Physaria filiformis</i>	\$500	\$0
	Bladderpod, Short's -- Wherever found	<i>Physaria globosa</i>	\$1,200	\$0
	Blazingstar, scrub -- Wherever found	<i>Liatris ohlingerae</i>	\$5,500	\$0
	Blue-star, Kearney's -- Wherever found	<i>Amsonia kearneyana</i>	\$500	\$0
	Bonamia, Florida -- Wherever found	<i>Bonamia grandiflora</i>	\$3,000	\$0
	Brickell-bush, Florida -- Wherever found	<i>Brickellia mosieri</i>	\$1,200	\$0
	Buckwheat, scrub -- Wherever found	<i>Eriogonum longiternum</i> var. spachii	\$12,000	\$0
	Bulrush, Northeastern -- Wherever found	<i>Scirpus ancistrochaetus</i>	\$852	\$0
	Bush-clover, prairie -- Wherever found	<i>Lespedeza leptostachya</i>	\$1,300	\$0
	Butterfly plant, Colorado -- Wherever found	<i>Gaura neomexicana</i> var. <i>coloradensis</i>	\$100	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Butterwort, Godfrey's -- Wherever found	Pinguicula ionantha	\$1,400	\$0
	Cactus, Acuna -- Wherever found	Echinomastus erectocentrus var. acunensis	\$2,500	\$0
	Cactus, Arizona hedgehog -- Wherever found	Echinocereus triglochmatus var. arizonensis	\$2,000	\$0
	Cactus, black lace -- Wherever found	Echinocereus reichenbachii var. albertii	\$25,936	\$0
	Cactus, Brady pincushion -- Wherever found	Pediocactus bradyi	\$2,500	\$0
	Cactus, Cochise pincushion -- Wherever found	Coryphantha robbinsiorum	\$500	\$0
	Cactus, Colorado hookless -- Wherever found	Sclerocactus glaucus	\$1,518	\$0
	Cactus, Fickeisen plains -- Wherever found	Pediocactus peeblesianus fickeiseniae	\$2,000	\$0
	Cactus, Florida semaphore -- Wherever found	Consolea corallicola	\$600	\$0
	Cactus, Key tree -- Wherever found	Pilosocereus robinii	\$12,000	\$0
	Cactus, Knowlton's -- Wherever found	Pediocactus knowltonii	\$3,125	\$0
	Cactus, Nichol's Turk's head -- Wherever found	Echinocactus horizontalis var. nicholii	\$500	\$0
	Cactus, Peebles Navajo -- Wherever found	Pediocactus peeblesianus var. peeblesianus	\$2,000	\$0
	Cactus, Pima pineapple -- Wherever found	Coryphantha scheeri var. robustispina	\$3,500	\$0
	Cactus, Siler pincushion -- Wherever found	Pediocactus (=Echinocactus, =Utama) sileri	\$1,500	\$0
	Campion, fringed -- Wherever found	Silene polypetala	\$6,700	\$0
	Catchfly, Spalding's -- Wherever found	Silene spaldingii	\$46,970	\$0
	Chaffseed, American -- Wherever found	Schwalbea americana	\$2,500	\$0
	Cliffrose, Arizona -- Wherever found	Purshia (=Cowania) subintegra	\$500	\$0
	Clover, running buffalo -- Wherever found	Trifolium stoloniferum	\$30,784	\$0
	Coneflower, smooth -- Wherever found	Echinacea laevigata	\$5,000	\$0
	Cycladenia, Jones -- Wherever found	Cycladenia humilis var. jonesii	\$500	\$0
	Dropwort, Canby's -- Wherever found	Oxypolis canbyi	\$15,000	\$25,000
	Flax, Carter's small-flowered -- Wherever found	Linum carteri carteri	\$700	\$0
	Flax, sand -- Wherever found	Linum arenicola	\$1,300	\$0
	Fleabane, Zuni -- Wherever found	Erigeron rhizomatus	\$500	\$0
	Four-o'clock, MacFarlane's -- Wherever found	Mirabilis macfarlanei	\$4,983	\$0
	Fringe-tree, pygmy -- Wherever found	Chionanthus pygmaeus	\$3,000	\$0
	Gerardia, sandplain -- Wherever found	Agalinis acuta	\$650	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Goldenrod, Short's -- Wherever found	<i>Solidago shortii</i>	\$1,000	\$0
	Gooseberry, Miccosukee -- Wherever found	<i>Ribes echinellum</i>	\$200	\$0
	Gourd, Okeechobee -- Wherever found	<i>Cucurbita okeechobeensis</i> ssp. <i>okeechobeensis</i>	\$200	\$0
	Grass, Tennessee yellow-eyed -- Wherever found	<i>Xyris tennesseensis</i>	\$5,000	\$0
	Harebells, Avon Park -- Wherever found	<i>Crotalaria avonensis</i>	\$2,000	\$0
	Harperella -- Wherever found	<i>Ptilimnium nodosum</i>	\$22,417	\$0
	Howellia, water -- Wherever found	<i>Howellia aquatilis</i>	\$5,169	\$0
	Hypericum, highlands scrub -- Wherever found	<i>Hypericum cumulicola</i>	\$7,500	\$0
	Jacquemontia, beach -- Wherever found	<i>Jacquemontia reclinata</i>	\$5,000	\$0
	ladies-tresses, Canelo Hills -- Wherever found	<i>Spiranthes delitescens</i>	\$1,500	\$0
	Ladies'-tresses, Ute -- Wherever found	<i>Spiranthes diluvialis</i>	\$1,208	\$0
	Lead-plant, Crenulate -- Wherever found	<i>Amorpha crenulata</i>	\$8,000	\$0
	Leather flower, Alabama -- Wherever found	<i>Clematis socialis</i>	\$2,000	\$0
	Lousewort, Furbish -- Wherever found	<i>Pedicularis furbishiae</i>	\$6,500	\$0
	Lupine, scrub -- Wherever found	<i>Lupinus aridorum</i>	\$21,000	\$0
	mallow, Gierisch -- Wherever found	<i>Sphaeralcea gierischii</i>	\$1,500	\$0
	Meadowrue, Cooley's -- Wherever found	<i>Thalictrum cooleyi</i>	\$2,200	\$0
	Milkpea, Small's -- Wherever found	<i>Galactia smallii</i>	\$700	\$0
	Milk-vetch, Holmgren -- Wherever found	<i>Astragalus holmgreniorum</i>	\$1,500	\$0
	Milk-vetch, Jesup's -- Wherever found	<i>Astragalus robbinsii</i> var. <i>jesupi</i>	\$13,000	\$0
	Milk-vetch, Sentry -- Wherever found	<i>Astragalus cremnophylax</i> var. <i>cremnophylax</i>	\$1,000	\$0
	Milkweed, Mead's -- Wherever found	<i>Asclepias meadii</i>	\$10,800	\$0
	Milkweed, Welsh's -- Wherever found	<i>Asclepias welshii</i>	\$1,000	\$0
	Mint, Garrett's -- Wherever found	<i>Dicerandra christmanii</i>	\$700	\$0
	Mint, Lakela's -- Wherever found	<i>Dicerandra immaculata</i>	\$12,000	\$0
	Mint, longspurred -- Wherever found	<i>Dicerandra cornutissima</i>	\$6,000	\$0
	Mint, scrub -- Wherever found	<i>Dicerandra frutescens</i>	\$1,000	\$0
	Monkshood, northern wild -- Wherever found	<i>Aconitum noveboracense</i>	\$500	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Mustard, Carter's -- Wherever found	Warea carteri	\$4,500	\$0
	No common name -- Wherever found	Geocarpon minimum	\$400	\$0
	Orchid, eastern prairie fringed -- Wherever found	Platanthera leucophaea	\$3,100	\$0
	Orchid, western prairie fringed -- Wherever found	Platanthera praeclara	\$1,800	\$0
	Orchid, white fringeless --	Platanthera integrilabia	\$4,000	\$0
	Pawpaw, beautiful -- Wherever found	Deeringothamnus pulchellus	\$5,000	\$0
	Pawpaw, four-petal -- Wherever found	Asimina tetramera	\$900	\$0
	Pawpaw, Rugel's -- Wherever found	Deeringothamnus rugelii	\$1,500	\$0
	Pea, Big Pine partridge -- Wherever found	Chamaecrista lineata keyensis	\$700	\$0
	Penstemon, blowout -- Wherever found	Penstemon haydenii	\$10,100	\$0
	Peppergrass, Slickspot --	Lepidium papilliferum	\$45,040	\$0
	Phacelia, DeBeque -- Wherever found	Phacelia submutica	\$1,108	\$0
	Pigeon wings -- Wherever found	Clitoria fragrans	\$6,000	\$0
	Pinkroot, gentian -- Wherever found	Spigelia gentianoides	\$7,500	\$0
	Pink, swamp -- Wherever found	Helonias bullata	\$6,000	\$0
	Plum, scrub -- Wherever found	Prunus geniculata	\$5,000	\$0
	Pogonia, small whorled -- Wherever found	Isotria medeoloides	\$6,200	\$0
	Polygala, Lewton's -- Wherever found	Polygala lewtonii	\$17,500	\$0
	Polygala, tiny -- Wherever found	Polygala smallii	\$200	\$0
	Pondberry -- Wherever found	Lindera melissifolia	\$5,300	\$0
	Prickly-apple, aboriginal -- Wherever found	Harrisia (=Cereus) aboriginum (=gracilis)	\$200	\$0
	Prickly-apple, fragrant -- Wherever found	Cereus eriophorus var. fragrans	\$700	\$0
	Ragwort, San Francisco Peaks -- Wherever found	Packera franciscana	\$500	\$0
	Rattleweed, hairy -- Wherever found	Baptisia arachnifera	\$20,000	\$0
	Rhododendron, Chapman -- Wherever found	Rhododendron chapmanii	\$400	\$0
	Rockcress, Georgia -- Wherever found	Arabis georgiana	\$2,000	\$0
	rock cress, Shale barren -- Wherever found	Arabis serotina	\$6,683	\$0
	Rose-mallow, Neches River -- Wherever found	Hibiscus dasycalyx	\$18,671	\$0
	Rosemary, Apalachicola -- Wherever found	Conradina glabra	\$12,500	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
	Rosemary, Etonia -- Wherever found	<i>Conradina etonia</i>	\$6,500	\$0
	Rosemary, short-leaved -- Wherever found	<i>Conradina brevifolia</i>	\$6,000	\$0
	Sandlace -- Wherever found	<i>Polygonella myriophylla</i>	\$3,500	\$0
	Seagrass, Johnson's -- Wherever found	<i>Halophila johnsonii</i>	\$200	\$0
	Sedge, Navajo -- Wherever found	<i>Carex specuicola</i>	\$500	\$0
	silverbush, Blodgett's -- Wherever found	<i>Argythamnia blodgettii</i>	\$5,000	\$0
	Skullcap, Florida -- Wherever found	<i>Scutellaria floridana</i>	\$400	\$0
	Skullcap, large-flowered -- Wherever found	<i>Scutellaria montana</i>	\$10,000	\$0
	Skyrocket, Pagosa -- Wherever found	<i>Ipomopsis polyantha</i>	\$31,530	\$0
	Snakeroot -- Wherever found	<i>Eryngium cuneifolium</i>	\$900	\$0
	Spiraea, Virginia -- Wherever found	<i>Spiraea virginiana</i>	\$10,325	\$0
	Spurge, deltoid -- Wherever found	<i>Chamaesyce deltoidea</i> ssp. <i>deltoidea</i>	\$200	\$0
	Spurge, Garber's -- Wherever found	<i>Chamaesyce garberi</i>	\$3,500	\$0
	Spurge, telephus -- Wherever found	<i>Euphorbia telephioides</i>	\$7,000	\$0
	Spurge, wedge -- Wherever found	<i>Chamaesyce deltoidea serpyllum</i>	\$700	\$0
	Sumac, Michaux's -- Wherever found	<i>Rhus michauxii</i>	\$3,000	\$0
	Sunflower, whorled --	<i>Helianthus verticillatus</i>	\$5,000	\$0
	Thistle, Pitcher's -- Wherever found	<i>Cirsium pitcheri</i>	\$500	\$0
	Thoroughwort, Cape Sable -- Wherever found	<i>Chromolaena frustrata</i>	\$2,000	\$0
	Trillium, persistent -- Wherever found	<i>Trillium persistens</i>	\$6,000	\$0
	Trillium, relict -- Wherever found	<i>Trillium reliquum</i>	\$3,000	\$0
	Twinpod, Dudley Bluffs -- Wherever found	<i>Physaria obcordata</i>	\$1,108	\$0
	Warea, wide-leaf -- Wherever found	<i>Warea amplexifolia</i>	\$50,500	\$0
	Water-plantain, Kral's -- Wherever found	<i>Sagittaria secundifolia</i>	\$500	\$0
	Water-umbel, Huachuca -- Wherever found	<i>Lilaeopsis schaffneriana</i> var. <i>recurva</i>	\$2,500	\$0
	Water-willow, Cooley's -- Wherever found	<i>Justicia cooleyi</i>	\$2,000	\$0
	Whitlow-wort, papery -- Wherever found	<i>Paronychia chartacea</i>	\$2,000	\$0
	wild buckwheat, clay-loving -- Wherever found	<i>Eriogonum pelinophilum</i>	\$6,808	\$0
	Wireweed -- Wherever found	<i>Polygonella basiramia</i>	\$1,700	\$0
	Ziziphus, Florida -- Wherever found	<i>Ziziphus celata</i>	\$31,000	\$0

Group Name	Common Name	Scientific Name	General Expenditures	Land Expenditures
Flowering Plants subtotal			\$808,343	\$25,000
Conifers and Cycads	Torreya, Florida -- Wherever found	Torreya taxifolia	\$8,500	\$0
Conifers and Cycads subtotal			\$8,500	\$0
Ferns and Allies	fern, Florida bristle -- Wherever found	Trichomanes punctatum ssp. floridanum	\$700	\$0
	Quillwort, black spored -- Wherever found	Isoetes melanospora	\$2,000	\$0
	Quillwort, mat-forming -- Wherever found	Isoetes tegetiformans	\$3,000	\$0
Ferns and Allies subtotal			\$5,700	\$0
Lichens	Cladonia, Florida perforate -- Wherever found	Cladonia perforata	\$1,900	\$0
Lichens subtotal			\$1,900	\$0
Multi-Species Expenditures			\$7,560,418	\$383,300
Total			\$43,197,611	\$4,457,700