

Land Protection Plan

San Luis Valley Conservation Area

Colorado and New Mexico

December 2015

Prepared by

San Luis Valley National Wildlife Refuge Complex
8249 Emperius Road
Alamosa, CO 81101
719 / 589 4021

U.S. Fish and Wildlife Service
Region 6, Mountain-Prairie Region
Branch of Refuge Planning
134 Union Boulevard, Suite 300
Lakewood, CO 80228
303 / 236 8145

CITATION for this document:

U.S. Fish and Wildlife Service. 2015. Land protection plan for the San Luis Valley Conservation Area. Lakewood, CO: U.S. Department of the Interior, U.S. Fish and Wildlife Service. 151 p.

In accordance with the National Environmental Policy Act and U.S. Fish and Wildlife Service policy, an environmental assessment and land protection plan have been prepared to analyze the effects of establishing the San Luis Valley Conservation Area in southern Colorado and northern New Mexico.

The environmental assessment (appendix A) analyzes the environmental effects of establishing the San Luis Valley Conservation Area.

The San Luis Valley Conservation Area land protection plan describes the priorities for acquiring up to 250,000 acres through voluntary conservation easements and up to 30,000 acres in fee title.

Note: Information contained in the maps is approximate and does not represent a legal survey. Ownership information may not be complete.

Contents

Abbreviations	vii
Chapter 1—Introduction and Project Description	1
Purpose of the San Luis Valley Conservation Area	2
<i>Vision for the San Luis Valley National Wildlife Refuge Complex</i>	4
<i>Purpose of the Alamosa and Monte Vista National Wildlife Refuges</i>	4
<i>Purpose of the Baca national wildlife refuge</i>	4
<i>Purpose of the Sangre de Cristo Conservation Area</i>	4
Issues Identified and Selected for Analysis	4
Public Review of and Comments on the Draft EA and LPP	5
National Wildlife Refuge System and Authorities	6
Related Actions and Activities	7
<i>San Luis Valley National Wildlife Refuge Complex</i>	7
<i>U.S. Fish and Wildlife Service—Partners for Fish and Wildlife Program</i>	7
<i>USDA Forest Service</i>	7
<i>Bureau of Land Management</i>	7
<i>National Park Service</i>	9
<i>Natural Resources Conservation Service</i>	9
<i>State of Colorado</i>	9
<i>Land Trusts</i>	9
Habitat Protection and the Easement Acquisition Process	9
Conservation Easements and Other Acquisitions	10
Chapter 2—Area Description and Resources	11
Physical Environment	11
<i>Geology</i>	11
<i>Minerals</i>	11
<i>Water and Hydrology</i>	12
<i>Climate</i>	14
Biological Environment	14
<i>Plant Communities</i>	14
<i>Wildlife</i>	17
Cultural Resources	21
<i>Prehistory</i>	21
<i>Early History</i>	23
<i>Political Boundaries, Land Grants, and Public Lands</i>	25
<i>Native Peoples</i>	26
<i>Settlement</i>	26
<i>Summary of Known Historic Resources</i>	28
Socioeconomic Environment	28
<i>Socioeconomic Profile</i>	28
<i>Land Use and Ownership Changes Surrounding the conservation area</i>	30
<i>San Luis Valley conservation area land conservation efforts</i>	34
<i>Water Law</i>	36

Chapter 3—Threats to and Status of Resources	39
Threats to Resources	39
<i>Development</i>	39
<i>Fragmentation</i>	39
<i>Invasive Species</i>	40
<i>Water Resources</i>	40
<i>Cultural Resources</i>	41
<i>Climate Change</i>	41
Effects on the Natural and Human Environment	42
Chapter 4—Project Implementation	43
Land Protection Choices	43
<i>No Action</i>	43
<i>Conservation easements and limited fee-title acquisition (preferred alternative)</i>	43
Project Objectives and Actions	44
<i>Collaboration with Land Protection Organizations</i>	44
<i>Easement Terms and Requirements</i>	44
<i>Contaminants or Hazardous Materials</i>	45
<i>Acquisition money</i>	45
Incorporating Science and Strategic Habitat Conservation in the SLVCA	45
<i>Strategic Growth of the National Wildlife Refuge System</i>	45
<i>Strategic Habitat Conservation</i>	50
Protection Priorities	51
<i>Species-Habitat Mapping Methodology</i>	51
<i>Landscape Prioritization</i>	57
<i>Evaluation of Easement Potential</i>	61
Socioeconomic Considerations	61
Public Involvement and Coordination	61
<i>Scoping</i>	61
<i>Public Review of the Draft Environmental Assessment and Land Protection Plan</i>	62
Distribution and Availability	62
Glossary	63
Appendix A	65
Appendix B	79
Appendix C	81
Appendix D	91
Appendix E	133
Literature Cited	145

FIGURES

Figure 1. Overview map of the general land cover of the San Luis Valley Conservation Area in Colorado and New Mexico 3

Figure 2. Map of the San Luis Valley Conservation Area, which will be part of a broader network of public and private conservation lands in Colorado and New Mexico 8

Figure 3. Map of the San Luis Valley and Sangre de Cristo Conservation Areas in Colorado and New Mexico, which capture the upper headwaters of the Rio Grande, the fourth longest river in the United States. 13

Figure 4. Spatial graphic of predicted distribution of the New Mexico meadow jumping mouse in the 20th century and the late 21st century in the San Luis Valley Conservation Area, Colorado and New Mexico 47

Figure 5. Map of southwestern willow flycatcher and Canada lynx habitat in the San Luis Valley Conservation Area, Colorado and New Mexico 52

Figure 6. Map of Rio Grande cutthroat trout habitat in the San Luis Valley Conservation Area, Colorado and New Mexico 54

Figure 7. Map of present Gunnison sage-grouse habitat in and around the San Luis Valley Conservation Area, Colorado and New Mexico, with historical range indicated for reference 55

Figure 8. Map of Wilson’s phalarope habitat in the San Luis Valley Conservation Area, Colorado and New Mexico . . . 56

Figure 9. Map of American bittern habitat in the San Luis Valley Conservation Area, Colorado and New Mexico . . . 58

Figure 10. Map of sage thrasher habitat in the San Luis Valley Conservation Area, Colorado and New Mexico 59

Figure 11. Map of spatial conservation prioritization within the San Luis Valley Conservation Area, Colorado and New Mexico 60

TABLES

Table 1. Population statistics for counties in Colorado and New Mexico that contain the San Luis Valley Conservation Area. 29

Table 2. Income, education, unemployment, and poverty rates for counties in Colorado and New Mexico that contain the San Luis Valley Conservation Area. 30

Table 3. Percentage employment by sector for counties in Colorado and New Mexico that contain the San Luis Valley region 31

Table 4. Resident, migratory, and breeding birds of concern within the San Luis Valley Conservation Area, Colorado and New Mexico, with their status and objectives from a relevant continental or regional plan or list 49

Abbreviations

BLM	Bureau of Land Management
CCP	Comprehensive conservation plan
CFR	Code of Federal Regulations
DOE	Department of Energy
EA	Environmental assessment
LPP	Land protection plan
NASA	National Aeronautics and Space Administration
NEPA	National Environmental Policy Act
NPS	National Park Service
NRCS	Natural Resource Conservation Service
SCCA	Sangre de Cristo Conservation Area
Service	U.S. Fish and Wildlife Service
SLVCA	San Luis Valley Conservation Area
U.S.C.	United States Code
USFS	USDA Forest Service
USFWS	U.S. Fish and Wildlife Service
USGS	U.S. Geological Survey

Chapter 1—Introduction and Project Description

In short, this view combined the sublime and beautiful: the great and lofty mountains covered with eternal snows, seemed to surround the luxuriant vale, crowned with perennial flowers, like a terrestrial paradise, shut out from the view of man.

—Captain Zebulon Pike, on a hill overlooking the San Luis Valley, February 5, 1807

Through the San Luis Valley Conservation Area (SLVCA), we, the U.S. Fish and Wildlife Service (Service) will join our partners in protecting the remarkable ecological values and working landscapes of the high mountain desert that were so eloquently described during Pike’s Expedition to the Southwest in the early 19th century. The SLVCA is a landscape-level strategic habitat conservation initiative within the boundaries of the Southern Rockies Landscape Conservation Cooperative. The conservation area encompasses the headwaters of the Rio Grande in southern Colorado and a small part of northern New Mexico.

Despite substantial changes that have altered its landscape and ecology, the San Luis Valley is a remarkable North American example of the compatibility of wildlife and agriculture. Due to the low human population associated with the mostly agricultural economy,

the existing wetland, riparian, and upland habitats still retain a significant portion of their biological value, particularly for migratory birds. We hope to bring more resources and tools to the San Luis Valley to complement the ongoing efforts of other organizations to maintain and improve the relationship between agriculture and wildlife, especially migratory birds and endangered species.

The San Luis Valley is a large intermountain valley bounded by the San Juan and Sangre de Cristo Mountains, whose rain shadows result in high-desert conditions. However, the complex hydrology of the valley and the snowmelt runoff from the mountains have created a variety of dynamic wetlands and riparian corridors on the valley floor, resulting in a diverse assemblage of plants and wildlife. The valley provides habitat for many Federal trust species, including the southwestern willow flycatcher, western snowy plover, numerous species of migrating and nesting waterfowl, and 95 percent of the Rocky Mountain population of greater sandhill cranes.

Anthropogenic (human-caused) practices including agriculture, changes in fire regime, and climate change have resulted in substantial changes to the hydrology and historical vegetation of the San Luis Valley.



© Joe Zimm

The wetlands and fields of the San Luis Valley Conservation Area are an important stopover habitat for migrating sandhill cranes, an important tourist draw to the area.

In this desert environment, the timing and amount of water flowing from the adjoining mountains is the biggest influence on the environmental and economic conditions. The mid- to late-1800s brought extensive development of irrigation infrastructure to support a growing ranching and farming industry that relied on flood-irrigation techniques. This resulted in significant alteration in stream and river hydrology and the associated, naturally occurring, wetland habitats. However, the introduction of large amounts of irrigation water onto newly established and expanded hay meadows and crop fields provided substantial areas of habitat for migratory waterbirds that did not historically exist.

As the demand for surface water exceeded the supply, technology allowed development of ground water sources resulting in continued growth of the agriculture industry. By 1900, more than 1,000 irrigation wells had been drilled into the confined aquifer. Between 1930 and 1970, 6,000 more wells were drilled in both the confined and unconfined aquifers, allowing more land to be irrigated. These free-flowing wells are from only the confined aquifer and cover less than half the irrigated acreage. The amount of land under irrigation has not increased much, if at all, since the 1930s. Concerns about effects caused by stream depletions prompted the Colorado State Engineer in the early 1970s to issue a moratorium on new well development in the confined aquifer and all tributary wells except the Closed Basin. A similar moratorium was issued for the Closed Basin in 1981. The 1960s and 1970s saw dramatic conversion from flood-irrigation practices to the use of center-pivot sprinklers. This new technology offered substantial labor- and water-saving efficiencies but is of little value to migratory waterbirds except for the continued availability of waste grain eaten by geese and sandhill cranes.

Due to the hydrologic connection between surface and ground water, and the compounding effects brought by the chronic drought conditions of 2002 to present, wetland basins throughout the San Luis Valley are experiencing a very dry period. In 2003, farmers north of Alamosa began voluntary efforts to address well depletions. In 2004, the Colorado legislature provided guidelines for setting up management plans to regulate ground water use in the valley to ensure ground water sustainability in a way that does not harm senior water users. The State engineer adopted two rules, “Groundwater and Irrigation Season Rules for Water Division No. 3” and “Confined Aquifer New Use Rules for Division 3”, in 2015. Although the task has been challenging, the outcome will help the remaining agricultural operations and wetland and riparian habitats.

The San Luis Valley is lucky to have an active conservation community of local, State, Federal, and national organizations working to protect many of the same habitats valuable to migratory birds and

endangered species. As the cost of agriculture rises, so do the economic incentives to sell irrigation water rights and convert them to residential, industrial, and municipal uses, which increases the challenge of these organizations to keep water on the land to produce commodities and wildlife habitat.

The SLVCA will conserve a network of vital wildlife habitat through voluntary conservation easements and a limited amount of fee-title acquisition. The SLVCA acquisitions will focus on the protection of wetlands and associated uplands in the valley through the use of up to 250,000 acres of conservation easements. Up to an additional 30,000 acres of fee-title acquisition from willing sellers will be used (such as for boundary simplification and surface water rights acquisition) where it will benefit the management and objectives of the valley’s existing three Alamosa, Baca, and Monte Vista National Wildlife Refuges. We will acquire land in fee title only when Service objectives could not be accomplished with conservation easements. An overview of the project area is provided in figure 1.

The SLVCA will be the second phase of a larger conservation vision for the San Luis Valley that began in 2010. While the project was initially envisioned as a single broad conservation area, we recognized the need for accelerated planning in the southeastern portion of the project area, which resulted in the establishment of the Sangre de Cristo Conservation Area (SCCA) in 2012. The SLVCA encompasses the remainder of the original project area.

This document—a land protection plan (LPP)—sets out the basis, priorities, and general actions for the SLVCA. The environmental assessment (EA) in appendix A documents the analysis for the LPP that was conducted by the Service’s planning team (refer to “Appendix B—List of Preparers and Reviewers”).

The Service found that the SLVCA project would have no significant effect on the human environment; thus a finding of no significant impact was documented in “Appendix C—Environmental Compliance for the Land Protection Plan,” where other compliance documents and approvals are also found.

Purpose of the San Luis Valley Conservation Area

The purpose of the SLVCA is to protect Federal trust species and other plants and wildlife of the San Luis Valley while ensuring the long-term function and resilience of its diverse ecosystems. Acquisition and administration of the SLVCA will focus on protecting the land and water supporting the riparian areas, wetlands, and key uplands that complement and connect existing protected areas. This purpose is in alignment with, but does not supersede, the vision and statutory




Figure 1. Overview map of the general land cover of the San Luis Valley Conservation Area in Colorado and New Mexico.

purposes of the three existing refuges within the San Luis Valley National Wildlife Refuge Complex or the SCCA, as described below.



USFWS

The San Luis Valley Conservation Area contains a rich mosaic of working ranch lands and important wildlife habitat.

VISION FOR THE SAN LUIS VALLEY NATIONAL WILDLIFE REFUGE COMPLEX

The San Luis Valley National Wildlife Refuge Complex, set in a high expansive desert valley, is cradled between the snowcapped peaks of the San Juan and Sangre de Cristo Ranges. Mountain snowmelt feeds the Rio Grande, numerous streams, and a dynamic ground water system, creating a diverse mix of playas, wet meadows, and willow and cottonwood riparian corridors that are in stark contrast with the surrounding arid landscape. As reflected by 12,000 years of human history in the valley, the San Luis Valley National Wildlife Refuge Complex attracts many people. Visitors experience the ancient song of the sandhill crane, witness evening flights of thousands of waterfowl, and listen to bugling elk. Through ever changing conditions like climate change, the refuges support and foster a collaborative spirit between their neighbors and partners to conserve the valley's treasured resources.

PURPOSE OF THE ALAMOSA AND MONTE VISTA NATIONAL WILDLIFE REFUGES

The Alamosa and Monte Vista National Wildlife Refuges were established under the authority of the Migratory Bird Conservation Act “for use as inviolate sanctuaries, or for any other management purpose, for migratory birds.”

PURPOSE OF THE BACA NATIONAL WILDLIFE REFUGE

The purpose of the Baca National Wildlife Refuge shall be to restore, enhance, and maintain wetland, upland, riparian, and other habitats for native wildlife, plant, and fish species in the San Luis Valley. In administering the Baca National Wildlife Refuge, the Secretary shall, to the maximum extent practicable (A) emphasize migratory bird conservation; (B) take into consideration the role of the refuge in broader landscape conservation efforts; and (C) subject to any agreement in existence as of the date of enactment of this paragraph, and to the extent consistent with the purposes of the refuge, use decreed water rights on the refuge in approximately the same manner that the water rights have been used historically.

PURPOSE OF THE SANGRE DE CRISTO CONSERVATION AREA

The purpose of the SCCA is to protect the high-elevation wildlife habitat of the Sangre de Cristo Mountains and the uplands of the southeastern San Luis Valley, with an emphasis on migratory birds and imperiled species. Acquisitions within, and administration of, the SCCA will focus on promoting the adaptive capacity and resilience of these ecosystems by ensuring connectivity between existing protected areas and by protecting wildlife movement corridors, particularly riparian areas.

Issues Identified and Selected for Analysis

We solicited comments about the SLVCA from the public through direct mailings, news releases, public meetings, and direct contacts. These comments were incorporated into this land protection plan.

- On March 15, 2011, the Service opened a scoping period for the public with the publication of a notice of intent in the Federal Register (FR Doc. 2011–5924). The notice of intent notified the public of the Service’s intention to begin the coplanning and NEPA review for the CCP and LPP for the San Luis Valley National Wildlife Refuge Complex.

- Public scoping meetings were held on March 29, 2011, in Alamosa, Colorado; March 30, 2011, in Monte Vista, Colorado; and March 31, 2011, in Moffat, Colorado. The scoping meetings were attended by approximately 50 people, many of whom provided input for the scoping process. Additionally, 14 written comments were received from organizations and members of the public.
- A press event and public meeting was held at Adams State College in Alamosa, Colorado, on January 4, 2012, at which the Secretary of the Interior, Ken Salazar, organized the presentation of several complementary initiatives for the San Luis Valley and Sangre de Cristo Mountains. One of these initiatives was landscape scale conservation, which Dan Ashe, the Director of the Service, presented as being embodied by the SLVCA. Questions were answered and comments taken at a breakout session following the main meeting.
- The project's planning Web site (www.fws.gov/alamosa/planning/lpp/co/slv/slv.html) was established in early March of 2011. The site provides information about meetings and downloadable versions of public documents. Individuals can also sign up to be on the project mailing list through the Web site.

During scoping, the CCP and LPP were still being planned simultaneously. However, the two plans have since been separated and the LPP process has been moved up to take advantage of conservation opportunities that may not exist in the future. As such, many of the issues identified during scoping are not specific or relevant to the LPP. The applicable issues and questions identified during the scoping process and during internal conversations among the SLVCA planning team follow:

- The SLVCA must protect the wildlife habitat, specifically wetlands, riparian corridors, grasslands, and shrublands, of the San Luis Valley, while also supporting the rural agricultural qualities that define the region.
- What role can the conservation area play in protecting listed species and species of concern?
- How will the SLVCA affect water use in the valley?
- The SLVCA should not negatively affect private property rights in the valley.
- The Service needs to build on existing partnerships for land protection.
- How will the public be able to use lands protected under the SLVCA?
- The Service needs to make sure that the SLVCA planning process incorporates the importance of protecting cultural resources.
- How will the SLVCA increase the capacity to adapt to climate change on the existing refuges and habitat throughout the valley?
- How will the SLVCA protect the water resources of the San Luis Valley from efforts to move water to other areas of the State or elsewhere?
- The plan should account for air, soil, sound, and visibility effects.

Public Review of and Comments on the Draft EA and LPP

This final LPP is based on the draft EA and LPP for the SLVCA, which the Service released for comment on May 9, 2012 for a 30-day public review period. The draft documents were made available to the public via the project Web site, as well as to Federal officials and agencies, State officials and agencies, 17 Native American tribes with aboriginal interests, and members of the public who had asked to be added to the project mailing list. In addition, three public meetings were held in Alamosa, San Luis, and Moffat, Colorado on May 14, 15, and 16, 2012, respectively. Approximately 50 members of the public attended these three meetings. In addition to several comments voiced at these public meetings, the Service received eight written comments from government agencies and nongovernmental organizations and corporations, and six written comments from individuals. Comments were reviewed and incorporated into the administrative record.

The substantive comments were published with Service responses in appendix D of the final LPP for the SCCA, which was published in August 2012. Refer to the online document at http://www.fws.gov/mountain-prairie/refuges/lpp_PDFs/sdc_lpp_appendix.pdf.

Of those comments that showed a clear opinion in favor or opposed to the project, 75 percent were supportive in nature. The comments generally in favor of the SLVCA mention items such as the following:

- The SLVCA is entirely a willing-seller program, not an imposition.
- The Service has emphasized the collaborative nature of the project. The SLVCA is one of many conservation initiatives.
- The plan was comprehensive.

- The Service’s conservation goals are complementary with those of residents in Crestone and Baca Grande who would like to see perpetual conservation easements in those municipalities.
- The prioritization strategy emphasizes promoting capacity for climate change adaptation.
- Easement language should allow changes in water use only if beneficial to wildlife. Similar comments argued for a more aggressive stance by the Service to restore historical hydrology in the San Luis Valley.
- An emphasis on sustainability in the San Luis Valley could help bolster its already strong or growing nonagricultural sectors such as finance, services, and tourism.
- The land protection strategy is transparent and guided by habitat needs for identified trust species.
- Conservation easements are effective and more popular than new Federal land acquisition.
- There is appreciation of the landscape-scale nature of the project.
- In addition to full-market value, the Service should consider bargain sales for easements.
- There is appreciation for the gradual nature of and phased approach to the SLVCA.
- The Service needs to include more areas of northern New Mexico (Chama Peaks area and Jicarilla Apache lands) in the project boundary.
- In addition to habitat value, the Service should consider other qualities such as historical resources, open space, and public access.
- The easement program should accommodate small parcels, such as the vara strips associated with acequia irrigation practices (community-owned water distribution).
- The program will protect both wildlife and agriculture.

Comments not in support of the project identified the following concerns:

- There have been bad past experiences with easements restricting changes in agricultural operations.
- The Service should consider impacts of easement restrictions on the ability of utility companies to promote electrical reliability and renewable energy.
- Industry was not reached during scoping.

- There is general dissatisfaction with the impact of the Federal Government on land access and quality of life.
- The Rio Grande Water Conservation District is concerned about potential competition between the Service’s easement program and their attempts to acquire land for mitigation for the San Luis Valley Habitat Conservation Plan.

The following substantive questions were raised that were neither in opposition to nor in support of the SLVCA:

- How will being within the SLVCA boundary influence decisionmaking (for example, grazing permits) by other Federal agencies?
- How will the presence of an easement on an adjacent property affect a landowner who chooses not to sell an easement?
- How will the SLVCA’s establishment affect traditional use rights for Hispanos in Costilla County?

National Wildlife Refuge System and Authorities

The SLVCA will be part of the National Wildlife Refuge System, whose mission is “to administer a national network of lands and waters for the conservation, management, and where proper, restoration of the fish, wildlife, and plant resources and their habitats within the United States for the benefit of present and future generations of Americans” (National Wildlife Refuge System Improvement Act of 1997). National wildlife refuges provide important habitat for native plants and many species of mammals, birds, fish, insects, amphibians, and reptiles. They also play a vital role in conserving threatened and endangered species. Refuges offer a wide variety of wildlife-dependent recreational opportunities, and many have visitor centers, wildlife trails, and environmental education programs.

Conservation of more wildlife habitat in the SLVCA will be consistent with the following policies and management plans:

- Migratory Bird Treaty Act (1918)
- Migratory Bird Hunting and Conservation Stamp Act (1934)
- U.S. Fish and Wildlife Act (1956)
- Bald and Golden Eagle Protection Act (1962)
- Land and Water Conservation Fund Act (1965)

- Endangered Species Act (1973)
- Migratory Non-Game Birds of Management Concern in the U.S. (2002)
- Alamosa and Monte Vista National Wildlife Refuge Complex Comprehensive Conservation Plan (2003)
- Baca National Wildlife Refuge Conceptual Management Plan (2005)

The acquisition authorities for the easements and property acquisition are the U.S. Fish and Wildlife Act of 1956 (16 United States Code [U.S.C.] 742a–j) and the National Wildlife Refuge System Administration Act of 1966 (16 U.S.C. 668dd–ee), as amended. Land will be acquired with the use of the Land and Water Conservation Fund, which is derived primarily from oil and gas leases on the Outer Continental Shelf, motorboat fuel taxes, and the sale of surplus Federal property. As proper, the Service could also buy land interest through the use of Federal Duck Stamp revenue from the Migratory Bird Hunting and Conservation Stamp Act of 1934. There could also be more money to acquire lands, water, and interests for fish and wildlife conservation purposes as identified by the U.S. Congress or donations from nonprofit organizations. Any acquisition from willing sellers will be subject to available money.

Related Actions and Activities

The San Luis Valley and surrounding mountains contain many public lands and private protected areas, some of which are contiguous with other protected areas and some of which are isolated. Several existing State, Federal, and private land trust programs promote the conservation of habitats in the SLVCA; these public lands and their relationship to the SLVCA and SCCA are shown in figure 2.

SAN LUIS VALLEY NATIONAL WILDLIFE REFUGE COMPLEX

The San Luis Valley National Wildlife Refuge Complex includes three existing refuges and two conservation areas: the Alamosa, Baca, and Monte Vista National Wildlife Refuges and the SCCA and the SLVCA.

- The refuges were established for different purposes, as outlined earlier in this chapter, and protect 12,026 acres, 92,500 acres, and 14,800 acres, respectively. All three refuges currently contain a variety of habitats, with a special emphasis on wetlands and riparian systems. Management practices include vegetation manipulation and the artificial movement of water. An updated comprehensive

- conservation plan for the three refuges was finalized in 2015.
- The SCCA has an approved acquisition boundary of approximately 1 million acres, of which we are authorized to acquire up to 250,000 acres of conservation easements; nearly 172,000 acres have been conserved through this initiative to date.
- The SLVCA has an approved acquisition boundary of 4.2 million acres within which the Service can acquire up to 250,000 acres of conservation easements and 30,000 acres in fee title.

U.S. FISH AND WILDLIFE SERVICE—PARTNERS FOR FISH AND WILDLIFE PROGRAM

Since 1990, the Service has been working with private landowners in the San Luis Valley to conduct habitat improvement projects on private lands. These projects are with willing landowners and target the needs of trust species, such as migratory birds and imperiled species. Projects have focused on restoring wetlands, improving riparian habitats, and restoring habitat for Rio Grande sucker, chub, and cutthroat trout.

Agreements with landowners include technical and financial assistance for restoration practices in exchange for habitat assurance with a 10-year agreement for maintenance and management. Since the program has been active in the San Luis Valley, 285 agreements have been established to improve habitat on 16,680 acres of wetlands and riparian areas; 12,720 acres of uplands, and 98 river miles. An objective of the SLVCA is to continue working with the same landowners and use conservation easements to permanently protect the habitat improvements they have made to their property through the Partners for Fish and Wildlife Program.

USDA FOREST SERVICE

The Rio Grande, San Isabel, and Carson National Forests border the SLVCA to the north, east, and south. These forests contain nearly 4.5 million acres of public lands in the Sangre de Cristo, Saguache, and San Juan Mountains. The forests contain habitat ranging from pinyon-juniper savanna in the lower areas up to alpine tundra and scree fields at elevations more than 14,000 feet. Much of this is designated wilderness. These national forests are important habitat for Federal trust species, including Canada lynx and Rio Grande cutthroat trout, and for unlisted but climate change-imperiled species, such as American pika and white-tailed ptarmigan.

BUREAU OF LAND MANAGEMENT

Much of the land between the national forest boundaries and the largely private valley floor is administered by the Bureau of Land Management (BLM) as




Figure 2. Map of the San Luis Valley Conservation Area which will be part of a broader network of public and private conservation lands in Colorado and New Mexico.

the San Luis Resource Area. The BLM is actively working to restore the historical playa wetlands in the South San Luis Lakes and Blanca Wetlands areas, the latter of which has been designated as an Area of Critical Environmental Concern. These intermittent wetlands are particularly important for migratory shorebirds, some of which nest in the valley, and are also a priority habitat for the Service. The BLM also manages the 242,455-acre Rio Grande del Norte National Monument, which overlaps the southern tip of the project area in northern New Mexico. Rio Grande del Norte National Monument was established in 2013 by the President of the United States to protect the scientific and historical resources of the Rio Grande Gorge and surrounding landscape.

NATIONAL PARK SERVICE

Within the SLVCA boundary is the Great Sand Dunes National Park and Preserve. Together, these comanaged National Park Service units protect approximately 150,000 acres, from valley floor rabbitbrush scrub and the tallest sand dunes in North America to peaks more than 13,000 feet in the Sangre de Cristo Mountains.

NATURAL RESOURCES CONSERVATION SERVICE

The Natural Resources Conservation Service (NRCS) actively works in the San Luis Valley through its Wetlands Reserve Program, a voluntary easement program offering landowners the opportunity to protect, restore, and enhance wetlands on their property. They do not own land in fee title, but rather provide technical and financial support to help landowners with wetlands restoration efforts.

STATE OF COLORADO

The State of Colorado owns thousands of acres throughout the region and administers State Wildlife Areas and State Habitat Areas on many private lands. There are several school sections, managed by the State Land Board to provide revenue for K–12 education in the State. Some of these State Land Board parcels, such as La Jara Reservoir, allow recreational use as part of the Public Access program with Colorado Parks and Wildlife. There are a handful of regionally important wetlands and riparian corridors managed as state wildlife areas, including Russel Lakes; San Luis Lakes; and Rio Grande, Higel, and Hot Creek State Wildlife Areas. South of Baca National Wildlife Refuge and west of Great Sand Dunes National Park and Preserve is San Luis Lakes State Park, which provides important habitat for migratory birds as well as opportunities for wildlife-dependent recreation and watersports.

In addition, the State of Colorado has made significant investments in land conservation in the SLVCA through the additional programs of Great Outdoors

Colorado, Colorado Parks and Wildlife, and Colorado Water Conservation Board.

LAND TRUSTS

Tens of thousands of acres are protected in either fee title and easement programs paid for, or administered by, several conservation and land trust organizations, including but not limited to the Wetlands America Trust, The Nature Conservancy, the Rocky Mountain Elk Foundation, the Colorado Open Lands, the American Farmland Trust, Ducks Unlimited, the Rio Grande Headwaters Land Trust, and the Colorado Cattleman's Agricultural Land Trust. These accomplishments were made possible by substantial funding from State, private, and other Federal sources. These organizations have many different objectives; some focus on the preservation of undeveloped agricultural land to provide resources for the future, some are interested in protecting specific wildlife resources such as wetlands, and some have cultural or recreational objectives. The efforts of each of these organizations complement each other as well as efforts being undertaken by public agencies, including the Service. The locations of easements on private land are largely confidential, but there are some important land trust properties held in fee title as well, such as The Nature Conservancy's Medano-Zapata Ranch, which borders Baca National Wildlife Refuge and Great Sand Dunes National Park and Preserve. This property is a 103,000-acre working ranch and is home to a herd of 2,500 bison that are managed to mimic natural grazing patterns in the high desert shrub and grasslands.

Habitat Protection and the Easement Acquisition Process

Habitat protection will occur through the purchase of conservation easements and limited fee-title lands when necessary to further the management of existing refuges. It is the Service's long-established policy to acquire the minimum interest in land from willing sellers to achieve habitat protection goals, and conservation easements are an effective tool for achieving these goals.

The acquisition authority for the SLVCA is the U.S. Fish and Wildlife Act of 1956 (16 U.S.C. 742a–j). The Federal money used to acquire conservation easements will largely come from the Land and Water Conservation Fund, which is derived from oil and gas leases on the Outer Continental Shelf, motorboat fuel tax revenues, and the sale of surplus Federal property. There could be more money to acquire interests in habitat and water through direct congressional appropriations, donations, and the Federal Land Transaction Facilitation Act if the U.S. Congress votes to reauthorize it.

Conservation Easements and Other Acquisitions

An easement is a conservation tool that has been extensively employed in the SLVCA project area and throughout the larger region by other organizations. Easements involve the acquisition of certain rights to the property, such the right to subdivide or develop certain types of new infrastructure, while leaving the land title in the hands of the private property owner. Easements tend to be a cost-effective and socially acceptable means of habitat conservation. Many of the current agricultural land use practices are consistent with wildlife resource protection, and the use of easements will help ensure a strong and vibrant rural lifestyle.

There may be circumstances in which management objectives cannot be achieved, such as small boundary adjustments to existing refuges. In these cases, the Service will consider the limited use of fee-title acquisition, not to exceed 30,000 acres, as was described in the preliminary project proposal for the SLVCA.

Chapter 2—Area Description and Resources

This chapter describes the biological, cultural, and socioeconomic resources of the SLVCA that could be affected by its establishment. The SLVCA consists of approximately 4 million acres within the Southern Rockies and the Arizona and New Mexico Plateau ecoregions (U.S. Environmental Protection Agency 2011). The project encompasses all or most of Alamosa, Conejos, Hinsdale, Mineral, Rio Grande, and Saguache Counties in Colorado, as well as a portion of northern Rio Arriba and Taos Counties, New Mexico. About half of the project area is publicly owned.

Because of the nearly 7,000 feet in elevation change across the project area, the SLVCA contains a diverse array of plant communities, ranging from rabbitbrush scrub and sagebrush on the valley floor to alpine tundra and scree fields on the peaks of the surrounding mountains. As described in detail in this chapter, the habitats of the valley and surrounding mountains are crucial to the breeding and migration of migratory birds, and provide important opportunities for persistence or reintroduction of populations of imperiled species that are protected under the Endangered Species Act.

Physical Environment

The physical environment of the conservation area is described below in terms of its geology, minerals, water, hydrology, and climate.

GEOLOGY

The San Luis Valley is part of the much larger Rio Grande Rift Zone that extends from southern New Mexico northward through the San Luis and Upper Arkansas valleys to its northern termination near Leadville, Colorado (McCalpin 1996). The San Luis Valley is bordered on the east by the linear Sangre de Cristo Mountains, which were created by extensive block faulting during the Laramide Orogeny. The north-northwest part of the valley is bordered by the southernmost reach of the Sawatch Mountains. The west side of the valley is flanked by the San Juan Mountains, the result of extensive Tertiary-aged volcanism. In sharp contrast to the steeply rising mountains on the eastern side of the valley floor, the Oligocene volcanic rocks of the San Juan Mountains dip gently eastward into the valley floor, where they are interbedded with valley-fill deposits. Valley-fill



© Joe Zimm

Dozens of species of migratory waterbirds forage or nest in seasonal and temporary wetlands.

deposits consist of sedimentary rocks that interfinger with volcanic deposits. Quaternary deposits include alluvium, sand dunes, and pediments along the mountain fronts (USFWS 2011).

MINERALS

Sand and gravel are the major mineral commodities that are mined near the San Luis Valley. Rock, sand, and gravel mines are scattered throughout the valley, but are concentrated around the cities of Alamosa and Monte Vista and the town of Del Norte, Colorado. No coal mining permits are active in the SLVCA (Colorado Division of Reclamation, Mining,

and Safety 2012). Other minerals that are mined in the area include gold, silver, peat, and limestone. There is also nascent oil and gas exploration in the valley (USFWS 2011).

WATER AND HYDROLOGY

The SLVCA contains the upper headwaters of the Rio Grande watershed (figure 3). Because of its position in a high-mountain desert, the valley floor receives little precipitation, and most surface and ground water is a result of runoff from the surrounding mountains. There are numerous perennial and intermittent drainages that descend from the Sangre de Cristo and San Juan Mountains. Some of the larger waterways include the Alamosa, Conejos, Rio Grande, and San Antonio Rivers.

A portion of the northern valley, known as the Closed Basin (a sump) does not directly contribute water to the Rio Grande. The Closed Basin may have formed in the middle Pleistocene when the lake that filled the valley began to dry up, resulting in an environment of swamps and organic-rich sediments. Mayo et al. (2006, as cited in USFWS 2011) refer to the Closed Basin of Pleistocene time as the “ancestral sump.” Currently, the Closed Basin covers approximately 2,940 square miles in the northern part of the valley and is separated from the rest of the valley by a low alluvial fan. The Closed Basin is composed of the drainage basins of several small but important waterways, including but not limited to Carnero, Cottonwood, Crestone, La Garita, Medano, Saguache, San Luis, Spanish, and Willow Creeks. Water enters the Closed Basin through precipitation and snowmelt from the 4,700 square miles of watershed in the surrounding mountains and substantial inflow of water diverted from several ditches off the Rio Grande, including the Farmer’s Union Canal, Rio Grande Canal, and others. Water exits primarily through evapotranspiration and through exports from the U.S. Bureau of Reclamation’s Closed Basin Project (USFWS 2011). The Closed Basin Project extracts ground water from the unconfined aquifer in the sump. This water is carried in the Closed Basin conveyance channel, which starts in the central part of the basin, passes through the Baca Refuge, and delivers water to the Rio Grande on the Alamosa Refuge. Water from the Closed Basin Project helps Colorado meet its interstate compact obligations with New Mexico and Texas (USFWS 2012a). The Bureau of Reclamation closely monitors changes in the water table caused by the Closed Basin Project and continually adjusts the location and volume of pumping to ensure operations are within the project’s authorizing guidelines. The effects of other pumping in the sump and the chronic lack of runoff and aquifer recharge have required the Bureau of Reclamation to carefully assess its monitoring data to distinguish these effects from other factors.

The rest of the project area is in the San Luis Valley portion of the Rio Grande aquifer system. The San Luis Valley is the northernmost portion of a system of discrete and separate aquifers that stretches from Saguache County, Colorado, to western Texas (Robson and Banta 1995). The thick basin-fill deposits in the San Luis Valley consist of interbedded clay, silt, sand, gravel, and volcanic rock. These form many separate aquifer systems, which are generally grouped into two major aquifers—a shallow unconfined aquifer and a deep confined aquifer—although the lines between these features are not absolute. The unconfined aquifer is separated, but not totally disconnected, from the confined aquifer by clay layers and lava flows. The unconfined aquifer is recharged through infiltration of precipitation, irrigation water, runoff, and upward seepage of ground water from the confining bed. Discharge from the unconfined aquifer is from ground water withdrawals, ground water flow to the south, discharge to streams or drains, and evapotranspiration. Water levels in the unconfined aquifer respond to local climatic events and fall or rise with the availability of precipitation. Wells drilled into the deep confined aquifer are artesian and are buffered from climatic conditions. The confined aquifer is recharged from precipitation and snowmelt in the high San Juan Mountains and the Sangre de Cristo Mountains. Discharge from the confined aquifer is principally from ground water withdrawals and upward leakage through the confining bed (USFWS 2012b).

Alamosa, Baca, and Monte Vista National Wildlife Refuges all depend on extensive ground and surface water rights, acquired with respective property acquisitions to provide wetland and wildlife habitat. The beneficial uses of all surface rights on the three refuges are decreed by the State of Colorado as “irrigation” and must be used to grow wetland plants within legal criteria. The seniority of surface water rights on the three refuges range from junior rights, with recent appropriation dates, to senior rights appropriated in the late 1800s.

Surface water use on the refuges varies widely between years, depending on availability, and averages 20,795 acre-feet per year and 6,875 acre-feet per year on the Alamosa and Monte Vista Refuges, respectively. Surface water use on the Baca Refuge is poorly documented due to the historical lack of instrumentation. Measurement devices were installed on streams flowing onto the Baca Refuge in 2013 and will provide important data over time.

Flowing at approximately 2,800 gallons per minute, the Mumm Well on the Alamosa Refuge is the largest unconfined aquifer well in the San Luis Valley. The decree for this well limits diversions to 1,540 acre-feet per year. Water from this well represents the vast majority of ground water use on the Alamosa Refuge.




Figure 3. Map of the San Luis Valley and Sangre de Cristo Conservation Areas, Colorado and New Mexico, which capture the upper headwaters of the Rio Grande, the fourth longest river in the United States.

The Monte Vista Refuge uses significant quantities of ground water to meet wildlife management objectives. Although there is substantial variation between years, ground water use on the refuge averages approximately 8,122 acre-feet per year.

The U.S. Fish and Wildlife Service is a major user of both ground and surface water in the San Luis Valley. Federal law requires each Federal agency to participate in the water adjudication process of the associated State.

CLIMATE

The climate of the San Luis Valley is consistent with its high mountain desert setting, with substantial 24-hour temperature swings because of cold air drainage from the surrounding mountains. This cold air also creates winter overnight temperatures that are often much lower than at many other places at similar elevations and latitudes. The mid-January high averages 34 °F while the low averages -2 °F, and the mid-July high averages 83 °F while the low averages 37 °F. The montane and alpine parts of the SLVCA have much cooler weather because of their 10,000- to 14,000-foot elevations.

Precipitation in the valley is strongly influenced by the surrounding mountains. The windward side of the mountain ranges, particularly the San Juan Mountains, receives a substantial amount of orographic

precipitation, which is caused when air masses rise and subsequently cool, dumping their precipitation at higher elevations. This results in a marked rain shadow effect on the lee side of the mountains, with annual precipitation in Alamosa averaging 7.25 inches per year (National Weather Service 2012).

Biological Environment

Plant communities in the San Luis Valley form a variety of vegetated habitats—wetlands, riparian areas, and uplands—used by wildlife in the area. Refer to appendix D for a complete list of plant and animal species in the SLVCA.

PLANT COMMUNITIES

The vegetation across the project area varies greatly, depending on hydrology, slope, aspect, and elevation. The San Luis Valley's hydrology is strongly influenced by surface runoff and ground water flows from the surrounding mountains. These conditions have created a network of riparian corridors and wetlands that break up large expanses of associated desert and upland habitats across a 7,000-foot elevation gradient, resulting in high plant diversity. The broader San Luis Valley ecosystem contains 1,132 species of



© Joe Zimm

The riparian corridors of the San Luis Valley serve as wildlife corridors and provide nesting habitat for the endangered southwestern willow flycatcher.

plants (appendix D; Colorado State University Herbarium 2012), which is more than a third of the total plant species present in Colorado.

Wetlands

Wetlands in the conservation area are wet meadows, seasonal wetlands, and semipermanent wetlands. Irrigation practices on private and public lands support most of these wetlands.

Wet Meadows. Wet meadow habitat is naturally present in the San Luis Valley in both areas that have shallow water tables and areas that are periodically shallowly inundated early in the growing season. Wet meadows are the most widespread wetland type in the San Luis Valley. Dominant plants include Baltic rush, hair grass, and sedges. Most of the naturally occurring wet meadows have been modified by changes in water use, but in some areas this has resulted in an expansion of wet meadow areas because of artificial irrigation for hay fields and cattle grazing. These agricultural uses create habitat for a variety of wildlife (USFWS 2005).

The combination of plant structure and density coupled with water depth and duration creates rich habitat diversity within each larger area of wet meadow. This richness of habitat creates tremendous foraging and nesting opportunities for a variety of bird species. Among these are many species of ducks and geese as

well as sora, Virginia rail, white-faced ibis, American avocet, Wilson's snipe, and Wilson's phalarope. Wet meadows provide critical roosting and foraging areas for the Rocky Mountain population of greater sandhill cranes, which migrate through the valley in the spring and fall. Wet meadows also provide habitat for a variety of regionally rare or unusual amphibian species, such as northern leopard frog and Plains spadefoot toad (USFWS 2005). Also present in this habitat, particularly in areas of alkali soils, is the somewhat rare slender spiderflower, which once had a wide range in the southern Rocky Mountains but now occurs almost exclusively in the San Luis Valley.

Seasonal and Semipermanent Wetlands. Seasonal and semipermanent wetlands have hydrologic regimes that typically allow for the persistence of water throughout the growing season. Water in these areas is often deeper than 1 foot. Semipermanent wetlands may have substantial areas of open water with aquatic vegetation beds, and are often fringed by tall emergent vegetation. Tall emergent wetlands can also be seasonal and are typically dominated by bulrush and cattails.

Swimming birds, including grebes, coots, and waterfowl, as well as aerial species such as swallows and terns, use open-water areas of these wetlands for foraging. Emergent vegetation provides breeding habitat for diving and dabbling ducks, Canada



© Trinchera Ranch

The semidesert shrublands and sagebrush habitats of the San Luis Valley Conservation Area, while stark in appearance, are important habitat for declining bird species such as the sage sparrow and sage thrasher.



Steve Torbit / USFWS

The American pika “hays” the alpine grasses as a means of surviving the harsh winters above the treeline in the surrounding mountains.

geese, American bitterns, snowy and cattle egrets, black-crowned night-herons, white-faced ibis, and marsh passerines (songbirds) such as marsh wrens, common yellowthroats, and yellow-headed blackbirds. Northern harriers and short-eared owls will also nest in residual patches of tall emergent vegetation. Tall emergent wetlands with a high density of sedges and a shallow seasonal water regime host rails and provide nesting sites for dabbling ducks.

Floodplains and Playa Wetlands. Botanical studies and limited historical accounts indicate floodplain depressions and playa wetlands in the Closed Basin of the San Luis Valley exhibited dynamic flooding regimes and periods of equally dramatic drying. These wetlands received water from surface sources including snowmelt and monsoonal rains. During a series of wetter years, the result was emergent plant communities, including spikerush, scarlet smartweed, and pondweed. During extended dry periods, these same wetlands plant communities converted back to desert shrub vegetation, such as greasewood, rabbitbrush, and saltgrass, during extended periods of low precipitation (Heitmeyer and Aloia, 2013).

Riparian Habitats

Riparian habitat has trees, shrubs, and other streamside vegetation and is associated with intermittent and perennial waterways. This community may flood every year. The historical extent on the valley floor has been reduced by the diversion of surface water and depletions from ground water use. Woody

riparian habitat is sensitive to excessive grazing and browsing from both domestic and wild ungulates, which limits regeneration of the dominant willows and narrowleaf cottonwood trees. Shrubs that contribute to the structural diversity of riparian habitat include redosier dogwood and greasewood. These shrublands and forests provide important stopover habitat for migratory passerines, as well as nesting habitat for species such as Lewis’ woodpecker, willow flycatcher, and possibly yellow-billed cuckoo. In addition, the shade and streambank stabilization provided by riparian vegetation is important in keeping temperature and water quality in streams and rivers for species such as the endemic Rio Grande cutthroat trout, Rio Grande chub, and Rio Grande sucker.

Upland Vegetation

Upland vegetation in the conservation area ranges from semidesert shrublands and grasslands to montane forests and above treeline.

Semidesert Shrublands and Grasslands. Shrublands are the most common natural vegetation on the San Luis Valley floor. Many of the plants within these communities are drought resistant and tolerant of high soil salinity. These shrublands are characterized by an open to moderately dense assemblage of rubber rabbitbrush, greasewood, fourwing saltbush, shadscale, and winterfat. Also present in these communities are yucca, cactus, and various grasses. At slightly higher elevations, rabbitbrush shrublands transition to desert scrub and shrub-steppe habitats that have a substantial cover of big sagebrush or sand sagebrush and that intergrade with the pinyon-juniper woodlands above. Grasses in these areas include Indian ricegrass, alkali sacaton, western wheat grass, and blue grama.

Bird diversity and density tend to be relatively low in semidesert shrublands because of structural and floristic simplicity (Wiens and Rotenberry 1981). Species common to this habitat include the horned lark, mourning dove, western meadowlark, and loggerhead shrike. Upland grassland habitats have the potential to support grassland-dependent species such as burrowing owl, long-billed curlew, and a variety of sparrows. The sagebrush-dominated habitats are also home to the declining sage thrasher and the federally proposed as endangered Gunnison sage-grouse.

Montane Forests. Above the semidesert shrubland, the vegetation transitions into pinyon-juniper woodland. This open-canopy forest is dominated by pinyon pines and junipers, with an understory consisting of shrubs and grasses. According to the Colorado Natural Heritage Program, this woodland’s threat status is “fair” and its protection status is “poor-fair.” Pinyon-juniper woodland is particularly threatened by the spread of invasive grasses that increase its susceptibility to fire (Colorado Natural Heritage Program and

The Nature Conservancy 2008). Much of the existing pinyon-juniper woodland in the San Luis Valley is managed by BLM, though there are extensive stands on private lands in Costilla County. Pinyon jays are obligate nesters in the pinyon-juniper woodlands, and although their population is stable in Colorado, they are effective indicators of forest health and are therefore a priority species for Partners in Flight throughout the Intermountain West (Colorado Partners in Flight 2000). Other pinyon-juniper associated species include black-throated gray warbler and juniper titmouse.

As the elevation increases, the forest becomes a mixed conifer forest, which is sometimes part aspen, and finally becomes a subalpine spruce-fir forest. These forests are home to some bird species, including olive-sided flycatcher, yellow warbler, and mountain chickadee; they also provide habitat and migration corridors for some important large mammals such as elk, black bear, and the threatened Canada lynx.

Above Treeline. The highest elevations in the SLVCA are dominated by alpine tundra, scree fields, and bare stone, which can have the appearance of being stark or even lifeless. Upon closer inspection, however, one observes a remarkable diversity of plants adapted to this cold and arid environment, including impressive displays of summer wildflowers. These plants provide the foundation for an ecosystem containing a suite of charismatic fauna, many of which are imperiled by habitat shifts because of climate change such as the American pika. The high elevations are also home to State game species such as bighorn sheep.

WILDLIFE

The diverse mix of wetland, riparian, shrubland, forest, and alpine habitats throughout the SLVCA provide for the habitat needs of many assemblages of reptiles and amphibians, aquatic species, birds, and mammals, including several species of special concern. Appendix D lists the wildlife species found in the San Luis Valley and surrounding mountains.

Amphibians and Reptiles

The San Luis Valley is a cold desert, so it supports only a limited number of reptiles and amphibians. The large areas of semidesert shrubland and the scattered wetlands and riparian areas are home to a handful of snakes and lizards as well as the snapping turtle. The arid nature of the region restricts amphibians largely to wetlands and riparian corridors; these areas provide habitat for tiger salamander and seven species of frogs, toads, and spadefoot toads. Among the latter group is the boreal toad, a high-elevation toad that appears to have declined substantially because of infection by *Batrachochytrium dendrobatidis*, a pathogenic fungus. This species is State listed as endangered by both Colorado and New Mexico (Colorado Parks and Wildlife 2012).



The Rio Grande cutthroat trout, once found throughout the Rio Grande and Pecos River watersheds, is now found only in scattered cold-water, high-elevation streams.

USFWS

Fish and Aquatic Species

The project area contains the headwaters of the Rio Grande. The Rio Grande and its tributaries and the valley's marshes are home to several native fish as well as a range of introduced species. Most of the challenges faced by these aquatic species are due at least in part to anthropogenic causes such as competition with exotic species and water diversions. These impacts have been magnified by persistent drought conditions since the 1990s. In some cases, conservation easements offered by the SLVCA may be used to protect land use practices on watersheds that are beneficial to water quality and temperature. Easements that tie water use to the land will have neutral effects on aquatic species.

The Rio Grande chub is thought to have once been the most common fish throughout the Rio Grande drainage and in the San Luis Closed Basin, but it has been extirpated from much of its range, including from the main stem of the Rio Grande. The Rio Grande chub is now found in several small streams in the San Luis Valley, including Crestone Creek on Baca National Wildlife Refuge. The Colorado Natural Heritage Program considers the Rio Grande chub to be an S1 (critically imperiled) species. It is thought to have declined because of habitat fragmentation by impoundments for diversions, habitat destruction because of poor land use practices, and predation by, and competition with, introduced fish species (Rees et al. 2005a).

The Rio Grande sucker once had a historical range similar to that of the Rio Grande chub, and faces similar threats. It appears to have been particularly hard hit by competition with the introduced white sucker. At one point, the Rio Grande sucker was reduced to a single population in Hot Creek in Conejos County, Colorado, but it has since been discovered in Crestone Creek and reintroduced to several more streams. It is considered a State endangered fish in Colorado (Rees et al. 2005b).

In historical times, Rio Grande cutthroat trout¹ were found in large numbers in the main stem of the Rio Grande and its major tributaries, such as the Conejos River; one account from the Conejos River in 1877 states that “fishing was so successful ... our catch amounted to over a hundred pounds by midafternoon,” which the fishermen shipped off to a restaurant in Denver (Sanford 1933). Now, the native trout are restricted to high-elevation streams descending from the San Juan and Sangre de Cristo Mountains. The Rio Grande cutthroat trout occupies approximately 10 percent of its historical range. Threats to the species include competition and hybridization with, and predation by, introduced trout; reduction in habitat quality because of water diversions and other hydrological changes; and changes in stream temperature because of human water use and global climate change. Formerly identified as a candidate species under the Federal Endangered Species Act, it was recently removed from consideration in late 2014.

Some 57 species of nonnative fish have been introduced to the San Luis Valley, either as naturalized aquarium fish, escaped aquaculture species, or intentionally introduced sport fish. The latter category has brook, brown, golden, and rainbow trout; northern pike; bluegill; pumpkinseed; yellow bullhead; common carp; largemouth and smallmouth bass; blue, channel, and flathead catfish; walleye; and yellow perch. Nongame species such as American eel, grass carp, Mozambique tilapia, white suckers, and even neotropical tetras and armored catfish have become naturalized in the Rio Grande drainage as well (USGS 2012).

Birds

The diverse range of habitats along the elevational and hydrologic gradient of the SLVCA provide habitat for at least 274 species of birds. Some of these birds are year-round residents, but many migrate through the valley on their way to and from wintering and breeding grounds while others come to the valley to breed or spend the winter. Among the migratory species are neotropical migrants that winter in Central and South America and breed in North America. Riparian corridors and forests are particularly important to these species.

Cordilleran flycatchers breed in forested areas of the SLVCA, including cottonwood riparian forest. These gallery riparian forests are also thought to host a limited number of yellow-billed cuckoos, recently federally listed as threatened. Olive-sided flycatchers breed in the coniferous forests of the mountains surrounding the valley. The southwestern willow flycatcher, a subspecies of the more widespread willow flycatcher,



C. Suzanne Langbridge / USGS

The endangered southwestern willow flycatcher nests in the willows along the Rio Grande and its tributaries.

breeds in shrub riparian and tree riparian with a willow understory; the southwestern willow flycatcher is federally and State listed as endangered. Examples of other neotropical migrants in the SLVCA include two species of phoebe, several more flycatchers, western tanager, gray catbird, Bullock’s oriole, and many species of warblers.

Passerines are not the only migrants to make use of the area. Black-necked stilts and American avocets are shorebirds that migrate from winter ranges in Mexico, Central America, and South America to breed in the wetlands of the San Luis Valley. At least 25 other species of shorebirds use these wetlands as either stopover or breeding habitat. Six of these shorebirds, including the snowy plover, which breeds in the playa wetlands of the Closed Basin, are either focal species for the Service’s Migratory Bird Program or are the Service’s Region 6 Birds of Conservation Concern.

Given the scarcity of water in high desert and mountain environments, it is perhaps not surprising that the San Luis Valley is regionally important for both resident and migrant waterbirds. The marshes of the valley support 27 species of waterfowl. Approximately 30 percent of the cinnamon teal that summer in Colorado breed in the valley (Stephanie Jones, USFWS Migratory Birds, personal communication 2012). The secretive American bittern breeds in the valley, and has experienced population declines throughout its range, likely because of wetland disturbance. The white-faced ibis breeds in wet meadows and makes extensive use of natural and agricultural habitats in the valley. Nearly the entire Rocky Mountain population of sandhill cranes uses the San Luis Valley as migratory stopover habitat, particularly on and around

¹76 Federal Register No. 207, Wednesday, October 26, 2011. Endangered and Threatened Wildlife and Plants; Review of Native Species That Are Candidates for Listing as Endangered or Threatened; Annual Notice of Findings on Resubmitted Petitions; Annual Description of Progress on Listing Actions. 66403.

the Monte Vista National Wildlife Refuge, where they are the focus of an annual crane festival and a draw for thousands of tourists every year. Rookeries of great blue herons, snowy egrets, and black-crowned night-herons are also present. Conservation of wet meadow, playa, and emergent wetland habitat is crucial for these species.

The San Luis Valley hosts an array of diurnal raptors and owls throughout the year. Prairie falcons are common year-round residents and use uplands extensively for feeding and resting. The trees and snags along waterways are nesting sites for great horned owls, long-eared owls, red-tailed hawks, American kestrels, and Swainson's hawks (USFWS 2011). The latter species is a bird of conservation concern in the Service's Region 6 and is known to be sensitive to habitat fragmentation. Northern harriers and short-eared owls nest in wet meadows and emergent wetlands. These two species as well as ferruginous hawks, rough-legged hawks, golden eagles, and bald eagles overwinter in the valley, where they forage for small mammals and other prey in riparian areas, uplands, and short-emergent wetlands where cover is abundant (USFWS 2011). The higher elevation parts of the project area are home to the northern goshawk, a generalist predator of rodents and birds that inhabits the montane forests of the surrounding mountains. It is probable that the forested canyons above the valley floor provide habitat for the Mexican spotted owl; these species are both State (Colorado) and federally listed as threatened, although no designated critical habitat for the species occurs in the project area.

The San Luis Valley is also in the eastern corner of the sagebrush region of the Intermountain West (Pitkin and Quattrini 2010) and, as such, has some strongly sagebrush-associated or sagebrush-obligate bird species, meaning those species whose life history needs cannot be met in other habitats. The Gunnison sage-grouse has a small population at the north end of the San Luis Valley (D. Reinkensmeyer, personal communication with M. Dixon, February 2012). This

species is now listed as threatened under the Federal Endangered Species Act and is a species of special concern in Colorado. Gunnison sage-grouse likely had much broader distribution than they do now (Schroeder et al. 2004). The Colorado Parks and Wildlife identified that some of this former range has potential habitat for the species (Gunnison Sage-Grouse Rangewide Steering Committee 2005). Sage sparrows have similar habitat associations, preferring sagebrush-dominated habitats with open to closed canopies (Williams et al. 2011). Sage thrasher is another denizen of the upland shrub habitats of the valley, including sagebrush and rabbitbrush scrub. It is one of the Service's migratory bird focal species and one of the Service's Region 6 species of concern. It is thought that the primary reasons for the decline of Gunnison sage-grouse are the loss and fragmentation of sagebrush habitat (Oyler-McCance et al. 2001), so this species is likely to benefit from the protection of remaining potential habitat that the project will provide. Given the overlap in habitat needs of sage-grouse and other sagebrush obligates (Rowland et al. 2006), species like sage thrasher and sage sparrow will likely benefit from conservation of sagebrush and steppe habitat as well.

Mammals

The arid uplands, wetlands, and stream and river corridors of the SLVCA provide habitat for large game species, including pronghorn, elk, and mule deer. The higher elevations hold Rocky Mountain bighorn sheep. American bison were once an important part of both the San Luis Valley ecosystem and the socioeconomic system of the Ute and Pueblo peoples; however, the last bison were extirpated from the San Luis Valley by 1870 (Colville 1995). The Nature Conservancy now manages a bison herd on their Medano-Zapata Ranch as a means of simulating natural grazing regimes; however, their stated goal is to introduce a free-ranging genetically pure bison herd of at least 3,000 animals to the valley by 2015 (The Nature Conservancy 2008).



Steve Torbit / USFWS

The mountains surrounding the San Luis Valley provide important habitat for the threatened Canada lynx.



USFWS

The San Luis Valley and surrounding mountains are home to thousands of elk.

The bison and other megafauna provide opportunities for wildlife-dependent recreation, but are not without controversy. Perceived overpopulation of elk, in particular, is contentious among farmers and ranchers in the valley who are concerned about the crop damage and competition for forage between elk and cattle. Additionally, managers of the Baca and Alamosa Refuges are concerned about the documented effects of elk overbrowsing riparian vegetation important to migratory birds.

The elk herd on the east side of the valley has been estimated to number approximately 5,000 animals (R. Rivale, wildlife biologist, Colorado Parks and Wildlife, personal communication, cited in USFWS 2005). A recent study of elk carrying capacity in the Great Sand Dunes ecosystem found that, under current management practices, the carrying capacity of the region should be 6,104 elk (Wockner et al. 2010). Development of plans for elk management in the valley is ongoing.

Small mammals in the SLVCA are those typical of the greater Southern Rockies ecosystem. Riparian areas and marshes provide resources for beaver and common muskrat. Forested areas are home to North American porcupine and snowshoe hare. Uplands contain other rabbits, such as white-tailed jackrabbits and mountain cottontails, as well as the Ord's kangaroo rat. In the highest reaches of the project area, primarily above the tree line, are the charismatic American pika and the vocal and inquisitive yellow-bellied marmot. The Gunnison's prairie dog is a species which inhabits the valley floor. This species has suffered a sharp decline for reasons that include human persecution and outbreaks of plague. Formerly, it was listed

as a candidate for protection under the Endangered Species Act, but was recently removed following a genetic reevaluation of its taxonomic status.¹

The aforementioned species serve as prey for several predator species in the project area. Black bear is a generalist omnivore whose flexibility makes it common in many habitat types in the valley. The coyote is often found hunting small mammals and occasionally larger prey throughout the study area. Similarly, both mountain lion and bobcat are quite catholic in their habitat needs, though the mountain lion has much larger home ranges and tends to specialize in hunting ungulates, whereas the bobcat is more opportunistic. In contrast to those two cats, the State endangered and federally threatened Canada lynx is largely a specialist predator of snowshoe hare; in the SLVCA, lynx is primarily found in the spruce-fir forests of the Sangre de Cristo and San Juan Mountains, where its preferred prey are found.

The grizzly bear once roamed the mountains of the area but was extirpated from Colorado in the early 20th century; the San Luis Valley grizzlies are remembered now as the mascot of Adams State University in Alamosa, Colorado. Similarly, the gray wolf historically hunted in the San Luis Valley and surrounding mountains, but was extirpated from Colorado by 1945 (though it is still State and federally listed as endangered in Colorado). A mounting body of research shows the potential ecological benefits of natural or

¹Federal Register 78, No. 220. November 14, 2013. Endangered and Threatened Wildlife and Plants; 12-Month Finding on a Petition to List the Gunnison's Prairie Dog as an Endangered or Threatened Species. 68660

human-facilitated reintroduction of wolves, particularly on vegetation adversely affected by unnaturally high elk browsing (Ripple and Beschta 2012). Neither State nor Federal agencies presently plan to reintroduce wolves to the project area.

Finally, the SLVCA is home to nine species of bats. All are insectivorous and hunt primarily by capturing insects in flight. The hoary bat and silver-haired bat are solitary tree-roosting bats that are present during the summer and migrate to warmer climates during the winter. The presence of mature cottonwood riparian forests likely supports their presence on the valley floor. The migratory Mexican free-tailed bat has an exceptionally large summer colony of approximately 100,000 individuals (Freeman and Wunder 1988) in the historic Orient Mine in the northern San Luis Valley outside the SLVCA, though there are certainly other old mines within the project area that may provide roosts for smaller colonies. The remaining species are either resident or regionally migratory hibernators.

Cultural Resources

On the hottest days it is cool in the shade, and on the very coldest days it is comfortable in the sunshine.

—Geologist C.E. Siebenthal, describing the San Luis Valley in 1910

Humans have inhabited the San Luis Valley and surrounding mountains for more than 12,000 years. Their uses of the land reflect both the traditions of those who moved to the valley and local adaptations. The following summary of the prehistory and history of the valley provides an overview of some of the major themes and events that illustrate the human interaction with the land. There is an abundance of prehistoric evidence as well as early historical accounts, records, photographs, and local histories for the valley. This synopsis provides only a glimpse into the resources and information available with an emphasis on environmental references.

PREHISTORY

The prehistory of the San Luis Valley has four stages: Paleo-Indian, Archaic, Late Prehistoric, and Protohistoric.

Paleo-Indian Stage

Current archaeological evidence shows that the earliest humans, called the paleo-Indians, migrated to the region near the close of the last ice age approximately 12,000 years ago. These people had a highly mobile lifestyle that depended on the hunting of large, now-extinct mammals, including mammoths and a huge ancient bison. The hallmark of most paleo-Indian sites

are the beautiful but deadly spear points that were launched with the aid of a simple yet expertly engineered spear-thrower called an atlatl. These projectile points are generally recovered as isolated occurrences or in association with animal kills, butchering sites, or small temporary camps. Although the timing of this stage varies throughout the region and is constantly being refined as more data become available, the stage generally lasted until about 7,500 years ago.

Information from the Colorado Office of Archaeology and Historic Preservation shows that 62 paleo-Indian resources have been identified in the San Luis Valley and surrounding mountains. These sites are often located near wetlands and along the shorelines of ancient lakes, reflecting the use of abundant floral and faunal resources available in these locations. Several paleo-Indian sites in the valley and surrounding mountains have been excavated, including the high altitude Black Mountain Site (5HN55) located at 10,000 feet in the San Juan Mountains south of Lake City in the SLVCA. This campsite dates from approximately 10,000 to 7,000 years ago and has yielded a variety of stone tools suggesting animal procurement and processing (Jodry 1999a).

Several paleo-Indian sites on the valley floor have been excavated and provide an extensive record of the early occupations. Three of these sites, the Cattle Guard site (5AL101), the Linger site (5AL91), and the Zapata site (5AL90), are located just south of Great Sand Dunes National Park and Preserve and represent camps with an abundance of bison bone and associated stone tools (Cassells 1997, Jodry 1999a). The Reddin site (5SH77) near the town of Hooper yielded nearly 500 paleo-Indian artifacts suggesting a variety of activities and uses (Cassells 1997, Jodry 1999a).

Climatic fluctuations during the Holocene Epoch (which started about 12,000 years ago and has continued to the present) are often reflected in the archaeological record. Pollen remains, faunal assemblages, and geomorphological deposits suggest periods of significant and rather abrupt vegetation changes and variations in the amount of moisture (Jodry 1999b, Martorano 1999a). Bison remains associated with archaeological sites on the southern plains also show oscillations in bison numbers in response to climatic conditions (Creel et al. 1990). Although more research is needed and archaeologists' ability to recover and interpret the prehistoric record is continually improving, these preliminary studies are an intriguing look into the evidence for and the consequences of long-term climatic change.

Archaic Stage

There was a gradual but definite shift in the pattern of human use of the region that began about 7,500 years ago and continued until approximately 1,500 years ago. The changes were the result of a



Meg Van Ness / USFWS

The San Luis Valley contains archaeological sites extending thousands of years into prehistory.

combination of regional climatic fluctuations and an increasing population coupled with technological innovation and regional influences. Although the Archaic stage is better represented in the archaeological record than the preceding paleo-Indian stage, the identification and interpretation of the remains continues to be expanded and refined. Evidence of a greater diversity of tools and the use of a larger variety of plants and animals than during the preceding paleo-Indian stage is found on many sites.

There have been 618 Archaic stage resources recorded in the Colorado part of the study area. As with the earlier inhabitants, the Archaic peoples made extensive use of the valley's wetland resources and occupied the rock shelters and several high-altitude locations found in the surrounding mountains. Speaking of Archaic sites in the northeastern part of the valley, Hoefler states: "Most of the Closed Basin archaeological sites are open camps containing debitage and fire-cracked rock scatters, approximately half of which contain ground stone implements such as metate fragments or manos. Many of these sites are located around seasonal wetland marshes and lakes" (Hoefler 1999).

The use of the atlatl with spear points continued and basketry, cloth, and cordage came into use. Although still mobile, the population increasingly made short-term use of small groupings of structures with storage features. Former hunting blinds and other rock structures are fairly common but often difficult to interpret. Archaic Stage rock art is scattered throughout the region and the influences of surrounding regions, particularly the Plains and the Great Basin, are identifiable at several sites.

Late Prehistoric Stage

Beginning approximately 1,500 years ago, several innovations greatly influenced life in the valley (Martorano 1999b). Although these changes were adopted at different rates and degrees throughout the area, the advent of pottery and the bow and arrow coupled with a larger and more sedentary population defines the period until approximately 600 years ago. Early archaeological research in the valley identified many regional influences, with several sites exhibiting pueblo-inspired attributes (Renaud 1942). In 1694, Don Diego de Vargas documented his visit to the valley, thus providing an early historical written account and ushering in the historical period.

The 442 Late Prehistoric resources in the Office of Archaeology and Historic Preservation database are listed under a variety of designations for this stage, but all date to about the same time period. The distribution of Late Prehistoric sites in the valley reinforces the trend of intensive use of wetland habitats (Martorano 1999b). This is not surprising as the available resources, both floral and faunal, would have continued to be abundant in these areas. Site types include camps, stone tool scatters, rock art, rock alignments and enclosures, and quarries where the lithic material for stone tools was collected.

Protohistoric Stage

By the late 1600s, Spanish incursions into the valley were beginning to affect the lives of the native populations. The Utes, who, based on archaeological evidence, came to the valley sometime after A.D. 1100 (Reed 1994) and were the most prevalent occupants of the valley, quickly acquired horses and other trade items. Although many other Native American groups probably visited or traveled through the valley, the Comanche, Apache, Navajo, Arapaho, Cheyenne, and several northern Pueblos also had a significant if not sustained presence (Martorano 1999c).

The 59 recorded Office of Archaeology and Historic Preservation sites from this stage include the traditional stone tools and ceramics mixed with used or flaked glass, trade beads, and metal projectile points. Wickiups (conical timbered structures) and trees with peeled bark (indicating the harvesting of the edible cambium layer) were common, as is rock art with motifs and depictions of postcontact goods.

EARLY HISTORY

The historical period for the valley began with the reoccurring contact of the native peoples with people of European descent and ended in the mid-twentieth century. This interaction generally followed many years of occasional contact, often for the exchange of trade goods. The narrative below briefly summarizes some of the major historical influences, patterns, and themes in the region.

Early Exploration and Trade

I take and seize one, two, and three times, one, two, and three times, one, two, and three times, and all those which I can and ought, the Royal tenancy and possession, actual, civil, and criminal, at this aforesaid River of the North, without excepting anything and without any limitation, with the meadows, glens, and their pastures and watering places. And I take this aforesaid possession, and I seize upon it, in the voice and name of the other lands, towns, cities, villas, castles, and strong houses and dwellings, which are now founded

in the said kingdoms and provinces of New Mexico, and those neighboring to them, and shall in future time be founded in them, with their mountains, glens, watering places, and all its Indian natives

—Capitán Gaspar Pérez de Villagrán in *La Historia de la Nuevo Mexico*, 1610

With these bold words in 1598, Spain claimed all lands, structures, and people along the Rio Grande—including the San Luis Valley—forever. This followed several years of sporadic Spanish incursions into northern New Mexico and southern Colorado, which ushered in several decades of trade, conflict, and settlement. Many Spanish traveled along the North Branch of the Spanish Trail, which had both western and eastern routes through the valley. Although the Spanish relinquished ownership of the valley in 1821, their influence survives as a vital part of the landscape and people today.

There are many explorers and settlers who left a legacy of journals, maps, and other accounts of their time in the San Luis Valley. These documents offer a wide variety of historical and environmental information. The examples summarized below provide a glimpse into the types of information and insight available in these early accounts.

Don Diego de Vargas, 1694. The 1694 journal of Don Diego de Vargas survives as the earliest written account of the San Luis Valley. The journal is a wealth of information about the native peoples, topography, and environment (Colville 1995). After leaving Santa Fe, De Vargas followed the North Branch of the Spanish Trail northward, traveling east of the Rio Grande, and entering the valley just southeast of Ute Mountain. From there he continued north, crossing what would become the New Mexico and Colorado State line and paralleling the western side of San Pedro Mesa before heading west along Culebra Creek. When he reached the Rio Grande, he turned south and crossed the river about five miles south of the confluence. His return trip to Santa Fe took him along the Rio San Antonito on the west side of the Rio Grande, exiting the valley on the west side of San Antonio Mountain (Colville 1995).

His six days in the valley included contact, trade, and occasional skirmishes with the Utes and confrontations with Taos Puebloans. He also documented large herds of bison and some “very large deer.” This reference is the earliest known historical account of bison in the valley (Colville 1995), the last being a brief mention of bison by Juan Bautista Silva along the Rio San Antonio south of present-day Antonito in the spring of 1859 (Kessler 1998). During de Vargas’s travels, the use of sign language and smoke signals for communication is well documented, as is the need to be near water during midsummer.

Notable features of the de Vargas journal include the advantageous yet temporary alliance of de Vargas's men with the Utes and Apaches to combat a mutual enemy: the Comanche. As he traveled along the west side of the valley, de Vargas refers to the San Juan Mountains by their early Spanish name: Sierra de la Grulla, or Mountains of the Cranes. And, in an interesting meteorological observation, de Vargas states on August 24 that, "From the beginning of the march we suffered from bitter cold"—this during a month that now has an average daytime high temperature in the upper 70s.

Juan Bautista de Anza, 1779. Eighty-five years later in 1779, Juan Bautista de Anza, the Governor and Military Commander of New Mexico, left Santa Fe and headed north to quell the Comanche raids that were devastating Spanish settlements in the region. Traveling by night to avoid detection, de Anza followed the North Branch of the Spanish Trail along the eastern foothills of the San Juan Mountains, crossed Poncha Pass, and then headed east to the plains near Pikes Peak. From there he headed south along the foothills, through the areas that would become Colorado Springs and Pueblo, where he fought several victorious battles with the Comanche. He concluded his campaign by crossing back into the valley at Sangre de Cristo Pass (which is also known as La Veta Pass) and taking the eastern route of the North Branch of the Spanish trail back to Santa Fe (Kessler 1998). He initially entered the valley on August 19, 1779, and by September 4 of that year he had reentered the valley near Fort Garland on his return trip to Santa Fe.

Zebulon Montgomery Pike, 1807. Unlike the earlier Spanish explorers, Captain Zebulon Montgomery Pike entered the San Luis Valley from the east, having traveled west from St. Louis across Missouri, Kansas, and the plains of Colorado. Pike's mission was to map and describe the southern parts of the newly acquired Louisiana Purchase. On January 27, 1807, he and most of his men (except five that were left along the trail because they were unable to walk on their frozen feet) crossed the Sangre de Cristo Mountains and entered the valley near the Great Sand Dunes (Carter 1978, Hart and Hulbert 2006, Ubbelohde et al. 2001). Pike built a simple stockade near where the current town of Sanford is located and stayed there until February 26, when Spanish officials took him prisoner and escorted him down to Santa Fe because "it was necessary his Excellency should receive an explanation of my business on his frontier" (Zebulon Pike, Thursday, February 26, 1807).

Although Pike's journal in the days preceding the ascent into the valley often mentions seeing "a gang of buffalo," including in the Wet Valley, there is no mention of buffalo after he enters the San Luis Valley. In

contrast, deer are often mentioned in the valley and goose was a part of at least one meal. Pike grew fond of the valley and concluded that "it was at the same time one of the most sublime and beautiful prospects ever presented to the eyes of man" (Zebulon Pike, Thursday, February 5, 1807).

Jacob Fowler, 1821 to 1822. The journal of Jacob Fowler, which dates from 1821 to 1822 and which *The New York Times* referred to as "quaint and interesting" (*The New York Times* 1898), is a wealth of information about the environment and the interactions between the various peoples who occupied the valley (Coues 1965). *The New York Times* further describes the journal, just published by noted ornithologist Elliott Coues, as "... a notable contribution to our knowledge of early adventure and pioneering in the Great West. His style is straightforward and his wonderful power of observation has made the narrative very attractive."

Fowler was a fur trader who left Fort Smith, Arkansas, in September 1821 and entered the valley via La Veta Pass on February 4, 1822. For the next 3 months, he traveled between Taos and the central part of the valley, going as far north as near where Fort Garland would be later established. Many animals are noted in the valley, including beaver, elk, deer, bear, pronghorn, otter, bighorn sheep, wild horses, geese, ducks, and a wolf. Although great herds of "buffelaw" were noted as the party crossed the Plains, and as far west as the Wet Valley, there is no mention of them once they reach the San Luis Valley. As with the references to animals, the descriptions of plants, particularly the distribution (or lack thereof) of cottonwoods and willows along specific creeks, is frequent and often detailed. These descriptions are mixed with wonderful accounts of life in the many small Spanish settlements that dotted the landscape and interactions with the native peoples.

Fowler recorded an exceptionally astute observation while crossing the southern part of the valley on February 18, 1822:

I Have no doubt but the River from the Head of those Rocks up for about one Hundred miles has once been a lake of about from forty to fifty miles Wide and about two Hundred feet deep—and that the running and dashing of the Water Has Worn a Way the Rocks So as to form the present Chanel.

With this Robert Fowler had speculated about some of the complex geological processes that formed the valley—processes that were studied and confirmed a hundred years later.

Many other explorers and settlers visited the valley and left behind journals of varying detail (Hart and Hulbert 2006, Kessler 1998, Preuss 1958, Richmond 1990, Sanchez 1997). Among these are the following:

- George Frederick Ruxton, 1846
- John C. Fremont, 1848 to 1849
- Charles Preuss, 1848 to 1849 (traveling with Fremont)
- Gwinn Harris Heap, 1853
- John Williams Gunnison, 1853
- John Heinrich Schiel, 1853 (traveling with Gunnison)
- Randolph Barnes Marcy, 1858
- William Wing Loring, 1858
- Juan Bautista Silva, 1859

POLITICAL BOUNDARIES, LAND GRANTS, AND PUBLIC LANDS

The San Luis Valley has endured many changes in governance over the last 300 years. Following nearly 12,000 years of sovereignty by various Native Americans, the control (or at least the declared control) and political boundaries of the region shifted continually until Colorado and New Mexico obtained statehood. The brief timeline below summarizes some of these changes in “ownership” of the San Luis Valley:

1598 Don Juan de Onate claims the San Luis Valley and surrounding areas for Spain.

1763 The Treaty of Paris at the end of the French and Indian War divides much of the North American interior between Spain and France. The San Luis Valley is considered Spanish territory.

1803 The Louisiana Purchase is negotiated between the United States and France but the western boundaries are not clarified and remain ambiguous.

1819 The United States negotiates the Adams-Onís Treaty with Spain to clarify the boundaries of the Louisiana Purchase. The San Luis Valley remains part of Spain’s New Mexico Territory.

1821 Mexican War of Independence (1810 to 1821). The valley becomes a part of the new nation of Mexico.

1836 The Republic of Texas achieves independence from Mexico. Texas claims the land in the valley east and north of the Rio Grande. Mexico does not recognize the Republic, disputes this boundary, and continues to claim the entire valley.

1837 The United States recognizes the Republic of Texas, including the San Luis Valley.

1845 The United States annexes Texas, including the San Luis Valley, and Texas achieves statehood.

1848 Following the Mexican-American War (1846 to 1848), the Treaty of Guadalupe Hidalgo establishes the present Mexico-United States border except for the later 1853 Gadsden Purchase (southern Arizona and southern New Mexico).

1850 Amid much controversy about the admittance of free versus slave States, and as a result of the Compromise of 1850, Texas surrenders its claim to New Mexico, and the New Mexico Territory, including the San Luis Valley generally south of the Rio Grande (38th parallel), is established.

1854 The Kansas Territory, which includes the northern part of the San Luis Valley (above the 38th parallel), is established out of unorganized lands of the Louisiana Purchases.

1861 The Colorado Territory is created by the Colorado Organic Act with the same boundaries that would later become the State of Colorado.

1876 Colorado becomes a State.

1912 New Mexico becomes a State.

Beginning in 1833, many Mexican land grants were issued in the valley as a direct result of the political turmoil noted above and the desire for Mexico City to keep control over the distant northern borderlands of their newly independent nation. These land grants were intended to encourage Mexican settlement in the borderlands, thereby dissuading any thoughts of Texas independence and discouraging encroachment by American fur traders.

The first grants consisted of many small parcels along the Conejos River in Colorado in 1833 (Athearn 1985). These small grants were ineffective in establishing permanent settlement, but the much larger 1842 Conejos Grant proved to have more success in persuading the founding of farms and towns. This grant covered more than 2.5 million acres and included all of what would become the Colorado counties of Conejos and Rio Grande with parts of the counties of Mineral, Saguache, and Alamosa. As with other Mexican land grants in the valley, the grants were considered invalid following the Mexican-American War. The Court of Private Land Claims in 1900 ruled against the grantees and negated the claim (Colorado State Archives 2001).

The Sangre de Cristo grant included all of what is now Costilla County and extended a short distance into the current State of New Mexico. The grant consisted

of 1 million acres and was originally awarded to two Mexican nationals in 1844, but following their deaths during the Pueblo Revolt of 1847, the land was sold to Charles (Carlos) Beaubien. Unlike the Conejos Grant, Beaubien's claim to the land was upheld by the courts in 1860. The land was later sold to William Gilpin (Colorado's first Territorial Governor) in 1864. Large tracts of the grant have been sold to various developers and disputes over the rights of local people to use the land have continued through 2009 (The Center for Grant Studies 2003, The Pueblo Chieftain 2009).

The Baca "Land Grant" in the San Luis Valley was the result of a land dispute. The Baca land patents, of which there are five, were granted to the heirs of Luis Maria Baca in replacement for his 1825 grant near Las Vegas, New Mexico, which was also claimed by Juan de Dios Maiese in 1835. These conflicting claims came to light when the United States took control of the lands in the mid-1840s. The Baca claim was settled in 1860 and patented in 1903, when the Baca heirs were given five parcels of land: two in New Mexico, two in Arizona, and one in the San Luis Valley—Baca #4. In various configurations and sizes, the Baca #4 lands have changed hands many times over the ensuing hundred years, with a large part established as the Baca National Wildlife Refuge in 2000.

The broader San Luis Valley region contains more than 40 percent public land. This includes large parts of the Rio Grande and the Pike-San Isabel National Forests in Colorado, with small sections of the Carson National Forest in New Mexico. The National Forest System was established at the turn of the 20th century as the American public became alarmed at the destruction of forests by timber and mining interests. The BLM was established in 1946 as a result of combining several agencies and policies into one bureau and now owns large parcels of land in the area, primarily in the western and northern parts of the valley floor. Great Sand Dunes National Park and Preserve was initially established as a national monument in 1932 and was expanded to include many upland parcels in 2004. Three national wildlife refuges, Monte Vista (1953), Alamosa (1962), and Baca (2003), were established to protect wetland habitat for migratory birds along the central flyway. Additional lands are owned by the Bureau of Reclamation and the State of Colorado.

NATIVE PEOPLES

The postcontact history of Native Americans in the San Luis Valley involves both cooperation and conflict and ends with the establishment of reservations outside of the valley. Although several Native American tribes are now represented in the valley, today they compose less than 1 percent of the current population.

The Utes consist of several bands and at the time of contact were the primary Native American inhabitants of much of Utah, central and western Colorado,

and parts of northern New Mexico. Increased settlement after the United States gained possession of the valley in 1848 and the surrounding gold rush of 1859 brought new people to the valley and ushered in several decades of escalating pressure to remove the Utes (Ellis 1996). Fort Massachusetts (1852–58) and Fort Garland (1858–83) were established in the valley primarily to protect settlers from Ute attacks. The 1863 and 1868 treaties between the United States and the Utes gave parts of Colorado, including the San Luis Valley, to the United States. Over the next four decades, a series of treaties and agreements continued to reduce Ute lands and relocate the Ute peoples, with the eventual establishment of three reservations in southwestern Colorado and northern Utah by the early years of the 20th century.

Many other Native Americans visited or lived in the valley, including the Apache, Arapaho, Cheyenne, Comanche, Kiowa, and Navajo (NPS 2011). Early historical accounts frequently mention various members of pueblos along the Rio Grande coming north into the central San Luis Valley to hunt bison, causing occasional confrontations with the Utes (Carson 1998, Colville 1995). The first Pueblo revolt of 1680, a response to the expanding Spanish control in northern New Mexico, effectively ceased Spanish rule in the region until Don Diego de Vargas reestablished control over the pueblos in 1692 and 1696. The Taos Pueblo rebelled against the occupation of United States troops during the Mexican-American War in 1847, but the rebellion was soon repelled, effectively ending major conflicts in the region.

SETTLEMENT

Settlement of the San Luis Valley reflects cultural, economic, and political influences as well as creative adaptation to a unique environment. Following the 1610 establishment of Santa Fe as the capital of the New Mexico province, explorers and traders slowly made their way north into the central San Luis Valley. Jacob Fowler encountered several small Spanish settlements during his travels north of Taos and into southern Colorado in 1821 and 1822 (Coues 1965).

The Catholic Church, which was a primary influence during the initial exploration of the region, continued to play a major role in the establishment of settlements and in the day-to-day lives of most of the inhabitants. Members of various church orders were often part of the early explorations, such as the 22 Franciscans who accompanied de Onate during his 1598 exploration and settlement in northern New Mexico (Athearn 1989). The church was instrumental not only in matters of faith, but also as educators, trade coordinators, keepers of public records, and builders of comparatively grand architecture. On the other hand, the oppressive condemnation and suppression of the Native American religious practices were a major contributor to



Meg Van Ness / USFWS

The ranching heritage extends back into the 17th century and is an important part of the San Luis Valley's history and culture, as evidenced by the National Historic Register-listed Trujillo Homestead on Baca Refuge.

the unrest that led to the Pueblo Revolt of 1680 and the destruction of several missions. Nonetheless, the Catholic church began the 18th century as one of the few institutions in the area to prosper, and soon missions were established throughout the region (Athearn 1989). The journals of a Jesuit order near Conejos from 1871 to 1875 reveal days full of baptisms, marriages, deaths, prayers, attending to the sick, and rituals, with a persistent concern for obtaining basic supplies (Stoller and Steele 1982).

In her 1997 book on the San Luis Valley, Olibama Lopez-Tushar describes the first attempted settlement of the valley as that of George Gold (Gould) near the town of Costilla in 1848 (Lopez-Tushar 1997). This settlement was found to be in trespass of the lands held by the Sangre de Cristo Grant and Gold was evicted before establishing a colony, although the town of San Luis de Culebra was established on the land grant 3 years later (Athearn 1985, Wyckoff 1999). The establishment of towns on the land grants was encouraged and within a few years the towns of San Pedro, San Acacio, Chama, and San Francisco were on the Sangre de Cristo Grant and the towns of Conejos, Guadalupe, Ortiz, and Magote were on the Conejos Grant.

Early settlements in the valley were established based on the traditional pattern of the Spanish plaza with homes, churches, and public buildings clustered around a central square and long narrow fields radiating out around the buildings and fronting a nearby creek, sometimes referred to as cordillera or plaza farming (Colville 1995). The extensive systems of early irrigation canals and water control structures supported small grain fields and gardens, some of which are still in use today. Several large canals and their associated laterals, including the Travelers Canal, the Empire Canal, and the Monte Vista Canal, were built in the 1880s in response to the increasing demand for

the valley's beans, corn, grains, and other vegetables. The extensive irrigation in the valley was recognized early as a source of future problems as noted by Major John Wesley Powell in his 1890 testimony before the Senate Special Committee on Irrigation and Reclamation of Arid Lands:

Passing into New Mexico, then, the water that practically heads in the high mountains of Colorado is largely, almost wholly, cut off from the Rio Grande, so that no portion of the water that heads in these mountains where there is great precipitation will cross the line into New Mexico (in the dry season). In a dry season, nothing can be raised in the lower region and sometimes the dry seasons come two or three together. (Siebenthal 1910)

The mining boom in the surrounding mountains in 1859, the completion of the Denver and Rio Grande Railroad over the Sangre de Cristo Mountains and into the valley in 1877, and a vigorous advertising effort by land speculators led to a slow but steady increase in population in the latter half of the 19th century. Before the discovery of gold in 1859, the valley was the home of Colorado's largest non-Native American population, and by 1870 the population of Conejos, Costilla, and Saguache Counties is estimated to have been approximately 5,000 (Wyckoff 1999). Speculators capitalized on the increasing number of immigrants heading west from the eastern United States and Europe, as is illustrated by the description of the valley in a 1884 promotional brochure:

Society is very good. The intelligence of average western people is far above those of the eastern States. Under the duck or buckskin coat of many a miner, farmer or stockman of Colorado is concealed diplomas from the best colleges of the east and Europe.

The climate is almost perfect. Extremes of heat or cold are unknown, and the land is one of almost perpetual sunshine by day, and cloudless skies at night. The healthfulness of the country is notorious, sickness almost unknown. No malaria, no cyclones, no deluges, and when the orchards of small fruits, apples, cherries and plums, and groves of shade trees are planted, the country will be as fruitful and beautiful as the land of Italy. (The Republican Publishing Company 1884)

By the early 1870s, the effect of hunting and development was already taking a toll on Colorado's wildlife. In 1872, the Colorado Territorial Governor Edward N. Cook passed the first game laws to protect certain birds, bison, deer, elk, and bighorn sheep

(Colville 1995). His words sounded the alarm that the wildlife needed protection:

I desire to say a word in favor of protecting our game—birds, beasts, and fishes—all of which are being wastefully destroyed ... and unless some law is passed ... the buffalo, elk, deer antelope and trout will soon become extinct, and Colorado will be robbed of the many attractions she today possesses.

SUMMARY OF KNOWN HISTORIC RESOURCES

Information about the recorded resources in the Colorado part of the San Luis Valley is summarized from data obtained from the Colorado Office of Archaeology and Historic Preservation in February 2012. Similar trends can be extrapolated for the New Mexico part of the area. The Office of Archaeology and Historic Preservation data represent the efforts of hundreds of agencies, organizations, and individuals to document and study the past. The counts include sites, buildings, structures, and isolated finds; however, an individual resource may have many of these elements and may represent more than one time period (multicomponent) and therefore may be counted more than once. It is also important to note that the distribution of the known resources often shows where modern activities have mandated cultural resource surveys and may also potentially show recorder bias as much as actual prehistoric or historic settlement or use patterns.

A total of 6,490 cultural resource sites or properties have been recorded in the Colorado part of the San Luis Valley. Another 2,740 isolated artifacts or features have also been recorded in this area. These resources include 4,719 prehistoric components, 4,091 historic components, 62 components lacking a temporal designation, and 3 paleontological locations, with some resources representing multiple components.

Nearly 20 percent of the prehistoric components are lithic scatters. These locations consist of stone tools or the remains associated with stone tool manufacture. Camps, which are lithic scatters in association with the remains of a campfire, are only slightly less common and have been recorded at approximately 19 percent of the sites. The third most frequent prehistoric site type, representing 4 percent of the sites, is architectural, and generally consist of stone circles or alignments. Other relatively frequent site types found in the valley but never consisting of more than 1 percent include peeled trees, rock art, and human burials. More than half of the prehistoric components on sites in the valley have not been classified into a particular type.

The 4,091 historic components include standing buildings or structures or historic archaeological deposits. Many of these are homes, commercial

buildings, or public buildings within the towns in the valley, with 100 or more each recorded in Alamosa, San Luis, and Monte Vista. Rural sites with historical components often include water control structures (111 recorded), cabins or homesteads (68 recorded), roads or trails (62 recorded), and railroad-related features (28 recorded). The 1,635 historical archaeology components include both isolated rubbish scatters and small features in addition to artifacts or deposits associated with a building or structure.

Two resources in the valley have been designated as National Historic Landmarks. These include Pike's Stockade (5CN75) from 1808 and the Pedro Trujillo Homestead (5AL706) from the late 19th century. Approximately 100 cultural resources in the valley are listed on the National or State Register of Historic Places. Another 435 resources are officially eligible to be listed on the National or State Registers but have yet to be formally nominated.

Socioeconomic Environment

This section describes the socioeconomic profile, land use and ownership, SLVCA land conservation efforts, and water law in the area.

SOCIOECONOMIC PROFILE

The following narrative contains information about the area's population and economics, including employment and the major industries of agriculture, recreation, and tourism.

Population

The combined SLVCA and SCCA spans nine counties: Alamosa, Conejos, Costilla, Hinsdale, Mineral, Rio Grande, and Saguache Counties in Colorado and Rio Arriba and Taos Counties in New Mexico. Table 1 lists population statistics for these counties. The nine-county region has a population of roughly 120,000 people (U.S. Census Bureau 2010a). The counties in Colorado are home to approximately 47,000 residents. Most Hinsdale County residents live in the Gunnison river drainage. Consequently, four of the six counties in the Rio Grande drainage experienced population declines from 2000 to 2010. Slow growth may be the result of increasing unemployment, decreasing non-residential construction, and declining prices of key agriculture commodities (such as barley, alfalfa, and potatoes in 2009) (Colorado Legislative Council Staff 2011). From 2000 to 2010, the nine-county region experienced a 2-percent increase in population, representing slow growth relative to the statewide figures for Colorado (which had a 17-percent increase from 2000 levels) and New Mexico (which had a 13-percent increase from 2000 levels).

Table 1. Population statistics for counties in Colorado and New Mexico that contain the San Luis Valley Conservation Area.

	<i>Residents (2010)</i>	<i>Persons per square mile (2010)</i>	<i>Percentage population change (2000–2010)</i>	<i>Percentage population change (2010–2025)†</i>
Colorado	5,029,196	48.5	17	26
Alamosa County	15,445	21.4	3	25
Conejos County	8,256	6.4	–2	10
Costilla County	3,524	2.9	–4	8
Hinsdale County	843	0.8	7	26
Mineral County	712	0.8	–14	16
Rio Grande County	11,982	13.1	–3	7
Saguache County	6,108	1.9	3	18
New Mexico	2,059,179	17	13	19
Rio Arriba County	40,246	6.9	–2	11
Taos County	32,937	15	10	17

Sources: U.S. Census Bureau 2010a and †Colorado Department of Local Affairs 2002, University of New Mexico 2002.

Population growth in the San Luis Valley region is expected to continue at a slow pace over the next decade. From 2010 to 2025, the population of the local area is projected to increase by 14 percent, indicating slow growth compared to the projected statewide figures for Colorado (which has a projected 26-percent increase) and New Mexico (which has a projected 19-percent increase) (Colorado Department of Local Affairs 2002, University of New Mexico 2002). In the SLVCA, the smallest projected increases are in Costilla County (8 percent) (Colorado Department of Local Affairs 2002, University of New Mexico 2002).

Race, Ethnicity, and Education

Hispanic and Latino residents (57 percent of the total population) represent the largest ethnicity in the nine-county San Luis Valley region. The prevalence of this ethnic group is because of the presence of two large Hispanic communities in the local area. The region is home to a large population of White residents who consider themselves to be of Hispanic or Latino ethnicity. This is particularly true in Alamosa, Conejos, Costilla, Saguache, Rio Arriba, and Taos Counties, where, collectively, White Hispanics represent 32 percent of the county-wide population on average (U.S. Census Bureau 2010a). The occurrence of this race-ethnicity pairing in the San Luis Valley may be because of residents of Hispano heritage (such as descendants from Spaniards) (Sangre de Cristo National Heritage Area 2012). Hispanics of Mexican descent also represent a substantial share of the population in Costilla County (34 percent) (U.S. Census Bureau 2010a).

Whites (including Whites of Hispanic and Latino origin) represent the largest race in the nine-county region (66 percent of the total population). Native Americans and Alaska Natives account for 8 percent

of the total population of the region. Collectively, Black or African-American residents, Asians, and native Hawaiians and other Pacific islanders account for about 1 percent of the total population of the region (U.S. Census Bureau 2010a).

Table 2 shows the percent of the population that has obtained a bachelor's degree or higher within each of the SLVCA States and counties. Of the two States, Colorado has the highest percentage of individuals with a bachelor's degree or higher (36 percent of the population), followed by New Mexico (26 percent) (U.S. Census Bureau 2010a). Costilla County residents were less likely to hold a bachelor's degree than the average Colorado resident; in New Mexico, the opposite is true for Taos County (30 percent of the county-wide population with a bachelor's degree or higher) relative to the State average (26 percent) (U.S. Census Bureau 2010a).

Regional Economy, Employment, and Income

Table 2 also shows median household income and poverty rates for each of the SLVCA States and counties. Among the two States, Colorado had the highest median household income in 2010 (\$56,456 per year), followed by New Mexico (\$43,820 per year) (U.S. Census Bureau 2010b). At a statewide level, New Mexico had the highest poverty rate at 18.4 percent, and Colorado had the lowest at 12.2 percent. However, the San Luis Valley is one of the most impoverished regions of Colorado. Costilla County has the highest poverty level, more than twice the State average, and its median household income is less than half the State average at \$24,388 per year. Taos County has somewhat higher median household income (\$35,441 per year) and its poverty level is lower than the State of New Mexico's. (U.S. Census Bureau 2010b).

Table 2. Income, education, unemployment, and poverty rates for counties in Colorado and New Mexico that contain the San Luis Valley Conservation Area.

	Median household income (average \$2006–2010)†	Percentage bachelor's degree or higher†	Percentage unemployed‡		Percentage of individuals below poverty (average 2006–2010)†
			2008	2011	
Colorado	56,456	36	4.8	7.9	12
Alamosa County	35,935	27	5.2	7.5	24
Conejos County	33,627	19	7.3	9.5	18
Costilla County	24,388	14	7.7	12.4	28
Hinsdale County	74,659	42	3.4	6.1	4
Mineral County	53,438	39	5.4	7.3	8
Rio Grande County	39,871	19	5.6	7.8	17
Saguache County	30,430	19	7.2	9.9	24
New Mexico	43,820	26	4.5	6.6	18
Costilla County	41,437	16	5.4	8.9	20
Taos County	35,441	30	5.5	10.4	17

Sources: †U.S. Census Bureau 2010b and ‡Bureau of Labor Statistics 2011a, Bureau of Labor Statistics 2011b, Bureau of Labor Statistics 2008.

Table 3 shows the percent of employment by sector within the San Luis Valley region. The combined nine-county region had a total employment of more than 62,000 individuals in 2009 (U.S. Department of Commerce 2009). The highest percentage of total employment in 2009 was in public administration (18 percent of total local employment), the second highest was in the arts, entertainment, recreation, and accommodation and food services (11 percent), and the third highest was in agriculture, forestry, fishing, hunting, and mining (11 percent) (U.S. Department of Commerce 2009).

Agriculture, Recreation, and Tourism

Agriculture is a prominent industry in the San Luis Valley. Crops grown in the valley include alfalfa, native grass hay, wheat, barley, sorghum, canola, spinach, lettuce, carrots, and potatoes (Colorado Division of Wildlife 2010). Agriculture, forestry, fishing, hunting, and mining accounted for roughly 11 percent of the total jobs in the region in 2009 (U.S. Census Bureau 2009). The total number of agricultural jobs in the local area increased from about 3,700 jobs in 1970 to 4,446 in 2009 (U.S. Department of Commerce 2010a). Costilla County, Colorado, had the largest percentage of employment in agriculture in the region (22 percent) (U.S. Department of Commerce 2010a, U.S. Department of Commerce 2010b; data compiled using the EPS–HDT system [Headwaters Economics 2011]). Approximately 29 percent of the land in the nine-county region is in agriculture (U.S. Department of Agriculture 2009, data compiled using the EPS–HDT system [Headwaters Economics 2011]).

Tourism is a cornerstone of the local economy, and the tourism industry in the San Luis Valley shows strong development potential. With a diverse collection of natural and heritage assets, the local tourism industry is able to cater to a variety of recreationists, including outdoor recreationists; visitors to the Great Sand Dunes National Park and Preserve; resort tourists; vacation and second home owners; ecotourists; heritage, arts, and cultural tourists; and visitors who pass through the area on their way to other regional attractions (Center for Rural Entrepreneurship 2008). According to the 2006 National Survey of Fishing, Hunting, and Wildlife-Associated Recreation, approximately 3.1 million residents took part in wildlife-associated recreation activities in Colorado and New Mexico in 2006 (USFWS 2008). It was estimated that residents and visitors combined spent \$3.8 billion on wildlife-associated recreational activities in 2006 in the two States combined, with Colorado accounting for approximately 79 percent of this spending. Among participants, wildlife watching was the most frequently reported activity, followed by fishing and hunting. In Colorado, 82 percent of individuals surveyed watched wildlife, 30 percent fished, and 12 percent hunted, while in New Mexico, 83 percent watched wildlife, 26 percent fished, and 10 percent hunted (USFWS 2008).

LAND USE AND OWNERSHIP CHANGES SURROUNDING THE CONSERVATION AREA

The current land use and changes in land use are described below.

Table 3. Percentage employment by sector for counties in Colorado and New Mexico that contain the San Luis Valley region.

<i>Employment sectors</i>	<i>Percentage of nine-county region employed</i>
Total employment in 2009 ^a was 62,121	
Agriculture, forestry, fishing, hunting, and mining	11
Arts, entertainment, recreation, and accommodation and food services	11
Construction	6
Educational services, health care, and social aid	8
Finance and insurance, real estate, rental, and leasing	6
Information	1
Manufacturing	2
Other services, except public administration	4
Professional, scientific, management, administrative, and waste-management services	5
Public administration	18
Retail trade	10
Transportation and warehousing, and utilities	2
Wholesale trade	2

Source: U.S. Department of Commerce 2009

^aNot every sector category for every county was fully disclosed because of confidentiality requirements; the table reflects the best and most correct information available.

Current Land Use

The San Luis Valley is a large intermountain basin covering approximately 3,200 square miles of land in southern Colorado and northern New Mexico. The valley is bordered by the Sangre de Cristo Mountains to the east and northeast, the San Juan and La Garita Mountains to the west and northwest, and the Taos Plateau to the south. Snowmelt from the mountains around the valley is responsible for most of the area's stream flow in the associated watershed, including the Rio Grande and Conejos Rivers (Emery [no date]). The valley floor is primarily grassland and shrubland, while the hills surrounding the valley are forested. Collectively, grasslands (40 percent of all land cover in the nine-county region), forests (30 percent), and shrublands (22 percent) account for most of the land cover in the local area (NASA 2006; data compiled using the EPS-HDT system [Headwaters Economics 2011]). Approximately 56 percent (2,944,353 acres) of the project area is in private ownership. The remaining acres are protected and managed by the Service, USFS, BLM, National Park Service, and State of Colorado. Most of the private land and wetland habitat occurs on the valley floor, creating one of the largest intermountain valleys in the world (USFWS 2010a).

The nine-county region is relatively rural, and population densities in the San Luis Valley are among the lowest in Colorado. Only 2 percent of land cover in the region area is urban (NASA 2006, data compiled using the EPS-HDT system [Headwaters Economics 2011]), U.S. Census Bureau 2010a). Major municipalities

in the region include Alamosa, Crestone, Del Norte, Monte Vista, Saguache, and San Luis.

Changes in Land Use

The SLVCA contains a rich diversity of trust species and habitat types. The San Luis Valley is the southernmost significant waterbird production area in the central flyway and is the most important waterfowl production area in Colorado. According to Partners in Flight, riparian habitats in the region support the highest bird diversity of any western habitat type (USFWS 2010a).

Historically, land use remained unchanged in the San Luis Valley until the early 1800s, when Euro-American settlement began to alter the presettlement landscape (USFWS 2010a). During this period, livestock grazing, farming, and water development began to affect ecosystem processes such as the natural hydrological regime. Since then, nearly 50 percent of Colorado's wetlands have been lost (Dahl 1990, 2000).

The highest remaining concentration of wetlands in Colorado occurs in the San Luis Valley, and protection of every remaining wetland acre is a high priority (USFWS 2010a). Manipulation of the natural hydrological cycle in the San Luis Valley for agricultural purposes likely has resulted in a significant conversion of upland habitat to wetland habitat through the extensive development of flood-irrigation infrastructure. These conversions have benefitted many wetland-dependent species, with tradeoffs. The results of these conversions of uplands to wetlands included the

shortening of some streams and the alteration of annual hydrographs in many stream reaches, which have lowered the peak in the hydrograph yet increased flows after the irrigation season due to unconsumed water returning to the stream system. Stream depletions from the use of wells have compounded these effects. Chronically reduced levels of runoff from the San Juan and Sangre de Cristo Mountains have increased all of these effects on streams. Most wetland habitat in the SLVCA is on private ranch and farm lands and relies on surface and ground water diversions to maintain its value to wetland-dependent wildlife. Through the SLVCA, we hope to maintain these land management practices with willing landowners to support priority wildlife habitat.

In recent years, human population growth has occurred in rural areas causing fragmentation of wetland habitat types throughout the SLVCA. Statewide, Colorado's population grew by more than 36.9 percent between 1990 and 2003. During that same time, Mineral County's population grew by 57.9 percent and Saguache County's by 45.2 percent. While population densities are still low in these counties, the amount of habitat loss and fragmentation resulting from subdivision of lands for residential and commercial development is a significant threat to trust wildlife species in the SLVCA.

Exurban housing development typically occurs within riparian areas in the SLVCA. This is detrimental

to wildlife habitat because riparian habitats account for only a limited portion of Colorado's landscape, but contain high species richness and abundance. Additionally, most of the wetland and riparian habitat in the SLVCA occurs on private ranchland and farmland. The recent downturn in the economy coupled with depressed agricultural markets have increased the rate of exurban development, as some ranchers and farmers have been forced to subdivide their properties to continue operating (USFWS 2010a).

As agricultural lands are subdivided, the resulting fragmentation can affect habitat use for a wide array of waterfowl, shorebird, colonial waterbird, and songbird species. Many of these species require specific habitat conditions for successful reproduction and building energy reserves for breeding and migration. As habitats are lost, the spatial juxtaposition of available habitat is altered, disrupting wildlife movement, dispersal, and migration patterns.

In addition to the direct loss of wildlife habitat from fragmentation, the water rights associated with these properties are subject to sale, potentially resulting in not only the loss of wetland habitat and wetland functions on the subdivided property but also on adjoining lands as the water is redistributed off of the property. Keeping the current connectedness of habitat through permanent protection will limit the risk for disruption of species' movement patterns because of fragmentation. Connectivity will also keep important migration



USFWS


Agricultural practices such as haying and grazing are a primary part of the economy in the San Luis Valley, and often provide habitat for wildlife as well.

corridors and linkages between seasonal ranges necessary to meet the life history needs for many species (USFWS 2010a).

Currently, the landscape of the San Luis Valley has not been altered to the extent found in many other western regions with more rapid population growth. The downturn in the national and regional economy has slowed population growth and development pressures. Although the overall population of the project area only increased by 2 percent between 2000 and 2010, it is projected to increase by 16 percent between 2010 and 2025, with greater increases in Alamosa, Hinsdale, and Taos Counties (U.S. Census bureau 2010a). Any pressures on the agricultural economy of the San Luis Valley can result in conversion of agricultural land to other uses potentially resulting in fragmentation and loss of critical wildlife habitat, including riparian habitat. Whether or not agricultural lands change ownership, there is potential for water to be sold and transferred off the property, often to the detriment of wildlife habitat, but only after a change of water right is filed and approved by the court.

As explained previously, the overall population in the SLVCA increased by only 2 percent between 2000 and 2010. The largest increase in population growth occurred in Hinsdale County (7 percent increase from 2000 levels) and Taos County (10 percent increase from 2000 levels), while five of the nine counties in the region (Conejos, Costilla, Mineral, Rio Arriba, and Rio Grande) experienced population declines during these years (U.S. Census Bureau 2010a). From 2010 to 2025, the population in the SLVCA is projected to increase by 16 percent, indicating slower growth relative to the projected State-level increases of 26 percent for Colorado and 18 percent for New Mexico (Colorado Department of Local Affairs 2002, University of New Mexico 2002). However, the population is projected to increase at rates similar to the Colorado State average in Alamosa County (25 percent increase) and Hinsdale County (26 percent increase) and above the New Mexico State average in Taos County (21 percent increase).

In 2000, the American Farmland Trust identified 4.9 million acres of prime ranchlands in Colorado and 2.6 million acres in New Mexico as being vulnerable to low-density development by the year 2020. Within the Rocky Mountain region (which includes 263 counties in Idaho, Montana, Wyoming, Utah, Colorado, Arizona, and New Mexico), Saguache County, Colorado, and Rio Arriba County, New Mexico (both close to the project area), ranked in the top 25 counties for acres of strategic ranchland at risk (American Farmland Trust 2000). While population densities are still low in these counties, development has been occurring within sensitive riparian areas in the valley floor. Taking added steps to conserve wildlife habitat in the San Luis Valley now, while land prices are still



© Joe Zimm

Much of the plant and animal diversity of the San Luis Valley is dependent on its sensitive wetland habitats.

affordable and irreplaceable habitat has not been lost, may be proper. Protecting this land from residential and industrial development is the only way to ensure the long-term resiliency of the ecosystem and support viable wildlife populations and habitats in the face of climate change and other threats (USFWS 2010a).

Water quantity, quality, and use issues are major threats to the sustainability of wetland and riparian habitats in the SLVCA. Changes in water quality and quantity have adverse effects on the function of the wetland complex located in the valley floor. There are, for example, growing concerns about the impacts of new contaminants, such as endocrine-disrupting chemicals, that can affect water quality on both private and public lands (USFWS 2010a).

Ground water usage, especially artesian well development, started during the early 1900s. The result has been the construction of more than 7,000 wells in the San Luis Valley and development of one of the world's largest concentrations of center pivot irrigation systems, some of which depend solely upon ground water. As a consequence, water users and regulators have acknowledged that annual ground water use chronically exceeds recharge under the current hydrologic cycle. New ground water use rules were developed by the Colorado Division of Water Resources in late 2015 and may soon be applied to well owners in the San Luis Valley (USFWS 2010a).

Once the new ground water rules are carried out, ground water users will be responsible for eliminating injury to senior water rights through a process with the State (USFWS 2010a). The Rio Grande Water Conservation District, who is working with ground and surface water users, is offering an alternative approach of establishing ground water management subdistricts. These self-taxing entities will be responsible for collectively replacing injurious depletions to senior water right holders caused by the ground water use of its members and ensuring ground water use within the subdistricts is sustainable. In most cases, this will require ground water users to acquire and, in many cases, remove senior water rights from other properties to replace depletions from well use.

These circumstances threaten healthy riparian systems along the tributaries of the Rio Grande, including in the SLVCA, where senior water rights are now used to irrigate meadows and grasslands in the floodplain. The evolving economic and regulatory environment in the SLVCA may likely result in the acquisition of some of these water rights to augment distant wells, moving water out of the floodplain and degrading migratory bird habitat (USFWS 2010a).

SAN LUIS VALLEY CONSERVATION AREA LAND CONSERVATION EFFORTS

Land protection is a relatively new practice in the San Luis Valley, as most conservation easements have been completed within the last 10 years. However, during this short timeframe, more than 232,000 acres of land have been protected in the region, which suggests that public support for land protection in the SLVCA is strong (USFWS 2010a). In fact, there are so many landowners interested in entering into conservation easements that organizations like the Rio Grande Headwaters Land Trust, The Nature Conservancy, Ducks Unlimited, and NRCS are challenged to meet the demand (USFWS 2010a). Citizens of the San Luis Valley understand that the rural lifestyle and wildlife habitat is what makes this area unique and have voiced their concern over the loss of these values. They recognize that conservation easements are a tool to keep both ranches and wildlife habitat intact (USFWS 2010a).

The Service plans to conserve up to 280,000 acres to protect the remaining expanses of wildlife habitat in the SLVCA. This will be accomplished primarily through the purchase of conservation easements by the Service on a voluntary basis from private landowners. Other Federal, State, and nongovernmental partners may assist in acquiring conservation easements or fee title to a lesser extent. On a limited basis, we may use fee-title acquisition to acquire land and water to facilitate more efficient management of the three national wildlife refuges. For example, we hope to explore the feasibility of acquiring more land

on the west side of the Rio Grande next to Alamosa Refuge to reduce fencing problems and enable consistent management of this important stretch of riparian habitat. Examples of water acquisitions we are considering are (1) those needed to maintain current ground water rights on the refuges, (2) those rights associated with any fee-title land acquisitions, and (3) purchase of more shares in mutual ditch companies that currently serve the Alamosa and Monte Vista Refuges because of the need to also replace depletions from existing ground water uses.

Acquisition of these lands will occur over a period assumed to range from 15 to 20 years but, based on past acquisition rates, could reasonably be expected to occur over a longer period, possibly up to 100 years.

Conservation Easements

One of the Service's high-priority objectives is to guide residential and commercial development away from high-priority conservation areas by securing strategic conservation easements. The SLVCA will focus on the protection of wetlands, riparian areas, montane forests, and sagebrush habitats on private land within the area through acquisition of conservation easements from willing sellers. Conservation easements leave land in private ownership, protecting private property rights, while providing the Service with a cost-effective conservation strategy that enables the conservation of large blocks of habitat.

A conservation easement is a voluntary legal agreement entered into between a landowner and a conservation entity. Conservation easements are binding in perpetuity; the landowner reserves the right to sell or bequeath the property, but the easement and its associated restrictions remain with the property forever. Owners of land that does not contain a conservation easement have a set of rights associated with their land. For example, landowners have the right to run cattle, grow crops, harvest trees, build structures, and subdivide and sell their land. Under a conservation easement, landowners keep ownership of their property, but transfer some of their ownership rights to the conservation entity. The most common right transferred under a conservation easement is the right to develop or subdivide the land.

Conservation easements in the SLVCA may require the transfer of more rights. A conservation easement on a parcel of land may have restrictions for all types of human development, such as surface disturbance from solar, mineral, or wind energy development, depending upon the particular wildlife values of the habitat.

In most cases, a conservation easement acquired for wetland values will be associated with appurtenant irrigation water rights that have resulted in desirable wildlife habitat. Doing anything less may often result in separation of water use from the land, reducing the easement's value to trust wildlife species.

Several nonprofit organizations active in the SLVCA that hold many easements are already acquiring the property rights to separate water from the land subject to the easements. Water use must respect State requirements; therefore, water issues will need to be addressed individually for each easement. In all cases, the terms of a conservation easement must be mutually agreed-upon by the landowner and the easement holder. Conservation easements acquired from private landowners will not affect their property rights beyond those purchased through conservation easement.

Subsurface rights are often severed from the surface rights of a parcel of land. In such cases, conservation easements apply only to surface rights; therefore, the mineral interest may be extracted at any time by the person who holds the qualified mineral right (Byers and Ponte 2005). For this reason, the Service is unlikely to enter into a conservation easement agreement for a parcel of land that has a viable subsurface mineral interest. Exceptions may be made if the parcel has high habitat value and the probability of mineral extraction is low as determined by a formal remoteness survey.

Fee-Title Purchases

Within the SLVCA, the Service may purchase limited property in fee title at fair-market value to protect up to 30,000 acres of significant wildlife habitat and maintain wildlife populations, plant communities, and ecosystem processes in perpetuity (USFWS 2010a). Under fee-title purchases, full ownership of the land,

including the underlying title, is transferred to another party. This gives the new owner maximum interest in the purchased land and allows the new owner to manage the land in any manner that is consistent with local, State, and Federal laws. For fee-title acquisitions, the Service intends to evaluate the purchase of water rights with each property. We will consider acquiring land in fee title only where the Service's conservation objectives could not be met with conservation easements. The Service will consider fee-title acquisition under the following scenarios:

1. A priority of this plan is the protection of wetlands still further west of Alamosa Refuge—identified as the Alamosa Marshes by the 1874, 1875, and 1877 Wheeler Expedition (U.S. Army Corps of Engineers 1878). The area still provides one of the largest intact wetland complexes in the San Luis Valley and provides significant migration and breeding habitat for waterfowl, shorebirds, and wading birds on average or above water years. We envision that conservation easements—with nonprofit organizations, the Service, or other agencies—will be the primary method of protecting these wetlands. We also recognize that some landowners may be willing to sell property for wildlife conservation purposes but unable or unwilling to pursue easements. To allow for this circumstance, the Service will have the ability to acquire fee-title interest in lands within the Alamosa Marshes.



Sue Olievira / USFWS

The tributaries of the Rio Grande are some of the last refuges of genetically pure populations of Rio Grande cutthroat trout, and provide important nesting and migration habitat for countless birds.

1. An important acquisition idea is to expand the western boundary of the Alamosa Refuge to include riparian habitat along the Rio Grande that is not already part of the refuge. Currently, some of the refuge boundary runs down the middle of the Rio Grande. Other parts of the boundary are located on the east side of the river, while the remainder includes land only on the west side of the river. This configuration makes livestock management very difficult on both sides of the refuge boundary. By including the entire riparian zone within the refuge, we will be able to improve management for riparian habitat species like the southwestern willow flycatcher.
2. The Service needs to have the ability to meet administrative challenges such as those resulting from upcoming ground water regulation that may require the Service to acquire surface water for ground water augmentation purposes. Although Colorado allows water to be sold separately from the land, the owner may choose not to do so. In this situation, the Service's best alternative may be to buy from willing sellers both the land and the associated water rights to achieve our ground water management goals. The Service also wants the ability to make small boundary adjustments to all three national wildlife refuges to prevent refuge management conflicts.

The total lands acquired in fee title by the Service over the life of this plan depend on several variables including the following:

- the priorities and success of our partners in protecting these lands
- the willingness of private landowners to sell any property rights
- private landowner preference of whether to sell fee-title or much more limited property rights
- the availability of refuge operation and maintenance funds
- the availability of land acquisition funding.

This plan places a maximum cap of fee-title acquisition by the Service at 30,000 acres, but these influencing factors will inhibit or preclude the need for such acquisition and dictate the amount of land acquired over the life of the plan.

WATER LAW

To address water rights affecting the SLVCA, the following narrative describes water law in Colorado and New Mexico. In either State, water rights that the Service owns or acquires must conform to the

respective State water adjudication process as required by the McCarran Amendment (43 U.S.C. § 666), 1952.

Colorado

Colorado is divided into seven water divisions using watershed boundaries. Each division has a Water Court and a division engineer who administers water rights by priority. The Rio Grande is in division 3. This method of administration has been in use since 1969, when the original statutes governing water were revised and rewritten.

Water rights in Colorado are subject to the prior appropriation doctrine, which is enshrined in the Colorado Constitution and dates from before statehood in 1876. The first entity to claim the water right has the first right to use the full amount of water they claimed for beneficial use. The prior appropriation doctrine allows State officials to properly manage and distribute water according to the decreed priority dates. There are four elements of a water right under the prior appropriation doctrine: intent, diversion, beneficial use, and priority. An applicant must show that there is intent to use the water, construct the diversion works, and put the water to beneficial use, which establishes a priority date. In Colorado, every water right must be adjudicated through the Water Court. Colorado law now recognizes certain beneficial uses that do not require a diversion, such as instream flows.

If there is not enough water to satisfy all water right holders in a particular stream, the State must shut off junior rights as necessary to make sure that senior water right holders receive their full appropriation. The Rio Grande basin in Colorado is considered overappropriated, meaning that more water rights exist than can be continuously served by the average amount of available water supply.

Ground water in Colorado is designated as either tributary or nontributary. Tributary ground water is water contained in aquifers that have a direct hydraulic connection to surface water. The unconfined and confined aquifers in the San Luis Valley are both considered tributary ground water. Tributary ground water is treated administratively the same as a surface water diversion.

Water rights in Colorado can be transferred from one entity to another, but a change application must be filed and approved by the Water Court. The amount available for transfer is limited to the consumptive use part of the right. Water rights in Colorado are considered real property and they may be bought or sold. A water right can be conveyed either as part of a piece of property or separate from a property. If a water right is transferred separately from the land, it is necessary to file a change application through the Water Court to use it at any other location.

In 1973, the Colorado Legislature passed Senate Bill 97, creating the State's Instream Flow Program. This program, one of the first of its kind, vested the Colorado Water Conservation Board with exclusive authority to protect streamflow through a reach of stream rather than just at a point, and to protect levels in natural lakes. Until this law was passed, all appropriations of water in Colorado were required to divert water from the natural stream.

Since about 1990, Colorado clarified the Colorado Water Conservation Board's authority to acquire existing, decreed senior water rights on a voluntary basis from willing owners for instream flow uses. New appropriations are junior water rights claimed by the board to preserve the natural environment. New appropriations are considered by the board each year and are filed annually with the Water Court for adjudication. New appropriations are generally limited to the minimum amount necessary to protect the natural environment to a reasonable degree.

New Mexico

New Mexico's water law is also based on the doctrine of prior appropriation. All waters in New Mexico are declared to be public and subject to appropriation for beneficial use. Apart from water rights acquired before 1907 and small-scale stock watering (10 acre-feet or less), a permit from the State engineer is required to appropriate water, change the point of diversion, change the location of wells in declared basins, divert or store water, or change the place or purpose of water use. There is a new requirement in New Mexico that before obtaining a water right involving the use of public lands, the person seeking the right must prove that he or she actually has a permit to use the public lands.

The New Mexico ground water code was enacted in 1931. Ground water procedures closely parallel those for surface water, with several important differences. A permit to drill a well and appropriate water is not required in areas outside of declared "underground-water basins." Within underground water basins, however, use is regulated by the State engineer. The State engineer has the authority to establish these basins when regulation is necessary to protect prior appropriations, make sure that water is put to beneficial use, and support orderly development of the State's water resources. There are now 33 declared underground water basins throughout New Mexico.

Water rights in New Mexico can be transferred from one entity to another, but a change application must be filed and approved by the State engineer. Water rights in New Mexico are considered real property and they may be bought or sold. A water right can be conveyed as part of a piece of property or separate from a property, as long as that water right has been severed from the land by an approved application through the State engineer. Water rights in New

Mexico are considered real property and they may be bought or sold. A water right can be conveyed as part of a piece of property or separate from a property, as long as that water right has been severed from the land by an approved application through the State engineer.

New Mexico has had adjudicated water rights since 1907. In an adjudication suit, each claimant has an opportunity to present evidence of water right to the court. The completion of adjudication results in a court decree outlining the priority, amount, purpose (determination of use), periods, and place of water use.

New Mexico's instream flow program is complex, unclear, and continually evolving. New Mexico does not have a legislated instream flow program, and instream flow is not a recognized beneficial use. Recent case law, however, has allowed the development of an instream flow program in New Mexico. In 1998, the New Mexico Attorney General issued a legal opinion concluding that the transfer of a consumptive water right to an instream flow right is allowable under State law. The legal opinion found that instream uses such as recreation and fish and wildlife habitat are beneficial uses, and that transfers of existing water rights to instream flows are not expressly prohibited. Before this opinion, New Mexico was the only State that did not recognize instream flow as a beneficial use.

The 1998 Attorney General's opinion is limited to the transfer of existing water rights. The opinion notes that new appropriations of water for instream flow are not subject to this precedent. Although the opinion concludes that there are no legal barriers to the transfer of existing water rights to an instream flow right, the State engineer still has the responsibility for approving such a transfer. Although instream flow in itself is not recognized as a beneficial use, it appears that water can be dedicated to instream flow for the purpose of recreation or fish and wildlife habitat.

The Attorney General's opinion does not explicitly address the issue of ownership of instream flow rights. Since ownership of other types of water rights are not limited, it could be interpreted that instream flow rights could be held by a public or private entity. Current law is unclear and continues to develop.

Chapter 3—Threats to and Status of Resources

Threats to Resources

The land cover of the San Luis Valley was largely unaltered, except by natural processes, until the 19th century, when human land use associated with settlers of European origin began to alter the landscape. During this period, livestock grazing, farming, and water development also began to affect ecosystem processes such as the historical hydrological regime. Since then, Colorado has lost nearly 50 percent of its wetlands (Dahl 1990, 2000). The highest remaining concentration of wetlands in Colorado occurs in the San Luis Valley, and their protection is a high conservation priority. Sagebrush-dependent birds are often sensitive to vertical structure in their habitat, and thus the protection of these habitats from development is a priority.

DEVELOPMENT

Population growth, primarily exurban development, led to habitat fragmentation in the San Luis Valley in the latter part of the 20th century and the first part of the 21st century. Colorado's population increased by nearly 17 percent between 2000 and 2010 (U.S. Census Bureau 2010a). The absolute population numbers and densities are still low in the project area, as with the population growth. However, habitat loss and fragmentation because of residential and commercial development remain a threat to trust species in the SLVCA based on population growth projections. This rapid statewide growth has lessened somewhat during the current economic downturn, with relatively stable populations in the counties of the San Luis Valley from 2000 to 2010 (U.S. Census Bureau 2010a). However, subdivision of ranchlands and farmlands will continue and may increase as ground water regulation goes into effect and changes the economic viability of some properties currently in agriculture. It is not the intent of the SLVCA to freeze all residential development at today's level but to give owners of important wildlife habitat an alternative to such development in the effort to protect the highest priority habitat.

As the country increases its dependence on renewable energy sources, we need to take care to minimize the effects on high-priority wildlife habitat. The effects on wildlife populations from solar energy development are of particular concern in the San Luis Valley, as interest in industrial solar electric-generating facilities has increased during the last decade. The San Luis Valley is already the site of two, large photovoltaic projects. Economically viable wind energy potential is generally quite low in most of the valley (Hanser 2010) and thus unlikely to be an issue in the near term. Hydrocarbon potential is low throughout the valley (Copeland et al. 2009), although some oil has been found during mineral exploration (Watkins et al. 1995). There is potential for further oil and gas exploration in this region, which the Service has found is unlikely to have significant impacts on the living resources of the valley (USFWS 2011).

FRAGMENTATION

Changes in land cover because of exurban development, energy development, roads, and changes in agricultural land use (such as transition from flood irrigation to center-pivot irrigation) not only cause a loss of habitat, they also fragment the remaining habitat. There is a robust body of literature on the effects



Steve Torbit / USFWS

The forests along the mountains are a corridor for movement of wide-ranging species such as mountain lions.

of habitat fragmentation, summarized eloquently by Collinge (2009). Countless manipulative and observational studies have shown that habitat area and connectivity among types of similar habitat are important for everything from soil decomposers (Rantalainen et al. 2005) to passerine birds (Telleria and Santos 1995). Corridors between fragments promote use of, and persistence in, those habitats by migratory birds (Haas 1995), large carnivores (Shepherd and Whittington 2006, Tremblay 2001), and ungulates (Tremblay 2001) that are native to the SLVCA. Perhaps the most obvious way to protect corridors throughout the SLVCA, while protecting valuable habitat at the same time, is to focus on the conservation of the riparian corridors that cross and connect existing protected areas. This action will protect wildlife movement corridors for both seasonal migration and colonization following large-scale disturbance or environmental change.

INVASIVE SPECIES

Increased human disturbance associated with development has also been shown to negatively affect adjoining habitat because of the invasion and establishment of invasive plant species. Invasive plants can have many detrimental effects; besides displacing native vegetation, they can alter nutrient cycling and soil chemistry, change hydrology, increase erosion, and

change fire regimes (Dukes and Mooney 2004). Noxious weeds, such as tall whitetop, Canada thistle, and Russian knapweed, can have severe negative effects on wildlife habitat (such as reducing the quality of nesting and foraging areas) when these weed species begin to replace native vegetation. The San Luis Valley already has one of the densest concentrations of Russian knapweed in the State of Colorado (Goslee et al. 2003). Other invasive species that could threaten resources in the SLVCA include New Zealand mudsnail, quagga and zebra mussels, and Asian clam. Diseases such as white-nose syndrome, chytrid fungus, whirling disease, and chronic wasting disease also threaten wildlife and fish in the San Luis Valley.

WATER RESOURCES

In addition to the threats of the direct loss of habitat and fragmentation that accompany subdivision for exurban development, water rights can be sold with the property, or can be severed and sold to other landholders. This can result in the loss of wetland habitat and wetland functions not only on the property, but also on adjoining lands as the water is redistributed off of the property, directly affecting wildlife populations that depend on the wetlands to complete their life cycle. As fragmentation increases, remaining habitats become geographically isolated and wildlife



USFWS

Water is a critical resource to breeding and migratory birds in the high desert. Many wetlands and riparian areas have been lost due to ground water pumping and surface water diversion.

populations with limited dispersal abilities may potentially become genetically and spatially isolated.

Another threat to the sustainability of wetland and riparian habitat in the SLVCA is the chronic overuse of ground water. In 2004, the Colorado Legislature directed the State Engineer to develop ground water rules and regulations for the San Luis Valley to bring use to sustainable levels and protect senior surface water users. The complicated and contentious process of developing and carrying out regulations is well underway. However, there have been delays because of court challenges and daunting technical hurdles in developing the ground water model that will be used to quantify depletions to individual streams caused by ground water use throughout the valley.

Ground water usage, especially artesian well development, started during the early 1900s. The result has been construction of more than 7,000 wells and development of one of the world's largest concentration of center pivot irrigation systems, many of which depend solely upon ground water. As a consequence, water users and regulators have acknowledged that annual ground water use during a period of below-average precipitation exceeds recharge. It is important to note that in addition to traditional agricultural irrigation, the existing national wildlife refuges in the San Luis Valley also use ground water adjudicated for both wildlife and irrigation extensively for irrigation and impoundments to create wildlife habitat. Ways to reduce reliance on ground water are being explored in the planning process for the comprehensive conservation plan for the San Luis Valley National Wildlife Refuge Complex.

CULTURAL RESOURCES

The SLVCA is considered an important area for cultural resources because of the abundance of cultural sites that date to almost 12,000 years ago that are located throughout the valley. However, much of the archaeological research associated with the San Luis Valley has been conducted on public lands, such as the Closed Basin, San Juan National Forest, and Great Sand Dunes National Park and Preserve (Jones 2000). Permanent protection of wildlife habitat on private land will benefit the preservation of cultural sites from future disturbance on all acquired lands.

CLIMATE CHANGE

Climate change has quickly moved to the forefront of conservation challenges during the 21st century, and the Service has made it a high priority in conservation planning (USFWS 2010b). Mountain ecosystems in the western United States are expected to be especially sensitive to climate change. In fact, data show that many places in the Rocky Mountains have experienced

three times the global average temperature increase over the past century. Measurements have shown that Colorado's temperature has increased by approximately 2 °F between 1977 and 2006 (Ray et al. 2008). The western United States has seen a shift toward earlier spring snowmelt (Karl et al. 2009). The Rio Grande Basin is predicted to experience significantly warmer temperatures, decreasing snowpack, earlier runoff, increased intensity of droughts and floods, and an overall decrease in water availability (Bureau of Reclamation 2013).

Wetland and riparian habitats, such as those found in the SLVCA, that are dependent on snowmelt from surrounding high mountain ecosystems are expected to be more acutely affected than other ecosystems. The San Luis Valley is predicted to have a 10- to 20-percent reduction in runoff by midcentury compared to the 1900 to 1970 baseline (Karl et al. 2009). As with many areas across the West, it is difficult to predict what the specific effects of climate change may be in a given area, particularly because of the complex interplay between the timing of temperature change and precipitation. The Western Water Assessment predicted that Colorado's ecosystems will be affected by climate change in nine broad ways: increased frequency and severity of forest-insect interactions; increased frequency and severity of wildfires; changes in the hydrologic cycle that affect aquatic species, including reduction in overall streamflow, shift to earlier spring runoff, and warming of water temperatures; northward and upward shift in animal ranges, causing shifts in ecosystem composition; increased range and spread of wildlife pathogens; increase in tree mortality because of drought stress; increased risk of desertification in dryland ecosystems; and an overall reduction in biodiversity because of the above impacts (Averyt et al. 2011). We must be cognizant of the potential impacts that climate change may have on wetland, riparian, and upland habitat in the SLVCA.

The SLVCA intends to support and restore habitat connectivity to promote San Luis Valley and Southern Rockies ecosystems that will be robust in the face of climate change. Protection of large intact expanses of wetland habitat types where natural ecosystem processes can be sustained will help wetland-dependent species resist some of the impacts of a changing climate. Some of these may not be the same type of wetland in the future, but the use of hydrogeomorphic modeling to assess historical hydrology should allow us to predict where and what kind of wetlands will persist in a potentially warmer and more arid future. We will respond by targeting these habitats for acquisition in the SLVCA. Besides intrinsically providing habitat for wildlife, riparian areas also serve as corridors, as do the montane forests along the flanks of the surrounding mountains. Protection of such corridors will preserve a network through which wildlife can recolonize or

disperse following disturbance, making the ecosystem more resilient to short-term change and increasing its adaptive capacity to long-term change.

Effects on the Natural and Human Environment

For a thorough discussion of the effects of the easement program, see the EA (appendix A) in this volume. Effects of the land protection strategy discussed in this volume are analyzed as alternative B in the EA (appendix A).

Chapter 4—Project Implementation



© Joe Zimm

Willows and cottonwood trees line a streamside at the base of the San Juan Mountains in Colorado.

Land Protection Choices

The following summarizes the two alternatives in the EA (appendix A) that led to the Service's approval of the SLVCA land protection project.

NO ACTION

Under the no-action alternative, the areas outside of existing protected areas would largely remain in private ownership and subject to changes in land use and land cover. Some protection in addition to the SLVCA is likely because of ongoing conservation easement initiatives in the San Luis Valley by public entities such as NRCS and nongovernmental organizations such as Rio Grande Headwaters Land Trust and Colorado Open Lands.

CONSERVATION EASEMENTS AND LIMITED FEE-TITLE ACQUISITION (PREFERRED ALTERNATIVE)

It is the Service's policy to acquire the minimum interest in a property necessary to accomplish its

conservation objectives. It can be possible to achieve most of these objectives with conservation easements. The preservation of working landscapes such as farms and rangeland is more cost effective, socially acceptable, and politically popular than acquiring fee-title land, and it often promotes the preservation of unfragmented, quality habitat. Under this project, the Service and potential partners will protect through conservation easements up to 250,000 acres to address habitat priorities described in this plan.

There are instances when the management and objectives of the three refuges in the San Luis Valley refuge complex may be simplified with small-scale acquisitions, but not with conservation easements. In such circumstances (for example, boundary simplification or surface water rights acquisition for an existing refuge), the Service will consider the acquisition of up to a total of 30,000 acres of fee title under the SLVCA. This acreage cap represents an estimate of the effort required to protect most of the first-priority habitat for the eight focal species described later in the document. Although this land protection plan in no way

binds or directs other land protection organizations, if their efforts result in protection of the same habitats identified in this plan, our efforts may be correspondingly reduced.

Project Objectives and Actions

The SLVCA sits in the San Luis Valley of central southern Colorado and northern New Mexico. The project area contains land in Alamosa, Conejos, Costilla, Hinsdale, Mineral, Rio Grande, and Saguache Counties in Colorado, as well as a small portion of Rio Arriba and Taos Counties in New Mexico. The SLVCA boundary approximates the headwaters and upper watershed of the upper Rio Grande. Within the project boundary, the Service will strategically identify and acquire from willing sellers an appropriate interest in upland, wetland, and riparian habitats on privately owned lands.

The Service plans to buy or receive donated conservation easements or fee-title lands on identified areas within the project boundaries. These easements and limited fee-title acquisitions will connect and expand existing lands under public and private conservation protection. Based on the area of privately held priority habitat and the amount protected by other organizations in the San Luis Valley, the SLVCA will protect up to 250,000 acres of uplands, wetlands, and riparian areas through easements and up to 30,000 acres through fee title.

COLLABORATION WITH LAND PROTECTION ORGANIZATIONS

The San Luis Valley is fortunate to have many active land conservation organizations. Although the mission of each organization may be unique, they all share many common objectives. In general, the Service's goal is protecting land with priority habitat from development and, in some cases, preventing water transfer when that water is supporting valuable habitat. We recognize that other organizations are doing the same thing with the same result, at times on land supporting important habitat for focal species identified in this plan. Our intent is to participate in land protection in the San Luis Valley as a partner, simply adding capacity to the collective effort to protect these habitats.

We met twice with the following organizations to discuss joint collaboration on the land protection plan:

- Trust for Public Land
- The Nature Conservancy
- Natural Resources Conservation Service

- Rio Grande Water Conservation District
- Colorado Cattlemen's Agricultural Land Trust
- Rio Grande Headwaters Land Trust
- Colorado Open Lands

We are excited to work with these and other entities interested in the objectives of the land protection plan. We are committed to institutionalizing a process that provides a suite of options for landowners and that results in easements being held by the most appropriate organization to meet specific circumstances. For example, concern has been expressed that Service-held easements may potentially interfere with the ability of the Rio Grande Water Conservation District to meet mitigation requirements of the habitat conservation plan for the southwestern willow flycatcher. This is a valid concern since the only conservation easements that can be credited to mitigation goals are those funded from non-Federal sources. Such conflict would be counterproductive for endangered species conservation. It is clear that land conservation in potential southwestern willow flycatcher habitat would need to be carried out in coordination with other conservation parties.

EASEMENT TERMS AND REQUIREMENTS

The Service has successfully carried out easements in many projects, and existing language and guidelines will contribute substantially to the drafting of the SLVCA easement language. Given the Service's conservation goals in the SLVCA, the easements will be drafted with standard language to preclude subdivision and development and conversion of native vegetation to cropland, as well as to protect existing wetlands from being drained or filled.

In addition, because of the scarcity of water resources in the valley and impending changes to ground water law in the State of Colorado, there may be provisions about water use. The types of wetland and associated upland habitats in which we are interested are largely supported by current water use practices. Easements may include a stipulation that any sale of water cannot adversely affect the quality of habitats that we seek to protect in the easements, and that water rights now owned for use on a property under an easement cannot be sold or transferred for use on other properties unless such a transfer was deemed beneficial to wildlife. These will be new easement terms for the Service and require further investigation before they can be carried out as part of the SLVCA program.

The protection of riparian corridors is important in the SLVCA, particularly because much of the lower elevation habitat has, or has the potential to have, the constituent elements of critical habitat for the

southwestern willow flycatcher¹. While easement language will not prescribe specific management practices on these lands, landowners with suitable or potentially suitable riparian habitat will be encouraged to work with the Partners for Fish and Wildlife Program or the new Working Lands for Wildlife Program (NRCS 2012) to develop alternative strategies such as fencing of riparian corridors and off-river stock watering to best manage grazing of regenerating riparian vegetation.

CONTAMINANTS OR HAZARDOUS MATERIALS

Level 1 pre-acquisition site assessments will be conducted on individual tracts before the purchase of any land interests. The Service's environmental contaminants specialists from the Ecological Services offices in Colorado and New Mexico will be contacted to make sure that policies and guidelines are followed before acquisition of conservation easements or fee title.

ACQUISITION MONEY

The Service will acquire easements in the SLVCA primarily through Land and Water Conservation Fund monies. These monies are derived primarily through revenue generated from oil and gas leases on the Outer Continental Shelf, motorboat fuel taxes, and the sale of surplus Federal property. These monies are not derived from general taxes. While Land and Water Conservation Fund monies are intended for land and water conservation projects, payment is subject to annual appropriations by the U.S. Congress for specific acquisition projects. If it is reauthorized by the U.S. Congress, the Federal Land Transaction Facilitation Act could also be used to pay for specific acquisitions. This act is a law that allows the BLM to dispose of certain public lands to generate revenue for strategic conservation of habitat not now in Federal trust.

The SLVCA project area has several other government and nongovernmental organizations with overlapping conservation objectives. In the development of the SLVCA, land for acquisition has been ranked by the Service, but the LPP may also guide acquisitions for conservation by the NRCS Wetland Reserve Program, The Nature Conservancy, Colorado Open Lands, and the Rio Grande Headwaters Land Trust, among others.

Incorporating Science and Strategic Habitat Conservation in the SLVCA

The Service uses the best available science to help figure out which areas are appropriate to consider adding

to the National Wildlife Refuge System. Continuing to look at the resources and objectives and gathering more data will help us be strategic in ongoing and future conservation efforts.

STRATEGIC GROWTH OF THE NATIONAL WILDLIFE REFUGE SYSTEM

The SLVCA encompasses approximately 4.2 million acres in a region where demand for conservation easements already far exceeds available funding. Given the likelihood that there may be more land available for conservation easements than appropriated funding, it is important to ensure that the money that is available is spent in a way that maximizes returns for trust species or helps ensure the connectivity, resiliency, and long-term function of the ecosystems in the project area, or both. In recognition of the limited resources available for land protection, the National Wildlife Refuge System is narrowing our focus to acquiring interests in land that contributes to the goals of one or more of the following national conservation objectives. We describe our contribution to these objectives below.

Aid in the Recovery of Threatened and Endangered Species

The SLVCA provides habitat for six listed species (southwestern willow flycatcher, Mexican spotted owl, and Canada lynx, Gunnison sage-grouse, yellow-billed cuckoo, and New Mexico meadow jumping mouse).

A section 7 biological evaluation for federally listed species found that the actions of the SLVCA “may affect but [are] not likely to adversely affect” listed species or designated critical habitat (appendix E). Descriptions of several of these species follow.

Southwestern Willow Flycatcher. This is a genetically distinct subspecies (Paxton 2000) of willow flycatcher that inhabits the woody riparian corridors of the desert southwest. Its population has declined significantly because of habitat loss, and it is listed as endangered by the States of Colorado and New Mexico as well as under the Federal Endangered Species Act. The recovery plan for the southwestern willow flycatcher specifically lists conservation easements and land acquisition as ways to implement recovery criterion 2: “Provide protection and create/secure sufficient habitat to assure maintenance of these populations and/or habitats over time.” The recovery plan also identifies segments of the Rio Grande and Conejos River within the project area as having substantial recovery value (USFWS 2002). As of 2009, there were 15,128 acres of woody (willow- or cottonwood-dominated) riparian vegetation in the San Luis Valley (ERO 2012). Of these, approximately 11,475 acres are on unprotected private lands within the SLVCA boundary. Many of these acres are in very small, scattered parcels and

¹FR 76(157), 50542–629. Endangered and Threatened Wildlife and Plants; Designation of Revised Critical Habitat for Southwestern Willow Flycatcher. Agency: U.S. Fish and Wildlife Service. Action: Proposed Rule. August 15, 2011

may not be suitable for hosting breeding birds, but there is documentation of willow flycatchers on the Rio Grande breeding in patch sizes as small as 0.25 acre (Cooper 1997, as cited in USFWS 2002). Using the estimate of 11 acres per territory that was used in the Service's Biological Opinion for the San Luis Valley Regional Habitat Conservation Plan (ERO 2012), it is likely that the SLVCA could help conserve habitat for several dozen more pairs of flycatchers in addition to the 68 pairs known as of 2005 (T. Ireland, U.S. Fish and Wildlife Service, personal communication, March 2013). This could substantially enhance the redundancy of breeding pairs in the San Luis Valley management unit of the Rio Grande recovery unit, which requires a minimum number of 50 territories for downlisting.

Canada Lynx. The lynx is federally listed as threatened and State-listed in Colorado as endangered. Canada lynx range through the montane forests of the Rocky Mountains. They are resident in the San Juan and Sangre de Cristo Mountains, and the junction between the Sangre de Cristo Range and the Culebra Range of the Sangre de Cristo Mountains has been identified as a particularly important corridor for the species (L. Ellwood, the Service's Ecological Services Colorado Field Office, personal communication, January 2012). This land falls within the adjoining SCCA. Within the SLVCA, the vast majority of lynx habitat is already protected, largely in Rio Grande, San Isabel, and Carson National Forests. However, the Canada lynx could benefit from the SLVCA, particularly from protection of the willow riparian corridors that the lynx prefer (Mowat and Slough 2003), which also surround the habitat of the Rio Grande cutthroat trout.

Gunnison Sage-Grouse. The sage-grouse has been listed as threatened, wherever it is found, primarily due to "habitat loss, degradation, and fragmentation due to residential, exurban, and commercial development and associated infrastructure such as roads and power lines" (USFWS 2013c). These are precisely the threats that the SLVCA's conservation easement program is designed to counter. The smallest of the existing sage-grouse populations is found in the sagebrush habitat on the south side of Poncha Pass at the far northern end of the SLVCA. Although the area was originally proposed as critical habitat for this population which covers 48,292 acres, of which 15,921 acres are unprotected private lands (USFWS 2013b), it was not included in the final critical habitat designation because of concerns that this small population may need to be augmented by translocation or release of captive-reared birds to be sustainable (USFWS 2013b). Nonetheless, both occupied and adjacent suitable but unoccupied habitat in the SLVCA could be protected under this plan, in part due to the high landowner support in this portion of the project area.

Yellow-billed Cuckoo. The San Luis Valley Regional Habitat Conservation Plan and its associated woody riparian vegetation mapping also inform estimates of the SLVCA's potential conservation benefits to the federally threatened yellow-billed cuckoo. Cuckoos breed in cottonwood-dominated riparian forest and occupy territories that average 54.4 acres in size. The extent of breeding by yellow-billed cuckoos in the San Luis Valley has not been well documented, but the birds have been observed along the Conejos River and Rio Grande. Of the 10,019 acres of cottonwood-dominated riparian forest in the valley, 8,239 acres is on unprotected private lands within the SLVCA boundary. There are seven patches greater than 54.4 acres, which could host at least 10 pairs of cuckoos.

New Mexico Meadow Jumping Mouse. Listed as endangered under the Endangered Species Act, the New Mexico meadow jumping mouse is a strongly genetically divergent lineage (King et al. 2006) of the more widespread meadow jumping mouse that inhabits the southwestern United States. The mouse prefers riparian habitat and requires permanent free-flowing water (Morrison 1990). Poor grazing management is one of the mouse's primary stressors, although drought, development, and other factors are influencing its decline (Frey and Malaney 2009). The presence of the New Mexico meadow jumping mouse in the SLVCA is unknown, but it was historically found along the Rio Grande and in the Jemez and Sangre de Cristo Mountains just south of the project area (Morrison 1992). Because the SLVCA is at the northern boundary of the mouse's range, we performed a cursory analysis of its potential for range expansion in the project area using the correlative environmental niche modeling technique implemented in Maxent (Phillips et al. 2006). A thorough overview of this method is beyond the scope of this plan, but essentially the program uses known locality data (species occurrence records from museum specimens) and candidate correlative variables (in this case, 30-arc-second resolution bioclimatic data) to define the probability of the species' occurrence across the landscape. This was then graphically displayed as a map of the species' potential range. It is important to recognize that this method does not account for physical factors (such as mountains that prevent dispersal to suitable habitat) or ecological factors (like similar species already occupying the same niche) that constrain the species' actual distribution; nor does the model account for the mouse's restrictive habitat requirements. The resulting model was then used to forecast the potential distribution of the mouse under climate change, specifically in an A2 emissions scenario in the year 2080 (www.worldclim.org/download). It is apparent from figure 4 that the New Mexico meadow jumping mouse may undergo both expansion and contraction of in different portions of





Figure 4. Spatial graphic of predicted distribution of the New Mexico meadow jumping mouse in the 20th century (top) and the late 21st century (bottom) in the San Luis Valley Conservation Area, Colorado and New Mexico.

(Note the contraction of its southern range and expansion into a more northern distribution.)

its range, and most relevant to the SLVCA, it has the potential to expand into the riparian corridors of the San Juan and northern Sangre de Cristo Mountains within the SLVCA boundary. Protection of riparian habitats in this region could give the species the adaptive capacity to persist in the face of likely range contractions in more southerly portions of its range.

Rio Grande Cutthroat Trout. Recently removed as a candidate species under the Endangered Species Act, the Rio Grande cutthroat trout is primarily threatened by introduction of nonnative trout. However, these trout are also susceptible in their present range to anthropogenic habitat disturbance due to things such as “grazing, logging, mining, road construction, and water extraction” (Pritchard and Cowley 2006). Some of these disturbances, such as construction, could be reduced or prevented on streams that cross properties with SLVCA conservation easements. Of the approximately 168 miles of designated “conservation population” (managed for less than 10-percent genetic introgression from other trout) stream segments in the SLVCA, just over 50 miles are on private lands without protection. These stream segments range from less than 50 fish per mile to more than 400 fish per mile (USFWS 2008b), so minimization of disturbance in these watersheds could protect significant numbers of these fish.

Conserve Migratory Birds in Decline

The San Luis Valley hosts 32 species which were identified as declining or of conservation concern by staff of the Region 6 Migratory Birds Program (table 4). Of these species, the Partners in Flight North American Landbird Conservation Plan identifies the following species’ considerations:

- Gunnison sage-grouse is a watch list species with multiple causes for concern across its entire range.
- Swainson’s hawk, short-eared owl, olive-sided flycatcher, and willow flycatcher are watch list species that are moderately abundant or widespread with declines or high threats.
- Lewis’ woodpecker is a watch list species because of its restricted distribution or low population size.
- Sage thrasher and Williamson’s sapsucker are “Additional Stewardship Species” because a high percent of their global populations occur in a single biome (Rich et al. 2004).

For those species with a significant portion of their global population in the Intermountain West biome (which contains the project area), estimates of the proportion of their breeding and wintering populations are provided in table 4.



© Joe Zinn

Consistent monitoring is essential for assessing whether we are being successful in meeting our conservation objectives, particularly in a dynamic environment.

Table 4. Resident, migratory, and breeding birds of concern within the San Luis Valley Conservation Area, Colorado and New Mexico, with their status and objectives from a relevant continental or regional conservation plan or list.

Common name	Partners in Flight percent of global population in Colorado	North American Landbird Conservation Plan		U.S. Shorebird Conservation Plan	
		Watch list (WL) and stewardship (S) species and population objective	Percent of global breeding and winter population in Intermountain West avifaunal biome	Relative importance of Intermountain West to species ²	Conservation category ³
American avocet				B, M, W	3
American bittern ^{1,a,b}					
Baird's sandpiper				M	2
Black-bellied plover				M, W	3
Black-headed grosbeak ¹	3.2				
Black-necked stilt				B, M, W	2
Cinnamon teal ^{1,b}					
Cordilleran flycatcher ¹	14.6				
Greater yellowlegs				M, W	3
Gunnison sage-grouse ^{1,a}		WL, increase 100%	100/100		
House wren ¹	2.2				
Least sandpiper				M, W	3
Lesser yellowlegs				M, w	3
Lewis's woodpecker ^{1,a,b}	16.1	WL, maintain/increase	87/52		
Long-billed curlew ^{a,b,c}				B, M, W	5
Long-billed dowitcher				M, W	2
Marbled godwit ^{b,c}				B, M, W	4
Northern goshawk ^{1,a}	0.9				
Olive-sided flycatcher ^{1,a,b}	1.8	WL, increase 100%	21/0		
Pectoral sandpiper ^a				<i>m</i>	2
Sage thrasher ^{1,b}	5.7	S, maintain	99/31		
Sanderling				<i>m</i>	4
Semipalmated plover				M, w	3
Short-billed dowitcher ^b				<i>m</i>	4
Short-eared owl ^{1,a,b}		WL, increase 100%			
Snowy plover ^{a,b,c}				B, M, W	5
Solitary sandpiper ^b				<i>m</i>	4
Stilt sandpiper ^a				<i>m</i>	3
Swainson's hawk ^{1,a,b}	9.8	WL, maintain/increase	15/0		
Warbling vireo ¹	4.4				
Western sandpiper				M, W	4
Willet				B, M, W	3
Williamson's sapsucker ^{1,a,b}	9.3	S, maintain	94/15		
Willow flycatcher ^{1,a,b}	0.7	WL, increase 50%			
Wilson's phalarope				B, M	4

^a Colorado Species of Greatest Conservation Need, Tier 1 and 2 (CPW 2015)

^b Service's Region 6 Bird of Conservation Concern (FWS 2008c).

^c Service's Migratory Bird Program focal species (FWS 2015).

¹ Known to breed in the San Luis Valley Conservation Area.

² B=breeding, M=migration, W=wintering: common or locally abundant, region important to the species.

B=breeding, M=migration, W=wintering: high concentration, region extremely important to the species relative to the majority of other regions.

b=breeding, m=migration, w=wintering: uncommon to fairly common, region within species range but occurs in low relative abundance relative to other regions.

³ Low (1) to high (5) priority.

The U.S. Shorebird Plan identifies the Intermountain West region, of which the SLVCA is part, as important for several species of shorebird. For example, some 90 percent of the world's adult Wilson's phalaropes stage in the Intermountain West during migration, many of them in the San Luis Valley. The biggest threat to shorebirds in this region is human-driven competition for water, which is why the preservation of wetlands is a primary focus of this project. The importance of the region to these species, as well as their conservation category, is shown in table 4.

Implement the North American Waterfowl Management Plan

While there are certainly many species of waterfowl that use the San Luis Valley for breeding, wintering, or migration, none of them do so in numbers significant at a continental scale. However, there is a regionally significant breeding population of cinnamon teal in the project area, and these are identified as a priority species by the Intermountain West Joint Venture.

STRATEGIC HABITAT CONSERVATION

To ensure that our efforts to deliver conservation are effective, the SLVCA will incorporate the elements of strategic habitat conservation. Strategic habitat conservation is based on an adaptive management framework and entails starting with strategic conservation planning, followed by conservation design, conservation delivery, and monitoring and research to assess results.

Strategic Biological Planning

Biological planning requires the identification of specific biological objectives or focal species so that the relative success of a strategy can be assessed following implementation. The focal species identified to guide prioritization of the SLVCA were chosen because of the Service's obligations to them as Federal trust species (candidate, threatened, and endangered species and migratory birds). Another factor is that the land protection undertaken to benefit focal species is likely to have conservation benefits for other species of conservation concern, such as species that are federally or State-listed as threatened or endangered, the Service's Region 6 birds of conservation concern, and the Service's migratory bird focal species. For example, protection of cottonwood riparian habitat for Lewis' woodpecker, a conspicuous regional bird of conservation concern, may also protect habitat for the more elusive yellow-billed cuckoo, recently listed as threatened under the Endangered Species Act. Because of a lack of systematic nesting surveys for these species in the project area, assumptions were made based on scientific literature and expert opinion regarding which types of habitat were important for maintaining viable populations of the focal species. In

particular, given the limited amount of quality wetland and riparian habitat present compared to presettlement conditions, it was assumed that the continued presence of those riparian types was a limiting resource in the life history of species that are thought to be obligate breeders in such habitat.

These focal species were chosen with the knowledge that there are gaps in existing data and that the habitat in the project area is likely to evolve over time in the face of environmental change and changes in human water use. As new data become available or as conditions change to the point that this conservation strategy is no longer effective, biological planning will be revisited.

Conservation Design and Delivery

Preventing loss of habitats identified for the diverse suite of focal species is the goal of the prioritization scheme outlined in section 4.4. Decisions about how to rank competing parcels with limited available funds will follow the outline described in that section.

The recovery plan for southwestern willow flycatcher requires a minimum of 50 occupied breeding territories in the San Luis Valley (USFWS 2002), and specific reaches of the Rio Grande were identified as critical habitat to maintain that level (USFWS 2013a). As previously discussed, this habitat will be granted high priority for land protection. Some of these reaches have poor or nonexistent habitat but were included in the designation due to their potential and proximity to known concentrations of southwestern willow flycatchers. All easement opportunities within the priority lands for southwestern willow flycatcher should be considered in the interest of providing redundancy to currently occupied habitat, even if the priority lands are unoccupied.

In the absence of specific population goals for the remaining focal species, no population targets or breeding pair densities have been selected. Following the principle that between 25 and 75 percent of a region must be conserved to meet targets for biodiversity (Noss et al. 2012), the initial targets for easement acquisition are to protect 50 percent of existing priority habitat for these focal species. As survey data for the valley informs the role of the SLVCA in meeting specific regional or continental population objectives for other species, the acquisition of easements and limited lands in fee title can be adjusted accordingly.

Monitoring and Research

Essential to the success of strategic habitat conservation is an effective monitoring program to ensure that conservation delivery is resulting in net positive benefits for the focal species around which the project was designed. While the consensus conservation model is primarily meant to guide effective easement acquisition, the individual species maps are intended to

guide conservation delivery for those species. Monitoring of populations will help ensure the efficacy of the program. If negative population trends for those species are detected within the project area or at a regional or continental scale, then further literature review or targeted research, or both, can be applied to adjust conservation planning for the SLVCA. Some of the monitoring phase of strategic habitat conservation can be carried out using the capacity of the refuge biologist and Service's Inventory and Monitoring assistance. However, it is important to recognize that similar monitoring will be carried out by partner agencies, and communication among these agencies is crucial for effective monitoring in the face of limited personnel and financial resources. Further, Service staff should leverage biological expertise at regional academic institutions to facilitate basic and applied research while addressing research gaps as they are identified.

Specifically, monitoring and research should include the following:

- *Developing, improving, and assessing landscape models for focal species.* Emphasis will be placed on the highest priority species with the greatest degree of uncertainty regarding limiting factors and the effectiveness of management actions, including acquisition under the SLVCA program, at minimizing and reducing the limiting factors for those species. Data from existing surveys such as Breeding Bird Survey routes in the project area will be evaluated and incorporated into spatial models. When necessary, additional data will be collected to evaluate assumptions used in the modeling process and assessments will be adjusted accordingly. These methods will provide an estimate of the population response of trust species on easement lands and on noneasement properties. Similar modeling approaches may be developed or incorporated for priority nontrust species in cooperation with partners such as State wildlife agencies, nongovernmental organizations, and universities.
- *Evaluating assumptions and addressing uncertainties identified through the biological planning, conservation design, and conservation delivery elements.* When warranted, the Service will evaluate assumptions such as increased redundancy of occupied southwestern willow flycatcher habitat through protection of riparian vegetation.
- *Identifying appropriate population goals for focal species and assessing the contribution of land protection toward meeting the population goals.* This will allow the Service and conservation partners to evaluate the contribution of the program to meeting the population goals and to refine conservation delivery to ensure maximum effectiveness.

- *Determining how changing environmental conditions may influence the effectiveness of this conservation design.* The Service will look at how increased evapotranspiration, social and economically driven changes in water use, and evolution of the type and timing of precipitation and runoff influence the hydrology of the SLVCA.

Protection Priorities

The Service, in consultation with internal divisions (Migratory Birds, Fisheries, Ecological Services), nongovernmental organization partners, Colorado Parks and Wildlife, and BLM, selected eight focal species whose habitat needs have driven the prioritization within the SLVCA. Each of these focal species represents a group of species that are vulnerable to the same threat processes (Lambeck 1997):

- Canada lynx
- Rio Grande cutthroat trout
- Southwestern willow flycatcher
- Lewis' woodpecker
- Wilson's phalarope
- American bittern
- Gunnison sage-grouse
- sage thrasher

All of these are Federal trust species or have State or regional conservation status, or both, making them worthy of protection on their own. However, conserving habitat for these species will also protect habitat for other species with similar habitat requirements.

SPECIES-HABITAT MAPPING METHODOLOGY

Some of the chosen species, by virtue of their having special conservation status, had already been the subject of detailed habitat mapping in the project area. For others, simple conceptual models were developed based on literature reviews.

The willow and cottonwood riparian habitats necessary for southwestern willow flycatcher breeding in the San Luis Valley have been mapped in detail as part of the development of the San Luis Valley Habitat Conservation Plan for that species (ERO Resources, unpublished data). The data also capture the mature cottonwood habitat needed for both the Lewis' woodpecker in this portion of its range and for the breeding habitat of the yellow-billed cuckoo. The existing data were used to define core habitat in this prioritization




Figure 5. Map of southwestern willow flycatcher and Canada lynx habitat in the San Luis Valley Conservation Area, Colorado and New Mexico.



Steve Torbit / USFWS

Protection of habitat for Federal trust species would also ensure connectivity for State-managed species such as the American black bear.

scheme; as a second priority, a 656-foot (200-meter) buffer was used to minimize disturbance of the core habitat (Terry Ireland, the Service's Ecological Services, personal communication, February 2012). These priorities are illustrated in figure 5.

Canada lynx habitat in the project area has already been mapped by Colorado Parks and Wildlife and the U.S. Forest Service. A small portion of the project area in northern New Mexico had not been covered by previous mapping but is known to be actively used by lynx. Therefore, a minimum convex polygon for this region was created that captured the land cover that largely comprises the Colorado Parks and Wildlife habitat (Rocky Mountain aspen forest and woodland, Rocky Mountain lodgepole pine forest, Southern Rocky Mountain mesic montane mixed conifer forest and woodland, and Rocky Mountain subalpine dry-mesic spruce-fir forest and woodland) using 98-foot (30-meter) Landfire data (USGS 2010). Lynx habitat is identified in figure 5.

The habitat of the Endangered Species Act candidate Rio Grande cutthroat trout has been mapped throughout the species' range; in addition, information on barriers to fish passage and data on genetic integrity has been incorporated into a spatial database. Because interbreeding has been a problem for cutthroat trout species, the signatory parties to the 2009 Rio Grande Cutthroat Trout Conservation Agreement identified populations with less than 10-percent

genetic introgression and defined them as conservation populations (Rio Grande Cutthroat Trout Conservation Team 2009). These conservation populations were chosen as representing priority habitat for the species in this land protection plan (figure 6).

The range of the Gunnison sage-grouse is much more geographically limited than it once was. The Gunnison Sage-Grouse Steering Committee revised early, coarse-scale, historical range mapping for the species (Schroeder et al. 2004) and identified current and suitable but unoccupied habitat (Gunnison Sage-grouse Rangewide Steering Committee 2005). In the project area, there is a small lek at Poncha Pass and some adjacent suitable but unoccupied habitat. Current range polygons were selected to represent priority habitat for this species; the historical range is also displayed for reference (figure 7).

The San Luis Valley represents a regionally important breeding habitat for the Wilson's phalarope (Scott Miller, San Luis Valley Refuge Complex, personal communication, January 2012) as well as habitat for many other species of migratory shorebirds. Because an applicable statistical or conceptual model for migratory shorebird breeding in the Southern Rockies was unavailable, a conceptual model based on published habitat associations of Wilson's phalarope was developed. A study of waterbird nesting in the San Luis Valley found that phalaropes preferred seasonal and short-emergent wetlands, probably because these habitats have the highest invertebrate biomass of the habitats available to them (Laubhan and Gammonley 2000). Wetlands classified by the National Wetland Inventory as temporary and seasonal were given the highest priority, followed by areas of saturated soils, as these wetland classes most closely match the definitions of seasonal and short emergent. Because Wilson's phalaropes are known to be sensitive to encroachment by woody vegetation (Cunningham and Johnson 2006), wetlands in the first and second priority classes were downgraded to third priority if they occurred within 328 feet (100 meters) of woody vegetation. In Colorado, Wilson's phalaropes typically breed in intermountain valleys between 7,000 and 10,000 feet (Kingsley 1998); however, the Service's Migratory Birds staff believe that most breeding likely occurs below 8,000 feet (S. Jones, the Service's Migratory Birds, personal communication, February 24, 2012), and so more conservative criteria were used for characterizing important phalarope habitat in the SLVCA (figure 8).

The secretive American bittern is an important representative species for a suite of waterbirds in the project area. Like the Wilson's phalarope, neither San Luis Valley-specific habitat mapping nor applicable modeling from elsewhere were available. A review of American bittern biology demonstrates that the species will nest in a wide variety of wetland and




Figure 6. Map of Rio Grande cutthroat trout habitat in the San Luis Valley Conservation Area, Colorado and New Mexico.

Conservation populations are those stream reaches with Rio Grande cutthroat trout that have less than 10-percent genetic introgression from other species. Barriers indicate impediments to upstream movement, many of which prevent invasion by nonnative fish.




Figure 7. Map of present Gunnison sage-grouse habitat in and around the San Luis Valley Conservation Area, Colorado and New Mexico, with historical range indicated for reference.




Figure 8. Map of Wilson's phalarope habitat in the San Luis Valley Conservation Area, Colorado and New Mexico.

associated upland types (Dechant et al. 2004). However, research has consistently shown a preference for tall, dense cover (Duebber and Lokemoen 1977, Riffell et al. 2001), particularly bulrush- and cattail-dominated wetlands (Azure 1998; Bent 1963; Brininger 1996; Faanes 1981, as cited in Dechant et al. 2004; Weber 1978; Weber et al. 1982). Bitterns are also found occasionally in wet meadows (Faanes 1981), particularly those with some cattails (Middleton 1949). Therefore, wetlands that were classified by the National Wetlands Inventory as permanent, semipermanent, or seasonal (those with a tall emergent vegetation component) were selected as the highest priority for American bitterns. Because bitterns are area sensitive (Brown and Dinsmore 1986, Riffell et al. 2001) and prefer wetlands of greater than 7.4 acres (3 hectares) (Daub 1993, as cited in Dechant et al. 2004), that area was used as a threshold. To define first-priority wetlands, wetlands of these types covering less than 7.4 acres were assigned second priority. Temporary and saturated wetlands, which are often wet meadows, were designated as third priority. In Colorado, American bitterns are residents of marshes between 3,500 and 8,000 feet (Bailey and Niedrach 1967), so the latter elevation was used to constrain bittern habitat in the SLVCA. These priorities are illustrated in figure 9.

Sage thrasher is a migratory bird that has been declining throughout its range due to habitat loss and degradation, and is one of the Service's Region 6 birds of conservation concern as well as a migratory bird focal species. A rangewide conceptual model for the species was developed by the American Bird Conservancy based on Rocky Mountain Bird Observatory sampling data (Beason et al. 2005) and ReGap land cover data. The population estimates they assign to these land cover classes are further stratified based on the classification of vegetation quality as good, fair, or poor, which was in turn derived from shrub cover density and prevalence of invasive plants. In the absence of data on vegetation quality for the San Luis Valley, "fair" quality was selected for all land cover types. The model developers determined that Intermountain basins big sagebrush shrubland, Intermountain basins montane sagebrush steppe, and Colorado Plateau mixed low sagebrush shrubland will support, on average, 0.0528252 birds per acre; this group of vegetation types was selected as the first priority in the sage thrasher-specific map (figure 10). Intermountain basins mixed salt desert scrub, Intermountain basins greasewood flat, and Intermountain basins semidesert shrub steppe support 0.009348 birds per acre; these vegetation classes were selected as the second priority for the species. Within these two priority levels, only polygons greater than 247 acres (100 hectares) in area were included because sage thrasher are known to be somewhat area sensitive and are found most commonly in patches of that size or greater (Knick and Rotenberry 1995).

LANDSCAPE PRIORITIZATION

The species-specific maps are useful for determining where in the landscape the key habitats are for the identified focal species. However, the maps do not assist decisionmakers with determining which areas will provide the most effective conservation returns overall. In addition to the presence or absence of habitat for individual species, it is important to take into account issues such as connectivity, cost, and unequal conservation need for each species. Therefore, the simulated-annealing algorithm implemented in the software package Marxan (Ball et al. 2009) was used to identify optimal solutions for conservation prioritization within the SLVCA. Marxan permits the user to specify individual conservation targets for conservation features (in this case, area of focal species habitat) and species-specific penalties on models that do not meet conservation targets. This allows the user to individually weight features, such as upweight penalties for not including enough habitat for species of higher conservation concern, or reduce the amount of land necessary for generalist widespread species. By designating a boundary-length modifier, the user can generate a more compact reserve system. The landscape can also be classified by cost, which can be made as simple as land area or more complex and meaningful by accounting for variables like land costs.

Because of the degree of flexibility allowed by Marxan, the values for these parameters need to be optimized by successive iterations of the program. For this analysis, hexagonal planning units were selected, as these have been shown to result in less fragmented, more efficient reserve networks (Nhancale and Smith 2011). Hexagons were 49.4 acres (20 hectares) in area, which provides resolution that is sufficient for making land protection decisions while covering the SLVCA in few enough planning units to not be computationally overwhelming. Hexagons already in a permanent protected status (existing conservation easements) were locked in to the model and counted toward the targets for the individual conservation features (focal species). Marxan was run for 1,000 runs at 100 million iterations. The species-specific data were included as features in the Marxan model. A boundary length modifier of 0.0001 was used to create a slightly more compact reserve network. Increasing that value to 0.001 oversimplified the reserve network and did not meet the intent of the SLVCA. Targets for protection were set at 50 percent of the land holding a particular conservation feature. Two exceptions were Rio Grande cutthroat trout, which was set at 75 percent because the linearity of its habitat promotes connectivity between existing protected areas, and Gunnison sage-grouse, which was set at 90 percent because of




Figure 9. Map of American bittern habitat in the San Luis Valley Conservation Area, Colorado and New Mexico.




Figure 10. Map of sage thrasher habitat in the San Luis Valley Conservation Area, Colorado and New Mexico.




Figure 11. Map of spatial conservation prioritization within the San Luis Valley Conservation Area, Colorado and New Mexico.

The relative frequency at which an individual hexagon was selected is indicated by its color, with warmer colors denoting more important locations.

its relatively small size and the high landowner support and interest in that portion of the project area. The frequency with which individual hexagons were selected in the final solution for each of the 1,000 models is shown in figure 11.

EVALUATION OF EASEMENT POTENTIAL

As described in section 4.1, acquisition of conservation easements is not a new tool for achieving conservation objectives within the SLVCA; the NRCS has a small number of easements, and nongovernmental organizations hold tens of thousands of acres of easements in the project area. These organizations have overlapping, but not identical missions to the Service. The Service currently holds only three conservation easements in the project area; however, we have more than 50 years of experience acquiring conservation easements in other parts of the country.

The landscape modeling described above has generated maps of species-specific conservation priorities for each of the focal species, as well as a consensus map that shows where conservation returns for Federal funds could be maximized for the suite of species examined. Biologists and realty specialists will work cooperatively to use these tools to identify parcels whose conservation will result in the greatest benefit to trust species.

When a willing seller approaches the Service or if the Service wishes to proactively seek out sellers, the following criteria will guide our decisionmaking:

- *Overall Conservation Value.* Is the property located, in whole or in part, in an area that was selected frequently in Marxan, as indicated by figure 11?
- *Trust Species Value.* Does the parcel contain priority habitat that was identified in any of the species-specific maps in the previous section?
- *Previously Unidentified Conservation Value.* If neither of the preceding thresholds are reached, is there another compelling reason (for example securing of important water rights, promoting critical habitat connectivity, identification of new species of conservation concern, simplified management of an existing refuge unit, or donation of intact or easily restored habitat) that justifies the property's protection?
- *Interest of Collaborators.* Is there a consensus among collaborators that a potential easement is best held by a land protection organization other than the Service? (See "Collaboration with Land Protection Organizations" under "4.2 Project Objectives and Actions.")

Nothing in these guidelines is intended to limit the appropriate exercising of discretion and professional judgment by realty specialists and refuge staff.

Acquisition will comply with realty policy and potential acquisitions will be subject to scrutiny to determine that the habitat for which the property was identified as a priority is, in fact, present on the parcel. As mentioned in the third criterion, there may also be additional reasons why acquisition of interest in a parcel is justified, even if it did not rank highly in models for selected priority trust species at the time that this plan was approved.

Socioeconomic Considerations

As discussed in detail earlier, the population in the project area is relatively low. Much of the land is cropland or rangeland. Landownership patterns vary widely, from dense 5- to 10-acre parcel subdivisions to ranches of more than 90,000 acres. Some facets of the agricultural economy are likely to be challenged by new ground water augmentation laws. The potential infusion of capital from the SLVCA conservation easement program may provide landowners with resources to invest that would allow them to continue operation. That money will largely be invested within the San Luis Valley, so there will be short-term benefits to the local economy as well. Local governments are supportive of the initiative for these reasons, and because the program is easement based and therefore should not significantly affect revenues.

Because the wildlife resources for which the SLVCA was designed already occur in these agricultural lands, sustaining this cornerstone of the regional economy is important to the mission of the Service. Maintaining these working, agricultural lands will also retain the character that defines the San Luis Valley.

Public Involvement and Coordination

Public involvement is important throughout the planning process, including scoping before plan development as well as public review of the draft environmental assessment and land protection plan.

SCOPING

At the beginning of the planning process, the planning for the SLVCA was conducted in tandem with that for the San Luis Valley National Wildlife Refuge Complex CCP, at the time in the context of a broader, valley-wide conservation area. Public scoping meetings were held on March 29, 2011, in Alamosa, Colorado; March 30, 2011, in Monte Vista, Colorado;

and March 31, 2011, in Moffat, Colorado. The scoping meetings were attended by approximately 50 people, many of whom provided input for the scoping process. Additionally, 14 written comments were received from organizations and members of the public. A press event and public meeting was held at Adams State College in Alamosa, Colorado, on January 4, 2012, at which the Secretary of the Interior, Ken Salazar, organized the presentation of several complementary initiatives for the San Luis Valley and Sangre de Cristo Mountains. One of these initiatives was landscape-scale conservation, which the Director of the Service presented as being embodied by the then SLVCA. Questions were answered and comments taken at a breakout session following the main meeting. The meeting was attended by more than 300 members of the public.

Together, these meetings and subsequent feedback helped the Service to see the questions and concerns of the public, as well as to refine the project boundary.

PUBLIC REVIEW OF THE DRAFT ENVIRONMENTAL ASSESSMENT AND LAND PROTECTION PLAN

The Service released the draft EA and LPP on May 9, 2012, for a 30-day public review period. The draft documents were made available to Federal elected officials and agencies, State elected officials and agencies, 17 Native American tribes with aboriginal interests, and other members of the public who asked to be added to our mailing list.

In February and May of 2012, refuge staff met with members of the land protection community in the San Luis Valley to discuss conservation priorities in the region. At these meetings, the Service discussed the SLVCA with representatives from entities including Rio Grande Headwaters Trust, Colorado Open Lands, Orient Land Trust, The Nature Conservancy, Colorado Cattlemen's Agricultural Land Trust, the Natural Resources Conservation Service, Trust for Public Lands, and Colorado Parks and Wildlife. Positive, constructive feedback received at those meetings guided the Service in the development of the draft and final LPP and EA.

In addition, three public meetings were held in Alamosa, San Luis, and Moffat, Colorado on May 14, 15, and 16, 2012, respectively. Approximately 50 residents and representatives of elected officials attended the 3 meetings.

Additional copies of the document are available from the following offices and contacts:

U.S. Fish and Wildlife Service
Region 6, Division of Refuge Planning
P.O. Box 25486-DFC
Denver, CO 80225
303 / 236 8132

<http://www.fws.gov/mountain-prairie/refuges/slv.php>

U.S. Fish and Wildlife Service
San Luis Valley National Wildlife Refuge
Complex
8249 Emperius Road
Alamosa, CO 81101
719 / 589 4021

Distribution and Availability

Copies of the land protection plan and environmental assessment were made available to Federal and State legislative delegations, tribes, agencies, landowners, private groups, and other interested individuals.

Glossary

- acequia**—A community-owned water distribution system found in Spain or former Spanish colonies; in the United States, found primarily in northern New Mexico and southern Colorado.
- adaptive strategy**—The ability of an ecosystem to keep ecological function while adjusting to long-term changes in the environment, or shifting to a new normal (such as climate change, established invasive species).
- anthropogenic**—Caused by human activity.
- BLM**—Bureau of Land Management, an agency of the U.S. Department of the Interior.
- candidate species**—A species of plant or animal for which the U.S. Fish and Wildlife Service has sufficient information on its biological status and threats to propose it for listing as endangered or threatened under the Endangered Species Act, but for which development of a proposed listing regulation is precluded by other higher priority listing activities.
- CCP**—See comprehensive conservation plan.
- CFR**—See Code of Federal Regulations.
- Code of Federal Regulations (CFR)**—Codification of the general and permanent rules published in the Federal Register by the executive departments and agencies of the Federal Government. Each volume of the CFR is updated once each calendar year.
- comprehensive conservation plan (CCP)**—A 15-year plan providing overall management guidance to a unit or complex of the National Wildlife Refuge System.
- conservation easement**—A legally enforceable encumbrance or transfer of property rights to a government agency or land trust for the purposes of conservation. Rights transferred could include discretion to subdivide or develop land, to change current land use practices, to sever water rights, or others as proper, and are specified by contract between the landowner and the conservation entity.
- DOE**—Department of Energy.
- EA**—See environmental assessment.
- ecological resilience**—The ability of an ecosystem to rebound from short-term changes to a landscape (such as wildfires, floods, pest outbreaks).
- EIS**—Environmental impact statement.
- endangered species**—A species of plant or animal that is in danger of extinction throughout all or a substantial part of its range.
- Endangered Species Act**—A United States law passed by the U.S. Congress in 1973 with the purpose of protecting and recovering imperiled species and the ecosystems on which they depend.
- environmental assessment (EA)**—A National Environmental Policy Act (NEPA) compliance document that analyzes whether to prepare an environmental impact statement or a finding of no significant impact, facilitates compliance when no environmental impact statement (EIS) is necessary, or facilitates preparation of an EIS when one is necessary.
- focal species**—Species that represent a group of species vulnerable to similar threats.
- HUC**—Hydrologic unit code, a hierarchical system created by the U.S. Geological Survey to find locations and regions by hydrology.
- LPP**—See land protection plan.
- land protection plan (LPP)**—A document required by policy of the U.S. Fish and Wildlife Service before the establishment of new units of the National Wildlife Refuge System, or major expansions of existing units.
- landscape conservation cooperative (LCC)**—A public-private partnership intended to facilitate cross-political boundary conservation in the face of a changing environment through application of science.
- Marxan**—A software package used as a decision support tool for spatial conservation prioritization.
- NEPA**—National Environmental Policy Act.
- NPS**—National Park Service, an agency of the U.S. Department of the Interior.
- NRCS**—Natural Resources Conservation Service, an agency of the U.S. Department of Agriculture.
- Region 6**—An administrative unit of the U.S. Fish and Wildlife Service known as the Mountain–Prairie Region, which covers eight States: Colorado, Kansas, Montana, Nebraska, North Dakota, South Dakota, Utah, and Wyoming.
- SCCA**—Sangre de Cristo Conservation Area.
- Service**—U.S. Fish and Wildlife Service, an agency of the U.S. Department of the Interior.
- SLVCA**—San Luis Valley Conservation Area.
- strategic habitat conservation**—An iterative adaptive management framework designed to make sure that decisionmaking and management within the Service is science-based. Consists of four stages: biological planning, conservation design, delivery of conservation action, and monitoring and research.

threatened species—A species of plant or animal that is likely to become endangered in the future.

trust species—Species for which the Federal Government has statutory responsibility, including threatened and endangered species, migratory birds, marine mammals, and interjurisdictional fish.

U.S.C.—United States Code.

USFS—U.S. Department of Agriculture Forest Service.

USFWS—See Service.

USGS—United States Geological Survey, an agency of the U.S. Department of the Interior.

Appendix A

Environmental Assessment

SECTION 1—PURPOSE OF AND NEED FOR ACTION

This EA documents the purpose of and the issues, alternatives, and analysis for the SLVCA. The Service has already established the SCCA within the SLVCA boundary, as proposed in the draft EA. The SLVCA would be located largely in southern Colorado, but a small portion would be in northern New Mexico (figure EA-1). Section 1 provides background information and describes the conditions that led the Service to propose creation of the SLVCA for the protection of important wetland and upland habitats, primarily through conservation easements with willing landowners.

Introduction

The SLVCA is a landscape-level strategic habitat conservation initiative within the Southern Rockies Landscape Conservation Cooperative. The SLVCA would encompass the headwaters of the Rio Grande including its namesake San Luis Valley. The San Luis Valley is a large intermountain valley bounded by the San Juan and Sangre de Cristo mountain ranges, whose rain shadows create high desert conditions in the region. However, the complex hydrology of the valley, as well as the snowmelt runoff from the mountains, have created a variety of dynamic wetlands and riparian corridors on the mountain slopes and valley floor. These wetland areas support a diverse assemblage of plants and wildlife, including habitat for many trust species such as the southwestern willow flycatcher, western snowy plover, many species of migrating and nesting waterfowl, and 95 percent of the Rocky Mountain population of greater sandhill crane. The mountains themselves are also ecologically important, providing habitat for imperiled species such as Canada lynx and Mexican spotted owl, as well as serving as migration corridors for wildlife in this southernmost extension of the Rocky Mountains.

Anthropogenic practices including agriculture, changes in fire regime, and climate change have changed the historical vegetation of the San Luis Valley. Low human population density associated with the largely agricultural economy of the valley have resulted in the San Luis Valley and surrounding mountains keeping

substantial parts of their biological value, particularly for migratory birds. However, rising agricultural costs, including those resulting from the recent State of Colorado requirement to augment surface flows to offset the impacts of ground water use, have led to an unsettled agricultural economy. The risk of second-home development could substantially reduce the quality of that habitat for sagebrush-dependent species. Surface and ground water diversions have significantly changed the amounts and timing of flows in most valley streams. In addition, ground water use has exceeded recharge rates in large portions of the valley. These factors, plus the impact of chronic drought, have resulted in a net loss of wetland habitat. The potential for farmers and ranchers to sell water rights from their lands or to convert current land use practices from agricultural to residential, industrial, or municipal uses would continue to grow and threaten the biological integrity of the San Luis Valley.

Proposed Action

The Service is moving to create the approximately 4-million-acre SLVCA to conserve vital wildlife habitats and migration corridors through voluntary conservation easements. The SLVCA acquisitions would focus on the protection of riparian corridors, wetlands, sagebrush, and montane forests in the valley through the purchase of up to 250,000 acres of conservation easements. The lands protected via easement would remain in private ownership. These lands could continue to be grazed, hayed, farmed, or otherwise managed in accordance with current practices. However, subdivision and development would be restricted, subject to stipulations agreed-upon by the landowner and the Service. Furthermore, some easements may include stipulations that the exercise of water rights associated with these lands could be changed only if the proposed changes would be beneficial to wildlife.

Up to an additional 30,000 acres of fee-title acquisition from willing sellers has been proposed as part of the preliminary project proposal for this project. However, the present intent is to use fee-title acquisition only in




Figure EA-1. Map of the San Luis Valley Conservation Area, which would be part of a broader network of public and private conservation lands.

limited circumstances to simplify the management of existing units of the National Wildlife Refuge System and when conservation objectives of those existing refuges clearly cannot be met using easements alone.

Unlike some other conservation areas of the National Wildlife Refuge System, in which objectives and the setting of priorities are largely based on modeling for one species or a guild of species, the SLVCA is intended to meet all the objectives of a complex geographic, ecological, and political environment. It therefore has a diverse range of goals:

- conserve, restore, enhance, and protect wetland and riparian habitat, an important breeding and foraging resource in the high mountain desert for migratory shorebirds, waterfowl, and neotropical passerine birds
- support the recovery and protection of threatened and endangered species that occur in the SLVCA, and reduce the likelihood of future listings under the Endangered Species Act by prioritizing key habitat for listed species and species that are candidates for listing
- protect the integrity of these habitats by preventing fragmentation and conversion of native vegetation
- conserve working landscapes based on ranching and farming activities that support a viable agricultural industry
- promote ecological resiliency and adaptive capacity by connecting together the existing network of public and private conservation lands

Decisions to Be Made

Based on the analysis provided in this final EA, the Regional Director of the Service will make two decisions:

1. Figure out if the Service should establish the SLVCA, in accordance with its land protection planning policy.
2. If yes, figure out if the selected alternative would have a significant impact on the quality of the human environment. This decision is required by the NEPA. If the quality of the human environment would not be affected, a “finding of no significant impact” will be signed and will be made available to the public. If the preferred alternative would have a significant impact, an environmental impact statement will be prepared to further address those impacts.

Issues Identified and Selected for Analysis

A description of issues identified and selected for analysis is in chapter 1.

Related Actions and Activities

A description of related actions and activities is in chapter 1.

SECTION 2—ALTERNATIVES

This section describes the two alternatives identified for this project:

- no-action alternative
- proposed action, giving the Service the authority to create the SLVCA

These alternatives were developed according to NEPA §102(2)(E) requirements to “study, develop, and describe proper alternatives to recommend courses of action in any proposal that involves unresolved conflicts concerning alternatives uses of available resources.” The alternatives consider the effects of a conservation easement program with limited fee-title acquisition within the project area boundary identified in this EA.

In addition, alternatives that were dropped from detailed study are briefly discussed.

Alternative A (No Action)

Under the no-action alternative, the areas outside of existing protected areas would largely remain in private ownership and subject to changes in land use or habitat type. Some added protection is likely because of ongoing conservation easement initiatives in the San Luis Valley by public entities such as the NRCS and nongovernmental organizations such as the Colorado Cattleman’s Agricultural Land Trust and Colorado Open Lands.

Alternative B (Preferred Alternative)

Under the preferred alternative, the Service would establish the SLVCA in southern Colorado and northern New Mexico. The project boundary encompasses approximately 4 million acres. Within this boundary, the Service would strategically acquire from willing sellers perpetual conservation easements on up to 250,000 acres through purchase or donation. The Service would also consider fee-title acquisition on up to 30,000 acres.

Conservation easements are both a cost-effective and politically effective means of land protection. They stem from the “bundle of rights” concept of land ownership (Merenlender et al. 2004), wherein, like severed surface and mineral rights for a given parcel, a part of the land title is severed and transferred to a land trust or public agency for conservation purposes. They are

quite popular for a variety of reasons. Because they allow the property owner to continue using the land, subject to agreed-upon stipulations, they protect working landscapes, which is a priority of the America’s Great Outdoors initiative. Perpetual conservation easements provide a one-time source of income to the seller or a tax incentive to the donor, and can even be an estate planning tool (Engel 2007). In many cases, they can meet the conservation objectives of the Service without our incurring the costs associated with managing fee-title land; furthermore, the land remains on the county tax rolls. In the SLVCA, the Service seeks to protect up to 250,000 acres through conservation easements and 30,000 acres in fee title.

Potential easements or fee-title lands would be prioritized based on wildlife needs in the project area, which include wetland, riparian, montane forest, and upland habitats. The Service may also investigate the possibility of acquiring properties with water rights whose protection may benefit habitat elsewhere in the valley. Chapter 4 describes these priorities in detail.

Alternatives Considered but Dropped from Further Analysis

The reasons for dropping three alternatives from further analysis are described by each nonconsidered alternative below.

Voluntary Landowner Zoning or County Zoning

Under this alternative, landowners would voluntarily petition their county commissioners to create a zoning district to direct the types of development that can occur in an area. An example of citizen-initiated zoning is when landowners would petition the county government to zone an area as agricultural, precluding certain types of nonagricultural development, such as residential subdivision or construction of a solar energy facility. However, zoning decisions are easily changed and thus do not ensure perpetual habitat protection. Also, agricultural zoning would be inadequate because water has become an increasingly expensive and limiting resource and it thus would not in itself stop continued conversion from flood-irrigated vegetation to less biologically diverse cultivated crops.

This conversion has often been accompanied by the replacement of flood irrigation practices with center-pivot irrigation. Although center-pivot irrigation offers onsite water efficiency, it results in land cover that is far less suitable to wildlife than native vegetation or even flood-irrigated agriculture. Because of these reasons, this alternative was not investigated further.

MANAGEMENT BY OTHERS

Some governmental and nongovernmental organizations are active in promoting conservation within the SLVCA and the broader San Luis Valley region. Current land managers include the Colorado Parks and Wildlife, Colorado State Land Board, BLM, National Park Service, USFS, and the Service. Additional land is conserved in fee title by The Nature Conservancy, and conservation easements are held by Ducks Unlimited, Rio Grande Headwaters Trust, the NRCS, and the Colorado Parks and Wildlife, among others.

Although the mission and focus of these organizations may be different than the mission of the Refuge System, they have made tremendous strides in protecting wildlife habitat in the San Luis Valley. The Service hopes to add capacity to these existing efforts and provide the ability to focus on habitats important for Federal trust species through the SLVCA.

FEE-TITLE ACQUISITION ONLY

Much of the publicly owned land mentioned in the previous section has been managed for conservation purposes for decades; indeed, Great Sand Dunes National Park and Preserve was originally established in 1932 as a National Monument. Fee-title ownership allows the strongest protection for the habitat and allows the greatest flexibility for adaptive management in response to new data or changing conditions. However, acquisition of new public land on the scale of the SLVCA is politically untenable and, given the low appropriation of Land and Water Conservation Fund monies, it is also financially unrealistic. For these reasons as well as the expense of managing more public lands, it is the Service's policy to acquire the minimum interest necessary to reach conservation objectives.

SECTION 3—AFFECTED ENVIRONMENT

Discussions of the resources and affected environment are in chapter 2.

SECTION 4—ENVIRONMENTAL CONSEQUENCES

For alternatives A and B described earlier, the following narrative documents the analysis of environmental effects expected to occur from carrying out each of the alternatives.

Effects on the Physical Environment

The estimated effects of each alternative on mineral, soil, and water resources, and on the Service's ability to address climate change, are described below.

ALTERNATIVE A (NO ACTION)

Development and associated habitat loss could continue on lands outside of existing protected areas; in riparian areas, development may cause erosion and sedimentation that ultimately could adversely affect aquatic species like the Rio Grande cutthroat trout. Further land protection would be limited to the efforts of other agencies and organizations. The Service's role would be limited to programs such as Partners for Fish and Wildlife; no Land and Water Conservation Fund monies would be expended in the project area by the Service for further land protection outside of the immediate vicinity of existing refuge units. Important water-dependent wildlife habitat would remain vulnerable to reallocation of surface water offsite or changes to how existing water rights are exercised.

ALTERNATIVE B (PREFERRED ALTERNATIVE)

The implementation of the goals of the SLVCA would primarily support current land use practices, and is therefore unlikely to substantially affect soil resources in the valley. There may be some reduction in erosion and sedimentation because of the prevention of subdivision and development. The SLVCA would not supersede existing third party mineral rights, and the program is therefore unlikely to affect mineral resources. If the mineral estate has not been severed, the easement may include restrictions on surface occupancy, but the Service would not, and cannot, prevent a mineral owner from accessing their minerals. The Service is unlikely to pursue acquisition of interests

in lands with outstanding surface mineral leases or rights because the associated destruction of surface vegetation and need for reclamation diminishes the wildlife value of such land. In some circumstances, habitat that depends on continuation of current water use practices would be protected from degradation caused by the sale of surface water rights or substantial changes to water use.

Effects on the Biological Environment

This section describes the likely effects of the project on species and their habitats.

ALTERNATIVE A (NO ACTION)

The Service's Partners for Fish and Wildlife Program would remain active within the project area, where it works cooperatively with landowners to voluntarily improve habitat on private land. Habitats would continue to be protected because of the ongoing efforts of agency partners and nongovernmental organizations, primarily through easements paid for by private donations, the NRCS Wetland Reserve Program, and North American Wetlands Conservation Act grants. These efforts are laudable and have conserved valuable habitat, particularly wetlands. However, they tend to underrepresent nonwetland riparian forest and uplands such as sagebrush steppe, both of which are particularly important for federally listed species and candidates for listing in the project area. Further, the demand for money under both the North American Wetlands Conservation Act and the Wetland Reserve Program is much higher than for historically available money. Also, unlike a Land and Water Conservation Fund easement program, the North American Wetlands Conservation Act requires matching funds, which are currently available through funding sources such as Great Outdoors Colorado and State tax incentives. Therefore, there would likely continue to be erosion of habitat quality and a decrease in ecological resiliency because of land

cover changes and associated fragmentation, introduction of exotic species, and construction of structures that are incompatible with habitat use by some wildlife.

Outright habitat loss because of conversion of land to other uses is a chronic, long-term threat to wildlife in many parts of the San Luis Valley. In the SLVCA, this can take the form of changes in irrigation schemes and residential and industrial development. This habitat destruction, along with construction of associated infrastructure such as water diversion structures, can result in the fragmentation of habitat. The effects of fragmentation on wildlife have been intensively studied in ecology and wildlife biology (for a conceptual review, see Collinge 2009).

Both the loss and fragmentation of riparian habitat are real concerns in the SLVCA. Riparian areas are necessary for the maintenance of medium and large mammal diversity in agricultural landscapes (for example, Hilty and Merenlender 2004), and for both breeding and stopover habitat for neotropical migratory songbirds in human-altered landscapes (Pennington et al. 2008). Riparian areas provide nest habitat for the threatened southwestern willow flycatcher and the threatened yellow-billed cuckoo, and the slow but continued loss of this habitat under alternative A would have an impact not just on regional species diversity but also on the potential persistence of imperiled species.

Besides providing habitat in and of themselves, riparian areas also serve as corridors for animal movement. Facilitating animal movement across complex mosaic landscapes is essential in a time of global environmental change. One of the greatest ecological threats of climate change is that species and varieties that are adapted to specific environmental conditions may die out because they are isolated from habitats that may have those conditions in the future (Loss et al. 2011). Under alternative A, there would be continued risk of development in previously contiguous riparian corridors.

ALTERNATIVE B (PREFERRED ALTERNATIVE)

Establishment of the SLVCA, combined with the existing SCCA, would enable the Service to permanently protect up to 530,000 acres of vital wildlife habitat in addition to that already held in Alamosa, Baca, and Monte Vista Refuges. While there are conservation initiatives by other government agencies and private land trusts underway in the project area, the SLVCA specifically targets habitat that is necessary for migration or breeding of Federal trust species, namely migratory birds and a handful of federally listed and candidate nonbird species. The conservation area should complement and enhance the ecological benefits of existing public and private conservation lands and habitat improvement programs by capturing habitats

not included in these programs and by helping to link together the existing protected area.

The use of easements to protect and buffer riparian habitats under alternative B would benefit both obligate riparian species like the southwestern willow flycatcher, bats like the Yuma myotis, and species that simply use the riparian areas as corridors to move from point to point, like bobcat and black bear. Of particular interest are the willow and cottonwood riparian forests along the tributaries of the Rio Grande, which are used by dozens of species of migratory songbirds. In the rivers and tributaries themselves, the use of easements could support conditions suitable for fish such as the Rio Grande cutthroat trout, Rio Grande chub, and Rio Grande sucker by preventing development of houses and roads, which can cause siltation and changes in water chemistry and temperature. Easements would also prevent conversion of shrub steppe near riparian areas to cropland, which can lead to increases in sediment, nitrogen loads, and temperatures in associated streams.

The presence of mesic (wet) habitats in the midst of a high-mountain desert provides an irreplaceable resource to regional, and in some cases continental, populations of breeding and migrating shorebirds, wading birds, and waterfowl. Water costs in the San Luis Valley are increasing because of restrictions on the use of ground water, and water is likely to become an increasingly complex issue because of projected changes in runoff timing and uncertainty about future precipitation trends (Ray et al. 2008). Many wetlands in the SLVCA that are important to migratory birds rely on current land use practices. We hope to work with willing landowners to maintain these practices to benefit important habitat. Consequently, on some properties with important water-dependent habitat, the easements may restrict the sale of water off the property and prevent conversion of mesic, native plant communities to domestic crops.

Sagebrush shrubland and steppe are not widespread in the project area, but are found in a ring above the desert scrubland and below the pinyon-juniper woodland in the far northern, southeastern, and southwestern portions of the valley. The Poncha Pass area was historically home to a small lek of the federally threatened Gunnison sage-grouse. Sage-grouse, as well as other sagebrush obligates, are particularly sensitive to disturbance, especially the construction of vertical structures in their habitat, which could happen if homes and associated power lines were constructed. With the lack of attention being given to that habitat type by conservation partners at present, land protection under alternative B is likely to play an important role in preventing modification of this important ecosystem.

Effects on Cultural Resources

The estimated effects of each alternative on cultural resources are described below.

ALTERNATIVE A (NO ACTION)

Some cultural resources could be adversely affected by activities such as development and road construction on lands outside of existing public and private conservation lands. While the rate of development is not rapid now, the San Luis Valley is rich with millennia of human history, and much of the valley's history is poorly documented. There are legitimate concerns that important sites may be destroyed or irreparably disturbed in the absence of protection.

ALTERNATIVE B (PREFERRED ALTERNATIVE)

There is the potential for greater protection of cultural resources than under alternative A because the easement terms that prevent development of land in ways that could adversely affect wildlife could also prevent destruction of Native American, Hispano, and other historic American sites.

Effects on the Socioeconomic Environment

This section describes the estimated effects of the alternatives on land use, ecosystem services, land ownership, and the regional economy.

ALTERNATIVE A (NO ACTION)

Landownership patterns would continue to change in accordance with market forces, as will resulting modification of ecosystem services and changes in cost of public service delivery by local government. Landowner compensation through conservation easements would remain available through other Federal programs and the efforts of nongovernmental organizations.

ALTERNATIVE B (PREFERRED ALTERNATIVE)

Following are the social and economic impacts of conservation easements and fee-title acquisitions.

Conservation easements provide public benefits for local residents, communities, and governments. Easements and fee-title acquisitions also reshape future development patterns, affect property values, and inject new money into local communities. There are many dynamic variables at play when considering the social and economic effects of conservation easements, especially given that potential purchases may span decades. Because of future uncertainty surrounding such factors as the likelihood and timing of

easements; the availability of Service money to buy lands; and population growth, land values, and agricultural commodity prices, the social and economic impacts of the easements cannot be quantified in this analysis. However, these impacts can be described qualitatively. This analysis discusses the following effects of conservation easements in the SLVCA:

- conservation values in the region
- benefits to local communities
- landowner compensation
- effects to local government net revenue

Table A, located at the end of this section, provides a summary of the social and economic impacts of conservation easements and fee-title acquisitions in the SLVCA.

Conservation Value

Conservation easements can protect values associated with biodiversity and wildlife abundance, keep aesthetic beauty, and protect social and culturally significant features of landscapes and livelihoods (Millennium Ecosystem Service Assessment 2005, Ehrlich and Ehrlich 1992, Daily 1997). Ecosystem services, such as water purification, oxygen production, pollination, and waste breakdown, are also supported for local residents through land preservation (Millennium Ecosystem Service Assessment 2005). The primary public benefit of Service conservation easements is enhanced and preserved wildlife habitat. As development stressors increase over time, many key off-refuge habitat areas may become less available because of conversion to nonwildlife habitat uses. Habitat preservation has been shown to stabilize and increase wildlife populations (Reynolds et al. 2001). Conservation easements on private lands strengthen the resiliency of species habitat and provide opportunities for wildlife movement and adaptation for years to come.

Benefits to Local Communities

Although local residents may not be able to explicitly use or access land protected by conservation easements, protected lands act as a buffer that benefits residents through increased biodiversity, recreational quality, and hunting opportunities on publicly accessible wildlife refuges and on some private lands (Rissman et al. 2007). It is well documented that open space carries positive values to local residents and communities, as well as to passersby (McConnell and Walls 2005). This is evidenced by the success of open space preservation ballot initiatives at the local, county, and State levels. Banzhaf et al. (2006) point out that between 1997 and 2004, more than 75 percent of the more than 1,100 referenda on open space conservation that appeared on ballots across the United States passed, most by a wide margin.

It is also well documented that open space and protected natural areas can increase surrounding property values (see McConnell and Walls 2005 for a comprehensive review). The reciprocating value of open space on property values varies depending on landscape characteristics and location attributes (for example, distance to the conserved area) (Kroger 2008). The permanence of the open space is also an influencing factor. Typically, open space that is permanently protected (such as refuge lands and lands protected with perpetual conservation easements) generates a higher enhancement value to local properties than land that has the potential for future development (Geoghegan et al. 2003). Location and demographic factors in the region can also influence the relative level of property enhancement value. For instance, open space may generate larger amenity premiums for property in more urbanized areas and where median incomes are higher (Netusil et al. 2000), which is not to say there is not the chance for property values to increase substantially in rural areas as well (Crompton 2001, Phillips 2000, Thorsnes 2002, Vrooman 1978).

Conservation easements would also inject new money into the local economy. The sale of conservation easements provides landowners with more revenue. Some percentage of this money may be spent in the local economy, including purchasing new real estate, consumer goods, or services in the local area. Conservation easements may also help keep the character of a region by protecting a traditional and historical way of life and the associated working landscape. Land with historical commercial use, such as ranching, forestry, and farming, is often compatible with or beneficial to wildlife refuge objectives (Jordan et al. 2007, Rissman et al. 2007). Conservation easements provide financial benefits for landowners that may enable them to preserve the natural and historical value of their farm, ranch, and open space lands, and to pass this legacy on to their children and grandchildren. In addition to supporting a cultural heritage, the preservation of farming and ranching operations can result in economic benefits to the local economy. Farmers' costs for equipment, supplies, and materials may be spent in the local economy, thus stimulating local businesses and supporting local employment. Farm workers would also spend their salaries in the local economy, thus supporting further local employment. Conservation easements may also result in increased recreation-related spending by visitors.

Landowner Compensation

The Service would buy conservation easements from willing sellers at fair-market value. The fair-market value of a conservation easement is found through an appraisal process. An appraiser estimates how much the land would sell for unencumbered by the conservation easement (the “before” value) and how much the

land would sell for with the conservation easement in place (the “after” value). The value of the conservation easement is equal to the before value minus the after value, or the difference in the fair-market value of the property with and without the easement. Landowners may also choose to donate conservation easements to the Service. The donation of a conservation easement may qualify as a tax-deductible charitable donation, which may result in Federal income tax benefits. The sale of a conservation easement for less than its fair-market value (called a “bargain sale”) may also qualify for tax deductions. Landowners may be able to claim a charitable income tax donation equal to the difference between the fair-market value and the bargain sale price of their easement. Income from the sale of a conservation easement may be taxable. Please note that the Service does not give tax advice. Landowners considering entering into a conservation agreement with the Service should consult a tax advisor or attorney for advice on how a conservation easement would affect their taxes and estate. Additionally, the Service pays for appraisals and closing costs for easements negotiated under the SLVCA.

Conservation easements reduce the value of the encumbered property. A conservation easement would reduce the fair-market value of an estate because the easement permanently removes some of the estate's development potential. The reduction in value depends on the potential development value of the land and the level of restriction agreed-upon in the easement. In general, an easement on land located in an area with high-development pressure would have a greater effect on the value of the land than an easement on land located in an area with low development pressure, and an easement that was more restrictive would have a greater effect on the value of the land than an easement that is less restrictive. The Service would buy easements at their appraised fair-market value; therefore, easements on lands with high-development pressure would receive higher payments.

Effects on Local Government Net Revenue

The effects of conservation easements on the net revenue of local government are complex and speculative; many variables are at play, and realizing the effects often requires time. Local governments collect revenue through intergovernmental transfers, property taxes, sales taxes, personal income taxes, and other charges, such as permitting. These revenues are then spent to provide community services such as fire and police services, schools, infrastructure, and public spaces. Conservation easements affect the location of future development, and therefore affect both future revenues and costs for local governments. The following sections describe the possible effects to local government revenues and costs. Overall, the SLVCA conservation easement program is expected to have

negligible effects on local government net revenues (revenues minus costs).

Effects on Local Government Revenues. Property taxes constitute the largest source of local governments' own revenue (Urban Institute and Brookings Institution 2008), and are not expected to be substantially affected by conservation easements in the SLVCA. Property taxes are assessed based on the value of property. For most types of properties, county assessors use fair-market value to find property tax liabilities; however, agricultural land is often assessed differently. In many States, the assessed value of agricultural land is based on the productive value of the land rather than on the fair-market value of the property. The fair-market value of land is the amount that a property is estimated to sell for. This value includes both the productive value of the land and any speculative value associated with the possibility of developing the land. Conservation easements reduce the fair-market value of property by removing the speculative value associated with possible development; however, conservation easements generally do not affect the productive value of agricultural land.

The SLVCA would include land in two States: Colorado and New Mexico. In both States, property taxes for agricultural land are assessed based on the productive value of the land or farm income¹ (Colorado Division of Property Taxation 2006; New Mexico Taxation and Revenue Department 2011). In the SLVCA, most properties that would enter into conservation easement agreements with the Service would be classified as agricultural land; thus, there would be little effect on the current property tax base for the nine-county area. Some of the lands in the SLVCA that would enter into easements are now fallow and do not classify as agricultural lands. For these properties, assessors may assess the fair-market value of the land based only on the uses allowed by the easement. This could result in a small reduction in property tax revenue in some counties within the region. The reduction in property taxes would be dependent on the percent of easement acres that are bought on fallow land (versus agricultural land), and on the reduction in the market value of the fallow lands.

The donation or purchase of any fee-title lands would reduce the amount of property tax revenue collected by local governments because the Service is exempt from taxation on its property holdings. Under Federal fee-title ownership, counties would qualify for reimbursement of some property tax revenue foregone under the Refuge Revenue Sharing Act of 1935, which allows the Service to make annual payments to local governments in areas where fee-title purchases have removed land from the tax rolls. Under provisions of

¹Special rules and statutes apply in each State to figure out if land in agricultural production and land in conservation easements is eligible to be assessed as agricultural land

the Refuge Revenue Sharing Act, local counties receive an annual payment for lands that have been bought by full fee-title acquisition by the Service. Payments are based on the greater of 75 cents per acre or 0.75 percent of the fair-market value. The exact amount of the annual payment depends on congressional appropriations, which in recent years have tended to be substantially less than the amount required to fulfill the authorized level of payments. In fiscal year 2010, actual Refuge Revenue Sharing payments were 22 percent of authorized levels.

Local government revenue associated with personal income is expected to remain relatively constant within the nine-county area. Conservation easements and fee-title acquisitions in the SLVCA would affect the location and distribution of development, but are not expected to change the rate or density of human population growth. Redistribution of population growth could affect the distribution of personal income-related revenues across the counties, but is expected to have little effect on total revenues within the nine-county area. There would be a one-time increase in landowner income as the Service buys the easements.

Effects on Local Government Costs. Land protection through conservation easements could result in a reduction in future expenditures for local governments and municipalities. New residential developments require local governments to provide services such as fire protection, police services, and schools, and to construct new infrastructure such as roads, parks, and water and electric-delivery systems. The costs to provide government services for new residential developments often exceed new revenues derived from the developments. This is especially true for rural residences, which tend to have higher costs for county governments and school districts than urban residences. In 2001, the American Farmland Trust found that, on average, the cost to provide community services to new residential developments was \$1.15 for every \$1.00 of revenue generated by those developments (American Farmland Trust 2001; Coupal et al. 2002). A study conducted in Wyoming found that community service costs averaged \$2.01 for every \$1.00 of revenue for rural residential lands; in contrast, the average cost to provide services for lands under agricultural production averaged \$0.54 for every \$1.00 of revenue (Taylor and Coupal 2000).

Impacts on Federal Permitting and Property Rights of Nonparticipants

Neither the authorization nor the establishment of the SLVCA would affect the administration of lands by other Federal agencies; the SLVCA boundary is simply an acquisition boundary within which the Service could acquire easements or property. Landowners who choose to take part in the program would sell or donate certain property rights to the Service. There would be no impact on adjacent property owners.

Unavoidable Adverse Impacts

This section describes adverse effects which may be unavoidable when carrying out alternatives A and B.

ALTERNATIVE A (NO ACTION)

Loss of wetland, riparian, and upland vegetation and their associated habitat values would continue because of development of areas outside of those protected by partner agencies and land trusts.

ALTERNATIVE B (PREFERRED ALTERNATIVE)

No direct or indirect unavoidable adverse impacts to the environment would result from choosing alternative B. An easement program would not result in adverse impacts on the physical or biological environment. Choosing an approved boundary for the SLVCA and concurrent authorization to go forward with an easement program would not, by itself, affect land ownership or value, or other aspects of the socioeconomic environment.

Irreversible and Irrecoverable Commitment of Resources

Any commitments of resources that may be irreversible or irretrievable because of carrying out alternatives A or B are described below.

ALTERNATIVE A (NO ACTION)

There would be no commitment of resources by the Service if alternative A were selected. The Service could still exercise its authority to acquire inholdings or for minor expansions of existing refuges, but would not be obligated to do so.

ALTERNATIVE B (PREFERRED ALTERNATIVE)

The establishment of the SLVCA would not, of itself, constitute an irreversible or irretrievable commitment of resources. However, if interests in land were acquired through the use of Land and Water Conservation Fund or donations, the administration of the easement provisions or donated property would require an irreversible and irretrievable commitment of resources. The amount of annual monitoring activity and associated compliance actions will be directly related to the number and size of easements acquired over time. We anticipate these efforts will represent a minor increase in Service costs in the near future. However, these costs could grow to very significant levels if and when the conservation area grows and

the Service holds easements on acreages approaching the limits set by the Director under this plan.

Short-Term versus Long-Term Productivity

Following is a discussion of short- and long-term effects of the alternatives.

ALTERNATIVE A (NO ACTION)

Continued efforts to conserve habitats would be ongoing through the efforts of Service activities like Partners for Fish and Wildlife and the efforts of other agency and nonprofit partners. Important wetland and upland habitats would be expected to continue to be lost at current rates of conversion, which would have long-term negative implications on the maintenance of the ecological communities they support.

ALTERNATIVE B (PREFERRED ALTERNATIVE)

The Service would be authorized to buy perpetual easements only from willing sellers, providing an immediate short-term economic benefit to landowners. This may provide capital for expansion of agricultural operations, or simply allow struggling operators to stay in business. This is particularly relevant given the changes to Colorado water law, which now requires ground water users to buy increasingly expensive surface water to decrease their impact on senior surface water users. This infusion of capital at an opportune time would likely have important long-term benefits to the economy of the San Luis Valley. The conservation of habitats under this program would also have important short- and long-term ecological benefits. The program would preserve habitat now used by wildlife, including federally protected species. This would result in the preservation of the area's biodiversity, which is important for long-term ecosystem stability and function in arid environments (Maestre et al. 2012). By preventing fragmentation, particularly in wildlife corridors like riparian areas and along the Sangre de Cristo Mountains, the program would promote long-term ecological resiliency to habitat perturbations such as large wildfires and climate change.

Cumulative Impacts

As defined by NEPA regulations, a cumulative impact on the environment “results from the incremental impact of the action when added to other past, present, and reasonably foreseeable future actions regardless of what agency or person undertakes such other actions” (40 Code of Federal Regulations [CFR]

1508.7). The following describes the past, present, and reasonably foreseeable actions related to the SLVCA. A discussion follows about the cumulative impacts of these actions in combination with the actions of alternatives A and B.

PAST, PRESENT, AND REASONABLY FORESEEABLE FUTURE ACTIONS

Some private and public organizations have successfully conducted land protection programs in the San Luis Valley through negotiation of conservation easements with willing landowners. One specific example is a coalition of local governments, landowners, and nonprofit organizations that is working to conserve land as part of their mitigation strategy in the San Luis Valley Habitat Conservation Plan that was released in November 2012. The Service assumes this would likely continue in the future.

The State of Colorado is carrying out new laws about ground water augmentation, where landowners who use ground water may have to buy surface water rights to offset any adverse impacts on downstream users. In response to these upcoming regulations, the community, largely through the leadership and support of the Rio Grande Water Conservation District and county governments, is developing locally managed subdistricts of the Rio Grande Water Conservation District. The subdistricts are meant to be locally designed and managed programs that will provide well users alternatives to individual augmentation plans.

There is ongoing interest in the San Luis Valley for renewable energy development. There are small-scale commercial solar facilities now deployed in the San Luis Valley, and the Department of Energy and the BLM are studying the impacts of more facilities being developed on public land (BLM and DOE 2010). The BLM is now reviewing the potential impacts of expanded geothermal leasing on public lands in the San Luis Valley (BLM 2012).

ALTERNATIVE A (NO ACTION)

Under this alternative, there would be no cumulative impacts on the environment because the Service would not undertake any more land protection measures.

ALTERNATIVE B (PROPOSED ACTION)

The continuing land protection efforts of others, combined with the proposed action, may have nonlinear, positive effects on wildlife populations. Because this alternative focuses on federally regulated species (such as priority migratory bird species and species listed or being considered for listing under the Endangered Species Act), implementation would result in accelerated protection of habitats for those species. Service seeks to coordinate its land protection efforts by promoting active communication with our conservation partners on land protection opportunities as they arise so that the organization whose program is

most proper can seek the acquisition of a particular land interest. The public and private conservation entities in the San Luis Valley have a longstanding friendly relationship and view each other's conservation objectives as largely complementary. However, there are specific instances where potential conflict could arise without this communication, such as riparian habitat of the southwestern willow flycatcher. The Service does not intend to compromise the ability of local government to meet their mitigation targets in the San Luis Valley Habitat Conservation Plan. To this end, the Service would not undertake any acquisition of southwestern willow flycatcher habitats along the Rio Grande or its tributaries without discussing the opportunity with our conservation partners. The Service would defer to partners in all instances where they need to seek an interest in the land first.

The impacts of new Colorado water law on water availability and cost may be cumulative with the impacts of the Service's easements, which, depending on the habitat present on a specific property, may include language restricting the sale of surface water rights from lands protected under this program. Because the easements would keep current water use practices on lands where an interest is acquired, these impacts are unlikely to be significant.

The presence of a Service interest in land could preclude construction of commercial energy production or transmission infrastructure on that property if such activity is deemed to be incompatible with the purpose of the SLVCA. This would result in unknown effects because of potentially limiting where such facilities could be sited, but the impacts of such limitations on economics and the attainment State and Federal renewable energy requirements are speculative at best, and are outside of the scope of this analysis.

Any impacts of the proposed action that are cumulative with the actions of others would largely be decided by (1) the number of landowners willing to enter into easement agreements with the Service and (2) the amount of money available for acquisition of these easements.

Table A. Social and economic impacts of conservation easements and fee-title acquisitions.

<i>Issue</i>	<i>Social and economic impacts</i>	
	<i>Conservation easements</i>	<i>Fee-title acquisitions</i>
Conservation value	<ul style="list-style-type: none"> ■ Migration corridors and habitat for deer, elk, moose, and migratory birds would be preserved. 	<ul style="list-style-type: none"> ■ Same as for easements plus the conservation value of fee-title lands may be greater than easement lands because the Service has the ability to increase conservation value through projects on the land.
Affects to local communities	<ul style="list-style-type: none"> ■ The public would enjoy increased biodiversity, recreational quality, and hunting opportunities on nearby publicly accessible refuges and some private lands. ■ Neighboring property values may increase. ■ Positive economic impacts may result from new landowner money injected into the local economy. ■ Traditional and historical ranching and farming landscapes would be preserved. 	<ul style="list-style-type: none"> ■ Same as for easements except traditional and historical ranching and farming landscapes may not be preserved. ■ Positive economic impacts may also result from increased Service habitat improvement expenditures injected into the local economy. ■ Possible increase in refuge visitation and associated impacts of visitor spending in the local economy. However, neighbors and other public may be affected by increased accesses to refuge lands.
Landowner compensation	<ul style="list-style-type: none"> ■ Landowners would be compensated for the fair-market value of the easement. ■ Easements would reduce the fair-market value of the encumbered property. ■ Landowners keep most use rights, but forfeit their right to develop or subdivide the land. Other possible restrictions include development of vertical structures, diversion or sale of water rights. 	<ul style="list-style-type: none"> ■ Landowners would be compensated for the fair-market value of the land. ■ Landowners would forfeit all rights of ownership and turn the property over to the Service.
Affects on local government net revenue	<ul style="list-style-type: none"> ■ No changes to property tax revenues are expected for agricultural lands. ■ Property tax revenues from fallow lands would decrease. ■ Other government revenues, such as personal income tax, may be redistributed throughout the region. ■ Land protection through conservation easements could result in reduced future service costs for local governments and municipalities. 	<ul style="list-style-type: none"> ■ The Service does not pay property taxes on land they own; thus, county tax revenue would decline. ■ Lost property tax revenues would be partially replaced with Refuge Revenue Sharing payments.

SECTION 5—COORDINATION AND ENVIRONMENTAL REVIEW

This section describes how the Service coordinated with others and conducted environmental reviews of various aspects of the project proposal and analysis. Additional coordination and review will be needed to carry out the preferred alternative.

Agency Coordination

The Service has discussed the establishment of the SLVCA with other Federal (USFS, National Park Service, BLM, and NRCS), State of Colorado (Colorado Parks and Wildlife, Colorado Water Conservation Board), local county governments, and regional entities (Rio Grande Water Conservation District) through a series of meetings and correspondence. Tribes with an aboriginal interest in the San Luis Valley and surrounding mountains (Pueblo of Picuris, Cochiti Pueblo, Jemez Pueblo, Jicarilla Apache Nation, Navajo Nation, San Juan Pueblo, Pueblo of Acoma, Pueblo of Jemez, Pueblo of Laguna, Pueblo of Ildefonso, Pueblo of Santa Ana, Pueblo of Santa Clara, Pueblo of Taos, Pueblo of Zuni, Southern Ute Tribe, Uintah and Ouray Ute Indian Tribe, and Ute Mountain Ute Tribe) were invited to take part or formally consult in the planning process. The Service's regional archaeologist consulted with the State Historic Preservation Officer, and was intimately involved with the development of this EA. Some nongovernmental organizations that are active in and around the San Luis Valley were also consulted, including Colorado Cattleman's Agricultural Land Trust, Colorado Open Lands, Colorado Water Trust, Ducks Unlimited, The Nature Conservancy, Orient Land Trust, Rio Grande Headwaters Land Trust, and Trust for Public Land.

The Service coordinated internally in the development of this EA as well. Region 6 refuge planning staff and San Luis Valley National Wildlife Refuge Complex staff conducted the analysis and prepared this document, as well as the LPP. An intra-Service Endangered Species Act section 7 consultation was conducted and resulted in a finding of "May affect but not likely to affect" for Endangered Species Act-protected or candidate species (appendix E). Region 6 Migratory Birds staff guided the development of our focal species list, and both that office and staff from the Region 6 Fisheries office reviewed the document (See "Appendix B—List of Preparers and Reviewers").

Contaminants and Hazardous Waste

The Service is required to invest in healthy lands. At a minimum, a level 1 pre-acquisition site assessment by the Service's Ecological Services, Colorado Field Office or New Mexico Field Office, as appropriate, would be required before acquisition.

National Environmental Policy Act

The Service conducted this analysis under the authority of and in compliance with NEPA, which requires an evaluation of reasonable alternatives that will meet stated objectives, and an assessment of the possible effects on the natural and human environment.

ENVIRONMENTAL ASSESSMENT

The EA will be the basis for determining whether the implementation of the proposed action would constitute a major Federal action significantly affecting the quality of natural and human environments. It involved other government agencies and the public in naming issues and alternatives for the project.

DISTRIBUTION AND AVAILABILITY

The Service made available this final EA (with the associated LPP in the same volume) to the project mailing list, which includes Federal and State legislative delegations; tribes; Federal, State, and local agencies; nongovernmental organizations; and interested individuals. Copies can be requested from the Service's Region 6 office. The documents are also available electronically on the refuge planning Web site.

- Project Web site: <http://www.fws.gov/mountain-prairie/refuges/slv.php>

- Project email: slvrefugesplanning@fws.gov

Region 6 Division of Biological Resources
Attn: SLVCA EA
Branch of Refuge Planning
U.S. Fish and Wildlife Service
P.O. Box 25486, Denver Federal Center
Denver, CO 80225
303 / 236 8132

Appendix B

List of Preparers and Reviewers

<i>Author's name</i>	<i>Position</i>	<i>Work unit</i>
Mike Blenden	Project leader	USFWS, San Luis Valley Refuge Complex, Alamosa, CO
Mike Dixon	Land protection planner	USFWS, Region 6, Planning Division, Lakewood, CO
Mark Ely	Geographic Information Systems specialist	USFWS, Region 6, Planning Division, Lakewood, CO
Meg Estep	Chief, water resources	USFWS, Region 6, Water Resources Division, Lakewood, CO
Lynne Koontz	Economist	USGS, Fort Collins Science Center, Policy and Science Analysis Assistance, Fort Collins, CO
Erik Larsen	Student intern	USGS, Fort Collins Science Center, Policy and Science Analysis Assistance, Fort Collins, CO
Leslie Richardson	Economist	USGS, Fort Collins Science Center, Policy and Science Analysis Assistance, Fort Collins, CO
Catherine M. Cullinane Thomas	Economist	USGS, Fort Collins Science Center, Policy and Science Analysis Assistance, Fort Collins, CO
Meg Van Ness	Regional archaeologist	USFWS, Region 6, Archaeology Division, Alamosa, CO

<i>Reviewer's name</i>	<i>Position</i>	<i>Work unit</i>
David Lucas	Chief, Division of Refuge Planning	USFWS, Region 6, Planning Division, Lakewood, CO
Brooke McDonald	Writer-editor	North State Resources, Inc., Redding, CA
Kathryn McDonald	Managing writer-editor	North State Resources, Inc., Redding, CA
Sue Oliveira	Chief, Division of Realty	USFWS, Region 6, Planning Division, Lakewood, CO
Laurie Shannon	Conservation planner	USFWS, Region 6, Planning Division, Lakewood, CO
Mitch Werner	Writer-editor	USFWS, Region 6, Planning Division, Lakewood, CO

Appendix C

Environmental Compliance

This appendix contains several environmental compliance documents:

- finding of no significant impact
- environmental action statement
- environmental compliance certificate
- Service Director's approval

Finding of No Significant Impact

U.S. Department of the Interior, U.S. Fish and Wildlife Service
Region 6, Lakewood, Colorado

San Luis Valley Conservation Area

Alamosa, Conejos, Costilla, Hinsdale, Mineral, Rio Grande, and Saguache Counties, Colorado
Rio Arriba and Taos Counties, New Mexico

The U.S. Fish and Wildlife Service (Service) has completed the San Luis Valley Conservation Area Land Protection Plan and Environmental Assessment. This planning process considered the authorization of a new unit of the National Wildlife Refuge System, the San Luis Valley Conservation Area. The Service conducted a National Environmental Policy Act (NEPA) review of the proposed easement program. The resulting environmental assessment (EA) evaluates two alternatives—alternative A, a no-action alternative; and alternative B, the preferred alternative—to establish the San Luis Valley Conservation Area.

Alternative B, the preferred alternative, was selected for implementation because it best meets the Service's mission to sustain fish and wildlife populations and to conserve a network of lands that provide their habitats. The San Luis Valley Conservation Area will use conservation easements, with limited fee-title acquisition where it benefits the management of existing refuges, to conserve these habitats in a working agricultural landscape by maintaining current land management practices while preventing the conversion of native vegetation to other uses. In so doing, this project will protect habitat for Federal trust species of wildlife, the water quality of the upper Rio Grande, and the rural agricultural aesthetic that defines the region.

PUBLIC INVOLVEMENT

As part of the public scoping process associated with this action, comments were solicited from the public through news releases and public meetings. On March 15, 2011, a news release was issued by the USFWS Region 6, which announced the beginning of the NEPA review and solicited scoping comments from the public. Public scoping meetings were held March 29, 30, and 31, 2012 in Alamosa, Monte Vista, and Moffat, Colorado, respectively. Everyone in attendance was given an opportunity to express their ideas and concerns. Approximately 50 people attended these meetings, and an additional 14 written comments were received and used to define the scope of the NEPA review and identify potential issues. On January

4, 2012, in Alamosa, Colorado, the Secretary of the Interior held a public and media event to highlight San Luis Valley projects under the America's Great Outdoors initiative, of which the San Luis Valley Conservation Area was a component. This was followed by a news release on May 9, 2012, which announced the release of a draft EA and land protection plan (LPP) for 30 days of public comment. Formal public meetings were held on May 14, 15, and 16, 2012, in Alamosa, San Luis, and Moffat, Colorado, respectively. The public comment period ended on June 8, 2012. In addition to comments presented by some of the approximately 50 people who attended these meetings, another 14 written comments were received from individuals, nongovernmental organizations, and agencies. Public comments and responses are in appendix D of the LPP.

EFFECTS OF THE PROPOSED ACTION

The EA has taken a hard look at the environmental impacts to inform the public and ourselves about the consequences of the proposed action (the Service's preferred alternative).

In determining whether this project is a major action significantly affecting the quality of the human environment, we looked at both the context and intensity of the action (40 CFR § 1508.27, 40 CFR § 1508.14) as required by NEPA. In terms of context, the preferred alternative will occur in the San Luis Valley in south-central Colorado and north-central New Mexico, but we have evaluated whether it will have effects on the human environment on a broader scale. The project will be implemented over time, dependent on the Service's ability to obtain funding needed for easement acquisition. Of the roughly 4.2 million acres within the overall project boundary, easements may be purchased by the Service only from willing sellers on a strictly voluntary basis on up to 250,000 acres, plus up to 30,000 acres of fee title purchased from willing sellers. Because the human environment is interpreted by NEPA to mean the natural and physical environment and the relationship of people with that environment (40 CFR § 1508.14), in addition to our thorough analysis of physical environmental

effects, we carefully assessed the manner in which the local people and natural resources relate to the environment in the San Luis Valley, though economic or social effects are not intended by themselves to require the preparation of an environmental impact statement (40 CFR § 1508.14).

Establishment of the San Luis Valley Conservation Area will enable the Service to seek permanent protection for important wildlife habitat for federal trust species including the federally endangered southwestern willow flycatcher, the federally threatened Gunnison sage-grouse, and migratory birds such as Wilson’s phalarope and American bittern, as well as more than 300 other species of birds. This protection will also positively benefit state-managed species such as elk.

Conservation easements in the San Luis Valley Conservation Area will increase the capacity for ecosystems in that region to adapt to climate change and increase their resiliency to temporary environmental disturbances such as drought and fire. These benefits will result from ensuring connectivity between permanently protected areas within and around the conservation area, which will allow migration and colonization by variants better adapted to changing conditions. Easements will also prevent the negative local effects of habitat fragmentation resulting from land cover changes due to subdivision and infrastructure development.

The easements will be a source of capital for local landowners. It is likely that much of this money will be reinvested locally, so easements in the San Luis Valley Conservation Area may be a one-time positive benefit to the local economy. By placing restrictions on where willing landowners could build structures, the San Luis Valley Conservation Area will affect the location and distribution, but not the rate or density, of human population growth in the project area. There may be benefits to nonparticipating landowners due to the preservation of habitat that may provide for wildlife-dependent recreation off easement lands and due to the preservation of the open-space aesthetic on participating properties. The purchase of an easement will reduce the sale value of a property, which could result in a minor reduction in tax revenues if the land was being taxed on its market, rather than agricultural value. However, such reduction in revenue is likely to be more than offset by a reduction in local government costs associated with providing services to lands converted to residential development.

The establishment of the San Luis Valley Conservation Area will not impact how other State and Federal agencies manage their lands or how they allot permits for things such as grazing on public lands. The purchase of an easement will not affect the rights of third parties to exercise their preexisting legal rights on that property (for example, third-party mineral owners).

Unless explicitly stated in the easement due to the requirement of a participating land owner, the San

Luis Valley Conservation Area will not necessarily preclude the development of certain energy infrastructure. The proposed development will be subject to a compatibility determination by the refuge manager. If the proposed development was found compatible, the Service would work with the landowner and developer to minimize the negative environmental effects of the development.

Conservation easements purchased on private land will not change the landowners’ rights to manage public access to their properties. Private landowners will retain full control over their property access rights, including allowing or restricting recreational access.

DECISION AND FINDING OF NO SIGNIFICANT IMPACT

The analysis indicates that there will not be a significant impact¹, individually or cumulatively, on the quality of the human environment² as a result of this proposed action. I agree with this conclusion and therefore find that an EIS need not be prepared. This determination is based on the following factors:

1. Environmental consequences will be beneficial to wildlife habitat, migratory bird populations, and water quality. Based on informal intra-Service section 7 consultation, the proposed action will not result in the jeopardy of any federally threatened or endangered species, or adversely modify existing designated critical habitat. The proposed conservation area may permanently protect tens of thousands of acres of habitat for Endangered Species Act listed and candidate species.
2. The proposed action will pose no known risk to public health and safety.
3. The effects on the quality of the human environment are not highly controversial. There is the potential for some restriction in the potential siting

¹ 40 CFR § 1508.27 “Significantly” as used in NEPA requires considerations of both context and intensity: (a) Context. This means that the significance of an action must be analyzed in several contexts such as society as a whole (human, national), the affected region, the affected interests, and the locality. Significance varies with the setting of the proposed action. For instance, in the case of a site-specific action, significance would usually depend upon the effects in the locale rather than in the world as a whole. Both short- and long-term effects are relevant; and (b) Intensity. This refers to the severity of impact. Responsible officials must bear in mind that more than one agency may make decisions about partial aspects of a major action.

² 40 CFR § 1508.14 “Human environment” shall be interpreted comprehensively to include the natural and physical environment and the relationship of people with that environment. (See the definition of “effects” (40 CFR § 1508.8).) This means that economic or social effects are not intended by themselves to require preparation of an environmental impact statement. When an environmental impact statement is prepared and economic or social and natural or physical environmental effects are interrelated, then the environmental impact statement will discuss all of these effects on the human environment.

for energy infrastructure, but there is unlikely to be substantial conflict over this land use issue because the San Luis Valley Conservation Area easement program is a voluntary initiative with willing sellers only.

4. The proposed easements will not affect districts, sites, highways, structures, or objects listed in or eligible for listing in the National Register of Historic Places, nor will they likely cause any loss or destruction of significant scientific, cultural, or historic resources.
5. No significant cumulative effects were identified through this assessment. The EA discussed the cumulative effects on and off the refuge with those actions proposed by others.
6. The proposed action will be in compliance with all federal, state, and local laws.

Therefore, in light of the compelling science in support of the project, and my review of the information contained in the supporting reference, I have determined that authorizing the San Luis Valley Conservation Area is not a major Federal action that would significantly affect the quality of the human environment within the meaning of Section 102(2)(C) of NEPA.

The Finding of No Significant Impact and supporting NEPA analysis will be available to the public on request. Copies of the EA are available for all

affected landowners, agencies, private groups, and other interested parties. These documents are on file at the U.S. Fish and Wildlife Service, National Wildlife Refuge System, Division of Refuge Planning, P.O. Box 25486-DFC, Denver, Colorado 80225 (telephone: 303 / 236 8145).

SUPPORTING REFERENCE

U.S. Fish and Wildlife Service. 2015. Land protection plan for the San Luis Valley Conservation Area. Lakewood, CO: U.S. Department of the Interior, U.S. Fish and Wildlife Service. 151 p.



Matt Hogan
Acting Regional Director
U.S. Fish and Wildlife Service, Region 6
Lakewood, Colorado

3-19-15
Date



Environmental Action Statement

U.S. Department of the Interior, U.S. Fish and Wildlife Service
Region 6, Lakewood, Colorado

Within the spirit and intent of the Council on Environmental Quality's regulations for implementing the National Environmental Policy Act and other statutes, orders, and policies that protect fish and wildlife resources, I have established the following administrative record and have determined that the action to establish the San Luis Valley Conservation Area and associated easement and land acquisition program:

- is a categorical exclusion as provided by 516 DM 2, appendices 1 and 2, and 516 DM 6, appendix 1. No further documentation will be made.
- is found not to have significant environmental effects as determined by the attached Finding of No Significant Impact and environmental assessment.
- is found to have special environmental conditions as described in the attached environmental assessment. The attached Finding of No Significant Impact will not be final nor any actions taken pending a 30-day period for public review [40 CFR 1501.4(e)(2)].
- is found to have significant effects and, therefore, a notice of intent will be published in the Federal Register to prepare an environmental impact statement before the project is considered further.
- is denied because of environmental damage, Service policy, or mandate.
- is an emergency situation. Only those actions necessary to control the immediate impacts of the emergency will be taken. Other related actions remain subject to National Environmental Policy Act review.

Other supporting document: U.S. Fish and Wildlife Service. 2015. Land protection plan for the San Luis Valley Conservation Area. Lakewood, CO: U.S. Department of the Interior, U.S. Fish and Wildlife Service. 151 p.

 <hr style="border: 0; border-top: 1px solid black;"/>	3/12/15 <hr style="border: 0; border-top: 1px solid black;"/>	 <hr style="border: 0; border-top: 1px solid black;"/>	3-19-15 <hr style="border: 0; border-top: 1px solid black;"/>
Will Meeks Assistant Regional Director National Wildlife Refuge System, Region 6 Lakewood, Colorado	Date	Matt Hogan Acting Regional Director U.S. Fish and Wildlife Service, Region 6 Lakewood, Colorado	Date

Environmental Compliance Certificate

U.S. Department of the Interior, U.S. Fish and Wildlife Service
Region 6, Lakewood, Colorado

Project: San Luis Valley Conservation Area

State: Colorado and New Mexico

Action (indicate if not applicable)	Date
National Environmental Policy Act (indicate one)	
<i>Categorical Exclusion</i>	N/A
<i>Environmental Assessment and Finding of No Significant Impact</i>	3/19/2015
<i>Environmental Impact Statement and Record of Decision</i>	N/A
Executive Order 11593—Protection of Historical, Archaeological, and Scientific Properties	8/1/2012
Executive Order 11988—Floodplain Management	8/1/2012
Executive Order 11990—Protection of Wetlands	8/1/2012
Executive Order 12372—Intergovernmental Review of Federal Programs	8/1/2012
Executive Order 12898—Federal Actions to Address Environmental Justice in Minority and Low-Income Populations	8/1/2012
Executive Order 12996—Management and General Public Use of the National Wildlife Refuge System	8/1/2012
Endangered Species Act, Section 7	6/28/2012
Coastal Zone Management Act, Section 307	N/A
Uniform Relocation Assistance and Real Property Acquisition Policies Act	Various
Level 1 Contaminants and Hazardous Waste (Secretarial Order 3127: 602 DM 2)	Various

I hereby certify that all requirements of the law, rules, and Service regulations or policies applicable to planning for the above project have met with compliance. I approve the establishment of the San Luis Valley Conservation Area to be administered and managed as part of the National Wildlife Refuge System.



 Matt Hogan
 Acting Regional Director
 U.S. Fish and Wildlife Service, Region 6
 Lakewood, Colorado

3-19-15
 Date

Statement of Compliance

The following Executive orders and legislative acts have been reviewed as they apply to the establishment of the San Luis Valley Conservation Area:

1. *Executive Order 11593—Protection of Historical, Archaeological, and Scientific Properties.* Per the regional archaeologist, the creation of this document constitutes an “undertaking” as defined by the National Historic Preservation Act (36 CFR 800.16(y)). It is an undertaking that has no potential to cause effects on historic properties and therefore there are no further review obligations under the act. If, in the future, there are undertakings planned that would potentially cause adverse effects on historic properties, including ground disturbance or alterations to buildings or structures over 50 years of age, those projects should be reviewed under section 106 of the act before the start of the project.
2. *Executive Order 11988—Floodplain Management.* No structures that could be damaged by or that would significantly influence the movement of floodwater are planned for construction by the U.S. Fish and Wildlife Service on land acquired as part of this project.
3. *Executive Order 11990—Protection of Wetlands.* Conveyance of the lands and interests herein shall not exempt such lands and interests from all Federal, State, and local laws and regulations as applicable thereto by virtue of their characteristics as wetlands, subject to Executive Order 11990 (May 24, 1990).
4. *Executive Order 12372—Intergovernmental Review.* The Service has discussed or offered to discuss the proposal to establish the San Luis Valley Conservation Area with landowners; conservation organizations; State, Federal, and county agencies; tribes; and other interested groups and individuals. At the Federal level, the Service staff has coordinated with the U.S. Forest Service, the Bureau of Land Management, the National Park Service, and the Natural Resource Conservation Service, as well as the congressional delegations for the affected region. At the State level, the Service has worked with the Colorado Water Conservation Board and Colorado Parks and Wildlife. The Service has consulted representatives from local governments including Costilla County, Colorado, and Taos County, New Mexico. In addition, the Service has provided information to seventeen tribes with potential interest in this project.
5. *Executive Order 12898—Federal Actions to Address Environmental Justice in Minority and Low-Income Populations.* Establishing the San Luis Valley Conservation Area will not have a disproportionately high or adverse human health or environmental effect on minority or low-income populations. Therefore, this action complies with this Executive order.
6. *Executive Order 12996—Management and General Public Use of the National Wildlife Refuge System.* The public has been invited to participate in the planning process and has been very engaged. The Service held three public scoping meetings and three public comment meetings, and released the draft environmental assessment and land protection plan for 30 days to get input on the project. The Service received four written public comments on the draft EA. Comments and issues raised by the public have been incorporated into the EA and a copy of the final document will be sent to all interested landowners, agencies, private groups, and other parties. While the San Luis Valley Conservation Area will be, by definition, a unit of the National Wildlife Refuges System, the project is largely focused on conservation easements, and the Service will not manage or have control over public access to private lands. This right will remain with the private landowner. Management of any fee-title lands purchased or donated will be in accordance with the comprehensive conservation plan for the San Luis Valley National Wildlife Refuge Complex.
7. *Endangered Species Act, section 7.* An informal intra-Service section 7 consultation with the Ecological Services field offices in Colorado and New Mexico concluded with their concurrence that the establishment of the San Luis Valley Conservation Area may affect, but is not likely to adversely affect, species protected under the Endangered Species Act.
8. *Coastal Zone Management Act.* Due to the location of the project area, compliance with this act was determined not to be needed.
9. *Uniform Relocation Assistance and Real Property Acquisition Policies Act.* The relevant portions of the act relating to tax reimbursements, etc., will be implemented on a case-by-case basis as appropriate.
10. *Secretarial Order 3127—Contaminants and Hazardous Waste.* A level 1 pre-acquisition contaminant survey will be completed before the purchase of any easement.

I hereby certify that the Service has complied with all requirements of law, rules, or regulations applicable to pre-acquisition planning for the above project. I approve the establishment of an acquisition boundary for the San Luis Valley Conservation Area and the subsequent acquisition of up to 250,000 acres of easements and 30,000 acres of fee-title from willing sellers.


_____ 3-19-15
Date

Matt Hogan
Acting Regional Director
U.S. Fish and Wildlife Service, Region 6
Lakewood, Colorado

Service Director's Approval



United States Department of the Interior

FISH AND WILDLIFE SERVICE
Washington, D.C. 20240



DEC - 2 2015

In Reply Refer To:
FWS/ANRS/NRCP/060075

Memorandum

To: Regional Director, Region 6

From: ^{Deputy} Director *Steph Jantz*

Subject: Approval of San Luis Valley Conservation Area Land Protection Plan (LPP)

I approve the Land Protection Plan for the San Luis Valley Conservation Area (SLVCA). The SLVCA will be part of a landscape conservation strategy to protect Federal trust species, conserve other plants and wildlife, and ensure the long-term functioning and resilience of diverse ecosystems. It will encompass approximately 5,249,151 acres in southern Colorado and northern New Mexico.

The SLVCA provides habitats for such well-known surrogates for biodiversity conservation as the greater sandhill crane, American bittern, pronghorn, bighorn sheep, and Rio Grande cutthroat trout. Wetland protection in the SLVCA will help fulfill population objectives for waterfowl species listed in the North American Waterfowl Management Plan. Six species of federally listed threatened and endangered species are protected in the SLVCA: southwestern willow flycatcher, Mexican spotted owl, Canada lynx, Gunnison sage-grouse, yellow-billed cuckoo, and New Mexico meadow jumping mouse.

The SLVCA provides connectivity of habitats with three National Wildlife Refuges (NWRs) and the Sangre de Cristo Conservation Area as well as National Park Service, Bureau of Land Management, U.S. Forest Service, State, and The Nature Conservancy lands. Spatially explicit models linking habitat characteristics to population objectives for focal species will be used to identify priorities for protection. Conservation easements, primarily, and limited fee title lands will be acquired from willing sellers for SLVCA purposes.

The SLVCA will serve as an outstanding example of Strategic Habitat Conservation in the Southern Rockies. It will be managed in collaboration with public and private partners of the Southern Rockies Landscape Conservation Cooperative.

Attachment

Appendix D

Species List for the San Luis Valley Conservation Area

Sources: Colorado Natural Diversity Information Source, San Luis Valley National Wildlife Refuge Complex Species List, USGS Nonindigenous Aquatic Species Database, NRCS Plants Database.

* Nonnative (Because of the number of plant species in the project area, introduced plants are not shown).

Scientific name	Common name	Status
Birds		
<i>Recurvirostra americana</i>	American avocet	
<i>Botaurus lentiginosus</i>	American bittern	
<i>Fulica americana</i>	American coot	
<i>Corvus brachyrhynchos</i>	American crow	
<i>Cinclus mexicanus</i>	American dipper	
<i>Carduelis tristis</i>	American goldfinch	
<i>Pluvialis dominica</i>	American golden plover	
<i>Falco sparverius</i>	American kestrel	
<i>Falco peregrinus anatum</i>	American peregrine falcon	
<i>Anthus rubescens</i>	American pipit	
<i>Turdus migratorius</i>	American robin	
<i>Spizella arborea</i>	American tree sparrow	
<i>Pelecanus erythrorhynchos</i>	American white pelican	
<i>Anas americana</i>	American wigeon	
<i>Calypte anna</i>	Anna's hummingbird	
<i>Gavia arctica</i>	Arctic loon	
<i>Myiarchus cinerascens</i>	Ash-throated flycatcher	
<i>Calidris bairdii</i>	Baird's sandpiper	
<i>Haliaeetus leucocephalus</i>	Bald eagle	State special concern
<i>Columba fasciata</i>	Band-tailed pigeon	
<i>Riparia riparia</i>	Bank swallow	
<i>Tyto alba</i>	Barn owl	
<i>Hirundo rustica</i>	Barn swallow	
<i>Bucephala islandica</i>	Barrow's goldeneye	
<i>Ceryle alcyon</i>	Belted kingfisher	
<i>Thryomanes bewickii</i>	Bewick's wren	
<i>Sayornis nigricans</i>	Black phoebe	
<i>Leucosticte atrata</i>	Black rosy finch	
<i>Cypseloides niger</i>	Black swift	
<i>Chlidonias niger</i>	Black tern	
<i>Mniotilta varia</i>	Black-and-white warbler	
<i>Pluvialis squatarola</i>	Black-bellied plover	
<i>Pica pica</i>	Black-billed magpie	
<i>Poecile atricapillus</i>	Black-capped chickadee	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Archilochus alexandri</i>	Black-chinned hummingbird	
<i>Nycticorax nycticorax</i>	Black-crowned night-heron	
<i>Pheucticus melanocephalus</i>	Black-headed grosbeak	
<i>Himantopus mexicanus</i>	Black-necked stilt	
<i>Dendroica striata</i>	Blackpoll warbler	
<i>Dendroica caerulescens</i>	Black-throated blue warbler	
<i>Dendroica nigrescens</i>	Black-throated gray warbler	
<i>Amphispiza bilineata</i>	Black-throated sparrow	
<i>Guiraca caerulea</i>	Blue grosbeak	
<i>Dendragapus obscurus</i>	Blue grouse	
<i>Cyanocitta cristata</i>	Blue jay	
<i>Poliophtila caerulea</i>	Blue-gray gnatcatcher	
<i>Anas discors</i>	Blue-winged teal	
<i>Dolichonyx oryzivorus</i>	Bobolink	
<i>Bombycilla garrulus</i>	Bohemian waxwing	
<i>Larus philadelphia</i>	Bonaparte's gull	
<i>Aegolius funereus</i>	Boreal owl	
<i>Euphagus cyanocephalus</i>	Brewer's blackbird	
<i>Spizella breweri</i>	Brewer's sparrow	
<i>Selasphorus platycercus</i>	Broad-tailed hummingbird	
<i>Certhia americana</i>	Brown creeper	
<i>Toxostoma rufum</i>	Brown thrasher	
<i>Leucosticte australis</i>	Brown-capped rosy finch	
<i>Molothrus ater</i>	Brown-headed cowbird	
<i>Bucephala albeola</i>	Bufflehead	
<i>Icterus bullockii</i>	Bullock's oriole	
<i>Athene cunicularia</i>	Burrowing owl	State threatened
<i>Psaltriparus minimus</i>	Bushtit	
<i>Branta hutchinsii</i>	Cackling goose	
<i>Larus californicus</i>	California gull	
<i>Stellula calliope</i>	Calliope hummingbird	
<i>Branta canadensis</i>	Canada goose	
<i>Aythya valisineria</i>	Canvasback	
<i>Pipilo fuscus</i>	Canyon towhee	
<i>Catherpes mexicanus</i>	Canyon wren	
<i>Hydroprogne caspia</i>	Caspian tern	
<i>Carpodacus cassinii</i>	Cassin's finch	
<i>Tyrannus vociferans</i>	Cassin's kingbird	
<i>Aimophila cassinii</i>	Cassin's sparrow	
<i>Bubulcus ibis</i>	Cattle egret	
<i>Bombycilla cedrorum</i>	Cedar waxwing	
<i>Dendroica pensylvanica</i>	Chestnut-sided warbler	
<i>Corvus cryptoleucus</i>	Chihuahuan raven	
<i>Spizella passerina</i>	Chipping sparrow	
<i>Anas cyanoptera</i>	Cinnamon teal	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Aechmophorus clarkii</i>	Clark's grebe	
<i>Nucifraga columbiana</i>	Clark's nutcracker	
<i>Spizella pallida</i>	Clay-colored sparrow	
<i>Petrochelidon pyrrhonota</i>	Cliff swallow	
<i>Bucephala clangula</i>	Common goldeneye	
<i>Quiscalus quiscula</i>	Common grackle	
<i>Gavia immer</i>	Common loon	
<i>Mergus merganser</i>	Common merganser	
<i>Chordeiles minor</i>	Common nighthawk	
<i>Phalaenoptilus nuttallii</i>	Common poorwill	
<i>Corvus corax</i>	Common raven	
<i>Carduelis flammea</i>	Common redpoll	
<i>Gallinago gallinago</i>	Common snipe	
<i>Sterna hirundo</i>	Common tern	
<i>Geothlypis trichas</i>	Common yellowthroat	
<i>Accipiter cooperii</i>	Cooper's hawk	
<i>Empidonax occidentalis</i>	Cordilleran flycatcher	
<i>Junco hyemalis</i>	Dark-eyed junco	
<i>Spiza americana</i>	Dickcissel	
<i>Phalacrocorax auritus</i>	Double-crested cormorant	
<i>Picoides pubescens</i>	Downy woodpecker	
<i>Empidonax oberholseri</i>	Dusky flycatcher	
<i>Dendragapus obscurus</i>	Dusky grouse	
<i>Podiceps nigricollis</i>	Eared grebe	
<i>Tyrannus tyrannus</i>	Eastern kingbird	
<i>Streptopelia decaocto</i>	Eurasian collared-dove*	
<i>Sturnus vulgaris</i>	European starling*	
<i>Coccythraustes vespertinus</i>	Evening grosbeak	
<i>Buteo regalis</i>	Ferruginous hawk	State special concern
<i>Otus flammeolus</i>	Flammulated owl	
<i>Sterna forsteri</i>	Forster's tern	
<i>Passerella iliaca</i>	Fox sparrow	
<i>Larus pipixcan</i>	Franklin's gull	
<i>Anas strepera</i>	Gadwall	
<i>Callipepla gambelii</i>	Gambel's quail	
<i>Larus glaucescens</i>	Glaucous-winged gull	
<i>Aquila chrysaetos</i>	Golden eagle	
<i>Regulus satrapa</i>	Golden-crowned kinglet	
<i>Dendroica graciae</i>	Grace's warbler	
<i>Ammodramus savannarum</i>	Grasshopper sparrow	
<i>Dumetella carolinensis</i>	Gray catbird	
<i>Empidonax wrightii</i>	Gray flycatcher	
<i>Perisoreus canadensis</i>	Gray jay	
<i>Leucosticte tephrocotis</i>	Gray-crowned rosy finch	
<i>Ardea herodias</i>	Great blue heron	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Ardea alba</i>	Great egret	
<i>Bubo virginianus</i>	Great horned owl	
<i>Geococcyx californianus</i>	Greater roadrunner	
<i>Grus canadensis tabida</i>	Greater sandhill crane	State special concern
<i>Aythya marila</i>	Greater scaup	
<i>Anser albifrons</i>	Greater white-fronted goose	
<i>Tringa melanoleuca</i>	Greater yellowlegs	
<i>Quiscalus mexicanus</i>	Great-tailed grackle	
<i>Butorides virescens</i>	Green heron	
<i>Pipilo chlorurus</i>	Green-tailed towhee	
<i>Anas crecca</i>	Green-winged teal	
<i>Centrocercus minimus</i>	Gunnison sage-grouse	Federally endangered
<i>Picoides villosus</i>	Hairy woodpecker	
<i>Empidonax hammondi</i>	Hammond's flycatcher	
<i>Zonotrichia querula</i>	Harris' sparrow	
<i>Catharus guttatus</i>	Hermit thrush	
<i>Larus argentatus</i>	Herring gull	
<i>Lophodytes cucullatus</i>	Hooded merganser	
<i>Podiceps auritus</i>	Horned grebe	
<i>Eremophila alpestris</i>	Horned lark	
<i>Carpodacus mexicanus</i>	House finch	
<i>Passer domesticus</i>	House sparrow*	
<i>Troglodytes aedon</i>	House wren	
<i>Passerina cyanea</i>	Indigo bunting	
<i>Baeolophus griseus</i>	Juniper titmouse	
<i>Charadrius vociferus</i>	Killdeer	
<i>Calcarius lapponicus</i>	Lapland longspur	
<i>Calamospiza melanocorys</i>	Lark bunting	
<i>Chondestes grammacus</i>	Lark sparrow	
<i>Passerina amoena</i>	Lazuli bunting	
<i>Ixobrychus exilis</i>	Least bittern	
<i>Empidonax minimus</i>	Least flycatcher	
<i>Calidris minutilla</i>	Least sandpiper	
<i>Sternula antillarum</i>	Least tern	Federally and State endangered
<i>Carduelis psaltria</i>	Lesser goldfinch	
<i>Aythya affinis</i>	Lesser scaup	
<i>Tringa flavipes</i>	Lesser yellowlegs	
<i>Melanerpes lewis</i>	Lewis' woodpecker	
<i>Melospiza lincolni</i>	Lincoln's sparrow	
<i>Egretta caerulea</i>	Little blue heron	
<i>Lanius ludovicianus</i>	Loggerhead shrike	
<i>Numenius americanus</i>	Long-billed curlew	State special concern
<i>Limnodromus scolopaceus</i>	Long-billed dowitcher	
<i>Asio otus</i>	Long-eared owl	
<i>Oporornis tolmiei</i>	MacGillivray's warbler	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Eugenes fulgens</i>	Magnificent hummingbird	
<i>Anas platyrhynchos</i>	Mallard	
<i>Limosa fedoa</i>	Marbled godwit	
<i>Cistothorus palustris</i>	Marsh wren	
<i>Falco columbarius</i>	Merlin	
<i>Strix occidentalis lucida</i>	Mexican spotted owl	Federally and State threatened
<i>Sialia currucoides</i>	Mountain bluebird	
<i>Poecile gambeli</i>	Mountain chickadee	
<i>Charadrius montanus</i>	Mountain plover	State special concern
<i>Zenaida macroura</i>	Mourning dove	
<i>Vermivora ruficapilla</i>	Nashville warbler	
<i>Cardinalis cardinalis</i>	Northern cardinal	
<i>Colaptes auratus</i>	Northern flicker	
<i>Accipiter gentilis</i>	Northern goshawk	
<i>Circus cyaneus</i>	Northern harrier	
<i>Mimus polyglottos</i>	Northern mockingbird	
<i>Parula americana</i>	Northern parula	
<i>Anas acuta</i>	Northern pintail	
<i>Glaucidium gnoma</i>	Northern pygmy-owl	
<i>Stelgidopteryx serripennis</i>	Northern rough-winged swallow	
<i>Aegolius acadicus</i>	Northern saw-whet owl	
<i>Anas clypeata</i>	Northern shoveler	
<i>Lanius excubitor</i>	Northern shrike	
<i>Seiurus noveboracensis</i>	Northern waterthrush	
<i>Contopus cooperi</i>	Olive-sided flycatcher	
<i>Vermivora celata</i>	Orange-crowned warbler	
<i>Pandion haliaetus</i>	Osprey	
<i>Seiurus aurocapillus</i>	Ovenbird	
<i>Gavia pacifica</i>	Pacific loon	
<i>Calidris melanotos</i>	Pectoral sandpiper	
<i>Falco peregrinus</i>	Peregrine falcon	State special concern
<i>Podilymbus podiceps</i>	Pied-billed grebe	
<i>Pinicola enucleator</i>	Pine grosbeak	
<i>Carduelis pinus</i>	Pine siskin	
<i>Gymnorhinus cyanocephalus</i>	Pinyon jay	
<i>Vireo plumbeus</i>	Plumbeous vireo	
<i>Falco mexicanus</i>	Prairie falcon	
<i>Porphyrio martinica</i>	Purple gallinule	
<i>Progne subis</i>	Purple martin	
<i>Sitta pygmaea</i>	Pygmy nuthatch	
<i>Loxia curvirostra</i>	Red crossbill	
<i>Mergus serrator</i>	Red-breasted merganser	
<i>Sitta canadensis</i>	Red-breasted nuthatch	
<i>Vireo olivaceus</i>	Red-eyed vireo	
<i>Aythya americana</i>	Redhead	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Melanerpes erythrocephalus</i>	Red-headed woodpecker	
<i>Sphyrapicus nuchalis</i>	Red-naped sapsucker	
<i>Phalaropus lobatus</i>	Red-necked phalarope	
<i>Buteo jamaicensis</i>	Red-tailed hawk	
<i>Agelaius phoeniceus</i>	Red-winged blackbird	
<i>Larus delawarensis</i>	Ring-billed gull	
<i>Aythya collaris</i>	Ring-necked duck	
<i>Phasianus colchicus</i>	Ring-necked pheasant*	
<i>Columba livia</i>	Rock pigeon*	
<i>Salpinctes obsoletus</i>	Rock wren	
<i>Pheucticus ludovicianus</i>	Rose-breasted grosbeak	
<i>Chen rossii</i>	Ross' goose	
<i>Buteo lagopus</i>	Rough-legged hawk	
<i>Regulus calendula</i>	Ruby-crowned kinglet	
<i>Oxyura jamaicensis</i>	Ruddy duck	
<i>Selasphorus rufus</i>	Rufous hummingbird	
<i>Aimophila ruficeps</i>	Rufous-crowned sparrow	
<i>Xema sabini</i>	Sabine's gull	
<i>Centrocercus urophasianus</i>	Sage-grouse	
<i>Amphispiza belli</i>	Sage sparrow	
<i>Oreoscoptes montanus</i>	Sage thrasher	
<i>Calidris alba</i>	Sanderling	
<i>Grus canadensis</i>	Sandhill crane	
<i>Passerculus sandwichensis</i>	Savannah sparrow	
<i>Sayornis saya</i>	Say's phoebe	
<i>Tyrannus forficatus</i>	Scissor-tailed flycatcher	
<i>Cistothorus platensis</i>	Sedge wren	
<i>Charadrius semipalmatus</i>	Semipalmated plover	
<i>Calidris pusilla</i>	Semipalmated sandpiper	
<i>Accipiter striatus</i>	Sharp-shinned hawk	
<i>Limnodromus griseus</i>	Short-billed dowitcher	
<i>Asio flammeus</i>	Short-eared owl	
<i>Chen caerulescens</i>	Snow goose	
<i>Egretta thula</i>	Snowy egret	
<i>Charadrius alexandrinus</i>	Snowy plover	State special concern
<i>Tringa solitaria</i>	Solitary sandpiper	
<i>Melospiza melodia</i>	Song sparrow	
<i>Porzana carolina</i>	Sora	
<i>Empidonax traillii extimus</i>	Southwestern willow flycatcher	Federally and State endangered
<i>Strix occidentalis</i>	Spotted owl	
<i>Actitis macularia</i>	Spotted sandpiper	
<i>Pipilo maculatus</i>	Spotted towhee	
<i>Cyanocitta stelleri</i>	Steller's jay	
<i>Calidris himantopus</i>	Stilt sandpiper	
<i>Melanitta perspicillata</i>	Surf scoter	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Piranga rubra</i>	Summer tanager	
<i>Buteo swainsoni</i>	Swainson's hawk	
<i>Catharus ustulatus</i>	Swainson's thrush	
<i>Melospiza georgiana</i>	Swamp sparrow	
<i>Vermivora peregrina</i>	Tennessee warbler	
<i>Picoides tridactylus</i>	Three-toed woodpecker	
<i>Myadestes townsendi</i>	Townsend's solitaire	
<i>Dendroica townsendi</i>	Townsend's warbler	
<i>Tachycineta bicolor</i>	Tree swallow	
<i>Cygnus columbianus</i>	Tundra swan	
<i>Cathartes aura</i>	Turkey vulture	
<i>Ixoreus naevius</i>	Varied thrush	
<i>Catharus fuscescens</i>	Veery	
<i>Pyrocephalus rubinus</i>	Vermilion flycatcher	
<i>Poocetes gramineus</i>	Vesper sparrow	
<i>Tachycineta thalassina</i>	Violet-green swallow	
<i>Rallus limicola</i>	Virginia rail	
<i>Vermivora virginiae</i>	Virginia's warbler	
<i>Vireo gilvus</i>	Warbling vireo	
<i>Sialia mexicana</i>	Western bluebird	
<i>Athene cunicularia</i>	Western burrowing owl	
<i>Aechmophorus occidentalis</i>	Western grebe	
<i>Tyrannus verticalis</i>	Western kingbird	
<i>Sturnella neglecta</i>	Western meadowlark	
<i>Calidris mauri</i>	Western sandpiper	
<i>Otus kennicottii</i>	Western screech-owl	
<i>Aphelocoma californica</i>	Western scrub jay	
<i>Charadrius alexandrinus nivosus</i>	Western snowy plover	
<i>Piranga ludoviciana</i>	Western tanager	
<i>Contopus sordidulus</i>	Western wood-pewee	
<i>Numenius phaeopus</i>	Whimbrel	
<i>Eudocimus albus</i>	White ibis	
<i>Sitta carolinensis</i>	White-breasted nuthatch	
<i>Zonotrichia leucophrys</i>	White-crowned sparrow	
<i>Plegadis chihi</i>	White-faced ibis	
<i>Calidris fuscicollis</i>	White-rumped sandpiper	
<i>Lagopus leucurus</i>	White-tailed ptarmigan	
<i>Zonotrichia albicollis</i>	White-throated sparrow	
<i>Aeronautes saxatalis</i>	White-throated swift	
<i>Loxia leucoptera</i>	White-winged crossbill	
<i>Melanitta fusca</i>	White-winged scoter	
<i>Meleagris gallopavo</i>	Wild turkey	
<i>Grus americana</i>	Whooping crane	Federally and State endangered
<i>Catoptrophorus semipalmatus</i>	Willet	
<i>Sphyrapicus thyroideus</i>	Williamson's sapsucker	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Empidonax traillii</i>	Willow flycatcher	
<i>Phalaropus tricolor</i>	Wilson's phalarope	
<i>Gallinago delicata</i>	Wilson's snipe	
<i>Wilsonia pusilla</i>	Wilson's warbler	
<i>Aix sponsa</i>	Wood duck	
<i>Helmitheros vermivorum</i>	Worm-eating warbler	
<i>Hylocichla mustelina</i>	Wood thrush	
<i>Dendroica petechia</i>	Yellow warbler	
<i>Coccyzus americanus</i>	Yellow-billed cuckoo	Federally threatened
<i>Icteria virens</i>	Yellow-breasted chat	
<i>Xanthocephalus xanthocephalus</i>	Yellow-headed blackbird	
<i>Dendroica coronata</i>	Yellow-rumped warbler	
Amphibians		
<i>Bufo boreas boreas</i>	Boreal toad	State endangered
<i>Rana catesbeiana</i>	Bullfrog*	
<i>Hyla arenicolor</i>	Canyon treefrog	
<i>Bufo cognatus</i>	Great Plains toad	
<i>Spea multiplicata</i>	New Mexico spadefoot	
<i>Rana pipiens</i>	Northern leopard frog	State special concern
<i>Spea bombifrons</i>	Plains spadefoot	
<i>Ambystoma tigrinum</i>	Tiger salamander	
<i>Pseudacris triseriata</i>	Western chorus frog	
<i>Bufo woodhousii</i>	Woodhouse's toad	
Mammals		
<i>Sciurus aberti</i>	Abert's squirrel	
<i>Taxidea taxus</i>	American badger	
<i>Castor canadensis</i>	American beaver	
<i>Cervus elaphus</i>	American elk	
<i>Martes americana</i>	American marten	
<i>Ochotona princeps</i>	American pika	
<i>Eptesicus fuscus</i>	Big brown bat	
<i>Ovis canadensis</i>	Bighorn sheep	
<i>Ursus americanus</i>	Black bear	
<i>Mustela nigripes</i>	Black-footed ferret	Federally and State endangered
<i>Lepus californicus</i>	Black-tailed jackrabbit	
<i>Lynx rufus</i>	Bobcat	
<i>Thomomys bottae</i>	Botta's pocket gopher	State special concern
<i>Tadarida brasiliensis</i>	Brazilian free-tailed bat	
<i>Neotoma cinerea</i>	Bushy-tailed woodrat	
<i>Tamias quadrivittatus</i>	Colorado chipmunk	
<i>Conepatus mesoleucus</i>	Common hog-nosed skunk	
<i>Ondatra zibethicus</i>	Common muskrat	
<i>Erethizon dorsatum</i>	Common porcupine	
<i>Canis latrans</i>	Coyote	
<i>Peromyscus maniculatus</i>	Deer mouse	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Sylvilagus audubonii</i>	Desert cottontail	
<i>Mustela erminea</i>	Ermine	
<i>Myotis thysanodes</i>	Fringed myotis	
<i>Spermophilus lateralis</i>	Golden-mantled ground squirrel	
<i>Urocyon cinereoargenteus</i>	Gray fox	
<i>Cynomys gunnisoni</i>	Gunnison's prairie dog	
<i>Phenacomys intermedius</i>	Heather vole	
<i>Lasiurus cinereus</i>	Hoary bat	
<i>Mus musculus</i>	House mouse*	
<i>Tamias minimus</i>	Least chipmunk	
<i>Myotis lucifugus</i>	Little brown myotis	
<i>Myotis evotis</i>	Long-eared myotis	
<i>Myotis volans</i>	Long-legged myotis	
<i>Microtus longicaudus</i>	Long-tailed vole	
<i>Mustela frenata</i>	Long-tailed weasel	
<i>Lynx canadensis</i>	Canada lynx	Federally threatened, State endangered
<i>Sorex cinereus</i>	Masked shrew	
<i>Microtus pennsylvanicus</i>	Meadow vole	
<i>Neotoma mexicana</i>	Mexican woodrat	
<i>Mustela vison</i>	Mink	
<i>Sorex monticolus</i>	Montane shrew	
<i>Microtus montanus</i>	Montane vole	
<i>Alces alces</i>	Moose	
<i>Sylvilagus nuttallii</i>	Mountain cottontail	
<i>Oreamnos americanus</i>	Mountain goat	
<i>Felis concolor</i>	Mountain lion	
<i>Odocoileus hemionus</i>	Mule deer	
<i>Onychomys leucogaster</i>	Northern grasshopper mouse	
<i>Thomomys talpoides</i>	Northern pocket gopher	State special concern
<i>Lutra canadensis</i>	Northern river otter	
<i>Peromyscus nasutus</i>	Northern rock mouse	
<i>Dipodomys ordii</i>	Ord's kangaroo rat	
<i>Tamiasciurus hudsonicus</i>	Pine squirrel	
<i>Perognathus flavescens</i>	Plains pocket mouse	
<i>Antilocapra americana</i>	Pronghorn	
<i>Procyon lotor</i>	Raccoon	
<i>Vulpes vulpes</i>	Red fox	
<i>Bassariscus astutus</i>	Ringtail	
<i>Perognathus flavus</i>	Silky pocket mouse	
<i>Lasionycteris noctivagans</i>	Silver-haired bat	
<i>Lepus americanus</i>	Snowshoe hare	
<i>Clethrionomys gapperi</i>	Southern red-backed vole	
<i>Mephitis mephitis</i>	Striped skunk	
<i>Spermophilus tridecemlineatus</i>	Thirteen-lined ground squirrel	
<i>Plecotus townsendii</i>	Townsend's big-eared bat	State special concern

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Sorex palustris</i>	Water shrew	
<i>Reithrodontomys megalotis</i>	Western harvest mouse	
<i>Zapus princeps</i>	Western jumping mouse	
<i>Myotis ciliolabrum</i>	Western small-footed myotis	
<i>Spilogale gracilis</i>	Western spotted skunk	
<i>Odocoileus virginianus</i>	White-tailed deer	
<i>Lepus townsendii</i>	White-tailed jackrabbit	
<i>Gulo gulo</i>	Wolverine	State endangered
<i>Spermophilus elegans</i>	Wyoming ground squirrel	
<i>Marmota flaviventris</i>	Yellow-bellied marmot	
<i>Myotis yumanensis</i>	Yuma myotis	
Reptiles		
<i>Sceloporus undulatus</i>	Fence lizard	
<i>Pituophis catenifer</i>	Gopher snake	
<i>Eumeces multivirgatus</i>	Many-lined skink	
<i>Crotalus viridis concolor</i>	Midget faded rattlesnake	State special concern
<i>Lampropeltis triangulum</i>	Milk snake	
<i>Phrynosoma hernandesi</i>	Short-horned lizard	
<i>Liochlorophis vernalis</i>	Smooth green snake	
<i>Eumeces gaigeae</i>	Variable skink	
<i>Crotalus viridis</i>	Western rattlesnake	
<i>Thamnophis elegans</i>	Western terrestrial garter snake	
Fish		
<i>Anguilla rostrata</i>	American eel*	
<i>Thymallus arcticus</i>	Arctic grayling*	
<i>Ameiurus melas</i>	Black bullhead	
<i>Pomoxis nigromaculatus</i>	Black crappie*	
<i>Gymnocorymbus ternetzi</i>	Black tetra*	
<i>Ictalurus furcatus</i>	Blue catfish*	
<i>Oreochromis aureus</i>	Blue tilapia*	
<i>Lepomis macrochirus</i>	Bluegill*	
<i>Culaea inconstans</i>	Brook stickleback*	
<i>Salvelinus fontinalis</i>	Brook trout*	
<i>Salmo trutta</i>	Brown trout*	
<i>Ictalurus punctatus</i>	Channel catfish*	
<i>Oncorhynchus clarkii pleuriticus</i>	Colorado River cutthroat*	
<i>Cyprinus carpio</i>	Common carp*	
<i>Corydoras</i> sp.	Corydoras catfish*	
<i>Oncorhynchus clarkii × mykiss</i>	Cutbow trout (hybrid)*	
<i>Pimephales promelas</i>	Fathead minnow	
<i>Oncorhynchus clarkii carmichaeli</i>	Fine-spotted Snake River cutthroat*	
<i>Pylodictis olivaris</i>	Flathead catfish*	
<i>Platygobio gracilis</i>	Flathead chub*	
<i>Pterophyllum</i> sp.	Freshwater angelfish*	
<i>Oncorhynchus aguabonita</i>	Golden trout*	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Carassius auratus</i>	Goldfish*	
<i>Ctenopharyngodon idella</i>	Grass carp*	
<i>Xiphophorus hellerii</i>	Green swordtail*	
<i>Lepomis cyanellus</i>	Green sunfish	
<i>Poecilia reticulata</i>	Guppy*	
<i>Hemigrammus ocellifer</i>	Head-and-tailight tetra*	
<i>Oncorhynchus nerka</i>	Kokanee*	
<i>Salvelinus namaycush</i>	Lake trout*	
<i>Salmo salar sebago</i>	Landlocked Atlantic salmon*	
<i>Micropterus salmoides</i>	Largemouth bass*	
<i>Rhinichthys cataractae</i>	Long-nose dace	
<i>Catostomus catostomus</i>	Longnose sucker*	
<i>Cottus bairdii</i>	Mottled sculpin*	
<i>Oreochromis mossambicus</i>	Mozambique tilapia*	
<i>Paracheirodon innesi</i>	Neon tetra*	
<i>Esox lucius</i>	Northern pike*	
<i>Fundulus zebrinus</i>	Plains killifish*	
<i>Fundulus sciadicus</i>	Plains topminnow*	
<i>Lepomis gibbosus</i>	Pumpkinseed*	
<i>Oncorhynchus mykiss</i>	Rainbow trout*	
<i>Symphysodon discus</i>	Red discus*	
<i>Gila pandora</i>	Rio Grande chub	State special concern
<i>Oncorhynchus clarki virginalis</i>	Rio Grande cutthroat trout	State special concern
<i>Catostomus plebeius</i>	Rio Grande sucker	State endangered
<i>Poecilia latipinna</i>	Sailfin molly*	
<i>Poecilia mexicana</i>	Shortfin molly*	
<i>Micropterus dolomieu</i>	Smallmouth bass*	
<i>Xiphophorus maculatus</i>	Southern platyfish*	
<i>Hypostomus</i> sp.	Suckermouth catfish*	
<i>Otocinclus</i> sp.	Suckermouth catfish*	
<i>Tinca tinca</i>	Tench*	
<i>Dorosoma petenense</i>	Threadfin shad*	
<i>Xiphophorus variatus</i>	Variable platyfish*	
<i>Pterygoplichthys disjunctivus</i>	Vermiculated sailfin*	
<i>Sander vitreus</i>	Walleye*	
<i>Lepomis gulosus</i>	Warmouth*	
<i>Oncorhynchus clarkii lewisi</i>	West slope cutthroat*	
<i>Gambusia affinis</i>	Western mosquitofish*	
<i>Catostomus commersonii</i>	White sucker*	
<i>Ameiurus natalis</i>	Yellow bullhead*	
<i>Perca flavescens</i>	Yellow perch*	
<i>Oncorhynchus clarkii bouvieri</i>	Yellowstone cutthroat*	
Plants		
<i>Abies concolor</i>	White fir	
<i>Abies lasiocarpa</i>	Subalpine fir	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Abies lasiocarpa</i> var. <i>arizonica</i>	Corkbark fir	
<i>Abies lasiocarpa</i> var. <i>lasiocarpa</i>	Subalpine fir	
<i>Acer glabrum</i>	Rocky Mountain maple	
<i>Achillea millefolium</i>	Common yarrow	
<i>Achillea millefolium</i> var. <i>occidentalis</i>	Western yarrow	
<i>Achnatherum</i> × <i>bloomeri</i>	—	
<i>Achnatherum hymenoides</i>	Indian ricegrass	
<i>Achnatherum lettermanii</i>	Letterman's needlegrass	
<i>Achnatherum nelsonii</i>	Columbia needlegrass	
<i>Achnatherum nelsonii</i> ssp. <i>nelsonii</i>	Columbia needlegrass	
<i>Achnatherum robustum</i>	Sleepygrass	
<i>Achnatherum scribneri</i>	Scribner needlegrass	
<i>Aconitum columbianum</i>	Columbian monkshood	
<i>Aconitum columbianum</i> ssp. <i>columbianum</i>	Columbian monkshood	
<i>Acroptilon repens</i>	Hardheads	
<i>Actaea rubra</i>	Red baneberry	
<i>Actaea rubra</i> ssp. <i>arguta</i>	Red baneberry	
<i>Adoxa moschatellina</i>	Muskroot	
<i>Agastache pallidiflora</i>	Bill Williams Mountain giant hyssop	
<i>Agastache pallidiflora</i> ssp. <i>pallidiflora</i>	Bill Williams Mountain giant hyssop	
<i>Agastache pallidiflora</i> ssp. <i>pallidiflora</i> var. <i>greenei</i>	Bill Williams Mountain giant hyssop	
<i>Agoseris aurantiaca</i>	Orange agoseris	
<i>Agoseris glauca</i>	Pale agoseris	
<i>Agrostis exarata</i>	Spike bentgrass	
<i>Agrostis gigantea</i>	Redtop	
<i>Agrostis humilis</i>	Alpine bentgrass	
<i>Agrostis scabra</i>	Rough bentgrass	
<i>Agrostis variabilis</i>	Mountain bentgrass	
<i>Aletes anisatus</i>	Rocky Mountain Indian parsley	
<i>Aliciella pinatifida</i>	Sticky gilia	
<i>Alisma gramineum</i>	Narrowleaf water plantain	
<i>Alisma triviale</i>	Northern water plantain	
<i>Allium cernuum</i>	Nodding onion	
<i>Allium geyeri</i>	Geyer's onion	
<i>Allium geyeri</i> var. <i>tenerum</i>	Bulbil onion	
<i>Almutaster pauciflorus</i>	Alkali marsh aster	
<i>Alnus incana</i>	Gray alder	
<i>Alnus incana</i> ssp. <i>tenuifolia</i>	Thinleaf alder	
<i>Alopecurus aequalis</i>	Shortawn foxtail	
<i>Alopecurus aequalis</i> var. <i>aequalis</i>	Shortawn foxtail	
<i>Alopecurus alpinus</i>	Boreal alopecurus	
<i>Alyssum simplex</i>	Alyssum	
<i>Amaranthus albus</i>	Prostrate pigweed	
<i>Amaranthus blitoides</i>	Mat amaranth	
<i>Amaranthus retroflexus</i>	Redroot amaranth	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Ambrosia acanthicarpa</i>	Flatspine bur ragweed	
<i>Amelanchier alnifolia</i>	Saskatoon serviceberry	
<i>Amelanchier alnifolia</i> var. <i>alnifolia</i>	Saskatoon serviceberry	
<i>Amelanchier utahensis</i>	Utah serviceberry	
<i>Amelanchier utahensis</i> var. <i>utahensis</i>	Utah serviceberry	
<i>Anaphalis margaritacea</i>	Western pearly everlasting	
<i>Androsace chamaejasme</i>	Sweetflower rockjasmine	
<i>Androsace chamaejasme</i> ssp. <i>carinata</i>	Sweetflower rockjasmine	
<i>Androsace occidentalis</i>	Western rockjasmine	
<i>Androsace septentrionalis</i>	Pygmyflower rockjasmine	
<i>Anemone canadensis</i>	Canadian anemone	
<i>Anemone multifida</i>	Pacific anemone	
<i>Angelica ampla</i>	Giant angelica	
<i>Angelica grayi</i>	Gray's angelica	
<i>Antennaria anaphaloides</i>	Pearly pussytoes	
<i>Antennaria corymbosa</i>	Flat-top pussytoes	
<i>Antennaria marginata</i>	Whitemargin pussytoes	
<i>Antennaria media</i>	Rocky Mountain pussytoes	
<i>Antennaria microphylla</i>	Littleleaf pussytoes	
<i>Antennaria parvifolia</i>	Small-leaf pussytoes	
<i>Antennaria rosea</i>	Rosy pussytoes	
<i>Antennaria rosulata</i>	Kaibab pussytoes	
<i>Antennaria umbrinella</i>	Umber pussytoes	
<i>Apocynum androsaemifolium</i>	Spreading dogbane	
<i>Apocynum cannabinum</i>	Indianhemp	
<i>Aquilegia coerulea</i>	Colorado blue columbine	
<i>Aquilegia elegantula</i>	Western red columbine	
<i>Arabis</i> × <i>divaricarpa</i>	Spreadingpod rockcress	
<i>Arabis drummondii</i>	Drummond's rockcress	
<i>Arabis fendleri</i>	Fendler's rockcress	
<i>Arabis fendleri</i> var. <i>fendleri</i>	Fendler's rockcress	
<i>Arabis gunnisoniana</i>	Gunnison's rockcress	
<i>Arabis hirsuta</i>	Hairy rockcress	
<i>Arabis hirsuta</i> var. <i>pycnocarpa</i>	Creamflower rockcress	
<i>Arabis holboellii</i>	Holboell's rockcress	
<i>Arabis holboellii</i> var. <i>pinetorum</i>	Holboell's rockcress	
<i>Arabis lignifera</i>	Desert rockcress	
<i>Arabis oxylobula</i>	Glenwood Springs rockcress	
<i>Arctostaphylos uva-ursi</i>	Kinnikinnick	
<i>Arenaria fendleri</i>	Fendler's sandwort	
<i>Arenaria fendleri</i> var. <i>fendleri</i>	Fendler's sandwort	
<i>Arenaria hookeri</i>	Hooker's sandwort	
<i>Arenaria hookeri</i> ssp. <i>hookeri</i>	Hooker's sandwort	
<i>Arenaria lanuginosa</i>	Spreading sandwort	
<i>Arenaria lanuginosa</i> ssp. <i>saxosa</i>	Spreading sandwort	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Argentina anserina</i>	Silverweed cinquefoil	
<i>Argyrochosma fendleri</i>	Fendler's false cloak fern	
<i>Aristida purpurea</i>	Purple threeawn	
<i>Aristida purpurea</i> var. <i>longiseta</i>	Fendler threeawn	
<i>Aristida purpurea</i> var. <i>purpurea</i>	Purple threeawn	
<i>Arnica chamissonis</i>	Chamisso arnica	
<i>Arnica chamissonis</i> ssp. <i>foliosa</i>	Chamisso arnica	
<i>Arnica chamissonis</i> ssp. <i>foliosa</i> var. <i>andina</i>	Chamisso arnica	
<i>Arnica cordifolia</i>	Heartleaf arnica	
<i>Arnica mollis</i>	Hairy arnica	
<i>Artemisia biennis</i>	Biennial wormwood	
<i>Artemisia biennis</i> var. <i>biennis</i>	Biennial wormwood	
<i>Artemisia bigelovii</i>	Bigelow sage	
<i>Artemisia campestris</i>	Field sagewort	
<i>Artemisia campestris</i> ssp. <i>borealis</i>	Field sagewort	
<i>Artemisia campestris</i> ssp. <i>borealis</i> var. <i>borealis</i>	Field sagewort	
<i>Artemisia campestris</i> ssp. <i>borealis</i> var. <i>scouleriana</i>	Field sagewort	
<i>Artemisia campestris</i> ssp. <i>caudata</i>	Field sagewort	
<i>Artemisia cana</i>	Silver sagebrush	
<i>Artemisia cana</i> ssp. <i>cana</i>	Silver sagebrush	
<i>Artemisia carruthii</i>	Carruth's sagewort	
<i>Artemisia dracunculus</i>	Tarragon	
<i>Artemisia franserioides</i>	Ragweed sagebrush	
<i>Artemisia frigida</i>	Prairie sagewort	
<i>Artemisia longifolia</i>	Longleaf wormwood	
<i>Artemisia ludoviciana</i>	White sagebrush	
<i>Artemisia ludoviciana</i> ssp. <i>albula</i>	White sagebrush	
<i>Artemisia ludoviciana</i> ssp. <i>incompta</i>	White sagebrush	
<i>Artemisia ludoviciana</i> ssp. <i>ludoviciana</i>	White sagebrush	
<i>Artemisia michauxiana</i>	Michaux's wormwood	
<i>Artemisia parryi</i>	Parry's wormwood	
<i>Artemisia scopulorum</i>	Alpine sagebrush	
<i>Artemisia tridentata</i>	Big sagebrush	
<i>Artemisia tridentata</i> ssp. <i>tridentata</i>	Basin big sagebrush	
<i>Artemisia tridentata</i> ssp. <i>vaseyana</i>	Mountain big sagebrush	
<i>Asclepias hallii</i>	Hall's milkweed	
<i>Asclepias speciosa</i>	Showy milkweed	
<i>Asparagus officinalis</i>	Garden asparagus	
<i>Asplenium septentrionale</i>	Forked spleenwort	
<i>Aster alpinus</i>	Alpine aster	
<i>Aster alpinus</i> var. <i>vierhapperi</i>	Vierhapper's aster	
<i>Astragalus agrestis</i>	Purple milkvetch	
<i>Astragalus allochrous</i>	Halfmoon milkvetch	
<i>Astragalus allochrous</i> var. <i>playanus</i>	Halfmoon milkvetch	
<i>Astragalus alpinus</i>	Alpine milkvetch	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Astragalus alpinus</i> var. <i>alpinus</i>	Alpine milkvetch	
<i>Astragalus bisulcatus</i>	Twogrooved milkvetch	
<i>Astragalus bodinii</i>	Bodin's milkvetch	
<i>Astragalus brandegeei</i>	Brandegee's milkvetch	
<i>Astragalus ceramicus</i>	Painted milkvetch	
<i>Astragalus ceramicus</i> var. <i>ceramicus</i>	Painted milkvetch	
<i>Astragalus cerussatus</i>	Powdery milkvetch	
<i>Astragalus crassicaarpus</i>	Groundplum milkvetch	
<i>Astragalus crassicaarpus</i> var. <i>crassicaarpus</i>	Groundplum milkvetch	
<i>Astragalus drummondii</i>	Drummond's milkvetch	
<i>Astragalus flexuosus</i>	Flexile milkvetch	
<i>Astragalus flexuosus</i> var. <i>flexuosus</i>	Flexile milkvetch	
<i>Astragalus hallii</i>	Hall's milkvetch	
<i>Astragalus hallii</i> var. <i>hallii</i>	Hall's milkvetch	
<i>Astragalus kentrophyta</i>	Spiny milkvetch	
<i>Astragalus kentrophyta</i> var. <i>tegetarius</i>	Mat milkvetch	
<i>Astragalus laxmannii</i>	Laxmann's milkvetch	
<i>Astragalus laxmannii</i> var. <i>robustior</i>	Prairie milkvetch	
<i>Astragalus miser</i>	Timber milkvetch	
<i>Astragalus miser</i> var. <i>oblongifolius</i>	Timber milkvetch	
<i>Astragalus pattersonii</i>	Patterson's milkvetch	
<i>Astragalus ripleyi</i>	Ripley's milkvetch	
<i>Astragalus scopulorum</i>	Rocky Mountain milkvetch	
<i>Astragalus tenellus</i>	Looseflower milkvetch	
<i>Atriplex</i> × <i>aptera</i>	Moundscale	
<i>Atriplex argentea</i>	Silverscale saltbush	
<i>Atriplex canescens</i>	Fourwing saltbush	
<i>Atriplex canescens</i> var. <i>canescens</i>	Fourwing saltbush	
<i>Atriplex patula</i>	Spear saltbush	
<i>Atriplex rosea</i>	Tumbling saltweed	
<i>Atriplex truncata</i>	Wedgescale saltbush	
<i>Atriplex wolfii</i>	Wolf's saltweed	
<i>Bahia dissecta</i>	Ragleaf bahia	
<i>Balsamorhiza sagittata</i>	Arrowleaf balsamroot	
<i>Bassia hyssopifolia</i>	Fivehorn smotherweed	
<i>Bassia scoparia</i>	Burningbush	
<i>Beckmannia syzigachne</i>	American sloughgrass	
<i>Berberis fendleri</i>	Colorado barberry	
<i>Besseyia alpina</i>	Alpine besseyia	
<i>Besseyia plantaginea</i>	White River coraldrops	
<i>Betula occidentalis</i>	Water birch	
<i>Bidens cernua</i>	Nodding beggartick	
<i>Bidens frondosa</i>	Devil's beggartick	
<i>Bidens tenuisecta</i>	Slimlobe beggarticks	
<i>Bidens vulgata</i>	Big devils beggartick	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Blepharoneuron tricholepis</i>	Pine dropseed	
<i>Botrychium hesperium</i>	Western moonwort	
<i>Botrychium pinnatum</i>	Northern moonwort	
<i>Botrychium simplex</i>	Little grapefern	
<i>Bouteloua gracilis</i>	Blue grama	
<i>Bouteloua simplex</i>	Matted grama	
<i>Brassica juncea</i>	India mustard	
<i>Brassica napus</i>	Rape	
<i>Brickellia eupatorioides</i>	False boneset	
<i>Brickellia eupatorioides</i> var. <i>chlorolepis</i>	False boneset	
<i>Brickellia grandiflora</i>	Tasselflower brickellbush	
<i>Bromus ciliatus</i>	Fringed brome	
<i>Bromus ciliatus</i> var. <i>ciliatus</i>	Fringed brome	
<i>Bromus inermis</i>	Smooth brome	
<i>Bromus inermis</i> ssp. <i>inermis</i>	Smooth brome	
<i>Bromus inermis</i> ssp. <i>inermis</i> var. <i>inermis</i>	Smooth brome	
<i>Bromus lanatipes</i>	Woolly brome	
<i>Bromus porteri</i>	Porter brome	
<i>Bromus tectorum</i>	Cheatgrass	
<i>Calamagrostis canadensis</i>	Bluejoint	
<i>Calamagrostis purpurascens</i>	Purple reedgrass	
<i>Calamagrostis purpurascens</i> var. <i>purpurascens</i>	Purple reedgrass	
<i>Calamagrostis stricta</i>	Slimstem reedgrass	
<i>Callitriche palustris</i>	Vernal water-starwort	
<i>Calochortus gunnisonii</i>	Gunnison's mariposa lily	
<i>Calochortus gunnisonii</i> var. <i>gunnisonii</i>	Gunnison's mariposa lily	
<i>Caltha leptosepala</i>	White marsh marigold	
<i>Caltha leptosepala</i> ssp. <i>leptosepala</i>	White marsh marigold	
<i>Caltha leptosepala</i> ssp. <i>leptosepala</i> var. <i>leptosepala</i>	White marsh marigold	
<i>Camelina microcarpa</i>	Littlepod false flax	
<i>Campanula parryi</i>	Parry's bellflower	
<i>Campanula parryi</i> var. <i>parryi</i>	Parry's bellflower	
<i>Campanula rotundifolia</i>	Bluebell bellflower	
<i>Campanula uniflora</i>	Arctic bellflower	
<i>Capsella bursa-pastoris</i>	Shepherd's purse	
<i>Cardamine cordifolia</i>	Heartleaf bittercress	
<i>Cardamine cordifolia</i> var. <i>incana</i>	Heartleaf bittercress	
<i>Cardaria chalepensis</i>	Lenspod whitetop	
<i>Cardaria draba</i>	Whitetop	
<i>Cardaria pubescens</i>	Hairy whitetop	
<i>Carex albonigra</i>	Blackandwhite sedge	
<i>Carex aquatilis</i>	Water sedge	
<i>Carex aquatilis</i> var. <i>aquatilis</i>	Water sedge	
<i>Carex atherodes</i>	Wheat sedge	
<i>Carex aurea</i>	Golden sedge	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Carex bella</i>	Southwestern showy sedge	
<i>Carex brunnescens</i>	Brownish sedge	
<i>Carex brunnescens</i> ssp. <i>sphaerostachya</i>	Brownish sedge	
<i>Carex canescens</i>	Silvery sedge	
<i>Carex canescens</i> ssp. <i>canescens</i>	Silvery sedge	
<i>Carex diandra</i>	Lesser paniced sedge	
<i>Carex disperma</i>	Softleaf sedge	
<i>Carex douglasii</i>	Douglas' sedge	
<i>Carex duriuscula</i>	Needleleaf sedge	
<i>Carex ebenea</i>	Ebony sedge	
<i>Carex elynoides</i>	Blackroot sedge	
<i>Carex geophila</i>	White Mountain sedge	
<i>Carex geyeri</i>	Geyer's sedge	
<i>Carex hallii</i>	Deer sedge	
<i>Carex haydeniana</i>	Cloud sedge	
<i>Carex heteroneura</i>	Different-nerve sedge	
<i>Carex heteroneura</i> var. <i>brevisquama</i>	Different-nerve sedge	
<i>Carex heteroneura</i> var. <i>chalciolepis</i>	Holm sedge	
<i>Carex inops</i>	Long-stolon sedge	
<i>Carex inops</i> ssp. <i>heliophila</i>	Sun sedge	
<i>Carex microptera</i>	Smallwing sedge	
<i>Carex nebrascensis</i>	Nebraska sedge	
<i>Carex nelsonii</i>	Nelson's sedge	
<i>Carex nigricans</i>	Black alpine sedge	
<i>Carex norvegica</i>	Norway sedge	
<i>Carex norvegica</i> ssp. <i>stevenii</i>	Steven's sedge	
<i>Carex nova</i>	Black sedge	
<i>Carex obtusata</i>	Obtuse sedge	
<i>Carex occidentalis</i>	Western sedge	
<i>Carex parryana</i>	Parry's sedge	
<i>Carex parryana</i> var. <i>parryana</i>	Parry's sedge	
<i>Carex peltita</i>	Woolly sedge	
<i>Carex perglobosa</i>	Globe sedge	
<i>Carex phaeocephala</i>	Dunhead sedge	
<i>Carex praegracilis</i>	Clustered field sedge	
<i>Carex praticola</i>	Meadow sedge	
<i>Carex pyrenaica</i>	Pyrenean sedge	
<i>Carex pyrenaica</i> ssp. <i>pyrenaica</i>	Pyrenean sedge	
<i>Carex scopulorum</i>	Mountain sedge	
<i>Carex siccata</i>	Dryspike sedge	
<i>Carex simulata</i>	Analogue sedge	
<i>Carex utriculata</i>	Northwest Territory sedge	
<i>Carex vernacula</i>	Native sedge	
<i>Carex vesicaria</i>	Blister sedge	
<i>Carex vesicaria</i> var. <i>vesicaria</i>	Blister sedge	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Carum carvi</i>	Caraway	
<i>Castilleja flava</i>	Yellow Indian paintbrush	
<i>Castilleja flava</i> var. <i>flava</i>	Yellow Indian paintbrush	
<i>Castilleja haydenii</i>	Hayden's Indian paintbrush	
<i>Castilleja integra</i>	Wholeleaf Indian paintbrush	
<i>Castilleja integra</i> var. <i>integra</i>	Wholeleaf Indian paintbrush	
<i>Castilleja linariifolia</i>	Wyoming Indian paintbrush	
<i>Castilleja miniata</i>	Giant red Indian paintbrush	
<i>Castilleja miniata</i> ssp. <i>miniata</i>	Giant red Indian paintbrush	
<i>Castilleja occidentalis</i>	Western Indian paintbrush	
<i>Castilleja rhexiifolia</i>	Splitleaf Indian paintbrush	
<i>Castilleja sulphurea</i>	Sulphur Indian paintbrush	
<i>Ceanothus fendleri</i>	Fendler's ceanothus	
<i>Ceanothus velutinus</i>	Snowbrush ceanothus	
<i>Ceanothus velutinus</i> var. <i>velutinus</i>	Snowbrush ceanothus	
<i>Cerastium arvense</i>	Field chickweed	
<i>Cerastium arvense</i> ssp. <i>strictum</i>	Feld chickweed	
<i>Cerastium beeringianum</i>	Bering chickweed	
<i>Cerastium beeringianum</i> ssp. <i>earlei</i>	Bering chickweed	
<i>Cercocarpus montanus</i>	Alderleaf mountain mahogany	
<i>Chaenactis douglasii</i>	Douglas' dustymaiden	
<i>Chaenactis douglasii</i> var. <i>alpina</i>	Alpine dustymaiden	
<i>Chaetopappa ericoides</i>	Rose heath	
<i>Chamaerhodos erecta</i>	Little rose	
<i>Chamaerhodos erecta</i> ssp. <i>nuttallii</i>	Nuttall's little rose	
<i>Chamaesyce serpyllifolia</i>	Thymeleaf sandmat	
<i>Chamaesyce serpyllifolia</i> ssp. <i>serpyllifolia</i>	Thymeleaf sandmat	
<i>Chamerion angustifolium</i>	Fireweed	
<i>Chamerion angustifolium</i> ssp. <i>circumvagum</i>	Fireweed	
<i>Cheilanthes feei</i>	Slender lipfern	
<i>Cheilanthes fendleri</i>	Fendler's lipfern	
<i>Chenopodium album</i>	Lambsquarters	
<i>Chenopodium atrovirens</i>	Pinyon goosefoot	
<i>Chenopodium berlandieri</i>	Pitseed goosefoot	
<i>Chenopodium botrys</i>	Jerusalem oak goosefoot	
<i>Chenopodium desiccatum</i>	Aridland goosefoot	
<i>Chenopodium foliosum</i>	Leafy goosefoot	
<i>Chenopodium fremontii</i>	Fremont's goosefoot	
<i>Chenopodium fremontii</i> var. <i>fremontii</i>	Fremont's goosefoot	
<i>Chenopodium glaucum</i>	Oakleaf goosefoot	
<i>Chenopodium graveolens</i>	Fetid goosefoot	
<i>Chenopodium leptophyllum</i>	Narrowleaf goosefoot	
<i>Chenopodium pratericola</i>	Desert goosefoot	
<i>Chenopodium rubrum</i>	Red goosefoot	
<i>Chenopodium watsonii</i>	Watson's goosefoot	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Chionophila jamesii</i>	Rocky Mountain snowlover	
<i>Chrysothamnus greenei</i>	Greene's rabbitbrush	
<i>Chrysothamnus vaseyi</i>	Vasey's rabbitbrush	
<i>Chrysothamnus viscidiflorus</i>	Yellow rabbitbrush	
<i>Chrysothamnus viscidiflorus</i> ssp. <i>lanceolatus</i>	Yellow rabbitbrush	
<i>Cicuta maculata</i>	Spotted water hemlock	
<i>Cirsium arvense</i>	Canada thistle	
<i>Cirsium canescens</i>	Prairie thistle	
<i>Cirsium centaureae</i>	Fringed thistle	
<i>Cirsium ochrocentrum</i>	Yellowspine thistle	
<i>Cirsium ochrocentrum</i> ssp. <i>ochrocentrum</i>	Yellowspine thistle	
<i>Cirsium pallidum</i>	Pale thistle	
<i>Cirsium parryi</i>	Parry's thistle	
<i>Cirsium parryi</i> ssp. <i>parryi</i>	Parry's thistle	
<i>Cirsium scariosum</i>	Meadow thistle	
<i>Cirsium scopulorum</i>	Mountain thistle	
<i>Claytonia megarhiza</i>	Alpine springbeauty	
<i>Claytonia megarhiza</i> var. <i>megarhiza</i>	Alpine springbeauty	
<i>Clematis columbiana</i>	Rock clematis	
<i>Clematis columbiana</i> var. <i>columbiana</i>	Rock clematis	
<i>Clematis hirsutissima</i>	Hairy clematis	
<i>Clematis hirsutissima</i> var. <i>scottii</i>	Scott's clematis	
<i>Clematis ligusticifolia</i>	Western white clematis	
<i>Clematis ligusticifolia</i> var. <i>ligusticifolia</i>	Western white clematis	
<i>Cleome multicaulis</i>	Slender spiderflower	
<i>Cleome serrulata</i>	Rocky Mountain beeplant	
<i>Collomia linearis</i>	Tiny trumpet	
<i>Comandra umbellata</i>	Bastard toadflax	
<i>Comandra umbellata</i> ssp. <i>pallida</i>	Pale bastard toadflax	
<i>Comarum palustre</i>	Purple marshlocks	
<i>Conioselinum scopulorum</i>	Rocky Mountain hemlockparsley	
<i>Convolvulus arvensis</i>	Field bindweed	
<i>Conyza canadensis</i>	Canadian horseweed	
<i>Corallorhiza maculata</i>	Summer coralroot	
<i>Corallorhiza striata</i>	Hooded coralroot	
<i>Corallorhiza trifida</i>	Yellow coralroot	
<i>Coreopsis tinctoria</i>	Golden tickseed	
<i>Coreopsis tinctoria</i> var. <i>tinctoria</i>	Golden tickseed	
<i>Corispermum americanum</i>	American bugseed	
<i>Corispermum americanum</i> var. <i>rydbergii</i>	American bugseed	
<i>Corispermum villosum</i>	Hairy bugseed	
<i>Cornus canadensis</i>	Bunchberry dogwood	
<i>Cornus sericea</i>	Redosier dogwood	
<i>Cornus sericea</i> ssp. <i>sericea</i>	Redosier dogwood	
<i>Corydalis aurea</i>	Scrambled eggs	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Corydalis caseana</i>	Sierra fumewort	
<i>Corydalis caseana</i> ssp. <i>brandegeei</i>	Brandegee's fumewort	
<i>Corydalis curvisiliqua</i>	Curvepod fumewort	
<i>Corydalis curvisiliqua</i> ssp. <i>occidentalis</i>	Curvepod fumewort	
<i>Crataegus rivularis</i>	River hawthorn	
<i>Crepis occidentalis</i>	Largeflower hawksbeard	
<i>Crepis occidentalis</i> ssp. <i>occidentalis</i>	Largeflower hawksbeard	
<i>Crepis runcinata</i>	Fiddleleaf hawksbeard	
<i>Crepis runcinata</i> ssp. <i>runcinata</i>	Fiddleleaf hawksbeard	
<i>Cryptantha bakeri</i>	Baker's cryptantha	
<i>Cryptantha cinerea</i>	James' cryptantha	
<i>Cryptantha cinerea</i> var. <i>jamesii</i>	James' cryptantha	
<i>Cryptantha cinerea</i> var. <i>pustulosa</i>	James' cryptantha	
<i>Cryptantha fendleri</i>	Sanddune cryptantha	
<i>Cryptantha minima</i>	Little cryptantha	
<i>Cryptantha weberi</i>	Weber's cryptantha	
<i>Cryptogramma acrostichoides</i>	American rockbrake	
<i>Cycloloma atriplicifolium</i>	Winged pigweed	
<i>Cymopterus acaulis</i>	Plains springparsley	
<i>Cymopterus montanus</i>	Mountain springparsley	
<i>Cynoglossum officinale</i>	Gypsyflower	
<i>Cyperus squarrosus</i>	Bearded flatsedge	
<i>Cystopteris fragilis</i>	Brittle bladderfern	
<i>Cystopteris reevesiana</i>	Reeves' bladderfern	
<i>Dalea leporina</i>	Foxtail prairie clover	
<i>Danthonia californica</i>	California oatgrass	
<i>Danthonia intermedia</i>	Timber oatgrass	
<i>Danthonia parryi</i>	Parry's oatgrass	
<i>Dasiphora fruticosa</i>	Shrubby cinquefoil	
<i>Dasiphora fruticosa</i> ssp. <i>floribunda</i>	Shrubby cinquefoil	
<i>Delphinium alpestre</i>	Colorado larkspur	
<i>Delphinium barbeyi</i>	Subalpine larkspur	
<i>Delphinium nuttallianum</i>	Twolobe larkspur	
<i>Delphinium ramosum</i>	Mountain larkspur	
<i>Delphinium robustum</i>	Wahatoya Creek larkspur	
<i>Deschampsia cespitosa</i>	Tufted hairgrass	
<i>Descurainia incana</i>	Mountain tansymustard	
<i>Descurainia incana</i> ssp. <i>incisa</i>	Mountain tansymustard	
<i>Descurainia incana</i> ssp. <i>viscosa</i>	Mountain tansymustard	
<i>Descurainia pinnata</i>	Western tansymustard	
<i>Descurainia pinnata</i> ssp. <i>filipes</i>	Western tansymustard	
<i>Descurainia ramosissima</i>	Villa Grove tansymustard	
<i>Distichlis spicata</i>	Saltgrass	
<i>Dodecatheon pulchellum</i>	Darkthroat shootingstar	
<i>Dodecatheon pulchellum</i> ssp. <i>pulchellum</i>	Darkthroat shootingstar	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Draba aurea</i>	Golden draba	
<i>Draba crassa</i>	Thickleaf draba	
<i>Draba crassifolia</i>	Snowbed draba	
<i>Draba fladnizensis</i>	Austrian draba	
<i>Draba grayana</i>	Gray's draba	
<i>Draba helleriana</i>	Heller's draba	
<i>Draba helleriana</i> var. <i>helleriana</i>	Heller's draba	
<i>Draba rectifruta</i>	Mountain draba	
<i>Draba smithii</i>	Smith's draba	
<i>Draba spectabilis</i>	Showy draba	
<i>Draba streptobrachia</i>	Alpine tundra draba	
<i>Draba streptocarpa</i>	Pretty draba	
<i>Dracocephalum parviflorum</i>	American dragonhead	
<i>Dryas octopetala</i>	Eightpetal mountain-avens	
<i>Dryas octopetala</i> ssp. <i>hookeriana</i>	Hooker's mountain-avens	
<i>Dryopteris filix-mas</i>	Male fern	
<i>Dyssodia papposa</i>	Fetid marigold	
<i>Echinocereus triglochidiatus</i>	Kingcup cactus	
<i>Echinocereus triglochidiatus</i> var. <i>triglochidiatus</i>	Kingcup cactus	
<i>Echinocereus viridiflorus</i>	Nylon hedgehog cactus	
<i>Echinocereus viridiflorus</i> var. <i>viridiflorus</i>	Nylon hedgehog cactus	
<i>Echinochloa crus-galli</i>	Barnyardgrass	
<i>Echinocystis lobata</i>	Wild cucumber	
<i>Elaeagnus commutata</i>	Silverberry	
<i>Eleocharis acicularis</i>	Needle spikerush	
<i>Eleocharis palustris</i>	Common spikerush	
<i>Eleocharis palustris</i> var. <i>palustris</i>	Common spikerush	
<i>Eleocharis quinqueflora</i>	Fewflower spikerush	
× <i>Elyhordeum macounii</i>	Macoun's barley	
<i>Elymus canadensis</i>	Canada wildrye	
<i>Elymus elymoides</i>	Squirreltail	
<i>Elymus elymoides</i> ssp. <i>brevifolius</i>	Squirreltail	
<i>Elymus lanceolatus</i>	Thickspike wheatgrass	
<i>Elymus lanceolatus</i> ssp. <i>lanceolatus</i>	Thickspike wheatgrass	
<i>Elymus repens</i>	Quackgrass	
<i>Elymus scribneri</i>	Spreading wheatgrass	
<i>Elymus trachycaulus</i>	Slender wheatgrass	
<i>Elymus trachycaulus</i> ssp. <i>trachycaulus</i>	Slender wheatgrass	
<i>Epilobium brachycarpum</i>	Tall annual willowherb	
<i>Epilobium ciliatum</i>	Fringed willowherb	
<i>Epilobium ciliatum</i> ssp. <i>glandulosum</i>	Fringed willowherb	
<i>Epilobium halleanum</i>	Glandular willowherb	
<i>Epilobium hornemannii</i>	Hornemann's willowherb	
<i>Epilobium hornemannii</i> ssp. <i>hornemannii</i>	Hornemann's willowherb	
<i>Epilobium saximontanum</i>	Rocky Mountain willowherb	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Equisetum arvense</i>	Field horsetail	
<i>Equisetum hyemale</i>	Scouringrush horsetail	
<i>Equisetum hyemale</i> var. <i>affine</i>	Scouringrush horsetail	
<i>Equisetum laevigatum</i>	Smooth horsetail	
<i>Equisetum pratense</i>	Meadow horsetail	
<i>Equisetum variegatum</i>	Variegated scouringrush	
<i>Equisetum variegatum</i> var. <i>variegatum</i>	Variegated scouringrush	
<i>Eragrostis pilosa</i>	Indian lovegrass	
<i>Ericameria nauseosa</i>	Rubber rabbitbrush	
<i>Ericameria nauseosa</i> ssp. <i>consimilis</i>	Rubber rabbitbrush	
<i>Ericameria nauseosa</i> ssp. <i>consimilis</i> var. <i>oreophila</i>	Rubber rabbitbrush	
<i>Ericameria nauseosa</i> ssp. <i>nauseosa</i>	Rubber rabbitbrush	
<i>Ericameria nauseosa</i> ssp. <i>nauseosa</i> var. <i>bigelovii</i>	Rubber rabbitbrush	
<i>Ericameria nauseosa</i> ssp. <i>nauseosa</i> var. <i>glabrata</i>	Rubber rabbitbrush	
<i>Ericameria nauseosa</i> ssp. <i>nauseosa</i> var. <i>nauseosa</i>	Rubber rabbitbrush	
<i>Ericameria parryi</i>	Parry's rabbitbrush	
<i>Ericameria parryi</i> var. <i>affinis</i>	Parry's rabbitbrush	
<i>Ericameria parryi</i> var. <i>parryi</i>	Parry's rabbitbrush	
<i>Erigeron acris</i>	Bitter fleabane	
<i>Erigeron acris</i> ssp. <i>debilis</i>	Bitter fleabane	
<i>Erigeron acris</i> ssp. <i>politus</i>	Bitter fleabane	
<i>Erigeron canus</i>	Hoary fleabane	
<i>Erigeron colomexicanus</i>	Running fleabane	
<i>Erigeron compositus</i>	Cutleaf daisy	
<i>Erigeron coulteri</i>	Large mountain fleabane	
<i>Erigeron divergens</i>	Spreading fleabane	
<i>Erigeron elatior</i>	Tall fleabane	
<i>Erigeron engelmannii</i>	Engelmann's fleabane	
<i>Erigeron engelmannii</i> var. <i>engelmannii</i>	Engelmann's fleabane	
<i>Erigeron eximius</i>	Sprucefir fleabane	
<i>Erigeron flagellaris</i>	Trailing fleabane	
<i>Erigeron formosissimus</i>	Beautiful fleabane	
<i>Erigeron glabellus</i>	Streamside fleabane	
<i>Erigeron leiomerus</i>	Rockslide yellow fleabane	
<i>Erigeron lonchophyllus</i>	Shortray fleabane	
<i>Erigeron melanocephalus</i>	Blackhead fleabane	
<i>Erigeron peregrinus</i>	Subalpine fleabane	
<i>Erigeron peregrinus</i> ssp. <i>callianthemus</i>	Subalpine fleabane	
<i>Erigeron peregrinus</i> ssp. <i>callianthemus</i> var. <i>callianthemus</i>	Subalpine fleabane	
<i>Erigeron philadelphicus</i>	Philadelphia fleabane	
<i>Erigeron philadelphicus</i> var. <i>philadelphicus</i>	Philadelphia fleabane	
<i>Erigeron pinnatisectus</i>	Featherleaf fleabane	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Erigeron pumilus</i>	Shaggy fleabane	
<i>Erigeron pumilus</i> ssp. <i>pumilus</i>	Shaggy fleabane	
<i>Erigeron simplex</i>	Onestem fleabane	
<i>Erigeron speciosus</i>	Aspen fleabane	
<i>Erigeron speciosus</i> var. <i>speciosus</i>	Aspen fleabane	
<i>Erigeron subtrinervis</i>	Threenerve fleabane	
<i>Erigeron subtrinervis</i> var. <i>subtrinervis</i>	Threenerve fleabane	
<i>Erigeron ursinus</i>	Bear River fleabane	
<i>Erigeron vetensis</i>	Early bluetop fleabane	
<i>Erigeron vreelandii</i>	Vreeland's erigeron	
<i>Eriodictyon angustifolium</i>	Narrowleaf yerba santa	
<i>Eriogonum alatum</i>	Winged buckwheat	
<i>Eriogonum alatum</i> var. <i>alatum</i>	Winged buckwheat	
<i>Eriogonum cernuum</i>	Nodding buckwheat	
<i>Eriogonum cernuum</i> var. <i>cernuum</i>	Nodding buckwheat	
<i>Eriogonum coloradense</i>	Colorado buckwheat	
<i>Eriogonum effusum</i>	Spreading buckwheat	
<i>Eriogonum effusum</i> var. <i>effusum</i>	Spreading buckwheat	
<i>Eriogonum jamesii</i>	James' buckwheat	
<i>Eriogonum jamesii</i> var. <i>flavescens</i>	James' buckwheat	
<i>Eriogonum jamesii</i> var. <i>jamesii</i>	James' buckwheat	
<i>Eriogonum jamesii</i> var. <i>xanthum</i>	James' buckwheat	
<i>Eriogonum lachnogynum</i>	Woollycup buckwheat	
<i>Eriogonum microthecum</i>	Slender buckwheat	
<i>Eriogonum racemosum</i>	Redroot buckwheat	
<i>Eriogonum umbellatum</i>	Sulphur-flower buckwheat	
<i>Eriogonum umbellatum</i> var. <i>aureum</i>	Sulphur-flower buckwheat	
<i>Eriogonum umbellatum</i> var. <i>majus</i>	Sulphur-flower buckwheat	
<i>Eriogonum umbellatum</i> var. <i>umbellatum</i>	Sulphur-flower buckwheat	
<i>Eriophorum angustifolium</i>	Tall cottongrass	
<i>Eriophorum angustifolium</i> ssp. <i>angustifolium</i>	Tall cottongrass	
<i>Eritrichium nanum</i>	Arctic alpine forget-me-not	
<i>Erysimum capitatum</i>	Sanddune wallflower	
<i>Erysimum capitatum</i> var. <i>capitatum</i>	Sanddune wallflower	
<i>Erysimum cheiranthoides</i>	Wormseed wallflower	
<i>Erysimum inconspicuum</i>	Shy wallflower	
<i>Erysimum inconspicuum</i> var. <i>inconspicuum</i>	Shy wallflower	
<i>Escobaria vivipara</i>	Spinystar	
<i>Escobaria vivipara</i> var. <i>vivipara</i>	Spinystar	
<i>Euphorbia brachycera</i>	Horned spurge	
<i>Euthamia graminifolia</i>	Flat-top goldentop	
<i>Euthamia graminifolia</i> var. <i>graminifolia</i>	Flat-top goldentop	
<i>Euthamia occidentalis</i>	Western goldentop	
<i>Fallugia paradoxa</i>	Apache plume	
<i>Festuca arizonica</i>	Arizona fescue	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Festuca brachyphylla</i>	Alpine fescue	
<i>Festuca brachyphylla</i> ssp. <i>coloradensis</i>	Colorado fescue	
<i>Festuca earlei</i>	Earle's fescue	
<i>Festuca idahoensis</i>	Idaho fescue	
<i>Festuca idahoensis</i> ssp. <i>idahoensis</i>	Idaho fescue	
<i>Festuca minutiflora</i>	Smallflower fescue	
<i>Festuca rubra</i>	Red fescue	
<i>Festuca saximontana</i>	Rocky Mountain fescue	
<i>Festuca sororia</i>	Ravine fescue	
<i>Festuca thurberi</i>	Thurber's fescue	
<i>Fragaria vesca</i>	Woodland strawberry	
<i>Fragaria vesca</i> ssp. <i>bracteata</i>	Woodland strawberry	
<i>Fragaria virginiana</i>	Virginia strawberry	
<i>Fragaria virginiana</i> ssp. <i>glauca</i>	Virginia strawberry	
<i>Frasera speciosa</i>	Elkweed	
<i>Gaillardia aristata</i>	Blanketflower	
<i>Galium boreale</i>	Northern bedstraw	
<i>Galium trifidum</i>	Threepetal bedstraw	
<i>Galium trifidum</i> ssp. <i>subbiflorum</i>	Threepetal bedstraw	
<i>Gaura coccinea</i>	Scarlet beeblossom	
<i>Gayophytum diffusum</i>	Spreading groundsmoke	
<i>Gayophytum diffusum</i> ssp. <i>parviflorum</i>	Spreading groundsmoke	
<i>Gayophytum ramosissimum</i>	Pinyon groundsmoke	
<i>Gentiana affinis</i>	Pleated gentian	
<i>Gentiana algida</i>	Whitish gentian	
<i>Gentiana fremontii</i>	Moss gentian	
<i>Gentiana parryi</i>	Parry's gentian	
<i>Gentiana prostrata</i>	Pygmy gentian	
<i>Gentianella amarella</i>	Autumn dwarf gentian	
<i>Gentianella amarella</i> ssp. <i>acuta</i>	Autumn dwarf gentian	
<i>Gentianella amarella</i> ssp. <i>heterosepala</i>	Autumn dwarf gentian	
<i>Gentianella tenella</i>	Dane's dwarf gentian	
<i>Gentianella tenella</i> ssp. <i>tenella</i>	Dane's dwarf gentian	
<i>Gentianopsis barbellata</i>	Perennial fringed gentian	
<i>Gentianopsis thermalis</i>	Rocky Mountain fringed gentian	
<i>Geranium caespitosum</i>	Pineywoods geranium	
<i>Geranium caespitosum</i> var. <i>caespitosum</i>	Pineywoods geranium	
<i>Geranium richardsonii</i>	Richardson's geranium	
<i>Geum aleppicum</i>	Yellow avens	
<i>Geum macrophyllum</i>	Largeleaf avens	
<i>Geum macrophyllum</i> var. <i>perincisum</i>	Largeleaf avens	
<i>Geum rivale</i>	Purple avens	
<i>Geum rossii</i>	Ross' avens	
<i>Geum rossii</i> var. <i>turbinatum</i>	Ross' avens	
<i>Geum triflorum</i>	Old man's whiskers	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Geum triflorum</i> var. <i>triflorum</i>	Old man's whiskers	
<i>Glaux maritima</i>	Sea milkwort	
<i>Glyceria grandis</i>	American mannagrass	
<i>Glyceria grandis</i> var. <i>grandis</i>	American mannagrass	
<i>Glyceria striata</i>	Fowl mannagrass	
<i>Glycyrrhiza lepidota</i>	American licorice	
<i>Gnaphalium uliginosum</i>	Marsh cudweed	
<i>Goodyera oblongifolia</i>	Western rattlesnake plantain	
<i>Goodyera repens</i>	Lesser rattlesnake plantain	
<i>Gratiola neglecta</i>	Clammy hedgehyssop	
<i>Grindelia decumbens</i>	Reclined gumweed	
<i>Grindelia decumbens</i> var. <i>decumbens</i>	Reclined gumweed	
<i>Grindelia nuda</i>	Curlytop gumweed	
<i>Grindelia nuda</i> var. <i>aphanactis</i>	Curlytop gumweed	
<i>Grindelia squarrosa</i>	Curlycup gumweed	
<i>Gutierrezia sarothrae</i>	Broom snakeweed	
<i>Gymnocarpium dryopteris</i>	Western oakfern	
<i>Hackelia floribunda</i>	Manyflower stickseed	
<i>Halogeton glomeratus</i>	Saltlover	
<i>Hedysarum occidentale</i>	Western sweetvetch	
<i>Helianthella parryi</i>	Parry's dwarf-sunflower	
<i>Helianthella quinquenervis</i>	Fivenerve helianthella	
<i>Helianthus annuus</i>	Common sunflower	
<i>Helianthus nuttallii</i>	Nuttall's sunflower	
<i>Helianthus petiolaris</i>	Prairie sunflower	
<i>Heliomeris multiflora</i>	Showy goldeneye	
<i>Heliotropium curassavicum</i>	Salt heliotrope	
<i>Heliotropium curassavicum</i> var. <i>obovatum</i>	Seaside heliotrope	
<i>Heracleum maximum</i>	Common cowparsnip	
<i>Hesperostipa comata</i>	Needle and thread	
<i>Hesperostipa comata</i> ssp. <i>comata</i>	Needle and thread	
<i>Hesperostipa neomexicana</i>	New Mexico feathergrass	
<i>Heterotheca fulcrata</i>	Rockyscree false goldenaster	
<i>Heterotheca pumila</i>	Alpine false goldenaster	
<i>Heterotheca villosa</i>	Hairy false goldenaster	
<i>Heterotheca villosa</i> var. <i>minor</i>	Hairy false goldenaster	
<i>Heterotheca villosa</i> var. <i>nana</i>	Hairy false goldenaster	
<i>Heterotheca villosa</i> var. <i>villosa</i>	Hairy false goldenaster	
<i>Heuchera parvifolia</i>	Littleleaf alumroot	
<i>Heuchera parvifolia</i> var. <i>parvifolia</i>	Littleleaf alumroot	
<i>Hieracium gracile</i>	Slender hawkweed	
<i>Hieracium gracile</i> var. <i>gracile</i>	Slender hawkweed	
<i>Hierochloa hirta</i>	Northern sweetgrass	
<i>Hierochloa hirta</i> ssp. <i>arctica</i>	Northern sweetgrass	
<i>Hoffmannseggia glauca</i>	Indian rushpea	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Holodiscus dumosus</i>	Rockspirea	
<i>Hordeum brachyantherum</i>	Meadow barley	
<i>Hordeum brachyantherum</i> ssp. <i>brachyantherum</i>	Meadow barley	
<i>Hordeum jubatum</i>	Foxtail barley	
<i>Hordeum jubatum</i> ssp. <i>jubatum</i>	Foxtail barley	
<i>Humulus lupulus</i>	Common hop	
<i>Humulus lupulus</i> var. <i>neomexicanus</i>	Common hop	
<i>Hydrophyllum fendleri</i>	Fendler's waterleaf	
<i>Hydrophyllum fendleri</i> var. <i>fendleri</i>	Fendler's waterleaf	
<i>Hymenopappus filifolius</i>	Fineleaf hymenopappus	
<i>Hymenopappus filifolius</i> var. <i>cinereus</i>	Fineleaf hymenopappus	
<i>Hymenopappus filifolius</i> var. <i>parvulus</i>	Fineleaf hymenopappus	
<i>Hymenopappus newberryi</i>	Newberry's hymenopappus	
<i>Hymenoxyis helenioides</i>	Intermountain rubberweed	
<i>Hymenoxyis hoopesii</i>	Owl's-claws	
<i>Hymenoxyis richardsonii</i>	Pingue rubberweed	
<i>Hymenoxyis richardsonii</i> var. <i>richardsonii</i>	Pingue rubberweed	
<i>Hyoscyamus niger</i>	Black henbane	
<i>Hypericum scouleri</i>	Scouler's St. Johnswort	
<i>Hypericum scouleri</i> ssp. <i>nortoniae</i>	Norton's St. Johnswort	
<i>Ipomopsis aggregata</i>	Scarlet gilia	
<i>Ipomopsis aggregata</i> ssp. <i>candida</i>	Scarlet gilia	
<i>Ipomopsis aggregata</i> ssp. <i>collina</i>	Scarlet gilia	
<i>Ipomopsis longiflora</i>	Flaxflowered ipomopsis	
<i>Ipomopsis longiflora</i> ssp. <i>longiflora</i>	Flaxflowered ipomopsis	
<i>Ipomopsis multiflora</i>	Manyflowered ipomopsis	
<i>Iris missouriensis</i>	Rocky Mountain iris	
<i>Iva axillaris</i>	Povertyweed	
<i>Ivesia gordonii</i>	Gordon's ivesia	
<i>Jamesia americana</i>	Fivepetal cliffbush	
<i>Jamesia americana</i> var. <i>americana</i>	Fivepetal cliffbush	
<i>Juncus arcticus</i>	Arctic rush	
<i>Juncus arcticus</i> ssp. <i>littoralis</i>	Mountain rush	
<i>Juncus bufonius</i>	Toad rush	
<i>Juncus bufonius</i> var. <i>bufonius</i>	Toad rush	
<i>Juncus castaneus</i>	Chestnut rush	
<i>Juncus castaneus</i> ssp. <i>castaneus</i>	Chestnut rush	
<i>Juncus castaneus</i> ssp. <i>castaneus</i> var. <i>castaneus</i>	Chestnut rush	
<i>Juncus drummondii</i>	Drummond's rush	
<i>Juncus interior</i>	Inland rush	
<i>Juncus longistylis</i>	Longstyle rush	
<i>Juncus longistylis</i> var. <i>longistylis</i>	Longstyle rush	
<i>Juncus mertensianus</i>	Mertens' rush	
<i>Juncus saximontanus</i>	Rocky Mountain rush	
<i>Juncus torreyi</i>	Torrey's rush	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Juniperus communis</i>	Common juniper	
<i>Juniperus communis</i> var. <i>depressa</i>	Common juniper	
<i>Juniperus scopulorum</i>	Rocky Mountain juniper	
<i>Kalmia microphylla</i>	Alpine laurel	
<i>Kobresia myosuroides</i>	Bellardi bog sedge	
<i>Koeleria macrantha</i>	Prairie Junegrass	
<i>Krascheninnikovia lanata</i>	Winterfat	
<i>Lactuca tatarica</i>	Blue lettuce	
<i>Lactuca tatarica</i> var. <i>pulchella</i>	Blue lettuce	
<i>Lappula occidentalis</i>	Flatspine stickseed	
<i>Lappula occidentalis</i> var. <i>occidentalis</i>	Flatspine stickseed	
<i>Lathyrus eucosmus</i>	Bush vetchling	
<i>Lathyrus lanszwertii</i>	Nevada pea	
<i>Lathyrus lanszwertii</i> var. <i>leucanthus</i>	Nevada pea	
<i>Lathyrus latifolius</i>	Perennial pea	
<i>Lemna minuta</i>	Least duckweed	
<i>Lemna turionifera</i>	Turion duckweed	
<i>Lepidium alyssoides</i>	Mesa pepperwort	
<i>Lepidium alyssoides</i> var. <i>alyssoides</i>	Mesa pepperwort	
<i>Lepidium densiflorum</i>	Common pepperweed	
<i>Lepidium latifolium</i>	Broadleaved pepperweed	
<i>Lepidium ramosissimum</i>	Manybranched pepperweed	
<i>Leptochloa fusca</i>	Malabar sprangletop	
<i>Leptochloa fusca</i> ssp. <i>fascicularis</i>	Bearded sprangletop	
<i>Leptosiphon nuttallii</i>	Nuttall's linanthus	
<i>Leptosiphon nuttallii</i> ssp. <i>nuttallii</i>	Nuttall's linanthus	
<i>Lesquerella montana</i>	Mountain bladderpod	
<i>Levisticum officinale</i>	Garden lovage	
<i>Lewisia pygmaea</i>	Alpine lewisia	
<i>Leymus ambiguus</i>	Colorado wildrye	
<i>Leymus cinereus</i>	Basin wildrye	
<i>Leymus triticoides</i>	Beardless wildrye	
<i>Liatris punctata</i>	Dotted blazing star	
<i>Ligusticum porteri</i>	Porter's licorice-root	
<i>Ligusticum porteri</i> var. <i>porteri</i>	Porter's licorice-root	
<i>Limosella aquatica</i>	Water mudwort	
<i>Linanthus pungens</i>	Granite prickly phlox	
<i>Linnaea borealis</i>	Twinflower	
<i>Linnaea borealis</i> ssp. <i>americana</i>	Twinflower	
<i>Linum australe</i>	Southern flax	
<i>Linum australe</i> var. <i>australe</i>	Southern flax	
<i>Linum lewisii</i>	Lewis flax	
<i>Linum lewisii</i> var. <i>lewisii</i>	Prairie flax	
<i>Listera cordata</i>	Heartleaf twayblade	
<i>Listera cordata</i> var. <i>nephrophylla</i>	Heartleaf twayblade	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Lithophragma tenellum</i>	Slender woodland-star	
<i>Lithospermum incisum</i>	Narrowleaf stoneseed	
<i>Lithospermum multiflorum</i>	Manyflowered stoneseed	
<i>Lloydia serotina</i>	Common alplily	
<i>Lloydia serotina</i> var. <i>serotina</i>	Common alplily	
<i>Lonicera involucrata</i>	Twinberry honeysuckle	
<i>Lonicera involucrata</i> var. <i>involucrata</i>	Twinberry honeysuckle	
<i>Lupinus argenteus</i>	Silvery lupine	
<i>Lupinus bakeri</i>	Baker's lupine	
<i>Lupinus bakeri</i> ssp. <i>bakeri</i>	Baker's lupine	
<i>Lupinus caespitosus</i>	Stemless dwarf lupine	
<i>Lupinus caespitosus</i> var. <i>caespitosus</i>	Stemless dwarf lupine	
<i>Lupinus caudatus</i>	Taileup lupine	
<i>Lupinus kingii</i>	King's lupine	
<i>Lupinus pusillus</i>	Rusty lupine	
<i>Lupinus pusillus</i> ssp. <i>pusillus</i>	Rusty lupine	
<i>Lupinus sericeus</i>	Silky lupine	
<i>Lupinus sericeus</i> ssp. <i>sericeus</i>	Silky lupine	
<i>Luzula parviflora</i>	Smallflowered woodrush	
<i>Luzula spicata</i>	Spiked woodrush	
<i>Lycopus asper</i>	Rough bugleweed	
<i>Lygodesmia juncea</i>	Rush skeletonplant	
<i>Machaeranthera bigelovii</i>	Bigelow's tansyaster	
<i>Machaeranthera bigelovii</i> var. <i>bigelovii</i>	Bigelow's tansyaster	
<i>Machaeranthera canescens</i>	Hoary tansyaster	
<i>Machaeranthera canescens</i> ssp. <i>glabra</i>	Hoary tansyaster	
<i>Machaeranthera canescens</i> ssp. <i>glabra</i> var. <i>glabra</i>	Hoary tansyaster	
<i>Machaeranthera coloradoensis</i>	Colorado tansyaster	
<i>Machaeranthera coloradoensis</i> var. <i>coloradoensis</i>	Colorado tansyaster	
<i>Machaeranthera parviflora</i>	Smallflower tansyaster	
<i>Machaeranthera pinnatifida</i>	Lacy tansyaster	
<i>Machaeranthera pinnatifida</i> ssp. <i>pinnatifida</i>	Lacy tansyaster	
<i>Machaeranthera pinnatifida</i> ssp. <i>pinnatifida</i> var. <i>pinnatifida</i>	Lacy tansyaster	
<i>Machaeranthera tanacetifolia</i>	Tanseyleaf tansyaster	
<i>Mahonia repens</i>	Creeping barberry	
<i>Maianthemum racemosum</i>	Feathery false lily of the valley	
<i>Maianthemum racemosum</i> ssp. <i>amplexicaule</i>	Feathery false lily of the valley	
<i>Maianthemum stellatum</i>	Starry false lily of the valley	
<i>Malva neglecta</i>	Common mallow	
<i>Marsilea vestita</i>	Hairy waterclover	
<i>Medicago sativa</i>	Alfalfa	
<i>Medicago sativa</i> ssp. <i>sativa</i>	Alfalfa	
<i>Melilotus officinalis</i>	Sweetclover	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Mentha arvensis</i>	Wild mint	
<i>Mentzelia albicaulis</i>	Whitestem blazingstar	
<i>Mentzelia multiflora</i>	Adonis blazingstar	
<i>Mentzelia multiflora</i> var. <i>multiflora</i>	Adonis blazingstar	
<i>Mentzelia nuda</i>	Bractless blazingstar	
<i>Mentzelia rusbyi</i>	Rusby's blazingstar	
<i>Mentzelia speciosa</i>	Jeweled blazingstar	
<i>Menyanthes trifoliata</i>	Buckbean	
<i>Mertensia alpina</i>	Alpine bluebells	
<i>Mertensia brevistyla</i>	Shortstyle bluebells	
<i>Mertensia ciliata</i>	Tall fringed bluebells	
<i>Mertensia ciliata</i> var. <i>ciliata</i>	Tall fringed bluebells	
<i>Mertensia franciscana</i>	Franciscan bluebells	
<i>Mertensia lanceolata</i>	Prairie bluebells	
<i>Mertensia lanceolata</i> var. <i>lanceolata</i>	Prairie bluebells	
<i>Mertensia oblongifolia</i>	Oblongleaf bluebells	
<i>Mimulus floribundus</i>	Manyflowered monkeyflower	
<i>Mimulus glabratus</i>	Roundleaf monkeyflower	
<i>Mimulus guttatus</i>	Seep monkeyflower	
<i>Minuartia obtusiloba</i>	Twinflower sandwort	
<i>Minuartia rubella</i>	Beautiful sandwort	
<i>Mirabilis linearis</i>	Narrowleaf four o'clock	
<i>Mirabilis multiflora</i>	Colorado four o'clock	
<i>Mirabilis oxybaphoides</i>	Smooth spreading four o'clock	
<i>Mitella pentandra</i>	Fivestamen miterwort	
<i>Mitella stauropetala</i>	Smallflower miterwort	
<i>Mitella stauropetala</i> var. <i>stenopetala</i>	Drywoods miterwort	
<i>Moehringia lateriflora</i>	Bluntleaf sandwort	
<i>Moehringia macrophylla</i>	Largeleaf sandwort	
<i>Monarda fistulosa</i>	Wild bergamot	
<i>Monarda fistulosa</i> ssp. <i>fistulosa</i>	Wild bergamot	
<i>Monarda fistulosa</i> ssp. <i>fistulosa</i> var. <i>menthifolia</i>	Mintleaf bergamot	
<i>Monarda pectinata</i>	Pony beebalm	
<i>Moneses uniflora</i>	Single delight	
<i>Monolepis nuttalliana</i>	Nuttall's povertyweed	
<i>Monotropa hypopithys</i>	Pinesap	
<i>Montia chamissoi</i>	Water minerslettuce	
<i>Muhlenbergia andina</i>	Foxtail muhly	
<i>Muhlenbergia asperifolia</i>	Scratchgrass	
<i>Muhlenbergia brevis</i>	Short muhly	
<i>Muhlenbergia filiculmis</i>	Slimstem muhly	
<i>Muhlenbergia filiformis</i>	Pullup muhly	
<i>Muhlenbergia minutissima</i>	Annual muhly	
<i>Muhlenbergia montana</i>	Mountain muhly	
<i>Muhlenbergia pungens</i>	Sandhill muhly	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Muhlenbergia richardsonis</i>	Mat muhly	
<i>Muhlenbergia torreyi</i>	Ring muhly	
<i>Munroa squarrosa</i>	False buffalograss	
<i>Myriophyllum sibiricum</i>	Shortspike watermilfoil	
<i>Nassella viridula</i>	Green needlegrass	
<i>Nasturtium officinale</i>	Watercress	
<i>Neoparrya lithophila</i>	Bill's neoparrya	
<i>Noccaea montana</i>	Alpine pennycress	
<i>Noccaea montana</i> var. <i>montana</i>	Alpine pennycress	
<i>Nuphar lutea</i>	Yellow pond-lily	
<i>Nuphar lutea</i> ssp. <i>polysepala</i>	Rocky Mountain pond-lily	
<i>Oenothera albicaulis</i>	Whitest evening primrose	
<i>Oenothera caespitosa</i>	Tufted evening primrose	
<i>Oenothera caespitosa</i> ssp. <i>caespitosa</i>	Tufted evening primrose	
<i>Oenothera coronopifolia</i>	Crownleaf evening primrose	
<i>Oenothera elata</i>	Hooker's evening primrose	
<i>Oenothera elata</i> ssp. <i>hirsutissima</i>	Hooker's evening primrose	
<i>Oenothera flava</i>	Yellow evening primrose	
<i>Oenothera flava</i> ssp. <i>flava</i>	Yellow evening primrose	
<i>Oenothera pallida</i>	Pale evening primrose	
<i>Oenothera pallida</i> ssp. <i>runcinata</i>	Pale evening primrose	
<i>Oenothera villosa</i>	Hairy evening primrose	
<i>Oenothera villosa</i> ssp. <i>strigosa</i>	Hairy evening primrose	
<i>Opuntia polyacantha</i>	Plains pricklypear	
<i>Opuntia polyacantha</i> var. <i>polyacantha</i>	Hairspine pricklypear	
<i>Oreochrysum parryi</i>	Parry's goldenrod	
<i>Oreoxis alpina</i>	Alpine oreoxis	
<i>Oreoxis alpina</i> ssp. <i>alpina</i>	Alpine oreoxis	
<i>Oreoxis alpina</i> ssp. <i>puberulenta</i>	Alpine oreoxis	
<i>Oreoxis bakeri</i>	Baker's alpineparsley	
<i>Orobanche fasciculata</i>	Clustered broomrape	
<i>Orthilia secunda</i>	Sidebells wintergreen	
<i>Orthocarpus luteus</i>	Yellow owl's-clover	
<i>Oryzopsis asperifolia</i>	Roughleaf ricegrass	
<i>Osmorhiza depauperata</i>	Bluntseed sweetroot	
<i>Oxypolis fendleri</i>	Fendler's cowbane	
<i>Oxyria digyna</i>	Alpine mountainsorrel	
<i>Oxytropis campestris</i>	Field locoweed	
<i>Oxytropis deflexa</i>	Nodding locoweed	
<i>Oxytropis deflexa</i> var. <i>sericea</i>	Blue nodding locoweed	
<i>Oxytropis lambertii</i>	Purple locoweed	
<i>Oxytropis lambertii</i> var. <i>lambertii</i>	Purple locoweed	
<i>Oxytropis parryi</i>	Parry's oxytrope	
<i>Oxytropis sericea</i>	White locoweed	
<i>Oxytropis sericea</i> var. <i>sericea</i>	White locoweed	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Oxytropis splendens</i>	Showy locoweed	
<i>Packera cana</i>	Woolly groundsel	
<i>Packera crocata</i>	Saffron ragwort	
<i>Packera dimorphophylla</i>	Splitleaf groundsel	
<i>Packera dimorphophylla</i> var. <i>intermedia</i>	Splitleaf groundsel	
<i>Packera fendleri</i>	Fendler's ragwort	
<i>Packera neomexicana</i>	New Mexico groundsel	
<i>Packera neomexicana</i> var. <i>mutabilis</i>	New Mexico groundsel	
<i>Packera pseudaurea</i>	Falsegold groundsel	
<i>Packera pseudaurea</i> var. <i>pseudaurea</i>	Falsegold groundsel	
<i>Packera streptanthifolia</i>	Rocky Mountain groundsel	
<i>Packera tridenticulata</i>	Threetooth ragwort	
<i>Packera wernerifolia</i>	Hoary groundsel	
<i>Parietaria pensylvanica</i>	Pennsylvania pellitory	
<i>Parnassia palustris</i>	Marsh grass of Parnassus	
<i>Parnassia palustris</i> var. <i>montanensis</i>	Mountain grass of Parnassus	
<i>Paronychia pulvinata</i>	Rocky Mountain nailwort	
<i>Paronychia sessiliflora</i>	Creeping nailwort	
<i>Parthenium tetraeuris</i>	Arkansas River feverfew	
<i>Pascopyrum smithii</i>	Western wheatgrass	
<i>Pastinaca sativa</i>	Wild parsnip	
<i>Paxistima myrsinites</i>	Oregon boxleaf	
<i>Pectis angustifolia</i>	Lemonscent	
<i>Pectis angustifolia</i> var. <i>angustifolia</i>	Narrowleaf pectis	
<i>Pedicularis canadensis</i>	Canadian lousewort	
<i>Pedicularis canadensis</i> ssp. <i>fluvialilis</i>	Canadian lousewort	
<i>Pedicularis crenulata</i>	Meadow lousewort	
<i>Pedicularis groenlandica</i>	Elephanthead lousewort	
<i>Pedicularis parryi</i>	Parry's lousewort	
<i>Pedicularis parryi</i> ssp. <i>parryi</i>	Parry's lousewort	
<i>Pedicularis procera</i>	Giant lousewort	
<i>Pedicularis racemosa</i>	Sickletop lousewort	
<i>Pedicularis racemosa</i> ssp. <i>alba</i>	Sickletop lousewort	
<i>Pediocactus simpsonii</i>	Mountain ball cactus	
<i>Penstemon barbatus</i>	Beardlip penstemon	
<i>Penstemon caespitosus</i>	Mat penstemon	
<i>Penstemon griffinii</i>	Griffin's beardtongue	
<i>Penstemon hallii</i>	Hall's beardtongue	
<i>Penstemon procerus</i>	Littleflower penstemon	
<i>Penstemon procerus</i> var. <i>procerus</i>	Pincushion beardtongue	
<i>Penstemon rydbergii</i>	Rydberg's penstemon	
<i>Penstemon secundiflorus</i>	Sidebells penstemon	
<i>Penstemon strictus</i>	Rocky Mountain penstemon	
<i>Penstemon unilateralis</i>	Oneside penstemon	
<i>Penstemon whippleanus</i>	Whipple's penstemon	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Pericome caudata</i>	Mountain tail-leaf	
<i>Petasites frigidus</i>	Arctic sweet coltsfoot	
<i>Petasites frigidus</i> var. <i>sagittatus</i>	Arrowleaf sweet coltsfoot	
<i>Phacelia alba</i>	White phacelia	
<i>Phacelia bakeri</i>	Baker's phacelia	
<i>Phacelia glandulosa</i>	Glandular phacelia	
<i>Phacelia glandulosa</i> var. <i>glandulosa</i>	Glandular phacelia	
<i>Phacelia heterophylla</i>	Varileaf phacelia	
<i>Phacelia heterophylla</i> ssp. <i>heterophylla</i>	Varileaf phacelia	
<i>Phacelia sericea</i>	Silky phacelia	
<i>Phacelia sericea</i> ssp. <i>sericea</i>	Silky phacelia	
<i>Phalaris arundinacea</i>	Reed canarygrass	
<i>Phleum alpinum</i>	Alpine timothy	
<i>Phleum pratense</i>	Timothy	
<i>Phlox austromontana</i>	Mountain phlox	
<i>Phlox condensata</i>	Dwarf phlox	
<i>Phlox hoodii</i>	Spiny phlox	
<i>Phlox pulvinata</i>	Cushion phlox	
<i>Physaria floribunda</i>	Oointtip twinpod	
<i>Physocarpus monogynus</i>	Mountain ninebark	
<i>Picea engelmannii</i>	Engelmann spruce	
<i>Picea engelmannii</i> var. <i>engelmannii</i>	Engelmann spruce	
<i>Picea pungens</i>	Blue spruce	
<i>Picradeniopsis oppositifolia</i>	Oppositeleaf bahia	
<i>Pinus aristata</i>	Bristlecone pine	
<i>Pinus edulis</i>	Twoneedle pinyon	
<i>Pinus flexilis</i>	Limber pine	
<i>Pinus ponderosa</i>	Ponderosa pine	
<i>Pinus ponderosa</i> var. <i>brachyptera</i>	Ponderosa pine	
<i>Pinus ponderosa</i> var. <i>scopulorum</i>	Ponderosa pine	
<i>Pinus strobiformis</i>	Southwestern white pine	
<i>Piptatherum micranthum</i>	Littleseed ricegrass	
<i>Piptatherum pungens</i>	Mountain ricegrass	
<i>Plagiobothrys scouleri</i>	Scouler's popcornflower	
<i>Plagiobothrys scouleri</i> var. <i>hispidulus</i>	Sleeping popcornflower	
<i>Plantago eriopoda</i>	Redwool plantain	
<i>Plantago major</i>	Common plantain	
<i>Platanthera dilatata</i>	Scentbottle	
<i>Platanthera dilatata</i> var. <i>albiflora</i>	Scentbottle	
<i>Platanthera obtusata</i>	Bluntleaved orchid	
<i>Platanthera obtusata</i> ssp. <i>obtusata</i>	Bluntleaved orchid	
<i>Platanthera sparsiflora</i>	Sparse-flowered bog orchid	
<i>Platanthera sparsiflora</i> var. <i>ensifolia</i>	Sparse-flowered bog orchid	
<i>Poa alpina</i>	Alpine bluegrass	
<i>Poa annua</i>	Annual bluegrass	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Poa arctica</i>	Arctic bluegrass	
<i>Poa arctica</i> ssp. <i>aperta</i>	Arctic bluegrass	
<i>Poa compressa</i>	Canada bluegrass	
<i>Poa fendleriana</i>	Muttongrass	
<i>Poa glauca</i>	Glaucous bluegrass	
<i>Poa glauca</i> ssp. <i>rupicola</i>	Timberline bluegrass	
<i>Poa leptocoma</i>	Marsh bluegrass	
<i>Poa lettermanii</i>	Letterman's bluegrass	
<i>Poa nemoralis</i>	Wood bluegrass	
<i>Poa nemoralis</i> ssp. <i>interior</i>	Inland bluegrass	
<i>Poa palustris</i>	Fowl bluegrass	
<i>Poa pratensis</i>	Kentucky bluegrass	
<i>Poa pratensis</i> ssp. <i>pratensis</i>	Kentucky bluegrass	
<i>Poa reflexa</i>	Nodding bluegrass	
<i>Poa secunda</i>	Sandberg bluegrass	
<i>Podistera eastwoodiae</i>	Eastwood's podistera	
<i>Polemonium brandegeei</i>	Brandegee's Jacob's-ladder	
<i>Polemonium confertum</i>	Rocky Mountain Jacob's-ladder	
<i>Polemonium foliosissimum</i>	Towering Jacob's-ladder	
<i>Polemonium occidentale</i>	Western polemonium	
<i>Polemonium occidentale</i> ssp. <i>occidentale</i>	Western polemonium	
<i>Polemonium pulcherrimum</i>	Jacob's-ladder	
<i>Polemonium pulcherrimum</i> ssp. <i>delicatum</i>	Jacob's-ladder	
<i>Polemonium viscosum</i>	Sticky polemonium	
<i>Polygonum amphibium</i>	Water knotweed	
<i>Polygonum amphibium</i> var. <i>emersum</i>	Longroot smartweed	
<i>Polygonum arenastrum</i>	Oval-leaf knotweed	
<i>Polygonum argyrocoleon</i>	Silversheath knotweed	
<i>Polygonum bistortoides</i>	American bistort	
<i>Polygonum douglasii</i>	Douglas' knotweed	
<i>Polygonum pensylvanicum</i>	Pennsylvania smartweed	
<i>Polygonum persicaria</i>	Spotted ladythumb	
<i>Polygonum viviparum</i>	Alpine bistort	
<i>Populus</i> × <i>acuminata</i>	Lanceleaf cottonwood	
<i>Populus angustifolia</i>	Narrowleaf cottonwood	
<i>Populus tremuloides</i>	Quaking aspen	
<i>Portulaca oleracea</i>	Little hogweed	
<i>Potamogeton alpinus</i>	Alpine pondweed	
<i>Potamogeton foliosus</i>	Leafy pondweed	
<i>Potamogeton foliosus</i> ssp. <i>foliosus</i>	Leafy pondweed	
<i>Potamogeton nodosus</i>	Longleaf pondweed	
<i>Potamogeton pusillus</i>	Small pondweed	
<i>Potamogeton pusillus</i> ssp. <i>pusillus</i>	Small pondweed	
<i>Potentilla ambigens</i>	Silkleaf cinquefoil	
<i>Potentilla concinna</i>	Elegant cinquefoil	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Potentilla concinna</i> var. <i>concinna</i>	Elegant cinquefoil	
<i>Potentilla diversifolia</i>	Varileaf cinquefoil	
<i>Potentilla diversifolia</i> var. <i>diversifolia</i>	Varileaf cinquefoil	
<i>Potentilla gracilis</i>	Slender cinquefoil	
<i>Potentilla hippiana</i>	Woolly cinquefoil	
<i>Potentilla hippiana</i> var. <i>hippiana</i>	Woolly cinquefoil	
<i>Potentilla norvegica</i>	Norwegian cinquefoil	
<i>Potentilla norvegica</i> ssp. <i>monspeliensis</i>	Norwegian cinquefoil	
<i>Potentilla paradoxa</i>	Paradox cinquefoil	
<i>Potentilla pensylvanica</i>	Pennsylvania cinquefoil	
<i>Potentilla pensylvanica</i> var. <i>pensylvanica</i>	Pennsylvania cinquefoil	
<i>Potentilla plattensis</i>	Platte River cinquefoil	
<i>Potentilla pulcherrima</i>	Beautiful cinquefoil	
<i>Potentilla rivalis</i>	Brook cinquefoil	
<i>Potentilla subjuga</i>	Colorado cinquefoil	
<i>Potentilla uniflora</i>	Oneflower cinquefoil	
<i>Primula angustifolia</i>	Alpine primrose	
<i>Primula parryi</i>	Parry's primrose	
<i>Prunella vulgaris</i>	Common selfheal	
<i>Prunella vulgaris</i> ssp. <i>lanceolata</i>	Lance selfheal	
<i>Prunus pensylvanica</i>	Pin cherry	
<i>Prunus pensylvanica</i> var. <i>pensylvanica</i>	Pin cherry	
<i>Prunus virginiana</i>	Chokecherry	
<i>Prunus virginiana</i> var. <i>melanocarpa</i>	Black chokecherry	
<i>Psathyrostachys juncea</i>	Russian wildrye	
<i>Pseudocymopterus montanus</i>	Alpine false springparsley	
<i>Pseudotsuga menziesii</i>	Douglas-fir	
<i>Pseudotsuga menziesii</i> var. <i>glauca</i>	Rocky Mountain Douglas-fir	
<i>Psoralidium lanceolatum</i>	Lemon scurfpea	
<i>Pteridium aquilinum</i>	Western brackenfern	
<i>Pteridium aquilinum</i> var. <i>pubescens</i>	Hairy brackenfern	
<i>Pterospora andromedea</i>	Woodland pinedrops	
<i>Pteryxia hendersonii</i>	Henderson's wavewing	
<i>Puccinellia nuttalliana</i>	Nuttall's alkaligrass	
<i>Pulsatilla patens</i>	Eastern pasqueflower	
<i>Pulsatilla patens</i> ssp. <i>multifida</i>	Cutleaf anemone	
<i>Pyrola asarifolia</i>	Liverleaf wintergreen	
<i>Pyrola asarifolia</i> ssp. <i>asarifolia</i>	Liverleaf wintergreen	
<i>Pyrola chlorantha</i>	Greenflowered wintergreen	
<i>Pyrola minor</i>	Snowline wintergreen	
<i>Pyrrcoma clementis</i>	Tranquil goldenweed	
<i>Pyrrcoma clementis</i> var. <i>clementis</i>	Tranquil goldenweed	
<i>Pyrrcoma lanceolata</i>	Lanceleaf goldenweed	
<i>Pyrrcoma lanceolata</i> var. <i>lanceolata</i>	Lanceleaf goldenweed	
<i>Pyrrcoma uniflora</i>	Plantain goldenweed	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Pyrrcoma uniflora</i> var. <i>uniflora</i>	Plantain goldenweed	
<i>Quercus gambelii</i>	Gambel oak	
<i>Quercus gambelii</i> var. <i>gambelii</i>	Gambel oak	
<i>Ranunculus abortivus</i>	Littleleaf buttercup	
<i>Ranunculus alismifolius</i>	Plantainleaf buttercup	
<i>Ranunculus alismifolius</i> var. <i>montanus</i>	Waterplantain buttercup	
<i>Ranunculus cardiophyllus</i>	Heartleaf buttercup	
<i>Ranunculus cymbalaria</i>	Alkali buttercup	
<i>Ranunculus gmelinii</i>	Gmelin's buttercup	
<i>Ranunculus hyperboreus</i>	High northern buttercup	
<i>Ranunculus inamoenus</i>	Graceful buttercup	
<i>Ranunculus macauleyi</i>	Rocky Mountain buttercup	
<i>Ranunculus macounii</i>	Macoun's buttercup	
<i>Ranunculus sceleratus</i>	Cursed buttercup	
<i>Ranunculus sceleratus</i> var. <i>multifidus</i>	Cursed buttercup	
<i>Ranunculus sceleratus</i> var. <i>sceleratus</i>	Cursed buttercup	
<i>Ranunculus trichophyllus</i>	Threadleaf crowfoot	
<i>Ranunculus trichophyllus</i> var. <i>trichophyllus</i>	Threadleaf crowfoot	
<i>Ranunculus uncinatus</i>	Woodland buttercup	
<i>Redfieldia flexuosa</i>	Blowout grass	
<i>Rhinanthus minor</i>	Little yellow rattle	
<i>Rhinanthus minor</i> ssp. <i>minor</i>	Little yellow rattle	
<i>Rhodiola integrifolia</i>	Ledge stonecrop	
<i>Rhodiola rhodantha</i>	Redpod stonecrop	
<i>Rhus trilobata</i>	Skunkbush sumac	
<i>Rhus trilobata</i> var. <i>trilobata</i>	Skunkbush sumac	
<i>Ribes aureum</i>	Golden currant	
<i>Ribes cereum</i>	Wax currant	
<i>Ribes cereum</i> var. <i>pedicellare</i>	Whisky currant	
<i>Ribes inerme</i>	Whitestem gooseberry	
<i>Ribes inerme</i> var. <i>inerme</i>	Whitestem gooseberry	
<i>Ribes laxiflorum</i>	Trailing black currant	
<i>Ribes leptanthum</i>	Trumpet gooseberry	
<i>Ribes montigenum</i>	Gooseberry currant	
<i>Ribes wolfii</i>	Wolf's currant	
<i>Rorippa alpina</i>	Alpine yellowcress	
<i>Rorippa curvipes</i>	Bluntleaf yellowcress	
<i>Rorippa curvipes</i> var. <i>curvipes</i>	Bluntleaf yellowcress	
<i>Rorippa curvipes</i> var. <i>truncata</i>	Bluntleaf yellowcress	
<i>Rorippa palustris</i>	Bog yellowcress	
<i>Rorippa palustris</i> ssp. <i>hispida</i>	Hispid yellowcress	
<i>Rorippa sinuata</i>	Spreading yellowcress	
<i>Rorippa sphaerocarpa</i>	Roundfruit yellowcress	
<i>Rosa acicularis</i>	Prickly rose	
<i>Rosa acicularis</i> ssp. <i>sayi</i>	Prickly rose	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Rosa woodsii</i>	Woods' rose	
<i>Rosa woodsii</i> var. <i>ultramontana</i>	Woods' rose	
<i>Rubus deliciosus</i>	Delicious raspberry	
<i>Rubus idaeus</i>	American red raspberry	
<i>Rubus idaeus</i> ssp. <i>strigosus</i>	Grayleaf red raspberry	
<i>Rubus parviflorus</i>	Thimbleberry	
<i>Rubus parviflorus</i> var. <i>parviflorus</i>	Thimbleberry	
<i>Rudbeckia hirta</i>	Blackeyed Susan	
<i>Rudbeckia hirta</i> var. <i>pulcherrima</i>	Blackeyed Susan	
<i>Rudbeckia laciniata</i>	Cutleaf coneflower	
<i>Rudbeckia laciniata</i> var. <i>ampla</i>	Cutleaf coneflower	
<i>Rumex aquaticus</i>	Western dock	
<i>Rumex aquaticus</i> var. <i>fenestratus</i>	Western dock	
<i>Rumex densiflorus</i>	Denseflowered dock	
<i>Rumex maritimus</i>	Golden dock	
<i>Rumex salicifolius</i>	Willow dock	
<i>Rumex salicifolius</i> var. <i>mexicanus</i>	Mexican dock	
<i>Rumex venosus</i>	Veiny dock	
<i>Sagina saginoides</i>	Arctic pearlwort	
<i>Sagittaria cuneata</i>	Arumleaf arrowhead	
<i>Salix amygdaloides</i>	Peachleaf willow	
<i>Salix bebbiana</i>	Bebb willow	
<i>Salix brachycarpa</i>	Shortfruit willow	
<i>Salix brachycarpa</i> var. <i>brachycarpa</i>	Shortfruit willow	
<i>Salix drummondiana</i>	Drummond's willow	
<i>Salix exigua</i>	Narrowleaf willow	
<i>Salix geyeriana</i>	Geyer willow	
<i>Salix ligulifolia</i>	Strapleaf willow	
<i>Salix lucida</i>	Shining willow	
<i>Salix lucida</i> ssp. <i>caudata</i>	Greenleaf willow	
<i>Salix monticola</i>	Park willow	
<i>Salix nivalis</i>	Snow willow	
<i>Salix orestera</i>	Sierra willow	
<i>Salix petrophila</i>	Alpine willow	
<i>Salix planifolia</i>	Diamondleaf willow	
<i>Salix planifolia</i> ssp. <i>planifolia</i>	Diamondleaf willow	
<i>Salix scouleriana</i>	Scouler's willow	
<i>Salix wolfii</i>	Wolf's willow	
<i>Salsola tragus</i>	Prickly Russian thistle	
<i>Salvia reflexa</i>	Lanceleaf sage	
<i>Sambucus racemosa</i>	Red elderberry	
<i>Sambucus racemosa</i> var. <i>racemosa</i>	Red elderberry	
<i>Sarcobatus vermiculatus</i>	Greasewood	
<i>Saxifraga bronchialis</i>	Yellowdot saxifrage	
<i>Saxifraga bronchialis</i> ssp. <i>austromontana</i>	Matted saxifrage	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Saxifraga caespitosa</i>	Tufted alpine saxifrage	
<i>Saxifraga caespitosa</i> ssp. <i>delicatula</i>	Tufted alpine saxifrage	
<i>Saxifraga cernua</i>	Nodding saxifrage	
<i>Saxifraga chrysantha</i>	Goldbloom saxifrage	
<i>Saxifraga flagellaris</i>	Whiplash saxifrage	
<i>Saxifraga flagellaris</i> ssp. <i>crandallii</i>	Crandall's saxifrage	
<i>Saxifraga odontoloma</i>	Brook saxifrage	
<i>Saxifraga rhomboidea</i>	Diamondleaf saxifrage	
<i>Saxifraga rivularis</i>	Weak saxifrage	
<i>Schedonnardus paniculatus</i>	Tumblegrass	
<i>Schizachyrium scoparium</i>	Little bluestem	
<i>Schizachyrium scoparium</i> var. <i>scoparium</i>	Little bluestem	
<i>Schkuhria multiflora</i>	Manyflower false threadleaf	
<i>Schoenocrambe linearifolia</i>	Slimleaf plainsmustard	
<i>Schoenoplectus acutus</i>	Hardstem bulrush	
<i>Schoenoplectus acutus</i> var. <i>acutus</i>	Hardstem bulrush	
<i>Schoenoplectus maritimus</i>	Cosmopolitan bulrush	
<i>Schoenoplectus pungens</i>	Common threesquare	
<i>Schoenoplectus pungens</i> var. <i>longispicatus</i>	Common threesquare	
<i>Schoenoplectus tabernaemontani</i>	Softstem bulrush	
<i>Scirpus microcarpus</i>	Panicled bulrush	
<i>Scirpus nevadensis</i>	Nevada bulrush	
<i>Scrophularia lanceolata</i>	Lanceleaf figwort	
<i>Scutellaria galericulata</i>	Marsh skullcap	
<i>Sedum lanceolatum</i>	Spearleaf stonecrop	
<i>Sedum lanceolatum</i> ssp. <i>lanceolatum</i>	Spearleaf stonecrop	
<i>Selaginella densa</i>	Lesser spikemoss	
<i>Selaginella weatherbiana</i>	Weatherby's spikemoss	
<i>Senecio amplexens</i>	Showy alpine ragwort	
<i>Senecio amplexens</i> var. <i>amplexens</i>	Showy alpine ragwort	
<i>Senecio amplexens</i> var. <i>holmii</i>	Holm's ragwort	
<i>Senecio atratus</i>	Tall blacktip ragwort	
<i>Senecio bigelovii</i>	Nodding ragwort	
<i>Senecio bigelovii</i> var. <i>hallii</i>	Hall's ragwort	
<i>Senecio crassulus</i>	Thickleaf ragwort	
<i>Senecio eremophilus</i>	Desert ragwort	
<i>Senecio eremophilus</i> var. <i>kingii</i>	King's ragwort	
<i>Senecio fremontii</i>	Dwarf mountain ragwort	
<i>Senecio fremontii</i> var. <i>blitoides</i>	Dwarf mountain ragwort	
<i>Senecio pudicus</i>	Bashful ragwort	
<i>Senecio soldanella</i>	Colorado ragwort	
<i>Senecio spartioides</i>	Broom-like ragwort	
<i>Senecio spartioides</i> var. <i>multicapitatus</i>	Broom-like ragwort	
<i>Senecio taraxacoides</i>	Dandelion ragwort	
<i>Senecio triangularis</i>	Arrowleaf ragwort	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Senecio wootonii</i>	Wooton's ragwort	
<i>Sesuvium verrucosum</i>	Verrucose seapurslane	
<i>Setaria viridis</i>	Green bristlegrass	
<i>Setaria viridis</i> var. <i>viridis</i>	Green bristlegrass	
<i>Shepherdia canadensis</i>	Russet buffaloberry	
<i>Sibbaldia procumbens</i>	Creeping sibbaldia	
<i>Sidalcea candida</i>	White checkerbloom	
<i>Sidalcea neomexicana</i>	Salt spring checkerbloom	
<i>Sidalcea neomexicana</i> ssp. <i>neomexicana</i>	Salt spring checkerbloom	
<i>Silene acaulis</i>	Moss campion	
<i>Silene acaulis</i> var. <i>subacaulescens</i>	Moss campion	
<i>Silene drummondii</i>	Drummond's campion	
<i>Silene drummondii</i> var. <i>drummondii</i>	Drummond's campion	
<i>Silene menziesii</i>	Menzies' campion	
<i>Silene menziesii</i> ssp. <i>menziesii</i>	Menzies' campion	
<i>Silene menziesii</i> ssp. <i>menziesii</i> var. <i>menziesii</i>	Menzies' campion	
<i>Silene scouleri</i>	Simple campion	
<i>Silene scouleri</i> ssp. <i>hallii</i>	Simple campion	
<i>Sisymbrium altissimum</i>	Tall tumbled mustard	
<i>Sisyrinchium demissum</i>	Stiff blue-eyed grass	
<i>Sisyrinchium montanum</i>	Strict blue-eyed grass	
<i>Sisyrinchium montanum</i> var. <i>montanum</i>	Strict blue-eyed grass	
<i>Sisyrinchium pallidum</i>	Pale blue-eyed grass	
<i>Sium suave</i>	Hemlock waterparsnip	
<i>Smelowskia calycina</i>	Alpine smelowskia	
<i>Smelowskia calycina</i> var. <i>americana</i>	American false candytuft	
<i>Solanum triflorum</i>	Cutleaf nightshade	
<i>Solidago canadensis</i>	Canada goldenrod	
<i>Solidago missouriensis</i>	Missouri goldenrod	
<i>Solidago multiradiata</i>	Rocky Mountain goldenrod	
<i>Solidago multiradiata</i> var. <i>scopulorum</i>	Manyray goldenrod	
<i>Solidago simplex</i>	Mt. Albert goldenrod	
<i>Solidago simplex</i> ssp. <i>simplex</i>	Mt. Albert goldenrod	
<i>Solidago simplex</i> ssp. <i>simplex</i> var. <i>simplex</i>	Mt. Albert goldenrod	
<i>Solidago velutina</i>	Threenerve goldenrod	
<i>Sonchus arvensis</i>	Field sowthistle	
<i>Sonchus arvensis</i> ssp. <i>uliginosus</i>	Moist sowthistle	
<i>Sophora nuttalliana</i>	Silky sophora	
<i>Spartina gracilis</i>	Alkali cordgrass	
<i>Sphaeralcea coccinea</i>	Scarlet globemallow	
<i>Sphaeralcea coccinea</i> ssp. <i>coccinea</i>	Scarlet globemallow	
<i>Sphaerophysa salsula</i>	Alkali swainsonpea	
<i>Sphenopholis obtusata</i>	Prairie wedgescale	
<i>Spiranthes romanzoffiana</i>	Hooded lady's tresses	
<i>Sporobolus airoides</i>	Alkali sacaton	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Sporobolus contractus</i>	Spike dropseed	
<i>Sporobolus cryptandrus</i>	Sand dropseed	
<i>Stachys pilosa</i>	Hairy hedgenettle	
<i>Stachys pilosa</i> var. <i>pilosa</i>	Hairy hedgenettle	
<i>Stellaria calycantha</i>	Northern starwort	
<i>Stellaria crassifolia</i>	Fleshy starwort	
<i>Stellaria crassifolia</i> var. <i>crassifolia</i>	Fleshy starwort	
<i>Stellaria longifolia</i>	Longleaf starwort	
<i>Stellaria longifolia</i> var. <i>longifolia</i>	Longleaf starwort	
<i>Stellaria longipes</i>	Longstalk starwort	
<i>Stellaria longipes</i> ssp. <i>longipes</i>	Chickweed, starwort	
<i>Stellaria umbellata</i>	Umbrella starwort	
<i>Stephanomeria pauciflora</i>	Brownplume wirelettuce	
<i>Streptopus amplexifolius</i>	Claspleaf twistedstalk	
<i>Streptopus amplexifolius</i> var. <i>chalazatus</i>	Tubercle twistedstalk	
<i>Suaeda calceoliformis</i>	Pursh seepweed	
<i>Suaeda moquinii</i>	Mojave seablite	
<i>Swertia perennis</i>	Felwort	
<i>Symphoricarpos occidentalis</i>	Western snowberry	
<i>Symphoricarpos rotundifolius</i>	Roundleaf snowberry	
<i>Symphoricarpos rotundifolius</i> var. <i>rotundifolius</i>	Roundleaf snowberry	
<i>Symphyotrichum ascendens</i>	Western aster	
<i>Symphyotrichum boreale</i>	Northern bog aster	
<i>Symphyotrichum eatonii</i>	Eaton's aster	
<i>Symphyotrichum ericoides</i>	White heath aster	
<i>Symphyotrichum ericoides</i> var. <i>ericoides</i>	White heath aster	
<i>Symphyotrichum falcatum</i>	White prairie aster	
<i>Symphyotrichum falcatum</i> var. <i>falcatum</i>	White prairie aster	
<i>Symphyotrichum foliaceum</i>	Alpine leafybract aster	
<i>Symphyotrichum frondosum</i>	Short-rayed alkali aster	
<i>Symphyotrichum lanceolatum</i>	White panicle aster	
<i>Symphyotrichum lanceolatum</i> ssp. <i>hesperium</i>	White panicle aster	
<i>Symphyotrichum lanceolatum</i> ssp. <i>hesperium</i> var. <i>hesperium</i>	White panicle aster	
<i>Symphyotrichum spathulatum</i>	Western mountain aster	
<i>Symphyotrichum spathulatum</i> var. <i>spathulatum</i>	Western mountain aster	
<i>Taraxacum lyratum</i>	Harp dandelion	
<i>Taraxacum officinale</i>	Common dandelion	
<i>Taraxacum officinale</i> ssp. <i>ceratophorum</i>	Common dandelion	
<i>Tetradymia canescens</i>	Spineless horsebrush	
<i>Tetrameuris acaulis</i>	Stemless four-nerve daisy	
<i>Tetrameuris acaulis</i> var. <i>acaulis</i>	Stemless four-nerve daisy	
<i>Tetrameuris acaulis</i> var. <i>caespitosa</i>	Caespitose four-nerve daisy	
<i>Tetrameuris brandegeei</i>	Brandegee's four-nerve daisy	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Tetraneuris grandiflora</i>	Graylocks four-nerve daisy	
<i>Teucrium canadense</i>	Canada germander	
<i>Teucrium canadense</i> var. <i>occidentale</i>	Western germander	
<i>Thalictrum alpinum</i>	Alpine meadow-rue	
<i>Thalictrum fendleri</i>	Fendler's meadow-rue	
<i>Thalictrum fendleri</i> var. <i>fendleri</i>	Fendler's meadow-rue	
<i>Thalictrum sparsiflorum</i>	Fewflower meadow-rue	
<i>Thalictrum sparsiflorum</i> var. <i>saximontanum</i>	Fewflower meadow-rue	
<i>Thelesperma filifolium</i>	Stiff greenthread	
<i>Thelesperma filifolium</i> var. <i>intermedium</i>	Stiff greenthread	
<i>Thelesperma subnudum</i>	Navajo tea	
<i>Thelesperma subnudum</i> var. <i>subnudum</i>	Navajo tea	
<i>Thermopsis divaricarpa</i>	Spreadfruit goldenbanner	
<i>Thermopsis montana</i>	Mountain goldenbanner	
<i>Thermopsis montana</i> var. <i>montana</i>	Mountain goldenbanner	
<i>Thermopsis rhombifolia</i>	Prairie thermopsis	
<i>Thlaspi arvense</i>	Field pennycress	
<i>Tonestus pygmaeus</i>	Pygmy goldenweed	
<i>Townsendia eximia</i>	Tall Townsend daisy	
<i>Townsendia exscapa</i>	Stemless Townsend daisy	
<i>Townsendia grandiflora</i>	Largeflower Townsend daisy	
<i>Townsendia hookeri</i>	Hooker's Townsend daisy	
<i>Townsendia leptotes</i>	Common Townsend daisy	
<i>Tragopogon porrifolius</i>	Salsify	
<i>Trautvetteria caroliniensis</i>	Carolina bugbane	
<i>Trautvetteria caroliniensis</i> var. <i>occidentalis</i>	Western bugbane	
<i>Trifolium attenuatum</i>	Rocky Mountain clover	
<i>Trifolium brandegeei</i>	Brandegee's clover	
<i>Trifolium dasyphyllum</i>	Alpine clover	
<i>Trifolium dasyphyllum</i> ssp. <i>dasyphyllum</i>	Alpine clover	
<i>Trifolium hybridum</i>	Alsike clover	
<i>Trifolium longipes</i>	Longstalk clover	
<i>Trifolium longipes</i> ssp. <i>pygmaeum</i>	Pygmy clover	
<i>Trifolium nanum</i>	Dwarf clover	
<i>Trifolium parryi</i>	Parry's clover	
<i>Trifolium parryi</i> ssp. <i>salictorum</i>	Parry's clover	
<i>Trifolium repens</i>	White clover	
<i>Trifolium wormskioldii</i>	Cows clover	
<i>Triglochin maritima</i>	Seaside arrowgrass	
<i>Triglochin palustris</i>	Marsh arrowgrass	
<i>Tripterocalyx micranthus</i>	Smallflower sandverbena	
<i>Trisetum spicatum</i>	Spike trisetum	
<i>Trollius laxus</i>	American globeflower	
<i>Trollius laxus</i> ssp. <i>albiflorus</i>	American globeflower	
<i>Typha latifolia</i>	Broadleaf cattail	

<i>Scientific name</i>	<i>Common name</i>	<i>Status</i>
<i>Urtica dioica</i>	Stinging nettle	
<i>Urtica dioica</i> ssp. <i>gracilis</i>	California nettle	
<i>Utricularia ochroleuca</i>	Yellowishwhite bladderwort	
<i>Vaccinium cespitosum</i>	Dwarf bilberry	
<i>Vaccinium myrtillos</i>	Whortleberry	
<i>Vaccinium scoparium</i>	Grouse whortleberry	
<i>Valeriana acutiloba</i>	Sharpleaf valerian	
<i>Valeriana acutiloba</i> var. <i>acutiloba</i>	Sharpleaf valerian	
<i>Valeriana arizonica</i>	Arizona valerian	
<i>Valeriana edulis</i>	Tobacco root	
<i>Valeriana edulis</i> var. <i>edulis</i>	Tobacco root	
<i>Veratrum tenuipetalum</i>	Colorado false hellebore	
<i>Verbena bracteata</i>	Bigbract verbena	
<i>Verbena macdougalii</i>	MacDougal verbena	
<i>Verbesina encelioides</i>	Golden crownbeard	
<i>Verbesina encelioides</i> ssp. <i>encelioides</i>	Golden crownbeard	
<i>Verbesina encelioides</i> ssp. <i>exauriculata</i>	Golden crownbeard	
<i>Veronica americana</i>	American speedwell	
<i>Veronica peregrina</i>	Neckweed	
<i>Veronica peregrina</i> ssp. <i>xalapensis</i>	Hairy purslane speedwell	
<i>Veronica serpyllifolia</i>	Thymeleaf speedwell	
<i>Veronica serpyllifolia</i> ssp. <i>humifusa</i>	Brightblue speedwell	
<i>Veronica wormskjoldii</i>	American alpine speedwell	
<i>Veronica wormskjoldii</i> var. <i>wormskjoldii</i>	American alpine speedwell	
<i>Vicia americana</i>	American vetch	
<i>Vicia americana</i> ssp. <i>americana</i>	American vetch	
<i>Vicia sativa</i>	Garden vetch	
<i>Vicia sativa</i> ssp. <i>nigra</i>	Garden vetch	
<i>Viola adunca</i>	Hookedspur violet	
<i>Viola adunca</i> var. <i>adunca</i>	Hookedspur violet	
<i>Viola biflora</i>	Arctic yellow violet	
<i>Viola biflora</i> ssp. <i>biflora</i>	Arctic yellow violet	
<i>Viola canadensis</i>	Canadian white violet	
<i>Viola canadensis</i> var. <i>scopulorum</i>	Canadian white violet	
<i>Viola labradorica</i>	Alpine violet	
<i>Viola macloskeyi</i>	Small white violet	
<i>Viola macloskeyi</i> ssp. <i>pallens</i>	Smooth white violet	
<i>Viola nephrophylla</i>	Northern bog violet	
<i>Viola renifolia</i>	White violet	
<i>Woodsia oregana</i>	Oregon cliff fern	
<i>Woodsia oregana</i> ssp. <i>cathcartiana</i>	Oregon cliff fern	
<i>Woodsia scopulina</i>	Rocky Mountain woodsia	
<i>Yucca glauca</i>	Soapweed yucca	
<i>Zigadenus elegans</i>	Mountain deathcamas	
<i>Zigadenus elegans</i> ssp. <i>elegans</i>	Mountain deathcamas	

Appendix E

Section 7 Biological Evaluation

Section 7 of the Endangered Species Act requires that Federal agencies, including the Service, consider the effects of their actions on threatened and endangered species. Our Colorado Ecological Services Field Office is responsible for section 7 consultation for the SLVCA project area.

We conducted informal intra-Service consultation during development of the draft EA and LPP in 2012, and the Colorado Ecological Services Field Office concurred with our determination that authorization of the SLVCA “may affect, but is not likely to adversely affect” Endangered Species Act-protected species. Because of changes to the status of candidate species that were the subject of that consultation, in November 2013 we sought and received concurrence that the project still meets the same determination.



United States Department of the Interior

FISH AND WILDLIFE SERVICE
Mountain-Prairie Region



IN REPLY REFER TO:
FWS/R6/NWRS
Mail Stop 60130

MAILING ADDRESS:
Post Office Box 25486
Denver Federal Center
Denver, Colorado 80225-0486

STREET LOCATION:
134 Union Boulevard
Lakewood, Colorado 80228-1807

Memorandum

To: Field Supervisor, Colorado Ecological Services Field Office

From: *Acting* Assistant Regional Director, National Wildlife Refuge System, Region 6 *Rankin*

Subject: Request for Reaffirmation of Concurrence on the San Luis Valley
Conservation Area

On June 28, 2012, the Colorado Ecological Services Field Office concurred with the San Luis Valley National Wildlife Refuge Complex Project Leader's determination that the establishment of the San Luis Valley Conservation Area (SLVCA) "may affect, but is not likely to adversely affect" Endangered Species Act (ESA) protected species and their critical habitats (Attachment 1). As detailed in the informal consultation form (TAILS ID 2012-I-0396), the SLVCA is a voluntary conservation easement program with limited fee-title land acquisition in southern Colorado which is focused on protecting key habitats, including habitat for species which were the subject of the consultation. There have been minor changes since the initial concurrence was signed.

First, the project area boundary has been split into two (Attachment 2). The Sangre de Cristo Conservation Area was established in September 2012, and we are now proceeding with the authorization of the SLVCA in the remaining original project area. There is no change in the overall boundary encompassed by the two conservation areas, and the overall acreage target is the same as in the initial consultation.

Second, the status of four of the species or their critical habitats have changed:

- Southwestern willow flycatcher – Revised critical habitat designated
- Yellow-billed cuckoo (western DPS) – Was Candidate, now Proposed Threatened
- Gunnison sage-grouse – Was Candidate, now Proposed Endangered
- New Mexico meadow jumping mouse – Was Candidate, now Proposed Endangered

The potential impacts (all beneficial) of the proposed SLVCA on each of these species was discussed in the initial consultation.

2

We do not believe that the separation of the original project into two units, nor the changes in the ESA status of the species, alters our determination that of "may affect, but not likely to adversely affect." We request that the Colorado Ecological Services Field Office concur with this determination.

Concur Do Not Concur


Colorado Ecological Services Field Supervisor

11/27/2013
Date

Intra-Service Section 7 Biological Evaluation Form - Region 6

Originating Person: Mike Blenden

Date Submitted: 19March2012

Telephone Number: 303-289-0350

I. **Service Program and Geographic Area or Station Name:**

San Luis Valley NWR Complex

II. **Flexible Funding Program** (e.g. Joint Venture, etc) if applicable:

N/A

III. **Location:** Location of the project including County, State and TSR (township, section & range):

Saguache, Alamosa, Rio Grande, Mineral, Hinsdale, Conejos, and Costilla Counties, CO; and Rio Arriba and Taos Counties, NM

The San Luis Valley Conservation Area (SLVCA) is a new, proposed unit of the National Wildlife Refuge System which encompasses the headwaters HUC6 of the Rio Grande River in south-central Colorado and northern New Mexico. It is comprised of the San Luis Valley and the slopes of the Sangre de Cristo and San Juan Mountains which border the valley. The elevation of the SLVCA extends from approximately 7500' on the valley floor to 14,345' at Blanca Peak. Please see attached map.

IV. **Species/Critical Habitat:** List federally endangered, threatened, proposed, and candidate species or designated or proposed critical habitat that may occur within the action area.

Mexican spotted owl (*Strix occidentalis lucida*) (Threatened)

Southwestern willow flycatcher (*Empidonax traillii extimus*) (Endangered; proposed critical habitat present)

Yellow-billed cuckoo (*Coccyzus americanus*) (Candidate)

Gunnison sage-grouse (*Centrocercus minimus*) (Candidate)

Rio Grande cutthroat trout (*Oncorhynchus clarki virginalis*) (Candidate)

Canada lynx (*Lynx canadensis*) (Threatened)

Gunnison's prairie dog (*Cynomys gunnisoni*) (Candidate)

New Mexico meadow jumping mouse (*Zapus hudsonius luteus*) (Candidate; presence in project area uncertain)

V. **Project Description:** Describe proposed project or action or, if referencing other documents, prepare an executive summary (attach additional pages as needed):

The Service proposes to establish the SLVCA, a new, 5.2 million acre unit of the San Luis Valley NWR Complex. Within the proposed acquisition boundary of the SLVCA, the Service will seek to acquire an interest in up to 430,000 acres. Acquisition will primarily be in the form of conservation easements which will restrict subdivision, development, and sale of surface water rights; however, the acquisition of up to 30,000 acres in fee-title was approved in the preliminary project proposal, and this tool would be used judiciously as appropriate. Any lands acquired in fee would be managed in accordance with the Comprehensive Conservation Plan for the existing 3 refuges in the complex.

The San Luis Valley is a high desert basin which forms the headwaters of the Rio Grande River. Much of the valley is irrigated for agriculture. Most of the natural vegetation on the valley floor is semi-desert shrubland. However, the precipitation captured in the surrounding valley contributes to a network of surface and groundwater-driven wetlands and riparian corridors which provide breeding and migration stopover habitat for dozens of species of migratory birds. At higher elevations, the habitat transitions to sagebrush scrub, pinon juniper woodland, montane forests, and finally alpine vegetation.



Acquisition within the SLVCA boundary has been prioritized based upon the needs of 8 focal species: southwestern willow flycatcher, Lewis' woodpecker, Rio Grande cutthroat trout, sage thrasher, Gunnison sage grouse, Canada lynx, Wilson's phalarope, and American bittern.



VI. Determination of Effects:


(A) Description of Effects: Describe the action(s) that may affect the species and critical habitats listed in item IV. Your rationale for the Section 7 determinations made below (B) should be fully described here.

We anticipate the effects of our proposed action to be entirely beneficial to ESA listed and candidate species. In the spatial conservation prioritization for the project, the highest weights were granted to protection of lands in a 200m buffer around southwestern willow flycatcher and yellow-billed cuckoo habitat, as identified by mapping for the draft SLV HCP; and to protection of a 100m buffer around designated conservation populations of Rio Grande cutthroat trout. Included in the Marxan model for spatial conservation prioritization were the occupied, vacant, and potential habitat for Gunnison sage-grouse identified in the GUSG Rangewide Conservation Plan, which may allow for future expansion of the Poncha Pass population and potential reintroduction elsewhere in the valley. Also included was the Canada lynx habitat identified by Colorado Parks and Wildlife (and extended into New Mexico using the land cover classes which comprise most of that habitat). The conservation area includes the montane forests of Costilla County, of which much is currently unprotected. Bringing them into the conservation estate will ensure that this area, identified as an important movement corridor (Leslie Ellwood, CO ES Field Office, pers. comm. with Mike Dixon), will remain largely undeveloped and unfragmented. Please see attached maps.

No explicit population goals are associated with the conservation prioritization strategy at present. However, we expect that the permanent protection of these habitats will allow these imperiled species to maintain their existing populations and/or provide opportunities for future growth.







(B) Determination: Determine the anticipated effects of the proposed project on species and critical habitats listed in item IV. Check all applicable boxes and list the species (or attach a list) associated with each determination.

	<u>Determination</u>
<p><i>No Effect:</i> This determination is appropriate when the proposed project will not directly or indirectly affect (neither negatively nor beneficially) individuals of listed/proposed/candidate species or designated/proposed critical habitat of such species. No concurrence from ESFO required.</p>	_____
<p><i>May Affect but Not Likely to Adversely Affect:</i> This determination is appropriate when the proposed project is likely to cause insignificant, discountable, or wholly beneficial effects to individuals of listed species and/or designated critical habitat. Concurrence from ESFO required.</p>	_____ X
<p><i>May Affect and Likely to Adversely Affect:</i> This determination is appropriate when the proposed project is likely to adversely impact individuals of listed species and/or designated critical habitat. Formal consultation with ESFO required.</p>	_____
<p><i>May affect but Not Likely to Jeopardize candidate or proposed species/critical habitat:</i> This determination is appropriate when the proposed project may affect, but is not expected to jeopardize the continued existence of a species proposed for listing or a candidate species, or adversely modify an area proposed for designation as critical habitat. Concurrence from ESFO optional.</p>	_____
<p><i>Likely to Jeopardize candidate or proposed species/critical habitat:</i> This determination is appropriate when the proposed project is reasonably expected to jeopardize the continued existence of a species proposed for listing or a candidate species, or adversely modify an area proposed for designation as critical habitat. Conferencing with ESFO required.</p>	_____

Signature 
[Supervisor at originating station]

Date 3/14/2012

Reviewing Ecological Services Office Evaluation (check all that apply):

A. Concurrence Nonconcurrency
Explanation for nonconcurrency: _____

B. Formal consultation required
List species or critical habitat unit _____

C. Conference required
List species or critical habitat unit _____

Name of Reviewing ES Office Susan C. Linner
(Colorado) _____

Signature

Susan C. Linner

Date 6/28/2012

Name of Reviewing ES Office _____
(New Mexico)

Signature

Date

Reviewing Ecological Services Office Evaluation (check all that apply):

A **Concurrence** **Nonconcurrence**
Explanation for nonconcurrence _____

B Formal consultation required
List species or critical habitat unit _____

C. Conference required
List species or critical habitat unit _____

Name of Reviewing ES Office _____
(Colorado)


Signature _____

Date _____

Name of Reviewing ES Office George D. Dennis
(New Mexico)

Consultation Number 02ENNM00-2012-I-0057

Signature _____



5/10/2012
Date

Literature Cited

- American Farmland Trust. 2000. Strategic ranchland in the Rocky Mountain west—mapping the threats to prime ranchland in seven western States: Washington DC: American Farmland Trust. <www.farmland.org/resources/rockymtn/default.asp>.
- . 2001. Cost of community service studies. Washington DC: American Farmland Trust.
- Athearn, F.J. 1989. A forgotten kingdom: the Spanish frontier in Colorado and New Mexico, 1540–1821. Denver, CO: Bureau of Land Management. Cultural Resource Series Number 29.
- Athearn, F.J. 1985. Land of contrast: A history of southeast Colorado. Denver, CO: Bureau of Land Management. Cultural Resources Series Number 17.
- Averyt, K.; Cody, K.; Gordon, E.; [and others]. 2011. Colorado climate preparedness project: final report. Prepared by the Western Water Assessment for the State of Colorado. Boulder, CO: University of Colorado.
- Azure, D.A. 1998. Aspects of American bittern ecology in northwest Minnesota [master's thesis]. Grand Forks, ND: University of North Dakota. 139 p.
- Bailey, A.M.; Niedrach, R.J. 1967. Pictorial checklist of Colorado birds: with brief notes on the status of each species in neighboring States of Nebraska, Kansas, New Mexico, Utah, and Wyoming. Denver, CO: Denver Museum of Natural History. 168 p.
- Ball, I.R.; Possingham, H.P.; Watts, M. 2009. Marxan and relatives: software for spatial conservation prioritisation. Chapter 14: p. 185–95 in Spatial conservation prioritisation: quantitative methods and computational tools. Moilanen, A.; Wilson, K. A.; Possingham, H.P., editors. Oxford, UK: Oxford University Press. 10 p.
- Banzhaf, S.; Oates, W.; Sanchirico, J.; [and others]. 2006. Voting for conservation: What is the American electorate revealing? Resources, volume 160, p. 8–12.
- Bayne, E.M.; Dale, B.C. 2011. Effects of energy development on songbirds. In: Naugle, D.E. Energy development and wildlife conservation in western North America. Washington, DC: Island Press.
- Beason, J.; Leivad, R.; Leukering, T. 2005. Monitoring Colorado's birds: the 2004 field season report. Unpublished report, RMBO, Brighton, CO.
- Bent, A.C. 1963. Life histories of North American marsh birds. New York: Dover Publications, Inc. 392 p.
- Bringer, W.L., Jr. 1996. The ecology of the American bittern in northwest Minnesota [master's thesis]. Saint Cloud, MN: Saint Cloud State University. 70 p.
- Brown, M.; Dinsmore, J.J. 1986. Implications of marsh size and isolation for marsh bird management. Journal of Wildlife Management 50:392–97.
- Bureau of Labor Statistics. 2008. Labor force data by county: 2008 annual averages. Washington, DC: Bureau of Labor Statistics, U.S. Department of Labor. <[ftp://ftp.bls.gov/pub/special.requests/la/laucnty08.txt](http://ftp.bls.gov/pub/special.requests/la/laucnty08.txt)> accessed November 14, 2011.
- . 2011b. Labor force data by county (not seasonally adjusted): July 2010–August 2011. Washington, DC: Bureau of Labor Statistics, U.S. Department of Labor. <www.bls.gov/lau/laucntycur14.txt> accessed November 14, 2011.
- . 2011a. Local area unemployment statistics. Washington, DC: Bureau of Labor Statistics, U.S. Department of Labor. <www.bls.gov/web/laus/laumstrk.htm> accessed November 14, 2011.
- [BLM] Bureau of Land Management. 2012. Draft geothermal leasing RMP amendment EA. [City unknown], CO: U.S. Department of the Interior, Bureau of Land Management, San Luis Valley Field Office.
- [BLM and DOE] Bureau of Land Management and U.S. Department of Energy. 2010. Solar energy development draft programmatic environmental impact statement. [City, State unknown].
- Bureau of Reclamation. 2013. West-wide climate risk assessment: Upper Rio Grande impact assessment. Albuquerque, NM: U.S. Department of the Interior, Bureau of Reclamation, Upper Colorado Region, Albuquerque Area Office. 138 p.
- Byers, E.; Ponte, K. 2005. The conservation easement handbook. Washington, DC: Land Trust Alliance and San Francisco, CA: The Trust for Public Land. 555 p.
- Carson, P. 1998. Across the northern frontier: Spanish explorations in Colorado. Boulder, CO: Johnson Books.
- Carter, C. J. 1978. Pike in Colorado. Fort Collins, CO: The Old Army Press.
- Cassells, E. S. 1997. The archaeology of Colorado. Boulder, CO: Johnson Books.
- The Center for Grant Studies. 2003. <www.southwestbooks.org/sangredecristo.htm> accessed on March 17, 2012.

- Center for Rural Entrepreneurship. 2008. Final report: San Luis Valley targeted industry study. <<http://scseed.org/wb/media/8-25-08%20SLV%20Final%20Report.pdf>> accessed March 1, 2012.
- Collinge, S. K. 2009. Ecology of fragmented landscapes. Baltimore, MD: Johns Hopkins University Press.
- Colorado Department of Local Affairs. 2002. Preliminary population projections for Colorado, (table 3 C). Colorado Demography Section. <www.epa.gov/ttn/naaqs/ozone/areas/pop/popp_co.pdf> accessed February 29, 2012.
- . 2011. Colorado planning and management region report: region 8—Alamosa, Conejos, Costilla, Mineral, Rio Grande, and Saguache Counties. San Luis Valley Development Resources Group, Inc and Colorado Demography Office. <<http://dola.colorado.gov/dlg/demog/profiles/reports/region8.pdf>> accessed February 29, 2012.
- Colorado Division of Property Taxation. 2006. How agricultural property is valued in Colorado, Department of Local Affairs 15–DPT–AR PUB B5. <www.dola.state.co.us/propertytax> accessed March 6, 2012.
- Colorado Division of Reclamation, Mining, and Safety. 2012. Division of Reclamation, Mining, and Safety map. Department of Natural Resources. <<http://drmsmaps.state.co.us>> accessed February 6, 2012.
- Colorado Division of Wildlife. 2010. San Luis Valley: habitat partnership program habitat management plan 2010–2014. <<http://wildlife.state.co.us/SiteCollectionDocuments/DOW/Education/pdf/HPP/SLValleyApprovedPlan.pdf>> accessed February 29, 2012.
- Colorado Legislative Council Staff. 2011. Focus Colorado: economic and revenue forecast. Denver, CO: Colorado General Assembly. <www.colorado.gov/legcouncil/Forecast/11JuneForecast.pdf> accessed February 29, 2012.
- Colorado Natural Heritage Program. 2011. CNHP maps for download. Fort Collins, CO: Colorado State University. <www.cnhp.colostate.edu/download/gis.asp> accessed July 23, 2012.
- Colorado Natural Heritage Program and The Nature Conservancy. 2008. A biodiversity scorecard for Colorado. Unpublished report, draft dated October 20, 2008. Fort Collins, CO: Colorado Natural Heritage Program, Colorado State University and Boulder, CO: The Nature Conservancy.
- Colorado Parks and Wildlife. 2012. Boreal toad. <<http://wildlife.state.co.us/Research/Aquatic/BorealToad/Pages/BorealToad.aspx>> accessed March 7, 2012.
- . 2015. State Wildlife Action Plan. A strategy for conserving wildlife in Colorado (in draft), Denver, CO. 448p.
- Colorado Partners in Flight. 2000. Colorado land bird conservation plan. [City unknown], CO.
- Colorado State Archives. 2001. Spanish-Mexican land grants. <www.colorado.gov/dpa/doit/archives/mlg/mlg.html> accessed March 17, 2011.
- Colorado State University Herbarium. 2012. Species lists for each county in Colorado. <http://herbarium.biology.colostate.edu/colorado_map.html> accessed January 17, 2012.
- Colorado Tourism Office. 2012. Town of San Luis. <www.colorado.com/Articles.aspx?aid=42178> accessed March 1, 2012.
- Colville, R.M. 1995. La Vereda: a trail through time. Alamosa, CO: San Luis Valley Historical Society.
- Copeland, H.E.; Doherty, K.E.; Naugle, D.E.; [and others]. 2009. Mapping oil and gas development potential in the US Intermountain West and estimating impacts to species. PLoS One 4:e7400
- Coues, Elliot. 1965. The journal of Jacob Fowler: narrating an adventure from Arkansas through the Indian Territory, Oklahoma, Kansas, Colorado and New Mexico to the sources of Rio Grande Del Norte, 1821–1822. Edited with notes by Elliott Coues, copyright 1898 by Francis P. Harper. Minneapolis, MN: Ross and Haines, Inc.
- Coupal, R.; Taylor, D.T.; McLeod, D. 2002. The cost of community services for rural residential development in Wyoming. Laramie, WY: Wyoming Open Spaces. 4 p.
- Creel, D.; Scott, R.F., IV; Collins, M.B. 1990. A faunal record from west central Texas and its bearing on late Holocene bison population changes in the southern Plains. Plains Anthropologist 35(127):55–69.
- Crompton, J. 2001. The impact of parks on property values: a review of the empirical literature. Journal of Leisure Research 33(1):1–31.
- Cunningham, M.A.; Johnson, D.H. 2006. Proximate and landscape factors influence grassland bird distributions. Ecological Applications 16:1062–75.
- Dahl, Thomas E. 1990. Wetlands losses in the United States 1780s to 1980s. Washington, DC: U.S. Fish and Wildlife Service.
- . 2000. Report to Congress on the status and trends of wetlands in the conterminous United States 1986 to 1997. Washington, DC: U.S. Fish and Wildlife Service. 82 p.
- Daily, G. 1997. Nature's services. Washington, DC: Island Press.
- Daub, B.C. 1993. Effects of marsh area and characteristics on avian diversity and nesting success [master's thesis]. Ann Arbor, MI: University of Michigan. 37 p.
- Dechant, J.A.; Sondreal, M.L.; Johnson, D.H.; [and others]. 2004. Effects of management practices on grassland birds: American bittern. USGS Northern Prairie Wildlife Research Center. Paper 126. <<http://digitalcommons.unl.edu/usgsnpwrc/126>>.
- Duebbert, H.F.; Lokemoen, J.T. 1977. Upland nesting of American bitterns, marsh hawks, and short-eared

- owls. Jamestown, ND: U.S. Fish and Wildlife Service, Northern Prairie Wildlife Research Center.
- Dukes, J.S.; Mooney, H.A. 2004. Disruption of ecosystem processes in western North America by invasive species. *Revista Chilena de Historia Natural* 77:411–37.
- Ehrlich, P.; Ehrlich, A. 1992. The value of biodiversity. *Ambio* 21(3):219–26.
- Ellis, R. 1996. Blair, R. editor. *The Utes in the western San Juan Mountains: their geology, ecology and human history*. [City unknown], CO: University Press of Colorado.
- Emery, Philip A. [date unknown]. *Hydrogeology of the San Luis Valley, Colorado: an overview and a look at the future*. Report compiled for the National Park Service. <www.nps.gov/grsa/naturescience/upload/Trip2023.pdf> accessed March 1, 2012.
- Engel, J.B. 2007. Development, status, and viability of the conservation easement as a private land conservation tool in the eastern United States. *Urban Lawyer* 39:19–74.
- ERO Resources Corporation. 2012. *San Luis Valley regional habitat conservation plan*, November 2012. Prepared for Rio Grande Water Conservation District. 136 p.
- Faanes, C.A. 1981. *Birds of the St. Croix River Valley: Minnesota and Wisconsin*. North American Fauna 73. Washington, DC: U.S. Fish and Wildlife Service. 196 p.
- Freeman, J.; Wunder, L. 1988. Observations at a colony of Brazilian free-tailed bat (*Tadarida brasiliensis*) in southern Colorado. *The Southwestern Naturalist* 33:102–104.
- Frey, J.K.; Malaney, J.L. 2009. Decline of the meadow jumping mouse (*Zapus hudsonius luteus*) in two mountain ranges in New Mexico. *The Southwestern Naturalist* 54:31–44.
- Galbraith, K. 2010. Colorado increases renewables requirements. *New York Times*, March 22, 2010, edition. <<http://green.blogs.nytimes.com/2010/03/22/colorado-to-boost-renewables-requirements>> accessed February 29, 2012.
- Geoghegan, J.; Lynch, L.; Bucholtz, S. 2003. Capitalization of open spaces into housing values and the residential property tax revenue impacts of agricultural easement programs. *Agricultural and Resource Economics Review* 32(1):33–45.
- Goslee, S.C.; Beck, K.G.; Peters, D.P.C. 2003. Distribution of Russian knapweed in Colorado: climate and environmental factors. *Journal of Range Management* 56:206–12.
- Gunnison Sage-Grouse Rangewide Steering Committee. 2005. *Gunnison sage-grouse rangewide conservation plan*. Denver, CO: Colorado Division of Wildlife.
- Haas, C.A. 1995. Dispersal and use of corridors by birds in wooded patches on an agricultural landscape. *Conservation Biology* 9:845–54.
- Hanser, S. 2010. *Western U.S. wind resource at 50 meters above ground level*. Boise, ID: U.S. Geological Survey. Raster data. <http://sagemap.wr.usgs.gov/wus_50mwind.zip>.
- Hart, S.H.; Hulbert, A.B. 2006. *Hart, Hulbert, editors. The southwestern journals of Zebulon Pike: 1806–1807*. Albuquerque, NM: University of New Mexico Press.
- Headwaters Economics. 2011. *Economic profile system-human dimensions toolkit version 6.01 (EPS-HDT)*. <<http://headwaterseconomics.org/tools/eps-hdt>> accessed February 27, 2012.
- Hebblewhite, M. 2011. Effects of energy development on ungulates. In: Naugle, D.E. *Energy development and wildlife conservation in western North America*. Washington, DC: Island Press.
- Heitmeyer, M.E.; Aloia, C.M. 2013. *Hydrogeomorphic evaluation of ecosystem restoration and management options for Baca National Wildlife Refuge*. Greenbrier Wetland Services Report No. 13–08. Bloomfield, MO: Blue Heron Conservation Design and Printing LLC.
- Hilty J.A.; Merenlender, A.M. 2004. Use of riparian corridors and vineyards by mammalian predators in northern California. *Conservation Biology* 18:126–35.
- Hofer, T. 1999. Archaic stage. In: Martorano, M.A.; Hofer, T., III; Jodry, M.A.; [and others]; editors. *Colorado prehistory: a context for the Rio Grande basin*. [City unknown], CO: Colorado Council of Professional Archaeologists. 115–28.
- Jaffe, M. 2011. *Solar-energy industry taking a shine to San Luis*. *Denver Post*, March 6, 2010, edition. <www.denverpost.com/business/ci_17544881> accessed February 29, 2012.
- Jodry, M.A. 1999a. Paleoindian stage. In: Martorano, M.A.; Hofer, T., III; Jodry, M.A.; [and others]; editors. *Colorado prehistory: a context for the Rio Grande basin*. [City unknown], CO: Colorado Council of Professional Archaeologists. 45–114.
- . 1999b. Paleoindian stage paleoecological records. In: Martorano, M.A.; Hofer, T., III; Jodry, M.A.; [and others]; editors. *Colorado prehistory: a context for the Rio Grande basin*. [City unknown], CO: Colorado Council of Professional Archaeologists. 12–26.
- Jones, B.A. 2000. *Archeological overview and assessment for the Alamosa/Monte Vista National Wildlife Refuge Complex, Alamosa, Costilla, and Rio Grande Counties, Colorado*. Prepared for the Mountain-Prairie Region, U.S. Fish and Wildlife Service. Lincoln, NE: National Park Service, Midwest Archaeological Center.

- Jordan, N.; Boody, G.; Broussard W.; [and others]. 2007. Sustainable development of the agricultural bio-economy. *Science* 316:1570–71.
- Karl, R.T.; Melillo, J.M.; Peterson, T.C. 2009. Global climate change impacts in the United States. New York: Cambridge University Press.
- Kessler, R. 1998. Old Spanish Trail North Branch, and its travelers. Santa Fe, NM: Sunstone Press.
- King, T.L.; Switzer, J.F.; Morrison, C.L.; [and others]. 2006. Comprehensive genetic analyses reveal evolutionary distinction of a mouse (*Zapus hudsonius preblei*) proposed for delisting from the U.S. Endangered Species Act. *Molecular Ecology* 15:4331–59.
- Kingsley, H.E. 1998. Colorado breeding bird atlas. [City unknown], CO: Colorado Division of Wildlife. 648 p.
- Knick, S.T.; Rotenberry, J.T. 1995. Characteristics of fragmented shrubsteppe habitats and breeding passerine birds. *Conservation Biology* 9:1059–71.
- Kroger, T. 2008. Open space property value premium analysis. Washington, DC: National Council for Science and the Environment, Wildlife Habitat Policy Research Program, Conservation Economics Program, Defenders of Wildlife.
- Lambeck, R.J. 1997. Focal species: a multi-species umbrella for nature conservation. *Conservation Biology* 11:849–56.
- Laubhan, M.K.; Gammonley, J.H. 2000. Density and foraging habitat selection of waterbirds breeding in the San Luis Valley of Colorado. *Journal of Wildlife Management* 64:808–19.
- Lopez-Tushar, O. 1997. The people of El Valle: a history of the Spanish colonials in the San Luis Valley. Pueblo, CO: El Escritorio.
- Loss S.R.; Terwilliger L.A.; Peterson A.C. 2011. Assisted colonization: integrating conservation strategies in the face of climate change. *Biological Conservation* 144:92–100.
- Maestre, F. T.; Quero, J.L.; Gotelli, N.J.; [and others]. 2012. Plant species richness and ecosystem multifunctionality in global drylands. *Science* 335:214–18.
- Martorano, M. 1999a. Post paleoindian paleoenvironmental studies. In: Martorano, M.A.; Hoefler, T., III; Jodry, M.A.; [and others]. *Colorado prehistory: a context for the Rio Grande basin*. [City unknown], CO: Colorado Council of Professional Archaeologists. 27–30.
- . 1999b. Late prehistoric/ceramic stage. In: Martorano, M.A.; Hoefler, T., III; Jodry, M.A.; [and others]. *Colorado prehistory: a context for the Rio Grande basin*. [City unknown], CO: Colorado Council of Professional Archaeologists. 129–37.
- . 1999c. Protohistoric stage. In: Martorano, M.A.; Hoefler, T., III; Jodry, M.A.; [and others]. *Colorado prehistory: a context for the Rio Grande basin*. [City unknown], CO: Colorado Council of Professional Archaeologists. 138–45.
- Mayo, A.L.; Davey, A.; Christiansen, D. 2006. Ground-water flow patterns in the San Luis Valley, Colorado, USA revisited: an evaluation of solute and isotope data. [City, State unknown]: [publisher unknown].
- McCalpin, J.P. 1996. General geology of the north-east San Luis Valley, Colorado in Geological Society of America—Colorado geological survey field trip guidebook: geologic excursions to the Rocky Mountains and beyond. Annual meeting, October 1996. Denver, CO: Geological Society of America.
- McConnell, V.; Walls, M. 2005. The value of open space—evidence from studies of nonmarket benefits: Washington, DC: Resources for the Future. 79 p.
- Merenlender, A.M.; Huntsinger, L.; Guthey, G.; Fairfax, S.K. 2004. Land trusts and conservation easements: who is conserving what for whom? *Conservation Biology* 18:65–75.
- Middleton, D.S. 1949. Close proximity of two nests of American bitterns. *Wilson Bulletin* 61:113.
- Millennium Ecosystem Service Assessment. 2005. Ecosystems and human well-being: synthesis report. Washington, DC: Island Press.
- Morrison, J. 1990. The meadow jumping mouse in New Mexico: habitat preferences and management recommendations. In: Krausman, P.R.; Smith, N.S.; editors. *Managing wildlife in the Southwest: proceedings of the symposium*. [Date and location of symposium unknown]. Phoenix, AZ: Arizona Chapter of the Wildlife Society. 136–43.
- Morrison, J. 1992. Persistence of the meadow jumping mouse, *Zapus hudsonius luteus*, in New Mexico. *The Southwestern Naturalist* 37:308–11.
- Mowat, G.; Slough, B. 2003. Habitat preference of Canada lynx through a cycle in snowshoe hare abundance. *Canadian Journal of Zoology* 81:1736–45.
- [NASA] National Aeronautics and Space Administration. 2006. MODIS land cover type, compiled using Headwaters Economics Economic profile system-human dimensions toolkit version 6.01. [City, State unknown].
- [NPS] National Park Service. 2011. San Luis Valley and Central Sangre de Cristo Mountains: Reconnaissance Survey Report (draft). Denver, CO: Intermountain Regional Office, National Park Service.
- National Renewable Energy Laboratory. 2007a. Concentration solar resource of the Southwest United States. Washington, DC: U.S. Department of Energy. <www.nrel.gov/csp/pdfs/csp_sw.pdf> accessed February 29, 2012.
- . 2007b. Photovoltaic solar resource of the United States. Washington, DC: U.S. Department of Energy. <www.nrel.gov/gis/images/map_pv_national_lo-res.jpg> accessed February 29, 2012.
- National Weather Service. 2012. Alamosa, Colorado climate normals (1971–2000). <www.crh.noaa.gov/pub/?n=climate/cli/alamosa.php> accessed January 17, 2012.

- [NRCS] Natural Resources Conservation Service. 2012. Working lands for wildlife. <www.fws.gov/workinglandsforwildlife.html> accessed March 11, 2012.
- The Nature Conservancy. 2008. The Great Sand Dunes project: measures of conservation success. Project summary. Boulder, CO: The Nature Conservancy.
- Naugle, D.E.; Doherty, K.E.; Walker, B.L.; [and others]. 2011. Sage-grouse and cumulative impacts of energy development. In: Naugle, D.E. Energy development and wildlife conservation in western North America. Washington, DC: Island Press.
- Netusil, N.R.; Boyd, E.; Griffen, Z.V.; [and others]. 2000. Can open space be self-financing? *Choices* 15(2):21–23.
- New Mexico Taxation and Revenue Department. 2011. Regulations pertaining to the Property Tax Code, Sections 7–35–1 to 7–28–93 NMSA 1978. <www.tax.newmexico.gov/SiteCollectionDocuments/Tax-Library/Statutes-and-Department-Directives/Recent-Regulation-Changes/Property_Tax_Code.pdf> accessed March 6, 2012.
- The New York Times. 1898. Up the Rio Grande in 1821: Jacob Fowler's quaint and interesting journal of the trip. June 18, 1898. New York: The New York Times.
- Nhancale, B.A.; Smith, R.J. 2011. The influence of planning unit characteristics on the efficiency and spatial pattern of systematic conservation planning assessments. *Biodiversity Conservation* 20:1821–35.
- Noss, R.F.; Dobson, A.P.; Baldwin, R.; [and others]. 2012. Bold conservation. *Conservation Biology* 26:1–4.
- Oyler-McCance, S.J.; Burnham, K.P.; Braun, C.E. 2001. Influence of changes in sagebrush on Gunnison sage-grouse in southwestern Colorado. *The Southwestern Naturalist* 46:323–31.
- Paxton, E.H. 2000. Molecular genetic structuring and demographic history of the willow flycatcher (*Epidomax traillii*) [master's thesis]. Flagstaff, AZ: Northern Arizona University.
- Pennington D.N.; Hansel J.; Blair R.B. 2008. The conservation value of urban riparian areas for landbirds during spring migration: landcover, scale, and vegetation effects. *Biological Conservation* 141:1235–48.
- Phillips, S. 2000. Windfalls for wilderness: land protection and land value in the Green Mountains. In: McCool, S.F.; Cole, D.N.; Borrie, W.T.; O'Loughlin, J. Wilderness science in a time of change conference. Volume 2: wilderness in the context of larger systems. Proceedings RMRS–P–15–VOL–2:258–67, Missoula, MT, May 23–27, 1999. Ogden, UT: Rocky Mountain Research Station, USDA Forest Service.
- Phillips, S.J.; Anderson, R.P.; Shapire, R. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modeling* 190:231–59.
- Pitkin, M.; Quattrini, L. 2010. Pocket guide to sagebrush birds. Brighton, CO: Rocky Mountain Bird Observatory and Petaluma, CA: Point Reyes Bird Observatory.
- Preuss, C. 1958. Exploring with Fremont. Gudde, E.G.; Gudde, E.K.; editors. Norman, OK: University of Oklahoma Press.
- Pritchard, V.L.; Cowley, D.E. 2006. Rio Grande cutthroat trout (*Oncorhynchus clarkia virginalis*): a technical conservation assessment. USDA Forest Service, Rocky Mountain Region. <www.fs.fed.us/r2/projects/scp/assessments/riograndecutthroattrout.pdf> accessed March 7, 2013.
- The Pueblo Chieftain. 2009. Taylor Ranch owners appeal court's handling of access proceedings. September 12, 2009. <www.chieftain.com/news/local/article_3a47173d-a18f-5af5-9ad7-545ff94a6a7c.html> accessed on March 17, 2012.
- Rantalainen, M.L.; Fritze, H.; Haimi, J.; [and others]. 2005. Species richness and food web structure of soil decomposer community as affected by the size of habitat fragmentation and habitat corridors. *Global Change Biology* 11:1614–27.
- Ray, A.; Barsugli, J.; Averyt, K.; [and others]. 2008. Climate change in Colorado: a synthesis to support water resources management and adaptation. Intermountain West Climate Summary 4:2–4.
- Reed, A.D. 1994. The numic occupation of western Colorado and eastern Utah during the prehistoric and protohistoric periods. In: Madsen, D.B.; Rohde, D.; editors. Across the West: human population movement and the expansion of the numa, Salt Lake City, UT: University of Utah Press. 188–99.
- Rees, D.E.; Carr, R.J.; Miller, W.J. 2005a. Rio Grande chub (*Gila pandora*): a technical conservation assessment. USDA Forest Service, Rocky Mountain Region. <www.fs.fed.us/r2/projects/scp/assessments/riograndechub.pdf> accessed January 31, 2012.
- Rees, D.E.; Miller, W.J. 2005b. Rio Grande sucker (*Catostomus plebeius*): a technical conservation assessment. USDA Forest Service, Rocky Mountain Region. <www.fs.fed.us/r2/projects/scp/assessments/riograndesucker.pdf> accessed January 31, 2012.
- Renaud, E.B. 1942. Reconnaissance work in the upper Rio Grade Valley: Colorado and New Mexico. Department of Anthropology Archaeological Series: Third Paper. Denver, CO: University of Denver.
- The Republican Publishing Company. 1884. Immigrants' guide to the Great San Luis Park: where can be had the best farming lands in the world. Denver, CO: The Republican Publishing Company.
- Reynolds, R.; Shaffer, T.; Renner, R.; [and others]. 2001. Impact of the conservation reserve program on duck recruitment in the U.S. prairie pothole region. *Journal of Wildlife Management* 65(4):765–80.
- Rich, T.D.; Beardmore, C.J.; Berlanga, H.; [and others]. 2004. Partners in Flight North American

- landbird conservation plan. Ithaca, NY: Cornell Lab of Ornithology.
- Richmond, P. J. 1990. Trail to disaster. Denver, CO: Colorado Historical Society.
- Riffell, S.K.; Keas, B.E.; Burton, T.M. 2001. Area and habitat relationships of birds in Great Lakes coastal wet meadows. *Wetlands* 21:492–507.
- Rio Grande Cutthroat Trout Conservation Team. 2009. Conservation agreement for Rio Grande cutthroat trout (*Oncorhynchus clarkia virginalis*) in the States of Colorado and New Mexico. Fort Collins, CO: Colorado Division of Wildlife. 24 p.
- Rio Grande Water Conservation District. 2012. The closed basin project. <www.rgwcd.org/page21.html> accessed March 7, 2012.
- Ripple, W.J.; Beschta, R.L. 2012. Trophic cascades in Yellowstone: the first 15 years after wolf reintroduction. *Biological Conservation* 145:205–13.
- Rissman, A.; Lozier, L.; Comendant, T.; Kareiva, P.; Kiesecker, J.; Shaw, M.; et al. 2007. Conservation easements: biodiversity protection and private use. *Conservation Biology* 21(3):709–18.
- Robson, S.; Banta, E. 1995. Ground water atlas of the United States: Arizona, Colorado, New Mexico, Utah: Rio Grande aquifer system. [City, State unknown]: U.S. Geological Survey.
- Rowland, M.M.; Wisdom, M.J.; Suring, L.H.; Meinke, C.W. 2006. Greater sage-grouse as an umbrella species for sagebrush-associated vertebrates. *Biological Conservation* 129:323–35.
- Sanchez, J.P. 1997. Explorers, Traders, and Slavers: Forging the Old Spanish Trail, 1678–1850. Salt Lake City, UT: University of Utah Press.
- Sanford, A.B. 1933. Recollections of a trip to the San Luis Valley in 1877. In: Grinstead, S.; Fogelberg, B.; editors. *Western voices: 125 years of Colorado writing*. 2004. Colorado Historical Society. Golden, CO: Fulcrum Publishing.
- Sangre de Cristo National Heritage Area. 2012. Hispano culture, folklore, religion, and language. <<http://sdcnha.org/js/hispano-culture-sdcnha.html>> accessed March 6, 2012.
- Schroeder, M.A.; Aldridge, C.L.; Apa, A.D.; [and others]. 2004. Distribution of sage-grouse in North America. *Condor* 106:363–76.
- Shepherd, B.; Whittington, J. 2006. Response of wolves to corridor restoration and human use management. *Ecology and Society* 11: 1. <www.ecologyandsociety.org/vol11/iss2/art1/>.
- Siebenthal, C.E. 1910. Geology and water resources of the San Luis Valley. United States Geological Survey Water-Supply Paper 240. Washington, DC: Government Printing Office.
- Stoller, M.L.; Steele, T.J. 1982. Diary of the Jesuit residence of Our Lady of Guadalupe Parish, Conejos, Colorado: December 1871–December 1875. The Colorado College Studies, Number Nineteen. Colorado Springs, CO: The Colorado College.
- Taylor, D.T.; Coupal, R.H. 2000. The cost of rural community services in Laramie, Wyoming. Laramie, WY: Department of Agricultural and Applied Economics, University of Wyoming.
- Telleria, J.L.; Santos, T. 1995. Effects of forest fragmentation on a guild of wintering passerines: the role of habitat selection. *Biological Conservation* 71:61–67.
- Thorsnes, P. 2002. The value of a suburban forest preserve: estimates from sales of vacant residential building lots. *Land Economics* 78(3):426–41.
- Tremblay, Marie A. 2001. Modeling and management of potential movement for elk (*Cervus elaphus*), bighorn sheep (*Ovis canadensis*) and grizzly bear (*Ursus arctos*) in the Radium Hot Springs area, British Columbia. Davis, CA: Road Ecology Center, University of California, Davis. <<http://escholarship.org/uc/item/1vt0p0h8>> accessed October 20, 2011.
- Ubbelodhe, C.; Benson, M.; Smith, D.A.; 2001. A Colorado History. Boulder, CO: Pruett Publishing Company.
- University of New Mexico. 2002. Projected population: New Mexico counties 2000 to 2030. Bureau of Business and Economic Research. <www.epa.gov/ttn/naaqs/ozone/areas/pop/popp_nm.htm> accessed February 29, 2012.
- Urban Institute and Brookings Institution. 2008. State and local tax policy: What are the sources of revenue for local governments? Washington, DC: Tax Policy Center. <www.taxpolicycenter.org/briefing-book/state-local/revenues/local_revenue.cfm> accessed July 18, 2011.
- U.S. Army Corps of Engineers. 1878. Land classification map of part of southwestern Colorado atlas sheet, number 61(D). U.S. Army Corps of Engineers. [City, State unknown].
- U.S. Census Bureau. 2010a. American factfinder. Washington, DC: U.S. Census Bureau. <<http://factfinder2.census.gov/main.html>> accessed February 28, 2012.
- . 2010b. State and county quickfacts. Washington, DC: U.S. Census Bureau. <<http://quickfacts.census.gov/qfd/index.html>> accessed February 28, 2012.
- U.S. Department of Agriculture. 2009. Census of agriculture, (table 8). Compiled using Headwaters Economics Economic profile system-human dimensions toolkit version 6.01. Washington, DC: National Agricultural Statistics Service.
- U.S. Department of Commerce. 2009. Regional economic information system, (table CA25N). Washington, DC: Bureau of Economic Analysis. <www.bea.gov/regional/reis> accessed February 28, 2012.
- . 2010a. Regional economic information system, (table CA25N and CA25N). Compiled using Headwaters Economics Economic profile system-human

- dimensions toolkit version 6.01. Washington, DC: Bureau of Economic Analysis.
- . 2010b. County business patterns. Compiled using Headwaters Economics Economic profile system-human dimensions toolkit version 6.01. Washington, DC: Census Bureau.
- [DOE] U.S. Department of Energy. 2011. Department of Energy finalizes \$90.6 million loan to guarantee to Cogentrix of Alamosa, LLC, to support Colorado solar project. <<http://energy.gov/articles/department-energy-finalizes-906-million-loan-guarantee-cogentrix-alamosa-llc-support>> accessed February 29, 2012.
- U.S. Environmental Protection Agency. 2011. Level 3 ecoregions of the Continental United States. national health and environmental effects laboratory. <ftp://ftp.epa.gov/wed/ecoregions/us/Eco_Level_III_US.pdf> accessed January 13, 2012.
- [USFWS] U.S. Fish and Wildlife Service. 2002. Southwestern willow flycatcher recovery plan. Albuquerque, NM. i–ix + 210 p., Appendices A–O.
- . 2005. Baca National Wildlife Refuge conceptual management plan. Lakewood, CO: Region 6, U.S. Fish and Wildlife Service.
- . 2008. 2006 National survey of fishing, hunting, and wildlife-associated recreation: Washington, DC: U.S. Fish and Wildlife Service and U.S. Census Bureau. 164 p.
- . 2008b. Endangered and threatened wildlife and plants; status review for Rio Grande cutthroat trout. Federal Register 73(94):27900–26.
- . 2008c. Birds of Conservation Concern, Arlington Virginia. 93 p.
- . 2010a. Preliminary project proposal: San Luis Valley Conservation Area. Alamosa, CO: San Luis Valley National Wildlife Complex.
- . 2010b. Rising to the urgent challenge: strategic plan for responding to accelerating climate change. Washington, DC: U.S. Fish and Wildlife Service. 32 p.
- . 2011. Environmental assessment of proposed oil and gas exploration, Baca National Wildlife Refuge, Saguache County, Colorado. Lakewood, CO: Region 6, U.S. Fish and Wildlife Service.
- . 2012a. Draft Baca National Wildlife Refuge water resource inventory and assessment. Lakewood, CO: Region 6, U.S. Fish and Wildlife Service.
- . 2012b. Nonindigenous aquatic species HUC–6 query (Rio Grande headwaters). <<http://nas.er.usgs.gov/queries/SpeciesList.aspx?Group=&Status=0&FMB=0&pathway=0&Sortby=1&Size=50&HUCNumber=130100>> accessed January 31, 2012.
- . 2013a. Endangered and threatened wildlife and plants; designation of critical habitat for southwestern willow flycatcher; final rule. Federal Register 78(2):343–44.
- . 2013b. Endangered and threatened wildlife and plants; designation of critical habitat for Gunnison sage-grouse; final rule. Federal Register 79(224):69312–69362.
- . 2013c. Endangered and threatened wildlife and plants; endangered status for Gunnison sage-grouse; final rule. Federal Register 70(224):69192–69310.
- [USGS] U.S. Geological Survey. 2010. Landfire 1.1.0. Existing vegetation type layer. U.S. Department of the Interior, U.S. Geological Survey. <<http://landfire.cr.usgs.gov/viewer>>.
- . 2015. Migratory Birds-Focal Species <<http://www.fws.gov/mountain-prairie/migbirds/migBirdsUpdate/focal-species.php>> accessed March 2015.
- Vrooman, D. 1978. An empirical analysis of determinants of land values in the Adirondack Park. American Journal of Economics and Sociology 37(2):165–77.
- Watkins, T.; Belcher, J.; Gries, R.; Longacre, M. 1995. The surprising discovery of oil and Mesozoic sediments in the San Luis Basin, Saguache County, Colorado. Colorado Scientific Society Abstracts. <www.coloscisoc.org/abstracts/abstracts95.html> accessed October 20, 2011.
- Weber, M.J. 1978. Non-game birds in relation to habitat variation on South Dakota wetlands [master's thesis]. Brookings, SD: South Dakota State University. 54 p.
- Weber, M.J.; Vohs, P.A., Jr.; Flake, L.D. 1982. Use of prairie wetlands by selected bird species in South Dakota. The Wilson Bulletin 94:550–54.
- Wiens, J.A.; Rotenberry, J.T. 1981. Habitat associations and community structure of birds in shrub steppe environments. Ecological Monographs 51:21–41.
- Williams, M.I.; Paige, G.B.; Thuro, T.L.; [and others]. 2011. Songbird relationships to shrub-steppe ecological site characteristics. Rangeland Ecology and Management 64:109–18.
- Wockner, G.; Boone, R.; Schoenecker, K.; Zeigenfuss, L. 2010. The Great Sand Dunes ecosystem elk and bison carrying capacity model: description and scenario results. Fort Collins, CO: Natural Resource Ecology Laboratory, Colorado State University and Biological Resources Division, U.S. Geological Survey.
- Wyckoff, W. 1999. Creating Colorado: the making of a western American landscape, 1860–1940. New Haven, CT: Yale University Press.

