

**NEAT (North East Atlantic Taxa):  
South Scandinavian marine CRUSTACEA Check-List  
compiled at TMBL (Tjärnö Marine Biological Laboratory) by:**

Hans G. Hansson 1990-02-18 / small revisions until April 1996,  
when it for the 2:nd time was published on Internet, and again after some revisions August 1998.

Citation suggested: Hansson, H.G. (Comp.), 1998. NEAT (North East Atlantic Taxa): South Scandinavian marine Crustacea Check-List. Internet pdf Ed., Aug. 1998. [http://www.tmbi.gu.se].

Denotations: (TM) = Genotype @ = Associated to \* = General note

N.B.: This is one of several preliminary check-lists, covering S. Scandinavian marine animal (and partly marine protist) taxa. Some financial support from (or via) NKMB (Nordiskt Kollegium för Marin Biologi), during the last years of the existence of this organisation (until 1993), is thankfully acknowledged. The primary purpose of these checklists is to facilitate for everyone, trying to identify organisms from the area, to know which species that earlier have been encountered there, or in neighbouring areas. A secondary purpose is to facilitate for non-experts to find as correct names as possible for organisms, including names of authors and years of description. So far these checklists are very preliminary. Due to restricted access to literature there are (some known, and probably many unknown) omissions in the lists. Certainly also several errors may be found, especially regarding taxa like Plathelminthes and Nematoda, where the experience of the compiler is very rudimentary. The credibility of some other lists, containing taxa like e.g. Porifera, where, at least in certain subtaxa, taxonomic confusion seems to prevail, is certainly also suboptimal so far.

Species found in South Scandinavia, as well as from neighbouring areas, chiefly the British Isles, have been considered, as some of them may show to have a slightly more northern distribution, than what is known today. However, species with a typical Lusitanian distribution, with their northern distribution limit around France or Southern British Isles, have as a rule been omitted here, albeit a few species with probable northern limits around the British Isles are listed here until distribution patterns are better known. Arctic Taxa, known only from areas north of N Norway are probably not as well represented in this list as they should be. The compiler would be very grateful for every correction of presumptive lapses and omissions an initiated reader could make.

In the context below, Crustacea is ranked as a subphylum (within phylum Arthropoda) with the following superclasses: Maxillopoda, Remipedia, Phyllopoda (Branchiopoda), Ostracoda & Malacostraca. I am fully aware of the incompleteness of this list, but for the moment I cannot advance much further because of limited library resources. Intentionally taxa from neighbouring areas have been listed, as some of them probably will show to occur in Scandinavian waters.

N.B. This document is intended to be completely unavailable regarding nomenclature within the meaning of ICZN.

**ARTHROPODA** Siebold & Stannius, 1845

**CRUSTACEA** Pennant, 1777 (sensu Manton, 1972)

**MAXILLOPODA** Dahl, 1956

\* Some authors are considering Ostracoda as part of this taxon. As this is not yet fully accepted, I have in this context chosen to treat Ostracoda as a taxon of the same rank as Maxillopoda.

**MYSTACOCARIDA** Pennak & Zinn, 1943

**DEROCHEILOCARIDA**

= **MYSTACOCARIDIDA** Pennak & Zinn, 1943

**Derocheilocaris** Pennak & Zinn, 1943

**D. remanei** Delamare Deboutteville & Chappuis, 1951 **biscayensis** Delamare Deboutteville, 1953  
Bay of Biscay-Portugal-Italy

**BRANCHIURA** Thorell, 1864

**ARGULOIDA** Raffinesque-Schmaltz, 1815

**Argulidae** Leach, 1819

@ Chiefly on freshwater fish. Only marine genus is *Argulus* with 30-40 marine species

**Argulus** O.F. Müller, 1785

**A. foliaceus** (Linnaeus, 1758)

@ teleosts (e.g. Salmonidae, Gobidae & Gasterosteidae) (& frogs tadpole larvae)  
Freshwaters in Fennoscandia & Baltic estuarine water

**A. coregoni** Thorell, 18??

@ teleosts (chiefly Salmonidae)  
Freshwaters in Fennoscandia & ?Baltic estuarine water

**A. arcassonensis** Cuénot, 1912

@ on skin of a diversity of fish and on skin of *Sepia officinalis*  
Arcachon

**THECOSTRACA** Gruvel, 1905

**CIRRIPEDIA** Burmeister, 1834

**ACROTHORACICA** Gruvel, 1905

**PYGOPHORIDA** Berndt, 1907

**Lithoglyptidae** Aurivillius, 1892

**Weltneria** Berndt, 1907

**W. exargilla** Newman, 1974

Bay of Biscay at ca 1500 m

**Cryptohialidae** Gerstaecker, 1866

\* No taxa within the area?

**APYGOPHORIDA** Berndt, 1907

**Trypetidae** Stebbing, 1910

**Trypetesa** Norman, 1903

= *Alcippe* Hancock, 1849, non Blyth, 1844

**T. lampas** (Hancock, 1849)

Öresund-Bohuslän-S Norway-North Sea-Br. Isles, W Mediterranean

**THORACICA** Darwin, 1854

**PEDUNCULATA** de Lamarck, 1818

**SCALPELLOMORPHA**

**Scalpellioidea** Pilsbry, 1916

**Scalpellidae** Pilsbry, 1916

**Calanticinae**

**Calantica** Gray, 1825

**C. (Scillaelepas** Seguenza, 1876) **gemma** C.W. Aurivillius, 1894  
off W Iceland

**Scalpellinae** Pilsbry, 1916

**Scalpellum** Leach, 1818

**S. (Strictoscalpellum** Broch, 1924) **scalpellum** (Linnaeus, 1767)

Öresund-Bohuslän-N Norway-Norw. Sea-Faeroes-Br. Isles-W Africa-Mediterranean

**S. (Strictoscalpellum)** **stroemi** M. Sars, 1859

Bohuslän-N Norway-Barents Sea-Spitsbergen-Iceland-Greenland-Faeroes-Shetlands

**S. (Strictoscalpellum)** **nymphocola** Hoek, 1883

Shetlands-Norw. Sea-E Greenland-Spitsbergen & Arctic Seas

**S. (Strictoscalpellum)** **cornutum** G.O. Sars, 1879

Faeroes-Norw. Sea-Spitsbergen & Arctic Seas

**S. (Strictoscalpellum)** **hamatum** G.O. Sars, 1879

Shetlands-off Norway S of Lofoten-off SW Spitsbergen

**S. (Arcoscalpellum** Hoek, 1907) **hispidum** G.O. Sars, 1890

Trondheim-N Norway

**S. (Episcalpellum** Broch, 1924) **striolatum** G.O. Sars, 1877

off NE Faeroes-Norw. Sea-S Spitsbergen & Arctic Seas

**Pollicipes** Leach, 1817

= *Mitella* Oken, 1815 (nomen rejicendum, ICZN Op. 417)

**P. pollicipes** (Gmelin, 1789)

= *Lepas pollicipes* Gmelin, 1789

SW Cornwall - French & Iberian coasts

**LEPADOMORPHA** Pilsbry, 1916

**Lepadoida** Darwin, 1851

**Lepadidae** Darwin, 1851

**Lepas** Linnaeus, 1758

**L. (Lepas** Pilsbry, 1907) **anatifera** Linnaeus, 1758

Öresund-Bohuslän-N Norway-Spitsbergen-Iceland-Faeroes-Shetlands-Br. Isles-southwards

**L. (Lepas)** **anserifera** Linnaeus, 1767

Faeroe Channel-Norw. Sea; on ships to Denm. & Holland

**L. (Lepas)** **hilli** (Leach, 1818)

carried by ships fr. tropical-subtropical waters to Öresund-Bohuslän-Lofoten-Faeroes-Shetlands-North Sea-southwards

**L. (Lepas)** **pectinata** Spengler, 1793

carried by ships to Copenhagen, Bergen, Faeroes, S North Sea, by floatsome to Bohuslän

**L. (Dosima** Gray, 1825) **fascicularis** Ellis & Solander, 1786

on small floating objects from tropical and subtropical latitudes to Belt Sea-Bohuslän-North Sea-Trondheim-Shetlands-Iceland

**Anelasma** Darwin, 1851

**A. squalicola** (Lovén, 1845)

@ *Etmopterus spinax*

Oslofjord-Skagerrak-Lofoten-W Br. Isles-Bay of Biscay

**Conchoderma** Olfers, 1814

**C. virgatum** (Spengler, 1790)

\* attached to living and dead things, like ships, driftwood, sea-snakes, whales and big fishes  
Bohuslän & Copenhagen (ships), Oslofjord, Bergen, Lofoten, Iceland, Faeroes & S North Sea

**C. auritum** (Linnaeus, 1776)

@ *Coronula diadema* (& *C. reginae*)

Bohuslän, Belgium, Lofoten-N Norway-Norw. Sea-Iceland-Faeroes-Shetlands-E Scotland-Atlantic

**Alepes** Sander Rang, 1829

**A. parasita** Sander Rang, 1829

@ medusae (*Cyanea*)

?Falmouth, Atlantic Ocean & Mediterranean

**CYPRILEPADOMORPHA**

**IBLOMORPHA**

**HETERALEPADOMORPHA**

**PRAELEPADOMORPHA**

**SESSILIA** de Lamarck, 1818

**BRACHYLEPADOMORPHA** Withers, 1923

\* One Pacific species, else Extinct.

**VERRUCOMORPHA** Pilsbry, 1916

**Verrucidae** Darwin, 1854

**Verruca** Schumacher, 1817

**V. stroemia** (O.F. Müller, 1776)

Öresund-Bohuslän-North Sea-N Norway-Barents Sea-Spitsbergen-E Greenland-Iceland-Faeroes-Shetlands-Br. Isles-Portugal-Mediterranean

**BALANOMORPHA** Pilsbry, 1916

**Chthamaloidea** Darwin, 1854

**Chthamalidae** Darwin, 1854

**Chthamalinae** Darwin, 1854

**Chthamalus** Ranzani, 1817

**C. stellatus** (Poli, 1791)

Anholt, Helgoland, Faeroes-Shetlands-W Br. Isles-Bretagne-Canaries-Cap Verde-Mediterranean

**C. montagui** Southward, 1976

= *C. punctatus* : Montagu, 1803, non (Martini & Chemnitz, 1788)

Bay of St Malo-Br. Isles-Bay of Biscay-W Morocco

**Euraphiinae** Newman & Ross, 1976

\* Not in the area.

Notochthamalinae Foster & Newman,1987  
\* Not in the area.

Catophragmidae Utinomi,1968  
\* Not in the area.

Coronuloidea Leach,1817

Coronulidae Leach,1817

Coronula de Lamarck,1802

C. diadema (Linnaeus,1767)  
@ whales, often *Megaptera novaeangliae* (& e.g. *Hyperoodon*)  
N Norway-Barents Sea, Iceland-Faeroes-Shetlands

C. reginae Darwin,1854  
@ whales, often *Megaptera novaeangliae* (& e.g. *Balaenoptera* spp.)  
Skagerrak, N Norway, Iceland

Xenobalanus Steenstrup,1852

X. globicipitis Steenstrup,1852  
@ whales, (e.g. *Balaenoptera* spp., *Globicephala melaena*, *Orcinus orca*, *Delphinus delphis* *Tursiops truncatus* & *Grampus griseus*)

Platylepadidae Newman & Ross,1976

Platylepas J.E. Gray,1825

P. hexastylus (O. Fabricius,1798)  
= *Lepas hexastylus* O. Fabricius,1798  
@ a variety of sirenian & chelonian hosts, e.g. *Chelonia mydas* & *Caretta caretta*, but not *Dermochelys coriacea*, while the somewhat suspect "species" *P. multidecorata* is thought to be confined to the last species, which may indicate that it may be a synonym of *P. hexastylus*.  
?Britain & warmer seas

Stomatolepas Pilsbry,1910

S. elegans (Costa,1838)  
@ embedded in the soft skin portion of chelonians, chiefly *Dermochelys coriacea* & *Lepidichelys olivacea* (non-European), but also occasionally *Chelonia mydas* & *Caretta caretta*.  
Norway, Netherlands, Scotland, England, Eire - temperate & warmer seas  
N Norway, Faeroes, Shetlands, Bay of Biscay, W Mediterranean

Chelonibiidae Pilsbry,1916  
\* Not in the area.

Emersoniidae Ross,1967  
\* Not in the area.

Tetraclitoidae Gruvel,1903

Tetraclitidae Gruvel,1903  
\* Not in the area.

Tetraclitinae Gruvel,1903

Tetraclitellinae Newman & Ross,1976

Austrobalaninae Newman & Ross,1976

Bathylasmatidae Newman & Ross,1971

Bathylasmatinae Newman & Ross,1971

Bathylasma Neman & Ross,1971

B. hirsutum (Hoek,1883)  
off Faeroes-southwards in Atlantic-Azores

Hexelasmatinae Newman & Ross,1976  
\* Not in the area.

Balanoidea Leach,1817

Archaeobalanidae Newman & Ross,1976

Archaeobalaninae Newman & Ross,1976

Chirona Gray,1835

C. hameri (Ascanius,1767)  
= *Lepas hameri* Ascanius,1767  
= *Lepas tulipa* O.F. Müller,1776  
= *Balanus candidus* Brown,1827  
Kattegatt-Bohuslän-North Sea-Norway-Barents Sea-Iceland-Faeroes-Shetlands-Br. Isles

C. amaryllis (Darwin,1854)  
= *Balanus amaryllis* Darwin,1854  
Indian Ocean species brought to Europe (?Britain) on ships' bottoms

Acasta Leach,1817

A. spongites (Poli,1791)  
@ embedded in *Dysidea fragilis*  
Br. Isles-France-Portugal-S. Africa-Mediterranean

Elmininae Foster,1982

Elminius Leach,1825

E. modestus Darwin,1854  
Bohuslän (on floatsome)-W Jutland-Engl. Channel-Br. Isles-Bay of Biscay-NW Spain (immigrated fr. Australia-New Zealand area)

Semibalaninae Newman & Ross,1976

Semibalanus Pilsbry,1916

S. balanoides (Linnaeus,1767)  
Öresund & Belt Sea-Bohuslän-North Sea-Norway-Barents Sea-Spitsbergen-Arctic Seas-Iceland-Faeroes-Br. Isles-NW Portugal

Pyrgomatidae Gray,1825

Pyrgomatinae Gray,1825  
=? Bosciinae Newman & Ross,1976

Pyrgoma Leach,1818 ex Savigny MS  
=? *Boscia* Férussac,1822, non Leach,1814 (Amphipoda)

P. anglica G.B. Sowerby II,1823  
@ *Caryophyllia*  
Br. Isles-Ireland-W France-W Africa-Cap Verde-Mediterranean

Ceratoconchinae Newman & Ross,1976  
\* Not in the area.

Megatreminae Holthuis,1982  
\* Not in the area.

Balanidae Leach,1817

Balaninae Leach,1817

Balanus da Costa,1778

B. balanus (Linnaeus,1758)  
SW Baltic-Bohuslän-North Sea-Norway-Barents Sea & Spitsbergen-Arctic Seas-Iceland-Faeroes-Shetlands-Br. Isles-NW Portugal

B. crenatus Bruguière,1789  
Blekinge-Bohuslän-North Sea-Norway-Barents Sea-Spitsbergen-Arctic Seas-Iceland-Faeroes-Br. Isles-Azores-NW Spain-W Algeria

B. improvisus Darwin,1854  
Bothnian Sea-Bohuslän-North Sea-Bergen-W Br. Isles-Morocco-E Mediterranean-Black Sea

B. spongicola Brown,1844  
= *B. dollfusii* Broch,1927  
S Wales-SW England-Portugal-S Africa-Azores-Madeira

B. perforatus perforatus Bruguière,1789  
SW Br. Isles-France-Spain-NW Africa-Mediterranean-Black Sea

B. perforatus angustus (Gmelin,1789)  
SW Br. Isles-France-Spain-Mediterranean Africa

B. perforatus fistulosus (Poli,1791)  
?Denmark, Morocco, Canaries

B. perforatus mirabilis Darwin,1854  
La Rochelle (France)

B. amphitrite amphitrite Darwin,1854  
occasionally on ships in Scandinavia; Iberian Peninsula-southw.

Solidobalanus Hoek,1913  
\* Correct subfamily?

S. fallax (Broch,1927)  
= *Balanus fallax* Broch,1927  
\* Epizoic on upper shells of *Aequipecten opercularis*.  
W Engl. Channel, Algeria-Angola

Megabalaninae Newman,1979

Megabalanus Hoek,1913

M. tintinnabulum (Linnaeus,1758)  
occasionally on ships in Scandinavia; Mediterranean-W & S Africa

M. nigrescens (Lamarck,1818)  
occasionally on ships in Scandinavia; Australia

**RHIZOCEPHALA** F. Müller,1862

Briarosaccus ??,1???  
\* Which taxon?

B. callosus ?,1???  
@ a variety of lithodids  
cosmopolitan

KENTROGONIDA Delage,1884

Peltogastridae Lilljeborg,1861

Peltogaster Rathke,1842

P. paguri Rathke,1842 (TM - Gäsörännan, Bohuslän)  
@ *Pagurus bernhardus*, *P. cuanensis*, *Anapagurus chiroacanthus*, *A. laevis* & some foreign species  
Öresund-Bohuslän-North Sea-N Norway-Spitsbergen, W Ireland, Br. Isles-Bay of Biscay & W Mediterranean

P. curvatus Kossmann,1874  
@ *Pagurus cuanensis*, *P. prideaux* & *P. alatus*  
N Kattegatt-Bohuslän-Sogn, SW Ireland, SW Br. Isles-W Mediterranean

Peltogasterella Krüger,1912 (TM *P. socialis* Krüger,1912, non Müller,1862  
= *P. gracilis* (Boschma,1927))

P. sulcata (Lilljeborg,1859)  
= *Peltogaster sulcatus* Lilljeborg,1859  
= *Peltogaster microstoma* Lilljeborg,1859  
= *Peltogaster socialis* Müller,1862  
@ *Pagurus cuanensis*, *P. prideaux*, *P. alatus*, *Anapagurus laevis*, *A. chiroacanthus* & *A. hyndmanni*  
Kattegatt-Bohuslän-Bergen, W Ireland, W Scotland, SW Br. Isles-Canaries-Mediterranean

Galatheascus Boschma,1929

G. striatus Boschma,1929 (TM - Isle of Man)  
= *G. minutus* Boschma,1933  
@ *Galathea strigosa*, *G. intermedia*, *G. nexa* & *G. dispersa*  
Öresund-Bohuslän-Bergen, W Ireland, W Britain, Engl. Channel-W Mediterranean

Tortugaster Reinhard,1948 (TM *T. fistulatus* Reinhard,1948)

T. boschmai (Brinkmann,1936)  
= *Triangulus boschmai* Brinkmann,1936  
@ *Munida tenuimana*, *M. sarsi* (& *M. rugosa*)  
Skagerrak-W Norway

Parthenopea Kossmann,1874

P. subterranea Kossmann,1874 (TM - Gulf of Naples)  
@ *Calocarides coronatus* & *Callianassa subterranea*  
Skagerrak, Mediterranean & Black Sea

Trachelosaccus Boschma,1928

T. hymenodora (G.O. Sars,1879)  
= *Sylon hymenodora* G.O. Sars,1879 (TM - Norwegian Sea, in 3350m)  
@ *Hymenodora glacialis*

Norw. Sea

*Cyphosaccus* Reinhard, 1958 (TM *C. chacei* Reinhard, 1958)

*C. norvegicus* Boschma, 1962  
@ *Munidopsis serricornis* (Lovén)  
Bohuslän-Bergen-Trondheim-Vesterålen

*Septosaccus* Duboscq, 1912

*S. rodriguezii* (Fraisie, 1876)  
@ ?  
Roscoff

Sacculinidae Lilljeborg, 1861

*Sacculina* Thompson, 1836

*S. carcini* Thompson, 1836 (TM - England)  
@ *Portunidae* & *Pirimelidae*  
SW Baltic-Bohuslän-S North Sea-Br. Isles-Bay of Biscay-Mediterranean-Black Sea

*S. inflata* Leuckart, 1859  
= *S. lyadis* Malm, 1881  
@ *Hyas araneus* & *H. coarctatus*  
Kattegatt-Bohuslän-Helgoland

*S. triangularis* Anderson, 1862  
@ *Cancer pagurus*  
Bohuslän, British Isles-Ireland-Bretagne

*S. gibbsii* (Hesse, 1867)  
= *Sacculinidia gibbsii* Hesse, 1867  
= *Sacculina pisae* Hoek, 1888  
@ *Pisa* spp.  
Engl. Channel, Mediterranean

*S. atlantica* Boschma, 1927  
@ *Dorhynchus thomsoni*  
off SW Ireland, in 1200m

*S. lobata* Boschma, 1965  
@ *Ebalia tuberosa*  
NW Bretagne

*S. bourdoni* Boschma, 1960  
@ *Xantho pilipes*  
W Bretagne

*S. gerbei* Bonnier, 1887  
@ *Xantho incisus*, *Ethusa mascarone*, *Pilumnus hirtellus* & *Atelecycclus rotundatus*  
W Br. Isles-W Engl. Channel-Bay of Biscay-Adriatic

*S. gonoplaxae* Guérin-Ganivet, 1911  
@ *Goneplax angulata*  
Gulf of Cadiz

*S.* sp.  
\* not referable to any of the *Sacculina* spp. above  
@ *Euryome aspera*  
Väderöarna (Bohuslän)

*Drepanorthis* Boschma, 1927

*D. neglecta* (Fraisie, 1877)  
= *Sacculina neglecta* Fraisse, 1877 (TM - Gulf of Naples)  
= *Sacculina phalangi* Hoek, 1878  
= *Sacculina stenorhynchi* Malm, 1881  
@ *Macropodia* spp. & *Inachus* spp.  
Öresund-Bohuslän-Trondheim-Faeroes-S North Sea-Br. Isles-Gibraltar-Adriatic

Lernaeodiscidae Boschma, 1928

*Lernaeodiscus* F. Müller, 1862 (TM *L. porcellanae* F. Müller, 1862)

*L. ingolfi* Boschma, 1928  
@ *Munida sarsi*, (*M. tenuimana* & *M. rugosa*)  
Oslofjord-Bergen-Senja-off SW Iceland-W Br. Isles

*L. squamiferae* Pérez, 1922  
@ *Galathea squamifera*  
Norway, Canaries, French Mediterranean

*Triangulus* Smith, 1906

*T. munidae* Smith, 1906 (TM - Sætreosen, W Norway)  
= *Lernaeodiscus munidae* : Auclt.  
@ *Munida sarsi*, *M. tenuimana* & *M. rugosa*  
Bohuslän-Lofoten-Shetlands-Irish Sea-Canaries

*T. galathea* (Norman & Scott, 1906)  
= *Sacculina galathea* Norman & Scott, 1906  
= *Lernaeodiscus galathea* : Auclt.  
@ *G. intermedia*, (*G. nexa* & *G. dispersa*)  
Kattegatt-Bohuslän-Bergen-Br. Isles-Engl. Channel-Bay of Biscay-W Mediterranean

"*T. perezi*" Veillet, 194? (nomen nudum)  
@ *Galathea squamifera*  
Bretagne

AKENTROGONIDA Häfele, 1911

Clistosaccidae Boschma, 1928  
= Syloniidae Boschma, 1928

*Clistosaccus* Lilljeborg, 1860

*C. paguri* Lilljeborg, 1860 (TM - Humlesäcken, Bohuslän)  
= *Apeltes paguri* Lilljeborg, 1860  
@ *Pagurus bernhardus*, *P. pubescens* & some foreign species  
Öresund-Bohuslän-N Norway-Spitsbergen-Iceland-Faeroes-Greenland

*Sylon* Krøyer, 1855

*S. hippolytes* M. Sars, 1870 (TM - Drøbak)  
= *S. pandali* M. Sars, 1870  
= *S. schneideri* Hoek, 1888  
= *S. challengerii* Hoek, 1888  
@ Hippolytidae, Pandalidae & *Crangon almani*  
N Kattegatt-Bohuslän-N Norway-Spitsbergen-Iceland-Greenland-Shetlands

Chthamalophilidae J. Bocquet-Védrine, 1961

*Chthamalophilus* J. Bocquet-Védrine, 1957

*C. delagei* J. Bocquet-Védrine, 1957 (TM - Roscoff)  
@ *Chthamalus stellatus*  
Roscoff

*Boschmaella* J. Bocquet-Védrine, 1968  
= *Microgaster* J. Bocquet-Védrine, 1967, non Latreille, 1804, nec Swinson, 1839, nec Doumerc, 1855

*B. balani* (J. Bocquet-Védrine, 1967)  
= *Microgaster balani* J. Bocquet-Védrine, 1967 (TM - Arcachon)  
@ *Balanus improvisus*  
Arcachon

Duplorbidae

*Duplorbis* Smith, 1906

*D. calathurae* Smith, 1906 (TM - Forsblads Fjord, E Greenland)  
@ *Calathura brachiata*  
E Greenland

*Cryptogaster* ?, 19??

*Arcturosaccus* ?, 19??

Thompsoniidae Høeg & Rybakov, 1992  
\* No NE Atlantic species

*Thylacoplethus* Coutière, 1902 (TM *Thylacoplethus edwardsi* Coutière, 1902 - Australia)

*Diplothyllacus* Høeg & Lützen, 1993 (TM *Thompsonia sinensis* Keppen, 1877 - China)

*Thompsonia* Kossmann, 1874 (TM *Thompsonia globosa* Kossmann, 1874 - Philippines)

*Pottisia* Høeg & Lützen, 1993 (TM *Pottisia plesionikae* Høeg & Lützen, 1993 - Florida)

Mycetomorphidae

*Mycetomorpha* Potts, 1912

Incertae sedis

*Polysaccus* Høeg & Lützen, 1993

*P. mediterraneus* (Caroli, 1929)  
= *Thompsonia mediterranea* Caroli, 1929 (TM)  
@ *Callinassa subterranea* (Montagu) & *C. truncata* Giard & Bonnier  
Mediterranean

*Pirusaccus* Lützen, 1985

FACETOTECTA Grygier, 1984  
\* Inhabiting "Hansen's Y-larvae", puzzling youth stages from e.g. Scandinavian waters of presumed Cirripedian relatives.

ASCOTHORACIDA Lacaze-Duthiers, 1880

SYNAGOGOIDA Vagin, 1976  
@ Echinodermata, except 2 species of *Synagoga*, which occur on Anthozoa

Synagogidae Gruvel, 1905

*Ascothorax* Djakonov, 1914

*A. ophiocentis* Djakonov, 1914 (TM)  
@ bursae of *Ophiocent sericeum*  
Faeroes-Iceland-Barents Sea, Kara Sea

*A. pilocaudatus* Grygier, 1983  
@ bursae of *Ophiophalma armigerum*  
NE Atlantic

*Isidascus* Moysse, 1983

*I. bassindalei* Moysse, 1983 (TM)  
@ *Acanella arbuscula* (Johnson) (a gorgonian coral)  
NE Atlantic at 2000 m.

*Synagoga* Norman, 1888

*S. mira* Norman, 1888 (TM)  
@ *Antipathes larix* Ellis  
off Naples

Dendrogasteridae Gruvel, 1905

*Ulophysema* Brattström, 1936

*U. oeresundense* Brattström, 1936 (TM)  
@ *Brissopsis lyrifera*, *Echinocardium cordatum* & *E. flavescens*  
Öresund-Bohuslän-W Norway

*U. pourtalesiae* Brattström, 1937  
@ coelomic cavity of *Pourtalesia jeffreysi*  
Norw. Sea (between Norway, Spitsbergen, Greenland & Faeroes)

*Dendrogaster* Knipowitsch, 1891

*D. astericola* Knipowitsch, 1891 (TM)  
@ coelomic cavity of *Henricia*  
White Sea

*D. murmanensis* Kluge, 1910  
@ coelomic cavity of *Crossaster papposus* & *Solaster endeca*  
Kolafjord

*D. hymenasteri* Stone, 1987  
@ coelomic cavity of *Hymenaster membranaceus*  
NE Atlantic

*D. pontasteri* Stone, 1987  
@ coelomic cavity of *Pontaster tenuispinus*  
NE Atlantic

*D. psilasteri* Stone, 1987  
@ coelomic cavity of *Psilaster andromeda*  
NE Atlantic

*D. zoroasteri* Stone, 1987  
@ coelomic cavity of *Zoroaster fulgens*  
NE Atlantic

*D. arbusculus* Fisher, 1911  
@ coelomic cavity of *Hippasteria californica* & ?*H. phrygiana*

California & ?St Lawrence Estuary

**Myriocladus** Okada, 1925 (™ *Dendrogaster arborescens* le Roi, 1905 - S Africa @ *Dipsacaster sladeni*)

**LAURODIDA** Vagin, 1976  
@ Anthozoa

**Lauridae** Gruvel, 1905

**Laura** Lacaze-Duthiers, 1866 (short note & 1883)

**L. gerardiae** Lacaze-Duthiers, 1866 (short note & 1883) (™)  
@ in gall-like buds in the antipatharian *Gerardia lamarki* (Milne-Edwards & Haine)  
Mediterranean

**Baccalaureus** Broch, 1929 (™ *B. japonicus* Broch, 1929 - Japan @ *Zoantheria*)

**Ctenosculidae** Thiele, 1925

**Petraridae** Gruvel, 1905

**Petrarca** Fowler, 1889 (™ *P. bathyactidis* Fowler, 1889 - deep Pacific @ *Bathyactis symmetrica*)

**TANTULOCARIDA** Boxshall & Lincoln, 1983  
\* Zool. Rec. covered until December 1997.

**TANTULOCARIDIDA** Boxshall & Lincoln, 1983

**Cumonicus** Bonnier, 1903 (™ *C. kruppi* Bonnier, 1903 @ Cumacea, Leuconidae - Biscay)  
\* Which family?

**Deoterthridae** Boxshall & Lincoln, 1987, emend. Huys, 1990  
@ Primarily copepods

**Boreotantulus** Huys & Boxshall, 1988

**B. kunzi** Huys & Boxshall, 1988  
@ *Cylindropsyllus laevis*  
Bohuslän

**Tantulacus** Huys, Andersen & Kristensen, 1992

**T. hoegi** Huys, Andersen & Kristensen, 1992  
@ not known  
Faeroes

**Amphitantulus** Boxshall & Vader, 1993

**A. harpiniacheres** Boxshall & Vader, 1993 (™)  
@ *Harpinia antennaria*  
North Sea

**Deoterthron** Bradford & Hewitt, 1980 (™ *D. dentatum* Bradford & Hewitt, 1980)  
= *Austrotantulus* Boxshall, 1988 (™ *A. lincolni* Boxshall, 1988 @ *Stenhetia* - Tasmania)

**Campyloxiphos** Huys, 1990 (™ *C. dineti* Huys, 1990 @ the harpacticoid *Peresime abyssalis* - SE Atlantic)

**Dicrotrichura** Huys, 1989 (™ *D. tricincta* Huys, 1989 - Corsica)

**Coralliotantulus** Huys, 1990 (™ *C. coomansi* Huys, 1990 - Queensland)

**Aphotocentor** Huys, 1991 (™ *A. styx* Huys, 1991 - Corsica)

**Itoitantulus** Huys, Ohtsuka & Boxshall, 1992 (™ *I. misophricola* Huys, Ohtsuka & Boxshall, 1992 @ *Misophriopsis okinawensis* - Japan)

**Microdajidae** Boxshall & Lincoln, 1987

**Microdajus** Greve, 1965

**M. langi** Greve, 1965  
@ *Typhlotanais aequiremis*, *Leptognathia brevimis* & *Haplocope angusta*  
W Norway

**M. gaelicus** Boxshall & Lincoln, 1987  
@ *Typhlotanais pulcher*  
North Sea

**M. pectinatus** Boxshall, Huys & Lincoln, 1989  
@ *Typhlotanais*  
NE Atlantic

**Xenalytus** Huys, 1991 (™ *X. scotophilus* Huys, 1991 - Corsica)

**Onceroxenidae** Huys, 1990

**Onceroxenus** Boxshall & Lincoln, 1987

**O. curtus** Boxshall & Lincoln, 1987  
@ *Leptognathia zezinae*  
North Sea

**O. birdi** Boxshall & Lincoln, 1987  
@ *Paranarthura insignis*  
North Sea

**Basipodellidae** Boxshall & Lincoln, 1983, emend. Huys, 1990

**Basipodella** Becker, 1975 (™ *B. harpacticola* Becker, 1975)

**Stygotantulus** Boxshall & Huys, 1989 (™ *S. stocki* Boxshall & Huys, 1989 @ harpacticoida - Canaries)

**Nipponotantulus** Huys, Ohtsuka & Boxshall, 1994 (™ *N. heteroxenus* Huys, Ohtsuka & Boxshall, 1994 @ Pseudocyclopiidae - Japan)

**Doryphallophoridae** Huys, 1990

**Doryphallophora** Huys, 1990

**D. harrisoni** (Boxshall & Lincoln, 1987)  
= *Deoterthron harrisoni* Boxshall & Lincoln, 1987 (™)  
@ *Macrostylis magnifica*  
North Sea

**COPEPODA** H. Milne-Edwards, 1840

**PROGYMNOPLA** Lang, 1948

**PLATYCOPIOIDA** Fosshagen, 1985

**Platycopiidae** G.O. Sars, 1911

\* Bottom dwellers

**Platycofia** G.O. Sars, 1911

**P. perplexa** G.O. Sars, 1911 (™)  
S Norway (Korshavn)

**P. pygmaea** G.O. Sars, 1921  
S Norway (Korshavn)

**NEOCOPEPODA** Huys & Boxshall, 1991

**GYMNOPLA** Giesbrecht, 1882

**CALANOIDA** G.O. Sars, 1903

**Megacalanoida** Sewell, 1947

**Calanidae** Dana, 1849

**Calanus** Leach, 1816  
= *Cetochilus* Claus, 1863

**C. hyperboreus** Krøyer, 1838  
Kattegat-Bohuslän-North Sea-W Norway-Norw. Sea-Iceland-Faeroes-Shetlands-Ireland

**C. helgolandicus** (Claus, 1863)  
= *Cetochilus helgolandicus* Claus, 1863  
Kattegat-Bohuslän-North Sea-W Norway-(Spitsbergen)-Iceland-Faeroes-Shetlands-Br. Isles-Mediterranean-Black Sea-Azores-Canaries

**C. finmarchicus** (Gunnerus, 1765)  
= *Monoculus finmarchicus* (™)  
Baltic-Bohuslän-North Sea-W Norway-Norw. Sea-Arctic Seas-Iceland-Faeroes-Shetlands-Br. Isles-N Portugal-NE Atlantic

**C. glacialis** Jaschnov, 1955  
N Norway-Faeroes-Iceland-Arctic Seas

**C. minor** Claus, 1863 (™ of *Namocalanus* G.O. Sars, 1925)  
= *Calanus valgus* Brady, 1883  
Cold & temperate Atlantic, Mediterranean

**C. propinquus** Brady, 1883  
?NE Atlantic

**Calanoides** Brady, 1883

**C. carinatus** (Krøyer, 1849)  
= *Calanus brevicornis* Lubbock, 1856  
\* Atlantic deep water species  
N Atlantic

**Mesocalanus** Bradford & Jillet, 1974

**M. tenuicornis** (Dana, 1849)  
= *Calanus tenuicornis* Dana, 1849  
= *Neocalanus tenuicornis* G.O. Sars, 1925  
North Sea-Faeroes-Shetlands-Ireland-NE Atlantic-Mediterranean

**Neocalanus** G.O. Sars, 1925

**N. gracilis** (Dana, 1849)  
= *Calanus gracilis* Dana, 1849  
?S North Sea, Ireland-Biscay, Mediterranean

**N. robustior** (Giesbrecht, 1888)  
= *C. robustior* Giesbrecht, 1888  
Coald & temperate Atlantic, Mediterranean

**Megacalanidae** Sewell, 1947

**Megacalanus** Wolfenden, 1904

**M. princeps** Wolfenden, 1904  
= *Macrocalanus longicornis* G.O. Sars, 1905  
Iceland-Faeroes-Ireland-NE Atlantic

**Bathycalanus** G.O. Sars, 1905

**B. princeps** (Brady, 1883)  
Iceland-Faeroes-Shetlands-Ireland-NE Atlantic

**B. bradyi** (Wolfenden, 1905)  
Iceland-Faeroes-Ireland-NE Atlantic

**B. richardi** G.O. Sars, 1905  
\* Is not this and *B. bradyi* (Wolfenden) synonyms? If this is the case, which of them has priority?  
NE Atlantic

**Bradycalanus** A. Scott, 1909

**B. sarsi** (Farran, 1939)  
NE Atlantic

**Paracalanidae** Giesbrecht, 1892

**Paracalanus** Boeck, 1864

**P. parvus** (Claus, 1863)  
Baltic-Bohuslän-North Sea-W Norway-Norw. Sea-Iceland-Faeroes-Shetlands-Br. Isles-Engl. Channel-NE Atlantic

**P. pygmaeus** (Claus, 1863)  
\* May be a synonym of *P. parvus* (Claus)  
NE Atlantic

**P. nanus** G.O. Sars, 1907  
?NE Atlantic

**Calocalanidae** Bernard, 1958

**Calocalanus** Giesbrecht, 1888

**C. contractus** Farran, 1926  
NE Atlantic-Biscay-Mediterranean

**C. pavo** (Dana, 1849)  
Atlantic 60° N-40° S

**C. pseudocontractus** Bernard, 1958  
S & W Ireland, NE Atlantic, Mediterranean

**C. styliremis** Giesbrecht, 1888  
Worldwide distribution except polar sea

- Ischnocalanus* Bernard,1963
- I. plumulosus* (Claus,1863)  
Temperate-tropical Atlantic, Mediterranean
- Mecynoceridae Andronov,1973
- Mecynocera* I.C. Thompson,1888
- M. clausii* I.C. Thompson,1888  
\* Oceanic epiplankton  
NE Atlantic
- Eucalanoidea Giesbrecht,1892
- Eucalanidae Giesbrecht,1892
- Eucalanus* Dana,1853 (<sup>TM</sup>*E. attenuatus* (Dana,1849) - W. Pacific)
- E. elongatus* (Dana,1849)  
\* According to Fleminger,1973,*E. elongatus* (Dana) is a tropical Indian Ocean to W Pacific species, while *E. hyalinus* (Claus,1866) is tropical-subtropical circumglobal, i.e. the only species in the *elongatus* group, thought to be found also in the Atlantic. They may perhaps have been mixed up earlier?  
Skagerrak-N North Sea-Norw. Sea-Iceland-Faeroes-Ireland-NE Atlantic
- E. crassus* Giesbrecht,1888  
S & W Ireland, NE Atlantic
- E. sewelli* Fleminger,1973  
= *E. attenuatus* : Auctt., non (Dana,1849)  
\* According to Fleminger,1973 this is the only N Atlantic species of the *attenuatus* group.  
S & W Ireland, NE Atlantic
- Rhincalanus* Dana,1853
- R. nasutus* Giesbrecht,1888  
Koster Channel, off Väderöbod, Skagerrak, N North Sea-W Norway-Norw. Sea-?Barents Sea-Iceland-Faeroes-Scotland-Ireland-NE Atlantic
- R. cornutus* (Dana,1849)  
?NE Atlantic
- Clausocalanoidea Giesbrecht,1892
- Spinocalanidae Vervoort,1951
- Spinocalanus* Giesbrecht,1888 (<sup>TM</sup>*S. abyssalis* Giesbrecht,1888 - abyssal Pacific)
- S. longicornis* G.O. Sars,1900  
= *S. abyssalis* : G.O. Sars,1903, non Giesbrecht,1888  
= *S. schaudinni* Mrazek,1902  
non = *S. longicornis* : G.O. Sars,1903  
Central Arctic Ocean
- S. brevicaudatus* Brodsky,1950  
= *S. longicornis* : G.O. Sars,1903, non G.O. Sars,1900  
N North Sea-W Norway-Norw. Sea-Barents Sea-Iceland-Faeroes-Shetlands-Ireland-NE Atlantic
- S. magnus* Wolfenden,1904  
= *S. latifrons* G.O. Sars,1907  
Norw. Sea-Barents Sea-Iceland-Ireland-NE Atlantic
- S. angusticeps* G.O. Sars,1920  
\* Oceanic deep water species  
S & W Ireland, NE Atlantic
- S. spinosus* Farran,1908  
\* Oceanic deep water species  
S & W Ireland, NE Atlantic
- Drepanopsis* Wolfenden,1911  
= *Farrania* G.O. Sars,1920
- D. frigidus* Wolfenden,1911 (<sup>TM</sup>)  
\* Oceanic deep water species. As *D. oblongus* has been considered a synonym of this species, the distributions are somewhat mixed up.  
?Biscay, NE Atlantic
- D. oblongus* (G.O. Sars,1920)  
= *Farrania oblonga* G.O. Sars,1920 (<sup>TM</sup> of *Farrania* G.O. Sars,1920)  
NE Atlantic
- Mimocalanus* Farran,1908
- M. cultrifer* Farran,1908  
\* Oceanic deep water species  
S & W Ireland, NE Atlantic
- M. nudus* Farran,1908  
\* Oceanic deep water species  
S & W Ireland, NE Atlantic
- Monacilla* G.O. Sars,1905
- M. tenera* G.O. Sars,1907  
\* Oceanic deep water species  
S & W Ireland, NE Atlantic
- M. typica* G.O. Sars,1905  
\* Oceanic deep water species  
S & W Ireland
- Aetideidae Giesbrecht,1892
- Bradyetes* Farran,1905
- B. matthei* Johannessen,1976  
Bergen (bottom-living)
- B. brevis* Farran,1936  
\* Living close to bottom in moderately deep water  
Barents Sea, E Greenland
- B. inermis* Farran,1905  
\* Living close to bottom in moderately deep water  
S & W Ireland, NE Atlantic
- Chiridiella* G.O. Sars,1907
- C. macrodactyla* G.O. Sars,1907 (<sup>TM</sup>)  
\* Deep water of temperate Atlantic  
S & W Ireland, NE Atlantic
- C. reducta* Brodsky,1950  
Central part of deep Arctic Ocean
- Aetideinae Giesbrecht,1892
- Gaidius* Giesbrecht,1895
- G. tenuispinus* (G.O. Sars,1900)  
= *Chiridius tenuispinus* G.O. Sars,1900  
N North Sea-Norw. Sea-Barents Sea-Iceland-Faeroes-Shetlands-Ireland-NE Atlantic
- G. brevispinus* (G.O. Sars,1900)  
= *Chiridius brevispinus* G.O. Sars,1900  
N North Sea-Norw. Sea-Arctic Seas-Iceland-Faeroes-Ireland-NE Atlantic
- G. affinis* G.O. Sars,1905  
S & W Ireland, NE Atlantic
- G. minutus* G.O. Sars,1905  
NE Atlantic
- G. pungens* Giesbrecht,1895 (<sup>TM</sup> - Pacific)  
N Atlantic
- Gaetanus* Giesbrecht,1888
- G. armiger* Giesbrecht,1888  
N North Sea-Norw. Sea-Barents Sea-Iceland-Faeroes-Shetlands-NE Atlantic
- G. miles* Giesbrecht,1888 (<sup>TM</sup>)  
N North Sea-Faeroes-Shetlands-Ireland-NE Atlantic
- G. latifrons* G.O. Sars,1905  
N North Sea-Ireland-NE Atlantic
- G. minor* Farran,1905  
Iceland-Faeroes-Ireland-NE Atlantic
- G. kruppii* Giesbrecht,1903  
Iceland-Faeroes-Shetlands-Ireland-NE Atlantic
- G. brachyurus* G.O. Sars,1907  
Iceland-Faeroes-NE Atlantic
- G. pileatus* Farran,1903  
\* May be a synonym of *G. caudani* Canu  
Iceland-Faeroes-Ireland-NE Atlantic
- G. curvicornis* G.O. Sars,1905  
\* Atlantic & Arctic deep water  
NE Atlantic
- G. caudani* Canu,1896  
Biscay
- Aetideopsis* G.O. Sars,1903  
= *Faroella* Wolfenden,1904
- A. rostrata* G.O. Sars,1903 (<sup>TM</sup>)  
Norw. Sea-Barents Sea-Iceland-Faeroes-Shetlands-off W Greenland-NE Atlantic
- A. multiserrata* (Wolfenden,1904)  
= *Faroella multiserrata* Wolfenden,1904  
Iceland-Faeroes-Ireland-NE Atlantic-central Norwegian Sea
- Undinopsis* G.O. Sars,1884  
= *Bradyidius* Giesbrecht,1897
- U. similis* G.O. Sars,1903  
\* hyperbenthic  
W Norway-Norw. Sea-NE Atlantic
- U. luluae* (Grize,1972)  
Bergen (bottom-living)
- U. bradyi* G.O. Sars,1884 (<sup>TM</sup>)  
= *U. armatus* (Brady,1878), non (Boeck,1872)  
\* hyperbenthic  
Oslofjord-N North Sea-Bergen (bottom-living)-Aalesund-Norw. Sea-Barents Sea-W Br. Isles-NE Atlantic
- Aetideus* Brady,1883
- A. armatus* (Boeck,1872)  
= *Pseudocalanus armatus* Boeck,1872  
Oslofjord-North Sea-Bergen (bottom-living)-Norw. Sea-?Barents Sea-Iceland-Faeroes-Shetlands-W Br. Isles-NE Atlantic
- Pseudaeetideus* Wolfenden,1903
- P. armatus* (Boeck,1872)  
= *Euchaeta armata* Boeck,1872 (<sup>TM</sup>)  
= *Chiridius armatus* : Auctt.  
Bohuslän-N North Sea-Bergen (bottom-living)-Norw. Sea-Barents Sea-Iceland-Faeroes-Shetlands-W Br. Isles-NE Atlantic
- Chiridius* Giesbrecht,1892
- C. obtusifrons* G.O. Sars,1903  
= *C. armatus* G.O. Sars,1900, non Boeck,1872  
N Norw. Sea-Barents Sea-Iceland-Faeroes-NE Atlantic
- C. tenuispinus* G.O. Sars,18??  
Ofotenfjord, Arctic Seas
- C. poppei* Giesbrecht,1892  
S & W Ireland, NE Atlantic, Mediterranean
- C. gracilis* Farran,1908  
S & W Ireland, NE Atlantic
- Comantenna* Wilson,1924  
= *Bryaxis* G.O. Sars,1903, ex Boeck MS, non Kugelann,1794
- C. brevicornis* G.O. Sars,1903, ex Boeck MS  
\* hyperbenthic  
N North Sea-Bergen-Barents Sea-Ireland-NE Atlantic
- C. minor* (Farran,1905)  
= *Bryaxis minor* Farran,1905  
\* hyperbenthic in deep water  
S & W Ireland, NE Atlantic
- Euaetideus* G.O. Sars,1925

- E. giesbrechti* (Cleve,1904)  
Faeroes-Shetlands-Ireland-NE Atlantic
- Derjuginia* Jaschnov,1947
- D. tolli* (Linko,1913)  
= *Scolecithrix tolli* Linko,1913  
All marginal seas of the Polar Basin
- Pseudogaetanus* Brodsky,1950
- P. robustus* (G.O. Sars,1905)  
off Ireland Azores
- Euchirellinae Brodsky,1950
- Euchirella* Giesbrecht,1888
- E. messinensis* (Claus,1863)  
= *Undina messinensis* Claus,1863  
?N North Sea-Norw. Sea-Iceland-Faeroes-Ireland-NE Atlantic- Mediterranean
- E. rostrata* (Claus,1863)  
= *Undina rostrata* Claus,1863  
= *Euchaeta hessei* Brady,1883  
N North Sea-Norw. Sea-Iceland-Faeroes-Shetlands-Ireland-NE Atlantic-Mediterranean
- E. bitumida* With,1915  
?N North Sea-Norw. Sea-Iceland-Faeroes-NE Atlantic
- E. curticauda* Giesbrecht,1888  
N North Sea-Iceland-Faeroes-Ireland-Biscay-NE Atlantic
- E. maxima* Wolfenden,1905  
Iceland-Faeroes-Ireland-NE Atlantic
- E. pulchra* (Lubbock,1856)  
?Iceland-?Faeroes-Azores-NE Atlantic
- E. brevis* G.O. Sars,1905  
Biscay-NE Atlantic
- E. intermedia* With,1915  
\* Deep water species  
NE Atlantic
- Pseudochirella* G.O. Sars,1920
- P. spectabilis* (G.O. Sars,1900)  
= *Undeuchaeta spectabilis* G.O. Sars,1900  
Central part of Arctic Seas
- P. notacantha* (G.O. Sars,1905)  
= *Gaidius notacanthus* G.O. Sars,1905  
?Iceland-?Faeroes-Ireland-NE Atlantic
- P. pustulifera* (G.O. Sars,1905)  
= *Undeuchaeta pustulifera* G.O. Sars,1905  
Iceland-Faeroes-Ireland-NE Atlantic
- P. superba* (With,1915)  
= *Undeuchaeta superba* With,1915  
Iceland-Faeroes-Ireland-NE Atlantic
- P. dubia* (G.O. Sars,1905)  
= *Undeuchaeta dubia* G.O. Sars,1905  
?Iceland-?Faeroes-NE Atlantic
- P. cryptospina* (G.O. Sars,1905)  
= *Gaidius cryptospinus* G.O. Sars,1905  
\* Temperate & N Atlantic deep water species  
S & W Ireland, NE Atlantic
- P. obtusa* (G.O. Sars,1905)  
= *Undeuchaeta obtusa* G.O. Sars,1905  
\* Temperate Atlantic deep water species  
S & W Ireland, NE Atlantic
- P. divaricata* (G.O. Sars,1905)  
= *Gaidius divaricanus* G.O. Sars,1905  
Biscay, temperate Atlantic
- P. scopularis* (G.O. Sars,1905)  
= *Undeuchaeta scopularis* G.O. Sars,1905  
Temperate Atlantic
- Chirundina* Giesbrecht,1895
- C. streetsii* Giesbrecht,1895 (TM)  
Iceland-Faeroes-Shetlands-Ireland-Biscay-NE Atlantic
- Undeuchaeta* Giesbrecht,1888
- U. plumosa* (Lubbock,1856)  
= *Undina plumosa* Lubbock,1856  
Norw. Sea-Iceland-Faeroes-Ireland-E Engl. Channel-NE Atlantic
- U. major* Giesbrecht,1888 (TM)  
Norw. Sea-Iceland-Faeroes-Ireland-?E Engl. Channel-Biscay-NE Atlantic
- Pseudeuchaeta* G.O. Sars,1905
- P. brevicauda* G.O. Sars,1905  
\* Temperate Atlantic deep water species  
S & W Ireland, NE Atlantic
- Valdiviella* Steuer,1904 (TM V. *oligartha* Steuer,1904)
- V. insignis* Farran,1908  
Gibraltar, Canaries, temperate Atlantic
- V. brevicornis* G.O. Sars,1905  
temperate deep Atlantic
- Euchaetidae Giesbrecht,1892
- Pareuchaeta* A. Scott,1909  
= *Euchaeta* : Auctt.
- P. norvegica* (Boeck,1872)  
= *Euchaeta norvegica* Boeck,1872 (TM)  
Bohuslän-W Norway-Arctic Seas-Iceland-Scotland
- P. glacialis* (Hansen,1886)  
N Norw. Sea-Iceland-Arctic Seas
- P. scotti* (Farran,1909)  
Iceland-W Ireland-NE Atlantic
- P. acuta* (Giesbrecht,1892)  
= *Euchaeta acuta* Giesbrecht,1892  
N Atlantic, Mediterranean
- P. bisinuata* (G.O. Sars,1907)  
Biscay, temperate Atlantic, Azores, Canaries
- P. bradyi* (With,1915)  
= *Euchaeta Bradyi* With,1915  
Biscay, N & temperate Atlantic
- P. gracilis* (G.O. Sars,1905)  
= *Euchaeta quadrata* Farran,1909  
off Ireland, Biscay, N Atlantic
- P. hanseni* (With,1915)  
= *Euchaeta Hanseni* With,1915  
N & temperate Atlantic
- P. hebes* (Giesbrecht,1888)  
= *Euchaeta hebes* Giesbrecht,1888  
N Atlantic, Gibraltar, Mediterranean
- P. spinosa* (Giesbrecht,1892)  
= *Euchaeta spinosa* Giesbrecht,1892  
Temperate Atlantic, Mediterranean
- P. tonsa* (Giesbrecht,1895)  
= *Euchaeta tonsa* Giesbrecht,1895  
Biscay, N & temperate Atlantic
- P. sarsi* (Farran,1909)  
Biscay, N Atlantic
- P. pubera* (G.O. Sars,1907)  
= *Euchaeta pubera* G.O. Sars,1907  
= *Euchaeta marina* : Wolfenden,1907, non (Prestandrea,1833)  
= *Euchaeta Wolfendeni* A Scott,1907  
Portugal, temperate Atlantic
- P. rubicunda* (Farran,1908)  
= *Euchaeta rubicunda* Farran,1908  
off Ireland, temperate Atlantic
- P. media* (Giesbrecht,1888)  
= *Euchaeta media* Giesbrecht,1888  
off Ireland & Portugal, temperate Atlantic
- P. polaris* Brodsky,1950  
Central part of Arctic Ocean
- Euchaeta* Philippi,1843
- E. barbata barbata* Brady,1883  
\* Should perhaps be moved to *Pareuchaeta* A. Scott,1909  
SW Norw. Sea, Faeroes, S. Iceland, Rockall, Atlantic W off central Europe
- E. barbata farrani* With,1915  
\* Should perhaps be moved to *Pareuchaeta* A. Scott,1909  
W Norway-off Lofoten-Iceland-W Greenland-Barents Sea-Arctic Seas
- E. marina* (Prestandrea,1833)  
= *Cyclops marinus* Prestandrea,1833 (TM)  
temperate Atlantic, Mediterranean
- Phaennidae G.O. Sars,1902
- Cornucalanus* Wolfenden,1905
- C. chelififer* (J.C. Thompson,1903)  
= *Scolecithrix chelififer* J.C. Thompson,1903  
= *Cornucalanus magnus* Wolfenden,1905  
off W Ireland, off SW Faeroes, Iceland-Faeroe Channel, off S Iceland, Denmark Strait
- Onchocalanus* G.O. Sars,1905
- O. magnus* (Wolfenden,1906)  
= *Xanthocalanus magnus* Wolfenden,1906  
= *Onchocalanus frigidus* Wolfenden,1911  
Arctic Seas
- O. cristatus* (Wolfenden,1904)  
= *Xanthocalanus cristatus* Wolfenden,1904  
= *Onchocalanus trigoniceps* G.O. Sars,1905  
=? *Xanthocalanus similis* Esterly,1906  
off W Ireland, off SW Faeroes, Iceland-Faeroe Channel, off S Iceland, Denmark Strait
- O. hirtipes* G.O. Sars,1905  
off W & NW Ireland
- O. affinis* With,1915  
Temperate & boreal Atlantic, Azores
- O. trigoniceps* G.O. Sars,1905 (TM)  
= *Xanthocalanus magnus* Wolfenden,1909, non Wolfenden,1906  
Biscay, off Portugal, Mediterranean
- Oothrix* Farran,1905
- O. borealis* Wiborg,19??  
W Norwegian fjords
- Pseudophaenna* G.O. Sars,1903
- P. typica* G.O. Sars,1903  
Oslofjord-N Norway
- Talacalanus* Wolfenden,1911
- T. grenii* (Farran,1905)  
= *Xanthocalanus grenii* Farran,1905  
=? *X. calaminus* Wolfenden,1906  
\* Is correct spelling greeni ?  
off S Iceland, off W Ireland, Biscay
- Xanthocalanus* Giesbrecht,1892
- X. minor* Giesbrecht,1892  
= *X. fragilis* Aurivillius,1898  
Koster area (between Hällsöarna), S Norway-Bergen (bottom-living)-Trondheim

- X. fragilis* Aurivillius, 1898  
Skagerrak
- X. simplex* Aurivillius, 1898  
Skagerrak
- X. borealis* G.O. Sars, 1900  
W Norway-Kara Sea-Greenland Sea-Arctic Seas-Shetlands, Biscay, off Azores
- X. propinquus* G.O. Sars, 1903  
W-N Norway
- X. fallax* G.O. Sars, 1919  
Bergen (bottom-living), off W Ireland
- X. pinguis* Farran, 1905  
off W Norway, off W Ireland
- X. obtusus* Farran, 1905  
off W Ireland
- X. subagilis* Wolfenden, 1904  
off W Ireland
- X. muticus* G.O. Sars, 1905  
Biscay, Azores
- X. agilis* Giesbrecht, 1892 (TM)  
Mediterranean
- X. polaris* Brodsky, 1950  
Abyssal Arctic Ocean
- Cephalophanes* G.O. Sars, 1907
- C. refulgens* G.O. Sars, 1907 (TM - deep Atlantic)  
off W Ireland & Britain - off W Spain & Africa
- Amalophora* T. Scott, 1894
- A. typica* (T. Scott, 1892)  
off W Ireland
- Heteremalla* G.O. Sars, 1907
- H. dubia* (T. Scott, 1894)  
= *Amalophora dubia* T. Scott, 1894  
Temperate Atlantic, Mediterranean
- Brachycalanus* Farran, 1905
- B. atlanticus* (Wolfenden, 1904)  
= *Xanthocalanus atlanticus* Wolfenden, 1904  
off W Ireland
- Oothrix* Farran, 1905
- O. bidentata* Farran, 1905  
off W Ireland
- Phaenna* Claus, 1863
- P. spinifera* Claus, 1863  
=? *P. zelandica* T. Scott, 1909  
off Svinyø (North of Cape Stadt), ?Shetlands, Atlantic, Mediterranean
- Scolecithricidae Giesbrecht, 1892
- Scottocalanus* G.O. Sars, 1905
- S. securifrons* (T. Scott, 1894)  
= *Scolecithrix securifrons* T. Scott, 1894  
= *Scottocalanus acutus* G.O. Sars, 1905  
Iceland, off W Ireland, Biscay, Canaries, Azores
- S. persecanus* (Giesbrecht, 1892)  
= *Scolecithrix persecanus* Giesbrecht, 1892 (TM)  
= *Scottocalanus Helenae* Scott, 1909  
= *Scottocalanus Thorii* With, 1915  
off W Ireland, Biscay, off Gibraltar, Canaries, Azores
- Scolecithricella* G.O. Sars, 1903
- S. minor* (Brady, 1883) v. *occidentalis* Brodsky, 1950  
= *Scolecithrix minor* Brady, 1883 (TM)  
S Skagerrak - off Väderöbod-Oslofjord-W Norway-Lofoten-Iceland
- S. ovata* (Farran, 1905)  
= *Scolecithrix ovata* Farran, 1905  
off W Ireland, off E & W Greenland
- S. auropecten* (Giesbrecht, 1892)  
= *Scolecithrix auropecten* Giesbrecht, 1892  
off W Ireland, Biscay, Mediterranean
- S. dentata* (Giesbrecht, 1892)  
= *Scolecithrix dentata* Giesbrecht, 1892  
off W Ireland, Biscay, Mediterranean
- S. vittata* (Giesbrecht, 1892)  
= *Scolecithrix vittata* Giesbrecht, 1892  
Biscay, Mediterranean
- S. abyssalis* (Giesbrecht, 1892)  
= *Scolecithrix abyssalis* Giesbrecht, 1892  
off Faeroes, Temperate Atlantic, Mediterranean
- Scaphocalanus* G.O. Sars, 1900  
= *Amalophora* G.O. Sars, 1903, non Scott, 1894
- S. magnus* (Scott, 1894)  
= *Amalophora magna* T. Scott, 1894 (TM)  
= *Scolecithrix cristata* Giesbrecht, 1895  
= *Scaphocalanus acrocephalus* G.O. Sars, 1903  
= *Amalophora magna* : G.O. Sars, 1903  
W Norway-Arctic Seas-Iceland
- S. brevicornis* (G.O. Sars, 1900)  
= *Scolecithrix brevicornis* G.O. Sars, 1900  
= *Scaphocalanus gracilipes* Farran, 1908  
N Norw. Sea-Arctic Seas-Shetlands, Biscay
- S. curtus* (Farran, 1926)  
= *Scolecithrix curta* Farran, 1926
- off W Ireland, Biscay
- S. echinatus* (Farran, 1905)  
off W Ireland, Biscay
- S. robustus* (T. Scott, 1893)  
= *Scolecithrix robusta* T. Scott, 1893  
off W Ireland
- S. medius* (G.O. Sars, 1907)  
= *Amalophora media* G.O. Sars, 1907  
Azores, Temperate Atlantic
- S. affinis* (G.O. Sars, 1905)  
= *Amalophora affinis* G.O. Sars, 1905  
Biscay, off Gibraltar, Canaries, Azores
- S. angulifrons* G.O. Sars, 1920  
off Lisboa
- S. polaris* Brodsky, 1950  
Central part of abyssal Arctic Ocean
- Scolecithrix* Brady, 1883
- S. danae* (Lubbock, 1856)  
= *Undina Danae* Lubbock, 1856 (TM)  
Atlantic, Mediterranean
- S. bradyi* Giesbrecht, 1888  
Atlantic, Mediterranean
- Lophothrix* Giesbrecht, 1895
- L. insignis* G.O. Sars, 1920  
off NW France
- L. humilifrons* G.O. Sars, 1905  
temperate Atlantic
- L. frontalis* Giesbrecht, 1895 (TM)  
off W Ireland, Biscay
- L. latipes* (T. Scott, 1893)  
= *Scolecithrix latipes* T. Scott, 1893  
= *Scolecithrix angusta* Esterly, 1911  
= *Scolecithrix acuta* Wolfenden, 1911  
Biscay, temperate Atlantic
- Amalothrix* G.O. Sars, 1925
- A. arcuata* (G.O. Sars, 1920)  
= *Scolecithricella arcuata* G.O. Sars, 1920  
off W Scotland
- A. gracilis* (G.O. Sars, 1905)  
= *Scolecithricella gracilis* G.O. Sars, 1905 (TM)  
= *Scolecithrix globiceps* Farran, 1909  
off W Ireland, off NW France, off Lisboa, Azores
- A. emarginata* (Farran, 1905)  
off W Ireland, Biscay, off Gibraltar, Canaries, Azores
- A. valida* (Farran, 1909)  
= *Scolecithrix valida* Farran, 1909  
off W Ireland
- A. valens* (Farran, 1926)  
= *Scolecithrix valens* Farran, 1926  
off W Ireland, Biscay
- A. laminata* (Farran, 1926)  
= *Scolecithrix laminata* Farran, 1926  
off W Ireland, Biscay
- A. lobata* (G.O. Sars, 1920)  
= *Scolecithricella lobata* G.O. Sars, 1920  
off NW France, Biscay
- A. falcifer* (Farran, 1926)  
= *Scolecithrix falcifer* Farran, 1926  
Biscay
- A. propinqua* (G.O. Sars, 1920)  
= *Scolecithricella propinqua* G.O. Sars, 1920  
off Lisboa
- A. curticauda* (A. Scott, 1909)  
= *Scolecithricella curticauda* A. Scott, 1909  
off Gibraltar, Canaries
- A. obtusifrons* (G.O. Sars, 1905)  
= *Scolecithricella Tydemani* Scott, 1909  
Biscay, off Gibraltar, Canaries, Azores
- Diaixidae G.O. Sars, 1902
- Diaixis* G.O. Sars, 1902
- D. hibernica* (A. Scott, 1896)  
= *Scolecithrix hibernica* A. Scott, 1896  
Oslofjord-W Norway-Trondheim
- D. pygmaea* (T. Scott, 1899)  
= *Scolecithrix pygmaea* T. Scott, 1899  
Scotland, Ireland, Mediterranean
- Mesaiokeratidae Matthews, 1961
- Mesaiokeras* Matthews, 1961
- M. nanseni* Matthews, 1961  
Bergen-N Norway (bottom-living)
- Stephidae G.O. Sars, 1902
- Stephos* T. Scott, 1892
- S. lamellatus* G.O. Sars, 1903  
Christiansund-N Norway
- S. scotti* G.O. Sars, 1903  
Kieler Bucht-Kattegatt-W Norway-Scotland

- S. rustadi* Strömgen, 1969  
Bergen
- S. minor* (T. Scott, 1892)  
S Norway-Scotland
- S. fultoni* (T. & A. Scott, 1898)  
Scotland
- Parastephos* G.O. Sars, 1903
- P. pallidus* G.O. Sars, 1903  
S-W Norway-Scotland
- Tharybidae* G.O. Sars, 1902
- Undinella* G.O. Sars, 1905  
= *Paratharybia* Tanaka, 1937
- U. oblonga* (G.O. Sars, 1900) (TM)  
off Greenland, Arctic Seas, off Ireland
- U. simplex* (Wolfenden, 1906)  
= *Xanthocalanus simplex* Wolfenden, 1906, non Aurivillius, 1898  
= *Undinella brevipes* Farran, 1906  
off W Ireland, Biscay
- Tharybis* G.O. Sars, 1903
- T. macrophthalma* G.O. Sars, 1903  
Oslofjord-W Norway-Trondheim
- Pseudocyclopiidae* G.O. Sars, 1902
- Pseudocyclopia* T. Scott, 1892
- P. stephoides* J.C. Thompson, 1895  
Norway (Christiansund)-Liverpool Bay
- P. crassicornis* T. Scott, 1892  
S-W Norway-Scotland
- P. giesbrechti* Wolfenden, 1902  
S-W Norway-Shetlands-Scotland
- P. caudata* T. Scott, 1894  
Aberdeen-Firth of Forth
- P. minor* T. Scott, 1892  
Firth of Forth
- Clausocalanidae* Giesbrecht, 1892  
= *Pseudocalanidae* G.O. Sars, 1903
- Clausocalanus* Giesbrecht, 1888
- C. arcuicornis* (Dana, 1849)  
W Norway, Norw. Sea, S & W Ireland, Irish Sea, W Engl. Channel
- C. pargens* Farran, 1926  
\* Probably a form of *C. arcuicornis* (Dana)  
S & W Ireland, NE Atlantic
- C. paululus* Farran, 1926  
\* Probably a form of *C. arcuicornis* (Dana)  
S & W Ireland, NE Atlantic
- Pseudocalanus* Boeck, 1872
- P. elongatus* (Boeck, 1865)  
= *Clausia elongata* Boeck, 1865  
= *Calanus clausii* Brady, 1865  
Bothnian Sea-Kattegatt-Bohuslän-North Sea-N Norway-Arctic Seas-Engl. Channel
- P. minutus* (Krøyer, 1845)  
= *P. gracilis* G.O. Sars, 1903  
Faeroes-Iceland-N Norway-N Norw. Sea-Arctic Seas
- P. acuspes* (Giesbrecht, 1881)  
= *Lucillus acuspes* Giesbrecht, 1881  
Baltic, Katt, Bohus-N Norway-Arctic Seas
- P. major* G.O. Sars, 1900  
Arctic Seas, ?Kara Sea, ?Spitsbergen
- Microcalanus* G.O. Sars, 1901
- M. pygmaeus* (G.O. Sars, 1900)  
= *Pseudocalanus pygmaeus* G.O. Sars, 1900  
North Sea-W Norway-Arctic Seas
- M. pusillus* G.O. Sars, 1903  
\* Sometimes considered to be a form of *M. pygmaeus* (G.O. Sars)  
Kattegatt-North Sea-W Norway
- Drepanopus* Brady, 1883
- D. bungei* G.O. Sars, 1898  
Barents Sea, Kara Sea, Laptev Sea
- Ctenocalanus* Giesbrecht, 1888
- C. vanus* Giesbrecht, 1888  
S & W Ireland, NE Atlantic
- Centropagoidea* Giesbrecht, 1892
- Temoridae* Giesbrecht, 1892
- Temora* Baird, 1850
- T. longicornis* (O.F. Müller, 1785)  
= *Cyclops longicornis* O.F. Müller, 1785  
Finland-Kattegatt-Bohuslän-North Sea-W & N Norway-Iceland-Engl. Channel-Mediterranean
- T. stylifera* (Dana, 1849)  
= *Calanus stylifer* Dana, 1849  
= *Temora armata* Claus, 1863  
Temperate & chilly Atlantic, Mediterranean
- Eurytemora* Giesbrecht, 1881  
= *Temorella* Claus, 1881
- E. velox* (Liljeborg, 1853)
- Brackish water-limnic; e.g. Baltic, S Norwegian & Scottish rock-pools
- E. affinis* (Poppe, 1880)  
= *E. hirundo* Giesbrecht, 1881 (TM)  
= *E. hirundoides* Nordquist, 1888  
\* See Buch & Brenning, 1992. *Crustaceana* 62: 13-38 regarding synonymy.  
Brackish water; river mouths in Finland, E Sweden, Skåne, Isefjord, Oslofjord-N Norway, Zuidoostzee, Germany, Br. Isles
- E. americana* L.W. Williams, 1906  
= ? *E. thompsoni* Willey, 1923  
\* Estuarine species living in freshened waters  
?Europe, Woods Hole area
- Temoropia* T. Scott, 1893
- T. mayumbaensis* T. Scott, 1893  
off W Ireland, Biscay, temperate & chilly Atlantic
- Centropagidae* Giesbrecht, 1892
- Limnocalanus* G.O. Sars, 1863
- L. macrurus* G.O. Sars, 1863 *grimaldi* (de Guerne, 1886)  
= *Centropages grimaldi* de Guerne, 1886  
\* *L. macrurus* is a freshwater species in Sw., Norw., Finl., etc. and *L. grimaldi* a brackish water form in Baltic, Caspian & Arctic Seas.  
Baltic, E Greenland, Spitsbergen, Kara Sea, Laptev Sea
- Isias* Boeck, 1865
- I. clavipes* Boeck, 1865  
Kattegatt-Gullmarfjord-Hafstensund-Oslofjord-W Norway-Trondheim-Br. Isles-Mediterranean
- Centropages* Krøyer, 1849
- C. typicus* Krøyer, 1849 (TM)  
Åland-Kattegatt-Bohuslän-Trondheim-North Sea-Iceland-Faeroes-Shetlands-Br. Isles-E Atlantic
- C. hamatus* (Liljeborg, 1853)  
Bothnian Sea-Kattegatt-Bohuslän-North Sea-W Norway-Barents Sea-Iceland-Faeroes-Shetlands-Br. Isles-E Atlantic
- C. bradyi* Wheeler, 1899  
S & W Ireland, temperate Atlantic, Mediterranean
- C. violaceus* (Claus, 1863)  
S & W Ireland
- C. chierchiai* Giesbrecht, 1889  
Warm water part of NE Atlantic
- C. kroeyeri* Giesbrecht, 1892  
?NE Atlantic
- Candaciidae* Giesbrecht, 1892
- Candacia* Dana, 1846
- C. norvegica* (Boeck, 1865)  
Oslofjord-Skagerrak-W Norway-N North Sea-Lofoten-Iceland-Faeroes-Shetlands-Ireland
- C. armata* (Boeck, 1865)  
Oslofjord-Skagerrak-North Sea-W Norway-Faeroes-Shetlands-W Br. Isles-Engl. Channel
- C. elongata* (Boeck, 1872)  
= *C. rotundata* (Wolfenden, 1904)  
= *C. obtusa* G.O. Sars, 1905  
Norw. Sea-Iceland-Faeroes-Shetlands-Ireland
- C. tenuimana* (Giesbrecht, 1889)  
\* Oceanic epiplankton  
S & W Ireland, NE Atlantic
- C. varicans* (Giesbrecht, 1892)  
\* Oceanic epiplankton  
S & W Ireland, NE Atlantic
- C. longimana* (Claus, 1863)  
\* Oceanic epiplankton  
NE Atlantic
- Paracandacia* Grice, 1963
- P. bispinosa* (Claus, 1863)  
= *Candacia bispinosa* Claus, 1863  
Temperate Atlantic, Mediterranean
- P. simplex* (Giesbrecht, 1892)  
= *Candacia simplex* Giesbrecht, 1892  
Temperate Atlantic, Mediterranean
- Pontellidae* Dana, 1853
- Anomalocera* Templeton, 1837
- A. patersoni* Templeton, 1837  
Kattegatt-Bohuslän-North Sea-W Norway-Iceland-Br. Isles- Mediterranean-Black Sea
- Labidocera* Lubbock, 1853
- L. wollastoni* (Lubbock, 1857)  
= *Pontella wollastoni* Lubbock, 1857  
= *Pontella helgolandica* Claus, 1863  
Måseskär, S & E Skagerrak-Oslofjord-Bergen-North Sea-Engl. Channel-Mediterranean
- L. kroeyeri* (Brady, 1883)  
= *Pontella kroeyeri* Brady, 1883  
off W Ireland
- L. acutifrons* (Dana, 1849)  
= *Pontella acutifrons* Dana, 1849  
= *Pontia Edwardsii* Krøyer, 1849  
Temperate Atlantic, Mediterranean
- L. brunescens* (Czerniavsky, 1868)  
= *Pontella brunescens* Czerniavsky, 1868  
Temperate Atlantic, Mediterranean
- L. nerii* (Krøyer, 1849)  
= *Pontia Nerii* Krøyer, 1849  
off Cape Finisterre
- Pontella* Dana, 1846

- P. atlantica* (H. Milne-Edwards, 1840)  
= *Pontia atlantica* H. Milne-Edwards, 1840  
Temperate Atlantic, Mediterranean
- P. lobiancoi* (Canu, 1888)  
= *Pontellina Lo Biancoi* Canu, 1888  
Engl. Channel, Mediterranean
- P. mediterranea* (Claus, 1863)  
= *Pontellina mediterranea* Claus, 1863  
Mediterranean, Black Sea
- Pontellina* Dana, 1852  
= *Calanops* Claus, 1863
- P. plumata* (Dana, 1849)  
= *Calanops messinensis* Claus, 1863  
Temperate Atlantic, Azores, Mediterranean
- Pontellopsis* Brady, 1883
- P. regalis* (Dana, 1849)  
= *Pontella regalis* Dana, 1849  
Temperate Atlantic, Mediterranean
- P. villosa* Brady, 1883  
Temperate Atlantic, Mediterranean
- Parapontellidae Giesbrecht, 1892
- Parapontella* Brady, 1878
- P. brevicornis* (Lubbock, 1857)  
W Norway-Br. Isles-France-Mediterranean
- Acartiidae G.O. Sars, 1903
- Acartia* Dana, 1846  
= *Dias* Liljeborg, 1853
- A. (Acanthacartia) bifilosa* Giesbrecht, 1881  
Bothnian Sea-Bohuslän-North Sea-Iceland-Engl. Channel
- A. (Acanthacartia) tonsa* Dana, 1848  
Bay of Finland-Belt Sea, S North Sea-E Engl. Channel
- A. (Acartiura) Steuer*, 1915) *clausii* Giesbrecht, 1889  
Belt Sea-Bohuslän-North Sea-W Norway-Iceland-Faeroes-Shetlands-W Br. Isles-W Engl. Channel-Mediterran.
- A. (Acartiura) longiremis* (Liljeborg, 1853)  
Bay of Finland-Bohuslän-Engl. Channel-North Sea-W & N Norway-Barents Sea-Iceland-Faeroes-Shetlands
- A. (Acartiura) discoidata* (Giesbrecht, 1881)  
Belt Sea-Kattegatt, N North Sea-W Norway-W Br. Isles-Engl. Channel
- A. (Paracartia) T. Scott*, 1894) *grani* (G.O. Sars, 1904)  
= *Paracartia grani* G.O. Sars, 1904  
W Norway
- A. (Paracartia) latisetosa* (Kricagin, 1873)  
Mediterranean, Black Sea
- A. lefevreae* Bradford, 1976  
?NE Atlantic
- Diaptomidae Baird, 1850
- Pseudodiaptomidae G.O. Sars, 1902
- Sulcanidae Nicholls, 1945
- Tortanidae G.O. Sars, 1902
- Bathypontioidea Brodsky, 1950
- Bathypontiididae Brodsky, 1950
- Sognocalanus* Fosshagen, 1967
- S. confertus* Fosshagen, 1967  
Sognefjord & Hardangerfjord
- Damkaeria* Fosshagen, 1983
- D. falcifera* Fosshagen, 1983 (TM)  
Norw. Sea (bottom-living, deep)
- Bathypontia* G.O. Sars, 1905
- B. elongata* G.O. Sars, 1905 (TM)  
off W Ireland, Biscay, off Portugal
- Temorites* G.O. Sars, 1900  
\* Previously included in Temoridae
- T. brevis* G.O. Sars, 1900 (TM)  
= *Temoropsis simplex* Wolfenden, 1911  
Arctic Seas, Mediterranean
- Augaptiloidea G.O. Sars, 1905
- Metridinidae G.O. Sars, 1902
- Metridia* Boeck, 1865
- M. lucens* Boeck, 1865  
= *Paracalanus hibernicus* Giesbrecht, 1892  
Kattegatt-Bohuslän-German Bight-North Sea-Lofoten-Iceland-Faeroes-Shetlands-Br. Isles-Engl. Channel, Biscay, Mediterranean
- M. longa* (Lubbock, 1854)  
= *Calanus longus* Lubbock, 1854  
= *Metridia armata* Boeck, 1865 (TM)  
Kattegatt-Bohuslän-North Sea-N Norway-Norw. Sea-Arctic Seas-Iceland-Faeroes-Shetlands-Ireland
- M. princeps* Giesbrecht, 1892  
Norw. Sea-Iceland-Faeroes-Ireland, Biscay
- M. venusta* Giesbrecht, 1892  
Iceland-Faeroes-Shetlands-Ireland
- M. brevicauda* Giesbrecht, 1889
- Biscay, E Atlantic
- M. macrura* G.O. Sars, 1905  
E Atlantic
- Pleuromamma* Giesbrecht, 1898
- P. robusta* (F. Dahl, 1893)  
W Norway-Norw. Sea-Iceland-Faeroes-Shetlands-Ireland-W Engl. Channel
- P. abdominalis* (Lubbock, 1856)  
= *Diaptomus abdominalis* Lubbock, 1856 (TM)  
W Norway, Iceland-Faeroes-Shetlands-Ireland
- P. xiphias* (Giesbrecht, 1889)  
= *Pleuromma xiphias* Giesbrecht, 1889  
Iceland-Faeroes-Shetlands-Ireland-W Engl. Channel
- P. borealis* (F. Dahl, 1893)  
= *Pleuromma borealis* F. Dahl, 1893  
Faeroes-Shetlands-Ireland
- P. gracilis* (Claus, 1863)  
= *Pleuromma gracilis* Claus, 1863  
S & W Ireland, E Atlantic
- P. piseki* Farran, 1929  
?NE Atlantic
- Heterorhabdidae G.O. Sars, 1902
- Heterorhabdus* Giesbrecht, 1898  
= *Heterochaeta* Claus, 1862, non Westwood, 1843 (Orthoptera), nee Claparède, 1863 (Oligochaeta)
- H. (Paraheterorhabdus) Brodsky*, 1950) *compactus* (G.O. Sars, 1900)  
= *Heterochaeta compacta* G.O. Sars, 1900  
off Ireland, off W Greenland, Biscay, Arctic Seas-N Atlantic
- H. (Paraheterorhabdus) robustus* Farran, 1909  
Iceland-Faeroes-Ireland
- H. (Paraheterorhabdus) vipera* (Giesbrecht, 1889)  
= *Heterochaeta vipera* Giesbrecht, 1889  
E Atlantic
- H. (Euheterorhabdus) Brodsky*, 1950) *norvegicus* (Boeck, 1872)  
= *Heterochaeta profunda* Dahl, 1894  
= *Heterochaeta norvegica* : G.O. Sars, 1900  
Bohuslän (fairly common in deepest Koster Channel)-N North Sea-Lofoten-Iceland-Faeroes-Shetlands-Ireland, Greenland, Arctic Seas
- H. (Euheterorhabdus) abyssalis* (Giesbrecht, 1889)  
N North Sea-Iceland-Faeroes-Shetlands-Ireland
- H. (Euheterorhabdus) spinifrons* (Claus, 1863)  
= *Heterochaeta spinifrons* Claus, 1863 (TM)  
Norw. Sea, S & W Ireland, E Atlantic, Mediterranean
- H. (Euheterorhabdus) papilliger* (Claus, 1863)  
= *Heterochaeta papilligera* Claus, 1863  
S & W Ireland, E Atlantic, Mediterranean
- H. (Euheterorhabdus) clausii* (Giesbrecht, 1889)  
= *Heterochaeta clausii* Giesbrecht, 1889  
Biscay, temperate Atlantic
- Heterostylites* G.O. Sars, 1920
- H. longicornis* (Giesbrecht, 1892)  
= *Heterochaeta longicornis* Giesbrecht, 1892  
Iceland-Faeroes-Shetlands-Ireland
- H. major* (F. Dahl, 1894)  
= *Heterochaeta major* F. Dahl, 1894  
= *Alloiorhabdus medius* Wolfenden, 1911  
Norw. Sea-Iceland-Faeroes-Ireland, Biscay, Azores
- Hemirhabdus* Wolfenden, 1911
- H. grimaldii* (Richard, 1893)  
\* Oceanic deep water species  
S & W Ireland, NE Atlantic
- Disseta* Giesbrecht, 1889
- D. palumbai* Giesbrecht, 1889 (TM)  
\* Oceanic deep water species  
S & W Ireland, Biscay, NE Atlantic
- Mesorhabdus* G.O. Sars, 1905
- M. angustus* G.O. Sars, 1907  
?NE Atlantic
- M. brevicaudatus* (Wolfenden, 1905)  
= *Heterorhabdus brevicaudatus* Wolfenden, 1905  
\* Oceanic deep water species  
S & W Ireland, NE Atlantic
- Augaptilidae G.O. Sars, 1905
- Augaptilus* Giesbrecht, 1892
- A. glacialis* G.O. Sars, 1900  
= *Augaptilus zetesios* Wolfenden, 1902  
Shetlands-Faeroes-Iceland-Greenland-Spitsbergen-Arctic Seas-NE Atlantic
- A. longicaudatus* (Claus, 1863)  
= *Hemicalanus longicaudatus* Claus, 1863 (TM)  
off W Ireland, Biscay, temperate & boreal Atlantic, Mediterranean
- A. megalurus* Giesbrecht, 1892  
off W Ireland, temperate Atlantic
- A. anceps* Farran, 1909  
off W Ireland, Mediterranean
- A. spinifrons* G.O. Sars, 1907  
Temperate Atlantic, off Gibraltar
- Augaptilina* G.O. Sars, 1920
- A. scopifera* G.O. Sars, 1920  
Biscay

- Centraugaptilus* G.O. Sars,1920
- C. cucullatus* (G.O. Sars,1905)  
off W Ireland, Biscay, Azores, Canaries, temperate Atlantic
- C. horridus* (Farran,1908)  
off W Ireland, temperate Atlantic
- C. rattrayi* (T. Scott,1893) (TM)  
off W Ireland, Biscay, Azores, Canaries, temperate Atlantic
- Euaugaptilus* G.O. Sars,1920
- E. affinis* G.O. Sars,1920  
Biscay, temperate Atlantic
- E. angustus* (G.O. Sars,1905)  
= *Augaptilus angustus* G.O. Sars,1905  
Canaries, Azores, temperate Atlantic, Mediterranean
- E. bullifer* (Giesbrecht,1892)  
= *Augaptilus bullifer* Giesbrecht,1892  
off W Ireland, Biscay, temperate Atlantic, Azores, Canaries
- E. clavatus* (G.O. Sars,1907)  
= *Augaptilus clavatus* G.O. Sars,1927  
off Portugal, Azores, temperate Atlantic
- E. digitatus* G.O. Sars,1920  
off W Bretagne
- E. elongatus* (G.O. Sars,1905)  
= *Augaptilus elongatus* G.O. Sars,1905  
~~non~~ = *E. elongatus* : Sewell,1947  
off W Ireland, Iceland, Biscay, temperate Atlantic, Azores, Canaries
- E. facilis* (Farran,1908)  
= *Augaptilus facilis* Farran,1908  
off W Ireland, temperate Atlantic, Canaries
- E. filigerus* (Claus,1863)  
= *Hemicalanus filigerus* Claus,1863  
Canaries, Azores, temperate Atlantic, Mediterranean
- E. gibbus* (Wolfenden,1904)  
= *Augaptilus gibbus* Wolfenden,1904  
Biscay, off Gibraltar, Azores, temperate Atlantic
- E. hecticus* (Giesbrecht,1889)  
= *Augaptilus hecticus* Giesbrecht,1889  
Biscay, temperate Atlantic, Mediterranean
- E. humilis* Farran,1926  
Biscay, temperate Atlantic
- E. laticeps* (G.O. Sars,1905)  
= *Augaptilus laticeps* G.O. Sars,1905  
off W Ireland, Biscay, temperate Atlantic, Azores, Canaries, Mediterranean
- E. magnus* (Wolfenden,1904)  
Engl. Channel, off W Ireland, Biscay, temperate Atlantic
- E. nodifrons* (G.O. Sars,1905)  
= *Augaptilus nodifrons* G.O. Sars,1905  
off W Ireland, Biscay, temperate Atlantic, Azores, Canaries
- E. oblongus* (G.O. Sars,1905)  
= *Augaptilus oblongus* G.O. Sars,1905  
Biscay, temperate Atlantic, Azores, Canaries
- E. palumboi* (Giesbrecht,1892)  
= *Augaptilus Palumbii* Giesbrecht,1892  
Biscay, temperate Atlantic
- E. penicillatus* G.O. Sars,1920  
off Gibraltar, temperate Atlantic
- E. similis* (Farran,1908)  
off W Ireland, temperate Atlantic
- E. squamatus* (Giesbrecht,1889)  
= *Augaptilus squamatus* Giesbrecht,1889  
= *Augaptilus brevicaudatus* G.O. Sars,1905  
off W Ireland, off Portugal, temperate Atlantic, Azores, Canaries
- E. truncatus* (G.O. Sars,1905)  
= *Augaptilus truncatus* G.O. Sars,1905  
off W Ireland, temperate Atlantic, Azores, Canaries
- E. hyperboreus* Brodsky,1950  
Abyssal central Arctic Ocean
- Haloptilus* Giesbrecht,1898
- H. longicornis* (Claus,1863)  
= *Hemicalanus longicornis* Claus,1863 (TM)  
Between Finnmark & Bear Island-Bay of Biscay-Mediterranean
- H. acutifrons* (Giesbrecht,1892)  
= *Hemicalanus acutifrons* Giesbrecht,1892  
= *Haloptilus spinifrons* G.O. Sars,1900  
? North Sea-Between Jan Mayen & Norway-Iceland-Biscay-Mediterranean, Greenland
- H. fons* Farran,1908  
off W Ireland, temperate Atlantic
- H. tenuis* Farran,1908  
off W Ireland, temperate Atlantic, Mediterranean
- H. angusticeps* G.O. Sars,1907  
Biscay, temperate Atlantic, Mediterranean
- H. bulliceps* Farran,1905  
Biscay, temperate Atlantic
- H. plumosus* (Claus,1863)  
= *Hemicalanus plumosus* Claus,1863  
Biscay, temperate Atlantic, Mediterranean
- H. mucronatus* (Claus,1863)  
= *Hemicalanus mucronatus* Claus,1863
- Temperate Atlantic, Mediterranean
- H. ornatus* (Giesbrecht,1892)  
= *Hemicalanus ornatus* Giesbrecht,1892  
Temperate Atlantic, Mediterranean
- H. spiniceps* (Giesbrecht,1892)  
= *Hemicalanus spiniceps* Giesbrecht,1892  
Temperate Atlantic, Mediterranean
- H. validus* G.O. Sars,1920  
Temperate Atlantic, Mediterranean
- Heteroptilus* G.O. Sars,1920
- H. acutilobus* (G.O. Sars,1905)  
= *Pontoptilus acutilobus* G.O. Sars,1905  
Biscay, Canaries, Azores, temperate Atlantic
- H. attenuatus* (G.O. Sars,1905)  
= *Pontoptilus attenuatus* G.O. Sars,1905  
Biscay, Canaries, Azores, temperate Atlantic
- Pontoptilus* G.O. Sars,1905
- P. muticus* G.O. Sars,1905  
off W Ireland, temperate Atlantic
- Pachyptilus* G.O. Sars,1920
- P. abbreviatus* (G.O. Sars,1905)  
= *Pontoptilus abbreviatus* G.O. Sars,1905  
Biscay, Canaries, Azores, temperate Atlantic
- P. erygnathus* G.O. Sars,1920  
off Lisboa, Azores, temperate Atlantic
- Pseudaugaptilus* G.O. Sars,1907 (TM *P. longiremis* G.O. Sars,1907 - off Azores)
- P. polaris* Brodsky,1950  
Abyssal central Arctic Ocean
- Discoidea Gordejeva,1975
- Disco* Grice & Hulsemann,1965
- D. fiordicus* Fosshagen,1967  
Hardangerfjord
- Arietellidae G.O. Sars,1902
- Arietellinae G.O. Sars,1902
- Paramisophria* T. Scott,1897
- P. cluthae* T. Scott,1897  
W Norway-Scotland-Mediterranean
- P. spooneri* Krishnaswamy,1959  
SW Br. Isles (shell gravel living)
- Scottula* G.O. Sars,1903
- S. inaequicornis* G.O. Sars,1903  
Oslofjord-W Norway
- S. abyssalis* G.O. Sars,1905  
off Lisboa, Azores, temperate Atlantic
- Arietellus* Giesbrecht,1892
- A. pavoninus* G.O. Sars,1905  
off W Ireland, Azores
- A. plumifer* G.O. Sars,1905  
off W Ireland, off Portugal, Canaries, Azores
- A. simplex* G.O. Sars,1905  
off W Iceland, W Greenland, off W Ireland, Azores
- A. giesbrechti* G.O. Sars,1905  
off Lisboa & Gibraltar, Canaries, temperate Atlantic
- A. setosus* Giesbrecht,1892 (TM)  
NE? Atlantic, Mediterranean
- Paraugaptilus* Wolfenden,1904
- P. buchani* Wolfenden,1904 (TM)  
off W Ireland, Canaries, Azores, temperate Atlantic
- Epacteriscidae Fosshagen,1973
- Lucicutiidae G.O. Sars,1902
- Lucicutia* Giesbrecht, in Giesbrecht & Schmeil,1898 (n. cons., Op. 1613, ICZN)  
= *Leuckartia* Claus,1862, ~~non~~ Agassiz,1862 (Coelenterata)  
= *Rudolfia* Wilson,1924
- L. atlantica* Wolfenden,1904  
= *Lucicutia magna* Wolfenden,1920  
= *Lucicutia gracilis* G.O. Sars,1905  
Iceland, off W Ireland, Biscay, Azores, temperate Atlantic, Biscay
- L. curta* Farran,1905  
off W Ireland, Biscay, northern & temperate Atlantic
- L. anomala* Brodsky,1950  
Arctic Seas
- L. polaris* Brodsky,1950  
Arctic Seas
- L. pseudopolaris* Heptner,1969  
Arctic Seas
- L. grandis* (Giesbrecht,1895)  
= *Leuckartia grandis* Giesbrecht,1895  
Engl. Channel, W Ireland, N & temperate Atlantic
- L. lucida* Farran,1908  
= *Lucicutia pera* A. Scott,1909  
off W Ireland, temperate Atlantic, Mediterranean

- L. macrocera* G.O. Sars,1920  
off W Bretagne, temperate Atlantic
- L. maxima* (Steuer,1904)  
= *Leuckartia maxima* Steuer,1904  
off W Bretagne, off Lisboa, temperate Atlantic
- L. simulans* G.O. Sars,1920  
Biscay, temperate Atlantic, Mediterranean
- L. longiserrata* (Giesbrecht,1892)  
= *Leuckartia longiserrata* Giesbrecht,1892  
Gibraltar, Canaries, temperate Atlantic, Mediterranean
- L. clausi* (Giesbrecht,1889)  
= *Leuckartia clausi* Giesbrecht,1889  
Atlantic, Mediterranean
- L. flavicornis* (Claus,1863)  
= *Leuckartia flavicornis* Claus,1863 (TM)  
Atlantic, Mediterranean
- L. gemina* Farran,1926  
Biscay, temperate Atlantic
- L. longicornis* (Giesbrecht,1889)  
= *Leuckartia longicornis* Giesbrecht,1889  
Biscay
- L. ovalis* Wolfenden,1911  
Biscay, temperate & tropical Atlantic
- L. tenuicauda* G.O. Sars,1907  
off Lisboa, Canaries, temperate Atlantic
- L. polaris* Brodsky,1950  
Central part of Arctic Ocean
- L. anomala* Brodsky,1950  
Central part of Arctic Ocean
- Isochaeta* Giesbrecht,1898
- I. ovalis* Giesbrecht,1898 (TM)  
Biscay, temperate Atlantic
- Phylloporidae Brodsky,1950
- Phyllopus* Brady,1883
- P. bidentatus* Brady,1883 (TM)  
off Ireland
- P. helgae* Farran,1908  
off Ireland & Gibraltar, Biscay, Azores, temperate Atlantic
- P. impar* Farran,1908  
off Ireland & Gibraltar, Azores, temperate Atlantic
- P. aequalis* G.O. Sars,1920  
off Gibraltar, temperate Atlantic
- Pseudocyclopoidea Giesbrecht,1893
- Pseudocycloporidae Giesbrecht,1893
- Pseudocyclops* Brady,1876
- P. crassiremis* Brady,1873  
S Norway (Korshavn)-Br. Isles
- P. obtusatus* Brady & Robertson,1873  
Christiansund-Scotland-Arctic Seas
- Ridgewayidae M.S. Wilson,1958
- Ridgewayia* Thompson & Scott,1903
- Exumella* Fosshagen,1970
- Brattstromia* Fosshagen, in Fosshagen & Iliffe,1991
- Placocalanus* Ohtsuka, Fosshagen & Soh,1996 (TM *P. insularis* Fosshagen,1970 - Bahamas)  
= *Placocalanus* Fosshagen,1970 (nom. nud.)
- Ryocalanoidea Andronov,1974
- Ryocalanidae Andronov,1974  
\* No species within the area?
- PODOPLEA Giesbrecht,1882
- CYCLOPOIDA Burmeister,1834
- GNATHOSTOMATINA Thorell,1860
- Oithonidae Dana,1853
- Paraoithona* Farran,1908
- P. parvula* Farran,1908  
W Norway-W Ireland
- Oithona* Baird,1843  
\* The nomenclature within this genus should be checked more.
- O. atlantica* Farran,1908  
= *O. spinirostris* : G.O. Sars,1918  
=? *O. spinirostris* Claus,1863  
=? *O. plumifera* Baird,1843  
\* Should probably be named *O. spinirostris* Claus,1863, while Baird's species may be another.  
Kattegatt-Skagerrak-Trondheim-Norw. Sea-Iceland-Br. Isles-Mediterranean
- O. minuta* Kritchaguine,1873  
= *O. nana* Giesbrecht,1892  
Kattegatt-Skagerrak-S North Sea
- O. helgolandica* Claus,1863  
= *O. similis* Claus,1863  
\* Has been considered a synonym of both *O. nana* & *O. similis*  
Åland-Isefjord-Bohuslän, Oslofjord-N Norway-Iceland-Arctic Seas-S North Sea-Yorkshire-Engl. Channel-Mediterranean, Atlantic Canada
- O. linearis* Giesbrecht,1891  
Temperate Atlantic, Adriatic
- O. setigera* (Dana,1849)  
= *Oithona tropica* Wolfenden,1905  
= *Oithona pelagica* Farran,1908  
Biscay, Cape Verde, tropical & temperate Atlantic, Mediterranean
- O. robusta* Giesbrecht,1891  
Cape Verde
- Cyclopinidae G.O. Sars,1913
- Pterinopsyllinae Kiefer,1927-28
- Pterinopsyllus* Brady,1880
- P. insignis* (Brady,1878)  
Oslofjord-Skagerrak-Norw. Sea-Trondheim-Br. Isles-Mediterranean
- Cyclopininae G.O. Sars,1913
- Cyclopetta* G.O. Sars,1918
- C. difficilis* G.O. Sars,1918  
S Norway (Risør)-North Sea
- Cyclopinula* Strand,1929  
= *Cyclopina* Claus,1862, non Agassiz,1846
- C. schneideri* (T. Scott,1903)  
= *C. brevifurca* (G.O. Sars,1918)  
Bohuslän-N Norway-Spitsbergen-Arctic Seas-Canada-?Br. Isles
- C. pygmaea* (G.O. Sars,1918)  
= *Cyclopina pygmaea* G.O. Sars,1918  
North Sea-S & SW Norway-Br. Isles-Mediterranean
- C. brachystylis* (G.O. Sars,1921)  
= *Cyclopina brachystylis* G.O. Sars,1921  
Norway-SE North Sea
- C. norvegica* (Boeck,1865)  
= *Cyclopina norvegica* Boeck,1865  
\* Sensu Sars, G.O. 1918 is this a synonyme of *C. gracilis* (Claus)  
S Baltic-W Sweden-North Sea-N Norway-Arctic Seas-Br. Isles-?Mediterranean
- C. gracilis* (Claus,1862)  
= *Cyclopina gracilis* Claus,1862  
S Baltic-W Sweden-North Sea-Norway-Br. Isles-Mediterranean-Black Sea
- C. kieferi* (Schäfer,1936)  
SE North Sea, Mediterranean
- C. latipes* (Schäfer,1936)  
SE North Sea
- C. rotundipes* (Herbst,1952)  
= *Cyclopina rotundipes* Herbst,1952  
Sylt
- ? *C. phallusiae* (Hansen,1923)  
= *Cyclopina phallusiae* Hansen,1925  
@ *Ascidia obliqua*  
Faeroes
- C. esilis* (Brian,1958)  
= *Cyclopina esilis* Brian,1958  
?Britain
- C. tuberculata* (Herbst,1962)  
= *Cyclopina tuberculata* Herbst,1962  
?North Sea area
- Cyclopidina* Steuer,1940
- C. euacantha* (G.O. Sars,1918)  
Norway
- Cyclopinella* G.O. Sars,1918
- C. tumidula* G.O. Sars,1918  
S Norway (Risør & Lillesand)-North Sea-Trondheim-Norw. Sea
- Cyclopinodes* Wilson,1932
- C. elegans* (Scott,1894)  
Bohuslän-SW Norway (Farsund)-North Sea-Br. Isles-Italy
- Cyclopinoides* Lindberg,1953
- C. longicornis* (Boeck,1872)  
Skagerrak-S North Sea-N Norway-Arctic Seas-Br. Isles-Mediterranean
- C. dilatata* (G.O. Sars,1921)  
North Sea-S Norway (Korshavn)
- C. littoralis* (Brady,1872)  
Bohuslän-North Sea-W Norway-Br. Isles-France-Italy
- Hemicyclopinia* Herbst,1953
- H. setifera* (Herbst,1952)  
Sylt
- Metacyclopinia* Lindberg,1953
- M. harpacticoidea* (Klie,1949)  
Helgoland
- M. brevisetosa* Herbst,1974  
Sylt
- M. roscoffensis* Bozic,1953  
Roscoff
- Microcyclopinia* Plesa,1961
- Herbstina* Huys & Boxshall,1990
- H. exigua* (Herbst,1974)  
= *Microcyclopinia exigua* (Herbst,1974) (TM)  
Sylt
- Neocyclopinia* Herbst,1952

*N. reducta* Herbst, 1952  
Sylt, Amrum

*Pseudocyclopinodes* Lang, 1946

*P. (Parapseudocyclopinodes)* Lindberg, 1961 *neglecta* (Herbst, 1952)  
Sylt

*P. (Parapseudocyclopinodes)* Lindberg, 1961 *dacunhai* Lindberg, 1961  
W Portugal

*Smirnovipina* Arbizu, 1997

*S. barentsiana* (Smirnov, 1931) Arbizu, 1997  
= *Cyclopina barentsiana* Smirnov, 1931 (TM)  
= *Cyclopinodes barentsiana* (Smirnov, 1931) Smirnov, 1935  
off NE Spitsbergen in 197 m

*Erythropolites* Huys & Boxshall, 1990 (TM *Microcyclopina semireducta* Herbst, 1964)

*Glareolina* Huys & Boxshall, 1990 (TM *Hemicyclopina kliei* Herbst, 1953)

*Cyclopidae* Dana, 1853

*Halicyclopinæ* Kiefer, 1927

*Euryte* Philippi, 1843

*E. longicauda* Philippi, 1843  
Bohuslän-Oslofjord-N Norway-Faeroes-Br. Isles-Mediterranean-Black Sea

*E. minor* (T. Scott, 1905)  
\* possibly a subspecies of *E. longicauda* Philippi  
Oslofjord-North Sea-W Norway-Norw. Sea-Scotland-Plymouth

*E. robusta* Giesbrecht, 1901?  
S Norway (Korshavn & Kleven)-Raunefjorden-Christiansund, Mediterranean

*E. curticornis* G.O. Sars, 1918  
Lofoten, Arctic Seas

*Halicyclops* Norman, 1903

*H. magniceps* (Liljeborg, 1853)  
= *H. christianseni* (Boeck, 1872?)  
= *H. aequoreus* (Fischer, 1860)  
S Finland-Oresund-S & W Norway, Br. Isles-France-Mediterranean

*H. neglectus* Kiefer, 1936?  
= *H. aequoreus* : (Brady, 1878), non (Fischer, 1860)  
Baltic-North Sea

*H. neglectus septentrionalis* ?Kiefer, 1936?  
= *H. thermophilus septentrionalis* Kiefer, 1936?  
Baltic-North Sea

*H. brevispinosus brevispinosus* Herbst, 1952  
Sylt

*H. incognitus* Herbst, 1962  
?North Sea area

*Neocyclops* Gurney, 1927

*N. remanei* (Herbst, 1952)  
Sylt

*Diacyclops* Kiefer, 1927

*D. nanus* (G.O. Sars, 1862)  
?NE Atlantic

*Cyclopinae* Dana, 1853  
\* Freshwater species

*Megacyclops* Kiefer, 1927

*M. viridis* (Jurine, 1820)  
= *Cyclops vulgaris* Koch, 1835  
Fresh water-Bothnian Sea

*Mesocyclops* G.O. Sars, 1918

*M. leuckarti* (Claus, 1862?)  
= *Mesocyclops obsoletus* G.O. Sars 1918, ?non (Koch, 1835)  
Fresh water-Bothnian Sea

*Thermocyclops* Kiefer, 1927

*T. oithonoides* (G.O. Sars, 1862)  
Fresh water-Bothnian Sea

NOTODELPHYOINA

*Chordeumiidae* Boxshall, 1988

*Chordeumium* Stephensen, 1918  
= *Chordeuma* Jungersen, 1912, non Koch, 1847

*C. obesum* (Jungersen, 1912)  
@ In gonad galls of *Asteronyx loveni*  
Skagerrak

*Parachordeumium* le Calvez, 1938  
= *Ophioithys* Heegard, 1951  
= *Amphiurophilus* Delamere Deboutteville, 1962

*P. tetracerus* le Calvez, 1938  
@ genital cavity & coelom of *Axiognathus squamatus*  
\* Not mentioned by Gotto, 1993  
Roscoff-French Mediterranean

*P. amphiuuræ* (Hérouard, 1906)  
= *Philichthys amphiuuræ* Hérouard, 1906  
= *Ophioithys amphiuuræ* (Hérouard, 1906)  
= *Amphiurophilus amphiuuræ* : Aucct.  
@ bursae (genital sacs) of *Axiognathus squamatus*  
Bohuslän, Bergen, Engl. Channel, Isle of Man, Roscoff, ?E USA

*P. bocqueti* Goudey-Perrière, 1979  
@ bursae of *Axiognathus squamatus*  
\* Not mentioned by Gotto, 1993  
NE Atlantic

*Ophioika* Stephensen, 1934

*O. appendiculata* Stephensen, 1935  
@ in bursae of *Ophiomitrella clavigera* & *Ophiacantha bidentata*  
\* Not mentioned by Gotto, 1993  
W Hebrides-S Iceland-S Greenland

*Arthrochordeumium* Stephensen, 1918  
\* W Pacific

*Ophioicodes* Heegard, 1951

*Lernaeosaccus* Heegard, 1951 (TM *L. ophiacanthæ* Heegard, 1951)

*Thespesiopsyllidae* Wilson, 1924  
= *Thaumatopsyllidae* G.O. Sars, 1913  
\* Probably more related to Cancerillidae than close to Monstrillidae, where they formerly were put, sensu Fossahagen, 1970, but Huys & Boxshall 1991 removed the family to the Cyclopoida. McKinnon, 1994 is of the opinion that Wilson, 1924 erroneously changed the genus name of our genus.

*Thespesiopsyllus* Wilson, 1924  
= *Thaumatopsyllus* G.O. Sars, 1913, non T. Scott, 1894

*T. paradoxus* (G.O. Sars, 1913) (TM)  
@ nauplius in stomach folds of *Ophiotrix fragilis*, *Ophiopholis aculeata* & *Ophiura albida*  
\* 50% of Gullmarfjord *Ophiopholis* population infested, according to Bresciani & Lützen, 1962  
Bohuslän-Oslofjord-Trondheim

*Cucumaricoidae* Bouligand & Delamare-Deboutteville, 1959

*Cucumaricola* Paterson, 1958

*Notodelphyoidea* Dana, 1853

*Notodelphyidae* Dana, 1853  
@ Ascidians, Octocorals  
\* Some genera should perhaps go to a family of their own, Doropygidae Brady, 1878

*Notodelphys* Allman, 1847

*N. allmani* Thorell, 1859  
= *N. ascidicola* Allman, 1847 (p.p.) (TM)  
@ *Ascidia* spp., *Ascidietta* spp., *Phallusia* spp. & *Ciona* spp.  
Isefjord, Bohuslän-N Norway-Br. Isles-Mediterranean-Black Sea-S Africa

*N. rufescens* Thorell, 1859  
@ *Ascidia conchilega*, *A. obliqua*, *Ascidietta aspersa* & *A. scabra*  
Bohuslän-S Norway-North Sea-Britain-N Ireland-Faeroes-Mediterranean

*N. caerulea* Thorell, 1859  
@ *Ascidia virginea*, ?*A. mentula*, *Ciona intestinalis* & *Corella parallelogramma*  
Isefjord, Bohuslän-S Norway-Shetlands-Br. Isles

*N. agilis* Thorell, 1859  
@ *Ascidia* spp., *Ascidietta* spp., *Molgula* spp., *Polycarpa* spp., *Ciona intestinalis* & *Corella parallelogramma*  
Kattegat-Bohuslän-N Norway-Br. Isles-Mediterranean

*N. tenera* Thorell, 1859  
@ *Ascidia mentula*, *A. obliqua*, *Ciona intestinalis* & *Phallusia mammilata*  
Kattegat-Bohuslän-NW Norway-Mediterranean

*N. elegans* Thorell, 1859  
@ *Ciona intestinalis* & *Ascidietta aspersa*  
Isefjord, Bohuslän-S Norway-Br. Isles-Mediterranean

*N. prasina* Thorell, 1859  
= *N. pusilla* Buchholz, 1869  
@ *Ascidia mentula*, *Ascidietta aspersa*, *Phallusia mammilata* & *Ciona intestinalis*  
Bohuslän-S Norway-Trondheim-Shetlands-Br. Isles-Mediterranean

*N. dentata* Schellenberg, 1921  
@ *Rhopalaea nordgaardii* Hartm.  
Norway

*N. canui* Roland, 1962  
@ *Molgula echinosiphonica*  
Engl. Channel

*N. cryptopyge* Bocquet & Stock, 1960  
@ *Pyura savignyi*  
Engl. Channel

*N. transatlantica* Bocquet & Stock, 1960  
@ *Clavelina lepadiformis*  
N Ireland, Engl. Channel

*N. platymera* Illg & Dudley, 1961  
@ ?*Ascidia mentula*  
Engl. Channel

*Agnathaner* Canu, 1891

*A. typicus* Canu, 1891 (TM)  
@ *Dendrodoa grossularia* & *Polycarpa rustica*  
S Norway (Grimstad)-Br. Isles-Roscoff

*A. freemani* Hamond, 1968  
@ ?  
Norfolk

*A. minutus* Canu, 1891  
@ *Sidnyum turbinatum*  
N France

*Doropygus* Thorell, 1859  
\* Ca 20 indeterminate species, especially erected by Hesse from the coast of France, exist.

*D. pulex* Thorell, 1859 (TM)  
= *D. pullus* Buchholz, 1869  
@ *Ascidietta scabra* & many other ascidians & *Antedon petasus*  
Bohuslän-Norway-Faeroes-Br. Isles-Mediterranean, NE America

*Doropygella* G.O. Sars, 1921

*D. psyllus* (Thorell, 1859)  
@ *Ascidia* spp., *Ascidietta* spp. & *Phallusia fumigata*  
Bohuslän-W Norway-Br. Isles-N France-Mediterranean

*D. porciauda* (Brady, 1878)  
@ *Corella parallelogramma* & *Ascidia conchilega*  
SE Norway (Hvalør)-W Norway-Clyde Sea-Ireland, Biscay, Mediterranean

*D. thorelli* (Aurivillius, 1882)

- = *Doropygus thorelli* Aurivillius (TM)  
 @ *Ascidia mentula* & *A. obliqua*  
 Bohuslän-S & W Norway-Faeroes-Iceland, Greenland, Davis Strait
- D. normani** (Brady, 1878)  
 @ *Styela rustica*, *Cynthia papillosa* (Linnaeus), *Cynthia* sp. & ?*Ascidia virginea*  
 SW Scotland, NE & W Ireland, S England, Mediterranean
- Doropygopsis** G.O. Sars, 1921
- D. longicauda** (Aurivillius, 1882)  
 = *Doropygus longicauda* Aurivillius, 1882 (TM)  
 @ *Ascidia mentula*, *A. obliqua*, *A. prunum*, *A. callosa* & *Botenia echinata*  
 Bohuslän-S & W Norway-Spitsbergen-Kara Sea-Iceland-Faeroes-Greenland-Davis Strait-Maine
- Pachypygus** G.O. Sars, 1921
- P. gibber** (Thorell, 1859)  
 = *Doropygus gibber* Thorell, 1859 (TM)  
 @ *Ascidia* spp., *Ascidella* spp., *Molgula* spp., *Polycarpa* spp., *Ciona intestinalis*, *Corella parallelogramma*, *Clavelina lepadiformis* etc.  
 Bohuslän-S & W Norway-Br. Isles-Mediterranean
- Notopterophorus** Leuckart, 1859  
 = *Notopterophorus* Costa, 1840 (nom. nud.)
- N. auritus** (Thorell, 1859)  
 @ *Ascidia* spp. & *Ciona intestinalis*  
 Bohuslän-Oslofjord-Trondheim-Shetlands-Br. Isles-Ireland
- N. papilio** Hesse, 1864  
 @ *Ascidia* spp. & *Ciona intestinalis*  
 Bohuslän-W Norway-Shetlands-Br. Isles-Mediterranean
- N. micropterus** G.O. Sars, 1921  
 @ *Ascidia mentula*  
 \* Not mentioned in Gotto, 1993. Could it be a synonym of *N. elatus* Giesbrecht?  
 W Norway
- N. elongatus** Buchholz, 1869  
 = *N. elongatus* Costa, 1840 (nom. nud.)  
 = *N. elatus* Costa, 1840 (nom. nud.)  
 = *N. veranyi* Leuckart, 1859 (TM)  
 @ *Ascidia mentula*, *A. gelatinosa*, *Ascidella aspersa*, *Phallusia mamillata* & *Clavelina lepadiformis*  
 \* Probably confined to *Phallusia mamillata* according to Gotto, 1993  
 ?W Sweden-?Britain-Mediterranean
- N. elatus** Giesbrecht, 1892  
 @ *Ascidia mentula*  
 North Sea, Engl. Channel, Mediterranean
- N. sp.** Schimkewitsch, 1896  
 @ *Ascidia sanguinolenta* & *Ciona intestinalis*  
 Roscoff
- N. bombyx** Hesse, 1865  
 \* indeterminate species.  
 @ *Ciona intestinalis*  
 Coast of France
- Gunenotophorus** Buchholz, 1869 (Spelling emended)  
 = *Gunenotophorus* O.G. Costa, 1840 (nom. nud.)  
 = *Sphaeronotus* Claus, 1864, **non** Laporte, 1832  
 = *Gunenotophorus* Buchholz, 1869 & : Auctt.
- G. globularis** Buchholz, 1869 (TM)  
 = *G. globularis* O.G. Costa, 1840 (nom. nud.)  
 = *Sphaeronotus thorellii* Claus, 1864  
 \* Should not the correct name be *G. thorellii* (Claus, 1864)?  
 @ *Ascidia* spp., *Styela* spp., *Polycarpa* spp., *Ciona intestinalis*, *Molgula manhattensis*, *Dendrodoa grossularia* etc.  
 Bohuslän-Trondheim-Britain-France-Mediterranean, S Africa, Florida
- Botachus** Thorell, 1859
- B. cylindratus** Thorell, 1859 (TM)  
 = *B. fusiformis* Buchholz, 1869  
 @ *Ascidia* spp., *Ascidella scabra* & *Ciona intestinalis*  
 Kattegatt-Bohuslän-Norway-Shetlands-Br. Isles-Mediterranean
- B. fulvus** Hesse, 1869  
 \* indeterminate species  
 @ *Ascidia canina*  
 Coast of France
- B. macroone** Hesse, 1869  
 \* indeterminate species  
 @ *Ascidia canina*  
 Coast of France
- Ophioseides** Gerstaecker, 1871
- O. cardiacephalus** Gerstaecker, 1871 (TM)  
 =? *O. joubini* Chatton, 1909  
 @ *Distomus variolosus*, *Stolonica socialis*, *Polycarpa pomaria*, *Styela* sp. & *Microcosmus sulcatus*  
 Engl Channel, European Atlantic
- Ophiodelphys** Bocquet & Stock, 1961
- O. illgi** Bocquet & Stock, 1961 (TM)  
 @ an ascidian of the family Polyclinidae  
 Engl. Channel
- O. apoda** (Giard, 1873)  
 = *Ophioseides apoda* Giard, 1873  
 @ *Astellium perspicuum*  
 Roscoff
- Ooneides** Chatton & Brément, 1915
- O. amela** Chatton & Brément, 1915  
 @ *Didemnum* spp.  
 W Scotland, Mediterranean
- Bremenia** Chatton & Brément, 1915
- B. balneolensis** Chatton & Brément, 1915  
 @ *Didemnum commune*, *Polysyncrator bilobatum* & *Polysyncrator canatensis*  
 European Atlantic-Mediterranean
- Scolecimorpha** G.O. Sars, 1926
- S. joubini** (Chatton, 1909)
- = *Ophioseides joubini* Chatton, 1909  
 @ *Styela rustica*, *Polycarpa pomaria*, *Microcosmus sulcatus* (= *M. sabatieri*), *Pyura trita* Sluiter & *Cnemidocarpa cerea* Sluiter  
 Bohuslän-?W Norway-French Mediterranean
- S. insignis** G.O. Sars, 1926 (TM)  
 @ *Polycarpa pomaria*  
 Trondheim Fjord
- Doroixys** Kerschner, 1879
- D. uncinata** Kerschner, 1879 (TM)  
 @ *Aplidium* spp., *Sidnyum* spp., *Polyclinum aurantium*, *Morchellium argus*, *Diazona violacea*, *Botrylloides* spp. etc.  
 Br. Isles-NE Ireland-N France-Mediterranean
- Bonnierilla** Canu, 1891  
 = *Paryphes* Kerschner, 1879, **non** Burmeister, 1835
- B. alterata** Stock, 1967  
 = *B. longipes* : (Kerschner, 1879) Canu, 1891, **non** *Paryphes longipes* Kerschner, 1879  
 @ *Clavelina lepadiformis*, *Polycarpa rustica* & *Pyura lurida*  
 W Ireland-N & W France-Mediterranean
- B. filipes** Stock, 1967  
 = *B. armata* : Illg & Dudley, 1961, **non** Schellenberg, 1922  
 @ *Dendrodoa grossularia*, *Polycarpa rustica* & *Ciona roulei*  
 W Ireland, Mediterranean
- B. similis** Illg & Dudley, 1961  
 @ *Pyura squamulosa*  
 NW Ireland, Mediterranean
- Archinotodelphyidae** Lang, 1949  
 \* No taxa in N Europe?  
 @ Ascidiaceans
- Mantridae** Leigh-Sharpe, 1934
- Mantra** Leigh-Sharpe, 1934 (TM, *M. speciosa* Leigh-Sharpe, 1934)
- Ascidicoidea** Thorell, 1859
- Ascidicolidae** Thorell, 1859  
 @ Ascidiaceans
- Ascidicolinae Thorell, 1859  
 @ Ascidiaceans
- Ascidicola** Thorell, 1859
- A. rosea** Thorell, 1859 (TM)  
 @ *Ascidia* spp., *Ascidella* spp., *Corella parallelogramma*, *Ciona intestinalis*, *Pyura squamulosa*, *Polycarpa pomaria* & *Halocynthia papillosa*  
 Isefjord, Bohuslän-S Norway-North Sea-Br. Isles-Mediterranean
- Styelicola** Lützen, 1968
- S. bahusia** Lützen, 1968 (TM)  
 @ *Styela atlantica* (Van Name), *S. gelatinosa* Traustedt & *Ciona gelatinosa*  
 \* So far found only below 80 m.  
 Koster Channel, Biscay
- Buprorinae Thorell, 1859  
 @ Ascidiaceans
- Buprorus** Thorell, 1859
- B. loveni** Thorell, 1859 (TM)  
 @ *Ascidia mentula*, *A. obliqua* & *Ascidella aspersa*  
 Bohuslän-N Norway-Faeroes
- B. nordgaardi** G.O. Sars, 1921  
 @ *Amaroucium* sp.  
 \* Not mentioned in Gotto, 1993  
 Trondheimsfjord
- Enterocolinae Della Valle, 1883
- Lequerrea** Chatton & Harant, 1924
- L. perezii** Chatton & Harant, 1924 (TM)  
 @ intestine of *Polycarpa gracilis* Heller  
 Plymouth, Brest
- L. canui** Illg & Dudley, 1980  
 @ gut of *Polycarpa rustica* (Linnaeus)  
 Plymouth
- Enterocola** van Beneden, 1860  
 = *Biocryptus* Hesse, 1865
- E. bilamellata** G.O. Sars, 1921  
 @ not known (found free)  
 \* Not mentioned by Gotto, 1993  
 SW Norway (Farsund)
- E. pterophora** Chatton & Brément, 1909  
 @ *Didemnum fulgens*, *Trididemnum tenerum* & various other colonial ascidiaceans  
 NE Ireland, S England, Mediterranean
- E. fulgens** van Beneden, 1860 (TM)  
 = *E. betencourti* Canu, 1891  
 @ *Aplidium* spp., *Polyclinum aurantium*, *Synoicum pulmonaria*, *Sidnyum turbinatum* & ?*Botryllus schlosseri*  
 Belgium-Br. Isles-NE Ireland-N France-Mediterranean-S Africa
- E. megalova** Gotto, 1962  
 @ *Polyclinum aurantium* Milne-Edwards  
 NE Ireland
- E. clavelinae** Chatton & Harant, 1924  
 @ *Polycitor (Clavelina) nana* Lahille  
 N France, Mediterranean
- E. hessei** Chatton & Harant, 1924  
 @ Unidentified compound ascidian  
 N France
- E. sydnii** Chatton & Harant, 1924  
 @ *Sidnyum turbinatum*, *S. elegans* & *Morchellium argus*  
 N Ireland, S England, France (Cotes du Nord)
- E. setifera** Hansen, 1923

- @ unknown  
\* Not mentioned by Gotto, 1993  
off SW Iceland
- E. brementi** Illg & Dudley, 1980  
@ *Aplidium pallidum* (Verrill)  
N France
- E. africanus** López-González, Conradi & García-Gómez, 1993  
@ *Synoicum* sp.  
Gibraltar area
- E.** sp. B Chatton & Brément, 1909  
@ unidentified ascidian  
Scotland
- Enteropsinae** Aurivillius, 1885  
@ Ascidians
- Mychophilus** Hesse, 1865  
= *Mycophilus* : Auctt.  
= *Mychephilus* : Canu, 1892  
= *Michophilus* : Chatton & Harant, 1922
- M. roseus** Hesse, 1865 (TM)  
= *M. pachygaster* Hesse, 1865  
= *M. curvatus* Chatton & Brément, 1909  
= *M. rosovula* Gray, 1933  
= *Enteropsis vararensis* T. Scott, 1901  
@ *Botryllus schlosseri* (Pallas) & *Botrylloides leachii* (Savigny)  
Bohuslän-W Norway-Br. Isles-Ireland-W France-Mediterranean
- Enteropsis** Aurivillius, 1885
- E. sphinx** Aurivillius, 1885 (TM)  
= *Haligyrops aculeatus* Aurivillius, 1885  
@ *Molgula manhattensis* (De Kay) & *Dendrodoa grossularia* (van Beneden)  
\* Not mentioned by Gotto, 1993  
Siberian Arctic, Alaska
- E. rosoffensis** Chatton & Brément, 1909  
@ *Dendrodoa grossularia* (van Beneden), *Pyura microcosmus* (Savigny) & *Tethyum savignyi*  
W Ireland, Plymouth, N France-Mediterranean, NW USA
- E. chattoni** Monniot, 1961  
@ *Microcosmus vulgaris* Heller, *Halocynthia papillata* (Linnaeus) & *Diazona violacea* Savigny  
S England, Mediterranean
- E. pilosus** Canu, 1886  
@ *Diazona violacea*  
\* Excluded from the genus in the revision by Illg & Dudley, 1980. Not mentioned by Gotto, 1993.  
W France
- Haplostominae** Chatton & Harant, 1924
- Haplostoma** (Canu, 1886)  
= *Aplostoma* Canu, 1886, non Mouquin-Tandon, 1856  
= *Cryptopodus* Hesse, 1865  
= *Tranostoma* Wilson, 1924  
= *Haplostoma* Chatton & Brément, 1915
- H. brevicauda** (Canu, 1886) (TM)  
@ *Synoicum pulmonaria*, *Polyclinum aurantium*, *Sidnyum turbinatum*, *Morchellium argus* & *Aplidium nordmanni*  
Bohuslän-W Norway-E England-NE & SW Ireland-N France
- H. eruca** (Norman, 1869)  
@ *Ciona intestinalis*  
Oslofjord-Shetlands-E Scotland-NE Ireland
- H. banyulensis** Brément, 1909  
@ *Trididemnum tenerum* & *Didemnum maculosum*  
NE Ireland, French Mediterranean
- H. canui** Chatton & Harant, 1924  
@ *Polyclinum aurantium*  
W France
- H. mizoulei** Monniot, 1962  
@ stomach of the small intestinal form of *Sidnyum turbinatum*  
SW Ireland, Mediterranean
- Haplostomides** Chatton & Harant, 1924
- H. scotti** Chatton & Harant, 1924 (TM)  
@ *Polyclinum aurantium*  
NE Ireland-N France
- H. hibernicus** (T. & A. Scott, 1895)  
@ *Polyclinum aurantium*  
NE & SW Ireland-N France
- ? **H. beaumonti** (T. & A. Scott, 1895)  
= *Enterocola beaumonti* T. & A. Scott, 1895  
@ unknown ascidian  
\* Considered as Haplostominae, sp. incertae sedis, by Ooishi & Illg, 1977. Not mentioned by Gotto, 1993.  
Ireland (Valencia)
- Haplosaccus** Chatton & Harant, 1924
- H. sacculus** (Chatton & Brément, 1910) (TM)  
@ ?  
\* Not mentioned by Gotto, 1993  
? Britain
- Haplostomella** Chatton & Harant, 1924  
= *Rhabdomorpha* Fukui, 1965 (TM) *R. halocynthiae* Fukui, 1965 (Japan)
- H. malacocera** Chatton & Harant, 1924 (TM)  
@ ?  
\* Not mentioned by Gotto, 1993  
?
- Botryllophilinae** Illg & Dudley, 1980  
@ Ascidians
- Botryllophilus** Hesse, 1864  
= *Kossmechthrus* Della Valle, 1883 (TM) *K. notopus* Della Valle, 1883)  
= *Blakeanus* Wilson, 1921 (TM) *B. corniger* Wilson, 1921)
- B. ruber** Hesse, 1864 (TM)  
@ *Botryllus schlosseri*, *Botrylloides leachi*, *Aplidium punctum*, *Morchellium argus*, *Sidnyum* spp.,  
*Synoicum pulmonaria* & *Clavelina lepadiformis*
- \* This taxon is problematic and need revision. According to Gotto, 1993 probably several species have been lumped in this taxon.  
? Bohuslän-? W Norway-? Arctic Seas-? Br. Isles-? Engl. Channel-Brest-? Mediterranean
- B. brevipes** G.O. Sars, 1921  
non = *B. brevipes* Brément, 1909  
\* may possibly be a synonyme of *B. bergensis* Schellenberg, 1921  
@ *Botryllus* sp.  
Bergen
- B. macropus** Canu, 1891  
@ *Lithonephria eugyranda* Giard  
France (Wimereux)
- B. bergensis** Schellenberg, 1921  
@ *Leptoclinides faeroensis* (Bjerkan)  
\* Not mentioned by Gotto, 1993.  
Bergen
- B. norvegicus** Schellenberg, 1921  
= *Blakeanus corniger* Wilson, 1921  
= *Blakeanus groenlandicus* Hansen, 1923  
= *Pelonia corrugata* Goodsir, *Cynthia carnea* Verrill & *Cynthia rustica* Linnaeus  
\* Not mentioned by Gotto, 1993.  
Trondheim-W Greenland-Long Island
- B. aspinosus** Schellenberg, 1922  
@ *Polycarpa pomaria* (Savigny) & *Sryela hupferi* Michaelsen  
\* Not mentioned by Gotto, 1993.  
Plymouth, Angola
- B. virescens** Hesse, 1864  
\* indeterminate. Sp. obl.  
@ ?  
Brest
- Schizoproctinae** Aurivillius, 1887  
\* Neither family nor genus is mentioned by Gotto, 1993.
- Schizoproctus** Aurivillius, 1885  
= *Pteropygus* G.O. Sars, 1921
- S. inflatus** Aurivillius, 1885 (TM)  
@ *Ascidia obliqua* and other ascidians  
N Norway-Spitsbergen-Arctic Seas-E Greenland
- S. vestitus** (G.O. Sars, 1921)  
= *Pteropygus vestitus* G.O. Sars, 1921  
@ *Ascidia obliqua*  
S Norway (Risør)
- Enterognathinae** Illg & Dudley, 1980
- Enterognathus** Giesbrecht, 1900
- E. comatulae** Giesbrecht, 1900 (TM)  
@ guts of ? *Antedon mediterranea* Lamarck, ? *A. rosacea* (Linck) & *A. bifida* (Pennant)  
E Scotland, E Ireland, Hebrides, Plymouth, Mediterranean
- LERNAEOLINA**
- Lernaeidae** Cobbold, 1879  
\* Only fresh-water species recognized from Scandinavia  
@ Fish
- POECILOSTOMATOIDA** Thorell, 1859
- Lateracanthus** ??, 19??  
\* Which family?
- L. quadripedis** ??, 19??  
@ *Coryphaenoides rupestris*  
NW Atlantic
- L. curtus** Kabata, 1993  
@ *Macrourus*  
NE Atlantic
- Bomolochidae** Sumpf, 1871  
@ Fish
- Bomolochus** von Nordmann, 1832  
@ Marine Teleosts
- B. soleae** Claus, 1864 (TM)  
@ in the nose of *Solea solea*  
Helgoland-Holland-Belgium-W France-Mediterranean
- B. bellones** Burmeister, 1835 (TM of *Parabomolochus* Vervoort, 1962)  
@ Belonidae  
Helgoland-Br. Isles-Mediterranean
- Holobomolochus** Vervoort, 1969 (TM *H. nothrus* (Wilson, 1913))
- H. confusus** (Stock, 1953)  
= *Bomolochus soleae* : T. Scott, 1893 & : Auctt., non Claus, 1864  
@ nasal capsules of Gadidae, *Conger*, *Cyclopterus* & *Labrus bergylta*  
Bohuslän-North Sea-Br. Isles-Eur. Atlantic
- Taeniacanthidae** Wilson, 1911  
@ Fish, chiefly teleosts & *Echiniosocius* Humes & Cressey, 1959 & *Echinirus* Humes & Cressey, 1959 on Echinoderms
- Taeniacanthus** Sumpf, 1871 (TM *T. carchariae* Sumpf, 1871)  
= *Anchistrotos* Brian, 1906
- T. wilsoni** A. Scott, 1929  
@ *Raja fullonica*, *R. montagui* & ? *R. naevus*  
Irish Sea, Falmouth Bay & Yorkshire
- T. lucipetus** (Holmes, 1985)  
= *Anchistrotos lucipetus* Holmes, 1985  
@ ?  
Ireland
- T. onosi** (T. Scott, 1902)  
= *Bomolochus onosi* T. Scott, 1902  
@ *Ciliata mustela*, *C. septentrionalis* & *Rhinonemus cimbricus*  
Shetlands-Scotland
- T. laqueus** (Leigh-Sharp, 1935)  
= *Anchistrotos laqueus* Leigh-Sharp, 1935  
= *Anchistrotos hamatus* Rounds, 1960  
@ *Ciliata mustela*, *Serranus cabrilla* & *S. scriba*

W Baltic, Plymouth, Mediterranean

*T. zeugopteri* (T. Scott,1902)  
= *Bomolochus zeugopteri* T. Scott,1902  
@ *Zeugopterus punctatus*  
Clyde Sea-Roscoff

**Ergasilidae** von Nordmann,1832  
@ Marine & limnic teleosts (& gen. *Ostrincola* in bivalve mantle cavities)

**Ergasilus** von Nordmann,1832  
= *Ergasiloides* G.O. Sars,

**E. briani** Markevich,1933  
= *E. minor* Halisch,1934  
@ *Perca fluviatilis*, *Leuciscus rutilus*, *Abramis brama* etc.  
Baltic, Black Sea, Caspian Sea & several European fresh water systems

**E. gibbus** von Nordmann,1832  
@ gills of *Anguilla anguilla* (species specific)  
Brackish Baltic & North Sea waters

**E. lizae** Krøyer,1863  
= *E. nanus* van Beneden,1870  
@ limnic fishes, Mugilidae, *Alosa* & *Engraulis*  
Aberdeen-Mediterranean-Black Sea, Azov Sea (limnic & brackish waters)

**E. mugilis** Vogt,1877 (sp. inquirenda)  
@ Mugilidae  
?

**E. sieboldi** von Nordmann,1832 (TM)  
= *E. tristaceus* von Nordmann,1832  
=? *E. esocis* Sumpf,1871  
= *E. surbecki* Baumann,1912  
= *E. hoferi* Borodin,1915  
= *E. baicalensis* Messjatzeff,1926  
@ on gills of limnic fishes & E Baltic *Anguilla anguilla* & *Platichthys flesus*  
Baltic-Br. Isles-Eurasia

**Thersitina** Norman,1905

**T. gasterostei** (Pagenstecher,1861)  
= *Thersites gasterostei* Pagenstecher,1861 (TM)  
= *Ergasilus biuncinatus* Gadd,1901  
=? *Thersitina inopinata* Percival,1937  
@ branchiae of *Gasterosteus Pungitius* a.o.  
Scandinavia (e.g. Øresund, S Baltic & Furusund)-Br. Isles-Faeroes-Greenland-W France-Caspian Sea

**Chondracanthidae** H. Milne-Edwards,1840  
@ Ectoparasites on fish, chiefly teleosts

**Chondracanthus** Delaroche,1811

**C. zei** Delaroche,1811 (TM)  
= *Lernaeacanthus delarochiana* de Blainville,1822  
@ On gills of *Zeus faber* & other Zeidae  
Plymouth-Mediterranean

**C. neali** Leigh-Sharpe,1930  
@ *Malacocephalus laevis*  
off SW Ireland-S Africa

**C. lophii** Johnston,1836  
= *Lernentoma Dufresnii* de Blainville,1822  
= *Chondracanthus Lophius* Risso,1826  
= *Chondracanthus Delarochiana* Cuvier,1829  
= *Chondracanthus gibbosus* Krøyer,1837  
= *Chondracanthus abdominalis* Heegaard,1943  
@ On gills of *Lophius* spp.  
?Øresund-Kattegat-Norw. Sea-Faeroes-North Sea-Br. Isles-Mediterranean

**C. merluccii** (Holten,1802)  
= *Lernaea merluccii* Holten,1802  
= *Chondracanthus xiphiæ* Guérin-Meneville,1829  
= *Chondracanthus stramineus* Wilson,1923  
@ On gills of *Merluccius* spp & *Trisopterus luscus*  
Norw. Sea-Br. Isles-S Africa

**C. nodosus** (O.F. Müller,1776)  
= *Lernaea nodosa* O.F. Müller,1776  
= *Chondracanthus williamsoni* T. Scott,1909  
@ *Sebastes* spp.  
?Øresund-North Sea-Norw. Sea-Barents Sea-Greenland-Iceland-Faeroes-Br. Isles

**C. ornatus** T. Scott,1900  
@ *Callionymus maculatus* & *C. festivus*  
E & W Scotland-Mediterranean

**Acanthochondria** Oakley,1927

**A. cornuta cornuta** (O.F. Müller,1776)  
= *Lernaea cornuta* O.F. Müller,1776 (TM)  
= *Chondracanthus fluræ* Krøyer,1863  
= *Chondracanthus depressus* T. Scott,1905  
@ *Glyptocephalus*, *Pleuronectes*, *Hippoglossoides*, *Platichthys* & possibly other flatfishes  
Kattegat-North Sea-Norw. Sea-Faeroes-North Atlantic

**A. clavata** (Bassett-Smith,1896)  
= *Chondracanthus clavatus* Bassett-Smith,1896  
@ *Microstomus kitt*  
North Sea-Norw. Sea-White Sea-Iceland-Faeroes-Br. Isles-Bay of Biscay

**A. limandæ** (Krøyer,1863)  
= *Chondracanthus limandæ* Krøyer,1863  
@ *Limanda limanda*  
North Sea-Murmansk-Faeroes-Br. Isles

**A. soleae** (Krøyer,1838)  
= *Chondracanthus soleae* Krøyer,1838  
@ *Solea solea* & *Scophthalmus rhombus*  
North Sea-Faeroes-Br. Isles-Mauretania

**Acanthochondrites** Oakley,1930

**A. annulatus** (Olsson,1869)  
= *Chondracanthus annulatus* Olsson,1869 (TM)  
= *Chondracanthus inflatus* Bainbridge,1909  
= *Chondracanthus japonicus* Gusev,1951  
=? *Chondracanthus pallidus* van Beneden,1851  
@ *Raja* spp.  
North Sea-Norw. Sea-Br. Isles-N Atlantic

**Lernentoma** de Blainville,1822

**L. asellina** (Linnaeus,1758)  
=? *Chondracanthus gurnardi* : Olsson,1869  
@ Triglidae & possibly other marine teleosts  
North Sea-S Norw. Sea-Br. Isles-NE Atlantic-W Mediterranean

**Diocus** Krøyer,1863

**D. frigidus** Hansen,1923  
@ *Lycodes frigidus*  
Iceland

**Philichthyidae** Vogt,1877, emend. Delamare Deboutteville,1962  
= **Sarcotacidae** Yamaguti,1963  
@ Inhabiting the subcutaneous mucus ducts or lateral line canals of Teleosts

**Sarcotaces** Olsson,1872

**S. arcticus** Collett,18??  
@ endoparasitic in "*Molva abyssorum*". Blood from engorged females may stain the flesh of the host.  
North Sea

**Philichthys** Steenstrup,1862

**P. xiphiæ** Steenstrup,1862 (TM)  
@ *Xiphias gladius*  
Kattegat-North Sea-Br. Isles-Mediterranean- North Atlantic

**Colobomatus** Hesse,1873

**C. bergyltæ** Hesse,1873  
@ *Labrus bergylta*  
\* Not mentioned by Kabata,1992  
France

**C. mugilis** Raibaut, Caillet & Ben Hassine,1978  
@ *Liza aurata*  
Mediterranean

**Leposphilus** Hesse,1866

**L. labrei** Hesse,1886  
@ subcutaneous ducts in the frontal region of the head of *Centrolabrus exoletus* & *Cosseyphus diplotaenia*  
\* Not mentioned by Kabata,1992  
France

**Oncaeiidae** Giesbrecht,1892, emend. Boxshall,1977  
@ probably pelagic semiparasites, but hosts not known

**Oncaea** Philippi,1843

**O. borealis** G.O. Sars,1918  
= *O. conifera*: G.O. Sars, non Giesbrecht,1891  
Oslofjord-S & W Norway-Arctic Seas-Br. Isles

**O. conifera** Giesbrecht,1891  
off W Norway, Cape Verde

**O. similis** G.O. Sars,1918  
Norway (Bergen area)

**O. minuta** Giesbrecht,1892  
Gullmarfjord, Skagerrak, Norway (Bergen area)

**O. subtilis** Giesbrecht,1892  
Skagerrak-Norway-Ireland-Mediterran.

**O. obscura** Farran,1908  
Irish Atlantic slope

**O. ornata** Giesbrecht,1891  
Engl. Channel-Ireland, Cape Verde

**O. mediterranea** (Claus,1863)  
= *Antaria mediterranea* Claus,1863  
Mediterranean, Cape Verde

**O. rotundata** Boxshall,1977  
Cape Verde

**O. notopus** Giesbrecht,1891  
Cape Verde

**O. media** Giesbrecht,1891  
= *O. curta* : Estrada & Genicio,1970, non G.O. Sars,1916  
Cape Verde

**O. ivlevi** Schmeleva,1966  
?NE Atlantic, Adriatic

**O. curta** G. O. Sars,1916  
= *O. ovalis* Shmeleva,1966  
= *O. longiseta* Schmeleva,1969  
= *O. latimana* Gordeyeva,1975  
= O. sp.1 Ferrari,1975  
= O. sp.2 Ferrari,1975  
Cape Verde

**O. venusta** Philippi,1843 (TM)  
= *Antaria obtusa* Dana,1852  
= *Oncaea pyriformis* Lubbock,1860  
= *Antaria coerulescens* Claus,1866  
Cape Verde

**Lubbockia** Claus,1863

**L. glacialis** G.O. Sars,1900  
High arctic regions, Cape Verde

**L. brevis** Farran,1908  
NE Atlantic

**L. extenuata** Boxshall,1977  
Cape Verde

**L. minuta** Wolfenden,1905  
NE Atlantic

**L. squillimana** Claus,1863 (TM)  
Cape Verde

**L. aculeata** Giesbrecht,1891  
Cape Verde

**Pseudolubbockia** G.O. Sars,1909

- P. dilatata* G.O. Sars, 1909  
?NE Atlantic
- Conaea* Giesbrecht, 1891
- C. gracilis* (Dana, 1852) T. Scott, 1894  
= *Antaria gracilis* Dana, 1852  
= *Conaea rapax* Giesbrecht, 1891 (TM)  
Norway (Sognesjøen - according to Wiborg 1954), Cape Verde
- Pachos* Stebbing, 1910  
= *Pachysoma* Claus, 1863 (preoccupied)
- P. punctatum* (Claus, 1863) (TM)  
= *Pachysoma punctata* Claus, 1863  
Cape Verde
- P. tuberosum* (Giesbrecht, 1891)  
= *Pachysoma tuberosum* Giesbrecht, 1891  
= *Pachysoma dentatum* Mori, 1932  
Cape Verde
- Epicalymna* Heron, 1977
- E. exigua* (Farran, 1908)  
= *Oncaea exigua* Farran, 1908  
off Ireland
- Paralubbockiidae Boxshall & Huys, 1990
- Paralubbockia* Boxshall, 1977
- P. longipedia* Boxshall, 1977 (TM)  
Cape Verde
- Corycaecidae Dana, 1852  
@ probably pelagic semiparasites on gelatinous zooplankton like appendicularians and salps.
- Corycaeus* Dana, 1845
- C. (Ditrichocorycaeus) M. Dahl, 1912) anglicus* Lubbock, 1855  
Kattegatt-Bohuslän-Oslofjord-W Norway-North Sea-Br. Isles-Engl. Channel-?Biscay
- C. (Corycaeus) clausi* F. Dahl, 1894  
N temperate & tropical Atlantic, Mediterranean
- C. (Corycaeus) speciosus* Dana, 1849  
= *Corycaeus remiger* Dana, 1849-52  
= *Corycaeus varius* Dana, 1849-52  
= *Corycaeus longicaudatus* Dana, 1849-52  
= *Corycaeus Sutherlandi* Lubbock, 1856  
= *Corycaeus styliferus* Lubbock, 1856  
Atlantic, Mediterranean
- C. (Agetus* Krøyer, 1849) *flaccus* Giesbrecht, 1891  
N temperate & tropical Atlantic, Mediterranean
- C. (Urocorycaeus* M. Dahl, 1912) *furcifer* Claus, 1863  
Biscay, temperate & tropical Atlantic, Mediterranean
- C. (Onchocorycaeus* M. Dahl, 1912) *giesbrechti* F. Dahl, 1894  
Temperate & tropical Atlantic, Mediterranean
- C. venustus* Dana, 1849  
?NE Atlantic
- Corycella* Farran, 1911
- C. rostrata* (Claus, 1863)  
= *Corycaeus rostratus* Claus, 1863  
= *Corycaeus parvus* Claus, 1863  
= *Corycaeus pellucidus* Brady, 1883 (p.p.)  
Boreal & temperate Atlantic, Mediterranean
- Sapphirinidae Thorell, 1859  
@ Salps
- Sapphirina* J.V. Thompson, 1829
- S. iris* Dana, 1849  
= *Sapphirina salpae* Claus, 1859  
= *Sapphirina scalaris* Fisher, 1860  
= *Sapphirina longifurca* Scott, 1909  
@ Salps  
?W Norway-NE Atlantic-Mediterranean
- S. angusta* Dana, 1849  
= *Sapphirina Clausi* Haeckel, 1864  
= *Sapphirina Danae* Lubbock, 1856  
= *Sapphirina opaca* Lubbock, 1856  
Boreal Atlantic, Mediterranean
- S. nigromaculata* Claus, 1863  
= *Sapphirina loma* Esterly, 1905  
Temperate & tropical Atlantic, Biscay, Mediterranean
- S. opalina* Dana, 1849  
= *Sapphirina versicolor* Dana, 1849  
= *Sapphirina Darwini* Haeckel, 1864  
= *Sapphirina Thompsoni* Lubbock, 1860  
= *Sapphirina pachygaster* Claus, 1863  
= *Sapphirina aureofurca* Giesbrecht, 1891  
Temperate & tropical Atlantic, Mediterranean
- S. ovatolanceolata* Dana, 1849  
= *Sapphirina gemma* Dana, 1849  
= *Sapphirina Edwardsii* Haeckel, 1864  
= *Sapphirina Gegenbaueri* Haeckel, 1864  
Gibraltar, temperate & tropical Atlantic, Mediterranean
- S. sali* Farran, 1929  
= *Sapphirina pseudolactens* Lehnhofer, 1929  
= *Sapphirina gibba* Rose, 1929  
Temperate & tropical Atlantic, Biscay
- Copilia* Dana, 1849  
= *Sapphirinella* Claus, 1863  
= *Hyalophyllum* Haeckel, 1864
- C. vitrea* Haeckel, 1864  
N temperate & tropical Atlantic, Mediterranean
- Corina* Giesbrecht, 1891
- C. granulosa* Giesbrecht, 1891  
off W Ireland, Mediterranean
- Urocopiidae Humes & Stock, 1972  
@ not known, but probably gelatinous zooplankton
- Urocopia* G.O. Sars, 1917
- U. singularis* G.O. Sars, 1917  
pelagic S of Iceland
- Clausiidae Giesbrecht, 1895  
@ Polychaeta & possibly other invertebrates like terebratulid bivalves
- Clausia* Claparède, 1863
- C. lubbocki* Claparède, 1863  
@ ?  
E England, Engl. Channel
- C. uniseta* Bocquet & Stock, 1960  
@ ?  
E England
- Rhodicola* Levensen, 1878  
= *Seridium* Giesbrecht, 1895  
@ Maldanidae
- R. elongata* Levensen, 1878  
@ *Rhodine gracilior*  
Øresund, SE Scotland, NE England, ?W Ireland
- R. gibbosa* Bresciani, 1964  
@ *Praxillella praetermissa*  
\* Not mentioned by Gotto, 1993  
Øresund
- Conchocheres* G.O. Sars, 1918  
\* Tentatively put in this fam. Systematic position not fully clear.
- C. malleolatus* G.O. Sars, 1918  
@ pallial cavity of *Cuspidaria obesa*  
\* Not mentioned by Gotto, 1993  
W Norway
- Megaclausia* O'Reilly, 1995
- M. mirabilis* O'Reilly, 1995 (TM)  
@ In tubes of *Rhodine gracilior*  
SE Scotland, NE England
- Mesnilia* Canu, 1898
- M. martinensis* Canu, 1898  
@ *Polydora flava*  
Engl. Channel
- M. cluthae* (T. & A. Scott, 1896)  
@ *Polydora ciliata* & *P. flava*  
NE, E & S England, SW Scotland, N France
- Pseudoclausia* Bocquet & Stock, 1960
- P. longiseta* Bocquet & Stock, 1963  
@ *Hymeniacion sanguinea* & ?*Aphysina aerophoba*  
Engl. Channel, Mediterranean
- Clausidiidae Embleton, 1901  
@ Sponges, Octocorals, Polychaets, Bivalves, Sipunculids, Nudibranchs, Stomatopods, Thalassinideans
- Conchyluria* Bocquet & Stock, 1957  
@ Bivalvia
- C. cardii* Gooding, 1957  
@ Mantle cavity of *Acanthocardia* spp., *Tapes decussatus*, *Venerupis* spp., *Scrobicularia plana*, *Petricola pholadiformis*, *Barnea candida*, *Meretrix chione*  
\* Two subspecies are recognized: *C. cardii cardii* Gooding, 1957 @ *Acanthocardia echinata* (Linnaeus) & *Meretrix chione* (Linnaeus) and *C. cardii tapetis* Bocquet & Stock, 1958 @ *Venerupis aurea*.  
E & SW Ireland, SW England, N & W France
- C. solenis* Bocquet & Stock, 1957 (TM)  
@ Mantle cavity of *Solen marginatus* Don.  
Roscoff
- Giardella* Canu, 1888
- G. callianassae* Canu, 1888  
@ Burrows of *Callianassa subterranea*  
NW England, Engl. Channel
- G. thompsoni* A. Scott, 1906  
@ ?  
Irish Sea
- Hemicyclops* Boeck, 1872
- H. purpureus* Boeck, 1872  
@ not known  
Bohuslän-Oslofjord-SE Scotland-SW England
- H. cylindracea* (Pelseneer, 1929)  
@ Mantle cavity of *Loripes lacteus*  
N & W France
- H. aberdonensis* (T & A. Scott, 1892)  
@ ?  
E England, NE Scotland, E & SW Ireland, W France
- Hersiliodes* Canu, 1888  
@ Maldanidae
- H. latericia* (Grube, 1869)  
@ in tube of *Leiochone clypeata*  
E England, W & SW Ireland, N & W France, Mediterranean
- Hippomolgus* G.O. Sars, 1918
- H. furcifer* G.O. Sars, 1918  
@ ?  
S Norway (Risør)-Stavanger

- Leptinogaster** Pelseneer,1929
- L. histria** (Pelseneer,1929)  
@ *Macoma balthica* (Linnaeus) & *Abra alba*  
German Bight, Engl. Channel
- L. pholadis** Pelseneer,1929  
@ *Pholas dactylus*  
Brest, Naples
- L.** sp.  
@ Digestive tract of *Echinocardium cordatum*  
Netherlands
- Pseudopsyllus** T. Scott,1902
- P. elongatus** T. Scott,1902  
@ ?  
?Britain
- Eunicolidae** G.O. Sars,1918  
@ Eunicid Polychaeta
- Eunicola** Kurz,1877
- E. clausi** Kurz, sensu G.O. Sars, ?non Kurz,1877  
@ *Eunice claparedi*  
\* Sensu Gotto,1993 is the North Sea form recorded by Sars differing from the true Mediterranean *E. clausi*  
Kurz in several significant details.  
Oslofjord-S Norway (Risør)-Faeroes-Br. Isles-Mediterranean
- E. insolens** (T. & A. Scott,18??)  
@ *Eunice harassii* Audouin & Milne-Edwards  
SW Scotland- NE Ireland
- Catiniidae** Bocquet & Stock,1957  
@ Sipunculids
- Catinia** Bocquet & Stock,1957
- C. plana** Bocquet & Stock,1957 (TM)  
@ On skin of *Sipunculus nudus*  
Engl. Channel, W France
- Myzomolgus** Bocquet & Stock,1957
- M. stupendus** Bocquet & Stock,1957 (TM)  
@ On skin of *Sipunculus nudus*  
\* Ho,1984 suggests a closer affinity to Catiniidae than to Clausidiidae.  
Engl. Channel, W France
- Anomopsyllidae** G.O. Sars,1921
- Anomopsyllus** G.O. Sars,1921
- A. pranizoides** G.O. Sars,1921  
@ ?  
Oslofjord
- Nereicolidae** Claus,1875  
@ Ectoassociates of errant Polychaeta
- Nereicola** Keferstein,1863
- N. ovatus** Keferstein,1863  
@ *Platynereis dumerilii*, *Perinereis cultrifera*, *Nereis rava*, *Nereis zonata*, *N.* sp.  
Engl. Channel, Mediterranean, Black Sea
- Flabelliphilus** Bresciani & Lützen,1962  
\* Systematic position not clear. Nereicolid affinity suggested by Stock,1968.
- F. inersus** Bresciani & Lützen,1962  
@ Among gelatinous protuberances of the jelly coat covering the body of *Flabelligera affinis*  
Bohuslän
- Sigecheres** Bresciani,1964
- S. brittae** Bresciani,1964  
=? *Nereicola concinna* T. Scott,1902  
@ *Sige fusigera*  
Öresund-Kattegat-Bohuslän, ?W Scotland
- Selioides** Levinsen,1878
- S. bolbraei** Levinsen,1878  
= *Cancerina confusa* T. Scott,1900  
@ *Harmothoe imbricata*, *Gattyana cirrosa*, *Eunoe nodosa*, *Lagisca rarispina*, *Antinoe sarsi* & *Hermadion rouchi*  
Öresund-Bohuslän-North Sea-Britain-Arctic Seas-Iceland
- S. bocqueti** Carton,1963  
@ *Scalisetosus assimilis*  
Roscoff
- Spiophanicolidae** Ho,1984  
\* Included in the "Nereicolidiform group" by Ho 1984.
- Spiophanicola** Ho,1984 (TM *S. spinulosus* Ho,1984 - California @ *Spiophanes* spp.)
- Entobiidae** Ho,1984  
@ Terebellidae  
\* This family is not recognized by Huys & Boxshall 1991, but included in the "Nereicolidiform group" by Ho 1984.
- Entobius** Dogiel,1908
- E. loimiae** Dogiel,1908  
@ ?*Loimia* sp.  
\* Not mentioned by Gotto,1993  
?
- E. hamondi** Gotto,1966  
@ *Polycirrus caliendrum*  
SW England
- Mytilicolidae** Bocquet & Stock,1957  
@ Bivalves & Gastropods
- Mytilicola** Steuer,1902
- M. intestinalis** Steuer,1902  
@ *Mytilus edulis* & *M. galloprovincialis* (& *Cerastoderma edule*, *Ostrea edulis* & *Tapes decussatus*)  
Limfjorden-North Sea-Br. Isles-Portugal-Mediterranean
- M. orientalis** Mori,1935  
= *M. ostreae* C.B. Wilson,1938  
=? *M. macrae* Hoshina & Kuwabara,1959  
@ *Crassostrea gigas* & *Mytilus crassitesta* & ? *Maetra veneriformis*  
\* Introduced from Japan to Pacific N America, there infesting e.g. *Mytilus edulis*, *M. californicus*, *Ostrea lurida* & *Crepidula fornicata*, later introduced in W France with imported *Crassostrea gigas*, there infesting also *Mytilus* spp. & *Ostrea edulis* - Not mentioned by Gotto,1993  
Arcachon, NE Pacific, Japan
- Trochicola** Dollfus,1914
- T. entericus** Dollfus,1914  
@ Rectum of *Gibbula cineraria*, *G. varia*, *Calliostoma zizyphinum* & *Monodonta mutabilis*  
N France, Mediterranean
- T. pectinidarum** Tuzet & Ormières,1961  
@ *Aequipecten opercularis* & *Pecten glaber*  
\* Not mentioned by Gotto,1993  
French Mediterranean
- Anthessiidae** Humes,1986  
@ Mollusca
- Anthessius** Della Valle,1880  
= *Pseudomolgus* G.O. Sars,1916
- A. dilatatus** (G.O. Sars,1918)  
*Pseudomolgus dilatatus* G.O. Sars,1918  
@ ?  
\* Not mentioned by Gotto,1993  
W Norway (Romsdalfjord)
- A. arenicolus** (Brady,1872)  
= *Lichomolgus arenicolus* Brady,1872  
@ *Dosinia exoleta*  
NE England, SW Scotland, SW Ireland, Engl. Channel, W France
- A. leptostylis** (G.O. Sars,1916)  
= *Pseudomolgus leptostylis* G.O. Sars,1916  
@ *Buccinum undatum*  
Bohuslän-Oslofjord-Trondheim-Spitsbergen, E England
- A. arcuatus** López-González, Conradi, Naranjo & García-Gómez,1992  
@ *Berthella stellata* (Risso,1826)  
Cádiz, Spain
- A. teissieri** Bocquet & Stock,1958  
= *Pseudomolgus arenicola* : G.O. Sars,1921, non *Lichomolgus arenicolus* Brady,1872  
@ *Buccinum undatum*  
Hvalør, E England, Engl. Channel
- A. pleurobranchae** Della Valle,1880  
@ *Pleurobranchaea* sp.  
Napoli
- Mycolidae** Yamaguti,1936  
@ Molluscs
- Myicola** Wright,1885
- M. ostreae** ??,1???  
@ *Crassostrea gigas*, *Ostrea edulis* & *Crassostrea angulata*  
Arcachon
- Pseudomyicola** Yamaguti,1936
- P. spinosus** (Raffaele & Monticelli,1885)  
@ *Mytilus galloprovincialis*, *M. edulis*, *Perna pectinata*, *Ostrea stentata* & *Pteria hirundo*  
Arcachon-Mediterranean-Black Sea-Bermudas-Caribbean
- Rhinomolgus** G.O. Sars,1918  
\* Humes & Stock,1973 put this genus in this fam. (Mycolidae)!
- R. anomalus** G.O. Sars,1918  
@ *Antedon*  
Trondheimfjord
- Ventriculinidae** Leigh-Sharp,1934  
@ Sipunculoids & Gastropods
- Endocheres** Bocquet & Stock,1956
- E. obscurus** Bocquet & Stock,1956  
@ In rectal gland or in galls developed in the pallial cavity of *Calliostoma zizyphinum* (Linnaeus)  
N France
- Antheacheridae** M. Sars,1870  
\* This family includes perhaps (sensu Ho,1981) three genera parasiting sipunculans: *Akessonia*, *Coelotropus* & *Siphonobius*  
@ Actinian mesenteries ?& Sipunculids
- Antheacheres** M. Sars,1857
- A. duebeni** M. Sars,1857  
@ In rounded galls formed from the mesentery walls of *Bolocera tuedie*  
Bohuslän-W & N Norway
- Akessonia** Bresciani & Lützen,1962  
\* Family placement probably not definitive, but Humes,1987 thinks this for the moment might be a proper family to keep this genus in.
- A. occulta** Bresciani & Lützen,1962  
@ In coelomic cavity of *Golfingia minuta*  
Öresund-Bohuslän
- Gastroecus** Hansen,1886
- G. arcticus** Hansen,1886 (TM)  
@ *Actinostola intermedia* & *Actinostola spetsbergensis* Carlgren  
\* Not mentioned by Gotto,1993  
Kara Sea
- Staurosoma** Will,1844
- S. parasiticum** Will,1844  
@ In galls developed on the mesenteries of *Anemonia sulcata*  
Normandie-Mediterranean
- Lamippidae** Joliet,1882  
@ Octocorals
- Lamippe** Bruzelius,1858

- L. rubra* Bruzelius, 1858  
@ *Pennatula phosphorea*  
Skagerrak, S Norway, Mediterranean
- Lamipella* Bouligand & Delamare Deboutteville, 1959
- L. faurei* Bouligand & Delamare Deboutteville, 1959  
@ In coenosarc canal of *Swifitia rosea* & other octocorals  
Bohuslän, Mediterranean
- Enalcyonium* Olsson, 1869
- E. rubicundum* Olsson, 1869  
@ In coenosarc canal & sometimes within host polyps of *Alcyonium digitatum*, *A. palmatum* & *A. acaule*  
Bohuslän-Br. Isles, NE Ireland, N France, ?Mediterranean
- E. olssoni* (De Zulueta, 1908)  
@ *Alcyonium digitatum*  
Bohuslän
- E. forbesi* (T. Scott, 1901)  
@ *Alcyonium digitatum*  
SE Scotland, Isle of Man-Wales-N France
- Mesoglicolidae de Zulueta, 1911  
\* Order uncertain. Lamippidae and Ascidicolidae have been discussed as possibly related.  
@ Anthozoans
- Mesoglicola* Quidor, 1906
- M. delagei* Quidor, 1906  
@ In mesogloea and in galls of *Corynactis viridis* Allman  
Roscoff & Banyuls
- Jeanella* T. Scott, 1904
- J. minor* (T. Scott, 1902)  
@ ?  
\* Correct family?  
?Britain
- Splanchnotrophidae Norman & Scott, 1906  
@ Molluscs (chiefly Nudibranchs) & Pterobranchia
- Splanchnotrophus* Hancock & Norman, 1863  
=? *Lomanotocola* T. & A. Scott, 1895  
\* Not yet seen revision by Delamare-Deboutteville, 1950
- S. brevipes* Hancock & Norman, 1863  
@ *Coryphella verrucosa*, *Facelina curta*, *Doto* spp., *Cuthona genovae*  
Isefjord, Kattegatt-Bohuslän-Br. Isles, Ireland
- S. gracilis* Hancock & Norman, 1863  
@ Endoparasitic in *Acanthodoris pilosa*, *Okenia aspersa*, *Eubranchius pallidus* & *Doto coronata*  
SE Scotland, E & SW England, Isle of Man, W Ireland
- S. insolens* (T. & A. Scott, 1895)  
= *Lomanotocola insolens* T. & A. Scott, 1895  
@ *Lomanotus genei*  
\* Of uncertain status  
SW Ireland
- S. willemi* ?, 19??  
@ *Facelina coronata*  
\* Not mentioned by Gotto, 1993  
?
- S. angulatus* ?, 19??  
@ *Aeolidia papillosa*, *Aeolidiella glauca*  
\* Not mentioned by Gotto, 1993  
?
- Gastrodelphyidae List, 1890
- Gastrodelphys* Graeffe, 1883
- G. clausii* Graeffe  
@ *Bispira volutacornis*  
Engl. Channel, Mediterranean
- Anomoclausidae Gotto, 1964
- Corallovexiidae Stock, 1975
- Echiurophilidae Delamare-Deboutteville & Nunes-Ruivo, 1955
- Pharodidae Illg, 1948
- Philoblennidae Izawa, 1976  
@ on a cowry
- Rhynchomolgidae Humes & Stock, 1972
- Serpulidicolidae Stock, 1979
- Serpulidicola* Southword, 1963
- S. placostegi* Southword, 1963  
@ *Placostegus tridentatus*  
W entrance of Engl. Channel in 450-760 m
- Shiinoidea Cressey, 1975
- Synaptiphilidae Bocquet, 1953  
@ Holothuroidea
- Synaptiphilus* Canu & Cuénot, 1892  
= *Colaceutes* Hartman, 1856 (nomen oblitum)  
= *Remigulus* T. & A. Scott, 1893  
\* This genus was originally placed in a separate family, Synaptiphilidae Bocquet, 1952, moved to Clausiidae by Bocquet & Stock, 1957 but suggested by Ho, 1984 to have closer affinities to Clausiidae.  
@ Echinoderms
- S. luteus* Canu & Cuénot, 1892 (TM)  
= *Colaceutes mulleri* Hartmann, 1856 (nomen oblitum)  
@ *Labidoplax digitata* (Montagu), *Leptosynapta gallienae* (Herapath), *L. inhaerens* (Müller) & *L. bergensis* Östergren  
SW Br. Isles-N & W France-Mediterranean
- S. cantacuzenei* Bocquet & Stock, 1957  
@ *Labidoplax digitata*  
NW France-Mediterranean
- S. tridens* (T. & A. Scott, 1893)  
= *Remigulus tridens* T. & A. Scott, 1893  
@ *Leptosynapta inhaerens* (Müller), *Leptosynapta cruenta* Cherbonnier & *Labidoplax media*  
W Scotland, Ireland, NW France
- Presynaptiphilus* Bocquet & Stock, 1960  
\* Correct family?
- P. acrocnidae* Bocquet & Stock, 1960  
@ *Acrocnida brachiata*  
N France-SW England
- Telsidae Ho, 1967
- Tuccidae Vervoort, 1962
- Vahiniidae Humes, 1967
- Xarifiidae Humes, 1960
- Lichomolgoidea Humes & Stock, 1972
- Lichomolgidae Kossmann, 1877  
@ very wide host spectrum
- Acaenomolgus* Humes & Stock, 1972
- A. protulae* (Stock, 1959)  
@ *Protula tubularia* & *Protula intestinum*  
NE Ireland-SW Scotland, Mediterranean
- A. serpulae* (Stock, 1960)  
@ *Serpula vermicularis*  
Mediterranean France
- Astericola* Rosoll, 1889
- A. clausi* Rosoll, 1889 (TM)  
@ *Marthasteria glacialis*, *Asterina gibbosa* & *Astropecten aurantium*  
Roscoff-European Atlantic, Mediterranean
- Critomolgus* Humes & Stock, 1983
- C. actinae* (Della Valle, 1880)  
= *Doridicola actinae* (Della Valle, 1880)  
= *Lichomolgus actinae* Della Valle, 1880  
@ *Anemonia sulcata* & *Actinia* spp.  
Plymouth-NE Ireland-Roscoff-Mediterranean
- C. bulbipes* (Stock & Kleeton, 1963)  
@ ?  
N? Europe
- Doridicola* Leydig, 1853  
=? *Metaxymolgus* Humes & Stock, 1972
- D. agilis* Leydig, 1853  
= *Eolidicola tenax* M. Sars, 1862  
= *Lichomolgus agilis* : Auctt.  
= *Doridicola agilis* : Odhner, 1922  
@ many different Nudibranchs (& *Aplysia*, *Hiatella* & *Polynoe* sp.) and ? *Sagittatus sagittatus*  
Bohuslän-North Sea-S & W Norway-Br. Isles-France-Mediterranean-Senegal
- D. chlamydis* (Humes & Stock, 1973)  
@ *Chlamys varia*  
European Atlantic
- D. longicauda* (Claus, 1860)  
= *Metaxymolgus longicauda* : (Claus, 1860) Humes & Stock, 1972  
@ gills & mantle cavity of *Sepia officinalis* & *S. officinalis* var. *filliouxii*  
Netherlands, S England, Arcachon, Biscay, Mediterranean
- D. hirsutipes* (T. Scott, 1893)  
= *Macrocheiron hirsutipes* : G.O. Sars, 1917  
= *Metaxymolgus hirsutipes* : (T. Scott, 1893) Humes & Stock, 1972  
@ ?tubes of *Sabella pavonina*  
S & W Norway, Novaya Semlja, SE Scotland, NW England
- Epimolgus* Bocquet & Stock, 1956
- E. trochi* (Canu, 1899)  
@ *Gibbula* spp. (e.g. *G. cineraria*) & *Monodonta*  
Engl. Channel-W Mediterranean
- Lichomolgella* G.O. Sars, 1918
- L. pusilla* G.O. Sars, 1918  
@ ?  
S-SW Norway, SE Ireland
- Lichomolgides* Gotto, 1954
- L. cuanensis* Gotto, 1954  
@ cloacal cavities of *Trididemnum tenerum* (Verrill)  
NE Ireland
- Lichomolgus* Thorell, 1859
- L. albens* Thorell, 1859  
@ Ascidians, often *Corella* or e.g. *Molgula manhattensis*  
Isefjord, Bohuslän-North Sea-S & W Norway-Br. Isles-N & W France
- L. canui* G.O. Sars, 1917  
@ many different Ascidians, e.g. *Molgula*, *Stryela* & *Dendrodoa*  
Kiel-Norway-Holland-N & S Ireland-France-Sicily
- L. diazoniae* Gotto, 1961  
@ *Diazonia violacea* Savigny  
Plymouth
- L. forficula* Thorell, 1859  
=? *L. elongatus* Buchholz, 1869  
@ *Ascidia mentula* (& *Psammechinus miliaris*)  
Bohuslän-S & W Norway-Br. Isles-N France-Mediterranean
- L. furcillatus* Thorell, 1859  
@ *Ciona* (& *Corella parallelogramma*)  
Bohuslän-Norway-Shetlands-Br. Isles-Mediterranean
- L. leptodermatus* Gooding, 1957  
@ gills of *Laevicardium crassum*  
Engl. Channel & SW England
- L. marginatus* Thorell, 1859  
@ *Ascidia* spp.

Bohuslän-Holland-Norway-N France-NE Ireland

**Macrochiron** Brady,1872

**M. fucicolum** Brady,1872

@ Generally associated with algae.  
?Norway-S North Sea-Br. Isles, Ireland, NE USA, Mediterranean

**M. mutatum** Stock,1957

= *Macrocheiron fucicolum* : G.O. Sars,1917, non Brady,1872 (p.p.)  
Oslofjord-S & W Norway

**Octopicola** Humes, in Delamare-Deboutteville, Humes & Paris,1957

**O. superbus superbus** Humes,1957

@ *Octopus vulgaris*  
Roscoff-Mediterranean

**Zygomolgus** Humes & Stock,1972

**Z. tenuifurcatus** (G.O. Sars,1917)  
@ *Labidoplax digitata* & *Diplosoma listerianum*  
S & W Norway-NE & SW Ireland

**Z. curtiramus** (Bocquet & Stock,1962)

@ the alga *Lithophyllum incrustans*  
N France

**Z. didemni** (Gotto,1956)

@ *Didemnum maculosum* (Milne-Edwards)  
NE Ireland

**Z. poucheti** (Canu,1891)

= *Lichomolgus poucheti* Canu,1891  
@ Ascidians, e.g. *Sidnyum*, *Morchellium* & *Fragarium*  
S & W Norway-W Ireland-N & NW France

**Stellicola** Kossmann,1877

\* Not mentioned by Gotto,1993.

**Pseudanthessidae** Humes & Stock,1972

@ Wide host spectrum

**Heteranthessius** T. Scott,1904

**H. dubius** (T. Scott,1903)

@ ?  
Clyde Sea

**H. scotti** Bocquet, Stock & Bénard,1959

@ ?  
Bretagne, among *Lithophyllum incrustans*

**H. furcatus** Stock,1971

@ *Microcosmus sabatieri*  
N France, Mediterranean

**Pseudanthessius** Claus,1889

**P. gracilis** Claus,1889

@ *Pomatoceros*, *Filograna*, *Spisula* & *Palmaria palmata*  
Bohuslän-S & W Norway-Br. Isles-SW Ireland-N & W France-Mediterranean

**P. assimilis** G.O. Sars,1917

@ *Echinus elegans*  
\* Not mentioned by Gotto,1993  
Norway (Hardangerfjord & Romsdalsfjord)

**P. dubius** G.O. Sars,1918

@ *Asterias rubens*  
\* Not mentioned by Gotto,1993  
Bohuslän-W Norway

**P. liber** (Brady,1880)

@ *Echinus esculentus*, *E. acutus*, *Psammechinus miliaris*, *Paracentrotus lividus* & *Strongylocentrotus droebakiensis*  
Bohuslän-S & W Norway-Br. Isles-Faeroes-N France

**P. sauvaigi** Canu,1892

@ *Echinocardium cordatum*, *E. flavescens*, *Spatangus purpureus* (& *Echinus esculentus*)  
Bohuslän-S North Sea-S Norway-Br. Isles-N France-Mediterranean

**P. thorellii** (Brady,1880)

@ *Swiftia pallida*, *Cladocora caespitosa* & ?*Asterias rubens*  
Bohuslän-Br. Isles-SW Ireland-?Italy

**Nemerthessius** Stock,1985

**N. nemertophilus** (Gallien,1936)

= *Pseudanthessius nemertophilus* Gallien,1936  
@ *Lineus longissimus* Sowerby  
Wimereux (NE France), W France

**Phyllocolidae** Delamare-Deboutteville & Laubier,1961

@ Phyllocolidae

**Cyclorhiza** Heegaard,1942

**C. eteonicola** Heegaard,1942

@ *Eteone longa*  
Trondheimsfjord-?Isle of Man, E North America

**C. megalova** Gotto & Leahy,1988

@ *Eteone longa*  
W Ireland

**Sabelliphilidae** Gurney,1927

@ Wide host spectrum

**Lichomolgidium** Kossmann,1877

**L. cynthiae** (Brian,1924)

=?? *L. sardum* Kossmann,1877  
@ *Syela clava* & *Pyura microcosmus*  
Plymouth-Roscoff-Mauretania, ??Mediterranean

**Modiolicola** Aurivillius,1882

**M. insignis** Aurivillius,1882

@ *Modiolus modiolus* & *Mytilus edulis* & *M. galloprovincialis*  
Bohuslän-SE Norway (Hvalør)-Netherlands-Br. Isles-Engl. Channel-Mediterranean-Black Sea

**M. inermis** Canu,1892

@ *Aequipecten opercularis*  
E Scotland, W Br. Isles-Engl. Channel-W France-W Mediterranean

**M. maximus** (Thompson,1893)

@ *Pecten maximus* (& *Aequipecten opercularis* f. *audouini*)  
W Br. Isles-NE & SW Ireland,N & W France, Mediterranean

**Myxomolgus** Humes & Stock,1972

**M. myxicolae** (Bocquet & Stock,1958)

@ *Myxicola infundibulum* (Rénier)  
S England-Finistère

**M. proximus** Humes & Stock,1973

@ *Myxicola aesthetica* (Claparède)  
Finistère

**Paranthessius** Claus,1889

**P. anemoniae** Claus,1889

=? *Doricicola antheae* Ridley,1879  
@ *Anemonia sulcata* (Pennant)  
N France-E England-NE & S-W Ireland-Mediterranean

? **P. dubia** (G.O. Sars,1921)

= *Herrmannella dubia* G.O. Sars,1921

@ ?  
\* Not mentioned by Gotto,1993.  
Norway (Drøbak)

**Herrmannella** Canu,1891

= *Paranthessius* : Auctt., non Claus,1889

**H. valida** G.O. Sars,1918

@ ?  
\* Not mentioned by Gotto,1993.  
Norway (Stavanger)

**H. parva** Norman & T. Scott,1905

@ ?*Eunicella stricta*

\* The position of this species in *Herrmannella* is questionable according to Gotto,1993  
S & W Norway-Engl. Channel-W Mediterranean

**H. rostrata** Canu,1891

@ many different bivalves, including *Cerastoderma edule*, *Spisula* spp., *Venerupis pullastra*, *Macoma balthica* (& *Ascidia mentula*)  
Elbe mouth-Holland-Engl. Channel-Br. Isles-Irish Sea-France-Mediterranean

**H. barneae** (Pelseneer,1929)

@ *Barnea*, *Pholas* & *Ostrea*  
E England, N & W France

**H. haploceras** (Bocquet & Stock,1959)

= *Paranthessius colmani* Reddiah,1960  
@ *Laevicardium crassum*  
Isle of Man, Engl. Channel, European Atlantic

**H. pecteni** (Sowinski,1884)

@ Pectenidae & *Anomia ephippium*  
W Br. Isles-NW & W France-Mediterranean & Black Sea

**H. duggani** Holmes & Minchin,1991

@ ?  
\* Not mentioned by Gotto,1993  
?Britain

**Sabelliphilus** M. Sars,1862

**S. elongatus** M. Sars,1862 (TM)

@ branchial crown of *Sabella pavonina* & *Spirographis*  
Bohuslän-S, W & NW Norway-Br. Isles-Engl. Channel-NE Ireland-W France-Mediterranean

**S. sarsi** Claparède,1870

@ body of *Spirographis spallanzanii* & *Sabella*  
Engl. Channel-W France-Mediterranean

**Scambicornus** Heegaard,1944

= *Prehermannella* Sewell,1949

**S. finmarchicus** (T. Scott,1903)

= *Herrmannella finmarchica* : Auctt.  
= *S. armoricanus* (Bocquet, Stock & Kleeton,1963)  
@ *Eupolymnia nebulosa*  
S, W & N Norway, SW Ireland, Roscoff

**S. prehensilis** (G.O. Sars,1918)

= *Herrmannella prehensilis* G.O. Sars,1918

@ ?  
\* Not mentioned by Gotto,1993.  
Norway (outside Sognefjord)

**S. tenuicaudis** (G.O. Sars,1918)

= *Herrmannella tenuicaudis* G.O. Sars,1918

@ ?  
\* Not mentioned by Gotto,1993.  
W Norway

**Calypsina** Humes & Stock,1972

\* Not mentioned by Gotto,1993.

**Calypsarion** Humes & Ho,1969

\* Not mentioned by Gotto,1993.

**Caribulus** Humes & Stock,1972

\* Not mentioned by Gotto,1993.

**Chauliobion** Humes,1975

\* Not mentioned by Gotto,1993.

**Diogenella** Stock,1968

\* Not mentioned by Gotto,1993.

**Diogenidium** Edwards,1891

\* Not mentioned by Gotto,1993.

**Lecanurius** Kossmann,1877

\* Not mentioned by Gotto,1993.

**Lichothuria** Stock,1968

\* Not mentioned by Gotto,1993.

**Synapticola** Voigt,1892

\* Not mentioned by Gotto,1993.

**Ventriculinidae** Leigh-Sharp,1934

**Family uncertain**

**Gomphopodarion** Humes,1974

**G. byssoicum** Humes,1974  
@ *Oneirophanta mutabilis* Theel  
off W Ireland in ca 4430 m

SIPHONOSTOMATOIDA Latreille,1829

Caligidae Burmeister,1835  
@ Fish (& perhaps Cephalopods)

**Caligus** O.F. Müller,1785  
\* Are *C. belones* Krøyer,1863 & *C. angustatus* ?Krøyer,18?? good species or synonyms of other species.  
Both taxa are described from the North Sea? *C. belones* is found by P. Olsson in Skagerrak "ad corpus *Belones vulgaris*".

**C. curtus** O.F. Müller,1785 (TM)  
= *C. mulleri* Leach,1816  
= *C. bicuspidatus* von Nordmann,1832  
= *C. americanus* Pickering & Dana,1838  
= *C. elegans* van Beneden,1851  
= *C. fallax* Krøyer,1863  
= *C. aeglefini* Krøyer,1863  
@ Teleosts & Elasmobranchs, chiefly Gadoids  
Öresund-Kattegat-Skagerrak-Br. Isles-Norw. Sea-Faeroes-Iceland-boreoartic Atlantic

**C. bonito** Wilson,1905  
= *C. kuroshio* Shiino,1959  
= *C. krishnai* Thomas,1967  
@ chiefly Scombridae, e.g. *Sarda* spp.  
Br. Isles-S Africa-Mediterranean

**C. brevicaudatus** A. Scott,1901  
@ Triglidae  
Irish Sea-Barents Sea-Mediterranean-S Africa

**C. centrodoni** Baird,1850  
= *C. abbreviatus* Krøyer,1863  
@ *Pagellus bogaraveo*, Labridae & *Sebastes marinus*  
Bergen-North Sea-Br. Isles, Black Sea

**C. coryphaenae** Steenstrup & Lütken,1861  
=? *C. scutatus* H. Milne-Edwards,1840  
=? *C. thymni* Dana,1852  
= *C. aliuncus* Wilson,1905  
= *C. elongatus* Heegaard,1943, non von Nordmann,1832  
= *C. tesserifer* Shiino,1952  
non = *C. coryphaenae* : Brian,1935, neg : Yamaguti,1936, neg : Barnard,1955  
@ chiefly Scombridae & Carangidae, e.g. *Thunnus*  
North Sea-Tropical Seas

**C. diaphanus** von Nordmann,1832  
@ Triglidae & several other teleost taxa  
Bohuslän-Norway-Br. Isles-Spain-Mediterranean

**C. elongatus** von Nordmann,1832  
= *C. kroeyeri* H. Milne-Edwards,1840  
= *C. rissoanus* H. Milne-Edwards,1840  
= *Caligus rapax* : Baird,1850 (p.p.), non H. Milne-Edwards,1840  
= *C. gurnardi* Krøyer,1863 (p.p.)  
= *C. lumpi* Krøyer,1863  
= *C. trachipteri* Kollar, in Krøyer,1863  
= *C. latifrons* Wilson,1905  
= *C. rabidus* Leigh-Sharp,1936  
= *C. arcticus* Brandes,1956  
non = *C. elongatus* Heegaard,1943  
@ many teleost (e.g. *Scomber*) & elasmobranch species  
\* Probably the most common parasitic copepod in British waters, sensu Kabata,1979  
Bohuslän-Norw. Sea-Iceland-Br. Isles-Ireland-southwards

**C. gurnardi** Krøyer,1863  
= *C. nanus* Krøyer,1863  
non = *C. gurnardi* : Brian,1898, neg : Wilson,1908, neg : Fraser,1920  
@ Triglidae  
North Sea-Br. Isles-European Atlantic

**C. labracis** T. Scott,1902  
@ *Labrus bergylla* & *L. mixtus*  
Irish Sea, S & SW England

**C. minimus** Otto,1821  
= *C. minutus* : Auctt.  
@ *Dicentrarchus*, *Mugil*, *Pagellus* & *Umbrina*  
North Sea-Br. Isles-Spain-Mediterranean

**C. pelamydis** Krøyer,1863  
= *C. scomberi* Bassett-Smith,1896, non *C. scombri* : J.V. Thompson, in W. Thompson,1856  
@ *Scomber*, other Scomridae & several other teleosts  
Skagerrak-North Sea-Br. Isles-S Atlantic-Mediterranean-Black Sea

**C. productus** Dana,1852  
@ chiefly big Scombridae, but also other fish taxa  
Atlantic

**C. zeii** Norman & T. Scott,1906  
@ Usually in gill chamber of *Zeus faber*  
S Br. Isles-southwards

**C. pageti** Russel,1925  
@ *Liza aurata*  
Mediterranean

**Pseudocaligus** A. Scott,1901

**P. brevipedis** (Bassett-Smith,1896) (TM)  
@ usually gadid fish, but also e.g. *Gobius*  
Br. Isles-Bay of Biscay

**P. apodus** Brian,1924  
@ *Liza aurata*  
Mediterranean

**Sciaenophilus** van Beneden,1852

**S. tenuis** van Beneden,1852 (TM)  
= *Caligus (Sciaenophilus) benedeni* Bassett-Smith,1898  
@ Sciaenidae  
?Baltic-North Sea-Br. Isles-S Atlantic-Mediterranean

**Lepeophtheirus** von Nordmann,1832

**L. pectoralis** (Müller,1776) (TM)  
= *Lernaea pectoralis* O.F. Müller,1776

@ inner surface of pectoral fins of chiefly Pleuronectidae, but also other hosts, e.g. *Gadus morhua* & *Raja radiata*  
Bohuslän-Norw. Sea-Faeroes-Iceland-White Sea-Br. Isles

**L. hippoglossi** (Krøyer,1837)  
= *Caligus hippoglossi* Krøyer,1837  
= *Lepeophtheirus obscurus* Baird,1850 (male)  
@ *Hippoglossus hippoglossus* & Reinhardtius hippoglossoides; occasionally other flatfishes  
Kattegat-Denmark-North Sea-Barents Sea-Iceland-Faeroes-Br. Isles

**L. nordmanni** (H. Milne-Edwards,1840)  
= *Caligus nordmanni* H. Milne-Edwards,1840  
= *Caligus ornatus* H. Milne-Edwards,1840  
= *Lepeophtheirus insignis* Wilson,1908  
@ *Mola mola*  
North Sea, around Britain, "Europe"-Iceland-S Atlantic-Mediterranean

**L. pollachius** Bassett-Smith,1896  
= *L. innominatus* Wilson,1905  
@ chiefly *Pollachius* spp. & *Molva molva*  
around Br. Isles, North Sea

**L. salmonis** (Krøyer,1837) "Laxlus"  
= *Caligus salmonis* Krøyer,1837  
= *Caligus vespa* H. Milne-Edwards,1840  
= *Caligus strömii* Baird,1848  
= *Caligus pacificus* Gisler,1883  
= *Lepeophtheirus uenoi* Yamaguti,1939  
@ chiefly *Salmo*, *Oncorhynchus* & *Salvelinus* spp.  
\* A marine copepod with capability to survive limnic water for up to two weeks during host spawning.  
Br. Isles-N Hemisphere, Norwegian Salmon farms, rivers in all nordic countries

**L. sturionis** (Krøyer,1838)  
= *Caligus sturionis* Kr.,1838  
@ *Acipenser sturio* & *A. transmontanus*  
Baltic-Denmark-Skagerrak-North Sea-Br. Isles, Pacific N America

**L. thompsoni** Baird,1850  
= *Caligus gracilis* van Beneden,1851  
= *Lepeophtheirus rhombi* Krøyer,1863  
= *Lepeophtheirus gibbosus* Krøyer,1863  
= *Lepeophtheirus graciliscens* Krøyer,1863  
=? *Lepeophtheirus obscurus* Bassett-Smith,1896  
@ *Scophthalmus maximus*  
W Baltic-Bohuslän-North Sea-Norw. Sea-Faeroes-Br. Isles-W Mediterranean

**L. europaeus** Zeddarn, Berbebi, Renaud, Raibaut & Gabrion,1988  
= *L. thompsoni* : Auctt. (p.p.)  
@ *Scophthalmus rhombus* & *Platichthys flesus*  
?Scandinavia, Britain

**Euryphoridae** Wilson,1905  
@ Fish

**Euryphorus** H. Milne-Edwards,1840 (TM *E. nordmanni* H. Milne-Edwards,1840)

**E. brachypterus** (Gerstaecker,1853)  
=? *Caligeria bella* Dana,1852  
= *Elytrophora brachyptera* Gerstaecker,1853  
= *Arnaeus thymni* Krøyer,1863  
= *Dysgamus longifurcatus* Wilson,1923  
= *Elytrophora atlantica* Wilson,1932  
= *Elytrophora indica* Shiino,1958  
= *Dysgamus sagamiensis* Shiino,1958  
@ big Scombridae  
E England, Engl. Channel-all oceans inhabited by "tunas"

**Trebiidae** Wilson,1905  
@ Fish

**Trebius** Krøyer,1838

**T. caudatus** Krøyer,1838  
@ *Raja* spp. & some other elasmobranchs  
Skagerrak-North Sea-Norw. Sea-Br. Isles-Iceland-S Africa-Mediterranean

**Pandaridae** H. Milne-Edwards,1840  
@ Fish

**Pandarus** Leach,1816

**P. bicolor** Leach,1816 (TM)  
= *Pandarus bosci* Leach,1816  
= *Pandarus fissifrons* H. Milne-Edwards,1840  
= *Pandarus lividus* Frey & Leuckart,1847  
= *Nogagus angustulus* Gerstaecker,1854  
= *Nogagus latus* T. Scott,1907  
@ many different sharks, e.g. *Squalus acanthias*  
Bohuslän-Norw. Sea-Iceland-Br. Isles-southwards

**Dinemoura** Latreille,1829

**D. producta** (O.F. Müller,1785) (TM)  
= *Caligus productus* O.F. Müller,1785  
= *Dinemoura lamnae* Baird,1830  
= *Dinematura affinis* Thomsen,1949  
@ many different sharks, e.g. *Squalus acanthias*  
Öresund-Norw. Sea-Br. Isles-southwards

**D. ferax** (Krøyer,1838)  
@ sharks, e.g. *Carcharias glaucus* (Linnaeus)  
Faeroes-Iceland-Arctic Seas

**Echthrogaleus** Steenstrup & Lütken,1861

**E. caloptratus** (Guerin-Meneville,1837) (TM)  
= *Dinemoura affinis* H. Milne-Edwards,1840  
= *Dinematura braccata* Dana,1852  
= *Nogagus lutkenii* Norman,1868  
= *E. perspicax* Olsson,1869  
@ many different sharks, e.g. *Squalus acanthias*  
Bohuslän-Norw. Sea-Faeroes-Shetlands-Br. Isles-southwards

**Demoleus** Heller,1865

**D. heptapus** (Otto,1821) (TM)  
= *Caligus heptapus* Otto,1821  
= *Caligus paradoxus* Otto,1821  
= *Binoculus sexsetaceus* von Nordmann,1832  
@ *Hexanchus* spp.  
Br. Isles-Atlantic, Mediterranean, Pacific

**Phyllothyreus** Norman,1903

**P. cornutus** (H. Milne-Edwards,1840)  
@ *Prionace glauca* & other big sharks  
Br. Isles-Atlantic

**Nogaus** Leach,1816  
= *Nogaus* : Auctt. & *Nessipus* : Auctt.  
\* An assemblage of pandarid males, often known only from free-swimming records.

"N." **borealis** Steenstrup & Lütken,1861  
= *Nessipus borealis* (Steenstrup & Lütken,1861) : Wilson,1907  
@ *Isurus oxyrinchus* - & in plankton  
Br. Isles-Atlantic-Iceland-Barents Sea

"N." **ambiguus** (T. Scott,1907)  
@ *Squalus acanthias*  
North Sea-Plymouth

**Cecropidae** Dana,1852  
@ Fish

**Cecrops** Leach,1816

**C. latreillii** Leach,1816 (TM)  
@ gills of *Mola mola* & *Mola lanceolata*  
Öresund-Norw. Sea-Iceland-Br. Isles-southwards

**Orthogoriscicola** Poche,1902

**O. muricatus** (Krøyer,1837) (TM)  
= *Laemargus muricatus* Krøyer,1837  
= *Orthogoriscicola szidati* Stekhoven & Stekhoven,1956  
= *Orthogoriscicola wilsoni* Stekhoven & Stekhoven,1956  
@ *Mola mola*  
Norw. Sea-Iceland-Br. Isles-Atlantic-Mediterranean

**Philorthogoriscus** Horst,1897

**P. serratus** (Krøyer,1863) (TM)  
= *Dinemoura serrata* Krøyer,1863  
@ *Mola mola*  
Br. Isles-Atlantic-Mediterranean

**Luetkenia** Claus,1864

**L. asterodermi** Claus,1864 (TM)  
= *Cecropsina glabra* Heller,1865  
@ *Larus imperialis*  
Br. Isles-Mediterranean-Atlantic

**Dichelesthiidae** Dana,1853  
@ Fish

**Dichelesthium** Hermann,1804

**D. oblongum** (Abildgaard,1794)  
= *Caligus oblongus* Abildgaard,1894  
= *Dichelesthium sturionis* Hermann,1804 (TM)  
@ *Acipenser* spp. & *Huso huso*  
North Sea-Br. Isles-Mediterranean-Black Sea

**Anthosoma** Leach,1816

**A. crassum** (Abildgaard,1794)  
= *Caligus crassus* Abildgaard,1794  
= *Caligus imbricatus* Risso,1816  
= *Anthosoma smithii* Leach,1816 (TM)  
@ Half embedded in flesh of Galeoidea & *Mola mola*, only poserior half partially protruding  
Öresund-Belgium-Br. Isles-Atlantic-Mediterranean

**Eudactylinidae** Yamaguti,1963  
@ Gills of Fish

**Eudactylina** van Beneden,1853

**E. acuta** van Beneden,1853 (TM)  
@ *Squatina squatina* & ?*Squalus acanthias*  
?Skagerrak-Br. Isles-Mediterranean-Canaries

**E. similis** T. Scott,1902  
@ *Raja* spp. & *Torpedo nobilitiana*  
North Sea-Br. Isles-Mediterranean

**E. acanthii** A. Scott,1901  
@ *Squalus acanthias* & *Mustelus asterias*  
British Isles-N Atlantic-Angola

**E. insolens** T. & A. Scott,1913  
@ *Galeorhinus galeus*  
Norfolk, Irish Sea

**E. minuta** T. Scott,1904  
@ *Dasyatis pastinaca*  
Scotland (Dornoch Firth)

**Eudactylinella** Wilson,1932

**E. alba** Wilson,1932 (TM - Woods Hole)  
@ *Aetobatus narinari* & *Dasyatis* spp.  
Falmouth Bay, Mediterranean, New England-Gulf of Mexico

**Nemesis** Risso,1826 (TM *N. lamna* Risso,1826)

**N. lamna** Risso,1826 **vermi** A. Scott,1929 : Hewitt,1969  
@ *Cetorhinus maximus*  
Firth of Clyde, Concarneau, New Zealand, Japan

**N. robusta** (van Beneden,1851)  
= *Ergasilina robusta* van Beneden,1851  
= *Pagodina robusta* van Beneden,1853  
= *Nemesis pallida* Wilson,1932  
= *Nemesis aggregatus* Cressey,1967  
@ a number of shark species & some few rays  
Br. Isles-Atlantic-Mediterranean

**Kroyeriidae** Kabata,1979  
@ Elasmobranchs (except for the Holocephalan gill parasite *Prokroyeria meridionalis*)

**Kroyeria** van Beneden,1853

**K. lineata** van Beneden,1853 (TM)  
=? *Lomchidium aculeatum* Gerstaecker,1854  
= *Kroyeria sublineata* Yamaguti & Yamasu,1959  
@ several shark species  
S North Sea-Br. Isles-European Atlantic-Mediterranean, E USA, Japan

**Pseudocycnidae** Wilson,1922, emend. Kabata,1979  
@ Fish

**Pseudocycnus** Heller,1865

**P. appendiculatus** Heller,1865 (TM)  
= *P. spinosus* Pearse,1952  
= *P. thunnus* Brandes,1955  
@ *Thunnus thynnus* & other big Scombridae; some other teleosts  
Bohuslän-North Sea-Br. Isles

**Hatschekiidae** Kabata,1979  
@ Gills of Teleosts

**Hatschekia** Poche,1902

**H. hippoglossi** (Guérin,[1831])  
= *Clavella hippoglossi* Guérin,[1831] (TM)  
@ *Hippoglossus hippoglossus* & *Reinhardtius hippoglossoides*  
Skagerrak-Br. Isles-Iceland-Barents Sea-Greenland-Europ. Atlantic

**H. cluthae** (T. Scott,1902)  
= *Clavella cluthae* T. Scott,1902  
@ *Ctenolabrus rupestris* (& *Labrus bergylta*)  
Br. Isles

**H. labracis** (van Beneden,1871)  
= *Clavella labracis* van Beneden,1871  
= *Cycnus Labri mixti* Hesse,1879  
@ *Labrus mixtus*, *L. bergylta* & *L. trimaculatus*  
Br. Isles & adjacent coastal regions of Europe

**H. nulli** (van Beneden,1851)  
= *Clavella nulli* van Beneden,1851  
= *Clavella crassa* Valle,1879  
@ *Mullus surmuletus* & related mullids  
Belgium-Br. Isles-Bay of Biscay-Mediterranean

**H. pagellibogneravei** (Hesse,1879)  
= *Cycnus Pagelli Bogneravei* Hesse,1879  
=? *Clavella sargi* Valle,1882  
= *H. cornigera* T. Scott,1909  
@ *Pagellus Dentex* & *Sargus*  
North Sea-Br. Isles-Senegal-Mediterranean

**H. pygmaea** T. & A. Scott,1913  
@ *Ctenolabrus melops*, *Ctenolabrus rupestris* & ?*Labrus bergylta*  
Br. Isles-Roscoff

**Congericola** van Beneden,1854

**C. pallidus** van Beneden,1854 (TM)  
@ *Conger conger*  
S North Sea-Br. Isles-Eur. Atlantic-W Mediterranean

**Lernanthropidae** Kabata,1979  
@ Fish

**Lernanthropus** de Blainville,1822 (TM *L. musca* de Blainville,1822)

**L. gisleri** van Beneden,1852  
= *L. Thompsoni* Brian,1898  
@ Sciaenidae, Scatophagidae, Trachinotidae, Centropomidae & Polynemidae  
Belgium-Br. Isles-W Africa-Mediterranean

**L. kroyeri** van Beneden,1851  
@ *Dicentrarchus labrax* & related teleosts  
North Sea-Br. Isles-W Africa-Mediterranean

**L. mugilis** Brian,1898  
@ *Liza aurata*  
Mediterranean

**Pennelliidae** Burmeister,1835 emend. Kabata,1979  
@ Fish & Whales. They have normally a 2-host cycle.

**Lernaocera** de Blainville,1822

**L. branchialis** (Linnaeus,1767)  
= *Lernaea branchialis* Linnaeus,1767 (TM)  
= *Lernaea gobina* O.F. Müller,1776  
= *Lernaocera megacephala* A. Scott,1929  
= *Lernaocera wilsoni* Schuurmans Stekhoven,1936  
= *L. obtusa* Kabata,1957  
=? *Lernaocera caparti* Machado Cruz,1959  
@ 13 intermediate hosts, 8 of them flatfish, but also e.g. *Callionymus lyra*, *Cyclopterus lumpus*, *Agonus cataphractus* & *Myoxocephalus scorpius*, 21 definitive hosts, of which ca half of them are Gadidae  
Kattegat-Bohuslän-North Sea-Iceland-Faeroes-Br. Isles-Greenland-N Atlantic-Mediterranean

**L. lusci** (Bassett-Smith,1896)  
= *Lernaea lusci* Bassett-Smith,1896  
= *Lernaocera phycidis* Leigh-Sharpe,1933  
= *Lernaocera nulli* Leigh-Sharpe,1935  
=? *Lernaocera longicollis* Schuurmans Stekhoven,1935  
@ *Solea solea* is intermediate host; *Trisopterus*, other gadoids & several other teleosts are definitive hosts.  
Kattegat-North Sea-Barents Sea-Br. Isles-Morocco

**L. minuta** (T. Scott,1900)  
@ *Pomatoschistus minutus* & *Myoxocephalus quadricornis* (definitive hosts)  
\* May be a dwarf form of *L. lusci*, living on small hosts.  
Baltic-North Sea-Br. Isles

**Haemobaphes** Steenstrup & Lütken,1861

**H. cycloptera** (O. Fabricius,1780)  
= *Lernaea cycloptera* O.F. Müller,1776 (nom. nud.)  
= *Lernaea cycloptera* O. Fabricius,1780 (TM)  
@ broad range of definitive hosts, e.g. Cottidae, Zoarcidae, Liparidae & *Agonus*; intermediate host unknown  
Kattegat-Norw. Sea-Iceland-Faeroes-Br. Isles-N Atlantic-Arctic Seas

**H. ambiguus** T. Scott,1900  
@ *Callionymus maculatus*, *C. festivus* & *C. belenus*  
W Scotland-W Mediterranean

**Lernaenicus** Le Sueur,1824 (TM *L. radiatus* Le Sueur,1824)

**L. sprattae** (Sowerby,1806)  
= *Lernaea sprattae* Sowerby,1806  
= *Lernaea cyclophora* de Blainville,1822  
= *Lernaocera surtrensensis* de Blainville,1823  
= *Lernaea ocularis* Cuvier,1830  
= *Lernaeonema monillar* H. Milne-Edwards,1840  
= *Lernaeonema sprattae* Baird,1850  
= *Lernaeonema bairdi* Salter,1850  
= *Lernaeonema monillar* Steenstrup & Lütken,1861

- = *Lernaeenicus sardinae* Baudouin, 1904  
 @ eyes of *Sprattus sprattus*, *Sardina pilchardus* (& occasionally *Clupea harengus*)  
 Bohuslän-Oslofjord-North Sea-Norw. Sea-Br. Isles-W Mediterranean
- L. encrasicoli** (Turton, 1807)  
 = *Lernaea encrasicoli* Turton, 1807  
 @ *Sprattus sprattus*, *Engraulis encrasicolus*, *Sardina pilchardus* & *Alosa alosa*  
 Distribution like its congener above
- L. neglectus** Richiardi, 1877  
 @ *Liza aurata*  
 Mediterranean
- Pennella** Oken, 1815 (™ *Pennatula sagitta* Linnaeus, 1758 @ *Histrio histrio* - NW Atlantic)  
 @ 1:rst intermediate host : Cephalopoda  
 \* Nom. cons.; ICZN Op. 1817.
- P. filosa** (Linnaeus, 1758)  
 = *Lernaea cirrhosa* La Martiniere, 1798  
 = *Pennella orthogorisci* Wright, 1870  
 = *Pennella rubra* Brian, 1906  
 = *Pennella plumosa* Linton, 1905  
 = *Pennella germonia* Leigh-Sharpe, 1931  
 @ *Thunnus* spp. and other big Scombridae & *Mola* spp & *Xiphias gladius*.  
 Br. Isles-Atlantic- Fundy Bay
- P. balaenopterae** Koren & Danielssen, 1877  
 = *P. crassicornis* Steenstrup & Lütken, 1861  
 = *P. antarctica* Quidor, 1913  
 = *P. anthonyi* Quidor, 1913  
 = *P. cetti* Quidor, 1913  
 = *P. charcoti* Quidor, 1913  
 @ *Balaenoptera* spp., *Hyperoodon rostratus* (O.F. Müller) and other big whales  
 \* Why Hogans 1987 used Koren & Danielssen's younger name instead of Steenstrup & Lütken's name in his revision of this species is unclear.  
 Norw. Sea-Iceland, Faeroes
- P. instructa** Wilson, 1917  
 @ *Xiphias gladius*  
 NW Atlantic
- P. varians** Steenstrup & Lütken, 1861  
 @ *Eledone moschata*, *Todaropsis eblanae*, *Loligo vulgaris*, *Alloteuthis subulata*, *Sepia officinalis* & *S. elegans*  
 \* This is probably a synonym of another species.  
 Engl. Channel, Mediterranean
- Peniculus** von Nordmann, 1832
- P. clavatus** (O.F. Müller, 1776)  
 @ *Sebastes marinus*  
 Oslofjorden, North Sea-Norw. Sea-Iceland
- P. fistula** von Nordmann, 1832? (™)  
 =? *P. fissipes* Wilson, 1917  
 @ *Zeus faber*  
 Naples
- P. sp.** : Schram, 1991  
 @ *Scomber scombrus*  
 Celtic Sea
- Sarcotretes** Jungersen, 1911
- S. scopeli** Jungersen, 1911  
 @ *Benthosema glaciale*  
 \* the copepod may be fouled by a hydroide, *Ichthyocodium sarcotretis* Jungersen, 1911  
 North Sea-W Norway-Norw. Sea-Faeroes
- Peroderma** Heller, 1865
- P. cylindricum** Heller, 1865?  
 @ various organs of *Sardina pilchardus*  
 Spain, France, Tunisia
- Sphyridae** Wilson, 1919  
 @ Fish
- Sphyrion** Cuvier, 1830 (™ *S. laevigatum* (Quoy & Gaimard, 1824))
- S. lumpi** (Krøyer, 1845)  
 = *Lestes lumpi* Krøyer, 1845  
 =? *Sphyrion norvegicum* Thor, 1900  
 @ Gadiformes, Scorpaeniformes, Anarhichas & Reinhardtius  
 Kattegatt-N Norway-Iceland-Br. Isles-S Atlantic
- Lophoura** Kölliker, 1853
- L. edwardsi** Kölliker, 1853 (™)  
 @ Macrouridae, chiefly *Coelorinchus coelorinchus*  
 W Br. Isles-Angola-W Mediterranean
- Tripaphylus** Richiardi, 1878
- T. musteli** (van Beneden, 1851)  
 = *Lernaeonema musteli* van Beneden, 1851 (™)  
 @ *Mustelus* spp.  
 Br. Isles-Irish Sea-Eur. Atlantic-Italy
- Lernaeopodidae** Olsson, 1869  
 @ Marine & limnic Fish
- Achtheres** von Nordmann, 1832
- A. percarum** von Nordmann, 1832 (™)  
 = *A. sandrae* Gadd, 1904  
 = *A. sibiricus* Messjatzeff, 1926  
 @ *Perca fluviatilis*, *Lucioperca* spp. & *Acerina cernua*  
 Brackish water at river mouths in the Baltic, Black Sea, Aral Sea, Sea of Azov & limnic waters
- Salmincola** Wilson, 1915
- S. salmoneus** (Linnaeus, 1758)  
 = *Lernaea salmonea* Linnaeus, 1758 (™)  
 @ *Salmo salar*, *S. gairdneri*, *Onkorrhynchus keta* & *Thymallus thymallus*  
 Limnic-marine; Baltic-Br. Isles-"Europe"
- S. gordonii** Gurney, 1933  
 = *S. salmoneus* : Kabata, 1979 (p.p.)  
 @ *Trutta trutta*  
 Yorkshire, Scotland, Port Erin, Ireland, ?Orkneys
- S. extensus** (Kessler, 1868)
- @ *Coregonus*  
 \* Not mentioned by Kabata, 1992  
 Baltic & limnic
- S. thymalli** (Kessler, 1868)  
 @ *Thymallus* & other salmonids  
 Limnic-"freshened estuarine"; Sweden, Finland, ?Britain
- Ommatokoita** Leigh-Sharpe, 1926
- O. elongata** (Grant, 1827)  
 = *Lernaea elongata* Grant, 1827  
 = *Lernaeopoda elongata* : Auctt.  
 = *Ommatokoita superba* Leigh-Sharpe, 1926 (™)  
 @ cornea of eyes of *Somniosus microcephalus* (& *Lamna* & *Mustelus*)  
 Skagerrak-off Belgium-Br. Isles-Norw. Sea-Faeroes-Iceland-Greenland-Barents Sea
- Vanbenedenia** Malm, 1860
- V. kroeyeri** Malm, 1860 (™)  
 @ dorsal spine of *Chimaera monstrosa*  
 Kattegatt-Skagerrak-North Sea-W Norway-Faeroes
- Charopinus** Krøyer, 1863
- C. dalmanni** (Retzius, 1829)  
 = *Lernaea dalmanni* Retzius, 1829 (™)  
 = *Strylophorus hipposcephalus* Hesse, 1879  
 @ On gills, in nasal fossae & on inner surfaces of the spiracles of *Raja batis* (& possibly other ray species)  
 North Sea-Norw. Sea-Iceland-Barents Sea-Br. Isles-off Eur. Atl.-Mediterranean-Black Sea
- C. dubius** T. Scott, 1900  
 @ Attached within branchial & buccal cavities of *Raja batis*, *R. maculata*, *R. circularis*, *R. oxyrinchus*, *R. montagui*, *R. naevius*, *R. fullonica* & *Laeviraja macrorhynchus*  
 North Sea-Br. Isles-E North Atlantic
- Pseudocharopinus** Kabata, 1964 (™ *P. dentatus* Wilson, 1912))
- P. bicaudatus** (Krøyer, 1837)  
 = *Lernaeopoda bicaudata* Krøyer, 1837  
 @ spiracles of *Squalus acanthias*, *S. fernandinus*, *Trygon* sp. (& possibly other elasmobranchs)  
 North Sea-Br. Isles-W Africa
- P. malleus** (Rudolphi, in von Nordmann, 1832)  
 = *Brachiella malleus* Rudolphi, in von Nordmann, 1832  
 = *Brachiella lophii* H. Milne-Edwards, 1840  
 = *Brachiella pastinacae* van Beneden, 1851  
 @ *Dasyatis pastinaca*, *Trygon* spp. (& *Torpedo*, *Myliobatis* & *Rhinoptera*)  
 Br. Isles-Senegal-Mediterranean
- Schistobranchia** Kabata, 1964
- S. ramosa** (Krøyer, 1863)  
 = *Charopinus ramosus* Krøyer, 1863 (™)  
 @ *Raja radiata*, *R. clavata*, *R. fullonica* & *R. maculata* (+ ?*R. batis* & ?*R. scabrata*)  
 North Sea-Barents Sea-Iceland-Br. Isles-?Black Sea
- Lernaeopodina** Wilson, 1915 (™ *L. relata* Wilson, 1915)
- L. longimana** (Olsson, 1869)  
 = *Lernaeopoda longimana* Olsson, 1869  
 = *Lernaeopoda cluthae* T. Scott, 1900  
 =? *Lernaeopoda similis* Scott & Scott, 1913  
 @ *Raja fullonica*, *R. radiata*, *R. batis*, *R. erinacea*, *R. laevis*, *R. naevius* & *R. montagui*  
 Skagerrak-North Sea-Barents Sea-Iceland-Br. Isles
- Lernaeopoda** de Blainville, 1822 (™ *L. brogniartii* de Blainville, 1822)
- L. galei** Krøyer, 1837  
 = *Achtheres selachiorum* Kurz, 1877  
 = *Lernaeopoda scyllicola* Leigh-Sharpe, 1916  
 @ Most often in cloacal region of *Scylliorhinus*, *Galeorhinus*, *Mustelus* (& other elasmobranchs)  
 Br. Isles-Barents Sea-Iceland-S Africa-Mediterranean-Black Sea
- L. bidiscalis** Kane, 1892  
 @ *Galeorhinus* (& *Mustelus* & *Scylliorhinus*)  
 North Sea-Br. Isles-Iceland-Eur. Atlantic-Mediterranean
- L. mustelicola** Leigh-Sharpe, 1919 (sp. inquire.)  
 @ *Mustelus vulgaris*  
 \* Probably a synonyme of *L. galei* Krøyer  
 Plymouth-?Sweden
- Albionella** Kabata, 1979
- A. globosa** (Leigh-Sharpe, 1918)  
 = *Lernaeopoda globosa* Leigh-Sharpe, 1918 (™)  
 @ nasal fossae of *Scylliorhinus caniculus*  
 Br. Isles
- Clavellistes** Shiino, 1963 (™ *C. shoyoe* Shiino, 1963)
- C. lampri** (Scott & Scott, 1913)  
 = *Lernaeopoda lampri* Scott & Scott, 1913  
 @ *Lampris guttatus*  
 North Sea-Faeroes-Shetlands-Atlantic
- Brachiella** Cuvier, 1830
- B. thynni** Cuvier, 1830 (™)  
 = *Thynnica ziegleri* Miculicich, 1905  
 @ big Scombridae, *Argyrosomus regium* & *Pomatomus saltatrix*  
 Br. Isles-southwards
- Neobrachiella** Kabata, 1979
- N. merluccii** (Bassett-Smith, 1896)  
 = *Brachiella merluccii* Bassett-Smith, 1896 (™)  
 @ Chiefly on the tips of the gill rakers of *Merluccius* spp.  
 Br. Isles-E North Atlantic-Mediterranean
- N. bispinosa** (von Nordmann, 1832)  
 = *Brachiella bispinosa* von Nordmann, 1832  
 = *Lernaeopoda obesa* Krøyer, 1837  
 = *Anchorella ovalis* : Auctt., non Krøyer, 1837  
 @ Triglidae  
 Skagerrak-Br. Isles-W Mediterranean
- N. chevreuxii** (van Beneden, 1891)  
 = *Brachiella chevreuxii* van Beneden, 1891  
 = *Clavella sciaenae* Brian, 1906  
 = *Brachiella macrura* Wilson, 1920  
 @ Scianidae & Lutjanidae, e.g. *Argyrosomum regium*  
 Br. Isles-Eur. Atlantic-Mediterranean

- N. impudica* (von Nordmann, 1832)  
= *Brachiella impudica* von Nordmann, 1832  
= *Brachiella parva* Nunes-Ruivo, 1957  
@ Triglidae  
North Sea-Br. Isles-W Africa-Mediterranean
- N. insidiosa insidiosa* (Heller, 1865)  
= *Brachiella insidiosa* Heller, 1865  
= *Parabrachiella australis* Wilson, 1923  
= *Brachiella lageniformis* Szidat, 1955  
@ *Merluccius* spp.  
Br. Isles-Eur. & Afr. Atlantic, Mediterranean
- N. rostrata* (Krøyer, 1837)  
= *Brachiella rostrata* Krøyer, 1837  
@ *Hippoglossus hippoglossus*, *Reinhardtius hippoglossoides* (& *Scophthalmus maximus*)  
Kattegat-North Sea-Br. Isles-Faeroes-Iceland-Barents Sea-Greenland
- N. triglae* (Claus, 1860)  
= *Brachiella triglae* Claus, 1860  
@ Triglidae  
North Sea-Br. Isles-Atl. France-Mediterranean
- Clavella* Oken, 1815  
\* Nom. cons.; ICZN Op. 1817.
- C. adunca* (Strøm, 1762)  
= *Lernaea adunca* Strøm, 1762 (TM)  
= *Lernaea cyclopteryna* O.F. Müller, 1776  
= *Lernaea uncinata* O.F. Müller, 1776  
= *Anchorella lagenula* Cuvier, 1830  
= *Anchorella uncinata* von Nordmann, 1832  
= *Anchorella brevicollis* H. Milne-Edwards, 1840  
= *Anchorella agilis* Krøyer, 1863  
= *Anchorella quadrata* Bassett-Smith, 1896  
= *Clavella dubia* T. & A. Scott, 1913  
= *Clavella sciatherica* Leigh-Sharpe, 1918  
= *Clavella iadda* Leigh-Sharpe, 1920  
= *Clavella devastatrix* Leigh-Sharpe, 1925  
= *Clavella invicta* Leigh-Sharpe, 1925  
= *Clavella typica* Leigh-Sharpe, 1925  
= *Clavella deliciosa* Leigh-Sharpe, 1933  
= *Clavellina uncinata* Dogiel, 1936  
= *Clavellina brevicollis* Dogiel, 1936  
@ chiefly Gadidae, but several other hosts known  
Øresund-Skagerrak-North Sea-Br. Isles-N Atlantic-Iceland-Faeroes-Mediterranean-Antarctic
- C. alata* Brian, 1906  
@ *Phycis* spp.  
North Sea-Br. Isles-Eur. Atlantic-Mediterranean
- C. stellata* (Krøyer, 1838)  
= *Anchorella stellata* Krøyer, 1838  
@ along rims of opercula of *Merluccius merluccius*  
North Sea-Br. Isles-Eur. Atlantic
- C. stichaei* (Krøyer, 1863)  
@ *Lumpenus*, *Stichaeus* etc.  
\* Not mentioned by Kabata, 1992  
Iceland
- Clavellodes* Wilson, 1915
- C. rugosa* (Krøyer, 1837)  
= *Anchorella rugosa* Krøyer, 1837 (TM)  
@ *Anarhichas* spp.  
North Sea-Br. Isles-Faeroes-Iceland-Greenland-Norw. Sea-Barents Sea
- Clavellisa* Wilson, 1915 (TM *C. spinosa* Wilson, 1915)
- C. emarginata* (Krøyer, 1837)  
= *Anchorella emarginata* Krøyer, 1837  
= *Anchorella ovalis* Krøyer, 1837  
@ *Alosa* spp., (*Caspialosa* & *Clupeonella*)  
North Sea-Br. Isles-Mediterranean-Black & Azov Seas
- C. scombri* (Kurz, 1877)  
= *Anchorella scombri* Kurz, 1877  
@ first gill arch of *Scomber scombrus* & *S. colias*  
Skagerrak-North Sea-Br. Isles-Mediterranean-SW Africa
- Alella* Leigh-Sharpe, 1925
- A. pagelli* (Krøyer, 1863)  
= *Anchorella pagelli* Krøyer, 1863 (TM)  
= *Anchorella canthari* Heller, 1865  
= *Anchorella centrodonii* van Beneden, 1871  
@ Sparidae, e.g. *Pagellus* spp. & *Spondyliosoma cantharus*  
North Sea-Br. Isles-Eur. Atlantic-Mediterranean
- Advena* Kabata, 1979
- A. paradoxa* (van Beneden, 1851)  
= *Anchorella paradoxa* van Beneden, 1851 (TM)  
@ *Scomber scombrus*  
Skagerrak, Belgium-Br. Isles-Ireland-Mediterranean
- Eubrachiella* Wilson, 1915
- E. mugilis* Kabata, Raibaut & Ben Hassine, 1971  
@ *Liza aurata*  
Mediterranean
- Herpyllobiidae Hansen, 1892  
@ Polynoids
- Herpyllobius* Steenstrup & Lütken, 1861  
= *Silenium* Krøyer, 1863
- H. arcticus* Steenstrup & Lütken, 1861  
= *H. crassirostris* M. Sars, 18??  
@ On elyptrophores only in the anterior half of *Harmothoe* spp., *Gattyana* spp. & *Austrolaenilla mollis*  
Øresund-Bohuslän-Oslofjord-Arctic Seas-Iceland-Faeroes-Plymouth
- H. polynoies* (Krøyer, 1863)  
@ On dorsal surface of prostomium of *Harmothoe* spp., *Gattyana cirrosa*, *G. amondseni*, *Eunoe nodosa*, *Austrolaenilla mollis*, *Gaudichaudius iphionellodes*, *Antinoe sarsi* & *Antinoe badia*  
Øresund-Bohuslän-North Sea-N Norway-Arctic Seas-Iceland-Faeroes, NE USA, Alaska, NE Pacific
- Eurysilenium* M. Sars, 1870
- E. truncatum* M. Sars, 1870  
@ On dorsal surface of *Gattyana cirrosa* & *Harmothoe imbricata*  
Øresund-Bohuslän-Oslofjord-"Norway"
- Phallusiella* Leigh-Sharpe, 1926
- P. psalliota* Leigh-Sharpe, 1926  
@ On dorsal side of *Harmothoe extenuata*  
Plymouth
- P. vera* Leigh-Sharpe, 1926  
@ On dorsal surface of *Malmgrenia castanea*  
Plymouth
- Nicothoidae Dana, 1852  
= Choniostomatidae Hansen, 1886  
@ Crustaceans
- Rhizorhina* Hansen, 1892
- R. ampeliseae* Hansen, 1892  
@ gill sacs of *Ampelisca brevicornis* & *A. diadema*, with only egg-sacs protruding  
Kattegat-Norway-Clyde Sea-Engl. Channel, Black Sea
- Nicothoe* Audouin & Milne-Edwards, 1826
- N. astaci* Audouin & Milne-Edwards, 1826  
@ On gills of *Homarus gammarus* & *Palinurus vulgaris*  
Bohuslän-North Sea-Br. Isles-Orkneys-Ireland-Mediterranean
- Choniomyzon* Pillai, 1962  
@ decapods  
\* No Scandinavian species
- Hadrothoe* Humes, 1975  
@ decapods  
\* No Scandinavian species
- Paranicothoe* Carton, 1970 (TM *Paranicothoe cladocera* Carton, 1970 - Mozambique Channel)  
@ on epicarideans parasitizing penaeids  
\* Warm regions of the Indopacific only.
- Pseudonicothoe* Avdeev & Avdeev, 1978 (TM *Pseudonicothoe branchialis* Avdeev & Avdeev, 1978)  
@ on the gills of pandalid decapods and on epicarideans parasitizing such shrimps  
\* No Scandinavian species
- Choniosphaera* Connolly, 1929
- C. maenadis* (Bloch & Gallien, 1933)  
@ Among eggs of *Carcinus maenas* & *Maia squinado*  
E England, Isle of Man, N France
- Choniostoma* Hansen, 1886
- C. mirabile* Hansen, 1886  
@ *Lebbeus polaris*, *Eualus gaimardi* & ?*Spirontocaris* spp.  
\* Not mentioned by Gotto, 1993  
?N Norway-Kara Sea-Iceland-E Greenland
- C. hanseni* Giard & Bonnier, 1889  
@ *Spirontocaris* spp.  
\* Not mentioned by Gotto, 1993  
?N Norway-Kara Sea-Iceland-W Greenland
- C. rotundatum* Stock, 1958  
@ *Spirontocaris lilljeborgi*  
Skagerrak, North Sea
- Mysidion* Hansen, 1897
- M. commune* Hansen, 1897  
@ In marsupium of *Erythroops* spp. & *Parerythroops obesus* G.O. Sars  
Norway, SW Scotland
- M. abyssorum* Hansen, 1897  
@ *Erythroops abyssorum* G.O. Sars  
\* Not mentioned by Gotto, 1993  
Norway
- Aspidoecia* Giard & Bonnier, 1889
- A. normani* Giard & Bonnier, 1889  
@ Externally attached to the body of *Erythroops* spp.  
Norway, SE & SW Scotland & SW England
- Stenothocheres* Hansen, 1897
- S. sarsi* Hansen, 1897  
@ *Stenothoe marina*  
S or W Norway
- S. egregius* Hansen, 1897  
@ *Metopa bruzelii* (Goes)  
Greenland
- S.* sp.  
= *S. egregius* T. Scott, 1904, ?non Hansen, 1897  
@ *Metopa borealis*  
Aberdeen
- Homoeoscelis* Hansen, 1897
- H. minuta* Hansen, 1897  
@ *Diastylis lucifera* (Krøyer)  
Øresund-Kattegat, SE Scotland
- H. frigida* Hansen, 1923  
@ *Diastylis polaris* G.O. Sars  
\* Not mentioned by Gotto, 1993  
S of Jan Mayen, deep water
- H. longipes* Hansen, 1923  
@ *Leptostylis villosa* G.O. Sars  
\* Not mentioned by Gotto, 1993  
W of S Iceland
- H. mediterranea* Hansen, 1897  
@ Branchial cavity of *Iphinoe trispinosa*  
SW England, Mediterranean
- Sphaeronella* Salensky, 1868
- S. atyli* Hansen, 1897  
@ Marsupium of *Atylus swammerdami* (Milne-Edwards)  
Kattegat, W Ireland
- S. danica* Hansen, 1897  
@ Marsupium of *Corophium crassicornis* Bruzelius, *Apherusa bispinosa* & *Erichthonius punctatus*

Kattegatt-SE Scotland, SW England, W Ireland, Engl. Channel

*S. vestita* Hansen, 1897

@ *Microprotopus maculatus* Norman  
\* Not mentioned by Gotto, 1993  
Denmark

*S. paradoxa* Hansen, 1897

@ Marsupium of *Bathyporeia* spp. & *Periculodes longimanus* (Bate)  
Kattegatt, Netherlands, NE & SE Scotland, NE & E England, E & SW Ireland

*S. abyssi* Hansen, 1897

@ *Asityra abyssi* Boeck  
\* Not mentioned by Gotto, 1993  
N Norway

*S. intermedia* Hansen, 1897

@ *Bruzelia typica*  
\* Not mentioned by Gotto, 1993  
Norway

*S. gitanopsidis* Hansen, 1897

@ *Gitanopsis arctica* G.O. Sars  
\* Not mentioned by Gotto, 1993  
N Norway

*S. giardii* Hansen, 1897

@ *Protomedea fasciata* Krøyer  
\* Not mentioned by Gotto, 1993  
Kattegatt

*S. longipes* Hansen, 1897

@ Marsupium of *Ampelisca tenuicornis* Liljeborg  
Kattegatt-SE Scotland, NE England, W Ireland, Engl. Channel

*S. amphiloichi* Hansen, 1897

@ *Paramphilochooides odontonyx* & *Amphilochox neapolitanus*  
Denmark, Scotland

*S. dulichiae* Hansen, 1897

@ *Dulichia monacantha* Metzger  
\* Not mentioned by Gotto, 1993  
Denmark

*S. frontalis* Hansen, 1897

@ Marsupium of *Ampelisca macrocephala* Liljeborg & *A. tenuicornis*  
Kattegatt, Clyde Sea

*S. microcephala* Giard & Bonnier, 1893

@ Marsupium of *Ampelisca typica* Bate & *A. tenuicornis* Liljeborg  
Kattegatt & W Bretagne

*S. modesta* Hansen, 1897

@ *Eudorella emarginata* (Krøyer)  
\* Not mentioned by Gotto, 1993  
Denmark

*S. dispar* Hansen, 1897

@ Marsupium of *Eudorella truncatula* (Bate)  
Kattegatt, SW England

*S. insignis* Hansen, 1897

@ *Diastylis cornuta* Boeck & *D. laevis* Norman  
\* Not mentioned by Gotto, 1993  
Denmark

*S. affinis* Hansen, 1897

@ *Amira maculosa* Leach  
\* Not mentioned by Gotto, 1993  
Norway (Herøy)

*S. pleurogonii* Hansen, 1923

@ *Pleurogonium spinosissimum*  
\* Not mentioned by Gotto, 1993  
Iceland

*S. leuckartii* Salensky, 1868

= *S. elegantula* Hansen, 1897  
@ In marsupium of *Cheirocratus sundevalli* (Rathke), *Microdeutopus gryllotalpa* Costa & *Aora typica* Krøyer, *A. gracilis*, *Lembos longipes*, *Apherusa hispidosa*, *Corophium bonelli* & *C. volutator*  
Kattegatt-E England, Clyde Sea-Ireland-Engl. Channel-Naples

*S. irregularis* Hansen, 1897

@ *Metopa rubrovittata* G.O. Sars  
\* Not mentioned by Gotto, 1993  
?Denmark

*S. minuta* T. Scott, 1904

= *S. pikei* Green, 1958  
@ *Periculodes longimanus* & *Pontocrates arenarius*  
NE & SE Scotland, SW Ireland, Engl. Channel

*S. valida* Green, 1958

= *S. minuta* var. *valida* T. Scott, 1905  
@ Marsupium of *Megamphopus cornutus* Norman  
Norfolk-Firth of Forth

*S. melphidippae* Hansen, 1923

@ *Melphidippa borealis* Boeck  
\* Not mentioned by Gotto, 1993  
NW Iceland

*S. monothrix* (Bowman & Kornicker, 1967)

@ Marsupium of *Cylindroleberis mariae*  
SE Scotland, NE USA

*S. aeginae* Hansen, 1923

@ *Aegina echinata*  
\* Not mentioned by Gotto, 1993  
between Iceland & Faeroes

*S. decorata* Hansen, 1897

@ *Diastylis rathkei*  
\* Not mentioned by Gotto, 1993  
Kara Sea-W Greenland-Davis Strait

*S. devosae* Stock, 1960

@ Marsupium of *Corophium arenarium*  
Netherlands

*S. ecaudata* Stock, 1960

@ Marsupium of *Corophium arenarium*  
Netherlands

*S. rotundata* Hansen, 1923

@ *Hemilamprops cristata* G.O. Sars

\* Not mentioned by Gotto, 1993  
off SW Faeroes

*S. munnopsidis* Hansen, 1897

@ *Munnopsis typica* M. Sars  
\* Not mentioned by Gotto, 1993  
Kara Sea-E Greenland

*S. munnae* Hansen, 1923

@ *Munna spinifera* Hansen  
\* Not mentioned by Gotto, 1993  
off E Iceland, deep water

*S. nannonisci* Hansen, 1923

@ *Nannoniscus simplex* Hansen  
\* Not mentioned by Gotto, 1993  
off W Iceland, deep water

*S. callisomae* T. Scott, 1904

@ ?  
\* Inadequately described according to Gotto, 1993  
Britain

*S. cluthae* T. Scott, 1904

@ ?  
\* Inadequately described according to Gotto, 1993  
Britain

*S. pygmaea* T. Scott, 1904

@ ?  
\* Inadequately described according to Gotto, 1993  
Britain

*S. vararensis* T. Scott, 1905

@ ?  
\* Inadequately described according to Gotto, 1993  
Britain

Melinnacheridae Bresciani & Lützen, 1975

= Saccopsidae Lützen, 1964  
@ Terebellomorph Polychaets  
\* Not recognized by Huys & Boxshall 1991

*Melinnacheres* M. Sars, 1870

= *Saccopsis* Levisen, 1878  
= *Oestrella* McIntosh, 1885

*M. terebellidis* (Levisen, 1878)

@ *Terebellides stroemi*  
\* Not mentioned by Gotto, 1993  
Greenland-N America

*M. steenstrupi* (Bresciani & Lützen, 1961)

@ Blood-feeding from the base of the gills of *Terebellides stroemi*  
Øresund-Bohuslän-SE Scotland-Iceland-E & W Greenland-E Canada, Mediterranean

*M. ergasiloides* M. Sars, 1870

@ On parapodia of the last thoracic segments of *Melinna cristata*  
N Kattegatt-Bohuslän-Oslofjord, Massachusetts

*M. alleni* (Brumpt, 1987)

= *Saccopsis alleni* Brumpt, 1987  
@ *Polycirrus aurantiacus* Grube  
\* Not mentioned by Gotto, 1993  
?France

Asterocheridae Giesbrecht, 1899

= Ascomyzontidae Thorell, 1859?  
@ Sponges, Anthozoans, Echinoderms, Ascidians

*Asterocheres* Boeck, 1859

= *Ascomyzon* Thorell, 1859  
= *Echinocheres* Claus, 1889

*A. siphonatus* Giesbrecht, 1899

= *A. thorelli* Bresciani & Lützen, 1962  
= *A. liljeborgi* (Thorell, 1859), non Boeck, 1859  
= *Ascomyzon liljeborgi* : G.O. Sars, 1918, non *Asterocheres liljeborgi* Boeck, 1859  
@ *Corella parallelogramma*, *Ascidia virginia* & *Anseropoda membranacea*  
Bohuslän-S Norway-Norw. Sea-Iceland-?Mediterranean

*A. echincola* (Norman, 1868)

= *Ascomyzon parvum* : G.O. Sars, 1918, non *Asterocheres parvus* Giesbrecht, 1897  
@ *Haliclona oculata*, *Haliclondria* spp., *Grantia compressa*, *Ciocalypa penicillus*, ?*Echinus esculentus* & other sponges  
Bohuslän-S Norway-North Sea-Shetlands-Br. Isles, N Ireland, Engl. Channel

*A. flustrae* Ivanenko & Smurov, 1997

@ *Flustra foliacea*  
White Sea

*A. parvus* Giesbrecht, 1897

@ *Clathrina lacunosa*, *C. primordialis* & *Spongelia fragilis*  
N France, Mediterranean

*A. latus* (Brady, 1880)

@ *Clathrina primordialis* & perhaps algae  
Skagerrak, S Norway, W Britain, SW Ireland, Mediterranean

*A. boeckii* (Brady, 1880)

non = *A. boeckii* : Giesbrecht, 1899 (which is the Mediterranean *A. complexus* Stock, 1960)  
@ *Haliclondria panicea*, *Haliclona oculata*  
SW & W Norway-Spitsbergen-Iceland-Br. Isles-W & SW Ireland

*A. liljeborgi* Boeck, 1859

= *Ascomyzon asterocheres* G.O. Sars, 1918  
@ *Henricia*, *Asterias*, *Crossaster*, *Luidia*, *Antedon* etc.  
Bohuslän-S & W Norway-North Sea-Britain-W Ireland-Engl. Channel-Davis Strait

*A. violaceus* (Claus, 1889)

@ regular & irregul. Echinoidea, Ophiuroids, Asteroidea & *Antedon*  
Bohuslän-North Sea-W Norway-Shetlands-Br. Isles-France-Mediterranean

*A. suberitis* Giesbrecht, 1897

@ *Suberites domuncula*  
SW England, ?SW Scotland, Mediterranean

*A. stimulans* Giesbrecht, 1897

@ ?  
N France, Mediterranean

*A. simulans* T. Scott, 1894

@ sponges  
S Norway, SW Scotland, SW Ireland

- A. ellisi* Hamond, 1968  
@ *Halichondria panicea*, *Ciocalypa penicillus*  
Norfolk
- A. renaudi* Canu, 1891  
@ *Reniera* sp.  
\* Doubtful if this is placed in correct genus according to Gotto, 1993  
N France
- A. simplex* ??, 1???  
@ *Clathrina clathrus* & *Clathrina lacunosa*  
\* Not mentioned by Gotto, 1993  
Banyuls-sur-mer
- A. mucronipes* ??, 1???  
@ *Oscarella lobularis*  
\* Not mentioned by Gotto, 1993  
Banyuls-sur-mer
- A. corneliae* ??, 1???  
@ *Clathrina primordialis*  
\* Not mentioned by Gotto, 1993  
Banyuls-sur-mer
- Dermatomyzon* Claus, 1889
- D. nigripes* (Brady & Robertson, 1876)  
@ ? (so far found together with *Laminaria* - or free)  
Bohuslän-N Norway-Arctic Seas-Faeroes-Shetlands-Br. Isles-NE & SW Ireland, N France, Mediterranean
- Rhynchomyzon* Giesbrecht, 1895
- R. purpurocinctum* (Scott, 1893)  
@ ?  
Aalesund-Christiansund-Br. Isles-off Irish coast, Mediterranean
- R. rubrovittatum* G.O. Sars, 1918  
@ ?  
\* Not mentioned by Gotto, 1993  
W Norway (Kallevaag)
- R. falco* Giesbrecht, 1895  
@ *Clathrina contorta*  
\* Not mentioned by Gotto, 1993  
S Norway-Mediterranean
- Acontiophorus* Brady, 1880
- A. scutatus* (Brady & Robertson, 1873)  
@ *Spongelia fragilis* var. *ramosa*  
W Norway, Faeroes, Ireland, Britain, N France, Mediterranean
- A. ornatus* (Brady & Robertson, 1876)  
= *A. armatus* Brady, 1880  
@ ?  
S Norway (Risør)-NE England, Davis Strait, Mediterranean
- A. antennatus* Hansen, 1923  
@ *Clathrina lacunosa* & *Oscarella lobularis*  
around Greenland-?Mediterranean
- Orecturus* Humes, 1992 (™ *O. grandisetiger* Humes, 1992 - Madagascar @ *Antipathes*)
- Collocheres* Canu, 1893  
= *Leptomyzon* G.O. Sars, 1915
- C. gracilicauda* (Brady, 1880)  
@ *Ophiolithrix fragilis*, (*Antedon* spp. & *Echinus esculentus*)  
Bohuslän-S Norway-Br. Isles-NE & SW Ireland-Roscoff-Mediterranean
- C. gracilipes* Stock, 1966  
= *C. gracilicauda* : Canu, 1893, : G.O. Sars, 1915, non (Brady, 1880)  
@ ? (possibly a crinoid)  
S Norway (Risør), Boulogne
- C. elegans* A. Scott, 1896  
@ arm base of *Ophiocomina nigra*  
W Norway-W Br. Isles
- Scottomyzon* Giesbrecht, 1897
- S. gibberum* (T. & A. Scott, 1894)  
@ making galls in rosettes of pedicellariae in *Asterias rubens*, *Marthasterias glacialis* & *Spatangus purpureus*  
Belt Sea-Bohuslän-S & SW Norway-North Sea-Br. Isles-N France
- Mesocheres* Norman & Scott, 1905
- M. anglicus* Norman & Scott, 1905  
@ ?  
Oslofjord-SW England
- Scottocheres* Giesbrecht, 1897
- S. elongatus* (T. & A. Scott, 1894)  
@ *Dysidea tupa*, *Ulosa stiposa*, *Sigmatosella annexa*, *Halichondria* sp. & *Haliclona* sp.  
W Norway-North Sea-Br. Isles-SW Ireland-N France-Mediterranean
- Brychiopontiidae Humes, 1974
- Brychiopontius* Humes, 1974
- B. falcatus* Humes, 1974  
@ *Oneirophanta mutabilis* Theel  
off W Ireland in ca 4430m
- Myzopontiidae G.O. Sars, 1915  
@ ?Sponges  
\* Not recognized in Huys & Boxshall 1991
- Myzopontius* Giesbrecht, 1895
- M. pungens* Giesbrecht, 1895  
@ *Clathrina* spp., especially *C. contorta* ? & *Stichopus regalis* (Cuvier)  
S Norway (Korshavn)-Arctic Seas-Mediterranean
- Neopontius* T. Scott, 1898
- N. angularis* T. Scott, 1898  
@ ?  
W Norway-SW Scotland
- Dyspontiidae Giesbrecht, 1895  
@ sponges, scleractinians, ascidians  
\* Not represented as a family of its own in Gotto, 1993, but included in Artotrogidae Brady, 1880
- \* Not recognized in Huys & Boxshall 1991
- Dyspontius* Thorell, 1859
- D. striatus* Thorell, 1859  
@ "small ascidians"  
?Öresund-Bohuslän-N Norway-Br. Isles-Ireland, N France, Mediterranean
- D. curticaudatus* T. Scott, 1905  
@ ?  
SE Scotland
- Cryptopontius* Giesbrecht, 1895
- C. brevifurcatus* (Giesbrecht, 1895)  
@ *Echinus esculentus*  
S Norway (Korshavn)-Mediterranean
- Arctopontius* G.O. Sars, 1915  
\* Not mentioned by Gotto, 1993
- A. expansus* G.O. Sars, 1915  
@ ?  
N Norway (Hammerfest)-Davis Strait
- Bradyopontius* Giesbrecht, 1895
- B. magniceps* (Brady, 1880)  
@ ? , possibly sponges  
Aalesund-Hammerfest-Br. Isles, Ireland, N France
- B. papillatus* (T. Scott, 1888)  
@ probably *Pomatoceros triquetus*  
Kieler Bucht-Öresund-Bohuslän-W Norway-Br. Isles-Mediterranean
- B. major* G.O. Sars, 1918  
@ ?  
\* Not mentioned by Gotto, 1993  
Aalesund-Trondheim
- B. caudatus* G.O. Sars, 1918  
@ ?  
\* Not mentioned by Gotto, 1993  
NW Norway (Nordland)
- B. unidens* Hansen, 1923  
@ ?  
\* Not mentioned by Gotto, 1993  
E Greenland
- Cribropontius* Giesbrecht, 1899
- C. normani* (Brady & Robertson, 1876)  
@ ?  
W Norway-Br. Isles-Madeira
- Metapontius* Hansen, 1923  
\* Not mentioned by Gotto, 1993
- M. latispinis* Hansen, 1923  
@ ?  
Faeroes, deep water
- Artotrogidae Brady, 1880  
@ Nudibranchs
- Artotrogus* Boeck, 1859
- A. orbicularis* Boeck, 1859  
@ Nudibranchs ("Doridae")  
Kieler Bucht, Bohuslän-S, W & N Norway-Kara Sea-Br. Isles-E Ireland
- Undescribed family with presumed Artotrogid affinities
- Apodomyzon* Stock, 1970
- A. brevicorne* Stock, 1970  
@ In tissue of *Haliclona indistincta* (Bowerbank)  
Roscoff
- A. longicorne* Stock, 1970  
@ In tissue of *Haliclona indistincta* (Bowerbank)  
Roscoff
- Cancerillidae Giesbrecht, 1897  
@ Ophiuroids
- Parartotrogus* T. & A. Scott, 1893
- P. arcticus* T. Scott, 1901  
@ ?  
\* Not mentioned by Gotto, 1993  
NW Norway (Nordland)-off E Greenland-Arctic Seas
- P. richardi* T. & A. Scott, 1893  
@ *Ophiura albida* & ?*O. ophiura* ("Ophioglypha lacertosa")  
W Norway-SE & SW Scotland-Isle of Man-Mediterranean
- Cancerilla* Dalyell, 1851
- C. tubulata* Dalyell, 1851 (™)  
= *Caligidum vagabundum* Claus, 1889  
= ?? *Phitiopsis emilii* Giard, 1879  
@ Generally on oral disk surface, near arms, of *Axiognathus squamatus*, *Ophiocomina nigra*, ?*Ophiolithrix fragilis* & *Ophiostila aranea*  
S Baltic-Bohuslän-S Norway-Trondheim-Barents Sea-Br. Isles-N & W France-Mediterranean
- Micropontiidae Gooding, 1957  
@ Irregular Echinoids
- Micropontius* Gooding, 1957
- M. ovoides* Gooding, 1957  
@ *Spatangus purpureus*, *Echinocardium* spp. & *Brissopsis lyrifera*  
Bohuslän-Br. Isles-N & W France-W Mediterranean
- Nanaspididae Humes & Cressey, 1959  
= Allantogynidae Changeux  
@ Holothurians
- Nanaspis* Humes & Cressey, 1959
- N. ninae* Bresciani & Lützen, 1962  
@ On surface of *Parastichopus tremulus*  
Bohuslän-Trondheim, off W Ireland in 480m

- Allantogynus* Changeux, 1958  
\* Not mentioned by Gotto, 1993
- Xenocoelomatidae Bresciani & Lützen, 1966
- Aphanodomus* Wilson, 1920  
= *Crypsidomus* Levinsen, 1878, non Guenther, 1864
- A. terebellae* (Levinsen, 1878)  
@ *Thelepus cincinnatus* & other Terebellidae  
\* Not mentioned by Gotto, 1993  
Iceland-Greenland-Kara Sea
- Xenocoeloma* Caullery & Mesnil, 1915
- X. allenii* (Brumpt, 1897)  
@ *Polycirrus caliendrum* Claparède  
Engl. Channel
- X. brumpti* Caullery & Mesnil, 1915  
\* May be conspecific with *X. allenii* (Brumpt); considered to be so by Gotto, 1993 with some doubt.  
\* *Polycirrus arenivorus* Caullery  
?Engl. Channel
- Spongiocnizontidae Stock & Kleeton, 1964
- Spongiocnizon* Stock & Kleeton, 1964
- S. petiti* Stock & Kleeton, 1964 (TM)  
@ *Hemimyscale columella*  
French Mediterranean
- S. vermiformis* Stock, 1967  
@ *Iophon hyndmani* var. *junis*  
French Mediterranean
- Sponginticolidae Topsent, 1928  
\* May possibly be related to fam. Spongiocnizontidae.
- Sponginticola* Topsent, 1928  
= *Clionophilus* Silén, 1963
- S. uncifer* Topsent, 1928 (TM - French Channel coast)  
= *Clionophilus vermularis* Silén, 1963 (TM of *Clionophilus* Silén, 1963 - Bohuslän)  
@ *Cliona celata*, *Sylopus coriaceus* & *Myscale maclenata*  
Bohuslän - NE Ireland - Engl. Channel, Mediterranean
- Calverocheidae Stock, 1968  
@ Echinoida
- Dinopontiidae Murnane, 1967  
@ Porifera & Actiniaria
- Dirivulitidae Humes & Dojiri, 1980  
@ vestimentiferans
- Dissonidae Yamaguti, 1963
- Entomolepididae Brady, 1899  
@ Porifera
- Hyponeoidae Heegaard, 1962
- Megapontiidae Heptner, 1968  
@ not known
- Naobranchiidae Yamaguti, 1939  
\* not recognized by Huys & Boxshall 1991.
- Pontoeciellidae Giesbrecht, 1895
- Rataniidae Giesbrecht, 1897
- Ratania* Giesbrecht, 1892
- R. atlantica* Farran, 1926  
off Ireland, Biscay
- R. flava* Giesbrecht, 1892  
deep Atlantic & Mediterranean
- Stellicomitidae Humes & Cressey, 1958
- Tanypleuridae Kabata, 1969
- ?Poecilo-?Siphonostomatoida not yet referred to any Family?
- Gonophysemia* Bresciani & Lützen, 1960  
\* Possibly another sp. in *Heterostigma reptans* Monniot & Monniot
- G. gullmarensis* Bresciani & Lützen, 1960  
@ *Ascidella aspersa*, *A. scabra* & ?*Distomus variolosus*  
Kattegat-Bohuslän, ?Mediterranean
- Axinophilus* Bresciani & Ockelmann, 1966  
\* May be related to Mycolidae?
- A. thyasirae* Bresciani & Ockelmann, 1966  
@ *Thyasira flexuosa* & *T. sarsi* (& in *T. gouldii* from Spitsbergen & W Greenland, if this is not another species)  
Öresund-Bohuslän-S Norway-Bergen-SE Scotland
- Phyllocola* Delamare-Deboutteville & Laubier, 1960
- P. petiti* Delamare-Deboutteville & Laubier, 1960  
@ *Eulalia expusilla* & other phyllocolidae  
Britain?
- MONSTRILLOIDA G.O. Sars, 1903
- Monstrillidae Giesbrecht, 1892  
@ not known for most species; Polychaeta & Mollusca
- Monstrilla* Dana, 1849 (TM *M. viridis* Dana, 1849) (publ. September 4) (n. cons. ICZN Op. 1869)  
= *Thaumatoessa* Krøyer, in Gaimard, 1842-1845  
\* Grygier, 1993 has submitted an application to ICZN to suppress this older name for purposes of the Principle of Priority.  
= *Thaumaleus* Krøyer, 1849 (publ. December 31) (n. cons. ICZN Op. 1869)
- M. longicornis* I.C. Thompson, 1890  
=? *Thaumatoessa typica* Krøyer, in Gaimard, 1842-45
- = *M. intermedia* Aurivillius, 1898, ?non Kriczagin, 1877 (Kriczagins name is suppressed for Law of Priority by ICZN Op. 1175)  
=? *M. clavata* G.O. Sars, 1921  
Skagerrak-S & W Norway-Br. Isles-Mediterranean, Maine
- M. longiremis* Giesbrecht, 1892  
Hvaler-Ofsfjord-NW Norway-North Sea-Cornwall-Mediterranean, Maine
- M. conjunctiva* Giesbrecht, 1902  
= *M. leucopis* G.O. Sars, 1921  
NW Norway-North Sea-Cornwall
- M. gracilicauda* Giesbrecht, 1892  
\* Male unknown; may be *M. anglica* Lubbock  
NW Norway-Arctic-Br. Isles-Engl. Channel-Mediterranean
- M. anglica* Lubbock, 1857  
\* Female unknown; may be *M. gracilicauda* Giesbrecht  
Br. Isles-Engl. Channel, Atlantic Canada, "almost world-wide"
- M. helgolandica* Claus, 1863  
= *M. canadensis* McMurrich, 1917 (male)  
@ *Brachystomia rissoides* (Hanley) & *Odosstomia scalaris* MacGillivray  
Skagerrak-North Sea-Norway (Stavanger, Christiansund & Nordland)-Cornwall-Mediterranean-Hudson Strait, Maine
- M. serricornis* G.O. Sars, 1921  
North Sea, Norw. Sea
- M. minuta* Isaac, 1974  
Isle of Man
- M. obesa* Isaac, 1975 (nomen nudum)  
Channel Islands
- Cymbasoma* J.C. Thompson, 1888  
= *Thaumaleus* : Auctt., non Krøyer, 1849  
@ Annelida
- C. rigidus* J.C. Thompson, 1888 (TM)  
= *Thaumaleus germanicus* Timm, 1896  
@ *Salmacina dysteri* (Huxley), *Polydora cilata* & *P. giardi* Mesnil  
S & W Norway-E North Sea-Br. Isles-Engl. Channel-Mediterranean
- C. thompsoni* (Giesbrecht, 1892)  
= *Thaumaleus thompsoni* Giesbrecht, 1892  
Belt Sea-S & W Norway-Engl. Channel-"more or less world-wide"
- C. longispinosus* (Bourne, 1890)  
= *Monstrilla longispinosa* Bourne, 1890  
Norway-German Bight-North Sea-Engl. Channel-"more or less world-wide"
- C. claparedi* (Giesbrecht, 1892)  
= *Thaumaleus claparedi* Giesbrecht, 1892  
North Sea-Mediterranean
- C. reticulatus* (Giesbrecht, 1892)  
= *Thaumaleus reticulatus* Giesbrecht, 1892  
North Sea-Mediterranean
- C. filigranarum* (Malaquin, 1896)  
= *Thaumaleus filigranarum* Malaquin, 1896  
\* Should perhaps be classified as a *Monstrilla* Dana?  
@ *Filigrana implexa* Berkeley  
Engl. Channel
- C. roscovita* (Malaquin, 1901)  
= *Haemocera roscovita* Malaquin, 1901  
= *Thaumaleus roscovita* : Auctt.  
\* Should perhaps be classified as a *Monstrilla* Dana?  
@ *Filigrana setosa* Langerhans  
British waters-?Roscoff
- C. frondipes* (T. Scott, 1904)  
\* Should the author be written within parenthesis?  
E Br. Isles
- C. rostratus* (T. Scott, 1904)  
= *Thaumaleus rostratus* T. Scott, 1904  
\* Male unknown; may be *Thaumaleus similirostratus* Isaac  
Shetlands
- C. similirostratus* (Isaac, 1974)  
= *Thaumaleus similirostratus* Isaac, 1974  
\* Female unknown; may be *T. rostratus* T. Scott  
Cornwall
- C. zetlandicus* (T. Scott, 1904)  
= *Thaumaleus zetlandicus* T. Scott, 1904  
Shetlands, Engl. Channel
- C. malaquini* Caullery & Mesnil, 1914  
= *Thaumaleus malaquini* : Auctt.  
@ *Syllis gracilis* Grube  
Engl. Channel
- C. pallidus* (Isaac, 1974)  
= *Thaumaleus pallidus* Isaac, 1974  
Cornwall
- C. tenuis* (Isaac, 1974)  
= *Thaumaleus tenuis* Isaac, 1974  
Cornwall
- C. striatus* (Isaac, 1974)  
= *Thaumaleus striatus* Isaac, 1974  
Cornwall-Channel Islands
- Monstrillopsis* G.O. Sars, 1921
- M. dubia* (T. Scott, 1904)  
Norway (Raunefjorden & outside Trondheimsfjord)-Scotland-Cornwall-Mediterranean, Maine
- M. sarsi* Isaac, 1974  
Cornwall-Bristol Channel
- M. gracilis* (Gurney, 1927)  
Bristol & Engl. Channels-Portugal-Suez-Red Sea
- Strilloma* Isaac, 1975 (nomen nudum)
- S. grandis* (Giesbrecht, 1891)  
= *Monstrilla grandis* Giesbrecht, 1891  
German Bight-North Sea-Engl. Channel-Cornwall-Norfolk, China
- S. scotti* Isaac, 1975  
= *Monstrilla grandis* : T. Scott, 1904, non Giesbrecht, 1891

- Clyde Sea
- S. lata* (Desai & Bal, 1963)  
Bombay, Firth of Clay
- Thaumatohessia* Giard, 1900
- T. armoricana* (Hesse, 1868)  
Brest
- MISOPHROIDA Gurney, 1933
- Misophriidae Brady, 1878
- Misophria* Boeck, 1865
- M. pallida* Boeck, 1865  
W Norway-Arctic Seas-Br. Isles-W Mediterranean
- Arcticomisophria* Martínez Arbizu & Seifried, 1996
- A. bathylaptevensis* Martínez Arbizu & Seifried, 1996 (TM - deep Laptev Sea)  
Laptev Sea in 2330-3237 m
- Expansophria* Boxshall & Iliffe, 1987
- E. dimorpha* Boxshall & Iliffe, 1987 (TM)  
Canaries
- Palpophria* Boxshall & Iliffe, 1987
- P. aestetha* Boxshall & Iliffe, 1987 (TM)  
Canaries
- Dimisophria* Boxshall & Iliffe, 1987
- D. cavernicola* Boxshall & Iliffe, 1987 (TM)  
Canaries
- Benthomisophria* G.O. Sars, 1909
- B. palliata* G.O. Sars, 1909  
?Mediterranean
- B.* sp?  
\* One bathypelagic N. Atlantic species described by Hulsemann & Grice, 1964
- MORMONILLOIDA Boxshall, 1979
- Mormonillidae Giesbrecht, 1892
- Mormonilla* Giesbrecht, 1891  
\* males of the two species below described by Huys, Boxshall & Böttger-Schnack 1992
- M. phasma* Giesbrecht, 1891  
off W Ireland-Bay of Biscay-W Mediterranean
- M. minor* Giesbrecht, 1891  
= *M. polaris* G.O. Sars, 1900  
= *M. atlantica* Wolfenden, 1905  
temperate Atlantic-Bay of Biscay-off W Ireland-Arctic Seas
- HARPACTICOIDA G.O. Sars, 1903
- Superornatiremidae Huys, 1997
- POLYARTHRA Lang, 1948
- Longipediidae Brady, 1880
- Longipedia* Claus, 1863
- L. coronata* Claus, 1863  
Bohuslän-S North Sea-Frierfjord/Langesundsfjord-Trondheim-Br. Isles-Mediterranean
- L. minor* T. & A. Scott, 1893  
= *L. pontica* : Auct., ?non Kriczagin, 1873  
Bohuslän-S North Sea-W Norway-Trondheim-Br. Isles-Mediterranean
- L. helgolandica* Klie, 1949  
= *L. minor helgolandica* Klie, 1949  
Helgoland, Norfolk
- L. scotti* G.O. Sars, 1903  
= *L. australica* Nicholls, 1941 (according to Wells 1980)  
Bohuslän-S North Sea-Trondheim-Br. Isles, Australia
- L. rosea* G.O. Sars, 1903 (nom. dub. according to Wells 1980)  
\* Not included in Huys & al. 1996  
Bergen-Trondheim-?Mediterranean
- L. weberi* A. Scott, 1909  
\* Not included in Huys & al. 1996, but redescribed by Wells 1980 and Itô 1980  
Norfolk-S Ireland-Roscoff-Mediterranean, Antarctic
- Canuellidae Lang, 1944
- Sunaristes* Hesse, 1867
- S. paguri* Hesse, 1867 (TM)  
@ Usually in shell inhabited by *Pagurus bernhardus*, *Clibanarius erythropus* & *Diogenes pugilator*  
?Öresund-Bohuslän-S Norway-German Bight-Br. Isles-W Ireland-Engl. Channel, Mediterranean, Black Sea
- Canuella* T. & A. Scott, 1893
- C. perplexa* T. & A. Scott, 1893 (TM)  
@ *Pagurus bernhardus* & *Maia squinado* - or often found free.  
Bohuslän-S Norway-S North Sea-Br. Isles-Engl. Channel-Mediterranean
- C. furcigera* G.O. Sars, 1903  
Bohuslän-Oslofjord-S Norway-German Bight-Iceland-Greenland-Mediterranean
- C.* sp. : Huys & al. 1996  
Belgium
- Canuellopsis* Lang, 1936
- C. swedmarki* Por, 1964  
off N Koster (in 150m)-Skagerrak-Bergen
- Brianola* Monard, 1926
- B. stebleri* Monard, 1926  
?Roscoff, W Mediterranean
- B.* sp. : Huys & al. 1996
- Cornwall
- Microcanuella* Mielke, 1994
- Ellucana* Sewell, 1940
- Parasunaristes* Fiers, 1982
- Intersunaristes* Huys, 1995
- Galapacanuella* Mielke, 1979
- Canuellina* Gurney, 1927
- Nathaniella* Por, 1984
- Ifanella* Vervoort, 1964
- Intercanuella* Becker & Schriever, 1979
- Coullana* Por, 1984
- Scottolana* Por, 1964
- Elanella* Por, 1984
- Echinosunaristes* Huys, 1995
- OLIGOARTHRA Lang, 1948  
\* not a monophyletic taxon
- Cancrincolidae Fiers, 1990  
@ gill chambers of aquatic Decapoda & land crabs
- Cancrincola* Wilson, 1913 (*C. jamaicensis* Wilson, 1913)
- Abscondicola* Fiers, 1990 (TM *A. humesi* Fiers, 1990 - New Guinea @ land crab)
- Hamondiidae Huys, 1990
- Hamondia* Huys, 1990 (TM *H. superba* Huys, 1990 - Australia @ sponge)
- MAXILLIPEDASPHAELEA Lang, 1948
- Cervinioidea G.O. Sars, 1903
- Aegisthidae Giesbrecht, 1892
- Aegisthus* Giesbrecht, 1891
- A. mucronatus* Giesbrecht, 1891  
= *Aegisthus atlanticus* Wolfenden, 1902  
\* "The only harpacticoid species for which bioluminescence has been demonstrated with confidence" according to Huys & Böttger-Schnack, 1994  
Faeroe Channel-off Ireland-Bay of Biscay-"Cosmopolitan, Planktonic"
- A. spinulosus* Farran, 1905  
off W Ireland, pelagic
- A. aculeatus* Giesbrecht, 1891  
Biscay
- Cerviniidae G.O. Sars, 1903, emend. Lang, 1948  
\* A deep water species family.
- Cerviniella* Smirnov, 1946
- Paracerviniella* Brodskaya, 1963
- Expansicervinia* Montagna, 1981
- Pontostratiotes* Brady, 1883
- Tonpostratiotes* Itô, 1982
- Stratiopontotes* Soyer, 1970
- Herdmaniopsis* Brodskaya, 1963  
= *Ameliotes* Por, 1969 (according to Itô 1982)
- Cerviniinae Brodskaya, 1963
- Eucanuella* T. Scott, 1900
- E. spinifera* T. Scott, 1900 (TM)  
Bohuslän-W Norway-Iceland-Shetlands-Scotland
- Cervinia* Brady, 1878 ex Norman MS  
= *Pseudocervinia* Brodskaya, 1963 (according to Montagna 1981)
- C. bradyi* Brady, 1878 ex Norman MS (TM)  
Bohuslän-W Norway-Br. Isles-Mediterranean
- C. synarthra* G.O. Sars, 1910  
Bohuslän-Lofoten-Arctic Seas-Iceland-Br. Isles
- C. pilosa* Lang, 1948  
\* Not mentioned in Huys & al. 1996  
Bohuslän
- Cerviniopsinae Brodskaya, 1963
- Hemicervinia* Lang, 1935  
= *Herdmania* I.C. Thompson, 1893, non Lahille, 1888
- H. stylifera* (I.C. Thompson, 1893)  
= *H. ryforsi* Lang, 1935 (TM)  
Bohuslän-Isle of Man
- Cerviniopsis* G.O. Sars, 1903
- C. clavicornis* G.O. Sars, 1903  
Bohuslän-Lofoten
- C. longicaudata* G.O. Sars, 1903  
W Norway
- Rotundiclipeidae Huys, 1988
- Rotundiclipeus* (TM *R. canariensis* Huys, 1988 - Canaries in anchialine caves)
- Ectinosomatoidea G.O. Sars, 1903

*Ectinosomatidae* G.O. Sars, 1903 (p.p.); Olofsson, 1917

*Tetanopsis* Brady, 1910  
\* Doubtful genus.

*Oikopus* Wells, 1967

*Ectinosomoides* Nicholls, 1945

*Bradyellopsis* Brian, 1924

*Peltobradya* Médioni & Soyer, 1967

*Sigmatidium* Giesbrecht, 1881  
\* According to Huys & al. 1996 are several unnamed species present in the area.

*S. difficile* Giesbrecht, 1881  
Kieler Bucht-Elbe Mouth-Aberdeen

*Pseudectinosoma* Kunz, 1935

*P. minor* Kunz, 1935  
= *Sigmatidium minor* : Autctt.  
\* Brackish marsh species.  
Kieler Bucht-Hiddensee

*Pseudobradya* G.O. Sars, 1904

*P. pulchella* G.O. Sars, 1920  
Öresund-Bohuslän-S Norway

*P. elegans* (T. & A. Scott, 1894)  
Bohuslän-W Norway-Scotland

*P. minor* (T. & A. Scott, 1894)  
Belt Sea-Helgoland-Trondheim-Arctic Seas-Br. Isles-?Roscoff

*P. similis* (T. & A. Scott, 1894)  
Bohuslän-Trondheim-Br. Isles-W France

*P. hirsuta* (T. & A. Scott, 1894)  
Bohuslän-SW Norway-Br. Isles

*P. fusca* (T. & A. Scott, 1894)  
SW Norway-Br. Isles

*P. acuta* G.O. Sars, 1904  
off Gullmarn, Trondheim-?Italy

*P. robusta* G.O. Sars, 1910  
SW Norway

*P. digitata* G.O. Sars, 1920  
S Norway

*P. leptognatha* G.O. Sars, 1920  
Bohuslän-S Norway

*P. scabriuscula* G.O. Sars, 1920  
S Norway

*P. attenuata* G.O. Sars, 1920  
Hällö, off N Koster, S Norway

*P. tenella* G.O. Sars, 1920  
S Norway

*P. parvula* G.O. Sars, 1920  
S Norway

*P. exilis* G.O. Sars, 1920  
Trondheim

*P. pygmaea* G.O. Sars, 1920  
SW Norway

*P. ambigua* G.O. Sars, 1920  
S Norway-Trondheim

*P. maxima* Lang, 1935  
Skagerrak

*P. beduina* Monard, 1935  
Elbe mouth-Bretagne-SW Mediterranean

*P. distinctum* Wells, 1968  
Scilly Islands

*P. rhea* Bodin, 1977  
W France

*Bradya* Boeck, 1872

*B. (Bradya) typica* Boeck, 1872  
?Kieler Bucht-Öresund-Bohuslän-Trondheim-Arctic Seas-Br. Isles

*B. (Bradya) scotti* G.O. Sars, 1920  
Bohuslän-Bergen-Br. Isles

*B. (Bradya) macrochaeta* G.O. Sars, 1920  
Bohuslän-Bergen

*B. (Bradya) furcata* G.O. Sars, 1920  
Bohuslän-S Norway

*B. (Bradya) congenera* G.O. Sars, 1920  
Bohuslän-S Norway, Isle of Man

*B. (Bradya) simulans* G.O. Sars, 1920  
S Norway, Britain

*B. (Bradya) proxima* T. Scott, 1912  
North Sea, South Georgia

*B. (Parabradya) dilatata* G.O. Sars, 1904  
Bohuslän-Bergen-Iceland

*B. (Parabradya) atlantica* Bodin, 1968  
Biscay

*Ectinosomella* G.O. Sars, 1910

*E. nitidula* G.O. Sars, 1910  
\* Pelagic? in 55-91 m  
SW Norway

*Microsetella* Brady & Robertson, 1873

*M. norvegica* (Boeck, 1865)  
= *Microsetella atlantica* Brady & Robertson, 1873  
Pelagic around all European coasts, fr. Arctic Seas-Mediterranean, e.g. S Baltic & Bergen

*M. rosea* (Dana, 1848)  
Pelagic fr. Kattegat-North Sea-Br. Isles-Bay of Biscay-Mediterranean

*Halophytophilus* Brian, 1918 (& 1917 (nom. nudum))  
= *Halophytophilus* Brian, 1921

*H. spiniornis* (G.O. Sars, 1920)  
Bohuslän-SE Norway

*H. similis* (Lang, 1948)  
Bohuslän, Scotland

*Kliesoma* Hincks & Schriever, 1985  
= *Kliella* Hincks & Schriever, 1983, non Schäfer, 1945

*K. triarticulatus* (Klie, 1949)  
= *Halophytophilus? triarticulatus* Klie, 1949  
Helgoland, Britain

*Arenosetella* C.B. Wilson, 1932  
= *Pararenosetella* Lang, 1948 (p.p.)

*A. germanica germanica* Kunz, 1937  
SW Baltic-German Bight-Aberdeen-Scilly Isles-Pontomediterranean

*A. tenuissima* (Klie, 1929)  
= *Ectinosoma tenuissima* Klie, 1929  
= *Hastigerella palpabra* Nicholls, 1935  
= *Arenosetella monensis* Moore, 1976  
Baltic-Sylt-North Sea-Brittany-Br. Isles

*A. littoralis* Bodin, 1977  
W France

*A. duriensis* Galhano, 1970  
Atl. Portugal

*A. smithi* (Perkins, 1956)  
= *Tetanopsis* Brady, 1910 *smithi* Perkins, 1956  
\* Specific validity questionable according to Huys & al. 1996  
England (Kent)

*A. medius* (Perkins, 1956)  
= *Tetanopsis* Brady, 1910 *medius* Perkins, 1956  
\* Specific validity questionable according to Huys & al. 1996  
England (Kent)

*Hastigerella* Nicholls, 1935  
= *Pararenosetella* Lang, 1948 (p.p.)

*H. abotti abotti* Lang, 1965  
?Irish Sea-W Mediterranean

*H. leptoderma* (Klie, 1929)  
Kieler Bucht-German Bight-Br. Isles-Portugal-W Mediterranean

*H. psammae* (Noodt, 1955)  
= *Pararenosetella psammae* Noodt, 1955  
W France-Scotland

*H. scheibeli* Mielke, 1975  
Sylt

*H. unisetosa* (Wells, 1965)  
= *Arenosetella unisetosa* Wells, 1965  
Scotland

*H. grandimandibularis* Wells, 1967  
?Atlantic Europe, Mozambique

*H. bozici* Soyer, 1974  
Finistère, French Mediterranean

*Halectinosoma* Lang, 1944

*H. erythrops* (Brady & Robertson, 1875)  
= *Ectinosoma gracile* : G.O. Sars, 1904, non T. & A. Scott, 1894  
W Norway-S North Sea-Br. Isles & Ireland-N France

*H. gracile* (T. & A. Scott, 1894)  
Br. Isles & Ireland

*H. longicorne* (T. & A. Scott, 1894)  
Scotland, Roscoff

*H. tenuireme* (T. & A. Scott, 1894)  
= *Pararenosetella tenuiremis* (T. & A. Scott, 1894)  
off Gullmarn, Hällö, Norway-Helgoland-Scotland-Scilly Isles-Roscoff

*H. oblongum* (Kunz, 1949)  
Helgoland

*H. sarsi* (Boeck, 1872)  
\* Sp. incertae sedis according to Clement & Moore, 1995  
Bohuslän-German Bight-N Norway-Arctic Seas-Br. Isles-N France

*H. chrySTALLI* (T. & A. Scott, 1894)  
= *H. propinquum* (T. & A. Scott, 1894)  
Bohuslän-N Norway-Arctic Seas-Scotland & Ireland

*H. herdmanni* (T. & A. Scott, 1896)  
\* First intermediate host for *Acanthogyrus (Acanthocephalus) lizae* Orecchia, Paggi & Radujkovic, 1988  
(Acanthocephala)  
Kieler Bucht-Bohuslän-Amrum-Trondheim-Br. Isles-Mediterranean & Black Sea

*H. armiferum* (T. & A. Scott, 1894)  
Bohuslän-Norway-Scotland

*H. neglectum* (G.O. Sars, 1904)  
Bohuslän-S & N Norway-Arctic Seas-?Faeroes-Iceland & Greenland; ?Mediterranean

*H. elongatum* (G.O. Sars, 1904)  
\* Possibly a synonym of *H. herdmanni* (T. & A. Scott) sensu Por, 1964  
Kieler Bucht-Bohuslän-S Norway-S North Sea-Trondheim

*H. clavatum* (G.O. Sars, 1920)  
= *Ectinosoma clavatum* G.O. Sars, 1920  
off Gullmarn, Kosterfjord, S Norway-Arctic Seas

*H. brunneum* (Brady, 1905)  
Northumberland

- H. proximum* (G.O. Sars,1919)  
Öresund-Bohuslän-Trondheim
- H. angulifrons* (G.O. Sars,1919)  
S Norway
- H. tenerum* (G.O. Sars,1920)  
Bohuslän-S Norway-Scilly Isles
- H. curticorne* (Boeck,1872)  
Finland & Askö-Bohuslän-S North Sea-S & N Norway-Arctic Seas-Br. Isles-W Mediterranean
- H. gothicops* (Giesbrecht,1881)  
= *Ectinosoma pygmaeum* T. & A. Scott,1894  
Kieler Bucht-Bohuslän-German Bight-S Norway-Trondheim-Br. Isles-Roscoff
- H. mixtum* (G.O. Sars,1904)  
Belt Sea-Bohuslän-Oslofjord; ?Mediterranean
- H. brevirostre* (G.O. Sars,1904)  
Bohuslän-S Norway, Scotland
- H. abrau* (Kricagin,1877)  
= *Bradya edwardsi* Richard,1890  
Freshwater species that tolerates brackish water; N Baltic-Kiel & German Bight
- H. distinctum* (G.O. Sars,1920)  
Bohuslän-S Norway
- H. finmarchium* (T. Scott,1903)  
Belt Sea-North Norway-Spitsbergen & Arctic Seas
- H. spinicauda* Wells,1961  
Scilly Islands
- H. pterinum* Moore,1974  
Isle of Man-W France
- H. paraspinicauda* Bodin,1979  
W France
- H. gascognense* Bodin,1968  
Bay of Biscay
- H. canaliculatum* (Por,1964)  
= *Ectinosoma canaliculatum* Por,1964  
Atlantic Europe
- H. cooperatum* Bodin, Bodiou & Soyier,1972  
W & S France, Biscay
- H. valeriae* Soyier,1973  
?Atlantic Europe & French Mediterranean
- H. argyllensis* Clement & Moore,1995  
Scotland
- H. crenulatum* Clement & Moore,1995  
Scotland
- H. denticulatum* Clement & Moore,1995  
Scotland
- H. pseudosarsi* Clement & Moore,1995  
Scotland
- Ectinosoma* Boeck,1865
- E. melaniceps* Boeck,1865  
Kieler Bucht-Bohuslän-North Sea-W Norway-Arctic Sea-Iceland-Br. Isles-Mediterranean & Black Sea
- E. normani* T. & A. Scott,1894  
Bohuslän-Sylt-?Helgoland-N Norway-Arctic Seas-Br. Isles-Mediterranean
- E. tenuipes* T. & A. Scott,1894  
Bohuslän-S Norway-Br. Isles & Ireland
- E. compressum* G.O. Sars,1920  
Kieler Bucht-Bohuslän-S Norway-Helgoland
- E. obtusum* G.O. Sars,1920  
Kieler Bucht-S Norway-Wales
- E. reductum* Bocic,1954  
?Helgoland
- E. reductum listensis* Mielke,1975  
Sylt
- E. dentatum* Steuer,1940  
?Europe, SE Mediterranean
- Noodtiella* Wells,1965  
= *Noodtia* Lang,1965
- N. gracile* Mielke,1975  
Sylt
- N. lusitanica* Wells,1965  
Portugal
- Lineosoma* Wells,1965
- L. iscensis* Wells,1965  
Exmouth
- Neobradoidea Olofsson,1917
- Chappuisiidae Chappuis,1940  
\* Freshwater species only.
- Phyllognathopodiidae Gurney,1932  
\* Freshwater species only.
- Neobradidae Olofsson,1917  
= Marsteiniidae Drzymiski,1969
- Neobradia* T. Scott,1892
- N. pectinifera* T. Scott,1892 (TM)  
Kattegatt-Bohuslän-Bergen-Br. Isles
- Marsteinia* Drzymiski,1968
- M. typica* Drzymiski,1968  
Bergen
- M. similis* Drzymiski,1968  
Bergen
- Antarcticobradia* Huys,1987
- Darcythompsoniidae Lang,1936
- Darcythompsonia* T. Scott,1906
- D. fairliensis* (T. Scott,1899)  
= *D. scotti* Gurney,1920  
W Norway-Shetlands-Scotland-Ireland
- D. neglecta* Redeke,1953  
Netherlands
- Leptocaris* T. Scott,1899  
= *Horsicella* Gurney,1920
- L. brevicornis* (van Douwe,1904)  
Freshwater species which may be found in Baltic brackish water, German North Sea coast
- L. biscayensis* (Noodt,1955)  
= *L. adriatica* Petkovski,1955  
W France-Mediterranean
- L. trisetosus* (Kunz,1935)  
Finland-Kieler Bucht-Amrum
- L. ignavus* (Noodt,1953)  
= *Horsicella ignava* Noodt,1953  
\* May be a synonym of *L. minutus* T. Scott,1899  
Belt Sea-S Ireland
- L. minutus* T. Scott,1899 (TM)  
W Scotland, Devon
- Kristensenia* Por,1983
- EXANECHENTERA Lang,1948
- Tachidioidea Boeck,1864
- Euterpinidae Brian,1921
- Euterpina* Norman,1903
- E. acutifrons* (Dana,1848)  
= *Laophonte sagenarum* Oliveira,1945 (male)  
= *Tachidius hydrobiologicus* Oliveira,1945 (female)  
@ *Astropecten irregularis*  
\* Pelagic, with worldwide distrib. in warm-boreotemperate seas  
Bohuslän-North Sea-SW Norway-Br. Isles-Mediterranean
- Thompsonuliidae Lang,1948
- Thompsonula* T. Scott,1905
- T. hyaenae* (L.C. Thompson,1889)  
Kieler Bucht, German Bight-Br. Isles, Algeria
- Caribbula* Huys & Gee,1990 (TM *Thompsonula hyaenae elongata* (Gee,1988))
- Tachidiidae Boeck,1865  
= Microarthridioninae Lang,1948
- Cithadius* Bowman,1972
- Tachidius* Liljeborg,1853
- T. (Tachidius) discipes* Giesbrecht,1881  
Åland & Finland-Askö-Öresund-Belt Sea-German North Sea coast-Oslofjord-N Norway-Spitsbergen-Iceland-Br. Isles-N France, Italy
- Geeopsis* Huys,1996
- G. incisipes* (Klie,1913)  
= *Tachidius incisipes* Klie,1913  
= *Tachidius longicornis* Olofsson,1917  
Elbe & Weser mouths-Norfolk-Spitsbergen-Novaja Semlja
- Microarthridion Lang,1944
- M. littorale* (Poppe,1881)  
= *T. crassicornis* T. Scott,1893  
S Finland-Öresund, N Norway-Novaja Semlja, Br. Isles-S North Sea coasts-N France
- M. reductum* (Monard,1935)  
= *Microarthridion perkinsi* Bodin,1970  
Galway, Roscoff
- M. fallax* Perkins,1956  
Belt Sea-SE North Sea-Kent-Wales
- M. berberum* (Monard,1936)  
Portugal
- Harpacticidae Dana,1846
- Discoharpacticus* Noodt,1954
- Paratigiropus* Itô,1969
- Zausopsis* Lang,1934
- Harpacticellinae Lang,1948
- Harpacticella* G.O. Sars,1908
- Harpacticus* H. Milne-Edwards,1840
- H. chelifera* (O.F. Müller,1776)  
= *Cyclops chelifera* O.F. Müller,1776  
= *Harpacticus helgolandicus* Poppe,1884  
Kieler Bucht-Bohuslän-Helgoland-S & N Norway-Iceland-Greenland-Faeroes-Br. Isles-N France-?Mediterranean
- H. uniremis* Krøyer,1842  
Kieler Bucht-Bohuslän-Helgoland-S & N Norway-Arctic Seas-Iceland-Greenland-Br. Isles-Roscoff
- H. gracilis* Claus,1863  
= *Harpacticus elongatus* Boeck,1864

- \* Not mentioned in Huys & al. 1996  
Kieler Bucht-Bohuslän-German Bight-Trondheim-Faeroes-Br. Isles-N France-Mediterranean-Black Sea
- H. flexus*** (Brady & Robertson, 1873)  
Kieler Bucht-Bohuslän-S North Sea-S & N Norway-Arctic Seas-Br. Isles-N France-Mediterranean-Black Sea
- H. obscurus*** T. Scott, 1895  
= *H. gracilis* : G.O. Sars, 1911, non Claus, 1863  
\* Very difficult to separate from *H. littoralis* and *H. giesbrechti*  
Kieler Bucht-Bohuslän-German Bight-Br. Isles-Roscoff
- H. littoralis*** G.O. Sars, 1910  
\* Very difficult to separate from *H. lgiesbrechti* and *H. obscurus*  
Bohuslän-W Norway-S North Sea-Br. Isles-Mediterranean-Black Sea
- H. tenellus*** G.O. Sars, 1920  
\* Not mentioned in Huys & al. 1996  
Oslofjord-S & W Norway-Ireland-Wales
- H. islandicus*** Klie, 1939  
Iceland
- H. pulvinatus*** Brady, 1910  
Portugal
- H. compsonyx*** Monard, 1926  
Scilly Islands-Roscoff-Mediterranean, world-wide in warmer seas
- H. giesbrechti*** Klie, 1927  
= *Harpacticus chelifer* : Giesbrecht, 1881, non (O.F. Müller, 1776)  
\* Very difficult to separate from *H. littoralis* and *H. obscurus*  
Helgoland
- H. septentrionalis*** Klie, 1939  
Iceland
- Tigriopus*** Norman, 1868
- T. brevicornis*** (O.F. Müller, 1776)  
= *T. fulvus* : Auctt., non Fischer, 1860  
=? *T. liljeborgi* Norman, 1868 *northumbriensis* Mistakidis, 1949  
Öresund-Bohuslän-N Norway-Br. Isles-N & W France- W Spain
- Perissocope*** Brady, 1910
- P. adiastaltus*** Wells, 1968  
Scilly Islands
- P.*** sp. : Huys & al. 1996  
=? *P. xenus* Monard, 1926 (nom. dub.)  
NE England, ?Banyuls
- Zausodiinae** Lang, 1948
- Zaus*** Goodsir, 1845
- Z. spinatus*** Goodsir, 1845 (TM)  
Öresund-Bohuslän-Helgoland-S & N Norway-Arctic Seas-Greenland-Faeroes-Shetlands-Br. Isles-N France
- Z. goodsiri*** Brady, 1880  
\* According to Huys & al. 1996 this probably belongs to a new genus.  
S North Sea-W Norway-Arctic Seas-Greenland-Br. Isles
- Z. abbreviatus*** G.O. Sars, 1904  
Öresund-Bohuslän-W Norway-Barents Sea-Arctic Seas-Iceland-Faeroes-Ireland
- Z. aurelii*** Poppe, 1884  
Arctic Seas, ?Iceland
- Z. caeruleus*** Campbell, 1929  
= *Zaus aurelii* : G.O. Sars, 1909, non Poppe, 1884  
= *Zaus schaeferi* Klie, 1949  
Helgoland, Bergen, Norfolk, Vancouver Island
- Zausodes*** C.B. Wilson, 1932
- Tisboidea** Stebbing, 1910
- Tisbidae** Stebbing, 1910, emend. Lang, 1948
- Tisbinae** Stebbing, 1910
- Tisbe*** Liljeborg, 1853  
= *Idya* Philippi, 1843, non Fréminville, 1809  
= *Idyaea* G.O. Sars, 1909
- T. furcata*** (Baird, 1837)  
Stockholm archipelago-Bohuslän-S North Sea-S & N Norway-Arctic Seas-Iceland-Faeroes-Shetlands-Br. Isles-?Mediterranean
- T. ensifer*** Fischer, 1860  
Bohuslän-Helgoland-N Norway-Faeroes-Br. Isles-Mediterranean & ?Madeira
- T. gracilis*** (T. Scott, 1895)  
Bohuslän-Helgoland-Belgium-N Norway-Arctic Seas-Br. Isles-N France
- T. longicornis*** (T. & A. Scott, 1895)  
Kristiansund-Shetlands-Br. Isles-Roscoff-Mediterranean
- T. minor*** (T. & A. Scott, 1896)  
Kattegatt-Bohuslän-Helgoland.-W Norway-Arctic Seas-Iceland-Br. Isles-Roscoff
- T. elegantula*** (G.O. Sars, 1905)  
Kristiansund-Roscoff-Algeria
- T. angusta*** (G.O. Sars, 1905)  
Norway (Herö)-Ireland-Roscoff-W Mediterran.
- T. tenella*** (G.O. Sars, 1910)  
S Norway-Helgoland-Algeria
- T. graciloides*** (G.O. Sars, 1920)  
Bohuslän-S Norway
- T. compacta*** (G.O. Sars, 1920)  
S Norway-Helgoland
- T. dilatata*** Klie, 1949  
= *Tisbe bulbisetosa* Volkmann-Rocco, 1972  
Helgoland, Belgium
- T. holothuriae*** Humes, 1957
- = *T. helgolandica* Uhlig & Noodt, 1966  
@ freelifving or on *Holothuria (Holothuria) stellati* Delle Chiaie & *Holothuria (Holothuria) tubulosa* Gmelin  
Helgoland-Belgium-Plymouth-Portugal-Italy, E USA
- T. elongata*** (A. Scott, 1896)  
@ Among gills of *Homarus gammarus* - or found free in mud  
Orkneys-Scotland, NW & SW England, Isle of Man
- T. cluthae*** (T. Scott, 1899)  
Shetlands-Scotland
- T. tenera*** (G.O. Sars, 1905)  
Stavanger-N Norway-Iceland-Ireland-Roscoff
- T. finnarhica*** (G.O. Sars, 1905)  
N Norway
- T. battagliai*** Volkmann-Rocco, 1972  
Belgium
- T. cucumariae*** Humes, 1957  
@ freelifving or on *Ocnus planci* (Brandt)  
Arcachon, Portugal, Mediterranean, Bermuda
- T. gurneyi*** (Lang, 1934)  
= *Idyaea gurneyi* Lang, 1934  
Atlantic Europe, Campbell Island
- T. reticulata*** Bocquet, 1951  
?Atlantic Europe
- T. clodiensis*** Battaglia & Fava, 1968  
?Atlantic Europe, Mediterranean
- Scutellidium*** Claus, 1866
- S. longicauda*** (Philippi, 1840)  
= *Scutellidium tishoides* Claus, 1866  
= *Psamate sargassi* G.O. Sars, 1916  
Bohuslän-W Norway-Iceland-Br. Isles-N France-Mediterranean-Black Sea
- S. hippolytes*** (Krøyer, 1863)  
= *Machairopus minutus* G.O. Sars, 1905  
Öresund-Bohuslän-Helgoland-N Norway-Greenland-Iceland-Br. Isles
- S. arthuri*** Poppe, 1884  
Iceland
- S. ligusticum*** (Brian, 1920)  
Roscoff-Mediterranean
- Sacodiscus*** C.B. Wilson, 1924  
= *Aspidiscus* Norman, 1868, non Koenig, 1825
- S. littoralis*** (G.O. Sars, 1904)  
Bohuslän-W Norway-Iceland-Faeroes-Shetlands-Br. Isles-N France
- S. fasciatus*** (Norman, 1868)  
W Norway-Trondheim-Shetlands-Ireland-Roscoff
- S. humesi*** Stock, 1960  
@ *Holothuria (Holothuria) tubulosa* Gmelin  
W France, Banyuls
- Cholidyinae** Boxshall, 1979
- Cholidya*** Farran, 1914  
= *Cholydia* : Bresciani, 1970
- C. polypi*** Farran, 1914 (TM)  
@ between arms of *Benthocopus ergasticus*. Also reported from e.g. Bathypolypus  
\* Not mentioned by Gotto, 1993  
W Ireland
- Cholidyella*** Avdeev, 1982  
\* Originally spelled *Cholydiella* but corrected by Avdeev, 1986
- C. intermedia*** (Bresciani, 1970) Avdeev, 1982  
= *Cholydia intermedia* Bresciani, 1970  
@ Cirrothetidae  
Faeroe-Shetland Channel
- Idyanthinae** Lang, 1948
- Tachidiella*** G.O. Sars, 1909
- T. minuta*** G.O. Sars, 1911  
Bohuslän-German Bight-Norway (Skutesnaes)-Roscoff
- Zosime*** Boeck, 1872
- Z. valida*** G.O. Sars, 1919  
Kieler Bucht-Bohuslän-N Norway-Spitsbergen-Arctic Seas-Faeroes-Br. Isles
- Z. bergensis*** Drzycimski, 1968  
Bergen
- Z. typica*** Boeck, 1872  
Bohuslän-Bergen
- Z. incrassata*** G.O. Sars, 1910  
Bohuslän-Bergen
- Z. major*** G.O. Sars, 1919  
Bohuslän-S Norway
- Z. gisleni*** Lang, 1948  
Bohuslän-Bergen
- Idyella*** G.O. Sars, 1905
- I. pallidula*** G.O. Sars, 1905  
Bohuslän-Bergen, Arctic Seas
- I. exigua*** G.O. Sars, 1905  
Bohusl-Bergen, Bermudas, Algeria
- I. major*** G.O. Sars, 1920  
Bohuslän-S Norway
- Idyellopsis*** Lang, 1948

- I. typica* Lang,1948  
Bohuslän-Bergen
- Idyanthe* G.O. Sars,1909  
= *Idyopsis* G.O. Sars,1905, non Agassiz,1860
- I. pusilla* (G.O. Sars,1905)  
N Norway, W Mediterranean
- I. dilatata* (G.O. Sars,1905)  
Sauesund-Bergen-Spitsbergen
- Tachidiopsis* G.O. Sars,1911
- T. cyclopoidea* G.O. Sars,1911  
S Norway-Bergen
- T. bozici* Bodin,1968  
Biscay in 1200 m
- Idyanthopsis* Bocquet & Bozic,1955  
\* Dubious genus and species. Kunz,1962 considered this species as a synonym of *Diarthroedella orbiculata* Klie,1949. The systematic position of *Diarthroedella* is also dubious. Different workers locate that genus in Paramesochridae and Thalestridae.
- I. psammophila* Bocquet & Bozic,1955  
Roscoff
- Porcellidiidae Brady,1880
- Porcellidium* Claus,1860
- P. fimbriatum* Claus,1863  
= *Porcellidium viride* : Auctt., ?non *Thyone viridis* Philippi,1840  
?Kattegatt-?Bohuslän-?Trondheim-?Shetlands-?Br. Isles-Scilly Islands-N France-Canaries-Mediterranean
- P. sarsi* Bocquet,1848  
= *Porcellidium fimbriatum* : Sars,1904, non Claus,1863  
\* This species, *P. lecanoides* Claus,1989 & *P. fimbriatum* have been so confused, so it is difficult to encounter their respective distribution (see *P. fimbriatum*)
- P. tenuicauda* Claus,1860  
Ireland-Isle of Man-Scilly Islands-Roscoff-Mediterranean
- P. lecanoides* Claus,1889  
\* See note under *P. sarsi*  
Ireland, Devon, Roscoff
- P. lecanoides* var. *roscoffensis* Bocquet,1948  
Roscoff
- P. ovatum* Haller,1879  
SW Ireland, Mediterranean
- Peltidiidae Claus,1860
- Alteutha* Baird,1845
- A. depressa* (Baird,1837)  
Br. Isles-Roscoff-Mediterranean-Black Sea
- A. interrupta* (Goodsir,1845)  
Bohuslän-S North Sea-Trondheim-Shetlands-Br. Isles-N France-Mediterranean
- A. oblonga* (Goodsir,1845)  
Bohuslän-S North Sea-W Norway-Arctic Seas-Shetlands-Br. Isles-N France
- A. sarsi* Monard,1935  
\* Not mentioned in Huys & al. 1996  
S Ireland
- A. roeae* Hicks,1982  
Ireland, South Africa
- Peltidium* Philippi,1839
- P. purpureum* Philippi,1839  
Bohuslän-Trondheim-Br. Isles-Mediterranean
- P. robustum* (Claus,1889)  
S Ireland-Engl. Channel-Mediterranean
- Eupelte* Claus,1860
- E.* sp. : Huys & Al.,1996  
S. England
- E. gracilis* Claus,1860  
?Scilly Islands, Mediterranean
- Alteuthella* A. Scott,1909
- Alteuthoides* Hicks,1986
- Parapeltidium* A. Scott,1909
- Alteuthellopsis* Lang,1944
- Neopeltopsis* Hicks,1976
- Clytemnestridae A. Scott,1909  
= Pseudopeltidiidae Poppe,1891
- Clytemnestra* Dana,1848  
\* Pelagic
- C. scutellata* Dana,1848  
Swedish westcoast, Oslofjord-Br. Isles-Bay of Biscay-Mediterranean
- C. rostrata* (Brady,1883)  
Engl. Channel-Bay of Biscay-Mediterranean
- Tegastidae G.O. Sars,1904
- Tegastes* Norman,1903
- T. grandimanus* G.O. Sars,1904  
Öresund-W Norway (Eggesboenaes)-Br. Isles
- T. longimanus* (Claus,1863)  
SW Norway-Helgoland
- T. falcatus* (Norman,1868)  
Oslofjord-N Norway-Arctic Seas-Shetlands-Orkneys-Br. Isles, Ireland
- T. nanus* G.O. Sars,1904  
Kieler Bucht-W Norway (Christiansund)-Ireland
- T. clausi* G.O. Sars,1904  
W Norway (Eggesboenaes)-German Bight, Britain
- T. calcaratus* G.O. Sars,1910  
W Norway
- T. flavidus* G.O. Sars,1904  
W Norway
- T. satyrus* (Claus,1860)  
= *T. harpacticoides* : G.O. Sars,1910  
SW Norway (Skutesnaes), Mediterranean
- Parategastes* G.O. Sars,1904
- P. sphaericus* (Claus,1863)  
Kieler Bucht-Bohuslän-Helgoland-W Norway-Br. Isles-Roscoff-Mediterranean
- Arawella* Cottarelli & Baldari,1987
- Feregastes* Fiers,1986
- Syngastes* Monard,1924
- PODOGENNONTA Lang,1948
- Thalestroidea G.O. Sars,1905, emend. Lang,1948
- Thalestridae G.O. Sars,1905, emend. Lang,1948
- Thalestrinae G.O. Sars,1905, emend. Lang,1948
- Thalestris* Claus,1863
- T. longimana* Claus,1863  
@ *Ophiolithrix fragilis*, *Ophiolithrix aculeata* & *Buccinum undatum*  
Kieler Bucht-Bohuslän-Helgoland-W Norway-Greenland-Faeroes-Br. Isles-N & W France-Mediterranean
- T. rufoviolascens* Claus,1866  
Kattegatt-Norway (Herøy)-Scotland-French Mediterranean
- T. brunnea* G.O. Sars,1905  
W Norway-Shetlands-Br. Isles
- T. purpurea* G.O. Sars,1905  
Kieler Bucht-Kattegatt-W Norway-Iceland-Greenland
- T. gibba* (Krøyer,1842)  
N Norway-Arctic Seas-Iceland
- T. frigida* T. Scott,1898  
S Norway & Arctic Seas
- T. normani* T. Scott,1903  
N Norway
- T. longipes* Boeck,1865  
\* May belong in *Parathalestris*  
Oslofjord
- T. curticornis* Boeck,1865  
\* May belong in *Dactylopodia*  
Oslofjord
- T. northumbrica* (Norman & Brady,1909, ex Brady MS)  
\* May be an extreme specimen of another species  
Northumberland
- T. denti* Brady,1905  
\* May belong in *Parathalestris*  
England
- T. rhodymeniae* (Brady,1894)  
@ *Palmaria palmata*  
W Scotland
- Amenophia* Boeck,1865
- A. peltata* Boeck,1865  
Kieler Bucht-Bohuslän-Helgoland-Trondheim-Greenland-Br. Isles
- A. pulchella* G.O. Sars,1906  
Bergen-Christiansund
- Parathalestris* Brady & Robertson,1873
- P. harpacticoides* (Claus,1863)  
@ *Crossaster papposus*  
Kieler Bucht-Bohuslän-North Sea-N Norway-Iceland-Faeroes-Shetlands-Br. Isles-Roscoff-Mediterranean
- P. clausi* (Norman,1868)  
Kieler Bucht-Bohuslän-North Sea-N Norway-Shetlands-Br. Isles-N France
- P. hibernica* (Brady & Robertson,1873)  
Stavanger-Christiansund-Br. Isles-Ireland-N France
- P. jacksoni* (T. Scott,1898)  
S-N Norway-Arctic Seas-Greenland-Iceland
- P. intermedia* Gurney,1930  
= *Parathalestris plumiseta* C.G. Moore,1976  
Schleswig-Holstein North Sea coast-Norfolk-Ireland-Isle of Man-Roscoff
- P. cronii* (Krøyer,1842)  
= *Halithalestris cronii* : Auctt.  
W-N Norway-Spitsbergen-Arctic Seas-Iceland-Greenland-Br. Isles-Engl. Channel; -- Pelagic - often in company with floating seaweed -
- P. irelandica* Roe,1958  
S Ireland
- P. cambriensis* Wells,1964  
Wales
- P. dovi* Marcus,1966  
W France-Black Sea
- Phyllothalestris* G.O. Sars,1905
- P. mysis* (Claus,1863)

Bohuslän-Trondheim-Faeroes-Br. Isles-N France-Mediterranean-Black Sea

Rhynchothalestrinae Lang,1948

**Rhynchothalestris** G.O. Sars,1905

**R. helgolandica** Claus,1863, emend. Huys,1990

Kieler Bucht-Bohuslän-North Sea-N Norway-Arctic Seas-Greenland-Iceland-Br. Isles-Ireland

Dactylopusiinae

= Dactylopodinae Lang,1936

**Diarthodes** G.M. Thomson,1882

**D. nobilis** (Baird,1845)

Kieler Bucht-Bohuslän-Helgoland-Lofoten-Arctic Seas-Shetlands-Br. Isles-N France-W Mediterranean-Madeira

**D. major** (T. & A. Scott,1895)

Haugesund-Vadsø-Helgoland-Br. Isles & Ireland

**D. pygmaeus** (T. & A. Scott,1895)

Kieler Bucht-Bohuslän-North Sea-W Norway-Br. Isles-Roscoff-Mediterranean

**D. minutus** (Claus,1863)

S Norway-Trondheim-North Sea-Ireland-Roscoff-Mediterranean-Black Sea

**D. assimilis** (G.O. Sars,1909)

Norway

**D. ponticus** (Kricagin,1873)

Ireland-Roscoff-Mediterranean-Black Sea

**D. andrewi** (T. Scott,1894)

Br. Isles

**D. roscoffensis** (Monard,1935)

Roscoff

**D. feldmanni** Bocquet,1953

@ Rhodophyceae

Roscoff

**D. intermedius** (T. Scott,1912)

= *Pseudothalestris intermedia* T. Scott,1912

Atlantic Europe, South Orkneys

**D. aegideus** (Brian,1927)

= *Westwoodia pygmaea* var. *aegidea* Brian,1927

?European Atlantic, Mediterranean

**Dactylopusia** Norman,1903

= *Dactylopus* Claus,1862, non Gill,1959

= *Dactylopodia* Lang,1948

**D. vulgaris** G.O. Sars,1905

Kieler Bucht-Bohuslän-Helgoland-N Norway-Arctic Seas-Iceland-Br. Isles-Roscoff-Mediterranean

**D. vulgaris** var. *dissimilis* Brian,1921

Atlantic Europe, Mediterranean

**D. euryhalina** Monard,1935

= *Dactylopodia vulgaris holsatica* Noodt,1953

Kiel Channel

**D. neglecta** (G.O. Sars,1905)

Trondheim-Spitsbergen-Br. Isles-Ireland-Roscoff

**D. tisboides** (Claus,1863)

Kieler Bucht-Bohuslän-N Norway-Arctic Seas-Shetlands-Br. Isles-N France-Madeira-Mediterranean-Black Sea

**D. micronyx** G.O. Sars,1905

Kieler Bucht-Kattegat-Trondheim-Ireland-Devon-Roscoff

**D. longyerbyensis** (Mielke,1974)

= *Dactylopodia longyerbyensis* Mielke,1974

Spitsbergen

**Dactylopusioides** Brian,1828

**D. macrolabris** (Claus,1866)

= *Dactylopus macrolabris* Claus,1863

= *Dactylopusioides stampaliae* Brian,1928 (TM)

?Atlantic Europe, Mediterranean

**Paradactylopodia** Lang,1948

= *Dactylopusia* Norman,1903 (p.p.)

**P. latipes** (Boeck,1865)

Kieler Bucht-Bohuslän-Helgoland-W Norway-Ireland

**P. brevicornis** (Claus,1866)

@ *Limnoria lignorum*

Kieler Bucht-Bohuslän-Helgoland-S & N Norway-Br. Isles-Engl. Channel-Mediterranean-Black Sea

**Dactylopodopsis** G.O. Sars,1911

**D. dilatata** G.O. Sars,1911

Bohuslän-Lofoten-Iceland

Pseudotachidiinae Lang,1936

**Pseudotachidius** T. Scott,1897

**P. coronatus** T. Scott,1897

Bohuslän-W Norway-Scotland

**P. vikingus** Drzycimski,1968

Bergen

**P. similis** T. Scott,1902

Faeroe Channel

**Idomene** Philippi,1843

**I. forficata** Philippi,1843

Öresund-Kattegat-Øslofjord-Trondheim-Br. Isles-Ireland-Italy

**I. coronata** (T. Scott,1894)

S Norway-Arctic Seas-Scotland

**I. pectinata** (T. & A. Scott,1898)

Scotland

**I. purpurocineta** (Norman & T. Scott,1905)

England (Devon)

**I. borealis** G.O. Sars,1911

Lofoten

**I. aberrans** Por,1964

= *Idomene peresi* Bodin,1964

\* Not belonging in this genus or even family?

?Atlantic Europe, Mediterranean

**Dactylopedella** G.O. Sars,1905

**D. flava** (Claus,1866)

= *Dactylopedella ornata* (Norman & T. Scott,1905)

Bohuslän-Helgoland-W Norway-Arctic Seas-Br. Isles-Mediterranean

**D. clypeata** G.O. Sars,1911

S Norway-Bergen

**D. rostrata** (T. Scott,1893)

Scotland

Donsiellinae Lang,1948

**Donsiella** Stephensen,1936

**D. limnoriae** Stephensen,1936

@ Usually on ventral surface, especially around the bases of the legs on *Limnoria lignorum* (Rathke), *L. tripunctata*, *L. quadripunctata* & *L. japonica*

Trondheim-Southampton, SE Ireland (British records may perhaps have been *D. anglica*), E Canada

**D. anglica** Hicks,1988

Britain

Ambunguipedidae Huys,1990

**Ambungipes** Huys,1990

**A. rufocincta** (Brady,1880, ex Norman MS)

= *Thalestris rufocincta* Brady,1880, ex Norman MS (TM)

= *Rhynchothalestris rufocincta* (Brady,1880, ex Norman MS)

Bohuslän-Trondheim-Br. Isles-N France-Madeira-Mediterranean

**Lucayostratiotes** Huys,1990 (TM *Rhynchothalestris cornuta* Geddes,1969)

Balaenophilidae G.O. Sars,1910

**Balaenophilus** Christian Aurivillius,1879

**B. unisetosus** Christian Aurivillius,1879

@ On baleen plates of the *Balaenoptera* spp.

N Norway, Isle of Man, W Ireland

Parastenheliidae Lang,1936

**Parastenhelia** Thompson & Scott,1903

**P. spinosa** (Fischer,1860)

= *Thalestris forficula* Claus,1863

= *Microthalestris forficula* : G.O. Sars,1911

Kieler Bucht-Bohuslän-W & N Norway-Arctic Seas-Faeroes-Br. Isles-Roscoff-Mediterranean

**P. spinosa bulbosa** (Bozic,1954)

S England

**P. littoralis** (G.O. Sars,1911)

= *Microthalestris littoralis* G.O. Sars,1911

\* Distribution not ascertained, because of presumed identity of this species with *P. spinosa* (Fischer)

Bohuslän-Bergen

**P. anglica** Norman & T. Scott,1905

Cornwall-Scilly Isles

**P. ornatisima** (Monard,1937)

Roscoff

Diosaccidae G.O. Sars,1906

**Psammotopa** Pennak,1942

= *Protoleptastacus* Noodt,1952

**P. phyllosetosa** (Noodt,1952)

= *Protoleptastacus phyllosetosus* Noodt,1952 (TM of *Protoleptastacus* Noodt,1952 - Syla)

Nordfriesland-Belgium-W Br. Isles-France-Portugal

**P. polyphylla** Noodt,1955

Arcachon-Portugal

**P. vulgaris** Pennak,1942

SW Br. Isles-Arcachon-E Mediterranean

**P. chappuisi** Noodt,1955

= *P. vulgaris* : Chappuis,1954, non Pennak,1942

W France

**Protopsammotopa** Geddes,1968

**P. norvegica** Geddes,1968

Bergen

**Robertsonia** Brady,1880

**R. tenuis** (Brady & Robertson, in Brady,1880)

= *Ectinosoma tenuis* Brady & Robertson,1875 (nomen nudum)

= *R. tenuis kielensis* Becker,1970

Belt Sea-Öresund-Bohuslän-SW Norway-Arctic Seas-Br. Isles

**R. diademata** Monard,1926

Roscoff

**R. celtica** (Monard,1935)

= *Teissierella celtica* Monard,1935

Roscoff-S Ireland

**R. propinqua** (T. Scott,1893)

= *Dactylopus propinquus* T. Scott,1893

= *Amphiascus scotti* Sewell,1924

European Atlantic, Bermudas, India

**R. monardi** (Klie,1937)

= *Varnaia monardi* Klie,1937

?European Atlantic, Bulgaria, Romania

**Schizopera** G.O. Sars,1905

**S. (Schizopera) clandestina** (Klie,1924)

Finland-Elbe estuary-Aberdeen-Black Sea

- S. (Schizopera) compacta* de Lint, 1922  
Kiel Channel-Elbe mouth-Zuidersee
- S. (Neoschizopera) Apostolov, 1982) pratensis* Noodt, 1958  
Belt Sea-German North Sea coast
- S. (Neoschizopera) Apostolov, 1982) meridionalis* Petkovski, 1954  
= *S. inornata* Noodt, 1955  
Swedish Baltic
- S. (Neoschizopera) Apostolov, 1982) meridionalis listensis* Mielke, 1975  
Sylt
- S. baltica* Lang, 1965  
Askö area (Baltic) & S Norway
- S. ornata* Noodt & Purasjoki, 1953  
Finnish & German Baltic, S Norway & Spitsbergen
- S. minuta* Noodt, 1955  
W France
- S. nana* Noodt, 1955  
W France
- Eoschizopera* Wells & Rao, 1976
- E. (Eoschizopera) syltensis* (Mielke, 1973)  
= *Schizopera syltensis* Mielke, 1973  
Sylt
- Pseudodiosaccus* T. Scott, 1906
- P. propinquus* (T. & A. Scott, 1893)  
Scotland
- Pseudomesochra* T. Scott, 1902  
= *Stenheliospis* G.O. Sars, 1906  
= *Jonestella* Monard, 1927
- P. longifurcata* T. Scott, 1902  
Bohuslän-S Norway-Scotland
- P. divaricata* (G.O. Sars, 1906)  
Bohuslän-W Norway
- P. similis* Lang, 1935  
Bohuslän
- P. latifurca* (G.O. Sars, 1911)  
Norway (Lyngfjord)
- P. media* (G.O. Sars, 1911)  
Bohuslän-SW Norway
- P. tatananae* Drzycimski, 1968  
Bergen
- P. scheibeli* Schriever, 1982  
Iceland-Faeroer-Ridge
- Stenheliospis* Boeck, 1865
- S. (Stenheliospis) gibba* Boeck, 1865  
@ *Crossaster papposus* & *Musculus tumidus*  
Kieler Bucht-Bohuslän-Helgoland-N Norway-Arctic Seas-Br. Isles
- S. (Stenheliospis) aemula* (T. Scott, 1893)  
Bohuslän-SW Norway-Br. Isles-Roscoff
- S. (Stenheliospis) proxima* G.O. Sars, 1909  
Kieler Bucht-Bohuslän-SW Norway-Aberdeen
- S. (Delavalia) Brady, 1868) palustris* (Brady, 1868)  
\* Tube building & home finding species.  
Finland-Bohuslän-Oslofjord-S Norway-German Bight-Br. Isles-Zuidersee-Roscoff
- S. (Delavalia) palustris bispinosa* Bodin, 1970  
W France
- S. (Delavalia) longicaudata* Boeck, 1872  
= *Delavalia robusta* Brady & Robertson, 1875  
Kieler Bucht-Bohuslän-N Norway-Arctic Seas-Br. Isles
- S. (Delavalia) longicaudata finnarchica* (T. Scott, 1903)  
= *Delavalia robusta* var. *finnarchica* T. Scott, 1903  
N Norway
- S. (Delavalia) reflexa* Brady & Robertson, 1880  
Belt Sea-Bohuslän-SW Norway-Scotland-Br. Isles
- S. (Delavalia) giesbrechti* T. & A. Scott, 1896  
SW Norway-Br. Isles-Roscoff
- S. (Delavalia) normani* (T. Scott, 1905)  
= *Delavalia Normani* T. Scott, 1905  
S Norway-Br. Isles-Mediterranean
- S. (Delavalia) hanstroemi* Lang, 1948  
Bohuslän-Bergen
- S. (Delavalia) confluens* Lang, 1948  
Bohuslän-Bergen
- S. (Delavalia) pygmaea* (Brady, 1905)  
= *Delavalia pygmaea* Brady, 1905  
Northumberland
- S. (Delavalia) mastigochaeta* Wells, 1965  
Scotland
- S. (Delavalia) noodti* Schriever, 1982  
Iceland-Faeroer-Ridge
- S. (Delavalia) islandica* Schriever, 1982  
Iceland-Faeroer-Ridge
- S. (Delavalia) arctica* (T. Scott, 1898)  
= *Delavalia arctica* T. Scott, 1898  
off N Koster in 110m, Franz Joseph Land, King Karls Land
- S. (Delavalia) elisabethae* Por, 1960  
?Britain, Black Sea
- Diosaccus* Boeck, 1872  
= *Paradiosaccus* Lang, 1948
- D. tenuicornis* (Claus, 1863)  
Belt Sea-Bohuslän-North Sea-S & N Norway-Br. Isles-Engl. Channel-Canaries-Mediterranean
- D. varicolor* (Farran, 1913)  
Ireland-Roscoff-Mediterranean
- Robertgurneya* Lang, 1948
- R. spinulosa* (G.O. Sars, 1911)  
= *Amphiascus spinulosus* G.O. Sars, 1911  
Bohuslän-S & W Norway
- R. similis* (A. Scott, 1896)  
= *Amphiascus scotti* Williamson, 1958  
off Gullmar (Bonden & Mittskären)-S-W Norway-Ireland-England-Roscoff-Bermudas-Mediterranean
- R. ilievecensis* (Monard, 1935)  
= *Amphiascus ilievecensis* Monard, 1935  
N Ireland-Roscoff
- R. remanei* Klie, 1950  
Helgoland
- R. intermedia* Bozic, 1954  
\* Dubious species  
Roscoff
- R. dictydiophora* (Monard, 1924)  
= *Amphiascus dictydiophorus* Monard, 1924  
?European Atlantic, Mediterranean
- R. erythraeus* (A. Scott, 1902)  
= *Stenheliospis erythraea* A. Scott, 1902  
?European Atlantic, Massachusetts, Red Sea
- R. oligochaeta* Noodt, 1955  
Isle of Man, Marmara Sea
- R. rostrata* (Gurney, 1927)  
= *Amphiascus rostratus* Gurney, 1927  
Celtic Sea, Bermudas, Mediterranean
- R. simulans* (Norman & T. Scott, 1905)  
= *Stenhelios simulans* Norman & T. Scott, 1905  
SW Britain
- Haloschizopera* Lang, 1948
- H. exigua* (G.O. Sars, 1906)  
Bohuslän-SW Norway
- H. bulbifer* (G.O. Sars, 1911)  
Kosterfjorden in 72-110m-S Norway
- H. conspicua* Por, 1964  
off Gullmar (Mittskären in 27-30m)
- H. minima* Por, 1964  
off Gullmar (Mittskären in 27-30m)
- H. pygmaea* (Norman & T. Scott, 1905)  
= *Haloschizopera junodi* (Monard, 1935)  
= *Haloschizopera marmarae* Noodt, 1955  
Bohuslän-Helgoland-Bergen-Roscoff-Br. Isles
- H. pauciseta* Por, 1959  
?European Atlantic, Mediterranean
- H. ruthorum* Por, 1959  
?Isle of Man, Mediterranean
- Paramphiascella* Lang, 1948
- P. intermedia* (T. Scott, 1896)  
Kieler Bucht-Bohuslän-Trondheim-Arctic Seas-Faeroe Channel-Br. Isles
- P. vararensis* (T. Scott, 1903)  
= *Amphiascus affinis* G.O. Sars, 1906  
Kieler Bucht-Bohuslän-Helgoland-W Norway-Arctic Seas-Scotland-S Ireland
- P. hispida* (Brady, 1880)  
Bohuslän-N Norway-Arctic Seas-Iceland-Br. Isles-Roscoff
- P. fulvofasciata* Rosenfield & Coull, 1974  
@ ? *Limnoria lignorum*  
Helgoland-Norfolk
- P. hyperborea* (T. Scott, 1903)  
N Norway-Greenland-Arctic Seas
- P. roberti* (Monard, 1935)  
\* Dubious species  
Roscoff
- P. bodini* Marcotte, 1974  
= *P. mediterranea* : Bodin, 1964, non Lang, 1948  
Celtic Sea, Mediterranean
- Amphiascoides* Nicholls, 1941  
= *Amphiascella* Lang, 1948
- A. debilis* (Giesbrecht, 1881)  
=? *A. limicola* (Brady, 1900)  
Finland-Bohuslän-S North Sea-S Norway-Trondheim-Br. Isles-Ireland-Roscoff
- A. dispar* (T. & A. Scott, 1894)  
Belt Sea-Bohuslän-Scotland
- A. neglecta* (Norman & T. Scott, 1905)  
= *Amphiascella neglecta* : Lang, 1948  
SW Norway-England-Scilly Isles-Roscoff-Mediterranean
- A. nanus* (G.O. Sars, 1906)  
= *Amphiascella nana* : Lang, 1948  
Hällö in 6m-S Norway-Roscoff-Mediterranean
- A. littoralis* (T. Scott, 1903)  
Scotland
- A. nanoides* (G.O. Sars, 1911)  
S Norway
- A. brevifurca* (Czerniavski, 1868)  
S Ireland-Roscoff-Mediterranean-Black Sea

- A. subdehilis* (Willey, 1835)  
Kosterfjorden in 72-240m-Skagerrak-Bergen-S Ireland-Roscoff-Mediterranean-Bermudas
- Metamphiascopsis** Lang, 1948
- M. hirsutus* (I.C. Thompson & A. Scott, 1903)  
= *Dactylophus hirsutus* I.C. Thompson & A. Scott, 1903  
?European Atlantic, Mediterranean
- Typhlamphiascus** Lang, 1948
- T. blanchardi* (T. & A. Scott, 1895)  
Bohuslän-SW Norway-Scotland-Scilly Isles
- T. gracilis* Por, 1963  
off N Koster-Skagerrak
- T. confusus confusus* (T. Scott, 1902)  
Bohuslän-Helgoland-Bergen-Br. Isles-Roscoff-Mediterranean
- T. confusus gullmaricus* Por, 1964  
Bohuslän-Bergen-Mediterranean
- T. typhloides* (G.O. Sars, 1911)  
S-SW Norway
- T. typhlops* (G.O. Sars, 1906)  
Kieler Bucht-Bohuslän-SW Norway-Arctic Seas
- T. lamellifer* (G.O. Sars, 1911)  
Kieler Bucht-Bohuslän-S Norway-Barents Sea-Iceland
- T. hirsutus* (Thompson, 1893)  
= *Sienhelia hirsuta* Thompson, 1893  
\* Problematic species, which may be identical with e.g. *T. confusus*  
N Norway-Br. Isles
- Bulbamphiascus** Lang, 1948
- B. imus* (Brady, 1872)  
= *Canthocampus imus* Brady, 1872  
= *Sienhelia reflexa* T. Scott, 1895, non Brady & Robertson, 1880  
= *Amphiascus Normani* G.O. Sars, 1911  
Bohuslän-W & N? Norway-Arctic Seas-Br. Isles-N France-Mediterranean
- B. denticulatus* (Thompson, 1893)  
off Gullmarn (Bonden)-S Norway-Helgoland-Br. Isles
- B. angustifolius* Klie, 1950  
Helgoland
- Rhyncholagena** Lang, 1948
- R. lagenirostris* (G.O. Sars, 1911)  
S Norway
- R. spinifer* (Farran, 1913)  
Ireland
- R. pestai* (Monard, 1935)  
Roscoff-Algeria
- Pararobertsonia** Lang, 1948
- P. abyssi* (Boeck, 1872)  
Kosterfjorden in 92-150m-Oslofjord-W Norway
- Dactylopodamphiascopsis** Lang, 1948
- D. latifolius* (G.O. Sars, 1909)  
Christiansund-Lofoten-Iceland
- Amphiascus** G.O. Sars, 1905, emend. Lang, 1948
- A. minutus* (Claus, 1863)  
\* A species complex  
Kieler Bucht-Bohuslän-North Sea-N Norway-Arctic Seas-Iceland-Faeroes-Br. Isles-Mediterranean
- A. tenuiremis* (Brady & Robertson, in Brady, 1880)  
=? *A. minutus* sp. 1, Lang, 1965 (according to Mielke, 1974, though Scheibel, 1973 was of the opinion that this form is a synonym of *A. graciloides trisetosus* Klie)  
Kieler Bucht-Bohuslän-German Bight-N Norway-Arctic Seas-Br. Isles-Roscoff
- A. varians* (Norman & T. Scott, 1905)  
Kosterfjorden in 240m-S-W Norway-Helgoland-England-Roscoff
- A. propinquus* G.O. Sars, 1906  
= *A. angustipes* Gurney, 1927  
Kieler Bucht-Bohuslän-W Norway-?Scilly Islands
- A. tenellus* G.O. Sars, 1906  
Bohuslän-W Norway-Roscoff-Algeria
- A. parvus* G.O. Sars, 1906  
= *Amphiascus humphriesi* Roe, 1960  
S-W Norway-Helgoland-Iceland-S Ireland-Roscoff
- A. sinuatus* G.O. Sars, 1906  
Bohuslän-W Norway-Scotland
- A. amblyops* G.O. Sars, 1911  
SW Norway
- A. polaris* G.O. Sars, 1909  
Bergen-Arctic Seas
- A. graciloides* Klie, 1950  
Belt Sea-Helgoland
- A. graciloides* Klie, 1950 *trisetosus* Klie, 1950  
?Helgoland
- A. longarticulatus* A. Marcus, 1974  
Galway Bay-Scilly Isles-Roscoff-Black Sea
- A. hirtus* Gurney, 1927  
?Atlantic Europe, Mediterranean, Campbell Island
- Pseudamphiascopsis** Lang, 1948
- P. attenuatus* (G.O. Sars, 1906)  
= *Pseudamphiascopsis ismaelensis* (Monard, 1936)  
off Gullmarn (Bonden & Mittskären)-SW Norway-Ireland-Scilly Islands
- P. herdmani* (A. Scott, 1896)  
Northumberland-Isle of Man
- Paramphiascopsis** Lang, 1948
- P. longirostris* (Claus, 1863)  
= *Amphiascus paracaudaepinosus* Roe, 1958  
Kieler Bucht-Bohuslän-Sylt-Helgoland-N Norway-Iceland-Faeroes-Br. Isles-S. Ireland-Roscoff-Mediterranean
- P. giesbrechti* (G.O. Sars, 1906)  
Kieler Bucht-Bohuslän-Helgoland-W Norway-Greenland
- P. pallidus* (G.O. Sars, 1906)  
Christiansund
- P. triarticulatus* Moore, 1976  
Isle of Man
- P. paramolae* Soyer, 1973  
@ gills of *Paramola cuvieri* (Risso)  
?Mediterranean
- Amphiascopsis** Gurney, 1927, emend. Lang, 1948
- A. cinctus* (Claus, 1866)  
= *Amphiascopsis obscurus* : Lang, 1948, non (G.O. Sars, 1906)  
Bohuslän-Helgoland-W Norway-Ireland-Roscoff-Bermudas-Mediterranean
- A. obscurus* (G.O. Sars, 1906)  
S Norway
- A. thalestroides* (G.O. Sars, 1911)  
SW Norway-Helgoland-Shetlands-Br. Isles-N France-Mediterranean
- Amonardia** Lang, 1948
- A. normani* (Brady, 1872)  
= *Amphiascus similis* : Auct., non (Claus, 1866)  
Bohuslän-Helgoland-S- Norway-Trondheim-Faeroes-Br. Isles-Ireland
- A. arctica* (T. Scott, 1898)  
Kieler Bucht-Bohuslän-N Norway-Arctic Seas-Iceland
- A. phyllopus* (G.O. Sars, 1906)  
S Norway-Ireland-Roscoff-Mediterranean
- A. similis* (Claus, 1866)  
S Ireland-Roscoff-Canaries-Mediterranean
- Pholenota** Vervoort, 1964 (<sup>TM</sup> *P. spatulifera* Vervoort, 1964)
- Cladorostrata* Shen & Tai, 1963 (<sup>TM</sup> *C. brevipoda* Shen & Tai, 1963 or? *C. longipoda* Shen & Tai, 1963 - fresh water (Pearl River), Cina)
- Balucopsylla** Chandrasekhara Rao, 1972 (<sup>TM</sup> *B. similis* Chandrasekhara Rao, 1972 - India)
- Miraciidae** Dana, 1846  
= Macrosetellidae A. Scott, 1909  
\* Family holoplanktic, with species associated to pelagic colonies of in tropical and subtropical seas distributed species of *Trichodesmium* (formerly marine *Oscillatoria*).
- Distiocolus** Huys & Böttger-Schnack, 1994 (<sup>TM</sup> *Miracia minor* T. Scott, 1894 - Gulf of Guinea)
- Macrosetella** A. Scott, 1909  
= *Setella* Dana, 1846, non Schrank, 1802 (Lepidoptera)  
= *Dwightia* C.B. Wilson, 1924
- M. gracilis** (Dana, 1847)  
= *Setella gracilis* Dana, 1847 (<sup>TM</sup> - Pacific)  
= *Setella longicauda* Dana, 1847  
= *Setella tenuicornis* Dana, 1847  
= *Setella crassicornis* Dana, 1847  
= *Setella aciculus* Dana, 1847  
= *Setella tenuis* Lubbock, 1860  
= *Setella mexsinensis* Claus, 1863  
"Cosmopolitan, planktonic", off W Ireland-southwards
- Miracia** Dana, 1846 (<sup>TM</sup> *Miracia efferata* Dana, 1849 - Equatorial E Atlantic)
- Oculosetella** Dahl, 1895 (<sup>TM</sup> *Miracia gracilis* Dana, 1849 - S Pacific)
- Metoidea** Boeck, 1872
- Metidae** Boeck, 1872
- Metis** Philippi, 1843
- M. ignea** Philippi, 1843  
= *Ilyosyllus coriaceus* Brady & Robertson, 1873  
Bohuslän-W Norway-German Bight-Br. Isles-Roscoff-Mediterranean-Black Sea
- M. holothuriae** (Edwards, 1891)  
N France-Mediterranean, Bahamas, New England
- Ameiroidea** Monard, 1927
- Ameiridae** Monard, 1927, emend. Lang, 1963
- Ameirinae** Monard, 1927
- Ameira** Boeck, 1865
- A. scotti* G.O. Sars, 1911  
Kieler Bucht-Bohuslän-Helgoland-W Norway-Roscoff-Mediterranean
- A. longipes* Boeck, 1865  
= *A. curviseta* Willey, 1929  
Kieler Bucht-Bohuslän-Helgoland-N Norway-Arctic Seas-Br. Isles-Roscoff-Mediterranean
- A. minuta* Boeck, 1865  
Øresund-Oslofjord-Telemark-Bergen-Helgoland-Scotland-Ireland
- A. parvula* (Claus, 1866)  
= *Canthocampus parvulus* Claus, 1866  
= *Nitocra tau* Giesbrecht, 1881  
= *Ameira parvuloides* Lang, 1965  
Kieler Bucht-Bohuslän-S North Sea-Telemark-Trondheim-Arctic Seas-Br. Isles-Roscoff-Mediterranean-Black Sea
- A. tenuicornis* T. Scott, 1902  
Bohuslän-S & W Norway-Scotland-Scilly Isles
- A. tenella* G.O. Sars, 1907  
Helgoland-S & SW Norway-Ireland-Scilly Isles
- A. usitata* Klie, 1950  
Helgoland

- A. brevipes* Kunz, 1954  
Helgoland
- A. brevipes pestae* Petkovski, 1955  
= *Ameira pestae* Petkovski, 1955  
?European Atlantic, NE Mediterranean
- A. speciosa* Monard, 1935  
Roscoff
- A. attenuata* I.C. Thompson, 1893  
\* May be identical with *A. speciosa* Monard  
Isle of Man
- A. exigua* T. Scott, 1894  
\* Belongs probably in *Proameira*  
Br. Isles-Arctic Seas
- A. pusilla* T. Scott, 1903  
Scotland-Scilly Isles-?Roscoff-W France
- A. listensis* Mielke, 1975  
Sylt
- A. divagans* Nicholls, 1940  
Kieler Bucht, N Bretagne
- A. hyalina* (Noodt, 1952)  
= *Psammameira hyalina* Noodt, 1952 (TM of *Psammameira* Noodt, 1952 - Sylt)  
German North Sea coast
- A. lusitanica* Galhano, 1970  
Portugal
- A. intermedia* Galhano, 1970  
Portugal
- A. faeroensis* Schriever, 1982  
Iceland-Faeroer-Ridge
- Proameira* Lang, 1948
- P. arenicola* (Lang, 1935)  
Öresund-Helgoland
- P. hiddenseensis* (Schäfer, 1936)  
Finland-Helgoland
- P. simplex* (Norman & T. Scott, 1905)  
Kieler Bucht-Bohuslän-S Norway-Bergen-Barents Sea-England
- P. dubia* (G.O. Sars, 1920)  
Bohuslän-S Norway
- P. signata* Por, 1964  
off Gullmarn (Mittskären in 27-30m)-Kosterfjorden-Skagerrak
- P. phaedra* (Monard, 1935)  
Roscoff
- P. psammophila* Wells, 1963  
W Br. Isles & Ireland
- P. echinipes* Soyer, 1975  
?European Atlantic, Mediterranean
- Nitokra* Boeck, 1865  
= *Nitocra* Giesbrecht, 1882
- N. typica* Boeck, 1865  
= *Nitocra oligochaeta* Giesbrecht, 1881  
= *Ameira amphibia* Brady, 1902  
= *Ameira littoralis* Nicholls, 1939  
Finland-Kieler Bucht-Bohuslän-S North Sea-S Norway-Trondheim-Br. Isles-N France-Mediterranean-Black Sea
- N. spinipes* Boeck, 1865  
= *N. parafragilis* (Roe, 1958)  
Finland-Öresund-Oslofjord-Karmøy-Telemark-S North Sea-Shetlands-Br. Isles-S Ireland-Roscoff-Mediterranean-Black Sea
- N. lacustris* (Schmankewitsch, 1875)  
Freshwater species, entering weak brackish water: e.g. S Finland, Br. Isles
- N. hibernica* (Brady, 1880)  
Freshwater species, entering weak brackish water: Baltic
- N. pusilla* (G.O. Sars, 1911)  
= *Nitocra pusilla* G.O. Sars, 1911  
SW Norway-S Ireland
- N. elegans* (T. Scott, 1904)  
= *Ameira elegans* T. Scott, 1904  
Scotland-Roscoff
- N. fallaciosa* Klie, 1937  
= *N. psammophila* Noodt, 1952  
German ?North Sea coast
- N. fallaciosa* (Klie, 1937) *baltica* Lang, 1965  
Finland-S Baltic estuarine sand interstitium species
- N. reducta* (Schäfer, 1936)  
Finland-S Baltic brackish water & Elbe estuary
- N. reducta* (Schäfer, 1936) *fluviatilis* Galhano, 1968  
Portugal
- N. affinis* Gurney, 1927  
S Norway-S Ireland-Bermudas-Mediterranean
- N. hamata* Bodin, 1970  
W France
- N. sewelli* Gurney, 1927 *husmanni* Kunz, 1976  
N Germany
- Psyllocamptus* T. Scott, 1899
- P. (Psyllocamptus) minutus* G.O. Sars, 1911  
Bohuslän-S Norway-Scilly Islands
- P. (Psyllocamptus) minutus* G.O. Sars, 1911 *gelatinosus* (Kunz, 1951)  
Sylt
- P. (Psyllocamptus) propinquus* (T. Scott, 1895)  
Helgoland-Scotland
- Ameiropsis* G.O. Sars, 1907
- A. mixta* G.O. Sars, 1907  
Bohuslän-W Norway-Ireland-Roscoff
- A. longicornis* G.O. Sars, 1907  
Skagerrak-S & SW Norway-Helgoland-Ireland
- A. brevicornis* G.O. Sars, 1907  
S & W Norway-Helgoland-Roscoff-Br. Isles
- A. nobilis* G.O. Sars, 1911  
S & W Norway-Ireland-Mediterranean
- A. angulifera* G.O. Sars, 1911  
S Norway
- A. abbreviata* G.O. Sars, 1911  
Skagerrak-S & W Norway
- A. minor* (G.O. Sars, 1920)  
off N Koster in 150m-S & W Norway
- Pseudameira* G.O. Sars, 1911
- P. crassicornis* G.O. Sars, 1911  
Bohuslän-S & W Norway-Scotland-W France
- P. furcata* G.O. Sars, 1911  
Belt Sea-Bohuslän-S Norway
- P. gracilis* G.O. Sars, 1920  
Bohuslän-S Norway
- P. mixta* G.O. Sars, 1920  
Bohuslän-S Norway
- P. reflexa* (T. Scott, 1894)  
Br. Isles-Arctic Seas
- P. reducta* Klie, 1950  
Helgoland-N Ireland
- P. breviseta* Klie, 1950  
Helgoland
- P. perplexa* Soyer, 1975  
?European Atlantic, Mediterranean
- Leptomesochra* G.O. Sars, 1911
- L. confluens* G.O. Sars, 1911  
Bohuslän-S Norway-Helgoland
- L. macintoshi* (T. & A. Scott, 1895)  
Kieler Bucht-German Bight-Br. Isles
- L. nasuta* Sewell, 1940  
W France
- L. attenuata* (Nicholls, 1940)  
= *L. nicholli* Noodt, 1955  
non = *L. attenuata* (A. Scott, 1896) (= *Interleptomesochra attenuata* (A. Scott))  
W France
- Sicameira* Klie, 1950
- S. leptoderma* Klie, 1950  
Helgoland
- S. gracilis* (A. Scott, 1896)  
= *Ameira gracilis* A. Scott, 1896  
off Gullmarn (Bonden)-Cornwall-Isle of Man
- Pseudoleptomesochrella* Lang, 1965
- P. halophila* (Noodt, 1952)  
= *Nitocrella halophila* Noodt, 1952  
= *Nitocrella halophila brevifurca* Wells, 1961  
Sylt-SW Br. Isles-Bulgaria
- Interleptomesochra* Lang, 1965
- I. attenuata* (A. Scott, 1896)  
off Gullmarn (Bonden)-S Norway-W Br. Isles-Brittany
- I. elongata* (Bozic, 1954)  
Brittany
- I. eulitoralis* (Noodt, 1952)  
= *Leptomesochra eulitoralis* Noodt, 1952  
German Bight-Belgium-W Br. Isles-Portugal-Israel
- I. noodti* Galhano, 1968  
NW Portugal
- I. tenuicornis* (G.O. Sars, 1911)  
Korshavn (S Norway)-Engl. Channel
- Sarsameira* Wilson, 1924  
= *Parameira* G.O. Sars, 1907, non Seidlitz, 1868  
= *Pseudosarsameira* Lang, 1948
- S. parva* (Boeck, 1872)  
= *Ameira longiremis* var. *intermedia* T. Scott, 1894  
Bohuslän-Trondheim-W Br. Isles-Bay of Biscay
- S. longiremis* (T. Scott, 1894)  
Bohuslän-S & W Norway-Arctic Seas-Scotland
- S. propinqua* (T. Scott, 1902)  
off Gullmarn (Bonden)-S & W Norway-Scotland
- S. sarsi* Lang, 1948  
= *Parameira intermedia* : G.O. Sars, 1920, non (T. Scott, 1894)  
Bohuslän-S Norway
- S. major* (G.O. Sars, 1907)  
SW Norway-Arctic Seas, Celtic Sea
- S. giraulti* Monard, 1935  
Roscoff-Scilly Islands
- S. peresi* Bodin, 1970  
W France
- S. exilis* (T. & A. Scott, 1894)

Bohuslän-S Norway-Br. Isles

*Parapseudoleptomesochra* Lang,1965

*P. polychaeta* (Noodt,1952)  
= *Nitocrella polychaeta* Noodt,1952  
Sylt-Exmouth

*Parevansula* Guille & Soyer,1966

*P. vermiformis* Moore,1976  
Isle of Man

*P. mediterranea* Guille % Soyer,1966  
?European Atlantic, Mediterranean

*Ameiropsyllus* Bodin,1977

*A. monardi* Bodin,1977  
W France

*Stenocopiinae* Lang,1948

*Stenocopia* G.O. Sars,1907

*S. longicaudata* (T. Scott,1892)  
Bohuslän-SW & W Norway-Helgoland-Br. Isles-Roscoff

*S. setosa* G.O. Sars,1907  
Bohuslän-SW & W Norway

*S. spinosa* (T. Scott,1892)  
off Gullmarn (Bonden)-Skagerrak-S & W Norway-Scotland-Ireland

*Malacopsyllus* G.O. Sars,1911

*M. fragilis* G.O. Sars,1911  
S & W Norway

*Anoplosoma* G.O. Sars,1911

*A. sordidum* G.O. Sars,1911  
S Norway

*Paramesochridae* Lang,1944, emend. Lang,1965

*Paramesochrinae* Lang,1944

*Paramesochra* T. Scott,1892

*P. acutata* Klie,1935  
Kieler Bucht-German North Sea Coast

*P. borealis* Geddes,1981  
Tromsø

*P. brevifurca* Galhano,1970  
NW Portugal-Sardinia

*P. dubia* T. Scott,1892  
off Gullmarn (Bonden)-S Norway-Helgoland-Br. Isles-W Ireland-Brittany

*P. helgolandica* Kunz,1936  
Belt Sea-W Sweden-Sylt-Helgoland-Belgium-Irish Sea-Bulgaria

*P. pterocaudata* Kunz,1936  
Bonden (Bohuslän)-Helgoland

*P. similis* Kunz,1936  
Helgoland-Isle of Man-Bulgaria

*P. longicaudata* Nicholls,1945  
?European Atlantic, Red Sea, W Australia

*P. mielkei* Huys,1987  
Netherlands

*Leptopsyllus* T. Scott,1894

*L. harveyi* Wells,1963  
Exmouth

*L. paratypicus* Nicholls,1939  
W Scotland

*L. reductus* Lang,1948  
Bohuslän

*L. typicus* T. Scott,1894  
W Scotland-W Mediterranean

*L. elongatus* Drzycimski,1967  
Bergen

*L. celticus* Bodin & Jackson,1987  
Bretagne-Galway Bay (Ireland)

*Remanea* Klie,1929

*R. arenicola* Klie,1929  
Gdansk-Simrishamn-Sylt-W Br. Isles-Portugal

*Kliopsyllus* Kunz,1962

*K. coelebs* (Monard,1935)  
= *Paramesochra coelebs* Monard,1935  
Exmouth-Wales-Brittany-Mediterranean

*K. constrictus constrictus* (Nicholls,1935)  
German Baltic-Sylt-Br. Isles-Portugal-Bulgaria

*K. constrictus orotavae* (Noodt,1958)  
= *Paramesochra orotavae* Noodt,1958  
?European Atlantic, Canaries, E Mediterranean

*K. holsaticus holsaticus* (Klie,1929)  
= *Paramesochra holsatica* Klie,1929 (TM)  
= *Paramesochra pygmaea* Nicholls,1939  
Kieler Bucht-Sylt-W Br. Isles-Portugal

*K. holsaticus longicaudatus* Galhano,1970  
NW Portugal

*K. laurenticus* (Nicholls,1939)  
= *Paramesochra laurentica* Nicholls,1939  
Exmouth

*K. longifurcatus* Scheibel,1975  
Kieler Bucht-Sylt

*K. major* (Nicholls,1939)  
= *Paramesochra major* Nicholls,1939  
Exmouth

*K. paraholsaticus* Mielke,1975  
Sylt-Belgium

*K. perhariensis* (Wells,1963)  
= *Paramesochra perhariensis* Wells,1963  
= *Paramesochra coelebs* : Bozic,1953, ?non Monard,1935  
Br. Isles-Roscoff-Angola

*K. longisetosus* (Krishnaswamy,1951)  
= *Paramesochra longisetosa* Krishnaswamy,1951  
?European Atlantic, Madras coast

*Scottopsyllus* Kunz,1962

*S. (S.) minor* (T. & A. Scott,1895) Kunz,1962  
= *Leptopsyllus minor* T. & A. Scott,1895  
= *Paramesochra minor* : Auctt.  
Kieler Bucht-Hällö in 6-8m-Sylt-Holland-Br. Isles-France-Bulgaria

*S. (S.) robertsoni* (T. & A. Scott,1895) Kunz,1962  
= *Leptopsyllus robertsoni* T. & A. Scott,1895  
= *Paramesochra robertsoni* : Auctt.  
Scotland-Bulgaria

*S. (S.) herdmanni* (I.C. Thompson & A. Scott,1900) Kunz,1962  
= *Leptopsyllus herdmanni* I.C. Thompson & A. Scott,1900  
= *Paramesochra herdmanni* (I.C. Thompson & A. Scott,1900)  
Gdansk-Kieler Bucht-S Norway-Sylt-W Br. Isles-Black Sea

*S. (Wellsopsyllus* Kunz,1981) *gigas* (Wells,1965)  
= *Paramesochra gigas* Wells,1965 (TM of *Wellsopsyllus* Kunz,1981)  
= *Kliopsyllus gigas* (Wells,1965)  
Scotland

*S. (Intermedopsyllus* Kunz,1962) *intermedius* (T. & A. Scott,1895)  
= *Leptopsyllus intermedius* T. & A. Scott,1895  
= *Paramesochra intermedia* (T. & A. Scott,1895)  
Kieler Bucht-Öresund-Hällö in 8m-Sylt-Helgoland-Netherlands-Br. Isles-Portugal-Bulgaria

*S. (Intermedopsyllus* Kunz,1962) *minutus* (Nicholls,1939)  
= *Paramesochra minuta* Nicholls,1939  
W Scotland

*Apodopsyllus* Kunz,1962

*A. africanus* Kunz,1962  
?European Atlantic

*A. africanus* Kunz,1962 *listensis* Mielke,1975  
Sylt

*A. arenicolus* (Chappuis,1954) Kunz,1962  
= *Leptopsyllus arenicola* Chappuis,1954  
W France-Algeria-Mozambique

*A. littoralis* (Nicholls,1939) Kunz,1962  
= *Leptopsyllus littoralis* Nicholls,1939  
W Scotland-Scilly Isles-Tunisia

*A. spinipes* (Nicholls,1939) Kunz,1962  
= *Leptopsyllus spinipes* Nicholls,1939  
W Scotland-W Mediterranean

*A. reductus* (Petkovski,1955) Kunz,1962  
= *Leptopsyllus reductus* Petkovski,1955  
= *Leptopsyllus perplexus* Wells,1963  
Br. Isles

*A. madrasensis* (Krishnaswamy,1951) Lang,1965  
= *Leptopsyllus madrasensis* Krishnaswamy,1951  
?European Atlantic

*Diarthrodellinae*

*Diarthrodella* Klie,1949  
= *Idyanthopsis* Bocquet & Bozic,1955

*D. orbiculata* Klie,1949  
=? *Idyanthopsis psammophila* Bocquet & Bozic,1955  
Helgoland-W Mediterranean (& Roscoff if the synonym is correct)

*D. parorbiculata parorbiculata* Wells,1963  
Ireland

*D. secunda secunda* Kunz,1954  
?Kieler Bucht-Sylt-Belgium-W Mediterranean

*Tetragonepsidae* Lang,1944, emend. Coull,1973

*Diagoniceps* Willey,1930

*D. brevicauda* Huys,1995 (nom. nud.?)  
France (La Rochelle)

*Aigondiceps* Fiers,1995

*A. bocki* (Lang,1948)  
= *Diagoniceps bocki* Lang,1948 (TM)  
Bohuslän-Skagerrak

*A. bodini* Fiers,1995  
France

*Paraschizopera* Wells,1981 (TM *P. beckeri* Wells,1981)  
= *Nidiagoceps* Fiers,1995

*P. menaensis* (Geddes,1968) Huys,1995  
= *Diagoniceps menaensis* Geddes,1968 (TM of *Nidiagoceps* Fiers,1995)  
W Br. Isles-W France

*Tetragoneiceps* Brady,1880

*T. scotti* G.O. Sars,1911  
S Norway-Scotland

*T. brevicauda* T. Scott,1899  
Scotland

*T. malleolatus* Brady,1880

Ireland-Scilly Isles-Roscoff-W Mediterranean

*T. bergensis* Por,1965  
Bergen

*Pteropsyllus* T. Scott,1906

*P. consimilis* (T. Scott,1894)  
= *P. plebeius plebeius* Monard,1935  
S & W Norway-Br. Isles-Ireland-Roscoff

*P. plebeius* Monard,1935 *furcatus* Kunz,1938  
\* Possibly conspecific with *P. consimilis* (T. Scott), but Klie,1971 won't agree with that synonymy. Should, if not identical, probably be named *P. consimilis furcatus* off Gullmarn (Bonden)-Helgoland

*Phyllopodopsyllus* T. Scott,1906  
= *Paraphyllopodopsyllus* Lang,1948

*P. bradyi* (T. Scott,1892)  
off Gullmarn (Bonden)-S & W Norway-Helgoland-Br. Isles

*P. furciger* G.O. Sars,1907  
= *P. furcifer*: Auctt.  
SW Norway-W Mediterranean

*P. hibernicus* (Roe,1955)  
= *Paraphyllopodopsyllus hibernicus* Roe,1955  
S Ireland

*P. hardingi* (Roe,1955)  
= *Paraphyllopodopsyllus hardingi* Roe,1955  
S Ireland

*P. longicaudatus* A. Scott,1909  
Scilly Isles (according to Wells,1961) & Indonesia

*Fearia* Coull,1971 (<sup>TM</sup> *F. prima* Coull,1971 - North Carolina)

Orthopsyllidae Huys,1991

*Orthopsyllus* Brady & Robertson,1873

*O. sarsi* Klie,1941  
= *O. linearis*: G.O. Sars,1909, non (Claus,1866)  
Norway (Skjærstadfjord)-Aberdeen

*O. linearis* (Claus,1866)  
= *O. agnatus* Klie,1950  
= *O. propinquus* Monard,1926  
Helgoland-Br. Isles-N France-Mediterranean

*O. linearis setosus* Boer,1971  
NW France

*O. linearis major* Klie,1939  
Iceland

*O. linearis illgi* (Chappuis,1958)  
= *Enhydrosoma illgi* Chappuis,1958  
?NW France

Canthocamptidae Brady,1880

*Barbaracletoles* Becker, Noodt & Schriever,1979

*Mesochra* Boeck,1865

*M. pygmaea* (Claus,1863)  
Kieler Bucht-Bohuslän-Sylt-Helgoland-S & N Norway-Arctic Seas-Br. Isles-Roscoff-Mediterranean-Black Sea

*M. lilljeborgi* Boeck,1865  
Finland-Askö-Bohuslän-S North Sea-S Norway-Trondheim-Br. Isles-N France-Mediterranean-Black Sea-Caspian Sea

*M. rapiens* (Schmeil,1894)  
Örnsköldsvik-Askö-S Baltic, S Norwegian & North Sea Estuaries, Br. Isles-S Ireland

*M. aestuarii* Gurney,1921  
Finland-German Baltic & North Sea river mouths, Br. Isles, Black Sea

*M. inconspicua* (T. Scott,1899)  
Belt Sea-S Norway-Sylt-Scotland

*M. armoricana* Monard,1935  
Roscoff, among *Enteromorpha* -S Ireland

*M. xenopoda* Monard,1935  
Roscoff-Mediterranean-Black Sea

*M. anomala* Klie,1950  
Helgoland-W France

*M. heldti* Monard,1935  
Exe Estuary (S England)-W France

*M. schmidti* Mielke,1974  
Spitsbergen

*M. pontica* Marcus,1965  
W France-Black Sea

*M. lindbergi* Petkovski,1964  
\* Is this a marine species?  
Portugal

*M. reducta* Klie,1950  
\* Dubious species.  
Helgoland

*Nannomesochra* Wilson,1932  
= *Pseudomesochra* Gurney,1927, non Scott,1902

*N. arupinensis* (Brian,1925)  
Belt Sea-Ireland-Roscoff-Mediterranean-Black Sea

*Itunella* Brady,1896

*I. tenuiremis* (T. Scott,1893)  
Bohuslän-Scotland

*I. muelleri* (Gagern,1922)  
= *I. bacescui* Chappuis & Serban,1953  
Brackish water sp.: Finland-Sweden-Germany-Ireland-France

*Bryocamptus* Chappuis,1928

*B. (Bryocamptus) pygmaeus* (G.O. Sars,1863)  
Freshwater species, which enters Baltic estuarine water

*B. (Bryocamptus) minutus* (Claus,18??)  
Freshwater species, which enters Baltic estuarine water

*Psammocamptus* Mielke,1975

*P. axi* Mielke,1975  
Sylt

*Hemimesochra* G.O. Sars,1920

*H. clavularis* G.O. Sars,1920  
S Norway

*H. nympa* Por,1964  
Kosterfjord in 200m-Skagerrak

*Pusillargillus* Huys & Thistle,1989

*P. nixe* (Por,1964)  
= *Hemimesochra nixe* Por,1964  
Kosterfjord in 200m-Skagerrak-Bergen

*Mesopsyllus* Por,1960

*M. secunda* (Wells,1965)  
= *Hemimesochra secunda* Wells,1965  
Scotland

*Heteropsyllus* T. Scott,1894  
= *Cletomesochra* G.O. Sars,1920

*H. nanus* (G.O. Sars,1920)  
Bohuslän-S Norway

*H. major* (G.O. Sars,1920)  
Belt Sea-Bohuslän-S & W Norway-W France

*H. exiguus* (G.O. Sars,1911)  
SW & W Norway

*H. masculus* Kunz,1971  
Helgoland

*H. curticaudatus* T. Scott,1894  
Br. Isles

*H. rostratus* (G.O. Sars,1920)  
Kosterfjord & off N Koster in 150-200m-S Norway-Arctic Seas

*H. meridionalis* Soyer,1975  
?European Atlantic, Mediterranean

*H. confluens* Soyer,1975  
?European Atlantic, Mediterranean

*Bathycamptus* Huys & Thistle,1989

*B. minutus* (Wells,1965)  
= *Heteropsyllus minutus* Wells,1965  
Scotland

*Cletocamptus* Schmankewitsch,1875  
\* Should possibly be included in Cletodidae.

*C. confluens* (Schmeil,1894)  
S Baltic river mouths

*Boreolimella* Huys & Thistle,1989

*B. dubia* (Wells,1965)  
= *Leimia* Willey,1925 *dubia* Wells,1965  
Scotland

*Parepactophanes* Kunz,1935  
\* Should possibly be included in Cletodidae.

*P. minuta* Kunz,1935  
Kieler Bucht-German North Sea Coast

Cylindropsyllidae G.O. Sars,1909, emend. Lang,1948

*Cylindropsyllus* Brady,1880  
= *Cylindrosoma* Brady,1878, non Tschudi,1838, nec Gray, in Todd, 1843, nec Rondani,1856

*C. laevis* Brady,1880 (<sup>TM</sup>)  
Belt Sea-Bohuslän-S & W Norway-Helgoland-Br. Isles-N & W France-Mediterranean

*C. remanei* Kunz,1949  
Helgoland-Ireland-Isle of Man-Portugal

*C. kunzi* Huys,1988  
Netherlands Delta region

*C. ibericus* Huys & Willems,1993  
Portugal

*C. govaerei* Huys & Willems,1993  
Corsica

*Evansula* T. Scott,1906

*E. incerta* (T. Scott,1892)  
Kieler Bucht-off Gullmarn (Bonden)-Hällö-German Bight-Lofoten-Br. Isles-Arcachon

*E. pygmaea* (T. Scott,1903)  
Kieler Bucht-Sylt-Helgoland-Belgium-SW England-Scotland

*Stenocarix* G.O. Sars,1909

*S. baltica* Artl,1983  
S Baltic

*S. gracilis* G.O. Sars,1909  
Korshavn (S Norway)-Aberdeen

*S. kliei* Kunz,1936  
Helgoland-Isle of Man

*S. minor* (T. Scott,1892)  
Bonden (Bohuslän)-Korshavn (S Norway)-Helgoland-W Scotland-Roscoff-Bulgaria

*S. minuta* Nicholls,1935

- Poland-Kieler Bucht-S Norway-Sylt-W Br. Isles-Bulgaria
- S. pygmaea* Noodt,1955  
Aberdeen-Arcachon-W France-Biscay
- S. pontica* Chappuis & Serban,1953  
Black Sea
- Boreopontia* Willems,1981
- B. heipi* Willems,1981 (TM - Belgium)  
Netherlands, Belgium, Finisterre
- Willemsia* Huys & Conroy-Dalton,1993
- W. calceola* Huys & Conroy-Dalton,1993 (TM - S North Sea)  
Netherlands, Belgium
- Stenocaropsis* Apostolov,1982
- S. pristina* (Wells,1968)  
= *Stenocaris pristina* Wells,1968  
Scilly Isles-Black Sea
- Cylinula* Coull,1971 (TM *Cylinula proxima* Coull,1971 - N Carolina)
- C. arganoi* (Cottarelli & Venanzetti,1989)  
Corsica, Italy
- Leptopontiidae Lang,1944
- Leptopontia* T. Scott,1902, emend. Huys & Conroy-Dalton,1996
- L. curvicauda* T. Scott,1902, emend Huys & Conroy-Dalton,1996  
?Kieler Bucht-off Gullmarn (Bonden)-Skagerrak-Helgoland-W Br. Isles-Bulgaria
- L. doypori* Huys & Conroy-Dalton,1996  
England
- L. flandrica* Huys & Conroy-Dalton,1996  
North Sea
- L. punctata* Huys & Conroy-Dalton,1996  
Sweden
- Syrticola* Willems & Claeys,1982
- S. flandricus* Willems & Claeys,1982 (TM)  
Belgium
- S. trispinosus* (A. Scott,1896)  
= *Tetragoneiceps trispinosus* A. Scott,1896  
= *Evansula trispinosus* : Auctt.  
= *Evansia trispinosus* : Auctt.  
\* Considered a species inquirenda by Huys & Ohtsuka 1993  
Isle of Man
- Psammopsyllinae Krishnaswamy,1956
- Psammopsyllus* Nicholls,1945  
= *Paulianicaris* Chappuis,1952
- P. arenarius* Enckell,1965  
Spain
- Ichnusella* Cottarelli,1971 (TM *I. eione* Cottarelli,1971 - Italy, interstitial)
- Arenopontiidae
- Arenopontia* Kunz,1937
- A. (Arenopontia) stygia* Noodt,1955  
Arcachon-W France-NW Portugal-Bulgaria
- A. (Arenopontia) subterranea* Kunz,1937  
S Baltic-German Bight-Scotland-Scilly Isles-Mediterranean-?Black Sea
- A. (Neoleptastacus* Nicholls,1945) *spiniacaudata* (Nicholls,1945)  
W France
- Pararenopontia* Bodiou & Colomines,1986
- P. breviarticulata* (Mielke,1975)  
= *Arenopontia (Arenopontia) breviarticulata* Mielke,1975  
Sylt
- Leptastacidae Lang,1944 (Huys, 1992 upgraded this to full family rank)
- Leptastacus* T. Scott,1892
- L. aberrans* Chappuis,1954  
Arcachon-W Mediterranean
- L. laticaudatus laticaudatus* Nicholls,1935  
Belgium-W Br. Isles-Arcachon
- L. laticaudatus intermedius* Kunz,1938  
?Kieler Bucht-Hällö in 6-8m-Helgoland-Belgium-W Br. Isles-Brittany-Bulgaria
- L. macronyx* (T. Scott,1892)  
S Baltic-off Gullmarn (Bonden)-North Sea-S Norway-Br. Isles-Bulgaria
- L. minutus* Chappuis,1954  
Belgium-Portugal
- L. rostratus* Nicholls,1939  
Tromsö-Canada
- L. taurica* Marinov,1974  
\* Not included in Bodin's catalogue from 1979.  
?European Atlantic, Black Sea
- Paraleptastacus* C.B. Wilson,1932
- P. espinulatus* Nicholls,1935  
Kieler Bucht-Belgium-Br. Isles
- P. spinicauda* (T. & A. Scott,1895)  
Finland-Estonia-Askö-Kattegatt-German Bight-Br. Isles-Roscoff-Portugal-Black Sea
- P. holsaticus* Kunz,1937  
Belt Sea-North Sea-Scotland
- P. supralitoralis* Mielke,1975  
Sylt
- P. kliei* (Gagern,1923)  
S Norway
- Psammastacus* Nicholls,1935
- P. brevicaudatus* Nicholls,1935  
W Scotland
- P. confluentis* Nicholls,1935  
?Belt Sea-Nordfriesland-E & W Scotland-N & W France-Algeria-Bulgaria
- P. perplexus* Wells,1965  
Scotland-W Portugal
- P. remanei* Noodt,1964  
Belgium
- Psamathea* Cottarelli & Venanzetti,1989 (TM *P. nautarum* Cottarelli & Venanzetti,1989)
- P. britannica* Huys, Bodiou & Bodin,1996  
Brittany
- Arenocaris* Nicholls,1935
- A. bifida* Nicholls,1935  
German Baltic-Nordfriesland-Engl. Channel-Scotland
- Schizothrix* Huys,1992
- Parastenocarididae Chappuis,1933  
\* A freshwater species family.
- Parastenocaris* Kessler,1913
- P. fontinalis fontinalis* Schnitter & Chappuis,1915  
Freshwater species; may be found in Baltic estuaries
- P. fontinalis borea* Kiefer,1960  
Freshwater species; may be found in estuaries in N Europe
- P. phyllura* Kiefer,1938  
Baltic brackish water (Swe., Denm., Finl.), e.g. Askö-Nordfriesland
- P. vicesima* Klie,1935  
Finland-Askö-S Baltic-S Norway-Nordfriesland-Anglesey-Aberdeen
- Louriniidae Monard,1927  
\* No species closer than in the Mediterranean?
- Cletodoidea T. Scott,1904
- Rhizothricidae Por,1986
- Rhizothrix* G.O. Sars,1909  
= *Rhizothrix* Brady & Robertson,1875 (nom. nud.?)
- R. curvata* Brady & Robertson,1880  
Öresund-Bohuslän-SW Norway-Helgoland-Br. Isles-Brittany-Mediterranean-Black Sea
- R. gracilis* (T. Scott,1903)  
German Baltic-S Norway-Sylt-Helgoland-Br. Isles-W Mediterranean
- R. minuta* (T. Scott,1903)  
Kieler Bucht-Öresund-Helgoland-Brittany-Br. Isles & Ireland
- R. reducta reducta* Noodt,1952  
Northfriesland-Arcachon-?S France
- R. reducta noodi* Galhano,1970  
Sylt-Portugal
- R. wilsoni* Bodin,1979  
W France
- Tryphoema* Monard,1926  
= *Adelopoda* Pennak,1942
- T. bocqueti* (Bozic,1953)  
= *Rhizothrix bocqueti* Bozic,1953  
Hällö in 6-8m-Sylt-W Br. Isles-Roscoff-Arcachon
- T. lusitanica* (Wells & Clark,1965)  
W Br. Isles-W Portugal
- T. porca* Monard,1926  
off Gullmarn (Mittskären in 27-30m)-Aberdeen-Isle of Man-Exmouth-Roscoff-Marseille
- T. scillonensis* (Wells,1968)  
Scilly Isles
- Huntemanniidae Por,1986
- Huntemannia* Poppe,1884
- H. jadensis* Poppe,1884  
Finland-Askö-Öresund, Nordfriesland, Trondheim, Scotland
- H. micropus* Monard,1935  
W Br. Isles-Roscoff
- Pseudocletodes* T. & A. Scott,1893
- P. vararensis* T. & A. Scott,1893  
Scotland
- Metahuntemannia* Smirnov,1946  
= *Apodella* Por,1965
- M. drzycimskii* Soyer,1970  
= *M. smirnovi* Drzycimski,1968, non Bodin,1968  
Bergen
- M. spinosa* (Klie,1941)  
Iceland
- M. crassa* (Por,1965)  
= *Apodella crassa* Por,1965 (TM of *Apodella* Por,1965)  
Bergen
- Pontopolites* T. Scott,1894
- P. typicus* T. Scott,1894  
Kieler Bucht-Bohuslän-SW Norway-Nordfriesland-Br. Isles
- Nannopus* Brady,1880

- N. palustris* Brady, 1880  
Finland & Stockholm skerries-S Baltic-Oslofjord-Telemark-German Bight-Netherlands-Iceland-Br. Isles
- Danielsseniidae* Huys & Gee, 1996  
= *Paranannopidae* Por, 1986
- Carolinicola* Huys & Thistle, 1989  
\* Placed as incertae sedis in family.
- Cylindronannopus* Coull, 1973
- Mucrosenia* Gee & Huys, 1994
- M. kendalli* Gee & Huys, 1994 (TM - Spitsbergen)  
Spitsbergen
- M. kliei* (Smirnov, 1946)  
= *Psammis kliei* Smirnov, 1946  
?
- Bathysammis* Huys & Gee, 1993 (TM *Psammis longifurca* Bodin, 1968)
- Leptotachidia* Becker, 1974
- Paranannopus* Huys & Gee, 1996  
= *Paranannopus* Lang, 1936 (nom. nud.?)
- P. abyssi* (G.O. Sars, 1920)  
S Norway
- P. bahusiense* Por, 1964  
off N Koster in 110m
- P. triarticulatus* Wells, 1965  
= *P. langi* Wells, 1965  
\* The female was described as *P. triarticulatus* and the male as *P. langi*  
Scotland
- P. aff. philistinus* Por, 1964  
Celtic Sea
- P. sarsi* Lang, 1936  
Spitsbergen
- Danielssenia* Boeck, 1872
- D. typica* Boeck, 1872  
= *D. fusiformis* : G.O. Sars, 1910, non Brady, 1880  
Poland-Bohuslän-W & N Norway-Spitsbergen-Arctic Seas, Helgoland-Br. Isles
- D. quadriseta* Gee, 1988  
Norway
- D. spitsbergensis* Gee & Huys, 1994  
Spitsbergen
- Jonesiella* Brady, 1880  
= *Sentirenia* Huys & Gee, 1992
- J. fusiformis* Brady, 1880  
= *Danielssenia perezii* Monard, 1935 (TM of *Sentirenia* Huys & Gee, 1992)  
= *Danielssenia paraperezii* Soyer, 1970  
Firth of Clyde-Liverpool Bay-Scilly Islands-Roscoff-Mediterranean
- Paradanielssenia* Soyer, 1970 (TM *P. kunzi* Soyer, 1970 - Mediterranean)
- P. biclavata* Gee, 1988  
S Celtic Sea
- P. christineae* Gee & Huys, 1994  
Spitsbergen
- P. kathleenae* Gee & Huys, 1994  
Spitsbergen
- Fladenia* Gee & Huys, 1990
- F. robusta* (G.O. Sars, 1921)  
= *Danielssenia robusta* G.O. Sars, 1921  
= *Danielssenia intermedia* Wells, 1965 (TM)  
Bohuslän-Bergen, Scotland
- Psammis* G.O. Sars, 1910
- P. longisetosa* G.O. Sars, 1910  
Bohuslän-Bergen, Scotland
- Anapophysia* Huys & Gee, 1996
- A. borealis* (Klie, 1939)  
= *Psammis borealis* Klie, 1939 (TM)  
Iceland
- Micropsammis* Mielke, 1975
- M. noodti* Mielke, 1975 (TM)  
Sylt
- Telopsammis* Gee & Huys, 1991
- T. secunda* (Mielke, 1975)  
= *Micropsammis secunda* Mielke, 1975 (TM)  
Sylt
- Archisenia* Huys & Gee, 1992
- A. sibirica* (G.O. Sars, 1898)  
= *Danielssenia sibirica* G.O. Sars, 1898 (TM)  
= *Archisenia stefanssonii* (Willey, 1920)  
?Scotland, Arctic Seas
- Afrosenia* Huys & Gee, 1996 (TM *Danielssenia spinipes* Wells, 1967)
- Sentiropsis* Huys & Gee, 1996 (TM *Danielssenia minuta* Coull, 1969)
- Peltsenia* Huys & Gee, 1996 (TM *Idomene aberrans* Por, 1964)
- Prionos* Huys & Gee, 1996 (TM *P. ornata* Huys & Gee, 1996 - Malaysia)
- Cithadius* Bowman, 1972 (TM *C. cythurae* Bowman, 1972 - ? - @ *Cyathura polita*)
- Argestidae* Por, 1986
- Neoargestes* Drzycimski, 1967
- N. variabilis* Drzycimski, 1967  
Bergen
- Parargestes* Lang, 1948
- P. tenuis* (G.O. Sars, 1921)  
Bohuslän-S & W Norway-Spitsbergen
- Argestes* G.O. Sars, 1910
- A. mollis* G.O. Sars, 1910  
Bohuslän-W Norway-Arctic Seas
- Fultonia* T. Scott, 1902
- F. hirsuta* T. Scott, 1902  
Bohuslän-SW & W Norway-Scotland
- Leptocletodes* G.O. Sars, 1920
- L. debilis* G.O. Sars, 1920  
Bohuslän-S & W Norway
- Eurycletodes* G.O. Sars, 1909
- E. (Eurycletodes) laticauda* (Boeck, 1872)  
Bohuslän-Haugesund
- E. (Eurycletodes) serratus* G.O. Sars, 1920  
Bohuslän
- E. (Oligocletodes) langi* (Lang, 1948) *latus* (T. Scott, 1892)  
SW & W-?N Norway-Br. Isles
- E. (Oligocletodes) major* G.O. Sars, 1909  
Øresund-Bohuslän-S Norway-Arctic Seas
- E. (Oligocletodes) similis* (T. Scott, 1895)  
Bohuslän-Helgoland-Lofoten-Arctic Seas-Br. Isles-Roscoff
- E. (Oligocletodes) minutus* G.O. Sars, 1920  
Bohuslän-Bergen
- E. (Oligocletodes) aculeatus* G.O. Sars, 1920  
Bohuslän-Skagen-S & W Norway
- E. (Oligocletodes) oblongus* G.O. Sars, 1920  
Bohuslän-Skagen-S Norway
- E. (Oligocletodes) pori* Drzycimski, 1969  
= *E. cf. echinatus* : Por, 1965  
Bergen
- E. (Oligocletodes) irlandica* Roe, 1959  
S Ireland
- Mesocletodes* G.O. Sars, 1909
- M. monensis* (L.C. Thompson, 1893)  
Bohuslän-S & W Norway-Br. Isles
- M. irrasus* (T. & A. Scott, 1894)  
Bohuslän-Bergen-Scotland
- M. inermis* G.O. Sars, 1920  
S Norway
- M. arenicola* Noodt, 1952  
Sylt
- M. abyssicola* (T. & A. Scott, 1901)  
Bohuslän-Bergen-Spitsbergen
- M. robustus* Por, 1965  
Bergen
- M. glaber* Por, 1964  
Skagerrak in 300-425m-Bergen
- M. fladensis* Wells, 1965  
Scotland
- M. duosetosus* Schriever, 1985  
Iceland-Faeroe-ridge (500-1850 m)
- M. thieli* Schriever, 1985  
Iceland-Faeroe-ridge (500-1850 m)
- M. faeroensis* Schriever, 1985  
Iceland-Faeroe-ridge (500-1850 m)
- M. kunzi* Schriever, 1985  
Iceland-Faeroe-ridge (500-1850 m)
- M. quadrispinosa* Schriever, 1985  
Iceland-Faeroe-ridge (500-1850 m)
- M. carpinei* Soyer, 1975  
Celtic Sea, Mediterranean
- M. katharinae* Soyer, 1964  
Celtic Sea, Mediterranean
- Argestigens* Willey, 1935
- A. uniremis* Willey, 1935 (TM - Bermudas)  
?European Atlantic, Bermudas
- Cletodidae* T. Scott, 1904
- Monocletodes* Lang, 1936
- M. varians* (T. Scott, 1903)  
= *Thieliella nordatlantica* Schriever, 1982 (TM of *Thieliella* Schriever, 1982)  
= *Thieliella reducta* Schriever, 1982  
Norway (Bøgfjord), Iceland-Faeroer-Ridge
- Acrenhydrosoma* Lang, 1944
- A. perplexum* (T. Scott, 1899)  
= *Cletodes perplexa* T. Scott, 1899  
= *Enhydrosoma perplexum* (T. Scott, 1899) Lang, 1936  
Bohusl-Norway (Bøgfjord)-Br. Isles

- Stylectodes* Lang, 1936
- S. longicaudatus* (Brady & Robertson, 1880)  
= *S. numidicus* (Monard, 1935)  
Bohuslän-S Norway-Arctic Seas-Br. Isles
- S. reductus* Wells, 1965  
Scotland
- Cletodes* Brady, 1872
- C. endopodita* (Schriever, 1984)  
= *Thieliella endopodita* Schriever, 1984  
Norw. Sea
- C. pusillus* G.O. Sars, 1920  
Öresund-Bohuslän-Helgoland-S & W Norway
- C. tenuipes* T. Scott, 1896  
Öresund & Belt Sea-Bohuslän-Helgoland-N Norway-Arctic Seas-Br. Isles-Ireland
- C. longifurca* Lang, 1948  
Bohuslän-Bergen
- C. longicaudatus* (Boeck, 1872)  
Bohuslän-SW & W Norway-Br. Isles-Roscoff-Mediterranean
- C. limicola* Brady, 1872  
Bohuslän-SW Norway-Br. Isles-Mediterranean
- C. latirostris* Drzycimski, 1967  
=? *Enhydrosoma wellsii* Bodin, 1968  
Bergen & ?W France
- C. smirnovi* Bodin, 1970  
W France
- C. spinulatus* Por, 1967  
Bretagne
- Enhydrosoma* Boeck, 1872
- E. sarsi* (T. Scott, 1904)  
Öresund & Belt Sea-Bohuslän-S & W Norway-Scotland-W France
- E. longifurcatum* G.O. Sars, 1909  
\* Is together with *E. sordidum* & *E. gariene* constituting a natural group  
Kieler Bucht-Bohuslän-Nordfriesland-SW & W Norway-Scotland-Ireland-N & W France
- E. sordidum* Monard, 1926  
Scotland-Roscoff-Banyuls
- E. gariene* Gurney, 1930  
Nordfriesland-England (Norfolk)
- E. propinquum* (Brady, 1880)  
= *Cletodes propinqua* Brady, 1880  
Bohuslän-S Norway-German Bight-Br. Isles-Roscoff-Mediterranean
- E. curticauda* Boeck, 1872 (TM)  
= *Cletodes hirsutipes* T. Scott, 1896  
Kieler Bucht-Bohuslän-N Norway-White Sea-German Bight-Zuidersee-Br. Isles-Ireland
- E. curvirostre* (T. Scott, 1894)  
\* This will possibly be transferred to a new genus in a forthcoming paper (sensu Gee, 1994)  
Kieler Bucht-Bohuslän-N Norway-Br. Isles
- Strongylacron* Gee & Huys, 1996
- S. buchholzi* (Boeck, 1872)  
= *Cletodes buchholzii* Boeck, 1872 (TM)  
= *Enhydrosoma buchholzii* (Boeck, 1872)  
Kieler Bucht-Bohuslän-German North Sea coast-Haugesund, Anglesey, Ireland, S England, W France
- Enhydrosomella* Monard, 1935
- Scintis* Por, 1986  
\* *Cletodidae* incertae sedis.
- Interclotodes* Fiers, 1987
- Schizacron* Gee & Huys, 1996 (TM *Enhydrosoma barnishi* Wells, 1987 - Mozambique)
- Triathrix* Gee & Burgess, 1997 (TM *T. montagni* Gee & Burgess, 1997 - California)
- Sphingothrix* Fiers, 1997 (TM *S. goldi* Fiers, 1997 - Yucatan shelf)
- Laophontopsidae* Huys & Willems, 1989
- Laophontopsis* G.O. Sars, 1908  
= *Cleta* Claus, 1863, *non* Duponchel, 1845
- L. lamellifera* (Claus, 1863)  
Bohuslän-S Norway-Br. Isles-N France-Mediterranean
- L. borealis* Huys & Willems, 1989  
Hvaler, Netherlands, Norfolk
- L. monardi* Huys & Willems, 1989  
France
- Aculeopsis* Huys & Willems, 1989 (TM *A. longisetosa* Huys & Willems, 1989 - Mediterranean France)
- Telodocus* Huys & Willems, 1989 (TM *Cleta secunda* Sewell, 1924)
- Normanellidae Lang, 1944  
\* Not recognized by Huys & Boxshall 1991 but raised from subfamily to family by Huys & Willems 1989
- Normanellinae Lang, 1944
- Normanella* Brady, 1880
- N. similis* Lang, 1936  
Öresund & Belt Sea-Bohuslän-Skagen
- N. mucronata* G.O. Sars, 1909  
Bohuslän-Bergen-Roscoff
- N. tenuifurca* G.O. Sars, 1909  
Bohuslän-Bergen-Bukken-Roscoff
- N. minuta* (Boeck, 1872)  
Bohuslän-Helgoland-Trondheim-Ireland-Wales-Roscoff-Bermudas-Mediterranean
- N. quarta* Monard, 1935
- Roscoff-Algeria
- N. dubia* Brady & Robertson, in Brady, 1880  
\* Dubious species  
Br. Isles-Ireland
- Cletopsyllinae* Huys & Willems, 1989
- Cletopsyllus* Willey, 1935
- Pseudocletopsyllus* Vervoort, 1964
- Adenopleurellidae* Huys, 1990
- Sarsocletodes* Wilson, 1924  
= *Pseudocletodes* G.O. Sars, 1920, *non* Scott, 1893
- S. typicus* (G.O. Sars, 1920)  
Bohuslän-Bergen
- Adenopleurella* Huys, 1990 (TM *A. brevipes* Huys, 1990 - Queensland)
- Miroslavia* Apostolov, 1980
- Proceropes* Huys, 1990 (TM *Sarsocletodes secundus* Smirnov, 1946)
- Laophontidae* T. Scott, 1904
- Laophonte* Philippi, 1840
- L. inopinata* T. Scott, 1892  
Hällö- off N Koster-S-N Norway-Helgoland-Iceland-White Sea-Br. Isles
- L. denticornis* T. Scott, 1894  
Bohuslän-SW Norway-Helgoland-Br. Isles-Ireland
- L. nordgaardi* G.O. Sars, 1908  
Norway (Saltfjord)
- L. brevifurca* G.O. Sars, 1920  
Norway (Hvalör)-Black Sea
- L. baltica* Klie, 1929  
?Kieler Bucht-Kattegatt-Helgoland-S Ireland & White Sea
- L. setosa* Boeck, 1865  
?German Baltic-Oslofjord-W Norway-German North Sea coast-Iceland-Br. Isles-N France-Mediterranean
- L. thoracica* Boeck, 1865  
Kieler Bucht-Bohuslän-Helgoland-S Norway-Trondheim-Br. Isles-N France-Mediterranean
- L. longicaudata* Boeck, 1865  
= *Laophonte Hodgii* Brady, 1872  
Bohuslän-Trondheim-Br. Isles-Ireland
- L. longicaudata reducta* Lang, 1936  
Öresund
- L. elongata* Boeck, 1872  
Bohuslän-W Norway-Helgoland-Ireland-Roscoff-Mediterranean
- L. inornata* A. Scott, 1902  
S England-Roscoff-Mediterranean
- L. parvuloides* Monard, 1935  
Roscoff
- L. parvula* G.O. Sars, 1908  
= *Laophonte sima* Gurney, 1927  
Bohuslän-S Norway-Helgoland-W France-Mediterranean
- L. serrata* (Claus, 1863)  
= *Cleta serrata* Claus, 1863  
Bohuslän-Trondheim-Helgoland-Br. Isles-Ireland
- L. dominicalis* Monard, 1935  
Roscoff
- L. depressa* T. Scott, 1894  
W Norway-Arctic Seas-Iceland-Br. Isles
- L. cornuta* Philippi, 1840  
Bohuslän-Trondheim-Shetlands-Br. Isles-N France-Canaries-Madeira-Mediterranean
- L. trilobata* Willey, 1929  
= *Laophonte foxi* Harding, 1956  
Sylt-Helgoland-Norfolk-North America
- L. drachi* Médioni & Soyser, 1967  
= *L. danversae* Hamond, 1969  
@ *Schismopora armata* (Hincks, 1860)  
Norfolk-France
- ? *L. royi* Jakubisiak, 1932  
@ ? *Maia squinado*  
Roscoff
- L. farrani* Roe, 1958  
\* May belong to a not yet described genus  
S Ireland
- L. commensalis* Raibaut, 1961  
@ among pilose setae, especially on the ventral surface of *Xantho pilipes*, *X. floridus* & *X. rivulosus*  
N France & Mediterranean
- L. adamsiae* Raibaut, 1966  
@ *Adamsia carcinopados*  
French Mediterranean
- Laophontina* Norman & T. Scott, 1905
- L. acantha* Noodt, 1955  
W France-Portugal
- L. dubia* Norman & T. Scott, 1905  
Anglesey-Scilly Isles
- Platyhelipus* Brady, 1880
- P. laophontoides* G.O. Sars, 1908  
Kieler Bucht-Bohuslän-Oslofjord-Helgoland-NE Br. Isles
- P. littoralis* Brady, 1880  
Oslofjord-German Bight-N Norway-Arctic Seas-Br. Isles-Roscoff
- Pseudolaophonte* A. Scott, 1896

- P. spinosa* (I.C. Thompson, 1893)  
Bohuslän-S Norway-Helgoland-Scotland-Isle of Man-Scilly Isles
- P. proteus* Klie, 1950  
Belt Sea
- P. glemareci* Bodin, 1977  
Roscoff
- Pilifera* Noodt, 1952
- P. gracilis* (T. Scott, 1903)  
= *Laophonte gracilis* T. Scott, 1903 (TM)  
= *Pseudonyhocamptus gracilis* : Lang, 1948  
Sylt, Firth of Forth
- Harrietella* T. Scott, 1906  
@ external body associates of crustaceans
- H. simulans* (T. Scott, 1894)  
@ Attached to especially the telson of *Limnoria lignorum* (Rathke) & *L. tripunctata* Menzies or in their excavated galleries.  
Oslofjord-Trondheim-SE & SW Scotland-Wales-Netherlands, NE Ireland, Mediterranean, E USA
- Hemilaophonte* Jakubisiak, 1932
- H. janinae* Jakubisiak, 1932  
@ On carapace of *Maia squinado*  
Roscoff
- Esola* Edwards, 1891
- E. typhlops* (G.O. Sars, 1908)  
Bohuslän-Flekkerøy-Bergen-Scotland
- E. longicauda* Edwards, 1891  
Scotland-Ireland-Bahamas-Bermudas-Mediterranean
- E. longiremis* (T. Scott, 1904)  
Scotland-Scilly Isles
- E. bulligera* (Farran, 1913)  
Ireland
- Onychocamptus* Daday, 1903
- O. mohammed* (Blanchard & Richardson, 1891)  
Finland-S Baltic, Elbe mouth, Zuidersee and other Eur. brackish water sites
- O. chathemensis* (G.O. Sars, 1905)  
S Africa-Chatham Islands-?Europe
- Klieonyhocamptus* Noodt, 1958
- K. kliei* (Monard, 1935)  
\* Lang, 1965 doubt that this is a natural genus  
Scilly Isles-Roscoff-Yugoslavia
- Echinolaophonte* Nicholls, 1941
- E. horrida* (Norman, 1876)  
= *Onychocamptus horridus* : Auctt.  
Kieler Bucht-Bohuslän-Helgoland-S & N Norway-Arctic Seas-Greenland-Iceland-Br. Isles-Mediterranean
- E. brevispinosa* (G.O. Sars, 1908)  
S Norway-Mediterranean
- Asellopsis* Brady & Robertson, 1873
- A. hispida* Brady & Robertson, 1873  
Kieler Bucht-Bohuslän-S Norway-Bergen-Nordfriesland-Br. Isles-Roscoff-Mediterranean
- A. intermedia* (T. Scott, 1895)  
Kieler Bucht-Hällö- off N Koster-German Bight-Arctic Seas-Br. Isles-Roscoff-Algeria
- Heterolaophonte* Lang, 1944
- H. minuta* (Boeck, 1872)  
= *Laophonte subsalsa* Brady, 1903  
Belt Sea-Bohuslän-Oslofjord-Telemark-Helgoland-Scotland-Wales-Ireland-Scilly Isles & White Sea
- H. stroemii* (Baird, 1837) (original spelling emended)  
= *Cyclops Stromii* Baird, 1837  
Kattegat-German Bight-W Norway-Arctic Seas-Greenland-Shetlands-Br. Isles-N France
- H. stroemi paraminuta* Noodt, 1955  
European Atlantic, Mediterranean, Marmara Sea
- H. stroemi* var. *brevicaudata* (Monard, 1928)  
?European Atlantic, Canaries, Mediterranean
- H. littoralis* (T. & A. Scott, 1893)  
Kieler Bucht-Bohuslän-Helgoland-Bergen-Br. Isles-Mediterranean
- H. longisetigera* Klie, 1950  
= *H. littoralis longisetigera* Klie, 1950  
Belt Sea-Helgoland-NW France
- H. mendax* (Klie, 1939)  
= *Laophonte mendax* Klie, 1939  
Iceland
- H. uncinata* (Czerniavski, 1868)  
= *Laophonte vietzi* Klie, 1941  
Roscoff-Mediterranean-Black Sea
- H. phycobates* (Monard, 1935)  
Roscoff
- H. norvegica* Drzycimski, 1968  
Bergen
- H. hamondi* Hicks, 1975  
= *Heterolaophonte bisetosa* Mielke, 1975  
Sylt, Norfolk
- H. denticulata* Roe, 1958  
S Ireland
- H. brevipes* Roe, 1958  
S Ireland
- Quinquelaophonte* Wells, Hicks & Coull, 1982
- Q. quinquispinosa* (Sewell, 1924)  
= *Laophonte quinquispinosa* Sewell, 1924  
= *Heterolaophonte sigmoides* : Wells, 1963, non (Willey, 1930)
- ?Exe estuary, Mediterranean, India
- Pseudonyhocamptus* Lang, 1944
- P. koreni* (Boeck, 1872)  
Bohuslän-S & W Norway-White Sea-German Bight-Ireland-Scilly Isles-Tunesia
- P. proximus* (G.O. Sars, 1908)  
Kieler Bucht-Sylt-W Norway-Scotland-Isle of Man
- P. abbreviatus* (G.O. Sars, 1920)  
S & W Norway
- P. carthyi* Hamond, 1968  
Norfolk
- ? *P. feroensis* (T. Scott, 1902)  
= *Laophonte feroensis* T. Scott, 1902  
Faeroe Channel
- Paralaophonte* Lang, 1944
- P. brevisrostris* (Claus, 1863)  
= *Laophonte herdmanni* A. Scott, 1902  
Bohuslän-W Norway-Br. Isles-Roscoff-Mediterranean-Black Sea
- P. congenera* (G.O. Sars, 1908)  
@ *Limnoria*  
Bohuslän-W Norway-Ireland-Roscoff-Bermudas-Mediterranean
- P. perplexa* (T. Scott, 1898)  
W-N Norway-Spitsbergen-Arctic Seas-Iceland
- P. macera* (G.O. Sars, 1908)  
S-W Norway-Arctic Seas
- P. karmensis* (G.O. Sars, 1911)  
Norway (Skutesnæs)-White Sea
- P. tenera* (G.O. Sars, 1920)  
S Norway
- P. hyperborea* (G.O. Sars, 1909)  
Spitsbergen-White Sea
- P. spitzbergensis* Mielke, 1974  
Spitsbergen
- Paronyhocamptus* Lang, 1944
- P. curticaudatus* (Boeck, 1865)  
= *Laophonte curticaudata* Boeck, 1964  
Kieler Bucht-Bohuslän-Karmøya-Telemark-Sylt-Helgoland-Scotland-Wales-Ireland-Roscoff-Mediterranean
- P. nanus* (G.O. Sars, 1908)  
= *Laophonte nana* G.O. Sars, 1908  
Finland-Bohuslän-Oslofjord-Nordfriesland-Zuidersee-N France-Mediterranean
- P. exiguus* (G.O. Sars, 1905)  
= *Laophonte exigua* G.O. Sars, 1905  
?European Atlantic, Chatham Island
- Namakosiramia* Ho & Perkins, 1977  
@ one sp. *N. californiensis* Ho & Perkins, 1977 on *Stichopus parvimensis* (Clark)  
\* Huys 1988 moved this to Laophontidae (Harpacticoida) from Namakosiramiidae (Cyclopoida)
- Cristacoxidae* Huys, 1990
- Cristacoxa* Huys, 1990 (TM *C. petkovskii* Huys, 1990 - Lesser Antilles)
- Cubanocleta* Petkovski, 1977 (TM *C. noodti* Petkovski, 1977 - Cuba)
- Noodtortopsyllus* Lang, 1965
- Ancorabolidae* G.O. Sars, 1909
- Laophontodinae* Lang, 1948
- Laophontodes* T. Scott, 1894  
= *Merope* Thomson, 1883, non Newman, 1838 (Mecoptera), nec Adams, 1856 (Mollusca), nec Martens, 1860  
ex Albers MS (Mollusca)  
= *Meropia* Wilson, 1932
- L. gracilipes* Lang, 1936  
Bohuslän
- L. expansus* G.O. Sars, 1908  
Bohuslän-Norway (Saltenfjord)-?Ireland
- L. bicornis* A. Scott, 1896  
Bohuslän-W Norway-Br. Isles-Roscoff-Mediterranean
- L. typicus* T. Scott, 1894  
Bohuslän-Helgoland-W Norway-Arctic Seas-Br. Isles-Roscoff
- L. armatus* Lang, 1936  
W Scotland, Bermudas, Malvinas
- Ancorabolinae* G.O. Sars, 1909
- Echinopsyllus* G.O. Sars, 1909
- E. normani* G.O. Sars, 1909  
Bohuslän-Bergen-Scotland
- Arthropsoyllus* G.O. Sars, 1909
- A. serratus* G.O. Sars, 1909  
Kieler Bucht-S Norway-Trondheim-White Sea
- Echinocletodes* Lang, 1936  
\* Should possibly be transferred to *Cletodidae*?
- E. armatus* (T. Scott, 1902)  
Faeroe Channel
- Ancorabolus* Norman, 1903
- A. mirabilis* Norman, 1903  
Bohuslän-N Norway-Scotland
- Ceratonotus* G.O. Sars, 1909
- C. pectinatus* G.O. Sars, 1909  
Bohuslän-Norway (Flekkerøy)

*C. pectinatus elaphus* Por,1965  
Bergen

*Dorsiceratus* Drzycimski,1967

*D. octocornis* Drzycimski,1967  
Bergen

SUBORDO incertae sedis

Latremidae Bozic,1969

\* Dubious family. Should be named Delamarellidae if it shows to be a good taxon.

*Delamarella* Chappuis,1954  
= *Latiremus* Bozic,1969

*D. arenicola* Chappuis,1954  
W France

*D. eximia* (Bozic,1969)  
?France

GELYELLOIDA Huys,1988

Gelyellidae Rouch & Lescher-Moutoué,1977

\* Only stygobiontic species from European karstic systems.

REMIPEDIA Yager,1981

\* Atlantic marine troglobionts with one recent order, Nectiopoda Schram,1986, not yet found close to Scandinavia, and one extinct order Enantiopoda Birshtein,1960

NECTIPODA Schram,1986

PHYLLOPODA Latreille,1825

PHYLLOCARIDA Packard,1879

LEPTOSTRACA Claus,1880

\* The only recent order. 4 extinct orders.

Nebaliidae Baird,1850

*Nebalia* Leach,1814

*N. borealis* Dahl,1985  
Bohuslän-Troms & Shetlands

*N. bipes* (O. Fabricius,1780)  
Trondheimsfjord-N Norway-Spitsbergen-Iceland & E Greenland

*N. herbstii* Leach,1814  
Shetlands-N Spain

*N. strausi* Risso,1826  
Engl. Channel-Adriatic Sea

*Nebaliella* Thiele,1904

*N. caboti* Clark,1932  
off W Ireland-Rockall Trough

*Sarsinebalia* Dahl,1985

*S. typhlops* (G.O. Sars,1870)  
Lofoten-Bergen area-European Atlantic-W Mediterranean

Nebaliopsidae Hessler,1984

*Nebaliopsis* G.O. Sars,1887

*N. typica* G.O. Sars,1887  
off W Ireland-Rockall Trough-southwards to 50° S

CALMANOSTRACA Tasch,1969

NOTOSTRACA G.O. Sars,1867

\* Three Scandinavian (non marine) species in smaller, sometimes temporary, water systems.

Triop(s)idae Kielhack,1910

\* Only limnic species.

SARSOSTRACA Tasch,1969

ANOSTRACA H. Milne-Edwards,1840

\* Five Scandinavian (non marine) species in temporary land-locked waters.

Chirocephalidae Daday,1910

\* Only limnic species.

Branchiopodidae Burmeister,1843 (Simon,1886)

\* Only limnic species.

Artemiidae Grochowski,1896

\* The well-known genus *Artemia* from e.g. Mediterranean Salines belongs to this group.

Branchinectidae Daday,1910

Polyartemiidae Simon,1886

Streptocephalidae Daday,1910

Thamnocephalidae Simon,1886

DIPOSTRACA Gerstaecker,1866

CONCHOSTRACA G.O. Sars,1867

\* Two Scandinavian (non marine) species in smaller, sometimes temporary, water systems.

Cyclestheriidae G.O. Sars,1899

Cyzicidae Stebbing,1910

Leptestheriidae Daday,1923

Limmadiidae Baird,1849

\* Only limnic species.

LAEVICAUDATA Linder,1945

Lynceidae Stebbing,1910

\* Only limnic species.

CLADOCERA Latreille,1829

\* Most species limnic. Some more limnic species, than the few treated here, may be found in inner Baltic archipelagos.

EUCLADOCERA Eriksson,1934

Sidoidea Baird,1850 (Brooks,1959)

= Ctenopoda G.O. Sars,1865

Sididae Baird,1850

*Penilia* Dana,1852

*P. avirostris* Dana,1852

Last week of September and first weeks of October 1997 this species was very common in N Bohuslän, but its northernmost records else is the S Nort Sea (German Bight & off Rotterdam)

Holopedidae G.O. Sars,1865

\* Only limnic species in the area.

Chydoroidea Stebbing,1902 (Brooks,1959)

= Daphnoidea Straus,1820

= Anomopoda G.O. Sars,1865

Bosminidae Baird,1845

*Bosmina* Baird,1845 (™ *B. coregoni* Baird,1857)

*B. (Eubosmina) longispina* Leydig,1860 *maritima* P.E. Müller,1867  
Gulf of Bothnia-Öresund-Kattegat (around Laesoe)-S. Skagerrak (rare)

Moinidae Goulden,1968

\* Only limnic species in the area.

Daphniidae Straus,1820

*Daphnia* O.F. Müller,1785

*D. (Ctenodaphnia) dybowskii* & *Grochowski*,1895 *magna* Strauss,1820 (™ of *Ctenodaphnia* D. & G.,1895

Often in fresh-water rock-pools above the marine littoral, in Denm., Finl., Swe. & Norw.

*D. (Daphnia) longispina* O.F. Müller,1785

Fresh-brackish waters in Denm., Finl., Swe., Norw. & Icel.

Macrothricidae Norman & Brady,1867

Ilyocryptinae Smirnov,1976

\* Only limnic species in the area.

Macrothricinae Norman & Brady,1867

\* Only limnic species in the area.

Chydoridae Stebbing,1902

Eurycercinae Kurz,1875

\* Only limnic species in the area.

Aloninae Frey,1967

\* Only limnic species in the area.

Chydorinae Stebbing,1902

\* Only limnic species in the area.

Polyphemoidae Baird,1845 (Brooks,1959)

= Onychopoda G.O. Sars,1865

Polyphemidae Baird,1845

*Polyphemus* O.F. Müller,1776

*P. pediculus* (Linnaeus,1761)

= *Monoculus pediculus* Linnaeus,1761 (™)

= *Polyphemus oculus* O.F. Müller,1785

= *Monoculus polyphemus* Jurine,1820

= *Polyphemus stagnorum* Fischer,1848

= *Polyphemus kochii* Schoedler,1863

Fresh-brackish waters in Denm., Finl., Swe., Norw. & Icel.

Podonidae Mordukhai-Boltovskoi,1968

*Podon* Liljeborg,1853

*P. polyphemoides* Leuckart,1859

\* Mordukhai-Boltovskoi already in 1978 moved this species to the genus *Pleopsis* Dana,1853 and Gieskes,

W.W.C.,1971 also propose a removal of this species to *Pleopsis*.

Bothnian Sea-Bohuslän-North Sea-Norw. Sea-Br. Isles-Mediterranean & Black Sea

*P. leuckarti* G.O. Sars,1862

N Baltic-Bohuslän-North Sea-Norw. Sea-Barents Sea-Iceland-Br. Isles-Engl. Channel, Black Sea

*P. intermedius* Liljeborg,1853 (™)

Bothnian Sea-Bohuslän-North Sea-Norw. Sea-Iceland-Br. Isles-Mediterranean & Black Sea

*Evadne* Lovén,1836

*E. nordmanni* Lovén,1836 (™)

Bothnian Sea-Bohuslän-North Sea-Norw. Sea-Barents Sea-Br. Isles-Mediterranean & Black Sea

*E. spinifera* P.E. Müller,1868

S Baltic-Bohuslän-North Sea-60° N-40° S

*E. tergestina* Claus,1877

Central North Sea-southwards

Cercopagidae Mordukhai-Boltovskoi,1966

*Cercopagis* G.O. Sars,1897

*C. pengoi* (Ostroumov,1891)

\* Native in fresh and brackish water in the Ponto-Caspian area, but successful immigration into the Baltic was detected in 1992.

HAPLOPODA G.O. Sars,1865

Leptodoridae Liljeborg,1900

*Leptodora* Liljeborg,1861

***L. kindti*** (Focke,1845)

Fresh water in Denm., Finl., Swe. &amp; Norw. &amp; Baltic archipelagos

**CEPHALOCARIDA** Sanders,1955

\* No European species so far recognized in this taxon with it's only recent order Brachypoda Birshstein,1960

**OSTRACODA** Latreille,1806

\* Here considered to be a taxon separate from Maxillopoda, whithin which they sometimes are grouped

**OSTRACODA** Latreille,1806= **OSTRAPODA** H.E. Straus,1821**BRADORIIDA** Matthew,1902

\* Extinct taxon.

**LEPERDITICOPIDA** Scott,1961

\* Extinct taxon.

**PHOSPHATOCOPIIDA** K.J. Müller,1964

\* Extinct taxon

**PALAEOCOPA** Henningsmoen,1953**PALAEOCOPIIDA** Henningsmoen,1953**KIRKBYOCOPINA** Gründel,1969**Puncioidea** Hornibrook,1949**Punciidae** Hornibrook,1949**MYODOCOPA** G.O. Sars,1866**MYODOCOPIDA** G.O. Sars,1866**Cypridinoidea** Baird,1850**Cylindroleberididae** Müller,1906**Cylindroleberidinae** Müller,1906**Synasterope** Kornicker,1975 (™ *Synasterope implumis* Paulsen,1965)= *Asterope* : Auctt.= *Synasterope* Paulsen,1965 (nom. nud.)***S. abyssicola*** (G.O. Sars,1870)

\* Correct genus?

off Lofoten-off N- Norway

***S. norvegica*** (G.O. Sars,1869)\* Kornicker,1987 attributes this species to his genus *Priantoleberis* Kornicker,1974 Bohuslän (e.g. the Koster area)-W Norway-Trondheim-N North Sea-Shetlands-NE Scotland***Parasterope*** Kornicker,1975= *Parasterope* Paulsen,1965 (nom. nud.)***P. muelleri*** (Skogsberg,1920)= *Asterope muelleri* Skogsberg,1920 (™)= ? *Cypridina teres* Norman,1861

NE Atlantic-?Br. Isles-Mediterranean

***P. aberrata*** (Skogsberg,1920)

Strangford Lough (N Ireland)

***Cylindroleberis*** Brady,1867= ? *Asterope* Philippi,1840, non Müller & Trotschel,1840= *Asteropina* Poulsen,1967***C. mariae*** (Baird,1850)= *Cypridina mariae* Baird,1850

SW Br. Isles

***Bruniella*** Poulsen,1965 (™ *B. breviata* Poulsen,1965 - off Thailand)***Heptonema*** Cohen & Kornicker,1975 (™ *H. serratum* Poulsen,1965 - Virgin Island)***Diasterope*** Kornicker,1975 (™ *D. pilosa* Poulsen,1965 - ?)***Bathyleberis*** Kornicker,1975 (™ *B. grossmani* Kornicker,1975 - ?Antarctic)***Postasterope*** Kornicker,1986 (™ *P. abaco* Kornicker,1986 - Bahamas)***Archasterope*** Poulsen,1965 (™ *A. dentata* Poulsen,1965 - S Oceans)***Domromeus*** Kornicker,1989 (™ *D. heptathrix* Kornicker,1989 - Biscay)***Priantoleberis*** Kornicker,1974 (™ *P. gyion* Kornicker,1974 - off Mauretania)**Cyclasteropinae** Poulsen,1965

\* Not represented in the area

**Asteropteroinae** Kornicker,1981

\* Not represented in the area

**Sarsiellidae** G.S. Brady & A.M. Norman,1896***Sarsiella*** Norman,1869***S. capsula*** Norman,1869 (™)

off Shetlands

***Eusarsiella*** Cohen & Kornicker,1975***E. zostericola*** (Cushman,1906)

Br. Isles (introduced)

**Cypridinidae** Baird,1850**Cypridininae** Baird,1850**Cypridini** Baird,1850***Doloria*** Skogsberg,1920 (™ *Cypridina (Doloria) levis* Skogsberg,1920 - South Georgia)***D. sarsi*** Kornicker,1987

\* Furca with 11 claws.

Koster Channel in 180m

***Skogsbergia*** Poulsen,1962= *Cypridina* : Auctt., non H. Milne-Edwards,1840

\* Furca with 9 claws.

***S. megalops*** (G.O. Sars,1871)

off W Norway-Hardangerfjord

***Vargula*** Skogsberg,1920= *Cypridina* : Auctt., non H. Milne-Edwards,1840

\* Furca with 9 claws.

***V. norvegica*** (Baird,1860)= *Cypridina norvegica* Baird,1860 (™)

Koster Channel-W Skagerrak-N Norway-Shetlands-Faeroes-Iceland

***Macrocypridina*** Skogsberg,1920***M. castanea*** (Brady,1897)= *Cypridina castanea* Brady,1897 (™)

N Atlantic-56° N

**Gigantocypridini** Hartmann,1974***Gigantocypris*** G.W. Müller,1895 (™ *G. agassizii* G.W. Müller,1895)***G. muelleri*** Skogsberg,1920

N Atlantic-64° N

**Azygocypridinae** Kornicker,1970***Azygocypridina*** Sylvester-Bradley,1950 (™ *Crossophorus imperator* Brady,1880 (SW Pacific))***A. imperialis*** (Stebbing,1901)

E Atlantic between 21°N-60°N at 1884-2930m

**Philomedidae** G.W. Müller,1906**Philomedinae** G.W. Müller,1906***Philomedes*** Liljeborg,1853 (™ *P. longicornis* Liljeborg,1853)***P. brenda*** (Baird,1850)= *Cypridina brenda* Baird,1850= *P. globosus* (Liljeborg,1853)

Öresund-Bohuslän-North Sea-Barents Sea &amp; Spitsbergen-Iceland-Faeroes-Shetlands-Br. Isles

***P. liljeborgi*** (G.O. Sars,1866)= *P. bonneti* Kornicker & Caraiion,1977

Kattegatt-Bohuslän-Skagerrak-Trondheim-Lofoten-Bjornoja-Iceland-Faeroes, Biscay

***Euphilomedes*** Poulsen,1962***E. interpuncta*** (Baird,1850)= ? *E. sinister* Kornicker,1974?

E Atlantic 30-65° N

**Pseudophilomedinae** Kornicker,1967

\* Not recorded N of Biscay

**HALOCYPRIDA** Dana,1853**HALOCYPRIDINA** Dana,1853**Halocyprididae** Dana,1852**Conchoecinae*****Conchoecia*** Dana,1849***C. (Boroecia) Poulsen,1973 borealis*** G.O. Sars,1866

Koster area-Lofoten-Arctic Seas, Iceland

***C. (Discoconchoecia) Martens,1979 elegans*** G.O. Sars,1866

N Kattegatt-Bohuslän-Lofoten-Br. Isles-Atlantic-80° N, Iceland

***C. (Conchoecilla) Poulsen,1973 daphnoides*** (Claus,1890)

Atlantic-60° N

***C. (Orthoconchoecia) Poulsen,1973 haddoni*** Brady & Norman,1896

Atlantic-60° N

***C. (Conchoecissa) Poulsen,1973 imbricata*** Brady,1880

N Atlantic

***C. (Conchoecissa) ametra*** G.W. Müller,1906

Iceland

***C. (Obtusocia) Martens,1979 obtusata*** G.O. Sars,1866

Bohuslän-Trondheim-boreal-subarctic Atlantic

***C. (Metaconchoecia) Poulsen,1973 rotundata*** G.W. Müller,1891= *C. skogsbergi* Iles,1953\* Of this "species" 15 different forms are recognized by Angel & Fasham,1975, of which one, "*C skogsbergi*": Angel,1968, non Iles,1953, is referred to here.

Norw. Sea

***C. (Conchoecia) lophura*** G.W. Müller,1906

Iceland

**Euconchoecinae**\* The other genus of the subfamily, *Euconchoecia* Müller,1890 is not known from the area.***Bathyconchoecia*** Deevey,1968***B. darcythompsoni*** (Scott,1909)= *Euconchoecia darcythompsoni* Scott,1909

NW Scotland

**Halocypridinae** Dana,1852\* The third genus of the subfamily, *Fellia* is not known from the area.***Halocypria*** Claus,1874***H. globosa*** Claus,1874

Atlantic-62° N

***Halocypris*** Dana,1852***H. inflata*** Dana,1852= *H. brevisiridis* Dana,18??

Atlantic, occasionally N to S &amp; W Britain

**CLADOCOPIDINA** G.O. Sars,1866

- Polycopoidea G.O. Sars,1866
- Polycopidae G.O. Sars,1866
- Polycope** G.O. Sars,1866
- P. areolata* G.O. Sars,1923  
Bohuslän-Korshavn (S Norway)-Plymouth
- P. clathrata* G.O. Sars,1923  
off Väderöarna & Grisbådarna-Korshavn (S Norway)-Trondheim
- P. dentata* Brady,1868  
?North Sea
- P. difficilis* Klie,1936  
Helgoland
- P. elongata* Hartmann,1954  
Sylt
- P. helgolandica* Klie,1936  
Helgoland
- P. orbicularis* G.O. Sars,1866  
Öresund-Skagerrak-Trondheim-SW Br. Isles
- P. punctata* G.O. Sars,1870  
off Väderöarna-off Lofoten-Tromsø, ?Br. Isles
- P. pustulata* G.O. Sars,1890  
W Norway
- P. schulzi* Klie,1950  
Kieler Bucht
- P. sublaevis* G.O. Sars,1923  
Bohuslän-Risör-Korshavn (S Norway)-Trondheim
- Parapolycope** Klie,1936
- P. germanica* Klie,1936  
Helgoland
- Polycopsis** G.W. Müller,1894
- P. compressa* (Brady & Robertson,1869)  
Korshavn (S Norway)-SW Br. Isles-Mediterranean
- PODOCOPA** G.W. Müller,1894
- PLATYCOPIA G.O. Sars,1866
- Cytherelloidea G.O. Sars,1866
- Cytherellidae G.O. Sars,1866
- Cytherella** Jones,1850
- C. abyssorum* G.O. Sars,1866  
Kattegatt-Bohuslän-Trondheim-Lofoten-Iceland-Br. Isles-?Mediterranean
- C. serrulata* Brady & Norman,1896  
Britain
- PODOCOPIDA G.O. Sars,1866
- METACOPINA** Sylvester-Bradley,1961
- Darwinuloididea  
= Darwinulacea Brady & Norman,1889
- Darwinulidae Brady & Norman,1889
- Darwinula** Brady & Norman,1889
- D. stevensoni* (Brady & Robertson,1870)  
Tvärminne area, Stadil Fjord (oligohaline lagoon in W Jutland)
- Sigillioidea  
= Sigillidacea Mandelstam,1960  
\* No European species.
- Saipanettidae McKenzie,1968
- PODOCOPINA G.O. Sars,1866
- Bairdioidea  
= Bairdiacea G.O. Sars,1888
- Bairdiidae G.O. Sars,1888
- Bairdiinae G.O. Sars,1888
- Bairdia** McCoy,1844
- Neonesidea** Maddocks,1969
- N. inflata* (Norman,1862)  
= *Cythere inflata* Norman,1862  
= *Bairdia inflata* (Norman,1862)  
W Norway-N Norway-Br. Isles-NW France
- N. acanthigera* (Brady,1868)  
= *Bairdia acanthigera* Brady,1868  
Engl. Channel
- Bythocyprididae Maddocks,1969
- Bythocypris** Brady,1880
- B. bosquetiana* (Brady,1866)  
W Norway-Trondheim-Shetlands-SW Br. Isles-Mediterranean
- B. obtusata* (G.O. Sars,1866)  
S Norway (shells)-Bergen-Br. Isles
- Anchistrocheles** Brady & Norman,1889
- A. acerosa* (Brady,1868)
- = *Cythere (?) acerosa* Brady,1868  
Plymouth
- Cytheroidea Baird,1850  
= Cytheracea Baird,1850
- Limnocytheridae Klie,1938
- Limnocythere** Brady,1868
- L. inopinata* (Baird,1843)  
Baltic coasts
- Krithidae Mandelstam,1958
- Kritha** Brady, Crosskey & Robertson,1874
- K. angusta* Brady & Norman,1889?  
Skagerrak-North Sea
- K. praetexta* (G.O. Sars,1866)  
= *K. bartonensis* : Auctt., non (Jones,1857)  
= *Ilyobates praetexta* G.O. Sars,1866 (TM)  
Öresund-Bohuslän-Oslofjord-Lofoten-N Norway-Iceland-Shetlands-NE Britain
- K. producta* Brady,1880  
Spitsbergen-?
- Paracyprideis** Klie,1929
- P. fennica* (Hirschmann,1909)  
Gävle & Tvärminne-S Baltic, Arctic Seas
- Cytherideidae G.O. Sars,1925
- Cytheridea** Bosquet,1852
- C. angustata* (Baird,1850)  
Öresund-Bohuslän-Helgoland-Holland-Br. Isles-?Iceland-Bay of Biscay-?Mediterranean
- Cyprideis** Jones,1857  
= *Anomocytheridea* Stephenson,1938
- C. torosa* (Jones,1850)  
= *C. torosa teres* Brady & Robertson,1870  
= *C. litoralis* (Brady,1868)  
= *Cytheridea pedaschenkoi* von Daday,1909  
Tvärminne-Öresund-Bohuslän-Oslofjord estuaries-Holland-E Br. Isles-France-Mediterranean
- Sarsicytheridea** Athersuch,1982  
= *Cytheridea sensu* Sars,1925, non Bosquet,1851  
= *Cytheridea (Eucytheridea)* Bronstein,1930  
=? *Cyamocytheridea* Oertli,1956
- S. bradii* (Norman,1865)  
= *Cythere bradii* Norman,1865 (TM)  
= *Cyprideis bradii* G.O. Sars,1866  
= *Cytheridea papillosa* : Auctt., non Bosquet,1852  
S Baltic-Bohuslän-North Sea-N Norway-Iceland & Spitzbergen-Novaja Zemlja-Faeroes-N Br. Isles
- S. punctillata* (Brady,1865)  
= *Cyprideis proxima* G.O. Sars,1866  
Ålands Sea-Öres.-Bohuslän-North Sea-N Norway-Iceland & Arctic Seas-?Faeroes-N Br. Isles
- Neocytherideidae Puri,1957
- Neocytherideis** Puri,1952  
= *Sahnia* Puri,1952 (p.p.)
- N. crenulata* (Klie,1929)  
\* Is this identical with *N. subulata* var. *crenulata*?  
S Baltic-Belt Sea & Öresund, North Sea
- ? *N. baltica* (Klie,1938)  
S Baltic
- N. subulata* (Brady,1868)  
= *Cytherideis subulata* var. *fasciata* Brady & Robertson,1874  
= *Cytherideis subulata* var. *crenulata* Brady,1911  
= *N. elongatus* Puri,1952 (TM)  
SW Br. Isles-W Scotland, W France
- Sahnicythere* Athersuch,1982  
= *Sahnia* Puri,1952 (p.p.)
- S. retroflexa* (Klie,1936)  
= *Cytherideis retroflexa* Klie,1936 (TM)  
Helgoland-Amrum, Britain
- Heterocyprideis** Elofson,1941
- H. sorbyana* (Jones,1856)  
Gävle & Tvärminne-S Baltic-Bohuslän (shells)-W Norway (shells)-N Norway-Spitsbergen-Scotland (shells)-Ireland (shells)
- Cushmanideidae Puri,1974
- Pontocythere** Dubowsky,1939 (TM *P. tschernjanskii* Dubowski,1939)  
= *Hemicytherideis* Ruggieri,1952  
= *Cytherideis sensu* Puri,1952  
= *Hulungia* Puri,1958
- P. elongata* (Brady,1868)  
Skagerrak, SW Br. Isles-Thames estuary, N France
- Trachyleberididae Sylvester-Bradley,1948
- Buntonia** Howe & Chambers,1935  
= *Semicythereis* Elofson,1943
- B. corpulenta* (Brady & Norman,1889)  
Bergen-Trondheim
- Acanthocythereis** Howe,1963 (TM *A. arenosa* Howe,1963)  
= *Trachyleberis* : Auctt., non Brady,1898
- A. dunelmensis* (Norman,1865)  
S Baltic-Öres.-Bohuslän-N Norway-Iceland & Arctic Seas-Faeroes-Shetlands-Br. Isles, E Greenland
- A. horrida* (G.O. Sars,1866)  
\* Elofsons (1941) larger form of *Cythereis dunelmensis* is probably this species sensu Athersuch & al.,1989.  
?Skagerrak

- Pterygocythereis* Blake, 1933
- P. jonesii* (Baird, 1850)  
= *Cythereis jonesii* Baird, 1850 (TM)  
Öresund-Bohuslän-SE North Sea-S & W Norway-Shetlands-Br. Isles-Bay of Biscay-Mediterranean-Black Sea
- P. mucronata* (G.O. Sars, 1866)  
Trondheim-off Lofoten-Shetlands
- Echinocythereis* Puri, 1954
- E. echinata* (G.O. Sars, 1866)  
Kattegat-Koster Channel-Oslofjord-Lofoten-Atlantic
- E. mirabilis* (Brady, 1868)  
Bohuslän (shells)-Scotland (shells)-N Norway-Arctic Seas
- Cythereis* Jones, 1849
- C. leioderma* (Norman, 1869)  
Bohuslän (shells)-W Norway-Shetlands-Iceland-Arctic Seas
- Carinocythereis* Ruggieri, 1956
- C. carinata* (Roemer, 1838)  
= *Cytherina carinata* Roemer, 1838 (TM)  
= *Cythereis antiquata* Baird, 1850  
W & SW Br. Isles, W France, Mediterranean
- C. whitei* (Baird, 1850)  
= *Cythereis aspera* Brady, 1865  
= *Carinocythereis bairdii* Uliczny, 1969  
S Britain, W France, Mediterranean
- Robertsonites* Swain, 1963
- R. tuberculata* (G.O. Sars, 1866)  
= *Cythereis tuberculata* G.O. Sars, 1866  
= *Robertsonites gubikensis* Swain, 1963 (TM)  
= *R. tuberculatina* Swain, 1963  
SW Baltic-Öres.-Bohuslän-N Norway-Spitsbergen-Iceland-Faeroes-Scotland & E England-Thames estuary
- Celtia* Neale, 1973
- C. quadridentata* (Baird, 1850)  
= *Cythere quadridentata* Baird, 1850 (TM)  
Risør (S Norway)-Hardangerfjord-Shetlands-Br. Isles-Bay of Biscay
- Costa* Neviani, 1928 (TM *Cytherina edwardsii* Roemer, 1838)
- C. runcinata* (Baird, 1850)  
S Britain, W France, W. Spain, Mediterranean-Turkey
- Hiltermannicythere* Bassiouni, 1970 (TM *Cythereis turbida* G.W. Müller, 1894)  
= *Falunia* Grekoff & Moyes, 1956 (p.p.)  
= *Falunia rugosa* group Sissingh, 1971
- H. emaciata* (Brady, 1868)  
SW Br. Isles, Cardigan Bay, Ireland
- Basslerites* Howe, 1937
- B. berchoni* (Brady, 1869)  
Arcachon-Mediterranean
- Eucytheridae Puri, 1954
- Eucythere* Brady, 1868  
= *Cytheropsis* G.O. Sars, 1866, non M'Coy, 1849
- E. argus* (G.O. Sars, 1866)  
= *E. undulata* Klie, 1929  
Kieler Bucht-Öres.-Bohuslän-Oslofjord-W & N Norway-Shetlands-SW Br. Isles-Ireland
- E. declivis* (Norman, 1865)  
= *Cythere declivis* Norman, 1865 (TM)  
=? *Cytheropsis tenuitesta* G.O. Sars, 1866  
Belt Sea-Oslofjord-N Norway-Arctic Seas-SW Br. Isles
- E. anglica* Brady, 1868  
NE England
- E. prava* Brady & Robertson, 1869  
W Ireland, S England
- Paracytherideidae Puri, 1957
- Paracytheridea* G.W. Müller, 1894 (TM *P. depressa* G.W. Müller, 1894)  
= *Xenocythere* G.O. Sars, 1925 (TM *Cythere cuneiformis* Brady, 1868)
- P. cuneiformis* (Brady, 1868)  
Öresund & Belt Sea-Bohuslän-Bergen-Scotland-Ireland-Channel Islands
- Cytheridae Baird, 1850
- Cythere* O.F. Müller, 1785  
= *Cytherina* Lamarck, 1818 (TM *Cythere lutea* O.F. Müller, 1785)
- C. lutea* O.F. Müller, 1785 (TM)  
= *C. reniformis* Baird, 1835  
Kieler Bucht-Öres.-Bohuslän-Bergen-N Norway-Iceland & Arctic Seas-Shetlands-SW Br. Isles, W France
- C. globulifera* Brady, 1868  
\* Not mentioned in S.B.F. (N.S.) 43, 1989. May be a synonym of another species?  
W Scotland, Roundstone
- C. acuta* Baird, 1850  
\* Not mentioned in S.B.F. (N.S.) 43, 1989. May be a synonym of another species?  
Arran
- Palmenella* Hirschmann, 1916  
= *Kyphocythere* G.O. Sars, 1925 (TM *Cythere limicola* Norman, 1865)
- P. limicola* (Norman, 1865)  
= *Cythere limicola* Norman, 1865 (TM)  
= *Cythere nodosa* G.O. Sars, 1866  
= *Cythere areolata* Brady, 1866  
S Baltic-Öres.-Bohuslän-Lofoten-Arctic Seas-N Br. Isles
- Cuneocytheridae Mandelstam, 1959
- Cuneocythere* Lienenklaus, 1894 (TM *Cuneocythere truncata* Lienenklaus, 1894)
- C. semipunctata* (Brady, 1868)  
SW Br. Isles, Ireland, SW France
- Microcytheridae Klie, 1938
- Microcythere* G.W. Müller, 1894
- M. bahusiensis* Elofson, 1944  
Bohuslän
- M. helgolandica* Klie, 1936  
Helgoland
- M. minuta* Klie, 1936  
Helgoland
- M. monstrosa* Elofson, 1944  
Bohuslän
- M. navicula* (Norman, 1869)  
Shetlands-SW Br. Isles
- M. parva* Klie, 1936  
Helgoland
- M. producta* Elofson, 1944  
Bohuslän
- M. schyttei* Hartmann, 1957  
Sylt
- Psammocytheridae Klie, 1938
- Psammocythere* Klie, 1936
- P. remanei* Klie, 1936  
Helgoland
- Cobanocytheridae Schornikov, 1975
- Cobanocythere* Hartmann, 1959
- C. angularis* Gottwald & Scherner, 1980  
S? North Sea
- C. syltensis* Gottwald, 1983  
Sylt
- Leptocytheridae Hanai, 1957
- Leptocythere* G.O. Sars, 1925
- L. baltica* Klie, 1929  
Tvärminne-Öresund-Bohuslän-Sylt, Britain-Biscay
- L. castanea* (G.O. Sars, 1866)  
Dalarö & Tvärminne-Öresund-Bohuslän-Oslofjord-North Sea-W & N Norway-SW Br. Isles-Biscay, S Greenland
- L. confusa* (Brady & Norman, 1889)  
S Baltic
- L. lacertosa* (Hirschmann, 1912)  
Ålands Sea & Tvärminne-Öres.-Bohuslän-Holland, Britain, Biscay, Iceland, White Sea
- L. macallana* (Brady & Robertson, 1869)  
= *Cythere levis* G.W. Müller, 1894  
Oslofjord-Skagerrak-Hardangerfjord-Holland-Scotland-W Ireland-SW Br. Isles-Mediterranean
- L. major* Hartmann, 19??  
Öresund, German North Sea Coast
- L. pellucida* (Baird, 1850)  
= *Cythere pellucida* Baird, 1850 (TM)  
= *Cythere confusa* Brady & Norman, 1889  
Ålands Sea & Tvärminne-Öres.-Bohuslän-Oslofjord-North Sea-N Norway-Shetlands-SW Br. Isles-Bay of Biscay-Mediterranean
- L. porcellanea* (Brady, 1869)  
= *L. llyophila* (Hirschmann, 1909)  
Dalarö & Tvärminne-S Baltic, SW Br. Isles, Biscay
- L. tenera* (Brady, 1868)  
S Baltic-Oslofjord-Risør-Hardangerfjord-Holland-Br. Isles-Bay of Biscay
- L. psammophila* Guillaume, 1976  
Baltic, Iceland, Britain, Roscoff, W France
- Callistocythere* Ruggieri, 1953
- C. badia* (Norman, 1861?)  
SW Br. Isles, Cyprus
- C. littoralis* (G.W. Müller, 1894)  
= *Cythere littoralis* G.W. Müller, 1894 (TM)  
= *C. crispata* : Auctt., non (Brady, 1868) (Bradys species is restricted to the Mediterranean)  
Bohuslän-Oslofjord-S & W Norway-Ireland-Br. Isles-Mediterranean
- C. murrayi* Whittaker, 1978  
Wales, S England, Norfolk
- Hemicytheridae Puri, 1953
- Heterocythereis* Elofson, 1941
- H. albomaculata* (Baird, 1838)  
=? *Cythere alba* Baird, 1835 (nom. dub.)  
= *Cythere albo-maculata* Baird, 1838 (TM)  
Bohuslän-Helgoland-Hardangerfjord-N Norway-Iceland-Shetlands-SW Br. Isles-Bay of Biscay-Mediterranean, Azores
- Urocycythereis* Ruggieri, 1950 (TM *Cytherina favosa* Roemer, 1838)
- U. britannica* Athersuch, 1977  
= *Cythere oblonga* Brady, 1868, non Brady, 1866, nec M'Coy, 1844  
\* The Mediterranean *Cythere oblonga* Brady, 1866 is a junior objective homonym of M'Coy's species and was renamed *U. distinguenda* (Neviani, 1928).  
Öresund-Skagerrak-S Norway-Engl. Channel-Wales-Bay of Biscay-N Spain

- Elofsonella** Pokorny,1955  
= *Paracythereis* Elofson,1941, non Delachaux,1928
- E. concinna** (Jones,1857)  
= *Cythere concinna* Jones,1857 (TM)  
= *Cythereis clavata* G.O. Sars,1866  
Kieler Bucht-Öresund-Bohuslän-Bergen-Spitsbergen-Iceland-Faeroes-Shetlands-N Br. Isles
- Muellerina** Bassiouni,1965 (TM *Cythere latimarginata* Speyer,1863)
- M. abyssicola** (G.O. Sars,1866)  
= *Hemicythere latimarginata* : G.O. Sars,1925, non (Speyer,1863)  
Bohuslän-Bergen-Lofoten-Spitsbergen-Iceland-Shetland area-N Britain
- Aurila** Pokorny,1955  
= *Auris* Neviani,1928 (p.p.)  
= *Eucythereis* Klie,1940 (p.p.)  
=? *Mutilus* Neviani,1930
- A. convexa** (Baird,1850)  
= *Cythere convexa* Baird,1850 (TM)  
Bohuslän-Hardangerfjord-Shetlands-Br. Isles-Mediterranean
- A. arborescens** (Brady,1865)  
= *Cythere woodwardii* Brady,1868  
Wales-Mediterranean
- A. woutersi** Home,1986  
S Britain
- Hemicythere** G.O. Sars,1925  
= *Auris* Neviani,1928 (p.p.)  
= *Eucythereis* Klie,1940 (p.p.)
- H. costata** (Brady,1866)  
Shetlands-Spitsbergen
- H. crenulata** (G.O. Sars,1866)  
Koster Channel-S & W Norway-Shetlands-?Faeroes-Arctic Seas-Br. Isles
- H. emarginata** (G.O. Sars,1866)  
Bohuslän-Bergen-Lofoten-N Norway-Arctic Seas-Iceland-Shetland-W Scotland (empty shells in Britain)
- H. villosa** (G.O. Sars,1866)  
= *Cythereis villosa* G.O. Sars,1866 (TM)  
Kieler Bucht-Öres.-Bohuslän-North Sea-N Norway-Iceland-Bear Island-Br. Isles-Bay of Biscay
- H. rubida** (Brady,1868)  
W Scotland, S England, Ireland
- Procythereis** Skogsberg,1928
- ?**P. marginata** (Norman,1861?)  
SW Br. Isles
- Finmarchinella** Swain,1963
- F. (Finmarchinella) finmarchica** (G.O. Sars,1866)  
Korshavn (S Norway)-N Norway-S North Sea-Br. Isles (empty shells in Britain)
- F. (Barentsova) angulata** (G.O. Sars,1866)  
S Baltic-Öres.-Bohuslän-Oslofjord-N Norway-Iceland-Arctic Seas-Ireland-Br. Isles (empty shells in Britain)
- Baffinicythere** ?,19??
- B. howei** Hazel,1???  
Britain (empty shells in Britain)
- Cythereis** Jones,1849
- C. navicula** (Norman,18??)  
= *Cythere navicula* Norman,18??  
= *Cythere fiducula* Brady & Robertson,1889  
\* Not mentioned in S.B.F. (N.S.), No. 43, 1989. May possibly be a synonym of another species?  
Bergen, Shetlands, Northumberland, Firth of Forth, Maas & Schelde estuaries
- Xestoleberididae** G.O. Sars,1928
- Xestoleberis** G.O. Sars,1866
- X. nitida** (Liljeborg,1853)  
= *Cythere nitida* Liljeborg,1853 (TM)  
= *X. aurantia* : G.O. Sars,1928, non (Baird,1838)  
SW Finland & Gräsö-Öres.-Bohuslän-North Sea-Bergen-N Norway-Shetlands-SW Br. Isles-Arcachon
- X. aurantia** (Baird,1838)  
= *X. pusilla* Elofson,1941  
Bohuslän-Bergen, Britain, Bretagne
- X. depressa** G.O. Sars,1866  
S Baltic-Öres.-Bohuslän-Bergen-Lofoten-Arctic Seas-Shetlands-Br. Isles-Bay of Biscay-?Mediterranean
- X. labiata** Brady & Robertson,1874  
SW Br. Isles, N France
- X. rubens** Whittaker,1978  
Dorset, W France
- Microxestoleberis** G.W. Müller,1894
- M. nana** G.W. Müller,1894  
Helgoland-Mediterranean
- M. rostrata** Klie,1936  
Helgoland
- Loxoconchidae** G.O. Sars,1925
- Hirschmannia** Elofson,1941
- H. viridis** (O.F. Müller,1785)  
= *Cythere viridis* O.F. Müller,1785 (TM)  
= *Cythere cyamos* Norman,1905  
= *Loxoconcha sarsii* Hirschmann,1912  
non = *Cythere viridis* : Liljeborg,1853 (which is = *Loxoconcha elliptica* Brady,1868)  
Tvärminne-Öresund-Bohuslän-Helgoland-Bergen-Murmansk-SW Br. Isles-SW France
- Cytheromorpha** Hirschmann,1909
- C. claviformis** Hirschmann,1909?  
?S Baltic?
- C. fuscata** (Brady,1869)  
Dalarö & Tvärminne-Öresund, Stadil Fjord (oligohaline lagoon in W Jutland), Drammen, Holland river mouths, Thames estuary (now only empty shells in Britain) & Caspian Sea
- C. n.sp. A. Rosenfeld,1977  
Belt Sea
- Bonnyannella** Athersuch,1982
- B. robertsoni** (Brady,1868)  
= *Cythere robertsoni* Brady,1868 (TM)  
Belt Sea-Oslofjord-SW Br. Isles-W France
- Elofsonia** Wagner,1957
- E. baltica** (Hirschmann,1909)  
= *Loxoconcha baltica* Hirschmann,1909 (TM)  
= *Cythere gibbosa* Brady & Robertson,1869, non H. Milne-Edwards,1840  
Dalarö & Tvärminne-Öres.-Holland, N Norway, Britain-W Ireland-Arcachon
- E. pusilla** (Brady & Robertson,1870)  
= *Loxoconcha pusilla* Brady & Robertson,1870  
Tvärminne-Öresund-Bohuslän-Sylt-Holland-Br. Isles
- Palmococoncha** Swain & Gilby,1974 (TM *P. laevimarginata* Swain & Gilby,1974)  
= *Lindisfarnia* Horne & Kilenyi,1981 (TM *Cythere laevata* Norman,1865)
- P. guttata** (Norman,1865)  
= *Cythere impressa* Baird,1850, non M'Coy,1844  
= *Cythere guttata* Norman,1865  
= *Loxoconcha granulata* G.O. Sars,1866  
Kieler Bucht-Öres.-Bohuslän-N Norway-Br. Isles-Ireland
- P. laevata** (Norman,1865)  
= *Cythere laevata* Norman,1865  
= *Loxoconcha longipes* G.O. Sars,1866  
= *Loxoconcha tamarindus* : Aucutt., non (Jones,1856)  
S Baltic-Öres.-Bohuslän-N Norway-Iceland-Shetlands-Br. Isles & Ireland-Bay of Biscay
- Loxoconcha** G.O. Sars,1866  
= *Normania* Brady,1866 (published 3 months after Sars' name)
- L. emelwardensis** Redeke,1936  
S North Sea
- L. elliptica** Brady,1868  
= *L. gauthieri* Klie,1929  
Dalarö & Finland-Öres.-North Sea-Br. Isles & ?Ireland, all around Mediterranean
- L. helgolandica** Klie,1929  
Helgoland
- L. rhomboidea** (Fischer,1855)  
= *L. impressa* : Brady,1868, non *Cythere impressa* Baird,1850, non M'Coy,1844  
= *Cythere rhomboidea* Fischer,1855 (TM)  
= *Cythere (Normania) carinata* Brady,1865  
S Baltic-Öres.-Bohuslän-North Sea-N Norway-Shetlands-Br. Isles-W France-Mediterranean-Madeira & Canaries
- Sagmatocythere** Athersuch,1976 (TM *Loxoconcha napoliana* Puri,1963)
- S. multiflora** (Norman,1866)  
= *Loxoconcha multiflora* : Elofsson,1938  
Koster Channel-Holland-Trondheim-Shetlands-around Br. Isles-Bay of Biscay
- Phlyctocythere** Keij,1958
- P. fragilis** (G.O. Sars,1866)  
= *Loxoconcha f.* : Aucutt.  
Kieler Bucht-Belt Sea-Bohuslän-Oslofjord-Lofoten-Br. Isles-W Ireland
- Nannocythere** Schäfer,1953
- N. pavo** (Malcolmson,1886)  
= *N. remane* Schäfer,1953 (TM)  
Helgoland, Engl. Channel, E Ireland, W Scotland
- Taberoloxoconcha** Hartmann,1973  
\*The British records may possibly be an undescribed species of the genus.
- T. tuberosa** (Hartmann,1954)  
?Britain
- Bythocytheridae** G.O. Sars,1866
- Pseudocythere** G.O. Sars,1866
- P. caudata** G.O. Sars,1866 (TM)  
Bohuslän-Oslofjord-Trondheim-N Norway-Arctic Seas-Shetlands-Br. Isles-?Bay of Biscay
- P. britannica** Horne,1986  
NE England, Isles of Scilly
- P. norvegica** Horne,1986  
Norway
- Jonesia** Brady,1866  
= *Macrocythere* G.O. Sars,1926
- J. acuminata** (G.O. Sars,1866)  
= *Cythere simplex* Norman,1865 (TM), non *Cythere auriculata* var. *simplex* Cornuel,1848  
S Baltic-Öres.-Bohuslän-North Sea-Lofoten-Troms-Arctic Seas-Iceland-Faeroes-Shetlands-Br. Isles, W Greenland
- Bythocythere** G.O. Sars,1866
- B. bradyi** G.O. Sars,1926  
Stavanger-SW Br. Isles
- B. constricta** G.O. Sars,1866  
Oslofjord-Skagerrak-W & N Norway-Spitsbergen
- B. dromedaria** G.O. Sars,1866  
Koster Channel-Oslofjord-Trondheim
- B. insignis** G.O. Sars,1870  
off Lofoten
- B. intermedia** Elofson,1938  
Bohuslän-S Norway, S North Sea, Britain-Biscay
- B. turgida** G.O. Sars,1866 (TM)

Koster Channel-Oslofjord-S & W Norway-Spitsbergen-?Shetlands-Ireland-?Bay of Biscay-?Mediterranean

**B. bradleyi** Athersuch, Horne & Whittaker, 1983  
= *B. turgida* : Sylvester-Bradley, 1947, non G.O. Sars, 1866  
SW Ireland

**B. robinsoni** Athersuch, Horne & Whittaker, 1983  
= *B. turgida* : Brady, 1868 & 1870, non G.O. Sars, 1866  
Britain

**B. zetlandica** Athersuch, Horne & Whittaker, 1983  
= *B. constricta* : Brady, 1868 (p.p.), non G.O. Sars, 1866  
Shetlands, N Britain

**Scleerochilus** G.O. Sars, 1866

**S. contortus** (Norman, 1861)  
= *Cythere contorta* Norman, 1861 (TM)  
Cardigan Bay, Engl. Channel

**S. abbreviatus** Brady & Robertson, 1869  
W Ireland, W Britain

**S. bradyi** Rudjakov, 1962  
NE England, Shetlands, Clyde Sea

**S. rudjakovi** Athersuch & Horne, 1987  
S Baltic-Öres.-Bohuslän-S North Sea-White Sea, Shetlands, Britain

**S. gewemuelleri** Dubowsky, 1939  
NE & S England, Wales, Mediterranean, Black Sea

**S. hicksi** Athersuch & Horne, 1987  
NE England

**S. schornikovi** Athersuch & Horne, 1987  
W Scotland, Bristol Channel

**S. truncatus** (Malcolmson, 1886)  
= *Paradoxostoma truncatum* Malcolmson, 1886  
NE England, N Wales

**S. whatleyi** Athersuch & Horne, 1987  
W Scotland

**S. n.sp.** : Athersuch & al., 1989  
= *S. contortus* : G.O. Sars, 1926, non (Norman, 1861)  
Norway-Spitzbergen

**Paradoxostomatidae** Brady & Norman, 1889

**Cytherois** G.W. Müller, 1884

**C. arenicola** Klie, 1929  
Tvärminne-Kattegatt-S North Sea, White Sea

**C. fischeri** (G.O. Sars, 1866)  
= *Scleerochilus (?) gracilis* Brady & Robertson, 1869  
= *Cytherois virens* G.W. Müller, 1884 (TM)  
?S Finland-S Baltic-Öres.-Bohuslän-S North Sea-S & ?W Norway-Br. Isles-W France-Mediterranean

**C. pusilla** G.O. Sars, 1928  
S Baltic-Öresund-Bohuslän-SW Norway-North Sea-Bergen-Britain-NW France

**C. vitrea** (G.O. Sars, 1866)  
Kieler Bucht-Öresund-Bohuslän-Oslofjord-North Sea-Lofoten-Shetlands-Faeroes-SW Br. Isles

**C. stephanidesi** Klie, 1938  
Britain (Cristchurch Harbour)-SW France-Mediterranean

**C. n.sp.** : Athersuch & al., 1989  
Britain

**Paradoxostoma** Fischer, 1855 (TM *P. dispar* Fischer, 1855)  
= *Xiphichilus* Brady, 1870 (TM *Bythocythere tenuissimum* Norman, 1869)

**P. abbreviatus** G.O. Sars, 1866  
S Baltic-Öres.-Bohuslän-Holland-Bergen-Shetlands-Br. Isles-N France-?Bay of Biscay

**P. robinhoodi** Horne & Whittaker, 1985  
= *P. normani* : Wagner, 1957, non Brady, 1868  
Britain

**P. bradyi** G.O. Sars, 1928  
= *P. obliquum* : Brady, 1868, non G.O. Sars, 1866  
= *P. hibernicum* : G.O. Sars, 1928, non Brady, 1868  
Bohuslän-Norway-Faeroes-Shetlands-SW Br. Isles-Ireland-N France

**P. ensiforme** Brady, 1868  
Bohuslän-Holland-?S & W Norway-Shetlands-SW Br. Isles-N France-?Bay of Biscay

**P. fleetense** Horne & Whittaker, 1985  
S England, Channel Islands

**P. angliorum** Horne & Whittaker, 1985  
= *P. ensiforme* Brady, 1868 (p.p.)  
E Anglia & Thames estuary

**P. hibernicum** Brady, 1868  
SW Br. Isles-NW Ireland

**P. normani** Brady, 1868  
Öresund & Belt Sea-Bohuslän-SW Norway-S North Sea-Faeroes-Shetlands-Br. Isles-?Bay of Biscay

**P. nealei** Horne & Whittaker, 1985  
around English & Irish coasts

**P. obliquum** G.O. Sars, 1866  
Lofoten-N Norway

**P. orcadense** Brady & Robertson, 1874?  
SW Br. Isles

**P. pulchellum** G.O. Sars, 1866  
Öresund-Bohuslän-Oslofjord-Bergen-SW Br. Isles-Ireland

**? P. rostratum** G.O. Sars, 1866  
N Norway

**P. sarniense** Brady, 1868  
W Scotland-SW England-Channel Islands-Normandie-Bretagne

**P. variabile** (Baird, 1850)  
Dalarö-Öres.-Bohuslän-North Sea-Bergen-N Norway-White-Sea-Iceland-Shetlands-Br. Isles-N Bretagne

**P. arcticum** Elofson, 1941  
Arctic Seas

**P. tenuissimum** (Norman, 1869)  
= *Bythocythere tenuissimum* Norman, 1869  
= *Xiphichilus tenuissimus* : Brady, 1870  
= *Machaerina tenuissima* : Brady & Norman, 1889  
off Scotland at 50-100 m

**P. trieri** Horne & Whittaker, 1985  
Britain-SW France

**P. porlockense** Horne & Whittaker, 1985  
around Britain

**P. amygdaloides** (Brady, 1870)  
\* Not mentioned in S.B.F. (N.S) 43, 1989  
Shetlands, Biscay, Mediterranean

**Paracytherois** G.W. Müller, 1894 (TM *P. striata* G.W. Müller, 1894)  
\* This genus is poorly known, but there seems to be at least 4 British species sensu Athersuch & al., 1989.

**P. arcuata** (Brady, 1868)  
Kieler Bucht-Öresund-Bohuslän-Bergen-Iceland-Shetlands-Br. Isles

**P. flexuosa** (Brady, 1867)  
= *Bythocythere? flexuosa* Brady, 1867  
=? *P. striata* G.W. Müller, 1894  
? non *P. flexuosa* (Brady) sensu G.O. Sars, 1928  
?Bohuslän-?W Norway-?Holland-SW Br. Isles-?Bay of Biscay-?Mediterranean

**P. producta** (Brady & Norman, 1889)  
Bergen

**Aspidoconcha** de Vos, 1953

**A. limnoriae** de Vos, 1953  
\* Not mentioned in S.B.F. (N.S) 43, 1989  
?NE Atlantic

**Sphaeromicola** Paris, 1916

**S. dudichi** Klie, 1938  
\* Not mentioned in S.B.F. (N.S) 43, 1989  
?NE Atlantic

**Cytheruridae** G.W. Müller, 1894

**Cytheropterinae** G.O. Sars, 1866

**Cytheropteron** G.O. Sars, 1866  
= *Kobayashina* Hanai, 1957  
= *Lobosocytheropteron* Ishizaki & Gunther, 1974

**C. alatum** G.O. Sars, 1866  
Bohuslän-Oslofjord-Risör (S Norway)-Trondheim-Shetlands-N Br. Isles & Ireland

**C. angulatum** (Brady, 1868)  
Öresund-Risör (S Norway)-Br. Isles-Spitsbergen

**C. crassipinnatum** Brady & Norman, 1889  
Risör (S Norway)-W Ireland

**C. depressum** Brady & Norman, 1889  
SW Br. Isles

**C. hamatum** G.O. Sars, 1870  
Stoksund (W Norway)-Lofoten-Spitsbergen, Britain

**C. inflatum** Brady, 1867  
Korshavn (S Norway)-Br. Isles

**C. latissimum** (Norman, 1865)  
= *Cythere latissima* Norman, 1865 (TM)  
= *Cytheropteron convexum* G.O. Sars, 1866  
S Baltic-Öres.-Bohuslän-North Sea-N Norway-Spitsbergen-Iceland-Shetlands-SW Br. Isles

**C. nodosum** Brady, 1868  
Korshavn (S Norway)-W Norway-Spitsbergen-Br. Isles-Bay of Biscay

**C. punctatum** Brady, 1868  
Korshavn (S Norway)-Bergen-Br. Isles-Bay of Biscay

**C. pyramidale** Brady, 1868  
Bohuslän-Oslofjord-N Norway-Arctic Seas

**C. subcircinatum** G.O. Sars, 1866  
Risör (S Norway)-Hardangerfjord

**C. testudo** G.O. Sars, 1870  
Koster Channel-Hardangerfjord-Lofoten-Spitsbergen

**C. inornatum** Brady & Robertson, 1872  
NW Britain

**C. dorsocostatum** Whatley & Masson, 1980  
S Norway-around Britain-NW Spain

**C. volantium** Whatley & Masson, 1980  
off Britain

**C. humile** Brady & Norman, 18??  
\* not mentioned in SoBF (NS) 43.  
@ burrows of *Limnoria* & *Chelura*  
?Britain

**Cytherurinae** G.W. Müller, 1894

**Hemicytherura** Elofson, 1941

**H. cellulosa** (Norman, 1865)  
= *Cythere cellulosa* Norman, 1865 (p.p.) (TM)  
= *Cytherura nana* G.O. Sars, 1866  
Kieler Bucht-Öres. & Belt Sea-Bohuslän-North Sea-S Norway-Bergen-Faeroes-Shetlands-all around Br. Isles-Bay of Biscay

**H. hoskini** Horne, 1981  
= *Cythere cellulosa* Norman, 1865 (p.p.)  
W Ireland, Bristol Channel

- H. clathrata* (G.O. Sars, 1866)  
= *Cytherura clathrata* G.O. Sars, 1866  
= *Cytheropteron clathratum* ? Auct.  
Öresund-North Sea-W Br. Isles-Trondheim-Lofoten-Spitsbergen-Br. Isles
- Cytherura* G.O. Sars, 1866
- C. gibba* (O.F. Müller, 1785)  
= *Cythere gibba* O.F. Müller, 1785 (female) (TM)  
= *Cythere gibbera* O.F. Müller, 1785 (male)  
= *Cytherura robertsoni* Brady, 1868  
SW Finland-Öres.-Skagerrak-Oslofjord-North Sea-around Br. Isles
- C. atra* G.O. Sars, 1866  
S Baltic-Lofoten-Shetlands
- C. undata* G.O. Sars, 1865  
Kieler Bucht-Bohuslän, Oslofjord-Kola peninsula, Iceland, Spitsbergen, Franz Joseph Land, Britain, Ireland
- C. concentrica* Brady, Crosskey & Robertson, 1874  
\* Not mentioned in S.B.F. (N.S.) 43, 1989  
Bergen, Hardangerfjord, Florø, Shetlands, Northumberland, Yorkshire, Durham, Spitsbergen
- Microcytherura* G.W. Müller, 1894
- M. affinis* Klie, 1938  
S Baltic
- M. fulva* (Brady & Robertson, 1874)  
S Baltic-Öres.-Bohuslän-S Norway-North Sea-SW Br. Isles-Bay of Biscay
- Semicytherura* Wagner, 1957
- S. acuticostata* (G.O. Sars, 1866)  
S Baltic-Bohuslän-Holland-Trondheim-Shetlands-around Br. Isles
- S. affinis* (G.O. Sars, 1866)  
W Norway-Spitsbergen
- S. angulata* (Brady, 1868)  
= *Cytherura insolita* Brady, 1870  
Kieler Bucht-Belt Sea-Bohuslän-Holland-Hardangerfjord-Iceland-Shetlands-SW Br. Isles
- S. cornuta* (Brady, 1868)  
= *Cytherura lineata* Brady, 1868  
= *Cytherura intumescens* G.O. Sars, 1925  
Risør (S Norway)-Br. Isles & Ireland, Bretagne
- ? *S. bidens* (Klie, 1929)  
Kieler Bucht-Öresund, Helgoland
- ? *S. brevifistulosa* (Klie, 1936)  
Helgoland
- ? *S. cochlearis* (Klie, 1929)  
Kieler Bucht
- S. concentrica* (Brady, Crosskey & Robertson, 1874)  
off Norwegian coast
- S. exserta* (Brady & Norman, 1889?)  
off Norwegian coast
- S. inconspicua* (Klie, 1934)  
Kieler Bucht-Öresund-Bergen
- ? *S. nasuta* (Klie, 1934)  
Kieler Bucht
- S. nigrescens* (Baird, 1838)  
= *Cythere nigrescens* Baird, 1838 (TM)  
Tvärminne-Öres.-Bohuslän-North Sea-Bergen-N Norway-Br. Isles-Bay of Biscay, SW Greenland
- S. producta* (Brady, 1866)  
Korshavn (S Norway)-W Norway-Holland-SW Br. Isles
- S. sella* (G.O. Sars, 1866)  
= *Cytherura cuneata* Brady, 1868 (male)  
= *Cytherura flavescens* Brady, 1869 (female)  
S Baltic-Öres.-Skagerrak-Oslofjord-North Sea-W Norway-W Br. Isles-Bay of Biscay
- S. similis* (G.O. Sars, 1866)  
S Baltic-Öres.-Bohuslän-Oslofjord-Helgoland & Holland-N Norway-Shetlands-SW Br. Isles-?Bay of Biscay
- S. simplex* (Brady & Norman, 1889)  
= *Cytherura hamata* Klie, 1936  
S Baltic-SW Br. Isles, W. Ireland, Helgoland
- S. striata* (G.O. Sars, 1866)  
= *Cytherura quadrata* Norman, 1869  
S Baltic-Öres.-Bohuslän (shells)-Oslofjord-Holland-W Norway-Faeroes-Shetlands-Br. Isles-Channel Islands-Biscay-?Mediterranean
- S. undata* (G.O. Sars, 1866)  
= *Cytherura humilis* Brady, 1867  
= *Cytherura pumila* Brady, 1874  
S Baltic-Öres.-Bohuslän-North Sea-Lofoten-Spitsbergen-Shetlands-N Br. Isles
- S. tela* Horne & Whitaker, 1980  
S Britain, Arcachon
- Cytheromatidae Elofson, 1939
- Cytheroma* G.W. Müller, 1894
- C. latiantennata* Elofson, 1938  
Bohuslän
- Terrestrialcytheroidea Schornikov, 1969  
\* Known from NW Pacific isles only
- Cypridoidea  
= Cypridacea Baird, 1845
- Macrocyprididae G.W. Müller, 1912
- Macrocypris* Brady, 1868
- M. minna* (Baird, 1850)  
= *Cythere minna* Baird, 1850 (TM)
- Kattegatt-Koster Channel-N North Sea-Lofoten-Shetlands
- Macrocypris* G.O. Sars, 1923
- M. angusta* (G.O. Sars, 1866)  
Kattegatt-Koster Channel-Oslofjord-Trondheim
- Pontocyprididae G.W. Müller, 1894
- Argilloecia* G.O. Sars, 1866
- A. conoidea* G.O. Sars, 1923  
Bohuslän-Risør (S Norway)
- A. cylindrica* G.O. Sars, 1866  
Bohuslän-Oslofjord-Hammerfest-SW Br. Isles
- A. propinqua* Brady, 1868?  
off Britain
- Propontocypris* Sylvester-Bradley, 1947
- P. trigonella* (G.O. Sars, 1866)  
= *Pontocypris trigonella* G.O. Sars, 1866 (TM)  
Koster Channel-Lofoten-N Norway-Shetlands-SW Br. Isles-?Mediterranean-?Cape Verde
- P. pirifera* (G.W. Müller, 1894)  
Scotland, S England, W France, Mediterranean
- Pontocypris* G.O. Sars, 1866 (p.p.), non G.W. Müller, 1912, nec G.O. Sars, 1923  
= *Erythrocypris* G.W. Müller, 1894
- P. hispida* G.O. Sars, 1866  
Bohuslän-Oslofjord-Risør (S Norway)-Trondheim-Shetlands-Br. Isles
- P. mytiloides* (Norman, 1862)  
= *Cypris serrulata* G.O. Sars, 1863 (TM)  
= *Cythere avena* Norman, 1865  
Korshavn & Risør (S Norway)-Møre-Br. Isles-Mediterranean
- P. pallida* (G.W. Müller, 1894)  
Korshavn & Risør (S Norway)-Mediterranean
- P. acupunctata* Brady, 1866  
\* Not mentioned in S.B.F. (N.S.) 43, 1989  
Florø, Shetlands, Northumberland, Firth of Forth, Durham
- P. frequens* (G.W. Müller, 1???)  
\* Not mentioned in S.B.F. (N.S.) 43, 1989  
?Mediterranean, ?Britain
- Cypridopsidae Kaufmann, 1910
- Cypridopsis* Brady, 1868
- C. aculeata* (Costa, 1852)  
Finnish archipel.-Norwegian estuaries-SW Br. Isles
- C. vidua* (O.F. Müller, 1785)  
Scandinavian freshwater-Thames estuary
- C. newtoni* Brady & Robertson, 1870  
Finnish rock-pools
- Potamocypris* Brady, 1870
- P. villosa* (Jurine, 1820)  
Finnish rock-pools
- P. humilis* (G.O. Sars, 18??)  
Finnish rock-pools
- Ilvocyprididae Kaufmann, 1900
- Ilvocypris* Brady & Norman, 1889
- I. gibba* (Rahmdohr, 1808)  
Freshwater in Sw., Norw., Denm. & Germ., estuaries in Arcachon-Mediterranean-Black Sea
- Cyprididae Baird, 1845
- Cyprinotinae Bronstein, 1947
- Cyprinotus* Brady, 1886
- C. incongruens* (Ramdohr, 1808)  
Baltic rock-pools
- C. salinus* (Brady, 1868)  
Baltic rock-pools-, Oslofjord estuaries
- C. reptans* (Baird, 1835)  
Fresh-water species, which can be found in North Sea "weak brackish water"
- Heterocypris* Claus, 1892
- H. salinus* (Brady, 1868)  
= *Cypris salina* Brady, 1868  
Baltic rock pools, coastal N Europe, Britain (in salt water pools (0.4-13 ppt S) & salt marshes)
- Candonidae Kaufmann, 1900
- Paracypridinae G.O. Sars, 1923
- Paracypris* G.O. Sars, 1866
- P. polita* G.O. Sars, 1866 (TM)  
Bohuslän-Trondheim-Shetlands-Br. Isles-Bay of Biscay-?Mediterranean
- Candoninae Kaufmann, 1900
- Candona* Baird, 1845
- C. angulata* G.W. Müller, 1900  
Fresh-water species, which can be found in Baltic "weak brackish water"
- C. candida* (O.F. Müller, 1785)  
Fresh-water species, which can be found in Baltic "weak brackish water", e.g. Ömsköldsvik
- C. compressa* (Koch, 1837)  
Fresh-water species, which can be found in Baltic "weak brackish water", e.g. Stockholm archipelago

*C. neglecta* G.O. Sars, 1887  
Fresh-water species, which can be found in Baltic brackish water, e.g. Ålands Sea & Central Baltic; Stadil Fjord (oligohaline lagoon in W Jutland)

*C. pratensis* Hartwig, 1901  
Fresh-water species, which can be found in Baltic "weak brackish water"

*C. sucki* Hartwig, 1901  
Fresh-water species, which can be found in North Sea "weak brackish water"

*Cyclocypris* Brady & Norman, 1889

*C. ovum* (Jurine, 1820)  
Fresh-water species, which can be found in Baltic "weak brackish water", e.g. Örnköldsvik

*C. laevis* (O.F. Müller, 1791)  
Fresh-water species, which can be found in Baltic "weak brackish water", e.g. Dalarö

*C. globosa* (G.O. Sars, 1863)  
Fresh-water species, which can be found in Baltic "weak brackish water"

*Cypria* Zenker, 1854

*C. ophthalmica* (Jurine, 1820)  
Fresh-water species, which can be found in Baltic "weak brackish water", e.g. Dalarö

*C. exculpta* (Fischer, 1855)  
Fresh-water species, which can be found in Baltic "weak brackish water"

## **MALACOSTRACA** Latreille, 1806

### **HOPLOCARIDA** Calman, 1904

**STOMATOPODA** Latreille, 1817

**UNIPELTATA** Latreille, 1825

**Squilloidea** Latreille, 1803

**Squillidae** Latreille, 1803

*Meiosquilla* Manning, 1977

*M. desmaresti* (Risso, 1816)  
S Br. Isles-Mediterranean

**Lysiosquilloidea** Giesbrecht, 1910

**Nannosquillidae** Manning, 1980

*Platysquilla* Manning, 1977

*P. eusebia* (Risso, 1816)  
S Br. Isles-Mediterranean

## **EUMALACOSTRACA** Grobben, 1892

**SYNCARIDA** Packard, 1885  
\* no marine species in the area.  
(Anaspidacea Calman, 1904 and Stygocarididae Noodt, 1963 are confined to the S Hemisphere & Bathynellacea Chappuis, 1915 is confined to Limno-terrestrial habitats)

**EUPHAUSIACEA** Dana, 1852

**Bentheuphausiidae** Colosi, 1917

*Bentheuphausia* G.O. Sars, 1885

*B. amblyops* G.O. Sars, 1885  
Atlantic, usually below 40° N

**Euphausiidae** Dana, 1852

*Euphausia* Dana, 1850

*E. krohnii* (Brandt, 1851)  
off W Norway-off Iceland-E Shetlands-NE Scotland-off W Br. Isles-southwards

*Meganyctiphanes* Holt & Tattersall, 1905

*M. norvegica* (M. Sars, 1857)  
Katteg.-Bohuslän-Barents Sea-Spitsbergen-Greenland-W Br. Isles-Azores-Mediterranean

*Nyctiphanes* G.O. Sars, 1884

*N. couchii* (Bell, 1853)  
?S Baltic-W Skagerrak-Møre-Shetlands-W Br. Isles-Azores-Mediterranean

*Thysanoessa* Brandt, 1851

*T. inermis* (Krøyer, 1846)  
Katteg.-Bohuslän-N North Sea-Barents Sea-E Greenland-off W Br. Isles-south to 50° N

*T. raschii* (M. Sars, 1864)  
Katteg.-Bohuslän-N North Sea-N Scotland-Iceland-Greenland-Spitsbergen & Kara Sea-Arctic Seas

*T. longicaudata* (Krøyer, 1846)  
Koster Channel, Skagerrak-N North Sea-Barents Sea-E Greenland-off W Br. Isles-off Portugal

*T. gregaria* G.O. Sars, 1883  
Atlantic, between 20-40° N & W Mediterranean

*Thysanopoda* Milne-Edwards, 1830

*T. acutifrons* Holt & Tattersall, 1906  
W-SW Norway-Shetland-Faeroe Channel-Iceland-off W Ireland-off W Gibraltar

*Nematoscelis* G.O. Sars, 1884, non Wollaston, 1867  
\* Does a proper name exist?

*N. megalops* G.O. Sars, 1884  
S Greenland-S Iceland-S & N Norway-North Sea-off W Br. Isles-Azores

*Nematobranchion* Calman, 1905

*N. boopis* (Calman, 1896)  
S Greenland-S Iceland-Faeroe-Shetland Channel-off W Br. Isles-NW Africa

*Stylocheiron* G.O. Sars, 1884

*S. elongatum* G.O. Sars, 1884

S Iceland-N, W & SW Br. Isles-southwards

*S. longicornis* G.O. Sars, 1884  
S Iceland-Bergen-off W Br. Isles-southwards

*S. maximum* Hansen, 1908  
off SE Greenland-off S Iceland-Faeroes-Shetlands-off W Ireland-southwards

*S. abbreviatum* G.O. Sars, 1884  
NW Norway-Barents Sea-Iceland-SE Greenland-Faeroe-Shetland Channel-off W Ireland

**AMPHIONIDACEA** Williamson, 1973

\* With a circumglobal marine planktonic species, *Amphionides reynaudii* (H. Milne-Edwards) as only species. (Not in N Europe)

**DECAPODA** Latreille, 1803

**DENDROBRANCHIATA** Bate, 1888

\* This subordo has been grouped together with Stenopodidea, Procarididea & Caridea in the next subordos, together constituting the older subordo Natantia Boas, 1880

**Penaeoidea** Rafinesque-Schmaltz, 1815

**Penaeidae** Rafinesque-Schmaltz, 1815

**Penaeinae** Rafinesque-Schmaltz, 1815

Tribe: Peneini Burkenroad, 1936

*Penaeus* Fabricius, 1798

*P. kerathurus* (Forskål, 1775)

\* "Caramote prawn or shrimp", "Tiger shrimp", "Striped shrimp", "Langostino"  
Mediterranean, E Atlantic up to W Engl. Channel

*P. semisulcatus* W. De Haan, in Siebold, 1849  
E Atlantic

*Funchalia* Johnson, 1867

= *Pelagopenaeus* Burkenroad, 1934

*F. woodwardi* Johnson, 1867

Hebrides-southwards in Atlantic, Mediterranean

*F. balboae* (Faxon, 1893)

= *Pelagopenaeus balboae* (Faxon, 1893)  
?Mediterranean

*Heteropeneus* De Man, 1896

Tribe: Parapeneini Burkenroad, in Schram, 1983

*Metapenaeopsis* Bouvier, 1905

*M. miersi* Holthuis, 1952  
E Atlantic off W Africa

*Parapenaeus* Smith, 1885

*P. longirostris* (H. Lucas, 1846)

= *Penaeus longirostris* H. Lucas, 1846  
\* "Deep water rose shrimp", "Gamba"  
E Atlantic up to N Spain, Mediterranean

*Penaeopsis* Bate, 1881

= *Peneopsis* Faxon, 1895

*Artemesia* Bate, 1888

Tribe: Trachypeneini Burkenroad, in Schram, 1983

*Parapenaeopsis* Alcock, 1901

*P. atlantica* Balss, 1926  
E Atlantic off W Africa

*Mangalura* Miers, 1878

*Macropetasma* Stebbing, 1914

*Trachypeneopsis* Burkenroad, 1934

*Atypopeneus* Alcock, 1905

*Protrachypene* Burkenroad, 1934

*Xiphopeneus* S.I. Smith, 1886

*Trachypeneus* Alcock, 1901

**Sicyoniinae** Ortman, 1898

*Sicyonia* H. Milne-Edwards, 1840

**Aristeidae** Wood-Mason, 1891

**Aristeinae** Wood-Mason, 1891

Tribus: Benthysycimini Bouvier, 1908

*Gennadas* Bate, 1881

*G. elegans* (S.I. Smith, 1882)

= *Amatopeneus elegans* S.I. Smith, 1882  
SW Greenland-Iceland-between Iceland & Faeroes-off W Ireland-Bay of Biscay, Mediterranean

*Benthysycymus* Bate, 1881

*Bentheogenemma* Burkenroad, 1936

Tribus: Aristeini Bouvier, 1908

*Plesiopenaeus* Bate, 1881

= *Plesiopeneus* Faxon, 1895

*P. edwardsianus* (Johnson, 1867)

\* "Scarlet shrimp"  
Along E Atlantic coasts up to S Biscay

*Hemipenaeus* Bate, 1881

*Aristeus* Duvernoy, 1840

- Hepomadus** Bate,1881
- Aristeomorpha** Wood-Mason,1891
- Solenocerinae** Wood-Mason & Alcock,1891
- Solenocera** Lucas,1849
- S. membranacea** (Risso,1816)  
= *Peneus membranaceus* Risso,1816  
= *Peneus siphonoceros* Philippi,1840  
?European Atlantic, African Atlantic, Mediterranean
- Hymenopeneus** S.I. Smith,1882
- Haliporus** Bate,1881
- Sergestoidea** Dana,1852
- Sergestidae** Dana,1852
- Sergestinae** Dana,1852
- Sergestes** H. Milne-Edwards,1830
- S. arcticus** Krøyer,1855  
Kattegatt-W Norway-Norw. Sea
- S. mollis** S.I. Smith,1884  
?European Atlantic
- Sergia** Ortmann,1893
- S. robustus** (S.I. Smith,1882)  
= *Sergestes robustus* S.I. Smith,1882  
N North Sea-Norw. Sea-Faeroes-Iceland-Cape Verde Isles
- Sicyonellinae** Burkenroad,1983
- Luciferidae** Burkenroad,1983  
\* No species within the area?
- EUKYPHIDA** Boas,1880
- PROCARIDIDEA** Chace & Manning,1972  
\* No representing species within the area. 3 sp. known from caves in Bermuda, Hawaii & Ascension
- CARIDEA** Dana,1852
- Oplophoroidea** Dana,1852
- Oplophoridae** Dana,1852
- Meningodora** S.I. Smith,1882 (TM *Meningodora mollis* S.I. Smith,1882)  
=? *Notostomus* A. Milne-Edwards,1881
- M. vesca** S.I. Smith,1882?  
Faeroe-Iceland area
- Acanthephyra** A. Milne-Edwards,1881 (TM *Acanthephyra armata* A. Milne-Edwards,1881)  
= *Ephyra* P. Roux,1831, non Péron & Lesueur,1810 (Cnidaria), nec Duponchel,1829 (Lepidoptera)  
= *Miersia* Kingsley,1880 (Replacement name for *Ephyra* P. Roux, priority suppressed by ICZN)
- A. purpurea** A. Milne-Edwards,1881  
Shetland-Faeroe area-W Ireland-Bay of Biscay
- A. pelagica** (Risso,1816)  
= *Alpheus pelagicus* Risso,1816  
=? *Hoplocarcyphus similis* Coutière,1907  
Shetland-Faeroe-Iceland area-W Ireland-Bay of Biscay
- Hymenodora** G.O. Sars,1877
- H. glacialis** (Buchholz,1874)  
= *Pasiphaë glacialis* Buchholz,1874 (TM)  
Barents & Norw. Sea-Shetland-Faeroe-Iceland area-W Ireland-?Bay of Biscay
- H. gracilis** (S.I. Smith,1882)  
= *Miersia gracilis* S.I. Smith,1882  
off W & S Ireland-Bay of Biscay
- Systellaspis** Bate,1888 (TM *Systellaspis lanceocaudata* Bate,1888 - "Challenger" material)  
= *Hoplopassiphaea* Yokoya & Shibata,1965 (TM *Hoplopassiphaea philippinensis* Yokoya & Shibata,1965)
- S. debilis** (A. Milne-Edwards,1881)  
= *Hoplopassiphaea philippinensis* Yokoya & Shibata,1965  
Shetland-Faeroe-Iceland area-W Ireland-Bay of Biscay
- Atyoidea** de Haan,1849
- Atyidae** de Haan,1849  
\* Predominantly freshwater distribution, although 2 species occur in brackish waters
- Atyaephyra** de Brito Capello,1867
- A. desmarestii desmarestii** (Millet,1832)  
= *Hyppolyte Desmarestii* Millet,1832  
?European Atlantic, Madeira, Mediterranean
- Bresilioidea** Calman,1896
- Bresiliidae** Calman,1896  
= **Disciidae** Rathbun,1902
- Bresilia** Calman,1896
- B. atlantica** Calman,1896 (TM)  
off SW Ireland
- Nematocarcinoidea** S.I. Smith,1884
- Rhynchocinetidae** Ortmann,1890  
\* No species within the area?
- Nematocarcinidae** S.I. Smith,1884
- Nematocarcinus** A. Milne-Edwards,1881 (TM *Nematocarcinus cursor* A. Milne-Edwards,1881)  
= *Eumiersia* S.I. Smith,1882 (TM *Eumiersia ensifera* S.I. Smith,1882)  
= *Stochasmus* Bate,1888 (TM *Stochasmus exilis* Bate,1888)
- N. ensifer** (S.I. Smith,1882) **exilis** (Bate,1888)  
= *Stochasmus exilis* Bate,1888  
Shetland-Faeroe-Iceland area, Bay of Biscay
- Styrodactyloidea** Bate,1888  
\* No species within the area?
- Styrodactylidae** Bate,1888
- Pasiphaeidea** Dana,1852
- Pasiphaeidae** Dana,1852
- Pasiphaea** Savigny,1816  
= *Phye* Wood Mason,1892 (TM *Parapasiphaë alcocki* Wood Mason & Alcock,1891)
- P. tarda** Krøyer,1845  
Koster Channel-Skagerrak-Norw. Sea-Barents Sea-Faeroe Shetland Iceland area
- P. multidentata** Esmark,1866  
Bohuslän-Skagerrak-N North Sea-Norw. Sea-Faeroe Shetland Iceland area-W Ireland-Bay of Biscay
- P. sivado** (Risso,1816)  
= *Alpheus sivado* Risso,1816 (TM)  
Kattegatt-Bohuslän-Norw. Sea-Shetland Faeroe area-W (& S) Br. Isles-Bay of Biscay
- Parapasiphae** S.I. Smith,1884  
= *Orphanina* Bate,1888, non Fischer,1853 (Orthoptera)  
= *Dantecia* Caullery,1896
- P. sulcatifrons** S.I. Smith,1884 (TM)  
= *Orphanina tenuimana* Bate,1888 (TM of *Orphanina* Bate)  
= *Dantecia caudant* Caullery,1896 (TM of *Dantecia* Caullery)  
Faeroe Iceland area-off W Ireland
- P. cristata** S.I. Smith,1884  
Shetland Faeroe area-off W Ireland
- Palaemonidea** Rafinesque-Schmaltz,1815
- Palaemonidae** Rafinesque-Schmaltz,1815
- Palaemoninae** Rafinesque-Schmaltz,1815
- Palaemon** Weber,1795
- P. (Palaemon) Holthuis,1950 elegans** Rathke,1837 (TM of *Palaemon* Holthuis,1950)  
= *Leander squilla* - Czerniavsky,1884, non (Linnaeus,1758)  
S Baltic-Bohuslän-North Sea-Shetland Faeroe area-W Br. Isles-Engl. Channel-Bay of Biscay
- P. adpersus** Rathke,1837 (TM)  
= *P. fabricii* Rathke,1843 & *Cancer Squilla* Linnaeus,1758  
Åland-Bohuslän-North Sea-Norw. Sea-Br. Isles-Bay of Biscay
- P. serratus** (Pennant,1777)  
Öresund, ?Skagerrak-North Sea-Engl. Channel-W Br. Isles-Bay of Biscay
- P. longirostris** Milne-Edwards,1837  
S North Sea-Engl-Channel, W Ireland-Bay of Biscay
- Palaemonetes** Heller,1869
- P. varians** (Leach,1814)  
= *Palaemon varians* Leach,1814 (TM)  
Fresh water in Sw., Nor. & Denm., S Baltic, ?Kattegatt-?Skagerrak-North Sea-Engl. Channel-W Br. Isles-Bay of Biscay
- Leander** E. Desmarest,1849
- L. tenuicornis** (Say,1818)  
= *Palaemon tenuicornis* Say,1818  
= *Leander erraticus* E. Desmarest,1849 (TM)  
?European Atlantic
- Pontoniinae** Kingsley,1878
- Typton** Costa,1844
- T. spongicola** Costa,1844 (TM)  
Engl. Channel-Bay of Biscay
- Periclimenes** Costa,1844
- P. sagittifer** (Norman,1861)  
Roscoff
- Gnathophyllidae** Dana,1852  
\* No species within the area?
- Hymenoceridae** Ortmann,1890  
\* No species within the area?
- Campylonotoidea** Sollaud,1913
- Campylonotidae** Sollaud,1913  
\* No species within the area?
- Psalidopodoidea** Wood-Mason & Alcock,1892  
\* The only 2 species known are W Atlantic & Indo-pacific in distribution
- Psalidopodidae** Wood-Mason & Alcock,1892
- Alpheoidea** Rafinesque-Schmaltz,1815
- Alpheidae** Rafinesque-Schmaltz,1815
- Athanas** Leach,1814
- A. nitescens** (Leach,1814)  
= *Palaemon nitescens* Leach,1814 (TM)  
Öresund-Bohuslän-W Norway-S North Sea-Engl. Channel-W Br. Isles-Mediterranean & Black Sea
- Alpheus** Weber,1795 (TM *Alpheus avarus* Fabricius,1798)
- A. glaber** (Olivieri,1792)  
Engl. Channel-W Br. Isles-Bay of Biscay
- A. macrocheles** (Hailstone,1835)  
Engl. Channel-Bay of Biscay
- Hippolytidae** Bate,1888

- Cryptocheles* G.O. Sars, 1870
- C. pygmaea* G.O. Sars, 1870 (TM)  
Bohuslän-N North Sea-Norw. Sea
- Hippolyte* Leach, 1814  
= *Virbius* Stimpson, 1860 (TM *Hippolyte acuminata* Dana, 1852)
- H. varians* Leach, 1814 (TM)  
@ *Alcyonium digitatum*  
Øresund-Bohuslän-North Sea-Norw. Sea-Shetland Faeroe area-W Br. Isles-Engl. Channel-Bay of Biscay
- H. inermis* Leach, 1815  
= *Hippolyte prideuxiana* Leach, 1817  
Engl. Channel-W Br. Isles-Bay of Biscay
- H. longirostris* (Czerniavsky, 1868)  
Engl. Channel-Bay of Biscay
- H. hunti* (Gosse, 1877)  
@ *Antedon bifida*  
Britain
- Bythocaris* G.O. Sars, 1870
- B. simplicirostris* G.O. Sars, 1870 (TM)  
Norw. Sea-Barents Sea-Shetland-Faeroes area
- B. leucopsis* G.O. Sars, 1879  
Norw. Sea-Barents Sea-Faeroes-Iceland area
- B. payeri* (Heller, 1875)  
Norw. Sea-Barents Sea-Shetland-Faeroes-Iceland area
- B. gracilis* S.I. Smith, 1885  
Shetland-Faeroe-Iceland area-off W Ireland
- Thoralus* Holthuis, 1947  
= *Lysippe* Kinahan, 1858, non Malmgren, 1865 (Polychaeta) (TM *Hippolyte cranchii* Leach, 1817 - generic name totally suppressed, ICZN Op. 671)
- T. cranchii* (Leach, 1817)  
= *Hippolyte cranchii* Leach, 1817 (TM)  
Bohuslän-North Sea-Norw. Sea-Shetland Faeroe area-W Br. Isles-Engl. Channel-Bay of Biscay
- Spirontocaris* Bate, 1888
- S. liljeborgii* (Danielsen, 1859)  
= *S. liljeborgii* : Auctt.  
\* *Spirontocaris securifrons* (Norman, 1862)  
\* Vilhelm or Wilhelm Liljeborg changed his name to William Liljeborg (with a double l) around 1860, when he for a while was living in USA. This spelling should be reflected when citing him as an author (single l before 1860 and double after), but the original spelling of taxon names based upon his name must be used.  
S Baltic-Bohuslän-N North Sea-W Br. Isles & Engl. Channel-Shetland Faeroe Iceland are-Norw. Sea-Barents Sea
- S. spinus* (James Sowerby, 1805)  
= *Cancer spinus* James Sowerby, 1805 (TM)  
?Øresund-Skagerrak-N North Sea-W Br. Isles-Shetland Faeroe Iceland area-Norw. Sea-Barents Sea
- S. phippisii* (Krøyer, 1841)  
= *Spirontocaris turgida* (Krøyer, 1841)  
?Bohuslän-Skagerrak-N North Sea-Norw. Sea-Barents Sea
- Lebbeus* White, 1847
- L. polaris* (Sabine, 1824)  
= *Alpheus polaris* Sabine, 1824  
= *Lebbeus orthorhynchus* (Leach MS) White, 1847 (TM)  
Kattegat-Bohuslän-N North Sea-Shetland Faeroe Iceland area-Norw. Sea-Barents Sea
- Eualus* Thallwitz, 1892
- E. gaimardii* (H. Milne-Edwards, 1837)  
= *Hippolyte gaimardii* H. Milne-Edwards, 1837  
= *Eualus obses* Thallwitz, 1892 (TM)  
S Baltic-Bohuslän-North Sea-NW Br. Isles-Faeroe Iceland area-Norw. Sea-Barents Sea
- E. pusiolus* (Krøyer, 1841)  
Øresund-Bohuslän-North Sea-W Br. Isles-Shetland Faeroe Iceland area-Norw. Sea-Barents Sea
- E. occultus* (Lebour, 1936)  
Bohuslän-North Sea-Engl. Channel-W Br. Isles-Bay of Biscay
- Caridion* Goës, 1863  
= *Doryphorus* Norman, 1861, non Cuvier, 1829 (Reptilia)
- C. gordonii* (Bate, 1858)  
= *Hippolyte gordonii* Bate, 1858 (TM)  
Kattegat-Bohuslän-North Sea-Barents Sea-Iceland-Faeroes-Br. Isles-Bay of Biscay
- C. steveni* Lebour, 1930  
Skagerrak-North Sea-Engl. Channel & W Br. Isles
- Lysmata* Risso, 1816  
= *Melicerta* Risso, 1816, non Schrank, 1803 (Vermes), non Péron & Lesueur, 1810 (Cnidaria)
- L. seticaudata* (Risso, 1816)  
= *Melicerta seticaudata* Risso, 1816 (TM)  
Engl. Channel-Bay of Biscay
- Ogyridae* Hayard Share, 1918  
\* No species within the area?
- Processoidea* Ortmann, 1890
- Processidae* Ortmann, 1890
- Processa* Leach, 1815  
= *Nika* Risso, 1816 (TM *Nika edulis* Risso, 1816)  
\* At least 2 species in Scandinavia. Which ones?
- P. canaliculata* Leach, 1815 (TM)  
= *Nika mediterranea* Parisi, 1915 & *P. mediterranea* : Nouvel & Holthuis, 1957  
N North Sea-W Br. Isles-Engl. Channel-Bay of Biscay
- P. edulis* (Risso, 1816) *crassipes* Nouvel & Holthuis, 1957  
S North Sea-Engl. Channel-W Br. Isles, Roscoff
- P. edulis arcassonensis* Nouvel & Holthuis, 1957  
Arcachon
- P. modica* D.I. Williamson & Rochanaburanon, 1979 *modica* D.I. Williamson & Rochanaburanon, 1979  
= *P. parva* : Holthuis, 1954, : Nouvel & Holthuis, 1957, non Holthuis, 1951 & *P. aequimana* : Auctt., non (Paulson, 1875)  
S North Sea-Engl. Channel-W Br. Isles-Bay of Biscay
- P. nouveli* Al-Adhub & D.I. Williamson, 1975 *holthuisi* Al-Adhub & D.I. Williamson, 1975  
= *P. canaliculata* : Nouvel & Holthuis, 1957, non Leach, 1815  
Bohuslän-North Sea-W Br. Isles-Bay of Biscay
- P. elegantula* Nouvel & Holthuis, 1957  
Roscoff, Mediterranean
- P. parva* Holthuis, 1951  
?European Atlantic, tropical W Africa
- Pandaloidae* Haworth, 1825
- Pandalidae* Haworth, 1825
- Pandalus* Leach, 1814  
= *Dymas* Krøyer, 1861
- P. borealis* Krøyer, 1838  
= *Dymas typus* Krøyer, 1861 (TM of *Dymas* Krøyer, 1861)  
Kattegat-Bohuslän-N North Sea-Norw. Sea-Barents Sea-Shetland Faeroe Iceland area
- P. montagui* Leach, 1814 (TM)  
S Baltic-Bohuslän-North Sea-Norw. Sea-Barents Sea-Shetland Faeroe Iceland area-W Br. Isles-Engl. Channel
- P. propinquus* G.O. Sars, 1870  
Bohuslän-Norw. Sea-Shetland Faeroe Iceland area-W Br. Isles-Bay of Biscay
- Dichelopandalus* Caullery, 1896
- D. bonnierii* Caullery, 1896 (TM)  
Bohuslän-Norw. Sea-Shetland Faeroe Iceland area-W Br. Isles-Bay of Biscay
- Pandalina* Calman, 1899
- P. brevirostris* (Rathke, 1843)  
= *Pandalus brevirostris* Rathke, 1843 (TM)  
Øresund-Bohuslän-North Sea-Norw. Sea-Shetland Faeroe area-W Br. Isles-Engl. Channel-Bay of Biscay
- P. profunda* Holthuis, 1946  
Skagerrak-Norw. Sea-Shetland Faeroe area-W Ireland-?Bay of Biscay
- Plesionika* Bate, 1888 (TM *P. uniproductata* Bate, 1888 (Junior subjective synonym of *Acanthephyra ensis* A. Milne-Edwards, 1881))
- P. edwardsii* (Brandt, 1851)  
\* "Striped soldier shrimp", "Soldier striped shrimp"  
From E tropical Atlantiv to S Biscay, Mediterranean
- Thalassocarididae* Bate, 1888  
\* Only 2 indo-pacific species known
- Physetocaridoidea* Chase, 1940  
\* Only one bathypelagic species known from 1000-1200 m depths off Bermuda
- Physetocarididae* Chase, 1940
- Crangonoidea* Haworth, 1825
- Crangonidae* Haworth, 1825
- Crangon* J.C. Fabricius, 1798
- C. crangon* (Linnaeus, 1758)  
= *Cancer crangon* Linnaeus, 1758 (TM)  
Åland-Bohuslän-North Sea-Barents Sea-Iceland-Faeroes-Br. Isles-Bay of Biscay
- C. allmani* Kinahan, 1857  
Øresund-Bohuslän-North Sea-Barents Sea-Iceland-Faeroes-Br. Isles-Bay of Biscay
- Sclerocrangon* G.O. Sars, 1883
- S. boreas* (Phipps, 1774)  
= *Cancer boreas* Phipps, 1774 (TM)  
Norw. Sea-Faeroe-Iceland area-Barents Sea
- S. ferox* (G.O. Sars, 1877)  
Shetland-Faeroe area-off W Ireland-Norw. Sea-Barents Sea, N Greenland
- S. jacqueti* (A. Milne-Edwards, 18??)  
Shetland-Faeroe area-off W Ireland
- Philocheras* Stebbing, 1900
- P. echinulatus* (M. Sars, 1861)  
Bohuslän-N North Sea-Norw. Sea-Shetland Faeroe area-W Br. Isles-Bay of Biscay
- P. trispinosus* (Hailstone, 1835)  
North Sea-Shetland Faeroe area-W Br. Isles-Engl. Channel-Bay of Biscay
- P. bispinosus* (Hailstone, 1835)  
= *Pontophilus bispinosus* Hailstone, 1835  
= *Crangon nanus* Krøyer, 1842 (TM)  
Kattegat-Bohuslän-North Sea-Norw. Sea-Iceland-Br. Isles-Bay of Biscay
- P. sculptus* (Bell, 1853)  
?N North Sea, Engl Channel-W Br. Isles-Bay of Biscay
- P. fasciatus* (Risso, 1816)  
North Sea-?Norw. Sea-Engl. Channel-W Br. Isles-Bay of Biscay
- Pontophilus* Leach, 1817  
= *Cheraphilus* Kinahan, 1862 (TM *Crangon spinosus* Leach, 1815)
- P. norvegicus* (M. Sars, 1861)  
Bohuslän-Norw. Sea-Iceland-Barents Sea
- P. spinosus* (Leach, 1815)  
= *Crangon spinosus* Leach, 1815 (TM)  
Kattegat-Bohuslän-North Sea-Norw. Sea-Iceland-Faeroes-Br. Isles-Bay of Biscay
- Sabinea* J.C. Ross, 1835
- S. sarsii* Smith, 1879  
Bohuslän-Norw. Sea-Iceland-Barents Sea

- S. septemcarinatus* (Sabine,1824)  
= *Crangon septemcarinatus* Sabine,1824 (TM)  
Norw. Sea-Iceland-Barents Sea, N Greenland
- S. hystrix* (A. Milne-Edwards,1881)  
off SW Iceland
- Aegaeon* Agassiz,1846
- A. lacazei* (Gouret,1887)  
?European Atlantic
- Glyphocrangonidae Smith,1884  
\* No species within the area?
- EUZYGIDA Burkenroad,1981  
\* An extinct infraorder, Uncinidea Beurlen,1930 is also belonging here
- STENOPODIDEA Bate,1888
- Stenopodidae Claus,1872
- Richardina* A. Milne-Edwards,1881
- R. spinicincta* A. Milne-Edwards,1881 (TM)  
off W Ireland-Bay of Biscay
- Spongiocolidae Schram,1986
- Spongicolidoides* Hansen,1908
- S. profundus* Hansen,1908 (TM)  
off SW Iceland
- REPTANTIA Boas,1880
- ASTACIDEA Latreille,1803
- Astacidae Latreille,1802  
\* Some other fresh-water taxa exists in the area
- Astacus* J.C. Fabricius,1775
- A. astacus* (Linnaeus,1758)  
= *A. fluviatilis* J.C. Fabricius,1775  
Fresh water, in Sw., Denm., Finl., & Norw.
- Nephronidae Dana,1852
- Nephrops* Leach,1814
- N. norvegicus* (Linnaeus,1758)  
Öresund-Bohuslän-North Sea-N Norway-Iceland-Br. Isles-Mediterranean
- Nephropsis* Wood-Mason,1873
- N. atlantica* Norman,1882  
off N Scotland-Shetland Faeroe Channel-Bay of Biscay
- Homarus* Weber,1795
- H. gammarus* (Linnaeus,1758)  
= *H. vulgaris* H. Milne-Edwards,1837  
Kattegatt-Bohuslän-North Sea-Troms-Br. Isles-Mediterranean
- Thaumastocheilidae Bate,1888  
\* Only 2 species known, from off central America & off Japan
- Cambaridae Hobbs,1942  
\* Only freshwater species
- Parastacidae Huxley,1879  
\* Freshwater species in the Indo-pacific area & S South America
- THALASSINIDEA Latreille,1831
- Thalassinidae Latreille,1831  
\* No species within the area?
- Axiidae Huxley,1879
- Calocaris* Bell,1846
- C. macandreae* Bell,1846  
Kattegatt-Bohuslän-North Sea-Troms-Iceland-Br. Isles-Adriatic
- Calocarides* Wollebaek,1909
- C. coronatus* (Trybom,1904)  
= *Eiconaxius coronatus* Trybom,1904 (TM)  
Bohuslän-Norw. Sea-Finmark-N Br. Isles
- Axius* Leach,1815
- A. styrnchus* Leach,1815  
Br. Isles-Mediterranean
- A. nodulosus* Meinert,1877  
off W Jutland
- Laomediidae Borradaile,1903
- Jaxea* Nardo,1847
- J. nocturna* Nardo,1847 ex Chiereghin MS  
\* Jim Atkinson at Millport, when studying video tapes from the Swedish west coast with burrows of *Nephrops* & *Calocaris* in 50-60 m. depth, also found burrows typical of *Jaxea* (in fax to Mats Ulmestrand autumn 1996)  
SW Br. Isles & Irish Sea-Adriatic
- Callianassidae Dana,1852
- Callianassa* Leach,1814
- C. subterranea* (Montagu,1808) (TM)  
N Kattegatt-Bohuslän-Skagerrak-Engl. Channel-W Br. Isles-Mediterranean
- C. tyrrhena* (Petagna,1792)  
= *C. stebbingi* Borradaile,1903  
S North Sea-Scotland-Engl. Channel-Mediterranean
- Upogebiidae Borradaile,1903
- Upogebia* Leach,1814
- U. stellata* (Montagu,1814)  
Kattegatt-Bohuslän-North Sea-W Norway-W Br. Isles-Engl. Channel-Mediterranean
- U. deltaura* (Leach,1815)  
Kattegatt-Bohuslän-North Sea-W Norway-W Br. Isles-Engl. Channel-Mediterranean
- U. pusilla* (Petagna,1792)  
Sponvika, Norway, Britain, Spain, Mediterranean
- Callianideidae Kossman,1880  
\* Indo-Pacific & West Indian distribution
- Axianassidae Schmitt,1924  
\* No species within the area?
- PALINURA Latreille,1803
- Palinuridae Latreille,1803
- Palinurus* Weber,1795
- P. elephas* (J.C. Fabricius,1787)  
= *P. vulgaris* Latreille,1808  
Bergen-North Sea-W Br. Isles-Engl Channel-Mediterranean
- P. mauritanicus* Gruvel,1911  
off SW Ireland-Senegal, Mediterranean
- Polychelidae Wood-Mason,1874
- Polycheles* Heller,1862
- P. typhlops* Heller,1862  
= *Polycheles agassizi* Faxon,19??  
= *Polycheles doederleini* Riggio,1885  
= *Pentacheles hextii* Alcock,1901  
SW Iceland-SW Ireland-Bay of Biscay-Cap Verde Isles & Mediterranean
- P. granulatus* Faxon,1893  
off SW Ireland-Madeira & Canaries
- Stereomastis* Bate,1888
- S. sculpta* (S.I. Smith,1880)  
= *Polycheles sculptus* S.I. Smith,1880 (Apr.)  
= *Pentacheles spinosus* A. Milne-Edwards,1880 (Dec.)  
off SW Iceland-W Ireland-Biscay-Mediterranean
- S. grimaldi* (Bouvier,19??)  
off Iceland-off W Ireland-Biscay
- Glypheidae Winckler,1883  
\* The only recent species is known from the vicinity of the Philippines
- Scyllaridae Latreille,1825
- Scyllarus* J.C. Fabricius,1775
- S. arctus* (Linnaeus,1758)  
W Engl. Channel, Madeira, Azores, Mediterranean
- Synaxidae Bate,1881  
\* Monotypical family from the West Indies
- ANOMALA Boas,1880  
\* This infraorder has often been combined with Thalassinidea in a larger taxon Anomura MacLeay,1838
- Diogenidae Ortmann,1892
- Diogenes* Dana,1852 (TM *Pagurus miles* J.C. Fabricius,1787)
- D. pugilator pugilator* (Roux,1829)  
= *Pagurus pugilator* Roux,1829  
German Bight, Netherlands, Swansea-Engl. Channel-Bay of Biscay-Mediterranean
- Clibanarius* Dana,1852 (TM *Cancer clibanarius* Herbst,1791)
- C. erythropus* (Latreille,1818)  
= *Pagurus erythropus* Latreille,1818  
= *Pagurus misanthropus* Roux,1828-30  
N Cornwall & Engl. Channel-Bay of Biscay-Azores-Mediterranean
- Paguridae Latreille,1803
- Pagurus* J.C. Fabricius,1775
- P. bernardus* (Linnaeus,1758)  
= *Cancer bernardus* Linnaeus,1758 (TM)  
Öresund-Bohuslän-North Sea-N Norway-Barents Sea-area S of Spitsbergen-Shetland Faeroe Iceland area-Br. Isles-Bay of Biscay-Portugal
- P. pubescens* Krøyer,1838  
= *Pagurus thompsoni* Bell,1853  
Kattegatt-Bohuslän-N North Sea-N Norway-Barents Sea & Spitsbergen-Shetland Faeroe Iceland area-W Br. Isles-Engl. Channel
- P. cuanensis* Bell,1845  
Öresund-Bohuslän-N North Sea-W Norway (Bergen area)-?Shetland Faeroe area-W Br. Isles-Engl. Channel-Bay of Biscay
- P. prideaux* Leach,1815  
= *Pagurus prideauxi* : Auctt.  
= *Pagurus solitarius* Roux,1828-30  
?Skagerrak-S & N North Sea-Trøndelag-Faeroe Shetland area-W Br. Isles-Engl. Channel-Bay of Biscay-Mediterranean
- P. alatus* J.C. Fabricius,1775  
= *Pagurus angulatus* Risso,1816  
= *Pagurus tricarinus* Norman,1869  
= *Eupagurus variabilis* A. Milne-Edwards & Bouvier,1892  
W Norway-N North Sea-W Norway-Shetland Faeroe area-Iceland-W Ireland-Bay of Biscay-Mediterranean
- P. forbesii* Bell,1846  
= *P. sculpiimanus* Lucas,1846  
Engl. Channel-Irish Sea-W Ireland-Bay of Biscay-Portugal, Madeira, Mediterranean
- P. carneus* Pocock,1889  
Faeroes in >500m, off W Ireland

- P. chevreuxi* (Bouvier, 1896)  
= *Eupagurus chevreuxi* Bouvier, 1896  
N Bretagne-Roscoff-Mediterranean
- P. pubescentulus* (A. Milne-Edwards & Bouvier, 1892)  
= *Eupagurus pubescentulus* A. Milne-Edwards & Bouvier, 1892  
46° 09' N, 3° 38' E - Biscay - Angola, Azores
- P. anachoretus* Risso, 1826  
= *Pagurus pictus* H. Milne-Edwards, 1836  
N Portugal - southwards
- Anapagurus* Henderson, 1886
- A. chiroacanthus* (Liljeborg, 1855)  
= *Pagurus chiroacanthus* Liljeborg, 1855  
= *Anapagurus brevicarpus* A. Milne-Edwards & Bouvier, 1892  
Kattegat-Bohuslän-N North Sea-W Norway (Bergen area)-Shetland Faeroe area-W Br. Isles-Engl. Channel-Biscay- Azores-Mediterranean
- A. laevis* (Bell, 1845)  
= *Pagurus laevis* Bell, 1845 (TM)  
= *Pagurus laevis* W. Thompson, 1844 (nomen nudum)  
Kattegat-Bohuslän-North Sea-W Norway-Shetland Faeroe area-W Br. Isles-Engl. Channel-Bay of Biscay-Mediterranean
- A. hyndmanni* (Bell, 1845)  
= *Pagurus hyndmanni* Bell, 1845  
= *Pagurus hyndmanni* W. Thompson, 1844 (nomen nudum)  
N North Sea-Shetland Faeroe area-W Br. Isles-Engl. Channel-Bay of Biscay
- Cestopagurus* Bouvier, 1897 (TM C. *coutieri* Bouvier, 1897)
- C. timidus* (Roux, 1828-30)  
= *Pagurus timidus* Roux, 1828-30  
= *Capaguroides* Milne-Edwards & Bouvier, 1892 *timidus* (Roux, 1828-30)  
Clyde Sea area, Ireland N Bretagne-Bay of Biscay-Mediterranean
- Nematopagurus* A. Milne-Edwards & Bouvier, 1892
- N. longicornis* A. Milne-Edwards & Bouvier, 1892 (TM)  
off Galway & W Ireland, N Spain-Azores-Madeira-Angola-Mediterranean
- Lithodidae Samouelle, 1819
- Lithodes* Latreille, 1806
- L. maja* (Linnaeus, 1758)  
= *Cancer maja* Linnaeus, 1758  
Oresund-Bohuslän-North Sea-N Norway-Barents Sea & Spitsbergen-Shetland Faeroe Iceland area-W Br. Isles
- Parapaguridae Smith, 1882
- Parapagurus* S.I. Smith, 1879
- P. pilosimanus pilosimanus* S.I. Smith, 1879  
off S Iceland-off SW Faeroes-off W Ireland-Bay of Biscay
- Galatheidae Samouelle, 1819
- Galathea* J.C. Fabricius, 1793
- G. intermedia* Liljeborg, 1851  
Oresund-Bohuslän-North Sea-Lofoten-Shetland Faeroe area-Br. Isles-Bay of Biscay
- G. strigosa* (Linnaeus, 1761)  
Kattegat-Bohuslän-North Sea-Trøndelag-Shetland Faeroe area-W Br. Isles-Engl. Channel-Bay of Biscay
- G. squamifera* Leach, 1814  
Oresund-Bohuslän-North Sea-Trøndelag-W Br. Isles-Engl. Channel-Bay of Biscay
- G. nexa* Embleton, 1835  
?Kattegat-Bohuslän-N North Sea-Norw. Sea-?Barents Sea-W Br. Isles-Engl. Channel
- G. dispersa* Bate, 1859  
Bohuslän-N North Sea-Norw. Sea-Shetland Faeroe Iceland area-W Br. Isles-Engl. Channel-Bay of Biscay
- G. machadoi* Barrois, 1888  
off W France
- Munida* Leach, 1820
- M. tenuimana* G.O. Sars, 1872  
?Bohuslän-Skagerrak-Norw. Sea-Barents Sea-Shetland Faeroe Iceland area-W Br. Isles, W Greenland, Iberia, Mediterranean
- M. sarsi* Huus, 1935  
= *M. bamffica* : Auctt. & *M. rugosa* : Auctt.  
Bohuslän-N North Sea-Norw. Sea-Barents Sea-Shetland Faeroe Iceland area - Biscay - N Spain
- M. rugosa* (J.C. Fabricius, 1775)  
= *M. bamffica* (Pennant, 1777)  
Bohuslän-N North Sea-Sognefjord-Shetland Faeroe area-W Br. Isles-Bay of Biscay - Madeira - Mediterranean
- M. intermedia* A. Milne-Edwards & Bouvier, 1899  
E Atlantic north to S part of Porcupine Seabight
- M. microphthalmia* A. Milne-Edwards, 1880  
off S Iceland, Porcupine Seabight, Cape Verde, Mid Atlantic Bight, W Atlantic
- Munidopsis* Whiteaves, 1874  
= *Galathodes* A. Milne-Edwards, 1880
- M. serricornis* (Lovén, 1853)  
= *M. tridentatus* (Esmark, 1857)  
Bohuslän-N Norway-Norw. Sea-W Ireland-Bay of Biscay
- M. curvirostra* Whiteaves, 1874  
off S Iceland-off W Ireland
- Chirostylidae Ortmann, 1892
- Uroptychus* Henderson, 1888
- U. rubrovittatus* (A. Milne-Edwards, 1881)  
off S Iceland-off W Ireland-Bay of Biscay
- Porcellanidae Haworth, 1825
- Pisidia* Leach, 1820
- P. longicornis* (Linnaeus, 1767)  
Oresund-Bohuslän-North Sea-Norw. Sea-Shetland Faeroe area-Br. Isles-Bay of Biscay
- Porcellana* Lamarck, 1801, non Müller, 1766
- P. platycheles* (Pennant, 1777)  
North Sea-Shetland Faeroe area-Br. Isles-Bay of Biscay
- Pomatocheilidae Miers, 1879  
\* Distribution chiefly tropical - subtropical
- Coenobitidae Dana, 1851  
\* Terrestrial with tropical distribution (e.g. *Birgus*)
- Lomisidae Bouvier, 1895  
\* The only known species with E Australian distribution
- Aeglidae Dana, 1852  
\* Freshwater in temperate South America only
- Albunidae Stimpson, 1858  
\* No species within the area?
- Hippidae Latreille, 1825  
\* No species within the area?
- BRACHYURA Latreille, 1803
- DROMIACEA De Haan, 1833
- Superfamily?
- Prosonidae von Meyer, 1860  
\* No species within the area?
- Dynomenidae Ortmann, 1892  
\* Only in tropical seas
- Homolodromidae Alcock, 1895  
Deep water (>600m). No species within the area?
- Dromioidea De Haan, 1833
- Dromiidae De Haan, 1833
- Dromia* Weber, 1795
- D. personata* (Linnaeus, 1758)  
S North Sea-Engl. Channel-Bay of Biscay-Mediterranean
- ARCHAEOBRACHYURA Guinot, 1977
- Homolidea De Haan, 1839
- Homolidae De Haan, 1839
- Paromola* Wood-Mason & Alcock, 1891
- P. cuvieri* (Risso, 1816)  
N North Sea-off W Norway-Orkneys-off W Ireland-Bay of Biscay-Mediterranean
- Tymoloidae Alcock, 1896
- Cymonomidae Bouvier, 1897
- Cymonomus* A. Milne-Edwards, 1880
- C. granulatus* (Thompson, 1873)  
Porcupine & Rockall areas
- C. normani* Lancaster, 1903  
Faeroes, deep water-Bailey & Rosemary Banks (SW of Faeroes)
- Tymolidae Alcock, 1896  
\* No species within the area?
- Latreilliidae Stimpson, 1858  
\* No species within the area?
- Raniniidae de Haan, 1839  
\* No species within the area?
- EUBRACHYURA de St. Laurent, 1980
- HETEROTREMATA Guinot, 1977
- (OXYSTOMATA H. Milne-Edwards, 1837)  
\* Older systematic unit
- Calappoidea De Haan, 1833
- Leucosiidae Samouelle, 1819
- Ebaliinae Stimpson, 1870
- Ebalia* Leach, 1817
- E. tumefacta* (Montagu, 1808)  
Kattegat-Bohuslän-S & W Norway-Shetlands-Br. Isles-Portugal
- E. cranchii* Leach, 1817  
Kattegat-Bohuslän-S & W Norway-Br. Isles
- E. tuberosa* (Pennant, 1777)  
Bohuslän-S & W Norway-Shetlands-Br. Isles-Mediterranean
- E. granulosa* H. Milne-Edwards, 1837  
Scotland-Ireland
- E. nux* A. Milne-Edwards, 1883  
SW Ireland-Porcupine Bank
- Calappidae De Haan, 1833  
\* No species within the area?
- (BRACHYRHYNCHA Borradaile, 1903)  
\* Older systematic unit
- Cancroidea Latreille, 1803

- Corvridae Samouelle, 1819
- Corystes* Bosc, 1802
- C. cassivelaunus* (Pennant, 1777)  
Bohuslän-S Norway-North Sea-Br. Isles-Mediterranean
- Atelecyclidae Ortmann, 1893
- Atelecyclus* Leach, 1814
- A. rotundatus* (Olivier, 1792)  
Bohuslän-S & W Norway-Faeroes-Shetlands-Br. Isles-Mediterranean
- A. undecimdentatus* (Herbst, 1783)  
NW France
- Thiidae Dana, 1852
- Thia* Leach, 1815
- T. scutellata* (Fabricius, 1793)  
Bohuslän-W Norway-S North Sea-Br. Isles-Portugal-Mediterranean
- Pirimelidae Alcock, 1899
- Pirimela* Leach, 1816
- P. denticulata* (Montagu, 1808)  
N Kattegatt-Bohuslän-S & W Norway-Br. Isles-NW Africa-Mediterranean
- Canceridae Latreille, 1803
- Cancer* Linnaeus, 1758
- C. pagurus* Linnaeus, 1758  
Öresund-Bohuslän-North Sea-Troms-Shetlands-Br. Isles-Portugal-Mediterranean
- C. bellianus* Johnson, 1861  
SE Iceland-Shetlands-SW Ireland-Bay of Biscay-Portugal-Azores-Canaries
- Xanthoidea MacLeay, 1838
- Goneplacidae H. Milne-Edwards, 1837
- Goneplacinae H. Milne-Edwards, 1837
- Goneplax* Leach, 1814
- G. rhomboides* (Linnaeus, 1758)  
Ireland-W Br. Isles-Engl. Channel
- Xanthidae MacLeay, 1838
- Pilumnus* Leach, 1815
- P. hirtellus* (Linnaeus, 1761)  
W Norway-S North Sea-Engl. Channel-W Br. Isles-Cape Verde-Mediterranean-Black Sea
- P. spinifer* H. Milne-Edwards, 1834  
\* Sometimes considered to be a subsp. of *P. hirtellus* (Linnaeus)  
Black Sea-Mediterranean-Azores-Mauretania-Portugal. Many specimens were found 1826 in Uddevalla harbour (Bohuslän).
- Pilumnoides* H. Milne-Edwards & Lucas, 1843
- P.inglei* Guinot & Macpherson, 1987  
= *P. perlatus* : Auctt., non (Poeppig, 1836) (This is a South American species)  
Recorded a few times in SW Br. Isles
- Neopanope* A. Milne-Edwards, 1880
- N. sayi* (Smith, 1869)  
North American species introduced into Swansea area
- Monodaeus* Guinot, 1967
- M. couchi* (Couch, 1851)  
Ireland-W Br. Isles-W Engl. Channel
- Xantho* Leach, 1814
- X. pilipes* A. Milne-Edwards, 1867  
N Kattegatt-Bohuslän-W Norway-Shetlands-Br. Isles-SW North Sea-Engl. Channel-Senegal-Mediterranean
- X. incisus* Leach, 1814  
Ireland-W Br. Isles-Engl. Channel-W France-southwards
- Rhithropanopeus* Rathbun, 1898
- R. harrisi* (Gould, 1841)  
E North American species, which has been established in S North Sea and has also been found fr. Poland-Belt Sea-Öresund, Norway, N & W France, Black Sea & Caspian Sea
- Portunoidae Rafinesque-Schmaltz, 1815
- Ceryoniidae Colosi, 1923
- Geryon* Krøyer, 1837
- G. trispinosus* (Herbst, 1803)  
= *G. tridens* Krøyer, 1837 (TM)  
Kattegatt-Bohuslän-S & W Norway-Tromsö-N North Sea-Iceland-off N Scotland-Bay of Biscay-?Morocco-?Mediterranean
- Chaeon* Manning & Holthuis, 1989
- C. affinis* (A. Milne-Edwards & Bouvier, 1894)  
off W Norway-off S Iceland-off Rockall-Azores
- C.inglei* Manning & Holthuis, 1989  
Porcupine Seabight
- Portunidae Rafinesque-Schmaltz, 1815
- Polybiinae Ortmann, 1893
- Polybius* Leach, 1820
- P. henslowii* Leach, 1820  
Skagerrak-N North Sea-Engl. Channel-Morocco-Mediterranean
- Bathynectes* Stimpson, 1870
- B. longipes* (Risso, 1816)  
W Br. Isles-Engl. Channel-NW France
- B. maravigna* (Prestandrea, 1839)  
= *Portunus superbus* Costa, 1853  
off W Norway (Storeggen)-Faeroes-Shetlands-Bay of Biscay-Angola-Mediterranean
- Macropipus* Prestandrea, 1833
- M. tuberculatus* (Roux, 1830)  
= *Portunus tuberculatus* Roux, 1830 (TM)  
W Norway (Stavanger-off Storeggen)-Shetlands-W Br. Isles-Engl. Channel-Bay of Biscay-Azores-Mediterranean
- Necora* Holthuis, 1987
- N. puber* (Linnaeus, 1767)  
= *Cancer puber* Linnaeus, 1767 (TM)  
W Norway-S North Sea-Br. Isles-NW Africa-Mediterranean
- Liocarcinus* Stimpson, 1870
- L. arcuatus* (Leach, 1814)  
Öresund-Bohuslän-S & W Norway-North Sea-Br. Isles-Mediterranean-Black Sea
- L. corrugatus* (Pennant, 1777)  
?Yorkshire-W Br. Isles-Engl. Channel-NW France
- L. zariquieyi* (Gordon, 1968)  
??Shetlands-SW Br. Isles-Mediterranean
- L. pusillus* (Leach, 1815)  
Öresund-Bohuslän-S & W Norway-Lofoten-North Sea-Shetlands-Br. Isles-NW Africa-Mediterranean
- L. depurator* (Linnaeus, 1758)  
Öresund-Bohuslän-North Sea-Lofoten-Shetlands-Br. Isles-Nw. Africa-Mediterranean
- L. holsatus* (Fabricius, 1798)  
= *Portunus holsatus* Fabricius, 1798 (TM)  
Öresund-Bohuslän-North Sea-N Norway-Iceland-Faeroes-Shetlands-Br. Isles-Portugal-Canaries
- L. marmoreus* (Leach, 1814)  
Helgoland-Netherlands-Br. Isles-S Portugal-Azores
- Carcininae MacLeay, 1838
- Carcinus* Leach, 1814
- C. maenas* (Linnaeus, 1758)  
= *Cancer maenas* Linnaeus, 1758  
S Baltic-Bohuslän-Troms-Iceland-Faeroes-Shetlands-Br. Isles-S Portugal-Mauritania
- Xaiva* MacLeay, 1838
- X. biguttata* (Risso, 1816)  
Engl. Channel
- Portumnus* Leach, 1814
- P. latipes* (Pennant, 1777)  
German North Sea coast-Engl. Channel-Br. Isles-Morocco-Mediterranean
- Portuninae Rafinesque-Schmaltz, 1815
- Callinectes* Stimpson, 1860
- C. sapidus* Rathbun, 1896  
E American (Nova Scotia - Uruguay) species, which has been established in E Mediterranean and a few specimens has been found in Öresund, S North Sea & W France
- Pinnotheroidea De Haan, 1833
- Pinnotheridae De Haan, 1833
- Pinnotherinae De Haan, 1833
- Pinnotheres* Bosc, 1802
- P. pisum* (Linnaeus, 1767)  
Öresund-Bohuslän-S Norway-North Sea-Br. Isles-Mauretania-Mediterranean
- P. pinnotheres* (Linnaeus, 1758)  
Yorkshire-Engl. Channel-W Br. Isles-W France-?southwards
- Asthenognathinae Stimpson, 1858
- Asthenognathus* Stimpson, 1858
- A. atlanticus* Monod, 1933  
NW France
- (OXYRHYNCHA Latreille, 1803)  
\* Older systematic unit
- Majidae Samouelle, 1819
- Majinae Samouelle, 1819
- Maja* Lamarck, 1801
- M. squinado* (Herbst, 1788)  
Helgoland-Engl. Channel-W Br. Isles-Guinea-Mediterranean
- Oregoninae Garth, 1958
- Hyas* Leach, 1814
- H. araneus* (Linnaeus, 1758)  
Öresund & Belt Sea-Bohuslän-North Sea-N Norway-Spitsbergen-Iceland-Faeroes-Shetlands-Br. Isles-N Engl. Channel
- H. coarctatus* Leach, 1815  
Öresund & Belt Sea-Bohuslän-North Sea-N Norway-Spitsbergen-Barents Sea-Iceland-Faeroes-Shetlands-Br. Isles-Engl. Channel-Bay of Biscay
- Inachinae MacLeay, 1838
- Achaeus* Leach, 1817

- A. cranchii* Leach, 1817  
Belgium-Engl. Channel-W Br. Isles-Azores-Mediterranean
- Inachus* Weber, 1795
- I. dorsettensis* (Pennant, 1777)  
Öresund & Belt Sea-Bohuslän-North Sea-North Tröndelag-Shetlands-Br. Isles-S Africa-Mediterranean
- I. phalangium* (Fabricius, 1775)  
= *Inachus dorynchus* Leach, 1815  
N Kattegat-Bohuslän-S North Sea-W Norway-Shetlands-Br. Isles-Cape Verde-Mediterranean
- I. leptochirus* Leach, 1817  
off W Jutland-Shetlands-off W Faeroes-Br. Isles-W Africa-Mediterranean
- Dorhynchus* Thomson, 1873
- D. thomsoni* Thomson, 1873  
W Norway-off W Shetlands-S Iceland-between Faeroes & Scotland-Cape Verde-Mediterranean
- Macropodia* Leach, 1814
- M. rostrata* (Linnaeus, 1761)  
Öresund & Belt Sea-Bohuslän-North Sea-Norway (65° N)-Shetlands-Br. Isles-Azores-S Africa-Mediterranean
- M. parva* Van Noort & Adema, 1985  
Bohuslän, Oslofjord, S North Sea, S Portugal
- M. linaresi* Forest & Zariquiey Alvarez, 1964  
SW North Sea, Engl. Channel-NW France, Mediterranean
- M. deflexa* Forest, 1978  
Ireland-W Br. Isles-Engl. Channel-NW & SW France
- M. tenuirostris* (Leach, 1814)  
Netherlands-Br. Isles-Shetlands-off W Faeroes-Engl. Channel-Portugal
- Pisinae* Dana, 1853
- Pisa* Leach, 1814
- P. armata* (Latreille, 1803)  
Belgium-Engl. Channel-W Br. Isles-Angola-Mediterranean
- P. tetraodon* (Pennant, 1777)  
Thames estuary-Engl. Channel-W Br. Isles-Ireland-W France-southwards
- Eurynome* Leach, 1814
- E. aspera* (Pennant, 1777)  
Öresund-Bohuslän-S & W Norway-?Shetlands-Br. Isles-NW Africa-Mediterranean
- E. spinosa* Hailstone, 1835  
Bohuslän-S & W Norway-W Br. Isles-Engl. Channel-NW Spain-Mediterranean
- Rochinia* A. Milne-Edwards, 1875
- R. carpenteri* (Thomson, 1873)  
off W Norway-between Norway & Shetlands-off S Iceland-between Faeroes & Scotland-off Ireland-Bay of Biscay-NW Africa
- Parthenopidae* MacLeay, 1838
- Parthenopinae* MacLeay, 1838
- Parthenope* Weber, 1795
- P. massena* (Roux, 1830)  
NW France
- Mimilambridae* Williams, 1979  
\* The only known recent species collected in West Indies (Tobago)
- Bellidae* Dana, 1852  
\* S South American & New Zealand distribution
- Bythograeidae* Williams, 1980  
\* The only known species is collected from vicinity of thermal vents on the Galapagos Rift
- Deckeniidae* Bott, 1970  
\* Freshwater crabs from E Africa & the Seychelles
- Platyxanthidae* Guinot, 1977  
\* Occurs only in S South America
- Potamidae* Ortmann, 1896  
\* Eurasian freshwater crabs
- Retroplumidae* Gill, 1894  
\* Strictly Indian Ocean
- THORACOTREMATA Guinot, 1977
- Grapsoidea* MacLeay, 1838
- Grapsidae* MacLeay, 1838
- Grapsinae* MacLeay, 1838
- Planes* Bowdich, 1825
- P. minutus* (Linnaeus, 1758)  
Sargasso Sea-(occasionally) S North Sea-Engl. Channel-Mediterranean
- Pachygrapsus* Randall, 1840
- P. marmoratus* (Fabricius, 1787)  
NW France-southwards
- P. transversus* (Gibbes, 1850)  
warm Atlantic-Mediterranean-(once on a ship) Copenhagen
- Varuninae* H. Milne-Edwards, 1853
- Eriocheir* De Haan, 1835
- E. sinensis* H. Milne-Edwards, 1854  
River mouths in Bothnian Bay-S Baltic-Göteborg-S North Sea-Thames-W France
- Brachyotus* De Haan, 1833
- B. sexdentatus* (Risso, 1826)
- W Br. Isles-Mediterranean
- Hemigrapsus* Dana, 1851
- H. penicillatus* (de Haan, 1835)  
\* Should possibly be transferred to genus *Brachyotus*.  
Shore-living invader from NW Pacific, which first was observed in La Rochelle, Biscay coast of France in March 1994. It is spreading rapidly (probably >100 km/year) both southwards and northwards and will possibly reach Scandinavian shores within a few years.
- Gecarcinidae* MacLeay, 1838  
\* Tropical American, W African & Indo-Pacific terrestrial crabs
- Gecarcinucidae* Rathbun, 1904  
\* Old world freshwater crabs
- Hapalocarcinidae* Calman, 1900  
\* Distribution like reef corals, in which they produce "galls"
- Hexapodidae* Miers, 1886  
\* No species within the area?
- Hymenosomatidae* MacLeay, 1838  
\* No species within the area?
- Isolapotamidae* Bott, 1970  
\* Freshwater crabs from Indonesia, Philippines & Coastal Asia
- Ocypodidae* Rafinesque-Schmaltz, 1815  
\* Predominantly warm & tropical areas, e.g. *Uca*
- Palciidae* Rathbun, 1898  
\* No species within the area?
- Parathelphusidae* Alcock, 1910  
\* Large, old world freshwater crabs
- Potamocarcinidae* Ortmann, 1899  
\* New world freshwater crabs
- Potamonautidae* Bott, 1920  
\* African freshwater crabs
- Pseudothelphusidae* Ortmann, 1893  
\* New world freshwater crabs
- Sinopotamidae* Bott, 1970  
\* Freshwater crabs from China & Burma
- Sundathelphusidae* Bott, 1969  
\* Old world freshwater crabs
- Trichodactylidae* Milne-Edwards, 1853  
\* New world freshwater crabs
- MYSIDA* Boas, 1883
- Petalophtalmidae* Willemoës-Suhm, 1875
- Hansenomysis* Stebbing, 1893
- H. fyllae* (Hansen, 1887)  
Faeroe Channel-SW Iceland-off W Ireland
- Petalophtalmus* Willemoës-Suhm, 1875
- P. armiger* Willemoës-Suhm, 1875  
off W Ireland-off W Africa
- Mysidae* Latreille, 1803
- Boreomysinae* Holt & Tattersall, 1905
- Boreomysis* G.O. Sars, 1869
- B. tridens* G.O. Sars, 1870  
Skagerrak-W Norway-Malangen-Faeroes-Iceland-W Br. Isles-Bay of Biscay
- B. arctica* (Krøyer, 1861)  
Koster Channel-Lofoten-Arctic Seas-Faeroe Channel-W Br. Isles-Mediterranean
- B. microps* G.O. Sars, 1883  
Arctic Seas-Iceland-Faeroe Channel-off Ireland-Azores
- B. megalops* G.O. Sars, 1872  
Hardangerfjord-Finmark-off Scotland & W Ireland, ?Mediterranean
- B. scyphops* G.O. Sars, 1879  
\* Not listed by Mauchline, 1980. Synonymized?  
Arctic Seas-Faeroe-Iceland area
- B. nobilis* G.O. Sars, 1885  
Arctic Seas-Faeroe-Iceland area
- B. semicaeca* Hansen, 1905  
Bay of Biscay
- Siriellinae* Norman
- Siriella* Dana, 1850
- S. armata* (Milne-Edwards, 1837)  
Helgoland-Br. Isles-Mediterranean
- S. norvegica* G.O. Sars, 1869  
Kattegat-Bohuslän-Hardangerfjord-Cape Stadt-Br. Isles-Mediterranean
- S. clausii* G.O. Sars, 1876  
Holland-W Scotland-Engl. Channel-Mediterranean
- S. jaltensis* Czerniavsky, 1868  
E & W Scotland-Engl. Channel-Black Sea & W Africa
- S. jaltensis brooki* Norman, 1887  
= *S. gordonae* Zimmer, 1932  
W Scotland & SW North Sea
- Gastrosaccinae* Norman
- Gastrosaccus* Norman, 1868
- G. sanctus* (van Beneden, 1861)

?Skagerrak, Nordland-Finmark, W Denmark-SW North Sea-S & W Br. Isles-W Africa-Black Sea

*G. spinifer* (Goës, 1864)  
S Baltic-Bohuslän-SW Norway-North Sea-Br. Isles-W Africa-Black Sea

*G. normani* G.O. Sars, 1877  
Engl. Channel-Bay of Biscay, Ireland, ?SW North Sea, W Br. Isles

*G. lobatus* Nouvel, 1951  
Engl. Channel-Bay of Biscay, ?SW North Sea, ?W Br. Isles

*Anchialina* Norman & Scott, 1906  
\* Correct subfamily?

*A. agilis* (G.O. Sars, 1877)  
W Scotland & Holland-Engl. Channel-Mediterranean

*Mysinae* Latreille, 1803

Erythropini

*Erythroops* G.O. Sars, 1869

*E. erythropthalma* (Goës, 1864)  
= *E. goesi* (G.O. Sars, 1866)  
Öresund-Bohuslän-all Norway-Spitsbergen-E Scotland-Ireland-W England, N Greenland

*E. abyssorum* G.O. Sars, 1869  
Skagerrak-all Norway-Arctic Seas-Faeroe Channel, N Greenland

*E. serrata* (G.O. Sars, 1863)  
Bohuslän-Porsangerfjord-E Scotland-Ireland-Isle of Man-Iceland-?Arctic Seas

*E. elegans* (G.O. Sars, 1863)  
Öresund-Skagerrak-Trondheimsfjord-North Sea-Br. Isles-Mediterranean

*E. microps* (G.O. Sars, 1864)  
Skagerrak-Lofoten & Arctic Seas, W Ireland

*E. glacialis* G.O. Sars, 1885  
Between Norway & Iceland-Arctic Seas

*Parerythroops* G.O. Sars, 1869

*P. spectabilis* G.O. Sars, 1885  
W-N Norway-Arctic Seas-between Norway & Iceland

*P. abyssicola* G.O. Sars, 1876  
\* Not listed in Mauchline, 1980. Synonymized?  
W-N Norway

*P. obesa* (G.O. Sars, 1864)  
Koster Channel-all Norway-Iceland-W Ireland-?Mediterranean

*Metererythroops* S.I. Smith, 1879

*M. picta* Holt & Tattersall, 1905  
E North Sea-Br. Isles-Iceland & Greenland-Bay of Biscay

*M. robusta* S.I. Smith, 1879  
Bodö-N Norway-Iceland-Arctic Seas

*Katerythroops* Holt & Tattersall, 1905

*K. oceanae* Holt & Tattersall, 1905  
W Ireland-southwards in Atlantic

*Hypererythroops* Holt & Tattersall, 1905

*H. serriventer* Holt & Tattersall, 1905  
W Ireland

*Dactylerythroops* Holt & Tattersall, 1905

*D. dactylops* Holt & Tattersall, 1905  
Faeroe Channel-off W & SW Ireland

*D. bidigitata* W. Tattersall, 1907  
off W Ireland

*D. gracilura* W. Tattersall, 1907  
off W Ireland

*Pseudomma* G.O. Sars, 1870

*P. affine* G.O. Sars, 1870  
Koster Area, Skagerrak-Lofoten-Greenland-Iceland-Faeroes-Ireland-Bay of Biscay

*P. roseum* G.O. Sars, 1870  
Kostersäcken-Skagerrak-Malangen-Iceland-Faeroes

*P. truncatum* S.I. Smith, 1879  
off Cape Stadt, off N Faeroes, Lyngenfjord-Arctic Seas & Iceland

*P. frigidum* Hansen, 1908  
off Cape Stadt, off E & N Faeroes, Arctic Seas

*P. nanum* Holt & Tattersall, 1906  
Faeroe Channel-off W & SW Ireland

*P. jasi* Melan & Brattegard, 1995  
off SW Faeroes in 1319 m.

*Parapseudomma* Nouvel & Lagardère, 1976

*P. calloplura* (Holt & Tattersall, 1905)  
Faeroe Channel-off W & SW Ireland, Mediterranean

*Michthyops* W. Tattersall, 1911

*M. parva* (Vanhöffen, 1897)  
W Greenland-off W Ireland

*M. theeli* (Ohlin, 1901)  
Arctic Seas

*Amblyops* G.O. Sars, 1869

*A. abbreviata* (G.O. Sars, 1869)  
Koster Channel-Skagerrak-all Norway-Arctic Seas-Iceland-Faeroes-Ireland-Bay of Biscay

*A. kempii* (Holt & Tattersall, 1905)

off Faeroes & Hebrides- off W Ireland

*A. tenuicauda* W. Tattersall, 1911  
Faeroe Channel-off SW Ireland

*Paramblyops* Holt & Tattersall, 1905

*P. rostrata* Holt & Tattersall, 1905  
S Iceland-Faeroes-W & SW Ireland-Mediterranean

*P. bidigitata* W. Tattersall, 1911  
off SW Ireland

*Dactylamblyops* Holt & Tattersall, 1906

*D. thaumatops* W. Tattersall, 1907  
Faeroe Channel-off SW Ireland

*D. goniops* W. Tattersall, 1907  
Faeroe Channel-off SW Ireland

*D. sarsi* (Ohlin, 1901)  
Arctic Seas

*Metamblyops* W. Tattersall, 1907

*M. oculata* W. Tattersall, 1907  
Faeroe Channel-off SW Ireland

*Longithorax* Illig, 1906

*L. fuscus* Hansen, 1908  
off S Iceland-off SW Ireland-Bay of Biscay

*Euchaetomera* G.O. Sars, 1883

*E. tenuis* G.O. Sars, 1883  
off SW Ireland-Eur. Atlantic-Mediterranean

*Chunomysis* Holt & Tattersall, 1905

*C. diadema* Holt & Tattersall, 1905  
Faeroe Channel-off SW Ireland

*Arachnomysis* Chun, 1887

*A. leuckartii* Chun, 1887  
off-SW Ireland-Eur. & Afr. Atlantic & Mediterranean

Leptomysini Czerniavsky, 1882

*Leptomysis* G.O. Sars, 1869

*L. gracilis* (G.O. Sars, 1864)  
Öresund-Bohuslän-S Norway-Hardangerfjord-North Sea-Shetlands-Br. Isles-Mediterranean

*L. lingvura* (G.O. Sars, 1866)  
Farsund-Molde-Br. Isles-Black Sea

*L. mediterranea* G.O. Sars, 1877  
SW North Sea & Ireland-Suez & Adriatic Sea

*Bathymysis* W. Tattersall, 1907

*B. helgae* W. Tattersall, 1907  
Faeroe Channel-off W Ireland

*Mysidetes* Holt & Tattersall, 1906

*M. farrani* (Holt & Tattersall, 1905)  
deep Skagerrak, off W Ireland-Bay of Biscay-Mediterranean

*Mysideis* G.O. Sars, 1869

*M. insignis* (G.O. Sars, 1864)  
Koster Channel-W Finmark-Iceland-Faeroes-Br. Isles-Bay of Biscay

*Mysidopsis* G. O. Sars, 1864

*M. gibbosa* G.O. Sars, 1864  
Öresund-Bohuslän-North Sea-Ålesund-Br. Isles-Mediterran.

*M. angusta* G.O. Sars, 1864  
Öresund-Skagerrak-North Sea-Ålesund-Br. Isles-Mediterran.

*M. didelphys* (Norman, 1863)  
Öresund-Skagerrak-Lofoten-Iceland-Br. Isles-W France

*Pseudomysis* G.O. Sars, 1879

*P. abyssi* G.O. Sars, 1879  
Arctic Seas & between Norway & Iceland

*Limysis* Nouvel, 1966

*L. atlantica* (Nouvel, 1942)  
= *Tenagomysis atlantica* Nouvel, 1942  
Bay of Biscay

Mysini Dana, 1850

*Hemimysis* G.O. Sars, 1869

*H. anomala* G.O. Sars, 1907  
\* inhabitant of sublittoral crevices from the Caspian area. Observed in the Baltic from 1992.  
Gulf of Finland

*H. lamornae* (R.Q. Couch, 1856)  
= *Mysis lamornae* R.Q. Couch, 1856  
Kattegatt-Bohuslän-North Sea-Lofoten-Iceland-Br. Isles-Black Sea

*H. abyssicola* G.O. Sars, 1869  
Koster Channel, Skagerrak-W Norway (Tranöydyppet)

*Mysis* Latreille, 1802

*M. relicta* Lovén, 1862  
Limnic in Lakes of Sw., No., Fi., De. etc., Bothn. Bay-Baltic (N of 58°), Arctic Seas, e.g. E & N Greenland

*M. oculata* (Fabricius, 1780)  
?Bothnian Sea-?C. Baltic, Lofoten-Arctic Seas & Iceland

- M. litoralis* (Banner, 1948)  
Troms (Rossfjordvatnet), Arctic Seas-Iceland
- Michtheimysis* Norman, 1902
- M. mixta* (Liljeborg, 1852)  
= *Mysis mixta* Liljeborg, 1852  
Bothnian Sea-Bohuslän, Trondheimsfjord-Norw. Sea-Arctic Seas & Iceland
- Paramysis* Czerniavsky, 1882
- P. (Paramysis) helleri* (G.O. Sars, 1877)  
\* Strictly Mediterranean according to Labat, 1953. It has been confused with *P. noveli* Labat, 1953  
?W Scotland & ?E England-?Black Sea
- P. (Paramysis) arenosa* (G.O. Sars, 1877)  
Ireland-W Scotland-Engl. Channel-Mediterranean
- P. (Paramysis) portzicensis* Nouvel, 1950  
Bay of Biscay
- P. (Paramysis) noveli* Labat, 1953  
Roscoff, Devon
- P. lacustris* (Czerniavsky, 1882)  
= *Mesomysis kowalevskiy* ? 1977?  
Caspian Sea, Baltic (introduced via Newman River in Lithuania, where it was used as fish food)
- P. (Pseudoparamysis* Bacesco, 1940?) *bacescoi* Labat, 1953  
Roscoff, Brest, Finistère
- Schistomysis* Norman, 1892
- S. spiritus* (Norman, 1860)  
?S Baltic-Oresund-Bohuslän-North Sea-North Trøndelag-Faeroes-Br. Isles-W France-?Bay of Biscay
- S. ornata* (G.O. Sars, 1864)  
S Baltic-Bohuslän-Lofoten-North Sea-Faeroes-Iceland-Br. Isles-W Africa
- S. kervillei* (G.O. Sars, 1885)  
Helgoland-Holland-S & W England-Ireland
- S. parkeri* Norman, 1892  
S England & W Morocco
- Praunus* Leach, 1814
- P. inermis* (Rathke, 1843)  
Åland-Bohuslän-all Norway-White Sea-Br. Isles-N Brittany
- P. flexuosus* (O.F. Müller, 1776)  
Bothnian Sea-Bohuslän-Foldenfjord-?Arctic Seas-Br. Isles-Bay of Biscay
- P. neglectus* (G.O. Sars, 1869)  
Öresund-Bohuslän-Lofoten-Br. Isles-N Brittany
- Mesopodopsis* Czerniavsky, 1882
- M. slabberi* (van Beneden, 1861)  
Kiel area-Oresund-Bohuslän-S Norway-Br. Isles-Black Sea
- Neomysis* Czerniavsky, 1882
- N. integer* (Leach, 1814)  
= *Mysis vulgaris* Thompson, 1828  
Bothnian Bay-Bohuslän-Nordland, ?White Sea, Br. Isles-Spain
- Acanthomysis* Czerniavsky, 1882
- A. longicornis* (Milne-Edwards, 1837)  
Ireland-S & W England & SW North Sea-Mediterranean
- Stilomysis* Norman, 1892
- S grandis* (Goës, 1863)  
Troms-Finmark-Iceland-Arctic Seas
- Heteromysini Norman, 1892
- Heteromysis* S.I. Smith, in Verrill, 1874
- H. formosa* S.I. Smith, in Verrill, 1874  
= *H. norvegica* G.O. Sars, 1882  
Bergen, E & W Scotland-Engl. Channel
- H. microps* (G.O. Sars, 1877)  
Engl. Channel, Tunisian Gulf
- H. armoricana* Nouvel, 1940  
Engl. Channel
- Mysidellinae* Czerniavsky, 1882-83
- Mysidella* G.O. Sars, 1872
- M. typica* G.O. Sars, 1872  
W Norway-Ålesund, off SW Ireland
- M. typhlops* G.O. Sars, 1872  
W Norway-Florø
- Lepidomysidae Fage, 1924  
\* In fresh & brackish water caves or land crab burrows. No species within the area.
- Strygiomysidae Caroli, 1937  
\* Cave inhabitants from West Indies & S Italy
- LOPHOGASTRIDA Boas, 1883  
\* Was formerly treated as part of "Mysidacea"
- Eucopiidae* Dana, 1852
- Eucopia* Dana, 1852
- E. unguiculata* (Willemoes-Suhm, 1875)  
= *E. hanseni* Nouvel, 1942  
Faeroe Channel-European Atlantic & Mediterranean
- E. grimaldii* Nouvel, 1942  
off W Iceland-off W Ireland & European Atlantic
- E. sculpticauda* Faxon, 1893  
off W Hebrides-European Atlantic
- Lophogastridae* M. Sars, 1857
- Gnathopausia* Willemoes-Suhm, 1875
- G. zoea* Willemoes-Suhm, 1875  
off Norway (below 650 m), off W Iceland-Faeroe Channel-European Atlantic
- G. gigas* Willemoes-Suhm, 1875  
Greenland-off W Br. Isles & Ireland
- Lophogaster* M. Sars, 1857
- L. typicus* M. Sars, 1857  
Koster Channel-Skagerrak-Trondheimsfjorden-Br. Isles-Mediterranean
- MICTACEA Bowman, Garner, Hessler, Iliffe & Sanders, 1985  
\* no species in the area
- EDRIOPHTHALMA Leach, 1815
- ISOPODA Latreille, 1817
- GNATHIIDEA Hansen, 1916
- Gnathiidae Harger, 1880
- Akidognathia* Stebbing, 1912
- A. oedipus* Stebbing, 1913  
Rockall Bank
- Paragnathia* Omer-Cooper & Omer-Cooper, 1916
- P. formica* (Hesse, 1864)  
= *Anceus formica* Hesse, 1864  
Ireland, W & E England, Holland
- Gnathia* Leach, 1814
- G. abyssorum* (G.O. Sars, 1872)  
= *Anceus abyssorum* G.O. Sars, 1872  
@ *Stryphnus ponderosus*, *Tragosia infundibuliformis*, *Axinella arctica*, *Phakellia robusta*, *Stelletta normani*  
Faeroes-Norw. Sea-Iceland & Arctic Seas
- G. albescens* Hansen, 1916  
off SW Faeroes
- G. arctica* Gurjanova, 1929  
Novaja Zemlja
- G. bicolor* Hansen, 1916  
Norw. Sea-off S Iceland-Arctic Seas
- G. cristatipes* Stebbing, 1913  
deep W Engl. Channel
- G. dentata* (G.O. Sars, 1872)  
= *Anceus dentatus* G.O. Sars, 1872  
Kattegatt-Bohuslän-Trondheim, Holland
- G. elongata* (Krøyer, 1849)  
= *Anceus elongatus* Krøyer, 1849  
Lofoten-N Norway-Iceland & Arctic Seas, e.g. N Greenland
- G. hirsuta* (G.O. Sars, 1877)  
= *Anceus hirsutus* G.O. Sars, 1877  
Faeroes-Norw. Sea-Iceland & Arctic Seas
- G. maxillaris* (Montagu, 1804)  
= *Cancer maxillaris* Montagu, 1804  
Ireland-W & S Br. Isles & S North Sea
- G. oxyuraea* (Liljeborg, 1855)  
Öresund-Oslofjord-Lofoten-Br. Isles
- G. robusta* (G.O. Sars, 1879)  
Faeroes-Norw. Sea-Iceland & Arctic Seas
- G. schistifrons* Stebbing, 1912  
Ireland in 380 m
- G. stygia* (G.O. Sars, 1877)  
= *Anceus stygius* G.O. Sars, 1877  
Faeroes-Norw. Sea-Iceland & Arctic Seas
- G. vorax* (Lucas, 1849)  
Greenland-Br. Isles-Mediterranean
- ANTHURIDEA Leach, 1814
- Antheluridae* Poore & Lew Ton, 1988
- Anthelura* Norman & Stebbing, 1886
- A. elongata* Norman & Stebbing, 1886 (TM)  
NE Atlantic in 1100-1590 m
- A. truncata* (Hansen, 1916)  
N Atlantic
- Anathura* Barnard, 1925
- A. sulcatauda* Barnard, 1925  
off SW & NE Iceland in 216-1505 m
- Anthuridae* Leach, 1814
- Anthura* Leach, 1814
- A. gracilis* (Montagu, 1808)  
= *Oniscus gracilis* Montagu, 1808 (TM)  
S Baltic-North Sea-S & W Br. Isles-Mediterranean
- Cyathura* Norman & Stebbing, 1886
- C. carinata* (Krøyer, 1847)  
= *Anthura carinata* Krøyer, 1847  
S Baltic(Karlskrona)-Kungsbacka, North Sea-England-Morocco, Greenland

*Haliophasma* Haswell,1881

*H. mjoelniri* Negoescu & Svavarson,1997  
off SW Iceland in 340-508 m

*Quantanthurus* Menzies & George,1972

*Q. tyri* Negoescu & Svavarson,1997  
off SW Iceland in 304-1212 m

*Paranthuridae* Menzies & Glynn,1968

*Calathura* Norman & Stebbing,1886

*C. brachiata* (Stimpson,1853)  
= *Anthura brachiata* Stimpson,1853 (™ - Bay of Fundy)  
= *Paranthura norvegica* G.O. Sars,1873  
Outer Skagerrak-W & N Norway-Faeroes-off Iceland in 19-1392 m-Arctic Seas-N Greenland-Shetlands-W Br. Isles-off W Bretagne

*Leptanthura* G.O. Sars,1897

*L. tenuis* (G.O. Sars,1873)  
= *Paranthura tenuis* G.O. Sars,1873  
Skagerrak-Trondheim & Portugal

*L. affinis* (Bonnier,1896)  
off SE, S & SW Iceland in 501-1105 m

*L. chardy* Negoescu,1992  
off SW & W Iceland in 304-1295 m

*L. micrura* Kensley,1982  
off W Iceland in 1295 m, off SE Biscay in 1894-1950 m

*Paranthura* Bate & Westwood,1866

*P. costana* Bate & Westwood,1866 (™)  
Channel Islands

*Hyssuridae* Wägele,1981  
\* No species within the area.

VALVIFERA G.O. Sars,1882

*Chaetiliidae* Dana,1853

*Chiridotea* Harger,1878

*C. caeca* (Say,1818)  
= *Idotea caeca* Say,1818  
NE Atlantic

*C. tuftsii* (Stimpson,1853)  
= *Idotea tuftsii* Stimpson,1853  
NE Atlantic

*Saduria* Adams,1852  
= *Idotea* Lockington,1877  
= *Mesidotea* Richardson,1905  
= *Glyptonotus* Eights,1853 (p.p.)

*S. entomon* (Linnaeus,1758)  
= *Oniscus entomon* Linnaeus,1758  
Bothnian Bay-Öresund

*S. megalura* (G.O. Sars,1879)  
Faeroes-Norw. Sea

*S. sabini* (Krøyer,1849)  
= *Idotea sabini* Krøyer,1849  
NE Atlantic

*S. sibirica* (Birula,1896)  
= *Glyptonotus sibirica* Birula,1896  
Siberia

*Idoteidae* Samouelle,1819

*Idotea* J.C. Fabricius,1798  
= *Idothea* J.C. Fabricius,1799

*I. linearis* (Linnaeus,1766)  
= *Oniscus linearis* Linnaeus,1766  
Kattegat-North Sea-Br. Isles-Canaries-Mediterranean

*I. emarginata* (J.C. Fabricius,1793)  
= *Cymothoa emarginata* J.C. Fabricius,1793  
S Baltic-Bohuslän-North Sea-N Norway-Faeroes-Iceland-Br. Isles-W Mediterranean

*I. metallica* Bosc,1802  
Br. Isles (occasionally adrifted fr. America), Holland, Iceland?

*I. balthica* (Pallas,1772)  
= *Oniscus balthicus* Pallas,1772  
S Bothnian Bay-Bohuslän-North Sea-Tromsø-Faeroes-Iceland-Shetlands-Br. Isles-Morocco-Mediterranean-Black Sea

*I. pelagica* Leach,1815  
Öresund-S & W Norway-Faeroes-Iceland-Br. Isles-N & W France

*I. granulosa* Rathke,1843  
Åland-Bohuslän-N Norway-Faeroes-Iceland-Br. Isles-NW France

*I. chelipes* (Pallas,1766)  
= *Oniscus chelipes* Pallas,1766  
= *Oniscus viridis* Slabber,1775  
= *I. angusta* : G.O. Sars,1897  
Åland-Bohuslän-Trøndelag-Faeroes-Br. Isles-Ireland-France-NW Spain

*I. neglecta* G.O. Sars,1897  
= *Idothea neglecta* G.O. Sars,1897  
Öresund-Bohuslän-N Norway-Bear Island-Iceland-Faeroes-Br. Isles-France

*Synidotea* Harger,1878

*S. bathyalis* Gurjanova,1955  
Arctic Sea

*S. nodulosa* (Krøyer,1846)  
= *Idotea nodulosa* Krøyer,1846  
Greenland

*Synisoma* Collinge,1917  
= *Stenosoma* Leach,1815 (p.p.)

*S. lanceifer* (Miers, 1881)  
= *Stenosoma lanceifera* Miers,1881  
= *Idotea appendiculata* : Bate & Westwood,1868  
S & W Br. Isles-W France-Mediterranean

*S. acuminatum* (Leach,1815)  
= *Stenosoma acuminata* Leach,1815  
Ireland, SW Scotland-S England-Mediterranean

*Holognathidae* G. Thomson,1904

*Cleantis* Dana,1849  
= *Zenobiana* Stebbing,1895  
= *Zenobia* Risso,1826, non Oken,1815 (n. nud.), nec Gray,1821

*C. prismatica* (Risso,1826)  
= *Zenobia prismatica* Risso,1826  
Firth of Clyde, S & W Engl., W France- Mediterranean

*Arcturidae* Bate & Westwood,1868

*Astacilla* Cordiner,1793

*A. longicornis* (Sowerby,1806)  
= *Oniscus longicornis* Sowerby,1806  
Öresund-Bohuslän-North Sea-N Norway-Faeroes-Iceland-Br. Isles-Engl. Channel

*A. pusilla* (G.O. Sars,1873)  
= *Arcturus pusilla* G.O. Sars,1873  
Skagerrak-Lofoten-off SW Faeroes

*A. arietina* G.O. Sars,1883  
Hardangerfjord-Norw. Sea-?Davis Strait

*A. intermedia* (Goodsir,1841)  
= *Leachia intermedia* Goodsir,1841  
= *A. affinis* G.O. Sars,1897  
Skagerrak-Lofoten-S Iceland-Scotland-off W & SW Ireland

*A. granulata* (G.O. Sars,1877)  
= *Leachia granulata* G.O. Sars,1877  
off W & N Norway-off W Faeroes-between Faeroes & Iceland

*A. deshaysi* (Norman & Scott,1906)  
\* Does this species exist, or is the name a synonym of another?  
Br. Isles

*Arcturella* G.O. Sars,1897

*A. dilatata* (G.O. Sars,1883)  
= *Astacilla dilatata* G.O. Sars,1883  
Kattegat-Bohuslän-W Norway-Shetlands-Mediterranean

*A. danmoniensis* (Stebbing,1874)  
= *Arcturus danmoniensis* Stebbing,1874  
Devon & Pembrokeshire

*Arcturus* Latreille,1829

*A. scabrosus* Norman,1904  
off SW Faeroes

*A. baffini* (Sabine,1824)  
= *Idothea baffini* Sabine,1824  
Faeroes-Iceland-Davis Strait

*A. tuberosus* G.O. Sars,1885?  
\* Does this species exist? Not in the Smithsonian Internet list.  
Faeroes

*Pleuropion* Zur Strassen,1902

*P. hystrix* (G.O. Sars,1877)  
= *Arcturus hystrix* G.O. Sars,1877  
off Lofoten-Norw. Sea-Faeroes

*P. frigidum* Hansen,1916  
Greenland

*Spectarcturus* Schultz,1981

*S. multispinatus* Schultz,1981 (™ - Greenland)  
Greenland

*Amesopodidae* Stebbing,1905  
\* No species within the area?

*Pseudidotheidae* Ohlin,1901  
\* No species within the area?

*Xenarcturidae* Sheppard,1957  
\* No species within the area?

FLABELLIFERA G.O. Sars,1882

*Limnoriidae* Harger,1879

*Limnoria* Leach,1814

*L. (Limnoria) lignorum* (Rathke,1799)  
= *Cymothoa lignorum* Rathke,1799  
W Baltic-Bohuslän-North Sea-Lofoten-Faeroes-Br. Isles

*L. (Limnoria) tripunctata* Menzies,1951  
S Wales & S England-tropical waters, California

*L. (Limnoria) quadripunctata* Holthuis,1949  
Netherlands & S Br. Isles

*L. borealis* Kussakin,1963  
Iceland, White Sea

*Sphaeromatidae* H. Milne-Edwards,1840

*Bathycopea* Tattersall,1905

*B. typhlops* Tattersall,1905 (™ - Ireland)  
@ ?

off SW Faeroes-off W Ireland

*Sphaeroma* Bosc, 1802

*S. prideauxianum* Leach, 1818  
England

*S. curtum* (Leach, 1818)  
= *Oniscus curtum* Leach, 1818  
England

*S. serratum* (J.C. Fabricius, 1787)  
= *Oniscus serratum* J.C. Fabricius, 1787  
Holland, S & W England, Wales & Ireland-Mediterranean

*Lekanesphaera* Verhoeff, 1943

*L. rugicauda* (Leach, 1814)  
= *Sphaeroma rugicauda* Leach, 1814  
Kalmar Sound-Öresund-N Jutland, Br. Isles-Morocco

*L. hookeri hookeri* (Leach, 1814)  
= *Sphaeroma hookeri* Leach, 1814  
Kalmar Sound-Kattegat

*L. monodi* (Arcangeli, 1934)  
= *Sphaeroma monodi* Arcangeli, 1934  
Belgium, S & SE England, Wales & E Ireland-Spain, Libya

*L. teissieri* (Bocquet & Lejuez, 1967)  
= *Sphaeroma teissieri* Bocquet & Lejuez, 1967  
France

*L. levii* (Argano & Ponticelli, 1981)  
= *Sphaeroma levii* Argano & Ponticelli, 1981  
France

*Dynamene* Leach, 1814

*D. bidentata* (Adams, 1800)  
= *Oniscus bidentata* Adams, 1800 (TM)  
S & W Br. Isles & Ireland-Portugal

*D. edwardsi* (Lucas, 1849)  
= *Naesea edwardsi* Lucas, 1849  
?European Atlantic, Tunisia

*D. magnitorata* Holdich, 1968  
?European Atlantic, S France, NW Spain

*Campeopea* Leach, 1814

*C. hirsuta* (Montagu, 1804)  
= *Oniscus hirsutus* Montagu, 1804  
= *Campeopea cranchii* Leach, 1814 (TM)  
Wales, Cornwall & Devon-W Africa

*Cymodoce* Leach, 1814

*C. truncata* Leach, 1814 (TM)  
S & W Br. Isles-Shetlands, NW Spain

*C. emarginata* Leach, 1818  
S & W Br. Isles-Shetlands

*C. granulata* Miers, 1876  
off Ireland, New Zealand

*Cirolanidae* Dana, 1852

*Eurydice* Leach, 1815

*E. affinis* Hansen, 1905  
Holland, S & W Br. Isles-Mediterranean

*E. caeca* Hansen, 1916  
\* Listed as belonging to a genus incertae sedis in the Smithsonian Internet list.  
off S Iceland

*E. grimaldii* Dollfus, 1888  
Bergen-N Norway-Norw. Sea-Faeroes-Iceland-Br. Isles-Portugal-Azores

*E. inermis* Hansen, 1890  
North Sea, England

*E. pulchra* Leach, 1815  
S Baltic-Bohuslän-Trondheim-Norw. Sea-W France

*E. rotundicauda* Norman, 1906  
NE Atlantic

*E. spinigera* Hansen, 1890  
Netherlands, West Indies

*E. truncata* (Norman, 1868)  
= *Cirolana truncata* Norman, 1868  
North Sea-Norw. Sea

*Faucheria* Dollfus & Vire, 1905

*F. faucheri* (Dollfus & Vire, 1900)  
= *Caecosphaeroma faucheri* Dollfus & Vire, 1900  
France

*Cirolana* Leach, 1818

*C. cranchii* Leach, 1818 (TM - England)  
Firth of Clyde-Plymouth-Mediterranean

*Metacirolana* Nierstrasz, 1931

*M. hanseni* (Bonnier, 1896)  
= *Cirolana hanseni* Bonnier, 1896  
Faeroes-off W Ireland-Bay of Biscay

*Natolana* Bruce, 1981

*N. borealis* (Liljeborg, 1851)  
= *Cirolana borealis* Liljeborg, 1851  
Kieler Bucht-Öresund-Bohuslän-North Sea-Trondheim-off S Lofoten-Faeroes-Shetlands-Br. Isles-W France-Mediterranean, in (5) 35-694 m

*N. gallica* (Hansen, 1905)  
= *Cirolana gallica* Hansen, 1905  
Roscoff, NE Atlantic in 195-210 m

*N. imicola* (Dollfus, 1903)  
= *Cirolana neglecta* var. *imicola* Dollfus, 1903  
= *Cirolana schmidtii* Hansen, 1905  
Faeroes-Davis Strait in 872-2450 m

*Politolana* Bruce, 1981

*P. microphthalma* (Hoek, 1882)  
= *Cirolana microphthalma* Hoek, 1882  
North Sea-Arctic Seas

*Conilera* Leach, 1818

*C. cylindracea* (Montagu, 1804)  
= *Oniscus cylindracea* Montagu, 1804  
British Isles

*Aegidae* Leach, 1815

*Syscenus* Harger, 1880

*S. infelix* Harger, 1880 (TM - Massachusetts)  
= *Harponyx pranzoides* G.O. Sars, 1883  
@ (adult, ecto) *Merlangius merlangus*  
N Kattegatt-Bohuslän-W Norway-Faeroes-Mediterranean

*S. liljeborgii* (Bovallius, 1885)  
= *Rocinela liljeborgii* Bovallius, 1885  
\* Listed in the Smithsonian Internet list as a species of its own, but considered as a synonym of *S. infelix*  
Hargrave, 1880 by Wahrberg.  
Sweden

*S. atlanticus* Kononenko, 1988  
Atlantic

*Rocinela* Leach, 1818

*R. danmoniensis* Leach, 1818 (TM - England)  
@ (facultative, ecto) *Gadidae*  
Öresund-Bohuslän-Trondheim-Faeroes-Iceland-Shetlands-Br. Isles-Bay of Biscay

*R. dumerilii* (Lucas, 1849)  
= *Acherusia dumerilii* Lucas, 1849  
@ ?  
?European Atlantic, Algeria

*Aega* Leach, 1815  
= *Barybrotus* Schiödte & Meinert, 1879 (TM B. indus Schiödte & Meinert, 1879 - India)

*A. (Aega) arctica* Lütken, 1859  
@ sharks  
N Norway-Faeroes, Iceland & Arctic Seas-off W Ireland

*A. (Aega) crenulata* Lütken, 1859  
@ *M. byrkelange* & *Somniosus microcephalus*  
Bohuslän-S, W & N Norway-Faeroes-Iceland-Shetlands-Scotland-W Ireland

*A. (Rhamphion* ? 18???) *gracilipes* Hansen, 1895  
off SW Iceland-off NW Scotland

*A. loveni* Bovallius, 1886  
\* A synonym of *A. ventrosa* M. Sars, according to Wahrberg, but included in the Smithsonian internet list.  
Sweden

*A. (Rhamphion* ? 18???) *microphthalma* Dana, 1853  
\* Probably a synonym of another species, but included in the Smithsonian internet list.  
North Sea (England)

*A. (Aega) monophthalma* Johnston, 1834  
@ *Anarhichas* & *Gadus*  
Bohuslän-Trondheim-Faeroes-Iceland-N Br. Isles

*A. nordenskjoeldi* (Bovallius, 1885)  
= *Aegiochus nordenskjoeldi* Bovallius, 1885  
\* A synonym of *A. ventrosa* M. Sars, according to Wahrberg, but included in the Smithsonian internet list.  
Arctic Ocean

*A. (Aega) psora* (Linnaeus, 1758)  
= *Oniscus psora* Linnaeus, 1758  
@ *Raja*, *Cottus*, *Gadus*, *Molva*, *Hippoglossus*, *Anarhichas* etc.  
Kattegat-Oslofjord-N Norway-Faeroes-off W Iceland-Arctic Seas-Br. Isles

*A. (Rhamphion* ? 18???) *stroemii* Lütken, 1859  
@ different fish species  
Bohuslän-Lofoten-Faeroes-Iceland-N Br. Isles-off SW Ireland

*A. (Aega) tridens* Leach, 1815  
@ *Gadus morhua*  
Oslofjord-Skagerrak-W Norway-Faeroes-Shetlands-Scotland-Belfast

*A. (Rhamphion* ? 18???) *ventrosa* M. Sars, 1859  
@ *G. morhua*, *Brosme*, *M. byrkelange*, *Hippoglossus*  
Bohuslän-N Norway-Faeroes-Iceland-off Shetlands-off W Ireland

*A. (Aega) bicarinata* Leach, 1818  
@ ?  
?European Atlantic

*A. (Aega) rosacea* (Risso, 1816)  
= *Cymothoa rosacea* Risso, 1816  
@ ?  
?European Atlantic, S France

*Xenuraega* Tattersall, 1909

*X. ptilocera* Tattersall, 1909 (TM - off Azores)  
Ireland, S England, W Biscay, Azores

*Cymothoidae* Leach, 1818

*Anilocra* Leach, 1818

*A. physodes* (Linnaeus, 1758)  
= *Oniscus physodes* Linnaeus, 1758  
@ Labridae & other fishes  
Holland & Engl. Channel-Mediterranean

*A. frontalis* H. Milne-Edwards, 1840

?European Atlantic, Mediterranean

**Glossobius** Schiødt & Meinert, 1883

**G. impressus** (Say, 1818)  
= *Cymothoa impressa* Say, 1818  
@ different flying fish species etc.  
N Atlantic, e.g. Køge Bugt (Denmark)

**G. auritus** Bovallius, 1885  
N. Atlantic

**Cymothoa** Fabricius, 1787

**C. scopulorum** (Linnaeus, 1758)  
= *Oniscus scopulorum* Linnaeus, 1758  
Norw. Sea

**Nerocila** Leach, 1818

**N. orbignyi** (Guérin, 1832)  
= *Ichthyophilus orbignyi* Guérin, 1832  
\* Félix Edouard Guérin did not change his last name to Guérin-Méneville until 1836, so citations before that date should use the short name  
?European Atlantic, Greece

**N. maculata** H. Milne-Edwards, 1840  
\* This name is probably now synonymized with another. Which?  
?European Atlantic

**N. neapolitana** Schiødt & Meinert, 1879  
\* This name is probably now synonymized with another. Which?  
?European Atlantic

**Mothocya** Costa, 1851

**M. nana** (Schiødt & Meinert, 1884)  
= *Irona* Schiødt & Meinert, 1884  
*nana* Schiødt & Meinert, 1884  
?European Atlantic, Lesser Antilles

**Anuropodidae** Stebbing, 1893  
\* No species within the area?

**Bathynataliidae** Kensley, 1978  
\* No species within the area?

**Bathynomidae** Wood-Mason & Alcock, 1891  
\* No species within the area?

**Corallanidae** Hansen, 1890  
\* No species within the area.

**Keuphyliidae** Bruce, 1980  
\* No species within the area?

**Phoratosopodidae** Hale, 1925  
\* One S Australian species only known.

**Plakarhriidae** Hansen, 1890  
\* Only two species known, none of them NE Atlantic.

**Protognathiidae** Wägele & Brandt, 1988  
\* Only one Antarctic species known.

**Serolidae** Dana, 1853  
\* No species within the area. Chiefly confined to S hemisphere.

**Tecticipitidae** Bruce, 1993  
\* Pacific family.

**Tridentellidae** Bruce, 1984  
\* No species within the area.

MICROCERBERIDEA Lang, 1961

**Microcerberidae** Karaman, 1933

**Microcerberus** Karaman, 1933

**M. remani** Chappuis, Delamare-Deboutteville & Paulian, 1956  
**lusitanicus** Galhano, 1970  
Portugal

ASELLOTA Latreille, 1803

Familia Incertae sedis.

**Tole** Ortmann, 1901  
= *Iolella* Richardson, 1905

**T. laciniata** (G.O. Sars, 1872)  
= *Janthe laciniata* G.O. Sars, 1872  
off NW & N Norway-off Iceland-Faeroes-Davis Strait

**T. spinosa** (Harger, 1879)  
= *Ianira spinosa* Harger, 1879  
off SE & NW Iceland-Davis Strait

**T. spinosissima** (Stephens, 1936)  
= *Ianira spinosissima* Stephensen, 1935  
Greenland

**T. libbeyi** (Ortmann, 1901)  
= *Jolanthe libbeyi* Ortmann, 1901  
Greenland

**T. vilhelminae** (Stephens, 1913)  
= *Ianira vilhelminae* Stephensen, 1913  
W Greenland

Aselloidea Rafinesque-Schmaltz, 1815

**Asellidae** Rafinesque-Schmaltz, 1815  
\* Only freshwater & groundwater species. *Asellus* Geoffroy St Hilaire, 1764 *aquaticus* (Linnaeus, 1758) is common in all N Europe and some species of *Proasellus* Dudich, 1925 may be found as well.

**Atlantassellidae** Sket, 1980  
\* No species within the area?

**Stenasellidae** Dudich, 1924  
\* Only cave & groundwater species

**Stenetrioidea** Hansen, 1905

**Stenetriidae** Hansen, 1905  
No species within the area? Usually tropical.

**Janiroidea** G.O. Sars, 1899  
= **Paraselloidea**

**Janiridae** G.O. Sars, 1899

**Ianira** Leach, 1814

**I. maculosa** Leach, 1814  
Öresund-Bohuslän-N Norway-Faeroes-Iceland-Greenland-Morocco

**I. alta** (Stimpson, 1853)  
off W Iceland-E North America

**I. tricornis** (Krøyer, in Gaimard, 1846)  
Jan Mayen-Spitsbergen-Davis Strait

**Ianiropsis** G.O. Sars, 1897

**I. breviremis** (G.O. Sars, 1882)  
Öresund-Bohuslän-Bergen-Faeroes-Iceland-Br. Isles

**Jaera** Leach, 1814

**J. albifrons albifrons** Leach, 1814  
Bothnian Bay-France-Norw. Sea-White Sea

**J. albifrons syei** Bocquet, 1950  
Bothnian Bay-Norw. Sea

**J. praehirsuta** Forsman, 1949  
Öresund-Norw. Sea, NW France, Britain, Greenland, New England

**J. posthirsuta** Forsman, 1949  
Britain, New England

**J. ischiosetosa** Forsman, 1949  
Bothnian Bay-Norw. Sea-White Sea, Britain

**J. forsmanni** Bocquet, 1950  
Britanny, S & W Br. Isles

**J. nordmanni** (Rathke, 1837) **nordica** Lemerrier, 1958  
\* in fresh or brackish water  
W Scotland-Mediterranean-Black Sea

**J. hopeana** Costa, 1853  
@ (ectocommensally) *Sphaeroma serratum*  
Devon-Normandy-Mediterranean-Black Sea

**Janiralata** Menzies, 1951 (TM *J. davisii* Menzies, 1951 - California)

**J. pulchra** (Hansen, 1916) Brandt, 1993  
= *Ianthopsis pulchra* (Hansen, 1916) Wolff, 1962  
= *Janira pulchra* Hansen, 1916  
off SW & N Iceland-between Iceland & Greenland-Davis Strait

**Microjaera** Bocquet & Lévi, 1955

**M. anisopoda** Bocquet & Lévi, 1955  
\* Interstitial  
Roscoff, Scilly Isles, Eddystone, W Mediterranean

**Caecijaera** Menzies, 1951

**C. borealis** Kussakin, 1962  
@ *Limnoria* spp. (ectocommensalistic)  
Iceland

**Microparasellidae** Karaman, 1933  
= **Micraniridae**

**Microcharon** Karaman, 1933

**M. teissieri** (Lévi, 1951)  
Roscoff

**M. harrisi** Spooner, 1959  
Plymouth

**M. monnotti** Bocquet, 1970  
Roscoff

**M. coineauae** Galhano, 1970  
Portugal

**Acanthaspidiidae** Menzies, 1962

**Acanthaspidia** Stebbing, 1898

**A. typhlops** (G.O. Sars, 1879)  
off Lofoten-between Iceland & Greenland-Davis Strait

**Ianthopsis** Beddard, 1886

**I. pulchra** (Hansen, 1916)  
= *Janira pulchra* Hansen, 1916  
Greenland

**Janirellidae** Menzies, 1956

**Janirella** Bonnier, 1896

**J. spongicola** Hansen, 1916  
off SW Iceland

**Katianiridae** Svavarsson, 1987

**Katianira** Hansen, 1916

**K. chelifera** Hansen, 1916  
off SW Iceland

**K. biloba** Gurjanova, 1930  
Faeroes, Spitsbergen, E Arctic ocean

**K. cornigera** Gurjanova, 1930  
Spitsbergen

- K. sadko* Gurjanova, 1946  
E Arctic ocean
- Haplonicidae Hansen, 1916
- Haploniciscus* Richardson, 1909
- H. ingolfi* Wolff, 1962  
Faeroes, S of Jan Mayen in 2465 m
- H. bicuspis bicuspis* (G.O. Sars, 1877)  
Faeroes-off W Norway-off Iceland, Faeroes & Jan Mayen
- H. bicuspis tedipus* Wolff, 1962  
SW of Iceland in 1505 m
- H. spinifer* Hansen, 1916  
off SW Iceland-Davis Strait
- Chauliodoniscus* Lincoln, 1985
- C. armadilloides* (Hansen, 1916)  
= *Haploniciscus armadilloides* Hansen, 1916  
off S Iceland
- Joeropsidae Nordenstam, 1933
- Joeropsis* Koehler, 1885
- J. brevicornis brevicornis* Koehler, 1885  
S & W Br. Isles, Norway
- Munnidae G.O. Sars, 1899
- Munna* Krøyer, 1839
- M. acanthifera* Hansen, 1916  
Faeroes-around Iceland-Davis Strait
- M. armoricana* Carton, 1962  
France
- M. boeckii* Krøyer, 1839  
Bohuslän-North Sea-W Norway-Faeroes-Iceland & Arctic Seas-Scotland
- M. coeca* Gurjanova, 1930  
E of Spitsbergen
- M. fabricii* Krøyer, 1846  
non = *M. fabricii*: G.O. Sars, 1899  
Belt Sea-North Sea-W Norway-Iceland & Arctic Seas-Normandie
- M. groenlandica* Hansen, 1916  
W & E Greenland
- M. hanseni* Stappers, 1911  
\* possibly a synonyme of *M. limicola*  
Faeroes-around Iceland-Arctic Seas
- M. kroeyeri* Goodsir, 1842  
Bay of Kiel-S & W Norway-North Sea-Iceland-Br. Isles-Engl. Channel
- M. limicola* G.O. Sars, 1866  
Bohuslän-North Sea-N Norway-Faeroes-Iceland & Arctic Seas-Ireland-Engl. Channel
- M. minuta* Hansen, 1916  
= *M. fabricii*: G.O. Sars, 1899, non Krøyer, 1846  
Öresund-Bohuslän-W & N Norway-Faeroes-Iceland & Arctic Seas-Ireland
- M. palmata* Liljeborg, 1851  
S Norway
- M. pellucida* Gurjanova, 1930  
Murman coast
- M. roemeri* Gurjanova, 1930  
?Spitsbergen-arctic Seas
- M. spitzbergensis* Gurjanova, 1930  
Spitsbergen-arctic Seas
- Uromunna* Menzies, 1962
- U. petiti* (Amar, 1948)  
= *Munna petiti* Amar, 1948  
Scilly Islands-Mediterranean
- Paramunnidae Vanhöffen, 1914
- Paramunna* G.O. Sars, 1866
- P. bilobata* G.O. Sars, 1866 (TM)  
Bohusl-Sognefjord-Br. Isles
- P. typica* (Tattersall, 1905)  
= *Metamunna typica* Tattersall, 1905  
W of Ireland in 220-300 m
- Pleurogonidae Nordenstam, 1933
- Pleurogonium* G.O. Sars, 1882  
= *Pleuracantha* G.O. Sars, 1864 (homonym)
- P. inerme* G.O. Sars, 1882  
Öresund-Bohuslän-North Sea-W Norway-Faeroes-Iceland & Arctic Seas-Br. Isles
- P. rubicundum* (G.O. Sars, 1864)  
= *Pleuracantha rubicunda* G.O. Sars, 1864  
S Baltic-Bohuslän-N Norway-Faeroes-Arctic Seas-N Br. Isles
- P. spinosissimum* (G.O. Sars, 1868)  
= *Pleuracantha spinosissima* G.O. Sars, 1868  
Öresund-Bohuslän-N Norway-Faeroes-Iceland & Arctic Seas-N Br. Isles
- P. intermedium* Hansen, 1916  
Faeroes-N Iceland
- P. pulchrum* Hansen, 1916  
Faeroes-W Iceland
- P. polaris* Svavarson, 1988  
\* Is this a synonym of another species?
- abyssal depths in North Polar Sea
- P. latimanum* Hansen, 1916  
Davis Strait
- Dendroitiidae Vanhöffen, 1914
- Dendromunna* Menzies, 1962 (TM) *D. spinipes* Menzies, 1962 - S Atlantic)
- D. compsa* Lincoln & Boxshall, 1983  
NE Atlantic in 1271-2925 m
- Dendrotion* G.O. Sars, 1872
- D. elegans* Lincoln & Boxshall, 1983  
NE Atlantic in 1600-2200 m
- D. setosum* Lincoln & Boxshall, 1983  
NE Atlantic in 1160-2076 m
- D. spinosum* G.O. Sars, 1872 (TM)  
Skagerrak-N North Sea-Hardangerfjord-Faeroes-Iceland & Arctic Seas
- D. paradoxum* Hansen, 1916  
Faeroes-off SW Iceland
- Acanthomunna* Beddard, 1886  
= *Pseudomunna* Hansen, 1916
- A. hystrix* (Hansen, 1916)  
= *Pseudomunna hystrix* Hansen, 1916 (TM of *Pseudomunna* Hansen, 1916)  
off SW Iceland
- Mictosomatidae Wolff, 1965
- Mictosoma* Wolff, 1965  
= *Schistosoma* Hansen, 1916, non Weinland, 1858, nec Brady, 1877
- M. ramosum* (Hansen, 1916)  
off S Iceland
- Macrostylidae Hansen, 1916
- Macrostylis* G.O. Sars, 1864
- M. longiremis* (Meinert, 1890)  
Bohuslän-Skagerrak-SW Iceland
- M. spinifera* G.O. Sars, 1864  
Kattegatt-Bohuslän-W Norway-Iceland & Arctic Seas-off SW Iceland-Shetlands
- M. subinermis* Hansen, 1916  
Faeroes-off N & E Iceland
- M. elongata* Hansen, 1916  
= *M. longipes* Hansen, 1916 (male)  
off S & E Iceland
- M. abyssicola* Hansen, 1916  
E Arctic ocean, Davis Strait
- Ischnomesidae Hansen, 1916
- Ischnomesus* Richardson, 1908
- I. bispinosus* (G.O. Sars, 1868)  
= *Ischnosoma bispinosus* G.O. Sars, 1868  
Bohuslän-Lofoten-Faeroes-W Ireland-Mediterranean
- I. norvegicus* Svavarson, 1984  
the slopes of Norw. Sea
- Haplomesus* Richardson, 1909
- H. quadrispinosus* (G.O. Sars, 1879)  
Faeroes-off Iceland-off Lofoten, S of Jan Mayen, E Arctic ocean
- H. angustus* Hansen, 1916  
off N Iceland, SE Greenland, E Arctic ocean
- Helomesus* Wolff, 1962 (TM) *Stylomesus menziesi* Birstein, 1960 - abyssal Pacific)
- H. gorbunovi* (Gurjanova, 1946)  
= *Haplomesus gorbunovi* Gurjanova, 1946  
depths of Norw., North Polar & Greenland Seas
- Heteromesus* Richardson, 1909
- H. dentatus* Hansen, 1916  
off SW Iceland
- H. schmidtii* Hansen, 1916  
off S Iceland
- H. frigidus* Hansen, 1916  
Faeroes-off N & E Iceland
- H. longiremis* Hansen, 1916  
E Arctic Ocean, Davis Strait
- H. greeni* (Tattersall, 1905)  
off N Ireland
- Nannoniscidae Hansen, 1916
- Nannoniscus* G.O. Sars, 1869
- N. aequiremis* Hansen, 1916  
off S Jan Mayen
- N. affinis* Hansen, 1916  
off SW Iceland
- N. analis* Hansen, 1916  
Davis Strait
- N. arcticus* Hansen, 1916  
off Jan Mayen-E Greenland
- N. arctoabyssalis* Just, 1980  
abyssal Arctic Seas

- N. bicuspis* G.O. Sars, 1877  
Norw. Sea
- N. inermis* Hansen, 1916  
Davis Strait
- N. laticeps* Hansen, 1916  
off N Iceland
- N. minutus* Hansen, 1916  
Davis Strait
- N. oblongus* G.O. Sars, 1870  
off Lindesnæs-off Lofoten-Faeroes-off W & SW Iceland-Arctic Seas
- N. plebejus* Hansen, 1916  
off SW Iceland
- N. profundus* Svavarson, 1982  
deep part of Norw. Sea & Arctic Basins
- N. reticulatus* Hansen, 1916  
off N Iceland, N Greenland, Kara Sea
- N. simplex* Hansen, 1916  
off W & SW Iceland
- N. spiniornis* Hansen, 1916  
off S Jan Mayen in 2465 m
- Nannoniscoides* Hansen, 1916
- N. angulatus* Hansen, 1916 (TM)  
Faeroes, N Greenland, Kara Sea
- Rapaniscus* Siebenaller & Hessler, 1981
- R. crassipes* (Hansen, 1916)  
= *Nannoniscus crassipes* Hansen, 1916  
off Lofoten
- Regabellator* Siebenaller & Hessler, 1981
- R. armatus* (Hansen, 1916)  
Davis Strait in 3521 m
- Exilinisca* Siebenaller & Hessler, 1981
- E. hanseni* (Just, 1970)  
= *Nannoniscus hanseni* Just, 1970  
N Greenland (Jørgen Brønlund Fjord) in 80-90m
- Austroniscus* Vanhöffen, 1914
- A. norbi* Svavarson, 1982  
deep part of Norw. Sea & Greenland Sea
- A. groenlandicus* (Hansen, 1916)  
= *Nannoniscella groenlandica* Hansen, 1916  
Greenland
- Pseudomesidae Hansen, 1916
- Pseudomesus* Hansen, 1916
- P. brevicornis* Hansen, 1916 (TM)  
off N Iceland
- P. sp. (nov. sp.)* : Just, 1970  
N Greenland
- Desmosomatidae G.O. Sars, 1897
- Desmosomatinae G.O. Sars, 1897
- Mirabilicoxa* Hessler, 1970
- M. fletcheri* Paul & George, 1975  
deep Arctic ocean
- Whoia* Hessler, 1970
- W. angusta* (G.O. Sars, 1899)  
= *Desmosoma angustum* G.O. Sars, 1899  
Bohuslän-Skagerrak-Trondheim Fjord
- W. dumbshafensis* Svavarson, 1988  
deep parts of Norw. Sea
- Desmosoma* G.O. Sars, 1864
- D. lineare* G.O. Sars, 1864 (TM)  
Bohuslän-N Norway-Arctic Seas, e.g. N Greenland
- D. strombergi* Svavarson, 1987  
Faeroes-slopes & depth of Norw. Sea
- Chelator* Hessler, 1970
- C. chelatum* (Stephensen, 1915)  
= *Desmosoma chelator* Stephensen, 1915  
\* pelagic species.  
W Mediterranean
- C. insignis* (Hansen, 1916)  
= *Desmosoma insignis* Hansen, 1916  
Davis Strait
- Eugerdia* Meinert, 1890
- E. arctica* Svavarson, 1988  
deep part of Norw. Sea & Arctic Basins
- E. fragilis* (Kussakin, 1965)  
NE Atlantic
- E. reticulata* (Gurjanova, 1946)  
= *Desmosoma reticulata* Gurjanova, 1946  
Faeroes-depths of Norw. Sea & North Polar Sea
- E. tenuimana* (G.O. Sars, 1868)  
= *Desmosoma tenuimanum* G.O. Sars, 1868
- = *Eugerdia globiceps* Meinert, 1890 (TM)  
non = *Eugerdia tenuimana* : G.O. Sars, 1897  
Öresund-Bohuslän-Skagerrak-W Norway-Faeroes-Iceland-E Greenland
- Pseudogerdia* Kussakin, 1965
- P. filipes* (Hult, 1936)  
= *Desmosoma filipes* Hult, 1936  
Kattegat-Bohuslän-S Norway-Ireland-Mediterranean
- P. intermedia* (Hult, 1936)  
= *Desmosoma intermedium* Hult, 1936  
Kattegat-Bohuslän-W Norway-Faeroes-W Greenland
- P. latipes* (Hansen, 1916)  
= *Desmosoma latipes* Hansen, 1916  
Davis Strait in 200 m
- Echinopleura* G.O. Sars, 1897
- E. aculeata* (G.O. Sars, 1864)  
Bohuslän-Arctic Seas
- Eugerdellatinae Hessler, 1970
- Desmosomella* Kussakin, 1965
- D. armata* (G.O. Sars, 1864)  
= *Desmosoma armatum* G.O. Sars, 1864  
\* May belong to *Eugerdella* Kussakin, 1965?  
Bohuslän-Trondheim-E & N Greenland
- Eugerdella* Kussakin, 1965
- E. coarctata* (G.O. Sars, 1899)  
= *Desmosoma coarctata* G.O. Sars, 1899  
Bohuslän-Trondheim-Davis Strait
- E. hessleri* Just, 1980  
slopes of Norw. Sea-depths of North Polar Sea in 800 m
- E. polita* (Hansen, 1916)  
= *Desmosoma politum* Hansen, 1916  
off W & SW Iceland
- E. zenkewitschi* (Gurjanova, 1946)  
= *Desmosoma zenkewitschi* Gurjanova, 1946  
sublittoral Arctic Seas
- Prochelator* Hessler, 1970
- P. lateralis* (G.O. Sars, 1899)  
= *Eugerdia lateralis* G.O. Sars, 1899  
Bohuslän-Skagerrak-Trondheim-Davis Strait
- Oecidiobranchus* Hessler, 1970
- O. nansenii* Just, 1980  
Faeroes-slopes and depths of Norw., North Polar & Greenland Seas
- O. plebejum* (Hansen, 1916)  
= *Desmosoma plebejum* Hansen, 1916  
off E & NE Iceland, N Greenland
- O. polare* (Gurjanova, 1946)  
= *Desmosoma polaris* Gurjanova, 1946  
sublittoral Arctic Seas
- Cryodesma* Svavarson, 1988
- C. agnari* Svavarson, 1988  
deep part of Norw. Sea & Arctic Basins
- Ilyarachnidae Hansen, 1916
- Ilyarachna* G.O. Sars, 1869
- I. longicornis* (G.O. Sars, 1864)  
= *Mesosstenus longicornis* G.O. Sars, 1869  
= *I. plunketti* Tattersall, 1905  
Bohuslän-W Norway-Iceland & Arctic Seas, e.g. E & N Greenland
- I. torleivi* Svavarson, 1988  
Faeroes-slopes of Norw. Sea
- I. bergendahli* Ohlin, 1901  
Faeroes-Spitsbergen-E Greenland
- I. hirticeps* G.O. Sars, 1870  
= *I. denticulata* G.O. Sars, 1899  
Stavanger-N Norway-Spitsbergen-off Iceland-Faeroes-Davis Strait
- I. dubia* Hansen, 1916  
off NE Iceland in 1666-1902m
- I. thori* Wulff, 1962  
SW of Faeroes, S of Iceland (in 216-820m)
- I. derjugini* Gurjanova, 1946  
abyssal Arctic Seas
- I. frami* Just, 1980  
abyssal Arctic Seas
- Pseudarachna* G.O. Sars, 1897
- P. hirsuta* (G.O. Sars, 1864)  
Kattegat-Bohuslän-Trondheim-Faeroes-N Br. Isles
- Aspidarachna* G.O. Sars, 1897
- A. clypeata* (G.O. Sars, 1870)  
= *Ilyarachna clypeata* G.O. Sars, 1870  
N Norway, Faeroes-Arctic Seas
- Echinozone* G.O. Sars, 1897
- E. coronata* (G.O. Sars, 1870)  
= *Ilyarachna coronata* G.O. Sars, 1870  
W & N Norway-between Norway & Shetlands-Faeroes-off S & SW Iceland-Davis Strait
- E. arctica* Hansen, 1916

= *Ilyarachna arctica* (Hansen,1916) : Wolff,1962  
Faeroes-Jan Mayen, N Greenland, Kara Sea

Eurycopidae Hansen,1916

*Eurycope* G.O. Sars,1864

*E. cornuta* G.O. Sars,1864  
Bohuslän-N Norway-Faeroes-Iceland & Arctic Seas

*E. producta* G.O. Sars,1868  
Bohuslän-N Norway-Faeroes-Iceland & Arctic Seas-off W Ireland

*E. inermis* Hansen,1916  
Faeroes-Norw. Sea-Arctic Seas

*E. hanseni* Ohlin,1901  
Faeroes-Norw. Sea-Arctic Seas

*E. brevisrostris* Hansen,1916  
Faeroes-off Iceland, N Greenland

*E. parva* Bonnier,1896  
off SW Faeroes-Bay of Biscay-Davis Strait

*E. cryoabyssalis* Just,1980  
deep water in North Polar Sea

*E. dahli* Svavarson,1987  
deep part of Norw. Sea & Arctic Basins

*E. ratmanovi* Gurjanova,1946  
deep Arctic waters

*Munneurycope* Stephensen,1912

*M. murrayi* (Walker,1903)  
= *Eurycope murrayi* Walker,1903  
\* pelagic species.  
off SW Faeroes-off S Iceland- off W Ireland

*M. incisa* (Gurjanova,1946)  
the depths of Norw. Sea, off NW Spitsbergen

*Munnopsurus* Richardson,1912

*M. giganteus giganteus* (G.O. Sars,1877)  
off W Norway - off N & SE Faeroes-off E & N Iceland-Arctic Seas

*M. longipes* (Tattersall,1905)  
off SW Faeroes-off W Ireland-Davis Strait

*Tythocope* Wilson & Hessler,1981

*T. megalura* (G.O. Sars,1872)  
= *Eurycope megalura* G.O. Sars,1872  
Stavanger-Hardangerfjord-off Faeroes-off SW Iceland

*T. pygmaea* (G.O. Sars,1870)  
= *Eurycope pygmaea* G.O. Sars,1870  
Bohuslän-Lofoten

*T. laktionovi* (Gurjanova,1946)  
Slopes of North Polar Seas

*T. neupokoevi* (Gurjanova,1946)  
= *Eurycope neupokoevi* Gurjanova,1946  
slopes of North Polar Seas

*Disconectes* Wilson & Hessler,1981

*D. phalangium* (G.O. Sars,1864)  
= *Eurycope phalangium* G.O. Sars,1864  
Bohuslän-N Norway-Faeroes-W Ireland-Scotland-Davis Strait

*D. furcatus* (G.O. Sars,1870)  
= *Eurycope furcata* G.O. Sars,1870  
E Skagerrak-Stavanger-off Lofoten-Faeroes-off SW Iceland-Davis Strait

*D. latirostris* (G.O. Sars,1882)  
= *Eurycope latirostris* G.O. Sars,1882  
Norw. Sea - W of Ireland

*D. coxalis* Kussakin,1983  
abyssal Arctic Seas

*Baeonectes* Wilson,1982

*B. muticus* (G.O. Sars,1864)  
= *Eurycope mutica* G.O. Sars,1864  
Öresund-Bohuslän-N Norway-Faeroes-Iceland & Arctic Seas, e.g. N Greenland-Br. Isles

Munnopsidae G.O. Sars,1869

*Lipomera* Tattersall,1905

*L. (Lipomera) lamellata* Tattersall,1905 (TM)  
Ireland

*Munnopsis* M. Sars,1861  
= *Munnopsoides* Tattersall,1905 (p.p.)  
= *Pseudomunnopsis* Hansen,1916

*M. typica* M. Sars,1861 (TM)  
Bohuslän-N Norway-Faeroes-Iceland & Arctic Seas

*M. beddardi* (Tattersall,1905)  
= *Munnopsoides beddardi* Tattersall,1905  
Faeroes-W Ireland in 364 m

*M. eximius* (Hansen,1916)  
= *Munnopsoides eximius* Hansen,1916  
Faeroes-off S Iceland

*Paramunnopsis* Hansen,1916

*P. oceanica* (Tattersall,1905)  
= *Munnopsis oceanica* Tattersall,1905  
\* pelagic species.  
off W Ireland-Davis Strait

*P. justii* Svavarson,1988

deepest parts of Greenland & North Polar Seas

Abysianiridae Menzies,1956  
\* No species within the area?

Echinothambematidae Menzies,1956  
\* No species within the area?

Haplomunnidae Wilson,1976  
\* No species within the area?

Mesosignidae Schultz,1969  
\* No species within the area?

Pleurocopidae Fresi & Schiecke,1972  
\* No species within the area?

Thambematidae Stebbing,1912

*Thambema* Stebbing,1912

*T. amicorum* Stebbing,1912 (TM)  
W of Ireland in 1993 m

*T. fiatum* Harrison,1987  
Rockall Trench in 1800 m

*T. golanachum* Harrison,1987  
S Rockall Trench in 1330 m

*T. tanum* Harrison,1987  
S Rockall Trench in 2875 m

Gnathostenroidoidea Kussakin,1967  
\* No species within the area?

Gnathostenroidoidea Kossakin,1967

CALABOZOIDEA van Lieshout,1983  
\* Not known from Europe.

Calabozoidae van Lieshout,1983

ONISC(O)IDEA Latreille,1803

TYLOMORPHA Vandel,1943  
\* Only terrestrial?

Tyliidae H. Milne-Edwards,1940

LIGIAMORPHA Vandel,1943

DIPLOCHAETA Vandel,1957

Ligiidae Brandt,1883

*Ligia* J.C. Fabricius,1798

*L. oceanica* (Linnaeus,1767)  
Terrestrial sea-shore inhabitant in Sweden, Norw., Denm., Icel., Faeroes, Br. Isles etc.

*L. exotica* Roux,1828  
Britain, Mediterranean shores

Mesoniscidae Verhoeff,1908  
\* Only terrestrial?

SYNOCHAETA Legrand,1946  
\* 2 superfamilies, 6 families. Only terrestrial?

CRNOCHAETA Legrand,1946  
\* 2 superfamilies, 25 families. Only terrestrial?

PHREATOICIDEA Stebbing,1893  
\* 3 recent families. Not marine & not European.

EPICARIDEA Latreille,1831  
\* Yip,1984 found an unidentified miktoniscus larvae of an Epicaridean Isopod in *Pleurobrachia pileus* at Ireland

Bopyroidea Rafinesque-Schmaltz,1815

Bopyridae Rafinesque-Schmaltz,1815  
@ branchial cavity of Decapods

*Anisarthrus* Giard,1907

*A. pelseeneeri* Giard,1907  
@ abdomen of *Athanas nitescens* (Leach)  
Galway area, NW France

*Asymmetrione* R. & M. Codreanu & Pike,1965  
= *Megachelione* Bourdon,1968

*A. dardani* Bourdon,1968  
@ *Dardanus arrosor* (Herbst)  
Morocco

*A. foresti* (Bourdon,1968)  
= *Megachelione foresti* Bourdon,1968  
@ *Paguristes oculatus* Fabricius  
Balears, Sète, Tunisia

*Athelges* Hesse,1861

*A. bilobus* G.O. Sars,1898  
=? *A. lorifera* Hesse,187?  
@ abdomen of *Pagurus cauanensis* Bell  
N Kattegatt, Bohuslän, Skagerrak, Br. Isles, ?France

*A. cladophorus* Gerstaecker,1862  
@ ?  
Engl. Channel

*A. paguri* (Rathke,1843)  
@ abdomen of *Pagurus bernhardus* (Linnaeus)  
Öresund-Bohuslän-Skagerrak- North Sea, Britain, NW France

*A. tenuicaudis* G.O. Sars,1898  
@ abdomen of *Anapagurus chiroacanthus* (Liljeborg)  
N Kattegatt, Bohuslän, S Norway, N Norway, North Sea-Norw. Sea-Scotland

**A. prideauxi** Giard & Bonnier,1890  
@ *Pagurus prideauxi* (Leach)  
Br. Isles, Mediterranean

**Bopyrus** Latreille,1802

**B. squillarum** Latreille,1802  
= *Monoculus crangorum* O. Fabricius,1798 (nomen oblitum?)  
= *Bopyrus fougerouxii* Giard & Bonnier,1890  
= *Bopyrus rathkei* Giard & Bonnier,1890 (nom. nudum)  
= *Bopyrus helleri* Giard & Bonnier,1890  
= *Bopyrus treillanus* Giard & Bonnier,1890 (nom. nudum)  
= *Bopyrus xiphias* Giard & Bonnier,1890  
@ *Palaemon serratus* (Pennant), *P. elegans* Rathke, *P. adpersus* Rathke, *P. squilla* (Linnaeus) & *P. xiphias* (Risso)  
Öresund, Netherlands, Britain, Ireland, Roscoff, Brest, Arcachon, Portugal, Spain, Mediterranean, Black Sea

**Bopyrina** Kossman,1881

**B. ocellata** (Czerniavsky,1868)  
= *Bopyrus ocellatus* Czerniavsky,1868  
= *Bopyrus virbii* Walz,1881  
= *Bopyrina giardi* Bonnier,1900  
@ *Hippolyte varians* Leach, *H. inermis* Leach & *H. longirostris* Czerniavsky  
SW England, Galway, Dinard, Roscoff, Arcachon, Mediterranean, Black Sea

**Bopyrissa** Nierstrasz & Brender a Brandis,1931  
= *Urocryptella* R. & M. Codreanu,1963

**B. fraisei** (Carayon,1943)  
= *Pseudione fraisei* Kossman,1886 (nomen nudum)  
= *Pseudione fraisei* Carayon,1943  
@ *Clibanarius erythropus* (Latreille)  
Arcachon

**B. diogeni** (Popov,1927)  
= *Pseudione diogeni* Popov,1927  
@ *Diogenes pugilator* (Roux)  
Jersey, Roscoff, Arcachon, Mediterranean, Black Sea

**Bopyroides** Stimpson,1864

**B. hippolytes** (Krøyer,1838)  
= *Bopyrus hippolytes* Krøyer,1838  
= *Bopyroides acutimarginatus* Stimpson,1864  
= *Bopyroides sarsi* Bonnier,1900  
@ *Spirotocaris lilleborgi* (Danielssen), *S. spinus* (Sowerby), *Eualus gaimardi* (H. Milne-Edwards), *Lebbeus polaris* (Sabine) & ?*Pandalus* spp.  
Bohuslän-Skagerrak-Trondheim-Norw. Sea-Iceland-Davis Strait

**B. cluthae** (Scott,1902)  
= *Pleurocrypta cluthae* Scott,1902  
= *Pleurocrypta patiencei* Scott,1902  
= ?*Bopyroides furcata* Norman,1905 (nomen nudum)  
= *Bopyroides sarsi* : Allen,1965, non Bonnier,1900  
@ *Pandalina brevisrostris* (Rathke) & *Caridion gordoni* Bate  
Northumberland, Plymouth, Clyde Sea, Minch, Roscoff, Galway Bay

**Cancericepon** Giard & Bonnier,1887

**C. elegans** (Giard & Bonnier,1886)  
= *Cepon elegans* Giard & Bonnier,1886  
@ *Pilumnus hirtellus* (Linnaeus)  
Norfolk, The Wash, Roscoff, Arcachon

**C. pilula** (Giard & Bonnier,1886)  
= *Cepon pilula* Giard & Bonnier,1886  
@ *Xantho incisus* Leach  
Roscoff, Concarneau

**Epipenaeon** Nobili,1906

**E. ingens ingens** Nobili,1906  
@ *Penaeus semisulcatus* De Haan  
Turkish Mediterranean, Red Sea

**Ergyne** Risso,1816

**E. cervicornis** Risso,1816 (TM)  
@ *Liocarcinus arcuatus* (Leach)  
Roscoff, Spain, Mediterranean

**Grapsicepon** Giard & Bonnier,1887

**G. edwardsi** Giard & Bonnier,1888  
@ *Planes minutus* (Linnaeus)  
Roscoff, Sargasso Sea

**Gyge** Cornalia & Panceri,1861

**G. branchialis** Cornalia & Panceri,1861 (TM)  
= *G. galathea* Bate & Westwood,1868  
@ *Upogebia stellata* (Leach), *U. deltaura* Leach & *U. pusilla* (Petagna)  
Frederikshavn, Skagerrak, Engl. Channel, Clyde Sea, Irish Sea, Arcachon, Mediterranean, Black Sea

**Hemiarthrus** Giard & Bonnier,1887  
= *Phryxus* Rathke,1843, non Rafinesque-Schmaltz,1815, nec Huebner,1819

**H. abdominalis** (Krøyer,1841)  
= *Bopyrus abdominalis* Krøyer,1841  
@ abdomen of e.g. *Pandalus*, *Spirotocaris* & *Crangon allmani*  
Öresund-Bohuslän-North Sea-N Norway-Iceland-Arctic Seas

**Ione** Latreille, in Cuvier,1817

**I. thoracica** (Montagu,1808)  
= *Oniscus thoracica* Montagu,1808  
@ *Callianassa subterranea* (Montagu) & *C. tyrrenna* (Petagna)  
Belgium, Ireland-S & W Br. Isles-Mediterranean

**Pleurocrypta** Hesse,1865

**P. galathea** Hesse,1865 (TM -Mediterranean)  
= *Pleurocrypta nexa* Stebbing, in Herdman,1894  
= *Pleurocrypta marginata* G.O. Sars,1899  
= *Pleurocrypta hessei* Nierstrasz & Brender-à-Brandis,1931  
@ *Galathea squamifera* Leach, *G. nexa* Embleton & *G. dispersa* Bate  
Bohuslän, Skagerrak, North Sea, Bergen, Trondheim, Shetlands, Br. Isles, Ireland, Bretagne, Mediterranean

**P. longibranchiata** (Bate & Westwood,1868)  
= *Phryxus longibranchiatus* Bate & Westwood,1868  
@ *Galathea squamifera* Leach, *G. nexa* Embleton & *G. dispersa* Bate

Bohusl-Skagerrak-North Sea, Manger, Br. Isles, Bretagne, S France

**P. microbranchiata** G.O. Sars,1898  
= *P. intermedia* Giard & Bonnier,1890 (nomen nudum)  
= *P. intermedia* Bonnier,1900  
@ *Galathea intermedia* Lilleborg  
Kattegatt-Bohuslän-Skagerrak-S Norway-Bergen-North Sea, Shetlands, Br. Isles, Ireland, Bretagne, Spain, Adriatic

**P. piriformis** Bourdon,1968  
@ *Galathea strigosa* (Linnaeus)  
Spain (Bae de Cadaqués)

**P. porcellanae** Hesse,1876  
= *P. porcellanae longicornis* Hesse,1876  
= *P. perezi* Nierstrasz & Brender-à-Brandis,1926  
= *Pseudione convergens* Stock,1960  
@ *Pisidia longicornis* (Linnaeus)  
Northumberland, W Br. Isles, Ireland, Bretagne, Morocco

**P. strigosa** Bourdon,1968, ex Giard & Bonnier MS  
= *Pleurocrypta strigosa* Giard & Bonnier,1890 (nomen nudum)  
@ *Galathea strigosa* (Linnaeus)  
Jersey, Roscoff, Spain, Mediterranean

**Pleurocryptella** Bonnier,1900

**P. formosa** (Giard & Bonnier,1888)  
= *Pleurocrypta formosa* Giard & Bonnier,1888  
@ *Psychogaster formosus* A. Milne-Edwards  
off Ireland (Porcupine Bank) & Canaries

**P. superba** Bourdon,1981  
@ ?  
Denmark

**Pliophrixus** Caroli,1930

\* This genus & species is not listed in the Smithsonian Internet list. Bourdon 1968 does not mention it either.

**P. philonika** (Caroli,1930)  
@ *Processa* spp.  
Britain

**Pseudione** Kossman,1881

**P. hyndmanni** (Bate & Westwood,1868)  
= *Phryxus hyndmanni* Bate & Westwood,1868  
= *Pseudione proxima* Bonnier,1900  
@ *Pagurus bernhardus* (Linnaeus), *P. pubescens* (Krøyer), *P. prideauxi* (Leach), *P. cuanensis* (Bell), *Anapagurus laevis* (Bell) & *A. chiroacanthus* (Lilleborg)  
Öresund-Kattegat-Bohuslän-Skagerrak, Trondheim-Tromsø, Iceland-Faeroes-Br. Isles-Ireland, Belgium-Bretagne, ?Mediterranean

**P. affinis** (G.O. Sars,1882)  
= *Pleurocrypta affinis* G.O. Sars,1882  
= *Pseudione hoylei* Bonnier,1900  
@ *Pandalus montagui* Leach, *Dichelopandalus bonnieri* (Kinahan) & *Plesionika* spp.  
Drøbak, Skagerrak-North Sea, Trondheim, Tromsø, Br. Isles, Ireland, Engl. Channel, Biscay, Mediterranean

**P. crenulata** G.O. Sars,1898  
= *Palaegyge insignis* Giard & Bonnier,1890 (nomen nudum)  
= *Pseudione insignis* Bonnier,1900  
@ *Munida tenuimana* M. Sars & *M. rugosa* (Fabricius)  
Skagerrak, in 500m-North Sea, Bergen, Br. Isles, France

**P. dohrni** Bonnier,1900  
= *Palaegyge dohrni* Giard & Bonnier,1890 (nomen nudum)  
= ?*P. callianassae* Kossman,1881 (nomen nudum)  
@ *Callianassa truncata* Giard & Bonnier  
Gulf of Naples

**P. borealis** Caspers,1939  
= ?*P. callianassae* Kossman,1881 (nomen nudum)  
@ *Callianassa tyrrenna* (Petagna,1792)  
Helgoland

**P. caspersi** Gruner,1966  
= *P. tuberculata* Caspers,1939, non Richardson,1904  
= ?*P. callianassae* Kossman,1881 (nomen nudum)  
@ *Callianassa tyrrenna* (Petagna,1792)  
\* This species is not mentioned in the Smithsonian internet list, but *P. callianassae* Kossman,1881 is.  
Helgoland

**P. confusa confusa** (Norman,1886)  
= *Gyge confusa* Norman,1886  
@ *Galathea dispersa* Bate  
Shetlands, Ireland, Portugal

**Progebiophilus** Codreanu & Codreanu,1963

**P. euxinicus** (Popov,1927)  
= *Pseudione euxinica* Popov,1927  
= *Gyge branchialis* var. *arcassonensis* Carayon,1943  
@ *Upogebia deltaura* Leach, *U. pusilla* (Petagna) & *U. stellata* (Montagu)  
Clyde Sea, Irish Sea, Galway Bay, Roscoff, Arcachon, Mediterranean, Black Sea

**Scyracepon** Tattersall,1905

**S. tuberculosa** Tattersall,1905 (TM - Ireland)  
@ *Rochinia carpenteri* (Norman)  
Ireland & ? France

**Urobopyrus** Richardson,1904

**U. processae** Richardson,1904 (TM - Brazil)  
= *U. provisorius* Rouch & Taberly,1961  
@ *Processa edulis* (Risso), *P. acutirostris* Nouvel & Holthuis & other *Processa* spp.  
Roscoff, Arcachon, Mediterranean, Florida, Brazil

Entoniscidae F. Müller,1871  
@ visceral cavity of Decapoda

**Portunio** Giard & Bonnier,1887

**P. maenadis** (Giard,1886)  
= *Entoniscus maenadis* Giard,1886  
@ visceral cavity of crabs and anomurans, usually *Carcinus maenas* (Linnaeus)  
Isefjord, Bohuslän-Skagerrak, Engl. Channel, W France, Gulf of Naples

**P. kossmani** (Giard & Bonnier,1886)  
= *Entoniscus kossmani* Giard & Bonnier,1886  
@ *Portunus latipes* (Pennant)

Roscoff

**P. moniezi** (Giard,1878)  
= *Entoniscus moniezi* Giard,1878  
@ *Liocarcinus puber* (Linnaeus)  
Roscoff

**P. pusillus** Pérez,19??  
@ *Liocarcinus pusillus* (Leach)  
Roscoff

**P. salvatoris** (Kossmann,1881)  
= *Entione salvatoris* Kossmann,1881  
@ *Liocarcinus arcuatus* (Leach)  
Roscoff

**Priapon** ?,1???

**P. fraisei** (Giard & Bonnier,1886)  
= *Entoniscus fraisei* Giard & Bonnier,1886  
@ *Liocarcinus holsatus* (Fabricius)  
Roscoff

**Entionella** Miyashita,1941 (TM *E. fluviatilis* Miyashita,1941 - Japan. @ *Eriocheir japonicus* de Haan)

**E. monensis** Hartnoll,1960  
@ *Eurynome aspera* (Pennant)  
Br. Isles

**Pinnotherion** Giard & Bonnier,1889

**P. vermiforme** Giard & Bonnier,1889?  
@ *Pinnotheres pisum* (Linnaeus)  
Cornwall, Bretagne

**Cancrion** Giard & Bonnier,1887

**C. miser** Giard & Bonnier,1886  
@ *Pilumnus hirtellus* (Linnaeus)  
Roscoff

**C. floridus** Giard & Bonnier,1886  
@ *Xantho incisus* Leach  
Roscoff

**C. pilipedi** Veillet & Bourdon,19??  
@ *Xantho pilipes* A. Milne-Edwards  
Roscoff

**Entoniscus** Fritz Müller,1862 (TM *E. porcellanae* F. Müller,1862 - Brazil @ *Porcellana* sp.)

**E. muelleri** Giard & Bonnier,1886  
@ *Pisidia longicornis* (Pennant)  
Concarneau

**Synalpheion** Coutière,1908 (TM *S. giardi* Coutière,1908)

**Daididae** G.O. Sars,1882  
@ Eucarida

**Dajus** Krøyer,1849

**D. mysidis** Krøyer,1842  
@ *Mysis oculata* (Fabricius), *M. relicta* Lovén & *M. mixta* Lilljeborg  
N Norway-Iceland, E & N Greenland

**D. profundus** Hansen,1916  
@ *Paramblyops rostrata* Holt & Tattersall  
off SW Faeroes

**Branchiophryxus** Caullery,1897

**B. nyctiphanae** Caullery,1897  
@ gills of *Meganyctiphanes norvegica* (M. Sars)  
Biscay

**Prodajus** Bonnier,1903

**P. lobiancoi** Bonnier,1903  
@ *Gastrosaccus lobatus* Nouvel,1951  
Roscoff

**P. ostendensis** Gilson,1909  
@ *Gastrosaccus spinifer* (Goës,1864)  
S & SE North Sea

**Aegophila** Bresciani,1966

**A. socialis** Bresciani,1966  
@ *Aega ventrosa* M. Sars  
Greenland

**Aspidophryxus** G.O. Sars,1882

**A. peltatus** G.O. Sars,1882  
@ dorsum of mysids, e.g. *Erythrops* spp., *Parerythrops obesa* (G.O. Sars) & *Mysidopsis didelphys* (Norman)  
Bohuslän-Skagerrak-Lofoten-North Sea-Iceland & Arctic Seas, Scotland, Ireland

**Notophryxus** G.O. Sars,1882

**N. ovoides** G.O. Sars,1882  
@ dorsum (usually) of mysids & euphausiids, e.g. *Amblyops abbreviata* (M. Sars)  
Hardangerfjord, Trondheimsfjord, North Sea-Norw. Sea

**N. clypeatus** G.O. Sars,1885  
@ dorsum of *Pseudomma roseum* & *Pseudomma truncatum*  
off N Iceland

**Holophryxus** Richardson,1905

**H. richardii** Koehler,1911  
@ *Sergestes arcticus* Krøyer  
Bergen-Iceland-W France-Azores

**H. acanthephyrae** Stephensen,1912  
@ *Acanthephyra purpurea* A. Milne-Edwards  
Cape Farewell

**Heterophryxus** G.O. Sars,1885

**Zonophryxus** Richardson,1903

**Arthropryxus** Richardson,1908

**Colophryxus** Richardson,1908

**Prophryxus** Richardson,1909

**Allophryxus** Koehler,1911

**Paradajus** Nierstrasz & Brender à Brandis,1931

**Streptodajus** Nierstrasz & Brender à Brandis,1931

**Cryptoniscina**

@ non Eucaridean Crustacea

\* Sometimes considered to contain just one family (Cryptoniscidae), but usually split up in different (host depending) families.

**Hemioniscidae**

@ Cirripedia

\* Not listed by Schram,1986

**Hemioniscus** Buchholz,1866

=? *Cryptothir* Dana,1852

**H. balani** (Bate,1860)

@ different balanid species

\* Should the author name be Buchholz,1866 (if (Bate,1860) is considered a nomen nudum)?  
Bohuslän, Kristiansand, North Sea, N Norway, Ireland, Scotland, Engl. Channel

**Leponiscus** Giard,1887

**L. anatiferae** Giard,1887

@ *Lepus anatifera*

Roscoff

**Crinoniscidae** Bonnier,1900

@ Balanidae

\* May perhaps be synonymized with Hemioniscidae, but the the single species within this family, *C. equiptans* is said to differ much from that other family.

**Crinoniscus** Pérez,1900

**C. equiptans** Pérez,1900 (TM)

@ *Balanus perforatus* ? & *B. tintinnabulum*

?European Atlantic

**Cyproniscidae**

@ Ostracoda

\* Not listed by Schram,1986

**Cyproniscus** Kossmann,1884

**C. cypridinae** (G.O. Sars,1882)

@ *Vargula norvegica* & *Philomedes brenda*

Bohuslän-Skagerrak-Lofoten-Norw. Sea

**Liriopsidae** Bonnier,1900

@ Rhizocephala

**Liriopsis** Max Schultze,1859

= *Liriopse* Rathke,1843, *non* Meigen,1800, *nec* Lesson,1837

**L. pygmaea** (Rathke,1843)

@ *Peltoaster paguri*

N Kattegatt-Bohuslän-W Norway, North Sea, White Sea, Black Sea

**Liriopsis** sp. cf. **L. monophthalma** Fraisse,1878 / **pygmaea** Rathke

@ rhizocephalans on *Munida* sp.

\* vide Brinkmann, A. jr.,1984. *Sarsia* 69 : 211-213

W Norway

**Liriopsis** sp.

@ *Galathea minor* Boschma on *Galathea nexa* sp.

\* vide Brinkmann, A. jr.,1984. *Sarsia* 69 : 211-213

W Norway

**Danalia** Giard,1887

**D. larvaeformis** (Giard,1887?)

@ *Saccalana carcini* / *Carcinus maenas* (Linnaeus)

Roscoff

**Danalia** sp.

@ *Sylon hippolytes* from *Pandalina* ?*profunda*

\* vide Brinkmann, A. jr.,1984. *Sarsia* 69 : 211-213

W Norway

**Asconiscidae**

@ Mysida

\* Not listed by Schram,1986

**Asconiscus** G.O. Sars,1899

**A. simplex** G.O. Sars,1899

@ *Boreomysis arctica* (Krøyer)

Bohuslän-Oslofjord-Skagerrak, Trondheim, North Sea, Ireland

**Podasconidae**

@ Amphipods

\* Not listed by Schram,1986

**Parapodascon** Hansen,1916

**P. stebbingi** (Giard & Bonnier,1895)

= *Podascon Stebbingi* Giard & Bonnier,1895

@ *Boeckosimus plautus* (Krøyer) & *B. normani* (G.O. Sars)

Bergen, North Sea

**Cabiropsidae**

@ Isopods

\* Not listed by Schram,1986

**Clypeoniscus** Giard & Bonnier,1895

**C. hanseni** Giard & Bonnier,1895

@ *Idotea* spp.

S Baltic-Bohuslän-W Norway, Scotland, Engl. Channel, France

**C. meinerti** Giard & Bonnier,1895

@ *Pleuropriion murdochi* & *Synidothea nodulosa* Krøyer

?Faeroes or ?Iceland-Kara Sea-W Greenland

- Munnoniscus* Giard & Bonnier, 1895
- M. marsupialis* (G.O. Sars, 1882) <sup>(TM)</sup>  
@ *Eurycope cornuta* G.O. Sars & *Ilyarachna longicornis* (G.O. Sars)  
Bohuslän-Skagerrak-Norway-North Sea
- Cironiscus* Nielsen, 1967
- C. dahli* Nielsen, 1967  
@ *Natatalana borealis* (Lilljeborg) & *Politolana microphthalmia* (Hoek)  
W Baltic-Skagerrak-North Sea-Norw. Sea
- Ancyroniscus* Caullery & Mesnil, 1919
- A. bonnieri* Caullery & Mesnil, 1919  
@ *Dynamene bidentata* (Adams)  
Roscoff, S & W Britain, Ireland?
- Arcturocheres* Hansen, 1916
- A. pulchripes* Hansen, 1916  
@ *Astacilla granulata* (G.O. Sars) & *Arcturus hystrix* G.O. Sars  
off SE Iceland-off W Faeroes
- Astacilloechus* Hansen, 1916
- A. ingolfi* Hansen, 1916  
@ marsupium of *Astacilla granulata* (G.O. Sars)  
off SE Iceland
- Cabtirops* Kossman, 1884  
= *Cabira* Kossman, 1872, *non* Sodovsky, 1837 (Lepidoptera), *nc* Gistel, 1847 (Coelenterata)
- C. marsupialis* (Caroli, 1953)  
= *Paracabtirops marsupialis* Caroli, 1953  
@ hyperparasite on *Gyge branchialis* Cornalia & Panceri & *Bopyrina ocellata* (Czerniavsky)  
Gulf of Naples, S France
- C. ibizae* Bourdon, 1966  
@ hyperparasite on *Megachelione foresti* Bourdon, 1968  
Sète, Balears
- C. perezi* Carayon, 1942  
@ hyperparasite on *Bopyrissa fraissei* (Carayon)  
Arcachon
- C. codreanui* Bourdon, 1966  
@ hyperparasite on *Bopyrissa diogeni* (Popov)  
Engl. Channel
- C. sp.* - very close to *C. ibizae* Bourdon : Bourdon, 1968  
@ hyperparasite on *Pleurocrypta microbranchiata* G.O. Sars, 1899  
Mediterranean
- "*Cryponiscid No. 2*" : G.O. Sars, 1899  
@ *Aega ventrosa*  
?Norway
- Cryponiscidae s.l.
- Cumochus* Hansen, 1916
- C. insignis* Hansen, 1916  
@ *Diatylis polaris* G.O. Sars, *D. echinata* Bate & *Hemilamprops cristata* G.O. Sars  
Faeroes-Jan Mayen
- AMPHIPODA Latreille, 1816
- CAPRELLIDEA Leach, 1814
- CYAMIDA Bousfield, 1979
- Cyamoidea White, 1847
- Cyamidae White, 1847
- Isocyamus* Gervais & van Beneden, 1859
- delphinii* (Guérin-Méneville, 1836)  
= *Cyamus globicipitis* Lütken, 1873  
@ *Phocaena phocaena* & other species (e.g. *Globicephala melaena*, *G. melas*, *Mesoplodon grayi*, *Pseudorca crassidens*)  
Bohuslän, Netherlands, Faeroes, Mediterranean, Australia
- I. kogiae* Sedlak-Weinstein, 1992  
@ *Kogia breviceps* (de Blainville, 1838)  
Queensland
- Cyamus* Latreille, 1796  
= *Paracyamus* G.O. Sars, 1895
- C. boopis* Lütken, 1873  
@ *Megaptera novaeangliae* (Borowski) "Humpback whale"  
Skagerrak-Arctic Seas & Iceland
- C. erraticus* Roussel de Vauzème, 1834  
@ *Eubalaena glacialis* Müller  
Shetland, Rockall-Iceland & Arctic Seas
- C. ovalis* Roussel de Vauzème, 1834  
@ *Eubalaena glacialis* Müller & *Physeter catodon* Linnaeus  
Shetlands-Faeroes-Iceland
- C. gracilis* (Roussel de Vauzème, 1834)  
@ *Eubalaena glacialis* Müller  
British Isles
- C. oreini* Leung, 1970  
@ *Orcinus orca* (Linnaeus)  
South Africa
- C. ceti* (Linnaeus, 1758) Lamarck, 1801  
@ *Balaena mysticetus* Linnaeus & *Eschrichtius gibbosus* (Erleben)  
Greenland, Scotland
- C. balaenoptera* Barnard, 1931  
@ *Balaenoptera* spp.  
South Africa
- C. monodontis* Lütken, 1873  
@ *Monodon monoceros* Linnaeus & *Delphinapterus leucas* (Pallas)
- Greenland
- C. nodosus* Lütken, 1860  
@ *Monodon monoceros* Linnaeus & *Delphinapterus leucas* (Pallas)  
Greenland
- C. catodontis* Margolis, 1954  
@ *Physeter catodon* Linnaeus  
Netherlands, S America
- C. bahamondei* Buzeta, 1963  
@ *Physeter catodon* Linnaeus  
Bahamas?
- C. antarcticensis* Berzin & Vlasova, 1982  
@ *Orcinus orca* (Linnaeus)  
Antarctic
- C. orubraedon* Waller, 1989  
@ *Berardius bairdii*  
Japan
- Platyeyamus* Lütken, 1873
- P. thompsoni* (Gosse, 1855)  
@ *Hyperoodon ampullatus* Forster, *Mesoplodon grayi*  
Netherlands, England, Spitsbergen, Australia
- P. flaviscutatus* Waller, 1989  
@ *Berardius bairdii*  
Japan
- Scutocyamus* Lincoln & Hurley, 1974
- S. parvus* Lincoln & Hurley, 1974  
@ *Lagenorhynchus albirostris* Gray  
Netherlands
- S. antipodensis* Lincoln & Hurley, 1980  
@ *Cephalorhynchus hectori*  
New Zealand
- Syncyamus* Bowman, 1955
- S. pseudorcae* Bowman, 1955  
@ *Pseudorca crassidens* (Owen)  
The Caribbean, Australia
- S. aequus* Lincoln & Hurley, 1981  
@ *Stenella coeruleoalba*, *S. longirostris*, *Delphinus delphis*, *Tursiops aduncus*, *T. truncatus*, *Mesoplodon grayi*  
Mediterranean, South Africa, Australia
- S. sp.* Leung, 1970  
@ *Delphinus delphis* Linnaeus  
Gibraltar
- Neocyamus* Margolis, 1955
- N. physeteris* (Pouchet, 1888)  
@ *Physeter catodon* Linnaeus & *Phocoenoides truei* Andrews  
S America
- CAPRELLIDA Bousfield, 1979
- Phisicoidea Vassilenko, 1968
- Phisicidae Vassilenko, 1968
- Phtisica* Slabber, 1769
- P. marina* Slabber, 1769  
Öresund-Bohuslän-W & N Norway-Br. Isles-Senegal-Azores-Canaries-Black Sea
- Protellidae McCain, 1970  
\* Correct superfamily?
- Protellina* Stephensen, 1944
- P. ingolfi* Stephensen, 1944  
Faeroes, Iceland, between Iceland & Norway
- Paracercopidae Vassilenko, 1968  
\* No species within the area?
- Dodecadidae Vassilenko, 1968  
\* No species within the area?
- Caprelloidea White, 1847
- Caprellidae White, 1847
- Caprella* Lamarck, 1801
- C. linearis* (Linnaeus, 1767)  
W Baltic-Bohuslän-Iceland & Arctic Seas
- C. septentrionalis* Krøyer, 1838  
Öresund-Bohuslän-Iceland & Arctic Seas
- C. monocera* G.O. Sars, 1895  
\* this variety of *C. septentrionalis* should be treated as a separate species until more information about it have been gained, sensu Laubitz, 1972  
Finnmark?
- C. equilibra* Say, 1818  
= *Caprella aequilibra* : Aucet.  
Kattegatt & Trondheimsfjord-"almost cosmopolitan"
- C. ciliata* G.O. Sars, 1883  
Öresund-Bohuslän-S Norway-off W Norway-Iceland
- C. microtuberculata* G.O. Sars, 1879  
N Norway-Arctic Seas-off SW & NW Faeroes
- C. acanthifera* Leach, 1814  
North Sea
- C. rinki* Stephensen, 1916  
off SW Faeroes-off W Iceland-SW Greenland
- C. penantis* Leach, 1814

- = *Caprella acutifrons* Latreille, 1816  
Engl. Channel, Mediterranean
- C. tuberculata* Bate & Westwood, 1866  
Helgoland, E Britain, Engl. Channel, W France
- C. fretensis* Stebbing, 1878  
Engl. Channel, W Britain, Ireland
- C. erethizon* Mayer, in Gadeau de Kerville, 1901  
Engl. Channel
- C. macho* Platvoet, De Bruyne & Gmelig-Meyling, 1995  
Netherlands
- C. andreae* Mayer, 1999  
Ireland
- Metacaprella* Mayer, 1903
- M. horrida* (G.O. Sars, 1885)  
round Faeroes-Arctic Seas
- Proaeginina* Stephensen, 1940
- P. norvegica* (Stephensen, 1931)  
N Norway-S Iceland
- Aeginina* Norman, 1905
- A. longicornis* (Krøyer, 1842-43)  
= *A. echinata* (Boeck, 1861)  
Öresund, W Jutland, E Scotland, Bergen, Faeroes, Iceland, Arctic Seas
- Parvipalpa* Stephensen, 1944
- P. verrucosa* Stephensen, 1944  
off SW Iceland
- Parvipalpus* Mayer, 1890
- P. capillaceus* (Chevreux, 1887)  
= *Aegina capillacea* Chevreux, 1887  
Ireland (Valencia harbour), Concarneau
- Thorina* Stephensen, 1944
- T. spinosa* Stephensen, 1944  
off SW Faeroes
- Pariambidae Laubitz, 1993
- Pariambus* Stebbing, 1888  
= *Podalirius* Krøyer, 1844, non Latreille, 1802, non Swainson, 1833
- P. typicus* (Krøyer, 1845)  
= *Podalirius typicus* Krøyer, 1845  
\* Associated with *Asterias rubens* & *Crossaster papposus*  
Öresund-Bohuslän-Norw. Sea, Br. Isles
- Pseudoprotella* Mayer, 1890
- P. phasma* (Montagu, 1806)  
Kattegat-Bohuslän-North Sea, Br. Isles
- Aeginellidae Vassilenko, 1968
- Aeginella* Boeck, 1861
- A. spinosa* Boeck, 1861  
Haugesund-W & N Norway-Arctic Seas-Iceland-off W & NW Faeroes
- Caprogammaridae Kudrjaschov & Vassilenko, 1966  
\* No species within the area?
- HYPERIIDEA H. Milne-Edwards, 1830
- PHYSOSOMATA Pirlot, 1929
- Scinoidea Stebbing, 1888
- Scinidae Stebbing, 1888
- Scina* Prestandrea, 1833
- S. borealis* (G.O. Sars, 1882)  
= *Tyro Clausii* Bovallius, 1885  
Koster Channel-Skagerrak-W & N Norway-Arctic Seas-off S Iceland-Faeroe Channel
- S. crassicornis* (J.C. Fabricius, 1775)  
= *Tyro Sarsi* Bovallius, 1885  
off S Iceland-Davis Strait
- S. ratrayi* Stebbing, 1895  
off S Iceland-W Scotland
- S. tullbergi* (Bovallius, 1885)  
= *Scina pacifica* Stebbing, 1895  
Iceland
- S. marginata* (Bovallius, 1885)  
= *S. lepisma* Chun, 1889  
?European Atlantic
- S. submarginata* Tattersall, 1906  
?European Atlantic
- S. vosseleri* Tattersall, 1906  
?European Atlantic
- S. spinosa* Vosseler, 1901  
= *Scina uniceps* Stebbing, 1895  
?European Atlantic
- S. oedicarpus* Stebbing, 1895  
?European Atlantic
- Acanthoscina* Vosseler, 1900
- A. acanthodes* (Stebbing, 1895)  
off S Iceland-off W Ireland-N & S Atlantic
- Parascina* Stebbing, 1904  
\* Is probably synonymous with another genus (& may belong to another family) as this genus is not mentioned in the revision of Bowman & Gruner, 1973.
- P. fowleri* Stebbing, 1904  
off S Faeroes-off S Iceland-Davis Strait
- Mimonectidae Bovallius, 1885
- M. loveni* Bovallius, 1885  
off SW Iceland-Sargasso Sea
- Mimonectes* Bovallius, 1885
- M. steenstrupii* Bovallius, 1885  
off S Iceland-Davis Strait
- M. gausi* (Woltereck, 1904)  
?European Atlantic
- M. loveni* Bovallius, 1885  
?European Atlantic
- Proscinidae Pirlot, 1933  
\* No species in the area?
- Archaeoscinidae Stebbing, 1904  
\* No species in the area?
- Lanceoloidea Bovallius, 1887
- Lanceolidae Bovallius, 1887
- Lanceola* Say, 1818 (<sup>TM</sup> *L. pelagica* Say, 1818)
- L. sayana* Bovallius, 1885  
off S Iceland-southwards in the Atlantic
- L. murrayi* Norman, 1900  
Faeroe Channel-southwards in the Atlantic
- L. serrata* Bovallius, 1885  
off S Iceland-Davis Strait-N Atlantic
- L. pacifica* Stebbing, 1888  
off S Iceland-W Ireland-Gibraltar
- L. loveni* Bovallius, 1885  
= *Lanceola aestiva* Stebbing, 1888  
off S Iceland-Davis Strait-SW Ireland-Bay of Biscay-N Atlantic
- L. borealis* Bate & Westwood, 1888  
?European Atlantic
- Chuneolidae Woltereck, 1909  
\* No species in the area?
- Microphasmidae Stephenson & Pirlot, 1931  
\* No species in the area?
- PHYSOCEPHALATA Bowman & Gruner, 1973
- Vibilioidea Dana, 1952
- Vibilidae Dana, 1852
- Vibilia* H. Milne-Edwards, 1830 (<sup>TM</sup> *V. peronii* H. Milne-Edwards, 1830)
- V. borealis* Bate & Westwood, 1868  
= *Vibilia Krøyeri* Bovallius, 1887  
Faeroes-W Greenland-Scotland-N Ireland-Gibraltar-W Mediterranean
- V. armata* Bovallius, 1887  
European Atlantic N to 50°, Mediterranean
- V. cultripes* Vosseler, 1901  
European Atlantic N to 50°, Mediterranean
- V. jeangerardii* Lucas, 1849  
@ Salps  
European Atlantic N to 43° (52°), Mediterranean
- V. propinqua* Stebbing, 1888  
European Atlantic N to 45° (60°), Mediterranean
- V. pyripes* Bovallius, 1887  
European Atlantic N to 40° (50°)
- V. viatrix* Bovallius, 1887  
@ Salps  
European Atlantic N to 45°, Mediterranean
- V. gibbosa* Bovallius, 1887  
?European Atlantic, Mediterranean
- Cystisomatidae Willemoës-Suhm, 1875
- Cystisoma* Guérin-Méneville, 1842
- C. fabricii* Stebbing, 1888  
European Atlantic N to 50°
- C. latipes* (Stephensen, 1925)  
= *Thaumatops latipes* Stephensen, 1925  
European Atlantic N to 50°
- C. pellucida* (Willemoës-Suhm, 1874)  
= *Cystisoma parkinsoni* Stebbing, 1888  
European Atlantic N to 50° & 58-59° N
- C. neptunus* Guérin-Méneville, 1842 (<sup>TM</sup>)  
?European Atlantic
- C. spinosum* (J.C. Fabricius, 1775)  
?European Atlantic
- Paraphronimidae Bovallius, 1887
- Paraphronima* Claus, 1879
- P. crassipes* Claus, 1879  
European Atlantic N to 36° (40°), Mediterranean

- P. gracilis* Claus, 1879 (TM)  
European Atlantic N to 40°, Mediterranean
- Platysceloidea Bate, 1862
- Lycaeidae Claus, 1879  
\* No species in the area?
- Tryphanidae
- Tryphana* Boeck, 1870  
= *Tryphaena* Bovallius, 1887
- T. malmi* Boeck, 1870 (TM)  
= *Tryphaena Nordenskiöldii* Bovallius, 1887  
= *Tryphaena Boeckii* Stebbing, 1888  
Hardangerfjord-off W Norway-Iceland-Faeroes-Shetlands-Scotland
- Brachyscelidae
- Brachyscelus* Bate, 1861
- B. cruscum* Bate, 1861 (TM)  
= *Thamyris mediterranea* Claus, 1887  
off SW Iceland-off SW Faeroes-Atlantic Ocean & Mediterranean
- Anapronoidea Bowman & Grüner, 1973  
\* No species in the area?
- Oxycephalidae Bate, 1862
- Glossocephalus* Bovallius, 1887
- G. milneedwardsi* Bovallius, 1887 (TM)  
?European Atlantic, Mediterranean
- Parascelidae Bovallius, 1887
- Thyropus* Dana, 1852 (TM *T. diaphanus* Dana, 1852)  
= *Parascelus* Claus, 1879
- T. typhoides* (Claus, 1887)  
= *Parascelus typhoides* Claus, 1887  
European Atlantic in 40-42° N, Mediterranean
- Platyscelidae Bate, 1862
- Platyscelus* Bate, 1861
- P. ovoides* (Risso, 1816)  
= *Typhis ovoides* Risso, 1816 (TM)  
European Atlantic N to 35° (50°), Mediterranean
- P. serratulus* Stebbing, 1888  
European Atlantic N to 35° (50°), Mediterranean
- Tetrathyrus* Claus, 1879
- T. forcipatus* Claus, 1879 (TM)  
European Atlantic N to 35°, Mediterranean
- Pronoidea Claus, 1879
- Euphranoë* Claus, 1879
- E. minuta* Claus, 1879  
European Atlantic N to 35°, Mediterranean
- Paralycaea* Claus, 1879
- P. gracilis* Claus, 1879  
European Atlantic N to 35°, Mediterranean
- Phronimoidea Dana, 1853
- Phronimidae Dana, 1853
- Phronima* Latreille, 1802
- P. sedentaria* (Forskål, 1775)  
= *Cancer sedentarius* Forskål, 1775 (TM)  
= *Phronima atlantica* Guérin-Méneville, 1836  
\* Associated with salps, pyrosomes & dolioleids  
North Sea-off SW Faeroes-off SW Iceland, Mediterranean
- P. colletti* Bovallius, 1887  
European Atlantic N to 40° (45°), Mediterranean
- P. stebbingii* Vosseler, 1901  
European Atlantic N to 36°, Mediterranean
- Hyperiididae Dana, 1852
- Themisto* Guérin, 1828, non Oken, 1807, non Oken, 1815 (nomina rejicenda)  
= *Parathemisto* Boeck, 1871  
= *Euthemisto* Bovallius, 1887
- T. abyssorum* (Boeck, 1871)  
= *Parathemisto abyssorum* Boeck, 1870  
= *Parathemisto oblivia* G.O. Sars, 1895  
Kattegatt-Bohuslän-W & N Norway-Arctic Seas-Jan Mayen-Iceland-Faeroes
- T. compressa* Goës, 1865  
= *Parathemisto gracilipes* (Norman, 1869)  
= *Parathemisto gaudichaudi* : Auct., non (Guérin, 1825)  
= *Themisto bispinosa* Boeck, 1871  
@ different hydromedusae; also feeding on other zooplankton, e.g. copepods & chaetognaths  
Bohuslän-North Sea-N Norway-Jan Mayen-Iceland-Faeroes-W Ireland
- T. libellula* (Mandt, 1822)  
N Norway-Arctic Seas, e.g. N Greenlandoff W, N & E Iceland-off N Faeroes
- Hyperoche* Bovallius, 1887
- H. medusarum* (Krøyer, 1838)  
= *Meteocus medusarum* Krøyer, 1838  
= *Hyperoche krøyeri* Bovallius, 1887 (TM)  
\* Associated with *Pleurobrachia pileus* (O.F. Müller)  
Øresund-Bohuslän-S, W & N Norway-Arctic Seas-Jan Mayen-Iceland-off W Faeroes
- Hyperia* Latreille, in Desmarest, 1823
- H. galba* (Montagu, 1813)  
= *Cancer gammarus galba* Montagu, 1813  
@ *Aurelia* (*Cyanea* etc.)  
Åland-Bohuslän-North Sea-W & N Norway-Iceland-Faeroes-Shetlands-Br. Isles, N Greenland
- H. medusarum* (O.F. Müller, 1776)  
= *Hyperia Sueri* Latreille, in Desmarest, 1823 (TM)  
= *Cancer medusarum* O.F. Müller, 1776  
@ *Cyanea*, *Aurelia* etc.  
Øresund-Bohuslän-W & N Norway-Spitsbergen-Iceland-Faeroes-W Ireland
- H. spinigera* Bovallius, 1889  
@ ?  
\* Possibly a synonyme of *H. galba* (Montagu)  
off N Norway-Spitsbergen, off S Iceland, off W Ireland & S England
- Hyperoidea Chevreux, 1900
- H. longipes* Chevreux, 1900 (TM)  
W France, off Biscay, Azores, Mediterranean
- Lestrigonus* H. Milne-Edwards, 1830 (TM *L. fabrei* H. Milne-Edwards, 1830)
- L. latissimus* (Bovallius, 1887)  
= *Hyperia latissima* Bovallius, 1887  
?European & SE Atlantic, Mediterranean
- Dairellidae Bovallius, 1987
- Dairella* Bovallius, 1887 (TM *Paraphronima californica* Bovallius, 1885 - Pacific)
- D. latissima* Bovallius, 1887  
European Atlantic N to 40° (57°)
- Phrosinidae Dana, 1853
- Anchylomera* H. Milne-Edwards, 1830
- A. blossevillii* H. Milne-Edwards, 1830 (TM)  
European Atlantic N to 36°, Mediterranean
- Phrosina* Risso, 1822
- P. semilunata* Risso, 1822 (TM)  
European Atlantic N to 40° (50°), Mediterranean
- Primno* Guérin-Méneville, 1836 (TM *P. macropus* Guérin-Méneville, 1836)
- P. evansi* Shearer, 1986  
= *Primno macropus* : Auct., non Guérin-Méneville, 1836  
European Atlantic N to 40° (50°)
- Lycaeoidea Chevreux, 1913
- Lycaeoidea Chevreux, 1913
- Lycaeopsis* Claus, 1879
- L. themistoides* Claus, 1879 (TM)  
N Atlantic, Mediterranean, Antarctic
- INGOLFIELLIDEA Hansen, 1903
- Ingolfiellidae Hansen, 1903
- Ingolfiella* Hansen, 1903
- I. (Hanseniella) Stock*, 1979) *britannica* Spooner, 1960  
SW England at 45 m
- Metaingolfiellidae Ruffo, 1969  
\* No species within the area
- GAMMARIDEA Latreille, 1803
- Lysianassoidea Dana, 1849
- Lysianassidae Dana, 1849
- Trischizostoma* Boeck, 1861
- T. raschii* Boeck, 1871  
=?? *T. nicagense* (Costa, 1853)  
@ skin of *Etmopterus spinax*  
Skagerrak-W & N Norway-off S Iceland-off SW Ireland
- Sophrosyne* Stebbing, 1888
- S. robertsoni* Stebbing & Robertson, 1891  
W Scotland
- Normanion* Bonnier, 1893
- N. sarsi* Stebbing, 1906  
= *N. quadrimanus* : G.O. Sars, 1895, non (Bate & Westwood, 1868)  
@ skin of fishes (both living & dead)  
S Norway-Trondheim
- N. quadrimanus* (Bate & Westwood, 1868)  
= *N. amblyops* G.O. Sars, 1895  
@ skin of *Etmopterus spinax* & *Melanogrammus aeglefinus*  
Trondheimsfjord-W Scotland-N & W France
- Cheirimedon* Stebbing, 1888
- C. latimanus* (G.O. Sars, 1882)  
?Skagerrak-Stavanger-off SW Iceland
- Podopriionella* G.O. Sars, 1895
- P. norvegica* G.O. Sars, 1895  
Trondheimsfjord
- Opisa* Boeck, 1876
- O. eschrichti* (Krøyer, 1842)  
= *O. typica* (Krøyer, 1846)  
Skagerrak-N Norway-Arctic Seas-Iceland-Faeroes-Ireland
- Lepidepecreum* Bate & Westwood, 1868

- L. longicorne* (Bate & Westwood, 1861)  
= *L. carinatum* Bate & Westwood, 1868  
Kattegatt-S Norway-North Sea-Br. Isles-Portugal-Mediterranean
- L. umbo* (Goës, 1866)  
NW & N Norway-Arctic Seas-Iceland-between Iceland & Faeroes
- L. serratum* Stephensen, 1925  
off SW Iceland-off SW Faeroes
- L. typhlops* Bonnier, 1896  
off SW Faeroes-Bay of Biscay
- Perrierella* Chevreux & Bouvier, 1892
- P. audouiniana* (Bate, 1857)  
Kattegatt-Bohuslän-North Sea-W Norway-Br. Isles-W Africa-Mediterranean
- Ambasia* Boeck, 1871
- A. atlantica* (Milne-Edwards, 1830)  
= *A. danielsseni* Boeck, 1871  
Skagerrak-W & N Norway-Greenland-Iceland-Faeroes-off SW Ireland
- A. murmanica* Brügger, 1905  
N Norway-Murmansk-E Greenland-off SW Faeroes
- Metambasia* Stephensen, 1923
- M. faeroensis* Stephensen, 1925  
off SW Faeroes
- Schisturella* Norman, 1900
- S. pulchra* (Hansen, 1887)  
off NW Scotland-W Greenland
- Aristias* Boeck, 1871
- A. neglectus* Hansen, 1887  
@ *Bolocera tuediae*, *Macandrevia cranium*, other Brachiopods & Sponges  
Kattegatt-Bohuslän-W & N Norway-Shetlands-Br. Isles-Mediterranean
- A. microps* G.O. Sars, 1895  
?Skagerrak-Trondheim-N Norway-off W & SW Iceland-off SW Faeroes
- A. megalops* G.O. Sars, 1895  
\* Possibly the male of *A. microps* G.O. Sars  
Trondheim-N Norway
- A. tumidus* (Krøyer, 1846)  
N Norway-Jan Mayen-Arctic Seas
- A. falcatus* Stephensen, 1923  
off SW Faeroes
- Lysianella* G.O. Sars, 1882
- L. petalocera* G.O. Sars, 1882  
Koster Channel-S, W & N Norway
- Tryphosites* G.O. Sars, 1891
- T. longipes* (Bate & Westwood, 1861)  
Kattegatt-Bohuslän-W & N Norway-Faeroes-Rockall-Canaries-Mediterranean
- T. alleni* Sexton, 1911  
off SW Faeroes-off W Ireland-Bay of Biscay
- Scopelocheirus* Bate, 1856
- S. hopei* (Costa, 1851)  
= *S. kroeyeri* (Bruzellius, 1859)  
= *S. crenatus* Bate, 1857  
\* Lincoln, 1979 synonymized *S. crenatus* with *S. hopei*  
Kattegatt-Bohuslän-North Sea-W & N Norway-Iceland-Br. Isles-W France-Mediterranean
- Tmetonyx* Stebbing, 1906  
= *Hoplonyx* G.O. Sars, 1891 (homonym)
- T. cicada* (O. Fabricius, 1780)  
Öresund-Bohuslän-W & N Norway-Arctic Seas-Iceland-off Shetlands-Br. Isles-Jersey
- T. similis* (G.O. Sars, 1895)  
= *T. leucophthalmus* (G.O. Sars, 1895)  
Kattegatt-Skagerrak-W & N Norway-Arctic Seas-Iceland-Br. Isles-NW France-Mediterranean
- T. acutus* (G.O. Sars, 1895)  
Bohuslän-Skagerrak-N Norway-off SW Faeroes
- T. albidus* (G.O. Sars, 1895)  
N Norway
- T. rotundatus* Chevreux, 1926  
off Lofoten
- T. caeculus* (G.O. Sars, 1895)  
\* Should be moved to a new genus, according to Oleröd, 1987  
Bohuslän-W & N Norway-SW Iceland
- T. longichela* Stephensen, 1925  
off SE Iceland
- T. gracilipes* Stephensen, 1925  
off N Faeroes
- T. trionyx* Stephensen, 1925  
off SW Iceland
- T. norbiensis* Oleröd, 1987  
abyssal depths in Norw. Sea
- Euonyx* Norman, 1867
- E. chelatus* Norman, 1867 (TM)  
@ *Echinus esculentus*  
N Norway-off SW Faeroes-Shetlands-W Scotland-Wales
- Stephonyx* Lowry & Stoddart, 1989
- S. biscayensis* (Chevreux, 1908)  
= *Euonyx biscayensis* Chevreux, 1908 (TM)
- off SW Faeroes-off SW Ireland-Bay of Biscay-S Africa
- S. talismani* (Chevreux, 1919)  
= *Euonyx talismani* Chevreux, 1919  
off SW Faeroes-off SW Ireland
- Kerguelenia* Stebbing, 1888
- K. borealis* G.O. Sars, 1895  
Hardangerfjord-N Norway
- Acidostoma* Lilljeborg, 1865
- A. nodiferum* Stephensen, 1923  
= *Acidostoma obesum* Auctores, non (Bate & Westwood, 1861)  
= *Acidostoma sarsi* Lincoln, 1979  
@ *A. nodiferum* occurs together with *Actinostola callosa*  
Öresund-Bohuslän-North Sea-W & N Norway-Iceland-?off SW Faeroes
- A. laticorne* G.O. Sars, 1879  
Arctic Seas, ?Roscoff
- A. obesum* (Bate & Westwood, 1861)  
= *Acidostoma laticorne* : Oldevig, 1933, non G.O. Sars, 1879  
= *Acidostoma neglectum* Dahl, 1964  
@ *Peachia hastata* & *Cereactis aurantiaca*  
Kattegatt-North Sea-Br. Isles-NW France
- Socarnes* Boeck, 1871
- S. bidenticulatus* (Bate, 1858)  
N Norway-Arctic Seas-Jan Mayen-?Iceland
- S. vahli* (Krøyer, 1838)  
Haugesund-N Norway-Arctic Seas-NE Iceland
- S. erythrophthalmus* Robertson, 1892  
W Br. Isles
- Socarnopsis* Chevreux, 1911
- S. filicornis* (Heller, 1867)  
= *Socarnopsis crenulatus* Chevreux, 1911  
Shetland, W Br. Isles
- Nannonyx* G.O. Sars, 1890
- N. goesii* (Boeck, 1871)  
North Sea
- N. spinimanus* Walker, 1895  
NE France & Br. Isles
- Menigrates* Boeck, 1871
- M. obtusifrons* (Boeck, 1861)  
Skagerrak-North Sea-W & N Norway-Arctic Seas-Iceland-Br. Isles-Guernsey
- Menigratopsis* Dahl, 1945
- M. svennilsoni* Dahl, 1945  
Öresund-North Sea
- Tetronychia* Stephensen, 1923
- T. abyssalis* Stephensen, 1923  
off S Iceland
- Koroga* Holmes, 1909
- K. megalops* Holmes, 1909  
?off S Iceland-off S Alaska
- Centromedon* G.O. Sars, 1895
- C. pumilus* (Lilljeborg, 1865)  
Bohuslän-Skagerrak-W & N Norway-Arctic Seas & Iceland
- C. typhlops* (G.O. Sars, 1885)  
off NE Iceland-between Jan Mayen & Finnmark
- C. calcaratus* (G.O. Sars, 1885)  
between Iceland & Jan Mayen-Arctic Seas
- Ichnopus* Costa, 1853
- I. spinicornis* Boeck, 1861  
Bohuslän-Trondheim-North Sea-Br. Isles-Mediterranean
- Lysianassa* Milne-Edwards, 1830
- L. plumosa* Boeck, 1871  
Haugesund-Trondheim-Shetlands-Br. Isles-NW France-Senegal-Mediterranean
- L. ceratina* (Walker, 1889)  
W Br. Isles
- L. insperata* Lincoln, 1979  
Guernsey
- Hippomedon* Boeck, 1871
- H. holboelli* (Krøyer, 1846)  
Skagerrak, ?N Norway-Arctic Seas, e.g. N Greenland & Iceland-off N Faeroes
- H. denticulatus* (Bate, 1857)  
Öresund-Bohuslän-W & N Norway-Kolafjord-Shetlands-Br. Isles-Mediterranean
- H. propinquus* G.O. Sars, 1895  
?Bohuslän-Skagerrak-W & N Norway-Arctic Seas-Iceland
- H. robustus* G.O. Sars, 1895  
Skagerrak-W & N Norway-Iceland-Azores
- H. striolatus* Stephensen, 1923  
off SW Faeroes
- H. reticulatus* Stephensen, 1923  
off SW Faeroes
- H. nasutus* Stephensen, 1923  
off SW Iceland

- H. frigidus* Stephensen, 1923  
off NE Iceland
- H. serratipes* Stephensen, 1923  
off S Iceland
- Thoriella* Stephensen, 1915
- T. islandica* Stephensen, 1915  
off S Iceland
- Crybelocephalus* Tattersall, 1906
- C. megalurus* Tattersall, 1906  
off S Iceland-off W Ireland
- Cyclocaris* Stebbing, 1888
- C. guilelmi* Chevreux, 1899  
= *Cyclocaris feroensis* Norman, 1900  
off Lofoten-Arctic Seas-off E Jan Mayen-off N & S Faeroes
- Cyphocaris* Lütken, in Boeck, 1871
- C. anonyx* Boeck, 1871  
= *Cyphocaris micronyx* Chevreux, 1900  
off Lofoten-off W & S Iceland-off NW & SW Faeroes-off N & W Ireland
- C. bouvieri* Chevreux, 1916  
off N Faeroes-N Atlantic
- C. richardii* Chevreux, 1905  
?European Atlantic
- Metacyphocaris* Tattersall, 1906
- M. helgae* Tattersall, 1906  
off S Iceland-off SW Ireland
- Paracyphocaris* Chevreux, 1905
- P. praedator* Chevreux, 1905  
off S Iceland-Bay of Biscay-Azores
- Amaryllis* Haswell, 1879
- A. pulchella* Bonnier, 1896  
off S Iceland-off SW Faeroes-Bay of Biscay
- Onisimus* Boeck, 1871 (™ *Anonyx litoralis* Krøyer, 1845)  
= *Onesimus* Boeck, 1876 (invalid emendation of *Onisimus*)  
= *Pseudalibrotus* Della Valle, 1893 (™ *Anonyx litoralis* Krøyer, 1845)  
= *Paronesimus* Stebbing, 1894 (™ *Paronesimus barentsi* Stebbing, 1894)  
= *Boeckosimus* Barnard, 1969 (™ *Anonyx edwardsii* Krøyer, 1846)  
\* Lowry & Stoddart, 1992 synonymized the genera above
- O. affinis* Hansen, 1887  
Jan Mayen-Arctic Seas
- O. barentsi* (Stebbing, 1894)  
off W Novaja Zemlja, circum-Arctic
- O. brevicaudatus* Hansen, 1887  
N Greenland
- O. edwardsii* (Krøyer, 1846)  
= *Anonyx edwardsii* Krøyer, 1846  
Kattegat-off W Norway-N Norway-Arctic Seas-Jan Mayen-Iceland
- O. glacialis* (G.O. Sars, 1900)  
= *Pseudalibrotus glacialis* G.O. Sars, 1900  
Arctic Seas, e.g. N Greenland-Jan Mayen-Iceland-Orkneys
- O. leucopsis* G.O. Sars, 1879  
between Norway & Iceland-between Iceland & Jan Mayen
- O. litoralis* (Krøyer, 1845)  
= *Anonyx litoralis* Krøyer, 1845 (™)  
N Norway-Arctic Seas, including NE Greenland & Iceland
- O. nansenii* (G.O. Sars, 1900)  
Arctic Seas-Iceland
- O. normani* G.O. Sars, 1891  
@ *Bolocera tuediae*  
\* Should possibly be placed in a new genus, according to Lowry & Stoddart, 1992.  
Oslofjord-Skagerrak-between S Norway & Shetlands-W & N Norway-Arctic Seas
- O. plautus* (Krøyer, 1845)  
= *Anonyx plautus* Krøyer, 1845  
Bohuslän-Haugesund-Arctic Seas-Jan Mayen-Iceland
- O. sextoni* Chevreux, 1926  
off Lofoten Islands
- Anonyx* Krøyer, 1838  
= *Chironesimus* G.O. Sars, 1895  
= *Lakota* Holmes, 1908
- A. nugax* (Phipps, 1774)  
= *Lakota chelata* Chevreux, 1926  
off Lofoten-N Norway-Faeroes-Arctic Seas
- A. lilljeborgi* Boeck, 1871  
Norwegian Sea
- A. sarsi* Steele & Brunel, 1968  
= *A. nugax* Sars, 1891, non (Phipps, 1774)  
Bohuslän?
- A. debruyni* Hoek, 1882  
N Norway-Arctic Seas
- A. pacificus* Gurjanova, 1962  
= *A. nugax pacificus* Gurjanova, 1962  
N Greenland
- Tryphosella* Bonnier, 1893  
= *Tryphosa* : G.O. Sars, 1891, non Boeck, 1871
- T. sarsi* Bonnier, 1893
- = *T. nana* (G.O. Sars, 1891), non Boeck, 1861  
Kattegat-Bohuslän-N Norway-Shetlands-Plymouth
- T. hoeringii* (Boeck, 1871)  
Bohuslän-North Sea-W & N Norway-off SW Faeroes-Br. Isles
- T. nanooides* (Lilljeborg, 1865)  
Bohuslän-W & N Norway-Arctic Seas-Iceland-Shetlands
- T. angulata* (G.O. Sars, 1895)  
Trondheim-N Norway-Iceland
- T. compressa* (G.O. Sars, 1895)  
Iceland-Arctic Seas
- T. spitzbergensis* (Chevreux, 1926)  
?N Norway-Spitsbergen
- T. schneideri* (Stephensen, 1921)  
N Norway-Iceland-Arctic Seas
- T. triangula* (Stephensen, 1925)  
Iceland, Arctic Seas
- T. rotundata* (Stephensen, 1925)  
off S Iceland-off SW Faeroes
- T. insignoides* (Stephensen, 1925)  
off SW Faeroes
- T. abyssalis* (Stephensen, 1925)  
off S Jan Mayen (deep water)-abyssal depths of Norw. Sea
- T. abyssi* Norman, 1900  
Faeroe Channel
- Orchomene* Boeck, 1871  
= *Tryphosa* Boeck, 1871  
= *Orchomenella* G.O. Sars, 1891  
= *Orchomenopsis* G.O. Sars, 1895
- O. serrata* (Boeck, 1861)  
Bohuslän-W & N Norway-Arctic Seas-Iceland-Faeroes
- O. humilis* (Costa, 1853)  
= *O. baei* G.O. Sars, 1882  
= *O. hansenii* Meinert, 1890  
Kattegat-Bohuslän-Norwegian Sea
- O. pectinata* G.O. Sars, 1882  
Bergen-N Norway-Arctic Seas-Iceland
- O. crispata* (Goës, 1866)  
Bohuslän-N Norway-Spitsbergen-?off S Iceland
- O. amblyops* G.O. Sars, 1895  
Koster area-Skagerrak-W & N Norway-off NE Iceland-off N & SW Faeroes
- O. thorii* Stephensen, 1923  
off SW Faeroes
- O. laevipes* Stephensen, 1923  
off W Iceland
- O. faeroensis* Stephensen, 1923  
off SW Faeroes
- O. obtusa* (G.O. Sars, 1895)  
= *Orchomenopsis o.* G.O. Sars, 1895  
Bohuslän-Skagerrak-SW & N Norway-?off SW Iceland
- O. minuta* (Krøyer, 1846)  
= *Orchomenella m.* : G.O. Sars, 1895  
Öresund-Bohuslän-W & N Norway-Arctic Seas-Iceland
- O. nana* (Krøyer, 1846)  
= *Tryphosa ciliata* G.O. Sars, 1882  
= *Orchomenella n.* : Auctt.  
Öresund-Bohuslän-S & N Norway-North Sea-?Murmansk-Br. Isles-Spain-Senegal
- O. pinguis* Boeck, 1871  
= *Orchomenella p.* : G.O. Sars, 1895  
Skagerrak-Helgoland-W & N Norway-Arctic Seas-Mediterranean
- O. groenlandica* (Hansen, 1887)  
= *Orchomenella g.* : G.O. Sars, 1895  
N Norway-Arctic Seas
- O. crenata* (Chevreux & Fages, 1925)  
Roscoff
- O. similis* Chevreux, 1912  
W France
- O. abyssorum* Stebbing, 1888  
?European Atlantic
- Eurytenes* Lilljeborg, 1865  
= *Eurythenes* Barnard, 1932  
= *Kalius* Chevreux, 1905
- E. gryllus* (Mandt, 1822)  
= *Kalius obesus* Chevreux, 1905  
N Norway-Arctic Sea-off S Iceland-between Iceland & Greenland
- Parachevreuxiella* Andres, 1987
- P. lobata* Andres, 1987  
?European Atlantic
- Uristidae Hurley, 1963
- Uristes* Dana, 1849
- U. umbonatus* (G.O. Sars, 1882)  
W Skagerrak-W & N Norway-Iceland-off SW Faeroes
- Leucothoidea Dana, 1852
- Pleustidae Buchholz, 1874  
= Paramphithoidea

*Pleustes* Bate,1858

*P. panopla* (Krøyer,1838)  
Bergen-N Norway-Arctic Seas

*Pleusymtes* Barnard,1969

*P. glaber* (Boeck,1861)  
= *Amphitopsis glaber* Boeck,1861  
= *Parapleustes glaber* : G.O. Sars,1893  
= *Sympleustes glaber* : Auct.  
Kattegat-Bohuslän-S & N Norway-Arctic Seas-Iceland-Br. Isles-Bay of Biscay

*Parapleustes* Buchholz,1874

*P. gracilis* Buchholz,1874  
N Norway-Spitsbergen-Arctic Seas-Iceland-E Greenland

*P. assimilis* (G.O. Sars,1882)  
= *Paramphithoe assimilis* G.O. Sars,1882  
Bohuslän-Helgoland-W & N Norway-Iceland-Greenland-Faeroes-N Br. Isles

*P. pulchellus* (Krøyer,1846)  
Bohuslän-N Norway-Arctic Seas & Iceland

*P. monocuspis* G.O. Sars,1895  
\* According to Lincoln,1979, is this a synonym of *P. bicuspis* (Krøyer)  
Bohuslän-Skagerrak-N Norway-Greenland-Iceland-Faeroes

*P. bicuspis* (Krøyer,1838)  
= *Amphithoe bicuspis* Krøyer,1838  
Kattegat-Bohuslän-N Norway-Arctic Seas-Iceland-Br. Isles-NW France

*Stenopleustes* G.O. Sars,1893  
= *Sympleustes* Stebbing,1899

*S. malmgreni* (Boeck,1871)  
\* Often on *Paragorgia rosea*  
Koster Channel-Skagerrak-N Norway-Iceland & Arctic Seas

*S. nodifer* (G.O. Sars,1882)  
= *Amphitopsis nodifera* G.O. Sars,1882  
Bohuslän-S Norway-Trondheim-Faeroes-Br. Isles-NW France

*S. latipes* (M. Sars,1858)  
= *Amphithoe latipes* M. Sars,1858  
@ dead ?*Paramuricea*  
Bohuslän-North Sea-N Norway-Iceland & Arctic Seas-Br. Isles-Bay of Biscay

*S. eldingi* Gurjanova,1930  
Tromsø area-Barents Sea

*S. pulchellus* (G.O. Sars,1876)  
N Norway-off E Iceland-Arctic Seas

*S. megacheir* (Walker,1897)  
=? *S. grandimanus* (Chevreux,1887)  
off W Iceland-slope of Engl. Channel

Colomastigidae Stebbing,1899

*Colomastix* Grube,1861  
= *Cratippus* Bate,1862  
= *Exunguia* Norman,1869

*C. pusilla* Grube,1861  
= *Cratippus tenuipes* Bate,1862  
= *Exunguia stillipes* Norman,1869  
\* usually associated with Sponges  
Atlantic coast of Europe-Mediterranean

Amphiloichidae Boeck,1872

*Stegoplax* G.O. Sars,1882

*S. longirostris* G.O. Sars,1882  
Hardangerfjord-off Lofoten-off SW Iceland

*Amphiloichus* Bate,1862

*A. tenuimanus* Boeck,1871  
W Skagerrak-W & N Norway-off Iceland-Br. Isles

*A. manudens* Bate,1862  
= *A. boeckii* Meinert,1???  
Öresund-Bohuslän-W & N Norway-Murmansk-Iceland-Faeroes-Bay of Biscay

*A. borealis* Enequist,1949  
Skagerrak

*A. spencebatei* (Stebbing,1876)  
= *Probolium Spence-Batei* Stebbing,1876  
=? *A. brunneus* Della Valle,1893  
= *A. anomalus* Chevreux,1900  
Irish Sea-Mediterranean

*A. neapolitanus* Della Valle,1893  
= *A. melanops* Walker,1894  
Liverpool Bay-Mediterranean

*Amphiloichopsis* Stephensen,1925

*A. hamatus* Stephensen,1925  
off S Jan Mayen-off W Greenland-?off W, SW & S Iceland-?off N Faeroes

*Amphiloichoides* G.O. Sars,1892

*A. boeckii* G.O. Sars,1892  
= *A. odontonyx* : G.O. Sars,1892, non (Boeck,1871)  
Bohuslän-W Norway & North Sea

*A. serratipes* (Norman,1869)  
= *Probolium serratipes* Norman,1869  
Öresund-Bohuslän-Skagerrak-Trondheim-Shetlands-W Scotland

*Paramphiloichoides* Lincoln,1979

*P. odontonyx* (Boeck,1871)  
= *Amphiloichus odontonyx* Boeck,1871  
= *Amphiloichoides pusillus* G.O. Sars,1892  
= *Amphiloichoides odontonyx* : G.O. Sars,1895  
?Öresund-?Bohuslän-Skagerrak-W & N Norway-Br. Isles

*P. intermedius* (Scott,1896)  
= *Amphiloichoides intermedius* Scott,1896  
Gullmar, Firth of Forth

*Gitanopsis* G.O. Sars,1895

*G. bispinosa* (Boeck,1871)  
= *Amphiloichus bispinosus* Boeck,1871  
Oslofjord-?Skagerrak-W & N Norway-Clyde Sea

*G. inermis* (G.O. Sars,1882)  
= *Amphiloichus inermis* G.O. Sars,1882  
N Norway-Arctic Seas-Jan Mayen-E Iceland-Northumberland

*G. arctica* G.O. Sars,1895  
N Norway-Arctic Seas-E Iceland-S Greenland

*Gitana* Boeck,1871

*G. rostrata* Boeck,1871  
Skagerrak-W & N Norway

*G. abyssicola* G.O. Sars,1895  
?Koster area-W & N Norway-?Scotland

*G. sarsi* Boeck,1871  
= *Amphiloichus sabrinae* Stebbing,1878  
Öresund-Bohuslän-W & N Norway-Spitsbergen-Br. Isles-Mediterranean

Cyproideidae

*Peltacoxa* Catta,1875

*P. dammoniensis* (Stebbing,1885)  
= *Cyproidea dammoniensis* Stebbing,1885  
W British Isles

*P. brevisrostris* (Scott & Scott,1893)  
= *Cyproidea brevisrostris* Scott & Scott,1893  
W Br. Isles & Bay of Biscay

Leucothoidae Dana,1852

*Leucothoe* Leach,1814

*L. lilljeborgii* Boeck,1861  
= *Leucothoe imparicornis* Norman,1889  
Öresund-Bohuslän-North Sea

*L. spinicarpa* (Abildgaard,1789)  
= *Gammarus spinicarpus* Abildgaard,1789  
= *Leucothoe articularia* Leach,1814  
Öresund-Bohuslän-W & N Norway-Iceland-Br. Isles-N France

*L. procera* Bate,1857  
= *L. richiardi* : Lincoln,1979, non Lessona,1865  
Engl. Channel, Scotland, Ireland

*L. incisa* Robertson,1892  
Scotch North Sea coast-Engl. Channel-Mediterranean

Laphystiopsidae Stebbing,1899

*Laphystiopsis* G.O. Sars,1893

*L. planifrons* G.O. Sars,1893  
@ various fish species (half parasite)  
Bohuslän-N Norway-Iceland-off Faeroes

Cressidae Stebbing,1899

*Cressa* Boeck,1871  
= *Danaia* Bate,1857 (Homonym according to Lincoln,1979, but Neave,1939 does not mention an earlier name)

*C. dubia* (Bate,1857)  
= *Danaia dubia* Bate,1857  
= *Cressa sholdtei* Boeck,1871  
Öresund-Bohuslän-Trondheim-Br. Isles & W Mediterranean

*C. minuta* Boeck,1871  
Bohuslän-Skagerrak-W Norway-Iceland-E Scotland

*C. abyssicola* (G.O. Sars,1885)  
between Norway & Bear Island-off N Iceland-W Greenland

*C. carinata* Stephensen,1931  
Faeroes-Iceland-off S Jan Mayen-W Greenland

*C. quinquedentata* Stephensen,1931  
off SW Iceland

*Cressina* Stephensen,1931

*C. monocuspis* Stephensen,1931  
off SW Faeroes-off S Iceland-off W Greenland

Stenothoidae Boeck,1871

*Metopella* G.O. Sars,1892

*M. longimana* (Boeck,1871)  
Oslofjord-Skagerrak-Haugesund-Iceland-Greenland

*M. neglecta* (Hansen,1887)  
Haugesund-Greenland

*Hardametopa* Barnard & Karaman,1991

*H. nasuta* (Boeck,1871)  
= *Metopa nasuta* Boeck,1871  
= *Metopella nasuta* : Auct.  
Bohuslän-N Norway-N Br. Isles-Iceland-W Greenland

*Proboloidea* Della Valle,1893

*P. gregarius* (G.O. Sars,1882)  
Bohuslän-W & N Norway-Normandie-Bay of Biscay

*P. calcaratus* (G.O. Sars,1882)  
Skagerrak-North Sea-N Norway

- P. glacialis* (Krøyer, 1842)  
Iceland-Spitsbergen-Greenland
- Parametopa* Chevreux, 1901
- P. kervillei* Chevreux, 1901  
= *Metopa sarniensis* Norman, 1907  
Engl. Channel
- Metopa* Boeck, 1871
- M. quadrangulata* Reibish, 1905-1906  
Skagerrak
- M. robusta* G.O. Sars, 1895  
Skagerrak-N Norway-Greenland-off Faeroes-Scotland
- M. propinqua* G.O. Sars, 1892  
Bohuslän-North Sea-W & N Norway-Barents Sea-Scotland
- M. alderi* (Bate, 1857)  
= *Montagua alderi* Bate, 1857  
= *Metopa spectabilis* G.O. Sars, 1879  
@ may be found on *Tima bairdii*  
Kattegatt-Bohuslän-North Sea-W & N Norway-Iceland-N Br. Isles
- M. affinis* Boeck, 1871  
Oslofj.-W & N Norway
- M. norvegica* (Liljeborg, 1851)  
= *Leucothoe norvegica* Liljeborg, 1851  
= *Montagua pollexiana* Bate, 1857  
Bohuslän-North Sea-W & N Norway-Iceland-Shetlands-Br. Isles
- M. boeckii* G.O. Sars, 1895  
Skagerrak-Haugesund-N Norway-Iceland
- M. longicornis* Boeck, 1871  
Oslofjord-Skagerrak-Lofoten-W Greenland
- M. pusilla* G.O. Sars, 1892  
Belt Sea-Bohuslän-Trondheim-Arctic Seas-Wales
- M. borealis* G.O. Sars, 1882  
@ may be found on *Phialidium* sp.  
Skagerrak-N Norway-Br. Isles-Engl. Channel
- M. soelsbergi* Schneider, 1884  
@ *Actinobla* (= *Metridium*)  
Øresund-Kattegatt-Hardangerfjord-Tromsø-Iceland-Clyde Sea
- M. bruzelii* (Grøes, 1866)  
Kattegatt-Bohuslän-North Sea-W & N Norway-Spitsbergen-Iceland-Faeroes-Br. Isles
- M. tenuimana* G.O. Sars, 1892  
W Norway-Shetlands
- M. leptocarpa* G.O. Sars, 1895  
Christiansund-off SW Iceland
- M. sinuata* G.O. Sars, 1895  
N Norway-Spitsbergen-Iceland-off W Faeroes-between Norway & Scotland-Davis Strait
- M. invalida* G.O. Sars, 1895  
N Norway-Iceland
- M. latimana* Hansen, 1887  
Scotland-Greenland
- M. palmata* G.O. Sars, 1895  
Iceland, Hammerfest
- M. glacialis* (Krøyer, 1842)  
= *Proboloides glacialis* (Krøyer, 1842)  
Iceland-Arctic Seas
- M. abyssalis* Stephensen, 1931  
off N Faeroes-off NE Iceland
- Stenula* Barnard, 1962
- S. rubrovittata* (G.O. Sars, 1882)  
= *Metopa r.* & *Stenothoides latipes* Chevreux & Fage, 1925  
@ typically associated with *Pagurus bernhardus* (Linnaeus)  
Kattegatt-Bohuslän-North Sea-W & N Norway-Faeroes-Shetlands-Br. Isles-Bay of Biscay
- Stenothoe* Dana, 1852
- S. monoculoides* (Montagu, 1815)  
Kattegatt-Bohuslän-N Norway-Iceland-Faeroes-Azores-Senegal-Black Sea
- S. tenella* G.O. Sars, 1882  
Bergen-Trondheim
- S. megacheir* (Boeck, 1871)  
Bohuslän-W & N Norway
- S. microps* G.O. Sars, 1895  
Skagerrak-North Sea
- S. marina* (Bate, 1856)  
@ *Alcyonium digitatum*  
Bohuslän-North Sea-N Norway-?Faeroes-W Africa-Mediterranean
- S. brevicornis* G.O. Sars, 1882  
S-N Norway
- S. ? macrophthalma* Stephensen, 1931  
off SW Faeroes
- S. ? aequicornis* Stephensen, 1931  
between Faeroes & Iceland
- S. valida* Dana, 1852  
Engl. Channel-Mediterranean
- S. tergestina* (Nebeski, 1881)  
= *S. spinimana* Chevreux, 1911  
Engl. Channel-Mediterranean
- S. crassicornis* Walker, 1896
- Isle of Man
- S. setosa* Norman, 1900  
Plymouth
- S. gallensis* Walker, 1904  
= *S. cattaui* Stebbing, 1906  
Engl. Channel-Mediterranean
- S. elachistoides* Myers & McGrath, 1980  
Ireland
- Anamixidae* Stebbing, 1897  
\* No species within the area?
- Maxillipiidae* Ledoyer, 1973  
\* No species within the area?
- Pagetinidae* K.H. Barnard, 1932  
\* No species within the area?
- Nihotungidae* Barnard, 1972  
\* No species within the area?
- Thaumetelsonidae* Gurjanova, 1938  
\* No species within the area?
- Phoxocephaloidea* G.O. Sars, 1891
- Urothoidea* Bousfield, 1969
- Urothoe* Dana, 1852  
= *Egidia* Costa, 1853
- U. elegans* (Bate, 1856)  
= *Gammarus elegans* Bate, 1856  
= *U. norvegica* Boeck, 1861  
Kattegatt-Skagerrak-W & N Norway-Kolafjord-Iceland-Faeroes-Engl. Channel-Senegal-Mediterranean
- U. poseidonis* Reibish, 1905  
= *U. grimaldii* var. *poseidonis*: Schellenberg, 1942  
Kattegatt-North Sea
- U. brevicornis* Bate, 1862  
Engl. Channel, North Sea
- U. marina* (Bate, 1857)  
= *Sulcator marinus* Bate, 1857  
\* Associated with Synaptidae and *Echinocardium cordatum*  
Shetlands-Bay of Biscay
- U. pulchella* (Costa, 1853)  
= *Egidia pulchella* Costa, 1853  
E Scotland-Engl. Channel-Mediterranean
- Phoxocephalidae* G.O. Sars, 1891
- Paraphoxus* G.O. Sars, 1891
- P. oculatus* (G.O. Sars, 1879)  
Skagerrak-W & N Norway-Jan Mayen-Iceland-Shetlands-Isle of Man
- P. maculatus* Chevreux, 1900  
?European Atlantic
- Leptophoxus* G.O. Sars, 1895
- L. falcatus* (G.O. Sars, 1882)  
Koster Channel-W & N Norway-W Greenland
- Phoxocephalus* Stebbing, 1888  
= *Phoxus* Krøyer, 1842, *non* Billberg, 1820 (Coleoptera)
- P. holboelli* (Krøyer, 1842)  
= *Phoxus holböllii* Krøyer, 1842  
S Baltic-Bohuslän-N Norway-White Sea-Iceland-W Greenland-Br. Isles
- P. tenuipes* Stephensen, 1925  
off W & SW Iceland-off W Greenland
- Metaphoxus* Bonnier, 1896
- M. pectinatus* (Walker, 1896)  
= *Phoxocephalus pectinatus* Walker, 1896  
Ireland, Engl. Channel-Mediterranean
- M. simplex* (Bate, 1857)  
= *Phoxus simplex* Bate, 1857  
\* Lincoln, 1979 does not comment on his synonymy of this species with *M. pectinatus* (Walker), but Bate's name is the oldest and should be used if they are considered as synonyms.  
Britain
- Parametaphoxus* Gurjanova, 1977
- P. fultoni* (T. Scott, 1890)  
= *Phoxocephalus fultoni* T. Scott, 1890  
= *Metaphoxus fultoni*: Auctt.  
= *Phoxocephalus chelatus* Della Valle, 1893  
Ireland, Engl. Channel-Mediterranean
- Harpiniopsis* Stephensen, 1925
- H. similis* Stephensen, 1925  
off N & NE Iceland-off N Faeroes-off S Jan Mayen-off W Greenland
- Harpinia* Boeck, 1871
- H. mucronata* G.O. Sars, 1879  
North Sea-N Norway-Iceland-off N Faeroes-E Greenland
- H. crenulata* (Boeck, 1871)  
Øresund-Bohuslän-W & N Norway-Iceland-Br. Isles-Bay of Biscay-W Mediterranean
- H. pectinata* G.O. Sars, 1891  
Bohuslän-North Sea-W & N Norway-Br. Isles-W Africa-W Mediterranean
- H. serrata* G.O. Sars, 1879  
N Norway-Arctic Seas, ?Northumberland
- H. antennaria* Meinert, 1890  
= *H. neglecta* G.O. Sars, 1895  
Øresund-Bohuslän-W & N Norway-Barents Sea-Iceland-Br. Isles-Bay of Biscay

- H. propinqua* G.O. Sars, 1895  
Kattegatt, Lofoten-Jan Mayen-off N Spitsbergen-Iceland
- H. plumosa* (Krøyer, 1842)  
Arctic Seas & Iceland
- H. truncata* G.O. Sars, 1895  
Koster Channel-W & N Norway-N Iceland
- H. abyssi* G.O. Sars, 1879  
off W Norway-off N Norway-Arctic Seas-off E Iceland-off Faeroes
- H. laevis* G.O. Sars, 1891  
Kattegatt-Bohuslän-W Norway-Isle of Man-off SW Iceland
- H. amundseni* Gurjanova, 1946  
N Greenland
- Platyschnopidae Barnard & Drummond, 1979  
\* No species within the area?
- Stegocephaloidea Dana, 1952
- Stegocephalidae Dana, 1852
- Andaniexis* Stebbing, 1906  
= *Andania* Boeck, 1871 (homonym of a Lepidoptera)
- A. abyssi* (Boeck, 1871) Stebbing, 1906  
= *Andania abyssi* Boeck, 1871 (TM)  
Bohuslän-Oslofjord-Skagerrak-W & N Norway-Arctic Seas-Rockall-off W Africa
- A. lupus* Berge & Vader, 1997  
N Norway, Faeroe area, Iceland, Arctic
- A. eilae* Berge & Vader, 1997  
Faeroe area, Iceland
- Stegocephalus* Krøyer, 1842
- S. inflatus* Krøyer, 1842 (TM)  
Bohuslän-W & N Norway-Arctic Seas, e.g. N Greenland-S Iceland-Faeroe Channel
- Phippsiella* Schellenberg, 1925
- P. similis* (G.O. Sars, 1891)  
= *Stegocephalus similis* G.O. Sars, 1891  
Skagerrak-W & N Norway-Faeroes-Spitsbergen-Arctic Seas-off NE Iceland-Davis Strait
- P. bioice* Berge & Vader, 1997  
off Iceland
- P. minima* Stephensen, 1925  
W Greenland-?off N Faeroes
- Stegocephaloides* G.O. Sars, 1891
- S. christianiensis* (Boeck, 1871) G.O. Sars, 1891  
= *Stegocephalus christianiensis* (TM)  
Bohuslän-North Sea-W Norway-Trondheimsfjord-Br. Isles-W Mediterranean
- S. auratus* (G.O. Sars, 1883) G.O. Sars, 1891  
Bohuslän-Skagerrak-W Norway-Trondheimsfjord-Iceland-off S Faeroes, Biscay, ?Kara Sea
- S. barnardi* Berge & Vader, 1997  
off SW Iceland in 1042-1407 m, E Mediterranean
- S. wagini* (Gurjanova, 1936) Barnard & Karaman, 1991  
= *Stegocephalopsis wagini* Gurjanova, 1936  
Faeroe area, Iceland, Kara Sea
- Phippsia* Stebbing, 1906  
= *Aspidopleurus* G.O. Sars, 1891 (homonym of a fish)
- P. gibbosa* (G.O. Sars, 1883) Stebbing, 1906  
= *Stegocephalus gibbosus* G.O. Sars, 1883 (TM)  
W & N Norway, off Iceland
- P. roemeri* Schellenberg, 1925  
Faeroe area, Iceland, E Greenland, off N Spitsbergen
- Stegocephalina* Stephensen, 1925
- S. ingolffi* Stephensen, 1925 (TM)  
off SW Iceland
- ?*S. biofar* Berge & Vader, 1997  
off SE Faeroes-off SW Iceland and off E Greenland in 417-1525 m
- ?*S. idae* Berge & Vader, 1997  
off Iceland in 501-1042 m
- Stegocephalopsis* Schellenberg, 1925
- S. ampulla* (Phipps, 1774)  
= *Cancer ampulla* Phipps, 1774 (TM)  
off Iceland, Arctic circumpolar
- Andaniella* G.O. Sars, 1895
- A. pectinata* (G.O. Sars, 1882)  
= *Andania pectinata* G.O. Sars, 1883 (TM)  
Trondheimsfjord-N Norway-Spitsbergen-Arctic Seas-Faeroes-Iceland-off S Jan Mayen-W Greenland-Newfoundland
- Andaniopsis* G.O. Sars, 1895
- A. nordlandica* (Boeck, 1871) G.O. Sars, 1895  
= *Andania nordlandica* Boeck, 1871 (TM)  
Bohuslän-Oslofjord-Skagerrak-W & N Norway
- Metandania* Stephensen, 1925
- M. islandica* Stephensen, 1925  
off W Iceland
- Euandania* Stebbing, 1899
- E. gigantea* (Stebbing, 1883) Stebbing, 1899  
= *Andania gigantea* Stebbing, 1883 (TM)  
\* Pelagic species
- Faeroe area, off Rockall, "cosmopolitan"
- Iphimediidae Boeck, 1871  
= Acanthonotozomatidae Stebbing, 1906  
= Paramphithoidea Stebbing, 1906
- Clenardopsis* Barnard, 1916
- C. carinata* Barnard, 1916  
off SW Faeroes-S Africa
- Epimeria* Costa, 1851
- E. parasitica* (M. Sars, 1859)  
@ *Parastichopus tremulus* & *Porania pulvillus* (semiparasitic feeder of outer tissues)  
Oslofjord-Skagerrak-W & N Norway & North Sea-Bay of Biscay
- E. loricata* G.O. Sars, 1879  
off W Norway & Shetlands-N Norway-Arctic Seas & Iceland
- E. cornigera* (J.C. Fabricius, 1779)  
= *Acanthonotus testudo* Bate, 1862  
= *Acanthonotus owenii* Bate & Westwood, 1863  
Bohuslän-North Sea-N Norway-Br. Isles-Iceland-Mediterranean
- E. tuberculata* G.O. Sars, 1893  
?Bohuslän-Skagerrak-W & N Norway-Scotland-Iceland
- Paramphithoe* Bruzelius, 1859
- P. hystrix* (J.C. Ross, 1835)  
W & N Norway-Arctic Seas-off NW Iceland
- Iphimedia* Rathke, 1843  
= *Panoploea* Thompson, 1880
- I. minuta* G.O. Sars, 1882  
Öresund-Bohuslän-North Sea-N Norway-Iceland-Br. Isles-Senegal-Cape Verde
- I. obesa* Rathke, 1843  
Öresund-Bohusl-W & N Norway-Faeroes-Br. Isles & N France, Mediterranean
- I. eblanae* Bate, 1857  
Ireland-Atl. coast of France-Mediterranean
- I. spatula* Myers, McGrath & Costello, 1987  
Atl. coast of France-Shetlands
- I. nexa* Myers, McGrath & Costello, 1987  
Atl. coast of France-Ireland
- I. perplexa* Myers, McGrath & Costello, 1987  
W Scotland-W Ireland
- Odius* Lilljeborg, 1865
- O. carinatus* (Bate, 1862)  
Skagerrak-N Norway-Arctic Seas-Engl. Channel
- Acanthonotozoma* Boeck, 1876
- A. cristatum* (J.C. Ross, 1835)  
N North Sea-Norw. Sea-N Norway-Arctic Seas & Iceland
- A. serratum* (O. Fabricius, 1780)  
?Bohuslän-?Skagerrak-N Norway-Arctic Seas & Iceland
- A. rusanovae* Bryazgin, 1974  
NW Norway-Iceland-Arctic Seas
- A. sinuatum* Just, 1978  
off W Norway-Lofoten-Jan Mayen & Arctic Seas
- Lafystiidae G.O. Sars, 1893
- Lafystius* Krøyer, 1842
- L. sturionis* Krøyer, 1842  
@ various marine fishes, especially cod, *Raja batís* & *Lophius piscatorius*  
Öresund-Trondheim-Iceland-Br. Isles-Mediterranean
- Phliantidae Stebbing, 1899, emend. ?Barnes
- Pereionotus* Bate & Westwood, 1862
- P. testudo* (Montagu, 1808)  
Engl. Channel-Mediterranean
- Ochlesidae Stebbing, 1910  
\* No species within the area?
- Synopioidea Dana, 1853
- Synopidae Dana, 1853  
= Syrrhoidea G.O. Sars, 1895  
= Tironidae Stebbing, 1906
- Syrrhoites* G.O. Sars, 1895
- S. serratus* (G.O. Sars, 1879)  
Skagerrak-Lofoten-Arctic Seas-off SW Iceland
- S. pusillus* Enequist, 1950  
Skagerrak-W Norway
- Syrrhoë* Goës, 1866
- S. crenulata* Goës, 1866  
Oslofjord-Skagerrak-N Norway-Arctic Seas & Iceland
- S. affine* Chevreux, 1908  
?W Norway, Britain
- Austrosyrrhoë* K.H. Barnard, 1925
- A. fimbriatus* (Stebbing & Robertson, 1891)  
W Britain & Ireland-off N Spitsbergen
- A. ? septentrionalis* Stephensen, 1931  
between Iceland & Greenland-W Greenland-?off E Jan Mayen
- Tiron* Lilljeborg, 1865

- T. spiniferum* (Stimpson, 1853)  
= *T. acanthurus* Liljeborg, 1865  
?Kattegatt-Bohuslän-Skagerrak-W & N Norway-Arctic Seas-Iceland-Br. Isles
- Bruzelia** Boeck, 1871
- B. typica* Boeck, 1871  
?Skagerrak-W & N Norway-Br. Isles
- B. tuberculata* G.O. Sars, 1882  
Trondheim-off Lofoten-off SW Faeroes
- B. dentata* Stephensen, 1931  
off N Faeroes-off E & NE Iceland-off S Jan Mayen
- Argissidae Walker, 1904
- Argissa** Boeck, 1871
- A. hamatipes* (Norman, 1869)  
= *A. typica* Boeck, 1871  
Öresund-Bohuslän-N Norway-Greenland-Iceland-Shetlands-N Br. Isles
- Ampeliscoidea Bate, 1861
- Ampeliscidae Bate, 1861
- Haploops** Liljeborg, 1856
- H. tubicola* Liljeborg, 1856  
= *H. carinata* Liljeborg, 1856  
= *H. spinosa* Shoemaker, 1931  
Öresund-Bohuslän-North Sea- Norway- Spitzbergen, Br. Isles-W Bretagne, Iceland, E & W Greenland, Barents Sea, Kara Sea
- H. setosa* Boeck, 1871  
= *H. robusta* G.O. Sars, 1891  
Bohuslän-Skagerrak-W & N Norway-Spitzbergen-Iceland-Faeroes-E Scotland, E & W Greenland
- H. tenuis* Kannerworf, 1966  
Öresund-Kattegatt, W & N Norway
- H. vallifera* Stephensen, 1925  
off S, SW & W Iceland
- H. laevis* Hoek, 1882  
Barents Sea, Kara Sea, Davis Strait
- H. nira* Kaïm-Malka, 1976  
NW Bretagne - Morocco, Mediterranean
- Byblis** Boeck, 1871
- B. erythrops* G.O. Sars, 1882  
W & N Norway, Ireland
- B. abyssii* G.O. Sars, 1879  
off W & N Norway-S Jan Mayen-Arctic Seas
- B. gaimardii* (Krøyer, 1846)  
Öresund-Bohuslän-North Sea-W & N Norway-Arctic Seas-Iceland-Br. Isles
- B. longicornis* G.O. Sars, 1895  
off Lofoten area-off N Norway-Arctic Seas
- B. crassicornis* Metzger, 1875  
Oslofjord-W & N Norway & Iceland
- B. affinis* G.O. Sars, 1895  
W Norway
- B. minuticornis* G.O. Sars, 1895  
off N Faeroes-off N Iceland-Arctic Seas
- B. arcticus* Just, 1970  
N & E Greenland
- Ampelisca** Krøyer, 1842
- A. odontoplax* G.O. Sars, 1895  
W Skagerrak-Hardangerfjord-N Norway-off W & S Iceland-between Faeroes & Shetlands, Biscay
- A. eschrichti* Krøyer, 1842  
?Bohuslän-Bergen-N Norway-Arctic Seas-Iceland-Faeroe Channel-off Ireland
- A. brevicornis* (Costa, 1853)  
Öresund-Bohuslän-North Sea-W Norway-Lofoten-Br. Isles-S Africa-Mediterran.
- A. macrocephala* Liljeborg, 1852  
Öresund-Bohuslän-N North Sea-W & N Norway-Iceland-Arctic Seas-W Ireland
- A. gibba* G.O. Sars, 1882  
Kattegatt-Bohuslän-W & N Norway-off S Iceland-off SW Faeroes-off W Ireland-off Engl. Channel
- A. amblyops* G.O. Sars, 1895  
Bohuslän-W Norway-off W Greenland
- A. pusilla* G.O. Sars, 1895  
\* synonymized with *A. amblyops* by Enequist, 1949  
Bohuslän-W & N Norway
- A. anomala* G.O. Sars, 1882  
Bohuslän-S & W Norway-Angola
- A. spinipes* Boeck, 1860  
Kattegatt-Bohuslän-S Norway-North Sea-Lofoten-Faeroes-Shetlands-Br. Isles-Bretagne-Senegal
- A. aequicornis* Bruzelius, 1859  
Bohuslän-North Sea-W & N Norway-Iceland-Faeroes-Br. Isles-Canaries-Gulf of Guinea
- A. typica* (Bate, 1856)  
Kattegatt-Bohuslän-North Sea-W & N Norway-Br. Isles-Senegal-Mediterranean
- A. tenuicornis* Liljeborg, 1855  
\* synonymized with *A. diadema* by Enequist, 1949  
Öresund-Bohuslän-North Sea-Trondheim-Br. Isles-Senegal-Mediterranean
- A. diadema* (Costa, 1853)  
= *Ampelisca assimilis* Boeck, 1870  
Kattegatt-Bohuslän-North Sea-Trondheim-Br. Isles-W Africa-Mediterran-Black Sea
- A. spinifer* Reid, 1951  
Ireland-Mediterranean & Liberia
- A. spooneri* Dauvin & Bellan-Santini, 1982  
Firth of Clyde, Engl. Channel -Dakar area
- A. dalmatina* Karaman, 1975  
Ireland, Mediterranean
- A. provincialis* Bellan-Santini & Kaïm-Malka, 1977  
Celtic Sea, Mediterranean
- A. toule Monti* Dauvin & Bellan-Santini, 1982  
Celtic Sea & N Bay of Biscay
- A. armoricana* Bellan-Santini & Dauvin, 1981  
Engl. Channel-Dakar
- A. sarsi* Chevreux, 1887  
N Brittany-Senegal
- A. spinimana* Chevreux, 1887  
N Brittany-Gulf of Guinea
- A. compacta* Norman, 1882  
off S Faeroes-?off S Iceland
- Pontoporeioidea G.O. Sars, 1882
- Haustoriidae Stebbing, 1906
- Haustorius** S. Müller, 1775
- H. arenarius* (Slabber, 1769)  
S Baltic-Isefjord-North Sea-Br. Isles-Atl. coast of France
- Pontoporeiidae G.O. Sars, 1882
- Bathyporeia** Lindström, 1855
- B. guilliamsoniana* (Bate, 1856)  
= *B. norvegica* G.O. Sars, 1891  
Kattegatt-Bohuslän-North Sea
- B. pelagica* (Bate, 1856)  
Isefjord, around British Isles
- B. elegans* Watkin, 1938  
= *B. pelagica*: Sars, 1891  
Kattegatt-Bohuslän-North Sea-Engl. Channel
- B. tenuipes* Meinert, 1877  
Kattegatt-Bohuslän-North Sea-Engl. Channel
- B. gracilis* G.O. Sars, 1891  
North Sea
- B. pilosa* Lindström, 1855  
Bothnian Sea-Isefjord-North Sea
- B. sarsi* Watkin, 1938  
= *B. robertsoni*: G.O. Sars, 1891, non Bate  
Isefjord, Kattegatt-Arctic Seas
- B. nana* Toulmond, 1966  
Engl. Channel, Yorkshire, Ireland
- Pontoporeia** Krøyer, 1842
- P. femorata* Krøyer, 1842  
= *P. sinuata* Ekman, 1913  
Åland-Gullmarsfjord, (W) & N Norway-Arctic Seas, e.g. N Greenland & Iceland
- Monoporeia** Bousfield, 1989
- M. affinis* (Lindström, 1989)  
= *Pontoporeia affinis* Lindström, 1855 (TM)  
= *Pontoporeia weltneri* Ekman, 1913  
Bothnian Bay-Öresund, Lakes in Sw., Norw., Denm. & Finl.
- Priscillina** Stebbing, 1888
- P. armata* (Boeck, 1861)  
?? W Norway-Arctic Seas
- Talitroidea Costa, 1857
- Talitridae Costa, 1857, emend. Bulycheva, 1957
- Talitrus** Bosc, 1802
- T. saltator* (Montagu, 1808)  
S Baltic-Bohuslän-S & W Norway-North Sea-Br. Isles-W Mediterranean-Azores
- Talitroides** Bonnier, 1898
- T. alluaudi* (Chevreux, 1896)  
Sweden & Europe in greenhouses
- Arcitalitrus** Hurley, 1975
- A. dorrieni* (Hunt, 1925)  
= *Talitroides dorrieni* Hunt, 1925  
\* Terrestrial immigrant, presumably from Australasia.  
SW England, Ireland
- Talorchestia** Dana, 1852
- T. deshayesii* (Audouin, in Savigny, 1826)  
Gotska Sandön-Denmark-North Sea-Br. Isles-Black Sea
- Pseudorchestioidea** Bousfield, 1982
- P. brito* (Stebbing, 1891)  
= *Talorchestia brito* Stebbing, 1891  
= *Orchestioidea* Nicolet, 1849 *brito*: Auct.  
W Denmark-SW Norway-S North Sea-Engl. Channel-Morocco-W Mediterranean-Black Sea
- Platorchestia** Bousfield, 1982
- P. platensis* (Krøyer, 1845)  
SE Skåne & Denmark-Bohuslän-Oslofjorden-Br. Isles

- Orchestia* Leach, 1814
- O. gammarellus* (Pallas, 1766)  
Västervik-Bohuslän-W & N Norway-Br. Isles-Azores-Madeira-Canaries-Mediterranean-Black Sea
- O. bottae* Milne-Edwards, 1830?  
Holland-France
- O. cavimana* Heller, 1865  
S North Sea area-Br. Isles-Mediterranean
- O. mediterranea* Costa, 1857  
Br. Isles-Mediterranean
- O. roffensis* (Wildish, 1969)  
Medway Estuary (Br. Isles)
- O. aestuarensis* Wildish, 1987  
Medway Estuary (Br. Isles)
- Hyalidae Bulychева, 1957
- Hyale* Rathke, 1837
- H. prevostii* Milne-Edwards, 1830  
= *H. nilssonii* Rathke, 1843  
Kattegatt-Bohuslän-W & N Norway-Murmansk-Iceland-Faeroes-SW France
- H. pontica* Rathke, 1837  
= *H. lubbockiana*: G.O. Sars, 1890  
Skagerrak-North Sea-N Norway-Br. Isles-Portugal-Mediterranean-Black Sea
- H. stebbingi* Chevreux, 1888  
Shetlands-Mediterranean-Azores-Canaries
- H. perieri* (Lucas, 1846)  
SW Britain-Mediterranean
- H. grimaldii* Chevreux, 1891  
\* Associated with driftwood  
Galway Bay-Mediterranean-Azores
- Dogielinotidae Gurjanova, 1954  
\* No species within the area?
- Hyalellidae Bulychева, 1957  
\* No species within the area?
- Najnidae Barnard, 1972  
\* No species within the area?
- Ceinidae Barnard, 1972  
\* No species within the area?
- Eophliantidae Sheard, 1938  
\* No species within the area?
- Temnophliantidae Griffiths, 1975  
\* No species within the area?
- Kuriidae Barnard, 1964  
\* No species within the area?
- Corophioidea Dana, 1849
- Isaidae Dana, 1853  
= Photidae Barnard, 1969
- Isaea* Milne-Edwards, 1830
- I. montagui* Milne-Edwards, 1830  
\* on mouth-parts of *Maia squinado*  
W Ireland-Mediterranean
- I. elmhirsti* Patience, 1909  
\* on mouth-parts of *Homarus gammarus*  
Ireland-Bay of Biscay
- Microprotopus* Norman, 1867
- M. maculatus* Norman, 1867  
Isefjord, Kattegatt-Bohuslän-Norw. Sea
- M. longimanus* Chevreux, 1887  
Engl. Channel & Atl. coast of France
- Photis* Krøyer, 1842
- P. reinhardi* Krøyer, 1842  
Öresund-Bohuslän-W & N Norway-Iceland & Arctic Seas-Br. Isles
- P. longicaudata* (Bate & Westwood, 1862)  
Öresund-Bohuslän-North Sea
- P. tenuicornis* G.O. Sars, 1882  
N Norway-Iceland & Arctic Seas
- P. pollex* A.O. Walker, 1895  
\* easily confused with *P. reinhardi* (see Myers & McGrath, 1981)  
Atlantic America & Ireland
- Protomeдея* Krøyer, 1842
- P. fasciata* Krøyer, 1842  
Öresund-Bohuslän-W & N Norway-Arctic Seas, e.g. S, W & N Greenland-Iceland-Faeroes-NE Br. Isles
- P. grandimana* Brügger, 1905  
N Norway-Arctic Seas & Iceland
- Gammaropsis* Liljeborg, 1855  
= *Eurytheus* Bate, 1857  
= *Podocerospis* Boeck, 1861  
= *Megamphopus* Norman, 1869
- G. cornuta* (Norman, 1869)  
= *Megamphopus cornutus* Norman, 1869 (™ of *Megamphopus* Norman, 1869)  
Öresund-Bohuslän-Skagerrak-W & N Norway-Br. Isles-Dakar
- G. palmata* (Stebbing & Robertson, 1891)  
= *G. nana* G.O. Sars, 1894  
Kattegatt-Bohuslän-North Sea
- G. melanops* G.O. Sars, 1882  
Öresund-Bohuslän-W & N Norway-Arctic Seas-Iceland-Faeroes
- G. maculata* (Johnston, 1828)  
= *G. erythrophthalma*: G.O. Sars, 1894  
Öresund-Bohuslän-S, W & N Norway-Faeroes-Br. Isles-Senegal-Mediterranean
- G. nitida* (Stimpson, 1853)  
= *Noenia excavata* Bate, 1862  
@ Paguridae  
Öresund-Bohuslän-North Sea-W & N Norway-Iceland-off SE Faeroes-Engl. Channel
- G. sophiae* (Boeck, 1861)  
= *Podocerospis sophiae* Boeck, 1861  
Kattegatt-Bohuslän-W & N Norway-Br. Isles-Canaries-W Mediterranean
- G. lobata* (Chevreux, 1920)  
Eddystone-Ireland-W France
- G. abyssalis* (Stephensen, 1944)  
off SW Faeroes
- Bonnierella* Chevreux, 1900
- B. abyssi* (Chevreux, 1887)  
off Faeroes-off W Africa
- Goesia* Boeck, 1871
- G. depressa* (Goës, 18??)  
N Greenland
- Ischyroceridae Stebbing, 1899
- Parajassa* Stebbing, 1899  
= *Janassa* Boeck, 1871 (homonym)
- P. pelagica* (Leach, 1814)  
= *Jassa pelagica* Leach, 1814  
= *Podocerus capillatus* Rathke, 1843  
Kalmarsund, Bohuslän-W & N Norway-Iceland-Faeroes-Shetlands-Br. Isles-Portugal
- Jassa* Leach, 1814
- J. falcata* (Montagu, 1808)  
= *Cancer (Gammarus) falcatus* Montagu, 1808 (™)  
= *J. pulchella* Leach, 1814  
Kalmarsund, Kattegatt-Gullmarfjord-Väderöarna-Frederikshavn-Helgoland-W & N Norway-Iceland-Faeroes-Br. Isles-Spain-Portugal
- J. pusilla* (G.O. Sars, 1894)  
= *Podocerus minutus* G.O. Sars, 1882 (Nomen oblitum)  
= *Podocerus pusillus* G.O. Sars, 1894  
= *Podocerus odontomyx* G.O. Sars, 1894  
Gullmarfjord-Väderöarna-W & N Norway-Iceland-Faeroes-Br. Isles-W France-NW Spain
- J. marmorata* Holmes, 1903  
= *J. falcata*: Chevreux & Fage, 1925  
Tjärnö, Hirtshals, Helgoland, Engl. Channel, Millport, Ireland, Mediterranean, E USA & Canada
- J. herdmani* (Walker, 1893)  
= *Podocerus herdmani* Walker, 1893  
W Norway, Denmark (Agger), Netherlands, Britain, Isle of Man, Calais, Spain, Portugal
- J. ocia* (Bate, 1862)  
= *Podocerus ocia* Bate, 1862  
non = *Podocerus dentex* Czerniavski, 1868  
\* Will be ascribed to another genus, sensu Conlan, 1990  
?Norway-W Br. Isles-W France-Mediterranean-Black Sea
- Microjassa* Stebbing, 1899
- M. cumbrensis* (Stebbing & Robertson, 1891)  
W Britain-Mediterranean
- Cerapus* Say, 1817
- C. crassicornis* (Bate, 1856)  
= *Siphonocetes c.*  
Skagerrak-SW Norway-North Sea-Faeroes-N Br. Isles
- Erichthonius* Milne-Edwards, 1830  
= *Erichthonius* Burmeister, 1837
- E. punctatus* (Bate, 1857)  
= *E. brasiliensis* Auctores, non (Dana, 1852)  
= *E. abditus*: G.O. Sars, 1894, non Templeton, 1836  
Öresund-Bohuslän-W & N Norway-Arctic Seas-Br. Isles-Mediterranean
- E. rubricornis* (Stimpson, 1853)  
= *E. hunteri* (Bate, 1862)  
\* dark gray speckled in living state  
Öresund-Bohuslän-W Norway-Arctic Seas-Iceland-Faeroes-Br. Isles-?Mediterran.-?Black Sea
- E. difformis* Milne-Edwards, 1830  
\* often associated with *Zostera marina*  
Öresund-Bohuslän-W Norway-Arctic Seas-Faeroes-Br. Isles-?Mediterran.-?Black Sea
- E. fasciatus* (Stimpson, 1853)  
\* yellowish with one single dark reddish brown band per segment  
Scandinavia, Br. Isles
- E. megalops* (G.O. Sars, 1879)  
Faeroes-Iceland-Greenland & Arctic Seas
- E. stephenseni* Myers & McGrath, 1984  
Faeroes & Greenland
- Ischyrocerus Krøyer, 1838
- I. anguipes* Krøyer, 1838  
= *I. minutus* Liljeborg, 1855  
S Baltic-Bohuslän-W & N Norway-Arctic Seas-Iceland-Faeroes-Br. Isles
- I. megalops* G.O. Sars, 1895  
N Norway-Arctic Seas-off SW Faeroes
- I. megacheir* (Boeck, 1871)  
Bohuslän-Skagerrak-W & N Norway-Arctic Seas-Iceland-off SW Faeroes
- I. assimilis* (G.O. Sars, 1885)

off S & SW Faeroes-off W Iceland-Arctic Seas

*I. brevicornis* (G.O. Sars, 1885)  
off S Iceland-off N Shetlands-Arctic Seas

*I. brusilovi* Gurjanova, 1933  
off E & NE Iceland-Arctic Seas

*I. nanoides* (Hansen, 1887)  
off SW Iceland-Arctic Seas

*I. hanseni* Stephensen, 1844  
off W Iceland

Amphithoidae Stebbing, 1899

*Sunamphithoe* Bate, 1856

*S. pelagica* (Milne-Edwards, 1830)  
Kattegatt-Bohuslän-North Sea-W Norway-Br. Isles-Sargasso Sea

*Amphithoe* Leach, 1814  
= *Pleonexes* Bate, 1856

*A. (Amphithoe) rubricata* (Montagu, 1808)  
S Baltic-Bohuslän-all of Norway-Iceland-Arctic Seas-Faeroes-Br. Isles-NW France

*A. (Amphithoe) ramondi* Audouin, 1821  
N Atlantic, e.g. Br. Isles & NW Spain

*A. (Pleonexes) gammaroides* (Bate, 1856)  
Bohuslän-North Sea-W & N Norway

*A. (Pleonexes) helleri* G. Karaman, 1975  
= *A. (Pleonexes) neglecta* Lincoln, 1976  
\* for long time confused with *P. gammaroides*.  
Brit. Isles, NW Spain

*Amphitolina* Ruffo, 1953

*A. cuniculus* (Stebbing, 1874)  
S & W British Isles & Atl. coast of France, NW Spain

Podoceridae Stebbing, 1906

*Podocerus* Leach, 1814

*P. variegatus* Leach, 1814  
Shetlands-Mediterranean

*Laetmatophilus* Bruzelius, 1859

*L. tuberculatus* Bruzelius, 1859  
= *L. armatus* (Norman, 1869)  
= *L. spinosissimus* Boeck, 1871  
\* Enequist, 1949 synonymized *L. armatus* with *L. tuberculatus*  
Bohuslän-W & N Norway-Iceland-off W Faeroes-Shetlands-Bay of Biscay-Canaries

*Xenodice* Boeck, 1871

*X. frauenfeldti* Boeck, 1871  
Kattegatt-Skagerrak-W & N Norway-Iceland-Faeroes

*Paradulichia* Boeck, 1871

*P. typica* Boeck, 1871  
Hardangerfjord-W & N Norway-Iceland-Arctic Seas

*Dyopedos* Bate, 1857

*D. monacanthus* (Metzger, 1875)  
= *Dulichia monacantha*: Auctt.  
Öresund-Bohuslän-Skagerrak-North Sea-W & N Norway-Iceland

*D. normani* (G.O. Sars, 1895)  
= *Dulichia normani* G.O. Sars, 1895  
Trondheimsfjord

*D. porrectus* Bate, 1857  
= *Dulichia porrecta*: Auctt.  
Öresund-Bohuslän-W & N Norway-Iceland-Arctic Seas-Shetlands-Br. Isles-NW France

*Dulichioipsis* Laubitz, 1977

*D. nordlandica* (Boeck, 1871)  
= *Dulichia n.*  
Skagerrak-W & N Norway-Iceland

*Dulichia* Krøyer, 1845

*D. falcata* (Bate, 1857)  
Öresund-Bohuslän-W & N Norway-Murmansk-Iceland-N Br. Isles

*D. tuberculata* Boeck, 1871  
= *D. curticauda* Boeck, 1871  
Koster Channel-Skagerrak-W & N Norway-Iceland-Arctic Seas-Br. Isles

*D. spinosissima* Krøyer, 1845  
N Norway-Iceland & Arctic Seas

*D. knipowitschi* Gurjanova, 1933  
Iceland

*D. macera* G.O. Sars, 1885  
Jan Mayen-Arctic Seas

*D. hirticornis* G.O. Sars, 1885  
off SE Faeroes-Arctic Seas

*D. spinosa* Stephensen, 1944  
off SW Faeroes

*D. abyssi* Stephensen, 1944  
off W Iceland-W Greenland

Cheluridae Allman, 1847

*Chelura* Philippi, 1839

*C. terebrans* Philippi, 1839 (TM - Mediterranean)  
Oslofjord-North Sea-W Norway-Br. Isles-Mediterranean

Aoridae Stebbing, 1899 (revised)

*Leptocheirus* Zaddach, 1844  
= *Pilocheirus* Stimpson, 1853  
= *Boeckia* Malm, 1871

*L. pilosus* Zaddach, 1844  
Kattegatt-Central Baltic & S Finland

*L. hirsutimanus* (Bate, 1862)  
= *Protomeдея hirsutimana* Bate, 1862  
= *L. pilosus*: G.O. Sars, 1894  
Kattegatt-Bohuslän-North Sea

*L. pectinatus* (Norman, 1869)  
= *Protomeдея pectinata* Norman, 1869  
= *Leptocheirus dellavalli* Stebbing, 1999  
Mediterranean-Shetlands

*L. tricristatus* (Chevreux, 1887)  
= *Pilocheirus tricristatus* Chevreux, 1887  
Atlantic coast of Europe-Engl. Channel

*L. hispinosus* Norman, 1908  
W France-Mediterranean

*Aora* Krøyer, 1845

*A. gracilis* Bate, 1857  
= *A. typica*: Auctt., non Krøyer, 1845  
Öresund-Bohuslän-Trondheim-Br. Isles-Mediterranean & Senegal

*A. spincornis* Afonso, 1976  
SW Ireland-Canary Island & Mediterranean

*Microdeutopus* Costa, 1853

*M. gryllotalpa* Costa, 1853  
S Baltic-Katteg.-Bohuslän-North Sea-SW & ?N Norway-Br. Isles-Mediterranean

*M. damnoniensis* (Bate, 1856)  
W Engl. Channel-Mediterranean

*M. anomalus* (Rathke, 1843)  
= *M. propinquus* G.O. Sars, 1894  
Öresund-Bohuslän-North Sea-Trondheim-Iceland-Shetlands-Ireland-Br. Isles-Canaries-Black Sea

*M. versiculatus* (Bate, 1856)  
Shetlands-Br. Isles-Mediterranean

*M. chelifera* (Bate, 1862)  
Belfast-Engl. Channel-Mediterranean

*M. stationis* Della Valle, 1893  
Engl. Channel-Mediterranean

*Lembos* Bate, 1857

*L. megacheira* (G.O. Sars, 1879)  
Skagerrak?-Stavanger-N Norway

*L. websteri* Bate, 1856  
?Öresund-S & W Norway-Shetlands-Br. Isles-Mediterranean

*Autonoe* Bruzelius, 1859

*A. longipes* (Liljeborg, 1852)  
= *Lembos longipes*: Auctt.  
Öresund-Bohuslän-W & N Norway-Kara Sea-Ireland-NW France

*A. denticarpus* (Myers & McGrath, 1978)  
= *Lembos denticarpus* Myers & McGrath, 1978  
Galway (Ireland)

*Uncinotarsus* l'Hardy & Truchot, 1964

*U. pellucidus* l'Hardy & Truchot, 1964  
Roscoff, interstitial at 50 m in coarse shell sand

*Unciola* Say, 1818

*U. planipes* Norman, 1867  
Belt Sea & Öresund-?Skagerrak, Helgoland-S, W & N Norway-Arctic Seas-Iceland-Br. Isles-N Bay of Biscay

*U. leucopis* (Krøyer, 1845)  
N Norway-Arctic Seas & Iceland

*U. crenatipalmata* (Bate, 1862)  
Atlantic coast of France-North Sea

*U. laticornis* Hansen, 1887  
off SW Iceland

*U. crassipes* Hansen, 1887  
off NW Faeroes-off S Iceland

*U. petalocera* (G.O. Sars, 1885)  
off W & N Norway-off N Faeroes-Arctic Seas

Corophiidae Dana, 1849 (revised)

*Corophium* Latreille, 1806

*C. volutator* (Pallas, 1766)  
Bothnian Bay-Bohuslän-North Sea-W & N Norway-Shetlands-Br. Isles-Mediterranean-Black Sea

*C. insidiosum* Crawford, 1937  
S Baltic-Bohuslän-?Norway-North Sea-Br. Isles

*C. bonnellii* (Milne-Edwards, 1830)  
Öresund-Bohuslän-W & N Norway-Arctic Seas-Iceland-Br. Isles-Mediterranean

*C. affine* Bruzelius, 1859  
Öresund-Bohuslän-W & N Norway-Faeroes-Shetlands-N Br. Isles

*C. crassicornis* Bruzelius, 1859  
S Baltic-Bohuslän-W & N Norway-Arctic Seas-Iceland-Faeroes-Br. Isles-Black Sea

*C. curvispinum* G.O. Sars, 1895  
River systems emptying into S Baltic & North Sea

- C. lacustre* Vanhöffen, 1911  
Central Baltic, S Finland, Br. Isles (in low salinity waters)
- C. arenarium* Crawford, 1937  
Mediterranean-W France-Br. Isles-North Sea
- C. acherusicum* (Costa, 1851)  
Mediterranean-Br. Isles-North Sea
- C. multisetosum* Stock, 1952  
\* in fresh and slightly brackish water  
S North Sea-Holland-Germany-Poland-Br. Isles
- C. sextonae* Crawford, 1937  
S Norway (between Risør & Mandal), North Sea-Br. Isles-Mediterranean
- C. acutum* Chevreux, 1908  
Mediterranean-Engl. Channel
- C. clarencense* Shoemaker, 1949  
N Greenland, Alaska
- Neohela* Smith, 1881
- N. monstrosa* (Boeck, 1861)  
Kattegatt-Bohuslän-W & N Norway-Faeroes-Arctic Seas & Iceland
- Siphonoecetes* Krøyer, 1845
- S. kroeyeranus* Bate, 1856  
= *S. colleti* Boeck, 1862  
Kattegatt-Bohuslän-W & N Norway
- S. pallidus* G.O. Sars, 1882  
??Skagerrak-W & N Norway
- S. striatus* Myers & McGrath, 1979  
All coasts of France-W Norway
- Biancolinidae Barnard, 1972  
\* No species within the area?
- Neomegamphopidae Myers, 1981  
\* No species within the area?
- Liljeborgioidea Stebbing, 1888
- Liljeborgiidae Stebbing, 1888
- Liljeborgia* Bate, 1862
- L. fissicornis* (M. Sars, 1858)  
Bohuslän-N Norway-Iceland & Arctic Seas-Bay of Biscay
- L. pallida* (Bate, 1857)  
= *L. brevicornis* (Bruzelius, 1859)  
Bohuslän-N Norway-Iceland & Arctic Seas-W Africa
- L. kinahani* Bate, 1862  
Bohuslän-Skagerrak-N Norway-Br. Isles-S Bretagne
- L. macronyx* (G.O. Sars, 1894)  
Bohuslän-Skagerrak-W & N Norway
- Idunella* G.O. Sars, 1895
- I. aequicornis* (G.O. Sars, 1876)  
N Norway-off N Iceland-between Iceland & Faeroes-Jan Mayen-Arctic Seas
- Listriella* Barnard, 1959
- L. picta* (Norman, 1889)  
Engl. Channel & all coasts of France
- L. mollis* Myers & McGrath, 1983  
Ireland, ?Eddystone
- L. dentipalma* Dauvin & Gentil, 1983  
W France
- L. spinifera* Dauvin & Gentil, 1983  
W France
- Sextonia* Chevreux, 1920
- S. longirostris* Chevreux, 1920  
Bretagne
- Pardaliscidae Boeck, 1871
- Pardaliscella* G.O. Sars, 1895
- P. boeckii* (Malm, 1871)  
Bohuslän-Oslofj.-Skagerrak-W Norway-W Greenland
- P. axelii* Stebbing, 1906?  
\* dubious species?  
Oslofjord-Skagerrak
- Halice* Boeck, 1871
- H. abyssi* Boeck, 1871  
Koster Channel-Skagerrak-N Norway-Arctic Seas-Iceland
- Pardalisca* Krøyer, 1842
- P. tenuipes* G.O. Sars, 1895  
Koster Channel-W Norway-Spitsbergen Area-off SW Faeroes
- P. abyssi* Boeck, 1871  
Skagerrak-W & N Norway-off E Iceland-Arctic Seas
- P. cuspidata* Krøyer, 1842  
Bohuslän-N Norway-Arctic Seas-Iceland
- Nicippe* Bruzelius, 1859
- N. tumida* Bruzelius, 1859  
Säcken area-Skagerrak-Lofoten-Shetlands-Iceland-Greenland-Scotland
- Pardaliscoides* Stebbing, 1888
- P. tenellus* Stebbing, 1888  
off SW Iceland
- Astyridae Pirlot, 1934
- Astyra* Boeck, 1871
- A. abyssi* Boeck, 1871  
Koster Channel-Skagerrak-N Norway-Norw. Sea-Arctic Seas-Iceland-off SW Faeroes
- Stilipediidae Holmes, 1908  
\* No species within the area?
- Hyperopsidae Bovallius, 1886  
\* No species within the area?
- Vitjazianidae Birstein & Vinogradov, 1955  
\* No species within the area. Abyssopelagic in NW Pacific & the Caribbean
- Sebidae Walker, 1908  
\* No species within the area?
- Salentinellidae Bousfield, 1977  
\* No species within the area?
- Paracrangonvctidae Bousfield, 1982  
\* No species within the area?
- Eusiroidea Stebbing, 1888
- Eusiridae Stebbing, 1888
- Eusirus* Krøyer, 1845
- E. propinquus* G.O. Sars, 1895  
Trondheimsfjord-N Norway-Arctic Seas
- E. cuspidatus* Krøyer, 1845  
N Norway-Arctic Seas & Iceland
- E. longipes* Boeck, 1861  
Öresund-Bohuslän-Skagerrak-W & N Norway-Br. Isles-Mediterranean
- E. leptocarpus* G.O. Sars, 1895  
Skagerrak in 200-250 m, Hardangerfjord & Trondheimsfjord
- E. minutus* G.O. Sars, 1895  
Trondheimsfjord-N Norway
- E. biscayensis* Bonnier, 1896  
off SW Faeroes-Bay of Biscay
- E. holmi* Hansen, 1886  
Jan Mayen-off NE Iceland-off N & E Faeroes-between Norway & Iceland
- E. ? abyssi* Stephensen, 1944  
off SW Iceland
- Eusirella* Chevreux, 1908
- E. elegans* Chevreux, 1908  
off SW Faeroes-Azores
- Apherusa* Walker, 1891
- A. jurinei* (Milne-Edwards, 1830)  
Kattegatt-Bohuslän-N Norway-Iceland-Arctic Seas-Faeroes-Br. Isles-Portugal
- A. tridentata* (Bruzelius, 1859)  
Lofoten-N Norway-Arctic Seas & Iceland
- A. cirrus* (Bate, 1862)  
= *Halirages borealis* Boeck, 1871  
N Norway-SE Spitsbergen-Engl. Channel-Scotland-Iceland-NW France
- A. sarsi* Shoemaker, 1930  
= *Halirages megalops* : G.O. Sars, 1882, non (Buchholz)  
N Norway-Spitsbergen-Iceland-Arctic Seas
- A. bispinosa* (Bate, 1856)  
S Baltic-Öresund-Bohuslän-N Norway-White Sea-Iceland-Faeroes-Engl. Channel-Canaries-Black Sea
- A. ovalipes* Norman & T. Scott, 1906  
Skagerrak-Bohuslän-North Sea-Engl. Channel
- A. clevei* G.O. Sars, 1904  
Kattegatt-Skagerrak-North Sea-Engl. Channel
- A. henneguyi* Chevreux & Fage, 1925  
Ireland-Engl. Channel-Bay of Biscay
- A. megalops* (Buchholz, 1874)  
= *Paramphithoe megalops* Buchholz, 1874  
E, N & W Greenland
- Cleonardo* Stebbing, 1906
- C. appendiculatus* (G.O. Sars, 1885)  
off Troms
- C. microdactylus* Stephensen, 1913  
off SW Iceland-W Greenland
- Rhachotropis* Smith, 1883
- R. aculeata* (Lepechin, 1780)  
N Norway-Arctic Seas & Iceland
- R. leucophthalma* G.O. Sars, 1895  
Skagerrak-Trondheim
- R. oculata* (Hansen, 1887)  
Iceland-Arctic Seas
- R. inflata* (G.O. Sars, 1882)  
= *R. tumida* : G.O. Sars, 1895  
Oslofjord-Skagerrak-W & N Norway & Arctic Seas
- R. helleri* (Boeck, 1871)  
Bohuslän-W & N Norway-Arctic Seas
- R. macropus* G.O. Sars, 1895

Koster Channel-W & N Norway-Arctic Seas, e.g. N Greenland

*R. grimaldi* Chevreux, 1900  
off SW Faeroes-off S Iceland-S Africa-Mediterranean

*R. rostrata* Bonnier, 1896  
off SW Faeroes-Mediterranean

*R. distincta* (Holmes, 1908)  
off S Iceland-Bay of Biscay

*R. faeroensis* Stephensen, 1944  
off SW Faeroes

Pontogeniidae Stebbing, 1906

*Pontogenia* Boeck, 1871

*P. inermis* (Krøyer, 1838)  
?SW & ?W Norway-Arctic Seas

Calliopiidae G.O. Sars, 1893

*Laothoes* Boeck, 1871

*L. meinerti* Boeck, 1871  
W-N Norway, Iceland & Faeroes

*Amphithopsis* Boeck, 1861

*A. longicaudata* Boeck, 1861  
Bohuslän-W & N Norway-Iceland-Greenland

*Halirages* Boeck, 1871

*H. fulvocinctus* (M. Sars, 1858)  
= *H. bidentatus* Stephensen, 1916  
Bohuslän-N Norway-Arctic Seas-Greenland-Iceland

*H. quadridentatus* G.O. Sars, 1876  
off NW Norway-off Spitsbergen-off Jan Mayen-off SE Faeroes

*H. elegans* (Norman, 1882)  
off N Faeroes-off W Iceland-off S Jan Mayen-off W Spitsbergen-Kara Sea

*H. mixtus* Hansen, 1931  
E Greenland

*Leptamphopus* G.O. Sars, 1895

*L. sarsi* Vanhoeffen, 1897  
= *L. longimanus* : G.O. Sars, 1895, non (Boeck, 1871)  
W Norway-off SW Iceland

*Calliopi* Lilljeborg, 1865

*C. laeviusculus* (Krøyer, 1838)  
= *C. rathkei* (Zaddach, 1844)  
Åland-Bohuslän-N Norway-Arctic Seas-Iceland-Faeroes-Br. Isles-NW France

*Cleippides* Boeck, 1871

*C. quadricuspis* Heller, 1875  
off NW Norway-off N Faeroes-off N & E Iceland-off S Jan Mayen-Arctic Seas

*C. bicuspis* Stephensen, 1931  
off S & W Iceland

*Haliragoides* G.O. Sars, 1895

*H. inermis* (G.O. Sars, 1882)  
W Skagerrak-W & N Norway-off N Faeroes

Gammarellidae Bousfield, 1977

*Gammarellus* Herbst, 1793

*G. homari* (J.C. Fabricius, 1779)  
Baltic-Bohuslän-all of Norway-Arctic Seas & Iceland-W France

*G. angulosus* (Rathke, 1843)  
Baltic-Bohuslän-all of Norway-Arctic Seas & Iceland-Mediterranean

Amathillopsidae Pirlot, 1934

*Amathillopsis* Heller, 1875

*A. spinigera* Heller, 1875  
off Lofoten-Arctic Seas-off E Iceland-between Iceland & Norway

*A. atlantica* Chevreux, 1908  
off S Iceland

Gammaracanthidae Bousfield, 1989

*Relictacanthus* Bousfield, 1989  
= *Gammaracanthus* : Auctt., non Bate, 1862

*R. relictus* G.O. Sars, 1895  
= *Gammaracanthus relictus* G.O. Sars, 1895 (TM of *Relictacanthus* Bousfield, 1989)  
= *Gammaracanthus loricatus* : Auctt., non (Sabine, 1821)  
\* *Gammaracanthus loricatus* (Sabine, 1821) is an Arctic species from e.g. N Greenland. The other species, which has been transferred to *Relictacanthus* is *Gammaracanthus lacustris* G.O. Sars, 1867  
lakes in Sweden (Vättern, Vänern, Stora Le), Norway (Mjøsa), Finland, Russia (Ladoga, Onega), Denmark?

Bateidae Stebbing, 1906

\* No species within the area. Restricted to Gulf of California & the Caribbean

Oedicerotoidea Lilljeborg, 1865

Oedicerotidae Lilljeborg, 1865

*Oedicerus* Krøyer, 1842

*O. saginatus* Krøyer, 1842  
W & N Norway-Spitsbergen-Barents Sea-Iceland-W Greenland

*O. borealis* Boeck, 1876  
?N Norway-Spitsbergen-Barents Sea-Iceland-E & W Greenland

*Periculodes* G.O. Sars, 1892

*P. longimanus* (Bate & Westwood, 1868)  
Öresund-Bohuslän-W Norway (& Finnmark)-Br. Isles-W Africa-Mediterranean

*Pontocrates* Boeck, 1871  
\* See P.G. Moore & D.J. Beare, 1993 (J. mar. biol. Ass. U. K. 73:609-615)

*P. altamarinus* (Bate & Westwood, 1862)  
Öresund-Skagerrak-W Norway-North Sea-Br. Isles-N France

*P. arcticus* G.O. Sars, 1893  
W & N Norway, Millport, Arctic Seas & Iceland

*P. arenarius* (Bate, 1858)  
= *Kroyera arenaria* Bate, 1858  
= *Pontocrates norvegicus* Boeck, 1871 : G.O. Sars, 1893  
Bohuslän-Skagerrak-North Sea-SW Norway-Br. Isles-W Mediterranean

*Synchelidium* G.O. Sars, 1892

*S. haplocheles* (Grube, 1864)  
= *S. brevicarpum* (Bate & Westwood, 1868)  
Kattegatt-Bohuslän-W Norway-Trondheim

*S. tenuimanum* Norman, 1895  
= *S. haplocheles* : G.O. Sars, 1892, non (Grube, 1864)  
Bohuslän-North Sea-W Norway-Trondheim

*S. intermedium* G.O. Sars, 1892  
Koster Channel-Skagerrak-W Norway-Trondheim

*S. maculatum* Stebbing, 1906  
Engl. Channel-Mediterranean

*Paroediceros* G.O. Sars, 1892

*P. curvirostris* (Hansen, 1887)  
? S Iceland

*P. lyncaeus* (M. Sars, 1858)  
off S Lofoten-Arctic Seas-Iceland

*P. propinquus* (Goës, 1866)  
N Norway-Arctic Seas-Iceland-?off N Faeroes

*P. macrocheir* (G.O. Sars, 1885)  
off NE Iceland

*P. intermedius* ?Stebbing, 1906  
= *Oedicerus microps* Hansen, 1887  
?N Greenland

*Halicieion* Boeck, 1871

*H. aequicornis* (Norman, 1869)  
= *H. longicaudatum* Boeck, 1871  
?Skagerrak-Bergen-Trondheim-Shetland, Mediterranean

*Arrhis* Stebbing, 1906

*A. phyllonyx* (M. Sars, 1858)  
Kattegatt-Bohuslän-Skagerrak-W & N Norway-Orkneys-Iceland & Arctic Seas, e.g. N Greenland

*Westwoodilla* Bate, 1862

*W. caecula* (Bate, 1856)  
= *W. hyalina* Bate, 1862  
= *W. acutifrons* G.O. Sars, 1895  
Öresund-Bohuslän-Narvik-Br. Isles & N France

*W. megalops* (G.O. Sars, 1882)  
\* another synonyme of *W. caecula* (Bate), according to Dunbar, 1954  
N Norway-Arctic Seas & Iceland

*W. brevicarcar* (Goës, 1866)  
N Norway-Iceland-Arctic Seas, e.g. E & N Greenland

*W. rectirostris* (Della Valle, 1893)  
W France

*Oediceropsis* Lilljeborg, 1865

*O. brevicornis* Lilljeborg, 1865  
Koster Channel-W & N Norway

*Acanthostepheia* Boeck, 1871

*A. malmgreni* (Goës, 1866)  
Iceland-Spitsbergen-Arctic Seas

*Aceroides* G.O. Sars, 1895

*A. latipes* (G.O. Sars, 1882)  
= *Aceros distingendus* Hansen, 1887  
N Norway-Spitsbergen-Arctic Seas, e.g. E, N & W Greenland

*Bathymedon* G.O. Sars, 1892

*B. obtusifrons* (Hansen, 1887)  
Skagerrak-N Norway-Arctic Seas & Iceland

*B. saussurei* (Boeck, 1871)  
Bohuslän (Koster Channel)-N Norway-W Greenland

*B. longimanus* (Boeck, 1871)  
Kattegatt-Bohuslän-W & N Norway

*Monoculopsis* G.O. Sars, 1892

*M. longicornis* (Boeck, 1871)  
?Skagerrak, ?W & N Norway-Jan Mayen-Iceland-Arctic Seas

*Monoculodes* Stimpson, 1853

*M. carinatus* (Bate, 1856)  
Kattegatt-Bohuslän-off W Norway-N Norway-Mediterranean

*M. borealis* Boeck, 1871  
Trondheim-N Norway-Iceland-Arctic Seas-Scotland

*M. tuberculatus* Boeck, 1871  
N Norway-Clyde Sea-Iceland-Arctic Seas

- M. gibbosus* Chevreux, 1888  
Clyde Sea-Bay of Biscay
- M. subnudus* Norman, 1889  
= *M. falcatus* G.O. Sars, 1892  
Bohuslän-Skagerrak-W & N Norway-Iceland-Shetlands-N Bretagne
- M. packardii* Boeck, 1871  
= *M. tenuirostratus* Boeck, 1871  
Öresund-Bohuslän-W & N Norway-Iceland & Arctic Seas-Jan Mayen-Scotland
- M. kroeyeri* Boeck, 1871  
?Skagerrak, SW & N Norway-W Greenland
- M. longirostris* (Goës, 1866)  
?Kattegatt, N Norway-off N Faeroes-Jan Mayen-Iceland & Arctic Seas
- M. pallidus* G.O. Sars, 1895  
W Norway-Iceland-off SW Faeroes-Arctic Seas
- M. schneideri* G.O. Sars, 1895  
N Norway-Arctic Seas, e.g. N & W Greenland
- M. latimanus* (Goës, 1866)  
Bergen-N Norway-Iceland & Arctic Seas
- M. tessellatus* Schneider, 1884  
N Norway-Iceland-Faeroes-Arctic Seas
- M. norvegicus* (Boeck, 1861)  
Kattegatt-Bohuslän-Skagerrak-S, W & N Norway-Iceland & Arctic Seas
- M. rostratus* Stephensen, 1931  
off S Iceland
- Oedicerina* Stephensen, 1931
- O. ingolfi* Stephensen, 1931  
between Faeroes & Jan Mayen
- Exoedicerotidae Barnard & Karaman, 1983  
\* No species within the area?
- Paracullispiidae Barnard & Karaman, 1983  
\* No species within the area?
- Dexaminoidea Leach, 1814
- Lepechinellidae Schellenberg, 1926
- Lepechinella* Stebbing, 1908
- L. chrysotheras* Stebbing, 1908  
off SW Faeroes-off S Shetlands
- Dexaminiidae Leach, 1814  
= Atylidae Lilljeborg, 1865
- Atylus* Leach, 1815  
= *Nototropis* Costa, 1853 & *Paratylus* G.O. Sars, 1893
- A. falcatus* Metzger, 1871  
?Bohuslän-Skagerrak-North Sea-N Norway-Br. Isles-NW France
- A. swammerdami* (Milne-Edwards, 1830)  
S Baltic-Bohuslän-N Norway-White Sea-Faeroes-Senegal-Azores-Mediterranean
- A. smitti* (Goës, 1866)  
N Norway-Iceland-Jan Mayen-Spitsbergen-Arctic Seas
- A. guttatus* (Costa, 1851)  
S North Sea-Irish Sea- Mediterranean
- A. vedlomensis* (Bate & Westwood, 1862)  
Kattegatt-Bohuslän-Trondheim-Shetlands-Br. Isles
- A. nordlandicus* Boeck, 1871  
?Skagerrak-W Norway-Iceland & Arctic Seas
- A. carinatus* (J.C. Fabricius, 1793)  
N Norway-Spitsbergen-Arctic Seas, e.g. N Greenland
- Dexamine* Leach, 1814
- D. spinosa* (Montagu, 1813)  
Öresund-Bohuslän-W & N Norway-Br. Isles-Mediterranean-Senegal-Canaries-Azores
- D. thea* Boeck, 1861  
Kattegatt-Bohuslän-W & N Norway-Iceland-Br. Isles.-W France, W Mediterranean
- Tritaeta* Boeck, 1876
- T. gibbosa* (Bate, 1862)  
Bohuslän (e.g. Segelskären)-Skagerrak-W Norway-North Sea-Br. Isles-Mediterranean-Black Sea-Senegal
- Guernea* Chevreux, 1887
- G. coalita* (Norman, 1868)  
North Sea-W Mediterranean
- Anatylidae Bulycheva, 1955  
\* No species within the area?
- Prophiliantidae Nicholls, 1940  
\* No species within the area?
- Melphidippoidae Stebbing, 1899
- Melphidippidae Stebbing, 1899
- Melphidippella* G.O. Sars, 1894
- M. macra* (Norman, 1869)  
Öresund-Bohuslän-S & N Norway-North Sea-Br. Isles-Senegal
- Melphidippa* Boeck, 1871
- M. goesii* Stebbing, 1899  
= *Gammarus spinosus* Goës, 1866 (homonym)  
Trondheim-N Norway-Arctic Seas-Iceland-Scotland
- M. macrura* G.O. Sars, 18
- W-N Norway
- M. borealis* Boeck, 1871  
Koster Channel-Skagerrak-Oslofjord-N Norway
- Megaluroidea J.L. Barnard, 198?
- Megaluropus* Hoek, 1889
- M. agilis* Hoek, 1889  
Kattegatt-Bohuslän-Skagerrak, Engl. Channel
- "*Hornellia-Cheirocratus* family group"  
\* Should also *Cheirocratella* & other genera be moved to this undescribed (1986) family?
- Cheirocratus* Norman, 1867
- C. sundevallii* (Rathke, 1843)  
S Baltic-Bohuslän-W & N Norway-Bosporus
- C. intermedius* G.O. Sars, 1895  
Katteg.-Bohuslän-W & N Norway-Br. Isles-NW France
- C. assimilis* (Lilljeborg, 1852)  
Katteg.-Bohuslän-S, W & N Norway-Mediterranean
- C. robustus* G.O. Sars, 1895  
Bohuslän-Skagerr.-Trondheimsfjord
- Phreatogammaridae Bousfield, 1982  
\* No species within the area. Confined to fresh & brackish waters on and around New Zealand
- Gammaroidea Leach, 1814  
\* Several (9) other families, chiefly from fresh water, belong here
- Gammaridae Leach, 1814
- Gammarus* Fabricius, 1775 (TM *G. pulex* Linnaeus)  
= *Rivulogammarus* Karaman, 1931 (TM *G. pulex* Linnaeus)
- G. duebenii* Lilljeborg, 1852  
= *Rivulogammarus duebenii* (Lilljeborg, 1852)  
Bothnian Bay-Bohuslän-all coastal Norway-Iceland-Br. Isles-France
- G. lacustris* G.O. Sars, 1863  
= *Rivulogammarus lacustris* (G.O. Sars, 1863)  
\*limnic species usually found in lakes  
Sweden, Denmark, Norway, Iceland, Finland
- G. finmarchicus* Dahl, 1938  
all of Norway & Iceland
- G. locusta* (Linnaeus, 1758)  
Åland-Bohuslän-N Norway-?Iceland
- G. oceanicus* Segerstråle, 1947  
Bothnian Sea-Bohuslän-all coast of Norway-Arctic Seas & Iceland
- G. pulex* (Linnaeus, 1758)  
= *Rivulogammarus pulex* (Linnaeus, 1758)  
\*limnic species usually found in running waters  
Sweden, Denmark, Finland, Britain
- G. zaddachi* Sexton, 1912  
Bothnian Sea-North Sea-W & N Norway
- G. inaequicauda* Stock, 1966  
= *G. campylops* : G.O. Sars, 1895, non Leach, 1814  
\* May possibly be synonym. with *G. insensibilis* Stock, 1966  
S Baltic, Oslofjord, Grimstad (S Norway)
- G. insensibilis* Stock, 1966  
Engl. Channel-Mediterranean
- G. salinus* Spooner, 1947  
Bothnian Bay-Oslofjord-W Norway-North Sea
- G. crinicornis* Stock, 1966  
Holland-Mediterranean
- G. tigrinus* Sexton, 1939  
\* American brackish-fresh-water species introduced in Europe  
Baltic, Germany, Holland, British Isles, Ireland
- G. chevreuxi* Sexton, 1913  
N Wales-N Africa
- G. setosus* Dementieva, 1931  
N Norway-Spitsbergen & Arctic Seas, e.g. N Greenland
- G. wilkitzkii* (Birula, 1897)  
Spitsbergen-Arctic Seas-Iceland
- Echinogammarus* Stebbing, 1899  
= *Chaetogammarus* Martynov, 1925  
= *Marinogammarus* Schellenberg, 1937
- C. marinus* (Leach, 1815)  
Öresund-Bohuslän-all of Norway-Iceland
- C. stoerensis* (Reid, 1938)  
Öresund-probably all Norway-Murmansk-Iceland
- C. pirloti* (Sexton & Spooner, 1940)  
Oslofjord-W Norway
- Eulimnogammarus* Bazikalova, 1945
- E. obtusatus* (Dahl, 1938)  
probably all Norwegian coast-Iceland
- Pectenogammarus* Reid, 1940
- P. planicrurus* Reid, 1940  
Mediterranean-Brit. Isles
- Weyprechtia* Stuxberg, 1880
- W. pinguis* (Krøyer, 18??)  
Spitsbergen, Murmansk, White Sea, Novaja Zemlja, Franz Joseph Land, N. Greenland
- W. heuglini* (Buchholz, 18??)

= *W. mirabilis* Stuxberg, 1882  
Spitsbergen, Novaja Zemlja, Kara Sea

Hadzioidae Karaman, 1932

Melitidae Bousfield, 1973

*Eriopisa* Stebbing, 1890

*E. elongata* (Bruzellius, 1859)  
Katteg.-Bohuslän-Skagerr.-W & N Norway-Iceland-Faeroes-Scotland-Mediterranean

*Eriopisella* Chevreux, 1920

*E. pusilla* Chevreux, 1920 (TM)  
N France

*Cheirocratella* Stephensen, 1940

*C. thori* Stephensen, 1940  
Iceland

*Gammarella* Bate, 1857

*G. fucicola* (Leach, 1814)  
Shetlands-Mediterranean

*Melita* Leach, 1814

*M. palmata* (Montagu, 1804)  
C. Baltic-Bohuslän-North Sea

*M. formosa* Murdoch, 1885  
N Norway-Iceland-Arctic Seas

*M. dentata* (Krøyer, 1842)  
Öresund-Kattegatt-Bohuslän-W & N Norway-Iceland-Arctic Seas-Br. Isles-Cameroun

*M. hergensis* Reid, 1939  
\* often mistaken for *M. palmata*  
British Isles

*M. richardi* Chevreux, 1900  
off S Iceland (deep water)

*Abludomelita* Karaman, 1981

*A. obtusata* (Montagu, 1813)  
= *Melita obtusata* (Montagu, 1813)  
@ apparently commensalistic with *Anemonia sulcata*  
Öresund-Bohuslän-North Sea

*A. gladiosa* (Bate, 1862)  
= *Melita gladiosa* Bate, 1862  
Engl. Channel-Mediterranean

*Allomelita* Stock, 1984

*A. pellucida* (G.O. Sars, 1882)  
= *Melita pellucida* G.O. Sars, 1882  
?Skagerrak-N Ice Sea

*Maera* Leach, 1814

*M. loveni* (Bruzellius, 1859)  
Öresund-Bohuslän-W & N Norway-Arctic Seas-Iceland-Scotland

*M. othonis* (Milne-Edwards, 1830)  
Kattegatt-Bohuslän-North Sea

*M. grossimana* (Montagu, 1808)  
Bristol Channel-Mediterranean

*M. inaequipes* (A. Costa, 1851)  
?W France, Azores, Canaries, Mediterranean

*Maerella* Chevreux, 1911

*M. tenuimana* (Bate, 1862)  
North Sea-Mediterranean

*Elasmopus* Costa, 1853

*E. rapax* Costa, 1853  
Skagerrak

*Ceradocus* Costa, 1853

*C. torelli* (Goës, 1866)  
Iceland-Arctic Seas

*C. semiserratus* (Bate, 1862)  
Brit. Isles-Mediterranean

*Pallasea* Bate, 1862

*P. quadrispinosa* G.O. Sars, 1867  
NE Bothnian Bay & lakes in Sweden, Norway, Denmark & Finland & White Sea

Hadziidae Karaman, 1932  
\* No species within the area. Fresh & brackish waters in warm-temperate & tropical regions.

Carangoliopsidae Bousfield, 1977  
\* No species within the area. Only species known from Mediterranean hadal zone

Bogidielloidea Hertzog, 1936

Bogidiellidae Hertzog, 1936  
\* No species in the area. Chiefly fresh water, but some species are coastal marine with a distribution similar to Ingolfiellidae

Artesiidae Holsinger, 1980  
\* No species in the area. Confined to artesian fresh water wells

Crangonyctoidea Bousfield, 1977  
\* Strictly fresh water living, i.a. Niphargidae S. Karaman, 1962

THERMOSBAENACEA Monod, 1927  
\* no marine species

HEMICARIDEA Schram, 1981

SPELAEOGRIPHACEA Gordon, 1957  
\* no marine species

CUMACEA Krøyer, 1846

Leuconidae G.O. Sars, 1878

*Leucon* Krøyer, 1846

*L. siphonatus* Calman, 1905  
Iceland-off W Ireland-Bay of Biscay-Mediterranean

*L. nasicoides* Liljeborg, 1855  
Öresund-Bohuslän-W Norway-Iceland-Arctic Seas, e.g. N Greenland

*L. nasica* (Krøyer, 1841)  
Öresund-Bohuslän-North Sea-N Norway-Br. Isles-Arctic Seas & Iceland

*L. nathorstii* Ohlin, 1901  
Iceland-Spitsbergen-Arctic Seas

*L. acutirostris* G.O. Sars, 1865  
Öresund-Bohuslän-N Norway-Newfoundland

*L. fulvus* G.O. Sars, 1865  
Lofoten-Arctic Seas & Iceland

*L. pallidus* G.O. Sars, 1865  
Koster Channel-Oslofjord-Skagerrak-N Norway-Arctic Seas, e.g. N Greenland & Iceland

*L. tener* Hansen, 1920  
Skagerrak-off SW Iceland-Davis Strait

*L. spiniventris* Hansen, 1920  
off S Iceland

*L. spinulosus* Hansen, 1920  
off S Jan Mayen-Davis Strait, Kara Sea, N Greenland

*Eudorellopsis* G.O. Sars, 1883

*E. deformis* (Krøyer, 1846)  
Belt Sea-Skagerrak-Helgoland-W Norway-N & W Br. Isles-Arctic Seas & Iceland

*Eudorella* Norman, 1867

*E. emarginata* (Krøyer, 1846)  
Öresund-Bohuslän-N Norway-Arctic Seas & Iceland

*E. hirsuta* (G.O. Sars, 1869)  
Koster Channel-Skagerrak-Lofoten-Arctic Seas-between Rockall & Ireland

*E. truncatula* (Bate, 1856)  
Öresund-Bohuslän-N Norway-Arctic Seas-Br. Isles-Mediterranean

Bodotriidae T. Scott, 1901

Vauntomponiinae Hale, 1944

*Vauntomponia* Bate, 1858

*V. cristata* Bate, 1858  
N North Sea-Shetlands-Faeroes-W Br. Isles-Bay of Biscay-Morocco-Mediterran.

*Cumopsis* G.O. Sars, 1878

*C. goodsiri* (van Beneden, 1861)  
Kattegatt, North Sea-Br. Isles & Ireland-Bay of Biscay

*C. longipes* (Dohrn, 1869)  
Engl. Channel-Bay of Biscay-Mediterranean

*C. fagei* Bacescu, 1956  
Ireland-Engl. Channel, Morocco

*Bathycuma* Hansen, 1895  
\* Correct subfamily?

*B. brevirostris* (Norman, 1879)  
off S Iceland-off Ireland-Bay of Biscay

Bodotriinae Hale, 1944

*Iphinoe* Bate, 1856, non Rafinesque-Schmaltz, 1815, nee Adams, 1854  
= *Cyrianassa* Bate, 1856  
= *Iphithoe* Norman, 1867

*I. trispinosa* (Goodsir, 1843)  
Kattegatt-Bohuslän-North Sea-Br. Isles-Bay of Biscay-?Norw. Sea-Morocco & Canaries-Mediterranean

*I. tenella* G.O. Sars, 1879  
Engl. Channel-Bay of Biscay-Mediterranean

*I. serrata* Norman, 1867  
Shetlands-W Br. Isles-Mediterranean

*Bodotria* Goodsir, 1843

*B. pulchella* (G.O. Sars, 1879)  
Kattegatt-North Sea-W Br. Isles-Bay of Biscay-Morocco-Mediterranean

*B. scorpioides* (Montagu, 1804)  
= *Cuma edwardsii* G.O. Sars, 1899  
Öresund-Bohuslän-North Sea-W Norway-Engl. Channel & Ireland-Bay of Biscay -Mediterranean

*B. arenosa arenosa* (Goodsir, 1842)  
= *Cuma scorpioides* G.O. Sars, 1899, non Montagu, 1804  
Skagerrak-North Sea-Shetlands-Faeroes-W Br. Isles

*B. armoricana* Le Loeuff & Intes, 1977  
Concarneau

*Eocuma* Marcusen, 1894

*E. ferox* (P. Fischer, 1872)  
Bay of Biscay-Mediterranean

*E. dollfusi* Calman, 1907  
Engl. Channel-Bay of Biscay

*Cyclaspis* G.O. Sars, 1865

- C. longicaudata* G.O. Sars, 1865  
W Skagerrak-North Sea-Lofoten-Arctic Seas-Iceland-off SW Faeroes-Bay of Biscay-Azores-Mediterranean
- Nannastacidae Bate, 1866
- Nannastacus* Bate, 1865
- N. unguiculatus* (Bate, 1859)  
Shetlands-Bay of Biscay-Mediterranean
- N. brevicaudatus* Calman, 1905  
W Ireland-Morocco
- N. longirostris* G.O. Sars, 1879  
Concarneau
- Campylaspis* G.O. Sars, 1865
- C. rubicunda* (Liljeborg, 1855)  
Öresund-Bohuslän-Skagerrak-North Sea-W Br. Isles-N Norway-Arctic Seas, e.g. N Greenland & Iceland
- C. glabra* G.O. Sars, 1879  
Skagerrak-N North Sea-W Norway-Iceland-W Br. Isles-Bay of Biscay-Mediterranean
- C. costata* G.O. Sars, 1865  
Kattegatt-Bohuslän-Skagerrak-North Sea-N Norway-Shetlands-Faeroes-W Br. Isles
- C. undata* G.O. Sars, 1865  
off Lofoten-between Faeroes & Shetlands
- C. sulcata* G.O. Sars, 1870  
Skagerrak-N North Sea-Lofoten-Faeroes-Shetlands-W Br. Isles-Mediterranean
- C. horrida* G.O. Sars, 1870  
W Skagerrak-N Norway-Arctic Seas-Iceland-Faeroes
- C. affinis* G.O. Sars, 1870  
off Lofoten-Firth of Forth
- C. verrucosa* G.O. Sars, 1866  
Skagerrak-Lofoten-Iceland-Faeroes-Ireland-Bay of Biscay-Mediterranean
- C. macrophthalma* G.O. Sars, 1879  
North Sea, Engl. Channel
- C. globosa* Hansen, 1920  
Lofoten-Davis Strait
- C. rostrata* Calman, 1905  
off SW Faeroes-off SW Ireland-off W Africa-Mediterranean
- C. serratipes* Hansen, 1920  
off W & SW Iceland
- C. laticarpa* Hansen, 1920  
off SW Faeroes
- C. alba* Hansen, 1920  
off SW Faeroes
- C. intermedia* Hansen, 1920  
off S Jan Mayen-Davis Strait
- C. legendrei* Fage, 1951  
W Br. Isles-Bay of Biscay
- C. stephensi* Just, 1970  
N Greenland
- Procampylaspis* Bonnier, 1896
- P. armata* Bonnier, 1896  
= *P. echinata* Bonnier, 1896  
Shetlands-SW Ireland-Bay of Biscay-Azores-Mediterranean
- P. macronyx* Hansen, 1920  
off W Iceland-off SW Faeroes
- P. bituberculata* Hansen, 1920  
off SW Faeroes
- Cumella* G.O. Sars, 1865
- C. pygmaea* G.O. Sars, 1865  
Oslofjord-North Sea-Lofoten-Engl. Channel-W Br. Isles-Bay of Biscay-Shetlands-Faeroes
- C. tarda* Hansen, 1920  
between Shetlands & Faeroes
- Cumellopsis* Calman, 1905
- C. helgae* Calman, 1905  
Iceland-off SW Faeroes
- Pseudocumatidae G.O. Sars, 1878
- Petalosarsia* Stebbing, 1893
- P. declivis* (G.O. Sars, 1865)  
Skagerrak-North Sea-W Br. Isles-N Norway-Arctic Seas & Iceland
- Pseudocuma* G.O. Sars, 1865
- P. similis* G.O. Sars, 1900  
Kattegatt-Bohuslän-North Sea-W Norway-Engl. Channel & Ireland-Bay of Biscay
- P. longicornis* (Bate, 1858)  
= *P. cercaria* G.O. Sars, 1879  
Öresund-Bohuslän-North Sea-N Norway-Engl. Channel & W Br. Isles-Bay of Biscay
- P. gilsoni* Bacescu, 1950  
\* Possibly a neotenus form of *P. longicornis* (Bate)  
W Br. Isles-Belgium
- Lampropiidae G.O. Sars, 1878
- Platysympus* Stebbing, 1912  
= *Platyspis* G.O. Sars, 1870, non Agassiz, 1846
- P. typicus* (G.O. Sars, 1870)  
Skagerrak-W Norway, off W Ireland, Mediterranean
- P. tricarinatus* Hansen, 1920  
Skagerrak, Lofoten, S Iceland, Spitsbergen
- Platytyphlops* Stebbing, 1913
- P. orbicularis* (Calman, 1905)  
off SW Faeroes-off W Ireland
- Lamprops* G.O. Sars, 1863
- L. fasciata* G.O. Sars, 1863  
Öresund-Katteg., Oslofjord-N Norway & Arctic Seas, North Sea-W Br. Isles-Engl. Channel
- L. fuscata* G.O. Sars, 1865  
Lofoten-Arctic Seas
- Hemilamprops* G.O. Sars, 1883
- H. rosea* (Norman, 1863)  
Danish Belt Sea-Bohuslän-N North Sea-N Norway-Faeroes-Shetlands-W Br. Isles
- H. cristata* (G.O. Sars, 1870)  
Skagerrak-NW Norway-Iceland-Faeroes-Br. Isles
- H. assimilis* G.O. Sars, 1883  
off N Norway, SW Faeroes-Irish Sea
- H. uniplicata* (G.O. Sars, 1872)  
Hardangerfjord-Lofoten, Norw. Sea, SW Faeroes, off W Ireland
- Diastylidae Bate, 1856
- Diastylis* Say, 1818
- D. echinata* Bate, 1865  
between Gotland & Landsort-Koster Channel-North Sea-N Norway-Arctic Seas & Iceland
- D. rugosa* G.O. Sars, 1865  
\* often among algae!  
Kattegatt-Bohuslän-Trondheimfjord-North Sea-Engl. Channel & W Br. Isles-Bay of Biscay-Mediterranean
- D. goodsiri* (Bell, 1855)  
Skagerrak??. N Norway-Arctic Seas, e.g. E & N Greenland & Faeroe Iceland area
- D. cornuta* (Boeck, 1864)  
Kattegatt-Bohuslän-Skagerrak-N North Sea-Lofoten-Br. Isles-Engl. Channel-Ireland-Bay of Biscay-Mediterranean
- D. boeckii* Zimmer, 1930  
S Baltic-Bohuslän-N North Sea
- D. tumida* (Liljeborg, 1855)  
Öresund-Bohuslän-N North Sea-Lofoten-W Br. Isles
- D. lucifera* (Krøyer, 1841)  
Öresund-Bohuslän-North Sea-Engl. Channel-W Br. Isles-Shetlands-Faeroes-N Norway-Arctic Seas & ?Iceland
- D. glabra* Zimmer, 1900  
Faeroe-Iceland-area-Barents Sea-Arctic Seas
- D. rathkei* (Krøyer, 1841) *typica* Zimmer, 1926  
off NE Gotska Sandön-Bohuslän-North Sea-W Br. Isles-N Norway-Arctic Seas & Iceland
- D. rathkei* (Krøyer, 1841) *sarsi* Norman, 1905  
N Norway-Spitsbergen-Arctic Seas
- D. oxyrhyncha* Zimmer, 1926  
N Norway-Shetland-Faeroe-area-Norw. Sea-Arctic Seas & Iceland
- D. laevis* Norman, 1869  
= *D. rostrata* : G.O. Sars, 1900, non (Goodsir, 1843)  
Öresund-Bohuslän-SW Norway-North Sea-Engl. Channel-W Br. Isles-Ivory Coast
- D. bradyi* Norman, 1879  
Isefjord, Skagerrak-S North Sea-Engl. Channel-W Br. Isles-Bay of Biscay, ?Norw. Sea
- D. scorpioides* (Lepechin, 1780)  
Kara Sea-Jan Mayen-Arctic Seas, e.g. N Greenland & Faeroe Iceland area
- D. spinulosa* Heller, 1875  
N Norway-Arctic Seas, e.g. E, N & W Greenland & Kara Sea
- D. lepechini* Zimmer, 1926  
= *D. scorpioides* : G.O. Sars, 1900, non (Lepechin 1780)  
N Norway-Arctic Seas & Iceland
- D. edwardsi* (Krøyer, 1841)  
Arctic Seas, e.g. N Greenland & ?Iceland
- D. sulcata* Calman, 1912  
W Br. Isles-Norw. Sea-Arctic Seas
- D. polaris* G.O. Sars, 1872  
= *D. stygia* G.O. Sars, 1872  
off Faeroes-off Iceland-Jan Mayen-Arctic Seas
- Leptostylis* G.O. Sars, 1869
- L. macrura* G.O. Sars, 1870  
W-N Norway & Norw. Sea-Spitsbergen-Ireland-Mediterranean
- L. longimana* (G.O. Sars, 1865)  
Kattegatt-Bohuslän-N North Sea-Lofoten-W Br. Isles & Faeroe-Iceland area
- L. ampullacea* (Liljeborg, 1855)  
Öresund & Belt Sea-Oslofjord-Skagerrak-N North Sea-N Norway-Faeroe-Shetland area-Iceland
- L. villosa* G.O. Sars, 1869  
Öresund-Bohuslän-N North Sea-N Norway-W Br. Isles-Faeroe-Shetland area-Iceland & Arctic Seas
- Diastylodes* G.O. Sars, 1900
- D. serrata* (G.O. Sars, 1865)  
Kattegatt-Bohuslän-N North Sea-N Norway-Ireland-Mediterranean
- D. biplicata* (G.O. Sars, 1865)  
Kattegatt-Bohuslän-N North Sea-N Norway-Faeroe-Shetland area-W Br. Isles-Engl. Channel
- D. scabra* Hansen, 1920  
off SW Faeroes

**Brachydiastylis** Stebbing,1912

**B. resima** (Krøyer,1846)  
Kattegatt-Skagerrak-N North Sea-N Norway-Arctic Seas & Iceland

**B. nimia** Hansen,1920  
N Greenland, arctic Canada

**Makrocyliandrus** Stebbing,1912  
= *Makrocyliandrus* : Auuct.

**M. longipes** (G.O. Sars,1871)  
= *M. spiniventris* Hansen,1920  
Shetland Faeroe Channel-Bay of Biscay-Mediterranean

**M. insignis** (G.O. Sars,1871)  
W Scotland & Ireland-Portugal-Mediterranean

**M. josephinae** (G.O. Sars,1871)  
off SW Faeroes-Bay of Biscay-Portugal

**M. longicaudatus** (Bonnier,1896)  
off SW Faeroes-off Arcachon-Azores

**M. tubulicaudata** (Calman,1905)  
off N Scotland-off W Ireland-Bay of Biscay

**M. sp.** Forsman,1938  
Skagerrak

Gynodiastylidae Stebbing,1912  
\* No species within the area?

**Ceratocumatidae** Calman,1905  
\* No species within the area?

**TANAIDACEA** Dana,1853

APSEUDOMORPHA Sieg,1980  
\* Only one of 10 families represented in Scandinavian waters (see e.g. Schram,1986)

**Apseuideoidea** Leach,1814 (Schram, Sieg & Malzahn,1986)

**Apseuididae** Leach,1814

**Apseudes** Leach,1814

**A. spinosus** (M. Sars,1858)  
Öresund-Bohuslän-N Norway-Arctic Seas-Iceland-Faeroes

**A. latreillii** (Milne-Edwards,1828)  
S & NE England, E Scotland, Engl. Channel, France, Italy

**A. talpa** (Montagu,1808)  
Holland, France, S & W Br. Isles, Ireland, N Africa

**A. grossimanus** Norman, in Norman & Stebbing,1886  
off SW Ireland, Bay of Biscay, off Portugal

**A. vicinus** Hansen,1913  
off S Iceland

**A. gracillimus** Hansen,1913  
off S Iceland

**Calozodion** ?,19??

**C. dollfusi** Gutu,1989  
Channel Islands

**Sphyrapidae** Gutu,1980

**Sphyrapus** G.O. Sars,1882 ex Norman MS

**S. anomalus** (G.O. Sars,1869)  
Oslofjord-Skagerrak-N Norway-Arctic Seas-Iceland

**S. serratus** G.O. Sars,1882  
between Norway & Iceland-off N Iceland-Jan Mayen-Arctic Seas

**S. tudes** Norman & Stebbing,1886  
off SW Faeroes-off S Rockall

**S. malleolus** Norman & Stebbing,1886  
S of Rockall, Bay of Biscay, off Portugal & Greenland

NEOTANAIDOMORPHA Sieg,1980

**Neotanaidae** Lang,1956

**Neotanais** Beppard,1886  
= *Alaotanais* Norman & Stebbing,1886

**N. serratispinosus** (Norman & Stebbing,1886)  
off NW & W Ireland-off S Greenland, very deep water

TANAIDOMORPHA Sieg,1980

**Tanaoidea** Dana,1849

**Tanaidae** Dana,1849

**Tanais** Latreille,1831  
= *Tanais* Audouin & M.-Edw.,1829 (n.nud.) & M.-Edw,1830 (n.nud.)

**T. dulongii** (Audouin,1826)  
= *T. cavolinii* H. Milne-Edwards, in Audouin & H. Milne-Edwards,1840  
= *T. tomentosus* Krøyer,1842  
Öresund, S-N Norway, Faeroes-Br. Isles & Mediterranean

**Parasinelobus** Sieg,1980

**P. chevreuxi** (Dollfus,1898)  
W Ireland, Wales, Devon, atl. coast of France, Spain & Morocco

**Zeuxo** Kossmann,1873, non Templeton,1840 (Amphipoda), nec Gistel,1847 (Echinodermata)  
= *Zeuxokoma* Kossmann,1???

**Z. holdichii** Bamber,1990  
Biscay

**Paratanaoidea** Lang,1949

**Paratanaidae** Lang,1949

**Heterotanais** G.O. Sars,1882

**H. oerstedii** (Krøyer,1842)  
= *Tanais curculio* Krøyer,1842  
= *Tanais balticus* F. Müller,1852  
C. Baltic-Bohuslän, Norway, Wales, S & E England, N France

**Leptocheilia** Dana,1849

**L. dubia** (Krøyer,1842)  
= *Leptocheilia savignyi* (Krøyer,1842)  
Holland, SW England, S & W Ireland, Brittany-Senegal

**Paratanais** ?,1???

**P. hessleri** Kudinova-Pasternak,1985  
NE Atlantic

**Typhlotanaidae** Sieg,1984

**Typhlotanais** G.O. Sars,1882

**T. aequiremis** (Lilljeborg,1864)  
Bohuslän- Skagerrak-Tromsø, Iceland, N North Sea

**T. cornutus** (G.O. Sars,1879)  
N Norway-between Finnmark & Bear Island-Davis Strait

**T. finmarchicus** G.O. Sars,1882  
N Norway-Iceland-Jan Mayen-White Sea-Arctic Seas, e.g. E, N & W Greenland

**T. tenuimanu** (Lilljeborg,1864)  
Koster area-Skagerrak-Lofoten

**T. trispinosus** Hansen,1913  
Skagerrak-Davis Strait

**T. assimilis** G.O. Sars,1882  
Skagerrak-N Norway

**T. tenuicornis** G.O. Sars,1882  
Gullmarn, Koster area, Oslofjord, Skagerrak-Norw. Sea, E Scotland

**T. penicillatus** G.O. Sars,1882  
W Norway-off SW Iceland

**T. brevicornis** (Lilljeborg,1864)  
Koster Channel, Skagerrak-Norw. Sea, Scotland

**T. microcheles** G.O. Sars,1882  
Norw. Sea-Arctic Seas, W Scotland & Ireland, Isle of Man

**T. pulcher** Hansen,1913  
W Scotland & Davis Straits

**T. irregularis** Hansen,1913  
off Faeroes-off Iceland-off Jan Mayen-Kara Sea, N Greenland

**T. gracilipes** Hansen,1913  
off S Iceland

**T. eximius** Hansen,1913  
off SW Iceland

**T. mucronatus** Hansen,1913  
off NE Iceland-off S Jan Mayen

**T. inermis** Hansen,1913  
off N Faeroes-off E Iceland, N Greenland

**T. variabilis** Hansen,1913  
off N Faeroes-off E Iceland-Spitsbergen

**T. grandis** Hansen,1913  
off S Iceland

**T. solidus** Hansen,1913  
off SW Iceland

**T. proctagon** Tattersall,1904  
off W Ireland, off Hebrides

**T. sandersi** Kudinova-Pasternak,1985  
NE Atlantic

**Peraeospinosus** Sieg,1986

**P. mixtus** (Hansen,1913)  
= *Typhlotanais mixtus* Hansen,1913 (TM)  
off N Faeroes-off Iceland-Davis Strait

**Meromonakantha** Sieg,1986

**M. macrocephala** (Hansen,1913)  
= *Typhlotanais macrocephala* Hansen,1913 (TM)  
off SW Iceland

**Leptognathiidae** Sieg,1973

**Nematotanais** Bird & Holdich,1986  
\* Which subtaxon?

**N. mirabilis** Bird & Holdich,1985 (TM)  
\* tube building  
Rockall Trough

**Tanabornmia** Gutu,1986  
\* Which subtaxon?

**T. cornicauda** Gutu,1986 (TM)  
Biscay

**Robustochelia** Kudinova-Pasternak,1983  
\* Which subtaxon?

**R. longa** Kudinova-Pasternak,1983 (TM)  
NE Atlantic

*Paraleptognathia* Kudinova-Pasternak, 1981 (TM *P. typica* Kudinova-Pasternak, 1981 - N Pacific)  
\* Which subtaxon?

*P. bacescui* Kudinova-Pasternak, 1985  
NE Atlantic

Akanthophoreinae Sieg, 1986

*Akanthophoreus* Sieg, 1986

*A. gracilis* (Krøyer, 1842)  
= *Tanais gracilis* Krøyer, 1842 (TM)  
= *Leptognathia gracilis* : Auctt.  
= *Tanais islandicus* G.O. Sars, 1877  
= *L. longiremis* : G.O. Sars, 1896 (p.p.)  
= *L. sarsii* Hansen, 1909  
= *L. henseni* Vanhöffen, 1907  
Öresund-Bohuslän-N Norway-White Sea-Arctic Seas-Iceland-Faeroes-E & W Br. Isles, W Ireland

*A. longiremis* (Lilljeborg, 1864)  
= *Leptognathia longiremis* (Lilljeborg, 1864)  
~~non~~ = *L. longiremis* : G.O. Sars, 1896  
Kattegat-Bohuslän-W Norway-Iceland-Jan Mayen-Davis Strait

*A. inermis* (Hansen, 1913)  
= *Leptognathia inermis* Hansen, 1913  
off N & E Iceland-E Greenland

*A. multiserratus* (Hansen, 1913)  
= *Leptognathia multiserrata* Hansen, 1913  
off E Iceland

*Araphura* Bird & Holdich, 1985

*A. filiformis* (Lilljeborg, 1864)  
= *Leptognathia filiformis* (Lilljeborg, 1864)  
Öresund-S, W & N Norway, E Br. Isles

*A. brevimana* (Lilljeborg, 1864)  
= *Leptognathia brevimana* (Lilljeborg, 1864) (TM)  
Öresund-Bohuslän-N Norway, W Ireland & Scotland, E Br. Isles, Italy

*Araphuroides* Sieg, 1986 (TM *A. parabreviremis* Sieg, 1986 - Antarctic)

*A. brevispina* (Bird & Holdich, 1985)  
= *Araphura brevispina* Bird & Holdich, 1985  
NE Atlantic

*Haplocope* G.O. Sars, 1882

*H. angusta* G.O. Sars, 1882  
S Norway-Trondheim-North

*Cryptocope* G.O. Sars, 1880

*C. abbreviata* (G.O. Sars, 1865)  
Oslofjord-Skagerrak-W Norway

*C. arctica* Hansen, 1886  
Kara Sea-Spitsbergen-Jan Mayen-around Iceland-E & N Greenland-Davis Strait

*Tanaella* Norman & Stebbing, 1886

*T. unguicillata* Norman & Stebbing, 1886  
off SW Iceland-W slope of Engl. Channel

*Collettea* Lang, 1973  
= *Strongylura* G.O. Sars, 1882, ~~non~~ van Hasselt, 1824

*C. cylindrata* (G.O. Sars, 1882)  
Gullmar Fjord-S & W Norway-Arctic Seas

*C. minima* (Hansen, 1913)  
off S Iceland-Davis Strait

*C. rotundicauda* Kudinova-Pasternak, 1983  
NE Atlantic

*Strongylurella* Hansen, 1913

*S. indivisa* Hansen, 1913  
off SW Iceland (deep water)

*Leptoganthiopsis* Holdich & Bird, 1986

*L. attenuata* Holdich & Bird, 1986 (TM - Scotland)  
Br. Isles

*Subulella* Holdich & Bird, 1986

*S. scotti* Holdich & Bird, 1986 (TM - Scotland)  
Br. Isles

*Pseudoparatanais* Lang, 1973  
\* Leptognathiinae?

*P. batei* (G.O. Sars, 1882)  
Norw. Sea, Shetlands, Faeroes, S Iceland, Br. Isles, France, Mediterranean

*Tanaopsis* G.O. Sars, 1896  
\* Leptognathiinae?

*T. graciloides* (Lilljeborg, 1864)  
= *Leptognathia laticaudata* G.O. Sars, 1882  
Kattegat-Oslofjord-S Norway-North Sea-Shetlands-Br. Isles-N France-Italy

*Stenotanais* Bird & Holdich, 1985  
\* Leptognathiinae?

*S. crassiseta* Bird & Holdich, 1985 (TM)  
NE Atlantic

*S. hamicauda* Bird & Holdich, 1985  
NE Atlantic

Leptognathiinae Sieg, 1973

*Leptognathia* G.O. Sars, 1882

*L. manca* G.O. Sars, 1882  
Skagerrak-W Norway-S Iceland, Isle of Man, Davis Strait

*L. dentifera* G.O. Sars, 1896  
Oslo Fjord, Skagerrak

*L. breviremis* (Lilljeborg, 1864)  
Kattegat-Bohuslän-S & W Norway-off Iceland-off Faeroes-off Jan Mayen-Greenland-Ireland

*L. voeringii* (G.O. Sars, 1877) Lang, 1971  
off W Norway-off S Jan Mayen-off N Faeroes

*L. hastata* Hansen, 1913  
off E & N Iceland

*L. latiremis* Hansen, 1913  
off E & N Iceland

*L. ventralis* Hansen, 1913  
off S & E Iceland-Jan Mayen-Davis Strait

*L. paramanca* Lang, 1958  
N France, Isle of Man

*L. rigida* (Bate & Westwood, 1868)  
Br. Isles?

*L. amdrupi* Hansen, 1913  
E & N Greenland

*L. armata* ?, 19??  
White Sea

*Leptognathiella* Hansen, 1913

*L. abyssi* Hansen, 1913 (TM)  
NE Atlantic

*L. subaequalis* (Hansen, 1913)  
= *Leptognathia subaequalis* Hansen, 1913 ö  
off E Iceland-off N Faeroes-Davis Strait

*L. spinicauda* Bird & Holdich, 1985  
NE Atlantic

*L. clivicola* Bird & Holdich, 1985  
Biscay

*Leptognathioides* Bird & Holdich, 1985

*L. polita* (Hansen, 1913)  
= *Leptognathia polita* Hansen, 1913 (TM)  
off N Faeroes, N Greenland

*L. vicina* (Hansen, 1913)  
= *Leptognathia vicina* Hansen, 1913  
NE Atlantic

*L. potens* Bird & Holdich, 1985  
Biscay

Anarthruridae Lang, 1971

Anarthrurinae Lang, 1971

*Anarthrura* G.O. Sars, 1882

*A. simplex* G.O. Sars, 1882  
Skagerrak-N North Sea-N Norway-E Scotland, W France

Agathotanaeinae Lang, 1971

*Agathotanais* Hansen, 1913

*A. ingolfi* Hansen, 1913  
off Northumberland, Rockall, off S Iceland,

*Allodaposia* Sieg, 1986

*Metagathotanais* Bird & Holdich, 1988

*M. insulcatus* Bird & Holdich, 1988 (TM)  
Deep NE Atlantic

*Paragathotanais* ?, 19??

*P. gracilis* Bird & Holdich, 1988  
Deep NE Atlantic

*P. robustus* Bird & Holdich, 1988  
Deep NE Atlantic

*P. nanus* Bird & Holdich, 1988  
Deep Biscay

*Paranarthrura* Hansen, 1913

*P. borealis* Bird & Holdich, 1989  
Deep NE Atlantic

*P. crassa* Bird & Holdich, 1989  
Deep NE Atlantic

*P. tridens* Bird & Holdich, 1989  
Deep NE Atlantic

*P. lusitanus* Bird & Holdich, 1989  
Deep Biscay

*P. intermedia* Kudinova-Pasternak, 1982  
? deep NE Atlantic, deep Mediterranean

Pseudotanaidae Sieg, 1973

*Mystriocentrus* Bird & Holdich, 1989

*M. serratus* Bird & Holdich, 1989  
NE Atlantic

*Parapseudotanais* Bird & Holdich, 1989

*P. abyssalis* Bird & Holdich, 1989 (TM)  
Biscay

Pseudotanaeinae Sieg, 1973

- Pseudotanaïs* G.O. Sars, 1882
- P. macrocheles* G.O. Sars, 1882  
Oslofj.-W & N Norway
- P. forcipatus* (Lilljeborg, 1864)  
Öresund-Bohuslän-N Norway-N North Sea, E & W Scotland, N & E Iceland
- P. lilljeborgi* G.O. Sars, 1882  
N Norway-Barents Sea-Iceland-Jan Mayen-White Sea-Arctic Seas, e.g. E, N & W Greenland
- P. jonesi* Sieg, 1977  
Isle of Man, W Scotland
- P. (P.) affinis* Hansen, 1886  
= *P. crassicornis* Hansen, 1886  
off Faeroes-off Iceland-Arctic Seas, e.g. N Greenland
- P. longipes* Hansen, 1913  
off SW Iceland
- P. mediterraneus* G.O. Sars, 1882  
Biscay, Mediterranean
- P. similis* Sieg, 1973  
Biscay
- P. siegi* Kudinova-Pasternak, 1985  
NE Atlantic
- P. (P.) denticulatus* Bird & Holdich, 1989  
NE Atlantic
- P. (P.) falcicula* Bird & Holdich, 1989  
NE Atlantic
- P. (P.) scalpellum* Bird & Holdich, 1989  
NE Atlantic
- P. (P.) spatula* Bird & Holdich, 1989  
NE Atlantic
- P. (P.) spicatus* Bird & Holdich, 1989  
NE Atlantic
- P. (P.) vulsella* Bird & Holdich, 1989  
NE Atlantic
- P. (P.) longispinus* Bird & Holdich, 1989  
Biscay
- P. (P.) colonus* Bird & Holdich, 1989  
Biscay
- Nototanaidæ Sieg, 1973
- Androtanaïs* Sieg, 1976  
\* Correct family?
- Nototanaïs* Richardson, 1906
- Tanaïssus* Norman & Scott, 1906
- T. lilljeborgi* (Stebbing, 1891)  
= *Leptochelia danica* Hansen, 1909  
S Baltic-Kattegat, Holland, N France, around Br. Isles
- T. elongatus* Jones & Holdich, 1983  
Shetlands
- Protanaïssus* Sieg, 1983 (™ *Typhlotanaïs longidactylus* Shiino, 1970)
- Bathytanaïssus* Bird & Holdich, 1989
- B. spinulosus* Bird & Holdich, 1989 (™)  
Biscay
- Pseudonototanaïs* Lang, 1973
- Nototanaoides* Sieg & Heard, 1985 (™ *N. trifurcatus* Sieg & Heard, 1985 - Gulf of Mexico)
- Pseudozeuxidæ Sieg, 1982  
\* No species within the area?
- Pseudozeuxo* Sieg, 1982 (™ *P. belizensis* Sieg, 1982 - Belize)
- Heterotanaoides* Sieg, 1973