(c)2001 Mineral Data Publishing, version 1.2

Crystal Data: Orthorhombic. *Point Group:* 2/m 2/m 2/m. Anhedral plates, flattened on $\{010\}$; as aggregates, to 15 mm.

Physical Properties: Cleavage: Good on $\{010\}$; poor on $\{101\}$, intersecting at 83°. Hardness = 5.5-6 D(meas.) = 2.55 D(calc.) = 2.53

Optical Properties: Transparent. Color: Pink with a violet hue. Luster: Vitreous, pearly on $\{010\}$ cleavage.

Optical Class: Biaxial (+). Orientation: X=a; Y=c; Z=b. $\alpha=1.522(1)$ $\beta=1.522(1)$ $\gamma=1.543(1)$ $2V(\text{meas.})=0^\circ-9^\circ$

Cell Data: Space Group: Pcab. a = 10.213(2) b = 15.878(4) c = 9.058(1) Z = 4

X-ray Powder Pattern: Ilímaussaq intrusion, Greenland. 3.117 (10), 4.355 (7), 3.880 (7), 4.447 (6), 3.388 (6), 3.313 (5), 2.576 (5)

Chemistry:

	(1)
SiO_2	65.00
${ m TiO}_2$	< 0.02
$\mathrm{Al_2O_3}$	< 0.02
Y_2O_3	0.40
FeO	0.05
MnO	1.59
MgO	< 0.02
CaO	8.49
Na_2O	22.06
K_2O	< 0.02
Total	97.59

(1) Ilímaussaq intrusion, Greenland; by electron microprobe, average of 16 analyses on two grains; TGA gave 4% weight loss at 600 °C; corresponds to $Na_{3.96}(Ca_{0.84}Mn_{0.12}Y_{0.02})_{\Sigma=0.98}Si_{6.01}O_{14}(OH)_2$.

Occurrence: As patches, streaks, and veinlets cutting arfvedsonite-bearing nepheline syenite; probably of late-stage hydrothermal origin.

Association: Villiaumite, analcime.

Distribution: On the Kvanefjeld Plateau, in the Ilímaussag intrusion, southern Greenland.

Name: For the type locality, which is Danish for the mountain where the angelica grows.

Type Material: University of Copenhagen, Copenhagen, Denmark, 1983.136, GGU 47887; Harvard University, Cambridge, Massachusetts, 127475; National Museum of Natural History, Washington, D.C., USA, 162234.

References: (1) Petersen, O.V. and O. Johnsen (1984) Kvanefjeldite, a new mineral species from the Ilímaussaq alkaline complex, southwest Greenland. Can. Mineral., 22, 465–467. (2) Johnsen, O., E.S. Leonardsen, L. Fälth, and H. Annehed (1983) Crystal structure of kvanefjeldite: the introduction of ${}_{\infty}^{2}$ [Si₃O₇OH] layers with eight-membered rings. Neues Jahrb. Mineral., Monatsh., 505–512. (3) (1985) Amer. Mineral., 70, 873 (abs. refs. 1 and 2).