

Trophic structure of megabenthic food webs along depth gradients in the South China Sea and off northeastern Taiwan

H. Y. Lin¹, P. Y. Lin¹, N. N. Chang¹, J. C. Shiao^{1,*}, S. J. Kao^{1,2}

¹Institute of Oceanography, College of Science, National Taiwan University No. 1, Sec. 4, Roosevelt Road, Taipei 106, Taiwan

²Research Center for Environmental Changes, Academia Sinica, No. 128, Academia Road, Sec. 2, Nankang, Taipei 115, Taiwan

*Corresponding author. Email: jcsiao@ntu.edu.tw

Marine Ecology Progress Series 501: 53–66 (2014)

Supplement. Additional data.

Table S1. Isotopic values of diets (stomach contents [S]) and muscle tissues [M]) of fish. n: number of individuals for isotope analysis; $\Delta\delta X$: $\delta X(M) - \delta X(S)$; SG is the major stomach content group. Ch: chyme; P: Pisces; Cr: crustacean; M: mollusc. For sample sizes >3, only mean value \pm SD are reported

Station	Species	n	M $\delta^{13}C$ (‰)	M $\delta^{15}N$ (‰)	S $\delta^{13}C$ (‰)	S $\delta^{15}N$ (‰)	$\Delta\delta^{13}C$ (‰)	$\Delta\delta^{15}N$ (‰)	SG
Stn 1	<i>Barathronus maculatus</i>	1	-19.2	14.6	-22.9	11.1	3.7	3.5	Ch
	<i>Aldrovandia phalacra</i>	1	-18.4	14.0	-19.2	12.0	0.8	2.0	Ch
Stn 2	<i>Chlorophthalmus acutifrons</i>	1	-18.9	11.2	-20.9	9.8	2.0	1.4	P
	<i>Tydemanina navigatoris</i>	1	-18.5	13.1	-22.2	8.5	3.7	4.6	Ch
	<i>Physiculus roseus</i>	1	-18.0	12.3	-18.8	10.3	0.8	1.9	Cr
	<i>Coelorinchus acutirostris</i>	1	-17.7	13.8	-22.3	5.6	4.6	8.2	Cr, M
	<i>Spicomacrurus kuronumai</i>	5	-18.6 \pm 0.2	11.6 \pm 0.2	-21.3 \pm 0.4	7.7 \pm 0.9	2.7 \pm 0.4	3.9 \pm 0.9	Cr, M
	<i>Parazen pacificus</i>	1	-18.6	11.0	-20.9	8.5	2.3	2.6	Cr
Stn 4	<i>Apogon semilineatus</i>	2	-17.1, -17.2	11.3, 11.2	-20.6, -22.1	9.4, 8.9	3.5, 4.9	1.9, 2.3	Cr
	<i>Rogadius asper</i>	4	-17.2 \pm 0.3	11.1 \pm 0.8	-20.9 \pm 1.2	7.6 \pm 0.9	3.7 \pm 0.9	3.5 \pm 1.0	Cr, P
	<i>Suggrundus meerdervoorti</i>	2	-17.1, -17.1	11.6, 11.7	-18.5, -20.4	9.9, 9.0	1.3, 3.3	1.7, 2.6	Cr

	<i>Saurida</i> sp.	1	-17.4	12.0	-21.8	8.7	4.4	3.3	Ch
	<i>Synodus lobeli</i>	3	-17.7 ± 0.3	11.5 ± 0.3	-19.0 ± 0.1	9.9 ± 0.4	1.3 ± 0.2	1.6 ± 0.5	Cr, P
	<i>Trachinocephalus myops</i>	3	-17.9 ± 0.3	11.3 ± 0.1	-20.8 ± 0.8	9.9 ± 0.2	2.9 ± 0.6	1.4 ± 0.1	Ch
	<i>Cynoglossus interruptus</i>	3	-17.1 ± 0.5	11.5 ± 0.5	-21.0 ± 0.6	9.0 ± 0.5	3.9 ± 0.8	2.5 ± 0.5	Ch
	<i>Nemipterus bathybius</i>	3	-17.0 ± 0.3	11.8 ± 0.2	-21.0 ± 0.2	7.9 ± 0.6	4.0 ± 0.4	3.9 ± 0.8	Cr, M
	<i>Parapercis sexfasciata</i>	1	-16.6	11.6	-20.4	8.5	3.8	3.1	Ch
	<i>Halieutaea stellata</i>	1	-17.3	12.6	-21.8	6.0	4.5	6.6	Cr
	<i>Engyprosoxon macroptera</i>	3	-18.0 ± 0.0	10.8 ± 0.3	-21.4 ± 0.3	8.6 ± 1.7	2.6 ± 1.5	1.9 ± 1.7	Ch
	<i>Laeops tungkongensis</i>	1	-17.8	10.1	-21.7	5.7	3.9	4.4	Ch
	<i>Scorpaenopsis neglecta</i>	1	-15.6	11.7	-17.7 ±	10.6	2.1	1.0	Cr
	<i>Gymnothorax reticularis</i>	2	-16.5	12.8	-18.9 ±	10.8	2.4	2.0	Cr, P
	<i>Aesopia cornuta</i>	1	-16.8	11.9	-19.7	9.6	2.9	2.3	Ch
Stn 5	<i>Alepocephalus bicolor</i>	1	-19.2	13.3	-20.8	9.3	1.7	4.0	Ch
	<i>Coryphaenoides marginatus</i>	1	-18.7	13.7	-21.0	8.3	2.4	5.4	Ch
	<i>Hymenocephalus lethoemus</i>	1	-18.8	11.9	-21.5	8.7	2.7	3.2	Ch
	<i>Hoplostethus melanopus</i>	1	-18.9	13.5	-21.0	10.2	2.1	3.3	Ch
	<i>Symphurus strictus</i>	1	-22.2	13.9	-26.6	6.3	4.4	7.6	Ch
	<i>Dicrolene tristis</i>	1	-18.2	14.6	-22.0	6.8	3.8	7.8	Ch
	<i>Monomitopus pallidus</i>	1	-18.2	13.7	-22.1	8.0	3.9	5.7	Ch
	<i>Anacanthobatis borneensis</i>	1	-19.0	12.8	-21.1	10.0	2.1	2.8	Ch
	<i>Bathyraja matsubarai</i>	1	-18.0	12.1	-21.6	8.7	3.6	3.5	Ch
	<i>Nettastoma parviceps</i>	1	-18.4	11.3	-20.3	7.3	1.9	4.0	Cr
Stn 6	<i>Ilyophis brunneus</i>	1	-17.4	12.4	-21.4	11.4	4.0	1.0	Ch
	<i>Synaphobranchus affinis</i>	1	-17.8	12.6	-18.8	11.4	1.0	1.3	Ch
Stn 7	<i>Aldrovandia affinis</i>	1	-25.7	13.9	-25.9	11.1	0.2	2.8	M
	<i>Alepocephalus umbriceps</i>	1	-26.4	13.0	-25.2	10.6	-1.2	2.4	P
Stn 8	<i>Synaphobranchus kaupii</i>	3	-19.6 ± 1.4	15.6 ± 0.9	-19.8 ± 0.5	13.8 ± 0.3	0.2 ± 2.2	1.8 ± 0.7	Ch
	<i>Conocara krefftii</i>	1	-17.3	12.7	-19.5	8.4	2.2	4.3	Ch
	<i>Monomitopus kumae</i>	1	-17.7	14.0	-19.8	12.7	2.1	1.3	Ch
	<i>Dicrolene tristis</i>	1	-16.9	15.6	-19.2	13.3	2.2	2.3	Cr
	<i>Halosauropsis macrochir</i>	5	-17.0 ± 0.2	14.3 ± 0.3	-19.3 ± 0.7	12.8 ± 0.5	2.3 ± 0.9	1.5 ± 0.3	Cr
	<i>Aldrovandia affinis</i>	4	-17.0 ± 0.2	15.6 ± 0.5	-19.9 ± 0.4	12.1 ± 1.3	2.9 ± 0.4	3.5 ± 0.8	Cr

Table S2. Isotopic and biological information of the benthic megafauna. n: sample size for isotope analysis, bf: benthic feeder, zf: zooplankton feeder, sf: suspension feeder; df: deposit feeder. Species code was used in Fig. 2. For sample sizes >3, only mean value \pm SD are reported

Station (depth)/ Taxonomic group	Species	n	Species code	Feeding type	Length (cm)	Weight (g)	$\delta^{13}\text{C}$ (‰)	$\delta^{15}\text{N}$ (‰)	C/N ratio
Stn 1 (1442 m)									
Alcyonacea	not identified	1		sf			-20.2	11.8	5.0
Decapoda	<i>Galacantha valdiviae</i>	1	Gv	bf		2.1	-18.0	13.1	3.7
Decapoda	<i>Heterocarpus longirostris</i>	1	Hl	bf		0.5	-19.3	9.2	3.7
Decapoda	<i>Hymenopenaeus neptunus</i>	2	Hn	bf		1.0	-18.6, -18.9	12.8, 13.3	3.7, 3.8
Decapoda	<i>Heterocarpus tricarinatus</i>	2	Ht	bf		4.15	-18.7, -18.7	10.8, 11.0	3.6, 3.7
Decapoda	<i>Nephropsis suhmi</i>	1	Ns	bf		3.6	-18.0	11.1	3.7
Decapoda	<i>Acantheephyra quadrispinosa</i>	1	Aq	zf		0.6	-19.1	9.4	3.7
Decapoda	<i>Lebbeus</i> sp.	1	Lsp	bf		0.7	-19.1	12.9	3.8
Decapoda	<i>Nematocarcinus</i> sp.	2	Nsp	bf		4.5	-18.4, -18.5	10.3, 10.7	3.6, 3.6
Decapoda	<i>Parapagurus richeri</i>	1	Pr	bf		0.81	-18.1	11.7	3.9
Decapoda	<i>Uroptychus nigricapillis</i>	3	Un	bf		0.42	-19.7 \pm 0.5	12.8 \pm 1.3	4.5
Ophiuroidea	not identified	1		df			-22.8	11.5	12.5
Holothuroidea	not identified	5		df			-17.6 \pm 1.6	13.9 \pm 1.3	4.5
Asteroidea	not identified	1		bf			-16.2	15.5	4.4
Teleostei	<i>Aldrovandia phalacra</i>	1	Ap	bf	20.3	6	-18.4	14.0	3.3
Teleostei	<i>Barathronus maculates</i>	1	Bm	bf	8.4	2	-19.2	14.6	3.5
Teleostei	<i>Porogadus guentheri</i>	1	Pg	bf	73.1	1201	-17.2	15.8	3.4
Stn 2 (342 m)									
Decapoda	<i>Hadropenaeus lucasii</i>	2	Hlu	bf		2.8	-18.4, -18.9	10.2, 10.3	-
Decapoda	<i>Paramunida cristata</i>	7	Pc	bf		1.1	-18.6 \pm 0.4	10.2 \pm 0.3	-
Decapoda	<i>Solenocera comata</i>	1	Sc	bf		1.3	-18.7	11.1	-
Teleostei	<i>Chlorophthalmus acutifrons</i>	1	Ca	bf	11.8	20	-18.9	11.2	3.5
Teleostei	<i>Coelorinchus acutirostris</i>	1	Cac	bf	20.8	34	-17.7	13.8	3.4
Teleostei	<i>Spicomacrus kuronumai</i>	5	Sku	bf	10.5 \pm 0.6	3.8 \pm 0.4	-18.6 \pm 0.2	11.6 \pm 0.2	3.5
Teleostei	<i>Parazen pacificus</i>	1	Pp	zf	8.5	11	-18.6	11.0	3.5
Teleostei	<i>Physiculus roseus</i>	1	Pro	bf	25.5	179	-17.7	13.8	3.4
Teleostei	<i>Tydemania navigatoris</i>	1	Tn	bf	9.5	25	-18.5	13.1	3.5
St.3 (3458 m)									
Bivalvia	<i>Cuspidaria abyssopacifica</i>	2	Cab	bf			-16.2, -17.4	15.3, 16.4	3.5, 3.6

Bivalvia	<i>Cetoconcha</i> sp.	2	Csp			-17.8, -17.8	13.5, 14.9	3.9, 39	
Bivalvia	<i>Pseudomalletia</i> sp.	7	Psp			-17.6 ± 0.7	15.5 ± 0.6	3.9 ± 0.3	
Gastropoda	<i>Bathyaemaea</i> sp.	1	Bsp			-22.2	2.4	3.7	
Gastropoda	<i>Daphnella</i> sp.	2	Dsp			-19.0, -19.5	7.7, 11.3	3.8, 3.9 ± 0.1	
Scaphopoda	<i>Fissidentalium horikoshii</i>	1	Fh			-18.1	13.5	3.7	
Decapoda	not identified	1	Csp1			-18.3	14.9		
Infraorder Caridea									
Decapoda	<i>Munidopsis centrina</i>	2	Mc	bf		-23.8, -23.8	2.3, 2.7	4.0, 4.0	
Decapoda	<i>Munidopsis protunda</i>	1	Mp	bf		-21.6	6.0		
Decapoda	<i>Parastylodactylus</i> sp.	5	Pasp	bf		-17.8 ± 0.5	15.2 ± 0.8	3.4 ± 0.0	
Decapoda	<i>Sergia talismani</i>	1	St	zf		-19.0	8.6	3.5	
Decapoda	<i>Gennadas</i> sp.	1	Gsp	zf		-19.5	8.1	3.6	
Decapoda	<i>Gnathophausia</i> sp.	2	Gnsp	zf		-19.8, -19.9	10.7, 10.8	4.3, 4.3	
Teleostei	<i>Dysomma</i> sp.	1	Dysp	bf	30.6	24	-17.6	13.6	3.5
Stn 4 (91 m)									
Actiniaria	not identified	1		sf			-17.2	9.1	3.1
Actiniaria	not identified	3		sf			-18.1 ± 1.0	8.9 ± 0.4	3.0 ± 0.1
Gastropoda	<i>Microgaza</i> sp.	1	Msp				-18.9	6.6	3.4
Gastropoda	<i>Ficus subintermedia</i>	3	Fs				-16.8 ± 0.2	8.8 ± 0.3	3.2 ± 0.0
Decapoda	<i>Izanami curtispina</i>	3	Ic	bf	3.0		-18.1 ± 0.2	10.9 ± 0.2	3.4 ± 0.0
Decapoda	<i>Metapenaeopsis dalei</i>	4	Md	bf			-18.3 ± 0.2	10.3 ± 0.1	3.6 ± 0.1
Decapoda	<i>Nikoides</i> sp.	2	Nisp	bf			-17.7, -17.9	11.1, 11.3	3.4, 3.5
Decapoda	<i>Quollastria imperialis</i>	3	Qi	bf	0.8		-21.5 ± 0.2	10.2 ± 0.4	7.7 ± 0.6
Decapoda	<i>Calappa philargius</i>	2	Cp	bf	6.8		-17.5, -17.9	10.3, 10.3	3.3, 3.5
Decapoda	<i>Charybdis variegata</i>	3	Cv	bf			-17.6 ± 0.4	10.7 ± 0.5	3.5 ± 0.1
Decapoda	<i>Lissocarcinus laevis</i>	1	Ll	bf	0.7		-18.3	10.1	3.7
Decapoda	<i>Portunus haanii</i>	4	Ph	bf	9.0		-17.3 ± 0.2	11.2 ± 0.5	3.4 ± 0.0
Decapoda	<i>Portunus argentatus</i>	4	Pa	bf	1.9		-17.4 ± 0.6	11.2 ± 0.2	3.3 ± 0.0
Decapoda	<i>Scyllarus martensii</i>	3	Sm	bf	0.6		-18.6 ± 0.2	9.7 ± 0.3	3.5 ± 0.0
Decapoda	<i>Sicyonia cristata</i>	2	Scr	bf	2.5		-17.2, -17.7	10.9, 11.0	3.5, 3.5
Decapoda	<i>Solenocera alticarinata</i>	4	Sa	bf			-18.0 ± 0.2	10.9 ± 0.1	3.5 ± 0.2
Decapoda	<i>Solenocera pectinulata</i>	1	Sp	bf			-18.2	10.9	3.6
Asteroidea	not identified	2		bf			-14.6, -14.7	11.0, 11.2	3.1, 3.1
Teleostei	<i>Psettina iijimae</i>	2	Pi	bf	2.1, 2.2	1.3, 1.5	-17.3, -17.6	9.3, 9.5	3.6, 3.6
Teleostei	<i>Laeops tungkongensis</i>	1	Lt	bf	12.1	24	-17.8	10.1	3.4
Teleostei	<i>Ablabys taenianotus</i>	1	At	bf	4.0	2.4	-16.4	10.2	3.4

Teleostei	<i>Engyproson macroptera</i>	5	Em	zf	11.1 ± 2.3	25.2 ± 16	-18.1 ± 0.1	10.6 ± 0.4	3.3 ± 0.2
Teleostei	<i>Callionymus planus</i>	5	Cpl	zf	9.5 ± 0.7		-17.6 ± 0.5	10.7 ± 0.2	3.4 ± 0.1
Teleostei	<i>Pseudorhombus pentophthalmus</i>	1	Ppe	bf	11.5	19	-17.7	10.9	3.3
Teleostei	<i>Upeneus japonicus</i>	6	Uj	bf	8.3 ± 2.6	12 ± 13	-18.0 ± 0.3	11.0 ± 0.6	3.3 ± 0.1
Teleostei	<i>Apogon semilineatus</i>	4	As	zf	6.3 ± 0.6	4.5 ± 1.7	-17.4 ± 0.4	11.0 ± 0.3	3.2 ± 0.1
Teleostei	<i>Apogon kiensis</i>	5	Ak	zf	4.9 ± 0.6	1.7 ± 0.9	-17.5 ± 0.5	11.6 ± 0.3	3.3 ± 0.1
Teleostei	<i>Rogadius asper</i>	4	Ra	bf	9.8 ± 5.7	23.3 ± 38	-17.2 ± 0.3	11.1 ± 0.8	3.2 ± 0.1
Teleostei	<i>Trachinocephalus myops</i>	3	Tm	bf	13.5 ± 0.9	29 ± 5.6	-17.9 ± 0.3	11.3 ± 0.1	3.2 ± 0.1
Teleostei	<i>Cynoglossus interruptus</i>	5	Ci	bf	12.0 ± 0.3	15.0 ± 12	-17.2 ± 0.4	11.5 ± 0.3	3.4 ± 0.0
Teleostei	<i>Aploactis aspera</i>	2	Aas	bf			-16.8, -17.1	11.5, 11.6	3.4, 3.4
Teleostei	<i>Synodus lobeli</i>	3	Sl	bf	10.8 ± 1.6	9.3 ± 3.5	-17.7 ± 0.3	11.5 ± 0.3	3.3 ± 0.1
Teleostei	<i>Parapercis sexfasciata</i>	1	Ps	bf	6.9	7	-16.6	11.5	3.3
Teleostei	<i>Suggrundus meerdervoorti</i>	2	Sme	bf	17.2, 17	67, 60	-17.1, -17.1	11.6, 11.7	3.3, 3.3
Teleostei	<i>Scorpaenopsis neglecta</i>	1	Sn	bf	14.5	163	-15.6	11.7	3.2
Teleostei	<i>Aesopia cornuta</i>	1	Ac	bf	17.3	78	-16.8	11.9	3.3
Teleostei	<i>Saurida</i> sp.	1	Ssp	bf	10.4	10	-17.4	12.0	3.4
Teleostei	<i>Nemipterus bathybius</i>	5	Nb	bf	7.4 ± 0.8	9.2 ± 3.1	-17.0 ± 0.4	12.0 ± 0.3	3.3 ± 0.0
Teleostei	<i>Halieutaea stellata</i>	1	Hs	bf	9.7	47	-17.2	12.6	3.2
Teleostei	<i>Gymnothorax reticularis</i>	2	Gr	bf	43.7, 41.5	93, 94	-16.4, -16.6	12.5, 13.1	3.2, 3.3
Stn 5 (595 m)									
Elasmobranchii	<i>Sinobatis borneensis</i>	1	Sb	bf	25.7	73	-19.0	12.8	3.7
Teleostei	<i>Alepocephalus bicolor</i>	1	Ab	bf	23.5	154	-19.2	13.3	4.5
Teleostei	<i>Coryphaenoides marginatus</i>	1	Cm	bf	2.15	81	-18.7	13.7	3.4
Teleostei	<i>Bathygadus nipponicus</i>	1	Bn	bf	27.5	70	-18.4	12.7	3.4
Teleostei	<i>Hymenocephalus lethoemus</i>	1	Hle	bf	11.4	5	-18.8	11.9	3.5
Teleostei	<i>Hoplostethus melanopus</i>	1	Hm	bf	8.5	25	-18.9	13.5	3.4
Teleostei	<i>Symphurus strictus</i>	1	Ss	bf	10.1	15	-22.2	13.9	3.4
Teleostei	<i>Dicrolene tristis</i>	1	Dt	bf	17	33	-18.2	14.7	3.7
Teleostei	<i>Monomitopus pallidus</i>	1	Mpa	bf	18.5	39	-18.2	13.7	3.4
Teleostei	<i>Bathyraja matsubarai</i>	1	Bma	bf	16.4	14	-18.0	12.1	3.5
Teleostei	<i>Nettastoma parviceps</i>	1	Np	zf	69	76	-18.4	11.3	3.5
Stn 6 (803 m)									
Teleostei	<i>Ilyophis brunneus</i>	1	Ib	bf	45.5	78	-17.4	12.4	3.4
Teleostei	<i>Synaphobranchus affinis</i>	1	Saf	bf	52	192	-17.8	12.6	3.4
Teleostei	<i>Avocettina infans</i>	1	Ai	zf	46	13	-18.4	10.6	3.5

Stn 7 (1525 m)

Decapoda	<i>AcanthePHYra examia</i>	1	Ae	zf		4.7	-24.5	11.2	3.3
Mysidacea	<i>Gnathophausia</i> sp.	1	Gnsp	zf		0.2	-18.8	10.9	3.7
Teleostei	<i>Aldrovandia affinis</i>	1	Aa	bf	53	109	-25.7	13.9	3.3
Teleostei	<i>Alepocephalus umbriceps</i>	1	Au	bf	29.2	232	-26.4	13.0	3.3
Teleostei	<i>Elassodiscus obscurus</i>	1	Eo	bf	22.4		-23.9	12.4	3.5

Stn 8 (1242 m)

Actiniaria	not identified	1		sf			-18.6	11.6	4.4
Actiniaria	not identified	4		sf			-20.7 ± 0.3	11.4 ± 0.3	6.2 ± 0.7
Gastropoda	<i>Borsonia</i> sp.	3	Bosp				-15.3 ± 0.4	14.7 ± 0.2	3.6 ± 0.1
Gastropoda	<i>Ceratoxancus</i> sp.	3	Cesp				-17.5 ± 0.1	12.6 ± 0.3	3.9 ± 0.0
Gastropoda	<i>Comitas</i> sp.	2	Cosp				-14.7, -14.8	12.8, 13.2	3.8, 3.9
Gastropoda	<i>Daphnella</i> sp.	2	Dsp				-15.8, -15.9	11.6, 13.2	3.5, 3.5
Gastropoda	<i>Manaria</i> sp.1	1	Msp1				-16.7	11.7	3.6
Gastropoda	<i>Manaria</i> sp.2	1	Msp2				-15.5	11.8	3.5
Gastropoda	<i>Spergo</i> sp.	1	Spsp				-15.9	13.5	3.8
Decapoda	<i>Parapagurus richeri</i>	4	Pr	bf			-18.0 ± 0.5	11.3 ± 0.9	3.6 ± 0.0
Decapoda	<i>Benthesicymus investigatoris</i>	4	Bi	zf			-18.3 ± 0.2	10.6 ± 0.4	3.6 ± 0.2
Decapoda	<i>Galacantha valdiviae</i>	2	Gv	bf	2.2, 3.5		-18.5, -18.5	12.2, 12.4 1	3.7, 3.8
Decapoda	<i>Heterocarpus longirostris</i>	2	Hlo	bf	8.1, 9.2		-18.3, -19.3	9.4, 11.3	3.4, 3.7
Decapoda	<i>Heterocarpus tricarinatus</i>	2	Ht	bf	12.7, 11.8		-17.8, -17.9	12.4, 12.4	3.5, 3.5
Decapoda	<i>Lamoha superciliosa</i>	2	Ls	bf	4.0, 5.0		-17.9, -18.2	11.6, 11.9	3.5, 3.5
Decapoda	<i>Nephropsis suhmi</i>	3	Ns	bf	6.1 ± 4.4		-17.7 ± 0.2	11.2 ± 0.1	3.5 ± 0.1
Decapoda	<i>Polycheles helleri</i>	2	Phe	bf	0.7		-18.5, -19.7	12.5, 12.6	3.6, 4.1
Holothuroidea	not identified	4		df			-18.1 ± 1.5	11.4 ± 1.0	5.0 ± 1.8
Ophiuroidea	not identified	1		df			-20.6	8.0	6.6
Asteroidea	not identified	1		bf			-14.1	14.8	3.4
Echinoidea	not identified	1		bf			-16.7	13.5	4.1
Teleostei	<i>Aldrovandia affinis</i>	4	Aa	bf	42.8 ± 1.7	45 ± 7	-17.0 ± 0.2	15.6 ± 0.5	3.4 ± 0.1
Teleostei	<i>Conocara krefftii</i>	1	Ck	bf	28.6	147	-17.3	12.7	3.3
Teleostei	<i>Dicrolene tristis</i>	1	Dt	bf	24	76	-16.9	15.6	3.3
Teleostei	<i>Halosauropsis macrochir</i>	6	Hma	bf	40.6 ± 5.6	111 ± 27	-17.0 ± 0.1	14.4 ± 0.3	3.3 ± 0.1
Teleostei	<i>Monomitopus kumae</i>	1	Mk	bf	18.2	44	-17.7	14.0	3.2
Teleostei	<i>Synphobranchus kaupii</i>	4	Sk	bf	63.8 ± 7.8	414 ± 223	-19.7 ± 1.4	15.3 ± 0.9	4.3 ± 1.6