

Full Length Research Paper

Antioxidant activities of phenolic components from various plants of *Desmodium* species

Jen-Chieh Tsai¹, Guan-Jhong Huang², Tai-Hui Chiu¹, Shyh-Shyun Huang², Shun-Chieh Huang¹, Tai-Hung Huang¹, Shang-Chih Lai³ and Chao-Ying Lee^{1*}

¹School of Pharmacy, College of Pharmacy, China Medical University, Taichung, Taiwan, Republic of China.

²Graduate Institute of Chinese Pharmaceutical Sciences, College of Pharmacy, China Medical University, Taichung, Taiwan, Republic of China.

³Department of Health and Nutrition Biotechnology, College of Health Science, Asia University, Taichung, Taiwan, Republic of China.

Accepted 10 April, 2017

The aim of this study was to examine the antioxidant activities and phenolic components of the crude extracts of 10 *Desmodium* species from Taiwan. In this study, DPPH free radical scavenging activity, ABTS radical monocation scavenging activity, ferric-reducing antioxidant power (FRAP) and reducing power of the 10 *Desmodium* species were evaluated for their antioxidant activities. The results showed that, of all the samples, *Desmodium sequax* was the most active in ABTS, DPPH, FRAP and reducing power assays. The total polyphenol, total flavonoid and total flavonol contents of the crude extract were calculated. The correlation coefficient (R^2) values of TEAC with phenolic compounds indicated strong positive correlations, except for total flavonoid content. Furthermore, HPLC chromatographic fingerprints were established, and chlorogenic acid and vitexin in *D. sequax* were quantified. Chlorogenic acid was confirmed to express strong antioxidant activities in ABTS, DPPH, FRAP and reducing power assays. The present study indicated the antioxidant activities of the 10 *Desmodium* species were related to their phenolic components. *D. sequax* is a potent antioxidant medicinal plant, and chlorogenic acid may be an important factor in the antioxidant activity of this plant.

Key words: *Desmodium*, antioxidant activity, chlorogenic acid, vitexin.

INTRODUCTION

As we know, overproduction of reactive oxygen species (ROS) such as superoxide anion (O_2^-), hydrogen peroxide, hydroxyl radical (OH^\bullet) and peroxy radical (ROO^\bullet) may induce oxidative stress in the human body, in consequence causing degenerative and pathological damages, such as aging, cancer, cardiovascular diseases, Alzheimer's disease and inflammation (Ames, 1983; Smith et al., 1996). Certain environmental factors, such as stress, cigarette smoking, and some drugs are also associated with elevation of free radicals in the human body. Therefore, antioxidants play an important role in protecting the body from oxidative stress by scavenging free radicals.

In recent years, antioxidant activities of herbs and

health foods are investigated comprehensively (Cai et al., 2004). Some medicinal plants or plant derived chemical compounds with anti-inflammatory, anti-necrotic, hepato-protective effect or other pharmaceutical activities have also been demonstrated to possess antioxidant and/or radical scavenging mechanisms, at least partly (Jimoh et al., 2010). The genus *Desmodium*, of the Fabaceae family, includes about 350 species distributed in tropical and subtropical zones worldwide. In Taiwan, there are 18 known species that belong to the *Desmodium* genus, found mostly in regions of medium and low altitudes. Other than being processed into green manures and forages, some of them can be used as herbal medicines. For example, *Desmodium gangeticum* has been demonstrated to possess antioxidant, anti-nociceptive, anti-inflammatory (Govindarajan et al., 2007; Rathi et al., 2004), antiemetic (Joshi and Parle, 2007), cardio-

*Corresponding author. E-mail: cylee@mail.cmu.edu.tw.
Tel/Fax: (+886) 4-2203-1075.

protective (Kurian and Philip, 2005), and anti-ulcer effects (Dharmani et al., 2005).

Phytochemical studies have progressively isolated gangetin (Purushothaman et al., 1971), salicylic acid, rutin (Mishra et al., 2005), desmodin, gangetinin (Purushothaman et al., 1975), chlorogenic acid and caffeic acid from *Desmodium gangeticum* (Govindarajan et al., 2006). *Desmodium triflorum* has been reported to express antioxidant (Lai et al., 2010), analgesic and anti-inflammatory activities (Lai et al., 2009), and 2"-O-glucosylvitexin, vitexin, isovitexin, apigenin, and (+)-pinitol have been isolated from this plant (Adinarayana and Syamasundar, 1982). Another plant from this genus, *Desmodium uncinatum*, has been shown to possess genistein, uncinanone A, uncinanone B, uncinanone C (Tsanuo et al., 2003), uncinanone D and uncinanone E (Guchu et al., 2007) with phytochemical studies. However, no studies to date have been able to demonstrate antioxidant effect of any *Desmodium* species other than *D. gangeticum* and *D. triflorum*.

The aim of this study was to investigate the antioxidant activities and phenolic contents of the crude extracts of *Desmodium gangeticum* (L.) DC. (DG), *Desmodium heterocarpon* (L.) DC. (DH), *Desmodium intortum* (DC.) Urb. (DI), *Desmodium microphyllum* (Thunb ex Murray) DC. (DM), *Desmodium renifolium* (L.) Schindl. (DR), *Desmodium scorpiurus* (Sw.) Desv. (DSC), *D. sequax* Wall. (DSE), *Desmodium tortuosum* (Sw.) DC. (DTO), *D. triflorum* (L.) DC. (DTR) and *D. uncinatum* DC. (DU). The antioxidant activities of the 10 *Desmodium* species were estimated by various experimental models, such as DPPH, ABTS, FRAP and reducing power assays. Additionally, HPLC fingerprints of the extracts were also established.

MATERIALS AND METHODS

Plant materials

Mature whole plants of the 10 *Desmodium* species were collected from various places in Taiwan. Voucher specimens were deposited in the National Museum of Natural Science (TNM), Taichung. They were authenticated in many aspects, including morphology (flowers, fruits, seeds and pollens), histological microscopic examination (leaves and stems) and ITS (Internal Transcribed Spacer) regions of rDNA. The sequential data was submitted to the National Center for Biotechnology Information (NCBI) genbank.

Preparation of plant extracts

Whole plants were dried in a circulating air oven, then ground and crushed into a coarse powder. The powder (500 g) was extracted with 2 L of 70% ethanol three times. The filtrates were collected and concentrated under reduced pressure with a vacuum rotary evaporator. The remaining solution was lyophilized before the final extract was obtained. The extract was stored in -20°C before use.

Chemicals

Folin-Ciocalteu's phenol reagent, 1,1-diphenyl-2-picrylhydrazyl radical (DPPH), sodium carbonate (Na₂CO₃), potassium ferricyanide (K₃Fe(CN)₆), ferric chloride (FeCl₃), aluminum chloride

hexahydrate (AlCl₃·6H₂O), trichloroacetic acid (TCA), 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt (ABTS), potassium peroxydisulfate (K₂S₂O₈), sodium phosphate dibasic (Na₂HPO₄), sodium phosphate monobasic (NaH₂PO₄), (+)-catechin, chlorogenic and vitexin were purchased from Sigma Aldrich Ltd (Steinheim, Germany). Trolox (6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid) and ascorbate were used as antioxidant standards. 2,4,6-tris(2-pyridyl)-s-triazine (TPTZ) and all solvents used were purchased from Merck (Darmstadt, Germany).

Fingerprint analysis by HPLC

HPLC fingerprint profiles were established for 0.1 mg/ml of the standards (chlorogenic acid, rutin and vitexin) and 10 mg/ml of the 10 samples. The HPLC instruments used in this study consisted of a Hitachi L-7200 autosampler, a Hitachi L-7100 HPLC solvent delivery pump, and a Hitachi L-7455 diode array detector. Analyses were performed with a LiChroCART Purospher Star RP-18e column (250 mm × 4 mm, i.d. 5 µm; Merck, Germany). 20 µl of each sample was filtered through a 0.22 µm minipore filter and injected into the column. The mobile phase consisted of (A) acetonitrile and (B) 0.2% aqueous formic acid, using a gradient elution of 5 to 14% A at 0 to 10 min, 14% A at 10 to 20 min, 14 to 28% A at 20 to 60 min and 28 to 55% A at 60 to 85 min. The flow rate was 0.7 ml/min, and the detection wavelength was 330 nm.

Determination of antioxidant activity by ABTS assay

The ABTS assay was performed as reported by Re et al. (1999). The antioxidant abilities of the extracts were indicated as Trolox Equivalent Antioxidant Capacity (TEAC). Briefly, an aqueous solution of ABTS (7 mM) was oxidized with potassium peroxydisulfate (2.45 mM) for 16 h in the dark. The ABTS solution was diluted with 95% ethanol to an absorbance of 0.75 ± 0.05 at 734 nm (Beckman UV-Vis spectrophotometer, Model DU640B). The reference standard Trolox was diluted into concentrations of 0, 15.625, 31.25, 62.5, 125, 250, 500 µM, and a standard curve was constructed. 20 µl of the samples (125 µg/ml) or Trolox were each mixed with 180 µl of ABTS solution and the absorbance was read. TEAC was expressed as micromole Trolox / mg dry weight of the extract.

Ferric reducing antioxidant power assay

The ferric reducing antioxidant power (FRAP) of the crude extracts was evaluated according to the method of Benzie and Strain (1999). At 593 nm, the formation of a blue colored ferrous tripyridyltriazine (Fe²⁺-TPTZ) from a yellow ferric tripyridyltriazine (Fe³⁺-TPTZ) due to reduction was measured. To prepare the FRAP reagent, 0.1 M acetate buffer (pH 3.6), 10 mM 2,4,6-tris(2-pyridyl)-s-triazine (TPTZ) and 20 mM ferric chloride (10:1:1, v/v/v) were mixed. 10 µl of each sample (250 µg/ml) was mixed with 300 µl of FRAP reagent, and the absorbance was read at 593 nm after 15 min. FeSO₄·7H₂O was used as the standard reference. A standard curve was established for FeSO₄·7H₂O at concentrations of 0, 31.25, 62.5, 125, 250, 500, 1000 µg/ml. The FRAP values were expressed in µmole Fe²⁺/mg dry weight of the extract.

Determination of antioxidant activity by DPPH radical scavenging ability

The capacity of the samples in scavenging 1,1-diphenyl-2-picrylhydrazyl radical (DPPH) stable free radicals was measured according to the method of Yamaguchi et al. (1998). The stock solutions (10 mg/ml) were diluted with 70% ethanol into different

concentrations (2000, 1000, 500, 250, 125 and 62.5 µg/ml). The wells in a 96-well plate were pipetted with 25 µl of sample solutions and 175 µl of 0.3 mM DPPH solution, and then left to stand at room temperature for 30 min in the dark before the absorbance was measured at 517 nm. The inhibition percentage (%) of radical scavenging activity was calculated according to the following equation:

$$\text{Inhibition (\%)} = (A_0 - A_s)/A_0 \times 100$$

where A_0 is the absorbance of the control and A_s is the absorbance of the sample at 517 nm.

IC₅₀ value was the effective concentration at which 50% of DPPH radicals were scavenged and was obtained by interpolation from linear regression analysis. A lower IC₅₀ value indicated a greater antioxidant activity.

Reducing power assay

The reducing power of the crude extracts was determined according to the method described by Wu et al. (2007). The standard material, ascorbic acid, was diluted with methanol into 250, 125, 62.5, 31.3 and 15.6 µg/ml concentrations. 25 µl of each sample (250 µg/ml) was mixed with 50 µl of 50 µM phosphate buffer (pH 6.6) and 50 µl of 0.1% (w/v) potassium ferricyanide. The mixture was incubated in a water bath at 50°C for 20 min. Following this, 100 µl of 1% (w/v) trichloroacetic acid solution was added before the mixture was centrifuged at 3000 rpm for 10 min. An aliquot of 175 µl of the upper layer was combined with 25 µl of 5 mM ferric chloride, and the absorbance of the reaction mixture was measured at 700 nm. The reducing power was expressed as µg of ascorbate equivalent per mg dry weight of the extract.

Determination of total polyphenol content

The total phenolic compound contents of the extracts were estimated by the Folin-Ciocalteu method (Ragazzi and Veronese, 1973). 20 µl of the sample extracts (250 µg/ml) were added into the wells of a 96-well plate, followed by adding 40 µl of Folin-Ciocalteu's phenol reagent and 200 µl of distilled water into the wells. The mixtures were left to stand at room temperature for 5 min before 40 µl of 20% sodium carbonate was added. The absorbance was recorded at 680 nm. (+)-Catechin was used for the construction of the standard curve. The total phenolic compound contents were expressed in µg of (+)-catechin equivalents per mg of dry weight.

Determination of total flavonoid content

The total flavonoid content of the crude extracts was determined according to the method of Lamaison and Carnet (1990). 100 µl of each sample extract was added to an equal volume of 2% AlCl₃·6H₂O solution (2 g in 100 ml methanol). The mixtures were vigorously shaken, and after 10 min of incubation, the absorbance was read at 430 nm. Rutin was used as the standard for the calibration curve, and the total flavonoid content was derived from the linear equation of this calibration curve. The total flavonoid content was expressed as µg of rutin equivalent per mg of dry weight.

Determination of total flavonol content

The total flavonol content of the crude extracts was estimated by

the method of Arnous et al. (2001). 40 µl of each sample extract (500 µg/ml) was added to 200 µl of 0.1% *p*-dimethylaminocinnamaldehyde (DMACA) in methanol/HCl (3:1, V/V). The mixtures were shaken and then left to stand at room temperature for 10 min. The absorbance of the reaction solutions was measured at 640 nm. (+)-Catechin was used as the standard for the calibration curve. The total flavonol content was calibrated using the linear equation based on the calibration curve. The total flavonol content was expressed as µg (+)-catechin equivalent/mg dry weight.

Statistical analysis

Experimental results were presented as the mean ± standard deviation (SD) of three parallel measurements. The statistical analyses were performed by one-way ANOVA, followed by Dunnett's *t* test. Differences were considered statistically significant when the *p* values were less than 0.05.

RESULTS AND DISCUSSION

Extraction yield

The extraction yields of the 10 *Desmodium* species are shown in Table 1. The extraction yields ranged from 6.74 to 29.62%, and *D. microphyllum* had the highest extraction yield.

HPLC fingerprint analysis of the 10 *Desmodium* species

HPLC chromatographic fingerprint profiles of the 10 *Desmodium* samples are shown in Figure 1. Variations in fingerprint profiles among the species signify differences in chemical constituents, which in turn indicate discrepant pharmaceutical activities. Furthermore, different fingerprint profiles could help in identifying these species effectively. Regarding the compounds identified, chlorogenic acid and rutin has already been reported to be present in *D. gangeticum*, and vitexin has been isolated from *D. triflorum*. Our chromatographic fingerprinting profiles were in agreement with the findings from previous studies.

Additionally, chlorogenic acid and vitexin were found in *D. sequax* by comparing the retention times and UV spectra of authentic standards analyzed under identical analytical conditions, and also confirmed by spiking the extracts with pure standards (standard addition method). The retention times of chlorogenic acid and vitexin were 17.36 and 35.41 min respectively. The quantitative data of chlorogenic acid and vitexin in *D. sequax* were calculated using their respective concentration vs. peak area calibration curves. According to the calibration curve, the contents of chlorogenic acid and vitexin were 7.96 mg/g of extract and 2.85 mg/g of extract, respectively.

Table 1. Extraction yields, DPPH IC₅₀ values, and reducing powers of the extracts of 10 *Desmodium* species.

Species and positive controls	Extraction yield ^a (%)	DPPH ^b IC ₅₀ (ig/ml)	Reducing power ^b (ig ascorbate/ mg sample)
<i>D. gangeticum</i> (DG)	13.91	688.73 ± 5.67	66.92 ± 2.54
<i>D. heterocarpon</i> (DH)	21.68	258.21 ± 7.31	110.37 ± 2.53
<i>D. intortum</i> (DI)	20.19	451.76 ± 6.48	71.16 ± 2.47
<i>D. microphyllum</i> (DM)	29.62	286.94 ± 5.54	103.88 ± 0.64
<i>D. renifolium</i> (DR)	22.78	>4000	22.34 ± 0.69
<i>D. scorpiurus</i> (DSC)	10.13	852.87 ± 20.21	48.40 ± 1.56
<i>D. sequax</i> (DSE)	8.68	201.19 ± 12.77	148.37 ± 3.48
<i>D. tortuosum</i> (DTO)	14.75	3390.93 ± 53.81	28.84 ± 0.68
<i>D. triflorum</i> (DTR)	13.74	1179.31 ± 21.72	41.30 ± 1.38
<i>D. uncinatum</i> (DU)	6.74	446.56 ± 10.44	80.01 ± 0.39
Chlorogenic acid	—	167.47 ± 5.64	270.48 ± 3.89
Vitexin	—	3537.94 ± 68.72	18.59 ± 0.21

^a Dried weight basis; ^b Values represented mean ± S.D. of three parallel measurements (P<0.05).




Figure 1. HPLC chromatographs of the 10 *Desmodium* species and the standards.




Figure 2. TEAC values of the extracts of the 10 *Desmodium* species. Each value represents the mean \pm S.D. of three parallel measurements ($P < 0.05$).

DPPH and ABTS scavenging activity

DPPH and ABTS assays are commonly used for the assessment of free radical-scavenging abilities of herbal medicines and health foods due to their simplicity, stability, accuracy and reproducibility (Vijaya et al., 2010). ABTS is a synthetic radical that can be used to estimate scavenging activities for both polar and non-polar samples (Shi et al., 2009), and it is often expressed in TEAC values. In this study, TEAC values of the 10 *Desmodium* species were determined from the Trolox calibration curve as shown in Figure 2. The antioxidant activities of the 10 *Desmodium* species were in the following decreasing order: DSE > DH > DM > DU > DI > DG > DSC > DTR > DTO > DR.

DPPH is a stable radical which can accept hydrogen from an antioxidant to form reduced DPPH (Moon and Shibamoto, 2009). IC_{50} values of the 10 *Desmodium* species were investigated using the DPPH colorimetric method (Table 1). The low IC_{50} values of *D. sequax*, *D. heterocarpon* and *D. microphyllum* indicated that they expressed strong antioxidant activities in the DPPH assay. The aforesaid results revealed that ABTS and DPPH assays were similar. Since both DPPH and ABTS are used to evaluate free radical-scavenging abilities of the samples, our findings revealed consistent results between these two assays.

Previous studies have reported that *D. gangeticum* showed antioxidant activity in the DPPH assay (Govindarajan et al., 2003), and a similar result was obtained in this study. However we discovered that *D.*

sequax, *D. heterocarpon*, *D. microphyllum*, *D. uncinatum* and *D. intortum* showed even stronger antioxidant activities in comparison with *D. gangeticum* in both ABTS and DPPH assays. Of all the investigated plant species, *D. sequax* had the highest free radical scavenger activity. *D. sequax* and *D. heterocarpon* are used in folk medicine, and *D. microphyllum* is a traditional Chinese herbal medicine. Therefore, it is worthy to investigate the pharmacological activities of these medicines in the future.

Ferric reducing antioxidant power assay

As shown in Figure 3, the total antioxidant powers (FRAP) of the 10 *Desmodium* species differed significantly. *D. sequax* expressed the highest FRAP value, followed by DM, DH, DG, DU, DI, DSC, DTR, DTO and DR. Therefore *D. sequax*, *D. microphyllum*, and *D. heterocarpon* expressed strong activities, and *D. sequax* had the strongest activity. In the present study, FRAP, DPPH and ABTS assays were used to estimate the total antioxidant power.

The relationship between FRAP and ABTS and between FRAP and DPPH ($1/IC_{50}$) values of the 10 *Desmodium* species were appraised and expressed as correlation coefficients (R^2). R^2 values of FRAP/ABTS (Figure 4A) and FRAP/DPPH (Figure 4B) were 0.8447 and 0.9224 respectively. The results showed high positive correlations between these assays, and thus revealed high similarity between these assays.




Figure 3. FRAP values of the extracts of the 10 *Desmodium* species. Each value represents the mean \pm S.D. of three parallel measurements ($P < 0.05$).




Figure 4. Correlation coefficients (R^2) of (A) FRAP and TEAC and (B) FRAP and DPPH of the extracts of the 10 *Desmodium* species.

Measurement of reducing power

Previous researches have conceived that antioxidant activities of herbal medicines and health foods are probably related to their reducing powers (Wu et al., 2007). The reducing powers of the 10 samples are shown

in Table 1, and the results indicated large variations in antioxidant activities. The reducing powers of the 10 samples ranged from 22.34 ± 0.69 to 148.37 ± 3.48 μg ascorbate/mg and were in the following decreasing order: DSE>DH>DM>DU>DI>DG>DSC>DTR>DTO> DR.

Table 2. Total polyphenol, flavonoid, and flavonol contents of the extracts of 10 *Desmodium* species^a.

Species	Total polyphenols (ig CE/mg) ^b	Total flavonoids (ig RE/mg) ^c	Total flavonols (ig CE/mg) ^b
<i>D. gangeticum</i> (DG)	173.28 ± 3.07	127.49 ± 2.73	2.62 ± 0.01
<i>D. heterocarpon</i> (DH)	266.50 ± 6.10	75.79 ± 2.66	88.54 ± 0.72
<i>D. intortum</i> (DI)	144.31 ± 5.96	68.64 ± 1.74	57.43 ± 0.18
<i>D. microphyllum</i> (DM)	249.85 ± 3.02	387.73 ± 11.58	45.95 ± 0.26
<i>D. renifolium</i> (DR)	25.50 ± 0.42	45.01 ± 0.52	1.56 ± 0.05
<i>D. scorpiurus</i> (DSC)	90.04 ± 2.28	42.24 ± 1.26	37.01 ± 0.50
<i>D. sequax</i> (DSE)	368.05 ± 8.99	153.32 ± 1.50	138.28 ± 0.92
<i>D. tortuosum</i> (DTO)	20.92 ± 0.19	122.52 ± 1.40	0.93 ± 0.01
<i>D. triflorum</i> (DTR)	85.50 ± 0.77	80.33 ± 0.38	56.48 ± 0.14
<i>D. uncinatum</i> (DU)	178.88 ± 3.17	203.93 ± 2.93	64.42 ± 0.09

^a Values represented mean ± S.D. of three parallel measurements. ^b Data expressed in ig (+)-catechin equivalent/mg dry weight (ig CE/mg). ^c Data expressed in ig rutin equivalent/mg dry weight (ig RE/mg).

These results suggested that *D. sequax*, *D. heterocarpon* and *D. microphyllum* had strong abilities to react with free radicals and could convert them into more stable nonreactive forms that eventually led radical chain reactions to termination. *D. sequax* was also demonstrated to possess the highest reducing activity. Similar results were observed for the reducing power assay as compared to the ABTS and DPPH assays, and these were consistent with the free radical scavenging capacities.

Total polyphenol, flavonoid, and flavonol contents

The total polyphenol, flavonoid, and flavonol contents of the 10 *Desmodium* species are shown in Table 2. The results revealed the total polyphenol contents of the 10 *Desmodium* species extracts varied from 20.92 to 368.05 mg (+)-catechin equivalent/ig dry weight, in the following decreasing order: DSE > DH > DM > DU > DG > DI > DSC > DTR > DR > DTO. The total flavonoid contents varied from 42.24 to 387.73 mg rutin equivalent/ig of dry weight, in the following decreasing order: DM > DU > DSE > DG > DTO > DTR > DH > DI > DR > DSC. The total flavonol contents of the 10 *Desmodium* species extracts varied from 0.93 to 138.28 mg (+)-catechin equivalent/ig dry weight, in the following descending order: DSE > DH > DU > DI > DTR > DM > DSC > DG > DR > DTO.

Relationships between total antioxidant power and total polyphenol, flavonoid, and flavonol contents

Phenolic compounds play important roles in stabilizing lipid oxidation and may be attributed directly to antioxidative activities (Yen et al., 1993). Both flavonoids and flavonols are polyphenolic compounds. The

correlation coefficients (R^2) between TEAC and total polyphenol, TEAC and total flavonoid, and TEAC and total flavonol of the 10 sample extracts were evaluated in this study. As shown in Figure 5, the R^2 values between TEAC and total polyphenol content and TEAC and total flavonol content of the 10 *Desmodium* species extracts were 0.9383 and 0.8199 respectively. However, the R^2 value between TEAC and total flavonoid (Figure 5B) was much lower ($R^2=0.1169$). The above results suggested strong linear relationships between TEAC and total polyphenol/total flavonol; therefore the higher the TEAC activity, the higher the total polyphenol and flavonol contents in the samples.

Antioxidant activities of chlorogenic acid and vitexin

In this study, the extract of *D. sequax* was found with the highest antioxidant activity, HPLC analysis also revealed the presence of chlorogenic acid and vitexin. The antioxidant activities of these two compounds were determined. In ABTS assay, the TEAC values of chlorogenic acid and vitexin were 2.04 ± 0.02 imole Trolox / mg and 0.14 ± 0.00 imole Trolox / mg respectively (Figure 3). In DPPH assay, chlorogenic acid had a lower IC_{50} value in comparison with vitexin (Table 1).

In FRAP assay, chlorogenic acid expressed a higher FRAP value than vitexin (Figure 2). In the reducing power assay, we found that chlorogenic acid was more active in comparison with vitexin (Table 1). Chlorogenic acid has been reported to exhibit antioxidant, anti-bacterial, anti-carcinogenic and anti-inflammatory activities (Shan et al., 2009). Our results of antioxidant activities were in agreement with the previous reports. Therefore, since chlorogenic acid possesses strong antioxidant activities, it may be an essential ingredient in the antioxidant activity of *D. sequax*.

Figure 5




Figure 5. Correlation coefficients (R^2) of (A) TEAC and total polyphenol content; (B) TEAC and total flavonoids content, and (C) TEAC and total flavonol content of the 10 *Desmodium* species.

CONCLUSIONS

In the present study, the antioxidant capacities of the 10 *Desmodium* species were examined with *in vitro* experiments, including ABTS radical scavenging assay, FRAP method, DPPH radical scavenging assay and reducing power method. The results demonstrated that most samples expressed strong activities and *D. sequax* exhibited the highest antioxidant potency. There are significant relationships between the antioxidant activities and amount of phenolic compounds of the *Desmodium* species, except for flavonoids. The chromatographic fingerprints of the 10 *Desmodium* species were established, and some phytochemicals were found. Furthermore, chlorogenic acid and vitexin were found in *D. sequax*, and the contents of these two compounds were examined by HPLC. Chlorogenic acid has been shown to be strongly antioxidative and may be an important component in the antioxidant activity of *D. sequax*.

ACKNOWLEDGMENTS

The authors would like to acknowledge Dr. Hsin-Fu Yen from the National Museum of Natural Science (TNM), Taichung, Taiwan for kindly helping with the collection of plants.

REFERENCES

- Adinarayana D, Syamasundar KV (1982). Occurrence of a rare diholosylflavone 2"-O-glucosylvitexin in *Desmodium triflorum*. *Curr. Sci.*, 51: 936-937.
- Ames BN (1983). Dietary carcinogens and anticarcinogens: oxygen radicals and degenerative diseases. *Sci.*, 221: 1256-1264.
- Arnous A, Makris DP, Kefalas P (2001). Effect of principal polyphenolic components in relation to antioxidant characteristics of aged red wines. *J. Agric. Food Chem.*, 49: 5736-5742.
- Benzie IFF, Strain JJ (1996). The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": the FRAP assay. *Anal. Biochem.*, 239: 70-76.
- Cai Y, Luo Q, Sun M, Corke H (2004). Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci.*, 74: 2157-2184.
- Dharmani P, Mishra PK, Maurya R, Chauhan VS, Palit G (2005). *Desmodium gangeticum*: a potent anti-ulcer agent. *Indian J. Exp. Biol.*, 43: 517-521.
- Govindarajan R, Rastogi S, Vijayakumar M, Shirwaikar A, Rawat AKS, Mehrotra S, Pushpangadan P (2003). Studies on the antioxidant activities of *Desmodium gangeticum*. *Biol. Pharm. Bull.*, 26: 1424-1427.
- Govindarajan R, Vijayakumar M, Shirwaikar A, Rawat AKS, Mehrotra S, Pushpangadan P (2006). Antioxidant activity of *Desmodium gangeticum* and its phenolics in arthritic rats. *Acta Pharm.*, 56: 489-496.
- Govindarajan R, Vijayakumar M, Rao CV, Shirwaikar A, Kumar S, Rawat AKS, Pushpangadan P (2007). Anti-inflammatory and antioxidant activities of *Desmodium gangeticum* fractions in carrageenan-induced inflamed rats. *Phytother. Res.*, 21: 975-979.
- Guchu SM, Yenesew A, Tsanuo MK, Gikonyo NK, Pickett JA, Hooper AM, Hassanali A (2007). C-methylated and C-prenylated isoflavonoids from root extract of *Desmodium uncinatum*. *Phytochemistry*, 68: 646-651.

- Jimoh FO, Adedapo AA, Afolayan AJ (2010). Assessing the polyphenolic, nutritive and biological activities of acetone, methanol and aqueous extracts of *Rumex sagittatus* Thunb. Afr. J. Pharm. Pharmacol., 4: 629-635.
- Joshi H, Parle M (2007). Pharmacological evidences for the anti-amnesic effects of *Desmodium gangeticum* in mice. Iranian J. Pharm. Res., 6: 199-207.
- Kurian GA, Philip S, Varghese T (2005). Effect of aqueous extract of the *Desmodium gangeticum* DC root in the severity of myocardial infarction. J. Ethnopharmacol., 97: 457-461.
- Lai SC, Peng WH, Huang SC, Ho YL, Huang TH, Lai ZR, Chang YS (2009). Analgesic and anti-inflammatory activities of methanol extract from *Desmodium triflorum* DC in mice. Am. J. Chin. Med., 37: 573-588.
- Lai SC, Ho YL, Huang SC, Huang TH, Lai ZR, Wu CR, Lian KY, Chang YS (2010). Antioxidant and antiproliferative activities of *Desmodium triflorum* (L.) DC. Am. J. Chin. Med., 38: 329-342.
- Lamaison JLC, Carnet A (1990). Teneurs en principaux flavonoids des fleurs de *Crataegus monogyna* Jacq et de *Crataegus laevigata* (Poiret D. C) en fonction de la vegetation. Pharm. Acta. Helv., 65: 315-320.
- Mishra PK, Singh N, Ahmad G, Dube A, Maurya R (2005). Glycolipids and other constituents from *Desmodium gangeticum* with antileishmanial and immunomodulatory activities. Bioorg. Med. Chem. Lett., 15: 4543-4546.
- Moon JK, Shibamoto T (2009). Antioxidant assays for plant and food component. J. Agric. Food Chem., 57: 1655-1666.
- Purushothaman KK, Kishore VM, Narayanaswami V, Connolly JD (1971). The structure and stereochemistry of gangetin, a new pterocarpan from *Desmodium gangeticum* (Leguminosae). J. Chem. Soc. C., 20: 2420-2422.
- Purushothaman KK, Chandrasekharan S, Balakrishna K, Connolly JD (1975). Gangetinin and desmodin, two minor pterocarpanoids of *Desmodium gangeticum*. Phytochem., 14: 1129-1130.
- Ragazzi E, Veronese G (1973). Quantitative analysis of phenolics compounds after thin-layer chromatographic separation. J. Chromatogr., 77: 369-375.
- Rathi A, Rao CV, Ravishankar B, De S, Mehrotra S (2004). Anti-inflammatory and anti-nociceptive activity of the water decoction *Desmodium gangeticum*. J. Ethnopharmacol., 95: 259-263.
- Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C (1999). Antioxidant activity applying an improved ABTS radical cation decolourization assay. Free Radic. Biol. Med., 26: 1231-1237.
- Shan J, Fu J, Zhao Z, Kong X, Huang H, Luo L, Yin Z (2009). Chlorogenic acid inhibits lipopolysaccharide-induced cyclooxygenase-2 expression in RAW264.7 cells through suppressing NF- κ B and JNK/AP-1 activation. Int. Immunopharmacol., 9: 1042-1048.
- Shi JY, Gong JY, Liu JE, Wu XQ, Zhang Y (2009). Antioxidant capacity of extract from edible flowers of *Prunus mume* in China and its active components. LWT-Food Sci. Technol., 42: 477-482.
- Smith MA, Perry G, Richey PL, Sayre LM, Anderson VE, Beal MF, Kowal N (1996). Oxidative damage in Alzheimer's. Nature, 382: 120-121.
- Tsanuo MK, Hassanali A, Hooper AM, Khan Z, Kaberia F, Pickett JA, Wadhams LJ (2003). Isoflavanones from the allelopathic aqueous root exudate of *Desmodium uncinatum*. Phytochem., 64: 265-273.
- Vijaya Kumar Reddy C, Sreeramulu D, Raghunath M (2010). Antioxidant activity of fresh and dry fruits commonly consumed in India. Food Res. Int., 43: 285-288.
- Wu CR, Huang MY, Lin YT, Ju Y, Ching H (2007). Antioxidant properties of *Cortex Fraxini* and its simple coumarins. Food Chem., 104: 1464-1471.
- Yamaguchi T, Takamura H, Matoba T, Terao J (1998). HPLC method for evaluation of the free radical-scavenging activity of foods by using 1,1-diphenyl-2-picrylhydrazyl. Biosci. Biotechnol. Biochem., 62: 1201-1204.
- Yen GC, Duh PD, Tsai CL (1993). Relationship between antioxidant activity and maturity of peanut hulls. J. Agric. Food Chem., 41:67-70.